



(51) International Patent Classification:

C07D 498/18 (2006.01) A61P 35/00 (2006.01)

C07D 519/00 (2006.01)

(21) International Application Number:

PCT/US2019/029737

(22) International Filing Date:

29 April 2019 (29.04.2019)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/665,435 01 May 2018 (01.05.2018) US

62/752,874 30 October 2018 (30.10.2018) US

62/836,036 18 April 2019 (18.04.2019) US

(71) Applicant: REVOLUTION MEDICINES, INC.

[US/US]; 700 Saginaw Drive, Redwood City, California 94063 (US).

(72) Inventors: PITZEN, Jennifer; c/o Revolution Medicines,

Inc., 700 Saginaw Drive, Redwood City, California 94063 (US). GLIEDT, Micah James Evans; c/o Revolution Medicines, Inc., 700 Saginaw Drive, Redwood City, California 94063 (US). BURNETT, G. Leslie; c/o Revolution Medicines, Inc., 700 Saginaw Drive, Redwood City, California 94063 (US). AGGEN, James Bradley; c/o Revolution Medicines, Inc., 700 Saginaw Drive, Redwood City, California 94063 (US). KISS, Gert; c/o Revolution Medicines, Inc., 700 Saginaw Drive, Redwood City, California 94063 (US). CREGG, James Joseph; c/o Revolution Medicines, Inc., 700 Saginaw Drive, Redwood City, California 94063 (US). SEMKO, Christopher Michael; c/o Revolution Medicines, Inc., 700 Saginaw Drive, Redwood City, California 94063 (US). WON, Walter; c/o Revolution Medicines, Inc., 700 Saginaw Drive, Redwood City, California 94063 (US). WANG, Gang; c/o Revolution Medicines, Inc., 700 Saginaw Drive, Redwood City, California 94063 (US). LEE, Julie Chu-Li; c/o Revolution Medicines, Inc., 700 Saginaw Drive, Redwood City, California 94063 (US). THOTTUMKARA, Arun P.; c/o Revolution Medicines, Inc., 700 Saginaw Drive, Redwood City, California 94063 (US). GILL, Adrian Liam; c/o Revolution Medicines, Inc., 700 Saginaw Drive, Redwood City, California 94063 (US). MELLEME, Kevin T.; c/o Revolution Medicines, Inc., 700 Saginaw Drive, Redwood City, California 94063 (US).

(74) Agent: TONG, Connie C. et al.; Cooley LLP, 1299 Penn-

sylvania Avenue, NW, Suite 700, Washington, District of Columbia 20004-2400 (US).

(81) Designated States (unless otherwise indicated, for every

kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every

kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))

Published:

— with international search report (Art. 21(3))
 — before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))
 — with sequence listing part of description (Rule 5.2(a))

(54) Title: C40-, C28-, AND C-32-LINKED RAPAMYCIN ANALOGS AS MTOR INHIBITORS

(57) Abstract: The present disclosure relates to mTOR inhibitors. Specifically, the embodiments are directed to compounds and compositions inhibiting mTOR, methods of treating diseases mediated by mTOR, and methods of synthesizing these compounds.



WO 2019/212990 A1

C40-, C28-, AND C-32-LINKED RAPAMYCIN ANALOGS AS MTOR INHIBITORS**CROSS REFERENCE TO RELATED APPLICATIONS**

[0001] This application claims the benefit of U.S. Provisional Application No. 62/665,435, filed May 1, 2018 and U.S. Provisional Application No. 62/752,874, filed October 30, 2018 and U.S. Provisional Application No. 62/836,036, filed April 18, 2019, the contents of which are incorporated herein by reference in their entireties.

REFERENCE TO A SEQUENCE LISTING

[0002] The Sequence Listing associated with this application is provided in text format in lieu of a paper copy, and is hereby incorporated by reference into the specification. The name of the text file containing the Sequence Listing is REME_008_01WO_SeqList_ST25.txt. The text file is about 40 Kilo Bytes, was created on April 26, 2019, and is being submitted electronically via EFS-Web.

FIELD OF THE DISCLOSURE

[0003] The present disclosure relates to mTOR inhibitors. Specifically, the embodiments are directed to compounds and compositions inhibiting mTOR, methods of treating diseases mediated by mTOR, and methods of synthesizing these compounds.

BACKGROUND OF THE DISCLOSURE

[0004] The mammalian target of rapamycin (mTOR) is a serine-threonine kinase related to the lipid kinases of the phosphoinositide 3-kinase (PI3K) family. mTOR exists in two complexes, mTORC1 and mTORC2, which are differentially regulated, have distinct substrate specificities, and are differentially sensitive to rapamycin. mTORC1 integrates signals from growth factor receptors with cellular nutritional status and controls the level of cap-dependent mRNA translation by modulating the activity of key translational components such as the cap-binding protein and oncogene eIF4E.

[0005] mTOR signaling has been deciphered in increasing detail. The differing pharmacology of inhibitors of mTOR has been particularly informative. The first reported inhibitor of mTOR, Rapamycin is now understood to be an incomplete inhibitor of mTORC1. Rapamycin is a selective mTORC1 inhibitor through the binding to the FK506 Rapamycin Binding (FRB) domain of mTOR kinase with the aid of FK506 binding protein 12 (FKBP12). The FRB domain of mTOR is accessible in the mTORC1 complex, but less so in the mTORC2 complex. Interestingly, the potency of inhibitory activities against downstream

substrates of mTORC1 by the treatment of Rapamycin is known to be diverse among the mTORC1 substrates. For example, Rapamycin strongly inhibits phosphorylation of the mTORC1 substrate S6K and, indirectly, phosphorylation of the downstream ribosomal protein S6 which control ribosomal biogenesis. On the other hand, Rapamycin shows only partial inhibitory activity against phosphorylation of 4E-BP1, a major regulator of eIF4E which controls the initiation of CAP-dependent translation. As a result, more complete inhibitors of mTORC1 signaling are of interest.

[0006] A second class of “ATP-site” inhibitors of mTOR kinase were reported. This class of mTOR inhibitors will be referred to as TORi (ATP site TOR inhibitor). The molecules compete with ATP, the substrate for the kinase reaction, in the active site of the mTOR kinase (and are therefore also mTOR active site inhibitors). As a result, these molecules inhibit downstream phosphorylation of a broader range of substrates.


[0007] Although mTOR inhibition may have the effect of blocking 4E-BP1 phosphorylation, these agents may also inhibit mTORC2, which leads to a block of Akt activation due to inhibition of phosphorylation of Akt S473.

[0008] Disclosed herein, *inter alia*, are mTOR inhibitors. In some embodiments, compounds disclosed herein are more selective inhibitors of mTORC1 versus mTORC2. In some embodiments, compounds disclosed herein are more selective inhibitors of mTORC2 versus mTORC1. In some embodiments, compounds disclosed herein exhibit no selectivity difference between mTORC1 and mTORC2.

SUMMARY OF THE DISCLOSURE

[0009] The present disclosure relates to compounds capable of inhibiting the activity of mTOR. The present disclosure further provides a process for the preparation of compounds of the present disclosure, pharmaceutical preparations comprising such compounds and methods of using such compounds and compositions in the management of diseases or disorders mediated by mTOR.

[0010] The present disclosure provides a compound of Formula Ic:



(Ic)

or a pharmaceutically acceptable salt or tautomer thereof, wherein:

R^{32} is $-H$, $=O$, $-OR^3$, $-N_3$, or $-O-C(=Z^1)-R^{32a}$;

R^{28} is $-H$, $(C_1-C_6)alkyl$, or $-C(=Z^1)-R^{28a}$;


R^{40} is $-H$ or $-C(=Z^1)-R^{40a}$;

wherein when R^{28} and R^{40} are H , then R^{32} is not $=O$;

each Z^1 is independently O or S ;

R^{28a} , R^{32a} , and R^{40a} are independently $-A^1-L^1-A^2-B$; $-A^1-A^2-B$; $-L^2-A^1-L^1-A^2-L^3-B$; $-O-(C_1-C_6)alkyl$; or $-O-(C_6-C_{10})aryl$; wherein the aryl of $-O-(C_6-C_{10})aryl$ is unsubstituted or substituted with 1-5 substituents selected from $-NO_2$ and halogen;

A^1 and A^2 are independently absent or are independently selected from



wherein the bond on the left side of A^1 , as drawn, is bound to $-C(=Z^1)-$ or L^2 ; and
 wherein the bond on the right side of the A^2 moiety, as drawn, is bound to B or L^3 ;

each Q is independently 1 to 3 rings selected from arylene, cycloalkylene, heteroarylene, and heterocyclylene;

each X is independently absent or 1 to 2 rings selected from arylene, cycloalkylene, heteroarylene, and heterocyclylene;

each X^1 is independently a heteroarylene or heterocyclylene ring;


each W is independently absent or 1 to 2 rings selected from arylene, cycloalkylene, heteroarylene, and heterocyclylene;


each W^1 is independently a heteroarylene or heterocyclylene ring;

each G is independently absent or a ring selected from arylene, cycloalkylene, heteroarylene, and heterocyclylene;


each G^1 and G^2 are independently heteroarylene or heterocyclylene ring;

each L^1 is independently selected from






L^2 and L^3 are independently absent or are independently selected from







each B is independently selected from



each B¹ is independently selected from $-\text{NR}^3-(\text{C}(\text{R}^3)_2)_n-$, $-\text{NR}^3-(\text{C}(\text{R}^3)_2)_n-(\text{C}_6-\text{C}_{10})\text{arylene}-(\text{C}(\text{R}^3)_2)_n-$, $-\text{NR}^3-(\text{C}(\text{R}^3)_2)_n-\text{heteroarylene}-$, $-\text{NR}^3-(\text{C}(\text{R}^3)_2)_n-\text{heteroarylene}-(\text{C}(\text{R}^3)_2)_n-$, $-(\text{C}_6-\text{C}_{10})\text{arylene}-$, $-\text{NR}^3-(\text{C}(\text{R}^3)_2)_n-\text{NR}^3\text{C}(\text{O})-$,

$^{-3}\text{-NR}^3\text{-(C(R}^3\text{)}_2)_n\text{-heteroarylene-heterocyclylene-(C}_6\text{-C}_{10}\text{)arylene-}$, $^{-3}\text{-heteroarylene-}$

$\text{heterocyclylene-(C}_6\text{-C}_{10}\text{)arylene-}$, $\text{---C(=O)-(C(R}^3\text{)}_2)_p\text{---}$, $\text{---C(=O)-(C(R}^3\text{)}_2)_p\text{-heteroarylene-}$,



$\text{S(O)}_2\text{-arylene-C(O)-}$, and $^{-3}\text{-NR}^3\text{-(C(R}^3\text{)}_2)_n\text{-S(O)}_2\text{-arylene-(C(R}^3\text{)}_2)_n\text{-}$, wherein the $^{-3}\text{-}$ bond on the left side of B^1 , as drawn, is bound to A^2 , L^3 , or L^1 ; and wherein the heteroarylene, heterocyclylene, and arylene are each independently optionally substituted with alkyl, hydroxyalkyl, haloalkyl, alkoxy, halogen, or hydroxyl;

each R^3 is independently H or $(\text{C}_1\text{-C}_6)\text{alkyl}$;

each R^4 is independently H, $(\text{C}_1\text{-C}_6)\text{alkyl}$, halogen, 5-12 membered heteroaryl, 5-12 membered heterocyclyl, $(\text{C}_6\text{-C}_{10})\text{aryl}$, wherein the heteroaryl, heterocyclyl, and aryl are each independently optionally substituted with $-\text{N(R}^3\text{)}_2$, $-\text{OR}^3$, halogen, $(\text{C}_1\text{-C}_6)\text{alkyl}$, $-(\text{C}_1\text{-C}_6)\text{alkylene-heteroaryl}$, $-(\text{C}_1\text{-C}_6)\text{alkylene-CN}$, $-\text{C(O)NR}^3\text{-heteroaryl}$, or $-\text{C(O)NR}^3\text{-heterocyclyl}$;

each R^5 is independently H, $(\text{C}_1\text{-C}_6)\text{alkyl}$, $-\text{C(O)OR}^3$, or $-\text{N(R}^3\text{)}_2$, wherein the alkyl of $(\text{C}_1\text{-C}_6)\text{alkyl}$ is each independently optionally substituted with $-\text{N(R}^3\text{)}_2$ or $-\text{OR}^3$;


each R^6 is independently H, $(\text{C}_1\text{-C}_6)\text{alkyl}$, $-\text{C(O)OR}^3$, or $-\text{N(R}^3\text{)}_2$, wherein the alkyl of $(\text{C}_1\text{-C}_6)\text{alkyl}$ is each independently optionally substituted with $-\text{N(R}^3\text{)}_2$ or $-\text{OR}^3$;

each R^7 is independently H, $(\text{C}_1\text{-C}_6)\text{alkyl}$, $-\text{C(O)OR}^3$, or $-\text{N(R}^3\text{)}_2$, wherein the alkyl of $(\text{C}_1\text{-C}_6)\text{alkyl}$ is each independently optionally substituted with $-\text{N(R}^3\text{)}_2$ or $-\text{OR}^3$;

each R^8 is independently H, $(\text{C}_1\text{-C}_6)\text{alkyl}$, $-\text{C(O)OR}^3$, or $-\text{N(R}^3\text{)}_2$, wherein the alkyl of $(\text{C}_1\text{-C}_6)\text{alkyl}$ is each independently optionally substituted with $-\text{N(R}^3\text{)}_2$ or $-\text{OR}^3$;

each Y is independently $-\text{C}(\text{R}^3)_2$ or a bond;
 each n is independently an integer from one to 12;
 each o is independently an integer from zero to 30;
 each p is independently an integer from zero to 12;
 each q is independently an integer from zero to 30; and
 each r is independently an integer from one to 6.

[0011] The present disclosure provides a compound of Formula Ia:



or a pharmaceutically acceptable salt or tautomer thereof, wherein:

R^{32} is $-\text{H}$, $=\text{O}$, $-\text{OR}^3$, $-\text{N}_3$, or $-\text{O}-\text{C}(=\text{Z}^1)-\text{R}^{32a}$;

R^{28} is $-\text{H}$, $(\text{C}_1-\text{C}_6)\text{alkyl}$, or $-\text{C}(=\text{Z}^1)-\text{R}^{28a}$;


R^{40} is $-\text{H}$ or $-\text{C}(=\text{Z}^1)-\text{R}^{40a}$;


wherein when R^{28} and R^{40} are H , then R^{32} is not $=\text{O}$;

each Z^1 is independently O or S ;

R^{28a} , R^{32a} , and R^{40a} are independently $-\text{A}^1-\text{L}^1-\text{A}^2-\text{B}$; $-\text{A}^1-\text{A}^2-\text{B}$; $-\text{L}^2-\text{A}^1-\text{L}^1-\text{A}^2-\text{L}^3-\text{B}$; $-\text{O}-(\text{C}_1-\text{C}_6)\text{alkyl}$; or $-\text{O}-(\text{C}_6-\text{C}_{10})\text{aryl}$; wherein the aryl of $-\text{O}-(\text{C}_6-\text{C}_{10})\text{aryl}$ is unsubstituted or substituted with 1-5 substituents selected from $-\text{NO}_2$ and halogen;

A^1 and A^2 are independently absent or are independently selected from





wherein the bond on the left side of A¹, as drawn, is bound to -C(=Z¹)- or L²; and
 wherein the bond on the right side of the A² moiety, as drawn, is bound to B or L³;

each Q is independently 1 to 3 rings selected from arylene, cycloalkylene, heteroarylene, and heterocyclylene;

each X is independently absent or 1 to 2 rings selected from arylene, cycloalkylene, heteroarylene, and heterocyclylene;

each X¹ independently is a heteroarylene or heterocyclylene ring;


each W is independently absent or 1 to 2 rings selected from arylene, cycloalkylene, heteroarylene, and heterocyclylene;

each W¹ independently is a heteroarylene or heterocyclylene ring;


each G is independently absent or a ring selected from arylene, cycloalkylene, heteroarylene, and heterocyclylene;

each G¹ and G² are independently heteroarylene or heterocyclylene ring;



each L¹ is independently selected from



L^2 and L^3 are independently absent or are independently selected from



each B is independently selected from



each R^3 is independently H or (C_1-C_6) alkyl;

each R^4 is independently H, (C_1-C_6) alkyl, halogen, 5-12 membered heteroaryl, 5-12 membered heterocyclyl, (C_6-C_{10}) aryl, wherein the heteroaryl, heterocyclyl, and aryl are each independently optionally substituted with $-N(R^3)_2$, $-OR^3$, halogen, (C_1-C_6) alkyl,

-(C₁-C₆)alkylene-heteroaryl, -(C₁-C₆)alkylene-CN, -C(O)NR³-heteroaryl, or -C(O)NR³-heterocyclyl;

each R⁵ is independently H, (C₁-C₆)alkyl, -C(O)OR³, or -N(R³)₂, wherein the alkyl of (C₁-C₆)alkyl is each independently optionally substituted with -N(R³)₂ or -OR³;

each R⁶ is independently H, (C₁-C₆)alkyl, -C(O)OR³, or -N(R³)₂, wherein the alkyl of (C₁-C₆)alkyl is each independently optionally substituted with -N(R³)₂ or -OR³;

each R⁷ is independently H, (C₁-C₆)alkyl, -C(O)OR³, or -N(R³)₂, wherein the alkyl of (C₁-C₆)alkyl is each independently optionally substituted with -N(R³)₂ or -OR³;

each R⁸ is independently H, (C₁-C₆)alkyl, -C(O)OR³, or -N(R³)₂, wherein the alkyl of (C₁-C₆)alkyl is each independently optionally substituted with -N(R³)₂ or -OR³;

each Y is independently -C(R³)₂ or a bond;

each n is independently an integer from one to 12;


each o is independently an integer from zero to 30;

each p is independently an integer from zero to 12;

each q is independently an integer from zero to 30; and

each r is independently an integer from one to 6.

[0012] The present disclosure provides a compound of Formula I:



or pharmaceutically acceptable salt or tautomer thereof, wherein:

R³² is -H, =O, or -OR³;

R²⁸ is -H or -C(=Z¹)-R^{28a};


R⁴⁰ is -H or -C(=Z¹)-R^{40a};

wherein at least one of R²⁸ and R⁴⁰ is not H;

Z¹ is independently O or S;

R^{28a} and R^{40a} are independently $-A^1-L^1-A^2-B$; $-A^1-A^2-B$; $-L^2-A^1-L^1-A^2-L^3-B$; $-O-(C_1-C_6)alkyl$; or $-O-(C_6-C_{10})aryl$; wherein the aryl of $-O-(C_6-C_{10})aryl$ is unsubstituted or substituted with 1-5 substituents selected from $-NO_2$ and halogen;

A^1 and A^2 are independently absent or are independently selected from



wherein the bond on the left side of A¹, as drawn, is bound to -C(=Z¹)- or L²; and wherein the bond on the right side of the A² moiety, as drawn, is bound to B or L³;

each Q is independently 1 to 3 rings selected from arylene, cycloalkylene, heteroarylene, and heterocyclylene;

each X is independently absent or 1 to 2 rings selected from arylene, cycloalkylene, heteroarylene, and heterocyclylene;

each X¹ is independently a heteroarylene or heterocyclylene ring;


each W is independently absent or 1 to 2 rings selected from arylene, cycloalkylene, heteroarylene, and heterocyclylene;

each W¹ independently is a heteroarylene or heterocyclylene ring;


each G is independently absent or a ring selected from arylene, cycloalkylene, heteroarylene, and heterocyclylene;

each G¹ and G² are independently heteroarylene or heterocyclylene ring;


each L^1 is independently selected from




L^2 and L^3 are independently absent or are independently selected from



each B is independently selected from





each R³ is independently H or (C₁–C₆)alkyl;

each R⁴ is independently H, (C₁–C₆)alkyl, halogen, 5-12 membered heteroaryl, 5-12 membered heterocyclyl, (C₆–C₁₀)aryl, wherein the heteroaryl, heterocyclyl, and aryl are each independently optionally substituted with $\text{---N}(\text{R}^3)_2$, ---OR^3 , halogen, (C₁–C₆)alkyl, $\text{---}(\text{C}_1\text{---C}_6)\text{alkylene---heteroaryl}$, $\text{---}(\text{C}_1\text{---C}_6)\text{alkylene---CN}$, $\text{---C}(\text{O})\text{NR}^3\text{---heteroaryl}$, or $\text{---C}(\text{O})\text{NR}^3\text{---heterocyclyl}$;

each R⁵ is independently H, (C₁–C₆)alkyl, $\text{---C}(\text{O})\text{OR}^3$, or $\text{---N}(\text{R}^3)_2$, wherein the alkyl of (C₁–C₆)alkyl is optionally substituted with $\text{---N}(\text{R}^3)_2$ or ---OR^3 ;

each R⁶ is independently H, (C₁–C₆)alkyl, $\text{---C}(\text{O})\text{OR}^3$, or $\text{---N}(\text{R}^3)_2$, wherein the alkyl is of (C₁–C₆)alkyl optionally substituted with $\text{---N}(\text{R}^3)_2$ or ---OR^3 ;


each R⁷ is independently H, (C₁–C₆)alkyl, $\text{---C}(\text{O})\text{OR}^3$, or $\text{---N}(\text{R}^3)_2$, wherein the alkyl of (C₁–C₆)alkyl is optionally substituted with $\text{---N}(\text{R}^3)_2$ or ---OR^3 ;

each R⁸ is independently H, (C₁–C₆)alkyl, $\text{---C}(\text{O})\text{OR}^3$, or $\text{---N}(\text{R}^3)_2$, wherein the alkyl of (C₁–C₆)alkyl is optionally substituted with $\text{---N}(\text{R}^3)_2$ or ---OR^3 ;

each Y is independently $\text{---C}(\text{R}^3)_2$ or a bond;

each n is independently an integer from one to 12;
 each o is independently an integer from zero to 30;
 each p is independently an integer from zero to 12;
 each q is independently an integer from zero to 30; and
 each r is independently an integer from one to 6.

[0013] The present disclosure provides a compound of Formula II:



or a pharmaceutically acceptable salt or tautomer thereof, wherein:

R^{32} is $-H$, $=O$ or $-OR^3$;

R^{28} is $-H$ or $-C(=Z^1)-R^{28a}$;


R^{40} is $-H$ or $-C(=Z^1)-R^{40a}$;


wherein at least one of R^{28} and R^{40} is not H ;

Z^1 is independently O or S ;

R^{28a} and R^{40a} are independently $-A^1-L^1-A^2-B$; $-A^1-A^2-B$; $-O-(C_1-C_6)alkyl$; or $-O-(C_6-C_{10})aryl$; wherein the aryl of $-O-(C_6-C_{10})aryl$ is unsubstituted or substituted with 1-5 substituents selected from $-NO_2$ and halogen;

A^1 and A^2 are independently absent or are independently selected from





wherein the bond on the left side of A¹, as drawn, is bound to -C(=Z¹)-; and wherein the bond on the right side of the A² moiety, as drawn, is bound to B;

each Q is independently 1 to 3 rings selected from arylene, cycloalkylene, heteroarylene, and heterocyclylene;

each X is independently absent or 1 to 2 rings selected from arylene, cycloalkylene, heteroarylene, and heterocyclylene;

each X¹ is independently a heteroarylene or heterocyclylene ring;


each W is independently absent or 1 to 2 rings selected from arylene, cycloalkylene, heteroarylene, and heterocyclylene;

each W¹ is independently a heteroarylene or heterocyclylene ring;


each G is independently absent or a ring selected from arylene, cycloalkylene, heteroarylene, and heterocyclylene;


each G¹ and G² are independently heteroarylene or heterocyclylene ring;







each L¹ is independently selected from



each B is independently selected from





each B¹ is independently selected from $\text{---}\text{NR}^3\text{---}(\text{C}(\text{R}^3)_2)_n\text{---}$,
 $\text{---}\text{NR}^3\text{---}(\text{C}(\text{R}^3)_2)_n\text{---}(\text{C}_6\text{---C}_{10})\text{arylene---}(\text{C}(\text{R}^3)_2)_n\text{---}$, $\text{---}\text{NR}^3\text{---}(\text{C}(\text{R}^3)_2)_n\text{---heteroarylene---}$,
 $\text{---}(\text{C}_6\text{---C}_{10})\text{arylene---}$, $\text{---}\text{NR}^3\text{---}(\text{C}(\text{R}^3)_2)_n\text{---NR}^3\text{C(O)---}$,
 $\text{---}\text{NR}^3\text{---}(\text{C}(\text{R}^3)_2)_n\text{---heteroarylene---heterocyclylene---}(\text{C}_6\text{---C}_{10})\text{arylene---}$,
 $\text{---heteroarylene---heterocyclylene---}(\text{C}_6\text{---C}_{10})\text{arylene---}$, $\text{---}\text{C}(=\text{O})\text{---}(\text{C}(\text{R}^3)_2)_p\text{---}$, $\text{---}\text{C}(=\text{O})\text{---}(\text{C}(\text{R}^3)_2)_p\text{---heteroarylene---}$,
,
,
,
,
,
,
,
 and

$^{-3-}\text{NR}^3\text{-(C(R}^3\text{))}_n\text{-S(O)}_2\text{-arylene-C(O)-}$, wherein the $^{-3-}$ bond on the left side of B¹, as drawn, is bound to A² or L¹; and wherein the heteroarylene, heterocyclylene, and arylene are each independently optionally substituted with alkyl, hydroxyalkyl, haloalkyl, alkoxy, halogen, or hydroxyl;

each R³ is independently H or (C₁-C₆)alkyl;

each R⁴ is independently H, (C₁-C₆)alkyl, halogen, 5-12 membered heteroaryl, 5-12 membered heterocyclyl, (C₆-C₁₀)aryl, wherein the heteroaryl, heterocyclyl, and aryl are each independently optionally substituted with -N(R³)₂, -OR³, halogen, (C₁-C₆)alkyl, -(C₁-C₆)alkylene-heteroaryl, -(C₁-C₆)alkylene-CN, -C(O)NR³-heteroaryl, or -C(O)NR³-heterocyclyl;

each R^5 is independently H, (C₁-C₆)alkyl, -C(O)OR³, or -N(R³)₂, wherein the alkyl of (C₁-C₆)alkyl is optionally substituted with -N(R³)₂ or -OR³;

each R^6 is independently H, (C₁-C₆)alkyl, -C(O)OR³, or -N(R³)₂, wherein the alkyl of (C₁-C₆)alkyl is optionally substituted with -N(R³)₂ or -OR³;

each R^7 is independently H, (C₁-C₆)alkyl, -C(O)OR³, or -N(R³)₂, wherein the alkyl of (C₁-C₆)alkyl is optionally substituted with -N(R³)₂ or -OR³;

each R^8 is independently H, (C₁-C₆)alkyl, -C(O)OR³, or -N(R³)₂, wherein the alkyl of (C₁-C₆)alkyl is optionally substituted with -N(R³)₂ or -OR³;

each Y is independently C(R³)₂ or a bond;

each n is independently an integer from one to 12;


each o is independently an integer from zero to 30;

each p is independently an integer from zero to 12;

each q is independently an integer from zero to 30; and


each r is independently an integer from one to 6.

[0014] In some embodiments, a compound of Formula I or II is represented by the structure of Formula I-28:




or a pharmaceutically acceptable salt or tautomer thereof.

[0015] In some embodiments, a compound of Formula Ia, Ic, I, or II is represented by the structure of Formula I-28b:




or a pharmaceutically acceptable salt or a tautomer thereof.

[0016] In some embodiments, a compound of Formula I or II is represented by the structure of Formula I-40:



or a pharmaceutically acceptable salt or tautomer thereof.


[0017] In some embodiments, a compound of Formula Ia, Ic, I or II is represented by the structure of Formula I-40b:



(I-40b)

or a pharmaceutically acceptable salt or a tautomer thereof.

[0018] In some embodiments, a compound of Formula Ia, Ic, I or II is represented by the structure of Formula I-32b:



(I-32b)

or a pharmaceutically acceptable salt or a tautomer thereof.

[0019] The present disclosure provides a method of treating a disease or disorder mediated by mTOR comprising administering to the subject suffering from or susceptible to developing a disease or disorder mediated by mTOR a therapeutically effective amount of one or more disclosed compounds. The present disclosure provides a method of preventing a disease or disorder mediated by mTOR comprising administering to the subject suffering from or susceptible to developing a disease or disorder mediated by mTOR a therapeutically effective amount of one or more disclosed compounds. The present disclosure provides a method of reducing the risk of a disease or disorder mediated by mTOR comprising administering to the subject suffering from or susceptible to developing a disease or disorder mediated by mTOR a therapeutically effective amount of one or more disclosed compounds.

[0020] Another aspect of the present disclosure is directed to a pharmaceutical composition comprising a compound of Formula I, Ia, Ib, Ic, II, or IIb, or a pharmaceutically acceptable salt or tautomer of any of the foregoing, and a pharmaceutically acceptable carrier. The pharmaceutically acceptable carrier can further comprise an excipient, diluent, or surfactant. The pharmaceutical composition can be effective for treating, preventing, or reducing the risk of a disease or disorder mediated by mTOR in a subject in need thereof.

[0021] Another aspect of the present disclosure relates to a compound of Formula I, Ia, Ib, Ic, II, or IIb, or a pharmaceutically acceptable salt or tautomer of any of the foregoing, for use in treating, preventing, or reducing the risk of a disease or disorder mediated by mTOR in a subject in need thereof.

[0022] Another aspect of the present disclosure relates to the use of a compound of Formula I, Ia, Ib, Ic, II, or IIb, or a pharmaceutically acceptable salt or tautomer of any of the foregoing, in the manufacture of a medicament for in treating, preventing, or reducing the risk of a disease or disorder mediated by mTOR in a subject in need thereof.

[0023] The present disclosure also provides compounds that are useful in inhibiting mTOR.

DETAILED DESCRIPTION OF THE DISCLOSURE

[0024] The present disclosure relates to mTOR inhibitors. Specifically, the embodiments are directed to compounds and compositions inhibiting mTOR, methods of treating diseases mediated by mTOR, and methods of synthesizing these compounds.

[0025] The details of the disclosure are set forth in the accompanying description below. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present disclosure, illustrative methods and materials are now described. Other features, objects, and advantages of the disclosure will be apparent from the description and from the claims. In the specification and the appended claims, the singular forms also may include the plural unless the context clearly dictates otherwise. Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this disclosure belongs. All patents and publications cited in this specification are incorporated herein by reference in their entireties.

Terms

[0026] The articles “a” and “an” are used in this disclosure and refers to one or more than one (i.e., to at least one) of the grammatical object of the article, unless indicated otherwise. By way of example, “an element” may mean one element or more than one element, unless indicated otherwise.

[0027] The term “or” means “and/or” unless indicated otherwise. The term “and/or” means either “and” or “or”, or both, unless indicated otherwise.

[0028] The term “optionally substituted” unless otherwise specified means that a group may be unsubstituted or substituted by one or more (e.g., 0, 1, 2, 3, 4, or 5 or more, or any range derivable therein) of the substituents listed for that group in which said substituents may be the same or different. In an embodiment, an optionally substituted group has 1 substituent. In another embodiment an optionally substituted group has 2 substituents. In another embodiment an optionally substituted group has 3 substituents. In another embodiment an optionally substituted group has 4 substituents. In another embodiment an optionally substituted group has 5 substituents.

[0029] The term “alkyl,” by itself or as part of another substituent, means, unless otherwise stated, a straight (i.e., unbranched) or branched non-cyclic carbon chain (or carbon), or combination thereof, which may be fully saturated, mono- or polyunsaturated and can include di- and multivalent radicals, having the number of carbon atoms designated (i.e., C₁-C₁₀ means one to ten carbons). Examples of saturated hydrocarbon radicals include, but are not limited to, groups such as methyl, ethyl, n-propyl, isopropyl, n-butyl, t-butyl, isobutyl, sec-butyl, (cyclohexyl)methyl, homologs and isomers of, for example, n-pentyl, n-hexyl, n-heptyl, n-octyl, and the like. An unsaturated alkyl group is one having one or more double bonds or triple bonds. Examples of unsaturated alkyl groups include, but are not limited to, vinyl, 2-propenyl, crotyl, 2-isopentenyl, 2-(butadienyl), 2,4-pentadienyl, 3-(1,4-pentadienyl), ethynyl, 1- and 3-propynyl, 3-butyne, and the higher homologs and isomers.

[0030] The term “alkylene,” by itself or as part of another substituent, means, unless otherwise stated, a divalent radical derived from an alkyl. Typically, an alkyl (or alkylene) group will have from 1 to 24 carbon atoms, such as those groups having 10 or fewer carbon atoms.

[0031] The term “alkenyl” means an aliphatic hydrocarbon group containing a carbon—carbon double bond and which may be straight or branched having about 2 to about 6 carbon

atoms in the chain. Certain alkenyl groups have 2 to about 4 carbon atoms in the chain. Branched may mean that one or more lower alkyl groups such as methyl, ethyl, or propyl are attached to a linear alkenyl chain. Exemplary alkenyl groups include ethenyl, propenyl, n-butenyl, and i-butenyl. A C₂-C₆ alkenyl group is an alkenyl group containing between 2 and 6 carbon atoms.

[0032] The term “alkenylene,” by itself or as part of another substituent, means, unless otherwise stated, a divalent radical derived from an alkene.

[0033] The term “alkynyl” means an aliphatic hydrocarbon group containing a carbon—carbon triple bond and which may be straight or branched having about 2 to about 6 carbon atoms in the chain. Certain alkynyl groups have 2 to about 4 carbon atoms in the chain. Branched may mean that one or more lower alkyl groups such as methyl, ethyl, or propyl are attached to a linear alkynyl chain. Exemplary alkynyl groups include ethynyl, propynyl, n-butenyl, 2-butenyl, 3-methylbutynyl, and n-pentynyl. A C₂-C₆ alkynyl group is an alkynyl group containing between 2 and 6 carbon atoms.

[0034] The term “alkynylene,” by itself or as part of another substituent, means, unless otherwise stated, a divalent radical derived from an alkyne.

[0035] The term “cycloalkyl” means a monocyclic or polycyclic saturated or partially unsaturated carbon ring containing 3-18 carbon atoms. Examples of cycloalkyl groups include, without limitation, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctanyl, norboranyl, norbornyl, bicyclo[2.2.2]octanyl, or bicyclo[2.2.2]octenyl. A C₃-C₈ cycloalkyl is a cycloalkyl group containing between 3 and 8 carbon atoms. A cycloalkyl group can be fused (e.g., decalin) or bridged (e.g., norbornane).

[0036] A “cycloalkylene,” alone or as part of another substituent, means a divalent radical derived from a cycloalkyl.

[0037] The terms “heterocyclyl” or “heterocycloalkyl” or “heterocycle” refers to a monocyclic or polycyclic 3 to 24-membered ring containing carbon and at least one heteroatom selected from oxygen, phosphorous, nitrogen, and sulfur and wherein there is not delocalized π electrons (aromaticity) shared among the ring carbon or heteroatom(s). Heterocyclyl rings include, but are not limited to, oxetanyl, azetadiny, tetrahydrofuranyl, pyrrolidinyl, oxazolinyl, oxazolidinyl, thiazolinyl, thiazolidinyl, pyranyl, thiopyranyl, tetrahydropyranyl, dioxalinyl, piperidinyl, morpholinyl, thiomorpholinyl, thiomorpholinyl S-oxide, thiomorpholinyl S-dioxide, piperazinyl, azepinyl, oxepinyl, diazepinyl, tropanyl, and

homotropanyl. A heterocyclyl or heterocycloalkyl ring can also be fused or bridged, e.g., can be a bicyclic ring.

[0038] A “heterocyclylene” or “heterocycloalkylene,” alone or as part of another substituent, means a divalent radical derived from a “heterocyclyl” or “heterocycloalkyl” or “heterocycle.”

[0039] The term “aryl” means, unless otherwise stated, a polyunsaturated, aromatic, hydrocarbon substituent, which can be a single ring or multiple rings (preferably from 1 to 3 rings) that are fused together (i.e., a fused ring aryl) or linked covalently. A fused ring aryl may refer to multiple rings fused together wherein at least one of the fused rings is an aryl ring.

[0040] An “arylene,” alone or as part of another substituent, means a divalent radical derived from an aryl.

[0041] The term “heteroaryl” refers to an aryl group (or rings) that contains at least one heteroatom such as N, O, or S, wherein the nitrogen and sulfur atom(s) are optionally oxidized, and the nitrogen atom(s) is optionally quaternized. Thus, the term “heteroaryl” includes fused ring heteroaryl groups (i.e., multiple rings fused together wherein at least one of the fused rings is a heteroaromatic ring). A 5,6-fused ring heteroarylene refers to two rings fused together, wherein one ring has 5 members and the other ring has 6 members, and wherein at least one ring is a heteroaryl ring. Likewise, a 6,6-fused ring heteroarylene refers to two rings fused together, wherein one ring has 6 members and the other ring has 6 members, and wherein at least one ring is a heteroaryl ring. And a 6,5-fused ring heteroarylene refers to two rings fused together, wherein one ring has 6 members and the other ring has 5 members, and wherein at least one ring is a heteroaryl ring. A heteroaryl group can be attached to the remainder of the molecule through a carbon or heteroatom. Non-limiting examples of aryl and heteroaryl groups include phenyl, 1-naphthyl, 2-naphthyl, 4-biphenyl, 1-pyrrolyl, 2-pyrrolyl, 3-pyrrolyl, 3-pyrazolyl, 2-imidazolyl, 4-imidazolyl, pyrazinyl, 2-oxazolyl, 4-oxazolyl, 2-phenyl-4-oxazolyl, 5-oxazolyl, 3-isoxazolyl, 4-isoxazolyl, 5-isoxazolyl, 2-thiazolyl, 4-thiazolyl, 5-thiazolyl, 2-furyl, 3-furyl, 2-thienyl, 3-thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 2-pyrimidyl, 4-pyrimidyl, 5-benzothiazolyl, purinyl, 2-benzimidazolyl, 5-indolyl, 1-isoquinolyl, 5-isoquinolyl, 2-quinoxaliny, 5-quinoxaliny, 3-quinolyl, and 6-quinolyl. Substituents for each of the above noted aryl and heteroaryl ring systems are selected from the group of acceptable substituents described herein.

[0042] The term “heteroaryl” may also include multiple condensed ring systems that have at least one such aromatic ring, which multiple condensed ring systems are further described below. The term may also include multiple condensed ring systems (e.g., ring systems comprising 2, 3 or 4 rings) wherein a heteroaryl group, as defined above, can be condensed with one or more rings selected from heteroaryls (to form for example a naphthyridinyl such as 1,8-naphthyridinyl), heterocycles, (to form for example a 1, 2, 3, 4-tetrahydronaphthyridinyl such as 1, 2, 3, 4-tetrahydro-1,8-naphthyridinyl), carbocycles (to form for example 5,6,7, 8-tetrahydroquinolyl) and aryls (to form for example indazolyl) to form the multiple condensed ring system. The rings of the multiple condensed ring system can be connected to each other via fused, spiro and bridged bonds when allowed by valency requirements. It is to be understood that the individual rings of the multiple condensed ring system may be connected in any order relative to one another. It is also to be understood that the point of attachment of a multiple condensed ring system (as defined above for a heteroaryl) can be at any position of the multiple condensed ring system including a heteroaryl, heterocycle, aryl or carbocycle portion of the multiple condensed ring system and at any suitable atom of the multiple condensed ring system including a carbon atom and heteroatom (e.g., a nitrogen).

[0043] A “heteroarylene,” alone or as part of another substituent, means a divalent radical derived from a heteroaryl.

[0044] Non-limiting examples of aryl and heteroaryl groups include pyridinyl, pyrimidinyl, thiophenyl, thienyl, furanyl, indolyl, benzoxadiazolyl, benzodioxolyl, benzodioxanyl, thianaphthanyl, pyrrolopyridinyl, indazolyl, quinolinyl, quinoxalinyl, pyridopyrazinyl, quinazolinonyl, benzoisoxazolyl, imidazopyridinyl, benzofuranyl, benzothienyl, benzothiophenyl, phenyl, naphthyl, biphenyl, pyrrolyl, pyrazolyl, imidazolyl, pyrazinyl, oxazolyl, isoxazolyl, thiazolyl, furylthienyl, pyridyl, pyrimidyl, benzothiazolyl, purinyl, benzimidazolyl, isoquinolyl, thiadiazolyl, oxadiazolyl, pyrrolyl, diazolyl, triazolyl, tetrazolyl, benzothiadiazolyl, isothiazolyl, pyrazolopyrimidinyl, pyrrolopyrimidinyl, benzotriazolyl, benzoxazolyl, or quinolyl. The examples above may be substituted or unsubstituted and divalent radicals of each heteroaryl example above are non-limiting examples of heteroarylene. A heteroaryl moiety may include one ring heteroatom (e.g., O, N, or S). A heteroaryl moiety may include two optionally different ring heteroatoms (e.g., O, N, or S). A heteroaryl moiety may include three optionally different ring heteroatoms (e.g., O, N, or S). A heteroaryl moiety may include four optionally different ring heteroatoms (e.g., O,

N, or S). A heteroaryl moiety may include five optionally different ring heteroatoms (e.g., O, N, or S). An aryl moiety may have a single ring. An aryl moiety may have two optionally different rings. An aryl moiety may have three optionally different rings. An aryl moiety may have four optionally different rings. A heteroaryl moiety may have one ring. A heteroaryl moiety may have two optionally different rings. A heteroaryl moiety may have three optionally different rings. A heteroaryl moiety may have four optionally different rings. A heteroaryl moiety may have five optionally different rings.

[0045] The terms “halo” or “halogen,” by themselves or as part of another substituent, mean, unless otherwise stated, a fluorine, chlorine, bromine, or iodine atom. Additionally, terms such as “haloalkyl” may include monohaloalkyl and polyhaloalkyl. For example, the term “halo(C₁-C₄)alkyl” may include, but is not limited to, fluoromethyl, difluoromethyl, trifluoromethyl, 2,2,2-trifluoroethyl, 4-chlorobutyl, 3-bromopropyl, 1-fluoro-2-bromoethyl, and the like.

[0046] The term “hydroxyl,” as used herein, means -OH.

[0047] The term “hydroxyalkyl” as used herein, means an alkyl moiety as defined herein, substituted with one or more, such as one, two or three, hydroxy groups. In certain instances, the same carbon atom does not carry more than one hydroxy group. Representative examples include, but are not limited to, hydroxymethyl, 2-hydroxyethyl, 2-hydroxypropyl, 3-hydroxypropyl, 1-(hydroxymethyl)-2-methylpropyl, 2-hydroxybutyl, 3-hydroxybutyl, 4-hydroxybutyl, 2,3-dihydroxypropyl, 2-hydroxy-1-hydroxymethylethyl, 2,3-dihydroxybutyl, 3,4-dihydroxybutyl and 2-(hydroxymethyl)-3-hydroxypropyl.

[0048] The term “oxo,” as used herein, means an oxygen that is double bonded to a carbon atom.

[0049] A substituent group, as used herein, may be a group selected from the following moieties:

(A) oxo, halogen, -CF₃, -CN, -OH, -OCH₃, -NH₂, -COOH, -CONH₂, -NO₂, -SH, -SO₃H, -SO₄H, -SO₂NH₂, -NHNH₂, -ONH₂, -NHC(=O)NHNH₂, -NHC(=O)NH₂, -NHSO₂H, -NHC(=O)H, -NHC(O)-OH, -NHOH, -OCF₃, -OCHF₂, -OCH₂F, unsubstituted alkyl, unsubstituted cycloalkyl, unsubstituted heterocycloalkyl, unsubstituted aryl, unsubstituted heteroaryl, and

(B) alkyl, cycloalkyl, heterocyclyl, aryl, heteroaryl, substituted with at least one substituent selected from:

(i) oxo, halogen, -CF₃, -CN, -OH, -OCH₃, -NH₂, -COOH, -CONH₂, -NO₂, -SH, -SO₃H, -SO₄H, -SO₂NH₂, -NHNH₂, -ONH₂, -NHC=(O)NHNH₂, -NHC=(O)NH₂, -NH₂SO₂H, -NHC=(O)H, -NHC(O)-OH, -NHOH, -OCF₃, -OCHF₂, -OCH₂F, unsubstituted alkyl, unsubstituted heteroalkyl, unsubstituted cycloalkyl, unsubstituted heterocycloalkyl, unsubstituted aryl, unsubstituted heteroaryl, and

(ii) alkyl, heteroalkyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, substituted with at least one substituent selected from:

(a) oxo, halogen, -CF₃, -CN, -OH, -OCH₃, -NH₂, -COOH, -CONH₂, -NO₂, -SH, -SO₃H, -SO₄H, -SO₂NH₂, -NHNH₂, -ONH₂, -NHC=(O)NHNH₂, -NHC=(O)NH₂, -NH₂SO₂H, -NHC=(O)H, -NHC(O)-OH, -NHOH, -OCF₃, -OCHF₂, -OCH₂F, unsubstituted alkyl, unsubstituted heteroalkyl, unsubstituted cycloalkyl, unsubstituted heterocycloalkyl, unsubstituted aryl, unsubstituted heteroaryl, and

(b) alkyl, heteroalkyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, substituted with at least one substituent selected from: oxo, halogen, -CF₃, -CN, -OH, -OCH₃, -NH₂, -COOH, -CONH₂, -NO₂, -SH, -SO₃H, -SO₄H, -SO₂NH₂, -NHNH₂, -ONH₂, -NHC=(O)NHNH₂, -NHC=(O)NH₂, -NH₂SO₂H, -NHC=(O)H, -NHC(O)-OH, -NHOH, -OCF₃, -OCHF₂, -OCH₂F, unsubstituted alkyl, unsubstituted heteroalkyl, unsubstituted cycloalkyl, unsubstituted heterocycloalkyl, unsubstituted aryl, unsubstituted heteroaryl.

[0050] An “effective amount” when used in connection with a compound is an amount effective for treating or preventing a disease in a subject as described herein.

[0051] The term “carrier”, as used in this disclosure, encompasses carriers, excipients, and diluents and may mean a material, composition or vehicle, such as a liquid or solid filler, diluent, excipient, solvent or encapsulating material, involved in carrying or transporting a pharmaceutical agent from one organ, or portion of the body, to another organ, or portion of the body of a subject.

[0052] The term “treating” with regard to a subject, refers to improving at least one symptom of the subject’s disorder. Treating may include curing, improving, or at least partially ameliorating the disorder.

[0053] The term “prevent” or “preventing” with regard to a subject refers to keeping a disease or disorder from afflicting the subject. Preventing may include prophylactic treatment. For instance, preventing can include administering to the subject a compound disclosed herein before a subject is afflicted with a disease and the administration will keep the subject from being afflicted with the disease.


[0054] The term “disorder” is used in this disclosure and means, and is used interchangeably with, the terms disease, condition, or illness, unless otherwise indicated.

[0055] The term “administer”, “administering”, or “administration” as used in this disclosure refers to either directly administering a disclosed compound or pharmaceutically acceptable salt or tautomer of the disclosed compound or a composition to a subject, or administering a prodrug derivative or analog of the compound or pharmaceutically acceptable salt or tautomer of the compound or composition to the subject, which can form an equivalent amount of active compound within the subject’s body.

[0056] A “patient” or “subject” is a mammal, e.g., a human, mouse, rat, guinea pig, dog, cat, horse, cow, pig, or non-human primate, such as a monkey, chimpanzee, baboon or rhesus.


Compounds

[0057] The present disclosure provides a compound having the structure of Formula Ic,




or a pharmaceutically acceptable salt or tautomer thereof, wherein R^{32} , R^{28} , and R^{40} are described as above.

[0058] The present disclosure provides a compound having the structure of Formula Ia,




or a pharmaceutically acceptable salt or tautomer thereof, wherein R³², R²⁸, and R⁴⁰ are described as above.

[0059] The present disclosure provides a compound having the structure of Formula I,



or a pharmaceutically acceptable salt or tautomer thereof, wherein R³², R²⁸, and R⁴⁰ are described as above.


[0060] The present disclosure provides a compound having the structure of Formula Ib:



(Ib)

or a pharmaceutically acceptable salt and or tautomer thereof, wherein R³², R²⁸, and R⁴⁰ are described as above for Formula I.


[0061] The present disclosure provides a compound having the structure of Formula II,



(II)


or a pharmaceutically acceptable salt or tautomer thereof, wherein R³², R²⁸, and R⁴⁰ are described as above.

[0062] The present disclosure provides a compound having the structure of Formula IIb:



or a pharmaceutically acceptable salt or tautomer thereof, wherein R^{32} , R^{28} , and R^{40} are described as above for Formula II.

[0063] In certain embodiments, a compound has the following formula:




or a pharmaceutically acceptable salt or tautomer thereof.

[0064] In certain embodiments, R^{32} is $=O$. In certain embodiments, R^{32} is $-OR^3$. In certain embodiments, R^{32} is H . In certain embodiments, R^{32} is $-N_3$.

[0065] As described above, each R^3 is independently H or (C_1-C_6) alkyl. In certain embodiments, R^3 is H . In certain embodiments, R^3 is (C_1-C_6) alkyl. In certain embodiments, R^3 is methyl.


[0066] In certain embodiments, R^{28} is H . In certain embodiments, R^{28} is (C_1-C_6) alkyl. In certain embodiments, R^{40} is H .

[0067] In certain embodiments, a compound is represented by the structure of Formula I-40b:



or a pharmaceutically acceptable salt or tautomer thereof, wherein R^{32} and R^{40} are described as above for Formula Ia, Ic, I, or II.

[0068] In certain embodiments, a compound is represented by the structure of Formula I-40:



or a pharmaceutically acceptable salt or tautomer thereof, wherein R^{32} and R^{40} are described as above.


[0069] In certain embodiments, R^{40} is $-C(=Z^1)-R^{40a}$. In certain embodiments, Z^1 is O. In certain embodiments, Z^1 is S.

[0070] In certain embodiments, R^{40a} is $-O-(C_1-C_6)\text{alkyl}$ or $-O-(C_6-C_{10})\text{aryl}$; wherein the aryl is unsubstituted or substituted with 1-5 substituents selected from NO_2 and halogen.

[0071] In certain embodiments, R^{40a} is $-A^1-L^1-A^2-B$. In certain embodiments, R^{40a} is $-A^1-A^2-B$. In certain embodiments, R^{40a} is $-L^2-A^1-L^1-A^2-L^3-B$.


[0072] In certain embodiments, R^{40a} is $-A^1-L^1-A^2-B$, wherein A^1 and A^2 are absent. In certain embodiments, R^{40a} is $-A^1-L^1-A^2-B$, wherein A^2 is absent. In certain embodiments, R^{40a} is $-A^1-L^1-A^2-B$, wherein A^1 is absent. In certain embodiments, R^{40a} is $-A^1-L^1-A^2-B$. In certain embodiments, R^{40a} is $-A^1-A^2-B$. In certain embodiments, R^{40a} is $-L^2-A^1-L^1-A^2-L^3-B$, wherein L^2 and A^1 are absent. In certain embodiments, R^{40a} is $-L^2-A^1-L^1-A^2-L^3-B$, wherein L^2 is absent. In certain embodiments, R^{40a} is $-L^2-A^1-L^1-A^2-L^3-B$, wherein L^3 is absent.

[0073] In certain embodiments, a compound is represented by the structure of Formula I-28b:



or a pharmaceutically acceptable salt or tautomer thereof, wherein R^{32} and R^{28} are described as above for Formula Ia, Ic, I, or II.

[0074] In certain embodiments, a compound is represented by the structure of Formula I-28:



or a pharmaceutically acceptable salt or tautomer thereof, wherein R^{32} and R^{28} are described as above.


[0075] In certain embodiments, R^{28} is $-C(=Z^1)-R^{28a}$. In certain embodiments, Z^1 is O. In certain embodiments, Z^1 is S.

[0076] In certain embodiments, R^{28a} is $-O-(C_1-C_6)\text{alkyl}$ or $-O-(C_6-C_{10})\text{aryl}$; wherein the aryl is unsubstituted or substituted with 1-5 substituents selected from NO_2 and halogen.

[0077] In certain embodiments, R^{28a} is $-A^1-L^1-A^2-B$. In certain embodiments, R^{28a} is $-A^1-A^2-B$. In certain embodiments, R^{28a} is $-L^2-A^1-L^1-A^2-L^3-B$.

[0078] In certain embodiments, R^{28a} is $-A^1-L^1-A^2-B$, wherein A^1 and A^2 are absent. In certain embodiments, R^{28a} is $-A^1-L^1-A^2-B$, wherein A^2 is absent. In certain embodiments, R^{28a} is $-A^1-L^1-A^2-B$, wherein A^1 is absent. In certain embodiments, R^{28a} is $-A^1-L^1-A^2-B$. In certain embodiments, R^{28a} is $-A^1-A^2-B$. In certain embodiments, R^{28a} is $-L^2-A^1-L^1-A^2-L^3-B$, wherein L^2 and A^1 are absent. In certain embodiments, R^{28a} is $-L^2-A^1-L^1-A^2-L^3-B$, wherein L^2 is absent. In certain embodiments, R^{28a} is $-L^2-A^1-L^1-A^2-L^3-B$, wherein L^3 is absent.

[0079] In certain embodiments, the compounds are represented by the structure of Formula I-32b:



or a pharmaceutically acceptable salt or tautomer thereof, wherein R^{32} is described as above for Formula Ia, Ic, I, or II.


[0080] In certain embodiments, R^{32} is $-O-C(=Z^1)-R^{32a}$. In certain embodiments, Z^1 is O. In certain embodiments, Z^1 is S.

[0081] In certain embodiments, R^{32a} is -O-(C₁-C₆)alkyl or -O-(C₆-C₁₀)aryl; wherein the aryl is unsubstituted or substituted with 1-5 substituents selected from NO₂ and halogen.


[0082] In certain embodiments, R^{32a} is $-A^1-L^1-A^2-B$. In certain embodiments, R^{32a} is $-A^1-A^2-B$. In certain embodiments, R^{32a} is $-L^2-A^1-L^1-A^2-L^3-B$.

[0083] In certain embodiments, R^{32a} is $-A^1-L^1-A^2-B$, wherein A^1 and A^2 are absent. In certain embodiments, R^{32a} is $-A^1-L^1-A^2-B$, wherein A^2 is absent. In certain embodiments, R^{32a} is $-A^1-L^1-A^2-B$, wherein A^1 is absent. In certain embodiments, R^{32a} is $-A^1-L^1-A^2-B$. In certain embodiments, R^{32a} is $-A^1-A^2-B$. In certain embodiments, R^{32a} is $-L^2-A^1-L^1-A^2-L^3-B$, wherein L^2 and A^1 are absent. In certain embodiments, R^{32a} is $-L^2-A^1-L^1-A^2-L^3-B$, wherein L^2 is absent. In certain embodiments, R^{32a} is $-L^2-A^1-L^1-A^2-L^3-B$, wherein L^3 is absent.


[0084] As described above, each L^1 is independently selected from




[0085] As described above for Formula Ia, each L^1 is independently selected from




[0086] As described above for Formula Ic, each L^1 is independently selected from





[0087] In certain embodiments, L^1 is





[0088] In certain embodiments, L^1 is






[0089] In certain embodiments, L^1 is .


[0090] In certain embodiments, L^1 is . In certain


embodiments, L^1 is . In certain embodiments, L^1 is



. In certain embodiments, L^1 is


. In certain embodiments, L^1 is .

In certain embodiments, L^1 is .


In certain embodiments, L^1 is .

[0091] In certain embodiments, L^1 is . In certain


embodiments, L^1 is . In certain embodiments, L^1 is .


In certain embodiments, L^1 is .

[0092] As described above, L^2 and L^3 are independently absent or are independently selected from




[0093] As described above for Formula Ia and Ic, L^2 and L^3 are independently absent or are independently selected from







[0094] In certain embodiments, L^2 is absent. In certain embodiments, L^2 is




In certain embodiments, L^2 is




certain embodiments, L^2 is





certain embodiments, L^2 is




certain embodiments, L^2 is




[0095] In certain embodiments, L^2 is


embodiments, L^2 is

[0096] In certain embodiments, L^2 is





[0097] In certain embodiments, L^2 is





[0098] In certain embodiments, L^2 is 


[0099] In certain embodiments, L^3 is absent. In certain embodiments, L^3 is





In certain embodiments, L^3 is  . In


certain embodiments, L^3 is  . In certain


embodiments, L^3 is  . In certain

embodiments, L^3 is 


[00100] In certain embodiments, L^3 is  . In certain

embodiments, L^3 is  . In certain embodiments, L^3 is In certain


embodiments, L^2 is  . In certain embodiments, L^2 is




[00101] In certain embodiments, L^3 is




[00102] In certain embodiments, L^3 is



[00103] In certain embodiments, L^3 is



[00104] As described above, A^1 and A^2 are independently absent or are independently selected from



each Q is independently 1 to 3 rings selected from arylene, cycloalkylene, heteroarylene, and heterocyclylene;

each X is independently absent or 1 to 2 rings selected from arylene, cycloalkylene, heteroarylene, and heterocyclylene;

each X¹ is independently a heteroarylene or heterocyclylene ring;


each W is independently absent or 1 to 2 rings selected from arylene, cycloalkylene, heteroarylene, and heterocyclylene;


each W¹ is independently a heteroarylene or heterocyclylene ring;

each G is independently absent or a ring selected from arylene, cycloalkylene, heteroarylene, and heterocyclylene;

each G¹ and G² are independently heteroarylene or heterocyclylene ring.

[00105] As described above for Formula Ia, A¹ and A² are independently absent or are independently selected from





each Q is independently 1 to 3 rings selected from arylene, cycloalkylene, heteroarylene, and heterocyclylene;

each X is independently absent or 1 to 2 rings selected from arylene, cycloalkylene, heteroarylene, and heterocyclylene;

each X¹ is independently a heteroarylene or heterocyclylene ring;


each W is independently absent or 1 to 2 rings selected from arylene, cycloalkylene, heteroarylene, and heterocyclylene;


each W¹ is independently a heteroarylene or heterocyclylene ring;

each G is independently absent or a ring selected from arylene, cycloalkylene, heteroarylene, and heterocyclylene;

each G¹ and G² are independently heteroarylene or heterocyclylene ring.

[00106] As described above for Formula Ic, A¹ and A² are independently absent or are independently selected from





each Q is independently 1 to 3 rings selected from arylene, cycloalkylene, heteroarylene, and heterocyclylene;

each X is independently absent or 1 to 2 rings selected from arylene, cycloalkylene, heteroarylene, and heterocyclylene;

each X¹ is independently a heteroarylene or heterocyclylene ring;

each W is independently absent or 1 to 2 rings selected from arylene, cycloalkylene, heteroarylene, and heterocyclylene;

each W¹ is independently a heteroarylene or heterocyclylene ring;


each G is independently absent or a ring selected from arylene, cycloalkylene, heteroarylene, and heterocyclylene;

each G¹ and G² are independently heteroarylene or heterocyclylene ring;


[00107] For Formula I, the bond on the left side of A¹, as drawn, is bound to -C(=Z¹)- or L²; and the bond on the right side of the A² moiety, as drawn, is bound to B or L³. For Formula II, the bond on the left side of A¹, as drawn, is bound to -C(=Z¹)-; and the bond on the right side of the A² moiety, as drawn, is bound to B. For Formula Ia and Ic, the bond on

the left side of A¹, as drawn, is bound to -C(=Z¹)- or L²; and wherein the bond on the right side of the A² moiety, as drawn, is bound to B or L³.


[00108] In certain embodiments, A¹ is absent. In certain embodiments, A¹ is



. In certain embodiments, A¹ is




. In certain



embodiments, A¹ is

. In certain embodiments, A¹ is



[00109] In certain embodiments, A¹ is $\text{N}(\text{R}^3)\text{C}(\text{R}^3)_2\text{Q}\text{C}(\text{R}^3)_2\text{C}(\text{O})$, wherein each Q is independently 1 to 3 rings selected from arylene, cycloalkylene, heteroarylene, and heterocyclylene.

[00110] In certain embodiments, A¹ is $\text{N}(\text{R}^3)\text{C}(\text{R}^3)_2\text{Q}\text{C}(\text{R}^3)_2\text{N}(\text{R}^3)$, wherein each Q is independently 1 to 3 rings selected from arylene, cycloalkylene, heteroarylene, and heterocyclylene.


[00111] In certain embodiments, A¹ is $\text{N}(\text{R}^3)\text{C}(\text{R}^3)_2\text{X}\text{X}^1\text{N}$, wherein each X is independently absent or 1 to 2 rings selected from arylene, cycloalkylene, heteroarylene, and heterocyclylene; and each X¹ is a heteroarylene or heterocyclylene ring.


[00112] In certain embodiments, A¹ is $\text{N}\text{W}^1\text{W}\text{C}(\text{R}^3)_2\text{C}(\text{O})$, wherein each W is independently absent or 1 to 2 rings selected from arylene, cycloalkylene, heteroarylene, and heterocyclylene; and each W¹ is a heteroarylene or heterocyclylene ring.



[00113] In certain embodiments, A^1 is $\text{---}\text{N}(\text{W}^1)\text{---W---}\left[\text{C}(\text{R}^3)_2\right]_p\text{---N}(\text{R}^3)\text{---}$, wherein each W is independently absent or 1 to 2 rings selected from arylene, cycloalkylene, heteroarylene, and heterocyclylene; and each W^1 is a heteroarylene or heterocyclylene ring.


[00114] In certain embodiments, A^1 is $\text{---}\text{N}(\text{G}^1)\text{---}$, wherein each G is independently absent or a ring selected from arylene, cycloalkylene, heteroarylene, and heterocyclylene.

[00115] In certain embodiments, A^1 is $\text{---}\text{N}(\text{G}^1)\text{---N---G---N}(\text{G}^2)\text{---}$, wherein each G is independently absent or a ring selected from arylene, cycloalkylene, heteroarylene, and heterocyclylene; and each G^1 and G^2 are independently heteroarylene or heterocyclylene ring.






[00117] In certain embodiments, A¹ is . In certain embodiments, A¹ is


. In certain embodiments, A¹ is . In certain

embodiments, A¹ is .

[00118] In certain embodiments, A² is absent. In certain embodiments, A² is

. In certain embodiments, A² is . In certain

embodiments, A² is . In certain embodiments, A² is

.

[00119] In certain embodiments, A² is $\text{N}(\text{R}^3)\text{C}(\text{R}^3)_2\text{Q}\text{C}(\text{R}^3)_2\text{C}(\text{O})$, wherein each Q is independently 1 to 3 rings selected from arylene, cycloalkylene, heteroarylene, and heterocyclylene.

[00120] In certain embodiments, A² is $\text{N}(\text{R}^3)\text{C}(\text{R}^3)_2\text{Q}\text{C}(\text{R}^3)_2\text{N}(\text{R}^3)$, wherein each Q is independently 1 to 3 rings selected from arylene, cycloalkylene, heteroarylene, and heterocyclylene.

[00121] In certain embodiments, A² is , wherein each X is independently absent or 1 to 2 rings selected from arylene, cycloalkylene, heteroarylene, and heterocyclylene; and each X¹ is independently a heteroarylene or heterocyclylene ring.


[00122] In certain embodiments, A² is , wherein each W is independently absent or 1 to 2 rings selected from arylene, cycloalkylene, heteroarylene, and heterocyclylene; and each W¹ is independently a heteroarylene or heterocyclylene ring.


[00123] In certain embodiments, A² is , wherein each W is independently absent or 1 to 2 rings selected from arylene, cycloalkylene, heteroarylene, and heterocyclylene; and each W¹ is independently a heteroarylene or heterocyclylene ring.


[00124] In certain embodiments, A² is , wherein each G is independently absent or a ring selected from arylene, cycloalkylene, heteroarylene, and heterocyclylene.


[00125] In certain embodiments, A² is , wherein each G is independently absent or a ring selected from arylene, cycloalkylene, heteroarylene, and heterocyclylene; and each G¹ and G² are independently a heteroarylene or heterocyclylene ring.

[00126] In certain embodiments, A² is





[00127] In certain embodiments, A² is . In certain embodiments, A² is




embodiments, A² is .


[00128] As described above, each B is independently selected from




[00129] In certain embodiments, B is .



[00130] In certain embodiments, B is




[00131] In certain embodiments, B is




In certain embodiments, B is

[00132] As described above, each B¹ is independently selected from $^{-3}-NR^3-(C(R^3)_2)_n$, $^{-3}-NR^3-(C(R^3)_2)_n-(C_6-C_{10})\text{arylene}-(C(R^3)_2)_n$, $^{-3}-NR^3-(C(R^3)_2)_n\text{-heteroarylene-}$, $^{-3}-(C_6-C_{10})\text{arylene-}$, $^{-3}-NR^3-(C(R^3)_2)_n-NR^3C(O)-$, $^{-3}-NR^3-(C(R^3)_2)_n\text{-heteroarylene-}$, $\text{heterocyclylene}-(C_6-C_{10})\text{arylene-}$, $^{-3}\text{-heteroarylene-heterocyclylene}-(C_6-C_{10})\text{arylene-}$,




and $^{-3}-NR^3-(C(R^3)_2)_n-S(O)_2\text{-arylene-C(O)-}$, wherein the $^{-3}$ bond on the left side of B¹, as drawn, is bound to A² or L¹; and wherein the heteroarylene, heterocyclylene, and arylene are each independently optionally substituted with alkyl, hydroxyalkyl, haloalkyl, alkoxy, halogen, or hydroxyl.


[00133] As described above for Formula Ic, each B¹ is independently selected from $^{-\text{S}}-\text{NR}^3-(\text{C}(\text{R}^3)_2)_n-$, $^{-\text{S}}-\text{NR}^3-(\text{C}(\text{R}^3)_2)_n-(\text{C}_6-\text{C}_{10})\text{arylene}-(\text{C}(\text{R}^3)_2)_n-$, $^{-\text{S}}-\text{NR}^3-(\text{C}(\text{R}^3)_2)_n-$ heteroarylene-, $^{-\text{S}}-\text{NR}^3-(\text{C}(\text{R}^3)_2)_n$ -heteroarylene-(C(R³)₂)_n-, $^{-\text{S}}-(\text{C}_6-\text{C}_{10})\text{arylene}-$, $^{-\text{S}}-\text{NR}^3-(\text{C}(\text{R}^3)_2)_n-\text{NR}^3\text{C}(\text{O})-$, $^{-\text{S}}-\text{NR}^3-(\text{C}(\text{R}^3)_2)_n$ -heteroarylene-heterocyclylene-(C₆-



S(O)₂-arylene-C(O)-, and $^{-\text{S}}-\text{NR}^3-(\text{C}(\text{R}^3)_2)_n-\text{S}(\text{O})_2\text{-arylene}-(\text{C}(\text{R}^3)_2)_n-$; wherein the $^{-\text{S}}-$ bond on the left side of B¹, as drawn, is bound to A², L³, or L¹; and wherein the heteroarylene, heterocyclylene, and arylene are each independently optionally substituted with alkyl, hydroxyalkyl, haloalkyl, alkoxy, halogen, or hydroxyl.


[00134] In certain embodiments, B¹ is $^{-\text{S}}-\text{NR}^3-(\text{C}(\text{R}^3)_2)_n-$.


[00135] In certain embodiments, B¹ is . In certain embodiments,


B¹ is , wherein arylene is optionally substituted with haloalkyl.


[00136] In certain embodiments, B¹ is $^{-\text{S}}-\text{NR}^3-(\text{C}(\text{R}^3)_2)_n-$, $^{-\text{S}}-\text{NR}^3-(\text{C}(\text{R}^3)_2)_n-(\text{C}_6-\text{C}_{10})\text{arylene}-(\text{C}(\text{R}^3)_2)_n-$, $^{-\text{S}}-\text{NR}^3-(\text{C}(\text{R}^3)_2)_n$ -heteroarylene-, $^{-\text{S}}-(\text{C}_6-\text{C}_{10})\text{arylene}-$, $^{-\text{S}}-\text{NR}^3-(\text{C}(\text{R}^3)_2)_n-\text{NR}^3\text{C}(\text{O})-$, $^{-\text{S}}-\text{NR}^3-(\text{C}(\text{R}^3)_2)_n$ -heteroarylene-heterocyclylene-(C₆-C₁₀)arylene-, or

$\text{---}\overset{\text{Z}}{\text{---}}$ heteroarylene-heterocyclylene-(C₆-C₁₀)arylene-. In certain embodiments, B¹ is




[00137] In certain embodiments, B¹ is . In certain


embodiments, B¹ is . In certain embodiments, B¹ is



[00138] In certain embodiments, B¹ is $\text{---}\overset{\text{Z}}{\text{---}}\text{NR}^3\text{---}(\text{C}(\text{R}^3)_2)_n\text{---heteroarylene---}(\text{C}(\text{R}^3)_2)_n\text{---}$. In



certain embodiments, B¹ is . In certain embodiments, B¹ is



[00139] In certain embodiments, in B¹, the heteroaryl, heterocyclyl, and arylene are optionally substituted with alkyl, hydroxyalkyl, haloalkyl, alkoxy, halogen, or hydroxyl.

[00140] In certain embodiments, R³ is H. In certain embodiments, R³ is (C₁-C₆)alkyl.

[00141] In certain embodiments, R⁴ is H. In certain embodiments, R⁴ is (C₁-C₆)alkyl. In certain embodiments, R⁴ is halogen. In certain embodiments, R⁴ is 5-12 membered heteroaryl, 5-12 membered heterocyclyl, or (C₆-C₁₀)aryl, wherein the heteroaryl, heterocyclyl, and aryl are optionally substituted with $\text{---N(R}^3)_2$, ---OR^3 , halogen, (C₁-C₆)alkyl, $\text{---(C}_1\text{---C}_6\text{)alkylene---heteroaryl}$, $\text{---(C}_1\text{---C}_6\text{)alkylene---CN}$, or $\text{---C(O)NR}^3\text{---heteroaryl}$. In certain embodiments, R⁴ is $\text{---C(O)NR}^3\text{---heterocyclyl}$. In certain embodiments, R⁴ is 5-12 membered heteroaryl, optionally substituted with $\text{---N(R}^3)_2$ or ---OR^3 .

[00142] As described above, each R⁵ is independently H, (C₁-C₆)alkyl, ---C(O)OR^3 , or $\text{---N(R}^3)_2$, wherein the alkyl of (C₁-C₆)alkyl is optionally substituted with $\text{---N(R}^3)_2$ or ---OR^3 . In certain embodiments, R⁵ is H. In certain embodiments, R⁵ is (C₁-C₆)alkyl, wherein the alkyl

is optionally substituted with $-N(R^3)_2$ or $-OR^3$. In certain embodiments, R^5 is $-C(O)OR^3$. In certain embodiments, R^5 is $-N(R^3)_2$.

[00143] As described above, each R^6 is independently H, (C_1-C_6) alkyl, $-C(O)OR^3$, or $-N(R^3)_2$, wherein the alkyl of (C_1-C_6) alkyl is optionally substituted with $-N(R^3)_2$ or $-OR^3$. In certain embodiments, R^6 is H. In certain embodiments, R^6 is (C_1-C_6) alkyl, wherein the alkyl is optionally substituted with $-N(R^3)_2$ or $-OR^3$. In certain embodiments, R^6 is $-C(O)OR^3$. In certain embodiments, R^6 is $-N(R^3)_2$.

[00144] As described above, each R^7 is independently H, (C_1-C_6) alkyl, $-C(O)OR^3$, or $-N(R^3)_2$, wherein the alkyl of (C_1-C_6) alkyl is optionally substituted with $-N(R^3)_2$ or $-OR^3$. In certain embodiments, R^7 is H. In certain embodiments, R^7 is (C_1-C_6) alkyl, wherein the alkyl is optionally substituted with $-N(R^3)_2$ or $-OR^3$. In certain embodiments, R^7 is $-C(O)OR^3$. In certain embodiments, R^7 is $-N(R^3)_2$.

[00145] As described above, each R^8 is independently H, (C_1-C_6) alkyl, $-C(O)OR^3$, or $-N(R^3)_2$, wherein the alkyl of (C_1-C_6) alkyl is optionally substituted with $-N(R^3)_2$ or $-OR^3$. In certain embodiments, R^8 is H. In certain embodiments, R^8 is (C_1-C_6) alkyl, wherein the alkyl is optionally substituted with $-N(R^3)_2$ or $-OR^3$. In certain embodiments, R^8 is $-C(O)OR^3$. In certain embodiments, R^8 is $-N(R^3)_2$.

[00146] As described above, each Y is independently $C(R^3)_2$ or a bond. In certain embodiments, Y is $C(R^3)_2$. In certain embodiments, Y is CH_2 . In certain embodiments, Y is a bond.

[00147] In certain embodiments, n is 1, 2, 3, 4, 5, 6, 7, or 8, or any range derivable therein. In certain embodiments, n is 1, 2, 3, or 4. In certain embodiments, n is 5, 6, 7, or 8. In certain embodiments, n is 9, 10, 11, or 12.

[00148] In certain embodiments, o is an integer from zero to 10, or any range derivable therein. In certain embodiments, o is 0, 1, 2, 3, 4, or 5. In certain embodiments, o is 6, 7, 8, 9, or 10. In certain embodiments, o is one to 7. In certain embodiments, o is one to 8. In certain embodiments, o is one to 9. In certain embodiments, o is 3 to 8.

[00149] In certain embodiments, o is an integer from zero to 30, or any range derivable therein. In certain embodiments, o is an integer from zero to 30, 29, 28, 27, or 26. In certain embodiments, o is an integer from zero to 25, 24, 23, 22, or 21. In certain embodiments, o is an integer from zero to 20, 19, 18, 17, or 16. In certain embodiments, o is an integer from zero to 15, 14, 13, 12, or 11.


[00150] In certain embodiments, p is 0, 1, 2, 3, 4, 5, or 6, or any range derivable therein. In certain embodiments, p is 7, 8, 9, 10, 11, or 12. In certain embodiments, p is 0, 1, 2, or 3. In certain embodiments, p is 4, 5, or 6.

[00151] In certain embodiments, q is an integer from zero to 10, or any range derivable therein. In certain embodiments, q is 0, 1, 2, 3, 4, or 5. In certain embodiments, q is 6, 7, 8, 9, or 10. In certain embodiments, q is one to 7. In certain embodiments, q is one to 8. In certain embodiments, q is one to 9. In certain embodiments, q is 3 to 8.

[00152] In certain embodiments, q is an integer from zero to 30, or any range derivable therein. In certain embodiments, q is an integer from zero to 30, 29, 28, 27, or 26. In certain embodiments, q is an integer from zero to 25, 24, 23, 22, or 21. In certain embodiments, q is an integer from zero to 20, 19, 18, 17, or 16. In certain embodiments, q is an integer from zero to 15, 14, 13, 12, or 11.

[00153] As described above, r is an integer from one to 6. In certain embodiments, r is one. In certain embodiments, r is 2. In certain embodiments, r is 3. In certain embodiments, r is 4. In certain embodiments, r is 5. In certain embodiments, r is 6.

[00154] As described above, when R^{28} and R^{40} are H, then R^{32} is not =O. In certain embodiments, the compound is not rapamycin, as shown below:



In certain embodiments, in Formula Ia or Ic, R^{32} is $-O-C(=Z^1)-R^{32a}$. In certain embodiments, R^{32} is $-O-C(=Z^1)-R^{32a}$; wherein R^{32a} is $-A^1-L^1-A^2-B$; $-A^1-A^2-B$; or $-L^2-A^1-L^1-A^2-L^3-B$. In certain embodiments, in Formula Ia or Ic, R^{28} is $-C(=Z^1)-R^{28a}$. In certain embodiments, R^{28} is $-C(=Z^1)-R^{28a}$; wherein R^{28a} is $-A^1-L^1-A^2-B$; $-A^1-A^2-B$; or $-L^2-A^1-L^1-A^2-L^3-B$. In certain embodiments, in Formula Ia or Ic, R^{40} is $-C(=Z^1)-R^{40a}$. In certain embodiments, R^{40} is $-C(=Z^1)-R^{40a}$; wherein R^{40a} is $-A^1-L^1-A^2-B$; $-A^1-A^2-B$; or $-L^2-A^1-L^1-A^2-L^3-B$.

[00155] The present disclosure provides a compound of Formula Ia or Ic, or a pharmaceutically acceptable salt or tautomer thereof, having one, two, or three of the following features:

- a) R^{32} is $-O-C(=Z^1)-R^{32a}$;
- b) R^{28} is $-C(=Z^1)-R^{28a}$;
- c) R^{40} is $-C(=Z^1)-R^{40a}$.

[00156] The present disclosure provides a compound of Formula Ia or Ic, or a pharmaceutically acceptable salt or tautomer thereof, having one, two, or three of the following features:



- a) R^{32} is $-O-C(=Z^1)-R^{32a}$, wherein R^{32a} is $-A^1-L^1-A^2-B$; $-A^1-A^2-B$; or $-L^2-A^1-L^1-A^2-L^3-B$;
- b) R^{28} is $-C(=Z^1)-R^{28a}$, wherein R^{28a} is $-A^1-L^1-A^2-B$; $-A^1-A^2-B$; or $-L^2-A^1-L^1-A^2-L^3-B$;
- c) R^{40} is $-C(=Z^1)-R^{40a}$, wherein R^{40a} is $-A^1-L^1-A^2-B$; $-A^1-A^2-B$; or $-L^2-A^1-L^1-A^2-L^3-B$.


[00157] The present disclosure provides a compound of Formula Ia or Ic, or a pharmaceutically acceptable salt or tautomer thereof, having one, two, or three of the following features:

- a) R^{40} is $-C(=Z^1)-R^{40a}$;
- b) R^{40a} is $-A^1-L^1-A^2-B$; $-A^1-A^2-B$; $-L^2-A^1-L^1-A^2-L^3-B$;
- c) R^{32} is $-OR^3$, such as $-OH$.


[00158] The present disclosure provides a compound of Formula Ia or Ic, or a pharmaceutically acceptable salt or tautomer thereof, having one, two, three, or four of the following features:

- a) one of R^{28a} , R^{32a} , and R^{40a} is $-A^1-L^1-A^2-B$;
- b) A^1 is absent;
- c) A^2 is absent;






- f) B^1 is $-\text{NR}^3-(\text{C}(\text{R}^3)_2)_n-$ or ;
 g) R^4 is 5-12 membered heteroaryl, optionally substituted with $-\text{N}(\text{R}^3)_2$ or $-\text{OR}^3$.

[00159] The present disclosure provides a compound of formula:



or a pharmaceutically acceptable salt or tautomer thereof, having one, two, three, or four of the following features:


- a) Z^1 is O;
 b) A^1 is absent;



- f) R^4 is 5-12 membered heteroaryl, optionally substituted with $-\text{N}(\text{R}^3)_2$ or $-\text{OR}^3$; and
 g) R^{32} is =O.

In the above, R^{40a} can be $-\text{A}^1-\text{L}^1-\text{A}^2-\text{B}$; $-\text{A}^1-\text{A}^2-\text{B}$; or $-\text{L}^2-\text{A}^1-\text{L}^1-\text{A}^2-\text{L}^3-\text{B}$.

[00160] The present disclosure provides a compound of formula:



or a pharmaceutically acceptable salt or tautomer thereof, having one, two, three, or four of the following features:

a) Z^1 is O;

b) A^1 is absent;

c) L^1 is ;

d) B is


e) B^1 is $-\text{S}-\text{NR}^3-(\text{C}(\text{R}^3)_2)_n-$ or ;

f) R^4 is 5-12 membered heteroaryl, optionally substituted with $-\text{N}(\text{R}^3)_2$ or $-\text{OR}^3$; and

g) R^{32} is $-\text{OH}$.





In the above, R^{40a} can be $-\text{A}^1-\text{L}^1-\text{A}^2-\text{B}$; $-\text{A}^1-\text{A}^2-\text{B}$; or $-\text{L}^2-\text{A}^1-\text{L}^1-\text{A}^2-\text{L}^3-\text{B}$.

[00161] The present disclosure provides a compound of formula:



or a pharmaceutically acceptable salt or tautomer thereof, having one, two, three, or four of the following features:

a) Z^1 is O;




f) R^4 is 5-12 membered heteroaryl, optionally substituted with $-N(R^3)_2$ or $-OR^3$; and

g) R^{32} is =O.

In the above, R^{40a} can be $-A^1-L^1-A^2-B$; $-A^1-A^2-B$; or $-L^2-A^1-L^1-A^2-L^3-B$.


[00162] The present disclosure provides a compound of formula:





or a pharmaceutically acceptable salt or tautomer thereof, having one, two, three, or four of the following features:

a) Z^1 is O;

b) A^1 is absent;






f) R⁴ is 5-12 membered heteroaryl, optionally substituted with $-\text{N}(\text{R}^3)_2$ or $-\text{OR}^3$; and

g) R³² is -OH.




In the above, R⁴⁰ᵃ can be $-\text{A}^1-\text{L}^1-\text{A}^2-\text{B}$; $-\text{A}^1-\text{A}^2-\text{B}$; or $-\text{L}^2-\text{A}^1-\text{L}^1-\text{A}^2-\text{L}^3-\text{B}$.



[00163] The present disclosure provides a compound of formula:



or a pharmaceutically acceptable salt or tautomer thereof, having one, two, three, or four of the following features:

a) Z¹ is O;





$-\text{NR}^3-(\text{C}(\text{R}^3)_2)_n-\text{S}(\text{O})_2\text{-arylene-C}(\text{O})-$, wherein the arylene is optionally substituted with alkyl, hydroxyalkyl, haloalkyl, alkoxy, halogen, or hydroxyl;


g) R⁴ is 5-12 membered heteroaryl, optionally substituted with $-\text{N}(\text{R}^3)_2$ or $-\text{OR}^3$; and


h) R³² is -OH.

In the above, R⁴⁰ᵃ can be $-\text{A}^1-\text{L}^1-\text{A}^2-\text{B}$; $-\text{A}^1-\text{A}^2-\text{B}$; or $-\text{L}^2-\text{A}^1-\text{L}^1-\text{A}^2-\text{L}^3-\text{B}$.


[00164] In certain embodiments, in Formula Ia or Ic, R⁴⁰ᵃ is any organic moiety, which may have a molecular weight (e.g. the sum of the atomic masses of the atoms of the substituent) of less than 15 g/mol, 50 g/mol, 100 g/mol, 150 g/mol, 200 g/mol, 250 g/mol, 300 g/mol, 350 g/mol, 400 g/mol, 450 g/mol, or 500 g/mol.

[00165] In certain embodiments, the present disclosure provides for a compound selected from below or a pharmaceutically acceptable salt or tautomer thereof,







Example 3




Example 4




Example 5




Example 6




Example 7




Example 8




Example 9




Example 10




Example 11




Example 12




Example 13




Example 14




Example 15




Example 16




Example 17




Example 18




Example 19




Example 20




Example 21




Example 22




Example 23




Example 24




Example 25




Example 26




Example 27




Example 28




Example 29




Example 30




Example 31




Example 32




Example 33




Example 34




Example 35




Example 36




Example 37




Example 38




Example 39




Example 40




Example 41




Example 42




Example 43




Example 44




Example 45




Example 46




Example 47




Example 48




Example 49




Example 50




Example 51




Example 52




Example 53




Example 54




Example 55




Example 56




Example 57




Example 58




Example 59




Example 60




Example 61




Example 62




Example 63




Example 64




Example 65




Example 66




Example 67




Example 68




Example 69




Example 70




[00166] In certain embodiments, the present disclosure provides for a compound selected from below or a pharmaceutically acceptable salt or tautomer thereof,




Example 71




Example 72




Example 73




Example 74




Example 75




Example 76




Example 77




Example 78




Example 79




Example 80




Example 81




Example 82




Example 83




Example 84




Example 85




Example 86




Example 87




Example 88




Example 89




Example 90




Example 91




Example 92




Example 93




Example 94




Example 95




Example 96




Example 97




Example 98




Example 99




Example 100




Example 101




Example 102




Example 103




Example 104




Example 105




Example 106




Example 107




Example 108




Example 109




Example 110




Example 111




Example 112




Example 113




Example 114




Example 115




Example 116




Example 117




Example 118




Example 119




Example 120




Example 121




Example 122




Example 123




Example 124




Example 125




Example 126




Example 127




Example 128




Example 129




Example 130




Example 131




Example 132




Example 133




Example 134




Example 135




[00167] In certain embodiments, the present disclosure provides for a compound selected from below or a pharmaceutically acceptable salt or tautomer thereof,




Example 136




Example 137




Example 138




Example 139




Example 140




Example 141




Example 142




Example 143




Example 144




Example 145




Example 146




Example 147




Example 148




Example 149



Example 150



Example 151



[00168] The compounds of the disclosure may include pharmaceutically acceptable salts of the compounds disclosed herein. Representative “pharmaceutically acceptable salts” may include, e.g., water-soluble and water-insoluble salts, such as the acetate, amsonate (4,4-diaminostilbene-2,2-disulfonate), benzenesulfonate, benzonate, bicarbonate, bisulfate, bitartrate, borate, bromide, butyrate, calcium, calcium edetate, camsylate, carbonate, chloride, citrate, clavulariate, dihydrochloride, edetate, edisylate, estolate, esylate, fiunarate, gluceptate, gluconate, glutamate, glycolylarsanilate, hexafluorophosphate, hexylresorcinate, hydrabamine, hydrobromide, hydrochloride, hydroxynaphthoate, iodide, sethionate, lactate, lactobionate, laurate, magnesium, malate, maleate, mandelate, mesylate, methylbromide, methylnitrate, methylsulfate, mucate, napsylate, nitrate, N-methylglucamine ammonium salt, 3-hydroxy-2-naphthoate, oleate, oxalate, palmitate, pamoate, 1,1-methene-bis-2-hydroxy-3-naphthoate, einbonate, pantothenate, phosphate/diphosphate, picrate, polygalacturonate, propionate, p-toluenesulfonate, salicylate, stearate, subacetate, succinate, sulfate, sulfosalicylate, suramate, tannate, tartrate, teoclate, tosylate, triethiodide, and valerate salts.

[00169] “Pharmaceutically acceptable salt” may also include both acid and base addition salts. “Pharmaceutically acceptable acid addition salt” may refer to those salts which retain the biological effectiveness and properties of the free bases, which are not biologically or otherwise undesirable, and which may be formed with inorganic acids such as, but are not limited to, hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid and the like, and organic acids such as, but not limited to, acetic acid, 2,2-dichloroacetic acid, adipic acid, alginic acid, ascorbic acid, aspartic acid, benzenesulfonic acid, benzoic acid, 4-acetamidobenzoic acid, camphoric acid, camphor-10-sulfonic acid, capric acid, caproic acid, caprylic acid, carbonic acid, cinnamic acid, citric acid, cyclamic acid, dodecylsulfuric acid, ethane-1,2-disulfonic acid, ethanesulfonic acid, 2-hydroxyethanesulfonic acid, formic acid, fumaric acid, galactaric acid, gentisic acid, glucoheptonic acid, gluconic acid, glucuronic acid, glutamic acid, glutaric acid, 2-oxo-glutaric acid, glycerophosphoric acid, glycolic acid, hippuric acid, isobutyric acid, lactic acid, lactobionic acid, lauric acid, maleic acid, malic

acid, malonic acid, mandelic acid, methanesulfonic acid, mucic acid, naphthalene-1,5-disulfonic acid, naphthalene-2-sulfonic acid, 1-hydroxy-2-naphthoic acid, nicotinic acid, oleic acid, orotic acid, oxalic acid, palmitic acid, pamoic acid, propionic acid, pyroglutamic acid, pyruvic acid, salicylic acid, 4-aminosalicylic acid, sebacic acid, stearic acid, succinic acid, tartaric acid, thiocyanic acid, p-toluenesulfonic acid, trifluoroacetic acid, undecylenic acid, and the like.

[00170] “Pharmaceutically acceptable base addition salt” may refer to those salts that retain the biological effectiveness and properties of the free acids, which are not biologically or otherwise undesirable. These salts may be prepared from addition of an inorganic base or an organic base to the free acid. Salts derived from inorganic bases may include, but are not limited to, the sodium, potassium, lithium, ammonium, calcium, magnesium, iron, zinc, copper, manganese, aluminum salts and the like. For example, inorganic salts may include, but are not limited to, ammonium, sodium, potassium, calcium, and magnesium salts. Salts derived from organic bases may include, but are not limited to, salts of primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines and basic ion exchange resins, such as ammonia, isopropylamine, trimethylamine, diethylamine, triethylamine, tripropylamine, diethanolamine, ethanolamine, deanol, 2-dimethylaminoethanol, 2-diethylaminoethanol, dicyclohexylamine, lysine, arginine, histidine, caffeine, procaine, hydrabamine, choline, betaine, benethamine, benzathine, ethylenediamine, glucosamine, methylglucamine, theobromine, triethanolamine, tromethamine, purines, piperazine, piperidine, N-ethylpiperidine, polyamine resins and the like.

[00171] Unless otherwise stated, structures depicted herein may also include compounds which differ only in the presence of one or more isotopically enriched atoms. For example, compounds having the present structure except for the replacement of a hydrogen atom by deuterium or tritium, or the replacement of a carbon atom by ^{13}C or ^{14}C , or the replacement of a nitrogen atom by ^{15}N , or the replacement of an oxygen atom with ^{17}O or ^{18}O are within the scope of the disclosure. Such isotopically labeled compounds are useful as research or diagnostic tools.

[00172] In some embodiments, one or more deuterium atoms may be introduced into the PEG moiety of any compound of the present invention. Mechanisms for such modifications are known in the art starting from commercially available starting materials, such as isotopically enriched hydroxylamine building blocks. In some embodiments, a tritium or a

deuterium may be introduced at the C32 position of compounds of the present invention using, for example, a commercially available isotopically pure reducing agent and methods known to those in the art. In some embodiments, ^{14}C may be introduced into the C40 carbamate moiety of compounds of the present invention using commercially available materials and methods known to those of skill in the art. In some embodiments, an isotope such as deuterium or tritium may be introduced into the R^{40a} substituent of a compound of Formula Ia, Ic, I or II, using commercially available starting materials and methods known to those of skill in the art.

Methods of Synthesizing Disclosed Compounds

[00173] The compounds of the present disclosure may be made by a variety of methods, including standard chemistry. Suitable synthetic routes are depicted in the schemes given below.

[00174] The compounds of any of the formulae described herein may be prepared by methods known in the art of organic synthesis as set forth in part by the following synthetic schemes and examples. In the schemes described below, it is well understood that protecting groups for sensitive or reactive groups are employed where necessary in accordance with general principles of chemistry. Protecting groups are manipulated according to standard methods of organic synthesis (T. W. Greene and P. G. M. Wuts, "Protective Groups in Organic Synthesis", Third edition, Wiley, New York 1999). These groups are removed at a convenient stage of the compound synthesis using methods that are readily apparent to those skilled in the art. The selection processes, as well as the reaction conditions and order of their execution, shall be consistent with the preparation of compounds of Formula I, Ia, Ib, II, or IIb, or a pharmaceutically acceptable salt or tautomer of any of the foregoing.

[00175] The compounds of any of the formulae described herein may be prepared by methods which avoid the use of metal-mediated cycloaddition reactions which require the use of azide-containing compounds. Azide containing compounds present potential safety hazards associated with their preparation and storage (e.g., explosion due to high energy decomposition). Also, the reaction schemes herein can avoid the use of copper or ruthenium metals in the penultimate or ultimate synthetic steps, which can be advantageous. Avoiding the use of copper or ruthenium metals in the penultimate or ultimate synthetic steps reduces the potential for contamination of the final compounds with undesirable metal impurities.

[00176] As rapamycin can be an expensive starting material, good yields on reactions are advantageous. The reaction schemes herein provide better yields than other reaction schemes. In the reaction schemes herein, there is no need to alkylate at the C40-hydroxyl of rapamycin, which is advantageous for providing as much as a 5-fold improved overall yield in preparing bivalent compounds from rapamycin compared to other reaction schemes.

[00177] There is an additional synthetic improvement associated with better yields. Avoiding the need to alkylate at the C40-hydroxyl gives as much as a 5-fold improved overall yield in preparing bivalent compounds from rapamycin.

[00178] Those skilled in the art will recognize if a stereocenter exists in any of the compounds of the present disclosure. Accordingly, the present disclosure may include both possible stereoisomers (unless specified in the synthesis) and may include not only racemic compounds but the individual enantiomers and/or diastereomers as well. When a compound is desired as a single enantiomer or diastereomer, it may be obtained by stereospecific synthesis or by resolution of the final product or any convenient intermediate. Resolution of the final product, an intermediate, or a starting material may be effected by any suitable method known in the art. See, for example, "Stereochemistry of Organic Compounds" by E. L. Eliel, S. H. Wilen, and L. N. Mander (Wiley-Interscience, 1994).


Preparation of Compounds

[00179] The compounds described herein may be made from commercially available starting materials or synthesized using known organic, inorganic, and/or enzymatic processes.

[00180] The compounds of the present disclosure can be prepared in a number of ways well known to those skilled in the art of organic synthesis. By way of example, compounds of the disclosure can be synthesized using the methods described below, together with synthetic methods known in the art of synthetic organic chemistry, or variations thereon as appreciated by those skilled in the art. These methods may include but are not limited to those methods described below.

[00181] The term "tautomers" may refer to a set of compounds that have the same number and type of atoms, but differ in bond connectivity and are in equilibrium with one another. A "tautomer" is a single member of this set of compounds. Typically a single tautomer is drawn but it may be understood that this single structure may represent all possible tautomers that might exist. Examples may include enol-ketone tautomerism. When a ketone is drawn it may be understood that both the enol and ketone forms are part of the disclosure.


[00182] In addition to tautomers that may exist at all amide, carbonyl, and oxime groups within compounds of Formula I, Ia, Ib, Ic, II, or IIb, compounds in this family readily interconvert via a ring-opened species between two major isomeric forms, known as the pyran and oxepane isomers (shown below). This interconversion can be promoted by magnesium ions, mildly acidic conditions, or alkylamine salts, as described in the following references: i) Hughes, P. F.; Musser, J.; Conklin, M.; Russo, R. 1992. *Tetrahedron Lett.* 33(33): 4739-32. ii) Zhu, T. 2007. U.S. Patent 7,241,771; Wyeth. iii) Hughes, P.F. 1994. U.S. Patent 5,344,833; American Home Products Corp. The scheme below shows an interconversion between the pyran and oxepane isomers in compounds of Formula I, Ia, Ib, Ic, II, or IIb.



[00183] As this interconversion occurs under mild condition, and the thermodynamic equilibrium position may vary between different members of compounds of Formula I, Ia, Ib, Ic, II, or IIb, both isomers are contemplated for the compounds of Formula I, Ia, Ib, Ic, II, or IIb. For the sake of brevity, the pyran isomer form of all intermediates and compounds of Formula I, Ia, Ib, Ic, II, or IIb is shown.

General Assembly Approaches For Bifunctional Rapalogs

[00184] With reference to the schemes below, rapamycin is Formula RAP,




where R^{16} is $-OCH_3$; R^{26} is $=O$; R^{28} is $-OH$; R^{32} is $=O$; and R^{40} is $-OH$. A “rapalog” refers to an analog or derivative of rapamycin. For example, with reference to the schemes below, a rapalog can be rapamycin that is substituted at any position, such as R^{16} , R^{26} , R^{28} , R^{32} , or R^{40} . An active site inhibitor (AS inhibitor) is an active site mTOR inhibitor. In certain embodiments, AS inhibitor is depicted by B, in Formula I, Ia, Ib, Ic, II, or IIb.

Series 1 bifunctional rapalogs

[00185] A general structure of Series 1 bifunctional rapalogs is shown in Scheme 1 below. For these types of bifunctional rapalogs, the linker may include variations where $q = 0$ to 30, such as $q = 1$ to 7, and $r = 1$ to 6. The linker amine can include substitutions, such as $R = H$ and C1–C6 alkyl groups. The carbamate moiety, where $Z^1 = O$ or S, can be attached to the rapalog at R^{40} or R^{28} (Formula I, Ia, Ib, Ic, II, or IIb), including variations found in Table 1 in the Examples Section. An mTOR active site inhibitor can attach to the linker via a primary or secondary amine, and may include variations found in Table 2 in the Examples Section.

Scheme 1. Series 1 bifunctional rapalogs.




Series 2 bifunctional rapalogs

[00186] A general structure of Series 2 bifunctional rapalogs is shown in Scheme 2 below. For these types of bifunctional rapalogs, the linker may include variations where $q = 0$ to 30, such as $q = 1$ to 7. The linker amine can include substitutions, such as $R = H$ and C1–C6

alkyl groups. The pre-linker amine can include substitutions, such as $R^2 = H$, C1–C6 alkyl groups, and cycloalkyl including 4 to 8-membered rings. The carbamate moiety, where $Z^1 = O$ or S, can be attached to the rapalog at R^{40} or R^{28} (Formula I, Ia, Ib, Ic, II, or IIb), including variations found in Table 1 in the Examples Section. An mTOR active site inhibitor can attach to the linker via a primary or secondary amine, and may include variations found in Table 2 in the Examples Section.


Scheme 2. Series 2 bifunctional rapalogs.



Series 3 bifunctional rapalogs

[00187] A general structure of Series 3 bifunctional rapalogs is shown in Scheme 3 below. For these types of bifunctional rapalogs, the linker may include variations where $q = 0$ to 30, such as $q = 1$ to 7. The linker amine can include substitutions, such as $R = H$ and C1–C6 alkyl groups. The post-linker amine can include substitutions, such as $R^2 = H$, C1–C6 alkyl groups, and cycloalkyl including 4 to 8-membered rings. The carbamate moiety, where $Z^1 = O$ or S, can be attached to the rapalog at R^{40} or R^{28} (Formula I, Ia, Ib, Ic, II, or IIb), including variations found in Table 1 in the Examples Section. An mTOR active site inhibitor can attach to the linker via a primary or secondary amine, and may include variations found in Table 2 in the Examples Section.


Scheme 3. Series 3 bifunctional rapalogs



Series 4 bifunctional rapalogs


[00188] A general structure of Series 4 bifunctional rapalogs is shown in Scheme 4 below. For these types of bifunctional rapalogs, the linker may include variations where $q = 0$ to 30, such as $q = 1$ to 7. The linker amine can include substitutions, such as $R = H$ and C1–C6 alkyl groups. The pre- and post-linker amines can each include substitutions, such as $R^2 = H$, C1–C6 alkyl groups, and cycloalkyl including 4 to 8-membered rings. The carbamate moiety, where $Z^1 = O$ or S, can be attached to the rapalog at R^{40} or R^{28} (Formula I, Ia, Ib, Ic, II, or IIb), including variations found in Table 1 in the Examples Section. An mTOR active site inhibitor can attach to the linker via a primary or secondary amine, and may include variations found in Table 2 in the Examples Section.

Scheme 4. Series 4 bifunctional rapalogs

Series 5 bifunctional rapalogs

[00189] A general structure of Series 5 bifunctional rapalogs is shown in Scheme 5 below. For these types of bifunctional rapalogs, the pre-linker amine can include substitutions, such as $R^2 = H$, C1–C6 alkyl groups, and cycloalkyl including 4 to 8-membered rings. The carbamate moiety, where $Z^1 = O$ or S, can be attached to the rapalog at R^{40} or R^{28} (Formula I, Ia, Ib, Ic, II, or IIb), including variations found in Table 1 in the Examples Section. An mTOR active site inhibitor can attach to the linker via a primary or secondary amine, and may include variations found in Table 2 in the Examples Section.


Scheme 5. Series 5 bifunctional rapalogs



Series 6 bifunctional rapalogs


[00190] A general structure of Series 6 bifunctional rapalogs is shown in Scheme 6 below. For these types of bifunctional rapalogs, the linker may include variations where $q = 0$ to 30, such as $q = 1$ to 7. The linker amine can include substitutions, such as $R = H$ and C1–C6 alkyl groups. The post-linker amine can include substitutions, such as $R^2 = H$, C1–C6 alkyl groups, and cycloalkyl including 4 to 8-membered rings. The carbamate moiety, where $Z^1 = O$ or S , can be attached to the rapalog at R^{40} or R^{28} (Formula I, Ia, Ib, Ic, II, or IIb), including variations found in Table 1 in the Examples Section. An mTOR active site inhibitor can attach to the linker via a primary or secondary amine, and may include variations found in Table 2 in the Examples Section.

Scheme 6. Series 6 bifunctional rapalogs.

Series 7 bifunctional rapalogs


[00191] A general structure of Series 7 bifunctional rapalogs is shown in Scheme 7 below. For these types of bifunctional rapalogs, the linker may include variations where $q = 0$ to 30, such as $q = 1$ to 7. The linker amine can include substitutions, such as $R = H$ and C1–C6 alkyl groups. The pre- and post-linker amines can each include substitutions, such as $R^2 = H$, C1–C6 alkyl groups, and cycloalkyl including 4 to 8-membered rings. The carbamate moiety, where $Z^1 = O$ or S , can be attached to the rapalog at R^{40} or R^{28} (Formula I, Ia, Ib, Ic, II, or IIb), including variations found in Table 1 in the Examples Section. An mTOR active site inhibitor can attach to the linker via a primary or secondary amine, and may include variations found in Table 2 in the Examples Section.

Scheme 7. Series 7 bifunctional rapalogs

Series 8 bifunctional rapalogs

[00192] A general structure of Series 8 bifunctional rapalogs is shown in Scheme 8 below. For these types of bifunctional rapalogs, the linker may include variations where $q = 0$ to 30 , such as $q = 1$ to 7 . The linker amine can include substitutions, such as $R = H$ and C1–C6 alkyl groups. The post-linker amine can include substitutions, such as $R^2 = H$, C1–C6 alkyl groups, and cycloalkyl including 4 to 8-membered rings. The carbamate moiety, where $Z^1 = O$ or S , can be attached to the rapalog at R^{40} or R^{28} (Formula I, Ia, Ib, Ic, II, or IIb), including variations found in Table 1 in the Examples Section. An mTOR active site inhibitor can attach to the linker via a primary or secondary amine, and may include variations found in Table 2 in the Examples Section.

Scheme 8. Series 8 bifunctional rapalogs

**Pharmaceutical Compositions**

[00193] Another aspect provides a pharmaceutical composition including a pharmaceutically acceptable excipient and a compound of the present invention, or pharmaceutically acceptable salt or tautomer thereof.

[00194] In embodiments of the pharmaceutical compositions, a compound of the present invention, or a pharmaceutically acceptable salt or tautomer thereof, may be included in a therapeutically effective amount.

[00195] Administration of the disclosed compounds or compositions can be accomplished via any mode of administration for therapeutic agents. These modes may include systemic or local administration such as oral, nasal, parenteral, transdermal, subcutaneous, vaginal, buccal, rectal, topical, intrathecal, or intracranial administration modes.

[00196] In certain embodiments, administering can include oral administration, administration as a suppository, topical contact, intravenous, parenteral, intraperitoneal, intramuscular, intralesional, intrathecal, intracranial, intranasal or subcutaneous administration, or the implantation of a slow-release device, e.g., a mini-osmotic pump, to a subject. Administration can be by any route, including parenteral and transmucosal (e.g., buccal, sublingual, palatal, gingival, nasal, vaginal, rectal, or transdermal). Parenteral administration includes, e.g., intravenous, intramuscular, intra-arteriole, intradermal, subcutaneous, intraperitoneal, intraventricular, and intracranial. Other modes of delivery include, but are not limited to, the use of liposomal formulations, intravenous infusion, transdermal patches, etc. The compositions of the present disclosure can be delivered by transdermally, by a topical route, formulated as applicator sticks, solutions, suspensions, emulsions, gels, creams, ointments, pastes, jellies, paints, powders, and aerosols. Oral preparations include tablets, pills, powder, dragees, capsules, liquids, lozenges, cachets, gels, syrups, slurries, suspensions, etc., suitable for ingestion by the patient. Solid form preparations include powders, tablets, pills, capsules, cachets, suppositories, and dispersible granules. Liquid form preparations include solutions, suspensions, and emulsions, for example, water or water/propylene glycol solutions. The compositions of the present disclosure may additionally include components to provide sustained release and/or comfort. Such components include high molecular weight, anionic mucomimetic polymers, gelling polysaccharides and finely-divided drug carrier substrates. These components are discussed in greater detail in U.S. Pat. Nos. 4,911,920; 5,403,841; 5,212,162; and 4,861,760. The entire contents of these patents are incorporated herein by reference in their entirety for all purposes. The compositions of the present disclosure can also be delivered as microspheres for slow release in the body. For example, microspheres can be administered via intradermal injection of drug-containing microspheres, which slowly release subcutaneously (see Rao, J. Biomater Set Polym. Ed. 7:623-645, 1995; as biodegradable and injectable gel formulations (see, e.g., Gao Pharm. Res. 12:857-863, 1995); or, as microspheres for oral administration (see, e.g., Eyles, J. Pharm. Pharmacol. 49:669-674, 1997). In another embodiment, the formulations of the compositions of the present disclosure can be delivered by the use of

liposomes which fuse with the cellular membrane or are endocytosed, i.e., by employing receptor ligands attached to the liposome, that bind to surface membrane protein receptors of the cell resulting in endocytosis. By using liposomes, particularly where the liposome surface carries receptor ligands specific for target cells, or are otherwise preferentially directed to a specific organ, one can focus the delivery of the compositions of the present invention into the target cells *in vivo*. (See, e.g., Al-Muhammed, J. Microencapsul. 13:293-306, 1996; Chonn, Curr. Opin. Biotechnol. 6:698-708, 1995; Ostro, Am. J. Hosp. Pharm. 46: 1576-1587, 1989). The compositions of the present disclosure can also be delivered as nanoparticles.

[00197] Depending on the intended mode of administration, the disclosed compounds or pharmaceutical compositions can be in solid, semi-solid or liquid dosage form, such as, for example, injectables, tablets, suppositories, pills, time-release capsules, elixirs, tinctures, emulsions, syrups, powders, liquids, suspensions, or the like, sometimes in unit dosages and consistent with conventional pharmaceutical practices. Likewise, they can also be administered in intravenous (both bolus and infusion), intraperitoneal, intrathecal, subcutaneous or intramuscular form, and all using forms well known to those skilled in the pharmaceutical arts.

[00198] Illustrative pharmaceutical compositions are tablets and gelatin capsules comprising a compound of the disclosure and a pharmaceutically acceptable carrier, such as a) a diluent, e.g., purified water, triglyceride oils, such as hydrogenated or partially hydrogenated vegetable oil, or mixtures thereof, corn oil, olive oil, sunflower oil, safflower oil, fish oils, such as EPA or DHA, or their esters or triglycerides or mixtures thereof, omega-3 fatty acids or derivatives thereof, lactose, dextrose, sucrose, mannitol, sorbitol, cellulose, sodium, saccharin, glucose and/or glycine; b) a lubricant, e.g., silica, talcum, stearic acid, its magnesium or calcium salt, sodium oleate, sodium stearate, magnesium stearate, sodium benzoate, sodium acetate, sodium chloride and/or polyethylene glycol; for tablets also; c) a binder, e.g., magnesium aluminum silicate, starch paste, gelatin, tragacanth, methylcellulose, sodium carboxymethylcellulose, magnesium carbonate, natural sugars such as glucose or beta-lactose, corn sweeteners, natural and synthetic gums such as acacia, tragacanth or sodium alginate, waxes and/or polyvinylpyrrolidone, if desired; d) a disintegrant, e.g., starches, agar, methyl cellulose, bentonite, xanthan gum, alginic acid or its sodium salt, or effervescent mixtures; e) absorbent, colorant, flavorant and sweetener; f) an emulsifier or dispersing agent, such as Tween 80, Labrasol, HPMC, DOSS, caproyl 909, labrafac, labrafil,

peceol, transcitol, capmul MCM, capmul PG-12, captex 355, gelucire, vitamin E TGPS or other acceptable emulsifier; and/or g) an agent that enhances absorption of the compound such as cyclodextrin, hydroxypropyl-cyclodextrin, PEG400, PEG200.

[00199] Liquid, particularly injectable, compositions can, for example, be prepared by dissolution, dispersion, etc. For example, the disclosed compound is dissolved in or mixed with a pharmaceutically acceptable solvent such as, for example, water, saline, aqueous dextrose, glycerol, ethanol, and the like, to thereby form an injectable isotonic solution or suspension. Proteins such as albumin, chylomicron particles, or serum proteins can be used to solubilize the disclosed compounds.

[00200] The disclosed compounds can be also formulated as a suppository that can be prepared from fatty emulsions or suspensions; using polyalkylene glycols such as propylene glycol, as the carrier.

[00201] The disclosed compounds can also be administered in the form of liposome delivery systems, such as small unilamellar vesicles, large unilamellar vesicles and multilamellar vesicles. Liposomes can be formed from a variety of phospholipids, containing cholesterol, stearylamine or phosphatidylcholines. In some embodiments, a film of lipid components is hydrated with an aqueous solution of drug to a form lipid layer encapsulating the drug, as described for instance in U.S. Pat. No. 5,262,564, the contents of which are hereby incorporated by reference.

[00202] Disclosed compounds can also be delivered by the use of monoclonal antibodies as individual carriers to which the disclosed compounds are coupled. The disclosed compounds can also be coupled with soluble polymers as targetable drug carriers. Such polymers can include polyvinylpyrrolidone, pyran copolymer, polyhydroxypropylmethacrylamide-phenol, polyhydroxyethylaspanamidephenol, or polyethyleneoxidepolylysine substituted with palmitoyl residues. Furthermore, the disclosed compounds can be coupled to a class of biodegradable polymers useful in achieving controlled release of a drug, for example, polylactic acid, polyepsilon caprolactone, polyhydroxy butyric acid, polyorthoesters, polyacetals, polydihydropyrans, polycyanoacrylates and cross-linked or amphipathic block copolymers of hydrogels. In one embodiment, disclosed compounds are not covalently bound to a polymer, e.g., a polycarboxylic acid polymer, or a polyacrylate.

[00203] Parenteral injectable administration is generally used for subcutaneous, intramuscular or intravenous injections and infusions. Injectables can be prepared in conventional forms, either as liquid solutions or suspensions or solid forms suitable for dissolving in liquid prior to injection.

[00204] Another aspect of the disclosure relates to a pharmaceutical composition comprising a compound, or a pharmaceutically acceptable salt or tautomer thereof, of the present disclosure and a pharmaceutically acceptable carrier. The pharmaceutically acceptable carrier can further include an excipient, diluent, or surfactant.

[00205] Compositions can be prepared according to conventional mixing, granulating or coating methods, respectively, and the present pharmaceutical compositions can contain from about 0.1% to about 99%, from about 5% to about 90%, or from about 1% to about 20% of the disclosed compound by weight or volume.

[00206] The compounds described herein can be used in combination with one another, with other active agents known to be useful in treating cancer, autoimmune disease, inflammatory disease, metabolic disease, neurodegenerative disease, fungal infection, or transplant rejection, or with adjunctive agents that may not be effective alone, but may contribute to the efficacy of the active agent. The compounds described herein can be used in combination with other active agents known to be longevity agents or anti-aging agents.

[00207] In embodiments of the pharmaceutical compositions, the pharmaceutical composition may include a second agent (e.g., therapeutic agent). In embodiments of the pharmaceutical compositions, the pharmaceutical composition may include a second agent (e.g., therapeutic agent) in a therapeutically effective amount. In certain embodiments, the second agent is an anti-cancer agent. In certain embodiments, the second agent is an immunotherapeutic agent. In certain embodiments, the second agent is an immune-oncological agent. In certain embodiments, the second agent is an anti-autoimmune disease agent. In embodiments, the second agent is an anti-inflammatory disease agent. In certain embodiments, the second agent is an anti-neurodegenerative disease agent. In certain embodiments, the second agent is an anti-metabolic disease agent. In certain embodiments, the second agent is an anti-cardiovascular disease agent. In certain embodiments, the second agent is an anti-aging agent. In certain embodiments, the second agent is a longevity agent. In certain embodiments, the second agent is an agent for treating or preventing transplant rejection. In certain embodiments, the second agent is an agent for treating or preventing

fungal infection. In certain embodiments, the second agent is immune system repressor. In certain embodiments, the second agent is an mTOR modulator. In certain embodiments, the second agent is an mTOR inhibitor. In certain embodiments, the second agent is an active site mTOR inhibitor. In certain embodiments, the second agent is a rapamycin. In certain embodiments, the second agent is a rapamycin analog. In certain embodiments, the second agent is an mTOR pathway inhibitor. In certain embodiments, the second agent is CDK4/6 inhibitor; anti-PD1/PD-L1, PI3K inhibitor; or Ras inhibitor.

[00208] “Anti-cancer agent” or “anti-cancer drug” is used in accordance with its plain ordinary meaning and refers to a composition (e.g. compound, drug, antagonist, inhibitor, modulator) having antineoplastic properties or the ability to inhibit the growth or proliferation of cells. In some embodiments, an anti-cancer agent is a chemotherapeutic. In some embodiments, an anticancer agent is an agent approved by the FDA or similar regulatory agency of a country other than the USA, for treating cancer. Examples of anti-cancer agents include, but are not limited to, rapamycin, rapamycin analog, bevacizumab, PP242, FN 128, MLN0128, anti-androgens (e.g., Casodex, Flutamide, MDV3100, or ARN-509), MEK (e.g. MEK1, MEK2, or MEK1 and MEK2) inhibitors (e.g. XL518, CI-1040, PD035901, selumetinib/ AZD6244, GSK1 120212/ trametinib, GDC-0973, ARRY-162, ARRY-300, AZD8330, PD0325901, U0126, PD98059, TAK-733, PD318088, AS703026, BAY 869766), alkylating agents (e.g., cyclophosphamide, ifosfamide, chlorambucil, busulfan, melphalan, mechlorethamine, uramustine, thiotepa, nitrosoureas, nitrogen mustards (e.g., mechloroethamine, cyclophosphamide, chlorambucil, melphalan), ethylenimine and methylmelamines (e.g., hexamethylmelamine, thiotepa), alkyl sulfonates (e.g., busulfan), nitrosoureas (e.g., carmustine, lomustine, semustine, streptozocin), triazenes (decabazine)), anti-metabolites (e.g., 5- azathioprine, leucovorin, capecitabine, fludarabine, gemcitabine, pemetrexed, raltitrexed, folic acid analog (e.g., methotrexate), pyrimidine analogs (e.g., fluorouracil, floxouridine, Cytarabine), purine analogs (e.g., mercaptopurine, thioguanine, pentostatin), etc.), plant alkaloids (e.g., vincristine, vinblastine, vinorelbine, vindesine, podophyllotoxin, paclitaxel, docetaxel, etc.), topoisomerase inhibitors (e.g., irinotecan, topotecan, amsacrine, etoposide (VP 16), etoposide phosphate, teniposide, etc.), antitumor antibiotics (e.g., doxorubicin, adriamycin, daunorubicin, epirubicin, actinomycin, bleomycin, mitomycin, mitoxantrone, plicamycin, etc.), platinum-based compounds (e.g. cisplatin, oxaloplatin, carboplatin), anthracenedione (e.g., mitoxantrone), substituted urea (e.g., hydroxyurea), methyl hydrazine derivative (e.g., procarbazine), adrenocortical suppressant

(e.g., mitotane, aminoglutethimide), epipodophyllotoxins (e.g., etoposide), antibiotics (e.g., daunorubicin, doxorubicin, bleomycin), enzymes (e.g., L-asparaginase), inhibitors of mitogen-activated protein kinase signaling (e.g. U0126, PD98059, PD184352, PD0325901, ARRY-142886, SB239063, SP600125, BAY 43-9006, wortmannin, or LY294002), mTOR inhibitors, antibodies (e.g., rituxan), 5-aza-2'-deoxycytidine, doxorubicin, vincristine, etoposide, gemcitabine, imatinib (Gleevec.RTM.), geldanamycin, 17-N-Allylamino-17-Demethoxygeldanamycin (17-AAG), bortezomib, trastuzumab, anastrozole; angiogenesis inhibitors; antiandrogen, antiestrogen; antisense oligonucleotides; apoptosis gene modulators; apoptosis regulators; arginine deaminase; BCR/ABL antagonists; beta lactam derivatives; bFGF inhibitor; bicalutamide; camptothecin derivatives; casein kinase inhibitors (ICOS); clomifene analogues; cytarabine dacliximab; dexamethasone; estrogen agonists; estrogen antagonists; etanidazole; etoposide phosphate; exemestane; fadrozole; finasteride; fludarabine; fluorodaunorubicin hydrochloride; gadolinium texaphyrin; gallium nitrate; gelatinase inhibitors; gemcitabine; glutathione inhibitors; hepsulfam; immunostimulant peptides; insulin-like growth factor- 1 receptor inhibitor; interferon agonists; interferons; interleukins; letrozole; leukemia inhibiting factor; leukocyte alpha interferon; leuprolide+estrogen+progesterone; leuprorelin; matrilysin inhibitors; matrix metalloproteinase inhibitors; MIF inhibitor; mifepristone; mismatched double stranded RNA; monoclonal antibody;; mycobacterial cell wall extract; nitric oxide modulators; oxaliplatin; panomifene; pentrozole; phosphatase inhibitors; plasminogen activator inhibitor; platinum complex; platinum compounds; prednisone; proteasome inhibitors; protein A-based immune modulator; protein kinase C inhibitor; protein tyrosine phosphatase inhibitors; purine nucleoside phosphorylase inhibitors; ras farnesyl protein transferase inhibitors; ras inhibitors; ras-GAP inhibitor; ribozymes; signal transduction inhibitors; signal transduction modulators; single chain antigen-binding protein; stem cell inhibitor; stem-cell division inhibitors; stromelysin inhibitors; synthetic glycosaminoglycans; tamoxifen methiodide; telomerase inhibitors; thyroid stimulating hormone; translation inhibitors; tyrosine kinase inhibitors; urokinase receptor antagonists; steroids (e.g., dexamethasone), finasteride, aromatase inhibitors, gonadotropin-releasing hormone agonists (GnRH) such as goserelin or leuprolide, adrenocorticosteroids (e.g., prednisone), progestins (e.g., hydroxyprogesterone caproate, megestrol acetate, medroxyprogesterone acetate), estrogens (e.g., diethylstilbestrol, ethinyl estradiol), antiestrogen (e.g., tamoxifen), androgens (e.g., testosterone propionate, fluoxymesterone), antiandrogen (e.g., flutamide), immunostimulants (e.g., Bacillus Calmette-Guerin (BCG), levamisole, interleukin-2, alpha-interferon, etc.), monoclonal antibodies (e.g.,

anti-CD20, anti-HER2, anti-CD52, anti-HLA-DR, and anti-VEGF monoclonal antibodies), immunotoxins (e.g., anti-CD33 monoclonal antibody-calicheamicin conjugate, anti-CD22 monoclonal antibody-pseudomonas exotoxin conjugate, etc.), radioimmunotherapy (e.g., anti-CD20 monoclonal antibody conjugated to ^{111}In , ^{90}Y , or ^{131}I , etc.), triptolide, homoharringtonine, dactinomycin, doxorubicin, epirubicin, topotecan, itraconazole, vindesine, cerivastatin, vincristine, deoxyadenosine, sertraline, pitavastatin, irinotecan, clofazimine, 5-nonyloxytryptamine, vemurafenib, dabrafenib, erlotinib, gefitinib, EGFR inhibitors, epidermal growth factor receptor (EGFR)-targeted therapy or therapeutic (e.g. gefitinib (IressaTM), erlotinib (TarcevaTM), cetuximab (ErbixTM), lapatinib (TykerbTM), panitumumab (VectibixTM), vandetanib (CaprelsaTM), afatinib/BIBW2992, CI-1033/canertinib, neratinib/HKI-272, CP-724714, TAK-285, AST-1306, ARRY334543, ARRY-380, AG-1478, dacomitinib/PF299804, OSI-420/desmethyl erlotinib, AZD8931, AEE788, pelitinib/EKB-569, CUDC-101, WZ8040, WZ4002, WZ3146, AG-490, XL647, PD153035, BMS-599626), sorafenib, imatinib, sunitinib, dasatinib, pyrrolo benzodiazepines (e.g. tomaymycin), carboplatin, CC-1065 and CC-1065 analogs including amino-CBIs, nitrogen mustards (such as chlorambucil and melphalan), dolastatin and dolastatin analogs (including auristatins: eg. monomethyl auristatin E), anthracycline antibiotics (such as doxorubicin, daunorubicin, etc.), duocarmycins and duocarmycin analogs, enediynes (such as neocarzinostatin and calicheamicins), leptomycin derivatives, maytansinoids and maytansinoid analogs (e.g. mertansine), methotrexate, mitomycin C, taxoids, vinca alkaloids (such as vinblastine and vincristine), epothilones (e.g. epothilone B), camptothecin and its clinical analogs topotecan and irinotecan, GSK128, PP242, PP121, MLN0128, AZD8055, AZD2014, VP-BEZ235, BGT226, SFI 126, Torin 1, Torin 2, WYE 687, WYE 687 salt (e.g., hydrochloride), PF04691502, PI- 103, CC-223, OSI-027, XL388, KU-0063794, GDC-0349, PKI-587, rapamycin, deforolimus (AP23573, MK-8669, ridaforolimus), temsirolimus (CCI-779), ABT478, everolimus (RAD001) or the like.

mTOR and Methods of Treatment

[00209] The term “mTOR” refers to the protein “mechanistic target of rapamycin (serine/threonine kinase)” or “mammalian target of rapamycin.” The term “mTOR” may refer to the nucleotide sequence or protein sequence of human mTOR (e.g., Entrez 2475, Uniprot P42345, RefSeq NM_004958, or RefSeq NP_004949) (SEQ ID NO: 1). The term “mTOR” may include both the wild-type form of the nucleotide sequences or proteins as well as any mutants thereof. In some embodiments, “mTOR” is wild-type mTOR. In some

embodiments, “mTOR” is one or more mutant forms. The term “mTOR” XYZ may refer to a nucleotide sequence or protein of a mutant mTOR wherein the Y numbered amino acid of mTOR that normally has an X amino acid in the wildtype, instead has a Z amino acid in the mutant. In embodiments, an mTOR is the human mTOR. In embodiments, the mTOR has the nucleotide sequence corresponding to reference number GL206725550 (SEQ ID NO:2). In embodiments, the mTOR has the nucleotide sequence corresponding to RefSeq NM_004958.3 (SEQ ID NO:2). In embodiments, the mTOR has the protein sequence corresponding to reference number GL4826730 (SEQ ID NO: 1). In embodiments, the mTOR has the protein sequence corresponding to RefSeq NP_004949.1 (SEQ ID NO: 1). In embodiments, the mTOR has the following amino acid sequence:

MLGTGPAAATTAATTSSNVSVLQQFASGLKSRNEETRAKAAKELQHYVTMELREMSQEESTRFYDQLNHHI
 FELVSSSDANERKGGILAIASLIGVEGGNATRIGRFANYLRNLLPSNDPWMEMASKAIGRLAMAGDTF
 TAEYVEFEVKRALEWLGAADRNEGRRHA AVLVLRELAI SVPTFFFQVQPFDFNIFVAVWDPKQAIREGAV
 AALRACLILTTQREPKEKMPQWYRHTFEEAEKGFDETLAKEKGMNRDDRIHGALLILNELVRISSMEGE
 RLREEMEEITQQQLVHDKYCKDLMGFGTKPRHITPFTSFQAVQPQQSNALVGLLGYSSHQGLMGFGTSPS
 PAKSTLVESRCCRDLMEEKFDQVCQWVLKCRNSKNSLIQMTILNLLPRLAAFRPSAFTDTQYLQDTMNHV
 LSCVKKEKERTAAAFQALGLLSVAVRSEFKVYLPRVLDIIRAALPPKDFAHKRQKAMQVDATVFTCISMLA
 RAMGPGIQQDIKELLEPLAVGLSPALTAVLYDLRQIPQLKKDIQDGLLKMLSLVLMHKPLRHPGMPKG
 LAHQLASPLTTLPEASDVGSITLALRTLGSFEFEGHSLTQFVRHCADHFLNSEHKEIRMEAAARTCSRL
 TPSIHLISGHAHVVSQTAVQVADVLSKLLWGITDPDPDIRYCVLASLDERFDAHLAQENLQALFVAL
 NDQVFEIRELAICTVGRLLSSMNPAPFVMPFLRKMLIQILTELEHSGIGRIKEQSARMLGHLVSNAPRLIRP
 YMEPILKALILKLKDPDPDPNPGVINNVLATIGELAQVSGLEMRKWVDELFIIMDMLQDSSLLAKRQVA
 LWTGLQVLVASTGYVVEPYRKYPTLLEVLNLFKTEQNQGTTRREAIRVLGGLGALDPYKHKVNIGMIDQSR
 DASAVSLSEKSSQDSSDYSTSEMLVNMGNLPLDEFYPAVSMVALMRIFRDQSLSHHTMVVQAITFIFK
 SLGLKCVQFLPQVMPTFLNVIRVCDGAIREFLFQQLGMLVSFVKSHIRPYMDEIVTLMREFWVMNTSIQS
 TIILLIEQIVVALGGEFKLYLPQLIPHMLRVFMHDNSPGRIVSIKLLAAIQLFGANLDDYLHLLLPPIVK
 LFDAPEAPLPSRKAALLETVDRLTESLDFTDYASRIIPIVRTLDQSPELRSTAMDTLSSLVFQLGKKYQI
 FIPMVNKKVLVRHRINHQRVDVLCRIVKGYTLADEEEDPLIYQHRMLRSGQGDALASGPVETGPMKKLHV
 STINLQKAWGAARRVSKDDWLEWLRRLSLELLKDSSPSLRSCWALAQAYNPMARDLFNAAFVSCWSELN
 EDQQDELIRSIELALTSQDIAEVTQTLLNLAEFMEHSDKGPLPLRDDNGIVLLGERAAKCRAYAKALHYK
 ELEFQKGPTPAILESISINNKLQQPEAAAGVLEYAMKHFGELIQA TWYEKLHEWEDALVAYDKKMDTN
 KDDPELMLGRMRCLEALGEWGQLHQQCCEKWTLVNDETQAKMARMAAAAAWGLGQWDSMEEYTCMIP
 RDTHDGA FYRAVLALHQDLFSLAQQCIDKARDLLDAELTAMAGESYSRAYGAMV SCHMLSELEEVIQYKL
 VPERREIIRQIWWERLQGCQRIVEDWQKILMVRSLVSPHEDMRTWLKYASLCGKSGRLALAHKTLVLLG
 VDPSRQLDHPLPTVHPQVTYAYMKNMWKSARKIDAFQHMQHVFVQTMQQAQHA IATEDQQHKQELHKL
 MARCFLKLGEWQLNLQGINESTIPKVLQYYSAATEHDSWYKAWHAWAVMNFEAVLHYKHQNQARDEK
 KKL RHASGANITNATTAATTAATATTTASTEGSNSSEEAESTENSPTPSPLQKKVTEDLSKTLTLMYTPVAVQG
 FFRSISLSRGNLQDTLRVLTLWFDYGHWPDVNEALVEGVKAIQIDTWLQVIPQLIARIDTPRPLVGRLIHQ
 L TDIGRYHPQALYPLTVASKSTTTARHNAANKILKNMCEHSNTLVQQAMMVSEELIRVAILWHEMWHEG

LEEASRLYFGERNVKGMFEVLEPLHAMMERGPQTLKETSFNQAYGRDLMEAQEWCRKYMKSGNVKDLTQ
 A WDLYYHVFRISKQLPQLTSLELQYVSPKLLMCRDLELAVPGTYDPNQPIRIQSIAPSLQVITSKQRPR
 KLTLMGSGNGHEFVFLKGHEDLRQDERVMQLFGLVNTLLANDPTSLRKNLSIQRYAVIPLSTNSGLIGWV
 PHCDTLHALIRDYREKKKILLNIEHRIMLRMAPDYDHLTLMQKVEVFEHAVNNTAGDDLAKLLWLKSPSS
 EVWFDRRTNYTRSLAVMSMVGYLGLGDRHPSNMLDRLSGKILHIDFGDCFEVAMTREKFPEKIPFRLT
 RMLTNAMEVTGLDGNYRITCHTVMEVLREHKDSVMAVLEAFVYDPLLNWRLMDTNTKGNKRSRTRTDSY
 S AGQSVEILDGVELGEPAHKKTGTTVPESHSFIGDGLVKPEALNKKAIQIINRVRDKLTGRDFSHDDTLD
 VPTQVELLIKQATSHENLCQCYIGWCPFW

(SEQ ID NO: 1)

[00210] In embodiments, the mTOR is a mutant mTOR. In embodiments, the mutant mTOR is associated with a disease that is not associated with wildtype mTOR. In embodiments, the mTOR may include at least one amino acid mutation (e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, or 30 mutations, or any range derivable therein) compared to the sequence above.

[00211] The term “mTORC1” refers to the protein complex including mTOR and Raptor (regulatory-associated protein of mTOR). mTORC1 may also include MLST8 (mammalian lethal with SEC 13 protein 8), PRAS40, and/or DEPTOR. mTORC1 may function as a nutrient/energy/redox sensor and regulator of protein synthesis. The term “mTORC1 pathway” or “mTORC1 signal transduction pathway” may refer to a cellular pathway including mTORC1. An mTORC1 pathway includes the pathway components upstream and downstream from mTORC1. An mTORC1 pathway is a signaling pathway that is modulated by modulation of mTORC1 activity. In embodiments, an mTORC1 pathway is a signaling pathway that is modulated by modulation of mTORC1 activity but not by modulation of mTORC2 activity. In embodiments, an mTORC1 pathway is a signaling pathway that is modulated to a greater extent by modulation of mTORC1 activity than by modulation of mTORC2 activity.

[00212] The term “mTORC2” refers to the protein complex including mTOR and RICTOR (rapamycin-insensitive companion of mTOR). mTORC2 may also include GβL, mSIN1 (mammalian stress-activated protein kinase interacting protein 1), Protor 1/2, DEPTOR, TTI1, and/or TEL2. mTORC2 may regulate cellular metabolism and the cytoskeleton. The term “mTORC2 pathway” or “mTORC2 signal transduction pathway” may refer to a cellular pathway including mTORC2. An mTORC2 pathway includes the pathway components upstream and downstream from mTORC2. An mTORC2 pathway is a signaling pathway that is modulated by modulation of mTORC2 activity. In embodiments, an

mTORC2 pathway is a signaling pathway that is modulated by modulation of mTORC2 activity but not by modulation of mTORC1 activity. In embodiments, an mTORC2 pathway is a signaling pathway that is modulated to a greater extent by modulation of mTORC2 activity than by modulation of mTORC1 activity.

[00213] The term “rapamycin” or “sirolimus” refers to a macrolide produced by the bacteria *Streptomyces hygroscopicus*. Rapamycin may prevent the activation of T cells and B cells. Rapamycin has the IUPAC name (3*S*,6*R*,7*E*,9*R*, 10*R*, 12*R*, 14*S*, 15*E*, 17*E*, 19*E*,21*S*,23*S*,26*R*,27*R*,34*aS*)- 9, 10, 12, 13, 14,21,22,23,24,25,26,27,32,33,34,34*a*-hexadecahydro-9,27-dihydroxy-3-[(1*R*)-2-[(1*S*,3*R*,4*R*)-4-hydroxy-3-methoxycyclohexyl] -1-methylethyl] - 10,21 -dimethoxy-6,8, 12, 14,20,26-hexamethyl-23,27-epoxy-3*H*-pyrido[2,1-*c*][1,4]-oxaazacyclohentriacontine-1,5, 11,28,29(4*H*,6*H*,31*H*)-pentone. Rapamycin has the CAS number 53123-88-9. Rapamycin may be produced synthetically (e.g., by chemical synthesis) or through use of a production method that does not include use of *Streptomyces hygroscopicus*.

[00214] “Analog” is used in accordance with its plain ordinary meaning within chemistry and biology and refers to a chemical compound that is structurally similar to another compound (i.e., a so-called “reference” compound) but differs in composition, e.g., in the replacement of one atom by an atom of a different element, or in the presence of a particular functional group, or the replacement of one functional group by another functional group, or the absolute stereochemistry of one or more chiral centers of the reference compound, including isomers thereof.

[00215] The term “rapamycin analog” or “rapalog” refers to an analog or derivative (e.g., a prodrug) of rapamycin.

[00216] The terms “active site mTOR inhibitor” and “ATP mimetic” refers to a compound that inhibits the activity of mTOR (e.g., kinase activity) and binds to the active site of mTOR (e.g., the ATP binding site, overlapping with the ATP binding site, blocking access by ATP to the ATP binding site of mTOR). Examples of active site mTOR inhibitors include, but are not limited to, GSK128, PP242, PP121, MLN0128, AZD8055, AZD2014, NVP-BEZ235, BGT226, SF1126, Torin 1, Torin 2, WYE 687, WYE 687 salt (e.g., hydrochloride), PF04691502, PI-103, CC-223, OSI-027, XL388, KU-0063794, GDC-0349, and PKI-587. In embodiments, an active site mTOR inhibitor is an asTORi. In some embodiments, “active site inhibitor” may refer to “active site mTOR inhibitor.”

[00217] The term “FKBP” refers to the protein Peptidyl-prolyl cis-trans isomerase. For non-limiting examples of FKBP, see Cell Mol Life Sci. 2013 Sep;70(18):3243-75. In embodiments, “FKBP” may refer to “FKBP-12” or “FKBP 12” or “FKBP 1 A.” In embodiments, “FKBP” may refer to the human protein. Included in the term “FKBP” is the wildtype and mutant forms of the protein. In embodiments, “FKBP” may refer to the wildtype human protein. In embodiments, “FKBP” may refer to the wildtype human nucleic acid. In embodiments, the FKBP is a mutant FKBP. In embodiments, the mutant FKBP is associated with a disease that is not associated with wildtype FKBP. In embodiments, the FKBP includes at least one amino acid mutation (e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, or 30 mutations, or any range derivable therein) compared to wildtype FKBP.

[00218] The term “FKBP-12” or “FKBP 12” or “FKBP1A” may refer to the protein “Peptidyl-prolyl cis-trans isomerase FKBP 1 A.” In embodiments, “FKBP-12” or “FKBP 12” or “FKBP 1 A” may refer to the human protein. Included in the term “FKBP-12” or “FKBP 12” or “FKBP 1 A” are the wildtype and mutant forms of the protein. In embodiments, “FKBP-12” or “FKBP 12” or “FKBP 1 A” may refer to the protein associated with Entrez Gene 2280, OMIM 186945, UniProt P62942, and/or RefSeq (protein) NP_000792 (SEQ ID NO:3). In embodiments, the reference numbers immediately above may refer to the protein, and associated nucleic acids, known as of the date of filing of this application. In embodiments, “FKBP-12” or “FKBP 12” or “FKBP 1 A” may refer to the wildtype human protein. In embodiments, “FKBP-12” or “FKBP 12” or “FKBP1A” may refer to the wildtype human nucleic acid. In embodiments, the FKBP-12 is a mutant FKBP-12. In embodiments, the mutant FKBP-12 is associated with a disease that is not associated with wildtype FKBP-12. In embodiments, the FKBP-12 may include at least one amino acid mutation (e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, or 30 mutations, or any range derivable therein) compared to wildtype FKBP-12. In embodiments, the FKBP-12 has the protein sequence corresponding to reference number GI:206725550. In embodiments, the FKBP-12 has the protein sequence corresponding to RefSeq NP_000792.1 (SEQ ID NO:3).

[00219] The term “4E-BP1” or “4EBP1” or “EIF4EBP1” refers to the protein “Eukaryotic translation initiation factor 4E-binding protein 1.” In embodiments, “4E-BP1” or “4EBP1” or “EIF4EBP 1” may refer to the human protein. Included in the term “4E-BP 1” or “4EBP 1” or “EIF4EBP1” are the wildtype and mutant forms of the protein. In embodiments, “4E-BP1”

or “4EBP1” or “EIF4EBP1” may refer to the protein associated with Entrez Gene 1978, OMIM 602223, UniProt Q13541, and/or RefSeq (protein) NP_004086 (SEQ ID NO:4). In embodiments, the reference numbers immediately above may refer to the protein, and associated nucleic acids, known as of the date of filing of this application. In embodiments, “4E-BP 1” or “4EBP1” or “EIF4EBP1” may refer to the wildtype human protein. In embodiments, “4E-BP1” or “4EBP1” or “EIF4EBP1” may refer to the wildtype human nucleic acid. In embodiments, the 4EBP1 is a mutant 4EBP1. In embodiments, the mutant 4EBP1 is associated with a disease that is not associated with wildtype 4EBP1. In embodiments, the 4EBP1 may include at least one amino acid mutation (e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, or 30 mutations, or any range derivable therein) compared to wildtype 4EBP1. In embodiments, the 4EBP1 has the protein sequence corresponding to reference number GL4758258. In embodiments, the 4EBP1 has the protein sequence corresponding to RefSeq NP_004086.1 (SEQ ID NO:4).

[00220] The term “Akt” refers to the serine/threonine specific protein kinase involved in cellular processes such as glucose metabolism, apoptosis, proliferation, and other functions, also known as “protein kinase B” (PKB) or “Akt1.” In embodiments, “Akt” or “AM” or “PKB” may refer to the human protein. Included in the term “Akt” or “Akt1” or “PKB” are the wildtype and mutant forms of the protein. In embodiments, “Akt” or “Akt1” or “PKB” may refer to the protein associated with Entrez Gene 207, OMIM 164730, UniProt P31749, and/or RefSeq (protein) NP_005154 (SEQ ID NO:5). In embodiments, the reference numbers immediately above may refer to the protein, and associated nucleic acids, known as of the date of filing of this application. In embodiments, “Akt” or “Akt1” or “PKB” may refer to the wildtype human protein. In embodiments, “Akt” or “Akt1” or “PKB” may refer to the wildtype human nucleic acid. In embodiments, the Akt is a mutant Akt. In embodiments, the mutant Akt is associated with a disease that is not associated with wildtype Akt. In embodiments, the Akt may include at least one amino acid mutation (e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, or 30 mutations, or any range derivable therein) compared to wildtype Akt. In embodiments, the Akt has the protein sequence corresponding to reference number GI: 62241011. In embodiments, the Akt has the protein sequence corresponding to RefSeq NP_005154.2 (SEQ ID NO:5).

[00221] The present disclosure provides a method of treating a disease or disorder mediated by mTOR comprising administering to the subject suffering from or susceptible to developing a disease or disorder mediated by mTOR a therapeutically effective amount of one or more disclosed compositions or compounds. The present disclosure provides a method of preventing a disease or disorder mediated by mTOR comprising administering to the subject suffering from or susceptible to developing a disease or disorder mediated by mTOR a therapeutically effective amount of one or more disclosed compositions or compounds. The present disclosure provides a method of reducing the risk of a disease or disorder mediated by mTOR comprising administering to the subject suffering from or susceptible to developing a disease or disorder mediated by mTOR a therapeutically effective amount of one or more disclosed compositions or compounds.

[00222] In some embodiments, the disease is cancer or an immune-mediated disease. In some embodiments, the cancer is selected from brain and neurovascular tumors, head and neck cancers, breast cancer, lung cancer, mesothelioma, lymphoid cancer, stomach cancer, kidney cancer, renal carcinoma, liver cancer, ovarian cancer, ovary endometriosis, testicular cancer, gastrointestinal cancer, prostate cancer, glioblastoma, skin cancer, melanoma, neuro cancers, spleen cancers, pancreatic cancers, blood proliferative disorders, lymphoma, leukemia, endometrial cancer, cervical cancer, vulva cancer, prostate cancer, penile cancer, bone cancers, muscle cancers, soft tissue cancers, intestinal or rectal cancer, anal cancer, bladder cancer, bile duct cancer, ocular cancer, gastrointestinal stromal tumors, and neuro-endocrine tumors. In some embodiments, the disorder is liver cirrhosis. In some embodiments, the immune-mediated disease is selected from resistance by transplantation of heart, kidney, liver, medulla ossium, skin, cornea, lung, pancreas, intestinum tenue, limb, muscle, nerves, duodenum, small-bowel, or pancreatic-islet-cell; graft-versus-host diseases brought about by medulla ossium transplantation; rheumatoid arthritis, systemic lupus erythematosus, Hashimoto's thyroiditis, multiple sclerosis, myasthenia gravis, type I diabetes, uveitis, allergic encephalomyelitis, and glomerulonephritis. In certain embodiments, the disease is tuberous sclerosis complex (TSC). In certain embodiments, the disease is pancreatic neuroendocrine tumor (PNET), mantle cell lymphoma (MCL), colorectal cancer or bowel cancer (CRC), uterine cancer, ovarian cancer, bladder cancer, genitourinary tract cancer, or renal cell carcinoma (RCC).

[00223] The present disclosure provides a method of treating cancer comprising administering to the subject a therapeutically effective amount of one or more disclosed

compositions or compounds. In some embodiments, the cancer is selected from brain and neurovascular tumors, head and neck cancers, breast cancer, lung cancer, mesothelioma, lymphoid cancer, stomach cancer, kidney cancer, renal carcinoma, liver cancer, ovarian cancer, ovary endometriosis, testicular cancer, gastrointestinal cancer, prostate cancer, glioblastoma, skin cancer, melanoma, neuro cancers, spleen cancers, pancreatic cancers, blood proliferative disorders, lymphoma, leukemia, endometrial cancer, cervical cancer, vulva cancer, prostate cancer, penile cancer, bone cancers, muscle cancers, soft tissue cancers, intestinal or rectal cancer, anal cancer, bladder cancer, bile duct cancer, ocular cancer, gastrointestinal stromal tumors, and neuro-endocrine tumors. In some embodiments, the disorder is liver cirrhosis. In certain embodiments, the disease is tuberous sclerosis complex (TSC). In certain embodiments, the disease is pancreatic neuroendocrine tumor (PNET), mantle cell lymphoma (MCL), colorectal cancer or bowel cancer (CRC), uterine cancer, ovarian cancer, bladder cancer, genitourinary tract cancer, or renal cell carcinoma (RCC).

[00224] In certain embodiments, cancer includes human cancers and carcinomas, sarcomas, adenocarcinomas, lymphomas, leukemias, etc., including solid and lymphoid cancers, kidney, breast, lung, bladder, colon, ovarian, prostate, pancreas, stomach, brain, head and neck, skin, uterine, testicular, glioma, esophagus, and liver cancer, including hepatocarcinoma, lymphoma, including B-acute lymphoblastic lymphoma, non-Hodgkin's lymphomas (e.g., Burkitt's, Small Cell, and Large Cell lymphomas), Hodgkin's lymphoma, leukemia (including AML, ALL, and CML), or multiple myeloma. In certain embodiments, the disease is multiple myeloma. In certain embodiments, the disease is breast cancer. In certain embodiments, the disease is triple negative breast cancer.

[00225] In certain embodiments, cancer includes cancer, neoplasm or malignant tumors found in mammals (e.g. humans), including leukemia, carcinomas and sarcomas. Exemplary cancers that may be treated with a compound or method provided herein include cancer of the prostate, thyroid, endocrine system, brain, breast, cervix, colon, head and neck, liver, kidney, lung, non-small cell lung, melanoma, mesothelioma, ovary, sarcoma, stomach, uterus, Medulloblastoma, colorectal cancer, pancreatic cancer. Additional examples may include, Hodgkin's Disease, Non-Hodgkin's Lymphoma, multiple myeloma, neuroblastoma, glioma, glioblastoma multiforme, ovarian cancer, rhabdomyosarcoma, primary thrombocytosis, primary macroglobulinemia, primary brain tumors, malignant pancreatic insulanoma, malignant carcinoid, urinary bladder cancer, premalignant skin lesions, testicular cancer,

lymphomas, thyroid cancer, neuroblastoma, esophageal cancer, genitourinary tract cancer, malignant hypercalcemia, endometrial cancer, adrenal cortical cancer, neoplasms of the endocrine or exocrine pancreas, medullary thyroid cancer, medullary thyroid carcinoma, melanoma, colorectal cancer, papillary thyroid cancer, hepatocellular carcinoma, or prostate cancer.

[00226] In certain embodiments, the disease is leukemia. The term “leukemia” refers broadly to progressive, malignant diseases of the blood-forming organs and is generally characterized by a distorted proliferation and development of leukocytes and their precursors in the blood and bone marrow. Leukemia is generally clinically classified on the basis of (1) the duration and character of the disease-acute or chronic; (2) the type of cell involved; myeloid (myelogenous), lymphoid (lymphogenous), or monocytic; and (3) the increase or non-increase in the number of aberrant cells in the blood-leukemic or aleukemic (subleukemic). Exemplary leukemias that may be treated with a compound or method provided herein include, for example, acute nonlymphocytic leukemia, chronic lymphocytic leukemia, acute granulocytic leukemia, chronic granulocytic leukemia, acute promyelocytic leukemia, adult T-cell leukemia, aleukemic leukemia, a leukocythemetic leukemia, basophylic leukemia, blast cell leukemia, bovine leukemia, chronic myelocytic leukemia, leukemia cutis, embryonal leukemia, eosinophilic leukemia, Gross' leukemia, hairy-cell leukemia, hemoblastic leukemia, hemocytoblastic leukemia, histiocytic leukemia, stem cell leukemia, acute monocytic leukemia, leukopenic leukemia, lymphatic leukemia, lymphoblastic leukemia, lymphocytic leukemia, lymphogenous leukemia, lymphoid leukemia, lymphosarcoma cell leukemia, mast cell leukemia, megakaryocyte leukemia, micromyeloblastic leukemia, monocytic leukemia, myeloblasts leukemia, myelocytic leukemia, myeloid granulocytic leukemia, myelomonocytic leukemia, Naegeli leukemia, plasma cell leukemia, multiple myeloma, plasmacytic leukemia, promyelocytic leukemia, Rieder cell leukemia, Schilling's leukemia, stem cell leukemia, subleukemic leukemia, or undifferentiated cell leukemia.

[00227] In certain embodiments, the disease is sarcoma. The term “sarcoma” generally refers to a tumor which is made up of a substance like the embryonic connective tissue and is generally composed of closely packed cells embedded in a fibrillar or homogeneous substance. Sarcomas that may be treated with a compound or method provided herein include a chondrosarcoma, fibrosarcoma, lymphosarcoma, melanosarcoma, myxosarcoma, osteosarcoma, Abemethy's sarcoma, adipose sarcoma, liposarcoma, alveolar soft part

sarcoma, ameloblastic sarcoma, botryoid sarcoma, chloroma sarcoma, chorio carcinoma, embryonal sarcoma, Wilms' tumor sarcoma, endometrial sarcoma, stromal sarcoma, Ewing's sarcoma, fascial sarcoma, fibroblastic sarcoma, giant cell sarcoma, granulocytic sarcoma, Hodgkin's sarcoma, idiopathic multiple pigmented hemorrhagic sarcoma, immunoblastic sarcoma of B cells, lymphoma, immunoblastic sarcoma of T-cells, Jensen's sarcoma, Kaposi's sarcoma, Kupffer cell sarcoma, angiosarcoma, leukosarcoma, malignant mesenchymoma sarcoma, parosteal sarcoma, reticulocytic sarcoma, Rous sarcoma, serocystic sarcoma, synovial sarcoma, or telangiectaltic sarcoma.

[00228] In certain embodiments, the disease is melanoma. The term “melanoma” is taken to mean a tumor arising from the melanocyte system of the skin and other organs.

Melanomas that may be treated with a compound or method provided herein include, for example, acral-lentiginous melanoma, amelanotic melanoma, benign juvenile melanoma, Cloudman's melanoma, S91 melanoma, Harding-Passey melanoma, juvenile melanoma, lentigo maligna melanoma, malignant melanoma, nodular melanoma, subungal melanoma, or superficial spreading melanoma.

[00229] In certain embodiments, the disease is carcinoma. The term “carcinoma” refers to a malignant new growth made up of epithelial cells tending to infiltrate the surrounding tissues and give rise to metastases. Exemplary carcinomas that may be treated with a compound or method provided herein include, for example, medullary thyroid carcinoma, familial medullary thyroid carcinoma, acinar carcinoma, acinous carcinoma, adenocystic carcinoma, adenoid cystic carcinoma, carcinoma adenomatosum, carcinoma of adrenal cortex, alveolar carcinoma, alveolar cell carcinoma, basal cell carcinoma, carcinoma basocellulare, basaloid carcinoma, basosquamous cell carcinoma, bronchioalveolar carcinoma, bronchiolar carcinoma, bronchogenic carcinoma, cerebriiform carcinoma, cholangiocellular carcinoma, chorionic carcinoma, colloid carcinoma, comedo carcinoma, corpus carcinoma, cribriform carcinoma, carcinoma en cuirasse, carcinoma cutaneum, cylindrical carcinoma, cylindrical cell carcinoma, duct carcinoma, carcinoma durum, embryonal carcinoma, encephaloid carcinoma, epiermoid carcinoma, carcinoma epitheliale adenoides, exophytic carcinoma, carcinoma ex ulcere, carcinoma fibrosum, gelatiniformi carcinoma, gelatinous carcinoma, giant cell carcinoma, carcinoma gigantocellulare, glandular carcinoma, granulosa cell carcinoma, hair-matrix carcinoma, hematoid carcinoma, hepatocellular carcinoma, Hurthle cell carcinoma, hyaline carcinoma, hypernephroid carcinoma, infantile embryonal carcinoma, carcinoma in situ, intraepidermal carcinoma,

intraepithelial carcinoma, Krompecher's carcinoma, Kulchitzky-cell carcinoma, large-cell carcinoma, lenticular carcinoma, carcinoma lenticulare, lipomatous carcinoma, lymphoepithelial carcinoma, carcinoma medullare, medullary carcinoma, melanotic carcinoma, carcinoma molle, mucinous carcinoma, carcinoma muciparum, carcinoma mucocellulare, mucoepidermoid carcinoma, carcinoma mucosum, mucous carcinoma, carcinoma myxomatodes, nasopharyngeal carcinoma, oat cell carcinoma, carcinoma ossificans, osteoid carcinoma, papillary carcinoma, periportal carcinoma, preinvasive carcinoma, prickle cell carcinoma, pultaceous carcinoma, renal cell carcinoma of kidney, reserve cell carcinoma, carcinoma sarcomatodes, schneiderian carcinoma, scirrhous carcinoma, carcinoma scroti, signet-ring cell carcinoma, carcinoma simplex, small-cell carcinoma, solanoid carcinoma, spheroidal cell carcinoma, spindle cell carcinoma, carcinoma spongiosum, squamous carcinoma, squamous cell carcinoma, string carcinoma, carcinoma telangiectaticum, carcinoma telangiectodes, transitional cell carcinoma, carcinoma tuberosum, tuberos carcinoma, verrucous carcinoma, or carcinoma villosum.

[00230] The present disclosure provides a method of treating an immune-mediated disease comprising administering to the subject a therapeutically effective amount of one or more disclosed compositions or compounds. In some embodiments, the immune-mediated disease is selected from resistance by transplantation of heart, kidney, liver, medulla ossium, skin, cornea, lung, pancreas, intestinum tenue, limb, muscle, nerves, duodenum, small-bowel, or pancreatic-islet-cell; graft-versus-host diseases brought about by medulla ossium transplantation; rheumatoid arthritis, systemic lupus erythematosus, Hashimoto's thyroiditis, multiple sclerosis, myasthenia gravis, type I diabetes, uveitis, allergic encephalomyelitis, and glomerulonephritis.

[00231] In certain embodiments, the disease is autoimmune disease. As used herein, the term “autoimmune disease” refers to a disease or condition in which a subject's immune system has an aberrant immune response against a substance that does not normally elicit an immune response in a healthy subject. Examples of autoimmune diseases that may be treated with a compound, pharmaceutical composition, or method described herein include Acute Disseminated Encephalomyelitis (ADEM), Acute necrotizing hemorrhagic leukoencephalitis, Addison's disease, Agammaglobulinemia, Alopecia areata, Amyloidosis, Ankylosing spondylitis, Anti-GBM/Anti-TBM nephritis, Antiphospholipid syndrome (APS), Autoimmune angioedema, Autoimmune aplastic anemia, Autoimmune dysautonomia, Autoimmune hepatitis, Autoimmune hyperlipidemia, Autoimmune immunodeficiency,

Autoimmune inner ear disease (AIED), Autoimmune myocarditis, Autoimmune oophoritis, Autoimmune pancreatitis, Autoimmune retinopathy, Autoimmune thrombocytopenic purpura (ATP), Autoimmune thyroid disease, Autoimmune urticaria, Axonal or neuronal neuropathies, Balo disease, Behcet's disease, Bullous pemphigoid, Cardiomyopathy, Castleman disease, Celiac disease, Chagas disease, Chronic fatigue syndrome, Chronic inflammatory demyelinating polyneuropathy (CIDP), Chronic recurrent multifocal osteomyelitis (CRMO), Churg-Strauss syndrome, Cicatricial pemphigoid/benign mucosal pemphigoid, Crohn's disease, Cogans syndrome, Cold agglutinin disease, Congenital heart block, Coxsackie myocarditis, CREST disease, Essential mixed cryoglobulinemia, Demyelinating neuropathies, Dermatitis herpetiformis, Dermatomyositis, Devic's disease (neuromyelitis optica), Discoid lupus, Dressier's syndrome, Endometriosis, Eosinophilic esophagitis, Eosinophilic fasciitis, Erythema nodosum, Experimental allergic encephalomyelitis, Evans syndrome, Fibromyalgia, Fibrosing alveolitis, Giant cell arteritis (temporal arteritis), Giant cell myocarditis, Glomerulonephritis, Goodpasture's syndrome, Granulomatosis with Polyangiitis (GPA) (formerly called Wegener's Granulomatosis), Graves' disease, Guillain-Barre syndrome, Hashimoto's encephalitis, Hashimoto's thyroiditis, Hemolytic anemia, Henoch-Schonlein purpura, Herpes gestationis, Hypogammaglobulinemia, Idiopathic thrombocytopenic purpura (ITP), IgA nephropathy, IgG4-related sclerosing disease, Immunoregulatory lipoproteins, Inclusion body myositis, Interstitial cystitis, Juvenile arthritis, Juvenile diabetes (Type 1 diabetes), Juvenile myositis, Kawasaki syndrome, Lambert-Eaton syndrome, Leukocytoclastic vasculitis, Lichen planus, Lichen sclerosus, Ligneous conjunctivitis, Linear IgA disease (LAD), Lupus (SLE), Lyme disease, chronic, Meniere's disease, Microscopic polyangiitis, Mixed connective tissue disease (MCTD), Mooren's ulcer, Mucha-Habermann disease, Multiple sclerosis, Myasthenia gravis, Myositis, Narcolepsy, Neuromyelitis optica (Devic's), Neutropenia, Ocular cicatricial pemphigoid, Optic neuritis, Palindromic rheumatism, PANDAS (Pediatric Autoimmune Neuropsychiatric Disorders Associated with Streptococcus), Paraneoplastic cerebellar degeneration, Paroxysmal nocturnal hemoglobinuria (PNH), Parry Romberg syndrome, Parsonnage-Turner syndrome, Pars planitis (peripheral uveitis), Pemphigus, Peripheral neuropathy, Perivenous encephalomyelitis, Pernicious anemia, POEMS syndrome, Polyarteritis nodosa, Type I, II, & III autoimmune polyglandular syndromes, Polymyalgia rheumatica, Polymyositis, Postmyocardial infarction syndrome, Postpericardiotomy syndrome, Progesterone dermatitis, Primary biliary cirrhosis, Primary sclerosing cholangitis, Psoriasis, Psoriatic arthritis, Idiopathic pulmonary fibrosis, Pyoderma gangrenosum, Pure red

cell aplasia, Raynauds phenomenon, Reactive Arthritis, Reflex sympathetic dystrophy, Reiter's syndrome, Relapsing polychondritis, Restless legs syndrome, Retroperitoneal fibrosis, Rheumatic fever, Rheumatoid arthritis, Sarcoidosis, Schmidt syndrome, Scleritis, Scleroderma, Sjogren's syndrome, Sperm & testicular autoimmunity, Stiff person syndrome, Subacute bacterial endocarditis (SBE), Susac's syndrome, Sympathetic ophthalmia, Takayasu's arteritis, Temporal arteritis/Giant cell arteritis, Thrombocytopenic purpura (TTP), Tolosa-Hunt syndrome, Transverse myelitis, Type 1 diabetes, Ulcerative colitis, Undifferentiated connective tissue disease (UCTD), Uveitis, Vasculitis, Vesiculobullous dermatosis, Vitiligo, or Wegener's granulomatosis (i.e., Granulomatosis with Polyangiitis (GPA)).

[00232] The present disclosure provide a method of treating an age related condition comprising administering to the subject a therapeutically effective amount of one or more disclosed compositions or compounds. In certain embodiments, the age related condition is selected from sarcopenia, skin atrophy, muscle wasting, brain atrophy, atherosclerosis, arteriosclerosis, pulmonary emphysema, osteoporosis, osteoarthritis, high blood pressure, erectile dysfunction, dementia, Huntington's disease, Alzheimer's disease, cataracts, age-related macular degeneration, prostate cancer, stroke, diminished life expectancy, impaired kidney function, and age-related hearing loss, aging-related mobility disability (e.g., frailty), cognitive decline, age-related dementia, memory impairment, tendon stiffness, heart dysfunction such as cardiac hypertrophy and systolic and diastolic dysfunction, immunosenescence, cancer, obesity, and diabetes.

[00233] In certain embodiments, the disclosed compositions or compounds can be used with regard to immunosenescence. Immunosenescence may refer to a decrease in immune function resulting in impaired immune response, e.g., to cancer, vaccination, infectious pathogens, among others. It involves both the host's capacity to respond to infections and the development of long-term immune memory, especially by vaccination. This immune deficiency is ubiquitous and found in both long- and short-lived species as a function of their age relative to life expectancy rather than chronological time. It is considered a major contributory factor to the increased frequency of morbidity and mortality among the elderly. Immunosenescence is not a random deteriorative phenomenon, rather it appears to inversely repeat an evolutionary pattern and most of the parameters affected by immunosenescence appear to be under genetic control. Immunosenescence can also be sometimes envisaged as the result of the continuous challenge of the unavoidable exposure to a variety of antigens

such as viruses and bacteria. Immunosenescence is a multifactorial condition leading to many pathologically significant health problems, e.g., in the aged population. Age-dependent biological changes such as depletion of hematopoietic stem cells, an increase in PD1+ lymphocytes, a decline in the total number of phagocytes and NK cells and a decline in humoral immunity contribute to the onset of immunosenescence. In one aspect, immunosenescence can be measured in an individual by measuring telomere length in immune cells (See, e.g., U.S. Pat. No. 5,741,677). Immunosenescence can also be determined by documenting in an individual a lower than normal number of naive CD4 and/or CD8 T cells, T cell repertoire, the number of PD1-expressing T cells, e.g., a lower than normal number of PD-1 negative T cells, or response to vaccination in a subject greater than or equal to 65 years of age. In certain embodiments, mTOR selective modulation of certain T-cell populations may improve vaccine efficacy in the aging population and enhance effectiveness of cancer immunotherapy. The present disclosure provides a method of treating immunosenescence comprising administering to the subject a therapeutically effective amount of one or more disclosed compositions or compounds.

[00234] In certain embodiments, a disease that may be treated with a compound, pharmaceutical composition, or method described herein is organ or tissue transplant rejection (e.g. heart, lung, combined heart- lung, liver, kidney, pancreatic, skin or corneal transplants; graft-versus-host disease), restenosis, Hamartoma syndromes (e.g., tuberous sclerosis or Cowden Disease), Lymphangi leiomyomatosis, Retinitis pigmentosa, encephalomyelitis, insulin-dependent diabetes mellitus, lupus, dermatomyositis, arthritis, rheumatic diseases, Steroid-resistant acute Lymphoblastic Leukemia, fibrosis, scleroderma, pulmonary fibrosis, renal fibrosis, cystic fibrosis, Pulmonary hypertension, Multiple sclerosis, VHL syndrome, Carney complex, Familial adenomatous polyposis, Juvenile polyposis syndrome, Birt-Hogg-Duke syndrome, Familial hypertrophic cardiomyopathy, Wolf-Parkinson-White syndrome, Parkinson's disease, Huntingtin's disease, Alzheimer's disease, dementias caused by tau mutations, spinocerebellar ataxia type 3, motor neuron disease caused by SOD1 mutations, neuronal ceroid lipofuscinoses/Batten disease (pediatric neurodegeneration), wet macular degeneration, dry macular degeneration, muscle wasting (atrophy, cachexia), myopathies (e.g., Danon's disease), bacterial infection, viral infection, M. tuberculosis, group A streptococcus, HSV type I, HIV infection, Neurofibromatosis (e.g., Neurofibromatosis type 1), or Peutz-Jeghers syndrome.

[00235] In certain embodiments, the disease is neurodegenerative disease. As used herein, the term “neurodegenerative disease” refers to a disease or condition in which the function of a subject’s nervous system becomes impaired. Examples of neurodegenerative diseases that may be treated with a compound, pharmaceutical composition, or method described herein include Alexander's disease, Alper's disease, Alzheimer's disease, Amyotrophic lateral sclerosis, Ataxia telangiectasia, Batten disease (also known as Spielmeyer-Vogt-Sjogren-Batten disease), Bovine spongiform encephalopathy (BSE), Canavan disease, Cockayne syndrome, Corticobasal degeneration, Creutzfeldt- Jakob disease, frontotemporal dementia, Gerstmann-Straussler-Scheinker syndrome, Huntington's disease, HIV-associated dementia, Kennedy's disease, Krabbe's disease, kuru, Lewy body dementia, Machado- Joseph disease (Spinocerebellar ataxia type 3), Multiple sclerosis, Multiple System Atrophy, Narcolepsy, Neuroborreliosis, Parkinson's disease, Pelizaeus-Merzbacher Disease, Pick's disease, Primary lateral sclerosis, Prion diseases, Refsum's disease, Sandhoff's disease, Schilder's disease, Subacute combined degeneration of spinal cord secondary to Pernicious Anaemia, Schizophrenia, Spinocerebellar ataxia (multiple types with varying characteristics), Spinal muscular atrophy, Steele-Richardson-Olszewski disease, or Tabes dorsalis.

[00236] In certain embodiments, the disease is metabolic disease. As used herein, the term “metabolic disease” refers to a disease or condition in which a subject's metabolism or metabolic system (e.g., function of storing or utilizing energy) becomes impaired. Examples of metabolic diseases that may be treated with a compound, pharmaceutical composition, or method described herein include diabetes (e.g., type I or type II), obesity, metabolic syndrome, or a mitochondrial disease (e.g., dysfunction of mitochondria or aberrant mitochondrial function).

[00237] In certain embodiments, the disease is fungal disease. As used herein, the term “fungal disease” refers to a disease or condition associated with a fungus infection of the subject. Examples of fungal diseases that may be treated with a compound, pharmaceutical composition, or method described herein include infection with *Mucor circinelloides*, zygomycetes, *Cryptococcus neoformans*, *Candida albicans*, yeast, and *Saccharomyces cerevisiae* among others.

[00238] In certain embodiments, the disease is inflammatory disease. As used herein, the term “inflammatory disease” refers to a disease or condition characterized by aberrant inflammation (e.g. an increased level of inflammation compared to a control such as a healthy person not suffering from a disease). Examples of inflammatory diseases include traumatic

brain injury, arthritis, rheumatoid arthritis, psoriatic arthritis, juvenile idiopathic arthritis, multiple sclerosis, systemic lupus erythematosus (SLE), myasthenia gravis, juvenile onset diabetes, diabetes mellitus type 1, Guillain-Barre syndrome, Hashimoto's encephalitis, Hashimoto's thyroiditis, ankylosing spondylitis, psoriasis, Sjogren's syndrome, vasculitis, glomerulonephritis, auto-immune thyroiditis, Behcet's disease, Crohn's disease, ulcerative colitis, bullous pemphigoid, sarcoidosis, ichthyosis, Graves ophthalmopathy, inflammatory bowel disease, Addison's disease, Vitiligo, asthma, allergic asthma, acne vulgaris, celiac disease, chronic prostatitis, inflammatory bowel disease, pelvic inflammatory disease, reperfusion injury, sarcoidosis, transplant rejection, interstitial cystitis, atherosclerosis, and atopic dermatitis.

[00239] In certain embodiments, the disease is cardiovascular disease. As used herein, the term “cardiovascular disease” refers to a disease or condition in which the function of a subject's cardiovascular system becomes impaired. Examples of cardiovascular diseases that may be treated with a compound, pharmaceutical composition, or method described herein include congestive heart failure; arrhythmogenic syndromes (e.g., paroxysmal tachycardia, delayed after depolarizations, ventricular tachycardia, sudden tachycardia, exercise-induced arrhythmias, long QT syndromes, or bidirectional tachycardia); thromboembolic disorders (e.g., arterial cardiovascular thromboembolic disorders, venous cardiovascular thromboembolic disorders, or thromboembolic disorders in the chambers of the heart); atherosclerosis; restenosis; peripheral arterial disease; coronary bypass grafting surgery; carotid artery disease; arteritis; myocarditis; cardiovascular inflammation; vascular inflammation; coronary heart disease (CHD); unstable angina (UA); unstable refractory angina; stable angina (SA); chronic stable angina; acute coronary syndrome (ACS); myocardial infarction (first or recurrent); acute myocardial infarction (AMI); myocardial infarction; non-Q wave myocardial infarction; non-STE myocardial infarction; coronary artery disease; ischemic heart disease; cardiac ischemia; ischemia; ischemic sudden death; transient ischemic attack; stroke; peripheral occlusive arterial disease; venous thrombosis; deep vein thrombosis; thrombophlebitis; arterial embolism; coronary arterial thrombosis; cerebral arterial thrombosis, cerebral embolism; kidney embolism; pulmonary embolism; thrombosis (e.g., associated with prosthetic valves or other implants, indwelling catheters, stents, cardiopulmonary bypass, hemodialysis); thrombosis (e.g., associated with atherosclerosis, surgery, prolonged immobilization, arterial fibrillation, congenital

thrombophilia, cancer, diabetes, hormones, or pregnancy); or cardiac arrhythmias (e.g., supraventricular arrhythmias, atrial arrhythmias, atrial flutter, or atrial fibrillation).

[00240] In an aspect is provided a method of treating a disease associated with an aberrant level of mTOR activity in a subject in need of such treatment. The disease may be caused by an upregulation of mTOR. The method may include administering to the subject one or more compositions or compounds described herein. The method may include administering to the subject a therapeutically effective amount of one or more compositions or compounds described herein (e.g., an mTOR modulator (e.g., inhibitor) as described above).

[00241] In an aspect is provided one or more compositions or compounds as described herein for use as a medicament. In embodiments, the medicament is useful for treating a disease caused by an upregulation of mTOR. The use may include administering to the subject one or more compositions or compounds described herein. The use may include administering to the subject a therapeutically effective amount of one or more compositions or compounds described herein (e.g., an mTOR modulator (e.g., inhibitor) as described above).

[00242] In an aspect is provided one or more compositions or compounds as described herein for use in the treatment of a disease caused by aberrant levels of mTORC1 activity in a subject in need of such treatment. The disease may be caused by an upregulation of mTORC1. The use may include administering to the subject one or more compositions or compounds described herein. The use may include administering to the subject a therapeutically effective amount of one or more compositions or compounds described herein (e.g., an mTORC1 modulator (e.g., inhibitor) as described above).

[00243] Upregulation of mTOR can result in an increased amount of mTOR activity compared to normal levels of mTOR activity in a particular subject or a population of healthy subjects. The increased amount of mTOR activity may result in, for example, excessive amounts of cell proliferation thereby causing the disease state.

[00244] The subject of treatment for the disease is typically a mammal. The mammal treated with the compound (e.g., compound described herein, mTOR modulator (e.g., inhibitor)) may be a human, nonhuman primate, and/or non-human mammal (e.g., rodent, canine).

[00245] In another aspect is provided a method of treating an mTOR activity-associated disease in a subject in need of such treatment, the method including administering one or

more compositions or compounds as described herein, including embodiments (e.g., a claim, embodiment, example, table, figure, or claim) to the subject.

[00246] In another aspect is provided one or more compositions or compounds as described herein for use as a medicament. In embodiments, the medicament may be useful for treating an mTORC1 activity-associated disease in a subject in need of such treatment. In embodiments, the use may include administering one or more compositions or compounds as described herein, including embodiments (e.g., an aspect, embodiment, example, table, figure, or claim) to the subject.

[00247] In another aspect is provided one or more compositions or compounds for use in the treatment of an mTOR activity-associated disease in a subject in need of such treatment. In embodiments, the use may include administering one or more compositions or compounds as described herein, including embodiments (e.g., an aspect, embodiment, example, table, figure, or claim) to the subject.

[00248] In embodiments, the mTOR activity-associated disease or disease associated with aberrant levels of mTOR activity is cancer. In embodiments, the mTOR activity-associated disease or disease associated with aberrant levels of mTOR activity is an autoimmune disease. In embodiments, the mTOR activity-associated disease or disease associated with aberrant levels of mTOR activity is an inflammatory disease. In embodiments, the mTOR activity-associated disease or disease associated with aberrant levels of mTOR activity is a neurodegenerative disease. In embodiments, the mTOR activity-associated disease or disease associated with aberrant levels of mTOR activity is a metabolic disease. In embodiments, the mTOR activity-associated disease or disease associated with aberrant levels of mTOR activity is transplant rejection. In embodiments, the mTOR activity-associated disease or disease associated with aberrant levels of mTOR activity is fungal infection. In embodiments, the mTOR activity-associated disease or disease associated with aberrant levels of mTOR activity is a cardiovascular disease.

[00249] In embodiments, the mTOR activity-associated disease or disease associated with aberrant levels of mTOR activity is aging. In embodiments, the mTOR activity-associated disease or disease associated with aberrant levels of mTOR activity is dying of an age-related disease. In embodiments, the mTOR activity-associated disease or disease associated with aberrant levels of mTOR activity is an age-related condition. In certain embodiments, the age related condition is selected from the group consisting of sarcopenia, skin atrophy, muscle

wasting, brain atrophy, atherosclerosis, arteriosclerosis, pulmonary emphysema, osteoporosis, osteoarthritis, high blood pressure, erectile dysfunction, dementia, Huntington's disease, Alzheimer's disease, cataracts, age-related macular degeneration, prostate cancer, stroke, diminished life expectancy, impaired kidney function, and age-related hearing loss, aging-related mobility disability (e.g., frailty), cognitive decline, age-related dementia, memory impairment, tendon stiffness, heart dysfunction such as cardiac hypertrophy and systolic and diastolic dysfunction, immunosenescence, cancer, obesity, and diabetes. In certain embodiments, mTOR selective modulation of certain T-cell populations may improve vaccine efficacy in the aging population and enhance effectiveness of cancer immunotherapy. The present disclosure provides a method of treating immunosenescence comprising administering to the subject a therapeutically effective amount of one or more disclosed compounds.

[00250] In embodiments, the mTOR activity-associated disease or disease associated with aberrant levels of mTOR activity is cancer (e.g., carcinomas, sarcomas, adenocarcinomas, lymphomas, leukemias, solid cancers, lymphoid cancers; cancer of the kidney, breast, lung, bladder, colon, gastrointestinal, ovarian, prostate, pancreas, stomach, brain, head and neck, skin, uterine, esophagus, liver; testicular cancer, glioma, hepatocarcinoma, lymphoma, including B-acute lymphoblastic lymphoma, non-Hodgkin's lymphomas (e.g., Burkitt's, Small Cell, and Large Cell lymphomas), Hodgkin's lymphoma, leukemia (including AML, ALL, and CML), multiple myeloma, and breast cancer (e.g., triple negative breast cancer)).

[00251] In embodiments, the mTOR activity-associated disease or disease associated with aberrant levels of mTOR activity is Acute Disseminated Encephalomyelitis (ADEM), Acute necrotizing hemorrhagic leukoencephalitis, Addison's disease, Agammaglobulinemia, Alopecia areata, Amyloidosis, Ankylosing spondylitis, Anti-GBM/Anti-TBM nephritis, Antiphospholipid syndrome (APS), Autoimmune angioedema, Autoimmune aplastic anemia, Autoimmune dysautonomia, Autoimmune hepatitis, Autoimmune hyperlipidemia, Autoimmune immunodeficiency, Autoimmune inner ear disease (AIED), Autoimmune myocarditis, Autoimmune oophoritis, Autoimmune pancreatitis, Autoimmune retinopathy, Autoimmune thrombocytopenic purpura (ATP), Autoimmune thyroid disease, Autoimmune urticaria, Axonal or neuronal neuropathies, Balo disease, Behcet's disease, Bullous pemphigoid, Cardiomyopathy, Castleman disease, Celiac disease, Chagas disease, Chronic fatigue syndrome, Chronic inflammatory demyelinating polyneuropathy (CIDP), Chronic recurrent multifocal osteomyelitis (CRMO), Churg-Strauss syndrome, Cicatricial

pemphigoid/benign mucosal pemphigoid, Crohn's disease, Cogans syndrome, Cold agglutinin disease, Congenital heart block, Coxsackie myocarditis, CREST disease, Essential mixed cryoglobulinemia, Demyelinating neuropathies, Dermatitis herpetiformis, Dermatomyositis, Devic's disease (neuromyelitis optica), Discoid lupus, Dressier's syndrome, Endometriosis, Eosinophilic esophagitis, Eosinophilic fasciitis, Erythema nodosum, Experimental allergic encephalomyelitis, Evans syndrome, Fibromyalgia, Fibrosing alveolitis, Giant cell arteritis (temporal arteritis), Giant cell myocarditis, Glomerulonephritis, Goodpasture's syndrome, Granulomatosis with Polyangiitis (GPA) (formerly called Wegener's Granulomatosis), Graves' disease, Guillain-Barre syndrome, Hashimoto's encephalitis, Hashimoto's thyroiditis, Hemolytic anemia, Henoch- Schonlein purpura, Herpes gestationis, Hypogammaglobulinemia, Idiopathic thrombocytopenic purpura (ITP), IgA nephropathy, IgG4-related sclerosing disease, Immunoregulatory lipoproteins, Inclusion body myositis, Interstitial cystitis, Juvenile arthritis, Juvenile diabetes (Type 1 diabetes), Juvenile myositis, Kawasaki syndrome, Lambert-Eaton syndrome, Leukocytoclastic vasculitis, Lichen planus, Lichen sclerosus, Ligneous conjunctivitis, Linear IgA disease (LAD), Lupus (SLE), Lyme disease, chronic, Meniere's disease, Microscopic polyangiitis, Mixed connective tissue disease (MCTD), Mooren's ulcer, Mucha-Habermann disease, Multiple sclerosis, Myasthenia gravis, Myositis, Narcolepsy, Neuromyelitis optica (Devic's), Neutropenia, Ocular cicatricial pemphigoid, Optic neuritis, Palindromic rheumatism, PANDAS (Pediatric Autoimmune Neuropsychiatry Disorders Associated with Streptococcus), Paraneoplastic cerebellar degeneration, Paroxysmal nocturnal hemoglobinuria (PNH), Parry Romberg syndrome, Parsonnage -Turner syndrome, Pars planitis (peripheral uveitis), Pemphigus, Peripheral neuropathy, Perivenous encephalomyelitis, Pernicious anemia, POEMS syndrome, Polyarteritis nodosa, Type I, II, & III autoimmune polyglandular syndromes, Polymyalgia rheumatica, Polymyositis, Postmyocardial infarction syndrome, Postpericardiotomy syndrome, Progesterone dermatitis, Primary biliary cirrhosis, Primary sclerosing cholangitis, Psoriasis, Psoriatic arthritis, Idiopathic pulmonary fibrosis, Pyoderma gangrenosum, Pure red cell aplasia, Raynauds phenomenon, Reactive Arthritis, Reflex sympathetic dystrophy, Reiter's syndrome, Relapsing polychondritis, Restless legs syndrome, Retroperitoneal fibrosis, Rheumatic fever, Rheumatoid arthritis, Sarcoidosis, Schmidt syndrome, Scleritis, Scleroderma, Sjogren's syndrome, Sperm & testicular autoimmunity, Stiff person syndrome, Subacute bacterial endocarditis (SBE), Susac's syndrome, Sympathetic ophthalmia, Takayasu's arteritis, Temporal arteritis/Giant cell arteritis, Thrombocytopenic purpura (TTP), Tolosa-Hunt syndrome, Transverse myelitis, Type 1 diabetes, Ulcerative colitis,

Undifferentiated connective tissue disease (UCTD), Uveitis, Vasculitis, Vesiculobullous dermatosis, Vitiligo, Wegener's granulomatosis (i.e., Granulomatosis with Polyangiitis (GPA), traumatic brain injury, arthritis, rheumatoid arthritis, psoriatic arthritis, juvenile idiopathic arthritis, multiple sclerosis, systemic lupus erythematosus (SLE), myasthenia gravis, juvenile onset diabetes, diabetes mellitus type 1, Guillain-Barre syndrome, Hashimoto's encephalitis, Hashimoto's thyroiditis, ankylosing spondylitis, psoriasis, Sjogren's syndrome, vasculitis, glomerulonephritis, auto-immune thyroiditis, Behcet's disease, Crohn's disease, ulcerative colitis, bullous pemphigoid, sarcoidosis, ichthyosis, Graves ophthalmopathy, inflammatory bowel disease, Addison's disease, Vitiligo, asthma, allergic asthma, acne vulgaris, celiac disease, chronic prostatitis, inflammatory bowel disease, pelvic inflammatory disease, reperfusion injury, sarcoidosis, transplant rejection, interstitial cystitis, atherosclerosis, atopic dermatitis, Alexander's disease, Alper's disease, Alzheimer's disease, Amyotrophic lateral sclerosis, Ataxia telangiectasia, Batten disease (also known as Spielmeyer-Vogt-Sjogren-Batten disease), Bovine spongiform encephalopathy (BSE), Canavan disease, Cockayne syndrome, Corticobasal degeneration, Creutzfeldt- Jakob disease, frontotemporal dementia, Gerstmann-Straussler-Scheinker syndrome, Huntington's disease, HTV-associated dementia, Kennedy's disease, Krabbe's disease, kuru, Lewy body dementia, Machado-Joseph disease (Spinocerebellar ataxia type 3), Multiple sclerosis, Multiple System Atrophy, Narcolepsy, Neuroborreliosis, Parkinson's disease, Pelizaeus-Merzbacher Disease, Pick's disease, Primary lateral sclerosis, Prion diseases, Refsum's disease, Sandhoff's disease, Schilder's disease, Subacute combined degeneration of spinal cord secondary to Pernicious Anaemia, Schizophrenia, Spinocerebellar ataxia (multiple types with varying characteristics), Spinal muscular atrophy, Steele-Richardson-Olszewski disease, Tabes dorsalis, diabetes (e.g., type I or type II), obesity, metabolic syndrome, a mitochondrial disease (e.g., dysfunction of mitochondria or aberrant mitochondrial function), fungal infection, transplant rejection, or a cardiovascular disease (e.g., congestive heart failure; arrhythmogenic syndromes (e.g., paroxysmal tachycardia, delayed after depolarizations, ventricular tachycardia, sudden tachycardia, exercise-induced arrhythmias, long QT syndromes, or bidirectional tachycardia); thromboembolic disorders (e.g., arterial cardiovascular thromboembolic disorders, venous cardiovascular thromboembolic disorders, or thromboembolic disorders in the chambers of the heart); atherosclerosis; restenosis; peripheral arterial disease; coronary bypass grafting surgery; carotid artery disease; arteritis; myocarditis; cardiovascular inflammation; vascular inflammation; coronary heart disease (CHD); unstable angina (UA); unstable refractory angina; stable angina (SA); chronic stable angina; acute coronary syndrome (ACS);

myocardial infarction (first or recurrent); acute myocardial infarction (AMI); myocardial infarction; non-Q wave myocardial infarction; non-STE myocardial infarction; coronary artery disease; ischemic heart disease; cardiac ischemia; ischemia; ischemic sudden death; transient ischemic attack; stroke; peripheral occlusive arterial disease; venous thrombosis; deep vein thrombosis; thrombophlebitis; arterial embolism; coronary arterial thrombosis; cerebral arterial thrombosis, cerebral embolism; kidney embolism; pulmonary embolism; thrombosis (e.g., associated with prosthetic valves or other implants, indwelling catheters, stents, cardiopulmonary bypass, hemodialysis); thrombosis (e.g., associated with atherosclerosis, surgery, prolonged immobilization, arterial fibrillation, congenital thrombophilia, cancer, diabetes, hormones, or pregnancy); or cardiac arrhythmias (e.g., supraventricular arrhythmias, atrial arrhythmias, atrial flutter, or atrial fibrillation).

[00252] In an aspect is provided a method of treating a disease including administering an effective amount of one or more compositions or compounds as described herein. In an aspect is provided one or more compositions or compounds as described herein for use as a medicament (e.g., for treatment of a disease). In an aspect is provided one or more compositions or compounds as described herein for use in the treatment of a disease (e.g., including administering an effective amount of one or more compositions or compounds as described herein). In embodiments, the disease is cancer. In embodiments, the disease is an autoimmune disease. In embodiments, the disease is an inflammatory disease. In embodiments, the disease is a neurodegenerative disease. In embodiments, the disease is a metabolic disease. In embodiments, the disease is fungal infection. In embodiments, the disease is transplant rejection. In embodiments, the disease is a cardiovascular disease.

[00253] In embodiments, the disease is cancer (e.g., carcinomas, sarcomas, adenocarcinomas, lymphomas, leukemias, solid cancers, lymphoid cancers; cancer of the kidney, breast, lung, bladder, colon, ovarian, prostate, pancreas, stomach, brain, head and neck, skin, uterine, esophagus, liver; testicular cancer, glioma, hepatocarcinoma, lymphoma, including B-acute lymphoblastic lymphoma, non-Hodgkin's lymphomas (e.g., Burkitt's, Small Cell, and Large Cell lymphomas), Hodgkin's lymphoma, leukemia (including AML, ALL, and CML), multiple myeloma, and breast cancer (e.g., triple negative breast cancer)).

[00254] In embodiments, the disease is Acute Disseminated Encephalomyelitis (ADEM), Acute necrotizing hemorrhagic leukoencephalitis, Addison's disease, Agammaglobulinemia, Alopecia areata, Amyloidosis, Ankylosing spondylitis, Anti-GBM/Anti-TBM nephritis, Antiphospholipid syndrome (APS), Autoimmune angioedema, Autoimmune aplastic anemia,

Autoimmune dysautonomia, Autoimmune hepatitis, Autoimmune hyperlipidemia, Autoimmune immunodeficiency, Autoimmune inner ear disease (AIED), Autoimmune myocarditis, Autoimmune oophoritis, Autoimmune pancreatitis, Autoimmune retinopathy, Autoimmune thrombocytopenic purpura (ATP), Autoimmune thyroid disease, Autoimmune urticaria, Axonal or neuronal neuropathies, Balo disease, Behcet's disease, Bullous pemphigoid, Cardiomyopathy, Castleman disease, Celiac disease, Chagas disease, Chronic fatigue syndrome, Chronic inflammatory demyelinating polyneuropathy (CIDP), Chronic recurrent multifocal osteomyelitis (CRMO), Churg-Strauss syndrome, Cicatricial pemphigoid/benign mucosal pemphigoid, Crohn's disease, Cogans syndrome, Cold agglutinin disease, Congenital heart block, Coxsackie myocarditis, CREST disease, Essential mixed cryoglobulinemia, Demyelinating neuropathies, Dermatitis herpetiformis, Dermatomyositis, Devic's disease (neuromyelitis optica), Discoid lupus, Dressler's syndrome, Endometriosis, Eosinophilic esophagitis, Eosinophilic fasciitis, Erythema nodosum, Experimental allergic encephalomyelitis, Evans syndrome, Fibromyalgia, Fibrosing alveolitis, Giant cell arteritis (temporal arteritis), Giant cell myocarditis, Glomerulonephritis, Goodpasture's syndrome, Granulomatosis with Polyangiitis (GPA) (formerly called Wegener's Granulomatosis), Graves' disease, Guillain-Barre syndrome, Hashimoto's encephalitis, Hashimoto's thyroiditis, Hemolytic anemia, Henoch-Schonlein purpura, Herpes gestationis, Hypogammaglobulinemia, Idiopathic thrombocytopenic purpura (ITP), IgA nephropathy, IgG4-related sclerosing disease, Immunoregulatory lipoproteins, Inclusion body myositis, Interstitial cystitis, Juvenile arthritis, Juvenile diabetes (Type 1 diabetes), Juvenile myositis, Kawasaki syndrome, Lambert-Eaton syndrome, Leukocytoclastic vasculitis, Lichen planus, Lichen sclerosus, Ligneous conjunctivitis, Linear IgA disease (LAD), Lupus (SLE), Lyme disease, Meniere's disease, Microscopic polyangiitis, Mixed connective tissue disease (MCTD), Mooren's ulcer, Mucha-Habermann disease, Multiple sclerosis, Myasthenia gravis, Myositis, Narcolepsy, Neuromyelitis optica (Devic's), Neutropenia, Ocular cicatricial pemphigoid, Optic neuritis, Palindromic rheumatism, PANDAS (Pediatric Autoimmune Neuropsychiatric Disorders Associated with Streptococcus), Paraneoplastic cerebellar degeneration, Paroxysmal nocturnal hemoglobinuria (PNH), Parry Romberg syndrome, Parsonnage-Turner syndrome, Pars planitis (peripheral uveitis), Pemphigus, Peripheral neuropathy, Perivenous encephalomyelitis, Pernicious anemia, POEMS syndrome, Polyarteritis nodosa, Type I, II, & III autoimmune polyglandular syndromes, Polymyalgia rheumatica, Polymyositis, Postmyocardial infarction syndrome, Postpericardiotomy syndrome, Progesterone dermatitis, Primary biliary cirrhosis, Primary sclerosing cholangitis,

Psoriasis, Psoriatic arthritis, Idiopathic pulmonary fibrosis, Pyoderma gangrenosum, Pure red cell aplasia, Raynauds phenomenon, Reactive Arthritis, Reflex sympathetic dystrophy, Reiter's syndrome, Relapsing polychondritis, Restless legs syndrome, Retroperitoneal fibrosis, Rheumatic fever, Rheumatoid arthritis, Sarcoidosis, Schmidt syndrome, Scleritis, Scleroderma, Sjogren's syndrome, Sperm & testicular autoimmunity, Stiff person syndrome, Subacute bacterial endocarditis (SBE), Susac's syndrome, Sympathetic ophthalmia, Takayasu's arteritis, Temporal arteritis/Giant cell arteritis, Thrombocytopenic purpura (TTP), Tolosa-Hunt syndrome, Transverse myelitis, Type 1 diabetes, Ulcerative colitis, Undifferentiated connective tissue disease (UCTD), Uveitis, Vasculitis, Vesiculobullous dermatosis, Vitiligo, Wegener's granulomatosis (i.e., Granulomatosis with Polyangiitis (GPA), traumatic brain injury, arthritis, rheumatoid arthritis, psoriatic arthritis, juvenile idiopathic arthritis, multiple sclerosis, systemic lupus erythematosus (SLE), myasthenia gravis, juvenile onset diabetes, diabetes mellitus type 1, Guillain-Barre syndrome, Hashimoto's encephalitis, Hashimoto's thyroiditis, ankylosing spondylitis, psoriasis, vasculitis, glomerulonephritis, auto-immune thyroiditis, Behcet's disease, Crohn's disease, ulcerative colitis, bullous pemphigoid, sarcoidosis, ichthyosis, Graves ophthalmopathy, inflammatory bowel disease, Addison's disease, Vitiligo, asthma, allergic asthma, acne vulgaris, celiac disease, chronic prostatitis, inflammatory bowel disease, pelvic inflammatory disease, reperfusion injury, sarcoidosis, transplant rejection, interstitial cystitis, atherosclerosis, atopic dermatitis, Alexander's disease, Alper's disease, Alzheimer's disease, Amyotrophic lateral sclerosis, Ataxia telangiectasia, Batten disease (also known as Spielmeyer-Vogt-Sjogren-Batten disease), Bovine spongiform encephalopathy (BSE), Canavan disease, Cockayne syndrome, Corticobasal degeneration, Creutzfeldt- Jakob disease, frontotemporal dementia, Gerstmann-Straussler-Scheinker syndrome, Huntington's disease, HTV-associated dementia, Kennedy's disease, Krabbe's disease, kuru, Lewy body dementia, Machado-Joseph disease (Spinocerebellar ataxia type 3), Multiple sclerosis, Multiple System Atrophy, Narcolepsy, Neuroborreliosis, Parkinson's disease, Pelizaeus-Merzbacher Disease, Pick's disease, Primary lateral sclerosis, Prion diseases, Refsum's disease, Sandhoff's disease, Schilder's disease, Subacute combined degeneration of spinal cord secondary to Pernicious Anaemia, Schizophrenia, Spinocerebellar ataxia (multiple types with varying characteristics), Spinal muscular atrophy, Steele-Richardson-Olszewski disease, Tabes dorsalis, diabetes (e.g., type I or type II), obesity, metabolic syndrome, a mitochondrial disease (e.g., dysfunction of mitochondria or aberrant mitochondrial function), fungal infection, transplant rejection, or a cardiovascular disease (e.g., congestive heart failure; arrhythmogenic syndromes (e.g.,

paroxysmal tachycardia, delayed after depolarizations, ventricular tachycardia, sudden tachycardia, exercise-induced arrhythmias, long QT syndromes, or bidirectional tachycardia); thromboembolic disorders (e.g., arterial cardiovascular thromboembolic disorders, venous cardiovascular thromboembolic disorders, or thromboembolic disorders in the chambers of the heart); atherosclerosis; restenosis; peripheral arterial disease; coronary bypass grafting surgery; carotid artery disease; arteritis; myocarditis; cardiovascular inflammation; vascular inflammation; coronary heart disease (CHD); unstable angina (UA); unstable refractory angina; stable angina (SA); chronic stable angina; acute coronary syndrome (ACS); myocardial infarction (first or recurrent); acute myocardial infarction (AMI); myocardial infarction; non-Q wave myocardial infarction; non-STE myocardial infarction; coronary artery disease; ischemic heart disease; cardiac ischemia; ischemia; ischemic sudden death; transient ischemic attack; stroke; peripheral occlusive arterial disease; venous thrombosis; deep vein thrombosis; thrombophlebitis; arterial embolism; coronary arterial thrombosis; cerebral arterial thrombosis, cerebral embolism; kidney embolism; pulmonary embolism; thrombosis (e.g., associated with prosthetic valves or other implants, indwelling catheters, stents, cardiopulmonary bypass, hemodialysis); thrombosis (e.g., associated with atherosclerosis, surgery, prolonged immobilization, arterial fibrillation, congenital thrombophilia, cancer, diabetes, hormones, or pregnancy); or cardiac arrhythmias (e.g., supraventricular arrhythmias, atrial arrhythmias, atrial flutter, or atrial fibrillation). In embodiments, the disease is a polycystic disease. In embodiments, the disease is polycystic kidney disease. In embodiments, the disease is stenosis. In embodiments, the disease is restenosis. In embodiments, the disease is neointimal proliferation. In embodiments, the disease is neointimal hyperplasia.

[00255] In another aspect is provided a method of treating aging in a subject in need of such treatment, the method including administering one or more compositions or compounds as described herein, including embodiments (e.g., a claim, embodiment, example, table, figure, or claim) to the subject. The present disclosure provides a method of treating immunosenescence comprising administering to the subject a therapeutically effective amount of one or more disclosed compounds or compositions.

[00256] In another aspect is provided one or more compositions or compounds as described herein for use as a medicament. In embodiments, the medicament may be useful for treating aging in a subject in need of such treatment. In embodiments, the use may include

administering one or more compositions or compounds as described herein, including embodiments (e.g., an aspect, embodiment, example, table, figure, or claim) to the subject.

[00257] In another aspect is provided one or more compositions or compounds disclosed herein for use in the treatment of aging in a subject in need of such treatment. In embodiments, the use may include administering one or more compositions or compounds as described herein, including embodiments (e.g., an aspect, embodiment, example, table, figure, or claim) to the subject.

[00258] In another aspect is provided a method of extending life span or inducing longevity in a subject in need of such treatment, the method including administering one or more compositions or compounds as described herein, including embodiments (e.g., a claim, embodiment, example, table, figure, or claim) to the subject.

[00259] In another aspect is provided one or more compositions or compounds as described herein for use as a medicament. In embodiments, the medicament may be useful for extending life span or inducing longevity in a subject in need of such treatment. In embodiments, the use may include administering one or more compositions or compounds as described herein, including embodiments (e.g., an aspect, embodiment, example, table, figure, or claim) to the subject.

[00260] In another aspect is provided one or more compositions or compounds for use in extending life span or inducing longevity in a subject in need of such treatment. In embodiments, the use may include administering one or more compositions or compounds as described herein, including embodiments (e.g., an aspect, embodiment, example, table, figure, or claim) to the subject.

[00261] In an aspect is provided a method of treating a polycystic disease in a subject in need of such treatment. The polycystic disease may be polycystic kidney disease. The method may include administering to the subject one or more compositions or compounds described herein. The method may include administering to the subject a therapeutically effective amount of one or more compositions or compounds described herein (e.g., an mTOR modulator (e.g., inhibitor) as described above).

[00262] In an aspect is provided one or more compositions or compounds as described herein for use as a medicament. In embodiments, the medicament is useful for treating a polycystic disease. The polycystic disease may be polycystic kidney disease. The use may include administering to the subject one or more compositions or compounds described

herein. The use may include administering to the subject a therapeutically effective amount of one or more compositions or compounds described herein (e.g., an mTOR modulator (e.g., inhibitor) as described above).

[00263] In an aspect is provided one or more compositions or compounds as described herein for use in the treatment of a polycystic disease in a subject in need of such treatment. The polycystic disease may be polycystic kidney disease. The use may include administering to the subject one or more compositions or compounds described herein. The use may include administering to the subject a therapeutically effective amount of one or more compositions or compounds described herein (e.g., an mTOR modulator (e.g., inhibitor) as described above).

[00264] In an aspect is provided a method of treating stenosis in a subject in need of such treatment. The stenosis may be restenosis. The method may include administering to the subject one or more compositions or compounds described herein. In embodiments the one or more compositions or compounds are administered in a drug eluting stent. The method may include administering to the subject a therapeutically effective amount of one or more compositions or compounds described herein (e.g., an mTOR modulator (e.g., inhibitor) as described above).

[00265] In an aspect is provided one or more compositions or compounds as described herein for use as a medicament. In embodiments, the medicament is useful for treating stenosis. The stenosis may be restenosis. The use may include administering to the subject one or more compositions or compounds described herein. In embodiments the compound is administered in a drug eluting stent. The use may include administering to the subject a therapeutically effective amount of one or more compositions or compounds described herein (e.g., an mTOR modulator (e.g., inhibitor) as described above).

[00266] In an aspect is provided one or more compositions or compounds as described herein for use in the treatment of stenosis in a subject in need of such treatment. The stenosis may be restenosis. The use may include administering to the subject one or more compositions or compounds described herein. In embodiments the one or more compositions or compounds are administered in a drug eluting stent. The use may include administering to the subject a therapeutically effective amount of one or more compositions or compounds described herein (e.g., an mTOR modulator (e.g., inhibitor) as described above).

[00267] In embodiments, the disease is a disease described herein and the compound is a compound described herein and the composition is a composition described herein.

Methods of Modulating mTOR

[00268] In some embodiments, compounds disclosed herein are more selective inhibitors of mTORC1 versus mTORC2. In some embodiments, compounds disclosed herein are more selective inhibitors of mTORC2 versus mTORC1. In some embodiments, compounds disclosed herein exhibit no selectivity difference between mTORC1 and mTORC2.

[00269] In another aspect is provided a method of modulating mTORC1 activity in a subject in need thereof, including administering to the subject an effective amount of a compound as described herein, or a pharmaceutically acceptable salt thereof. In embodiments, the method includes inhibiting mTORC1 activity. In embodiments, the method includes inhibiting mTORC1 activity and not inhibiting mTORC2 activity.

[00270] In embodiments, the method includes inhibiting mTORC1 activity more than inhibiting mTORC2 activity. In embodiments, the method includes inhibiting mTORC1 activity at least 1.1 fold as much as inhibiting mTORC2 activity (e.g., at least 1.1, 1.2, 1.3, 1.4, 1.5, 1.6, 1.7, 1.8, 1.9, 2, 3, 4, 5, 6, 7, 8, 9, 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000, 2000, 3000, 4000, 5000, 6000, 7000, 8000, 9000, 10000, 10000, 20000, 30000, 40000, 50000, 60000, 70000, 80000, 90000, 100000, 100000, 200000, 300000, 400000, 500000, 600000, 700000, 800000, 900000, or 1000000 fold).

[00271] In another aspect is provided a method of modulating mTORC2 activity in a subject in need thereof, including administering to the subject an effective amount of a compound as described herein, or a pharmaceutically acceptable salt thereof. In embodiments, the method includes inhibiting mTORC2 activity. In embodiments, the method includes inhibiting mTORC2 activity and not inhibiting mTORC1 activity.


[00272] In embodiments, the method includes inhibiting mTORC2 activity more than inhibiting mTORC1 activity. In embodiments, the method includes inhibiting mTORC2 activity at least 1.1 fold as much as inhibiting mTORC1 activity (e.g., at least 1.1, 1.2, 1.3, 1.4, 1.5, 1.6, 1.7, 1.8, 1.9, 2, 3, 4, 5, 6, 7, 8, 9, 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000, 2000, 3000, 4000, 5000, 6000, 7000, 8000, 9000, 10000, 10000, 20000, 30000, 40000, 50000, 60000, 70000, 80000, 90000, 100000, 100000, 200000, 300000, 400000, 500000, 600000, 700000, 800000, 900000, or 1000000 fold).

[00273] In some embodiments, the mTOR is in a cell. In some embodiments, the cell is a mammalian cell, such as a human cell. The cell may be isolated *in vitro*, form part of a tissue *in vitro*, or may form part of an organism.

Exemplary Embodiments

[00274] Some embodiments of this disclosure are Embodiment I, as follows:

[00275] Embodiment I-1. A compound of Formula I:



or a pharmaceutically acceptable salt or tautomer thereof, wherein:

R^{32} is $-H$, $=O$ or $-OR^3$;

R^{28} is $-H$ or $-C(=Z^1)-R^{28a}$;


R^{40} is $-H$ or $-C(=Z^1)-R^{40a}$;


wherein at least one of R^{28} and R^{40} is not H ;

Z^1 is O or S ;

R^{28a} and R^{40a} are independently $-A^1-L^1-A^2-B$; $-A^1-A^2-B$; $-L^2-A^1-L^1-A^2-L^3-B$; $-O-(C_1-C_6)alkyl$; or $-O-(C_6-C_{10})aryl$; wherein the aryl is unsubstituted or substituted with 1-5 substituents selected from $-NO_2$ and halogen;

A^1 and A^2 are independently absent or are independently selected from





wherein the bond on the left side of A¹, as drawn, is bound to -C(=Z¹)- or L²; and
 wherein the bond on the right side of the A² moiety, as drawn, is bound to B or L³;

each Q is independently 1 to 3 rings selected from arylene, cycloalkylene, heteroarylene, and heterocyclylene;

each X is independently absent or 1 to 2 rings selected from arylene, cycloalkylene, heteroarylene, and heterocyclylene;


each X¹ is a heteroarylene or heterocyclylene ring;

each W is independently absent or 1 to 2 rings selected from arylene, cycloalkylene, heteroarylene, and heterocyclylene;


each W¹ is a heteroarylene or heterocyclylene ring;


each G is independently absent or a ring selected from arylene, cycloalkylene, heteroarylene, and heterocyclylene;

each G¹ and G² are independently heteroarylene or heterocyclylene ring;




L² and L³ are independently absent or are independently selected from






B is selected from



B^1 is selected from $^{-}\text{NR}^3\text{-(C(R}^3\text{)}_2)_n\text{-}$, $^{-}\text{NR}^3\text{-(C(R}^3\text{)}_2)_n\text{-(C}_6\text{-C}_{10}\text{)arylene-(C(R}^3\text{)}_2)_n\text{-}$, $^{-}\text{NR}^3\text{-(C(R}^3\text{)}_2)_n\text{-heteroarylene-}$, $^{-}\text{(C}_6\text{-C}_{10}\text{)arylene-}$, $^{-}\text{NR}^3\text{-(C(R}^3\text{)}_2)_n\text{-NR}^3\text{C(O)-}$, $^{-}\text{NR}^3\text{-(C(R}^3\text{)}_2)_n\text{-heteroarylene-heterocyclylene-(C}_6\text{-C}_{10}\text{)arylene-}$, $^{-}\text{heteroarylene-heterocyclylene-(C}_6\text{-C}_{10}\text{)arylene-}$, $\text{C(=O)-C(R}^3\text{)}_2\text{p-}$, $\text{C(=O)-C(R}^3\text{)}_2\text{p-heteroarylene-}$, $\text{N-methylpiperazine-C(R}^3\text{)}_2\text{p-}$, $\text{N-methylpiperidine-C(R}^3\text{)}_2\text{p-}$, $\text{N-methylpyrrolidine-C(R}^3\text{)}_2\text{p-}$, and



$\text{---}^3\text{---NR}^3\text{---}(\text{C}(\text{R}^3)_2)_n\text{---S}(\text{O})_2\text{---arylene---C}(\text{O})\text{---}$, wherein the $\text{---}^3\text{---}$ bond on the left side of B^1 , as drawn, is bound to A^2 , L^3 , or L^1 ; and wherein the heteroarylene, heterocyclylene, and arylene are optionally substituted with alkyl, hydroxyalkyl, haloalkyl, alkoxy, halogen, or hydroxyl;

each R^3 is independently H or $(\text{C}_1\text{---C}_6)\text{alkyl}$;

each R^4 is independently H, $(\text{C}_1\text{---C}_6)\text{alkyl}$, halogen, 5-12 membered heteroaryl, 5-12 membered heterocyclyl, $(\text{C}_6\text{---C}_{10})\text{aryl}$, wherein the heteroaryl, heterocyclyl, and aryl are optionally substituted with $\text{---N}(\text{R}^3)_2$, ---OR^3 , halogen, $(\text{C}_1\text{---C}_6)\text{alkyl}$, $\text{---}(\text{C}_1\text{---C}_6)\text{alkylene---heteroaryl}$, $\text{---}(\text{C}_1\text{---C}_6)\text{alkylene---CN}$, $\text{---C}(\text{O})\text{NR}^3\text{---heteroaryl}$; or $\text{---C}(\text{O})\text{NR}^3\text{---heterocyclyl}$;

each R^5 is independently H, $(\text{C}_1\text{---C}_6)\text{alkyl}$, $\text{---C}(\text{O})\text{OR}^3$, or $\text{---N}(\text{R}^3)_2$, wherein the alkyl is optionally substituted with $\text{---N}(\text{R}^3)_2$ or ---OR^3 ;

each R^6 is independently H, $(\text{C}_1\text{---C}_6)\text{alkyl}$, $\text{---C}(\text{O})\text{OR}^3$, or $\text{---N}(\text{R}^3)_2$, wherein the alkyl is optionally substituted with $\text{---N}(\text{R}^3)_2$ or ---OR^3 ;

each R^7 is independently H, $(\text{C}_1\text{---C}_6)\text{alkyl}$, $\text{---C}(\text{O})\text{OR}^3$, or $\text{---N}(\text{R}^3)_2$, wherein the alkyl is optionally substituted with $\text{---N}(\text{R}^3)_2$ or ---OR^3 ;

each R^8 is independently H, $(\text{C}_1\text{---C}_6)\text{alkyl}$, $\text{---C}(\text{O})\text{OR}^3$, or $\text{---N}(\text{R}^3)_2$, wherein the alkyl is optionally substituted with $\text{---N}(\text{R}^3)_2$ or ---OR^3 ;

each Y is independently $\text{C}(\text{R}^3)_2$ or a bond;

each n is independently a number from one to 12;


each o is independently a number from zero to 30;

each p is independently a number from zero to 12;

each q is independently a number from zero to 30; and

each r is independently a number from one to 6.

[00276] Embodiment I-2. A compound of Formula II:



(II)

or a pharmaceutically acceptable salt or tautomer thereof, wherein:

R^{32} is $-H$, $=O$ or $-OR^3$;

R^{28} is $-H$ or $-C(=Z^1)-R^{28a}$;


R^{40} is $-H$ or $-C(=Z^1)-R^{40a}$;

wherein at least one of R^{28} and R^{40} is not H ;

Z^1 is O or S ;

R^{28a} and R^{40a} are independently $-A^1-L^1-A^2-B$; $-A^1-A^2-B$; $-O-(C_1-C_6)alkyl$; or $-O-(C_6-C_{10})aryl$; wherein the aryl is unsubstituted or substituted with 1-5 substituents selected from $-NO_2$ and halogen;

A^1 and A^2 are independently absent or are independently selected from



wherein the bond on the left side of A¹, as drawn, is bound to -C(=Z¹)-; and wherein the bond on the right side of the A² moiety, as drawn, is bound to B;

each Q is independently 1 to 3 rings selected from arylene, cycloalkylene, heteroarylene, and heterocyclylene;

each X is independently absent or 1 to 2 rings selected from arylene, cycloalkylene, heteroarylene, and heterocyclylene;



each X¹ is a heteroarylene or heterocyclylene ring;

each W is independently absent or 1 to 2 rings selected from arylene, cycloalkylene, heteroarylene, and heterocyclylene;

each W¹ is a heteroarylene or heterocyclylene ring;

each G is independently absent or a ring selected from arylene, cycloalkylene, heteroarylene, and heterocyclylene;


each G¹ and G² are independently heteroarylene or heterocyclylene ring;



[00277] Embodiment I-3. The compound of Embodiment I-1 or I-2, wherein R^{32} is =O.

[00278] Embodiment I-4. The compound of Embodiment I-1 or I-2, wherein R^{32} is $-OR^3$.

[00279] Embodiment I-5. The compound of any one of Embodiments I-1 to I-4, wherein the compounds are represented by the structure of Formula I-40:



or a pharmaceutically acceptable salt or tautomer thereof.

[00280] Embodiment I-6. The compound of Embodiment I-5, wherein Z^1 is O.

[00281] Embodiment I-7. The compound of Embodiment I-5, wherein Z^1 is S.

[00282] Embodiment I-8. The compound of any one of Embodiments I-5 to I-7, wherein R^{40a} is $-A^1-L^1-A^2-B$, wherein A^1 and A^2 are absent.

[00283] Embodiment I-9. The compound of any one of Embodiments I-5 to I-7, wherein R^{40a} is $-A^1-L^1-A^2-B$, wherein A^2 is absent.

[00284] Embodiment I-10. The compound of any one of Embodiments I-5 to I-7, wherein R^{40a} is $-A^1-L^1-A^2-B$, wherein A^1 is absent.

[00285] Embodiment I-11. The compound of any one of Embodiments I-5 to I-7, wherein R^{40a} is $-A^1-L^1-A^2-B$.

[00286] Embodiment I-12. The compound of any one of Embodiments I-5 to I-7, wherein R^{40a} is $-A^1-A^2-B$.


[00287] Embodiment I-13. The compound of any one of Embodiments I-5 to I-7, wherein R^{40a} is $--L^2-A^1-L^1-A^2-L^3-B$, wherein L^2 and A^1 are absent.

[00288] Embodiment I-14. The compound of any one of Embodiments I-5 to I-7, wherein R^{40a} is $--L^2-A^1-L^1-A^2-L^3-B$, wherein L^2 is absent.

[00289] Embodiment I-15. The compound of any one of Embodiments I-5 to I-7, wherein R^{40a} is $-L^2-A^1-L^1-A^2-L^3-B$, wherein L^3 is absent.

[00290] Embodiment I-16. The compound of any one of Embodiments I-5 to I-7, wherein R^{40a} is $-O-(C_1-C_6)alkyl$ or $-O-(C_6-C_{10})aryl$; wherein the aryl is unsubstituted or substituted with 1-5 substituents selected from $-NO_2$ and halogen.

[00291] Embodiment I-17. The compound of any one of Embodiments I-1 to I-4, wherein the compounds are represented by the structure of Formula I-28:



or a pharmaceutically acceptable salt or tautomer thereof.

[00292] Embodiment I-18. The compound of Embodiment I-17, wherein Z^1 is O.

[00293] Embodiment I-19. The compound of Embodiment I-17, wherein Z^1 is S.

[00294] Embodiment I-20. The compound of any one of Embodiments I-17 to I-19, wherein R^{28a} is $-A^1-L^1-A^2-B$, wherein A^1 and A^2 are absent.

[00295] Embodiment I-21. The compound of any one of Embodiments I-17 to I-19, wherein R^{28a} is $-A^1-L^1-A^2-B$, wherein A^2 is absent.

[00296] Embodiment I-22. The compound of any one of Embodiments I-17 to I-19, wherein R^{28a} is $-A^1-L^1-A^2-B$, wherein A^1 is absent.

[00297] Embodiment I-23. The compound of any one of Embodiments I-17 to I-19, wherein R^{28a} is $-A^1-L^1-A^2-B$.

[00298] Embodiment I-24. The compound of any one of Embodiments I-17 to I-19, wherein R^{28a} is $-A^1-A^2-B$.


[00299] Embodiment I-25. The compound of any one of Embodiments I-17 to I-19, wherein R^{28a} is $-L^2-A^1-L^1-A^2-L^3-B$, wherein L^2 and A^1 are absent.

[00300] Embodiment I-26. The compound of any one of Embodiments I-17 to I-19, wherein R^{28a} is $-L^2-A^1-L^1-A^2-L^3-B$, wherein L^2 is absent.

[00301] Embodiment I-27. The compound of any one of Embodiments I-17 to I-19, wherein R^{28a} is $-L^2-A^1-L^1-A^2-L^3-B$, wherein L^3 is absent.


[00302] Embodiment I-28. The compound of any one of Embodiments I-17 to I-19, wherein R^{28a} is $-O-(C_1-C_6)\text{alkyl}$ or $-O-(C_6-C_{10})\text{aryl}$; wherein the aryl is unsubstituted or substituted with 1-5 substituents selected from $-\text{NO}_2$ and halogen.

[00303] Embodiment I-29. The compound of any one of Embodiments I-1 to I-11, I-13 to




I-15, I-17 to I-23, and I-25 to I-27, wherein L^1 is

[00304] Embodiment I-30. The compound of any one of Embodiments I-1 to I-11, I-13 to




I-15, I-17 to I-23, and I-25 to I-27, wherein L^1 is

[00305] Embodiment I-31. The compound of any one of Embodiments I-1 to I-11, I-13 to I-15, I-17 to I-23, and I-25 to I-27, wherein L^1 is




[00306] Embodiment I-32. The compound of any one of Embodiments I-1 to I-11, I-13 to




I-15, I-17 to I-23, and I-25 to I-27, wherein L^1 is

[00307] Embodiment I-33. The compound of any one of Embodiments I-1 to I-11, I-13 to



I-15, I-17 to I-23, and I-25 to I-27, wherein L^1 is


[00308] Embodiment I-34. The compound of any one of Embodiments I-1 to I-7, I-15, I-17



to I-19, and I-27, wherein L^2 is


[00309] Embodiment I-35. The compound of any one of Embodiments I-1 to I-7, I-13 to I-

14, I-17 to I-19, and I-25 to I-26, wherein L^3 is




[00310] Embodiment I-36. The compound of any one of Embodiments I-1 to I-8, I-10, I-13, I-17 to I-19, I-20, I-22, I-25 and I-29 to I-35, wherein A^1 is absent.


[00311] Embodiment I-37. The compound of any one of Embodiments I-1 to I-7, I-9, I-11 to I-12, I-14 to I-15, I-17 to I-19, I-21, I-23 to I-24, I-26 to I-27, and I-29 to I-35, wherein A^1




[00312] Embodiment I-38. The compound of any one of Embodiments I-1 to I-7, I-9, I-11 to I-12, I-14 to I-15, I-17 to I-19, I-21, I-23 to I-24, I-26 to I-27, and I-29 to I-35, wherein A^1




[00313] Embodiment I-39. The compound of any one of Embodiments I-1 to I-7, I-9, I-11 to I-12, I-14 to I-15, I-17 to I-19, I-21, I-23 to I-24, I-26 to I-27, and I-29 to I-35, wherein A^1




[00314] Embodiment I-40. The compound of any one of Embodiments I-1 to I-7, I-9, I-11 to I-12, I-14 to I-15, I-17 to I-19, I-21, I-23 to I-24, I-26 to I-27, and I-29 to I-35, wherein A^1




[00315] Embodiment I-41. The compound of any one of Embodiments I-1 to I-7, I-9, I-11 to I-12, I-14 to I-15, I-17 to I-19, I-21, I-23 to I-24, I-26 to I-27, and I-29 to I-35, wherein A¹




[00316] Embodiment I-42. The compound of any one of Embodiments I-1 to I-7, I-9, I-11 to I-12, I-14 to I-15, I-17 to I-19, I-21, I-23 to I-24, I-26 to I-27, and I-29 to I-35, wherein A¹



[00317] Embodiment I-43. The compound of any one of Embodiments I-1 to I-7, I-9, I-11 to I-12, I-14 to I-15, I-17 to I-19, I-21, I-23 to I-24, I-26 to I-27, and I-29 to I-35, wherein A¹




[00318] Embodiment I-44. The compound of any one of Embodiments I-1 to I-7, I-9, I-11 to I-12, I-14 to I-15, I-17 to I-19, I-21, I-23 to I-24, I-26 to I-27, and I-29 to I-35, wherein A¹




[00319] Embodiment I-45. The compound of any one of Embodiments I-1 to I-9, I-17 to I-21, and I-29 to I-44, wherein A² is absent.

[00320] Embodiment I-46. The compound of any one of Embodiments I-1 to I-7, I-10 to I-




15, I-17 to I-19, I-22 to I-27 and I-29 to I-44, wherein A² is


[00321] Embodiment I-47. The compound of any one of Embodiments I-1 to I-7, I-10 to I-




[00322] Embodiment I-48. The compound of any one of Embodiments I-1 to I-7, I-10 to I-




[00323] Embodiment I-49. The compound of any one of Embodiments I-1 to I-7, I-10 to I-




[00324] Embodiment I-50. The compound of any one of Embodiments I-1 to I-7, I-10 to I-




[00325] Embodiment I-51. The compound of any one of Embodiments I-1 to I-7, I-10 to I-




[00326] Embodiment I-52. The compound of any one of Embodiments I-1 to I-7, I-10 to I-




[00327] Embodiment I-53. The compound of any one of Embodiments I-1 to I-7, I-10 to I-



[00328] Embodiment I-54. The compound of any one of Embodiments I-1 to I-53, wherein




[00329] Embodiment I-55. The compound of any one of Embodiments I-1 to I-53, wherein



[00330] Embodiment I-56. The compound of any one of Embodiments I-1 to I-53, wherein




[00331] Embodiment I-57. The compound of any one of Embodiments I-1 to I-53, wherein




[00332] Embodiment I-58. The compound of any one of Embodiments I-1 to I-57, wherein R^4 is 5-12 membered heteroaryl, optionally substituted with $-N(R^3)_2$, $-OR^3$, halogen, (C_1-C_6) alkyl, $-(C_1-C_6)$ alkylene-heteroaryl, $-(C_1-C_6)$ alkylene-CN, or $-C(O)NR^3$ -heteroaryl.


[00333] Embodiment I-59. The compound of any one of Embodiments I-1 to I-58, or a pharmaceutically acceptable salt or tautomer thereof, wherein compound has the following formula:




[00334] Embodiment I-60. A compound selected from the group consisting of:




Example 1




Example 2




Example 3




Example 4




Example 5




Example 6




Example 7




Example 8




Example 9




Example 10




Example 11




Example 12




Example 13




Example 14




Example 15




Example 16




Example 17




Example 18




Example 19




Example 20




Example 21




Example 22




Example 23




Example 24




Example 25




Example 26




Example 27




Example 28



Example 29




Example 30




[illegible][illegible][illegible]CNc1nc2cc3[nH]cnc3c2n1CCCCNC(=O)OCCOCCOCCOCCOCCOCCOCCOCC(=O)Nc1ccc2nc3c(ncn3C2)c1


160




Example 36




Example 37




Example 38




Example 39




Example 40




Example 41




Example 42




Example 43




Example 44




Example 45




Example 46




Example 47




Example 48




Example 49




[illegible][illegible][illegible]CN1C=NC2=C1N=CN2C3=CC=C4C(=C3)OC(=N4)N5C=CC=C5C6=CC=CC=C6C7=CC=CC=C7C8=CC=CC=C8C9=CC=CC=C9C10=CC=CC=C10C11=CC=CC=C11C12=CC=CC=C12C13=CC=CC=C13C14=CC=CC=C14C15=CC=CC=C15C16=CC=CC=C16C17=CC=CC=C17C18=CC=CC=C18C19=CC=CC=C19C20=CC=CC=C20C21=CC=CC=C21C22=CC=CC=C22C23=CC=CC=C23C24=CC=CC=C24C25=CC=CC=C25C26=CC=CC=C26C27=CC=CC=C27C28=CC=CC=C28C29=CC=CC=C29C30=CC=CC=C30C31=CC=CC=C31C32=CC=CC=C32C33=CC=CC=C33C34=CC=CC=C34C35=CC=CC=C35C36=CC=CC=C36C37=CC=CC=C37C38=CC=CC=C38C39=CC=CC=C39C40=CC=CC=C40C41=CC=CC=C41C42=CC=CC=C42C43=CC=CC=C43C44=CC=CC=C44C45=CC=CC=C45C46=CC=CC=C46C47=CC=CC=C47C48=CC=CC=C48C49=CC=CC=C49C50=CC=CC=C50C51=CC=CC=C51C52=CC=CC=C52C53=CC=CC=C53C54=CC=CC=C54C55=CC=CC=C55C56=CC=CC=C56C57=CC=CC=C57C58=CC=CC=C58C59=CC=CC=C59C60=CC=CC=C60C61=CC=CC=C61C62=CC=CC=C62C63=CC=CC=C63C64=CC=CC=C64C65=CC=CC=C65C66=CC=CC=C66C67=CC=CC=C67C68=CC=CC=C68C69=CC=CC=C69C70=CC=CC=C70C71=CC=CC=C71C72=CC=CC=C72C73=CC=CC=C73C74=CC=CC=C74C75=CC=CC=C75C76=CC=CC=C76C77=CC=CC=C77C78=CC=CC=C78C79=CC=CC=C79C80=CC=CC=C80C81=CC=CC=C81C82=CC=CC=C82C83=CC=CC=C83C84=CC=CC=C84C85=CC=CC=C85C86=CC=CC=C86C87=CC=CC=C87C88=CC=CC=C88C89=CC=CC=C89C90=CC=CC=C90C91=CC=CC=C91C92=CC=CC=C92C93=CC=CC=C93C94=CC=CC=C94C95=CC=CC=C95C96=CC=CC=C96C97=CC=CC=C97C98=CC=CC=C98C99=CC=CC=C99C100=CC=CC=C100C101=CC=CC=C101C102=CC=CC=C102C103=CC=CC=C103C104=CC=CC=C104C105=CC=CC=C105C106=CC=CC=C106C107=CC=CC=C107C108=CC=CC=C108C109=CC=CC=C109C110=CC=CC=C110C111=CC=CC=C111C112=CC=CC=C112C113=CC=CC=C113C114=CC=CC=C114C115=CC=CC=C115C116=CC=CC=C116C117=CC=CC=C117C118=CC=CC=C118C119=CC=CC=C119C120=CC=CC=C120C121=CC=CC=C121C122=CC=CC=C122C123=CC=CC=C123C124=CC=CC=C124C125=CC=CC=C125C126=CC=CC=C126C127=CC=CC=C127C128=CC=CC=C128C129=CC=CC=C129C130=CC=CC=C130C131=CC=CC=C131C132=CC=CC=C132C133=CC=CC=C133C134=CC=CC=C134C135=CC=CC=C135C136=CC=CC=C136C137=CC=CC=C137C138=CC=CC=C138C139=CC=CC=C139C140=CC=CC=C140C141=CC=CC=C141C142=CC=CC=C142C143=CC=CC=C143C144=CC=CC=C144C145=CC=CC=C145C146=CC=CC=C146C147=CC=CC=C147C148=CC=CC=C148C149=CC=CC=C149C150=CC=CC=C150C151=CC=CC=C151C152=CC=CC=C152C153=CC=CC=C153C154=CC=CC=C154C155=CC=CC=C155C156=CC=CC=C156C157=CC=CC=C157C158=CC=CC=C158C159=CC=CC=C159C160=CC=CC=C160C161=CC=CC=C161C162=CC=CC=C162C163=CC=CC=C163C164=CC=CC=C164C165=CC=CC=C165C166=CC=CC=C166C167=CC=CC=C167C168=CC=CC=C168C169=CC=CC=C169C170=CC=CC=C170C171=CC=CC=C171C172=CC=CC=C172C173=CC=CC=C173C174=CC=CC=C174C175=CC=CC=C175C176=CC=CC=C176C177=CC=CC=C177C178=CC=CC=C178C179=CC=CC=C179C180=CC=CC=C180C181=CC=CC=C181C182=CC=CC=C182C183=CC=CC=C183C184=CC=CC=C184C185=CC=CC=C185C186=CC=CC=C186C187=CC=CC=C187C188=CC=CC=C188C189=CC=CC=C189C190=CC=CC=C190C191=CC=CC=C191C192=CC=CC=C192C193=CC=CC=C193C194=CC=CC=C194C195=CC=CC=C195C196=CC=CC=C196C197=CC=CC=C197C198=CC=CC=C198C199=CC=CC=C199C200=CC=CC=C200C201=CC=CC=C201C202=CC=CC=C202C203=CC=CC=C203C204=CC=CC=C204C205=CC=CC=C205C206=CC=CC=C206C207=CC=CC=C207C208=CC=CC=C208C209=CC=CC=C209C210=CC=CC=C210C211=CC=CC=C211C212=CC=CC=C212C213=CC=CC=C213C214=CC=CC=C214C215=CC=CC=C215C216=CC=CC=C216C217=CC=CC=C217C218=CC=CC=C218C219=CC=CC=C219C220=CC=CC=C220C221=CC=CC=C221C222=CC=CC=C222C223=CC=CC=C223C224=CC=CC=C224C225=CC=CC=C225C226=CC=CC=C226C227=CC=CC=C227C228=CC=CC=C228C229=CC=CC=C229C230=CC=CC=C230C231=CC=CC=C231C232=CC=CC=C232C233=CC=CC=C233C234=CC=CC=C234C235=CC=CC=C235C236=CC=CC=C236C237=CC=CC=C237C238=CC=CC=C238C239=CC=CC=C239C240=CC=CC=C240C241=CC=CC=C241C242=CC=CC=C242C243=CC=CC=C243C244=CC=CC=C244C245=CC=CC=C245C246=CC=CC=C246C247=CC=CC=C247C248=CC=CC=C248C249=CC=CC=C249C250=CC=CC=C250C251=CC=CC=C251C252=CC=CC=C252C253=CC=CC=C253C254=CC=CC=C254C255=CC=CC=C255C256=CC=CC=C256C257=CC=CC=C257C258=CC=CC=C258C259=CC=CC=C259C260=CC=CC=C260C261=CC=CC=C261C262=CC=CC=C262C263=CC=CC=C263C264=CC=CC=C264C265=CC=CC=C265C266=CC=CC=C266C267=CC=CC=C267C268=CC=CC=C268C269=CC=CC=C269C270=CC=CC=C270C271=CC=CC=C271C272=CC=CC=C272C273=CC=CC=C273C274=CC=CC=C274C275=CC=CC=C275C276=CC=CC=C276C277=CC=CC=C277C278=CC=CC=C278C279=CC=CC=C279C280=CC=CC=C280C281=CC=CC=C281C282=CC=CC=C282C283=CC=CC=C283C284=CC=CC=C284C285=CC=CC=C285C286=CC=CC=C286C287=CC=CC=C287C288=CC=CC=C288C289=CC=CC=C289C290=CC=CC=C290C291=CC=CC=C291C292=CC=CC=C292C293=CC=CC=C293C294=CC=CC=C294C295=CC=CC=C295C296=CC=CC=C296C297=CC=CC=C297C298=CC=CC=C298C299=CC=CC=C299C300=CC=CC=C300C301=CC=CC=C301C302=CC=CC=C302C303=CC=CC=C303C304=CC=CC=C304C305=CC=CC=C305C306=CC=CC=C306C307=CC=CC=C307C308=CC=CC=C308C309=CC=CC=C309C310=CC=CC=C310C311=CC=CC=C311C312=CC=CC=C312C313=CC=CC=C313C314=CC=CC=C314C315=CC=CC=C315C316=CC=CC=C316C317=CC=CC=C317C318=CC=CC=C318C319=CC=CC=C319C320=CC=CC=C320C321=CC=CC=C321C322=CC=CC=C322C323=CC=CC=C323C324=CC=CC=C324C325=CC=CC=C325C326=CC=CC=C326C327=CC=CC=C327C328=CC=CC=C328C329=CC=CC=C329C330=CC=CC=C330C331=CC=CC=C331C332=CC=CC=C332C333=CC=CC=C333C334=CC=CC=C334C335=CC=CC=C335C336=CC=CC=C336C337=CC=CC=C337C338=CC=CC=C338C339=CC=CC=C339C340=CC=CC=C340C341=CC=CC=C341C342=CC=CC=C342C343=CC=CC=C343C344=CC=CC=C344C345=CC=CC=C345C346=CC=CC=C346C347=CC=CC=C347C348=CC=CC=C348C349=CC=CC=C349C350=CC=CC=C350C351=CC=CC=C351C352=CC=CC=C352C353=CC=CC=C353C354=CC=CC=C354C355=CC=CC=C355C356=CC=CC=C356C357=CC=CC=C357C358=CC=CC=C358C359=CC=CC=C359C360




Example 55




Example 56




Example 57




Example 58




Example 59




Example 60




Example 61




Example 62




Example 63




Example 64



Example 65



Example 66



or a pharmaceutically acceptable salt or isomer thereof.

[00335] Embodiment I-61. A pharmaceutical composition comprising a compound of any one of Embodiments I-1 to I-60, or a pharmaceutically acceptable salt thereof, and at least one of a pharmaceutically acceptable carrier, diluent, or excipient.

[00336] Embodiment I-62. A method of treating a disease or disorder mediated by mTOR comprising administering to the subject suffering from or susceptible to developing a disease or disorder mediated by mTOR a therapeutically effective amount of one or more compounds of any one of Embodiments I-1 to I-60, or a pharmaceutically acceptable salt thereof.

[00337] Embodiment I-63. A method of preventing a disease or disorder mediated by mTOR comprising administering to the subject suffering from or susceptible to developing a disease or disorder mediated by mTOR a therapeutically effective amount of one or more

compounds of any one of Embodiments I-1 to I-60, or a pharmaceutically acceptable salt thereof.

[00338] Embodiment I-64. A method of reducing the risk of a disease or disorder mediated by mTOR comprising administering to the subject suffering from or susceptible to developing a disease or disorder mediated by mTOR a therapeutically effective amount of one or more compounds of any one of Embodiments I-1 to I-60, or a pharmaceutically acceptable salt thereof.

[00339] Embodiment I-65. The method of any one of Embodiments I-62 to I-64, wherein the disease is cancer or an immune-mediated disease.

[00340] Embodiment I-66. The method of Embodiment I-65, wherein the cancer is selected from brain and neurovascular tumors, head and neck cancers, breast cancer, lung cancer, mesothelioma, lymphoid cancer, stomach cancer, kidney cancer, renal carcinoma, liver cancer, ovarian cancer, ovary endometriosis, testicular cancer, gastrointestinal cancer, prostate cancer, glioblastoma, skin cancer, melanoma, neuro cancers, spleen cancers, pancreatic cancers, blood proliferative disorders, lymphoma, leukemia, endometrial cancer, cervical cancer, vulva cancer, prostate cancer, penile cancer, bone cancers, muscle cancers, soft tissue cancers, intestinal or rectal cancer, anal cancer, bladder cancer, bile duct cancer, ocular cancer, gastrointestinal stromal tumors, and neuro-endocrine tumors.

[00341] Embodiment I-67. The method of Embodiment I-65, wherein the immune-mediated disease is selected from resistance by transplantation of heart, kidney, liver, medulla ossium, skin, cornea, lung, pancreas, intestinum tenue, limb, muscle, nerves, duodenum, small-bowel, or pancreatic-islet-cell; graft-versus-host diseases brought about by medulla ossium transplantation; rheumatoid arthritis, systemic lupus erythematosus, Hashimoto's thyroiditis, multiple sclerosis, myasthenia gravis, type I diabetes, uveitis, allergic encephalomyelitis, and glomerulonephritis.

[00342] Embodiment I-68. A method of treating cancer comprising administering to the subject a therapeutically effective amount of one or more compounds of any one of Embodiments I-1 to I-60, or a pharmaceutically acceptable salt thereof.

[00343] Embodiment I-69. The method of Embodiment I-68, wherein the cancer is selected from brain and neurovascular tumors, head and neck cancers, breast cancer, lung cancer, mesothelioma, lymphoid cancer, stomach cancer, kidney cancer, renal carcinoma, liver cancer, ovarian cancer, ovary endometriosis, testicular cancer, gastrointestinal cancer,

prostate cancer, glioblastoma, skin cancer, melanoma, neuro cancers, spleen cancers, pancreatic cancers, blood proliferative disorders, lymphoma, leukemia, endometrial cancer, cervical cancer, vulva cancer, prostate cancer, penile cancer, bone cancers, muscle cancers, soft tissue cancers, intestinal or rectal cancer, anal cancer, bladder cancer, bile duct cancer, ocular cancer, gastrointestinal stromal tumors, and neuro-endocrine tumors.

[00344] Embodiment I-70. A method of treating an immune-mediated disease comprising administering to the subject a therapeutically effective amount of one or more compounds of any one of Embodiments I-1 to I-60, or a pharmaceutically acceptable salt thereof.

[00345] Embodiment I-71. The method of Embodiment I-70, wherein the immune-mediated disease is selected from resistance by transplantation of heart, kidney, liver, medulla ossium, skin, cornea, lung, pancreas, intestinum tenue, limb, muscle, nerves, duodenum, small-bowel, or pancreatic-islet-cell; graft-versus-host diseases brought about by medulla ossium transplantation; rheumatoid arthritis, systemic lupus erythematosus, Hashimoto's thyroiditis, multiple sclerosis, myasthenia gravis, type I diabetes, uveitis, allergic encephalomyelitis, and glomerulonephritis.

[00346] Embodiment I-72. A method of treating an age related condition comprising administering to the subject a therapeutically effective amount of one or more compounds of any one of Embodiments I-1 to I-60, or a pharmaceutically acceptable salt thereof.

[00347] Embodiment I-73. The method of Embodiment I-72, wherein the age related condition is selected from sarcopenia, skin atrophy, muscle wasting, brain atrophy, atherosclerosis, arteriosclerosis, pulmonary emphysema, osteoporosis, osteoarthritis, high blood pressure, erectile dysfunction, dementia, Huntington's disease, Alzheimer's disease, cataracts, age-related macular degeneration, prostate cancer, stroke, diminished life expectancy, impaired kidney function, and age-related hearing loss, aging-related mobility disability (e.g., frailty), cognitive decline, age-related dementia, memory impairment, tendon stiffness, heart dysfunction such as cardiac hypertrophy and systolic and diastolic dysfunction, immunosenescence, cancer, obesity, and diabetes.

[00348] Embodiment I-74. A compound of any one of Embodiments I-1 to I-60, or a pharmaceutically acceptable salt thereof, for use in treating, preventing, or reducing the risk of a disease or condition mediated by mTOR.

[00349] Embodiment I-75. Use of a compound of any of Embodiments I-1 to I-60, or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament for treating, preventing, or reducing the risk of a disease or disorder mediated by mTOR.

[00350] Embodiment I-76. A compound of any one of Embodiments I-1 to I-60, or a pharmaceutically acceptable salt thereof, for use in treating cancer.

[00351] Embodiment I-77. Use of a compound of any one of Embodiments I-1 to I-60, or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament for treating cancer.

[00352] Embodiment I-78. A compound of any one of Embodiments I-1 to I-60, or a pharmaceutically acceptable salt thereof, for use in treating an immune-mediated disease.


[00353] Embodiment I-79. Use of a compound of any one of Embodiments I-1 to I-60, or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament for treating an immune-mediated disease.

[00354] Embodiment I-80. A compound of any one of Embodiments I-1 to I-60, or a pharmaceutically acceptable salt thereof, for use in treating an age related condition.

[00355] Embodiment I-81. Use of a compound of any one of Embodiments I-1 to I-60, or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament for treating an age related condition.

[00356] Some embodiments of this disclosure are Embodiment II, as follows:

[00357] Embodiment II-1. A compound of Formula Ic:



or a pharmaceutically acceptable salt or tautomer thereof, wherein:

R^{32} is $-H$, $=O$, $-OR^3$, $-N_3$, or $-O-C(=Z^1)-R^{32a}$;

R^{28} is $-H$, $(C_1-C_6)alkyl$, or $-C(=Z^1)-R^{28a}$;


R^{40} is $-H$ or $-C(=Z^1)-R^{40a}$;

wherein when R^{28} and R^{40} are H , then R^{32} is not $=O$;


each Z^1 is independently O or S ;

R^{28a} , R^{32a} , and R^{40a} are independently $-A^1-L^1-A^2-B$; $-A^1-A^2-B$; $-L^2-A^1-L^1-A^2-L^3-B$; $-O-(C_1-C_6)alkyl$; or $-O-(C_6-C_{10})aryl$; wherein the aryl of $-O-(C_6-C_{10})aryl$ is unsubstituted or substituted with 1-5 substituents selected from $-NO_2$ and halogen;


A^1 and A^2 are independently absent or are independently selected from




wherein the bond on the left side of A^1 , as drawn, is bound to $-C(=Z^1)-$ or L^2 ; and
wherein the bond on the right side of the A^2 moiety, as drawn, is bound to B or L^3 ;




each B is independently selected from



each B¹ is independently selected from - NR^3 -(C(R³)₂)_n-, - NR^3 -(C(R³)₂)_n-(C₆-C₁₀)arylene-(C(R³)₂)_n-, - NR^3 -(C(R³)₂)_n-heteroarylene-, - NR^3 -(C(R³)₂)_n-heteroarylene-(C(R³)₂)_n-, - NR^3 -(C(R³)₂)_n-NR³C(O)-, - NR^3 -(C(R³)₂)_n-heteroarylene-heterocyclylene-(C₆-C₁₀)arylene-, -heteroarylene-heterocyclylene-(C₆-





each R^3 is independently H or $(\text{C}_1\text{--C}_6)\text{alkyl}$;

each R^4 is independently H, $(\text{C}_1\text{--C}_6)\text{alkyl}$, halogen, 5-12 membered heteroaryl, 5-12 membered heterocyclyl, $(\text{C}_6\text{--C}_{10})\text{aryl}$, wherein the heteroaryl, heterocyclyl, and aryl are optionally substituted with $\text{---N}(\text{R}^3)_2$, ---OR^3 , halogen, $(\text{C}_1\text{--C}_6)\text{alkyl}$, $\text{---}(\text{C}_1\text{--C}_6)\text{alkylene--heteroaryl}$, $\text{---}(\text{C}_1\text{--C}_6)\text{alkylene--CN}$, $\text{---C}(\text{O})\text{NR}^3\text{---heteroaryl}$, or $\text{---C}(\text{O})\text{NR}^3\text{---heterocyclyl}$;

each R^5 is independently H, $(\text{C}_1\text{--C}_6)\text{alkyl}$, $\text{---C}(\text{O})\text{OR}^3$, or $\text{---N}(\text{R}^3)_2$, wherein the alkyl of $(\text{C}_1\text{--C}_6)\text{alkyl}$ is optionally substituted with $\text{---N}(\text{R}^3)_2$ or ---OR^3 ;

each R^6 is independently H, $(\text{C}_1\text{--C}_6)\text{alkyl}$, $\text{---C}(\text{O})\text{OR}^3$, or $\text{---N}(\text{R}^3)_2$, wherein the alkyl of $(\text{C}_1\text{--C}_6)\text{alkyl}$ is optionally substituted with $\text{---N}(\text{R}^3)_2$ or ---OR^3 ;

each R^7 is independently H, $(\text{C}_1\text{--C}_6)\text{alkyl}$, $\text{---C}(\text{O})\text{OR}^3$, or $\text{---N}(\text{R}^3)_2$, wherein the alkyl of $(\text{C}_1\text{--C}_6)\text{alkyl}$ is optionally substituted with $\text{---N}(\text{R}^3)_2$ or ---OR^3 ;

each R^8 is independently H, $(\text{C}_1\text{--C}_6)\text{alkyl}$, $\text{---C}(\text{O})\text{OR}^3$, or $\text{---N}(\text{R}^3)_2$, wherein the alkyl of $(\text{C}_1\text{--C}_6)\text{alkyl}$ is optionally substituted with $\text{---N}(\text{R}^3)_2$ or ---OR^3 ;

each Y is independently $\text{C}(\text{R}^3)_2$ or a bond;

each n is independently an integer from one to 12;


each o is independently an integer from zero to 30;

each p is independently an integer from zero to 12;

each q is independently an integer from zero to 30; and

each r is independently an integer from one to 6.

[00358] Embodiment II-1A. A compound of Formula Ia:



(Ia)

or a pharmaceutically acceptable salt or tautomer thereof, wherein:

R^{32} is $-H$, $=O$, $-OR^3$, $-N_3$, or $-O-C(=Z^1)-R^{32a}$,

R^{28} is $-H$, $(C_1-C_6)alkyl$, or $-C(=Z^1)-R^{28a}$,

R^{40} is $-H$ or $-C(=Z^1)-R^{40a}$,

wherein when R^{28} and R^{40} are H , then R^{32} is not $=O$;

each Z^1 is independently O or S ;

R^{28a} , R^{32a} , and R^{40a} are independently $-A^1-L^1-A^2-B$; $-A^1-A^2-B$; $-L^2-A^1-L^1-A^2-L^3-B$; $-O-(C_1-C_6)alkyl$; or $-O-(C_6-C_{10})aryl$; wherein the aryl of $-O-(C_6-C_{10})aryl$ is unsubstituted or substituted with 1-5 substituents selected from $-NO_2$ and halogen;

A^1 and A^2 are independently absent or are independently selected from

wherein the bond on the left side of A¹, as drawn, is bound to -C(=Z¹)- or L²; and
 wherein the bond on the right side of the A² moiety, as drawn, is bound to B or L³;

each Q is independently 1 to 3 rings selected from arylene, cycloalkylene, heteroarylene, and heterocyclylene;

each X is independently absent or 1 to 2 rings selected from arylene, cycloalkylene, heteroarylene, and heterocyclylene;

each X¹ is independently a heteroarylene or heterocyclylene ring;


each W is independently absent or 1 to 2 rings selected from arylene, cycloalkylene, heteroarylene, and heterocyclylene;

each W¹ is independently a heteroarylene or heterocyclylene ring;


each G is independently absent or a ring selected from arylene, cycloalkylene, heteroarylene, and heterocyclylene;

each G¹ and G² are independently heteroarylene or heterocyclylene ring;


each L¹ is independently selected from




L^2 and L^3 are independently absent or are independently selected from




each B is independently selected from



B^1 is selected from $^{-}\text{NR}^3\text{-(C(R}^3\text{)}_2)_n$, $^{-}\text{NR}^3\text{-(C(R}^3\text{)}_2)_n\text{-(C}_6\text{-C}_{10}\text{)arylene-(C(R}^3\text{)}_2)_n$, $^{-}\text{NR}^3\text{-(C(R}^3\text{)}_2)_n\text{-heteroarylene-}$, $^{-}\text{(C}_6\text{-C}_{10}\text{)arylene-}$, $^{-}\text{NR}^3\text{-(C(R}^3\text{)}_2)_n\text{-NR}^3\text{C(O)-}$, $^{-}\text{NR}^3\text{-(C(R}^3\text{)}_2)_n\text{-heteroarylene-heterocyclylene-(C}_6\text{-C}_{10}\text{)arylene-}$,





$^{-3}-NR^3-(C(R^3)_2)_n-S(O)_2-arylene-C(O)-$, wherein the $^{-3}-$ bond on the left side of B^1 , as drawn, is bound to A^2 , L^3 , or L^1 ; and wherein the heteroarylene, heterocyclylene, and arylene are optionally substituted with alkyl, hydroxyalkyl, haloalkyl, alkoxy, halogen, or hydroxyl;

each R^3 is independently H or (C_1-C_6) alkyl;

each R^4 is independently H, (C_1-C_6) alkyl, halogen, 5-12 membered heteroaryl, 5-12 membered heterocyclyl, (C_6-C_{10}) aryl, wherein the heteroaryl, heterocyclyl, and aryl are optionally substituted with $-N(R^3)_2$, $-OR^3$, halogen, (C_1-C_6) alkyl, $-(C_1-C_6)$ alkylene-heteroaryl, $-(C_1-C_6)$ alkylene-CN, $-C(O)NR^3$ -heteroaryl, or $-C(O)NR^3$ -heterocyclyl;

each R^5 is independently H, (C_1-C_6) alkyl, $-C(O)OR^3$, or $-N(R^3)_2$, wherein the alkyl of (C_1-C_6) alkyl is optionally substituted with $-N(R^3)_2$ or $-OR^3$;

each R^6 is independently H, (C_1-C_6) alkyl, $-C(O)OR^3$, or $-N(R^3)_2$, wherein the alkyl of (C_1-C_6) alkyl is optionally substituted with $-N(R^3)_2$ or $-OR^3$;

each R^7 is independently H, (C_1-C_6) alkyl, $-C(O)OR^3$, or $-N(R^3)_2$, wherein the alkyl of (C_1-C_6) alkyl is optionally substituted with $-N(R^3)_2$ or $-OR^3$;

each R^8 is independently H, (C_1-C_6) alkyl, $-C(O)OR^3$, or $-N(R^3)_2$, wherein the alkyl of (C_1-C_6) alkyl is optionally substituted with $-N(R^3)_2$ or $-OR^3$;

each Y is independently $C(R^3)_2$ or a bond;

each n is independently an integer from one to 12;

each o is independently an integer from zero to 30;

each p is independently an integer from zero to 12;


each q is independently an integer from zero to 30; and

each r is independently an integer from one to 6.

[00359] Embodiment II-2. The compound of Embodiment II-1, wherein R^{32} is =O.

[00360] Embodiment II-3. The compound of Embodiment II-1, wherein R^{32} is $-OR^3$.

[00361] Embodiment II-4. The compound of any one of Embodiments II-1 to II-3, or a pharmaceutically acceptable salt or tautomer thereof, wherein the compound is represented by the structure of Formula (I-40b):




(I-40b)

wherein R^{40} is $-C(=Z^1)-R^{40a}$.

- [00362] Embodiment II-5. The compound of Embodiment II-4, wherein Z^1 is O.
- [00363] Embodiment II-6. The compound of Embodiment II-4, wherein Z^1 is S.
- [00364] Embodiment II-7. The compound of any one of Embodiments II-4 to II-6, wherein R^{40a} is $-A^1-L^1-A^2-B$, wherein A^1 and A^2 are absent.
- [00365] Embodiment II-8. The compound of any one of Embodiments II-4 to II-6, wherein R^{40a} is $-A^1-L^1-A^2-B$, wherein A^2 is absent.
- [00366] Embodiment II-9. The compound of any one of Embodiments II-4 to II-6, wherein R^{40a} is $-A^1-L^1-A^2-B$, wherein A^1 is absent.
- [00367] Embodiment II-10. The compound of any one of Embodiments II-4 to II-6, wherein R^{40a} is $-A^1-L^1-A^2-B$.
- [00368] Embodiment II-11. The compound of any one of Embodiments II-4 to II-6, wherein R^{40a} is $-A^1-A^2-B$.
- [00369] Embodiment II-12. The compound of any one of Embodiments II-4 to II-6, wherein R^{40a} is $-L^2-A^1-L^1-A^2-L^3-B$, wherein L^2 and A^1 are absent.
- [00370] Embodiment II-13. The compound of any one of Embodiments II-4 to II-6, wherein R^{40a} is $-L^2-A^1-L^1-A^2-L^3-B$, wherein L^2 is absent.
- [00371] Embodiment II-14. The compound of any one of Embodiments II-4 to II-6, wherein R^{40a} is $-L^2-A^1-L^1-A^2-L^3-B$, wherein L^3 is absent.

[00372] Embodiment II-15. The compound of any one of Embodiments II-4 to II-6, wherein R^{40a} is $-O-(C_1-C_6)alkyl$ or $-O-(C_6-C_{10})aryl$; wherein the aryl of $-O-(C_6-C_{10})aryl$ is unsubstituted or substituted with 1-5 substituents selected from $-NO_2$ and halogen.

[00373] Embodiment II-16. The compound of any one of Embodiments II-1 to II-3, or a pharmaceutically acceptable salt or tautomer thereof, wherein the compounds are represented by the structure of Formula (I-28b):



wherein R^{28} is $-C(=Z^1)-R^{28a}$.

[00374] Embodiment II-17. The compound of Embodiment II-16, wherein Z^1 is O.

[00375] Embodiment II-18. The compound of Embodiment II-16, wherein Z^1 is S.

[00376] Embodiment II-19. The compound of any one of Embodiments II-16 to II-18, wherein R^{28a} is $-A^1-L^1-A^2-B$, wherein A^1 and A^2 are absent.

[00377] Embodiment II-20. The compound of any one of Embodiments II-16 to II-18, wherein R^{28a} is $-A^1-L^1-A^2-B$, wherein A^2 is absent.

[00378] Embodiment II-21. The compound of any one of Embodiments II-16 to II-18, wherein R^{28a} is $-A^1-L^1-A^2-B$, wherein A^1 is absent.

[00379] Embodiment II-22. The compound of any one of Embodiments II-16 to II-18, wherein R^{28a} is $-A^1-L^1-A^2-B$.

[00380] Embodiment II-23. The compound of any one of Embodiments II-16 to II-18, wherein R^{28a} is $-A^1-A^2-B$.


[00381] Embodiment II-24. The compound of any one of Embodiments II-16 to II-18, wherein R^{28a} is $-L^2-A^1-L^1-A^2-L^3-B$, wherein L^2 and A^1 are absent.

[00382] Embodiment II-25. The compound of any one of Embodiments II-16 to II-18, wherein R^{28a} is $-L^2-A^1-L^1-A^2-L^3-B$, wherein L^2 is absent.

[00383] Embodiment II-26. The compound of any one of Embodiments II-16 to II-18, wherein R^{28a} is $-L^2-A^1-L^1-A^2-L^3-B$, wherein L^3 is absent.

[00384] Embodiment II-27. The compound of any one of Embodiments II-16 to II-18, wherein R^{28a} is $-O-(C_1-C_6)alkyl$ or $-O-(C_6-C_{10})aryl$; wherein the aryl of $-O-(C_6-C_{10})aryl$ is unsubstituted or substituted with 1-5 substituents selected from $-NO_2$ and halogen.

[00385] Embodiment II-28. The compound of Embodiment II-1, or a pharmaceutically acceptable salt or tautomer thereof, wherein the compound is represented by the structure of Formula (I-32b):



wherein R^{32} is $-O-C(=Z^1)-R^{32a}$.

[00386] Embodiment II-29. The compound of Embodiment II-28, wherein Z^1 is O.

[00387] Embodiment II-30. The compound of Embodiment II-28, wherein Z^1 is S.

[00388] Embodiment II-31. The compound of any one of Embodiments II-28 to II-30, wherein R^{32a} is $-A^1-L^1-A^2-B$, wherein A^1 and A^2 are absent.

[00389] Embodiment II-32. The compound of any one of Embodiments II-28 to II-30, wherein R^{32a} is $-A^1-L^1-A^2-B$, wherein A^2 is absent.

[00390] Embodiment II-33. The compound of any one of Embodiments II-28 to II-30, wherein R^{32a} is $-A^1-L^1-A^2-B$, wherein A^1 is absent.

[00391] Embodiment II-34. The compound of any one of Embodiments II-28 to II-30, wherein R^{32a} is $-A^1-L^1-A^2-B$.

[00392] Embodiment II-35. The compound of any one of Embodiments II-28 to II-30, wherein R^{32a} is $-A^1-A^2-B$.


[00393] Embodiment II-36. The compound of any one of Embodiments II-28 to II-30, wherein R^{32a} is $-L^2-A^1-L^1-A^2-L^3-B$, wherein L^2 and A^1 are absent.

[00394] Embodiment II-37. The compound of any one of Embodiments II-28 to II-30, wherein R^{32a} is $-L^2-A^1-L^1-A^2-L^3-B$, wherein L^2 is absent.

[00395] Embodiment II-38. The compound of any one of Embodiments II-28 to II-30, wherein R^{32a} is $-L^2-A^1-L^1-A^2-L^3-B$, wherein L^3 is absent.


[00396] Embodiment II-39. The compound of any one of Embodiments II-28 to II-30, wherein R^{32a} is $-O-(C_1-C_6)alkyl$ or $-O-(C_6-C_{10})aryl$; wherein the aryl of $-O-(C_6-C_{10})aryl$ is unsubstituted or substituted with 1-5 substituents selected from $-NO_2$ and halogen.

[00397] Embodiment II-40. The compound of any one of Embodiments II-1 to II-




10, II-12 to II-22, II-24 to II-35, and II-36 to II-39, wherein L^1 is

[00398] Embodiment II-41. The compound of any one Embodiments II-1 to II-10,




II-12 to II-22, II-24 to II-35, and II-36 to II-39, wherein L^1 is

[00399] Embodiment II-42. The compound of any one of Embodiments II-1 to II-10, II-12 to II-22, 24-35, and 36-39, wherein L^1 is




[00400] Embodiment II-43. The compound of any one of Embodiments II-1 to II-




10, II-12 to II-22, II-24 to II-35, and II-36 to II-39, wherein L^1 is

[00401] Embodiment II-44. The compound of any one of Embodiments II-1 to II-




10, II-12 to II-22, II-24 to II-35, and II-36 to II-39, wherein L^1 is

[00409] Embodiment II-52. The compound of any one of Embodiments II-1 to II-6, II-8, II-10 to II-11, II-13 to II-18, II-20, II-22 to II-23, II-25 to II-30, II-32, II-34 to II-35, and




II-37 to II-45, wherein A¹ is

[00410] Embodiment II-53. The compound of any one of Embodiments II-1 to II-6, II-8, II-10 to II-11, II-13 to II-18, II-20, II-22 to II-23, II-25 to II-30, II-32, II-34 to II-35, and




II-37 to II-45, wherein A¹ is

[00411] Embodiment II-54. The compound of any one of Embodiments II-1 to II-6, II-8, II-10 to II-11, II-13 to II-18, II-20, II-22 to II-23, II-25 to II-30, II-32, II-34 to II-35, and




II-37 to II-45, wherein A¹ is

[00412] Embodiment II-55. The compound of any one of Embodiments II-1 to II-6, II-8, II-10 to II-11, II-13 to II-18, II-20, II-22 to II-23, II-25 to II-30, II-32, II-34 to II-35, and



II-37 to II-45, wherein A¹ is.


[00413] Embodiment II-56. The compound of any one of Embodiments II-1 to II-6, II-8, II-10 to II-11, II-13 to II-18, II-20, II-22 to II-23, II-25 to II-30, II-32, II-34 to II-35, and



II-37 to II-45, wherein A¹ is


[00414] Embodiment II-57. The compound of any one of Embodiments II-1 to II-8, II-15 to II-20, II-27 to II-32, and II-39 to II-45, wherein A² is absent.

[00415] Embodiment II-58. The compound of any one of Embodiments II-1 to II-6,




II-9 to II-18, II-21 to II-30, and II-33 to II-45, wherein A² is

[00416] Embodiment II-59. The compound of any one of Embodiments II-1 to II-6,




II-9 to II-18, II-21 to II-30, and II-33 to II-45, wherein A² is

[00417] Embodiment II-60. The compound of any one of Embodiments II-1 to II-6,




II-9 to II-18, II-21 to II-30, and II-33 to II-45, wherein A² is

[00418] Embodiment II-61. The compound of any one of Embodiments II-1 to II-6,




II-9 to II-18, II-21 to II-30, and II-33 to II-45, wherein A² is

[00419] Embodiment II-62. The compound of any one of Embodiments II-1 to II-6,




II-9 to II-18, II-21 to II-30, and II-33 to II-45, wherein A² is

[00420] Embodiment II-63. The compound of any one of Embodiments II-1 to II-6,




II-9 to II-18, II-21 to II-30, and II-33 to II-45, wherein A² is

[00421] Embodiment II-64. The compound of any one of Embodiments II-1 to II-6,




II-9 to II-18, II-21 to II-30, and II-33 to II-45, wherein A² is.

[00422] Embodiment II-65. The compound of any one of Embodiments II-1 to II-6,




II-9 to II-18, II-21 to II-30, and II-33 to II-45, wherein A² is

[00423] Embodiment II-66. The compound of any one of Embodiments II-1 to II-



65, wherein B is

[00424] Embodiment II-67. The compound of any one of Embodiments II-1 to II-




65, wherein B is

[00425] Embodiment II-68. The compound of any one of Embodiments II-1 to II-

65, wherein B¹ is $\text{NR}^3\text{-(C(R}^3\text{))}_n\text{-}$.

[00426] Embodiment II-69. The compound of any one of Embodiments II-1 to II-




65, wherein B¹ is

[00427] Embodiment II-70. The compound of any one of Embodiments II-1 to II-


69, wherein R⁴ is 5-12 membered heteroaryl, optionally substituted with -N(R³)₂, -OR³, halogen, (C₁-C₆)alkyl, -(C₁-C₆)alkylene-heteroaryl, -(C₁-C₆)alkylene-CN, or -C(O)NR³-heteroaryl.

[00428] Embodiment II-71. The compound of any one of Embodiments II-1 to II-


70, or a pharmaceutically acceptable salt or tautomer thereof, wherein compound has the following formula:




[00429] Embodiment II-72. A compound selected from the group consisting of:




Example 1




Example 2




Example 3




Example 4




Example 5




Example 6




Example 7




Example 8




Example 9




Example 10




Example 11




Example 12




Example 13




Example 14




Example 15




Example 16




Example 17




Example 18




Example 19




Example 20




Example 21




Example 22




Example 23




Example 24




Example 25




Example 26




Example 27




Example 28




Example 29




Example 30




Example 31




Example 32




Example 33




Example 34




Example 35




Example 36




Example 37




Example 38




Example 39




Example 40




Example 41




Example 42




Example 43




Example 44




Example 45




Example 46




Example 47




Example 48




Example 49




Example 50




Example 51




Example 52




Example 53




Example 54




Example 55




Example 56




Example 57




Example 58




Example 59




Example 60




Example 61




Example 62




Example 63




Example 64




Example 65




Example 66




Example 67



Example 68








Example 69




or a pharmaceutically acceptable salt or tautomer thereof.

[00430] Embodiment II-73. A compound selected from the group consisting of:







Example 75




Example 76




Example 77




Example 78




Example 79




Example 80




Example 81




Example 82




Example 83




Example 84




Example 85




Example 86




Example 87




Example 88




Example 89




Example 90




Example 91




Example 92




Example 93




Example 94




Example 95




Example 96




Example 97




Example 98




Example 99




Example 100




Example 101




Example 102




Example 103




Example 104




Example 105




Example 106




Example 107




Example 108




Example 109




Example 110




Example 111




Example 112




Example 113




Example 114




Example 115




Example 116




Example 117




Example 118




Example 119




Example 120




Example 121




Example 122




Example 123




Example 124




Example 125




Example 126




Example 127




Example 128




Example 129




Example 130




Example 131




Example 132



Example 133




Example 134




Example 135

or a pharmaceutically acceptable salt or tautomer thereof.


[00431] Embodiment II-74. A compound selected from the group consisting of:




Example 136




Example 137




Example 138




Example 139




Example 140




Example 141




Example 142




Example 143




Example 144




Example 145




Example 146




Example 147




Example 148




Example 149



Example 150



Example 151



Example 152

or a pharmaceutically acceptable salt or tautomer thereof.

[00432] Embodiment II-75. A pharmaceutical composition comprising a compound of any one of Embodiments II-1 to II-74, or a pharmaceutically acceptable salt thereof, and at least one of a pharmaceutically acceptable carrier, diluent, or excipient.

[00433] Embodiment II-76. A method of treating a disease or disorder mediated by mTOR comprising administering to the subject suffering from or susceptible to developing a disease or disorder mediated by mTOR a therapeutically effective amount of one or more

compounds of any one of Embodiments II-1 to II-74, or a pharmaceutically acceptable salt thereof.

[00434] Embodiment II-77. A method of preventing a disease or disorder mediated by mTOR comprising administering to the subject suffering from or susceptible to developing a disease or disorder mediated by mTOR a therapeutically effective amount of one or more compounds of any one of Embodiments II-1 to II-74, or a pharmaceutically acceptable salt thereof.

[00435] Embodiment II-78. A method of reducing the risk of a disease or disorder mediated by mTOR comprising administering to the subject suffering from or susceptible to developing a disease or disorder mediated by mTOR a therapeutically effective amount of one or more compounds of any one of Embodiments II-1 to II-74, or a pharmaceutically acceptable salt thereof.

[00436] Embodiment II-79. The method of any one of Embodiments II-76 to II-78, wherein the disease is cancer or an immune-mediated disease.

[00437] Embodiment II-80. The method of Embodiment II-79, wherein the cancer is selected from brain and neurovascular tumors, head and neck cancers, breast cancer, lung cancer, mesothelioma, lymphoid cancer, stomach cancer, kidney cancer, renal carcinoma, liver cancer, ovarian cancer, ovary endometriosis, testicular cancer, gastrointestinal cancer, prostate cancer, glioblastoma, skin cancer, melanoma, neuro cancers, spleen cancers, pancreatic cancers, blood proliferative disorders, lymphoma, leukemia, endometrial cancer, cervical cancer, vulva cancer, prostate cancer, penile cancer, bone cancers, muscle cancers, soft tissue cancers, intestinal or rectal cancer, anal cancer, bladder cancer, bile duct cancer, ocular cancer, gastrointestinal stromal tumors, and neuro-endocrine tumors.

[00438] Embodiment II-81. The method of Embodiment II-79, wherein the immune-mediated disease is selected from resistance by transplantation of heart, kidney, liver, medulla ossium, skin, cornea, lung, pancreas, intestinum tenue, limb, muscle, nerves, duodenum, small-bowel, or pancreatic-islet-cell; graft-versus-host diseases brought about by medulla ossium transplantation; rheumatoid arthritis, systemic lupus erythematosus, Hashimoto's thyroiditis, multiple sclerosis, myasthenia gravis, type I diabetes, uveitis, allergic encephalomyelitis, and glomerulonephritis.

[00439] Embodiment II-82. A method of treating cancer comprising administering to the subject a therapeutically effective amount of one or more compounds of any one of Embodiments II-1 to II-74, or a pharmaceutically acceptable salt thereof.

[00440] Embodiment II-83. The method of Embodiment II-82, wherein the cancer is selected from brain and neurovascular tumors, head and neck cancers, breast cancer, lung cancer, mesothelioma, lymphoid cancer, stomach cancer, kidney cancer, renal carcinoma, liver cancer, ovarian cancer, ovary endometriosis, testicular cancer, gastrointestinal cancer, prostate cancer, glioblastoma, skin cancer, melanoma, neuro cancers, spleen cancers, pancreatic cancers, blood proliferative disorders, lymphoma, leukemia, endometrial cancer, cervical cancer, vulva cancer, prostate cancer, penile cancer, bone cancers, muscle cancers, soft tissue cancers, intestinal or rectal cancer, anal cancer, bladder cancer, bile duct cancer, ocular cancer, gastrointestinal stromal tumors, and neuro-endocrine tumors.

[00441] Embodiment II-84. A method of treating an immune-mediated disease comprising administering to the subject a therapeutically effective amount of one or more compounds of any one of Embodiments II-1 to II-74, or a pharmaceutically acceptable salt thereof.

[00442] Embodiment II-85. The method of Embodiment II-84, wherein the immune-mediated disease is selected from resistance by transplantation of heart, kidney, liver, medulla ossium, skin, cornea, lung, pancreas, intestinum tenue, limb, muscle, nerves, duodenum, small-bowel, or pancreatic-islet-cell; graft-versus-host diseases brought about by medulla ossium transplantation; rheumatoid arthritis, systemic lupus erythematosus, Hashimoto's thyroiditis, multiple sclerosis, myasthenia gravis, type I diabetes, uveitis, allergic encephalomyelitis, and glomerulonephritis.

[00443] Embodiment II-86. A method of treating an age related condition comprising administering to the subject a therapeutically effective amount of one or more compounds of any one of Embodiments II-1 to II-74, or a pharmaceutically acceptable salt thereof.

[00444] Embodiment II-87. The method of Embodiment II-86, wherein the age related condition is selected from sarcopenia, skin atrophy, muscle wasting, brain atrophy, atherosclerosis, arteriosclerosis, pulmonary emphysema, osteoporosis, osteoarthritis, high blood pressure, erectile dysfunction, dementia, Huntington's disease, Alzheimer's disease, cataracts, age-related macular degeneration, prostate cancer, stroke, diminished life

expectancy, impaired kidney function, and age-related hearing loss, aging-related mobility disability (e.g., frailty), cognitive decline, age-related dementia, memory impairment, tendon stiffness, heart dysfunction such as cardiac hypertrophy and systolic and diastolic dysfunction, immunosenescence, cancer, obesity, and diabetes.

[00445] Embodiment II-88. A compound of any one of Embodiments II-1 to II-74, or a pharmaceutically acceptable salt thereof, for use in treating, preventing, or reducing the risk of a disease or condition mediated by mTOR.

[00446] Embodiment II-89. Use of a compound of any of Embodiments II-1 to II-74, or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament for treating, preventing, or reducing the risk of a disease or disorder mediated by mTOR.

[00447] Embodiment II-90. A compound of any one of claims 1-74, or a pharmaceutically acceptable salt thereof, for use in treating cancer.

[00448] Embodiment II-91. Use of a compound of any one of Embodiments II-1 to II-74, or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament for treating cancer.

[00449] Embodiment II-92. A compound of any one of Embodiments II-1 to II-74, or a pharmaceutically acceptable salt thereof, for use in treating an immune-mediated disease.

[00450] Embodiment II-93. Use of a compound of any one of Embodiments II-1 to II-74, or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament for treating an immune-mediated disease.

[00451] Embodiment II-94. A compound of any one of Embodiments II-1 to II-74, or a pharmaceutically acceptable salt thereof, for use in treating an age related condition.

[00452] Embodiment II-95. Use of a compound of any one of Embodiments II-1 to II-74, or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament for treating an age related condition.

Examples

[00453] The disclosure is further illustrated by the following examples and synthesis examples, which are not to be construed as limiting this disclosure in scope or spirit to the specific procedures herein described. It is to be understood that the examples are provided to illustrate certain embodiments and that no limitation to the scope of the disclosure is intended thereby. It is to be further understood that resort may be had to various other embodiments,

modifications, and equivalents thereof which may suggest themselves to those skilled in the art without departing from the spirit of the present disclosure and/or scope of the appended claims.


[00454] Definitions used in the following examples and elsewhere herein are:

CH ₂ Cl ₂ , DCM	Methylene chloride, Dichloromethane
CH ₃ CN, MeCN	Acetonitrile
DIPEA	Diisopropylethyl amine or Hunig's base
DMA	Dimethylacetamide
DME	Dimethoxyethane
DMF	N,N-Dimethylformamide
EDCI	1-Ethyl-3-(3-dimethylaminopropyl)carbodiimide
EtOAc	Ethyl acetate
h	hour
H ₂ O	Water
HCl	Hydrochloric acid
HOBt	Hydroxybenzotriazole
HPLC	High-performance liquid chromatography
LCMS	Liquid chromatography–mass spectrometry
MeOH	Methanol
MTBE	Methyl <i>tert</i> -butyl ether
Na ₂ SO ₄	Sodium sulfate
PEG	Polyethylene glycol
TBDMS	<i>tert</i> -butyldimethylsilyl
TFA	Trifluoroacetic acid
THF	Tetrahydrofuran
TMS	Tetramethylsilane

Series 1 bifunctional rapalogs

[00455] A general structure of Series 1 bifunctional rapalogs is shown in Scheme 1 below. For these types of bifunctional rapalogs, the linker may include variations where q = 0 to 30, such as q = 1 to 7, and r = 1 to 6. The linker amine can include substitutions, such as R = H and C1–C6 alkyl groups. The carbamate moiety, where Z¹ = O or S, can be attached to the rapalog at R⁴⁰ or R²⁸ (Formula I and II), including variations found in Table 1 in the Examples Section. An mTOR active site inhibitor can attach to the linker via a primary or secondary amine, and may include variations found in Table 2 in the Examples Section.


Scheme 1. Series 1 bifunctional rapalogs.



Series 2 bifunctional rapalogs

[00456] A general structure of Series 2 bifunctional rapalogs is shown in Scheme 2 below. For these types of bifunctional rapalogs, the linker may include variations where $q = 0$ to 30, such as $q = 1$ to 7. The linker amine can include substitutions, such as $R = H$ and C1–C6 alkyl groups. The pre-linker amine can include substitutions, such as $R^2 = H$, C1–C6 alkyl groups, and cycloalkyl including 4 to 8-membered rings. The carbamate moiety, where $Z^1 = O$ or S, can be attached to the rapalog at R^{40} or R^{28} (Formula I and II), including variations found in Table 1 in the Examples Section. An mTOR active site inhibitor can attach to the linker via a primary or secondary amine, and may include variations found in Table 2 in the Examples Section.


Scheme 2. Series 2 bifunctional rapalogs.



Series 3 bifunctional rapalogs


[00457] A general structure of Series 3 bifunctional rapalogs is shown in Scheme 3 below. For these types of bifunctional rapalogs, the linker may include variations where $q = 0$ to 30, such as $q = 1$ to 7. The linker amine can include substitutions, such as $R = H$ and C1–C6 alkyl groups. The post-linker amine can include substitutions, such as $R^2 = H$, C1–C6 alkyl groups, and cycloalkyl including 4 to 8-membered rings. The carbamate moiety, where $Z^1 = O$ or S, can be attached to the rapalog at R^{40} or R^{28} (Formula I and II), including variations found in Table 1 in the Examples Section. An mTOR active site inhibitor can attach to the linker via a primary or secondary amine, and may include variations found in Table 2 in the Examples Section.

Scheme 3. Series 3 bifunctional rapalogs

**Series 4 bifunctional rapalogs**


[00458] A general structure of Series 4 bifunctional rapalogs is shown in Scheme 4 below. For these types of bifunctional rapalogs, the linker may include variations where $q = 0$ to 30, such as $q = 1$ to 7. The linker amine can include substitutions, such as $R = H$ and C1–C6 alkyl groups. The pre- and post-linker amines can each include substitutions, such as $R^2 = H$, C1–C6 alkyl groups, and cycloalkyl including 4 to 8-membered rings. The carbamate moiety, where $Z^1 = O$ or S, can be attached to the rapalog at R^{40} or R^{28} (Formula I and II), including variations found in Table 1 in the Examples Section. An mTOR active site inhibitor can attach to the linker via a primary or secondary amine, and may include variations found in Table 2 in the Examples Section.

Scheme 4. Series 4 bifunctional rapalogs

**Series 5 bifunctional rapalogs**


[00459] A general structure of Series 5 bifunctional rapalogs is shown in Scheme 5 below. For these types of bifunctional rapalogs, the pre-linker amine can include substitutions, such as $R^2 = H$, C1–C6 alkyl groups, and cycloalkyl including 4 to 8-membered rings. The carbamate moiety, where $Z^1 = O$ or S, can be attached to the rapalog at R^{40} or R^{28} (Formula I and II), including variations found in Table 1 in the Examples Section. An mTOR active site inhibitor can attach to the linker via a primary or secondary amine, and may include variations found in Table 2 in the Examples Section.

Scheme 5. Series 5 bifunctional rapalogs

**Series 6 bifunctional rapalogs**

[00460] A general structure of Series 6 bifunctional rapalogs is shown in Scheme 6 below. For these types of bifunctional rapalogs, the linker may include variations where $q = 0$ to 30, such as $q = 1$ to 7. The linker amines can include substitutions, such as $R = H$ and C1–C6 alkyl groups. The post-linker amine can include substitutions, such as $R^2 = H$, C1–C6 alkyl groups, and cycloalkyl including 4 to 8-membered rings. The carbamate moiety, where $Z^1 = O$ or S, can be attached to the rapalog at R^{40} or R^{28} (Formula I and II), including variations found in Table 1 in the Examples Section. An mTOR active site inhibitor can attach to the linker via a primary or secondary amine, and may include variations found in Table 2 in the Examples Section.


Scheme 6. Series 6 bifunctional rapalogs.

**Series 7 bifunctional rapalogs**

[00461] A general structure of Series 7 bifunctional rapalogs is shown in Scheme 7 below. For these types of bifunctional rapalogs, the linker may include variations where $q = 0$ to 30, such as $q = 1$ to 7. The linker amine can include substitutions, such as $R = H$ and C1–C6 alkyl groups. The pre- and post-linker amines can each include substitutions, such as $R^2 = H$, C1–C6 alkyl groups, and cycloalkyl including 4 to 8-membered rings. The carbamate moiety, where $Z^1 = O$ or S, can be attached to the rapalog at R^{40} or R^{28} (Formula I or II), including variations found in Table 1 in the Examples Section. An mTOR active site inhibitor can

attach to the linker via a primary or secondary amine, and may include variations found in Table 2 in the Examples Section.

Scheme 7. Series 7 bifunctional rapalogs



Series 8 bifunctional rapalogs

[00462] A general structure of Series 8 bifunctional rapalogs is shown in Scheme 8 below. For these types of bifunctional rapalogs, the linker may include variations where $q = 0$ to 30, such as $q = 1$ to 7. The linker amine can include substitutions, such as $R = H$ and C1–C6 alkyl groups. The post-linker amine can include substitutions, such as $R^2 = H$, C1–C6 alkyl groups, and cycloalkyl including 4 to 8-membered rings. The carbamate moiety, where $Z^1 = O$ or S, can be attached to the rapalog at R^{40} or R^{28} (Formula I or II), including variations found in Table 1 in the Examples Section. An mTOR active site inhibitor can attach to the linker via a primary or secondary amine, and may include variations found in Table 2 in the Examples Section.

Scheme 8. Series 8 bifunctional rapalogs


















Table 1. Carbonate and thiocarbonate containing rapalog monomers.

Carbonate containing rapalog	Carbonate containing rapalog
 Monomer 1	 Monomer 2
 Monomer 3	 Monomer 4
 Monomer 5	 Monomer 6
 Monomer 7	 Monomer 8

Carbonate containing rapalog	Carbonate containing rapalog
 Monomer 9	 Monomer 10
 Monomer 11	 Monomer 12
 Monomer 13	 Monomer 14




























Carbonate containing rapalog	Carbonate containing rapalog
 Monomer 15	 Monomer 16
 Monomer 17	









Table 2. Active Site inhibitor monomers.

Active Site inhibitor monomers	Active Site inhibitor monomers
 Monomer A	 Monomer B

Active Site inhibitor monomers	Active Site inhibitor monomers
 Monomer C	 Monomer D
 Monomer E	 Monomer F
 Monomer G	 Monomer H
 Monomer I	 Monomer J

Active Site inhibitor monomers	Active Site inhibitor monomers
 Monomer K	 Monomer L
 Monomer M	 Monomer N
 Monomer O	 Monomer P
 Monomer Q	 Monomer R

Active Site inhibitor monomers	Active Site inhibitor monomers
 Monomer S	 Monomer T
 Monomer U	 Monomer V
 Monomer W	 Monomer X

Active Site inhibitor monomers	Active Site inhibitor monomers
 Monomer Y	 Monomer Z
 Monomer AA	 Monomer AB
 Monomer AC	 Monomer AD
 Monomer AE	 Monomer AF






Active Site inhibitor monomers	Active Site inhibitor monomers
 Monomer AG	 Monomer AH
 Monomer AI	

Table 3. Active Site inhibitor monomers

Active Site Inhibitor	Active Site Inhibitor
 Monomer AJ	 Monomer AK











































Active Site Inhibitor	Active Site Inhibitor
 Monomer AL	 Monomer AM
 Monomer AN	 Monomer AO















Table 4. Amine containing pre- and post-linkers.


Amide containing block	Amide containing block
 Building block A	 Building block B
 Building block C	 Building block D

 Building block E	 Building block F
 Building block G	 Building block H
 Building block I	 Building block J
 Building block K	 Building block L
 Building block M	 Building block N
 Building block O	 Building block P

 Building block Q	 Building block R
 Building block S	 Building block T
 Building block U	 Building block V
 Building block W	 Building block X
 Building block Y	 Building block Z


 Building block AA	 Building block AB
 Building block AC	 Building block AD
 Building block AE	 Building block AF
 Building block AG	 Building block AH
 Building block AI	 Building block AJ
 Building block AK	 Building block AL

 Building block AM	 Building block AN
 Building block AO	 Building block AP
 Building block AQ	 Building block AR
 Building block AS	 Building block AT
 Building block AU	 Building block AV
 Building block AW	 Building block AX
 Building block AY	 Building block AZ



Preparation of Active Site Inhibitor Monomers

Monomer A. 5-(4-amino-1-(4-(aminomethyl)benzyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)benzo[d]oxazol-2-amine trifluoroacetic acid salt.



Step 1: Synthesis of *tert*-butyl 4-((4-amino-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl)benzylcarbamate

[00463] To a solution of 3-iodo-1H-pyrazolo[3,4-d]pyrimidin-4-amine (3.8 g, 14.56 mmol, 1.0 equiv) in DMF (20 mL) was added NaH (582.27 mg, 14.56 mmol, 60 wt.%, 1.0 equiv) at 0 °C and the reaction solution was stirred at this temperature for 30 min, then *tert*-butyl 4-(bromomethyl)benzylcarbamate (4.59 g, 15.29 mmol, 1.05 equiv) was added to the reaction at 0 °C and the reaction solution was stirred at room temperature for 2 h. The solution was poured into H₂O (80 mL) and the solid that precipitated out was filtered. The solid cake was washed with H₂O (2 x 10 mL) and then dried under reduced pressure to give *tert*-butyl 4-((4-

amino-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl)benzylcarbamate (5 g, 53% yield) as a yellow solid. LCMS (ESI) m/z : [M + Na] calcd for C₁₈H₂₁IN₆O₂: 503.07; found 503.2.


Step 2: Synthesis of tert-butyl 4-((4-amino-3-(2-aminobenzo[d]oxazol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl)benzylcarbamate

[00464] To a bi-phasic suspension of *tert*-butyl 4-((4-amino-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl)benzylcarbamate (5 g, 7.68 mmol, 1.0 equiv), 5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzo[d]oxazol-2-amine (2.40 g, 9.22 mmol, 1.2 equiv) and Pd(PPh₃)₄ (887.66 mg, 768.16 μmol, 0.1 equiv) in DME (100 mL) and H₂O (50 mL) was added Na₂CO₃ (1.91 g, 23.04 mmol, 3.0 equiv) at room temperature under N₂. The mixture was stirred at 110 °C for 3 h. The reaction mixture was cooled to room temperature and filtered, the filtrate was extracted by EtOAc (3 x 50 mL). The organic phases were combined and washed with brine (10 mL), dried over Na₂SO₄, filtered, and the filtrate was concentrated under reduced pressure to give a residue. The residue was purified by silica gel chromatography (0→20% MeOH/EtOAc) to give *tert*-butyl 4-((4-amino-3-(2-aminobenzo[d]oxazol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl)benzylcarbamate (4.5 g, 82% yield) as a yellow solid. LCMS (ESI) m/z : [M + H] calcd for C₂₅H₂₆N₈O₃: 487.22; found 487.2.

Step 3: Synthesis of 5-(4-amino-1-(4-(aminomethyl)benzyl)-1H-pyrazolo[3,4-d] pyrimidin-3-yl)benzo[d]oxazol-2-amine

[00465] To a solution of *tert*-butyl 4-((4-amino-3-(2-aminobenzo[d]oxazol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl)benzylcarbamate (4.5 g, 6.29 mmol, 1.0 equiv) in DCM (50 mL) was added TFA (30.80 g, 270.12 mmol, 20 mL, 42.95 equiv) at 0 °C. The reaction solution was stirred at room temperature for 2 h. The reaction solution was concentrated under reduced pressure to give a residue, which was dissolved in 10 mL of MeCN, then poured into MTBE (100 mL). The solid that precipitated was then filtered and the solid cake was dried under reduced pressure to give 5-[4-amino-1-[[4-(aminomethyl)phenyl]methyl]pyrazolo[3,4-d]pyrimidin-3-yl]-1,3-benzoxazol-2-amine (2.22 g, 71% yield, TFA) as a yellow solid. LCMS (ESI) m/z : [M + H] calcd for C₂₀H₁₈N₈O: 387.16; found 387.1.

Monomer B. 2-(4-amino-1-(4-aminobutyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-1H-indol-6-ol trifluoroacetic acid salt.



Step 1: Synthesis of tert-butyl N-(4-{4-amino-3-[6-(benzyloxy)-1H-indol-2-yl]-1H-pyrazolo[3,4-d]pyrimidin-1-yl}butyl)carbamate

[00466] To a mixture of *tert*-butyl (4-(4-amino-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-1-yl)butyl)carbamate (300 mg, 694 μ mol, 1.0 equiv) and (6-(benzyloxy)-1-(*tert*-butoxycarbonyl)-1H-indol-2-yl)boronic acid (763 mg, 2.08 mmol, 3.0 equiv) in DMF (2.6 mL), EtOH (525 μ L), and H₂O (350 μ L) were added Pd(OAc)₂ (15.5 mg, 69 μ mol, 0.1 equiv), triphenylphosphine (36.1 mg, 138 μ mol, 0.2 equiv), and sodium carbonate (440 mg, 4.16 mmol, 6.0 equiv). The reaction was heated at 80 °C for 20 h, cooled to room temperature, and quenched with H₂O (10 mL) and EtOAc (10 mL). The mixture was transferred to a separatory funnel and the aqueous phase was extracted with EtOAc (3 x 20 mL). The combined organic phase was washed with sat. aq. NaCl (1 x 20 mL), dried over Na₂SO₄, filtered, and concentrated under reduced pressure. The crude material was purified by silica gel chromatography (20→85% EtOAc/heptane) to provide the product (201 mg, 46% yield) as an orange solid. LCMS (ESI) *m/z*: [M + H] calcd for C₂₉H₃₃N₇O₃: 528.27; found 528.2.

Step 2: Synthesis of tert-butyl (4-(4-amino-3-(6-hydroxy-1H-indol-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)butyl)carbamate


[00467] To a solution of *tert*-butyl N-(4-{4-amino-3-[6-(benzyloxy)-1H-indol-2-yl]-1H-pyrazolo[3,4-d]pyrimidin-1-yl}butyl)carbamate (1.0 equiv) in EtOH is added Pd/C (10 mol%). The reaction is purged with H₂ and the reaction allowed to stir under an atmosphere of H₂ until consumption of starting material, as determined by LCMS. The reaction is then

diluted with EtOAc, filtered over Celite, and concentrated under reduced pressure. The resultant residue is purified by silica gel chromatography to afford the desired product.

Step 3: Synthesis of 2-(4-amino-1-(4-aminobutyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-1H-indol-6-ol

[00468] To a solution of *tert*-butyl (4-(4-amino-3-(6-hydroxy-1H-indol-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)butyl)carbamate (1.0 equiv) in anhydrous DCM is added TFA (50 equiv.) dropwise at 0 °C. The reaction is stirred at 0 °C and warmed to room temperature. Once the reaction is complete, as determined by LCMS, the reaction is concentrated under reduced pressure. The residue is triturated with MeCN, then dripped into MTBE over 10 min. The supernatant is removed and the precipitate is collected by filtration under N₂ to give 2-(4-amino-1-(4-aminobutyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-1H-indol-6-ol.

Monomer C. 5-(4-amino-1-((1,2,3,4-tetrahydroisoquinolin-6-yl)methyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)benzo[d]oxazol-2-amine trifluoroacetic acid salt.



Step 1: Synthesis of tert-butyl 6-((4-amino-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl)-3,4-dihydroisoquinoline-2(1H)-carboxylate

[00469] To a suspension of 3-iodo-1H-pyrazolo[3,4-d]pyrimidin-4-amine (5 g, 19.16 mmol, 1.0 equiv) in DMF (50.0 mL) was added NaH (766.22 mg, 19.16 mmol, 60 wt.%, 1.0 equiv) at 4 °C. The mixture was stirred at 4 °C for 30 min. To the reaction mixture was added *tert*-butyl 6-(bromomethyl)-3,4-dihydroisoquinoline-2(1H)-carboxylate (6.87 g, 21.07 mmol, 1.1 equiv) in DMF (30 mL) at 4 °C. The mixture was stirred at room temperature for 2 h. The mixture was then cooled to 4 °C and H₂O (400 mL) was added and the mixture was stirred for 30 min. The resulting precipitate was collected by filtration to give crude *tert*-butyl 6-((4-amino-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl)-3,4-dihydroisoquinoline-2(1H)-carboxylate (9.7 g, 76% yield) as a light yellow solid. The crude product was used for the next step directly.


Step 2: Synthesis of tert-butyl 6-((4-amino-3-(2-aminobenzo[d]oxazol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl)-3,4-dihydroisoquinoline-2(1H)-carboxylate

[00470] To a bi-phasic suspension of *tert*-butyl 6-((4-amino-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl)-3,4-dihydroisoquinoline-2(1H)-carboxylate (9.7 g, 14.63 mmol, 1.0 equiv), 5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzo[d]oxazol-2-amine (4.57 g, 17.55 mmol, 1.2 equiv), and Na₂CO₃ (7.75 g, 73.14 mmol, 5.0 equiv) in DME (120.0 mL) and H₂O (60 mL) was added Pd(PPh₃)₄ (1.69 g, 1.46 mmol, 0.1 equiv) at room temperature under N₂. The mixture was stirred at 110 °C for 3 h. The reaction mixture was then cooled to room temperature and partitioned between EtOAc (100 mL) and H₂O (100 mL). The aqueous layer was separated and extracted with EtOAc (2 x 60 mL). The organic layers were combined, washed with brine (80 mL) and dried over anhydrous Na₂SO₄, filtered and the filtrate was concentrated under reduced pressure. The residue was purified by silica gel chromatography (1→100% EtOAc/petroleum ether, then 20→50% MeOH/EtOAc) to afford *tert*-butyl 6-((4-amino-3-(2-aminobenzo[d]oxazol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl)-3,4-dihydroisoquinoline-2(1H)-carboxylate (4.5 g, 58% yield) as a light yellow solid.

Step 3: Synthesis of 5-(4-amino-1-((1,2,3,4-tetrahydroisoquinolin-6-yl)methyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)benzo[d]oxazol-2-amine

[00471] To neat TFA (32.5 mL, 438.97 mmol, 50.0 equiv) was added *tert*-butyl 6-((4-amino-3-(2-aminobenzo[d]oxazol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl)-3,4-dihydroisoquinoline-2(1H)-carboxylate (4.5 g, 8.78 mmol, 1.0 equiv) at room temperature. The mixture was stirred for 30 min and then concentrated under reduced pressure. The oily residue was triturated with MeCN (8 mL), then dripped into MTBE (350 mL) over 10 min. The supernatant was removed and then the precipitate was collected by filtration under N₂ to give 5-(4-amino-1-((1,2,3,4-tetrahydroisoquinolin-6-yl)methyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)benzo[d]oxazol-2-amine (5.72 g, over 100% yield, TFA) as a light pink solid. LCMS (ESI) *m/z*: [M + H]⁺ calcd for C₂₂H₂₀N₈O: 413.18; found 413.2.

Monomer D. 2-(4-amino-1-(4-aminobutyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-1H-indol-7-ol trifluoroacetic acid salt.



Step 1: Synthesis of *tert*-butyl 2-(4-amino-1-(4-((*tert*-butoxycarbonyl)amino)butyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-7-methoxy-1H-indole-1-carboxylate

[00472] To a mixture of *tert*-butyl (4-(4-amino-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-1-yl)butyl)carbamate (1.0 equiv) and (1-(*tert*-butoxycarbonyl)-7-methoxy-1H-indol-2-yl)boronic acid (3.0 equiv) in DME and H₂O is added Pd(PPh₃)₄ (0.1 equiv) and sodium carbonate (6.0 equiv). The reaction is heated at 80 °C until completion, as determined by LCMS and TLC analysis. The reaction is then quenched with H₂O and EtOAc. The mixture is transferred to a separatory funnel and the aqueous phase is extracted with EtOAc. The organic phase is washed with sat. aq. NaCl, dried over Na₂SO₄, filtered, and concentrated under reduced pressure. The desired product is isolated after chromatography on silica gel.

Step 2: Synthesis of *tert*-butyl 2-(4-amino-1-(4-((*tert*-butoxycarbonyl)amino)butyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-7-hydroxy-1H-indole-1-carboxylate


[00473] To a solution of *tert*-butyl 2-(4-amino-1-(4-((*tert*-butoxycarbonyl)amino)butyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-7-methoxy-1H-indole-1-carboxylate (1.0 equiv) in DCM at -10 °C is added BBr₃ (2.0 equiv). The reaction is allowed to stir until consumption of starting material, as determined by LCMS. The reaction is quenched by slow addition of sat. aq. NaHCO₃, transferred to a separatory funnel and the mixture is extracted with DCM. The organic phase is washed with sat. aq. NaCl, dried over Na₂SO₄, filtered, and concentrated under reduced pressure. The desired product is isolated after chromatography on silica gel.

Step 3: Synthesis of 2-(4-amino-1-(4-aminobutyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-1H-indol-7-ol

[00474] To a solution of *tert*-butyl 2-(4-amino-1-(4-((*tert*-butoxycarbonyl)amino)butyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-7-hydroxy-1H-indole-1-carboxylate (1.0 equiv) in DCM at 0 °C is added TFA dropwise. The reaction is stirred at 0 °C and warmed to room temperature. Once the reaction is complete, as determined by LCMS, the reaction is concentrated under reduced pressure. The residue is triturated with MeCN, then dripped into

MTBE over 10 min. The supernatant is removed and the precipitate is collected by filtration under N₂ to give 2-(4-amino-1-(4-aminobutyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-1H-indol-7-ol.

Monomer E. 5-(4-amino-1-(piperidin-4-ylmethyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)benzo[d]oxazol-2-amine trifluoroacetic acid salt.



Step 1: Synthesis of tert-butyl 4-((4-amino-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl)piperidine-1-carboxylate

[00475] To a solution of 3-iodo-1H-pyrazolo[3,4-d]pyrimidin-4-amine (3 g, 11.49 mmol, 1.0 equiv) in DMA (30 mL) was added *tert*-butyl 4-(bromomethyl)piperidine-1-carboxylate (3.36 g, 12.07 mmol, 1.05 equiv) and K₂CO₃ (4.77 g, 34.48 mmol, 3.0 equiv), then the reaction was stirred at 80 °C for 3 h. The reaction mixture was filtered to remove K₂CO₃ and the filtrate was poured into H₂O (200 mL). A solid precipitated was then filtered to give *tert*-butyl 4-((4-amino-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl)piperidine-1-carboxylate (3 g, 57% yield) as a light yellow solid. LCMS (ESI) *m/z*: [M + H] calcd for C₁₆H₂₃IN₆O₂: 459.10; found 459.1.

Step 2: Synthesis of tert-butyl 4-((4-amino-3-(2-aminobenzo[d]oxazol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl)piperidine-1-carboxylate


[00476] To a bi-phasic suspension of *tert*-butyl 4-((4-amino-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl)piperidine-1-carboxylate (3 g, 6.55 mmol, 1.0 equiv) and 5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzo[d]oxazol-2-amine (2.04 g, 7.86 mmol, 1.2 equiv) and Na₂CO₃ (3.47 g, 32.73 mmol, 5.0 equiv) in DME (60 mL) and H₂O (30 mL) was added Pd(PPh₃)₄ (756.43 mg, 654.60 μmol, 0.1 equiv) at room temperature under N₂. The mixture was stirred at 110 °C for 3 h and the two batches were combined together. The reaction mixture was cooled and partitioned between EtOAc (500 mL) and H₂O (500 mL). The aqueous layer was separated and extracted with EtOAc (3 x 300 mL). All the organic layers were combined, washed with brine (20 mL), dried over anhydrous Na₂SO₄, filtered, and the

filtrate was concentrated under reduced pressure to give *tert*-butyl 4-((4-amino-3-(2-aminobenzo[d]oxazol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl)piperidine-1-carboxylate (4.5 g, 74% yield) as a yellow solid. LCMS (ESI) *m/z*: [M + H] calcd for C₂₃H₂₈N₈O₃: 465.24; found 465.2.

Step 3: Synthesis of 5-(4-amino-1-(piperidin-4-ylmethyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)benzo[d]oxazol-2-amine

[00477] A solution of *tert*-butyl 4-((4-amino-3-(2-aminobenzo[d]oxazol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl)piperidine-1-carboxylate (2.5 g, 5.38 mmol, 1.0 equiv) in TFA (25 mL) was stirred at room temperature for 30 min. The reaction solution was concentrated under reduced pressure to remove TFA. The residue was added to MTBE (400 mL) and a solid precipitated, which was then filtered to give 5-(4-amino-1-(piperidin-4-ylmethyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)benzo[d]oxazol-2-amine (2.7 g, over 100 % yield, TFA) as a yellow solid. LCMS (ESI) *m/z*: [M + H] calcd for C₁₈H₂₀N₈O: 365.18; found 365.1.

Monomer F. 2-(4-amino-1-(4-aminobutyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-1H-indol-5-ol trifluoroacetic acid salt.



Step 1: Synthesis of *tert*-butyl 4-(4-amino-3-(5-((*tert*-butyldimethylsilyl)oxy)-1H-indol-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)butyl)carbamate

[00478] To a solution of *tert*-butyl 4-(4-amino-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-1-yl)butyl)carbamate (1.0 g, 2.31 mmol, 1.0 equiv) in dioxane (10.5 mL) and H₂O (3.5 mL) was added (1-(*tert*-butoxycarbonyl)-5-((*tert*-butyldimethylsilyl)oxy)-1H-indol-2-yl)boronic acid (1.54 g, 2.78 mmol, 1.2 equiv), K₃PO₄ (1.47 g, 6.94 mmol, 3.0 equiv), Pd₂(dba)₃ (211.84 mg, 231.34 μmol, 0.1 equiv), and SPhos (189.95 mg, 462.69 μmol, 0.2 equiv) at room temperature under N₂. The sealed tube was heated at 150 °C for 20 min in a microwave. This was repeated for 9 additional batches. The 10 batches were combined and the reaction mixture was cooled and partitioned between EtOAc (60 mL) and H₂O (80 mL). The aqueous layer was separated and extracted with EtOAc (2 x 50 mL). The organic layers were

combined, washed with brine (60 mL) and dried over anhydrous Na₂SO₄. The suspension was filtered and the filtrate was concentrated under reduced pressure. The crude material was purified by silica gel chromatography (1→75% EtOAc/petroleum ether). The desired fractions were combined and evaporated under reduced pressure to give *tert*-butyl (4-(4-amino-3-(5-((*tert*-butyldimethylsilyl)oxy)-1H-indol-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)butyl)carbamate (10 g, 60% yield) as a light yellow solid.


Step 2: Synthesis of tert-butyl (4-(4-amino-3-(5-hydroxy-1H-indol-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)butyl)carbamate

[00479] To a mixture of *tert*-butyl (4-(4-amino-3-(5-((*tert*-butyldimethylsilyl)oxy)-1H-indol-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)butyl)carbamate (10 g, 18.12 mmol, 1.0 equiv) in THF (100 mL) was added TBAF•3H₂O (1 M, 54.37 mL, 3.0 equiv) in one portion at room temperature under N₂. The mixture was stirred for 1 h and then H₂O (100 mL) was added to the reaction mixture. The layers were separated and the aqueous phase was extracted with EtOAc (2 x 80 mL). The combined organic phase was washed with brine (100 mL), dried with anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by silica gel chromatography (1→67% EtOAc/ petroleum ether) to afford *tert*-butyl (4-(4-amino-3-(5-hydroxy-1H-indol-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)butyl)carbamate (7 g, 87% yield) as a light pink solid.

Step 3: Synthesis of 2-[4-amino-1-(4-aminobutyl)pyrazolo[3,4-d]pyrimidin-3-yl]-1H-indol-5-ol

[00480] To TFA (50.0 mL, 675.26 mmol, 38.9 equiv) was added *tert*-butyl (4-(4-amino-3-(5-hydroxy-1H-indol-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)butyl)carbamate (7.6 g, 17.37 mmol, 1.0 equiv) at room temperature. The mixture was stirred for 40 min and was then concentrated under reduced pressure. The oily residue was triturated with MeCN (20 mL), then added dropwise into MTBE (300 mL) for 10 min. The supernatant was removed and then the precipitate was collected by filtration under N₂ to give 2-[4-amino-1-(4-aminobutyl)pyrazolo[3,4-d]pyrimidin-3-yl]-1H-indol-5-ol (7.79 g, 91% yield, TFA) as light yellow solid. LCMS (ESI) *m/z*: [M + H] calcd for C₁₇H₁₉N₇O: 338.17; found 338.2.

Monomer G. 5-(4-amino-1-(azetidin-3-ylmethyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)benzo[d]oxazol-2-amine trifluoroacetic acid salt.



Step 1: Synthesis of tert-butyl 3-((4-amino-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl)azetidine-1-carboxylate

[00481] To a solution of 3-iodo-1H-pyrazolo[3,4-d]pyrimidin-4-amine (4 g, 15.32 mmol, 1.0 equiv), *tert*-butyl 3-(hydroxymethyl)azetidine-1-carboxylate (3.01 g, 16.09 mmol, 1.05 equiv) and PPh₃ (6.03 g, 22.99 mmol, 1.5 equiv) in THF (80 mL) cooled to 0 °C was added DIAD (4.47 mL, 22.99 mmol, 1.5 equiv), dropwise. After the addition was complete, the reaction was stirred at room temperature for 14 h. The reaction was poured into H₂O (200 mL) and then extracted with EtOAc (3 x 50 mL). The organic layers were combined and washed with brine (2 x 50 mL). The organic phase was dried over Na₂SO₄, filtered, the filtrate was concentrated under reduced pressure to give a residue. The residue was purified by silica gel chromatography (0→100% EtOAc/petroleum ether) to give *tert*-butyl 3-((4-amino-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl)azetidine-1-carboxylate (4.2 g, 64% yield) as a white solid. LCMS (ESI) *m/z*: [M + H]⁺ calcd for C₁₄H₁₉IN₆O₂: 431.07; found 431.0.

Step 2: Synthesis of tert-butyl 3-((4-amino-3-(2-aminobenzo[d]oxazol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl)azetidine-1-carboxylate


[00482] To a bi-phasic suspension of *tert*-butyl 3-((4-amino-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl)azetidine-1-carboxylate (4 g, 9.30 mmol, 1.0 equiv), 5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzo[d]oxazol-2-amine (2.90 g, 11.16 mmol, 1.2 equiv) and Na₂CO₃ (4.93 g, 46.49 mmol, 5.0 equiv) in DME (100 mL) and H₂O (50 mL) was added Pd(PPh₃)₄ (1.07 g, 929.71 μmol, 0.1 equiv) at room temperature under N₂. The mixture was stirred at 110 °C for 3 h. The reaction mixture was then cooled to room temperature and filtered, and the filtrate was extracted by EtOAc (3 x 50 mL). The organic layers were combined and washed with brine (10 mL), dried over Na₂SO₄, filtered and the

filtrate was concentrated under reduced pressure to give a residue. The residue was purified by silica gel chromatography (0→20% MeOH/EtOAc) to give *tert*-butyl 3-((4-amino-3-(2-aminobenzo[d]oxazol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl)azetidine-1-carboxylate (3.5 g, 80% yield) as a yellow solid. LCMS (ESI) *m/z*: [M + H] calcd for C₂₁H₂₄N₈O₃: 437.20; found 437.2.

Step 3: Synthesis of 5-(4-amino-1-(azetidin-3-ylmethyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)benzo[d]oxazol-2-amine


[00483] To a solution of *tert*-butyl 3-((4-amino-3-(2-aminobenzo[d]oxazol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl)azetidine-1-carboxylate (3.29 g, 6.87 mmol, 1.0 equiv) in DCM (20 mL) was added TFA (7.50 mL, 101.30 mmol, 14.7 equiv) at 0 °C. The reaction was warmed to room temperature and stirred for 2 h. The reaction solution was concentrated under reduced pressure to give a residue. The residue was dissolved in MeCN (6 mL) and then poured into MTBE (80 mL). A solid precipitated, which was filtered and the solid cake was dried under reduced pressure to give 5-[4-amino-1-(azetidin-3-ylmethyl)pyrazolo[3,4-d]pyrimidin-3-yl]-1,3-benzoxazol-2-amine (4.34 g, over 100% yield, TFA) as a yellow solid. LCMS (ESI) *m/z*: [M + H] calcd for C₁₆H₁₆N₈O: 337.15; found 337.1.

Monomer H. 5-(4-amino-1-(4-aminobutyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)benzo[d]oxazol-2-amine trifluoroacetic acid salt.



[00484] This monomer was synthesized following the procedures outlined in *Nature* 2015, 534, 272-276, which is incorporated by reference in its entirety.

Monomer I. 5-(4-amino-1-(pyrrolidin-3-ylmethyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)benzo[d]oxazol-2-amine trifluoroacetic acid salt.



Step 1: Synthesis of tert-butyl 3-((4-amino-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl)pyrrolidine-1-carboxylate

[00485] A suspension of 3-iodo-1H-pyrazolo[3,4-d]pyrimidin-4-amine (4.5 g, 17.24 mmol, 1.0 equiv), *tert*-butyl 3-(bromomethyl)pyrrolidine-1-carboxylate (4.78 g, 18.10 mmol, 1.05 equiv) and K_2CO_3 (7.15 g, 51.72 mmol, 3.0 equiv) in DMA (40 mL) was heated to 85 °C. The reaction was stirred at 85 °C for 3 h, at which point the solution was cooled to room temperature. Then, H_2O (80 mL) was added to the reaction, and a solid precipitated out. The mixture was filtered, and the solid cake was washed with H_2O (2 x 40 mL), and then dried under reduced pressure to give *tert*-butyl 3-((4-amino-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl)pyrrolidine-1-carboxylate (6 g, 78% yield) as a yellow solid. LCMS (ESI) m/z : $[M + H]^+$ calcd for $C_{15}H_{21}IN_6O_2$: 445.08; found 445.1.


Step 2: Synthesis of tert-butyl 3-[[4-amino-3-(2-amino-1,3-benzoxazol-5-yl)pyrazolo[3,4-d]pyrimidin-1-yl]methyl]pyrrolidine-1-carboxylate

[00486] To a bi-phasic suspension of *tert*-butyl 3-((4-amino-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl)pyrrolidine-1-carboxylate (4 g, 9.00 mmol, 1.0 equiv), 5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzo[d]oxazol-2-amine (2.81 g, 10.80 mmol, 1.2 equiv) and Na_2CO_3 (4.77 g, 45.02 mmol, 5.0 equiv) in DME (120 mL) and H_2O (60 mL) was added $Pd(PPh_3)_4$ (1.04 g, 900.35 μ mol, 0.1 equiv) at room temperature under N_2 . The mixture was stirred at 110 °C for 3 h. The reaction mixture was cooled to room temperature and filtered and the filtrate was extracted with EtOAc (3 x 50 mL). The organic phases were combined and washed with brine (50 mL), dried over Na_2SO_4 , filtered and concentrated under reduced pressure to give a residue. The residue was purified by silica gel chromatography (0→20% MeOH/EtOAc) to give *tert*-butyl 3-((4-amino-3-(2-aminobenzo[d]oxazol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl)pyrrolidine-1-carboxylate (3 g, 64% yield) as a yellow solid. LCMS (ESI) m/z : $[M + H]^+$ calcd for $C_{22}H_{26}N_8O_3$: 451.21, found 451.2.

Step 3: Synthesis of 5-(4-amino-1-(pyrrolidin-3-ylmethyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)benzo[d]oxazol-2-amine

[00487] To a solution of *tert*-butyl 3-((4-amino-3-(2-aminobenzo[d]oxazol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl)pyrrolidine-1-carboxylate (3 g, 6.66 mmol, 1.0 equiv) in DCM (40 mL) was added TFA (20 mL) at 0 °C, dropwise. The reaction mixture was warmed to room temperature and stirred for 2 h. The reaction solution was then concentrated under reduced pressure to give a residue. The residue was dissolved in MeCN (4 mL), then poured into MTBE (100 mL), and a solid precipitated out. The solid was filtered and the cake was dried under reduced pressure to give 5-(4-amino-1-(pyrrolidin-3-ylmethyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)benzo[d]oxazol-2-amine (4.00 g, over 100% yield, TFA) as a yellow solid. LCMS (ESI) *m/z*: [M + H] calcd for C₁₇H₁₈N₈O: 351.17; found 351.2.

Monomer J. 1-(4-aminobutyl)-3-(7-methoxy-1H-indol-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-aminetrifluoroacetic acid salt.




Step 1: Synthesis of *tert*-butyl 2-(4-amino-1-(4-((*tert*-butoxycarbonyl)amino)butyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-7-methoxy-1H-indole-1-carboxylate

[00488] To a mixture of *tert*-butyl (4-(4-amino-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-1-yl)butyl)carbamate (1.0 equiv) and (1-(*tert*-butoxycarbonyl)-7-methoxy-1H-indol-2-yl)boronic acid (3.0 equiv) in DME and H₂O is added Pd(PPh₃)₄ (0.1 equiv) and sodium carbonate (6.0 equiv). The reaction is heated at 80 °C until completion, as determined by LCMS and TLC analysis. The reaction is then quenched with H₂O and EtOAc. The mixture is transferred to a separatory funnel and the aqueous phase is extracted with EtOAc. The organic phase is washed with sat. aq. NaCl, dried over Na₂SO₄, filtered, and concentrated under reduced pressure. The desired product is isolated after chromatography on silica gel.

Step 2: Synthesis of 1-(4-aminobutyl)-3-(7-methoxy-1H-indol-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-amine

[00489] To a solution of *tert*-butyl 2-(4-amino-1-(4-((*tert*-butoxycarbonyl)amino)butyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-7-hydroxy-1H-indole-1-carboxylate (1.0 equiv) in DCM at 0 °C is added TFA dropwise. The reaction is stirred at 0 °C and warmed to room temperature. Once the reaction is complete, as determined by LCMS, the reaction is concentrated under reduced pressure. The residue is triturated with MeCN, then dripped into MTBE over 10 min. The supernatant is removed and the precipitate is collected by filtration under N₂ to give 1-(4-aminobutyl)-3-(7-methoxy-1H-indol-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-amine.

Monomer K. Synthesis of 1-(4-aminobutyl)-1H-pyrazolo[3,4-d]pyrimidin-4-amine trifluoroacetic acid salt.



Step 1: Synthesis of *tert*-butyl (4-(4-amino-1H-pyrazolo[3,4-d]pyrimidin-1-yl)butyl)carbamate


[00490] To a mixture of *tert*-butyl (4-(4-amino-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-1-yl)butyl)carbamate (300 mg, 694 μ mol, 1.0 equiv) in MeOH (14 mL) at 0 °C was added zinc dust (226 mg, 3.46 mmol, 5.0 equiv). Sat. aq. NH₄Cl (14 mL) was added to the reaction mixture and the reaction was warmed to room temperature and stirred for 18 h. The reaction was quenched by EtOAc (40 mL) and H₂O (10 mL) and the mixture was transferred to a separatory funnel. The aqueous phase was extracted with EtOAc (3 x 20 mL) and the combined organic phases were washed with sat. aq. NaHCO₃ (15 mL), dried over Na₂SO₄, filtered, and concentrated under reduced pressure to provide the product (210 mg, 99% yield) as a light yellow solid that was used without further purification. LCMS (ESI) *m/z*: [M + H] calcd for C₁₄H₂₂N₆O₂: 307.19; found 307.1.

Step 2: Synthesis of 1-(4-aminobutyl)-1H-pyrazolo[3,4-d]pyrimidin-4-amine

[00491] To a solution of *tert*-butyl (4-(4-amino-1H-pyrazolo[3,4-d]pyrimidin-1-yl)butyl)carbamate (210 mg, 691 μ mol) in DCM (3.5 mL) at 0 °C was added TFA (3.5 mL), dropwise. After 3 h, the reaction was warmed to room temperature and concentrated under reduced pressure to provide the trifluoroacetate salt of the product (220 mg, 99% yield) as a


brown oil, which was used without further purification. LCMS (ESI) m/z : $[M + H]^+$ calcd for $C_9H_{14}N_6$: 207.13; found 207.1.

Monomer L. 1-[4-(piperazin-1-yl)-3-(trifluoromethyl)phenyl]-9-(quinolin-3-yl)-1H,2H-benzo[h]1,6-naphthyridin-2-one



[00492] The preparation of this monomer has been previously reported in the literature. See the following references: i) Liu, Qingsong; Chang, Jae Won; Wang, Jinhua; Kang, Seong A.; Thoreen, Carson C.; Markhard, Andrew; Hur, Wooyoung; Zhang, Jianming; Sim, Taebo; Sabatini, David M.; et al From Journal of Medicinal Chemistry (2010), 53(19), 7146-7155. ii) Gray, Nathanael; Chang, Jae Won; Zhang, Jianming; Thoreen, Carson C.; Kang, Seong Woo Anthony; Sabatini, David M.; Liu, Qingsong From PCT Int. Appl. (2010), WO 2010044885A2, which are incorporated by reference in their entirety.

Monomer M. 5-(1-(4-aminobutyl)-4-(dimethylamino)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)benzo[d]oxazol-2-amine trifluoroacetic acid salt.



Step 1: Synthesis of 3-iodo-1H-pyrazolo[3,4-d]pyrimidin-4-amine

[00493] A suspension of 3-iodo-1H-pyrazolo[3,4-d]pyrimidin-4-amine (10.5 g, 40.23 mmol, 1.0 equiv) in DMF (170.0 mL) was treated with CS_2CO_3 (19.7 g, 60.34 mmol, 1.5 equiv) and [chloro(diphenyl)methyl]benzene (13.5 g, 48.27 mmol, 1.2 equiv) at room temperature. The reaction mixture was stirred at 70 °C for 4 h under a nitrogen atmosphere.

The reaction mixture was added to H₂O (1200 mL). The precipitate was filtered and washed with H₂O. The residue was purified by silica gel chromatography (0→60% EtOAc/petroleum ether) to afford 3-iodo-1-trityl-1H-pyrazolo[3,4-d]pyrimidin-4-amine (15.40 g, 73.5% yield) as a white solid.

Step 2: Synthesis of 3-iodo-N,N-dimethyl-1-trityl-1H-pyrazolo[3,4-d]pyrimidin-4-amine

[00494] To a suspension of NaH (2.98 g, 74.50 mmol, 60 wt.%, 2.5 equiv) in DMF (150 mL) was added the solution of 3-iodo-1-trityl-1H-pyrazolo[3,4-d]pyrimidin-4-amine (15.0 g, 29.80 mmol, 1.0 equiv) in DMF (50 mL) at 0 °C. The mixture was stirred at 0 °C for 10 min. To the reaction mixture was then added iodomethane (16.92 g, 119.20 mmol, 7.42 mL, 4.0 equiv) at 0 °C. The mixture was stirred at room temperature for 2 h, at which point H₂O (1400 mL) was added at 0 °C. The mixture was stirred for an additional 10 min at 0 °C. The resulting precipitate was collected by filtration to give crude product, which was purified by silica gel chromatography (1%→25% EtOAc/petroleum ether) twice to afford 3-iodo-*N,N*-dimethyl-1-trityl-1H-pyrazolo[3,4-d]pyrimidin-4-amine (9.0 g, 89% yield) as a white solid.

Step 3: Synthesis of 3-iodo-N,N-dimethyl-1H-pyrazolo[3,4-d]pyrimidin-4-amine

[00495] To a cooled solution of TFA (19.1 mL, 258.1 mmol, 15.0 equiv) in DCM (100.0 mL) was added 3-iodo-*N,N*-dimethyl-1-trityl-1H-pyrazolo[3,4-d]pyrimidin-4-amine (9.10 g, 17.12 mmol, 1.0 equiv) at 4 °C. The mixture was stirred at room temperature for 1 h. The residue was poured into H₂O (100 mL) and the aqueous phase was extracted with DCM (2 x 50 mL). To the aqueous phase was then added a saturated aqueous solution of NaHCO₃ until the solution was pH 8. The resulting precipitate was collected by filtration to give 3-iodo-*N,N*-dimethyl-1H-pyrazolo[3,4-d]pyrimidin-4-amine (3.40 g, 68.7% yield) as a white solid.

Step 4: Synthesis of tert-butyl (4-(4-(dimethylamino)-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-1-yl)butyl)carbamate

[00496] To a suspension of 3-iodo-*N,N*-dimethyl-1H-pyrazolo[3,4-d]pyrimidin-4-amine (1.7 g, 5.88 mmol, 1.0 equiv) in DMF (20 mL) was added NaH (247 mg, 6.17 mmol, 60 wt.%, 1.05 equiv) at 4 °C. The mixture was stirred at 4 °C for 30 min. To the reaction mixture was then added *tert*-butyl N-(4-bromobutyl)carbamate (2.22 g, 8.82 mmol, 1.81 mL, 1.5 equiv) in DMF (10 mL) at 4 °C. The mixture was stirred at room temperature for 2 h. To the mixture was then added H₂O (100 mL) at 4 °C. The mixture was stirred for an additional 30 min at 4 °C and the resulting precipitate was collected by filtration to give crude product.

The residue was purified by silica gel chromatography (0→75% EtOAc/petroleum ether) to afford *tert*-butyl(4-(4-(dimethylamino)-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-1-yl)butyl)carbamate (2.0 g, 56% yield) as a white solid.

Step 5: Synthesis of tert-butyl (4-(3-(2-aminobenzo[d]oxazol-5-yl)-4-(dimethylamino)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)butyl)carbamate

[00497] To a bi-phasic suspension of *tert*-butyl (4-(4-(dimethylamino)-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-1-yl)butyl)carbamate (4.0 g, 8.69 mmol, 1.0 equiv), 5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzo[d]oxazol-2-amine (3.4 g, 13.03 mmol, 1.5 equiv), and Na₂CO₃ (4.6 g, 43.45 mmol, 5.0 equiv) in DME (80.0 mL) and H₂O (40.0 mL) was added Pd(PPh₃)₄ (1.0 g, 868.98 μmol, 0.1 equiv) at room temperature under N₂. The mixture was stirred at 110 °C for 3 h. The reaction mixture was then cooled and partitioned between EtOAc (300 mL) and H₂O (600 mL). The aqueous layer was separated and extracted with EtOAc (2 x 100 mL). The organic layers were combined, washed with brine (2 x 60 mL) and dried over anhydrous Na₂SO₄, filtered, and concentrated under reduced pressure. The crude material was purified by silica gel column chromatography (50% EtOAc/hexanes followed by 20% MeOH/EtOAc). The desired fractions were combined and concentrated under reduced pressure to give *tert*-butyl (4-(3-(2-aminobenzo[d]oxazol-5-yl)-4-(dimethylamino)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)butyl)carbamate (3.2 g, 78.9% yield) as a light brown solid.


Step 6: Synthesis of 5-(1-(4-aminobutyl)-4-(dimethylamino)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)benzo[d]oxazol-2-amine

[00498] To TFA (20.82 mL, 281.27 mmol, 36.5 equiv) was added *tert*-butyl (4-(3-(2-aminobenzo[d]oxazol-5-yl)-4-(dimethylamino)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)butyl)carbamate (3.6 g, 7.72 mmol, 1.0 equiv) at room temperature. The mixture was stirred for 30 min, at which point the mixture was concentrated under reduced pressure. The oily residue was triturated with MeCN (8 mL) and MTBE (60 mL) for 10 min. The supernatant was removed and then the precipitate was collected by filtration under N₂ to give 5-(1-(4-aminobutyl)-4-(dimethylamino)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)benzo[d]oxazol-2-amine (4.0 g, crude, TFA) as a light brown solid.

[00499] To 1 M NaOH (107.2 mL, 14.7 equiv) was added 5-(1-(4-aminobutyl)-4-(dimethylamino)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)benzo[d]oxazol-2-amine (3.5 g, crude, TFA) at room temperature. The mixture was stirred for 10 min and then the aqueous phase

was extracted with DCM (3 x 50 mL). The combined organic phase was washed with brine (50 mL), dried with anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. TFA (539.37 μ L, 7.28 mmol, 1.0 equiv) was added and concentrated under reduced pressure. MeCN (10 mL) was then added, followed by MTBE (150 mL). The resulting precipitate was collected by filtration to give 5-(1-(4-aminobutyl)-4-(dimethylamino)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)benzo[d]isoxazol-2-amine (1.3 g, 36.6% yield, TFA) as a light brown product. LCMS (ESI) *m/z*: [M + H] calcd for C₁₈H₂₂N₈O: 367.19; found 367.1.

Monomer N. 6-(4-amino-1-(4-aminobutyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)benzo[d]isoxazol-3-amine trifluoroacetic acid salt.



Step 1: Synthesis of *tert*-butyl (6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzo[d]isoxazol-3-yl)carbamate

[00500] To a solution of *tert*-butyl (6-bromobenzo[d]isoxazol-3-yl)carbamate (1.0 equiv) in dioxane is added Pd(PPh₃)₄ (0.1 equiv), sodium carbonate (6.0 equiv), and bis(pinacolato)diboron (3.0 equiv). The reaction mixture is stirred and heated until completion, as determined by LCMS and TLC analysis. The reaction is cooled to room temperature, quenched with sat. aq. NaHCO₃, and the mixture transferred to a separatory funnel. The aqueous phase is extracted with EtOAc and the organic phase is washed with sat. aq. NaCl, dried over Na₂SO₄, filtered, and concentrated under reduced pressure. The desired product is isolated after purification by silica gel chromatography.

Step 2: Synthesis of *tert*-butyl (4-(4-amino-3-(*tert*-butoxycarbonyl)amino)benzo[d]isoxazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)butyl)carbamate


[00501] To a mixture of *tert*-butyl (4-(4-amino-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-1-yl)butyl)carbamate (1.0 equiv) and *tert*-butyl (6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-

yl)benzo[d]isoxazol-3-yl)carbamate (3.0 equiv) in DME and H₂O is added Pd(PPh₃)₄ (0.1 equiv) and sodium carbonate (6.0 equiv). The reaction is heated at 80 °C until completion, as determined by LCMS and TLC analysis. The reaction is then quenched with H₂O and EtOAc. The mixture is transferred to a separatory funnel and the aqueous phase is extracted with EtOAc. The organic phase is washed with sat. aq. NaCl, dried over Na₂SO₄, filtered, and concentrated under reduced pressure. The desired product is isolated after chromatography on silica gel.

Step 3: Synthesis of 6-(4-amino-1-(4-aminobutyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)benzo[d]isoxazol-3-amine

[00502] To a solution of *tert*-butyl (4-(4-amino-3-(3-((*tert*-butoxycarbonyl)amino)benzo[d]isoxazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)butyl)carbamate (1.0 equiv) in DCM at 0 °C is added TFA, dropwise. The reaction is stirred at 0 °C and warmed to room temperature. Once the reaction is complete, as determined by LCMS, the reaction is concentrated under reduced pressure. The residue is triturated with MeCN, then added dropwise into MTBE over 10 min. The supernatant is removed and the precipitate is collected by filtration under N₂ to give 6-(4-amino-1-(4-aminobutyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)benzo[d]isoxazol-3-amine.

Monomer O. 4-(5-(4-morpholino-1-(1-(pyridin-3-ylmethyl)piperidin-4-yl)-1H-pyrazolo[3,4-d]pyrimidin-6-yl)-1H-indol-1-yl)butan-1-amine trifluoroacetic acid salt.




[00503] The synthesis of this monomer proceeds by alkylation of WAY-600 (CAS# 1062159-35-6) with *tert*-butyl (4-bromobutyl)carbamate under basic conditions, followed by Boc-deprotection using TFA to produce the TFA salt.

[00504] Reference for preparation of WAY-600: Discovery of Potent and Selective Inhibitors of the Mammalian Target of Rapamycin (mTOR) Kinase: Nowak, P.; Cole, D.C.; Brooijmans, N.; Bursavich, M.G.; Curran, K.J.; Ellingboe, J.W.; Gibbons, J.J.; Hollander, I.;

Hu, Y.; Kaplan, J.; Malwitz, D.J.; Toral-Barza, L.; Verheijen, J.C.; Zask, A.; Zhang, W.-G.; Yu, K. 2009; Journal of Medicinal Chemistry Volume 52, Issue 22, 7081-89, which is incorporated by reference in its entirety.

Monomer P. 2-(4-(8-(6-(aminomethyl)quinolin-3-yl)-3-methyl-2-oxo-2,3-dihydro-1H-imidazo[4,5-c]quinolin-1-yl)phenyl)-2-methylpropanenitrile trifluoroacetic acid salt.




[00505] The synthesis of this monomer proceeds first by synthesis of the Suzuki reaction coupling partner (3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolane)quinolin-6-yl)-*N*-boc-methanamine starting from methyl 3-bromoquinoline-6-carboxylate. Reduction of the methyl ester with lithium aluminum hydride followed by Mitsunobu reaction with phthalimide and hydrazine cleavage provides the benzylic amine. Protection of the benzylic amine with di-*tert*-butyl dicarbonate followed by a Miyaura borylation reaction provides (3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolane)quinolin-6-yl)-*N*-boc-methanamine.

[00506] An $\text{S}_{\text{N}}\text{Ar}$ reaction of 2-(4-aminophenyl)-2-methylpropanenitrile with 6-bromo-4-chloro-3-nitroquinoline provides the substituted amino-nitro-pyridine. Reduction of the nitro

group with Raney-Ni under a hydrogen atmosphere followed by cyclization with trichloromethyl chloroformate provides the aryl-substituted urea. Substitution of the free N-H of the urea with methyl iodide mediated by tetrabutylammonium bromide and sodium hydroxide followed by Suzuki coupling of (3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolane)quinolin-6-yl)-*N*-*tert*-butoxycarbonyl-methanamine and then Boc-deprotection using TFA produces the TFA salt.


[00507] Reference for preparation of 2-[4-(8-bromo-3-methyl-2-oxo-2,3-dihydro-imidazo[4,5-c]quinolin-1-yl)-phenyl]-2-methyl-propionitrile: Vannucchi, A.M.; Bogani, C.; Bartalucci, N. 2016. JAK PI3K/mTOR combination therapy. US9358229. Novartis Pharma AG, Incyte Corporation, which is incorporated by reference in its entirety.

Monomer Q. 8-(6-methoxypyridin-3-yl)-3-methyl-1-[4-(piperazin-1-yl)-3-(trifluoromethyl)phenyl]-1H,2H,3H-imidazo[4,5-c]quinolin-2-one



[00508] This monomer is a commercially available chemical known as BGT226 (CAS# 1245537-68-1). At the time this application was prepared, it was available for purchase from several vendors as the free amine.

Monomer R. 3-(4-amino-1-(4-aminobutyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-(4,5-dihydrothiazol-2-yl)benzamide trifluoroacetic acid salt.



Step 1: Synthesis of *tert*-butyl (4-(4-amino-3-(3-((4,5-dihydrothiazol-2-yl)carbamoyl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)butyl)carbamate


[00509] To a solution of 3-((4,5-dihydrothiazol-2-yl)carbamoyl)phenylboronic acid (500 mg, 1.15 mmol, 1.0 equiv) and *tert*-butyl (4-(4-amino-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-1-yl)butyl)carbamate (575 mg, 2.30 mmol, 2.0 equiv) in dioxane (19.1 mL), EtOH (3.8 mL), and H₂O (2.3 mL) was added Pd(PPh₃)₄ (265 mg, 230 μmol, 0.2 equiv) and sodium carbonate

(730 mg, 6.89 mmol, 6.0 equiv). The reaction mixture was sonicated until formation of a clear, yellow solution, which was subsequently heated at 80 °C for 14 h. The reaction was then diluted with sat. aq. NaCl (30 mL) and the mixture transferred to a separatory funnel. The aqueous phase was extracted with DCM (3 x 25 mL). The combined organic phases were dried over Na₂SO₄, filtered, and concentrated under reduced pressure. The desired product was isolated as a yellow solid (324 mg, 53% yield) after silica gel chromatography (0→15% MeOH/DCM). LCMS (ESI) *m/z*: [M + H] calcd for C₂₄H₃₀N₈O₃S: 511.22; found 511.2.

Step 2: Synthesis of 3-(4-amino-1-(4-aminobutyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-(4,5-dihydrothiazol-2-yl)benzamide

[00510] To a solution of *tert*-butyl (4-(4-amino-3-(3-((4,5-dihydrothiazol-2-yl)carbamoyl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)butyl)carbamate (324 mg, 614 μmol) in DCM (4.1 mL) at 0 °C was added TFA (1.5 mL), dropwise. After 1 h, the reaction was warmed to room temperature and concentrated under reduced pressure to provide the trifluoroacetate salt of the product as a yellow solid (320 mg, 99% yield). Used without further purification. LCMS (ESI) *m/z*: [M + H] calcd for C₁₉H₂₂N₈OS: 411.16; found 411.1


Monomer S. 2-(5-(4-morpholino-1-(1-(pyridin-3-ylmethyl)piperidin-4-yl)-1H-pyrazolo[3,4-d]pyrimidin-6-yl)-1H-indol-3-yl)ethan-1-amine.



[00511] The synthesis of this monomer proceeds by condensation of 2,4,6-trichloropyrimidine-5-carbaldehyde with 3-((4-hydrazineylpiperidin-1-yl)methyl)pyridine hydrochloride. Reaction of the product with morpholine followed by a Suzuki reaction with boronic ester gives the Boc-protected amine. Final deprotection with TFA gives the


monomer. This synthesis route follows closely to the reported preparation of highly related structures in the following references: i) Nowak, Pawel; Cole, Derek C.; Brooijmans, Natasja; Curran, Kevin J.; Ellingboe, John W.; Gibbons, James J.; Hollander, Irwin; Hu, Yong Bo; Kaplan, Joshua; Malwitz, David J.; et al From Journal of Medicinal Chemistry (2009), 52(22), 7081-7089. ii) Zask, Arie; Nowak, Pawel Wojciech; Verheijen, Jeroen; Curran, Kevin J.; Kaplan, Joshua; Malwitz, David; Bursavich, Matthew Gregory; Cole, Derek Cecil; Ayral-Kaloustian, Semiramis; Yu, Ker; et al From PCT Int. Appl. (2008), WO 2008115974 A2 20080925, which are incorporated by reference in their entirety.

Monomer T. 1-(4-aminobutyl)-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-4-amine trifluoroacetic acid salt.



[00512] To a mixture of *tert*-butyl (4-(4-amino-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-1-yl)butyl)carbamate (496 mg, 1.14 mmol, 1.0 equiv) in DCM (5.7 mL) at 0 °C was added TFA (1.5 mL) dropwise. The reaction was allowed to stir at 0 °C for 1 h, at which time the reaction was concentrated under reduced pressure to provide a yellow solid (505 mg, 99% yield) which was taken on without further purification. LCMS (ESI) *m/z*: [M + H] calcd for C₉H₁₃IN₆: 333.02; found 332.9.

Monomer U. 5-(4-amino-1-(4-(methylamino)butyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)benzo[d]oxazol-2-amine trifluoroacetic acid salt.



Step 1: Synthesis of tert-butyl (4-hydroxybutyl)(methyl)carbamate

[00513] To a solution of 4-(methylamino)butan-1-ol (0.5 g, 4.85 mmol, 104.2 mL, 1.0 equiv) in DCM (10 mL) at room temperature was added Boc₂O (1.06 g, 4.85 mmol, 1.11 mL, 1.0 equiv). The mixture was stirred for 3 h at room temperature and then the mixture was

concentrated under reduced pressure at 30 °C. The residue was purified by silica gel chromatography (100/1 to 3/1 petroleum ether/EtOAc) to afford *tert*-butyl (4-hydroxybutyl)(methyl)carbamate (0.9 g, 91.4% yield) as a colorless oil.

Step 2: Synthesis of tert-butyl (4-bromobutyl)(methyl)carbamate

[00514] To a solution of *tert*-butyl (4-hydroxybutyl)(methyl)carbamate (0.9 g, 4.43 mmol, 1.0 equiv) in THF (20 mL) at room temperature was added PPh₃ (2.21 g, 8.41 mmol, 1.9 equiv) and CBr₄ (2.79 g, 8.41 mmol, 1.9 equiv). The mixture was stirred for 1 h and then the reaction mixture was filtered and concentrated. The residue was purified by silica gel chromatography (1/0 to 4/1 petroleum ether/EtOAc) to afford *tert*-butyl (4-bromobutyl)(methyl) carbamate (1.1 g, 93.3% yield) as a colorless oil.

Step 3: Synthesis of tert-butyl (4-(4-amino-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-1-yl) butyl) (methyl)carbamate

[00515] To a suspension of 3-iodo-1H-pyrazolo[3,4-d]pyrimidin-4-amine (0.9 g, 3.45 mmol, 1.0 equiv) in DMF (10 mL) at 4 °C was added NaH (137.92 mg, 3.45 mmol, 60 wt.%, 1.0 equiv). The mixture was stirred at 4 °C for 30 min and then a solution of *tert*-butyl (4-bromobutyl)(methyl)carbamate (1.01 g, 3.79 mmol, 25.92 mL, 1.1 equiv) in DMF (3 mL) was added. The mixture was stirred at room temperature for 3 h, at which point H₂O (100 mL) was added. The aqueous phase was extracted with EtOAc (3 x 30 mL) and the combined organic phases were washed with brine (20 mL), dried with anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by silica gel chromatography (1/0 to 0/1 petroleum ether/EtOAc) to afford *tert*-butyl (4-(4-amino-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-1-yl)butyl) (methyl) carbamate (1.2 g, 78% yield) as a white solid. LCMS (ESI) *m/z*: [M + H] calcd for C₁₅H₂₃IN₆O₂: 447.10; found 447.1.

Step 4: Synthesis of tert-butyl (4-(4-amino-3-(2-aminobenzo[d]oxazol-5-yl)-1H- pyrazolo[3,4-d] pyrimidin-1-yl)butyl)(methyl)carbamate


[00516] To a bi-phasic suspension of *tert*-butyl (4-(4-amino-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-1-yl)butyl)(methyl)carbamate (1.2 g, 2.69 mmol, 1.0 equiv), 5-(4,4,5,5-tetramethyl-1,3,2- dioxaborolan-2-yl)benzo[d]oxazol-2-amine (1.19 g, 3.23 mmol, 1.2 equiv), and Na₂CO₃ (1.42 g, 13.44 mmol, 5.0 equiv) in DME (20 mL) and H₂O (10 mL) at room temperature was added Pd(PPh₃)₄ (310.71 mg, 268.89 μmol, 0.1 equiv) under N₂. The mixture was stirred at 110 °C for 3 h and then the reaction mixture was cooled and partitioned between EtOAc (20 mL) and H₂O (15 mL). The aqueous layer was separated and

extracted with EtOAc (3 x 20 mL). The combined organic layers were washed with brine (2 x 20 mL), dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. The crude product was purified by silica gel chromatography (1/0 to 4/1 EtOAc/MeOH) to give *tert*-butyl (4-(4-amino-3-(2-aminobenzo[d]oxazol-5-yl)-1H-pyrazolo [3,4-d]pyrimidin-1-yl)butyl)(methyl) carbamate (0.78 g, 62.5% yield) as an orange solid.

Step 5: Synthesis of 5-(4-amino-1-(4-(methylamino)butyl)-1H-pyrazolo[3,4-d] pyrimidin-3-yl) benzo[d]oxazol-2-amine

[00517] A solution of *tert*-butyl(4-(4-amino-3-(2-aminobenzo[d]oxazol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)butyl)(methyl)carbamate (0.78 g, 1.72 mmol, 1.0 equiv) in TFA (5 mL) at room temperature was stirred for 30 min. The solution was concentrated under reduced pressure and the oily residue was triturated with MeCN (1 mL) and then added to MTBE (100 mL). The supernatant was removed and then the precipitate was collected by filtration under N₂ to give 5-(4-amino-1-(4-(methylamino) butyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)benzo[d]oxazol -2-amine bis-trifluorosulfonate (0.959 g, 93% yield) as an orange solid. LCMS (ESI) *m/z*: [M + H] calcd for C₁₇H₂₀N₈O: 353.18; found 353.1.

Monomer V. 1-(4-(4-(5-(aminomethyl)pyrimidin-2-yl)piperazin-1-yl)-3-(trifluoromethyl)phenyl)-8-(6-methoxypyridin-3-yl)-3-methyl-1,3-dihydro-2H-imidazo[4,5-c]quinolin-2-one.



Step 1: Synthesis of *tert*-butyl N-*tert*-butoxycarbonyl-N-[(2-chloropyrimidin-5-yl)methyl] carbamate

[00518] To a solution of *tert*-butyl N-*tert*-butoxycarbonylcarbamate (7.33 g, 33.74 mmol, 1.0 equiv) in DMF (80 mL) was added NaH (1.62 g, 40.49 mmol, 60 wt.%, 1.2 equiv) at 0 °C. The mixture was stirred at 0 °C for 30 min and then 5-(bromomethyl)-2-chloro-

pyrimidine (7 g, 33.74 mmol, 1 equiv) was added. The reaction mixture was stirred at room temperature for 1.5 h and then the mixture was poured into sat. NH_4Cl (300 mL) and stirred for 5 min. The aqueous phase was extracted with EtOAc (3 x 80 mL) and the combined organic phases were washed with brine (50 mL), dried over anhydrous Na_2SO_4 , filtered and concentrated under reduced pressure. The residue was purified by silica gel chromatography (20:1 to 1:1 petroleum ether/EtOAc) to afford *tert*-butyl *N*-*tert*-butoxycarbonyl-*N*-[(2-chloropyrimidin-5-yl)methyl]carbamate (7.0 g, 60.3% yield) as a white solid. LCMS (ESI) m/z : $[\text{M} + \text{H}]$ calcd for $\text{C}_{15}\text{H}_{22}\text{ClN}_3\text{O}_4$: 344.14; found 344.2.

Step 2: Synthesis of tert-butyl N-tert-butoxycarbonyl-N-[[2-[4-[4-[8-(6-methoxy-3-pyridyl)-3-methyl-2-oxo-imidazo[4,5-c]quinolin-1-yl]-2-(trifluoromethyl)phenyl]piperazin-1-yl]pyrimidin-5-yl]methyl]carbamate


[00519] To a solution of 8-(6-methoxy-3-pyridyl)-3-methyl-1-[4-piperazin-1-yl]-3-(trifluoromethyl)phenyl]imidazo[4,5-c]quinolin-2-one (0.4 g, 748.32 μmol , 1.0 equiv) in MeCN (7 mL) was added *tert*-butyl *N*-*tert*-butoxycarbonyl-*N*-[(2-chloropyrimidin-5-yl)methyl]carbamate (514.55 mg, 1.50 mmol, 2.0 equiv) and K_2CO_3 (413.69 mg, 2.99 mmol, 4 equiv) at room temperature. The reaction mixture was stirred at 80 °C for 14 h and then the mixture was cooled to room temperature, filtered and concentrated under reduced pressure. The residue was purified by washing with MTBE (5 mL) to give *tert*-butyl *N*-*tert*-butoxycarbonyl-*N*-[[2-[4-[4-[8-(6-methoxy-3-pyridyl)-3-methyl-2-oxo-imidazo[4,5-c]quinolin-1-yl]-2-(trifluoromethyl)phenyl]piperazin-1-yl]pyrimidin-5-yl]methyl]carbamate (0.57 g, 90.5% yield) as a light yellow solid. LCMS (ESI) m/z : $[\text{M} + \text{H}]$ calcd for $\text{C}_{43}\text{H}_{46}\text{F}_3\text{N}_9\text{O}_6$: 842.36; found 842.7

Step 3: Synthesis of 1-[4-[4-[5-(aminomethyl)pyrimidin-2-yl]piperazin-1-yl]-3-(trifluoromethyl)phenyl]-8-(6-methoxy-3-pyridyl)-3-methyl-imidazo[4,5-c]quinolin-2-one

[00520] A solution of *tert*-butyl *N*-*tert*-butoxycarbonyl-*N*-[[2-[4-[4-[8-(6-methoxy-3-pyridyl)-3-methyl-2-oxo-imidazo[4,5-c]quinolin-1-yl]-2-(trifluoromethyl)phenyl]piperazin-1-yl]pyrimidin-5-yl]methyl]carbamate (0.95 g, 1.13 mmol, 1 equiv) in TFA (10 mL) was stirred at room temperature for 1 h, at which point the solvent was concentrated. The residue was dissolved in MeCN (10 mL) and then the solution was added to MTBE (150 mL), dropwise. The precipitate was collected to give 1-[4-[4-[5-(aminomethyl)pyrimidin-2-yl]piperazin-1-yl]-3-(trifluoromethyl)phenyl]-8-(6-methoxy-3-pyridyl)-3-methyl-imidazo[4,5-c]quinolin-2-

one trifluoromethanesulfonate (0.778 g, 84.8% yield) as a yellow solid. LCMS (ESI) m/z : $[M + H]$ calcd for $C_{33}H_{30}F_3N_9O_2$: 642.26; found 642.4

Monomer W. 1-(4-aminobutyl)-3-(1H-pyrrolo[2,3-b]pyridin-5-yl)pyrazolo[3,4-d]pyrimidin-4-amine.



Step 1: Synthesis of tert-butyl N-[4-[4-amino-3-(1H-indol-5-yl)pyrazolo[3,4-d]pyrimidin-1-yl]butyl]carbamate


[00521] To a bi-phasic suspension of *tert*-butyl N-[4-(4-amino-3-iodo-pyrazolo[3,4-d]pyrimidin-1-yl)butyl]carbamate (8 g, 18.51 mmol, 1 equiv), 5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrrolo[2,3-b]pyridine (5.42 g, 22.21 mmol, 1.2 equiv) and Na_2CO_3 (9.81 g, 92.54 mmol, 5 equiv) in diglyme (160 mL) and H_2O (80 mL) was added $Pd(PPh_3)_4$ (2.14 g, 1.85 mmol, 0.1 equiv) at room temperature under N_2 . The mixture was stirred at 110 °C for 3 h. The reaction mixture was cooled to room temperature, filtered and the filtrate was partitioned between EtOAc (500 mL) and H_2O (500 mL). The aqueous layer was separated and extracted with EtOAc (3 x 300 mL). The organic layers were combined, washed with brine (20 mL) and dried over anhydrous Na_2SO_4 , then filtered and the filtrate was concentrated under reduced pressure. The residue was purified by silica gel chromatography (1/0 to 0/1 petroleum ether/EtOAc then 4/1 EtOAc/MeOH) to give *tert*-butyl N-[4-[4-amino-3-(1H-indol-5-yl)pyrazolo[3,4-d]pyrimidin-1-yl]butyl]carbamate (6.6 g, 84.6% yield) as a yellow solid. LCMS (ESI) m/z : $[M + H]$ calcd for $C_{22}H_{27}N_7O_2$: 422.22; found 423.3.

Step 2: Synthesis of 1-(4-aminobutyl)-3-(1H-pyrrolo[2,3-b]pyridin-5-yl)pyrazolo[3,4-d]pyrimidin-4-amine

[00522] To *tert*-butyl N-[4-[4-amino-3-(1H-indol-5-yl)pyrazolo[3,4-d]pyrimidin-1-yl]butyl]carbamate (6.6 g, 15.66 mmol, 1 equiv) was added TFA (66 mL), which was then stirred at room temperature for 30 min. The reaction solution was concentrated under reduced pressure to remove TFA and then MTBE (400 mL) was added to the residue. The suspension was stirred for 15 min, at which point the yellow solid was filtered, and the solid cake dried

under reduced pressure to give 1-(4-aminobutyl)-3-(1H-pyrrolo[2,3-b]pyridin-5-yl)pyrazolo[3,4-d]pyrimidin-4-amine (10.2 g, 97.1% yield) as a yellow solid. LCMS (ESI) m/z : [M + H] calcd for C₁₆H₁₈N₈: 323.17; found 323.1.

Monomer X. 2-(4-amino-1-((1,2,3,4-tetrahydroisoquinolin-6-yl)methyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-1H-indol-5-ol 2,2,2-trifluoroacetate.



Step 1: Synthesis of tert-butyl 6-((4-amino-3-(5-((tert-butyldimethylsilyl)oxy)-1H-indol-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl)-3,4-dihydroisoquinoline-2(1H)-carboxylate

[00523] To a solution of *tert*-butyl 6-((4-amino-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl)-3,4-dihydroisoquinoline-2(1H)-carboxylate (1 g, 1.97 mmol, 1.0 equiv) in dioxane (10.5 mL) and H₂O (3.5 mL) was added (1-(*tert*-butoxycarbonyl)-5-((*tert*-butyldimethylsilyl)oxy)-1H-indol-2-yl)boronic acid (1.16 g, 2.96 mmol, 1.5 equiv), K₃PO₄ (1.26 g, 5.92 mmol, 3.0 equiv), Pd₂(dba)₃ (180.85 mg, 197.50 μmol, 0.1 equiv), and SPhos (162.16 mg, 394.99 μmol, 0.2 equiv) at room temperature under N₂. The sealed tube was heated at 150 °C for 20 min under microwave. The reaction mixture was then cooled and 6 separate batches were combined together. The reaction mixture was partitioned between EtOAc (100 mL) and H₂O (100 mL). The aqueous layer was separated and extracted with EtOAc (3 x 80 mL). The organic layers were combined, washed with brine (100 mL) and dried over anhydrous Na₂SO₄. The solution was filtered and the filtrate was concentrated under reduced pressure. The crude material was purified by silica gel column chromatography (100/1 to 1/4 petroleum ether/EtOAc) to give *tert*-butyl 6-((4-amino-3-(5-((*tert*-butyldimethylsilyl)oxy)-1H-indol-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl)-3,4-dihydroisoquinoline-2(1H)-carboxylate (6.17 g, 82.9% yield) as a light yellow solid.

Step 2: Synthesis of tert-butyl 6-((4-amino-3-(5-hydroxy-1H-indol-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl)-3,4-dihydroisoquinoline-2(1H)-carboxylate


[00524] To a mixture of *tert*-butyl 6-((4-amino-3-(5-((*tert*-butyldimethylsilyl)oxy)-1H-indol-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl)-3,4-dihydroisoquinoline-2(1H)-

carboxylate (6.17 g, 9.86 mmol, 1.0 equiv) in THF (100 mL) was added tetrabutylammonium fluoride trihydrate (1 M, 10.84 mL, 1.1 equiv) in one portion at 0 °C under N₂. The mixture was stirred at 0 °C for 1 h and was then added to H₂O (100 mL). The aqueous phase was extracted with EtOAc (3 x 80 mL) and the combined organic phase was washed with brine (2 x 80 mL), dried with anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by silica gel chromatography (1/1 to 0/1 petroleum ether/EtOAc) to afford *tert*-butyl 6-((4-amino-3-(5-hydroxy-1H-indol-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl)-3,4-dihydroisoquinoline-2(1H)-carboxylate (4 g, 79.3% yield) as a light pink solid. LCMS (ESI) *m/z*: [M + H] calcd for C₂₈H₂₉N₇O₃: 512.24; found 512.3.

Step 3: Synthesis of 2-(4-amino-1-((1,2,3,4-tetrahydroisoquinolin-6-yl)methyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-1H-indol-5-ol 2,2,2-trifluoroacetate

[00525] To a solution of *tert*-butyl 6-((4-amino-3-(5-hydroxy-1H-indol-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl)-3,4-dihydroisoquinoline-2(1H)-carboxylate (4.5 g, 8.80 mmol, 1.0 equiv) in MeOH (50 mL) was added HCl in MeOH (4 M, 50 mL, 22.7 equiv) at room temperature. The mixture was stirred at room temperature overnight and was then concentrated under reduced pressure. To the crude product was added EtOAc (100 mL) and the resulting precipitate was collected by filtration under N₂ to give 2-(4-amino-1-((1,2,3,4-tetrahydroisoquinolin-6-yl)methyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-1H-indol-5-ol 2,2,2-trifluoroacetate (4.1 g, 85.0% yield, 3HCl) as a light yellow solid. LCMS (ESI) *m/z*: [M + H] calcd for C₂₃H₂₁N₇O: 412.19; found 412.1.

Monomer Y. 3-(1H-pyrrolo[2,3-b]pyridin-5-yl)-1-((1,2,3,4-tetrahydroisoquinolin-6-yl)methyl)-1H-pyrazolo[3,4-d]pyrimidin-4-amine 2,2,2-trifluoroacetate.



Step 1: Synthesis of *tert*-butyl 6-(bromomethyl)-3,4-dihydroisoquinoline-2(1H)-carboxylate

[00526] A solution of NBS (34.07 g, 191.39 mmol, 4 equiv) in THF (200 mL) was added in portions to a solution of *tert*-butyl 6-(hydroxymethyl)-3,4-dihydroisoquinoline-2(1H)-carboxylate (12.6 g, 47.85 mmol, 1.0 equiv) and triphenylphosphine (37.65 g, 143.55 mmol, 3.0 equiv) in THF (200 mL) at 0 °C. After the addition was complete, the mixture was stirred for 1 h at room temperature. EtOAc (150 mL) was added and the mixture was washed with H₂O (200 mL) and brine (150 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The residue was purified by silica gel chromatography (100/1 to 10/1 petroleum ether/EtOAc) to afford *tert*-butyl 6-(bromomethyl)-3,4-dihydroisoquinoline-2(1H)-carboxylate (8.56 g, 54.8% yield) as a light yellow solid.

Step 2: Synthesis of *tert*-butyl 6-((4-amino-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl)-3,4-dihydroisoquinoline-2(1H)-carboxylate

[00527] To a suspension of 3-iodo-1H-pyrazolo[3,4-d]pyrimidin-4-amine (9.5 g, 36.40 mmol, 1.0 equiv) in DMF (110 mL) was added NaH (1.46 g, 36.40 mmol, 60 wt.%, 1.0 equiv) at 0 °C. The mixture was stirred at 0 °C for 30 min at which point a solution of *tert*-butyl 6-(bromomethyl)-3,4-dihydroisoquinoline-2(1H)-carboxylate (12.47 g, 38.22 mmol, 1.05 equiv) in DMF (40 mL) was added at 0 °C. The mixture was stirred at room temperature for 1 h and then H₂O (1000 mL) was added at 0 °C. The mixture stirred at 0 °C for 30 min and then the resulting precipitate was collected by filtration to give *tert*-butyl 6-((4-amino-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl)-3,4-dihydroisoquinoline-2(1H)-carboxylate

(17.8 g, 76.3% yield) as a light yellow solid, which was used the next step directly. LCMS (ESI) m/z : $[M + H]$ calcd for $C_{20}H_{23}N_6O_2$: 507.10; found 507.1.


Step 3: Synthesis of tert-butyl 6-((4-amino-3-(1H-pyrrolo[2,3-b]pyridin-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl)-3,4-dihydroisoquinoline-2(1H)-carboxylate

[00528] To a bi-phasic suspension of *tert*-butyl 6-((4-amino-3-iodo-1H-pyrazolo [3,4-d]pyrimidin-1-yl)methyl)-3,4-dihydroisoquinoline-2(1H)-carboxylate (6.5 g, 10.14 mmol, 1.0 equiv), 5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrrolo [2,3-b] pyridine (2.97 g, 12.16 mmol, 1.2 equiv), and Na_2CO_3 (5.37 g, 50.68 mmol, 5.0 equiv) in diglyme (100 mL) and H_2O (50 mL) was added $Pd(PPh_3)_4$ (1.17 g, 1.01 mmol, 0.1 equiv) at room temperature under N_2 . The mixture was stirred at 110 °C for 3 h. The reaction mixture was then cooled and partitioned between EtOAc (100 mL) and H_2O (100 mL). The aqueous layer was separated and extracted with EtOAc (2 x 100 mL). The combined organic phase was washed with brine (100 mL), dried with anhydrous Na_2SO_4 , filtered and concentrated under reduced pressure. The residue was purified by silica gel chromatography (0/1 to 1/4 MeOH/EtOAc) to afford *tert*-butyl 6-((4-amino-3-(1H-pyrrolo[2,3-b]pyridin-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl) methyl)-3,4-dihydroisoquinoline-2(1H)-carboxylate (3.77 g, 72.1% yield) as a light yellow solid. LCMS (ESI) m/z : $[M + H]$ calcd for $C_{27}H_{28}N_8O_2$: 497.24; found 497.3.

Step 4: Synthesis of 3-(1H-pyrrolo[2,3-b]pyridin-5-yl)-1-((1,2,3,4-tetrahydroiso quinolin-6-yl)methyl)-1H-pyrazolo[3,4-d]pyrimidin-4-amine 2,2,2-trifluoroacetate

[00529] *tert*-Butyl 6-((4-amino-3-(1H-pyrrolo[2,3-b]pyridin-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl)-3,4-dihydroisoquinoline-2(1H)-carboxylate (3.77 g, 7.59 mmol, 1.0 equiv) was added to TFA (85.36 mL, 1.15 mol, 151.8 equiv) at room temperature. The reaction mixture was stirred for 1 h. It was then concentrated under reduced pressure and the oily residue was triturated with MeCN (3 mL), then dripped into MTBE (200 mL) for 5 min. The supernatant was removed and then the precipitate was collected by filtration under N_2 to give the product, which was dissolved in MeCN (20 mL), and finally concentrated under reduced pressure to give 3-(1H-pyrrolo[2,3-b]pyridin-5-yl)-1-((1,2,3,4-tetrahydroisoquinolin-6-yl)methyl)-1H-pyrazolo[3,4-d]pyrimidin-4-amine 2,2,2-trifluoroacetate (4.84 g, 85.0% yield, 3TFA) as a light yellow solid. LCMS (ESI) m/z : $[M + H]$ calcd for $C_{22}H_{20}N_8$: 397.19; found 397.2.

Monomer Z. (4-((2-aminoethyl)sulfonyl)-3-fluoro-2-methylphenyl)(7- (6-aminopyridin-3-yl)-2,3-dihydrobenzo[f][1,4]oxazepin-4(5H)-yl)methanone 2,2,2-trifluoroacetate.



Step 1: Synthesis of methyl 3,4-difluoro-2-methylbenzoate

[00530] To a solution of 3,4-difluoro-2-methylbenzoic acid (2 g, 11.62 mmol, 1.0 equiv) in DMF (20 mL) was added K_2CO_3 (4.82g, 34.86 mmol, 3.0 equiv) and iodomethane (3.26 mL, 52.29 mmol, 4.5 equiv) at room temperature. The mixture was stirred at room temperature for 3 h. The solution of methyl 3,4-difluoro-2-methylbenzoate in DMF (20 mL) was used directly in the next step.

Step 2: Synthesis of methyl 4-((2-((tert-butoxycarbonyl)amino)ethyl)thio)-3-fluoro-2-methylbenzoate

[00531] To a solution of methyl 3,4-difluoro-2-methylbenzoate (2.16 g, 11.28 mmol, 1.0 equiv) in DMF (20 mL) was added *tert*-butyl (2-mercaptoethyl)carbamate (2.0 g, 11.28 mmol, 1 equiv) and K_2CO_3 (3.12 g, 22.56 mmol, 2.0 equiv) at room temperature. The reaction was stirred at 110 °C for 12 h, at which point the mixture was added to H_2O (50 mL). The aqueous solution was then extracted with EtOAc (3 x 30 mL) and the organic phase

was combined and concentrated under reduced pressure. The residue was purified by silica gel chromatography (1/0 to 3/1 petroleum ether/EtOAc) to afford methyl 4-((2-((*tert*-butoxycarbonyl)amino)ethyl)thio)-3-fluoro-2-methylbenzoate (3.0 g, 76% yield) as light yellow solid.

*Step 3: Synthesis of methyl 4-((2-((*tert*-butoxycarbonyl)amino)ethyl)sulfonyl)-3-fluoro-2-methylbenzoate*

[00532] To a solution of methyl 4-((2-((*tert*-butoxycarbonyl)amino)ethyl)thio)-3-fluoro-2-methylbenzoate (3.3 g, 9.61 mmol, 1.0 equiv), NaOH (2 M, 4.80 mL, 1.0 equiv), and NaHCO₃ (2.42 g, 28.83 mmol, 3.0 equiv) in acetone (30 mL) was added potassium peroxymonosulfate (12.35 g, 20.08 mmol, 2.1 equiv). The mixture was stirred for 12 h at room temperature and then the mixture was acidified to pH 5 by addition of 1N HCl. The aqueous layer was extracted with EtOAc (3 x 30 mL) and the combined organic phase was washed with brine (20 mL), dried with anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by silica gel chromatography (1/0 to 3/1 petroleum ether/EtOAc) to afford methyl 4-((2-((*tert*-butoxycarbonyl)amino)ethyl)sulfonyl)-3-fluoro-2-methylbenzoate (2.1 g, 58.2% yield) as a yellow solid. LCMS (ESI) *m/z*: [M-56 + H] calcd for C₁₆H₂₂FNO₆S: 320.12; found 320.1

*Step 4: Synthesis of 4-((2-((*tert*-butoxycarbonyl)amino)ethyl)sulfonyl)-3-fluoro-2-methylbenzoic acid*

[00533] To a solution of methyl 4-((2-((*tert*-butoxycarbonyl)amino)ethyl)sulfonyl)-3-fluoro-2-methylbenzoate (2.1 g, 5.59 mmol, 1.0 equiv) in THF (20 mL), MeOH (10 mL) and H₂O (10 mL) was added LiOH•H₂O (704.16 mg, 16.78 mmol, 3.0 equiv) at room temperature. The reaction mixture was stirred at 40 °C for 4 h. The mixture was then concentrated under reduced pressure to remove THF and MeOH. The aqueous phase was neutralized with 0.5N HCl and was then extracted with EtOAc (5 x 20 mL). The combined organic phase was washed with brine (2 x 20 mL), dried with anhydrous Na₂SO₄, filtered and concentrated under reduced pressure to give 4-((2-((*tert*-butoxycarbonyl)amino)ethyl)sulfonyl)-3-fluoro-2-methylbenzoic acid (2.01 g, 97.1% yield) as a white solid. LCMS (ESI) *m/z*: [M-100 + H] calcd for C₁₅H₂₀FNO₆S: 262.11; found 262.1.

*Step 5: Synthesis of (4-(*tert*-butoxycarbonyl)-2,3,4,5-tetrahydrobenzo[f][1,4]oxazepin-7-yl)boronic acid*

[00534] To a solution of *tert*-butyl 7-bromo-2,3-dihydrobenzo[f][1,4]oxazepine-4(5H)-carboxylate (4 g, 12.19 mmol, 1.0 equiv) in THF (80 mL) at -60 °C was added B(OiPr)₃ (4.58 g, 24.38 mmol, 5.60 mL, 2.0 equiv) followed by dropwise addition of *n*-BuLi (2.5 M, 12.19 mL, 2.5 equiv) in *n*-hexane. The reaction was stirred at -65 °C for 1 h. The reaction mixture was quenched with 1N HCl (12.25 mL) and allowed to warm to room temperature. The reaction mixture was extracted with EtOAc (3 x 30 mL), dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure to give (4-(*tert*-butoxycarbonyl)-2,3,4,5-tetrahydrobenzo[f][1,4]oxazepin-7-yl)boronic acid (3.5 g, crude) as light yellow oil, which was used to the next step directly. LCMS (ESI) *m/z*: [M-100 + H] calcd for C₁₄H₂₀BNO₅: 194.15; found 194.2.

Step 6: Synthesis of tert-butyl 7-(6-aminopyridin-3-yl)-2,3-dihydrobenzo[f][1,4] oxazepine-4(5H)-carboxylate

[00535] To a solution of (4-(*tert*-butoxycarbonyl)-2,3,4,5-tetrahydrobenzo[f][1,4]oxazepin-7-yl)boronic acid (4.2 g, 14.33 mmol, 1.0 equiv) in H₂O (20 mL) and dioxane (60 mL) was added 5-bromopyridin-2-amine (2.48 g, 14.33 mmol, 1.0 equiv), Pd(dppf)Cl₂•DCM (1.17 g, 1.43 mmol, 0.1 equiv) and Et₃N (4.35 g, 42.99 mmol, 5.98 mL, 3.0 equiv) at room temperature. The mixture was stirred at 85 °C for 12 h. The mixture was then cooled to room temperature and the residue was poured into H₂O (15 mL). The aqueous phase was extracted with EtOAc (3 x 40 mL) and the combined organic phase was washed with brine (2 x 40 mL), dried with anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by silica gel chromatography (1/0 to 1/8 petroleum ether/EtOAc) to afford *tert*-butyl 7-(6-aminopyridin-3-yl)-2,3-dihydrobenzo[f][1,4]oxazepine-4(5H)-carboxylate (3.3 g, 65.0% yield) as light yellow solid. LCMS (ESI) *m/z*: [M + H] calcd for C₁₉H₂₃N₃O₃: 342.18; found 342.2.

Step 7: Synthesis of 5-(2,3,4,5-tetrahydrobenzo[f][1,4]oxazepin-7-yl)pyridin-2-amine

[00536] To a solution of *tert*-butyl 7-(6-aminopyridin-3-yl)-2,3-dihydrobenzo[f][1,4]oxazepine-4(5H)-carboxylate (3.3 g, 9.67 mmol, 1.0 equiv) in THF (40 mL) was added HCl in EtOAc (4 M, 100 mL, 41.38 equiv) at room temperature. The mixture was stirred for 3 h. The reaction mixture was filtered and the filter cake was washed with EtOAc (3 x 15 mL) and then dried under reduced pressure to give 5-(2,3,4,5-tetrahydrobenzo[f][1,4]oxazepin-7-yl)pyridin-2-amine (3 g, 95.1% yield, 2HCl) as a light yellow solid.


Step 8: Synthesis of tert-butyl (2-((4-(7-(6-aminopyridin-3-yl)-2,3,4,5-tetrahydrobenzo[f][1,4]oxazepine-4-carbonyl)-2-fluoro-3-methylphenyl)sulfonyl)ethyl)carbamate

[00537] To a solution of 4-((2-((*tert*-butoxycarbonyl)amino)ethyl)sulfonyl)-3-fluoro-2-methylbenzoic acid (690.08 mg, 1.91 mmol, 1.0 equiv) in DMF (10 mL) was added HATU (1.09 g, 2.86 mmol, 1.5 equiv) and DIPEA (1.66 mL, 9.55 mmol, 5 equiv). The reaction was stirred at room temperature for 30 min and then 5-(2,3,4,5-tetrahydrobenzo[f][1,4]oxazepin-7-yl)pyridin-2-amine (0.6 g, 1.91 mmol, 1.0 equiv, 2HCl) was added. The mixture was stirred for 2 h, at which point H₂O (40 mL) was added. The mixture was stirred for 5 min and the resulting precipitate was collected by filtration to give the crude product. The residue was purified by silica gel chromatography (1/0 to 10/1 EtOAc/MeOH) to afford *tert*-butyl (2-((4-(7-(6-aminopyridin-3-yl)-2,3,4,5-tetrahydrobenzo[f][1,4]oxazepine-4-carbonyl)-2-fluoro-3-methylphenyl)sulfonyl)ethyl)carbamate (0.538 g, 47.4% yield) as a light yellow solid. LCMS (ESI) *m/z*: [M + H] calcd for C₂₉H₃₃FN₄O₆S: 585.22; found 585.3.

Step 9: Synthesis of (4-((2-aminoethyl)sulfonyl)-3-fluoro-2-methylphenyl)(7-(6-aminopyridin-3-yl)-2,3-dihydrobenzo[f][1,4]oxazepin-4(5H)-yl)methanone 2,2,2-trifluoroacetate

[00538] A solution *tert*-butyl (2-((4-(7-(6-aminopyridin-3-yl)-2,3,4,5-tetrahydrobenzo[f][1,4]oxazepine-4-carbonyl)-2-fluoro-3-methylphenyl)sulfonyl)ethyl)carbamate (0.538 g, 920.20 μmol, 1.0 equiv) in TFA (10.35 mL, 139.74 mmol, 151.85 equiv) was stirred at room temperature for 2 h. The solution was then concentrated under reduced pressure. The oily residue was triturated with MeCN (1 mL) and then dripped into MTBE (30 mL) for 10 min. The supernatant was removed and then the precipitate was collected by filtration under N₂ to give (4-((2-aminoethyl)sulfonyl)-3-fluoro-2-methylphenyl)(7-(6-aminopyridin-3-yl)-2,3-dihydrobenzo[f][1,4]oxazepin-4(5H)-yl)methanone 2,2,2-trifluoroacetate (0.50 g, 87.4% yield) as light brown solid. LCMS (ESI) *m/z*: [M + H] calcd for C₂₄H₂₅FN₄O₄S: 485.17; found 485.1.

Monomer AA. 5-(4-amino-1-(6-(piperazin-1-yl)pyrimidin-4-yl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)benzo[d]oxazol-2-amine trifluoroacetic acid salt.



Step 1: Synthesis of 1-(6-chloropyrimidin-4-yl)-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-4-amine

[00539] To a suspension of 3-iodo-1H-pyrazolo[3,4-d]pyrimidin-4-amine (5 g, 19.16 mmol, 1.0 equiv) in DMF (60 mL) was added NaH (804.53 mg, 20.11 mmol, 60 wt.%, 1.05 equiv) at 0 °C. The mixture was stirred at 0 °C for 30 min. To the reaction mixture was then added 4,6-dichloropyrimidine (3.42 g, 22.99 mmol, 1.2 equiv) at 0 °C. The mixture was stirred at room temperature for 2.5 h, at which point the reaction mixture was added to H₂O (600 mL). The suspension was then filtered to give the product (7.1 g, 99.2% yield) as yellow solid. LCMS (ESI) *m/z*: [M + H] calcd for C₉H₅ClIN₇: 373.94; found 373.9.

Step 2: Synthesis of tert-butyl 4-(6-(4-amino-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-1-yl)pyrimidin-4-yl)piperazine-1-carboxylate

[00540] To a solution of 1-(6-chloropyrimidin-4-yl)-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-4-amine (5 g, 13.39 mmol, 1.0 equiv) and *tert*-butyl piperazine-1-carboxylate (2.99 g, 16.06 mmol, 1.2 equiv) in DMF (50 mL) was added K₂CO₃ (3.70 g, 26.77 mmol, 2.0 equiv). The reaction mixture was stirred at 100 °C for 4 h, at which point it was added to H₂O (500 mL). The suspension was then filtered to give the product (6.2 g, 88.5% yield) as yellow solid. LCMS (ESI) *m/z*: [M + H] calcd for C₁₈H₂₂IN₉O₂: 524.09; found 524.2.


Step 3: Synthesis of tert-butyl 4-(6-(4-amino-3-(2-aminobenzo[d]oxazol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)pyrimidin-4-yl)piperazine-1-carboxylate

[00541] To a bi-phasic suspension of 5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzo[d]oxazol-2-amine (3.08 g, 11.85 mmol, 1.0 equiv), *tert*-butyl 4-(6-(4-amino-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-1-yl)pyrimidin-4-yl)piperazine-1-carboxylate (6.2 g, 11.85 mmol, 1.0 equiv) and Na₂CO₃ (6.28 g, 59.24 mmol, 5.0 equiv) in H₂O (100 mL) and DME (200 mL) was added Pd(PPh₃)₄ (1.37 g, 1.18 mmol, 0.1 equiv) at room temperature under N₂. The mixture was stirred at 110 °C for 24 h and then the mixture was filtered to give a solid cake. The solid was added to dioxane (20 mL) and stirred at 110 °C for 60 min, then filtered to give the product (3.5 g, 55.8% yield) as brown solid. LCMS (ESI) *m/z*: [M + H] calcd for C₂₅H₂₇N₁₁O₃: 530.24; found 530.3.

Step 4: Synthesis of 5-(4-amino-1-(6-(piperazin-1-yl)pyrimidin-4-yl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)benzo[d]oxazol-2-amine trifluoroacetic acid salt

[00542] A solution of *tert*-butyl 4-(6-(4-amino-3-(2-aminobenzo[d]oxazol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)pyrimidin-4-yl)piperazine-1-carboxylate (3.5 g, 6.61 mmol, 1.0 equiv) in TFA (35 mL) was stirred at room temperature for 1 h. The reaction solution was concentrated under reduced pressure and the resulting crude material was dissolved in MeCN (20 mL) and added dropwise to MTBE (500 mL). The resulting solid was then filtered to give the product (5.5 g, 91.9% yield) as brown solid. LCMS (ESI) *m/z*: [M + H] calcd for C₂₀H₁₉N₁₁O: 430.19; found 430.1.

Monomer AB. 8-(6-methoxypyridin-3-yl)-3-methyl-1-(4-(4-(5,6,7,8-tetrahydropyrido[4,3-d]pyrimidin-2-yl)piperazin-1-yl)-3-(trifluoromethyl)phenyl)-1H-imidazo[4,5-c]quinolin-2(3H)-one trifluoroacetic acid salt.



Step 1: Synthesis of *tert*-butyl 2-(4-(4-(8-(6-methoxypyridin-3-yl)-3-methyl-2-oxo-2,3-dihydro-1H-imidazo[4,5-c]quinolin-1-yl)-2-(trifluoromethyl)phenyl)piperazin-1-yl)-7,8-dihydropyrido[4,3-d]pyrimidine-6(5H)-carboxylate


[00543] To a mixture of 8-(6-methoxypyridin-3-yl)-3-methyl-1-(4-(piperazin-1-yl)-3-(trifluoromethyl)phenyl)-1H-imidazo[4,5-c]quinolin-2(3H)-one (0.3 g, 561.24 μmol, 1.0

equiv) and *tert*-butyl 2-chloro-7,8-dihydropyrido[4,3-*d*]pyrimidine-6(5*H*)-carboxylate (151.38 mg, 561.24 μ mol, 1.0 equiv) in DMF (5 mL) was added K₂CO₃ (193.92 mg, 1.40 mmol, 2.5 equiv). The mixture was stirred at 100 °C for 14 h, at which point H₂O (20 mL) was added. The aqueous layer was extracted with EtOAc (3 x 40 mL) and the combined organic layers were concentrated under reduced pressure. The crude material was purified by column chromatography (30/1 to 15/1 DCM/MeOH) to give the product (0.30 g, 69.6% yield) as a light-yellow solid. LCMS (ESI) *m/z*: [M + H] calcd for C₄₀H₄₀F₃N₉O₄: 768.33; found 768.5.

Step 2: Synthesis of 8-(6-methoxypyridin-3-yl)-3-methyl-1-(4-(4-(5,6,7,8-tetrahydropyrido[4,3-*d*]pyrimidin-2-yl)piperazin-1-yl)-3-(trifluoromethyl)phenyl)-1*H*-imidazo[4,5-*c*]quinolin-2(3*H*)-one

[00544] A solution of *tert*-butyl 2-(4-(4-(8-(6-methoxypyridin-3-yl)-3-methyl-2-oxo-2,3-dihydro-1*H*-imidazo[4,5-*c*]quinolin-1-yl)-2-(trifluoromethyl)phenyl)piperazin-1-yl)-7,8-dihydropyrido[4,3-*d*]pyrimidine-6(5*H*)-carboxylate (0.8 g, 1.04 mmol, 1.0 equiv) in TFA (8 mL) was stirred at room temperature for 2 h. The solvent was concentrated and the residue was dissolved in MeCN (5 mL), then the solution was added dropwise to MTBE (150 mL). The precipitate was filtered and the solid was dried under reduced pressure to give the product (600 mg, 70.6% yield) as a yellow solid. LCMS (ESI) *m/z*: [M + H] calcd for C₃₅H₃₂F₃N₉O₂: 668.27; found 668.3.

Monomer AC. 5-(4-amino-1-(piperidin-4-ylmethyl)-1*H*-pyrazolo[3,4-*d*]pyrimidin-3-yl)benzo[d]oxazol-2-amine trifluoroacetic acid salt.



Step 1: Synthesis of tert-butyl 4-((methylsulfonyl)oxy)piperidine-1-carboxylate

[00545] To a solution of *tert*-butyl 4-hydroxypiperidine-1-carboxylate (4 g, 19.87 mmol, 1.0 equiv) and Et₃N (3.87 mL, 27.82 mmol, 1.4 equiv) in DCM (40 mL) was added MsCl (2.15 mL, 27.82 mmol, 1.4 equiv) at 0 °C. Then the reaction mixture was stirred at room temperature for 1 h. H₂O (50 mL) was added and the aqueous phase was extracted with DCM (3 x 50 mL). The combined organic phase was washed with brine, dried with anhydrous Na₂SO₄, filtered and concentrated under reduced pressure to give the product (5.62 g, 101% crude yield) as yellow solid which was used directly in the next step.

Step 2: Synthesis of tert-butyl 4-(4-amino-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-1-yl)piperidine-1-carboxylate

[00546] To a suspension of 3-iodo-1H-pyrazolo[3,4-d]pyrimidin-4-amine (5 g, 19.16 mmol, 1.0 equiv) and *tert*-butyl 4-((methylsulfonyl)oxy)piperidine-1-carboxylate (5.62 g, 20.11 mmol, 1.05 equiv) in DMF (100 mL) was added K₂CO₃ (5.29 g, 38.31 mmol, 2.0 equiv). The mixture was stirred at 80 °C for 12 h. The reaction mixture was then added to H₂O (400 mL) at 0 °C. The resulting precipitate was filtered to give the product (5.0 g, 58.8% yield) as yellow solid. LCMS (ESI) *m/z*: [M + H] calcd for C₁₅H₂₁IN₆O₂: 445.09; found 445.1.


Step 3: Synthesis of tert-butyl 4-(4-amino-3-(2-aminobenzo[d]oxazol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)piperidine-1-carboxylate

[00547] To a suspension of *tert*-butyl 4-(4-amino-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-1-yl)piperidine-1-carboxylate (5 g, 11.25 mmol, 1.0 equiv), 5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzo[d]oxazol-2-amine (3.51 g, 13.51 mmol, 1.2 equiv) and Na₂CO₃ (5.96 g, 56.27 mmol, 5.0 equiv) in H₂O (50 mL) and DME (100 mL) was added Pd(PPh₃)₄ (1.30 g, 1.13 mmol, 0.1 equiv) at room temperature under N₂. The mixture was stirred at 110 °C for 3 h. The reaction mixture was then cooled to room temperature and filtered. The filtrate was partitioned between EtOAc (100 mL) and H₂O (100 mL) and then the aqueous layer was separated and extracted with EtOAc (3 x 100 mL). The combined organic layer was washed with brine (20 mL) and dried over anhydrous Na₂SO₄, filtered, and concentrated under reduced pressure. The residue was triturated with EtOAc (30 mL) and filtered to give the product (3.6 g, 71% yield) as yellow solid. LCMS (ESI) *m/z*: [M + H] calcd for C₂₂H₂₆N₈O₃: 451.22; found 451.3.

Step 4: Synthesis of 5-(4-amino-1-(piperidin-4-yl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)benzo[d]oxazol-2-amine trifluoroacetic acid salt

[00548] A solution of *tert*-butyl 4-(4-amino-3-(2-aminobenzo[d]oxazol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)piperidine-1-carboxylate (1.4 g, 3.11 mmol, 1.0 equiv) in TFA (10 mL) was stirred at room temperature for 30 min. The reaction solution was concentrated under reduced pressure and the crude solid was dissolved in MeCN (20 mL). The solution was added dropwise to MTBE (100 mL) and the resulting solid was filtered to give the product (1.6 g, 85.8% yield) as yellow solid. LCMS (ESI) *m/z*: [M + H] calcd for C₁₇H₁₈N₈O₃: 351.17; found 351.1.

Monomer AD. 1-(piperidin-4-yl)-3-(1H-pyrrolo[2,3-b]pyridin-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-amine trifluoroacetic acid salt.




Step 1: Synthesis of *tert*-butyl 4-(4-amino-3-(1H-pyrrolo[2,3-b]pyridin-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)piperidine-1-carboxylate

[00549] To a suspension of 5-(4,4,5-trimethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrrolo[2,3-b]pyridine (857.12 mg, 3.51 mmol, 1.2 equiv), *tert*-butyl 4-(4-amino-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-1-yl)piperidine-1-carboxylate (1.3 g, 2.93 mmol, 1.0 equiv) and Na₂CO₃ (1.55 g, 14.63 mmol, 5.0 equiv) in DME (20 mL) and H₂O (10 mL) was added Pd(PPh₃)₄ (338.13 mg, 292.62 μmol, 0.1 equiv) at room temperature under N₂. The mixture was stirred at 110 °C for 3 h. The reaction mixture was then cooled to room temperature and filtered. The filtrate was partitioned between EtOAc (50 mL) and H₂O (50 mL) and the aqueous layer was separated and extracted with EtOAc (3 x 50 mL). The combined organic layer were washed with brine, dried over anhydrous Na₂SO₄, filtered, and concentrated under reduced pressure. The residue was triturated with EtOAc (10 mL), filtered, the solid cake was dried under reduced pressure to give the product (1.0 g, 78.7% yield) as yellow solid.

Step 2: Synthesis of 1-(piperidin-4-yl)-3-(1H-pyrrolo[2,3-b]pyridin-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-amine trifluoroacetic acid salt

[00550] A solution of *tert*-butyl 4-(4-amino-3-(1H-pyrrolo[2,3-b]pyridin-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)piperidine-1-carboxylate (1.5 g, 3.45 mmol, 1.0 equiv) in TFA (10 mL) was stirred at room temperature for 30 min. The reaction solution was concentrated under reduced pressure and the crude residue was dissolved in MeCN (20 mL). The solution was added dropwise to MTBE (100 mL) and the resulting solid was filtered to give the product (1.19 g, 74.2% yield) as light yellow solid. LCMS (ESI) m/z : [M + H] calcd for C₁₇H₁₈N₈: 335.18; found 335.1.

Monomer AE. (4-((2-aminoethyl)sulfonyl)-2-methylphenyl)(7-(6-aminopyridin-3-yl)-2,3-dihydrobenzo[f][1,4]oxazepin-4(5H)-yl)methanone.



Step 1: Synthesis of methyl 4-fluoro-2-methylbenzoate

[00551] To a solution of 4-fluoro-2-methylbenzoic acid (86 g, 557.94 mmol, 1.0 equiv) in DMF (900 mL) was added K₂CO₃ (231.33 g, 1.67 mol, 3.0 equiv) and iodomethane (79.19 g, 557.94 mmol, 34.73 mL, 1.0 equiv). The mixture was stirred at room temperature for 1 h. The solution of methyl 4-fluoro-2-methylbenzoate in DMF (900 mL) was used directly in the next step.

Step 2: Synthesis of methyl 4-((2-((*tert*-butoxycarbonyl)amino)ethyl)thio)-2-methylbenzoate

[00552] To a solution of methyl 4-fluoro-2-methylbenzoate (93.8 g, 557.94 mmol, 1.0 equiv) in DMF (900 mL) was added *tert*-butyl (2-mercaptoethyl)carbamate (98.91 g, 557.97 mmol, 1.0 equiv) and K₂CO₃ (154.23 g, 1.12 mol, 2.0 equiv). The reaction was stirred at 110 °C for 12 h, at which point the mixture was cooled to room temperature and added to H₂O (1000 mL). The aqueous layer was then extracted with EtOAc (3 x 600 mL) and the combined organic layers were washed with brine, dried, and concentrated under reduced pressure. Purification by silica gel chromatography (0→25% EtOAc/petroleum ether) afforded the desired product as a colorless oil (144 g, 79% yield).

*Step 3: Synthesis of methyl 4-((2-((*tert*-butoxycarbonyl)amino)ethyl)sulfonyl)-2-methylbenzoate*

[00553] To two separate batches containing a solution of methyl 4-((2-((*tert*-butoxycarbonyl)amino)ethyl)thio)-2-methylbenzoate (72 g, 221.25 mmol, 1.0 equiv), NaOH (2 M, 110.6 mL, 1.0 equiv), and NaHCO₃ (55.76 g, 663.75 mmol, 3.0 equiv) in acetone (750 mL) was added potassium peroxymonosulfate (284.28 g, 462.41 mmol, 2.1 equiv). The mixture was stirred for 12 h at room temperature, at which point the two batches were combined and then the mixture was acidified to pH 5 by addition of 1N HCl. The aqueous layer was extracted with EtOAc (3 x 1500 mL) and the combined organic phases were washed with brine (2 x 500 mL), dried, and concentrated under reduced pressure. Purification by silica gel chromatography (0→25% EtOAc/petroleum ether) afforded the desired product as a white solid (120 g, 76% yield).

*Step 4: Synthesis of 4-((2-((*tert*-butoxycarbonyl)amino)ethyl)sulfonyl)-2-methylbenzoic acid*

[00554] To a solution of methyl 4-((2-((*tert*-butoxycarbonyl)amino)ethyl)sulfonyl)-2-methylbenzoate (35 g, 97.92 mmol, 1.0 equiv) in THF (200 mL), MeOH (100 mL) and H₂O (100 mL) was added LiOH•H₂O (12.33 g, 293.77 mmol, 3.0 equiv) at room temperature. The reaction mixture was stirred at 40 °C for 1 h. The mixture was then concentrated under reduced pressure to remove THF and MeOH. The aqueous phase was neutralized with 0.5N HCl and the resulting precipitate was isolated by filtration. The solid cake was washed with H₂O (3 x 20 mL) to afford the desired product as a white solid (25 g, 74% yield).

*Step 5: Synthesis of *tert*-butyl (2-((4-(7-(6-aminopyridin-3-yl)-2,3,4,5-tetrahydrobenzo[f][1,4]oxazepine-4-carbonyl)-3-methylphenyl)sulfonyl)ethyl)carbamate*


[00555] To a solution of 4-((2-((*tert*-butoxycarbonyl)amino)ethyl)sulfonyl)-2-methylbenzoic acid (9.7 g, 28.25 mmol, 1.0 equiv) and 5-(2,3,4,5-

tetrahydrobenzo[f][1,4]oxazepin-7-yl)pyridin-2-amine (8.88 g, 28.25 mmol, 1.0 equiv, 2HCl) in DMF (120 mL) was added HATU (16.11 g, 42.37 mmol, 1.5 equiv) and DIPEA (18.25 g, 141.24 mmol, 24.60 mL, 5.0 equiv). The reaction was stirred at room temperature for 1 h, at which point the reaction mixture was poured into H₂O (1000 mL). The mixture was stirred for 5 min and the resulting precipitate was collected by filtration to give the crude product. The crude product was triturated with EtOAc (100 mL), filtered, and the solid cake was dried under reduced pressure to afford the desired product as a white solid (14 g, 87% yield).

Step 6: Synthesis of (4-((2-aminoethyl)sulfonyl)-2-methylphenyl)(7-(6-aminopyridin-3-yl)-2,3-dihydrobenzo[f][1,4]oxazepin-4(5H)-yl)methanone

[00556] A solution *tert*-butyl (2-((4-(7-(6-aminopyridin-3-yl)-2,3,4,5-tetrahydrobenzo[f][1,4]oxazepine-4-carbonyl)-3-methylphenyl)sulfonyl)ethyl)carbamate (19 g, 33.53 mmol, 1.0 equiv) in TFA (100 mL) was stirred at room temperature for 30 min. The solution was then concentrated under reduced pressure. The residue was triturated with MeCN (30 mL) and then dripped into MTBE (600 mL) and stirred for 20 min. The suspension was filtered and the resulting solid was dissolved in MeCN (30 mL) and concentrated under reduced pressure to afford the desired product as a light yellow solid (24 g, TFA salt). LCMS (ESI) *m/z*: [M + H] calcd for C₂₄H₂₆N₄O₄S: 467.18; found 467.1.

Monomer AF. 5-(4-amino-1-((5,6,7,8-tetrahydropyrido[4,3-d]pyrimidin-2-yl)methyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)benzo[d]oxazol-2-amine.



Step 1: Synthesis of (*Z*)-*tert*-butyl 3-((dimethylamino)methylene)-4-oxopiperidine-1-carboxylate

[00557] A solution of *tert*-butyl 4-oxopiperidine-1-carboxylate (15 g, 75.28 mmol, 1.0 equiv) and 1,1-dimethoxy-*N,N*-dimethylmethanamine (11.00 mL, 82.81 mmol, 1.1 equiv) in

DMF (105 mL) was stirred at 95 °C for 12 h. The reaction mixture was then concentrated under reduced pressure and the resulting residue was dissolved in EtOAc (30 mL) and washed with brine (3 x 30 mL). The aqueous phase was extracted with EtOAc (50 mL), and the combined organic phases were dried and concentrated under reduced pressure to afford the desired product as a yellow solid (10.1 g, 53% yield).

Step 2: Synthesis of tert-butyl 2-(hydroxymethyl)-7,8-dihydropyrido[4,3-d]pyrimidine-6(5H)-carboxylate

[00558] To a solution of NaOEt (1.98 g, 29.10 mmol, 1.0 equiv) in EtOH (70 mL) was added (*Z*)-*tert*-butyl 3-((dimethylamino)methylene)-4-oxopiperidine-1-carboxylate (7.4 g, 29.10 mmol, 1.0 equiv) and 2-hydroxyacetimidamide hydrochloride (3.54 g, 32.01 mmol, 1.1 equiv). The reaction mixture was heated to 90 °C for 12 h, at which point the mixture was cooled to room temperature and concentrated under reduced pressure. The residue was partitioned with EtOAc (40 mL) and washed with sat. NaHCO₃ (40 mL). The aqueous phase was extracted with EtOAc (3 x 20 mL) and the combined organic phases were washed with brine (2 x 50 mL), dried, and concentrated under reduced pressure. Purification by silica gel chromatography (25% EtOAc/petroleum ether) afforded the desired product as a yellow solid (7.24 g, 94% yield).

Step 3: Synthesis of tert-butyl 2-(bromomethyl)-7,8-dihydropyrido[4,3-d]pyrimidine-6(5H)-carboxylate

[00559] To a solution of *tert*-butyl 2-(hydroxymethyl)-7,8-dihydropyrido[4,3-d]pyrimidine-6(5H)-carboxylate (6.24 g, 23.52 mmol, 1.0 equiv) and PPh₃ (12.34 g, 47.04 mmol, 2.0 equiv) in DCM (140 mL) was added CBr₄ (14.82 g, 44.69 mmol, 1.9 equiv). The mixture was stirred at room temperature for 3 h, at which point mixture was concentrated under reduced pressure. The residue was partitioned between EtOAc (20 mL) and H₂O (20 mL), the aqueous phase was extracted with EtOAc (3 x 20 mL). The combined organic phases were washed with brine (2 x 50 mL), dried, and concentrated under reduced pressure. Purification by silica gel chromatography (14% EtOAc/petroleum ether) afforded the desired product as a yellow solid (3.6 g, 47% yield).

Step 4: Synthesis of tert-butyl 2-((4-amino-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl)-7,8-dihydropyrido[4,3-d]pyrimidine-6(5H)-carboxylate

[00560] To a solution of 3-iodo-1H-pyrazolo[3,4-d]pyrimidin-4-amine (1.59 g, 6.09 mmol, 1.0 equiv) in DMF (15 mL) was added NaH (243.73 mg, 6.09 mmol, 60 wt.%, 1.0 equiv) at 0

°C. The suspension was stirred for 30 min and then *tert*-butyl 2-(bromomethyl)-7,8-dihydropyrido[4,3-d]pyrimidine-6(5H)-carboxylate (2.2 g, 6.70 mmol, 1.1 equiv) was added. The reaction mixture was warmed to room temperature and stirred for 3 h. The mixture was poured into H₂O at 0 °C and the precipitate was collected by filtration to afford the desired product as a brown solid (2.5 g, 66% yield).


Step 5: Synthesis of tert-butyl 2-((4-amino-3-(2-aminobenzo[d]oxazol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl)-7,8-dihydropyrido[4,3-d]pyrimidine-6(5H)-carboxylate

[00561] To a solution of *tert*-butyl 2-((4-amino-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl)-7,8-dihydropyrido[4,3-d]pyrimidine-6(5H)-carboxylate (4.55 g, 8.95 mmol, 1.0 equiv), 5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzo[d]oxazol-2-amine (2.79 g, 10.74 mmol, 1.2 equiv) and Na₂CO₃ (4.74 g, 44.76 mmol, 5.0 equiv) in dioxane (70 mL) and H₂O (35 mL) was added Pd(PPh₃)₄ (1.03 g, 895.11 μmol, 0.1 equiv). The reaction mixture was heated to 110 °C for 3 h, at which point the mixture was cooled to room temperature and poured into H₂O at 0 °C. The precipitate was filtered, and the solid cake was dried under reduced pressure. The crude product was washed with EtOAc (50 mL) to afford the desired product as light yellow solid (3.14 g, 68% yield).

Step 6: Synthesis of 5-(4-amino-1-((5,6,7,8-tetrahydropyrido[4,3-d]pyrimidin-2-yl)methyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)benzo[d]oxazol-2-amine

[00562] A solution of *tert*-butyl 2-((4-amino-3-(2-aminobenzo[d]oxazol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl)-7,8-dihydropyrido[4,3-d]pyrimidine-6(5H)-carboxylate (3.14 g, 6.10 mmol, 1.0 equiv) in TFA (20 mL) was stirred at room temperature for 30 min. The mixture was concentrated under reduced pressure and the resulting residue was added dissolved in MeCN (7 mL) and added to MTBE (700 mL). The precipitate was collected by filtration to afford the desired product as a brown solid (4.25 g, 92% yield, 3 TFA). LCMS (ESI) *m/z*: [M + H] calcd for C₂₀H₁₈N₁₀O: 415.18; found 415.1.

Monomer AG. 5-(4-amino-1-((2-((2-aminoethyl)sulfonyl)-1,2,3,4-tetrahydroisoquinolin-6-yl)methyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)benzo[d]oxazol-2-amine.



Step 1: Synthesis of N-Boc taurine tetrabutylammonium salt

[00563] To a solution of 2-aminoethanesulfonic acid (10.00 mL, 79.91 mmol, 1.0 equiv) in THF (60 mL) and aqueous NaOH (2 M, 40 mL, 1.0 equiv) was added Boc_2O (18.31 g, 83.90 mmol, 1.05 equiv). The mixture was stirred at room temperature for 15 h, at which point the mixture was extracted with EtOAc (10 mL). The aqueous phase was diluted with H_2O (450 mL), treated with $\text{LiOH}\cdot\text{H}_2\text{O}$ (3.35 g, 79.83 mmol, 1.0 equiv) and $n\text{Bu}_4\text{NH}_4\text{SO}_4$ (27.13 g, 79.90 mmol, 1.0 equiv) and stirred for 30 min. This mixture was extracted with DCM (3 x 80 mL), and the combined organic phases were dried and concentrated under reduced pressure to afford the desired product as a colorless oil (34.26 g, 91% yield).

Step 2: Synthesis of tert-butyl (2-(chlorosulfonyl)ethyl)carbamate

[00564] To a solution of *N*-Boc taurine tetrabutylammonium salt (4.7 g, 10.05 mmol, 1.0 equiv) in DCM (42 mL) was added DMF (77.32 μL , 1.00 mmol, 0.1 equiv) followed by a solution of triphosgene (0.5 M, 8.04 mL, 0.4 equiv) in DCM at 0 °C. The mixture was warmed to room temperature and stirred for 30 min. The solution of *tert*-butyl (2-(chlorosulfonyl)ethyl)carbamate (2.45 g, crude) in DCM was used directly in the next step.


Step 3: Synthesis of tert-butyl (2-(((6-((4-amino-3-(2-aminobenzo[d]oxazol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl)-3,4-dihydroisoquinolin-2(1H)-yl)sulfonyl)ethyl)carbamate

[00565] To a solution of 5-(4-amino-1-((1,2,3,4-tetrahydroisoquinolin-6-yl)methyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)benzo[d]oxazol-2-amine (6.04 g, 9.44 mmol, 1.0 equiv, 2TFA) in DMF (40 mL) was added Et₃N (7.88 mL, 56.63 mmol, 6.0 equiv). A solution of *tert*-butyl (2-(chlorosulfonyl)ethyl)carbamate in DCM (42 mL) at 0 °C was added. The mixture was warmed to room temperature and stirred 16 h. The reaction mixture was concentrated under reduced pressure to remove DCM and the resulting solution was purified by reverse phase chromatography (15→45% MeCN/H₂O) to afford the desired product as a white solid (5.8 g, 83% yield, TFA). LCMS (ESI) *m/z*: [M + H] calcd for C₂₉H₃₃N₉O₅S: 620.24; found 620.3.

Step 4: Synthesis of 5-(4-amino-1-((2-((2-aminoethyl)sulfonyl)-1,2,3,4-tetrahydroisoquinolin-6-yl)methyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)benzo[d]oxazol-2-amine

[00566] A solution of *tert*-butyl (2-(((6-((4-amino-3-(2-aminobenzo[d]oxazol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl)-3,4-dihydroisoquinolin-2(1H)-yl)sulfonyl)ethyl)carbamate (5.8 g, 9.36 mmol, 1.0 equiv) in TFA (48 mL) was stirred at room temperature for 0.5 h, at which point the reaction mixture was concentrated under reduced pressure. The crude product dissolved in MeCN (30 mL) and was added dropwise into MTBE (200 mL). The mixture was stirred for 5 min and filtered, the filter cake was dried under reduced pressure to afford the desired product as a yellow solid (3.6 g, 62% yield, 2.2TFA). LCMS (ESI) *m/z*: [M + H] calcd for C₂₄H₂₅N₉O₃S: 520.19; found 520.1.

Monomer AH. *tert*-butyl ((5-((4-amino-3-(2-aminobenzo[d]oxazol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl)pyrimidin-2-yl)methyl)carbamate.



Step 1: Synthesis of (2-(((*tert*-butoxycarbonyl)amino)methyl)pyrimidin-5-yl)methyl methanesulfonate

[00567] To a solution of *tert*-butyl ((5-(hydroxymethyl)pyrimidin-2-yl)methyl)carbamate (4.2 g, 17.55 mmol, 1.0 equiv) in DCM (42 mL) at 0 °C was added Et₃N (7.33 mL, 52.66 mmol, 3.0 equiv) followed by MsCl (2.41 g, 21.06 mmol, 1.63 mL, 1.2 equiv). The mixture was stirred at 0 °C for 10 min, and then H₂O (15 mL) was added. The reaction mixture was extracted with DCM (5 x 10 mL) and the combined organic phases were washed with brine (5 mL), dried, filtered, and concentrated under reduced pressure to afford the desired product (5.5 g, 98.7% yield) as a colorless solid.

Step 2: Synthesis of *tert*-butyl ((5-((4-amino-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl)pyrimidin-2-yl)methyl)carbamate

[00568] To a solution of (2-(((*tert*-butoxycarbonyl)amino)methyl)pyrimidin-5-yl)methyl methanesulfonate (5.47 g, 17.24 mmol, 1.2 equiv) and 3-iodo-1H-pyrazolo[3,4-d]pyrimidin-4-amine (3.75 g, 14.37 mmol, 1.0 equiv) in DMF (55 mL) at room temperature was added K₂CO₃ (5.96 g, 43.10 mmol, 3 equiv). The mixture was stirred at 80 °C for 5 h, at which point H₂O (100 mL) and brine (20 mL) were poured into the reaction mixture. The solution was extracted with EtOAc (10 x 30 mL) and the combined organic phases were dried, filtered, and concentrated under reduced pressure. Purification by silica gel chromatography (0→30% EtOAc/MeOH) afforded the desired product (2 g, 28.9% yield) as a yellow solid.


Step 3: Synthesis of tert-butyl ((5-((4-amino-3-(2-aminobenzo[d]oxazol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl)pyrimidin-2-yl)methyl)carbamate

[00569] To a solution of *tert*-butyl ((5-((4-amino-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl)pyrimidin-2-yl)methyl)carbamate (2 g, 4.15 mmol, 1.0 equiv), 5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1,3-benzoxazol-2-amine (1.13 g, 4.35 mmol, 1.05 equiv) and Na₂CO₃ (688.39 mg, 8.29 mmol, 2.0 equiv) in dioxane (20 mL) and H₂O (10 mL) was added Pd(PPh₃)₄ (479.21 mg, 414.70 μmol, 0.1 equiv). The mixture was stirred at 110 °C for 1 h, at which time the mixture was cooled to room temperature, filtered, and the solid cake washed with MeOH (3 x 10 mL). The filtrate was concentrated under reduced pressure to remove MeOH and then added dropwise into H₂O (50 mL). The resulting suspension was filtered, and the filter cake was washed with H₂O (3 x 10 mL). The solid cake was stirred in MeOH (20 mL) for 30 min. The resulting suspension was filtered, and the filter cake washed with MeOH (3 x 8 mL). The filter cake was dried under reduced pressure to afford the desired product (1.03 g, 48.9% yield) as a white solid. LCMS (ESI) *m/z*: [M + H] calcd for C₂₃H₂₄N₁₀O₃: 489.21; found 489.2.

Step 4: Synthesis of 5-(4-amino-1-{{2-(aminomethyl)pyrimidin-5-yl}methyl}-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-1,3-benzoxazol-2-amine

[00570] To *tert*-butyl ((5-((4-amino-3-(2-aminobenzo[d]oxazol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl)pyrimidin-2-yl)methyl)carbamate (100 mg, 0.205 mmol, 1.0 equiv) was added con. HCl (850 μL, 10.2 mmol, 50 equiv). The reaction was stirred for 1 h and was then poured into acetone (3 mL). The resulting precipitate was filtered, washed with acetone, and dried under reduced pressure to afford the desired product (80 mg, 92% yield) as a brown solid. LCMS (ESI) *m/z*: [M + H] calcd for C₁₈H₁₆N₁₀O: 389.16; found 389.0.

Monomer AI. 5-(4-(dimethylamino)-1-((1,2,3,4-tetrahydroisoquinolin-6-yl)methyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)benzo[d]oxazol-2-amine trifluoroacetic acid salt.



Step 1: Synthesis of tert-butyl 6-((4-(dimethylamino)-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl)-3,4-dihydroisoquinoline-2(1H)-carboxylate

[00571] To a solution of 3-iodo-*N,N*-dimethyl-1H-pyrazolo[3,4-d]pyrimidin-4-amine (3.6 g, 12.45 mmol, 1.0 equiv) in DMF (36 mL) at 0 °C was added NaH (523.00 mg, 13.08 mmol, 60 wt.%, 1.05 equiv). The mixture was stirred at 0 °C for 30 min. To the reaction mixture was then added a solution of *tert*-butyl 6-(bromomethyl)-3,4-dihydroisoquinoline-2(1H)-carboxylate (4.47 g, 13.70 mmol, 1.1 equiv) in DMF (18 mL) at 0 °C. The mixture was stirred at room temperature for 2 h. The reaction mixture was then added to cold H₂O (200 mL) and stirred for 30 min. The resulting precipitate was collected by filtration to afford the desired product (6 g, 71.9% yield) as a white solid.


Step 2: Synthesis of tert-butyl 6-((3-(2-aminobenzo[d]oxazol-5-yl)-4-(dimethylamino)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl)-3,4-dihydroisoquinoline-2(1H)-carboxylate

[00572] To a solution of *tert*-butyl 6-((4-(dimethylamino)-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl)-3,4-dihydroisoquinoline-2(1H)-carboxylate (2 g, 2.96 mmol, 1.0 equiv) and 5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzo[d]oxazol-2-amine (922.81 mg, 3.55 mmol, 1.2 equiv) in dioxane (24 mL) and H₂O (12 mL) was added Na₂CO₃ (1.57 g, 14.78 mmol, 5.0 equiv) and Pd(PPh₃)₄ (341.66 mg, 295.66 μmol, 0.1 equiv). The mixture was stirred at 110 °C for 12 h. The reaction mixture was then poured into cold H₂O (200 mL) and stirred for 30 min. The resulting precipitate was collected by filtration. Purification by silica gel chromatography (5→100% petroleum ether/EtOAc) afforded the desired product (1.2 g, 72.3% yield) as a yellow solid.

Step 3: Synthesis of 5-(4-(dimethylamino)-1-((1,2,3,4-tetrahydroisoquinolin-6-yl)methyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)benzo[d]oxazol-2-amine


[00573] A solution of *tert*-butyl 6-((3-(2-aminobenzo[d]oxazol-5-yl)-4-(dimethylamino)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl)-3,4-dihydroisoquinoline-2(1H)-carboxylate (1.7 g, 3.14 mmol, 1.0 equiv) in TFA (10 mL) was stirred at room temperature for 30 min. The reaction mixture was then concentrated under reduced pressure. The residue was added to MeCN (10 mL) and the solution was added dropwise into MTBE (200 mL). The resulting solid was dissolved in MeCN (30 mL) and the solution was concentrated under reduced pressure to afford the desired product (1.67 g, 92.9% yield,) as a yellow solid. LCMS (ESI) *m/z*: [M + H] calcd for C₂₄H₂₄N₈O: 441.22; found 441.2.

Monomer AJ. 4-amino-5-(2-aminobenzo[d]oxazol-5-yl)-5H-pyrimido[5,4-b]indole-7-carboxylic acid.




[00574] This monomer can be prepared from 7-methyl-5H-pyrimido[5,4-b]indol-4-ol by benzylic oxidation to the carboxylic acid, conversion to the ethyl ester, followed by *O*-ethylation with triethyloxonium tetrafluoroborate. Palladium-mediated arylation followed by ester hydrolysis and final ammonia-olysis provides the monomer.

Monomer AK. 4-amino-5-(2-aminobenzo[d]oxazol-5-yl)-5H-pyrimido[5,4-b]indole-8-carboxylic acid.



[00575] This monomer can be prepared following a similar route as that to prepare the previous monomer, but using the isomeric starting material from 8-methyl-5H-pyrimido[5,4-b]indol-4-ol. Benzylic oxidation to the carboxylic acid, conversion to the ethyl ester, followed by *O*-ethylation with triethyloxonium tetrafluoroborate and palladium-mediated arylation, followed by ester hydrolysis and final ammonia-olysis provides the monomer.

Monomer AL. 3-(2,4-bis((*S*)-3-methylmorpholino)-4a,8a-dihydropyrido[2,3-d]pyrimidin-7-yl)benzoic acid.



Step 1: Synthesis of (3*S*)-4-[7-chloro-2-[(3*S*)-3-methylmorpholin-4-yl]pyrido[2,3-d]pyrimidin-4-yl] 3-methyl-morpholine

[00576] To a solution of 2,4,7-trichloropyrido[2,3-d]pyrimidine (4.0 g, 17.06 mmol, 1.0 equiv) in DMA (10 mL) was added (3*S*)-3-methylmorpholine (4.31 g, 42.65 mmol, 2.5 equiv) and DIPEA (5.51 g, 42.65 mmol, 7.43 mL, 2.5 equiv). The reaction solution was heated to 70 °C for 48 h. The reaction suspension was cooled to room temperature, poured into cold H₂O (50 mL) to precipitate out a solid. The solid was filtered and the filter cake was rinsed with H₂O, and dried under reduced pressure to give the crude product, which was purified by column chromatography on silica gel (0→100% petroleum ether/EtOAc) to give (3*S*)-4-[7-chloro-2-[(3*S*)-3-methylmorpholin-4-yl]pyrido[2,3-d]pyrimidin-4-yl] 3-methyl-morpholine (3.5 g, 56.4% yield) as a yellow solid. LCMS (ESI) *m/z*: [M + H] calcd for C₁₇H₂₂ClN₅O₂: 364.15; found 364.2


Step 2: Synthesis of 3-[2,4-bis[(3*S*)-3-methylmorpholin-4-yl]pyrido[2,3-d]pyrimidin-7-yl]benzoic acid

[00577] To a solution of (3*S*)-4-[7-chloro-2-[(3*S*)-3-methylmorpholin-4-yl]pyrido[2,3-d]pyrimidin-4-yl] 3-methyl-morpholine (2 g, 5.50 mmol, 1.0 equiv) and 3-boronobenzoic acid (1.09 g, 6.60 mmol, 1.2 equiv) in 1,4-dioxane (40 mL) was added a solution of K₂CO₃ (911.65 mg, 6.60 mmol, 1.2 equiv) in H₂O (4 mL), followed by Pd(PPh₃)₄ (317.60 mg, 274.85 μmol, 0.05 equiv). The solution was degassed for 10 min and refilled with N₂, then the reaction mixture was heated to 100 °C under N₂ for 5 h. The reaction was cooled to room

temperature and filtered. The filtrate was acidified by HCl (2N) to pH 3, and the aqueous layer was washed with EtOAc (3 x 20 mL). The aqueous phase was concentrated under reduced pressure to give a residue, which was purified by column chromatography on silica gel (50%→100% petroleum ether/EtOAc) to give 3-[2,4-bis[(3*S*)-3-methylmorpholin-4-yl]pyrido[2,3-*d*]pyrimidin-7-yl]benzoic acid hydrochloride (2.5 g, 89.9% yield) as a yellow solid. LCMS (ESI) *m/z*: [M + H] calcd for C₂₄H₂₇N₅O₄: 450.21; found 450.2.


[00578] Reference for preparation of this monomer: Menear, K.; Smith, G.C.M.; Malagu, K.; Duggan, H.M.E.; Martin, N.M.B.; Leroux, F.G.M. 2012. Pyrido-, pyrazo- and pyrimido-pyrimidine derivatives as mTOR inhibitors. US8101602. Kudos Pharmaceuticals, Ltd, which is incorporated by reference in its entirety.

Monomer AM. (1*r*,4*r*)-4-[4-amino-5-(7-methoxy-1*H*-indol-2-yl)imidazo[4,3-*f*][1,2,4]triazin-7-yl]cyclohexane-1-carboxylic acid




[00579] This monomer, also known as OSI-027 (CAS# = 936890-98-1), is a commercially available compound. At the time this application was prepared, it was available for purchase from several vendors.

Monomer AN. 2-(4-(4-(8-(6-methoxypyridin-3-yl)-3-methyl-2-oxo-2,3-dihydro-1*H*-imidazo[4,5-*c*]quinolin-1-yl)-2-(trifluoromethyl)phenyl)piperazin-1-yl)pyrimidine-5-carboxylic acid.



[00580] Preparation of this monomer proceeds by reaction of BGT226 with methyl 2-chloropyrimidine-5-carboxylate, followed by ester hydrolysis, to give the titled Monomer.


Monomer AO. 4-amino-5-{1H-pyrrolo[2,3-b]pyridin-5-yl}-5H-pyrimido[5,4-b]indole-8-carboxylic acid.



[00581] This monomer can be prepared from 7-methyl-5H-pyrimido[5,4-b]indol-4-ol by benzylic oxidation to the carboxylic acid, conversion to the ethyl ester, followed by *O*-ethylation with triethyloxonium tetrafluoroborate. Palladium-mediated arylation followed by ester hydrolysis and final ammonia-olysis provides the monomer.

Preparation of pre- and post-Linkers

Building block A. *tert*-butyl N-[(*tert*-butoxy)carbonyl]-N-[2-(piperazin-1-yl)pyrimidin-5-yl]methyl carbamate.



Step 1: Synthesis of 5-(bromomethyl)-2-chloropyrimidine

[00582] To a solution of 2-chloro-5-methylpyrimidine (92 g, 715.62 mmol, 1.0 equiv) in CCl₄ (1000 mL) was added NBS (178.31 g, 1.00 mol, 1.4 equiv) and benzoyl peroxide (3.47 g, 14.31 mmol, 0.02 equiv). The mixture was stirred at 76 °C for 18 h. The reaction mixture was then cooled to room temperature and concentrated under reduced pressure. The reaction mixture was filtered and the solid cake was washed with DCM (150 mL). The resulting solution was concentrated under reduced pressure to give the crude product. The residue was purified by silica gel chromatography (1/0 to 0/1 petroleum ether/EtOAc) to give the product (70.8 g, 47.7% crude yield) as yellow oil, which was used directly for the next step. LCMS (ESI) *m/z*: [M + H] calcd for C₈H₄BrClN₂: 206.93; found 206.9.

Step 2: Synthesis of tert-butyl N-tert-butoxycarbonyl-N-((2-piperazin-1-ylpyrimidin-5-yl)methyl)carbamate

[00583] To a solution of *tert*-butyl *N-tert*-butoxycarbonylcarbamate (36.89 g, 169.79 mmol, 0.74 equiv) in DMF (750 mL) was added NaH (6.88 g, 172.09 mmol, 60 wt.%, 0.75 equiv) at 0 °C. The mixture was stirred at 0 °C for 30 min. Then, 5-(bromomethyl)-2-chloropyrimidine (47.6 g, 229.45 mmol, 1.0 equiv) was added at 0 °C. The reaction mixture was stirred at room temperature for 15.5 h. The mixture was then poured into H₂O (1600 mL) and the aqueous phase was extracted with EtOAc (3 x 300 mL). The combined organic phase was washed with brine (2 x 200 mL), dried with anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by silica gel chromatography (1/0 to 0/1 petroleum ether/EtOAc) to give the product (70 g, crude) as a yellow solid, which was used to next step directly.


Step 3: Synthesis of tert-butyl N-tert-butoxycarbonyl-N-[(2-piperazin-1-ylpyrimidin-5-yl)methyl]carbamate

[00584] To a solution of 1-benzylpiperazine (30.44 g, 122.16 mmol, 1.0 equiv, 2HCl) in MeCN (550 mL) was added *tert*-butyl *N-tert*-butoxycarbonyl-N-((2-chloropyrimidin-5-yl)methyl)carbamate (42 g, 122.16 mmol, 1.0 equiv) and K₂CO₃ (84.42 g, 610.81 mmol, 5.0 equiv). The mixture was stirred at 80 °C for 61 h. The reaction mixture was then diluted with EtOAc (150 mL) and the mixture was filtered. The resulting solution was concentrated under reduced pressure to give the crude product. The residue was purified by silica gel chromatography (1/0 to 0/1 petroleum ether/EtOAc) to give the product (45 g, 74% yield) as a white solid.

Step 4: Synthesis of tert-butyl N-tert-butoxycarbonyl-N-[(2-piperazin-1-yl)pyrimidin-5-yl)methyl]carbamate

[00585] To a solution of *tert*-butyl N-[[2-(4-benzylpiperazin-1-yl)pyrimidin-5-yl)methyl]-N-*tert*-butoxycarbonyl-carbamate (24 g, 49.63 mmol, 1.0 equiv) in MeOH (600 mL) was added Pd/C (24 g, 47.56 mmol, 10 wt.%, 1.0 equiv) under argon. The mixture was degassed under reduced pressure and purged with H₂ three times. The mixture was stirred under H₂ (50 psi) at 50 °C for 19 h. The reaction mixture was cooled to room temperature, filtered, and the filter cake was washed with MeOH (500 mL). The resulting solution was concentrated under reduced pressure. The residue was purified by silica gel chromatography (1/0 to 0/1 EtOAc/MeOH) to give the product (25.5 g, 68% yield) as a white solid.

Building block B. 2-(4-(5-(((*tert*-butoxycarbonyl)amino)methyl)pyrimidin-2-yl)piperazin-1-yl)pyrimidine-5-carboxylic acid.



*Step 1: Synthesis of ethyl 2-(4-(5-((bis(*tert*-butoxycarbonyl)amino)methyl)pyrimidin-2-yl)piperazin-1-yl)pyrimidine-5-carboxylate*


[00586] To a solution of ethyl 2-chloropyrimidine-5-carboxylate (2.37 g, 12.71 mmol, 1.0 equiv) and *tert*-butyl N-*tert*-butoxycarbonyl-N-((2-piperazin-1-yl)pyrimidin-5-yl)methylcarbamate (5 g, 12.71 mmol, 1.0 equiv) in MeCN (80 mL) was added K₂CO₃ (5.27 g, 38.12 mmol, 3.0 equiv). The mixture was stirred at 80 °C for 16 h. The reaction mixture was then poured into H₂O (200 mL) and the suspension was filtered. The filtrate was washed with H₂O (80 mL) and dried under reduced pressure to give the product (6.1 g, 87% yield) as a white solid.

*Step 2: Synthesis of 2-(4-(5-(((*tert*-butoxycarbonyl)amino)methyl)pyrimidin-2-yl)piperazin-1-yl)pyrimidine-5-carboxylic acid*

[00587] To a solution of ethyl 2-(4-(5-((bis(*tert*-butoxycarbonyl)amino)methyl)pyrimidin-2-yl)piperazin-1-yl)pyrimidine-5-carboxylate (5 g, 9.20 mmol, 1.0 equiv) in H₂O (50 mL), EtOH (15 mL) and THF (50 mL) was added LiOH•H₂O (1.54 g, 36.79 mmol, 4.0 equiv). The reaction mixture was stirred at 55 °C for 16 h. The mixture was then concentrated to remove

THF and EtOH and then the mixture was diluted with H₂O (55 mL) and was acidified (pH=3) with aqueous HCl (1 N). The mixture was filtered and the filter cake was washed with H₂O (36 mL). The filter cake was dried under reduced pressure to give the product (2.7 g, 69.3%) as a white solid. LCMS (ESI) *m/z*: [M + H] calcd for C₁₉H₂₅N₇O₄: 416.21; found 416.1.

Building block C. *tert*-butyl 2-(piperazin-1-yl)-7,8-dihydropyrido[4,3-d]pyrimidine-6(5H)-carboxylate.




*Step 1: Synthesis of *tert*-butyl 2-(4-benzylpiperazin-1-yl)-7,8-dihydropyrido[4,3-d]pyrimidine-6(5H)-carboxylate*

[00588] To a solution of *tert*-butyl 2-chloro-7,8-dihydropyrido[4,3-d]pyrimidine-6(5H)-carboxylate (15 g, 55.61 mmol, 1.0 equiv) in MeCN (150 mL) was added 1-benzylpiperazine (11.76 g, 66.73 mmol, 1.2 equiv) and K₂CO₃ (46.12 g, 333.67 mmol, 6.0 equiv). The mixture was stirred at 80 °C for 27 h. The reaction mixture was diluted with EtOAc (200 mL), filtered and concentrated under reduced pressure. The crude product was purified by silica gel chromatography (1/0 to 0/1 petroleum ether/EtOAc) to give the product (20.2 g, 80% yield) as a white solid. LCMS (ESI) *m/z*: [M + H] calcd for C₂₃H₃₁N₅O₂: 410.26; found 410.1.

*Step 2: Synthesis of *tert*-butyl 2-(piperazin-1-yl)-7,8-dihydropyrido[4,3-d]pyrimidine-6(5H)-carboxylate*

[00589] To a solution of *tert*-butyl 2-(4-benzylpiperazin-1-yl)-7,8-dihydropyrido[4,3-d]pyrimidine-6(5H)-carboxylate (8 g, 19.53 mmol, 1.0 equiv) in MeOH (200 mL) was added Pd/C (8 g, 19.53 mmol, 10 wt.%, 1.0 equiv) under argon. The mixture was degassed and purged with H₂ three times. The mixture was stirred under H₂ (50 psi) at 50 °C for 19 h. The reaction mixture was cooled to room temperature, filtered through a pad of Celite and the filter cake was washed with MeOH (150 mL). The resulting solution was concentrated under reduced pressure. The crude product was washed with petroleum ether (60 mL) to give the product (9.25 g, 72% yield) as a white solid. LCMS (ESI) *m/z*: [M + H] calcd for C₁₆H₂₅N₅O₂: 320.21; found 320.2.

Building block D. 2-(4-(6-(*tert*-butoxycarbonyl)-5,6,7,8-tetrahydropyrido[4,3-d]pyrimidin-2-yl)piperazin-1-yl)pyrimidine-5-carboxylic acid.




Step 1: Synthesis of tert-butyl 2-(4-(5-(ethoxycarbonyl)pyrimidin-2-yl)piperazin-1-yl)-7,8-dihydropyrido[4,3-d]pyrimidine-6(5H)-carboxylate

[00590] To a solution of ethyl 2-chloropyrimidine-5-carboxylate (4.09 g, 21.92 mmol, 1.0 equiv) in dioxane (80 mL) was added *tert*-butyl 2-(piperazin-1-yl)-7,8-dihydropyrido[4,3-d]pyrimidine-6(5H)-carboxylate (7 g, 21.92 mmol, 1.0 equiv) and Et₃N (9.15 mL, 65.75 mmol, 3.0 equiv). The mixture was stirred at 90 °C for 64 h. The solution was poured into H₂O (200 mL) and then the mixture was filtered and the filter cake was washed with H₂O (100 mL) followed by petroleum ether (60 mL). The filter cake was dried under reduced pressure to give the product (10.1 g, 92% yield) as a brown solid. LCMS (ESI) *m/z*: [M + H] calcd for C₂₃H₃₁N₇O₄: 470.25; found 470.4.

Step 2: Synthesis of 2-(4-(6-(tert-butoxycarbonyl)-5,6,7,8-tetrahydropyrido[4,3-d]pyrimidin-2-yl)piperazin-1-yl)pyrimidine-5-carboxylic acid

[00591] To a solution of *tert*-butyl 2-(4-(5-(ethoxycarbonyl)pyrimidin-2-yl)piperazin-1-yl)-7,8-dihydropyrido[4,3-d]pyrimidine-6(5H)-carboxylate (6.0 g, 12.78 mmol, 1.0 equiv) in THF (40 mL), EtOH (20 mL) and H₂O (40 mL) was added LiOH•H₂O (1.07 g, 25.56 mmol, 2.0 equiv). The reaction mixture was stirred at 35 °C for 15 h. The mixture was then concentrated under reduced pressure to remove THF and EtOH. The mixture was then diluted with H₂O (500 mL) and was adjusted to pH 3 with aqueous HCl (1 N). The mixture was filtered and the filter cake was washed with H₂O (80 mL) followed by petroleum ether (80 mL). The filter cake was dried under reduced pressure to give the product (3.8 g, 65% yield) as a white solid. LCMS (ESI) *m/z*: [M + H] calcd for C₂₁H₂₇N₇O₄: 442.22; found 442.3.

Building block E. *tert*-butyl methyl((2-(piperazin-1-yl)pyrimidin-5-yl)methyl)carbamate.



Step 1: Synthesis of (2-chloropyrimidin-5-yl)methanamine

[00592] To a solution of *tert*-butyl *N*-*tert*-butoxycarbonyl-*N*-((2-chloropyrimidin-5-yl)methyl)carbamate (28 g, 81.44 mmol, 1.0 equiv) in EtOAc (30 mL) was added HCl in EtOAc (260 mL). The reaction mixture was stirred at room temperature for 5 h. The reaction mixture was filtered and the filter cake was washed with EtOAc (100 mL). The solid cake was dried under reduced pressure to give the product (14.3 g, 96.6% yield, HCl) as a white solid.

Step 2: Synthesis of tert-butyl ((2-chloropyrimidin-5-yl)methyl)carbamate

[00593] To a solution of (2-chloropyrimidin-5-yl)methanamine (13 g, 72.21 mmol, 1.0 equiv, HCl) in DCM (130 mL) was added DIPEA (20.41 mL, 144.42 mmol, 1.8 equiv) and Boc₂O (16.59 mL, 72.21 mmol, 1.0 equiv), then the mixture was stirred at room temperature for 3 h. The reaction mixture was added to H₂O (100 mL) and then the aqueous layer was separated and extracted with DCM (2 x 100 mL). Then combined organic phase was washed with sat. NH₄Cl (2 x 200 mL) and brine (2 x 200 mL), dried with anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by silica gel chromatography (1/0 to 1/1 petroleum ether/EtOAc) to give the product (12 g, 68.2% yield) as a white solid.

Step 3: Synthesis of tert-butyl ((2-chloropyrimidin-5-yl)methyl)(methyl)carbamate

[00594] To a solution of *tert*-butyl ((2-chloropyrimidin-5-yl)methyl)carbamate (11 g, 45.14 mmol, 1.0 equiv) and MeI (14.05 mL, 225.70 mmol, 5.0 equiv) in THF (150 mL) was added NaH (1.99 g, 49.65 mmol, 60 wt.%, 1.1 equiv) at 0 °C. The mixture was stirred at 0 °C for 3 h and then the reaction was quenched with H₂O (100 mL). The aqueous phase was extracted with EtOAc (3 x 150 mL) and the combined organic phase was washed with brine (50 mL), dried with anhydrous Na₂SO₄, filtered and concentrated under reduced pressure.

The residue was purified by silica gel chromatography (1/0 to 3/1 petroleum ether/EtOAc) to give the product (9 g, 77.4% yield) as a white solid.


Step 4: Synthesis of tert-butyl ((2-(4-benzylpiperazin-1-yl)pyrimidin-5-yl)methyl)(methyl)carbamate

[00595] To a solution of *tert*-butyl ((2-chloropyrimidin-5-yl)methyl)(methyl)carbamate (9 g, 34.92 mmol, 1.0 equiv) in MeCN (90 mL) was added 1-benzylpiperazine (8.70 g, 34.92 mmol, 1.0 equiv, 2HCl), and K₂CO₃ (24.13 g, 174.61 mmol, 5.0 equiv). The reaction mixture was stirred at 80 °C for 20 h. The mixture was then filtered and concentrated under reduced pressure. The residue was purified by silica gel chromatography (1/0 to 1/1 petroleum ether/EtOAc) to give the product (12 g, 86.4% yield) as a yellow oil.

Step 5: Synthesis of tert-butyl methyl((2-(piperazin-1-yl)pyrimidin-5-yl)methyl)carbamate

[00596] To a solution of *tert*-butyl ((2-(4-benzylpiperazin-1-yl)pyrimidin-5-yl)methyl)(methyl)carbamate (12 g, 30.19 mmol, 1.0 equiv) in MeOH (120 mL) was added Pd/C (2 g, 10 wt.%). The suspension was degassed and purged with H₂ and then the mixture was stirred under H₂ (15 psi) at room temperature for 3 h. The reaction mixture was filtered through Celite and the filtrate was concentrated under reduced pressure. The residue was purified by silica gel chromatography 1/0 to 1/1 petroleum ether/EtOAc) to give semi-pure material (9 g) as a yellow oil. Petroleum ether was added to the residue and the solution was stirred at -60 °C until solid appeared. The suspension was filtered and the filtrate was concentrated under reduced pressure to give the product (4.07 g, 55.6% yield) as a yellow oil. LCMS (ESI) *m/z*: [M + H] calcd for C₁₅H₂₅N₅O₂: 308.21; found 308.1.

Building block F. 2-(4-(5-(((*tert*-butoxycarbonyl)(methyl)amino)methyl) pyrimidin-2-yl)piperazin-1-yl)pyrimidine-5-carboxylic acid.



*Step 1: Synthesis of ethyl 2-(4-(5-(((*tert*-butoxycarbonyl)(methyl)amino)methyl) pyrimidin-2-yl)piperazin-1-yl)pyrimidine-5-carboxylate*


[00597] To a mixture of *tert*-butyl methyl((2-(piperazin-1-yl)pyrimidin-5-yl)methyl)carbamate (4.3 g, 13.99 mmol, 1.0 equiv) and ethyl 2-chloropyrimidine-5-

carboxylate (2.87 g, 15.39 mmol, 1.1 equiv) in MeCN (20 mL) was added K₂CO₃ (3.87 g, 27.98 mmol, 2.0 equiv). The mixture was stirred at 80 °C for 12 h. The reaction mixture then cooled to room temperature and was filtered. The filtrate was concentrated under reduced pressure and the crude product was purified by silica gel chromatography (1/0 to 1/1 petroleum ether/EtOAc) to give the product (4.7 g, 71.3% yield) as a white solid.

*Step 2: Synthesis of 2-(4-(5-(((*tert*-butoxycarbonyl)(methyl)amino)methyl)pyrimidin-2-yl)piperazin-1-yl)pyrimidine-5-carboxylic acid*

[00598] To a solution of ethyl 2-(4-(5-(((*tert*-butoxycarbonyl)(methyl)amino)methyl)pyrimidin-2-yl)piperazin-1-yl)pyrimidine-5-carboxylate (6 g, 13.11 mmol, 1.0 equiv) in THF (100 mL), EtOH (30 mL), and H₂O (30 mL) was added LiOH•H₂O (1.10 g, 26.23 mmol, 2.0 equiv). The mixture was stirred at room temperature for 16 h. The mixture was then concentrated under reduced pressure to remove THF and EtOH and then neutralized by the addition of 1N HCl. The resulting precipitate was collected by filtration to give the product (5.11 g, 90.1% yield) as a white solid. LCMS (ESI) *m/z*: [M + H] calcd for C₂₀H₂₇N₇O₄: 430.22; found 430.2.

Building block G. *tert*-butyl N-*tert*-butoxycarbonyl-N-((2-(2-(((*tert*-butyl(diphenyl)silyl)oxymethyl)piperazin-1-yl)pyrimidin-5-yl)methyl)carbamate.



Step 1: Synthesis of tert-butyl N-((2-(4-benzyl-2-(hydroxymethyl)piperazin-1-yl)pyrimidin-5-yl)methyl)-N-tert-butoxycarbonyl-carbamate

[00599] To a solution of *tert*-butyl N-*tert*-butoxycarbonyl-N-((2-chloropyrimidin-5-yl)methyl)carbamate (18.33 g, 53.32 mmol, 1.1 equiv) and (4-benzylpiperazin-2-yl)methanol

(10 g, 48.48 mmol, 1.0 equiv) in DMF (100 mL) was added K₂CO₃ (13.40 g, 96.95 mmol, 2.0 equiv). The mixture was stirred at 100 °C for 12 h. The reaction mixture was then cooled to room temperature and H₂O (100 mL) was added. The aqueous layer was extracted with EtOAc (2 x 150 mL) and the combined organic layer was washed with brine (20 mL), dried with Na₂SO₄, filtered and the filtrate was concentrated under reduced pressure to give the product (7.3 g, 29.3% yield) as a yellow oil. LCMS (ESI) *m/z*: [M + H] calcd for C₂₇H₃₉N₅O₅: 514.31; found 514.5


Step 2: Synthesis of tert-butyl N-((2-(4-benzyl-2-((tert-butyl(diphenyl)silyl)oxymethyl)piperazin-1-yl)pyrimidin-5-yl)methyl)-N-tert-butoxycarbonyl-carbamate

[00600] To a solution of *tert*-butyl N-((2-(4-benzyl-2-(hydroxymethyl)piperazin-1-yl)pyrimidin-5-yl)methyl)-N-*tert*-butoxycarbonyl-carbamate (2.3 g, 4.48 mmol, 1.0 equiv) in DCM (30 mL) was added imidazole (609.69 mg, 8.96 mmol, 2.0 equiv) and TBDPSCl (1.73 mL, 6.72 mmol, 1.5 equiv). The reaction mixture was stirred at room temperature for 2 h. The mixture was then washed with H₂O (100mL) and the aqueous phase extracted with EtOAc (2 x 60 mL). The combined organic phase was washed with brine (20 mL), dried with Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by silica gel chromatography (20/1 to 3/1 petroleum ether/EtOAc) to give the product (4 g, 59.4% yield) as a yellow oil. LCMS (ESI) *m/z*: [M + H] calcd for C₄₃H₅₇N₅O₅Si: 752.42; found 752.4.

Step 3: Synthesis of tert-butyl N-tert-butoxycarbonyl-N-((2-(2-((tert-butyl(diphenyl)silyl)oxymethyl)piperazin-1-yl)pyrimidin-5-yl)methyl)carbamate

[00601] To a solution of *tert*-butyl N-((2-(4-benzyl-2-((tert-butyl(diphenyl)silyl)oxymethyl)piperazin-1-yl)pyrimidin-5-yl)methyl)-N-*tert*-butoxycarbonyl-carbamate (3.3 g, 4.39 mmol, 1.0 equiv) in EtOH (10 mL) was added Pd(OH)₂/C (1 g, 10 wt.%). The mixture was heated to 50 °C under H₂ (30 psi) for 30 h. The mixture was then cooled to room temperature, filtered through Celite, and concentrated under reduced pressure. The residue was purified by silica gel chromatography (20/1 to 3/1 EtOAc/EtOH) to give the product (1.44 g, 45.6% yield) as a yellow solid. LCMS (ESI) *m/z*: [M + H] calcd for C₃₆H₅₁N₅O₅Si: 662.38; found 662.3.

Building block H. 2-(4-(5-(((*tert*-butoxycarbonyl)amino)methyl)pyrimidin-2-yl)-3-(hydroxymethyl)piperazin-1-yl)pyrimidine-5-carboxylic acid.



Step 1: Synthesis of *tert*-butyl N-*tert*-butoxycarbonyl-N-((2-(2-(hydroxymethyl)piperazin-1-yl)pyrimidin-5-yl)methyl)carbamate

[00602] To a solution of *tert*-butyl N-((2-(4-benzyl-2-(hydroxymethyl)piperazin-1-yl)pyrimidin-5-yl)methyl)-N-*tert*-butoxycarbonyl-carbamate (3 g, 5.84 mmol, 1.0 equiv) in EtOH (40 mL) was added Pd/C (2 g, 10 wt.%). The suspension was degassed and purged with H₂, then stirred under H₂ (50 psi) at 30 °C for 16 h. The reaction mixture was cooled to room temperature and filtered through Celite and then concentrated under reduced pressure to give the product (1.6 g, crude) as a yellow oil. LCMS (ESI) *m/z*: [M + H] calcd for C₂₀H₃₃N₅O₅: 424.26; found 424.3.

Step 2: Synthesis of ethyl 2-(4-(5-((bis(*tert*-butoxycarbonyl)amino)methyl)pyrimidin-2-yl)-3-(hydroxymethyl)piperazin-1-yl)pyrimidine-5-carboxylate


[00603] To a solution of *tert*-butyl N-*tert*-butoxycarbonyl-N-((2-(2-(hydroxymethyl)piperazin-1-yl)pyrimidin-5-yl)methyl)carbamate (1.4 g, 3.31 mmol, 1.0 equiv) in MeCN (20 mL) was added K₂CO₃ (2.28 g, 16.53 mmol, 5.0 equiv) and ethyl 2-chloropyrimidine-5-carboxylate (616.84 mg, 3.31 mmol, 1.0 equiv). The solution was stirred at 80 °C for 4 h. The mixture was cooled to room temperature and poured into H₂O (30 mL). The aqueous layer was extracted with EtOAc (2 x 30 mL) and the combined organic layer was washed with brine (20 mL), dried with Na₂SO₄, filtered and concentrated under reduced pressure. The mixture was purified by silica gel chromatography (20/1 to 3/1 petroleum

ether/EtOAc) to give the product (1.6 g, 66.7% yield) as a light yellow solid. LCMS (ESI) m/z : $[M + H]$ calcd for $C_{27}H_{39}N_7O_7$: 574.30; found 574.4.

Step 3: Synthesis of 2-(4-(5-(((*tert*-butoxycarbonyl)amino)methyl)pyrimidin-2-yl)-3-(hydroxymethyl)piperazin-1-yl)pyrimidine-5-carboxylic acid

[00604] To a solution of ethyl 2-(4-(5-((bis(*tert*-butoxycarbonyl)amino)methyl)pyrimidin-2-yl)-3-(hydroxymethyl)piperazin-1-yl)pyrimidine-5-carboxylate (1.4 g, 2.44 mmol, 1.0 equiv) in THF (6 mL) and EtOH (6 mL) at 0 °C was added a solution of LiOH·H₂O (512.07 mg, 12.20 mmol, 5.0 equiv) in H₂O (3 mL). The reaction mixture was warmed to room temperature and stirred for 2 h. The mixture was then concentrated under reduced pressure to remove THF and EtOH. The aqueous phase was adjusted to pH 3 with 0.1 M HCl and the resulting suspension was filtered. The solid cake was dried under reduced pressure to give the product (613.14 mg, 55.6% yield) as a white solid. LCMS (ESI) m/z : $[M + H]$ calcd for $C_{20}H_{27}N_7O_5$: 446.22; found 446.2.

Building block I. *tert*-butyl N-[(*tert*-butoxy)carbonyl]-N-({2-[(3*R*)-3-(hydroxymethyl)piperazin-1-yl]pyrimidin-5-yl)methyl}carbamate.




Step 1: Synthesis of (*R*)-*tert*-butyl N-*tert*-butoxycarbonyl-((2-(3-(((*tert*-butyldiphenylsilyl)oxy)methyl)piperazin-1-yl)pyrimidin-5-yl)methyl)carbamate

[00605] To a solution of *tert*-butyl N-*tert*-butoxycarbonyl-((2-chloropyrimidin-5-yl)methyl)carbamate (24.24 g, 70.51 mmol, 1.0 equiv) in MeCN (300 mL) was added (*R*)-2-(((*tert*-butyldiphenylsilyl)oxy)methyl)piperazine (25 g, 70.51 mmol, 1.0 equiv) and K₂CO₃ (29.24 g, 211.53 mmol, 3.0 equiv). The mixture was stirred at 80 °C for 16 h. The reaction mixture was then cooled to room temperature, diluted with EtOAc (200 mL), filtered and the filtrate was concentrated under reduced pressure. Purification by silica gel chromatography (0→100% EtOAc/petroleum ether) afforded the desired product (46.5 g, 94% yield) as a white solid.

Step 2: Synthesis of *tert*-butyl N-[(*tert*-butoxy)carbonyl]-N-({2-[(3*R*)-3-(hydroxymethyl)piperazin-1-yl]pyrimidin-5-yl)methyl}carbamate

[00606] To a solution of (*R*)-*tert*-butyl-N-*tert*-butoxycarbonyl-((2-(3-(((*tert*-butyldiphenylsilyl)oxy)methyl)piperazin-1-yl)pyrimidin-5-yl)methyl)carbamate (12 g, 18.13 mmol, 1.0 equiv) in THF (120 mL) was added TBAF (1 M, 23.93 mL, 1.3 equiv). The mixture was stirred at room temperature for 2 h. The reaction mixture was then poured into H₂O (300 mL) and the aqueous phase was extracted with EtOAc (3 x 80 mL). The combined organic phases were combined, washed with brine (80 mL), dried, filtered and the filtrate was concentrated under reduced pressure. Purification by silica gel chromatography (0→20% MeOH/DCM) afforded the desired product (5 g, 64% yield) as a yellow solid.

Building block J. 2-{4-[5-({[(*tert*-butoxy)carbonyl]amino)methyl]pyrimidin-2-yl]piperazin-1-yl}pyrimidine-5-carboxylic acid.



Step 1: Synthesis of (*R*)-ethyl 2-(4-(5-(((di-*tert*-butoxycarbonyl)amino)methyl)pyrimidin-2-yl)-2-(((*tert*-butyldiphenylsilyl)oxy)methyl)piperazin-1-yl)pyrimidine-5-carboxylate

[00607] To a solution of (*R*)-*tert*-butyl-N-*tert*-butoxycarbonyl-N-((2-(3-(((*tert*-butyldiphenylsilyl)oxy)methyl)piperazin-1-yl)pyrimidin-5-yl)methyl)carbamate (31.5 g, 45.21 mmol, 1.0 equiv) in MeCN (350 mL) was added ethyl 2-chloropyrimidine-5-carboxylate (8.44 g, 45.21 mmol, 1.0 equiv) and K₂CO₃ (18.75 g, 135.63 mmol, 3.0 equiv). The mixture was stirred at 80 °C for 16 h. The reaction mixture was then cooled to room temperature, diluted with EtOAc (150 mL), and filtered to remove inorganic salts. The filtrate was then concentrated under reduced pressure. Purification by silica gel chromatography (0→100% EtOAc/petroleum ether) afforded the desired product (33.5 g, 89% yield).


Step 2: Synthesis of (*R*)-ethyl 2-(4-(5-(((di-*tert*-butoxycarbonyl)amino)methyl)pyrimidin-2-yl)-2-(hydroxymethyl)piperazin-1-yl)pyrimidine-5-carboxylate

[00608] To a solution of (*R*)-ethyl 2-(4-(5-(((di-*tert*-butoxycarbonyl)amino)methyl)pyrimidin-2-yl)-2-(((*tert*-butyldiphenylsilyl)oxy)methyl)piperazin-1-yl)pyrimidine-5-carboxylate (36.5 g, 44.95 mmol, 1.0 equiv) in THF (300 mL) was added TBAF (1 M, 59.33 mL, 1.32 equiv). The mixture was stirred at room temperature for 6 h, at which point the reaction mixture was poured into H₂O (500 mL). The aqueous phase was separated and extracted with EtOAc (3 x 150 mL) and the combined organic layers were washed with brine (150 mL), dried, filtered, and the filtrate was concentrated under reduced pressure. Purification by silica gel chromatography (0→100% EtOAc/petroleum ether) afforded the desired product (17 g, 64% yield) as a yellow oil.

Step 3: Synthesis of (R)-2-(4-(5-(((tert-butoxycarbonyl)amino)methyl)pyrimidin-2-yl)-2-(hydroxymethyl)piperazin-1-yl)pyrimidine-5-carboxylic acid


[00609] To a solution of (*R*)-ethyl 2-(4-(5-(((di-*tert*-butoxycarbonyl)amino)methyl)pyrimidin-2-yl)-2-(hydroxymethyl)piperazin-1-yl)pyrimidine-5-carboxylate (17 g, 29.64 mmol, 1.0 equiv) in H₂O (160 mL), EtOH (80 mL) and THF (160 mL) was added LiOH•H₂O (4.97 g, 118.54 mmol, 4.0 equiv). The reaction mixture was stirred at 55 °C for 16 h. To the mixture was then added LiOH•H₂O (1.01 g, 24.00 mmol, 0.81 equiv) and the reaction mixture was stirred at 55 °C for an additional 9 h. The mixture was cooled to room temperature, diluted with H₂O (150 mL), and concentrated under reduced pressure to remove THF and EtOH. The mixture was acidified (pH = 5) with 1 N HCl, filtered, and the filter cake washed with H₂O (2 x 30 mL). The filter cake was dried under reduced pressure to afford the desired product (9.2 g, 67% yield) as a white solid. LCMS (ESI) *m/z*: [M + H] calcd for C₂₀H₂₇N₇O₅: 446.22; found 446.1.

Building block K. *tert*-butyl N-[(*tert*-butoxy)carbonyl]-N-(2-[(3*S*)-3-(hydroxymethyl)piperazin-1-yl]pyrimidin-5-yl)methyl)carbamate.




[00610] This building block is prepared by a process similar to that for Building block I by utilizing [(2*S*)-piperazin-2-yl]methanol.

Building block L. 2-[(2*S*)-4-[5-({[(*tert*-butoxy)carbonyl]amino}methyl)pyrimidin-2-yl]-2-(hydroxymethyl)piperazin-1-yl]pyrimidine-5-carboxylic acid.



[00611] This building block is prepared from Building block K by a process similar to that for Building block J.

Building block M. *tert*-butyl 2-[(3*R*)-3-(hydroxymethyl)piperazin-1-yl]-5*H*,6*H*,7*H*,8*H*-pyrido[4,3-*d*]pyrimidine-6-carboxylate.



Step 1: Synthesis of (*R*)-*tert*-butyl 2-(3-(((*tert*-butyldiphenylsilyl)oxy)methyl)piperazin-1-yl)-7,8-dihydropyrido[4,3-*d*]pyrimidine-6(5*H*)-carboxylate


[00612] To a solution of (*R*)-2-(((*tert*-butyldiphenylsilyl)oxy)methyl)piperazine (25 g, 70.51 mmol, 1.0 equiv) in MeCN (250 mL) was added K₂CO₃ (29.24 g, 211.53 mmol, 3.0 equiv) and *tert*-butyl 2-chloro-7,8-dihydropyrido[4,3-*d*]pyrimidine-6(5*H*)-carboxylate (17.12 g, 63.46 mmol, 0.9 equiv). The mixture was stirred at 80 °C for 17 h. The reaction mixture was then cooled to room temperature, filtered, and the filtrate was concentrated under reduced pressure. Purification by silica gel chromatography (0→100% EtOAc/petroleum ether) afforded the desired product (31 g, 73.5% yield) as a white solid. LCMS (ESI) *m/z*: [M + H] calcd for C₃₃H₄₅N₅O₃Si: 588.34; found 588.2.

Step 2: Synthesis of (*R*)-*tert*-butyl 2-(3-(hydroxymethyl)piperazin-1-yl)-7,8-dihydropyrido[4,3-*d*]pyrimidine-6(5*H*)-carboxylate

[00613] To a mixture of (*R*)-*tert*-butyl 2-(3-(((*tert*-butyldiphenylsilyl)oxy)methyl)piperazin-1-yl)-7,8-dihydropyrido[4,3-*d*]pyrimidine-6(5*H*)-carboxylate (12 g, 20.41 mmol, 1.0 equiv) in THF (120 mL) was added TBAF (1.0 M, 24.50


mL, 1.2 equiv). The mixture was stirred at room temperature for 5 h. The mixture was poured into H₂O (100 mL), and the aqueous phase was extracted with EtOAc (2 x 100 mL). The combined organic phases were washed with brine (100 mL), dried, filtered, and the filtrate was concentrated under reduced pressure. Purification by silica gel chromatography (0→10% MeOH/DCM) afforded the desired product (6 g, 84.1% yield) as a white solid. LCMS (ESI) *m/z*: [M + H] calcd for C₁₇H₂₇N₅O₃: 350.22; found 350.2.

Building block N. 2-[(2*R*)-4-{6-[(*tert*-butoxy)carbonyl]-5*H*,6*H*,7*H*,8*H*-pyrido[4,3-*d*]pyrimidin-2-yl}-2-(hydroxymethyl)piperazin-1-yl]pyrimidine-5-carboxylic acid.




[00614] This building block is prepared from Building block M by a process similar to that for Building block J.

Building block O. *tert*-butyl 2-[(3*S*)-3-(hydroxymethyl)piperazin-1-yl]-5*H*,6*H*,7*H*,8*H*-pyrido[4,3-*d*]pyrimidine-6-carboxylate.




[00615] This building block is prepared by a process similar to that for Building block I by utilizing *tert*-butyl 2-chloro-7,8-dihydropyrido[4,3-*d*]pyrimidine-6(5*H*)-carboxylate and [(2*S*)-piperazin-2-yl]methanol.

Building block P. 2-[(2*S*)-4-{6-[(*tert*-butoxy)carbonyl]-5*H*,6*H*,7*H*,8*H*-pyrido[4,3-*d*]pyrimidin-2-yl}-2-(hydroxymethyl)piperazin-1-yl]pyrimidine-5-carboxylic acid.



[00616] This building block is prepared from Building block O by a process similar to that for Building block J.

Building block Q. *tert*-butyl N-[(*tert*-butoxy)carbonyl]-N-({2-[(3*S*)-3-[(dimethylamino)methyl]piperazin-1-yl]pyrimidin-5-yl}methyl)carbamate.



Step 1: Synthesis of (R)-dibenzyl 2-(dimethylcarbamoyl)piperazine-1,4-dicarboxylate

[00617] To a solution of CDI (12.21 g, 75.30 mmol, 1.2 equiv) in DCM (300 mL) at 0 °C was added (*R*)-1,4-bis((benzyloxy)carbonyl)piperazine-2-carboxylic acid (25 g, 62.75 mmol, 1.0 equiv). The mixture was stirred at 0 °C for 0.5 h, at which time dimethylamine (8.51 mL, 92.87 mmol, 1.5 equiv, HCl) was added. The reaction mixture was warmed to room temperature and stirred for 12 h. The reaction mixture was then added to H₂O (200 mL), and the aqueous layer was separated and extracted with DCM (2 x 200 mL). The combined organic phases were washed with brine (2 x 50 mL), dried, filtered, and the filtrate was concentrated under reduced pressure. Purification by silica gel chromatography (50→100% EtOAc/petroleum ether) afforded the desired product (23.5 g, 88.0% yield) as a yellow oil.

Step 2: Synthesis of (S)-dibenzyl 2-((dimethylamino)methyl)piperazine-1,4-dicarboxylate

[00618] To a solution of (*R*)-dibenzyl 2-(dimethylcarbamoyl)piperazine-1,4-dicarboxylate (28 g, 65.81 mmol, 1.0 equiv) in THF (300 mL) at 0 °C was added BH₃•Me₂S (10 M, 13.16 mL, 2.0 equiv). The reaction mixture was then stirred at 80 °C for 3 h. The reaction mixture was cooled to room temperature and then MeOH (50 mL) was added. After stirring for an additional 1 h the mixture was concentrated under reduced pressure. Purification by silica gel chromatography (50→100% EtOAc/petroleum ether) afforded the desired product (18 g, 66.5% yield) as a yellow oil.


Step 3: Synthesis of (R)-N,N-dimethyl-1-(piperazin-2-yl)methanamine

[00619] To a solution of (*S*)-dibenzyl 2-((dimethylamino)methyl)piperazine-1,4-dicarboxylate (18 g, 43.74 mmol, 1.0 equiv) in EtOAc (200 mL) was added Pd/C (1.5 g, 10 wt.%). The suspension was degassed under reduced pressure and purged with H₂ three times. The suspension was stirred under H₂ (30 psi) at 30 °C for 5 h. The reaction mixture was then filtered through celite and the filtrate was concentrated under reduced pressure to afford the desired product (6 g, 95.8% yield) as a yellow solid.

*Step 4: Synthesis of tert-butyl N-tert-butoxycarbonyl-N-((2-((3*S*)-3-((dimethylamino)methyl)piperazin-1-yl)pyrimidin-5-yl)methyl)carbamate*

[00620] To a solution of (*R*)-*N,N*-dimethyl-1-(piperazin-2-yl)methanamine (2.8 g, 19.55 mmol, 1.0 equiv) in MeCN (40 mL) was added *tert*-butyl *N-tert*-butoxycarbonyl-*N*-((2-chloropyrimidin-5-yl)methyl)carbamate (6.72 g, 19.55 mmol, 1.0 equiv) and K₂CO₃ (5.40 g, 39.10 mmol, 2.0 equiv). The mixture was stirred at 80 °C for 24 h. The mixture was then cooled to room temperature, filtered, and the filter cake washed with EtOAc (3 x 10 mL). The filtrate was then concentrated under reduced pressure. Purification by silica gel chromatography (0→100% MeOH/EtOAc) afforded the desired product (5.3 g, 57.8% yield) as a yellow oil. LCMS (ESI) *m/z*: [M + H] calcd for C₂₂H₃₈N₆O₄: 451.31; found 451.2.

Building block R. 2-[(2*S*)-4-[5-({[(*tert*-butoxy)carbonyl]amino}methyl)pyrimidin-2-yl]-2-[(dimethylamino)methyl]piperazin-1-yl]pyrimidine-5-carboxylic acid.




*Step 1: Synthesis of (S)-ethyl 2-(4-(5-(((*tert*-butoxycarbonyl)amino)methyl) pyrimidin-2-yl)-2-((dimethylamino)methyl)piperazin-1-yl)pyrimidine-5-carboxylate*

[00621] To a solution of (*S*)-*tert*-butyl-*N-tert*-butoxycarbonyl ((2-(3-((dimethylamino)methyl) piperazin-1-yl)pyrimidin-5-yl)methyl)carbamate (3.26 g, 7.24 mmol, 1.0 equiv) in DMF (30 mL) was added Et₃N (3.02 mL, 21.71 mmol, 3.0 equiv) and ethyl 2-chloropyrimidine-5-carboxylate (1.47 g, 7.86 mmol, 1.1 equiv). The mixture was stirred at 50 °C for 3 h and then concentrated under reduced pressure to afford the desired product (4.35 g, crude) as a solution in DMF (30 mL), which was used directly in the next step. LCMS (ESI) *m/z*: [M + H] calcd for C₂₉H₄₄N₈O₆: 601.35; found 601.5.

Step 2: Synthesis of (*S*)-2-(4-(5-(((*tert*-butoxycarbonyl)amino)methyl)-pyrimidin-2-yl)-2-((dimethylamino)methyl)piperazin-1-yl)pyrimidine-5-carboxylic acid


[00622] To a solution of (*S*)-ethyl 2-(4-(5-(((*bi-tert*-butoxycarbonyl)amino)methyl)-pyrimidin-2-yl)-2-((dimethylamino)methyl)piperazin-1-yl)pyrimidine-5-carboxylate (4.35 g, 7.24 mmol, 1.0 equiv) in DMF (30 mL) was added DMF (50 mL), EtOH (30 mL), and H₂O (30 mL). To the solution was then added LiOH•H₂O (3 g, 71.50 mmol, 9.9 equiv) at 50 °C. The reaction was stirred at 50 °C for 36 h. The mixture was then cooled to room temperature, neutralized with 0.5 N HCl, and concentrated under reduced pressure. Purification by reverse phase chromatography (2→30% MeCN/H₂O) afforded the desired product (1.15 g, 34% yield) as a white solid. LCMS (ESI) *m/z*: [M + H] calcd for C₂₂H₃₂N₈O₄: 473.26; found 473.3.

Building block S. *tert*-butyl N-[(*tert*-butoxy)carbonyl]-N-({2-[(3*R*)-3-[(dimethylamino)methyl]piperazin-1-yl]pyrimidin-5-yl}methyl)carbamate.




[00623] This building block is prepared by a process similar to that for Building block I by utilizing dimethyl({[(2*S*)-piperazin-2-yl]methyl})amine.

Building block T. 2-[(2*R*)-4-[5-({[(*tert*-butoxy)carbonyl]amino}methyl)pyrimidin-2-yl]-2-[(dimethylamino)methyl]piperazin-1-yl]pyrimidine-5-carboxylic acid.




[00624] This building block is prepared from Building block S by a process similar to that for Building block J.

Building block U. *tert*-butyl 2-[(3*S*)-3-[(dimethylamino)methyl]piperazin-1-yl]-5H,6H,7H,8H-pyrido[4,3-d]pyrimidine-6-carboxylate.



[00625] To a solution of *tert*-butyl 2-chloro-7,8-dihydropyrido[4,3-d]pyrimidine-6(5H)-carboxylate (4.80 g, 17.80 mmol, 1.4 equiv) in MeCN (45 mL) was added K₂CO₃ (10.42 g, 75.40 mmol, 3.0 equiv) and (*R*)-N,N-dimethyl-1-(piperazin-2-yl)methanamine (3.6 g, 25.13 mmol, 1.0 equiv). The mixture was stirred at 80 °C for 8 h. The mixture was then cooled to room temperature, filtered, and the filter cake was washed with EtOAc (50 mL). To the organic phase was added H₂O (50 mL) and the aqueous phase was extracted with EtOAc (3 x 50 mL). The combined organic phases were washed with brine (5 mL), dried, filtered and concentrated under reduced pressure. Purification by silica gel chromatography (8→67% EtOAc/petroleum ether) afforded the desired product (6.5 g, 63.5% yield) as a yellow oil.

Building block V. 2-[(2*S*)-4-{6-[(*tert*-butoxy)carbonyl]-5H,6H,7H,8H-pyrido[4,3-d]pyrimidin-2-yl}-2-[(dimethylamino)methyl]piperazin-1-yl]pyrimidine-5-carboxylic acid.



Step 1: Synthesis of (*S*)-*tert*-butyl 2-(3-((dimethylamino)methyl)-4-(5-(ethoxycarbonyl)pyrimidin-2-yl)piperazin-1-yl)-7,8-dihydropyrido[4,3-d]pyrimidine-6(5H)-carboxylate


[00626] To a solution of (*S*)-*tert*-butyl 2-(3-((dimethylamino)methyl)piperazin-1-yl)-7,8-dihydropyrido[4,3-d]pyrimidine-6(5H)-carboxylate (3 g, 7.97 mmol, 1.0 equiv) in DMF (70 mL) at 0 °C was added NaH (382.44 mg, 9.56 mmol, 60 wt.%, 1.2 equiv). The suspension was stirred at 0 °C for 0.5 h, then ethyl 2-chloropyrimidine-5-carboxylate (1.49 g, 7.97 mmol, 1 equiv) in DMF (50 mL) was added, dropwise. The mixture was warmed to room temperature and stirred for 5 h. The mixture was then cooled to 0 °C and poured into H₂O

(360 mL). The suspension was filtered, and the filter cake washed with H₂O (30 mL) and dried under reduced pressure. Purification by silica gel chromatography (6%→33% EtOAc/petroleum ether) afforded the desired product (1.8 g, 39.6% yield) as a brown oil.

Step 2: Synthesis of (S)-2-(4-(6-(tert-butoxycarbonyl)-5,6,7,8-tetrahydropyrido[4,3-d]pyrimidin-2-yl)-2-((dimethylamino)methyl)piperazin-1-yl)pyrimidine-5-carboxylic acid


[00627] To a solution of (*S*)-*tert*-butyl 2-(3-((dimethylamino)methyl)-4-(5-(ethoxycarbonyl)pyrimidin-2-yl)piperazin-1-yl)-7,8-dihydropyrido[4,3-d]pyrimidine-6(5H)-carboxylate (1.1 g, 2.09 mmol, 1.0 equiv) in THF (5 mL), EtOH (2.5 mL), and H₂O (2.5 mL) was added LiOH•H₂O (175.30 mg, 4.18 mmol, 2.0 equiv). The mixture was stirred at room temperature for 2 h, at which point the pH was adjusted to 7 by the addition of 1 N HCl at 0 °C. The mixture was concentrated under reduced pressure to remove THF and MeOH. The resulting suspension was filtered, and the filter cake was washed with H₂O (5 mL) and dried under reduced pressure to afford the desired product (680 mg, 65.3% yield) as a white solid. LCMS (ESI) *m/z*: [M + H] calcd for C₂₄H₃₄N₈O₄: 499.28; found 499.2.

Building block W. *tert*-butyl 2-[(3*R*)-3-[(dimethylamino)methyl]piperazin-1-yl]-5H,6H,7H,8H-pyrido[4,3-d]pyrimidine-6-carboxylate.




[00628] This building block is prepared by a process similar to that for Building block I by utilizing *tert*-butyl 2-chloro-7,8-dihydropyrido[4,3-d]pyrimidine-6(5H)-carboxylate and dimethyl({[(2*S*)-piperazin-2-yl]methyl})amine.

Building block X. 2-[(2*R*)-4-{6-[(*tert*-butoxy)carbonyl]-5H,6H,7H,8H-pyrido[4,3-d]pyrimidin-2-yl}-2-[(dimethylamino)methyl]piperazin-1-yl]pyrimidine-5-carboxylic acid.



[00629] This building block is prepared from Building block W by a process similar to that for Building block J.

Building block Y. *tert*-butyl (2*R*)-4-{6-[(*tert*-butoxy)carbonyl]-5*H*,6*H*,7*H*,8*H*-pyrido[4,3-*d*]pyrimidin-2-yl}piperazine-2-carboxylate.



Step 1: Synthesis of (*R*)-1,4-bis((benzyloxy)carbonyl)piperazine-2-carboxylic acid

[00630] To a solution of (*R*)-piperazine-2-carboxylic acid (70 g, 344.71 mmol, 1.0 equiv, 2HCl) in dioxane (1120 mL) and H₂O (700 mL) was added 50% aq. NaOH until the solution was pH=11. Benzyl chloroformate (156.82 mL, 1.10 mol, 3.2 equiv) was added and the reaction mixture was stirred at room temperature for 12 h. To the solution was then added H₂O (1200 mL) and the aqueous layer was washed with MTBE (3 x 800 mL). The aqueous layer was adjusted to pH=2 with concentrated HCl (12N) and extracted with EtOAc (2 x 1000 mL). The combined organic extracts were dried, filtered and the filtrate was concentrated under reduced pressure to afford the desired product (137 g, 99.8% yield) as a yellow solid. LCMS (ESI) *m/z*: [M + H] calcd for C₂₁H₂₂N₂O₆: 399.16; found 399.2.

Step 2: Synthesis of (*R*)-1,4-dibenzyl 2-*tert*-butyl piperazine-1,2,4-tricarboxylate

[00631] To a solution of (*R*)-1,4-bis((benzyloxy)carbonyl)piperazine-2-carboxylic acid (50 g, 125.50 mmol, 1.0 equiv) in toluene (500 mL) at 80 °C was added 1,1-di-*tert*-butoxy-N,N-dimethylmethanamine (57.17 mL, 238.45 mmol, 1.9 equiv). The solution was stirred at 80 °C for 2 h, at which point the reaction mixture was cooled to room temperature and partitioned between EtOAc (300 mL) and H₂O (500 mL). The aqueous layer was extracted with EtOAc (2 x 500 mL) and the combined organic layers were dried, filtered and the filtrate was concentrated under reduced pressure. Purification by silica gel chromatography (0%→25% EtOAc/petroleum ether) afforded the desired product (35 g, 61.2% yield) as a white solid. LCMS (ESI) *m/z*: [M + Na] calcd for C₂₅H₃₀N₂O₆: 477.20; found 477.1.


Step 3: Synthesis of (R)-tert-butyl piperazine-2-carboxylate

[00632] To a solution of (*R*)-1,4-dibenzyl 2-*tert*-butyl piperazine-1,2,4-tricarboxylate (35 g, 77.01 mmol, 1.0 equiv) in EtOAc (350 mL) was added Pd/C (10 g, 10 wt.%). The suspension was degassed under reduced pressure and purged with H₂ three times. The mixture was stirred under H₂ (30 psi) at 30 °C for 4 h. The reaction mixture was then filtered through celite, the residue was washed with MeOH (5 x 200 mL), and the filtrate was concentrated under reduced pressure to afford the desired product (14 g, 79.6% yield) as yellow oil. LCMS (ESI) *m/z*: [M + H] calcd for C₉H₁₈N₂O₂: 187.15; found 187.1.

Step 4: Synthesis of (R)-tert-butyl 2-(3-(tert-butoxycarbonyl)piperazin-1-yl)-7,8-dihydropyrido[4,3-d]pyrimidine-6(5H)-carboxylate

[00633] To a solution of *tert*-butyl (2*R*)-piperazine-2-carboxylate (12 g, 64.43 mmol, 1.0 equiv) in MeCN (200 mL) was added K₂CO₃ (17.81 g, 128.86 mmol, 2.0 equiv) and *tert*-butyl 2-chloro-7,8-dihydropyrido[4,3-d]pyrimidine-6(5H)-carboxylate (17.38 g, 64.43 mmol, 1.0 equiv). The reaction mixture was heated to 80 °C and stirred for 12 h. The reaction mixture was then cooled to room temperature and filtered, the residue was washed with EtOAc (3 x 150 mL), and the filtrate was concentrated under reduced pressure. Purification by silica gel chromatography (0%→100% EtOAc/petroleum ether) afforded the desired product (19 g, 69.2% yield) as a yellow solid. LCMS (ESI) *m/z*: [M + H] calcd for C₂₁H₃₃N₅O₄: 420.26; found 420.2.

Building block Z. 4-amino-2-[(2*R*)-2-[(*tert*-butoxy)carbonyl]-4-{6-[(*tert*-butoxy)carbonyl]-5*H*,6*H*,7*H*,8*H*-pyrido[4,3-d]pyrimidin-2-yl}piperazin-1-yl]pyrimidine-5-carboxylic acid.



Step 1: Synthesis of (R)-tert-butyl 2-(4-(4-amino-5-(ethoxycarbonyl)pyrimidin-2-yl)-3-(tert-butoxycarbonyl)piperazin-1-yl)-7,8-dihydropyrido[4,3-d]pyrimidine-6(5H)-carboxylate


[00634] To a stirred solution of (*R*)-*tert*-butyl 2-(3-(*tert*-butoxycarbonyl)piperazin-1-yl)-7,8-dihydropyrido[4,3-d]pyrimidine-6(5H)-carboxylate (12 g, 28.60 mmol, 1.0 equiv) in MeCN (150 mL) was added K₂CO₃ (7.91 g, 57.20 mmol, 2.0 equiv) and ethyl 4-amino-2-

chloropyrimidine-5-carboxylate (6.92 g, 34.32 mmol, 1.2 equiv). The reaction mixture was stirred at 80 °C for 12 h, at which point the reaction mixture was filtered and the filtrate was concentrated under reduced pressure. Purification by silica gel chromatography (0%→17% EtOAc/petroleum ether) afforded the desired product (16 g, 91.6% yield) as a yellow solid. LCMS (ESI) m/z : [M + H] calcd for C₂₈H₄₀N₈O₆: 585.32; found 585.1.

Step 2: Synthesis of (*R*)-4-amino-2-(2-(*tert*-butoxycarbonyl)-4-(6-(*tert*-butoxycarbonyl)-5,6,7,8-tetrahydropyrido[4,3-*d*]pyrimidin-2-yl)piperazin-1-yl)pyrimidine-5-carboxylic acid


[00635] To two separate batches run in parallel each containing a solution of (*R*)-*tert*-butyl 2-(4-(4-amino-5-(ethoxycarbonyl)pyrimidin-2-yl)-3-(*tert*-butoxycarbonyl)piperazin-1-yl)-7,8-dihydropyrido[4,3-*d*]pyrimidine-6(5H)-carboxylate (7 g, 11.97 mmol, 1.0 equiv) in THF (70 mL), EtOH (35 mL) and H₂O (35 mL) was added LiOH•H₂O (2.01 g, 47.89 mmol, 4.0 equiv). The mixtures were stirred at 60 °C for 3 h, at which point the two reaction mixtures were combined, and were adjusted to pH=7 with 1 N HCl. The mixture was concentrated under reduced pressure to remove THF and EtOH, filtered, and the residue was dried under reduced pressure. The residue was stirred in MTBE (100 mL) for 10 min, filtered, and the residue was dried under reduced pressure to afford the desired product (8.02 g, 55.1% yield) as a white solid. LCMS (ESI) m/z : [M + H] calcd for C₂₆H₃₆N₈O₆: 557.29; found 557.3.

Building block AA. *tert*-butyl (2*S*)-4-{6-[(*tert*-butoxy)carbonyl]-5H,6H,7H,8H-pyrido[4,3-*d*]pyrimidin-2-yl}piperazine-2-carboxylate.




[00636] This building block is prepared by a process similar to that for Building block I by utilizing *tert*-butyl 2-chloro-7,8-dihydropyrido[4,3-*d*]pyrimidine-6(5H)-carboxylate and *tert*-butyl (2*S*)-piperazine-2-carboxylate.

Building block AB. 4-amino-2-[(2*S*)-2-[(*tert*-butoxy)carbonyl]-4-{6-[(*tert*-butoxy)carbonyl]-5*H*,6*H*,7*H*,8*H*-pyrido[4,3-*d*]pyrimidin-2-yl}piperazin-1-yl]pyrimidine-5-carboxylic acid.



[00637] This building block is prepared from Building block AA by a process similar to that for Building block J by utilizing ethyl 4-amino-2-chloropyrimidine-5-carboxylate.

Building block AC. 4-amino-2-(4-{6-[(*tert*-butoxy)carbonyl]-5*H*,6*H*,7*H*,8*H*-pyrido[4,3-*d*]pyrimidin-2-yl}piperazin-1-yl)pyrimidine-5-carboxylic acid.



Step 1: Synthesis of *tert*-butyl 2-(4-(4-amino-5-(ethoxycarbonyl)pyrimidin-2-yl)piperazin-1-yl)-7,8-dihydropyrido[4,3-*d*]pyrimidine-6(5*H*)-carboxylate


[00638] To a solution of *tert*-butyl 2-(piperazin-1-yl)-7,8-dihydropyrido[4,3-*d*]pyrimidine-6(5*H*)-carboxylate (8.3 g, 25.99 mmol, 1.0 equiv) and ethyl 4-amino-2-chloropyrimidine-5-carboxylate (5.24 g, 25.99 mmol, 1.0 equiv) in MeCN (100 mL) was added to K₂CO₃ (7.18 g, 51.97 mmol, 2.0 equiv). The reaction was stirred at 80 °C for 12 h. The reaction was then cooled to room temperature, DCM (100 mL) was added, and the reaction mixture was stirred for 30 min. The suspension was filtered, and the filter cake was washed with DCM (6 x 100 mL). The filtrate was concentrated under reduced pressure and the residue was triturated with EtOAc (30 mL), filtered and then the filter cake was dried under reduced pressure to afford the desired product (8.7 g, 65.9% yield) as light yellow solid.

Step 2: Synthesis of 4-amino-2-(4-(6-(*tert*-butoxycarbonyl)-5,6,7,8-tetrahydropyrido[4,3-*d*]pyrimidin-2-yl)piperazin-1-yl)pyrimidine-5-carboxylic acid

[00639] To a solution of *tert*-butyl 2-(4-(4-amino-5-(ethoxycarbonyl)pyrimidin-2-yl)piperazin-1-yl)-7,8-dihydropyrido[4,3-*d*]pyrimidine-6(5*H*)-carboxylate (8.7 g, 17.95

mmol, 1.0 equiv) in THF (120 mL), EtOH (60 mL), and H₂O (60 mL) was added LiOH•H₂O (1.51 g, 35.91 mmol, 2.0 equiv). The mixture was stirred at 55 °C for 12 h. The reaction mixture was then concentrated under reduced pressure to remove EtOH and THF, and the reaction mixture was adjusted to pH=6 by the addition of 1 N HCl. The precipitate was filtered, and the filter cake was washed with H₂O (3 x 50 mL) and then dried under reduced pressure to afford the desired product (7.3 g, 89.1% yield) as light yellow solid. LCMS (ESI) *m/z*: [M + H] calcd for C₂₁H₂₈N₈O₄: 457.23; found 457.2.

Building block AD. 4-amino-2-{4-[5-({[(*tert*-butoxy)carbonyl]amino)methyl]pyrimidin-2-yl]piperazin-1-yl}pyrimidine-5-carboxylic acid.



Step 1: Synthesis of ethyl 4-amino-2-(4-(5-(((di-*tert*-butoxycarbonyl)amino)methyl)pyrimidin-2-yl)piperazin-1-yl)pyrimidine-5-carboxylate


[00640] To a solution of *tert*-butyl-*N*-*tert*-butoxycarbonyl-*N*-((2-(piperazin-1-yl)pyrimidin-5-yl)methyl)carbamate (8.3 g, 21.09 mmol, 1.0 equiv) in MeCN (100 mL) was added ethyl 4-amino-2-chloropyrimidine-5-carboxylate (4.04 g, 20.04 mmol, 0.95 equiv) and K₂CO₃ (8.75 g, 63.28 mmol, 3.0 equiv). The mixture was stirred at 80 °C for 3 h. The reaction was then cooled to room temperature, DCM (150 mL) was added, and the reaction mixture was stirred for 30 min. The suspension was filtered, the filter cake was washed with DCM (3 x 100 mL), and the filtrate was concentrated under reduced pressure. Purification by silica gel chromatography (0%→100% EtOAc/petroleum ether) afforded the desired product (8.35 g, 67% yield) as a white solid.

Step 2: Synthesis of 4-amino-2-(4-(5-(((*tert*-butoxycarbonyl)amino)methyl)pyrimidin-2-yl)piperazin-1-yl)pyrimidine-5-carboxylic acid

[00641] To a solution of ethyl 4-amino-2-(4-(5-(((di-*tert*-butoxycarbonyl)amino)methyl)pyrimidin-2-yl)piperazin-1-yl)pyrimidine-5-carboxylate (8.3 g, 14.86 mmol, 1.0 equiv) in H₂O (70 mL), EtOH (36 mL) and THF (80 mL) was added LiOH•H₂O (2.49 g, 59.43 mmol, 4.0 equiv). The reaction mixture was stirred at 55 °C for 16 h. The mixture was then concentrated under reduced pressure to remove THF and EtOH. The mixture was diluted with H₂O (55 mL) and was adjusted to pH=6 by the addition of 1 N HCl.

The mixture was filtered, and the filter cake was washed with H₂O (2 x 20 mL). The solid cake was dried under reduced pressure to afford the desired product (5.5 g, 84% yield) as a white solid. LCMS (ESI) *m/z*: [M + H] calcd for C₁₉H₂₆N₈O₄: 431.22; found 431.4.

Building block AE. 4-amino-2-[(2*R*)-4-{6-[(*tert*-butoxy)carbonyl]-5*H*,6*H*,7*H*,8*H*-pyrido[4,3-*d*]pyrimidin-2-yl}-2-(hydroxymethyl)piperazin-1-yl]pyrimidine-5-carboxylic acid.



Step 1: Synthesis of (*R*)-*tert*-butyl 2-(4-(4-amino-5-(ethoxycarbonyl)pyrimidin-2-yl)-3-(((*tert*-butyldiphenylsilyl)oxy)methyl)piperazin-1-yl)-7,8-dihydropyrido[4,3-*d*]pyrimidine-6(5*H*)-carboxylate

[00642] To a solution of (*R*)-*tert*-butyl 2-(3-(((*tert*-butyldiphenylsilyl)oxy)methyl)piperazin-1-yl)-7,8-dihydropyrido[4,3-*d*]pyrimidine-6(5*H*)-carboxylate (17.2 g, 29.26 mmol, 1.0 equiv) in MeCN (200 mL) was added K₂CO₃ (12.13 g, 87.78 mmol, 3.0 equiv) and ethyl 4-amino-2-chloropyrimidine-5-carboxylate (6.37 g, 31.60 mmol, 1.08 equiv). The mixture was stirred at 80 °C for 18 h. The reaction mixture was then cooled to room temperature, filtered and the filtrate was concentrated under reduced pressure. Purification by silica gel chromatography (0%→33% EtOAc/petroleum ether) afforded the desired product (20.3 g, 90.6% yield) as a white solid. LCMS (ESI) *m/z*: [M + H] calcd for C₄₀H₅₂N₈O₅Si: 753.39; found 753.4.

Step 2: Synthesis of (*R*)-4-amino-2-(4-(6-(*tert*-butoxycarbonyl)-5,6,7,8-tetrahydropyrido[4,3-*d*]pyrimidin-2-yl)-2-(hydroxymethyl)piperazin-1-yl)pyrimidine-5-carboxylic acid


[00643] To a solution of (*R*)-*tert*-butyl 2-(4-(4-amino-5-(ethoxycarbonyl)pyrimidin-2-yl)-3-(((*tert*-butyldiphenylsilyl)oxy)methyl)piperazin-1-yl)-7,8-dihydropyrido[4,3-*d*]pyrimidine-6(5*H*)-carboxylate (20.3 g, 26.96 mmol, 1.0 equiv) in THF (200 mL) was added TBAF (1.0 M, 50.75 mL, 1.9 equiv). The reaction mixture was stirred at room temperature for 5 h. The mixture was then poured into H₂O (200 mL) and the aqueous phase was extracted with

EtOAc (2 x 150 mL). The combined organic phases were washed with brine (2 x 100 mL), dried, filtered and concentrated under reduced pressure. Purification by silica gel chromatography (0%→20% EtOAc/petroleum ether) afforded the desired product (12 g, 85.7% yield) as a white solid. LCMS (ESI) m/z : $[M + H]^+$ calcd for $C_{24}H_{34}N_8O_5$: 515.28; found 515.4.

Step 3: Synthesis of (R)-4-amino-2-(4-(6-(tert-butoxycarbonyl)-5,6,7,8-tetrahydropyrido[4,3-d]pyrimidin-2-yl)-2-(hydroxymethyl)piperazin-1-yl)pyrimidine-5-carboxylic acid


[00644] To a solution of (R)-4-amino-2-(4-(6-(tert-butoxycarbonyl)-5,6,7,8-tetrahydropyrido[4,3-d]pyrimidin-2-yl)-2-(hydroxymethyl)piperazin-1-yl)pyrimidine-5-carboxylic acid (12 g, 23.32 mmol, 1.0 equiv) in THF (100 mL), EtOH (30 mL), and H₂O (30 mL) was added LiOH•H₂O (5.87 g, 139.92 mmol, 6.0 equiv). The mixture was stirred at 50 °C for 22 h. The mixture was then concentrated under reduced pressure to remove THF and EtOH. The aqueous phase was neutralized with 1 N HCl and the resulting precipitate was filtered. The filter cake was washed with H₂O (50 mL) and dried under reduced pressure. The filtrate was extracted with DCM (8 x 60 mL) and the combined organic phases were washed with brine (2 x 50 mL), dried, filtered, and concentrated under reduced pressure. The resulting residue was combined with the initial filter cake and the solid was dissolved in DCM (150 mL) and concentrated under reduced pressure to afford the desired product (9.76 g, 85.2% yield) as a white solid. LCMS (ESI) m/z : $[M + H]^+$ calcd for $C_{22}H_{30}N_8O_5$: 487.24; found 487.2.

Building block AF. 4-amino-2-[(2S)-4-{6-[(tert-butoxy)carbonyl]-5H,6H,7H,8H-pyrido[4,3-d]pyrimidin-2-yl}-2-(hydroxymethyl)piperazin-1-yl]pyrimidine-5-carboxylic acid




[00645] This building block is prepared from Building block O by a process similar to that for Building block J by utilizing ethyl 4-amino-2-chloropyrimidine-5-carboxylate.

Building block AG. 2-((2-(4-(5-((di-*tert*-butoxycarbonyl)amino)methyl)pyrimidin-2-yl)piperazin-1-yl)-2-oxoethyl)(methyl)amino)acetic acid.



[00646] To a solution of *tert*-butyl *N*-*tert*-butoxycarbonyl-*N*-((2-piperazin-1-ylpyrimidin-5-yl)methyl)carbamate (4.88 g, 12.39 mmol, 1.0 equiv) in EtOAc (40 mL) was added 4-methylmorpholine-2,6-dione (1.6 g, 12.39 mmol, 1.0 equiv). The reaction was stirred at room temperature for 2 h then reaction mixture was concentrated under reduced pressure to give the crude product. The residue was triturated with EtOAc (15 mL) and filtered to give the product (5.65 g, 87.2% yield) as a white solid. LCMS (ESI) *m/z*: [M + H] calcd for C₂₄H₃₉N₆O₇: 523.28; found 523.3.

Building block AH. *tert*-butyl *N*-*tert*-butoxycarbonyl-*N*-((2-(4-(3-(2-piperazin-1-ylethoxy)propanoyl)piperazin-1-yl)pyrimidin-5-yl)methyl)carbamate.



Step 1: Synthesis of benzyl 4-(2-(3-(*tert*-butoxy)-3-oxopropoxy)ethyl)piperazine-1-carboxylate

[00647] To a solution of *tert*-butyl 3-(2-bromoethoxy)propanoate (35 g, 138.27 mmol, 1.0 equiv) and benzyl piperazine-1-carboxylate (31.14 mL, 138.27 mmol, 1.0 equiv, HCl) in MeCN (420 mL) was added K₂CO₃ (57.33 g, 414.80 mmol, 3.0 equiv). The reaction was stirred at 80 °C for 20 h. The reaction mixture was cooled to room temperature and the suspension was filtered. The filter cake was washed with EtOAc (3 x 50 mL) and the combined filtrates were concentrated under reduced pressure to give crude product. The

residue was purified by silica gel chromatography (5/1 to 0/1 petroleum ether/EtOAc) to give the product (46 g, 84.8% yield) as a yellow oil.

Step 2: Synthesis of 3-(2-(4-((benzyloxy)carbonyl)piperazin-1-yl)ethoxy)propanoic acid


[00648] A solution of benzyl 4-(2-(3-(*tert*-butoxy)-3-oxopropoxy)ethyl)piperazine-1-carboxylate (21 g, 53.50 mmol, 1.0 equiv) in TFA (160 mL) was stirred at room temperature for 2 h and then concentrated under reduced pressure. The residue was purified by silica gel chromatography (1/0 to 4/1 EtOAc/MeOH) to give the product (20.4 g, 84.7% yield) as a yellow oil. LCMS (ESI) *m/z*: [M + H] calcd for C₁₇H₂₄N₂O₅: 337.18; found 337.1.

*Step 3: Synthesis of benzyl 4-(2-(3-(4-(5-(((di-*tert*-butoxycarbonyl)amino)methyl)pyrimidin-2-yl)piperazin-1-yl)-3-oxopropoxy)ethyl)piperazine-1-carboxylate*

[00649] To a solution of 3-(2-(4-((benzyloxy)carbonyl)piperazin-1-yl)ethoxy)propanoic acid (20.2 g, 44.85 mmol, 1.0 equiv, TFA) in DCM (500 mL) was added HATU (25.58 g, 67.27 mmol, 1.5 equiv) and DIPEA (17.39 g, 134.55 mmol, 23.44 mL, 3.0 equiv). The reaction was stirred at room temperature for 30 min, and then *tert*-butyl *N-tert*-butoxycarbonyl-*N*-((2-piperazin-1-yl)pyrimidin-5-yl)methyl)carbamate (14.12 g, 35.88 mmol, 0.8 equiv) was added. The reaction mixture was stirred at for 2 h and then quenched with sat. NH₄Cl (500 mL). The aqueous phase was extracted with DCM (3 x 300 mL) and the combined organic phase was washed with brine (30 mL), dried with anhydrous Na₂SO₄, filtered and concentrated under reduced pressure to give crude product. The residue was purified by silica gel chromatography (0/1 petroleum ether/EtOAc to 10/1 DCM/MeOH) to give the product (29 g, 90.8% yield) as a yellow oil. LCMS (ESI) *m/z*: [M + H] calcd for C₃₆H₅₃N₇O₈: 712.41; found 712.4.

Step 4: Synthesis of tert-butyl N-tert-butoxycarbonyl-N-((2-(4-(3-(2-piperazin-1-ylethoxy)propanoyl)piperazin-1-yl)pyrimidin-5-yl)methyl)carbamate

[00650] To a solution of 4-(2-(3-(4-(5-(((di-*tert*-butoxycarbonyl)amino)methyl)pyrimidin-2-yl)piperazin-1-yl)-3-oxopropoxy)ethyl)piperazine-1-carboxylate (5 g, 7.02 mmol, 1.0 equiv) in EtOAc (150 mL) was added Pd/C (2 g, 10 wt.%). The suspension was degassed and purged with H₂ and then stirred under H₂ (30 psi) at 30 °C for 3 h. The suspension was then cooled to room temperature and filtered through Celite. The filter cake was washed with MeOH (15 x 100 mL) and the combined filtrates were concentrated under reduced pressure to give the product (12 g, 89.9% yield) as a light yellow oil. LCMS (ESI) *m/z*: [M + H] calcd for C₂₈H₄₇N₇O₆: 578.37; found 578.5.


Building block AI. ethyl 2-(piperazin-1-yl)pyrimidine-5-carboxylate.

Step 1: Synthesis of ethyl 2-(4-(tert-butoxycarbonyl)piperazin-1-yl)pyrimidine-5-carboxylate

[00651] To a solution of *tert*-butyl piperazine-1-carboxylate (11.94 g, 53.59 mmol, 1.0 equiv, HCl) and ethyl 2-chloropyrimidine-5-carboxylate (10 g, 53.59 mmol, 1.0 equiv) in MeCN (100 mL) was added K₂CO₃ (7.41 g, 53.59 mmol, 1.0 equiv). The mixture was stirred at 80 °C for 17 h and then poured into H₂O (200 mL). The mixture was filtered and the filter cake was washed with H₂O (80 mL) and dried under reduced pressure to give the product (15.76 g, 82% yield) as a white solid.

Step 2: Synthesis of ethyl 2-(piperazin-1-yl)pyrimidine-5-carboxylate

[00652] To a solution of ethyl 2-(4-(*tert*-butoxycarbonyl)piperazin-1-yl)pyrimidine-5-carboxylate (15.7 g, 46.67 mmol, 1.0 equiv) in EtOAc (150 mL) was added HCl/EtOAc (150 mL) at 0 °C. The resulting mixture was stirred at room temperature for 9 h. The reaction mixture was filtered and the filter cake was washed with EtOAc (100 mL). The solid was dried under reduced pressure to give the product (12.55 g, 96% yield, HCl) as a white solid. LCMS (ESI) *m/z*: [M + H] calcd for C₁₁H₁₆N₄O₂: 237.14; found 237.3.

Building block AJ. 2-(4-(2-(3-(4-(5-(((di-*tert*-butoxycarbonyl)amino)methyl)pyrimidin-2-yl)piperazin-1-yl)-3-oxopropoxy)ethyl)piperazin-1-yl)pyrimidine-5-carboxylic acid.

Step 1: Synthesis of ethyl 2-(4-(2-(3-(tert-butoxy)-3-oxopropoxy)ethyl)piperazin-1-yl)pyrimidine-5-carboxylate

[00653] To a solution of ethyl 2-piperazin-1-ylpyrimidine-5-carboxylate (17.92 g, 75.85 mmol, 1.2 equiv) and *tert*-butyl 3-(2-bromoethoxy)propanoate (16 g, 63.21 mmol, 1.0 equiv) in MeCN (200 mL) was added K₂CO₃ (17.47 g, 126.42 mmol, 2.0 equiv). The reaction was stirred at 80 °C for 12 h and then the reaction mixture was filtered, and the filtrate was concentrated under reduced pressure. The crude product was suspended in petroleum ether (200 mL) and stirred for 20 min at 0 °C and then filtered. The solid was dried under reduced pressure to give the product (19.4 g, 75.1% yield) as a yellow solid.

Step 2: Synthesis of 3-(2-(4-(5-(ethoxycarbonyl)pyrimidin-2-yl)piperazin-1-yl)ethoxy)propanoic acid

[00654] A solution of ethyl 2-(4-(2-(3-(*tert*-butoxy)-3-oxopropoxy)ethyl)piperazin-1-yl)pyrimidine-5-carboxylate (19.4 g, 47.49 mmol, 1.0 equiv) in TFA (200 mL) was stirred at room temperature for 30 min. The reaction mixture was then concentrated under reduced pressure and the residue was purified by silica gel chromatography (50/1 to 1/1 EtOAc/MeOH) to give the product (18 g, 81.3% yield) as a yellow oil.

*Step 3: Synthesis of ethyl 2-(4-(2-(3-(4-(5-(((di-*tert*-butoxycarbonyl)amino)methyl)pyrimidin-2-yl)piperazin-1-yl)-3-oxopropoxy)ethyl)piperazin-1-yl)pyrimidine-5-carboxylate*


[00655] To a solution of 3-(2-(4-(5-(ethoxycarbonyl)pyrimidin-2-yl)piperazin-1-yl)ethoxy)propanoic acid (13 g, 27.87 mmol, 1.0 equiv) in DCM (200 mL) was added HATU (15.90 g, 41.81 mmol, 1.5 equiv) and DIPEA (19.42 mL, 111.49 mmol, 4.0 equiv). The reaction was then stirred at room temperature for 30 min and then *tert*-butyl *N-tert*-butoxycarbonyl-*N*-[(2-piperazin-1-ylpyrimidin-5-yl)methyl]carbamate (10.97 g, 27.87 mmol, 1.0 equiv) was added. The mixture was stirred at for 2 h and then poured into a sat. NH₄Cl solution (200 mL). The aqueous phase was extracted with DCM (2 x 200 mL) and the combined organic phase was washed with brine (2 x 20 mL), dried with anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by silica gel chromatography (100/1 to 9/1 EtOAc/MeOH) to give the product (17 g, 79.0% yield) as a yellow oil.

*Step 4: Synthesis of 2-(4-(2-(3-(4-(5-(((di-*tert*-butoxycarbonyl)amino)methyl)pyrimidin-2-yl)piperazin-1-yl)-3-oxopropoxy)ethyl)piperazin-1-yl)pyrimidine-5-carboxylic acid*

[00656] To a solution of ethyl 2-(4-(2-(3-(4-(5-(((di-*tert*-butoxycarbonyl)amino)methyl)pyrimidin-2-yl)piperazin-1-yl)-3-oxopropoxy)ethyl)piperazin-


1-yl)pyrimidine-5-carboxylate (11 g, 15.11 mmol, 1.0 equiv) in THF (40 mL), EtOH (10 mL), and H₂O (20 mL) was added LiOH•H₂O (1.27 g, 30.23 mmol, 2.0 equiv). The mixture was then stirred at 35 °C for 1.5 h. The reaction mixture was extracted with EtOAc (30 mL) and the aqueous phase was adjusted to pH = 7 by addition of HCl (1 N). The mixture was then concentrated under reduced pressure. The crude product was purified by reversed-phase chromatography (20/1 to 3/1 H₂O/MeCN) to give the product (6.1 g, 67.3% yield) as a white solid. LCMS (ESI) *m/z*: [M + H] calcd for C₃₃H₄₉N₉O₈: 700.38; found 700.4.

Building block AK. 2-(4-(2-(3-(4-(5-(((*tert*-butoxycarbonyl)amino)methyl)pyrimidin-2-yl)piperazin-1-yl)-3-oxopropoxy)ethyl)piperazin-1-yl)pyrimidine-5-carboxylic acid.



[00657] A solution of ethyl 2-(4-(2-(3-(4-(5-(((*tert*-butoxycarbonyl)amino)methyl)pyrimidin-2-yl)piperazin-1-yl)-3-oxopropoxy)ethyl)piperazin-1-yl)pyrimidine-5-carboxylate (5.4 g, 7.42 mmol, 1.0 equiv) in THF (40 mL), EtOH (10 mL), and H₂O (10 mL) was added LiOH•H₂O (933.92 mg, 22.26 mmol, 3.0 equiv). The mixture was then stirred at 30 °C for 12 h. The reaction mixture was then extracted with EtOAc (2 x 50 mL) and the aqueous phase was adjusted to pH = 7 by addition of HCl (1 N). The solution was then concentrated under reduced pressure. The crude product was purified by reversed-phase chromatography (20/1 to 3/1 H₂O/MeCN) to give the product (1.01 g, 22.5% yield) as a white solid. LCMS (ESI) *m/z*: [M + H] calcd for C₂₈H₄₁N₉O₆: 600.33; found 600.2.


Building block AL. 4-{4-[2-(3-{4-[5-(((*tert*-butoxy)carbonyl]amino)methyl)pyrimidin-2-yl]piperazin-1-yl}-3-oxopropoxy)ethyl]piperazin-1-yl}-4-oxobutanoic acid.



[00658] To a solution of *tert*-butyl *N*-*tert*-butoxycarbonyl-*N*-((2-(4-(3-(2-piperazin-1-ylethoxy)propanoyl)piperazin-1-yl)pyrimidin-5-yl)methyl)carbamate (1.0 equiv) in DCM is added succinic anhydride (1.2 equiv) and Et₃N (2.0 equiv). The reaction is stirred at room

temperature until consumption of starting material, as determined by LCMS analysis. The reaction mixture is then concentrated under reduced pressure to give the crude product. The residue is purified by silica gel chromatography to afford the product.

Building block AM. 2-(4-(4-(4-(5-(((*tert*-butoxycarbonyl)amino)methyl)pyrimidin-2-yl)piperazin-1-yl)-4-oxobutyl)piperazin-1-yl)pyrimidine-5-carboxylic acid.



Step 1: Synthesis of ethyl 2-(4-(4-(*tert*-butoxy)-4-oxobutyl)piperazin-1-yl)pyrimidine-5-carboxylate

[00659] To a solution of ethyl 2-(piperazin-1-yl)pyrimidine-5-carboxylate hydrochloride (10 g, 36.67 mmol, 1.0 equiv, HCl) and *tert*-butyl 4-bromobutanoate (8.18 g, 36.67 mmol, 1.0 equiv) in DMF (100 mL) was added Et₃N (15.31 mL, 110.00 mmol, 3.0 equiv). The mixture was stirred at 130 °C for 14 h. The mixture was then poured into H₂O (400 mL) and the solution was extracted with EtOAc (3 x 150 mL). The combined organic layer was washed with brine (200 mL), dried over Na₂SO₄ and concentrated under reduced pressure. The residue was purified by silica gel chromatography (5/1 to 1/1 petroleum ether/EtOAc) to give the product (9.5 g, 68.5% yield) as a yellow solid. LCMS (ESI) *m/z*: [M + H] calcd for C₁₉H₃₀N₄O₄: 379.24; found 379.2, 380.2.

Step 2: Synthesis of 4-(4-(4-(5-(ethoxycarbonyl)pyrimidin-2-yl)piperazin-1-yl)butanoic acid hydrochloride

[00660] To a solution of ethyl 2-(4-(4-(*tert*-butoxy)-4-oxobutyl)piperazin-1-yl)pyrimidine-5-carboxylate (9.5 g, 25.10 mmol, 1.0 equiv) in EtOAc (100 mL) was added HCl/EtOAc (500 mL). The mixture was stirred at room temperature for 10 h and then the solution was concentrated under reduced pressure to give the product (9.6 g, 96.8% yield) as a white solid. LCMS (ESI) *m/z*: [M + H] calcd for C₁₅H₂₂N₄O₄: 323.17; found 323.2.


Step 3: Synthesis of ethyl 2-(4-(4-(4-(5-(((di-*tert*-butoxycarbonyl)amino)methyl)pyrimidin-2-yl)piperazin-1-yl)-4-oxobutyl)piperazin-1-yl)pyrimidine-5-carboxylic acid

[00661] To a solution of 4-(4-(5-(ethoxycarbonyl)pyrimidin-2-yl)piperazin-1-yl)butanoic acid hydrochloride (5 g, 15.51 mmol, 1.0 equiv) and *tert*-butyl N-*tert*-butoxycarbonyl-N-((2-piperazin-1-yl)pyrimidin-5-yl)methyl)carbamate (6.10 g, 15.51 mmol, 1.0 equiv) in DMF (150 mL) was added DIPEA (8.11 mL, 46.53 mmol, 3.0 equiv) and HATU (7.08 g, 18.61 mmol, 1.2 equiv). The mixture was stirred at room temperature for 3 h and then the solution was poured into H₂O (600 mL). The aqueous layer was extracted with EtOAc (3 x 200 mL) and then the combined organic layer was washed with brine (100 mL), dried with Na₂SO₄ and concentrated under reduced pressure. The residue was purified by silica gel chromatography (50/1 to 15/1 DCM/MeOH) to give the product (6.3 g, 58.2% yield) as a yellow solid. LCMS (ESI) *m/z*: [M + H] calcd for C₃₄H₅₁N₉O₇: 698.40; found 698.6.

Step 4: Synthesis of 2-(4-(4-(4-(5-(((*tert*-butoxycarbonyl)amino)methyl)pyrimidin-2-yl)piperazin-1-yl)-4-oxobutyl)piperazin-1-yl)pyrimidine-5-carboxylic acid

[00662] To a solution of ethyl 2-(4-(4-(4-(5-((bis(*tert*-butoxycarbonyl)amino)methyl)pyrimidin-2-yl)piperazin-1-yl)-4-oxo-butyl)piperazin-1-yl)pyrimidine-5-carboxylate (4.5 g, 6.45 mmol, 1.0 equiv) in EtOH (7 mL) and THF (28 mL) was added a solution of LiOH•H₂O (541.17 mg, 12.90 mmol, 2.0 equiv) in H₂O (7 mL). The mixture was stirred at 30 °C for 8 h, then additional LiOH•H₂O (541 mg, 12.90 mmol, 2.0 equiv) was added. After stirring for an additional 8 h at 30 °C, the solution was concentrated under reduced pressure. H₂O (20 mL) was added and solution was adjusted to pH 3 with 1N HCl. The suspension was filtered and the solid dried under reduced pressure to give the product (3.2 g, 79.1% yield) as a white solid. LCMS (ESI) *m/z*: [M + H] calcd for C₂₇H₃₉N₉O₅: 570.32; found 570.3.

Building Block AN. 2-(4-(2-(2-(4-(6-(*tert*-butoxycarbonyl)-5,6,7,8-tetrahydropyrido[4,3-d]pyrimidin-2-yl)piperazin-1-yl)ethoxy)ethyl)piperazin-1-yl)pyrimidine-5-carboxylic acid.



Step 1: Synthesis of benzyl 4-(2-hydroxyethyl)piperazine-1-carboxylate

[00663] To a solution of benzyl piperazine-1-carboxylate hydrochloride (41.09 g, 160.04 mmol, 1.0 equiv, HCl) in MeCN (200 mL) was added K₂CO₃ (66.36 g, 480.13 mmol, 3.0 equiv) and 2-bromoethanol (20 g, 160.04 mmol, 1.0 equiv). The reaction mixture was stirred at 80 °C for 16 h, at which point it was cooled to room temperature and filtered. The filter cake was washed with EtOAc (100 mL) and the filtrate then washed with H₂O (100 mL). The aqueous phase was extracted with EtOAc (3 x 50 mL) and the combined organic phases were washed with brine (50 mL), dried, and concentrated under reduced pressure. Purification by silica gel chromatography (5→25% MeOH/EtOAc) afforded the desired product as a yellow solid (20 g, 47% yield). LCMS (ESI) *m/z*: [M + H] calcd for C₁₄H₂₀N₂O₃: 265.16; found 264.9.

Step 2: Synthesis of tert-butyl 4-(2-hydroxyethyl)piperazine-1-carboxylate

[00664] To a solution of *tert*-butyl piperazine-1-carboxylate (198.72 g, 1.07 mol, 1.0 equiv) in MeCN (1500 mL) was added 2-bromoethanol (240 g, 1.92 mol, 1.8 equiv) and K₂CO₃ (221.19 g, 1.60 mol, 1.5 equiv). The reaction mixture was stirred at 80 °C for 16 h, at which point the mixture was cooled to room temperature, filtered, and the filtrate was concentrated under reduced pressure. Purification by silica gel chromatography (0→14% MeOH/EtOAc) afforded the desired product as a white solid (146 g, 59% yield).

Step 3: Synthesis of tert-butyl 4-(2-bromoethyl)piperazine-1-carboxylate

[00665] To a solution of *tert*-butyl 4-(2-hydroxyethyl)piperazine-1-carboxylate (45 g, 195.39 mmol, 1.0 equiv) in THF (600 mL) was added triphenylphosphine (97.38 g, 371.25 mmol, 1.9 equiv) and CBr₄ (116.64 g, 351.71 mmol, 1.8 equiv). The mixture was stirred at room temperature for 3 h. Two separate batches were combined, and the reaction mixture was filtered, and the filtrate concentrated under reduced pressure. Purification by silica gel chromatography (1→25% EtOAc/petroleum ether) afforded the desired product as a light-yellow solid (31 g, 27% yield).

Step 4: Synthesis of benzyl 4-(2-(2-(4-(tert-butoxycarbonyl)piperazin-1-yl)ethoxy)ethyl)piperazine-1-carboxylate

[00666] To a solution of benzyl 4-(2-hydroxyethyl)piperazine-1-carboxylate (18 g, 68.10 mmol, 1.0 equiv) in toluene (200 mL) was added NaNH₂ (26.57 g, 680.99 mmol, 10.0 equiv). *tert*-Butyl 4-(2-bromoethyl)piperazine-1-carboxylate (25 g, 85.27 mmol, 1.25 equiv) was added and the mixture was heated to 90 °C for 18 h. The mixture was cooled to room temperature and poured into H₂O (700 mL) at 0 °C. The aqueous phase was extracted with EtOAc (3 x 240 mL) and the combined organic phases were washed successively with H₂O (350 mL) and sat. brine (2 x 200 mL), dried, filtered, and concentrated under reduced pressure. Purification by silica gel chromatography (0→12% MeOH/EtOAc) afforded the desired product as a light-yellow oil (20 g, 62% yield).

Step 5: Synthesis of tert-butyl 4-(2-(2-(piperazin-1-yl)ethoxy)ethyl)piperazine-1-carboxylate

[00667] To a solution of benzyl 4-(2-(2-(4-(tert-butoxycarbonyl)piperazin-1-yl)ethoxy)ethyl)piperazine-1-carboxylate (20 g, 41.96 mmol, 1.0 equiv) in EtOAc (180 mL) was added Pd/C (8 g, 10 wt.%). The suspension was degassed under reduced pressure and purged with H₂ three times. The mixture was stirred under H₂ (30 psi) at 35 °C for 12 h. The reaction mixture was then filtered, and the filtrate was concentrated under reduced pressure.

Purification by silica gel chromatography (0→100% MeOH/EtOAc) afforded the desired product as a colorless oil (10.8 g, 75% yield).

Step 6: Synthesis of ethyl 2-(4-(2-(2-(4-(*tert*-butoxycarbonyl)piperazin-1-yl)ethoxy)ethyl)piperazin-1-yl)pyrimidine-5-carboxylate

[00668] To a solution of *tert*-butyl 4-(2-(2-(piperazin-1-yl)ethoxy)ethyl)piperazine-1-carboxylate (10.8 g, 31.54 mmol, 1.0 equiv) in MeCN (100 mL) was added K₂CO₃ (13.08 g, 94.61 mmol, 3.0 equiv) and ethyl 2-chloropyrimidine-5-carboxylate (5.88 g, 31.54 mmol, 1.0 equiv). The mixture was stirred at 80 °C for 12 h, at which point the reaction was cooled to room temperature, filtered, and the filtrate concentrated under reduced pressure. Purification by silica gel chromatography (0→9% MeOH/DCM) afforded the desired product as a white solid (13.6 g, 85% yield).

Step 7: Synthesis of 2-(4-(2-(2-(4-(6-(*tert*-butoxycarbonyl)-5,6,7,8-tetrahydropyrido-[4,3-d]pyrimidin-2-yl)piperazin-1-yl)ethoxy)ethyl)piperazin-1-yl)pyrimidine-5-carboxylic acid

[00669] To a solution of ethyl 2-(4-(2-(2-(4-(*tert*-butoxycarbonyl)piperazin-1-yl)ethoxy)ethyl)piperazin-1-yl)pyrimidine-5-carboxylate (13.6 g, 27.61 mmol, 1.0 equiv) in MeOH (50 mL) was added a solution of HCl in MeOH (4 M, 150 mL, 21.7 equiv). The reaction was stirred at room temperature for 4 h, at which point the mixture was concentrated under reduced pressure to afford the crude desired product as a white solid (13.8 g, 4HCl) that was taken directly onto the next step. LCMS (ESI) *m/z*: [M + H] calcd for C₁₉H₃₂N₆O₃: 393.26; found 393.3.

Step 8: Synthesis of *tert*-butyl 2-(4-(2-(2-(4-(5-(ethoxycarbonyl)pyrimidin-2-yl)-piperazin-1-yl)ethoxy)ethyl)piperazin-1-yl)-7,8-dihydropyrido[4,3-d]pyrimidine-6(5H)-carboxylate


[00670] To a stirred solution of 2-(4-(2-(2-(4-(6-(*tert*-butoxycarbonyl)-5,6,7,8-tetrahydropyrido[4,3-d]pyrimidin-2-yl)piperazin-1-yl)ethoxy)ethyl)piperazin-1-yl)pyrimidine-5-carboxylic acid (10.2 g, 18.95 mmol, 1.0 equiv, 4HCl) and DIPEA (16.50 mL, 94.74 mmol, 5.0 equiv) in DMF (100 mL) was added *tert*-butyl 2-chloro-7,8-dihydropyrido[4,3-d]pyrimidine-6(5H)-carboxylate (5.11 g, 18.95 mmol, 1.0 equiv). The reaction mixture was stirred at 90 °C for 12 h. The reaction mixture was then cooled to room temperature and added to EtOAc (200 mL) and H₂O (400 mL). The aqueous phase was extracted with EtOAc (2 x 100 mL) and the combined organic phases were washed with aqueous NH₄Cl (4 x 100 mL), brine (2 x 100 mL), dried, filtered and concentrated under

reduced pressure. Purification by silica gel chromatography (0→9% MeOH/DCM) afforded the desired product as a white solid (5.4 g, 45% yield). LCMS (ESI) m/z : $[M + H]$ calcd for $C_{31}H_{47}N_9O_5$: 626.38; found 626.3.

*Step 9: Synthesis of 2-(4-(2-(2-(4-(6-(*tert*-butoxycarbonyl)-5,6,7,8-tetrahydropyrido[4,3-d]pyrimidin-2-yl)piperazin-1-yl)ethoxy)ethyl)piperazin-1-yl)pyrimidine-5-carboxylic acid*


[00671] To a solution of *tert*-butyl 2-(4-(2-(2-(4-(5-(ethoxycarbonyl)pyrimidin-2-yl)piperazin-1-yl)ethoxy)ethyl)piperazin-1-yl)-7,8-dihydropyrido[4,3-d]pyrimidine-6(5H)-carboxylate (5.4 g, 8.63 mmol, 1.0 equiv) in THF (50 mL), EtOH (20 mL), and H₂O (20 mL) was added LiOH•H₂O (1.09 g, 25.89 mmol, 3.0 equiv). The reaction mixture was stirred at 35 °C for 12 h, at which point the mixture was concentrated under reduced pressure to remove THF and EtOH. The aqueous phase was neutralized to pH = 7 with 0.5N HCl and concentrated under reduced pressure. Purification by reverse phase chromatography afforded the desired product as a white solid (4.72 g, 92% yield). LCMS (ESI) m/z : $[M + H]$ calcd for $C_{29}H_{43}N_9O_5$: 598.35; found 598.3.

Building block AO. 1'-[(*tert*-butoxy)carbonyl]-[1,4'-bipiperidine]-4-carboxylic acid.



[00672] At the time of this application this building block was commercially available (CAS # 201810-59-5).

Building block AP. 2-((2-(piperazin-1-yl)pyrimidin-5-yl)methyl)isoindoline-1,3-dione hydrochloride salt.



Step 1: Synthesis of 2-chloro-5-(dibromomethyl)pyrimidine

[00673] To a solution of 2-chloro-5-methylpyrimidine (100 g, 777.85 mmol, 1.0 equiv) in CCl₄ (1200 mL) was added NBS (304.58 g, 1.71 mol, 2.2 equiv) and AIBN (51.09 g, 311.14 mmol, 0.4 equiv). The mixture was stirred at 80 °C for 16 h. The reaction solution was then cooled to room temperature, filtered, and the filtrate was poured into H₂O (1500 mL). The solution was diluted with DCM (3 x 250 mL) and the organic layer washed with brine (300 mL), dried with anhydrous Na₂SO₄, filtered and concentrated under reduced pressure to give the crude product as a brown oil, which was used directly in the next step.

Step 2: Synthesis of 5-(bromomethyl)-2-chloropyrimidine

[00674] To a solution of 2-chloro-5-(dibromomethyl)pyrimidine (229 g, 799.72 mmol, 1.0 equiv) in THF (600 mL) was added DIPEA (111.44 mL, 639.77 mmol, 0.8 equiv) and 1-ethoxyphosphonyloxyethane (82.57 mL, 639.77 mmol, 0.8 equiv). The mixture was stirred at room temperature for 19 h. The mixture was then poured into H₂O (1200 mL) and the aqueous phase was extracted with EtOAc (3 x 300 mL). The combined organic phase was washed with brine (300 mL), dried with anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by silica gel chromatography (1/0 to 0/1 petroleum ether/EtOAc) to give the product as a brown oil, which was used directly for the next step.

Step 3: Synthesis of 2-((2-chloropyrimidin-5-yl)methyl)isoindoline-1,3-dione

[00675] To a mixture of isoindoline-1,3-dione (15 g, 101.95 mmol, 1.0 equiv) in DMF (126 mL) was added NaH (4.89 g, 122.34 mmol, 60 wt.%, 1.2 equiv) at 0 °C. The mixture was stirred at 0 °C for 30 min, then a solution of 5-(bromomethyl)-2-chloro-pyrimidine (30.21 g, 101.95 mmol, 1.0 equiv) in DMF (24 mL) was added dropwise to the above mixture at room temperature. The mixture was stirred at room temperature for 2 h and was then cooled to 0 °C and quenched with sat. NH₄Cl (600 mL). The suspension was filtered and the solid dried under reduced pressure to give the crude product (27.4 g, 98.2% yield) as a grey solid, which was used directly in the next step. LCMS (ESI) *m/z*: [M + H] calcd for C₁₃H₈ClN₃O₂: 274.04; found 274.0.

Step 4: Synthesis of tert-butyl 4-(5-((1,3-dioxoisoindolin-2-yl)methyl)pyrimidin-2-yl)piperazine-1-carboxylate


[00676] To a solution of 2-((2-chloropyrimidin-5-yl)methyl)isoindoline-1,3-dione (27 g, 98.66 mmol, 1.0 equiv) and *tert*-butyl piperazine-1-carboxylate (20.21 g, 108.52 mmol, 1.1 equiv) in DMF (270 mL) was added K₂CO₃ (34.09 g, 246.64 mmol, 2.5 equiv). The mixture

was stirred at 80 °C for 3 h and then the reaction was cooled to room temperature and poured into H₂O (1200 mL). The suspension was filtered and the solid was dried under reduced pressure to give the crude product (35.58 g, 85.2% yield) as a white solid, which was used directly in the next step.

Step 5: Synthesis of 2-((2-(piperazin-1-yl)pyrimidin-5-yl)methyl)isoindoline-1,3-dione


[00677] A solution of *tert*-butyl 4-(5-((1,3-dioxoisindolin-2-yl)methyl)pyrimidin-2-yl)piperazine-1-carboxylate (15 g, 35.42 mmol, 1 equiv) in HCl/EtOAc (150 mL) was stirred at room temperature for 2 h. The mixture was filtered and then the filter cake was washed with EtOAc (20 mL) and dried under reduced pressure to give the product (42.53 g, 92.5% yield) as a white solid.

Building block AQ. 2-[(2-{4-[2-(3-{4-[5-({bis[(*tert*-butoxy)carbonyl]amino}methyl)pyrimidin-2-yl]piperazin-1-yl}-3-oxopropoxy)ethyl]piperazin-1-yl}-2-oxoethyl)(methyl)amino]acetic acid.



[00678] To a solution of *tert*-butyl N-[(*tert*-butoxy)carbonyl]-N-{[2-(4-{3-[2-(piperazin-1-yl)ethoxy]propanoyl}piperazin-1-yl)pyrimidin-5-yl]methyl} carbamate (300 mg, 519 μmol, 1.0 equiv) in pyridine (8 mL) at 0 °C was added 4-methylmorpholine-2,6-dione (80.3 mg, 622 μmol, 1.2 equiv). The reaction mixture was stirred at 0 °C for 1 h and then warmed to room temperature and stirred for an additional 12 h. The solvent was concentrated under reduced pressure and the solid was partitioned between DCM and H₂O. The organic layer was separated, dried over MgSO₄ and the solvent was concentrated under reduced pressure to give the product (23.0 mg, 6.28% yield). LCMS (ESI) *m/z*: [M + H] calcd for C₃₃H₅₄N₈O₉: 707.41; found 707.4.

Building Block AR. 2-(4-(2-(3-(4-(6-(*tert*-butoxycarbonyl)-5,6,7,8-tetrahydropyrido[4,3-d]pyrimidin-2-yl)piperazin-1-yl)-3-oxopropoxy)ethyl)piperazin-1-yl)pyrimidine-5-carboxylic acid.



Step 1: Synthesis of *tert*-butyl 2-(4-(3-(2-(4-(5-(ethoxycarbonyl)pyrimidin-2-yl)piperazin-1-yl)ethoxy)propanoyl)piperazin-1-yl)-7,8-dihydropyrido[4,3-d]pyrimidine-6(5H)-carboxylate


[00679] To a solution of 3-(2-(4-(5-(ethoxycarbonyl)pyrimidin-2-yl)piperazin-1-yl)ethoxy)propanoic acid (6 g, 12.86 mmol, 1.0 equiv, TFA) in DMF (55 mL) was added HATU (6.36 g, 16.72 mmol, 1.3 equiv) and DIPEA (11.20 mL, 64.32 mmol, 5.0 equiv). After 0.5 h, *tert*-butyl 2-(piperazin-1-yl)-7,8-dihydropyrido[4,3-d]pyrimidine-6(5H)-carboxylate (4.11 g, 12.86 mmol, 1.0 equiv) was added. The mixture was stirred for 3 h, at which point it was filtered and the solid cake was dried under reduced pressure to afford the desired product as a white solid (7.5 g, 89% yield). LCMS (ESI) *m/z*: [M + H] calcd for C₃₂H₄₇N₉O₆: 654.37; found 654.4.

Step 2: Synthesis of 2-(4-(2-(3-(4-(6-(*tert*-butoxycarbonyl)-5,6,7,8-tetrahydropyrido[4,3-d]pyrimidin-2-yl)piperazin-1-yl)-3-oxopropoxy)ethyl)piperazin-1-yl)pyrimidine-5-carboxylic acid

[00680] To a solution of *tert*-butyl 2-(4-(3-(2-(4-(5-(ethoxycarbonyl)pyrimidin-2-yl)piperazin-1-yl)ethoxy)propanoyl)piperazin-1-yl)-7,8-dihydropyrido[4,3-d]pyrimidine-6(5H)-carboxylate (7.2 g, 11.01 mmol, 1.0 equiv) in THF (72 mL), EtOH (36 mL) and H₂O (36 mL) was added LiOH•H₂O (1.85 g, 44.05 mmol, 4.0 equiv). The reaction mixture was stirred at room temperature for 2.5 h, at which point the mixture was filtered and the filtrate was concentrated under reduced pressure to remove THF and EtOH. The aqueous phase was neutralized to pH = 7 with 1N HCl, and then concentrated under reduced pressure. Purification by reverse phase chromatography (30% MeCN/H₂O) afforded the desired

product as a white solid (3.85 g, 54% yield). LCMS (ESI) m/z : $[M + H]^+$ calcd for $C_{30}H_{43}N_9O_6$: 626.34; found 626.3.

Building Block AS. 2-(4-(2-(2-(3-(4-(5-((di-*tert*-butoxycarbonylamino)methyl)pyrimidin-2-yl)piperazin-1-yl)-3-oxo-propoxy)ethoxy)ethyl)piperazin-1-yl)pyrimidine-5-carboxylic acid.



Step 1: Synthesis of 3-(2-(2-(4-(5-(ethoxycarbonyl)pyrimidin-2-yl)piperazin-1-yl)ethoxy)ethoxy)propanoic acid

[00681] A solution of ethyl 2-(4-(2-(2-(3-(*tert*-butoxy)-3-oxopropoxy)ethoxy)ethyl)piperazin-1-yl)pyrimidine-5-carboxylate (4 g, 8.84 mmol, 1.0 equiv) in TFA (12.29 mL, 166.00 mmol, 18.8 equiv) was stirred at room temperature for 3 h. The reaction mixture was concentrated under reduced pressure. Purification by silica gel chromatography (0→20% MeOH/EtOAc) afforded the desired product as a brown oil (4.35 g, 95% yield, TFA salt).

*Step 2: Synthesis of ethyl 2-(4-(2-(2-(3-(4-(5-((bis(*tert*-butoxycarbonyl)amino)methyl)pyrimidin-2-yl)piperazin-1-yl)-3-oxopropoxy)ethoxy)ethyl)piperazin-1-yl)pyrimidine-5-carboxylate*


[00682] To a solution of 3-(2-(2-(4-(5-ethoxycarbonylpyrimidin-2-yl)piperazin-1-yl)ethoxy)ethoxy)propanoic acid (3.8 g, 7.44 mmol, 1.0 equiv, TFA) in DCM (30 mL) was added HATU (4.25 g, 11.17 mmol, 1.5 equiv) and DIPEA (6.48 mL, 37.22 mmol, 5.0 equiv). The reaction was stirred at room temperature for 30 min, and then *tert*-butyl *N-tert*-butoxycarbonyl-*N*-((2-piperazin-1-ylpyrimidin-5-yl)methyl)carbamate (2.93 g, 7.44 mmol, 1.0 equiv) was added. The mixture was stirred at room temperature for 3.5 h, at which point the reaction mixture was concentrated under reduced pressure. Purification by silica gel

chromatography (0→20% MeOH/EtOAc) afforded the desired product as a brown oil (4.14 g, 70% yield).

Step 3: Synthesis of 2-(4-(2-(2-(3-(4-(5-(((di-*tert*-butoxycarbonylamino)methyl)pyrimidin-2-yl)piperazin-1-yl)-3-oxo-propoxy)ethoxy)ethyl)piperazin-1-yl)pyrimidine-5-carboxylic acid

[00683] To a solution of ethyl 2-(4-(2-(2-(3-(4-(5-(((bis(*tert*-butoxycarbonyl)amino)methyl)pyrimidin-2-yl)piperazin-1-yl)-3-oxo-propoxy)ethoxy)ethyl)piperazin-1-yl)pyrimidine-5-carboxylate (1.4 g, 1.81 mmol, 1.0 equiv) in THF (28 mL), EtOH (14 mL) and H₂O (14 mL) was added LiOH•H₂O (304.44 mg, 7.25 mmol, 4.0 equiv). The mixture was stirred at 40 °C for 30 min, at which point the reaction mixture was concentrated under reduced pressure. Purification by reverse phase chromatography (10→40% MeCN/H₂O) afforded the desired product as a yellow solid (500 mg, 43% yield).

Building block AT. 2-{4-[2-(2-{4-[5-({[(*tert*-butoxy)carbonyl]amino}methyl)pyrimidin-2-yl]piperazin-1-yl};ethoxy)ethyl]piperazin-1-yl}pyrimidine-5-carboxylic acid.



Step 1: Synthesis of ethyl 2-(4-(2-(2-(4-(5-(((di-*tert*-butoxycarbonyl)amino)methyl)pyrimidin-2-yl)piperazin-1-yl)ethoxy)ethyl)piperazin-1-yl)pyrimidine-5-carboxylate


[00684] To a solution of ethyl 2-(4-(2-(2-(piperazin-1-yl)ethoxy)ethyl)piperazin-1-yl)pyrimidine-5-carboxylate hydrochloride (7.3 g, 13.56 mmol, 1.0 equiv, 4HCl) in DMF (75 mL) was added DIPEA (14.17 mL, 81.36 mmol, 6.0 equiv) and *tert*-butyl-*N*-*tert*-butoxycarbonyl-*N*-[(2-chloropyrimidin-5-yl)methyl]carbamate (5.59 g, 16.27 mmol, 1.2 equiv). The mixture was stirred at 80 °C for 12 h. The mixture was then cooled to room temperature and poured into H₂O (300 mL). The aqueous phase was extracted with EtOAc (3 x 80 mL). The combined organic phases were washed with sat. NH₄Cl (4 x 80 mL) and brine (150 mL), dried, filtered and the filtrate was concentrated under reduced pressure.

Purification by silica gel chromatography (0%→17% MeOH/EtOAc) afforded the desired product (7.7 g, 81.1% yield) as a light yellow oil. LCMS (ESI) m/z : $[M + Na]$ calcd for $C_{34}H_{53}N_9O_7$: 722.40; found 722.4.

*Step 2: Synthesis of 2-(4-(2-(2-(4-(5-(((*tert*-butoxycarbonyl)amino)methyl)pyrimidin-2-yl)piperazin-1-yl)ethoxy)ethyl)piperazin-1-yl)pyrimidine-5-carboxylic acid*

[00685] To a solution of ethyl 2-(4-(2-(2-(4-(5-(((*di-tert*-butoxycarbonyl)amino)methyl)pyrimidin-2-yl)piperazin-1-yl)ethoxy)ethyl)piperazin-1-yl)pyrimidine-5-carboxylate (7.7 g, 11.00 mmol, 1.0 equiv) in THF (80 mL), EtOH (20 mL), and H₂O (40 mL) was added LiOH•H₂O (2.31 g, 55.01 mmol, 5.0 equiv). The mixture was stirred at 50 °C for 26 h. The mixture was then concentrated under reduced pressure to remove THF and EtOH. The aqueous phase was neutralized with 0.5 N HCl, and concentrated under reduced pressure. Purification by reverse phase chromatography afforded the desired product (4.67 g, 74.3% yield) as a white solid. LCMS (ESI) m/z : $[M - H]$ calcd for $C_{27}H_{41}N_9O_5$: 570.31; found 570.3.

Building block AU. (*R*)-*tert*-butyl 4-(5-(((*tert*-butoxycarbonyl-*N-tert*-butoxycarbonyl)amino)methyl)pyrimidin-2-yl)piperazine-2-carboxylate.



*Step 1: Synthesis of (*R*)-1,4-bis((benzyloxy)carbonyl)piperazine-2-carboxylic acid*

[00686] To two separate batches containing a solution (*2R*)-piperazine-2-carboxylic acid (70 g, 344.71 mmol, 1 equiv, 2HCl) in H₂O (700 mL) and dioxane (1120 mL) was added 50% aq. NaOH until pH = 11. Benzyl chloroformate (156.82 mL, 1.10 mol, 3.2 equiv) was added and the reaction was stirred at room temperature for 12 h. The two reaction mixtures were combined and H₂O (1200 mL) was added. The aqueous layer was extracted with MTBE (3 x 1000 mL), adjusted to pH = 2 with con. HCl, and then extracted with EtOAc (2 x 1000 mL). The combined organic phases were dried, filtered, and concentrated under reduced

pressure to afford the desired product (280 g, 86% yield). LCMS (ESI) m/z : $[M + H]$ calcd for $C_{21}H_{22}N_2O_6$: 399.16; found 399.0.

Step 2: Synthesis of (R)-1,4-dibenzyl 2-tert-butyl piperazine-1,2,4-tricarboxylate

[00687] To a solution of (*R*)-1,4-bis((benzyloxy)carbonyl)piperazine-2-carboxylic acid (70 g, 175.70 mmol, 1.0 equiv) in toluene (700 mL) at 80 °C was added 1,1-di-*tert*-butoxy-*N,N*-dimethyl-methanamine (80.04 mL, 333.83 mmol, 1.9 equiv). The reaction was stirred at 80 °C for 2 h, at which point it was cooled to room temperature and partitioned between EtOAc (300 mL) and H₂O (500 mL). The aqueous layer was extracted with EtOAc (2 x 500 mL) and the combined organic layers were dried, filtered, and concentrated under reduced pressure. Purification by silica gel chromatography (0→25 EtOAc/petroleum ether) afforded the desired product as a white solid (50 g, 57% yield). LCMS (ESI) m/z : $[M + Na]$ calcd for $C_{25}H_{30}N_2O_6$: 477.20; found 476.9.


Step 3: Synthesis of (R)-tert-butyl piperazine-2-carboxylate

[00688] To a solution of (*R*)-1,4-dibenzyl 2-*tert*-butyl piperazine-1,2,4-tricarboxylate (50 g, 110.01 mmol, 1 equiv) in EtOAc (20 mL) was added Pd/C (15 g, 10 wt.%). The suspension was degassed under reduced pressure and purged with H₂ three times. The suspension was stirred under H₂ (30 psi) at 30 °C for 4 h. The reaction mixture was then filtered, the residue was washed with MeOH (5 x 200 mL), and the filtrate concentrated under reduced pressure to afford the desired product as a yellow oil (17 g, 81% yield). LCMS (ESI) m/z : $[M + H]$ calcd for $C_9H_{18}N_2O_2$: 187.15; found 187.1.

*Step 4: Synthesis of (R)-tert-butyl 4-(5-(((tert-butoxycarbonyl-*N*-tert-butoxycarbonyl)amino)methyl)pyrimidin-2-yl)piperazine-2-carboxylate*


[00689] To a suspension of (*R*)-*tert*-butyl piperazine-2-carboxylate (8 g, 23.27 mmol, 1.0 equiv) and *tert*-butyl-*N*-*tert*-butoxycarbonyl ((2-chloropyrimidin-5-yl)methyl)carbamate (5.20 g, 27.92 mmol, 1.2 equiv) in MeCN (100 mL) was added K₂CO₃ (6.43 g, 46.54 mmol, 2.0 equiv). The reaction mixture was heated to 80 °C for 12 h, at which point it was cooled to room temperature, filtered, and the filtrate was concentrated under reduced pressure. Purification by silica gel chromatography (0→100% EtOAc/petroleum ether) afforded the desired product as a yellow solid (9.2 g, 73% yield). LCMS (ESI) m/z : $[M + H]$ calcd for $C_{24}H_{39}N_5O_6$: 494.30; found 494.1.

Building block AV. (*S*)-*tert*-butyl 4-(5-(((*tert*-butoxycarbonyl-*N*-*tert*-butoxycarbonyl)amino)methyl)pyrimidin-2-yl)piperazine-2-carboxylate.



[00690] This building block is prepared by a process similar to that for Building block AU by utilizing (2*S*)-piperazine-2-carboxylic acid.

Building block AW. (*R*)-2-(4-(2-(3-(2-(*tert*-butoxycarbonyl)-4-(5-(((*tert*-butoxycarbonyl-*N*-*tert*-butoxycarbonyl)amino)methyl)pyrimidin-2-yl)piperazin-1-yl)-3-oxopropoxy)ethyl)piperazin-1-yl)pyrimidine-5-carboxylic acid.



Step 1: Synthesis of (*R*)-ethyl 2-(4-(2-(3-(2-(*tert*-butoxycarbonyl)-4-(5-(((*tert*-butoxycarbonyl-*N*-*tert*-butoxycarbonyl)amino)methyl)pyrimidin-2-yl)piperazin-1-yl)-3-oxopropoxy)ethyl)piperazin-1-yl)pyrimidine-5-carboxylate


[00691] To a solution of (*R*)-*tert*-butyl 4-(5-(((*tert*-butoxycarbonyl-*N*-*tert*-butoxycarbonyl)amino)methyl)pyrimidin-2-yl)piperazine-2-carboxylate (5.3 g, 11.36 mmol, 1.0 equiv, TFA) in DCM (80 mL) was added HATU (6.48 g, 17.05 mmol, 1.5 equiv) and DIPEA (7.92 mL, 45.45 mmol, 4.0 equiv). The reaction was stirred at room temperature for 30 min and then *tert*-butyl (2*R*)-4-(5-(((bis(*tert*-butoxycarbonyl)amino)methyl)pyrimidin-2-yl)piperazine-2-carboxylate (5.61 g, 11.36 mmol, 1.0 equiv) was added. The mixture was stirred for 1 h, at which point sat. NH₄Cl (80 mL) was added. The organic phase was washed

with sat. NH_4Cl (5 x 80 mL), dried, filtered, and concentrated under reduced pressure. Purification by silica gel chromatography (0→9% MeOH/EtOAc) afforded the desired product as a yellow solid (8.4 g, 85% yield).

Step 2: Synthesis of (R)-2-(4-(2-(3-(2-(tert-butoxycarbonyl)-4-(5-(((tert-butoxycarbonyl-N-tert-butoxycarbonyl)amino)methyl)pyrimidin-2-yl)piperazin-1-yl)-3-oxopropoxy)ethyl)piperazin-1-yl)pyrimidine-5-carboxylic acid


[00692] To two separate batches containing a solution a solution of (R)-ethyl 2-(4-(2-(3-(2-(tert-butoxycarbonyl)-4-(5-(((tert-butoxycarbonyl-N-tert-butoxycarbonyl)amino)methyl)pyrimidin-2-yl)piperazin-1-yl)-3-oxopropoxy)ethyl)piperazin-1-yl)pyrimidine-5-carboxylate (3.4 g, 4.11 mmol, 1.0 equiv) in THF (16 mL), EtOH (8 mL) and H_2O (8 mL) was added $\text{LiOH}\cdot\text{H}_2\text{O}$ (344.61 mg, 8.21 mmol, 2.0 equiv). The mixture was stirred at room temperature for 2 h. The two reaction mixtures were then combined and were adjusted to pH = 7 with 1N HCl. The solution was concentrated under reduced pressure to remove THF and EtOH. The solution was then filtered, and the resulting solid was purified by reverse phase chromatography to afford the desired product as a white solid (4 g, 59% yield). LCMS (ESI) m/z : $[\text{M} + \text{H}]$ calcd for $\text{C}_{38}\text{H}_{57}\text{N}_9\text{O}_{10}$: 800.43; found 800.3.

Building block AX. (S)-2-(4-(2-(3-(2-(tert-butoxycarbonyl)-4-(5-(((tert-butoxycarbonyl-N-tert-butoxycarbonyl)amino)methyl)pyrimidin-2-yl)piperazin-1-yl)-3-oxopropoxy)ethyl)piperazin-1-yl)pyrimidine-5-carboxylic acid.



[00693] This building block is prepared from Building block AV by a process similar to that for Building block AW.

Building block AY. 1'-(2-(3-(4-(5-(((*tert*-butoxycarbonyl)amino)methyl)pyrimidin-2-yl)piperazin-1-yl)-3-oxopropoxy)ethyl)-[1,4'-bipiperidine]-4-carboxylic acid.



Step 1: Synthesis of 1'-tert-butyl 4-ethyl [1,4'-bipiperidine]-1',4-dicarboxylate

[00694] To a solution of ethyl piperidine-4-carboxylate (30 g, 150.57 mmol, 1.0 equiv) and *tert*-butyl 4-oxopiperidine-1-carboxylate (23.67 g, 150.57 mmol, 1.0 equiv) in DCM (300 mL) was added HOAc (6.00 mL, 104.95 mmol, 0.7 equiv). The mixture was stirred at room temperature for 30 min, then NaBH(OAc)₃ (63.82 g, 301.13 mmol, 2.0 equiv) was added. The mixture was stirred for 16 h, at which point H₂O (50 mL) was added. The aqueous phase was extracted with DCM (3 x 15 mL) and the combined organic phases were washed with brine (10 mL), dried, filtered, and concentrated under reduced pressure. Purification by silica gel chromatography (8→100 MeOH/EtOAc) afforded the desired product as a yellow oil (30 g, 59% yield).

Step 2: Synthesis of ethyl [1,4'-bipiperidine]-4-carboxylate

[00695] To a solution of HCl in EtOAc (200 mL) was added 1'-*tert*-butyl 4-ethyl [1,4'-bipiperidine]-1',4-dicarboxylate (20 g, 58.74 mmol, 1.0 equiv). The mixture was stirred at room temperature for 3 h. The mixture was then concentrated under reduced pressure to afford the desired crude product as a white solid (15 g, HCl salt).

*Step 3: Synthesis of ethyl 1'-(2-(3-(4-(5-(((*tert*-butoxy)-3-oxopropoxy)ethyl)-[1,4'-bipiperidine]-4-carboxylate*

[00696] To a solution of *tert*-butyl 3-(2-bromoethoxy)propanoate (6.46 g, 25.54 mmol, 1.0 equiv) in DMF (240 mL) was added K₂CO₃ (10.59 g, 76.61 mmol, 3.0 equiv) and ethyl [1,4'-bipiperidine]-4-carboxylate (8 g, 25.54 mmol, 1.0 equiv, 2HCl). The mixture was stirred at 120 °C for 12 h, at which point the reaction was cooled to room temperature, filtered, the filter cake washed with H₂O (20 mL), and the filtrate was concentrated under reduced pressure. Purification by silica gel chromatography (0→11% MeOH/EtOAc) afforded the desired product as a yellow oil (6.6 g, 63% yield).

Step 4: Synthesis of 3-(2-(4-(ethoxycarbonyl)-[1,4'-bipiperidin]-1'-yl)ethoxy)propanoic acid

[00697] To the solution of HCl in EtOAc (70 mL) was added ethyl 1'-(2-(3-(*tert*-butoxy)-3-oxopropoxy) ethyl)-[1,4'-bipiperidine]-4-carboxylate (6.6 g, 16.00 mmol, 1.0 equiv). The mixture was stirred at room temperature for 3 h, at which point the reaction was concentrated under reduced pressure to afford the desired product as a white solid (6.5 g, 95% yield, 2HCl).

*Step 5: Synthesis of ethyl 1'-(2-(3-(4-(5-(((*N,N*-di-*tert*-butoxycarbonyl)amino)methyl)pyrimidin-2-yl)piperazin-1-yl)-3-oxopropoxy)ethyl)-[1,4'-bipiperidine]-4-carboxylate*


[00698] To a solution of *tert*-butyl-*tert*-butoxycarbonyl((2-(piperazin-1-yl)pyrimidin-5-yl)methyl)carbamate (2.49 g, 6.33 mmol, 1.5 equiv) in DMF (40 mL) was added DIPEA (9.74 mL, 55.89 mmol, 6.0 equiv) and HATU (5.31 g, 13.97 mmol, 1.5 equiv). The mixture was stirred at room temperature for 30 min, and then 3-(2-(4-(ethoxycarbonyl)-[1,4'-bipiperidin]-1'-yl)ethoxy) propanoic acid (4 g, 9.32 mmol, 1.0 equiv, 2HCl) was added. The mixture was stirred at for 1.5 h, at which point H₂O (5 mL) and EtOAc (20 mL) were added. The aqueous phase was extracted with EtOAc (3 x 10 mL) and the combined organic phases were washed with brine (5 mL), dried, filtered and concentrated under reduced pressure. Purification by reverse phase chromatography afforded the desired product as a brown oil (1.6 g, 23% yield). LCMS (ESI) *m/z*: [M + H] calcd for C₃₇H₆₁N₇O₈: 732.47; found 732.6.

*Step 6: Synthesis of 1'-(2-(3-(4-(5-(((*tert*-butoxycarbonyl)amino)methyl)pyrimidin-2-yl)piperazin-1-yl)-3-oxopropoxy)ethyl)-[1,4'-bipiperidine]-4-carboxylic acid*

[00699] To a solution of ethyl 1'-(2-(3-(4-(5-(((*N,N*-di-*tert*-butoxycarbonyl)amino)methyl)pyrimidin-2-yl)piperazin-1-yl)-3-oxopropoxy)ethyl)-[1,4'-bipiperidine]-4-carboxylate (1.4 g, 1.91 mmol, 1.0 equiv) in THF (7.5 mL), EtOH (3.8 mL), and H₂O (3.8 mL) was added LiOH•H₂O (321.07 mg, 7.65 mmol, 4.0 equiv). The mixture

was stirred at room temperature for 2 h, at which point the mixture was concentrated under reduced pressure. Purification by reverse phase chromatography (5→38% MeCN/H₂O) afforded the desired product as a yellow solid (325 mg, 22% yield). LCMS (ESI) *m/z*: [M + H] calcd for C₃₀H₄₉N₇O₆: 604.38; found 604.3.

Building block AZ. 1-(4-{2-[2-(2-[[[(benzyloxy)carbonyl]amino]ethoxy]ethoxy]ethyl}piperazin-1-yl)-3,6,9,12-tetraoxapentadecan-15-oic acid.



Step 1: Synthesis of benzyl (2-(2-(2-bromoethoxy)ethoxy)ethyl)carbamate

[00700] To a solution of benzyl (2-(2-(2-hydroxyethoxy)ethoxy)ethyl)carbamate (10 g, 35.30 mmol, 1.0 equiv) in DCM (300 mL) at 0 °C was added PPh₃ (13.79 g, 52.59 mmol, 1.49 equiv) and CBr₄ (17.44 g, 52.59 mmol, 1.49 equiv). Then the mixture was warmed to room temperature and stirred for 12 h. The reaction mixture was then filtered, and the filtrate was concentrated under reduced pressure. Purification by silica gel chromatography (1%→25% EtOAc/petroleum ether) afforded the desired product (10.8 g, 88.4% yield) as yellow oil.

Step 2: Synthesis of tert-butyl 4-(3-oxo-1-phenyl-2,7,10-trioxa-4-azadodecan-12-yl)piperazine-1-carboxylate

[00701] To a solution of benzyl (2-(2-(2-bromoethoxy)ethoxy)ethyl)carbamate (10.8 g, 31.19 mmol, 1.0 equiv) and *tert*-butyl piperazine-1-carboxylate (5.81 g, 31.19 mmol, 1.0 equiv) in MeCN (100 mL) was added K₂CO₃ (4.31 g, 31.19 mmol, 1.0 equiv). The mixture was stirred at 80 °C for 1 h. The reaction mixture was then filtered, and the filtrate was concentrated under reduced pressure. Purification by silica gel chromatography (0%→50% MeOH/EtOAc) afforded the desired product (13.1 g, 93.0% yield) as yellow oil.

Step 3: Synthesis of benzyl (2-(2-(2-(piperazin-1-yl)ethoxy)ethoxy)ethyl)carbamate

[00702] A solution of *tert*-butyl 4-(3-oxo-1-phenyl-2,7,10-trioxa-4-azadodecan-12-yl)piperazine-1-carboxylate (5.64 g, 12.49 mmol, 1.0 equiv) in HCl/EtOAc (50 mL, 4 M) was stirred at room temperature for 1 h. The reaction mixture was then concentrated under reduced pressure to afford the desired product (5.23 g, crude, HCl salt) as yellow oil.


Step 4: Synthesis of tert-butyl 1-(4-(3-oxo-1-phenyl-2,7,10-trioxa-4-azadodecan-12-yl)piperazin-1-yl)-3,6,9,12-tetraoxapentadecan-15-oate

[00703] A solution of benzyl (2-(2-(2-(piperazin-1-yl)ethoxy)ethoxy)ethyl)carbamate (13.3 g, 31.34 mmol, 1.0 equiv, 2HCl) and *tert*-butyl 1-bromo-3,6,9,12-tetraoxapentadecan-15-oate in MeCN (150 mL) was added K₂CO₃ (21.66 g, 156.71 mmol, 5.0 equiv). The mixture was stirred at 80 °C for 12 h. The reaction mixture was then filtered, and the filtrate was concentrated under reduced pressure. Purification by silica gel chromatography (1%→17% MeOH/DCM) afforded the desired product (5.4 g, 26.3% yield) as a yellow oil.

Step 5: Synthesis of 1-(4-(3-oxo-1-phenyl-2,7,10-trioxa-4-azadodecan-12-yl)piperazin-1-yl)-3,6,9,12-tetraoxapentadecan-15-oic acid

[00704] A solution of *tert*-butyl 1-(4-(3-oxo-1-phenyl-2,7,10-trioxa-4-azadodecan-12-yl)piperazin-1-yl)-3,6,9,12-tetraoxapentadecan-15-oate (2.4 g, 3.66 mmol, 1.0 equiv) in TFA (20 mL) was stirred at room temperature for 30 min. The reaction mixture was then concentrated under reduced pressure to afford the desired product (3.03 g, TFA salt) as yellow oil.

Building Block BA. (*R*)-2-(2-(*tert*-butoxycarbonyl)-4-(6-(*tert*-butoxycarbonyl)-5,6,7,8-tetrahydropyrido[4,3-*d*]pyrimidin-2-yl)piperazin-1-yl)pyrimidine-5-carboxylic acid.



*Step 1: Synthesis of (*R*)-tert-butyl 2-(3-(*tert*-butoxycarbonyl)-4-(5-(ethoxycarbonyl)pyrimidin-2-yl)piperazin-1-yl)-7,8-dihydropyrido[4,3-*d*]pyrimidine-6(5H)-carboxylate*


[00705] To two separate batches run in parallel each containing a solution of (*R*)-*tert*-butyl 2-(3-(*tert*-butoxycarbonyl)piperazin-1-yl)-7,8-dihydropyrido[4,3-*d*]pyrimidine-6(5H)-

carboxylate (6 g, 14.30 mmol, 1.0 equiv) and K_2CO_3 (3.95 g, 28.60 mmol, 2.0 equiv) in MeCN (80 mL) was added ethyl 2-chloropyrimidine-5-carboxylate (3.20 g, 17.16 mmol, 1.2 equiv). The reaction mixtures were stirred at 80 °C for 12 h. The two reactions mixtures were combined and filtered, the residue was washed with EtOAc (3 x 50 mL), and the filtrate was concentrated under reduced pressure. Purification by silica gel chromatography (0%→17% MeOH/EtOAc) afforded the desired product (15 g, 91.5% yield) as a yellow solid. LCMS (ESI) m/z : $[M + H]$ calcd for $C_{28}H_{39}N_7O_6$: 570.31; found 570.1.

Step 2: Synthesis of (R)-2-(2-(tert-butoxycarbonyl)-4-(6-(tert-butoxycarbonyl)-5,6,7,8-tetrahydropyrido[4,3-d]pyrimidin-2-yl)piperazin-1-yl)pyrimidine-5-carboxylic acid


[00706] To a solution of (*R*)-*tert*-butyl 2-(3-(*tert*-butoxycarbonyl)-4-(5-(ethoxycarbonyl)pyrimidin-2-yl)piperazin-1-yl)-7,8-dihydropyrido[4,3-d]pyrimidine-6(5H)-carboxylate (15 g, 26.33 mmol, 1.0 equiv) in THF (80 mL), EtOH (40 mL) and H_2O (40 mL) was added $LiOH \cdot H_2O$ (2.21 g, 52.66 mmol, 2.0 equiv). The mixture was stirred at room temperature for 6 h. The reaction mixture was then adjusted to pH=6 with 1 N HCl. The resulting suspension was filtered, and the solid cake was dried under reduced pressure to afford the desired product (10.87 g, 75.9% yield) as a white solid. LCMS (ESI) m/z : $[M + H]$ calcd for $C_{26}H_{35}N_7O_6$: 542.27; found 542.1.

Building Block BB. (*S*)-2-(2-(tert-butoxycarbonyl)-4-(6-(tert-butoxycarbonyl)-5,6,7,8-tetrahydropyrido[4,3-d]pyrimidin-2-yl)piperazin-1-yl)pyrimidine-5-carboxylic acid.




[00707] This building block is prepared from Building block AA by a process similar to that for Building block BA.

Building Block BC. 2-[(2*R*)-2-[(*tert*-butoxy)carbonyl]-4-[5-({[(*tert*-butoxy)carbonyl]amino}methyl)pyrimidin-2-yl]piperazin-1-yl]pyrimidine-5-carboxylic acid.




[00708] This building block is prepared from Building block AU by a process similar to that for Building block BA.

Building Block BD. 2-[(2*S*)-2-[(*tert*-butoxy)carbonyl]-4-[5-({[(*tert*-butoxy)carbonyl]amino}methyl)pyrimidin-2-yl]piperazin-1-yl]pyrimidine-5-carboxylic acid.



[00709] This building block is prepared from Building block AV by a process similar to that for Building block BA.

Building block BE. 15-(6-((4-amino-3-(2-aminobenzo[d]oxazol-5-yl)-1*H*-pyrazolo[3,4-*d*]pyrimidin-1-yl)methyl)-3,4-dihydroisoquinolin-2(1*H*)-yl)-1-((1*S*,4*S*)-5-(2-(2-(2-aminoethoxy)ethoxy)ethyl)-2,5-diazabicyclo[2.2.1]heptan-2-yl)-3,6,9,12-tetraoxapentadecan-15-one.



Step 1: Synthesis of (1*S*,4*S*)-*tert*-butyl 5-(3-oxo-1-phenyl-2,7,10-trioxa-4-azadodecan-12-yl)-2,5-diazabicyclo[2.2.1]heptane-2-carboxylate

[00710] To a solution of (1*S*,4*S*)-*tert*-butyl 2,5-diazabicyclo[2.2.1]heptane-2-carboxylate (2.85 g, 14.37 mmol, 1.0 equiv) in MeCN (50 mL) was added K₂CO₃ (3.97 g, 28.75 mmol, 2.0 equiv) and benzyl (2-(2-(2-bromoethoxy)ethoxy)ethyl)carbamate (4.98 g, 14.37 mmol, 1.0 equiv). The mixture was stirred at 80 °C for 24 h. The reaction mixture was then filtered, and the filtrate was concentrated under reduced pressure. Purification by silica gel chromatography (0→10% MeOH/EtOAc) afforded the desired product (6.2 g, 93.0% yield) as colorless oil. LCMS (ESI) *m/z*: [M + H] calcd for C₂₄H₃₇N₃O₆: 464.27; found 464.2.

*Step 2: Synthesis of benzyl (2-(2-(2-((1*S*,4*S*)-2,5-diazabicyclo[2.2.1]heptan-2-yl)ethoxy)ethoxy)ethyl)carbamate*

[00711] To a solution of (1*S*,4*S*)-*tert*-butyl 5-(3-oxo-1-phenyl-2,7,10-trioxa-4-azadodecan-12-yl)-2,5-diazabicyclo[2.2.1]heptane-2-carboxylate (6.2 g, 13.37 mmol, 1.0 equiv) in DCM (60 mL) was added TFA (20.7 mL, 279.12 mmol, 20.9 equiv). The reaction was stirred for 2 h, at which point the mixture was concentrated under reduced pressure at 45 °C to afford the desired crude product (10.5 g, 4TFA) as light brown oil, which was used the next step directly. LCMS (ESI) *m/z*: [M + H] calcd for C₁₉H₂₉N₃O₄: 364.22; found 364.2.


*Step 3: Synthesis of tert-butyl 1-((1*S*,4*S*)-5-(3-oxo-1-phenyl-2,7,10-trioxa-4-azadodecan-12-yl)-2,5-diazabicyclo[2.2.1]heptan-2-yl)-3,6,9,12-tetraoxapentadecan-15-oate*

[00712] To a solution of benzyl (2-(2-(2-((1*S*,4*S*)-2,5-diazabicyclo[2.2.1]heptan-2-yl)ethoxy)ethoxy)ethyl)carbamate (5 g, 6.10 mmol, 1.0 equiv, 4TFA) in MeCN (80 mL) was added K₂CO₃ (5.06 g, 36.61 mmol, 6.0 equiv) and *tert*-butyl 1-bromo-3,6,9,12-tetraoxapentadecan-15-oate (2.35 g, 6.10 mmol, 1.0 equiv). The reaction mixture was stirred at 80 °C for 12 h. The reaction mixture was then filtered, and the filtrate was concentrated under reduced pressure. Purification by silica gel chromatography (0→15% MeOH/EtOAc) afforded the desired product (5.2 g, 92.8% yield) as light yellow oil. LCMS (ESI) *m/z*: [M + H] calcd for C₃₄H₅₇N₃O₁₀: 668.4; found 668.4.

*Step 4: Synthesis of 1-((1*S*,4*S*)-5-(3-oxo-1-phenyl-2,7,10-trioxa-4-azadodecan-12-yl)-2,5-diazabicyclo[2.2.1]heptan-2-yl)-3,6,9,12-tetraoxapentadecan-15-oic acid*

[00713] A solution of *tert*-butyl 1-((1*S*,4*S*)-5-(3-oxo-1-phenyl-2,7,10-trioxa-4-azadodecan-12-yl)-2,5-diazabicyclo[2.2.1]heptan-2-yl)-3,6,9,12-tetraoxapentadecan-15-oate (5.2 g, 5.66 mmol, 1.0 equiv) in TFA (47.3 mL, 638.27 mmol, 112.75 equiv) was stirred at room temperature for 30 min. The mixture was then concentrated under reduced pressure at 45 °C. Purification by reverse phase chromatography (2→35% MeCN/H₂O(0.05% NH₄OH)) afforded the desired product (1.88 g, 54.3% yield) as light brown oil. LCMS (ESI) *m/z*: [M + H] calcd for C₃₀H₄₉N₃O₁₀: 612.34; found 612.3.

Building block BF. 21-(6-((4-amino-3-(2-aminobenzo[d]oxazol-5-yl)-1*H*-pyrazolo[3,4-d]pyrimidin-1-yl)methyl)-3,4-dihydroisoquinolin-2(1*H*)-yl)-1-(piperazin-1-yl)-


3,6,9,12,15,18-hexaoxahenicosan-21-one.

Step 1: Synthesis of benzyl 4-(23,23-dimethyl-21-oxo-3,6,9,12,15,18,22-heptaooxatetracosyl) piperazine-1-carboxylate

[00714] To a solution of *tert*-butyl 1-bromo-3,6,9,12,15,18-hexaoxahenicosan-21-oate (5 g, 10.56 mmol, 1.0 equiv) and benzyl piperazine-1-carboxylate (2.62 mL, 11.62 mmol, 1.1 equiv, HCl) in MeCN (50 mL) was added K₂CO₃ (4.38 g, 31.69 mmol, 3.0 equiv). The reaction mixture was stirred at 80 °C for 10 h. The mixture was then filtered, the solid cake washed with EtOAc (3 x 3 mL), and the filtrate concentrated under reduced pressure. Purification by silica gel chromatography (0→10% MeOH/EtOAc) afforded the desired product (4 g, 61.8% yield) as a red liquid.


Step 2: Synthesis of 1-(4-((benzyloxy)carbonyl)piperazin-1-yl)-3,6,9,12,15,18-hexaoxahenicosan-1-oic acid

[00715] To a solution of benzyl 4-(23,23-dimethyl-21-oxo-3,6,9,12,15,18,22-heptaooxatetracosyl)piperazine-1-carboxylate (1.8 g, 2.94 mmol, 1.0 equiv) in DCM (10 mL) was added TFA (10 mL). The solution was stirred for 0.5 h. The solution was then concentrated under reduced pressure. To the residue was added DCM (30 mL) and then the solution was concentrated under reduced pressure to afford the desired product (1.6 g, 2.87 mmol, TFA) as a red liquid.

Preparation of Rapamycin Monomers.**Monomer 1. 40(*R*)-*O*-(4-nitrophenyl)carbonate rapamycin.**

[00716] To a solution of rapamycin (30.10 g, 32.92 mmol, 1.0 equiv) in DCM (148.9 mL) was added pyridine (29.6 mL, 367 mmol, 11.1 equiv). The solution was cooled to -78 °C and then *p*-nitrophenyl chloroformate (12.48 g, 61.92 mmol, 1.9 equiv) was added. The reaction was stirred at -78 °C for 2 h. To the reaction mixture was then added DCM and the solution was then poured into H₂O. The aqueous layer was extracted with DCM and the combined organic layers were dried over MgSO₄, and concentrated under reduced pressure. The crude material was purified by silica gel chromatography (0→50% EtOAc/hexanes) to provide the product (23.1 g, 59.2% yield) as a white solid. LCMS (ESI) *m/z*: [M + Na] calcd for C₅₈H₈₂N₂O₁₇: 1101.55; found 1101.6.

Monomer 2. 32(*R*)-hydroxy 40(*R*)-*O*-(4-nitrophenyl)carbonate rapamycin.



Step 1: Synthesis of 32(*R*)-hydroxy rapamycin


[00717] A solution of 32(*R*)-hydroxy-28,40-bistriethylsilyl rapamycin (3.64 g, 3.18 mmol, 1 equiv) in THF (41.8 mL) was treated with pyridine (20.8 mL, 258 mmol, 81 equiv) and the reaction mixture was cooled to 0 °C. The solution was treated dropwise with 70% HF-pyridine (4.60 mL, 159 mmol, 50 equiv) and the reaction mixture was stirred at 0 °C for 20 min followed by warming to room temperature. After 5 h, the reaction mixture was cooled back to 0 °C and carefully added to ice cold sat. NaHCO₃ solution (400 mL). The mixture

was extracted with EtOAc (2 x 100 mL) and the organic phases were washed with 75 mL portions of H₂O, sat. NaHCO₃ solution and brine. The organic solution was dried over Na₂SO₄, filtered and concentrated to yield a light yellow oil that produced a stiff foam under reduced pressure. The crude material was purified by silica gel chromatography (20→40% acetone/hexanes) to yield the desired product as a white amorphous solid (1.66 g, 57% yield). LCMS (ESI) *m/z*: [M + Na] calcd for C₅₁H₈₁NO₁₃: 938.56; found 938.7; *m/z*: [M - H] calcd for C₅₁H₈₁NO₁₃: 914.56; found 914.7.

Step 2: Synthesis of 32(R)-hydroxy 40(R)-O-(4-nitrophenyl)carbonate rapamycin

[00718] To a suspension of powdered 4Å molecular sieves (6.0 g) in DCM (130 mL) was added 32(*R*)-hydroxy rapamycin (6.00 g, 6.55 mmol, 1.0 equiv). After stirring at room temperature for 45 min, pyridine (5.99 mL, 74.0 mmol, 11.3 equiv) was added. The suspension was cooled to -15 °C and then 4-nitrophenylchloroformate (1.78 g, 8.84 mmol, 1.4 equiv) was then added. The reaction mixture was stirred at -10 °C for 2 h and then filtered, and the filter pad washed with DCM (140 mL). The filtrate was washed with sat. NaHCO₃ (130 mL), H₂O (130 mL) and brine (130 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The crude material was purified by silica gel chromatography (20→50% EtOAc/hexanes) to give the product (4.44 g, 63% yield) as an off-white stiff foam. LCMS (ESI) *m/z*: [M + Na] calcd for C₅₈H₈₄N₂O₁₇: 1103.57; found 1103.5.

Monomer 3. 32(*R*)-methoxy 40(*R*)-O-(4-nitrophenyl)carbonate rapamycin.



Step 1: Synthesis of 32(R)-methoxy-28,40-bistriethylsilyl rapamycin

[00719] To a stirred solution of 32(R)-hydroxy-28,40-bistriethylsilyl rapamycin (3.83 g, 3.34 mmol, 1.0 equiv) in chloroform (95.8 mL) was added Proton Sponge® (7.17 g, 33.5 mmol, 10.0 equiv) along with freshly dried 4 Å molecular sieves (4 g). The solution was stirred for 1 h prior to the addition of trimethyloxonium tetrafluoroborate (4.95 g, 33.5 mmol, 10.0 equiv, dried by heating under reduced pressure at 50°C for 1 h before use) at room temperature. The reaction mixture was stirred for 18 h, and then the reaction mixture was diluted with DCM and filtered through Celite. The filtrate was washed sequentially with aqueous 1 M HCl (2x), sat. aqueous NaHCO_3 solution, then dried and concentrated under reduced pressure. Purification by silica gel chromatography (10 $\rightarrow$ 20% EtOAc/hexanes) afforded the desired product as a yellow oil that was contaminated with 3 wt.% Proton Sponge®. The residue was taken up in MTBE and washed with aqueous 1 M HCl, sat. aqueous NaHCO_3 solution, dried, and then concentrated under reduced pressure to furnish a yellow foam (3.15 g, 81.2% yield). LCMS (ESI) m/z : $[\text{M} - \text{TES} + \text{H}_2\text{O}]$ calcd for $\text{C}_{64}\text{H}_{111}\text{NO}_{13}\text{Si}_2$: 1061.68; found 1061.9.


Step 2: Synthesis of 32(R)-methoxy rapamycin

[00720] To a stirred solution of 32(R)-methoxy-28,40-bistriethylsilyl rapamycin (1.11 g, 0.958 mmol, 1.0 equiv) in THF (12.6 mL) and pyridine (6.30 mL) in a plastic vial was added

70% HF-pyridine (2.22 mL, 76.6 mmol, 80.0 equiv) dropwise at 0 °C. The reaction mixture was stirred at 0 °C for 20 min before being warmed to room temperature for 3 h, when HPLC showed complete consumption of starting material. The reaction mixture was cooled to 0 °C and poured slowly into ice cold sat. aqueous NaHCO₃ solution (50 mL). The aqueous layer was extracted with EtOAc (3x) and the combined organics were washed with sat. aqueous NaHCO₃ solution, brine, dried, and concentrated under reduced pressure. The yellow residue was dissolved in MeOH (5 mL) and added dropwise to H₂O (50 mL) to produce a white precipitate. After stirring for 15 min the slurry was filtered on a medium porosity funnel and the cake washed with H₂O (2x). The solids were then dissolved in MeCN (50 mL) and lyophilized overnight to provide the product as a white solid (780 mg, 87% yield). LCMS (ESI) *m/z*: [M + Na] calcd for C₅₂H₈₃NO₁₃: 952.58; found 952.4.

Step 3: Synthesis of 32(R)-methoxy 40(R)-O-(4-nitrophenyl)carbonate rapamycin

[00721] To a solution of 32(*R*)-methoxy rapamycin (4.50 g, 4.84 mmol, 1.0 equiv) in DCM (180 mL) was added powdered 4Å molecular sieves (6.0 g). The mixture was stirred at room temperature for 1 h and then pyridine (3.91 mL, 48.4 mmol, 10 equiv) was added. The mixture was cooled to -10 °C and 4-nitrophenylchloroformate (0.990 g, 4.91 mmol, 1.0 equiv) was added in one portion. The reaction was allowed to slowly warm to room temperature and after 3 h the reaction mixture was cooled to 0 °C and 4-nitrophenylchloroformate (250 mg, 1.24 mmol, 0.3 equiv) was added. The mixture was warmed to room temperature and after 1 h the reaction mixture was filtered through a pad of celite and the pad was washed with DCM (140 mL). The filtrate was washed with H₂O (120 mL) and sat NaHCO₃ (2 x 120 mL). The organic phase was dried over Na₂SO₄, filtered and concentrated under reduced pressure. The crude material was purified by flash chromatography (20→50% EtOAc/hex) to yield a white stiff foam. The material was taken up in MeCN during which time a white solid formed. The solid was filtered, washed with additional MeCN and allowed to air dry to provide the product (4.51 g, 85% yield). LCMS (ESI) *m/z* [M + Na] calcd for C₅₉H₈₆N₂O₁₇: 1117.58; found 1117.6.

Monomers 4. 32(*R*)-ethoxy 40(*R*)-*O*-(4-nitrophenyl)carbonate rapamycin.*Step 1: Synthesis of 32(R)-ethoxy-28,40-bistriethylsilyl rapamycin*

[00722] A solution of 32(*R*)-hydroxy-28,40-bistriethylsilyl rapamycin (773 mg, 0.675 mmol, 1.0 equiv) in chloroform (19 mL) was treated with *N,N,N',N'*-tetramethyl-1,8-naphthalenediamine (1.85 g, 8.63 mmol, 12.8 equiv) along with freshly dried 4Å molecular sieves. The mixture was stirred for 1 h at room temperature and treated with triethyloxonium tetrafluoroborate (1.51 g, 7.95 mmol, 11.8 equiv) in one portion at room temperature. The reaction mixture was stirred for 3 h, at which point the reaction mixture was diluted with DCM and filtered through Celite, washing the filter pad with additional DCM. The combined filtrates were washed twice with 1 M HCl, once with saturated NaHCO_3 solution, and dried over Na_2SO_4 . The solution was filtered and concentrated to a residue. The crude residue was treated with MTBE and filtered to remove polar insoluble material. The filtrate was concentrated and purified by silica gel chromatography (5→25% EtOAc/hex) to afford the product as a foam (516 mg, 65% yield). LCMS (ESI) m/z : $[\text{M} + \text{Na}]$ calcd for $\text{C}_{65}\text{H}_{113}\text{NO}_{13}\text{Si}_2$ 1194.77; found 1194.6.


Step 2: Synthesis of 32(R)-ethoxy rapamycin

[00723] To a solution of 32(*R*)-ethoxyethoxy-28,40-bistriethylsilyl rapamycin (131 mg, 0.112 mmol, 1.0 equiv) in THF (1.3 mL) at 0 °C was added pyridine (271 μ L, 3.35 mmol, 3.4 equiv) followed by 70% HF-pyridine (51 μ L, 1.8 mmol, 1.8 equiv). The reaction flask was capped and stored in the fridge for 3 days, at which point the reaction mixture was poured into cold saturated NaHCO₃ (20 mL). The aqueous layer extracted with EtOAc (3 x 20 mL) and the combined organic layers were washed with 1 M HCl (2 x 20 mL), saturated NaHCO₃ solution (20 mL), and brine. The solution was dried over Na₂SO₄, filtered, and concentrated. The residue was taken up in MeOH (1.5 mL) and added dropwise to H₂O (20 mL). The solids were filtered and washed with additional H₂O to provide the product (53 mg, 51% yield) as a white powder. LCMS (ESI) *m/z*: [M + Na] calcd for C₅₃H₈₅NO₁₃: 966.59; found 966.5.

Step 3: Synthesis of 32(R)-ethoxy 40(R)-O-(4-nitrophenyl)carbonate rapamycin

[00724] To a 0.03 M solution of 32(*R*)-ethoxy rapamycin (1.0 equiv) in DCM is added powdered 4Å molecular sieves. The mixture is stirred at room temperature for 1 h and then pyridine (10 equiv) is added. The mixture is cooled to -10 °C and 4-nitrophenylchloroformate (1.0 equiv) is added in one portion. The reaction is warmed to room temperature and stirred until consumption of 32(*R*)-ethoxy rapamycin, as determined by LCMS analysis. The mixture is filtered through a pad of celite and the pad washed with DCM. The filtrate is washed with H₂O and sat NaHCO₃. The organic phase is then dried over Na₂SO₄, filtered and concentrated under reduced pressure. The crude material is purified by flash chromatography (20→50% EtOAc/hex) to provide the product.


Monomer 5. 40(*R*)-O-(4-nitrophenyl)thiocarbonate rapamycin.



[00725] To a solution of rapamycin (4.00 g, 4.38 mmol, 1.0 equiv) in DCM (20 mL) at -78 °C was added pyridine (4.0 mL, 49 mmol, 11.2 equiv), followed by a solution of *O*-(4-nitrophenyl)chlorothiocarbonate (1.19 g, 5.47 mmol, 1.3 equiv) in DCM (8.0 mL). The reaction mixture was warmed to -20 °C and stirred for 48 h. Hexane (40 mL) was then added

and the resulting suspension was purified by silica gel chromatography (15/25/60 EtOAc/DCM/hexane then 20/25/55 EtOAc/DCM/hexane) to provide the product (3.09 g, 64.4% yield) as an off-white solid. LCMS (ESI) m/z : $[M + Na]$ calcd for $C_{58}H_{82}N_2O_{16}S$: 1117.53; found 1117.5.

Monomer 6. 28-*O*-(4-nitrophenyl)carbonate rapamycin.



*Step 1: Synthesis of 40-*O*-tert-butyldimethylsilyl rapamycin*

[00726] To a solution of rapamycin (1.00 g, 1.09 mmol, 1.0 equiv) in DMF (4 mL) at room temperature was added imidazole (0.22 g, 3.2 mmol, 2.9 equiv) followed by *tert*-butyldimethylsilyl chloride (0.176 g, 1.17 mmol, 1.07 equiv). The reaction mixture was stirred for 18 h. The reaction mixture was then diluted with DCM (100 mL) and washed with 20% aq. LiCl (3 x 20 mL). The organic layer was dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The residue was purified by silica gel chromatography (20→40% EtOAc/hexanes) to give the product (950 mg, 75% yield) as a faint yellow glass. LCMS (ESI) m/z : $[M + H_2O]$ calcd for $C_{57}H_{93}NO_{13}Si$: 1045.65; found 1045.9.

*Step 2: Synthesis of 28-*O*-(4-nitrophenoxycarbonyl)-40-*O*-(tert-butyldimethylsilyl) rapamycin*


[00727] To a solution of 40-*O*-*tert*-butyldimethylsilyl rapamycin (0.845 g, 0.822 mmol, 1.0 equiv) in DCM (10 mL) at room temperature was added pyridine (0.9 mL, 10 mmol, 12.1 equiv) followed by 4-nitrophenyl chloroformate (0.373 g, 1.85 mmol, 2.25 equiv). The reaction mixture was stirred for 2 h. The reaction mixture was then diluted with DCM (150

mL) and the solution sequentially washed with sat. NaHCO_3 (20 mL), 10% citric acid (2 x 20 mL), and brine (20 mL). The organic layer was dried over Na_2SO_4 , filtered, and concentrated under reduced pressure. The residue was purified by reverse phase chromatography (30 $\rightarrow$ 100% MeCN/ H_2O) to give the product (930 mg, 95% yield) as a pale yellow foam. LCMS (ESI) m/z : $[\text{M} + \text{H}]$ calcd for $\text{C}_{64}\text{H}_{96}\text{N}_2\text{O}_{17}\text{Si}$: 1193.66; found 1193.7.

Step 3: Synthesis of 28-O-(4-nitrophenoxy carbonyl) rapamycin

[00728] To a solution of 28-*O*-(4-nitrophenoxy carbonyl)-40-*O*-(*tert*-butyldimethylsilyl)rapamycin (0.930 g, 0.779 mmol, 1.0 equiv) in THF (10.7 mL) was added pyridine (3.78 mL, 46.8 mmol, 60.1 equiv) followed by the dropwise addition of 70% HF-pyridine (0.91 mL, 31.2 mmol, 40.0 equiv). The reaction mixture was stirred at room temperature for 48 h. The mixture was then poured slowly into ice cold sat. aqueous NaHCO_3 (20 mL). The aqueous layer was extracted with EtOAc (3 x 20 mL) and the combined organic layer was washed with sat. NaHCO_3 (10 mL) and brine (10 mL), dried over Na_2SO_4 , filtered, and concentrated under reduced pressure. The residue was purified by reverse phase chromatography (30 $\rightarrow$ 100% MeCN/ H_2O) to give the product (200 mg, 24% yield) as a faint yellow powder. LCMS (ESI) m/z : $[\text{M} + \text{Na}]$ calcd for $\text{C}_{58}\text{H}_{82}\text{N}_2\text{O}_{17}$: 1101.55; found 1101.3.


Monomer 7. 32(*R*)-hydroxy 40(*R*)-*O*-(4-nitrophenyl)thiocarbonate rapamycin.



[00729] To a solution of 32(*R*)-hydroxy rapamycin (2.80 g, 3.06 mmol, 1.0 equiv) in DCM (28 mL) was added pyridine (27.6 mL, 34 mmol, 11 equiv) and dried 4Å molecular sieves (2.8 g). The suspension was stirred at room temperature for 1 h, at which point the mixture was cooled to -25 °C and a solution of *O*-(4-nitrophenyl)chloroformate (0.798 g, 3.67 mmol, 1.2 equiv) in DCM (6 mL) was added. The reaction was warmed to room temperature and after 21 h was filtered through Celite. The filtrate was partitioned between DCM and H_2O and the aqueous layer was extracted with DCM. The combined organic layers were dried and concentrated under reduced pressure. Purification by silica gel chromatography


(EtOAc/hexanes) afforded the desired product as a white solid (2.15 g, 64% yield). LCMS (ESI) m/z : $[M + Na]$ calcd for $C_{58}H_{84}N_2O_{16}S$: 1119.54; found 1120.0.

Monomer 8. 32(*R*)-methoxy 40(*R*)-*O*-(4-nitrophenyl)thiocarbonate rapamycin.



[00730] To a solution of 32(*R*)-methoxy rapamycin (6.69 g, 7.19 mmol, 1.0 equiv) in DCM (67 mL) was added pyridine (6.6 mL, 81 mmol, 11 equiv) and dried 4 Å molecular sieves (6.7 g). The suspension was stirred at room temperature for 1 h, at which point the mixture was cooled to -25 °C and a solution of *O*-(4-nitrophenyl)chlorothioformate (1.88 g, 8.63 mmol, 1.20 equiv) in DCM (13 mL) was added. The reaction was warmed to room temperature and after 21 h was filtered through Celite. The filtrate was partitioned between DCM and H₂O and the aqueous layer was extracted with DCM. The combined organic layers were dried and concentrated under reduced pressure. Purification by silica gel chromatography (EtOAc/hexanes) afforded the desired product as a white solid (5.1 g, 64% yield). LCMS (ESI) m/z : $[M + Na]$ calcd for $C_{59}H_{86}N_2O_{16}S$: 1133.56; found 1134.1.

Monomer 9. 32-deoxy 40(*R*)-*O*-(4-nitrophenyl)carbonate rapamycin.




[00731] To a solution of 32-deoxy rapamycin (0.623 g, 0.692 mmol, 1.0 equiv) in DCM (24.7 mL) was added 4 Å molecular sieves (600 mg). The suspension was stirred for 1 h and then pyridine (557 μL, 6.92 mmol, 10 equiv) was added. The reaction mixture was cooled to 0 °C and then *O*-(4-nitrophenyl)chloroformate (175 mg, 1.03 mmol, 1.7 equiv) was added. The reaction warmed to room temperature and stirred for 2 h, at which point the reaction mixture was concentrated under reduced pressure. Purification by silica gel chromatography


(0→10% MeOH/DCM) afforded the desired product as a white solid (0.61 g, 82% yield).

LCMS (ESI) m/z : $[M + Na]$ calcd for $C_{58}H_{84}N_2O_{16}$: 1087.57; found 1087.6.

Monomer 10. 32-deoxy 40(R)-O-(4-nitrophenyl)thiocarbonate rapamycin.



[00732] To a 0.2 M solution of 32-deoxy rapamycin (1.0 equiv) in DCM at -78 °C is added pyridine (11.2 equiv), followed by a 0.7 M solution of *O*-(4-nitrophenyl)chlorothiocarbonate (1.3 equiv) in DCM. The reaction mixture is warmed to -20 °C and stirred until consumption of the starting material, as determined by LCMS analysis. Hexane is then added and the resulting suspension is purified by silica gel chromatography to provide the product.

Monomer 11. 28(R)-methoxy 32(R)-hydroxy 40(R)-(4-nitrophenyl)carbonate rapamycin.*Step 1: Synthesis of 28(R)-methoxy 40(R)-O-tert-butyldimethylsilyl rapamycin*

[00733] To a solution of 40(R)-O-*tert*-butyldimethylsilyl rapamycin (4.00 g, 4.89 mmol, 1.0 equiv) in chloroform (67 mL) was added proton sponge (11.2 mL, 52.3 mmol, 13 equiv) and dried 4 Å molecular sieves (5.8 g). The suspension was stirred at room temperature for 1 h, at which point trimethyloxonium tetrafluoroborate (7.21 g, 48.8 mmol, 12.5 equiv) was added. After 4 h the suspension was filtered through Celite. The filtrate was washed sequentially with aqueous 2 N HCl, H₂O, sat. aqueous NaHCO₃, then dried and concentrated under reduced pressure. Purification by silica gel chromatography (EtOAc/hexane) afforded the desired product as a white solid (2.1 g, 52% yield). LCMS (ESI) *m/z*: [M + Na] calcd for C₅₈H₉₅NO₁₃Si: 1064.65; found 1065.26.

Step 2: Synthesis of 28(R)-methoxy 32(R)-hydroxy 40(R)-O-tert-butyldimethylsilyl rapamycin

[00734] To a solution of 28(*R*)-methoxy 40(*R*)-*O*-*tert*-butyldimethylsilyl rapamycin (2.13 g, 2.04 mmol, 1.0 equiv) in THF (31 mL) at -20 °C was added a solution of lithium tri-*tert*-butoxyaluminum hydride in THF (1 M, 4.09 mL, 4.09 mmol, 2.0 equiv), dropwise. The reaction mixture was warmed to room temperature and after 3 h was added to a solution of H₂O (4 mL), EtOAc (31 mL), and 2M aqueous citric acid (4 mL) at 0 °C. After 5 min the mixture was partitioned, and the aqueous layer was extracted with EtOAc. The combined organic layers were poured into a sat. aqueous NaHCO₃ solution (60 mL) at 0 °C. The layers were partitioned, and the organic layer was dried and concentrated under reduced pressure to provide a crude white solid (2.32 g). The crude solid was dissolved in DCM (12 mL) and then pyridine (241 µL, 2.98 mmol, 1.5 equiv), dried 4 Å molecular sieves (2.1 g), and cupric acetate (0.27 g, 1.49 mmol, 0.7 equiv) were added. The suspension was stirred at room temperature for 1 h. The suspension was sparged with O₂ and then kept under an O₂ atmosphere for 30 min. After 2 h the mixture was filtered through Celite and the filtrate was concentrated under reduced pressure. Purification by silica gel chromatography (EtOAc/hexane) afforded the desired product as a white solid (307 mg, 14% yield). LCMS (ESI) *m/z*: [M + Na] calcd for C₅₈H₉₇NO₁₃Si: 1066.66; found 1067.0.

Step 3: Synthesis of 28(R)-methoxy 32(R)-hydroxy rapamycin


[00735] To a solution of 28(*R*)-methoxy 32(*R*)-hydroxy 40(*R*)-*O*-*tert*-butyldimethylsilyl rapamycin (0.307 g, 0.294 mmol, 1.0 equiv) in THF (4 mL) in a polypropylene vial at 0 °C was added pyridine (1.42 mL, 17.6 mmol, 60.0 equiv) followed by 70% HF-pyridine (0.34 mL, 11.7 mmol, 40 equiv). The solution was warmed to room temperature and stirred for 21 h, at which point the solution was poured into sat. aqueous NaHCO₃ at 0 °C. The aqueous layer was extracted with EtOAc (2x) and the combined organic layers were washed with sat. aqueous NaHCO₃ and brine, then dried and concentrated under reduced pressure. Purification by silica gel chromatography (EtOAc/hexane) afforded the desired product as a white solid (146 mg, 53% yield). LCMS (ESI) *m/z*: [M + Na] calcd for C₅₂H₈₃NO₁₃: 952.58; found 952.8.

Step 4: Synthesis of 28(R)-methoxy 32(R)-hydroxy 40(R)-(4-nitrophenyl)carbonate rapamycin

[00736] To a solution of 28(*R*)-methoxy 32(*R*)-hydroxy rapamycin (0.66 g, 0.71 mmol, 1.0 equiv) in DCM (3 mL) was added pyridine (0.64 mL, 7.9 mmol, 11 equiv) and dried 4 Å molecular sieves (0.66 g). The suspension was stirred at room temperature for 1 h, at which

point the mixture was cooled to $-35\text{ }^{\circ}\text{C}$ and *O*-(4-nitrophenyl)chloroformate (0.17 g, 0.85 mmol, 1.2 equiv) was added. After 3 h, DCM (5 mL) was added and the suspension was filtered through Celite. The filtrate was washed with H_2O , dried, and concentrated under reduced pressure. Purification by silica gel chromatography (EtOAc/hexanes) afforded the desired product as a white solid (0.44 g, 57% yield). LCMS (ESI) m/z : $[\text{M} + \text{Na}]$ calcd for $\text{C}_{59}\text{H}_{86}\text{N}_2\text{O}_{17}$: 1117.58; found 1118.0.

Monomer 12. 28(*R*)-methoxy 32(*R*)-methoxy 40(*R*)-(4-nitrophenyl)carbonate rapamycin.



Step 1: Synthesis of 28(*R*)-methoxy 32(*R*)-methoxy 40(*R*)-*O*-*tert*-butyldimethylsilyl rapamycin


[00737] To a solution of 28(*R*)-methoxy 32(*R*)-hydroxy 40(*R*)-*O*-*tert*-butyldimethylsilyl rapamycin (1.15 g, 1.10 mmol, 1.0 equiv) in chloroform (19 mL) was added proton sponge (3.22 mL, 15.0 mmol, 14 equiv) and dried 4Å molecular sieves (2.3 g). The suspension was stirred at room temperature for 1 h, at which point trimethyloxonium tetrafluoroborate (2.07 g, 14.0 mmol, 12.7 equiv) was added. After 4 h the suspension was filtered through Celite and the filtrate was washed with 1N HCl, H_2O , and sat. aqueous NaHCO_3 , dried and concentrated under reduced pressure. Purification by silica gel chromatography (EtOAc/hexane) afforded the desired product as a white solid. LCMS (ESI) m/z : $[\text{M} + \text{Na}]$ calcd for $\text{C}_{59}\text{H}_{99}\text{NO}_{13}\text{Si}$: 1080.68; found 1081.2.

Step 2: Synthesis of 28(*R*)-methoxy 32(*R*)-methoxy rapamycin

[00738] To a solution of 28(*R*)-methoxy 32(*R*)-methoxy 40(*R*)-*O*-*tert*-butyldimethylsilyl rapamycin in THF (4 mL) in a polypropylene vial at 0 °C was added pyridine (1.13 mL, 14.2 mmol, 12.9 equiv) followed by 70% HF-pyridine (0.27 mL, 9.42 mmol, 8.6 equiv). The solution was warmed to room temperature and stirred for 41 h, at which point the solution was poured into sat. aqueous NaHCO₃ at 0 °C. The aqueous layer was extracted with EtOAc (2x) and the combined organic layers were washed with sat. aqueous NaHCO₃ and brine, then dried and concentrated under reduced pressure. Purification by silica gel chromatography (EtOAc/hexane) afforded the desired product as a white solid (516 mg, 49% yield 2-steps). LCMS (ESI) *m/z*: [M + Na] calcd for C₅₃H₈₅NO₁₃: 966.59; found 967.0.

Step 3: Synthesis of 28(R)-methoxy 32(R)-methoxy 40(R)-(4-nitrophenyl)carbonate rapamycin

[00739] To a solution of 28(*R*)-methoxy 32(*R*)-methoxy rapamycin (0.30 g, 0.32 mmol, 1.0 equiv) in DCM (1.4 mL) was added pyridine (0.29 mL, 3.5 mmol, 11 equiv) and dried 4 Å molecular sieves (0.30 g). The suspension was stirred at room temperature for 1 h, at which point it was cooled to -35 °C and *O*-(4-nitrophenyl)chloroformate (0.08 g, 0.38 mmol, 1.2 equiv) was added. After 3 h, DCM (2 mL) was added and the suspension was filtered through Celite. The filtrate was washed with H₂O, dried and concentrated under reduced pressure. Purification by silica gel chromatography (EtOAc/hexanes) afforded the desired product as an off-white solid (0.20 g, 57% yield). LCMS (ESI) *m/z*: [M + Na] calcd for C₆₀H₈₈N₂O₁₇: 1131.60; found 1132.1.

Monomer 13. 32(R)-(4-nitrophenyl)carbonate rapamycin.*Step 1: Synthesis of 28,40-O-bis(triethylsilyl) 32(R)-(4-nitrophenyl)carbonate rapamycin*


[00740] To a solution of 28,40-*O*-bis(triethylsilyl) 32(R)-hydroxy rapamycin (0.602 g, 0.526 mmol, 1.0 equiv) in DCM (16 mL) at -20 °C was added pyridine (0.82 mL, 10 mmol, 19 equiv) followed by *O*-(4-nitrophenyl)chloroformate (0.36 g, 1.8 mmol, 3.4 equiv). The reaction mixture was warmed to room temperature and stirred for 1 h, at which point the solution was diluted with DCM (50 mL) and poured into H₂O (30 mL). The aqueous layer was extracted with DCM (50 mL) and the combined organic layers were washed with brine (20 mL), dried and concentrated under reduced pressure to afford a faint yellow foam that was used directly in the next step.

Step 2: Synthesis of 32(R)-(4-nitrophenyl)carbonate rapamycin

[00741] To a solution of 28,40-*O*-bis(triethylsilyl) 32(R)-(4-nitrophenyl)carbonate rapamycin in THF (10 mL) in a polypropylene vial at 0 °C was added pyridine (1.70 mL, 21.0 mmol, 40.0 equiv) followed by 70% HF-pyridine (0.46 mL, 15.8 mmol, 30.0 equiv). The solution was warmed to room temperature and stirred overnight, at which point the solution was poured into sat. aqueous NaHCO₃ at 0 °C. The aqueous layer was extracted with

EtOAc (3x) and the combined organic layers were washed with sat. aqueous NaHCO₃ and brine, then dried and concentrated under reduced pressure. Purification by reverse phase chromatography (20→100% MeCN/H₂O) afforded the desired product as an off-white powder (420 mg, 74% yield 2-steps). LCMS (ESI) *m/z*: [M + Na] calcd for C₅₈H₈₄N₂O₁₇: 1103.57; found 1104.0.

Monomer 14. 32(*S*)-azido 40-(4-nitrophenyl)carbonate rapamycin.



Step 1: Synthesis of 28,40-O-bis(triethylsilyl) 32(R)-O-methanesulfonyl rapamycin

[00742] To a solution of 28,40-*O*-bis(triethylsilyl) 32(*R*)-hydroxy rapamycin (2.50 g, 2.18 mmol, 1.0 equiv) in DCM (25 mL) at 0 °C was added Et₃N (0.912 mL, 6.54 mmol, 3.0 equiv) followed by methanesulfonyl chloride (0.338 mL, 4.36 mmol, 2.0 equiv). The solution was stirred at 0 °C for 3 h, at which point the EtOAc was added and the solution was washed with sat. aqueous NaHCO₃. The combined organic layers were washed with brine, dried and

concentrated under reduced pressure to give a yellow oil which was used directly in the next step.

Step 2: Synthesis of 28-O-triethylsilyl 32(S)-azido rapamycin


[00743] To a solution of 28,40-*O*-bis(triethylsilyl) 32(*R*)-*O*-methanesulfonyl rapamycin in THF (40 mL) was added DIPEA (0.761 mL, 4.37 mmol, 2.0 equiv) and tetrabutylammonium azide (3.72 g, 13.1 mmol, 6.0 equiv). The reaction solution heated to reflux for 5.5 h and then cooled to room temperature. The solution was diluted with EtOAc and washed with sat. aqueous NaHCO₃. The combined organic layers were washed with brine, dried and concentrated under reduced pressure. Purification by reverse phase chromatography (30→100% MeCN/H₂O) afforded the desired product as a clear glass (746 mg, 33% yield 2-steps). LCMS (ESI) *m/z*: [M + Na] calcd for C₅₇H₉₄N₄O₁₂Si: 1077.65; found 1077.8.

Step 3: Synthesis of 28-O-triethylsilyl 32(S)-azido 40(R)-(4-nitrophenyl)carbonate rapamycin

[00744] To a solution of 28-*O*-triethylsilyl 32(*S*)-azido rapamycin (0.505 g, 0.478 mmol, 1.0 equiv) in DCM (15 mL) was added pyridine (0.75 mL, 9.3 mmol, 19 equiv) and 4 Å molecular sieves. The suspension was cooled to -20 °C and *O*-(4-nitrophenyl)chloroformate (0.32 g, 1.6 mmol, 3.4 equiv) was added. The suspension was stirred at -20 °C for 2 h, at which point the it was diluted with DCM (50 mL), filtered and poured into H₂O (20 mL). The aqueous layer was extracted with DCM (50 mL) and the combined organic layers were washed with brine (20 mL), dried and concentrated under reduced pressure to give a pale-yellow foam which was used directly in the next step.

Step 4: Synthesis of 32(S)-azido 40-O-(4-nitrophenyl)carbonate rapamycin

[00745] To a solution of 28-*O*-triethylsilyl 32(*S*)-azido 40(*R*)-(4-nitrophenyl)carbonate rapamycin in THF (10 mL) in a polypropylene vial at 0 °C was added pyridine (1.55 mL, 19.1 mmol, 40.0 equiv) followed by 70% HF-pyridine (0.42 mL, 14.4 mmol; 30.0 equiv). The solution was warmed to room temperature and stirred overnight, at which point the solution was poured into sat. aqueous NaHCO₃ at 0 °C. The aqueous layer was extracted with EtOAc (3x) and the combined organic layers were washed with sat. aqueous NaHCO₃ and brine, then dried and concentrated under reduced pressure. Purification by reverse phase chromatography (30→100% MeCN/H₂O) afforded the desired product as an off-white powder (410 mg, 77% yield 2-steps). LCMS (ESI) *m/z*: [M + Na] calcd for C₅₈H₈₃N₅O₁₆: 1128.57; found 1129.0.

Monomer 15. 28-methoxy-40-*O*-(4-nitrophenoxy)carbonyl rapamycin.*Step 1: Synthesis of 28-methoxy rapamycin*


[00746] To a solution of 28-methoxy-40-*O*-(*tert*-butyldimethyl)silyl rapamycin (0.500 g, 0.480 mmol, 1.0 equiv) in MeOH (1.6 mL) at -20 °C was added H₂SO₄ (1.28 μL, 0.024 mmol, 0.05 equiv). The reaction mixture was stirred at -20 °C for 48 h. The reaction mixture was then poured into sat. aqueous NaHCO₃ (4 mL)/H₂O (4 mL). The aqueous layer was extracted with MTBE (2 x 6 mL), and the combined organic phases were dried, filtered, and concentrated under reduced pressure. Purification by reverse phase chromatography (30→100% MeCN/H₂O) afforded the desired product as a yellow powder (270 mg, 61% yield). LCMS (ESI) *m/z*: [M + Na] calcd for C₅₂H₈₁NO₁₃: 950.5; found 950.7.

*Step 2: Synthesis of 28-methoxy-40-*O*-(4-nitrophenoxy)carbonyl rapamycin*

[00747] To a solution of 28-methoxy rapamycin (0.210 g, 0.226 mmol, 1.0 equiv) in DCM (7.1 mL) at -20 °C was added pyridine (0.35 mL, 4.4 mmol, 19 equiv) and then *p*-nitrophenyl chloroformate (0.15 g, 0.76 mmol, 3.4 equiv). The reaction mixture was stirred at -20 °C for 30 min and then warmed to room temperature. After stirring overnight, *p*-nitrophenyl chloroformate (0.15 g, 0.76 mmol, 3.4 equiv) was added and the reaction was stirred for an additional 2 h. The reaction mixture was diluted with DCM (20 mL) and poured into H₂O (10 mL). The aqueous layer was extracted with DCM (20 mL), and the combined organic layers were washed with brine (9 mL), dried, filtered and concentrated under reduced pressure. Purification by reverse phase chromatography (50→100% MeCN/H₂O) afforded the desired


product as a pale yellow powder (200 mg, 81% yield). LCMS (ESI) m/z : $[M + Na]$ calcd for $C_{59}H_{84}N_2O_{17}$: 1115.6; found 1115.8.

Monomer 16. 32(*R*),40-*O,O*-bis[(4-nitrophenoxy)carbonyl] rapamycin.



[00748] To a solution of 32(*R*)-*O*-[(4-nitrophenoxy)carbonyl] rapamycin (675 mg, 0.624 mmol, 1.0 equiv) in DCM (13 mL) was added powdered 4Å molecular sieves (675 mg). The suspension was stirred for 1 h, at which point pyridine (0.56 mL, 6.90 mmol, 11.1 equiv) was added. The mixture was cooled to -15 °C and then *p*-nitrophenyl chloroformate (132 mg, 0.655 mmol, 1.05 equiv) was added in one portion. The mixture was warmed to 0 °C, stirred for 4 h, and then warmed to room temperature. The reaction mixture was filtered and washed with DCM (25 mL). The filtrate was washed with sat. aqueous $NaHCO_3$ (15 mL), H_2O (15 mL), and brine (10 mL), dried, filtered, and concentrated under reduced pressure. Purification by silica gel chromatography (25→45% EtOAc/hexanes) afforded the desired product as a faint yellow solid (566 mg, 73% yield). LC-MS (ESI) m/z : $[M + Na]$ calcd for $C_{65}H_{87}N_3O_{21}$: 1268.57; found 1269.3.

Monomer 17. 28(*R*),32(*R*),40(*R*)-*O,O,O*-tris[(4-nitrophenoxy)carbonyl] rapamycin.



[00749] To a solution of 32(*R*)-hydroxy rapamycin (1.00 g, 1.09 mmol, 1.0 equiv) in DCM (22 mL) was added powdered 4Å molecular sieves (1.0 g). The suspension was stirred for 45

min, at which point pyridine (0.97 mL, 12.0 mmol, 11.0 equiv) was added. The mixture was cooled to -15 °C and then *p*-nitrophenyl chloroformate (550 mg, 2.73 mmol, 2.5 equiv) was added in one portion. The mixture was warmed to room temperature over 4 h and stirred overnight. The mixture was cooled to 0 °C and additional *p*-nitrophenyl chloroformate (220 mg, 1.09 mmol, 1.0 equiv) was added in one portion. The reaction mixture was stirred for 1 h, warmed to room temperature, and then stirred for 2 h. The mixture was once again cooled to 0 °C and additional *p*-nitrophenyl chloroformate (660 mg, 3.27 mmol, 3.0 equiv) was added. The reaction mixture was stirred for 15 min and then at room temperature for 1 h. The reaction mixture was filtered and washed with DCM (25 mL). The filtrate was washed with sat. aqueous NaHCO₃ (20 mL), H₂O (20 mL), and brine (15 mL), dried, filtered, and concentrated under reduced pressure. Purification by silica gel chromatography (5→15% EtOAc/DCM) afforded the desired product as a faint yellow solid (550 mg, 36% yield). LC-MS (ESI) *m/z*: [M + Na] calcd for C₇₂H₉₀N₄O₂₅: 1433.58; found 1434.3.

General Procedures and Specific Examples.

General Procedure 1: Coupling of a carboxylic acid and an amine followed by *N*-Boc deprotection.




Step 1:

[00750] To a 0.1 M solution of carboxylic acid (1.0 equiv) in DMA was added an amine (1.2 equiv), DIPEA (4.0 equiv) and PyBOP (1.3 equiv). The reaction was allowed to stir until consumption of the carboxylic acid, as indicated by LCMS. The reaction mixture was then purified by silica gel chromatography to afford the product.

Step 2:

[00751] To a 0.07 M solution of *N*-Boc protected amine (1.0 equiv) in dioxane was added HCl (4 M in dioxane) (50 equiv). The reaction was allowed to stir until consumption of *N*-Boc protected amine, as indicated by LCMS. Then the reaction was concentrated to an oil, which was then dissolved in H₂O and lyophilized to afford the product.

Intermediate A1-7. 1-amino-27-(6-{[4-amino-3-(2-amino-1,3-benzoxazol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl]methyl}-1,2,3,4-tetrahydroisoquinolin-2-yl)-3,6,9,12,15,18,21,24-octaoxaheptacosan-27-one


















Step 1: Synthesis of *tert*-butyl N-[27-(6-{[4-amino-3-(2-amino-1,3-benzoxazol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl]methyl}-1,2,3,4-tetrahydroisoquinolin-2-yl)-27-oxo-3,6,9,12,15,18,21,24-octaoxaheptacosan-1-yl]carbamate









[00752] To a solution of 1-{[(*tert*-butoxy)carbonyl]amino}-3,6,9,12,15,18,21,24-octaoxaheptacosan-27-oic acid (102 mg, 189 μ mol, 1.0 equiv) and 6-{[4-amino-3-(2-amino-1,3-benzoxazol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl]methyl}-1,2,3,4-tetrahydroisoquinolin-2-ium (120 mg, 227 μ mol, 1.2 equiv) in DMA (1.88 mL) was added DIPEA (131 μ L, 756 μ mol, 4.0 equiv) followed by PyBOP (127 mg, 245 μ mol, 1.3 equiv). The reaction was stirred at room temperature. After 2 h, the reaction mixture was concentrated under reduced pressure and the crude residue was purified by silica gel chromatography (0 $\rightarrow$ 20% MeOH/DCM) to give the product (161.5 mg, 91% yield) as a pale yellow oil. LCMS (ESI) *m/z*: [M + H] calcd for C₄₆H₆₅N₉O₁₂: 936.49; found 936.3.









Step 2: Synthesis of 1-amino-27-(6-{[4-amino-3-(2-amino-1,3-benzoxazol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl]methyl}-1,2,3,4-tetrahydroisoquinolin-2-yl)-3,6,9,12,15,18,21,24-octaoxaheptacosan-27-one









[00753] To a solution of *tert*-butyl N-[27-(6-{[4-amino-3-(2-amino-1,3-benzoxazol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl]methyl}-1,2,3,4-tetrahydroisoquinolin-2-yl)-27-oxo-3,6,9,12,15,18,21,24-octaoxaheptacosan-1-yl]carbamate (0.9 g, 0.9614 mmol, 1.0 equiv) in dioxane (3.20 mL) was added HCl (4 M in dioxane, 2.40 mL, 9.61 mmol, 10.0 equiv). The reaction stirred for 2 h and then was concentrated under reduced pressure to an oil. The oil was azeotroped with DCM (3 x 15 mL) to provide the product (881 mg, 105% yield, HCl) as a tan solid, which was used directly in the next step. LCMS (ESI) *m/z*: [M + H] calcd for C₄₁H₅₇N₉O₁₀: 836.43; found 836.3.


Intermediate A1-6 	$C_{41}H_{57}N_9O_{10}$	$[M + H] = 836.43$	$[M + H] = 836.3$
Intermediate A1-7 	$C_{37}H_{49}N_9O_8$	$[M + H] = 748.38$	$[M + H] = 748.2$
Intermediate A1-8 	$C_{33}H_{41}N_9O_6$	$[M + H] = 660.33$	$[M + H] = 660.2$
Intermediate A1-9 	$C_{31}H_{37}N_9O_5$	$[M + H] = 616.30$	$[M + H] = 616.3$
Intermediate A1-10 	$C_{29}H_{33}N_9O_4$	$[M + H] = 572.28$	$[M + H] = 572.3$
Intermediate A1-11 	$C_{27}H_{29}N_9O_3$	$[M + H] = 528.25$	$[M + H] = 528.2$
Intermediate A1-12 	$C_{35}H_{55}N_9O_9$	$[M + H] = 746.42$	$[M + H] = 746.4$
Intermediate A1-13 			

 Intermediate A1-14	$C_{31}H_{47}N_9O_7$	$[M + H] = 658.37$	$[M + H] = 658.3$
 Intermediate A1-15	$C_{27}H_{39}N_9O_5$	$[M + H] = 570.32$	$[M + H] = 570.2$
 Intermediate A1-16	$C_{23}H_{31}N_9O_3$	$[M + H] = 482.26$	$[M + H] = 482.3$
 Intermediate A1-17	$C_{21}H_{27}N_9O_2$	$[M + H] = 438.24$	$[M + H] = 438.4$
 Intermediate A1-18	$C_{49}H_{73}N_9O_{13}$	$[M + H] = 996.54$	$[M + H] = 996.4$
 Intermediate A1-19	$C_{45}H_{65}N_9O_{11}$	$[M + H] = 908.49$	$[M + H] = 908.3$
 Intermediate A1-20	$C_{41}H_{57}N_9O_9$	$[M + H] = 820.44$	$[M + H] = 820.3$
 Intermediate A1-21	$C_{37}H_{49}N_9O_7$	$[M + H] = 732.39$	$[M + H] = 732.3$

Intermediate A1-21		$C_{35}H_{45}N_9O_6$	$[M + H] = 688.36$	$[M + H] = 688.3$
Intermediate A1-22		$C_{33}H_{41}N_9O_5$	$[M + H] = 644.33$	$[M + H] = 644.3$
Intermediate A1-23		$C_{31}H_{37}N_9O_4$	$[M + H] = 600.31$	$[M + H] = 600.4$
Intermediate A1-24		$C_{36}H_{55}N_9O_{10}$	$[M + H] = 774.42$	$[M + H] = 774.7$
Intermediate A1-25		$C_{32}H_{47}N_9O_8$	$[M + H] = 686.36$	$[M + H] = 686.4$
Intermediate A1-26		$C_{40}H_{63}N_9O_{11}$	$[M + H] = 846.47$	$[M + H] = 846.8$
Intermediate A1-27		$C_{43}H_{54}F_3N_7O_9$	$[M + H] = 870.40$	$[M + H] = 870.4$
Intermediate A1-28		$C_{52}H_{67}F_3N_{10}O_{11}$	$[M + H] = 1065.50$	$[M + H] = 1065.4$
Intermediate A1-29				







 Intermediate A1-30	$C_{48}H_{59}F_3N_{10}O_9$	$[M + H] = 977.45$	$[M + H] = 977.4$
 Intermediate A1-31	$C_{39}H_{57}F_3N_{12}O_{12}$	$[M + H] = 853.43$	$[M + H] = 853.4$
 Intermediate A1-32	$C_{40}H_{55}N_9O_{10}$	$[M + H] = 822.42$	$[M + H] = 822.2$
 Intermediate A1-33	$C_{39}H_{55}N_9O_{10}$	$[M + H] = 810.42$	$[M + H] = 810.3$
 Intermediate A1-34	$C_{43}H_{63}N_5O_{13}S$	$[M + H] = 890.42$	$[M + H] = 890.3$
 Intermediate A1-35	$C_{39}H_{54}FN_5O_{11}S$	$[M + H] = 820.36$	$[M + H] = 820.3$
 Intermediate A1-36	$C_{43}H_{62}FN_5O_{13}S$	$[M + H] = 908.41$	$[M + H] = 908.3$
 Intermediate A1-37	$C_{47}H_{62}F_3N_7O_{11}$	$[M + H] = 958.46$	$[M + H] = 958.3$

 Intermediate A1-38	$C_{34}H_{43}N_9O_6$	$[M + H] = 674.34$	$[M + H] = 674.3$
 Intermediate A1-39	$C_{41}H_{64}N_8O_{11}$	$[M + H] = 845.48$	$[M + H] = 845.3$
 Intermediate A1-40	$C_{37}H_{56}N_8O_9$	$[M + H] = 757.43$	$[M + H] = 757.3$
 Intermediate A1-41	$C_{27}H_{36}N_8O_4$	$[M + H] = 537.29$	$[M + H] = 537.2$
 Intermediate A1-42	$C_{37}H_{53}N_{11}O_{10}$	$[M + H] = 812.41$	$[M + H] = 812.3$
 Intermediate A1-43	$C_{36}H_{56}N_8O_{10}$	$[M + H] = 761.42$	$[M + H] = 761.3$
 Intermediate A1-44	$C_{43}H_{61}N_9O_{10}$	$[M + H] = 864.46$	$[M + H] = 864.4$
 Intermediate A1-45	$C_{43}H_{62}N_{10}O_{12}S$	$[M + H] = 943.44$	$[M + H] = 943.3$


 Intermediate A1-46	$C_{39}H_{55}N_{11}O_{10}$	$[M + H] = 838.42$	$[M + H] = 838.3$

[00755] Following General Procedure 1, but using the appropriate Intermediate A1 in Table 5 and PEG carboxylic acid, the Intermediates A2 in Table 6 were prepared:


Table 6. Additional amines prepared

Structure	Molecular Formula	Calculated MW	Observed MW
 Intermediate A2-1	$C_{38}H_{60}N_{10}O_{11}$	$[M + H] = 833.45$	$[M + H] = 833.8$
 Intermediate A2-2	$C_{44}H_{62}N_{10}O_{11}$	$[M + H] = 907.47$	$[M + H] = 908.0$
 Intermediate A2-3	$C_{38}H_{60}N_{10}O_{10}$	$[M + H] = 817.46$	$[M + H] = 817.4$
 Intermediate A2-4	$C_{48}H_{70}N_{10}O_{12}$	$[M + H] = 979.53$	$[M + H] = 979.9$
 Intermediate A2-5	$C_{46}H_{66}N_{10}O_{11}$	$[M + H] = 935.50$	$[M + H] = 935.3$
 Intermediate A2-6	$C_{40}H_{61}N_9O_{10}$	$[M + H] = 828.46$	$[M + H] = 828.3$

General Procedure 2: Coupling of a 4-nitrophenyl carbonate containing rapamycin monomer and an active site inhibitor containing intermediate having a primary or

secondary amine.









[00756] To a 0.02 M solution of 4-nitrophenyl carbonate containing rapamycin monomer (1.0 equiv) and an active site inhibitor containing intermediate (2.0 equiv) in DMA was added DIPEA (4.0 equiv). The resulting solution was stirred at room temperature under nitrogen. Upon completion as determined by LCMS analysis, the crude reaction mixture was purified by preparative HPLC to provide the product.







Example 2: Synthesis of Series 1 bivalent rapamycin compound.









[00757] To a solution of 40(*R*)-*O*-(4-nitrophenyl carbonate) rapamycin (25 mg, 23.16 μmol , 1.0 equiv) and Intermediate A1-7 (42.0 mg, 46.32 μmol , 2.0 equiv) in DMA (1.15 mL) was added DIPEA (16.0 μL , 92.64 μmol , 4.0 equiv). The reaction was stirred for 18 h, at which point the reaction mixture purified by reverse phase chromatography (10 $\rightarrow$ 40 $\rightarrow$ 95% MeCN + 0.1% formic acid/H₂O + 0.1% formic acid) to give the product (9.92 mg, 24% yield) as a white solid. LCMS (ESI) *m/z*: [M + H]⁺ calcd for C₉₃H₁₃₄N₁₀O₂₄: 1775.97; found 1775.7.






[00758] Following General Procedure 2, but using the appropriate 4-nitrophenyl carbonate containing rapamycin monomer in Table 1 and Intermediates A1 and A2 from Tables 5 and 6, the Series 1 bivalent analogs in Table 7 were synthesized:









Table 7. Series 1 Bivalent Compounds:










Structure	Molecular Formula	Calculated MW	Observed MW
 Example 1	$C_{87}H_{132}N_{10}O_{24}$	$[M + H] = 1701.95$	$[M + H] = 1701.8$
 Example 2	$C_{93}H_{134}N_{10}O_{24}$	$[M + H] = 1775.97$	$[M + H] = 1775.7$
 Example 3	$C_{89}H_{126}N_{10}O_{22}$	$[M + H] = 1687.91$	$[M + H] = 1687.8$
 Example 4	$C_{87}H_{132}N_{10}O_{23}$	$[M + H] = 1684.95$	$[M + H] = 1685.1$
 Example 5	$C_{101}H_{150}N_{10}O_{27}$	$[M + H] = 1936.08$	$[M + H] = 1936.4$
 Example 6	$C_{97}H_{142}N_{10}O_{25}$	$[M + H] = 1848.02$	$[M + H] = 1848.3$
 Example 7	$C_{93}H_{134}N_{10}O_{23}$	$[M + H] = 1759.97$	$[M + H] = 1760.1$
 Example 8	$C_{87}H_{134}N_{10}O_{24}$	$[M + H] = 1703.97$	$[M + H] = 1703.7$









 Example 9	$C_{93}H_{136}N_{10}O_{24}$	$[M + H] = 1777.98$	$[M + H] = 1777.7$
 Example 10	$C_{93}H_{136}N_{10}O_{23}$	$[M + H] = 1761.99$	$[M + H] = 1761.7$
 Example 11	$C_{88}H_{136}N_{10}O_{24}$	$[M + H] = 1717.98$	$[M + H] = 1718.0$
 Example 12	$C_{94}H_{138}N_{10}O_{24}$	$[M + H] = 1792.00$	$[M + H] = 1791.9$
 Example 13	$C_{91}H_{133}N_{13}O_{24}$	$[M + H] = 1792.97$	$[M + H] = 1793.0$
 Example 14	$C_{90}H_{137}N_{11}O_{25}$	$[M + H] = 1772.99$	$[M + H] = 1772.9$
 Example 15	$C_{96}H_{139}N_{11}O_{25}$	$[M + H] = 1847.00$	$[M + H] = 1847.0$
 Example 16	$C_{96}H_{141}N_{11}O_{25}$	$[M + H] = 1849.02$	$[M + H] = 1848.9$








 Example 17	$C_{97}H_{143}N_{11}O_{25}$	$[M + H] = 1863.04$	$[M + H] = 1863.0$
 Example 18	$C_{90}H_{137}N_{11}O_{24}$	$[M + H] = 1756.99$	$[M + H] = 1756.8$
 Example 19	$C_{88}H_{132}N_{10}O_{24}$	$[M + H] = 1713.95$	$[M + H] = 1713.9$
 Example 20	$C_{92}H_{140}N_{10}O_{25}$	$[M + H] = 1786.01$	$[M + H] = 1786.0$
 Example 21	$C_{100}H_{147}N_{11}O_{26}$	$[M + H] = 1919.06$	$[M + H] = 1919.0$
 Example 22	$C_{93}H_{134}N_{10}O_{23}S$	$[M + H] = 1791.94$	$[M + H] = 1791.8$
 Example 23	$C_{93}H_{134}N_{10}O_{24}$	$[M + H] = 1775.97$	$[M + H] = 1775.9$
 Example 24	$C_{104}H_{144}F_3N_{11}O_2$ 5	$[M + 2H]/2 = 1003.03$	$[M + 2H]/2 = 1003.5$







Example 24 	$C_{104}H_{146}F_3N_{11}O_2$ 5	$[M + 2H]/2 = 1004.03$	$[M + 2H]/2 = 1004.5$
Example 25 	$C_{105}H_{148}F_3N_{11}O_2$ 5	$[M + 2H]/2 = 1011.04$	$[M + 2H]/2 = 1011.5$
Example 26 	$C_{92}H_{132}N_{10}O_{24}$	$[M + H] = 1761.95$	$[M + H] = 1761.8$
Example 71 	$C_{92}H_{134}N_{10}O_{24}$	$[M + H] = 1763.97$	$[M + H] = 1763.8$
Example 72 	$C_{93}H_{136}N_{10}O_{23}$	$[M + H] = 1761.99$	$[M + H] = 1762.0$
Example 73 	$C_{93}H_{135}N_{13}O_{23}$	$[M + H] = 1802.99$	$[M + H] = 1802.9$
Example 74 	$C_{94}H_{138}N_{10}O_{24}$	$[M + H] = 1792.00$	$[M + H] = 1791.7$
Example 75 			

 Example 76	$C_{95}H_{140}N_{10}O_{24}$	$[M + H] = 1806.01$	$[M + H] = 1805.8$
 Example 77	$C_{93}H_{136}N_{10}O_{23}S$	$[M + H] = 1793.96$	$[M + H] = 1793.9$
 Example 78	$C_{94}H_{138}N_{10}O_{23}S$	$[M + H] = 1807.97$	$[M + H] = 1807.9$
 Example 79	$C_{91}H_{132}N_{10}O_{24}$	$[M + H] = 1749.95$	$[M + H] = 1749.9$
 Example 80	$C_{91}H_{134}N_{10}O_{24}$	$[M + H] = 1751.97$	$[M + H] = 1751.9$
 Example 81	$C_{92}H_{136}N_{10}O_{24}$	$[M + H] = 1765.98$	$[M + H] = 1765.8$
 Example 82	$C_{91}H_{133}N_{13}O_{23}$	$[M + H] = 1776.97$	$[M + H] = 1776.9$
 Example 83	$C_{91}H_{133}N_{10}O_{23}S$	$[M + H] = 1766.93$	$[M + H] = 1766.8$
 Example 84	$C_{95}H_{140}N_6O_{27}S$	$[M + H] = 1829.96$	$[M + H] = 1830.0$


 Example 85	$C_{95}H_{142}N_6O_{27}S$	$[M + H] = 1831.97$	$[M + H] = 1831.9$
 Example 86	$C_{96}H_{144}N_6O_{27}S$	$[M + H] = 1845.99$	$[M + H] = 1846.0$
 Example 87	$C_{95}H_{142}N_6O_{26}S$	$[M + H] = 1815.98$	$[M + H] = 1815.8$
 Example 88	$C_{95}H_{141}N_9O_{26}S$	$[M + H] = 1856.98$	$[M + H] = 1856.8$
 Example 89	$C_{95}H_{140}N_6O_{26}S_2$	$[M + H] = 1845.93$	$[M + H] = 1846.0$
 Example 90	$C_{95}H_{142}N_6O_{26}S_2$	$[M + H] = 1847.95$	$[M + H] = 1847.9$
 Example 91	$C_{96}H_{144}N_6O_{26}S_2$	$[M + H] = 1861.96$	$[M + H] = 1861.7$
 Example 92	$C_{95}H_{139}FN_6O_{27}S$	$[M + H] = 1847.95$	$[M + H] = 1848.0$
 Example 93	$C_{95}H_{141}FN_6O_{27}S$	$[M + H] = 1849.96$	$[M + H] = 1850.0$

 Example 94	$C_{96}H_{143}FN_6O_{27}S$	$[M + H] = 1863.98$	$[M + H] = 1864.0$
 Example 95	$C_{95}H_{141}FN_6O_{26}S$	$[M + H] = 1833.97$	$[M + H] = 1833.9$
 Example 96	$C_{95}H_{139}FN_6O_{26}S$ 2	$[M + H] = 1863.92$	$[M + H] = 1863.8$
 Example 97	$C_{95}H_{141}FN_6O_{26}S$ 2	$[M + H] = 1865.94$	$[M + H] = 1865.8$
 Example 98	$C_{96}H_{143}FN_6O_{26}S$ 2	$[M + H] = 1879.96$	$[M + H] = 1879.9$
 Example 99	$C_{93}H_{136}N_{10}O_{24}$	$[M + H] = 1777.98$	$[M + H] = 1777.8$
 Example 100	$C_{97}H_{144}N_{10}O_{24}$	$[M + H] = 1834.04$	$[M + H] = 1833.8$
 Example 101	$C_{99}H_{141}F_3N_8O_{25}$	$[M + H] = 1900.00$	$[M + H] = 1899.9$

 Example 102	$C_{99}H_{147}N_{11}O_{25}$	$[M + H] = 1891.06$	$[M + H] = 1890.9$
 Example 136	$C_{89}H_{135}N_9O_{23}$	$[M + H] = 1698.97$	$[M + H] = 1698.8$
 Example 137	$C_{93}H_{143}N_9O_{25}$	$[M + H] = 1787.03$	$[M + H] = 1786.7$
 Example 138	$C_{93}H_{141}N_9O_{25}$	$[M + H] = 1785.01$	$[M + H] = 1784.9$
 Example 139	$C_{89}H_{132}N_{12}O_{24}$	$[M + H] = 1753.96$	$[M + H] = 1753.7$
 Example 140	$C_{93}H_{136}N_{10}O_{24}$	$[M + H] = 1777.98$	$[M + H] = 1777.7$
 Example 141	$C_{88}H_{135}N_9O_{24}$	$[M + H] = 1702.97$	$[M + H] = 1702.9$
 Example 142	$C_{95}H_{140}N_{10}O_{24}$	$[M + H] = 1806.01$	$[M + H] = 1805.8$

 Example 143	$C_{97}H_{143}N_{13}O_{24}$	$[M + H] = 1875.04$	$[M + H] = 1874.9$
 Example 144	$C_{95}H_{141}N_{11}O_{26}S$	$[M + H] = 1884.98$	$[M + H] = 1884.9$
 Example 145	$C_{91}H_{134}N_{12}O_{24}$	$[M + H] = 1779.97$	$[M + H] = 1779.8$
 Example 146	$C_{135}H_{191}N_{19}O_{35}$	$[M + 2H]/2 = 1320.19$	$[M + 2H]/2 = 1320.8$
 Example 147	$C_{93}H_{143}N_9O_{24}$	$[M + H] = 1771.03$	$[M + H] = 1770.8$
 Example 148	$C_{92}H_{138}N_{10}O_{24}$	$[M + H] = 1768.00$	$[M + H] = 1767.8$

General Procedure 3: Coupling of a halide containing PEG ester and an amine containing pre-linker followed by ester deprotection.




Step 1:

[00759] To a 0.1 M solution of amine containing pre-linker (1.0 equiv) in MeCN was added K_2CO_3 (2.0 equiv) followed by halide containing PEG ester (1.0 equiv). The reaction was stirred at 80 °C until consumption of amine containing pre-linker, as indicated by LCMS analysis. The reaction was then purified by silica gel chromatography to afford the product.

Step 2:

[00760] To a 0.1 M solution of PEG *tert*-butyl ester (1.0 equiv) in EtOAc was added a solution of HCl in EtOAc. The resulting suspension was stirred at room temperature until consumption of the PEG ester, as indicated by LCMS analysis. The reaction was then concentrated under reduced pressure to afford the product.

Intermediate B1-1. 1-(4-(5-((1,3-dioxoisindolin-2-yl)methyl)pyrimidin-2-yl)piperazin-1-yl)-3,6,9,12-tetraoxapentadecan-15-oic acid




Step 1: Synthesis of *tert*-butyl 1-(4-(5-((1,3-dioxoisindolin-2-yl)methyl)pyrimidin-2-yl)piperazin-1-yl)-3,6,9,12-tetraoxapentadecan-15-oate

[00761] To a mixture of 2-((2-(piperazin-1-yl)pyrimidin-5-yl)methyl)isindoline-1,3-dione (7.97 g, 24.66 mmol, 1.0 equiv) in MeCN (200 mL) was added K_2CO_3 (6.82 g, 49.31 mmol, 2.0 equiv) followed by *tert*-butyl 1-bromo-3,6,9,12-tetraoxapentadecan-15-oate (9.5 g, 24.66 mmol, 1.0 equiv). The reaction mixture was heated to 85 °C and stirred for 15 h. The mixture was then cooled to room temperature and filtered. The filtrate was concentrated under reduced pressure and the residue was purified by silica gel chromatography (0→20% EtOAc/MeOH) to give the product (11.5 g, 74.3% yield) a light yellow liquid.


Step 2: Synthesis of 1-(4-(5-((1,3-dioxoisindolin-2-yl)methyl)pyrimidin-2-yl)piperazin-1-yl)-3,6,9,12-tetraoxapentadecan-15-oic acid

[00762] To a solution of *tert*-butyl 1-(4-(5-((1,3-dioxisoindolin-2-yl)methyl)pyrimidin-2-yl)piperazin-1-yl)-3,6,9,12-tetraoxapentadecan-15-oate (3.5 g, 5.58 mmol, 1.0 equiv) in EtOAc (50 mL) was added a solution of HCl in EtOAc (500 mL). The mixture was stirred at room temperature for 3 h. The mixture was then concentrated under reduced pressure to give the product (5.3 g, 78.2% yield, HCl) as a white solid. LCMS (ESI) *m/z*: [M + H] calcd for C₂₈H₃₇N₅O₈: 572.27; found 572.4.

[00763] Following General Procedure 3, but using the appropriate halide containing PEG and amine containing pre-linkers in Table 4, the Intermediates B1 in Table 8 were prepared:
Table 8. Additional protected amines prepared

Structure	Molecular Formula	Calculated MW	Observed MW
 Intermediate B1-1	C ₂₈ H ₃₇ N ₅ O ₈	[M + H] = 572.27	[M + H] = 572.4

General Procedure 4: Coupling of a PEG carboxylic acid and an amine containing active site inhibitor followed by amine deprotection.



Step 1:


[00764] To a 0.15 M solution of PEG carboxylic acid (1.0 equiv) in DMF was added HATU (1.3 equiv) and DIPEA (5.0 equiv). After stirring for 30 min, the amine containing active site inhibitor (1.2 equiv) was added. The reaction was stirred at room temperature until consumption of PEG carboxylic acid, as indicated by LCMS. The reaction was then purified by reverse phase chromatography to afford the product.

Step 2:

[00765] To a 0.1 M solution of phthalimide protected amine (1.0 equiv) in MeOH at 0 °C was added NH₂NH₂•H₂O (4.0 equiv). The resulting mixture was stirred at 60 °C until

consumption of the phthalimide protected amine, as indicated by LCMS analysis. The reaction was then purified by reverse phase chromatography to afford the product.

Intermediate B2-1. N-(4-(4-amino-3-(2-aminobenzo[d]oxazol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)butyl)-1-(4-(5-(aminomethyl)pyrimidin-2-yl)piperazin-1-yl)-3,6,9,12-tetraoxapentadecan-15-amide



Step 1: Synthesis of N-(4-(4-amino-3-(2-aminobenzo[d]oxazol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)butyl)-1-(4-(5-((1,3-dioxoisindolin-2-yl)methyl)pyrimidin-2-yl)piperazin-1-yl)-3,6,9,12-tetraoxapentadecan-15-amide

[00766] To a mixture of 1-(4-(5-((1,3-dioxoisindolin-2-yl)methyl)pyrimidin-2-yl)piperazin-1-yl)-3,6,9,12-tetraoxapentadecan-15-oic acid (3 g, 4.93 mmol, 1.0 equiv, HCl) in DMF (30 mL) was added HATU (12.11 μ L, 6.41 mmol, 1.3 equiv) and DIPEA (4.30 mL, 24.67 mmol, 5.0 equiv). After 30 min, 5-(4-amino-1-(4-aminobutyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)benzo[d]oxazol-2-amine (4.03 g, 5.92 mmol, 1.2 equiv, 3TFA) was added. The mixture was stirred at room temperature for 3 h. The reaction mixture was then purified by prep-HPLC (MeCN/H₂O) to give the product (5.4 g, 81.2% yield, 4TFA) as a light red solid. LCMS (ESI) *m/z*: [M + 2H]²⁺ calcd for C₄₄H₅₃N₁₃O₈: 446.71; found 447.0.




Step 2: Synthesis of N-(4-(4-amino-3-(2-aminobenzo[d]oxazol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)butyl)-1-(4-(5-(aminomethyl)pyrimidin-2-yl)piperazin-1-yl)-3,6,9,12-tetraoxapentadecan-15-amide

[00767] To a mixture of N-(4-(4-amino-3-(2-aminobenzo[d]oxazol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)butyl)-1-(4-(5-((1,3-dioxoisindolin-2-yl)methyl)pyrimidin-2-yl)piperazin-1-yl)-3,6,9,12-tetraoxapentadecan-15-amide (4 g, 2.97 mmol, 1.0 equiv, 4TFA) in MeOH (25 mL) at 0 °C was added NH₂NH₂•H₂O (588.63 μ L, 11.87 mmol, 4.0 equiv). The mixture was stirred at 60 °C for 2 h. The mixture was then cooled to room temperature and filtered, and


the filter cake was washed with MeOH (5 mL). The filtrate was concentrated under reduced pressure and the residue was purified by prep-HPLC (MeCN/H₂O) to give the product (700 mg, 24.5% yield, TFA) as a white solid. LCMS (ESI) m/z : $[M + 2H]/2$ calcd for C₃₆H₅₁N₁₃O₆: 381.71; found 381.8.

[00768] Following General Procedure 4, but using the appropriate Intermediate B1 in Table 8 and amine containing active site inhibitors in Table 2, the Intermediates B2 in Table 9 were prepared:

Table 9. Additional amines prepared


Structure	Molecular Formula	Calculated MW	Observed MW
 Intermediate B2-1	C ₃₆ H ₅₁ N ₁₃ O ₆	$[M + 2H]/2 = 381.71$	$[M + 2H]/2 = 381.8$
 Intermediate B2-2	C ₄₂ H ₅₃ N ₁₃ O ₆	$[M + H] = 836.43$	$[M + H] = 836.4$
 Intermediate B2-3	C ₃₆ H ₅₁ N ₁₃ O ₅	$[M + 2H]/2 = 373.72$	$[M + 2H]/2 = 373.7$

General Procedure 5: Coupling of a halide containing PEG carboxylic acid and an amine containing active site inhibitor.



[00769] To a 0.1 M solution of amine containing active site inhibitor (1.0 equiv) and PEG containing carboxylic acid (1.2 equiv) in DMA was added DIPEA (4.0 equiv) followed by PyBOP (1.3 equiv). The reaction was stirred until consumption of amine containing active site inhibitor, as indicated by LCMS. The reaction was then purified by reverse phase HPLC to afford the product.

Intermediate B3-1. 18-{6-[(4-amino-3-{1H-pyrrolo[2,3-b]pyridin-5-yl}-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl]-1,2,3,4-tetrahydroisoquinolin-2-yl}-1-bromo-3,6,9,12,15-pentaoxaoctadecan-18-one




[00770] To a solution of 1-bromo-3,6,9,12,15-pentaoxaoctadecan-18-oic acid (105 mg, 282 μmol , 1.2 equiv) and 3-{1H-pyrrolo[2,3-b]pyridin-5-yl}-1-[(1,2,3,4-tetrahydroisoquinolin-6-yl)methyl]-1H-pyrazolo[3,4-d]pyrimidin-4-amine (120 mg, 235 μmol , 1.0 equiv) in DMA (2.34 mL) was added DIPEA (163 μL , 940 μmol , 4.0 equiv) followed by PyBOP (158 mg, 305 μmol , 1.3 equiv). The resulting solution was stirred at room temperature for 3 h then purified by reverse phase HPLC (10 $\rightarrow$ 98% MeCN + 0.1% formic acid/H₂O + 0.1% formic acid) to afford the product (82.7 mg, 47% yield). LCMS (ESI) m/z : [M + H] calcd for C₃₅H₄₃BrN₈O₆: 751.26; found 751.2.

[00771] Following General Procedure 5, but using the appropriate halide containing PEG carboxylic acid and amine containing active site inhibitors in Table 2, the Intermediates B3 in Table 10 were prepared:

Table 10. Additional PEG halides prepared

Structure	Molecular Formula	Calculated MW	Observed MW
Intermediate B3-1	C ₃₅ H ₄₃ BrN ₈ O ₆	[M + H] = 751.26	[M + H] = 751.2
Intermediate B3-2	C ₂₇ H ₂₇ BrN ₈ O ₃	[M + H] = 591.15	[M + H] = 591.2

General Procedure 6: Displacement of a PEG halide with an amine containing post linker and deprotection of the amine.




Step 1:

[00772] To a 0.1 M solution of halide containing PEG (1.0 equiv) in MeCN was added K_2CO_3 (3.0 equiv) followed by amine containing post linker (1.2 equiv). The resulting suspension was heated to 80 °C and stirred until consumption of the PEG halide, as indicated by LCMS analysis. The reaction was cooled to room temperature and then purified by silica gel chromatography to afford the product.

Step 2:

[00773] To a 0.07 M solution of *N*-Boc protected amine (1.0 equiv) in dioxane was added HCl (4 M in dioxane, 10.0 equiv). The reaction was stirred until consumption of *N*-Boc protected amine, as indicated by LCMS analysis. The reaction was then concentrated under reduced pressure to afford the product.

Intermediate B2-4. 18-{6-[(4-amino-3-{1H-pyrrolo[2,3-b]pyridin-5-yl}-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl]-1,2,3,4-tetrahydroisoquinolin-2-yl}-1-(4-{5H,6H,7H,8H-pyrido[4,3-d]pyrimidin-2-yl}piperazin-1-yl)-3,6,9,12,15-pentaoxaoctadecan-18-one



Step 1: Synthesis of *tert*-butyl 2-[4-(18-{6-[(4-amino-3-{1H-pyrrolo[2,3-b]pyridin-5-yl}-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl]-1,2,3,4-tetrahydroisoquinolin-2-yl}-18-oxo-3,6,9,12,15-pentaoxaoctadecan-1-yl)piperazin-1-yl]-5H,6H,7H,8H-pyrido[4,3-d]pyrimidine-6-carboxylate


[00774] To a suspension of 18-{6-[(4-amino-3-{1H-pyrrolo[2,3-b]pyridin-5-yl}-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl]-1,2,3,4-tetrahydroisoquinolin-2-yl}-1-bromo-3,6,9,12,15-pentaoxaoctadecan-18-one (82.7 mg, 110 μ mol, 1.0 equiv) in MeCN (1.09 mL) was added K_2CO_3 (45.6 mg, 330 μ mol, 3.0 equiv) followed by *tert*-butyl 2-(piperazin-1-yl)-5H,6H,7H,8H-pyrido[4,3-d]pyrimidine-6-carboxylate (42.1 mg, 132 μ mol, 1.2 equiv). The resulting suspension was heated to 80 °C for 8 h, then purified by silica gel chromatography (0→20% MeOH/DCM) to afford the product (75.1 mg, 70% yield). LCMS (ESI) *m/z*: [M + H] calcd for $C_{51}H_{67}N_{13}O_8$: 990.53; found 990.5.

Step 2: Synthesis of 18-{6-[(4-amino-3-{1H-pyrrolo[2,3-b]pyridin-5-yl}-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl]-1,2,3,4-tetrahydroisoquinolin-2-yl}-1-(4-{5H,6H,7H,8H-pyrido[4,3-d]pyrimidin-2-yl}piperazin-1-yl)-3,6,9,12,15-pentaoxaoctadecan-18-one

[00775] To a solution of *tert*-butyl 2-[4-(18-{6-[(4-amino-3-{1H-pyrrolo[2,3-b]pyridin-5-yl}-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl]-1,2,3,4-tetrahydroisoquinolin-2-yl}-18-oxo-3,6,9,12,15-pentaoxaoctadecan-1-yl)piperazin-1-yl]-5H,6H,7H,8H-pyrido[4,3-d]pyrimidine-6-carboxylate (75.1 mg, 75.8 μ mol, 1.0 equiv) in dioxane (1 mL) was added HCl (4 M in dioxane, 472 μ L, 1.89 mmol, 10.0 equiv). The solution was stirred at room temperature for 45 min, then concentrated under reduced pressure to afford the product. LCMS (ESI) *m/z*: [M + Na] calcd for $C_{46}H_{59}N_{13}O_6$: 912.46; found 912.5.


[00776] Following General Procedure 6, but using the appropriate PEG carboxylic acid and amine containing active site inhibitors in Table 2, the Intermediates B2 in Table 11 were prepared:

Table 11. Additional amines prepared

Structure	Molecular Formula	Calculated MW	Observed MW
 Intermediate B2-4	$C_{46}H_{59}N_{13}O_6$	[M + H] = 890.48	[M + H] = 890.5


[00777] Following General Procedure 1, but using the appropriate carboxylic acid PEG *tert*-butyl ester and amine containing active site inhibitors in Table 2, the Intermediates B4 in Table 12 were prepared:

Table 12. Additional amines prepared

Structure	Molecular Formula	Calculated MW	Observed MW
 Intermediate B4-1	C ₂₈ H ₃₈ N ₈ O ₈	[M + H] = 615.29	[M + H] = 615.1



[00778] Following General Procedure 1, but using the appropriate Intermediates B4 in Table 12 and amine containing pre-linkers in Table 4, the Intermediates B2 in Table 13 were prepared:








Table 13. Additional amines prepared








Structure	Molecular Formula	Calculated MW	Observed MW
 Intermediate B2-5	C ₃₉ H ₅₃ N ₁₃ O ₇	[M + H] = 816.43	[M + H] = 816.4









[00779] Following General Procedure 1, but using the appropriate Intermediates A1 and amine containing pre-linkers in Table 4, the Intermediates B2 in Table 14 were prepared:










Table 14. Additional amines prepared

Structure	Molecular Formula	Calculated MW	Observed MW
 Intermediate B2-6	C ₄₁ H ₅₂ N ₁₆ O ₆	[M + H] = 865.44	[M + H] = 865.2
 Intermediate B2-7	C ₃₉ H ₄₈ N ₁₆ O ₅	[M + H] = 821.41	[M + H] = 821.2

 Intermediate B2-8	$C_{37}H_{44}N_{16}O_4$	$[M + H] = 777.38$	$[M + H] = 777.3$
 Intermediate B2-9	$C_{49}H_{58}N_{16}O_7$	$[M + H] = 983.48$	$[M + H] = 983.4$
 Intermediate B2-10	$C_{43}H_{46}N_{16}O_4$	$[M + H] = 851.40$	$[M + H] = 851.4$
 Intermediate B2-11	$C_{43}H_{56}N_{16}O_6$	$[M + H] = 893.47$	$[M + H] = 893.3$
 Intermediate B2-12	$C_{49}H_{58}N_{16}O_6$	$[M + Na] = 989.46$	$[M + Na] = 989.4$
 Intermediate B2-13	$C_{47}H_{54}N_{16}O_5$	$[M + H] = 923.46$	$[M + H] = 923.4$
 Intermediate B2-14	$C_{39}H_{50}N_{16}O_6$	$[M + H] = 839.42$	$[M + H] = 839.3$

 Intermediate B2-15	$C_{47}H_{56}N_{16}O_7$	$[M + H] = 957.46$	$[M + H] = 957.7$
 Intermediate B2-16	$C_{45}H_{52}N_{16}O_6$	$[M + Na] = 935.42$	$[M + Na] = 935.3$
 Intermediate B2-17	$C_{43}H_{48}N_{16}O_5$	$[M + Na] = 891.39$	$[M + Na] = 891.4$
 Intermediate B2-18	$C_{41}H_{54}N_{16}O_6$	$[M + H] = 867.45$	$[M + H] = 867.3$
 Intermediate B2-19	$C_{47}H_{56}N_{16}O_6$	$[M + H] = 941.47$	$[M + H] = 941.2$
 Intermediate B2-20	$C_{48}H_{58}N_{16}O_6$	$[M + H] = 955.48$	$[M + H] = 955.2$
 Intermediate B2-21	$C_{45}H_{52}N_{16}O_5$	$[M + H] = 897.44$	$[M + H] = 897.3$
 Intermediate B2-22	$C_{48}H_{58}N_{16}O_8$	$[M + H] = 987.47$	$[M + H] = 987.42$






 Intermediate B2-23	C ₄₈ H ₅₈ N ₁₆ O ₇	[M + H] = 971.48	[M + H] = 971.31
 Intermediate B2-24	C ₄₄ H ₅₅ N ₁₃ O ₇	[M + H] = 878.44	[M + H] = 878.5
 Intermediate B2-25	C ₃₇ H ₅₁ N ₁₅ O ₆	[M + H] = 802.42	[M + H] = 802.4
 Intermediate B2-26	C ₄₃ H ₅₃ N ₁₅ O ₆	[M + H] = 876.44	[M + H] = 876.4
 Intermediate B2-27	C ₃₇ H ₅₁ N ₁₅ O ₅	[M + H] = 786.43	[M + H] = 786.5
 Intermediate B2-28	C ₅₀ H ₅₈ N ₁₆ O ₉	[M + H] = 1027.47	[M + H] = 1027.1
 Intermediate B2-29	C ₃₅ H ₄₃ N ₁₇ O ₄	[M + H] = 766.38	[M + H] = 766.3
 Intermediate B2-30	C ₄₁ H ₄₅ N ₁₇ O ₄	[M + H] = 840.39	[M + H] = 840.4







 Intermediate B2-31	C ₄₇ H ₅₇ N ₁₇ O ₇	[M + H] = 972.47	[M + H] = 972.5
 Intermediate B2-32	C ₄₈ H ₅₉ N ₁₇ O ₇	[M + H] = 986.49	[M + H] = 986.4
 Intermediate B2-33	C ₄₇ H ₅₇ N ₁₇ O ₆	[M + H] = 956.48	[M + H] = 956.3
 Intermediate B2-34	C ₄₉ H ₅₉ N ₁₇ O ₆	[M + H] = 982.49	[M + H] = 982.2
 Intermediate B2-35	C ₄₄ H ₆₁ N ₁₁ O ₉	[M + H] = 888.48	[M + H] = 888.3
 Intermediate B2-36	C ₄₄ H ₅₁ N ₁₇ O ₄	[M + H] = 882.44	[M + H] = 882.4
 Intermediate B2-37	C ₄₃ H ₅₂ N ₁₄ O ₄	[M + H] = 829.44	[M + H] = 829.3
 Intermediate B2-38	C ₄₁ H ₅₆ N ₁₀ O ₈	[M + H] = 817.44	[M + H] = 817.2
 Intermediate B2-39	C ₄₃ H ₅₉ N ₁₁ O ₉	[M + H] = 874.46	[M + H] = 874.3








Intermediate B2-39







[00780] Following General Procedure 2, but using the appropriate 4-nitrophenyl carbonate containing rapamycin monomer in Table 1 and Intermediates B2 from Tables 9, 11, and 13 and 14, the Series 2 bivalent analogs in Table 15 were synthesized:






Table 15. Series 2 Bivalent Compounds:

Structure	Molecular Formula	Calculated MW	Observed MW
 Example 27	$C_{99}H_{133}N_{17}O_{21}$	$[M + H] = 1896.99$	$[M + H] = 1896.7$
 Example 28	$C_{99}H_{135}N_{17}O_{21}$	$[M + H] = 1899.01$	$[M + H] = 1899.1$
 Example 29	$C_{100}H_{137}N_{17}O_{21}$	$[M + H] = 1913.03$	$[M + H] = 1913.0$
 Example 30	$C_{97}H_{129}N_{17}O_{20}$	$[M + H] = 1852.97$	$[M + H] = 1852.9$
 Example 31	$C_{95}H_{125}N_{17}O_{19}$	$[M + H] = 1808.94$	$[M + H] = 1809.0$

 Example 32	$C_{93}H_{131}N_{17}O_{20}$	$[M + H] = 1806.98$	$[M + H] = 1806.7$
 Example 33	$C_{99}H_{133}N_{17}O_{20}$	$[M + H] = 1881.00$	$[M + H] = 1881.0$
 Example 34	$C_{100}H_{135}N_{17}O_{20}$	$[M + H] = 1895.01$	$[M + H] = 1895.1$
 Example 35	$C_{95}H_{133}N_{17}O_{20}$	$[M + H] = 1833.00$	$[M + H] = 1832.9$
 Example 36	$C_{101}H_{135}N_{17}O_{20}$	$[M + H] = 1907.01$	$[M + H] = 1907.0$
 Example 37	$C_{96}H_{132}N_{14}O_{21}$	$[M + H] = 1817.98$	$[M + H] = 1817.9$


 Example 38	$C_{98}H_{136}N_{14}O_{20}$	$[M + H] = 1830.01$	$[M + H] = 1830.1$
 Example 39	$C_{101}H_{135}N_{17}O_{21}$	$[M + H] = 1923.01$	$[M + H] = 1923.1$
 Example 40	$C_{100}H_{135}N_{17}O_{22}$	$[M + H] = 1927.01$	$[M + H] = 1927.0$
 Example 41	$C_{100}H_{135}N_{17}O_{21}$	$[M + H] = 1911.01$	$[M + H] = 1911.0$
 Example 42	$C_{95}H_{130}N_{16}O_{20}$	$[M + H] = 1815.97$	$[M + H] = 1816.0$
 Example 43	$C_{94}H_{130}N_{14}O_{20}$	$[M + H] = 1775.97$	$[M + H] = 1775.9$
 Example 44	$C_{95}H_{134}N_{14}O_{20}$	$[M + H] = 1791.99$	$[M + H] = 1791.8$






 Example 45	$C_{89}H_{132}N_{14}O_{20}$	$[M + H] = 1717.98$	$[M + H] = 1717.9$
 Example 103	$C_{96}H_{134}N_{16}O_{20}$	$[M + H] = 1832.00$	$[M + H] = 1832.0$
 Example 104	$C_{90}H_{132}N_{16}O_{20}$	$[M + H] = 1757.99$	$[M + H] = 1757.9$
 Example 105	$C_{103}H_{139}N_{17}O_{23}$	$[M + H] = 1983.03$	$[M + H] = 1983.0$
 Example 106	$C_{100}H_{138}N_{18}O_{21}$	$[M + H] = 1928.04$	$[M + H] = 1927.9$
 Example 107	$C_{101}H_{140}N_{18}O_{21}$	$[M + H] = 1942.05$	$[M + H] = 1942.1$
 Example 108	$C_{99}H_{134}N_{18}O_{20}$	$[M + H] = 1896.01$	$[M + H] = 1896.0$



 Example 109	$C_{101}H_{136}N_{18}O_{20}$	$[M + H] = 1922.03$	$[M + H] = 1922.1$
 Example 110	$C_{96}H_{140}N_{12}O_{23}$	$[M + H] = 1830.02$	$[M + H] = 1829.9$
 Example 111	$C_{97}H_{142}N_{12}O_{23}$	$[M + H] = 1844.04$	$[M + H] = 1843.9$
 Example 149	$C_{93}H_{135}N_{11}O_{22}$	$[M + H] = 1758.99$	$[M + H] = 1758.9$
 Example 150	$C_{95}H_{138}N_{12}O_{23}$	$[M + H] = 1816.01$	$[M + H] = 1815.9$

[00781] Following General Procedure 1, but using the appropriate amine containing active site inhibitors in Table 2 and amine containing pre-linkers in Table 4, the Intermediates C1 in Table 16 were prepared:

Table 16. Additional amines prepared




Structure	Molecular Formula	Calculated MW	Observed MW
 Intermediate C1-1	$C_{36}H_{35}N_{15}O_2$	$[M + H] = 710.32$	$[M + H] = 710.2$





 Intermediate C1-2	$C_{30}H_{33}N_{15}O$	$[M + H] = 620.31$	$[M + H] = 620.2$
 Intermediate C1-3	$C_{38}H_{37}N_{15}O_2$	$[M + H] = 736.34$	$[M + H] = 736.2$
 Intermediate C1-4	$C_{25}H_{28}N_{12}O_2$	$[M + H] = 529.26$	$[M + H] = 529.5$
 Intermediate C1-5	$C_{33}H_{38}N_{10}O_2$	$[M + H] = 607.33$	$[M + H] = 607.3$
 Intermediate C1-6	$C_{31}H_{30}N_{12}O_2$	$[M + H] = 603.27$	$[M + H] = 603.3$

 Intermediate C1-7	$C_{31}H_{30}N_{12}O$	$[M + H] = 587.28$	$[M + H] = 587.3$
 Intermediate C1-8	$C_{29}H_{32}N_{10}O_2$	$[M + H] = 553.28$	$[M + H] =$

[00782] Following General Procedure 1, but using the PEG carboxylic acids and Intermediates C1 in Table 16, the Intermediates C2 in Table 17 were prepared:



Table 17. Additional amines prepared





Structure	Molecular Formula	Calculated MW	Observed MW
 Intermediate C2-1	$C_{47}H_{56}N_{16}O_7$	$[M + H] = 957.46$	$[M + H] = 957.7$
 Intermediate C2-2	$C_{41}H_{54}N_{16}O_6$	$[M + H] = 867.45$	$[M + H] = 867.2$
 Intermediate C2-3	$C_{43}H_{46}N_{16}O_4$	$[M + H] = 851.40$	$[M + H] = 851.2$

 Intermediate C2-4	$C_{36}H_{49}N_{13}O_7$	$[M + H] = 776.40$	$[M + H] = 776.3$
 Intermediate C2-5	$C_{30}H_{37}N_{13}O_4$	$[M + H] = 644.32$	$[M + H] = 644.3$
 Intermediate C2-6	$C_{48}H_{67}N_{11}O_9$	$[M + H] = 942.52$	$[M + H] = 943.2$
 Intermediate C2-7	$C_{44}H_{61}N_{11}O_9$	$[M + H] = 888.48$	$[M + H] = 888.3$

[00783] Following General Procedure 2, but using the appropriate 4-nitrophenyl carbonate containing rapamycin monomer in Table 1 and Intermediates C2 from Table 17, the Series 3 bivalent analogs in Table 18 were synthesized:


Table 18. Series 3 Bivalent Compounds

Structure	Molecular Formula	Calculated MW	Observed MW
 Example 46	$C_{99}H_{133}N_{17}O_{21}$	$[M + H] = 1896.99$	$[M + H] = 1897.3$
 Example 47	$C_{93}H_{131}N_{17}O_{20}$	$[M + H] = 1806.98$	$[M + H] = 1806.8$

 Example 48	$C_{100}H_{144}N_{12}O_{23}$	$[M + H] = 1882.06$	$[M + H] = 1882.1$
 Example 49	$C_{99}H_{135}N_{17}O_{21}$	$[M + H] = 1899.01$	$[M + H] = 1899.1$
 Example 50	$C_{100}H_{137}N_{17}O_{21}$	$[M + H] = 1913.03$	$[M + H] = 1913.1$
 Example 112	$C_{96}H_{140}N_{12}O_{23}$	$[M + H] = 1830.02$	$[M + H] = 1829.9$
 Example 113	$C_{100}H_{146}N_{12}O_{23}$	$[M + H] = 1884.07$	$[M + H] = 1883.7$






[00784] Following General Procedure 1, but using the appropriate Intermediates C2 in Table 17 and amine containing pre-linkers in Table 4, the Intermediates D1 in Table 19 were prepared:








Table 19. Additional amines prepared








Structure	Molecular Formula	Calculated MW	Observed MW
 Intermediate D1-1	$C_{44}H_{52}N_{20}O_5$	$[M + H] = 941.45$	$[M + H] = 941.5$



[00785] Following General Procedure 1, but using the appropriate amine containing active site inhibitors in Table 2 and amine containing pre-linkers in Table 4, the Intermediates D1 in Table 20 were prepared:

Table 20. Additional amines prepared

Structure	Molecular Formula	Calculated MW	Observed MW
 Intermediate D1-2	C ₃₉ H ₄₉ N ₁₇ O ₄	[M + H] = 820.43	[M + H] = 820.4
 Intermediate D1-3	C ₄₅ H ₅₁ N ₁₇ O ₄	[M + H] = 894.44	[M + H] = 894.4
 Intermediate D1-4	C ₃₉ H ₄₉ N ₁₇ O ₃	[M + H] = 804.43	[M + H] = 804.4
 Intermediate D1-5	C ₄₅ H ₅₁ N ₁₇ O ₃	[M + H] = 878.45	[M + H] = 878.4
 Intermediate D1-6	C ₄₀ H ₄₉ N ₁₇ O ₄	[M + H] = 832.43	[M + H] = 832.4



 Intermediate D1-7	<chem>C58H63F3N18O5</chem>	$[M + H] = 1149.53$	$[M + H] = 1149.4$
 Intermediate D1-8	<chem>C45H56N16O5</chem>	$[M + H] = 901.47$	$[M + H] = 901.4$
 Intermediate D1-9	<chem>C38H51N15O4</chem>	$[M + H] = 782.43$	$[M + H] = 782.4$
 Intermediate D1-10	<chem>C47H53N17O4</chem>	$[M + H] = 920.46$	$[M + H] = 920.4$
 Intermediate D1-11	<chem>C47H56FN13O7S</chem>	$[M + H] = 966.42$	$[M + H] = 966.3$
 Intermediate D1-12	<chem>C47H57N13O7S</chem>	$[M + H] = 948.43$	$[M + H] = 948.4$
 Intermediate D1-13	<chem>C47H55N17O4</chem>	$[M + H] = 922.47$	$[M + H] = 922.4$







 Intermediate D1-14	$C_{47}H_{53}N_{17}O_3$	$[M + H] = 904.46$	$[M + H] = 904.4$
 Intermediate D1-15	$C_{44}H_{51}N_{17}O_3$	$[M + H] = 866.45$	$[M + H] = 866.3$
 Intermediate D1-16	$C_{46}H_{53}N_{17}O_3$	$[M + H] = 892.46$	$[M + H] = 892.3$
 Intermediate D1-17	$C_{44}H_{51}N_{17}O_2$	$[M + H] = 850.45$	$[M + H] = 850.3$
 Intermediate D1-18	$C_{46}H_{53}N_{17}O_2$	$[M + H] = 876.47$	$[M + H] = 876.3$
 Intermediate D1-19	$C_{44}H_{53}N_{15}O_5$	$[M + H] = 872.45$	$[M + H] = 872.3$
 Intermediate D1-20	$C_{46}H_{51}N_{17}O_6$	$[M + H] = 938.61$	$[M + H] = 938.3$







 Intermediate D1-21	$C_{46}H_{51}N_{17}O_5$	$[M + H] = 922.44$	$[M + H] = 922.3$
 Intermediate D1-22	$C_{47}H_{59}N_{15}O_4$	$[M + H] = 898.50$	$[M + H] = 898.4$







[00786] Following General Procedure 2, but using the appropriate 4-nitrophenyl carbonate containing rapamycin monomer in Table 1 and Intermediates D1 from Tables 19 and 20, the Series 4 bivalent analogs in Table 21 were synthesized:






Table 21. Series 4 Bivalent Compounds




Structure	Molecular Formula	Calculated MW	Observed MW
 Example 51	$C_{97}H_{128}N_{18}O_{18}$	$[M + H] = 1833.98$	$[M + H] = 1834.1$
 Example 52	$C_{97}H_{128}N_{18}O_{17}$	$[M + H] = 1817.98$	$[M + H] = 1817.9$

 Example 53	$C_{96}H_{129}N_{21}O_{19}$	$[M + H] = 1880.99$	$[M + H] = 1881.0$
 Example 54	$C_{91}H_{126}N_{18}O_{18}$	$[M + H] = 1759.96$	$[M + H] = 1760.0$
 Example 55	$C_{97}H_{130}N_{18}O_{18}$	$[M + H] = 1835.99$	$[M + H] = 1835.8$
 Example 56	$C_{98}H_{132}N_{18}O_{18}$	$[M + H] = 1850.01$	$[M + H] = 1849.8$
 Example 57	$C_{92}H_{130}N_{18}O_{18}$	$[M + H] = 1775.99$	$[M + H] = 1775.9$
 Example 58	$C_{90}H_{128}N_{16}O_{18}$	$[M + H] = 1721.97$	$[M + H] = 1721.9$

Example 58 	$C_{100}H_{134}N_{18}O_{18}$	$[M + H] = 1876.02$	$[M + H] = 1876.1$
Example 114 	$C_{97}H_{130}N_{18}O_{17}$	$[M + H] = 1819.99$	$[M + H] = 1820.1$
Example 115 	$C_{97}H_{128}N_{18}O_{17}S$	$[M + H] = 1849.95$	$[M + H] = 1849.9$
Example 116 	$C_{97}H_{130}N_{18}O_{17}S$	$[M + H] = 1851.97$	$[M + H] = 1851.9$
Example 117 	$C_{98}H_{132}N_{18}O_{17}S$	$[M + H] = 1865.98$	$[M + H] = 1865.7$
Example 118 	$C_{99}H_{135}FN_{14}O_{21}S$	$[M + H] = 1907.97$	$[M + H] = 1908.0$



Example 119 	$C_{100}H_{137}FN_{14}O_{21}S$	$[M + H] = 1921.99$	$[M + H] = 1922.0$
Example 120 	$C_{99}H_{134}N_{14}O_{21}S$	$[M + H] = 1887.96$	$[M + H] = 1888.0$
Example 121 	$C_{99}H_{136}N_{14}O_{21}S$	$[M + H] = 1889.98$	$[M + H] = 1890.0$
Example 122 	$C_{99}H_{132}N_{18}O_{18}$	$[M + H] = 1862.00$	$[M + H] = 1861.9$
Example 123 	$C_{99}H_{130}N_{18}O_{17}$	$[M + H] = 1843.99$	$[M + H] = 1844.1$
Example 124 	$C_{97}H_{132}N_{18}O_{17}$	$[M + H] = 1822.01$	$[M + H] = 1822.0$
Example 125 			







 Example 126	$C_{99}H_{134}N_{18}O_{17}$	$[M + H] = 1848.03$	$[M + H] = 1848.0$
 Example 127	$C_{96}H_{128}N_{18}O_{16}$	$[M + H] = 1789.98$	$[M + H] = 1789.9$
 Example 128	$C_{98}H_{130}N_{18}O_{16}$	$[M + H] = 1816.00$	$[M + H] = 1815.9$
 Example 129	$C_{96}H_{132}N_{16}O_{19}$	$[M + H] = 1813.99$	$[M + H] = 1814.0$
 Example 130	$C_{97}H_{134}N_{16}O_{19}$	$[M + H] = 1828.01$	$[M + H] = 1828.0$
 Example 131	$C_{99}H_{132}N_{18}O_{20}$	$[M + H] = 1893.99$	$[M + H] = 1894.0$

 Example 132	$C_{98}H_{128}N_{18}O_{19}$	$[M + H] = 1861.97$	$[M + H] = 1861.9$
 Example 133	$C_{97}H_{128}N_{18}O_{18}$	$[M + H] = 1833.97$	$[M + H] = 1833.9$
 Example 151	$C_{97}H_{129}N_{21}O_{17}$	$[M + H] = 1861.00$	$[M + H] = 1860.8$

[00787] Following General Procedure 1, but using the appropriate Intermediates C1 in Table 16 and amine containing pre-linkers in Table 4, the Intermediates E1 in Table 22 were prepared:

Table 22. Additional amines prepared

Structure	Molecular Formula	Calculated MW	Observed MW
 Intermediate E1-1	$C_{39}H_{43}N_{19}O_3$	$[M + H] = 826.39$	$[M + H] = 826.5$
 Intermediate E1-2	$C_{45}H_{45}N_{19}O_3$	$[M + H] = 900.41$	$[M + H] = 900.2$

 Intermediate E1-3	$C_{45}H_{45}N_{19}O_2$	$[M + H] = 884.41$	$[M + H] = 884.4$
 Intermediate E1-4	$C_{47}H_{47}N_{19}O_3$	$[M + H] = 926.42$	$[M + H] = 926.6$
 Intermediate E1-5	$C_{47}H_{47}N_{19}O_2$	$[M + H] = 910.43$	$[M + H] = 910.2$
 Intermediate E1-6	$C_{38}H_{47}N_{17}O_3$	$[M + H] = 790.41$	$[M + H] = 790.4$
 Intermediate E1-7	$C_{44}H_{49}N_{17}O_3$	$[M + H] = 864.43$	$[M + H] = 864.3$
 Intermediate E1-8	$C_{39}H_{47}N_{17}O_3$	$[M + H] = 802.41$	$[M + H] = 802.3$

[00788] Following General Procedure 2, but using the appropriate 4-nitrophenyl carbonate containing rapamycin monomer in Table 1 and Intermediates E1 from Table 22, the Series 5 bivalent analogs in Table 23 were synthesized:

Table 23. Series 5 Bivalent Compounds










Structure	Molecular Formula	Calculated MW	Observed MW
 Example 59	$C_{91}H_{120}N_{20}O_{17}$	$[M + H] = 1765.92$	$[M + H] = 1765.9$
 Example 60	$C_{97}H_{122}N_{20}O_{17}$	$[M + H] = 1839.94$	$[M + H] = 1840.0$
 Example 61	$C_{97}H_{122}N_{20}O_{16}$	$[M + H] = 1823.94$	$[M + H] = 1823.9$
 Example 62	$C_{99}H_{124}N_{20}O_{17}$	$[M + H] = 1865.95$	$[M + H] = 1865.8$
 Example 134	$C_{99}H_{124}N_{20}O_{16}$	$[M + H] = 1849.96$	$[M + H] = 1850.0$

Table 24. Additional amines prepared

Structure	Molecular Formula	Calculated MW	Observed MW
 Intermediate F1-1	C ₄₃ H ₆₁ N ₁₁ O ₈	[M + H] = 860.48	[M + H] = 860.4
 Intermediate F1-2	C ₄₄ H ₆₁ N ₁₁ O ₈	[M + H] = 872.48	[M + H] = 872.2

[00789] Following General Procedure 2, but using the appropriate 4-nitrophenyl carbonate containing rapamycin monomer in Table 1 and Intermediates F1 from Table 24, the Series 6 bivalent analogs in Table 25 were synthesized:




Table 25. Series 6 Bivalent Compounds

Structure	Molecular Formula	Calculated MW	Observed MW
 Example 135	C ₉₅ H ₁₄₀ N ₁₂ O ₂₂	[M + H] = 1802.03	[M + H] = 1802.5
 Example 152	C ₉₆ H ₁₄₀ N ₁₂ O ₂₂	[M + H] = 1814.03	[M + H] = 1813.9

[00790] Following General Procedure 1, but using the appropriate Intermediates A1 in Table 5 and amine containing pre-linkers in Table 4, the Intermediates G1 in Table 26 were prepared:


Table 26. Additional amines prepared

Structure	Molecular Formula	Calculated MW	Observed MW

 Intermediate G1-1	$C_{44}H_{58}N_{18}O_6$	$[M + H] = 935.49$	$[M + H] = 935.5$
 Intermediate G1-2	$C_{50}H_{60}N_{18}O_6$	$[M + H] = 1009.50$	$[M + H] = 1009.5$
 Intermediate G1-3	$C_{44}H_{58}N_{18}O_5$	$[M + H] = 919.49$	$[M + H] = 919.5$

[00791] Following General Procedure 6, but using the appropriate Intermediates B3 in Table 10 and amine containing pre-linkers in Table 4, the Intermediates G1 in Table 27 were prepared:





Table 27. Additional amines prepared

Structure	Molecular Formula	Calculated MW	Observed MW
 Intermediate G1-4	$C_{45}H_{57}N_{15}O_5$	$[M + H] = 888.48$	$[M + H] = 888.4$

[00792] Following General Procedure 2, but using the appropriate 4-nitrophenyl carbonate containing rapamycin monomer in Table 1 and Intermediates G1 from Tables 26 and 27, the Series 7 bivalent analogs in Table 28 were synthesized:


Table 28. Series 7 Bivalent Compounds



Structure	Molecular Formula	Calculated MW	Observed MW

 Example 63	$C_{96}H_{135}N_{19}O_{20}$	$[M + H] = 1875.02$	$[M + H] = 1874.9$
 Example 64	$C_{102}H_{137}N_{19}O_{20}$	$[M + H] = 1949.04$	$[M + H] = 1949.0$
 Example 65	$C_{103}H_{141}N_{19}O_{20}$	$[M + H] = 1965.07$	$[M + H] = 1965.0$
 Example 66	$C_{96}H_{135}N_{19}O_{19}$	$[M + H] = 1859.03$	$[M + H] = 1859.0$

[00793] Following General Procedure 1, but using the appropriate Intermediates D1 in Tables 19 and 20 and PEG carboxylic acids, the Intermediates H1 in Table 29 were prepared:





Table 29. Additional amines prepared

Structure	Molecular Formula	Calculated MW	Observed MW
 Intermediate H1-1	$C_{44}H_{58}N_{18}O_6$	$[M + H] = 935.49$	$[M + H] = 935.5$

 Intermediate H1-2	$C_{50}H_{60}N_{18}O_6$	$[M + 2H]/2 = 505.23$	$[M + 2H]/2 = 505.4$
 Intermediate H1-3	$C_{44}H_{58}N_{18}O_5$	$[M + 2H]/2 = 460.25$	$[M + 2H]/2 = 460.3$

[00794] Following General Procedure 2, but using the appropriate 4-nitrophenyl carbonate containing rapamycin monomer in Table 1 and Intermediates H1 from Table 29, the Series 8 bivalent analogs in Table 30 were synthesized:

Table 30. Series 8 Bivalent Compounds:

Structure	Molecular Formula	Calculated MW	Observed MW
 Example 67	$C_{102}H_{137}N_{19}O_{20}$	$[M + H] = 1949.04$	$[M + H] = 1949.0$
 Example 68	$C_{96}H_{135}N_{19}O_{19}$	$[M + H] = 1859.03$	$[M + H] = 1859.0$
 Example 69	$C_{102}H_{139}N_{19}O_{20}$	$[M + H] = 1951.05$	$[M + H] = 1951.1$
 Example 70	$C_{103}H_{141}N_{19}O_{20}$	$[M + H] = 1965.07$	$[M + H] = 1965.1$

Biological Examples

Cell Based AlphaLISA Assays For Determining IC₅₀ For Inhibition of P-Akt (S473), P-4E-BP1 (T37/46), and P-P70S6K (T389) in MDA-MB-468 Cells

mTOR Kinase Cellular Assay

[00795] To measure functional activity of mTORC1 and mTORC2 in cells the phosphorylation of 4EBP1 (Thr37/46) and P70S6K (Thr389), and AKT1/2/3 (Ser473) was monitored using AlphaLisa SureFire Ultra Kits (Perkin Elmer). MDA-MB-468 cells (ATCC® HTB-132) were cultured in 96-well tissue culture plates and treated with compounds in the disclosure at concentrations varying from 0.017 – 1,000 nM for two to four hours at 37 °C. Incubations were terminated by removal of the assay buffer and addition of lysis buffer provided with the assay kit. Samples were processed according to the manufacturer's instructions. The Alpha signal from the respective phosphoproteins was measured in duplicate using a microplate reader (Envision, Perkin-Elmer or Spectramax M5, Molecular Devices). Inhibitor concentration response curves were analyzed using normalized IC₅₀ regression curve fitting with control based normalization.

[00796] As an example, measured IC₅₀ values for selected compounds are reported below:

	IC ₅₀ for Inhibition of mTORC1 and mTORC2 Substrate Phosphorylation (nM)		
Compound	p-P70S6K-(T389)	p-4E-BP1-(T37/46)	p-AKT1/2/3-(S473)
MLN-128	1.4	16	3.7
Rapamycin	0.2	>1,000	>1,000

[00797] As an example, measured pIC₅₀ values for selected compounds are reported below:

Example	pIC ₅₀ for Inhibition of mTORC1 and mTORC2 Substrate Phosphorylation		
	p-P70S6K-(T389)	p-4E-BP1-(T37/46)	p-AKT1/2/3-(S473)
1	+++	+++	++
2	+++	+++	++
3	+++	+++	++
4	+++	+++	++
5	+++	+++	+
6	+++	+++	++
7	+++	+++	++
8	+++	+++	++

Example	pIC ₅₀ for Inhibition of mTORC1 and mTORC2 Substrate Phosphorylation		
	p-P70S6K-(T389)	p-4E-BP1-(T37/46)	p-AKT1/2/3-(S473)
9	+++	+++	++
10	+++	-	-
11	+++	+++	++
12	+++	+++	+
13	+++	+++	++
14	++	+++	++
15	+++	+++	++
16	+++	+++	+++
17	+++	+++	++
18	+++	++	+
19	+++	+++	+++
20	+++	+++	+++
21	+++	+++	+
27	+++	+++	++
28	+++	+++	-
29	+++	+++	-
30	+++	+++	++
31	+++	+++	++
32	+++	+++	++
33	+++	+++	-
34	+++	-	-
35	+++	+++	++
36	+++	+++	-
37	+++	+++	-
38	+++	-	-
39	+++	+++	+++
40	+++	+++	++
41	+++	+++	-
42	+++	+++	++
43	+++	+++	++
46	+++	+++	++
47	+++	+++	++
48	+++	+++	++
49	+++	+++	++
50	+++	+++	+
51	+++	+++	+++
52	+++	+++	-
53	+++	+++	++
54	+++	+++	+++

Example	pIC ₅₀ for Inhibition of mTORC1 and mTORC2 Substrate Phosphorylation		
	p-P70S6K-(T389)	p-4E-BP1-(T37/46)	p-AKT1/2/3-(S473)
55	+++	+++	++
56	+++	+++	-
59	+++	+++	+++
60	+++	++	+
61	+++	-	-
62	++	-	-
63	+++	+++	+++
64	+++	+++	++
65	+++	+++	+
66	+++	+++	++
67	+	+	+
68	+	+	-
69	+++	+++	++
70	+++	+++	+
71	+++	+++	++
72	+++	+++	+++
73	+++	+++	+
74	+++	+++	+
75	+++	+++	++
76	+++	+++	++
77	+++	++	+
78	+++	+++	+++
79	+++	+++	+
80	+++	+++	+++
81	+++	++	-
82	+++	+++	-
83	+++	+++	++
84	+++	+++	++
85	+++	+++	++
86	+++	+++	++
87	+++	+++	++
88	+++	+++	++
89	+++	+++	-
90	+++	-	-
91	+++	+++	++
92	+++	+++	++
93	+++	+++	++
94	+++	+++	+++
95	+++	+++	++

Example	pIC ₅₀ for Inhibition of mTORC1 and mTORC2 Substrate Phosphorylation		
	p-P70S6K-(T389)	p-4E-BP1-(T37/46)	p-AKT1/2/3-(S473)
96	+++	+++	+
97	+++	+++	++
98	+++	+++	++
99	+++	+++	-
100	+++	+++	++
101	+++	+++	++
102	+++	+++	++
103	+++	-	-
104	+++	+++	-
105	+++	++	-
106	+++	++	-
107	+++	-	-
108	++	-	-
109	+	-	-
110	++	-	-
111	+++	+++	-
112	+++	-	-
113	+++	+++	++
114	+++	+++	-
115	+++	+++	+++
116	+++	+++	-
117	+++	+++	++
118	+++	+++	++
119	+++	+++	++
120	+++	+++	++
121	+++	+++	-
122	+++	+++	++
123	+++	+++	+
124	+++	+++	++
125	+++	+++	++
126	+++	++	-
127	+++	++	-
128	+++	++	+
129	+++	+++	-
130	+++	+++	-
131	+++	+++	++
132	+++	+++	++
133	+++	+++	++
134	+++	+++	-

Example	pIC ₅₀ for Inhibition of mTORC1 and mTORC2 Substrate Phosphorylation		
	p-P70S6K-(T389)	p-4E-BP1-(T37/46)	p-AKT1/2/3-(S473)
135	+++	-	-
136	+++	++	++
137	+++	+++	++
138	+++	+++	+++
139	+++	+++	+
140	+	+	-
141	+++	+++	+
142	++	-	-
143	+++	++	-
144	+++	+++	++
145	+++	++	+
146	-	-	-
147	+++	+++	++
148	+++	+++	++
149	+++	+++	++
150	+++	+++	+
151	+++	++	-
152	+++	+++	+

Note:


pIC ₅₀ (p-P70S6K-(T389))	
≥9	+++
9>pIC ₅₀ ≥8	++
8>pIC ₅₀ ≥6	+
<6	-
pIC ₅₀ (p-4E-BP1-(T37/46) or p-AKT1/2/3-(S473))	
≥8.5	+++
8.5>pIC ₅₀ ≥7.5	++
7.5>pIC ₅₀ ≥6.0	+
<6	-

Equivalents

[00798] While the present disclosure has been described in conjunction with the specific embodiments set forth above, many alternatives, modifications and other variations thereof will be apparent to those of ordinary skill in the art. All such alternatives, modifications and variations are intended to fall within the spirit and scope of the present disclosure.

CLAIMS

1. A compound of Formula Ic:



or a pharmaceutically acceptable salt or tautomer thereof, wherein:

R^{32} is $-H$, $=O$, $-OR^3$, $-N_3$, or $-O-C(=Z^1)-R^{32a}$;

R^{28} is $-H$, $(C_1-C_6)alkyl$, or $-C(=Z^1)-R^{28a}$;






R^{40} is $-H$ or $-C(=Z^1)-R^{40a}$;

wherein when R^{28} and R^{40} are H , then R^{32} is not $=O$;

each Z^1 is independently O or S ;

R^{28a} , R^{32a} , and R^{40a} are independently $-A^1-L^1-A^2-B$; $-A^1-A^2-B$; $-L^2-A^1-L^1-A^2-L^3-B$; $-O-(C_1-C_6)alkyl$; or $-O-(C_6-C_{10})aryl$; wherein the aryl of $-O-(C_6-C_{10})aryl$ is unsubstituted or substituted with 1-5 substituents selected from $-NO_2$ and halogen;

A^1 and A^2 are independently absent or are independently selected from

$\text{---} \text{N} \left[\text{C}(\text{R}^3)_2 \right]_p \text{---} \text{Q} \text{---} \left[\text{C}(\text{R}^3)_2 \right]_p \text{---} \text{C}(\text{O}) \text{---}$	$\text{---} \text{N} \left[\text{C}(\text{R}^3)_2 \right]_p \text{---} \text{Q} \text{---} \left[\text{C}(\text{R}^3)_2 \right]_p \text{---} \text{N} \text{---}$	$\text{---} \text{N} \left[\text{C}(\text{R}^3)_2 \right]_p \text{---} \text{X} \text{---} \text{X}^1 \text{---} \text{N} \text{---}$
$\text{---} \text{N} \text{---} \text{W}^1 \text{---} \text{W} \text{---} \left[\text{C}(\text{R}^3)_2 \right]_p \text{---} \text{C}(\text{O}) \text{---}$	$\text{---} \text{N} \text{---} \text{W}^1 \text{---} \text{W} \text{---} \left[\text{C}(\text{R}^3)_2 \right]_p \text{---} \text{N} \text{---}$	$\text{---} \text{N} \text{---} \text{G}^1 \text{---} \text{N} \text{---}$
$\text{---} \text{N} \text{---} \text{G}^1 \text{---} \text{N} \text{---} \text{G}^2 \text{---} \text{N} \text{---}$		
	 and	

wherein the bond on the left side of A^1 , as drawn, is bound to $-\text{C}(=\text{Z}^1)-$ or L^2 ; and
wherein the bond on the right side of the A^2 moiety, as drawn, is bound to B or L^3 ;

each Q is independently 1 to 3 rings selected from arylene, cycloalkylene, heteroarylene, and heterocyclylene;

each X is independently absent or 1 to 2 rings selected from arylene, cycloalkylene, heteroarylene, and heterocyclylene;

each X^1 is independently a heteroarylene or heterocyclylene ring;


each W is independently absent or 1 to 2 rings selected from arylene, cycloalkylene, heteroarylene, and heterocyclylene;


each W^1 is independently a heteroarylene or heterocyclylene ring;

each G is independently absent or a ring selected from arylene, cycloalkylene, heteroarylene, and heterocyclylene;


each G^1 and G^2 are independently heteroarylene or heterocyclylene ring;

each L^1 is independently selected from






L^2 and L^3 are independently absent or are independently selected from




each B is independently selected from



each B^1 is independently selected from $^{-3}\text{-NR}^3\text{-(C(R}^3)_2)_n\text{-}$, $^{-3}\text{-NR}^3\text{-(C(R}^3)_2)_n\text{-(C}_6\text{-C}_{10}\text{)arylene-(C(R}^3)_2)_n\text{-}$,

$^{-3}\text{-NR}^3\text{-(C(R}^3)_2)_n\text{-heteroarylene-}$, $^{-3}\text{-NR}^3\text{-(C(R}^3)_2)_n\text{-heteroarylene-(C(R}^3)_2)_n\text{-}$,

$\text{---}^3\text{---}(\text{C}_6\text{--C}_{10})\text{arylene-}$, $\text{---}^3\text{---NR}^3\text{---}(\text{C}(\text{R}^3)_2)_n\text{---NR}^3\text{C}(\text{O})\text{---}$, $\text{---}^3\text{---NR}^3\text{---}(\text{C}(\text{R}^3)_2)_n\text{---heteroarylene-}$
heterocyclylene- $(\text{C}_6\text{--C}_{10})\text{arylene-}$, $\text{---}^3\text{---heteroarylene-heterocyclylene-}(\text{C}_6\text{--C}_{10})\text{arylene-}$,



$\text{S}(\text{O})_2\text{---arylene-C}(\text{O})\text{---}$, and $\text{---}^3\text{---NR}^3\text{---}(\text{C}(\text{R}^3)_2)_n\text{---S}(\text{O})_2\text{---arylene-}(\text{C}(\text{R}^3)_2)_n\text{---}$, wherein the $\text{---}^3\text{---}$ bond on the left side of B^1 , as drawn, is bound to A^2 , L^3 , or L^1 ; and wherein the heteroarylene, heterocyclylene, and arylene are each independently optionally substituted with alkyl, hydroxyalkyl, haloalkyl, alkoxy, halogen, or hydroxyl;

each R^3 is independently H or $(\text{C}_1\text{--C}_6)\text{alkyl}$;

each R^4 is independently H, $(\text{C}_1\text{--C}_6)\text{alkyl}$, halogen, 5-12 membered heteroaryl, 5-12 membered heterocyclyl, $(\text{C}_6\text{--C}_{10})\text{aryl}$, wherein the heteroaryl, heterocyclyl, and aryl are optionally substituted with $\text{---N}(\text{R}^3)_2$, ---OR^3 , halogen, $(\text{C}_1\text{--C}_6)\text{alkyl}$, $\text{---}(\text{C}_1\text{--C}_6)\text{alkylene-heteroaryl}$, $\text{---}(\text{C}_1\text{--C}_6)\text{alkylene-CN}$, $\text{---C}(\text{O})\text{NR}^3\text{---heteroaryl}$, or $\text{---C}(\text{O})\text{NR}^3\text{---heterocyclyl}$;

each R^5 is independently H, $(\text{C}_1\text{--C}_6)\text{alkyl}$, $\text{---C}(\text{O})\text{OR}^3$, or $\text{---N}(\text{R}^3)_2$, wherein the alkyl of $(\text{C}_1\text{--C}_6)\text{alkyl}$ is optionally substituted with $\text{---N}(\text{R}^3)_2$ or ---OR^3 ;

each R^6 is independently H, $(\text{C}_1\text{--C}_6)\text{alkyl}$, $\text{---C}(\text{O})\text{OR}^3$, or $\text{---N}(\text{R}^3)_2$, wherein the alkyl of $(\text{C}_1\text{--C}_6)\text{alkyl}$ is optionally substituted with $\text{---N}(\text{R}^3)_2$ or ---OR^3 ;

each R^7 is independently H, $(\text{C}_1\text{--C}_6)\text{alkyl}$, $\text{---C}(\text{O})\text{OR}^3$, or $\text{---N}(\text{R}^3)_2$, wherein the alkyl of $(\text{C}_1\text{--C}_6)\text{alkyl}$ is optionally substituted with $\text{---N}(\text{R}^3)_2$ or ---OR^3 ;

each R^8 is independently H, $(\text{C}_1\text{--C}_6)\text{alkyl}$, $\text{---C}(\text{O})\text{OR}^3$, or $\text{---N}(\text{R}^3)_2$, wherein the alkyl of $(\text{C}_1\text{--C}_6)\text{alkyl}$ is optionally substituted with $\text{---N}(\text{R}^3)_2$ or ---OR^3 ;

each Y is independently $\text{C}(\text{R}^3)_2$ or a bond;

each n is independently an integer from one to 12;

each o is independently an integer from zero to 30;

each p is independently an integer from zero to 12;


each q is independently an integer from zero to 30; and

each r is independently an integer from one to 6.

2. The compound of claim 1, or a pharmaceutically acceptable salt or tautomer thereof, wherein R^{32} is =O.

3. The compound of claim 1, or a pharmaceutically acceptable salt or tautomer thereof, wherein R^{32} is -OR³.

4. The compound of any one of claims 1-3, or a pharmaceutically acceptable salt or tautomer thereof, wherein the compound is represented by the structure of Formula (I-40b):



wherein R^{40} is $-C(=Z^1)-R^{40a}$.

5. The compound of claim 4, or a pharmaceutically acceptable salt or tautomer thereof, wherein Z^1 is O.

6. The compound of claim 4, or a pharmaceutically acceptable salt or tautomer thereof, wherein Z^1 is S.

7. The compound of any one of claims 4-6, or a pharmaceutically acceptable salt or tautomer thereof, wherein R^{40a} is $-A^1-L^1-A^2-B$, wherein A^1 and A^2 are absent.

8. The compound of any one of claims 4-6, or a pharmaceutically acceptable salt or tautomer thereof, wherein R^{40a} is $-A^1-L^1-A^2-B$, wherein A^2 is absent.

9. The compound of any one of claims 4-6, or a pharmaceutically acceptable salt or tautomer thereof, wherein R^{40a} is $-A^1-L^1-A^2-B$, wherein A^1 is absent.

10. The compound of any one of claims 4-6, or a pharmaceutically acceptable salt or tautomer thereof, wherein R^{40a} is $-A^1-L^1-A^2-B$.

11. The compound of any one of claims 4-6, or a pharmaceutically acceptable salt or tautomer thereof, wherein R^{40a} is $-A^1-A^2-B$.


12. The compound of any one of claims 4-6, or a pharmaceutically acceptable salt or tautomer thereof, wherein R^{40a} is $-L^2-A^1-L^1-A^2-L^3-B$, wherein L^2 and A^1 are absent.

13. The compound of any one of claims 4-6, or a pharmaceutically acceptable salt or tautomer thereof, wherein R^{40a} is $-L^2-A^1-L^1-A^2-L^3-B$, wherein L^2 is absent.

14. The compound of any one of claims 4-6, or a pharmaceutically acceptable salt or tautomer thereof, wherein R^{40a} is $-L^2-A^1-L^1-A^2-L^3-B$, wherein L^3 is absent.

15. The compound of any one of claims 4-6, or a pharmaceutically acceptable salt or tautomer thereof, wherein R^{40a} is $-O-(C_1-C_6)\text{alkyl}$ or $-O-(C_6-C_{10})\text{aryl}$; wherein the aryl of $-O-(C_6-C_{10})\text{aryl}$ is unsubstituted or substituted with 1-5 substituents selected from $-\text{NO}_2$ and halogen.

16. The compound of any one of claims 1-3, or a pharmaceutically acceptable salt or tautomer thereof, wherein the compound is represented by the structure of Formula (I-28b):



wherein R^{28} is $-C(=Z^1)-R^{28a}$.

17. The compound of claim 16, or a pharmaceutically acceptable salt or tautomer thereof, wherein Z^1 is O.

18. The compound of claim 16, or a pharmaceutically acceptable salt or tautomer thereof, wherein Z^1 is S.

19. The compound of any one of claims 16-18, or a pharmaceutically acceptable salt or tautomer thereof, wherein R^{28a} is $-A^1-L^1-A^2-B$, wherein A^1 and A^2 are absent.

20. The compound of any one of claims 16-18, or a pharmaceutically acceptable salt or tautomer thereof, wherein R^{28a} is $-A^1-L^1-A^2-B$, wherein A^2 is absent.

21. The compound of any one of claims 16-18, or a pharmaceutically acceptable salt or tautomer thereof, wherein R^{28a} is $-A^1-L^1-A^2-B$, wherein A^1 is absent.

22. The compound of any one of claims 16-18, or a pharmaceutically acceptable salt or tautomer thereof, wherein R^{28a} is $-A^1-L^1-A^2-B$.

23. The compound of any one of claims 16-18, or a pharmaceutically acceptable salt or tautomer thereof, wherein R^{28a} is $-A^1-A^2-B$.


24. The compound of any one of claims 16-18, or a pharmaceutically acceptable salt or tautomer thereof, wherein R^{28a} is $-L^2-A^1-L^1-A^2-L^3-B$, wherein L^2 and A^1 are absent.

25. The compound of any one of claims 16-18, or a pharmaceutically acceptable salt or tautomer thereof, wherein R^{28a} is $-L^2-A^1-L^1-A^2-L^3-B$, wherein L^2 is absent.

26. The compound of any one of claims 16-18, or a pharmaceutically acceptable salt or tautomer thereof, wherein R^{28a} is $-L^2-A^1-L^1-A^2-L^3-B$, wherein L^3 is absent.

27. The compound of any one of claims 16-18, or a pharmaceutically acceptable salt or tautomer thereof, wherein R^{28a} is $-O-(C_1-C_6)\text{alkyl}$ or $-O-(C_6-C_{10})\text{aryl}$; wherein the aryl of $-O-(C_6-C_{10})\text{aryl}$ is unsubstituted or substituted with 1-5 substituents selected from $-\text{NO}_2$ and halogen.

28. The compound of claim 1, or a pharmaceutically acceptable salt or tautomer thereof, wherein the compound is represented by the structure of Formula (I-32b):



wherein R^{32} is $-O-C(=Z^1)-R^{32a}$.

29. The compound of claim 28, or a pharmaceutically acceptable salt or tautomer thereof, wherein Z^1 is O.

30. The compound of claim 28, or a pharmaceutically acceptable salt or tautomer thereof, wherein Z^1 is S.

31. The compound of any one of claims 28-30, or a pharmaceutically acceptable salt or tautomer thereof, wherein R^{32a} is $-A^1-L^1-A^2-B$, wherein A^1 and A^2 are absent.

32. The compound of any one of claims 28-30, or a pharmaceutically acceptable salt or tautomer thereof, wherein R^{32a} is $-A^1-L^1-A^2-B$, wherein A^2 is absent.

33. The compound of any one of claims 28-30, or a pharmaceutically acceptable salt or tautomer thereof, wherein R^{32a} is $-A^1-L^1-A^2-B$, wherein A^1 is absent.

34. The compound of any one of claims 28-30, or a pharmaceutically acceptable salt or tautomer thereof, wherein R^{32a} is $-A^1-L^1-A^2-B$.

35. The compound of any one of claims 28-30, or a pharmaceutically acceptable salt or tautomer thereof, wherein R^{32a} is $-A^1-A^2-B$.

36. The compound of any one of claims 28-30, or a pharmaceutically acceptable salt or tautomer thereof, wherein R^{32a} is $-L^2-A^1-L^1-A^2-L^3-B$, wherein L^2 and A^1 are absent.


37. The compound of any one of claims 28-30, or a pharmaceutically acceptable salt or tautomer thereof, wherein R^{32a} is $-L^2-A^1-L^1-A^2-L^3-B$, wherein L^2 is absent.

38. The compound of any one of claims 28-30, or a pharmaceutically acceptable salt or tautomer thereof, wherein R^{32a} is $-L^2-A^1-L^1-A^2-L^3-B$, wherein L^3 is absent.


39. The compound of any one of claims 28-30, or a pharmaceutically acceptable salt or tautomer thereof, or a pharmaceutically acceptable salt or tautomer thereof, wherein

R^{32a} is $-O-(C_1-C_6)\text{alkyl}$ or $-O-(C_6-C_{10})\text{aryl}$; wherein the aryl of $-O-(C_6-C_{10})\text{aryl}$ is unsubstituted or substituted with 1-5 substituents selected from $-\text{NO}_2$ and halogen.


40. The compound of any one of claims 1-10, 12-22, 24-35, and 36-39, or a pharmaceutically acceptable salt or tautomer thereof, wherein L^1 is




41. The compound of any one of claims 1-10, 12-22, 24-35, and 36-39, or a pharmaceutically acceptable salt or tautomer thereof, wherein L^1 is




42. The compound of any one of claims 1-10, 12-22, 24-35, and 36-39, or a pharmaceutically acceptable salt or tautomer thereof, wherein L^1 is




43. The compound of any one of claims 1-10, 12-22, 24-35, and 36-39, or a pharmaceutically acceptable salt or tautomer thereof, wherein L^1 is




44. The compound of any one of claims 1-10, 12-22, 24-35, and 36-39, or a pharmaceutically acceptable salt or tautomer thereof, wherein L^1 is




45. The compound of any one of claims 1-10, 12-22, 24-35, and 36-39, or a pharmaceutically acceptable salt or tautomer thereof, wherein L^1 is



46. The compound of any one of claims 1-6, 12-18, 24-30, and 36-45, or a pharmaceutically acceptable salt or tautomer thereof, wherein L^2 is




47. The compound of any one of claims 1-6, 12-18, 24-30, and 36-45, or a pharmaceutically acceptable salt or tautomer thereof, wherein L^3 is




48. The compound of any one of claims 1-7, 9, 12, 16-19, 21, 24, 28-31, 33, 36, and 39-45, or a pharmaceutically acceptable salt or tautomer thereof, wherein A^1 is absent.


49. The compound of any one of claims 1-6, 8, 10-11, 13-18, 20, 22-23, 25-30, 32, 34-35, and 37-45, or a pharmaceutically acceptable salt or tautomer thereof, wherein A^1 is




50. The compound of any one of claims 1-6, 8, 10-11, 13-18, 20, 22-23, 25-30, 32, 34-35, and 37-45, or a pharmaceutically acceptable salt or tautomer thereof, wherein A^1 is




51. The compound of any one of claims 1-6, 8, 10-11, 13-18, 20, 22-23, 25-30, 32, 34-35, and 37-45, or a pharmaceutically acceptable salt or tautomer thereof, wherein A^1 is




52. The compound of any one of claims 1-6, 8, 10-11, 13-18, 20, 22-23, 25-30, 32, 34-35, and 37-45, or a pharmaceutically acceptable salt or tautomer thereof, wherein A^1 is




53. The compound of any one of claims 1-6, 8, 10-11, 13-18, 20, 22-23, 25-30, 32, 34-35, and 37-45, or a pharmaceutically acceptable salt or tautomer thereof, wherein A¹ is




54. The compound of any one of claims 1-6, 8, 10-11, 13-18, 20, 22-23, 25-30, 32, 34-35, and 37-45, or a pharmaceutically acceptable salt or tautomer thereof, wherein A¹ is



55. The compound of any one of claims 1-6, 8, 10-11, 13-18, 20, 22-23, 25-30, 32, 34-35, and 37-45, or a pharmaceutically acceptable salt or tautomer thereof, wherein A¹ is




56. The compound of any one of claims 1-6, 8, 10-11, 13-18, 20, 22-23, 25-30, 32, 34-35, and 37-45, or a pharmaceutically acceptable salt or tautomer thereof, wherein A¹ is




57. The compound of any one of claims 1-8, 15-20, 27-32, and 39-45, or a pharmaceutically acceptable salt or tautomer thereof, wherein A² is absent.

58. The compound of any one of claims 1-6, 9-18, 21-30, and 33-45, or a pharmaceutically acceptable salt or tautomer thereof, wherein A² is




59. The compound of any one of claims 1-6, 9-18, 21-30, and 33-45, or a pharmaceutically acceptable salt or tautomer thereof, wherein A^2 is




60. The compound of any one of claims 1-6, 9-18, 21-30, and 33-45, or a

pharmaceutically acceptable salt or tautomer thereof, wherein A^2 is




61. The compound of any one of claims 1-6, 9-18, 21-30, and 33-45, or a

pharmaceutically acceptable salt or tautomer thereof, wherein A^2 is




62. The compound of any one of claims 1-6, 9-18, 21-30, and 33-45, or a

pharmaceutically acceptable salt or tautomer thereof, wherein A^2 is




63. The compound of any one of claims 1-6, 9-18, 21-30, and 33-45, or a

pharmaceutically acceptable salt or tautomer thereof, wherein A^2 is




64. The compound of any one of claims 1-6, 9-18, 21-30, and 33-45, or a


pharmaceutically acceptable salt or tautomer thereof, wherein A^2 is



65. The compound of any one of claims 1-6, 9-18, 21-30, and 33-45, or a pharmaceutically acceptable salt or tautomer thereof, wherein A^2 is




66. The compound of any one of claims 1-65, or a pharmaceutically acceptable



salt or tautomer thereof, wherein B is


67. The compound of any one of claims 1-65, or a pharmaceutically acceptable



salt or tautomer thereof, wherein B is

68. The compound of any one of claims 1-65, or a pharmaceutically acceptable salt or tautomer thereof, wherein B^1 is $^{-\xi}-NR^3-(C(R^3)_2)_n-$.


69. The compound of any one of claims 1-65, or a pharmaceutically acceptable




salt or tautomer thereof, wherein B^1 is

70. The compound of any one of claims 1-69, or a pharmaceutically acceptable salt or tautomer thereof, wherein R^4 is 5-12 membered heteroaryl, optionally substituted with $-N(R^3)_2$, $-OR^3$, halogen, (C_1-C_6) alkyl, $-(C_1-C_6)$ alkylene-heteroaryl, $-(C_1-C_6)$ alkylene-CN, or $-C(O)NR^3$ -heteroaryl.


71. The compound of any one of claims 1-70, or a pharmaceutically acceptable salt or tautomer thereof, or a pharmaceutically acceptable salt or tautomer thereof, wherein compound has the following formula:




72. A compound selected from the group consisting of:




Example 1




Example 2




Example 3




Example 4




Example 5




Example 6




Example 7




Example 8




Example 9




Example 10




Example 11




Example 12




Example 13




Example 14




Example 15




Example 16




Example 17




Example 18




Example 19




Example 20




Example 21




Example 22




Example 23




Example 24




Example 25




Example 26




Example 27




Example 28




Example 29




Example 30




Example 31




Example 32




Example 33




Example 34




Example 35




Example 36




Example 37




Example 38




Example 39




Example 40




Example 41




Example 42




Example 43




Example 44




Example 45




Example 46




Example 47




Example 48




Example 49




Example 50




Example 51




Example 52




Example 53




Example 54




Example 55




Example 56




Example 57




Example 58




Example 59




Example 60




Example 61




Example 62




Example 63




Example 64




Example 65




Example 66




Example 67



Example 68









Example 69




or a pharmaceutically acceptable salt or isomer thereof.


73. A compound selected from the group consisting of:









Example 80




Example 81




Example 82




Example 83




Example 84




Example 85




Example 86




Example 87




Example 88




Example 89




Example 90




Example 91




Example 92




Example 93




Example 94




Example 95




Example 96




Example 97




Example 98




Example 99




Example 100




Example 101




Example 102




Example 103




Example 104




Example 105




Example 106




Example 107




Example 108




Example 109




Example 110




Example 111




Example 112




Example 113




Example 114




Example 115




Example 116




Example 117




Example 118




Example 119




Example 120




Example 121




Example 122




Example 123




Example 124




Example 125




Example 126




Example 127




Example 128




Example 129




Example 130




Example 131




Example 132



Example 133




Example 134




Example 135

or a pharmaceutically acceptable salt or isomer thereof.


74. A compound selected from the group consisting of:




Example 136




Example 137




Example 138




Example 139




Example 140




Example 141




Example 142




Example 143




Example 144




Example 145




Example 146




Example 147




Example 148







Example 149



Example 150



Example 151



Example 152

or a pharmaceutically acceptable salt or isomer thereof.

75. A pharmaceutical composition comprising a compound of any one of claims 1-74, or a pharmaceutically acceptable salt thereof, and at least one of a pharmaceutically acceptable carrier, diluent, or excipient.

76. A method of treating a disease or disorder mediated by mTOR comprising administering to the subject suffering from or susceptible to developing a disease or disorder

mediated by mTOR a therapeutically effective amount of one or more compounds of any one of claims 1-74, or a pharmaceutically acceptable salt thereof.

77. A method of preventing a disease or disorder mediated by mTOR comprising administering to the subject suffering from or susceptible to developing a disease or disorder mediated by mTOR a therapeutically effective amount of one or more compounds of any one of claims 1-74, or a pharmaceutically acceptable salt thereof.

78. A method of reducing the risk of a disease or disorder mediated by mTOR comprising administering to the subject suffering from or susceptible to developing a disease or disorder mediated by mTOR a therapeutically effective amount of one or more compounds of any one of claims 1-74, or a pharmaceutically acceptable salt thereof.

79. The method of any one of claims 76-78, wherein the disease is cancer or an immune-mediated disease.

80. The method of claim 79, wherein the cancer is selected from brain and neurovascular tumors, head and neck cancers, breast cancer, lung cancer, mesothelioma, lymphoid cancer, stomach cancer, kidney cancer, renal carcinoma, liver cancer, ovarian cancer, ovary endometriosis, testicular cancer, gastrointestinal cancer, prostate cancer, glioblastoma, skin cancer, melanoma, neuro cancers, spleen cancers, pancreatic cancers, blood proliferative disorders, lymphoma, leukemia, endometrial cancer, cervical cancer, vulva cancer, prostate cancer, penile cancer, bone cancers, muscle cancers, soft tissue cancers, intestinal or rectal cancer, anal cancer, bladder cancer, bile duct cancer, ocular cancer, gastrointestinal stromal tumors, and neuro-endocrine tumors.

81. The method of claim 79, wherein the immune-mediated disease is selected from resistance by transplantation of heart, kidney, liver, medulla ossium, skin, cornea, lung, pancreas, intestinum tenue, limb, muscle, nerves, duodenum, small-bowel, or pancreatic-islet-cell; graft-versus-host diseases brought about by medulla ossium transplantation; rheumatoid arthritis, systemic lupus erythematosus, Hashimoto's thyroiditis, multiple sclerosis, myasthenia gravis, type I diabetes, uveitis, allergic encephalomyelitis, and glomerulonephritis.

82. A method of treating cancer comprising administering to the subject a therapeutically effective amount of one or more compounds of any one of claims 1-74, or a pharmaceutically acceptable salt thereof.

83. The method of claim 82, wherein the cancer is selected from brain and neurovascular tumors, head and neck cancers, breast cancer, lung cancer, mesothelioma, lymphoid cancer, stomach cancer, kidney cancer, renal carcinoma, liver cancer, ovarian

cancer, ovary endometriosis, testicular cancer, gastrointestinal cancer, prostate cancer, glioblastoma, skin cancer, melanoma, neuro cancers, spleen cancers, pancreatic cancers, blood proliferative disorders, lymphoma, leukemia, endometrial cancer, cervical cancer, vulva cancer, prostate cancer, penile cancer, bone cancers, muscle cancers, soft tissue cancers, intestinal or rectal cancer, anal cancer, bladder cancer, bile duct cancer, ocular cancer, gastrointestinal stromal tumors, and neuro-endocrine tumors.

84. A method of treating an immune-mediated disease comprising administering to the subject a therapeutically effective amount of one or more compounds of any one of claims 1-74, or a pharmaceutically acceptable salt thereof.

85. The method of claim 84, wherein the immune-mediated disease is selected from resistance by transplantation of heart, kidney, liver, medulla ossium, skin, cornea, lung, pancreas, intestinum tenue, limb, muscle, nerves, duodenum, small-bowel, or pancreatic-islet-cell; graft-versus-host diseases brought about by medulla ossium transplantation; rheumatoid arthritis, systemic lupus erythematosus, Hashimoto's thyroiditis, multiple sclerosis, myasthenia gravis, type I diabetes, uveitis, allergic encephalomyelitis, and glomerulonephritis.

86. A method of treating an age related condition comprising administering to the subject a therapeutically effective amount of one or more compounds of any one of claims 1-74, or a pharmaceutically acceptable salt thereof.

87. The method of claim 86, wherein the age related condition is selected from sarcopenia, skin atrophy, muscle wasting, brain atrophy, atherosclerosis, arteriosclerosis, pulmonary emphysema, osteoporosis, osteoarthritis, high blood pressure, erectile dysfunction, dementia, Huntington's disease, Alzheimer's disease, cataracts, age-related macular degeneration, prostate cancer, stroke, diminished life expectancy, impaired kidney function, and age-related hearing loss, aging-related mobility disability (e.g., frailty), cognitive decline, age-related dementia, memory impairment, tendon stiffness, heart dysfunction such as cardiac hypertrophy and systolic and diastolic dysfunction, immunosenescence, cancer, obesity, and diabetes.

88. A compound of any one of claims 1-74, or a pharmaceutically acceptable salt thereof, for use in treating, preventing, or reducing the risk of a disease or condition mediated by mTOR.

89. Use of a compound of any of claims 1-74, or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament for treating, preventing, or reducing the risk of a disease or disorder mediated by mTOR.

90. A compound of any one of claims 1-74, or a pharmaceutically acceptable salt thereof, for use in treating cancer.

91. Use of a compound of any one of claims 1-74, or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament for treating cancer.

92. A compound of any one of claims 1-74, or a pharmaceutically acceptable salt thereof, for use in treating an immune-mediated disease.

93. Use of a compound of any one of claims 1-74, or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament for treating an immune-mediated disease.

94. A compound of any one of claims 1-74, or a pharmaceutically acceptable salt thereof, for use in treating an age related condition.

95. Use of a compound of any one of claims 1-74, or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament for treating an age related condition.

INTERNATIONAL SEARCH REPORT

International application No

PCT/US2019/029737

A. CLASSIFICATION OF SUBJECT MATTER

INV. C07D498/18 C07D519/00 A61P35/00
ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

C07D A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 2016/279108 A1 (FORREST MARCUS LAIRD [US] ET AL) 29 September 2016 (2016-09-29) claims and examples -----	1,2, 4-15, 40-71, 75-95
Y	WO 2009/131631 A1 (PONIARD PHARMACEUTICALS INC [US]; SUN CONNIE L [US]; LI XIAOYUAN [US]) 29 October 2009 (2009-10-29) claims and examples -----	1,2, 4-15, 40-71, 75-95
A		72-74
X	WO 2007/068462 A2 (NOVARTIS AG [CH]; NOVARTIS PHARMA GMBH [AT]; MOORE HENRIETTA [AT]) 21 June 2007 (2007-06-21) p. 2, 1. 7-p. 3, 1. 9 ----- -/-	1,2,5,15



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

17 September 2019

Date of mailing of the international search report

23/09/2019

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Sahagún Krause, H

INTERNATIONAL SEARCH REPORT

International application No

PCT/US2019/029737

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 5 480 989 A (KAO WENLING [US] ET AL) 2 January 1996 (1996-01-02)	1,2, 4-15, 40-71, 75-95
A	claims and examples -----	72-74
X,P	WO 2018/204416 A1 (REVOLUTION MEDICINES INC [US]; SEMKO CHRISTOPHER [US] ET AL.) 8 November 2018 (2018-11-08)	1,2,16, 17,28,29
Y,P	claims and examples, particularly starting compound in [00488]	4-15, 18-27, 30-71, 75-95
A,P		72-74
Y	VANESSA S. RODRIK-OUTMEZGUINE ET AL: "Overcoming mTOR resistance mutations with a new-generation mTOR inhibitor", NATURE, vol. 534, no. 7606, 18 May 2016 (2016-05-18), pages 272-276, XP055499733, London ISSN: 0028-0836, DOI: 10.1038/nature17963	1,2, 4-15, 40-71, 75-95
A	whole document particularly Fig. 3 and extended Data Figure 3(a) -----	72-74
X	WO 94/25072 A1 (AMERICAN HOME PROD [US]) 10 November 1994 (1994-11-10)	1,2,4,5, 15, 40-71, 75-95
	claims and examples particularly examples 19 and 20 -----	
X	WO 94/11380 A1 (AMERICAN HOME PROD [US]) 26 May 1994 (1994-05-26)	1,2,4,5, 15, 40-71, 75-95
	claims 1, 12 and examples +A, 1B, 2A, 2B, 3, 4A, 4B, 5, 7a, 7b, 7e -----	
X	WO 95/14023 A1 (ABBOTT LAB [US]) 26 May 1995 (1995-05-26)	1-5, 15-17, 28,29, 40-71, 75-95
Y	claims and examples	6-14, 18-27, 30-39
A	----- -/--	72-74

INTERNATIONAL SEARCH REPORT

International application No

PCT/US2019/029737

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 6 342 507 B1 (NAICKER SELVARAJ [CA] ET AL) 29 January 2002 (2002-01-29) fig. 5	1,2,4,5, 15, 40-71, 75-95
X	WO 2009/122176 A2 (BIOINTERACTIONS LTD [GB]; RHODES ALAN [GB] ET AL.) 8 October 2009 (2009-10-08)	1,2,4,5, 15, 40-71, 75-95
Y	claims and examples	6-14
A,P	WO 2019/064182 A1 (NOVARTIS AG [CH]) 4 April 2019 (2019-04-04) claim 1 and examples; claim 12	1,3-15, 40-95
X	WO 2004/101583 A1 (ISOTECHNIKA INTERNAT INC [BB]; ABEL MARK [CA] ET AL.) 25 November 2004 (2004-11-25) 31-0- (4-nitrophenyloxycarbonyl) rapamycin in p. 39	1,2,16, 17,75-95
X	US 5 258 389 A (GOULET MARK [US] ET AL) 2 November 1993 (1993-11-02) claims 1 and 9-12	1,2,16, 17
X	JOERG A. KALLEN ET AL: "X-ray Crystal Structure of 28- O -Methylrapamycin complexed with FKBP12: Is the Cyclohexyl Moiety Part of the Effector Domain of Rapamycin?", JOURNAL OF THE AMERICAN CHEMICAL SOCIETY, vol. 118, no. 25, 1 January 1996 (1996-01-01), pages 5857-5861, XP055620146, US ISSN: 0002-7863, DOI: 10.1021/ja954328h 28-0-methylrapamycin	1,2,16, 17
X	NELSON F C ET AL: "Manipulation of the C922)-C(27) region of rapamycin: stability issues and biological implications", BIOORGANIC & MEDICINAL CHEMISTRY LETTERS, PERGAMON, AMSTERDAM, NL, vol. 9, no. 2, 18 January 1999 (1999-01-18), pages 295-300, XP004152620, ISSN: 0960-894X, DOI: 10.1016/S0960-894X(98)00735-5 compound 7	1,3,16, 17,28,29
	----- -/--	

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2019/029737

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	TOBIN STROM ET AL: "Structural identification of SAR-943 metabolites generated by human liver microsomes in vitro using mass spectrometry in combination with analysis of fragmentation patterns", JOURNAL OF MASS SPECTROMETRY., vol. 46, no. 7, 1 July 2011 (2011-07-01), pages 615-624, XP055620244, GB ISSN: 1076-5174, DOI: 10.1002/jms.1930 SAR-943 and metabolites thereof -----	1,2,28, 29
A	WO 95/04738 A1 (AMERICAN HOME PROD [US]) 16 February 1995 (1995-02-16) the whole document -----	1-95

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US2019/029737

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☒ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fees, this Authority did not invite payment of additional fees.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- ☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- ☒ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. claims: 1, 2, 4-15, 40-95(all partially)

compounds of formula (Ic) wherein R32 is =O and R40 is
-C(=Z1)-R40a

2. claims: 1, 3-15, 40-95(all partially)

compounds of formula (Ic) wherein R32 is other than =O and
R40 is -C(=Z1)-R40a

3. claims: 16-39(completely); 1-3, 40-70, 73, 75-95(partially)

compounds of formula (Ic) wherein R40 is -H

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2019/029737

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 2016279108	A1	29-09-2016	NONE
WO 2009131631	A1	29-10-2009	US 2011098241 A1 28-04-2011 WO 2009131631 A1 29-10-2009
WO 2007068462	A2	21-06-2007	NONE
US 5480989	A	02-01-1996	US 5480989 A 02-01-1996 US 5508399 A 16-04-1996 US 5530007 A 25-06-1996 US 5530121 A 25-06-1996 US 5559120 A 24-09-1996
WO 2018204416	A1	08-11-2018	NONE
WO 9425072	A1	10-11-1994	AU 6711994 A 21-11-1994 WO 9425072 A1 10-11-1994
WO 9411380	A1	26-05-1994	AT 151769 T 15-05-1997 AU 5322094 A 08-06-1994 CA 2149669 A1 26-05-1994 DE 69309954 D1 22-05-1997 DE 69309954 T2 24-07-1997 DK 0669923 T3 25-08-1997 EP 0669923 A1 06-09-1995 ES 2100572 T3 16-06-1997 GR 3023801 T3 30-09-1997 HK 109197 A 22-08-1997 JP H08503210 A 09-04-1996 KR 950704330 A 17-11-1995 SG 47595 A1 17-04-1998 US 5260300 A 09-11-1993 WO 9411380 A1 26-05-1994
WO 9514023	A1	26-05-1995	CA 2175215 A1 26-05-1995 EP 0729471 A1 04-09-1996 JP 4105761 B2 25-06-2008 JP H09505299 A 27-05-1997 US 5583139 A 10-12-1996 US 5672605 A 30-09-1997 WO 9514023 A1 26-05-1995
US 6342507	B1	29-01-2002	US 6342507 B1 29-01-2002 US 2002028827 A1 07-03-2002 US 2003139440 A1 24-07-2003 US 2004198762 A1 07-10-2004
WO 2009122176	A2	08-10-2009	US 2009253733 A1 08-10-2009 WO 2009122176 A2 08-10-2009
WO 2019064182	A1	04-04-2019	TW 201920189 A 01-06-2019 US 2019092788 A1 28-03-2019 UY 37900 A 30-04-2019 WO 2019064182 A1 04-04-2019
WO 2004101583	A1	25-11-2004	AT 367396 T 15-08-2007 AU 2004238408 A1 25-11-2004 BR PI0411171 A 18-07-2006

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2019/029737

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
		CA 2525907 A1	25-11-2004
		DE 602004007662 T2	22-11-2007
		DK 1644392 T3	12-11-2007
		EP 1644392 A1	12-04-2006
		EP 1785430 A1	16-05-2007
		ES 2286637 T3	01-12-2007
		HR P20070383 T3	31-10-2007
		JP 2007503452 A	22-02-2007
		KR 20060031603 A	12-04-2006
		MX PA05012376 A	25-05-2006
		NZ 543479 A	28-03-2008
		PT 1644392 E	03-08-2007
		TW 200511991 A	01-04-2005
		US 2004235762 A1	25-11-2004
		US 2006079466 A1	13-04-2006
		US 2007099848 A1	03-05-2007
		WO 2004101583 A1	25-11-2004

US 5258389	A	02-11-1993	NONE

WO 9504738	A1	16-02-1995	AT 163420 T 15-03-1998
		AU 676086 B2	27-02-1997
		BR 9407236 A	24-09-1996
		CA 2169277 A1	16-02-1995
		CN 1132512 A	02-10-1996
		CZ 291205 B6	15-01-2003
		DE 69408678 D1	02-04-1998
		DE 69408678 T2	18-06-1998
		DK 0713490 T3	11-05-1998
		EP 0713490 A1	29-05-1996
		ES 2115255 T3	16-06-1998
		GR 3026586 T3	31-07-1998
		JP H09501436 A	10-02-1997
		NZ 271822 A	27-05-1998
		PL 312991 A1	27-05-1996
		RU 2152946 C1	20-07-2000
		SG 47977 A1	17-04-1998
		US 5387680 A	07-02-1995
		WO 9504738 A1	16-02-1995
