



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁶ : A61K 38/00</p>	<p>A2</p>	<p>(11) International Publication Number: WO 98/06415</p> <p>(43) International Publication Date: 19 February 1998 (19.02.98)</p>
<p>(21) International Application Number: PCT/US97/13810</p> <p>(22) International Filing Date: 7 August 1997 (07.08.97)</p> <p>(30) Priority Data: 08/694,843 9 August 1996 (09.08.96) US</p> <p>(71) Applicant: XOMA CORPORATION [US/US]; 2910 Seventh Street, Berkeley, CA 94710-2737 (US).</p> <p>(72) Inventor: LAMBERT, Lewis, H., Jr.; 45928 Omega Drive, Fremont, CA 94539 (US).</p> <p>(74) Agent: BORUN, Michael, F.; Marshall, O'Toole, Gerstein, Murray & Borun, 6300 Sears Tower, 233 S. Wacker Drive, Chicago, IL 60606-6402 (US).</p>		<p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>Without international search report and to be republished upon receipt of that report.</i></p>
<p>(54) Title: ANTI-CHLAMYDIAL METHODS AND MATERIALS</p>		
<p>(57) Abstract</p> <p>The present invention relates to methods for treating chlamydial infection comprising administering to a subject suffering from a chlamydial infection a bactericidal/permeability-inducing (BPI) protein product.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakistan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

- 1 -

ANTI-CHLAMYDIAL METHODS AND MATERIALS

BACKGROUND OF THE INVENTION

The present invention relates generally to methods of treating chlamydial infections by administration of bactericidal/permeability-increasing (BPI) protein products.

BPI is a protein isolated from the granules of mammalian polymorphonuclear leukocytes (PMNs or neutrophils), which are blood cells essential in the defense against invading microorganisms. Human BPI protein has been isolated from PMNs by acid extraction combined with either ion exchange chromatography [Elsbach, *J. Biol. Chem.*, 254:11000 (1979)] or *E. coli* affinity chromatography [Weiss, et al., *Blood*, 69:652 (1987)]. BPI obtained in such a manner is referred to herein as natural BPI and has been shown to have potent bactericidal activity against a broad spectrum of gram-negative bacteria. The molecular weight of human BPI is approximately 55,000 daltons (55 kD). The amino acid sequence of the entire human BPI protein and the nucleic acid sequence of DNA encoding the protein have been reported in Figure 1 of Gray et al., *J. Biol. Chem.*, 264:9505 (1989), incorporated herein by reference. The Gray et al. amino acid sequence is set out in SEQ ID NO: 1 hereto.

BPI is a strongly cationic protein. The N-terminal half of BPI accounts for the high net positive charge; the C-terminal half of the molecule has a net charge of -3. [Elsbach and Weiss (1981), *supra*.] A proteolytic N-terminal fragment of BPI having a molecular weight of about 25 kD has an amphipathic character, containing alternating hydrophobic and hydrophilic regions. This N-terminal fragment of human BPI possesses the anti-bacterial efficacy of the naturally-derived 55 kD human BPI holoprotein. [Ooi et al., *J. Bio. Chem.*, 262: 14891-14894 (1987)]. In contrast to the N-terminal portion, the C-terminal region of the isolated human BPI protein displays only slightly detectable anti-bacterial activity against gram-negative organisms. [Ooi et al., *J. Exp. Med.*, 174:649

- 2 -

(1991).] An N-terminal BPI fragment of approximately 23 kD, referred to as "rBPI₂₃," has been produced by recombinant means and also retains anti-bacterial activity against gram-negative organisms. Gazzano-Santoro et al., *Infect. Immun.* 60:4754-4761 (1992).

5 The bactericidal effect of BPI has been reported to be highly specific to gram-negative species, e.g., in Elsbach and Weiss, *Inflammation: Basic Principles and Clinical Correlates*, eds. Gallin et al., Chapter 30, Raven Press, Ltd. (1992). This reported target cell specificity was believed to be the result of the strong attraction of BPI for

10 lipopolysaccharide (LPS), which is unique to the outer membrane (or envelope) of gram-negative organisms. Although BPI was commonly thought to be non-toxic for other microorganisms, including yeast, and for higher eukaryotic cells, it has recently been discovered that BPI protein products, as defined *infra*, exhibit activity against gram-positive bacteria,

15 mycoplasma, mycobacteria, fungi, and protozoa. [See allowed, co-owned, co-pending U.S. Patent Application Serial No. 08/372,783 filed January 13, 1995, the disclosures of which are incorporated herein by reference; co-owned, co-pending U.S. Patent Application Serial No. 08/626,646, the disclosures of which are incorporated herein by reference; co-owned, co-

20 pending U.S. Patent Application Serial No. 08/372,105, the disclosures of which are incorporated herein by reference; and co-owned, co-pending U.S. Patent Application Serial No. 08/273,470, the disclosures of which are incorporated herein by reference.] It has also been discovered that BPI protein products have the ability to enhance the activity of antibiotics

25 against bacteria. [See U.S. Patent No. 5,523,288, the disclosures of which are incorporated herein by reference, and allowed, co-owned, co-pending U.S. Patent Application Serial No. 08/372,783.]

 The precise mechanism by which BPI kills gram-negative bacteria is not yet completely elucidated, but it is believed that BPI must

30 first bind to the surface of the bacteria through electrostatic and

- 3 -

hydrophobic interactions between the cationic BPI protein and negatively charged sites on LPS. LPS has been referred to as "endotoxin" because of the potent inflammatory response that it stimulates, i.e., the release of mediators by host inflammatory cells which may ultimately result in irreversible endotoxic shock. BPI binds to lipid A, reported to be the most toxic and most biologically active component of LPS.

In susceptible gram-negative bacteria, BPI binding is thought to disrupt LPS structure, leading to activation of bacterial enzymes that degrade phospholipids and peptidoglycans, altering the permeability of the cell's outer membrane, and initiating events that ultimately lead to cell death. [Elsbach and Weiss (1992), *supra*]. BPI is thought to act in two stages. The first is a sublethal stage that is characterized by immediate growth arrest, permeabilization of the outer membrane and selective activation of bacterial enzymes that hydrolyze phospholipids and peptidoglycans. Bacteria at this stage can be rescued by growth in serum albumin supplemented media [Mannion et al., *J. Clin. Invest.*, 85:853-860 (1990)]. The second stage, defined by growth inhibition that cannot be reversed by serum albumin, occurs after prolonged exposure of the bacteria to BPI and is characterized by extensive physiologic and structural changes, including apparent damage to the inner cytoplasmic membrane.

Initial binding of BPI to LPS leads to organizational changes that probably result from binding to the anionic groups of LPS, which normally stabilize the outer membrane through binding of Mg^{++} and Ca^{++} . Attachment of BPI to the outer membrane of gram-negative bacteria produces rapid permeabilization of the outer membrane to hydrophobic agents such as actinomycin D. Binding of BPI and subsequent gram-negative bacterial killing depends, at least in part, upon the LPS polysaccharide chain length, with long O-chain bearing, "smooth" organisms being more resistant to BPI bactericidal effects than short O-chain bearing, "rough" organisms [Weiss et al., *J. Clin. Invest.* 65: 619-

- 4 -

628 (1980)]. This first stage of BPI action, permeabilization of the gram-negative outer envelope, is reversible upon dissociation of the BPI, a process requiring high concentrations of divalent cations and synthesis of new LPS [Weiss et al., *J. Immunol.* 132: 3109-3115 (1984)]. Loss of
5 gram-negative bacterial viability, however, is not reversed by processes which restore the envelope integrity, suggesting that the bactericidal action is mediated by additional lesions induced in the target organism and which may be situated at the cytoplasmic membrane (Mannion et al., *J. Clin. Invest.* 86: 631-641 (1990)). Specific investigation of this possibility has
10 shown that on a molar basis BPI is at least as inhibitory of cytoplasmic membrane vesicle function as polymyxin B (In't Veld et al., *Infection and Immunity* 56: 1203-1208 (1988)) but the exact mechanism as well as the relevance of such vesicles to studies of intact organisms has not yet been elucidated.

15 Chlamydia are nonmotile, gram-negative, obligate intracellular bacteria that have unusual biological properties which phylogenetically distinguish them from other families of bacteria. Chlamydiae are presently placed in their own order, the *Chlamydiales*, family *Chlamydiaceae*, with one genus, *Chlamydia*. [Schachter and
20 Stamm, *Chlamydia*, in *Manual of Clinical Microbiology*, pages 669-677, American Society for Microbiology, Washington, DC (1995).] There are four species, *Chlamydia trachomatis*, *C. pneumoniae*, *C. psittaci* and *C. pecorum*, which cause a wide spectrum of human diseases. In developing countries, *C. trachomatis* causes trachoma, the world's leading cause of
25 preventable blindness. Over 150 million children have active trachoma, and over 6 million people are currently blind from this disease. In industrialized countries, *C. trachomatis* is the most prevalent sexually transmitted disease, causing urethritis, cervicitis, epididymitis, ectopic pregnancy and pelvic inflammatory disease. Last year alone, an estimated
30 300 million people contracted sexually transmitted chlamydial infections.

- 5 -

Among the 250,000 cases of pelvic inflammatory disease per year in the United States, approximately 25,000 women are rendered infertile each year. Neonatal *C. trachomatis* infections, contracted at birth from infected mothers, cause hundreds of thousands of conjunctivitis cases per year, of which about half of these infected infants develop pneumonia. Recently, *C. pneumoniae* has been implicated as a common cause of epidemic human pneumonitis. Members of the genus are not only important human pathogens, but also cause significant morbidity in other mammals and birds. Thus, chlamydia are one of the most ubiquitous pathogens in the animal kingdom. [Zhang et al., *Cell*, 69:861-869 (1992).]

Their unique developmental cycle differentiates them from all other microorganisms. They are obligate intracellular parasites that are unable to synthesize ATP, and thus depend on the host cells' energy to survive. Unlike viruses, they always contain both DNA and RNA, divide by binary fission, contain ribosomes, and can synthesize proteins. Chlamydia have cell walls similar in structure to those of gram-negative bacteria, and all members of the genus carry a unique LPS-like antigen, termed complement fixation (CF) antigen, that may be analogous to the LPS of certain gram-negative bacteria. [Schachter and Stamm, *supra*.]

Chlamydia also carry a major outer membrane protein (MOMP) that contains both species and subspecies-specific antigens.

The infectious form of chlamydia is the elementary body (EB), which infects mammalian cells by attaching to the host cell and entering in a host-derived phagocytic vesicle (endosome), within which the entire growth cycle is completed. The target host cell *in vivo* is typically the columnar epithelial cell, and the primary mode of entry is believed to be receptor-mediated endocytosis. Once the EB has entered the cell, it reorganizes into a reticulate body (RB) that is larger than the EB and metabolically active, synthesizing DNA, RNA and proteins. The EBs are specifically adapted for extracellular survival, while the metabolically

- 6 -

active RBs do not survive well outside the host cell and seems adapted for an intracellular milieu. After approximately 8 hours, the RBs begin dividing by binary fission. As they replicate within the endosomes of host cells, they form characteristic intracellular inclusions that can be seen by light microscopy. After a period of growth and division, the RBs reorganize and condense to form infectious EBs. The developmental cycle is complete when host cell lysis or exocytosis of chlamydia occurs, releasing the EBs to initiate another cycle of infection. The length of the complete developmental cycle, as studied in cell culture models, is 48 to 72 hours and varies as a function of the infecting strain, host cell and environmental conditions. [Beatty et al., *Microbiol. Rev.*, 58(4):686-699 (1994).]

It has been demonstrated, at least for *C. trachomatis*, that attachment of the chlamydia organism to host cells is mediated by a heparan sulfate-like glycosaminoglycan (GAG) present on the surface of the chlamydia. Treatment of chlamydia with either purified heparin, heparin sulfate, or heparin receptor analogs (such as platelet factor 4 and fibronectin, both of which are known to bind heparin sulfate), inhibited the attachment and infectivity of chlamydia to host cells. Inhibition was not seen with non-heparin GAGs, such as hyaluronate, chondroitin sulfate, or keratin sulfate. Treatment of *C. trachomatis* with heparitinase reduced attachment and infectivity by greater than 90%; subsequent treatment with exogenous heparan sulfate was able to restore the ability of treated organisms to attach to host cells in a dose-dependent manner. Other GAGs such as hyaluronate, chondroitin sulfate, or keratin sulfate did not restore attachment ability. These data suggest that a heparin sulfate-like GAG mediates attachment of chlamydia to host cells by bridging mutual GAG receptors on the host cell surface and on the chlamydial outer membrane surface. [Zhang et al., *Cell*, 69:861-869 (1992).]

- 7 -

C. trachomatis is almost exclusively a human pathogen, and is responsible for trachoma, inclusion conjunctivitis, lymphogranuloma venereum (LGV), and genital tract diseases. [Schachter and Stamm, *supra.*] Within this species, serotypes A, B, Ba, and C have been associated with endemic trachoma, the most common preventable form of blindness in the world. Trachoma is a chronic inflammation of the conjunctiva and the cornea, which is not sexually transmitted. The potentially blinding sequelae of trachoma include lid distortion, trichiasis (misdirection of lashes), and entropion (inward deformation of the lid margin). These can cause corneal ulceration followed by loss of vision. Serotypes L1, L2, and L3 of *C. trachomatis* are associated with LGV. Untreated, lymphogranuloma venereum progresses through three stages, each more severe than the preceding one. The primary lesion, if present, appears on the genitals. The second stage is a bubonic state marked by regional lymphadenopathy, during which the buboes may suppurate and develop draining fistulas. Rectal strictures and lymphatic obstruction can appear in the tertiary stage. Lymphogranuloma venereum is a common problem in developing countries with tropical or subtropical climates, especially among the lower socioeconomic groups.

C. trachomatis is also the most common agent of sexually transmitted disease. In men, serotypes D through K are the major identifiable causes of nongonococcal urethritis, and also cause epididymitis, Reiter's syndrome, and proctitis. Chlamydial infections are not easily identified in men by clinical symptoms alone, because the infection may be asymptomatic and because other pathogens cause similar symptoms. Chlamydial urethritis occurs twice as frequently as gonococcal urethritis (gonorrhea) in some populations, and its incidence is on the increase. Even when *N. gonorrhoea* is shown to be present, the urethritis may be due to a dual or multiple infection involving a second organism. Concurrent *C. trachomatis* and *N. gonorrhoea* infections have been reported in about 25

percent of men with gonorrhea. Epididymitis is the most important complication of chlamydial urethritis in men. *C. trachomatis* causes one of every two cases of epididymitis in younger men in the United States, with sterility a possible result. Reiter's syndrome is another manifestation of chlamydial infection in men. It is a painful systemic illness that classically includes symptoms of urethritis, conjunctivitis and arthritis. Urethritis and arthritis are by far the most frequent combination; it appears that the chlamydial urethral infection may trigger the arthritis. *C. trachomatis* can also cause proctitis (anal inflammation), particularly in homosexual men.

In women, chlamydial infection with the sexually transmitted serotypes results in cervicitis, urethritis, endometritis, salpingitis, and proctitis; serious sequelae of salpingitis include tubal scarring, infertility, and ectopic pregnancy. Unrecognized chlamydial infections in women are common. Approximately 50 percent of women infected with chlamydia are asymptomatic. *C. trachomatis* causes mucopurulent cervicitis and the urethral syndrome, as well as endometritis and salpingitis. These upper genital tract chlamydial infections may cause sterility or predispose to ectopic pregnancies and are the gravest complications of chlamydial infections in women. Ten percent of all maternal deaths are due to ectopic pregnancies. *C. trachomatis* causes over 30 percent of the cases of mucopurulent cervicitis. As many as one-half of the women with gonococcal cervicitis have a concomitant chlamydial infection. If the gonococcal infection is treated with penicillin, the concomitant chlamydial cervicitis will continue undetected and untreated, and may progress to pelvic inflammatory disease (salpingitis), which can lead to sterility and ectopic pregnancies. *C. trachomatis* is a cause of the urethral syndrome in women. Chlamydial infections may ascend from the cervix to the endometrium, where *C. trachomatis* has been found in the epithelial lining of the uterine cavity. It is estimated that about one-half of all women with cervicitis have endometritis. Salpingitis, a major cause of ectopic

pregnancies and infertility, is the most serious complication of female genital infections. Upper abdominal pain is the predominant symptom of perihepatitis. Both *C. trachomatis* and *N. gonorrhoea* can cause perihepatitis. This condition occurs almost exclusively in women in whom
5 the infecting organisms spread to the surface of the liver from inflamed fallopian tubes.

Women infected with *C. trachomatis* may also pass the disease to their newborn as it passes through the infected birth canal. These newborns most often develop inclusion conjunctivitis or chlamydial
10 pneumonia, but may also develop vaginal, pharyngeal, or enteric infections. Though not blinding, inclusion conjunctivitis can become chronic, causing mild scarring and pannus formation if left untreated. During passage through the birth canal, up to two-thirds of babies born to mothers with chlamydial genital infections will also become infected. With
15 as many as one in ten pregnant women having chlamydial cervicitis in some parts of the world, the risk to newborns is considerable. Chlamydial pneumonia occurs in 10 percent to 20 percent of infants born to infected mothers. *C. trachomatis* is responsible for 20 percent to 60 percent of all pneumonias during the first 6 months of life.

20 *C. trachomatis* strains are sensitive to the action of tetracyclines, macrolides and sulfonamides and produce a glycogen-like material within the inclusion vacuole that stains with iodine.

C. psittaci strains infect many avian species and mammals, producing such diseases as psittacosis, ornithosis, feline pneumonitis, and
25 bovine abortion. [Schachter and Stamm, *supra*.] *C. psittaci* is ubiquitous among avian species, and infection in birds usually involves the intestinal tract. The organism is shed in the feces, contaminates the environment, and is spread by aerosol. *C. psittaci* is also common in domestic mammals. In some parts of the world, these infections have important
30 economic consequences, as *C. psittaci* is a cause of a number of systemic

- 10 -

and debilitating diseases in domestic mammals and, most important, can cause abortions. Human chlamydial infections from this agent usually result from exposure to an infected avian species, but may also occur after exposure to infected domestic mammals. This species is resistant to the
5 action of sulfonamides and produces inclusions that do not stain with iodine.

C. pneumoniae has less than 10% DNA relatedness to the other species and has pear-shaped rather than round elementary bodies (EBs). Like *C. trachomatis*, it appears to be exclusively a human pathogen
10 without an animal reservoir. *C. pneumoniae* has been identified as the cause of a variety of respiratory tract diseases and is distributed worldwide. [Schachter and Stamm, *supra*.] Infections appear to be commonly acquired in later childhood, adolescence, and early adulthood, resulting in seroprevalences of 40 to 50% in 30 to 40-year-old people. Manifestations
15 of infection include pharyngitis, bronchitis, and mild pneumonia, and transmission is primarily via respiratory secretions. In seroepidemiological studies, these infections have been linked with coronary artery disease, and their role in atherosclerosis is currently under intense scrutiny.

The role of *C. pecorum* as a pathogen is not clear, and
20 specialized reagents are required for its identification.

The recommended procedure for primary isolation of chlamydia is cell culture. Chlamydia will grow in the yolk sac of the embryonated hen egg, as well as in cell culture (with some variability). *C. trachomatis* can infect several cell lines, such as McCoy's heteroploid
25 murine cells, HeLa 229 cells, BHK-21 cells, or L-929 cells. HL cells and Hep-2 cells may be more sensitive for the recovery of *C. pneumoniae*. The most common technique involves inoculation of clinical specimens into cycloheximide-treated McCoy cells. The basic principle involves centrifugation of the inoculum onto the cell monolayer, incubation of the
30 monolayers for 48 to 72 hours, and demonstration of typical

- 11 -

intracytoplasmic inclusions by appropriate immunofluorescence, iodine or Giemsa staining procedures. Cell culture generally requires two to six days to complete because of the incubation time required.

Chlamydia may also be detected in samples by the direct
5 fluorescent antibody (DFA) test, in which slides are incubated with fluorescein-conjugated monoclonal antibodies, and fluorescing elementary bodies are detected using a fluorescent microscope. This test has approximately 80% to 90% sensitivity and 98% to 99% specificity compared with cell cultures when both tests are performed under ideal
10 circumstances. [Schachter and Stamm, *supra*.]

A number of commercially available products can detect chlamydial antigens in clinical specimens by using enzyme immunoassay (EIA) procedures. Most of these products detect chlamydial LPS, which is more soluble than MOMP. Without confirmation, the tests have a
15 specificity on the order of 97%. [Schachter and Stamm, *supra*.] Several nucleic acid probes are also commercially available. One commercially available probe test (GenProbe) utilizes DNA-RNA hybridization in an effort to increase sensitivity by detecting chlamydial RNA.

The complement fixation (CF) test is the most frequently
20 performed serological test, and measures serum level of complement-fixing antibody (antibody to the group CF antigen). It is useful for diagnosing psittacosis, in which paired acute- and convalescent-phase sera often show four-fold or greater increases in titer. The same seems to be true for many *C. pneumoniae* infections. Approximately 50% of these infections are CF-
25 positive, although it may take 24 weeks to detect seroconversion. CF testing may also be useful in diagnosing LGV, in which single-point titers greater than 1:64 are highly supportive of this clinical diagnosis. [Schachter and Stamm, *supra*.] High titers of complement-fixing antibodies are not found in chlamydial conjunctivitis or genital tract infections, and
30 therefore are not sensitive for these infections.

- 12 -

The microimmunofluorescence (micro-IF) method is a much more sensitive procedure for measuring anti-chlamydial antibodies. This indirect fluorescent antibody technique uses antigens prepared by infecting the yolk sacs of fertile chick embryos with each chlamydial serotype.

5 Serial dilutions of patient serum are added to the prepared antigens, and the level of antibody in the blood sample is determined with the use of immunofluorescence. Trachoma, inclusion conjunctivitis, and genital tract infections may be diagnosed by the micro-IF technique if appropriately timed paired sera can be obtained, but the procedure is of limited clinical
10 utility because diagnosis requires demonstration of a four-fold or greater change in antibody titer in paired specimens, and because patients with superficial genital infections such as urethritis may not have a change in titer. However, a high antibody titer in a single serum specimen from a patient with Reiter's syndrome and a high IgM titer in the serum of an
15 infant with pneumonia are helpful in establishing a diagnosis.

Strain-to-strain variation in antimicrobial susceptibility profiles and newly acquired drug resistance are both very infrequent among chlamydia. Among the drugs most active in vitro against *C. trachomatis*,
20 *C. pneumoniae*, and *C. psittaci* are the tetracyclines, such as tetracycline and doxycycline, the macrolides, such as erythromycin and azithromycin, the quinolones, such as ciprofloxacin and ofloxacin, chloramphenicol, rifampin, clindamycin and the sulfonamides. The tetracyclines and
macrolides have generally been the mainstays of therapy for infections due to chlamydia. [Schachter and Stamm, *supra*; Goodman and Gilman, *The
25 Pharmacological Basis of Therapeutics*, 9th ed., McGraw-Hill, New York, NY (1996).]

Antimicrobial susceptibility testing is infrequently performed for chlamydial infections, but may be conducted as follows. The organisms for testing are grown for at least two passages in cells cultured in
30 antibiotic-free media before being harvested. An adjusted inoculum of

- 13 -

~ 100 inclusion-forming units per microtiter well is then used to infect antibiotic-free cell monolayers. After centrifugation of the inoculum onto the monolayer, serial dilutions of the test antibiotic can be added either immediately or at various time intervals over the next 24 hours. After 48
5 hours, fluorescein-conjugated monoclonal antibodies are used to identify minimum inhibitory concentration (MIC), *i.e.*, the highest antibiotic dilution that inhibits intracellular inclusion formation. Generally, monolayers are also disrupted and further passaged to define the minimum bactericidal concentration (MBC), *i.e.*, the highest antibiotic dilution that
10 prevents viable chlamydia from being detected in passage (MBC).

SUMMARY OF THE INVENTION

The present invention provides methods of treating a subject suffering from a chlamydial infection by administering a therapeutically effective amount of a BPI protein product. This is based on the surprising
15 discoveries that BPI protein products inhibit the infectivity of chlamydia and inhibit the proliferation of chlamydia in an established intracellular infection. The BPI protein products may be administered alone or in conjunction with other known anti-chlamydial agents. When made the subject of adjunctive therapy, the administration of BPI protein products
20 may reduce the amount of non-BPI anti-chlamydial agent needed for effective therapy, thus limiting potential toxic response and/or high cost of treatment. Administration of BPI protein products may also enhance the effect of such agents, accelerate the effect of such agents, or reverse resistance of chlamydia to such agents.

25 In addition, the invention provides a method of killing or inhibiting growth of chlamydia comprising contacting the chlamydia with a BPI protein product. This method can be practiced *in vivo* or in a variety of *in vitro* uses such as use to decontaminate fluids and surfaces and to

- 14 -

sterilize surgical and other medical equipment and implantable devices, including prosthetic joints and indwelling invasive devices.

A further aspect of the invention involves use of a BPI protein product for the manufacture of a medicament for treatment of chlamydial infection. The medicament may include, in addition to a BPI protein product, other chemotherapeutic agents such as non-BPI anti-chlamydial agents.

Numerous additional aspects and advantages of the invention will become apparent to those skilled in the art upon considering the following detailed description of the invention, which describes the presently preferred embodiments thereof.

DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to the surprising discovery that a BPI protein product can be administered to treat subjects suffering from chlamydial infection, and provides methods of prophylactically or therapeutically treating such infections. Unexpectedly, BPI protein products were demonstrated to have anti-chlamydial activities, as measured, for example, by a reduction in the number of reproductive bodies seen in the host cells. A variety of chlamydial infections, including infections caused by *C. trachomatis*, *C. pneumoniae*, *C. psittaci* and *C. pecorum*, may be treated according to the invention.

The term "treating" or "treatment" as used herein encompasses both prophylactic and therapeutic treatment.

The BPI protein product may be administered systemically or topically. Systemic routes of administration include oral, intravenous, intramuscular or subcutaneous injection (including into depots for long-term release), intraocular or retrobulbar, intrathecal, intraperitoneal (e.g. by intraperitoneal lavage), transpulmonary using aerosolized or nebulized drug, or transdermal. Topical routes include administration in the form of

- 15 -

salves, creams, jellies, ophthalmic drops or ophthalmic ointments, ear drops, suppositories, such as vaginal suppositories, or irrigation fluids (for, e.g., irrigation of wounds).

When given parenterally, BPI protein product compositions
5 are generally injected in doses ranging from 1 $\mu\text{g}/\text{kg}$ to 100 mg/kg per day, preferably at doses ranging from 0.1 mg/kg to 20 mg/kg per day, and more preferably at doses ranging from 1 to 20 $\text{mg}/\text{kg}/\text{day}$. The treatment may continue by continuous infusion or intermittent injection or infusion, or a combination thereof, at the same, reduced or increased dose per day for as
10 long as determined by the treating physician. When given topically, BPI protein product compositions are generally applied in unit doses ranging from 1 $\mu\text{g}/\text{mL}$ to 1 gm/mL , and preferably in doses ranging from 1 $\mu\text{g}/\text{mL}$ to 100 mg/mL . Those skilled in the art can readily optimize effective dosages and monotherapeutic or concurrent administration regimens for BPI
15 protein product and/or other anti-chlamydial agents, as determined by good medical practice and the clinical condition of the individual patient.

The BPI protein product may be administered in conjunction with other anti-chlamydial agents presently known to be effective. Preferred anti-chlamydial agents for this purpose include the tetracyclines,
20 such as tetracycline and doxycycline, the macrolides, such as erythromycin and azithromycin, the quinolones, such as ciprofloxacin and ofloxacin, chloramphenicol, rifampin, clindamycin and the sulfonamides. Concurrent administration of BPI protein product with anti-chlamydial agents is expected to improve the therapeutic effectiveness of the anti-chlamydial
25 agents. This may occur through reducing the concentration of anti-chlamydial agent required to eradicate or inhibit chlamydial growth, e.g., replication. Because the use of some agents is limited by their systemic toxicity or prohibitive cost, lowering the concentration of anti-chlamydial agent required for therapeutic effectiveness reduces toxicity and/or cost of
30 treatment, and thus allows wider use of the agent. Concurrent

- 16 -

administration of BPI protein product and another anti-chlamydial agent may produce a more rapid or complete bactericidal or bacteriostatic effect than could be achieved with either agent alone. BPI protein product administration may reverse the resistance of chlamydia to anti-chlamydial agents. BPI protein product administration may also convert a
5 bacteriostatic agent into a bactericidal agent.

An advantage of the present invention is that the wide spectrum of activity of BPI protein products against a variety of organisms, and the use of BPI protein products as adjunctive therapy to enhance the activity of antibiotics makes BPI protein products an excellent choice for
10 treating dual or multiple infections with chlamydia and another organism, such as the gram-negative bacteria *N. gonorrhoea*. Thus, BPI protein products may be especially useful in inhibiting transmission of sexually transmitted diseases, which often involve dual gonococcal/chlamydial
15 infection. It is therefore contemplated that BPI protein products will be incorporated into contraceptive compositions and devices, *e.g.*, included in spermicidal creams or jellies, or coated on the surface of condoms.

Another advantage is the ability to treat chlamydia that have acquired resistance to known anti-chlamydial agents. A further advantage
20 of concurrent administration of BPI with an anti-chlamydial agent having undesirable side effects is the ability to reduce the amount of anti-chlamydial agent needed for effective therapy. The present invention may also provide quality of life benefits due to, *e.g.*, decreased duration of therapy, reduced stay in intensive care units or reduced stay overall in the
25 hospital, with the concomitant reduced risk of serious nosocomial (hospital-acquired) infections.

"Concurrent administration" as used herein includes administration of the agents together, or before or after each other. The BPI protein products and anti-chlamydial agents may be administered by
30 different routes. For example, the BPI protein product may be

- 17 -

administered intravenously while the anti-chlamydial agents are administered intramuscularly, intravenously, subcutaneously, orally or intraperitoneally. Alternatively, the BPI protein product may be administered intraperitoneally while the anti-chlamydial agents are administered intraperitoneally or intravenously, or the BPI protein product may be administered in an aerosolized or nebulized form while the anti-chlamydial agents are administered, e.g., intravenously. The BPI protein product and anti-chlamydial agents may be both administered intravenously. The BPI protein product and anti-chlamydial agents may be given sequentially in the same intravenous line, after an intermediate flush, or may be given in different intravenous lines. The BPI protein product and anti-chlamydial agents may be administered simultaneously or sequentially, as long as they are given in a manner sufficient to allow both agents to achieve effective concentrations at the site of infection.

Concurrent administration of BPI protein product and antibiotic is expected to provide more effective treatment of chlamydial infections. Concurrent administration of the two agents may provide greater therapeutic effects *in vivo* than either agent provides when administered singly. For example, concurrent administration may permit a reduction in the dosage of one or both agents with achievement of a similar therapeutic effect. Alternatively, the concurrent administration may produce a more rapid or complete bactericidal/bacteriostatic effect than could be achieved with either agent alone.

Therapeutic effectiveness is based on a successful clinical outcome, and does not require that the anti-chlamydial agent or agents kill 100% of the organisms involved in the infection. Success depends on achieving a level of anti-chlamydial activity at the site of infection that is sufficient to inhibit the chlamydia in a manner that tips the balance in favor of the host. When host defenses are maximally effective, the anti-chlamydial effect required may be minimal. Reducing organism load by

- 18 -

even one log (a factor of 10) may permit the host's own defenses to control the infection. In addition, augmenting an early bactericidal/bacteriostatic effect can be more important than long-term bactericidal/bacteriostatic effect. These early events are a significant and critical part of therapeutic success, because they allow time for host defense mechanisms to activate.

BPI protein product is thought to interact with a variety of host defense elements present in whole blood or serum, including complement, p15 and LBP, and other cells and components of the immune system. Such interactions may result in potentiation of the activities of BPI protein product. Because of these interactions, BPI protein products can be expected to exert even greater activity *in vivo* than *in vitro*. Thus, while *in vitro* tests are predictive of *in vivo* utility, absence of activity *in vitro* does not necessarily indicate absence of activity *in vivo*. For example, BPI has been observed to display a greater bactericidal effect on gram-negative bacteria in whole blood or plasma assays than in assays using conventional media. [Weiss et al., *J. Clin. Invest.* 90:1122-1130 (1992)]. This may be because conventional *in vitro* systems lack the blood elements that facilitate or potentiate BPI's function *in vivo*, or because conventional media contain higher than physiological concentrations of magnesium and calcium, which are typically inhibitors of the activity of BPI protein products.

Furthermore, in the host, BPI protein product is available to neutralize translocation of gram-negative bacteria and concomitant release of endotoxin, a further clinical benefit not seen in or predicted by *in vitro* tests.

It is also contemplated that the BPI protein product be administered with other products that potentiate the activity of BPI protein products, including the anti-chlamydial activity of BPI protein products. For example, serum complement potentiates the gram-negative bactericidal activity of BPI protein products; the combination of BPI protein product and serum complement provides synergistic bactericidal/growth inhibitory

- 19 -

effects. See, e.g., Ooi *et al. J. Biol. Chem.*, 265: 15956 (1990) and Levy
et al. J. Biol. Chem., 268: 6038-6083 (1993) which address naturally-
occurring 15 kD proteins potentiating BPI antibacterial activity. See also
co-owned, co-pending PCT Application No. US94/07834 filed July 13,
5 1994, which corresponds to U.S. Patent Application Serial No. 08/274,303
filed July 11, 1994 as a continuation-in-part of U.S. Patent Application
Serial No. 08/093,201 filed July 14, 1993. These applications, which are
all incorporated herein by reference, describe methods for potentiating
gram-negative bactericidal activity of BPI protein products by administering
10 lipopolysaccharide binding protein (LBP) and LBP protein products. LBP
protein derivatives and derivative hybrids which lack CD-14
immunostimulatory properties are described in PCT Application No.
US94/06931 filed June 17, 1994, which corresponds to co-owned, co-
pending U.S. Patent Application Serial No. 08/261,660, filed June 17,
15 1994 as a continuation-in-part of U.S. Patent Application Serial No.
08/079,510, filed June 17, 1993, the disclosures of all of which are hereby
incorporated by reference. It has also been observed that poloxamer
surfactants enhance the anti-bacterial activity of BPI protein products, as
described in Lambert, U.S. Application Serial No. 08/586,133 filed
20 January 12, 1996, which is a continuation-in-part of U.S. Application
Serial No. 08/530,599 filed September 19, 1995, which is a continuation-
in-part of U.S. Application Serial No. 08/372,104 filed January 13, 1995,
all of which correspond to PCT Application No. PCT/US96/01095;
poloxamer surfactants may also enhance the activity of anti-chlamydial
25 agents.

In addition, the invention provides a method of killing or
inhibiting growth of chlamydia comprising contacting the chlamydia with a
BPI protein product. This method can be practiced *in vivo* or in a variety
of *in vitro* uses such as to decontaminate fluids and surfaces or to sterilize
30 surgical and other medical equipment and implantable devices, including

- 20 -

prostheses and intrauterine devices. These methods can also be used for *in situ* sterilization of indwelling invasive devices such as intravenous lines and catheters, which are often foci of infection.

A further aspect of the invention involves use of a BPI protein product for the manufacture of a medicament for treatment of chlamydial infection. The medicament may include, in addition to a BPI protein product, other chemotherapeutic agents such as anti-chlamydial agents. The medicament can optionally comprise a pharmaceutically acceptable diluent, adjuvant or carrier.

As used herein, "BPI protein product" includes naturally and recombinantly produced BPI protein; natural, synthetic, and recombinant biologically active polypeptide fragments of BPI protein; biologically active polypeptide variants of BPI protein or fragments thereof, including hybrid fusion proteins and dimers; biologically active polypeptide analogs of BPI protein or fragments or variants thereof, including cysteine-substituted analogs; and BPI-derived peptides. The BPI protein products administered according to this invention may be generated and/or isolated by any means known in the art. U.S. Patent No. 5,198,541, the disclosure of which is incorporated herein by reference, discloses recombinant genes encoding, and methods for expression of, BPI proteins including recombinant BPI holoprotein, referred to as rBPI and recombinant fragments of BPI. Co-owned, copending U.S. Patent Application Ser. No. 07/885,501 and a continuation-in-part thereof, U.S. Patent Application Ser. No. 08/072,063 filed May 19, 1993 and corresponding PCT Application No. 93/04752 filed May 19, 1993, which are all incorporated herein by reference, disclose novel methods for the purification of recombinant BPI protein products expressed in and secreted from genetically transformed mammalian host cells in culture and discloses how one may produce large quantities of recombinant BPI products suitable for incorporation into stable, homogeneous pharmaceutical preparations.

- 21 -

Biologically active fragments of BPI (BPI fragments) include biologically active molecules that have the same or similar amino acid sequence as a natural human BPI holoprotein, except that the fragment molecule lacks amino-terminal amino acids, internal amino acids, and/or

5 carboxy-terminal amino acids of the holoprotein. Nonlimiting examples of such fragments include a N-terminal fragment of natural human BPI of approximately 25 kD, described in Ooi et al., *J. Exp. Med.*, 174:649 (1991), and the recombinant expression product of DNA encoding N-terminal amino acids from 1 to about 193 to 199 of natural human BPI,

10 described in Gazzano-Santoro et al., *Infect. Immun.* 60:4754-4761 (1992), and referred to as rBPI₂₃. In that publication, an expression vector was used as a source of DNA encoding a recombinant expression product (rBPI₂₃) having the 31-residue signal sequence and the first 199 amino acids of the N-terminus of the mature human BPI, as set out in Figure 1 of Gray

15 et al., *supra*, except that valine at position 151 is specified by GTG rather than GTC and residue 185 is glutamic acid (specified by GAG) rather than lysine (specified by AAG). Recombinant holoprotein (rBPI) has also been produced having the sequence (SEQ ID NOS: 145 and 146) set out in Figure 1 of Gray et al., *supra*, with the exceptions noted for rBPI₂₃ and

20 with the exception that residue 417 is alanine (specified by GCT) rather than valine (specified by GTT). Other examples include dimeric forms of BPI fragments, as described in co-owned and co-pending U.S. Patent Application Serial No. 08/212,132, filed March 11, 1994, and corresponding PCT Application No. PCT/US95/03125, the disclosures of

25 which are incorporated herein by reference. Preferred dimeric products include dimeric BPI protein products wherein the monomers are amino-terminal BPI fragments having the N-terminal residues from about 1 to 175 to about 1 to 199 of BPI holoprotein. A particularly preferred dimeric product is the dimeric form of the BPI fragment having N-terminal residues

30 1 through 193, designated rBPI₄₂ dimer.

- 22 -

Biologically active variants of BPI (BPI variants) include but are not limited to recombinant hybrid fusion proteins, comprising BPI holoprotein or biologically active fragment thereof and at least a portion of at least one other polypeptide, and dimeric forms of BPI variants.

5 Examples of such hybrid fusion proteins and dimeric forms are described by Theofan et al. in co-owned, copending U.S. Patent Application Serial No. 07/885,911, and a continuation-in-part application thereof, U.S. Patent Application Serial No. 08/064,693 filed May 19, 1993 and corresponding PCT Application No. US93/04754 filed May 19, 1993, which are all
10 incorporated herein by reference and include hybrid fusion proteins comprising, at the amino-terminal end, a BPI protein or a biologically active fragment thereof and, at the carboxy-terminal end, at least one constant domain of an immunoglobulin heavy chain or allelic variant thereof.

15 Biologically active analogs of BPI (BPI analogs) include but are not limited to BPI protein products wherein one or more amino acid residues have been replaced by a different amino acid. For example, co-owned, copending U.S. Patent Application Ser. No. 08/013,801 filed February 2, 1993 and corresponding PCT Application No. US94/01235
20 filed February 2, 1994, the disclosures of which are incorporated herein by reference, discloses polypeptide analogs of BPI and BPI fragments wherein a cysteine residue is replaced by a different amino acid. A stable BPI protein product described by this application is the expression product of DNA encoding from amino acid 1 to approximately 193 or 199 of the N-
25 terminal amino acids of BPI holoprotein, but wherein the cysteine at residue number 132 is substituted with alanine and is designated rBPI₂₁Δcys or rBPI₂₁. Other examples include dimeric forms of BPI analogs; e.g. co-owned and co-pending U.S. Patent Application Serial No. 08/212,132 filed March 11, 1994, and corresponding PCT Application No.

PCT/US95/03125, the disclosures of which are incorporated herein by reference.

Other BPI protein products useful according to the methods of the invention are peptides derived from or based on BPI produced by recombinant or synthetic means (BPI-derived peptides), such as those described in co-owned and copending PCT Application No. US94/10427 filed September 15, 1994, which corresponds to U.S. Patent Application Serial No. 08/306,473, filed September 15, 1994, and PCT Application No. US94/02465 filed March 11, 1994, which corresponds to U.S. Patent Application Serial No. 08/209,762, filed March 11, 1994, which is a continuation-in-part of U.S. Patent Application Serial No. 08/183,222, filed January 14, 1994, which is a continuation-in-part of U.S. Patent Application Ser. No. 08/093,202 filed July 15, 1993 (for which the corresponding international application is PCT Application No. US94/02401 filed March 11, 1994), which is a continuation-in-part of U.S. Patent Application Ser. No. 08/030,644 filed March 12, 1993, the disclosures of all of which are incorporated herein by reference.

Presently preferred BPI protein products include recombinantly-produced N-terminal fragments of BPI, especially those having a molecular weight of approximately between 21 to 25 kD such as rBPI₂₁ or rBPI₂₃, or dimeric forms of these N-terminal fragments (e.g., rBPI₄₂ dimer). Additionally, preferred BPI protein products include rBPI and BPI-derived peptides.

The administration of BPI protein products is preferably accomplished with a pharmaceutical composition comprising a BPI protein product and a pharmaceutically acceptable diluent, adjuvant, or carrier. The BPI protein product may be administered without or in conjunction with known surfactants, other chemotherapeutic agents or additional known anti-chlamydial agents. A stable pharmaceutical composition containing BPI protein products (e.g., rBPI, rBPI₂₃) comprises the BPI protein product

- 24 -

at a concentration of 1 mg/ml in citrate buffered saline (5 or 20 mM citrate, 150 mM NaCl, pH 5.0) comprising 0.1% by weight of poloxamer 188 (Pluronic F-68, BASF Wyandotte, Parsippany, NJ) and 0.002% by weight of polysorbate 80 (Tween 80, ICI Americas Inc., Wilmington, DE).

5 Another stable pharmaceutical composition containing BPI protein products (e.g., rBPI₂₁) comprises the BPI protein product at a concentration of 2 mg/ml in 5 mM citrate, 150 mM NaCl, 0.2% poloxamer 188 and 0.002% polysorbate 80. Such preferred combinations are described in co-owned, co-pending PCT Application No. US94/01239 filed February 2, 1994,

10 which corresponds to U.S. Patent Application Ser. No. 08/190,869 filed February 2, 1994 and U.S. Patent Application Ser. No. 08/012,360 filed February 2, 1993, the disclosures of all of which are incorporated herein by reference.

Other aspects and advantages of the present invention will be

15 understood upon consideration of the following illustrative examples. Example 1 addresses the use of BPI protein product to inhibit infection of host cells with chlamydia when administered at the same time as chlamydial challenge. Example 2 addresses the anti-chlamydial activity of BPI protein product in chlamydia-infected host cells.

20

Example 1

USE OF BPI PROTEIN PRODUCT TO INHIBIT INFECTION OF HOST CELLS WITH CHLAMYDIA

A. Preparation of *Chlamydia* Stock

Chlamydia trachomatis (Ct) serovar L2 stock was prepared

25 as follows. McCoy cells (ATCC Accession No. CRL 1696) were cultured overnight in growth medium [Eagles Medium Nutrient Mixture (MEM), M-3786, Sigma, St. Louis, MO] with 1% sodium pyruvate (S-8636, Sigma) and 10% fetal bovine serum (FBS, A115-L, Hyclone, Logan, VT).

- 25 -

The media was aspirated and a vial of *Ct* was rapidly thawed and mixed with 30 mL of Dulbecco's phosphate buffered saline (PBS, Sigma) and 7% sucrose (DPBS-7). Ten mL of the suspension were added to each of 3 T150 flasks and the flasks were incubated at 37°C while being rocked
5 periodically over the next two hours to distribute the inoculum. The DPBS-7 was aspirated from the flasks and 50 mL of growth media were added to each flask. After incubation for three days at 37°C in 5% CO₂, the *Ct* was harvested as follows. The growth media was aspirated from the flasks and glass beads were added to the flasks to a depth of ~0.25 inches.
10 Ten mL Eagles MEM (without FBS) was added to each flask and the beads were rocked over the monolayer until all the cells were dislodged. The beads and cell debris were collected in 50 mL screw-capped centrifuge tubes, the flasks were washed twice with PBS, and the washings were added to the bead suspension. Each tube was placed on ice and sonicated
15 for 60 seconds to disrupt the cells. The disrupted cells/bead suspension were centrifuged at low speed (~800 rpm). The supernatant was removed and collected in a 250 mL polycarbonate centrifuge bottle, then centrifuged for one hour at high speed (~25,000 x g). The pellet was resuspended in FBS (40 mL) by repeated passage through a #16 gauge needle and syringe.
20 One mL aliquots were distributed into NUNC® (Naperville, IL) cryovials and frozen at -70°C.

B. Titration of *Chlamydia* Stock

Three vials of *Ct* stock prepared as described above in Section A were rapidly thawed at 37°C and serially diluted in 10-fold
25 concentrations in Eagles MEM or DPBS-7 without serum. Twenty-four well plates with coverslips in each well containing 24-hour McCoy cell monolayers were prepared. The media was aspirated, the wells were washed once with PBS, and 1 mL of each *Ct* dilution in either Eagles MEM or DPBS-7 was added to quadruplicate sets of McCoy cells. The

- 26 -

plates were incubated at 37°C in 5% CO₂ for 2 hours, the media was aspirated, and 2 mL of growth media was added. The plates were then reincubated at 27 °C in 5%CO₂ for 3 days, fixed in methanol, and stained for 30 minutes in a moist chamber with an FITC-labelled mouse
5 monoclonal anti-chlamydia antibody (Syva MicroTrak® Chlamydia trachomatis Culture Confirmation Test). The stained coverslips were washed in water, air dried, inverted into a drop of mounting fluid (50% glycerol; 50% PBS) and viewed using a Leitz fluorescent microscope with a 25X objective (excitation wavelength 480nm, emission wavelength 520
10 nm). The inclusion bodies were counted and comparable results were obtained over the 10⁻² to 10⁻¹⁰ concentration range tested in the Eagles MEM and DPBS-7. The 10⁻⁵ dilution of the stock preparation gave 100-300 inclusion body-forming units/mL; this dilution was selected for use in all subsequent studies using this *Ct* stock. Additional media studies were
15 performed using Basal Medium Eagle (BME, Sigma), Eagles MEM (E-MEM, Sigma), RPMI-1640 with HEPES (Sigma), RPMI-1640 without HEPES (Sigma), F-12 (Gibco) and Dulbecco's Modified Eagle's Medium Nutrient Mixture F-12 Ham (DMEM/F-12, Gibco). DMEM/F-12 without FBS was selected for use in subsequent *Chlamydia* infectivity studies.
20 Media without FBS was selected for use because the addition of 10% FBS to the above tested media inhibited infection of McCoy cells by *Ct*.

C. Infection by *Chlamydia* in the Presence or Absence of BPI Protein Product

The BPI protein product tested was rBPI₂₁ [2 mg/mL in
25 5mM sodium citrate, 150 mM sodium chloride, pH 5.0, with 0.2% PLURONIC® P123 (BASF Wyandotte, Parsippany, NJ), 0.002% polysorbate 80 (TWEEN® 80, ICI Americas Inc., Wilmington, DE) and 0.05% EDTA]. Equal volumes of formulation buffer alone [5mM sodium citrate, 150 mM sodium chloride, pH 5.0, with 0.2% PLURONIC® P123,

- 27 -

0.002 % polysorbate 80 and 0.05 % EDTA] were used as a control. Serial dilutions of rBPI₂₁ or formulation buffer were prepared with DMEM/F-12 (without FBS) so that when the serial dilutions were added at a 9:1 ratio to 1 mL of a 10⁻⁴ dilution of *Ct* stock, the final concentration of *Ct* would be a 10⁻⁵ dilution of *Ct* stock and the final rBPI₂₁ concentrations would be 128, 64, 32, 16 and 8 μg/mL. Comparable (by volume) formulation buffer controls were also prepared. The final suspensions were incubated at 37°C for 30 minutes in a water bath.

McCoy cells in DMEM/F-12/10%FBS were seeded at 2 x 10⁵ cells/well into 24-well tissue culture plates (Corning #25820), incubated for 24 hours and the media aspirated. *Ct*, with and without BPI, was added in 1 mL to duplicate wells at each rBPI₂₁ concentration. The plates were centrifuged at 2500 rpm for 30 minutes, incubated for 2 hours at 37°C in 5% CO₂, and the wells aspirated. Each well received 2 mL of DMEM/F-12/10%FBS and 1 μg/mL cycloheximide (Sigma) and the plates reincubated for 3 days. After removal of the media, the wells were washed with phosphate buffered saline (PBS), air dried, fixed with methanol and stained with Gram's iodine. The cells may be alternatively stained with FITC-labelled anti-chlamydia antibodies as described in section B above.

Using an inverted microscope, 100% of each well was scanned for the presence of inclusion bodies, which stain brown with Gram's iodine due to the high concentration of glycogen in vacuoles produced by the reproductive bodies. Results are shown below in Table 1.

- 28 -

Table 1

Number of Inclusion Bodies per Well		
	with rBPI ₂₁ (mean of 4 wells)	without rBPI ₂₁ (value for 1 well)
	128 µg/mL	0
	64 µg/mL	110
5	32 µg/mL	0
	16 µg/mL	115
	8 µg/mL	1.5
	Positive Control (Ct only)	114
10	Negative Control (no Ct)	124
		151
		0

These representative results from one of three studies indicate that rBPI₂₁ can inhibit infection of permissive cells.

Example 2

15 **ANTI-CHLAMYDIAL ACTIVITY OF BPI PROTEIN**
PRODUCT AGAINST CHLAMYDIA-INFECTED HOST CELLS

20 *Chlamydia trachomatis* (Ct) serovar L2 stock prepared as described in Example 1 was diluted to 10⁻⁵ with Dulbecco's Modified Eagle's Medium Nutrient Mixture F-12 Ham (DMEM/F-12) with 10% fetal bovine serum (FBS).

McCoy cells in DMEM/F-12/10%FBS were seeded at 1 X 10⁵ cells/well into 24-well tissue culture plates (Corning #25820), incubated for 24 hours, and the media aspirated. Ct (1 mL of the 10⁻⁵ stock) was added to each well of four plates except for two negative control wells per

- 29 -

plate. The plates were centrifuged at 2500 rpm for 30 minutes, incubated for 24 hours at 37°C in 5% CO₂, and the wells aspirated.

rBPI₂₁ as described in Example 1 was diluted to final concentrations of 128, 64, 32, 16 and 8 µg/mL in DMEM/F-12 and 1.0 mL added to the appropriate duplicate wells on each plate. Comparable formulation buffer controls as described in Example 1 were also prepared. The plates were incubated for 2 hours, and 1 mL of DMEM/F-12/20%FBS and 2 µg/mL cycloheximide was added to all wells, causing the rBPI₂₁ concentration to decrease by a factor of two. The plates were reincubated for up to 5 days.

At 24, 48, 72 and 120 hours, the media was removed from a single plate, the wells washed with PBS and air dried, fixed with methanol and stained with Gram's iodine. Using an inverted microscope, 100% of each well was scanned for the presence of inclusion bodies. Results are shown in Table 2 below.

Table 2

Initial rBPI ₂₁ Concentration*	Number of Inclusion Bodies Per Well		
	at 24 hours	at 48 hours	at 72 hours
0	285.5	398	335.75
8	194.5	180	108
16	138	140.5	109.5
32	112.5	95	57.5
64	119.5	81	39
128	113	77.5	5

*This initial concentration, which was present for the first two hours of incubation, was decreased to half of the initial value for the remainder of the 5-day incubation.

- 30 -

These representative results from one of two studies show that rBPI₂₁ at initial concentrations ranging from 16 $\mu\text{g/mL}$ to 128 $\mu\text{g/mL}$ was able to reduce the number of intracellular inclusion bodies in *Ct*-infected cells when administered 24 hours after challenge with *Ct*.

5 Numerous modifications and variations in the practice of the invention are expected to occur to those skilled in the art upon consideration of the foregoing description on the presently preferred embodiments thereof. Consequently the only limitations which should be placed upon the scope of the present invention are those that appear in the
10 appended claims.

- 31 -

SEQUENCE LISTING

- (1) GENERAL INFORMATION:
- (i) APPLICANT: KOMA Corporation
 - (ii) TITLE OF INVENTION: Anti-Chlamydial Methods and Materials
 - (iii) NUMBER OF SEQUENCES: 2
 - (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: Marshall, O'Toole, Gerstein, Murray & Borun
 - (B) STREET: 6300 Sears Tower, 233 South Wacker Drive
 - (C) CITY: Chicago
 - (D) STATE: Illinois
 - (E) COUNTRY: United States of America
 - (F) ZIP: 60606-6402
 - (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: Floppy disk
 - (B) COMPUTER: IBM PC compatible
 - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
 - (D) SOFTWARE: PatentIn Release #1.0, Version #1.25
 - (vi) CURRENT APPLICATION DATA:
 - (A) APPLICATION NUMBER:
 - (B) FILING DATE:
 - (C) CLASSIFICATION:
 - (vii) PRIOR APPLICATION DATA:
 - (A) APPLICATION NUMBER:
 - (B) FILING DATE:
 - (C) CLASSIFICATION:
 - (viii) ATTORNEY/AGENT INFORMATION:
 - (A) NAME: Borun, Michael F.
 - (B) REGISTRATION NUMBER: 25,447
 - (C) REFERENCE/DOCKET NUMBER: 27129/33433
 - (ix) TELECOMMUNICATION INFORMATION:
 - (A) TELEPHONE: 312/474-6300
 - (B) TELEFAX: 312/474-0448
 - (C) TELEX:
- (2) INFORMATION FOR SEQ ID NO:1:
- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 1813 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
 - (ii) MOLECULE TYPE: cDNA
 - (ix) FEATURE:
 - (A) NAME/KEY: CDS
 - (B) LOCATION: 31..1491
 - (ix) FEATURE:
 - (A) NAME/KEY: mat_peptide

(B) LOCATION: 124..1491

(ix) FEATURE:

(A) NAME/KEY: misc_feature

(D) OTHER INFORMATION: "rBPI"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

CAGGCCTTGA GGTTTTGGCA GCTCTGGAGG ATG AGA GAG AAC ATG GCC AGG GGC	54
Met Arg Glu Asn Met Ala Arg Gly	
-31 -30 -25	
CCT TGC AAC GCG CCG AGA TGG GTG TCC CTG ATG GTG CTC GTC GCC ATA	102
Pro Cys Asn Ala Pro Arg Trp Val Ser Leu Met Val Leu Val Ala Ile	
-20 -15 -10	
GGC ACC GCC GTG ACA GCG GCC GTC AAC CCT GGC GTC GTG GTC AGG ATC	150
Gly Thr Ala Val Thr Ala Ala Val Asn Pro Gly Val Val Val Arg Ile	
-5 1 5	
TCC CAG AAG GGC CTG GAC TAC GCC AGC CAG CAG GGG ACG GCC GCT CTG	198
Ser Gln Lys Gly Leu Asp Tyr Ala Ser Gln Gln Gly Thr Ala Ala Leu	
10 15 20 25	
CAG AAG GAG CTG AAG AGG ATC AAG ATT CCT GAC TAC TCA GAC AGC TTT	246
Gln Lys Glu Leu Lys Arg Ile Lys Ile Pro Asp Tyr Ser Asp Ser Phe	
30 35 40	
AAG ATC AAG CAT CTT GGG AAG GGG CAT TAT AGC TTC TAC AGC ATG GAC	294
Lys Ile Lys His Leu Gly Lys Gly His Tyr Ser Phe Tyr Ser Met Asp	
45 50 55	
ATC CGT GAA TTC CAG CTT CCC AGT TCC CAG ATA AGC ATG GTG CCC AAT	342
Ile Arg Glu Phe Gln Leu Pro Ser Ser Gln Ile Ser Met Val Pro Asn	
60 65 70	
GTG GGC CTT AAG TTC TCC ATC AGC AAC GCC AAT ATC AAG ATC AGC GGG	390
Val Gly Leu Lys Phe Ser Ile Ser Asn Ala Asn Ile Lys Ile Ser Gly	
75 80 85	
AAA TGG AAG GCA CAA AAG AGA TTC TTA AAA ATG AGC GGC AAT TTT GAC	438
Lys Trp Lys Ala Gln Lys Arg Phe Leu Lys Met Ser Gly Asn Phe Asp	
90 95 100 105	
CTG AGC ATA GAA GGC ATG TCC ATT TCG GCT GAT CTG AAG CTG GGC AGT	486
Leu Ser Ile Glu Gly Met Ser Ile Ser Ala Asp Leu Lys Leu Gly Ser	
110 115 120	
AAC CCC ACG TCA GGC AAG CCC ACC ATC ACC TGC TCC AGC TGC AGC AGC	534
Asn Pro Thr Ser Gly Lys Pro Thr Ile Thr Cys Ser Ser Cys Ser Ser	
125 130 135	
CAC ATC AAC AGT GTC CAC GTG CAC ATC TCA AAG AGC AAA GTC GGG TGG	582
His Ile Asn Ser Val His Val His Ile Ser Lys Ser Lys Val Gly Trp	
140 145 150	
CTG ATC CAA CTC TTC CAC AAA AAA ATT GAG TCT GCG CTT CGA AAC AAG	630
Leu Ile Gln Leu Phe His Lys Lys Ile Glu Ser Ala Leu Arg Asn Lys	
155 160 165	

- 33 -

ATG AAC AGC CAG GTC TGC GAG AAA GTG ACC AAT TCT GTA TCC TCC AAG	678
Met Asn Ser Gln Val Cys Glu Lys Val Thr Asn Ser Val Ser Ser Lys	
170 175 180 185	
CTG CAA CCT TAT TTC CAG ACT CTG CCA GTA ATG ACC AAA ATA GAT TCT	726
Leu Gln Pro Tyr Phe Gln Thr Leu Pro Val Met Thr Lys Ile Asp Ser	
190 195 200	
GTG GCT GGA ATC AAC TAT GGT CTG GTG GCA CCT CCA GCA ACC ACG GCT	774
Val Ala Gly Ile Asn Tyr Gly Leu Val Ala Pro Pro Ala Thr Thr Ala	
205 210 215	
GAG ACC CTG GAT GTA CAG ATG AAG GGG GAG TTT TAC AGT GAG AAC CAC	822
Glu Thr Leu Asp Val Gln Met Lys Gly Glu Phe Tyr Ser Glu Asn His	
220 225 230	
CAC AAT CCA CCT CCC TTT GCT CCA CCA GTG ATG GAG TTT CCC GCT GCC	870
His Asn Pro Pro Pro Phe Ala Pro Pro Val Met Glu Phe Pro Ala Ala	
235 240 245	
CAT GAC CGC ATG GTA TAC CTG GGC CTC TCA GAC TAC TTC TTC AAC ACA	918
His Asp Arg Met Val Tyr Leu Gly Leu Ser Asp Tyr Phe Phe Asn Thr	
250 255 260 265	
GCC GGG CTT GTA TAC CAA GAG GCT GGG GTC TTG AAG ATG ACC CTT AGA	966
Ala Gly Leu Val Tyr Gln Glu Ala Gly Val Leu Lys Met Thr Leu Arg	
270 275 280	
GAT GAC ATG ATT CCA AAG GAG TCC AAA TTT CGA CTG ACA ACC AAG TTC	1014
Asp Asp Met Ile Pro Lys Glu Ser Lys Phe Arg Leu Thr Thr Lys Phe	
285 290 295	
TTT GGA ACC TTC CTA CCT GAG GTG GCC AAG AAG TTT CCC AAC ATG AAG	1062
Phe Gly Thr Phe Leu Pro Glu Val Ala Lys Lys Phe Pro Asn Met Lys	
300 305 310	
ATA CAG ATC CAT GTC TCA GCC TCC ACC CCG CCA CAC CTG TCT GTG CAG	1110
Ile Gln Ile His Val Ser Ala Ser Thr Pro Pro His Leu Ser Val Gln	
315 320 325	
CCC ACC GGC CTT ACC TTC TAC CCT GCC GTG GAT GTC CAG GCC TTT GCC	1158
Pro Thr Gly Leu Thr Phe Tyr Pro Ala Val Asp Val Gln Ala Phe Ala	
330 335 340 345	
GTC CTC CCC AAC TCC TCC CTG GCT TCC CTC TTC CTG ATT GGC ATG CAC	1206
Val Leu Pro Asn Ser Ser Leu Ala Ser Leu Phe Leu Ile Gly Met His	
350 355 360	
ACA ACT GGT TCC ATG GAG GTC AGC GCC GAG TCC AAC AGG CTT GTT GGA	1254
Thr Thr Gly Ser Met Glu Val Ser Ala Glu Ser Asn Arg Leu Val Gly	
365 370 375	
GAG CTC AAG CTG GAT AGG CTG CTC CTG GAA CTG AAG CAC TCA AAT ATT	1302
Glu Leu Lys Leu Asp Arg Leu Leu Glu Leu Lys His Ser Asn Ile	
380 385 390	
GGC CCC TTC CCG GTT GAA TTG CTG CAG GAT ATC ATG AAC TAC ATT GTA	1350
Gly Pro Phe Pro Val Glu Leu Leu Gln Asp Ile Met Asn Tyr Ile Val	
395 400 405	
CCC ATT CTT GTG CTG CCC AGG GTT AAC GAG AAA CTA CAG AAA GGC TTC	1398
Pro Ile Leu Val Leu Pro Arg Val Asn Glu Lys Leu Gln Lys Gly Phe	
410 415 420 425	

CCT CTC CCG ACG CCG GCC AGA GTC CAG CTC TAC AAC GTA GTG CTT CAG	1446
Pro Leu Pro Thr Pro Ala Arg Val Gln Leu Tyr Asn Val Val Leu Gln	
430 435 440	
CCT CAC CAG AAC TTC CTG CTG TTC GGT GCA GAC GTT GTC TAT AAA	1491
Pro His Gln Asn Phe Leu Leu Phe Gly Ala Asp Val Val Tyr Lys	
445 450 455	
TGAAGGCACC AGGGGTGCCG GGGGCTGTCA GCCGCACCTG TTCCTGATGG GCTGTGGGGC	1551
ACCGGCTGCC TTTCCCCAGG GAATCCTCTC CAGATCTTAA CCAAGAGCCC CTTGCAAACCT	1611
TCTTCGACTC AGATTCAGAA ATGATCTAAA CACGAGGAAA CATTATTCAT TGGAAAAGTG	1671
CATGGTGTGT ATTTTAGGGA TTATGAGCTT CTTTCAAGGG CTAAGGCTGC AGAGATATTT	1731
CCTCCAGGAA TCGTGTTCAT ATTGTAACCA AGAAATTTCC ATTTGTGCTT CATGAAAAAA	1791
AACTTCTGGT TTTTTTCATG TG	1813

(2) INFORMATION FOR SEQ ID NO:2:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 487 amino acids
 - (B) TYPE: amino acid
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met Arg Glu Asn Met Ala Arg Gly Pro Cys Asn Ala Pro Arg Trp Val	
-31 -30 -25 -20	
Ser Leu Met Val Leu Val Ala Ile Gly Thr Ala Val Thr Ala Ala Val	
-15 -10 -5 1	
Asn Pro Gly Val Val Val Arg Ile Ser Gln Lys Gly Leu Asp Tyr Ala	
5 10 15	
Ser Gln Gln Gly Thr Ala Ala Leu Gln Lys Glu Leu Lys Arg Ile Lys	
20 25 30	
Ile Pro Asp Tyr Ser Asp Ser Phe Lys Ile Lys His Leu Gly Lys Gly	
35 40 45	
His Tyr Ser Phe Tyr Ser Met Asp Ile Arg Glu Phe Gln Leu Pro Ser	
50 55 60 65	
Ser Gln Ile Ser Met Val Pro Asn Val Gly Leu Lys Phe Ser Ile Ser	
70 75 80	
Asn Ala Asn Ile Lys Ile Ser Gly Lys Trp Lys Ala Gln Lys Arg Phe	
85 90 95	
Leu Lys Met Ser Gly Asn Phe Asp Leu Ser Ile Glu Gly Met Ser Ile	
100 105 110	
Ser Ala Asp Leu Lys Leu Gly Ser Asn Pro Thr Ser Gly Lys Pro Thr	
115 120 125	

- 35 -

Ile Thr Cys Ser Ser Cys Ser Ser His Ile Asn Ser Val His Val His
 130 135 140 145
 Ile Ser Lys Ser Lys Val Gly Trp Leu Ile Gln Leu Phe His Lys Lys
 150 155 160
 Ile Glu Ser Ala Leu Arg Asn Lys Met Asn Ser Gln Val Cys Glu Lys
 165 170 175
 Val Thr Asn Ser Val Ser Ser Lys Leu Gln Pro Tyr Phe Gln Thr Leu
 180 185 190
 Pro Val Met Thr Lys Ile Asp Ser Val Ala Gly Ile Asn Tyr Gly Leu
 195 200 205
 Val Ala Pro Pro Ala Thr Thr Ala Glu Thr Leu Asp Val Gln Met Lys
 210 215 220 225
 Gly Glu Phe Tyr Ser Glu Asn His His Asn Pro Pro Pro Phe Ala Pro
 230 235 240
 Pro Val Met Glu Phe Pro Ala Ala His Asp Arg Met Val Tyr Leu Gly
 245 250 255
 Leu Ser Asp Tyr Phe Phe Asn Thr Ala Gly Leu Val Tyr Gln Glu Ala
 260 265 270
 Gly Val Leu Lys Met Thr Leu Arg Asp Asp Met Ile Pro Lys Glu Ser
 275 280 285
 Lys Phe Arg Leu Thr Thr Lys Phe Phe Gly Thr Phe Leu Pro Glu Val
 290 295 300 305
 Ala Lys Lys Phe Pro Asn Met Lys Ile Gln Ile His Val Ser Ala Ser
 310 315 320
 Thr Pro Pro His Leu Ser Val Gln Pro Thr Gly Leu Thr Phe Tyr Pro
 325 330 335
 Ala Val Asp Val Gln Ala Phe Ala Val Leu Pro Asn Ser Ser Leu Ala
 340 345 350
 Ser Leu Phe Leu Ile Gly Met His Thr Thr Gly Ser Met Glu Val Ser
 355 360 365
 Ala Glu Ser Asn Arg Leu Val Gly Glu Leu Lys Leu Asp Arg Leu Leu
 370 375 380 385
 Leu Glu Leu Lys His Ser Asn Ile Gly Pro Phe Pro Val Glu Leu Leu
 390 395
 Gln Asp Ile Met Asn Tyr Ile Val Pro Ile Leu Val Leu Pro Arg Val
 405 410 415
 Asn Glu Lys Leu Gln Lys Gly Phe Pro Leu Pro Thr Pro Ala Arg Val
 420 425 430
 Gln Leu Tyr Asn Val Val Leu Gln Pro His Gln Asn Phe Leu Leu Phe
 435 440 445
 Gly Ala Asp Val Val Tyr Lys
 450 455

WHAT IS CLAIMED ARE:

1. A method of treating chlamydial infections comprising administering to a subject suffering from a chlamydial infection a therapeutically effective amount of a bactericidal/permeability-increasing (BPI) protein product.
2. The method of claim 1 wherein the BPI protein product is an N-terminal fragment of BPI or dimeric form thereof.
3. The method of claim 3 wherein the N-terminal fragment has a molecular weight of approximately between 21 kD and 25 kD.
4. The method of claim 1 wherein the BPI protein product is BPI holoprotein, rBPI₂₃ or rBPI₂₁.
5. The method of claim 1 wherein the BPI protein product is administered orally or parenterally at a dose of between 1 μ g/kg to 100 mg/kg per day.
6. The method of claim 1 wherein the BPI protein product is administered topically at a unit dose of between 1 μ g/mL to 1 gm/mL.
7. The method of claim 1 wherein the chlamydial infection involves a chlamydial species selected from the group consisting of *C. trachomatis*, *C. pneumoniae*, *C. psittaci* and *C. pecorum* species.
8. The method of claim 6 wherein the chlamydial species is *C. trachomatis*.

- 37 -

9. The method of claim 1 comprising the additional step of administering a non-BPI anti-chlamydial agent.

10. A method of killing or inhibiting replication of chlamydia comprising contacting the chlamydia with a bactericidal/permeability-increasing (BPI) protein product.

11. The method of claim 10 further comprising contacting the chlamydial species with a non-BPI anti-chlamydial agent.