



(12) DEMANDE DE BREVET CANADIEN
CANADIAN PATENT APPLICATION

(13) A1

(86) Date de dépôt PCT/PCT Filing Date: 2019/12/20
(87) Date publication PCT/PCT Publication Date: 2020/06/25
(85) Entrée phase nationale/National Entry: 2021/06/15
(86) N° demande PCT/PCT Application No.: US 2019/067872
(87) N° publication PCT/PCT Publication No.: 2020/132455
(30) Priorités/Priorities: 2018/12/21 (US62/783,956);
2019/10/23 (US62/924,970); 2019/11/13 (US62/934,915)


(51) Cl.Int./Int.Cl. A61K 48/00 (2006.01),
C12N 15/113 (2010.01), C12N 15/864 (2006.01),
C12N 5/00 (2006.01)

(71) Demandeur/Applicant:
THE TRUSTEES OF THE UNIVERSITY OF
PENNSYLVANIA, US

(72) Inventeurs/Inventors:
HORDEAUX, JULIETTE, US;
WILSON, JAMES M., US

(74) Agent: GOWLING WLG (CANADA) LLP

(54) Titre : COMPOSITIONS POUR LA REDUCTION SPECIFIQUE DE DRG DE L'EXPRESSION DE TRANSGENE
(54) Title: COMPOSITIONS FOR DRG-SPECIFIC REDUCTION OF TRANSGENE EXPRESSION



(57) Abrégé/Abstract:

Provided herein is a recombinant AAV (rAAV) comprising an AAV capsid and a vector genome packaged therein, wherein the vector genome comprises an AAV 5' inverted terminal repeat (ITR), an engineered nucleic acid sequence encoding a gene product for expression in target cells, and miRNA target sequences which selectively repress expression in dorsal root ganglion (DRG) cells. Also provided is a pharmaceutical composition comprising a rAAV as described herein in a formulation buffer, and a method of treating a human subject with CNS -targeted gene therapy while selectively preventing expression in DRG cells.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
25 June 2020 (25.06.2020)

(10) International Publication Number
WO 2020/132455 A1

(51) International Patent Classification:

A61K 48/00 (2006.01) *C12N 15/113* (2010.01)
C12N 5/00 (2006.01) *C12N 15/864* (2006.01)

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(21) International Application Number:

PCT/US2019/067872

(22) International Filing Date:

20 December 2019 (20.12.2019)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/783,956 21 December 2018 (21.12.2018) US
62/924,970 23 October 2019 (23.10.2019) US
62/934,915 13 November 2019 (13.11.2019) US

(71) Applicant: **THE TRUSTEES OF THE UNIVERSITY OF PENNSYLVANIA** [US/US]; 3600 Civic Center Blvd. 9th Floor, Philadelphia, PA 19104 (US).

(72) Inventors: **HORDEAUX, Juliette**; 203 Head House Court, Philadelphia, PA 19147 (US). **WILSON, James, M.**; 1831 Delancey Street, Philadelphia, PA 19103 (US).


(74) Agent: **KODROFF, Cathy, A. et al.**; Howson & Howson LLP, 325 Sentry Parkway East, Five Sentry East, Suite 160, Blue Bell, PA 19422 (US).

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- with sequence listing part of description (Rule 5.2(a))

(54) Title: COMPOSITIONS FOR DRG-SPECIFIC REDUCTION OF TRANSGENE EXPRESSION



(57) Abstract: Provided herein is a recombinant AAV (rAAV) comprising an AAV capsid and a vector genome packaged therein, wherein the vector genome comprises an AAV 5' inverted terminal repeat (ITR), an engineered nucleic acid sequence encoding a gene product for expression in target cells, and miRNA target sequences which selectively repress expression in dorsal root ganglion (DRG) cells. Also provided is a pharmaceutical composition comprising a rAAV as described herein in a formulation buffer, and a method of treating a human subject with CNS -targeted gene therapy while selectively preventing expression in DRG cells.

COMPOSITIONS FOR DRG-SPECIFIC REDUCTION OF TRANSGENE EXPRESSION

BACKGROUND OF THE INVENTION

5 The vector platform of choice for in vivo gene therapy is based on primate-derived adeno-associated viruses (AAV). In the 1960s, gene-therapy products were derived from AAVs isolated from preparations of adenoviruses (Hoggan, M.D. et al. Proc Natl Acad Sci U S A 55:1467-1474, 1966). Although these vectors were safe, many programs failed in the clinic because of poor transduction. At the turn of the century, researchers discovered a
10 family of endogenous AAVs that, as vectors, achieved much higher transduction efficiencies while retaining favorable safety profiles (Gao, G., et al. J Virol 78:6381-6388, 2004).

Untoward responses of the host to AAV vectors have been minimal. In contrast to non-viral and adenoviral vectors, which elicit vibrant acute inflammatory responses (Raper, S.E., et al. Mol Genet Metab 80:148-158, 2003; Zhang, Y., et al. Mol Ther 3:697-707, 2001),
15 AAV vectors are not pro-inflammatory. Destructive adaptive immune responses to vector-transduced cells—such as cytotoxic T cells—have been minimal following AAV vector administration. There is evidence in animals and humans that AAV can induce tolerance to capsid or transgene products under certain circumstances depending on the serotype, dose, route of administration, and immune-suppression regimen (Gernoux, G., et al. Hum Gene Ther 28:338-349, 2017; Mays, L.E. & Wilson, J.M. Mol Ther 19:16-27, 2011; Manno, C.S., et al. Nat Med 12:342-347, 2006; Mingozi, F., et al. Blood 110:2334-2341, 2007).
20 However, given the current expansion of clinical applications of AAV gene therapy, we are beginning to see toxicities that can limit the clinical impact of this technology.

The most severe toxicities have occurred following intravenous administration of
25 high doses of AAV to target the CNS and musculoskeletal system. Studies in nonhuman primates (NHPs) showed the acute development of thrombocytopenia and transaminitis, which, in some cases, evolved into a lethal syndrome of hemorrhage and shock (Hordeaux, J., et al. Mol Ther 26:664-668, 2018; Hinderer, C., et al. Hum Gene Ther. 29(3):285-298, 2018). Acute elevations in liver enzymes and/or reductions in platelets have also been
30 observed in most high-dose AAV clinical trials (AveXis, I. ZOLGENSMA Prescribing Information, 2019; Solid Biosciences Provides SGT-001 Program Update, 2019; Pfizer, Pfizer Presents Initial Clinical Data on Phase 1b Gene Therapy Study for Duchenne Muscular Dystrophy (DMD), 2019; Flanigan, K.T. et al. Molecular Genetics and

Metabolism 126:S54, 2019). Although infrequent, severe toxicities were characterized by anemia, renal failure, and complement activation (Solid Biosciences, 2019; Pfizer, 2019).

More recently, the problem of degenerating neurons in the dorsal root ganglia (DRG) of NHPs and pigs that received AAV vector either into the cerebral spinal fluid (CSF) or at 5 high doses into the blood has been observed (Hinderer, C., et al. Hum Gene Ther. 29(3):285-298, 2018; Hordeaux, J., et al. Mol Ther Methods Clin Dev 10:68-78, 2018; Hordeaux, J., et al. Mol Ther Methods Clin Dev 10:79-88, 2018). This neuronal toxicity is associated with degeneration of both the peripheral axons in peripheral nerves and the central axons that ascend through the dorsal columns of the spinal cord.

10 A need in the art exists for compositions and methods for gene therapy which minimize expression of a gene product in cells that are more sensitive to toxicity.

SUMMARY OF THE INVENTION

15 In certain embodiments, compositions and methods are provided which repress transgene expression in DRG neurons. Advantageously, these compositions decrease neuronal degeneration and/or decrease secondary dorsal spinal cord axonal degeneration which can be caused by overexpression and or immune-mediated toxicity following intrathecal or systemic gene-therapy administration.

20 In certain embodiments, a composition for gene delivery is provided which specifically represses expression of a gene product in dorsal root ganglion (DRG) comprising an expression cassette. In certain embodiments, the expression cassettes is a nucleic acid sequence comprising: (a) a coding sequence for a gene product under the control of regulatory sequences which direct expression of the gene product in a cell containing the expression cassette; and (b) at least one target sequence specific for at least one of miR-183, 25 miR-182, or miR-96, the at least one target sequence being operably linked at the 3' end of the coding sequence (a). In certain embodiments, the expression cassette is carried by a non-viral vector, a viral vector, or a non-vector based delivery system. In certain embodiments, the composition comprises at least two tandem repeats of the targeting sequences which comprise at least a first miRNA target sequence and at least a second miRNA target 30 sequence which may be the same or different. In certain embodiments, the expression cassette comprises at least two miRNA tandem repeats that are located in 3' UTR. In certain embodiments, the expression cassette comprises a 3' UTR having three miRNA tandem repeats. In certain embodiments, the at least two DRG-specific miRNA target sequences are

located in both the 5' UTR and the 3' UTR. In certain embodiments, the expression cassette is carried by a viral vector selected from a recombinant parvovirus, a recombinant lentivirus, a recombinant retrovirus, or a recombinant adenovirus. In certain embodiments, the expression cassette is carried by a non-viral vector or delivery system selected from naked 5 DNA, naked RNA, an inorganic particle, a lipid particle, a polymer-based vector, or a chitosan-based formulation.

In certain embodiments, a composition comprising an expression cassette is provided, wherein the start of the first of the at least two miRNA tandem repeats is within 20 nucleotides from the 3' end of the gene coding sequence. In certain embodiments, the 10 composition comprises an expression cassette, wherein the start of the first of the at least two miRNA tandem repeats is at least 100 nucleotides from the 3' end of the gene coding sequence. In certain embodiments, a composition comprising an expression cassette is provided, wherein the 3' UTR and the miRNA tandem repeats comprise 200 to 1200 nucleotides in length. In certain embodiments, the expression cassette comprises four 15 miRNA target sequences located in the 3' UTR. In certain embodiments, a composition is provided, wherein the expression cassette further comprises at least one target sequence specific for miR-183, miR-182, or miR-96 in the 5' UTR. In certain embodiments, the expression cassette comprises at least two miRNA target sequences located in both the 5' UTR and the 3' UTR. In certain embodiments, the expression cassette comprises at least one 20 target sequence specific for miR-183, miR-182, or miR-96 in the 5' UTR. In certain embodiments, the expression cassette comprises at least two miRNA target sequences located in both the 5' UTR and the 3' UTR.

In certain embodiments, a composition comprising an expression cassette is provided, wherein two or more consecutive miRNA target sequences are continuous and not 25 separated by a spacer. In certain embodiments, two or more of the miRNA target sequences are separated by a spacer and each spacer is different. In certain embodiments, the spacer located between the miRNA target sequences is located 3' to the first miRNA target sequence and/or 5' to the last miRNA target sequence. In certain embodiments, the spacers between the miRNA target sequences are the same.

30 In certain embodiments, a recombinant AAV (rAAV) for delivery of a gene product to a patient in need thereof is provided which specifically represses expression of a gene product in DRG, the rAAV comprising a viral capsid having packaged therein an AAV vector genome, wherein the vector genome comprises: (a) a coding sequence for the gene

product under the control of regulatory sequences which direct expression of the gene product in a cell containing the vector genome; and (b) at least one miRNA target sequence specific for at least one of miR-183, miR-182, or miR-96.

In certain embodiments, the composition comprises the expression cassette or the 5 rAAV and a formulation buffer suitable for delivery via intracerebroventricular (ICV), intrathecal (IT), intracisternal, or intravenous (IV) injection.

In certain embodiments, a method for repressing transgene expression in DRG 10 neurons is provided. The method comprises delivering a composition containing the expression cassette and/or the rAAV to a patient. In certain embodiments, the method permits reduced dose or duration of immunosuppressive therapy as compared to gene 15 therapy without the miRNA tandem repeats.

In certain embodiments, a method for modulating neuronal degeneration and/or 15 decrease secondary dorsal spinal cord axonal degeneration following intrathecal or systemic gene therapy administration is provided. The method comprises delivering a composition containing the expression cassette and/or the rAAV to a patient. In certain embodiments, the method permits reduced dose or duration of immunosuppressive therapy as compared to gene 20 therapy without the miRNA tandem repeats.

In certain embodiments, a method for enhancing expression of a transgene in cells of 25 the central nervous system (CNS) following intrathecal or systemic gene therapy administration is provided. The method comprises delivering a composition containing the expression cassette and/or the rAAV to a patient. In certain embodiments, the expression cassette or rAAV vector genome comprises at least one miR183 target sequence. In certain embodiments, transgene expression is enhanced in cells of the CNS, including one or more of pyramidal neurons, purkinje neurons, granule cells, spindle neurons, interneuron cells, 20 astrocytes, oligodendrocytes, microglia, and ependymal cells.

Other aspects and advantages of the invention will be readily apparent from the 25 following detailed description of the invention.

BRIEF DESCRIPTION OF THE DRAWINGS

30 FIG. 1A – FIG. 1C show DRG toxicity and secondary axonopathy after AAV ICM administration. (FIG. 1A) DRG contain the cell bodies of sensory pseudo-unipolar neurons, which relay sensory messages from the periphery to the CNS through peripheral axons located in peripheral nerves and central axons located in the ascending dorsal white matter

tracts of the spinal cord. (FIG. 1B) Axonopathy and DRG neuronal degeneration. Axonopathy (upper left) manifests as clear vacuoles that are either empty (missing axon) or filled with macrophages digesting myelin and cellular debris (arrow). DRG lesions (upper right and lower left) consist of neuronal cell-body degeneration (arrow) with mononuclear cell infiltrate (circle). An eosinophilic (pink) cytoplasm due to the dissolution of the Nissl bodies (central chromatolysis) characterize degenerating neurons. Increased cellularity is due to the proliferation of satellite cells (satellitosis) and inflammatory cell infiltrates. Some mononuclear cells infiltrate and phagocytose the neuronal cell body (neuronophagia). Lower right picture shows immunostaining for the transgene encoded by AAV (GFP in this case).
5 The neurons displaying degenerative changes and mononuclear cell infiltrates are the ones that show the strongest protein expression (evidenced by dark brown staining on IHC). (FIG. 1C) Examples of grade 1 to grade 5 DRG lesion and grade 1 to grade 4 dorsal spinal cord axonopathy. Severity grades are defined as follows: 1 minimal (<10%), 2 mild (10-25%), 3 moderate (25-50%), 4 marked (50-95%), and 5 severe (>95%). Grade 5 was never observed
10 in spinal cord. Arrows and circles delineate neuronal degeneration with mononuclear cell infiltrates in DRG (left column) and axonopathy (right column).
15

FIG. 2A – FIG. 2B show overexpression-related toxicity model and mitigation strategy using miRNA-induced silencing. (FIG. 2A) Pseudo-unipolar sensory neuron cell bodies are located within DRG, surrounded by satellite cells and fenestrated capillaries. The peripheral axon of pseudo-unipolar sensory neurons is located in peripheral nerves and the central axon is located in the dorsal tracts of the spinal cord. AAV vectors hijack and overload the transcription and protein-synthesis machinery, thus leading to ER stress and secondary failure to maintain distal axons. Satellite cells undergo reactive proliferation and secrete cytokines, thereby attracting inflammatory cells such as lymphocytes. Those
20 reversible changes can culminate in cell death. Subsequently, glial cells and macrophages infiltrate and phagocytose the neuronal cell bodies. (FIG. 2B) An exemplary AAV expression cassette design for DRG-specific silencing. Four short tandem repeats of a miRNA reverse-complementary sequence (miR targets or target sequences) are introduced between the stop codon and the poly-A. In DRG neurons, miRNA such as miRNA 183 binds
25 the 3' untranslated region of the mRNA and recruits the RNA-induced silencing complex (RISC), which in turn leads to silencing through mRNA cleavage. In other cell types that do not express miRNA 183, translation and protein synthesis occur without any impact from the 3' UTR region.

FIG. 3A – FIG. 3D shows miR183 targets specifically silence transgene expression in vitro and in mice DRG neurons. (FIG. 3A) We transiently co-transfected 293 cells with GFP expressing AAV plasmids harboring miR183 or miR145 targets, and control or miR183-expression vector. We detected GFP protein levels 72 hrs post-transfection and 5 quantified the levels with Western blotting. Experiments were performed in triplicates. Error bars indicate standard deviation. (FIG. 3B) We injected C57BL6/J mice IV with AAV9.CB7.GFP control vector or AAV9.CB7.GFP-miR vectors at the dose of 4×10^{12} gc. We screened three DRG-enriched miR: miR183, miR145, and miR182. We harvested DRG 10 two weeks post-injection and stained for GFP using IHC. Using the ImageJ cell-counter tool, we counted the percentage of GFP-expressing neurons over total DRG neurons. Wilcoxon 15 test, * $p<0.05$, ** $p<0.01$, *** $p<0.001$. (FIG. 3C) Here we show representative pictures of GFP immunostainings from DRG quantified in panel FIG. 3B. (FIG. 3d) We injected C57BL6/J mice IV with AAV-PHP.B.CB7.GFP control vector or AAV-PHP.B.CB7.GFP-miR (miR183, miR145, miR182). We harvested CNS and liver three weeks post-injection 15 for direct GFP observation using fluorescent microscopy. Here we show representative pictures of cerebellum, cortex, and liver.

FIG. 4A – FIG. 4C show miR183 targets specifically silence GFP expression in DRG and decrease toxicity after AA Vhu68.GFP ICM administration to NHP. We injected adult rhesus macaques ICM with 3.5×10^{13} GC of AA Vhu68.CB7.GFP control vector (n=2) or 20 AA Vhu68.CB7.GFP-miR183 (n=4). Half of the animals were sacrificed two weeks post-injection for GFP expression analysis and the other half were sacrificed two months post-injection for GFP expression and histopathology. (FIG. 4A) Representative pictures of GFP-immunostained sections of DRG, spinal cord motor neurons, cerebellum, cortex, heart, and liver two weeks post-vector administration. (FIG. 4B) Quantification of GFP-positive cells in 25 DRG (sensory neurons), spinal cord (lower motor neurons), cerebellum, and cortex in NHP (n=2 AAV.GFP, n=4 AAV.GFP-miR183). We quantified a minimum of five 20x magnification fields per region per animal using the ImageJ cell-counter tool. Error bars indicate standard deviation. Wilcoxon test, * $p<0.05$, ** $p<0.01$, *** $p<0.001$. (FIG. 4C) 30 Histopathology two months after injection shows severity grades of dorsal spinal cord axonopathy, peripheral nerves axonopathy (median, peroneal and radial nerves), and DRG neuronal degeneration and mononuclear infiltration. A board-certified Veterinary Pathologist who was blinded to the vector group established severity grades, which were defined as follows: 1 minimal (<10%), 2 mild (10-25%), 3 moderate (25-50%), 4 marked (50-95%) and

5 severe (>95% - not observed). Each bar represents one animal. 0 represents absence of lesion.

FIG. 5 shows miR183 targets specifically silence hIDUA expression in DRG after AAVhu68.hIDUA ICM administration to NHP. We injected adult rhesus macaques ICM with either 1) 1 x 1013 GC of AAVhu68.CB7.hIDUA control vector (n=3); 2) AAVhu68.CB7.hIDUA control vector with prophylactic steroids treatment (1 mg/kg/day of prednisolone from day minus 7 to day 30 followed by progressive taper off, n=3); or 3) AAVhu68.CB7.hIDUA-miR183 (n=3). Animals were sacrificed three months post-injection to analyze transgene expression and histopathology. Representative pictures show the analysis of hIDUA expression by anti-hIDUA antibody immunofluorescence (DRG, first row), anti-hIDUA IHC (lower motor neurons, cerebellum, cortex), and anti-IDUA ISH (DRG last row). hIDUA ISH: exposure time is 200 ms for AAVhu68.hIDUA with and without steroids. Sensory neurons show massive transgene mRNA expression. Exposure time is 1 s for AAV.hIDUA-miR183. Sensory neurons have low ISH signal (mRNA) in the nucleus and cytoplasm. mRNA is visible in satellite cells that surround neurons at this higher exposure time.

FIG. 6A – FIG. 6C shows miR183 mediated silencing is specific to DRG neurons and fully prevents DRG toxicity in NHP treated ICM with AAVhu68.hIDUA. (FIG. 6A) Quantification of hIDUA-positive cells in DRG (sensory neurons), spinal cord (lower motor neurons), cerebellum, and cortex in NHP (n=3 per group). A minimal of five 20x magnification fields per region were quantified per animal. Error bars represent standard deviation. Wilcoxon test, * p<0.05, ** p<0.01, *** p<0.001. (FIG. 6B) Histopathology scoring three months post-injection: dorsal axonopathy cumulative scores (sum of severity grades from cervical, thoracic, and lumbar segments – maximal possible score 15); DRG cumulative score (sum of severity grades from cervical, thoracic, and lumbar segments – maximal possible score 15) and median nerve score (sum of axonopathy and fibrosis severity grades – maximal possible score 10). A board-certified Veterinary Pathologist who was blinded to the vector group established severity grades defined as follows: 1 minimal (<10%), 2 mild (10-25%), 3 moderate (25-50%), 4 marked (50-95%) and 5 severe (>95% - not observed). 0 represents absence of lesion. Error bars represent standard deviation. (FIG. 6C) ISH using hIDUA transgene-specific probes, high magnification of DRG sensory neurons and satellite cells; 1 s exposure time with blue DAPI nuclear counterstain. Arrows: DRG sensory neurons; arrowheads: satellite cells.

FIG. 7A – FIG. 7D show T cell and antibody responses to hIDUA in NHP. Adult rhesus macaques were injected ICM with either 1) 1×10^{13} GC of AAVhu68.CB7.hIDUA control vector (n=3); 2) AAVhu68.CB7.hIDUA control vector with prophylactic steroids treatment (1 mg/kg/day of prednisolone from day minus 7 to day 30 followed by progressive taper off, n=3); or 3) AAVhu68.CB7.hIDUA-miR183 (n=3). (FIG. 7A – FIG. 7C) Interferon gamma ELISPOT responses in lymphocytes isolated from PBMC, spleen, liver, and deep cervical lymph nodes 90 days post injection. Each animal has three values representing a different peptide pool (three overlapping peptide pools to cover the entire hIDUA sequence). Red indicates a positive ELISPOT response defined as >55 spot-forming units per 10⁶ lymphocytes and three times the medium negative control upon no stimulation. (FIG. 7D) anti-hIDUA antibody ELISA assay, serum dilution 1:1,000.

FIG. 8 shows concentration of cytokines/chemokines in the CSF. Samples were collected at time of vector administration (D0) and 24 hours (24h), 21 (D21) and 35 (D35) days after vector administration.

FIG. 9 shows vector biodistribution in brain, spinal cord, and DRG in NHP. Adult rhesus macaques were injected ICM with either 1) 1×10^{13} GC of AAVhu68.CB7.hIDUA control vector (n=3); 2) AAVhu68.CB7.hIDUA control vector with prophylactic steroids treatment (1 mg/kg/day of prednisolone from day minus 7 to day 30 followed by progressive taper off, n=3); or 3) AAVhu68.CB7.hIDUA-miR183 (n=3). We extracted NHP tissue DNA with a QIAamp DNA Mini Kit. We quantified vector genomes by real-time polymerase chain reaction using Taqman reagents and primers/probes that targeted the rBG polyadenylation sequence of the vectors. Results are expressed in genome copy per diploid genome. Error bars represent standard deviation.

25 DETAILED DESCRIPTION OF THE INVENTION

The compositions and methods provided herein are useful in therapies for gene delivery for repressing transgene expression in DRG neurons through the use of miRNA. As used herein, the term “repression” includes partial reduction or complete extinction or silencing of transgene expression. Transgene expression may be assessed using an assay suitable for the selected transgene. The compositions and methods provided decrease toxicity of the DRG characterized by neuronal degeneration, secondary dorsal spinal cord axonal degeneration, and/or mononuclear cell infiltrate. In certain embodiments, the expression cassette or vector genome comprises one or more miRNA target sequences in the

untranslated region (UTR) 3' to a gene product coding sequence. Suitably, two or more miRNA target sequences are provided in tandem, optionally separated by a spacer sequence. In certain embodiments, three or more miRNA target sequences are provided in tandem, optionally separated by a spacer sequence. In certain embodiments, three or more miRNA target sequences are provided in tandem, optionally separated by a spacer sequence. A variety of delivery systems may be used to deliver the expression cassette to a subject, e.g., a human patient. Such delivery systems may be a viral vector, a non-viral vector, or a non-vector-based system (e.g., a liposome, naked DNA, naked RNA, etc.). These delivery systems may be used for delivery directly to the central nervous system (CNS), peripheral nervous system (PNS), or for intravenous or an alternative route of delivery. In other embodiments, these compositions and methods are used for systemic delivery of gene therapy vectors (e.g., rAAV). In certain embodiments, these compositions and methods are useful where high doses of vector (e.g., rAAV) are delivered. In certain embodiments, the compositions and methods provided herein permit a reduced dose, reduced length, and/or reduced number of immunomodulators to be co-administered with a gene therapy vector (e.g., a rAAV-mediated gene therapy). In certain embodiments, the compositions and methods provided herein eliminate the need to co-administer immunosuppressants or immunomodulatory therapy prior to, with, and/or following administration of a viral vector (e.g. a rAAV).

20 A “5’ UTR” is upstream of the initiation codon for a gene product coding sequence. The 5’ UTR is generally shorter than the 3’ UTR. Generally, the 5’ UTR is about 3 nucleotides to about 200 nucleotides in length, but may optionally be longer.

25 A “3’ UTR” is downstream of the coding sequence for a gene product and is generally longer than the 5’ UTR. In certain embodiments, the 3’ UTR is about 200 nucleotides to about 800 nucleotides in length, but may optionally be longer or shorter.

As used herein, an “miRNA” refers to a microRNA which is a small non-coding RNA molecule which regulates mRNA and stops it from being translated to protein. The miRNA contains a “seed sequence” which is a region of nucleotides which specifically binds to mRNA by complementary base pairing, leading to destruction or silencing of the mRNA. In certain embodiments, the seed sequence is located on the mature miRNA (5’ to 3’) and is generally located at position 2 to 7 or 2 to 8 (from the 5’ end of the sense (+) strand) of the miRNA, although it may be longer than in length. In certain embodiments, the length of the seed sequence is no less than about 30% of the length of the miRNA sequence, which may

be at least 7 nucleotides to about 28 nucleotides in length, at least 8 nucleotides to about 28 nucleotides in length, 7 nucleotides to 28 nucleotides, 8 nucleotides to 18 nucleotides, 12 nucleotides to 28 nucleotides in length, about 20 to about 26 nucleotides, about 22 nucleotides, about 24 nucleotides, or about 26 nucleotides.

5 As used herein, an “miRNA target sequence” is a sequence located on the DNA positive strand (5’ to 3’) and is at least partially complementary to a miRNA sequence, including the miRNA seed sequence. The miRNA target sequence is exogenous to the untranslated region of the encoded transgene product and is designed to be specifically targeted by miRNA in cells in which repression of transgene expression is desired. The 10 term “miR183 cluster target sequence” refers to a target sequence that responds to one or members of the miR183 cluster (alternatively termed family), including miRs-183, -96 and -182 (as described by Dambal, S. et al. Nucleic Acids Res 43:7173-7188, 2015, which is incorporated herein by reference). Without wishing to be bound by theory, the messenger RNA (mRNA) for the transgene (encoding the gene product) is present in a cell type to 15 which the expression cassette containing the miRNA is delivered, such that specific binding of the miRNA to the 3’ UTR miRNA target sequences results in mRNA silencing and cleavage, thereby reducing or eliminating transgene expression only in the cells that express the miRNA.

Typically, the miRNA target sequence is at least 7 nucleotides to about 28 nucleotides in length, at least 8 nucleotides to about 28 nucleotides in length, 7 nucleotides to 28 nucleotides, 8 nucleotides to 18 nucleotides, 12 nucleotides to 28 nucleotides in length, about 20 to about 26 nucleotides, about 22 nucleotides, about 24 nucleotides, or about 26 nucleotides, and which contains at least one consecutive region (e.g., 7 or 8 nucleotides) which is complementary to the miRNA seed sequence. In certain embodiments, the target 25 sequence comprises a sequence with exact complementarity (100%) or partial complementarity to the miRNA seed sequence with some mismatches. In certain embodiments, the target sequence comprises at least 7 to 8 nucleotides which are 100% complementary to the miRNA seed sequence. In certain embodiments, the target sequence consists of a sequence which is 100% complementary to the miRNA seed sequence. In 30 certain embodiments, the target sequence contains multiple copies (e.g., two or three copies) of the sequence which is 100% complementary to the seed sequence. In certain embodiments, the region of 100% complementarity comprises at least 30% of the length of the target sequence. In certain embodiments, the remainder of the target sequence has at least

about 80 % to about 99% complementarity to the miRNA. In certain embodiments, in an expression cassette containing a DNA positive strand, the miRNA target sequence is the reverse complement of the miRNA.

In certain embodiments, provided herein are engineered expression cassettes or 5 vector genomes comprising at least one copy of an miR target sequence directed to one or more members of the miR-183 family or cluster operably linked to a transgene to repress expression of the transgene in DRG and/or reduce or eliminate DRG toxicity and/or axonopathy. In certain embodiments, the engineered expression cassette or vector genome comprises multiple miRNA target sequences, such that the number of miRNA target 10 sequences is sufficient to reduce or minimize transgene expression in DRG to reduce and/or eliminate DRG toxicity and/or axonopathy. The expression cassette or vector genome may be delivered via any suitable carrier system, viral vector or non-viral vector, via any route, but is particularly useful for intrathecal administration.

As used herein, the terms “intrathecal delivery” or “intrathecal administration” refer 15 to a route of administration via an injection into the spinal canal, more specifically into the subarachnoid space so that it reaches the cerebrospinal fluid (CSF). Intrathecal delivery may include lumbar puncture, intraventricular (including intracerebroventricular (ICV)), suboccipital/intracisternal, and/or C1-2 puncture. For example, material may be introduced for diffusion throughout the subarachnoid space by means of lumbar puncture. In another 20 example, injection may be into the cisterna magna.

As used herein, the terms “intracisternal delivery” or “intracisternal administration” refer to a route of administration directly into the cerebrospinal fluid of the cisterna magna cerebellomedularis, more specifically via a suboccipital puncture or by direct injection into the cisterna magna or via permanently positioned tube.

25 Unexpectedly, compositions comprising the miR-183 target sequences described herein for repressing expression in the DRG have been observed to provide enhanced transgene expression in one or more different cell types (other than the DRG) within the central nervous system, including, but not limited to, neurons (including, e.g., pyramidal, purkinje, granule, spindle, and interneuron cells) or glial cells (including, e.g., astrocytes, 30 oligodendrocytes, microglia, and ependymal cells). While this observation was made following an intrathecal delivery route, this CNS-enhancing effect is not limited to CNS-delivery routes and may be achieved using other routes, e.g., high dose intravenous, high dose intramuscular, or other systemic delivery routes.

In certain embodiments, one may wish to select miR-182 target sequences and/or miR-96 target sequences for expression cassettes comprising transgenes which are not targeted to the CNS, so as to avoid enhancing CNS expression of the transgene (while repressing DRG expression). For example, expression cassettes comprising transgenes for 5 delivery to skeletal muscle or the liver may wish to avoid any enhancement of CNS expression, but prevent DRG-toxicity and/or axonopathy which can be associated with the high doses which may be required.

In certain embodiments, the vector genome or expression cassette contains at least one miRNA target sequence that is a miR-183 target sequence. In certain embodiments, the 10 vector genome or expression cassette contains an miR-183 target sequence that includes AGTGAATTCTACCAGTGCCATA (SEQ ID NO:1), where the sequence complementary to the miR-183 seed sequence is underlined. In certain embodiments, the vector genome or expression cassette contains more than one copy (e.g. two or three copies) of a sequence that is 100% complementary to the miR-183 seed sequence. In certain embodiments, a miR-183 15 target sequence is about 7 nucleotides to about 28 nucleotides in length and includes at least one region that is at least 100% complementary to the miR-183 seed sequence. In certain embodiments, a miR-183 target sequence contains a sequence with partial complementarity to SEQ ID NO: 1 and, thus, when aligned to SEQ ID NO: 1, there are one or more mismatches. In certain embodiments, a miR-183 target sequence comprises a sequence 20 having at least 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 mismatches when aligned to SEQ ID NO: 1, where the mismatches may be non-contiguous. In certain embodiments, a miR-183 target sequence includes a region of 100% complementarity which also comprises at least 30% of the length of the miR-183 target sequence. In certain embodiments, the region of 100% complementarity includes a sequence with 100% complementarity to the miR-183 seed 25 sequence. In certain embodiments, the remainder of a miR-183 target sequence has at least about 80% to about 99% complementarity to miR-183. In certain embodiments, the expression cassette or vector genome includes a miR-183 target sequence that comprises a truncated SEQ ID NO: 1, i.e., a sequence that lacks at least 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 nucleotides at either or both the 5' or 3' ends of SEQ ID NO: 1. In certain embodiments, the 30 expression cassette or vector genome comprises a transgene and one miR-183 target sequence. In yet other embodiments, the expression cassette or vector genome comprises at least two, three or four miR-183 target sequences.

In certain embodiments, the expression cassette or vector genome includes a combination of miRNA target sequences. In certain embodiments, the combination of target sequences includes different target sequences with at least partial complementarity for the same miRNA (such as miR-183). In certain embodiments, the expression cassette or vector genome includes a combination of miRNA target sequences selected from miR-183, miR-182, and/or miR-96 target sequences as provided herein. In certain embodiments, the expression cassette or vector genome comprises a transgene and two, three, or four miR-96 target sequences. In certain embodiments, an expression cassette or vector genome comprises a transgene and two, three or four miR-182 target sequences. In certain embodiments, an expression cassette or vector genome comprises at least one, at least two, at least three, or at least four miR-183 target sequences, optionally in combination with at least one, at least two, at least three, or at least four miR-182 target sequences, and/or optionally in combination with at least one, at least two, at least three, or at least four miR-96 target sequences.

Compositions comprising a transgene and an miR-182 have been observed to minimize or eliminate dorsal root ganglia toxicity and/or prevent axonopathy. However, while effective for this purpose, the expression cassettes or vector genomes containing miR-182 target sequence have not been observed to enhance CNS expression as was unexpectedly found in the composited which had the miR-183 target sequence. Thus, these compositions may be desirable for genes to be targeted outside the CNS.

In certain embodiments, provided herein is an expression cassette or vector genome that comprises one or more miR-183 family target sequences and lacks a transgene (i.e. the miR-183 family target sequence(s) is not operably linked to a sequence encoding a heterologous gene product).

In certain embodiments, the vector genome or expression cassette contains at least one miRNA target sequence that is a miR-182 target sequence. In certain embodiments, the vector genome or expression cassette contains an miR-182 target sequence that includes AGTGTGAGTTCTACCATTGCCAAA (SEQ ID NO: 3). In certain embodiments, the vector genome or expression cassette contains more than one copy (e.g. two or three copies) of a sequence that is 100% complementary to the miR-182 seed sequence. In certain embodiments, a miR-182 target sequence is about 7 nucleotides to about 28 nucleotides in length and includes at least one region that is at least 100% complementary to the miR-182 seed sequence. In certain embodiments, a miR-182 target sequence contains a sequence with

partial complementarity to SEQ ID NO: 3 and, thus, when aligned to SEQ ID NO: 3, there are one or more mismatches. In certain embodiments, a miR-183 target sequence comprises a sequence having at least 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 mismatches when aligned to SEQ ID NO: 3, where the mismatches may be non-contiguous. In certain embodiments, a miR-182 target sequence includes a region of 100% complementarity which also comprises at least 30% of the length of the miR-182 target sequence. In certain embodiments, the region of 100% complementarity includes a sequence with 100% complementarity to the miR-182 seed sequence. In certain embodiments, the remainder of a miR-182 target sequence has at least about 80% to about 99% complementarity to miR-182. In certain embodiments, the expression cassette or vector genome includes a miR-182 target sequence that comprises a truncated SEQ ID NO: 3, i.e., a sequence that lacks at least 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 nucleotides at either or both the 5' or 3' ends of SEQ ID NO: 3. In certain embodiments, the expression cassette or vector genome comprises a transgene and one miR-182 target sequence. In yet other embodiments, the expression cassette or vector genome comprises at least two, three or four miR-182 target sequences.

In certain embodiments, an expression cassette or vector genome has two or more consecutive miRNA target sequences are continuous and not separated by a spacer. In certain embodiments, wherein two or more of the miRNA target sequences are separated by a spacer. In certain embodiments, the spacer is a non-coding sequence of about 1 to about 12 nucleotides, or about 2 to about 10 nucleotides in length, or about 3 to about 10 nucleotides, about 4 to about 6 nucleotide in length, or 3, 4, 5, 6, 7, 8, 9, 10 or 11 nucleotide in length.. Optionally, a single expression cassette may contain three or more miRNA target sequences, optionally having different spacer sequences therebetween. In certain embodiments, one or more spacer is independently selected from (i) GGAT (SEQ ID NO:5); (ii) CACGTG (SEQ ID NO: 6); or (iii) GCATGC (SEQ ID NO: 7). In certain embodiments, a spacer is located 3' to the first miRNA target sequence and/or 5' to the last miRNA target sequence. In certain embodiments, the spacers between the miRNA target sequences are the same.

In certain embodiments, an expression cassette comprises a transgene and one miR-183 target sequence and one or more different miRNA target sequences. In certain embodiments, expression cassettes contains miR-96 target sequence: mRNA and on DNA positive strand (5' to 3'): AGCAAAAATGTGCTAGTGCCAAA (SEQ ID NO: 2); miR-182 target sequence: mRNA and on DNA positive strand (5' to 3'): and/or AGTGTGAGTTCTACCATTGCCAAA (SEQ ID NO: 3).

Although miR-145 has been associated with brain in the literature, the studies to date have shown that miR-145 target sequences have no effect in reducing transgene expression in dorsal root ganglia. miR-145 target sequence: mRNA and on DNA positive strand (5' to 3'): AGGGATTCTGGGAAACTGGAC (SEQ ID NO: 4).

5 As provided herein, expression cassettes and vector genomes contain transgenes operably linked, or under the control, of regulatory sequences which direct expression of the transgene product in the target cell. In certain embodiments, the expression cassette or vector genome contains a transgene that is operably linked to one or more miRNA target sequences provided herein. In certain embodiments, the expression cassette or vector genome is 10 designed to contain multiple miRNA target sequences. The miRNA target sequences are incorporated into the UTR of the transgene (i.e., 3' or downstream of the gene open reading frame).

15 The term "tandem repeats" is used herein to refer to the presence of two or more consecutive miRNA target sequences. These miRNA target sequences may be continuous, i.e., located directly after one another such that the 3' end of one is directly upstream of the 5' end of the next with no intervening sequences, or vice versa. In another embodiment, two or more of the miRNA target sequences are separated by a short spacer sequence.

20 As used herein, as "spacer" is any selected nucleic acid sequence, e.g., of 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 nucleotides in length which is located between two or more consecutive miRNA target sequences. In certain embodiments, the spacer is 1 to 8 nucleotides in length, 2 to 7 nucleotides in length, 3 to 6 nucleotides in length, four nucleotides in length, 4 to 9 nucleotides, 3 to 7 nucleotides, or values which are longer. Suitably, a spacer is a non-coding sequence. In certain embodiments, the spacer may be of four (4) nucleotides. In certain embodiments, the spacer is GGAT. In certain embodiments, the spacer is six (6) nucleotides. In certain embodiments, the spacer is CACGTG or GCATGC.

25 In certain embodiments, the tandem repeats contain two, three, four or more of the same miRNA target sequence. In certain embodiments, the tandem repeats contain at least two different miRNA target sequences, at least three different miRNA target sequences, or at least four different miRNA target sequences, etc. In certain embodiments, the tandem repeats may contain two or three of the same miRNA target sequence and a fourth miRNA target sequence which is different.

In certain embodiments, there may be at least two different sets of tandem repeats in the expression cassette. For example, a 3' UTR may contain a tandem repeat immediately

downstream of the transgene, UTR sequences, and two or more tandem repeats closer to the 3' end of the UTR. In another example, the 5' UTR may contain one, two or more miRNA target sequences. In another example the 3' may contain tandem repeats and the 5' UTR may contain at least one miRNA target sequence.

5 In certain embodiments, the expression cassette contains two, three, four or more tandem repeats which start within about 0 to 20 nucleotides of the stop codon for the transgene. In other embodiments, the expression cassette contains the miRNA tandem repeats at least 100 to about 4000 nucleotides from the stop codon for the transgene.

“Comprising” is a term meaning inclusive of other components or method steps.

10 When “comprising” is used, it is to be understood that related embodiments include descriptions using the “consisting of” terminology, which excludes other components or method steps, and “consisting essentially of” terminology, which excludes any components or method steps that substantially change the nature of the embodiment or invention. It should be understood that while various embodiments in the specification are presented 15 using “comprising” language, under various circumstances, a related embodiment is also described using “consisting of” or “consisting essentially of” language.

It is to be noted that the term “a” or “an”, refers to one or more, for example, “a vector”, is understood to represent one or more vector(s). As such, the terms “a” (or “an”), “one or more,” and “at least one” is used interchangeably herein.

20 As used herein, the term “about” means a variability of plus or minus 10 % from the reference given, unless otherwise specified.

1. Expression Cassette

An “expression cassette” as described herein, includes a nucleic acid sequence 25 encoding a functional gene product operably linked to regulatory sequences which direct its expression in a target cell and miRNA target sequences in the UTR. As described herein, the miRNA target sequences are designed to be specifically recognized by miRNA present in cells in which transgene expression is undesirable and/or reduced levels of transgene expression are desired. In certain embodiments, the miRNA target sequences specifically 30 reduce expression of the transgene in dorsal root ganglion. In certain embodiments, the miRNA target sequences are located in the 3' UTR, 5' UTR, and/or in both 3' and 5' UTR. The discussion of the miRNA target sequences found in this specification is incorporated by reference herein.

In one embodiment, the expression cassette is designed for expression in a human subject while reducing or eliminating DRG-expression of the transgene product. In one embodiment, the expression cassette is designed for expression in the central nervous system (CNS), including the cerebral spinal fluid and brain. In certain embodiments, the expression cassette or vector genome is designed for expression or enhanced expression of the transgene in one or more cell type present in the CNS (excluding the dorsal root ganglia), including nerve cells (such as, pyramidal, purkinje, granule, spindle, and interneuron cells) and glia cells (such as astrocytes, oligodendrocytes, microglia, and ependymal cells). In certain embodiments, enhanced expression of the transgene is achieved in one or more cell type with little to no expression of the transgene in another cell type of the CNS. In certain embodiments, the expression cassette is useful for expression in cells other than those of the CNS.

As used herein, the term “expression” or “gene expression” refers to the process by which information from a gene is used in the synthesis of a functional gene product. The gene product may be a protein, a peptide, or a nucleic acid polymer (such as a RNA, a DNA or a PNA).

As used herein, the term “regulatory sequence”, or “expression control sequence” refers to nucleic acid sequences, such as initiator sequences, enhancer sequences, and promoter sequences, which induce, repress, or otherwise control the transcription of protein encoding nucleic acid sequences to which they are operably linked.

As used herein, the term “operably linked” refers to both expression control sequences that are contiguous with the nucleic acid sequence encoding a gene product and/or expression control sequences that act *in trans* or at a distance to control the transcription and expression thereof.

The term “exogenous” as used to describe a nucleic acid sequence or protein means that the nucleic acid or protein does not naturally occur in the position in which it exists in a chromosome, or host cell. An exogenous nucleic acid sequence also refers to a sequence derived from and inserted into the same host cell or subject, but which is present in a non-natural state, *e.g.* a different copy number, or under the control of different regulatory elements.

The term “heterologous” as used to describe a nucleic acid sequence or protein means that the nucleic acid or protein was derived from a different organism or a different species of the same organism than the host cell or subject in which it is expressed. The term

"heterologous" when used with reference to a protein or a nucleic acid in a plasmid, expression cassette, or vector, indicates that the protein or the nucleic acid is present with another sequence or subsequence which with which the protein or nucleic acid in question is not found in the same relationship to each other in nature.

5 In one embodiment, the regulatory sequence comprises a promoter. In one embodiment, the promoter is a chicken β -actin promoter. In a further embodiment, the promoter is a hybrid of a cytomegalovirus immediate-early enhancer and the chicken β -actin promoter (a CB7 promoter). In another embodiment, a suitable promoter may include without limitation, an elongation factor 1 alpha (EF1 alpha) promoter (see, e.g., Kim DW et 10 al, Use of the human elongation factor 1 alpha promoter as a versatile and efficient expression system. *Gene*. 1990 Jul 16;91(2):217-23), a Synapsin 1 promoter (see, e.g., Kügler S et al, Human synapsin 1 gene promoter confers highly neuron-specific long-term transgene expression from an adenoviral vector in the adult rat brain depending on the transduced area. *Gene Ther*. 2003 Feb;10(4):337-47), a neuron-specific enolase (NSE) 15 promoter (see, e.g., Kim J et al, Involvement of cholesterol-rich lipid rafts in interleukin-6-induced neuroendocrine differentiation of LNCaP prostate cancer cells. *Endocrinology*. 2004 Feb;145(2):613-9. Epub 2003 Oct 16), or a CB6 promoter (see, e.g., Large-Scale Production of Adeno-Associated Viral Vector Serotype-9 Carrying the Human Survival Motor Neuron Gene, *Mol Biotechnol*. 2016 Jan;58(1):30-6. doi: 10.1007/s12033-015-9899-5).

20 Suitable promoters may be selected, including but not limited to a constitutive promoter, a tissue-specific promoter or an inducible/regulatory promoter. Example of a constitutive promoter is chicken beta-actin promoter. A variety of chicken beta-actin promoters have been described alone, or in combination with various enhancer elements (e.g., CB7 is a chicken beta-actin promoter with cytomegalovirus enhancer elements; a CAG 25 promoter, which includes the promoter, the first exon and first intron of chicken beta actin, and the splice acceptor of the rabbit beta-globin gene; a CBh promoter, SJ Gray et al, *Hu Gene Ther*, 2011 Sep; 22(9): 1143-1153). Examples of promoters that are tissue-specific are well known for liver (albumin, Miyatake et al., (1997) *J. Virol.*, 71:5124-32; hepatitis B virus core promoter, Sandig et al., (1996) *Gene Ther.*, 3:1002-9; alpha-fetoprotein (AFP), 30 Arbuthnot et al., (1996) *Hum. Gene Ther.*, 7:1503-14), neuron (such as neuron-specific enolase (NSE) promoter, Andersen et al., (1993) *Cell. Mol. Neurobiol.*, 13:503-15; neurofilament light-chain gene, Piccioli et al., (1991) *Proc. Natl. Acad. Sci. USA*, 88:5611-5; and the neuron-specific vgf gene, Piccioli et al., (1995) *Neuron*, 15:373-84), and other

tissues. Alternatively, a regulatable promoter may be selected. See, e.g., WO 2011/126808B2, incorporated by reference herein.

In one embodiment, the regulatory sequence further comprises an enhancer. In one embodiment, the regulatory sequence comprises one enhancer. In another embodiment, the 5 regulatory sequence contains two or more expression enhancers. These enhancers may be the same or may be different. For example, an enhancer may include an Alpha mic/bik enhancer or a CMV enhancer. This enhancer may be present in two copies which are located adjacent to one another. Alternatively, the dual copies of the enhancer may be separated by one or more sequences.

10 In one embodiment, the regulatory sequence further comprises an intron. In a further embodiment, the intron is a chicken beta-actin intron. Other suitable introns include those known in the art may by a human β -globulin intron, and/or a commercially available Promega® intron, and those described in WO 2011/126808.

15 In one embodiment, the regulatory sequence further comprises a Polyadenylation signal (polyA). In a further embodiment, the polyA is a rabbit globin poly A. See, e.g., WO 2014/151341. Alternatively, another polyA, e.g., a human growth hormone (hGH) polyadenylation sequence, an SV40 polyA, or a synthetic polyA may be included in an expression cassette.

20 It should be understood that the compositions in the expression cassette described herein are intended to be applied to other compositions, regiments, aspects, embodiments and methods described across the Specification.

25 Expression cassettes can be delivered via any suitable non-viral vector delivery system or by a suitable viral vector. Suitable non-viral vector delivery systems are known in the art (see, e.g., Ramamoorth and Narvekar. J Clin Diagn Res. 2015 Jan; 9(1):GE01-GE06, which is incorporated herein by reference) and can be readily selected by one of skill in the art and may include, e.g., naked DNA, naked RNA, dendrimers, PLGA, polymethacrylate, an inorganic particle, a lipid particle, a polymer-based vector, or a chitosan-based formulation.

30 2. Vector

A “vector” as used herein is a biological or chemical moiety comprising a nucleic acid sequence which can be introduced into an appropriate target cell for replication or expression of said nucleic acid sequence. Examples of a vector includes but not limited to a

recombinant virus, a plasmid, Lipoplexes, a Polymersome, Polyplexes, a dendrimer, a cell penetrating peptide (CPP) conjugate, a magnetic particle, or a nanoparticle. In one embodiment, a vector is a nucleic acid molecule into which an exogenous or heterologous or engineered nucleic acid encoding a functional gene product, which can then be introduced 5 into an appropriate target cell. Such vectors preferably have one or more origin of replication, and one or more site into which the recombinant DNA can be inserted. Vectors often have means by which cells with vectors can be selected from those without, e.g., they encode drug resistance genes. Common vectors include plasmids, viral genomes, and “artificial chromosomes”. Conventional methods of generation, production, characterization 10 or quantification of the vectors are available to one of skill in the art.

In one embodiment, the vector is a non-viral plasmid that comprises an expression cassette described thereof, e.g., “naked DNA”, “naked plasmid DNA”, RNA, and mRNA; coupled with various compositions and nano particles, including, e.g., micelles, liposomes, cationic lipid - nucleic acid compositions, poly-glycan compositions and other polymers, 15 lipid and/or cholesterol-based - nucleic acid conjugates, and other constructs such as are described herein. See, e.g., X. Su et al, Mol. Pharmaceutics, 2011, 8 (3), pp 774–787; web publication: March 21, 2011; WO2013/182683, WO 2010/053572 and WO 2012/170930, all of which are incorporated herein by reference.

In certain embodiments, the vector described herein is a “replication-defective virus” 20 or a “viral vector” which refers to a synthetic or artificial viral particle in which an expression cassette containing a nucleic acid sequence encoding a functional gene product and the DRG-detargetting miRNA target sequence(s) packaged in a viral capsid or envelope, where any viral genomic sequences also packaged within the viral capsid or envelope are replication-deficient; *i.e.*, they cannot generate progeny virions but retain the ability to infect 25 target cells. In one embodiment, the genome of the viral vector does not include genes encoding the enzymes required to replicate (the genome can be engineered to be “gutless” - containing only the nucleic acid sequence encoding flanked by the signals required for amplification and packaging of the artificial genome), but these genes may be supplied during production. Therefore, it is deemed safe for use in gene therapy since replication and 30 infection by progeny virions cannot occur except in the presence of the viral enzyme required for replication.

As used herein, a recombinant viral vector is any suitable viral vector. The examples provide illustrative recombinant adeno-associated viruses (rAAV). Other suitable viral

vectors may include, e.g., an adenovirus, a poxvirus, a bocavirus, a hybrid AAV/bocavirus, a herpes simplex virus, or a lentivirus. In preferred embodiments, these recombinant viruses are replication incompetent.

As used herein, the term “host cell” may refer to the packaging cell line in which a vector (e.g., a recombinant AAV) is produced. A host cell may be a prokaryotic or eukaryotic cell (e.g., human, insect, or yeast) that contains exogenous or heterologous DNA that has been introduced into the cell by any means, e.g., electroporation, calcium phosphate precipitation, microinjection, transformation, viral infection, transfection, liposome delivery, membrane fusion techniques, high velocity DNA-coated pellets, viral infection and protoplast fusion. Examples of host cells may include, but are not limited to an isolated cell, a cell culture, an *Escherichia coli* cell, a yeast cell, a human cell, a non-human cell, a mammalian cell, a non-mammalian cell, an insect cell, an HEK-293 cell, a liver cell, a kidney cell, a cell of the central nervous system, a neuron, a glial cell, or a stem cell.

As used herein, the term “target cell” refers to any target cell in which expression of the functional gene product is desired. Examples of target cells may include, but are not limited to, a liver cell, a kidney cell, a cell of the central nervous system, a neuron, a glial cell, and a stem cell. In certain embodiments, the vector is delivered to a target cell *ex vivo*. In certain embodiments, the vector is delivered to the target cell *in vivo*.

As used herein, a “vector genome” refers to the nucleic acid sequence packaged inside a viral vector. In one example, a “vector genome” contains, at a minimum, from 5' to 3', a vector-specific sequence, a nucleic acid sequence encoding a functional gene product operably linked to regulatory control sequences which direct its expression in a target cell and miRNA target sequences in the untranslated region(s) and a vector-specific sequence. For example, an AAV vector genome contains inverted terminal repeat sequences and an expression cassette which comprises, e.g., a nucleic acid sequence encoding a functional gene product operably linked to regulatory control sequences which direct its expression in a target cell and miRNA target sequences in the untranslated region(s). As described herein, the miRNA target sequences are designed to be specifically recognized by miRNA sequences in cells in which transgene expression is undesirable (e.g., dorsal root ganglia) and/or reduced levels of transgene expression are desired.

It should be understood that the compositions in the vector described herein are intended to be applied to other compositions, regiments, aspects, embodiments and methods described across the Specification.

3. Adeno-associated Virus (AAV)

In one aspect, provided herein is a recombinant AAV (rAAV) comprising an AAV capsid and a vector genome packaged therein.

5 In one embodiment, the regulatory sequence is as described above. In one embodiment, the vector genome comprises an AAV 5' inverted terminal repeat (ITR), an expression cassette as described herein, and an AAV 3' ITR. In one embodiment, the vector genome refers to the nucleic acid sequence packaged inside a rAAV capsid forming an rAAV vector. Such a nucleic acid sequence contains AAV inverted terminal repeat sequences (ITRs) flanking an expression cassette. In one example, a "vector genome" contains, at a minimum, from 5' to 3', an AAV 5' ITR, a nucleic acid sequence encoding a functional gene product operably linked to regulatory control sequences which direct its expression in a target cell and miRNA target sequences in the untranslated region(s) and an AAV 3' ITR. In certain embodiments, the ITRs are from AAV2 and the capsid is from a 10 different AAV. Alternatively, other ITRs may be used. As described herein, the miRNA target sequences are designed to be specifically recognized by miRNA sequences in cells in which transgene expression is undesirable and/or reduced levels of transgene expression are 15 desired.

The ITRs are the genetic elements responsible for the replication and packaging of 20 the genome during vector production and are the only viral cis elements required to generate rAAV. In one embodiment, the ITRs are from an AAV different than that supplying a capsid. In a preferred embodiment, the ITR sequences from AAV2, or the deleted version thereof (Δ ITR), which may be used for convenience and to accelerate regulatory approval. However, ITRs from other AAV sources may be selected. Where the source of the ITRs is 25 from AAV2 and the AAV capsid is from another AAV source, the resulting vector may be termed pseudotyped. Typically, AAV vector genome comprises an AAV 5' ITR, the NAGLU coding sequences and any regulatory sequences, and an AAV 3' ITR. However, other configurations of these elements may be suitable. A shortened version of the 5' ITR, termed Δ ITR, has been described in which the D-sequence and terminal resolution site (trs) 30 are deleted. In other embodiments, the full-length AAV 5' and 3' ITRs are used.

The term "AAV" as used herein refers to naturally occurring adeno-associated viruses, adeno-associated viruses available to one of skill in the art and/or in light of the composition(s) and method(s) described herein, as well as artificial AAVs. An adeno-

associated virus (AAV) viral vector is an AAV DNase-resistant particle having an AAV protein capsid into which is packaged expression cassette flanked by AAV inverted terminal repeat sequences (ITRs) for delivery to target cells. An AAV capsid is composed of 60 capsid (cap) protein subunits, VP1, VP2, and VP3, that are arranged in an icosahedral symmetry in a ratio of approximately 1:1:10 to 1:1:20, depending upon the selected AAV. 5 Various AAVs may be selected as sources for capsids of AAV viral vectors as identified above. See, e.g., US Published Patent Application No. 2007-0036760-A1; US Published Patent Application No. 2009-0197338-A1; EP 1310571. See also, WO 2003/042397 (AAV7 and other simian AAV), US Patent 7790449 and US Patent 7282199 (AAV8), WO 10 2005/033321 and US 7,906,111 (AAV9), and WO 2006/110689, and WO 2003/042397 (rh.10). These documents also describe other AAV which may be selected for generating AAV and are incorporated by reference. Among the AAVs isolated or engineered from human or non-human primates (NHP) and well characterized, human AAV2 is the first AAV that was developed as a gene transfer vector; it has been widely used for efficient gene 15 transfer experiments in different target tissues and animal models. Unless otherwise specified, the AAV capsid, ITRs, and other selected AAV components described herein, may be readily selected from among any AAV, including, without limitation, the AAVs commonly identified as AAV1, AAV2, AAV3, AAV4, AAV5, AAV6, AAV6.2, AAV7, AAV8, AAV9, AAV8bp, AAVrh10, AAVhu37, AAV7M8 and AAVAnc80, variants of any 20 of the known or mentioned AAVs or AAVs yet to be discovered or variants or mixtures thereof. See, e.g., WO 2005/033321, which is incorporated herein by reference. In one embodiment, the AAV capsid is an AAV9 capsid or variant thereof. In certain embodiments, the capsid protein is designated by a number or a combination of numbers and letters following the term “AAV” in the name of the rAAV vector. 25 As used herein, relating to AAV, the term “variant” means any AAV sequence which is derived from a known AAV sequence, including those with a conservative amino acid replacement, and those sharing at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 99% or greater sequence identity over the amino acid or nucleic acid sequence. In another embodiment, the AAV capsid includes variants 30 which may include up to about 10% variation from any described or known AAV capsid sequence. That is, the AAV capsid shares about 90% identity to about 99.9 % identity, about 95% to about 99% identity or about 97% to about 98% identity to an AAV capsid provided herein and/or known in the art. In one embodiment, the AAV capsid shares at least 95%

identity with an AAV capsid. When determining the percent identity of an AAV capsid, the comparison may be made over any of the variable proteins (e.g., vp1, vp2, or vp3).

The ITRs or other AAV components may be readily isolated or engineered using techniques available to those of skill in the art from an AAV. Such AAV may be isolated, 5 engineered, or obtained from academic, commercial, or public sources (e.g., the American Type Culture Collection, Manassas, VA). Alternatively, the AAV sequences may be engineered through synthetic or other suitable means by reference to published sequences such as are available in the literature or in databases such as, e.g., GenBank, PubMed, or the like. AAV viruses may be engineered by conventional molecular biology techniques, 10 making it possible to optimize these particles for cell specific delivery of nucleic acid sequences, for minimizing immunogenicity, for tuning stability and particle lifetime, for efficient degradation, for accurate delivery to the nucleus, etc.

As used herein, the terms “rAAV” and “artificial AAV” used interchangeably, mean, without limitation, a AAV comprising a capsid protein and a vector genome packaged 15 therein, wherein the vector genome comprising a nucleic acid heterologous to the AAV. In one embodiment, the capsid protein is a non-naturally occurring capsid. Such an artificial capsid may be generated by any suitable technique, using a selected AAV sequence (e.g., a fragment of a vp1 capsid protein) in combination with heterologous sequences which may be obtained from a different selected AAV, non-contiguous portions of the same AAV, from a 20 non-AAV viral source, or from a non-viral source. An artificial AAV may be, without limitation, a pseudotyped AAV, a chimeric AAV capsid, a recombinant AAV capsid, or a “humanized” AAV capsid. Pseudotyped vectors, wherein the capsid of one AAV is replaced with a heterologous capsid protein, are useful in the invention. In one embodiment, AAV2/5 and AAV2/8 are exemplary pseudotyped vectors. The selected genetic element may be 25 delivered by any suitable method, including transfection, electroporation, liposome delivery, membrane fusion techniques, high velocity DNA-coated pellets, viral infection and protoplast fusion. The methods used to make such constructs are known to those with skill in nucleic acid manipulation and include genetic engineering, recombinant engineering, and synthetic techniques. See, e.g., Green and Sambrook, Molecular Cloning: A Laboratory 30 Manual, Cold Spring Harbor Press, Cold Spring Harbor, NY (2012).

As used herein, “AAV9 capsid” refers to the AAV9 having the amino acid sequence of (a) GenBank accession: AAS99264, is incorporated by reference herein and the AAV vp1 capsid protein is reproduced in SEQ ID NO: 17, and/or (b) the amino acid sequence encoded

by the nucleotide sequence of GenBank Accession: AY530579.1: (nt 1...2211) (reproduced in SEQ ID NO: 16). Some variation from this encoded sequence is encompassed by the present invention, which may include sequences having about 99% identity to the referenced amino acid sequence in GenBank accession: AAS99264 and US7906111 (also WO 5 2005/033321) (i.e., less than about 1% variation from the referenced sequence). Such AAV may include, e.g., natural isolates (e.g., hu68, hu31 or hu32), or variants of AAV9 having amino acid substitutions, deletions or additions, e.g., including but not limited to amino acid substitutions selected from alternate residues “recruited” from the corresponding position in any other AAV capsid aligned with the AAV9 capsid; e.g., such as described in US 10 9,102,949, US 8,927,514, US2015/349911; WO 2016/049230A11; US 9,623,120; US 9,585,971. However, in other embodiments, other variants of AAV9, or AAV9 capsids having at least about 95% identity to the above-referenced sequences may be selected. See, e.g., US Published Patent Application No. 2015/0079038. Methods of generating the capsid, coding sequences therefore, and methods for production of rAAV viral vectors have been 15 described. See, e.g., Gao, et al, Proc. Natl. Acad. Sci. U.S.A. 100 (10), 6081-6086 (2003) and US 2013/0045186A1.

AAVhu68 varies from another Clade F virus AAV9 by two encoded amino acids at positions 67 and 157 of vp1, SEQ ID NO: 9. In contrast, the other Clade F AAV (AAV9, hu31, hu31) have an Ala at position 67 and an Ala at position 157. Provided are novel 20 AAVhu68 capsids and/or engineered AAV capsids having valine (Val or V) at position 157 based on the numbering of SEQ ID NO: 9 and optionally, a glutamic acid (Glu or E) at position 67. See, also, WO 2018/160582, which is incorporate by reference herein in its entirety (which includes the sequence listing).

As used herein, the term “clade” as it relates to groups of AAV refers to a group of 25 AAV which are phylogenetically related to one another as determined using a Neighbor-Joining algorithm by a bootstrap value of at least 75% (of at least 1000 replicates) and a Poisson correction distance measurement of no more than 0.05, based on alignment of the AAV vp1 amino acid sequence. The Neighbor-Joining algorithm has been described in the literature. *See, e.g.,* M. Nei and S. Kumar, Molecular Evolution and Phylogenetics (Oxford 30 University Press, New York (2000). Computer programs are available that can be used to implement this algorithm. For example, the MEGA v2.1 program implements the modified Nei-Gojobori method. Using these techniques and computer programs, and the sequence of an AAV vp1 capsid protein, one of skill in the art can readily determine whether a selected

AAV is contained in one of the clades identified herein, in another clade, or is outside these clades. See, e.g., G Gao, et al, J Virol, 2004 Jun; 78(10): 6381-6388, which identifies Clades A, B, C, D, E and F, and provides nucleic acid sequences of novel AAV, GenBank Accession Numbers AY530553 to AY530629. See, also, WO 2005/033321.

5 In certain embodiments, an AAV68 capsid is further characterized by one or more of the following. AAV hu68 capsid proteins comprise: AAVhu68 vp1 proteins produced by expression from a nucleic acid sequence which encodes the predicted amino acid sequence of 1 to 736 of SEQ ID NO: 8, vp1 proteins produced from SEQ ID NO: 9, or vp1 proteins produced from a nucleic acid sequence at least 70% identical to SEQ ID NO: 8 which
10 encodes the predicted amino acid sequence of 1 to 736 of SEQ ID NO: 9; AA Vhu68 vp2 proteins produced by expression from a nucleic acid sequence which encodes the predicted amino acid sequence of at least about amino acids 138 to 736 of SEQ ID NO: 9, vp2 proteins produced from a sequence comprising at least nucleotides 412 to 2211 of SEQ ID NO: 8, or vp2 proteins produced from a nucleic acid sequence at least 70% identical to at least
15 nucleotides 412 to 2211 of SEQ ID NO: 8 which encodes the predicted amino acid sequence of at least about amino acids 138 to 736 of SEQ ID NO: 9, and/or AAVhu68 vp3 proteins produced by expression from a nucleic acid sequence which encodes the predicted amino acid sequence of at least about amino acids 203 to 736 of SEQ ID NO: 9, vp3 proteins produced from a sequence comprising at least nucleotides 607 to 2211 of SEQ ID NO: 8, or
20 vp3 proteins produced from a nucleic acid sequence at least 70% identical to at least nucleotides 607 to 2211 of SEQ ID NO: 8 which encodes the predicted amino acid sequence of at least about amino acids 203 to 736 of SEQ ID NO: 9.

The AA Vhu68 vp1, vp2 and vp3 proteins are typically expressed as alternative splice variants encoded by the same nucleic acid sequence which encodes the full-length vp1 amino acid sequence of SEQ ID NO: 9 (amino acid 1 to 736). Optionally the vp1-encoding sequence is used alone to express the vp1, vp2 and vp3 proteins. Alternatively, this sequence may be co-expressed with one or more of a nucleic acid sequence which encodes the AA Vhu68 vp3 amino acid sequence of SEQ ID NO: 9 (about aa 203 to 736) without the vp1-unique region (about aa 1 to about aa 137) and/or vp2-unique regions (about aa 1 to
25 about aa 202), or a strand complementary thereto, the corresponding mRNA or tRNA (about nt 607 to about nt 2211 of SEQ ID NO: 8), or a sequence at least 70% to at least 99% (e.g., at least 85%, at least 90%, at least 95%, at least 97%, at least 98% or at least 99%) identical to SEQ ID NO: 8 which encodes aa 203 to 736 of SEQ ID NO: 9. Additionally, or
30

alternatively, the vp1-encoding and/or the vp2-encoding sequence may be co-expressed with the nucleic acid sequence which encodes the AAVhu68 vp2 amino acid sequence of SEQ ID NO: 9 (about aa 138 to 736) without the vp1-unique region (about aa 1 to about 137), or a strand complementary thereto, the corresponding mRNA or tRNA (nt 412 to 2211 of SEQ 5 ID NO: 8), or a sequence at least 70% to at least 99% (e.g., at least 85%, at least 90%, at least 95%, at least 97%, at least 98% or at least 99%) identical to SEQ ID NO: 8 which encodes about aa 138 to 736 of SEQ ID NO: 9.

As described herein, a rAAVhu68 has a rAAVhu68 capsid produced in a production system expressing capsids from an AAVhu68 nucleic acid which encodes the vp1 amino acid sequence of SEQ ID NO: 9, and optionally additional nucleic acid sequences, e.g., encoding a vp 3 protein free of the vp1 and/or vp2-unique regions. The rAAVhu68 resulting from production using a single nucleic acid sequence vp1 produces the heterogenous populations of vp1 proteins, vp2 proteins and vp3 proteins. More particularly, the AAVhu68 capsid contains subpopulations within the vp1 proteins, within the vp2 proteins and within 15 the vp3 proteins which have modifications from the predicted amino acid residues in SEQ ID NO: 9. These subpopulations include, at a minimum, deamidated asparagine (N or Asn) residues. For example, asparagines in asparagine - glycine pairs are highly deamidated.

In one embodiment, the AAVhu68 vp1 nucleic acid sequence has the sequence of SEQ ID NO: 8, or a strand complementary thereto, e.g., the corresponding mRNA or tRNA. 20 In certain embodiments, the vp2 and/or vp3 proteins may be expressed additionally or alternatively from different nucleic acid sequences than the vp1, e.g., to alter the ratio of the vp proteins in a selected expression system. In certain embodiments, also provided is a nucleic acid sequence which encodes the AAVhu68 vp3 amino acid sequence of SEQ ID NO: 9 (about aa 203 to 736) without the vp1-unique region (about aa 1 to about aa 137) 25 and/or vp2-unique regions (about aa 1 to about aa 202), or a strand complementary thereto, the corresponding mRNA or tRNA (about nt 607 to about nt 2211 of SEQ ID NO: 8). In certain embodiments, also provided is a nucleic acid sequence which encodes the AAVhu68 vp2 amino acid sequence of SEQ ID NO: 9 (about aa 138 to 736) without the vp1-unique region (about aa 1 to about 137), or a strand complementary thereto, the corresponding 30 mRNA or tRNA (nt 412 to 2211 of SEQ ID NO:8).

However, other nucleic acid sequences which encode the amino acid sequence of SEQ ID NO: 9 may be selected for use in producing rAAVhu68 capsids. In certain embodiments, the nucleic acid sequence has the nucleic acid sequence of SEQ ID NO:8 or a

sequence at least 70% to 99% identical, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 99%, identical to SEQ ID NO: 8 which encodes SEQ ID NO: 9. In certain embodiments, the nucleic acid sequence has the nucleic acid sequence of SEQ ID NO: 8 or a sequence at least 70% to 99%, at least 75%, at least 80%, at 5 least 85%, at least 90%, at least 95%, at least 97%, at least 99%, identical to about nt 412 to about nt 2211 of SEQ ID NO: 8 which encodes the vp2 capsid protein (about aa 138 to 736) of SEQ ID NO: 9. In certain embodiments, the nucleic acid sequence has the nucleic acid sequence of about nt 607 to about nt 2211 of SEQ ID NO: 8 or a sequence at least 70% to 99%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at 10 least 99%, identical to nt SEQ ID NO: 8 which encodes the vp3 capsid protein (about aa 203 to 736) of SEQ ID NO: 9.

In certain embodiments, the AAVhu68 capsid is produced using a nucleic acid sequence of SEQ ID NO: 8 or a sequence at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 99%, which encodes the vp1 amino acid sequence of SEQ ID NO: 9 with a modification (e.g., deamidated amino acid) as described herein. In certain embodiments, the vp1 amino acid sequence is reproduced in SEQ ID NO: 9.

As used herein when used to refer to vp capsid proteins, the term “heterogenous” or any grammatical variation thereof, refers to a population consisting of elements that are not the same, for example, having vp1, vp2 or vp3 monomers (proteins) with different modified amino acid sequences. SEQ ID NO: 9 provides the encoded amino acid sequence of the AAVhu68 vp1 protein. The term “heterogenous” as used in connection with vp1, vp2 and vp3 proteins (alternatively termed isoforms), refers to differences in the amino acid sequence of the vp1, vp2 and vp3 proteins within a capsid. The AAV capsid contains subpopulations within the vp1 proteins, within the vp2 proteins and within the vp3 proteins which have modifications from the predicted amino acid residues. These subpopulations include, at a minimum, certain deamidated asparagine (N or Asn) residues. For example, certain subpopulations comprise at least one, two, three or four highly deamidated asparagines (N) positions in asparagine - glycine pairs and optionally further comprising other deamidated amino acids, wherein the deamidation results in an amino acid change and other optional modifications.

As used herein, a “subpopulation” of vp proteins refers to a group of vp proteins which has at least one defined characteristic in common and which consists of at least one group member to less than all members of the reference group, unless otherwise specified.

For example, a “subpopulation” of vp1 proteins is at least one (1) vp1 protein and 5 less than all vp1 proteins in an assembled AAV capsid, unless otherwise specified. A “subpopulation” of vp3 proteins may be one (1) vp3 protein to less than all vp3 proteins in an assembled AAV capsid, unless otherwise specified. For example, vp1 proteins may be a subpopulation of vp proteins; vp2 proteins may be a separate subpopulation of vp proteins, and vp3 are yet a further subpopulation of vp proteins in an assembled AAV capsid. In 10 another example, vp1, vp2 and vp3 proteins may contain subpopulations having different modifications, e.g., at least one, two, three or four highly deamidated asparagines, e.g., at asparagine - glycine pairs.

Unless otherwise specified, highly deamidated refers to at least 45% deamidated, at 15 least 50% deamidated, at least 60% deamidated, at least 65% deamidated, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 99%, or up to about 100% deamidated at a referenced amino acid position, as compared to the predicted amino acid sequence at the reference amino acid position (e.g., at least 80% of the asparagines at amino acid 57 based on the numbering of SEQ ID NO: 9 [AAVhu68] may be deamidated based on the total vp1 proteins may be deamidated based on the total vp1, vp2 20 and vp3 proteins). Such percentages may be determined using 2D-gel, mass spectrometry techniques, or other suitable techniques.

In the AAVhu68 capsid protein, 4 residues (N57, N329, N452, N512) routinely display levels of deamidation >70% and in most cases >90% across various lots. Additional asparagine residues (N94, N253, N270, N304, N409, N477, and Q599) also display 25 deamidation levels up to ~20% across various lots. The deamidation levels were initially identified using a trypsin digest and verified with a chymotrypsin digestion.

The AAVhu68 capsid contains subpopulations within the vp1 proteins, within the vp2 proteins and within the vp3 proteins which have modifications from the predicted amino acid residues in SEQ ID NO:9. These subpopulations include, at a minimum, certain 30 deamidated asparagine (N or Asn) residues. For example, certain subpopulations comprise at least one, two, three or four highly deamidated asparagines (N) positions in asparagine - glycine pairs in SEQ ID NO: 9 and optionally further comprising other deamidated amino

acids, wherein the deamidation results in an amino acid change and other optional modifications.

In other embodiments, the method involves increasing yield of a rAAV and thus, increasing the amount of an rAAV which is present in supernatant prior to, or without requiring cell lysis. This method involves engineering an AAV VP1 capsid gene to express a capsid protein having Glu at position 67, Val at position 157, or both based on an alignment having the amino acid numbering of the AAVhu68 vp1 capsid protein. In other embodiments, the method involves engineering the VP2 capsid gene to express a capsid protein having the Val at position 157. In still other embodiments, the rAAV has a modified capsid comprising both vp1 and vp2 capsid proteins Glu at position 67 and Val at position 157.

In certain embodiments, the rAAV as described herein is a self-complementary AAV. “Self-complementary AAV” refers a construct in which a coding region carried by a recombinant AAV nucleic acid sequence has been designed to form an intra-molecular double-stranded DNA template. Upon infection, rather than waiting for cell mediated synthesis of the second strand, the two complementary halves of scAAV will associate to form one double stranded DNA (dsDNA) unit that is ready for immediate replication and transcription. See, e.g., D M McCarty et al, “Self-complementary recombinant adeno-associated virus (scAAV) vectors promote efficient transduction independently of DNA synthesis”, Gene Therapy, (August 2001), Vol 8, Number 16, Pages 1248-1254. Self-complementary AAVs are described in, e.g., U.S. Patent Nos. 6,596,535; 7,125,717; and 7,456,683, each of which is incorporated herein by reference in its entirety.

In certain embodiments, the rAAV described herein is nuclease-resistant. Such nuclease may be a single nuclease, or mixtures of nucleases, and may be endonucleases or exonucleases. A nuclease-resistant rAAV indicates that the AAV capsid has fully assembled and protects these packaged genomic sequences from degradation (digestion) during nuclease incubation steps designed to remove contaminating nucleic acids which may be present from the production process. In many instances, the rAAV described herein is DNase resistant.

The recombinant adeno-associated virus (AAV) described herein may be generated using techniques which are known. See, e.g., WO 2003/042397; WO 2005/033321, WO 2006/110689; US 7588772 B2. Such a method involves culturing a host cell which contains a nucleic acid sequence encoding an AAV capsid; a functional rep gene; an expression

cassette as described herein flanked by AAV inverted terminal repeats (ITRs); and sufficient helper functions to permit packaging of the expression cassette into the AAV capsid protein. Also provided herein is the host cell which contains a nucleic acid sequence encoding an AAV capsid; a functional rep gene; a vector genome as described; and sufficient helper

5 functions to permit packaging of the vector genome into the AAV capsid protein. In one embodiment, the host cell is a HEK 293 cell. These methods are described in more detail in WO2017160360 A2, which is incorporated by reference herein.

Other methods of producing rAAV available to one of skill in the art may be utilized. Suitable methods may include without limitation, baculovirus expression system or 10 production via yeast. See, *e.g.*, Robert M. Kotin, Large-scale recombinant adeno-associated virus production. *Hum Mol Genet.* 2011 Apr 15; 20(R1): R2–R6. Published online 2011 Apr 29. doi: 10.1093/hmg/ddr141; Aucoin MG et al., Production of adeno-associated viral vectors in insect cells using triple infection: optimization of baculovirus concentration ratios. *Biotechnol Bioeng.* 2006 Dec 20;95(6):1081-92; SAMI S. THAKUR, Production of 15 Recombinant Adeno-associated viral vectors in yeast. Thesis presented to the Graduate School of the University of Florida, 2012; Kondratov O et al. Direct Head-to-Head Evaluation of Recombinant Adeno-associated Viral Vectors Manufactured in Human versus Insect Cells, *Mol Ther.* 2017 Aug 10. pii: S1525-0016(17)30362-3. doi: 10.1016/j.ymthe.2017.08.003. [Epub ahead of print]; Mietzsch M et al, OneBac 2.0: SF9 Cell 20 Lines for Production of AAV1, AAV2, and AAV8 Vectors with Minimal Encapsulation of Foreign DNA. *Hum Gene Ther Methods.* 2017 Feb;28(1):15-22. doi: 10.1089/hgtb.2016.164.; Li L et al. Production and characterization of novel recombinant adeno-associated virus replicative-form genomes: a eukaryotic source of DNA for gene transfer. *PLoS One.* 2013 Aug 1;8(8):e69879. doi: 10.1371/journal.pone.0069879. Print 25 2013; Galibert L et al, Latest developments in the large-scale production of adeno-associated virus vectors in insect cells toward the treatment of neuromuscular diseases. *J Invertebr Pathol.* 2011 Jul;107 Suppl:S80-93. doi: 10.1016/j.jip.2011.05.008; and Kotin RM, Large-scale recombinant adeno-associated virus production. *Hum Mol Genet.* 2011 Apr 15;20(R1):R2-6. doi: 10.1093/hmg/ddr141. Epub 2011 Apr 29.

30 A two-step affinity chromatography purification at high salt concentration followed by anion exchange resin chromatography are used to purify the vector drug product and to remove empty capsids. These methods are described in more detail in WO 2017/160360 entitled “Scalable Purification Method for AAV9”, which is incorporated by reference

herein. In brief, the method for separating rAAV9 particles having packaged genomic sequences from genome-deficient AAV9 intermediates involves subjecting a suspension comprising recombinant AAV9 viral particles and AAV 9 capsid intermediates to fast performance liquid chromatography, wherein the AAV9 viral particles and AAV9 intermediates are bound to a strong anion exchange resin equilibrated at a pH of 10.2, and subjected to a salt gradient while monitoring eluate for ultraviolet absorbance at about 260 and about 280. Although less optimal for rAAV9, the pH may be in the range of about 10.0 to 10.4. In this method, the AAV9 full capsids are collected from a fraction which is eluted when the ratio of A260/A280 reaches an inflection point. In one example, for the Affinity Chromatography step, the diafiltered product may be applied to a Capture SelectTM Poros-AAV2/9 affinity resin (Life Technologies) that efficiently captures the AAV2/9 serotype. Under these ionic conditions, a significant percentage of residual cellular DNA and proteins flow through the column, while AAV particles are efficiently captured.

Conventional methods for characterization or quantification of rAAV are available to one of skill in the art. To calculate empty and full particle content, VP3 band volumes for a selected sample (e.g., in examples herein an iodixanol gradient-purified preparation where # of GC = # of particles) are plotted against GC particles loaded. The resulting linear equation ($y = mx+c$) is used to calculate the number of particles in the band volumes of the test article peaks. The number of particles (pt) per 20 μ L loaded is then multiplied by 50 to give particles (pt) /mL. Pt/mL divided by GC/mL gives the ratio of particles to genome copies (pt/GC). Pt/mL–GC/mL gives empty pt/mL. Empty pt/mL divided by pt/mL and $\times 100$ gives the percentage of empty particles. Generally, methods for assaying for empty capsids and AAV vector particles with packaged genomes have been known in the art. See, e.g., Grimm et al., Gene Therapy (1999) 6:1322-1330; Sommer et al., Molec. Ther. (2003) 7:122-128. To test for denatured capsid, the methods include subjecting the treated AAV stock to SDS-polyacrylamide gel electrophoresis, consisting of any gel capable of separating the three capsid proteins, for example, a gradient gel containing 3-8% Tris-acetate in the buffer, then running the gel until sample material is separated, and blotting the gel onto nylon or nitrocellulose membranes, preferably nylon. Anti-AAV capsid antibodies are then used as the primary antibodies that bind to denatured capsid proteins, preferably an anti-AAV capsid monoclonal antibody, most preferably the B1 anti-AAV-2 monoclonal antibody (Wobus et al., J. Viral. (2000) 74:9281-9293). A secondary antibody is then used, one that binds to the primary antibody and contains a means for detecting binding with the primary antibody,

more preferably an anti-IgG antibody containing a detection molecule covalently bound to it, most preferably a sheep anti-mouse IgG antibody covalently linked to horseradish peroxidase. A method for detecting binding is used to semi-quantitatively determine binding between the primary and secondary antibodies, preferably a detection method capable of 5 detecting radioactive isotope emissions, electromagnetic radiation, or colorimetric changes, most preferably a chemiluminescence detection kit. For example, for SDS-PAGE, samples from column fractions can be taken and heated in SDS-PAGE loading buffer containing reducing agent (e.g., DTT), and capsid proteins were resolved on pre-cast gradient polyacrylamide gels (e.g., Novex). Silver staining may be performed using SilverXpress 10 (Invitrogen, CA) according to the manufacturer's instructions or other suitable staining method, i.e. SYPRO ruby or coomassie stains. In one embodiment, the concentration of AAV vector genomes (vg) in column fractions can be measured by quantitative real time PCR (Q-PCR). Samples are diluted and digested with DNase I (or another suitable nuclease) to remove exogenous DNA. After inactivation of the nuclease, the samples are further 15 diluted and amplified using primers and a TaqManTM fluorogenic probe specific for the DNA sequence between the primers. The number of cycles required to reach a defined level of fluorescence (threshold cycle, Ct) is measured for each sample on an Applied Biosystems Prism 7700 Sequence Detection System. Plasmid DNA containing identical sequences to that contained in the AAV vector is employed to generate a standard curve in the Q-PCR 20 reaction. The cycle threshold (Ct) values obtained from the samples are used to determine vector genome titer by normalizing it to the Ct value of the plasmid standard curve. End-point assays based on the digital PCR can also be used.

In one aspect, an optimized q-PCR method is used which utilizes a broad-spectrum serine protease, e.g., proteinase K (such as is commercially available from Qiagen). More 25 particularly, the optimized qPCR genome titer assay is similar to a standard assay, except that after the DNase I digestion, samples are diluted with proteinase K buffer and treated with proteinase K followed by heat inactivation. Suitably samples are diluted with proteinase K buffer in an amount equal to the sample size. The proteinase K buffer may be concentrated to 2 fold or higher. Typically, proteinase K treatment is about 0.2 mg/mL, but 30 may be varied from 0.1 mg/mL to about 1 mg/mL. The treatment step is generally conducted at about 55 °C for about 15 minutes, but may be performed at a lower temperature (e.g., about 37 °C to about 50 °C) over a longer time period (e.g., about 20 minutes to about 30 minutes), or a higher temperature (e.g., up to about 60 °C) for a shorter time period (e.g.,

about 5 to 10 minutes). Similarly, heat inactivation is generally at about 95 °C for about 15 minutes, but the temperature may be lowered (e.g., about 70 to about 90 °C) and the time extended (e.g., about 20 minutes to about 30 minutes). Samples are then diluted (e.g., 1000 fold) and subjected to TaqMan analysis as described in the standard assay.

5 Additionally, or alternatively, droplet digital PCR (ddPCR) may be used. For example, methods for determining single-stranded and self-complementary AAV vector genome titers by ddPCR have been described. See, e.g., M. Lock et al, *Hu Gene Therapy Methods*, *Hum Gene Ther Methods*. 2014 Apr;25(2):115-25. doi: 10.1089/hgtb.2013.131. Epub 2014 Feb 14.

10 Methods for determining the ratio among vp1, vp2 and vp3 of capsid protein are also available. See, e.g., Vamseedhar Rayaprolu et al, *Comparative Analysis of Adeno-Associated Virus Capsid Stability and Dynamics*, *J Virol*. 2013 Dec; 87(24): 13150–13160; Buller RM, Rose JA. 1978. Characterization of adenovirus-associated virus-induced polypeptides in KB cells. *J. Virol.* 25:331–338; and Rose JA, Maizel JV, Inman JK, Shatkin AJ. 1971. Structural proteins of adenovirus-associated viruses. *J. Virol.* 8:766–770.

15 It should be understood that the compositions in the rAAV described herein are intended to be applied to other compositions, regiments, aspects, embodiments and methods described across the Specification.

20 4. **Pharmaceutical Composition**

A pharmaceutical composition comprising the expression cassette comprising the transgene and the miRNA target sequences may be a liquid suspension, a lyophilized or frozen composition, or another suitable formulation. In certain embodiments, the composition comprises the expression cassette and a physiologically compatible liquid (e.g., 25 a solution, diluent, carrier) which form a suspension. Such a liquid is preferably aqueous based and may contain one or more: buffering agent(s), a surfactant(s), pH adjuster(s), preservative(s), or other suitable excipients. Suitable components are discussed in more detail below. The pharmaceutical composition comprises the aqueous suspending liquid and any selected excipients, and the expression cassette.

30 The expression cassette comprising the transgene and the miRNA target sequences is as described throughout this specification herein. For example, an expression cassette may be a nucleic acid sequence comprising: (a) a coding sequence for the gene product under the control of regulatory sequences which direct expression of the gene product in a cell

containing the recombinant virus; (b) regulatory sequences which direct expression of the gene product in a cell; (c) a 5' untranslated region (UTR) sequence which is 5' of the coding sequence; (d) a 3' UTR sequence which is 3' of the coding sequence; and e) at least two tandem dorsal root ganglion (DRG)-specific miRNA target sequences, wherein the at least 5 two miRNA target sequences comprise at least a first miRNA target sequence and at least a second miRNA target sequence which may be the same or different.

In certain embodiments, the pharmaceutical composition comprises the expression cassette comprising the transgene and the miRNA target sequences and a non-viral delivery system. This may include, e.g., naked DNA, naked RNA, an inorganic particle, a lipid or 10 lipid-like particle, a chitosan-based formulation and others known in the art and described for example by Ramamoorth and Narvekar, as cited above).

In other embodiments, the pharmaceutical composition is a suspension comprising the expression cassette comprising the transgene and the miRNA target sequences is a engineered in a non-viral or viral vector system. Such a non-viral vector system may 15 include, e.g., a plasmid or non-viral genetic element, or a protein-based vector.

In certain embodiments, the pharmaceutical composition comprises a non-replicating viral vector. Suitable viral vectors may include any suitable delivery vector, such as, e.g., a recombinant adenovirus, a recombinant lentivirus, a recombinant bocavirus, a recombinant adeno-associated virus (AAV), or another recombinant parvovirus. In certain embodiments, 20 the viral vector is a recombinant AAV for delivery of a gene product to a patient in need thereof.

In one embodiment, the pharmaceutical composition comprises the expression cassette comprising the transgene and the miRNA target sequences and a formulation buffer suitable for delivery via intracerebroventricular (ICV), intrathecal (IT), intracisternal or 25 intravenous (IV) injection. In one embodiment, the expression cassette comprising the transgene and the miRNA target sequences is in packaged a recombinant AAV.

In one embodiment, a composition as provided herein comprises a surfactant, preservative, excipients, and/or buffer dissolved in the aqueous suspending liquid. In one embodiment, the buffer is PBS. In another embodiment, the buffer is an artificial 30 cerebrospinal fluid (aCSF), e.g., Eliott's formulation buffer; or Harvard apparatus perfusion fluid (an artificial CSF with final Ion Concentrations (in mM): Na 150; K 3.0; Ca 1.4; Mg 0.8; P 1.0; Cl 155). Various suitable solutions are known including those which include one or more of: buffering saline, a surfactant, and a physiologically compatible salt or mixture of

salts adjusted to an ionic strength equivalent to about 100 mM sodium chloride (NaCl) to about 250 mM sodium chloride, or a physiologically compatible salt adjusted to an equivalent ionic concentration.

Suitably, the formulation is adjusted to a physiologically acceptable pH, e.g., in the 5 range of pH 6 to 8, or pH 6.5 to 7.5, pH 7.0 to 7.7, or pH 7.2 to 7.8. As the pH of the cerebrospinal fluid is about 7.28 to about 7.32, for intrathecal delivery, a pH within this range may be desired; whereas for intravenous delivery, a pH of 6.8 to about 7.2 may be desired. However, other pHs within the broadest ranges and these subranges may be selected for other routes of delivery.

10 A suitable surfactant, or combination of surfactants, may be selected from among non-ionic surfactants that are nontoxic. In one embodiment, a difunctional block copolymer surfactant terminating in primary hydroxyl groups is selected, e.g., such as Pluronic® F68 [BASF], also known as Poloxamer 188, which has a neutral pH, has an average molecular weight of 8400. Other surfactants and other Poloxamers may be selected, i.e., nonionic 15 triblock copolymers composed of a central hydrophobic chain of polyoxypropylene (poly (propylene oxide)) flanked by two hydrophilic chains of polyoxyethylene (poly (ethylene oxide)), SOLUTOL HS 15 (Macrogol-15 Hydroxystearate), LABRASOL (Polyoxy caprylic glyceride), polyoxy 10 oleyl ether, TWEEN (polyoxyethylene sorbitan fatty acid esters), ethanol and polyethylene glycol. In one embodiment, the formulation contains a poloxamer. 20 These copolymers are commonly named with the letter "P" (for poloxamer) followed by three digits: the first two digits x 100 give the approximate molecular mass of the polyoxypropylene core, and the last digit x 10 gives the percentage polyoxyethylene content. In one embodiment Poloxamer 188 is selected. The surfactant may be present in an amount up to about 0.0005 % to about 0.001% of the suspension.

25 In one example, the formulation may contain, e.g., buffered saline solution comprising one or more of sodium chloride, sodium bicarbonate, dextrose, magnesium sulfate (e.g., magnesium sulfate ·7H₂O), potassium chloride, calcium chloride (e.g., calcium chloride ·2H₂O), dibasic sodium phosphate, and mixtures thereof, in water. Suitably, for intrathecal delivery, the osmolarity is within a range compatible with cerebrospinal fluid 30 (e.g., about 275 to about 290); see, e.g., emedicine.medscape.com/article/2093316-overview. Optionally, for intrathecal delivery, a commercially available diluent may be used as a suspending agent, or in combination with another suspending agent and other optional excipients. See, e.g., Elliotts B® solution [Lukare Medical].

In other embodiments, the formulation may contain one or more permeation enhancers. Examples of suitable permeation enhancers may include, e.g., mannitol, sodium glycocholate, sodium taurocholate, sodium deoxycholate, sodium salicylate, sodium caprylate, sodium caprate, sodium lauryl sulfate, polyoxyethylene-9-laurel ether, or EDTA

5 Additionally provided is a pharmaceutical composition comprising a pharmaceutically acceptable carrier and a vector comprising a nucleic acid sequence as described herein. As used herein, “carrier” includes any and all solvents, dispersion media, vehicles, coatings, diluents, antibacterial and antifungal agents, isotonic and absorption delaying agents, buffers, carrier solutions, suspensions, colloids, and the like. The use of
10 such media and agents for pharmaceutical active substances is well known in the art. Supplementary active ingredients can also be incorporated into the compositions. Delivery vehicles such as liposomes, nanocapsules, microparticles, microspheres, lipid particles, vesicles, and the like, may be used for the introduction of the compositions of the present invention into suitable host cells. In particular, the rAAV vector may be formulated for
15 delivery either encapsulated in a lipid particle, a liposome, a vesicle, a nanosphere, or a nanoparticle or the like. In one embodiment, a therapeutically effective amount of said vector is included in the pharmaceutical composition. The selection of the carrier is not a limitation of the present invention. Other conventional pharmaceutically acceptable carrier, such as preservatives, or chemical stabilizers. Suitable exemplary preservatives include
20 chlorobutanol, potassium sorbate, sorbic acid, sulfur dioxide, propyl gallate, the parabens, ethyl vanillin, glycerin, phenol, and parachlorophenol. Suitable chemical stabilizers include gelatin and albumin.

25 The phrase “pharmaceutically-acceptable” refers to molecular entities and compositions that do not produce an allergic or similar untoward reaction when administered to a host.

As used herein, the term “dosage” or “amount” can refer to the total dosage or amount delivered to the subject in the course of treatment, or the dosage or amount delivered in a single unit (or multiple unit or split dosage) administration.

30 The aqueous suspension or pharmaceutical compositions described herein are designed for delivery to subjects in need thereof by any suitable route or a combination of different routes.

In one embodiment, the pharmaceutical composition is formulated for delivery via intracerebroventricular (ICV), intrathecal (IT), or intracisternal injection. In one

embodiment, the compositions described herein are designed for delivery to subjects in need thereof by intravenous injection. Alternatively, other routes of administration may be selected (e.g., oral, inhalation, intranasal, intratracheal, intraarterial, intraocular, intramuscular, and other parenteral routes).

5 As used herein, the terms “intrathecal delivery” or “intrathecal administration” refer to a route of administration for drugs via an injection into the spinal canal, more specifically into the subarachnoid space so that it reaches the cerebrospinal fluid (CSF). Intrathecal delivery may include lumbar puncture, intraventricular, suboccipital/intracisternal, and/or C1-2 puncture. For example, material may be introduced for diffusion throughout the
10 subarachnoid space by means of lumbar puncture. In another example, injection may be into the cisterna magna. Intracisternal delivery may increase vector diffusion and/or reduce toxicity and inflammation caused by the administration. See, e.g., Christian Hinderer et al, Widespread gene transfer in the central nervous system of cynomolgus macaques following delivery of AAV9 into the cisterna magna, Mol Ther Methods Clin Dev. 2014; 1: 14051.

15 Published online 2014 Dec 10. doi: 10.1038/mtm.2014.51.

As used herein, the terms “intracisternal delivery” or “intracisternal administration” refer to a route of administration for drugs directly into the cerebrospinal fluid of the brain ventricles or within the cisterna magna cerebellomedularis, more specifically via a suboccipital puncture or by direct injection into the cisterna magna or via permanently positioned tube.

In one aspect, provided herein is a pharmaceutical composition comprising a vector as described herein in a formulation buffer. In certain embodiments, the replication-defective virus compositions can be formulated in dosage units to contain an amount of replication-defective virus that is in the range of about 1.0×10^9 GC to about 1.0×10^{16} GC (to treat an
25 average subject of 70 kg in body weight) including all integers or fractional amounts within the range, and preferably 1.0×10^{12} GC to 1.0×10^{14} GC for a human patient. In one embodiment, the compositions are formulated to contain at least 1×10^9 , 2×10^9 , 3×10^9 , 4×10^9 , 5×10^9 , 6×10^9 , 7×10^9 , 8×10^9 , or 9×10^9 GC per dose including all integers or fractional amounts within the range. In another embodiment, the compositions are formulated to contain at least
30 1×10^{10} , 2×10^{10} , 3×10^{10} , 4×10^{10} , 5×10^{10} , 6×10^{10} , 7×10^{10} , 8×10^{10} , or 9×10^{10} GC per dose including all integers or fractional amounts within the range. In another embodiment, the compositions are formulated to contain at least 1×10^{11} , 2×10^{11} , 3×10^{11} , 4×10^{11} , 5×10^{11} , 6×10^{11} , 7×10^{11} , 8×10^{11} , or 9×10^{11} GC per dose including all integers or fractional amounts

within the range. In another embodiment, the compositions are formulated to contain at least 1×10^{12} , 2×10^{12} , 3×10^{12} , 4×10^{12} , 5×10^{12} , 6×10^{12} , 7×10^{12} , 8×10^{12} , or 9×10^{12} GC per dose including all integers or fractional amounts within the range. In another embodiment, the compositions are formulated to contain at least 1×10^{13} , 2×10^{13} , 3×10^{13} , 4×10^{13} , 5×10^{13} ,
5 6×10^{13} , 7×10^{13} , 8×10^{13} , or 9×10^{13} GC per dose including all integers or fractional amounts within the range. In another embodiment, the compositions are formulated to contain at least 1×10^{14} , 2×10^{14} , 3×10^{14} , 4×10^{14} , 5×10^{14} , 6×10^{14} , 7×10^{14} , 8×10^{14} , or 9×10^{14} GC per dose including all integers or fractional amounts within the range. In another embodiment, the compositions are formulated to contain at least 1×10^{15} , 2×10^{15} , 3×10^{15} , 4×10^{15} , 5×10^{15} ,
10 6×10^{15} , 7×10^{15} , 8×10^{15} , or 9×10^{15} GC per dose including all integers or fractional amounts within the range. In one embodiment, for human application the dose can range from 1×10^{10} to about 1×10^{12} GC per dose including all integers or fractional amounts within the range.

In one embodiment, provided is a pharmaceutical composition comprising a rAAV as described herein in a formulation buffer. In one embodiment, the rAAV is formulated at about 1×10^9 genome copies (GC)/mL to about 1×10^{14} GC/mL. In a further embodiment, the rAAV is formulated at about 3×10^9 GC/mL to about 3×10^{13} GC/mL. In yet a further embodiment, the rAAV is formulated at about 1×10^9 GC/mL to about 1×10^{13} GC/mL. In one embodiment, the rAAV is formulated at least about 1×10^{11} GC/mL. In one embodiment, the pharmaceutical composition comprising a rAAV as described herein is administrable at a dose of about 1×10^9 GC per gram of brain mass to about 1×10^{14} GC per gram of brain mass.

In certain embodiments, the composition may be formulated in a suitable aqueous suspension media (e.g., a buffered saline) for delivery by any suitable route. The compositions provided herein are useful for systemic delivery of high doses of viral vector.
25 For rAAV, a high dose may be at least 1×10^{13} GC or at least 1×10^{14} GC. However, for improved safety, the miRNA sequences provided herein may be included in expression cassettes and/or vector genomes which are delivered at other lower doses.

In certain embodiments, the composition is delivered by two different routes at essentially the same time.
30 It should be understood that the compositions in the pharmaceutical composition described herein are intended to be applied to other compositions, regimens, aspects, embodiments and methods described across the Specification.

5. Method of Treatment

In certain embodiments, the compositions provided herein are useful for delivery of a desired transgene product to patient, while for repressing transgene expression in dorsal root ganglion neurons. The method involves delivering a composition comprising an expression 5 cassette comprising the transgene and miRNA target sequences to a patient.

Examples of suitable transgenes useful in treatment of one or more neurodegenerative disorders. Such disorders may include, without limitation, transmissible spongiform encephalopathies (e.g., Creutzfeld-Jacob disease), Duchenne muscular dystrophy (DMD), myotubular myopathy and other myopathies, Parkinson's disease, amyotrophic 10 lateral sclerosis (ALS), multiple sclerosis, Alzheimer's Disease, Huntington disease, Canavan's disease, traumatic brain injury, spinal cord injury (ATI335, anti-nogo1 by Novartis), migraine (ALD403 by Alder Biopharmaceuticals; LY2951742 by Eli; RN307 by Labrys Biologics), lysosomal storage diseases, stroke, and infectious disease affecting the central nervous system. Examples of lysosomal storage disease include, e.g., Gaucher 15 disease, Fabry disease, Niemann-Pick disease, Hunter syndrome, glycogen storage disease II (Pompe disease), or Tay-Sachs disease. For certain of these conditions, e.g., DMD and myopathies, the compositions provided herein are useful in reducing or eliminating axonopathy associated with high doses of expression cassettes (e.g., carried by a viral vector) for transduction or invention of skeletal and cardiac muscle.

20 Still other nucleic acids may encode an immunoglobulin which is directed to leucine rich repeat and immunoglobulin-like domain-containing protein 1 (LINGO-1), which is a functional component of the Nogo receptor and which is associated with essential tremors in patients which multiple sclerosis, Parkinson's Disease or essential tremor. One such commercially available antibody is ocrelizumab (Biogen, BIIB033). See, e.g., US Patent 25 8,425,910. In one embodiment, the nucleic acid constructs encode immunoglobulin constructs useful for patients with ALS. Examples of suitable antibodies include antibodies against the ALS enzyme superoxide dismutase 1 (SOD1) and variants thereof (e.g., ALS variant G93A, C4F6 SOD1 antibody); MS785, which directed to Derlin-1-binding region); antibodies against neurite outgrowth inhibitor (NOGO-A or Reticulon 4), e.g., GSK1223249, 30 ozanezumab (humanized, GSK, also described as useful for multiple sclerosis). Nucleic acid sequences may be designed or selected which encode immunoglobulins useful in patients having Alzheimer's Disease. Such antibody constructs include, e.g., adumanucab (Biogen), Bapineuzumab (Elan; a humanised mAb directed at the amino terminus of A β);

Solanezumab (Eli Lilly, a humanized mAb against the central part of soluble A β); Gantenerumab (Chugai and Hoffmann-La Roche, is a full human mAb directed against both the amino terminus and central portions of A β); Crenezumab (Genentech, a humanized mAb that acts on monomeric and conformational epitopes, including oligomeric and protofibrillar forms of A β ; BAN2401 (Esai Co., Ltd, a humanized immunoglobulin G1 (IgG1) mAb that selectively binds to A β protofibrils and is thought to either enhance clearance of A β protofibrils and/or to neutralize their toxic effects on neurons in the brain); GSK 933776 (a humanised IgG1 monoclonal antibody directed against the amino terminus of A β); AAB-001, AAB-002, AAB-003 (Fc-engineered bapineuzumab); SAR228810 (a humanized mAb directed against protofibrils and low molecular weight A β); BIIB037/BART (a full human IgG1 against insoluble fibrillar human A β , Biogen Idec), an anti-A β antibody such m266, tg2576 (relative specificity for A β oligomers) [Brody and Holtzman, *Annu Rev Neurosci*, 2008; 31: 175-193]. Other antibodies may be targeted to beta-amyloid proteins, A β , beta secretase and/or the tau protein. In still other embodiments, an anti- β -amyloid antibody is derived from an IgG4 monoclonal antibodies to target β -amyloid in order to minimize effector functions, or construct other than an scFv which lacks an Fc region is selected in order to avoid amyloid related imaging abnormality (ARIA) and inflammatory response. In certain of these embodiments, the heavy chain variable region and/or the light chain variable region of one or more of the scFv constructs is used in another suitable immunoglobulin construct as provided herein. These scFV and other engineered immunoglobulins may reduce the half-life of the immunoglobulin in the serum, as compared to immunoglobulins containing Fc regions. Reducing the serum concentration of anti-amyloid molecules may further reduce the risk of ARIA, as extremely high levels of anti-amyloid antibodies in serum may destabilize cerebral vessels with a high burden of amyloid plaques, causing vascular permeability. Nucleic acids encoding other immunoglobulin constructs for treatment of patients with Parkinson's disease may be engineered or designed to express constructs, including, e.g., leucine-rich repeat kinase 2, dardarin (LRRK2) antibodies,; anti-synuclein and alpha-synuclein antibodies and DJ-1 (PARK7) antibodies,. Other antibodies may include, PRX002 (Prothena and Roche) Parkinson's disease and related synucleinopathies. These antibodies, particularly anti-synuclein antibodies may also be useful in treatment of one or more lysosomal storage disease.

One may engineer or select nucleic acid constructs encoding an immunoglobulin construct for treating multiple sclerosis. Such immunoglobulins may include or be derived

from antibodies such as natalizumab (a humanized anti-a4-ingrins, iNATA, Tysabri, Biogen Idec and Elan Pharmaceuticals), which was approved in 2006, alemtuzumab (Campath-1H, a humanized anti-CD52), rituximab (rituzin, a chimeric anti-CD20), daclizumab (Zenepax, a humanized anti-CD25), ocrelizumab (humanized, anti-CD20, Roche), ustekinumab (CINTO-1275, a human anti-IL12 p40+IL23p40); anti-LINGO-1, and ch5D12 (a chimeric anti-CD40), and rHIgM22 (a remyelinated monoclonal antibody; Acorda and the Mayo Foundation for Medical Education and Research). Still other anti-a4-integrin antibodies, anti-CD20 antibodies, anti-CD52 antibodies, anti-IL17, anti-CD19, anti-SEMA4D, and anti-CD40 antibodies may be delivered via the AAV vectors as described herein.

5 Antibodies against various infections of the central nervous system is also contemplated by the present invention. Such infectious diseases may include fungal diseases such as cryptococcal meningitis, brain abscess, spinal epidural infection caused by, e.g., Cryptococcus neoformans, Coccidioides immitis, order Mucorales, Aspergillus spp, and Candida spp; protozoal, such as toxoplasmosis, malaria, and primary amoebic

10 meningoencephalitis, caused by agents such as, e.g., Toxoplasma gondii, Taenia solium, Plasmodium falciparum, Spirometra mansonioides (sparaganoisis), Echinococcus spp (causing neuro hydatosis), and cerebral amoebiasis; bacterial, such as, e.g., tuberculosis, leprosy, neurosyphilis, bacterial meningitis, lyme disease (Borrelia burgdorferi), Rocky Mountain spotted fever (Rickettsia rickettsia), CNS nocardiosis (Nocardia spp), CNS tuberculosis

15 (Mycobacterium tuberculosis), CNS listeriosis (Listeria monocytogenes), brain abscess, and neuroborreliosis; viral infections, such as, e.g., viral meningitis, Eastern equine encephalitis (EEE), St Louis encephalitis, West Nile virus and/or encephalitis, rabies, California

20 encephalitis virus, La Crosse encephalitis, measles encephalitis, poliomyelitis, which may be caused by, e.g., herpes family viruses (HSV), HSV-1, HSV-2 (neonatal herpes simplex encephalitis), varicella zoster virus (VZV), Bickerstaff encephalitis, Epstein-Barr virus

25 (EBV), cytomegalovirus (CMV, such as TCN-202 is in development by Theraclone Sciences), human herpesvirus 6 (HHV-6), B virus (herpesvirus simiae), Flavivirus

encephalitis, Japanese encephalitis, Murray valley fever, JC virus (progressive multifocal leukoencephalopathy), Nipah Virus (NiV), measles (subacute sclerosing panencephalitis);

30 and other infections, such as, e.g., subacute sclerosing panencephalitis, progressive multifocal leukoencephalopathy; human immunodeficiency virus (acquired immunodeficiency syndrome (AIDS)); streptococcus pyogenes and other β - hemolytic Streptococcus (e.g., Pediatric Autoimmune Neuropsychiatric Disorders Associated with

Streptococcal Infection, PANDAS) and/or Sydenham's chorea, and Guillain-Barre syndrome, and prions.

Examples of suitable antibody constructs may include those described, e.g., in WO 2007/012924A2, Jan 29, 2015, which is incorporated by reference herein.

5 For example, other nucleic acid sequences may encode anti-prion immunoglobulin constructs. Such immunoglobulins may be directed against major prion protein (PrP, for prion protein or protease-resistant protein, also known as CD230 (cluster of differentiation 230). The amino acid sequence of PrP is provided, e.g.,
http://www.ncbi.nlm.nih.gov/protein/NP_000302, incorporated by reference herein. The
10 protein can exist in multiple isoforms, the normal PrPC, the disease-causing PrPSc, and an isoform located in mitochondria. The misfolded version PrPSc is associated with a variety of cognitive disorders and neurodegenerative diseases such as Creutzfeldt-Jakob disease, bovine spongiform encephalopathy, Gerstmann-Sträussler-Scheinker syndrome, fatal familial insomnia, and kuru.

15 Examples of suitable gene products may include those associated with familial hypercholesterolemia, muscular dystrophy, cystic fibrosis, and rare or orphan diseases. Examples of such rare disease may include spinal muscular atrophy (SMA), Huntingdon's Disease, Rett Syndrome (e.g., methyl-CpG-binding protein 2 (MeCP2); UniProtKB – P51608), Amyotrophic Lateral Sclerosis (ALS), Duchenne Type Muscular dystrophy,
20 Friedrichs Ataxia (e.g., frataxin), progranulin (PRGN) (associated with non-Alzheimer's cerebral degenerations, including, frontotemporal dementia (FTD), progressive non-fluent aphasia (PNFA) and semantic dementia), among others. Other useful gene products include, carbamoyl synthetase I, ornithine transcarbamylase (OTC), arginosuccinate synthetase, arginosuccinate lyase (ASL) for treatment of arginosuccinate lyase deficiency,
25 arginase, fumarylacetate hydrolase, phenylalanine hydroxylase, alpha-1 antitrypsin, rhesus alpha- fetoprotein (AFP), rhesus chorionic gonadotrophin (CG), glucose-6-phosphatase, porphobilinogen deaminase, cystathione beta-synthase, branched chain ketoacid decarboxylase, albumin, isovaleryl-coA dehydrogenase, propionyl CoA carboxylase, methyl malonyl CoA mutase, glutaryl CoA dehydrogenase, insulin, beta-glucosidase, pyruvate
30 carboxylate, hepatic phosphorylase, phosphorylase kinase, glycine decarboxylase, H-protein, T-protein, a cystic fibrosis transmembrane regulator (CFTR) sequence, and a dystrophin gene product [e.g., a mini- or micro-dystrophin]. Still other useful gene products include enzymes such as may be useful in enzyme replacement therapy, which is useful in a variety

of conditions resulting from deficient activity of enzyme. For example, enzymes that contain mannose-6-phosphate may be utilized in therapies for lysosomal storage diseases (e.g., a suitable gene includes that encoding β -glucuronidase (GUSB)).

Further illustrative genes which may be delivered via the rAAV include,

5 without limitation, glucose-6-phosphatase, associated with glycogen storage disease or deficiency type 1A (GSD1), phosphoenolpyruvate-carboxykinase (PEPCK), associated with PEPCK deficiency; cyclin-dependent kinase-like 5 (CDKL5), also known as serine/threonine kinase 9 (STK9) associated with seizures and severe neurodevelopmental impairment; galactose-1 phosphate uridyl transferase, associated with galactosemia; phenylalanine

10 hydroxylase, associated with phenylketonuria (PKU); branched chain alpha-ketoacid dehydrogenase, associated with Maple syrup urine disease; fumarylacetoacetate hydrolase, associated with tyrosinemia type 1; methylmalonyl-CoA mutase, associated with methylmalonic acidemia; medium chain acyl CoA dehydrogenase, associated with medium chain acetyl CoA deficiency; ornithine transcarbamylase (OTC), associated with ornithine

15 transcarbamylase deficiency; argininosuccinic acid synthetase (ASS1), associated with citrullinemia; lecithin-cholesterol acyltransferase (LCAT) deficiency; a methylmalonic acidemia (MMA); Niemann-Pick disease, type C1); propionic academia (PA); low density lipoprotein receptor (LDLR) protein, associated with familial hypercholesterolemia (FH); UDP-glucuronosyltransferase, associated with Crigler-Najjar disease; adenosine deaminase,

20 associated with severe combined immunodeficiency disease; hypoxanthine guanine phosphoribosyl transferase, associated with Gout and Lesch-Nyan syndrome; biotimidase, associated with biotimidase deficiency; alpha-galactosidase A (a-Gal A) associated with Fabry disease); ATP7B associated with Wilson's Disease; beta-glucocerebrosidase, associated with Gaucher disease type 2 and 3; peroxisome membrane protein 70 kDa,

25 associated with Zellweger syndrome; arylsulfatase A (ARSA) associated with metachromatic leukodystrophy, galactocerebrosidase (GALC) enzyme associated with Krabbe disease, alpha-glucosidase (GAA) associated with Pompe disease; sphingomyelinase (SMPD1) gene associated with Nieman Pick disease type A; argininosuccinate synthase associated with adult onset type II citrullinemia (CTLN2); carbamoyl-phosphate synthase 1 (CPS1)

30 associated with urea cycle disorders; survival motor neuron (SMN) protein, associated with spinal muscular atrophy; ceramidase associated with Farber lipogranulomatosis; b-hexosaminidase associated with GM2 gangliosidosis and Tay-Sachs and Sandhoff diseases; aspartylglucosaminidase associated with aspartyl-glucosaminuria; a-fucosidase associated

with fucosidosis; α -mannosidase associated with alpha-mannosidosis; porphobilinogen deaminase, associated with acute intermittent porphyria (AIP); alpha-1 antitrypsin for treatment of alpha-1 antitrypsin deficiency (emphysema); erythropoietin for treatment of anemia due to thalassemia or to renal failure; vascular endothelial growth factor, 5 angiopoietin-1, and fibroblast growth factor for the treatment of ischemic diseases; thrombomodulin and tissue factor pathway inhibitor for the treatment of occluded blood vessels as seen in, for example, atherosclerosis, thrombosis, or embolisms; aromatic amino acid decarboxylase (AADC), and tyrosine hydroxylase (TH) for the treatment of Parkinson's disease; the beta adrenergic receptor, anti-sense to, or a mutant form of, phospholamban, the 10 sarco(endo)plasmic reticulum adenosine triphosphatase-2 (SERCA2), and the cardiac adenylyl cyclase for the treatment of congestive heart failure; a tumor suppressor gene such as p53 for the treatment of various cancers; a cytokine such as one of the various interleukins for the treatment of inflammatory and immune disorders and cancers; dystrophin or 15 minidystrophin and utrophin or miniutrophin for the treatment of muscular dystrophies; and, insulin or GLP-1 for the treatment of diabetes. Additional genes and diseases of interest include, e.g., dystonin gene related diseases such as Hereditary Sensory and Autonomic Neuropathy Type VI (the DST gene encodes dystonin; dual AAV vectors may be required due to the size of the protein (~7570 aa); SCN9A related diseases, in which loss of function mutants cause inability to feel pain and gain of function mutants cause pain conditions, such 20 as erythromelagia. Another condition is Charcot-Marie-Tooth type 1F and 2E due to mutations in the NEFL gene (neurofilament light chain). characterized by a progressive peripheral motor and sensory neuropathy with variable clinical and electrophysiologic expression. In certain embodiments, the vectors described herein may be used in treatment of mucopolysaccharidoses (MPS) disorders. Such vectors may contain carry a nucleic acid 25 sequence encoding α -L-iduronidase (IDUA) for treating MPS I (Hurler, Hurler-Scheie and Scheie syndromes); a nucleic acid sequence encoding iduronate-2-sulfatase (IDS) for treating MPS II (Hunter syndrome); a nucleic acid sequence encoding sulfamidase (SGSH) for treating MPSIII A, B, C, and D (Sanfilippo syndrome); a nucleic acid sequence encoding N-acetylgalactosamine-6-sulfate sulfatase (GALNS) for treating MPS IV A and B (Morquio 30 syndrome); a nucleic acid sequence encoding arylsulfatase B (ARSB) for treating MPS VI (Maroteaux-Lamy syndrome); a nucleic acid sequence encoding hyaluronidase for treating MPSI IX (hyaluronidase deficiency) and a nucleic acid sequence encoding beta-

glucuronidase for treating MPS VII (Sly syndrome). See, e.g., www.orpha.net/consor/cgi-bin/Disease_Search_List.php; rarediseases.info.nih.gov/diseases.

Examples of other suitable genes may include, e.g., hormones and growth and differentiation factors including, without limitation, insulin, glucagon, glucagon-like peptide -1 (GLP1), growth hormone (GH), parathyroid hormone (PTH), growth hormone releasing factor (GRF), follicle stimulating hormone (FSH), luteinizing hormone (LH), human chorionic gonadotropin (hCG), vascular endothelial growth factor (VEGF), angiopoietins, angiostatin, granulocyte colony stimulating factor (GCSF), erythropoietin (EPO) (including, e.g., human, canine or feline epo), connective tissue growth factor (CTGF), neutrophic factors including, e.g., basic fibroblast growth factor (bFGF), acidic fibroblast growth factor (aFGF), epidermal growth factor (EGF), platelet-derived growth factor (PDGF), insulin growth factors I and II (IGF-I and IGF-II), any one of the transforming growth factor α superfamily, including TGF α , activins, inhibins, or any of the bone morphogenic proteins (BMP) BMPs 1-15, any one of the heregluin/neuregulin/ARIA/neu differentiation factor (NDF) family of growth factors, nerve growth factor (NGF), brain-derived neurotrophic factor (BDNF), neurotrophins NT-3 and NT-4/5, ciliary neurotrophic factor (CNTF), glial cell line derived neurotrophic factor (GDNF), neurturin, agrin, any one of the family of semaphorins/collapsins, netrin-1 and netrin-2, hepatocyte growth factor (HGF), ephrins, noggin, sonic hedgehog and tyrosine hydroxylase.

Other useful transgene products include proteins that regulate the immune system including, without limitation, cytokines and lymphokines such as thrombopoietin (TPO), interleukins (IL) IL-1 through IL-36 (including, e.g., human interleukins IL-1, IL-1 α , IL-1 β , IL-2, IL-3, IL-4, IL-6, IL-8, IL-12, IL-11, IL-12, IL-13, IL-18, IL-31, IL-35), monocyte chemoattractant protein, leukemia inhibitory factor, granulocyte-macrophage colony stimulating factor, Fas ligand, tumor necrosis factors α and β , interferons α , β , and γ , stem cell factor, flk-2/flt3 ligand. Gene products produced by the immune system are also useful in the invention. These include, without limitations, immunoglobulins IgG, IgM, IgA, IgD and IgE, chimeric immunoglobulins, humanized antibodies, single chain antibodies, T cell receptors, chimeric T cell receptors, single chain T cell receptors, class I and class II MHC molecules, as well as engineered immunoglobulins and MHC molecules. For example, in certain embodiments, the rAAV antibodies may be designed to delivery canine or feline antibodies, e.g., such as anti-IgE, anti-IL31, anti-CD20, anti-NGF, anti-GnRH. Useful gene products also include complement regulatory proteins such as complement regulatory

proteins, membrane cofactor protein (MCP), decay accelerating factor (DAF), CR1, CF2, CD59, and C1 esterase inhibitor (C1-INH). Still other useful gene products include any one of the receptors for the hormones, growth factors, cytokines, lymphokines, regulatory proteins and immune system proteins. The invention encompasses receptors for cholesterol 5 regulation and/or lipid modulation, including the low density lipoprotein (LDL) receptor, high density lipoprotein (HDL) receptor, the very low density lipoprotein (VLDL) receptor, and scavenger receptors. The invention also encompasses gene products such as members of the steroid hormone receptor superfamily including glucocorticoid receptors and estrogen receptors, Vitamin D receptors and other nuclear receptors. In addition, useful gene products 10 include transcription factors such as jun, fos, max, mad, serum response factor (SRF), AP-1, AP2, myb, MyoD and myogenin, ETS-box containing proteins, TFE3, E2F, ATF1, ATF2, ATF3, ATF4, ZF5, NFAT, CREB, HNF-4, C/EBP, SP1, CCAAT-box binding proteins, interferon regulation factor (IRF-1), Wilms tumor protein, ETS-binding protein, STAT, 15 GATA-box binding proteins, e.g., GATA-3, and the forkhead family of winged helix proteins.

Methods for sequencing a protein, peptide, or polypeptide (e.g., as an immunoglobulin) are known to those of skill in the art. Once the sequence of a protein is known, there are web-based and commercially available computer programs, as well as service based companies which back translate the amino acids sequences to nucleic acid 20 coding sequences. See, e.g., backtranseq by EMBOSS, available at www.ebi.ac.uk/Tools/st/; Gene Infinity, available at geneinfinity.org/sms/sms_backtranslation.html); ExPasy, available at expasy.org/tools/. In one embodiment, the RNA and/or cDNA coding sequences are designed for optimal expression in human cells.

25 In certain embodiments, the compositions provided herein are useful for a method for modulating neuronal degeneration and/or decrease secondary dorsal spinal cord axonal degeneration following intrathecal or systemic gene therapy administration. Thus, while the compositions provided herein are particularly useful for delivery of gene therapy to the CNS, they may also be useful for other routes of delivery, including e.g. systemic IV delivery, 30 where high doses of the gene therapy may result in DRG transduction and toxicity. The method involves delivering a composition comprising an expression cassette or vector genome comprising the transgene and miRNA target(s) to a patient.

In certain embodiments, the compositions provided herein are useful in methods for repressing transgene expression in the DRG. In certain embodiments, the method comprises delivering an expression cassette or vector genome that includes a miR-183 target sequence to repress transgene expression levels in the DRG. In certain embodiments, the method 5 enhances expression in one or more cells present in the CNS selected from one or more of pyramidal neurons, purkinje neurons, granule cells, spindle neurons, interneuron cells, astrocytes, oligodendrocytes, microglia, and/or ependymal cells.

In certain embodiments, provided is a method useful for delivering and/or enhancing expression of transgene in lower motor neurons the retina, inner ear, and olfactory receptors 10 comprising delivering an expression cassette or vector genome that includes a transgene operably linked to one or more miR-183 target sequences and/or more miR-183 target sequences. In certain embodiments, the cells or tissues may be one or more of liver, or heart.

In yet another embodiment, provided is a method comprising delivering an 15 expression cassette or vector genome to cells present in the CNS wherein the expression cassette or vector genome comprises one or more miR-183 target sequences and lacks a transgene (i.e. a sequence encoding a heterologous gene product). In such embodiments, delivery of miR-183 to cells of the CNS is achieved. In certain embodiments, delivery of an expression cassette or vector genome comprising miR-183 sequences results in repression of DRG expression and enhanced gene expression in certain other cells present in the CNS.

20 In certain embodiments, the compositions provided herein are useful in methods for enhancing expression of a transgene in a cell outside the CNS. In certain embodiments, methods for enhancing expression in a cell outside the CNS comprise delivering an expression cassette or vector genome that includes a miR-182 target sequence to a patient.

25 In one embodiment, the suspension has a pH of about 6.8 to about 7.32. Suitable volumes for delivery of these doses and concentrations may be determined by one of skill in the art. For example, volumes of about 1 μ L to 150 mL may be selected, with the higher volumes being selected for adults. Typically, for newborn infants a suitable volume is about 0.5 mL to about 10 mL, for older infants, about 0.5 mL to about 15 mL may be selected. For toddlers, a volume of about 0.5 mL to about 20 mL may be selected. For 30 children, volumes of up to about 30 mL may be selected. For pre-teens and teens, volumes up to about 50 mL may be selected. In still other embodiments, a patient may receive an intrathecal administration in a volume of about 5 mL to about 15 mL are selected, or about 7.5 mL to about 10 mL. Other suitable volumes and dosages may be determined. The

dosage will be adjusted to balance the therapeutic benefit against any side effects and such dosages may vary depending upon the therapeutic application for which the recombinant vector is employed.

In one embodiment, the composition comprising an rAAV as described herein is 5 administrable at a dose of about 1×10^9 GC per gram of brain mass to about 1×10^{14} GC per gram of brain mass. In certain embodiments, the rAAV is co-administered systemically at a dose of about 1×10^9 GC per kg body weight to about 1×10^{13} GC per kg body weight

In one embodiment, the subject is delivered a therapeutically effective amount of the expression cassettes described herein. As used herein, a “therapeutically effective amount” 10 refers to the amount of the expression cassette comprising the nucleic acid sequence encoding the gene product and the miRNA target sequences which delivers and expresses in the target cells and which specifically detargets DRG expression.

The use of rAAV for delivering for the treatment of various conditions have been 15 previously described. The expression cassettes for these rAAVs can be modified to include miRNA target sequences described herein (including, e.g., miR-183 target sequences, miR-182 target sequences and miR-96 target sequences, or combinations thereof) to, for example, reduce transgene expression in DRG and/or reduce or eliminate DRG toxicity and/or axonopathy. Examples of rAAV vector genomes that can be modified to include miRNA target sequences include the genes described in WO 2017/136500 (MPSI), WO 2017/181113 20 (MPSII), WO 2019/108857 (MPSIIIA), WO 2019/108856 (MPSIIB), WO 2017/106354 (SMN1), WO 2018/160585 (SMN1), WO 2018/209205 (Batten disease), WO 2015/164723 (AAV-mediated delivery of anti-HER2 antibody), WO2015/138348 (OTC), WO 2015/164778 (LDLR variants for FH); WO2017/106345 (Crigler-Najjar), WO 2017/106326 (anti-PCSK9 Abs), WO 2017/180857 (hemophilia A, Factor VIII), WO 2017/180861 25 (hemophilia B, Factor IX), as well as vectors in trials for treatment of Myotubular Myopathy (such as AT132, AAV8, Audentes).

In certain embodiments, an AAV.alpha-L-iduronidase (AAV.IDUA) gene therapy vector comprises a vector genome comprising at least one, at least two, at least three, or at 30 least four miR target sequences of the miRNA183 cluster (including miR-183, miR-182, and miR183 target sequences, or combinations thereof) operably linked to the coding sequence for the IDUA gene (see, e.g., nt 1938-3908 of SEQ ID NO: 15). In certain embodiments, the vector genome comprises multiple copies of the same miR target sequence each separated by a spacer which may be the same or which may differ from each other. In another

embodiment, the vector genome comprises three to six copies of a miR183 cluster target sequence, optionally wherein one or more of the target sequences is at least about 80% to about 99% complementarity to a miR-183 cluster member. In another embodiment, the vector comprises one, two, three, or four copies of a miR183 target sequence. Such a vector 5 genome may optionally contain additional target sequences that correspond to members of the miR183 cluster. In certain embodiments, the vector genome contains a single miR target sequence for a miR183 cluster member. In certain embodiments, the vector genome contains two miR target sequences for miR183 cluster members and optionally at least one spacer. In certain embodiments, the vector contains three miR target sequences for miR183 cluster 10 members and optionally at least two spacers. In certain embodiments, the vector genome contains two or more miR target sequences for the miR183 cluster which differ in sequence from one another. In certain embodiments, the vector genomes described herein are carried by a non-AAV vector.

In certain embodiments, an AAV.iduronate-2-sulfatase (IDS) (AAV.IDS) gene 15 therapy vector comprises a vector genome comprising at least one, at least two, at least three, or at least four miR target sequences of the miRNA 183 cluster (including miR-183, miR-182, and miR183 target sequences, or combinations thereof) operably linked to the coding sequence for the IDS gene. In certain embodiments, the vector genome comprises multiple copies of the same miR target sequence each separated by a spacer which may be the same or which may differ from each other. In another embodiment, the vector genome comprises 20 three to six copies of a miR183 cluster target sequence, optionally wherein one or more of the target sequences is at least about 80% to about 99% complementarity to a miR-183 cluster member. In another embodiment, the vector comprises one, two, three, or four copies of a miR183 target sequence. Such a vector genome may optionally contain additional target 25 sequences that correspond to members of the miR183 cluster. In certain embodiments, the vector genome contains a single miR target sequence for a miR183 cluster member. In certain embodiments, the vector genome contains two miR target sequences for miR183 cluster members and optionally at least one spacer. In certain embodiments, the vector contains three miR target sequences for miR183 cluster members and optionally at least two 30 spacers. In certain embodiments, the vector genome contains two or more miR target sequences for the miR183 cluster which differ in sequence from one another. In certain embodiments, the vector genomes described herein are carried by a non-AAV vector.

In certain embodiments, an AAV.N-sulfoglucosamine sulfohydrolase (AAV.SGSH) gene therapy vector comprises a vector genome comprising at least one, at least two, at least three, or at least four miR target sequences of the miRNA183 cluster (including miR-183, miR-182, and miR183 target sequences, or combinations thereof) operably linked to the 5 coding sequence for the SGSH gene. In certain embodiments, the vector genome comprises multiple copies of the same miR target sequence each separated by a spacer which may be the same or which may differ from each other. In another embodiment, the vector genome comprises three to six copies of a miR183 cluster target sequence, optionally wherein one or more of the target sequences is at least about 80% to about 99% complementarity to a miR- 10 183 cluster member. In another embodiment, the vector comprises one, two, three, or four copies of a miR183 target sequence. Such a vector genome may optionally contain additional target sequences that correspond to members of the miR183 cluster. In certain embodiments, the vector genome contains a single miR target sequence for a miR183 cluster member. In certain embodiments, the vector genome contains two miR target sequences for miR183 15 cluster members and optionally at least one spacer. In certain embodiments, the vector contains three miR target sequences for miR183 cluster members and optionally at least two spacers. In certain embodiments, the vector genome contains two or more miR target sequences for the miR183 cluster which differ in sequence from one another. In certain embodiments, the vector genomes described herein are carried by a non-AAV vector.

20 In certain embodiments, an AAV.N-acetyl-alpha-D-glucosaminidase (AAV.NAGLU) gene therapy vector comprises a vector genome comprising at least one, at least two, at least three, or at least four miR target sequences of the miRNA183 cluster (including miR-183, miR-182, and miR183 target sequences, or combinations thereof) operably linked to the coding sequence for the NAGLU gene. In certain embodiments, the 25 vector genome comprises multiple copies of the same miR target sequence each separated by a spacer which may be the same or which may differ from each other. In another embodiment, the vector genome comprises three to six copies of a miR183 cluster target sequence, optionally wherein one or more of the target sequences is at least about 80% to about 99% complementarity to a miR-183 cluster member. In another embodiment, the vector 30 comprises one, two, three, or four copies of a miR183 target sequence. Such a vector genome may optionally contain additional target sequences that correspond to members of the miR183 cluster. In certain embodiments, the vector genome contains a single miR target sequence for a miR183 cluster member. In certain embodiments, the vector genome contains

two miR target sequences for miR183 cluster members and optionally at least one spacer. In certain embodiments, the vector contains three miR target sequences for miR183 cluster members and optionally at least two spacers. In certain embodiments, the vector genome contains two or more miR target sequences for the miR183 cluster which differ in sequence 5 from one another. In certain embodiments, the vector genomes described herein are carried by a non-AAV vector.

In certain embodiments, an AAV.survival motor neuron 1 (AAV.SMN1) gene therapy vector comprises a vector genome comprising at least one, at least two, at least three, or at least four miR target sequences of the miRNA183 cluster (including miR-183, miR-10 182, and miR183 target sequences, or combinations thereof) operably linked to the coding sequence for the SMN1 gene. In certain embodiments, the vector genome comprises multiple copies of the same miR target sequence each separated by a spacer which may be the same or which may differ from each other. In another embodiment, the vector genome comprises three to six copies of a miR183 cluster target sequence, optionally wherein one or more of 15 the target sequences is at least about 80% to about 99% complementarity to a miR-183 cluster member. In another embodiment, the vector comprises one, two, three, or four copies of a miR183 target sequence. Such a vector genome may optionally contain additional target sequences that correspond to members of the miR183 cluster. In certain embodiments, the vector genome contains a single miR target sequence for a miR183 cluster member. In 20 certain embodiments, the vector genome contains two miR target sequences for miR183 cluster members and optionally at least one spacer. In certain embodiments, the vector contains three miR target sequences for miR183 cluster members and optionally at least two spacers. In certain embodiments, the vector genome contains two or more miR target sequences for the miR183 cluster which differ in sequence from one another. In certain 25 embodiments, the vector genomes described herein are carried by a non-AAV vector.

In certain embodiments, an AAV.tripeptidyl peptidase 1 (AAV.TPP1) gene therapy vector comprises a vector genome comprising at least one, at least two, at least three, or at least four miR target sequences of the miRNA183 cluster (including miR-183, miR-182, and miR183 target sequences, or combinations thereof) operably linked to the coding sequence 30 for the TPP1 gene. In certain embodiments, the vector genome comprises multiple copies of the same miR target sequence each separated by a spacer which may be the same or which may differ from each other. In another embodiment, the vector genome comprises three to six copies of a miR183 cluster target sequence, optionally wherein one or more of the target

sequences is at least about 80% to about 99% complementarity to a miR-183 cluster member. In another embodiment, the vector comprises one, two, three, or four copies of a miR183 target sequence. Such a vector genome may optionally contain additional target sequences that correspond to members of the miR183 cluster. In certain embodiments, the 5 vector genome contains a single miR target sequence for a miR183 cluster member. In certain embodiments, the vector genome contains two miR target sequences for miR183 cluster members and optionally at least one spacer. In certain embodiments, the vector contains three miR target sequences for miR183 cluster members and optionally at least two spacers. In certain embodiments, the vector genome contains two or more miR target 10 sequences for the miR183 cluster which differ in sequence from one another. In certain embodiments, the vector genomes described herein are carried by a non-AAV vector.

In certain embodiments, an AAV.anti-human epidermal growth factor receptor 2 antibody (AAV.anti-HER2) gene therapy vector comprises a vector genome comprising at least one, at least two, at least three, or at least four miR target sequences of the miRNA183 cluster (including miR-183, miR-182, and miR183 target sequences, or combinations thereof) operably linked to the coding sequence for the anti-HER2 antibody. In certain embodiments, the vector genome comprises multiple copies of the same miR target sequence each separated by a spacer which may be the same or which may differ from each other. In another embodiment, the vector genome comprises three to six copies of a miR183 cluster target sequence, optionally wherein one or more of the target sequences is at least about 80% to about 99% complementarity to a miR-183 cluster member. In another embodiment, the vector comprises one, two, three, or four copies of a miR183 target sequence. Such a vector genome may optionally contain additional target sequences that correspond to members of the miR183 cluster. In certain embodiments, the vector genome contains a single miR target 15 sequence for a miR183 cluster member. In certain embodiments, the vector genome contains two miR target sequences for miR183 cluster members and optionally at least one spacer. In certain embodiments, the vector contains three miR target sequences for miR183 cluster members and optionally at least two spacers. In certain embodiments, the vector genome contains two or more miR target sequences for the miR183 cluster which differ in sequence 20 from one another. In certain embodiments, the vector genomes described herein are carried by a non-AAV vector.

In certain embodiments, an AAV.ornithine transcarbamylase (AAV.OTC) gene therapy vector comprises a vector genome comprising at least one, at least two, at least three,

or at least four miR target sequences of the miRNA183 cluster (including miR-183, miR-182, and miR183 target sequences, or combinations thereof) operably linked to the coding sequence for the OTC gene. In certain embodiments, the vector genome comprises multiple copies of the same miR target sequence each separated by a spacer which may be the same 5 or which may differ from each other. In another embodiment, the vector genome comprises three to six copies of a miR183 cluster target sequence, optionally wherein one or more of the target sequences is at least about 80% to about 99% complementarity to a miR-183 cluster member. In another embodiment, the vector comprises one, two, three, or four copies of a miR183 target sequence. Such a vector genome may optionally contain additional target 10 sequences that correspond to members of the miR183 cluster. In certain embodiments, the vector genome contains a single miR target sequence for a miR183 cluster member. In certain embodiments, the vector genome contains two miR target sequences for miR183 cluster members and optionally at least one spacer. In certain embodiments, the vector contains three miR target sequences for miR183 cluster members and optionally at least two 15 spacers. In certain embodiments, the vector genome contains two or more miR target sequences for the miR183 cluster which differ in sequence from one another. In certain embodiments, the vector genomes described herein are carried by a non-AAV vector.

In certain embodiments, an AAV.low-density lipoprotein receptor (AAV.LDLR) gene therapy vector comprises a vector genome comprising at least one, at least two, at least 20 three, or at least four miR target sequences of the miRNA183 cluster (including miR-183, miR-182, and miR183 target sequences, or combinations thereof) operably linked to the coding sequence for the LDLR gene. In certain embodiments, the vector genome comprises multiple copies of the same miR target sequence each separated by a spacer which may be the same or which may differ from each other. In another embodiment, the vector genome comprises three to six copies of a miR183 cluster target sequence, optionally wherein one or 25 more of the target sequences is at least about 80% to about 99% complementarity to a miR-183 cluster member. In another embodiment, the vector comprises one, two, three, or four copies of a miR183 target sequence. Such a vector genome may optionally contain additional target sequences that correspond to members of the miR183 cluster. In certain embodiments, the vector genome contains a single miR target sequence for a miR183 cluster member. In certain 30 embodiments, the vector genome contains two miR target sequences for miR183 cluster members and optionally at least one spacer. In certain embodiments, the vector contains three miR target sequences for miR183 cluster members and optionally at least two

spacers. In certain embodiments, the vector genome contains two or more miR target sequences for the miR183 cluster which differ in sequence from one another. In certain embodiments, the vector genomes described herein are carried by a non-AAV vector.

In certain embodiments, an AAV.uridine diphosphate glucuronosyl transferase 1A1 (AAV.UGT1A1) gene therapy vector comprises a vector genome comprising at least one, at least two, at least three, or at least four miR target sequences of the miRNA183 cluster (including miR-183, miR-182, and miR183 target sequences, or combinations thereof) operably linked to the coding sequence for the UGT1A1 gene. In certain embodiments, the vector genome comprises multiple copies of the same miR target sequence each separated by a spacer which may be the same or which may differ from each other. In another embodiment, the vector genome comprises three to six copies of a miR183 cluster target sequence, optionally wherein one or more of the target sequences is at least about 80% to about 99% complementarity to a miR-183 cluster member. In another embodiment, the vector comprises one, two, three, or four copies of a miR183 target sequence. Such a vector genome may optionally contain additional target sequences that correspond to members of the miR183 cluster. In certain embodiments, the vector genome contains a single miR target sequence for a miR183 cluster member. In certain embodiments, the vector genome contains two miR target sequences for miR183 cluster members and optionally at least one spacer. In certain embodiments, the vector contains three miR target sequences for miR183 cluster members and optionally at least two spacers. In certain embodiments, the vector genome contains two or more miR target sequences for the miR183 cluster which differ in sequence from one another. In certain embodiments, the vector genomes described herein are carried by a non-AAV vector.

In certain embodiments, an AAV.anti-proprotein convertase subtilisin/kexin type 9 antibody (AAV.anti-PCSK9 Ab) gene therapy vector comprises a vector genome comprising at least one, at least two, at least three, or at least four miR target sequences of the miRNA183 cluster (including miR-183, miR-182, and miR183 target sequences, or combinations thereof) operably linked to the coding sequence for the anti-PCSK9 Ab. In certain embodiments, the vector genome comprises multiple copies of the same miR target sequence each separated by a spacer which may be the same or which may differ from each other. In another embodiment, the vector genome comprises three to six copies of a miR183 cluster target sequence, optionally wherein one or more of the target sequences is at least about 80% to about 99% complementarity to a miR-183 cluster member. In another

embodiment, the vector comprises one, two, three, or four copies of a miR183 target sequence. Such a vector genome may optionally contain additional target sequences that correspond to members of the miR183 cluster. In certain embodiments, the vector genome contains a single miR target sequence for a miR183 cluster member. In certain embodiments, 5 the vector genome contains two miR target sequences for miR183 cluster members and optionally at least one spacer. In certain embodiments, the vector contains three miR target sequences for miR183 cluster members and optionally at least two spacers. In certain embodiments, the vector genome contains two or more miR target sequences for the miR183 cluster which differ in sequence from one another. In certain embodiments, the vector 10 genomes described herein are carried by a non-AAV vector.

In certain embodiments, an AAV.Factor VIII (AAV.FVIII) gene therapy vector comprises a vector genome comprising at least one, at least two, at least three, or at least four miR target sequences of the miRNA183 cluster (including miR-183, miR-182, and miR183 target sequences, or combinations thereof) operably linked to the coding sequence for the 15 FVIII gene. In certain embodiments, the vector genome comprises multiple copies of the same miR target sequence each separated by a spacer which may be the same or which may differ from each other. In another embodiment, the vector genome comprises three to six copies of a miR183 cluster target sequence, optionally wherein one or more of the target sequences is at least about 80% to about 99% complementarity to a miR-183 cluster member. In another embodiment, the vector comprises one, two, three, or four copies of a 20 miR183 target sequence. Such a vector genome may optionally contain additional target sequences that correspond to members of the miR183 cluster. In certain embodiments, the vector genome contains a single miR target sequence for a miR183 cluster member. In certain embodiments, the vector genome contains two miR target sequences for miR183 cluster members and optionally at least one spacer. In certain embodiments, the vector 25 contains three miR target sequences for miR183 cluster members and optionally at least two spacers. In certain embodiments, the vector genome contains two or more miR target sequences for the miR183 cluster which differ in sequence from one another. In certain embodiments, the vector genomes described herein are carried by a non-AAV vector.

30 In certain embodiments, an AAV.Factor IX (AAV.IX) gene therapy vector comprises a vector genome comprising at least one, at least two, at least three, or at least four miR target sequences of the miRNA183 cluster (including miR-183, miR-182, and miR183 target sequences, or combinations thereof) operably linked to the coding sequence for the FIX

gene. In certain embodiments, the vector genome comprises multiple copies of the same miR target sequence each separated by a spacer which may be the same or which may differ from each other. In another embodiment, the vector genome comprises three to six copies of a miR183 cluster target sequence, optionally wherein one or more of the target sequences is at least about 80% to about 99% complementarity to a miR-183 cluster member. In another embodiment, the vector comprises one, two, three, or four copies of a miR183 target sequence. Such a vector genome may optionally contain additional target sequences that correspond to members of the miR183 cluster. In certain embodiments, the vector genome contains a single miR target sequence for a miR183 cluster member. In certain embodiments, 5 the vector genome contains two miR target sequences for miR183 cluster members and optionally at least one spacer. In certain embodiments, the vector contains three miR target sequences for miR183 cluster members and optionally at least two spacers. In certain embodiments, the vector genome contains two or more miR target sequences for the miR183 cluster which differ in sequence from one another. In certain embodiments, the vector 10 genomes described herein are carried by a non-AAV vector.

15

In certain embodiments, an AAV.myotubularin 1 (AAV.MTM1) gene therapy vector comprises a vector genome comprising at least one, at least two, at least three, or at least four miR target sequences of the miRNA183 cluster (including miR-183, miR-182, and miR183 target sequences, or combinations thereof) operably linked to the coding sequence for the MTM1 gene. In certain embodiments, the vector genome comprises multiple copies of the same miR target sequence each separated by a spacer which may be the same or which may differ from each other. In another embodiment, the vector genome comprises three to six copies of a miR183 cluster target sequence, optionally wherein one or more of the target sequences is at least about 80% to about 99% complementarity to a miR-183 cluster member. In another embodiment, the vector comprises one, two, three, or four copies of a miR183 target sequence. Such a vector genome may optionally contain additional target sequences that correspond to members of the miR183 cluster. In certain embodiments, the vector genome contains a single miR target sequence for a miR183 cluster member. In certain embodiments, the vector genome contains two miR target sequences for miR183 cluster members and optionally at least one spacer. In certain embodiments, the vector 20 contains three miR target sequences for miR183 cluster members and optionally at least two spacers. In certain embodiments, the vector genome contains two or more miR target 25 sequences for miR183 cluster members and optionally at least two spacers. In certain embodiments, the vector genome contains two or more miR target 30 sequences for miR183 cluster members and optionally at least two spacers.

sequences for the miR183 cluster which differ in sequence from one another. In certain embodiments, the vector genomes described herein are carried by a non-AAV vector.

In one embodiment, the expression cassette is in a vector genome delivered in an amount of about 1×10^9 GC per gram of brain mass to about 1×10^{13} genome copies (GC) per gram (g) of brain mass, including all integers or fractional amounts within the range and the endpoints. In another embodiment, the dosage is 1×10^{10} GC per gram of brain mass to about 1×10^{13} GC per gram of brain mass. In specific embodiments, the dose of the vector administered to a patient is at least about 1.0×10^9 GC/g, about 1.5×10^9 GC/g, about 2.0×10^9 GC/g, about 2.5×10^9 GC/g, about 3.0×10^9 GC/g, about 3.5×10^9 GC/g, about 4.0×10^9 GC/g, about 4.5×10^9 GC/g, about 5.0×10^9 GC/g, about 5.5×10^9 GC/g, about 6.0×10^9 GC/g, about 6.5×10^9 GC/g, about 7.0×10^9 GC/g, about 7.5×10^9 GC/g, about 8.0×10^9 GC/g, about 8.5×10^9 GC/g, about 9.0×10^9 GC/g, about 9.5×10^9 GC/g, about 1.0×10^{10} GC/g, about 1.5×10^{10} GC/g, about 2.0×10^{10} GC/g, about 2.5×10^{10} GC/g, about 3.0×10^{10} GC/g, about 3.5×10^{10} GC/g, about 4.0×10^{10} GC/g, about 4.5×10^{10} GC/g, about 5.0×10^{10} GC/g, about 5.5×10^{10} GC/g, about 6.0×10^{10} GC/g, about 6.5×10^{10} GC/g, about 7.0×10^{10} GC/g, about 7.5×10^{10} GC/g, about 8.0×10^{10} GC/g, about 8.5×10^{10} GC/g, about 9.0×10^{10} GC/g, about 9.5×10^{10} GC/g, about 1.0×10^{11} GC/g, about 1.5×10^{11} GC/g, about 2.0×10^{11} GC/g, about 2.5×10^{11} GC/g, about 3.0×10^{11} GC/g, about 3.5×10^{11} GC/g, about 4.0×10^{11} GC/g, about 4.5×10^{11} GC/g, about 5.0×10^{11} GC/g, about 5.5×10^{11} GC/g, about 6.0×10^{11} GC/g, about 6.5×10^{11} GC/g, about 7.0×10^{11} GC/g, about 7.5×10^{11} GC/g, about 8.0×10^{11} GC/g, about 8.5×10^{11} GC/g, about 9.0×10^{11} GC/g, about 9.5×10^{11} GC/g, about 1.0×10^{12} GC/g, about 1.5×10^{12} GC/g, about 2.0×10^{12} GC/g, about 2.5×10^{12} GC/g, about 3.0×10^{12} GC/g, about 3.5×10^{12} GC/g, about 4.0×10^{12} GC/g, about 4.5×10^{12} GC/g, about 5.0×10^{12} GC/g, about 5.5×10^{12} GC/g, about 6.0×10^{12} GC/g, about 6.5×10^{12} GC/g, about 7.0×10^{12} GC/g, about 7.5×10^{12} GC/g, about 8.0×10^{12} GC/g, about 8.5×10^{12} GC/g, about 9.0×10^{12} GC/g, about 9.5×10^{12} GC/g, about 1.0×10^{13} GC/g, about 1.5×10^{13} GC/g, about 2.0×10^{13} GC/g, about 2.5×10^{13} GC/g, about 3.0×10^{13} GC/g, about 3.5×10^{13} GC/g, about 4.0×10^{13} GC/g, about 4.5×10^{13} GC/g, about 5.0×10^{13} GC/g, about 5.5×10^{13} GC/g, about 6.0×10^{13} GC/g, about 6.5×10^{13} GC/g, about 7.0×10^{13} GC/g, about 7.5×10^{13} GC/g, about 8.0×10^{13} GC/g, about 8.5×10^{13} GC/g, about 9.0×10^{13} GC/g, about 9.5×10^{13} GC/g, or about 1.0×10^{14} GC/g brain mass.

In certain embodiments, the miR target sequence -containing compositions provided herein minimize the dose, duration, and/or amount of immunosuppressive co-therapy

required by the patient. Currently, immunosuppressants for such co-therapy include, but are not limited to, a glucocorticoid, steroids, antimetabolites, T-cell inhibitors, a macrolide (e.g., a rapamycin or rapalog), and cytostatic agents including an alkylating agent, an anti-metabolite, a cytotoxic antibiotic, an antibody, or an agent active on immunophilin. The 5 immune suppressant may include a nitrogen mustard, nitrosourea, platinum compound, methotrexate, azathioprine, mercaptopurine, fluorouracil, dactinomycin, an anthracycline, mitomycin C, bleomycin, mithramycin, IL-2 receptor- (CD25-) or CD3-directed antibodies, anti-IL-2 antibodies, ciclosporin, tacrolimus, sirolimus, IFN- β , IFN- γ , an opioid, or TNF- α (tumor necrosis factor-alpha) binding agent. In certain embodiments, the 10 immunosuppressive therapy may be started 0, 1, 2, 7, or more days prior to the gene therapy administration. Such therapy may involve co-administration of two or more drugs, the (e.g., prednisone, micophenolate mofetil (MMF) and/or sirolimus (i.e., rapamycin)) on the same day. One or more of these drugs may be continued after gene therapy administration, at the same dose or an adjusted dose. Such therapy may be for about 1 week (7 days), about 60 In 15 certain embodiments, the miR target sequence -containing compositions provided herein eliminate the need for immunosuppressive therapy prior to, during, or following delivery of a gene therapy (e.g., rAAV) vector.

In one embodiment, a composition comprising the expression cassette as described herein is administrated once to the subject in need. In certain embodiments, the expression 20 cassette is delivered via an rAAV.

It should be understood that the compositions in the method described herein are intended to be applied to other compositions, regiments, aspects, embodiments and methods described across the Specification.

25 6. Kit

In certain embodiments, a kit is provided which includes a concentrated expression cassette (e.g., in a viral or non-viral vector) suspended in a formulation (optionally frozen), optional dilution buffer, and devices and components required for intrathecal, intracerebroventricular or intracisternal administration. In another embodiment, the kit may 30 additional or alternatively include components for intravenous delivery. In one embodiment, the kit provides sufficient buffer to allow for injection. Such buffer may allow for about a 1:1 to a 1:5 dilution of the concentrated vector, or more. In other embodiments, higher or lower amounts of buffer or sterile water are included to allow for dose titration and other

adjustments by the treating clinician. In still other embodiments, one or more components of the device are included in the kit. Suitable dilution buffer is available, such as, a saline, a phosphate buffered saline (PBS) or a glycerol/PBS.

It should be understood that the compositions in kit described herein are intended to 5 be applied to other compositions, regiments, aspects, embodiments and methods described across the Specification.

7. Device

In one aspect, the compositions provided herein may be administered intrathecally 10 via the method and/or the device described, *e.g.*, in WO 2017/136500, which is incorporated herein by reference in its entirety. Alternatively, other devices and methods may be selected. In summary, the method comprises the steps of advancing a spinal needle into the cisterna magna of a patient, connecting a length of flexible tubing to a proximal hub of the spinal needle and an output port of a valve to a proximal end of the flexible tubing, and after said 15 advancing and connecting steps and after permitting the tubing to be self-primed with the patient's cerebrospinal fluid, connecting a first vessel containing an amount of isotonic solution to a flush inlet port of the valve and thereafter connecting a second vessel containing an amount of a pharmaceutical composition to a vector inlet port of the valve. After connecting the first and second vessels to the valve, a path for fluid flow is opened between 20 the vector inlet port and the outlet port of the valve and the pharmaceutical composition is injected into the patient through the spinal needle, and after injecting the pharmaceutical composition, a path for fluid flow is opened through the flush inlet port and the outlet port of the valve and the isotonic solution is injected into the spinal needle to flush the pharmaceutical composition into the patient. This method and this device may each 25 optionally be used for intrathecal delivery of the compositions provided herein.

Alternatively, other methods and devices may be used for such intrathecal delivery.

It should be understood that the compositions in the device described herein are intended to be applied to other compositions, regiments, aspects, embodiments and methods described across the Specification.

30

Examples

The invention is now described with reference to the following examples. These examples are provided for the purpose of illustration only and the invention should in no way

be construed as being limited to these examples but rather should be construed to encompass any and all variations that become evident as a result of the teaching provided herein.

Example 1: Methods

5 *Animals*

All animal procedures were approved by the Institutional Animal Care and Use Committee of the University of Pennsylvania. Rhesus macaques (*Macaca Mulatta*) were procured from Covance Research Products, Inc. (Alice, TX) and Primgen/Prelabs Primates (Hines, IL). Animals were housed in an Association for Assessment and Accreditation of 10 Laboratory Animal Care (AAALAC) International-accredited Nonhuman Primate Research Program facility at the University of Pennsylvania in stainless steel squeeze back cages. Animals received varied enrichments such as food treats, visual and auditory stimuli, manipulatives, and social interactions.

15 C56BL/6J mice (stock #000664) were purchased from the Jackson Laboratory. Animals were housed in an AAALAC International-accredited mouse barrier vivarium at the Gene Therapy Program, University of Pennsylvania, in standard caging of 2 to 5 animals per cage with enrichment (Nestlets nesting material). Cages, water bottles, and bedding substrates were autoclaved in the barrier facility and cages were changed once per week. An automatically controlled 12-hour light/dark cycle was maintained. Each dark period began at 20 1,900 hours (\pm 30 minutes). Irradiated laboratory rodent food was provided ad libitum.

17 Vectors

The AAV9.PHP.B trans plasmid (pAAV2/PHP.B) was generated with a QuikChange Lightning Site-Directed Mutagenesis Kit (Agilent Technologies, Santa Clara, CA, Cat #210515) using pAAV2/9 (Penn Vector Core) as the template, following the manufacturer's 25 manual. pAAV2/9 and pAAV2/hu68 were provided by the Penn Vector Core. AAV vectors were produced and titrated by the Penn Vector Core (as described previously by Lock, M., et al. *Hum Gene Ther* 21:1259-1271, 2010). Briefly, HEK293 cells were triple-transfected and the culture supernatant was harvested, concentrated, and purified with an iodixanol gradient. The purified vectors were titrated with droplet digital PCR using primers targeting the rabbit 30 beta-globin polyA sequence (as previously by Lock, M., et al. *Hum Gene Ther Methods* 25:115-125, 2014). The human alpha -L-iduronidase (hIDUA) sequence was obtained through back-translation and codon-optimization of the hIDUA isoform a precursor protein sequence NP_000194.2 and was cloned under the CB7 promoter (Penn Vector Core). Dorsal

root ganglion (DRG)-enriched microRNA sequences were selected from the public database available at mirbase.org. Four tandem repeats of the target for the DRG-enriched miR were cloned in the 3' untranslated region (UTR) of green fluorescent protein (GFP) or hIDUA cis plasmids.

5 *In vivo studies*

Mice received 1×10^{12} genome copies (GCs; 5×10^{13} GC/kg) of AAV-PHP.B, or 4×10^{12} GCs (2×10^{14} GC/kg) of AAV9 vectors encoding enhanced GFP (Penn Vector Core) with or without miR targets in 0.1 mL via the lateral tail vein and were euthanized by inhalation of CO₂ 21 days post injection. Tissues were promptly collected, starting with 10 brain, and immersion-fixed in 10% neutral buffered formalin for about 24 h, washed briefly in phosphate buffered saline (PBS), and equilibrated sequentially in 15% and 30% sucrose in PBS at 4°C. Tissues were then frozen in optimum cutting temperature embedding medium and cryosectioned for direct GFP visualization (brain were sectioned at 30 μ m, and other tissues at 8 μ m thickness). Images were acquired with a Nikon Eclipse Ti-E fluorescence 15 microscope. GFP expression in DRGs was analyzed by immunohistochemistry (IHC). Spinal columns with DRGs were fixed in formalin for 24 h, decalcified in 10% ethylenediaminetetraacetic acid (pH 7.5) until soft, and paraffin embedded following standard protocols. Sections were deparaffinized through an ethanol and xylene series, boiled for 6 min in 10 mM citrate buffer (pH 6.0) to perform antigen retrieval, blocked sequentially 20 with 2% H₂O₂ (15 min), avidin/biotin blocking reagents (15 min each; Vector Laboratories, Burlingame, CA), and blocking buffer (1% donkey serum in PBS + 0.2% Triton for 10 min) followed by incubation with primary (1 h at 37°C) and biotinylated secondary antibodies (diluted 1:500, 45 min; Jackson ImmunoResearch, West Grove, PA) diluted in blocking buffer. As rabbit antibody against GFP was used as the primary antibody (NB600-308, 25 Novus Biologicals, Centennial, CO; diluted 1:500). A Vectastain Elite ABC kit (Vector Laboratories, Burlingame, CA) with DAB as substrate allowed us to visualize bound antibodies as brown precipitate.

Non-human primates (NHP) received 3.5×10^{13} GCs of AA Vhu68.GFP vectors or 1×10^{13} GCs of AA Vhu68.hIDUA vectors in a total volume of 1 mL injected into the cisterna magna, under fluoroscopic guidance (as previously described by Katz, N., et al. Hum Gene Ther Methods 29:212-219, 2018). Period blood collection and cerebrospinal fluid (CSF) taps were performed for safety readouts. Serum chemistry, hematology, coagulation, and CSF analyses were performed by the contract facility Antech Diagnostics (Morrisville, NC).

Animals were euthanized with intravenous pentobarbital overdose and necropsied; the tissues were then harvested for comprehensive histopathologic examination. Collected tissues were immediately fixed in formalin and paraffin embedded. For histopathology, tissue sections were stained with hematoxylin and eosin following standard protocols. IHC
5 for GFP expression was carried out as described for the mouse studies but using a different antibody against GFP (goat antibody NB100-1770, Novus Biologicals; diluted 1:500, incubated overnight at 4°C). Immunostaining for hIDUA was performed using a sheep antibody against hIDUA (AF4119, R&D Systems, Minneapolis, MN; diluted 1:200) following the above protocol for IHC. In addition, sections were stained for hIDUA by
10 immunofluorescence (IF) using the same primary antibody. For IF, sections were deparaffinized and treated for antigen retrieval as described above, and then blocked with 1% donkey serum in PBS + 0.2% Triton for 15 min followed by sequential incubation with primary (2 h at room temperature, diluted 1:50) and FITC-labeled secondary (45 min; Jackson ImmunoResearch; diluted 1:100) antibodies diluted in blocking buffer. Sections
15 were mounted in Fluoromount G with DAPI as a nuclear counterstain.

In situ hybridization (ISH) was performed using probes specific for the codon-optimized RNA transcribed from the vector genome that do not bind to endogenous monkey IDUA RNA. Z-shaped probe pairs were synthesized by Life Technologies (Carlsbad, CA) and ISH was performed on paraffin sections using the Life Technologies ViewRNA ISH
20 Tissue Assay kit according to the manufacturer's protocol. The deposition of Fast Red precipitates indicating positive signals was imaged by fluorescence microscopy using a rhodamine filter set. Tissue sections with IDUA IHC were scanned for quantification purposes using an Aperio Versa slide scanner (Leica Biosystems, Buffalo Grove, IL).

Histopathology and morphometry
25 A board-certified Veterinary Pathologist who was blinded to the vector group established severity grades defined with 0 as absence of lesion, 1 as minimal (<10%), 2 mild (10-25%), 3 moderate (25-50%), 4 marked (50-95%), and 5 severe (>95%). Dorsal axonopathy scores were established in each animal from at least 3 cervical, 3 thoracic, and 3 lumbar sections; the DRG severity grades were established from at least 3 cervical, 3
30 thoracic, and 3 lumbar segments; and the median nerve score was the sum of axonopathy and fibrosis severity grades with a maximal possible score of 10 and was established on the distal and proximal portions of left and right nerves. For quantification of transgene expression, a board-certified Veterinary Pathologist counted cells immunostained with anti-GFP or anti-

hIDUA antibodies by comparing with signal from control slides obtained from untreated animals. The total number of positive cells per x20 magnification field was counted using the ImageJ cell counter tool on a minimum of five fields per structure and per animal.

Vector biodistribution

5 NHP tissue DNA was extracted with a QIAamp DNA Mini Kit (Qiagen, Germany, Cat #51306) and vector genomes were quantified by real-time PCR using Taqman reagents (Applied Biosystems, Life Technologies, Foster City, CA) and primers/probes targeting the rBG polyadenylation sequence of the vectors.

Immunology

10 Peripheral blood T-cell responses against hIDUA were measured by interferon gamma enzyme-linked immunosorbent spot assays according to previously published methods (Gao et al., 2009), using peptide libraries specific for the hIDUA transgene. Positive response criteria were >55 spot forming units per 10^6 lymphocytes and three times the medium negative control upon no stimulation. In addition, T-cell responses were assayed in 15 lymphocytes that were extracted from spleen, liver, and deep cervical lymph nodes after necropsy on study day 90. Antibodies to hIDUA were measured in serum (1:1,000 sample dilution) (as previously described by Hinderer, C., et al. Mol Ther 23:1298-1307, 2015).

20 Cytokine/Chemokine analysis: CSF samples were collected and stored at -80C until the time of analysis. CSF samples were analyzed using a Milliplex MAP kit containing the following analytes: sCD137, Eotaxin, sFasL, FGF-2, Fractalkine, Granzyme A, Granzyme B, IL-1 α , IL-2, IL-4, IL-6, IL-16, IL-17A, IL-17E/IL-25, IL-21, IL-22, IL-23, IL-28A, IL-31, IL-33, IP-10, MIP-3 α , Perforin, TNF β . CSF samples were evaluated in duplicate and 25 analyzed in a FLEXMAP 3D instrument using Luminex® xPONENT® 4.2; Bio-Plex Manager™ Software 6.1. Only samples with a %CV of less than 20% were included in the analysis.

In vitro studies

30 miR183 human microRNA expression plasmid was modified from Origene MI0000273 vector by deleting the KpnI-PstI fragment encoding GFP and partial internal ribosome entry sites. We confirmed the lack of GFP expression from the modified vector by transient transfection and anti-GFP immunoblotting. We performed polyethylenimine-mediated transient transfection in HEK293 cells with GFP cis-plasmids harboring microRNA binding sites located in the 3'-UTR of the GFP expression cassette. At 72 hours post-transfection, we lysed the cells in 50 mM Tris-HCl, pH 8.0, 150 mM NaCl, and 0.5%

Triton X-100 with protease inhibitors. A total of 13 µg of cell lysates was used for anti-GFP immunoblotting followed by electrochemiluminescence-based signal detection and quantification. We performed triplicate experiments for statistical analysis.

Statistical analysis

5 Statistical differences between groups were assessed using the Wilcoxon rank sum test.

Example 2: micro RNA mediated inhibition of transgene expression reduces dorsal root ganglia toxicity by AAV

10 Delivering adeno-associated virus (AAV) vectors into the CNS of non-human primates (NHP) via the blood or cerebral spinal fluid is associated with dorsal root ganglia (DRG) toxicity. This may be caused by high rates of transduction, which can cause endoplasmic reticulum stress from overproduction of the transgene product. We developed an approach to eliminate toxicity associated with CNS-directed AAV gene therapy by 15 introducing miRNA target sequences into the vector genome within the 3' untranslated region of the corresponding transgene mRNA. The expression cassette for ITR.CB7.CI.eGFP.miR145(four copies).rabbit beta globin, 3'ITR is provided in SEQ ID NO: 10, the expression cassette for ITR.CB7.CI.GFP.miR182(four copies).rabbit beta globin, 3'ITR is provided in SEQ ID NO: 11, the expression cassette for ITR.CB7.CI.GFP.miRNA96(four copies).rabbit beta globin, 3'ITR is provided in SEQ ID NO: 12, and the expression cassette for ITR.CB7.CI.GFP.miR183(four copies).rabbit beta globin, 3'ITR is provided in SEQ ID NO: 13.

20

AAV vectors cause DRG degeneration in NHPs

25 Based on our experience in DRG toxicity in NHPs, we developed a system to quantify the severity of toxicity. We evaluate cell bodies located along the spinal cord in the DRGs, the axons within the peripheral nerves, and the axons that ascend the dorsal white-matter tracts (FIG. 1B). We believe the primary lesion is degeneration of the sensory neuron cell body located in DRG. The lesion is histologically characterized by hypereosinophilia, irregular cell shapes, disruption of Nissl substance (central chromatolysis), and loss of 30 nuclear boundaries along with mononuclear cell infiltration (FIG. 1B). Cells expressing high levels of transgene protein are more likely to undergo degeneration as evidenced by transgene product immunostaining in animals that received an ICM administration of an AAV vector expressing green fluorescent protein (GFP; FIG. 1B). Secondary to the cell

body death is axonopathy, which is degeneration of the distal and proximal axons.

Axonopathy is characterized by missing axons, dilated myelin sheaths surrounding cell debris, and macrophages (FIG. 1B). FIG. 1C illustrates examples of different levels of DRG toxicity and spinal cord axonopathy. The grades are based on the proportion of affected tissue at high-power field histopathologic examination: 1 minimal (<10%), 2 mild (10-25%), 3 moderate (25-50%), 4 marked (50-95%) and 5 severe (>95%).

Our total experience of adolescent/adult NHPs administered AAV vectors into the CSF via ICM or lumbar puncture (LP) route totals 101 monkeys spanning 21 studies that encompasses previous published toxicology studies (Hordeaux, J., et al. Mol Ther Methods Clin Dev 10:68-78, 2018; Hordeaux, J., et al. Mol Ther Methods Clin Dev 10:79-88, 2018) and the two NHP experiments described in the Examples below as well as in a number of unpublished studies. This experience includes seven capsids, 12 transgenes, three promoters (CB7, UBC, hSyn), doses from 1×10^{12} GC to 5.7×10^{13} GC, vector purified by gradients or columns, three formulations (phosphate buffered saline and two different artificial CSF), and rhesus and cynomolgus macaques at various developmental stages. In every experimental group, we observed DRG toxicity and axonopathy. The pathology peaks about one month after injection and does not progress for up to six months, which is the longest period evaluated in mature macaques. In most cases, the pathology is mild to moderate. However, high doses of vectors expressing GFP injected ICM can lead to severe pathology associated with ataxia.

miRNAs specifically expressed in DRG neurons can ablate AAV transgene expression

Several mechanisms were evaluated when considering ways to mitigate DRG toxicity. In previous studies, we analyzed the role of destructive adaptive immune responses to the transduced DRGs by immune suppressing NHPs that were administered ICM AAV9 vectors expressing human IDUA or human IDS. Treatment with mycophenolate mofetil (MMF) and rapamycin blunted the adaptive immune response to the vector and transgene product but did not significantly impact DRG toxicity and axonopathy (Hordeaux, J., et al. Mol Ther Methods Clin Dev 10:68-78, 2018; Hordeaux, J., et al. Mol Ther Methods Clin Dev 10:79-88, 2018).

One possibility that has not been previously investigated is whether overexpression of transgene products in highly transduced DRGs is the cause neuronal injury and degeneration of the cell body and associated axons followed by a reactive inflammatory

response (FIG. 2A). Accordingly, to specifically ablate transgene expression in DRG we cloned miRNA targets that are solely expressed in DRG neurons into the 3' untranslated regions of the transgene (FIG. 2B). Any mRNA expressed from the vector would be destroyed by the endogenously expressed miRNA.

5 We used an *in vitro* assay to evaluate the activity and specificity of the miRNA strategy. We constructed AAV *cis* plasmids to include four repeat concatemers of the target miRNA sequences in the 3' untranslated region of the expression cassette (FIG. 2B). AAV *cis* plasmids were co-transfected into HEK293 cells with plasmids expressing miR183. Expression of the transgene GFP was reduced in the presence of miR183 only when it 10 contained the cognate recognition sequence (FIG. 3A).

The *in vivo* activity and specificity of potential miRNA targets within AAV vectors was screened in C57Bl/6J mice. We evaluated GFP-expressing vectors with or without miRNA targets from two members of the miRNA183 complex (miR182 and miR183) as well as miR145. We initially tested miR96, another member of the 183 complex, but 15 eliminated it due to decreased transgene expression in mice cortices (not shown). Animals received high-dose intravenous (IV) injections of AAV9 to target DRGs and high-dose AAV-PHP.B (AAV9-PHP.B.CB7.CI.GFP.rBG) injections to target the CNS. Animals were necropsied on day 21 and analyzed for GFP expression in DRGs by immunohistochemistry (IHC) and direct-fluorescence microscopy in brain and liver. Expression of GFP in DRG 20 neurons was substantially reduced with vectors containing miR183 and miR182 targets, however miR145 targets had no effect (FIG. 3B and FIG. 3C). There was no apparent reduction of expression in liver or other CNS compartments with vectors containing any of the miR targets. Expression seemed to be slightly enhanced in CNS with the miR183 vector (FIG. 3D). In this mouse experiment, we were unable to assess the impact of miR183 25 transgene suppression on pathology since the vector-induced DRG toxicity has only been observed in NHPs.

Restricted transgene expression by miR183 reduces DRG toxicity in NHPs

Based on the encouraging data in mice, we evaluated the GFP miR183 expression cassette in NHPs. We ICM injected AA Vhu68 vectors expressing GFP 30 (AAV9.CB7.CI.GFP.rBG) (N=2) or GFP miR183 (AAV9.CB7.CI.GFP.miR183.rBG) (N=4) from a CB7 promoter in rhesus macaques (3.5×10^{13} GC). Half of the animals were necropsied on day 14 for GFP expression (FIG. 4B - representative IHC for GFP expression; FIG. 4B - quantitation of expression). The remaining animals were necropsied on day 60 to

evaluate expression and DRG toxicity (FIG. 4C - DRG degeneration, dorsal spinal axonopathy, and peripheral nerve axonopathy). Animals tolerated the ICM-administered vector without clinical sequelae. There was a statistically significant reduction of GFP expression in DRG with the miR183 vector and enhancement or similar expression 5 elsewhere including lumbar motor neurons, cerebellum, cortex, heart, and liver (FIG. 4A and FIG. 4B). This was associated with a remarkable reduction of pathology across nine regions (DRG and dorsal spinal axonopathy at cervical, thoracic and lumbar spine and axonopathy of median, peroneal and radial nerves; FIG. 4C). Without miR183 targets in the vector, pathology was present in all regions and evenly distributed between grade 4, grade 2, and 10 grade 1. With the miR183 vector, the greatest pathology was grade 2 and was present in only 11% of regions; the remaining regions were either grade 1 (72%) or no pathology (16%).

These studies demonstrated that GFP expression is selectively repressed in DRG sensory neurons with vectors that contain miR183 targets. Other CNS neurons and peripheral organs were not affected. Accordingly, DRG toxicity and secondary axonopathy were 15 reduced from marked/severe to minimal levels in the context of a highly immunogenic/toxic transgene (GFP).

Example 3: Specific repression of therapeutic protein expression in dorsal root ganglia following delivery via AAV with a vector genome having miRNA target 20 sequences

We further evaluated miR183 target sequences in NHPs using vectors that expressed human IDUA—an enzyme deficient in patients with mucopolysaccharidosis I. Studies with this human transgene were the first to highlight DRG toxicity in NHPs (Hordeaux, J., et al. Mol Ther Methods Clin Dev 10:79-88, 2018). The experiment included three groups 25 (N=3/group): 1) group 1 – control vector alone without miR183 targets (AAVhu68.CB7.CI.hIDUAcoV1.rBG); 2) group 2 – control vector without miR183 targets (AAVhu68.CB7.CI.hIDUAcoV1.rBG) in animals treated with steroids (prednisolone 1 mg/kg/day from day minus 7 to day 30 followed by progressive taper off); and 3) group 3 - vector with miR183 targets (AAVhu68.CB7.CI.hIDUAcoV1.miR183.rBG). All vector 30 genomes included an hIDUA coding sequence under the control of a chicken β-actin promoter and CMV enhancer elements (referred to as the CB7 promoter), a chimeric intron (CI) consisting of a chicken β-actin splice donor (973 bp, GenBank: X00182.1) and a rabbit β-globin splice acceptor element, and a rabbit β-globin polyadenylation signal (rBG, 127 bp,

GenBank: V00882.1). The vector genome for ITR.CB7.CI.hIDUAcoV1.rBG.ITR is provided in SEQ ID NO: 14. The vector genome for ITR.CB7.CI.hIDUAcoV1.miRNA183.rBG.ITR is provided in SEQ ID NO: 15. All animals received an ICM injection of an AAVhu68 vector (1×10^{13} GC) expressing hIDUA from the 5 constitutive promoter CB7. Necropsies were conducted at day 90 to evaluate transgene expression and DRG-related toxicity.

Animals from all groups tolerated ICM vector with no vector-related clinical findings or abnormalities in clinical pathology (Table 1 and Table 2). Pleocytosis in CSF was very low and limited to one animal in group 2 and one animal in group 3 (Table 3). Both T-cell 10 responses (measured by ELISPOT) and antibodies to hIDUA were detected in all three groups (FIG. 7A – FIG. 7D). CSF cytokines were reduced in group 3 compared to group 1 at 21 and 35 days post-injection while levels were reduced in group 2 (steroids) at 24 hours (FIG. 8). Day 21-35 corresponds to peak expression of transgene when overexpression induced stress would be expected.

15 Using direct fluorescence and *in situ* hybridization (ISH), we observed high expression of hIDUA in DRG in groups 1 and 2 that used the control vector (without miR183 target; FIG. 5 and FIG. 6A). We detected moderate levels of hIDUA expression in other CNS compartments including lower motor neurons of the spinal cord and cerebellum and cortical neurons (FIG. 5 and FIG. 6A). Including the miR183 target into the vector 20 (group 3) ablated hIDUA expression in DRG neurons without decreasing expression in the CNS (FIG. 5 and FIG. 6A). Reduction of hIDUA expression in DRGs by miR183 was not due to decreased gene transfer since the biodistribution of vector throughout the CNS and DRGs was essentially the same across all groups (FIG. 9). Steroids moderately decreased expression in DRG and increased it in lower motor neurons compared with group 1 (FIG. 5 and FIG. 6A). As expected, group 1 exhibited pathology in DRGs, dorsal column and 25 peripheral nerve. However, these findings were completely absent when using vector with miR183 targets (group 3, FIG. 6B). Interestingly, co-treatment with steroids (group 2) did not reduce toxicity of the parent vector (i.e., not containing miR183). In fact, we noticed a trend of worsening toxicity (FIG. 6B).

30

Table 1. Blood chemistry in NHP injected ICM with AAV.hIDUA vectors

Animal # and group	Timepoint	Total Protein			A/G Ratio	AST	ALT	Alk Phosphatase	GGT	Total Bilirubin	BUN	Creatinine
		g/ dL	g/ dL	g/ dL								
17C024 AAVhu6 8.hIDUA	Baseline	5.6	3.8	1.8	2.1	33	40	703	64	0.1	27	0.6
	D0	6.0	3.7	2.3	1.6	28	36	682	61	0.1	23	0.4
	D7	6.1	3.9	2.2	1.8	30	33	680	64	0.1	30	0.5
	D21	5.9	3.7	2.2	1.7	25	22	825	66	0.1	21	0.5
	D35	5.9	3.4	2.5	1.4	32	24	777	74	0.1	22	0.5
	D60	5.9	3.7	2.2	1.7	28	30	815	74	0.1	28	0.4
	D90	5.5	3.6	1.9	1.9	29	25	689	76	0.1	24	0.6
17C031 AAVhu6 8.hIDUA	Baseline	5.7	3.8	1.9	2.0	35	39	539	36	0.1	27	0.5
	D0	6.5	4.0	2.5	1.6	37	43	570	47	0.2	23	0.4
	D7	6.3	3.9	2.4	1.6	27	39	547	45	0.1	26	0.5
	D21	6.6	3.9	2.7	1.4	33	34	476	39	0.1	25	0.6
	D35	6.3	3.4	2.9	1.2	32	31	591	42	0.1	24	0.5
	D60	6.2	4.0	2.2	1.8	32	33	545	45	0.1	25	0.4
	D90	6.0	3.6	2.4	1.5	34	27	478	43	0.1	24	0.5
17C029 AAVhu6 8.hIDUA	Baseline	6.8	3.6	3.2	1.1	41	43	574	74	0.1	27	0.7
	D0	6.9	4.0	2.9	1.4	27	40	607	79	0.1	18	0.7
	D7	7.1	3.8	3.3	1.2	29	40	503	71	0.1	24	0.6
	D21	6.9	3.9	3.0	1.3	49	38	557	77	0.1	15	0.6
	D35	6.6	3.6	3.0	1.2	30	33	555	76	0.1	19	0.6
	D60	6.3	4.0	2.3	1.7	38	43	545	86	0.1	22	0.6
	D90	6.2	3.7	2.5	1.5	31	23	506	81	0.1	20	0.6
17C016 AAVhu6 8.hIDUA +	Baseline	6.0	3.9	2.1	1.9	42	35	623	59	0.2	16	0.5
	D0	6.8	4.1	2.7	1.5	33	58	588	59	0.2	16	0.4
	D7	6.3	3.9	2.4	1.6	26	28	444	51	0.1	20	0.5
	D21	6.4	3.7	2.7	1.4	29	23	393	44	0.2	16	0.6

Animal # and group	Timepoint	Total Protein		Globulin	A/G Ratio	AST	ALT	Alk Phosphatase	GGT	Total Bilirubin	BUN	Creatinine
		g/dL	g/dL			-	/L	L				
steroids	D35	6.9	3.9	3.0	1.3	32	29	364	41	0.1	21	0.5
	D60	6.5	3.8	2.7	1.4	41	29	525	63	0.2	23	0.5
	D90	5.7	3.6	2.1	1.7	32	29	752	79	0.1	12	0.5
17C019 AAVhu6 8.hIDUA + Steroids	Baseline	5.5	3.4	2.1	1.6	35	42	338	49	0.1	31	0.6
	D0	6.3	3.7	2.6	1.4	32	39	383	49	0.1	19	0.6
	D7	6.1	3.6	2.5	1.4	26	34	322	47	0.1	30	0.7
	D21	6.0	3.3	2.7	1.2	28	32	332	53	0.1	19	0.7
	D35	5.9	3.5	2.4	1.5	42	38	378	49	0.1	27	0.6
	D60	5.6	3.6	2.0	1.8	35	36	437	61	0.2	23	0.5
	D90	5.6	3.6	2.0	1.8	41	34	650	73	0.2	21	0.6
17C020 AAVhu6 8.hIDUA + steroids	Baseline	6.1	4.1	2.0	2.1	26	33	641	87	0.1	21	0.7
	D0	6.5	4.1	2.4	1.7	21	25	538	61	0.1	12	0.6
	D7	6.9	4.0	2.9	1.4	21	28	463	58	0.1	21	0.7
	D21	6.6	4.0	2.6	1.5	23	29	456	55	0.1	13	0.5
	D35	6.2	3.7	2.5	1.5	22	26	309	54	0.1	20	0.6
	D60	6.2	4.1	2.1	2.0	26	27	516	90	0.1	19	0.5
	D90	6.1	3.8	2.3	1.7	23	22	605	82	0.1	16	0.6
17-167 AAVhu6 8.hIDUA- miR183	Baseline	6.6	3.9	2.7	1.4	24	21	764	112	0.1	18	0.5
	D0	6.7	4.1	2.6	1.6	27	21	606	88	0.1	20	0.4
	D7	6.6	3.9	2.7	1.4	31	27	606	87	0.1	24	0.6
	D21	6.8	4.2	2.6	1.6	29	27	584	102	0.1	22	0.5
	D35	6.6	4.0	2.6	1.5	35	34	642	93	0.1	16	0.5
	D60	6.0	3.7	2.3	1.6	37	25	689	79	0.1	15	0.6
	D90	6.6	3.9	2.7	1.4	24	21	764	112	0.1	18	0.5
17-215	Baseline	6.1	3.4	2.7	1.3	24	34	536	51	0.1	22	0.5
	D0	6.5	3.6	2.9	1.2	29	50	568	54	0.2	21	0.5

Animal # and group	Timepoint	Total Protein		Globulin	A/G Ratio	AST	ALT	Alk Phosphatase	GGT	Total Bilirubin	BUN	Creatinine
		g/dL	g/dL			-	/L	L	IU/L	mg/dL	mg/L	mg/dL
AAVhu6 8.hIDUA- miR183	D7	6.3	3.7	2.6	1.4	24	35	499	46	0.1	27	0.6
	D21	6.2	3.5	2.7	1.3	31	37	526	58	0.1	21	0.5
	D35	6.0	3.4	2.6	1.3	33	47	619	57	0.1	18	0.4
	D60	6.1	3.7	2.4	1.5	40	50	628	45	0.2	18	0.5
	D90	6.3	3.9	2.4	1.6	27	42	856	55	0.1	17	0.6
17-102	Baseline	6.7	3.8	2.9	1.3	24	31	613	57	0.1	20	6.7
AAVhu6 8.hIDUA- miR183	D0	7.0	3.6	3.4	1.1	32	35	632	57	0.1	13	7.0
	D7	6.9	3.9	3.0	1.3	31	25	601	57	0.1	16	6.9
	D21	7.0	3.8	3.2	1.2	31	30	611	65	0.1	20	7.0
	D35	6.8	3.8	3.0	1.3	28	29	684	59	0.1	20	6.8
	D60	6.4	3.8	2.6	1.5	34	36	588	44	0.1	20	6.4
	D90	7.0	4.0	3.0	1.3	27	28	576	41	0.1	18	7.0

Table 2. Complete blood count in NHP injected ICM with AAV.hIDUA vectors

Animal # and group	Timepoint	WBC	RBC	HGB	HCT	Platelet	Neutrophils	Lymphocytes	Monocytes	Eosinophils	Basophils
		x10 ³ / µL	x10 ⁶ / µL	g/dL	%	x10 ³ / µL	/µL	/µL	/µL	/µL	/µL
17C024 AAVhu68. hIDUA	Baseline	8.6	6.1	13.1	43	305	2838	5246	344	172	0
	D0	6.1	5.5	12.7	40	304	3294	2562	183	61	0
	D7	5.3	5.1	11.2	38	343	1060	3869	265	106	0
	D21	6.7	5.7	12.8	42	289	1742	4422	335	201	0
	D35	5.6	5.6	12.7	42	377	1792	3584	168	56	0
	D60	6.6	5.8	13.0	42	356	1716	4488	198	198	0

Animal # and group	Timepoint	WBC	RBC	HGB	HCT	Platelet	Neutrophils	Lymphocytes	Monocytes	Eosinophils	Basophils
		x10 ³ / µL	x10 ⁶ / µL	g/dL	%	x10 ³ / µL	/µL	/µL	/µL	/µL	/µL
		D90	5.4	5.7	13.1	41	399	1674	3456	162	108
17C031 AAVhu68. hIDUA	Baseline	11.9	5.3	12.6	41	386	4879	6188	476	357	0
	D0	5.5	5.2	12.5	41	366	2255	2970	165	110	0
	D7	14.4	4.8	11.2	39	357	8064	5040	864	432	0
	D21	10.7	5.4	12.8	43	410	5136	4815	428	321	0
	D35	9.7	5.3	12.8	43	267	4559	4559	388	194	0
	D60	8.3	5.3	12.7	42	390	2988	4731	332	249	0
	D90	8.4	4.9	12.1	39	414	4284	2940	672	420	84
17C029 AAVhu68. hIDUA	Baseline	15.5	5.9	12.7	43	561	4495	9610	775	620	0
	D0	15.1	5.7	12.6	41	493	9362	4681	755	302	0
	D7	11.2	5.8	12.7	43	576	2800	7280	784	336	0
	D21	10.6	5.8	12.6	42	496	4982	4770	530	318	0
	D35	12.0	5.8	13.1	44	511	2520	8280	600	600	0
	D60	11.6	5.8	13.6	43	497	3480	7192	696	232	0
	D90	19.6	5.7	12.9	42	283	14896	3528	980	196	0
17C016 AAVhu68. hIDUA + steroids	Baseline	10.2	5.9	13.5	44	235	1734	7752	306	408	0
	D0	8.9	5.6	12.9	44	353	3382	5073	356	89	0
	D7	9.6	5.5	12.3	41	346	2976	5952	576	96	0
	D21	11.9	5.4	12.4	41	424	2856	8330	595	119	0
	D35	7.9	5.4	12.7	43	352	4977	2528	316	79	0
	D60	8.7	5.6	12.8	43	380	2610	5655	348	87	0
	D90	8.4	5.4	12.7	41	357	2940	4956	420	84	0
17C019	Baseline	10.7	5.5	13.4	40	257	6099	4066	214	321	0
	D0	10.6	6.4	13.9	49	358	7208	2650	318	424	0

Animal # and group	Timepoint	WBC	RBC	HGB	HCT	Platelet	Neutrophils	Lymphocytes	Monocytes	Eosinophils	Basophils
		x10 ³ / µL	x10 ⁶ / µL	g/dL	%	x10 ³ / µL	/µL	/µL	/µL	/µL	/µL
AAVhu68. hIDUA + Steroids	D7	10.8	5.9	13.1	44	286	3564	6048	0	108	0
	D21	12.3	5.7	12.9	41	462	5658	5904	492	246	0
	D35	12.4	5.9	13.4	45	297	7812	3720	620	248	0
	D60	8.6	5.8	13.4	43	373	3612	4386	430	172	0
	D90	11.1	5.5	13.0	41	349	8214	2442	333	111	0
17C020 AAVhu68. hIDUA + steroids	Baseline	14.6	6.5	14.1	47	357	4818	8760	584	438	0
	D0	15.4	6.1	13.7	44	339	8778	5390	616	616	0
	D7	14.0	6.0	13.4	43	378	4060	8680	840	420	0
	D21	14.4	5.7	12.9	42	327	6336	7056	720	144	144
	D35	11.1	6.2	13.9	45	369	2664	7770	444	222	0
	D60	10.8	6.2	14.0	45	380	4104	5940	324	432	0
	D90	7.2	5.9	13.5	44	296	2376	4536	216	72	0
17-167 AAVhu68. hIDUA- miR183	Baseline	10.6	5.4	13.7	44	245	1802	8162	530	106	0
	D0	7.2	5.1	12.2	40	373	936	5904	288	72	0
	D7	7.6	4.9	12.2	39	258	1140	6080	304	76	0
	D21	10.2	5.5	13.4	45	270	3468	5712	714	102	204
	D35	10.6	5.3	13.2	42	275	5406	4664	424	106	0
	D60	9.5	5.2	13.4	42	115	3515	5510	380	95	0
	D90	10.6	5.4	13.7	44	245	1802	8162	530	106	0
17-215 AAVhu68. hIDUA- miR183	Baseline	12.9	5.8	12.7	44	380	2709	9546	387	258	0
	D0	10.0	5.5	11.9	40	374	3600	5900	300	200	0
	D7	11.2	5.4	12.0	40	423	2800	7616	560	224	0
	D21	11.8	5.2	11.3	39	375	2242	9086	236	118	118
	D35	12.5	5.6	12.4	42	337	5125	6625	500	250	0

Animal # and group	Timepoint	WBC	RBC	HGB	HCT	Platelet	Neutrophils	Lymphocytes	Monocytes	Eosinophils	Basophils
		x10 ³ / µL	x10 ⁶ / µL	g/dL	%	x10 ³ / µL	/µL	/µL	/µL	/µL	/µL
	D60	10.3	5.4	12.1	39	315	3811	5974	309	206	0
	D90	13.7	5.3	12.1	38	415	8494	4795	274	137	0
17-102 AAVhu68. hIDUA- miR183	Baseline	5.3	6.0	14.0	46	518	530	3816	371	530	53
	D0	12.4	5.9	13.5	44	478	6944	4588	496	372	0
	D7	9.2	5.8	13.1	44	501	3312	4968	368	552	0
	D21	9.0	5.9	14.0	45	510	2250	5940	540	270	0
	D35	9.4	6.4	14.5	49	409	3760	4888	564	188	0
	D60	9.1	6.2	14.6	46	410	3276	5005	455	364	0
	D90	7.8	5.9	14.4	44	528	3042	4212	312	234	0

Table 3. CSF white blood cell counts (cells per µL) in NHP injected ICM with AAV.hIDUA vectors

Group	Animal #	Day 0	Day 21	Day 35	Day 60	Day 90
AAVhu68. hIDUA	17C024	0	1	2	1	2
	17C031	0	1	2	1	1
	17C029	0	0	0	0	0
AAVhu68. hIDUA + steroids	17C016	0	0	0	0	3
	17C019	0	1	2	1	0
	17C020	0	0	5	1	Blood contamination
AAVhu68. hIDUA- miR183	17-167	0	2	3	1	1
	17-215	0	1	2	1	1
	17-102	0	3	7	2	0

Toxicity of DRGs is likely to occur in any therapy that relies on high systemic doses of vector or direct delivery of vector into the CSF. This safety concern is limited to primates

and has usually been asymptomatic. However, DRG toxicity can cause substantial morbidity such as ataxia due to proprioceptive defects (Hinderer, C., et al. Hum Gene Ther. 29(3):285-298, 2018) or intractable neuropathic pain. The U.S. Food & Drug Administration recently paused an intrathecal AAV9 clinical trial for late-onset SMA due to NHP DRG toxicity,
5 which underscores how this risk may limit the development of AAV therapies (Novartis. Novartis announces AVXS-101 intrathecal study update, 2019).

It was originally hypothesized that this toxicity was caused by destructive T-cell immunity to transduced neurons in DRGs directed towards foreign capsid or transgene epitopes. However, strong immune suppression regimens such as MMF and rapamycin did
10 not prevent the toxicity in toxicology studies (Hordeaux, J., et al. Mol Ther Methods Clin Dev 10:68-78, 2018; Hordeaux, J., et al. Mol Ther Methods Clin Dev 10:79-88, 2018), nor did steroids in this study. The time course of delayed but not progressive DRG degeneration did not support the notion that adaptive immunity played a role. If cytotoxic T cells were involved, DRG degeneration and mononuclear cell infiltrates that began early and progressed
15 over time would have been observed.

It may be that high levels of DRG transduction create cellular stress, which leads to degeneration in the highly transduced DRG neurons. Since toxicity can be prevented by suppressing transgene mRNA and protein expression, capsid or vector DNA cannot be the cellular stressors. Histological analysis demonstrated that degeneration was limited to DRG
20 neurons that expressed the highest level of transgene protein. The time course of delayed but self-limited DRG neuronal degeneration is consistent with the notion that non-immune toxicity is restricted to a subset of highly transduced cells. It is unclear whether the DRG toxicity and axonopathy are reversible. After following adult animals for six months, consistent reductions in pathology have not been observed. The only experiment where DRG
25 toxicity was observed in NHPs following ICM injection was when the vector was administered to one-month old macaques that were necropsied four years later (Hordeaux et al., 2019). It is possible that infant primates are resistant to DRG toxicity, or their DRG neurons have regenerative capacity, or the lesions regressed over this extended time period.

The findings presented support that DRG toxicity is caused by transgene
30 overexpression. Therefore, the severity of DRG toxicity should be influenced by dose, promoter strength, and the nature of the transgene. It is still not understood why sensory neurons are one of the most efficiently transduced cells in primates. DRGs are easily accessed by systemically administered vectors because they reside outside of the CNS and

have porous, fenestrated capillaries. Systemic vector could also access DRG neurons via retrograde transport after uptake from peripheral axons. The anatomy of sensory neuronal compartments that reside within the intrathecal space may promote high transduction of vectors delivered into the CSF. Axons of DRG neurons in the dorsal roots are exposed to

5 CSF providing easy access to vector following ICM/LP administration. Open access of the subarachnoid space to the extracellular fluid of the DRG should allow direct contact of ICM/LP vector to the neuronal cell bodies and other cells of the DRG. Suppression of transgene expression within DRG neurons with miR183 facilitated an analysis of transgene expression in other cells of the DRGs which should not be influenced by this miR. ISH

10 revealed transgene mRNA in surrounding glial satellite cells that could suggest direct transduction (FIG. 6C). The functional consequence of transgene mRNA in glial cells is unknown.

Selectively inhibiting vector transgene expression should reduce and potentially eliminate DRG toxicity. The key for achieving this is a strategy for specifically

15 extinguishing expression in DRG neurons without affecting expression elsewhere. There are currently no ways to achieve this specificity through capsid modifications or tissue-specific promoters. Including targets for miR183 into the vector achieved the desired result of reducing/eliminating DRG toxicity without affecting vector manufacturing, potency, or biodistribution. Included in the hIDUA NHP study above was a group that received non-

20 miR183 vector with concomitant steroids - a standard approach for mitigating immune-mediated toxicity in AAV trials. DRG toxicity was not reduced in the steroid-treated group; in fact, there was a trend toward worsening toxicity. This experiment demonstrates the limitations of prophylactic steroids in AAV gene-therapy trials.

The modularity of this approach for diminishing DRG toxicity suggests its use in any

25 AAV vector considered for CNS gene therapy where mitigating AAV-induced DRG toxicity is desirable. This approach can be used across a broad array of AAV vectors for therapeutic applications.

Example 4: *In vitro* assessment of expression constructs with miR183 cluster target sequences

An *in vitro* assay is used to evaluate the activity and specificity of constructs harboring miRNA target sequences. As described in Example 2 above, HEK293 cells (or another suitable cell line) are co-transfected with a cis plasmid having the GFP transgene and

plasmids expressing one or more miRNA, such as miR-182 and miR-183. The cis plasmids are designed with varying number of corresponding target miRNA sequences in the 3'UTR of the expression cassettes and alternative spacer sequences are introduced. At 72 hours post transfection, expression of GFP is quantified to determine relative levels of expression.

5 For example, constructs harboring one, two, three, or four copies of target miR183 sequences are tested. The individual target sequences are directly linked or separated by spacer sequences, such as those provided in SEQ ID NOs: 5-7. Based on results of *in vitro* study, the suitable combination of sequences (including number of repeats) and spacers that reduce or eliminate expression of GFP are identified. Candidates from this study are then
10 screened *in vivo* by delivering AAV vectors (e.g. AAV9 or AAV-PHP.B) having expression constructs with the same or similar arrangement of target miRNA sequences and spacers sequences. An exemplary *in vivo* mouse study to evaluate CNS expression levels, including, for example, detargeting of DRG (i.e. reduction of GFP expression), is provided in Example 2.

15 Similar studies are also performed using constructs having combinations of one, two, three, or four copies of target sequences for miR182 with and without various spacer sequences. Additionally, constructs having combinations and different arrangements of miR182 and miR183 recognition sequences are generated. The constructs having miR182 target sequences only and combinations of miR182 and miR183 target sequences that show
20 favorable reduced levels of expression *in vitro* are then evaluated *in vivo*, for example, following administration of AAV vectors to determine toxicity and levels of transgene expression (extent of detargeting) in cells of the CNS and DRG.

25 Alternatively, constructs are generated having one, two, three, or four copies of a combination of miR182 target sequence and/or other mir183 cluster target sequences (i.e. a target sequences corresponding to miR-183, -96, or -182). The combination miR182-mir183 cluster target sequence-harboring constructs are tested *in vitro* using a GFP expression assay such as that described in Example 2 above. As above, the tested expression cassettes have various number of miRNA target sequences that are or are not separated by spacer sequences. The activity of certain constructs having combinations of miR182 target
30 sequences and other mir183 cluster target sequences is then evaluated *in vivo* by generating AAV vectors that are then administered at high-dose IV. As above, expression of the AAV vector transgene is evaluated in various cells and tissues, including DRG and, in particular, in liver tissue.

Further, the effect of one, two, three, or four copies of miR182 target sequences of transgene expression is evaluated. As above, experimental constructs for *in vitro* testing are generated introducing miR182 target sequences into the 3'UTR of an expression cassette. Where multiple miR182 sequences are introduced, the sequences may be consecutive or, 5 alternatively, separated by any of various intervening spacer sequences. AAV vectors are generated having expression cassettes with any combination of miR182 target sequences and, where applicable, spacer sequences, and tested *in vivo*. In particular, in the case of expression cassettes having miR182 target sequences, transgene expression is evaluated in muscle tissue following high-dose IV administration of the AAV vector.

10

Example 5: Detargeting of a human iduronate-2-sulfatase (hIDS) transgene for treatment of Mucopolysaccharidosis Type II (MPS II)

One strategy for the treatment of MPS II (Hunter syndrome) is to functionally replace a patient's defective iduronate-2-sulfatase via rAAV-based CNS-directed gene 15 therapy (see, e.g., International Patent Application No. PCT/US2017/027770, which is incorporated by reference herein). To reduce DRG toxicity, AAV vector genomes for treatment of MPSII are modified by introducing miR target sequences. Accordingly, AAV vector genomes containing a hIDS coding sequence are designed with one, two, three, or four miR183 target sequences. The effectiveness of DRG detargeting *in vivo* is measured, 20 for example, following intrathecal administration of the AVV vector encoding hIDS to NHPs.

Example 6: Detargeting of a SMN1 transgene for treatment of spinal muscular atrophy (SMA)

25 SMA is an autosomal recessive disorder caused by mutations or deletion of the hSMN1 gene. Delivery of functional SMN protein via rAAV vectors has been effective for treatment of SMA but DRG toxicity has been observed. Suitable vectors include those described in International Patent Application No. PCT/US2018/019996, which is incorporated by reference herein, and Zolgensma®, an AAV9-based gene therapy). 30 Reduction or elimination of DRG toxicity following delivery of AAV vectors encoding human SMN1 is achieved by incorporating miRNA target sequences, such as those recognized by miR182 and miR183, into the vector genome. Accordingly, AAV vectors, including those with AAV9 or AA Vhu68 capsids, are generated having a nucleic acid

sequence encoding a hSMN1 transcript in combination with one, two, three, or four miRNA target sequences. The target sequences are selected, for example, from miR182 and miR183 target sequences, or a combination thereof. DRG toxicity following IV or intrathecal administration of a hSMN1-expressing AAV vectors is evaluated in a NHP model.

5

Example 7: Liver-directed gene therapy vectors having miRNA target sequences

Where improved expression of a transgene in liver tissue is desirable for gene therapy, AAV vector genomes can be modified to include miRNA target sequences. For example, a rAAV designed to express a functional low-density lipoprotein receptor (hLDLR) gene and bearing an AAV8 capsid is suitable for treatment of familial hypercholesterolemia (FH) (see, e.g., International Patent Application No. PCT/US2016/065984, which is incorporated herein by reference). Enhanced expression of the hLDLR transgene in liver tissue is achieved using an rAAV with a vector genome having a hLDLR coding sequence in combination with one, two, three, or four miR182 target sequences. Likewise, gene therapies for treatment of hemophilia A (Factor VIII) and hemophilia B (Factor IX) include vectors with tropism for the liver (see, e.g., International Patent Application No. PCT/US2017/027396 and International Patent Application No. PCT/US2017/027400, which are incorporated herein by reference). More effective delivery and expression of human factor VIII and factor IX in liver is achieved by delivering rAAVs with vectors genomes having one, two, three, or four miR182 target sequences in combination with the transgene.

(Sequence Listing Free Text)

The following information is provided for sequences containing free text under numeric identifier <223>.

SEQ ID NO: (containing free text)	Free text under <223>
1	<223> miR-183 target
2	<223> mirR-96 target
3	<223> miR-182 target
4	<223> miR-145 target

SEQ ID NO: (containing free text)	Free text under <223>
5	<223> Spacer (i)
6	<223> Spacer (ii)
7	<223> spacer iii
10	<223> ITR.CB7.CI.eGFP.miR145.rBG.ITR <220> <221> repeat_region <222> (1)..(130) <223> 5' - AAV2 - ITR <220> <221> misc_feature <222> (1)..(130) <223> 5' - AAV2 - ITR <220> <221> promoter <222> (198)..(579) <223> CMV IE promoter <220> <221> promoter <222> (582)..(863) <223> CB promoter <220> <221> misc_feature <222> (1979)..(2698) <223> eGFP gene <220> <221> misc_feature <222> (2705)..(2727) <223> miR145 <220> <221> misc_feature <222> (2728)..(2731) <223> spacer <220> <221> misc_feature <222> (2732)..(2754)

SEQ ID NO: (containing free text)	Free text under <223>
	<223> miR145 <220> <221> misc_feature <222> (2755)..(2760) <223> spacer <220> <221> misc_feature <222> (2761)..(2783) <223> miR145 <220> <221> misc_feature <222> (2784)..(2789) <223> spacer <220> <221> misc_feature <222> (2790)..(2812) <223> miR145 <220> <221> misc_feature <222> (2981)..(3198) <223> 3' ITR
11	<223> ITR.CB7.CI.eGFP.miR182.rGB.ITR <220> <221> misc_feature <222> (1)..(130) <223> 5' ITR (AAV2) <220> <221> misc_feature <222> (198)..(579) <223> CMV IE promoter <220> <221> misc_feature <222> (582)..(863) <223> CB promoter <220>

SEQ ID NO: (containing free text)	Free text under <223>
	<221> misc_feature <222> (958)..(1930) <223> chicken beta-actin intron <220> <221> misc_feature <222> (1979)..(2698) <223> eGFP coding sequence <220> <221> misc_feature <222> (2705)..(2728) <223> miR182 <220> <221> misc_feature <222> (2729)..(2732) <223> spacer <220> <221> misc_feature <222> (2733)..(2756) <223> miR182 <220> <221> misc_feature <222> (2757)..(2760) <223> spacer <220> <221> misc_feature <222> (2763)..(2786) <223> miR182 <220> <221> misc_feature <222> (2787)..(2792) <223> spacer <220> <221> misc_feature <222> (2793)..(2816) <223> miR182

SEQ ID NO: (containing free text)	Free text under <223>
	<220> <221> polyA_signal <222> (2858)..(2984) <220> <221> misc_feature <222> (3073)..(3202) <223> 3' ITR
12	<223> ITR.CB7.eGFP.miRNA96.rBG.ITR <220> <221> misc_feature <222> (1979)..(2699) <223> eGFP coding sequence <220> <221> misc_feature <222> (2705)..(2727) <223> miR96 <220> <221> misc_feature <222> (2728)..(2731) <223> spacer <220> <221> misc_feature <222> (2732)..(2754) <223> miR96 <220> <221> misc_feature <222> (2755)..(2760) <223> spacer <220> <221> misc_feature <222> (2761)..(2783) <223> miR96 <220> <221> misc_feature <222> (2784)..(2789) <223> spacer

SEQ ID NO: (containing free text)	Free text under <223>
	<220> <221> misc_feature <222> (2790)..(2812) <223> miR96
13	<223> ITR.CB7.CI.eGFP.miRNA183.rBG.ITR <220> <221> misc_feature <222> (1979)..(2698) <223> eGFP coding sequence <220> <221> misc_feature <222> (2705)..(2726) <223> miRNA183 <220> <221> misc_feature <222> (2727)..(2730) <223> spacer <220> <221> misc_feature <222> (2731)..(2752) <223> miRNA183 <220> <221> misc_feature <222> (2753)..(2758) <223> spacer <220> <221> misc_feature <222> (2781)..(2786) <223> spacer <220> <221> misc_feature <222> (2787)..(2808) <223> miRNA183
14	<223> ITR.CB7.CI.hIDUAcoV1.rBG.ITR

SEQ ID NO: (containing free text)	Free text under <223>
15	<223> ITR.CB7.CI.hIDUAcoV1.miR183.ITR <220> <221> misc_feature <222> (1938)..(3908) <223> hIDUAcoV1 <220> <221> misc_feature <222> (3915)..(3936) <223> miRNA183 <220> <221> misc_feature <222> (3937)..(3940) <223> spacer <220> <221> misc_feature <222> (3941)..(3962) <223> miRNA183 <220> <221> misc_feature <222> (3963)..(3968) <223> spacer <220> <221> misc_feature <222> (3969)..(3990) <223> miRNA183 <220> <221> misc_feature <222> (3991)..(3996) <223> spacer <220> <221> misc_feature <222> (3997)..(4018) <223> miRNA183

All publications cited in this specification are incorporated herein by reference in their entireties. US Provisional Patent Application No. 62/783,956, filed December 21, 2018, US Provisional Patent Application No. 62/924,970, filed October 23, 2019, and US Provisional Patent Application No. 62/934,915, filed November 13, 2019 are hereby incorporated by reference in their entireties. Similarly, the SEQ ID NOs which are referenced herein and which appear in the appended Sequence Listing are incorporated by reference. While the invention has been described with reference to particular embodiments, it will be appreciated that modifications can be made without departing from the spirit of the invention. Such modifications are intended to fall within the scope of the appended claims.

CLAIMS:

1. A composition for gene delivery which specifically represses expression of a gene product in dorsal root ganglion (DRG) comprising an expression cassette which is a nucleic acid sequence comprising:
 - (a) a coding sequence for a gene product under the control of regulatory sequences which direct expression of the gene product in a cell containing the expression cassette; and
 - (b) at least one target sequence specific for at least one of miR-183, miR-182, or miR-96, the at least one target sequence being operably linked at the 3' end of the coding sequence (a).
2. The composition according to claim 1, wherein the composition comprises at least two tandem repeats of the targeting sequences which comprise at least a first miRNA target sequence and at least a second miRNA target sequence which may be the same or different.
3. The composition according to claim 2, wherein the at least two miRNA tandem repeats are located in 3' UTR .
4. The composition according to any one of claims 1 to 3, wherein the expression cassette comprises a 3' UTR having three miRNA tandem repeats.
5. The composition according to any one of claims 2 to 4, wherein the tandem miRNA target sequences are continuous or are separated by a spacer of 1 to 10 nucleic acids, wherein said spacer is not a miRNA target sequence.
6. The composition according to any one of claims 1 to 5, wherein there are two sets of at least two miRNA target sequences located in the 3' UTR.
7. The composition according to any one of claims 2 to 6, wherein the start of the first of the at least two miRNA tandem repeats is within 20 nucleotides from the 3' end of the gene coding sequence.

8. The composition according to any one of claims 2 to 6, wherein the start of the first of the at least two miRNA tandem repeats is at least 100 nucleotides from the 3' end of the gene coding sequence.

9. The composition according to any one of claims 2 to 6, wherein the 3' UTR and the miRNA tandem repeats comprise 200 to 1200 nucleotides in length.

10. The composition according to any one of claims 1 to 9, wherein the expression cassette comprises four miRNA target sequences located in the 3' UTR.

11. The composition according to any one of claims 1 to 8, wherein the expression cassette further comprises at least one target sequence specific for miR-183, miR-182, or miR-96 in the 5' UTR.

12. The composition according to claim 11, wherein the expression cassette comprises at least two miRNA target sequences located in both the 5' UTR and the 3' UTR.

13. The composition according to any one of claims 1 to 12, wherein the at least one miRNA target sequence for the expression cassette mRNA or DNA positive strand is

- (a) AGTGAATTCTACCAGTGCCATA (SEQ ID NO: 1);
- (b) AGCAAAAATGTGCTAGTGCCAAA (SEQ ID NO: 2);
- (c) AGTGTGAGTTCTACCATTGCCAAA (SEQ ID NO: 3); or
- (d) AGGGATTCTGGAAAATGGAC (SEQ ID NO: 4).

14. The composition according to any one of claims 2 to 13, wherein two or more consecutive miRNA target sequences are continuous and not separated by a spacer.

15. The composition according to any one of claims 1 to 13, wherein two or more of the miRNA target sequences are separated by a spacer and each spacer is independently selected from one or more of (i) GGAT (SEQ ID NO: 5); (ii) CACGTG (SEQ ID NO: 6); or (iii) GCATGC (SEQ ID NO: 7).

16. The composition according to any one of claims 1 to 15, wherein the spacer located between the miRNA target sequences is located 3' to the first miRNA target sequence and/or 5' to the last miRNA target sequence.
17. The composition according to claim 15 or 16, wherein the spacers between the miRNA target sequences are the same.
18. The composition according to any one of claims 1 to 17, wherein the expression cassette is carried by a viral vector selected from a recombinant parvovirus, a recombinant lentivirus, a recombinant retrovirus, and a recombinant adenovirus.
19. The composition according to any one of claims 1 to 17, wherein the expression cassette is carried by a non-viral vector selected from naked DNA, naked RNA, an inorganic particle, a lipid particle, a polymer-based vector, and a chitosan-based formulation.
20. A recombinant AAV (rAAV) for delivery of a gene product to a patient in need thereof which specifically represses expression of a gene product in dorsal root ganglia, said rAAV comprising a viral capsid having packaged therein an AAV vector genome, wherein the vector genome comprises:
 - (a) a coding sequence for the gene product under the control of regulatory sequences which direct expression of the gene product in a cell containing the vector genome; and
 - (b) at least one miRNA target sequence specific for at least one of miR-183, miR-182, or miR-96.
21. A pharmaceutical composition comprising the rAAV according to claim 20 and a formulation buffer suitable for delivery via intracerebroventricular (ICV), intrathecal (IT), intracisternal or intravenous (IV) injection.
22. A method for repressing transgene expression in DRG neurons, said method comprising delivering a composition according to any one of claims 1 to 19, an rAAV according to claim 20, or a composition according to claim 21 to a patient.

23. A method for modulating neuronal degeneration and/or decreasing secondary dorsal spinal cord axonal degeneration following intrathecal or systemic gene therapy administration, said method comprising delivering a composition according to any one of claims 1 to 19, an rAAV according to claim 20, or a composition according to claim 21 to a patient.

24. A method for enhancing expression of a transgene in cells of the central nervous system (CNS), said method comprising delivering a composition according to any one of claims 1 to 19, an rAAV according to claim 20, or a composition according to claim 21 to a patient, wherein the miRNA target sequences comprise at least one miR183 target sequence.

25. The method of claim 24, wherein the cells of the CNS include one or more of pyramidal neurons, purkinje neurons, granule cells, spindle neurons, interneuron cells, astrocytes, oligodendrocytes, microglia, and ependymal cells.

FIG. 1A




FIG. 1B




FIG. 1C




FIG. 2A




FIG. 2B

AAV-transgene-miR target




FIG. 3A




FIG. 3B

DRG transduction




FIG. 3C




FIG. 3D




FIG. 4A




FIG. 4B




FIG. 4C




FIG. 5




FIG. 6A




FIG. 6B





FIG. 6C



AAVhu68.hIDUA-miR183

FIG. 7A




FIG. 7B





FIG. 7C




FIG. 7D

FIG. 8




FIG. 9

■ Group 1 AAVhu68.hNDUAcov1 (G1)
 ■ Group 2 AAVhu68.hNDUAcov1 steroids (G2)
 ■ Group 3 AAVhu68.hNDUAcov1.miR183 (G3)

