



(86) Date de dépôt PCT/PCT Filing Date: 2012/01/20
(87) Date publication PCT/PCT Publication Date: 2012/07/26
(85) Entrée phase nationale/National Entry: 2013/04/05
(86) N° demande PCT/PCT Application No.: US 2012/022053
(87) N° publication PCT/PCT Publication No.: 2012/100179
(30) Priorité/Priority: 2011/01/21 (US61/434,849)

(51) Cl.Int./Int.Cl. *C07D 307/22* (2006.01),
C07D 498/04 (2006.01), *C07D 513/04* (2006.01)


(71) Demandeur/Applicant:
EISAI R & D MANAGEMENT CO., LTD., JP

(72) Inventeurs/Inventors:
MITASEV, BRANKO, US;
KIM, DAE-SHIK, US;
ZHANG, HUIMING, US;
SCHNADERBECK, MATTHEW J., US;
FARTHING, CHRISTOPHER N., GB;
YOSHIZAWA, KAZUHIRO, JP


(74) Agent: SMART & BIGGAR

(54) Titre : PROCEDES ET COMPOSES UTILES DANS LA SYNTHESE DE DERIVES FUSIONNES
D'AMINODIHYDROTHIAZINE


(54) Title: METHODS AND COMPOUNDS USEFUL IN THE SYNTHESIS OF FUSED AMINODIHYDROTHIAZINE
DERIVATIVES




I




II




III




IV



V



VI



VII

(57) Abrégé/Abstract:

Provided are compounds and methods useful for the preparation of compounds useful as inhibitors of beta-site amyloid precursor protein (APP)-cleaving enzyme.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau(10) International Publication Number
WO 2012/100179 A1(43) International Publication Date
26 July 2012 (26.07.2012)

(51) International Patent Classification:

C07D 307/22 (2006.01) C07D 513/04 (2006.01)
C07D 498/04 (2006.01)

(21) International Application Number:

PCT/US2012/022053

(22) International Filing Date:

20 January 2012 (20.01.2012)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/434,849 21 January 2011 (21.01.2011) US

(71) Applicant (for all designated States except US): **EISAI R&D MANAGEMENT CO., LTD.** [JP/JP]; 6-10, Koishikawa 4-Chome, Bunkyo-ku, Tokyo 112-8088 (JP).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **MITASEV, Branko** [IT/US]; 3 Vine Street, North Reading, Massachusetts 01864 (US). **KIM, Dae-Shik** [KR/US]; 311 Lowell Street, Apt # 3113, Andover, Massachusetts 01810 (US). **ZHANG, Huiming** [CN/US]; 2 Dorset Circle, Andover, Massachusetts 01810 (US). **SCHNADERBECK, Matthew J.** [US/US]; 26 Morgan Drive, Methuen, Massachusetts 01844 (US). **FARTHING, Christopher N.** [GB/GB]; 29 Portland Rise, London Greater London N4 2PT (GB). **YOSHIZAWA, Kazuhiro** [JP/JP]; 2-11-4-101 Miyashita, Kashima, Ibaraki 314-0032 (JP).(74) Agent: **MURPHY, Sherry L.**; Myers Bigel Sibley & Sajovec, P.A., P.O. Box 37428, Raleigh, North Carolina 27627 (US).


(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: METHODS AND COMPOUNDS USEFUL IN THE SYNTHESIS OF FUSED AMINODIHYDROTHIAZINE DERIVATIVES



(57) Abstract: Provided are compounds and methods useful for the preparation of compounds useful as inhibitors of beta-site amyloid precursor protein (APP)-cleaving enzyme.

METHODS AND COMPOUNDS USEFUL IN THE SYNTHESIS OF FUSED AMINODIHYDROTHIAZINE DERIVATIVES

RELATED APPLICATIONS

This application claims the benefit under 35 U.S.C. § 119(e) of U.S. Provisional Patent Application No. 61/434,849, filed January 21, 2011.

FIELD OF THE INVENTION

The present invention relates to compounds and methods that are useful for the preparation of compounds useful as inhibitors of beta-site amyloid precursor protein (APP)-cleaving enzyme.

BACKGROUND OF THE INVENTION

Alzheimer's disease is a disease characterized by degeneration and loss of neurons as well as formation of senile plaques and neurofibrillary degeneration. Currently, Alzheimer's disease is treated only with symptomatic treatment using a symptom improving agent typified by an acetylcholinesterase inhibitor, and a fundamental remedy to inhibit progression of the disease has not yet been developed. It is necessary to develop a method for controlling the cause of the onset of pathology in order to create a fundamental remedy for Alzheimer's disease.

It is thought that A β -proteins as break-down products of amyloid precursor proteins (APP) are involved in the degeneration and loss of neurons and onset of symptoms of dementia (see, e.g., Klein et al., Alzheimer's disease-affected brain: Presence of oligomeric A β ligands (ADDLs) suggests a molecular basis for reversible memory loss, PNAS USA 2003, Sep 2; 100 (18), p. 10417-10422; Nitsch et al., Antibodies against β -amyloid slow cognitive decline in Alzheimer's disease, Neuron, 2003, May 22; 38, p. 547-554).

A β -proteins have, as main components, A β 40 consisting of 40 amino acids and A β 42 with two amino acids added at the C-terminal. The A β 40 and A β 42 are known to have high aggregability and to be main components of senile plaques (see, e.g., Jarrett et al., The carboxy terminus of the β amyloid protein is critical for the seeding of amyloid formation: Implications for the pathogenesis of Alzheimers' disease, Biochemistry, 1993, 32 (18), p. 4693-4697; Glenner et al., Alzheimer's disease: initial report of the purification and characterization of a novel cerebrovascular amyloid protein, Biochemical and biophysical research communications, 1984, May 16, 120 (3), p. 885-890; Masters et al., Amyloid plaque core protein in Alzheimer disease and Down syndrome, PNAS USA, 1985, Jun, 82 (12), p. 4245-4249). Further, it is known that


the A β 40 and A β 42 are increased by mutation in APP and presenilin genes which is observed in familial Alzheimer's disease (see, e.g., Gouras et al., Intraneuronal A β 42 accumulation in human brain, American Journal of Pathology, 2000, Jan, 156 (1), p. 15-20; Scheuner et al., Secreted amyloid β -protein similar to that in the senile plaques of Alzheimer's disease is increased in vivo by the presenilin 1 and 2 and APP mutations linked to familial Alzheimer's disease, Nature Medicine, 1996, Aug, 2 (8), p. 864-870; Forman et al., Differential effects of the swedish mutant amyloid precursor protein on β -amyloid accumulation and secretion in neurons and nonneuronal cells, The Journal of Biological Chemistry, 1997, Dec 19, 272 (51), p. 32247-32253). Accordingly, a compound that reduces production of A β 40 and A β 42 is expected to be a progression inhibitor or prophylactic agent for Alzheimer's disease.

A β is produced by cleaving APP by beta-secretase (BACE1) and subsequently by gamma-secretase. For this reason, attempts have been made to create gamma-secretase and beta-secretase inhibitors in order to inhibit A β production.

SUMMARY

The present invention relates to compounds and methods that are useful for the preparation of inhibitors of beta-site amyloid precursor protein (APP)-cleaving enzyme, also known as beta-secretase, or BACE1. BACE1 inhibitor compounds and methods of making the same have been described, for example, in WO 2009/091016, the contents of which is hereby incorporated by reference herein in its entirety.

Provided herein are compounds of **Formula I**:



I

wherein:

R₁ is alkyl, haloalkyl, hydroxyalkyl, or -(C₁₋₆alkyl)-OR', wherein R' is an oxygen protecting group; and

R₂ and R₃ are each independently hydrogen or halo (e.g., fluoro);

or a salt thereof.


In some embodiments, **Formula I** is subject to the proviso that when one of R₂ or R₃ is *ortho*-fluoro, and the other is hydrogen, R₁ is not -CH₂F.

In some embodiments, R_1 is trifluoromethyl, hydroxymethyl, monofluoromethyl, or triphenylmethoxymethyl.

In some embodiments, R_2 is fluoro, and/or R_3 is hydrogen or fluoro.

Also provided are methods of making a compound of **Formula I** as defined above. In some embodiments, the method includes the step of heating a mixture including:

(i) an oxime of **Formula A**:



A

wherein R_1 , R_2 and R_3 are as given above for **Formula I**;

(ii) hydroquinone; and


(iii) a hydrocarbon solvent (e.g., benzene, toluene, xylenes, etc.),

to a temperature of from 90, 95, 100, 105, or 110 °C, to 115, 120, 125, 130, 135, 140, 145, 150, 155, or 160 °C (for example, from 90°C to 160°C, or from 110°C to 140°C, or from 100°C to 120°C, or from 120°C to 140°C),

to make said compound of **Formula I**, or a salt thereof.

In some embodiments, the method includes the step of heating a mixture comprising:

(i) a compound of **Formula B**:



B

wherein R_1 , R_2 and R_3 are as given above;

(ii) an acetate;


(iii) hydroxylamine or a salt thereof; and

(iv) a hydrocarbon solvent (e.g., benzene, toluene, xylenes, etc.),

to a temperature of from 90, 95, 100, 105, or 110 °C, to 115, 120, 125, 130, 135, 140, 145, 150, 155, or 160 °C (for example, from 90°C to 160°C, or from 110°C to 140°C, or from 100°C to 120°C, or from 120°C to 140°C),

to make said compound of **Formula I**, or a salt thereof.

Further provided are compounds of **Formula II**:



II

wherein:

R_1 is alkyl, haloalkyl, hydroxyalkyl, or $-(C_{1-6})-OR'$, wherein R' is an oxygen protecting group; and

R_2 and R_3 are each independently hydrogen or halo (e.g., fluoro);


or a salt thereof.

In some embodiments, **Formula II** is subject to the proviso that when one of R_2 or R_3 is *ortho*-fluoro, and the other is hydrogen, R_1 is not $-CH_2F$.

In some embodiments, R_1 is trifluoromethyl, hydroxymethyl, monofluoromethyl, or triphenylmethoxymethyl.

In some embodiments, R_2 is fluoro, and/or R_3 is hydrogen or fluoro.

Also provided are methods of making a compound of **Formula II**:



II

wherein:

R_1 is C_{1-4} alkyl; and


R_2 and R_3 are each independently hydrogen or halo;

or a salt thereof,

comprising the steps of:

(a) heating a mixture comprising:

(i) a ketone of **Formula B**:



B

5 wherein R₁, R₂ and R₃ are as given above for **Formula II**;

(ii) hydroxylamine or a salt thereof;

(iii) a base (e.g., acetate, propionate, a tertiary amine base such as triethylamine or diethylisopropylamine, pyridine, or a salt thereof); and

(iv) a solvent selected from: water and an alcohol (e.g., a C₃₋₈ alcohol),

10 to at least 90 or 100° Celsius (e.g., to boiling such as by refluxing) produce a heated mixture,

(b) cooling the heated mixture to a temperature of at or below 60, 50 or 45° Celsius, and then,


(c) adding an organic solvent (e.g., tetrahydrofuran or an ether solvent such as diethyl ether or methyl *tert*-butyl ether), an acid (e.g., a carboxylic acid such as acetic acid, or a mineral acid such as hydrochloric acid or sulfuric acid), and zinc (e.g., zinc powder or zinc dust) to the mixture, wherein said mixture is maintained at or below 60, 50 or 45 degrees Celsius (e.g., between 25, 30 or 35, and 45, 50 or 60 degrees Celsius) during said adding,

to make said compound of **Formula II**.

20

In some embodiments, the heating is carried out for at least 30 hours.

Also provided is a method of making a stereochemically pure compound of **Formula IIa**:



IIa


25

wherein R₁, R₂ and R₃ are as given above for **Formula II**,

or a salt thereof,

including the steps of :

(a) adding a mixture of stereoisomers of the compound of **Formula II**:



II


wherein R_1 , R_2 and R_3 are as given above,

5 to a solvent (e.g., an alcohol such as ethanol or isopropanol, or a mixture such as a toluene – acetonitrile mixture, a toluene – acetone mixture, a toluene – tetrahydrofuran mixture, or an alcohol – water mixture), which solvent includes a chiral carboxylic acid compound (e.g., D-dibenzoyl tartaric acid, L-dibenzoyl tartaric acid, D-ditoluyl tartaric acid, and L-ditoluyl tartaric acid, etc.) to form a mixture of diastereomeric salts of the compound of **Formula II**; and
 10 then (b) crystallizing a single diastereomeric salt formed of the compound of **Formula II**, to make the stereochemically pure compound of **Formula IIa**, or a salt thereof.

In some embodiments, the stereochemically pure compound has greater than about 80% by weight of one stereoisomer of the compound and less than about 20% by weight of other stereoisomers of the compound, more preferably greater than about 90% by weight of one stereoisomer of the compound and less than about 10% by weight of the other stereoisomers of the compound, even more preferably greater than about 95% by weight of one stereoisomer of the compound and less than about 5% by weight of the other stereoisomers of the compound, and most preferably greater than about 97% by weight of one stereoisomer of the compound and less than about 3% by weight of the other stereoisomers of the compound greater than about 90% by
 20 weight of one stereoisomer of the compound and less than about 10% by weight of the other stereoisomers of the compound.

In some embodiments, the crystallizing step is performed by cooling the mixture of diastereomeric salts of the compound of **Formula II** (e.g., to a temperature of from 40, 45, 50, 55, or 60 °C to 80, 90, 100, or 110 °C, to a temperature of from -10, -5, 0, 5, or 10 °C to 15, 20, 29,
 25 or 35 °C).

Further provided are compounds of **Formula III**:



III

wherein:

5 R_1 is alkyl, haloalkyl, hydroxyalkyl, or $-(C_{1-6})-OR'$, wherein R' is an oxygen protecting group; and


R_2 and R_3 are each independently hydrogen or halo (e.g., fluoro);
or a salt thereof.

In some embodiments, **Formula III** is subject to the proviso that when one of R_2 or R_3 is *ortho*-fluoro, and the other is hydrogen, R_1 is not $-CH_2F$.

10 In some embodiments, R_1 is trifluoromethyl, hydroxymethyl, monofluoromethyl, or triphenylmethoxymethyl.

In some embodiments, R_2 is fluoro, and/or R_3 is hydrogen or fluoro.

Also provided are compounds of **Formula IV**:



IV

wherein:

15 R_1 is alkyl, haloalkyl, hydroxyalkyl, or $-(C_{1-6})-OR'$, wherein R' is an oxygen protecting group; and


20 R_2 and R_3 are each independently hydrogen or halo (e.g., fluoro);
or a salt thereof.

In some embodiments, **Formula IV** is subject to the proviso that when one of R_2 or R_3 is *ortho*-fluoro, and the other is hydrogen, R_1 is not $-CH_2F$.

25 In some embodiments, R_1 is trifluoromethyl, hydroxymethyl, monofluoromethyl, or triphenylmethoxymethyl.

In some embodiments, R_2 is fluoro, and/or R_3 is hydrogen or fluoro.

Further provided are compounds of **Formula V**:



5 wherein:

R_1 is alkyl, haloalkyl, hydroxyalkyl, or $-(C_{1-6})-OR'$, wherein R' is an oxygen protecting group; and

R_2 and R_3 are each independently hydrogen or halo (e.g., fluoro);
or a salt thereof.


10 In some embodiments, **Formula V** is subject to the proviso that when one of R_2 or R_3 is *ortho*-fluoro, and the other is hydrogen, R_1 is not $-CH_2F$.

In some embodiments, R_1 is trifluoromethyl, hydroxymethyl, monofluoromethyl, or triphenylmethoxymethyl.

In some embodiments, R_2 is fluoro, and/or R_3 is hydrogen or fluoro.

15

Also provided are methods for making a compound of **Formula V**:



20 wherein:

R_1 is C_{3-4} alkyl, halo- C_{1-4} alkyl, hydroxy- C_{1-4} alkyl, or $-C_{1-4}$ alkyl- OR' ,
wherein R' is an oxygen protecting group; and


R_2 and R_3 are each independently hydrogen or halo;
or a salt thereof,

25

comprising the steps of:

(a) providing a mixture of

(i) a compound of **Formula III**,



III

5 wherein R₁, R₂ and R₃ are as given above for **Formula V**;

(ii) an amine base (e.g., pyridine, or a substituted pyridine such as collidine); and

(iii) an organic solvent (e.g., an aromatic hydrocarbon solvent, toluene, xylenes, benzene, dichloromethane, etc.),

10 wherein said mixture is at a temperature at or below 0° Celsius,

(b) reacting the mixture with trifluoromethanesulfonic anhydride, 4-toluenesulfonyl chloride, methanesulfonyl chloride or methanesulfonyl anhydride, and then,

(c) adding:


(i) an alcohol (e.g., propanol, such as 2-propanol); and

15 (ii) a base, such as a strong base selected from: a hydroxide, an alkoxide, and ammonia,

to make said compound of **Formula V**.

20 In some embodiments, the reacting step is carried out at a temperature of from -15, -10, or -5, to 0° Celsius.

Also provided are compound of **Formula VI**:



VI

25 wherein:

WO 2012/100179

PCT/US2012/022053

R_1 is alkyl, haloalkyl, hydroxyalkyl, or $-(C_{1-6})-OR'$, wherein R' is an oxygen protecting group; and

R_2 and R_3 are each independently hydrogen or halo (e.g., fluoro);
or a salt thereof.


5 In some embodiments, **Formula VI** is subject to the proviso that when one of R_2 or R_3 is *ortho*-fluoro, and the other is hydrogen, R_1 is not $-CH_2F$.

In some embodiments, R_1 is trifluoromethyl, hydroxymethyl, monofluoromethyl, or triphenylmethoxymethyl.

In some embodiments, R_2 is fluoro, and/or R_3 is hydrogen or fluoro.

10

Further provided are compounds of **Formula VII**:



VII

wherein:

15 R_1 is alkyl, haloalkyl, hydroxyalkyl, or $-(C_{1-6})-OR'$, wherein R' is an oxygen protecting group; and


R_2 and R_3 are each independently hydrogen or halo (e.g., fluoro);
or a salt thereof.

In some embodiments, R_1 is methyl, trifluoromethyl, hydroxymethyl, monofluoromethyl,
20 or triphenylmethoxymethyl.

In some embodiments, R_2 is fluoro, and/or R_3 is hydrogen or fluoro.

Also provided are methods of making a compound of **Formula VII**, or a salt thereof,
including the step of:

25 reducing a compound of **Formula VI** to form a compound of **Formula VII**, as shown
below:



wherein:

R_1 is alkyl, haloalkyl, hydroxyalkyl, or $-(C_{1-6})-OR'$, wherein R' is an oxygen protecting group; and

R_2 and R_3 are each independently hydrogen or halo.

In some embodiments, R_1 is trifluoromethyl, hydroxymethyl, monofluoromethyl, or triphenylmethoxymethyl.


In some embodiments, R_2 is fluoro, and/or R_3 is hydrogen or fluoro.

In some embodiments, the reducing of the compound of **Formula VI** to form the compound of **Formula VII** may include one or more of the steps of: (a) adding powdered iron to a carbinol solvent to form a mixture thereof; (b) adding hydrochloric acid (HCl) (e.g., concentrated HCl) to the mixture of step (a); (c) heating the mixture of step (b) to a temperature of from 31, 40 or 50 °C to 65, 70, 75, 80, 85°C, or to the boiling point of the mixture of step (b) (e.g., for 0 - 4 hours) to form a heated mixture; and (d) adding the compound of **Formula VI**, or a salt thereof, to the heated mixture of step (c).

In some embodiments, the method further includes the steps of: (e) cooling the mixture of step (d) to a temperature of between 10 and 30°C to form a cooled mixture; and (f) optionally, stirring the cooled mixture (e.g., for 0 - 4 hours).

In some embodiments, the HCl is provided in an amount selected from the group consisting of: an amount of HCl stoichiometric with respect to the amount of the compound of **Formula VI**; and an amount of HCl in molar excess with respect to the amount of the compound of **Formula VI**.

Further provided are methods of making a compound of **Formula IX**:



wherein:

R_1 is alkyl, haloalkyl, hydroxyalkyl, or $-(C_{1-6})-OR'$, wherein R' is an oxygen protecting group; and


5 R_2 and R_3 are each independently hydrogen or halogen; and

R_4 is selected from the group consisting of alkyl, haloalkyl, alkoxy, and alkyl substituted with alkoxy;

or a salt thereof;

including the step of reacting a compound **Formula VII** with a pyrazine carboxylic acid of

10 **Formula VIII**:



wherein R_1 , R_2 , R_3 and R_4 are as given above,

to form said compound of **Formula IX**, or a salt thereof.

15 In some embodiments, the reacting may include one or more of the steps of: (a) adding thionyl chloride to a solution comprising a compound of **Formula VIII** in a solvent to make a formed mixture; and (b) adding a compound of **Formula VII** to the formed mixture of step (a), to form said compound of **Formula IX**, or a salt thereof.

In some embodiments, the solution comprising the compound of **Formula VIII** is a solvent is cooled prior to or during the adding step (a) (e.g., to a temperature of from 0 to 25 °C).

20 In some embodiments, the solvent may include N,N'-dimethylimidazoline-2-one, toluene, methylene chloride, dimethylformamide, N-methylpyrrolidine, or dimethylacetamide.

In some embodiments, the reacting may include one or more of the steps of: (a) adding ethyl acetate (EtOAc) to a compound of **Formula VIII** and a compound of **Formula VII** to make a


formed mixture; and (b) adding an alkyl phosphonic acid anhydride (e.g., 2-propyl phosphonic acid anhydride) to the formed mixture of step (a), to form said compound of Formula IX, or a salt thereof.

In some embodiments, the formed mixture may be maintained at a temperature of from 0°C to 35°C during said adding step (a).

BRIEF DESCRIPTION OF THE DRAWINGS

Figures 1A-1B present typical chromatograms from chiral HPLC upon diastereomeric resolution of a tetrahydrofuran of **Formula II** in accordance with Example 4, part C.

Figure 2 presents the x-ray crystal structure of ((2S,3R,4S)-4-Amino-4-(2-fluorophenyl)-2-((trityloxy)methyl)tetrahydrofuran-3-yl)methanol (2R,3R)-2,3-bis(benzoyloxy)succinate:



DETAILED DESCRIPTION

A. Definitions

Compounds of this invention include those described generally above, and are further illustrated by the embodiments, sub-embodiments, and species disclosed herein. As used herein, the following definitions shall apply unless otherwise indicated.

As described herein, compounds of the invention may optionally be substituted with one or more substituents, such as are illustrated generally above, or as exemplified by particular classes, subclasses, and species of the invention. In general, the term "substituted" refers to the replacement of hydrogen in a given structure with a specified substituent. Unless otherwise indicated, a substituted group may have a substituent at each substitutable position of the group, and when more than one position in any given structure may be substituted with more than one substituent selected from a specified group, the substituent may be either the same or different at every position. Combinations of substituents envisioned by this invention are preferably those that result in the formation of stable or chemically feasible compounds.

"Isomers" refer to compounds having the same number and kind of atoms and hence the same molecular weight, but differing with respect to the arrangement or configuration of the atoms.

"Stereoisomers" refer to isomers that differ only in the arrangement of the atoms in space.

"Diastereoisomers" refer to stereoisomers that are not mirror images of each other.

"Enantiomers" refers to stereoisomers that are non-superimposable mirror images of one another.

5 Enantiomers include "enantiomerically pure" isomers that comprise substantially a single enantiomer, for example, greater than or equal to 90%, 92%, 95%, 98%, or 99%, or equal to 100% of a single enantiomer.

"Enantiomerically pure" as used herein means a compound, or composition of a compound, that comprises substantially a single enantiomer, for example, greater than or equal to
10 90%, 92%, 95%, 98%, or 99%, or equal to 100% of a single enantiomer.

"Stereomerically pure" as used herein means a compound or composition thereof that comprises one stereoisomer of a compound and is substantially free of other stereoisomers of that compound. For example, a stereomerically pure composition of a compound having one chiral center will be substantially free of the opposite enantiomer of the compound. A
15 stereomerically pure composition of a compound having two chiral centers will be substantially free of diastereomers, and substantially free of the enantiomer, of the compound. A typical stereomerically pure compound comprises greater than about 80% by weight of one stereoisomer of the compound and less than about 20% by weight of other stereoisomers of the compound, more preferably greater than about 90% by weight of one stereoisomer of the compound and less
20 than about 10% by weight of the other stereoisomers of the compound, even more preferably greater than about 95% by weight of one stereoisomer of the compound and less than about 5% by weight of the other stereoisomers of the compound, and most preferably greater than about 97% by weight of one stereoisomer of the compound and less than about 3% by weight of the other stereoisomers of the compound. *See, e.g.,* US Patent No. 7,189,715.

25 "R" and "S" as terms describing isomers are descriptors of the stereochemical configuration at an asymmetrically substituted carbon atom. The designation of an asymmetrically substituted carbon atom as "R" or "S" is done by application of the Cahn-Ingold-Prelog priority rules, as are well known to those skilled in the art, and described in the International Union of Pure and Applied Chemistry (IUPAC) Rules for the Nomenclature of
30 Organic Chemistry. Section E, Stereochemistry.

"Enantiomeric excess" (ee) of an enantiomer is [(the mole fraction of the major enantiomer) minus (the mole fraction of the minor enantiomer)] x 100.

"Stable", as used herein, refers to compounds that are not substantially altered when subjected to conditions to allow for their production, detection, and preferably their recovery,
35 purification, and use for one or more of the purposes disclosed herein. In some embodiments, a

stable compound or chemically feasible compound is one that is not substantially altered when kept at a temperature of 40°C or less, in the absence of moisture or other chemically reactive conditions, for at least a week.

5 "Refluxing" as used herein refers to a technique in which vapors from a boiling liquid are condensed and returned to the mixture from which it came, typically by boiling the liquid in a vessel attached to a condenser.

"Powdered iron" or "iron powder" is iron having an average particle size of less than 0.1, 0.5, 1, 5, 10, 20, 50, 250, 500 or 1000 μm . Particle size can be measured using methods known in the art, e.g., mesh sizing, laser diffraction, etc.

10 "Zinc dust" is zinc having an average particle size of less than 0.001, 0.05, 0.1, 0.5, 1, 5, 10, 15 or 20 μm . "Zinc powder" is zinc having an average particle size of less than 200, 175, 150, 125, or 100 μm . Particle size can be measured using methods known in the art, e.g., mesh sizing, laser diffraction, etc.

15 An "organic" compound as used herein is a compound that contains carbon. Similarly, an "organic solvent" is a compound containing carbon that is useful as a solvent. An "inorganic" compound is a compound not containing carbon.

"Mineral acid" as used herein is the acid of an inorganic compound. Examples include, but are not limited to, hydrochloric acid (HCl), nitric acid (HNO₃), phosphoric acid (H₃PO₄), sulfuric acid (H₂SO₄), boric acid (B(OH)₃), hydrofluoric acid (HF), hydrobromic acid (HBr),
20 perchloric acid (HClO₄), etc.

A "hydrocarbon" is an organic compound consisting of carbon and hydrogen atoms. Examples of hydrocarbons useful as "hydrocarbon solvents" include, but are not limited to, an "aromatic hydrocarbon solvent" such as benzene, toluene, xylenes, etc., and an "aliphatic hydrocarbon solvent" such as pentane, hexane, heptane, etc.

25 An "amine" or "amine base" as used herein refers to an organic compound having a basic nitrogen atom (R-NR'R''), and may be a primary (R-NH₂), secondary (R-NHR') or tertiary (R-NR'R'') amine.

A "strong base" as used herein is a compound that is capable of deprotonating very weak acids. Examples of strong bases include, but are not limited to, hydroxides, alkoxides, and
30 ammonia.

A "hydroxide" is the commonly known diatomic anion OH⁻, or a salt thereof (typically an alkali metal or alkaline earth metal salt thereof). Examples of hydroxides include, but are not limited to, sodium hydroxide (NaOH), potassium hydroxide (KOH), lithium hydroxide (LiOH), and calcium hydroxide (CaOH).

An "alkoxide" is RO^- , the conjugate base of an alcohol. Examples include, but are not limited to, methoxide, ethoxide, and propoxide.

"Ar" or "aryl" refer to an aromatic carbocyclic moiety having one or more closed rings. Examples include, without limitation, phenyl, naphthyl, anthracenyl, phenanthracenyl, biphenyl, and pyrenyl.

"Heteroaryl" refers to a cyclic moiety having one or more closed rings, with one or more heteroatoms (for example, oxygen, nitrogen or sulfur) in at least one of the rings, wherein at least one of the rings is aromatic, and wherein the ring or rings may independently be fused, and/or bridged. Examples include without limitation quinolinyl, isoquinolinyl, indolyl, furyl, thienyl, pyrazolyl, quinoxalinyl, pyrrolyl, indazolyl, thieno[2,3-c]pyrazolyl, benzofuryl, pyrazolo[1,5-a]pyridyl, thiophenylpyrazolyl, benzothienyl, benzothiazolyl, thiazolyl, 2-phenylthiazolyl, and isoxazolyl.

"Alkyl" or "alkyl group," as used herein, means a straight-chain (i.e., unbranched), branched, or cyclic hydrocarbon chain that is completely saturated. In some embodiments, alkyl groups contain 1, 2, or 3, to 4, 5 or 6 carbon atoms (e.g., C_{1-4} , C_{2-4} , C_{3-4} , C_{1-5} , C_{2-5} , C_{3-5} , C_{1-6} , C_{2-6} , C_{3-6}). In some embodiments, alkyl groups contain 3-4 carbon atoms. In certain embodiments, alkyl groups contain 1-3 carbon atoms. In still other embodiments, alkyl groups contain 2-3 carbon atoms, and in yet other embodiments alkyl groups contain 1-2 carbon atoms. In certain embodiments, the term "alkyl" or "alkyl group" refers to a cycloalkyl group, also known as carbocycle. Non-limiting examples of exemplary alkyl groups include methyl, ethyl, propyl, isopropyl, butyl, cyclopropyl and cyclohexyl.

"Alkenyl" or "alkenyl group," as used herein, refers to a straight-chain (i.e., unbranched), branched, or cyclic hydrocarbon chain that has one or more double bonds. In certain embodiments, alkenyl groups contain 2-6 carbon atoms. In certain embodiments, alkenyl groups contain 2-4 carbon atoms. In still other embodiments, alkenyl groups contain 3-4 carbon atoms, and in yet other embodiments alkenyl groups contain 2-3 carbon atoms. According to another aspect, the term alkenyl refers to a straight chain hydrocarbon having two double bonds, also referred to as "diene." In other embodiments, the term "alkenyl" or "alkenyl group" refers to a cycloalkenyl group. Non-limiting examples of exemplary alkenyl groups include $-\text{CH}=\text{CH}_2$, $-\text{CH}_2\text{CH}=\text{CH}_2$ (also referred to as allyl), $-\text{CH}=\text{CHCH}_3$, $-\text{CH}_2\text{CH}_2\text{CH}=\text{CH}_2$, $-\text{CH}_2\text{CH}=\text{CHCH}_3$, $-\text{CH}=\text{CH}_2\text{CH}_2\text{CH}_3$, $-\text{CH}=\text{CH}_2\text{CH}=\text{CH}_2$, and cyclobutenyl.

"Alkoxy", or "alkylthio", as used herein, refers to an alkyl group, as previously defined, attached to the principal carbon chain through an oxygen ("alkoxy") or sulfur ("alkylthio") atom.

"Methylene", "ethylene", and "propylene" as used herein refer to the bivalent moieties $-\text{CH}_2-$, $-\text{CH}_2\text{CH}_2-$, and $-\text{CH}_2\text{CH}_2\text{CH}_2-$, respectively.

"Ethenylene", "propenylene", and "butenylene" as used herein refer to the bivalent moieties $-\text{CH}=\text{CH}-$, $-\text{CH}=\text{CHCH}_2-$, $-\text{CH}_2\text{CH}=\text{CH}-$, $-\text{CH}=\text{CHCH}_2\text{CH}_2-$, $-\text{CH}_2\text{CH}=\text{CH}_2\text{CH}_2-$, and $-\text{CH}_2\text{CH}_2\text{CH}=\text{CH}-$, where each ethenylene, propenylene, and butenylene group can be in the cis or trans configuration. In certain embodiments, an ethenylene, propenylene, or butenylene group can be in the trans configuration.

"Alkylidene" refers to a bivalent hydrocarbon group formed by mono or dialkyl substitution of methylene. In certain embodiments, an alkylidene group has 1-6 carbon atoms. In other embodiments, an alkylidene group has 2-6, 1-5, 2-4, or 1-3 carbon atoms. Such groups include propylidene ($\text{CH}_3\text{CH}_2\text{CH}=\text{}$), ethylidene ($\text{CH}_3\text{CH}=\text{}$), and isopropylidene ($\text{CH}_3(\text{CH}_3)\text{CH}=\text{}$), and the like.

"Alkenylidene" refers to a bivalent hydrocarbon group having one or more double bonds formed by mono or dialkenyl substitution of methylene. In certain embodiments, an alkenylidene group has 2-6 carbon atoms. In other embodiments, an alkenylidene group has 2-6, 2-5, 2-4, or 2-3 carbon atoms. According to one aspect, an alkenylidene has two double bonds. Exemplary alkenylidene groups include $\text{CH}_3\text{CH}=\text{C}=\text{}$, $\text{CH}_2=\text{CHCH}=\text{}$, $\text{CH}_2=\text{CHCH}_2\text{CH}=\text{}$, and $\text{CH}_2=\text{CHCH}_2\text{CH}=\text{CHCH}=\text{}$.

" C_{1-6} alkyl ester or amide" refers to a C_{1-6} alkyl ester or a C_{1-6} alkyl amide where each C_{1-6} alkyl group is as defined above. Such C_{1-6} alkyl ester groups are of the formula $(\text{C}_{1-6} \text{ alkyl})\text{OC}(=\text{O})-$ or $(\text{C}_{1-6} \text{ alkyl})\text{C}(=\text{O})\text{O}-$. Such C_{1-6} alkyl amide groups are of the formula $(\text{C}_{1-6} \text{ alkyl})\text{NHC}(=\text{O})-$ or $(\text{C}_{1-6} \text{ alkyl})\text{C}(=\text{O})\text{NH}-$.

" C_{2-6} alkenyl ester or amide" refers to a C_{2-6} alkenyl ester or a C_{2-6} alkenyl amide where each C_{2-6} alkenyl group is as defined above. Such C_{2-6} alkenyl ester groups are of the formula $(\text{C}_{2-6} \text{ alkenyl})\text{OC}(=\text{O})-$ or $(\text{C}_{2-6} \text{ alkenyl})\text{C}(=\text{O})\text{O}-$. Such C_{2-6} alkenyl amide groups are of the formula $(\text{C}_{2-6} \text{ alkenyl})\text{NHC}(=\text{O})-$ or $(\text{C}_{2-6} \text{ alkenyl})\text{C}(=\text{O})\text{NH}-$.

"Halo" refers to fluoro, chloro, bromo or iodo.

"Haloalkyl" refers to an alkyl group substituted with one or more halo atoms. For example, "fluoromethyl" refers to a methyl group substituted with one or more fluoro atoms (e.g., monofluoromethyl, difluoromethyl, trifluoromethyl).

"Hydroxyalkyl" refers to an alkyl group substituted with a hydroxyl group ($-\text{OH}$).

"Fluoromethoxy" as used herein, refers to a fluoromethyl group, as previously defined, attached to the principal carbon chain through an oxygen atom.

"Protecting group" as used herein, is meant that a particular functional moiety, e.g., O, S, or N, is temporarily blocked so that a reaction can be carried out selectively at another reactive site in a multifunctional compound. For example, in certain embodiments, as detailed herein, certain exemplary oxygen protecting groups are utilized. Oxygen protecting groups include, but

are not limited to, groups bonded to the oxygen to form an ether, such as methyl, substituted methyl (e.g., Trt (triphenylmethyl), MOM (methoxymethyl), MTM (methylthiomethyl), BOM (benzyloxymethyl), PMBM or MPM (p-methoxybenzyloxymethyl)), substituted ethyl (e.g., 2-(trimethylsilyl)ethyl), benzyl, substituted benzyl (e.g., para-methoxybenzyl), silyl (e.g., TMS (trimethylsilyl), TES (triethylsilyl), TIPS (triisopropylsilyl), TBDMS (t-butyldimethylsilyl), tribenzylsilyl, TBDPS (t-butyldiphenyl silyl), 2-trimethylsilylprop-2-enyl, t-butyl, tetrahydropyranyl, allyl, etc.

In some embodiments, the compounds may be provided as a salt, such as a pharmaceutically acceptable salt. Pharmaceutically acceptable salts are salts that retain the desired biological activity of the parent compound and do not impart undesired toxicological effects. Specific examples of pharmaceutically acceptable salts include inorganic acid salts (such as sulfates, nitrates, perchlorates, phosphates, carbonates, bicarbonates, hydrofluorides, hydrochlorides, hydrobromides and hydroiodides), organic carboxylates (such as acetates, oxalates, maleates, tartrates, fumarates and citrates), organic sulfonates (such as methanesulfonates, trifluoromethanesulfonates, ethanesulfonates, benzenesulfonates, toluenesulfonates and camphorsulfonates), amino acid salts (such as aspartates and glutamates), quaternary amine salts, alkali metal salts (such as sodium salts and potassium salts) and alkali earth metal salts (such as magnesium salts and calcium salts).

Unless indicated otherwise, nomenclature used to describe chemical groups or moieties as used herein follow the convention where, reading the name from left to right, the point of attachment to the rest of the molecule is at the right-hand side of the name. For example, the group "(C₁₋₃ alkoxy)C₁₋₃ alkyl," is attached to the rest of the molecule at the alkyl end. Further examples include methoxyethyl, where the point of attachment is at the ethyl end, and methylamino, where the point of attachment is at the amine end.


Unless indicated otherwise, where a mono or bivalent group is described by its chemical formula, including one or two terminal bond moieties indicated by "-", it will be understood that the attachment is read from left to right.

Unless otherwise stated, structures depicted herein are also meant to include all enantiomeric, diastereomeric, and geometric (or conformational) forms of the structure; for example, the R and S configurations for each asymmetric center, (Z) and (E) double bond isomers, and (Z) and (E) conformational isomers. Therefore, single stereochemical isomers as well as enantiomeric, diastereomeric, and geometric (or conformational) mixtures of the present compounds are within the scope of the invention. Unless otherwise stated, all tautomeric forms of the compounds of the invention are within the scope of the invention.

Additionally, unless otherwise stated, structures depicted herein are also meant to include compounds that differ only in the presence of one or more isotopically enriched atoms. For example, compounds having the present structures except for the replacement of hydrogen by deuterium or tritium, or the replacement of a carbon by a ^{13}C - or ^{14}C -enriched carbon are within the scope of this invention. Such compounds are useful, for example, as analytical tools or probes in biological assays.

B. Compounds and Chemical Synthesis

Provided herein are compounds of **Formula I**:



I

wherein:

R_1 is alkyl, haloalkyl, hydroxyalkyl, or $-(C_{1-6}\text{alkyl})-OR'$, wherein R' is an oxygen protecting group; and


R_2 and R_3 are each independently hydrogen or halo (e.g., fluoro); or a salt thereof.

In some embodiments, **Formula I** is subject to the proviso that when one of R_2 or R_3 is *ortho*-fluoro, and the other is hydrogen, R_1 is not $-\text{CH}_2\text{F}$.

In some embodiments, R_1 is trifluoromethyl, hydroxymethyl, monofluoromethyl, or triphenylmethoxymethyl.


In some embodiments, R_2 is fluoro, and/or R_3 is hydrogen or fluoro.

Examples of compounds of **Formula I** include:



Also provided are methods of making a compound of **Formula I** as defined above. In some embodiments, the method includes the step of heating a mixture including:

(i) an oxime of **Formula A**:



A

wherein R₁, R₂ and R₃ are as given above for **Formula I**;

(ii) hydroquinone; and


(iii) a hydrocarbon solvent (e.g., benzene, toluene, xylenes, etc.),

to a temperature of from 90, 95, 100, 105, or 110 °C, to 115, 120, 125, 130, 135, 140, 145, 150, 155, or 160 °C (for example, from 90°C to 160°C, or from 110°C to 140°C, or from 100°C to 120°C, or from 120°C to 140°C),

to make said compound of **Formula I**, or a salt thereof.

In some embodiments, the method includes the step of heating a mixture comprising:

(i) a compound of **Formula B**:



B

wherein R₁, R₂ and R₃ are as given above;

(ii) an acetate;


(iii) hydroxylamine or a salt thereof; and

(iv) a hydrocarbon solvent (e.g., benzene, toluene, xylenes, etc.),

to a temperature of from 90, 95, 100, 105, or 110 °C, to 115, 120, 125, 130, 135, 140, 145, 150, 155, or 160 °C (for example, from 90°C to 160°C, or from 110°C to 140°C, or from 100°C to 120°C, or from 120°C to 140°C),

to make said compound of **Formula I**, or a salt thereof.

Further provided are compounds of **Formula II**:

**II**

wherein:

5 R_1 is alkyl, haloalkyl, hydroxyalkyl, or $-(C_{1-6})-OR'$, wherein R' is an oxygen protecting group; and


R_2 and R_3 are each independently hydrogen or halo (e.g., fluoro);
or a salt thereof.

10 In some embodiments, **Formula II** is subject to the proviso that when one of R_2 or R_3 is *ortho*-fluoro, and the other is hydrogen, R_1 is not $-CH_2F$.


In some embodiments, R_1 is trifluoromethyl, hydroxymethyl, monofluoromethyl, or triphenylmethoxymethyl.

In some embodiments, R_2 is fluoro, and/or R_3 is hydrogen or fluoro.

15 Examples of compounds of **Formula II** include:



Also provided are methods of making a compound of **Formula II**:

**II**

wherein:

20


R_1 is C_{1-4} alkyl; and

R_2 and R_3 are each independently hydrogen or halo;
or a salt thereof,

5 comprising the steps of:

(a) heating a mixture comprising:

(i) a ketone of **Formula B**:



B

10 wherein R_1 , R_2 and R_3 are as given above for **Formula II**;

(ii) hydroxylamine or a salt thereof;

(iii) a base (e.g., acetate, propionate, a tertiary amine base such as triethylamine or diethylisopropylamine, pyridine, or a salt thereof); and

(iv) a solvent selected from: water and an alcohol (e.g., a C_{3-8} alcohol),

15 to at least 90 or 100° Celsius (e.g., to boiling such as by refluxing) produce a heated mixture,

(b) cooling the heated mixture to a temperature of at or below 60, 50 or 45° Celsius, and then,


(c) adding an organic solvent (e.g., tetrahydrofuran or an ether solvent such as diethyl ether or methyl *tert*-butyl ether), an acid (e.g., a carboxylic acid such as acetic acid, or a mineral acid such as hydrochloric acid or sulfuric acid), and zinc (e.g., zinc powder or zinc dust) to the mixture, wherein said mixture is maintained at or below 60, 50 or 45 degrees Celsius (e.g., between 25, 30 or 35, and 45, 50 or 60 degrees Celsius) during said adding,

20 to make said compound of **Formula II**.

25

In some embodiments, the heating is carried out for at least 30 hours.


Also provided is a method of making a stereochemically pure compound of **Formula IIa**:

**IIa**

wherein R_1 , R_2 and R_3 are as given above for **Formula II**,
or a salt thereof,

5 including the steps of :

(a) adding a mixture of stereoisomers of the compound of **Formula II**:

**II**

wherein R_1 , R_2 and R_3 are as given above,

10 to a solvent (e.g., an alcohol such as ethanol or isopropanol, or a mixture such as a toluene – acetonitrile mixture, a toluene – acetone mixture, a toluene – tetrahydrofuran mixture, or an alcohol – water mixture), which solvent includes a chiral carboxylic acid compound (e.g., D-dibenzoyl tartaric acid, L-dibenzoyl tartaric acid, D-ditoluyl tartaric acid, and L-ditoluyl tartaric acid, etc.) to form a mixture of diastereomeric salts of the compound of **Formula II**; and
15 then


(b) crystallizing a single diastereomeric salt formed of the compound of **Formula II**,
to make the stereochemically pure compound of **Formula IIa**, or a salt thereof.

In some embodiments, the stereochemically pure compound has greater than about 80%
20 by weight of one stereoisomer of the compound and less than about 20% by weight of other stereoisomers of the compound, more preferably greater than about 90% by weight of one stereoisomer of the compound and less than about 10% by weight of the other stereoisomers of the compound, even more preferably greater than about 95% by weight of one stereoisomer of the compound and less than about 5% by weight of the other stereoisomers of the compound, and
25 most preferably greater than about 97% by weight of one stereoisomer of the compound and less than about 3% by weight of the other stereoisomers of the compound greater than about 90% by

weight of one stereoisomer of the compound and less than about 10% by weight of the other stereoisomers of the compound.

In some embodiments, the crystallizing step is performed by cooling the mixture of diastereomeric salts of the compound of Formula II (e.g., to a temperature of from 40, 45, 50, 55, or 60 °C to 80, 90, 100, or 110 °C, to a temperature of from -10, -5, 0, 5, or 10 °C to 15, 20, 29, or 35 °C).

Further provided are compounds of **Formula III**:



III

wherein:

R₁ is alkyl, haloalkyl, hydroxyalkyl, or -(C₁₋₆)-OR', wherein R' is an oxygen protecting group; and


R₂ and R₃ are each independently hydrogen or halo (e.g., fluoro); or a salt thereof.

In some embodiments, **Formula III** is subject to the proviso that when one of R₂ or R₃ is *ortho*-fluoro, and the other is hydrogen, R₁ is not -CH₂F.


In some embodiments, R₁ is trifluoromethyl, hydroxymethyl, monofluoromethyl, or triphenylmethoxymethyl.

In some embodiments, R₂ is fluoro, and/or R₃ is hydrogen or fluoro.

Examples of compounds of **Formula III** include:



Also provided are compounds of **Formula IV**:



IV

5 wherein:

R_1 is alkyl, haloalkyl, hydroxyalkyl, or $-(C_{1-6})-OR'$, wherein R' is an oxygen protecting group; and

R_2 and R_3 are each independently hydrogen or halo (e.g., fluoro);
or a salt thereof.

10


In some embodiments, **Formula IV** is subject to the proviso that when one of R_2 or R_3 is *ortho*-fluoro, and the other is hydrogen, R_1 is not $-CH_2F$.

In some embodiments, R_1 is trifluoromethyl, hydroxymethyl, monofluoromethyl, or triphenylmethoxymethyl.

15


In some embodiments, R_2 is fluoro, and/or R_3 is hydrogen or fluoro.

Examples of compounds of **Formula IV** include:



20

Further provided are compounds of **Formula V**:



V

wherein:

R_1 is alkyl, haloalkyl, hydroxyalkyl, or $-(C_{1-6})-OR'$, wherein R' is an oxygen protecting group; and

R_2 and R_3 are each independently hydrogen or halo (e.g., fluoro);


or a salt thereof.

In some embodiments, **Formula V** is subject to the proviso that when one of R_2 or R_3 is *ortho*-fluoro, and the other is hydrogen, R_1 is not $-CH_2F$.


In some embodiments, R_1 is trifluoromethyl, hydroxymethyl, monofluoromethyl, or triphenylmethoxymethyl.

In some embodiments, R_2 is fluoro, and/or R_3 is hydrogen or fluoro.

Examples of compounds of **Formula V** include:



Also provided are methods for making a compound of **Formula V**:



wherein:


R_1 is C_{3-4} alkyl, halo- C_{1-4} alkyl, hydroxy- C_{1-4} alkyl, or $-C_{1-4}$ alkyl- OR' , wherein R' is an oxygen protecting group; and

R_2 and R_3 are each independently hydrogen or halo; or a salt thereof,

comprising the steps of:

(a) providing a mixture of

(i) a compound of **Formula III**,



III

wherein R₁, R₂ and R₃ are as given above for **Formula V**;

(ii) an amine base (e.g., pyridine, or a substituted pyridine such as collidine); and

(iii) an organic solvent (e.g., an aromatic hydrocarbon solvent, toluene, xylenes, benzene, dichloromethane, etc.),

wherein said mixture is at a temperature at or below 0° Celsius,

(b) reacting the mixture with trifluoromethanesulfonic anhydride, 4-toluenesulfonyl chloride, methanesulfonyl chloride or methanesulfonyl anhydride, and then,

(c) adding:

(i) an alcohol (e.g., propanol, such as 2-propanol); and


(ii) a base, such as a strong base selected from: a hydroxide, an alkoxide,

and ammonia,

to make said compound of **Formula V**.

In some embodiments, the reacting step is carried out at a temperature of from -15, -10, or -5, to 0° Celsius.

Also provided are compound of **Formula VI**:



VI

wherein:

R₁ is alkyl, haloalkyl, hydroxyalkyl, or -(C₁₋₆)-OR', wherein R' is an oxygen protecting group; and


R_2 and R_3 are each independently hydrogen or halo (e.g., fluoro);
or a salt thereof.

In some embodiments, **Formula VI** is subject to the proviso that when one of R_2 or R_3 is
5 *ortho*-fluoro, and the other is hydrogen, R_1 is not $-\text{CH}_2\text{F}$.


In some embodiments, R_1 is trifluoromethyl, hydroxymethyl, monofluoromethyl, or
triphenylmethoxymethyl.

In some embodiments, R_2 is fluoro, and/or R_3 is hydrogen or fluoro.

10 Examples of compounds of **Formula VI** include:



Further provided are compounds of **Formula VII**:



VII

15 wherein:


R_1 is alkyl, haloalkyl, hydroxyalkyl, or $-(\text{C}_{1-6})-\text{OR}'$, wherein R' is an oxygen protecting
group; and

R_2 and R_3 are each independently hydrogen or halo (e.g., fluoro);
20 or a salt thereof.

In some embodiments, R_1 is methyl, trifluoromethyl, hydroxymethyl, monofluoromethyl,
or triphenylmethoxymethyl.


In some embodiments, R_2 is fluoro, and/or R_3 is hydrogen or fluoro.

25 Examples of compound of **Formula VII** include:



Also provided are methods of making a compound of **Formula VII**, or a salt thereof, including the step of:

- 5 reducing a compound of **Formula VI** to form a compound of **Formula VII**, as shown below:



wherein:

- 10 R_1 is alkyl, haloalkyl, hydroxyalkyl, or $-(C_{1-6})-OR'$, wherein R' is an oxygen protecting group; and

R_2 and R_3 are each independently hydrogen or halo.

- 15 In some embodiments, R_1 is trifluoromethyl, hydroxymethyl, monofluoromethyl, or triphenylmethoxymethyl.

In some embodiments, R_2 is fluoro, and/or R_3 is hydrogen or fluoro.

In some embodiments, the reducing of the compound of **Formula VI** to form the compound of **Formula VII** may include one or more of the steps of:

- 20 (a) adding powdered iron to a carbinol solvent to form a mixture thereof;
 (b) adding hydrochloric acid (HCl) (e.g., concentrated HCl) to the mixture of step (a);
 (c) heating the mixture of step (b) to a temperature of from 31, 40 or 50 °C to 65, 70, 75, 80, 85°C, or to the boiling point of the mixture of step (b) (e.g., for 0 - 4 hours) to form a heated mixture; and
 25 (d) adding the compound of **Formula VI**, or a salt thereof, to the heated mixture of step (c).


In some embodiments, the method further includes the steps of:

(e) cooling the mixture of step (d) to a temperature of between 10 and 30°C to form a cooled mixture; and

5 (f) optionally, stirring the cooled mixture (e.g., for 0 - 4 hours).

In some embodiments, the HCl is provided in an amount selected from the group consisting of: an amount of HCl stoichiometric with respect to the amount of the compound of **Formula VI**; and an amount of HCl in molar excess with respect to the amount of the compound of **Formula VI**.

Further provided are methods of making a compound of **Formula IX**:



wherein:


15 R_1 is alkyl, haloalkyl, hydroxyalkyl, or $-(C_{1-6})-OR'$, wherein R' is an oxygen protecting group; and

R_2 and R_3 are each independently hydrogen or halogen; and

R_4 is selected from the group consisting of alkyl, haloalkyl, alkoxy, and alkyl substituted with alkoxy;

20 or a salt thereof;

including the step of reacting a compound **Formula VII** with a pyrazine carboxylic acid of **Formula VIII**:



wherein R₁, R₂, R₃ and R₄ are as given above,
to form said compound of **Formula IX**, or a salt thereof.

In some embodiments, the reacting may include one or more of the steps of:

- 5 (a) adding thionyl chloride to a solution comprising a compound of **Formula VIII** in a solvent to make a formed mixture; and
- (b) adding a compound of **Formula VII** to the formed mixture of step (a),
to form said compound of **Formula IX**, or a salt thereof.

- 10 In some embodiments, the solution comprising the compound of **Formula VIII** is a solvent is cooled prior to or during the adding step (a) (e.g., to a temperature of from 0 to 25°C).

In some embodiments, the solvent may include N,N'-dimethylimidazoline-2-one, toluene, methylene chloride, dimethylformamide, N-methylpyrrolidine, or dimethylacetamide.


15

In some embodiments, the reacting may include one or more of the steps of:

- (a) adding ethyl acetate (EtOAc) to a compound of **Formula VIII** and a compound of **Formula VII** to make a formed mixture; and
- (b) adding propane phosphonic acid anhydride (e.g., [®]T3P, Archimica, Germany) to the
- 20 formed mixture of step (a), to form said compound of **Formula IX**, or a salt thereof.

In some embodiments, the formed mixture may be maintained at a temperature of from 0°C to 35°C during said adding step (a).

- 25 These compounds, or salts thereof, as well as methods of making, are useful in methods of synthesis of fused aminodihydrothiazine derivatives, such as the method provided below in **Scheme I**.



Scheme I

In order that the invention described herein may be more fully understood, the following examples are set forth. It should be understood that these examples are for illustrative purposes only and are not to be construed as limiting this invention in any manner.

EXAMPLES

General:

Column chromatography was carried out using Biotage SP4. Solvent removal was carried out using either a Büchii rotary evaporator or a Genevac centrifugal evaporator. Preparative LC/MS was conducted using a Waters autopurifier and 19 x 100mm XTerra 5 micron MS C18 column under acidic mobile phase condition. NMR spectra were recorded using Varian 400MHz spectrometer.

When the term "inerted" is used to describe a reactor (e.g., a reaction vessel, flask, glass reactor, and the like) it is meant that the air in the reactor has been replaced with an essentially

moisture-free or dry, inert gas (such as nitrogen, argon, and the like). The term "equivalent" (abbreviation: eq) as used herein describes the stoichiometry (molar ratio) of a reagent or a reacting compound by comparison to a pre-established starting material. The term "weight" (abbreviation: wt) as used herein corresponds to the ratio of the mass of a substance or a group of substances by comparison to the mass of a particular chemical component of a reaction or purification specifically referenced in the examples below. The ratio is calculated as: g/g, or Kg/Kg. The term "volume" (abbreviation: vol) as used herein corresponds to the ratio of the volume of a given substance or a group of substances to the mass or volume of a pre-established chemical component of a reaction or purification. The units used in the equation involve matching orders of magnitude. For example, a ratio is calculated as: mL/mL, mL/g, L/L or L/Kg.

General methods and experimentals for preparing compounds of the present invention are set forth below. In certain cases, a particular compound is described by way of example. However, it will be appreciated that in each case a series of compounds of the present invention were prepared in accordance with the schemes and experimentals described below.


The following abbreviations are used herein:

Abbreviation	Definition
TMS	Trimethylsilyl
TBAF	Tetrabutylammonium fluoride
NaOH	Sodium hydroxide
Bu ₄ N HSO ₄	Tetrabutylammonium hydrogen sulfate
THF	Tetrahydrofuran
rt	Room temperature
h	Hour(s)
NaCl	Sodium chloride
HCOOH	Formic acid
V	Volumes
wt	Weights
CDI	N,N-Carbonyldiimidazole
DCM	Dichloromethane
Aq	Aqueous
Sat.	Saturated
HCl	Hydrochloric acid
HRMS	High Resolution Mass Spectrometry
nBuLi	n-butyl lithium
NH ₄ Cl	Ammonium chloride
MeOH	Methanol
EtOAc	Ethyl acetate
NaHCO ₃	Sodium bicarbonate
M	Molar (moles/liter)
T	Temperature
MTBE	Methyl <i>tert</i> -butyl ether
TLC	Thin layer chromatography

N	Normal (equivalents per liter)
iPrMgBr	Isopropyl magnesium bromide
LiCl	Lithium chloride
NaOAc	Sodium acetate
NH ₄ OH	Ammonium hydroxide
HPLC	High performance liquid chromatography
ee	Enantiomeric excess
DMI	1,3-Dimethyl-2-imidazolidinone
UV	Ultraviolet
RRT	Relative retention time
OROT	Optical rotation
Bz	Benzoyl
[®] T3P (Archimica)	n-propyl phosphonic acid anhydride

A. Preparation of Hexahydrofuroisoxazoles Compounds of Formula I


5 Example 1: Preparation of 6a-(2-fluorophenyl)-4-(trifluoromethyl)hexahydrofuro- [3,4-c]isoxazole.



tert-Butyl 2-(1,1,1-trifluorobut-3-en-2-yloxy)acetate. A reaction vessel was charged with
 10 toluene (3.2 L), THF (0.60 L) and acrolein (0.40 L, 5.985 mol) at rt under nitrogen.
 (Trifluoromethyl)trimethylsilane (1.003 kg, 7.059 mol) was added at 17 °C. The reaction mixture
 was cooled to 2.5 °C and TBAF (0.01 M in THF, 0.400 L, 0.004 mol) was added over 2 hours.
 During addition of TBAF, the temperature of the reaction mixture increased to 65 °C. The
 reaction mixture was cooled to 0 °C, and after 2h, tetra-*n*-butylammonium hydrogen sulfate
 15 (0.171 kg, 0.503 mol) was added, followed by *tert*-butyl bromoacetate (0.987 kg, 5.064 mol).
 Sodium hydroxide (50% wt in water, 4.2 kg, 52.6 mol) was added over 2h while maintaining the
 temperature under 10 °C. After 2h at 0-5 °C, to the reaction mixture was added water (2.9 L) and
 methyl *tert*-butyl ether (6.0 L). The aq. phase was extracted one more time with methyl *tert*-butyl
 ether (6.0 L). The organic phases were combined and washed with 14% aq. NaCl (3 x 1.6 L).
 20 The organics were concentrated under vacuum to afford the title compound **1** as an oil (1.150 kg,
 94.5%) which was used in the subsequent stage without additional purification.

tert-Butyl 2-(1,1,1-trifluorobut-3-en-2-yloxy)acetate: ¹H NMR (500 MHz, CDCl₃) δ 5.86 – 5.74 (m, 1H), 5.59 (d, *J* = 17.5 Hz, 1H), 5.56 (d, *J* = 10.9 Hz, 1H), 4.37 – 4.30 (m, 1H), 4.11 (d,

$J = 16.5$ Hz, 1H), 4.06 (d, $J = 16.4$ Hz, 1H), 1.40 (s, 9H); ^{13}C NMR (125 MHz, CDCl_3) δ 168.51, 128.49 (d, $J = 1.7$ Hz), 123.86, 123.71 (q, $J = 281.8$ Hz), 82.22, 78.67 (q, $J = 31.5$ Hz), 66.60, 28.02.



5


***N*-Methoxy-*N*-methyl-2-(1,1,1-trifluorobut-3-en-2-yloxy)acetamide.** To a reactor containing *tert*-butyl 2-(1,1,1-trifluorobut-3-en-2-yloxy)acetate **1** (1.150 kg, 4.788 mol) was added formic acid (6.2 kg) at rt. The reaction mixture was heated to 55-60 °C for 4-5 hours. The formic acid was evaporated under vacuum ($T = 40-45$ °C) and chased with toluene (2 x 3.0 L). To the residue was added CH_2Cl_2 (2.0 L) and further concentrated under vacuum. To the resulting residue was added CH_2Cl_2 (4.6 L) and the solution was cooled to 0 °C, followed by *N,N*-carbonyldiimidazole (1.05kg, 6.49 mol) in five portions. The mixture was stirred for 30 minutes, and *N,O*-dimethylhydroxylamine hydrochloride (0.67 kg, 6.72 mol) was added in portions while maintaining temperature below 10 °C. The reaction mixture was warmed to rt and stirred over 14 hours. The reaction mixture was cooled to 3.2 °C and imidazole (100.7 g, 1.48 mol) was charged in two portions. The reaction mixture was warmed to rt and water (1.4 kg) was added, followed by methyl *tert*-butyl ether (14.0 L). The organic phase was washed with 2.0 N aq. HCl (1.0 L and 0.7 L), followed by sat. aq. NaHCO_3 (1.2 L) and sat. aq. NaCl (1.20 L). The organics were concentrated to afford the title compound **2** as an oil (0.95kg, 87.2%).

20

N-Methoxy-*N*-methyl-2-(1,1,1-trifluorobut-3-en-2-yloxy)acetamide: ^1H NMR (500 MHz, CDCl_3) δ 5.85 – 5.76 (m, 1H), 5.62 (d, $J = 17.2$ Hz, 1H), 5.56 (d, $J = 10.4$ Hz, 1H), 4.49 – 4.34 (m, 3H), 3.68 (s, 3H), 3.67 (s, 1H), 3.18 (s, 3H), 3.08 (s, 1H); ^{13}C NMR (126 MHz, cdcl_3) δ 169.9*, 163.4*, 128.61, 123.87 (d, $J = 282.0$ Hz), 123.82, 78.54 (q, $J = 31.3$ Hz), 66.12, 61.52, 60.56, 36.20, 32.24. Note: this compound is a 3:1 mixture of amide bond rotamers. *Carbonyl chemical shifts estimated indirectly through ^1H - ^{13}C HMBC (heteronuclear multiple-bond correlation).

25


HRMS Calculated for $\text{C}_8\text{H}_{12}\text{F}_3\text{NO}_3$ $[\text{M}+\text{H}]^+$ 228.0848; found 228.0836.



1-(2-Fluorophenyl)-2-(1,1,1-trifluorobut-3-en-2-yloxy)ethanone. To a solution 1-bromo-2-fluorobenzene (0.967 kg, 5.527 mol) in THF (6.2 L) at -75 °C, was added n-butyllithium (2.50 M in hexane, 2.09 L, 5.22 mol) while maintaining temperature below -65 °C (ca. 100 min.). After 5 15 minutes, a solution of *N*-methoxy-*N*-methyl-2-(1,1,1-trifluorobut-3-en-2-yloxy)acetamide **2** (0.949 kg, 4.178 mol) in THF (1.6 L) was added while maintaining temperature below -65 °C (ca. 70 min.). After 2.5 h at -78 °C, the reaction was quenched by addition of sat. aq. NH₄Cl (3.0 L) and methyl *tert*-butyl ether (9.0 L). The reaction mixture was warmed to rt, the aq. phase was extracted again with methyl *tert*-butyl ether (2.5 L). The organic phases were combined, washed 10 with sat. aq. NaCl (2 x 0.3 L) and concentrated under vacuum to afford the title compound **3** as an oil (1.007kg, 80.0%).

1-(2-Fluorophenyl)-2-(1,1,1-trifluorobut-3-en-2-yloxy)ethanone: ¹H NMR (500 MHz, CDCl₃) δ 7.96 (td, *J* = 7.6, 1.8 Hz, 1H), 7.62 – 7.54 (m, 1H), 7.33 – 7.25 (m, 1H), 7.20 – 7.12 (m, 1H), 5.86 (ddd, *J* = 17.5, 10.4, 7.3 Hz, 1H), 5.60 (dd, *J* = 20.5, 13.8 Hz, 2H), 4.91 – 4.76 (m, 2H), 4.39 (dq, *J* = 12.8, 6.4 Hz, 1H); ¹³C NMR (125 MHz, CDCl₃) δ 193.55, 162.14 (d, *J*_{CF} = 254.1 Hz), 135.36 (d, *J*_{CF} = 9.2 Hz), 130.62 (d, *J*_{CF} = 3.2 Hz), 128.49, 124.85 (d, *J*_{CF} = 3.3 Hz), 123.89, 122.93, 122.72 (d, *J*_{CF} = 24.5 Hz), 116.50 (d, *J*_{CF} = 23.7 Hz), 78.97 (q, *J*_{CF} = 31.4 Hz), 74.56 (d, *J*_{CF} = 12.4 Hz).

HRMS Calculated for C₁₂H₁₀F₄O₂ [M+H]⁺ 263.0695; found 263.0709.




1-(2-Fluorophenyl)-2-(1,1,1-trifluorobut-3-en-2-yloxy)ethanone oxime. To a reactor was added hydroxylamine hydrochloride (0.34 kg, 4.95 mol), sodium acetate (0.47 kg, 5.70 mol) and MeOH (2.68 L). To this suspension was charged a solution of 1-(2-fluorophenyl)-2-(1,1,1-trifluorobut-3-en-2-yloxy)ethanone **3** (0.998 kg, 3.806 mol) in MeOH (1.8 L) and the reaction 25 mixture was heated to 40-50 °C. Upon completion (ca. 2 h) the reaction mixture was cooled to rt, and filtered over Celite (0.5 wt) and rinsed with EtOAc (3.0 L). The filtrate was concentrated under vacuum and to the resulting residue was added methyl *tert*-butyl ether (6.3 L), water (0.94

L) and sat. aq. NaHCO₃ (2.5 L). The organic phase was washed once with water (1.6 L) and sat. aq. NaCl (0.1 L). The organic phase was concentrated under vacuum to afford the title compound **4** as an oil (1.03 kg, 95.0%).

1-(2-Fluorophenyl)-2-(1,1,1-trifluorobut-3-en-2-yloxy)ethanone oxime: ¹H NMR (500 MHz, CDCl₃) δ 7.49 – 7.35 (m, 2H), 7.24 – 7.06 (m, 2H), 5.78 – 5.65 (m, 1H), 5.54 – 5.40 (m, 2H), 4.89 – 4.81 (m, 1H), 4.53 (d, *J* = 12.6 Hz, 1H), 4.47 (d, *J* = 12.6 Hz, 0.5H), 4.27 – 4.18 (m, 1H), 4.13 – 4.05 (m, 0.5H).

HRMS Calculated for C₁₂H₁₁F₄NO₂ [M+H]⁺ 278.0804; found 278.0780.


Note: Oxime **4** exists as an equilibrium of structural isomers, which accounts for the less-than-whole-number integral values.



6a-(2-Fluorophenyl)-4-(trifluoromethyl)hexahydrofuro[3,4-c]isoxazole. To a solution of 1-(2-fluorophenyl)-2-(1,1,1-trifluorobut-3-en-2-yloxy)ethanone oxime **4** (1.085 kg, 3.328 mol) in xylenes (6.9 L) was added hydroquinone (86.2g, 0.8 mol) at rt. The solution was heated to 128 °C (internal temperature) for 18 h. The solution was cooled to rt, and hexanes (7.0 L) was added, followed by 4.0 M aq. HCl (2.4 L). The reaction mixture was stirred for 1 h, and filtered. To the solid was added water (2.0 L), methyl *tert*-butyl ether (7.0 L) and 25% wt. aq. NaOH (0.4 L). The aq. layer was extracted once with methyl *tert*-butyl ether (7.0 L), the organics were combined, washed with 27% aq. NaCl (2.0 L) and concentrated under vacuum to a black oil **5** (512.0g, 56%).


6a-(2-Fluorophenyl)-4-(trifluoromethyl)hexahydrofuro[3,4-c]isoxazole: ¹H NMR (500 MHz, CDCl₃) δ 7.64 – 7.52 (m, 1H), 7.39 – 7.31 (m, 1H), 7.19 (td, *J* = 7.7, 1.2 Hz, 1H), 7.11 (ddd, *J* = 11.9, 8.2, 1.0 Hz, 1H), 4.54 (d, *J* = 10.1 Hz, 1H), 4.34 – 4.23 (m, 1H), 4.26 – 4.17 (m, 1H), 4.16 (d, *J* = 10.2 Hz, 1H), 4.10 (d, *J* = 8.5 Hz, 1H), 3.71 (d, *J* = 20.2 Hz, 1H); ¹³C NMR (125 MHz, CDCl₃) δ 160.59 (d, *J*_{CF} = 247.0 Hz), 130.50 (d, *J*_{CF} = 8.7 Hz), 128.72, 124.69 (d, *J*_{CF} = 3.3 Hz), 124.45 (q, *J*_{CF} = 281.8 Hz), 124.43 (d, *J*_{CF} = 11.9 Hz), 116.66 (d, *J*_{CF} = 22.7 Hz), 83.70 (q, *J*_{CF} = 32.1 Hz), 78.17 (d, *J*_{CF} = 3.1 Hz), 77.63, 54.53.

HRMS Calculated for C₁₂H₁₁F₄NO₂ [M+H]⁺ 278.0804; found 278.0802.

Example 2: Preparation of 6a-(2-fluorophenyl)-4-methylhexahydrofuro[3,4-c]isoxazole

tert-Butyl 2-(but-3-en-2-yloxy)acetate: A reactor was charged with tetrabutylammonium hydrogen sulfate (0.17 Wt, 0.10 eq) and toluene (2.6 Wt, 3.0 V). The mixture was stirred and cooled to 0-5 °C. While maintaining the internal temperature below 10 °C, 50 wt% aqueous sodium hydroxide (4.5 wt, 3.0 V, 10.5 eq; 50% wt sodium hydroxide made from 2.25 wt of sodium hydroxide and 2.25 wt of water) was added, followed by 3-buten-2-ol (0.45 Wt, 0.53 V, 1.20 eq). The mixture was stirred at 0-10 °C for 15 minutes. *tert*-Butyl bromoacetate (1.0 Wt, 1.00 eq) was added while maintaining the internal temperature between 0-10 °C. After the addition, the mixture was stirred between 0-5 °C for 1h and monitored for complete consumption of *tert*-butyl bromoacetate (target >98% conversion). The reactor was charged with water (3.0 Wt, 3.0 V) and MTBE (4.4 Wt, 6.00 V) and warmed to 20-25 °C. The biphasic mixture was stirred vigorously for 15 minutes, and then allowed to partition. The organic layer was washed with water (1.5 Wt, 1.5 V). The organic layer was concentrated under reduced pressure (T < 40 °C and not less than 15 torr or 20mbar). Heptane (1.0 wts, 1.5 V) and MTBE (1.1 wts, 1.5 V) were added. The solution was filtered to remove particulates, and rinsed with additional heptane:MTBE as necessary to clear the product. The solution was returned to the reactor and concentrated under reduced pressure to give **6** (0.91 Wt, 95.0%) as an oil with less than 1 volume of solvent.

tert-Butyl 2-(but-3-en-2-yloxy)acetate: ^1H NMR (500 MHz, CDCl_3) δ 5.78 – 5.66 (m, 1H), 5.23 – 5.12 (m, 2H), 4.00 – 3.85 (m, 3H), 1.47 (s, 11H), 1.31 (d, $J = 6.4$ Hz, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ 169.87, 139.30, 116.72, 81.20, 77.53, 65.83, 28.04, 27.92, 21.08.




2-(But-3-en-2-yloxy)-N-methoxy-N-methylacetamide: A reactor was charged with the toluene solution of **6** (1.0 Wt, 1.0V), and cooled to 0-5 °C. Formic acid (4.84 Wt, 4.0 V, 19.2 eq) was added while maintaining the internal temperature below 10 °C. After addition of formic acid, the

mixture was heated to 35-40 °C for three hours. The reaction was monitored for consumption of **6** (>90% conversion). Upon completion formic acid was removed under vacuum at 45 °C. Residual formic acid was removed by two azeotropic distillations with toluene (2.0-2.50 Wt, 2.3-3.0 V) at 45 °C under vacuum to give crude acid (0.67 Wt, 95%).


5 The crude acid was dissolved in dichloromethane (5.3 Wt, 4.0 V) and cooled at 0 °C. To the solution is added N,N-carbonyldiimidazole (1.15 Wt, 1.28eq) in 5-7 portions maintaining the reaction temperature between 0-10 °C. Note: Gas evolution occurs during the reaction. Upon complete addition of N,N-carbonyldiimidazole, the mixture was warmed to 15-20 °C and stirred for 30 minutes. The reaction mixture was then cooled back to 0-5 °C. N,O-
10 dimethylhydroxylamine hydrochloride (0.71 Wt, 1.33 eq) was added in portions maintaining the reaction between 0-10 °C. After the addition is complete, the reaction mixture was warmed to 15-20 °C. The reaction was monitored for complete consumption of intermediate acid (>98% conversion). The reactor was charged with water (1.2 Wt, 1.2 V) at 0-10 °C and then MTBE (9.1Wt, 12.3V). The biphasic mixture was stirred vigorously for 15 minutes, and then allowed to
15 partition. The aqueous layer was separated, , the combined organic layer was then washed with 2N HCl (1.8 Wt, 1.8 V), (check pH to make sure it is between 1-3 or check organic layer by TLC to make sure the polar material below product is removed. Charge more 2N HCl if necessary). The aqueous layer was separated, the organics were washed with saturated sodium bicarbonate (NaHCO₃, 0.1 Wt and water, 1.1 Wt, 1.2 V) and then 20% NaCl in water (1.2Wt, 1.2V). The
20 organic phase was concentrated under reduced pressure . Anhydrous THF (1.33 Wt, 1.50 V) was added, and the solution was filtered to remove particulates, rinsing with additional anhydrous THF as necessary to clear the product. The THF solution was returned to the reactor and concentrated under reduced pressure to give **7** (0.73 Wt, 75.0%) as an oil.

25 2-(But-3-en-2-yloxy)-N-methoxy-N-methylacetamide: ¹H NMR (500 MHz, CDCl₃) δ 5.80 – 5.70 (m, 1H), 5.21 (d, *J* = 17.3 Hz, 1H), 5.17 (d, *J* = 10.3 Hz, 1H), 4.28 (d, *J* = 15.6 Hz, 1H), 4.18 (d, *J* = 15.6 Hz, 1H), 4.03 – 3.96 (m, 1H), 3.68 (s, 3H), 3.18 (s, 3H), 1.32 (d, *J* = 6.4 Hz, 3H); ¹³C NMR (125 MHz, CDCl₃) δ 139.62, 116.73, 77.49, 65.34, 61.44, 32.36, 21.25.



2-(But-3-en-2-yloxy)-1-(2-fluorophenyl)ethanone. A reactor was charged with 1-bromo-2-fluorobenzene (1.31 Wt, 1.30 eq) and anhydrous THF (7.60 Wt, 8.6 V). The reaction mixture was cooled to below -70 °C. 2.5 M n-butyl lithium in hexanes (1.94 Wt, 2.89 V, 1.25 eq) was added while maintaining the internal temperature below -60 °C. Upon complete addition, the mixture was stirred at a temperature below -60 °C for 15 minutes and then a solution of amide 7 (1.0 Wt, 1.0 eq.) in THF (1.9 Wt, 2.2 V) was added while maintaining the internal temperature below -60 °C. After stirring the reaction for 60 minutes, the reaction was monitored for complete consumption of amide 7 (>90% conversion). The reactor was charged with 20 wt % aqueous ammonium chloride (1.0 Wt, 1.0 V), and warmed to 20–25 °C. The mixture was transferred to a work-up vessel containing additional 20 wt % aqueous ammonium chloride (3.0 Wt, 3.0 V). The mixture was stirred for 15 minutes, then stirring was stopped and the phases separated. The lower aqueous layer was returned to the reactor. MTBE (4.45 Wt, 6.0 V) was charged. The biphasic mixture was stirred vigorously for 15 minutes, and then allow to partition. The organic layers were combined and washed with 25 wt % aqueous sodium chloride (4.75 Wt, 4.75 V). The organic phase was concentrated under reduced pressure ($T < 40$ °C). Anhydrous THF (1.78 Wt, 2.0 V), was added and the mixture was filtered to remove particulates, rinsing with additional anhydrous THF as necessary to clear the product. The THF solution was returned to the reactor and concentrated under reduced pressure to give ketone 8 (0.8 Wt, 73.0 %) as an oil.


2-(But-3-en-2-yloxy)-1-(2-fluorophenyl)ethanone: ^1H NMR (500 MHz, CDCl_3) δ 7.93 (td, $J = 7.6, 1.8$ Hz, 1H), 7.57 – 7.49 (m, 1H), 7.24 (dt, $J = 14.6, 3.8$ Hz, 1H), 7.12 (ddd, $J = 11.0, 8.3, 0.8$ Hz, 1H), 5.21 (dd, $J = 17.3, 1.3$ Hz, 1H), 5.17 (dd, $J = 10.3, 0.8$ Hz, 1H), 4.69 (dd, $J = 18.1, 3.2$ Hz, 1H), 4.60 (dd, $J = 18.1, 3.4$ Hz, 1H); 4.00 (dq, $J = 12.9, 6.4$ Hz, 1H), 1.36 (d, $J = 6.4$ Hz, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ 195.24 (d, $J_{\text{CF}} = 5.3$ Hz), 162.03 (d, $J_{\text{CF}} = 253.9$ Hz), 139.42, 134.86 (d, $J_{\text{CF}} = 8.9$ Hz), 130.64 (d, $J_{\text{CF}} = 3.4$ Hz), 124.67 (d, $J_{\text{CF}} = 3.2$ Hz), 123.57 (d, $J_{\text{CF}} = 15.3$ Hz), 116.89, 116.45 (d, $J_{\text{CF}} = 23.7$ Hz), 77.93, 74.11 (d, $J_{\text{CF}} = 11.2$ Hz), 21.19.



2-(But-3-en-2-yloxy)-1-(2-fluorophenyl)ethanone oxime. A reactor was charged with crude ketone 8 (1.0 Wt, 1.0 eq), hydroxylamine hydrochloride (0.43 Wt, 1.30 eq.), sodium acetate (0.59 Wt, 1.50 eq.), and methanol (4.6 Wt, 5.8 V). The mixture was warmed to between 50–55 °C, and then continued stirring for 1-2 hr. The reaction was monitored for consumption of

ketone **8** (>95% conversion). The mixture was cooled to 15–20 °C, then filtered the reaction mixture to remove solids. The solids were rinsed with ethyl acetate until product is cleared (3.6 Wt, 4.0 V). The filtrates were returned to the reactor and concentrated under reduced pressure ($T < 40$ °C). MTBE (5.9 Wt, 8.0 V) and water (1.0 Wt, 1.0 V) were charged. The biphasic mixture was stirred vigorously for 15 minutes, and then partitioned. The organic phase was washed sequentially with 10 wt% aqueous sodium bicarbonate (3.2 Wt, 3.2 V), water (2.0 Wt, 2.0 V), and finally 25 wt% aqueous sodium chloride (1.0 Wt, 1.0 V). The organic phase was concentrated under reduced pressure ($T < 40$ °C). Anhydrous THF (1.78 Wt, 2.0 V) was added, and the mixture filtered to remove particulates, rinsing with additional anhydrous THF as necessary to clear the product. The THF solution was returned to the reactor and concentrated under reduced pressure to give oxime **9** (0.8-1.2 Wt, 90.0%) as an oil.

2-(But-3-en-2-yloxy)-1-(2-fluorophenyl)ethanone oxime: ^1H NMR (500 MHz, cdcl_3) δ 7.45 (ddd, $J = 9.2, 6.2, 2.5$ Hz, 1H), 7.38 – 7.30 (m, 1H), 7.20 – 7.03 (m, 2H), 5.73 – 5.58 (m, 1H), 5.21 – 5.05 (m, 2H), 4.67 (s, 2H), 4.38 (d, $J = 13.0$ Hz, 1H), 4.26 (d, $J = 13.0$ Hz, 1H), 3.97 – 3.88 (m, 1H), 3.79 – 3.72 (m, 1H), 1.18 (d, $J = 6.4$ Hz, 1H), 1.05 (d, $J = 6.4$ Hz, 2H); ^{13}C NMR (126 MHz, cdcl_3) δ 160.68 (d, $J_{\text{CF}} = 249.9$ Hz), 159.13 (d, $J_{\text{CF}} = 249.1$ Hz), 156.42, 151.80, 139.39, 130.78 (d, $J_{\text{CF}} = 8.2$ Hz), 130.67 (d, $J_{\text{CF}} = 8.4$ Hz), 130.27 (d, $J_{\text{CF}} = 3.7$ Hz), 129.77 (d, $J_{\text{CF}} = 4.2$ Hz), 123.96 (d, $J_{\text{CF}} = 3.5$ Hz), 123.86 (d, $J_{\text{CF}} = 3.4$ Hz), 122.51 (d, $J_{\text{CF}} = 14.1$ Hz), 120.08 (d, $J_{\text{CF}} = 17.1$ Hz), 116.59, 116.29, 115.72 (d, $J_{\text{CF}} = 7.4$ Hz), 115.55 (d, $J_{\text{CF}} = 7.7$ Hz), 77.45, 76.53, 68.48 (d, $J_{\text{CF}} = 1.7$ Hz), 77.45, 76.53, 68.48 (d, $J_{\text{CF}} = 1.7$ Hz), 62.01 (d, $J_{\text{CF}} = 2.4$ Hz), 20.99, 20.81. Note: there are many ^1H and ^{13}C signals because this compound is a mixture of four possible isomers.




6a-(2-fluorophenyl)-4-methylhexahydrofuro[3,4-c]isoxazole. A reactor was charged with oxime **9** (1.0 Wt, 1.0 eq), xylenes (6.8 Wt, 7.9 V) and then hydroquinone (0.10 Wt, 0.2 eq.). The mixture was heated to between 125–130 °C and monitored the reaction for consumption of oxime **9** (>90% conversion). The mixture was cooled to 20–25 °C. 2N aqueous hydrochloric acid (4.0 Wt, 4.0 V) was charged. The biphasic mixture was stirred vigorously for 15 minutes, and then partitioned the phases. The organic phase was extracted with 2N aqueous hydrochloric acid

(3.0 Wt, 3.0 V). Note: the product resides in the aqueous layers. The aqueous phases were returned to the reactor. Tert-butyl methyl ether (5.8 Wt, 7.9 V) was charged and then sodium hydroxide (0.55 Wt) to adjust the pH to 9-10. The biphasic mixture was stirred vigorously for 15 minutes, and then the phases were partitioned. The aqueous phase was extracted a second time with MTBE (5.8 Wt, 8.0 vol). The combined organic phases were washed with 25 wt% aqueous sodium chloride (1.0 Wt, 1.0 V). The organic phase was concentrated under reduced pressure ($T < 40\text{ }^{\circ}\text{C}$). Anhydrous THF (1.0 Wt, 1.1 V) was added, and filtered to remove particulates, rinsing with additional anhydrous THF as necessary to clear the product. The THF solution was returned to the reactor and concentrated under reduced pressure to give tetrahydroisoxazole **10** (0.6-0.8 Wt, 65.0%) as an oil.

6a-(2-fluorophenyl)-4-methylhexahydrofuro[3,4-c]isoxazole: ^1H NMR (400 MHz, CDCl_3) δ 7.61 (t, $J = 8.0$ Hz, 1H), 7.30 (ddd, $J = 15.0, 5.2, 1.8$ Hz, 1H), 7.19 – 7.03 (m, 2H), 4.36 (d, $J = 10.2$ Hz, 1H), 4.23 – 4.16 (m, 1H), 4.05 (dt, $J = 12.2, 6.2$ Hz, 1H), 3.98 – 3.87 (m, 2H), 3.19 – 3.09 (m, 1H), 1.40 (d, $J = 6.4$ Hz, 3H). Note: no ^{13}C -NMR data was available for this compound.


Alternative procedure #1 for preparation of ketone **8**:



tert-Butyl 2-(but-3-en-2-yloxy)-3-(2-fluorophenyl)-3-oxopropanoate. To a reactor under nitrogen atmosphere was added a toluene solution of ester **6** (1.00 Wt, 1.0 eq), methyl 2-fluorobenzoate (0.91 Wt, 0.75 Vol 1.10 eq), and THF (0.53 Wt, 1.8 Vol). The solution was cooled to below $-70\text{ }^{\circ}\text{C}$. Lithium hexamethyldisilazide (LiHMDS, 1.0M in THF, 7.6 Wt, 8.6 Vol, 1.6 eq) was added at such rate that the internal temperature does not exceed $-60\text{ }^{\circ}\text{C}$. After completion of the LiHMDS addition, the mixture was warmed (rate between $10\text{--}20\text{ }^{\circ}\text{C/h}$) and stirred at $-40\text{ }^{\circ}\text{C}$ for 1h. The reaction was monitored by GC (Conversion Target $\geq 95\%$). If after 4h the conversion is still under targeted amount, additional LiHMDS 1M in THF was added (1.1 eq based on unreacted starting material). After achieving the target conversion, the internal temperature was raised (rate between $20\text{--}30\text{ }^{\circ}\text{C/h}$) to -15 to $-20\text{ }^{\circ}\text{C}$. 25 wt% aqueous ammonium chloride (1.7 Wt, 1.6 Vol) was added to the reactor keeping the internal temperature below $0\text{ }^{\circ}\text{C}$. Upon completion of the addition, the mixture was warmed to $>0\text{ }^{\circ}\text{C}$, and transferred to a vessel

containing 25 wt% aqueous ammonium chloride (4.4 Wt, 4.1 Vol) and water (1.1 Wt, 1.1 Vol). The mixture was stirred for at least 30 min at 20-25 °C. The lower aqueous layer was removed and the organic layer was washed with 20 wt % aqueous sodium chloride (3.0 Wt, 1.4 Vol). The mixture was stirred for at least 15 minutes then allowed to settle. The lower aqueous layer was removed. The organic phase was concentrated under reduce pressure (Jacket < 25 °C) until distillation ceases. Toluene (6.6 Wt, 6.7 Vol) was added and the resulting mixture was washed with water (3.0 Wt, 3.0 Vol), stirred for at least 15 min, then allowed to settle. The organics were washed with water (3.0 Wt, 3.0 Vol) until pH is constant at pH=7-8. The organic phase was washed with 20% aqueous sodium chloride (3.0 Wt, 1.4 Vol). The organic phase was concentrated under reduced pressure (jacket <25°C) to afford ketoester **11** (2.1 Wt) as an orange oil.


tert-Butyl 2-(but-3-en-2-yloxy)-3-(2-fluorophenyl)-3-oxopropanoate: ¹H NMR (500 MHz, CDCl₃) δ 7.88 – 7.78 (m, 1H), 7.57 – 7.48 (m, 1H), 7.29 – 7.19 (m, 1H), 7.17 – 7.07 (m, 1H), 5.85 – 5.64 (m, 1H), 5.28 – 5.13 (m, 2H), 5.09 (s, 1H), 5.05 (s, 1H), 4.18 – 4.10 (m, 1H), 4.02 – 3.94 (m, 1H), 1.39 (s, 9H), 1.37 (s, 9H), 1.35 (d, *J* = 6.3 Hz, 3H), 1.28 (d, *J* = 6.4 Hz, 3H); ¹³C NMR (125 MHz, CDCl₃) δ 193.17, 193.14, 192.33, 192.30, 166.57, 166.16, 162.44, 162.36, 160.42, 160.33, 138.93, 138.80, 134.76, 134.68, 134.61, 131.17, 131.15, 131.05, 131.02, 124.50, 124.47, 124.40, 124.37, 117.86, 117.60, 116.37, 116.34, 116.19, 116.16, 82.95, 82.91, 82.76, 82.73, 82.38, 82.34, 78.75, 78.48, 27.82, 27.76, 21.21, 21.13. Note: this compound is a mixture of diastereomers.




2-(But-3-en-2-yloxy)-1-(2-fluorophenyl)ethanone. To a reactor under nitrogen atmosphere was added ketoester **11** (1.00 Wt, 1.0 eq) and formic acid (10.0 Wt, 8.0 Vol). The resulting mixture was stirred at 25-30°C for 12h then sampled for conversion (Target >97%). After achieving the target conversion, the mixture was concentrated under reduced pressure maintaining the internal temperature below 30°C. Toluene (1.4 Wt, 1.6 Vol) was charged and the mixture was concentrated under reduced pressure maintaining the internal temperature below 30 °C to azeotrope residual formic acid. Toluene (1.4 Wt, 1.6 Vol) was charged and the mixture was concentrated under reduced pressure maintaining the internal temperature below 30 °C. The crude ketone **8** (0.6 Wt, 77%) was used directly in the next stage.

2-(But-3-en-2-yloxy)-1-(2-fluorophenyl)ethanone: ^1H NMR (500 MHz, CDCl_3) δ 7.93 (td, $J = 7.6, 1.8$ Hz, 1H), 7.57 – 7.49 (m, 1H), 7.24 (dt, $J = 14.6, 3.8$ Hz, 1H), 7.12 (ddd, $J = 11.0, 8.3, 0.8$ Hz, 1H), 5.21 (dd, $J = 17.3, 1.3$ Hz, 1H), 5.17 (dd, $J = 10.3, 0.8$ Hz, 1H), 4.69 (dd, $J = 18.1, 3.2$ Hz, 1H), 4.60 (dd, $J = 18.1, 3.4$ Hz, 1H); 4.00 (dq, $J = 12.9, 6.4$ Hz, 1H), 1.36 (d, $J = 6.4$ Hz, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ 195.24 (d, $J_{\text{CF}} = 5.3$ Hz), 162.03 (d, $J_{\text{CF}} = 253.9$ Hz), 139.42, 134.86 (d, $J_{\text{CF}} = 8.9$ Hz), 130.64 (d, $J_{\text{CF}} = 3.4$ Hz), 124.67 (d, $J_{\text{CF}} = 3.2$ Hz), 123.57 (d, $J_{\text{CF}} = 15.3$ Hz), 116.89, 116.45 (d, $J_{\text{CF}} = 23.7$ Hz), 77.93, 74.11 (d, $J_{\text{CF}} = 11.2$ Hz), 21.19.

Alternative procedure #2 for preparation of ketone 8:




2-(But-3-en-2-yloxy)-1-morpholinoethanone. A reactor was charged with tetrabutylammonium hydrogen sulfate (0.10 eq) and toluene (4.0 V). The mixture was cooled to 10-20 °C. While maintaining the internal temperature below 20 °C, 50 wt% aqueous sodium hydroxide (2.2 V, 9.7 eq) and then 3-buten-2-ol (0.66 V, 1.25 eq) were added. The mixture was stirred at 10-20 °C for 15 minutes. 4-(chloroacetyl)morpholine (1.0 Wt, 1.00 eq) was added while maintaining the internal temperature between 10-20 °C. After the addition, the mixture was stirred between 10-20 °C for 1h and monitored the reaction for complete consumption of 4-(chloroacetyl)morpholine (target >98% conversion). The reactor was charged with water (3.0 V) and MTBE (6.00 V) while maintaining the internal temperature between 10-20 °C. The biphasic mixture was stirred vigorously for 15 minutes, and then allowed to partition. The aqueous layer was extracted with MTBE (3.0 V). The combined organic layers were washed with water twice (1.5V each). The water wash was extracted with MTBE twice (1.6 V each). The organic layers were combined and concentrated under reduced pressure ($T < 40$ °C). Heptane (1.0 V) and MTBE (1.0 V) were added. The solution was filtered to remove particulates, rinsing with additional heptane:MTBE as necessary to clear the product. The solution was returned to the reactor and concentrated under reduced pressure to give amide **12** (77%) as oil.




2-(But-3-en-2-yloxy)-1-(2-fluorophenyl)ethanone. A reactor was charged with 1.3 M of isopropylmagnesium chloride-lithium chloride in THF (5.15 V, 1.33 eq) and cooled to 0-5 °C. 1-Bromo-2-fluorobenzene (0.726 V, 1.33 eq) was added while maintaining the internal temperature between 0-10 °C. Upon complete addition, the mixture was stirred between 0-10 °C for 1h. A solution of amide **12** (1.0 Wt, 1.0 eq.) in THF (1.0 V) was added while maintaining the internal temperature between 0-10 °C. After stirring the reaction for 3h, the reaction was monitored for complete consumption of amide **12** (>97% conversion). The reaction mixture was transferred into 1M HCl (11 V, .2.2 eq) while maintaining the internal temperature below 20 °C. The mixture was stirred for 15 minutes and then the phases were separated. The lower aqueous layers were extracted with MTBE (5.0 V). All organic layers were combined and washed with 25 wt % aqueous sodium chloride (6.0 V). The organic phase was concentrated under reduced pressure (T < 40 °C) to give ketone **8** (93%) as oil.

Example 3: Preparation of 6a-(2,3-Difluorophenyl)-4-((trityloxy)methyl)hexahydrofuro[3,4-c]isoxazole



1-Morpholino-2-(1-(trityloxy)but-3-en-2-yloxy)ethanone. To a reactor with 1-(trityloxy)but-3-en-2-ol **13** (41.6 g, 0.111 mol, 1.0 equiv.) was charged toluene (146 mL). The resulting solution was cooled to 0-5 °C and tetra-n-butylammonium hydrogen sulfate (7.52 g, 0.0222 mol, 0.20 equiv.) was charged. 4-(Chloroacetyl)morpholine (18.1g, 0.111 mol, 1.00 equiv.) was added at 0-5 °C. Sodium hydroxide (50% wt. in water; 88.6g, 1.10 mol, 10 equiv.) was cooled to 15 °C and charged to the reaction mixture with T < 10 °C. The reaction mixture was stirred for 1 hr at T < 15 °C and monitored for consumption of 1-(trityloxy)but-3-en-2-ol (target >99%). To the reaction mixture was charged 2-methoxy-2-methylpropane (146 mL) and water (146 mL) with T < 20 °C. The organics were washed with 18% aq. NaCl (73 mL) and sat. aq NH4Cl (21 mL). The organics were filtered over Celite (10.4 g) to remove particulates and rinsed with 2-methoxy-2-methylpropane (83 mL). The solvents were evaporated under vacuum T < 30 °C to afford a white solid **14** on standing (55.0 g, 98.7% yield accounting for residual solvents).


1-Morpholino-2-(1-(trityloxy)but-3-en-2-yloxy)ethanone: ^1H NMR (500 MHz, CDCl_3) δ 7.46 – 7.41 (m, 6H), 7.32 – 7.25 (m, 6H), 7.25 – 7.18 (m, 3H), 5.79 – 5.68 (m, 1H), 5.28 (d, J = 17.0 Hz, 1H), 5.26 (d, J = 10.1 Hz, 1H), 4.20 (d, J = 12.7 Hz, 1H), 4.10 (d, J = 12.7 Hz, 1H), 3.97 – 3.88 (m, 1H), 3.69 – 3.47 (m, 8H), 3.25 (dd, J = 9.9, 6.5 Hz, 1H), 3.17 (dd, J = 9.9, 4.3 Hz, 1H); ^{13}C NMR (125 MHz, CDCl_3) δ 167.84, 143.84, 134.84, 128.67, 127.78, 127.04, 119.06, 86.73, 81.07, 68.78, 66.78, 66.34, 45.94, 42.16.



10 1-(2,3-Difluorophenyl)-2-(1-(trityloxy)but-3-en-2-yloxy)ethanone.


Isopropylmagnesiumchloride-LiCl complex (1.30 M in THF, 155.0 mL, 0.0715 mol) under nitrogen was cooled to 0-5 °C, and 2,3-difluorobromobenzene (13.8 g, 0.0715 mol, 1.50 equiv.) was added while $T < 10$ °C. After 1 h at 0-5 °C, a solution of 1-morpholino-2-(1-(trityloxy)but-3-en-2-yloxy)ethanone **14** (21.8 g, 0.048 mol, 1.0 equiv.) in THF (2.0 vols) was added while $T < 10$ °C. The reaction mixture was stirred for 2.5 hours and monitored for consumption of 1-morpholino-2-(1-(trityloxy)but-3-en-2-yloxy)ethanone (target >97%). The reaction mixture was quenched by charging into cold sat. aq. NH_4Cl (110 mL) and water (33 mL) while $T < 20$ °C. 2-Methoxy-2-methylpropane (218 mL) was added and the layers were separated. The organics were washed with sat. aq. NH_4Cl (65 mL) and 18% aq. NaCl (44 mL.). The organics were concentrated under vacuum $T < 25$ °C to a light yellow oil of ketone **15** (21.6 g).

1-(2,3-Difluorophenyl)-2-(1-(trityloxy)but-3-en-2-yloxy)ethanone: ^1H NMR (500 MHz, CDCl_3) δ 7.70 – 7.64 (m, 1H), 7.47 – 7.42 (m, 6H), 7.37 – 7.30 (m, 1H), 7.30 – 7.25 (m, 6H), 7.24 – 7.19 (m, 3H), 7.19 – 7.13 (m, 1H), 5.84 – 5.71 (m, 1H), 5.30 (d, J = 17.3 Hz, 1H), 5.25 (d, J = 10.4 Hz, 1H), 4.78 (dd, J = 17.7, 3.1 Hz, 1H), 4.70 (dd, J = 17.7, 3.1 Hz, 1H), 4.03 (dd, J = 11.8, 6.5 Hz, 1H), 3.35 (dd, J = 9.8, 6.4 Hz, 1H), 3.18 (dd, J = 9.8, 4.6 Hz, 1H); ^{13}C NMR (125 MHz, CDCl_3) δ 194.15 (dd, J_{CF} = 4.9, 2.4 Hz), 150.77 (dd, J_{CF} = 250.3, 14.1 Hz), 150.29 (dd, J_{CF} = 256.4, 13.9 Hz), 143.98, 135.34, 128.75, 127.76, 126.95, 125.70 (d, J_{CF} = 12.0 Hz), 125.16 (d, J_{CF} = 3.5 Hz), 124.55 (dd, J_{CF} = 6.4, 4.2 Hz), 121.62 (d, J_{CF} = 17.6 Hz), 118.83, 86.78, 81.35, 74.98 (d, J_{CF} = 9.5 Hz), 66.71.



1-(2,3-Difluorophenyl)-2-(1-(trityloxy)but-3-en-2-yloxy)ethanone oxime. To a solution of 1-(2,3-difluorophenyl)-2-(1-(trityloxy)but-3-en-2-yloxy)ethanone **15** (21.1 g, 0.0435 mol, 1.0 equiv.) in MeOH (106 mL) was added NaOAc (5.36 g, 0.0653 mol, 1.50 equiv.) at ambient temperature. After 5 min., NH₂OH HCl (3.33g, 0.048 mol, 1.10 equiv.) was added in portions. The suspension was warmed to 35 °C and stirred for 1 hour. The reaction mixture was cooled to 17.8 °C and stirred for 15 minutes, then filtered and rinsed with ethyl acetate (84 mL). The filtrates were concentrated under vacuum (T < 30 °C). To the resulting residue was charged 2-methoxy-2-methylpropane (169 mL) and water (21.1 mL). The organics were washed with sat. aq. NaHCO₃ (53 mL) and 18% aq. NaCl (21 mL). The organics were concentrated under vacuum to afford the title compound **16** as an oil (21.0 g).

1-(2,3-Difluorophenyl)-2-(1-(trityloxy)but-3-en-2-yloxy)ethanone oxime: ¹H NMR (500 MHz, CDCl₃) δ 7.42 – 7.32 (m, 6H), 7.30 – 7.15 (m, 10H), 7.16 – 7.06 (m, 1H), 7.03 – 6.95 (m, 1H), 5.72 – 5.58 (m, 1H), 5.24 – 5.13 (m, 2H), 4.80 (d, J = 14.6 Hz, 1H), 4.73 (d, J = 14.6 Hz, 1H), 4.45 (d, J = 12.4 Hz, 0.3H), 4.35 (d, J = 12.4 Hz, 0.3H), 3.99 – 3.91 (m, 0.3H), 3.91 – 3.83 (m, 1H), 3.20 (dd, J = 9.9, 6.8 Hz, 0.3H), 3.09 (dd, J = 9.8, 6.6 Hz, 1H), 3.02 (dd, J = 9.9, 4.2 Hz, 0.3H), 2.95 (dd, J = 9.8, 4.0 Hz, 1H); ¹³C NMR (125 MHz, CDCl₃) δ 155.40, 150.59 (dd, J_{CF} = 248.1, 13.0 Hz), 148.86 (dd, J_{CF} = 252.6, 13.5 Hz), 143.96, 135.10, 128.73, 128.69, 127.72, 127.68, 126.92, 126.85, 124.96, 124.56 (d, J_{CF} = 10.6 Hz), 124.03 – 123.86 (m), 118.64 (d, J_{CF} = 15.1 Hz), 117.92 (d, J_{CF} = 17.2 Hz), 86.66, 86.50, 81.30, 80.32, 69.20, 66.54, 66.48, 62.57.




6a-(2,3-Difluorophenyl)-4-((trityloxy)methyl)hexahydrofuro[3,4-c]isoxazole. To a solution of 1-(2,3-Difluorophenyl)-2-(1-(trityloxy)but-3-en-2-yloxy)ethanone oxime **16** (20.0 g, 0.0400 mol,

1.0 equiv.) in toluene (140 mL) was charged hydroquinone (0.0882 g, 0.008 mol, 0.2 equiv.). The reaction mixture was heated to reflux (110-115 °C) and held for 13 hours. The reaction mixture was cooled to 20-25 °C and then concentrated under vacuum ($T < 45$ °C). The solvents were chased with 2-propanol (100 mL). To the crude residue was charged isopropanol (140 mL) and the mixture was heated to 65-70 °C until a clear solution is formed. The solution was cooled to 0 °C at 10 °C/hr and then held for 1 hour. The resulting suspension was filtered and the solid was rinsed with cold 2-propanol (40.0). After drying the title compound **17** was obtained as a powder (13.2 g).

6a-(2,3-Difluorophenyl)-4-((trityloxy)methyl)hexahydrofuro[3,4-c]isoxazole: ^1H NMR (500 MHz, DMSO) δ 7.42 – 7.33 (m, 8H), 7.31 (t, $J = 7.6$ Hz, 6H), 7.28 – 7.21 (m, 3H), 7.20 – 7.12 (m, 1H), 6.39 (s, 1H), 4.13 (d, $J = 8.2$ Hz, 1H), 4.03 (s, 1H), 3.97 (s, 1H), 3.89 (d, $J = 9.2$ Hz, 1H), 3.81 – 3.70 (m, 1H), 3.36 – 3.29 (m, 1H), 3.26 (dd, $J = 10.1, 6.0$ Hz, 1H), 3.15 (dd, $J = 10.0, 3.5$ Hz, 1H); ^{13}C NMR (125 MHz, DMSO) δ 150.74 (dd, $J_{\text{CF}} = 246.4, 14.4$ Hz), 148.66 (dd, $J_{\text{CF}} = 248.3, 13.5$ Hz), 144.09, 130.58, 128.67, 128.32, 127.48, 124.75, 124.45, 117.18 (d, $J_{\text{CF}} = 17.0$ Hz), 86.57, 84.91, 78.16, 76.81, 64.82, 56.46.

Example 4: Preparation of 6a-(2-fluorophenyl)-4-((trityloxy)methyl)hexahydrofuro[3,4-c]isoxazole.




6a-(2-fluorophenyl)-4-((trityloxy)methyl)hexahydrofuro[3,4-c]isoxazole. A reactor was charged with 1-(2-fluorophenyl)-2-((1-(trityloxy)but-3-en-2-yl)oxy)ethanone (61.2g, 1.0 equiv), toluene (428mL) and sodium acetate (16.1g, 1.5 equiv). Upon complete addition, the reaction mixture was stirred at ambient temperature for 5 minutes. Hydroxylamine hydrochloride (10.0g, 1.1 Eq) was added and the suspension was heated to 111 °C. After stirring for >20 hours at 111 °C, the reaction mixture was allowed to cool down to ambient temperature, and the suspension was filtered and the solids rinsed with toluene (245mL) and then THF (10 volumes). The filtrate was concentrated under reduced pressure (bath temp. ≤ 40 °C) and chased with 2-propanol (275mL). 2-Propanol (428mL) was added to the solids and the suspension was heated to 83 °C. After stirring for 1 hour at 83 °C, the mixture was allowed to cool down to ambient temperature. The slurry was cooled to 0 °C over a 1 hour period and stirred at 0 °C for 1 hour.

The suspension was filtered and the solids rinsed in two portions with cold 2-propanol (490mL) to give 6a-(2-fluorophenyl)-4-((trityloxy)methyl)hexahydrofuro[3,4-c]isoxazole (42.3g) as a solid.

¹H NMR (400 MHz, CDCl₃) δ 7.62 (t, 1H), 7.41 (d, *J* = 7.2 Hz, 6H), 7.31 – 7.21 (m, 10H), 7.12 (m, 1H), 7.03 (dd, *J* = 12.0, 8.0 Hz, 1H), 4.29-4.04 (m, 3H), 3.91 (dd, *J* = 8.8, 1.2 Hz, 1H), 3.88 (m, 1H), 3.42 (dd, *J* = 9.2, 6.8 Hz, 1H), 3.29 (m, 1H), 3.22 (dd, *J* = 10.0, 4.8 Hz, 1H).

B. Preparation of Tetrahydrofurans of Formula II




10

4-Amino-4-(2-fluorophenyl)-2-(trifluoromethyl)tetrahydrofuran-3-yl)methanol. Zinc (389.2 g, 5.95 mol) was placed in a reaction vessel, and water was added (893 mL). Acetic acid was added while maintaining the temperature below 10 °C. After 15 min, 6a-(2-fluorophenyl)-4-(trifluoromethyl)hexahydrofuro[3,4-c]isoxazole **5** (550.0 g, 1.98 mol) was added as a solution in THF (665 mL). The reaction mixture was stirred over 16 h at rt. Methylene chloride (1.89 L) was added, followed by 28% aq. NH₄OH (552 mL) while the temperature was kept below 30 °C. The mixture was stirred for 30 min, and then filtered over Celite (80 g) rinsing with methylene chloride (378 mL). The aq. layer was extracted with methylene chloride (1.89 L). The organics were combined, washed with sat. aq. NaCl (1.0 L) and concentrated under vacuum to afford an aminoalcohol **18** (502 g, 90.6%). The crude residue was used in the following step without additional purification.

20


HRMS Calculated for C₁₂H₁₃F₄NO₂ [M+H]⁺ 280.0961; found 280.0972.



25

4-Amino-4-(2-fluorophenyl)-2-methyltetrahydrofuran-3-yl)methanol. A reactor was charged with tetrahydroisoxazole **10** (1.0 Wt, 1.0 eq) and glacial acetic acid (4.1 Wt, 6.0 V). The mixture was stirred at 20 °C, then in 3 portions, gradually charged zinc dust (1.17 Wt, 4.0 eq.) keeping the reaction temperature below 60 °C. Once the exotherm subsided, the mixture was stirred at 40 °C for 2 h and then monitored for complete consumption of tetrahydroisoxazole **10** (>99% conversion). The mixture was filtered to remove residual zinc. The reactor and solids were rinsed with ethyl acetate (4.5 Wt, 5 V). The filtrate was returned to the reactor and concentrated under reduced pressure ($T < 40\text{ }^{\circ}\text{C}$). Toluene (1.73 wt, 2.0 V) was charged and the mixture was concentrated under reduced pressure. Toluene (1.73 wt, 2.0 V) was charged and the mixture was concentrated under reduced pressure. Dichloromethane (6.81 wt, 5 V) and water (5.0 wt, 5.0 vol) were charged. Add 28% ammonium hydroxide until the pH reaches 8-9. The mixture was stirred for 15 minutes, then the phases were allowed to separate and the lower organic layer removed. Charged dichloromethane (5.4 wt, 4.0 V). Stirred the mixture for 15 minutes then separated the phases. The organic phases were combined then washed with 25% aqueous sodium chloride (2.0 Wt, 2.0V). The organic phase was concentrated under reduced pressure ($T < 40\text{ }^{\circ}\text{C}$). Anhydrous THF (2.0 Wt) was added, and the solution was filtered to remove particulates, rinsing with additional anhydrous THF as necessary to clear the product. The THF solution was returned to the reactor and concentrated under reduced pressure to give aminoalcohol **19** (0.93 Wt, 94.0%) as an oil.

4-Amino-4-(2-fluorophenyl)-2-methyltetrahydrofuran-3-yl)methanol: ^1H NMR (500 MHz, DMSO) δ 7.63 (td, $J = 8.6, 1.7\text{ Hz}$, 1H), 7.32 – 7.24 (m, 1H), 7.20 – 7.08 (m, 2H), 4.43 (s, 1H), 4.07 (dd, $J = 8.7, 1.3\text{ Hz}$, 1H), 4.02 (dq, $J = 8.5, 6.1\text{ Hz}$, 1H), 3.68 – 3.57 (m, 2H), 3.55 – 3.46 (m, 1H), 2.23 (dd, $J = 14.4, 6.7\text{ Hz}$, 1H), 1.24 (d, $J = 6.1\text{ Hz}$, 3H); ^{13}C NMR (125 MHz, DMSO) δ 160.69 (d, $J_{\text{CF}} = 244.7\text{ Hz}$), 133.57 (d, $J_{\text{CF}} = 11.3\text{ Hz}$), 128.95 (d, $J_{\text{CF}} = 8.8\text{ Hz}$), 128.81 (d, $J_{\text{CF}} = 4.8\text{ Hz}$), 124.33 (d, $J_{\text{CF}} = 3.2\text{ Hz}$), 116.32 (d, $J_{\text{CF}} = 23.5\text{ Hz}$), 79.58 (d, $J_{\text{CF}} = 4.9\text{ Hz}$), 78.52, 63.99 (d, $J_{\text{CF}} = 3.5\text{ Hz}$), 59.39, 56.91 (d, $J_{\text{CF}} = 2.2\text{ Hz}$), 21.37.




4-Amino-4-(2,3-difluorophenyl)-2-((trityloxy)methyl)tetrahydrofuran-3-yl)methanol. To 6a-(2,3-difluorophenyl)-4-((trityloxy)methyl)hexahydrofuro[3,4-c]isoxazole **17** (87.5 g, 0.175 mol,

1.0 equiv.) was added acetic acid (385 mL). The suspension was cooled to 15 °C, and zinc (88.4g, 1.35 mol, 7.71 equiv.) was added in small portions over 2-3 min. The reaction was allowed to warm up to 18 °C over 3-4 hours and stirred for 9 h. The reaction was filtered over celite (40 g) and rinsed with toluene (210 mL). The collected filtrate was concentrated under reduced pressure with bath temperature below 35 °C and chased with three portions of toluene (3 x 130 mL). The resulting oil was dissolved in CH₂Cl₂ (320 mL). Water (195 mL) was added followed by 28% NH₄OH (43.4 mL). The organic layer was separated and the remaining aqueous phase was extracted with CH₂Cl₂ (130 mL). The combined organic phases were washed with 27% aq. NaCl (87.5 mL). The solution was concentrated under reduced pressure, dissolved in THF (270 mL) and filtered through celite. The solids were rinsed with THF (130 mL) and the combined filtrate was concentrated under vacuum to produce title compound **20** (69.1 g) as a foam.

4-Amino-4-(2,3-difluorophenyl)-2-((trityloxy)methyl)tetrahydrofuran-3-yl)methanol: ¹H NMR (500 MHz, DMSO) δ 7.47 (t, *J* = 7.5 Hz, 1H), 7.41 (d, *J* = 7.5 Hz, 6H), 7.34 – 7.27 (m, 7H), 7.28 – 7.21 (m, 3H), 7.19 – 7.10 (m, 1H), 4.14 (d, *J* = 8.7 Hz, 1H), 4.15 – 4.09 (m, 1H), 3.77 (dd, *J* = 8.6, 2.6 Hz, 1H), 3.59 (dd, *J* = 11.1, 6.4 Hz, 1H), 3.46 (dd, *J* = 11.1, 6.6 Hz, 1H), 3.18 (dd, *J* = 9.9, 3.0 Hz, 1H), 3.07 (dd, *J* = 9.9, 5.8 Hz, 1H), 2.62 (dd, *J* = 14.6, 6.6 Hz, 1H); ¹³C NMR (125 MHz, DMSO) δ 150.68 (dd, *J*_{CF} = 244.4, 13.9 Hz), 148.46 (dd, *J*_{CF} = 246.7, 13.0 Hz), 144.36, 135.44 (d, *J*_{CF} = 8.1 Hz), 128.76, 128.23, 127.37, 124.31, 124.23, 116.12 (d, *J*_{CF} = 17.1 Hz), 86.29, 81.48, 78.92 (d, *J*_{CF} = 4.3 Hz), 66.03, 63.93, 59.49, 51.27

HRMS Calculated for C₃₁H₂₉F₂NO₃ [M+Na]⁺ 524.2013; found 524.2039.

Alternative procedure for the preparation of compounds of Formula II




4-Amino-4-(2-fluorophenyl)-2-methyltetrahydrofuran-3-yl)methanol (19)

A reactor was charged with ketone **8** (1.0 Wt, 1.0 eq), and water (10 Wt, 10 V), hydroxylamine hydrochloride (0.350 Wt, 1.05 eq) and sodium acetate (0.433 Wt, 1.10 eq), were added. The resulting mixture was stirred and warmed to between 101-102 °C for at least 40 h and monitored for complete conversion to tetrahydrooxazole **10** (>95%). The mixture was cooled to below 25°C and then charged with THF (1.8 wt, 2.0V) and acetic acid (0.86 Wt, 0.82 V, 3.0 eq). Zinc

dust (0.942 Wt, 3.00 eq) was added while maintaining the internal temperature below 45 °C. The resulting mixture was stirred between 35-45 °C for 1 h and the reaction was monitored for complete conversion to aminoalcohol **19** (>99%). The mixture was cooled to below 20 °C and dichloromethane (2.6 Wt, 2.0 V) was added. Aqueous ammonium hydroxide (28%; 1.3 wt, 1.5 V, 2.2 eq) was added until the pH reached 10-12. The resulting mixture was stirred for 15 minutes and the solids were then filtered, rinsing with dichloromethane (4.6 Wt, 3.5V). After allowing the filtrate layers to partition, the lower organic layer was set aside. The upper aqueous layer was extracted twice with dichloromethane (2.6 wt, 2.0 V each). The combined organic layers were concentrated under reduced pressure (jacket T <45 °C) to give aminoalcohol **19** (0.88 wt, 81%).

C. Diastereomeric Resolution of Tetrahydrofurans of Formula II



((2S,3R,4S)-4-Amino-4-(2-fluorophenyl)-2-(trifluoromethyl)tetrahydrofuran-3-yl)methanol (2S,3S)-2,3-bis(benzoyloxy)succinate. To a solution of 4-amino-4-(2-fluorophenyl)-2-(trifluoromethyl)tetrahydrofuran-3-yl)methanol **18** (0.502 kg, 1.798 mol) in ethanol (4.865 L) was added dibenzoyl-D-tartaric acid (0.642 kg, 1.798 mol). The resulting suspension was heated to 67 °C. Water (94.0 mL, 5.2 mol) was added over 15 min while maintaining temperature >66 °C. The resulting solution was cooled to 45 °C while precipitation occurred. The slurry was reheated to 60 °C, and then cooled to ambient temperature at 5 °C/hour. The slurry was filtered, and the solid was rinsed with premixed and cooled solution of ethanol (950 mL) and water (20 mL). The solid aminoalcohol dibenzoyltartaric acid salt **21** was dried until constant weight under vacuum (370 g, 97.6% ee).

((2S,3R,4S)-4-Amino-4-(2-fluorophenyl)-2-(trifluoromethyl)tetrahydrofuran-3-yl)methanol (2S,3S)-2,3-bis(benzoyloxy)succinate: ¹H NMR (500 MHz, CD₃OD) δ 8.13 (d, *J* = 7.2 Hz, 4H), 7.66 – 7.58 (m, 3H), 7.54 – 7.45 (m, 5H), 7.36 – 7.20 (m, 2H), 5.92 (s, 2H), 4.79 – 4.66 (m, 1H), 4.40 – 4.28 (m, 1H), 4.04 (dd, *J* = 12.1, 3.4 Hz, 1H), 3.92 (dd, *J* = 12.1, 5.4 Hz, 1H), 3.30 – 3.24 (m, 1H); ¹³C NMR (125 MHz, DMSO) δ 169.61, 165.81, 160.23 (d, *J* = 246.1

Hz), 133.00, 131.34 (d, $J = 9.1$ Hz), 129.65, 129.55, 128.08, 127.97 (d, $J = 3.5$ Hz), 124.95 (d, $J = 3.3$ Hz), 116.56 (d, $J = 23.5$ Hz), 77.48 (q, $J_{\text{CF}} = 31.0$ Hz), 76.33, 73.20, 65.61 (d, $J = 3.1$ Hz), 57.11.

HRMS Calculated for $\text{C}_{12}\text{H}_{13}\text{F}_4\text{NO}_2$ $[\text{M}+\text{H}]^+$ 280.0961; found 280.0967 (for amino alcohol).

The absolute stereochemistry of the amino alcohol **21** was assigned by comparison with a sample prepared starting from enantioenriched (*S*)-2-(trifluoromethyl)oxirane.

Chiral HPLC parameters:

10 Equipment, Reagents, and Mobile Phase:

Equipment:

HPLC column:	Chiralcel OD, 4.6 x 250 mm, 10 μm , Daicel Chemical Industries, Ltd., catalog no. 14025.
Solvent Delivery System:	Agilent 1100 HPLC ternary pump, low pressure mixing with in-line degasser, or equivalent.
Autosampler:	Agilent 1100 autosampler, 0.1 to 100 μL range, or equivalent.
Detector:	Agilent 1100 variable wavelength detector or equivalent.
Chromatographic Software:	Agilent ChemStation software version A.09.03 or higher for HPLC, Waters Empower 2 Build 2154 or equivalent.
Volumetric Glassware:	Class A.
Volumetric pipette:	Class A.
Pipettor:	Calibrated Eppendorf adjustable volume, or equivalent.
Balance:	Analytical balance, capable of weighing ± 0.1 mg.

Reagents:

Heptane:	HPLC grade, Baker (catalog no. 9177-03) or equivalent.
2-Propanol:	HPLC grade, Baker (catalog no. 9095-03) or equivalent.
Triethylamine:	$\geq 99\%$, Sigma-Aldrich (catalog no. T0886) or equivalent.

15 Mobile Phase:

Add 70 mL 2-propanol and 930 mL heptane (measured separately with a 100 mL and 1000-mL graduated cylinders) and 1.0 mL triethylamine (measured with volumetric glass pipette) to an appropriate flask and mix. Degas in-line during use.

Diluting Solution: 2-Propanol

HPLC Parameters:

HPLC column:	Chiralcel OD, 4.6 x 250 mm, 10 µm, Daicel Chemical Industries, Ltd., catalog no. 14025.
Temperature:	35 °C
Flow rate*:	0.8 mL/min
Gradient:	NA
Injection Volume:	5 µL
Detection:	262 nm UV
Data acquisition time:	30 min
Total run time:	30 min
Column Maximum Pressure:	35 Bar
Needle Wash:	2-propanol
*Flow rate may be adjusted ± 0.2 ml/min to obtain specified retention times.	


Retention Times for Analytes and Impurities:

Compound Peak	Retention Time (Relative Retention Time, RRT)
	20.6 min ± 10% (RRT 1.00)
 (Enantiomer)	19.2 min (RRT 0.93)

5

A typical chromatogram is presented in **Figure 1A**.

10




((2R,3R,4S)-4-Amino-4-(2-fluorophenyl)-2-methyltetrahydrofuran-3-yl)methanol

(2S,3S)-2,3-bis(benzoyloxy)succinate. A reactor was charged with aminoalcohol **19** (1.0 Wt, 1.0 eq) and ethanol (5.5 Wt, 7.0 V). The mixture was heated to internal temperature 65-70 °C. To the mixture is added a solution of dibenzoyl-D-tartaric acid (1.0 eq, 1.59 wt) in ethanol (1.97 Wt, 2.50 V) and water (0.25 Wt, 0.25V) over 30-60 minutes while maintaining the internal temperature above 60 °C. The vessel containing the dibenzoyltartaric acid was rinsed with minimum amount ethanol. The mixture was then cooled to 40 °C at a rate 3 °C/hour. After crystallization the mixture was cooled to 10 °C (at 10 °C/hour) and stirred for at least 2 hours at 10 °C. The crystals were then filtered and rinsed with pre-chilled ethanol (1.4 Wt, 1.8 V) and dried under vacuum to give amino alcohol dibenzoyltartaric acid salt **22** (0.86 Wt, 33.0%) as solid.

((2R,3R,4S)-4-Amino-4-(2-fluorophenyl)-2-methyltetrahydrofuran-3-yl)methanol

(2S,3S)-2,3-bis(benzoyloxy)succinate: ¹H NMR (500 MHz, CD₃OD) δ 8.14 – 8.10 (m, 4H), 7.60 (t, *J* = 7.4 Hz, 2H), 7.55 (td, *J* = 8.1, 1.3 Hz, 1H), 7.47 (t, *J* = 7.8 Hz, 4H), 7.46 – 7.42 (m, 1H), 7.30 – 7.19 (m, 2H), 5.90 (s, 2H), 4.39 – 4.26 (m, 2H), 4.15 (dd, *J* = 10.4, 2.4 Hz, 1H), 3.90 (ddd, *J* = 28.3, 11.9, 5.0 Hz, 2H), 2.52 (dt, *J* = 8.4, 5.0 Hz, 1H), 1.31 (d, *J* = 6.1 Hz, 3H); ¹³C NMR (126 MHz, CD₃OD) δ 169.96, 165.90, 160.35 (d, *J* = 245.9 Hz), 132.95, 131.05 (d, *J*_{CF} = 9.1 Hz), 129.75, 129.57, 128.06, 127.57 (d, *J*_{CF} = 3.5 Hz), 124.82 (d, *J*_{CF} = 3.2 Hz), 124.24 (d, *J*_{CF} = 11.5 Hz), 116.45 (d, *J*_{CF} = 23.1 Hz), 76.34, 74.32 (d, *J*_{CF} = 5.4 Hz), 73.50, 65.60 (d, *J*_{CF} = 2.8 Hz), 57.43, 54.88, 19.00.



23

(2S,3R,4S)-4-Amino-4-(2,3-difluorophenyl)-2-((trityloxy)methyl)tetrahydrofuran-3-yl)methanol (-)-dibenzoyl-L-tartaric acid. Racemic mixture 6a-(2,3-difluorophenyl)-4-

((trityloxy)methyl)hexahydrofuro [3,4-c]isoxazole **20** (69.1 g, 0.138 mol, 1.0 equiv.) and (-)-dibenzoyl-L-tartaric acid (49.4 g, 0.138 mol, 1.0 equiv) were dissolved in acetone (346 mL). The mixture was then heated to 61.9°C over 15 min. Toluene (415 mL) was added. The solution was then allowed to cool to room temperature. Then the reaction mixture was cooled to -5 °C and stirred for another hour at this temperature. The suspension was filtered and washed with a solution of toluene/acetone (V/V 6/5, 138 mL, cooled to 0 °C). The solid was dried under reduced pressure, and the title compound **23** (44.5 g) was isolated.

(2S,3R,4S)-4-Amino-4-(2,3-difluorophenyl)-2-((trityloxy)methyl)tetrahydrofuran-3-yl)methanol (-)dibenzoyl-L-tartaric acid: ¹H NMR (500 MHz, DMSO) δ 8.00 – 7.95 (m, 4H), 7.66 (t, J = 7.4 Hz, 2H), 7.53 (t, J = 7.8 Hz, 4H), 7.47 – 7.39 (m, 2H), 7.40 – 7.35 (m, 6H), 7.35 – 7.28 (m, 6H), 7.28 – 7.23 (m, 3H), 7.23 – 7.15 (m, 1H), 5.75 (s, 1H), 4.19 (d, J = 9.5 Hz, 2H), 4.02 (dd, J = 9.5, 2.1 Hz, 1H), 3.61 (dd, J = 11.4, 6.4 Hz, 1H), 3.52 (dd, J = 11.4, 6.2 Hz, 1H), 3.21 – 3.14 (m, 1H), 3.12 – 3.05 (m, 1H), 2.77 (dd, J = 14.3, 6.4 Hz, 1H); ¹³C NMR (125 MHz, DMSO) δ 168.07, 165.23, 150.68 (dd, J_{CF} = 245.2, 13.4 Hz), 148.27 (dd, J_{CF} = 248.0, 13.6 Hz), 144.18, 134.09, 129.75, 129.60, 129.24, 128.70, 128.28, 127.45, 124.88, 124.20, 117.40 (d, J_{CF} = 16.8 Hz), 86.40, 80.73, 76.73, 72.22, 65.42, 64.58, 58.63, 50.51.

HRMS Calculated for C₃₁H₂₉F₂NO₃ [M+Na]⁺ 524.2013; found 524.2047.

Chiral HPLC parameters:

Equipment, Reagents, and Mobile Phase:

Equipment:

HPLC column:	Chiralpak AD, 4.6 x 250 mm, 10 μm, Daicel Chemical Industries, Ltd., catalog no. 19025.
Solvent Delivery System:	Agilent 1100 HPLC ternary pump, low pressure mixing with an in-line degasser, or equivalent.
Autosampler:	Agilent 1100 autosampler, 0.1 to 100 μL range, or equivalent.
Detector:	Agilent 1100 variable wavelength detector or equivalent.
Chromatographic Software:	Agilent ChemStation software version A.09.03 or higher for HPLC, Waters Empower 2 Build 2154 or equivalent.
Volumetric Glassware:	Class A.
Volumetric pipette:	Class A.
Pipettor:	Calibrated Eppendorf adjustable volume, or equivalent.
Balance:	Analytical balance, capable of weighing ± 0.1 mg.

Reagents:

Heptane:	HPLC grade, Baker (catalog no. 9177-03) or equivalent.
2-Propanol:	HPLC grade, Baker (catalog no. 9095-03) or equivalent.
Triethylamine:	≥ 99%, Sigma-Aldrich (catalog no. T0886) or equivalent.


Mobile Phase:


- 5 Add 100 mL 2-propanol and 900 mL heptane (measured separately with a 100 mL and 1000-mL graduated cylinders) and 0.5 mL triethylamine (measured with volumetric glass pipette) to an appropriate flask and mix. Degas in-line during use.

Diluting Solution: 2-Propanol10 **HPLC Parameters:**

HPLC column:	Chiralpak AD, 4.6 x 250 mm, 10 μm, Daicel Chemical Industries, Ltd., catalog no. 19025.
Temperature:	35 °C
Flow rate*:	0.8 mL/min
Gradient:	NA
Injection volume:	5 μL
Detection:	260 nm UV
Data acquisition time:	40 min
Total run time:	40 min
Column Maximum Pressure:	35 Bar
Needle Wash:	2-propanol
*Flow rate may be adjusted ± 0.2 ml/min to obtain specified retention times.	

Retention Times for Analytes and Impurities:


Compound Peak	Retention Time (Relative Retention Time, RRT)
	15.5 min ± 10% (RRT 1.00)

Compound Peak	Retention Time (Relative Retention Time, RRT)
 (Enantiomer)	21.5 min (RRT 1.39)

A typical chromatogram is presented in **Figure 1B**.

5

D. Synthesis of Carbamothioyl Benzamides of Formula III




- 10 ***N*-((3*S*,4*R*,5*S*)-3-(2-fluorophenyl)-4-(hydroxymethyl)-5-(trifluoromethyl)-tetrahydrofuran-3-ylcarbamothioyl)benzamide.** To chiral salt **21** (0.361 kg, 0.556 mol) was added ethyl acetate (1.08 L) and the suspension was cooled to -3 °C. 1.0 N aq. NaOH (1.30 L) was added over 20 minutes while maintaining $T < 5$ °C. After 5 minutes, benzoyl isothiocyanate (80.0 mL, 594 mmol) was added over 8 minutes while maintaining $T < 5$ °C. After 1 hr, ethyl acetate (722 mL) was charged. The aq. layer was removed, and the organics were washed with sat. aq. NaHCO₃ (361 mL) and sat. aq. NaCl (361 mL). The organics were filtered over celite (90 g) and rinsed with ethyl acetate (360 mL). The organics were concentrated under vacuum to afford a residue which was re-dissolved into CH₂Cl₂ (1.1 L) and concentrated to afford the title compound **24** as yellow foam (261 g, 99% yield accounting for residual solvents) which was used in the following
- 15
- 20 step.

N-((3*S*,4*R*,5*S*)-3-(2-fluorophenyl)-4-(hydroxymethyl)-5-(trifluoromethyl)-tetrahydrofuran-3-ylcarbamothioyl)benzamide: ¹H NMR (500 MHz, DMSO) δ 12.04 (s, 2H), 11.20 (s, 2H), 7.95 (d, $J = 7.4$ Hz, 2H), 7.69 – 7.60 (m, 1H), 7.56 – 7.42 (m, 2H), 7.37 – 7.28 (m,


1H), 7.24 – 7.12 (m, 2H). 5.59 (t, $J = 4.5$ Hz, 1H), 5.03 (d, $J = 9.7$ Hz, 1H), 4.92 (d, $J = 9.7$ Hz, 1H), 4.75 – 4.63 (m, 1H). 3.92 – 3.74 (m, 2H), 2.77 – 2.66 (m, 1H); ^{13}C NMR (125 MHz, DMSO) δ 179.98, 167.85, 159.75 (d, $J_{\text{CF}} = 245.0$ Hz). 133.44, 132.58, 129.88, 129.81, 129.04, 128.85, 126.31 (d, $J_{\text{CF}} = 9.8$ Hz), 124.36, 116.83 (d, $J_{\text{CF}} = 23.4$ Hz), 76.11 (q, $J_{\text{CF}} = 31.0$ Hz).
 5 74.37 (d, $J_{\text{CF}} = 6.1$ Hz), 68.77 (d, $J_{\text{CF}} = 3.4$ Hz), 57.03, 52.23.

HRMS Calculated for $\text{C}_{20}\text{H}_{18}\text{F}_4\text{N}_2\text{O}_3\text{S}$ $[\text{M}+\text{H}]^+$ 441.0896; found 441.0818.



N-((3S,4R,5R)-3-(2-fluorophenyl)-4-(hydroxymethyl)-5-methyltetrahydrofuran-3-ylcarbamothioyl)benzamide. A reactor was charged with **22** (1 Wt, 1 eq) and ethyl acetate (2.7 Wt, 3.0 V). The mixture was cooled to 0 to 10 °C. 1.00 M aqueous NaOH (4.2 Wt, 4.0 V, 2.3 eq) was added. Benzoyl isothiocyanate (0.314 Wt, 0.259 V, 1.10 eq) was added with vigorous stirring while maintaining T between 0 to 10 °C. After completed addition, continued stirring at 0–5 °C over 2h and monitored for complete consumption of **22** (target >98% conversion). The
 10 reaction mixture was filtered and the filter cake was rinsed with: 1) water (2.0 Wt, 2.0 V); 2) a mixture of heptane (0.91 Wt, 1.3V) and Ethyl acetate (1.2 Wt, 1.3 V). The solid was dried under reduced pressure (T <40 °C) to give **25** (0.57 Wt, 85.4% yield).

N-((3S,4R,5R)-3-(2-fluorophenyl)-4-(hydroxymethyl)-5-methyltetrahydrofuran-3-ylcarbamothioyl)benzamide: ^1H NMR (500 MHz, CDCl_3) δ 11.78 (s, 1H), 8.88 (s, 1H), 7.88 – 7.82 (m, 2H), 7.67 (td, $J = 8.1, 1.6$ Hz, 1H), 7.65 – 7.59 (m, 1H), 7.50 (dd, $J = 10.8, 4.8$ Hz, 2H), 7.30 – 7.23 (m, 1H), 7.15 (td, $J = 7.8, 1.2$ Hz, 1H), 7.01 (ddd, $J = 12.3, 8.2, 1.1$ Hz, 1H), 4.73 (d, $J = 10.1$ Hz, 1H), 4.42 (dd, $J = 10.1, 1.7$ Hz, 1H), 4.10 – 4.03 (m, 1H), 4.03 – 3.91 (m, 2H), 2.80 (t, $J = 5.3$ Hz, 1H), 2.63 – 2.52 (m, 1H), 1.35 (d, $J = 6.1$ Hz, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ 179.42, 166.60, 160.35 (d, $J_{\text{CF}} = 247.4$ Hz), 133.60, 131.81, 129.27 (d, $J_{\text{CF}} = 8.9$ Hz), 129.20 (d, $J_{\text{CF}} = 3.7$ Hz), 129.15, 127.55, 127.39 (d, $J_{\text{CF}} = 10.0$ Hz), 123.96 (d, $J_{\text{CF}} = 3.3$ Hz), 116.18 (d, $J_{\text{CF}} = 23.0$ Hz), 76.90 (d, $J_{\text{CF}} = 2.8$ Hz), 76.27, 69.26, 59.49, 58.03, 20.04.
 20
 25



26

N-(((3S,4R,5S)-3-(2,3-difluorophenyl)-4-(hydroxymethyl)-5-

((trityloxy)methyl)tetrahydrofuran-3-yl)carbamothioyl)benzamide. A suspension of chiral

5 salt **23** (44.21 g, 0.051 mol, 1.0 equiv.) in ethyl acetate (133 mL) was cooled down to 1.7 °C. Aq. NaOH (1.0 M, 121 mL, 0.12 mol, 2.4 equiv.) was added while maintaining temperature below 5 °C. The reaction was stirred for 5 min then benzyl isothiocyanate (9.43 g, 0.058 mol, 1.12 equiv.) was added over 5 min. After 1.5 h, ethyl acetate (133 mL) was added. The organic phase was isolated and successively washed with saturated aq. NaHCO₃ (130 mL) and 18% aq. NaCl (44 mL). The combined aqueous phases were extracted with EtOAc (66 mL) and CH₂Cl₂ (30 mL). All the organic phase were combined, filtered and concentrated under reduced pressure. The title compound **26** (34.2 g) was isolated as a foam after chasing the residual solvent with CH₂Cl₂ (88 mL).


N-(((3S,4R,5S)-3-(2,3-difluorophenyl)-4-(hydroxymethyl)-5-

15 ((trityloxy)methyl)tetrahydrofuran-3-yl)carbamothioyl)benzamide: ¹H NMR (500 MHz, DMSO)

δ 11.97 (br s, 1H), 11.20 (br s, 1H), 7.94 (d, *J* = 7.4 Hz, 2H), 7.63 (t, *J* = 7.4 Hz, 1H), 7.51 (t, *J* = 7.8 Hz, 2H), 7.38 – 7.21 (m, 17H), 7.13 (dd, *J* = 13.4, 7.9 Hz, 1H), 5.13 (t, *J* = 4.4 Hz, 1H), 5.07 (d, *J* = 9.7 Hz, 1H), 4.47 (d, *J* = 9.8 Hz, 1H), 4.24 – 4.16 (m, 1H), 3.68 (t, *J* = 4.6 Hz, 2H), 3.19 (dd, *J* = 10.1, 2.9 Hz, 1H), 3.01 (dd, *J* = 10.1, 4.8 Hz, 1H), 2.63 – 2.52 (m, 1H); ¹³C NMR (125

20 MHz, DMSO) δ 180.06, 168.14, 150.75 (dd, *J*_{CF} = 244.6, 13.2 Hz), 147.93 (dd, *J*_{CF} = 247.5, 13.5 Hz), 144.15, 133.46, 132.55, 131.90 (d, *J*_{CF} = 6.4 Hz), 129.04, 128.86, 128.67, 128.28, 127.43, 124.41, 124.25, 116.12 (d, *J*_{CF} = 17.0 Hz), 86.27, 79.75, 75.03, 68.04, 64.63, 58.16, 53.35.

HRMS Calculated for C₃₉H₃₄F₂N₂O₄S [M+H]⁺ 663.2129; found 663.2200.



((2S,3R,4S)-4-Amino-4-(2-fluorophenyl)-2-((trityloxy)methyl)tetrahydrofuran-3-yl)methanol (2R,3R)-2,3-bis(benzoyloxy)succinate. A reactor was charged with zinc dust (<10 micron, 3.39 g 5.0 equiv.), **6a-(2-fluorophenyl)-4-((trityloxy)methyl)hexahydrofuro[3,4-c]isoxazole** (5.0 g, 1.0 equiv.) and THF (30.0mL). To the stirred suspension was charged acetic acid (2.97 mL, 5.0 equiv.). After addition, the reaction temperature was maintained between 20-30 °C for 1 hour. Then the reaction mixture was warmed to 35-40 °C and the temperature was maintained while monitoring for consumption of starting material. Upon completion, the reaction mixture was cooled to 20-25 °C and charged 28% aq NH₄OH (12.5 mL) while maintaining temperature <40 °C. The suspension was cooled to 20-25 °C and filtered over Celite (0.5 Wts). The solids were rinsed with MTBE (35.0 mL). The filtrate was transferred to a separatory funnel. The aqueous layer was removed, and the organic layer was washed with water (10.0 mL). The organics were concentrated under vacuum (bath temperature < 40 °C) to afford racemic aminoalcohol as a foam (95-105% crude yield). To the racemic amino alcohol was charged (-)-dibenzoyl-L-tartaric acid (3.74g, 1.0 equiv). To the mixture was charged water (0.76 mL, 4.0 equiv.) and then acetone (25.3 mL). The mixture was heated to 56-60 °C and then charged toluene (30.3 mL) while maintaining temperature above 50 °C. Upon completed addition, the temperature was held at 56-60 °C for at least 30 minutes and then cooled to 17-22 °C at 10 °C / hour. The suspension was stirred at 17-22 °C for at least two hours. The solids were filtered and washed with a solution of toluene/acetone in two portions (6/5 V/V, 4.0 volumes). The solid was dried under reduced pressure to obtain **((2S,3R,4S)-4-Amino-4-(2-fluorophenyl)-2-((trityloxy)methyl)tetrahydrofuran-3-yl)methanol (2R,3R)-2,3-bis(benzoyloxy) succinate** (3.4 g 38% yield) as a white powder. Typical ee% 96-97%.

¹H NMR (400 MHz, *d*₆-DMSO) δ 7.92 (d, *J* = 7.6Hz, 4H), 7.61-7.54 (m, 2H), 7.46 (t, *J* = 7.6Hz, 4H), 7.33-7.09 (m, 19H), 5.66 (s, 2H), 4.20 (s, 1H), 4.12 (d, *J* = 9.6 Hz, 1H), 4.07 (d, *J* = 9.6 Hz, 1H), 3.57 (dd, *J* = 11.2, 6.0, 1H), 3.48 (dd, *J* = 10.8, 6.4, 1H), 3.12 (d, *J* = 8.0 Hz, 1H), 3.03 (dd, *J* = 10.0, 5.2Hz, 1H), 2.77 (app q, *J* = 7.2Hz, 1H).

Analytical method for enantiomeric excess assessment:



HPLC Parameters:

Mobile Phase: Add 120 mL 2-propanol and 880 mL heptane (measured separately with a 100 mL and 1000-mL graduated cylinders) and 0.5 mL triethylamine (measured with volumetric glass pipette) to an appropriate flask and mix. Degas in-line during use.


HPLC column:	Chiralpak AD, 4.6 x 250 mm, 10 μm, Daicel Chemical Industries, Ltd., catalog no. 19025.
--------------	---

Temperature:	35 °C
Flow rate*:	0.8 mL/min
Gradient:	NA
Injection volume:	3 µL
Detection:	260 nm UV
Data acquisition time:	30 min
Total run time:	30 min
Column Maximum Pressure:	35 Bar
Needle Wash:	2-propanol
*Flow rate may be adjusted $\pm$ 0.2 ml/min to obtain specified retention times.	

Retention Times for Analytes and Impurities:


Compound Peak	Retention Time (Relative Retention Time, RRT)	Structure
((2S,3R,4S)-4-amino-4-(2-fluorophenyl)-2-((trityloxy)methyl)tetrahydrofuran-3-yl)methanol dibenzoate diformate	12.6 min $\pm$ 10% (RRT 1.00)	
Enantiomer: ((2R,3S,4R)-4-amino-4-(2-fluorophenyl)-2-((trityloxy)methyl)tetrahydrofuran-3-yl)methanol	19.6 min (RRT 1.56)	

An x-ray crystal structure of ((2S,3R,4S)-4-Amino-4-(2-fluorophenyl)-2-
5 ((trityloxy)methyl)tetrahydrofuran-3-yl)methanol (2R,3R)-2,3-bis(benzoyloxy)succinate:



was obtained and is presented in **Figure 2**.

E. Synthesis of Thiazyl Benzamides of Formula IV




5

N-((4a*S*,5*S*,7a*S*)-7a-(2-fluorophenyl)-5-(trifluoromethyl)-4a,5,7,7a-tetrahydro-4H-furo[3,4-d][1,3]thiazin-2-yl)benzamide. A solution of *N*-((3*S*,4*R*,5*S*)-3-(2-fluorophenyl)-4-(hydroxymethyl)-5-(trifluoromethyl)-tetrahydrofuran-3-ylcarbamothioyl)benzamide **24** (258.3 g, 583.8 mmol) in CH₂Cl₂ (1.55 L) was cooled to -19.4 °C. Pyridine (118 mL, 1.46 mol) was added while maintaining temperature at -20 °C, and then the reaction mixture was cooled to -24 °C. In another nitrogen purged vessel, CH₂Cl₂ (258 mL) was added followed by trifluoromethanesulfonic anhydride (108.0 mL, 642.2 mmol). The resulting solution was added to the reaction mixture over 30 min, while maintaining temperature below -19.7 °C. upon completed addition, the reaction mixture was stirred for 30 min at -20 °C to -15 °C, and then warmed to -11 °C over 20 min. Saturated aq. NH₄Cl (646 mL) and water (390 mL) was added. The mixture was warmed to ambient temperature and the aq. layer was removed. The organics were washed with premixed saturated aq. NH₄Cl (646 mL) and water (390 mL). The aq. layers were combined, and extracted once with CH₂Cl₂ (520 mL). The organics were combined, and concentrated under vacuum to afford **27** as a light orange foam (250 g, 100%). The residue was used in the next stage without purification.

N-((4a*S*,5*S*,7a*S*)-7a-(2-fluorophenyl)-5-(trifluoromethyl)-4a,5,7,7a-tetrahydro-4H-furo[3,4-d][1,3]thiazin-2-yl)benzamide: ¹H NMR (500 MHz, CDCl₃) δ 8.03 (d, *J* = 6.7 Hz, 2H), 7.52 (t, *J* = 7.0 Hz, 1H), 7.48 – 7.31 (m, 4H), 7.20 (t, *J* = 7.4 Hz, 1H), 7.12 (dd, *J* = 12.0, 8.4 Hz, 1H), 4.82 – 4.73 (m, 1H), 4.60 (d, *J* = 8.9 Hz, 1H), 4.03 (d, *J* = 8.3 Hz, 1H), 3.57 (d, *J* = 2.7 Hz, 1H), 3.20 (d, *J* = 13.6 Hz, 1H), 2.81 (dd, *J* = 13.8, 2.5 Hz, 1H); ¹³C NMR (125 MHz, CDCl₃) δ 171.50, 159.57 (d, *J*_{CF} = 247.2 Hz), 134.62, 132.49, 130.65 (d, *J*_{CF} *J* = 8.8 Hz), 129.77, 128.51, 128.45, 125.14 (q, *J*_{CF} = 281.8 Hz), 124.97 (d, *J*_{CF} = 3.0 Hz), 124.66 (d, *J*_{CF} = 10.3 Hz), 117.05 (d, *J*_{CF} = 23.5 Hz), 66.81 (d, *J*_{CF} = 5.2 Hz), 38.90, 23.20.

HRMS Calculated for C₂₀H₁₆F₄N₂O₂S [M+H]⁺ 425.0947; found 425.0945.

30




N-((4aS,5R,7aS)-7a-(2-fluorophenyl)-5-methyl-4a,5,7,7a-tetrahydro-4H-furo[3,4-

d][1,3]thiazin-2-yl)benzamide. A reactor was charged with **25** (1 Wt, 1 eq), methylene chloride (8.0 Wt, 6.0 V) and pyridine (0.509 Wt, 0.520 V, 2.50 eq) at 18–22 °C. The mixture was cooled to -20 to -15 °C. A solution of trifluoromethanesulfonic anhydride (0.807 Wt, 0.481 V, 1.10 eq) in methylene chloride (1.32 Wt, 1.00 V) was added while maintaining T at -15 to -5 °C. Upon complete addition, continued stirring at -5 to 0 °C over 0.5h and monitored for complete consumption of **25** (target >97% conversion). Saturated aqueous ammonium chloride (2.67 Wt, 2.50 V) and water (1.50 Wt, 1.50 V) were added. The biphasic mixture was stirred vigorously at 18–22 °C and allowed to partition. The organic layer was aside and the aqueous layer was extracted with methylene chloride (4.7 Wt, 3.5 V). The combined organic layers were concentrated under reduced pressure (T <40 °C) to give **28** (0.95 Wt, assumed 100% theory).

N-((4aS,5R,7aS)-7a-(2-fluorophenyl)-5-methyl-4a,5,7,7a-tetrahydro-4H-furo[3,4-

d][1,3]thiazin-2-yl)benzamide: ¹H NMR (500 MHz, CDCl₃) δ 8.11 – 8.07 (m, 2H), 7.56 – 7.49 (m, 1H), 7.49 – 7.33 (m, 4H), 7.26 – 7.19 (m, 1H), 7.18 – 7.11 (m, 1H), 4.55 (d, J = 9.6 Hz, 1H), 4.55 – 4.46 (m, 1H), 4.16 (s, 2H), 4.06 (dd, J = 9.5, 2.8 Hz, 1H), 3.26 (dd, J = 13.7, 3.7 Hz, 1H), 2.93 – 2.87 (m, 1H), 2.84 (dd, J = 13.7, 3.8 Hz, 1H), 1.42 (d, J = 6.1 Hz, 3H); ¹³C NMR (126 MHz, cdcl₃) δ 174.13, 160.15 (d, J_{CF} = 247.1 Hz), 148.20, 135.46, 132.50, 130.58 (d, J_{CF} = 8.9 Hz), 128.97, 128.47, 128.43 (d, J_{CF} = 3.5 Hz), 126.94 (d, J_{CF} = 10.7 Hz), 125.00 (d, J_{CF} = 3.4 Hz), 117.17 (d, J_{CF} = 23.1 Hz), 77.56 (d, J_{CF} = 5.7 Hz), 76.60, 66.88 (d, J_{CF} = 4.4 Hz), 46.16 (d, J_{CF} = 2.9 Hz), 23.36, 19.67.




N-((4aS,5S,7aS)-7a-(2,3-Difluorophenyl)-5-((trityloxy)methyl)-4a,5,7,7a-tetrahydro-4H-furo[3,4-d][1,3]thiazin-2-yl)benzamide. To a solution of N-(((3S,4R,5S)-3-(2,3-

difluorophenyl)-4-(hydroxymethyl)-5-(((trityloxy)methyl)tetrahydrofuran-3-

yl)carbamothioyl)benzamide **26** (34.0 g, 0.0511 mol, 1.0 equiv.) in CH₂Cl₂ (204 mL) at -20 °C, pyridine (10.3 mL, 0.128 mol, 2.50 equiv.) was added. A solution of trifluoromethanesulfonic anhydride (9.46 mL, 0.0563 mol, 1.10 equiv.) in CH₂Cl₂ (34 mL) was then added over 12 min while maintaining temperature below -17 °C. The reaction was stirred for 30 min at -20 °C then allowed to warm up slowly to 5 °C. Sat aq. NH₄Cl (85 mL) and water (32 mL) were added. The aqueous layer was then extracted with CH₂Cl₂ (68 mL). The organic phase was concentrated under reduced pressure providing the title compound **29** (33.6 g) as an orange foam.

HRMS Calculated for C₃₉H₃₂F₂N₂O₃S [M+H]⁺ 647.2180; found 647.2123.

10



N-(3-((4aS,5S,7aS)-2-amino-5-(fluoromethyl)-4a,5,7,7a-tetrahydro-4H-furo[3,4-d][1,3]thiazin-7a-yl)-4-fluorophenyl)-5-(difluoromethyl)pyrazine-2-carboxamide. 5-


(Difluoromethyl)pyrazine-2-carboxylic acid (30.0 g, 1.05 equiv) and (4aS,5S,7aS)-7a-(5-amino-2-fluorophenyl)-5-(fluoromethyl)-4a,5,7,7a-tetrahydro-4H-furo[3,4-d][1,3]thiazin-2-amine (49.2 g, 1.00 equiv.) were charged to a reactor and EtOAc (443 mL) was added to the mixture to give a suspension. A solution of ®T3P (105 g, 1.10 equiv, 50 wt% in EtOAc) (Archimica, Germany) was added at ambient temperature while controlling the internal temperature below 30 °C. The reaction mixture was stirred at 40-45 °C >3 hours and monitored by HPLC. The reaction mixture was cooled to ambient temperature and water (98 mL) was charged. After 10-15 minutes charged 28% ammonium hydroxide (137 mL) while controlling the temperature below 30 °C. Additional EtOAc (172 mL) was added and the reaction mixture was stirred for 30 minutes at ambient temperature. The aqueous phase was separated and back-extracted with EtOAc (246 mL). The organic phases were combined and washed with 15% aq. NaCl (98 mL) and water (98 mL). The organic layer was filtered over Celite (0.5 Wt) and concentrated under vacuum to obtain a beige solid (quantitative crude yield) which was recrystallized from 1-propanol to afford **N-(3-((4aS,5S,7aS)-2-amino-5-(fluoromethyl)-4a,5,7,7a-tetrahydro-4H-furo[3,4-d][1,3]thiazin-7a-yl)-4-fluorophenyl)-5-(difluoromethyl)pyrazine-2-carboxamide** as a tan solid (54.2g).

25

^1H NMR (500 MHz, DMSO) δ = 10.94 (s, 1H), 9.38 (d, J =0.9, 1H), 9.08 (s, 1H), 7.93 (dd, J =7.3, 2.7, 1H), 7.90 – 7.85 (m, 1H), 7.25 (t, J =54.0, 2H), 7.18 (dd, J =11.8, 8.8, 1H), 6.04 (s, 2H), 4.68 – 4.44 (m, 2H), 4.40 – 4.30 (m, 1H), 4.26 (d, J =8.2, 1H), 3.79 (dd, J =8.1, 3.1, 1H), 3.30 (s, 1H), 3.12 (dd, J =13.5, 3.6, 1H), 3.00 (dd, J =13.5, 3.8, 1H), 2.76 (dt, J =7.9, 3.7, 1H).

5 ^{13}C NMR (126 MHz, DMSO) δ 160.90 (s), 156.22 (d, J = 244.2 Hz), 149.16 (t, J = 24.7 Hz), 148.46 (s), 146.91 (s), 143.69 (s), 140.35 (t, J = 4.4 Hz), 134.02 (s), 130.27 (d, J = 12.3 Hz), 122.46 (d, J = 4.2 Hz), 121.18 (d, J = 8.7 Hz), 116.26 (d, J = 25.0 Hz), 112.49 (t, J = 239.1 Hz), 83.53 (d, J = 169.8 Hz), 78.52 (d, J = 18.2 Hz), 77.55 (s), 65.61 (s), 35.59 (s), 22.93 (s).

10 E. Synthesis of Amino Thiazines of Formula V




(4a*S*,5*S*,7a*S*)-7a-(2-Fluorophenyl)-5-(trifluoromethyl)-4a,5,7,7a-tetrahydro-4H-furo[3,4-

15 d][1,3]thiazin-2-amine. To a solution of *N*-((4a*S*,5*S*,7a*S*)-7a-(2-fluorophenyl)-5-(trifluoromethyl)-4a,5,7,7a-tetrahydro-4H-furo[3,4-d][1,3]thiazin-2-yl)benzamide **27** (250.2 g, 589.5 mmol) in methanol (1.25 L) was added K_2CO_3 (81.5 g, 590.0 mmol). The suspension was heated to 65 °C for 6 hours. Upon cooling to ambient temperature, the solvent was evaporated under vacuum. To the resulting residue, was added 1.0 N aq NaOH (1.18 L) and THF (502 mL).
 20 The heterogeneous mixture was heated to 45 °C for 1 hour. The mixture was cooled to ambient temperature, and EtOAc (1.38 L) was added. The aqueous layer was extracted with EtOAc (0.75 L). The organics were combined, washed with saturated aq. NaHCO_3 (500 mL) and saturated aq. NaCl (500 mL). The organics were concentrated under vacuum to afford the title compound **30** as a brown oil (184.1 g, 91.6 % yield accounting for residual solvents).

25 (4a*S*,5*S*,7a*S*)-7a-(2-Fluorophenyl)-5-(trifluoromethyl)-4a,5,7,7a-tetrahydro-4H-furo[3,4-d][1,3]thiazin-2-amine: ^1H NMR (500 MHz, DMSO) δ 7.49 – 7.42 (m, 1H), 7.40 – 7.33 (m, 1H), 7.26 – 7.15 (m, 2H), 6.26 (s, 2H), 4.77 – 4.54 (m, 1H), 4.40 (d, J = 8.0 Hz, 1H), 3.80 (dd, J = 7.9, 2.3 Hz, 1H), 3.24 – 3.17 (m, 1H), 3.00 (dd, J = 13.9, 3.2 Hz, 1H), 2.85 (dd, J = 13.9, 3.9 Hz, 1H); ^{13}C NMR (125 MHz, DMSO) δ 159.75 (d, J_{CF} = 245.1 Hz), 149.51, 131.31 (d, J_{CF} = 3.9

Hz), 130.13 (d, $J_{CF} = 8.8$ Hz), 128.08 (d, $J_{CF} = 10.4$ Hz), 128.28 (q, $J_{CF} = 282.1$ Hz), 124.87 (d, $J_{CF} = 3.0$ Hz), 116.80 (d, $J = 23.8$ Hz), 78.77, 76.80 (q, $J_{CF} = 30.8$ Hz), 66.31, 36.37, 23.27.
 HRMS Calculated for $C_{13}H_{12}F_4N_2OS$ $[M+H]^+$ 321.0685; found 321.0677.




5

(4aS,5R,7aS)-7a-(2-Fluorophenyl)-5-methyl-4a,5,7,7a-tetrahydro-4H-furo[3,4-d][1,3]thiazin-2-amine. A reactor was charged with **28** (1.0 Wt, 1.0 eq), potassium carbonate (0.19 Wt, 0.50 eq) and MeOH (4.0 Wt, 5.0 V). The reaction mixture was heated up to 60–65 °C. Continued stirring at 60–65 °C over 6h and monitored the reaction for complete consumption of **28** (target >97% conversion). The reaction mixture was concentrated under reduced pressure ($T < 40$ °C) to remove most of the methanol. Water (1.5 Wt, 1.5 V) was charged followed by a mixture of heptane (1.1 Wt, 1.6 V) and ethyl acetate (0.36 Wt, 0.40 V). The mixture was cooled to 0–5 °C and stirred vigorously for 1–2h. The mixture was filtered and rinsed the reactor and filter cake with water (1.0 Wt, 1.0 V) and a pre-cooled (0–5 °C) mixture of heptane (0.89 Wt, 1.3 V) and ethyl acetate (0.29 Wt, 0.32V). The solid was dried in vacuo ($T < 40$ °C) to give **31** (0.61 Wt, 85%, pale brown solid).

(4aS,5R,7aS)-7a-(2-fluorophenyl)-5-methyl-4a,5,7,7a-tetrahydro-4H-furo[3,4-d][1,3]thiazin-2-amine: ^1H NMR (500 MHz, CDCl_3) δ 7.41 (td, $J = 8.2, 1.6$ Hz, 1H), 7.28 – 7.20 (m, 1H), 7.15 – 7.09 (m, 1H), 7.03 (ddd, $J = 12.4, 8.1, 0.7$ Hz, 1H), 4.81 – 4.42 (s, 2H), 4.61 (dd, $J = 8.7, 0.5$ Hz, 1H), 4.40 – 4.27 (m, 1H), 3.80 (dd, $J = 8.7, 2.2$ Hz, 1H), 3.07 (dd, $J = 13.3, 3.9$ Hz, 1H), 2.70 (dd, $J = 13.3, 3.9$ Hz, 1H), 2.56 – 2.47 (m, 1H), 1.34 (d, $J = 6.1$ Hz, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ 160.03 (d, $J_{CF} = 247.0$ Hz), 149.41, 130.73 (d, $J_{CF} = 10.0$ Hz), 129.80 (d, $J_{CF} = 4.1$ Hz), 128.83 (d, $J_{CF} = 8.7$ Hz), 124.02 (d, $J_{CF} = 3.4$ Hz), 116.38 (d, $J_{CF} = 23.5$ Hz), 78.54 (d, $J_{CF} = 4.8$ Hz), 76.43, 66.97 (d, $J_{CF} = 4.6$ Hz), 44.50 (d, $J_{CF} = 3.7$ Hz), 23.70, 19.83.

25

Alternate synthesis of (4aS,5R,7aS)-7a-(2-Fluorophenyl)-5-methyl-4a,5,7,7a-tetrahydro-4H-furo[3,4-d][1,3]thiazin-2-amine (31).




A reactor was charged with thiourea **25** (1 Wt, 1 eq), pyridine (2.9 Wt, 3.0 V) and toluene (2.6 Wt, 3.0 V). The resulting mixture was cooled to below -15 °C. Trifluoromethanesulfonic anhydride (0.800 Wt, 0.476 V, 1.10 eq) was added while maintaining the temperature below 0 °C.

5 Upon complete addition, stirring between -5–0 °C was continued for 1.5 h and the reaction was monitored for complete conversion ($\geq 97.0\%$). Toluene (5.2 Wt, 6.0 V) and 20 wt% aqueous ammonium chloride solution (4.2 Wt, 4.0 V) were added. The resulting biphasic mixture was warmed to 15–25 °C, stirring for at least 15 min and then allowed to partition. The lower aqueous layer was separated, and the upper organic phase was removed and held. The aqueous

10 layer was extracted with toluene (2.6 Wt, 3.0 V). The combined organic layers were washed with water (2.0 Wt, 2.0 V) and concentrated under reduced pressure to give crude **28** as a thick brown oil. 2-propanol (2.4 Wt, 3.0 V) and 3.0 M aqueous sodium hydroxide (3.5 Wt, 3.0 V) were added, and the resulting reaction mixture was heated to 80 °C for 6 h and monitored for complete conversion ($>99.0\%$). Heptane (1.4 Wt, 2.0 V) and then water (10 Wt, 10 V) were then

15 added, while keeping the internal temperature above 60 °C. The resulting mixture was cooled to 0–5 °C and maintained at this temperature for at least 2 h, then filtered. The reactor and filter cake were rinsed with a mixture of water (1.8 Wt, 1.8V) and 2-propanol (0.16 Wt, 0.20 V) and then a mixture of heptane (1.2 Wt, 1.8 V) and ethyl acetate (0.18 Wt, 0.20 V). The solids were dried to give isothiurea **31** (0.57 Wt, 83% yield).

20




N-((4aS,5S,7aS)-7a-(2,3-Difluorophenyl)-5-(hydroxymethyl)-4a,5,7,7a-tetrahydro-4H-furo[3,4-d][1,3]thiazin-2-yl)benzamide. Formic acid (116 mL) was added to N-((4aS,5S,7aS)-7a-(2,3-difluorophenyl)-5-((trityloxy)methyl)-4a,5,7,7a-tetrahydro-4H-furo[3,4-d][1,3]thiazin-2-yl)benzamide **29** (33.30 g, 0.051 mol, 1.0 equiv.), the mixture was stirred at rt for 25 min. Water

25

(17 mL) was added drop wise then the reaction was stirred for another 30 min. The resulting suspension was filtrated; the filtrate was dried by azeotropic distillation under vacuum with toluene (360 mL). Methanol (110 mL) was added followed by triethylamine (22 mL, 0.15 mol, 3.0 equiv.) and the solution was stirred for 2 hours. The solvent was removed under vacuum and chased with toluene (66 mL). The residual oil was dissolved in CH₂Cl₂ (250 ml) and washed successively with aq.HCl (1M, 100 mL), sat aq.NaHCO₃ (67 mL) and 18% aq.NaCl (67 mL). The organic phase was concentrated under reduce pressure. Toluene (66 mL) and THF (67 mL) were added to the residue then the solution was filtered through celite and concentrated under reduced pressure to yield the title compound **32** (21 g).

¹H NMR (500 MHz, DMSO) δ 8.11 (dd, *J* = 16.4, 4.2 Hz, 2H), 7.70 – 7.58 (m, 1H), 7.57 – 7.44 (m, 3H), 7.43 – 7.34 (m, 1H), 7.34 – 7.26 (m, 1H), 4.43 (d, *J* = 9.5 Hz, 1H), 4.25 – 4.16 (m, 1H), 4.10 (d, *J* = 9.2 Hz, 1H), 3.59 (d, *J* = 4.5 Hz, 2H), 3.30 – 3.21 (m, 1H), 3.20 – 3.08 (m, 2H); ¹³C NMR (125 MHz, DMSO) δ 169.08, 150.95 (dd, *J*_{CF} = 245.8, 13.1 Hz), 148.22, 148.09 (dd, *J*_{CF} = 248.7, 13.2 Hz), 133.46, 129.11, 128.99, 128.62, 125.43, 124.70, 118.16 (d, *J*_{CF} = 16.9 Hz), 81.36, 75.29, 66.76, 62.39, 40.77, 24.25.

HRMS Calculated for C₂₀H₁₈F₂N₂O₃S [M+H]⁺ 405.1084; found 405.1081.




N-((4aS,5S,7aS)-7a-(2,3-difluorophenyl)-5-(fluoromethyl)-4a,5,7,7a-tetrahydro-4H-furo[3,4-d][1,3]thiazin-2-yl)benzamide.

N-((4aS,5S,7aS)-7a-(2,3-difluorophenyl)-5-(hydroxymethyl)-4a,5,7,7a-tetrahydro-4H-furo[3,4-d][1,3]thiazin-2-yl)benzamide **32** (21.0 g, 0.519 mol, 1.0 equiv.) was dissolved in dry THF (105 mL) under a nitrogen atmosphere and the solution was cooled at 0 °C. N,N-diisopropylethylamine (40.7 mL, 0.234 mol, 4.50 equiv.), triethylamine trihydrofluoride (14.0 mL, 0.0857 mol, 1.65 equiv.) and perfluorobutanesulfonyl fluoride (22.4 mL, 0.125 mol, 2.40 equiv.) were added while maintaining temperature under 5 °C. The reaction was stirred for 3 hours at 0 °C then slowly warmed to room temperature and stirred for 11 hours. To the reaction mixture was charged sat.NH₄Cl (100 mL) followed by 2-methoxy-2-methylpropane (100 mL). The organic phase was isolated and washed with aq. HCl (1.0 M, 100 ml), concentrated and redissolved in 2-methoxy-2-methylpropane (301 ml). The organic

phase was then washed with aq.HCl (1M, 63 mL), sat aq.NaHCO₃ (100 mL). The last aqueous layer was extracted with 2-methoxy-2-methylpropane. The combined organic phase was washed with 18% aq.NaCl (63 mL) and concentrated under reduced pressure to afford title compound **33** (19.40 g) as a foam.

5 N-((4aS,5S,7aS)-7a-(2,3-difluorophenyl)-5-(fluoromethyl)-4a,5,7,7a-tetrahydro-4H-furo[3,4-d][1,3]thiazin-2-yl)benzamide: ¹H NMR (500 MHz, DMSO) δ 8.02 (s, 2H), 7.61 – 7.51 (m, 1H), 7.51 – 7.38 (m, 3H), 7.39 – 7.18 (m, 2H), 4.72 – 4.49 (m, 2H), 4.48 – 4.37 (m, 1H), 4.41 (d, *J* = 9.1 Hz, 2H), 3.04 (s, 2H); ¹³C NMR (125 MHz, DMSO) δ 150.92 (dd, *J*_{CF} = 245.5, 13.3 Hz), 148.11 (dd, *J*_{CF} = 248.2, 13.4 Hz), 132.40, 128.99, 128.63, 125.16, 124.94, 117.61, 10
83.64 (d, *J*_{CF} = 170.2 Hz), 79.51 (d, *J*_{CF} = 18.4 Hz), 76.17, 66.27, 23.67.

HRMS Calculated for C₂₀H₁₇F₃N₂O₂S [M+H]⁺ 407.1041; found 407.1024.




15 (4aS,5S,7aS)-7a-(2,3-Difluorophenyl)-5-(fluoromethyl)-4a,5,7,7a-tetrahydro-4H-furo[3,4-d][1,3]thiazin-2-amine. N-((4aS,5S,7aS)-7a-(2,3-difluorophenyl)-5-(fluoromethyl)-4a,5,7,7a-tetrahydro-4H-furo[3,4-d][1,3]thiazin-2-yl)benzamide **33** (19.40 g, 0.048 mol, 1.0 equiv.) was dissolved in methanol (97 mL), under nitrogen. 1,8-diazabicyclo[5.4.0]undec-7-ene (8.92 mL, 0.060 mol, 1.25 equiv.) was added, and the solution was heated to 55-60 °C. After 8 h, the reaction mixture was concentrated under reduced pressure, and the residue purified by silica gel flash column chromatography (5% to 100% EtOAc in heptane) to afford the title compound **34** (9.01 g).

25 (4aS,5S,7aS)-7a-(2,3-difluorophenyl)-5-(fluoromethyl)-4a,5,7,7a-tetrahydro-4H-furo[3,4-d][1,3]thiazin-2-amine: ¹H NMR (500 MHz, DMSO) δ 7.38 – 7.28 (m, 1H), 7.26 – 7.14 (m, 2H), 6.12 (s, 2H), 4.66 – 4.41 (m, 2H), 4.37 – 4.27 (m, 2H), 4.29 (d, *J* = 8.2 Hz, 1H), 3.74 (dd, *J* = 8.2, 2.6 Hz, 1H), 3.30 (s, 1H), 3.06 – 2.90 (m, 2H), 2.79 – 2.71 (m, 1H); ¹³C NMR (125 MHz, DMSO) δ 150.84 (dd, *J* = 244.8, 14.0 Hz), 149.71, 148.07 (dd, *J*_{CF} = 247.8, 13.4 Hz), 133.08 (d, *J*_{CF} = 7.7 Hz), 125.24, 124.55 (dd, *J*_{CF} = 7.2, 4.3 Hz), 116.58 (d, *J*_{CF} = 17.2 Hz), 83.91 (d, *J*_{CF} = 170.0 Hz), 79.12 (d, *J*_{CF} = 18.2 Hz), 77.96 (d, *J*_{CF} = 4.9 Hz), 66.22, 36.41 (d, *J*_{CF} = 3.3 Hz), 23.40.
30

HRMS Calculated for C₁₃H₁₃F₃N₂OS [M+H]⁺ 303.0779; found 303.0767.


G. Synthesis of Nitrophenyl Thiazines of Formula VI



5 (4a*S*,5*S*,7a*S*)-7a-(2-Fluoro-5-nitrophenyl)-5-(trifluoromethyl)-4a,5,7,7a-tetrahydro-4H-furo[3,4-d][1,3]thiazin-2-amine hydrochloride. To a cooled vessel containing (4a*S*,5*S*,7a*S*)-7a-(2-fluorophenyl)-5-(trifluoromethyl)-4a,5,7,7a-tetrahydro-4H-furo[3,4-d][1,3]thiazin-2-amine **30** (184.1 g, 574.8 mmol) was added trifluoroacetic acid (0.954 kg) in portions while the temperature was maintained below 20 °C. The mixture was cooled to 3.5 °C and sulfuric acid (146 mL, 2.73 mol) was added over 20 min while the temperature was maintained below 5 °C. Fuming nitric acid (39.8 mL, 0.948 mol) was added over 30 min, while the temperature was maintained below 10 °C. After 1.5 h at 0-10 °C, the reaction mixture was slowly quenched by transferring into an aq. solution of NaOH (575 g, 14.4 mol) in water (4.6 L) cooled to 5 °C. The resulting suspension was stirred for 1 h at 21 °C. The suspension was then filtered and the solid rinsed with cold water (920 mL). The solid was dried under vacuum until constant weight, and then dissolved into ethanol (1.05 L). The solution was heated to 35 °C, and conc. HCl (55.6 mL, 0.690 mol) was added while maintaining temperature below 40 °C. The suspension was then cooled to -5 °C, held for 1 hr and filtered. The solid was rinsed with cold ethanol (420 mL) and dried until constant weight to obtain the title compound **35** (185.0 g, 87.3%).

20 (4a*S*,5*S*,7a*S*)-7a-(2-Fluoro-5-nitrophenyl)-5-(trifluoromethyl)-4a,5,7,7a-tetrahydro-4H-furo[3,4-d][1,3]thiazin-2-amine hydrochloride: ¹H NMR (500 MHz, DMSO) δ 11.80 (s, 2H), 8.45 – 8.36 (m, 1H), 8.31 (dd, *J* = 6.6, 2.5 Hz, 1H), 7.66 (dd, *J* = 11.1, 9.3 Hz, 1H), 4.96 – 4.72 (m, 1H), 4.58 (d, *J* = 10.0 Hz, 1H), 4.27 (d, *J* = 9.9 Hz, 1H), 3.76 – 3.66 (m, 1H), 3.39 (dd, *J* = 14.9, 3.6 Hz, 1H), 3.24 (dd, *J* = 14.3, 4.6 Hz, 1H); ¹³C NMR (125 MHz, DMSO) δ 168.34, 163.33 (d, *J*_{CF} = 257.8 Hz), 144.58, 127.61 (d, *J*_{CF} = 11.6 Hz), 125.84, 124.10, 119.28 (d, *J*_{CF} = 26.5 Hz), 77.38 (q, *J*_{CF} = 31.5 Hz), 75.99, 65.88 (d, *J*_{CF} = 4.8 Hz), 40.36, 23.98.


HRMS Calculated for C₁₃H₁₁F₄N₃O₃S [M+H]⁺ 366.0536; found 366.0523.



(4aS,5R,7aS)-7a-(2-fluoro-5-nitrophenyl)-5-methyl-4a,5,7,7a-tetrahydro-4H-furo[3,4-

d][1,3]thiazin-2-amine. A reactor (reactor 1) was charged with trifluoroacetic acid (3.7 Wt, 2.5 V) and cool down to 0–5 °C. **31** (1Wt, 1 eq) was added portionswise with stirring, while maintaining T below 20 °C. The holding vessel and reactor were rinsed with trifluoroacetic acid (1.5 Wt, 1.0 V). The mixture was cooled below 5 °C and then added sulfuric acid (1.3 Wt, 0.72 V) while maintaining T below 15 °C. The resulting solution was cooled down to 0–5 °C and then added fuming nitric acid (purity > 90%, 0.315 Wt, 0.210 V, 1.20 eq) while maintaining T below 25 °C. Upon completed addition, continued stirring the mixture with cooling (bath T. 0–5 °C) over 0.5h and monitored the reaction for complete consumption of **31** (target >98% conversion). Another reactor (reactor 2) was charged with 3.0 M NaOH (26 Wt, 23 V) and the mixture was cooled to 3–7 °C. The reaction mixture was transferred into the second reactor while maintaining the T below 30 °C. During the addition, a white precipitate was formed. Upon completed addition, continued stirring at 23–27 °C. The first reactor was rinsed with water (2.0 Wt, 2.0 V), and transfer the water wash into reactor 2. Additional 3M NaOH to make the pH > 12 was added if necessary. After stirring for 0.5–1 h at 23–27 °C, the mixture was filtered and the reactor and filter cake were rinsed with water (4.0 Wt, 4.0 V). The filter cake was dried under reduced pressure (T <30 °C) overnight to give **36** (1.12 Wt, 96.1% yield).

(4aS,5R,7aS)-7a-(2-fluoro-5-nitrophenyl)-5-methyl-4a,5,7,7a-tetrahydro-4H-furo[3,4-d][1,3]thiazin-2-amine: ¹H NMR (500 MHz, CDCl₃) δ 8.39 (dd, *J* = 6.9, 2.9 Hz, 1H), 8.20 – 8.14 (m, 1H), 7.21 (dd, *J* = 10.8, 9.0 Hz, 1H), 4.55 (dd, *J* = 9.1, 0.9 Hz, 1H), 4.41 – 4.31 (m, 1H), 3.80 (dd, *J* = 9.1, 1.8 Hz, 1H), 3.05 (dd, *J* = 13.5, 3.8 Hz, 1H), 2.74 (dd, *J* = 13.5, 4.1 Hz, 1H), 2.52 – 2.42 (m, 1H), 1.37 (d, *J* = 6.1 Hz, 3H); ¹³C NMR (125 MHz, CDCl₃) δ 163.56 (d, *J*_{CF} = 258.9 Hz), 150.91, 144.23, 133.73 (d, *J*_{CF} = 12.3 Hz), 126.05 (d, *J*_{CF} = 6.2 Hz), 124.76 (d, *J*_{CF} = 10.8 Hz), 117.53 (d, *J*_{CF} = 26.3 Hz), 78.83 (d, *J*_{CF} = 4.0 Hz), 76.96, 66.91 (d, *J*_{CF} = 4.8 Hz), 45.26 (d, *J*_{CF} = 3.1 Hz), 23.88, 19.63.



(4aS,5S,7aS)-7a-(2,3-Difluoro-5-nitrophenyl)-5-(fluoromethyl)-4a,5,7,7a-tetrahydro-4H-furo[3,4-d][1,3]thiazin-2-amine.


(4aS,5S,7aS)-7a-(2,3-difluorophenyl)-5-(fluoromethyl)-

5 4a,5,7,7a-tetrahydro-4H-furo[3,4-d][1,3]thiazin-2-amine **34** (9.10 g, 0.03 mol, 1.0 equiv.) was dissolved in trifluoroacetic acid (36.4 mL), and the solution was cooled to 0 °C. Sulfuric acid (conc., 12.0 mL) was added, followed by fuming nitric acid (6.9 mL) drop wise while maintaining the temperature below 5 °C. After stirring at 0-5 °C for 4 h, the reaction mixture was slowly charged into a vigorously stirred solution of aq.NaOH (43.3 g in 273 mL water) while
10 maintaining a temperature below 20 °C. The mixture was extracted with CH₂Cl₂ (1 x 94 mL, and 2 x 64 mL), and the combined organic phases were washed with sat. aq. NaCl (46 mL). Celite (15.0g) was added to the organics, and the mixture was filtered rinsing with CH₂Cl₂ (40 mL). The solvents were evaporated to afford the title compound **37** (8.3 g) which was used in the subsequent step without purification.

15 (4aS,5S,7aS)-7a-(2,3-difluoro-5-nitrophenyl)-5-(fluoromethyl)-4a,5,7,7a-tetrahydro-4H-furo[3,4-d][1,3]thiazin-2-amine: ¹H NMR (500 MHz, DMSO) δ 8.41– 8.30 (m, 1H), 8.22 – 8.13 (m, 1H), 6.36 (s, 2H), 4.70 – 4.46 (m, 2H), 4.43 – 4.31 (m, 1H), 4.35 (d, *J* = 8.7 Hz, 1H), 3.69 (dd, *J* = 8.5, 1.6 Hz, 1H), 3.04 – 2.91 (m, 2H), 2.86 – 2.76 (m, 1H); ¹³C NMR (125 MHz, DMSO) δ 152.36 (dd, *J*_{CF} = 246.8, 12.3 Hz), 151.13, 150.32 (dd, *J*_{CF} = 238.2, 12.2 Hz), 143.25 (dd, *J*_{CF} = 7.9, 2.5 Hz), 134.70 (d, *J*_{CF} = 9.5 Hz), 121.28, 113.02 (d, *J*_{CF} = 22.3 Hz), 83.74 (d, *J*_{CF} = 170.1 Hz), 79.44 (d, *J*_{CF} = 18.3 Hz), 78.02 (d, *J* = 3.9 Hz), 36.96, 23.28.

HRMS Calculated for C₁₃H₁₂F₃N₃O₃S [M+H]⁺ 348.0630; found 348.0614.


H. Synthesis of Diamino Thiazines of Formula VII



(4a*S*,5*S*,7a*S*)-7a-(5-amino-2-fluorophenyl)-5-(trifluoromethyl)-4a,5,7,7a-tetrahydro-4H-furo[3,4-d][1,3]thiazin-2-amine. Ethanol (0.975 L) was added to iron powder (62.5g, 1.12 mol) under nitrogen atmosphere. Concentrated HCl (9.03 mL) was added at ambient temperature and the suspension was heated to 65 °C for 1.5 h. The suspension was then cooled to 50 °C, and sat. aq. NH₄Cl (299 g) were added. The temperature of the reaction mixture was allowed to reach 50 °C, and (4a*S*,5*S*,7a*S*)-7a-(2-fluoro-5-nitrophenyl)-5-(trifluoromethyl)-4a,5,7,7a-tetrahydro-4H-furo[3,4-d][1,3]thiazin-2-amine hydrochloride **35** (75.0 g, 187.0 mol) was added in portions while maintaining temperature below 68 °C. After 30 min, ethanol (0.45 L) was added, and the reaction mixture was cooled to 20-25 °C over 1 h. The suspension was stirred for 2 h and filtered over Celite (75 g) rinsing with ethanol (0.972 L). The solution was concentrated under vacuum to a brown solid. Water (0.9 L) was added followed by 3.0 N NaOH (0.187 L, 560 mmol) while maintaining temperature below 35 °C. The resulting suspension was stirred for 1 h at 20-25 °C. The suspension was filtered, and the solid was rinsed with cold water (0.38 L). The solid was dried under vacuum at 40-45 °C over 24 h to obtain the title compound **38** (57.7 g, 95.5 %).

(4a*S*,5*S*,7a*S*)-7a-(5-Amino-2-fluorophenyl)-5-(trifluoromethyl)-4a,5,7,7a-tetrahydro-4H-furo[3,4-d][1,3]thiazin-2-amine: ¹H NMR (500 MHz, DMSO) δ 6.81 (dd, *J* = 12.5, 8.6 Hz, 1H), 6.62 (dd, *J* = 7.0, 2.9 Hz, 1H), 6.50 – 6.42 (m, 1H), 6.16 (s, 2H), 4.96 (s, 2H), 4.72 – 4.54 (m, 1H), 4.35 (d, *J* = 7.8 Hz, 1H), 3.74 (dd, *J* = 7.8, 2.5 Hz, 1H), 3.18 – 3.08 (m, 1H), 3.01 (dd, *J* = 13.9, 3.0 Hz, 1H), 2.84 (dd, *J* = 13.8, 3.8 Hz, 1H); ¹³C NMR (125 MHz, DMSO) δ 156.20 (d, *J*_{CF} = 243.0 Hz), 148.73, 145.49, 127.86 (d, *J*_{CF} = 11.0 Hz), 116.79 (d, *J*_{CF} = 24.8 Hz), 116.10 (d, *J*_{CF} = 3.3 Hz), 114.10 (d, *J*_{CF} = 8.0 Hz), 78.89, 76.57 (q, *J*_{CF} = 31.0 Hz), 66.35, 36.35, 23.11.


HRMS Calculated for C₁₃H₁₃F₄N₃OS [M+H]⁺ 336.0794; found 336.0789.



(4a*S*,5*R*,7a*S*)-7a-(5-amino-2-fluorophenyl)-5-methyl-4a,5,7,7a-tetrahydro-4H-furo[3,4-d][1,3]thiazin-2-amine. To a suspension of powder Iron (6 eq) in ethanol (8.0 V) at 18–22 °C was added conc. HCl (37 Wt%, 0.60 eq). The resulting mixture was warmed up to 65–70 °C and stirred for 2h. Cooled down to 50-55 °C and added saturated NH₄Cl solution (4.0 wt). In

another reactor, **36** (1.0 Wt, 1.0 eq) was dissolved in a mixture of ethanol (5.0 V) and conc. HCl (1.0 eq). The resulting solution was added into the Iron slurry reactor while T was maintained below 65 °C, rinsing with ethanol (1.0V). The reaction mixture was stirred at 55–65 °C until complete consumption of **36**. The reaction mixture was diluted with 6.0V ethanol and cooled down to 15-20 °C. After stirred for 1-2h, the mixture was filtrated through a Celite pad (1.0 Wt), rinsing with 15V ethanol. Concentration in vacuo gave orange solid residue, which was dissolved in water (8V). Some orange particles were generated and removed by a polish filtration. 2V of water was used for rinse. 3M NaOH (3V) was added over 45 min at 18-25 °C. The product precipitated out as a white solid. After 1-2h stirring at 18-22 °C, the mixture was filtered and the filter cake was rinsed with water three times (2V each). Drying in an oven vacuum at 45 °C overnight provided **39** as an off-white solid in 92% isolated yields.

(4aS,5R,7aS)-7a-(5-amino-2-fluorophenyl)-5-methyl-4a,5,7,7a-tetrahydro-4H-furo[3,4-d][1,3]thiazin-2-amine: ¹H NMR (500 MHz, DMSO) δ 6.76 (dd, *J* = 12.3, 8.6 Hz, 1H), 6.59 (dd, *J* = 7.1, 2.8 Hz, 1H), 6.45 – 6.36 (m, 1H), 5.91 (s, 2H), 4.87 (s, 2H), 4.29 (d, *J* = 8.2 Hz, 1H), 4.20 – 4.11 (m, 1H), 3.61 (dd, *J* = 8.0, 2.7 Hz, 1H), 2.93 (dd, *J* = 13.3, 3.7 Hz, 1H), 2.82 (dd, *J* = 13.3, 3.7 Hz, 1H), 2.37 – 2.28 (m, 1H), 1.23 (d, *J* = 6.1 Hz, 3H); ¹³C NMR (126 MHz, DMSO) δ 151.95 (d, *J*_{CF} = 232.6 Hz), 148.19, 145.02, 131.37 (d, *J*_{CF} = 11.3 Hz), 116.56 (d, *J*_{CF} = 24.6 Hz), 115.42 (d, *J*_{CF} = 3.4 Hz), 113.38 (d, *J*_{CF} = 7.9 Hz), 78.56 (d, *J*_{CF} = 5.4 Hz), 76.00, 66.55 (d, *J*_{CF} = 4.9 Hz), 43.73 (d, *J*_{CF} = 3.6 Hz), 22.96, 20.32.




(4aS,5S,7aS)-7a-(5-Amino-2,3-difluorophenyl)-5-(fluoromethyl)-4a,5,7,7a-tetrahydro-4H-furo[3,4-d][1,3]thiazin-2-amine. To iron (8.03 g, 0.144 mol, 6.0 equiv.) was added ethanol (66.56 mL), followed by conc. HCl (62%, 1.2 mL, 0.014 mol, 0.60 equiv.). The mixture was heated to 65 °C for 2 h, and then sat NH₄Cl (33%, 33.3 mL) was added and the reaction temperature was maintained at 55 °C. A solution of (4aS,5S,7aS)-7a-(2,3-difluoro-5-nitrophenyl)-5-(fluoromethyl)-4a,5,7,7a-tetrahydro-4H-furo[3,4-d][1,3]thiazin-2-amine **37** (8.32g, 0.024 mol, 1.0 equiv.) and conc. HCl (1.93 mL, 0.024 mol, 1.0 equiv.) in Ethanol (41.6 mL) was added to the iron suspension. The reaction was stirred for 30 min at 55 °C, then ethanol (50 mL) was added and the suspension was allowed to cool down to 20°C, filtered through celite

(8.3 g) and rinsed with ethanol (125 mL). The filtrate was concentrated under reduced pressure then water (66 mL) was added, followed by 3.0 M aq. NaOH (24 mL, 0.0179 mol, 3.0 equiv.). The resulting mixture was extracted with CH₂Cl₂ (2 x 83 mL). The organics phases were combined and filtered over celite and concentrated under reduce pressure to give the title compound **40** (7.60 g).

(4a*S*,5*S*,7a*S*)-7a-(5-Amino-2,3-difluorophenyl)-5-(fluoromethyl)-4a,5,7,7a-tetrahydro-4H-furo[3,4-*d*][1,3]thiazin-2-amine: ¹H NMR (500 MHz, DMSO) δ 6.42–6.35 (m, 2H), 6.02 (brs, 2H), 4.66–4.41 (m, 2H), 4.37–4.25 (m, 1H), 4.21 (d, *J* = 8.0 Hz, 1H), 3.69 (dd, *J* = 7.8, 2.2 Hz, 1H), 2.97 (qd, *J* = 13.5, 3.4 Hz, 3H), 2.74–2.64 (m, 1H); ¹³C NMR (125 MHz, DMSO) δ 151.08 (dd, *J*_{CF} = 240.1, 14.8 Hz), 148.88, 145.40 (d, *J*_{CF} = 10.7 Hz), 139.33 (dd, *J*_{CF} = 233.5, 13.8 Hz), 132.53 (d, *J*_{CF} = 8.1 Hz), 109.90, 100.70 (d, *J*_{CF} = 20.0 Hz), 83.99 (d, *J*_{CF} = 169.9 Hz), 78.79 (d, *J*_{CF} = 18.3 Hz), 77.99 (d, *J*_{CF} = 5.5 Hz), 66.22, 36.24, 23.14.

HRMS Calculated for C₁₃H₁₄F₃N₃OS [M+H]⁺ 318.0888; found 318.0874.

I. Synthesis of Compounds of Formula IX



N-(3-((4a*S*,5*S*,7a*S*)-2-Amino-5-(trifluoromethyl)-4a,5,7,7a-tetrahydro-4H-furo[3,4-*d*][1,3]thiazin-7a-yl)-4-fluorophenyl)-5-methoxypyrazine-2-carboxamide.

A suspension of 5-methoxypyrazine-2-carboxylic acid (26.29 g, 0.17 mol) in N,N'-dimethylimidazoline-2-one (160 mL) was stirred at ambient temperature for 15 min, then cooled to 2.2 °C. Thionyl chloride (14.7 mL, 0.202 mol) was added while maintaining temperature under 5 °C. The resulting suspension was stirred at 0-10 °C for 2 hours while it transitioned to a clear solution. In another vessel, (4a*S*,5*S*,7a*S*)-7a-(5-amino-2-fluorophenyl)-5-(trifluoromethyl)-4a,5,7,7a-tetrahydro-4H-furo[3,4-*d*][1,3]thiazin-2-amine **38** (52.0 g, 0.155 mol) was dissolved into N,N'-dimethylimidazoline-2-one (160 mL). The resulting solution was added to the solution of acyl chloride while maintaining temperature below 10 °C. The reaction mixture was stirred for 30 min. Water (780 mL) was charged while maintaining temperature below 30 °C. The resulting mixture was stirred for 30 min., and then EtOAc (780 mL) was added. To this mixture was added, 50% aq. NaOH

(84.8 g) until the pH of the aqueous layer reached 11. The aq. layer was extracted with EtOAc (260 mL). The organics were combined, washed with sat. aq. NaCl (260 mL) and water (260 mL). The organics were filtered over Celite pad (26 g) and rinsed with EtOAc (260 mL). The organics were concentrated under vacuum to afford a solid. To the solid was added 1-propanol (728 mL), and the suspension was heated to 75 °C until a clear solution formed. The solution was cooled to -10 °C and held for 1 hour. The solid was filtered, rinsed with cold 1-propanol (104 mL) and dried under vacuum (35 °C) until constant weight to afford the title compound **41** (62.1 g, 84.9 %).

N-(3-((4a*S*,5*S*,7a*S*)-2-Amino-5-(trifluoromethyl)-4a,5,7,7a-tetrahydro-4H-furo[3,4-
 10 d][1,3]thiazin-7a-yl)-4-fluorophenyl)-5-methoxypyrazine-2-carboxamide: ¹H NMR (500 MHz, DMSO) δ 10.56 (s, 2H), 8.88 (d, *J* = 1.2 Hz, 1H), 8.39 (d, *J* = 1.2 Hz, 1H), 7.95 – 7.83 (m, 2H), 7.18 (dd, *J* = 12.0, 8.8 Hz, 1H), 6.25 (s, 2H), 4.76 – 4.60 (m, 1H), 4.36 (d, *J* = 8.1 Hz, 1H), 4.01 (s, 3H), 3.88 (dd, *J* = 7.9, 2.3 Hz, 1H), 3.23 – 3.11 (m, 2H), 2.91 (dd, *J* = 13.8, 3.6 Hz, 1H); ¹³C NMR (125 MHz, DMSO) δ 162.11, 161.93, 156.13 (d, *J*_{CF} = 242.9 Hz), 149.38, 142.01, 138.35,
 15 135.09, 133.98, 128.53 (d, *J*_{CF} = 11.6 Hz), 126.06 (q, *J*_{CF} = 282.0 Hz), 123.32, 121.93 (d, *J*_{CF} = 8.6 Hz), 116.76 (d, *J*_{CF} = 25.1 Hz), 78.86 (d, *J*_{CF} = 6.9 Hz), 76.94 (q, *J*_{CF} = 30.5 Hz), 66.37, 54.75, 36.44, 23.53.

HRMS Calculated for C₁₉H₁₇F₄N₅O₃S [M+H]⁺ 472.1066; found 472.1052.

20 Specific optical rotation [α]_D +110.5 (c 0.584, MeOH)

Specific optical rotation parameters:

Equipment:

Polarimeter:	Perkin Elmer, model 341 or equivalent.
Cell:	Microglass cell, 100 mm pathlength, 1.0 mL capacity, Perkin-Elmer Cat. # B001-7047.
Balance:	Calibrated analytical balance capable of weighing ± 0.1 mg
Water Bath:	NESLAB RTE 1121 Chiller or equivalent.
Volumetric glassware:	Class A.
Quartz Standard	ID number 098799, or equivalent.
Polarimeter:	Perkin Elmer, model 341 or equivalent.


Reagents:

Methanol: HPLC grade, Baker (catalog no. 9093-03) or equivalent.

Instrument parameters:

Lamp: Na/Hal, Perkin-Elmer Cat. # B000-8754.
 Cell: Microcell (100mm), Perkin-Elmer Cat. #B004-1693.
 Cell Path: 100 mm (1 decimeter)
 Mode: OROT
 Wavelength: 589 nm
 Cell Temperature: 20°C
 Integration time: 2 seconds
 Aperture: MICRO
 Water bath temperature: 20 ± 1°C

5




N-(3-((4aS,5R,7aS)-2-amino-5-methyl-4a,5,7,7a-tetrahydro-4H-furo[3,4-d][1,3]thiazin-7a-yl)-4-fluorophenyl)-5-(difluoromethyl)pyrazine-2-carboxamide. 5-(Difluoromethyl)pyrazine-2-carboxylic acid **42** (0.68 wt, 1.1 eq) was charged into a reactor and if needed, dried by azeotropic distillation with toluene. DMI (2.0V) was added and the mixture was cooled down to 0-5 °C. Thionyl chloride (0.337 V, 1.30 eq) was added while keeping the internal T below 10 °C. The resulting solution was stirred at 4-10 °C until the conversion was >95.0 % by HPLC. The mixture was cooled down to 0 °C and a solution of **39** (1.0 Wt, 1.0 eq) in DMI (2.5V) was charged while keeping the internal temperature below 5 °C. The vessel was rinsed with DMI (0.50V) and the reaction mixture was stirred until the conversion is >99%. Water (12V) was charged and the resulting mixture is stirred at 15-20 °C for 0.5h. EtOAc (15V) and then 50% aq. NaOH (1.5 Wt, 5.3 eq) were added while keeping T below 30 °C. pH was monitored to make sure that it is above 10. The aqueous phase was separated and extracted with EtOAc (5.0V). The

15

organic layers were combined and washed with brine (5.0V) and water twice (5.0V each). The mixture was filtered through a Celite pad (0.50 Wt) rinsing with EtOAc (3.0V). The filtrate was concentrated under reduced pressure at 40-50 °C. 1-Propanol (15V) was charged and the mixture was warmed up to 90-100 °C while stirring until a clear solution was obtained. The mixture was cooled down to 0-5 °C over 2h and stirred over 1h. The mixture was filtered, rinsing with cold 1-propanol twice (2.5V each). Drying in an oven vacuum at 45 °C overnight provided the title compound **43** as an off-white solid in 81% isolated yield.

5-(difluoromethyl)pyrazine-2-carboxylic acid: ¹H NMR (500 MHz, DMSO) δ 4.72 (s, 1H), 4.57 (s, 1H), 4.32 (s, 1H), 2.06 (t, *J* = 54.2 Hz, 1H); ¹³C NMR (125 MHz, DMSO) δ 164.34, 151.67 (t, *J*_{CF} = 26.8 Hz), 145.45, 143.29, 141.04 (t, *J*_{CF} = 3.4 Hz), 112.59 (t, *J*_{CF} = 242.1 Hz).

N-(3-((4a*S*,5*R*,7a*S*)-2-amino-5-methyl-4a,5,7,7a-tetrahydro-4H-furo[3,4-*d*][1,3]thiazin-7a-yl)-4-fluorophenyl)-5-(difluoromethyl)pyrazine-2-carboxamide: ¹H NMR (500 MHz, CD₃OD) δ 9.38 (s, 1H), 9.00 (s, 1H), 7.89 – 7.77 (m, 2H), 7.14 (dd, *J* = 11.9, 8.8 Hz, 1H), 6.94 (t, *J* = 54.3 Hz, 1H), 4.80 (s, 2H), 4.58 (d, *J* = 8.9 Hz, 1H), 4.37 – 4.28 (m, 1H), 3.81 (dd, *J* = 8.8, 2.4 Hz, 1H), 3.14 (dd, *J* = 13.5, 4.0 Hz, 1H), 2.88 (dd, *J* = 13.5, 4.1 Hz, 1H), 2.61 – 2.54 (m, 1H), 1.33 (d, *J* = 6.1 Hz, 3H); ¹³C NMR (125 MHz, CD₃OD) δ 161.11, 156.94 (d, *J*_{CF} = 244.9 Hz), 154.91, 150.21 (t, *J*_{CF} = 25.8 Hz), 146.53, 143.30, 140.02 (t, *J*_{CF} = 4.0 Hz), 133.73 (d, *J*_{CF} = 2.6 Hz), 130.70 (d, *J*_{CF} = 11.6 Hz), 122.05 (d, *J*_{CF} = 3.9 Hz), 121.48 (d, *J*_{CF} = 8.8 Hz), 116.39 (d, *J*_{CF} = 25.3 Hz), 112.96 (t, *J*_{CF} = 239.8 Hz), 77.56 (d, *J*_{CF} = 5.3 Hz), 76.79, 66.49 (d, *J*_{CF} = 4.8 Hz), 45.16, 45.13, 22.60, 18.58.



N-(3-((4a*S*,5*S*,7a*S*)-2-Amino-5-(fluoromethyl)-4a,5,7,7a-tetrahydro-4H-furo[3,4-

d][1,3]thiazin-7a-yl)-4,5-difluorophenyl)-5-methoxypyrazine-2-carboxamide. A suspension of 5-methoxypyrazine-2-carboxylic acid (4.01 g, 0.026 mol, 1.10 equiv.) in N,N'-dimethylimidazoline-2-one (22.5 mL) was stirred at ambient temperature for 15 min, then cooled to 0 °C. Thionyl chloride (2.24 mL, 0.031 mol, 1.3 equiv.) was added while maintaining temperature under 10 °C. The resulting suspension was stirred at 0-10 °C for 2 hours while it transitioned to a clear solution. In another vessel, (4a*S*,5*S*,7a*S*)-7a-(5-amino-2,3-difluorophenyl)-

5-(fluoromethyl)-4a,5,7,7a-tetrahydro-4H-furo[3,4-d][1,3]thiazin-2-amine **40** (7.60 g, 0.024 mol, 1.0 equiv.) was dissolved into N,N'-dimethylimidazoline-2-one (22.5 mL). The resulting solution was added to the solution of acyl chloride while maintaining temperature below 10 °C. The reaction mixture was stirred for 30 min. Water (112 mL) was charged while maintaining temperature below 30 °C. The resulting mixture was stirred for 30 min., and then EtOAc (112 mL) was added. To this mixture was added, 50% aq. NaOH (10.0 g) until the pH of the aqueous layer reached 11. The aq. layer was extracted with EtOAc (75 mL). The organics were combined, washed with sat. aq. NaCl (38 mL) and water (38 mL). The organics were filtered over a pad of silica gel (15 g) and rinsed with EtOAc (37.5 mL). The organics were concentrated under vacuum to afford a solid. To the solid was added 1-propanol (112 mL), and the suspension was heated to 100 °C. The mixture was cooled to -10 °C and held for 1 hour. The solid was filtered, rinsed with cold 1-propanol (15 mL) and dried under vacuum (35 °C) until constant weight to afford the title compound **44** (8.18 g).

N-(3-((4aS,5S,7aS)-2-Amino-5-(fluoromethyl)-4a,5,7,7a-tetrahydro-4H-furo[3,4-d][1,3]thiazin-7a-yl)-4,5-difluorophenyl)-5-methoxypyrazine-2-carboxamide: ¹H NMR (500 MHz, DMSO) δ 10.73 (s, 1H), 8.88 (d, *J* = 0.8 Hz, 1H), 8.39 (d, *J* = 0.9 Hz, 1H), 8.11 – 7.87 (m, 1H), 7.73 (d, *J* = 5.0 Hz, 1H), 6.07 (s, 2H), 4.68 – 4.44 (m, 2H), 4.41 – 4.28 (m, 1H), 4.23 (d, *J* = 8.3 Hz, 1H), 4.01 (s, 3H), 3.82 (dd, *J* = 8.1, 2.4 Hz, 1H), 3.18 (dd, *J* = 13.5, 3.5 Hz, 1H), 3.01 (dd, *J* = 13.5, 3.8 Hz, 1H), 2.77 – 2.69 (m, 1H); ¹³C NMR (125 MHz, DMSO) δ 162.24, 162.20, 150.05 (dd, *J*_{CF} = 242.1, 14.4 Hz), 149.37, 144.42 (dd, *J*_{CF} = 245.5, 13.7 Hz), 142.18, 138.09, 134.73 (dd, *J*_{CF} = 10.3, 2.7 Hz), 134.03, 133.22 (d, *J*_{CF} = 9.1 Hz), 116.6, 108.42 (d, *J*_{CF} = 22.4 Hz), 83.98 (d, *J*_{CF} = 169.9 Hz), 79.21 (d, *J*_{CF} = 18.2 Hz), 77.96 (d, *J*_{CF} = 5.2 Hz), 66.26, 54.78, 36.23, 23.64.

HRMS Calculated for C₁₉H₁₈F₃N₅O₃S [M+H]⁺ 454.1161; found 454.1149.

Specific optical rotation [α]_D +115.3 (c 0.584, MeOH)

Specific optical rotation parameters:

Equipment:

Polarimeter:	Perkin Elmer, model 341 or equivalent.
Cell:	Microglass cell, 100 mm pathlength, 1.0 mL capacity, Perkin-Elmer Cat. # B001-7047.
Balance:	Calibrated analytical balance capable of weighing ± 0.1 mg

Water Bath:	NESLAB RTE 1121 Chiller or equivalent.
Volumetric glassware:	Class A.
Quartz Standard	ID number 098799, or equivalent.
Polarimeter:	Perkin Elmer, model 341 or equivalent.


Reagents:

Methanol:	HPLC grade, Baker (catalog no. 9093-03) or equivalent.
-----------	--

Instrument parameters:

Lamp:	Na/Hal, Perkin-Elmer Cat. # B000-8754.
Cell:	Microcell (100mm), Perkin-Elmer Cat. #B004-1693.
Cell Path:	100 mm (1 decimeter)
Mode:	OROT
Wavelength:	589 nm
Cell Temperature:	20°C
Integration time:	2 seconds
Aperture:	MICRO
Water bath temperature:	20 ± 1°C

5



N-(3-((4aS,5S,7aS)-2-amino-5-(trifluoromethyl)-4a,5,7,7a-tetrahydro-4H-furo[3,4-d][1,3]thiazin-7a-yl)-4-fluorophenyl)-5-(fluoromethyl)pyrazine-2-carboxamide. 5-

10 (Fluoromethyl)pyrazine-2-carboxylic acid (32.6 g, 1.05 equiv) and (4aS,5S,7aS)-7a-(5-amino-2-fluorophenyl)-5-(trifluoromethyl)-4a,5,7,7a-tetrahydro-4H-furo[3,4-d][1,3]thiazin-2-amine (70.0 g, 1.0 equiv) were charged to a reactor and EtOAc (630 mL) was added to the mixture to give a suspension. A solution of [®]T3P (146 g, 1.10 equiv, 50 wt% in EtOAc) (Archimica, Germany) was added at ambient temperature while controlling the internal temperature below 30 °C. The
15 reaction mixture was stirred at 40-45 °C >3 hours and monitored by HPLC. The reaction mixture was cooled to 15-20 °C and water (140 mL) was charged. After 10-15 minutes charged 28%


ammonium hydroxide (175 mL) while controlling the temperature below 30 °C. EtOAc (245 mL) was added and the reaction mixture was stirred for 30 minutes at ambient temperature. The aqueous phase was separated and back-extracted with EtOAc (490 mL). The organic phases were combined and washed with 15% aq. NaCl (140 mL) and water (140 mL). The organic layer was
5 filtered over Celite (1.0 Wt) and rinsed with EtOAc (140 mL). The solution was concentrated under vacuum to obtain a beige solid (quantitative crude yield) which was recrystallized from 1-propanol to afford **N-(3-((4a*S*,5*S*,7a*S*)-2-amino-5-(trifluoromethyl)-4a,5,7,7a-tetrahydro-4H-furo[3,4-*d*][1,3]thiazin-7a-yl)-4-fluorophenyl)-5-(fluoromethyl)pyrazine-2-carboxamide** as a white solid (70.0 g).

10 ¹H NMR (500 MHz, DMSO) δ 10.89 (s, 1H), 9.30 (s, 1H), 8.89 (s, 1H), 7.95 (dd, *J* = 7.3, 2.7 Hz, 1H), 7.94 – 7.89 (m, 1H), 7.21 (dd, *J* = 12.0, 8.8 Hz, 1H), 6.22 (s, 2H), 5.71 (d, *J* = 46.3 Hz, 2H), 4.77 – 4.61 (m, 1H), 4.37 (d, *J* = 8.1 Hz, 1H), 3.87 (dd, *J* = 8.0, 2.7 Hz, 1H), 3.20 (dt, *J* = 7.0, 3.5 Hz, 1H), 3.15 (dd, *J* = 13.9, 3.1 Hz, 1H), 2.91 (dd, *J* = 13.8, 3.8 Hz, 1H).

15 ¹³C NMR (126 MHz, DMSO) δ 161.32 (s), 155.82 (d, *J* = 243.4 Hz), 153.71 (d, *J* = 18.7 Hz), 148.77 (s), 144.71 (d, *J* = 1.9 Hz), 143.30 (s), 141.01 (d, *J* = 5.6 Hz), 134.36 (d, *J* = 2.0 Hz), 128.20 (d, *J* = 12.1 Hz), 125.57 (q, *J* = 283.0 Hz), 123.12 (d, *J* = 3.6 Hz), 121.64 (d, *J* = 8.6 Hz), 116.35 (d, *J* = 25.2 Hz), 82.55 (d, *J* = 165.8 Hz), 78.37 (s), 76.44 (q, *J* = 30.6 Hz), 65.89 (d, *J* = 5.3 Hz), 35.89 (s), 23.01 (s).

We claim:

1. A compound of **Formula I**:



wherein:

R₁ is C₃₋₄alkyl, halo-C₁₋₄alkyl, hydroxy-C₁₋₄alkyl, or -C₁₋₄alkyl-OR', wherein R' is an oxygen protecting group; and

R₂ and R₃ are each independently hydrogen or halo;

subject to the proviso that when one of R₂ or R₃ is *ortho*-fluoro, and the other is hydrogen, R₁ is not -CH₂F, or a salt thereof.

2. The compound of claim 1, wherein the oxygen protecting group is selected from the group consisting of: Trt (triphenylmethyl), MPM (p-methoxybenzyloxymethyl), TBDMS (t-butyldimethylsilyl), and TBDPS (t-butyldiphenyl silyl), or a salt thereof.

3. The compound of claim 1, wherein:

R₁ is trifluoromethyl, hydroxymethyl, monofluoromethyl, or triphenylmethoxymethyl, or a salt thereof.

4. The compound of claim 1, wherein:


R₁ is trifluoromethyl, hydroxymethyl, monofluoromethyl, or triphenylmethoxymethyl;

R₂ is fluoro; and

R₃ is hydrogen or fluoro,


or a salt thereof.

5. The compound of claim 1, wherein said compound is selected from the group consisting of:



or a salt thereof.

6. A method of making a compound of **Formula I**,




wherein:

R₁ is C₁₋₄alkyl, halo-C₁₋₄alkyl, hydroxy-C₁₋₄alkyl, or -C₁₋₄alkyl-OR',
wherein R' is an oxygen protecting group; and

R₂ and R₃ are each independently hydrogen or halo,
or a salt thereof,

said method comprising heating a mixture comprising:

(i) an oxime of **Formula A**:



wherein R₁, R₂ and R₃ are as given above;


(ii) hydroquinone; and

(iii) a hydrocarbon solvent,

to a temperature of from 90°C to 160°C,

to make said compound of **Formula I**, or a salt thereof.

7. The method of claim 6, wherein the hydrocarbon solvent is selected from the group consisting of: benzene, toluene, and xylenes.
8. The method of claim 6, wherein the hydrocarbon solvent is xylenes.
9. The method of any one of claims 6-8, wherein said heating is to a temperature of from 110°C to 140°C.
10. The method of any one of claims 6-8, wherein said heating is to a temperature of from 100°C to 120°C.
11. The method of any one of claims 6-8, wherein said heating is to a temperature of from 120°C to 140°C.
12. A compound of **Formula II**:



wherein:

R_1 is C_{3-4} alkyl, halo- C_{1-4} alkyl, hydroxy- C_{1-4} alkyl, or $-C_{1-4}$ alkyl-OR', wherein R' is an oxygen protecting group; and

R_2 and R_3 are each independently hydrogen or halo;

subject to the proviso that when one of R_2 or R_3 is *ortho*-fluoro and the other is hydrogen, R_1 is not $-CH_2F$,

or a salt thereof.

13. The compound of claim 12, wherein the oxygen protecting group is selected from the group consisting of: Trt (triphenylmethyl), MPM (p-methoxybenzyloxymethyl), TBDMS (t-butyldimethylsilyl), and TBDPS (t-butyldiphenyl silyl),
- or a salt thereof.

14. The compound of claim 12, wherein:

R_1 is trifluoromethyl, hydroxymethyl, monofluoromethyl, or triphenylmethoxymethyl, or a salt thereof.

15. The compound of claim 12, wherein:


R_1 is trifluoromethyl, hydroxymethyl, monofluoromethyl, or triphenylmethoxymethyl;

R_2 is fluoro; and

R_3 is hydrogen or fluoro,


or a salt thereof.

16. The compound of claim 12, wherein said compound is selected from the group consisting of:



or a salt thereof.

17. A method of making a stereochemically pure compound of **Formula IIa**:



IIa


wherein:

R_1 is C_{1-4} alkyl, halo- C_{1-4} alkyl, hydroxy- C_{1-4} alkyl, or $-C_{1-4}$ alkyl-OR', wherein R' is an oxygen protecting group; and

R_2 and R_3 are each independently hydrogen or halo, or a salt thereof,

said method comprising the steps of:

(a) adding a mixture of stereoisomers of the compound of **Formula II**:



II

wherein R₁, R₂ and R₃ are as given above,

to a solvent comprising a chiral carboxylic acid compound, to form a mixture of diastereomeric salts of the compound of **Formula II**; and then

(b) crystallizing a single diastereomeric salt formed of the compound of **Formula II**, to make said stereochemically pure compound of **Formula IIa**, or a salt thereof.

18. The method of claim 17, wherein said stereochemically pure compound has greater than about 90% by weight of one stereoisomer of the compound and less than about 10% by weight of the other stereoisomers of the compound.

19. The method of claim 17, wherein said stereochemically pure compound has greater than about 97% by weight of one stereoisomer of the compound and less than about 3% by weight of the other stereoisomers of the compound.

20. The method of any one of claims 17-19, wherein said crystallizing step is performed by cooling the mixture of diastereomeric salts of the compound of **Formula II**.

21. The method of claim 20, wherein said cooling is from a temperature of from 40°C to 80°C, to a temperature of from -10°C to 29°C.

22. The method of any one of claims 17-21, wherein the solvent is an alcohol.


23. The method of any one of claims 17-21, wherein the solvent is ethanol.

24. The method of any one of claims 17-21, wherein the solvent is a mixture of two or more solvents.

25. The method of claim 24, wherein the mixture is a toluene – acetonitrile mixture, a toluene – acetone mixture, a toluene – tetrahydrofuran mixture, or an alcohol – water mixture.

26. The method of any of claims 17-25, wherein the chiral carboxylic acid compound is selected from the group consisting of: D-dibenzoyl tartaric acid, L-dibenzoyl tartaric acid, D-ditoluyl tartaric acid, and L-ditoluyl tartaric acid.

27. A compound of **Formula III**:



wherein:

R_1 is C_{3-4} alkyl, halo- C_{1-4} alkyl, hydroxy- C_{1-4} alkyl, or $-C_{1-4}$ alkyl-OR', wherein R' is an oxygen protecting group; and

R_2 and R_3 are independently hydrogen or halo; subject to the proviso that when one of R_2 or R_3 is *ortho*-fluoro, and the other is hydrogen, R_1 is not $-CH_2F$, or a salt thereof.

28. The compound of claim 27, wherein the oxygen protecting group is selected from the group consisting of: Trt (triphenylmethyl), MPM (p-methoxybenzyloxymethyl), TBDMS (t-butyldimethylsilyl), and TBDPS (t-butyldiphenyl silyl), or a salt thereof.

29. The compound of claim 27, wherein:


R_1 is trifluoromethyl, hydroxymethyl, monofluoromethyl, or triphenylmethoxymethyl, or a salt thereof.

30. The compound of claim 27, wherein:

R_1 is trifluoromethyl, hydroxymethyl, monofluoromethyl, or triphenylmethoxymethyl;
 R_2 is fluoro; and
 R_3 is hydrogen or fluoro,


or a salt thereof.

31. The compound of claim 27, wherein said compound is selected from the group consisting of:



or a salt thereof.

32. A compound of **Formula IV**:



wherein:

R₁ is C₃₋₄alkyl, halo-C₁₋₄alkyl, hydroxy-C₁₋₄alkyl, or -C₁₋₄alkyl-OR', wherein R' is an oxygen protecting group; and

R₂ and R₃ are each independently hydrogen or halo; subject to the proviso that when one of R₂ or R₃ is *ortho*-fluoro, and the other is hydrogen, R₁ is not -CH₂F, or a salt thereof.

33. The compound of claim 32, wherein the oxygen protecting group is selected from the group consisting of: Trt (triphenylmethyl), MPM (p-methoxybenzyloxymethyl), TBDMS (t-butyldimethylsilyl), and TBDPS (t-butyldiphenyl silyl), or a salt thereof.

34. The compound of claim 32, wherein:

R₁ is trifluoromethyl, hydroxymethyl, monofluoromethyl, or triphenylmethoxymethyl; or a salt thereof.

35. The compound of claim 32, wherein:


R_1 is trifluoromethyl, hydroxymethyl, monofluoromethyl, or triphenylmethoxymethyl;

R_2 is fluoro; and

R_3 is hydrogen or fluoro,


or a salt thereof.

36. The compound of claim 32, wherein said compound is selected from the group consisting of:



or a salt thereof.

37. A compound of **Formula V**:



wherein:

R_1 is C_{3-4} alkyl, halo- C_{1-4} alkyl, hydroxy- C_{1-4} alkyl, or $-C_{1-4}$ alkyl-OR',
wherein R' is an oxygen protecting group; and

R_2 and R_3 are each independently hydrogen or halo;

subject to the proviso that when one of R_2 or R_3 is *ortho*-fluoro, and the other is
hydrogen, R_1 is not $-CH_2F$,

or a salt thereof.

38. The compound of claim 37, wherein the oxygen protecting group is selected from the group consisting of: Trt (triphenylmethyl), MPM (p-methoxybenzyloxymethyl), TBDMS (t-butyldimethylsilyl), and TBDPS (t-butyldiphenyl silyl), or a salt thereof.

39. The compound of claim 37, wherein:

R₁ is trifluoromethyl, hydroxymethyl, monofluoromethyl, or triphenylmethoxymethyl; or a salt thereof.

40. The compound of claim 37, wherein:


R₁ is trifluoromethyl, hydroxymethyl, monofluoromethyl, or triphenylmethoxymethyl;

R₂ is fluoro; and

R₃ is hydrogen or fluoro,


or a salt thereof.

41. The compound of claim 37, wherein said compound is selected from the group consisting of:



or a salt thereof.

42. A compound of **Formula VI**:



wherein:

R₁ is C₃₋₄alkyl, halo-C₁₋₄alkyl, hydroxy-C₁₋₄alkyl, or -C₁₋₄alkyl-OR', wherein R' is an oxygen protecting group; and


R_2 and R_3 are each independently hydrogen or halo;
 subject to the proviso that when one of R_2 or R_3 is *ortho*-fluoro, and the other is
 hydrogen, R_1 is not $-\text{CH}_2\text{F}$,
 or a salt thereof.

43. The compound of claim 42, wherein the oxygen protecting group is selected from
 the group consisting of: Trt (triphenylmethyl), MPM (p-methoxybenzyloxymethyl), TBDMS (t-
 butyldimethylsilyl), and TBDPS (t-butyldiphenyl silyl),
 or a salt thereof.

44. The compound of claim 42, wherein:
 R_1 is trifluoromethyl, hydroxymethyl, monofluoromethyl, or triphenylmethoxymethyl,
 or a salt thereof.


45. The compound of claim 42, wherein:
 R_1 is trifluoromethyl, hydroxymethyl, monofluoromethyl, or triphenylmethoxymethyl;
 R_2 is fluoro; and
 R_3 is hydrogen or fluoro,
 or a salt thereof.

46. The compound of claim 42, wherein said compound is selected from the group
 consisting of:



or a salt thereof.

47. A compound of **Formula VII**:



VII

wherein:

R_1 is C_{1-4} alkyl, halo- C_{1-4} alkyl, hydroxy- C_{1-4} alkyl, or $-C_{1-4}$ alkyl-OR',
wherein R' is an oxygen protecting group; and


R_2 and R_3 are each independently hydrogen or halo,
or a salt thereof.

48. The compound of claim 47, wherein the oxygen protecting group is selected from the group consisting of: Trt (triphenylmethyl), MPM (p-methoxybenzyloxymethyl), TBDMS (t-butyldimethylsilyl), and TBDPS (t-butyldiphenyl silyl),
or a salt thereof.

49. The compound of claim 47, wherein:
 R_1 is methyl, trifluoromethyl, hydroxymethyl, monofluoromethyl, or triphenylmethoxymethyl,
or a salt thereof.


50. The compound of claim 47, wherein:
 R_1 is methyl, trifluoromethyl, hydroxymethyl, monofluoromethyl, or triphenylmethoxymethyl;
 R_2 is fluoro; and
 R_3 is hydrogen or fluoro,
or a salt thereof.

51. The compound of claim 47, wherein said compound is selected from the group consisting of:



or a salt thereof.

52. A method of making a compound of **Formula VII**, or a salt thereof, comprising: reducing a compound of **Formula VI** to form a compound of **Formula VII**, as shown below



wherein:

R_1 is C_{1-4} alkyl, halo- C_{1-4} alkyl, hydroxy- C_{1-4} alkyl, or $-C_{1-4}$ alkyl-OR', wherein R' is an oxygen protecting group; and
 R_2 and R_3 are each independently hydrogen or halo,
 to make said compound of **Formula VII**, or a salt thereof.

53. The compound of claim 52, wherein the oxygen protecting group is selected from the group consisting of: Trt (triphenylmethyl), MPM (p-methoxybenzyloxymethyl), TBDMS (t-butyldimethylsilyl), and TBDPS (t-butyldiphenyl silyl), or a salt thereof.

54. The method of claim 52, wherein:

R_1 is trifluoromethyl, hydroxymethyl, monofluoromethyl, or triphenylmethoxymethyl.

55. The method of claim 52, wherein:

R_1 is trifluoromethyl, hydroxymethyl, monofluoromethyl, or triphenylmethoxymethyl;

R_2 is fluoro; and

R_3 is hydrogen or fluoro.

56. The method of any one of claims 52-55, wherein said reducing of the compound of **Formula VI** to form the compound of **Formula VII** comprises the steps of:

- (a) adding powdered iron to a carbinol solvent to form a mixture thereof;
- (b) adding hydrochloric acid (HCl) to the mixture of step (a);
- (c) heating the mixture of step (b) to a temperature of from 31°C to the boiling point of the mixture of step (b) to form a heated mixture; and

(d) adding the compound of **Formula VI**, or a salt thereof, to the heated mixture of step (c),
to thereby reduce said compound of **Formula VI** to form said compound of **Formula VII**, or a salt thereof.

57. The method of claim 56, wherein said heating of step (c) is to a temperature of from 31°C to 80°C.

58. The method of claim 56, wherein said heating of step (c) is to a temperature of from 31°C to 80°C for from 0 to 4 hours.

59. The method of claim 57 or claim 58, wherein said heating of step (c) is to a temperature of from 40°C to 75°C.

60. The method of claim 57 or claim 58, wherein said heating of step (c) is at a temperature of from 50°C to 65°C.

61. The method of any one of claims 56-60, wherein said method further comprises the step of:


(e) cooling the mixture of step (d) to a temperature of from 0°C to 30°C to form a cooled mixture.

62. The method of claim 61, wherein said cooling is from 0 to 4 hours.

63. The method of any of claims 56-62, wherein the HCl is concentrated hydrochloric acid.

64. The method of any of claims 56-63, wherein the HCl is provided in an amount selected from the group consisting of: an amount of HCl stoichiometric with respect to the amount of the compound of **Formula VI**; and an amount of HCl in molar excess with respect to the amount of the compound of **Formula VI**.

65. A method of making a compound of **Formula IX**




wherein:

R_1 is C_{1-4} alkyl, halo- C_{1-4} alkyl, hydroxy- C_{1-4} alkyl, or $-C_{1-4}$ alkyl-OR', wherein R' is an oxygen protecting group; and

R_2 and R_3 are each independently hydrogen or halo; and

R_4 is selected from the group consisting of C_{1-6} alkyl, halo- C_{1-6} alkyl, C_{1-6} alkoxy, and C_{1-4} alkyl substituted with C_{1-4} alkoxy, or a salt thereof,

comprising reacting a compound **Formula VII** with a pyrazine carboxylic acid of **Formula VIII**:



wherein R_1 , R_2 , R_3 and R_4 are as given above,
to form said compound of **Formula IX**, or a salt thereof.

66. The method of claim 65, wherein said reacting comprises the steps of:

(a) adding thionyl chloride to a solution comprising a compound of **Formula VIII** in a solvent to make a formed mixture; and

(b) adding a compound of **Formula VII** to the formed mixture of step (a),
to form said compound of **Formula IX**, or a salt thereof.

67. The method of claim 66, wherein said solution comprising a compound of **Formula VIII** in a solvent is cooled prior to or during said adding step (a).

68. The method of claim 66, wherein said solution comprising a compound of **Formula VIII** in a solvent is cooled to a temperature of from 0°C to 25°C prior to said adding step (a).

69. The method of any of claims 65-68, wherein the solvent is independently selected from the group consisting of: N,N'-dimethylimidazoline-2-one, toluene, methylene chloride, dimethylformamide, N-methylpyrrolidine, and dimethylacetamide.

70. The method of any of claims 65-68, wherein the solvent is N,N'-dimethylimidazoline-2-one.

71. The method of claim 65, wherein said reacting comprises the steps of:

(a) adding ethyl acetate (EtOAc) to a compound of **Formula VIII** and a compound of **Formula VII** to make a formed mixture; and


(b) adding an alkyl phosphonic acid anhydride to the formed mixture of step (a), to form said compound of **Formula IX**, or a salt thereof.

72. The method of claim 71, wherein said alkyl phosphonic acid anhydride is propane phosphonic acid anhydride.

73. The method of claim 71, wherein said alkyl phosphonic acid anhydride is n-propyl phosphonic acid anhydride.

74. The method of any of claims 71-73, wherein said formed mixture is maintained at a temperature of from 0°C to 35°C during said adding step (a).

75. A method of making a compound of **Formula II**



II

wherein:

R_1 is C_{1-4} alkyl; and


R_2 and R_3 are each independently hydrogen or halo;

or a salt thereof,

comprising the steps of:

(a) heating a mixture comprising:

(i) a ketone of **Formula B**:



B

wherein R_1 , R_2 and R_3 are as given above for **Formula II**;

(ii) hydroxylamine or a salt thereof;

(iii) a base; and

(iv) a solvent selected from: water and an alcohol,

to at least 90° Celsius produce a heated mixture,

(b) cooling the heated mixture to a temperature of at or below 60° Celsius, and then,

(c) adding an organic solvent, an acid, and zinc to the mixture, wherein said mixture is maintained at or below 60° Celsius during said adding,

to make said compound of **Formula II**.

76. The method of claim 75, wherein said heating is to a temperature of at least 100 ° Celsius.

77. The method of claim 75, wherein said heating comprises boiling.


78. The method of claim 75, wherein said heating comprises refluxing.

79. The method of claim 75, wherein said heating to at least 90° Celsius is carried out for at least 30 hours.

80. The method of any of claims 75-79, wherein said solvent is a C_{3-8} alcohol.

81. The method of any of claims 75-79, wherein said solvent is water.
82. The method of any of claims 75-81, wherein said cooling is to a temperature of at or below 50° Celsius.
83. The method of any of claims 75-81, wherein said cooling is to a temperature of at or below 45° Celsius.
84. The method of any of claims 75-83, wherein said mixture is maintained at or below 50° Celsius during said adding.
85. The method of any of claims 75-83, wherein said mixture is maintained at or below 45° Celsius during said adding.
86. The method of any of claims 75-83, wherein said mixture is maintained between 25 and 50° Celsius during said adding.
87. The method of any of claims 75-83, wherein said mixture is maintained between 30 and 45° Celsius during said adding.
88. The method of any of claims 75-83, wherein said mixture is maintained between 35 and 45° Celsius during said adding.
89. The method of any of claims 75-88, wherein the base is selected from the group consisting of: acetate, propionate, a tertiary amine base, pyridine, or a salt thereof.
90. The method of claim 89, wherein the tertiary amine base is triethylamine or diethylisopropylamine.
91. The method of any of claims 75-88, wherein the base is acetate.
92. The method of any of claims 75-91, wherein the organic solvent is tetrahydrofuran or an ether solvent.

93. The method of claim 92, wherein the ether solvent is diethyl ether or methyl *tert*-butyl ether.
94. The method of any of claims 75-93, wherein the acid is a carboxylic acid or a mineral acid.
95. The method of claim 94, wherein the mineral acid is hydrochloric acid or sulfuric acid.
96. The method of any of claims 75-93, wherein the acid is acetic acid.
97. The method of any of claims 75-96, wherein the zinc comprises zinc powder or zinc dust.
98. A method of making a compound of **Formula I**,




wherein:

- R_1 is C_{1-4} alkyl, halo- C_{1-4} alkyl, hydroxy- C_{1-4} alkyl, or $-C_{1-4}$ alkyl-OR', wherein R' is an oxygen protecting group; and
- R_2 and R_3 are each independently hydrogen or halo, or a salt thereof,

said method comprising heating a mixture comprising:

- (i) a compound of **Formula B**:



- wherein R_1 , R_2 and R_3 are as given above;
- (ii) acetate;

(iii) hydroxylamine or a salt thereof; and

(iv) a hydrocarbon solvent,

to a temperature of from 90°C to 160°C,

to make said compound of **Formula I**, or a salt thereof.

99. The method of claim 98, wherein the hydrocarbon solvent is selected from the group consisting of: benzene, toluene, and xylenes.


100. The method of claim 98, wherein the hydrocarbon solvent is toluene.

101. The method of any of claims 98-100, wherein said heating is to a temperature of from 110°C to 140°C.

102. The method of any of claims 98-100, wherein said heating is to a temperature of from 100°C to 120°C.

103. The method of any of claims 98-100, wherein said heating is to a temperature of from 120°C to 140°C.

104. A method for making a compound of **Formula V**



wherein:


R₁ is C₃₋₄alkyl, halo-C₁₋₄alkyl, hydroxy-C₁₋₄alkyl, or -C₁₋₄alkyl-OR', wherein R' is an oxygen protecting group; and

R₂ and R₃ are each independently hydrogen or halo;
or a salt thereof,

comprising the steps of:

(a) providing a mixture of

(i) a compound of **Formula III**,



III

wherein R₁, R₂ and R₃ are as given above for **Formula V**;

(ii) an amine base; and

(iii) an organic solvent,

wherein said mixture is at a temperature at or below 0° Celsius,

(b) reacting the mixture with trifluoromethanesulfonic anhydride, 4-toluenesulfonyl chloride, methanesulfonyl chloride or methanesulfonyl anhydride, and then,

(c) adding:

(i) an alcohol; and

(ii) a strong base selected from: a hydroxide, an alkoxide, and ammonia,

to make said compound of **Formula V**.

105. The method of claim 104, wherein the amine base is pyridine or a substituted pyridine.

106. The method of claim 105, wherein the substituted pyridine is collidine.

107. The method of any of claims 104-106, wherein the organic solvent is an aromatic hydrocarbon solvent.

108. The method of any of claims 104-106, wherein the organic solvent is toluene, xylenes, benzene, or dichloromethane.

109. The method of any of claims 104-106, wherein the organic solvent is toluene.

110. The method of any of claims 104-109, wherein said reacting step is carried out at a temperature of from -15 to 0° Celsius.

111. The method of any of claims 104-109, wherein said reacting step is carried out at a temperature of from -10 to 0° Celsius.

112. The method of any of claims 104-109, wherein said reacting step is carried out at a temperature of from -5 to 0° Celsius.

113. The method of any of claims 104-112, wherein the alcohol is propanol.

114. The method of any of claims 104-112, wherein the alcohol is 2-propanol.

115. The method of any of claims 104-114, wherein the strong base is a hydroxide.

116. The method of any of claims 104-114, wherein the strong base is a hydroxide selected from sodium hydroxide, potassium hydroxide, lithium hydroxide and calcium hydroxide.

117. The method of any of claims 104-114, wherein the strong base is an alkoxide.

118. The method of any of claims 104-114, wherein the strong base is an alkoxide selected from methoxide, ethoxide and propoxide.

1/3




FIGURE 1A




FIGURE 1B







FIGURE 2




I




II




III




IV



V



VI



VII