

(19) **DANMARK**

(10) **DK/EP 2588604 T3**



(12) **Oversættelse af
europæisk patentskrift**

Patent- og
Varemærkestyrelsen

-
- (51) Int.Cl.: **C 12 N 9/42 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2016-09-26**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2016-06-29**
- (86) Europæisk ansøgning nr.: **11736227.7**
- (86) Europæisk indleveringsdag: **2011-06-30**
- (87) Den europæiske ansøgnings publiceringsdag: **2013-05-08**
- (86) International ansøgning nr.: **US2011042682**
- (87) Internationalt publikationsnr.: **WO2012003379**
- (30) Prioritet: **2010-06-30 US 360351 P**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **Novozymes, Inc., 1445 Drew Avenue, Davis, CA 95616, USA**
- (72) Opfinder: **MORANT, Marc, Oesestads Boulevard 61D 7th, 2300 København S, Danmark**
PATKAR, Shamkant, Christoffers Alle 91, 2800 Kongens Lyngby, Danmark
DING, Hanshu, 2151 Caravaggion Drive, Davis, California 95616, USA
- (74) Fuldmægtig i Danmark: **Novozymes A/S, Krogshøjvej 36, 2880 Bagsværd, Danmark**
- (54) Benævnelse: **Polypeptider med beta-glucosidaseaktivitet og polynukleotider, der koder for dem**
- (56) Fremdragne publikationer:
WO-A1-2011/057140
KRISTIAN B R M KROGH ET AL: "Characterization and kinetic analysis of a thermostable GH3 beta-glucosidase from Penicillium brasilianum", APPLIED MICROBIOLOGY AND BIOTECHNOLOGY, SPRINGER, BERLIN, DE, vol. 86, no. 1, 16 September 2009 (2009-09-16), pages 143-154, XP019799835, ISSN: 1432-0614 -& DATABASE UniProt [Online] 29 May 2007 (2007-05-29), "Beta-glucosidase from Penicillium brasilianum;", XP000002659341, retrieved from EBI accession no. UNIPROT:A5A4M8 Database accession no. A5A4M8
COPA-PATINO J ET AL: "Purification and properties of a beta-glucosidase from Penicillium oxalicum autolysates", FEMS MICROBIOLOGY LETTERS, BLACKWELL PUBLISHING, AMSTERDAM, NL, vol. 67, no. 1-2, 15 January 1990 (1990-01-15), pages 191-196, XP023918497, ISSN: 0378-1097

DESCRIPTION

Statement as to Rights to Inventions Made Under Federally Sponsored Research and Development

[0001] This invention was made with Government support under Cooperative Agreement DE-FC36-08GO18080 awarded by the Department of Energy. The government has certain rights in this invention.

Reference to a Sequence Listing

[0002] This application contains a Sequence Listing in computer readable form.

Reference to a Deposit of Biological Material

[0003] This application contains a reference to a deposit of biological material.

Background of the Invention

Field of the Invention

[0004] The present invention relates to polypeptides having beta-glucosidase activity and polynucleotides encoding the polypeptides. The invention also relates to nucleic acid constructs, vectors, and host cells comprising the polynucleotides as well as methods of producing and using the polypeptides.

Description of the Related Art

[0005] Cellulose is a polymer of glucose linked by beta-1,4-bonds. Many microorganisms produce enzymes that hydrolyze beta-linked glucans. These enzymes include endoglucanases, cellobiohydrolases, and beta-glucosidases. Endoglucanases digest the cellulose polymer at random locations, opening it to attack by cellobiohydrolases. Cellobiohydrolases sequentially release molecules of cellobiose from the ends of the cellulose polymer. Cellobiose is a water-soluble beta-1,4-linked dimer of glucose. Beta-glucosidases hydrolyze cellobiose to glucose.

[0006] The conversion of lignocellulosic feedstocks into ethanol has the advantages of the ready availability of large amounts of feedstock, the desirability of avoiding burning or land filling the materials, and the cleanliness of the ethanol fuel. Wood, agricultural residues, herbaceous crops, and municipal solid wastes have been considered as feedstocks for ethanol production. These materials primarily consist of cellulose, hemicellulose, and lignin. Once the cellulose is converted to glucose, the glucose is easily fermented by yeast into ethanol. Since glucose is readily fermented to ethanol by a variety of yeasts while cellobiose is not, any cellobiose remaining at the end of the hydrolysis represents a loss of yield of ethanol. More importantly, cellobiose is a potent inhibitor of endoglucanases and cellobiohydrolases. The accumulation of cellobiose during hydrolysis is undesirable for ethanol production.

[0007] Cellobiose accumulation has been a major problem in enzymatic hydrolysis because cellulase-producing microorganisms may produce little beta-glucosidase. The low amount of beta-glucosidase results in a shortage of capacity to hydrolyze the cellobiose to glucose. Several approaches have been used to increase the amount of beta-glucosidase in cellulose conversion.

[0008] One approach is to produce beta-glucosidase using microorganisms that produce little cellulase, and add the beta-glucosidase exogenously to endoglucanase and cellobiohydrolase to enhance the hydrolysis.

[0009] A second approach is to carry out cellulose hydrolysis simultaneously with fermentation. This process is known as simultaneous saccharification and fermentation (SSF). In an SSF system, fermentation of the glucose removes it from solution.

[0010] A third approach to overcome the shortage of beta-glucosidase is to overexpress the beta-glucosidase in a host, thereby increasing the yield of beta-glucosidase.

[0011] It would be an advantage in the art to provide new beta-glucosidases for degrading cellulosic materials.

[0012] The present invention provides polypeptides having beta-glucosidase activity and polynucleotides encoding the polypeptides.

Summary of the Invention

[0013] The present invention relates to isolated polypeptides having beta-glucosidase activity selected from the group consisting of:

1. (a) a polypeptide having at least 95% sequence identity to the mature polypeptide of SEQ ID NO: 2;
2. (b) a variant of the mature polypeptide of SEQ ID NO: 2 comprising a substitution, deletion, and/or insertion at one or more (e.g., several) positions, wherein said variant has at least 85%, e.g. at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% sequence identity to the mature polypeptide of SEQ ID NO: 2 and wherein said variant has improved thermal stability over the mature polypeptide of SEQ ID NO:2.

[0014] The present invention also relates to isolated polynucleotides encoding the polypeptides of the present invention, nucleic acid constructs, recombinant expression vectors, and recombinant host cells comprising the polynucleotides, and methods of producing the polypeptides.

[0015] The present invention also relates to methods of using the polypeptides of the present invention.

[0016] The present disclosure also relates to a polynucleotide encoding a signal peptide comprising or consisting of amino acids 1 to 21 of SEQ ID NO: 2, which is operably linked to a gene encoding a protein; to nucleic acid constructs, expression vectors, and recombinant host cells comprising the polynucleotide; and to methods of producing a protein.

Brief Description of the Figures

[0017]

Figures 1A and 1B show the genomic DNA sequence (SEQ ID NO: 1) and the deduced amino acid sequence (SEQ ID NO: 2) of a *Penicillium oxalicum* strain IBT5387 GH3 beta-glucosidase gene. Introns are italicized.

Figure 2 shows a restriction map of pUC19 D55EX.

Definitions

[0018] Beta-glucosidase: The term "beta-glucosidase" means a beta-D-glucoside glucohydrolase (E.C. 3.2.1.21) that catalyzes the hydrolysis of terminal non-reducing beta-D-glucose residues with the release of beta-D-glucose. For purposes of the present invention, beta-glucosidase activity is determined according to the basic procedure described by Venturi et al., 2002, Extracellular beta-D-glucosidase from *Chaetomium thermophilum* var. *coprophilum*: production, purification and some biochemical properties, J. Basic Microbiol. 42: 55-66. One unit of beta-glucosidase is defined as 1.0 μ mole of p-nitrophenol produced per minute at 40°C, pH 5 from 0.8 mM p-nitrophenyl-beta-D-glucopyranoside as substrate in 50 mM sodium acetate containing 0.01% TWEEN® 20 (polyoxyethylene sorbitan monolaurate).

[0019] The polypeptides of the present invention have at least 20%, e.g., at least 40%, at least 50%, at least 60%, at least 70%,

at least 80%, at least 90%, at least 95%, or at least 100% of the beta-glucosidase activity of the mature polypeptide of SEQ ID NO: 2.

[0020] Acetylxyylan esterase: The term "acetylxyylan esterase" means a carboxylesterase (EC 3.1.1.72) that catalyzes the hydrolysis of acetyl groups from polymeric xylan, acetylated xylose, acetylated glucose, alpha-naphthyl acetate, and p-nitrophenyl acetate. For purposes of the present invention, acetylxyylan esterase activity is determined using 0.5 mM p-nitrophenylacetate as substrate in 50 mM sodium acetate pH 5.0 containing 0.01% TWEEN™ 20. One unit of acetylxyylan esterase is defined as the amount of enzyme capable of releasing 1 μ mole of p-nitrophenolate anion per minute at pH 5, 25°C.

[0021] Allelic variant: The term "allelic variant" means any of two or more alternative forms of a gene occupying the same chromosomal locus. Allelic variation arises naturally through mutation, and may result in polymorphism within populations. Gene mutations can be silent (no change in the encoded polypeptide) or may encode polypeptides having altered amino acid sequences. An allelic variant of a polypeptide is a polypeptide encoded by an allelic variant of a gene.

[0022] Alpha-L-arabinofuranosidase: The term "alpha-L-arabinofuranosidase" means an alpha-L-arabinofuranoside arabinofuranohydrolase (EC 3.2.1.55) that catalyzes the hydrolysis of terminal non-reducing alpha-L-arabinofuranoside residues in alpha-L-arabinosides. The enzyme acts on alpha-L-arabinofuranosides, alpha-L-arabinans containing (1,3)- and/or (1,5)-linkages, arabinoxylans, and arabinogalactans. Alpha-L-arabinofuranosidase is also known as arabinosidase, alpha-arabinosidase, alpha-L-arabinosidase, alpha-arabinofuranosidase, polysaccharide alpha-L-arabinofuranosidase, alpha-L-arabinofuranoside hydrolase, L-arabinosidase, or alpha-L-arabinanase. For purposes of the present invention, alpha-L-arabinofuranosidase activity is determined using 5 mg of medium viscosity wheat arabinoxylan (Megazyme International Ireland, Ltd., Bray, Co. Wicklow, Ireland) per ml of 100 mM sodium acetate pH 5 in a total volume of 200 μ l for 30 minutes at 40°C followed by arabinose analysis by AMINEX® HPX-87H column chromatography (Bio-Rad Laboratories, Inc., Hercules, CA, USA).

[0023] Alpha-glucuronidase: The term "alpha-glucuronidase" means an alpha-D-glucuronidate glucuronohydrolase (EC 3.2.1.139) that catalyzes the hydrolysis of an alpha-D-glucuronoside to D-glucuronate and an alcohol. For purposes of the present invention, alpha-glucuronidase activity is determined according to de Vries, 1998, J. Bacteriol. 180: 243-249. One unit of alpha-glucuronidase equals the amount of enzyme capable of releasing 1 μ mole of glucuronic or 4-O-methylglucuronic acid per minute at pH 5, 40°C.

[0024] Beta-xylosidase: The term "beta-xylosidase" means a beta-D-xyloside xylohydrolase (E.C. 3.2.1.37) that catalyzes the exo-hydrolysis of short beta (1→4)-xylooligosaccharides, to remove successive D-xylose residues from non-reducing termini. For purposes of the present invention, one unit of beta-xylosidase is defined as 1.0 μ mole of p-nitrophenolate anion produced per minute at 40°C, pH 5 from 1 mM p-nitrophenyl-beta-D-xyloside as substrate in 100 mM sodium citrate containing 0.01% TWEEN® 20.

[0025] cDNA: The term "cDNA" means a DNA molecule that can be prepared by reverse transcription from a mature, spliced, mRNA molecule obtained from a eukaryotic or prokaryotic cell. cDNA lacks intron sequences that may be present in the corresponding genomic DNA. The initial, primary RNA transcript is a precursor to mRNA that is processed through a series of steps, including splicing, before appearing as mature spliced mRNA.

[0026] Cellobiohydrolase: The term "cellobiohydrolase" means a 1,4-beta-D-glucan cellobiohydrolase (E.C. 3.2.1.91) that catalyzes the hydrolysis of 1,4-beta-D-glucosidic linkages in cellulose, cellobiosaccharides, or any beta-1,4-linked glucose containing polymer, releasing cellobiose from the reducing or non-reducing ends of the chain (Teeri, 1997, Crystalline cellulose degradation: New insight into the function of cellobiohydrolases, Trends in Biotechnology 15: 160-167; Teeri et al., 1998, Trichoderma reesei cellobiohydrolases: why so efficient on crystalline cellulose?, Biochem. Soc. Trans. 26: 173-178). For purposes of the present invention, cellobiohydrolase activity is determined according to the procedures described by Lever et al., 1972, Anal. Biochem. 47: 273-279; van Tilbeurgh et al., 1982, FEBS Letters, 149: 152-156; van Tilbeurgh and Claeysens, 1985, FEBS Letters, 187: 283-288; and Tomme et al., 1988, Eur. J. Biochem. 170: 575-581. In the present invention, the Lever *et al.* method can be employed to assess hydrolysis of cellulose in corn stover, while the methods of van Tilbeurgh *et al.* and Tomme *et al.* can be used to determine the cellobiohydrolase activity on a fluorescent disaccharide derivative, 4-methylumbelliferyl- β -D-lactoside.

[0027] Cellulolytic enhancing activity: The term "cellulolytic enhancing activity" means a biological activity catalyzed by a GH61 polypeptide that enhances the hydrolysis of a cellulosic material by enzyme having cellulolytic activity. For purposes of the present invention, cellulolytic enhancing activity is determined by measuring the increase in reducing sugars or the increase of the total of cellobiose and glucose from the hydrolysis of a cellulosic material by cellulolytic enzyme under the following conditions: 1-50 mg of total protein/g of cellulose in PCS, wherein total protein is comprised of 50-99.5% w/w cellulolytic enzyme protein and

0.5-50% w/w protein of a GH61 polypeptide having cellulolytic enhancing activity for 1-7 days at 50°C compared to a control hydrolysis with equal total protein loading without cellulolytic enhancing activity (1-50 mg of cellulolytic protein/g of cellulose in PCS). In a preferred aspect, a mixture of CELLULAST® 1.5L (Novozymes A/S, Bagsværd, Denmark) in the presence of 2-3% of total protein weight *Aspergillus oryzae* beta-glucosidase (recombinantly produced in *Aspergillus oryzae* according to WO 02/095014) or 2-3% of total protein weight *Aspergillus fumigatus* beta-glucosidase (recombinantly produced in *Aspergillus oryzae* as described in WO 2002/095014) of cellulase protein loading is used as the source of the cellulolytic activity.

[0028] The GH61 polypeptides having cellulolytic enhancing activity enhance the hydrolysis of a cellulosic material catalyzed by enzyme having cellulolytic activity by reducing the amount of cellulolytic enzyme required to reach the same degree of hydrolysis preferably at least 1.01-fold, more preferably at least 1.05-fold, more preferably at least 1.10-fold, more preferably at least 1.25-fold, more preferably at least 1.5-fold, more preferably at least 2-fold, more preferably at least 3-fold, more preferably at least 4-fold, more preferably at least 5-fold, even more preferably at least 10-fold, and most preferably at least 20-fold.

[0029] Cellulolytic enzyme or cellulase: The term "cellulolytic enzyme" or "cellulase" means one or more (e.g., several) enzymes that hydrolyze a cellulosic material. Such enzymes include endoglucanase(s), cellobiohydrolase(s), beta-glucosidase(s), or combinations thereof. The two basic approaches for measuring cellulolytic activity include: (1) measuring the total cellulolytic activity, and (2) measuring the individual cellulolytic activities (endoglucanases, cellobiohydrolases, and beta-glucosidases) as reviewed in Zhang et al., Outlook for cellulase improvement: Screening and selection strategies, 2006, Biotechnology Advances 24: 452-481. Total cellulolytic activity is usually measured using insoluble substrates, including Whatman N°1 filter paper, microcrystalline cellulose, bacterial cellulose, algal cellulose, cotton, pretreated lignocellulose, etc. The most common total cellulolytic activity assay is the filter paper assay using Whatman N°1 filter paper as the substrate. The assay was established by the International Union of Pure and Applied Chemistry (IUPAC) (Ghose, 1987, Measurement of cellulase activities, Pure Appl. Chem. 59: 257-68).

[0030] For purposes of the present invention, cellulolytic enzyme activity is determined by measuring the increase in hydrolysis of a cellulosic material by cellulolytic enzyme(s) under the following conditions: 1-20 mg of cellulolytic enzyme protein/g of cellulose in PCS for 3-7 days at 50°C compared to a control hydrolysis without addition of cellulolytic enzyme protein. Typical conditions are 1 ml reactions, washed or unwashed PCS, 5% insoluble solids, 50 mM sodium acetate pH 5, 1 mM MnSO₄, 50-65°C, 72 hours, sugar analysis by AMINEX® HPX-87H column (Bio-Rad Laboratories, Inc., Hercules, CA, USA).

[0031] Cellulosic material: The term "cellulosic material" means any material containing cellulose. The predominant polysaccharide in the primary cell wall of biomass is cellulose, the second most abundant is hemicellulose, and the third is pectin. The secondary cell wall, produced after the cell has stopped growing, also contains polysaccharides and is strengthened by polymeric lignin covalently cross-linked to hemicellulose. Cellulose is a homopolymer of anhydrocellobiose and thus a linear beta-(1-4)-D-glucan, while hemicelluloses include a variety of compounds, such as xylans, xyloglucans, arabinoxylans, and mannans in complex branched structures with a spectrum of substituents. Although generally polymorphous, cellulose is found in plant tissue primarily as an insoluble crystalline matrix of parallel glucan chains. Hemicelluloses usually hydrogen bond to cellulose, as well as to other hemicelluloses, which help stabilize the cell wall matrix.

[0032] Cellulose is generally found, for example, in the stems, leaves, hulls, husks, and cobs of plants or leaves, branches, and wood of trees. The cellulosic material can be, but is not limited to, agricultural residue, herbaceous material (including energy crops), municipal solid waste, pulp and paper mill residue, waste paper, and wood (including forestry residue) (see, for example, Wiseloge et al., 1995, in Handbook on Bioethanol (Charles E. Wyman, editor), pp.105-118, Taylor & Francis, Washington D.C.; Wyman, 1994, Bioresource Technology 50: 3-16; Lynd, 1990, Applied Biochemistry and Biotechnology 24/25: 695-719; Mosier et al., 1999, Recent Progress in Bioconversion of Lignocellulosics, in Advances in Biochemical Engineering/Biotechnology, T. Scheper, managing editor, Volume 65, pp.23-40, Springer-Verlag, New York). It is understood herein that the cellulose may be in the form of lignocellulose, a plant cell wall material containing lignin, cellulose, and hemicellulose in a mixed matrix. In a preferred aspect, the cellulosic material is any biomass material. In another preferred aspect, the cellulosic material is lignocellulose, which comprises cellulose, hemicelluloses, and lignin.

[0033] In one aspect, the cellulosic material is agricultural residue. In another aspect, the cellulosic material is herbaceous material (including energy crops). In another aspect, the cellulosic material is municipal solid waste. In another aspect, the cellulosic material is pulp and paper mill residue. In another aspect, the cellulosic material is waste paper. In another aspect, the cellulosic material is wood (including forestry residue).

[0034] In another aspect, the cellulosic material is arundo. In another aspect, the cellulosic material is bagasse. In another aspect, the cellulosic material is bamboo. In another aspect, the cellulosic material is corn cob. In another aspect, the cellulosic

material is corn fiber. In another aspect, the cellulosic material is corn stover. In another aspect, the cellulosic material is miscanthus. In another aspect, the cellulosic material is orange peel. In another aspect, the cellulosic material is rice straw. In another aspect, the cellulosic material is switchgrass. In another aspect, the cellulosic material is wheat straw.

[0035] In another aspect, the cellulosic material is aspen. In another aspect, the cellulosic material is eucalyptus. In another aspect, the cellulosic material is fir. In another aspect, the cellulosic material is pine. In another aspect, the cellulosic material is poplar. In another aspect, the cellulosic material is spruce. In another aspect, the cellulosic material is willow.

[0036] In another aspect, the cellulosic material is algal cellulose. In another aspect, the cellulosic material is bacterial cellulose. In another aspect, the cellulosic material is cotton linter. In another aspect, the cellulosic material is filter paper. In another aspect, the cellulosic material is microcrystalline cellulose. In another aspect, the cellulosic material is phosphoric-acid treated cellulose.

[0037] In another aspect, the cellulosic material is an aquatic biomass. As used herein the term "aquatic biomass" means biomass produced in an aquatic environment by a photosynthesis process. The aquatic biomass can be algae, emergent plants, floating-leaf plants, or submerged plants.

[0038] The cellulosic material may be used as is or may be subjected to pretreatment, using conventional methods known in the art, as described herein. In a preferred aspect, the cellulosic material is pretreated.

[0039] Coding sequence: The term "coding sequence" means a polynucleotide, which directly specifies the amino acid sequence of a polypeptide. The boundaries of the coding sequence are generally determined by an open reading frame, which begins with a start codon such as ATG, GTG, or TTG and ends with a stop codon such as TAA, TAG, or TGA. The coding sequence may be a genomic DNA, cDNA, synthetic DNA, or a combination thereof.

[0040] Control sequences: The term "control sequences" means nucleic acid sequences necessary for expression of a polynucleotide encoding a mature polypeptide of the present invention. Each control sequence may be native (*i.e.*, from the same gene) or foreign (*i.e.*, from a different gene) to the polynucleotide encoding the polypeptide or native or foreign to each other. Such control sequences include, but are not limited to, a leader, polyadenylation sequence, propeptide sequence, promoter, signal peptide sequence, and transcription terminator. At a minimum, the control sequences include a promoter, and transcriptional and translational stop signals. The control sequences may be provided with linkers for the purpose of introducing specific restriction sites facilitating ligation of the control sequences with the coding region of the polynucleotide encoding a polypeptide.

[0041] **Endoglucanase:** The term "endoglucanase" means an endo-1,4-(1,3;1,4)-beta-D-glucan 4-glucanohydrolase (E.C. 3.2.1.4) that catalyzes endohydrolysis of 1,4-beta-D-glycosidic linkages in cellulose, cellulose derivatives (such as carboxymethyl cellulose and hydroxyethyl cellulose), lichenin, beta-1,4 bonds in mixed beta-1,3 glucans such as cereal beta-D-glucans or xyloglucans, and other plant material containing cellulosic components. Endoglucanase activity can be determined by measuring reduction in substrate viscosity or increase in reducing ends determined by a reducing sugar assay (Zhang et al., 2006, Biotechnology Advances 24: 452-481). For purposes of the present invention, endoglucanase activity is determined using carboxymethyl cellulose (CMC) as substrate according to the procedure of Ghose, 1987, Pure and Appl. Chem. 59: 257-268, at pH 5, 40°C.

[0042] **Expression:** The term "expression" includes any step involved in the production of a polypeptide including, but not limited to, transcription, post-transcriptional modification, translation, post-translational modification, and secretion.

[0043] **Expression vector:** The term "expression vector" means a linear or circular DNA molecule that comprises a polynucleotide encoding a polypeptide and is operably linked to control sequences that provide for its expression.

[0044] **Family 3 glycoside hydrolase:** The term "Family 3 glycoside hydrolase" or "Family GH3" or "GH3" means a polypeptide falling into the glycoside hydrolase Family 3 according to Henrissat B., 1991, A classification of glycosyl hydrolases based on amino-acid sequence similarities, Biochem. J. 280: 309-316, and Henrissat B., and Bairoch A., 1996, Updating the sequence-based classification of glycosyl hydrolases, Biochem. J. 316: 695-696.

[0045] **Family 61 glycoside hydrolase:** The term "Family 61 glycoside hydrolase" or "Family GH61" or "GH61" means a polypeptide falling into the glycoside hydrolase Family 61 according to Henrissat B., 1991, A classification of glycosyl hydrolases based on amino-acid sequence similarities, Biochem. J. 280: 309-316, and Henrissat B., and Bairoch A., 1996, Updating the sequence-based classification of glycosyl hydrolases, Biochem. J. 316: 695-696. The enzymes in this family were originally classified as a glycoside hydrolase family based on measurement of very weak endo-1,4-beta-D-glucanase activity in one family

member. The structure and mode of action of these enzymes are non-canonical and they cannot be considered as bona fide glycosidases. However, they are kept in the CAZy classification on the basis of their capacity to enhance the breakdown of lignocellulose when used in conjunction with a cellulase or a mixture of cellulases.

[0046] Feruloyl esterase: The term "feruloyl esterase" means a 4-hydroxy-3-methoxycinnamoyl-sugar hydrolase (EC 3.1.1.73) that catalyzes the hydrolysis of 4-hydroxy-3-methoxycinnamoyl (feruloyl) groups from esterified sugar, which is usually arabinose in "natural" substrates, to produce ferulate (4-hydroxy-3-methoxycinnamate). Feruloyl esterase is also known as ferulic acid esterase, hydroxycinnamoyl esterase, FAE-III, cinnamoyl ester hydrolase, FAEA, cinnAE, FAE-I, or FAE-II. For purposes of the present invention, feruloyl esterase activity is determined using 0.5 mM *p*-nitrophenylferulate as substrate in 50 mM sodium acetate pH 5.0. One unit of feruloyl esterase equals the amount of enzyme capable of releasing 1 μ mole of *p*-nitrophenolate anion per minute at pH 5, 25°C.

[0047] Fragment: The term "fragment" means a polypeptide having one or more (*e.g.*, several) amino acids deleted from the amino and/or carboxyl terminus of a mature polypeptide; wherein the fragment has beta-glucosidase activity. In one aspect, a fragment contains at least 730 amino acid residues, *e.g.*, at least 770 amino acid residues or at least 810 amino acid residues.

[0048] Hemicellulolytic enzyme or hemicellulase: The term "hemicellulolytic enzyme" or "hemicellulase" means one or more (*e.g.*, several) enzymes that hydrolyze a hemicellulosic material. See, for example, Shallom, D. and Shoham, Y. Microbial hemicellulases. Current Opinion In Microbiology, 2003, 6(3): 219-228). Hemicellulases are key components in the degradation of plant biomass. Examples of hemicellulases include, but are not limited to, an acetylmannan esterase, an acetylxylan esterase, an arabinanase, an arabinofuranosidase, a coumaric acid esterase, a feruloyl esterase, a galactosidase, a glucuronidase, a glucuronoyl esterase, a mannanase, a mannosidase, a xylanase, and a xylosidase. The substrates of these enzymes, the hemicelluloses, are a heterogeneous group of branched and linear polysaccharides that are bound via hydrogen bonds to the cellulose microfibrils in the plant cell wall, crosslinking them into a robust network. Hemicelluloses are also covalently attached to lignin, forming together with cellulose a highly complex structure. The variable structure and organization of hemicelluloses require the concerted action of many enzymes for its complete degradation. The catalytic modules of hemicellulases are either glycoside hydrolases (GHs) that hydrolyze glycosidic bonds, or carbohydrate esterases (CEs), which hydrolyze ester linkages of acetate or ferulic acid side groups. These catalytic modules, based on homology of their primary sequence, can be assigned into GH and CE families. Some families, with an overall similar fold, can be further grouped into clans, marked alphabetically (*e.g.*, GH-A). A most informative and updated classification of these and other carbohydrate active enzymes is available in the Carbohydrate-Active Enzymes (CAZy) database. Hemicellulolytic enzyme activities can be measured according to Ghose and Bisaria, 1987, Pure & Appl. Chem. 59: 1739-1752.

[0049] Host cell: The term "host cell" means any cell type that is susceptible to transformation, transfection, transduction, or the like with a nucleic acid construct or expression vector comprising a polynucleotide of the present invention. The term "host cell" encompasses any progeny of a parent cell that is not identical to the parent cell due to mutations that occur during replication.

[0050] Isolated: The term "isolated" means a substance in a form or environment that does not occur in nature. Non-limiting examples of isolated substances include (1) any non-naturally occurring substance, (2) any substance including, but not limited to, any enzyme, variant, nucleic acid, protein, peptide or cofactor, that is at least partially removed from one or more or all of the naturally occurring constituents with which it is associated in nature; (3) any substance modified by the hand of man relative to that substance found in nature; or (4) any substance modified by increasing the amount of the substance relative to other components with which it is naturally associated (*e.g.*, multiple copies of a gene encoding the substance; use of a stronger promoter than the promoter naturally associated with the gene encoding the substance). An isolated substance may be present in a fermentation broth sample.

[0051] Mature polypeptide: The term "mature polypeptide" means a polypeptide in its final form following translation and any post-translational modifications, such as N-terminal processing, C-terminal truncation, glycosylation, phosphorylation, etc. In one aspect, the mature polypeptide is amino acids 22 to 883 of SEQ ID NO: 2 based on the SignalP program (Nielsen et al., 1997, Protein Engineering 10: 1-6) that predicts amino acids 1 to 21 of SEQ ID NO: 2 are a signal peptide.

[0052] Mature polypeptide coding sequence: The term "mature polypeptide coding sequence" means a polynucleotide that encodes a mature polypeptide having beta-glucosidase activity. In one aspect, the mature polypeptide coding sequence is nucleotides 64 to 2790 of SEQ ID NO: 1 based on the SignalP program (Nielsen et al., 1997, *supra*) that predicts nucleotides 1 to 63 of SEQ ID NO: 1 encode a signal peptide. In another aspect, the mature polypeptide coding sequence is the cDNA sequence of nucleotides 64 to 2790 of SEQ ID NO: 1.

[0053] Nucleic acid construct: The term "nucleic acid construct" means a nucleic acid molecule, either single- or double-stranded, which is isolated from a naturally occurring gene or is modified to contain segments of nucleic acids in a manner that would not otherwise exist in nature or which is synthetic. The term nucleic acid construct is synonymous with the term "expression cassette" when the nucleic acid construct contains the control sequences required for expression of a coding sequence of the present invention.

[0054] Operably linked: The term "operably linked" means a configuration in which a control sequence is placed at an appropriate position relative to the coding sequence of a polynucleotide such that the control sequence directs expression of the coding sequence.

[0055] Pretreated corn stover: The term "PCS" or "Pretreated Corn Stover" means a cellulosic material derived from corn stover by treatment with heat and dilute sulfuric acid, alkaline pretreatment, or neutral pretreatment.

[0056] Sequence identity: The relatedness between two amino acid sequences or between two nucleotide sequences is described by the parameter "sequence identity".

[0057] For purposes of the present invention, the sequence identity between two amino acid sequences is determined using the Needleman-Wunsch algorithm (Needleman and Wunsch, 1970, J. Mol. Biol. 48: 443-453) as implemented in the Needle program of the EMBOSS package (EMBOSS: The European Molecular Biology Open Software Suite, Rice et al., 2000, Trends Genet. 16: 276-277), preferably version 5.0.0 or later. The parameters used are gap open penalty of 10, gap extension penalty of 0.5, and the EBLOSUM62 (EMBOSS version of BLOSUM62) substitution matrix. The output of Needle labeled "longest identity" (obtained using the -nobrief option) is used as the percent identity and is calculated as follows:

$$(\text{Identical Residues} \times 100) / (\text{Length of Alignment} - \text{Total Number of Gaps in Alignment})$$

[0058] For purposes of the present invention, the sequence identity between two deoxyribonucleotide sequences is determined using the Needleman-Wunsch algorithm (Needleman and Wunsch, 1970, *supra*) as implemented in the Needle program of the EMBOSS package (EMBOSS: The European Molecular Biology Open Software Suite, Rice et al., 2000, *supra*), preferably version 5.0.0 or later. The parameters used are gap open penalty of 10, gap extension penalty of 0.5, and the EDNAFULL (EMBOSS version of NCBI NUC4.4) substitution matrix. The output of Needle labeled "longest identity" (obtained using the -nobrief option) is used as the percent identity and is calculated as follows:

$$(\text{Identical Deoxyribonucleotides} \times 100) / (\text{Length of Alignment} - \text{Total Number of Gaps in Alignment})$$

[0059] Subsequence: The term "subsequence" means a polynucleotide having one or more (*e.g.*, several) nucleotides deleted from the 5' and/or 3' end of a mature polypeptide coding sequence; wherein the subsequence encodes a fragment having beta-glucosidase activity. In one aspect, a subsequence contains at least 2190 nucleotides, *e.g.*, at least 2310 nucleotides or at least 2430 nucleotides.

[0060] Variant: The term "variant" means a polypeptide having beta-glucosidase activity comprising an alteration, *i.e.*, a substitution, insertion, and/or deletion, at one or more (*e.g.*, several) positions. A substitution means replacement of the amino acid occupying a position with a different amino acid; a deletion means removal of the amino acid occupying a position; and an insertion means adding an amino acid adjacent to and immediately following the amino acid occupying a position.

[0061] Xylan degrading activity or xylanolytic activity: The term "xylan degrading activity" or "xylanolytic activity" means a biological activity that hydrolyzes xylan-containing material. The two basic approaches for measuring xylanolytic activity include: (1) measuring the total xylanolytic activity, and (2) measuring the individual xylanolytic activities (*e.g.*, endoxylanases, beta-xylosidases, arabinofuranosidases, alpha-glucuronidases, acetylxylan esterases, feruloyl esterases, and alpha-glucuronoyl esterases). Recent progress in assays of xylanolytic enzymes was summarized in several publications including Biely and Puchard, Recent progress in the assays of xylanolytic enzymes, 2006, Journal of the Science of Food and Agriculture 86(11): 1636-1647; Spanikova and Biely, 2006, Glucuronoyl esterase - Novel carbohydrate esterase produced by Schizophyllum commune, FEBS Letters 580(19): 4597-4601; Herrmann, Vrsanska, Jurickova, Hirsch, Biely, and Kubicek, 1997, The beta-D-xylosidase of Trichoderma reesei is a multifunctional beta-D-xylan xylohydrolase, Biochemical Journal 321: 375-381.

[0062] Total xylan degrading activity can be measured by determining the reducing sugars formed from various types of xylan, including, for example, oat spelt, beechwood, and larchwood xylans, or by photometric determination of dyed xylan fragments released from various covalently dyed xylans. The most common total xylanolytic activity assay is based on production of reducing

sugars from polymeric 4-O-methyl glucuronoxylan as described in Bailey, Biely, Poutanen, 1992, Interlaboratory testing of methods for assay of xylanase activity, Journal of Biotechnology 23(3): 257-270. Xylanase activity can also be determined with 0.2% AZCL-arabinoxylan as substrate in 0.01% TRITON® X-100 (4-(1,1,3,3-tetramethylbutyl)phenyl-polyethylene glycol) and 200 mM sodium phosphate buffer pH 6 at 37°C. One unit of xylanase activity is defined as 1.0 μ mole of azurine produced per minute at 37°C, pH 6 from 0.2% AZCL-arabinoxylan as substrate in 200 mM sodium phosphate pH 6 buffer.

[0063] For purposes of the present invention, xylan degrading activity is determined by measuring the increase in hydrolysis of birchwood xylan (Sigma Chemical Co., Inc., St. Louis, MO, USA) by xylan-degrading enzyme(s) under the following typical conditions: 1 ml reactions, 5 mg/ml substrate (total solids), 5 mg of xylanolytic protein/g of substrate, 50 mM sodium acetate pH 5, 50°C, 24 hours, sugar analysis using p-hydroxybenzoic acid hydrazide (PHBAH) assay as described by Lever, 1972, A new reaction for colorimetric determination of carbohydrates, Anal. Biochem 47: 273-279.

[0064] Xylanase: The term "xylanase" means a 1,4-beta-D-xylan-xylohydrolase (E.C. 3.2.1.8) that catalyzes the endohydrolysis of 1,4-beta-D-xylosidic linkages in xylans. For purposes of the present invention, xylanase activity is determined with 0.2% AZCL-arabinoxylan as substrate in 0.01% TRITON® X-100 and 200 mM sodium phosphate buffer pH 6 at 37°C. One unit of xylanase activity is defined as 1.0 μ mole of azurine produced per minute at 37°C, pH 6 from 0.2% AZCL-arabinoxylan as substrate in 200 mM sodium phosphate pH 6 buffer.

Detailed Description of the Invention

Polypeptides Having Beta-Glucosidase Activity

[0065] The present invention relates to isolated polypeptides having beta-glucosidase activity selected from the group consisting of:

1. (a) a polypeptide having at least 95% sequence identity to the mature polypeptide of SEQ ID NO: 2;
2. (b) a variant of the mature polypeptide of SEQ ID NO: 2 comprising a substitution, deletion, and/or insertion at one or more (e.g., several) positions, wherein said variant has at least 85%, e.g. at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% sequence identity to the mature polypeptide of SEQ ID NO: 2 and wherein said variant has improved thermal stability over the mature polypeptide of SEQ ID NO:2.

[0066] The present invention relates to isolated polypeptides having a sequence identity to the mature polypeptide of SEQ ID NO: 2 of at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%, which have beta-glucosidase activity. In one aspect, the polypeptides differ by no more than 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, or 9, from the mature polypeptide of SEQ ID NO: 2.

[0067] A polypeptide of the present invention preferably comprises or consists of the amino acid sequence of SEQ ID NO: 2 or an allelic variant thereof; or is a fragment thereof having beta-glucosidase activity. In another aspect, the polypeptide comprises or consists of the mature polypeptide of SEQ ID NO: 2. In another preferred aspect, the polypeptide comprises or consists of amino acids 22 to 883 of SEQ ID NO: 2.

[0068] The disclosure also relates to isolated polypeptides having beta-glucosidase activity that are encoded by polynucleotides that hybridize under very low stringency conditions, low stringency conditions, medium stringency conditions, medium-high stringency conditions, high stringency conditions, or very high stringency conditions with (i) the mature polypeptide coding sequence of SEQ ID NO: 1, (ii) the cDNA sequence of the mature polypeptide coding sequence of SEQ ID NO: 1, or (iii) the full-length complement of (i) or (ii) (J. Sambrook, E.F. Fritsch, and T. Maniatis, 1989, Molecular Cloning, A Laboratory Manual, 2d edition, Cold Spring Harbor, New York).

[0069] The polynucleotide of SEQ ID NO: 1 or a subsequence thereof, as well as the polypeptide of SEQ ID NO: 2 or a fragment thereof, may be used to design nucleic acid probes to identify and clone DNA encoding polypeptides having beta-glucosidase activity from strains of different genera or species according to methods well known in the art. In particular, such probes can be used for hybridization with the genomic DNA or cDNA of a cell of interest, following standard Southern blotting procedures, in order to identify and isolate the corresponding gene therein. Such probes can be considerably shorter than the entire sequence,

but should be at least 14, e.g., at least 25, at least 35, or at least 70 nucleotides in length. Preferably, the nucleic acid probe is at least 100 nucleotides in length, e.g., at least 200 nucleotides, at least 300 nucleotides, at least 400 nucleotides, at least 500 nucleotides, at least 600 nucleotides, at least 700 nucleotides, at least 800 nucleotides, or at least 900 nucleotides in length. Both DNA and RNA probes can be used. The probes are typically labeled for detecting the corresponding gene (for example, with ^{32}P , ^3H , ^{35}S , biotin, or avidin). Such probes are encompassed by the present disclosure.

[0070] A genomic DNA or cDNA library prepared from such other strains may be screened for DNA that hybridizes with the probes described above and encodes a polypeptide having beta-glucosidase activity. Genomic or other DNA from such other strains may be separated by agarose or polyacrylamide gel electrophoresis, or other separation techniques. DNA from the libraries or the separated DNA may be transferred to and immobilized on nitrocellulose or other suitable carrier material. In order to identify a clone or DNA that hybridizes with SEQ ID NO: 1 or a subsequence thereof, the carrier material is used in a Southern blot.

[0071] For purposes of the present disclosure, hybridization indicates that the polynucleotide hybridizes to a labeled nucleic acid probe corresponding to the mature polypeptide coding sequence of SEQ ID NO: 1; the cDNA sequence of the mature polypeptide coding sequence of SEQ ID NO: 1; its full-length complement; or a subsequence thereof; under very low to very high stringency conditions. Molecules to which the nucleic acid probe hybridizes under these conditions can be detected using, for example, X-ray film, or any other detection means known in the art.

[0072] In one aspect, the nucleic acid probe is the mature polypeptide coding sequence of SEQ ID NO: 1 or the cDNA sequence thereof. In another aspect, the nucleic acid probe is nucleotides 64 to 2790 of SEQ ID NO: 1 or the cDNA sequence thereof. In another aspect, the nucleic acid probe is a polynucleotide that encodes the polypeptide of SEQ ID NO: 2 or a fragment thereof. In another preferred aspect, the nucleic acid probe is SEQ ID NO: 1 or the cDNA sequence thereof. In another aspect, the nucleic acid probe is the polynucleotide contained in plasmid pUC19 D55EX which is contained in *E. coli* NRRL B-50395, wherein the polynucleotide encodes a polypeptide having beta-glucosidase activity. In another aspect, the nucleic acid probe is the mature polypeptide coding region contained in plasmid pUC19 D55EX which is contained in *E. coli* NRRL B-50395.

[0073] For long probes of at least 100 nucleotides in length, very low to very high stringency conditions are defined as prehybridization and hybridization at 42°C in 5X SSPE, 0.3% SDS, 200 micrograms/ml sheared and denatured salmon sperm DNA, and either 25% formamide for very low and low stringencies, 35% formamide for medium and medium-high stringencies, or 50% formamide for high and very high stringencies, following standard Southern blotting procedures for 12 to 24 hours optimally. The carrier material is finally washed three times each for 15 minutes using 2X SSC, 0.2% SDS at 45°C (very low stringency), at 50°C (low stringency), at 55°C (medium stringency), at 60°C (medium-high stringency), at 65°C (high stringency), and at 70°C (very high stringency).

[0074] For short probes of about 15 nucleotides to about 70 nucleotides in length, stringency conditions are defined as prehybridization and hybridization at about 5°C to about 10°C below the calculated T_m using the calculation according to Bolton and McCarthy (1962, Proc. Natl. Acad. Sci. USA 48:1390) in 0.9 M NaCl, 0.09 M Tris-HCl pH 7.6, 6 mM EDTA, 0.5% NP-40, 1X Denhardt's solution, 1 mM sodium pyrophosphate, 1 mM sodium monobasic phosphate, 0.1 mM ATP, and 0.2 mg of yeast RNA per ml following standard Southern blotting procedures for 12 to 24 hours optimally. The carrier material is finally washed once in 6X SSC plus 0.1% SDS for 15 minutes and twice each for 15 minutes using 6X SSC at 5°C to 10°C below the calculated T_m .

[0075] The present disclosure also relates to isolated polypeptides having beta-glucosidase activity encoded by polynucleotides having a sequence identity to the mature polypeptide coding sequence of SEQ ID NO: 1 or the cDNA sequence thereof, of at least 85%, e.g., at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%.

[0076] The present disclosure also relates to variants of the mature polypeptide of SEQ ID NO: 2 comprising a substitution, deletion, and/or insertion at one or more (e.g., several) positions, or a homologous sequence thereof. Preferably, amino acid changes are of a minor nature, that is conservative amino acid substitutions or insertions that do not significantly affect the folding and/or activity of the protein; small deletions, typically of one to about 30 amino acids; small amino- or carboxyl-terminal extensions, such as an amino-terminal methionine residue; a small linker peptide of up to about 20-25 residues; or a small extension that facilitates purification by changing net charge or another function, such as a poly-histidine tract, an antigenic epitope or a binding domain.

[0077] Examples of conservative substitutions are within the groups of basic amino acids (arginine, lysine and histidine), acidic amino acids (glutamic acid and aspartic acid), polar amino acids (glutamine and asparagine), hydrophobic amino acids (leucine,

isoleucine and valine), aromatic amino acids (phenylalanine, tryptophan and tyrosine), and small amino acids (glycine, alanine, serine, threonine and methionine). Amino acid substitutions that do not generally alter specific activity are known in the art and are described, for example, by H. Neurath and R.L. Hill, 1979, In, *The Proteins*, Academic Press, New York. Common substitutions are Ala/Ser, Val/Ile, Asp/Glu, Thr/Ser, Ala/Gly, Ala/Thr, Ser/Asn, Ala/Val, Ser/Gly, Tyr/Phe, Ala/Pro, Lys/Arg, Asp/Asn, Leu/Ile, Leu/Val, Ala/Glu, and Asp/Gly.

[0078] Alternatively, the amino acid changes are of such a nature that the physico-chemical properties of the polypeptides are altered. For example, amino acid changes may improve the thermal stability of the polypeptide, alter the substrate specificity, change the pH optimum, and the like.

[0079] Essential amino acids in a polypeptide can be identified according to procedures known in the art, such as site-directed mutagenesis or alanine-scanning mutagenesis (Cunningham and Wells, 1989, *Science* 244: 1081-1085). In the latter technique, single alanine mutations are introduced at every residue in the molecule, and the resultant mutant molecules are tested for beta-glucosidase activity to identify amino acid residues that are critical to the activity of the molecule. See also, Hilton et al., 1996, *J. Biol. Chem.* 271: 4699-4708. The active site of the enzyme or other biological interaction can also be determined by physical analysis of structure, as determined by such techniques as nuclear magnetic resonance, crystallography, electron diffraction, or photoaffinity labeling, in conjunction with mutation of putative contact site amino acids. See, for example, de Vos et al., 1992, *Science* 255: 306-312; Smith et al., 1992, *J. Mol. Biol.* 224: 899-904; Wlodaver et al., 1992, *FEBS Lett.* 309: 59-64. The identity of essential amino acids can also be inferred from an alignment with a related polypeptide.

[0080] Single or multiple amino acid substitutions, deletions, and/or insertions can be made and tested using known methods of mutagenesis, recombination, and/or shuffling, followed by a relevant screening procedure, such as those disclosed by Reidhaar-Olson and Sauer, 1988, *Science* 241: 53-57; Bowie and Sauer, 1989, *Proc. Natl. Acad. Sci. USA* 86: 2152-2156; WO 95/17413; or WO 95/22625. Other methods that can be used include error-prone PCR, phage display (e.g., Lowman et al., 1991, *Biochemistry* 30: 10832-10837; U.S. Patent No. 5,223,409; WO 92/06204), and region-directed mutagenesis (Derbyshire et al., 1986, *Gene* 46: 145; Ner et al., 1988, *DNA* 7: 127).

[0081] Mutagenesis/shuffling methods can be combined with high-throughput, automated screening methods to detect activity of cloned, mutagenized polypeptides expressed by host cells (Ness et al., 1999, *Nature Biotechnology* 17: 893-896). Mutagenized DNA molecules that encode active polypeptides can be recovered from the host cells and rapidly sequenced using standard methods in the art. These methods allow the rapid determination of the importance of individual amino acid residues in a polypeptide.

[0082] The total number of amino acid substitutions, deletions and/or insertions of the mature polypeptide of SEQ ID NO: 2 is not more than 10, e.g., 1, 2, 3, 4, 5, 6, 7, 8 or 9.

[0083] The polypeptide may be hybrid polypeptide in which a portion of one polypeptide is fused at the N-terminus or the C-terminus of a portion of another polypeptide.

[0084] The polypeptide may be a fusion polypeptide or cleavable fusion polypeptide in which another polypeptide is fused at the N-terminus or the C-terminus of the polypeptide of the present invention. A fusion polypeptide is produced by fusing a polynucleotide encoding another polypeptide to a polynucleotide of the present invention. Techniques for producing fusion polypeptides are known in the art, and include ligating the coding sequences encoding the polypeptides so that they are in frame and that expression of the fusion polypeptide is under control of the same promoter(s) and terminator. Fusion polypeptides may also be constructed using intein technology in which fusion polypeptides are created post-translationally (Cooper et al., 1993, *EMBO J.* 12: 2575-2583; Dawson et al., 1994, *Science* 266: 776-779).

[0085] A fusion polypeptide can further comprise a cleavage site between the two polypeptides. Upon secretion of the fusion protein, the site is cleaved releasing the two polypeptides. Examples of cleavage sites include, but are not limited to, the sites disclosed in Martin et al., 2003, *J. Ind. Microbiol. Biotechnol.* 3: 568-576; Svetina et al., 2000, *J. Biotechnol.* 76: 245-251; Rasmussen-Wilson et al., 1997, *Appl. Environ. Microbiol.* 63: 3488-3493; Ward et al., 1995, *Biotechnology* 13: 498-503; and Contreras et al., 1991, *Biotechnology* 9: 378-381; Eaton et al., 1986, *Biochemistry* 25: 505-512; Collins-Racie et al., 1995, *Biotechnology* 13: 982-987; Carter et al., 1989, *Proteins: Structure, Function, and Genetics* 6: 240-248; and Stevens, 2003, *Drug Discovery World* 4: 35-48.

Sources of Polypeptides Having Beta-Glucosidase Activity

[0086] A polypeptide having beta-glucosidase activity of the present invention may be obtained from microorganisms of any genus. For purposes of the present invention, the term "obtained from" as used herein in connection with a given source shall mean that the polypeptide encoded by a polynucleotide is produced by the source or by a strain in which the polynucleotide from the source has been inserted. In one aspect, the polypeptide obtained from a given source is secreted extracellularly.

[0087] The polypeptide may be a bacterial polypeptide. For example, the polypeptide may be a Gram-positive bacterial polypeptide such as a *Bacillus*, *Clostridium*, *Enterococcus*, *Geobacillus*, *Lactobacillus*, *Lactococcus*, *Oceanobacillus*, *Staphylococcus*, *Streptococcus*, or *Streptomyces* polypeptide having beta-glucosidase activity, or a Gram-negative bacterial polypeptide such as a *Campylobacter*, *E. coli*, *Flavobacterium*, *Fusobacterium*, *Helicobacter*, *Ilyobacter*, *Neisseria*, *Pseudomonas*, *Salmonella*, or *Ureaplasma* polypeptide.

[0088] In one aspect, the polypeptide is a *Bacillus alkalophilus*, *Bacillus amyloliquefaciens*, *Bacillus brevis*, *Bacillus circulans*, *Bacillus clausii*, *Bacillus coagulans*, *Bacillus firmus*, *Bacillus lautus*, *Bacillus lentus*, *Bacillus licheniformis*, *Bacillus megaterium*, *Bacillus pumilus*, *Bacillus stearothermophilus*, *Bacillus subtilis*, or *Bacillus thuringiensis* polypeptide.

[0089] In another aspect, the polypeptide is a *Streptococcus equisimilis*, *Streptococcus pyogenes*, *Streptococcus uberis*, or *Streptococcus equi* subsp. *Zooepidemicus* polypeptide.

[0090] In another aspect, the polypeptide is a *Streptomyces achromogenes*, *Streptomyces avermitilis*, *Streptomyces coelicolor*, *Streptomyces griseus*, or *Streptomyces lividans* polypeptide.

[0091] The polypeptide may also be a fungal polypeptide. For example, the polypeptide may be a yeast polypeptide such as a *Candida*, *Kluyveromyces*, *Pichia*, *Saccharomyces*, *Schizosaccharomyces*, or *Yarrowia* polypeptide; or a filamentous fungal polypeptide such as an *Acremonium*, *Agaricus*, *Alternaria*, *Aspergillus*, *Aureobasidium*, *Botryosphaeria*, *Ceriporiopsis*, *Chaetomidium*, *Chrysosporium*, *Claviceps*, *Cochliobolus*, *Coprinopsis*, *Coptotermes*, *Corynascus*, *Cryphonectria*, *Cryptococcus*, *Diplodia*, *Exidia*, *Filibasidium*, *Fusarium*, *Gibberella*, *Holomastigotoidea*, *Humicola*, *Irpex*, *Lentinula*, *Leptosphaeria*, *Magnaporthe*, *Melanocarpus*, *Meripilus*, *Mucor*, *Myceliophthora*, *Neocallimastix*, *Neurospora*, *Paecilomyces*, *Penicillium*, *Phanerochaete*, *Piromyces*, *Poitrasia*, *Pseudoplectania*, *Pseudotrichonympha*, *Rhizomucor*, *Schizophyllum*, *Scytalidium*, *Talaromyces*, *Thermoascus*, *Thielavia*, *Tolypocladium*, *Trichoderma*, *Trichophaea*, *Verticillium*, *Volvariella*, or *Xylaria* polypeptide.

[0092] In another aspect, the polypeptide is a *Saccharomyces carlsbergensis*, *Saccharomyces cerevisiae*, *Saccharomyces diastaticus*, *Saccharomyces douglasii*, *Saccharomyces kluyveri*, *Saccharomyces norbensis*, or *Saccharomyces oviformis* polypeptide.

[0093] In another aspect, the polypeptide is an *Acremonium cellulolyticus*, *Aspergillus aculeatus*, *Aspergillus awamori*, *Aspergillus foetidus*, *Aspergillus fumigatus*, *Aspergillus japonicus*, *Aspergillus nidulans*, *Aspergillus niger*, *Aspergillus oryzae*, *Chrysosporium inops*, *Chrysosporium keratinophilum*, *Chrysosporium lucknowense*, *Chrysosporium merdarium*, *Chrysosporium pannicola*, *Chrysosporium queenslandicum*, *Chrysosporium tropicum*, *Chrysosporium zonatum*, *Fusarium bacridioides*, *Fusarium cerealis*, *Fusarium crookwellense*, *Fusarium culmorum*, *Fusarium graminearum*, *Fusarium graminum*, *Fusarium heterosporum*, *Fusarium negundi*, *Fusarium oxysporum*, *Fusarium reticulatum*, *Fusarium roseum*, *Fusarium sambucinum*, *Fusarium sarcochroum*, *Fusarium sporotrichioides*, *Fusarium sulphureum*, *Fusarium torulosum*, *Fusarium trichothecioides*, *Fusarium venenatum*, *Humicola grisea*, *Humicola insolens*, *Humicola lanuginosa*, *Irpex lacteus*, *Mucor miehei*, *Myceliophthora thermophila*, *Neurospora crassa*, *Penicillium funiculosum*, *Penicillium purpurogenum*, *Phanerochaete chrysosporium*, *Thielavia achromatica*, *Thielavia albomyces*, *Thielavia albopilosa*, *Thielavia australeinsis*, *Thielavia fimeti*, *Thielavia microspora*, *Thielavia ovispora*, *Thielavia peruviana*, *Thielavia setosa*, *Thielavia spededonium*, *Thielavia subthermophila*, *Thielavia terrestris*, *Trichoderma harzianum*, *Trichoderma koningii*, *Trichoderma longibrachiatum*, *Trichoderma reesei*, or *Trichoderma viride* polypeptide.

[0094] In another aspect, the polypeptide is a *Penicillium oxalicum* polypeptide, e.g., a polypeptide obtained from *Penicillium oxalicum* IBT5387.

[0095] It will be understood that for the aforementioned species the invention encompasses both the perfect and imperfect states, and other taxonomic equivalents, e.g., anamorphs, regardless of the species name by which they are known. Those skilled in the art will readily recognize the identity of appropriate equivalents.

[0096] Strains of these species are readily accessible to the public in a number of culture collections, such as the American Type Culture Collection (ATCC), Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH (DSMZ), Centraalbureau Voor Schimmelcultures (CBS), and Agricultural Research Service Patent Culture Collection, Northern Regional Research Center

(NRRL).

[0097] The polypeptide may be identified and obtained from other sources including microorganisms isolated from nature (*e.g.*, soil, composts, water, etc.) or DNA samples obtained directly from natural materials (*e.g.*, soil, composts, water, etc.) using the above-mentioned probes. Techniques for isolating microorganisms and DNA directly from natural habitats are well known in the art. A polynucleotide encoding the polypeptide may then be obtained by similarly screening a genomic DNA or cDNA library of another microorganism or mixed DNA sample. Once a polynucleotide encoding a polypeptide has been detected with the probe(s), the polynucleotide can be isolated or cloned by utilizing techniques that are known to those of ordinary skill in the art (see, *e.g.*, Sambrook *et al.*, 1989, *supra*).

Polynucleotides

[0098] The present invention also relates to isolated polynucleotides encode a polypeptide of the present invention.

[0099] The techniques used to isolate or clone a polynucleotide encoding a polypeptide are known in the art and include isolation from genomic DNA, preparation from cDNA, or a combination thereof. The cloning of the polynucleotides from genomic DNA can be effected, *e.g.*, by using the well known polymerase chain reaction (PCR) or antibody screening of expression libraries to detect cloned DNA fragments with shared structural features. See, *e.g.*, Innis *et al.*, 1990, PCR: A Guide to Methods and Application, Academic Press, New York. Other nucleic acid amplification procedures such as ligase chain reaction (LCR), ligation activated transcription (LAT) and polynucleotide-based amplification (NASBA) may be used. The polynucleotides may be cloned from a strain of *Penicillium*, or another or related organism, and thus, for example, may be an allelic or species variant of the polypeptide encoding region of the polynucleotide.

[0100] The present disclosure relates to isolated polynucleotides comprising or consisting of polynucleotides having a degree of sequence identity to the mature polypeptide coding sequence of SEQ ID NO: 1 or the cDNA sequence thereof of at least 85%, *e.g.*, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%, which encode a polypeptide having beta-glucosidase activity.

[0101] Modification of a polynucleotide encoding a polypeptide of the present invention may be necessary for synthesizing polypeptides substantially similar to the polypeptide. The term "substantially similar" to the polypeptide refers to non-naturally occurring forms of the polypeptide. These polypeptides may differ in some engineered way from the polypeptide isolated from its native source, *e.g.*, variants that differ in specific activity, thermostability, pH optimum, or the like. The variants may be constructed on the basis of the polynucleotide presented as the mature polypeptide coding sequence of SEQ ID NO: 1 or the cDNA sequence thereof, *e.g.*, a subsequence thereof, and/or by introduction of nucleotide substitutions that do not result in a change in the amino acid sequence of the polypeptide, but which correspond to the codon usage of the host organism intended for production of the enzyme, or by introduction of nucleotide substitutions that may give rise to a different amino acid sequence. For a general description of nucleotide substitution, see, *e.g.*, Ford *et al.*, 1991, Protein Expression and Purification 2: 95-107.

[0102] The present disclosure also relates to isolated polynucleotides encoding polypeptides of the present invention, which hybridize under very low stringency conditions, low stringency conditions, medium stringency conditions, medium-high stringency conditions, high stringency conditions, or very high stringency conditions with (i) the mature polypeptide coding sequence of SEQ ID NO: 1, (ii) the cDNA sequence of the mature polypeptide coding sequence of SEQ ID NO: 1, or (iii) the full-length complement of (i) or (ii); or allelic variants and subsequences thereof (Sambrook *et al.*, 1989, *supra*), as defined herein.

[0103] In one aspect, the polynucleotide comprises or consists of SEQ ID NO: 1, the mature polypeptide coding sequence of SEQ ID NO: 1, or the sequence contained in pUC19 D55EX which is contained in *E. coli* NRRL B-50395, or a subsequence of SEQ ID NO: 1 that encodes a fragment of SEQ ID NO: 2 having beta-glucosidase activity, such as the polynucleotide of nucleotides 64 to 2790 of SEQ ID NO: 1.

Nucleic Acid Constructs

[0104] The present invention also relates to nucleic acid constructs comprising a polynucleotide of the present invention operably linked to one or more (*e.g.*, several) control sequences that direct the expression of the coding sequence in a suitable host cell under conditions compatible with the control sequences.

[0105] A polynucleotide may be manipulated in a variety of ways to provide for expression of the polypeptide. Manipulation of the polynucleotide prior to its insertion into a vector may be desirable or necessary depending on the expression vector. The techniques for modifying polynucleotides utilizing recombinant DNA methods are well known in the art.

[0106] The control sequence may be a promoter, a polynucleotide recognized by a host cell for expression of a polynucleotide encoding a polypeptide of the present invention. The promoter contains transcriptional control sequences that mediate the expression of the polypeptide. The promoter may be any polynucleotide that shows transcriptional activity in the host cell including mutant, truncated, and hybrid promoters, and may be obtained from genes encoding extracellular or intracellular polypeptides either homologous or heterologous to the host cell.

[0107] Examples of suitable promoters for directing transcription of the nucleic acid constructs of the present invention in a bacterial host cell are the promoters obtained from the *Bacillus amyloliquefaciens* alpha-amylase gene (*amyQ*), *Bacillus licheniformis* alpha-amylase gene (*amyL*), *Bacillus licheniformis* penicillinase gene (*penP*), *Bacillus stearothermophilus* maltogenic amylase gene (*amyM*), *Bacillus subtilis* levansucrase gene (*sacB*), *Bacillus subtilis* *xylA* and *xylB* genes, *Bacillus thuringiensis* *cryIIIA* gene (Agaisse and Lereclus, 1994, Molecular Microbiology 13: 97-107), *E. coli lac* operon, *E. coli trc* promoter (Egon et al., 1988, Gene 69: 301-315), *Streptomyces coelicolor* agarase gene (*dagA*), and prokaryotic beta-lactamase gene (Villa-Kamaroff et al., 1978, Proc. Natl. Acad. Sci. USA 75: 3727-3731), as well as the *tac* promoter (DeBoer et al., 1983, Proc. Natl. Acad. Sci. USA 80: 21-25). Further promoters are described in "Useful proteins from recombinant bacteria" in Gilbert et al., 1980, Scientific American 242: 74-94; and in Sambrook et al., 1989, *supra*. Examples of tandem promoters are disclosed in WO 99/43835.

[0108] Examples of suitable promoters for directing transcription of the nucleic acid constructs of the present invention in a filamentous fungal host cell are promoters obtained from the genes for *Aspergillus nidulans* acetamidase, *Aspergillus niger* neutral alpha-amylase, *Aspergillus niger* acid stable alpha-amylase, *Aspergillus niger* or *Aspergillus awamori* glucoamylase (*glaA*), *Aspergillus oryzae* TAKA amylase, *Aspergillus oryzae* alkaline protease, *Aspergillus oryzae* triose phosphate isomerase, *Fusarium oxysporum* trypsin-like protease (WO 96/00787), *Fusarium venenatum* amyloglucosidase (WO 00/56900), *Fusarium venenatum* Daria (WO 00/56900), *Fusarium venenatum* Quinn (WO 00/56900), *Rhizomucor miehei* lipase, *Rhizomucor miehei* aspartic proteinase, *Trichoderma reesei* beta-glucosidase, *Trichoderma reesei* cellobiohydrolase I, *Trichoderma reesei* cellobiohydrolase II, *Trichoderma reesei* endoglucanase I, *Trichoderma reesei* endoglucanase II, *Trichoderma reesei* endoglucanase III, *Trichoderma reesei* endoglucanase IV, *Trichoderma reesei* endoglucanase V, *Trichoderma reesei* xylanase I, *Trichoderma reesei* xylanase II, *Trichoderma reesei* beta-xylosidase, as well as the NA2-tpi promoter (a modified promoter from an *Aspergillus* neutral alpha-amylase gene in which the untranslated leader has been replaced by an untranslated leader from an *Aspergillus* triose phosphate isomerase gene; non-limiting examples include modified promoters from an *Aspergillus niger* neutral alpha-amylase gene in which the untranslated leader has been replaced by an untranslated leader from an *Aspergillus nidulans* or *Aspergillus oryzae* triose phosphate isomerase gene); and mutant, truncated, and hybrid promoters thereof.

[0109] In a yeast host, useful promoters are obtained from the genes for *Saccharomyces cerevisiae* enolase (ENO-1), *Saccharomyces cerevisiae* galactokinase (GAL1), *Saccharomyces cerevisiae* alcohol dehydrogenase/glyceraldehyde-3-phosphate dehydrogenase (ADH1, ADH2/GAP), *Saccharomyces cerevisiae* triose phosphate isomerase (TPI), *Saccharomyces cerevisiae* metallothionein (CUP1), and *Saccharomyces cerevisiae* 3-phosphoglycerate kinase. Other useful promoters for yeast host cells are described by Romanos et al., 1992, Yeast 8: 423-488.

[0110] The control sequence may also be a transcription terminator, which is recognized by a host cell to terminate transcription. The terminator is operably linked to the 3'-terminus of the polynucleotide encoding the polypeptide. Any terminator that is functional in the host cell may be used in the present invention.

[0111] Preferred terminators for bacterial host cells are obtained from the genes for *Bacillus clausii* alkaline protease (*aprH*), *Bacillus licheniformis* alpha-amylase (*amyL*), and *Escherichia coli* ribosomal RNA (*rrnB*).

[0112] Preferred terminators for filamentous fungal host cells are obtained from the genes for *Aspergillus nidulans* anthranilate synthase, *Aspergillus niger* glucoamylase, *Aspergillus niger* alpha-glucosidase, *Aspergillus oryzae* TAKA amylase, and *Fusarium oxysporum* trypsin-like protease.

[0113] Preferred terminators for yeast host cells are obtained from the genes for *Saccharomyces cerevisiae* enolase, *Saccharomyces cerevisiae* cytochrome C (CYC1), and *Saccharomyces cerevisiae* glyceraldehyde-3-phosphate dehydrogenase. Other useful terminators for yeast host cells are described by Romanos et al., 1992, *supra*.

[0114] The control sequence may also be an mRNA stabilizer region downstream of a promoter and upstream of the coding

sequence of a gene which increases expression of the gene.

[0115] Examples of suitable mRNA stabilizer regions are obtained from a *Bacillus thuringiensis cryIIIA* gene (WO 94/25612) and a *Bacillus subtilis* SP82 gene (Hue et al., 1995, Journal of Bacteriology 177: 3465-3471).

[0116] The control sequence may also be a leader, a nontranslated region of an mRNA that is important for translation by the host cell. The leader is operably linked to the 5'-terminus of the polynucleotide encoding the polypeptide. Any leader that is functional in the host cell may be used.

[0117] Preferred leaders for filamentous fungal host cells are obtained from the genes for *Aspergillus oryzae* TAKA amylase and *Aspergillus nidulans* triose phosphate isomerase.

[0118] Suitable leaders for yeast host cells are obtained from the genes for *Saccharomyces cerevisiae* enolase (ENO-1), *Saccharomyces cerevisiae* 3-phosphoglycerate kinase, *Saccharomyces cerevisiae* alpha-factor, and *Saccharomyces cerevisiae* alcohol dehydrogenase/glyceraldehyde-3-phosphate dehydrogenase (ADH2/GAP).

[0119] The control sequence may also be a polyadenylation sequence, a sequence operably linked to the 3'-terminus of the polynucleotide and, when transcribed, is recognized by the host cell as a signal to add polyadenosine residues to transcribed mRNA. Any polyadenylation sequence that is functional in the host cell may be used.

[0120] Preferred polyadenylation sequences for filamentous fungal host cells are obtained from the genes for *Aspergillus nidulans* anthranilate synthase, *Aspergillus niger* glucoamylase, *Aspergillus niger* alpha-glucosidase *Aspergillus oryzae* TAKA amylase, and *Fusarium oxysporum* trypsin-like protease.

[0121] Useful polyadenylation sequences for yeast host cells are described by Guo and Sherman, 1995, Mol. Cellular Biol. 15: 5983-5990.

[0122] The control sequence may also be a signal peptide coding region that encodes a signal peptide linked to the N-terminus of a polypeptide and directs the polypeptide into the cell's secretory pathway. The 5'-end of the coding sequence of the polynucleotide may inherently contain a signal peptide coding sequence naturally linked in translation reading frame with the segment of the coding sequence that encodes the polypeptide. Alternatively, the 5'-end of the coding sequence may contain a signal peptide coding sequence that is foreign to the coding sequence. A foreign signal peptide coding sequence may be required where the coding sequence does not naturally contain a signal peptide coding sequence. Alternatively, a foreign signal peptide coding sequence may simply replace the natural signal peptide coding sequence in order to enhance secretion of the polypeptide. However, any signal peptide coding sequence that directs the expressed polypeptide into the secretory pathway of a host cell may be used.

[0123] Effective signal peptide coding sequences for bacterial host cells are the signal peptide coding sequences obtained from the genes for *Bacillus* NCIB 11837 maltogenic amylase, *Bacillus licheniformis* subtilisin, *Bacillus licheniformis* beta-lactamase, *Bacillus stearothermophilus* alpha-amylase, *Bacillus stearothermophilus* neutral proteases (*nprT*, *nprS*, *nprM*), and *Bacillus subtilis* *prsA*. Further signal peptides are described by Simonen and Palva, 1993, Microbiological Reviews 57: 109-137.

[0124] Effective signal peptide coding sequences for filamentous fungal host cells are the signal peptide coding sequences obtained from the genes for *Aspergillus niger* neutral amylase, *Aspergillus niger* glucoamylase, *Aspergillus oryzae* TAKA amylase, *Humicola insolens* cellulase, *Humicola insolens* endoglucanase V, *Humicola lanuginosa* lipase, and *Rhizomucor miehei* aspartic proteinase.

[0125] Useful signal peptides for yeast host cells are obtained from the genes for *Saccharomyces cerevisiae* alpha-factor and *Saccharomyces cerevisiae* invertase. Other useful signal peptide coding sequences are described by Romanos et al., 1992, *supra*.

[0126] The control sequence may also be a propeptide coding sequence that encodes a propeptide positioned at the N-terminus of a polypeptide. The resultant polypeptide is known as a proenzyme or propolypeptide (or a zymogen in some cases). A propolypeptide is generally inactive and can be converted to an active polypeptide by catalytic or autocatalytic cleavage of the propeptide from the propolypeptide. The propeptide coding sequence may be obtained from the genes for *Bacillus subtilis* alkaline protease (*aprE*), *Bacillus subtilis* neutral protease (*nprT*), *Myceliophthora thermophila* laccase (WO 95/33836), *Rhizomucor miehei* aspartic proteinase, and *Saccharomyces cerevisiae* alpha-factor.

[0127] Where both signal peptide and propeptide sequences are present, the propeptide sequence is positioned next to the N-terminus of a polypeptide and the signal peptide sequence is positioned next to the N-terminus of the propeptide sequence.

[0128] It may also be desirable to add regulatory sequences that regulate expression of the polypeptide relative to the growth of the host cell. Examples of regulatory systems are those that cause expression of the gene to be turned on or off in response to a chemical or physical stimulus, including the presence of a regulatory compound. Regulatory systems in prokaryotic systems include the *lac*, *tac*, and *trp* operator systems. In yeast, the ADH2 system or GAL1 system may be used. In filamentous fungi, the *Aspergillus niger* glucoamylase promoter, *Aspergillus oryzae* TAKA alpha-amylase promoter, and *Aspergillus oryzae* glucoamylase promoter may be used. Other examples of regulatory sequences are those that allow for gene amplification. In eukaryotic systems, these regulatory sequences include the dihydrofolate reductase gene that is amplified in the presence of methotrexate, and the metallothionein genes that are amplified with heavy metals. In these cases, the polynucleotide encoding the polypeptide would be operably linked with the regulatory sequence.

Expression Vectors

[0129] The present invention also relates to recombinant expression vectors comprising a polynucleotide of the present invention, a promoter, and transcriptional and translational stop signals. The various nucleotide and control sequences may be joined together to produce a recombinant expression vector that may include one or more (e.g., several) convenient restriction sites to allow for insertion or substitution of the polynucleotide encoding the polypeptide at such sites. Alternatively, the polynucleotide may be expressed by inserting the polynucleotide or a nucleic acid construct comprising the polynucleotide into an appropriate vector for expression. In creating the expression vector, the coding sequence is located in the vector so that the coding sequence is operably linked with the appropriate control sequences for expression.

[0130] The recombinant expression vector may be any vector (e.g., a plasmid or virus) that can be conveniently subjected to recombinant DNA procedures and can bring about expression of the polynucleotide. The choice of the vector will typically depend on the compatibility of the vector with the host cell into which the vector is to be introduced. The vector may be a linear or closed circular plasmid.

[0131] The vector may be an autonomously replicating vector, i.e., a vector that exists as an extrachromosomal entity, the replication of which is independent of chromosomal replication, e.g., a plasmid, an extrachromosomal element, a minichromosome, or an artificial chromosome. The vector may contain any means for assuring self-replication. Alternatively, the vector may be one that, when introduced into the host cell, is integrated into the genome and replicated together with the chromosome(s) into which it has been integrated. Furthermore, a single vector or plasmid or two or more vectors or plasmids that together contain the total DNA to be introduced into the genome of the host cell, or a transposon, may be used.

[0132] The vector preferably contains one or more (e.g., several) selectable markers that permit easy selection of transformed, transfected, transduced, or the like cells. A selectable marker is a gene the product of which provides for biocide or viral resistance, resistance to heavy metals, prototrophy to auxotrophs, and the like.

[0133] Examples of bacterial selectable markers are *Bacillus licheniformis* or *Bacillus subtilis* *dal* genes, or markers that confer antibiotic resistance such as ampicillin, chloramphenicol, kanamycin, neomycin, spectinomycin, or tetracycline resistance. Suitable markers for yeast host cells include, but are not limited to, ADE2, HIS3, LEU2, LYS2, MET3, TRP1, and URA3. Selectable markers for use in a filamentous fungal host cell include, but are not limited to, *amdS* (acetamidase), *argB* (ornithine carbamoyltransferase), *bar* (phosphinothricin acetyltransferase), *hph* (hygromycin phosphotransferase), *niad* (nitrate reductase), *pyrG* (orotidine-5'-phosphate decarboxylase), *sC* (sulfate adenylyltransferase), and *trpC* (anthranilate synthase), as well as equivalents thereof. Preferred for use in an *Aspergillus* cell are *Aspergillus nidulans* or *Aspergillus oryzae* *amdS* and *pyrG* genes and a *Streptomyces hygroscopicus* *bar* gene.

[0134] The vector preferably contains an element(s) that permits integration of the vector into the host cell's genome or autonomous replication of the vector in the cell independent of the genome.

[0135] For integration into the host cell genome, the vector may rely on the polynucleotide's sequence encoding the polypeptide or any other element of the vector for integration into the genome by homologous or non-homologous recombination. Alternatively, the vector may contain additional polynucleotides for directing integration by homologous recombination into the genome of the host cell at a precise location(s) in the chromosome(s). To increase the likelihood of integration at a precise

location, the integrational elements should contain a sufficient number of nucleic acids, such as 100 to 10,000 base pairs, 400 to 10,000 base pairs, and 800 to 10,000 base pairs, which have a high degree of sequence identity to the corresponding target sequence to enhance the probability of homologous recombination. The integrational elements may be any sequence that is homologous with the target sequence in the genome of the host cell. Furthermore, the integrational elements may be non-encoding or encoding polynucleotides. On the other hand, the vector may be integrated into the genome of the host cell by non-homologous recombination.

[0136] For autonomous replication, the vector may further comprise an origin of replication enabling the vector to replicate autonomously in the host cell in question. The origin of replication may be any plasmid replicator mediating autonomous replication that functions in a cell. The term "origin of replication" or "plasmid replicator" means a polynucleotide that enables a plasmid or vector to replicate *in vivo*.

[0137] Examples of bacterial origins of replication are the origins of replication of plasmids pBR322, pUC19, pACYC177, and pACYC184 permitting replication in *E. coli*, and pUB110, pE194, pTA1060, and pAM β 1 permitting replication in *Bacillus*.

[0138] Examples of origins of replication for use in a yeast host cell are the 2 micron origin of replication, ARS1, ARS4, the combination of ARS1 and CEN3, and the combination of ARS4 and CEN6.

[0139] Examples of origins of replication useful in a filamentous fungal cell are AMA1 and ANS1 (Gems et al., 1991, Gene 98: 61-67; Cullen et al., 1987, Nucleic Acids Res. 15: 9163-9175; WO 00/24883). Isolation of the AMA1 gene and construction of plasmids or vectors comprising the gene can be accomplished according to the methods disclosed in WO 00/24883.

[0140] More than one copy of a polynucleotide of the present invention may be inserted into a host cell to increase production of a polypeptide. An increase in the copy number of the polynucleotide can be obtained by integrating at least one additional copy of the sequence into the host cell genome or by including an amplifiable selectable marker gene with the polynucleotide where cells containing amplified copies of the selectable marker gene, and thereby additional copies of the polynucleotide, can be selected for by cultivating the cells in the presence of the appropriate selectable agent.

[0141] The procedures used to ligate the elements described above to construct the recombinant expression vectors of the present invention are well known to one skilled in the art (see, e.g., Sambrook et al., 1989, *supra*).

Host Cells

[0142] The present invention also relates to recombinant host cells, comprising a polynucleotide of the present invention operably linked to one or more (e.g., several) control sequences that direct the production of a polypeptide of the present invention. A construct or vector comprising a polynucleotide is introduced into a host cell so that the construct or vector is maintained as a chromosomal integrant or as a self-replicating extra-chromosomal vector as described earlier. The term "host cell" encompasses any progeny of a parent cell that is not identical to the parent cell due to mutations that occur during replication. The choice of a host cell will to a large extent depend upon the gene encoding the polypeptide and its source.

[0143] The host cell may be any cell useful in the recombinant production of a polypeptide of the present invention, e.g., a prokaryote or a eukaryote.

[0144] The prokaryotic host cell may be any Gram-positive or Gram-negative bacterium. Gram-positive bacteria include, but are not limited to, *Bacillus*, *Clostridium*, *Enterococcus*, *Geobacillus*, *Lactobacillus*, *Lactococcus*, *Oceanobacillus*, *Staphylococcus*, *Streptococcus*, and *Streptomyces*. Gram-negative bacteria include, but are not limited to, *Campylobacter*, *E. coli*, *Flavobacterium*, *Fusobacterium*, *Helicobacter*, *Ilyobacter*, *Neisseria*, *Pseudomonas*, *Salmonella*, and *Ureaplasma*.

[0145] The bacterial host cell may be any *Bacillus* cell including, but not limited to, *Bacillus alkalophilus*, *Bacillus amyloliquefaciens*, *Bacillus brevis*, *Bacillus circulans*, *Bacillus clausii*, *Bacillus coagulans*, *Bacillus firmus*, *Bacillus lautus*, *Bacillus lentus*, *Bacillus licheniformis*, *Bacillus megaterium*, *Bacillus pumilus*, *Bacillus stearothermophilus*, *Bacillus subtilis*, and *Bacillus thuringiensis* cells.

[0146] The bacterial host cell may also be any *Streptococcus* cell including, but not limited to, *Streptococcus equisimilis*, *Streptococcus pyogenes*, *Streptococcus uberis*, and *Streptococcus equi* subsp. *Zooepidemicus* cells.

[0147] The bacterial host cell may also be any *Streptomyces* cell including, but not limited to, *Streptomyces achromogenes*, *Streptomyces avermitilis*, *Streptomyces coelicolor*, *Streptomyces griseus*, and *Streptomyces lividans* cells.

[0148] The introduction of DNA into a *Bacillus* cell may be effected by protoplast transformation (see, e.g., Chang and Cohen, 1979, Mol. Gen. Genet. 168: 111-115), competent cell transformation (see, e.g., Young and Spizizen, 1961, J. Bacteriol. 81: 823-829, or Dubnau and Davidoff-Abelson, 1971, J. Mol. Biol. 56: 209-221), electroporation (see, e.g., Shigekawa and Dower, 1988, Biotechniques 6: 742-751), or conjugation (see, e.g., Koehler and Thorne, 1987, J. Bacteriol. 169: 5271-5278). The introduction of DNA into an *E. coli* cell may be effected by protoplast transformation (see, e.g., Hanahan, 1983, J. Mol. Biol. 166: 557-580) or electroporation (see, e.g., Dower et al., 1988, Nucleic Acids Res. 16: 6127-6145). The introduction of DNA into a *Streptomyces* cell may be effected by protoplast transformation, electroporation (see, e.g., Gong et al., 2004, Folia Microbiol. (Praha) 49: 399-405), conjugation (see, e.g., Mazodier et al., 1989, J. Bacteriol. 171: 3583-3585), or transduction (see, e.g., Burke et al., 2001, Proc. Natl. Acad. Sci. USA 98: 6289-6294). The introduction of DNA into a *Pseudomonas* cell may be effected by electroporation (see, e.g., Choi et al., 2006, J. Microbiol. Methods 64: 391-397) or conjugation (see, e.g., Pinedo and Smets, 2005, Appl. Environ. Microbiol. 71: 51-57). The introduction of DNA into a *Streptococcus* cell may be effected by natural competence (see, e.g., Perry and Kuramitsu, 1981, Infect Immun. 32: 1295-1297), protoplast transformation (see, e.g., Catt and Jollick, 1991, Microbios 68: 189-207), electroporation (see, e.g., Buckley et al., 1999, Appl. Environ. Microbiol. 65: 3800-3804), or conjugation (see, e.g., Clewell, 1981, Microbiol. Rev. 45: 409-436). However, any method known in the art for introducing DNA into a host cell can be used.

[0149] The host cell may also be a eukaryote, such as a mammalian, insect, plant, or fungal cell.

[0150] The host cell may be a fungal cell. "Fungi" as used herein includes the phyla Ascomycota, Basidiomycota, Chytridiomycota, and Zygomycota as well as the Oomycota and all mitosporic fungi (as defined by Hawksworth et al., in Ainsworth and Bisby's Dictionary of The Fungi, 8th edition, 1995, CAB International, University Press, Cambridge, UK).

[0151] The fungal host cell may be a yeast cell. "Yeast" as used herein includes ascosporogenous yeast (Endomycetales), basidiosporogenous yeast, and yeast belonging to the Fungi Imperfecti (Blastomycetes). Since the classification of yeast may change in the future, for the purposes of this invention, yeast shall be defined as described in Biology and Activities of Yeast (Skinner, Passmore, and Davenport, editors, Soc. App. Bacteriol. Symposium Series No. 9, 1980).

[0152] The yeast host cell may be a *Candida*, *Hansenula*, *Kluyveromyces*, *Pichia*, *Saccharomyces*, *Schizosaccharomyces*, or *Yarrowia* cell such as a *Kluyveromyces lactis*, *Saccharomyces carlsbergensis*, *Saccharomyces cerevisiae*, *Saccharomyces diastaticus*, *Saccharomyces douglasii*, *Saccharomyces kluyveri*, *Saccharomyces norbensis*, *Saccharomyces oviformis*, or *Yarrowia lipolytica* cell.

[0153] The fungal host cell may be a filamentous fungal cell. "Filamentous fungi" include all filamentous forms of the subdivision Eumycota and Oomycota (as defined by Hawksworth et al., 1995, *supra*). The filamentous fungi are generally characterized by a mycelial wall composed of chitin, cellulose, glucan, chitosan, mannan, and other complex polysaccharides. Vegetative growth is by hyphal elongation and carbon catabolism is obligately aerobic. In contrast, vegetative growth by yeasts such as *Saccharomyces cerevisiae* is by budding of a unicellular thallus and carbon catabolism may be fermentative.

[0154] The filamentous fungal host cell may be an *Acremonium*, *Aspergillus*, *Aureobasidium*, *Bjerkandera*, *Ceriporiopsis*, *Chrysosporium*, *Coprinus*, *Coriolus*, *Cryptococcus*, *Filibasidium*, *Fusarium*, *Humicola*, *Magnaporthe*, *Mucor*, *Myceliophthora*, *Neocallimastix*, *Neurospora*, *Paecilomyces*, *Penicillium*, *Phanerochaete*, *Phlebia*, *Piromyces*, *Pleurotus*, *Schizophyllum*, *Talaromyces*, *Thermoascus*, *Thielavia*, *Tolypocladium*, *Trametes*, or *Trichoderma* cell.

[0155] For example, the filamentous fungal host cell may be an *Aspergillus awamori*, *Aspergillus foetidus*, *Aspergillus fumigatus*, *Aspergillus japonicus*, *Aspergillus nidulans*, *Aspergillus niger*, *Aspergillus oryzae*, *Bjerkandera adusta*, *Ceriporiopsis aneirina*, *Ceriporiopsis caregiea*, *Ceriporiopsis gilvescens*, *Ceriporiopsis pannocinta*, *Ceriporiopsis rivulosa*, *Ceriporiopsis subrufa*, *Ceriporiopsis subvermispora*, *Chrysosporium inops*, *Chrysosporium keratinophilum*, *Chrysosporium lucknowense*, *Chrysosporium merdarium*, *Chrysosporium pannicola*, *Chrysosporium queenslandicum*, *Chrysosporium tropicum*, *Chrysosporium zonatum*, *Coprinus cinereus*, *Coriolus hirsutus*, *Fusarium bactridioides*, *Fusarium cerealis*, *Fusarium crookwellense*, *Fusarium culmorum*, *Fusarium graminearum*, *Fusarium graminum*, *Fusarium heterosporum*, *Fusarium negundi*, *Fusarium oxysporum*, *Fusarium reticulatum*, *Fusarium roseum*, *Fusarium sambucinum*, *Fusarium sarcochroum*, *Fusarium sporotrichioides*, *Fusarium sulphureum*, *Fusarium torulosum*, *Fusarium trichothecioides*, *Fusarium venenatum*, *Humicola insolens*, *Humicola lanuginosa*, *Mucor miehei*, *Myceliophthora thermophila*, *Neurospora crassa*, *Penicillium purpurogenum*, *Phanerochaete chrysosporium*, *Phlebia radiata*, *Pleurotus eryngii*, *Thielavia terrestris*, *Trametes villosa*, *Trametes versicolor*, *Trichoderma harzianum*, *Trichoderma koningii*, *Trichoderma longibrachiatum*, *Trichoderma reesei*, or *Trichoderma viride* cell.

[0156] Fungal cells may be transformed by a process involving protoplast formation, transformation of the protoplasts, and regeneration of the cell wall in a manner known *per se*. Suitable procedures for transformation of *Aspergillus* and *Trichoderma* host cells are described in EP 238023, Yelton et al., 1984, Proc. Natl. Acad. Sci. USA 81: 1470-1474, and Christensen et al., 1988, Bio/Technology 6: 1419-1422. Suitable methods for transforming *Fusarium* species are described by Malardier et al., 1989, Gene 78: 147-156, and WO 96/00787. Yeast may be transformed using the procedures described by Becker and Guarente, In Abelson, J.N. and Simon, M.I., editors, Guide to Yeast Genetics and Molecular Biology, Methods in Enzymology, Volume 194, pp 182-187, Academic Press, Inc., New York; Ito et al., 1983, J. Bacteriol. 153: 163; and Hinnen et al., 1978, Proc. Natl. Acad. Sci. USA 75: 1920.

Methods of Production

[0157] The present invention also relates to methods of producing a polypeptide of the present invention, comprising: (a) cultivating a cell, which in its wild-type form produces the polypeptide, under conditions conducive for production of the polypeptide; and (b) recovering the polypeptide. In a preferred aspect, the cell is of the genus *Penicillium*. In a more preferred aspect, the cell is *Penicillium oxalicum*. In a most preferred aspect, the cell is *Penicillium oxalicum* IBT5387.

[0158] The present invention also relates to methods of producing a polypeptide of the present invention, comprising: (a) cultivating a recombinant host cell of the present invention under conditions conducive for production of the polypeptide; and (b) recovering the polypeptide.

[0159] The host cells are cultivated in a nutrient medium suitable for production of the polypeptide using methods known in the art. For example, the cell may be cultivated by shake flask cultivation, or small-scale or large-scale fermentation (including continuous, batch, fed-batch, or solid state fermentations) in laboratory or industrial fermentors in a suitable medium and under conditions allowing the polypeptide to be expressed and/or isolated. The cultivation takes place in a suitable nutrient medium comprising carbon and nitrogen sources and inorganic salts, using procedures known in the art. Suitable media are available from commercial suppliers or may be prepared according to published compositions (e.g., in catalogues of the American Type Culture Collection). If the polypeptide is secreted into the nutrient medium, the polypeptide can be recovered directly from the medium. If the polypeptide is not secreted, it can be recovered from cell lysates.

[0160] The polypeptide may be detected using methods known in the art that are specific for the polypeptides. These detection methods include, but are not limited to, use of specific antibodies, formation of an enzyme product, or disappearance of an enzyme substrate. For example, an enzyme assay may be used to determine the activity of the polypeptide, as described herein.

[0161] The polypeptide may be recovered using methods known in the art. For example, the polypeptide may be recovered from the nutrient medium by conventional procedures including, but not limited to, collection, centrifugation, filtration, extraction, spray-drying, evaporation, or precipitation.

[0162] The polypeptide may be purified by a variety of procedures known in the art including, but not limited to, chromatography (e.g., ion exchange, affinity, hydrophobic, chromatofocusing, and size exclusion), electrophoretic procedures (e.g., preparative isoelectric focusing), differential solubility (e.g., ammonium sulfate precipitation), SDS-PAGE, or extraction (see, e.g., Protein Purification, Janson and Ryden, editors, VCH Publishers, New York, 1989) to obtain substantially pure polypeptides.

[0163] In an alternative aspect, the polypeptide is not recovered, but rather a host cell of the present invention expressing the polypeptide is used as a source of the polypeptide.

Plants

[0164] The present invention also relates to isolated plants, e.g., a transgenic plant, plant part, or plant cell, comprising an isolated polynucleotide of the present invention so as to express and produce a polypeptide in recoverable quantities. The polypeptide may be recovered from the plant or plant part. Alternatively, the plant or plant part containing the polypeptide may be used as such for improving the quality of a food or feed, e.g., improving nutritional value, palatability, and rheological properties, or to destroy an antinutritive factor.

[0165] The transgenic plant can be dicotyledonous (a dicot) or monocotyledonous (a monocot). Examples of monocot plants are

grasses, such as meadow grass (blue grass, *Poa*), forage grass such as *Festuca*, *Lolium*, temperate grass, such as *Agrostis*, and cereals, e.g., wheat, oats, rye, barley, rice, sorghum, and maize (corn).

[0166] Examples of dicot plants are tobacco, legumes, such as lupins, potato, sugar beet, pea, bean and soybean, and cruciferous plants (family Brassicaceae), such as cauliflower, rape seed, and the closely related model organism *Arabidopsis thaliana*.

[0167] Examples of plant parts are stem, callus, leaves, root, fruits, seeds, and tubers as well as the individual tissues comprising these parts, e.g., epidermis, mesophyll, parenchyme, vascular tissues, meristems. Specific plant cell compartments, such as chloroplasts, apoplasts, mitochondria, vacuoles, peroxisomes and cytoplasm are also considered to be a plant part. Furthermore, any plant cell, whatever the tissue origin, is considered to be a plant part. Likewise, plant parts such as specific tissues and cells isolated to facilitate the utilization of the invention are also considered plant parts, e.g., embryos, endosperms, aleurone and seed coats.

[0168] Also included within the scope of the present invention are the progeny of such plants, plant parts, and plant cells.

[0169] The transgenic plant or plant cell expressing the polypeptide may be constructed in accordance with methods known in the art. In short, the plant or plant cell is constructed by incorporating one or more expression constructs encoding the polypeptide into the plant host genome or chloroplast genome and propagating the resulting modified plant or plant cell into a transgenic plant or plant cell.

[0170] The expression construct is conveniently a nucleic acid construct that comprises a polynucleotide encoding a polypeptide operably linked with appropriate regulatory sequences required for expression of the polynucleotide in the plant or plant part of choice. Furthermore, the expression construct may comprise a selectable marker useful for identifying plant cells into which the expression construct has been integrated and DNA sequences necessary for introduction of the construct into the plant in question (the latter depends on the DNA introduction method to be used).

[0171] The choice of regulatory sequences, such as promoter and terminator sequences and optionally signal or transit sequences, is determined, for example, on the basis of when, where, and how the polypeptide is desired to be expressed. For instance, the expression of the gene encoding a polypeptide may be constitutive or inducible, or may be developmental, stage or tissue specific, and the gene product may be targeted to a specific tissue or plant part such as seeds or leaves. Regulatory sequences are, for example, described by Tague et al., 1988, Plant Physiology 86: 506.

[0172] For constitutive expression, the 35S-CaMV, the maize ubiquitin 1, or the rice actin 1 promoter may be used (Franck et al., 1980, Cell 21: 285-294; Christensen et al., 1992, Plant Mol. Biol. 18: 675-689; Zhang et al., 1991, Plant Cell 3: 1155-1165). Organ-specific promoters may be, for example, a promoter from storage sink tissues such as seeds, potato tubers, and fruits (Edwards and Coruzzi, 1990, Ann. Rev. Genet. 24: 275-303), or from metabolic sink tissues such as meristems (Ito et al., 1994, Plant Mol. Biol. 24: 863-878), a seed specific promoter such as the glutelin, prolamin, globulin, or albumin promoter from rice (Wu et al., 1998, Plant Cell Physiol. 39: 885-889), a *Vicia faba* promoter from the legumin B4 and the unknown seed protein gene from *Vicia faba* (Conrad et al., 1998, J. Plant Physiol. 152: 708-711), a promoter from a seed oil body protein (Chen et al., 1998, Plant Cell Physiol. 39: 935-941), the storage protein *napA* promoter from *Brassica napus*, or any other seed specific promoter known in the art, e.g., as described in WO 91/14772. Furthermore, the promoter may be a leaf specific promoter such as the *rbcs* promoter from rice or tomato (Kozuka et al., 1993, Plant Physiol. 102: 991-1000), the chlorella virus adenine methyltransferase gene promoter (Mitra and Higgins, 1994, Plant Mol. Biol. 26: 85-93), the *aldP* gene promoter from rice (Kagaya et al., 1995, Mol. Gen. Genet. 248: 668-674), or a wound inducible promoter such as the potato *pin2* promoter (Xu et al., 1993, Plant Mol. Biol. 22: 573-588). Likewise, the promoter may be induced by abiotic treatments such as temperature, drought, or alterations in salinity or induced by exogenously applied substances that activate the promoter, e.g., ethanol, oestrogens, plant hormones such as ethylene, abscisic acid, and gibberellic acid, and heavy metals.

[0173] A promoter enhancer element may also be used to achieve higher expression of a polypeptide in the plant. For instance, the promoter enhancer element may be an intron that is placed between the promoter and the polynucleotide encoding a polypeptide. For instance, Xu et al., 1993, *supra*, disclose the use of the first intron of the rice actin 1 gene to enhance expression.

[0174] The selectable marker gene and any other parts of the expression construct may be chosen from those available in the art.

[0175] The nucleic acid construct is incorporated into the plant genome according to conventional techniques known in the art,

including *Agrobacterium*-mediated transformation, virus-mediated transformation, microinjection, particle bombardment, biolistic transformation, and electroporation (Gasser et al., 1990, *Science* 244: 1293; Potrykus, 1990, *Bio/Technology* 8: 535; Shimamoto et al., 1989, *Nature* 338: 274).

[0176] *Agrobacterium tumefaciens*-mediated gene transfer is a method for generating transgenic dicots (for a review, see Hooykas and Schilperoort, 1992, *Plant Mol. Biol.* 19: 15-38) and for transforming monocots, although other transformation methods may be used for these plants. A method for generating transgenic monocots is particle bombardment (microscopic gold or tungsten particles coated with the transforming DNA) of embryonic calli or developing embryos (Christou, 1992, *Plant J.* 2: 275-281; Shimamoto, 1994, *Curr. Opin. Biotechnol.* 5: 158-162; Vasil et al., 1992, *Bio/Technology* 10: 667-674). An alternative method for transformation of monocots is based on protoplast transformation as described by Omirulleh et al., 1993, *Plant Mol. Biol.* 21: 415-428. Additional transformation methods include those described in U.S. Patent Nos. 6,395,966 and 7,151,204.

[0177] Following transformation, the transformants having incorporated the expression construct are selected and regenerated into whole plants according to methods well known in the art. Often the transformation procedure is designed for the selective elimination of selection genes either during regeneration or in the following generations by using, for example, co-transformation with two separate T-DNA constructs or site specific excision of the selection gene by a specific recombinase.

[0178] In addition to direct transformation of a particular plant genotype with a construct of the present invention, transgenic plants may be made by crossing a plant having the construct to a second plant lacking the construct. For example, a construct encoding a polypeptide can be introduced into a particular plant variety by crossing, without the need for ever directly transforming a plant of that given variety. Therefore, the present invention encompasses not only a plant directly regenerated from cells which have been transformed in accordance with the present invention, but also the progeny of such plants. As used herein, progeny may refer to the offspring of any generation of a parent plant prepared in accordance with the present invention. Such progeny may include a DNA construct prepared in accordance with the present invention. Crossing results in the introduction of a transgene into a plant line by cross pollinating a starting line with a donor plant line. Non-limiting examples of such steps are described in U.S. Patent No. 7,151,204.

[0179] Plants may be generated through a process of backcross conversion. For example, plants include plants referred to as a backcross converted genotype, line, inbred, or hybrid.

[0180] Genetic markers may be used to assist in the introgression of one or more transgenes of the invention from one genetic background into another. Marker assisted selection offers advantages relative to conventional breeding in that it can be used to avoid errors caused by phenotypic variations. Further, genetic markers may provide data regarding the relative degree of elite germplasm in the individual progeny of a particular cross. For example, when a plant with a desired trait which otherwise has a non-agronomically desirable genetic background is crossed to an elite parent, genetic markers may be used to select progeny which not only possess the trait of interest, but also have a relatively large proportion of the desired germplasm. In this way, the number of generations required to introgress one or more traits into a particular genetic background is minimized.

[0181] The present invention also relates to methods of producing a polypeptide of the present invention comprising: (a) cultivating a transgenic plant or a plant cell comprising a polynucleotide encoding the polypeptide under conditions conducive for production of the polypeptide; and (b) recovering the polypeptide.

Removal or Reduction of Beta-Glucosidase Activity

[0182] The present invention also relates to methods of producing a mutant of a parent cell, which comprises disrupting or deleting a polynucleotide, or a portion thereof, encoding a polypeptide of the present invention, which results in the mutant cell producing less of the polypeptide than the parent cell when cultivated under the same conditions.

[0183] The mutant cell may be constructed by reducing or eliminating expression of the polynucleotide using methods well known in the art, for example, insertions, disruptions, replacements, or deletions. In a preferred aspect, the polynucleotide is inactivated. The polynucleotide to be modified or inactivated may be, for example, the coding region or a part thereof essential for activity, or a regulatory element required for expression of the coding region. An example of such a regulatory or control sequence may be a promoter sequence or a functional part thereof, *i.e.*, a part that is sufficient for affecting expression of the polynucleotide. Other control sequences for possible modification include, but are not limited to, a leader, polyadenylation sequence, propeptide sequence, signal peptide sequence, transcription terminator, and transcriptional activator.

[0184] Modification or inactivation of the polynucleotide may be performed by subjecting the parent cell to mutagenesis and selecting for mutant cells in which expression of the polynucleotide has been reduced or eliminated. The mutagenesis, which may be specific or random, may be performed, for example, by use of a suitable physical or chemical mutagenizing agent, by use of a suitable oligonucleotide, or by subjecting the DNA sequence to PCR generated mutagenesis. Furthermore, the mutagenesis may be performed by use of any combination of these mutagenizing agents.

[0185] Examples of a physical or chemical mutagenizing agent suitable for the present purpose include ultraviolet (UV) irradiation, hydroxylamine, N-methyl-N'-nitro-N-nitrosoguanidine (MNNG), O-methyl hydroxylamine, nitrous acid, ethyl methane sulphonate (EMS), sodium bisulphite, formic acid, and nucleotide analogues.

[0186] When such agents are used, the mutagenesis is typically performed by incubating the parent cell to be mutagenized in the presence of the mutagenizing agent of choice under suitable conditions, and screening and/or selecting for mutant cells exhibiting reduced or no expression of the gene.

[0187] Modification or inactivation of the polynucleotide may be accomplished by insertion, substitution, or deletion of one or more nucleotides in the gene or a regulatory element required for transcription or translation thereof. For example, nucleotides may be inserted or removed so as to result in the introduction of a stop codon, the removal of the start codon, or a change in the open reading frame. Such modification or inactivation may be accomplished by site-directed mutagenesis or PCR generated mutagenesis in accordance with methods known in the art. Although, in principle, the modification may be performed *in vivo*, *i.e.*, directly on the cell expressing the polynucleotide to be modified, it is preferred that the modification be performed *in vitro* as exemplified below.

[0188] An example of a convenient way to eliminate or reduce expression of a polynucleotide is based on techniques of gene replacement, gene deletion, or gene disruption. For example, in the gene disruption method, a nucleic acid sequence corresponding to the endogenous polynucleotide is mutagenized *in vitro* to produce a defective nucleic acid sequence that is then transformed into the parent cell to produce a defective gene. By homologous recombination, the defective nucleic acid sequence replaces the endogenous polynucleotide. It may be desirable that the defective polynucleotide also encodes a marker that may be used for selection of transformants in which the polynucleotide has been modified or destroyed. In an aspect, the polynucleotide is disrupted with a selectable marker such as those described herein.

[0189] The present invention also relates to methods of inhibiting the expression of a polypeptide having beta-glucosidase activity in a cell, comprising administering to the cell or expressing in the cell a double-stranded RNA (dsRNA) molecule, wherein the dsRNA comprises a subsequence of a polynucleotide of the present invention. In a preferred aspect, the dsRNA is about 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25 or more duplex nucleotides in length.

[0190] The dsRNA is preferably a small interfering RNA (siRNA) or a micro RNA (miRNA). In a preferred aspect, the dsRNA is small interfering RNA for inhibiting transcription. In another preferred aspect, the dsRNA is micro RNA for inhibiting translation.

[0191] The present invention also relates to such double-stranded RNA (dsRNA) molecules, comprising a portion of the mature polypeptide coding sequence of SEQ ID NO: 1 for inhibiting expression of the polypeptide in a cell. While the present invention is not limited by any particular mechanism of action, the dsRNA can enter a cell and cause the degradation of a single-stranded RNA (ssRNA) of similar or identical sequences, including endogenous mRNAs. When a cell is exposed to dsRNA, mRNA from the homologous gene is selectively degraded by a process called RNA interference (RNAi).

[0192] The dsRNAs of the present invention can be used in gene-silencing. In one aspect, the invention provides methods to selectively degrade RNA using a dsRNAi of the present invention. The process may be practiced *in vitro*, *ex vivo* or *in vivo*. In one aspect, the dsRNA molecules can be used to generate a loss-of-function mutation in a cell, an organ or an animal. Methods for making and using dsRNA molecules to selectively degrade RNA are well known in the art, see, for example, U.S. Patent Nos. 6,489,127; 6,506,559; 6,511,824; and 6,515,109.

[0193] The present invention further relates to a mutant cell of a parent cell that comprises a disruption or deletion of a polynucleotide encoding the polypeptide or a control sequence thereof or a silenced gene encoding the polypeptide, which results in the mutant cell producing less of the polypeptide or no polypeptide compared to the parent cell.

[0194] The polypeptide-deficient mutant cells are particularly useful as host cells for expression of native and heterologous polypeptides. Therefore, the present invention further relates to methods of producing a native or heterologous polypeptide, comprising: (a) cultivating the mutant cell under conditions conducive for production of the polypeptide; and (b) recovering the polypeptide. The term "heterologous polypeptides" means polypeptides that are not native to the host cell, *e.g.*, a variant of a

native protein. The host cell may comprise more than one copy of a polynucleotide encoding the native or heterologous polypeptide.

[0195] The methods used for cultivation and purification of the product of interest may be performed by methods known in the art.

[0196] The methods of the present invention for producing an essentially beta-glucosidase-free product is of particular interest in the production of eukaryotic polypeptides, in particular fungal proteins such as enzymes. The beta-glucosidase-deficient cells may also be used to express heterologous proteins of pharmaceutical interest such as hormones, growth factors, receptors, and the like. The term "eukaryotic polypeptides" includes not only native polypeptides, but also those polypeptides, e.g., enzymes, which have been modified by amino acid substitutions, deletions or additions, or other such modifications to enhance activity, thermostability, pH tolerance and the like.

[0197] In a further aspect, the present invention relates to a protein product essentially free from beta-glucosidase activity that is produced by a method of the present invention.

Compositions

[0198] The present invention also relates to compositions comprising a polypeptide of the present invention. Preferably, the compositions are enriched in such a polypeptide. The term "enriched" indicates that the beta-glucosidase activity of the composition has been increased, e.g., with an enrichment factor of at least 1.1.

[0199] The composition may comprise a polypeptide of the present invention as the major enzymatic component, e.g., a mono-component composition. Alternatively, the composition may comprise multiple enzymatic activities, such as an aminopeptidase, amylase, carboxypeptidase, catalase, cellulase, chitinase, cutinase, cyclodextrin glycosyltransferase, deoxyribonuclease, esterase, alpha-galactosidase, beta-galactosidase, glucoamylase, alpha-glucosidase, beta-glucosidase, haloperoxidase, invertase, laccase, lipase, mannosidase, oxidase, pectinolytic enzyme, peptidoglutaminase, peroxidase, phytase, polyphenoloxidase, proteolytic enzyme, ribonuclease, transglutaminase, or xylanase. The additional enzyme(s) may be produced, for example, by a microorganism belonging to the genus *Aspergillus*, e.g., *Aspergillus aculeatus*, *Aspergillus awamori*, *Aspergillus foetidus*, *Aspergillus fumigatus*, *Aspergillus japonicus*, *Aspergillus nidulans*, *Aspergillus niger*, or *Aspergillus oryzae*; *Fusarium*, e.g., *Fusarium bacridioides*, *Fusarium cerealis*, *Fusarium crookwellense*, *Fusarium culmorum*, *Fusarium graminearum*, *Fusarium graminum*, *Fusarium heterosporum*, *Fusarium negundi*, *Fusarium oxysporum*, *Fusarium reticulatum*, *Fusarium roseum*, *Fusarium sambucinum*, *Fusarium sarcochroum*, *Fusarium sulphureum*, *Fusarium toruloseum*, *Fusarium trichothecioides*, or *Fusarium venenatum*; *Humicola*, e.g., *Humicola insolens* or *Humicola lanuginosa*; or *Trichoderma*, e.g., *Trichoderma harzianum*, *Trichoderma koningii*, *Trichoderma longibrachiatum*, *Trichoderma reesei*, or *Trichoderma viride*.

[0200] The compositions may be prepared in accordance with methods known in the art and may be in the form of a liquid or a dry composition. For instance, the composition may be in the form of a granulate or a microgranulate. The polypeptide may be stabilized in accordance with methods known in the art.

Uses

[0201] The present invention is also directed to the following methods for using the polypeptides having beta-glucosidase activity, or compositions thereof.

[0202] The present invention also relates to methods for degrading or converting a cellulosic material, comprising: treating the cellulosic material with an enzyme composition in the presence of a polypeptide having beta-glucosidase activity of the present invention. In a preferred aspect, the methods further comprise recovering the degraded or converted cellulosic material.

[0203] The present invention also relates to methods of producing a fermentation product, comprising: (a) saccharifying a cellulosic material with an enzyme composition in the presence of a polypeptide having beta-glucosidase activity of the present invention; (b) fermenting the saccharified cellulosic material with one or more (e.g., several) fermenting microorganisms to produce the fermentation product; and (c) recovering the fermentation product from the fermentation.

[0204] The present invention also relates to methods of fermenting a cellulosic material, comprising: fermenting the cellulosic

material with one or more (e.g., several) fermenting microorganisms, wherein the cellulosic material is saccharified with an enzyme composition in the presence of a polypeptide having beta-glucosidase activity of the present invention. In a preferred aspect, the fermenting of the cellulosic material produces a fermentation product. In another preferred aspect, the methods further comprise recovering the fermentation product from the fermentation.

[0205] The methods of the present invention can be used to saccharify a cellulosic material to fermentable sugars and convert the fermentable sugars to many useful substances, e.g., fuel, potable ethanol, and/or fermentation products (e.g., acids, alcohols, ketones, gases, and the like). The production of a desired fermentation product from the cellulosic material typically involves pretreatment, enzymatic hydrolysis (saccharification), and fermentation.

[0206] The processing of a cellulosic material according to the present invention can be accomplished using processes conventional in the art. Moreover, the methods of the present invention can be implemented using any conventional biomass processing apparatus configured to operate in accordance with the invention.

[0207] Hydrolysis (saccharification) and fermentation, separate or simultaneous, include, but are not limited to, separate hydrolysis and fermentation (SHF); simultaneous saccharification and fermentation (SSF); simultaneous saccharification and co-fermentation (SSCF); hybrid hydrolysis and fermentation (HHF); separate hydrolysis and co-fermentation (SHCF); hybrid hydrolysis and co-fermentation (HHCF); and direct microbial conversion (DMC), also sometimes called consolidated bioprocessing (CBP). SHF uses separate process steps to first enzymatically hydrolyze cellulosic material to fermentable sugars, e.g., glucose, cellobiose, cellobiose, and pentose monomers, and then ferment the fermentable sugars to ethanol. In SSF, the enzymatic hydrolysis of the cellulosic material and the fermentation of sugars to ethanol are combined in one step (Philippidis, G. P., 1996, Cellulose bioconversion technology, in Handbook on Bioethanol: Production and Utilization, Wyman, C. E., ed., Taylor & Francis, Washington, DC, 179-212). SSCF involves the co-fermentation of multiple sugars (Sheehan, J., and Himmel, M., 1999, Enzymes, energy and the environment: A strategic perspective on the U.S. Department of Energy's research and development activities for bioethanol, Biotechnol. Prog. 15: 817-827). HHF involves a separate hydrolysis step, and in addition a simultaneous saccharification and hydrolysis step, which can be carried out in the same reactor. The steps in an HHF process can be carried out at different temperatures, i.e., high temperature enzymatic saccharification followed by SSF at a lower temperature that the fermentation strain can tolerate. DMC combines all three processes (enzyme production, hydrolysis, and fermentation) in one or more (e.g., several) steps where the same organism is used to produce the enzymes for conversion of the cellulosic material to fermentable sugars and to convert the fermentable sugars into a final product (Lynd, L. R., Weimer, P. J., van Zyl, W. H., and Pretorius, I. S., 2002, Microbial cellulose utilization: Fundamentals and biotechnology, Microbiol. Mol. Biol. Reviews 66: 506-577). It is understood herein that any method known in the art comprising pretreatment, enzymatic hydrolysis (saccharification), fermentation, or a combination thereof, can be used in the practicing the methods of the present invention.

[0208] A conventional apparatus can include a fed-batch stirred reactor, a batch stirred reactor, a continuous flow stirred reactor with ultrafiltration, and/or a continuous plug-flow column reactor (Fernanda de Castilhos Corazza, Flávio Faria de Moraes, Gisella Maria Zanin and Ivo Neitzel, 2003, Optimal control in fed-batch reactor for the cellobiose hydrolysis, Acta Scientiarum. Technology 25: 33-38; Gusakov, A. V., and Sinitsyn, A. P., 1985, Kinetics of the enzymatic hydrolysis of cellulose: 1. A mathematical model for a batch reactor process, Enz. Microb. Technol. 7: 346-352), an attrition reactor (Ryu, S. K., and Lee, J. M., 1983, Bioconversion of waste cellulose by using an attrition bioreactor, Biotechnol. Bioeng. 25: 53-65), or a reactor with intensive stirring induced by an electromagnetic field (Gusakov, A. V., Sinitsyn, A. P., Davydkin, I. Y., Davydkin, V. Y., Protas, O. V., 1996, Enhancement of enzymatic cellulose hydrolysis using a novel type of bioreactor with intensive stirring induced by electromagnetic field, Appl. Biochem. Biotechnol. 56: 141-153). Additional reactor types include: fluidized bed, upflow blanket, immobilized, and extruder type reactors for hydrolysis and/or fermentation.

[0209] Pretreatment. In practicing the methods of the present invention, any pretreatment process known in the art can be used to disrupt plant cell wall components of a cellulosic material (Chandra et al., 2007, Substrate pretreatment: The key to effective enzymatic hydrolysis of lignocelluloses? Adv. Biochem. Engin./Biotechnol. 108: 67-93; Galbe and Zacchi, 2007, Pretreatment of lignocellulosic materials for efficient bioethanol production, Adv. Biochem. Engin. / Biotechnol. 108: 41-65; Hendriks and Zeeman, 2009, Pretreatments to enhance the digestibility of lignocellulosic biomass, Bioresource Technol. 100: 10-18; Mosier et al., 2005, Features of promising technologies for pretreatment of lignocellulosic biomass, Bioresource Technol. 96: 673-686; Taherzadeh and Karimi, 2008, Pretreatment of lignocellulosic wastes to improve ethanol and biogas production: A review, Int. J. of Mol. Sci. 9: 1621-1651; Yang and Wyman, 2008, Pretreatment: the key to unlocking low-cost cellulosic ethanol, Biofuels Bioproducts and Biorefining-Biofr. 2: 26-40).

[0210] The cellulosic material can also be subjected to particle size reduction, sieving, presoaking, wetting, washing, and/or conditioning prior to pretreatment using methods known in the art.

[0211] Conventional pretreatments include, but are not limited to, steam pretreatment (with or without explosion), dilute acid pretreatment, hot water pretreatment, alkaline pretreatment, lime pretreatment, wet oxidation, wet explosion, ammonia fiber explosion, organosolv pretreatment, and biological pretreatment. Additional pretreatments include ammonia percolation, ultrasound, electroporation, microwave, supercritical CO₂, supercritical H₂O, ozone, ionic liquid, and gamma irradiation pretreatments.

[0212] The cellulosic material can be pretreated before hydrolysis and/or fermentation. Pretreatment is preferably performed prior to the hydrolysis. Alternatively, the pretreatment can be carried out simultaneously with enzyme hydrolysis to release fermentable sugars, such as glucose, xylose, and/or cellobiose. In most cases the pretreatment step itself results in some conversion of biomass to fermentable sugars (even in absence of enzymes).

[0213] Steam Pretreatment: In steam pretreatment, the cellulosic material is heated to disrupt the plant cell wall components, including lignin, hemicellulose, and cellulose to make the cellulose and other fractions, e.g., hemicellulose, accessible to enzymes. The cellulosic material is passed to or through a reaction vessel where steam is injected to increase the temperature to the required temperature and pressure and is retained therein for the desired reaction time. Steam pretreatment is preferably performed at 140-230°C, more preferably 160-200°C, and most preferably 170-190°C, where the optimal temperature range depends on any addition of a chemical catalyst. Residence time for the steam pretreatment is preferably 1-15 minutes, more preferably 3-12 minutes, and most preferably 4-10 minutes, where the optimal residence time depends on temperature range and addition of a chemical catalyst. Steam pretreatment allows for relatively high solids loadings, so that the cellulosic material is generally only moist during the pretreatment. The steam pretreatment is often combined with an explosive discharge of the material after the pretreatment, which is known as steam explosion, that is, rapid flashing to atmospheric pressure and turbulent flow of the material to increase the accessible surface area by fragmentation (Duff and Murray, 1996, *Bioresource Technology* 85: 1-33; Galbe and Zacchi, 2002, *Appl. Microbiol. Biotechnol.* 59: 618-628; U.S. Patent Application No. 20020164730). During steam pretreatment, hemicellulose acetyl groups are cleaved and the resulting acid autocatalyzes partial hydrolysis of the hemicellulose to monosaccharides and oligosaccharides. Lignin is removed to only a limited extent.

[0214] Chemical Pretreatment: The term "chemical treatment" refers to any chemical pretreatment that promotes the separation and/or release of cellulose, hemicellulose, and/or lignin. Such a pretreatment can convert crystalline cellulose to amorphous cellulose. Examples of suitable chemical pretreatment processes include, for example, dilute acid pretreatment, lime pretreatment, wet oxidation, ammonia fiber/freezing explosion (AFEX), ammonia percolation (APR), ionic liquid, and organosolv pretreatments.

[0215] A catalyst such as H₂SO₄ or SO₂ (typically 0.3 to 3% w/w) is often added prior to steam pretreatment, which decreases the time and temperature, increases the recovery, and improves enzymatic hydrolysis (Ballesteros et al., 2006, *Appl. Biochem. Biotechnol.* 129-132: 496-508; Varga et al., 2004, *Appl. Biochem. Biotechnol.* 113-116: 509-523; Sassner et al., 2006, *Enzyme Microb. Technol.* 39: 756-762). In dilute acid pretreatment, cellulosic material is mixed with dilute acid, typically H₂SO₄, and water to form a slurry, heated by steam to the desired temperature, and after a residence time flashed to atmospheric pressure. The dilute acid pretreatment can be performed with a number of reactor designs, e.g., plug-flow reactors, counter-current reactors, or continuous counter-current shrinking bed reactors (Duff and Murray, 1996, *supra*; Schell et al., 2004, *Bioresource Technol.* 91: 179-188; Lee et al., 1999, *Adv. Biochem. Eng. Biotechnol.* 65: 93-115).

[0216] Several methods of pretreatment under alkaline conditions can also be used. These alkaline pretreatments include, but are not limited to, sodium hydroxide, lime, wet oxidation, ammonia percolation (APR), and ammonia fiber/freezing explosion (AFEX).

[0217] Lime pretreatment is performed with calcium oxide or calcium hydroxide at temperatures of 85-150°C and residence times from 1 hour to several days (Wyman et al., 2005, *Bioresource Technol.* 96: 1959-1966; Mosier et al., 2005, *Bioresource Technol.* 96: 673-686). WO 2006/110891, WO 2006/110899, WO 2006/110900, and WO 2006/110901 disclose pretreatment methods using ammonia.

[0218] Wet oxidation is a thermal pretreatment performed typically at 180-200°C for 5-15 minutes with addition of an oxidative agent such as hydrogen peroxide or over-pressure of oxygen (Schmidt and Thomsen, 1998, *Bioresource Technol.* 64: 139-151; Palonen et al., 2004, *Appl. Biochem. Biotechnol.* 117: 1-17; Varga et al., 2004, *Biotechnol. Bioeng.* 88: 567-574; Martin et al., 2006, *J. Chem. Technol. Biotechnol.* 81: 1669-1677). The pretreatment is preferably performed at 1-40% dry matter, more preferably 2-30% dry matter, and most preferably 5-20% dry matter, and often the initial pH is increased by the addition of alkali such as sodium carbonate.

[0219] A modification of the wet oxidation pretreatment method, known as wet explosion (combination of wet oxidation and steam explosion), can handle dry matter up to 30%. In wet explosion, the oxidizing agent is introduced during pretreatment after a certain

residence time. The pretreatment is then ended by flashing to atmospheric pressure (WO 2006/032282).

[0220] Ammonia fiber explosion (AFEX) involves treating the cellulosic material with liquid or gaseous ammonia at moderate temperatures such as 90-150°C and high pressure such as 17-20 bar for 5-10 minutes, where the dry matter content can be as high as 60% (Gollapalli et al., 2002, *Appl. Biochem. Biotechnol.* 98: 23-35; Chundawat et al., 2007, *Biotechnol. Bioeng.* 96: 219-231; Alizadeh et al., 2005, *Appl. Biochem. Biotechnol.* 121: 1133-1141; Teymouri et al., 2005, *Bioresource Technol.* 96: 2014-2018). During AFEX pretreatment cellulose and hemicelluloses remain relatively intact. Lignin-carbohydrate complexes are cleaved.

[0221] Organosolv pretreatment delignifies the cellulosic material by extraction using aqueous ethanol (40-60% ethanol) at 160-200°C for 30-60 minutes (Pan et al., 2005, *Biotechnol. Bioeng.* 90: 473-481; Pan et al., 2006, *Biotechnol. Bioeng.* 94: 851-861; Kurabi et al., 2005, *Appl. Biochem. Biotechnol.* 121: 219-230). Sulphuric acid is usually added as a catalyst. In organosolv pretreatment, the majority of hemicellulose and lignin is removed.

[0222] Other examples of suitable pretreatment methods are described by Schell et al., 2003, *Appl. Biochem. and Biotechnol.* Vol. 105-108, p. 69-85, and Mosier et al., 2005, *Bioresource Technology* 96: 673-686, and U.S. Published Application 2002/0164730.

[0223] In one aspect, the chemical pretreatment is preferably carried out as a dilute acid treatment, and more preferably as a continuous dilute acid treatment. The acid is typically sulfuric acid, but other acids can also be used, such as acetic acid, citric acid, nitric acid, phosphoric acid, tartaric acid, succinic acid, hydrogen chloride, or mixtures thereof. Mild acid treatment is conducted in the pH range of preferably 1-5, e.g., 1-4 or 1-2.5. In one aspect, the acid concentration is in the range from preferably 0.01 to 10 wt % acid, e.g., 0.05 to 5 wt % acid or 0.1 to 2 wt % acid. The acid is contacted with the cellulosic material and held at a temperature in the range of preferably 140-200°C, e.g., 165-190°C, for periods ranging from 1 to 60 minutes.

[0224] In another aspect, pretreatment takes place in an aqueous slurry. In preferred aspects, the cellulosic material is present during pretreatment in amounts preferably between 10-80 wt %, e.g., 20-70 wt % or 30-60 wt %, such as around 40 wt %. The pretreated cellulosic material can be unwashed or washed using any method known in the art, e.g., washed with water.

[0225] Mechanical Pretreatment or Physical Pretreatment: The term "mechanical pretreatment" or "physical pretreatment" refers to any pretreatment that promotes size reduction of particles. For example, such pretreatment can involve various types of grinding or milling (e.g., dry milling, wet milling, or vibratory ball milling).

[0226] The cellulosic material can be pretreated both physically (mechanically) and chemically. Mechanical or physical pretreatment can be coupled with steaming/steam explosion, hydrothermolysis, dilute or mild acid treatment, high temperature, high pressure treatment, irradiation (e.g., microwave irradiation), or combinations thereof. In one aspect, high pressure means pressure in the range of preferably about 100 to about 400 psi, e.g., about 150 to about 250 psi. In another aspect, high temperature means temperatures in the range of about 100 to about 300°C, e.g., about 140 to about 200°C. In a preferred aspect, mechanical or physical pretreatment is performed in a batch-process using a steam gun hydrolyzer system that uses high pressure and high temperature as defined above, e.g., a Sunds Hydrolyzer available from Sunds Defibrator AB, Sweden. The physical and chemical pretreatments can be carried out sequentially or simultaneously, as desired.

[0227] Accordingly, in a preferred aspect, the cellulosic material is subjected to physical (mechanical) or chemical pretreatment, or any combination thereof, to promote the separation and/or release of cellulose, hemicellulose, and/or lignin.

[0228] Biological Pretreatment: The term "biological pretreatment" refers to any biological pretreatment that promotes the separation and/or release of cellulose, hemicellulose, and/or lignin from the cellulosic material. Biological pretreatment techniques can involve applying lignin-solubilizing microorganisms and/or enzymes (see, for example, Hsu, T.-A., 1996, Pretreatment of biomass, in *Handbook on Bioethanol: Production and Utilization*, Wyman, C. E., ed., Taylor & Francis, Washington, DC, 179-212; Ghosh and Singh, 1993, Physicochemical and biological treatments for enzymatic/microbial conversion of cellulosic biomass, *Adv. Appl. Microbiol.* 39: 295-333; McMillan, J. D., 1994, Pretreating lignocellulosic biomass: a review, in *Enzymatic Conversion of Biomass for Fuels Production*, Himmel, M. E., Baker, J. O., and Overend, R. P., eds., ACS Symposium Series 566, American Chemical Society, Washington, DC, chapter 15; Gong, C. S., Cao, N. J., Du, J., and Tsao, G. T., 1999, Ethanol production from renewable resources, in *Advances in Biochemical Engineering/Biotechnology*, Scheper, T., ed., Springer-Verlag Berlin Heidelberg, Germany, 65: 207-241; Olsson and Hahn-Hagerdal, 1996, Fermentation of lignocellulosic hydrolysates for ethanol production, *Enz. Microb. Tech.* 18: 312-331; and Vallander and Eriksson, 1990, Production of ethanol from lignocellulosic materials: State of the art, *Adv. Biochem. Eng./Biotechnol.* 42: 63-95).

[0229] Saccharification. In the hydrolysis step, also known as saccharification, the cellulosic material, e.g., pretreated, is hydrolyzed to break down cellulose and/or hemicellulose to fermentable sugars, such as glucose, cellobiose, xylose, xylulose, arabinose, mannose, galactose, and/or soluble oligosaccharides. The hydrolysis is performed enzymatically by an enzyme composition in the presence of a polypeptide having beta-glucosidase activity of the present invention. The enzymes of the compositions can be added simultaneously or sequentially.

[0230] Enzymatic hydrolysis is preferably carried out in a suitable aqueous environment under conditions that can be readily determined by one skilled in the art. In a preferred aspect, hydrolysis is performed under conditions suitable for the activity of the enzyme(s), *i.e.*, optimal for the enzyme(s). The hydrolysis can be carried out as a fed batch or continuous process where the pretreated cellulosic material is fed gradually to, for example, an enzyme containing hydrolysis solution.

[0231] The saccharification is generally performed in stirred-tank reactors or fermentors under controlled pH, temperature, and mixing conditions. Suitable process time, temperature and pH conditions can readily be determined by one skilled in the art. For example, the saccharification can last up to 200 hours, but is typically performed for preferably about 12 to about 120 hours, e.g., about 16 to about 72 hours or about 24 to about 48 hours. The temperature is in the range of preferably about 25°C to about 70°C, e.g., about 30°C to about 65°C, about 40°C to about 60°C, or about 50°C to about 55°C. The pH is in the range of preferably about 3 to about 8, e.g., about 3.5 to about 7, about 4 to about 6, or about 5.0 to about 5.5. The dry solids content is in the range of preferably about 5 to about 50 wt %, e.g., about 10 to about 40 wt % or about 20 to about 30 wt %.

[0232] The enzyme compositions can comprise any protein useful in degrading the cellulosic material.

[0233] In one aspect, the enzyme composition comprises or further comprises one or more (*e.g.*, several) proteins selected from the group consisting of a cellulase, a GH61 polypeptide having cellulolytic enhancing activity, a hemicellulase, an esterase, an expansin, a laccase, a ligninolytic enzyme, a pectinase, a peroxidase, a protease, and a swollenin. In another aspect, the cellulase is preferably one or more (*e.g.*, several) enzymes selected from the group consisting of an endoglucanase, a cellobiohydrolase, and a beta-glucosidase. In another aspect, the hemicellulase is preferably one or more (*e.g.*, several) enzymes selected from the group consisting of an acetylmannan esterase, an acetylxylan esterase, an arabinanase, an arabinofuranosidase, a coumaric acid esterase, a feruloyl esterase, a galactosidase, a glucuronidase, a glucuronoyl esterase, a mannanase, a mannosidase, a xylanase, and a xylosidase.

[0234] In another aspect, the enzyme composition comprises one or more (*e.g.*, several) cellulolytic enzymes. In another aspect, the enzyme composition comprises or further comprises one or more (*e.g.*, several) hemicellulolytic enzymes. In another aspect, the enzyme composition comprises one or more (*e.g.*, several) cellulolytic enzymes and one or more (*e.g.*, several) hemicellulolytic enzymes. In another aspect, the enzyme composition comprises one or more (*e.g.*, several) enzymes selected from the group of cellulolytic enzymes and hemicellulolytic enzymes. In another aspect, the enzyme composition comprises an endoglucanase. In another aspect, the enzyme composition comprises a cellobiohydrolase. In another aspect, the enzyme composition comprises a beta-glucosidase. In another aspect, the enzyme composition comprises a polypeptide having cellulolytic enhancing activity. In another aspect, the enzyme composition comprises an endoglucanase and a polypeptide having cellulolytic enhancing activity. In another aspect, the enzyme composition comprises a cellobiohydrolase and a polypeptide having cellulolytic enhancing activity. In another aspect, the enzyme composition comprises a beta-glucosidase and a polypeptide having cellulolytic enhancing activity. In another aspect, the enzyme composition comprises an endoglucanase and a cellobiohydrolase. In another aspect, the enzyme composition comprises an endoglucanase and a beta-glucosidase. In another aspect, the enzyme composition comprises a cellobiohydrolase and a beta-glucosidase. In another aspect, the enzyme composition comprises an endoglucanase, a cellobiohydrolase, and a polypeptide having cellulolytic enhancing activity. In another aspect, the enzyme composition comprises an endoglucanase, a beta-glucosidase, and a polypeptide having cellulolytic enhancing activity. In another aspect, the enzyme composition comprises a cellobiohydrolase, a beta-glucosidase, and a polypeptide having cellulolytic enhancing activity. In another aspect, the enzyme composition comprises an endoglucanase, a cellobiohydrolase, a beta-glucosidase, and a polypeptide having cellulolytic enhancing activity.

[0235] In another aspect, the enzyme composition comprises an acetylmannan esterase. In another aspect, the enzyme composition comprises an acetylxylan esterase. In another aspect, the enzyme composition comprises an arabinanase (*e.g.*, alpha-L-arabinanase). In another aspect, the enzyme composition comprises an arabinofuranosidase (*e.g.*, alpha-L-arabinofuranosidase). In another aspect, the enzyme composition comprises a coumaric acid esterase. In another aspect, the enzyme composition comprises a feruloyl esterase. In another aspect, the enzyme composition comprises a galactosidase (*e.g.*, alpha-galactosidase and/or beta-galactosidase). In another aspect, the enzyme composition comprises a glucuronidase (*e.g.*, alpha-D-glucuronidase). In another aspect, the enzyme composition comprises a glucuronoyl esterase. In another aspect, the enzyme composition comprises a mannanase. In another aspect, the enzyme composition comprises a mannosidase (*e.g.*, beta-

mannosidase). In another aspect, the enzyme composition comprises a xylanase. In a preferred aspect, the xylanase is a Family 10 xylanase. In another aspect, the enzyme composition comprises a xylosidase (*e.g.*, beta-xylosidase).

[0236] In another aspect, the enzyme composition comprises an esterase. In another aspect, the enzyme composition comprises an expansin. In another aspect, the enzyme composition comprises a laccase. In another aspect, the enzyme composition comprises a ligninolytic enzyme. In a preferred aspect, the ligninolytic enzyme is a manganese peroxidase. In another preferred aspect, the ligninolytic enzyme is a lignin peroxidase. In another preferred aspect, the ligninolytic enzyme is a H₂O₂-producing enzyme. In another aspect, the enzyme composition comprises a pectinase. In another aspect, the enzyme composition comprises a peroxidase. In another aspect, the enzyme composition comprises a protease. In another aspect, the enzyme composition comprises a swollenin

[0237] In the methods of the present invention, the enzyme(s) can be added prior to or during fermentation, *e.g.*, during saccharification or during or after propagation of the fermenting microorganism(s).

[0238] One or more (*e.g.*, several) components of the enzyme composition may be wild-type proteins, recombinant proteins, or a combination of wild-type proteins and recombinant proteins. For example, one or more (*e.g.*, several) components may be native proteins of a cell, which is used as a host cell to express recombinantly one or more (*e.g.*, several) other components of the enzyme composition. One or more (*e.g.*, several) components of the enzyme composition may be produced as monocomponents, which are then combined to form the enzyme composition. The enzyme composition may be a combination of multicomponent and monocomponent protein preparations.

[0239] The enzymes used in the methods of the present invention may be in any form suitable for use, such as, for example, a crude fermentation broth with or without cells removed, a cell lysate with or without cellular debris, a semi-purified or purified enzyme preparation, or a host cell as a source of the enzymes. The enzyme composition may be a dry powder or granulate, a non-dusting granulate, a liquid, a stabilized liquid, or a stabilized protected enzyme. Liquid enzyme preparations may, for instance, be stabilized by adding stabilizers such as a sugar, a sugar alcohol or another polyol, and/or lactic acid or another organic acid according to established processes.

[0240] The optimum amounts of the enzymes and polypeptides having beta-glucosidase activity depend on several factors including, but not limited to, the mixture of component cellulolytic enzymes, the cellulosic material, the concentration of cellulosic material, the pretreatment(s) of the cellulosic material, temperature, time, pH, and inclusion of fermenting organism (*e.g.*, yeast for Simultaneous Saccharification and Fermentation).

[0241] In a preferred aspect, an effective amount of cellulolytic or hemicellulolytic enzyme to the cellulosic material is about 0.5 to about 50 mg, preferably at about 0.5 to about 40 mg, more preferably at about 0.5 to about 25 mg, more preferably at about 0.75 to about 20 mg, more preferably at about 0.75 to about 15 mg, even more preferably at about 0.5 to about 10 mg, and most preferably at about 2.5 to about 10 mg per g of the cellulosic material.

[0242] In another preferred aspect, an effective amount of a polypeptide having beta-glucosidase activity to the cellulosic material is about 0.01 to about 50.0 mg, preferably about 0.01 to about 40 mg, more preferably about 0.01 to about 30 mg, more preferably about 0.01 to about 20 mg, more preferably about 0.01 to about 10 mg, more preferably about 0.01 to about 5 mg, more preferably at about 0.025 to about 1.5 mg, more preferably at about 0.05 to about 1.25 mg, more preferably at about 0.075 to about 1.25 mg, more preferably at about 0.1 to about 1.25 mg, even more preferably at about 0.15 to about 1.25 mg, and most preferably at about 0.25 to about 1.0 mg per g of the cellulosic material.

[0243] In another preferred aspect, an effective amount of polypeptide(s) having beta-glucosidase activity to cellulolytic enzyme(s) is about 0.005 to about 1.0 g, preferably at about 0.01 to about 1.0 g, more preferably at about 0.15 to about 0.75 g, more preferably at about 0.15 to about 0.5 g, more preferably at about 0.1 to about 0.5 g, even more preferably at about 0.1 to about 0.5 g, and most preferably at about 0.05 to about 0.2 g per g of cellulolytic or hemicellulolytic enzyme.

[0244] The polypeptides having cellulolytic enzyme activity or hemicellulolytic enzyme activity as well as other proteins/polypeptides useful in the degradation of the cellulosic material, *e.g.*, GH61 polypeptides having cellulolytic enhancing activity (collectively hereinafter "polypeptides having enzyme activity") can be derived or obtained from any suitable origin, including, bacterial, fungal, yeast, plant, or mammalian origin. The term "obtained" also means herein that the enzyme may have been produced recombinantly in a host organism employing methods described herein, wherein the recombinantly produced enzyme is either native or foreign to the host organism or has a modified amino acid sequence, *e.g.*, having one or more (*e.g.*, several) amino acids that are deleted, inserted and/or substituted, *i.e.*, a recombinantly produced enzyme that is a mutant and/or

a fragment of a native amino acid sequence or an enzyme produced by nucleic acid shuffling processes known in the art. Encompassed within the meaning of a native enzyme are natural variants and within the meaning of a foreign enzyme are variants obtained recombinantly, such as by site-directed mutagenesis or shuffling.

[0245] A polypeptide having enzyme activity may be a bacterial polypeptide. For example, the polypeptide may be a gram positive bacterial polypeptide such as a *Bacillus*, *Streptococcus*, *Streptomyces*, *Staphylococcus*, *Enterococcus*, *Lactobacillus*, *Lactococcus*, *Clostridium*, *Geobacillus*, *Caldicellulosiruptor*, *Acidothermus*, *Thermobifidia*, or *Oceanobacillus* polypeptide having enzyme activity, or a Gram negative bacterial polypeptide such as an *E. coli*, *Pseudomonas*, *Salmonella*, *Campylobacter*, *Helicobacter*, *Flavobacterium*, *Fusobacterium*, *Ilyobacter*, *Neisseria*, or *Ureaplasma* polypeptide having enzyme activity.

[0246] In one aspect, the polypeptide is a *Bacillus alkalophilus*, *Bacillus amyloliquefaciens*, *Bacillus brevis*, *Bacillus circulans*, *Bacillus clausii*, *Bacillus coagulans*, *Bacillus firmus*, *Bacillus lautus*, *Bacillus lentus*, *Bacillus licheniformis*, *Bacillus megaterium*, *Bacillus pumilus*, *Bacillus stearothermophilus*, *Bacillus subtilis*, or *Bacillus thuringiensis* polypeptide having enzyme activity.

[0247] In another aspect, the polypeptide is a *Streptococcus equisimilis*, *Streptococcus pyogenes*, *Streptococcus uberis*, or *Streptococcus equi* subsp. *Zooepidemicus* polypeptide having enzyme activity.

[0248] In another aspect, the polypeptide is a *Streptomyces achromogenes*, *Streptomyces avermitilis*, *Streptomyces coelicolor*, *Streptomyces griseus*, or *Streptomyces lividans* polypeptide having enzyme activity.

[0249] The polypeptide having enzyme activity may also be a fungal polypeptide, and more preferably a yeast polypeptide such as a *Candida*, *Kluyveromyces*, *Pichia*, *Saccharomyces*, *Schizosaccharomyces*, or *Yarrowia* polypeptide having enzyme activity; or more preferably a filamentous fungal polypeptide such as an *Acremonium*, *Agaricus*, *Alternaria*, *Aspergillus*, *Aureobasidium*, *Botryosphaeria*, *Ceriporiopsis*, *Chaetomidium*, *Chrysosporium*, *Claviceps*, *Cochliobolus*, *Coprinopsis*, *Coptotermes*, *Corynascus*, *Cryphonectria*, *Cryptococcus*, *Diplodia*, *Exidia*, *Filibasidium*, *Fusarium*, *Gibberella*, *Holomastigotoides*, *Humicola*, *Irpex*, *Lentinula*, *Leptosphaeria*, *Magnaporthe*, *Melanocarpus*, *Meripilus*, *Mucor*, *Myceliophthora*, *Neocallimastix*, *Neurospora*, *Paecilomyces*, *Penicillium*, *Phanerochaete*, *Piromyces*, *Poitrasia*, *Pseudoplectania*, *Pseudotrichonympha*, *Rhizomucor*, *Schizophyllum*, *Scytalidium*, *Talaromyces*, *Thermoascus*, *Thielavia*, *Tolypocladium*, *Trichoderma*, *Trichophaea*, *Verticillium*, *Volvariella*, or *Xylaria* polypeptide having enzyme activity.

[0250] In one aspect, the polypeptide is a *Saccharomyces carlsbergensis*, *Saccharomyces cerevisiae*, *Saccharomyces diastaticus*, *Saccharomyces douglasii*, *Saccharomyces kluyveri*, *Saccharomyces norbensis*, or *Saccharomyces oviformis* polypeptide having enzyme activity.

[0251] In another aspect, the polypeptide is an *Acremonium cellulolyticus*, *Aspergillus aculeatus*, *Aspergillus awamori*, *Aspergillus fumigatus*, *Aspergillus foetidus*, *Aspergillus japonicus*, *Aspergillus nidulans*, *Aspergillus niger*, *Aspergillus oryzae*, *Chrysosporium keratinophilum*, *Chrysosporium lucknowense*, *Chrysosporium tropicum*, *Chrysosporium merdarium*, *Chrysosporium inops*, *Chrysosporium pannicola*, *Chrysosporium queenslandicum*, *Chrysosporium zonatum*, *Fusarium bactridioides*, *Fusarium cerealis*, *Fusarium crookwellense*, *Fusarium culmorum*, *Fusarium graminearum*, *Fusarium graminum*, *Fusarium heterosporum*, *Fusarium negundi*, *Fusarium oxysporum*, *Fusarium reticulatum*, *Fusarium roseum*, *Fusarium sambucinum*, *Fusarium sarcochroum*, *Fusarium sporotrichioides*, *Fusarium sulphureum*, *Fusarium torulosum*, *Fusarium trichothecioides*, *Fusarium venenatum*, *Humicola grisea*, *Humicola insolens*, *Humicola lanuginosa*, *Irpex lacteus*, *Mucor miehei*, *Myceliophthora thermophila*, *Neurospora crassa*, *Penicillium funiculosum*, *Penicillium purpurogenum*, *Phanerochaete chrysosporium*, *Thielavia achromatica*, *Thielavia albomyces*, *Thielavia albopilosa*, *Thielavia australeinsis*, *Thielavia fimeti*, *Thielavia microspora*, *Thielavia ovispora*, *Thielavia peruviana*, *Thielavia spededonium*, *Thielavia setosa*, *Thielavia subthermophila*, *Thielavia terrestris*, *Trichoderma harzianum*, *Trichoderma koningii*, *Trichoderma longibrachiatum*, *Trichoderma reesei*, *Trichoderma viride*, or *Trichophaea saccata* polypeptide having enzyme activity.

[0252] Chemically modified or protein engineered mutants of polypeptides having enzyme activity may also be used.

[0253] One or more (e.g., several) components of the enzyme composition may be a recombinant component, i.e., produced by cloning of a DNA sequence encoding the single component and subsequent cell transformed with the DNA sequence and expressed in a host (see, for example, WO 91/17243 and WO 91/17244). The host is preferably a heterologous host (enzyme is foreign to host), but the host may under certain conditions also be a homologous host (enzyme is native to host). Monocomponent cellulolytic proteins may also be prepared by purifying such a protein from a fermentation broth.

[0254] In one aspect, the one or more (e.g., several) cellulolytic enzymes comprise a commercial cellulolytic enzyme preparation. Examples of commercial cellulolytic enzyme preparations suitable for use in the present invention include, for example, CELLIC™

CTec (Novozymes A/S), CELLIC™ CTec2 (Novozymes A/S), CELLUCLAST™ (Novozymes A/S), NOVOZYM™ 188 (Novozymes A/S), CELLUZYME™ (Novozymes A/S), CEREFLO™ (Novozymes A/S), and ULTRAFLO™ (Novozymes A/S), ACCELERASE™ (Genencor Int.), LAMINEX™ (Genencor Int.), SPEZYME™ CP (Genencor Int.), FILTRASE® NL (DSM); METHAPLUS® S/L 100 (DSM), ROHAMENT™ 7069 W (Röhm GmbH), FIBREZYME® LDI (Dyadic International, Inc.), FIBREZYME® LBR (Dyadic International, Inc.), or VISCOSTAR® 150L (Dyadic International, Inc.). The cellulase enzymes are added in amounts effective from about 0.001 to about 5.0 wt % of solids, more preferably from about 0.025 to about 4.0 wt % of solids, and most preferably from about 0.005 to about 2.0 wt % of solids. The cellulase enzymes are added in amounts effective from about 0.001 to about 5.0 wt % of solids, more preferably from about 0.025 to about 4.0 wt % of solids, and most preferably from about 0.005 to about 2.0 wt % of solids.

[0255] Examples of bacterial endoglucanases that can be used in the methods of the present invention, include, but are not limited to, an *Acidothermus cellulolyticus* endoglucanase (WO 91/05039; WO 93/15186; U.S. Patent No. 5,275,944; WO 96/02551; U.S. Patent No. 5,536,655, WO 00/70031, WO 05/093050); *Thermobifida fusca* endoglucanase III (WO 05/093050); and *Thermobifida fusca* endoglucanase V (WO 05/093050).

[0256] Examples of fungal endoglucanases that can be used in the present invention, include, but are not limited to, a *Trichoderma reesei* endoglucanase I (Penttilä et al., 1986, Gene 45: 253-263, *Trichoderma reesei* Cel7B endoglucanase I (GENBANK™ accession no. M15665), *Trichoderma reesei* endoglucanase II (Saloheimo, et al., 1988, Gene 63:11-22), *Trichoderma reesei* Cel5A endoglucanase II (GENBANK™ accession no. M19373), *Trichoderma reesei* endoglucanase III (Okada et al., 1988, Appl. Environ. Microbiol. 64: 555-563, GENBANK™ accession no. AB003694), *Trichoderma reesei* endoglucanase V (Saloheimo et al., 1994, Molecular Microbiology 13: 219-228, GENBANK™ accession no. Z33381), *Aspergillus aculeatus* endoglucanase (Ooi et al., 1990, Nucleic Acids Research 18: 5884), *Aspergillus kawachii* endoglucanase (Sakamoto et al., 1995, Current Genetics 27: 435-439), *Erwinia carotovora* endoglucanase (Saarilahti et al., 1990, Gene 90: 9-14), *Fusarium oxysporum* endoglucanase (GENBANK™ accession no. L29381), *Humicola grisea* var. *thermoidea* endoglucanase (GENBANK™ accession no. AB003107), *Melanocarpus albomyces* endoglucanase (GENBANK™ accession no. MAL515703), *Neurospora crassa* endoglucanase (GENBANK™ accession no. XM_324477), *Humicola insolens* endoglucanase V, *Myceliophthora thermophila* CBS 117.65 endoglucanase, basidiomycete CBS 495.95 endoglucanase, basidiomycete CBS 494.95 endoglucanase, *Thielavia terrestris* NRRL 8126 CEL6B endoglucanase, *Thielavia terrestris* NRRL 8126 CEL6C endoglucanase, *Thielavia terrestris* NRRL 8126 CEL7C endoglucanase, *Thielavia terrestris* NRRL 8126 CEL7E endoglucanase, *Thielavia terrestris* NRRL 8126 CEL7F endoglucanase, *Cladorrhinum foecundissimum* ATCC 62373 CEL7A endoglucanase, and *Trichoderma reesei* strain No. VTT-D-80133 endoglucanase (GENBANK™ accession no. M15665).

[0257] Examples of cellobiohydrolases useful in the present invention include, but are not limited to, *Aspergillus aculeatus* cellobiohydrolase II (WO 2011/059740), *Chaetomium thermophilum* cellobiohydrolase I, *Chaetomium thermophilum* cellobiohydrolase II, *Humicola insolens* cellobiohydrolase I, *Myceliophthora thermophila* cellobiohydrolase II (WO 2009/042871), *Thielavia hyrcanie* cellobiohydrolase II (WO 2010/141325), *Thielavia terrestris* cellobiohydrolase II (CEL6A, WO 2006/074435), *Trichoderma reesei* cellobiohydrolase I, *Trichoderma reesei* cellobiohydrolase II, and *Trichophaea saccata* cellobiohydrolase II (WO 2010/057086).

[0258] Examples of beta-glucosidases useful in the present invention include, but are not limited to, beta-glucosidases from *Aspergillus aculeatus* (Kawaguchi et al., 1996, Gene 173: 287-288), *Aspergillus fumigatus* (WO 2005/047499), *Aspergillus niger* (Dan et al., 2000, J. Biol. Chem. 275: 4973-4980), *Aspergillus oryzae* (WO 2002/095014), *Penicillium brasilianum* IBT 20888 (WO 2007/019442 and WO 2010/088387), *Thielavia terrestris* (WO 2011/035029), and *Trichophaea saccata* (WO 2007/019442).

[0259] The beta-glucosidase may be a fusion protein. In one aspect, the beta-glucosidase is an *Aspergillus oryzae* beta-glucosidase variant BG fusion protein (WO 2008/057637) or an *Aspergillus oryzae* beta-glucosidase fusion protein (WO 2008/057637).

[0260] Other useful endoglucanases, cellobiohydrolases, and beta-glucosidases are disclosed in numerous Glycosyl Hydrolase families using the classification according to Henrissat B., 1991, A classification of glycosyl hydrolases based on amino-acid sequence similarities, Biochem. J. 280: 309-316, and Henrissat B., and Bairoch A., 1996, Updating the sequence-based classification of glycosyl hydrolases, Biochem. J. 316: 695-696.

[0261] Other cellulolytic enzymes that may be used in the present invention are described in EP 495,257, EP 531,315, EP 531,372, WO 89/09259, WO 94/07998, WO 95/24471, WO 96/11262, WO 96/29397, WO 96/034108, WO 97/14804, WO 98/08940, WO 98/012307, WO 98/13465, WO 98/015619, WO 98/015633, WO 98/028411, WO 99/06574, WO 99/10481, WO 99/025846, WO 99/025847, WO 99/031255, WO 2000/009707, WO 2002/050245, WO 2002/0076792, WO 2002/101078, WO

2003/027306, WO 2003/052054, WO 2003/052055, WO 2003/052056, WO 2003/052057, WO 2003/052118, WO 2004/016760, WO 2004/043980, WO 2004/048592, WO 2005/001065, WO 2005/028636, WO 2005/093050, WO 2005/093073, WO 2006/074005, WO 2006/117432, WO 2007/071818, WO 2007/071820, WO 2008/008070, WO 2008/008793, U.S. Patent No. 4,435,307, U.S. Patent No. 5,457,046, U.S. Patent No. 5,648,263, U.S. Patent No. 5,686,593, U.S. Patent No. 5,691,178, U.S. Patent No. 5,763,254, and U.S. Patent No. 5,776,757.

[0262] In the methods of the present invention, any GH61 polypeptide having cellulolytic enhancing activity can be used.

[0263] In a first aspect, the polypeptide having cellulolytic enhancing activity comprises the following motifs:

[ILMV]-P-X(4,5)-G-X-Y-[ILMV]-X-R-X-[EQ]-X(4)-[HNQ] (SEQ ID NO: 19 or SEQ ID NO: 20) and [FW]-[TF]-K-[AIV],

wherein X is any amino acid, X(4,5) is any amino acid at 4 or 5 contiguous positions, and X(4) is any amino acid at 4 contiguous positions.

[0264] The isolated polypeptide comprising the above-noted motifs may further comprise:

H-X(1,2)-G-P-X(3)-[YW]-[AILMV] (SEQ ID NO: 21 or SEQ ID NO: 22),

[EQ]-X-Y-X(2)-C-X-[EHQN]-[FILV]-X-[ILV] (SEQ ID NO: 23), or

H-X(1,2)-G-P-X(3)-[YW]-[AILMV] (SEQ ID NO: 24 or SEQ ID NO: 25) and [EQ]-X-Y-X(2)-C-X-[EHQN]-[FILV]-X-[ILV] (SEQ ID NO: 26),

wherein X is any amino acid, X(1,2) is any amino acid at 1 position or 2 contiguous positions, X(3) is any amino acid at 3 contiguous positions, and X(2) is any amino acid at 2 contiguous positions. In the above motifs, the accepted IUPAC single letter amino acid abbreviation is employed.

[0265] In a preferred embodiment, the isolated GH61 polypeptide having cellulolytic enhancing activity further comprises H-X(1,2)-G-P-X(3)-[YW]-[AILMV] (SEQ ID NO: 27 or SEQ ID NO: 28). In another preferred embodiment, the isolated GH61 polypeptide having cellulolytic enhancing activity further comprises [EQ]-X-Y-X(2)-C-X-[EHQN]-[FILV]-X-[ILV] (SEQ ID NO: 29). In another preferred embodiment, the isolated GH61 polypeptide having cellulolytic enhancing activity further comprises H-X(1,2)-G-P-X(3)-[YW]-[AILMV] (SEQ ID NO: 30 or SEQ ID NO: 31) and [EQ]-X-Y-X(2)-C-X-[EHQN]-[FILV]-X-[ILV] (SEQ ID NO: 32).

[0266] In a second aspect, isolated polypeptides having cellulolytic enhancing activity, comprise the following motif:

[ILMV]-P-X(4,5)-G-X-Y-[ILMV]-X-R-X-[EQ]-X(3)-A-[HNQ] (SEQ ID NO: 33 or SEQ ID NO: 34),

wherein X is any amino acid, X(4,5) is any amino acid at 4 or 5 contiguous positions, and X(3) is any amino acid at 3 contiguous positions. In the above motif, the accepted IUPAC single letter amino acid abbreviation is employed.

[0267] Examples of GH61 polypeptides having cellulolytic enhancing activity useful in the processes of the present invention include, but are not limited to, GH61 polypeptides from *Thielavia terrestris* (WO 2005/074647, WO 2008/148131, and WO 2011/035027), *Thermoascus aurantiacus* (WO 2005/074656 and WO 2010/065830), *Trichoderma reesei* (WO 2007/089290), *Myceliophthora thermophila* (WO 2009/085935, WO 2009/085859, WO 2009/085864, WO 2009/085868), *Aspergillus fumigatus* (WO 2010/138754), GH61 polypeptides from *Penicillium pinophilum* (WO 2011/005867), *Thermoascus* sp. (WO 2011/039319), *Penicillium* sp. (WO 2011/041397), and *Thermoascus crustaceus* (WO 2011/041504).

[0268] In one aspect, the GH61 polypeptide having cellulolytic enhancing activity is used in the presence of a soluble activating divalent metal cation according to WO 2008/151043, e.g., manganese sulfate.

[0269] In one aspect, the GH61 polypeptide having cellulolytic enhancing activity is used in the presence of a dioxy compound, a bicyclic compound, a heterocyclic compound, a nitrogen-containing compound, a quinone compound, a sulfur-containing compound, or a liquor obtained from a pretreated cellulosic material such as pretreated corn stover (PCS).

[0270] The dioxy compound may include any suitable compound containing two or more oxygen atoms. In some aspects, the dioxy compounds contain a substituted aryl moiety as described herein. The dioxy compounds may comprise one or more (e.g., several) hydroxyl and/or hydroxyl derivatives, but also include substituted aryl moieties lacking hydroxyl and hydroxyl derivatives. Non-limiting examples of the dioxy compounds include pyrocatechol or catechol; caffeic acid; 3,4-dihydroxybenzoic acid; 4-tert-butyl-5-methoxy-1,2-benzenediol; pyrogallol; gallic acid; methyl-3,4,5-trihydroxybenzoate; 2,3,4-trihydroxybenzophenone; 2,6-

dimethoxyphenol; sinapinic acid; 3,5-dihydroxybenzoic acid; 4-chloro-1,2-benzenediol; 4-nitro-1,2-benzenediol; tannic acid; ethyl gallate; methyl glycolate; dihydroxyfumaric acid; 2-butyne-1,4-diol; (croconic acid; 1,3-propanediol; tartaric acid; 2,4-pentanediol; 3-ethoxy-1,2-propanediol; 2,4,4'-trihydroxybenzophenone; cis-2-butene-1,4-diol; 3,4-dihydroxy-3-cyclobutene-1,2-dione; dihydroxyacetone; acrolein acetal; methyl-4-hydroxybenzoate; 4-hydroxybenzoic acid; and methyl-3,5-dimethoxy-4-hydroxybenzoate; or a salt or solvate thereof.

[0271] The bicyclic compound may include any suitable substituted fused ring system as described herein. The compounds may comprise one or more (e.g., several) additional rings, and are not limited to a specific number of rings unless otherwise stated. In one aspect, the bicyclic compound is a flavonoid. In another aspect, the bicyclic compound is an optionally substituted isoflavonoid. In another aspect, the bicyclic compound is an optionally substituted flavylum ion, such as an optionally substituted anthocyanidin or optionally substituted anthocyanin, or derivative thereof. Non-limiting examples of the bicyclic compounds include epicatechin; quercetin; myricetin; taxifolin; kaempferol; morin; acacetin; naringenin; isorhamnetin; apigenin; cyanidin; cyanin; kuromanin; keracyanin; or a salt or solvate thereof.

[0272] The heterocyclic compound may be any suitable compound, such as an optionally substituted aromatic or non-aromatic ring comprising a heteroatom, as described herein. In one aspect, the heterocyclic is a compound comprising an optionally substituted heterocycloalkyl moiety or an optionally substituted heteroaryl moiety. In another aspect, the optionally substituted heterocycloalkyl moiety or optionally substituted heteroaryl moiety is an optionally substituted 5-membered heterocycloalkyl or an optionally substituted 5-membered heteroaryl moiety. In another aspect, the optionally substituted heterocycloalkyl or optionally substituted heteroaryl moiety is an optionally substituted moiety selected from pyrazolyl, furanyl, imidazolyl, isoxazolyl, oxadiazolyl, oxazolyl, pyrrolyl, pyridyl, pyrimidyl, pyridazinyl, thiazolyl, triazolyl, thienyl, dihydrothieno-pyrazolyl, thianaphthenyl, carbazolyl, benzimidazolyl, benzothienyl, benzofuranyl, indolyl, quinolyl, benzotriazolyl, benzothiazolyl, benzooxazolyl, benzimidazolyl, isoquinolyl, isoindolyl, acridinyl, benzoisazolyl, dimethylhydantoin, pyrazinyl, tetrahydrofuranyl, pyrrolinyl, pyrrolidinyl, morpholinyl, indolyl, diazepinyl, azepinyl, thiepinyl, piperidinyl, and oxepinyl. In another aspect, the optionally substituted heterocycloalkyl moiety or optionally substituted heteroaryl moiety is an optionally substituted furanyl. Non-limiting examples of the heterocyclic compounds include (1,2-dihydroxyethyl)-3,4-dihydroxyfuran-2(5H)-one; 4-hydroxy-5-methyl-3-furanone; 5-hydroxy-2(5H)-furanone; [1,2-dihydroxyethyl]furan-2,3,4(5H)-trione; α -hydroxy- γ -butyrolactone; ribonic γ -lactone; aldohexuronicaldohexuronic acid γ -lactone; gluconic acid δ -lactone; 4-hydroxycoumarin; dihydrobenzofuran; 5-(hydroxymethyl)furfural; furin; 2(5H)-furanone; 5,6-dihydro-2H-pyran-2-one; and 5,6-dihydro-4-hydroxy-6-methyl-2H-pyran-2-one; or a salt or solvate thereof.

[0273] The nitrogen-containing compound may be any suitable compound with one or more nitrogen atoms. In one aspect, the nitrogen-containing compound comprises an amine, imine, hydroxylamine, or nitroxide moiety. Non-limiting examples of the nitrogen-containing compounds include acetone oxime; violuric acid; pyridine-2-aldoxime; 2-aminophenol; 1,2-benzenediamine; 2,2,6,6-tetramethyl-1-piperidinyl-oxo; 5,6,7,8-tetrahydrobiopterin; 6,7-dimethyl-5,6,7,8-tetrahydropterine; and maleamic acid; or a salt or solvate thereof.

[0274] The quinone compound may be any suitable compound comprising a quinone moiety as described herein. Non-limiting examples of the quinone compounds include 1,4-benzoquinone; 1,4-naphthoquinone; 2-hydroxy-1,4-naphthoquinone; 2,3-dimethoxy-5-methyl-1,4-benzoquinone or coenzyme Q₀; 2,3,5,6-tetramethyl-1,4-benzoquinone or duroquinone; 1,4-dihydroxyanthraquinone; 3-hydroxy-1-methyl-5,6-indolinedione or adrenochrome; 4-tert-butyl-5-methoxy-1,2-benzoquinone; pyrroloquinoline quinone; or a salt or solvate thereof.

[0275] The sulfur-containing compound may be any suitable compound comprising one or more sulfur atoms. In one aspect, the sulfur-containing comprises a moiety selected from thionyl, thioether, sulfinyl, sulfonyl, sulfamide, sulfonamide, sulfonic acid, and sulfonic ester. Non-limiting examples of the sulfur-containing compounds include ethanethiol; 2-propanethiol; 2-propene-1-thiol; 2-mercaptoethanesulfonic acid; benzenethiol; benzene-1,2-dithiol; cysteine; methionine; glutathione; cystine; or a salt or solvate thereof.

[0276] In one aspect, an effective amount of such a compound described above to cellulosic material as a molar ratio to glucosyl units of cellulose is about 10^{-6} to about 10, e.g., about 10^{-6} to about 7.5, about 10^{-6} to about 5, about 10^{-6} to about 2.5, about 10^{-6} to about 1, about 10^{-5} to about 1, about 10^{-5} to about 10^{-1} , about 10^{-4} to about 10^{-1} , about 10^{-3} to about 10^{-1} , or about 10^{-3} to about 10^{-2} . In another aspect, an effective amount of such a compound described above is about 0.1 μ M to about 1 M, e.g., about 0.5 μ M to about 0.75 M, about 0.75 μ M to about 0.5 M, about 1 μ M to about 0.25 M, about 1 μ M to about 0.1 M, about 5 μ M to about 50 mM, about 10 μ M to about 25 mM, about 50 μ M to about 25 mM, about 10 μ M to about 10 mM, about 5 μ M to about 5 mM, or about 0.1 mM to about 1 mM.

[0277] The term "liquor" means the solution phase, either aqueous, organic, or a combination thereof, arising from treatment of a lignocellulose and/or hemicellulose material in a slurry, or monosaccharides thereof, e.g., xylose, arabinose, mannose, etc., under conditions as described herein, and the soluble contents thereof. A liquor for cellulolytic enhancement of a GH61 polypeptide can be produced by treating a lignocellulose or hemicellulose material (or feedstock) by applying heat and/or pressure, optionally in the presence of a catalyst, e.g., acid, optionally in the presence of an organic solvent, and optionally in combination with physical disruption of the material, and then separating the solution from the residual solids. Such conditions determine the degree of cellulolytic enhancement obtainable through the combination of liquor and a GH61 polypeptide during hydrolysis of a cellulosic substrate by a cellulase preparation. The liquor can be separated from the treated material using a method standard in the art, such as filtration, sedimentation, or centrifugation.

[0278] In one aspect, an effective amount of the liquor to cellulose is about 10^{-6} to about 10 g per g of cellulose, e.g., about 10^{-6} to about 7.5 g, about 10^{-6} to about 5, about 10^{-6} to about 2.5 g, about 10^{-6} to about 1 g, about 10^{-5} to about 1 g, about 10^{-5} to about 10^{-1} g, about 10^{-4} to about 10^{-1} g, about 10^{-3} to about 10^{-1} g, or about 10^{-3} to about 10^{-2} g per g of cellulose.

[0279] In one aspect, the one or more (e.g., several) hemicellulolytic enzymes comprise a commercial hemicellulolytic enzyme preparation. Examples of commercial hemicellulolytic enzyme preparations suitable for use in the present invention include, for example, SHEARZYME™ (Novozymes A/S), CELLIC™ HTec (Novozymes A/S), CELLIC™ HTec2 (Novozymes A/S), VISCOZYME® (Novozymes A/S), ULTRAFLO® (Novozymes A/S), PULPZYME® HC (Novozymes A/S), MULTIFECT® Xylanase (Genencor), ACCELLERASE® XY (Genencor), ACCELLERASE® XC (Genencor), ECOPULP® TX-200A (AB Enzymes), HSP 6000 Xylanase (DSM), DEPOL™ 333P (Biocatalysts Limit, Wales, UK), DEPOL™ 740L (Biocatalysts Limit, Wales, UK), and DEPOL™ 762P (Biocatalysts Limit, Wales, UK).

[0280] Examples of xylanases useful in the methods of the present invention include, but are not limited to, xylanases from *Aspergillus aculeatus* (GeneSeqP:AAR63790; WO 94/21785), *Aspergillus fumigatus* (WO 2006/078256), *Penicillium pinophilum* (WO 2011/041405), *Penicillium* sp. (WO 2010/126772), *Thielavia terrestris* NRRL 8126 (WO 2009/079210), and *Trichophaea saccata* GH10 (WO 2011/057083).

[0281] Examples of beta-xylosidases useful in the methods of the present invention include, but are not limited to, beta-xylosidases from *Neurospora crassa* (SwissProt accession number Q7SOW4), *Trichoderma reesei* (UniProtKB/TrEMBL accession number Q92458), and *Talaromyces emersonii* (SwissProt accession number Q8X212).

[0282] Examples of acetylxyylan esterases useful in the methods of the present invention include, but are not limited to, acetylxyylan esterases from *Aspergillus aculeatus* (WO 2010/108918), *Chaetomium globosum* (Uniprot accession number Q2GWW4), *Chaetomium gracile* (GeneSeqP accession number AAB82124), *Humicola insolens* DSM 1800 (WO 2009/073709), *Hypocrea jecorina* (WO 2005/001036), *Myceliophthora thermophila* (WO 2010/014880), *Neurospora crassa* (UniProt accession number q7s259), *Phaeosphaeria nodorum* (Uniprot accession number Q0UHU1), and *Thielavia terrestris* NRRL 8126 (WO 2009/042846).

[0283] Examples of feruloyl esterases (ferulic acid esterases) useful in the methods of the present invention include, but are not limited to, feruloyl esterases from *Humicola insolens* DSM 1800 (WO 2009/076122), *Neosartorya fischeri* (UniProt Accession number A1D9T4), *Neurospora crassa* (UniProt accession number Q9HGR3), *Penicillium aurantiogriseum* (WO 2009/127729), and *Thielavia terrestris* (WO 2010/053838 and WO 2010/065448).

[0284] Examples of arabinofuranosidases useful in the methods of the present invention include, but are not limited to, arabinofuranosidases from *Aspergillus niger* (GeneSeqP accession number AAR94170), *Humicola insolens* DSM 1800 (WO 2006/114094 and WO 2009/073383), and *M. giganteus* (WO 2006/114094).

[0285] Examples of alpha-glucuronidases useful in the methods of the present invention include, but are not limited to, alpha-glucuronidases from *Aspergillus clavatus* (UniProt accession number alcc12), *Aspergillus fumigatus* (SwissProt accession number Q4VWW45), *Aspergillus niger* (Uniprot accession number Q96WX9), *Aspergillus terreus* (SwissProt accession number Q0CJP9), *Humicola insolens* (WO 2010/014706), *Penicillium aurantiogriseum* (WO 2009/068565), *Talaromyces emersonii* (UniProt accession number Q8X211), and *Trichoderma reesei* (Uniprot accession number Q99024).

[0286] The polypeptides having enzyme activity used in the methods of the present invention may be produced by fermentation of the above-noted microbial strains on a nutrient medium containing suitable carbon and nitrogen sources and inorganic salts, using procedures known in the art (see, e.g., Bennett, J.W. and LaSure, L. (eds.), More Gene Manipulations in Fungi, Academic Press, CA, 1991). Suitable media are available from commercial suppliers or may be prepared according to published

compositions (e.g., in catalogues of the American Type Culture Collection). Temperature ranges and other conditions suitable for growth and enzyme production are known in the art (see, e.g., Bailey, J.E., and Ollis, D.F., *Biochemical Engineering Fundamentals*, McGraw-Hill Book Company, NY, 1986).

[0287] The fermentation can be any method of cultivation of a cell resulting in the expression or isolation of an enzyme or protein. Fermentation may, therefore, be understood as comprising shake flask cultivation, or small- or large-scale fermentation (including continuous, batch, fed-batch, or solid state fermentations) in laboratory or industrial fermentors performed in a suitable medium and under conditions allowing the enzyme to be expressed or isolated. The resulting enzymes produced by the methods described above may be recovered from the fermentation medium and purified by conventional procedures.

[0288] Fermentation. The fermentable sugars obtained from the hydrolyzed cellulosic material can be fermented by one or more (e.g., several) fermenting microorganisms capable of fermenting the sugars directly or indirectly into a desired fermentation product. "Fermentation" or "fermentation process" refers to any fermentation process or any process comprising a fermentation step. Fermentation processes also include fermentation processes used in the consumable alcohol industry (e.g., beer and wine), dairy industry (e.g., fermented dairy products), leather industry, and tobacco industry. The fermentation conditions depend on the desired fermentation product and fermenting organism and can easily be determined by one skilled in the art.

[0289] In the fermentation step, sugars, released from the cellulosic material as a result of the pretreatment and enzymatic hydrolysis steps, are fermented to a product, e.g., ethanol, by a fermenting organism, such as yeast. Hydrolysis (saccharification) and fermentation can be separate or simultaneous, as described herein.

[0290] Any suitable hydrolyzed cellulosic material can be used in the fermentation step in practicing the present invention. The material is generally selected based on the desired fermentation product, i.e., the substance to be obtained from the fermentation, and the process employed, as is well known in the art.

[0291] The term "fermentation medium" is understood herein to refer to a medium before the fermenting microorganism(s) is(are) added, such as, a medium resulting from a saccharification process, as well as a medium used in a simultaneous saccharification and fermentation process (SSF).

[0292] "Fermenting microorganism" refers to any microorganism, including bacterial and fungal organisms, suitable for use in a desired fermentation process to produce a fermentation product. The fermenting organism can be hexose and/or pentose fermenting organisms, or a combination thereof. Both hexose and pentose fermenting organisms are well known in the art. Suitable fermenting microorganisms are able to ferment, i.e., convert, sugars, such as glucose, xylose, xylulose, arabinose, maltose, mannose, galactose, and/or oligosaccharides, directly or indirectly into the desired fermentation product. Examples of bacterial and fungal fermenting organisms producing ethanol are described by Lin et al., 2006, *Appl. Microbiol. Biotechnol.* 69: 627-642.

[0293] Examples of fermenting microorganisms that can ferment hexose sugars include bacterial and fungal organisms, such as yeast. Preferred yeast includes strains of *Candida*, *Kluyveromyces*, and *Saccharomyces*, e.g., *Candida sonorensis*, *Kluyveromyces marxianus*, and *Saccharomyces cerevisiae*.

[0294] Examples of fermenting organisms that can ferment pentose sugars in their native state include bacterial and fungal organisms, such as some yeast. Preferred xylose fermenting yeast include strains of *Candida*, preferably *C. sheatae* or *C. sonorensis*; and strains of *Pichia*, preferably *P. stipitis*, such as *P. stipitis* CBS 5773. Preferred pentose fermenting yeast include strains of *Pachysolen*, preferably *P. tannophilus*. Organisms not capable of fermenting pentose sugars, such as xylose and arabinose, may be genetically modified to do so by methods known in the art.

[0295] Examples of bacteria that can efficiently ferment hexose and pentose to ethanol include, for example, *Bacillus coagulans*, *Clostridium acetobutylicum*, *Clostridium thermocellum*, *Clostridium phytofermentans*, *Geobacillus* sp., *Thermoanaerobacter saccharolyticum*, and *Zymomonas mobilis* (Philippidis, 1996, *supra*).

[0296] Other fermenting organisms include strains of *Bacillus*, such as *Bacillus coagulans*; *Candida*, such as *C. sonorensis*, *C. methanosorbosa*, *C. diddensiae*, *C. parapsilosis*, *C. naedodendra*, *C. blankii*, *C. entomophila*, *C. brassicae*, *C. pseudotropicalis*, *C. boidinii*, *C. utilis*, and *C. sheatae*; *Clostridium*, such as *C. acetobutylicum*, *C. thermocellum*, and *C. phytofermentans*; *E. coli*, especially *E. coli* strains that have been genetically modified to improve the yield of ethanol; *Geobacillus* sp.; *Hansenula*, such as *Hansenula anomala*; *Klebsiella*, such as *K. oxytoca*; *Kluyveromyces*, such as *K. marxianus*, *K. lactis*, *K. thermotolerans*, and *K. fragilis*; *Schizosaccharomyces*, such as *S. pombe*; *Thermoanaerobacter*, such as *Thermoanaerobacter saccharolyticum*; and *Zymomonas*, such as *Zymomonas mobilis*.

[0297] In a preferred aspect, the yeast is a *Bretannomyces*. In a more preferred aspect, the yeast is *Bretannomyces clausenii*. In another preferred aspect, the yeast is a *Candida*. In another more preferred aspect, the yeast is *Candida sonorensis*. In another more preferred aspect, the yeast is *Candida boidinii*. In another more preferred aspect, the yeast is *Candida blankii*. In another more preferred aspect, the yeast is *Candida brassicae*. In another more preferred aspect, the yeast is *Candida diddensii*. In another more preferred aspect, the yeast is *Candida entomophili*. In another more preferred aspect, the yeast is *Candida pseudotropicalis*. In another more preferred aspect, the yeast is *Candida scheidtiae*. In another more preferred aspect, the yeast is *Candida utilis*. In another preferred aspect, the yeast is a *Clavispora*. In another more preferred aspect, the yeast is *Clavispora lusitanae*. In another more preferred aspect, the yeast is *Clavispora opuntiae*. In another preferred aspect, the yeast is a *Kluyveromyces*. In another more preferred aspect, the yeast is *Kluyveromyces fragilis*. In another more preferred aspect, the yeast is *Kluyveromyces marxianus*. In another more preferred aspect, the yeast is *Kluyveromyces thermotolerans*. In another preferred aspect, the yeast is a *Pachysolen*. In another more preferred aspect, the yeast is *Pachysolen tannophilus*. In another preferred aspect, the yeast is a *Pichia*. In another more preferred aspect, the yeast is a *Pichia stipitis*. In another preferred aspect, the yeast is a *Saccharomyces* spp. In a more preferred aspect, the yeast is *Saccharomyces cerevisiae*. In another more preferred aspect, the yeast is *Saccharomyces distaticus*. In another more preferred aspect, the yeast is *Saccharomyces uvarum*.

[0298] In a preferred aspect, the bacterium is a *Bacillus*. In a more preferred aspect, the bacterium is *Bacillus coagulans*. In another preferred aspect, the bacterium is a *Clostridium*. In another more preferred aspect, the bacterium is *Clostridium acetobutylicum*. In another more preferred aspect, the bacterium is *Clostridium phytofermentans*. In another more preferred aspect, the bacterium is *Clostridium thermocellum*. In another more preferred aspect, the bacterium is *Geobacillus* sp. In another more preferred aspect, the bacterium is a *Thermoanaerobacter*. In another more preferred aspect, the bacterium is *Thermoanaerobacter saccharolyticum*. In another preferred aspect, the bacterium is a *Zymomonas*. In another more preferred aspect, the bacterium is *Zymomonas mobilis*.

[0299] Commercially available yeast suitable for ethanol production include, e.g., BIOFERM™ AFT and XR (NABC - North American Bioproducts Corporation, GA, USA), ETHANOL RED™ yeast (Fermentis/Lesaffre, USA), FALI™ (Fleischmann's Yeast, USA), FERMOL™ (DSM Specialties), GERT STRAND™ (Gert Strand AB, Sweden), and SUPERSTART™ and THERMOSACC™ fresh yeast (Ethanol Technology, WI, USA).

[0300] In a preferred aspect, the fermenting microorganism has been genetically modified to provide the ability to ferment pentose sugars, such as xylose utilizing, arabinose utilizing, and xylose and arabinose co-utilizing microorganisms.

[0301] The cloning of heterologous genes into various fermenting microorganisms has led to the construction of organisms capable of converting hexoses and pentoses to ethanol (cofermentation) (Chen and Ho, 1993, Cloning and improving the expression of *Pichia stipitis* xylose reductase gene in *Saccharomyces cerevisiae*, Appl. Biochem. Biotechnol. 39-40: 135-147; Ho et al., 1998, Genetically engineered *Saccharomyces* yeast capable of effectively cofermenting glucose and xylose, Appl. Environ. Microbiol. 64: 1852-1859; Kotter and Ciriacy, 1993, Xylose fermentation by *Saccharomyces cerevisiae*, Appl. Microbiol. Biotechnol. 38: 776-783; Walfridsson et al., 1995, Xylose-metabolizing *Saccharomyces cerevisiae* strains overexpressing the TKL1 and TAL1 genes encoding the pentose phosphate pathway enzymes transketolase and transaldolase, Appl. Environ. Microbiol. 61: 4184-4190; Kuyper et al., 2004, Minimal metabolic engineering of *Saccharomyces cerevisiae* for efficient anaerobic xylose fermentation: a proof of principle, FEMS Yeast Research 4: 655-664; Beall et al., 1991, Parametric studies of ethanol production from xylose and other sugars by recombinant *Escherichia coli*, Biotech. Bioeng. 38: 296-303; Ingram et al., 1998, Metabolic engineering of bacteria for ethanol production, Biotechnol. Bioeng. 58: 204-214; Zhang et al., 1995, Metabolic engineering of a pentose metabolism pathway in ethanologenic *Zymomonas mobilis*, Science 267: 240-243; Deanda et al., 1996, Development of an arabinose-fermenting *Zymomonas mobilis* strain by metabolic pathway engineering, Appl. Environ. Microbiol. 62: 4465-4470; WO 2003/062430, xylose isomerase).

[0302] In a preferred aspect, the genetically modified fermenting microorganism is *Candida sonorensis*. In another preferred aspect, the genetically modified fermenting microorganism is *Escherichia coli*. In another preferred aspect, the genetically modified fermenting microorganism is *Klebsiella oxytoca*. In another preferred aspect, the genetically modified fermenting microorganism is *Kluyveromyces marxianus*. In another preferred aspect, the genetically modified fermenting microorganism is *Saccharomyces cerevisiae*. In another preferred aspect, the genetically modified fermenting microorganism is *Zymomonas mobilis*.

[0303] It is well known in the art that the organisms described above can also be used to produce other substances, as described herein.

[0304] The fermenting microorganism is typically added to the degraded cellulosic material or hydrolysate and the fermentation is performed for about 8 to about 96 hours, e.g., about 24 to about 60 hours. The temperature is typically between about 26°C to about 60°C, e.g., about 32°C or 50°C, and about pH 3 to about pH 8, e.g., pH 4-5, 6, or 7.

[0305] In one aspect, the yeast and/or another microorganism are applied to the degraded cellulosic material and the fermentation is performed for about 12 to about 96 hours, such as typically 24-60 hours. In another aspect, the temperature is preferably between about 20°C to about 60°C, e.g., about 25°C to about 50°C, about 32°C to about 50°C, or about 32°C to about 50°C, and the pH is generally from about pH 3 to about pH 7, e.g., about pH 4 to about pH 7. However, some fermenting organisms, e.g., bacteria, have higher fermentation temperature optima. Yeast or another microorganism is preferably applied in amounts of approximately 10^5 to 10^{12} , preferably from approximately 10^7 to 10^{10} , especially approximately 2×10^8 viable cell count per ml of fermentation broth. Further guidance in respect of using yeast for fermentation can be found in, e.g., "The Alcohol Textbook" (Editors K. Jacques, T.P. Lyons and D.R. Kelsall, Nottingham University Press, United Kingdom 1999).

[0306] For ethanol production, following the fermentation the fermented slurry is distilled to extract the ethanol. The ethanol obtained according to the methods of the invention can be used as, e.g., fuel ethanol, drinking ethanol, i.e., potable neutral spirits, or industrial ethanol.

[0307] A fermentation stimulator can be used in combination with any of the processes described herein to further improve the fermentation process, and in particular, the performance of the fermenting microorganism, such as, rate enhancement and ethanol yield. A "fermentation stimulator" refers to stimulators for growth of the fermenting microorganisms, in particular, yeast. Preferred fermentation stimulators for growth include vitamins and minerals. Examples of vitamins include multivitamins, biotin, pantothenate, nicotinic acid, meso-inositol, thiamine, pyridoxine, para-aminobenzoic acid, folic acid, riboflavin, and Vitamins A, B, C, D, and E. See, for example, Alfenore et al., Improving ethanol production and viability of *Saccharomyces cerevisiae* by a vitamin feeding strategy during fed-batch process, Springer-Verlag (2002). Examples of minerals include minerals and mineral salts that can supply nutrients comprising P, K, Mg, S, Ca, Fe, Zn, Mn, and Cu.

[0308] Fermentation products: A fermentation product can be any substance derived from the fermentation. The fermentation product can be, without limitation, an alcohol (e.g., arabinitol, n-butanol, isobutanol, ethanol, glycerol, methanol, ethylene glycol, 1,3-propanediol [propylene glycol], butanediol, glycerin, sorbitol, and xylitol); an alkane (e.g., pentane, hexane, heptane, octane, nonane, decane, undecane, and dodecane), a cycloalkane (e.g., cyclopentane, cyclohexane, cycloheptane, and cyclooctane), an alkene (e.g. pentene, hexene, heptene, and octene); an amino acid (e.g., aspartic acid, glutamic acid, glycine, lysine, serine, and threonine); a gas (e.g., methane, hydrogen (H_2), carbon dioxide (CO_2), and carbon monoxide (CO)); isoprene; a ketone (e.g., acetone); an organic acid (e.g., acetic acid, acetonetic acid, adipic acid, ascorbic acid, citric acid, 2,5-diketo-D-gluconic acid, formic acid, fumaric acid, glucaric acid, gluconic acid, glucuronic acid, glutaric acid, 3-hydroxypropionic acid, itaconic acid, lactic acid, malic acid, malonic acid, oxalic acid, oxaloacetic acid, propionic acid, succinic acid, and xylonic acid); and polyketide. The fermentation product can also be protein as a high value product.

[0309] In a preferred aspect, the fermentation product is an alcohol. It will be understood that the term "alcohol" encompasses a substance that contains one or more hydroxyl moieties. In a more preferred aspect, the alcohol is n-butanol. In another more preferred aspect, the alcohol is isobutanol. In another more preferred aspect, the alcohol is ethanol. In another more preferred aspect, the alcohol is methanol. In another more preferred aspect, the alcohol is arabinitol. In another more preferred aspect, the alcohol is butanediol. In another more preferred aspect, the alcohol is ethylene glycol. In another more preferred aspect, the alcohol is glycerin. In another more preferred aspect, the alcohol is glycerol. In another more preferred aspect, the alcohol is 1,3-propanediol. In another more preferred aspect, the alcohol is sorbitol. In another more preferred aspect, the alcohol is xylitol. See, for example, Gong, C. S., Cao, N. J., Du, J., and Tsao, G. T., 1999, Ethanol production from renewable resources, in *Advances in Biochemical Engineering/Biotechnology*, Scheper, T., ed., Springer-Verlag Berlin Heidelberg, Germany, 65: 207-241 ; Silveira, M. M., and Jonas, R., 2002, The biotechnological production of sorbitol, *Appl. Microbiol. Biotechnol.* 59: 400-408; Nigam, P., and Singh, D., 1995, Processes for fermentative production of xylitol - a sugar substitute, *Process Biochemistry* 30 (2): 117-124; Ezeji, T. C., Qureshi, N. and Blaschek, H. P., 2003, Production of acetone, butanol and ethanol by *Clostridium beijerinckii* BA101 and in situ recovery by gas stripping, *World Journal of Microbiology and Biotechnology* 19 (6): 595-603.

[0310] In another preferred aspect, the fermentation product is an alkane. The alkane can be an unbranched or a branched alkane. In another more preferred aspect, the alkane is pentane. In another more preferred aspect, the alkane is hexane. In another more preferred aspect, the alkane is heptane. In another more preferred aspect, the alkane is octane. In another more preferred aspect, the alkane is nonane. In another more preferred aspect, the alkane is decane. In another more preferred aspect, the alkane is undecane. In another more preferred aspect, the alkane is dodecane.

[0311] In another preferred aspect, the fermentation product is a cycloalkane. In another more preferred aspect, the cycloalkane is cyclopentane. In another more preferred aspect, the cycloalkane is cyclohexane. In another more preferred aspect, the cycloalkane is cycloheptane. In another more preferred aspect, the cycloalkane is cyclooctane.

[0312] In another preferred aspect, the fermentation product is an alkene. The alkene can be an unbranched or a branched alkene. In another more preferred aspect, the alkene is pentene. In another more preferred aspect, the alkene is hexene. In another more preferred aspect, the alkene is heptene. In another more preferred aspect, the alkene is octene.

[0313] In another preferred aspect, the fermentation product is an amino acid. In another more preferred aspect, the organic acid is aspartic acid. In another more preferred aspect, the amino acid is glutamic acid. In another more preferred aspect, the amino acid is glycine. In another more preferred aspect, the amino acid is lysine. In another more preferred aspect, the amino acid is serine. In another more preferred aspect, the amino acid is threonine. See, for example, Richard, A., and Margaritis, A., 2004, Empirical modeling of batch fermentation kinetics for poly(glutamic acid) production and other microbial biopolymers, *Biotechnology and Bioengineering* 87 (4): 501-515.

[0314] In another preferred aspect, the fermentation product is a gas. In another more preferred aspect, the gas is methane. In another more preferred aspect, the gas is H₂. In another more preferred aspect, the gas is CO₂. In another more preferred aspect, the gas is CO. See, for example, Kataoka, N., A. Miya, and K. Kiriya, 1997, Studies on hydrogen production by continuous culture system of hydrogen-producing anaerobic bacteria, *Water Science and Technology* 36 (6-7): 41-47; and Gunaseelan V.N. in *Biomass and Bioenergy*, Vol. 13 (1-2), pp. 83-114, 1997, Anaerobic digestion of biomass for methane production: A review.

[0315] In another preferred aspect, the fermentation product is isoprene.

[0316] In another preferred aspect, the fermentation product is a ketone. It will be understood that the term "ketone" encompasses a substance that contains one or more ketone moieties. In another more preferred aspect, the ketone is acetone. See, for example, Qureshi and Blaschek, 2003, *supra*.

[0317] In another preferred aspect, the fermentation product is an organic acid. In another more preferred aspect, the organic acid is acetic acid. In another more preferred aspect, the organic acid is acetic acid. In another more preferred aspect, the organic acid is adipic acid. In another more preferred aspect, the organic acid is ascorbic acid. In another more preferred aspect, the organic acid is citric acid. In another more preferred aspect, the organic acid is 2,5-diketo-D-gluconic acid. In another more preferred aspect, the organic acid is formic acid. In another more preferred aspect, the organic acid is fumaric acid. In another more preferred aspect, the organic acid is glucaric acid. In another more preferred aspect, the organic acid is gluconic acid. In another more preferred aspect, the organic acid is glucuronic acid. In another more preferred aspect, the organic acid is glutaric acid. In another preferred aspect, the organic acid is 3-hydroxypropionic acid. In another more preferred aspect, the organic acid is itaconic acid. In another more preferred aspect, the organic acid is lactic acid. In another more preferred aspect, the organic acid is malic acid. In another more preferred aspect, the organic acid is malonic acid. In another more preferred aspect, the organic acid is oxalic acid. In another more preferred aspect, the organic acid is propionic acid. In another more preferred aspect, the organic acid is succinic acid. In another more preferred aspect, the organic acid is xylonic acid. See, for example, Chen, R., and Lee, Y. Y., 1997, Membrane-mediated extractive fermentation for lactic acid production from cellulosic biomass, *Appl. Biochem. Biotechnol.* 63-65: 435-448.

[0318] In another preferred aspect, the fermentation product is polyketide.

[0319] Recovery. The fermentation product(s) can be optionally recovered from the fermentation medium using any method known in the art including, but not limited to, chromatography, electrophoretic procedures, differential solubility, distillation, or extraction. For example, alcohol is separated from the fermented cellulosic material and purified by conventional methods of distillation. Ethanol with a purity of up to about 96 vol.% can be obtained, which can be used as, for example, fuel ethanol, drinking ethanol, *i.e.*, potable neutral spirits, or industrial ethanol.

Signal Peptide

[0320] The present disclosure also relates to an isolated polynucleotide encoding a signal peptide comprising or consisting of amino acids 1 to 21 of SEQ ID NO: 2. The polynucleotide may further comprise a gene encoding a protein, which is operably linked to the signal peptide. The protein is preferably foreign to the signal peptide.

[0321] The present disclosure also relates to nucleic acid constructs, expression vectors and recombinant host cells comprising such a polynucleotide.

[0322] The present disclosure also relates to methods of producing a protein, comprising:

1. (a) cultivating a recombinant host cell comprising such a polynucleotide; and (b) recovering the protein.

[0323] The protein may be native or heterologous to a host cell. The term "protein" is not meant herein to refer to a specific length of the encoded product and, therefore, encompasses peptides, oligopeptides, and proteins. The term "protein" also encompasses two or more polypeptides combined to form the encoded product. The proteins also include hybrid polypeptides and fused polypeptides.

[0324] Preferably, the protein is a hormone or variant thereof, enzyme, receptor or portion thereof, antibody or portion thereof, or reporter. For example, the protein may be an oxidoreductase, transferase, hydrolase, lyase, isomerase, or ligase such as an aminopeptidase, amylase, carbohydrase, carboxypeptidase, catalase, cellulase, chitinase, cutinase, cyclodextrin glycosyltransferase, deoxyribonuclease, esterase, alpha-galactosidase, beta-galactosidase, glucoamylase, alpha-glucosidase, beta-glucosidase, invertase, laccase, another lipase, mannosidase, mutanase, oxidase, pectinolytic enzyme, peroxidase, phytase, polyphenoloxidase, proteolytic enzyme, ribonuclease, transglutaminase or xylanase.

[0325] The gene may be obtained from any prokaryotic, eukaryotic, or other source.

Examples

Strains

[0326] *Penicillium oxalicum* strain IBT5387 (Technical University of Denmark; NN005786) was used as the source of a Family GH3 beta-glucosidase gene. *Aspergillus oryzae* BECh2 was used as a host cell for expressing the *P. oxalicum* strain IBT5387 Family GH3 beta-glucosidase gene.

Media

[0327] LB medium was composed of 10 g of tryptone, 5 g of yeast extract, 5 g of sodium chloride, and deionized water to 1 liter.

[0328] LB agar plates were composed of 15 g of bacto agar per liter of LB medium

[0329] COVE selection plates were composed of 218 g of sorbitol, 20 g of agar, 20 ml of COVE salts solution, 10 mM acetamide, 15 mM CsCl, and deionized water to 1 liter. The solution was adjusted to pH 7.0 before autoclaving.

[0330] COVE salts solution was composed of 26 g of KCl, 26 g of MgSO₄·7H₂O, 76 g of KH₂PO₄, 50 ml of COVE trace metals solution, and deionized water to 1 liter.

[0331] COVE trace metals solution was composed of 0.04 g of Na₂B₄O₇·10H₂O, 0.4 g of CuSO₄·5H₂O, 1.2 g of FeSO₄·7H₂O, 0.7 g of MnSO₄·H₂O, 0.8 g of Na₂MoO₄·2H₂O, 10 g of ZnSO₄·7H₂O, and deionized water to 1 liter.

[0332] YP medium was composed of 10 g of yeast extract, 20 g of Bacto peptone, and deionized water to 1 liter.

Example 1: Cloning of a *Penicillium oxalicum* Family GH3 beta-glucosidase

[0333] A set of degenerate primers shown below were designed according to the strategy described by Rose et al., 1998,

Nucleic Acids Research 26: 1628-1635, for cloning a gene encoding a beta-glucosidase (EC 3.2.1.21) belonging to Family GH3.

GH3scree.f1:

atgaccctggccgaaaaagtaacytnacnacngg (SEQ ID NO: 3)

GH3scree.f2:

ggtggccggaactgggaaggctctsnccngaycc (SEQ ID NO: 4)

GH3scree.f5:

gagctgggcttcagggttgnatgwsngaytgg (SEQ ID NO: 5)

GH3scree.f6:

agcgcttggccggcctcgaytgwsnatgcc (SEQ ID NO: 6)

GH3scree.r1:

atcccagttgctcagggtcccknckngt (SEQ ID NO: 7)

GH3scree.r2:

aaaggtgtgtagctcagncrtgnccraaytc (SEQ ID NO: 8)

GH3scree.r3:

gtcaaagtggcggtagtcgatraanacnccytc (SEQ ID NO: 9)

GH3scree.r4:

ggtggcgagttgccgacgggttgactctgcrtanar (SEQ ID NO: 10)

GH3scree.r5:

gccgggcagaccggcccagaggatggcngtnacrtnngg (SEQ ID NO: 11)

GH3scree.r6:

caggacggggccaaccgagtgaatgacnacdatngtrtt (SEQ ID NO: 12)

[0334] PCR screening of *Penicillium oxalicum* strain IBT5387 was performed using two successive PCRs. The forward primers (GH3scree.f1, GH3scree.f2, GH3scree.f5, or GH3scree.f6) (0.33 µl of a 10 mM stock) were combined with the reverse primers (GH3scree.r1, GH3scree.r2, GH3scree.r3, GH3scree.r4, GH3scree.r5, or GH3scree.r6) (0.33 µl of a 10 mM stock) in a 10 µl mixture containing 0.33 µl of *P. oxalicum* genomic DNA and 5 µl of REDDYMIX™ Extensor PCR Master Mix 1 (ABgene Ltd., Surrey, United Kingdom). *P. oxalicum* genomic DNA was obtained according to the procedure described in the FastDNA® SPIN Kit (Q-BIOgene, Carlsbad, CA, USA). The PCR reaction was performed using a DYAD® Thermal Cycler (Bio-Rad Laboratories, Inc., Hercules, CA, USA) programmed for one cycle at 94°C for 2 minutes; 9 cycles each at 94°C for 15 seconds, 63°C for 30 seconds with a decrease of 1°C for each cycle, and 68°C for 1 minutes 45 seconds; 24 cycles each at 94°C for 15 seconds, 55°C for 30 seconds, and 68°C for 1 minutes 45 seconds; and an extension at 68°C for 7 minutes.

[0335] PCR products obtained during the first PCR were re-amplified with their corresponding primers by transferring 0.5 µl of the first PCR reaction to a second 20 µl mixture containing the same concentration of primers, dNTPs, DNA polymerase, and buffer as the first PCR reaction. The second PCR was performed using a DYAD® Thermal Cycler programmed for one cycle at 94°C for 2 minutes; 24 cycles each at 94°C for 15 seconds, 58°C for 30 seconds, and 68°C for 1 minutes 45 seconds; and extension at 68°C for 7 minutes.

[0336] PCR products obtained during the second amplification were analyzed by 1% agarose gel electrophoresis using 40 mM Tris base-20 mM sodium acetate-1 mM disodium EDTA (TAE) buffer. Single bands ranging from 2000 to 800 nucleotides in size were excised from the gel and purified using a GFX® PCR DNA and Gel Band Purification Kit (GE Healthcare, Hillerød Denmark) according to the manufacturer's instructions. Purified DNA samples were directly sequenced with primers used for amplification. Sequences were assembled in SeqMan v7.2.1 (DNA star, Madison, WI, USA) into a contig that was used for designing primers shown below, based on recommendations of length and temperature described in a GENE WALKING SPEEDUP™ Kit (Seegene, Inc., Seoul, Korea).

5786GH3-51TSP1f:

CACCAACACCGCAATCTAGC (SEQ ID NO: 13)

5786GH3-51 TSP2f:

GGTGACGAGGTTGTCCAACGTGACG (SEQ ID NO: 14)

5786GH3-51TSP1r:

CTTGAAGCCAAGGCGAGG (SEQ ID NO: 15)

5786GH3-51 TSP2r:

TCCGGTATTTCCTACACATGGTCC (SEQ ID NO: 16)

[0337] GeneWalking was based on the protocol from the GENE WALKING SPEEDUP™ Kit with some minor differences. Only two PCR amplifications were carried out and in both cases, the REDDYMIX™ Extensor PCR Master Mix 1 was used in place of the enzyme mix present in the Kit. GeneWalking PCR 1 was performed in a total volume of 15 µl by mixing 1.2 µl of primer 1 to 4 (2.5 mM) from the GENE WALKING SPEEDUP™ Kit with 0.3 µl of primer 5786GH3-51TSP1f or primer 5786GH3-51TSP1r (10 mM) in the presence of 7.5 µl of REDDYMIX™ Extensor PCR Master Mix 1, and 0.5 µl of *P. oxalicum* genomic DNA. The PCR was performed using a DYAD® Thermal Cycler programmed for one cycle at 94°C for 3 minutes followed by 1 minute at 42°C and 2 minutes at 68°C; 30 cycles each at 94°C for 30 seconds, 58°C for 30 seconds, and 68°C for 1 minute and 40 seconds; and elongation at 68°C for 7 minutes. A 0.5 µl aliquot of the amplification reaction was transferred to a second PCR tube containing a 20 µl mixture composed of 10 µl of REDDYMIX™ Extensor PCR Master Mix 1, 1 µl of primer 5 (10 mM) from the Kit, 1 µl of primer T5786GH3-51SP2f or 5786GH3-51 TSP2r (10 mM). The amplification was performed in a DYAD® Thermal Cycler programmed for denaturation at 94°C for 3 minutes; 35 cycles each at 94°C for 30 seconds, 58°C for 30 seconds, and 68°C for 1 minute and 40 seconds; and elongation at 68°C for 7 minutes.

[0338] The PCR products were analyzed by 1% agarose gel electrophoresis in TAE buffer. Single bands ranging from 700 to 1200 nucleotides in size were excised from the gel and purified using a GFX® PCR DNA and Gel Band Purification Kit according to manufacturer's instructions. Purified DNA samples were directly sequenced with the primers used for amplification.

[0339] Based on blastx analyses, the start and the stop codons of the gene were identified and the primers shown below were designed for cloning the gene into the expression vector pDAu109 (WO 2005042735) using an IN-FUSION™ Dry-Down PCR Cloning Kit (BD Biosciences, Palo Alto, CA, USA).

5786GH3-51-r1:

agatctcgagaagcttaccATGACTCCAATCGCGCGCTCAAGG (SEQ ID NO: 17)

5786GH3-51f1:

acacaactggggatccaccATGAGGAGCTCAACGACGGTTCTGGCC (SEQ ID NO: 18)

[0340] The *P. oxalicum* beta-glucosidase gene was amplified by PCR using the two cloning primers described above with *P. oxalicum* strain IBT5387 genomic DNA. The PCR was composed of 1 µl of *P. oxalicum* genomic DNA, 0.75 µl of primer 5786GH3-51f1 (10 µM), 0.75 µl of primer 5786GH3-51r1 (10 µM), 3 µl of 5X HF buffer (Finnzymes Oy, Finland), 0.25 µl of 50 mM MgCl₂, 0.3 µl of 10 mM dNTP, 0.15 µl of PHUSION® DNA polymerase (Finnzymes Oy, Finland), and PCR-grade water to 15 µl. The amplification reaction was performed using a DYAD® Thermal Cycler programmed for 2 minutes at 98°C followed by 10 touchdown cycles each at 98°C for 15 seconds, 70°C (-1°C/cycle) for 30 seconds, and 72°C for 2 minutes and 30 seconds; and 25 cycles each at 98°C for 15 seconds, 60°C for 30 seconds, 72°C for 2 minutes and 30 seconds, and 5 minutes at 72°C.

[0341] The reaction product was isolated by 1.0% agarose gel electrophoresis using TAE buffer where an approximately 2.8 kb PCR product band was excised from the gel and purified using a GFX® PCR DNA and Gel Band Purification Kit according to manufacturer's instructions. DNA corresponding to the *Penicillium oxalicum* beta-glucosidase gene was cloned into the expression vector pDAu109 (WO 2005042735) linearized with *Bam* HI and *Hind* III, using an IN-FUSION™ Dry-Down PCR Cloning Kit (BD Biosciences, Palo Alto, CA, USA) according to the manufacturer's instructions.

[0342] A 2.5 µl volume of the diluted ligation mixture was used to transform *E. coli* TOP10 chemically competent cells (Invitrogen, Carlsbad, CA, USA). Three colonies were selected on LB agar plates containing 100 µg of ampicillin per ml and cultivated overnight in 3 ml of LB medium supplemented with 100 µg of ampicillin per ml. Plasmid DNA was purified using an E.Z.N.A.® Plasmid Mini Kit (Omega Bio-Tek, Inc., Norcross, GA, USA) according to the manufacturer's instructions. The *Penicillium oxalicum* beta-glucosidase gene sequence was verified by Sanger sequencing before heterologous expression. One plasmid designated pIF113#1 was selected for expressing the *P. oxalicum* beta-glucosidase in an *Aspergillus oryzae* host cell.

Example 2: Characterization of the *Penicillium oxalicum* genomic DNA encoding a Family GH3 polypeptide having beta-glucosidase activity

[0343] The genomic DNA sequence (SEQ ID NO: 1) and deduced amino acid sequence (SEQ ID NO: 2) of the *Penicillium oxalicum* beta-glucosidase gene are shown in Figures 1A and 1B. The coding sequence is 2793 bp including the stop codon with 2 predicted introns of 82 bp (nucleotides 85 to 116) and 59 bp (nucleotides 346 to 404). The encoded predicted protein is 883 amino acids. Using the SignalP program (Nielsen et al., 1997, Protein Engineering 10: 1-6), a signal peptide of 21 residues was predicted. The predicted mature protein contains 862 amino acids with a predicted molecular mass of 94.7 kDa and an isoelectric point of 5.04. Based on Pozzo et al., 2010, J. Mol. Biol. 397: 724-739, two catalytic acids, D301 as the nucleophile and E531 as the general base/acid, were identified by sequence alignment. In addition, D113 and W302 were identified by sequence alignment as important for enzyme activity.

[0344] A comparative pairwise global alignment of amino acid sequences was determined using the Needleman-Wunsch algorithm (Needleman and Wunsch, 1970, J. Mol. Biol. 48: 443-453) as implemented in the Needle program of EMBOSS with gap open penalty of 10, gap extension penalty of 0.5, and the EBLOSUM62 matrix. The alignment showed that the deduced amino acid sequence of the mature polypeptide of the *P. oxalicum* Family GH3 polypeptide having beta-glucosidase activity shares 82.1% identity (excluding gaps) to the deduced amino acid sequence of a beta-glucosidase from *Penicillium brasilianum* (Uniprot Accession No. A5A4M8).

[0345] The *Penicillium oxalicum* beta-glucosidase gene was deposited with the Agricultural Research Service Patent Culture Collection (NRRL), Northern Regional Research Center, as *E. coli* D55EX containing pUC19 D55EX (Figure 2) on June 16, 2010 and assigned accession number NRRL B-50395.

Example 3: Expression of the *Penicillium oxalicum* beta-glucosidase gene in *Aspergillus oryzae*

[0346] Protoplasts of *Aspergillus oryzae* BECh2 (WO 2000/39322) were prepared according to WO 95/002043. One hundred μ l of protoplasts were mixed with 2.5-15 μ g of the *Aspergillus* expression vector pIF113#1 (Example 1) and 250 μ l of 60% PEG 4000 (polyethylene glycol, molecular weight 4,000; Applichem, Darmstadt, Germany), 10 mM CaCl_2 , and 10 mM Tris-HCl pH 7.5 and gently mixed. The mixture was incubated at 37°C for 30 minutes and the protoplasts were spread onto COVE plates for selection. After incubation for 4-7 days at 37°C spores of sixteen transformants were inoculated into 0.5 ml of YP medium supplemented with 2% maltodextrin in 96 deep-well plates. After 4 days cultivation at 30°C, the culture broths were analyzed to identify the best transformants producing large amounts of active *P. oxalicum* beta-glucosidase. The analysis was based on SDS-PAGE and activity of the enzyme on 4-nitrophenyl-beta-D-glucopyranoside (pNPG; (Sigma Aldrich Chemical Co., St. Louis, MO, USA). The pNPG assay was modified from Hägerdal et al., 1979, Biotechnology and Bioengineering 21: 345-355, as follows: 10 μ l of culture broth was mixed with 90 μ l of assay reagent containing 10 μ l of 0.1% TWEEN® 80K, 10 μ l of 1 M sodium citrate pH 5, 4 μ l of 100 mM pNPG substrate, previously solubilized in DMSO (0.4% final volume in stock solution), and filtered water. The pNPG assay was performed for 30 minutes at 37°C and the absorbance was measured at 405 nm before and after addition of 100 μ l of 1 M sodium carbonate pH 10. The highest absorbance values at 405 nm correlated to the SDS-PAGE data for selection of the best transformant.

[0347] Spores of the best transformant were spread on COVE plates containing 0.01% TRITON® X-100 in order to isolate single colonies. The spreading was repeated twice in total on COVE selection plates containing 10 mM sodium nitrate. Spores were then inoculated into 100 ml of YP medium supplemented with 2% maltodextrin in 250 ml shake flasks and incubated for 4 days at 30°C with shaking at 100 rpm.

Example 4: Purification of *Penicillium oxalicum* beta-glucosidase

[0348] Combined supernatants of the shake flask cultures of Example 3 were first filtered using a glass micro fiber filter with a 0.7 μ m pore size, and then sterile filtered using a filtration unit equipped with a PES (polyether sulfone) membrane with a 0.22 μ m pore size (Nalge Nunc International, New York, NY USA). The sterile filtered supernatant was then adjusted to a concentration of 2 M ammonium sulfate by slowly adding solid ammonium sulfate and dissolving by gentle stirring and then filtered using a glass micro fiber filter with a 0.7 μ m pore size. If there was any precipitate, the precipitate was discarded.

[0349] The filtered supernatant was applied to a 50 ml Toyopearl Phenyl-650M column (TOSOH Bioscience GmbH, Germany) equilibrated with 2 M ammonium sulfate in water and unbound material was eluted with 2 M ammonium sulfate until the UV absorbance at 280 nm was below 0.05. Bound protein was eluted with 50% ethanol as solution B using a step gradient. Ten ml fractions were collected and monitored by UV absorbance at 280 nm. The fractions were analyzed by SDS-PAGE and pooled the fractions containing protein with expected molecular weight. The pooled fractions were dialyzed using 50 mM HEPES pH 7.5 buffer, so the ionic strength was below 4 M Si and the pH was 7.5.

[0350] The pooled protein was applied to a 50 ml Q SEPHAROSE® Fast Flow column (GE Healthcare, Piscataway, NJ, USA) equilibrated with 50 mM HEPES pH 7.5 buffer and unbound material was eluted with 50 mM HEPES buffer pH 7.5. The bound protein was then eluted with a linear 20 column volume salt gradient in 50 mM HEPES buffer pH 7.5 containing 1 M NaCl as buffer B. Ten ml fractions of the eluate were collected and each were analyzed for purity of the protein by SDS-PAGE. The fractions with highest purity were pooled. Identity of the beta-glucosidase was confirmed by mass spectroscopy and protein identification was carried out by in gel digestion.

Example 5: Characterization of the *Penicillium oxalicum* GH3 beta-glucosidase in PCS hydrolysis

[0351] Corn stover was pretreated at the U.S. Department of Energy National Renewable Energy Laboratory (NREL) using dilute sulfuric acid. The conditions used for the pretreatment were 0.048 g sulfuric acid/ g dry biomass at 190°C and 25% w/w dry solids for around 1 minute. The water-insoluble solids in the pretreated corn stover (PCS) contained 52% cellulose, 3.6% hemicellulose and 29.8% lignin. Cellulose and hemicellulose were determined by a two-stage sulfuric acid hydrolysis with subsequent analysis of sugars by high performance liquid chromatography using NREL Standard Analytical Procedure #002. Lignin was determined gravimetrically after hydrolyzing the cellulose and hemicellulose fractions with sulfuric acid using NREL Standard Analytical Procedure #003. Prior to enzymatic hydrolysis, the PCS was ground (Multi Utility Grinder, Inno Concepts Inc., GA, USA) and sieved through a 450 µm screen (Retsch AS200). The *Penicillium oxalicum* GH3 beta-glucosidase broth was purified before the hydrolysis experiments as described in Example 4. The protein concentration was determined using a BCA Reagent Kit (Pierce, Rockford, IL, USA).

[0352] Synergistic effects between *Penicillium oxalicum* GH3 beta-glucosidase and CELLULAST® 1.5L (Novozymes A/S, Bagsvaerd, Denmark) were determined using a 1-gram ground-sieved, unwashed PCS (GS-PCS) hydrolysis assay at 50 and 55°C, pH 5. *Penicillium oxalicum* GH3 beta-glucosidase (1.5 mg/g cellulose) was added to CELLULAST® 1.5L (15 mg/g cellulose) for a total loading of 16.5 mg protein/g cellulose. Total insoluble solids loading of the GS-PCS was 50 g per liter of 50 mM sodium acetate pH 5.0 buffer containing 1 mM manganese sulfate. The total reaction volume was 1.0 ml in 96-well plates. Assays were run in triplicate. After 72 hour incubation at 50 and 55°C, samples were mixed well, centrifuged at 3000 rpm for 5 minutes, and supernatants were filtered using a 0.45 µm MULTISCREEN® 96-well filter plate (Millipore, Bedford, MA, USA) and filtrates analyzed for sugar content as described below. When not used immediately, filtered aliquots were frozen at -20°C. The sugar concentrations of samples diluted in 0.005 M H₂SO₄ were measured using a 4.6 x 250 mm AMINEX® HPX-87H column (Bio-Rad Laboratories, Inc., Hercules, CA, USA) by elution with 0.05% w/w benzoic acid-0.005 M H₂SO₄ at 65°C at a flow rate of 0.6 ml per minute, and quantitation by integration of the glucose, cellobiose, and xylose signals from refractive index detection (CHEMSTATION®, AGILENT® 1100 HPLC, Agilent Technologies, Santa Clara, CA, USA) calibrated by pure sugar samples. The resultant glucose and cellobiose equivalents were used to calculate the percentage of cellulose conversion for each reaction. Hydrolysis data were presented as % of total cellulose converted to glucose. The degree of cellulose conversion to reducing sugar was calculated using the following equation:

$$\text{Conversion (\%)} = \text{RS}_{(\text{mg/ml})} * 100 * 162 / (\text{Cellulose}_{(\text{mg/ml})} * 180) = \text{RS}_{(\text{mg/ml})} * 100 / (\text{Cellulose}_{(\text{mg/ml})} * 1.111)$$

In this equation, RS is the concentration of reducing sugar in solution measured in glucose equivalents (mg/ml), and the factor 1.111 reflects the weight gain in converting cellulose to glucose.

[0353] PCS hydrolysis by *Penicillium oxalicum* GH3 beta-glucosidase (1.5 mg/g cellulose) plus CELLULAST® 1.5L (15 mg/g cellulose) yielded a cellulose conversion of 64.2% and 63.5% after 72 hours at 50°C and 55°C, respectively, while PCS hydrolysis by CELLULAST® 1.5L alone at 15 mg/g cellulose yielded a cellulose conversion of 32.9% and 35.3% at 50 and 55°C respectively, and at 30 mg/g cellulose yielded a cellulose conversion of 51.6% and 53.3% at 50 and 55°C respectively. The results indicated that supplemented *Penicillium oxalicum* GH3 beta-glucosidase had a synergistic effect on CELLULAST® 1.5L in PCS hydrolysis at 50 and 55°C, pH 5.0.

Example 6: Characterization of the *Penicillium oxalicum* GH3 beta-glucosidase

[0354] Specific activity. 4-Nitrophenyl-beta-D-glucopyranoside (pNPG) was dissolved in DMSO to produce a 100 mM stock solution. Right before assay, the 100 mM pNPG stock solution was diluted 100-fold in 50 mM sodium acetate pH 5 with 0.01% TWEEN® 20 to 1 mM pNPG. Enzyme was diluted to a wide range of concentrations in the same buffer. To 20 µl of the enzyme dilutions (a range of concentration), 100 µl of the 1 mM pNPG solution were quickly added. The reaction was incubated at 40°C for 20 minutes before the reaction was stopped with 50 µl of 1 M sodium carbonate pH 10, and the absorbance at 405 nm was measured for 4-nitrophenolate production. The specific activity of the *Penicillium oxalicum* GH3 beta-glucosidase on pNPG was determined to be 72.3 ± 1.7 IU/mg, where one mg of enzyme produce 1 µmole of glucose per minute at 40°C, pH 5.

[0355] Thermostability. The *Penicillium oxalicum* GH3 beta-glucosidase was diluted in 50 mM sodium acetate pH 5 with 0.01% TWEEN® 20 to 1 g/L, and then incubated at 50°C for 3 days, and 60°C for 3 hours or 24 hours. The same sample was stored at 4°C to serve as control. After incubation, the residual activity of the samples was measured using pNPG as substrate, following the same assay protocol described above, but using one enzyme loading which yielded less than 5% conversion. The activity of the sample in 4°C was normalized to 100%, and the activities of samples at other incubation conditions were compared to the 4°C activity. The thermostability of the *Penicillium oxalicum* GH3 beta-glucosidase is shown below.

Incubation condition	Residual activity on pNPG
4°C	100%
50°C, 72 hours	100%
60°C, 3 hours	100%
60°C, 24 hours	100%
70°C, 24 hours	76%

It can be seen from the table above that the the *Penicillium brasiliense* GH3 (Uniprot: A5A4M8) from D1 has significantly reduced activity under the conditions tested after 50°C for 72 hours, at 60°C for 3 hours or 60°C for 24 hours. In comparison, the *Penicillium oxalicum* GH3 β-glucosidase (SEQ ID NO: 2) of the application retains 100% of the under all time points measured and is surprisingly more thermostable than the prior art sequence of D1.

[0356] pH profile. The pH activity profile of the *Penicillium oxalicum* GH3 beta-glucosidase was determined using the same protocol described above, except the beta-glucosidase was incubated at four different pHs (4, 5, 6, and 7) and one enzyme loading was used, which yielded less than 5% conversion. Britton Robinson buffer (100 mM) was used as the buffer system. The 100 mM Britton Robinson buffer was titrated to a various pH values in the range of 4-7 using 5 M sodium hydroxide and then diluted to 40 mM with deionized water. The pNP-glucose was prepared in the same buffer, and the activity was measured at 50°C. The highest activity was normalized to 100%, and activities at the other pH values were compared to the highest activity and expressed in % activity. The pH profile of the *Penicillium oxalicum* GH3 beta-glucosidase is shown below.

pH value	Activity
4.0	38%
5.0	99%
6.0	100%
7.0	36%

[0357] Inhibition by glucose. A Dixon plot was used to determine the half maximal inhibitory concentration (IC₅₀) of glucose toward the *Penicillium oxalicum* GH3 beta-glucosidase. pNPG was used as the substrate. An unsaturated pNPG concentration (0.5 mM) was used, with a series of glucose concentrations. The IC₅₀ for the *Penicillium oxalicum* GH3 beta-glucosidase was determined to be 8 mM glucose.

Deposit of Biological Material

[0358] The following biological material has been deposited under the terms of the Budapest Treaty with the Agricultural Research Service Patent Culture Collection (NRRL), Northern Regional Research Center, 1815 University Street, Peoria, IL, USA, and given the following accession number:

Deposit	Accession Number	Date of Deposit
<i>E. coli</i> D55EX	NRRL B-50395	June 16, 2010

[0359] The strain has been deposited under conditions that assure that access to the culture will be available during the pendency of this patent application to one determined by foreign patent laws to be entitled thereto. The deposit represents a substantially pure culture of the deposited strain. The deposit is available as required by foreign patent laws in countries wherein counterparts of the subject application, or its progeny are filed. However, it should be understood that the availability of a deposit does not constitute a license to practice the subject invention in derogation of patent rights granted by governmental action.

[0360] The present invention is further described by the following numbered paragraphs:

1. [1] An isolated polypeptide having beta-glucosidase activity, selected from the group consisting of: (a) a polypeptide having at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, and 100% sequence identity to the mature polypeptide of SEQ ID NO: 2; (b) a variant of the mature polypeptide of SEQ ID NO: 2 comprising a substitution, deletion, and/or insertion of one or more positions, wherein said variant has at least 85%, e.g. at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% sequence identity to the mature polypeptide of SEQ ID NO: 2 and wherein said variant has improved thermal stability over the mature polypeptide of SEQ ID NO:2.
2. [2] The polypeptide of paragraph 1, comprising or consisting of SEQ ID NO: 2.
3. [3] The polypeptide of any of paragraphs 1-2, comprising or consisting of the mature polypeptide of SEQ ID NO: 2.
4. [4] The polypeptide of paragraph 3, wherein the mature polypeptide is amino acids 22 to 883 of SEQ ID NO: 2.
5. [5] The polypeptide of paragraph 1, which is a variant of the mature polypeptide of SEQ ID NO: 2 comprising a substitution, deletion, and/or insertion at one or more positions.
6. [6] The polypeptide of any of paragraphs 1-5, which is a fragment of SEQ ID NO: 2, wherein the fragment has beta-glucosidase activity.
7. [7] The polypeptide of any of paragraphs 1-6, which is encoded by the polynucleotide contained in plasmid pUC19 D55EX which is contained in *E. coli* NRRL B-50395.
8. [8] A composition comprising the polypeptide of any of paragraphs 1-7.
9. [9] An isolated polynucleotide encoding the polypeptide of any of paragraphs 1-7.
10. [10] A nucleic acid construct or expression vector comprising the polynucleotide of paragraph 9 operably linked to one or more control sequences that direct the production of the polypeptide in an expression host.
11. [11] A recombinant host cell comprising the polynucleotide of paragraph 9 operably linked to one or more control sequences that direct the production of the polypeptide.
12. [12] A method of producing the polypeptide of any of paragraphs 1-7, comprising: (a) cultivating a cell, which in its wild-type form produces the polypeptide, under conditions conducive for production of the polypeptide; and (b) recovering the polypeptide.
13. [13] A method of producing a polypeptide having beta-glucosidase activity, comprising: (a) cultivating the host cell of paragraph 11 under conditions conducive for production of the polypeptide; and (b) recovering the polypeptide.
14. [14] A transgenic plant, plant part or plant cell transformed with a polynucleotide encoding the polypeptide of any of paragraphs 1-9.
15. [15] A method of producing the polypeptide having beta-glucosidase activity, comprising: (a) cultivating the transgenic plant or plant cell of paragraph 14 under conditions conducive for production of the polypeptide; and (b) recovering the polypeptide.
16. [16] A method of producing a mutant of a parent cell, comprising inactivating a polynucleotide encoding the polypeptide of any of paragraphs 1-7, which results in the mutant producing less of the polypeptide than the parent cell.
17. [17] A mutant cell produced by the method of paragraph 16.
18. [18] The mutant cell of paragraph 17, further comprising a gene encoding a native or heterologous protein.
19. [19] A method of producing a protein, comprising: (a) cultivating the mutant cell of paragraph 17 or 18 under conditions conducive for production of the protein; and (b) recovering the protein.
20. [20] A double-stranded inhibitory RNA (dsRNA) molecule comprising a subsequence of the polynucleotide of paragraph 9, wherein optionally the dsRNA is an siRNA or an miRNA molecule.
21. [21] The double-stranded inhibitory RNA (dsRNA) molecule of paragraph 20, which is about 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25 or more duplex nucleotides in length.
22. [22] A method of inhibiting the expression of a polypeptide having beta-glucosidase activity in a cell, comprising administering to the cell or expressing in the cell the double-stranded RNA (dsRNA) molecule of paragraph 20 or 21.
23. [23] A cell produced by the method of paragraph 22.
24. [24] The cell of paragraph 23, further comprising a gene encoding a native or heterologous protein.
25. [25] A method of producing a protein, comprising: (a) cultivating the cell of paragraph 23 or 24 under conditions conducive for production of the protein; and (b) recovering the protein.

26. [26] An isolated polynucleotide encoding a signal peptide comprising or consisting of amino acids 1 to 21 of SEQ ID NO: 2.
27. [27] A nucleic acid construct or expression vector comprising a gene encoding a protein operably linked to the polynucleotide of paragraph 26, wherein the gene is foreign to the polynucleotide encoding the signal peptide.
28. [28] A recombinant host cell comprising a gene encoding a protein operably linked to the polynucleotide of paragraph 26, wherein the gene is foreign to the polynucleotide encoding the signal peptide.
29. [29] A method of producing a protein, comprising: (a) cultivating a recombinant host cell comprising a gene encoding a protein operably linked to the polynucleotide of paragraph 26, wherein the gene is foreign to the polynucleotide encoding the signal peptide, under conditions conducive for production of the protein; and (b) recovering the protein.
30. [30] A method for degrading or converting a cellulosic material, comprising: treating the cellulosic material with an enzyme composition in the presence of the polypeptide having beta-glucosidase activity of any of paragraphs 1-7.
31. [31] The method of paragraph 30, wherein the cellulosic material is pretreated.
32. [32] The method of paragraph 30 or 31, wherein the enzyme composition comprises one or more enzymes selected from the group consisting of a cellulase, a GH61 polypeptide having cellulolytic enhancing activity, a hemicellulase, an esterase, an expansin, a laccase, a ligninolytic enzyme, a pectinase, a peroxidase, a protease, and a swollenin.
33. [33] The method of paragraph 32, wherein the cellulase is one or more enzymes selected from the group consisting of an endoglucanase, a cellobiohydrolase, and a beta-glucosidase.
34. [34] The method of paragraph 32, wherein the hemicellulase is one or more enzymes selected from the group consisting of a xylanase, an acetylxylan esterase, a feruloyl esterase, an arabinofuranosidase, a xylosidase, and a glucuronidase.
35. [35] The method of any of paragraphs 30-34, further comprising recovering the degraded cellulosic material.
36. [36] The method of paragraph 35, wherein the degraded cellulosic material is a sugar.
37. [37] The method of paragraph 36, wherein the sugar is selected from the group consisting of glucose, xylose, mannose, galactose, and arabinose.
38. [38] A method for producing a fermentation product, comprising: (a) saccharifying a cellulosic material with an enzyme composition in the presence of the polypeptide having beta-glucosidase activity of any of paragraphs 1-7; (b) fermenting the saccharified cellulosic material with one or more fermenting microorganisms to produce the fermentation product; and (c) recovering the fermentation product from the fermentation.
39. [39] The method of paragraph 38, wherein the cellulosic material is pretreated before saccharification.
40. [40] The method of paragraph 38 or 39, wherein the enzyme composition comprises one or more enzymes selected from the group consisting of a cellulase, a GH61 polypeptide having cellulolytic enhancing activity, a hemicellulase, an esterase, an expansin, a laccase, a ligninolytic enzyme, a pectinase, a peroxidase, a protease, and a swollenin.
41. [41] The method of paragraph 40, wherein the cellulase is one or more enzymes selected from the group consisting of an endoglucanase, a cellobiohydrolase, and a beta-glucosidase.
42. [42] The method of paragraph 40, wherein the hemicellulase is one or more enzymes selected from the group consisting of a xylanase, an acetylxylan esterase, a feruloyl esterase, an arabinofuranosidase, a xylosidase, and a glucuronidase.
43. [43] The method of any of paragraphs 38-42, wherein steps (a) and (b) are performed simultaneously in a simultaneous saccharification and fermentation.
44. [44] The method of any of paragraphs 38-43, wherein the fermentation product is an alcohol, an alkane, a cycloalkane, an alkene, an amino acid, a gas, isoprene, a ketone, an organic acid, or polyketide.
45. [45] A method of fermenting a cellulosic material, comprising: fermenting the cellulosic material with one or more fermenting microorganisms, wherein the cellulosic material is saccharified with an enzyme composition in the presence of a polypeptide having beta-glucosidase activity of any of paragraphs 1-7.
46. [46] The method of paragraph 45, wherein the fermenting of the cellulosic material produces a fermentation product.
47. [47] The method of paragraph 46, further comprising recovering the fermentation product from the fermentation.
48. [48] The method of any of paragraphs 45-47, wherein the cellulosic material is pretreated before saccharification.
49. [49] The method of any of paragraphs 45-48, wherein the enzyme composition comprises one or more enzymes selected from the group consisting of a cellulase, a GH61 polypeptide having cellulolytic enhancing activity, a hemicellulase, an esterase, an expansin, a laccase, a ligninolytic enzyme, a pectinase, a peroxidase, a protease, and a swollenin.
50. [50] The method of paragraph 49, wherein the cellulase is one or more enzymes selected from the group consisting of an endoglucanase, a cellobiohydrolase, and a beta-glucosidase.
51. [51] The method of paragraph 49, wherein the hemicellulase is one or more enzymes selected from the group consisting of a xylanase, an acetylxylan esterase, a feruloyl esterase, an arabinofuranosidase, a xylosidase, and a glucuronidase.
52. [52] The method of any of paragraphs 45-51, wherein the fermentation product is an alcohol, an alkane, a cycloalkane, an alkene, an amino acid, a gas, isoprene, a ketone, an organic acid, or polyketide.

SEQUENCE LISTING

[0361]

<110> Novozymes A/S Novozymes, Inc. Morant, Marc Patkar, Shamkant Ding, Hanshu

<120> Polypeptides having Beta-glucosidase activity and polynucleotides encoding same

<130> 11729-WO-PCT

<150> US 61/360,351

<151> 2010-06-30

<160> 34

<170> PatentIn version 3.5

<210> 1

<211> 2793

<212> DNA

<213> Penicillium oxalicum

<400> 1

atgaggagct caacgacggt tctggccttt gtgtcatacg ggagttttgg gattgcggca	60
gcgaagccca tacaagagca caaggtatga acggttttct ttaagggaga acgcgagata	120
taacgattcg agagtaacgg gcaactgacca ttttgttctc tcaaagcccg agcttctgaa	180
tagctttcac ggctttcaga ttgaagagcc gtattcgccg ccgtttttatc cctcgccatg	240
gatgaatcct cgtgctgaag gctgggagga tgcgtatcag aaggcgcttg catttgtctc	300
tcaactgacg ttactcgaga aggtcaatct gacgacgggt gttgggtgag tgattgccct	360
tttctctat gatctgtgag acagtgggct gaacatgttc aaagtggga aaacggacca	420
tgtgtaggaa ataccggatc ggtccctcgc cttggcttca aggggctttg cttgcaagac	480
tctctcaag gtgtgagatt tgcgactat gcactctgctt tcacgtccag tcagatggct	540
gctgccacat ttgacaagac ggttctttat gagcgtggtg ttgccatggc tcaagagcac	600
aagggaagg gcatacaggt gcaactgggc ccgctcgtg gccctttggg ccgcgcccct	660
gaaggaggcc gaaactggga aggattctcg ccgcatcccg tcctgaacgg cattgcgatg	720
gccgagacca tcaagggcat gcaagatacc ggagtcttgg cctgcgcgaa acattatgtt	780
ggaaacgaac aagagcactt ccgacaagca ggcgagctcg ccgggcatgg atacaacatt	840
gctgaatoga tctcgtcgaa cattgacgat cgtactatgc atgaaatgta cttgtggcct	900
tttgctgatg ccgtacgtgc tggagtggca tcttttatgt gctcctacaa ccagatcaac	960
aactcgtacg gatgcaaaaa cagtcacact ctaaacaaac tcctcaagag cgagttgggc	1020
ttccaagggt tcgtgatgag tgactggcaa gcacaacact ctggagtatc gtcggcgctg	1080
gccgggctcg atatgacaat gccaggagat actgaattcg acacgggact cagcttttgg	1140
ggatccaatc tcactattgc ggttctgaac ggcaccgtgc ccgagtggag gattgacgat	1200

```

atggcgatgc ggatcatggc cgcttatttc aaagtcggtc tgacaattga agatcagcct 1260
gacgtgaatt tcaactcatg gacctacgat acctatgggtt acaatactt ttacagcaag 1320
gagggttacg agaaagtga ctggcatgtc gacgtgcgtg caaaccacgc ccaacagatt 1380
cgtgagacgg ccgcaaaggg aactgtgctt ctcaaaaaca ccaatgggtg tcttcggctg 1440
aaacgacctt ccttcgtcgc cggtgttggc gaagatgcgg gtccaaatcc caagggtccg 1500
aatgggtgtc ctgacgtgg ctgcgatgag ggaaccctgg cgatgggatg gggatctggc 1560
tcgacccagt tcccgcttct tgttactcca gactctgcga ttcaagccaa ggtcttgag 1620
tatggtggcc gatatgagag tattttggac aactacgatg aaagtgccat ctccgcatg 1680
gtctcgcagc ctgatgtac ttgcacgtc tttggcaact cggattctgg cgaagcttcc 1740
atcacccgtc acggcaactg gggggatcgg aataacotca ctcttggca aaatgccgac 1800
gcggtgatcc gcaatgtcag tgctctgtgc agcaacacca ttgtcgttct tcacactgtc 1860
ggccctgtgc tgttgaccga ttactatgag caccocaaca tcacagccat tgtctgggcc 1920
ggcctacctg gcgaagaatc agggaatgct cttgtcgata tcctctgggg cgatgtcaac 1980
ccagccggtc gcacgccatt cacatgggcc aagagccgtg aagactatgg cacagacgtg 2040
atgtatgtgc ccaacaacgg cgaacaggca cccagcaat ccttctccga aggcattctc 2100
cttgactacc ggcattttga ccaagctgga attgatcca tctacagatt cggccacggg 2160
ctctcatata ccaatttctc atactctgac ctccgcacg tcaccaaaca cgttctgcc 2220
tatcgacca cgacaggcat gaccacgcaa gcacctatca tcggacccct cccagccca 2280
aacctcacgg agtatcaatt cctgtctact ttcaaataca tcccgccctt tatctatccc 2340
tacctcaaca gcacagatgc cctccgcgcc gcctcaaaag accctcacta cggcagcacc 2400
gcattcatte cccctcatgc cctcgacagc tcccacagc ctctcaacc agcgggtgac 2460
cccatcgctt ccggcggtaa cagtatgctc tatcaggagc tctacgaaat caccgcaaaa 2520
atcaccaaca ccggcaatct agccggtgac gaggttgctc aactgtacgt caacctcgt 2580
ggcgacaatc ctctcgaca gcttcgagga tttgaccgaa tccacctct gcccggccag 2640
agtaaggtgt tccgagcgac gttgacacgt cgggacttga gtaattggga cgttcggcg 2700
cagaattggc gcacatcag gtccgccaag gaggtttatg tcggacgctc aagtcgagat 2760
ctgcccttga gcgcgcgatt ggagtcatgg taa 2793

```

<210> 2

<211> 883

<212> PRT

<213> Penicillium oxalicum

<400> 2

```

Met Arg Ser Ser Thr Thr Val Leu Ala Phe Val Ser Tyr Gly Ser Phe
1          5          10          15

```

Gly Ile Ala Ala Ala Lys Pro Ile Gln Glu His Lys Pro Glu Leu Leu
 20 25 30
 Asn Ser Phe His Gly Phe Gln Ile Glu Glu Pro Tyr Ser Pro Pro Phe
 35 40 45
 Tyr Pro Ser Pro Trp Met Asn Pro Arg Ala Glu Gly Trp Glu Asp Ala
 50 55 60
 Tyr Gln Lys Ala Leu Ala Phe Val Ser Gln Leu Thr Leu Leu Glu Lys
 65 70 75 80
 Val Asn Leu Thr Thr Gly Val Gly Trp Glu Asn Gly Pro Cys Val Gly
 85 90 95
 Asn Thr Gly Ser Val Pro Arg Leu Gly Phe Lys Gly Leu Cys Leu Gln
 100 105 110
 Asp Ser Pro Gln Gly Val Arg Phe Ala Asp Tyr Ala Ser Ala Phe Thr
 115 120 125
 Ser Ser Gln Met Ala Ala Ala Thr Phe Asp Lys Thr Val Leu Tyr Glu
 130 135 140
 Arg Gly Val Ala Met Ala Gln Glu His Lys Gly Lys Gly Ile Thr Val
 145 150 155 160
 Gln Leu Gly Pro Val Ala Gly Pro Leu Gly Arg Ala Pro Glu Gly Gly
 165 170 175
 Arg Asn Trp Glu Gly Phe Ser Pro Asp Pro Val Leu Asn Gly Ile Ala
 180 185 190
 Met Ala Glu Thr Ile Lys Gly Met Gln Asp Thr Gly Val Leu Ala Cys
 195 200 205
 Ala Lys His Tyr Val Gly Asn Glu Gln Glu His Phe Arg Gln Ala Gly
 210 215 220
 Glu Ser Ala Gly His Gly Tyr Asn Ile Ala Glu Ser Ile Ser Ser Asn
 225 230 235 240
 Ile Asp Asp Arg Thr Met His Glu Met Tyr Leu Trp Pro Phe Ala Asp
 245 250 255
 Ala Val Arg Ala Gly Val Ala Ser Phe Met Cys Ser Tyr Asn Gln Ile
 260 265 270

Asn Asn Ser Tyr Gly Cys Gln Asn Ser His Thr Leu Asn Lys Leu Leu
 275 280 285
 Lys Ser Glu Leu Gly Phe Gln Gly Phe Val Met Ser Asp Trp Gln Ala
 290 295 300
 Gln His Ser Gly Val Ser Ser Ala Leu Ala Gly Leu Asp Met Thr Met
 305 310 315 320
 Pro Gly Asp Thr Glu Phe Asp Thr Gly Leu Ser Phe Trp Gly Ser Asn
 325 330 335
 Leu Thr Ile Ala Val Leu Asn Gly Thr Val Pro Glu Trp Arg Ile Asp
 340 345 350
 Asp Met Ala Met Arg Ile Met Ala Ala Tyr Phe Lys Val Gly Leu Thr
 355 360 365
 Ile Glu Asp Gln Pro Asp Val Asn Phe Asn Ser Trp Thr Tyr Asp Thr
 370 375 380
 Tyr Gly Tyr Lys Tyr Phe Tyr Ser Lys Glu Gly Tyr Glu Lys Val Asn
 385 390 395 400
 Trp His Val Asp Val Arg Ala Asn His Ala Gln Gln Ile Arg Glu Thr
 405 410 415
 Ala Ala Lys Gly Thr Val Leu Leu Lys Asn Thr Asn Gly Ala Leu Pro
 420 425 430
 Leu Lys Arg Pro Ser Phe Val Ala Val Val Gly Glu Asp Ala Gly Pro
 435 440 445
 Asn Pro Lys Gly Pro Asn Gly Cys Pro Asp Arg Gly Cys Asp Glu Gly
 450 455 460
 Thr Leu Ala Met Gly Trp Gly Ser Gly Ser Thr Gln Phe Pro Phe Leu
 465 470 475 480
 Val Thr Pro Asp Ser Ala Ile Gln Ala Lys Val Leu Glu Tyr Gly Gly
 485 490 495
 Arg Tyr Glu Ser Ile Leu Asp Asn Tyr Asp Glu Ser Ala Ile Ser Ala
 500 505 510
 Leu Val Ser Gln Pro Asp Ala Thr Cys Ile Val Phe Gly Asn Ser Asp
 515 520 525

Ser Gly Glu Ala Phe Ile Thr Val Asp Gly Asn Trp Gly Asp Arg Asn
 530 535 540
 Asn Leu Thr Leu Trp Gln Asn Ala Asp Ala Val Ile Arg Asn Val Ser
 545 550 555 560
 Ala Leu Cys Ser Asn Thr Ile Val Val Leu His Thr Val Gly Pro Val
 565 570 575
 Leu Leu Thr Asp Tyr Tyr Glu His Pro Asn Ile Thr Ala Ile Val Trp
 580 585 590
 Ala Gly Leu Pro Gly Glu Glu Ser Gly Asn Ala Leu Val Asp Ile Leu
 595 600 605
 Trp Gly Asp Val Asn Pro Ala Gly Arg Thr Pro Phe Thr Trp Ala Lys
 610 615 620
 Ser Arg Glu Asp Tyr Gly Thr Asp Val Met Tyr Val Pro Asn Asn Gly
 625 630 635 640
 Glu Gln Ala Pro Gln Gln Ser Phe Ser Glu Gly Ile Phe Leu Asp Tyr
 645 650 655
 Arg His Phe Asp Gln Ala Gly Ile Asp Pro Ile Tyr Glu Phe Gly His
 660 665 670
 Gly Leu Ser Tyr Thr Asn Phe Ser Tyr Ser Asp Leu Arg Ile Val Thr
 675 680 685
 Lys His Val Arg Pro Tyr Arg Pro Thr Thr Gly Met Thr Thr Gln Ala
 690 695 700
 Pro Ile Ile Gly Pro Leu Pro Ser Pro Asn Leu Thr Glu Tyr Gln Phe
 705 710 715 720
 Pro Ala Thr Phe Lys Tyr Ile Pro Ala Phe Ile Tyr Pro Tyr Leu Asn
 725 730 735
 Ser Thr Glu Ser Leu Arg Ala Ala Ser Lys Asp Pro His Tyr Gly Ser
 740 745 750
 Thr Ala Phe Ile Pro Pro His Ala Leu Asp Ser Ser Pro Gln Pro Leu
 755 760 765
 Asn Pro Ala Gly Asp Pro Ile Ala Ser Gly Gly Asn Ser Met Leu Tyr
 770 775 780
 Gln Glu Leu Tyr Glu Ile Thr Ala Lys Ile Thr Asn Thr Gly Asn Leu
 785 790 795 800
 Ala Gly Asp Glu Val Val Gln Leu Tyr Val Asn Leu Gly Gly Asp Asn
 805 810 815
 Pro Pro Arg Gln Leu Arg Gly Phe Asp Arg Ile His Leu Leu Pro Gly
 820 825 830
 Gln Ser Lys Val Phe Arg Ala Thr Leu Thr Arg Arg Asp Leu Ser Asn
 835 840 845
 Trp Asp Val Ala Ala Gln Asn Trp Arg Ile Ile Arg Ser Ala Lys Glu
 850 855 860
 Val Tyr Val Gly Arg Ser Ser Arg Asp Leu Pro Leu Ser Ala Arg Leu
 865 870 875 880
 Glu Ser Trp

<210> 3

<211> 35

<212> DNA

<213> Penicillium oxalicum

<220>

<221> misc_feature
 <222> (27) .. (27)
 <223> N=A,C,G or T

 <220>
 <221> misc_feature
 <222> (30) .. (33)
 <223> N=A,C,G or T

 <220>
 <221> misc_feature
 <222> (33) .. (33)
 <223> N=A,C,G or T

 <400> 3
 atgaccctgg ccgaaaaagt caacytnacn acngg 35

 <210> 4
 <211> 35
 <212> DNA
 <213> Penicillium oxalicum

 <220>
 <221> misc_feature
 <222> (27) .. (27)
 <223> N=A,C,G or T

 <220>
 <221> misc_feature
 <222> (30)..(30)
 <223> N=A,C,G or T

 <400> 4
 ggtggccgga actgggaagg cttctsnccn gaycc 35

 <210> 5
 <211> 36
 <212> DNA
 <213> Penicillium oxalicum

 <220>
 <221> misc_feature
 <222> (24) .. (24)
 <223> N=A,C,G or T

 <220>
 <221> misc_feature
 <222> (30) .. (30)
 <223> N=A,C,G or T

 <400> 5
 gagctgggct tccagggctt tgtnatgwsn gaytgg 36

 <210> 6
 <211> 32
 <212> DNA
 <213> Penicillium oxalicum

 <220>
 <221> misc_feature
 <222> (27) .. (27)
 <223> N=A,C,T,OR G

<400> 6
agcgcttgg ccggcctcga yatgwsnatg cc 32

<210> 7
<211> 27
<212> DNA
<213> Penicillium oxalicum

<220>
<221> misc_feature
<222> (22) .. (22)
<223> N=A,C,G ORT

<220>
<221> misc_feature
<222> (25) .. (25)
<223> N=A,C,G ORT

<400> 7
atcccagttg ctcaggctcc knckngt 27

<210> 8
<211> 33
<212> DNA
<213> Penicillium oxalicum

<220>
<221> misc_feature
<222> (19) .. (19)
<223> N=A, C, G, OR T

<220>
<221> misc_feature
<222> (25) .. (25)
<223> N=A, C, G, OR T

<400> 8
aaaggtgtg tagctcagnc crtgnccraa ytc 33

<210> 9
<211> 33
<212> DNA
<213> Penicillium oxalicum

<220>
<221> misc_feature
<222> (25) .. (25)
<223> N=A, C, T, OR G

<220>
<221> misc_feature
<222> (28) .. (28)
<223> N=A, C, T, OR G

<400> 9
gtcaaagtgg cggtagtcga traanacncc ytc 33

<210> 10
<211> 39
<212> DNA
<213> Penicillium oxalicum

<220>
 <221> misc_feature
 <222> (37) .. (37)
 <223> N=A,C,T, OR G

 <400> 10
 ggtgggcgag ttgccgacgg gggtgactct gcortanar 39

 <210> 11
 <211> 39
 <212> DNA
 <213> Penicillium oxalicum

 <220>
 <221> misc_feature
 <222> (28) .. (28)
 <223> N=A,C,G. OR T

 <220>
 <221> misc_feature
 <222> (31)..(31)
 <223> N=A,C,G. OR T

 <220>
 <221> misc_feature
 <222> (37) .. (37)
 <223> N=A,C,G. OR T

 <400> 11
 gccgggcaga ccggcccaga ggatggcngt nacrtngg 39

 <210> 12
 <211> 39
 <212> DNA
 <213> Penicillium oxalicum

 <220>
 <221> misc_feature
 <222> (28) .. (28)
 <223> N=A,C,G, OR T

 <220>
 <221> misc_feature
 <222> (34) .. (34)
 <223> N=A,C,G, OR T

 <400> 12
 caggacgggg ccaaccgagt gaatgacnac datngtrtt 39

 <210> 13
 <211> 21
 <212> DNA
 <213> Penicillium oxalicum

 <400> 13
 caccaacacc ggcaatctag c 21

 <210> 14
 <211> 25
 <212> DNA
 <213> Penicillium oxalicum

<400> 14
 ggtgacgagg ttgtccaact gtacg 25

 <210> 15
 <211> 18
 <212> DNA
 <213> Penicillium oxalicum

 <400> 15
 ctgaagcca aggcgagg 18

 <210> 16
 <211> 24
 <212> DNA
 <213> Penicillium oxalicum

 <400> 16
 tccggtattt cctacacatg gtcc 24

 <210> 17
 <211> 43
 <212> DNA
 <213> Penicillium oxalicum

 <400> 17
 agatctcgag aagcttacca tgactccaat cgcgcgctca agg 43

 <210> 18
 <211> 46
 <212> DNA
 <213> Penicillium oxalicum

 <400> 18
 acacaactgg ggaaccacca tgaggagctc aacgacggtt ctggcc 46

 <210> 19
 <211> 19
 <212> PRT
 <213> Thielavia terrestris

 <220>
 <221> MISC_FEATURE
 <222> (1) .. (1)
 <223> X=I,L,M, OR V

 <220>
 <221> misc_feature
 <222> (3) .. (6)
 <223> Xaa can be any naturally occurring amino acid

 <220>
 <221> misc_feature
 <222> (8) .. (8)
 <223> Xaa can be any naturally occurring amino acid

 <220>
 <221> MISC_FEATURE
 <222> (10) .. (10)
 <223> X=I,L,M, OR V

 <220>
 <221> misc_feature

<222> (11) .. (11)

<223> Xaa can be any naturally occurring amino acid

<220>

<221> misc_feature

<222> (13) .. (13)

<223> Xaa can be any naturally occurring amino acid

<220>

<221> MISC_FEATURE

<222> (14) .. (14)

<223> X=E OR Q

<220>

<221> misc_feature

<222> (15) .. (18)

<223> Xaa can be any naturally occurring amino acid

<220>

<221> MISC_FEATURE

<222> (19) .. (19)

<223> X=H,N, OR Q

<400> 19

Xaa Pro Xaa Xaa Xaa Xaa Gly Xaa Tyr Xaa Xaa Arg Xaa Xaa Xaa Xaa
1 5 10 15

Xaa Xaa Xaa

<210> 20

<211> 20

<212> PRT

<213> Thielavia terrestris

<220>

<221> MISC_FEATURE

<222> (1) .. (1)

<223> X=I,L,M, OR V

<220>

<221> misc_feature

<222> (3) .. (7)

<223> Xaa can be any naturally occurring amino acid

<220>

<221> misc_feature

<222> (9) .. (9)

<223> Xaa can be any naturally occurring amino acid

<220>

<221> MISC_FEATURE

<222> (11) .. (11)

<223> X=I,L,M, OR V

<220>

<221> misc_feature

<222> (12) .. (12)

<223> Xaa can be any naturally occurring amino acid

<220>

<221> misc_feature

<222> (14) .. (14)

<223> Xaa can be any naturally occurring amino acid

<220>
 <221> MISC_FEATURE
 <222> (15) .. (15)
 <223> X=E OR Q

<220>
 <221> misc_feature
 <222> (16) .. (19)
 <223> Xaa can be any naturally occurring amino acid

<220>
 <221> MISC_FEATURE
 <222> (20)..(20)
 <223> X=H,N, OR Q

<400> 20
 Xaa Pro Xaa Xaa Xaa Xaa Gly Xaa Tyr Xaa Xaa Arg Xaa Xaa Xaa
 1 5 10 15

Xaa Xaa Xaa Xaa
 20

<210> 21
 <211> 9
 <212> PRT
 <213> Thielavia terrestris

<220>
 <221> misc_feature
 <222> (2) .. (2)
 <223> Xaa can be any naturally occurring amino acid

<220>
 <221> misc_feature
 <222> (5) .. (7)
 <223> Xaa can be any naturally occurring amino acid

<220>
 <221> MISC_FEATURE
 <222> (8)..(8)
 <223> X= Y OR W

<220>
 <221> MISC_FEATURE
 <222> (9) .. (9)
 <223> X= A,I,L,M OR V

<400> 21
 His Xaa Gly Pro Xaa Xaa Xaa Xaa Xaa
 1 5

<210> 22
 <211> 10
 <212> PRT
 <213> Thielavia terrestris

<220>
 <221> misc_feature
 <222> (2) .. (3)
 <223> Xaa can be any naturally occurring amino acid

<220>
 <221> misc_feature

<222> (6) .. (8)
 <223> Xaa can be any naturally occurring amino acid

 <220>
 <221> MISC_FEATURE
 <222> (9) .. (9)
 <223> X= Y OR W

 <220>
 <221> MISC_FEATURE
 <222> (10)..(10)
 <223> X= Y OR W

 <400> 22
 His Xaa Xaa Gly Pro Xaa Xaa Xaa Xaa Xaa
 1 5 10

 <210> 23
 <211> 11
 <212> PRT
 <213> Thielavia terrestris

 <220>
 <221> MISC_FEATURE
 <222> (1) .. (1)
 <223> X= E OR Q

 <220>
 <221> misc_feature
 <222> (2) .. (2)
 <223> Xaa can be any naturally occurring amino acid

 <220>
 <221> misc_feature
 <222> (4) .. (5)
 <223> Xaa can be any naturally occurring amino acid

 <220>
 <221> misc_feature
 <222> (7) .. (7)
 <223> Xaa can be any naturally occurring amino acid

 <220>
 <221> MISC_FEATURE
 <222> (8) .. (8)
 <223> X= E,H,Q OR N

 <220>
 <221> MISC_FEATURE
 <222> (9) .. (9)
 <223> X=F,I,L, OR V

 <220>
 <221> misc_feature
 <222> (10) .. (10)
 <223> Xaa can be any naturally occurring amino acid

 <220>
 <221> MISC_FEATURE
 <222> (11) .. (11)
 <223> X=I,L,OR V

 <400> 23

Xaa Xaa Tyr Xaa Xaa Cys Xaa Xaa Xaa Xaa Xaa
 1 5 10

<210> 24

<211> 9

<212> PRT

<213> Thielavia terrestris

<220>

<221> misc_feature

<222> (2) .. (2)

<223> Xaa can be any naturally occurring amino acid

<220>

<221> misc_feature

<222> (5) .. (7)

<223> Xaa can be any naturally occurring amino acid

<220>

<221> MISC_FEATURE

<222> (8) .. (8)

<223> X= Y OR W

<220>

<221> MISC_FEATURE

<222> (9) .. (9)

<223> X= A,I,L,M OR V

<220>

<221> MISC_FEATURE

<222> (9) .. (9)

<223> X= A,I,L,M OR V

<400> 24

His Xaa Gly Pro Xaa Xaa Xaa Xaa Xaa
 1 5

<210> 25

<211> 10

<212> PRT

<213> Thielavia terrestris

<220>

<221> misc_feature

<222> (2)..(3)

<223> Xaa can be any naturally occurring amino acid

<220>

<221> misc_feature

<222> (6) .. (8)

<223> Xaa can be any naturally occurring amino acid

<220>

<221> MISC_FEATURE

<222> (9) .. (9)

<223> X= Y OR W

<220>

<221> MISC_FEATURE

<222> (10) .. (10)

<223> X= A,I,L,M OR V

<400> 25

His Xaa Xaa Gly Pro Xaa Xaa Xaa Xaa Xaa

1	5	10
<210> 26		
<211> 11		
<212> PRT		
<213> Thielavia terrestris		
<220>		
<221> MISC_FEATURE		
<222> (1) .. (1)		
<223> X= E OR Q		
<220>		
<221> misc_feature		
<222> (2) .. (2)		
<223> Xaa can be any naturally occurring amino acid		
<220>		
<221> misc_feature		
<222> (4) .. (5)		
<223> Xaa can be any naturally occurring amino acid		
<220>		
<221> misc_feature		
<222> (7) .. (7)		
<223> Xaa can be any naturally occurring amino acid		
<220>		
<221> MISC_FEATURE		
<222> (8) .. (8)		
<223> X= E,H,Q OR N		
<220>		
<221> MISC_FEATURE		
<222> (9) .. (9)		
<223> X=F,I,L, OR V		
<220>		
<221> misc_feature		
<222> (10) .. (10)		
<223> Xaa can be any naturally occurring amino acid		
<220>		
<221> MISC_FEATURE		
<222> (11) .. (11)		
<223> X=I,L,OR V		
<400> 26		
Xaa Xaa Tyr Xaa Xaa Cys Xaa Xaa Xaa Xaa Xaa		
1	5	10
<210> 27		
<211> 9		
<212> PRT		
<213> Thielavia terrestris		
<220>		
<221> misc_feature		
<222> (2) .. (2)		
<223> Xaa can be any naturally occurring amino acid		
<220>		

<221> misc_feature
 <222> (5) .. (7)
 <223> Xaa can be any naturally occurring amino acid

<220>
 <221> MISC_FEATURE
 <222> (8) .. (8)
 <223> X= Y OR W

<220>
 <221> MISC_FEATURE
 <222> (9) .. (9)
 <223> X= A,I,L,M OR V

<400> 27
 His Xaa Gly Pro Xaa Xaa Xaa Xaa Xaa
 1 5

<210> 28
 <211> 10
 <212> PRT
 <213> Thielavia terrestris

<220>
 <221> misc_feature
 <222> (2) .. (3)
 <223> Xaa can be any naturally occurring amino acid

<220>
 <221> misc_feature
 <222> (6) .. (8)
 <223> Xaa can be any naturally occurring amino acid

<220>
 <221> MISC_FEATURE
 <222> (9) .. (9)
 <223> X= Y OR W

<220>
 <221> MISC_FEATURE
 <222> (10) .. (10)
 <223> X= A,I,L,M OR V

<400> 28
 His Xaa Xaa Gly Pro Xaa Xaa Xaa Xaa Xaa
 1 5 10

<210> 29
 <211> 11
 <212> PRT
 <213> Thielavia terrestris

<220>
 <221> MISC_FEATURE
 <222> (1) .. (1)
 <223> X= E OR Q

<220>
 <221> misc_feature
 <222> (2) .. (2)
 <223> Xaa can be any naturally occurring amino acid

<220>

<221> misc_feature
 <222> (4) .. (5)
 <223> Xaa can be any naturally occurring amino acid

<220>
 <221> misc_feature
 <222> (7) .. (7)
 <223> Xaa can be any naturally occurring amino acid

<220>
 <221> MISC_FEATURE
 <222> (8) .. (8)
 <223> X= E,H,Q OR N

<220>
 <221> MISC_FEATURE
 <222> (9) .. (9)
 <223> X=F,I,L, OR V

<220>
 <221> misc_feature
 <222> (10) .. (10)
 <223> Xaa can be any naturally occurring amino acid

<220>
 <221> MISC_FEATURE
 <222> (11) .. (11)
 <223> X=I,L, OR V

<220>
 <221> MISC_FEATURE
 <222> (11) .. (11)
 <223> X=I,L, OR V

<400> 29
 Xaa Xaa Tyr Xaa Xaa Cys Xaa Xaa Xaa Xaa Xaa
 1 5 10

<210> 30
 <211> 9
 <212> PRT
 <213> Thielavia terrestris

<220>
 <221> misc_feature
 <222> (2) .. (2)
 <223> Xaa can be any naturally occurring amino acid

<220>
 <221> misc_feature
 <222> (5) .. (7)
 <223> Xaa can be any naturally occurring amino acid

<220>
 <221> MISC_FEATURE
 <222> (8) .. (8)
 <223> X= Y OR W

<220>
 <221> MISC_FEATURE
 <222> (9) .. (9)
 <223> X= A,I,L,M OR V

```

<220>
<221> MISC_FEATURE
<222> (9)..(9)
<223> X= A,I,L,M OR V

<400> 30
His Xaa Gly Pro Xaa Xaa Xaa Xaa Xaa
1           5

<210> 31
<211> 10
<212> PRT
<213> Thielavia terrestris

<220>
<221> misc_feature
<222> (2) .. (3)
<223> Xaa can be any naturally occurring amino acid

<220>
<221> misc_feature
<222> (6) .. (8)
<223> Xaa can be any naturally occurring amino acid

<220>
<221> MISC_FEATURE
<222> (9) .. (9)
<223> X= Y OR W

<220>
<221> MISC_FEATURE
<222> (10) .. (10)
<223> X= A,I,L,M OR V

<400> 31
His Xaa Xaa Gly Pro Xaa Xaa Xaa Xaa Xaa
1           5           10

<210> 32
<211> 11
<212> PRT
<213> Thielavia terrestris

<220>
<221> MISC_FEATURE
<222> (1) .. (1)
<223> X= E OR Q

<220>
<221> misc_feature
<222> (2) .. (2)
<223> Xaa can be any naturally occurring amino acid

<220>
<221> misc_feature
<222> (4) .. (5)
<223> Xaa can be any naturally occurring amino acid

<220>
<221> misc_feature
<222> (7) .. (7)
<223> Xaa can be any naturally occurring amino acid

```

<220>
 <221> MISC_FEATURE
 <222> (8) .. (8)
 <223> X= E,H,Q OR N

<220>
 <221> MISC_FEATURE
 <222> (9) .. (9)
 <223> X=F,I,L, OR V

<220>
 <221> misc_feature
 <222> (10) .. (10)
 <223> Xaa can be any naturally occurring amino acid

<220>
 <221> MISC_FEATURE
 <222> (11) .. (11)
 <223> X=I,L, OR V

<400> 32
 Xaa Xaa Tyr Xaa Xaa Cys Xaa Xaa Xaa Xaa Xaa
 1 5 10

<210> 33
 <211> 19
 <212> PRT
 <213> Thielavia terrestris

<220>
 <221> MISC_FEATURE
 <222> (1) .. (1)
 <223> X=I,L,M OR V

<220>
 <221> misc_feature
 <222> (3) .. (6)
 <223> Xaa can be any naturally occurring amino acid

<220>
 <221> misc_feature
 <222> (8) .. (8)
 <223> Xaa can be any naturally occurring amino acid

<220>
 <221> MISC_FEATURE
 <222> (10) .. (10)
 <223> X=I,L,M OR V

<220>
 <221> misc_feature
 <222> (11)..(11)
 <223> Xaa can be any naturally occurring amino acid

<220>
 <221> misc_feature
 <222> (13) .. (13)
 <223> Xaa can be any naturally occurring amino acid

<220>
 <221> MISC_FEATURE
 <222> (14) .. (14)

<223> X=I,L,M OR V

<220>

<221> MISC_FEATURE

<222> (14) .. (14)

<223> X= E OR Q

<220>

<221> misc_feature

<222> (15) .. (17)

<223> Xaa can be any naturally occurring amino acid

<220>

<221> MISC_FEATURE

<222> (19) .. (19)

<223> X= H,N, OR Q

<400> 33

Xaa	Pro	Xaa	Xaa	Xaa	Xaa	Gly	Xaa	Tyr	Xaa	Xaa	Arg	Xaa	Xaa	Xaa	Xaa
1				5					10					15	

Xaa Ala Xaa

<210> 34

<211> 20

<212> PRT

<213> Thielavia terrestris

<220>

<221> MISC_FEATURE

<222> (1) .. (1)

<223> X=I,L,M OR V

<220>

<221> misc_feature

<222> (3).. (7)

<223> Xaa can be any naturally occurring amino acid

<220>

<221> misc_feature

<222> (9) .. (9)

<223> Xaa can be any naturally occurring amino acid

<220>

<221> MISC_FEATURE

<222> (11) .. (11)

<223> X=I,L,M OR V

<220>

<221> misc_feature

<222> (12) .. (12)

<223> Xaa can be any naturally occurring amino acid

<220>

<221> misc_feature

<222> (14) .. (14)

<223> Xaa can be any naturally occurring amino acid

<220>

<221> MISC_FEATURE

<222> (15) .. (15)

<223> X=E OR Q

<220>

<221> misc_feature

<222> (16) .. (18)

<223> Xaa can be any naturally occurring amino acid

<220> <221> MISC_FEATURE <222> (20) .. (20) <223> X= H,N, OR Q

<400> 34

Xaa Pro Xaa Xaa Xaa Xaa Gly Xaa Tyr Xaa Xaa Arg Xaa Xaa Xaa
1 5 10 15

Xaa Xaa Ala Xaa
20

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- [WO02095014A \[0027\]](#)
- [WO2002095014A \[0027\] \[0258\]](#)
- [WO9517413A \[0080\]](#)
- [WO9522625A \[0080\]](#)
- [US5223409A \[0080\]](#)
- [WO9206204A \[0080\]](#)
- [WO9943835A \[0107\]](#)
- [WO9600787A \[0108\] \[0156\]](#)
- [WO0056900A \[0108\] \[0108\] \[0108\]](#)
- [WO9425612A \[0115\]](#)
- [WO9533836A \[0126\]](#)
- [WO0024883A \[0139\] \[0139\]](#)
- [EP238023A \[0156\]](#)
- [WO9114772A \[0172\]](#)
- [US6395966B \[0176\]](#)
- [US7151204B \[0176\] \[0176\]](#)
- [US6489127B \[0192\]](#)
- [US6506559B \[0192\]](#)
- [US6511824B \[0192\]](#)
- [US6515109B \[0192\]](#)
- [US20020164730A \[0213\] \[0222\]](#)
- [WO2006110891A \[0217\]](#)
- [WO2006110899A \[0217\]](#)
- [WO2006110900A \[0217\]](#)
- [WO2006110901A \[0217\]](#)
- [WO2006032282A \[0219\]](#)
- [WO9117243A \[0253\]](#)
- [WO9117244A \[0253\]](#)
- [WO9105039A \[0255\]](#)
- [WO9315186A \[0255\]](#)
- [US5275944A \[0255\]](#)
- [WO9602551A \[0255\]](#)

- [US5536655A \[0255\]](#)
- [WO0070031A \[0255\]](#)
- [WO05093050A \[0255\] \[0255\] \[0255\]](#)
- [WO2011059740A \[0257\]](#)
- [WO2009042871A \[0257\]](#)
- [WO2010141325A \[0257\]](#)
- [WO2006074435A \[0257\]](#)
- [WO2010057086A \[0257\]](#)
- [WO2005047499A \[0258\]](#)
- [WO2007019442A \[0258\] \[0258\]](#)
- [WO2010088387A \[0258\]](#)
- [WO2011035029A \[0258\]](#)
- [WO2008057637A \[0259\] \[0259\]](#)
- [EP495257A \[0261\]](#)
- [EP531315A \[0261\]](#)
- [EP531372A \[0261\]](#)
- [WO8909259A \[0261\]](#)
- [WO9407998A \[0261\]](#)
- [WO9524471A \[0261\]](#)
- [WO9611262A \[0261\]](#)
- [WO9629397A \[0261\]](#)
- [WO96034108A \[0261\]](#)
- [WO9714804A \[0261\]](#)
- [WO9808940A \[0261\]](#)
- [WO98012307A \[0261\]](#)
- [WO9813465A \[0261\]](#)
- [WO98015619A \[0261\]](#)
- [WO98015633A \[0261\]](#)
- [WO98028411A \[0261\]](#)
- [WO9906574A \[0261\]](#)
- [WO9910481A \[0261\]](#)
- [WO99025846A \[0261\]](#)
- [WO99025847A \[0261\]](#)
- [WO99031255A \[0261\]](#)
- [WO2000009707A \[0261\]](#)
- [WO2002050245A \[0261\]](#)
- [WO20020076792A \[0261\]](#)
- [WO2002101078A \[0261\]](#)
- [WO2003027306A \[0261\]](#)
- [WO2003052054A \[0261\]](#)
- [WO2003052055A \[0261\]](#)
- [WO2003052056A \[0261\]](#)
- [WO2003052057A \[0261\]](#)
- [WO2003052118A \[0261\]](#)
- [WO2004016760A \[0261\]](#)
- [WO2004043980A \[0261\]](#)
- [WO2004048592A \[0261\]](#)
- [WO2005001065A \[0261\]](#)
- [WO2005028636A \[0261\]](#)
- [WO2005093050A \[0261\]](#)
- [WO2005093073A \[0261\]](#)
- [WO2006074005A \[0261\]](#)
- [WO2006117432A \[0261\]](#)
- [WO2007071818A \[0261\]](#)
- [WO2007071820A \[0261\]](#)
- [WO2008008070A \[0261\]](#)
- [WO2008008793A \[0261\]](#)
- [US4435307A \[0261\]](#)

- [US5457046A \[0261\]](#)
- [US5648263A \[0261\]](#)
- [US5686593A \[0261\]](#)
- [US5691178A \[0261\]](#)
- [US5763254A \[0261\]](#)
- [US5776757A \[0261\]](#)
- [WO2005074647A \[0267\]](#)
- [WO2008148131A \[0267\]](#)
- [WO2011035027A \[0267\]](#)
- [WO2005074656A \[0267\]](#)
- [WO2010065830A \[0267\]](#)
- [WO2007089290A \[0267\]](#)
- [WO2009085935A \[0267\]](#)
- [WO2009085859A \[0267\]](#)
- [WO2009085864A \[0267\]](#)
- [WO2009085868A \[0267\]](#)
- [WO2010138754A \[0267\]](#)
- [WO2011005867A \[0267\]](#)
- [WO2011039319A \[0267\]](#)
- [WO2011041397A \[0267\]](#)
- [WO2011041504A \[0267\]](#)
- [WO2008151043A \[0268\]](#)
- [WO9421785A \[0280\]](#)
- [WO2006078255A \[0280\]](#)
- [WO2011041405A \[0280\]](#)
- [WO2010126772A \[0280\]](#)
- [WO2009079210A \[0280\]](#)
- [WO2011057083A \[0280\]](#)
- [WO2010108918A \[0282\]](#)
- [WO2009073709A \[0282\]](#)
- [WO2005001036A \[0282\]](#)
- [WO2010014880A \[0282\]](#)
- [WO2009042846A \[0282\]](#)
- [WO2009076122A \[0283\]](#)
- [WO2009127729A \[0283\]](#)
- [WO2010053838A \[0283\]](#)
- [WO2010065448A \[0283\]](#)
- [WO2006114094A \[0284\] \[0284\]](#)
- [WO2009073383A \[0284\]](#)
- [WO2010014706A \[0285\]](#)
- [WO2009068565A \[0285\]](#)
- [WO2003062430A \[0301\]](#)
- [WO2005042735A \[0339\] \[0341\]](#)
- [WO200039322A \[0346\]](#)
- [WO95002043A \[0346\]](#)
- [US61360351B \[0361\]](#)

Non-patent literature cited in the description

- **VENTURI et al.** Extracellular beta-D-glucosidase from *Chaetomium thermophilum* var. *coprophilum*: production, purification and some biochemical properties. *J. Basic Microbiol.*, 2002, vol. 42, 55-66 [\[0018\]](#)
- **VRIESJ.** *Bacteriol.*, 1998, vol. 180, 243-249 [\[0023\]](#)
- **TEERI** Crystalline cellulose degradation: New insight into the function of cellobiohydrolases *Trends in Biotechnology*, 1997, vol. 15, 160-167 [\[0026\]](#)

- **TEERI et al.** Trichoderma reesei cellobiohydrolases: why so efficient on crystalline cellulose? *Biochem. Soc. Trans.*, 1998, vol. 26, 173-178 [0026]
- **LEVER et al.** *Anal. Biochem.*, 1972, vol. 47, 273-279 [0026]
- **TILBEURGH et al.** *FEBS Letters*, 1982, vol. 149, 152-156 [0026]
- **TILBEURGHCLAESSSENS** *FEBS Letters*, 1985, vol. 187, 283-288 [0026]
- **TOMME et al.** *Eur. J. Biochem.*, 1988, vol. 170, 575-581 [0026]
- **ZHANG et al.** Outlook for cellulase improvement: Screening and selection strategies *Biotechnology Advances*, 2006, vol. 24, 452-481 [0029]
- **GHOSE** Measurement of cellulase activities *Pure Appl. Chem.*, 1987, vol. 59, 257-68 [0029]
- **WISELOGEL et al.** *Handbook on Bioethanol* 19950000105-118 [0032]
- **TAYLORFRANCIS** *Bioresource Technology*, 1994, vol. 50, 3-16 [0032]
- **LYND** *Applied Biochemistry and Biotechnology*, 1990, vol. 24/25, 695-719 [0032]
- Recent Progress in Bioconversion of Lignocellulosics **MOSIER et al.** *Advances in Biochemical Engineering/Biotechnology* Springer-Verlag 19990000vol. 65, 23-40 [0032]
- **ZHANG et al.** *Biotechnology Advances*, 2006, vol. 24, 452-481 [0041]
- **GHOSE** *Pure and Appl. Chem.*, 1987, vol. 59, 257-268 [0041]
- **HENRISSAT B.** A classification of glycosyl hydrolases based on amino-acid sequence similarities *Biochem. J.*, 1991, vol. 280, 309-316 [0044] [0260]
- **HENRISSAT B. BAIROCH A.** Updating the sequence-based classification of glycosyl hydrolases *Biochem. J.*, 1996, vol. 316, 695-696 [0044] [0045] [0260]
- **HENRISSAT A.** A classification of glycosyl hydrolases based on amino-acid sequence similarities *Biochem. J.*, 1991, vol. 280, 309-316 [0045]
- **SHALLOM, D. SHOHAM, Y.** Microbial hemicellulases *Current Opinion In Microbiology*, 2003, vol. 6, 3219-228 [0048]
- **GHOSE BISARIA** *Pure & Appl. Chem.*, 1987, vol. 59, 1739-1752 [0048]
- **NIELSEN et al.** *Protein Engineering*, 1997, vol. 10, 1-6 [0051] [0343]
- **NEEDLEMAN WUNSCH J.** *Mol. Biol.*, 1970, vol. 48, 443-453 [0057] [0344]
- **RICE et al.** *EMBOSS: The European Molecular Biology Open Software Suite* *Trends Genet.*, 2000, vol. 16, 276-277 [0057]
- **BIELY PUCHARD** Recent progress in the assays of xylanolytic enzymes *Journal of the Science of Food and Agriculture*, 2006, vol. 86, 111636-1647 [0061]
- **SPANIKOVABIELY** Glucuronoyl esterase - Novel carbohydrate esterase produced by *Schizophyllum commune* *FEBS Letters*, 2006, vol. 580, 194597-4601 [0061]
- **HERRMANN VRSANSKA JURICKOVA HIRSCHBIELY KUBICEK** The beta-D-xylosidase of *Trichoderma reesei* is a multifunctional beta-D-xylan xylohydrolase *Biochemical Journal*, 1997, vol. 321, 375-381 [0061]
- **BAILEYBIELY POUTANEN** Interlaboratory testing of methods for assay of xylanase activity *Journal of Biotechnology*, 1992, vol. 23, 3257-270 [0062]
- **LEVER** A new reaction for colorimetric determination of carbohydrates *Anal. Biochem.*, 1972, vol. 47, 273-279 [0063]
- **J. SAMBROOKE. F. FRITSCHT. MANIATIS** *Molecular Cloning, A Laboratory Manual* Cold Spring Harbor 19890000 [0068]
- **BOLTON MCCARTHY** *Proc. Natl. Acad. Sci. USA*, 1962, vol. 48, 1390- [0074]
- **H. NEURATH R. L. HILL** *In, The Proteins* Academic Press 19790000 [0077]
- **CUNNINGHAM WELLS** *Science*, 1989, vol. 244, 1081-1085 [0079]
- **HILTON et al.** *J. Biol. Chem.*, 1996, vol. 271, 4699-4708 [0079]
- **VOS et al.** *Science*, 1992, vol. 255, 306-312 [0079]
- **SMITH et al.** *J. Mol. Biol.*, 1992, vol. 224, 899-904 [0079]
- **WLODAVER et al.** *FEBS Lett.*, 1992, vol. 309, 59-64 [0079]
- **REIDHAAR-OLSON SAUER** *Science*, 1988, vol. 241, 53-57 [0080]
- **BOWIE SAUER** *Proc. Natl. Acad. Sci. USA*, 1989, vol. 86, 2152-2156 [0080]
- **LOWMAN et al.** *Biochemistry*, 1991, vol. 30, 10832-10837 [0080]
- **DERBYSHIRE et al.** *Gene*, 1986, vol. 46, 145- [0080]
- **NER et al.** *DNA*, 1988, vol. 7, 127- [0080]
- **NESS et al.** *Nature Biotechnology*, 1999, vol. 17, 893-896 [0081]
- **COOPER et al.** *EMBO J.*, 1993, vol. 12, 2575-2583 [0084]
- **DAWSON et al.** *Science*, 1994, vol. 266, 776-779 [0084]
- **MARTIN et al.** *J. Ind. Microbiol. Biotechnol.*, 2003, vol. 3, 568-576 [0085]
- **SVETINA et al.** *J. Biotechnol.*, 2000, vol. 76, 245-251 [0085]
- **RASMUSSEN-WILSON et al.** *Appl. Environ. Microbiol.*, 1997, vol. 63, 3488-3493 [0085]
- **WARD et al.** *Biotechnology*, 1995, vol. 13, 498-503 [0085]
- **CONTRERAS et al.** *Biotechnology*, 1991, vol. 9, 378-381 [0085]
- **EATON et al.** *Biochemistry*, 1986, vol. 25, 505-512 [0085]

- **COLLINS-RACIE et al.**Biotechnology, 1995, vol. 13, 982-987 [0085]
- **CARTER et al.**Proteins: Structure, Function, and Genetics, 1989, vol. 6, 240-248 [0085]
- **STEVENS**Drug Discovery World, 2003, vol. 4, 35-48 [0085]
- **INNIS et al.**PCR: A Guide to Methods and ApplicationAcademic Press19900000 [0099]
- **FORD et al.**Protein Expression and Purification, 1991, vol. 2, 95-107 [0101]
- **AGAISSELERECLUS**Molecular Microbiology, 1994, vol. 13, 97-107 [0107]
- **EGON et al.**Gene, 1988, vol. 69, 301-315 [0107]
- **VILLA-KAMAROFF et al.**Proc. Natl. Acad. Sci. USA, 1978, vol. 75, 3727-3731 [0107]
- **DEBOER et al.**Proc. Natl. Acad. Sci. USA, 1983, vol. 80, 21-25 [0107]
- **GILBERT et al.**Useful proteins from recombinant bacteriaScientific American, 1980, vol. 242, 74-94 [0107]
- **ROMANOS et al.**Yeast, 1992, vol. 8, 423-488 [0109]
- **HUE et al.**Journal of Bacteriology, 1995, vol. 177, 3465-3471 [0115]
- **GUOSHHERMAN**Mol. Cellular Biol., 1995, vol. 15, 5983-5990 [0121]
- **SIMONENPALVA**Microbiological Reviews, 1993, vol. 57, 109-137 [0123]
- **GEMS et al.**Gene, 1991, vol. 98, 61-67 [0139]
- **CULLEN et al.**Nucleic Acids Res., 1987, vol. 15, 9163-9175 [0139]
- **CHANGCOHEN**Mol. Gen. Genet., 1979, vol. 168, 111-115 [0148]
- **YOUNGSPIZEN**J. Bacteriol., 1961, vol. 81, 823-829 [0148]
- **DUBNAUDAVIDOFF-ABELSON**J. Mol. Biol., 1971, vol. 56, 209-221 [0148]
- **SHIGEKAWADOWER**Biotechniques, 1988, vol. 6, 742-751 [0148]
- **KOEHLERTHORNE**J. Bacteriol., 1987, vol. 169, 5271-5278 [0148]
- **HANAHAN**J. Mol. Biol., 1983, vol. 166, 557-580 [0148]
- **DOWER et al.**Nucleic Acids Res., 1988, vol. 16, 6127-6145 [0148]
- **GONG et al.**Folia Microbiol. (Praha), 2004, vol. 49, 399-405 [0148]
- **MAZODIER et al.**J. Bacteriol., 1989, vol. 171, 3583-3585 [0148]
- **BURKE et al.**Proc. Natl. Acad. Sci. USA, 2001, vol. 98, 6289-6294 [0148]
- **CHOI et al.**J. Microbiol. Methods, 2006, vol. 64, 391-397 [0148]
- **PINEDOSMETS**Appl. Environ. Microbiol., 2005, vol. 71, 51-57 [0148]
- **PERRYKURAMITSU**Infect Immun., 1981, vol. 32, 1295-1297 [0148]
- **CATTJOLLI**Microbios, 1991, vol. 68, 189-207 [0148]
- **BUCKLEY et al.**Appl. Environ. Microbiol., 1999, vol. 65, 3800-3804 [0148]
- **CLEWELL**Microbiol. Rev., 1981, vol. 45, 409-436 [0148]
- **HAWKSWORTH et al.**In, Ainsworth and Bisby's Dictionary of The FungiCAB International, University Press19950000 [0150]
- **Biology and Activities of Yeast**Soc. App. Bacteriol. Symposium Series19800000 [0151]
- **YELTON et al.**Proc. Natl. Acad. Sci. USA, 1984, vol. 81, 1470-1474 [0156]
- **CHRISTENSEN et al.**Bio/Technology, 1988, vol. 6, 1419-1422 [0156]
- **MALARDIER et al.**Gene, 1989, vol. 78, 147-156 [0156]
- **BECKERGUARENTE**Guide to Yeast Genetics and Molecular Biology, Methods in EnzymologyAcademic Press, Inc.vol. 194, 182-187 [0156]
- **ITO et al.**J. Bacteriol., 1983, vol. 153, 163- [0156]
- **HINNEN et al.**75Proc. Natl. Acad. Sci. USA, 1978, 1920- [0156]
- **Protein Purification**VCH Publishers19890000 [0162]
- **TAGUE et al.**86Plant Physiology, 1988, 506- [0171]
- **FRANCK et al.**Cell, 1980, vol. 21, 285-294 [0172]
- **CHRISTENSEN et al.**Plant Mol. Biol., 1992, vol. 18, 675-689 [0172]
- **ZHANG et al.**Plant Cell, 1991, vol. 3, 1155-1165 [0172]
- **EDWARDS**CORUZZIAnn. Rev. Genet., 1990, vol. 24, 275-303 [0172]
- **ITO et al.**Plant Mol. Biol., 1994, vol. 24, 863-878 [0172]
- **WU et al.**Plant Cell Physiol., 1998, vol. 39, 885-889 [0172]
- **CONRAD et al.**J. Plant Physiol., 1998, vol. 152, 708-711 [0172]
- **CHEN et al.**Plant Cell Physiol., 1998, vol. 39, 935-941 [0172]
- **KYOZUKA et al.**Plant Physiol., 1993, vol. 102, 991-1000 [0172]
- **MITRA**HIGGINSPlant Mol. Biol., 1994, vol. 26, 85-93 [0172]
- **KAGAYA et al.**Mol. Gen. Genet., 1995, vol. 248, 668-674 [0172]
- **XU et al.**Plant Mol. Biol., 1993, vol. 22, 573-588 [0172]
- **GASSER et al.**Science, 1990, vol. 244, 1293- [0175]

- POTRYKUSBio/Technology, 1990, vol. 8, 535- [0175]
- SHIMAMOTO et al.Nature, 1989, vol. 338, 274- [0175]
- HOOKASSCHILPEROORTPlant Mol. Biol., 1992, vol. 19, 15-38 [0176]
- CHRISTOUPlant J., 1992, vol. 2, 275-281 [0176]
- SHIMAMOTOCurr. Opin. Biotechnol., 1994, vol. 5, 158-162 [0176]
- VASIL et al.Bio/Technology, 1992, vol. 10, 667-674 [0176]
- OMIRULLEH et al.Plant Mol. Biol., 1993, vol. 21, 415-428 [0176]
- Cellulose bioconversion technologyPHILIPPIDIS, G. P.Handbook on Bioethanol: Production and UtilizationTaylor & Francis19960000179-212 [0207]
- SHEEHAN, J.HIMMEL, M.Enzymes, energy and the environment: A strategic perspective on the U.S. Department of Energy's research and development activities for bioethanolBiotechnol. Prog., 1999, vol. 15, 817-827 [0207]
- LYND, L. R.WEIMER, P. J.VAN ZYL, W. H.PRETORIUS, I. S.Microbial cellulose utilization: Fundamentals and biotechnologyMicrobiol. Mol. Biol. Reviews, 2002, vol. 66, 506-577 [0207]
- FERNANDA DE CASTILHOS CORAZAFLÁVIO FARIA DE MORAES GISELLA MARIA ZANINIVO NEITZELOptimal control in fed-batch reactor for the cellobiose hydrolysisActa Scientiarum. Technology, 2003, vol. 25, 33-38 [0208]
- GUSAKOV, A. V.SINITSYN, A. P.Kinetics of the enzymatic hydrolysis of cellulose: 1. A mathematical model for a batch reactor processEnz. Microb. Technol., 1985, vol. 7, 346-352 [0208]
- RYU, S. K.LEE, J. M.Bioconversion of waste cellulose by using an attrition bioreactorBiotechnol. Bioeng., 1983, vol. 25, 53-65 [0208]
- GUSAKOV, A. V.SINITSYN, A. P.DAVYDKIN, I. Y.DAVYDKIN, V. Y.PROTAS, O. V.Enhancement of enzymatic cellulose hydrolysis using a novel type of bioreactor with intensive stirring induced by electromagnetic fieldAppl. Biochem. Biotechnol., 1996, vol. 56, 141-153 [0208]
- CHANDRA et al.Substrate pretreatment: The key to effective enzymatic hydrolysis of lignocellulose?Adv. Biochem. Engin./Biotechnol., 2007, vol. 108, 67-93 [0209]
- GALBEZACCHI Pretreatment of lignocellulosic materials for efficient bioethanol productionBiochem. Engin. / Biotechnol., 2007, vol. 108, 41-65 [0209]
- HENDRIKSZEEMAN Pretreatments to enhance the digestibility of lignocellulosic biomassBioresource Technol., 2009, vol. 100, 10-18 [0209]
- MOSIER et al.Features of promising technologies for pretreatment of lignocellulosic biomassBioresource Technol., 2005, vol. 96, 673-686 [0209]
- TAHERZADEHKARIMI Pretreatment of lignocellulosic wastes to improve ethanol and biogas production: A reviewInt. J. of Mol. Sci., 2008, vol. 9, 1621-1651 [0209]
- YANGWYMAN Pretreatment: the key to unlocking low-cost cellulosic ethanolBiofuels Bioproducts and Biorefining-Biofpr., 2008, vol. 2, 26-40 [0209]
- DUFFMURRAY Bioresource Technology, 1996, vol. 855, 1-33 [0213]
- GALBEZACCHI Appl. Microbiol. Biotechnol., 2002, vol. 59, 618-628 [0213]
- BALLESTEROS et al. Appl. Biochem. Biotechnol., 2006, 496-508 [0215]
- VARGA et al. Appl. Biochem. Biotechnol., 2004, vol. 113-116, 509-523 [0215]
- SASSNER et al. Enzyme Microb. Technol., 2006, vol. 39, 756-762 [0215]
- SCHELL et al. Bioresource Technol., 2004, vol. 91, 179-188 [0215]
- LEE et al. Adv. Biochem. Eng. Biotechnol., 1999, vol. 65, 93-115 [0215]
- WYMAN et al. Bioresource Technol., 2005, vol. 96, 1959-1966 [0217]
- MOSIER et al. Bioresource Technol., 2005, vol. 96, 673-686 [0217]
- SCHMIDTTHOMSEN Bioresource Technol., 1998, vol. 64, 139-151 [0218]
- PALONEN et al. Appl. Biochem. Biotechnol., 2004, vol. 117, 1-17 [0218]
- VARGA et al. Biotechnol. Bioeng., 2004, vol. 88, 567-574 [0218]
- MARTIN et al. J. Chem. Technol. Biotechnol., 2006, vol. 81, 1669-1677 [0218]
- GOLLAPALLI et al. Appl. Biochem. Biotechnol., 2002, vol. 98, 23-35 [0220]
- CHUNDAWAT et al. Biotechnol. Bioeng., 2007, vol. 96, 219-231 [0220]
- ALIZADEH et al. Appl. Biochem. Biotechnol., 2005, vol. 121, 1133-1141 [0220]
- TEYMOURI et al. Bioresource Technol., 2005, vol. 96, 2014-2018 [0220]
- PAN et al. Biotechnol. Bioeng., 2005, vol. 90, 473-481 [0221]
- PAN et al. Biotechnol. Bioeng., 2006, vol. 94, 851-861 [0221]
- KURABI et al. Appl. Biochem. Biotechnol., 2005, vol. 121, 219-230 [0221]
- SCHELL et al. Appl. Biochem. and Biotechnol., 2003, vol. 105, 10869-85 [0222]
- MOSIER et al. Bioresource Technology, 2005, vol. 96, 673-686 [0222]
- Pretreatment of biomassHSU, T.-A. Handbook on Bioethanol: Production and Utilization19960000179-212 [0228]
- GHOSH SINGH Physicochemical and biological treatments for enzymatic/microbial conversion of cellulosic biomassAdv.

- Appl. Microbiol., 1993, vol. 39, 295-333 [0228]
- Pretreating lignocellulosic biomass: a review **MCMILLAN, J. D.** Enzymatic Conversion of Biomass for Fuels Production American Chemical Society 1994 0000 [0228]
 - Ethanol production from renewable resources **GONG, C. S. CAO, N. J. DU, J. TSAO, G. T.** Advances in Biochemical Engineering/Biotechnology Springer-Verlag Berlin Heidelberg 1999 0000 vol. 65, 207-241 [0228] [0309]
 - **OLSSON HAHN-HAGERDAL** Fermentation of lignocellulosic hydrolysates for ethanol production Enz. Microb. Tech., 1996, vol. 18, 312-331 [0228]
 - **VALLANDER ERIKSSON** Production of ethanol from lignocellulosic materials: State of the art Adv. Biochem. Eng./Biotechnol., 1990, vol. 42, 63-95 [0228]
 - **PENTTILA et al.** Gene, 1986, vol. 45, 253-263 [0256]
 - **SALOHEIMO et al.** Gene, 1988, vol. 63, 11-22 [0256]
 - **OKADA et al.** Appl. Environ. Microbiol., 1988, vol. 64, 555-563 [0256]
 - **SALOHEIMO et al.** Molecular Microbiology, 1994, vol. 13, 219-228 [0256]
 - **OOI et al.** Nucleic Acids Research, 1990, vol. 18, 5884- [0256]
 - **SAKAMOTO et al.** Current Genetics, 1995, vol. 27, 435-439 [0256]
 - **SAARILAHTI et al.** Gene, 1990, vol. 90, 9-14 [0256]
 - **KAWAGUCHI et al.** Gene, 1996, vol. 173, 287-288 [0256]
 - **DAN et al.** J. Biol. Chem., 2000, vol. 275, 4973-4980 [0256]
 - More Gene Manipulations in Fungi Academic Press 1991 0000 [0256]
 - **BAILEY, J. E. OLLIS, D. F.** Biochemical Engineering Fundamentals McGraw-Hill Book Company 1986 0000 [0286]
 - **LIN et al.** Appl. Microbiol. Biotechnol., 2006, vol. 69, 627-642 [0292]
 - Cloning and improving the expression of Pichia stipitis xylose reductase gene in Saccharomyces cerevisiae **CHEN HO** Appl. Biochem. Biotechnol. 1993 0000 vol. 39-40, 135-147 [0301]
 - **HO et al.** Genetically engineered Saccharomyces yeast capable of effectively cofermenting glucose and xylose Appl. Environ. Microbiol., 1998, vol. 64, 1852-1859 [0301]
 - **KOTTER CIRIACY** Xylose fermentation by Saccharomyces cerevisiae Appl. Microbiol. Biotechnol., 1993, vol. 38, 776-783 [0301]
 - **WALFRIDSSON et al.** Xylose-metabolizing Saccharomyces cerevisiae strains overexpressing the TKL1 and TAL1 genes encoding the pentose phosphate pathway enzymes transketolase and transaldolase Appl. Environ. Microbiol., 1995, vol. 61, 4184-4190 [0301]
 - **KUYPER et al.** Minimal metabolic engineering of Saccharomyces cerevisiae for efficient anaerobic xylose fermentation: a proof of principle FEMS Yeast Research, 2004, vol. 4, 655-664 [0301]
 - **BEALL et al.** Parametric studies of ethanol production from xylose and other sugars by recombinant Escherichia coli Biotech. Bioeng., 1991, vol. 38, 296-303 [0301]
 - **INGRAM et al.** Metabolic engineering of bacteria for ethanol production Biotechnol. Bioeng., 1998, vol. 58, 204-214 [0301]
 - **ZHANG et al.** Metabolic engineering of a pentose metabolism pathway in ethanologenic Zymomonas mobilis Science, 1995, vol. 267, 240-243 [0301]
 - **DEANDA et al.** Development of an arabinose-fermenting Zymomonas mobilis strain by metabolic pathway engineering Appl. Environ. Microbiol., 1996, vol. 62, 4465-4470 [0301]
 - **ALFENORE et al.** Improving ethanol production and viability of Saccharomyces cerevisiae by a vitamin feeding strategy during fed-batch process Springer-Verlag 2002 0000 [0307]
 - **SILVEIRA, M. M. JONAS, R.** The biotechnological production of sorbitol Appl. Microbiol. Biotechnol., 2002, vol. 59, 400-408 [0309]
 - **NIGAM, P. SINGH, D.** Processes for fermentative production of xylitol - a sugar substitute Process Biochemistry, 1995, vol. 30, 2117-124 [0309]
 - **EZEJI, T. C. QURESHI, N. BLASCHEK, H. P.** Production of acetone, butanol and ethanol by Clostridium beijerinckii BA101 and in situ recovery by gas stripping World Journal of Microbiology and Biotechnology, 2003, vol. 19, 6595-603 [0309]
 - **RICHARD, A. MARGARITIS, A.** Empirical modeling of batch fermentation kinetics for poly(glutamic acid) production and other microbial biopolymers Biotechnology and Bioengineering, 2004, vol. 87, 4501-515 [0313]
 - **KATAOKA, N. A. MIYAK. KIRIYAMA** Studies on hydrogen production by continuous culture system of hydrogen-producing anaerobic bacteria Water Science and Technology, 1997, vol. 36, 6-741-47 [0314]
 - **GUNASEELAN V. N.** Biomass and Bioenergy, 1997, vol. 13, 1-283-114 [0314]
 - **CHEN, R. LEE, Y. Y.** Membrane-mediated extractive fermentation for lactic acid production from cellulosic biomass Appl. Biochem. Biotechnol., 1997, vol. 63-65, 435-448 [0317]
 - **ROSE et al.** Nucleic Acids Research, 1998, vol. 26, 1628-1635 [0333]
 - **POZZO et al.** J. Mol. Biol., 2010, vol. 397, 724-739 [0343]
 - **HÄGERDAL et al.** Biotechnology and Bioengineering, 1979, vol. 21, 345-355 [0346]

Patentkrav

1. Isoleret polypeptid med beta-glucosidaseaktivitet, som er valgt fra gruppen bestående af:
 - 5 (a) et polypeptid, som udviser mindst 95%, f.eks. mindst 96%, mindst 97%, mindst 98%, mindst 99% og 100% sekvensidentitet med det modne polypeptid ifølge SEQ ID NO: 2; og
 - (b) en variant af det modne polypeptid ifølge SEQ ID NO: 2, som omfatter en substitution, deletion og/eller insertion i en eller flere positioner, hvor varianten udviser mindst 85%, f.eks. mindst 86%, mindst 87%, mindst 88%, mindst 89%, mindst 90%, mindst 91%,
 - 10 mindst 92%, mindst 93%, mindst 94%, mindst 95%, mindst 96%, mindst 97%, mindst 98%, mindst 99% sekvensidentitet med det modne polypeptid ifølge SEQ ID NO: 2, og hvor varianten har forbedret termisk stabilitet i forhold til det modne polypeptid ifølge SEQ ID NO: 2.
2. Polypeptid ifølge krav 1, hvilket polypeptid omfatter eller består af SEQ ID NO: 2 eller
- 15 det modne polypeptid deraf.
3. Polypeptid ifølge krav 1 eller 2, hvilket polypeptid kodes for af polynukleotidet indeholdt i plasmid pUC19 D55EX, hvilket plasmid er indeholdt i *E. coli* NRRL B-50395.
- 20 4. Sammensætning, som omfatter polypeptidet ifølge et hvilket som helst af kravene 1-3.
5. Isoleret polynukleotid, som koder for polypeptidet ifølge et hvilket som helst af kravene 1-3.
- 25 6. Fremgangsmåde til fremstilling af polypeptidet ifølge et hvilket som helst af kravene 1-3, hvilken fremgangsmåde omfatter:
 - (a) dyrkning af en celle, som i sin vildtypeform fremstiller polypeptidet, under betingelser, der er befordrende for fremstilling af polypeptidet; og
 - (b) indvinding af polypeptidet.
- 30 7. Fremgangsmåde til fremstilling af et polypeptid med beta-glucosidaseaktivitet, hvilken fremgangsmåde omfatter:
 - (a) dyrkning af en værtscelle, som omfatter polynukleotidet ifølge krav 5, under betingelser, der er befordrende for fremstilling af polypeptidet; og
 - 35 (b) indvinding af polypeptidet.
8. Transgen plante, plantedel eller plantecelle, som er transformeret med et polynukleotid,

hvilket polynukleotid koder for polypeptidet ifølge et hvilket som helst af kravene 1-3.

9. Fremgangsmåde til fremstilling af polypeptidet med beta-glucosidaseaktivitet, hvilken fremgangsmåde omfatter:

- 5 (a) dyrkning af den transgene plante eller plantecelle ifølge krav 8 under betingelser, som er befordrende for fremstilling af polypeptidet; og
- (b) indvinding af polypeptidet.

10 10. Fremgangsmåde til fremstilling af en mutant af en stamcelle, hvilken fremgangsmåde omfatter inaktivering af et polynukleotid, der koder for polypeptidet ifølge et hvilket som helst af kravene 1-3, hvilket resulterer i, at mutanten fremstiller mindre af polypeptidet end stamcellen.

15 11. Dobbeltstrengt hæmmende RNA (dsRNA)-molekyle, som omfatter en undersekvens af polynukleotidet ifølge krav 5, hvor dsRNA'et eventuelt er et siRNA- eller miRNA-molekyle.

 12. Fremgangsmåde til hæmning af ekspressionen af et polypeptid med beta-glucosidaseaktivitet i en celle, hvilken fremgangsmåde omfatter administration til cellen eller ekspression i cellen af det dobbeltstrengede RNA (dsRNA)-molekyle ifølge krav 11.

20 13. Fremgangsmåde til nedbrydning eller omdannelse af et celluloseholdigt materiale, hvilken fremgangsmåde omfatter: behandling af det celluloseholdige materiale med en enzymsammensætning i nærværelse af polypeptidet med beta-glucosidaseaktivitet ifølge et hvilket som helst af kravene 1-3.

25 14. Fremgangsmåde ifølge krav 13, hvilken fremgangsmåde endvidere omfatter indvinding af det nedbrudte celluloseholdige materiale.

 15. Fremgangsmåde til fremstilling af et fermenteringsprodukt, hvilken fremgangsmåde omfatter:

30 (a) forsukring af et celluloseholdigt materiale med en enzymsammensætning i nærværelse af polypeptidet med beta-glucosidaseaktivitet ifølge et hvilket som helst af kravene 1-3;

 (b) fermentering af det forsukrede celluloseholdige materiale med en eller flere fermenterende mikroorganismer for at fremstille fermenteringsproduktet; og

35 (c) indvinding af fermenteringsproduktet fra fermenteringen.

 16. Fremgangsmåde til fermentering af et celluloseholdigt materiale, hvilken fremgangsmåde omfatter: fermentering af det celluloseholdige materiale med en eller flere fermenterende mikroorganismer, hvor det celluloseholdige materiale er forsukret med en

enzym sammensætning i nærværelse af et polypeptid med beta-glucosidaseaktivitet ifølge et hvilket som helst af kravene 1-3.

17. Fremgangsmåde ifølge krav 16, hvor fermenteringen af det celluloseholdige materiale
5 frembringer et fermenteringsprodukt.

18. Fremgangsmåde ifølge krav 17, hvilken fremgangsmåde endvidere omfatter indvinding af fermenteringsproduktet fra fermenteringen.

10

DRAWINGS

M R S S T T V L A F V S Y G S F G I A A A K P I O E
 1 ATGAGGAGCTCAACCAAGCTTCTGGCCTTCTGTATACGGGAGTTTGGGATTCGGGCACCGAAGCCCATACAGAG
 H K
 79 CACAAGGTATGAACGGTTTCTTTAAGCGAGAACGGAGATATAACGATTTCGAGAGTAACGGGCACTGACCATTTTGT
 P E L L N S F H G F Q I E E P Y S P P F Y P S
 157 TCTCTCAAAGCCCGAGCTTCTGAATACCTTTCACCGCTTTCAGATTGAAGAGCCGTATTCGGCCGCTTTTATCCCTC
 P W M N P R A E G W E D A Y Q K A L A F V S Q L T L
 235 GCCATGGATGAATCTCTCTGCTGAAGCTTCGGAGGATCGGTATCAGAAGCGCTTGCATTGTCTCTCAACTGACGTT
 L E K V N L T T G V G
 313 ACTCGAGAAGGTCAATCTGACGACGGGTCTTGGGTGAGTGATTGCCCTTTTCCCTCTATGATCTGTGCGACAGTGGGCT
 W E N G P C V G N T G S V P R L G F K G L
 391 GAACATGTTCAAAGTTGGGAAACGGACCATCTGTAGGAAATACGGATCGGTCCCTCGCCTTGGCTTCAAGGCGCTT
 C L Q D S P Q G V R F A D Y A S A F T S S Q M A A A
 469 TGCTTGAAGACTTCTCTCAAGGTGTGAGATTGCGGACTATGCTGCTTTCACGTCAGTCAGATGGCTGCTGCC
 T F D K T V L Y E R G V A M A Q E H K G K G T T V Q
 547 ACATTTGACAAGACGGTTCTTTATGAGCGTGGTGTGGCATGGCTCAAGAGCACAAAGGCAAGGGCATCACGGTGCAA
 L G P V A G P L G R A P E G G R N W E G F S P D P V
 625 CTGGGCCCGCTGCTGGCCTTTGGGCGCGCCCTGAAGGAGCGAACTGGGAAGGATTCTCGCCCGATCCCGTC
 L N G I A M A E T I K G M Q D T G V L A C A K H Y V
 703 CTGAAGCGCATTCGGATGGCGGAGACCATCAAGGGCATGCAAGATACCGGAGTCTTGGCCTGCGGAAACATTATGTT
 G N E Q E H F R Q A G E S A G H G Y N I A E S I S S
 781 GGAAGCAACAGAGCACTTCCGACAAGCAGCGAGTCTGCGGCGATGGATACACATTGCTGAATCGATCTCTGCG
 N I D D R T M H E M Y L W P P A D A V R A C V A S P
 859 AACATGACGATGCTACATGACATGAATCTTGTGGCTTTTGTGATGCGGTACGTCGTCGAGTGGCATCTTTT
 M C S Y N N S Y G C O N S H T L N K L L K S E
 937 ATGTGCTCCTACACAGCTCAACAACTCGTACGGATGCCAAACAGTCACACTCTAAACAACTCTCAAGAGCGAG
 L G F Q G F V M S D W Q A Q H S G V S S A L A G L D
 1015 TTGGCTTCCAAGGTTCTGGTGTGAGTACTGGCAAGCACAACTCTCGAGTATCGTGGCGCTGCCCGGCTCGAT
 M T M P G D T E F D T G L S F W G S N L T I A V I N
 1093 ATGACAATGCCAGACATCTGAATTCGACACCGGACTCAGCTTTTGGGGATCCAATCTCACTATTGCGGTTCTGAAC
 G T V P E W R I D D M A M R I M A A Y F K V G L T I
 1171 GGCACCGTGCCCGATCGAGGATTGACGATATGGCGATGCGGATCATGGCCGCTTATTTCAAAGTCGGTCTGACAATT
 E D Q P D V N F N S W T Y D T Y G Y K Y F Y S K E G
 1249 GAAGATCAGCTGACGTGAATTTCAACTCATGGACCTACGATACCTATGGTTACAAATACCTTTTACAGCAAGGAGGT
 Y E K V N W H V D V R A N H A O Q I R E T A A K G T
 1327 TACGAGAAGTGAATGGCATGTGCGCTGCAAAACACGCCCAACAGATTCTGAGACCGCCCAAGGGAAC
 V L L K N T N G A L P L K R P S F V A V V G E D A G
 1405 GTGCTTCTCAAACACCAATGGTGTCTTCCGCTGAAACGACCTTCTTCTGTCGCGGTGTTGTTGGCGAAGATGCGGGT
 P N P K G P N G C P D R G C D E G T L A M G W G S G
 1483 CCAATGCCAAGGTTCCGAATGGGTGCTCTGATCGTGGCTGCGATGAGGGAACCTGCCATGGATGGGATCTGGC
 S T Q P P F L V T P D S A I Q A K V L E Y G C R Y E
 1561 TCGACCCAGTTCCTTCTTCTTACTCCAGACTCTGCGATTCAAGCCAAGGCTTTGGAGTATGGTGGCCGATATGAG
 S I L D N Y D E S A I S A L V S Q P D A T C I V F G
 1639 AGTATTTGGACAACACGATCAAGTGCATCTCCGCACTGGTCTGCGAGCTGATGCTACTTGCATCGCTTTTGGC
 N S D S G E A F I T V D G N W G D R N N L T L W Q N
 1717 AACTCGGATTTGCGGAAGCTTTCATCACCGTCGACGGCAACIGGGGGATCGGAATAACCTCACTCTTTGGCAAAAT
 A D A V G I R N V S A L C S N T I V V L H T V G P V L
 1795 GCCGACGCGGTGATCCGCAATGTCAGTGTCTGTGTCAGCAACACCATTTGCTTCTTCACTGTCGGCCCTGTGCTG
 L T D Y Y E H P N I T A I V W A G L P G E E S G N A
 1873 TTGACCGATTACTATGAGCACCCCAACATCACAGCCATTGTCTGGGCGGCTTACCTGGCGAAGATCAGGGAATGCT
 L V D I L W G D V N P A G R T P F T W A K S R E D Y
 1951 CTGTGATATCTCTGGGGCGATGTCAACCCAGCCGTCGACGCCATTACATGCGGCAAGAGCGGGAAGACTAT
 G T D V M Y V P N N G E Q A P Q Q S F S E G I F L D
 2029 GGCACAGCGTGTATGTGTCGCAACAGCGGCAACAGCCAGCAATCTTCTCGAAGGCATCTTCTTGTGAC
 Y R H F D Q A G I D P I Y E F G H G L S Y T N F S Y
 2107 TACCGGCAATTTGACCAAGTCGGCATGTATCCATCTACGAGTTCGGCCACGGTCTCTCATATACCAATTTCTCATAC

FIG. 1A

S D L R I V T K H V R P Y R P T T G M T T Q A P I I
 2185 TCTGACCTCCGCATCGTCACCAACACGTTGGTCCCTATCGACCCACGACAGGCATGACCACGCAAGCACCTATCATC
 G P L P S P N L T E Y Q F P A T F K Y I P A F I Y P
 2263 GGACCCCTCCCGAGCCCAAACTCACCGAGTATCAATTCCCTGCTACTTTCAAATACATCCCGCCCTTTATCTATCCC
 Y L N S T E S L R A A S K D P H Y G S T A F I P P H
 2341 TACCTCAACAGCAGACAGAGTCCCTCCGCGCGCCTCAAAAGACCTTACTACGGCAGCACCAGCAITTCATCCCGCTCAT
 A L D S S P Q P L N P A G D P I A S G G N S M L Y Q
 2419 GCGCTCGACAGTCCCGACAGCCTCTCAACCCAGCGGTGACCCCATCGCCTCCGCGCGTAACAGTATGCTCTATCAG
 E L Y E I T A K I T N T G N L A G D E V V Q L Y V N
 2497 GAGCTCTACGAAATCACCGCCAAAATCACCAACACCGGCAATCTAGCCGGTGACGAGGTTGTCCAACGTACGTCAAC
 L G G D N P P R Q L R G F D R I H L L P G Q S K V F
 2575 CTCGGTGGCGACAATCCTCTCGACAGCTTCGAGGATTTGACCGAATCCACCTCTGCCCGGCCAGAGTAAGGTGTTT
 R A T L T R R D L S N W D V A A Q N W R I I R S A X
 2653 CGAGCGACGTTGACACGTCGGGACTTGAGTAATTGGGACGTTGGGCGCAGAATTGGCGCATCATCAGSTCCGCCAAG
 E V Y V G R S S R D L P L S A R L E S W *
 2731 GAGGTTTATGTCGACGCTCAAGTCAGATCTCCCTTGAGCGCGCGATTGGAGTCATGGTAA

FIG. 1B




FIG. 2