



Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2329181 C 2011/02/22

(11)(21) **2 329 181**

(12) **BREVET CANADIEN
CANADIAN PATENT**

(13) **C**

(86) Date de dépôt PCT/PCT Filing Date: 1999/04/05
(87) Date publication PCT/PCT Publication Date: 1999/10/28
(45) Date de délivrance/Issue Date: 2011/02/22
(85) Entrée phase nationale/National Entry: 2000/10/18
(86) N° demande PCT/PCT Application No.: US 1999/007475
(87) N° publication PCT/PCT Publication No.: 1999/054319
(30) Priorité/Priority: 1998/04/21 (US60/082,564)

(51) Cl.Int./Int.Cl. *C07D 417/06* (2006.01),
C07D 277/24 (2006.01), *C07D 277/26* (2006.01),
C07D 313/00 (2006.01), *C07D 471/02* (2006.01),
C07D 491/02 (2006.01), *C07D 491/04* (2006.01),
C07D 493/04 (2006.01), *C07D 495/04* (2006.01),
C07D 498/02 (2006.01), *C07D 513/04* (2006.01)

(72) Inventeurs/Inventors:
VITE, GREGORY D., US;
KIM, SOONG-HOON, US;
HOFLE, GERHARD, DE

(73) Propriétaire/Owner:
BRISTOL-MYERS SQUIBB COMPANY, US

(74) Agent: GOWLING LAFLEUR HENDERSON LLP

(54) Titre : DERIVES D'EPOTHILONE MODIFIES EN POSITION 12,13
(54) Title: 12,13-MODIFIED EPOTHILONE DERIVATIVES

(57) Abrégé/Abstract:

The present invention relates to 12,13-position modified epothilone derivatives, methods of preparation of the derivatives and intermediates therefor.



**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 277/24, 277/26, 313/00, 471/02, 491/02, 498/02	A1	(11) International Publication Number: WO 99/54319 (43) International Publication Date: 28 October 1999 (28.10.99)
(21) International Application Number: PCT/US99/07475 (22) International Filing Date: 5 April 1999 (05.04.99) (30) Priority Data: 60/082,564 21 April 1998 (21.04.98) US (71) Applicant: BRISTOL-MYERS SQUIBB COMPANY [US/US]; P.O. Box 4000, Princeton, NJ 08543-4000 (US). (72) Inventors: VITE, Gregory, D.; 28 Continental Lane, Titusville, NJ 08560 (US). KIM, Soong-Hoon, Kim; 13126 East Run Drive, Lawrenceville, NJ 08648 (US). HOFLE, Gerhard; Alter Weg 12a, D-38124 Braunschweig (DE). (74) Agents: HOFFMAN, Frank, P. et al.; Bristol-Myers Squibb Company, P.O. Box 4000, Princeton, NJ 08543-4000 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: 12,13-MODIFIED EPOTHILONE DERIVATIVES (57) Abstract The present invention relates to 12,13-position modified epothilone derivatives, methods of preparation of the derivatives and intermediates therefor.		

The present invention relates to compounds of the formula

Q is selected from the group consisting of

G is selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl, heterocyclo,

1

W is O or NR₁₆;

X is O, S, CHR₁₇ or H, R₁₈;

Z is O; S; H, R₁₉ or R₂₀, R₂₁;

Y is selected from the group consisting of O; H, H; H, OR₂₂; OR₂₃,
 5 OR₂₃; NOR₂₄; H, NOR₂₅; H, NR₂₆R₂₇; NHNR₂₈R₂₉; H, NHNR₃₀R₃₁;
 or CHR₃₂; where OR₂₃, OR₂₃ can be a cyclic ketal;

B₁ and B₂ are selected from the group consisting of H, OR₃₃,
 OCOR₃₄, OCONR₃₅R₃₆, NR₃₇R₃₈ or NR₃₉CONR₄₀R₄₁;

D is selected from the group consisting of NR₄₂R₄₃ or heterocyclo;

10 M is selected from the group consisting of S, C=O, S=O, SO₂, NR₄₄,
 or CR₄₅R₄₆;

J, E, U, and V are selected from carbon, oxygen, nitrogen or sulfur; or
 V may be absent;

R₁, R₂, R₃, and R₄, are selected from H, lower alkyl;


15 R₅, R₈, R₉, R₁₀ and R₁₁ are selected from the group consisting of
 H, alkyl, substituted alkyl, aryl, substituted aryl, cycloalkyl, heterocyclo or
 substituted heterocyclo;

R₆ and R₇ are selected from the group consisting of H, alkyl,
 substituted alkyl, halogen, nitro, cyano, OR₄₇, NR₄₈R₄₉, R₅₀C=O;

20 R₁₇, R₁₈, R₂₂ and R₂₃ are selected from the group consisting of H,
 alkyl, and substituted alkyl;

$R_{20}, R_{21}, R_{24}, R_{25}, R_{26}, R_{28}, R_{30}, R_{32}, R_{33}, R_{34}, R_{35}, R_{36}, R_{37},$
 $R_{39}, R_{40}, R_{41}, R_{42}, R_{47}, R_{48}, R_{50}, R_{51}, R_{52}, R_{53}, R_{54}, R_{56}, R_{57}, R_{58},$
 R_{59} and R_{61} are selected from the group of H, alkyl, substituted alkyl,
 aryl or substituted aryl;

5 $R_{12}, R_{13}, R_{16}, R_{27}, R_{29}, R_{31}, R_{38}, R_{43}, R_{44}, R_{49}, R_{55}$ and R_{60}
 are selected from the group consisting of H, alkyl, substituted alkyl,
 substituted aryl, cycloalkyl, heterocyclo, $R_{51}C=O$, $R_{52}OC=O$, $R_{53}SO_2$,
 hydroxy, O-alkyl or O-substituted alkyl; when X is O then R_{16} is not
 $R_{51}C=O$, $R_{52}OC=O$, and $R_{53}SO_2$; and wherein R_{44} is further amino and



10


R_{14} and R_{15} are selected from the group consisting of H, halogen,
 alkyl, or substituted alkyl;

R_{19} is selected from the group consisting of H, alkyl, substituted
 alkyl, O-alkyl, O-substituted alkyl, $NR_{54}R_{55}$, $R_{56}C=O$; when L is O, S, or
 15 NR_{13} then R_{19} is not O-alkyl, O-substituted alkyl, $NR_{54}R_{55}$.

R_{45} and R_{46} are selected from the group consisting of H, halogen,
 alkyl, substituted alkyl, aryl, substituted aryl, cycloalkyl, heterocyclo,
 $R_{56}C=O$, $R_{57}OC=O$, $R_{58}NHC=O$, hydroxy, O-alkyl or O-substituted alkyl,
 $NR_{59}R_{60}$;

and any salts, solvates or hydrates thereof, with the proviso that

when Q is



and


M is $CR_{45}R_{46}$ then W is only NR_{16} .

5

Background of the Invention

Epothilones are macrolide compounds which find utility in the pharmaceutical field. For example, Epothilones A and B having the

10 structures:



Epothilone A R = H

Epothilone B R = Me

15 have been found to exert microtubule-stabilizing effects similar to TAXOL and hence cytotoxic activity against rapidly proliferating cells, such as, tumor cells or other hyperproliferative cellular disease, see Angew. Chem. Int. Ed. Engl., 1996, 35, No. 13/14.

Detailed Description of the Invention

Listed below are definitions of various terms used to describe this
5 invention. These definitions apply to the terms as they are used
throughout this specification, unless otherwise limited in specific
instances, either individually or as part of a larger group.

The term "alkyl" refers to straight or branched chain unsubstituted
hydrocarbon groups of 1 to 20 carbon atoms, preferably 1 to 7 carbon
10 atoms. The expression "lower alkyl" refers to unsubstituted alkyl groups
of 1 to 4 carbon atoms.

The term "substituted alkyl" refers to an alkyl group substituted by,
for example, one to four substituents, such as, halo, phenyl, substituted
phenyl, heterocyclo, trifluoromethyl, trifluoromethoxy, hydroxy, alkoxy,
15 cycloalkoxy, heterocylooxy, oxo, alkanoyl, aryloxy, alkanoyloxy, amino,
alkylamino, arylamino, aralkylamino, cycloalkylamino, heterocycloamino,
disubstituted amines in which the 2 amino substituents are selected from
alkyl, aryl or aralkyl, alkanoylamino, aroylamino, aralkanoylamino,
substituted alkanoylamino, substituted arylamino, substituted
20 aralkanoylamino, thiol, alkylthio, arylthio, aralkylthio, cycloalkylthio,
heterocyclothio, alkylthiono, arylthiono, aralkylthiono, alkylsulfonyl,
arylsulfonyl, aralkylsulfonyl, sulfonamido (e.g. SO₂NH₂), substituted

sulfonamido, nitro, cyano, carboxy, carbamyl (e.g. CONH_2), substituted carbamyl (e.g. CONH alkyl, CONH aryl, CONH aralkyl or cases where there are two substituents on the nitrogen selected from alkyl, aryl or aralkyl), alkoxycarbonyl, aryl, substituted aryl, guanidino and heterocyclos, such as, indolyl, imidazolyl, furyl, thienyl, thiazolyl, pyrrolidyl, pyridyl, pyrimidyl and the like. Where noted above where the substituent is further substituted it will be with halogen, alkyl, alkoxy, aryl or aralkyl.

The term "halogen" or "halo" refers to fluorine, chlorine, bromine and iodine.

The term "aryl" refers to monocyclic or bicyclic aromatic hydrocarbon groups having 6 to 12 carbon atoms in the ring portion, such as phenyl, naphthyl, biphenyl and diphenyl groups, each of which may be substituted.

The term "aralkyl" refers to an aryl group bonded directly through an alkyl group, such as benzyl.

The term "substituted aryl" refers to an aryl group substituted by, for example, one to four substituents such as alkyl; substituted alkyl, halo, trifluoromethoxy, trifluoromethyl, hydroxy, alkoxy, cycloalkyloxy, heterocyclooxy, alkanoyl, alkanoyloxy, amino, alkylamino, aralkylamino, cycloalkylamino, heterocycloamino, dialkylamino, alkanoylamino, thiol, alkylthio, cycloalkylthio, heterocyclothio, ureido, nitro, cyano, carboxy, carboxyalkyl, carbamyl, alkoxycarbonyl, alkylthiono, arylthiono, alkysulfonyl, sulfonamido, aryloxy and the like. The substituent may be

further substituted by halo, hydroxy, alkyl, alkoxy, aryl, substituted aryl, substituted alkyl or aralkyl.

The term "cycloalkyl" refers to optionally substituted, saturated cyclic hydrocarbon ring systems, preferably containing 1 to 3 rings and 3 to 7 carbons per ring which may be further fused with an unsaturated C₃-C₇ carbocyclic ring. Exemplary groups include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, cyclodecyl, cyclododecyl, and adamantyl. Exemplary substituents include one or more alkyl groups as described above, or one or more groups described above as alkyl substituents.

The terms "heterocycle", "heterocyclic" and "heterocyclo" refer to an optionally substituted, fully saturated or unsaturated, aromatic or nonaromatic cyclic group, for example, which is a 4 to 7 membered monocyclic, 7 to 11 membered bicyclic, or 10 to 15 membered tricyclic ring system, which has at least one heteroatom in at least one carbon atom-containing ring. Each ring of the heterocyclic group containing a heteroatom may have 1, 2 or 3 heteroatoms selected from nitrogen atoms, oxygen atoms and sulfur atoms, where the nitrogen and sulfur heteroatoms may also optionally be oxidized and the nitrogen heteroatoms may also optionally be quaternized. The heterocyclic group may be attached at any heteroatom or carbon atom.

Exemplary monocyclic heterocyclic groups include pyrrolidinyl, pyrrolyl, indolyl, pyrazolyl, oxetanyl, pyrazolinyl, imidazolyl, imidazolinyl, imidazolidinyl, oxazolyl, oxazolidinyl, isoxazolinyl, isoxazolyl, thiazolyl, thiadiazolyl, thiazolidinyl, isothiazolyl, isothiazolidinyl, furyl, 5 tetrahydrofuryl, thienyl, oxadiazolyl, piperidinyl, piperazinyl, 2-oxopiperazinyl, 2-oxopiperidinyl, 2-oxopyrrolidinyl, 2-oxazepinyl, azepinyl, 4-piperidonyl, pyridyl, N-oxo-pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, tetrahydropyranyl, tetrahydrothiopyranyl, tetrahydrothiopyranyl sulfone, morpholinyl, thiomorpholinyl, thiomorpholinyl sulfoxide, thiomorpholinyl 10 sulfone, 1,3-dioxolane and tetrahydro-1, 1-dioxothienyl, dioxanyl, isothiazolidinyl, thietanyl, thiiranyl, triazinyl, and triazolyl, and the like.

Exemplary bicyclic heterocyclic groups include benzothiazolyl, benzoxazolyl, benzothienyl, quinuclidinyl, quinolinyl, quinolinyl-N-oxide, tetrahydroisoquinolinyl, isoquinolinyl, benzimidazolyl, benzopyranyl, 15 indolizinyl, benzofuryl, chromonyl, coumarinyl, cinnolinyl, quinoxalinyl, indazolyl, pyrrolopyridyl, furopyridinyl (such as furo[2,3-c]pyridinyl, furo[3,1-b]pyridinyl] or furo[2,3-b]pyridinyl), dihydroisoindolyl, dihydroquinazolinyl (such as 3,4-dihydro-4-oxo-quinazolinyl), benzisothiazolyl, benzisoxazolyl, benzodiazinyl, benzofurazanyl, 20 benzothiopyranyl, benzotriazolyl, benzpyrazolyl, dihydrobenzofuryl, dihydrobenzothienyl, dihydrobenzothiopyranyl, dihydrobenzothiopyranyl sulfone, dihydrobenzopyranyl, indolinyl, isochromanyl, isoindolinyl, naphthyridinyl, phthalazinyl, piperonyl, purinyl, pyridopyridyl,

quinazoliny, tetrahydroquinoliny, thienofury, thienopyridy, thienothieny, and the like.

Exemplary substituents include one or more alkyl groups as described above or one or more groups described above as alkyl substituents. Also included are smaller heterocyclos, such as, epoxides and aziridines.

The term "heteroatoms" shall include oxygen, sulfur and nitrogen.

The compounds of formula I and II may form salts with alkali metals such as sodium, potassium and lithium, with alkaline earth metals such as calcium and magnesium, with organic bases such as dicyclohexylamine, tributylamine, pyridine and amino acids such as arginine, lysine and the like. Such salts can be obtained, for example, by exchanging the carboxylic acid protons, if they contain a carboxylic acid, in compounds of formula I and II with the desired ion in a medium in which the salt precipitates or in an aqueous medium followed by evaporation. Other salts can be formed as known to those skilled in the art.

The compounds for formula I and II form salts with a variety of organic and inorganic acids. Such salts include those formed with hydrogen chloride, hydrogen bromide, methanesulfonic acid, hydroxyethanesulfonic acid, sulfuric acid, acetic acid, trifluoroacetic acid, maleic acid, benzenesulfonic acid, toluenesulfonic acid and various others (e.g., nitrates, phosphates, borates, tartrates, citrates, succinates, benzoates, ascorbates, salicylates and the like). Such salts are formed by

reacting a compound of formula I and II in an equivalent amount of the acid in a medium in which the salt precipitates or in an aqueous medium followed by evaporation.

In addition, zwitterions ("inner salts") are formed.

- 5 Compounds of the formula I and II may also have prodrug forms. Any compound that will be converted in vivo to provide the bioactive agent (i.e., the compound for formula I and II) is a prodrug within the scope and spirit of the invention.

For example compounds of the formula I and II may form a
10 carboxylate ester moiety. The carboxylate esters are conveniently formed by esterifying any of the carboxylic acid functionalities found on the disclosed ring structure(s).

Various forms of prodrugs are well known in the art. For examples of such prodrug derivatives, see:

- 15 a) Design of Prodrugs, edited by H. Bundgaard, (Elsevier, 1985) and Methods in Enzymology, Vol. 42, p. 309-396, edited by K. Widder, et al. (Acamedic Press, 1985);
- b) A Textbook of Drug Design and Development, edited by Krosgaard-Larsen and H. Bundgaard, Chapter 5, "Design and Application of
20 Prodrugs." by H. Bundgaard, p. 113-191 (1991);
- c) H. Bundgaard, Advanced Drug Delivery Reviews, 8, 1-38 (1992);
- d) H. Bundgaard, et al., Journal of Pharmaceutical Sciences, 77, 285 (1988); and

e) N. Kakeya, et al., Chem Phar Bull, 32, 692 (1984).

It should further be understood that solvates (e.g., hydrates) of the compounds of formula I and II are also within the scope of the present
5 invention. Methods of solvation are generally known in the art.

Use and Utility

The compounds of formula I and II are microtubule-stabilizing
10 agents. They are thus useful in the treatment of a variety of cancers, including (but not limited to) the following;

- carcinoma, including that of the bladder, breast, colon, kidney, liver, lung, ovary, pancreas, stomach, cervix, thyroid and skin; including squamous cell carcinoma;
- 15 - hematopoietic tumors of lymphoid lineage, including leukemia, acute lymphocytic leukemia, acute lymphoblastic leukemia, B-cell lymphoma, T-cell lymphoma, Hodgkins lymphoma, non-Hodgkins lymphoma, hairy cell lymphoma and Burketts lymphoma;
- hematopoietic tumors of myeloid lineage, including acute and
20 chronic myelogenous leukemias and promyelocytic leukemia;
- tumors of mesenchymal origin, including fibrosarcoma and rhabdomyosarcoma;

- other tumors, including melanoma, seminoma, teratocarcinoma, neuroblastoma and glioma;
 - tumors of the central and peripheral nervous system, including astrocytoma, neuroblastoma, glioma, and schwannomas;
 - 5 - tumors of mesenchymal origin, including fibrosarcoma, rhabdomyosarcoma, and osteosarcoma; and
 - other tumors, including melanoma, xenoderma pigmentosum, keratoactanthoma, seminoma, thyroid follicular cancer and teratocarcinoma.
- 10 Compounds of formula I and II may also inhibit tumor angiogenesis, thereby affecting the growth of tumors. Such anti-angiogenesis properties of the compounds of formula I and II may also be useful in the treatment of certain forms of blindness related to retinal vascularization, arthritis, especially inflammatory arthritis, multiple sclerosis, retinosis and
- 15 psoriasis.
- Compounds of formula I and II may induce or inhibit apoptosis, a physiological cell death process critical for normal development and homeostasis. Alterations of apoptotic pathways contribute to the pathogenesis of a variety of human diseases. Compounds of formula I and
- 20 II, as modulators of apoptosis, will be useful in the treatment of a variety of human diseases with aberrations in apoptosis including cancer (particularly, but not limited to follicular lymphomas, carcinomas with p53 mutations, hormone dependent tumors of the breast, prostate and

ovary, and precancerous lesions such as familial adenomatous polyposis), viral infections (including but not limited to herpesvirus, poxvirus, Epstein-Barr virus, Sindbis virus and adenovirus), autoimmune diseases (including but not limited to systemic lupus erythematosus, immune

5 mediated glomerulonephritis, rheumatoid arthritis, psoriasis, inflammatory bowel diseases and autoimmune diabetes mellitus), neurodegenerative disorders (including but not limited to Alzheimer's disease, AIDS-related dementia, Parkinson's disease, amyotrophic lateral sclerosis, retinitis pigmentosa, spinal muscular atrophy and cerebellar

10 degeneration), AIDS, myelodysplastic syndromes, aplastic anemia, ischemic injury associated myocardial infarctions, stroke and reperfusion injury, arrhythmia, atherosclerosis, toxin-induced or alcohol induced liver diseases, hematological diseases (including but not limited to chronic anemia and aplastic anemia), degenerative diseases of the

15 musculoskeletal system (including but not limited to osteoporosis and arthritis), aspirin-sensitive rhinosinusitis, cystic fibrosis, multiple sclerosis, kidney diseases, and cancer pain.

The compounds of this invention are also useful in combination with known anti-cancer and cytotoxic agents and treatments, including

20 radiation. If formulated as a fixed dose, such combination products employ the compounds of this invention within the dosage range described below and the other pharmaceutically active agent within its approved dosage range. Compounds of formula I and II can be used sequentially with known

anticancer or cytotoxic agents and treatment, including radiation when a combination formulation is inappropriate. Especially useful are cytotoxic drug combinations wherein the second drug chosen acts in a different phase of the cell cycle, e.g. S phase, than the present compounds of formula
5 I and II which exert their effects at the G₂-M phase.

The present compounds may exist as multiple optical, geometric, and stereoisomers. Included within the present invention are all such isomers and mixtures thereof in the racemic form.


The compounds of this invention can be formulated with a
10 pharmaceutical vehicle or diluent for oral, intravenous or subcutaneous administration. The pharmaceutical composition can be formulated in a classical manner using solid or liquid vehicles, diluents and additives appropriate to the desired mode of administration. Orally, the compounds can be administered in the form of tablets, capsules, granules, powders
15 and the like. The compounds are administered in a dosage range of about 0.05 to 200 mg/kg/day, preferably less than 100 mg/kg/day, in a single dose or in 2 to 4 divided doses.

20

Methods of Preparation

Compounds of formula I and II are prepared by the following schemes.

Scheme 1



5


Compounds of formula I where X is O, W is O, and Q is an aziridine group (i.e., M is NR₄₄) can be prepared as outlined in Scheme 1.

Compounds of formula 1A can be obtained from a fermentation process (see Angew. Chem. Int. Ed. Engl., 1996, 35, No. 13/14). A compound of

formula 1B, where P_1 is an oxygen protecting group such as triethylsilyl, can be prepared from a compound of formula 1A by known methods (See for example: Corey, E. J.; Venkateswarlu, A., J. Am. Chem. Soc., (1972) 94, 6190). Compounds of formula 1C and 1D are prepared by treatment with
5 an azide, such as sodium azide, in polar solvents such as DMF. A compound of formula 1E can be prepared from compounds of formulas 1C and 1D by the Staudinger reaction (See for example: Ittah, Y., et al., J. Org. Chem., (1978) 43, 4271). A compound of formula 1F where R_{44} is not H can be prepared from a compound of formula 1E using methods known in
10 the art. Deprotection of a compound of formula 1F using, for example when P_1 is a triethylsilyl group, hydrogen fluoride in acetonitrile or tetra-n-butylammonium fluoride in THF provides a compound of formula I (1G) where X is O, W is O, Q is an aziridine group (M is NR_{44}), and R_1 , R_2 , R_3 , R_4 and R_5 are defined as described above.

15


20

Scheme 2

- Alternatively, a compound of formula 1E can be prepared as shown
- 5 in Scheme 2. A compound of formula 2A, where P₁ is an oxygen protecting group such as triethylsilyl, can be prepared from a compound of formula 1B by reaction of tungsten (VI) chloride and n-butyllithium (See for example: Sharpless, K. B., et al., J. Am. Chem. Soc., (1972) 94, 6538) in THF. A compound of formula 2B can be prepared from a compound of formula 2A
- 10 by addition of a N-toluenesulfonamido group according to the method of Evans (i.e., Evans, D. A., et al., J. Org. Chem., (1991) 56, 6744). Deprotection of a compound of formula 2B using samarium iodide in THF/1,3-dimethyl-3,4,5,6-tetrahydro-2(1H)-pyrimidinone provides a compound of formula 1E (See for example: Vedejs, E., et al., J. Org. Chem.,
- 15 (1994) 59, 1602). Furthermore, a compound of formula 1B can be prepared

from a compound of formula 2A by oxidation (See for example: Balog, A., et al., Angew. Chem. Int. Ed. Engl., (1996) 35, 2801).

5 Scheme 3




Compounds of formula I where X is O, W is O, and Q is a thiirane (i.e., M is S) can be prepared as outlined in Scheme 3. A compound of formula 4A can be prepared from a compound of formula 1B using

10 potassium thiocyanate in alcoholic solution (See for example: Culvenor, C. C. J., et al., J. Chem. Soc., (1946) 1050). Deprotection of a compound of formula 4A using, for example when P₁ is a triethylsilyl group, hydrogen fluoride in acetonitrile or tetra-n-butylammonium fluoride in THF provides a compound of formula I (4B) where X is O, W is O, Q is a thiirane

15 group (i.e., M is S), and R₁, R₂, R₃, R₄ and R₅ are defined as described

above. Mild oxidation of a compound of formula 4A using stoichiometric or excess 3-chloroperoxybenzoic acid provides compounds of formula I where M is a S=O or SO₂ group, respectively.

5 Scheme 4




Compounds of formula I where X is O, W is O, and Q is an oxetane can be prepared as outlined in Scheme 4. A compound of formula 5A can be prepared from a compound of formula 1B, where R₅ is a methyl group, by using lithium 2,2,6,6-tetramethylpiperidide and diethylaluminum chloride in benzene (See for example: Paquette, L. A., et al., J. Org. Chem., (1990) 55, 1589). A compound of formula 5B can be prepared from a

compound of formula 5A by hydroboration and oxidation (See for example: Uzarewicz, A., et al., Rocz. Chem., (1977) 51, 723). A compound of formula 5C can be prepared from a compound of formula 5B by treatment with p-toluenesulfonylchloride (TsCl) in pyridine. A compound of formula 5D can
5 be prepared from a compound of formula 5C by an intramolecular Williamson reaction according to the method of Moulines (i.e., Moulines, B. J., Leclercq, D., Picard, P., Synthesis, (1981) 550). Deprotection of a compound of formula 5D using, for example when P₁ is a triethylsilyl group, hydrogen fluoride in acetonitrile or tetra-n-butylammonium fluoride
10 in THF provides a compound of formula I (5E) where X is O, W is O, Q is an oxetane group, and R₁, R₂, R₃, R₄ and R₅ are defined as described above.

15

20

Scheme 5

- 5 Alternatively, compounds of formula I where Q is an oxetane can be prepared as outlined in Scheme 5. Compounds of formula 6B and 6C can be prepared from a compound of formula 2A by Paterno-Buchi reaction of a carbonyl compound of formula 6A (See for example: Arnold, D. R., et al., Org. Photochem. Synth. (1971) 1, 51). Deprotection of compounds of
- 10 formula 6B and 6C using, for example when P₁ is a triethylsilyl group,


hydrogen fluoride in acetonitrile or tetra-n-butylammonium fluoride in THF provides a compound of formula I (6D and 6E) where X is O, W is O, Q is an oxetane group, and R₁, R₂, R₃, R₄ and R₅ are defined as described above.

5

10

15

20

Scheme 6

Compounds of formula I where X is O, W is O, and Q is a cyclobutane (i.e., L is CR₁₄R₁₅) can be prepared as outlined in Scheme 6. Compounds of formula 7B and 7C can be prepared from a compound of formula 2A by [2+2] cycloaddition of a substituted ketene compound of formula 7A (See for example: Krapcho, A. P., J. Org. Chem., (1966) 31, 2030). Deprotection of compounds of formula 7B and 7C using, for example when P₁ is a triethylsilyl group, hydrogen fluoride in acetonitrile or tetra-n-butylammonium fluoride in THF provides a compound of formula I (7D and


7E) where X is O, W is O, Q is an cyclobutane group, and R₁, R₂, R₃, R₄ and R₅ are defined as described above.

5

10


15

20

Scheme 7

Compounds of formula I where X is O, W is O, and Q is a 5-membered ring heterocycle can be prepared as outlined in Scheme 7.

Compounds of formula 8A and 8B can be prepared from a compound of formula 1B, where R₅ is a hydrogen, by reaction of magnesium bromide diethyl etherate in dichloromethane. Compounds of formula 8C and 8D can be prepared from compounds of formula 8A and 8B by pyridinium chlorochromate oxidation in dichloromethane (See for example: White, J. D., et al., J. Org. Chem., (1995) 12, 3600). Compounds of formula 8F and 8G can be prepared from compounds of formula 8C and 8D by addition of a substituted thioamide of formula 8E (See for example: Cauvin, P. Compt. Rend., (1973) 276C, 1453). Deprotection of compounds of formula 8F and 8G using, for example when P₁ is a triethylsilyl group, hydrogen fluoride in acetonitrile or tetra-n-butylammonium fluoride in THF provides a compound of formula I (8H and 8I) where X is O, W is O, Q is a 5-membered heterocycle, and R₁, R₂, R₃, R₄ and R₅ are defined as described above.

Scheme 8


Compounds of formula I where X is O, W is O, and Q is a 6-membered ring heterocycle can be prepared as outlined in Scheme 8.

Compounds of formula 9B and 9C can be prepared from compounds of formula 8C and 8D by reaction of diethyl zinc in THF with a Michael acceptor of formula 9A according to the method of Hansen (i.e., Hansen, M.

M., et al., Organometallics, (1987) 6, 2069). Compounds of formula 9D and 9E can be prepared from compounds of formula 9B and 9C by addition of hydroxyl amine in ethanol (See for example: Chemistry of Heterocyclic Compounds, Wiley: New York, 1974; Vol. 14, Parts 1-5). Deprotection of

5 compounds of formula 9D and 9E using, for example when P₁ is a triethylsilyl group, hydrogen fluoride in acetonitrile or tetra-n-butylammonium fluoride in THF provides a compound of formula I (9F and 9G) where X is O, W is O, Q is a 6-membered heterocycle, and R₁, R₂, R₃, R₄ and R₅ are defined as described above.

10 Scheme 9




A compound of formula 10G that is used for the preparation of compounds of formula I where X is O, W is O, and G is as defined above can

be prepared as outlined in Scheme 9. Deprotection of a compound of formula 2A using, for example when P₁ is a triethylsilyl group, hydrogen fluoride in acetonitrile or tetra-n-butylammonium fluoride in THF provides a compound of formula 10A. A compound of formula 10B can be prepared from a compound of formula 10A by pig-liver esterase mediated hydrolysis (See for example: Ohno, M., et al., Tetrahedron, (1984) 40, 145). A compound of formula 10C can be prepared from a compound of formula 10B by reaction of (trimethylsilyl)diazomethane in methanol and toluene (See for example: Hashimoto, N., et al., Chem. Pharm. Bull., (1981) 1397).

10 A compound of formula 10D can be prepared from a compound of formula 10C by addition of t-butyldimethylsilyltriflate and 2,6-lutidine in dichloromethane (See for example: Askin, D., et al., J. Org. Chem., (1987) 52, 622). A compound of formula 10E can be prepared from a compound of formula 10D by reaction of AD-mix- α according to the method of Sharpless (i.e., Sharpless, K. B., et al., J. Org. Chem., (1992) 57, 2768). Compounds of formula 10F and 10G can be prepared from a compound of formula 10E by oxidative cleavage using lead tetraacetate in ethyl acetate (See for example: C. A. Bunton in K. B. Wiberg, ed., Oxidation in Organic Chemistry. Part A, Academic Press, New York, 1965, p. 367).

15 Alternatively, a compound of formula 10G can be prepared from a compound of formula 10D by ozonolysis.

Scheme 10


Compounds of formula I where X is O, W is O, and G is a 1,2-
 5 disubstituted olefin can be prepared as outlined in Scheme 10. A
 compound of formula 11C can be prepared from compound of formula 11A
 and 11B using standard Wittig olefination known in the art (See for

example: Meng, D., et al., J. Org. Chem., (1996) 61, 7999). A compound of formula 11D can be prepared from a compound of formula 11C by addition of an allyl metal reagent such as allylmagnesium bromide (Hoffmann, R. W. Angew. Chem. Int. Ed., Engl., (1982) 21, 555). A compound of formula 11E, where P₂ is a triethylsilyl group, can be prepared from a compound of formula 11D by reaction of triethylsilyl chloride in DMF according to the method of Corey (See for example: Corey, E. J.; Venkateswarlu, A., J. Am. Chem. Soc., (1972) 94, 6190). A compound of formula 11F can be prepared from a compound of formula 11E by reaction of AD-mix- α according to the method of Sharpless (i.e., Sharpless, K. B., et al., J. Org. Chem., (1992) 57, 2768). A compound of formula 11G can be prepared from a compound of formula 11F by oxidative cleavage using lead tetraacetate in ethyl acetate (See for example: C. A. Bunton in K. B. Wiberg, ed., Oxidation in Organic Chemistry, Part A, Academic Press, New York, 1965, p. 367). A compound of formula 11H can be prepared from a compound of formula 11I by using a reducing agent such as sodium borohydride in methanol. A compound of formula 11I can be prepared from a compound with a formula 11H by reaction of iodine, imidazole and triphenylphosphine in toluene. A compound of formula 11J can be prepared from compound of formula 11I by reaction of triphenylphosphine in refluxing acetonitrile. A compound of formula 11K can be prepared from compounds of formula 11J and 10G using standard Wittig olefination (See for example: Bestmann, K. H., et

al., Chem. Ber. (1979) 112, 1923). A compound of formula 11L can be prepared from a compound of formula 11K by selective deprotection using AcOH in THF. A compound of formula 11M can be prepared from a compound of formula 11L by reaction of LiOH in t-butanol and water. A
5 compound of formula 11P can be prepared from a compound of formula 11M using typical macrolactonization coupling reagents such as 2,4,6-trichlorobenzoyl chloride, triethylamine, and 4-dimethylaminopyridine (See for example: Inanaga, J., et al., Bull. Chem. Soc. Jpn., (1979) 52, 1989). A compound of formula 11Q can be prepared from a compound of
10 formula 11P by reaction of 1,1,1-trifluoroacetone according to the method of Yang (Yang, D., et al., J. Org. Chem., (1995) 60, 3887). A compound of formula I (11R) where X is O, W is O, G is a 1,2-disubstituted olefin, and R₁, R₂, R₃, R₄ and R₅ are defined as described above can be prepared from compounds of formula 11P and 11Q as described in Schemes 1-9.

15

20


Scheme 11

Compounds of formula I where X is O, W is O, and G is a N, N-

- 5 substituted amine as defined above can be prepared as outlined in Scheme
11. A compound of formula 12A can be prepared from a compound of
- formula 11R by addition of oxygen protecting groups (P_1), such as a t-
- butyldimethylsilyl, with methods known in the art, followed by ozonolysis
- in dichloromethane. Reductive amination of compound of formula 12A
- 10 with a compound of formula 12B using standard reagents such as sodium
- triacetoxyborohydride in acetic acid/dichloromethane provides a compound
- of formula 12C. Deprotection of a compound of formula 12C using, for
- example when P_1 is a t-butyldimethylsilyl group, hydrogen fluoride in
- acetonitrile or tetra-n-butylammonium fluoride in THF provides a
- 15 compound of formula I (12D) where X is O, W is O, G is N, N-disubstituted

amine (i.e., D is $\text{NR}_{42}\text{R}_{43}$), and R_1 , R_2 , R_3 , R_4 and R_5 are defined as described above.

Scheme 12




- 5 Compounds of formula I where X is O, W is O, and G is a N, N-substituted amide as defined above can be prepared as outlined in Scheme 12. A compound of formula 13A can be prepared from a compound of formula I where G is 1-methyl-2-(2-methyl-4-thiazolyl)ethenyl by protection with triethylsilyl chloride (P_1 is triethylsilyl) in DMF using
- 10 methods known in the art followed by ozonolysis in dichloromethane. A


compound of formula 13B can be prepared from a compound of formula 13C by reaction of sodium bis(trimethylsilyl)amide in THF followed by addition of chlorotrimethylsilane to quench the intermediate lithium enolate (See for example: House, H. O., et al., J. Org. Chem., (1971) 36, 2361). A compound of formula 13C can be prepared from a compound of formula 13B by ozonolysis. A compound of formula 13E can be prepared from compounds of formula 13C and 13D by standard amide bond coupling agents (i.e., DCC, BOP, EDC/HOBT, PyBrOP). Deprotection of a compound of formula 13E using, for example when P₁ is a triethylsilyl group, hydrogen fluoride in acetonitrile or tetra-n-butylammonium fluoride in THF provides a compound of formula I (13F) where X is O, W is O, G is N, N-disubstituted amide, and R₁, R₂, R₃, R₄ and R₅ are defined as described above.

15


20

Scheme 13

- As shown in Scheme 13, compounds of formula I where W is NR₁₆ can be prepared from compounds of formula I where W is O. A compound of formula 14A can be prepared from a compound of formula 11R by reaction of sodium azide and tetrakis(triphenylphosphine)palladium in THF/water. Reduction of a compound of formula 14A using Adam's catalyst (PtO₂) or a trialkylphosphine in ethanol provides a compound of formula 14B. Intramolecular cyclization of compound of formula 14B using diphenylphosphoryl azide and sodium bicarbonate or 1-hydroxybenzotriazole and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide in DMF provides a compound of formula I (14C), where W is NR₁₆ (R₁₆ is H).

Scheme 14


Compounds of formula II can be prepared from a compound of formula I, where B1 and B2 are hydroxyl groups, as shown in Scheme 14. A compound of formula 15A can be prepared from compounds of formula I by addition of formyl groups using standard conditions such as formic acid, triethylamine, acetic anhydride in dichloromethane. Elimination of a compound of formula 15A using 1,8-diazabicyclo[5.4.0]undec-7-ene in dichloromethane provides a compound of formula 15B. Deprotection of a compound of formula 15B using ammonia in methanol provides a compound of formula II (15C).

Scheme 15

As shown in Scheme 15, compounds of formula I where X is S or dihydrogen can be prepared from compounds of formula I where X is O. A compound of formula 16A, where X is S, can be prepared from a compound of formula I where X is O by reaction of triethylsilyl chloride(P₁) with
5 methods known in the art, followed by the reaction of Lawesson's reagent in toluene (See for example: Nicolaou, K. C., et al., J. Amer. Chem. Soc., (1990) 112. 6263). Reduction of a compound of formula I (16A) using triphenyltin hydride and 2,2'-azobisisobutyronitrile in benzene, followed by deprotection using standard methods such as acetic acid in THF (See for
10 example: Nicolaou, K. C., et al., Chem. Comm., (1995) 1583) provides a compound of formula I (16B), where X is dihydrogen.

15

20


Scheme 16

Alternatively, as shown in Scheme 16, compounds of formula I where X is H, R₁₈ can be prepared from compounds of formula I where X is O. A compound of formula 17A can be prepared from a compound of formula I by reaction of triethylsilyl chloride in DMF using standard methods known in the art. A compound of formula 17B can be prepared from a compound of formula 17A by reaction of diisobutylaluminum hydride followed by acetic anhydride according to the method of Dahanukar (i.e., Dahanukar, V. H., et al., J. Org. Chem. (1996) 61, 8317). Reaction of a compound of formula 17B with dialkylzinc, (R₁₈)₂Zn, in dichloromethane according to the method of Rychnovsky (Rychnovsky, S. D., et. al., J. Org. Chem. (1997) 62, 6460) provides a compound of formula 17C. Deprotection of a compound of formula 17C using, for example when P₁ is a triethylsilyl group, hydrogen fluoride in acetonitrile or tetra-n-butylammonium fluoride in THF

provides a compound of formula I (17D) where X is H, R₁₈ and R₁, R₂, R₃.

R₄ and R₅ are defined as described above.


Scheme 17




5

The hydroxyl groups of a compound of formula I, such as 1A (where R₁₋₄ are methyl and R₆ is 2-methyl-4-thiazolyl), can be optionally protected as, for example, triethylsilyl ethers using methods known in the art. A compound of formula 18B where X is halogen, can be prepared from a

compound for formula 18A by treatment with certain metal halide salts such as magnesium bromide. A compound of formula 18C can be prepared from a compound of formula 18B by treatment with a metal azide salt such as lithium azide. A compound of formula 18D can be prepared from a compound of formula 18C by Mitsunobu reaction using, for example, triphenylphosphine and a carboxylic acid such as 4-nitrobenzoic acid. A compound of formula 18E can be prepared from a compound of formula 18D by hydrolysis or ammoniolysis of the ester group using, for example, a solution of ammonia in methanol. Optionally, a compound of formula 18E where P1 is an oxygen protecting group such as triethylsilyl can be deprotected using trifluoroacetic acid in dichloromethane, or other methods known in the art. Reduction of the azido group and subsequent cyclization of a compound of formula 18E with a reducing agent such as a triaryl- or trialkylphosphine provides a compound of formula I, such as 18F (where R₁₋₄ are methyl and R₆ is 2-methyl-4-thiazolyl).

Scheme 18

Alternatively, a compound of formula 18E where P1 is an oxygen protecting group, can be converted to a alkyl or aryl sulfonyl chloride. Reduction of the azido group and subsequent cyclization of a compound of formula 19A using a reducing agent such as a triaryl- or trialkylphosphine or with hydrogen and Lindlar's catalyst (Pd, CaCo3/Pb) provides a compound of formula I such as 18F (where R_{1-4} are methyl and R_6 is 2-methyl-4-thiazolyl).

Scheme 19

- 5 Alternatively, a compound of formula 18C where P_1 is an oxygen protecting group, can be converted to a compound of formula 20A where X is a halogen by treatment with, for example, triphenylphosphine and a carbon tetrahalide. Reduction of the azido group and subsequent cyclization of a compound of formula 20A using a reducing agent such as a
- 10 triaryl- or trialkylphosphine or with hydrogen and Lindlar's catalyst (Pd , $CaCO_3/Pb$) provides a compound of formula I, such as 18F (where R_{1-4} are methyl and R_6 is 2-methyl-4-thiazolyl).

The *in vitro* assessment of biological activity of the compounds of Formula I and II was performed as follows:

15 **In vitro Tubulin Polymerization**

Twice cycled (2X) calf brain tubulin was prepared following the procedure of Williams and Lee (see Williams, R.C., Jr., and Lee, J. C. Preparation of tubulin from brain. Methods in Enzymology 85, Pt. D: 376-385, 1982) and stored in liquid nitrogen before use. Quantification of tubulin polymerization potency is accomplished following a modified procedure of Swindell, et al., (see Swindell, C.S., Krauss, N.E., Horwitz, S.B., and Ringel, I. Biologically active taxol analogues with deleted A-ring side chain substituents and variable C-2' configurations. J. Med. Chem. 34: 1176-1184, 1991). These modifications, in part, result in the expression of tubulin polymerization potency as an effective concentration for any given compound. For this method, different concentrations of compound in polymerization buffer (0.1M MES, 1mM EGTA, 0.5 mM MgCl₂, pH 6.6) are added to tubulin in polymerization buffer at 37° in microcuvette wells of a Beckman (Beckman Instruments) Model DU 7400 UV spectrophotometer. A final microtubule protein concentration of 1.0 mg/ml and compound concentration of generally 2.5, 5.0, and 10 µM are used. Initial slopes of OD change measured every 10 seconds were calculated by the program accompanying the instrument after initial and final times of the linear region encompassing at least 3 time points were manually defined. Under these conditions linear variances were generally <10⁻⁶, slopes ranged from 0.03 to 0.002 absorbance unit/minute, and maximum absorbance was 0.15 absorbance units. Effective concentration

(EC_{0.01}) is defined as the interpolated concentration capable of inducing an initial slope of 0.01 OD/minute rate and is calculated using the formula:

EC_{0.01} = concentration/slope. EC_{0.01} values are expressed as the mean with standard deviation obtained from 3 different concentrations. EC_{0.01} values

5 for the compounds in this invention fall in the range 0.01-1000 μ M.

Cytotoxicity (In-Vitro)

Cytotoxicity was assessed in HCT-116 human colon carcinoma cells by MTS (3-(4,5-dimethylthiazol-2-yl)-5-(3-carboxymethoxyphenyl)-2-(4-sulphenyl)-2H-tetrazolium, inner salt) assay as reported in T.L. Riss, et.

10 al., "Comparison of MTT, XTT, and a novel tetrazolium compound MTS for in vitro proliferation and chemosensitivity assays.," *Mol. Biol. Cell* 3

(Suppl.):184a, 1992. Cells were plated at 4,000 cell/well in 96 well microtiter plates and 24 hours later drugs were added and serial diluted.

The cells were incubated at 37° for 72 hours at which time the

15 tetrazolium dye, MTS at 333 μ g/ml (final concentration), in combination with the electron coupling agent phenazine methosulfate at 25 μ M (final concentration) was added. A dehydrogenase enzyme in live cells reduces the MTS to a form that absorbs light at 492nm which can be quantitated spectrophotometrically. The greater the absorbance the greater the


20 number of live cells. The results are expressed as an IC₅₀, which is the drug concentration required to inhibit cell proliferation (i.e. absorbance at

450nM) to 50% of that of untreated control cells. The IC₅₀ values for compounds of this invention fall in the range 0.01 - 1000 nM.


As preferred compounds there are formula I and II compounds wherein

Q is selected from the group consisting of

5



G is



10

X is O or S

and Y is O.


Most preferred are compounds of formulas I and II

15 Wherein

WO 99/54319


PCT/US99/07475

Q is




5

G is



The following examples illustrate the present invention.

Example 1



5

[1S-[1R*,3R*(E),7R*,10S*,11R*,12R*,16S*]]-7,11-Dihydroxy-8,8,10,12-tetramethyl-3-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-4-oxa-17-azabicyclo[14.1.0]heptadecane-5,9-dione.

10

A. [1R-[1R*,3S*(E),7S*,10R*,11S*,12S*,16S*]]-7,11-Dihydroxy-8,8,10,12-tetramethyl-3-[1-methyl-2(2-methyl-4-thiazolyl)ethenyl]-4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione. [12,13-Epi Epothilone A]

15 To solution of epothilone C (100 mg, 0.21 mmol) in CH₃CN (2.5 mL) and 0.0004 M Na₂EDTA (1.5 ml) at 0 °C were added excess CF₃COCH₃ (1 mL) followed by 2KHSO₅•KHSO₄•K₂SO₄ (potassium peroxymonosulfate) (325 mg, 0.53 mmol) and NaHCO₃ (75 mg, 0.89 mmol). The reaction mixture was stirred for 35 min then treated with dimethylsulfide (0.50


mL) and water (3.75 mL). The reaction mixture was warmed to room temperature, extracted with EtOAc (4 x 10 mL), dried over MgSO₄, filtered, and concentrated. The residue was chromatographed with 55% EtOAc/hexanes to give epothilone A (59 mg, 57%) and compound A (30 mg, 29%) as a white solid. m/z: 494(M+H)⁺.

B. [4S-[4R*,7S*,8R*,9R*,13S*,14S*,16R*(E)]]-13-Azido-4,8,14-trihydroxy-5,5,7,9-tetramethyl-16-[1-methyl-2(2-methyl-4-thiazolyl)ethenyl]-1-oxacyclohexadecane-2,6-dione.

To a solution of Compound A (30mg, 0.06mmol) in EtOH (2.5 mL) were added NaN₃ (24 mg, 0.36 mmol) and NH₄Cl (2 mg, 0.04 mmol). The reaction mixture was heated at 60 °C for 2 days. The reaction was concentrated, and the residue was chromatographed with 60% EtOAc/hexanes to give Compound B (19 mg, 59%) as a colorless oil. m/z:537 (M+H)⁺.

C. [1S-[1R*,3R*(E),7R*,10S*,11R*,12R*,16S*]]-7,11-Dihydroxy-8,8,10,12- tetramethyl-3-[1-methyl-2(2-methyl-4-thiazolyl)ethenyl]-4-oxa-17-azabicyclo[14.1.0]heptadecane-5,9-dione.


To a solution of Compound B (19mg, 0.035mmol) in THF (1.5 mL) was added PPh₃ (12 mg, 0.047 mmol), and the reaction mixture was heated at 60 °C for 14hr. The reaction was concentrated, and the residue was chromatographed with MeOH/EtOAc/Et₃N (1:9:0.3) to give the title compound (11 mg, 62%) as a white solid. m/z:493 (M+H)⁺.

Example 2

[1S-[1R*,3R*(E),7R*,10S*,11R*,12R*,16S*]]-17-Formyl-7,11-
5 Dihydroxy-8,8,10,12-tetramethyl-3-[1-methyl-2-(2-methyl-4-
thiazolyl)ethenyl]-4-oxa-17-azabicyclo[14.1.0]heptadecane-5,9-dione.


To solution of epothilone 1C (2mg, 0.004 mmol) in acetone (0.2mL)
 at 0 °C was added 4-formyl-2-methyl-1,3,4-thiadiazolin-5-thione (1 mg,
 10 0.006 mmol), and the reaction mixture was stirred for 1 hr (See for
 example: Yazawa, H.: Goto, S. Tetrahedron Lett., (1985) 26, 3703-3706).
 The reaction was concentrated, and the residue was chromatographed with
 65% EtOAc/hexanes to give the title compound (0.8 mg, 38%) as a white
 film. m/z: 521 (M+H)⁺.

15

Example 3

- 5 [1S-[1R*,3R*(E),7R*,10S*,11R*,12R*,16S*]]-7,11-dihydroxy-8,8,10,12-tetramethyl-3-{1-methyl-2(2-methyl-4-thiazolyl)ethenyl]-4-oxa-17-thiabicyclo[14.1.0]heptadecane-5,9-dione.

Compound 1A (50 mg, 0.1 mmol) and ammonium thiocyanate (25 mg, 0.33 mmol) were dissolved in acetonitrile (1.5 mL) and heated in a sealed vial to 120 °C for 30 min. The reaction mixture was separated by HPLC (C18, 45% acetonitrile/55% 50 mM ammonium acetate buffer, pH 6.8, with UV detection at 254 nm) to give the title compound (11 mg, 21%).
m/z: 553 (M+H)⁺.

Example 4


[1S-[1R*,3R*(E),7R*,10S*,11R*,12R*,16S*]]-17-

[Dimethylamino)acetyl]-7,11-dihydroxy-8,8,10,12- tetramethyl-3-[1-

5 methyl-2(2-methyl-4-thiazolyl)ethenyl]-4-oxa-17-


azabicyclo[14.1.0]heptadecane-5,9-dione.

To a solution of compound 1C (10 mg, 0.020 mmol) and N,N-
dimethylglycine (2.3 mg, 0.022 mmol) in methylene chloride (0.20 mL) was
added a catalytic amount of DMAP followed by the addition of DCC (8.3
10 mg, 0.040 mmol) at °C. The reaction mixture was warmed to room
temperature and stirred for 18 h under Ar. The mixture was directly
purified by flash chromatography (short SiO₂ plug, 0-2%
methanol/chloroform gradient elution), and then further purified by reverse
phase HPLC (YMC S-5 ODS 10 x 250 mm column, A-95% water, 5%
15 MeCN, B=95% MeCN, 5% water, 0-100% B/40 minutes/gradient, flow rate
= 3mL/min, retention time = 29 min) to afford 3 mg of the title compound
(26%) as white a lyophilizate along with 4 mg of starting material (40%).
FABHRMS m/z 578.3287 (C₃₀H₄₇N₃O₆S, requires 577.3186).

Example 5

**[1S-[1R*,3R*(E),7R*,10S*,11R*,12R*,16S*]]-7,11-Dihydroxy-
5 8,8,10,12-tetramethyl-3-[1-methyl-2-(2-methyl-4-
thiazolyl)ethenyl]-17-(3-methyl-1-oxobutyl)-4-oxa-17-
azabicyclo[14.1.0]heptadecane-5,9-dione.**

To a solution of compound 1C (10 mg, 0.020 mmol) and triethylamine (10
μL, 0.080 mmol) in methylene chloride (0.20 mL) was added isovaleryl
10 chloride (3.0 μL, 0.024 mmol) at 0°C. The reaction mixture was warmed to
room temperature and stirred for 1 h under Ar. The mixture was diluted with
methylene chloride (5 mL) and was quenched with saturated aqueous
sodium bicarbonate (1 mL) at 0°C. The organic layer was collected, dried
over sodium sulfate, and concentrated. The oily residue was purified by
15 flash chromatography (SiO₂, 4.5 x 30 cm, 25-50% EtOAc-hexanes gradient
elution) to afford the title compound (6 mg, 51%) as a white solid. FABHRMS
m/z 577.3343 (C₃₁H₄₈N₂O₆S, requires 576.3233).

Example 6


20

[1S-[1R*,3R*(E),7R*,10S*,11R*,12R*,16S*]]-7,11-Dihydroxy-8,8,10,12-tetramethyl-3-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-17-(methylsulfonyl)-4-oxa-17-azabicyclo[14.1.0]heptadecane-5,9-dione.

- 5 To a solution of compound 1C (10 mg, 0.020 mmol) and pyridine (8 μ l, 0.10 mmol) in methylene chloride (0.20 ml) under Ar was added methanesulfonyl chloride (1.6 μ l, 0.021 mmol) at -15°C. The reaction mixture was kept at -15°C and stirred for 30 minutes. The mixture was diluted with methylene chloride (5 mL) and was quenched with saturated aqueous sodium
- 10 bicarbonate (1 mL) at 0°C. The organic layer was collected, dried over sodium sulfate, and concentrated. The oily residue was purified by flash chromatography (SiO₂, 4.5 x 30 cm, 25-50% EtOAc-hexanes gradient elution) to afford the title compound (6 mg, 52%) as a white solid. FABHRMS m/z 571.2525 (C₂₇H₄₂N₂O₇S₂, requires 570.2433).

15

Example 7



- 20 **[1S-[1R*,3R*(E),7R*,10S*,11R*,12R*,16S*]]-N-Ethyl-7,11-dihydroxy-8,8,10,12-tetramethyl-3-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-5,9-dioxo-4-oxa-17-azabicyclo[14.1.0]heptadecane-17-carboxamide.**


To a solution of compound 1C (10 mg, 0.020 mmol) in EtOAc (0.20 mL) was

25 added ethyl isocyanate (1.9 μ L, 0.024 mmol) at 0°C under Ar. The reaction mixture was warmed to room temperature and stirred for 30 minutes. The

mixture was directly purified by flash chromatography (SiO₂, 4.5 x 30 cm, 0-3% methanol-chloroform gradient elution) to provide the title compound (5.5 mg, 66%) as white solid. FABHRMS m/z 564.3367 (C₂₉H₄₅N₃O₆S, requires 563.3029).

5


Example 8



- 10 **[1S-[1R*,3R*(E),7R*,10S*,11R*,12R*,16S*]]-7,11-Dihydroxy-8,8,10,12,17-pentamethyl-3-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-4-oxa-17-azabicyclo[14.1.0]heptadecane-5,9-dione**

- To a solution of compound 1C (10 mg, 0.020 mmol) in THF (0.20 mL) under
 15 Ar was added proton sponge (0.05 M solution in THF, 4.2 μL, 0.021 mmol) followed by dimethylsulfate (2.8 μL, 0.030 mmol) at -10°C. The reaction mixture was then stirred overnight at 4°C. The mixture was directly purified by flash chromatography (SiO₂, 4.5 x 30 cm, 0-3% methanol-chloroform gradient elution) to afford the title compound (5.0 mg, 40%) as white solid.
 20 FABHRMS m/z 507.2888 (C₂₇H₄₂N₂O₅S, requires 506.2814).

25


Example 9

- 5 **[1S-[1R*,3R*(E),7R*,10S*,11R*,12R*,16S*]]-7,11-Dihydroxy-8,8,10,12-tetramethyl-3-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-5,9-dioxo-4-oxa-17-azabicyclo[14.1.0]heptadecane-17-carboxylic acid, 1,1-dimethylethyl ester.**
- 10 Di-*tert*-butyl dicarbonate (0.008 mmol, 1.8 mg) was added to Compound 1C (0.004 mmol, 2 mg) and triethylamine (0.0048 mmol, 0.0007 ml) in THF (0.15 mL) at room temperature. After 30 min, the reaction was concentrated and purified by flash chromatography (50%EtOAc/Hexanes) to afford the title compound as a clear oil (1.8 mg). (M+H)⁺ 593.

15

20

25

Example 10

5 **[1S-[1R*,3R*(E),10S*,11R*,12R*,16S*]]-11-Hydroxy-8,8,10,12-tetramethyl-3-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-4,18-dioxabicyclo[14.2.0]octadec-6(E)-ene-5,9-dione**

10 **A. [1S-[1R*,3R*(E),10S*,11R*,12R*,16S*]]-11-Hydroxy-8,8,10,12-tetramethyl-3-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-4,17-dioxabicyclo[14.1.0]heptadec-6(E)-ene-5,9-dione**

HCO₂H (22.7 mmol, 0.85 mL) was added to a solution of epothilone B (4.5 mmol, 2.3g), NEt₃ (45.3 mmol, 6.3 mL) and *N,N*-dimethylaminopyridine (10 mmol, 1.2g) in CH₂Cl₂ (100 mL) at room temperature. The reaction mixture
 15 was cooled to -15°C, and (CH₃CO)₂O (22.7 mmol, 2.1 mL) was added dropwise. The reaction was maintained at -15°C for 15 minutes then at room temperature for 30 minutes. The reaction was quenched with pH 7 buffer (100 mL) and extracted with CH₂Cl₂ (3X100 mL). The combined organic phase was washed with 1N HCl (100 mL), saturated NaHCO₃ (100
 20 mL), dried over Na₂SO₄, and concentrated to give the bis-formate of epothilone B as a white solid, which was taken to next step without further purification. 1,8-Diazabicyclo[5.4.0]undec-7-ene (44.7 mmol, 6.7 mL) was added dropwise to the crude bis-formate in CH₂Cl₂ (125 mL) at room temperature. The reaction mixture was stirred for 1 hour, quenched with pH
 25 4 buffer (200 mL) and extracted with CH₂Cl₂ (3X200 mL). The combined organic phase was washed with saturated NaHCO₃ (200 mL), dried over

Na₂SO₄ and concentrated to give a viscous oil which was taken up in MeOH (95 mL) and treated with NH₃ (2M solution in MeOH, 21.9 mmol, 11 mL). The reaction mixture was maintained at room temperature for 4 hours, concentrated and purified by flash chromatography (60% EtOAc/Hexanes to neat EtOAc) to afford Compound A as a white solid (2.05g, 92% for 3 steps).
 5 (M+H)⁺ 490.

B. [1S-[1R*,3R*(E),10S*,11R*,12R*,16S*]]-11-Triethylsilyloxy-8,8,10,12-tetramethyl-3-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-4,17-dioxabicyclo[14.1.0]heptadec-6(E)-ene-5,9-dione
 10

Et₃SiCl (10.8 mmol, 1.8 mL) was added to Compound A (1.08 mmol, 529 mg), *N,N*-diisopropylethylamine (16.2 mmol, 2.8 mL), and *N,N*-dimethylaminopyridine (1.08 mmol, 132 mg) in CH₂Cl₂ (10 mL) at room temperature. The reaction mixture was then heated at 45°C for 12 hours, quenched with pH 4 buffer (50 ml), extracted with CH₂Cl₂ (3X25 mL), dried over Na₂SO₄, concentrated and purified by flash chromatography (hexanes to 20%EtOAc/hexanes) to afford Compound B as a white solid (590 mg).
 15 (M+H)⁺ 604.

20

C. [7R,8S,9S,14S,16S(E)]-14-Hydroxy-13-hydroxymethyl-8-triethylsilyloxy-5,5,7,9-tetramethyl-16-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-1-oxa-3(E)-cyclohexadecene-2,6-dione.

Diethylaluminum-2,2,6,6-tetramethylpiperidide [1.5 mL of 0.28M solution in benzene maintained at 0°C, 0.42 mmol; prepared by method of Paquette, L.A. *et. al. J. Org. Chem* 1990, 55, 1589; stepwise addition of *n*-BuLi (1.1 mmol, or 0.68 mL of 1.6 M solution in hexanes) and diethylaluminumchloride (1.1 mmol, 1.1 mL of 1.0 M solution in hexanes) to 2,2,6,6-tetramethylpiperidine (1.1 mmol, 0.15 mL) in benzene (2.0 ml) at 0°C] was
 25 added dropwise over 20 min to Compound B (0.18 mmol, 109 mg) in benzene (2.0 mL) at 0°C. The reaction mixture was quenched with saturated NH₄Cl (8 mL), extracted with CH₂Cl₂ (3X10 mL), dried over Na₂SO₄,
 30

concentrated and purified by flash chromatography (25% EtOAc/hexanes) to afford 13,14-allylic alcohol as a clear oil (70 mg, 64%). (M+H)⁺ 604.

BH₃•THF (0.35 mmol, 0.35 ml of 1.0 M solution in THF) was added dropwise to the 13,14-allylic alcohol intermediate in THF (4.0 mL) at -78°C. The
5 reaction mixture was stirred at -78°C for 10 minutes, then transferred to a 0°C bath. After 4.6 hours, the reaction mixture was stirred at room temperature for 10 minutes, diluted with THF/EtOH (4.0 mL), quenched by stepwise addition of pH 7 phosphate buffer (4.0 mL) and 30% H₂O₂ aqueous solution (4.0 mL). The resulting reaction mixture was stirred at room
10 temperature for 17 hours, extracted with CH₂Cl₂ (3X10 mL), dried over Na₂SO₄, concentrated and purified by flash chromatography (60-75% EtOAc/hexanes) to afford Compound C as a clear oil (51.2 mg, 2:1 mixture of 13R/13S diastereomers). (M+H)⁺ 622.

15 **D. [7R, 8S,9S,13R,14S,16S(E)]-14-Hydroxy-13-*p*-toluenesulfonyloxymethyl-8-triethylsilyloxy-5,5,7,9-tetramethyl-16-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-1-oxa-3(E)-cyclohexadecene-2,6-dione**

p-Toluenesulfonylchloride (0.1 mmol, 19 mg) was added to Compound C
20 (31 mg, 2:1 (trans:cis), 0.05 mmol) and pyridine (0.25 mmol, 0.02 mL) in CH₂Cl₂ (1.0 mL) at room temperature. The reaction mixture was heated at 35°C for 16 hours, then additional *p*-toluenesulfonylchloride (0.1 mmol, 19 mg) and pyridine (0.25 mmol, 0.02 mL) were added. After 4 hours, the reaction was quenched with pH 4 buffer (20 mL), extracted with CH₂Cl₂
25 (3X10 mL), dried over Na₂SO₄, concentrated and purified by flash chromatography (30% to 60% EtOAc/hexanes) to afford tosylate, as a clear oil (22.3 mg, 13R/13S=2) and unreacted starting material (3.8 mg). The 13S and 13R (Compound D) isomers were separated by flash chromatography (neat Et₂O). (M+H)⁺ 776 for both isomers.

30

E. [1S-[1R*,3R*(E),10S*,11R*,12R*,16S*]]-11-Triethylsilyloxy-8,8,10,12-tetramethyl-3-[1-methyl-2-(2-methyl-4-

thiazolyl)ethenyl]-4,18-dioxabicyclo[14.2.0]octadec-6(E)-ene-5,9-dione.


NaH (60% in nujol, a pinch, >0.053 mmol) was added to Compound D (41 mg, 0.053 mmol) in DMF (1.0 mL) at room temperature. After 20 minutes, the reaction was quenched with saturated NH_4Cl (10 mL), extracted with CH_2Cl_2 (3X50 mL), dried over Na_2SO_4 , concentrated and purified by flash chromatography (10% to 30% EtOAc/hexanes) to afford Compound E, as a clear oil (11.3 mg). $(\text{M}+\text{H})^+$ 604.

F. [1S-[1R*,3R*(E),10S*,11R*,12R*,16S*]]-11-Hydroxy-8,8,10,12-tetramethyl-3-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-4,18-dioxabicyclo[14.2.0]octadec-6(E)-ene-5,9-dione.

Trifluoroacetic acid (1 mL of 20% solution in CH_2Cl_2) was added to Compound E (0.011 mmol, 6.5 mg) in CH_2Cl_2 (1 mL) at -20°C . After 20 minutes, the reaction mixture was quenched with saturated NaHCO_3 (1mL), extracted with CH_2Cl_2 (3X10 mL), dried over Na_2SO_4 , concentrated and purified by flash chromatography (20% to 60% EtOAc/hexanes with 1% triethylamine) to afford the title compound, as a white solid (4.3 mg). $(\text{M}+\text{H})^+$ 490.

20

Example 11



[5S-[5R*(E),9R*,12S*,13R*,14R*]]-4,9,10,13,14,15,16,17-Octahydro-9,13-dihydroxy-2,10,10,12,14-pentamethyl-5-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-5H-oxacyclohexadecino[5,4-d]thiazole-7,11(8H,12H)-dione

25

A. [1S-[1R*,3R*(E),7R*,10S*,11R*,12R*,16S*]]-7,11-Bistriethylsilyloxy-8,8,10,12-tetramethyl-3-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione.

Et₃SiCl (4.15 mmol, 0.7 mL) was added to epothilone A (4.2 mmol, 205 mg), *N,N*-diisopropylethylamine (6.2 mmol, 1.1 mL), and imidazole (2.1 mmol, 141 mg) in DMF (5 mL) at room temperature. The reaction mixture was then heated at 40°C for 15.5 hours. Additional *N,N*-diisopropylethylamine (4.2 mmol, 0.720 mL) and Et₃SiCl (2.1 mmol, 0.35 mL) were added and the reaction was heated to 60°C. After 48 hours, the reaction mixture was concentrated and purified by flash chromatography (5 to 10% EtOAc/hexanes) to afford Compound A as a clear oil (264mg). (M+H)⁺ 722.

B. [4S-[4R*,7S*,8R*,9R*,13S*,14S*,16R*(E)]-14-Bromo-13-hydroxy-4,8-bistriethylsilyloxy-5,5,7,9-tetramethyl-16-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-1-oxacyclohexadecane-2,6-dione.

MgBr₂•OEt₂ (9.7 mmol, 2.5 g) was added to Compound A (6.45 mmol, 4.66 g) in CH₂Cl₂ (80 mL) at -78°C. After 2 hours, the reaction mixture was warmed to 50°C and additional MgBr₂•OEt₂ (9.7 mmol, 2.5 g) was added. The reaction mixture was again warmed to -30°C and maintained for 1 hour; and then warmed to -20°C, where additional MgBr₂•OEt₂ (6.45 mmol, 1.7 g) was added. After 1 hour, the reaction mixture was quenched with pH 7 phosphate buffer (20 mL), extracted with CH₂Cl₂ (3X100 mL), dried over Na₂SO₄, concentrated and purified by flash chromatography (30% EtOAc/hexanes) to afford Compound B as a white solid (2.95 g). (M+H)⁺ 802.

C. [4S-[4R*,7S*,8R*,9R*,13S*,14S*,16R*(E)]-14-Bromo-4,8-bistriethylsilyloxy-5,5,7,9-tetramethyl-16-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-1-oxacyclohexadecane-2,6,13-trione.

Pyridinium chlorochromate (1.48 mmol, 320 mg) was added to Compound B (0.37 mmol, 300 mg), pyridine (3.7 mmol, 0.3 mL) and small amount of

crushed 4A molecular sieves in CH₂Cl₂ (3 mL). After 5 hours, the reaction mixture was diluted with CH₂Cl₂ (50 mL), filtered through a short pad of silica gel, concentrated and purified by flash chromatography (5%EtOAc/Hexanes) to afford Compound C as a white solid (273 mg). (M+H)⁺ 800.

5

D. [5S-[5R*(E),9R*,12S*,13R*,14R*]]-4,9,10,13,14,15,16,17-Octahydro-9,13-bistriethylsilyloxy-2,10,10,12,14-pentamethyl-5-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-5H-oxacyclohexadecino[5,4-d]thiazole-7,11(8H,12H)-dione

10

Thioacetamide (0.03 mmol, 2.3 mg) was added to Compound C (0.015 mmol, 12 mg) and *N,N*-diisopropylethylamine (0.038 mmol, 0.006 mL) in ethanol (0.3 mL). The reaction mixture was heated at 80°C for 24 hours.

The reaction mixture was concentrated and purified by flash chromatography (50% to 75% EtOAc/Hexanes) to afford a hydroxythiazoline intermediate (4.6 mg, 39%). (M+H)⁺ 795.

Methanesulfonyl chloride (0.01 mmol, 0.0008 mL) and triethylamine (0.04 mmol, 0.006 mL) were added stepwise to the hydroxythiazoline intermediate (0.005 mmol, 4.2 mg) in CH₂Cl₂ (0.175 mL) at 0°C. The reaction flask was then removed from the ice bath and stirred at room temperature for 5 hours, concentrated and purified by flash chromatography (18% EtOAc/Hexanes) to afford Compound D (1.2 mg). (M+H)⁺ 777.

20

E. [5S-[5R*(E),9R*,12S*,13R*,14R*]]-4,9,10,13,14,15,16,17-Octahydro-9,13-dihydroxy-2,10,10,12,14-pentamethyl-5-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-5H-oxacyclohexadecino[5,4-d]thiazole-7,11(8H,12H)-dione


25

Trifluoroacetic acid (0.150 mL of 40% solution in CH₂Cl₂) was added to Compound D (0.0036 mmol, 2.8 mg) in CH₂Cl₂ (0.150 mL) at 0°C. After 20 minutes, the reaction mixture was concentrated and purified by flash

30

chromatography (80% EtOAc/Hexanes) to afford the title compound (1.8 mg).
(M+H)⁺ 549.

Example 12



5

[1S-[1R*,3R*(E),7R*,10S*,11R*,12R*,16S*]]-7,11-Dihydroxy-
8,8,10,12-tetramethyl-3-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-4-
oxa-17-azabicyclo[14.1.0]heptadecane-5,9-dione.

10 **A. [1S-[1R*,3R*(E),7R*,10S*,11S*,12R*,16S*]]-7,11-Bistriethylsilyloxy-8,8,10,12-tetramethyl-3-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione.**

Et₃SiCl (25 ml, 149 mmol) was added to epothilone A (10.39 g, 21 mmol),
15 *N,N*-diisopropylethylamine (55 ml, 315 mmol), and imidazole (7.15 g, 105 mmol,) in DMF (75 mL) at 25°C. The reaction mixture was then heated at 55°C for 6.5 h and concentrated *in vacuo*. The residue was then diluted with CH₂Cl₂ (100 mL) and the organic extracts were washed with NaCHO₃ (30 mL), dried over Na₂SO₄ and concentrated *in vacuo*. The residue was
20 purified by flash chromatography (SiO₂, 5.0 x 30 cm, hexanes to 15% EtOAc/hexanes gradient elution) to afford Compound A as a white solid [15.1 g, >95%. MS (ESI⁺):(M+H)⁺ 722.

B. [4S-[4R*,7S*,8R*,9R*,13S*,14S*,16R*(E)]-14-Bromo-13-hydroxy-4,8-bistriethylsilyloxy-5,5,7,9-tetramethyl-16-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-1-oxacyclohexadecane-2,6-dione.

25

To a solution of compound A (2.0 g, 2.8 mmol) in CH₃Cl₃ (30 mL) at -20 °C under Ar was added MgBr₂•OEt₂ (3 x 1.1g, 12 mmol total) in three portions every two hours while maintaining an internal temperature between -15 and
 5 -5 °C. After 7 h, the reaction mixture was quenched with pH7 phosphate buffer (40 mL) and brine (40 mL), carefully extracted with EtOAc (3 x 10 ml), dried (Na₂SO₄) and concentrated *in vacuo*. The residue was purified by flash chromatography (SiO₂, 4.5 x 25 cm, 10-20% EtOAc/hexanes gradient elution) to afford Compound B as a white solid [1.0 g, 45% (67% based on
 10 0.6g of recovered starting material; <2% of the other C13-OH/C12-Br regioisomer was detected.). MS (ESI⁺): (M+H)⁺ +802.

**C. [4S-[4R*,7S*,8R*,9R*,13S*,14R*,16R*(E)]-14-Azido-4,8-bistriethylsilyloxy-13-hydroxy-5,5,7,9-tetramethyl-16-[1-methyl-2-
 15 (2-methyl-4-thiazolyl)ethenyl]-1-oxacyclohexadecane-2,6,dione.**

To a solution of compound B (0.17 g, 0.21 mmol) in DMF (2 ml) Ar was added sodium azide (0.14 mg, 2.1 mmol) and the resulting suspension was warmed to 43 °C. After 36 h, the solvent was removed *in vacuo* and the
 20 residue was directly purified by flash chromatography (SiO₂, 2.5 x 15 cm, 10-20% EtOAc/hexanes gradient elution) to give Compound C (0.14 G, 88%) as a white foam. MS (ESI⁺): (M+H)⁺ 765

**D. [4S-[4R*,7S*,8R*,9R*,13R*,14R*,16R*(E)]-14-Azido-4,8-bistriethylsilyloxy-13-(4-nitrobenzoyloxy)-5,5,7,9-tetramethyl-16-
 25 [1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-1-oxacyclohexadecane-2,6-dione.**

To a solution of compound C (0.10 mg, 0.13 mmol) in THF under Ar
 30 was sequentially added 4-nitrobenzoic acid (55 mg, 0.33 mmol), triphenylphosphine (86 mg, 0.33 mmol and diethyl azodicarboxylate (52 mL, 0.33 mmol). The reaction mixture was stirred at 25 °C for 1.5 h, concentrated

in vacuo and the residue was purified by flash chromatography (SiO₂, 2.5 x 10 cm, 10-20% EtOAc/hexanes gradient elution) to afford Compound D (0.10 g, 85%) as a white foam. MS (ESI)⁺: 914.6 (M+H)⁺

- 5 **E. [4S-[4R*,7S*,8R*,9R*,13R*,14R*,16R*(E)]-14-Azido-4,8-bistriethylsilyloxy-13-hydroxy-5,5,7,9-tetramethyl-16-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-1-oxacyclohexadecane-2,6-dione.**

Compound D (0.10 mg, 0.11 mmol) was treated with 2.0 M ammonia
10 in methanol (1mL) at at 25 °C under Ar for 4 h. the solvent was removed *in vacuo* and the residue was directly purified by flash chromatography (SiO₂, 1.5 x 10 cm, 10-30% EtOAc/hexanes gradient elution) to afford Compound E (50 mg, 60%) as a white foam. MS (ESI)⁺: 763.3 (M-H)⁺.

- 15 **F. [4S-[4R*,7S*,8R*,9R*,13R*,14R*,16R*(E)]]-14-Azido-4,8-13-trihydroxy-5,5,7,9-tetramethyl-16-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-1-oxacyclohexadecane-2,6-dione**

Compound E (15 mg, 0.20 mmol) was treated with 20% trifluoroacetic
20 acid in methylene chloride (0.2 mL) at 0 °C under Ar for 10 min. The reaction mixture was concentrated under a constant stream of nitrogen at 0 °C and the residue was directly purified by flash chromatography (SiO₂, 1x 5 cm, 0-5% MeOH/CHCl₃ gradient elution) to afford Compound F (9 mg, 86%) as a film. MS (ESI)⁺: 537.3 (M-H)⁺.

25

- G. [1S-[1R*,3R*(E),7R*,10S*,11R*,12R*,16S*]]-7-11-Dihydroxy-8,8,10,12-tetramethyl-3-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-4-oxa-17-azabicyclo[14.1.0]heptadecane-5,9-dione**


30

To a solution of Compound F (9 mg, 17 μmol) THF (0.2 mL) under Ar was added triphenylphosphine (18 mg, 67 μmol). The reaction mixture was

warmed to 45 °C for 4 h, and the solvent was removed under a constant flow of nitrogen. The residue was purified by radial chromatography (1 mm SiO₂, GF rotor, 2-10% MeOH/CHCl₃ gradient elution) to afford the title compound (4 mg, 50%) as a film.

5

Example 13



10 **[1S-[1R*,3R*(E),7R*,10S*,11R*,12R*,16R*]]-7,11-Dihydroxy-8,8,10,12,16-pentamethyl-3-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-4-azabicyclo[14.1.0]heptadecane-5,9-dione**

A. [4S-[4R*,7S*,8R*,9R*,16R*(E)]]-4,8-Dihydroxy-5,5,7,9,13-pentamethyl-16-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-1-oxa-13(Z)-cyclohexadecene-2,6-dione. [Epothilone D]

15 To anhydrous THF (5 ml) at -78 °C under argon was added WCl₆ (198 mg, 0.5 mmol) followed by *n*BuLi (0.625 ml of 1.6 *M* solution in hexanes, 1.0 mmol). The reaction was allowed to warm to room temperature over a 20 min period. An aliquot (0.50 ml, 0.05 mmol) of the tungsten reagent was removed and added to epothilone B (9.0 mg, 0.018 mmol) under argon and the reaction stirred for 15 min then quenched by the addition of saturated NaHCO₃ (1 ml). The reaction was extracted with EtOAc (3 x 1 ml). The combined extracts dried (Na₂SO₄), and filtered. The volatiles removed under vacuum. The residue was chromatographed with 35% EtOAc/hexanes to give compound A (7.0 mg, 0.014 mmol) in 80% yield.

25 *m/z*: 492.3 (M+H)⁺.

B. [4S-[4R*,7S*,8R*,9R*,16R*(E)]]-4,8-Bistriethylsilyloxy-5,5,7,9,13-pentamethyl-16-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-1-oxa-13(Z)-cyclohexadecene-2,6-dione. [Bis-Triethylsilyl Epothilone D]

To a solution of compound A (30 mg, 0.061 mmol) in anhydrous
 5 CH₂Cl₂ (1.25 mL) under argon were added *N,N*-diisopropylethylamine (0.16 mL, 0.92 mmol, 15 eq) followed by triethylsilylchloride (0.10 mL, 0.61 mL, 10 eq). The reaction mixture was stirred for 18 hrs. The reaction mixture was cooled to 0 °C then 2,6-lutidine (0.021 mL, 0.18 mmol, 3 eq) was added followed by triethylsilyltrifluoromethanesulphonate (0.056 mL,
 10 0.24 mmol, 4 eq). The reaction was stirred for 0.5 hr then poured into a 1:1 mixture of H₂O/saturated NaHCO₃ (1 mL) and extracted with CH₂Cl₂ (3 x 1 mL). The combined organics were dried (Na₂SO₄), filtered, and the volatiles were removed. The residue was chromatographed with 1% Et₂O/CH₂Cl₂ to give 35 mg of compound B (80% yield) as a clear glass. *m/z*:
 15 720.5 (M+H)⁺.

C. [1S-[1R*,3R*(E),7R*,10S*,11R*,12R*,16R*]]-7,11-Dihydroxy-8,8,10,12,16-pentamethyl-3-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-4-oxabicyclo[14.1.0]heptadecane-5,9-dione

20 To a solution of diethylzinc (0.24 mL of 1.0 M in heptane, 0.24 mmol, 5 eq) in 1,2-dichloroethane (1.5 mL) at -15 °C under argon was added chloriodomethane (0.035 mL, 0.48 mmol, 10 eq), and the mixture was stirred for 10 min. A solution of compound B (35 mg, 0.048 mmol) in 1,2-dichloroethane (0.40 mL) was slowly added, and the reaction mixture
 25 was stirred for 1.5 hrs. The reaction was quenched by addition of saturated NH₄Cl (1.5 mL) and extracted with CH₂Cl₂ (3 x 2 mL). The combined organics were dried (Na₂SO₄), filtered, and the volatiles removed *in vacuo*. To the residue was added 15% trifluoroacetic acid/CH₂Cl₂ (0.50

mL) and the reaction stirred for 15 min. The volatiles were removed under a stream of air and the residue was chromatographed with 70% EtOAc/hexanes to give 2.2 mg of compound C (10% yield- two steps) as a white film; m/z : 506.3 ($M+H$)⁺.

- 5 Alternatively, sodium hydroxide (0.3 ml of 50% solution in H₂O) was added to compound 1B (109 mg, 0.15 mmol), PhCH₂(CH₃CH₂)₃NCl (0.7 mg, 0.002 mmol), and EtOH (0.03ml) in CHBr₃ (1.0 ml). The resulting mixture was heated at 40°C for 2 hr. The brown reaction mixture was diluted with H₂O (30 ml), extracted with CH₂Cl₂ (3X30 ml), dried over Na₂SO₄,
10 concentrated and purified by flash chromatography (stepwise gradient: 5 to 25% Et₂O/hexanes) to afford a dibromocyclopropane intermediate as a light brown oil (40mg, 30% yield). ($M+H$)⁺ 892.3

- Bu₃SnH (3.4 mmol, 0.91 ml) was added to the dibromocyclopropane intermediate (0.34 mmol, 305 mg) and 2,2-azobisisobutyronitrile (0.034
15 mmol, 6 mg) in hexanes (7.0 ml). The reaction mixture was heated at 70 °C for 5 hr. The reaction was concentrated and purified by flash chromatography (stepwise gradient: hexanes to 20%Et₂O/hexanes) to afford a reduced cyclopropane intermediate as a clear film (228 mg, 91%). ($M+H$)⁺ 734.7.

- 20 The preceding cyclopropane intermediate(0.31 mmol, 228 mg) was dissolved in CF₃CO₂H/CH₂Cl₂ (20% solution by volume, 10 ml) and stirred at -15°C for 1.5 hr. The reaction mixture was concentrated and purified by flash chromatography (70%EtoAc/Hexanes) to afford compound C as a clear oil (111 mg, 71%). ($M+H$)⁺ 506.3. ¹H NMR (CDCl₃, 400 MHz) d 7.04
25 (s, 1H), 6.64 (s, 1H), 5.16 (dd, J=8.0, 3.4 Hz, 1H), 4.17 (dd, J=9.5, 2.8 Hz, 1H), 3.79-3.83 (m, 1H), 3.23 (dq, J=6.7, 4.5 Hz, 1H), 2.79 (s, 3H), 2.52 (dd, J=15.1, 9.7 Hz, 1H), 2.41 (dd, J=15.2, 2.9 Hz, 1H), 1.98-2.02 (m, 1H), 2.00 (s, 3H), 1.63-1.73 (m, 1H), 1.40-1.58 (m, 5H), 1.36 (s, 3H), 1.20-1.33 (m,


1H), 1.11-1.17 (m, 1H), 1.15 (d, J=6.8 Hz, 3H), 1.08 (s, 3H), 0.96 (d, J=7.0 Hz, 3H), 0.94 (s, 3H), 0.40-0.54 (m, 1H), 0.37 (dd, J=8.8, 4.1 Hz, 1H), -0.14-(-0.10) (m, 1H).

5 **D. [1S-[1R*,3R*(E),7R*,10S*,11R*,12R*,16R*]]-7,11-Dihydroxy-8,8,10,12,16-pentamethyl-3-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-4-azabicyclo[14.1.0]heptadecane-5,9-dione**

A suspension of compound C (26 mg, 51 mmol) and sodium azide (6.7 mg, 10
10 0.10 mmol) in a THF-H₂O mixture (2:1, 1.0 mL) was degassed for 15 min with nitrogen. To this mixture was added a catalytic amount (12 mg, 10 mmol) of tetrakis(triphenylphosphine) palladium (0) under Ar. The reaction mixture was warmed to 45 °C for 1 hr and cooled to 25 °C.

The resulting yellow homogeneous solution was directly treated with
15 a 1.0 M solution of trimethylphosphine in THF (0.11 mL, 0.11 mmol) at 25 °C and the reaction mixture was stirred for 1 hr at ambient temperature.

The amino acid-containing mixture was then diluted with MeCN-DMF (12:1, 1.9 mL), cooled to 0 °C and treated with 1-hydroxybenzotriazole hydrate (7.0 mg, 51 mmol) followed by 1-(3-dimethylaminopropyl)-3-
20 ethylcarbodiimide hydrochloride (25 mg, 0.13 mmol). The reaction mixture was gradually warmed to 25 °C, stirred for 3 hr, diluted with H₂O (5 mL) and extracted with EtOAc (3 x 5 mL). The organic extracts were washed with H₂O (8 mL), saturated aqueous NaHCO₃ (2 x 8 mL), and saturated aqueous NaCl (8 mL). The organic extracts were dried (Na₂SO₄) and concentrated *in*
25 *vacuo*. The residue was purified by radial chromatography (1 mm SiO₂ rotor, 2 % MeOH-CHCl₃) to afford the title compound (film, 3.0 mg, 12 % overall yield), along with a minor amount of a diastereomer (5:1 ratio by ¹H NMR). (M+H)⁺ 505.4

Example 14

5 **[1S-[1R*,3R*(E),7R*,10S*,11R*,12R*,16S*]]-17-Dichloro-7,11-dihydroxy-8,8,10,12,16-pentamethyl-3-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-4-azabicyclo[14.1.0]heptadecane-5,9-dione**

A. **[1S-[1R*,3R*(E),7R*,10S*,11R*,12R*,16S*]]-7,11-Dihydroxy-8,8,10,12,16-pentamethyl-3-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-4-aza-17-oxabicyclo[14.1.0]heptadecane-5,9-dione**

A suspension of epothilone B (5.06 g, 9.97 mmol) and sodium azide (0.777 g, 12.0 mmol) in a THF-H₂O mixture (5:1, 96 mL) was degassed for 15-20 min with nitrogen and then treated with a catalytic amount (1.2 g, 0.997 mmol) of tetrakis(triphenylphosphine) palladium (0) under Ar. The reaction mixture was warmed to 45 °C for 20 min and cooled to 25 °C.

The resulting bright yellow homogeneous solution was directly treated with a 1.0 M solution of trimethylphosphine in THF (24.9 mL, 24.9 mmol) at 25 °C and the reaction mixture was stirred for 1-2 hr at ambient temperature.

The amino acid-containing mixture was then diluted with MeCN-DMF (20:1, 450 mL), cooled to 0 °C and treated with 1-hydroxybenzotriazole hydrate (1.35 g, 9.97 mmol) followed by 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (4.78 g, 24.9 mmol). The reaction mixture was warmed to 25 °C, stirred for 12 hr and extracted with EtOAc (4 x 200 mL). The organic extracts were washed with H₂O (400 mL), saturated

aqueous NaHCO₃ (400 mL), and saturated aqueous NaCl (400 mL). The organic extracts were dried (Na₂SO₄) and concentrated *in vacuo*. The residue was purified by flash chromatography (SiO₂, 5.0 x 25 cm, 2% MeOH-CHCl₃) and then HPLC (YMC S-15 ODS 50 x 500 mm column, 38 to 95% MeCN/H₂O, gradient (40 min), 50 mL/min flow rate). The appropriate fractions were concentrated *in vacuo* and the residue was lyophilized from aqueous acetonitrile to afford compound A (0.998 g, 20%), as a white lyophilizate. MS (ESI⁺): 507.2 (M+H)⁺; MS(ESI⁻): 505.4 (M-H)⁻.

10 **B. [4S-[4R*,7S*,8R*,9R*,16R*(E)]]-4,8-Dihydroxy-5,5,7,9,13-pentamethyl-16-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-1-aza-13(Z)-cyclohexadecene-2,6-dione**

Tungsten chloride (1.98 g, 5.00 mmol) was added in one portion to pre-cooled THF (50 ml) at -78 °C, and the resulting mixture was stirred for 5 min. n-BuLi (1.6 M in hexanes, 6.25 ml, 10.0 mmol) was added to the suspension which was kept at -78 °C for 10 min, then warmed to room temperature over 30 min. After 30 additional min at room temperature, the reaction mixture was added to compound A (0.351 g, 0.690 mmol) via syringe. After 15 min, the reaction mixture was cooled to 0 °C, diluted with ethyl acetate (10 ml), and quenched with saturated sodium bicarbonate (15 ml). The reaction mixture was extracted with ethyl acetate (5 x 20 ml), dried over sodium sulfate, concentrated and purified by reverse phase HPLC (solvent A = 5:95, acetonitrile:H₂O, solvent B = 95:5, acetonitrile:H₂O; gradient: 40-100% solvent B over 40 minutes; Flow rate: 40 ml/min, 50X500 mm YMC ODS S-15 column) to afford compound B (0.164 g, 36%) as a white solid. (M+H)⁺ 491.3

30 **C. [4S-[4R*,7S*,8R*,9R*,16R*(E)]]-4,8-Bis(t-butyldimethylsilyloxy)-5,5,7,9,13-pentamethyl-16-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-1-aza-13(Z)-cyclohexadecene-2,6-dione**

To a solution of compound B (19 mg, 0.039 mmol) in CH₂Cl₂ (1.5 ml) at 0 °C was added lutidine (0.023 mL, 0.20 mmol). This was followed by addition of *tert*-butyldimethylsilyl trifluoromethanesulfonate (0.045 mL, 0.20 mmol) over 2hr. The reaction mixture was poured into saturated NaHCO₃ (2 ml) and extracted with CH₂Cl₂ (3 x 2 ml). The combined organic extracts were dried (Na₂SO₄), filtered and concentrated. The residue was chromatographed with 20% EtOAc/hexanes to give compound C (19 mg, 68%) as a white solid. (M+H)⁺ 719

10 **D. 1S-[1R*,3R*(E),7R*,10S*,11R*,12R*,16S*]]-17-Dichloro-7,11-bis(*t*-butyldimethylsilyloxy)-8,8,10,12,16-pentamethyl-3-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-4-azabicyclo[14.1.0]heptadecane-5,9-dione**

15 To a solution of compound C (9.6 mg, 0.013 mmol) in CHCl₃ (0.10 ml) was added benzyltriethylammonium chloride (0.00026 mmol, <1mg), EtOH (0.002 ml), and 50% NaOH(aq) (0.015 ml, 0.29 mmol). The reaction mixture was vigorously stirred for 2 hr and then directly chromatographed (15% EtOAc/hexanes) to give compound D (2.4 mg, 22%) as a clear oil. (M+H)⁺ 801

20 **E. 1S-[1R*,3R*(E),7R*,10S*,11R*,12R*,16S*]]-17-Dichloro-7,11-dihydroxy-8,8,10,12,16-pentamethyl-3-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-4-azabicyclo[14.1.0]heptadecane-5,9-dione**

25 To compound D (2.2 mg, 0.0027 mmol) at -15 °C was added 20% trifluoroacetic acid/CH₂Cl₂ (0.20 ml). The reaction mixture was warmed to 0°C and stirred for 2 hr. The volatiles were removed under a stream of air and the residue was chromatographed with 60% EtOAc/hexanes to afford the title compound (1.2 mg, 75%) as a white solid. (M+H)⁺ 573

Example 15

30 The following compounds were prepared by the methods described above:

[1S-[1R*,3R*(E),7R*,10S*,11R*,12R*,16S*]]-7,11-Dihydroxy-8,8,10,12-tetramethyl-3-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-5,9-dioxo-4-oxa-17-

- azabicyclo[14.1.0]heptadecane-17-carboxylic acid phenyl ester, (M+H)⁺ 613;
- 5 [1S-[1R*,3R*(E),7R*,10S*,11R*,12R*,16S*]]-7,11-Dihydroxy-8,8,10,12-tetramethyl-3-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-5,9-dioxo-4-oxa-17-azabicyclo[14.1.0]heptadecane-17-carboxylic acid phenylmethyl ester, (M+H)⁺ 627;
- 10 [1S-[1R*,3R*(E),7R*,10S*,11R*,12R*,16S*]]-7,11-Dihydroxy-N,N,8,8,10,12-hexamethyl-3-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-5,9-dioxo-4-oxa-17-azabicyclo[14.1.0]heptadecane-17-sulfonamide, (M+H)⁺ 600;
- 15 [1S-[1R*,3R*(E),7R*,10S*,11R*,12R*,16S*]]-7,11-Dihydroxy-8,8,10,12-tetramethyl-3-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-17-(2-thienylsulfonyl)-4-oxa-17-azabicyclo[14.1.0]heptadecane-5,9-dione, (M+H)⁺ 639;
- 20 [1S-[1R*,3R*(E),7R*,10S*,11R*,12R*,16S*]]-7,11-Dihydroxy-8,8,10,12-tetramethyl-17-[(1-methyl-1H-imidazol-4-yl)sulfonyl]-3-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-4-oxa-17-azabicyclo[14.1.0]heptadecane-5,9-dione, (M+H)⁺ 637;
- 25 [1S-[1R*,3R*(E),7R*,10S*,11R*,12R*,16S*]]-7,11-Dihydroxy-8,8,10,12-tetramethyl-3-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-17-(propylsulfonyl)-4-oxa-17-azabicyclo[14.1.0]heptadecane-5,9-dione, (M+H)⁺ 599;
- 30 [1S-[1R*,3R*(E),7R*,10S*,11R*,12R*,16S*]]-17-Acetyl-7,11-dihydroxy-8,8,10,12-tetramethyl-3-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-4-oxa-17-azabicyclo[14.1.0]heptadecane-5,9-dione, (M+H)⁺ 535;
- [1S-[1R*,3R*(E),7R*,10S*,11R*,12R*,16S*]]-17-Benzoyl-7,11-dihydroxy-8,8,10,12-tetramethyl-3-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-4-oxa-17-azabicyclo[14.1.0]heptadecane-5,9-dione, (M+H)⁺ 597;
- 35 [1S-[1R*,3R*(E),7R*,10S*,11R*,12R*,16S*]]-7,11-Dihydroxy-8,8,10,12-tetramethyl-3-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-17-(2-thienylcarbonyl)-4-oxa-17-azabicyclo[14.1.0]heptadecane-5,9-dione, (M+H)⁺ 603;
- 40 [1S-[1R*,3R*(E),7R*,10S*,11R*,12R*,16S*]]-7,11-Dihydroxy-17-(methoxyacetyl)-8,8,10,12-tetramethyl-3-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-4-oxa-17-azabicyclo[14.1.0]heptadecane-5,9-dione, (M+H)⁺ 565;

[1S-[1R*,3R*(E),7R*,10S*,11R*,12R*,16S*]]-17-(2-Oxopropanoyl)-7,11-dihydroxy-8,8,10,12-tetramethyl-3-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-4-oxa-17-azabicyclo[14.1.0]heptadecane-5,9-dione, (M+H)⁺ 561; and

5

[1S-[1R*,3R*(E),7R*,10S*,11R*,12R*,16S*]]-7,11-Dihydroxy-8,8,10,12-tetramethyl-3-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-5,9-dioxo-4-oxa-17-azabicyclo[14.1.0]heptadecane-17-carboxylic acid ethyl ester, (M+H)⁺ 565.


10

15


What is Claimed.

1. A compound of the formula

5



or




I

II


wherein Q is

10



G is

15



W is O or NH;

X is O, S, CHR₁₇ or H, R₁₈;

5 Y is selected from the group consisting of O; H, H; H, OR₂₂; OR₂₃,
OR₂₃; =NOR₂₄; H, NOR₂₅; H, NR₂₆R₂₇; NHNR₂₈R₂₉; H, NHNR₃₀R₃₁;
and CHR₃₂; where OR₂₃, OR₂₃ can be a cyclic ketal;

B₁ and B₂ are selected from the group consisting of H, OR₃₃,
OCOR₃₄, OCONR₃₅R₃₆, NR₃₇R₃₈ and NR₃₉CONR₄₀R₄₁;

10

M is selected from the group consisting of S, C=O, S=O, SO₂, NR₄₄
and CR₄₅R₄₆;

15 R₁, R₂, R₃, and R₄ are H or lower alkyl;

R₅, R₈, and R₉ are selected from the group consisting of
H, alkyl, substituted alkyl, aryl, substituted aryl, cycloalkyl, heterocyclo and
substituted heterocyclo;

20

R_{17} , R_{18} , R_{22} and R_{23} are selected from the group consisting of H, alkyl, and substituted alkyl;

R_{24} , R_{25} , R_{26} , R_{28} , R_{30} , R_{32} , R_{33} , R_{34} , R_{35} , R_{36} , R_{37} ,

R_{39} , R_{40} , R_{41} ,

R_{51} , R_{52} , R_{53} ,

R_{56} , R_{57} , R_{58} ,

5 R_{59} and R_{61} are selected from the group of H, alkyl, substituted alkyl, aryl and substituted aryl;


R_{27} , R_{29} , R_{31} , R_{38} ,

R_{44} ,

and R_{60}

are selected from the group consisting of H, alkyl, substituted alkyl, substituted aryl, cycloalkyl, heterocyclo, $R_{51}C=O$, $R_{52}OC=O$, $R_{53}SO_2$,

10 hydroxy, O-alkyl and O-substituted alkyl; when X is O then R_{16} is not $R_{51}C=O$, $R_{52}OC=O$, and $R_{53}SO_2$; and wherein R_{44} is further amino or



15

R_{45} and R_{46} are selected from the group consisting of H, halogen, alkyl, substituted alkyl, aryl, substituted aryl, cycloalkyl, heterocyclo,

$R_{56}C=O$, $R_{57}OC=O$, $R_{58}NHC=O$, hydroxy, O-alkyl and O-substituted alkyl,

$NR_{59}R_{60}$:

wherein alkyl is a straight or branched chain unsubstituted hydrocarbon group of 1 to 20 carbon atoms;

aryl is a monocyclic or bicyclic aromatic hydrocarbon group having 6 to 12 carbon atoms in the ring portion;

5 heterocyclo is an optionally substituted, fully saturated or unsaturated, aromatic or non-aromatic cyclic group;

a substituted alkyl or aryl is substituted by one to four substituents;

and any salts, solvates or hydrates thereof, with the proviso that

when Q is



and

M is $CR_{45}R_{46}$ then W is only NH.

10

15

2. The compound of claim 1 wherein

X is O or S

and Y is O.

3. A compound selected from the group consisting of:

5 [1S-[1R*,3R*(E),7R*,10S*,11R*,12R*,16R*]]-7,11-Dihydroxy-8,8,10,12,16-pentamethyl-3-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-4-oxabicyclo[14.1.0]heptadecane-5,9-dione;

10 [1S-[1R*,3R*(E),7R*,10S*,11R*,12R*,16S*]]-17-Dichloro-7,11-dihydroxy-8,8,10,12,16-pentamethyl-3-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-4-oxabicyclo[14.1.0]heptadecane-5,9-dione;

15 [1S-[1R*,3R*(E),7R*,10S*,11R*,12R*,16S*]]-17-Dibromo-7,11-dihydroxy-8,8,10,12,16-pentamethyl-3-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-4-oxabicyclo[14.1.0]heptadecane-5,9-dione;

20 [1S-[1R*,3R*(E),7R*,10S*,11R*,12R*,16R*]]-7,11-Dihydroxy-8,8,10,12-tetramethyl-3-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-4-oxabicyclo[14.1.0]heptadecane-5,9-dione;

[1S-[1R*,3R*(E),7R*,10S*,11R*,12R*,16S*]]-17-Dichloro-7,11-dihydroxy-8,8,10,12,16-pentamethyl-3-[1-methyl-2-(2-methyl-3-oxo-4-thiazolyl)ethenyl]-4-oxabicyclo[14.1.0]heptadecane-5,9-dione;

[1S-[1R*,3R*(E),7R*,10S*,11R*,12R*,16R*]]-7,11-Dihydroxy-8,8,10,12,16-pentamethyl-3-[1-methyl-2-(2-methyl-3-oxo-4-thiazolyl)ethenyl]-4-oxabicyclo[14.1.0]heptadecane-5,9-dione;

- 5 [1S-[1R*,3R*(E),7R*,10S*,11R*,12R*,16S*]]-7,11-Dihydroxy-8,8,10,12-tetramethyl-3-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-5,9-dioxo-4-oxa-17-azabicyclo[14.1.0]heptadecane-17-carboxylic acid, 1,1-dimethylethyl ester;

10

[1S-[1R*,3R*(E),7R*,10S*,11R*,12R*,16R*]]-7,11-Dihydroxy-8,8,10,12,16-pentamethyl-3-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-4-

- 15 azabicyclo[14.1.0]heptadecane-5,9-dione;

[1S-[1R*,3R*(E),7R*,10S*,11R*,12R*,16S*]]-17-Dichloro-7,11-dihydroxy-8,8,10,12,16-pentamethyl-3-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-4-azabicyclo[14.1.0]heptadecane-5,9-dione;

- 20 [1S-[1R*,3R*(E),7R*,10S*,11R*,12R*,16S*]]-7,11-Dihydroxy-8,8,10,12-tetramethyl-3-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-5,9-dioxo-4-oxa-17-azabicyclo[14.1.0]heptadecane-17-carboxylic acid phenyl ester;

- 25 [1S-[1R*,3R*(E),7R*,10S*,11R*,12R*,16S*]]-7,11-Dihydroxy-8,8,10,12-tetramethyl-3-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-5,9-dioxo-4-oxa-17-azabicyclo[14.1.0]heptadecane-17-carboxylic acid phenylmethyl ester;

- [1S-[1R*,3R*(E),7R*,10S*,11R*,12R*,16S*]]-7,11-Dihydroxy-N,N,8,8,10,12-hexamethyl-3-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-5,9-dioxo-4-oxa-17-azabicyclo[14.1.0]heptadecane-17-sulfonamide;
- 5 [1S-[1R*,3R*(E),7R*,10S*,11R*,12R*,16S*]]-7,11-Dihydroxy-8,8,10,12-tetramethyl-3-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-17-(2-thienylsulfonyl)-4-oxa-17-azabicyclo[14.1.0]heptadecane-5,9-dione;
- 10 [1S-[1R*,3R*(E),7R*,10S*,11R*,12R*,16S*]]-7,11-Dihydroxy-8,8,10,12-tetramethyl-17-[(1-methyl-1H-imidazol-4-yl)sulfonyl]-3-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-4-oxa-17-azabicyclo[14.1.0]heptadecane-5,9-dione;
- 15 [1S-[1R*,3R*(E),7R*,10S*,11R*,12R*,16S*]]-7,11-Dihydroxy-8,8,10,12-tetramethyl-3-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-17-(propylsulfonyl)-4-oxa-17-azabicyclo[14.1.0]heptadecane-5,9-dione;
- 20 [1S-[1R*,3R*(E),7R*,10S*,11R*,12R*,16S*]]-17-Acetyl-7,11-dihydroxy-8,8,10,12-tetramethyl-3-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-4-oxa-17-azabicyclo[14.1.0]heptadecane-5,9-dione;
- [1S-[1R*,3R*(E),7R*,10S*,11R*,12R*,16S*]]-17-Benzoyl-7,11-dihydroxy-8,8,10,12-tetramethyl-3-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-4-oxa-17-azabicyclo[14.1.0]heptadecane-5,9-dione;
- 25 [1S-[1R*,3R*(E),7R*,10S*,11R*,12R*,16S*]]-7,11-Dihydroxy-8,8,10,12-tetramethyl-3-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-17-(2-thienylcarbonyl)-4-oxa-17-azabicyclo[14.1.0]heptadecane-5,9-dione;
- 30 [1S-[1R*,3R*(E),7R*,10S*,11R*,12R*,16S*]]-7,11-Dihydroxy-17-(methoxyacetyl)-8,8,10,12-tetramethyl-3-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-4-oxa-17-azabicyclo[14.1.0]heptadecane-5,9-dione;
- 35 [1S-[1R*,3R*(E),7R*,10S*,11R*,12R*,16S*]]-17-(2-Oxopropanoyl)-7,11-dihydroxy-8,8,10,12-tetramethyl-3-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-4-oxa-17-azabicyclo[14.1.0]heptadecane-5,9-dione; and
- 40 [1S-[1R*,3R*(E),7R*,10S*,11R*,12R*,16S*]]-7,11-Dihydroxy-8,8,10,12-tetramethyl-3-[1-methyl-2-(2-methyl-4-thiazolyl)ethenyl]-5,9-dioxo-4-oxa-17-azabicyclo[14.1.0]heptadecane-17-carboxylic acid ethyl ester.

4. **Use of a pharmaceutically effective amount of a compound of the formula I or II as
claimed in claim 1 for treating cancer.**
5. **Use of a pharmaceutically effective amount of a compound of the formula I or II as
5 claimed in claim 1 in the manufacture of a medicament for treating cancer.**