



Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2875478 A1 2005/02/24

(21) **2 875 478**

(12) **DEMANDE DE BREVET CANADIEN
CANADIAN PATENT APPLICATION**

(13) **A1**

(22) **Date de dépôt/Filing Date:** 2004/05/11

(41) **Mise à la disp. pub./Open to Public Insp.:** 2005/02/24

(62) **Demande originale/Original Application:** 2 524 194

(30) **Priorités/Priorities:** 2003/05/13 (US60/469,997);
2003/10/31 (US60/516,100)

(51) **Cl.Int./Int.Cl. C12Q 1/68** (2006.01),
C07H 21/04 (2006.01), **C12N 15/31** (2006.01)

(71) **Demandeur/Applicant:**
GEN-PROBE INCORPORATED, US

(72) **Inventeurs/Inventors:**
HOGAN, JAMES J., US;
KAPLAN, SHANNON K., US


(74) **Agent:** SMART & BIGGAR

(54) **Titre : PROCEDE ET TROUSSE PERMETTANT L'IDENTIFICATION DE MICRO-ORGANISMES RESISTANTS AUX
ANTIBIOTIQUES**

(54) **Title: METHOD AND KIT FOR IDENTIFYING ANTIBIOTIC-RESISTANT MICROORGANISMS**

(57) **Abrégé/Abstract:**

The invention provides a rapid sample-processing method for preparing hybridization reaction mixtures substantially depleted of RNA, and a method of identifying the methicillin-resistance status and vancomycin-resistance status of an organism.



ABSTRACT

The invention provides a rapid sample-processing method for preparing hybridization reaction mixtures substantially depleted of RNA, and a method of identifying the methicillin-resistance status and vancomycin-resistance status of an organism.

DEMANDES OU BREVETS VOLUMINEUX

**LA PRÉSENTE PARTIE DE CETTE DEMANDE OU CE BREVETS
COMPREND PLUS D'UN TOME.**

CECI EST LE TOME _____ DE _____

NOTE: Pour les tomes additionels, veuillez contacter le Bureau Canadien des Brevets.

JUMBO APPLICATIONS / PATENTS

**THIS SECTION OF THE APPLICATION / PATENT CONTAINS MORE
THAN ONE VOLUME.**

THIS IS VOLUME 1 OF 2

NOTE: For additional volumes please contact the Canadian Patent Office.

METHOD AND KIT FOR IDENTIFYING ANTIBIOTIC-RESISTANT MICROORGANISMS

Field of the Invention

5 The present invention relates to the field of nucleic acid diagnostics. More particularly, the invention relates to methods and compositions for detecting and identifying methicillin-resistant and/or vancomycin-resistant microorganisms.

Background of the Invention

10 Procedures for detecting and identifying infectious organisms are some of the most critical tasks performed in the clinical laboratory. Whereas laboratory diagnoses of infectious diseases formerly were made by experienced microbiologists using visual inspection of stained clinical material, more rapid and objective results are obtainable using modern techniques. Immunoassays, including radioimmunoassays, enzyme-linked immunoassays, and latex agglutination and immunoblotting assays have developed into powerful diagnostic tools having
15 utilities that are enhanced by the availability of monoclonal antibodies. Nucleic acid hybridization assays have been developed to detect microorganisms, and more recent advances in signal and target amplification have introduced the era of molecular diagnostics based on the use of oligonucleotide probes. Generally, a probe is a single-stranded polynucleotide having some degree of complementarity with a nucleic acid sequence that is to be detected ("target
20 sequence"). A double-stranded nucleic acid hybrid between the probe and the target sequence results if the target sequence is contacted under hybridization-promoting conditions with a probe having a sufficient number of contiguous bases complementary to the target sequence. DNA/DNA, RNA/DNA or RNA/RNA hybrids may thus be formed under appropriate conditions. Probes commonly are labeled with a detectable moiety such as a radioisotope, a
25 ligand, or a colorimetric, fluorometric or chemiluminescent moiety to facilitate the detection of hybrids.

 Indeed, clinical microbiologists now use an extensive array of techniques for identifying infectious organisms (see Manual of Clinical Microbiology Murray et al., eds., 6th edition, ASM Press (1995)). Automated substrate utilization systems typically rely on enzymatic
30 reactions that release chromogenic or fluorogenic compounds, tetrazolium-based indicators of

metabolic activity in the presence of different carbon sources, or detection of the acid products of metabolism. The patterns of positive and negative reactions with these substrates establish a biochemical profile that can be used to identify microorganisms isolated from clinical samples. The chromatographic profiles of the more than 300 fatty acids that contribute to the formation of lipids in bacteria and yeast have also been used to phenotype microorganisms. Despite the availability of these very powerful techniques, polynucleotide-based assays are rapidly gaining popularity in clinical laboratory practice.

The specificity of polynucleotide hybridization reactions, together with the extraordinary sensitivity afforded by nucleic acid amplification techniques, has made molecular diagnostics the method of choice for detecting and identifying microbes that are available in only very small quantities. Commonly used DNA probe hybridization formats include: solid phase hybridization, solution-phase hybridization and *in situ* hybridization. In solid phase hybridization methods, a sample containing microbial polynucleotides is immobilized to a solid support, denatured and then probed with a polynucleotide probe that harbors a detectable label. Unhybridized probe is removed from the system and specifically hybridized probe detected, for example, by autoradiography or direct visual observation. In solution-phase hybridization procedures, the target polynucleotide and the labeled probe are free to interact in an aqueous hybridization buffer. Specifically hybridized probe is then detected as an indicator of the presence of target polynucleotides in the mixture. *In situ* hybridization using formalin-fixed tissue sections is used for obtaining information about the physical distribution and abundance of microorganisms.

One example of an organism for which a number of polynucleotide-based assays have been described is *Staphylococcus aureus*. For example, Milliman in U.S. Patent No. 5,292,874 describes a hybridization-based assay to distinguish *S. aureus* from other *Staphylococcus spp.* which employs probes specific for 23S rRNA. The detection of *S. aureus* in biological samples is important as within the genus *Staphylococcus*, *S. aureus* is the most clinically significant species due to the incidence and severity of the infections it can cause (Morse, *Staphylococci*, in Medical Microbiology and Infectious Diseases, Abraham Braude, editor, W. B. Saunders Company, Philadelphia, Pa., 1981). Moreover, *S. aureus* is a prominent agent of nosocomial

infections, and methicillin resistant strains (MRSA) have emerged as a major epidemiological problem in hospitals throughout the United States.

Vancomycin resistant enterococci (VRE) represent another emerging class of drug resistant bacteria. Since these organisms were first identified in 1986, nearly 30 years after vancomycin was clinically introduced, it has been established that vancomycin resistance is primarily conferred by either of two functionally similar operons, VanA and VanB. These operons, transfer of which can be mediated by plasmids or transposons, are complex resistance determinants that may have evolved in other species and then been acquired by enterococci. Frequently identified risk factors for VRE colonization and infection include prolonged hospital stays, exposure to intensive care units, transplants, hematologic malignancies, and exposure to antibiotics. Notably, more than 95% of VRE recovered in the United States are *Enterococcus faecium*, and virtually all are resistant to high levels of ampicillin. (Rice, L., *Emerging Infectious Diseases* 7:183 (2001))

Because vancomycin resistance is transferrable by genetic means, there is the possibility that resistance to this important antibiotic can be acquired by other microorganisms, such as *Staphylococcus aureus*. Indeed, conjugative transfer of the VanA gene from *Enterococcus faecalis* to *S. aureus* has already been demonstrated *in vitro*. It has been speculated that such a transfer mechanism may underlie the appearance of vancomycin resistant *S. aureus* (VRSA). (Noble et al., *FEMS Microbiol Lett* 93:195 (1992)) Conceivably, there could emerge a strain of *S. aureus* which has acquired resistance to both methicillin and vancomycin (MVSA).

Accordingly, there is a continuing need for the rapid processing of clinical or biological samples, and for the rapid and accurate detection of pathogens and antibiotic resistance genes in clinical samples.

Summary

A first aspect of this disclosure relates to a device for detecting nucleic acids encoding resistance to an antibiotic. The invented device includes a solid support and a plurality of detectably labeled hybridization probes distributed among a plurality of loci on the solid support. The plurality of loci include a first locus that includes one or more probes that collectively hybridize to ribosomal nucleic acids from a plurality of species of bacteria in the genus *Staphylococcus*, including *Staphylococcus aureus* and *Staphylococcus epidermidis*, but

do not hybridize to ribosomal nucleic acids of bacteria in the genus *Enterococcus*. The device further includes a second locus that has one or more probes that collectively hybridize to ribosomal nucleic acids from a plurality of bacteria in the genus *Enterococcus*, *Enterococcus faecalis* and *Enterococcus faecium*, but that do not hybridize to *Staphylococcus aureus* or any other bacteria in the genus *Staphylococcus*. The second locus also has at least one probe that hybridizes to ribosomal nucleic acids from *Staphylococcus aureus* but not from other species in the genus *Staphylococcus* or bacteria in the genus *Enterococcus*. The device further includes a third locus having at least one probe that hybridizes to *mecA* nucleic acids. The device further includes a fourth locus that has one or more probes that collectively hybridize to VanA nucleic acids and to VanB nucleic acids. In one embodiment, each of the plurality of detectably labeled hybridization probes is a detectably labeled soluble hybridization probe. In another embodiment, each of the detectably labeled hybridization probes is labeled with a homogeneously detectable label. For example, the homogeneously detectable label may be a chemiluminescent label. In yet another embodiment, the device further includes a fifth locus that includes one or more probes that collectively hybridize to ribosomal nucleic acids from a plurality of species of Gram-positive bacteria (including the high(G+C) subset of Gram-positive bacteria), a plurality of species of bacteria in the family Enterobacteriaceae, a plurality of species of bacteria in the genus *Enterococcus*, and a plurality of species of bacteria in the genus *Staphylococcus*. In still yet another embodiment, the probes that collectively hybridize to VanA nucleic acids and to VanB nucleic acids include at least one probe that hybridizes to VanA nucleic acids and at least one probe that hybridizes to VanB nucleic acids. In a different embodiment, the probes that collectively hybridize to VanA nucleic acids and to VanB nucleic acids include a single probe that hybridizes both to VanA nucleic acids and VanB nucleic acids.

A second aspect of this disclosure also relates to a device for detecting nucleic acids encoding resistance to an antibiotic. The device includes a solid support and a plurality of detectably labeled hybridization probes distributed among a plurality of loci on the solid support. The plurality of loci includes a first locus having one or more probes that collectively hybridize to ribosomal nucleic acids from a plurality of species of bacteria in the genus *Staphylococcus*, including *Staphylococcus aureus* and *Staphylococcus epidermidis*. The device further includes a second locus having a probe that hybridizes to ribosomal nucleic acids from

Staphylococcus aureus but not other species in the genus *Staphylococcus*. The device further includes a third locus that comprises at least one probe that hybridizes to *mecA* nucleic acids. The device further includes a fourth locus that has one or more probes that collectively hybridize to ribosomal nucleic acids from a plurality of species of Gram-positive bacteria (including the high(G+C)subset of Gram-positive bacteria), a plurality of species of bacteria in the family Enterobacteriaceae, a plurality of species of bacteria in the genus Enterococcus, and a plurality of species of bacteria in the genus *Staphylococcus*. In one embodiment, the fourth locus includes a pan-bacterial probe having a base sequence that is either SEQ ID NO:16 or SEQ ID NO:19. In a preferred embodiment, the probes of the third locus have base sequences that are any of SEQ ID NO:2, SEQ ID NO:3 or SEQ ID NO:5.

A third aspect of this disclosure also relates to a device for detecting nucleic acids encoding resistance to an antibiotic. The device includes a solid support and a plurality of detectably labeled hybridization probes distributed among a plurality of loci on the solid support. The plurality of loci include a first locus having one or more probes that collectively hybridize to ribosomal nucleic acids from a plurality of bacteria in the genus Enterococcus, including *Enterococcus faecalis* and *Enterococcus faecium*. The device further includes a second locus having one or more probes that collectively hybridize to VanA nucleic acids and to VanB nucleic acids. The device further includes a third locus having one or more probes that collectively hybridize to ribosomal nucleic acids from a plurality of species of Gram-positive bacteria (including the high(G+C)subset of Gram-positive bacteria), a plurality of species of bacteria in the family Enterobacteriaceae, a plurality of species of bacteria in the genus Enterococcus, and a plurality of species of bacteria in the genus *Staphylococcus*. In one embodiment, the third locus includes a pan-bacterial probe having a base sequence that is either SEQ ID NO:16 or SEQ ID NO:19. In a preferred embodiment, the probes of the second locus have sequences selected from the group consisting of SEQ ID NO:27, SEQ ID NO:28 and SEQ ID NO:29. In another preferred embodiment, the probes of the second locus have sequences selected from the group consisting of SEQ ID NO:30, SEQ ID NO:40 and SEQ ID NO:41.

A fourth aspect of this disclosure relates to a method of detecting the presence of a target DNA sequence in a sample containing cells. The method includes a step for lysing cells contained in the sample to yield a lysate. There also is included a step for treating the lysate

with a basic composition to yield a basic mixture. This is followed by introducing one or more reagents to the basic mixture to yield a reaction mixture at a pH in the range of 4.5 - 5.5. Next, there is a step for hybridizing the reaction mixture under high stringency hybridization conditions with an oligonucleotide probe to form a hybrid between the oligonucleotide probe and the target DNA. Finally, there is a step for detecting the hybrid as an indicator of the presence of the target DNA sequence. In one embodiment, the oligonucleotide probe in the hybridizing step is complementary either to a *mecA* gene sequence, a VanA gene sequence or a VanB gene sequence. In a different embodiment, the treating step involves heating the basic mixture at a temperature up to 100°C for a time sufficient to substantially hydrolyze RNA that may be present in the lysate.

A fifth aspect of this disclosure relates to a probe mix for hybridizing *mecA* nucleic acids. The probe mix includes a first detectably labeled hybridization probe having the sequence of SEQ ID NO:2, a second detectably labeled hybridization probe having the sequence of SEQ ID NO:3, and a third detectably labeled hybridization probe having the sequence of SEQ ID NO:5. More preferably, the first detectably labeled hybridization probe has the length and sequence of SEQ ID NO:2 or the complement thereof, and optionally a non-complementary sequence that does not hybridize to the nucleic acids of the *mecA* gene; the second detectably labeled hybridization probe has the length and sequence of SEQ ID NO:3 or the complement thereof, and optionally a non-complementary sequence that does not hybridize to the nucleic acids of the *mecA* gene; and the third detectably labeled hybridization probe has the length and sequence of SEQ ID NO:5 or the complement thereof, and optionally a non-complementary sequence that does not hybridize to the nucleic acids of the *mecA* gene.

A sixth aspect of this disclosure relates to a probe mix for detecting VanA nucleic acids. The invented probe mix includes a first detectably labeled hybridization probe having the sequence of SEQ ID NO:27 or the complement thereof; a second detectably labeled hybridization probe having the sequence of SEQ ID NO:28 or the complement thereof; and a third detectably labeled hybridization probe having the sequence of SEQ ID NO:29 or the complement thereof. More preferably, the first detectably labeled hybridization probe has the length and sequence of SEQ ID NO:27 or the complement thereof, and optionally a non-complementary sequence that does not hybridize to the nucleic acids of the VanA gene; the

second detectably labeled hybridization probe has the length and sequence of SEQ ID NO:28 or the complement thereof, and optionally a non-complementary sequence that does not hybridize to the nucleic acids of the VanA gene; and the third detectably labeled hybridization probe has the length and sequence of SEQ ID NO:29 or the complement thereof, and optionally a non-complementary sequence that does not hybridize to the nucleic acids of the VanA gene.

A seventh aspect of this disclosure relates to a probe mix for detecting VanB nucleic acids. The invented probe mix includes a first detectably labeled hybridization probe having the sequence of SEQ ID NO:30 or the complement thereof; a second detectably labeled hybridization probe having the sequence of SEQ ID NO:40 or the complement thereof; and a third detectably labeled hybridization probe having the sequence of SEQ ID NO:41 or the complement thereof. More preferably, the first detectably labeled hybridization probe has the length and sequence of SEQ ID NO:30 or the complement thereof, and optionally a non-complementary sequence that does not hybridize to the nucleic acids of the VanB gene; the second detectably labeled hybridization probe has the length and sequence of SEQ ID NO:40 or the complement thereof, and optionally a non-complementary sequence that does not hybridize to the nucleic acids of the VanB gene; and the third detectably labeled hybridization probe has the length and sequence of SEQ ID NO:41 or the complement thereof, and optionally a non-complementary sequence that does not hybridize to the nucleic acids of the VanB gene.

Various embodiments of the claimed invention relate to a method for detecting the presence or absence of at least a first target nucleic acid and a second target nucleic acid in a sample, wherein the first target nucleic acid is a nucleic acid encoding resistance to an antibiotic and the second target nucleic acid is specific for one or more bacterial species, said method comprising: providing a sample suspected of containing the first and second target nucleic acids; contacting the sample with (i) a first oligonucleotide probe that hybridizes to the first target nucleic acid to form a first hybrid between the first oligonucleotide probe and the first target nucleic acid, and (ii) a second oligonucleotide probe that hybridizes to the second target nucleic acid to form a second hybrid between the second oligonucleotide probe and the second target nucleic acid; wherein the first and second oligonucleotide probes are respectively distributed at first and second loci on a solid support; and detecting the presence or absence of

the first and second hybrids as indicators of the presence or absence of the first and second target nucleic acids.

Various embodiments of the claimed invention relate to a method for detecting the presence or absence of a target nucleic acid encoding resistance to an antibiotic and a target
5 nucleic acid from a bacterial species, said method comprising: providing a sample suspected of containing at least one bacterial species; contacting the sample with oligonucleotide probes distributed at a plurality of loci on a solid support, wherein the oligonucleotide probes comprise (i) at a first locus, one or more probes that collectively hybridize to ribosomal nucleic acids from a plurality of species of bacteria in the genus *Staphylococcus*, comprising *Staphylococcus*
10 *aureus* and *Staphylococcus epidermis*, but not to ribosomal nucleic acids of bacteria in the genus *Enterococcus*, to form at least a first hybrid; (ii) at a second locus, (a) one or more oligonucleotide probes that collectively hybridize to ribosomal nucleic acids from a plurality of bacteria in the genus *Enterococcus*, comprising *Enterococcus faecalis* and *Enterococcus faecium*, but not *Staphylococcus aureus* or any other bacteria in the genus *Staphylococcus*, and
15 (b) at least one probe that hybridizes to a ribosomal nucleic acid from *Staphylococcus aureus* but not from other species in the genus *Staphylococcus* or bacteria in the genus *Enterococcus*, to form at least a second hybrid; (iii) at a third locus, a least one oligonucleotide probe that hybridizes to a *mecA* nucleic acid; and (iv) at a fourth locus, one or more oligonucleotide probes that collectively hybridize to VanA nucleic acids and to VanB nucleic acids; detecting
20 the presence or absence of the at least first, second, third, and fourth hybrids as indicators of the presence or absence of the an antibiotic resistance gene and a bacterial species in the sample.

Various embodiments of the claimed invention relate to a method for detecting the presence or absence of a target nucleic acid encoding resistance to an antibiotic and a target nucleic acid from a bacterial species, said method comprising: providing a sample suspected of
25 containing at least one bacterial species; contacting the sample with oligonucleotide probes distributed at a plurality of loci on a solid support, wherein the oligonucleotide probes comprise (i) at a first locus, one or more oligonucleotide probes that collectively hybridize to ribosomal nucleic acids from a plurality of species of bacteria in the genus *Staphylococcus*, comprising *Staphylococcus aureus* and *Staphylococcus epidermidis*, to form at least a first hybrid; (ii) at a
30 second locus, at least one oligonucleotide probe that hybridizes to a ribosomal nucleic acid

from *Staphylococcus aureus* but not other species in the genus *Staphylococcus*, to form at least a second hybrid; (iii) at a third locus, a least one oligonucleotide probe that hybridizes to a *mecA* nucleic acid, to form at least a third hybrid; and (iv) at a fourth locus, one or more oligonucleotide probes that collectively hybridize to ribosomal nucleic acids from a plurality of species of Gram positive bacteria comprising the high(G+C) subset thereof, a plurality of species of bacteria in the family Enterobacteriaceae, a plurality of species of bacteria in the genus Enterococcus, and a plurality of species of bacteria in the genus *Staphylococcus*, to form at least a fourth hybrid; and detecting the presence or absence of the at least first, second, third, and fourth hybrids as indicators of the presence or absence of the an antibiotic resistance gene and a bacterial species in the sample.

Various embodiments of the claimed invention relate to a method for detecting the presence or absence of a target nucleic acid encoding resistance to an antibiotic and a target nucleic acid from a bacterial species, said method comprising: providing a sample suspected of containing at least one bacterial species; contacting the sample with oligonucleotide probes distributed at a plurality of loci on a solid support, wherein the oligonucleotide probes comprise (i) at a first locus, one or more oligonucleotide probes that collectively hybridize to ribosomal nucleic acids from a plurality of bacteria in the genus Enterococcus, comprising Enterococcus faecalis and Enterococcus faecium, to form at least a first hybrid; (ii) at a second locus, one or more oligonucleotide probes that collectively hybridize to VanA nucleic acids and to VanB nucleic acids, to form at least a second hybrid; and (iii) at a third locus, one or more oligonucleotide probes that collectively hybridize to ribosomal nucleic acids from a plurality of species of Gram-positive bacteria comprising the high(G+C) subset thereof, a plurality of species of bacteria in the family Enterobacteriaceae, a plurality of species of bacteria in the genus Enterococcus, and a plurality of species of bacteria in the genus *Staphylococcus*, to form at least a third hybrid; and detecting the presence or absence of the at least first, second, and third hybrids as indicators of the presence or absence of the an antibiotic resistance gene and a bacterial species in the sample.

Various embodiments of the claimed invention relate to a method of detecting the presence of a target DNA sequence in a sample containing cells, comprising the steps of: lysing cells contained in the sample to yield a lysate; treating the lysate with a basic

composition to yield a basic mixture; introducing one or more reagents to the basic mixture to yield a reaction mixture at a pH in the range of 4.5 - 5.5; hybridizing the reaction mixture with an oligonucleotide probe to form a hybrid between the oligonucleotide probe and the target DNA; and detecting said hybrid as an indicator of the presence of said target DNA sequence, wherein said oligonucleotide probe is complementary either to a *mecA* gene sequence, a *VanA* gene sequence or a *VanB* gene sequence.

Various embodiments of the claimed invention relate to a method for detecting the presence or absence of a target DNA sequence in a sample containing cells, said method comprising: (a) treating the sample to yield a lysate, wherein the treating includes: (i) treating the lysate with a basic composition to yield a basic mixture; and (ii) introducing one or more reagents to the basic mixture to yield a reaction mixture at a pH in the range of 4.5 to 5.5; (b) hybridizing the reaction mixture with a plurality of oligonucleotide probes distributed at a plurality of loci on a solid support, wherein the solid support is configured to comprise: (i) a first locus containing one or more probes that collectively hybridize to ribosomal nucleic acids from a plurality of species of bacteria in the genus *Staphylococcus*, comprising *Staphylococcus aureus* and *Staphylococcus epidermis*, but not to ribosomal nucleic acids of bacteria in the genus *Enterococcus*, to form a hybrid with a target nucleic acid in the reaction mixture; (ii) a second locus containing, (a) one or more oligonucleotide probes that collectively hybridize to ribosomal nucleic acids from a plurality of bacteria in the genus *Enterococcus*, comprising *Enterococcus faecalis* and *Enterococcus faecium*, but not *Staphylococcus aureus* or any other bacteria in the genus *Staphylococcus*, to form a hybrid with a target nucleic acid in the reaction mixture, and (b) at least one probe that hybridizes to a ribosomal nucleic acid from *Staphylococcus aureus* but not from other species in the genus *Staphylococcus* or bacteria in the genus *Enterococcus*, to form a hybrid with a target nucleic acid in the reaction mixture; (iii) a third locus containing at least one oligonucleotide probe that hybridizes to a *mecA* nucleic acid, to form a hybrid with a target nucleic acid in the reaction mixture; and (iv) a fourth locus containing one or more oligonucleotide probes that collectively hybridize to *VanA* nucleic acids and to *VanB* nucleic acids, to form a hybrid with a target nucleic acid in the reaction mixture; and (c) detecting the presence or absence of a hybrid at each loci as an indicator of the presence or absence of the target DNA, and thereby determining the presence or absence of a

Staphylococcus organism, the presence or absence of an Enterococcus organism and the presence or absence of a drug resistant genotype.

Detailed Description of the Invention

Definitions

5 As used herein, the following terms have the given meanings unless expressly stated to the contrary.

 "Ribosomal nucleic acids" are rRNA and the rDNA that encodes the rRNA.

 A "locus" is a place in which something is situated. A locus can be a reaction tube; a single well for containing soluble polynucleotides in a multi-well plate; a single spot of
10 immobilized polynucleotides on a piece of nitrocellulose membrane or a dipstick; or a single spot of immobilized polynucleotides on a "DNA chip." Probe molecules disposed at one locus in a testing device do not mingle with probe molecules disposed at another locus in the device.

 An "address" refers to one or more polynucleotide probes at a single locus in a testing device, whereby a hybridization result at the address provides discrete information

WO 2005/017202

PCT/US2004/014690

about the presence or absence of complementary polynucleotide sequences among a collection of polynucleotides undergoing testing. For example, an address may provide information about the presence of rRNA that is of bacterial origin or the presence of one or more specific genes, such as a gene encoding a drug resistance phenotype. Such

5 information could be derived from a single probe or a cocktail of probes disposed at a single locus on a testing device. An address also may provide information about the presence of rRNA from one or more species of microorganism, such as any of *E. coli*, *S. aureus*, *C. albicans*, *P. aeruginosa* and *S. pneumoniae*. For convenience, an address that detects nucleic acids of a particular organism or type of organism is referred to by the name

10 or type of that organism. Thus, a positive hybridization signal at a "pan-bacterial" address would indicate the presence of ribosomal nucleic acids that are of bacterial origin.

A probe "matrix" is a collection of addresses useful for identifying an unknown microorganism, for narrowing the range of possible identities of an unknown organism, and/or the presence of one or more specific genes, in an organism. Probes of the matrix are

15 ordinarily disposed (either by containment of soluble probes or by physical immobilization) at a plurality of physical loci in a testing device where each locus specifically hybridizes nucleic acids from one or a plurality of microorganism species.

A "nucleotide" is a subunit of a nucleic acid conventionally comprising a purine or pyrimidine base, a 5-carbon sugar and a phosphate group. The 5-carbon sugar found in

20 RNA is ribose. In DNA, the 5-carbon sugar is 2'-deoxyribose. The term also includes analogs of such subunits, such as a methoxy group at the 2' position of ribose.

A "non-nucleotide unit" is a unit which does not significantly participate in hybridization of a polymer. Such units must not, for example, participate in any significant hydrogen bonding with a nucleotide, and would exclude units having as a component one of

25 the five nucleotide bases or analogs thereof.

An "oligonucleotide" is a polynucleotide having two or more nucleotide subunits covalently joined together. Oligonucleotides are generally about 10 to about 100 nucleotides in length, or more preferably 10 to 50 nucleotides in length. The sugar groups of the nucleotide subunits may be ribose, deoxyribose, or modified derivatives or analogs

30 thereof such as 2'-O-methyl ribose (referred to herein alternatively as "2'-O-Me", "2' methoxy", "2'-MeO"). The nucleotide subunits may be joined by linkages such as phosphodiester linkages, modified linkages or by non-nucleotide moieties that do not

WO 2005/017202

PCT/US2004/014690

prevent hybridization of the oligonucleotide to its complementary target nucleotide sequence. Modified linkages include those in which a standard phosphodiester linkage is replaced with a different linkage, such as a phosphorothioate linkage, a methylphosphonate linkage, or a neutral peptide linkage. Nitrogenous base analogs also may be components of oligonucleotides in accordance with the invention. Ordinarily, oligonucleotides will be synthesized by organic chemical methods and will be single-stranded unless specified otherwise. Oligonucleotides can be labeled with a detectable label.

A "target nucleic acid" is a nucleic acid comprising a target nucleic acid sequence.

A "target nucleic acid sequence," "target nucleotide sequence" or "target sequence" is a specific deoxyribonucleotide or ribonucleotide sequence that can be hybridized by an oligonucleotide. For instance, a "target nucleic acid sequence region" of bacteria in the *Staphylococcus* genus refers to a nucleic acid sequence present in nucleic acid or a sequence complementary thereto found in *Staphylococcal* bacteria, which is not present in nucleic acids of other species. Nucleic acids having nucleotide sequences complementary to a target sequence may be generated by target amplification techniques such as polymerase chain reaction (PCR) or transcription mediated amplification (e.g., see U.S. Patent No. 5,824,518).

A "probe" is a single-stranded polynucleotide that combines with a complementary single-stranded target polynucleotide to form a double-stranded hybrid. A probe may be an oligonucleotide or a nucleotide polymer, and may contain a detectable moiety which can be attached to the end(s) of the probe or can be internal to the sequence of the probe. The nucleotides which combine with the target polynucleotide need not be strictly contiguous as may be the case with a detectable moiety internal to the sequence of the probe.

A "pan-bacterial" probe is capable of forming double-stranded hybrids with complementary sequences that are found in the ribosomal nucleic acids of a plurality of species of bacteria. For example, a pan-bacterial probe can specifically hybridize to ribosomal nucleic acids from a plurality of species of Gram-positive bacteria (including the high(G+C)subset of Gram-positive bacteria known as the "Actinomycetes"), a plurality of species of bacteria in the family Enterobacteriaceae, a plurality of species of bacteria in the genus *Enterococcus*, and a plurality of species of bacteria in the genus *Staphylococcus*. A pan-bacterial probe does not hybridize to the nucleic acids of fungal organisms.

A "detectable moiety" is a label molecule attached to, or synthesized as part of, a

WO 2005/017202

PCT/US2004/014690

polynucleotide probe. This molecule should be uniquely detectable and will allow the probe to be detected as a result. These detectable moieties include but are not limited to radioisotopes, colorimetric, fluorometric or chemiluminescent molecules, enzymes, haptens, redox-active electron transfer moieties such as transition metal complexes, metal
5 labels such as silver or gold particles, or even unique oligonucleotide sequences.

A "hybrid" is the complex formed between two single-stranded polynucleotide sequences by Watson-Crick base pairings or non-canonical base pairings between the complementary bases. By "nucleic acid hybrid" or "probe:target duplex" is meant a structure that is a double-stranded, hydrogen-bonded structure, preferably 10 to 100
10 nucleotides in length, more preferably 14 to 50 nucleotides in length. The structure is sufficiently stable to be detected by means such as chemiluminescent or fluorescent light detection, colorimetry, autoradiography, electrochemical analysis or gel electrophoresis. Such hybrids include RNA:RNA, RNA:DNA, or DNA:DNA duplex molecules.

"Hybridization" is the process by which two complementary strands of
15 polynucleotide combine to form a stable double-stranded structure ("hybrid").

"Stable" means substantially resistant to chemical or biochemical degradation, reaction, decomposition, displacement or modification.

"Stability" means the resistance of a substance to chemical or biochemical degradation, reaction, decomposition, displacement or modification.

20 The term "stringency" is used to describe the temperature and solvent composition existing during hybridization and the subsequent processing steps. Under high stringency conditions only highly complementary nucleic acid hybrids will form; hybrids without a sufficient degree of complementarity will not form. Accordingly, the stringency of the assay conditions determines the amount of complementarity needed between two
25 polynucleotide strands forming a hybrid. Stringency conditions are chosen to maximize the difference in stability between the hybrid formed with the target and the non-target polynucleotide.

The term "probe specificity" refers to a characteristic of a probe which describes its ability to distinguish between target and non-target sequences. Probe specificity is
30 dependent on sequence and assay conditions and may be absolute (i.e., the probe can distinguish between target organisms and any non-target organisms), or it may be functional (i.e., the probe can distinguish between the target organism and any other organism

WO 2005/017202

PCT/US2004/014690

normally present in a particular sample). Many probe sequences can be used for either broad or narrow specificity determinations depending on the conditions of use.

"Polynucleotide" means either RNA or DNA, along with any synthetic nucleotide analogs or other molecules that may be present in the sequence and that do not prevent
5 hybridization of the polynucleotide with a second molecule having a complementary sequence. The term includes polymers containing analogs of naturally occurring nucleotides and particularly includes analogs having a methoxy group at the 2' position of the ribose (MeO).

A "helper oligonucleotide" is an oligonucleotide that binds a region of a target
10 polynucleotide other than the region that is bound by an assay probe. These oligonucleotides impose new secondary and tertiary structures on the targeted region of the single-stranded polynucleotide so that the rate of binding of the assay probe is accelerated. Although helper oligonucleotides are not labeled with a detectable label, they facilitate binding of labeled probes and so indirectly enhance hybridization signals.

15 A "biological sample" refers to a sample of material that is to be tested for the presence of microorganisms or nucleic acid thereof. The biological sample can be obtained from an organism such as a human patient, a laboratory mammal such as a mouse, rat, pig, monkey or other member of the primate family, by drawing a blood sample, sputum sample, spinal fluid sample, a urine sample, a nasal swab or a throat swab, or a culture of
20 such a sample. Ordinarily, the biological sample will contain hybridizable polynucleotides. These polynucleotides may have been released from organisms that comprise the biological sample, or alternatively can be released from the organisms in the sample using techniques such as sonic disruption or enzymatic or chemical lysis of cells to release polynucleotides so that they are available for hybridization with a polynucleotide probe.

25 As used herein "acridinium ester" means any of the family of chemiluminescent compounds based on the acridinium ring structure having an ester linkage at the C-9 position.

An "analyte" means any substance capable of undergoing a binding reaction with one or more specific binding partners, including, without limitation, antigens and antibodies
30 thereto, haptens and antibodies thereto; hormones, drugs, metabolites, vitamins, coenzymes and their binding partners, including receptors; polynucleotides, oligonucleotides, and hybrids of polynucleotides or oligonucleotides and antibodies and binding substances

WO 2005/017202

PCT/US2004/014690

thereto; polynucleotides or oligonucleotides and hybridizable polynucleotides or oligonucleotides thereto; metals and chelating agents thereto.

A "binding partner" means any molecule or substance capable of undergoing a specific binding reaction with an analyte.

5 "Bound" means a condition in which a binding interaction has been formed between a molecule and its specific binding partner.

"T_m" refers to the temperature at which 50% of the probe is converted from the hybridized to the unhybridized form.

One skilled in the art will understand that probes that substantially correspond to a
 10 reference sequence or region can vary from that reference sequence or region and still hybridize to the same target nucleic acid sequence. Probes of the present invention substantially correspond to a nucleic acid sequence or region if the percentage of identical bases or the percentage of perfectly complementary bases between the probe and its target sequence is from 100% to 80% or from 0 base mismatches in a 10 nucleotide target
 15 sequence to 2 bases mismatched in a 10 nucleotide target sequence. In preferred embodiments, the percentage is from 100% to 85%. In more preferred embodiments this percentage is from 90% to 100%; in other preferred embodiments, this percentage is from 95% to 100%. Probes that substantially correspond to a reference sequence or region include probes having any additions or deletions which do not prevent the probe from
 20 having its claimed property, such as being able to preferentially hybridize under high stringency hybridization conditions to its target nucleic acid over non-target nucleic acids.

By "sufficiently complementary" or "substantially complementary" is meant nucleic acids having a sufficient amount of contiguous complementary nucleotides to form, under high stringency hybridization conditions, a hybrid that is stable for detection.

25 By "antisense" is meant a nucleic acid molecule perfectly complementary to a reference (i.e., sense) nucleic acid molecule.

"RNA and DNA equivalents" refer to RNA and DNA molecules having the same complementary base pair hybridization properties. RNA and DNA equivalents have different sugar groups (i.e., ribose versus deoxyribose), and may differ by the presence of
 30 uracil in RNA and thymine in DNA. The difference between RNA and DNA equivalents do not contribute to differences in substantially corresponding nucleic acid sequences because the equivalents have the same degree of complementarity to a particular sequence.

Introduction to the Invention

The present invention relates to polynucleotide-based methods, compositions, kits and devices that can be used for detecting the nucleic acids that encode resistance to methicillin and/or vancomycin. More specifically, the invention provides for detection of the *mecA* gene, which is associated with methicillin-resistance of microorganisms, as well as for detection of the VanA and VanB genes, which are associated with vancomycin resistance of microorganisms. The present invention also relates to a rapid method for processing biological samples to obtain DNA, preferably which is substantially free of RNA, including ribosomal RNA (rRNA).

As described hereinbelow, a rapid sample processing method is provided which results in a sample that has denatured DNA and preferably is substantially free of rRNA. The method includes providing a sample, for instance, a cellular lysate which contains at least partially denatured DNA. In one embodiment, cells are treated with a lysing agent and then heat. In another embodiment, cells are treated with a lysing agent and heat simultaneously. The sample is then treated with a base and optionally heat to substantially decrease or eliminate RNA, including rRNA, in the sample. As used herein, a "substantial decrease" in rRNA in a sample means a decrease of at least 100-fold, preferably at least 500-fold, and more preferably at least 900-fold or more, relative to an untreated (control) sample. After base hydrolysis, the sample is treated with one or more reagents to yield a reaction mixture at a pH of about 4.5 to about 5.5, as well as reagents in an amount which results in a reaction mixture suitable for hybridizing polynucleotides (i.e., a mixture effective for a hybridization reaction with one or more probes and a target polynucleotide). While base treatment alone yielded reliable results, the combination of thermal and base denaturation yielded enhanced results. More specifically, the combination yielded enhanced rRNA degradation and denaturation of double-stranded DNA. Moreover, the combination of thermal and base denaturation resulted in reduced background signals and higher probe-specific signals relative to base denaturation alone. Further, if a probe in a hybridization reaction can bind to both RNA and DNA targets, the decrease in RNA in the sample reduces the level of RNA target relative to a genomic DNA target. In this way, a rDNA target can be used as an internal control and/or organism identifier in a hybridization reaction.

The above-described sample processing method was employed to identify an isolate

WO 2005/017202

PCT/US2004/014690

as a member of the *Staphylococcus* genus, and more specifically as *S. aureus*, coupled with the determination of its *mecA* status, in one hour. Indeed, 290 clinical and 24 ATCC isolates representing 303 *S. aureus* and 11 other *Staphylococcus spp.* were processed and hybridized with *mecA* gene probes in a hybridization protection assay (HPA). Results of the *mecA* HPA assay were compared to oxacillin (Ox) minimum inhibitory concentrations (MICs). Using the *mecA* probes disclosed herein, there was 100% correlation with 263 methicillin resistant *S. aureus* (MRSA) and methicillin sensitive *S. aureus* (MSSA). Additionally, 15 of 27 borderline resistant *S. aureus* (BORSA) were positive by the hybridization assay. *mecA* PCR and a latex agglutination assay (Oxoid, UK) results on the borderline isolates demonstrated 100% agreement with the probe results. Thus, by probing a sample with a ribosomal DNA probe and DNA probes specific for multiple regions within the *mecA* gene, a single processed sample yielded both isolate identification and antibiotic marker detection. This multi-probe system also serves as an internal control, as rDNA copy number is quite stable for a given organism, and so ensures that adequate target DNA is present to provide meaningful results. This advantageously can eliminate false-negative results. In one embodiment, testing for methicillin resistance status may be performed after a primary blood culture bottle turns positive, thereby eliminating the need for plating and incubating samples.

The invented method can lead to the appropriate choice of antibiotics to treat an infection, thereby reducing the occurrence of antibiotic resistance. This rapid probe-based assay allows the simultaneous identification and *mecA* gene detection of *Staphylococcus* directly from plates, and so is very useful in clinical laboratories and hospitals to identify MRSA and methicillin-resistant, coagulase-negative *Staphylococci* (MRCoNS). Moreover, it is even possible to perform simultaneous testing for methicillin-resistance and vancomycin-resistance, as described below.

Thus, one embodiment of the invention includes a method for detecting the *mecA* gene in a sample. The method includes lysing a sample comprising cells and treating the lysate with a basic composition in an amount and under conditions effective to substantially decrease or eliminate RNA in the sample, yielding a basic mixture. In one embodiment, the sample does not contain *in vitro* amplified nucleic acids. To denature the DNA in the sample, the sample may be treated with heat at a temperature of at least 95°C for at least 5 minutes, concurrent with, or subsequent to, addition of a lysing reagent. Alternatively,

WO 2005/017202

PCT/US2004/014690

DNA in a sample may be denatured by treatment with a basic composition, optionally with concurrent heat treatment, for example at 60°C. Preferably, the DNA in the sample is treated with heat, for example at a temperature of at least 95°C, concurrent with, or subsequent to, addition of a lysing reagent followed by treatment with a basic composition, as the combination of treatments was found to improve the sensitivity of target DNA detection. In one embodiment, the basic composition includes a hydroxide solution. The hydrolyzed sample is then contacted with one or more reagents in an amount that alters the pH of the basic mixture to a pH of about 4.0 to about 6.0, preferably a pH of about 4.5 to about 5.5, and result in a buffered mixture suitable for conducting a hybridization reaction.

In one embodiment, the one or more reagents added to alter the pH of the basic mixture include an acid, such as HCl, and optionally a buffering compound, e.g., a succinate salt. In one embodiment, the basic mixture is aliquoted into separate reaction vessels, each aliquot then contacted with one or more reagents in an amount that alters the pH of the basic mixture to a pH of about 4.0 to about 6.0 and results in a buffered hybridization reaction mixture. Each buffered hybridization reaction mixture can be contacted with at least one probe, such as a probe specific for *mecA* sequences or either or both of the VanA and VanB sequences, under conditions, preferably, high stringency conditions, effective to hybridize the probe to denatured target DNA. Optionally, at least one buffered hybridization reaction mixture is contacted with one or more probes that are different from the probe(s) in other hybridization reaction mixtures. In another embodiment, the basic mixture is contacted with one or more reagents in an amount that alters the pH of the basic mixture to a pH of about 4.0 to 6.0 and results in a buffered mixture suitable for conducting a hybridization reaction, and the buffered hybridization reaction mixture, or a portion thereof, is further contacted with at least one probe under conditions, preferably, high stringency conditions, effective to hybridize the probe to denatured target DNA. For instance, a portion of the hybridization reaction mixture may be added to a reaction vessel and one or more probes added to the vessel. Alternatively, a portion of the hybridization reaction mixture may be added to at least two reaction vessels and different probes added to each of those vessels. Alternatively, the hybridization reaction mixture or a portion thereof is added to one or more probes in a reaction vessel. Exemplary high stringency conditions include a hybridization reaction mixture including 0.48 M sodium phosphate buffer, 0.1% sodium dodecyl sulfate and 1 mM each of EDTA and EGTA, or including 0.6 M LiCl, 1% lithium

WO 2005/017202

PCT/US2004/014690

lauryl sulfate, 50 to 60 mM lithium succinate and 10 mM each EDTA and EGTA, and temperatures of at least 50°C.

In one preferred embodiment, the one or more probes used in the hybridization reaction includes a detectable moiety. Optionally, after the hybridization reaction is
5 subjected to conditions effective to result in hybrid formation, the hybridization reaction is subjected to conditions effective to decrease or eliminate the amount of unhybridized probe. Then the presence or amount of hybridized probe is detected or determined. Although the sample processing method described herein was employed to prepare samples to detect *mecA* DNA, the sample processing method may be employed to prepare a sample to detect
10 any DNA target sequence.

The invention also provides a method for detecting *mecA* sequences in a sample. The method comprises contacting a sample comprising denatured DNA with one or more *mecA* oligonucleotide probes under high stringency hybridization conditions effective to form a hybrid between the oligonucleotide probes and *mecA* DNA. The presence of hybrid
15 formation is detected or determined thereafter. The probes include sequences substantially corresponding to nucleotides 304 to 338 of the *mecA* gene, the complement thereof, or a portion thereof, sequences substantially correspond to nucleotides 501 to 531 of the *mecA* gene, the complement thereof, or a portion thereof, or sequences substantially correspond to nucleotides 1010 to 1044 of the *mecA* gene, the complement thereof, or a portion thereof.
20 Preferably, the probes are labeled, for example with a chemiluminescent label, a fluorescent label or a radioactive label. In one embodiment, at least three *mecA* oligonucleotide probes are employed to detect or determine the *mecA* status of a sample. In one preferred embodiment, at least one probe has at least one nucleotide that has a substituted ribose moiety, such as a methoxy group at the 2' position of the ribose moiety.

25 Optionally, one or more non-*mecA* gene probes, such as a probe specific for ribosomal nucleic acids, may be employed in addition to the *mecA* gene probes to identify the microorganism in the sample and/or to confirm that sufficient DNA was present in the sample to permit detection of the *mecA* gene (i.e., an internal control). For instance, a rDNA probe specific for one or more organisms, such as a probe specific for
30 *Staphylococcus* bacteria, may be used to identify the organism in the sample. In one embodiment, the sample processing method described herein is employed to prepare the sample prior to detecting or determining the *mecA* status of the sample. Although the *mecA*

WO 2005/017202

PCT/US2004/014690

gene is generally associated with *S. aureus*, other Staphylococcal species (coagulase-negative Staphylococci, "CoNS") may contain the *mecA* gene as the result of lateral transfer of genetic information between the microorganisms. Therefore, an identification or internal control reaction used in conjunction with a *mecA* reaction may employ a rRNA or rDNA probe for an organism other than *S. aureus*. Thus, in one embodiment, the method includes comparing the presence or amount of hybrid formation in a sample contacted with *mecA* gene probes to the presence or amount of hybrid formation in a corresponding sample contacted with a probe specific for rRNA or rDNA.

In another embodiment of the invention, one or more *mecA* gene probes are employed with a sample which contains amplified nucleic acid. The method includes contacting a sample comprising amplified *mecA* nucleic acid, with at least one *mecA* oligonucleotide probe under high stringency hybridization conditions effective to form a hybrid between the probe and the amplified *mecA* nucleic acid. In one embodiment, the amplified nucleic acid is RNA amplified using transcription mediated amplification. Optionally, prior to amplification, a cellular sample is contacted with methicillin to induce transcription from the *mecA* gene. Then the presence or amount of hybrid formation is detected or determined. Probes useful in this embodiment of the invention includes those with sequences substantially corresponding to nucleotides 304 to 338 of the *mecA* gene, the complement thereof, or a portion thereof, nucleotides 501 to 531 of the *mecA* gene, the complement thereof, or a portion thereof, nucleotides 1010 to 1044 of the *mecA* gene, the complement thereof, or a portion thereof.

The invention also includes a probe mix. The mix includes two or more of a first oligonucleotide substantially corresponding to nucleotides 304 to 338 of the *mecA* gene, the complement thereof, or a portion thereof, a second oligonucleotide substantially corresponding to nucleotides 501 to 531 of the *mecA* gene, the complement thereof, or a portion thereof, and a third oligonucleotide substantially corresponding to nucleotides 1010 to 1044 of the *mecA* gene, the complement thereof, or a portion thereof. Each oligonucleotide in the mix hybridizes under stringent hybridization conditions to *mecA* genomic DNA. In one embodiment, the mix includes a first oligonucleotide corresponding to nucleotides 304 to 338 of the *mecA* gene, a second oligonucleotide corresponding to the complement of nucleotides 501 to 531 of the *mecA* gene, and a third oligonucleotide corresponding to nucleotides 1010 to 1044 of the *mecA* gene. In another embodiment, the

WO 2005/017202

PCT/US2004/014690

mix includes a first oligonucleotide corresponding to nucleotides 304 to 338 of *mecA*, a second oligonucleotide corresponding to the complement of nucleotides 501 to 531 of the *mecA* gene, and a third oligonucleotide corresponding to the complement of nucleotides 1010 to 1044 of the *mecA* gene. In another embodiment, the mix includes a first
5 oligonucleotide corresponding to the complement of nucleotides 304 to 338 of *mecA*, a second oligonucleotide corresponding to the complement of nucleotides 501 to 531 of the *mecA* gene, and a third oligonucleotide corresponding to nucleotides 1010 to 1044 of the *mecA* gene. In yet another embodiment, the mix includes a first oligonucleotide corresponding to nucleotides 304 to 338 of the *mecA* gene, a second oligonucleotide
10 corresponding to nucleotides 501 to 531 of the *mecA* gene, and a third oligonucleotide corresponding to the complement of nucleotides 1010 to 1044 of the *mecA* gene. Preferably, at least one oligonucleotide in the mix comprises one or more nucleotide analogs comprising a 2'-O-methyl ribosyl analog rather than a ribosyl or deoxyribosyl moiety.

15 The invention further includes a kit with probes useful for detecting the *mecA* gene in a test sample. The kit includes one or more of a first oligonucleotide comprising sequences corresponding to nucleotides 304 to 338 of the *mecA* gene, the complement thereof, or a portion thereof, a second oligonucleotide comprising sequences corresponding to nucleotides 501 to 531 of the *mecA* gene, the complement thereof, or a portion thereof,
20 and a third oligonucleotide comprising sequences corresponding to nucleotides 1010 to 1044 of the *mecA* gene, the complement thereof, or a portion thereof, wherein each oligonucleotide hybridizes under stringent hybridization conditions to *mecA* genomic DNA. The kit optionally includes probes useful for detecting single-copy genes other than *mecA*, including other drug resistance genes, or to detect multiple copy genes, such as rDNA
25 genes.

The invention thus provides for the use of two or more probes, one of which is optionally employed as an identifier probe capable of providing information about the identity of an organism. For example, the identifier probe may be capable of providing information about the group, family, genus or species of an organism. Alternatively, an
30 identifier probe may be chosen to distinguish Gram-positive and Gram-negative bacteria, or to make a species- or strain-level identification. More preferably, the use of a "matrix" of polynucleotide hybridization probes could be used in conjunction with the antibiotic

resistance probes described herein. In one embodiment, certain hybridization probes are specific for a selected gene, such as the *mecA* gene, while other hybridization probes are specific for nucleic acids, such as rRNA and rDNA, of various species or taxonomically related groups of organisms. Using these probes, it is possible to determine the status of the organism, meaning whether the genome of the organism includes the gene or sequences related to the gene, and the species of the organism. The method is particularly suited for use in connection with automated systems.

Even negative results in a probe matrix hybridization procedure can be meaningful when taken in the context of other results from the matrix. For example, if it is determined from a two-locus matrix that a sample contains polynucleotides that give a positive hybridization signal at a first address that identifies *S. aureus*, and negative results at a second address that identifies the *mecA* gene, then the combination of results would indicate that the polynucleotides were derived from a *S. aureus* that is *mecA*-negative (*mecA*(-)). Alternatively, if it is determined from a two-locus matrix that a sample does not contain polynucleotides that hybridize at a first address that identifies nucleic acids of bacteria that are members of the *Staphylococcus* genus (i.e., a negative result at a *Staphylococcus* genus address) and a positive result at a second address that identifies the *mecA* gene, then the combination of results would indicate that the polynucleotides were derived from a non-*Staphylococcus* organism that is *mecA*-positive (*mecA*(+)).

Another aspect of a matrix-based method of gene and/or microbe identification relates to the use of a combination of probes in a single hybridization reaction, such as may be conducted in a single well of a multi-well plate. In one embodiment, a combination of polynucleotide probes that hybridize to a selected gene, for instance, the *mecA* gene, is employed with one or more probes for another gene, e.g., rDNA genes, which optionally may be employed as an internal control. In a preferred embodiment, probes for the *mecA* gene are labeled with a label that is different from the label on the internal control probe. This arrangement facilitates detection of both targets in a single hybridization reaction. In another preferred embodiment, the probes for detecting the *mecA* gene and the probes that are specific for ribosomal nucleic acids are labeled with the same detectable label, or with labels that are not distinguished from each other during a detection step which is used in a method employing the invented apparatus or device.

I. Probe Selection and Preparation

It is not always necessary to determine the entire nucleic acid sequence of a gene of interest in order to obtain a probe specific for that gene. In a highly preferred embodiment, all of the probes are particularly selected for use under a single set of hybridization
5 conditions, including a single temperature and/or a single set of ionic strength conditions. The following guidelines are useful for designing probes with desirable characteristics.

First, the stability of the probe:target polynucleotide hybrid is chosen to be compatible with the assay conditions. This may be accomplished by avoiding long A and T rich sequences, by terminating the hybrids with G:C base pairs and by designing the probe
10 in such a way that the T_m will be appropriate for standard conditions to be employed in the assay. The nucleotide sequence of the probe should be chosen so that the length and % G and % C result in a probe having a T_m about 2 to 10°C higher than the temperature at which the final assay is performed. The base composition of the probe is significant because G:C base pairs exhibit greater thermal stability when compared with A:T base pairs. Thus,
15 hybrids involving complementary polynucleotides having a high G:C content are generally stable at higher temperatures when compared with hybrids having a lower G:C content.

Second, the position at which the probe binds its target polynucleotide is chosen to minimize the stability of hybrids formed between probe:non-target polynucleotides. This may be accomplished by minimizing the length of perfect complementarity with
20 polynucleotides of non-target organisms, by avoiding G:C rich regions of homology with non-target sequences, and by positioning the probe to span as many destabilizing mismatches as possible. Whether a probe sequence is useful for detecting only a specific type of organism or gene depends largely on thermal stability differences between probe:target hybrids and probe:non-target hybrids. The differences in T_m should be as large
25 as possible to produce highly specific probes.

The length of the target polynucleotide sequence and the corresponding length of the probe sequence also are important factors to be considered when designing a probe. While it is possible for polynucleotides that are not perfectly complementary to hybridize to each other, the longest stretch of perfectly homologous base sequence is ordinarily the primary
30 determinant of hybrid stability.

Third, regions which are known to form strong internal structures inhibitory to hybridization of a probe are less preferred as targets. Probes having extensive self-

WO 2005/017202

PCT/US2004/014690

complementarity also should be avoided.

Once a presumptive unique sequence has been identified, corresponding oligonucleotides are produced. Defined oligonucleotides that can be used to practice the invention can be produced by any of several well-known methods, including automated
5 solid-phase chemical synthesis using phosphoramidite precursors (Barone et al., *Nucl. Acids Res.* **12**:4051 (1984)). Other well-known methods for construction of synthetic oligonucleotides may, of course, be employed (see Sambrook et al., Molecular Cloning: A Laboratory Manual, 11 (1989)). All of the oligonucleotides of the present invention may be modified with chemical groups to enhance their performance. Thus, it is to be understood
10 that references to "oligonucleotide probes" or "helper oligonucleotides" or simply "oligonucleotides" embrace polymers of native nucleotides as well as polymers that include at least one nucleotide analog. Backbone-modified oligonucleotides, such as those having phosphorothioate or methylphosphonate groups, are examples of analogs that can be used in conjunction with oligonucleotides of the present invention. These modifications render the
15 oligonucleotides resistant to the nucleolytic activity of certain polymerases or to nuclease enzymes. Other analogs that can be incorporated into the structures of the oligonucleotides include peptide nucleic acids, or "PNAs." The PNAs are compounds comprising ligands linked to a peptide backbone rather than to a phosphodiester backbone. Representative ligands include either the four main naturally occurring DNA bases (i.e., thymine, cytosine,
20 adenine or guanine) or other naturally occurring nucleobases (e.g., inosine, uracil, 5-methylcytosine or thiouracil) or artificial bases (e.g., bromothymine, azaadenines or azaguanines, etc.) attached to a peptide backbone through a suitable linker. PNAs are able to bind complementary single-stranded DNA (ssDNA) and RNA strands. Methods for making and using PNAs are disclosed in U.S. Patent No. 5,539,082. Another type of
25 modification that can be used to make oligonucleotides having the sequences described herein involves the use of non-nucleotide linkers (e.g., see U.S. Patent No. 6,031,091) between nucleotides in the nucleic acid chain which do not interfere with hybridization or optionally elongation of a primer.

Yet other analogs include those which increase the binding affinity of a probe to a
30 target nucleic acid and/or increase the rate of binding of the probe to the target nucleic acid relative to a probe without the analog. Such analogs include those with a modification (substitution) at the 2' position of a ribofuranosyl nucleotide. When employed in

hybridization reactions, probes containing one or more of these analogs, with their resulting changes in hybridization properties, may be shorter in length than corresponding DNA probes. The use of shorter oligonucleotides to specifically bind target nucleic acids at a given temperature has additional advantages. For instance, shorter oligonucleotides will generally have a greater ability to discriminate perfectly complementary targets from "mismatched" base sequence regions. Shorter oligonucleotides are also less likely to overlap undesirable base sequences. The use of higher hybridization temperatures kinetically drives the hybridization reaction, resulting in faster hybridization rates than would occur at lower temperatures. Further, modified oligonucleotides may result in faster hybridization rates than the unmodified versions, even when the temperature is not raised.

Thus, hybridization assay probes and/or helper oligonucleotides can all be designed to contain modified bases which, alone or in combination, may have the advantage of increasing the rate of target-specific hybridization.

Analogues having a modification at the 2' position of the ribose, e.g., an alkyl, and alkoxy or a halide substitution, are one preferred embodiment. In one preferred embodiment, oligonucleotides contain nucleotide analogs having 2'-O-methylribofuranosyl moieties linked to a nitrogenous base. Other substitutions at the 2' position of the sugar are expected to have similar properties so long as the substitution is not so large as to cause steric inhibition of hybridization.

Additionally, other modifications which increase the T_m of a modified oligonucleotide:target hybrid would reasonably be expected to contribute to increases in the rate of hybridization as well. Such modifications may occur at the 2' position (or other positions) of the deoxyribofuranosyl or ribofuranosyl moiety (such as 2' halide substitutions), on the nitrogenous bases (such as N-diisobutylaminomethylidene-5-(1-propynyl)-2'-deoxycytidine; a cytidine analog, or 5-(1-propynyl)-2'-deoxyuridine); a thymidine analog, or in the linkage moiety.

Preferably, probes are labeled. Essentially any labeling and detection system that can be used for monitoring specific nucleic acid hybridization can be used in conjunction with the probes disclosed herein when a labeled probe is desired. Included among the collection of useful labels are: radiolabels, enzymes, haptens, linked oligonucleotides, colorimetric, fluorometric or chemiluminescent molecules, and redox-active moieties that are amenable to electrochemical detection methods. Standard isotopic labels that can be

WO 2005/017202

PCT/US2004/014690

used to produce labeled oligonucleotides include ^3H , ^{35}S , ^{32}P , ^{125}I , ^{57}Co and ^{14}C . When using radiolabeled probes, hybrids can be detected by autoradiography, scintillation counting or gamma counting.

Non-isotopic materials can also be used for labeling oligonucleotide probes. These
5 non-isotopic labels can be positioned internally or at a terminus of the oligonucleotide probe. Modified nucleotides can be incorporated enzymatically or chemically with modifications of the probe being performed during or after probe synthesis, for example, by the use of non-nucleotide linker groups. Non-isotopic labels include colorimetric molecules, fluorescent molecules, chemiluminescent molecules, enzymes, cofactors,
10 enzyme substrates, haptens or other ligands. Acridinium esters are one example of chemiluminescent labels useful for detecting probe hybrids.

Indeed, any number of different non-isotopic labels can be used for preparing labeled oligonucleotides in accordance with the invention. Preferred chemiluminescent molecules include acridinium esters of the type disclosed in U.S. Patent No. 5,283,174 for
15 use in connection with homogenous protection assays, and of the type disclosed in U.S. Patent No. 5,656,207 for use in connection with assays that quantify multiple targets in a single reaction. The disclosures of these U.S. patents are hereby incorporated by reference. U.S. Patent No. 5,998,135 discloses yet another method that can be used for labeling and detecting probes using fluorimetry to detect fluorescence emission from lanthanide metal
20 labels disposed on probes, where the emission from these labels becomes enhanced when it is in close proximity to an energy transfer partner. Preferred electrochemical labeling and detection approaches are disclosed in U.S. Patent Nos. 5,591,578 and 5,770,369, and PCT/US98/12082, the disclosures of which are hereby incorporated by reference. Redox active moieties useful as electrochemical labels include transition metals such as Cd, Mg,
25 Cu, Co, Pd, Zn, Fe and Ru.

In some applications, probes exhibiting at least some degree of self-complementarity are desirable to facilitate detection of probe:target duplexes in a test sample without first requiring the removal of unhybridized probe prior to detection. By way of example, structures referred to as "molecular torches" are designed to include
30 distinct regions of self-complementarity (coined "the target binding domain" and "the target closing domain") which are connected by a joining region and which hybridize to one another under predetermined hybridization assay conditions. When exposed to denaturing

conditions, the two complementary regions (which may be fully or partially complementary) of the molecular torch melt, leaving the target binding domain available for hybridization to a target sequence when the predetermined hybridization assay conditions are restored. Molecular torches are designed so that the target binding domain
5 favors hybridization to the target sequence over the target closing domain. The target binding domain and the target closing domain of a molecular torch include interacting labels (e.g., fluorescent/quencher) positioned so that a different signal is produced when the molecular torch is self-hybridized as opposed to when the molecular torch is hybridized to a target nucleic acid, thereby permitting detection of probe:target duplexes in a test sample in
10 the presence of unhybridized probe having a viable label associated therewith. Molecular torches are fully described in U.S. Patent No. 6,361,945, the disclosure of which is hereby incorporated by reference.

Another example of a self-complementary hybridization assay probe that may be used in conjunction with the invention is a structure commonly referred to as a “molecular
15 beacon.” Molecular beacons comprise nucleic acid molecules having a target complementary sequence, an affinity pair (or nucleic acid arms) holding the probe in a closed conformation in the absence of a target nucleic acid sequence, and a label pair that interacts when the probe is in a closed conformation. Hybridization of the target nucleic acid and the target complementary sequence separates the members of the affinity pair,
20 thereby shifting the probe to an open conformation. The shift to the open conformation is detectable due to reduced interaction of the label pair, which may be, for example, a fluorophore and a quencher (e.g., DABCYL and EDANS). Molecular beacons are fully described in U.S. Patent No. 5,925,517, the disclosure of which is hereby incorporated by reference. Molecular beacons useful for detecting specific nucleic acid sequences may be
25 created by appending to either end of one of the probe sequences disclosed herein, a first nucleic acid arm comprising a fluorophore and a second nucleic acid arm comprising a quencher moiety. In this configuration, the specific probe sequence disclosed herein serves as the target-complementary “loop” portion of the resulting molecular beacon.

Molecular beacons preferably are labeled with an interactive pair of detectable
30 labels. Examples of detectable labels that are preferred as members of an interactive pair of labels interact with each other by FRET or non-FRET energy transfer mechanisms. Fluorescence resonance energy transfer (FRET) involves the radiationless transmission of

energy quanta from the site of absorption to the site of its utilization in the molecule, or system of molecules, by resonance interaction between chromophores, over distances considerably greater than interatomic distances, without conversion to thermal energy, and without the donor and acceptor coming into kinetic collision. The “donor” is the moiety
5 that initially absorbs the energy, and the “acceptor” is the moiety to which the energy is subsequently transferred. In addition to FRET, there are at least three other “non-FRET” energy transfer processes by which excitation energy can be transferred from a donor to an acceptor molecule.

When two labels are held sufficiently close that energy emitted by one label can be
10 received or absorbed by the second label, whether by a FRET or non-FRET mechanism, the two labels are said to be in “energy transfer relationship” with each other. This is the case, for example, when a molecular beacon is maintained in the closed state by formation of a stem duplex, and fluorescent emission from a fluorophore attached to one arm of the probe is quenched by a quencher moiety on the opposite arm.

15 Highly preferred label moieties for the invented molecular beacons include a fluorophore and a second moiety having fluorescence quenching properties (i.e., a “quencher”). In this embodiment, the characteristic signal is likely fluorescence of a particular wavelength, but alternatively could be a visible light signal. When fluorescence is involved, changes in emission are preferably due to FRET, or to radiative energy transfer
20 or non-FRET modes. When a molecular beacon having a pair of interactive labels in the closed state is stimulated by an appropriate frequency of light, a fluorescent signal is generated at a first level, which may be very low. When this same probe is in the open state and is stimulated by an appropriate frequency of light, the fluorophore and the quencher moieties are sufficiently separated from each other that energy transfer between them is
25 substantially precluded. Under that condition, the quencher moiety is unable to quench the fluorescence from the fluorophore moiety. If the fluorophore is stimulated by light energy of an appropriate wavelength, a fluorescent signal of a second level, higher than the first level, will be generated. The difference between the two levels of fluorescence is detectable and measurable. Using fluorophore and quencher moieties in this manner, the molecular
30 beacon is only “on” in the “open” conformation and indicates that the probe is bound to the target by emanating an easily detectable signal. The conformational state of the probe alters the signal generated from the probe by regulating the interaction between the label moieties.

WO 2005/017202

PCT/US2004/014690

Examples of donor/acceptor label pairs that may be used in connection with the invention, making no attempt to distinguish FRET from non-FRET pairs, include fluorescein/tetramethylrhodamine, IAEDANS/fluorescein, EDANS/DABCYL, coumarin/DABCYL, fluorescein/fluorescein, BODIPY FL/BODIPY FL, fluorescein/DABCYL, lucifer yellow/DABCYL, BODIPY/DABCYL, eosine/DABCYL, erythrosine/DABCYL, tetramethylrhodamine/DABCYL, Texas Red/DABCYL, CY5/BH1, CY5/BH2, CY3/BH1, CY3/BH2 and fluorescein/QSY7 dye. Those having an ordinary level of skill in the art will understand that when donor and acceptor dyes are different, energy transfer can be detected by the appearance of sensitized fluorescence of the acceptor or by quenching of donor fluorescence. When the donor and acceptor species are the same, energy can be detected by the resulting fluorescence depolarization. Non-fluorescent acceptors such as DABCYL and the QSY 7 dyes advantageously eliminate the potential problem of background fluorescence resulting from direct (i.e., non-sensitized) acceptor excitation. Preferred fluorophore moieties that can be used as one member of a donor-acceptor pair include fluorescein, ROX, and the CY dyes (such as CY5). Highly preferred quencher moieties that can be used as another member of a donor-acceptor pair include DABCYL and the BLACK HOLE QUENCHER moieties which are available from Biosearch Technologies, Inc., (Novato, CA).

Alternative procedures for detecting particular genes can be carried out using either labeled probes or unlabeled probes. For example, hybridization assay methods that do not rely on the use of a labeled probe are disclosed in U.S. Patent No. 5,945,286 which describes immobilization of unlabeled oligonucleotide probe analogs made of peptide PNAs, and detectably labeled intercalating molecules which can bind double-stranded PNA probe/target nucleic acid duplexes. In these procedures, as well as in certain electrochemical detection procedures, such as those disclosed in PCT/US98/12082, PCT/US98/12430 and PCT/US97/20014, the oligonucleotide probe is not required to harbor a detectable label.

II. Probe Specificity

High stringency conditions useful for conducting the hybridization procedures disclosed herein include conditions of 55°C to 65°C when the salt concentration is in the range of 0.6 to 0.9 M. Preferred salts include lithium chloride, but other salts such as sodium chloride and sodium citrate also can be used in the hybridization solution. Other

useful high stringency hybridization conditions are alternatively provided by 0.48 M sodium phosphate buffer, 0.1% sodium dodecyl sulfate, and 1 mM each of EDTA and EGTA, or by 0.6 M LiCl, 1 % lithium lauryl sulfate, 60 mM lithium succinate and 10 mM each of EDTA and EGTA. It is preferred that all of the probes for hybridization at about
 5 60°C will have T_m values in the range of from 63°C to about 78°C.

III. Exemplary Methods and Probes of the Invention

Nucleic acid probes specific for a gene of interest, such as a drug resistance gene, optionally in combination with one or more probes specific for a group of organisms find use in an assay for detecting the presence of the gene of interest in a biological sample that
 10 includes nucleic acids, and optionally for identifying a group of organisms and/or to serve as an internal control. For instance, a plurality of probes specific for the *mecA* gene and a probe specific for Staphylococcus rRNA or rDNA may be employed to determine whether a biological sample contains *mecA*(+) or *mecA*(-) organisms, such as *mecA*(+) or *mecA*(-) Staphylococci including *mecA*(+) or *mecA*(-) *S. aureus*. Likewise, a plurality of probes
 15 specific for the VanA and VanB genes encoding vancomycin resistance and one or more probes specific for the rRNA or rDNA of one or more Enterococcus species that include *E. faecalis* and *E. faecium* may be employed to determine whether a biological sample contains vancomycin-resistant Enterococcus. Including a probe specific for the rRNA or rDNA of *S. aureus* among the other probes of this latter panel provides a means for rapidly
 20 identifying vancomycin-resistant *S. aureus*.

A. Sample Processing Method

Since hybridization signals measured using certain probes intended for detecting DNA targets may actually be due to reactivity with RNA, which may be present in excess relative to the intended DNA target sequence, conditions which resulted in a substantial
 25 decrease or elimination of RNA were determined. Cells were lysed with a lysing agent in an amount which released polynucleotides from the cells, thus forming a lysate. Polynucleotides were denatured by heat treatment, for example, by heating at temperatures greater than 90°C, preferably at least 95°C, and more preferably at least 100°C to 105°C, for at least 5, and preferably at least 10 minutes and optionally 25 minutes or more. The
 30 lysing and denaturation steps may be conducted consecutively or simultaneously.

To remove RNA from the lysate, the lysate can be treated with base in an amount and under conditions which result in a substantial decrease in RNA in the sample. Any

base may be employed in this regard. In one embodiment, hydroxide is employed to remove RNA from the sample. Preferably, the hydroxide is added to a final concentration of about 0.2 to about 1.0 N and the resulting mixture heated at about 50-60°C for 5 minutes, and optionally up to 30 minutes or more. The mixture is then neutralized with an acidic composition to achieve a pH in the range of about 4.0-6.0, and reagents added to yield a buffered mixture containing, for example, monovalent ions, detergent, and succinate buffer at a final concentration of about 20 to about 100 mM. The neutralized mixture is suitable for conducting a hybridization reaction.

B1. *MecA* Probes

MecA sequences and structurally and/or functionally related sequences from a collection of organisms were aligned to identify candidate conserved sequences that could be used to distinguish *mecA*(+) organisms from *mecA*(-) organisms. Thus, by examining partial or complete sequences of *mecA*(+) genes of various organisms including the Staphylococcal organisms, aligning those sequences with structurally and/or functionally related sequences to reveal areas of maximum homology and areas of sequence variation, sequences were identified that are conserved among *mecA* genes but that exhibit mismatches with structurally and/or functionally related genes. Based on such considerations, the following regions of the *mecA* gene were selected for testing as probes: nucleotides 304 to 338, nucleotides 501 to 531, and nucleotides 1010 to 1044 of the *mecA* gene. Such conserved sequences were then tested against a panel of *mecA* standards and bacterial lysates to verify their utility as probes under laboratory conditions. In particular, probes that preferentially hybridized to a nucleic acid target region to form a detectable duplex were chosen for polynucleotide-based diagnostic assays. Preferably, two or more of such probes are employed to enhance the hybridization signal corresponding to formation of a duplex between labeled oligonucleotide probes and their complementary target nucleic acids. Use of multiple probes also minimizes false-negative results arising from decreased detection of spurious mutations in a single probe target region.

Preferred methods for detecting the presence of the *mecA* gene include the step of contacting a test sample under high stringency hybridization conditions with at least two, and preferably at least three, oligonucleotide probes that preferentially hybridize to the *mecA* gene, and optionally at least one oligonucleotide probe that preferentially hybridizes to a nucleic acid sequence, such as a DNA sequence characteristic of Staphylococcal

organisms, over a nucleic acid sequence of other organisms.

While oligonucleotide probes of different lengths and base composition may be used for detecting the *mecA* gene, preferred probes have lengths of up to 70 nucleotides, and more preferably have lengths of up to 60 nucleotides, and are sufficiently homologous to the target nucleic acid to permit hybridization under high stringency conditions. However, the specific probe sequences described below also may be provided in a nucleic acid cloning vector or transcript or other longer nucleic acid and still can be used for detecting the *mecA* gene. Thus, the probes may include sequences unrelated to the *mecA* gene, for instance at the 5' end, at the 3' end, or at both the 5' and 3' ends (as would be the case for molecular beacons and molecular torches). Preferred probes for detecting the *mecA* gene have sequences of up to 60 nucleotides in length and preferably have at least 17 contiguous nucleotides and more preferably 30 to 35 contiguous nucleotides corresponding to sequences in the *mecA* gene (see GENBANK Accession No. X52593) or the complement thereof. Preferred oligonucleotide sequences include RNA and DNA equivalents, and may include at least one nucleotide analog.

The probes described herein were tested in hybridization reactions using synthetic targets as well as biological samples, such as *Staphylococcus* isolates, to detect the *mecA* gene and optionally to identify the genus and/or species of a microorganism present in the biological sample. In one method of determining whether a biological sample contains *mecA* gene sequences and nucleic acids that would indicate the presence of organisms in the *Staphylococcus* genus, nucleic acids can be released from bacterial cells by addition of a lysing agent, such as a detergent, or by other known methods for disrupting cells including the use of enzymes, osmotic shock, chemical treatment, and vortexing, for instance, with glass beads, or sonic disruption, for example according to the method disclosed in U.S. Patent No. 5,374,522. Other appropriate methods suitable for liberating nucleic acids from cells have been described in U.S. Patent No. 5,837,452 and in U.S. Patent No. 5,364,763. In one preferred embodiment, cells are contacted with a lysis buffer containing a lysing agent.

Preferably, the *mecA*-specific probes hybridize to *mecA* DNA sequences only under conditions of high stringency. Under these conditions only highly complementary nucleic acid hybrids will form (i.e., those having at least 14 out of 17 bases in a contiguous series of bases being complementary). Hybrids will not form in the absence of a sufficient degree of

complementarity. Accordingly, the stringency of the assay conditions determines the amount of complementarity needed between two nucleic acid strands forming a hybrid. Stringency is chosen to maximize the difference in stability between the hybrid formed with target nucleic acid and non-target nucleic acid.

- 5 In one embodiment, probes used for detecting *mecA* sequences include at least one of SEQ ID NOs:1-6, which have the following sequences:
- GGTATGTGGAAGTTAGATTGGGATCATAGCG (SEQ ID NO:1);
- CGCTATGATCCCAATCTAACTTCCACATACC (SEQ ID NO:2);
- GCGATAATGGTGAAGTAGAAATGACTGAACGTCCG (SEQ ID NO:3);
- 10 CGGACGTTTCAGTCATTTCTACTTCACCATTATCGC (SEQ ID NO:4);
- GCTCCAACATGAAGATGGCTATCGTGTCACAATCG (SEQ ID NO:5); and
- CGATTGTGACACGATAGCCATCTTCATGTTGGAGC (SEQ ID NO:6), respectively.
- Of course, portions of any of these sequences which preferentially hybridize to the *mecA* gene or its complement can be used instead. In one embodiment, preferred probes comprise
- 15 one or more nucleotide analogs comprising a 2'-methoxy ribosyl analog substituted for one or more nucleotides comprising a deoxyribosyl moiety in one of SEQ ID NOs:1-6.

B2. VanA and VanB Probes

- Preferred methods for detecting the presence of VanA or VanB gene sequences, for example in an Enterococcal organism, include the step of contacting a test sample under
- 20 high stringency hybridization conditions with at least one, preferably at least two, and still more preferably at least three, oligonucleotide probes that preferentially hybridize, in aggregate, to the VanA and VanB genes, and optionally at least one oligonucleotide probe that preferentially hybridizes to a nucleic acid sequence, such as a DNA sequence, of Enterococcal organisms over a nucleic acid sequence of other organisms.

- 25 While oligonucleotide probes of different lengths and base composition may be used for detecting the VanA and VanB genes, preferred probes have lengths of up to 70 nucleotides, and more preferably have lengths of up to 60 nucleotides, and are sufficiently homologous to the target nucleic acid to permit hybridization under high stringency conditions. However, the specific probe sequences described below also may be provided
- 30 in a nucleic acid cloning vector or transcript or other longer nucleic acid and still can be used for detecting the VanA and VanB genes. Thus, the probes may include sequences unrelated to the VanA and VanB genes, for instance at the 5' end, at the 3' end, or at both

the 5' and 3' ends (as would be the case for molecular beacons and molecular torches). Preferred probes for detecting the VanA and VanB genes have sequences of up to 60 nucleotides in length and preferably have at least 17 contiguous nucleotides and more preferably 30 to 35 contiguous nucleotides corresponding to sequences in the VanA and/or VanB genes, or the complements thereof. Preferred oligonucleotide sequences include RNA and DNA equivalents, and may include at least one nucleotide analog.

In one method of determining whether a biological sample contains VanA and/or VanB gene sequences and nucleic acid that would indicate the presence of members of the Enterococcus group, nucleic acids can be released from bacterial cells by addition of a lysing agent, such as a detergent, or by other known methods for disrupting cells including the use of enzymes, osmotic shock, chemical treatment, and vortexing, for instance, with glass beads, or sonic disruption, for example according to the method disclosed in U.S. Patent No. 5,374,522. Other methods suitable for liberating nucleic acids from cells which can then be subjected to hybridization methods have been described in U.S. Patent No. 5,837,452 and in U.S. Patent No. 5,364,763. In one preferred embodiment, cells are contacted with a lysis buffer containing a lysing agent.

Preferably, the probes specifically hybridize to VanA and/or VanB in genomic DNA only under conditions of high stringency. Under these conditions only highly complementary nucleic acid hybrids will form (i.e., those having at least 14 out of 17 bases in a contiguous series of bases being complementary). Hybrids will not form in the absence of a sufficient degree of complementarity. Accordingly, the stringency of the assay conditions determines the amount of complementarity needed between two nucleic acid strands forming a hybrid. Stringency is chosen to maximize the difference in stability between the hybrid formed with target nucleic acid and non-target nucleic acid.

In one embodiment, the probes for detecting the VanA and/or VanB genes include the sequences presented under Example 10, or portions thereof which preferentially hybridize to one of the strands of the VanA and/or VanB genes, or their complements. In various preferred embodiments, probes comprise one or more nucleotide analogs, such as 2'-methoxy ribosyl analogs or base analogs.

30 C. rDNA Probes

rRNA sequences from a collection of related and unrelated organisms can be aligned to identify candidate sequences conserved within a genus that could be used to

WO 2005/017202

PCT/US2004/014690

distinguish Staphylococcal and/or Enterococcal organisms from other bacterial and eukaryotic organisms. Thus, by examination of partial or complete sequences of rRNA or rDNA of various organisms including the Staphylococcal or Enterococcal organisms and unrelated phylogenetic neighbors, aligning the sequences to reveal areas of maximum
5 homology and areas with sequence variation, rRNA or rDNA sequences are identified that are conserved among members of a genus but that exhibit mismatches with rRNA or rDNA sequences of other closely and distantly related genera. Such sequences are then tested against a panel of rRNA or rDNA standards and bacterial lysates to verify their utility as probes under laboratory conditions.

10 Polynucleotide sequences of rRNAs or rDNAs are most conveniently determined using a dideoxynucleotide sequencing procedure. For rRNA, oligonucleotide primers of about 10-100 bases in length and complementary to conserved regions of rRNA from any of the 5S, 16S or 23S ribosome subunits can be extended by reverse transcriptase. The resulting DNA extension products can then be sequenced by chemical degradation or
15 dideoxynucleotide sequencing (Lane et al., *Proc. Natl. Acad. Sci. USA*, **82**:6955 (1985)). According to another method, genomic sequences encoding the rRNA can also be determined.

 The strong interdependence of secondary structure and function of the rRNA molecules is well known. Indeed, evolutionary changes in the primary sequence of the
20 rRNA are effectively restricted such that secondary structure of the molecule will be maintained. For example, if a base is changed on one side of a helix of a rRNA molecule, then a compensating change will be made on the other side of the helix to preserve complementarity (this is referred to as covariance). This relationship allows two very different rRNA sequences to be "aligned" based on conserved primary sequence and
25 conserved elements of the secondary structure. Once the sequences have been aligned, it becomes possible to identify conserved and variable regions of the rRNA or rDNA sequence.

 Variable regions of rRNAs were identified by comparative analysis using published rRNA sequences and sequences that were determined during the development of the present
30 invention. Commercially available software can be used or adapted for the purposes disclosed herein. Since the sequence evolution at each of the variable regions (for example, spanning a minimum of 10 nucleotides) of rRNA is, for the most part, divergent and not

WO 2005/017202

PCT/US2004/014690

convergent, probes can be based on a few rRNA sequences which differ between the target organism and its phylogenetically closest relatives.

Preferred probes for detecting the rRNA or rDNA of bacteria in the genus *Staphylococcus* have sequences of up to 100 nucleotides in length and have at least 17 contiguous nucleotides, more preferably 30 contiguous nucleotides, and still more preferably 39 contiguous nucleotides contained in the sequence given by GCGATTCCAGCTTCATGTAGTCGAGTTGCAGACTACAATCCGAACTGAGAACA ACTTTATGGGATTTGCTTGACCTCGCGGTTTCG (SEQ ID NO:13), e.g., CCGAACTGAGAACAACCTTTATGGGATTTGC (SEQ ID NO:10), while CCACTCAAGAGAGACAACATTTTCGACTAC (SEQ ID NO:7) is specific for *S. aureus* rRNA or rDNA. Notably, these probes do not hybridize to ribosomal nucleic acids of bacteria in the genus *Enterococcus* under high stringency hybridization conditions. Preferred oligonucleotide sequences include RNA and DNA equivalents, and may include at least one nucleotide analog.

Preferred hybridization probes for detecting the rRNA or rDNA of bacteria in the group *Enterococcus* have the sequences of CTCCTAGGTGCCAGTCAAATTTTG (SEQ ID NO:14) or CATCATTCTCAATTCCGAGGC (SEQ ID NO:15). Notably, these probes do not hybridize to ribosomal nucleic acids of bacteria in the genus *Staphylococcus* under high stringency hybridization conditions. In certain embodiments of the invention, the hybridization probes are labeled with a detectable label which allows detection independent of the helper oligonucleotides. Preferred oligonucleotide sequences include RNA and DNA equivalents, and may include at least one nucleotide analog.

Preferred pan-bacterial hybridization probes for detecting the rRNA or rDNA of a plurality of bacterial species have the sequences of CGACAAGGAATTTTCGC (SEQ ID NO:16) (which may be synthesized using 2'-methoxy nucleotide analogs, and which may be used in conjunction with helper oligonucleotides having the sequences of TACCTTAGGACCGTTAT (SEQ ID NO:17) and CAGGTCGGAACCTTACC (SEQ ID NO:18)), or GGAACCTACCCGACAAGGAATTTTCGCTACCTTAGG (SEQ ID NO:19) (which may be used in conjunction with helper oligonucleotides having the sequences of ACCGTTATAGTTACGGCCGCGTTTACCGGGGCTTC (SEQ ID NO:20), GCCTGGCCATCATTACGCCATTCGTGCAGGTC (SEQ ID NO:21) and GCCCAAATCGTTACGCCTTTCGTGCGGGTC (SEQ ID NO:22)). Probes having the

sequences of SEQ ID No:16 and SEQ ID NO:19 are useful for detecting the ribosomal nucleic acids of a plurality of species of Gram-positive bacteria (including the high(G+C)subset of Gram-positive bacteria), a plurality of species of bacteria in the family Enterobacteriaceae, a plurality of species of bacteria in the genus Enterococcus, and a plurality of species of bacteria in the genus Staphylococcus. These probes do not hybridize to the nucleic acids of fungal organisms. In certain embodiments of the invention, the hybridization probes are labeled with a detectable label which allows detection independent of the helper oligonucleotides. Preferred oligonucleotide sequences include RNA and DNA equivalents, and may include at least one nucleotide analog.

10 IV. Helper Oligonucleotides

Hybridization between one or more labeled oligonucleotide probes and a target polynucleotide can be enhanced through the use of unlabeled "helper oligonucleotides" according to the procedure disclosed in U.S. Patent No. 5,030,557. Helper oligonucleotides bind a region of the target nucleic acid other than the region that is bound by the assay probe. This binding imposes new secondary and tertiary structures on the targeted region of the single-stranded nucleic acid and accelerates the rate of probe binding, and, in some cases, the extent of probe binding.

For example, rRNA possesses tertiary structure arising from the same kind of attractive forces that order duplex DNA into its now well known helical structure. A substantial portion of this secondary and tertiary structure is not lost under conditions normally employed for nucleic acid hybridization, including elevated temperature conditions, the presence of salt, the presence of accelerators, and the like. This residual structure can sterically inhibit, or even block hybrid formation between a nucleotide multimer, for example a DNA or RNA oligomer being used as a probe, and its complementary sequence in the ribosomal RNA or other single stranded nucleic acid such as mRNA or DNA which the probe targets. This inhibition can be reduced or eliminated by use of a "helper" oligonucleotide which binds to a portion of the RNA or DNA other than the portion being targeted by the probe. Interaction with the helper oligonucleotide imposes a new secondary and tertiary structure on the targeted region of the single stranded nucleic acid, and consequently accelerates the rate of probe binding. Thus, by using a properly selected helper oligonucleotide, the rate of hybridization between the probe and its complementary sequence in the targeted nucleic acid can be substantially increased and

WO 2005/017202

PCT/US2004/014690

even permit hybridization to occur at a rate and under conditions otherwise adequate for an assay where, without the use of the helper, no substantial hybridization can occur.

The use of a helper can raise the T_m of the hybrid of a relatively short probe and its intended target relative to the T_m of the hybrid of the probe and a nucleic acid sequence with which the probe is less complementary. As a result, highly specific assays for organisms
5 which occur in environments populated by closely related organisms can be obtained.

Helper oligonucleotides which can be used in combination with detectably labeled oligonucleotide probes are preferably 17 to 100 nucleotides in length. Helper oligonucleotides for use in combination with the above-described *Staphylococcus* genus
10 ribosomal nucleic acid probe include, but are not limited to: UUGACCUCGCGGUUUCG (SEQ ID NO:11) and GCGATTCCAGCTTCATGTAGTCGAGTTGCAGACTACAAT (SEQ ID NO:12) (see WO 00/667189). Helper oligonucleotides for use in combination with a ribosomal nucleic acid probe specific for *S. aureus*, include but are not limited to: GATGATTCGTCTAATGTCGACCTTTGTA ACTCC (SEQ ID NO:8) and
15 CGGAATTTACGTGCTCCGTCGTA CT CAGGAT (SEQ ID NO:9). Helper oligonucleotides that can be used in combination with the above-described probe of sequence CTCCTAGGTGCCAGTCAAATTTTG (SEQ ID NO:14) for detecting the ribosomal nucleic acid of members of the *Enterococcus* genus include, but are not limited to: TCTACGGGGCTTTTACCCTTTCTAGCAGACC (SEQ ID NO:23) and
20 CCTCGTGTTCCGCCGTA CT CAGGATC (SEQ ID NO:24). Helper oligonucleotides that can be used in combination with the above-described probe of sequence CATCATTCTCAATTCCGAGGC (SEQ ID NO:15) for detecting the ribosomal nucleic acid of members of the *Enterococcus* genus include, but are not limited to: TAGCCCTAAAGCTATTTTCGGAGAGAACCAGCTATCTCC (SEQ ID NO:25) and
25 CCCTAGTCCAAACAGTGCTCTACCTC (SEQ ID NO:26). Helper oligonucleotides that can be used in combination with the pan-bacterial probe having the sequence of CGACAAGGAATTTTCGC (SEQ ID NO:16) include, but are not limited to: TACCTTAGGACCGTTAT (SEQ ID NO:17) and CAGGTCGGAACTTACC (SEQ ID NO:18). Helper oligonucleotides that can be used in combination with the pan-bacterial
30 probe having the sequence of GGA ACTTACCCGACAAGGAATTTTCGCTACCTTAGG (SEQ ID NO:19) include, but are not limited to: ACCGTTATAGTTACGGCCGCCGTTTACCGGGGCTTC (SEQ ID NO:20),

WO 2005/017202

PCT/US2004/014690

GCCTGGCCATCATTACGCCATTCGTGCAGGTC (SEQ ID NO:21) and
 GCCCAAATCGTTACGCCTTTCGTGCGGGTC (SEQ ID NO:22). In certain
 embodiments of the invention, the hybridization probes are labeled with a detectable label
 which allows detection independent of the helper oligonucleotides. Preferred
 5 oligonucleotide sequences include RNA and DNA equivalents, and may include at least one
 nucleotide analog.

V. Probe Matrix

Results from a probe matrix hybridization procedure can be inputted into a
 computer or data processor ("computer"), either manually using a keyboard or directly
 10 through an interface from an automated device such as a plate reader, film scanner or
 luminometer. The computer can sort the positive and negative hybridization results for a
 particular sample to establish a profile that can be compared with a look-up table stored in a
 memory device linked to the computer. This facilitates associating the hybridization profile
 with hybridization results obtained using control organisms in order to determine the
 15 identity or candidate identity in the case of ambiguous results that are characteristic of more
 than one organism, and/or the presence or absence of a gene of interest, in the test
 organism.

It is generally true that the presence of a microorganism in a biological sample will
 be indicated if the biological sample also contains rRNA or rDNA that is characteristic of
 20 the microorganism. Thus, the presence of a particular rRNA or rDNA in a biological
 sample is diagnostic of the presence of a microorganism that produces that rRNA or rDNA.
 If a hybridization reaction gives a positive result with a probe specific for Gram-positive
 bacteria, that result would indicate the presence of one or more species of Gram-positive
 bacteria in the biological sample. In contrast, a negative result would indicate the absence
 25 of Gram-positive bacteria.

Staphylococcus genus probe can be used to identify organisms as members of the
 broad genus of Staphylococcal bacteria, and a species-specific probe can be used
 independently to identify *Staphylococcus aureus*. Likewise, one or more *mecA*-specific
 probes can be used to identify the *mecA* status of a sample.

30 *Enterococcus spp.* probes, such as those described herein, can be used for detecting
 bacterial species that include *E. faecalis* and *E. faecium*. One or more probes able to
 hybridize the VanA and/or VanB genes can be used to identify the vancomycin-resistance

WO 2005/017202

PCT/US2004/014690

status of a sample.

Of course, a series of positive and negative control hybridizations can be carried out in parallel to ensure validity of the testing results.

VI. Apparatus Useful for Conducting Hybridization Reactions

5 Examples of formats that can be used to conduct hybridization reactions include, but are by no means limited to: individual tubes each with a different probe or comprising a plurality of probes; the wells of a 96-well or other multi-well microtiter plate; and a solid support such as a dipstick or a "DNA chip" where polynucleotide probes are immobilized to the support at different addresses in a spaced-apart configuration. In one preferred
10 embodiment, the probes are soluble probes. In another preferred embodiment, the probes are immobilized to a solid support. Generally speaking, it is preferred for the various probes used in an apparatus for identifying antibiotic-resistant microorganisms to function under a common set of temperature conditions so that all of the probes can be disposed on a single device. In certain highly preferred embodiments, the probes are used under a single
15 set of hybridization conditions, including a single temperature and/or a single set of ionic strength conditions.

Identifying microorganisms and/or the presence of gene(s) of interest advantageously can be performed without requiring any *in vitro* amplification step. Alternatively, a preliminary amplification step, for example using transcription mediated
20 amplification (TMA), may be employed (see, e.g., U.S. Patent No. 5,399,491).

Regardless of whether the probes are soluble or immobilized, in certain preferred embodiments the different probe addresses are maintained spatially separated from each other. For example, probes for detecting Van A target nucleic acids can be disposed in a tube, microtiter well, or immobilized spot in an array that is physically distinct from the
25 tube, microtiter well, or immobilized spot having probes for detecting Van B target nucleic acids. Likewise, for example, probes for detecting the ribosomal nucleic acids of *Staphylococcus aureus* can be physically separated from the probes used for detecting the ribosomal nucleic acids of bacteria in the genus *Enterococcus*. When all of the probe addresses are maintained physically separated from each other, and when the probes are
30 soluble probes rather than probes immobilized in a microarray format, the number of hybridization reactions will be increased. Whereas four hybridization reactions were used to conduct the procedure illustrated in Example 13 (which included ambiguous probe

WO 2005/017202

PCT/US2004/014690

addresses for detecting either (a) ribosomal nucleic acids of *S. aureus* or bacteria in the genus *Enterococcus* and (b) Van A or Van B nucleic acids), the number of hybridization reactions would be increased to six if the different probe addresses were separated from each other. In this instance the probes of the six probe addresses would hybridize to nucleic acids from: (1) bacteria in the *Staphylococcus* genus; (2) *S. aureus* bacteria; (3) bacteria in the genus *Enterococcus*; (4) the *mecA* gene; (5) the Van A gene; and (6) the Van B gene. Of course, each of these six probe addresses could correspond to a spot in a microarray if the probes are immobilized, and then a single hybridization reaction could be conducted. For example, the different spots in the array could be contained in a single well of a 96-well microtiter plate.

According to one approach for conducting hybridization procedures, probes can be labeled with distinguishable labels. Examples of particularly preferred chemiluminescent labels that can be used for performing the methods described herein are the acridinium ester (AE) labels disclosed in U.S. Patent No. 5,756,011, the disclosure of which is hereby incorporated by reference. More particularly, a single tube, well, support or address may include distinct probes that are independently labeled with chemiluminescent labels that emit peak energy at different times after generating a light emission. Materials and methods that can be used for making and using distinguishable probes useful in connection with the present invention can be found in U.S. Patent No. 5,756,011, the disclosure of which is hereby incorporated by reference. Fluorescent labels that produce light at different wavelengths following excitation represent still other examples of distinguishable labels that can be used in connection with the procedures described herein. In this way, two probes that employ distinguishable labels can be distinguished from each other even though they are combined at the same locus of a testing device. Accordingly, it is possible to combine large numbers of different probes at a single address while still being able to distinguish the results of hybridization for the different probes or sets of probes.

In accordance with still another approach, probes with different specificities were labeled with the same label, or with labels that were not distinguished from each other during a step for detecting specific hybrids that include the labeled probe and target, thereby creating an ambiguous result when the probes were combined and when either probe gave a positive hybridization signal.

WO 2005/017202

PCT/US2004/014690

VII. Kits for Conducting Hybridization Procedures

The materials used for carrying out hybridization procedures in accordance with the invention may be incorporated into kits that can be used for conducting diagnostic procedures. The kits will include at least one device or container containing a plurality of probes for hybridizing nucleic acids from test organisms, and instructions for conducting a nucleic acid hybridization procedure using the probes. The kits optionally may include instructions for detecting specific hybrids between probes that comprise the various addresses, e.g., *Staphylococcus* and/or *Enterococcus* genes, and target genes of interest, e.g., *mecA*, *VanA* and/or *VanB*, obtained from biological samples that undergo testing. In certain embodiments, there also will be included one or more hybridization probes for detecting a broad range of bacterial species.

VIII. Exemplary Detection Assays

In certain preferred embodiments, Hybridization Protection Assays (HPA) may be used to detect an analyte in a medium, where the analyte is part of a specific binding pair. When the medium suspected of containing the analyte is combined with a binding partner, a detectable label attached to the binding partner is capable of undergoing a change in stability or differential degradation whenever the analyte binds to the specific binding partner. In a specific embodiment, single-stranded polynucleotide probes are modified to contain labels at virtually any desired position or location on the probe. In one embodiment, probe labels may be of different stability or susceptible to differential degradation depending on whether the target polynucleotide sequence is hybridized to the probe. In one embodiment, the label on the bound probe is stabilized relative to the label on the unbound probe.

First, binding partners comprising a binding substance and one or more binding partners are selected. These pairs may be polynucleotides or oligonucleotides. In one embodiment, the binding partners are a polynucleotide and one or more oligonucleotides.

Second, the assay format is selected. These may be selected from formats comprising direct binding assays, competition assays, sequential saturation methods, and sandwich assays.

Third, a label is selected for the assay to be performed. This may be a label which can be directly or indirectly detected by colorimetric, fluorimetric, chemiluminescent, or bioluminescent means. The label may have the property that it can be chemically or

WO 2005/017202

PCT/US2004/014690

biochemically degraded so as to modify its ability to be detected, said degradation being possible under conditions which do not adversely effect the binding between the labeled binding partner and its binding substance and other binding partners and binding substances which may participate in the reaction. Preferred labels are ones which are affected in their
5 ability to be detected after exposure to acids, bases, or selective oxidizing agents such as peroxidate, or enzymes.

Fourth, using chemical methods known in the art, the label is attached to the binding substance at a site such that the sensitivity of the label to chemical or biochemical degradation is modified upon interaction of the labeled binding partner with its specific
10 binding substance(s). In some cases several different sites may be tested for label attachment and the site which gives the best differential degradation may be used.

Fifth, the degradation conditions, be they chemical or biochemical, are optimized if needed to give the best detection discrimination of the labeled binding partner in the presence and absence of its binding substance.

15 Methods for preparing acridinium ester labeled DNA probes and conditions for differential hydrolysis are described in Arnold et al. (*Clin. Chem.* **35**:1588 (1989)) and U.S. Patent No. 6,004,745.

Finally, using the preselected assay format, the ability of the assay system to detect quantitatively or qualitatively the analyte generally employing the steps of: incubating,
20 selectively degrading, and detecting or simultaneously incubating and selectively degrading, and detecting.

Oligonucleotide probes labeled with chemiluminescent acridinium esters are particularly useful for the detection of sequence specific polynucleotides through hybridization. Acridinium esters may be attached at a number of different sites on DNA
25 probes and/or mixed nucleotide/non-nucleotide polymers as described in U.S. application Serial No. 07/099,050 filed September 21, 1987. This includes the ability to label the nucleotide bases, the phosphate backbone, the 3' terminus, and the 5' terminus of oligonucleotides as well as the non-nucleotide monomeric units of mixed nucleotide/non-nucleotide polymers. Such acridinium ester labeled probes can show significant differential
30 chemical stability when the probes to which they are attached are free in solution as compared to when they are hybridized.

In one embodiment, at least two probes in a single hybridization reaction are labeled

WO 2005/017202

PCT/US2004/014690

with detectable moieties which are not substantially distinguished from each other during a step for detecting specific hybrids following a hybridization reaction. For example, probes having different specificities may even be labeled with identical detectable labels when the probes are soluble probes that are mixed for use in a single hybridization reaction.

5 In another embodiment, at least two probes in a single hybridization reaction are labeled with detectable moieties which are distinguishable, e.g., each label comprises components capable of taking part in a distinguishable chemiluminescent reaction (see for instance, U.S. Patent Nos. 5,656,207 and 5,827,656). Different members of a number of classes of chemiluminescent molecules are capable of exhibiting differences in kinetic
10 and/or spectroscopic properties and can hence be used in the invention, including acridinium and related compounds (e.g., phenanthridinium compounds), phthalhydrazides and related compounds (e.g., naphthalhydrazides), oxalate esters and related compounds and also stabilized dioxetanes and dioxetanones. The variations of compounds within such groups are well-known to those of ordinary skill in the art, likewise it is known that the
15 quantum yield, kinetics and emission wavelengths of their chemiluminescent reactions are affected by their structure. Thus, compounds with high quantum yields, and which, relative to each other possess substantial differences in their reaction rates or their emission wavelengths, in order to maximize the resolution between the detection of these compounds, can be employed. For instance, aryl acridinium esters may be used as labels
20 with appropriate chemical modifications made to produce the desired kinetic and spectroscopic parameters (see, e.g., U.S. Patent No. 5,656,207).

Thus, probes specific for different genes can be labeled with different labels, the labeled probes can be mixed and allowed to hybridize to any nucleic acid contained in the test sample having a sequence sufficiently complementary to the probe sequence to allow
25 hybridization under appropriately selective conditions. In one embodiment, one or more reagents are added to the solution which will specifically alter the labeling reagent associated with unhybridized labeled probe while leaving the labeling reagent associated with the hybridized probes substantially unaltered. This allows each labeling compound to be differentially resistant to loss of chemiluminescent potential depending on whether the
30 label is associated with a hybridized or unhybridized probe. In a preferred embodiment, the hybridized probe associated label is so protected (see, for instance, U.S. Patent No. 5,827,656).

Such labeling reagents are particularly useful in, although not limited to, homogeneous assay systems in which analytes of interest may be detected and measured without the need for the analyte-bound label to be physically separated from the unbound label prior to detection. However, such reagents may be used in heterogeneous systems or in combinations of homogeneous and heterogeneous assay systems as well.

Notably, oligonucleotides, or the complements thereof, which are disclosed herein as probes also may be used as primers in amplification reactions. For example, primers having the target-complementary sequences of SEQ ID NO:4 and SEQ ID NO:6 were used in PCR reactions to amplify *mecA* sequences. Example 11 illustrates the use of probe sequences as primer sequences to amplify nucleic acids diagnostic of resistance to vancomycin.

The invention will be further described by the following non-limiting Examples.

Example 1

Initial Evaluation of Probes for Detecting the *mecA* Gene

Two probes were designed to hybridize at nucleotides 501 to 531 of the *mecA* gene sequence: probe 2, CGCTATGATCCCAATCTAACTTCCACATACC (SEQ ID NO:2) was designed to hybridize one strand of the *mecA* gene, and probe 1, GGTATGTGGAAGTTAGATTGGGATCATAGCG (SEQ ID NO:1), was designed to hybridize to the opposite strand of the *mecA* gene. The probes were synthesized by standard phosphoramidite procedures with an internal non-nucleotide linker inserted between nucleotide positions 21 and 22, and between positions 13 and 14, respectively using the procedure disclosed in U.S. Patent No. 5,656,744. The probes were labeled with an acridinium ester (AE) according to the method disclosed in U.S. Patent No. 5,185,439 and used for testing biological samples.

Two *S. aureus* isolates were obtained from an outside laboratory which reported one specimen (#99) to be *mecA*-positive by a diagnostic PCR assay (Murakami and Minamide, In: Diagnostic Molecular Microbiology, D. Pershing et al., Eds., American Society for Microbiology, 1993, pp. 539-541) and the other (#100) *mecA*-negative by PCR. Both specimens were reported to have an Oxacillin Minimum Inhibitory Concentration ("Ox MIC") of 4 µg/ml, a value considered unusually high for a *mecA*-negative specimen.

The specimens were lysed by suspending approximately 5×10^8 cells in a lysis buffer (0.1% lithium lauryl sulfate, 1 mM EDTA, 20 mM lithium succinate, pH 5.5) and

WO 2005/017202

PCT/US2004/014690

then incubating in a heat block set at 110°C for 20 minutes. Thermal lysis of a variety of other specimens was effected under a range of temperature (heat block set at about 95°C to 110°C) and time (about 10 to 25 minutes) conditions. The lysed samples were tested for the presence of target sequences complementary to the probes using a Hybridization Protection Assay (HPA) as described by Arnold et al., in *Clin. Chem.* **35**:1588(1989) and U.S. Patent No. 6,004,745. 50 µl of the heat-treated suspension was incubated with an equal volume of AE-labeled probe in 2x hybridization buffer (1.2 M LiCl, 2% lithium lauryl sulfate, 20 mM EDTA, 20 mM EGTA, 100 mM lithium succinate, pH 5.5) at 60°C for 30 minutes. The pH of succinate solutions may range from about 4.5 to about 5.5, although it is desirable to use a pH of 4.8 for succinate buffering. 300 µl of a selection reagent (0.6 M borate buffered solution, pH 8.5; note, pH may vary from 8.0 to 9.0) was added, the mixture was heated at 60°C for 10 minutes to effect differential hydrolysis, and the samples were read in a GEN-PROBE® LEADER® luminometer using sequential 300 µl injections of 0.1% H₂O₂ and 1 N NaOH solutions (see Arnold et al, *supra*; GEN-PROBE® Detection Reagent Kit, Cat. No. 1791 may be used).

Representative test results for the two probes with specimens #99 and #100 are provided in Table 1.

Table 1Probe Hybridization Confirms the Results of Diagnostic PCR Assays

Test Sample	<i>mecA</i> PCR	Ox MIC (µg/ml)	AE-probe Average RLU	
			Probe 1	Probe 2
No Specimen	--	--	431	3062
#100	Neg	4	590	5585
#99	Pos	4	2664	8072

The results presented in Table 1 showed that specimen #100 was *mecA*(-) and that specimen #99 was *mecA*(+), thereby confirming the results obtained using a diagnostic PCR assay. However, the cell load used in this Example was near the limit of detection for direct probe detection. This was more noticeable in the trial that included probe 2, as evidenced by a high background signal that made interpretation somewhat difficult at this level of cell load. Conversely, probe 1 was found to give more definitive results, perhaps as the result of a substantially lower background signal.

Example 2**Use of a Pan-Bacterial Probe as a Positive Control**

Procedures similar to those described in Example 1 were conducted using a large number of bacterial samples, except that parallel hybridization reactions employed a pan-bacterial probe which targeted rRNA (Probe Reagent 1, GEN-PROBE® MTC-NI Kit, catalog no. 4573). The PCR-negative specimens had abnormally high Ox MIC levels, and one PCR-positive specimen had a very low Ox MIC value. The pan-bacterial probe reagent served as a positive control for lysis of all of the specimens. In this procedure each sample was tested against a panel of the three probes contained in separate hybridization reactions. Aliquots of the samples tested using the pan-bacterial probe reagent were diluted 1:3000 prior to performing luminometry so that results fell within the linear range of the detector. The signals indicated lysis and detection of ribosomal nucleic acids in each of the samples. These values depended on the number of bacterial cells in the specimen, as well as on the extent of cell lysis and the release or accessibility of ribosomal nucleic acids from the cells to be detected by the pan-bacterial probe.

The results for various specimens are shown in Table 2.

Table 2Hybridization Results Obtained Using *mecA* and Pan-Bacterial Probes

Test Sample	<i>mecA</i> PCR	Ox MIC $\mu\text{g/ml}$	AE-Probe Signal (RLU)		
			Probe 1	Probe 2	Pan-Bacterial*
Negative Control	NA	NA	1817	1931	350
rRNA Pos. Control	NA	NA	NA	NA	52141
10	Neg	8	2918	3274	50408*
100	Neg	4	815	6186	24006*
101	Neg	4	2768	6266	45639*
103	Neg	4	1097	2366	27138*
104	Neg	2	959	2000	31212*
105	Neg	2	1086	2714	13699*
106	Neg	2	1327	2989	32059*
145	Neg	4	997	2788	19564*
155	Neg	4	730	2257	41384*
164	Neg	2	1546	3287	41698*
99	Pos	4	5822	13021	35354*
102	Pos	4	5623	13065	36721*
139	Pos	4	5755	9563	29994*
140	Pos	16	3143	4967	17506*
142	Pos	4	6845	10345	29905*
148	Pos	4	10483	15878	66145*
178	Pos	2	7772	12652	41643*
188	Pos	2	11856	20757	41605*
270	Pos	<0.25	9256	14965	23183*
296	Pos	Lab Pos control	5567	10585	23240*

*(diluted 1:3000 prior to luminometry)

The results presented in Table 2 showed that both of the *mecA*-specific probes gave

WO 2005/017202

PCT/US2004/014690

positive, but relatively weak, signals for specimens that were positive for the *mecA* gene by the diagnostic PCR assay. Equivocal results that were obtained using probe 2 with one PCR-positive and two PCR-negative specimens may have been due, in part, to the relatively high background of probe 2 and to the sensitivity limitation imposed by using a single probe to test the specimens at the levels of cells in the sample. Additionally, the *mecA* probes required denaturation and accessibility to ssDNA containing the gene target region to obtain the optimal detection signal. In contrast, rRNA detected by the pan-bacterial probe reagent is available in substantial quantities per cell compared to DNA, and the accessibility of the rRNA target region for hybridization and detection by the probe reagent was assisted by the inclusion of helper probes (Hogan et al., U.S. Patent No. 5,030,557) in that reagent. Detection of rRNA would not be indicative of the amount or reactivity of a ssDNA target region with regard to accessibility or kinetics of hybridization. Instead, it would indicate success of the lysis procedure.

Example 3

15 Base Treatment of Specimens

Treatment of specimens under alkaline conditions was investigated as a means for denaturing nucleic acids and making target sequences available for hybridization with the detection probe(s). Base treatment offered the further possible advantage of hydrolyzing RNA so that positive internal control hybridization probes more properly detect ribosomal DNA (rDNA) targets, thereby permitting more direct comparison between results obtained using the *mecA* probe and the control probe. Base treatment to lyse cells and degrade RNA depends on hydroxide concentration, temperature and incubation time. Notably, probes designed to detect rRNA are also capable of detecting the corresponding rDNA target.

The effect of base treatment on a *S. aureus* specimen (#102) was evaluated using the pan-bacterial probe reagent. A pelleted sample of the specimen was suspended in 120 μ l of lysis buffer and lysed by treatment for 20 minutes in a heat block set at 110°C. A 50 μ l aliquot of the lysed specimen was treated with 50 μ l of a 0.4 N LiOH solution at 55°C for 10 minutes, and then neutralized with the addition of 50 μ l of 0.4 N HCl and buffered with 50 μ l of a 1x hybridization buffer (0.6 M LiCl, 1% lithium lauryl sulfate, 10 mM EDTA, 10 mM EGTA, 50 mM lithium succinate, pH 5.5). The treated sample was hybridized with the AE-labeled pan-bacterial probe by incubating at 55°C for 1 hour. An HPA assay to detect specific probe hybrids was carried out using standard procedures that will be familiar to

those having an ordinary level of skill in the art. A control sample consisted of 50 μ l of the lysed specimen and volumes of hydrolysis, neutralization and buffered solutions equivalent to those of the treated sample.

The results of base treatment of a *S. aureus* specimen using a pan-bacterial probe reagent are shown in Table 3.

Table 3

Base Treatment Reduces the
Hybridization Signal Obtained Using the Pan-Bacterial Probe

Sample	Sample Volume Read	Pan-Bacterial Probe Signal (RLU)
#102, Control (no base treatment)	50 μ l †	483845
#102, Base Treated	300 μ l	134426

† 1/6 of the volume was used to bring the control signal into the range of the luminometer.

The results presented in Table 3 showed evidence for significant, but perhaps incomplete, degradation of the rRNA target under the condition tested since both rDNA and residual rRNA targets could have contributed to the chemiluminescent signal measured in the hydrolyzed sample. Higher base concentrations, longer incubation times, and higher temperatures showed increased degradation of the signal. For example, the effect of base treatment with 0.8 N LiOH for 10 minutes at 55°C was assessed using AE-probe 1. For these treatments, an equal volume of 0.8 N LiOH was added to the sample, and the reaction mixtures were neutralized with the addition of an equivalent volume of 0.8 N HCl prior to buffering and assaying by HPA. 300 μ l aliquots of each reaction mixture were read in the luminometer. These procedures employed a single-stranded DNA oligonucleotide having the same sequence that was present in the target bacteria, and that was fully complementary to the hybridization probe undergoing testing, as synthetic target.

Table 4Enhanced Base Treatment of the TargetDoes Not Compromise Hybridization of the *mecA* Probe

Sample	AE-probe 1 Signal (RLU)
Synthetic target, Control (no base treatment)	22195
Synthetic target, base treated	21283
#148, Control (no base treatment)	34677
#148, base treated	42627
Synthetic target + #148, Control	53367
Synthetic target + #148, base treated	58565

The results shown in Table 4 indicated that there was very little, if any, loss of signal due to degradation of target DNA sequences under the base treatment conditions that were tested.

Example 4Base Treatment and Lysis

The effect of base treatment with and without the heat treatment in lysis buffer was evaluated using a *S. aureus* specimen (GP1217, ATCC #33591). In this procedure the sample was treated with 0.8 N LiOH and thermal lysis was conducted by heat treatment for 15 minutes. Other conditions were as described above. Probe 1 labeled with acridinium ester was used in this procedure.

Table 5Bacterial Sample Preparation

Sample	Probe 1 RLU	Pan-Bacterial RLU
Negative Control	592	3891
Positive Control	nd	51752
GP1217, Thermal lysis 15 minutes at 100°C + Base treatment 15 minutes at 60 °C	38017	514600
GP1217, Base treatment 30 minutes at 60°C	16016	66871

The results presented in Table 5 indicated that base treatment under the specified

WO 2005/017202

PCT/US2004/014690

conditions was adequate to lyse cells and render the target sequence available for detection by the labeled hybridization probe.

A range of base treatment conditions were found to be effective for this purpose. For example, hydroxide at a final concentration of about 0.2-0.4 N, temperatures of about 55°C to about 60°C, and incubation times of about 5 to 30 minutes all gave good results in the procedure. Accordingly, a wide range of conditions may be used for base treatment of specimens including, but not limited to, final base concentrations of about 0.2 N to about 1 N in hydroxide, temperatures from about 50°C to about 60°C, and incubation times of about 5 minutes to about 30 minutes. Of course, conditions involving lower hydroxide concentrations and higher temperature incubations are contemplated as part of the invention. Generally speaking, a temperature range of from about room temperature up to about 100°C with different base concentrations are contemplated for sample preparation steps that fall within the scope of the invention.

Example 5

Internal Controls

Staphylococcus aureus probe and helpers

Probe

SauB338: CCACTCAAGAGAGACAACATTTTCGACTAC (SEQ ID NO:7).

SauB338-AE(20/21): This probe had the sequence of SEQ ID NO:7, and was labeled with acridinium ester attached to a non-nucleotide linker inserted between nucleotide positions 20 and 21

Helpers

SauB278: GATGATTCGTCTAATGTCGACCTTTGTA ACTCC (SEQ ID NO:8)

SauB368: CGGAATTTACGTGCTCCGTCGTACTCAGGAT (SEQ ID NO:9)

Staphylococcus genus probe and helpers

Probe

SauA1276: CCGAACTGAGAACA ACTTTATGGGATTTGC (SEQ ID NO:10).

SauA1276-AE(19/20): This probe had the sequence of SEQ ID NO:10, and was labeled with acridinium ester attached to a linker inserted between nucleotide positions 19 and 20.

Helpers

MeO-SauA1259: UUGACCUCGCGGUUUCG (SEQ ID NO:11)

(2-methoxy nucleotide analogs were used to synthesize this helper oligonucleotide)

WO 2005/017202

PCT/US2004/014690

SauA1306: GCGATTCCAGCTTCATGTAGTCGAGTTGCAGACTACAAT (SEQ ID NO:12)

Methicillin-resistant (*mecA*-positive) *S. aureus* ("MRSA") isolates GP1217 (ATCC #33591) and GP1218 (ATCC #43866), and methicillin-susceptible (*mecA*-negative) *S.*
 5 *aureus* ("MSSA") isolates GP1214 (ATCC #29247) and GP18 (ATCC #12600), were grown on TSA plates (Hardy Diagnostics, Cat. No. A-10) overnight at 37°C and used as specimens for testing the *S. aureus* probe and helpers identified by SEQ ID NOs:7-9 (Milliman, U.S. Patent No. 5,292,874), and the *Staphylococcus spp.* probe and helpers identified by SEQ ID NOs:10-12 (Hogan et al., U.S. Patent No. 6,376,186) as internal
 10 controls for adequate presence of target organisms in the specimen and to determine whether the specimen is *S. aureus* or a coagulase-negative *Staphylococcus spp.* ("CoNS").

The cultured *S. aureus* specimens were lysed by suspending 1 µl loops of bacteria in 150 µl of lysis buffer and then incubating at 105°C for 15 minutes. Next, 100 µl of 2 N LiOH base solution was added and RNA was hydrolyzed by incubating the mixtures at
 15 60°C for 15 minutes. Hydrolyzed samples were neutralized by the addition of 100 µl of a neutralizing reagent that included 2 N HCl and 100 mM succinate buffer (pH 5.5 prior to acidification with HCl). Specifically hybridized probe was detected by standard HPA, essentially as described in Example 1.

Test results using *S. aureus* and *Staphylococcus* genus probes with helper probes for
 20 four specimens are provided in Table 6.

Table 6

Taxonomic Probes Used in Combination with a *mecA*-Specific Probe Provide Greater Functionality in Probe Panels

Probe Composition	GP1217(Pos) RLU	GP1218(Pos) RLU	GP1214(Neg) RLU	GP18(Neg) RLU
25 probe 1 (<i>mecA</i>)	53,322	36,730	848	821
<i>S. aureus</i> probe with helpers	199,476	163,825	203,725	148,780
<i>Staphylococcus spp.</i> probe with helpers	225,772	203,488	234,286	170,434

30

The results presented in Table 6 exactly matched the expected hybridization patterns for all four samples. More specifically, the *Staphylococcus spp.* probe, that was specific for

WO 2005/017202

PCT/US2004/014690

a plurality of members of the *Staphylococcus* genus, gave positive hybridization signals for all samples. This indicated that all samples contained *Staphylococcus* bacteria, as expected. Additionally, all four samples gave positive hybridization signals when hybridized with the probe specific for *S. aureus*. This indicated that all samples contained *S. aureus* bacteria.

5 Finally, only the *mecA* positive specimens and not the *mecA*-negative specimens gave positive hybridization signals with *mecA*-specific probe 1.

These results defined a highly desirable panel of hybridization probes useful for determining the identity and methicillin resistance status of bacteria undergoing testing. In one embodiment, the invention includes a probe that is specific for a plurality of bacteria in
10 the *Staphylococcus* genus, including *S. aureus* and *S. epidermidis*. It is highly preferred that this probe is specific for ribosomal nucleic acids of the target bacteria. Also present in the panel is a probe that specifically hybridizes to the nucleic acids of *S. aureus*, but not to the nucleic acids of other bacteria in the *Staphylococcus* genus, such as *S. epidermidis*. It is highly preferred that this probe is specific for ribosomal nucleic acids of *S. aureus*. Further,
15 it is preferred that the panel include a probe that hybridizes to the *mecA* gene or the RNA transcript encoded by the *mecA* gene. This *mecA*-specific probe should not hybridize to other drug resistance markers, such as the genes encoding resistance to vancomycin.

As indicated in the following Example, alternative probe sequences can be used for detecting the *mecA* gene in a sample containing nucleic acids.

20

Example 6

Evaluation of Additional *mecA* Gene Probes

Probes

Probe 3: GCGATAATGGTGAAGTAGAAATGACTGAACGTCCG (SEQ ID NO:3)

25

Probe 4: CGGACGTTTCAGTCATTTCTACTTCACCATTATCGC (SEQ ID NO:4)

Probe 5: GCTCCAACATGAAGATGGCTATCGTGTCAACAATCG (SEQ ID NO:5)

Probe 6: CGATTGTGACACGATAGCCATCTTCATGTTGGAGC (SEQ ID NO:6)

30

To increase the sensitivity of *mecA* detection, additional probes were designed that corresponded to nucleotide bases 304-338 and 1010-1044 of the *mecA* gene. These probes

WO 2005/017202

PCT/US2004/014690

flanked probe 1 and probe 2. Probe 3 (SEQ ID NO:3, complementary to the antisense (-) strand of the *mecA* gene) was labeled with AE attached by a linker positioned between bases 16 and 17. Probe 4 (SEQ ID NO:4, complementary to the sense (+) strand of the *mecA* gene) was labeled with AE attached by a linker positioned between bases 15 and 16.

5 Probe 5 (SEQ ID NO:5, complementary to the antisense (-) strand of the *mecA* gene) was labeled with AE attached by a linker positioned between bases 20 and 21. Probe 6 (SEQ ID NO:6, complementary to the sense (+) strand of the *mecA* gene) was labeled with AE through a linker attached between bases 23 and 24.

Procedures

10 Specimens (1 μ l loops of bacteria) were treated and tested essentially as described in Example 5 with the following differences: incubation for lysis was for 15 minutes and incubation for hybridization was for 45 minutes for samples shown in Tables 7-8, while incubation for lysis was for 10 minutes and incubation for hybridization was for 20 minutes for samples shown in Table 9. All of the base-treated solutions were neutralized by the

15 addition of 250 μ l of 0.8 N HCl containing 100 mM succinate.

The specimen testing results with probe 3-6, alone and in mixtures, are presented in Tables 7-9.

Table 7

Testing of Additional *mecA*-Specific Probes

Probe Composition	Probe alone Avg RLU	GP1217(Pos) RLU	GP18(Neg) RLU
probe 1	3668	25928	757
probe 6	6117	38158	1195
probe 5	4088	37472	1250
25 probe 1 + probe 6	4098	63676	2193
probe 1 + probe 5	2821	72033	1771

Table 8Testing of Additional *mecA*-Specific Probes

Probe Composition	Probe alone Avg RLU	GP1217(Pos) RLU	GP18(Neg) RLU
probe 1	1123	11485	879
probe 4	690	20192	641
probe 3	445	36123	371
probe 1 + probe 4	1493	41537	928
probe 1 + probe 3	1296	63175	844

Table 9Testing of Additional *mecA*-Specific Probes

Probe Composition	Probe alone Avg RLU	GP1217(Pos) RLU
probe 1	2476	27256
probe 4	734	34352
probe 3	684	46566
probe 6	1004	55947
probe 5	2823	42816
probe 4 + probe 6	1443	93569
probe 3 + probe 5	12014	97324
probe 1 + probe 4 + probe 6	3064	115414
probe 1 + probe 3 + probe 5	11759	121633

The results presented in Tables 7-9 showed that probes directed to additional target regions gave good specificity with desirably low backgrounds. Mixtures of the probes showed enhanced signals, although some combinations with high backgrounds may indicate undesirable probe interactions.

Example 72'-Deoxy and 2'-Methoxy Probe Mixtures

2'-methoxy analogs of the AE-labeled DNA probes were prepared by standard phosphoramidite synthesis using 2' methoxy phosphoramidite analogs. Additional linker

WO 2005/017202

PCT/US2004/014690

positions, which are indicated by parentheses in Table 11, for probe 2 were also prepared for testing. Deoxy probes and 2'-methoxy probe analogs were tested essentially as described in Example 5, except as noted below. The results are provided in Tables 10-12. Probes synthesized using 2'-methoxy nucleotide analogs are particularly identified by the "MeO" designations. Brief summaries of the procedures used to obtain the results are appended to the tables.

Table 10

Probe Combinations and the Use of Nucleotide Analogs Can Further Improve Probe Performance

Probe Composition	Probe alone Avg RLU	GP1217(Pos) RLU	GP1214(Neg) RLU
probe 1	2131	37695	1844
MeO-probe 2	2378	69687	5242
MeO-probe 1	1879	62594	2308
MeO-probe 1 + probe 3	2732	94942	2504
MeO-probe 1 + probe 6	3915	89312	9797
probe 3 + probe 6	3745	105024	3606
MeO-probe 1 + probe 3 + probe 6	4247	146128	4684
1 µl loop <i>S. aureus</i> in 150 µl lysis reagent, incubate for 10 minutes at 105°C - base treatment: 100 µl 2 N hydroxide, incubate for 10 minutes at 60°C - neutralize: 100 µl 2 N HCl containing 200 mM succinate - hybridize: 45 minutes at 60°C			

Table 11
Numerous Labeling Positions Yield Useful Probes

Probe Composition	Probe alone Avg RLU	GP1217(Pos) RLU	GP1214(Neg) RLU
(13,14)-MeO-probe 1	779	33659	975
(4,5)-MeO-probe 2	665	52714	1852
(14,15)-MeO-probe 2	829	48954	6327
(17,18)-MeO-probe 2	825	55066	3406
(16,17)-probe 3	1098	36993	1303
(16,17)-MeO-probe 3	1287	60095	997
(15,16)-MeO-probe 4	579	56497	1185
(23,24)-probe 6	1540	43307	1994
(23,24)-MeO-probe 6	2370	70180	5015
(20,21)-MeO-probe 5	815	60009	1809
1 μl loop <i>S. aureus</i> in 150 μl lysis reagent, incubate for 10 minutes at 105°C - base treatment: 100 μl 2 N hydroxide, incubate for 10 minutes at 60°C - neutralize: 250 μl 0.8 N HCl containing 100 mM succinate - hybridize: 45 minutes at 60°C			

Further testing results for AE (4,5)-MeO-probe 2 are shown in Table 12.

Table 12Variously Positioned Labels Yield Useful Probe Compositions

	Probe Composition	Probe alone Avg RLU	GP1217(Pos) RLU	GP1214(Neg) RLU
5	(4,5) MeO-probe 2 + MeO-probe 4	4840	132167	6689
	(4,5)-MeO-probe 2 + probe 6	4349	132124	6059
	MeO-probe 4 + probe 6	2723	138838	5552
	(4,5)-MeO-probe 2 + probe 4 + probe 6	6841	196585	10082
	MeO-probe 3 + probe 5	992	133327	2461
10	(4,5)-MeO-probe 2 + probe 3 + probe 5	2260	208346	4563
	(4,5)-MeO-probe 2 + probe 3 + probe 6	3365	202877	6203
	(4,5)-MeO-probe 2 + probe 4 + probe 5	7537	209019	7742
15	• 1 µl loop <i>S. aureus</i> in 150 µl lysis reagent, incubate for 10 minutes at 105°C • base treatment: 100 µl 2 N hydroxide, incubate for 10 minutes at 60°C • neutralize: 250 µl 0.8 N HCl containing 100 mM succinate • hybridize: 20 minutes at 60°C			

These results showed that methoxy nucleotide analogs gave modestly increased signals with *mecA*-negative samples, but substantially higher signals when hybridized to nucleic acids prepared from methicillin-resistant organisms. Mixtures of the probes resulted in significantly higher signals. Various combinations of the probes gave good results.

Example 8Testing Clinical Specimens

The results of testing clinical isolates with a combination of *mecA* probes are shown in Table 13. Oxacillin MIC and Oxoid PBP2' Latex Agglutination ("OLA") test results were determined at the collection site prior to shipment of the samples for the determination of *mecA* gene status. 1 µl loops of the specimens at the collection site were suspended in 150 µl of lysis reagent and incubated for 10 minutes at 100°C, and then followed by freezing. For HPA determinations, the samples were thawed, vortexed and incubated at 100°C for 5 minutes. Each sample received a 100 µl aliquot of 2 N LiOH, and was subsequently incubated at 60°C for 10 minutes. Samples were neutralized by adding 100 µl of 2 N HCl containing 200 mM succinate buffer. Aliquots of the reaction mixture were

WO 2005/017202

PCT/US2004/014690

then assayed by HPA as described in Example 1. The Gen-Probe ("GP") HPA *Staphylococcus spp.* and *S. aureus* RLU values were determined using the probes and helper oligonucleotides described in Example 5, except that the *S. aureus* results were determined using an AE-labeled *S. aureus* probe (SEQ ID NO:7) having a linker inserted
5 between positions 22 and 23 instead of between positions 20 and 21. The *mecA* status was determined using a mixture of AE-labeled probes (i.e., (4,5)-2'-OMe-probe 2, (16,17)-probe 3 and (20,21)-probe 5).

Table 13Testing Clinical Specimens Using a Probe Panel

Sample	HPA Results (RLU)			Oxacillin MIC (μ g/ml)	Oxoid (OLA)	Identity
	<i>Staph Spp</i>	<i>S. aureus</i>	<i>mecA</i>			
5 Neg Control	1775	3472	1390	nd	nd	Hybridization Buffer
Pos Control GP1217	170530	162795	145716	>16	Pos	<i>S. aureus</i> , <i>mecA</i> (+) (MRSA)
GP21	373232	351348	3429	<0.25	Neg	<i>S. aureus</i> , <i>mecA</i> (-) (MSSA)
10 GP22	225141	7637	210788	0.5	Pos	CoNS, <i>mecA</i> (+)
GP23	301657	280428	3562	<0.25	Neg	<i>S. aureus</i> , <i>mecA</i> (-)
GP24	306718	274112	265780	>16	Pos	<i>S. aureus</i> , <i>mecA</i> (+)
GP25	456426	408938	456055	>16	Pos	<i>S. aureus</i> , <i>mecA</i> (+)
GP26	262267	241597	3309	<0.25	Neg	<i>S. aureus</i> , <i>mecA</i> (-)
15 GP27	182304	7057	3886	<0.25	Neg	CoNS, <i>mecA</i> (-)
GP28	246636	195759	41183	1	Neg	<i>S. aureus</i> , <i>mecA</i> (-)
GP29	388748	416372	440128	>16	Pos	<i>S. aureus</i> , <i>mecA</i> (+)
GP30	440128	399498	5660	<0.25	Neg	<i>S. aureus</i> , <i>mecA</i> (-)
GP31	349207	332750	322837	>16	Pos	<i>S. aureus</i> , <i>mecA</i> (+)
20 GP32	432041	414709	11706	<0.25	Neg	<i>S. aureus</i> , <i>mecA</i> (-)
GP33	160830	154183	142431	>16	Pos	<i>S. aureus</i> , <i>mecA</i> (+)
GP34	634079	497890	5191	<0.25	Neg	<i>S. aureus</i> , <i>mecA</i> (-)
GP35	543089	484580	370770	>16	Pos	<i>S. aureus</i> , <i>mecA</i> (+)
GP36	370898	356243	5587	<0.25	Neg	<i>S. aureus</i> , <i>mecA</i> (-)
25 GP37	397722	373448	406010	>16	Pos	<i>S. aureus</i> , <i>mecA</i> (+)
GP38	387613	367728	402740	nd	Pos	<i>S. aureus</i> , <i>mecA</i> (+)
GP39	280358	268665	4838	<0.25	Neg	<i>S. aureus</i> , <i>mecA</i> (-)
GP40	292407	283200	3617	<0.25	Neg	<i>S. aureus</i> , <i>mecA</i> (-)

30

The results presented in Table 13 demonstrated that the invented hybridization assay correctly identified the organism type and methicillin-resistance status for numerous clinical

WO 2005/017202

PCT/US2004/014690

isolates. The unusual result obtained for the GP28 sample, meaning an intermediate *mecA* hybridization signal coupled with an intermediate MIC reading, could not be repeated and was attributed to unreacted probe that was not efficiently hydrolyzed by the selection reagent. The high background HPA result would easily be explained by incomplete mixing
5 of the probe and selection reagent during the HPA procedure.

Example 9

Additional Specimen Testing

Additional testing of isolates is summarized in Table 14. Oxacillin MIC results were provided by an outside laboratory. The GP Probe results were determined as
10 described in Example 8.

Table 14Testing Clinical Specimens Using a Probe Panel

Organism	# Isolates	Oxacillin MIC (μ g/ml)	HPA Results (# positive for indicated probes)				OLA	
			<i>Staph.</i> <i>Spp</i>	<i>S.</i> <i>aureus</i>	<i>mecA</i> (+)	<i>mecA</i> (-)	Pos	Neg
Clinical MRSA	14	4	14	14	6	8	6	8
	13	8	13	13	9	4	9	4
	8	16	8	8	8	0	7	1
	107	>16	107	107	107	0	106	1
Clinical MSSA	99	≤ 0.25	99	99	1	98	1	98
	28	0.5	28	28	0	28	0	28
	13	1	13	13	0	13	0	13
	8	2	8	8	2	6	2	6
ATCC MRSA	1	8	1	1	1	0	1	0
	1	16	1	1	1	0	1	0
ATCC MSSA	6	≤ 0.25	6	6	0	6	0	6
	4	0.5	4	4	0	4	0	4
	1	1	1	1	0	1	0	1
<i>S. epidermidis</i>	2	≤ 0.25	2	0	0	2	0	2
	1	4	1	0	1	0	1	0
	1	>16	1	0	1	0	1	0
<i>S. intermedius</i>	1	≤ 0.25	1	0	0	1	0	1
<i>S. hyicus-hyicus</i>	1	≤ 0.25	1	0	0	1	0	1
<i>S. warneri</i>	1	≤ 0.25	1	0	0	1	1	0
<i>S. hemoliticus</i>	1	≤ 0.25	1	0	0	1	0	1
<i>S. saprophyticus</i>	1	0.5	1	0	0	1	0	1
<i>S. simulans</i>	1	≤ 0.25	1	0	0	1	0	1
<i>S. hominis</i>	1	≤ 0.25	1	0	0	1	0	1

The results in Table 14 showed that the Oxoid (OLA) test missed detecting some of the clinical *mecA*(+) *S. aureus* samples. The fact that false-negative results were obtained in the OLA assay was supported by the results of the MIC assay. Additionally, the OLA assay gave false-positive results with the *S. warneri* isolate. In aggregate, the results presented in the table represented evidence for the superior accuracy and excellent specificity of the invented probe-based assay.

Example 10Probes for Detecting Vancomycin Resistance Genes

The sequences of oligonucleotide probes that were used for detecting the VanA and VanB genes are presented below. Of course, because these probes are intended for
 5 detecting double-stranded DNA targets, probes having the complements of these sequences also can be used for detecting the VanA and VanB genes, and fall within the scope of the invention. Additionally, detection of mRNA targets falls within the scope of the invention.

VanA-specific probes

VanA(+)-464 GGGTTGCTCAGAGGAGCATGACGTATCGG (SEQ ID NO:27)
 10 VanA(-)-792 GCTGAGCTTTGAATATCGCAGCC (SEQ ID NO:28)
 VanA(-)-1325 CGTTCAGTACAATGCGGCCG (SEQ ID NO:29)

VanB-specific probes

VanB(+)-084 CCGCAGCTTGCATGGACAAATCACTGGC (SEQ ID NO:30)
 VanB(+)-428 CGCATCCATCAGGAAAACGAGCCGG (SEQ ID NO:31)
 15 VanB(-)-1006 CCAAGCACCCGATATACTTTCTTTGCC (SEQ ID NO:32)

VanA and VanB cross-reactive probes

Van A&B(-)-734 GAGCTTTGAATATCGCAGCCTAC (SEQ ID NO:33)

As will be apparent from inspection of these sequences, the VanA(-)-792 probe and
 20 Van A&B(-)-734 probe share a core sequence given by: GAGCTTTGAATATCGCAGCC (SEQ ID NO:34), which is contained within a composite sequence given by GCTGAGCTTTGAATATCGCAGCCTAC (SEQ ID NO:35). The composite represents a contiguous sequence of 26 bases having ends defined by the 5' boundary of the VanA(-)-792 probe and by the 3' boundary of the Van A&B(-)-734 probe. Each of the two probes has 20
 25 contiguous bases contained within the given composite sequence. Despite the substantial similarities between the VanA(-)-792 and Van A&B(-)-734 probe sequences, the results presented below proved that the Van A&B(-)-734 probe had a superior capacity for either or both of the VanA and VanB gene sequences in lysates derived from clinical samples. This unexpected property could not have been predicted in advance of this showing, and so
 30 illustrates one advantage of the Van A&B(-)-734 probe. Conversely, the VanA(-)-792 probe advantageously was strongly reactive with the VanA gene sequence, and substantially less

WO 2005/017202

PCT/US2004/014690

reactive with the VanB gene sequence, thereby providing a means for selectively detecting the VanA gene sequence in a highly efficient manner.

Each of the above-listed probes was synthesized using DNA precursors, and labeled with acridinium ester by a non-nucleotide linker joined according to the procedures described herein. The linker on the VanA(+)-464 probe was located between nucleotide positions 19 and 20. The linker on the VanA(-)-792 probe was located between nucleotide positions 8 and 9. The linker on the VanA(-)-1325 probe was located between nucleotide positions 13 and 14. The linker on the VanB(+)-084 probe was located between nucleotide positions 8 and 9. The linker on the VanB(+)-428 probe was located between nucleotide positions 8 and 9. The linker on the VanB(-)-1006 probe was located between nucleotide positions 18 and 19. The linker on the Van A&B(-)-734 probe was located between nucleotide positions 11 and 12. It should be understood that use of the chemiluminescent AE labeling system merely represents one embodiment of the invention that has been used for illustrative purposes. Other detectable species that will be familiar to those having an ordinary level of skill in the art can be used for labeling the probes. Of course, homogeneously detectable labels represent the most highly preferred detectable labels.

Example 11

Establishing the Genotypes of Cultured VRE Bacteria

Viable samples of VRE bacteria were obtained from the ATCC, propagated on appropriate growth media and their isolated nucleic acids tested by a PCR protocol to determine the identity of the gene responsible for the drug-resistant phenotype. Primers for amplifying the VanA gene had the sequences of VanA(+)-464Kpn (CGGGGTACCGGGTTGCTCAGAGGAGCATGACGTATCGG (SEQ ID NO:36)) and VanA(-)-1325Bam (CGCGGATCCGTTTCAGTACAATGCGGCCG (SEQ ID NO:37)). These primers had the target-complementary sequences of SEQ ID NO:27 and SEQ ID NO:29, respectively, positioned downstream of sequences that were not complementary to the target to be amplified. Primers for amplifying the VanB gene had the sequences of VanB(+)-084Kpn (CGGGGTACCGCAGCTTGTCATGGACAAATCACTGGC (SEQ ID NO:38)) and VanB(-)-1006Bam (CGCGGATCCAAGCACCCGATATACTTTCTTTGCC (SEQ ID NO:39)). These primers had the target-complementary sequences of SEQ ID NO:30 and SEQ ID NO:32, respectively, positioned downstream of sequences that were not complementary to the target to be amplified. In every instance, separate reactions were

WO 2005/017202

PCT/US2004/014690

conducted to amplify the VanA and VanB genes using a sample of genomic DNA from each isolate. Positive results indicating the presence of a particular vancomycin-resistance gene were judged by the appearance of a discrete band representing a PCR product on an agarose gel stained with ethidium bromide. Each VRE sample contained either the VanA or VanB gene, and none of the samples contained both genes. The results of these procedures are presented in Table 15.

Table 15Cultured VRE Bacteria Harbor VanA or VanBGene Sequences

ATCC #	Species	Genotype
51575	<i>E. faecalis</i>	VanB
700221	<i>E. faecium</i>	VanA
700802	<i>E. faecalis</i>	VanB
51299	<i>E. faecalis</i>	VanB

Example 12

Hybridization Assays for Detecting Genes Encoding
Vancomycin Resistance

Double-stranded plasmids harboring inserts of either VanA or VanB gene sequences were prepared for use as control templates. More specifically, VanA gene sequences were amplified in a PCR protocol using primers having the sequences of VanA(+)464Kpn and VanA(-)1325Bam. Similarly, VanB gene sequences were amplified using primers having the sequences of VanB(+)084Kpn and VanB(-)1006Bam. Genomic DNA samples from bacteria known to have VanA or VanB genotypes were used as templates in the amplification reactions. Blunt-ended amplification products containing VanA gene sequences were ligated into a pPCR-SCRIPTAMP SK(+) (Stratagene; La Jolla, CA) plasmid cloning vector, and propagated in *E. coli* host bacteria using procedures that will be familiar to those having an ordinary level of skill in the art. Double-stranded amplification products containing VanB gene sequences were first digested with KpnI and BamHI, and then ligated into a similarly cleaved pBLUESCRIPT-II KS(+) (Stratagene; La Jolla, CA) plasmid cloning vector. Isolated double-stranded plasmid DNA served as controls in hybridization assays.

WO 2005/017202

PCT/US2004/014690

2'-deoxy probes having the sequences presented under Example 10 were prepared by standard phosphoramidite synthesis. AE labeling of the probes also was performed using standard procedures, as described herein and as will be familiar to those having an ordinary level of skill in the art. Samples of four different cultured VRE bacteria described under the previous Example were lysed essentially as described above. RNA in the samples was hydrolyzed and DNA denatured by incubation of the lysate with hydroxide at elevated temperature. In parallel procedures, samples of two plasmid clones harboring either the VanA or VanB inserts (as described under the previous Example) were also treated to denature DNA. All samples were neutralized by addition of a buffered HCl reagent. AE-labeled probes were added to each of the samples and then hybridized at 55-60°C for 45 minutes to permit formation of complementary probe:target duplexes. The pan-bacterial probe (Probe Reagent 1, GEN-PROBE® MTC-NI Kit, catalog no. 4573) described under Example 2 was used as a positive control to confirm the presence of bacterial nucleic acid targets. Trials containing labeled probe in the absence of an added target (i.e., "probe alone") served as negative controls. The AE label joined to probe molecules that remained single-stranded following the hybridization procedure were destroyed by treatment under mild alkaline conditions, and the extent of specific hybrid formation quantified by luminometry as described in the foregoing Examples. Notably, the VanB plasmid clone did not fully contain the sequence complementary to the Van A&B(-)734 probe. Numerical results from these procedures (measured in RLU) are presented in Table 16.

Table 16
Detection of VanA and VanB Gene Sequences by
Probe Hybridization (in RLU)

Probe	ATCC 51575 (VanB)	ATCC 700221 (VanA)	ATCC 700802 (VanB)	ATCC 51299 (VanB)	VanA clone	VanB clone	Probe Alone
5 VanA(+)-464	696	10910	757	765	37434	817	374
VanA(-)-792	2254	10627	3653	2983	52729	505	256
VanA(-)-1325	1761	23523	931	1182	24378	942	368
10 VanA&B(-)-734	5166	8901	7425	5497	48999	718†	240
VanB(+)-084	6003	768	10119	7863	716	54711	371
VanB(+)-428	6307	683	9657	8113	684	49242	370
VanB(-)-1006	6300	1485	9116	8335	1708	27493	586
15 MTC-NI (+)-control	23497	33027	68973	29585	975	530	297

† The VanB plasmid clone does not fully include the complement of the Van A&B(-)-734 probe.

20 The results presented in Table 16 proved that each of the various probes specific for VanA, VanB, or the combination of VanA and/or VanB was able to detect and identify the appropriate vancomycin-resistance gene present in each of the four cultured bacterial samples. More particularly, the VanA(+)-464, VanA(-)-792 and VanA(-)-1325 probes gave very strong hybridization signals for samples containing nucleic acids from the single
25 bacterial isolate that was VanA-positive. Each of these probes produced substantially weaker signals when hybridized with samples that did not harbor the VanA target. Similarly, the VanB(+)-084, VanB(+)-428 and VanB(-)-1006 probes gave very strong signals when hybridized with samples containing nucleic acids from the VRE bacteria that were VanB-positive. Each of these probes produced substantially weaker signals when
30 hybridized with samples that did not harbor the VanB nucleic acid target. Finally, the Van

WO 2005/017202

PCT/US2004/014690

A&B(-)734 probe gave very strong signals when hybridized with nucleic acids from VRE bacteria that were either VanA-positive or VanB-positive.

Interestingly, the Van A&B(-)734 and VanA(-)792 probes exhibited strikingly different hybridization characteristics despite having target-complementary sequences that are 23 bases in length and sharing a common core sequences of 20 bases. Despite these similarities, the two probes differed from each other in their relative selectivity for VanA and VanB targets, as illustrated by the quantitative results obtained using lysates prepared from the different bacterial isolates. For example, comparison of background-corrected hybridization signals relative to negative controls obtained using the VanA-positive bacterial sample (ATCC # 700221) and either of the two probes were substantially comparable (being about 40 fold and about 36 fold greater than background signals for the VanA(-)792 and Van A&B(-)734 probes, respectively). A similar comparison of signals resulting from hybridization of the two probes with lysates of VanB-positive bacterial samples (ATCC Nos. 51575, 700802 and 51299) yielded very different results. In these instances the hybridization signals obtained using the VanA(-)792 probe averaged about 10.5 fold greater than background, while the hybridization signals obtained using the Van A&B(-)734 probe averaged about 24 greater than background. Under a conventional scenario wherein positive results are judged as being at least 10 fold greater than the negative control, the two probes uniformly detected the VanA target, but equivocally detected the VanB target. These findings emphasized the subtle effect of minor sequence variations on probe functionality.

The following Example describes a test device that fulfilled several preferred design goals. The device included a solid support harboring individual wells for containing defined collections of soluble hybridization probes in a spaced-apart configuration. This arrangement meant that all of the probes could be transported simultaneously, for example to a hybridization incubator or luminometer, and that all of the hybridization reactions could conveniently be conducted under the same temperature conditions. The device facilitated rapid identification of *S. aureus*, coagulase-negative Staphylococcal bacteria (CoNS), as well as Enterococcus bacteria. Additionally, the device was capable of detecting nucleic acid markers indicative of resistance to methicillin or vancomycin in any organism undergoing testing. Finally, the device was useful for conducting homogeneous assays. Accordingly, it was unnecessary to separate unhybridized probe from specifically

WO 2005/017202

PCT/US2004/014690

hybridized probe in order to detect specific hybrids which indicated the presence of a complementary target sequence.

The fact that the device is preferably used for testing DNA, and not RNA, to determine the presence or absence of the relevant targets imposed certain limitations, and defined a problem to be overcome. More particularly, when the amount of a DNA sample to be tested is limiting, the number of hybridization reactions needed to make the identification must be minimized. A preferred solution to the problem involved combining certain probes in a single hybridization reaction such that a positive hybridization result yielded an ambiguous determination. As described below, rather than conducting separate hybridization reactions using probes specific for the VanA and VanB genes, a single hybridization reaction was used to identify nucleic acids containing either of these targets without determining which target was present. Additionally, two probes that were independently specific for the nucleic acids of *S. aureus* and for Enterococcus bacteria harbored identical labels and were combined in a single hybridization reaction. A positive hybridization signal in this reaction indicated the presence of *S. aureus* or Enterococcus bacteria, without particularly identifying which organism was present.

When sufficient material is available for testing, the added use of a pan-bacterial probe as a positive control provides additional functionality to the invented device, and eliminates the requirement for foreknowledge about the identity of the organism undergoing testing. For example, because it is easily possible for an organism to be misidentified as a member of the Enterococcus group preliminary to testing with the invented device, negative results using an Enterococcus-specific hybridization probe (in the absence of a pan-bacterial address), together with negative results at other addresses in the device, would be uninformative because it would be possible that either insufficient material was included in the procedure, or that the organism had been misidentified prior to the probe hybridization procedure. This ambiguity can be overcome by the use of a pan-bacterial probe as a positive control. Accordingly, a positive hybridization signal observed in a hybridization reaction using a pan-bacterial probe of the type described herein will indicate that sufficient nucleic acid is present in the sample undergoing testing to provide meaningful results in the other hybridization reactions conducted using the same device.

Example 13Device for Conducting Simultaneous Bacterial Identification and
Antibiotic Resistance Testing

Aliquots of AE-labeled hybridization probes having the following specificities were
 5 dispensed into four wells of a 96-well plastic microtiter plate as follows. The first well of
 the device contained an aliquot of AE-labeled probe having the sequence of SEQ ID
 NO:10, together with helper oligonucleotides having the sequences of SEQ ID NOs:11 and
 12, and so was capable of hybridizing to the nucleic acids of a plurality of bacteria in the
 Staphylococcus genus, including *S. aureus*, *S. epidermidis* and *S. haemolyticus*, but not to
 10 the nucleic acids of bacteria in the genus Enterococcus. The oligonucleotide having the
 sequence of SEQ ID NO:11 was synthesized using 2'-methoxy nucleotide analogs. The
 second well contained a mixture of AE-labeled probes, one of the probes having the *S.*
aureus-specific sequence of SEQ ID NO:7, together with helper oligonucleotides having
 the sequences of SEQ ID NOs:8 and 9. The second well further included a first AE-labeled
 15 Enterococcus-specific probe having the sequence of SEQ ID NO:14, together with helper
 oligonucleotides having the sequences of SEQ ID NOs:23 and 24, and a second AE-labeled
 Enterococcus-specific probe having the sequence of SEQ ID NO:15, together with helper
 oligonucleotides having the sequences of SEQ ID NOs:25 and 26. The probes in the
 second well were capable of hybridizing to the nucleic acids of *S. aureus* and Enterococcus
 20 bacteria. Notably, the probes in the second well did not exhibit cross-reactivity, meaning
 that the probe specific for the nucleic acids of *S. aureus* did not hybridize to the nucleic
 acids of Enterococcus bacteria, and the probe specific for the nucleic acids of Enterococcus
 bacteria did not hybridize to the nucleic acids of *S. aureus* or any other member of the
 Staphylococcus genus. Also notably, the probe specific for the nucleic acids of *S. aureus*
 25 did not hybridize to the nucleic acids of other members of the Staphylococcus genus. Thus,
 the *S. aureus*-specific probe did not hybridize to, for example, *S. epidermidis* or *S.*
haemolyticus. The third well of the device contained a mixture of AE-labeled probes
 having the sequences of SEQ ID NOs:2, 3 and 5, and so was capable of hybridizing to
 nucleic acids of the *mecA* gene, but not to the nucleic acids of either the VanA or VanB
 30 genes. These three probes were all synthesized using 2'-methoxy nucleotide analogs. The
 fourth well of the device contained a mixture of AE-labeled probes having the sequences of
 SEQ ID NOs:27, 28, 29, 30. The mixture further included AE-labeled probes having the

WO 2005/017202

PCT/US2004/014690

complements of SEQ ID NO:31 (i.e., CCGGCTCGTTTTCTGATGGATGCG (SEQ ID NO:40)) and SEQ ID NO:32 (i.e., GGCAAAGAAAGTATATCGGGTGCTTGG (SEQ ID NO:41)). The probes in the fourth well were capable of hybridizing to the nucleic acids of the VanA and VanB genes, but not to the nucleic acids of the *mecA* gene. As indicated above, an optional fifth well of the device can contain a labeled probe that specifically hybridizes to the nucleic acids of a plurality of bacterial organisms, but not to the nucleic acids of fungal organisms. For example, a labeled probe having the sequence of SEQ ID NO:16 may be used for this purpose.

Numerous biological samples were tested using the sample preparation, probe hybridization and specific-hybrid detection methods described herein. The results uniformly conformed to the patterns of positive and negative hybridization results presented in Table 17. Indeed, when the device was used for hybridizing nucleic acids of a test organism, it was possible to determine the identity of the donor organism from the unique combination of positive and negative hybridization results using the information shown in the table. Notably, filled boxes in Table 17 indicate positive hybridization signals, and open boxes indicate the absence of a positive hybridization signal. Entries in the table for MRE (methicillin-resistant *Enterococcus*) and VRSA (vancomycin-resistant *S. aureus*) show the unique patterns of hybridization results that would be expected for these isolates, although these organisms were not available for testing. Since many VRSA isolates are also resistant to methicillin, bacteria having the dual antibiotic resistant phenotype would be expected to give positive hybridization signals at the *mecA* address, as well as at the VanA and/or VanB addresses. Methicillin-resistant CoNS isolates would be expected to give positive hybridization signals at the *Staphylococcus* genus address and at the *mecA* address, but not at the *S. aureus*/*Enterococcus* address or at the VanA/VanB address. Finally, all of the probes used in the device were hybridized at the same temperature to achieve the results presented in the following table.

Table 17

Decoding Hybridization Results for Bacterial Identification
and Antibiotic Resistance Testing

5 Probe Address	Hybridization Result and Corresponding Interpretation						
	<i>S. aureus</i>	CoNS	Entero- coccus	MRE	VRE	MRSA	VRSA
Staphylococcus genus							
<i>S. aureus</i> and Enterococcus genus							
10 <i>mecA</i>							
VanA and VanB							

An alternative approach for obtaining diagnostic information substantially equivalent to that presented in the foregoing table involves the use of six independent probe addresses. Table 18 shows the results that would be expected if hybridization reactions were conducted using the same hybridization probes and procedures as described above, with the exception that each hybridization reaction yielded an unambiguous result. To illustrate this aspect of the invention, the VanA and VanB results are shown in Table 18 as being coincident for the VRE and VRSA addresses for the vancomycin-resistant phenotypes. However, one (but not both) of the VanA and VanB addresses could give a negative hybridization result and still indicate a vancomycin-resistant phenotype. Stated differently, if either the VanA or VanB probe address gave a positive hybridization signal, that would be evidence for resistance to vancomycin. As indicated above, because many VRSA isolates are also resistant to methicillin, bacteria having the dual antibiotic resistant phenotype would be expected to give positive hybridization signals at the *mecA* address, as well as at the VanA and/or VanB addresses. Methicillin-resistant CoNS isolates would give positive hybridization signals at the Staphylococcus genus address and at the *mecA* address, but not at addresses for detecting the nucleic acids specific for *S. aureus*, Enterococcus, VanA or VanB.

Table 18

Decoding Hybridization Results for Bacterial Identification
and Antibiotic Resistance Testing

5	Probe Address	Hybridization Result and Corresponding Interpretation						
		<i>S. aureus</i>	CoNS	Entero-coccus	MRE	VRE	MRSA	VRSA
	Staphylococcus genus							
	<i>S. aureus</i>							
	Enterococcus genus							
10	<i>mecA</i>							
	VanA							
	VanB							

Finally, using the above-described hybridization assay techniques, it is possible to
 15 quantify the contributions of mixtures of different organisms based on the magnitudes of
 the measured hybridization signals.

This invention has been described with reference to a number of specific examples
 and embodiments thereof. Of course, a number of different embodiments of the present
 invention will suggest themselves to those having ordinary skill in the art upon review of
 20 the foregoing detailed description. Thus, the true scope of the present invention is to be
 determined upon reference to the appended claims.

DEMANDES OU BREVETS VOLUMINEUX

**LA PRÉSENTE PARTIE DE CETTE DEMANDE OU CE BREVETS
COMPREND PLUS D'UN TOME.**

CECI EST LE TOME _____ DE _____

NOTE: Pour les tomes additionels, veuillez contacter le Bureau Canadien des Brevets.

JUMBO APPLICATIONS / PATENTS

**THIS SECTION OF THE APPLICATION / PATENT CONTAINS MORE
THAN ONE VOLUME.**

THIS IS VOLUME 1 OF 2

NOTE: For additional volumes please contact the Canadian Patent Office.

What is claimed is:

1. A method for detecting the presence or absence of at least a first target nucleic acid and a second target nucleic acid in a sample, wherein the first target nucleic acid is a nucleic acid encoding resistance to an antibiotic and the second target nucleic acid is specific for one or more bacterial species, said method comprising:

providing a sample suspected of containing the first and second target nucleic acids;

contacting the sample with

(i) a first oligonucleotide probe that hybridizes to the first target nucleic acid to form a first hybrid between the first oligonucleotide probe and the first target nucleic acid, and

(ii) a second oligonucleotide probe that hybridizes to the second target nucleic acid to form a second hybrid between the second oligonucleotide probe and the second target nucleic acid;

wherein the first and second oligonucleotide probes are respectively distributed at first and second loci on a solid support; and

detecting the presence or absence of the first and second hybrids as indicators of the presence or absence of the first and second target nucleic acids.

2. The method of claim 1, wherein the first and second oligonucleotide probes are soluble oligonucleotide probes.

3. The method of claim 1 or 2, wherein the first target nucleic acid is a *mecA* gene, a VanA gene or a VanB gene.

4. The method of claim 1, 2 or 3, wherein the second target nucleic acid is from a *Staphylococcus* bacteria.

5. The method of claim 4, wherein the *Staphylococcus* bacteria is *Staphylococcus aureus*.

6. The method of any one of claims 1 to 5, wherein the contacting is under high stringency hybridization conditions.

7. A method for detecting the presence or absence of a target nucleic acid encoding resistance to an antibiotic and a target nucleic acid from a bacterial species, said method comprising:

providing a sample suspected of containing at least one bacterial species;

contacting the sample with oligonucleotide probes distributed at a plurality of loci on a solid support, wherein the oligonucleotide probes comprise

(i) at a first locus, one or more probes that collectively hybridize to ribosomal nucleic acids from a plurality of species of bacteria in the genus *Staphylococcus*, comprising *Staphylococcus aureus* and *Staphylococcus epidermis*, but not to ribosomal nucleic acids of bacteria in the genus *Enterococcus*, to form at least a first hybrid;

(ii) at a second locus, (a) one or more oligonucleotide probes that collectively hybridize to ribosomal nucleic acids from a plurality of bacteria in the genus *Enterococcus*, comprising *Enterococcus faecalis* and *Enterococcus faecium*, but not *Staphylococcus aureus* or any other bacteria in the genus *Staphylococcus*, and (b) at least one probe that hybridizes to a ribosomal nucleic acid from *Staphylococcus aureus* but not from other species in the genus *Staphylococcus* or bacteria in the genus *Enterococcus*, to form at least a second hybrid;

(iii) at a third locus, a least one oligonucleotide probe that hybridizes to a *mecA* nucleic acid; and

(iv) at a fourth locus, one or more oligonucleotide probes that collectively hybridize to VanA nucleic acids and to VanB nucleic acids;

detecting the presence or absence of the at least first, second, third, and fourth hybrids as indicators of the presence or absence of the an antibiotic resistance gene and a bacterial species in the sample.

8. The method of claim 7, wherein the oligonucleotide probes at each of said loci are soluble oligonucleotide probes.

9. The method of claim 7 or 8, wherein the contacting is under high stringency hybridization conditions.

10. The method of claim 7, 8 or 9, wherein the oligonucleotide probes further comprise (v) at a fifth locus, one or more probes that collectively hybridize to ribosomal nucleic acids from a plurality of species of Gram-positive bacteria comprising the high(G+C) subset thereof, a plurality of species of bacteria in the family Enterobacteriaceae, a plurality of species of bacteria in the genus Enterococcus, and a plurality of species of bacteria in the genus Staphylococcus.

11. The method of any one of claims 7 to 10, wherein the one or more probes that collectively hybridize to VanA nucleic acids and to VanB nucleic acids comprise at least one probe that hybridizes to VanA nucleic acids and at least one probe that hybridizes to VanB nucleic acids.

12. The method of any one of claims 7 to 10, wherein the one or more probes that collectively hybridize to VanA nucleic acids and to VanB nucleic acids comprise a single probe that hybridizes both to VanA nucleic acids and VanB nucleic acids.

13. A method for detecting the presence or absence of a target nucleic acid encoding resistance to an antibiotic and a target nucleic acid from a bacterial species, said method comprising:

providing a sample suspected of containing at least one bacterial species;

contacting the sample with oligonucleotide probes distributed at a plurality of loci on a solid support, wherein the oligonucleotide probes comprise

(i) at a first locus, one or more oligonucleotide probes that collectively hybridize to ribosomal nucleic acids from a plurality of species of bacteria

in the genus *Staphylococcus*, comprising *Staphylococcus aureus* and *Staphylococcus epidermidis*, to form at least a first hybrid;

- (ii) at a second locus, at least one oligonucleotide probe that hybridizes to a ribosomal nucleic acid from *Staphylococcus aureus* but not other species in the genus *Staphylococcus*, to form at least a second hybrid;
- (iii) at a third locus, a least one oligonucleotide probe that hybridizes to a *mecA* nucleic acid, to form at least a third hybrid; and
- (iv) at a fourth locus, one or more oligonucleotide probes that collectively hybridize to ribosomal nucleic acids from a plurality of species of Gram positive bacteria comprising the high(G+C) subset thereof, a plurality of species of bacteria in the family Enterobacteriaceae, a plurality of species of bacteria in the genus Enterococcus, and a plurality of species of bacteria in the genus *Staphylococcus*, to form at least a fourth hybrid; and

detecting the presence or absence of the at least first, second, third, and fourth hybrids as indicators of the presence or absence of the an antibiotic resistance gene and a bacterial species in the sample.

14. The method of claim 13, wherein the oligonucleotide probes at each of said loci are soluble oligonucleotide probes.

15. The method of claim 13 or 14, wherein the one or more probes at the fourth locus comprise a pan-bacterial probe selected from the group consisting of the base sequence of SEQ ID NO:16 and the base sequence of SEQ ID NO:19.

16. The method of claim 13, 14 or 15, wherein the at least one probe at the third locus has a base sequence selected from the group consisting of SEQ ID NO:2, SEQ ID NO:3, and SEQ ID NO:5.

17. The method of any one of claims 13 to 16, wherein the contacting is under high stringency hybridization conditions.

18. A method for detecting the presence or absence of a target nucleic acid encoding resistance to an antibiotic and a target nucleic acid from a bacterial species, said method comprising:

providing a sample suspected of containing at least one bacterial species;
contacting the sample with oligonucleotide probes distributed at a plurality of loci on a solid support, wherein the oligonucleotide probes comprise

- (i) at a first locus, one or more oligonucleotide probes that collectively hybridize to ribosomal nucleic acids from a plurality of bacteria in the genus *Enterococcus*, comprising *Enterococcus faecalis* and *Enterococcus faecium*, to form at least a first hybrid;
- (ii) at a second locus, one or more oligonucleotide probes that collectively hybridize to VanA nucleic acids and to VanB nucleic acids, to form at least a second hybrid; and
- (iii) at a third locus, one or more oligonucleotide probes that collectively hybridize to ribosomal nucleic acids from a plurality of species of Gram-positive bacteria comprising the high(G+C) subset thereof, a plurality of species of bacteria in the family Enterobacteriaceae, a plurality of species of bacteria in the genus *Enterococcus*, and a plurality of species of bacteria in the genus *Staphylococcus*, to form at least a third hybrid; and

detecting the presence or absence of the at least first, second, and third hybrids as indicators of the presence or absence of the an antibiotic resistance gene and a bacterial species in the sample.

19. The method of claim 18, wherein the oligonucleotide probes at each of said loci are soluble oligonucleotide probes.

20. The method of claim 18 or 19, wherein the one or more probes at the third locus comprise a pan-bacterial probe selected from the group consisting of SEQ ID NO:16 and SEQ ID NO:19.

21. The method of claim 18, 19 or 20, wherein the one or more probes at the second locus have base sequences selected from the group consisting of SEQ ID NO:27, SEQ ID NO:28, and SEQ ID NO:29.

22. The method of claim 18, 19 or 20, wherein the one or more probes at the second locus have base sequences selected from the group consisting of SEQ ID NO:30, SEQ ID NO:40, and SEQ ID NO:41.

23. The method of any one of claims 18 to 22, wherein each of said oligonucleotide probes is labeled with a homogeneously detectable label.

24. The method of claim 23, wherein the homogeneously detectable label is a chemiluminescent label.

25. The method of any one of claims 18 to 24, wherein the contacting is under high stringency hybridization conditions.

26. The method of any one of claims 1 to 25, wherein the sample comprises cells and wherein the method further comprises before said contacting:
lysing the cells to yield a lysate; treating the lysate with a basic composition to yield a basic mixture; and introducing one or more reagents to the basic mixture to yield a reaction mixture at a pH in the range of 4.5 - 5.5.

27. A method of detecting the presence of a target DNA sequence in a sample containing cells, comprising the steps of: lysing cells contained in the sample to yield a lysate; treating the lysate with a basic composition to yield a basic mixture; introducing one or more reagents to the basic mixture to yield a reaction mixture at a pH in the range of 4.5 - 5.5; hybridizing the reaction mixture with an oligonucleotide probe to form a hybrid between the oligonucleotide probe and the target DNA; and detecting said hybrid as an indicator of the presence of said target DNA sequence, wherein said oligonucleotide probe is complementary either to a *mecA* gene sequence, a *VanA* gene sequence or a *VanB* gene sequence.

28. The method of claim 27, wherein the hybridizing is under high stringency hybridization conditions.

29. A method for detecting the presence or absence of a target DNA sequence in a sample containing cells, said method comprising:

(a) treating the sample to yield a lysate, wherein the treating includes: (i) treating the lysate with a basic composition to yield a basic mixture; and (ii) introducing one or more reagents to the basic mixture to yield a reaction mixture at a pH in the range of 4.5 to 5.5;

(b) hybridizing the reaction mixture with a plurality of oligonucleotide probes distributed at a plurality of loci on a solid support, wherein the solid support is configured to comprise:

(i) a first locus containing one or more probes that collectively hybridize to ribosomal nucleic acids from a plurality of species of bacteria in the genus *Staphylococcus*, comprising *Staphylococcus aureus* and *Staphylococcus epidermis*, but not to ribosomal nucleic acids of bacteria in the genus *Enterococcus*, to form a hybrid with a target nucleic acid in the reaction mixture;

(ii) a second locus containing, (a) one or more oligonucleotide probes that collectively hybridize to ribosomal nucleic acids from a plurality of bacteria in the genus *Enterococcus*, comprising *Enterococcus faecalis* and *Enterococcus faecium*, but not *Staphylococcus aureus* or any other bacteria in the genus *Staphylococcus*, to form a hybrid with a target nucleic acid in the reaction mixture, and (b) at least one probe that hybridizes to a ribosomal nucleic acid from *Staphylococcus aureus* but not from other species in the genus *Staphylococcus* or bacteria in the genus *Enterococcus*, to form a hybrid with a target nucleic acid in the reaction mixture;

(iii) a third locus containing at least one oligonucleotide probe that hybridizes to a *mecA* nucleic acid, to form a hybrid with a target nucleic acid in the reaction mixture; and

(iv) a fourth locus containing one or more oligonucleotide probes that collectively hybridize to *VanA* nucleic acids and to *VanB* nucleic acids, to form a hybrid with a target nucleic acid in the reaction mixture; and

(c) detecting the presence or absence of a hybrid at each loci as an indicator of the presence or absence of the target DNA, and thereby determining the presence or absence of a Staphylococcus organism, the presence or absence of an Enterococcus organism and the presence or absence of a drug resistant genotype.

30. The method of claim 29, wherein the at least one probe at the third locus substantially corresponds to SEQ ID NOS:2, 3 or 5; or wherein the at least one probe at the forth locus substantially corresponds to SEQ ID NOS:27, 28, 29, 30, 40 or 41; or both.

31. The method of claim 29 or 30, wherein the hybridizing is under high stringency hybridization conditions.

32. The method of any one of claims 26 to 31, wherein said lysing comprises treating the cells with a lysing agent and then heat.

33. The method of any one of claims 26 to 31, wherein said lysing comprises treating the cells with a lysing agent and heat simultaneously.

34. The method of any one of claims 26 to 33, further comprising heating the basic mixture at a temperature and time sufficient to enhance degradation of RNA that may be present in the lysate, relative to treatment in the absence of heat.

35. The method of claim 34, wherein heating the basic mixture is to a temperature of up to 100°C.