

(19) **DANMARK**

(10) **DK/EP 2943567 T3**



(12) **Oversættelse af
europæisk patentskrift**

Patent- og
Varemærkestyrelsen

-
- (51) Int.Cl.: **C 12 N 7/00 (2006.01)** **A 61 K 39/245 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2018-01-02**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2017-09-13**
- (86) Europæisk ansøgning nr.: **14703459.9**
- (86) Europæisk indleveringsdag: **2014-01-07**
- (87) Den europæiske ansøgnings publiceringsdag: **2015-11-18**
- (86) International ansøgning nr.: **US2014010553**
- (87) Internationalt publikationsnr.: **WO2014110053**
- (30) Prioritet: **2013-01-08 US 201361750175 P**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **Genzyme Corporation, Genzyme Center , 500 Kendall Street, Cambridge, MA 02142, USA**
- (72) Opfinder: **PECHAN, Peter, 500 Kendall Square, Cambridge, MA 02141, USA**
ARDINGER, Jeffery, 500 Kendall Square, Cambridge, MA 02141, USA
SCARIA, Abraham, 500 Kendall Square, Cambridge, MA 02141, USA
WADSWORTH, Samuel, 500 Kendall Square, Cambridge, MA 02141, USA
- (74) Fuldmægtig i Danmark: **RWS Group, Europa House, Chiltern Park, Chiltern Hill, Chalfont St Peter, Bucks SL9 9FG, Storbritannien**
- (54) Benævnelse: **ANVENDELSE AF INOS-INHIBITORER TIL FORØGELSE AF VIRALT UDBYTTE I KULTUR**
- (56) Fremdragne publikationer:
WO-A2-2004/027048
MYSKIW CHAD ET AL: "Aurintricarboxylic acid inhibits the early stage of vaccinia virus replication by targeting both cellular and viral factors", JOURNAL OF VIROLOGY, vol. 81, no. 6, March 2007 (2007-03), pages 3027-3032, XP002725934, ISSN: 0022-538X
DATABASE BIOSIS [Online] BIOSCIENCES INFORMATION SERVICE, PHILADELPHIA, PA, US; 1992, NEYTS J ET AL: "POLYHYDROXYCARBOXYLATES AS SELECTIVE INHIBITORS OF CYTOMEGALOVIRUS AND HERPES SIMPLEX VIRUS REPLICATION", XP002725935, Database accession no. PREV199294082420
GIL-FERNANDEZ C ET AL: "Aurintricarboxylic acid: An inhibitor of herpes simplex types I and II, vaccinia and African swine fever viruses replication in vero cells", ANTIVIRAL RESEARCH, ELSEVIER BV, NL, vol. 15, 1 April 1991 (1991-04-01), page 75, XP023704130, ISSN: 0166-3542, DOI: 10.1016/0166-3542(91)90145-H [retrieved on 1991-04-01]
DATABASE BIOSIS [Online] BIOSCIENCES INFORMATION SERVICE, PHILADELPHIA, PA, US; 1992, GARCIA-VILLALON D ET AL: "ANIONIC COMPOUNDS AS INHIBITORS OF AFRICAN SWINE FEVER VIRUS REPLICATION IN VERO CELLS", XP002725936, Database accession no. PREV199293108974
DATABASE BIOSIS [Online] BIOSCIENCES INFORMATION SERVICE, PHILADELPHIA, PA, US; 1981,

Fortsættes ...

DOROKHOV YU L ET AL: "EFFECT OF AURINTRICARBOXYLIC-ACID ON TOBACCO MOSAIC VIRUS REPRODUCTION", XP002725937, Database accession no. PREV198375059919

DOROKHOV Y L ET AL: "Stimulation by aurintricarboxylic acid of tobacco mosaic virus-specific RNA synthesis and production of informosome-like infection-specific ribonucleoprotein", VIROLOGY, ELSEVIER, AMSTERDAM, NL, vol. 135, no. 2, 1 June 1984 (1984-06-01), pages 395-405, XP023048664, ISSN: 0042-6822, DOI: 10.1016/0042-6822(84)90195-8 [retrieved on 1984-06-01]

MOLECULAR THERAPY - METHODS AND CLINICAL DEVELOPMENT, vol. 1, no. 6, 19 February 2014 (2014-02-19), pages 1-5, XP002725938,

DESCRIPTION

TECHNICAL FIELD

[0001] The present disclosure relates generally to methods for producing viruses and recombinant virions in culture. In particular, the disclosure pertains to the use of the iNOS inhibitors, such as aurintricarboxylic acid, dexamethasone and valproic acid to increase the yield of a variety of viruses in culture, including recombinant herpesviruses which can in turn be used as helpers for the production of recombinant adeno-associated virus virions.

BACKGROUND

[0002] Herpesviruses are highly disseminated in nature and found in most animal species. At least 100 herpesviruses have been characterized, including several from humans, such as herpes simplex virus-1 (HSV-1) and herpes simplex virus-2 (HSV-2), varicella zoster virus (VZV), Epstein-Barr virus (EBV), cytomegalovirus (CMV) and other human herpesviruses such as HHV6 and HHV7. These viruses are responsible for a variety of human diseases, such as skin infections, genital herpes, viral encephalitis, and the like.

[0003] HSV-1 infection activates the host defense and innate immune system by inducing intracellular signaling pathways that lead to the expression of proteins with proinflammatory and microbicidal activities, including cytokines and interferons (INF) (Sainz and Halford, J. Virol. (2002) 76:11541-11550; Haller et al., Virology (2006) 344:119-130; Paludan et al., Nat. Rev. Immunol. (2011) 11:143-154). INF signaling is one of the most important cellular defense mechanism for viral clearance (Brandner & Mueller, Hoppe-Seyler's Zeitschrift für physiologische Chemie (1973) 354:1176; De Vries et al., Gene Ther. (2008) 15:545-552).

[0004] Investigators have reported antiviral activity of nitric oxide (NO) against several viruses such as vaccinia virus, vesicular stomatitis virus, and Japanese encephalitis virus, among others (Bi et al., J. Virol. (1995) 69:6466-6472; Harris et al., J. Virol. (1995) 69:910-915; Lin et al., J. Virol. (1997) 71:5227-5235; Pertile et al., Avian Dis. (1996) 40:342-348. NO is a free radical gaseous molecule and is a mediator of host defense (Croen K.D., J. Clin. Invest. (1993) 91 :2446-2452; Karupiah et al., Science (1993) 261:1445-1448; Rolph et al., Virol. (1996) 217:470-477; Amaro et al., J. Med. Virol. (1997) 51:326-331; Lane et al., J. Virol. (1997) 71:2202-2210. HSV-1 is known both to induce and evade host antiviral responses (Mossman et al., J. Virol. (2001) 75:750-758). HSV infection is capable of inducing expression of inducible nitric oxide synthase (iNOS), a gene encoding an inducible isoform of NOS that produces large amounts of NO.

[0005] Herpesviruses and recombinant proteins therefrom have been used in the manufacture of a number of vaccines. Besides adenoviruses, herpesviruses have been shown to provide complete helper virus functions for the production of recombinant adeno-associated virus virions (Buller, R.M.L., J. Virol. (1981) 40:241-247; Mishra et al., Virology (1990) 179:632-639). The minimal set of HSV-1 genes required for AAV replication and packaging has been identified as the early genes UL5, UL8, UL52 and UL29 (Weindler et al., J. Virol. (1991) 65:2476-2483). These genes encode components of the HSV-1 core replication machinery - the helicase, primase and primase accessory proteins (UL5, UL8 and UL52) and the single-stranded DNA binding protein (UL29).

[0006] Recombinant AAV (rAAV) vectors have been successfully used to achieve long-term, high level transduction *in vivo*. Despite the above advances, production of large quantities of clinical grade high-titer

rAAV virions for gene therapy continues to be challenging due to limitations in scalability of the cotransfection protocol. The process requires the efficient cellular delivery of three components: (1) a vector including the gene of interest flanked by AAV inverted terminal repeats (ITRs); (2) a vector including the AAV *rep* and *cap* genes; and (3) genes provided using a helper virus, such as adenovirus or herpes simplex virus or using virus-free helper plasmids (see, Muzyczka, N., Curr. Top. Microbiol. Immunol. (1992) 158:97-129). Thus, in rHSV-based rAAV manufacturing protocols, the yield of rAAV is limited by the maximal titer of helper rHSV vectors.

[0007] A replication-deficient HSV-1 vector, termed d27.1-rc, expresses AAV-2 *rep* and *cap* genes (Conway et al., Gene Ther. (1999) 6:986-993) and it has been engineered from original d27-1 virus (Rice et al., J. Virol. 1989 vol. 63 (8) pp. 3399-407), that does not produce ICP27, a protein required for HSV-1 replication. Although this vector is replication-defective, it does express the HSV-1 early genes required for rAAV replication and packaging (Conway et al., Gene Ther. (1999) 6:986-993).

[0008] Typically, one vector bearing the rAAV template and the other vector expressing the AAV *rep* and *cap* regions are co-infected into 293 cells in order to produce rAAV virions. Both HSV-1 vectors are replication-deficient and can therefore only be propagated in an ICP27-complementing cell line, V27 (Rice et al., J. Virol. 1989 vol. 63 (8) pp. 3399-407). In HSV-based AAV production protocol, 293 cells need to be infected with HSV-1 at a higher multiplicity of infection (MOI) of 12. This represents a limitation, because yields of d27-1-derived vectors in V27 cells are typically around 1×10^7 plaque forming units (PFU)/ml.

[0009] Several methods and reagents have been investigated in order to further increase HSV-1 titers (see, e.g., Wechuck et al., Biotechnol. Prog. (2000) 16:493-496; Ozuer et al., Biotechnol. Prog. (2002) 18:476-482; Erlandsson et al., J. Endocrinol., (2002) 175:165-176; Otsuki et al., Mol. Ther. (2008) 16:1546-1555). Both dexamethasone and valproic acid inhibited the host defense mechanism represented by several interferon (IFN)-responsive antiviral genes, augmented the transcriptional level of viral genes, and thus improved viral propagation and yield of HSV-1 (Erlandsson et al., J. Endocrinol. (2002) 175:165-176; Otsuki et al., Mol. Ther. (2008) 16:1546-1555).

[0010] Despite the above knowledge, more methods to inhibit host defense in order to improve viral production in culture are needed. As explained above, investigators have reported antiviral activity of nitric oxide (NO) against several viruses such as vaccinia virus, vesicular stomatitis virus, and Japanese encephalitis virus, among others (Bi et al., J. Virol. (1995) 69:6466-6472; Harris et al., J. Virol. (1995) 69:910-915; Lin et al., J. Virol. (1997) 71:5227-5235; Pertile et al., Avian Dis. (1996) 40:342-348. NO is a free radical gaseous molecule and is a mediator of host defense (Croen K.D., J. Clin. Invest. (1993) 91 :2446-2452; Karupiah et al., Science (1993) 261:1445-1448; Rolph et al., Virol. (1996) 217:470-477; Amaro et al., J. Med. Virol. (1997) 51:326-331; Lane et al., J. Virol. (1997) 71:2202-2210). As described above, HSV infection can induce expression of iNOS, a gene encoding an inducible isoform of NOS that produces large amounts of NO.

[0011] The presence of the iNOS inhibitor N-methyl-L-arginine (L-NMA) reversed the inhibition of viral replication for all three of these viruses (Karupiah et al., Science (1993) 261:1445-1448). For a review of iNOS inhibitors, see, Southan et al., Biochem. Pharmacol. (1996) 51:383-394. Another compound, aurintricarboxylic acid (ATA), has been shown to protect macrophages from cell death induced by bacterial lipopolysaccharide by downregulation of iNOS expression and thus decreasing the NO production (Chen et al., British Journal of Pharmacology (2002) vol. 137 (7) pp. 1011-20). ATA is a heterogeneous mixture of polymers accredited with an increasing number of biological activities, such as interaction with a number of enzymes including DNA polymerases, RNA polymerases, reverse transcriptase (RNA-dependent DNA polymerase), aminoacyl-tRNA-synthetase, ribonucleotide reductase, ribonucleases nuclease, protein synthesis inhibition, prevention of apoptosis and blocking DNA fragmentation in oligodendrocytes induced by oxidative stress (Tscherne and Pestka, Antimicrob. Agents Chemother. (1975) 8:479-487; Mikelens et al., Biochemical Pharmacology (1976) 25:821-827; Vollgraf et al., J. Neurochem.(1999) 73:2501-2509).

[0012] Aurintricarboxylic acid (ATA) has been also reported to prevent IFN-mediated transcriptional activation (Tsi et al., Mol. Pharmacol. (2002) 101:90-101; Chen et al., British J. Pharmacol. (2002) 137:1011-1020). ATA is known as an activator of the Raf/MEK/MAPK pathway, IGF-1 receptor and protein kinase C signaling (Beery et al., Endocrinology (2001) 142:3098-3107; Chen et al., J. Biol. Chem. (2001) 276:46722-46728). Antiviral antimicrobial and antiproliferative actions of cytokines such as interferons may be due to their ability to induce the expression of *iNOS*, a gene encoding an isoform of nitric oxide synthase (NOS) that produces large amounts of the radical gas, NO, from a guanidino nitrogen of L-arginine (Nathan, C., FASEB J. (1992) 6:3051; Werner-Felmayer et al., J. Exp. Med. (1990) 172:1599). It has been shown that treatment of macrophages with IFN- γ severely restricts replication of ectromelia virus (EV), vaccinia virus (VV) and HSV-1.

[0013] On one hand, ATA is also known as an antiviral agent against several viruses including HIV, herpesvirus HHV-7, SARS-CoV and others (Cushman et al., J. Med. Chem. (1991) 34:329-337; Zhang et al., Antiviral Res. (1999) 43:23-35; Yap et al., Computational Biol. and Chem. (2005) 29:212-219; De Clercq, Advances, and Adventures Med. Res. Rev. (2011) 31:118-160). ATA, however, did not block the replication of adenovirus type 5 (Ad5) in HEK-293 cells (He, Biochem. Biophys. Res. Comm. (2004) 320:1199-1203). Moreover, ATA has been reported to unexpectedly increase titer of a control adenovirus vector in 293 cells while at the same time having antiviral effects on vaccinia virus (Myskiw et al., J. Virol. (2007) 81:3027-3032).

SUMMARY OF DISCLOSURE / INVENTION

[0014] The present disclosure thus overcomes deficiencies in the prior art by addressing problems that limit viral production, such as low production of rHSV, which hampers efforts to produce sufficient quantities of rHSV for a variety of purposes, including for vaccine production, as well as for rAAV virion production in quantities necessary for efficient gene therapy procedures. Using the methods described herein, higher titers of a variety of viruses can be obtained, such as at least an order of magnitude greater than traditional methods.

[0015] In particular, the inventors herein have discovered that aurintricarboxylic acid (ATA) inhibits iNOS and increases HSV production. As shown in the examples herein, micromolar concentrations of ATA in the presence of fetal bovine serum (FBS) increased both HSV-1/d27-1 vector yield in V27 cells and wild-type (wt) HSV-1 virus in Vero, V27 and 293 cells. Other iNOS inhibitors, including dexamethasone and valproic acid, also increased HSV-1 titers in culture in certain instances. HSV-induced iNOS expression was shown to be reduced in HSV+ATA samples as analyzed by SABiosciences Microarray. Similarly, Affymetrix human genome array analysis confirmed that expression of HSV-up-regulated all three nitric oxide synthase genes (nNOS, iNOS and eNOS) were down-regulated in HSV+ATA samples. Affymetrix Gene Array also detected that genes involved in inflammatory IgE and IFN signaling, and general immune responses were upregulated by HSV-1 and suppressed after the addition of ATA. On the other hand, genes primarily involved in cell cycle G1/S, signal transduction in WNT development were significantly down-regulated by HSV and upregulated after addition of ATA.

[0016] These results are significant because of the demand for higher HSV-1 titers for rAAV virion production, as well as for prophylactic, therapeutic and diagnostic purposes. The invention to which the present specification pertains is set out in the claims appended to this description.

BRIEF DESCRIPTION OF THE FIGURES

[0017]

Figure 1 is a representation of a Western blot showing that aurintricarboxylic acid (ATA) inhibits iNOS expression in d27-1-infected V27 lysates.

Figures 2A-2C show the ATA-HSV protocol rationale (Figure 2A) and the optimization of d27-1 HSV-1 titers in V27 supernatants in order to determine which concentrations and conditions for ATA addition have impact on HSV yields. Figure 2B shows viral titers expressed as DNase resistant particles (DRP/ml) at various ATA concentrations in six well plates (Figure 2B) and T150 flasks (Figure 2C).

Figure 3 shows the importance of serum presence in ATA-HSV protocol. Additional optimization and the importance of FBS presence in ATA-HSV protocol was shown on d27-1 HSV-1 titers expressed as drp/ml and pfu/ml.

Figures 4A-4B show the effect of ATA on wild type HSV KOS and McIntyre strains in culture. ATA increased the yield of both virus types in Vero cells, however, ATA appeared to inhibit HSV-1 KOS growth in 293 cells. On the other hand, wtHSV-1 McIntyre strain reached the highest titers after ATA induction in 293 cells. Moreover, ATA appeared to inhibit BOTH types of HSV-1 viruses in HeLa cells.

Figures 5A-5B show the effect of ATA in HSV stocks on the production of rAAV Virions. rAAV titers were slightly increased when using the HSV stock prepared with 20 μ M ATA added during infection. ATA was also shown to increase rAAV titer when 10 μ M ATA was spiked directly into 293 cell media during 2 hrs of the HSV coinfection step.

Figure 6 shows the effect of dexamethasone (Dex) on d27-1/GFP HSV-1 viral titer. The final titers d27-1/GFP HSV-1 were, in general, slightly elevated after dex pretreatments or treatments as compared to untreated control.

Figure 7 shows the effect of pretreatment by valproic acid (VA) on d27-1/GFP HSV-1 viral titer. VA at a concentration 5 mM slightly elevated the titer of d27-1/GFP HSV-1, however the concentrations below and above of 5 mM appeared to have an inhibitory effect on d27-1/GFP HSV-1 titer, as compared to untreated control.

DETAILED DESCRIPTION OF THE INVENTION

[0018] The practice of the present invention will employ, unless otherwise indicated, conventional methods of chemistry, biochemistry, recombinant DNA techniques and immunology, within the skill of the art. Such techniques are explained fully in the literature. See, e.g., Fundamental Virology, 2nd Edition, vol. I & II (B.N. Fields and D.M. Knipe, eds.); Handbook of Experimental Immunology, Vols. I-IV (D.M. Weir and C.C. Blackwell eds., Blackwell Scientific Publications); T.E. Creighton, Proteins: Structures and Molecular Properties (W.H. Freeman and Company, 1993); A.L. Lehninger, Biochemistry (Worth Publishers, Inc., current addition); Sambrook, et al., Molecular Cloning: A Laboratory Manual (2nd Edition, 1989); Methods In Enzymology (S. Colowick and N. Kaplan eds., Academic Press, Inc.).

1. DEFINITIONS

[0019] In describing the present disclosure invention, the following terms will be employed, and are intended to be defined as indicated below.

[0020] It must be noted that, as used in this specification and the appended claims, the singular forms "a", "an" and "the" include plural referents unless the content clearly dictates otherwise. Thus, for example, reference to "a herpesvirus" includes a mixture of two or more such viruses, and the like.

[0021] The terms "recombinant HSV," "rHSV," and "rHSV vector" refer to isolated, genetically modified forms of herpes simplex virus (HSV) containing heterologous genes incorporated into the viral genome. By the term "rHSV/rc" or "rHSV/rc virus" or "rHSV helper function vector" is meant a rHSV in which the AAV *rep* and/or *cap* genes have been incorporated into the rHSV genome. The terms "rHSV expression virus," and "rHSV/AAV" denote a rHSV in which inverted terminal repeat (ITR) sequences from AAV have been incorporated into the rHSV genome.

[0022] The terms "polypeptide" and "protein" refer to a polymer of amino acid residues and are not limited to a minimum length of the product. Thus, peptides, oligopeptides, dimers, multimers, and the like, are included within the definition. Both full-length proteins and fragments thereof are encompassed by the definition. The terms also include postexpression modifications of the polypeptide, for example, glycosylation, acetylation, phosphorylation and the like. Furthermore, for purposes of the present invention, a "polypeptide" refers to a protein which includes modifications, such as deletions, additions and substitutions (generally conservative in nature), to the native sequence, so long as the protein maintains the desired activity. These modifications may be deliberate, as through site-directed mutagenesis, or may be accidental, such as through mutations of hosts which produce the proteins or errors due to PCR amplification. Depending on the expression system used, a polypeptide may include or lack an N-terminal methionine. Additionally, a polypeptide may or may not include the native signal sequence, if one is naturally present. If a signal sequence is not normally present, the protein can be produced with a heterologous sequence.

[0023] A "native" polypeptide refers to a polypeptide having the same amino acid sequence as the corresponding molecule derived from nature. Such native sequences can be isolated from nature or can be produced by recombinant or synthetic means. The term "native" sequence specifically encompasses naturally-occurring truncated or secreted forms of the specific molecule (e.g., an extracellular domain sequence), naturally-occurring variant forms (e.g., alternatively spliced forms) and naturally-occurring allelic variants of the polypeptide.

[0024] By "variant" is meant an active polypeptide as defined herein having at least about 80% amino acid sequence identity with the corresponding full-length native sequence, a polypeptide lacking the signal peptide, an extracellular domain of a polypeptide, with or without a signal peptide, or any other fragment of a full-length polypeptide sequence as disclosed herein. Such polypeptide variants include, for instance, polypeptides wherein one or more amino acid residues are added, or deleted, at the N- and/or C-terminus of the full-length native amino acid sequence. Ordinarily, a variant will have at least about 80% amino acid sequence identity, alternatively at least about 81% amino acid sequence identity, alternatively at least about 82% amino acid sequence identity, alternatively at least about 83% amino acid sequence identity, alternatively at least about 84% amino acid sequence identity, alternatively at least about 85% amino acid sequence identity, alternatively at least about 86% amino acid sequence identity, alternatively at least about 87% amino acid sequence identity, alternatively at least about 88% amino acid sequence identity, alternatively at least about 89% amino acid sequence identity, alternatively at least about 90% amino acid sequence identity, alternatively at least about 91% amino acid sequence identity, alternatively at least about 92% amino acid sequence identity, alternatively at least about 93% amino acid sequence identity, alternatively at least about 94% amino acid sequence identity, alternatively at least about 95% amino acid sequence identity, alternatively at least about 96% amino acid sequence identity, alternatively at least about 97% amino acid sequence identity, alternatively at least about 98% amino acid sequence identity and alternatively at least about 99% amino acid sequence identity to the corresponding full-length native sequence. Ordinarily, variant polypeptides are at least about 10 amino acids in length, such as at least about 20 amino acids in length, e.g., at least about 30 amino acids in length, alternatively at least about 40 amino acids in length, alternatively at least about 50 amino acids in

length, alternatively at least about 60 amino acids in length, alternatively at least about 70 amino acids in length, alternatively at least about 80 amino acids in length, alternatively at least about 90 amino acids in length, alternatively at least about 100 amino acids in length, alternatively at least about 150 amino acids in length, alternatively at least about 200 amino acids in length, alternatively at least about 300 amino acids in length, or more.

[0025] Particularly preferred variants include substitutions that are conservative in nature, i.e., those substitutions that take place within a family of amino acids that are related in their side chains. Specifically, amino acids are generally divided into four families: (1) acidic -- aspartate and glutamate; (2) basic -- lysine, arginine, histidine; (3) non-polar -- alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan; and (4) uncharged polar -- glycine, asparagine, glutamine, cysteine, serine, threonine, tyrosine. Phenylalanine, tryptophan, and tyrosine are sometimes classified as aromatic amino acids. For example, it is reasonably predictable that an isolated replacement of leucine with isoleucine or valine, an aspartate with a glutamate, a threonine with a serine, or a similar conservative replacement of an amino acid with a structurally related amino acid, will not have a major effect on the biological activity. For example, the polypeptide of interest may include up to about 5-10 conservative or non-conservative amino acid substitutions, or even up to about 15-25 or 50 conservative or non-conservative amino acid substitutions, or any number between 5-50, so long as the desired function of the molecule remains intact.

[0026] "Homology" refers to the percent identity between two polynucleotide or two polypeptide moieties. Two DNA, or two polypeptide sequences are "substantially homologous" to each other when the sequences exhibit at least about 50% , preferably at least about 75%, more preferably at least about 80%-85%, preferably at least about 90%, and most preferably at least about 95%-98% sequence identity over a defined length of the molecules. As used herein, substantially homologous also refers to sequences showing complete identity to the specified DNA or polypeptide sequence.

[0027] In general, "identity" refers to an exact nucleotide-to-nucleotide or amino acid-to-amino acid correspondence of two polynucleotides or polypeptide sequences, respectively. Percent identity can be determined by a direct comparison of the sequence information between two molecules by aligning the sequences, counting the exact number of matches between the two aligned sequences, dividing by the length of the shorter sequence, and multiplying the result by 100. Readily available computer programs can be used to aid in the analysis, such as ALIGN, Dayhoff, M.O. in Atlas of Protein Sequence and Structure M.O. Dayhoff ed., 5 Suppl. 3:353-358, National Biomedical Research Foundation, Washington, DC, which adapts the local homology algorithm of Smith and Waterman Advances in Appl. Math.2:482-489, 1981 for peptide analysis. Programs for determining nucleotide sequence identity are available in the Wisconsin Sequence Analysis Package, Version 8 (available from Genetics Computer Group, Madison, WI) for example, the BESTFIT, FASTA and GAP programs, which also rely on the Smith and Waterman algorithm. These programs are readily utilized with the default parameters recommended by the manufacturer and described in the Wisconsin Sequence Analysis Package referred to above. For example, percent identity of a particular nucleotide sequence to a reference sequence can be determined using the homology algorithm of Smith and Waterman with a default scoring table and a gap penalty of six nucleotide positions.

[0028] Another method of establishing percent identity in the context of the present disclosure is to use the MPSRCH package of programs copyrighted by the University of Edinburgh, developed by John F. Collins and Shane S. Sturrock, and distributed by IntelliGenetics, Inc. (Mountain View, CA). From this suite of packages the Smith-Waterman algorithm can be employed where default parameters are used for the scoring table (for example, gap open penalty of 12, gap extension penalty of one, and a gap of six). From the data generated the "Match" value reflects "sequence identity." Other suitable programs for calculating the percent identity or similarity between sequences are generally known in the art, for example, another alignment program is BLAST, used with default parameters. For example, BLASTN and BLASTP can be used using the following default parameters: genetic code = standard; filter = none; strand = both; cutoff = 60; expect = 10; Matrix =

BLOSUM62; Descriptions = 50 sequences; sort by = HIGH SCORE; Databases = non-redundant, GenBank + EMBL + DDBJ + PDB + GenBank CDS translations + Swiss protein + Spupdate + PIR. Details of these programs are well known in the art.

[0029] Alternatively, homology can be determined by hybridization of polynucleotides under conditions which form stable duplexes between homologous regions, followed by digestion with single-stranded-specific nuclease(s), and size determination of the digested fragments. DNA sequences that are substantially homologous can be identified in a Southern hybridization experiment under, for example, stringent conditions, as defined for that particular system. Defining appropriate hybridization conditions is within the skill of the art. See, e.g., Sambrook et al., *supra*; *DNA Cloning*, *supra*; *Nucleic Acid Hybridization*, *supra*.

[0030] By the term "degenerate variant" is intended a polynucleotide containing changes in the nucleic acid sequence thereof, that encodes a polypeptide having the same amino acid sequence as the polypeptide encoded by the polynucleotide from which the degenerate variant is derived.

[0031] A "coding sequence" or a sequence which "encodes" a selected polypeptide, is a nucleic acid molecule which is transcribed (in the case of DNA) and translated (in the case of mRNA) into a polypeptide *in vivo* when placed under the control of appropriate regulatory sequences. The boundaries of the coding sequence are determined by a start codon at the 5' (amino) terminus and a translation stop codon at the 3' (carboxy) terminus. A transcription termination sequence may be located 3' to the coding sequence.

[0032] By "vector" is meant any genetic element, such as a plasmid, phage, transposon, cosmid, chromosome, virus, virion, etc., which is capable of replication when associated with the proper control elements and which can transfer gene sequences to cells. Thus, the term includes cloning and expression vehicles, as well as viral vectors.

[0033] By "recombinant vector" is meant a vector that includes a heterologous nucleic acid sequence which is capable of expression *in vivo*.

[0034] By "recombinant virus" is meant a virus that has been genetically altered, e.g., by the addition or insertion of a heterologous nucleic acid construct into the particle.

[0035] The term "transgene" refers to a polynucleotide that is introduced into a cell and is capable of being transcribed into RNA and optionally, translated and/or expressed under appropriate conditions. In one aspect, it confers a desired property to a cell into which it was introduced, or otherwise leads to a desired therapeutic or diagnostic outcome.

[0036] The terms "genome particles (gp)," and "genome equivalents," as used in reference to a viral titer, refer to the number of virions containing the recombinant AAV DNA genome, regardless of infectivity or functionality. The number of genome particles in a particular vector preparation can be measured by procedures known in the art, such as described in for example, in Clark et al., *Hum. Gene Ther.* (1999) 10:1031-1039; and Veldwijk et al., *Mol. Ther.* (2002) 6:272-278.

[0037] The terms "infection unit (iu)," "infectious particle," or "replication unit," as used in reference to a viral titer, refer to the number of infectious recombinant AAV vector particles as measured by the infectious center assay, also known as replication center assay, as described, for example, in McLaughlin et al., *J. Virol.* (1988) 62:1963-1973.

[0038] The term "transducing unit (tu)" as used in reference to a viral titer, refers to the number of infectious recombinant AAV vector particles that result in the production of a functional transgene product as measured in functional assays such as described in, for example, in Xiao et al., *Exp. Neurobiol.* (1997) 144:1 13-124; or

in Fisher et al., J. Virol. (1996) 70:520-532 (LFU assay).

[0039] The term "transfection" is used to refer to the uptake of foreign DNA by a cell, and a cell has been "transfected" when exogenous DNA has been introduced inside the cell membrane. A number of transfection techniques are generally known in the art. See, e.g., Graham et al. (1973) Virology, 52 :456, Sambrook et al. (1989) Molecular Cloning, a laboratory manual, Cold Spring Harbor Laboratories, New York, Davis et al. (1986) Basic Methods in Molecular Biology, Elsevier, and Chu et al. (1981) Gene13:197. Such techniques can be used to introduce one or more exogenous DNA moieties into suitable host cells.

[0040] The term "heterologous" as it relates to nucleic acid sequences such as coding sequences and control sequences, denotes sequences that are not normally joined together, and/or are not normally associated with a particular cell. Thus, a "heterologous" region of a nucleic acid construct or a vector is a segment of nucleic acid within or attached to another nucleic acid molecule that is not found in association with the other molecule in nature. For example, a heterologous region of a nucleic acid construct could include a coding sequence flanked by sequences not found in association with the coding sequence in nature. Another example of a heterologous coding sequence is a construct where the coding sequence itself is not found in nature (e.g., synthetic sequences having codons different from the native gene). Similarly, a cell transformed with a construct which is not normally present in the cell would be considered heterologous for purposes of this invention. Allelic variation or naturally occurring mutational events do not give rise to heterologous DNA, as used herein.

[0041] A "nucleic acid" sequence refers to a DNA or RNA sequence. The term captures sequences that include any of the known base analogues of DNA and RNA such as, but not limited to 4-acetylcytosine, 8-hydroxy-N6-methyladenosine, aziridinylcytosine, pseudoisocytosine, 5-(carboxyhydroxyl-methyl) uracil, 5-fluorouracil, 5-bromouracil, 5-carboxymethylaminomethyl-2-thiouracil, 5-carboxymethyl-aminomethyluracil, dihydrouracil, inosine, N6-isopentenyladenine, 1-methyladenine, 1-methylpseudo-uracil, 1-methylguanine, 1-methylinosine, 2,2-dimethyl-guanine, 2-methyladenine, 2-methylguanine, 3-methyl-cytosine, 5-methylcytosine, N6-methyladenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxy-amino-methyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarbonylmethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid, oxybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, -uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid, pseudouracil, queosine, 2-thiocytosine, and 2,6-diaminopurine.

[0042] The term DNA "control sequences" refers collectively to promoter sequences, polyadenylation signals, transcription termination sequences, upstream regulatory domains, origins of replication, internal ribosome entry sites ("IRES"), enhancers, and the like, which collectively provide for the replication, transcription and translation of a coding sequence in a recipient cell. Not all of these control sequences need always be present so long as the selected coding sequence is capable of being replicated, transcribed and translated in an appropriate host cell.

[0043] The term "promoter" is used herein in its ordinary sense to refer to a nucleotide region comprising a DNA regulatory sequence, wherein the regulatory sequence is derived from a gene which is capable of binding RNA polymerase and initiating transcription of a downstream (3'-direction) coding sequence. Transcription promoters can include "inducible promoters" (where expression of a polynucleotide sequence operably linked to the promoter is induced by an analyte, cofactor, regulatory protein, etc.), "repressible promoters" (where expression of a polynucleotide sequence operably linked to the promoter is induced by an analyte, cofactor, regulatory protein, etc.), and "constitutive promoters".

[0044] "Operably linked" refers to an arrangement of elements wherein the components so described are configured so as to perform their usual function. Thus, control sequences operably linked to a coding sequence are capable of effecting the expression of the coding sequence. The control sequences need not

be contiguous with the coding sequence, so long as they function to direct the expression thereof. Thus, for example, intervening untranslated yet transcribed sequences can be present between a promoter sequence and the coding sequence and the promoter sequence can still be considered "operably linked" to the coding sequence.

[0045] By "isolated" when referring to a protein or nucleotide sequence, is meant that the indicated molecule is present in the substantial absence of other biological macromolecules of the same type. Thus, for example, an "isolated nucleic acid molecule which encodes a particular polypeptide" refers to a nucleic acid molecule which is substantially free of other nucleic acid molecules that do not encode the subject polypeptide; however, the molecule may include some additional bases or moieties which do not deleteriously affect the basic characteristics of the composition.

[0046] For the purpose of describing the relative position of nucleotide sequences in a particular nucleic acid molecule throughout the instant application, such as when a particular nucleotide sequence is described as being situated "upstream," "downstream," "3-prime (3′)" or "5-prime (5′)" relative to another sequence, it is to be understood that it is the position of the sequences in the "sense" or "coding" strand of a DNA molecule that is being referred to as is conventional in the art.

[0047] The term "about", particularly in reference to a given quantity, is meant to encompass deviations of plus or minus five percent.

2. MODES OF CARRYING OUT THE INVENTION

[0048] Before describing the present invention in detail, it is to be understood that this invention is not limited to particular formulations or process parameters as such may, of course, vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments of the invention only, and is not intended to be limiting.

[0049] Although a number of methods and materials similar or equivalent to those described herein can be used in the practice of the present invention, the preferred materials and methods are described herein.


[0050] Central to the present invention is the discovery that aurotricarboxylic acid (ATA) in micromolar concentrations increases HSV-1 vector yield. This finding is important for both large-scale HSV production, as well as rHSV and rAAV vector production. Moreover and surprisingly, as shown in the examples, the presence of ATA in rHSV-1 stocks did not negatively influence rAAV yield. This result is surprising as ATA in millimolar amounts and higher concentrations is known to be an antiviral agent (Cushman et al., J. Med. Chem. (1991) 34:329-337; Zhang et al., Antiviral Res. (1999) 43:23-35; Yap et al., Computational Biol. and Chem. (2005) 29:212-219; De Clercq, Advents, Advances, and Adventures Med. Res. Rev. (2011) 31:118-160).

[0051] As explained above, investigators have reported antiviral activity of nitric oxide (NO) against several viruses such as vaccinia virus, vesicular stomatitis virus, and Japanese encephalitis virus, among others (Bi et al., J. Virol. (1995) 69:6466-6472; Harris et al., J. Virol. (1995) 69:910-915; Lin et al., 1997; Pertile et al., Avian Dis. (1996) 40:342-348). NO is a free radical gaseous molecule is a mediator of host defense (Croen K.D., J. Clin. Invest. (1993) 91 :2446-2452; Karupiah et al., Science (1993) 261:1445-1448; Rolph et al., Virol. (1996) 217:470-477; Amaro et al., J. Med. Virol. (1997) 51:326-331; Lane et al., J. Virol. (1997) 71:2202-2210). HSV infection is capable of inducing the expression of inducible nitric oxide synthase (iNOS), a gene encoding an inducible isoform of NOS that produces large amounts of NO.

[0052] As shown herein, ATA suppresses HSV-up-regulated iNOS and thereby increases HSV titers in culture. Additional iNOS inhibitors, including dexamethasone and valproic acid, also have the same effect. In

one embodiment, then, the use of such iNOS inhibitors increases recombinant herpesvirus titers in culture, allowing the production of significantly more virus than produced in the absence of the particular inhibitor. Viruses produced by the method can be used in a variety of contexts, including for prophylactic, therapeutic and diagnostic purposes, as well as for producing recombinant constructs in sufficient quantity to use in the preparation of recombinant virions for gene delivery and gene therapy.

[0053] Aurintricarboxylic acid (ATA), 5-((3-carboxy-4-hydroxyphenyl)(3-carboxy-4-oxo-2,5-cyclohexadien-1-ylidene)methyl)-2-hydroxybenzoic acid, is a heterogeneous mixture of nonsulfated negatively-charged aromatic polymers that forms when salicylic acid is treated with formaldehyde, sulfuric acid and sodium nitrite (see Cushman, et al., (1991) J. Med. Chem.34:329-337; Cushman, et al., J. Med. Chem. 34:337-342). Aurintricarboxylic acid has the formula:



[0054] The heterogeneous mixture of ATA was described to inhibit protein nucleic acid interactions (Gonzalez et al., Biochim. Biophys. Acta, (1979) 562:534-545); to interact with steroid receptors at the nuclear uptake and the nuclear binding levels (Mellon, W. S., Biochem. Pharmacol. (1984) 33:1047-1057; Moudgillet al., J. Steroid Biochem. (1985) 23:125-132); to inhibit DNA polymerase (Nakane et al., Eur. J. Biochem. (1988) 177:91-96); and to act as a RNAase inhibitor (Skidmore et al., Biochem. J. (1989) 263:73-80).

[0055] ATA for addition to virus in culture can be used in the acidic form or can be provided as a salt, such as aurintricarboxylic acid trisodium salt; calcium salt; ammonium salt, etc.

[0056] Additional substances that will find use with the present methods include dexamethasone (Dex) and valproic acid (VA). Dex has been shown to inhibit iNOS expression in rat mesangial cells at transcriptional and post-transcriptional levels (Kunz et al., Biochem. J. (1994) 304:337-340; Kunz et al., Proc. Natl. Acad. Sci. USA (1996) 93:255-259). Dex has also been shown to enhance HSV-1 oriL DNA replication in PC12 cells. Sodium valproate, the sodium salt of VA, has been shown to stimulate replication of HSV-1, human cytomegalovirus, HIF-1, human herpes virus-8, measles virus and poliovirus type 1 (Motamedifar et al., Iran. J. Med. Sci. (2006) 31:28-32; Kuntz-Simon et al., J. Gen. Virol (1995) 76:1409-1415; Moog et al., J. Gen. Virol (1996) 77:1993-1999; Ylisastigui et al., AIDS (2004) 18:1101-1108; Shaw et al., AIDS (2000) 14:899-902; Kabiri et al. Iran J. Med. Sci. (2001) 26:55-61). Additionally, valproic acid has been shown to inhibit iNOS (Guo et al., Surgery (2007) 142:156-162). As with ATA, VA, or its salts, such as sodium, calcium, ammonium salts, etc., can be used in the present methods.

[0057] Although the use of ATA, Dex, and VA to produce rHSV-1 vectors at higher titers is exemplified herein, these iNOS inhibitors can be used to increase the titer of a variety of viruses in culture, such as but not limited to viruses of the families Adenoviridae, Picornaviridae (e.g., polioviruses, etc.); Caliciviridae; Togaviridae (e.g., rubella virus, dengue virus, etc.); Flaviviridae; Coronaviridae; Reoviridae; Birnaviridae; Rhabdoviridae (e.g., rabies virus, etc.); Poxviridae; Filoviridae; Paramyxoviridae (e.g., mumps virus, measles virus, respiratory syncytial virus, etc.); Orthomyxoviridae (e.g., influenza virus types A, B and C, etc.); Bunyaviridae; Arenaviridae; the various hepatitis viruses, such as HAV, HBV and HCV; papilloma viruses and rotaviruses; retroviruses; etc. See, e.g. Virology, 3rd Edition (W.K. Joklik ed. 1988); Fundamental Virology, 2nd Edition (B.N. Fields and D.M. Knipe, eds. 1991), for a description of these and other viruses. These viruses, or immunogens derived therefrom, can be used in the production of vaccines and diagnostics. Moreover, some

of these viruses, and in particular, the herpesviruses, can be used to produce recombinant vectors for the production of recombinant virions for use in gene delivery techniques described below.

[0058] Thus, ATA, Dex and VA can be used to increase the yield of any of the herpesviruses that are members of the family herpesviridae. This includes equine herpes virus, bovine herpes virus (BHV) and human herpes simplex virus (HSV) types 1 and 2, such as BHV-1, BHV-2, HSV-1 and HSV-2, varicella zoster virus (VZV), Epstein-Barr virus (EBV), cytomegalovirus (CMV), HHV6 and HHV7, among others. The herpesviruses can be derived from any of the many strains. For example, when the virus produced using the invention is HSV, the virus may be derived from, for example, HSV-1 or HSV-2, and may be from any of the various HSV strains, such as HSV-1 strain KOS, HSV-1 strain McIntyre, HSV-1 strain Patton, HSV-2 strain 333, HSV-2 strain G, and the like. Moreover, the viruses produced may be either wild-type viruses, or derivatives thereof, including recombinant viruses and inter-type recombinants containing DNA from HSV-1 and HSV-2. Derivatives preferably have at least 70% sequence homology to either the HSV-1 or HSV-2 genomes or portions thereof, more preferably at least 80%, even more preferably at least 90 or 95%. A derivative may have the sequence of a HSV-1 or HSV-2 genome modified by nucleotide substitutions, for example from 1, 2 or 3 to 10, 25, 50 or 100 substitutions. The HSV-1 or HSV-2 genome may alternatively or additionally be modified by one or more insertions and/or deletions and/or by an extension at either or both ends.

[0059] Other derivatives include strains that already have mutations in genes, particularly mutations in genes that result in attenuation of the virus. Examples of such viruses include strain 1716 (MacLean et al., J. Gen. Virol. (1991) 72:632-639), strains R3616 and R4009 (Chou and Roizman, Proc. Natl. Acad. Sci USA (1992) 89: 3266-3270) and R930 (Chou et al., J. Virol. (1994) 68:8304-8311) all of which have mutations in ICP34.5, strain d120 which has a deletion in ICP4 (DeLuca et al., J. Virol. (1985) 56:558-570), strain d27-1 (Rice and Knipe, J. Virol. (1990) 64:1704-1715) which has a deletion in ICP27 or strain d92 which has deletions in both ICP27 and ICP4 (Samaniego et al., J. Virol. (1995) 69:5705-5715). The terminology used in describing the various HSV genes is as found in, e.g., Coffin and Latchman (1996), In: Genetic Manipulation of the Nervous System (DS Latchman Ed.) pp 99-114: Academic Press, London.

[0060] As is readily apparent, any rHSV suitable for the intended purpose can be used in the invention. In certain embodiments, the rHSV used in the invention is replication-defective. For the production of rAAV virions, infection of producer cells with rHSV that is incapable of replication is preferred because in contrast to methods involving use of adenovirus, the rHSV does not become a significant contaminant of the rAAV product. This can serve to increase the final yield of rAAV virions by eliminating purification steps associated with removal of adenovirus. In a particular embodiment of the invention, the rHSV is constructed from a mutant of HSV-1 in which the inability to replicate is due to a deletion in the ICP27 gene. Any other suitable mutants of HSV exhibiting a replication-defective phenotype can also be used to construct the rHSV.

[0061] One particularly preferred recombinant mutant HSV-1 strain for rAAV production using the subject methods is HSV-1 strain d27-1. This strain can be prepared as described in e.g., Conway et al., Gene Ther. (1999) 6:973 985 and U.S. Patent No. 7,091,029, incorporated herein by reference in its entirety. As explained above, this mutant vector does not produce ICP27 and is advantageously used to produce rAAV virions as host cell splicing of messenger RNA is known to be inhibited by ICP27. ICP27 may also effect appropriate splicing of the AAV-2 *rep* and *cap* messages. This vector is replication defective and shows reduced cytotoxicity compared with wild type (wt) HSV-1. The virus d27-1 displays several other features that are advantageous for use as a helper virus for rAAV virion production. First, it expresses the early genes known to be required for rAAV production (Weindler et al., J. Virol. (1991) 65:2476-2483). In addition, d27.1 over-expresses ICP8, the single-stranded DNA binding protein that is the product of UL29, one of the HSV-1 genes essential for AAV replication and packaging (Weindler et al., J. Virol. (1991) 65:2476-2483).

[0062] The AAV genome is a linear, single-stranded DNA molecule containing about 4681 nucleotides. The AAV genome generally comprises an internal, nonrepeating genome flanked on each end by inverted terminal

repeats (ITRs). The ITRs are approximately 145 base pairs (bp) in length. The ITRs have multiple functions, including providing origins of DNA replication, and packaging signals for the viral genome. The internal nonrepeated portion of the genome includes two large open reading frames, known as the AAV replication (*rep*) and capsid (*cap*) genes. The *rep* and *cap* genes code for viral proteins that allow the virus to replicate and package into a virion. In particular, a family of at least four viral proteins are expressed from the AAV *rep* region, Rep 78, Rep 68, Rep 52, and Rep 40, named according to their apparent molecular weight. The AAV *cap* region encodes at least three proteins, VP1, VP2, and VP3.

[0063] By "AAV *rep* coding region" is meant the art-recognized region of the AAV genome which encodes the replication proteins Rep 78, Rep 68, Rep 52 and Rep 40. These Rep expression products have been shown to possess many functions, including recognition, binding and nicking of the AAV origin of DNA replication, DNA helicase activity and modulation of transcription from AAV (or other heterologous) promoters. The Rep expression products are collectively required for replicating the AAV genome. For a description of the AAV *rep* coding region, see, e.g., Muzyczka, N. (1992) Current Topics in Microbiol. and Immunol.158:97-129; and Kotin, R.M. (1994) Human Gene Therapy5:793-801. Suitable homologues of the AAV *rep* coding region include the human herpesvirus 6 (HHV-6) *rep* gene which is also known to mediate AAV-2 DNA replication (Thomson et al. (1994) Virology204:304-311).

[0064] By "AAV *cap* coding region" is meant the art-recognized region of the AAV genome which encodes the capsid proteins VP1, VP2, and VP3, or functional homologues thereof. These Cap expression products supply the packaging functions which are collectively required for packaging the viral genome. For a description of the AAV *cap* coding region, see, e.g., Muzyczka, N. and Kotin, R.M. (*supra*).

[0065] Typically two rHSV vectors will be used to produce rAAV virions. One is a rHSV helper function vector in which the AAV *rep* and/or *cap* genes have been incorporated into the rHSV genome. The other is a rHSV expression vector in which ITR sequences from AAV have been incorporated into the rHSV genome and flank a gene of interest.

[0066] Thus, in one embodiment, the iNOS inhibitor can be used to increase the yield of a first rHSV vector that contains the AAV *rep* and/or *cap* genes. Embodiments of the first rHSV vector of the method include but are not limited to gene constructs based on the *cap* gene found in various serotypes of AAV including AAV-1, AAV-2, AAV-3, AAV-4, AAV-5 and AAV-6, AAV-7 and AAV-8, caprine and bovine AAV (see, e.g., U.S. Publ. No. 20080292595, incorporated herein by reference in its entirety), and variants thereof. Also within the scope of the invention are *rep* and *cap* genes from novel AAV serotypes, and those modified by recombination or mutation of existing serotypes. The *rep* and *cap* genes of the AAV helper function vector can be derived from any of the known AAV serotypes, as explained above. For example, the rHSV helper function vector may have a *rep* gene derived from AAV-2 and a *cap* gene derived from AAV-6; one of skill in the art will recognize that other *rep* and *cap* gene combinations are possible, the defining feature being the ability to support rAAV virion production.

[0067] In certain embodiments, the AAV *rep* and *cap* genes in the rHSV helper function vector may be driven by their native promoters. The p5 and p19 promoters of AAV-2 control expression of Rep 78 and 68 and Rep 52 and 40, respectively. The p40 promoter controls expression of VP1, VP2 and VP3. Additionally, heterologous promoters may be used to drive expression of the AAV genes. Examples of other promoters that can be used in the disclosed methods include but are not limited to the SV40 early promoter, CMV promoter, HSV-1 thymidine kinase (HSV-1 tk) promoter, metallothionine inducible promoter, mouse mammary tumor virus promoter and chicken β -action promoter.

[0068] The gene construct can be inserted into any site or sites in the HSV genome suitable for integration of the *rep* and *cap* genes. In certain embodiments, the vector is constructed by homologous recombination of the AAV *rep* and *cap* genes into the thymidine kinase (*tk*) locus of the rHSV-1 virus, as described in Conway et

al., Gene Ther. (1999) 6:986-993 and U.S. Patent No. 7,091,029, incorporated herein by reference in its entirety.

[0069] As explained herein, the rHSV helper function vector encodes the "AAV helper function" sequences (i.e., *rep* and *cap*), which function *in trans* for productive AAV replication and encapsidation. Preferably, the rHSV helper function vector supports efficient rAAV virion production without generating any detectable wt AAV virions (i.e., AAV virions containing functional *rep* and *cap* genes). An example of such a vector is rHSV-1 d27.1rc. The vector and methods of producing the same are described herein in the examples, as well as in Conway et al., Gene Ther. (1999) 6:986-993; and U.S. Patent No. 7,091,029, incorporated herein by reference in its entirety.

[0070] The second rHSV vector is termed a rHSV expression vector and contains ITRs from AAV with one or more genes of interest driven by one or more promoters. In some embodiments, the gene of interest is inserted between a pair of ITRs. The heterologous gene is typically functionally linked to a heterologous promoter (constitutive, cell-specific, or inducible) capable of driving gene expression in the patient's target cells under appropriate conditions. Termination signals, such as polyadenylation sites, can also be included.

[0071] The nucleotide sequences of AAV ITR regions are known. See, e.g., Kotin, R.M. (1994) Human Gene Therapy 5:793-801; Berns, K.I. "Parvoviridae and their Replication" in Fundamental Virology, 2nd Edition, (B.N. Fields and D.M. Knipe, eds.) for the AAV-2 sequence. AAV ITRs used in the vectors of the invention need not have a wild-type nucleotide sequence, and may be altered, e.g., by the insertion, deletion or substitution of nucleotides. Additionally, AAV ITRs may be derived from any of several AAV serotypes, including without limitation, AAV-1, AAV-2, AAV-3, AAV-4, AAV-5, AAV-6, AAV-7 and AAV-8, caprine and bovine AAV (see, e.g., U.S. Publ. No. 20080292595, incorporated herein by reference in its entirety), and variants thereof. Furthermore, 5' and 3' ITRs which flank a selected nucleotide sequence in an expression vector need not necessarily be identical or derived from the same AAV serotype or isolate, so long as they function as intended, i.e., to allow for excision and rescue of the sequence of interest from a host cell genome or vector, and to allow integration of the DNA molecule into the recipient cell genome when AAV *rep* gene products are present in the cell.

[0072] AAV ITRs can be excised from the viral genome or from an AAV vector containing the same and fused 5' and 3' of a selected nucleic acid construct using standard ligation techniques, such as those described in Sambrook et al., *supra*. For example, ligations can be accomplished in 20 mM Tris-Cl pH 7.5, 10 mM MgCl₂, 10 mM DTT, 33 µg/ml BSA, 10 mM-50 mM NaCl, and either 40 µM ATP, 0.01-0.02 (Weiss) units T4 DNA ligase at 0°C (for "sticky end" ligation) or 1 mM ATP, 0.3-0.6 (Weiss) units T4 DNA ligase at 14°C (for "blunt end" ligation). Intermolecular "sticky end" ligations are usually performed at 30-100 µg/ml total DNA concentrations (5-100 nM total end concentration). AAV vectors which contain ITRs have been described in, e.g., U.S. Patent no. 5,139,941. In particular, several AAV vectors are described therein which are available from the American Type Culture Collection ("ATCC") under Accession Numbers 53222, 53223, 53224, 53225 and 53226.

[0073] The selected polynucleotide sequence is operably linked to control elements that direct the transcription or expression thereof in a subject's cells. Such control elements can comprise control sequences normally associated with the selected gene. Alternatively, heterologous control sequences can be employed. Useful heterologous control sequences generally include those derived from sequences encoding mammalian or viral genes. Examples include, but are not limited to, neuron-specific enolase promoter, a GFAP promoter, the SV40 early promoter, mouse mammary tumor virus LTR promoter; adenovirus major late promoter (Ad MLP); a herpes simplex virus (HSV) promoter, a cytomegalovirus (CMV) promoter such as the CMV immediate early promoter region (CMVIE), a rous sarcoma virus (RSV) promoter, synthetic promoters, hybrid promoters, and the like. In addition, sequences derived from nonviral genes, such as the murine metallothionein gene, will also find use herein. Such promoter sequences are commercially available from,

e.g., Stratagene (San Diego, CA), Invivogen (San Diego, CA) and others.

[0074] The gene of interest may be a gene likely to be of therapeutic value. Examples of therapeutic genes include but are not limited to α -1 antitrypsin, Factor VIII, Factor IX, GAA, erythropoietin and PEDF. When it is desirable to select for or to identify successful transgene expression, the gene of interest may be a reporter gene. Many examples of genes used as reporters or for selection are known, and can be used in the invention. These include but are not limited to the genes encoding β -galactosidase, neomycin, phosphotransferase, chloramphenicol acetyl transferase, thymidine kinase, luciferase, beta-glucuronidase, aminoglycoside, phosphotransferase, hygromycin B, xanthine-guanine phosphoribosyl, luciferase, DHFR/methotrexate, and green fluorescent protein (GFP).

[0075] The rHSV-1 expression virus can be produced in much the same manner as described above, namely, by homologous recombination into the HSV-1 tk gene, as described in, e.g., Conway et al., Gene Ther. (1999) 6:986-993 and U.S. Patent No. 7,091,029, incorporated herein by reference in its entirety.

[0076] Once produced, the rHSV vectors, or any other virus of interest, is propagated in culture in an appropriate cell line. For herpesviruses, such cell lines include, but are not limited to, Vero cells, 293 cells, HeLa cells, and the like, available from the American Type Culture Collection, Rockville, Md. If the HSV-1 d27.1 vector is used, this virus will typically be cultured in the ICP27-complementing cell line V27 (Rice et al., J. Virol. (1990) 64:1704-1715). Any suitable medium for the virus in question, with or without serum, such as fetal bovine serum, can be used, such as but not limited to RPMI 1640 medium, Dulbecco's Modified Eagles Medium (DMEM), F12 medium or a mixture of the latter (DF medium). If serum is present, the culture can include, for example 2% to 20% serum, more typically 5% to 15% serum, 7% to 12% serum, or any number within these ranges, such as 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, and the like. Moreover, if serum is used, it can be present in the initial culture, and/or in subsequent fresh media added to the cultures.

[0077] The amount of the iNOS inhibitor to be added in the method according to the invention can be varied and is to a certain extent dependent on the particular inhibitor used, the medium used and the virus to be cultured. For example, when herpesviruses are cultured in Vero cells, the concentration of ATA in the initial culture, if present, will typically be 5 μ M to 500 μ M, preferably 10 μ M to 250 μ M, such as 20 μ M to 100 μ M, i.e., 30 μ M to 75 μ M, such as 30...35...40...45...50...55...60...65...70...75, etc. or any integer within these stated ranges. Similarly, the concentration of dexamethasone if present in the initial culture will typically be 1 μ M to 500 μ M, preferably .5 μ M to 250 μ M, such as 1 μ M to 100 μ M, i.e., 5 μ M to 75 μ M, such as 1... 5... 15... 10... 20... 25... 30...35...40...45...50...55...60...65...70...75, etc. or any integer within these stated ranges. If valproic acid is used, the initial concentration will be .1 mM to 500 mM, preferably .5 mM to 250 mM, such as 1 mM to 100 mM, i.e., 5 mM to 75 mM, such as 1... 5... 15... 10... 25... 30... 35... 40... 45... 50... 55... 60... 65... 70... 75, etc. or any integer within these stated ranges.

[0078] In certain embodiments, virus-infected cells are initially cultured in media as described above for .5 hours to 24 hours, such as .75 hours to 12 hours, 1 hour to 5 hours, 1 hour to 2 hours, or any number of hours or fractions thereof within these ranges. The iNOS inhibitor and/or serum may or may not be present in the initial culture. Fresh media is subsequently added and cultures incubated for 24 to 120 hours, such as 48 to 96 hours, 50 to 80 hours, 60 to 75 hours, 70 to 74 hours, or any number of hours or fractions thereof within these ranges. The iNOS inhibitor and/or serum may or may not be present in the subsequent culture, with the proviso that the iNOS inhibitor is present in either one or both of the initial culture and the subsequent culture.

[0079] In some embodiments, the iNOS inhibitor in the initial culture is present at a higher concentration than in the subsequent culture. Thus, for example, if the iNOS inhibitor is ATA, it can be present in an amount of 30 to 75 μ M in the initial culture and then 5 to 25 μ M in the subsequent culture. Alternatively, the inhibitor can be present only in the initial or the subsequent culture.

[0080] As shown in the examples below, one particularly preferred method using ATA includes the presence of 50 μM ATA in the initial culture, with a reduction in the amount of ATA in the subsequent culture to 20 μM . Additionally, in the case of ATA, it is preferable to include serum at the same time that ATA is present. Thus, if ATA is added to the initial culture, it is advantageous to add fetal bovine serum (FBS) to the media. Likewise, if ATA is added to the subsequent culture, the addition of FBS is necessary to obtain higher viral yields.

[0081] The viruses are then cultured to obtain a desirable titer of virus. For example, in the case of the HSV-1 d27-1 vectors described herein, ATA increases the d27-1 yields in V27 cells supernatants 3- 5 times and the titers can be at least 1×10^8 PFU/ml or 4×10^8 DRP/ml. Similarly, ATA also increases the yields of wild type (wt) HSV-1 viruses strains McIntyre and KOS by 1 log, and the 293 or Vero titers can be up to 1×10^9 DRP/ml. The viruses are then harvested for further use.

[0082] For the purposes of the invention, suitable host cells for producing rAAV virions include microorganisms, yeast cells, insect cells, and mammalian cells, that can be, or have been, used as recipients of a heterologous DNA molecule and that are capable of growth in, for example, suspension culture, flasks, plates, a bioreactor, or the like. The term includes the progeny of the original cell which has been transfected. Thus, a "host cell" as used herein generally refers to a cell which has been transduced with an exogenous DNA sequence. If recombinant herpesviruses are produced in one type of cells for use in making rAAV virions, the harvested vectors are then transfected into another suitable host cell. 293 cells, originated from a stable human cell line, (readily available through, e.g., the American Type Culture Collection under Accession Number ATCC CRL1573) are preferred host cells to produce rAAV virions. Particularly, the human cell line 293 is a human embryonic kidney cell line that has been transformed with adenovirus type-5 DNA fragments (Graham et al. (1977) J. Gen. Virol. 36:59), and expresses the adenoviral E1a and E1b genes (Aiello et al. (1979) Virology 94:460). The 293 cell line is readily transfected, and provides a particularly convenient platform in which to produce rAAV virions.

[0083] AAV helper functions are introduced into the host cell by transducing the host cell with a rHSV helper function construct either prior to, or concurrently with, using the rHSV expression vector. rHSV helper constructs are thus used to provide at least transient expression of AAV *rep* and/or *cap* genes to complement missing AAV functions that are necessary for productive AAV infection. AAV helper constructs lack AAV ITRs and can neither replicate nor package themselves.

[0084] Following recombinant AAV replication, rAAV virions can be purified from the host cell using a variety of conventional purification methods, such as column chromatography, CsCl gradients, and the like. For example, a plurality of column purification steps can be used, such as purification over an anion exchange column, an affinity column and/or a cation exchange column. See, for example, International Publication No. WO 02/12455.

[0085] The resulting rAAV virions containing the nucleotide sequence of interest can then be used for gene delivery using techniques well known in the art and described in e.g., U.S. Patent Nos. 5,173,414 and 5,139,941; International Publication Nos. WO 92/01070 (published 23 January 1992) and WO 93/03769 (published 4 March 1993); Lebkowski et al., Molec. Cell. Biol. (1988) 8:3988-3996; Vincent et al., Vaccines 90 (1990) (Cold Spring Harbor Laboratory Press); Carter, B.J. Current Opinion in Biotechnology (1992) 3:533-539; Muzyczka, N. Current Topics in Microbiol. and Immunol. (1992) 158:97-129; Kotin, R.M. Human Gene Therapy (1994) 5:793-801; Shelling and Smith, Gene Therapy (1994) 1:165-169; and Zhou et al., J. Exp. Med. (1994) 179:1867-1875.

2. EXPERIMENTAL

[0086] Below are examples of specific embodiments for carrying out the present invention. The examples are offered for illustrative purposes only, and are not intended to limit the scope of the present invention in any way.

[0087] Efforts have been made to ensure accuracy with respect to numbers used (e.g., amounts, temperatures, etc.), but some experimental error and deviation should, of course, be allowed for.

Materials and Methods

Cells

[0088] Vero-derived V27 cells (Rice et al., J. Virol. (1989) 63:3399-3407) and human embryonic kidney cells (HEK)-derived 293 cells (Graham et al., J. Gen. Virol. (1977) 36:59-74) were obtained from the Applied Genetic Technologies Corporation (AGTC, Alachua, FL), Vero and HeLa cells were purchased from American Type Culture Collection (ATCC, Manassas, VA). All cells were maintained in Dulbecco's modified Eagle's medium (DMEM; HyClone, South Logan, UT) containing 10% fetal bovine serum (FBS; HyClone) and either Geneticin (50mg/ml; Invitrogen) for V27 cells or 1% penicillin/streptomycin (Cellgro Mediatech, Manassas, VA) for the other cells.

HSV-1 Production

[0089] The wtHSV-1 KOS strain and the ICP27-deficient derivatives of the wtHSV-1 KOS strain: vectors d27-1 (Rice and Knipe, J. Virol. (1990) 64:1704-1715), rHSV-rep2/cap2 and rHSV-EGFP (Kang et al., Gene Ther. (2009) 16:229-239) and their producer ICP27-complementing V27 cell line were obtained from AGTC (Alachua, FL). The wtHSV-1 MacIntyre strain, purchased from Advanced Biotechnologies Inc. (ABI, Columbia MD) and wtHSV-1 KOS strain were propagated in Vero, 293 or HeLa cell lines. Infectious vector particles were harvested 72 hr postinfection by recovering culture supernatant. The titers of HSV-1 stocks in DNase resistant particles/ml (DRP/ml) were determined by Taqman assay. Viral genomes within crude culture medium were quantified via treatment in the presence of DNase I (50 U/ml final; Promega) at 37°C for 60 min, followed by proteinase K (Invitrogen) digestion (1 U/ml) at 50°C for 60 min, and then denatured at 95°C for 30 min. Linearized plasmid pZero 195 UL36 (obtained from AGTC, Inc., Alachua, FL) was used to generate standard curves. The primer-probe set was specific for the vector genome UL36 sequence (HSV-UL36 F: 5'-GTTGGTTATGGGGGAGTGTGG (SEQ ID NO:1); HSV-UL36 R: 5'-TCCTTGCTCTGGGGTGTCTTCG (SEQ ID NO:2); HSV-UL36 Probe: 5'- 6FAM - CGACGAAGACTCCGACGCCACCTC-TAMRA (SEQ ID NO:3). Amplification of the PCR product was achieved with the following cycling parameters: 1 cycle at 50°C for 2 min, 1 cycle at 95°C for 10 min; 40 cycles of 95°C for 15 sec, and 60°C for 60 sec.

ATA Experiments

[0090] ATA (Sigma -A1895 Aurintricarboxylic acid practical grade, ≥85% (titration), powder) stock solution was generated as 500 μM concentration in 100 mM sodium bicarbonate water solution. ATA stock solution was further diluted in DMEM +/- 10% fetal bovine serum (FBS, Hyclone, Waltham, MA) into concentration ranges of 12-60 μM ATA (8.5-21 μgATA /ml). The HSV-1 infection at multiplicity of infection of 0.15 (MOI=0.15) (typically 6x10⁵ cells in a 6-well plate) was performed in 40% (2/5 vol) of the total final media volume for 1-2 hrs and the remaining media (60% or 3/5 of the total final volume) was added during the dilution step. The cells were then

incubated 72 hours and supernatant harvested to perform forming units per milliliter (PFU/ml) and DRP/ml titer assays.

Dexamethasone(Dex) Experiments

[0091] Dexamethasone (Sigma -D4902) was dissolved to 2 mg/ml in absolute alcohol. This was diluted with DMEM to achieve a 1M concentration and stored at -20°C.

[0092] V27 cells were seeded into six well plates the day prior to infection at 6×10^5 cells/well. Dexamethasone was added to achieve a concentration of 1 μ M into the seeding media. This was mixed well and added to the wells.

[0093] Media was aspirated and infectious HSV-1 d27-1 stock was added at MOI 0.15 (= cell seed density x 0.15/ pfu titer of HSV stock) per 1ml of DMEM (no additives). 1 ml of infectious inoculum was added per well. This was incubated for 1-2 hours at 37°C, 5% CO₂ incubator, after which time 1.5 ml of DMEM-10% FBS was added. The cultures were returned to the incubator for 70-74 hours.

[0094] Free media was harvested, vortexed, centrifuged at 1,100 x g for 10 minutes at 4°C. Supernatant was transferred to a new dispensing tube, vortexed, aliquoted and stocks were frozen at -80°C.

Valproic Acid (VA) Experiments

[0095] Valproic acid (Sigma -P4543) was dissolved to 1 M concentration in water. V27 cells were seeded into six well plates the day prior to infection at 6×10^5 cells/well. 1 M valproic acid was spiked into 1 ml DMEM-10% FBS to achieve a concentration of 5 μ M. This was vortexed well and added to the wells. Plates were incubated for six hours, aspirated and infectious HSV-1 d27-1 stock added at MOI 0.15 (= cell seed density x 0.15/ pfu titer of HSV stock) per 1ml of DMEM (no additives).

[0096] 1ml of infectious inoculum was added per well and incubated for 1-2 hours at 37°C, 5% CO₂ incubator after which time 1.5 ml of DMEM-10% FBS was added. The plates were returned to the incubator for 70-74 hours. Free media was harvested, vortexed, and centrifuged at 1,100 x g for 10 minutes at 4°C. Supernatant was transferred to a new dispensing tube, vortexed, and aliquoted. Stocks were frozen at -80°C.

rAAV Production

[0097] 293 cells (2.5×10^6) were simultaneously co-infected with both rHSV-rep2/cap2 and rHSV-EGFP vectors as described by Kang et al., Gene Ther (2009) 16:229-239. At 2-4 hr post infection, infectious medium was exchanged with DMEM +10% FBS equivalent to double the preinfection culture volume. At the time of harvest, the cell pellet was frozen at -80°C. DRP titers were quantified by real-time polymerase chain reaction (qPCR) in a 96-well block thermocycler (Applied Biosystems; 7500 Real Time PCR system). Crude samples were subjected to three cycles of freezing and thawing, then incubated in the presence of Benzonase (250U/ml), 2mM MgCl₂, 1% final concentration protein grade Tween 80 (Calbiochem) and incubated at 37°C for 60 min, followed by 0.25% Trypsin (Gibco) digestion at 50°C for 60 min. Finally, treatment with DNase I (50 U/ml; Promega) at 37°C for 30 min, and then denatured at 95°C for 20 min. Linearized plasmid pDC67/+SV40 was used to generate standard curves. The primer-probe set was specific for the simian virus 40 (SV40) poly(A) sequence: rAAV-F: 5'- AGCAATAGCATCACAAATTTACAA-3' (SEQ ID NO:4); rAAV-R: 5'-

GCAGACATGATAAGATACATTGATGAGTT-3' (SEQ ID NO:5) ;rAAV-Probe: 5' 6-FAM-AGCATTTCCTCACTGCATTCTAGTTGTGGTTTGTGTC-TAMRA-3' (SEQ ID NO:6). Amplification of the PCR product was achieved with the following cycling parameters: 1 cycle at 50°C for 2 min, 1 cycle at 95°C for 10 min; 40 cycles of 95°C for 15 sec, and 60°C for 60 sec.

Human Genome Array

[0098] Confluent 293 cells (2.4×10^6 cells in a 75 cm² flask) were cultured in DMEM + 10% FCS for approximately 20 hrs and were then infected with one plaque forming unit per cell (PFU/cell) of HSV-1 MacIntyre Strain for 90 min. ATA at a concentration 20 µM (final) was added 90 min post-infection. The infected cells were harvested at 24 h after infection. Total RNA of sufficient quality and quantity was isolated from suspension using RNeasy Plus Mini Kit (QIAGEN, Valencia, CA) according to the manufacturer's instructions. Whole genome expression profiling was performed on Affymetrix Human U133 Plus 2.0 Array by the Asuragen, Inc. 3 µg total RNA was provided as input material.

[0099] Overall, RNA quality assessed on the bioanalyzer had RIN values of >9. Array hybridization and scaling factors were within passing quality control range. After scanning, raw expression CEL files were processed with Affymetrix Expression Console vs.1.1.2 (affymetrix.com). Each CEL file was processed with Robust Multichip Analysis (RMA) and a summarized table was exported as a text file. All downstream genomics and statistical analyses were done with JMP Genomics (version 5) (jmp.com/software/genomics). First, a filtering step was applied to filter-out low expression, specifically, those values below 6 in log² mode in at least 2 out of 8 samples. Out of 54,675 total Affy U133 Plus2 probes, 41,569 (76%) probes remained after the cutoff. Two positive control samples were also seen, human brain and pooled human universal reference RNA, respectively, that hybridized well. For future analysis, these two samples were excluded.

[0100] Principle Component Analysis revealed two distinct populations, samples treated with and without HSV. The first principle component separation (91.7%) is therefore explained by the effect of HSV versus Vehicle. Data was median-normalized across each probe to have a median signal of 0. ANOVA was performed to find significant differential transcripts between HSV versus Vehicle, ATA versus Vehicle, HSV+ATA versus Vehicle, and HSV+ATA versus HSV alone. Multiple testing corrections were not included, and significant genes were considered less than p-value of 0.01.

[0101] A self-organized map was used to examine patterns of expression among Vehicle, HSV1, and HSV1+ATA. It was desired to find specific transcripts that were either up-regulated or down-regulated by the ATA treatment. A distinct cluster of genes that showed some stimulation in the presence of HSV was found, and ATA correction brought these genes back to vehicle baseline. Similarly, a cluster of genes that were repressed by HSV compared to the vehicle was also found, and subsequently activated by treatment with ATA. These were called Clusters A and B, respectively (see, Tables 1 and 2). Analysis was followed-up to interrogate the biological functions of these clusters using GeneGO software (genego.com).

RT² Profiler™ PCR Array (SABiosciences- QIAGEN)

[0102] 6×10^5 293 cells (from AGTC) were infected with wtHSV-1 McIntyre strain at MOI 1 in the presence of DMEM & 10% FBS with or without 50µM ATA and 1-2 hours and diluted to 40% with DMEM & 10% FBS (final ATA concentration of 20µM). Triplicate total RNA samples were harvested 24 hrs later using Qiagen RNeasy mini kit and DNase treated on the column and eluted. Eluate RNA triplicate samples were combined and photospectrometer O.D. readings were performed on the eluate at 260 and 280nm to determine concentration. Sample RNA was converted into template cDNA using SABiosciences RT2 first strand kit. The

cDNA was then used in the Human JAK / STAT Signaling Pathway PCR Array (PAHS-039A).

Example 1

Inhibition of HSV-Induced iNOS Levels in V27 Cells using ATA

[0103] V27 cells in Dulbecco's Modified Eagles Medium (DMEM) (Hyclone, Waltham, MA) were infected with the HSV-1 d27.1 vector at an MOI of 1 and treated with 30 μ M ATA (Sigma, St. Louis, MO), with or without 10% fetal bovine serum (FBS). Control experiments without ATA were also performed. V27 cell lysates were obtained 24 hours post infection (h.p.i.) and Western blots were run on the lysates and iNOS protein was detected by using purified rabbit anti-iNOS/NOS Type II pAb (BD Biosciences, Cat# 610332). Results are shown in Figure 1.

[0104] As shown, HSV infection induced iNOS expression in those cultures that did not include ATA (lanes 4 and 5). iNOS expression was inhibited in the presence of 30 μ M ATA (lanes 2 and 3). The presence or absence of 10% FBS did not have an effect on iNOS expression.

Example 2

Optimization of ATA-HSV Protocol

[0105] In order to test whether ATA could increase the rHSV-1 d27-1 (d27-1) vector yield in V27 cells, ATA was applied to the media during the infection (Step 1) or dilution steps (Step 2) (Figure 2A). The d27-1 infection at MOI=0.15 was performed in 2/5 of the final media volume and the remaining 3/5 of the media was added during dilution step.

[0106] ATA treatment delayed HSV-1 plaque formation or cell lysis in V27 cell monolayers. Cytopathic effect (CPE) at the time of harvest, 72 hours post-infection, (hpi) was between 20-60% as compared to 100% CPE in the absence of ATA. Treatment with ATA during the HSV-1 infection step (ATA at Step 1) showed increased d27-1 titers in V27 cells supernatants harvested at 72 hpi (Figure 4B). The optimal ATA concentration when added during the infection step (ATA I) was 50 μ M and this was further diluted to the final concentration (f.c.) of 20 μ M ATA during the dilution step by adding the remaining 3/5 vol of the media. In this case, ATA increased d27-1 titers in supernatant at the time of harvest (72 h.p.i.) about 10 times: from $4.0 \pm 0.3 \times 10^7$ DRP/ml or $1.4 \pm 0.2 \times 10^7$ PFU/ml to $3.7 \pm 0.2 \times 10^8$ DRP/ml or $1.2 \pm 0.3 \times 10^8$ PFU/ml (Figure 4B).

[0107] In order to determine which concentrations and conditions for ATA addition have impact on HSV yields, the following experiments were conducted. ATA at varying concentrations was added to V27 cultures infected with the HSV-1 d27-1 vectors in either six-well plates (Figure 2B) or T150 flasks (Figure 2C) in two steps as follows. In Step 1 of the protocol, V27 cells were infected with 0.15 MOI of the rHSV vector in 2/5 of the final volume of DMEM with 10% FBS and 0-60 μ M ATA concentrations. Cells were cultured for 1-2 hours at 37°C to complete Step 1. In Step 2, the concentration of ATA was reduced to a range between 12-24 μ M by the addition of 3/5 of final media volume. Cells were cultured for 70-74 hours at 37°C and the supernatant harvested. Viral titers were expressed as Dnase Resistant Particles (DRP/ml) or Plaque Forming Units (pfu/ml) per ml and were determined as described above.

[0108] Results are shown in both Figures 2B and 2C and optimal ATA concentrations are bolded. As can be seen, in both six-well plates (Figure 2B) and T150 flasks (Figure 2C), cultures with ATA added in either of Step 1 or Step 2 had significantly higher HSV titers than those without ATA. Moreover, the highest titers were seen at ATA concentrations of 50 μ M in Step 1, reduced to 20 μ M in Step 2 (Figure 2B), although all concentrations of ATA produced higher viral titers than those cultures that lacked ATA.

[0109] Additional experiments were conducted to determine optimal conditions and the effect of the presence or absence of FBS on HSV titer. The serum presence, specifically, 10% fetal bovine serum (FBS), in ATA-containing media was the most important parameter for the ATA-induced HSV-1 titer yield. A supplementation of ATA into serum-free media during the dilution step (ATA at Step 2) caused reduction of virus titer, where the "DRP" titer was reduced below control level and PFU/ml titer was below detection (Figure 3). An even more dramatic effect in serum-free media was seen when ATA, starting at 3 μ M concentration, was supplemented during the infection Step 1 and both DRP/ml and PFU/ml titers were below the detection limit.

[0110] In this experiment, V27 cells were seeded into six well plates the day prior to infection at 6×10^5 cells/well. HSV 2x infection solution was prepared as follows: HSV-1 d27.1 stock was added to DMEM (no FBS) at 2x titer that the cells would be infected at final MOI 0.15. Several 2xATA solution combinations were prepared containing either 100 μ M or 40 μ M ATA in DMEM and without FBS or with 20% FBS.

[0111] HSV 2x infection solution was mixed equal volumes of either DMEM +/- 20%FBS or desired 2xATA solution +/- 20% FBS, vortexed, and 1ml of infectious inoculum per well was added in Step 1. If wells contained ATA in Step 1, the concentration was 50 μ M and all wells were incubated for 1-2 hrs. In Step2, additional 1.5 ml of DMEM combinations +/- 40 μ M ATA and +/-20% FBS were added. If wells contained ATA whether in Step 1 or Step 2, the final ATA concentration was 20 μ M. The plates were returned to the incubator for 70-74 hours after which time free media was harvested, vortexed, and centrifuged at 1,100 x g for 10 minutes at 4°C. Supernatants were transferred to a new dispensing tube, vortexed, aliquoted and stocks were frozen at -80°C.

Example 3

Importance of Serum Presence in ATA-HSV Protocol

[0112] Additional experiments were conducted to determine optimal conditions and the effect of the presence or absence of FBS on HSV titer. The serum presence, specifically, 10% fetal bovine serum (FBS), in ATA-containing media was the most important parameter for the ATA-induced HSV-1 titer yield. A supplementation of ATA into serum-free media during the dilution step (ATA at Step 2) caused reduction of virus titer, where the "DRP" titer was reduced below control level and PFU/ml titer was below detection (Figure 3). An even more dramatic effect in serum-free media was seen when ATA, starting at 3 μ M concentration, was supplemented during the infection Step 1 (data not shown) and both DRP/ml and PFU/ml titers were below the detection limit.

[0113] In this experiment, V27 cells were seeded into six well plates the day prior to infection at 6×10^5 cells/well. HSV 2x infection solution was prepared as follows: HSV-1 d27.1 stock was added to DMEM (no FBS) at 2x titer that the cells would be infected at final MOI 0.15. Several 2x ATA solution combinations were prepared containing either 100 μ M or 40 μ M ATA in DMEM and without FBS or with 20% FBS. HSV 2x infection solution was mixed equal volumes of either DMEM +/- 20%FBS or desired 2xATA solution +/- 20% FBS, vortexed, and 1ml of infectious inoculum per well was added in Step 1. If wells contained ATA in Step 1, the concentration was 50 μ M and all wells were incubated for 1-2 hrs. In Step 2, additional 1.5 ml of DMEM

combinations \pm 40 μ M ATA and \pm 20% FBS were added. If wells contained ATA whether in Step 1 or Step 2, the final ATA concentration was 20 μ M. The plates were returned to the incubator for 70-74 hours after which time free media was harvested, vortexed, and centrifuged at 1,100 x g for 10 minutes at 4°C. Supernatants were transferred to a new dispensing tube, vortexed, aliquoted and stocks were frozen at -80°C.

Example 4

Effect of ATA on Wild Type HSV Titers in Culture

[0114] Figure 4A shows supernatant titers of both wtHSV-1 strains, KOS and McIntyre in cell lines permissive for wtHSV-1 propagation: 293 cells, HeLa and Vero cells. wtHSV-1 strains KOS and McIntyre were propagated in 293, HeLa and Vero cells using the ATA-HSV Protocol where 50 μ M ATA was added at Step 1 and the final concentration after Step 2 was diluted to 20 μ M ATA. Supernatants containing virus were harvested after three days as described above. ATA increased the yield of both virus types in Vero cells. wtHSV-1 McIntyre strain reached the highest titers after ATA induction in 293 cells, however, ATA appeared to inhibit HSV-1 KOS growth in 293 cells. Finally, ATA also appeared to inhibit both types of HSV-1 viruses in HeLa cells. Two-way ANOVA; Bonferroni test; - ATA vs. + ATA: ***: $p < 0.001$, ns: $p > 0.05$; $n = 4$ independent experiments.

[0115] For statistical calculations, a larger study with a set of ten independent experiments ($n = 10$) was conducted in 6-well plates with or without ATA added during the infection step (Step 1) comparing replication-deficient d27-GFP and also wtHSV-1 McIntyre strain in order to investigate, whether ATA could increase the titer in the wtHSV-1 strain as well (Figure 4B). ATA was added at 50 mM to confluent cell monolayers, V27 cells for d27-1 or 293 cells for wtHSV-1 McIntyre, during the infection step (Step 1) and the cells were infected with vectors at $MOI = 0.15$ while ATA was diluted 1 hr later to a final 20 mM concentration. After ATA addition, d27-1 titer significantly increased 6.1 times from 5.4×10^7 DRP/ml or 1.1×10^8 DRP/ml (** $P < 0.01$) and wtHSV-1 McIntyre significantly increased 9.1 times from 3.3×10^8 DRP/ml or 1.0×10^9 DRP/ml (*** $P < 0.001$) as analyzed by 2-way ANOVA and Bonferroni test (Figure 4B).

Example 5

Effect of ATA in HSV Stocks on the Production of rAAV Virions

[0116] In order to determine whether the presence of ATA during HSV production impacted the yield of rAAV virions produced using HSV vectors, the following experiment was performed showing rAAV titers affected by ATA residue present from rHSV-1 stocks produced in 293 cells (Figure 5A). The rAAV-GFP vector was produced by co-infection of rHSV-rep2/cap2 and rHSV-EGFP vectors in 293 cells 60mm plates using ATA-containing rHSV-1 stocks prepared under different concentrations ATA concentrations (12 μ M or 20 μ M f.c.) (see Material and Methods). rAAV DRP/ml titer was slightly increased by 1.3 times when using the rHSV-rep2/cap2 stock that was prepared with 20 μ M ATA added during infection (ATA Step 1) versus "No ATA" Control (* $p < 0.05$), where the ATA concentration during rAAV production was approximately 3 μ M. No significant rAAV titer increase was detected when using the rHSV-rep2/cap2 stock that was prepared with 12 μ M ATA (Figure 5A). One-way ANOVA; Tukey's-test: * $p < 0.05$; 20 μ M ATA vs. No ATA; $n = 4$ independent experiments.

[0117] In another experiment, ATA was shown to increase rAAV titer when 10 μ M ATA was spiked directly into

293 cells media during 2 hrs of rHSV-rep2/cap2 and rHSV-EGFP co-infection step. This effect was not observed when the ATA concentrations were higher than 20 μ M (Figure 5B).

Example 6

ATA Mechanism of Action

[0118] In order to illuminate the ATA mechanism of action by human genome array and the fact that Vero and V27 are monkey-derived cell lines, wtHSV-1 propagation was tested in several human cell lines (Figures 4A-4B). A propagation of two wild type HSV-1 (wtHSV-1) strains, KOS and McIntyre, were compared in three cell lines: human embryonic kidney 293 (293 cells), human cervical cancer HeLa cells and African green monkey kidney epithelial Vero cells. The experiments were carried out in 10%FBS-containing media following the ATA I protocol scheme, where 50 μ M ATA was supplemented during the infection step and further diluted to 20 μ M ATA concentration (see Material and Methods). In human-derived 293 cells, only the wtHSV-1 McIntyre virus titer increased significantly by ATA, from 1.4×10^8 DRP/ml to 1.2×10^9 DRP/ml ($***p < 0.001$). In Vero cells, ATA significantly increased titers only in wtHSV-1 KOS strain from 1.7×10^8 DRP/ml up to 9.1×10^8 DRP/ml ($***p < 0.001$; both statistics: Two-way ANOVA, Bonferroni test; $n=4$). Surprisingly, in HeLa cells, ATA seemed to inhibit propagation of both KOS and McIntyre HSV-1 strains.

[0119] To identify cell response gene signatures to wtHSV-1 McIntyre infection, transcriptional profiling was done using Affymetrix Human U133 Plus 2.0 Array with 293 cells that were either mock-treated, ATA-treated (ATA), wtHSV-1 McIntyre-infected (HSV) or ATA-treated and wtHSV-1 McIntyre-infected (HSV&ATA) for 24 h. Gene expression changes induced by infection or ATA treatment were identified by referencing the gene expression level for each probe to that corresponding in the uninfected cells sample. HSV-1 infection alone had strong impact on the gene expression profile of 293 cells when compared to untreated cells gene profile.

[0120] First, a filtering step was applied to filter out low expression and out of 54,675 total Affy U133 Plus2 probes, 41,569 (76%) probes remained after the cutoff. Principle Component Analysis revealed two distinct populations, samples treated with and without HSV. ANOVA was performed to find significant differential transcripts between HSV versus Vehicle, ATA versus Vehicle, HSV+ATA versus Vehicle, and HSV+ATA versus HSV alone. Significant genes were considered less than a p-value of 0.01. A self-organized map was used to examine patterns of expression among Vehicle, HSV1, and HSV1+ATA. A distinct cluster of genes was found, Cluster A and B (Tables 1 and 2), that showed changes in transcriptional profiles caused by HSV-1 and their correction by ATA that brought these genes back to vehicle baseline.

[0121] The biological functions of these clusters were analyzed in GeneGo. Cluster A (Table 2) represented 58 probes that showed up-regulation in HSV-1 and suppression after the addition of ATA. These genes were primarily involved in inflammatory IgE and IFN signaling, and general immune response. Cluster B (Table 1) represented 152 probes that showed down-regulation by HSV-1 and their subsequent ATA up-regulation by bringing them back to vehicle baseline. Genes in this cluster were primarily involved in cell cycle G1/S, signal transduction in PTEN, and WNT development. For example, ATA in the presence of HSV-1 up-regulated CDC25A, CDKN1A, CDKN1C, CCNK, CNM2 genes from cell cycle progression pathways and genes from Ras/Raf/MEK pathway, such as FOXC1, FOXD3, FOXO3 (See, Table 1).

[0122] Tables 3A and 3B show nNOS, iNOS and eNOS expression reduction in HSV ATA samples as analyzed by Affymetrix Gen Array and iNOS expression reduction in HSV + ATA samples as analyzed by Qiagen SAB Jak-Stat RT-PCR Microarray

Example 7**Effect of Dexamethasone on HSV Yields**

[0123] In order to determine whether the iNOS inhibitor dexamethasone also increased rHSV yields in culture, the following experiment was conducted (Figure 6). shows the effect of dexamethasone (Dex) on d27-1/GFP HSV-1 viral titer. The final titers d27-1/GFP HSV-1 were, in general, slightly elevated after dex pretreatments or treatments as compared to untreated control.

[0124] V27 cells were seeded into six well plates the day prior to infection at 6×10^5 cells/well and the next day, different concentrations of dexamethasone (dex) were added to the wells either before HSV-1 infection (pretreat) or during the infection (treat) in 2/5 vol of DMEM-10% FBS media. After 1-2 hours of incubation at 37°C, 3/5 vol. of DMEM-10% FBS was added. The cultures were returned to the incubator and supernatant was harvested after 70-74 hours. The data are shown as Mean value of titers \pm S.D. in pfu/ml (n=2).

Example 8**Effect of Valproic Acid on HSV Yields**

[0125] In order to determine whether the iNOS inhibitor valproic acid also increased rHSV yields in culture, the following experiment was conducted. Figure 7 shows the effect of pretreatment by valproic acid (VA) on d27-1/GFP HSV-1 viral titer. VA at concentration 5mM slightly elevated the titer of d27-1/GFP HSV-1, however the concentrations below and above of 5 mM appear to have an inhibitory effect on d27-1/GFP HSV-1 titer, as compared to untreated control.

[0126] V27 cells were seeded into six well plates the day prior to infection at 6×10^5 cells/well and the next day, different concentrations of VA were added to wells. Plates were incubated for 6 hrs, aspirated and 2/5 vol of media containing infectious HSV-1 d27-1 stock was added and incubated for 1-2 hours at 37°C, 5 after which time 3/5 vol of DMEM 10% FBS media was added. The cultures were returned to the incubator and supernatant was harvested after 70-74 hours.

[0127] As shown in the foregoing examples, ATA in micromolar concentrations increases HSV-1 vector yield. This finding is important for both large-scale HSV production, as well as rHSV and rAAV vector production. Moreover and surprisingly, the presence of ATA in rHSV-1 stocks did not negatively influence rAAV yield. This result is surprising as ATA in millimolar amounts and higher concentrations is known to be an antiviral agent (Cushman et al., J. Med. Chem. (1991) 34:329-337; Zhang et al., Antiviral Res. (1999) 43:23-35; Yap et al., Computational Biol. and Chem. (2005) 29:212-219; De Clercq, Advents, Advances, and Adventures Med. Res. Rev. (2011) 31:118-160). In serum-free media the inventors have also observed a possible antiviral effect of ATA at micromolar concentrations, but not in the presence of serum (10% FBS).

[0128] Also as shown herein, ATA treatment delayed HSV-1 plaque formation and cell lysis in V27 cells monolayers and cytopathic effect (CPE), which evidences antiapoptotic properties. The mechanism of action of ATA in the increase of HSV-1 yield appears to involve changes in factors required for HSV-1 production and reduction of cellular innate antiviral immune response responsible for a viral clearance. From the Human Genome Array analysis above, it was discovered that HSV-1 infection alone has strong impact on the gene

expression profile of 293 cells and the effect of ATA was mostly observed in cells treated with both HSV-1 and ATA. Genes involved in cell cycle G1/S, signal transduction in PTEN, and WNT development were significantly down-regulated by HSV-1 and up-regulated after addition of ATA.

[0129] Genes primarily involved in inflammatory IgE and IFN signaling, and general immune response were up-regulated by HSV-1 and suppressed after the addition of ATA. In the presence of HSV-1, ATA up-regulated CDC25A, CDKN1A, CDKN1C, CCNK, CNNM2 genes from the cell cycle progression pathways and genes from the Ras/Raf/MEK pathway, including FOXC1, FOXD3, FOXO3.

[0130] The finding that ATA can increase HSV-1 yield are important because of the need of higher yields of HSV-1 vectors for large-scale production for gene therapy and other applications.

[0131] Thus, methods for increasing viral yields using iNOS inhibitors are described. Although preferred embodiments of the subject invention have been described in some detail, it is understood that obvious variations can be made without departing from the scope of the invention as defined herein.

Table 1. Cluster B gene list: Genes suppressed by HSV and up-regulated by HSV+ATA

Probe_Set_ID	Gene Symbol	Gene_Title	Estimate of HSV+ATA vs HSV	p-Value for Estimate of HSV+ATA vs HSV
230304_at	---	---	2.1691	0.0001
225806_at	JUB	jub, ajuba homolog (Xenopus laevis)	2.126171	0.0004
202935_s_at	SOX9	SRY (sex determining region Y)-box 9	1.855105	0.000046
204790_at	SMAD7	SMAD family member 7	1.797951	0.0029
214633_at	SOX3	SRY (sex determining region Y)-box 3	1.797254	0.000007
210512_s_at	VEGFA	vascular endothelial growth factor A	1.754719	0.000021
216652_s_at	DR1	down-regulator of transcription 1, TBP-binding (negative cofactor 2)	1.74529	0.0036
205932_s_at	MSX1	msh homeobox 1	1.726634	6.90E-06
209348_s_at	MAF	v-maf musculoaponeurotic fibrosarcoma oncogene homolog (avian)	1.685705	0.0073
217028_at	CXCR4	chemokine (C-X-C motif) receptor 4	1.652962	0.0093
218251_at	MID1IP1	MID1 interacting protein 1 (gastrulation specific G12 homolog (zebrafish))	1.642017	0.0064
1553764_a_at	JUB	jub, ajuba homolog (Xenopus laevis)	1.623019	0.0017
214446_at	ELL2	elongation factor, RNA polymerase II, 2	1.602283	0.0001
227718_at	PURB	purine-rich element binding protein B	1.59314	0.0019
219624_at	BAG4	BCL2-associated athanogene 4	1.578527	0.0007
225642_at	KTI12	KTI12 homolog, chromatin associated (S. cerevisiae)	1.566082	0.0002
203705_s_at	FZD7	frizzled homolog 7 (Drosophila)	1.54382	0.0002
222696_at	AXIN2	axin 2	1.526076	0.0066
202007_at	NID1	nidogen 1	1.521713	0.0029

Probe_Set_ID	Gene Symbol	Gene_Title	Estimate of HSV+ATA vs HSV	p-Value for Estimate of HSV+ATA vs HSV
226858_at	CSNK1E	casein kinase 1, epsilon	1.51752	0.0002
57739_at	DND1	dead end homolog 1 (zebrafish)	1.515518	0.0005
1558290_a_at	PVT1	Pvt1 oncogene (non-protein coding)	1.515288	0.0016
215694_at	SPATA5L1	spermatogenesis associated 5-like 1	1.51283	0.0012
206302_s_at	NUDT4 /// NUDT4P1	nudix (nucleoside diphosphate linked moiety X)-type motif 4 /// nudix (nucleoside diphosphate linked moiety X)-type motif 4 pseudogene 1	1.49443	0.0073
218486_at	KLF11	Kruppel-like factor 11	1.493386	0.0006
201695_s_at	NP	nucleoside phosphorylase	1.492386	0.0004
203002_at	AMOTL2	angiomin like 2	1.489062	0.0093
227195_at	ZNF503	zinc finger protein 503	1.478008	0.0094
238012_at	DPP7	Dipeptidyl-peptidase 7	1.471194	0.0003
209905_at	HOXA9	homeobox A9	1.46661	0.0003
209188_x_at	DR1	down-regulator of transcription 1, TBP-binding (negative cofactor 2)	1.4541	0.0001
221168_at	PRDM13	PR domain containing 13	1.448075	0.0043
213338_at	TMEM158	transmembrane protein 158	1.436626	0.0001
209098_s_at	JAG1	jagged 1 (Alagille syndrome)	1.433177	0.0001
206300_s_at	PTH1H	parathyroid hormone-like hormone	1.430301	0.0004
220018_at	CBLL1	Cas-Br-M (murine) ecotropic retroviral transforming sequence-like 1	1.427214	0.00003
226284_at	ZBTB2	zinc finger and BTB domain containing 2	1.421748	6.50E-06
213348_at	CDKN1C	cyclin-dependent kinase inhibitor 1C (p57, Kip2)	1.413731	0.0001
209565_at	RNF113A	ring finger protein 113A	1.413412	7.80E-06
239696_at	---	---	1.413287	0.0006
212696_s_at	RNF4	ring finger protein 4	1.408882	0.0034
214651_s_at	HOXA9	homeobox A9	1.404355	0.000042
217741_s_at	ZFAND5	zinc finger, AN1-type domain 5	1.401465	0.0003
229309_at	ADRB1	adrenergic, beta-1-, receptor	1.398788	0.0008
228953_at	WHAMM	WAS protein homolog associated with actin, golgi membranes and microtubules	1.392654	0.0013
1554522_at	CNNM2	cyclin M2	1.389108	0.0049
204913_s_at	SOX11	SRY (sex determining region Y)-box 11	1.385923	0.0004
231901_at	C19orf52	chromosome 19 open reading frame 52	1.378798	0.0034
224314_s_at	EGLN1	egl nine homolog 1 (C. elegans)	1.375757	0.0001

Probe_Set_ID	Gene Symbol	Gene_Title	Estimate of HSV+ATA vs HSV	p-Value for Estimate of HSV+ATA vs HSV
209099_x_	JAG1	jagged 1 (Alagille syndrome)	1.374203	0.0029
205555_s_at	MSX2	msh homeobox 2	1.368114	0.00003
216035_x_at	TCF7L2	transcription factor 7-like 2 (T-cell specific, HMG-box)	1.368112	0.0003
225824_at	CCNK	cyclin K	1.363977	0.0076
209201_x_at	CXCR4	chemokine (C-X-C motif) receptor 4	1.360311	0.0001
235147_at	FU32063	Hypothetical LOC150538	1.357503	0.0008
204527_at	MYO5A	myosin VA (heavy chain 12, myoxin)	1.356721	0.0017
242963_at	SGMS2	sphingomyelin synthase 2	1.356199	0.0008
218247_s_at	MEX3C	mex-3 homolog C (C. elegans)	1.3496	0.0001
207654_x_at	DR1	down-regulator of transcription 1, TBP-binding (negative cofactor 2)	1.348295	9.30E-06
209357_at	CITED2	Cbp/p300-interacting transactivator, with Glu/Asp-rich carboxy-terminal domain, 2	1.347165	0.0002
202241_at	TRIB1	tribbles homolog 1 (Drosophila)	1.340674	0.0009
209372_x_at	TUBB2A /// TUBB2B	tubulin, beta 2A /// tubulin, beta 2B	1.339095	0.0001
205541_s_at	GSPT2	G1 to S phase transition 2	1.33517	0.005
1568815_a_at	DDX50	DEAD (Asp-Glu-Ala-Asp) box polypeptide 50	1.329483	0.0043
204805_s_at	H1FX	H1 histone family, member X	1.324347	0.0013
213152_s_at	SFRS2B	splicing factor, arginine/serine-rich 2B	1.322967	6.70E-06
225699_at	C7orf40	chromosome 7 open reading frame 40	1.321373	0.0014
218295_s_at	NUP50	nucleoporin 50kDa	1.318229	0.0014
224739_at	PIM3	pim-3 oncogene	1.31654	0.0001
214911_s_at	BRD2	bromodomain containing 2	1.313049	0.0015
208938_at	PRCC	papillary renal cell carcinoma (translocation-associated)	1.304724	0.000041
206907_at	TNFSF9	tumor necrosis factor (ligand) superfamily, member 9	1.295621	0.0032
202284_s_at	CDKN1A	cyclin-dependent kinase inhibitor 1A (p21, Cip1)	1.28534	0.0001
206915_at	NKX2-2	NK2 homeobox 2	1.280773	0.000021
215087_at	C15orf39	chromosome 15 open reading frame 39	1.277596	0.0013
202219_at	SLC6A8	solute carrier family 6 (neurotransmitter transporter, creatine), member 8	1.273131	0.0006
202704_at	TOB1	transducer of ERBB2, 1	1.272848	0.0016
213038_at	RNF19B	ring finger protein 19B	1.27284	0.0022

Probe_Set_ID	Gene Symbol	Gene_Title	Estimate of HSV+ATA vs HSV	p-Value for Estimate of HSV+ATA vs HSV
213150_at	HOXA10	homeobox A10	1.272477	0.008
204383_at	DGCR14	DiGeorge syndrome critical region gene 14	1.268562	0.000043
1553613_s_at	FOXC1	forkhead box C1	1.26852	0.0003
218398_at	MRPS30	mitochondrial ribosomal protein S30	1.266043	0.0057
202166_s_at	PPP1R2	protein phosphatase 1, regulatory (inhibitor) subunit 2	1.264115	0.00003
235004_at	RBM24	RNA binding motif protein 24	1.260011	0.0086
223742_at	MRPL4	mitochondrial ribosomal protein L4	1.251284	0.000049
211756_at	PTH1H	parathyroid hormone-like hormone	1.24883	0.0016
209211_at	KLF5	Kruppel-like factor 5 (intestinal)	1.241821	0.0003
225796_at	PXK	PX domain containing serine/threonine kinase	1.239504	0.0002
225434_at	DEDD2	death effector domain containing 2	1.23754	0.000044
208686_s_at	BRD2	bromodomain containing 2	1.229145	0.0001
217821_s_at	WBP11	WW domain binding protein 11	1.218077	0.0002
244519_at	ASXL1	additional sex combs like 1 (Drosophila)	1.208056	0.000044
208415_x_at	ING1	inhibitor of growth family, member 1	1.184054	0.0017
216511_s_at	TCF7L2	transcription factor 7-like 2 (T-cell specific, HMG-box)	1.174677	0.0024
235959_at	---	---	1.172362	0.0008
243707_at	---	---	1.171741	0.0005
227852_at	RP9	retinitis pigmentosa 9 (autosomal dominant)	1.171328	0.0002
1555772_a_at	CDC25A	cell division cycle 25 homolog A (S. pombe)	1.170244	0.0051
221841_s_at	KLF4	Kruppel-like factor 4 (gut)	1.16333	0.0006
214789_x_at	SFRS2B	splicing factor, arginine/serine-rich 2B	1.162135	0.0001
216997_x_at	TLE4	transducin-like enhancer of split 4 (E(sp1) homolog, Drosophila)	1.159916	0.0001
205780_at	BIK	BCL2-interacting killer (apoptosis-inducing)	1.158085	0.0001
212075_s_at	CSNK2A1	casein kinase 2, alpha 1 polypeptide	1.156022	0.0049
228820_at	XPNPEP3	X-prolyl aminopeptidase (aminopeptidase P) 3, putative	1.152448	0.0008
209653_at	KPNA4	karyopherin alpha 4 (importin alpha 3)	1.146903	0.0004
209457_at	DUSP5	dual specificity phosphatase 5	1.144663	0.0016
224671_at	MRPL10	mitochondrial ribosomal protein L10	1.144585	0.0061
200618_at	LASP1	LIM and SH3 protein 1	1.140405	0.0097

Probe_Set_ID	Gene Symbol	Gene_Title	Estimate of HSV+ATA vs HSV	p-Value for Estimate of HSV+ATA vs HSV
228931_at	COQ4	coenzyme Q4 homolog (S. cerevisiae)	1.136018	0.0008
224583_at	COTL1	coactosin-like 1 (Dictyostelium)	1.128103	0.001
238738_at	PSMD7	Proteasome (prosome, macropain) 26S subunit, non-ATPase, 7	1.128031	0.0062
202501_at	MAPRE2	microtubule-associated protein, RP/EB family, member 2	1.12791	0.0046
214321_at	NOV	nephroblastoma overexpressed gene	1.123481	0.0004
202936_s_at	SOX9	SRY (sex determining region Y)-box 9	1.12181	0.0054
222163_s_at	SPATA5L1	spermatogenesis associated 5-like 1	1.116794	0.0013
241612_at	FOXD3	forkhead box D3	1.11263	0.0041
202431_s_at	MYC	v-myc myelocytomatosis viral oncogene homolog (avian)	1.105333	0.0007
215933_s_at	HHEX	hematopoietically expressed homeobox	1.105321	0.0002
204132_s_at	FOXO3 /// FOXO3B	forkhead box O3 /// forkhead box O3B pseudogene	1.100606	0.0007
201041_s_at	DUSP1	dual specificity phosphatase 1	1.09043	0.0004
230233_at	---	---	1.087609	0.0005
215223_s_at	SOD2	superoxide dismutase 2, mitochondrial	1.087493	0.0001
225689_at	C3orf39	chromosome 3 open reading frame 39	1.080337	0.003
223132_s_at	TRIM8	tripartite motif-containing 8	1.073976	0.0065
203313_s_at	TGIF1	TGFB-induced factor homeobox 1	1.073363	0.0008
201461_s_at	MAPKAPK2	mitogen-activated protein kinase-activated protein kinase 2	1.070375	0.0088
236174_at	---	---	1.069052	0.0013
223679_at	CTNNB1	catenin (cadherin-associated protein), beta 1, 88kDa	1.060314	0.0044
204039_at	CEBPA	CCAAT/enhancer binding protein (C/EBP), alpha	1.052213	0.01
234302_s_at	ALKBH5	alkB, alkylation repair homolog 5 (E. coli)	1.049599	0.0043
223290_at	PDXP	pyridoxal (pyridoxine, vitamin B6) phosphatase	1.048187	0.0022
226644_at	MIB2	mindbomb homolog 2 (Drosophila)	1.041356	0.0011
206363_at	MAF	v-maf musculoaponeurotic fibrosarcoma oncogene homolog (avian)	1.037386	0.0008
1555639_a_at	RBM14	RNA binding motif protein 14	1.035626	0.0003

Probe_Set_ID	Gene Symbol	Gene_Title	Estimate of HSV+ATA vs HSV	p-Value for Estimate of HSV+ATA vs HSV
222527_s_at	RBM22	RNA binding motif protein 22	1.027056	0.0008
213419_at	APBB2	amyloid beta (A4) precursor protein-binding, family B, member 2	1.025879	0.0009
213360_s_at	POM121 /// POM121C	POM121 membrane glycoprotein (rat) /// POM121 membrane glycoprotein C	1.023715	0.0013
225832_s_at	DAGLB	diacylglycerol lipase, beta	1.001586	0.0001
212445_s_at	NEDD4L	neural precursor cell expressed, developmentally downregulated 4-like	0.997235	0.0022
224562_at	WASF2	WAS protein family, member 2	0.996687	0.0043
223389_s_at	ZNF581	zinc finger protein 581	0.99618	0.0004
231721_at	JAM3	junctional adhesion molecule 3	0.989941	0.0001
203140_at	BCL6	B-cell CLL/lymphoma 6	0.984695	0.0004
213823_at	HOXA11	homeobox A11	0.974615	0.0036
1552275_s_at	PXK	PX domain containing serine/threonine kinase	0.973552	0.0002
52731_at	AMBRA1	autophagy/beclin-1 regulator 1	0.970744	0.000027
238624_at	---	---	0.970064	0.0005
210479_s_at	RORA	RAR-related orphan receptor A	0.965295	0.0006
220941_s_at	C21orf91	chromosome 21 open reading frame 91	0.943651	0.0013
203234_at	UPP1	uridine phosphorylase 1	0.932337	0.0009
202102_s_at	BRD4	bromodomain containing 4	0.930432	0.0034
217775_s_at	RDH11	retinol dehydrogenase 11 (all-trans/9-cis/11-cis)	0.92863	0.0029

Table 2. Cluster A gene list: Genes up-regulated by HSV and downregulated by HSV+ATA

Probe_Set_ID	Gene Symbol	Gene_Title	Estimate of HSVI_ATA_vs_HSV1	p-Value for Estimate of HSVI_ATA_vs_HSV1
243160_at	---	---	-1.01645	0.0022
205247_at	NOTCH4	Notch homolog 4 (Drosophila)	-0.93759	0.0007
219256_s_at	SH3TC1	SH3 domain and tetratricopeptide repeats 1	-0.91997	0.0021
221631_at	CACNA1I	calcium channel, voltage-dependent, T type, alpha 1I subunit	-0.90394	0.0018
1569961_at	---	---	-0.89195	0.0013
220277_at	CXXC4	CXXC finger 4	-0.85343	0.0054
229611_at	LMLN	leishmanolysin-like (metallopeptidase M8 family)	-0.85329	0.0005

Probe_Set_ID	Gene Symbol	Gene_Title	Estimate of HSVI_ATA_vs_HS V1	p-Value for Estimate of HSVI_ATA_vs_HS V1
232341_x_at	HABP4	hyaluronan binding protein 4	-0.84077	0.0071
242219_at	---	---	-0.83521	0.0028
233767_at	HHLA1	HERV-H LTR-associating 1	-0.8339	0.0017
236925_at	LOC728288	hypothetical LOC728288	-0.81812	0.008
231905_at	C20orf96	chromosome 20 open reading frame 96	-0.80575	0.0041
233313_at	---	---	-0.80466	0.0011
1563507_at	---	---	-0.79441	0.0043
232456_at	C10orf71	chromosome 10 open reading frame 71	-0.78722	0.0067
203468_at	CDK10	cyclin-dependent kinase 10	-0.77888	0.0035
237456_at	---	---	-0.77558	0.0085
216127_at	PDIA2	protein disulfide isomerase family A, member 2	-0.76924	0.0049
207977_s_at	DPT	dermatopontin	-0.76077	0.0065
224510_s_at	CLPB	ClpB caseinolytic peptidase B homolog (E. coli)	-0.75625	0.006
208267_at	TRPV5	transient receptor potential cation channel, subfamily V, member 5	-0.73228	0.0091
221868_at	PAIP2B	poly(A) binding protein interacting protein 2B	-0.7284	1.10E-06
1559439_s_at	C21orf58	chromosome 21 open reading frame 58	-0.7256	0.0051
1552960_at	LRRC15	leucine rich repeat containing 15	-0.72214	0.0049
204413_at	TRAF2	TNF receptor-associated factor 2	-0.69678	0.0039
213888_s_at	TRAF3IP3	TRAF3 interacting protein 3	-0.69063	0.0043
1553945_at	GPHB5	glycoprotein hormone beta 5	-0.67838	0.0003
221312_at	GLP2R	glucagon-like peptide 2 receptor	-0.67682	0.0097
1554783_s_at	ARHGEF2	Rho/Rac guanine nucleotide exchange factor (GEF) 2	-0.67352	0.0005
205567_at	CHST1	carbohydrate (keratan sulfate Gal-6) sulfotransferase 1	-0.67335	0.0009
232473_at	PRPF18	PRP18 pre-mRNA	-0.66869	0.0064

Probe_Set_ID	Gene Symbol	Gene_Title	Estimate of HSVI_ATA_vs_HS V1	p-Value for Estimate of HSVI_ATA_vs_HS V1
		processing factor 18 homolog (S. cerevisiae)		
1565583_at	LOC100291336	hypothetical protein LOC100291336	-0.66137	0.0086
243638_at	---	---	-0.65654	0.0041
208608_s_at	SNTB1	syntrophin, beta 1 (dystrophin-associated protein A1, 59kDa, basic component 1)	-0.65372	0.005
1556202_at	SRGAP2	SLIT-ROBO Rho GTPase activating protein 2	-0.64525	0.0034
206266_at	GPLD1	glycosylphosphatidylinositol specific phospholipase D1	-0.64321	0.0015
234659_at	---	---	-0.63414	0.007
204708_at	MAPK4	mitogen-activated protein kinase 4	-0.633	0.0045
236604_at	BAHCC1	BAH domain and coiled- coil containing 1	-0.63262	0.0083
242129_at	SIN3B	SIN3 homolog B, transcription regulator (yeast)	-0.63025	0.0018
233029_at	OBSCN	obscurin, cytoskeletal calmodulin and titin- interacting RhoGEF	-0.62959	0.0049
203398_s_at	GALNT3	UDP-N-acetyl-alpha-D- galactosamine:polypeptide N- acetylgalactosaminyltransf- erase 3 (GalNAc-T3)	-0.62709	0.0016
242701_at	TBRG1	transforming growth factor beta regulator 1	-0.62074	0.0032
235083_at	LOC151009	hypothetical LOC151009	-0.61926	0.004
206372_at	MYF6	myogenic factor 6 (herculin)	-0.6146	0.0018
1567611_at	---	---	-0.61369	0.0061
236252_at	---	---	-0.61329	0.0041
206286_s_at	TDGF1 /// TDGF3	teratocarcinoma-derived growth factor 1 /// teratocarcinoma-derived growth factor 3, pseudogene	-0.61123	0.0012
1563263_at	PLCG2	Phospholipase C, gamma 2 (phosphatidylinositol- specific)	-0.60753	0.0003
214742_at	AZI1	5-azacytidine induced 1	-0.60464	0.0089

Probe_Set_ID	Gene Symbol	Gene_Title	Estimate of HSVI_ATA_vs_HS V1	p-Value for Estimate of HSVI_ATA_vs_HS V1
244656_at	RASL10B	RAS-like, family 10, member B	-0.6032	0.0072
1555665_at	---	---	-0.59953	0.0088
224291_at	CACNG6	calcium channel, voltage-dependent, gamma subunit 6	-0.59548	0.0068
235616_at	TSHZ2	teashirt zinc finger homeobox 2	-0.59219	0.0037
237461_at	NLRP7	NLR family, pyrin domain containing 7	-0.58969	0.0068
223693_s_at	RADIL	Ras association and DIL domains	-0.58969	0.0052
208454_s_at	PGCP	plasma glutamate carboxypeptidase	-0.58916	0.001
216426_at	TCEB1	transcription elongation factor B (Sill), polypeptide 1 (15kDa, elongin C)	-0.58852	0.002

Table 3A. nNOS, iNOS and eNOS expression reduction in HSV+ATA Samples analyzed by Affymetrix Gen Array; NOS1=nNOS, NOS2=iNOS, NOS3=eNOS

Probe_Set_ID	Pub_ID	Symbol	ATA	HSV	HSV + ATA	p-Val ATA	p-Val HSV	p-Val HSV+ATA
240911_at	AI733341	NOS1	1.02	1.71	1.5	0.872	0.009	0.024
207310_s_at	U31466	NOS1	1	2.79	1.67	0.981	0.003	0.033
AF049656	---	NOS2	-1.03	2.05	1.72	0.706	0.002	0.007
205581_s_at	NM_000603	NOS3	1.24	2.45	1.94	0.234	0.005	0.013

Table 3B. iNOS expression reduction in HSV+ATA Samples analyzed by SABiosciences (QIAGEN) Jak-Stat RT-PCR Microarray

RT2 Catalog	Pub_ID	Symbol	ATA	HSV	HSV + ATA	p-Val ATA	p-Val HSV	p-Val HSV+ATA
240911_at	AI733341	NOS1	1.02	1.71	1.5	0.872	0.009	0.024

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- [US7091029B](#) [0061] [0068] [0069] [0075]

- [US20080292595A \[0066\] \[0071\]](#)
- [US5139941A \[0072\] \[0085\]](#)
- [WO0212455A \[0084\]](#)
- [US5173414A \[0085\]](#)
- [WO9201070A \[0085\]](#)
- [WO9303769A \[0085\]](#)

Non-patent literature cited in the description

- **SAINZHALFORDJ.** Virol., 2002, vol. 76, 11541-11550 [\[0003\]](#)
- **HALLER et al.** Virology, 2006, vol. 344, 119-130 [\[0003\]](#)
- **PALUDAN et al.** Nat. Rev. Immunol., 2011, vol. 11, 143-154 [\[0003\]](#)
- **BRANDNERMUELLER** Hoppe-Seyler's Zeitschrift für physiologische Chemie, 1973, vol. 354, 1176- [\[0003\]](#)
- **DE VRIES et al.** Gene Ther., 2008, vol. 15, 545-552 [\[0003\]](#)
- **BI et al.** J. Virol., 1995, vol. 69, 6466-6472 [\[0004\] \[0010\] \[0051\]](#)
- **HARRIS et al.** J. Virol., 1995, vol. 69, 910-915 [\[0004\] \[0010\] \[0051\]](#)
- **LIN et al.** J. Virol., 1997, vol. 71, 5227-5235 [\[0004\] \[0010\]](#)
- **PERTILE et al.** Avian Dis., 1996, vol. 40, 342-348 [\[0004\] \[0010\] \[0051\]](#)
- **CROEN K.D.J.** Clin. Invest., 1993, vol. 91, 2446-2452 [\[0004\] \[0010\] \[0051\]](#)
- **KARUPIAH et al.** Science, 1993, vol. 261, 1445-1448 [\[0004\] \[0010\] \[0011\] \[0051\]](#)
- **ROLPH et al.** Virol., 1996, vol. 217, 470-477 [\[0004\] \[0010\] \[0051\]](#)
- **AMARO et al.** J. Med. Virol., 1997, vol. 51, 326-331 [\[0004\] \[0010\] \[0051\]](#)
- **LANE et al.** J. Virol., 1997, vol. 71, 2202-2210 [\[0004\] \[0010\] \[0051\]](#)
- **MOSSMAN et al.** J. Virol., 2001, vol. 75, 750-758 [\[0004\]](#)
- **BULLER, R.M.L.J.** Virol., 1981, vol. 40, 241-247 [\[0005\]](#)
- **MISHRA et al.** Virology, 1990, vol. 179, 632-639 [\[0005\]](#)
- **WEINDLER et al.** J. Virol., 1991, vol. 65, 2476-2483 [\[0005\] \[0061\] \[0061\]](#)
- **MUZYCZKA, N.** Curr. Top. Microbiol. Immunol., 1992, vol. 158, 97-129 [\[0006\]](#)
- **CONWAY et al.** Gene Ther., 1999, vol. 6, 986-993 [\[0007\] \[0007\] \[0068\] \[0069\] \[0075\]](#)
- **RICEJ.** Virol., 1989, vol. 63, 83399-407 [\[0007\] \[0008\]](#)
- **WECHUCKET** Biotechnol. Prog., 2000, vol. 16, 493-496 [\[0009\]](#)
- **OZUER et al.** Biotechnol. Prog., 2002, vol. 18, 476-482 [\[0009\]](#)
- **ERLANDSSON et al.** J. Endocrinol., 2002, vol. 175, 165-176 [\[0009\] \[0009\]](#)
- **OTSUKI et al.** Mol. Ther., 2008, vol. 16, 1546-1555 [\[0009\] \[0009\]](#)
- **SOUTHAN et al.** Biochem. Pharmacol., 1996, vol. 51, 383-394 [\[0011\]](#)
- **CHEN et al.** British Journal of Pharmacology, 2002, vol. 137, 71011-20 [\[0011\]](#)
- **TSCHERNEPESTKA** Antimicrob. Agents Chemother, 1975, vol. 8, 479-487 [\[0011\]](#)
- **MIKELENS et al.** Biochemical Pharmacology, 1976, vol. 25, 821-827 [\[0011\]](#)
- **VOLLGRAF et al.** J. Neurochem., 1999, vol. 73, 2501-2509 [\[0011\]](#)
- **TSI et al.** Mol. Pharmacol., 2002, vol. 101, 90-101 [\[0012\]](#)
- **CHEN et al.** British J. Pharmacol., 2002, vol. 137, 1011-1020 [\[0012\]](#)
- **BEERY et al.** Endocrinology, 2001, vol. 142, 3098-3107 [\[0012\]](#)
- **CHEN et al.** J. Biol. Chem., 2001, vol. 276, 46722-46728 [\[0012\]](#)
- **NATHAN, C.** FASAB J., 1992, vol. 6, 3051- [\[0012\]](#)
- **WERNER-FELMAYER et al.** J. Exp. Med., 1990, vol. 172, 1599- [\[0012\]](#)
- **CUSHMAN et al.** J. Med. Chem., 1991, vol. 34, 329-337 [\[0013\]](#)

- **ZHANG et al.**Antiviral Res., 1999, vol. 43, 23-35 [0013] [0050] [0127]
- **YAP et al.**Computational Biol. and Chem., 2005, vol. 29, 212-219 [0013] [0050] [0127]
- **DE CLERCQ**Advents, Advances, and Adventures Med. Res. Rev., 2011, vol. 31, 118-160 [0013] [0050] [0127]
- **HE**Biochem. Biophys. Res. Comm., 2004, vol. 320, 1199-1203 [0013]
- **MYSKIW et al.**J. Virol., 2007, vol. 81, 3027-3032 [0013]
- **Fundamental Virology**vol. I & II, [0018]
- **Handbook of Experimental Immunology**Blackwell Scientific Publicationsvol. I-IV, [0018]
- **T.E. CREIGHTON**Proteins: Structures and Molecular PropertiesW.H. Freeman and Company19930000 [0018]
- **A.L. LEHNINGER**BiochemistryWorth Publishers, Inc. [0018]
- **SAMBROOK et al.**Molecular Cloning: A Laboratory Manual19890000 [0018]
- **Methods In Enzymology**Academic Press, Inc. [0018]
- **National Biomedical Research Foundation****DAYHOFF, M.O.**Atlas of Protein Sequence and Structurevol. 5, 353-358 [0027]
- **SMITHWATERMAN**Advances in Appl. Math., 1981, vol. 2, 482-489 [0027]
- **CLARK et al.**Hum. Gene Ther., 1999, vol. 10, 1031-1039 [0036]
- **VELDWIJK et al.**Mol. Ther., 2002, vol. 6, 272-278 [0036]
- **MCLAUGHLIN et al.**J. Virol., 1988, vol. 62, 1963-1973 [0037]
- **XIAO et al.**Exp. Neurobiol., 1997, vol. 144, 1 13-124 [0038]
- **FISHER et al.**J. Virol., 1996, vol. 70, 520-532 [0038]
- **GRAHAM et al.**Virology, 1973, vol. 52, 456- [0039]
- **SAMBROOK et al.**Molecular Cloning, a laboratory manualCold Spring Harbor Laboratories19890000 [0039]
- **DAVIS et al.**Basic Methods in Molecular BiologyElsevier19860000 [0039]
- **CHU et al.**Gene, 1981, vol. 13, 197- [0039]
- **CUSHMAN et al.**J. Med. Chem., 1991, vol. 34, 329-337 [0050] [0053] [0127]
- **CUSHMAN et al.**J. Med. Chem., vol. 34, 337-342 [0053]
- **GONZALEZET**Biochim. Biophys. Acta, 1979, vol. 562, 534-545 [0054]
- **MELLON, W. S.**Biochem. Pharmacol., 1984, vol. 33, 1047-1057 [0054]
- **MOUDGILET**J. Steroid Biochem., 1985, vol. 23, 125-132 [0054]
- **NAKANEET**Eur. J. Biochem., 1988, vol. 177, 91-96 [0054]
- **SKIDMOREET**Biochem. J., 1989, vol. 263, 73-80 [0054]
- **KUNZ et al.**Biochem. J., 1994, vol. 304, 337-340 [0056]
- **KUNZ et al.**Proc. Natl. Acad. Sci. USA, 1996, vol. 93, 255-259 [0056]
- **MOTAMEDIFAR et al.**Iran. J. Med. Sci., 2006, vol. 31, 28-32 [0056]
- **KUNTZ-SIMON et al.**J. Gen. Virol, 1995, vol. 76, 1409-1415 [0056]
- **MOOG et al.**J. Gen. Virol, 1996, vol. 77, 1993-1999 [0056]
- **YLISASTIGUI et al.**AIDS, 2004, vol. 18, 1101-1108 [0056]
- **SHAW et al.**AIDS, 2000, vol. 14, 899-902 [0056]
- **KABIRI et al.**Iran J. Med. Sci., 2001, vol. 26, 55-61 [0056]
- **GUO et al.**Surgery, 2007, vol. 142, 156-162 [0056]
- **Virology**19880000 [0057]
- **Fundamental Virology**19910000 [0057]
- **MACLEAN et al.**J. Gen. Virol., 1991, vol. 72, 632-639 [0059]
- **CHOUROIZMAN**Proc. Natl. Acad. Sci USA, 1992, vol. 89, 3266-3270 [0059]
- **CHOU et al.**J. Virol., 1994, vol. 68, 8304-8311 [0059]
- **DELUCA et al.**J. Virol., 1985, vol. 56, 558-570 [0059]
- **RICEKNIPE**J. Virol., 1990, vol. 64, 1704-1715 [0059] [0089]
- **SAMANIEGO et al.**J. Virol., 1995, vol. 69, 5705-5715 [0059]
- **COFFINLATCHMAN**Genetic Manipulation of the Nervous SystemAcademic Press1996000099-

114 [0059]

- **CONWAY et al.** Gene Ther., 1999, vol. 6, 973-985 [0061]
- **MUZYCZKA, N.** Current Topics in Microbiol. and Immunol., 1992, vol. 158, 97-129 [0063] [0085]
- **KOTIN, R.M.** Human Gene Therapy, 1994, vol. 5, 793-801 [0063] [0071] [0085]
- **THOMSON et al.** Virology, 1994, vol. 204, 304-311 [0063]
- Parvoviridae and their Replication **BERNS, K.I.** Fundamental Virology [0071]
- **RICE et al.** J. Virol., 1990, vol. 64, 1704-1715 [0076]
- **GRAHAM et al.** J. Gen. Virol., 1977, vol. 36, 59- [0082]
- **AIELLO et al.** Virology, 1979, vol. 94, 460- [0082]
- **LEBKOWSKI et al.** Molec. Cell. Biol., 1988, vol. 8, 3988-3996 [0085]
- **VINCENT et al.** Vaccines Cold Spring Harbor Laboratory Press 1990 0000 vol. 90, [0085]
- **CARTER, B.J.** Current Opinion in Biotechnology, 1992, vol. 3, 533-539 [0085]
- **SHELLINGSMITH** Gene Therapy, 1994, vol. 1, 165-169 [0085]
- **ZHOU et al.** J. Exp. Med., 1994, vol. 179, 1867-1875 [0085]
- **RICE et al.** J. Virol., 1989, vol. 63, 3399-3407 [0088]
- **GRAHAM et al.** J. Gen. Virol., 1977, vol. 36, 59-74 [0088]
- **KANG et al.** Gene Ther., 2009, vol. 16, 229-239 [0089]
- **KANG et al.** Gene Ther., 2009, vol. 16, 229-239 [0097]

Patentkrav

1. Fremgangsmåde til fremstilling af herpesvirus, hvilken fremgangsmåde omfatter dyrkning af herpesvirus i en
5 cellekultur, der omfatter aurintricarboxylsyre og serum, hvor aurintricarboxylsyre er til stede i en mængde, der øger udbyttet af herpesvirus.
2. Fremgangsmåde ifølge krav 1, hvor herpesviruset er et
10 *Herpes simplex*-1-virus (HSV-1).
3. Fremgangsmåde ifølge krav 2, hvor HSV-1 er et vildtype-HSV-1.
- 15 4. Fremgangsmåde ifølge krav 2, hvor HSV-1 er en rekombinant HSV-1-vektor.
5. Fremgangsmåde ifølge krav 4, hvor den rekombinante HSV-1-vektor er en HSV-1-d27.1-vektor.
20
6. Fremgangsmåde ifølge et hvilket som helst af kravene 1-5, hvor viruset dyrkes i 293- eller Vero-celler.
7. Fremgangsmåde ifølge krav 6, hvor viruset dyrkes i V27-
25 celler.
8. Fremgangsmåde til dyrkning af en HSV-1-d27.1-vektor, som omfatter:
(a) inficering af V27-celler med en HSV-1-d27.1-vektor; og
30 (b) dyrkning af de inficerede V27-celler i en cellekultur, der omfatter aurintricarboxylsyre, valproesyre eller dexamethason i en mængde, der øger HSV-1-d27.1-virustiteren, og yderligere omfatter serum.
- 35 9. Fremgangsmåde ifølge krav 8, hvor cellekulturen omfatter aurintricarboxylsyre.
10. Fremgangsmåde ifølge et hvilket som helst af kravene 1

til 9, hvor serummet er føtalt bovinserum.

11. Anvendelse af aurintricarboxylsyre til forøgelse af udbyttet af herpesvirus, der dannes i cellekultur, hvor
5 cellekulturen yderligere omfatter serum, og hvor herpesviruset dyrkes i Vero-celler eller 293-celler.

12. Anvendelse ifølge krav 11, hvor said cellekulturen omfatter Vero-celler.
10

13. Anvendelse ifølge krav 12, hvor Vero-cellerne er V27-celler.

14. Anvendelse ifølge et hvilket som helst af kravene 11 til
15 13, hvor viruset er som defineret i et hvilket som helst af kravene 2-5.

15. Cellekultur, der omfatter aurintricarboxylsyre, serum, et herpesvirus og 293- eller Vero-celler, hvor
20 aurintricarboxylsyre er til stede i en mængde, der øger udbyttet af herpesvirus, der dannes i cellekulturen.

16. Cellekultur ifølge krav 15, der omfatter V27-celler.

25 17. Cellekultur ifølge krav 15 eller 16, hvor herpesviruset er som defineret i et hvilket som helst af kravene 2-5.

DRAWINGS




Fig. 1




Fig. 2A




Fig. 2B




Fig. 2C




Fig. 3





Fig. 4A



HSV-1 vector /cells	Control: - ATA [DRP/ml]	+ATA [DRP/ml]	Ratio
d27-GFPV/27 cells	(5.4 ± 2.7) x 10 ⁷	**(3.3 ± 1.2) x 10 ⁸	6.1 x
wHSV-1 McIntyre/293 cells	(1.1 ± 0.5) x 10 ⁸	*** (1.0 ± 0.3) x 10 ⁹	9.1 x

Fig. 4B





Fig. 5A



**ATA: ATA concentration in 293 cells during to 2 hrs HSV co-infection

Fig. 5B




Fig. 6




Fig. 7