



(22) Date de dépôt/Filing Date: 2002/10/11
(41) Mise à la disp. pub./Open to Public Insp.: 2003/08/07
(45) Date de délivrance/Issue Date: 2018/05/01
(62) Demande originale/Original Application: 2 463 476
(30) Priorités/Priorities: 2001/10/11 (US60/328,101);
2002/08/30 (US60/406,934)

(51) Cl.Int./Int.Cl. *C12N 15/31* (2006.01),
A61K 39/095 (2006.01), *A61K 39/395* (2006.01),
C07K 14/22 (2006.01), *C07K 16/12* (2006.01),
C12N 1/21 (2006.01), *C12N 15/13* (2006.01),
C12N 15/63 (2006.01), *C12N 5/10* (2006.01),
C12N 5/16 (2006.01), *C12P 21/02* (2006.01),
C12Q 1/18 (2006.01), *G01N 33/569* (2006.01)

(72) Inventeurs/Inventors:
ZLOTNICK, GARY W., US;
FLETCHER, LEAH D., US;
FARLEY, JOHN, US;
BERNFELD, LIESEL A., US;

...

(73) Propriétaire/Owner:

(54) Titre : ANTIGENES FONDES SUR LA PROTEINE ORF 2086 DE L'ESPECE NEISSERIA ET COMPOSITIONS
RENFERMANT DE TELS ANTIGENES
(54) Title: ANTIGENS BASED ON NEISSERIA SPECIES ORF 2086 PROTEIN AND COMPOSITIONS COMPRISING
SUCH ANTIGENS

(57) **Abrégé/Abstract:**

The present invention relates to Neisseria ORF 2086 proteins, crossreactive immunogenic proteins which can be isolated from nesseria strains or prepared recombinantly, including immunogenic portions thereof, biological equivalents thereof, antibodies that immunospecifically bind to the foregoing and nucleic acid sequences encoding each of the foregoing, as well as immunogenic compositions comprising the same.

(72) Inventeurs(suite)/Inventors(continued): ZAGURSKY, ROBERT J., US; METCALF, BENJAMIN J., US

(73) Propriétaires(suite)/Owners(continued): WYETH HOLDINGS LLC, US

(74) Agent: BERESKIN & PARR LLP/S.E.N.C.R.L., S.R.L.

ABSTRACT

The present invention relates to *Neisseria* ORF 2086 proteins, crossreactive immunogenic proteins which can be isolated from nesserial strains or prepared recombinantly, including immunogenic portions thereof, biological equivalents thereof, antibodies that immunospecifically bind to the foregoing and nucleic acid sequences encoding each of the foregoing, as well as immunogenic compositions comprising the same.

-1-

ANTIGENS BASED ON NEISSERIA SPECIES ORF 2086 PROTEIN AND COMPOSITIONS COMPRISING SUCH ANTIGENS

Field of the Invention

The present invention relates to *Neisseria* ORF2086 proteins (Subfamily A and Subfamily B), which may be isolated from bacterial strains such as those of *Neisseria species*, including strains of *Neisseria meningitidis* (serogroups A, B, C, D, W-135, X, Y, Z and 29E), *Neisseria gonorrhoeae*, and *Neisseria lactamica*, as well as immunogenic portions and/or biological equivalents of said proteins. The present invention also relates to antibodies that immunospecifically bind to said proteins, immunogenic portions and/or biological equivalents. Further, the present invention relates to isolated polynucleotides comprising nucleic acid sequences encoding any of the foregoing proteins, immunogenic portions, biological equivalents and/or antibodies. Additionally, the present invention relates to compositions comprising said proteins, polynucleotides or antibodies, as well as methods for preparing said compositions. This invention relates to both recombinant forms and forms isolated from a natural source, as well as both lipidated and non-lipidated forms.

Background of the Invention

Meningococcal meningitis is a devastating disease that can kill children and young adults within hours despite the availability of antibiotics. Pizza *et al.*, 2000, *Science* 287:1816-1820. Meningitis is characterized as an inflammation of the meninges resulting in an intense headache, fever, loss of appetite, intolerance to light and sound, rigidity of muscles, especially in the neck, and in severe cases convulsions, vomiting and delirium leading to death. The symptoms of meningococcal meningitis appear suddenly and culminate in meningococcal septicemia with its characteristic hemorrhagic rash. A rapid diagnosis and immediate

-2-

treatment with large doses of antibiotics is critical if there is to be any chance of survival. 2000. *Bantam Medical Dictionary, Third Edition* 302.

Meningococcal meningitis is caused by *Neisseria meningitidis* (the meningococcus), a Gram-negative, capsulated bacterium that has been classified into several pathogenic serogroups including A, B, C, D, W-135, X, Y, Z and 29E. Serogroup B strains of *N. meningitidis* are a major cause of meningococcal disease throughout the world. For example, it is reported in the medical literature that serogroup B is responsible for about 50% of bacterial meningitis in infants and children residing in the United States and Europe. No vaccine currently exists to prevent meningococcal disease caused by *N. meningitidis* serogroup B.

Developing an immunogenic composition for the prevention of serogroup B meningococcal disease has been a challenge to researchers since the work of Goldschneider *et al.* over thirty years ago. Goldschneider *et al.*, 1969, *J. Exp. Med* 129(6):1307-26; Goldschneider *et al.*, 1969, *J. Exp. Med* 129(6):1327-48; Gotschlich *et al.*, 1969, *J. Exp. Med.* 129(6):1385-95; and Gotschlich *et al.*, 1969, *J. Exp. Med.* 129(6):1367-84. Unlike serogroup A disease, which virtually disappeared from North America after World War II, Achtman, M., 1995, *Trends in Microbiology* 3(5):186-92, disease caused by serogroup B and C organisms remains endemic throughout much of the economically developed world. The incidence of disease varies from <1/100,000 where endemic disease is rare to 200/100,000 in high risk populations during epidemics.

Vaccines based on polysaccharide conjugates have been developed against *N. meningitidis* serogroups A and C and appear to be effective in preventing disease. Currently, an immunogenic composition made of capsular polysaccharide from serogroups A, C, Y, & W-135 is available. Ambrosch *et al.*, 1983, Immunogenicity and side-effects of a new tetravalent. *Bulletin of the World Health Organization* 61(2):317-23. However, this immunogenic composition elicits a T-cell independent immune response, is not effective in young children, and provides no coverage for serogroup B strains, which cause upwards of 50% of meningococcal disease.

-3-

Others have also attempted to develop immunogenic compositions using capsular polysaccharides. Recently, immunogenic compositions for serogroup C disease prepared by conjugating the serogroup C capsular material to proteins have been licensed for use in Europe. However, the serogroup B capsule may be
5 unsuitable as a vaccine candidate because the capsule polysaccharide is composed of polysialic acid which bears a similarity to carbohydrate moieties on developing human neural tissues. This sugar moiety is recognized as a self-antigen and is thus poorly immunogenic in humans.

Outer membrane proteins (OMP's) have been developed as alternative vaccine
10 antigens for serogroup B disease. Monoclonal antibody binding to the two variable regions of PorA define the serosubtyping scheme for meningococci. PorA proteins thus serve as the serosubtyping antigens (Abdillahi *et al.*, 1988, *Microbial Pathogenesis* 4(1):27-32) for meningococcal strains and are being actively investigated as components of a serogroup B immunogenic composition (Poolman,
15 1996, *Adv. Exp. Med. Biol.* 397:73-7), since they can elicit bactericidal antibodies (Saukkonen, 1987, *Microbial Pathogenesis* 3(4):261-7). Bactericidal antibodies are thought to be an indicator of protection and any new immunogenic composition candidate should elicit these functional antibodies.

Studies in humans as well as animals indicate that the serosubtyping antigen,
20 PorA, elicits bactericidal antibodies. However, the immune response to Por A is generally serosubtype specific. In particular, serosubtyping data indicate that an immunogenic composition made of PorAs may require a PorA for each serosubtype to be covered by such an immunogenic composition, perhaps as many as six to nine. Therefore, 6-9 PorAs will be needed to cover 70-80% of serogroup B strains. Thus,
25 the variable nature of this protein requires a multivalent vaccine composition to protect against a sufficient number of meningococcal serosubtype clinical isolates.

Developing an immunogenic composition for serogroup B meningococci has been so difficult that recently several groups have sequenced the genomes from strains representing both serogroups A and B to assist in identifying new
30 immunogenic composition candidates. Tettelin, 2000, *Science*, 287(5459):1809-15;

-4-

Pizza *et al.*, 2000, *Science* 287:1816-1820. Identifying new immunogenic composition candidates, even with the knowledge of the neisserial genome, is a challenging process for which adequate mathematical algorithms do not currently exist. In fact, a recent report indicates that despite identifying hundreds of open
 5 reading frames ("ORFs") containing theoretical membrane spanning domains, problems with expression, purification, and inducing surface reactive, and functionally active antibodies have led investigators to only seven candidates for a serogroup B meningococcal immunogenic composition. *See Id*. One of these was previously known.

10 Accordingly, there remains a need for compositions that elicit antibodies that (1) are bactericidal to multiple neisserial strains; (2) react with the surface of multiple strains; (3) confer passive protection against a live challenge; and/or (4) prevent colonization

Summary of the Invention

15 The present invention provides *Neisseria* ORF2086 proteins ("2086 proteins"), including 2086 Subfamily A proteins and 2086 Subfamily B proteins. Each of the 2086 proteins are proteins that can be isolated from native neisserial strains, including strains of *Neisseria meningitidis* (serogroups A, B, C, D, W-135, X, Y, Z, and 29E), *Neisseria gonorrhoeae*, and *Neisseria lactamica*. The 2086 proteins
 20 may also be prepared using recombinant technology.

In particular, the present invention provides the 2086 proteins, immunogenic portions thereof, and/or biological equivalents thereof, antibodies that immunospecifically bind to any of the foregoing, and polynucleotides comprising nucleic acid sequences that encode any of the foregoing. The present invention
 25 includes compositions, comprising said proteins, polynucleotides and antibodies, as well as methods for preparing said compositions. The 2086 proteins herein include recombinant forms and forms isolated from a natural source, as well as both lipidated and non-lipidated forms.

-5-

In particular, candidates were identified that in some embodiments have the ability to elicit antibodies that (1) are bactericidal to multiple neisserial strains, such as strains of *N. meningitidis*, *N. gonorrhoeae*, and/or *N. lactamica*; (2) react with the surface of multiple strains; (3) confer passive protection against a live challenge; and/or (4) prevent colonization, as well as methods of using said compositions and methods of preparing said compositions. Various embodiments of the invention are described below.

10 Brief Description of the Drawings

FIG. 1A depicts an SDS-PAGE gel that depicts the two major proteins of the protein fractions obtained from the experiments for identifying neisserial membrane protein extract that is capable of eliciting bactericidal antibodies against heterologous strains.

15 FIG. 1B depicts the results from the experiments from the identification of the two major proteins by analysis of TMAE Flow Through components by protease digestion and reverse Phase N-terminal sequencing.

FIG. 2 depicts the purification scheme and homogeneity as determined by SDS-PAGE of rLP2086.

20 FIG. 3 depicts the results from the experiments from the identification of the two major proteins and one minor protein by analysis of TMAE Flow Through components by LC-MS/MS and the corresponding SDS-PAGE.

FIG. 4 is an SDS-PAGE gel from the recombinant expression of 2086 protein.

25

-6-

FIG. 5 is a schematic diagram of plasmid pPX7340, as described in the examples herein.

FIG. 6 is a schematic diagram of plasmid pPX7328 as described in the examples herein.

5 FIG. 7 is a schematic diagram of plasmid pPX7343 as described in the examples herein.

FIG. 8 illustrates N-terminal regions of 2086 gene from various strains.

FIG. 9A is a flow chart showing the preliminary steps in the identification of an immunogenic component in a nesserial strain.

10 FIG. 9B is a flow chart showing the final steps in the identification of an immunogenic component in a nesserial strain.

FIG. 10A is a schematic diagram of the pBAD arabinose inducible promoter which drives the expression of the P4 signal/ORF2086 fusion protein to express a lipidated form of rP2086 as described in the examples herein.

15 FIG. 10B is a schematic diagram of the pET9a-T7 vector for recombinant expression of nonlipidated form of ORF2086.

FIG. 11A is a photograph representing whole cell lysates of *E. coli* B expressing the rLP2086 protein.

20 FIG. 11B is a photograph representing whole cell lysates of *E. coli* B expressing the rP2086 protein.

FIG. 12 is a phylogenetic tree showing the organization of the subfamilies and groups of ORF2086 proteins.

FIG. 13 is a graphical illustration of whole cell ELISA data for the rLP2086 Subfamily A antisera.

-7-

FIG. 14 is a graphical illustration of whole cell ELISA data for the rLP2086 Subfamily B antisera.

FIG. 15 is a graphical illustration of the results of the rLP2086 mixing study - WCE Titers.

5 FIG. 16 is a graphical illustration of the results of the rLP2086/rPorA mixing study - WCE Titers.

FIG. 17 is a Western Blot showing reactivity of rLP2086 mouse antisera to P2086 Subfamily B *N. meningitidis* whole cell lysates.

10 FIG. 18 is a Western Blot showing reactivity of rLP2086 mouse antisera to P2086 Subfamily A *N. meningitidis* and *N. lactamica* whole cell lysates.

Sequence Summary

SEQ ID NOS. For Studied Sequences:

15 SEQ ID NO:1 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from L3 6275 strain when combined with a native leader sequence.

SEQ ID NO:2 amino acid sequence for mature 2086 protein from L3 6275 strain prepared using a native leader sequence.

SEQ ID NO:3 nucleic acid sequence for encoding amino acid sequence for mature 2086 protein from L3 6275 when combined with a P4 leader sequence.

20 SEQ ID NO:4 amino acid sequence for mature 2086 protein from L3 6275 strain prepared using a P4 leader sequence.

SEQ ID NO:5 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from L3 6275 strain.

SEQ ID NO:6 amino acid sequence for mature 2086 protein from L3 6275 strain.

-8-

SEQ ID NO:7 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from CDC2369 strain when combined with a native leader sequence.

SEQ ID NO:8 amino acid sequence for mature 2086 protein from CDC2369 strain prepared using a native leader sequence.

- 5 SEQ ID NO:9 nucleic acid sequence for encoding amino acid sequence for mature 2086 protein from CDC2369 when combined with a P4 leader sequence.

SEQ ID NO:10 amino acid sequence for mature 2086 protein from CDC2369 strain prepared using a P4 leader sequence.

- 10 SEQ ID NO:11 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from CDC2369 strain.

SEQ ID NO:12 amino acid sequence for mature 2086 protein from CDC2369 strain.

SEQ ID NO:13 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from CDC1034 strain when combined with a native leader sequence.

- 15 SEQ ID NO:14 amino acid sequence for mature 2086 protein from CDC1034 strain prepared using a native leader sequence.

SEQ ID NO:15 nucleic acid sequence for encoding amino acid sequence for mature 2086 protein from CDC1034 when combined with a P4 leader sequence.

SEQ ID NO:16 amino acid sequence for mature 2086 protein from CDC1034 strain prepared using a P4 leader sequence.

- 20 SEQ ID NO:17 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from CDC1034 strain.

SEQ ID NO:18 amino acid sequence for mature 2086 protein from CDC1034 strain.

SEQ ID NO:19 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from L4 891 strain when combined with a native leader sequence.

SEQ ID NO:20 amino acid sequence for mature 2086 protein from L4 891 strain prepared using a native leader sequence.

SEQ ID NO:21 nucleic acid sequence for encoding amino acid sequence for mature 2086 protein from L4 891 when combined with a P4 leader sequence.

- 5 SEQ ID NO:22 amino acid sequence for mature 2086 protein from L4 891 strain prepared using a P4 leader sequence.

SEQ ID NO:23 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from L4 891 strain.

SEQ ID NO:24 amino acid sequence for mature 2086 protein from L4 891 strain.

- 10 SEQ ID NO:25 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from B16B6 strain when combined with a native leader sequence.

SEQ ID NO:26 amino acid sequence for mature 2086 protein from B16B6 strain prepared using a native leader sequence.

- 15 SEQ ID NO:27 nucleic acid sequence for encoding amino acid sequence for mature 2086 protein from B16B6 when combined with a P4 leader sequence.

SEQ ID NO:28 amino acid sequence for mature 2086 protein from B16B6 strain prepared using a P4 leader sequence.

SEQ ID NO:29 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from B16B6 strain.

- 20 SEQ ID NO:30 amino acid sequence for mature 2086 protein from B16B6 strain.

SEQ ID NO:31 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from W135 (ATCC35559) strain when combined with a native leader sequence.

WO 03/063766

PCT/US02/32369

-10-

SEQ ID NO:32 amino acid sequence for mature 2086 protein from W135 (ATCC35559) strain prepared using a native leader sequence.

SEQ ID NO:33 nucleic acid sequence for encoding amino acid sequence for mature 2086 protein from W135 (ATCC35559) when combined with a P4 leader sequence.

- 5 SEQ ID NO:34 amino acid sequence for mature 2086 protein from W135 (ATCC35559) strain prepared using a P4 leader sequence.

SEQ ID NO:35 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from W135 (ATCC35559) strain.

- 10 SEQ ID NO:36 amino acid sequence for mature 2086 protein from W135 (ATCC35559) strain.

SEQ ID NO:37 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from C11 strain when combined with a native leader sequence.

SEQ ID NO:38 amino acid sequence for mature 2086 protein from C11 strain prepared using a native leader sequence.

- 15 SEQ ID NO:39 nucleic acid sequence for encoding amino acid sequence for mature 2086 protein from C11 when combined with a P4 leader sequence.

SEQ ID NO:40 amino acid sequence for mature 2086 protein from C11 strain prepared using a P4 leader sequence.

- 20 SEQ ID NO:41 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from C11 strain.

SEQ ID NO:42 amino acid sequence for mature 2086 protein from C11 strain.

SEQ ID NO:43 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from Y (ATCC35561) strain when combined with a native leader sequence.

WO 03/063766

PCT/US02/32369

-11-

SEQ ID NO:44 amino acid sequence for mature 2086 protein from Y (ATCC35561) strain prepared using a native leader sequence.

SEQ ID NO:45 nucleic acid sequence for encoding amino acid sequence for mature 2086 protein from Y (ATCC35561) when combined with a P4 leader sequence.

- 5 SEQ ID NO:46 amino acid sequence for mature 2086 protein from Y (ATCC35561) strain prepared using a P4 leader sequence.

SEQ ID NO:47 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from Y (ATCC35561) strain.

- 10 SEQ ID NO:48 amino acid sequence for mature 2086 protein from Y (ATCC35561) strain.

SEQ ID NO:49 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from M98 250732 strain when combined with a native leader sequence.

SEQ ID NO:50 amino acid sequence for mature 2086 protein from M98 250732 strain prepared using a native leader sequence.

- 15 SEQ ID NO:51 nucleic acid sequence for encoding amino acid sequence for mature 2086 protein from M98 250732 when combined with a P4 leader sequence.

SEQ ID NO:52 amino acid sequence for mature 2086 protein from M98 250732 strain prepared using a P4 leader sequence.

- 20 SEQ ID NO:53 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from M98 250732 strain.

SEQ ID NO:54 amino acid sequence for mature 2086 protein from M98 250732 strain.

SEQ ID NO:55 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from M98 250771 strain when combined with a native leader sequence.

-12-

SEQ ID NO:56 amino acid sequence for mature 2086 protein from M98 250771 strain prepared using a native leader sequence.

SEQ ID NO:57 nucleic acid sequence for encoding amino acid sequence for mature 2086 protein from M98 250771 when combined with a P4 leader sequence.

- 5 SEQ ID NO:58 amino acid sequence for mature 2086 protein from M98 250771 strain prepared using a P4 leader sequence.

SEQ ID NO:59 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from M98 250771 strain.

- 10 SEQ ID NO:60 amino acid sequence for mature 2086 protein from M98 250771 strain.

SEQ ID NO:61 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from CDC1135 strain when combined with a native leader sequence.

SEQ ID NO:62 amino acid sequence for mature 2086 protein from CDC1135 strain prepared using a native leader sequence.

- 15 SEQ ID NO:63 nucleic acid sequence for encoding amino acid sequence for mature 2086 protein from CDC1135 when combined with a P4 leader sequence.

SEQ ID NO:64 amino acid sequence for mature 2086 protein from CDC1135 strain prepared using a P4 leader sequence.

- 20 SEQ ID NO:65 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from CDC1135 strain.

SEQ ID NO:66 amino acid sequence for mature 2086 protein from CDC1135 strain.

SEQ ID NO:67 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from M97 252153 strain when combined with a native leader sequence.

WO 03/063766

PCT/US02/32369

-13-

SEQ ID NO:68 amino acid sequence for mature 2086 protein from M97 252153 strain prepared using a native leader sequence.

SEQ ID NO:69 nucleic acid sequence for encoding amino acid sequence for mature 2086 protein from M97 252153 when combined with a P4 leader sequence.

- 5 SEQ ID NO:70 amino acid sequence for mature 2086 protein from M97 252153 strain prepared using a P4 leader sequence.

SEQ ID NO:71 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from M97 252153 strain.

- 10 SEQ ID NO:72 amino acid sequence for mature 2086 protein from M97 252153 strain.

SEQ ID NO:73 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from CDC1610 strain when combined with a native leader sequence.

SEQ ID NO:74 amino acid sequence for mature 2086 protein from CDC1610 strain prepared using a native leader sequence.

- 15 SEQ ID NO:75 nucleic acid sequence for encoding amino acid sequence for mature 2086 protein from CDC1610 when combined with a P4 leader sequence.

SEQ ID NO:76 amino acid sequence for mature 2086 protein from CDC1610 strain prepared using a P4 leader sequence.

- 20 SEQ ID NO:77 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from CDC1610 strain.

SEQ ID NO:78 amino acid sequence for mature 2086 protein from CDC1610 strain.

SEQ ID NO:79 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from CDC1492 strain when combined with a native leader sequence.

WO 03/063766

PCT/US02/32369

-14-

SEQ ID NO:80 amino acid sequence for mature 2086 protein from CDC1492 strain prepared using a native leader sequence.

SEQ ID NO:81 nucleic acid sequence for encoding amino acid sequence for mature 2086 protein from CDC1492 when combined with a P4 leader sequence.

- 5 SEQ ID NO:82 amino acid sequence for mature 2086 protein from CDC1492 strain prepared using a P4 leader sequence.

SEQ ID NO:83 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from CDC1492 strain.

SEQ ID NO:84 amino acid sequence for mature 2086 protein from CDC1492 strain.

- 10 SEQ ID NO:85 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from L8 M978 strain when combined with a native leader sequence.

SEQ ID NO:86 amino acid sequence for mature 2086 protein from L8 M978 strain prepared using a native leader sequence.

- 15 SEQ ID NO:87 nucleic acid sequence for encoding amino acid sequence for mature 2086 protein from L8 M978 when combined with a P4 leader sequence.

SEQ ID NO:88 amino acid sequence for mature 2086 protein from L8 M978 strain prepared using a P4 leader sequence.

SEQ ID NO:89 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from L8 M978 strain.

- 20 SEQ ID NO:90 amino acid sequence for mature 2086 protein from L8 M978 strain.

SEQ ID NO:91 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from M97 252988 strain when combined with a native leader sequence.

SEQ ID NO:92 amino acid sequence for mature 2086 protein from M97 252988 strain prepared using a native leader sequence.

WO 03/063766

PCT/US02/32369

-15-

SEQ ID NO:93 nucleic acid sequence for encoding amino acid sequence for mature 2086 protein from M97 252988 when combined with a P4 leader sequence.

SEQ ID NO:94 amino acid sequence for mature 2086 protein from M97 252988 strain prepared using a P4 leader sequence.

- 5 SEQ ID NO:95 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from M97 252988 strain.

SEQ ID NO:96 amino acid sequence for mature 2086 protein from M97 252988 strain.

- 10 SEQ ID NO:97 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from M97 252697 strain when combined with a native leader sequence.

SEQ ID NO:98 amino acid sequence for mature 2086 protein from M97 252697 strain prepared using a native leader sequence.

SEQ ID NO:99 nucleic acid sequence for encoding amino acid sequence for mature 2086 protein from M97 252697 when combined with a P4 leader sequence.

- 15 SEQ ID NO:100 amino acid sequence for mature 2086 protein from M97 252697 strain prepared using a P4 leader sequence.

SEQ ID NO:101 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from M97 252697 strain.

- 20 SEQ ID NO:102 amino acid sequence for mature 2086 protein from M97 252697 strain.

SEQ ID NO:103 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from 6557 strain when combined with a native leader sequence.

SEQ ID NO:104 amino acid sequence for mature 2086 protein from 6557 strain prepared using a native leader sequence.

-16-

SEQ ID NO:105 nucleic acid sequence for encoding amino acid sequence for mature 2086 protein from 6557 when combined with a P4 leader sequence.

SEQ ID NO:106 amino acid sequence for mature 2086 protein from 6557 strain prepared using a P4 leader sequence.

- 5 SEQ ID NO:107 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from 6557 strain.

SEQ ID NO:108 amino acid sequence for mature 2086 protein from 6557 strain.

SEQ ID NO:109 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from 2996 strain when combined with a native leader sequence.

- 10 SEQ ID NO:110 amino acid sequence for mature 2086 protein from 2996 strain prepared using a native leader sequence.

SEQ ID NO:111 nucleic acid sequence for encoding amino acid sequence for mature 2086 protein from 2996 when combined with a P4 leader sequence.

- 15 SEQ ID NO:112 amino acid sequence for mature 2086 protein from 2996 strain prepared using a P4 leader sequence.

SEQ ID NO:113 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from 2996 strain.

SEQ ID NO:114 amino acid sequence for mature 2086 protein from 2996 strain.

- 20 SEQ ID NO:115 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from M97 252976 strain when combined with a native leader sequence.

SEQ ID NO:116 amino acid sequence for mature 2086 protein from M97 252976 strain prepared using a native leader sequence.

SEQ ID NO:117 nucleic acid sequence for encoding amino acid sequence for mature 2086 protein from M97 252976 when combined with a P4 leader sequence.

-17-

SEQ ID NO:118 amino acid sequence for mature 2086 protein from M97 252976 strain prepared using a P4 leader sequence.

SEQ ID NO:119 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from M97 252976 strain.

- 5 SEQ ID NO:120 amino acid sequence for mature 2086 protein from M97 252976 strain.

SEQ ID NO:121 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from M97 251854 strain when combined with a native leader sequence.

- 10 SEQ ID NO:122 amino acid sequence for mature 2086 protein from M97 251854 strain prepared using a native leader sequence.

SEQ ID NO:123 nucleic acid sequence for encoding amino acid sequence for mature 2086 protein from M97 251854 when combined with a P4 leader sequence.

SEQ ID NO:124 amino acid sequence for mature 2086 protein from M97 251854 strain prepared using a P4 leader sequence.

- 15 SEQ ID NO:125 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from M97 251854 strain.

SEQ ID NO:126 amino acid sequence for mature 2086 protein from M97 251854 strain.

- 20 SEQ ID NO:127 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from CDC1521 strain when combined with a native leader sequence.

SEQ ID NO:128 amino acid sequence for mature 2086 protein from CDC1521 strain prepared using a native leader sequence.

SEQ ID NO:129 nucleic acid sequence for encoding amino acid sequence for mature 2086 protein from CDC1521 when combined with a P4 leader sequence.

WO 03/063766

PCT/US02/32369

-18-

SEQ ID NO:130 amino acid sequence for mature 2086 protein from CDC1521 strain prepared using a P4 leader sequence.

SEQ ID NO:131 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from CDC1521 strain.

- 5 SEQ ID NO:132 amino acid sequence for mature 2086 protein from CDC1521 strain.

SEQ ID NO:133 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from M98 250622 strain when combined with a native leader sequence.

SEQ ID NO:134 amino acid sequence for mature 2086 protein from M98 250622 strain prepared using a native leader sequence.

- 10 SEQ ID NO:135 nucleic acid sequence for encoding amino acid sequence for mature 2086 protein from M98 250622 when combined with a P4 leader sequence.

SEQ ID NO:136 amino acid sequence for mature 2086 protein from M98 250622 strain prepared using a P4 leader sequence.

- 15 SEQ ID NO:137 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from M98 250622 strain.

SEQ ID NO:138 amino acid sequence for mature 2086 protein from M98 250622 strain.

SEQ ID NO:139 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from 870446 strain when combined with a native leader sequence.

- 20 SEQ ID NO:140 amino acid sequence for mature 2086 protein from 870446 strain prepared using a native leader sequence.

SEQ ID NO:141 nucleic acid sequence for encoding amino acid sequence for mature 2086 protein from 870446 when combined with a P4 leader sequence.

WO 03/063766

PCT/US02/32369

-19-

SEQ ID NO:142 amino acid sequence for mature 2086 protein from 870446 strain prepared using a P4 leader sequence.

SEQ ID NO:143 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from 870446 strain.

- 5 SEQ ID NO:144 amino acid sequence for mature 2086 protein from 870446 strain.

SEQ ID NO:145 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from M97 253248 strain when combined with a native leader sequence.

SEQ ID NO:146 amino acid sequence for mature 2086 protein from M97 253248 strain prepared using a native leader sequence.

- 10 SEQ ID NO:147 nucleic acid sequence for encoding amino acid sequence for mature 2086 protein from M97 253248 when combined with a P4 leader sequence.

SEQ ID NO:148 amino acid sequence for mature 2086 protein from M97 253248 strain prepared using a P4 leader sequence.

- 15 SEQ ID NO:149 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from M97 253248 strain.

SEQ ID NO:150 amino acid sequence for mature 2086 protein from M97 253248 strain.

SEQ ID NO:151 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from M98 250809 strain when combined with a native leader sequence.

- 20 SEQ ID NO:152 amino acid sequence for mature 2086 protein from M98 250809 strain prepared using a native leader sequence.

SEQ ID NO:153 nucleic acid sequence for encoding amino acid sequence for mature 2086 protein from M98 250809 when combined with a P4 leader sequence.

-20-

SEQ ID NO:154 amino acid sequence for mature 2086 protein from M98 250809 strain prepared using a P4 leader sequence.

SEQ ID NO:155 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from M98 250809 strain.

- 5 SEQ ID NO:156 amino acid sequence for mature 2086 protein from M98 250809 strain.

SEQ ID NO:157 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from L5 M981 strain when combined with a native leader sequence.

- 10 SEQ ID NO:158 amino acid sequence for mature 2086 protein from L5 M981 strain prepared using a native leader sequence.

SEQ ID NO:159 nucleic acid sequence for encoding amino acid sequence for mature 2086 protein from L5 M981 when combined with a P4 leader sequence.

SEQ ID NO:160 amino acid sequence for mature 2086 protein from L5 M981 strain prepared using a P4 leader sequence.

- 15 SEQ ID NO:161 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from L5 M981 strain.

SEQ ID NO:162 amino acid sequence for mature 2086 protein from L5 M981 strain.

SEQ ID NO:163 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from NMB strain when combined with a native leader sequence.

- 20 SEQ ID NO:164 amino acid sequence for mature 2086 protein from NMB strain prepared using a native leader sequence.

SEQ ID NO:165 nucleic acid sequence for encoding amino acid sequence for mature 2086 protein from NMB when combined with a P4 leader sequence.

-21-

SEQ ID NO:166 amino acid sequence for mature 2086 protein from NMB strain prepared using a P4 leader sequence.

SEQ ID NO:167 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from NMB strain.

- 5 SEQ ID NO:168 amino acid sequence for mature 2086 protein from NMB strain.

SEQ ID NO:169 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from M98 250572 strain when combined with a native leader sequence.

SEQ ID NO:170 amino acid sequence for mature 2086 protein from M98 250572 strain prepared using a native leader sequence.

- 10 SEQ ID NO:171 nucleic acid sequence for encoding amino acid sequence for mature 2086 protein from M98 250572 when combined with a P4 leader sequence.

SEQ ID NO:172 amino acid sequence for mature 2086 protein from M98 250572 strain prepared using a P4 leader sequence.

- 15 SEQ ID NO:173 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from M98 250572 strain.

SEQ ID NO:174 amino acid sequence for mature 2086 protein from M98 250572 strain.

- 20 SEQ ID NO:175 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from A4 Sanford; M97 251836 PART; M97 251957; M97 251985; M97 252060; M97 251870; M97 251994; M98 250024; M97 251905; M97 251876; M97 251898; or M97 251830 strain when combined with a native leader sequence.

- 25 SEQ ID NO:176 amino acid sequence for mature 2086 protein from A4 Sanford; M97 251836 PART; M97 251957; M97 251985; M97 252060; M97 251870; M97 251994; M98 250024; M97 251905; M97 251876; M97 251898; or M97 251830 strain prepared using a native leader sequence.

-22-

SEQ ID NO:177 nucleic acid sequence for encoding amino acid sequence for mature 2086 protein from A4 Sanford; M97 251836 PART; M97 251957; M97 251985; M97 252060; M97 251870; M97 251994; M98 250024; M97 251905; M97 251876; M97 251898; or M97 251830 when combined with a P4 leader sequence.

- 5 SEQ ID NO:178 amino acid sequence for mature 2086 protein from A4 Sanford; M97 251836 PART; M97 251957; M97 251985; M97 252060; M97 251870; M97 251994; M98 250024; M97 251905; M97 251876; M97 251898; or M97 251830 strain prepared using a P4 leader sequence.

- 10 SEQ ID NO:179 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from A4 Sanford; M97 251836 part; M97 251957; M97 251985; M97 252060; M97 251870; M97 251994; M98 250024; M97 251905; M97 251876; M97 251898; or M97 251830 strain.

- 15 SEQ ID NO:180 amino acid sequence for mature 2086 protein from A4 Sanford; M97 251836 PART; M97 251957; M97 251985; M97 252060; M97 251870; M97 251994; M98 250024; M97 251905; M97 251876; M97 251898; or M97 251830 strain.

SEQ ID NO:181 nucleic acid sequence encoding partial amino acid sequence for mature 2086 protein from CDC937 strain when combined with a native leader sequence.

- 20 SEQ ID NO:182 amino acid sequence for mature 2086 protein from CDC937 strain prepared using a native leader sequence.

SEQ ID NO:183 nucleic acid sequence for encoding partial amino acid sequence for mature 2086 protein from CDC937 when combined with a P4 leader sequence.

- 25 SEQ ID NO:184 amino acid sequence for mature 2086 protein from CDC937 strain prepared using a P4 leader sequence.

SEQ ID NO:185 nucleic acid sequence encoding partial amino acid sequence for mature 2086 protein from CDC937 strain.

-23-

SEQ ID NO:186 amino acid sequence for mature 2086 protein from CDC937 strain.

SEQ ID NO:187 nucleic acid sequence encoding partial amino acid sequence for mature 2086 protein from M97 252097 strain when combined with a native leader sequence.

- 5 SEQ ID NO:188 amino acid sequence for mature 2086 protein from M97 252097 strain prepared using a native leader sequence.

SEQ ID NO:189 nucleic acid sequence for encoding partial amino acid sequence for mature 2086 protein from M97 252097 when combined with a P4 leader sequence.

- 10 SEQ ID NO:190 amino acid sequence for mature 2086 protein from M97 252097 strain prepared using a P4 leader sequence.

SEQ ID NO:191 nucleic acid sequence encoding partial amino acid sequence for mature 2086 protein from M97 252097 strain.

SEQ ID NO:192 amino acid sequence for mature 2086 protein from M97 252097 strain.

- 15 SEQ ID NO:193 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from 870227 strain when combined with a native leader sequence.

SEQ ID NO:194 amino acid sequence for mature 2086 protein from 870227 strain prepared using a native leader sequence.

- 20 SEQ ID NO:195 nucleic acid sequence for encoding amino acid sequence for mature 2086 protein from 870227 when combined with a P4 leader sequence.

SEQ ID NO:196 amino acid sequence for mature 2086 protein from 870227 strain prepared using a P4 leader sequence.

SEQ ID NO:197 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from 870227 strain.

-24-

SEQ ID NO:198 amino acid sequence for mature 2086 protein from 870227 strain.

SEQ ID NO:199 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from H355 strain when combined with a native leader sequence.

5 SEQ ID NO:200 amino acid sequence for mature 2086 protein from H355 strain prepared using a native leader sequence.

SEQ ID NO:201 nucleic acid sequence for encoding amino acid sequence for mature 2086 protein from H355 when combined with a P4 leader sequence.

SEQ ID NO:202 amino acid sequence for mature 2086 protein from H355 strain prepared using a P4 leader sequence.

10 SEQ ID NO:203 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from H355 strain.

SEQ ID NO:204 amino acid sequence for mature 2086 protein from H355 strain.

SEQ ID NO:205 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from H44_76 strain when combined with a native leader sequence.

15 SEQ ID NO:206 amino acid sequence for mature 2086 protein from H44_76 strain prepared using a native leader sequence.

SEQ ID NO:207 amino acid sequence for mature 2086 protein from H44_76 strain prepared using a P4 leader sequence.

20 SEQ ID NO:208 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from H44_76 strain.

SEQ ID NO:209 nucleic acid sequence for encoding amino acid sequence for mature 2086 protein from H44_76 when combined with a P4 leader sequence.

SEQ ID NO:210 amino acid sequence for mature 2086 protein from H44_76 strain.

WO 03/063766

PCT/US02/32369

-25-

SEQ ID NO:211 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from 8529 strain when combined with a native leader sequence.

SEQ ID NO:212 amino acid sequence for mature 2086 protein from 8529 strain prepared using a native leader sequence.

- 5 SEQ ID NO:213 nucleic acid sequence for encoding amino acid sequence for mature 2086 protein from 8529 when combined with a P4 leader sequence.

SEQ ID NO:214 amino acid sequence for mature 2086 protein from 8529 strain prepared using a P4 leader sequence.

- 10 SEQ ID NO:215 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from 8529 strain.

SEQ ID NO:216 amino acid sequence for mature 2086 protein from 8529 strain.

SEQ ID NO:217 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from 6940 strain when combined with a native leader sequence.

- 15 SEQ ID NO:218 amino acid sequence for mature 2086 protein from 6940 strain prepared using a native leader sequence.

SEQ ID NO:219 nucleic acid sequence for encoding amino acid sequence for mature 2086 protein from 6940 when combined with a P4 leader sequence.

SEQ ID NO:220 amino acid sequence for mature 2086 protein from 6940 strain prepared using a P4 leader sequence.

- 20 SEQ ID NO:221 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from 6940 strain.

SEQ ID NO:222 amino acid sequence for mature 2086 protein from 6940 strain.

SEQ ID NO:223 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from M982 strain when combined with a native leader sequence.

WO 03/063766

PCT/US02/32369

-26-

SEQ ID NO:224 amino acid sequence for mature 2086 protein from M982 strain prepared using a native leader sequence.

SEQ ID NO:225 nucleic acid sequence for encoding amino acid sequence for mature 2086 protein from M982 when combined with a P4 leader sequence.

- 5 SEQ ID NO:226 amino acid sequence for mature 2086 protein from M982 strain prepared using a P4 leader sequence.

SEQ ID NO:227 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from M982 strain.

SEQ ID NO:228 amino acid sequence for mature 2086 protein from M982 strain.

- 10 SEQ ID NO:229 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from 880049 strain when combined with a native leader sequence.

SEQ ID NO:230 amino acid sequence for mature 2086 protein from 880049 strain prepared using a native leader sequence.

- 15 SEQ ID NO:231 nucleic acid sequence for encoding amino acid sequence for mature 2086 protein from 880049 when combined with a P4 leader sequence.

SEQ ID NO:232 amino acid sequence for mature 2086 protein from 880049 strain prepared using a P4 leader sequence.

SEQ ID NO:233 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from 880049 strain.

- 20 SEQ ID NO:234 amino acid sequence for mature 2086 protein from 880049 strain.

SEQ ID NO:235 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from M97 253524, M97 251885, and M97 251926 strains when combined with a native leader sequence.

-27-

SEQ ID NO:236 amino acid sequence for mature 2086 protein from M97 253524, M97 251885, and M97 251926 strains prepared using a native leader sequence.

SEQ ID NO:237 nucleic acid sequence for encoding amino acid sequence for mature 2086 protein from M97 253524, M97 251885, and M97 251926 strains when
5 combined with a P4 leader sequence.

SEQ ID NO:238 amino acid sequence for mature 2086 protein from M97 253524, M97 251885, and M97 251926 strains prepared using a P4 leader sequence.

SEQ ID NO:239 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from M97 253524, M97 251885, and M97 251926 strains.

10 SEQ ID NO:240 amino acid sequence for mature 2086 protein from M97 253524, M97 251885, and M97 251926 strains.

SEQ ID NO:241 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from M98 250670 strain when combined with a native leader sequence.

SEQ ID NO:242 amino acid sequence for mature 2086 protein from M98 250670
15 strain prepared using a native leader sequence.

SEQ ID NO:243 nucleic acid sequence for encoding amino acid sequence for mature 2086 protein from M98 250670 when combined with a P4 leader sequence.

SEQ ID NO:244 amino acid sequence for mature 2086 protein from M98 250670 strain prepared using a P4 leader sequence.

20 SEQ ID NO:245 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from M98 250670 strain.

SEQ ID NO:246 amino acid sequence for mature 2086 protein from M98 250670 strain.

SEQ ID NO:247 nucleic acid sequence encoding amino acid sequence for mature
25 2086 protein from CDC1573 strain when combined with a native leader sequence.

-28-

SEQ ID NO:248 amino acid sequence for mature 2086 protein from CDC1573 strain prepared using a native leader sequence.

SEQ ID NO:249 nucleic acid sequence for encoding amino acid sequence for mature 2086 protein from CDC1573 when combined with a P4 leader sequence.

- 5 SEQ ID NO:250 amino acid sequence for mature 2086 protein from CDC1573 strain prepared using a P4 leader sequence.

SEQ ID NO:251 nucleic acid sequence encoding amino acid sequence for mature 2086 protein from CDC1573 strain.

SEQ ID NO:252 amino acid sequence for mature 2086 protein from CDC1573 strain.

- 10 SEQ ID NO:253 partial nucleic acid sequence encoding amino acid sequence for 2086 protein from a strain of *Neisseria lactamica*.

SEQ ID NOS:254 to 259 amino acid sequences associated with proteins of 2086 family of proteins.

- 15 SEQ ID NOS:260 to 278 amino acid sequences associated with proteins of 2086 Subfamily A.

SEQ ID NOS:279 to 299 amino acid sequences associated with proteins of 2086 Subfamily B.

SEQ ID NO:300 is the amino acid consensus sequence corresponding to the 2086 protein family ("2086 proteins") according to an embodiment of the present invention.

- 20 SEQ ID NO:301 is the amino acid consensus sequence corresponding to the 2086 protein Subfamily A according to an embodiment of the present invention.

SEQ ID NO:302 is the amino acid consensus sequence corresponding to the 2086 protein Subfamily B according to an embodiment of the present invention.

WO 03/063766

PCT/US02/32369

-29-

SEQ ID NO:303 nucleic acid sequence for a reverse primer with BamHI restriction site (Compound No. 4623).

SEQ ID NO:304 nucleic acid sequence for a forward primer with NdeI restriction site (Compound No. 4624).

5 SEQ ID NO:305 nucleic acid sequence for a forward primer (Compound No. 4625).

SEQ ID NO:306 nucleic acid sequence for a forward primer (Compound No. 5005).

SEQ ID NO:307 nucleic acid sequence for a reverse primer (Compound No. 5007).

SEQ ID NO:308 nucleic acid sequence for a reverse primer with BglII restriction site (Compound No. 5135).

10 SEQ ID NO:309 nucleic acid sequence for a forward primer with BamHI restriction site (Compound No. 5658).

SEQ ID NO:310 nucleic acid sequence for a reverse primer with SphI restriction site (Compound No. 5660).

15 SEQ ID NO:311 nucleic acid sequence for a forward primer with BamHI restriction site (Compound No. 6385).

SEQ ID NO:312 nucleic acid sequence for a forward primer with BglII and NdeI restriction sites (Compound No. 6406).

SEQ ID NO:313 nucleic acid sequence for a forward primer (Compound No. 6470).

SEQ ID NO:314 nucleic acid sequence for a reverse primer (Compound No. 6472).

20 SEQ ID NO:315 nucleic acid sequence for a forward primer with BamHI restriction site (Compound 6473).

SEQ ID NO:316 nucleic acid sequence for a forward primer with BglII and NdeI restriction sites (Compound No. 6474).

-30-

SEQ ID NO:317 nucleic acid sequence for a forward primer (Compound No. 6495).

SEQ ID NO:318 nucleic acid sequence for a reverse primer (Compound No. 6496).

SEQ ID NO:319 nucleic acid sequence for a reverse primer with SphI restriction site (Compound No. 6543).

- 5 SEQ ID NO:320 nucleic acid sequence for a reverse primer with BglII restriction site (Compound No. 6605).

SEQ ID NO:321 nucleic acid sequence for a forward primer with BglII and NdeI restriction sites (Compound No. 6721).

SEQ ID NO:322 nucleic acid sequence for the P4 leader sequence.

- 10 SEQ ID NO:323 nucleic acid sequence for native 2086 leader variant 1.

SEQ ID NO:324 nucleic acid sequence for native 2086 leader variant 2.

SEQ ID NO:325 nucleic acid sequence for native 2086 leader variant 3.

SEQ ID NO:326 nucleic acid sequence for native 2086 leader variant 4.

SEQ ID NO:327 is the amino acid sequence of P4431.

- 15 SEQ ID NO:328 is the amino acid sequence of P5163.

SEQ ID NO:329 is an amino acid sequence according to an embodiment of the present invention.

Detailed Description of the Invention

- 20 A new class of antigens with cross-functional bactericidal activity against *Neisseria meningitidis* serogroup B may potentially obviate the need for a multi-valent PorA approach to immunization against infection. A candidate antigen has been unexpectedly identified and is described and claimed herein. The presence of one such candidate antigen was first observed in a complex mixture of soluble outer

-31-

membrane proteins (sOMPs) from a meningococcal strain. The bactericidal activity of this antigen was followed through a series of fractionation and protein purification steps until the protein mixture of interest contained just a few proteins. The major proteins in this mixture were identified by N-terminal amino acid sequencing and peptide mapping. The protein of interest exhibiting bactericidal activity was identified as ORF2086 protein, a lipidated protein (also more specifically referred to as LP2086). "ORF2086 protein" refers to a protein encoded by open reading frame 2086 (ORF2086) of *Neisseria species*.

As described herein, new antigen candidates based on *Neisseria species* ORF2086 protein (also referred to as "2086 protein" or "ORF2086" protein, used interchangeably herein, or P2086 for the non-lipated proteins and LP2086 for the lipidated version of the proteins) isolated from *N. meningitidis* were identified by combining cell fractionation, differential detergent extraction, protein purification, with the preparation of antisera, and a bactericidal activity assay utilizing multiple strains.

According to an embodiment of the present invention, immunogenic agents based on 2086 protein, including isolated polypeptides, immunogenic portions thereof and/or biological equivalents thereof were unexpectedly identified as immunogenic candidates based on the ability of said agents to exhibit cross-reactivity or non-strain specificity. In particular, candidates were identified that in some embodiments have the ability to elicit antibodies that (1) are bactericidal to multiple neisserial and/or gonococcal strains; (2) react with the surface of multiple strains; (3) confer passive protection against a live challenge; and/or (4) prevent colonization. Accordingly, in some embodiments, the present invention provides immunogenic compositions comprising said immunogenic agents, including isolated polypeptides, immunogenic portions thereof, and/or biological equivalents thereof. (See Example 1 herein for the methodology used in the identification of the 2086 protein.)

As used herein, the term "non-strain specific" refers to the characteristic of an antigen to elicit an immune response against more than one strain of *N. meningitidis* (e.g., heterologous meningococcal strains). The term "cross-reactive" as it is used

-32-

herein is used interchangeably with the term "non-strain specific". The term "immunogenic non-strain specific *N. meningitidis* antigen," as used herein, describes an antigen that can be isolated from *N. meningitidis*, although it can also be isolated from another bacterium (e.g., other neisserial strains, such as gonococcal strains, for example), or prepared using recombinant technology.

The 2086 proteins of the present invention include lipidated and non-lipidated proteins. Further, the present invention also contemplates the use of the immature proteins or preproteins that correspond to each protein as intermediate compounds/compositions.

The present invention also provides antibodies that immunospecifically bind to the foregoing immunogenic agents, according to some embodiments of the invention. Further, the present invention relates to isolated polynucleotides comprising nucleic acid sequences encoding any of the foregoing. Additionally, the present invention provides compositions and/or immunogenic compositions, as well as methods for preparing said compositions.

Embodiments of compositions disclosed herein have been shown to be highly immunogenic and capable of eliciting the production of bactericidal antibodies. In some embodiments, these antibodies are cross-reactive to serogroup, serotype and serosubtype heterologous meningococcal strains. Accordingly, the present compositions may overcome deficiencies of previous *N. meningitidis* vaccine attempts by exhibiting the ability to elicit bactericidal antibodies to heterologous neisserial strains. Thus, the present invention may provide immunogenic compositions that can be compounded with fewer components with the aim to potentially elicit protection comparable to previously used agents. The compositions or immunogenic agents therein (e.g., polypeptides, immunogenic portions or fragments, and biological equivalents, etc., without limitation) may be used alone or in combination with other antigens or agents to provide immunogenicity against meningococcal infection, as well as potentially elicit immunological protection from infection caused by other pathogens. This may simplify the design of an immunogenic composition for use against meningococcal infection by potentially reducing the number of antigens

-33-

required for protection against multiple strains. In fact, the disclosure herein suggests that purified 2086 protein may reduce the number of proteins required to provide adequate immunogenic coverage of the strains responsible for meningococcal disease. The 2086 protein can be recombinantly expressed in *E. coli* as a lipoprotein, which is the wild type form of the protein, at levels much higher than in the native meningococci.

Because at least some embodiments of the antibodies directed against the 2086 protein from a single strain were found to kill unrelated (i.e., heterologous) strains, an attempt was made to characterize a large number of heterologous strains for the presence of a "2086 homolog", and to determine the level of sequence conservation. While about 70% of the strains tested by PCR had 2086 homologs that could be amplified using the primers that amplified the original 2086 gene from strain 8529, the remaining approximately 30% were negative by this test. These remaining approximately 30% were found to contain a 2086 homolog that has only about 60% amino acid sequence homology to the original 8529 derived 2086 gene. Other primers were identified that could amplify a 2086 homolog from these approximately 30% of strains. The *N. meningitidis* strains tested have been designated as belonging to Subfamily A or Subfamily B depending on which primer set can amplify a 2086 homolog. The details of these experiments are outlined in Example 5 below.

20 The presence of a 2086 protein in numerous serosubtypes.

To determine the level of sequence conservation of the 2086 gene between *N. meningitidis* strains, several representatives from Subfamilies A and B were cloned as full length genes and submitted for DNA sequence analysis. Using primers as disclosed herein, see, for example, Table IV, twenty four serogroup B meningococcal strains were identified, which express different serosubtype antigens and also express a shared protein, P2086. Examples of these sequences are provided herein and are shown as mature DNA sequences (i.e., all lipoprotein signal sequences have been cleaved at the cysteine residue). See, for example, the amino acid sequences of even numbered SEQ ID NOS: 2-252, without limitation.

-34-

Although the 2086 protein is not present in large amounts in wild type strains, it is a target for bactericidal antibodies. These antibodies, unlike those produced in response to the PorAs, are capable of killing strains expressing heterologous serosubtypes.

5 At least some embodiments of the antibodies to the 2086 protein also passively protect infant rats from challenge with meningococci. (see Table VII) Recombinant expression of 2086 protein thus may enable the use of 2086 protein as an immunogenic composition for the prevention of meningococcal disease. All of the recent meningococcal immunogenic composition candidates in clinical trials have
10 been complex mixtures or outer membrane protein preparations containing many different proteins. The PorA protein, that provides serosubtype specificity, will require the inclusion of 6 to 9 variants in an immunogenic composition to provide about 70-80% coverage of disease related serosubtypes. In contrast, in embodiments demonstrated herein, antisera to a single 2086 protein alone is able to kill
15 representatives of six serosubtypes responsible for about 65% of the disease isolates in western Europe and the United States. Therefore, purified 2086 protein has the potential to reduce the number of proteins required to provide adequate immunogenic composition coverage of the serosubtypes responsible for meningococcal disease.

Proteins, Immunogenic Portions and Biological Equivalents

20 The 2086 proteins provided by the present invention are isolated proteins. The term "isolated" means altered by the hand of man from the natural state. If an "isolated" composition or substance occurs in nature, it has been changed or removed from its original environment, or both. For example, a polypeptide or a
25

-35-

polynucleotide naturally present in a living animal is not “isolated,” but the same polypeptide or polynucleotide separated from the coexisting materials of its natural state is “isolated”, as the term is employed herein. Accordingly, as used herein, the term “isolated protein” encompasses proteins isolated from a natural source and
5 proteins prepared using recombinant technology, as well as such proteins when combined with other antigens and/or additives, such as pharmaceutically acceptable carriers, buffers, adjuvants, etc., for example.

A 2086 protein, immunogenic portion thereof and/or a biological equivalent thereof, according an embodiment of the invention, comprises any of the following
10 amino acid sequences:

ADIGxGLADA (SEQ ID NO:254), wherein x is any amino acid;

IGxGLADALT (SEQ ID NO:255), wherein x is any amino acid;

SLNTGKLKND (SEQ ID NO:256);

SLNTGKLKNDKxSRFDF (SEQ ID NO:257, wherein x is any amino acid);

15 SGEFQxYKQ (SEQ ID NO:258), wherein x is any amino acid; or

IEHLKxPE (SEQ ID NO:259), wherein x is any amino acid.

A 2086 Subfamily A protein, immunogenic portion thereof and/or biological equivalent thereof comprises any of the following amino acid sequences, in accordance with an embodiment of the present invention:

20 GGGVAADIGx (SEQ ID NO:260), wherein x is any amino acid;

SGEFQIYKQ (SEQ ID NO:261);

HSAVVALQIE (SEQ ID NO:262);

EKINNPDKID (SEQ ID NO:263);

SLINQRSFLV (SEQ ID NO:264);

-36-

SGLGGEHTAF (SEQ ID NO:265);

GEHTAFNQLP (SEQ ID NO:266);

SFLVSGLGGEH (SEQ ID NO:267);

EKINNPDKIDSLINQRSFLVSGLGGEHTAFNQLP (SEQ ID NO:268);

5 GKAEYHGKAF (SEQ ID NO:269);

YHGKAFSSDD (SEQ ID NO:270);

GKAEYHGKAFSSDD (SEQ ID NO:271);

IEHLKTPEQN (SEQ ID NO: 272);

KTPEQNVELA (SEQ ID NO:273);

10 IEHLKTPEQNVELA (SEQ ID NO:274);

AELKADEKSH (SEQ ID NO:275);

AVILGDTRYG (SEQ ID NO:276);

AELKADEKSHAVILGDTRYG (SEQ ID NO:277); or

EEKGTYHLAL (SEQ ID NO:278).

15 A 2086 Subfamily B protein, immunogenic portion thereof and/or biological equivalent thereof comprises any of the following amino acid sequences, in accordance with an embodiment of the present invention:

LITLESGEFQ (SEQ ID NO:279);

SALTALQTEQ (SEQ ID NO:280);

20 FQVYKQSHSA (SEQ ID NO:281);

LITLESGEFQVYKQSHSALTALQTEQ (SEQ ID NO:282);

-37-

IGDLAGEHTS (SEQ ID NO:283);

EHTSFDKLPK (SEQ ID NO:284);

IGDLAGEHTSFDKLPK (SEQ ID NO:285);

ATYRGTAFGS (SEQ ID NO:286);

5 DDAGGKLTYT (SEQ ID NO:287);

IDFAAKQGHG (SEQ ID NO:288);

KIEHLKSPEL (SEQ ID NO:289);

ATYRGTAFGSDDAGGKLTYTIDFAAKQGHGKIEHLKSPELNV (SEQ ID
NO: 290);

10 HAVISGSVLY (SEQ ID NO:291);

KGSYSLGIFG (SEQ ID NO:292);

VLYNQDEKGS (SEQ ID NO:293);

HAVISGSVLYNQDEKGSYSLGIFG (SEQ ID NO:294);

AQEVAGSAEV (SEQ ID NO:295);

15 IHHIGLAAKQ (SEQ ID NO:296);

VETANGIHHI (SEQ ID NO:297);

AQEVAGSAEVETANGIHHIGLAAKQ (SEQ ID NO:298); or

VAGSAEVETANGIHHIGLAAKQ (SEQ ID NO:299).

The 2086 protein comprises the following consensus sequence and/or
20 immunogenic portions thereof in accordance with an embodiment of the present
invention.

-38-

2086 Protein Consensus Sequence (SEQ ID NO: 300):

CSSG-----GGGVxADIGxGLADALTxPx DxKDKxLxSLTLxxSxxxNxxLxLxAQGA
 EKTxxxGD---SLNTGKLKNDKxSRFDFxxxIxVDGxxITLxSGEFQxYKQxHSAxx
 ALQxExxxxxxxxxxxxxRxFxxxxxGEHTxFxxLPxx-xAxYxGxAFxSDDxxGxLxYx
 5 IDExxKQGxGxIEHLKxPExNVxLAXxxxKxDIEKxHAVIxGxxxYxxxFKGxYxLxxx
 GxxAQFxAQxAvxxxxxxHxLxxAxKQ

In the foregoing consensus sequence, the "x" represents any amino acid. the
 region from amino acid position 5 to amino acid position 9 is any of 0 to 5 amino
 acids, the region from amino acid position 67 to amino acid position 69 is any of 0 to
 10 3 amino acids, and amino acid position 156 is any of 0 to 1 amino acid. The region
 from amino acid position 5 to amino acid position 9 preferably comprises 0, 4 or 5
 amino acids. The region from amino acid position 67 to amino acid position 69
 preferably comprises 0 or 3 amino acids. It should be particularly noted that this
 consensus sequence illustrates the high variability of the 2086 proteins. By way of
 15 theory, without intending to be bound thereto, it is believed that this high variability
 provides the advantageous and unexpected cross-reactivity.

According to some embodiments of the present invention, the 2086 proteins
 are characterized as being immunogenic, nonpathogenic and non-strain specific.
 Moreover, these proteins unexpectedly exhibit immunogenicity while being about 2%
 20 to about 40% nonconserved.

As used herein, the term "nonconserved" refers to the number of amino acids
 that may undergo insertions, substitution and/or deletions as a percentage of the total
 number of amino acids in a protein. For example, if a protein is 40% nonconserved
 and has, for example, 263 amino acids, then there are 105 amino acid positions in the
 25 protein at which amino acids may undergo substitution. Likewise, if a protein is 10%
 nonconserved and has, for example, about 280 amino acids, then there are 28 amino
 acid positions at which amino acids may undergo substitution. The 2086 proteins
 may also undergo deletion of amino acid residues, potentially without compromising
 the immunogenicity of the proteins.

-39-

Further, the 2086 proteins may be divided into subfamilies based upon homology at various regions. For example, without intending to be limited thereto, the consensus sequences for two such subfamilies, Subfamily A and Subfamily B, are provided below:

5

2086 Subfamily A sequence (SEQ ID 301)

CSSG---GGGVAADIGxGLADALTxPx DxKDKxLxSLTLxxSxxxNxxLxLxAQGA
 EKTxxxGD---SLNTGKLKNDKxSRFDFxxxLxVDGQxITLxSGEFQIYKQxHSAVV
 10 ALQIEKINNPDKIDSLINQRSFLVSGLGGEHTAFNQLPxGKAEYHGKAFSSDDx
 xGxLxYxIDFxxKQGxGxIEHLKTPEQNVELAxAELKADEKSHAVILGDTRYGxE
 EKGTYHLALxGDRAQEIAGxATVKIxEKVHEIxIAxKQ

The reference "x" is any amino acid.

15 The region from amino acid position 5 to amino acid position 8 is any of 0 to 4 amino acids.

The region from amino acid position 66 to amino acid position 68 is any of 0 to 3 amino acids.

20 The region from amino acid position 5 to amino acid position 8 preferably comprises 0 or 4 amino acids. The region from amino acid position 66 to amino acid position 68 preferably comprises 0 or 3 amino acids.

2086 Subfamily B (SEQ ID 302)

25 CSSGGGG-----VxADIGxGLADALTAPLDHKDKxLxSLTLxxSxxxNxxLxLxAQ
 GAECTYGNGLSLNTGKLKNDKVSRLFDFIRQIEVDGxLITLESGEFQVYKQSHS
 ALTALQTEQxQDxExSxKMVAKRxFxIGDIAGEHTSFDKLPKxxxATYRGTAFGS
 DDAGGKLTYTIDFAAKQGHGKIEHLKSPELNVxLAXxYIKPDEKxHAVISGSVL
 YNQDEKGSYSLGIFGxxAQEVAGSAEVETANGIHHIGLAAKQ

30 The reference "x" is any amino acid.

-40-

The region from amino acid position 8 to amino acid position 12 is any of 0 to 5 amino acids.

The region from amino acid position 8 to amino acid position 12 preferably comprises 0 or 5 amino acids.

5 According to implementations of the present invention, the 2086 protein subfamilies may be further subdivided into clusters. For example, according to an implementation of the present invention, the following clusters are provided: even numbered SEQ ID NOS:2-12; even numbered SEQ ID NOS:14-24; even numbered SEQ ID NOS:26-42; even numbered SEQ ID NOS:50-60; even numbered SEQ ID NOS:62-108; even numbered SEQ ID NOS:110-138; even numbered SEQ ID NOS:140-156; even numbered SEQ ID NOS:158-174; and even numbered SEQ ID NOS: 224-252.

15 A polypeptide sequence of the invention may be identical to the reference sequence of even numbered SEQ ID NOS: 2-252, that is, 100% identical, or it may include a number of amino acid alterations as compared to the reference sequence such that the % identity is less than 100%. Such alterations include at least one amino acid deletion, substitution, including conservative and non-conservative substitution, or insertion. The alterations may occur at the amino- or carboxy-terminal positions of the reference polypeptide sequence or anywhere between those terminal positions, interspersed either individually among the amino acids in the reference amino acid sequence or in one or more contiguous groups within the reference amino acid sequence.

25 Thus, the invention also provides proteins having sequence identity to the amino acid sequences contained in the Sequence Listing (i.e., even numbered SEQ ID NOS: 2-252). Depending on the particular sequence, the degree of sequence identity is preferably greater than 60% (e.g., 60%, 70%, 80%, 90%, 95%, 97%, 99%, 99.9% or more). These homologous proteins include mutants and allelic variants.

-41-

In preferred embodiments of the invention, the 2086 proteins or other 2086 polypeptides (e.g., immunological portions and biological equivalents) generate bactericidal antibodies to homologous and at least one heterologous strain of meningococci. Specifically, embodiments of the antibodies to the 2086 polypeptides
5 may passively protect infant rats from challenge, such as intranasal, with meningococci. In further preferred embodiments, the 2086 polypeptides may exhibit such protection for infants rats for homologous strains and at least one heterologous strain. The polypeptide may be selected from the Sequence Summary above, as set forth in the even numbered SEQ ID NOS: 2-252, or the polypeptide may be any
10 immunological fragment or biological equivalent of the listed polypeptides. Preferably, the polypeptide is selected from any of the even numbered SEQ ID NOS: 2-252 in the Sequence Summary above.

This invention also relates to allelic or other variants of the 2086 polypeptides, which are biological equivalents. Suitable biological equivalents may in some
15 embodiments have the ability to elicit antibodies that (1) are bactericidal to homologous strains and at least one heterologous neisserial strain and/or gonococcal strain; (2) react with the surface of homologous strains and at least one heterologous neisserial and/or gonococcal strain; (3) confer passive protection against a live challenge; and/or (4) prevent colonization.

Potentially suitable biological equivalents have at least about 60%, preferably at least about 70%, more preferably at least about 75%, even more preferably about 80%, even more preferably about 85%, even more preferably about 90%, even more preferably 95 % or even more preferably 98%, or even more preferably 99%
20 similarity to one of the 2086 polypeptides specified herein (i.e., the even numbered SEQ ID NOS: 2-252), provided the equivalent is capable of eliciting substantially the same immunogenic properties as one of the 2086 proteins of this invention.
25

Alternatively, the biological equivalents have substantially the same immunogenic properties of one of the 2086 proteins in the even numbered SEQ ID NOS: 2-252. According to some embodiments of the present invention, the biological

-42-

equivalents have the same immunogenic properties as the even numbered SEQ ID NOS 2-252.

The biological equivalents are obtained by generating variants and modifications to the proteins of this invention. These variants and modifications to the proteins are obtained by altering the amino acid sequences by insertion, deletion or substitution of one or more amino acids. The amino acid sequence is modified, for example by substitution in order to potentially create a polypeptide having substantially the same or improved qualities. A preferred means of introducing alterations comprises making predetermined mutations of the nucleic acid sequence of the polypeptide by site-directed mutagenesis.

Modifications and changes may be made in the structure of a polypeptide of the present invention and potentially still obtain a molecule having *N. meningitidis* immunogenicity. For example, without limitation, it is contemplated that certain amino acids can be substituted for other amino acids, including nonconserved and conserved substitution, in a sequence potentially without appreciable loss of immunogenicity. Because it is the interactive capacity and nature of a polypeptide that defines that polypeptide's biological functional activity, it is contemplated that a number of amino acid sequence substitutions can be made in a polypeptide sequence (or, of course, its underlying DNA coding sequence) and nevertheless potentially obtain a polypeptide with like properties. The present invention contemplates any changes to the structure of the polypeptides herein, as well as the nucleic acid sequences encoding said polypeptides, wherein the polypeptide retains immunogenicity. A person of ordinary skill in the art would be readily able to modify the disclosed polypeptides and polynucleotides accordingly, based upon the guidance provided herein.

For example, certain variable regions have been identified where substitution or deletion may be permissible. The 2086 consensus sequence, as previously discussed, shows conserved and nonconserved regions of the 2086 family of proteins according to an embodiment of the present invention.

In making such changes, any techniques known to persons of skill in the art may be utilized. For example, without intending to be limited thereto, the hydrophobic

-43-

index of amino acids can be considered. The importance of the hydropathic amino acid index in conferring interactive biologic function on a polypeptide is generally understood in the art. Kyte *et al.* 1982. *J. Mol. Bio.* 157:105-132.

Substitution of like amino acids can also be made on the basis of hydrophilicity, particularly where the biological functional equivalent polypeptide or peptide thereby created is intended for use in immunological embodiments. U.S. Pat. No. 4,554,101, states that the greatest local average hydrophilicity of a polypeptide, as governed by the hydrophilicity of its adjacent amino acids, correlates with its immunogenicity, *i.e.* with a biological property of the polypeptide.

Biological equivalents of a polypeptide can also be prepared using site-specific mutagenesis. Site-specific mutagenesis is a technique useful in the preparation of second generation polypeptides, or biologically functional equivalent polypeptides or peptides, derived from the sequences thereof, through specific mutagenesis of the underlying DNA. Such changes can be desirable where amino acid substitutions are desirable. The technique further provides a ready ability to prepare and test sequence variants, for example, incorporating one or more of the foregoing considerations, by introducing one or more nucleotide sequence changes into the DNA. Site-specific mutagenesis allows the production of mutants through the use of specific oligonucleotide sequences which encode the DNA sequence of the desired mutation, as well as a sufficient number of adjacent nucleotides, to provide a primer sequence of sufficient size and sequence complexity to form a stable duplex on both sides of the deletion junction being traversed. Typically, a primer of about 17 to 25 nucleotides in length is preferred, with about 5 to 10 residues on both sides of the junction of the sequence being altered.

In general, the technique of site-specific mutagenesis is well known in the art. As will be appreciated, the technique typically employs a phage vector which can exist in both a single stranded and double stranded form. Typically, site-directed mutagenesis in accordance herewith is performed by first obtaining a single-stranded vector which includes within its sequence a DNA sequence which encodes all or a portion of the *N. meningitidis* polypeptide sequence selected. An oligonucleotide

-44-

primer bearing the desired mutated sequence is prepared (*e.g.*, synthetically). This primer is then annealed to the single-stranded vector, and extended by the use of enzymes such as *E. coli* polymerase I Klenow fragment, in order to complete the synthesis of the mutation-bearing strand. Thus, a heteroduplex is formed wherein one strand encodes the original non-mutated sequence and the second strand bears the desired mutation. This heteroduplex vector is then used to transform appropriate cells such as *E. coli* cells and clones are selected which include recombinant vectors bearing the mutation. Commercially available kits come with all the reagents necessary, except the oligonucleotide primers.

2086 polypeptides include any protein or polypeptide comprising substantial sequence similarity and/or biological equivalence to a 2086 protein having an amino acid sequence from one of the even numbered SEQ ID NOS 2-252. In addition, a 2086 polypeptide of the invention is not limited to a particular source. Thus, the invention provides for the general detection and isolation of the polypeptides from a variety of sources. Also, the 2086 polypeptides can be prepared recombinantly, as is well within the skill in the art, based upon the guidance provided herein, or in any other synthetic manner, as known in the art.

It is contemplated in the present invention, that a 2086 polypeptide may advantageously be cleaved into fragments for use in further structural or functional analysis, or in the generation of reagents such as 2086-related polypeptides and 2086-specific antibodies. This can be accomplished by treating purified or unpurified *N. meningitidis* polypeptides with a peptidase such as endoproteinase glu-C (Boehringer, Indianapolis, IN). Treatment with CNBr is another method by which peptide fragments may be produced from natural *N. meningitidis* 2086 polypeptides. Recombinant techniques also can be used to produce specific fragments of a 2086 protein.

“Variant” as the term is used herein, is a polynucleotide or polypeptide that differs from a reference polynucleotide or polypeptide respectively, but retains essential properties. A typical variant of a polynucleotide differs in nucleotide sequence from another, reference polynucleotide. Changes in the nucleotide sequence

-45-

of the variant may or may not alter the amino acid sequence of a polypeptide encoded by the reference polynucleotide. Nucleotide changes may result in amino acid substitutions, additions, deletions, fusions and truncations in the polypeptide encoded by the reference sequence, as discussed below. A typical variant of a polypeptide
5 differs in amino acid sequence from another, reference polypeptide. Generally, differences are limited so that the sequences of the reference polypeptide and the variant are closely similar overall and, in many regions, identical (*i.e.*, biologically equivalent). A variant and reference polypeptide may differ in amino acid sequence by one or more substitutions, additions, deletions in any combination. A substituted
10 or inserted amino acid residue may or may not be one encoded by the genetic code. A variant of a polynucleotide or polypeptide may be a naturally occurring such as an allelic variant, or it may be a variant that is not known to occur naturally. Non-naturally occurring variants of polynucleotides and polypeptides may be made by mutagenesis techniques or by direct synthesis.

15 “Identity,” as known in the art, is a relationship between two or more polypeptide sequences or two or more polynucleotide sequences, as determined by comparing the sequences. In the art, “identity” also means the degree of sequence relatedness between polypeptide or polynucleotide sequences, as the case may be, as determined by the match between strings of such sequences. “Identity” and
20 “similarity” can be readily calculated by known methods, including but not limited to those described in Computational Molecular Biology, Lesk, A. M., ed., Oxford University Press, New York, 1988; Biocomputing: Informatics and Genome Projects, Smith, D. W., ed., Academic Press, New York, 1993; Computer Analysis of Sequence Data, Part I, Griffin, A. M., and Griffin, H. G., eds., Humana Press, New Jersey,
25 1994; Sequence Analysis in Molecular Biology, von Heinje, G., Academic Press, 1987; and Sequence Analysis Primer, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991; and Carillo, H., and Lipman, D., *SIAM J. Applied Math.*, 48:1073 (1988). Preferred methods to determine identity are designed to give the largest match between the sequences tested. Methods to determine identity and
30 similarity are codified in publicly available computer programs. Preferred computer program methods to determine identity and similarity between two sequences include,

-46-

but are not limited to, the GCG program package (Devereux, J., *et al* 1984), BLASTP, BLASTN, and FASTA (Altschul, S. F., *et al.*, 1990). The BLASTX program is publicly available from NCBI and other sources (BLAST Manual, Altschul, S., *et al.*, NCBI NLM NIH Bethesda, Md. 20894; Altschul, S., *et al.*, 1990). The well known
 5 Smith Waterman algorithm may also be used to determine identity.

By way of example, without intending to be limited thereto, an amino acid sequence of the present invention may be identical to the reference sequences, even numbered SEQ ID NOS: 2-252; that is be 100% identical, or it may include a number of amino acid alterations as compared to the reference sequence such that the %
 10 identity is less than 100%. Such alterations are selected from the group consisting of at least one amino acid deletion, substitution, including conservative and non-conservative substitution, or insertion, and wherein said alterations may occur at the amino- or carboxy-terminal positions of the reference polypeptide sequence or anywhere between those terminal positions, interspersed either individually among
 15 the amino acids in the reference sequence or in one or more contiguous groups within the reference sequence. The number of amino acid alterations for a given % identity is determined by multiplying the total number of amino acids in SEQ ID NOS:2-252 by the numerical percent of the respective percent identity (divided by 100) and then subtracting that product from said total number of amino acids in any of SEQ ID
 20 NOS:2-252, or:

$$n_a = x_a - (x_a \cdot y),$$

wherein n_a is the number of amino acid alterations, x_a is the total number of amino acids in SEQ ID NOS:2-252, and y is, for instance 0.70 for 70%, 0.80 for 80%, 0.85 for 85% etc., and wherein any non-integer product of $x_{\text{sub.a}}$ and y is rounded down
 25 to the nearest integer prior to subtracting it from x_a .

In preferred embodiments, the polypeptide above is selected from the proteins set forth in the even numbered SEQ ID NOS 2-252, such as mature processed form of a 2086 protein. The 2086 proteins or equivalents, etc. may be lipidated or non-lipidated.

-47-

ORF 2086 is expressible in *E. coli* with the native ORF 2086 signal sequence. However, it is desirable to find means to improve the expression of proteins. According to an embodiment of the present invention, a leader sequence produces a lipidated form of the protein. For example, the following describes the use of the
5 signal sequence of the nontypable *Haemophilus influenzae* P4 protein to enhance expression.

The processing of bacterial lipoproteins begins with the synthesis of a precursor or prolipoprotein containing a signal sequence, which in turn contains a consensus lipoprotein processing/modification site. This prolipoprotein initially
10 passes through the common Sec system on the inner membrane of Gram negative bacteria or on the membrane in Gram positive bacteria. Once placed in the membrane by the Sec system, the prolipoprotein is cleaved by signal peptidase II at the consensus site and the exposed N-terminal cysteine residue is glycerated and acylated. Hayashi et al. 1990. Lipoproteins in bacteria. *J. Bioenerg. Biomembr.* Jun;
15 22(3):451-71; Oudega et al. 1993. *Escherichia coli* SecB, SecA, and SecY proteins are required for expression and membrane insertion of the bacteriocin release protein, a small lipoprotein. *J. Bacteriol.* Mar;175(5):1543-7; Sankaran et al. 1995. Modification of bacterial lipoproteins. *Methods Enzymol.* 250:683-97.

In Gram negative bacteria, transport of the lipidated protein to the outer
20 membrane is mediated by a unique ABC transporter system with membrane specificity depending on a sorting signal at position 2 of the lipoprotein. Yakushi et al. 2000. A new ABC transporter mediating the detachment of lipid modified proteins from membranes. *Nat Cell Biol.* Apr;2(4):212-8.

Fusion with bacterial lipoproteins and their signal sequences has been used to
25 display recombinant proteins on the surface of bacteria. U.S. Patent Numbers 5,583,038 and 6,130,085. Exchanging lipoprotein signal sequences can increase the production of the lipoprotein. De et al. 2000. Purification and characterization of *Streptococcus pneumoniae* palmitoylated pneumococcal surface adhesin A expressed in *Escherichia coli*. *Vaccine.* Mar 6;18(17):1811-21.

Bacterial lipidation of proteins is known to increase or modify the immunological response to proteins. Erdile et al. 1993. Role of attached lipid in immunogenicity of *Borrelia burgdorferi* OspA. *Infect. Immun.* Jan;61(1):81-90; Snapper et al. 1995. Bacterial lipoproteins may substitute for cytokines in the humoral immune response to T cell-independent type II antigens. *J. Immunol.* Dec 15;155(12):5582-9. However, bacterial lipoprotein expression can be complicated by the stringency of the processing. Pollitt et al. 1986. Effect of amino acid substitutions at the signal peptide cleavage site of the *Escherichia coli* major outer membrane lipoprotein. *J. Biol. Chem.* Feb 5; 261(4):1835-7; Lunn et al. 1987. Effects of prolipoprotein signal peptide mutations on secretion of hybrid prolipo-beta-lactamase in *Escherichia coli*. *J. Biol. Chem.* Jun 15;262(17):8318-24; Klein et al. 1988. Distinctive properties of signal sequences from bacterial lipoproteins. *Protein Eng.* Apr; 2(1):15-20. Bacterial lipoprotein expression is also complicated by other problems such as toxicity and low expression levels. Gomez et al. 1994. Nucleotide The *Bacillus subtilis* lipoprotein LplA causes cell lysis when expressed in *Escherichia coli*. *Microbiology.* Aug;140 (Pt 8):1839-45; Hansson et al. 1995. Expression of truncated and full-length forms of the Lyme disease *Borrelia* outer surface protein A in *Escherichia coli*. *Protein Expr. Purif.* Feb; 6(1):15-24; Yakushi et al. 1997. Lethality of the covalent linkage between mislocalized major outer membrane lipoprotein and the peptidoglycan of *Escherichia coli*. *J. Bacteriol.* May; 179(9):2857-62.

The nontypable *Haemophilus influenzae* bacterium expresses a lipoprotein designated P4 (also known as protein "c"). The recombinant form of the P4 protein is highly expressed in *E. coli* using the native P4 signal sequence. U.S. Patent Number 5,955,580. When the native P4 signal sequence is substituted for the native ORF 2086 signal sequence in an expression vector in *E. coli*, the level of expression of ORF2086 is increased.

This concept of using the heterologous P4 signal sequence to increase expression is extendible to other bacterial lipoproteins. In particular, analysis of bacterial genomes leads to the identification of many ORFs as being of possible

-49-

interest. Attempting to express each ORF with its native signal sequence in a heterologous host cell, such as *E. coli*, may give rise to a variety of problems inherent in using a variety of signal sequences, including stability, compatibility and so forth. It may be possible to minimize these problems, by using the P4 signal sequence to
5 express each ORF of interest. As described above, the P4 signal sequence improves the expression of the heterologous 2086 ORF. An expression vector is constructed by deleting the native signal sequence of the ORF of interest, and ligating the P4 signal sequence to the ORF. A suitable host cell is then transformed, transfected or infected with the expression vector, and expression of the ORF is increased in comparison to
10 expression using the native signal sequence of the ORF.

The non-lipidated form is produced by a protein lacking the original leader sequence or a by a leader sequence which is replaced with a portion of sequence that does not specify a site for fatty acid acylation in a host cell.

The various forms of the 2086 proteins of this invention are referred to herein
15 as "2086" protein, unless otherwise specifically noted. Also "2086 polypeptide" refers to the 2086 proteins as well as immunogenic portions or biological equivalents thereof as noted above, unless otherwise noted.

The full length isolated and purified *N. meningitidis* 2086 protein has an apparent molecular weight of about 28 to 35 kDa as measured on a 10 % to 20%
20 gradient SDS polyacrylamide gel (SDS-PAGE). More specifically, this protein has a molecular weight of about 26,000 to 30,000 daltons as measured by mass spectrometry.

Antibodies

The proteins of the invention, including the amino acid sequences of SEQ ID
25 NOS: 2-252, their fragments, and analogs thereof, or cells expressing them, may also be used as immunogens to produce antibodies immunospecific for the polypeptides of the invention. In some embodiments, the invention includes antibodies to immunospecific polypeptides and the use of such antibodies to potentially detect the

-50-

presence of *N. meningitidis*, provide passive protection or measure the quantity or concentration of the polypeptides in a cell, a cell or tissue extract, or a biological fluid.

The antibodies of the invention may include polyclonal antibodies,
 5 monoclonal antibodies, chimeric antibodies, and anti-idiotypic antibodies. Polyclonal antibodies are heterogeneous populations of antibody molecules derived from the sera of animals immunized with an antigen. Monoclonal antibodies are a substantially homogeneous population of antibodies to specific antigens. Monoclonal antibodies may be obtained by methods known to those skilled in the art, e.g., Kohler and
 10 Milstein, 1975, *Nature* 256:495-497 and U.S. Patent Number 4,376,110. Such antibodies may be of any immunoglobulin class including IgG, IgM, IgE, IgA, GILD and any subclass thereof.

Chimeric antibodies are molecules, different portions of which are derived from different animal species, such as those having variable region derived from a
 15 murine monoclonal antibody and a human immunoglobulin constant region. Chimeric antibodies and methods for their production are known in the art (Cabilly *et al.*, 1984, *Proc. Natl. Acad. Sci. USA* 81:3273-3277; Morrison *et al.*, 1984, *Proc. Natl. Acad. Sci. USA* 81:6851-6855; Boulianne *et al.*, 1984, *Nature* 312:643-646; Cabilly *et al.*, European Patent Application 125023 (published November 14, 1984);
 20 Taniguchi *et al.*, European Patent Application 171496 (published February 19, 1985); Morrison *et al.*, European Patent Application 173494 (published March 5, 1986); Neuberger *et al.*, PCT Application WO 86/01533 (published March 13, 1986); Kudo *et al.*, European Patent Application 184187 (published June 11, 1986); Morrison *et al.*, European Patent Application 173494 (published March 5, 1986); Sahagan *et al.*,
 25 1986, *J. Immunol.* 137:1066-1074; Robinson *et al.*, PCT/US86/02269 (published May 7, 1987); Liu *et al.*, 1987, *Proc. Natl. Acad. Sci. USA* 84:3439-3443; Sun *et al.*, 1987, *Proc. Natl. Acad. Sci. USA* 84:214-218; Better *et al.*, 1988, *Science* 240:1041-1043).

An anti-idiotypic (anti-Id) antibody is an antibody which recognizes unique determinants generally associated with the antigen-binding site of an antibody. An
 30 anti-Id antibody is prepared by immunizing an animal of the same species and genetic

-51-

type (e.g., mouse strain) as the source of the monoclonal antibody with the monoclonal antibody to which an anti-Id is being prepared. The immunized animal will recognize and respond to the idiotypic determinants of the immunizing antibody by producing an antibody to these isotypic determinants (the anti-Id antibody).

5 Accordingly, monoclonal antibodies generated against the polypeptides of the present invention may be used to induce anti-Id antibodies in suitable animals. Spleen cells from such immunized mice can be used to produce anti-Id hybridomas secreting anti-Id monoclonal antibodies. Further, the anti-Id antibodies can be coupled to a carrier such as keyhole limpet hemocyanin (KLH) and used to immunize additional
10 BALB/c mice. Sera from these mice will contain anti-anti-Id antibodies that have the binding properties of the final mAb specific for an R-PTPase epitope. The anti-Id antibodies thus have their idiotypic epitopes, or "idiotopes" structurally similar to the epitope being evaluated, such as *Streptococcus pyogenes* polypeptides.

 The term "antibody" is also meant to include both intact molecules as well as
15 fragments such as Fab which are capable of binding antigen. Fab fragments lack the Fc fragment of intact antibody, clear more rapidly from the circulation, and may have less non-specific tissue binding than an intact antibody (Wahl *et al.*, 1983, *J. Nucl. Med.* 24:316-325). It will be appreciated that Fab and other fragments of the antibodies useful in the present invention may be used for the detection and
20 quantitation of *N. meningitidis* polypeptides according to the methods for intact antibody molecules.

 The anti-Id antibody may be used as an "immunogen" to induce an immune response in yet another animal, producing a so-called anti-anti-Id antibody. The anti-anti-Id may be epitopically identical to the original mAb which induced the anti-Id.
25 Thus, by using antibodies to the idiotypic determinants of a mAb, it may be possible to identify other clones expressing antibodies of identical specificity.

 The antibodies may potentially be used in a variety of ways, e.g., for confirmation that a protein is expressed, or to confirm where a protein is expressed. Labeled antibody (e.g., fluorescent labeling for FACS) can be incubated with intact

-52-

bacteria and the presence of the label on the bacterial surface confirms the location of the protein, for instance.

Antibodies generated against the polypeptides of the invention can be obtained by administering the polypeptides or epitope-bearing fragments, analogs, or cells to an animal using routine protocols. For preparing monoclonal antibodies, any technique which provides antibodies produced by continuous cell line cultures are used.

Polynucleotides

As with the proteins of the present invention, a polynucleotide of the present invention may comprise a nucleic acid sequence that is identical to any of the reference sequences of odd numbered SEQ ID NOS:1-253, that is be 100% identical, or it may include up to a number of nucleotide alterations as compared to the reference sequence. Such alterations are selected from the group consisting of at least one nucleotide deletion, substitution, including transition and transversion, or insertion, and wherein said alterations may occur at the 5' or 3' terminal positions of the reference nucleotide sequence or anywhere between those terminal positions, interspersed either individually among the nucleotides in the reference sequence or in one or more contiguous groups within the reference sequence. The number of nucleotide alterations is determined by multiplying the total number of nucleotides in any of odd numbered SEQ ID NOS:1-253 by the numerical percent of the respective percent identity (divided by 100) and subtracting that product from said total number of nucleotides in said sequence.

25

-53-

By way of example, without intending to be limited thereto, an isolated *N. meningitidis* polynucleotide comprising a polynucleotide sequence that has at least 70% identity to any nucleic acid sequence of SEQ ID NOS:1-253; a degenerate variant thereof or a fragment thereof, wherein the polynucleotide sequence may

5 include up to n_n nucleic acid alterations over the entire polynucleotide region of the nucleic acid sequence of SEQ ID NOS:1-253, wherein n_n is the maximum number of alterations and is calculated by the formula:

$$n_n = x_n - (x_n \cdot y),$$

in which x_n is the total number of nucleic acids of any of SEQ ID NOS:1-253 and y

10 has a value of 0.70, wherein any non-integer product of x_n and y is rounded down to the nearest integer prior to subtracting such product from x_n . Of course, y may also have a value of 0.80 for 80%, 0.85 for 85%, 0.90 for 90% 0.95 for 95%, *etc.*

Alterations of a polynucleotide sequence encoding the polypeptides comprising amino acid sequences of any of SEQ ID NOS:2-252 may create nonsense, missense or

15 frameshift mutations in this coding sequence and thereby alter the polypeptide encoded by the polynucleotide following such alterations.

Certain embodiments of the present invention relate to polynucleotides (herein referred to as the "2086 polynucleotides" or "ORF2086 polynucleotides") which encode the 2086 proteins and antibodies made against the 2086 proteins. In preferred

20 embodiments, an isolated polynucleotide of the present invention is a polynucleotide comprising a nucleotide sequence having at least about 95% identity to a nucleotide sequence chosen from one of the odd numbered SEQ ID NO: 1 and SEQ ID NOS: 253, a degenerate variant thereof, or a fragment thereof. As defined herein, a

"degenerate variant" is defined as a polynucleotide that differs from the nucleotide

25 sequence shown in the odd numbered SEQ ID NOS:1 and SEQ ID NOS:253 (and fragments thereof) due to degeneracy of the genetic code, but still encodes the same 2086 protein (*i.e.*, the even numbered SEQ ID NOS: 2-252) as that encoded by the nucleotide sequence shown in the odd numbered SEQ ID NOS: 1-253.

-54-

In other embodiments, the polynucleotide is a complement to a nucleotide sequence chosen from one of the odd numbered SEQ ID NOS: 1-253, a degenerate variant thereof, or a fragment thereof. In yet other embodiments, the polynucleotide is selected from the group consisting of DNA, chromosomal DNA, cDNA and RNA and may further comprises heterologous nucleotides. In another embodiment, an isolated polynucleotide hybridizes to a nucleotide sequence chosen from one of SEQ ID NOS: 1 -253, a complement thereof, a degenerate variant thereof, or a fragment thereof, under high stringency hybridization conditions. In yet other embodiments, the polynucleotide hybridizes under intermediate stringency hybridization conditions.

It will be appreciated that the 2086 polynucleotides may be obtained from natural, synthetic or semi-synthetic sources; furthermore, the nucleotide sequence may be a naturally occurring sequence, or it may be related by mutation, including single or multiple base substitutions, deletions, insertions and inversions, to such a naturally occurring sequence, provided always that the nucleic acid molecule comprising such a sequence is capable of being expressed as 2086 immunogenic polypeptide as described above. The nucleic acid molecule may be RNA, DNA, single stranded or double stranded, linear or covalently closed circular form. The nucleotide sequence may have expression control sequences positioned adjacent to it, such control sequences usually being derived from a heterologous source. Generally, recombinant expression of the nucleic acid sequence of this invention will use a stop codon sequence, such as TAA, at the end of the nucleic acid sequence.

The invention also includes polynucleotides capable of hybridizing under reduced stringency conditions, more preferably stringent conditions, and most preferably highly stringent conditions, to polynucleotides described herein. Examples of stringency conditions are shown in the Stringency Conditions Table below: highly stringent conditions are those that are at least as stringent as, for example, conditions A-F; stringent conditions are at least as stringent as, for example, conditions G-L; and reduced stringency conditions are at least as stringent as, for example, conditions M-R.

30

-55-

STRINGENCY CONDITIONS - TABLE I

Stringency Condition	Polynucleotide Hybrid	Hybrid Length (bp) ^I	Hybridization Temperature and Buffer ^{II}	Wash Temperature and Buffer ^{II}
A	DNA:DNA	> 50	65°C; 1xSSC -or- 42°C; 1xSSC, 50% formamide	65°C; 0.3xSSC
B	DNA:DNA	< 50	T _D ; 1xSSC	T _D ; 1xSSC
C	DNA:RNA	> 50	67°C; 1xSSC -or- 45°C; 1xSSC, 50% formamide	67°C; 0.3xSSC
D	DNA:RNA	< 50	T _D ; 1xSSC	T _D ; 1xSSC
E	RNA:RNA	> 50	70°C; 1xSSC -or- 50°C; 1xSSC, 50% formamide	70°C; 0.3xSSC
F	RNA:RNA	< 50	T _F ; 1xSSC	T _F ; 1xSSC
G	DNA:DNA	> 50	65°C; 4xSSC -or- 42°C; 4xSSC, 50% formamide	65°C; 1xSSC
H	DNA:DNA	< 50	T _H ; 4xSSC	T _H ; 4xSSC
I	DNA:RNA	> 50	67°C; 4xSSC -or- 45°C; 4xSSC, 50% formamide	67°C; 1xSSC
J	DNA:RNA	< 50	T _J ; 4xSSC	T _J ; 4xSSC
K	RNA:RNA	> 50	70°C; 4xSSC -or- 50°C; 4xSSC, 50% formamide	67°C; 1xSSC
L	RNA:RNA	< 50	T _L ; 2xSSC	T _L ; 2xSSC
M	DNA:DNA	> 50	50°C; 4xSSC -or- 40°C; 6xSSC, 50% formamide	50°C; 2xSSC
N	DNA:DNA	< 50	T _N ; 6xSSC	T _N ; 6xSSC
O	DNA:RNA	> 50	55°C; 4xSSC -or- 42°C; 6xSSC, 50% formamide	55°C; 2xSSC
P	DNA:RNA	< 50	T _P ; 6xSSC	T _P ; 6xSSC
Q	RNA:RNA	> 50	60°C; 4xSSC -or- 45°C; 6xSSC, 50% formamide	60°C; 2xSSC
R	RNA:RNA	< 50	T _R ; 4xSSC	T _R ; 4xSSC

-56-

bp^l: The hybrid length is that anticipated for the hybridized region(s) of the hybridizing polynucleotides. When hybridizing a polynucleotide to a target polynucleotide of unknown sequence, the hybrid length is assumed to be that of the hybridizing polynucleotide. When polynucleotides of known sequence are
 5 hybridized, the hybrid length can be determined by aligning the sequences of the polynucleotides and identifying the region or regions of optimal sequence complementarities.

buffer^H: SSPE (1xSSPE is 0.15M NaCl, 10mM NaH₂PO₄, and 1.25mM EDTA, pH 7.4) can be substituted for SSC (1xSSC is 0.15M NaCl and 15mM sodium
 10 citrate) in the hybridization and wash buffers; washes are performed for 15 minutes after hybridization is complete.

T_B through T_R: The hybridization temperature for hybrids anticipated to be less than 50 base pairs in length should be 5-10EC less than the melting temperature (T_m) of the hybrid, where T_m is determined according to the following equations. For
 15 hybrids less than 18 base pairs in length, T_m(EC) = 2(# of A + T bases) + 4(# of G + C bases). For hybrids between 18 and 49 base pairs in length, T_m(EC) = 81.5 + 16.6(log₁₀[Na⁺]) + 0.41(%G+C) - (600/N), where N is the number of bases in the hybrid, and [Na⁺] is the concentration of sodium ions in the hybridization buffer ([Na⁺] for 1xSSC = 0.165 M).

20 Additional examples of stringency conditions for polynucleotide hybridization are provided in Sambrook, J., E.F. Fritsch, and T. Maniatis, 1989, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, chapters 9 and 11, and *Current Protocols in Molecular Biology*, 1995, F.M. Ausubel *et al.*, eds., John Wiley & Sons, Inc., sections 2.10 and 6.3-6.4.

25 The invention also provides polynucleotides that are fully complementary to these polynucleotides and also provides antisense sequences. The antisense sequences of the invention, also referred to as antisense oligonucleotides, include both

-57-

internally generated and externally administered sequences that block expression of polynucleotides encoding the polypeptides of the invention. The antisense sequences of the invention comprise, for example, about 15-20 base pairs. The antisense sequences can be designed, for example, to inhibit transcription by preventing
5 promoter binding to an upstream nontranslated sequence or by preventing translation of a transcript encoding a polypeptide of the invention by preventing the ribosome from binding.

The polynucleotides of the invention are prepared in many ways (e.g., by chemical synthesis, from DNA libraries, from the organism itself) and can take
10 various forms (e.g., single-stranded, double-stranded, vectors, probes, primers). The term "polynucleotide" includes DNA and RNA, and also their analogs, such as those containing modified backbones.

According to further embodiments of the present invention, the polynucleotides of the present invention comprise a DNA library, such as a cDNA
15 library.

Fusion Proteins

The present invention also relates to fusion proteins. A "fusion protein" refers to a protein encoded by two, often unrelated, fused genes or fragments thereof. For
20 example, fusion proteins comprising various portions of constant region of immunoglobulin molecules together with another immunogenic protein or part thereof. In many cases, employing an immunoglobulin Fc region as a part of a fusion protein is advantageous for use in therapy and diagnosis resulting in, for example, improved pharmacokinetic properties (see, e.g., EP 0 232 262 A1). On the other
25 hand, for some uses it would be desirable to be able to delete the Fc part after the fusion protein has been expressed, detected and purified. The 2086 polynucleotides

-58-

of the invention may be used for the recombinant production of polypeptides of the present invention, the polynucleotide may include the coding sequence for the mature polypeptide, by itself, or the coding sequence for the mature polypeptide in reading frame with other coding sequences, such as those encoding a leader or secretory sequence, a pre-, or pro- or prepro- protein sequence, or other fusion peptide portions. For example, a marker sequence which facilitates purification of a 2086 polypeptide or fused polypeptide may be encoded (*see Gentz et al.*, 1989). Thus, contemplated in some embodiments of the present invention is the preparation of polynucleotides encoding fusion polypeptides permitting His-tag purification of expression products.

10 The polynucleotide may also contain non-coding 5' and 3' sequences, such as transcribed, non-translated sequences, splicing and polyadenylation signals. Such a fused polypeptide can be produced by a host cell transformed/transfected or infected or infected with a recombinant DNA cloning vehicle as described below and it can be subsequently isolated from the host cell to provide the fused polypeptide substantially

15 free of other host cell proteins.

Immunogenic Compositions

One aspect of the present invention provides immunogenic compositions which comprise at least one 2086 protein or a nucleic acid encoding said protein. In some embodiments, such proteins have the ability to elicit antibodies that (1) are bactericidal to multiple strains; (2) react with the surface of multiple strains; (3) confer passive protection against a live challenge; and/or (4) prevent colonization.

The formulation of such immunogenic compositions is well known to persons skilled in this field. Immunogenic compositions of the invention preferably include a pharmaceutically acceptable carrier. Suitable pharmaceutically acceptable carriers and/or diluents include any and all conventional solvents, dispersion media, fillers, solid carriers, aqueous solutions, coatings, antibacterial and antifungal agents, isotonic

-59-

and absorption delaying agents, and the like. Suitable pharmaceutically acceptable carriers include, for example, one or more of water, saline, phosphate buffered saline, dextrose, glycerol, ethanol and the like, as well as combinations thereof.

Pharmaceutically acceptable carriers may further comprise minor amounts of
 5 auxiliary substances such as wetting or emulsifying agents, preservatives or buffers, which enhance the shelf life or effectiveness of the antibody. The preparation and use of pharmaceutically acceptable carriers is well known in the art. Except insofar as any conventional media or agent is incompatible with the active ingredient, use thereof in the immunogenic compositions of the present invention is contemplated.

10 The immunogenic compositions of the invention can include one or more adjuvants, including, but not limited to aluminum hydroxide; aluminum phosphate; STIMULON™ QS-21 (Aquila Biopharmaceuticals, Inc., Framingham, MA); MPL™ (3-O-deacylated monophosphoryl lipid A; Corixa, Hamilton, MT), 529 (an amino alkyl glucosamine phosphate compound, Corixa, Hamilton, MT), IL-12 (Genetics
 15 Institute, Cambridge, MA); GM-CSF (Immunex Corp., Seattle, Washington); N-acetyl-muramyl-L-theronyl-D-isoglutamine (thr-MDP); N-acetyl-nor-muramyl-L-alanyl-D-isoglutamine (CGP 11637, referred to as nor-MDP); N-acetylmuramyl-L-alanyl-D-isoglutaminyl-L-alanine-2-(1'-2'-dipalmitoyl-sn-glycero-3-hydroxyphosphoryloxy-ethylamine) (CGP 19835A, referred to as MTP-PE); and cholera toxin.
 20 Others which may be used are non-toxic derivatives of cholera toxin, including its A subunit, and/or conjugates or genetically engineered fusions of the *N. meningitidis* polypeptide with cholera toxin or its B subunit ("CTB"), procholeraenoid, fungal polysaccharides, including schizophyllan, muramyl dipeptide, muramyl dipeptide ("MDP") derivatives, phorbol esters, the heat labile toxin of *E. coli*, block polymers
 25 or saponins.

In certain preferred embodiments, the proteins of this invention are used in an immunogenic composition for oral administration which includes a mucosal adjuvant. The mucosal adjuvant can be a cholera toxin; however, preferably, mucosal adjuvants other than cholera toxin which may be used in accordance with the present invention

-60-

include non-toxic derivatives of a cholera holotoxin, wherein the A subunit is mutagenized, chemically modified cholera toxin, or related proteins produced by modification of the cholera toxin amino acid sequence. For a specific cholera toxin which may be particularly useful in preparing immunogenic compositions of this invention, see the mutant cholera holotoxin E29H, as disclosed in Published International Application WO 00/18434. These may be added to, or conjugated with, the polypeptides of this invention. The same techniques can be applied to other molecules with mucosal adjuvant or delivery properties such as *Escherichia coli* heat labile toxin (LT). Other compounds with mucosal adjuvant or delivery activity may be used such as bile; polycations such as DEAE-dextran and polyornithine; detergents such as sodium dodecyl benzene sulphate; lipid-conjugated materials; antibiotics such as streptomycin; vitamin A; and other compounds that alter the structural or functional integrity of mucosal surfaces. Other mucosally active compounds include derivatives of microbial structures such as MDP; acridine and cimetidine.

15 STIMULON™ QS-21, MPL, and IL-12, as described above, may also be used.

The immunogenic compositions of this invention may be delivered in the form of ISCOMS (immune stimulating complexes), ISCOMS containing CTB, liposomes or encapsulated in compounds such as acrylates or poly(DL-lactide-co-glycoside) to form microspheres of a size suited to adsorption. The proteins of this invention may also be incorporated into oily emulsions.

20

-61-

Multiple Antigens

The immunogenic agents, including proteins, polynucleotides and equivalents of the present invention may be administered as the sole active immunogen in a immunogenic composition, or alternatively, the composition may include other active
5 immunogens, including other *Neisseria sp.* immunogenic polypeptides, or immunologically-active proteins of one or more other microbial pathogens (e.g. virus, prion, bacterium, or fungus, without limitation) or capsular polysaccharide. The compositions may comprise one or more desired proteins, fragments or pharmaceutical compounds as desired for a chosen indication. In the same manner,
10 the compositions of this invention which employ one or more nucleic acids in the immunogenic composition may also include nucleic acids which encode the same diverse group of proteins, as noted above.

Any multi-antigen or multi-valent immunogenic composition is contemplated by the present invention. For example, the compositions of the present invention may
15 comprise combinations of two or more 2086 proteins, a combination of 2086 protein with one or more Por A proteins, a combination of 2086 protein with meningococcus serogroup A, C, Y and W135 polysaccharides and/or polysaccharide conjugates, a combination of 2086 protein with meningococcus and pneumococcus combinations, or a combination of any of the foregoing in a form suitable for mucosal delivery.
20 Persons of skill in the art would be readily able to formulate such multi-antigen or multi-valent immunologic compositions.

The 2086 polypeptides, fragments and equivalents can be used as part of a conjugate immunogenic composition; wherein one or more proteins or polypeptides
25 are conjugated to a carrier in order to generate a composition that has immunogenic properties against several serotypes. Alternatively, one of the 2086 polypeptides may potentially be used as a carrier protein for other immunogenic polypeptides.

-62-

The present invention also relates to a method of inducing immune responses in a mammal comprising the step of providing to said mammal an immunogenic composition of this invention. The immunogenic composition is a composition which is antigenic in the treated animal or human such that the immunologically effective
5 amount of the polypeptide(s) contained in such composition brings about the desired immune response against *N. meningitidis* infection.

Another specific aspect of the present invention may relate to using as the immunogenic composition a vector or plasmid which expresses a protein of this invention, or an immunogenic portion thereof. Accordingly, a further aspect this
10 invention may provide a method of inducing an immune response in a mammal, which comprises providing to a mammal a vector or plasmid expressing at least one isolated 2086 polypeptide. The protein of the present invention may be delivered to the mammal using a live vector, in particular using live recombinant bacteria, viruses or other live agents, containing the genetic material necessary for the expression of
15 the polypeptide or immunogenic portion as a foreign polypeptide.

Viral and Non-Viral Vectors

Preferred vectors, particularly for cellular assays *in vitro* and *in vivo*, may be viral vectors, such as lentiviruses, retroviruses, herpes viruses, adenoviruses, adeno-
20 associated viruses, vaccinia virus, baculovirus, and other recombinant viruses with desirable cellular tropism. Thus, a nucleic acid encoding a 2086 protein or immunogenic fragment thereof may be introduced *in vivo*, *ex vivo*, or *in vitro* using a viral vector or through direct introduction of DNA. Expression in targeted tissues can potentially be effected by targeting the transgenic vector to specific cells, such as with
25 a viral vector or a receptor ligand, or by using a tissue-specific promoter, or both. Targeted gene delivery is described in PCT Publication No. WO 95/28494.

-63-

Viral vectors commonly used for *in vivo* or *ex vivo* targeting and therapy procedures are DNA-based vectors and retroviral vectors. Methods for constructing and using viral vectors are known in the art (e.g., Miller and Rosman, *BioTechniques*, 1992, 7:980-990). Preferably, the viral vectors are replication-defective, that is, they are unable to replicate autonomously in the target cell. Preferably, the replication defective virus is a minimal virus, *i.e.*, it retains only the sequences of its genome which are necessary for encapsulating the genome to produce viral particles.

DNA viral vectors include an attenuated or defective DNA virus, such as but not limited to herpes simplex virus (HSV), papillomavirus, Epstein Barr virus (EBV), adenovirus, adeno-associated virus (AAV), and the like. Defective viruses, which entirely or almost entirely lack viral genes, are preferred. Defective virus is not infective after introduction into a cell. Use of defective viral vectors allows for administration to cells in a specific, localized area, without concern that the vector can infect other cells. Thus, a specific tissue can be specifically targeted. Examples of particular vectors include, but are not limited to, a defective herpes virus 1 (HSV1) vector (Kaplitt *et al.*, *Molec. Cell. Neurosci.*, 1991, 2:320-330), defective herpes virus vector lacking a glyco-protein 1 gene, or other defective herpes virus vectors (PCT Publication Nos. WO 94/21807 and WO 92/05263); an attenuated adenovirus vector, such as the vector described by Stratford-Perricaudet *et al.* (*J. Clin. Invest.*, 1992, 90:626-630; see also La Salle *et al.*, *Science*, 1993, 259:988-990); and a defective adeno-associated virus vector (Samulski *et al.*, *J. Virol.*, 1987, 61:3096-3101; Samulski *et al.*, *J. Virol.*, 1989, 63:3822-3828; Lebkowski *et al.*, *Mol. Cell. Biol.*, 1988, 8:3988-3996).

Various companies produce viral vectors commercially, including, but not limited to, Avigen, Inc. (Alameda, CA; AAV vectors), Cell Genesys (Foster City, CA; retroviral, adenoviral, AAV vectors, and lentiviral vectors), Clontech (retroviral and baculoviral vectors), Genovo, Inc. (Sharon Hill, PA; adenoviral and AAV vectors), Genevec (adenoviral vectors), IntroGene (Leiden, Netherlands; adenoviral vectors), Molecular Medicine (retroviral, adenoviral, AAV, and herpes viral vectors), Norgen

-64-

(adenoviral vectors), Oxford BioMedica (Oxford, United Kingdom; lentiviral vectors), and Transgene (Strasbourg, France; adenoviral, vaccinia, retroviral, and lentiviral vectors).

Adenovirus vectors. Adenoviruses are eukaryotic DNA viruses that can be modified to efficiently deliver a nucleic acid of this invention to a variety of cell types. Various serotypes of adenovirus exist. Of these serotypes, preference is given, within the scope of the present invention, to using type 2 or type 5 human adenoviruses (Ad 2 or Ad 5) or adenoviruses of animal origin (*see* PCT Publication No. WO 94/26914). Those adenoviruses of animal origin which can be used within the scope of the present invention include adenoviruses of canine, bovine, murine (example: Mav1, Beard *et al.*, Virology, 1990, 75-81), ovine, porcine, avian, and simian (example: SAV) origin. Preferably, the adenovirus of animal origin is a canine adenovirus, more preferably a CAV2 adenovirus (*e.g.*, Manhattan or A26/61 strain, ATCC VR-800, for example). Various replication defective adenovirus and minimum adenovirus vectors have been described (PCT Publication Nos. WO 94/26914, WO 95/02697, WO 94/28938, WO 94/28152, WO 94/12649, WO 95/02697, WO 96/22378). The replication defective recombinant adenoviruses according to the invention can be prepared by any technique known to the person skilled in the art (Levrero *et al.*, *Gene*, 1991, 101:195; European Publication No. EP 185 573; Graham, EMBO J., 1984, 3:2917; Graham *et al.*, *J. Gen. Virol.*, 1977, 36:59). Recombinant adenoviruses are recovered and purified using standard molecular biological techniques, which are well known to persons of ordinary skill in the art.

Adeno-associated viruses. The adeno-associated viruses (AAV) are DNA viruses of relatively small size that can integrate, in a stable and site-specific manner, into the genome of the cells which they infect. They are able to infect a wide spectrum of cells without inducing any effects on cellular growth, morphology or differentiation, and they do not appear to be involved in human pathologies. The AAV genome has been cloned, sequenced and characterized. The use of vectors derived from the AAVs for transferring genes *in vitro* and *in vivo* has been described

-65-

(see, PCT Publication Nos. WO 91/18088 and WO 93/09239; U.S. Patent Nos. 4,797,368 and 5,139,941; European Publication No. EP 488 528). The replication defective recombinant AAVs according to the invention can be prepared by cotransfecting a plasmid containing the nucleic acid sequence of interest flanked by two AAV inverted terminal repeat (ITR) regions, and a plasmid carrying the AAV encapsidation genes (rep and cap genes), into a cell line which is infected with a human helper virus (for example an adenovirus). The AAV recombinants which are produced are then purified by standard techniques.

Retrovirus vectors. In another implementation of the present invention, the nucleic acid can be introduced in a retroviral vector, e.g., as described in U.S. Patent No. 5,399,346; Mann *et al.*, *Cell*, 1983, 33:153; U.S. Patent Nos. 4,650,764 and 4,980,289; Markowitz *et al.*, *J. Virol.*, 1988, 62:1120; U.S. Patent No. 5,124,263; European Publication Nos. EP 453 242 and EP178 220; Bernstein *et al.*, *Genet Eng.*, 1985, 7:235; McCormick, *BioTechnology*, 1985, 3:689; PCT Publication No. WO 95/07358; and Kuo *et al.*, *Blood*, 1993, 82:845. The retroviruses are integrating viruses that infect dividing cells. The retrovirus genome includes two LTRs, an encapsidation sequence and three coding regions (gag, pol and env). In recombinant retroviral vectors, the *gag*, *pol* and *env* genes are generally deleted, in whole or in part, and replaced with a heterologous nucleic acid sequence of interest. These vectors can be constructed from different types of retrovirus, such as, HIV, MoMuLV ("murine Moloney leukaemia virus" MSV ("murine Moloney sarcoma virus"), HaSV ("Harvey sarcoma virus"); SNV ("spleen necrosis virus"); RSV ("Rous sarcoma virus") and Friend virus. Suitable packaging cell lines have been described in the prior art, in particular the cell line PA317 (U.S. Patent No. 4,861,719); the PsiCRIP cell line (PCT Publication No. WO 90/02806) and the GP+envAm-12 cell line (PCT Publication No. WO 89/07150). In addition, the recombinant retroviral vectors can contain modifications within the LTRs for suppressing transcriptional activity as well as extensive encapsidation sequences which may include a part of the gag gene (Bender *et al.*, *J. Virol.*, 1987, 61:1639). Recombinant retroviral vectors are purified by standard techniques known to those having ordinary skill in the art.

-66-

Retroviral vectors can be constructed to function as infectious particles or to undergo a single round of transfection. In the former case, the virus is modified to retain all of its genes except for those responsible for oncogenic transformation properties, and to express the heterologous gene. Non-infectious viral vectors are manipulated to destroy the viral packaging signal, but retain the structural genes required to package the co-introduced virus engineered to contain the heterologous gene and the packaging signals. Thus, the viral particles that are produced are not capable of producing additional virus.

Retrovirus vectors can also be introduced by DNA viruses, which permits one cycle of retroviral replication and amplifies transfection efficiency (*see* PCT Publication Nos. WO 95/22617, WO 95/26411, WO 96/39036 and WO 97/19182).

Lentivirus vectors. In another implementation of the present invention, lentiviral vectors can be used as agents for the direct delivery and sustained expression of a transgene in several tissue types, including brain, retina, muscle, liver and blood. The vectors can efficiently transduce dividing and nondividing cells in these tissues, and effect long-term expression of the gene of interest. For a review, *see*, Naldini, *Curr. Opin. Biotechnol.*, 1998, 9:457-63; *see also* Zufferey, *et al.*, *J. Virol.*, 1998, 72:9873-80). Lentiviral packaging cell lines are available and known generally in the art. They facilitate the production of high-titer lentivirus vectors for gene therapy. An example is a tetracycline-inducible VSV-G pseudotyped lentivirus packaging cell line that can generate virus particles at titers greater than 10^6 IU/mL for at least 3 to 4 days (Kafri, *et al.*, *J. Virol.*, 1999, 73: 576-584). The vector produced by the inducible cell line can be concentrated as needed for efficiently transducing non-dividing cells *in vitro* and *in vivo*.

Non-viral vectors. In another implementation of the present invention, the vector can be introduced *in vivo* by lipofection, as naked DNA, or with other transfection facilitating agents (peptides, polymers, etc.). Synthetic cationic lipids can be used to prepare liposomes for *in vivo* transfection of a gene encoding a marker

-67-

(Felgner, *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 1987, 84:7413-7417; Felgner and Ringold, *Science*, 1989, 337:387-388; see Mackey, *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 1988, 85:8027-8031; Ulmer *et al.*, *Science*, 1993, 259:1745-1748). Useful lipid compounds and compositions for transfer of nucleic acids are described in PCT Patent Publication Nos. WO 95/18863 and WO 96/17823, and in U.S. Patent No. 5,459,127. Lipids may be chemically coupled to other molecules for the purpose of targeting (see Mackey, *et al.*, *supra*). Targeted peptides, *e.g.*, hormones or neurotransmitters, and proteins such as antibodies, or non-peptide molecules could be coupled to liposomes chemically.

10 Other molecules are also useful for facilitating transfection of a nucleic acid *in vivo*, such as a cationic oligopeptide (*e.g.*, PCT Patent Publication No. WO 95/21931), peptides derived from DNA binding proteins (*e.g.*, PCT Patent Publication No. WO 96/25508), or a cationic polymer (*e.g.*, PCT Patent Publication No. WO 95/21931).

15 It is also possible to introduce the vector *in vivo* as a naked DNA plasmid. Naked DNA vectors for vaccine purposes or gene therapy can be introduced into the desired host cells by methods known in the art, *e.g.*, electroporation, microinjection, cell fusion, DEAE dextran, calcium phosphate precipitation, use of a gene gun, or use of a DNA vector transporter (*e.g.*, Wu *et al.*, *J. Biol. Chem.*, 1992, 267:963-967; Wu and Wu, *J. Biol. Chem.*, 1988, 263:14621-14624; Canadian Patent Application No. 2,012,311; Williams *et al.*, *Proc. Natl. Acad. Sci. USA*, 1991, 88:2726-2730). Receptor-mediated DNA delivery approaches can also be used (Curiel *et al.*, *Hum. Gene Ther.*, 1992, 3:147-154; Wu and Wu, *J. Biol. Chem.*, 1987, 262:4429-4432). U.S. Patent Nos. 5,580,859 and 5,589,466 disclose delivery of exogenous DNA
20 sequences, free of transfection facilitating agents, in a mammal. Recently, a relatively low voltage, high efficiency *in vivo* DNA transfer technique, termed electrotransfer, has been described (Mir *et al.*, *C.P. Acad. Sci.*, 1988, 321:893; PCT Publication Nos. WO 99/01157; WO 99/01158; WO 99/01175). Accordingly, additional embodiments of the present invention may relate to a method of inducing an immune response in a

-68-

human comprising administering to said human an amount of a DNA molecule encoding a 2086 polypeptide of this invention, optionally with a transfection-facilitating agent, where said polypeptide, when expressed, retains immunogenicity and, when incorporated into an immunogenic composition and administered to a human, may provide immunogenicity without inducing enhanced disease upon subsequent infection of the human with *Neisseria sp.* pathogen, such as *N. meningitidis*. Transfection-facilitating agents are known in the art and include bupivacaine, and other local anesthetics (for examples see U.S. Patent No. 5,739,118) and cationic polyamines (as published in International Patent Application WO 96/10038).

15

-69-

The present invention also relates to an antibody, which may either be a monoclonal or polyclonal antibody, specific for 2086 polypeptides as described above. Such antibodies may be produced by methods which are well known to those skilled in the art.

5 Bacterial Expression Systems and Plasmids

This invention also provides a recombinant DNA molecule, such as a vector or plasmid, comprising an expression control sequence having promoter sequences and initiator sequences and a nucleotide sequence which codes for a polypeptide of this invention, the nucleotide sequence being located 3' to the promoter and initiator
10 sequences. In yet another aspect, the invention provides a recombinant DNA cloning vehicle capable of expressing a 2086 polypeptide comprising an expression control sequence having promoter sequences and initiator sequences, and a nucleotide sequence which codes for a 2086 polypeptide, the nucleotide sequence being located 3' to the promoter and initiator sequences. In a further aspect, there is provided a host
15 cell containing a recombinant DNA cloning vehicle and/or a recombinant DNA molecule as described above. Suitable expression control sequences and host cell/cloning vehicle combinations are well known in the art, and are described by way of example, in Sambrook *et al.* (1989).

Once recombinant DNA cloning vehicles and/or host cells expressing a
20 desired *a* polypeptide of this invention have been constructed by transforming, transfecting or infecting such cloning vehicles or host cells with plasmids containing the corresponding 2086 polynucleotide, cloning vehicles or host cells are cultured under conditions such that the polypeptides are expressed. The polypeptide is then isolated substantially free of contaminating host cell components by techniques well
25 known to those skilled in the art.

The following examples are included to demonstrate preferred embodiments of the invention. It should be appreciated by those skilled in the art that the techniques disclosed in the examples which follow represent techniques discovered by the inventors to function well in the practice of the invention, and thus can be

-70-

considered to constitute preferred modes for its practice. The scope of the claims should not be limited by the preferred embodiments and examples, but should be given the broadest interpretation consistent with the description as a whole.

EXAMPLES

5 Example 1

Identification of a neisserial membrane protein extract capable of eliciting bactericidal antibodies against heterologous strains:

Referring to Table II below, LOS-depleted outer membrane protein preparations have been shown to elicit bactericidal antibodies. These antibodies are often directed towards the PorA of the respective strain. LOS-depleted outer membrane preparations from serogroup B meningococcal strain 8529 (B:15:P1.7b,3) are unusual in this manner because they unexpectedly elicit bactericidal antibodies to several heterologous strains.

TABLE II

15 **BC Activity of Anti-sOMPS Against Different Strains of *N. meningitidis***

Anti-serum Week 6	H44/76	5315	H355	M982	880049	8529 ^a	NMB
Serotype	P1.7,16	P1.5	P1.15	P1.9	P1.4	P1.3	P1.5,2
sOMPs H44/76 25µg QS-21 20µg	1,000	< 50	< 50	< 50	< 50	980	< 50
sOMPs 5315 25µg QS-21 20µg	50	< 50	< 50	< 50	< 50	2170	< 50
sOMPs H355 25µg QS-21 20µg	< 50	< 50	450	< 50	< 50	860	< 50
sOMPs M982 25µg QS-21 20µg	92	< 50	< 50	300	< 50	1100	< 50
sOMPs 880049 25µg QS-21 20µg	50	< 50	< 50	< 50	< 50	1190	< 50
sOMPs 8529 25µg QS-21 20µg	1,000	< 50	450	50	215	>4050 (81.7)	< 50
sOMPs 2996 25µg QS-21 20µg	< 50	< 50	< 50	< 50	< 50	790	148
Whole-cell control serum 25µg 3DMP1, 25µg	450	50	100	500	150	>1350 (66.0)	952

-71-

To facilitate the isolation and characterization of the antigen(s) responsible for eliciting heterologous bactericidal antibodies, we sought to identify which detergent optimally extracted the antigen(s)

Strains and culture conditions.

- 5 *N. meningitidis* strain 8529 from a frozen vial was streaked onto a GC plate. (The meningococcal strain 8529 was received from The RIVM, Bilthoven, The Netherlands). The plate was incubated at 36C/5%CO₂ for 7.5h. Several colonies were used to inoculate a flask containing 50 mL of modified Frantz medium + GC supplement. The flask was incubated in an air shaker at 36°C and agitated at 200
- 10 RPM for 4.5h. 5 mL was used to inoculate a Fernbach flask containing 450 mL of modified Frantz medium + GC supplement. The flask was incubated in an air shaker at 36°C and agitated at 100 RPM for 11h. The entire 450 mL was used to inoculate 8.5 L of modified Frantz medium+ GC supplement in a 10 L fermentor.

15 Composition of Modified Frantz Medium:

	Glutamic acid	1.3 g/L
	Cysteine	0.02
	Sodium phosphate, dibasic, 7 hydrate	10
20	Potassium chloride	0.09
	Sodium chloride	6
	Ammonium chloride	1.25
	Dialyzed yeast extract (YE)	40 ml
	(25% YE soln. dialyzed against 5 volumes of dH ₂ O overnight, then	
25	autoclaved)	

GC supplement 100X, filter sterilize

30

-72-

Dextrose	400 g/L
Glutamic acid	10
Coccarboxylase	0.02
Ferric nitrate	0.5

5

The following parameters were controlled during fermentation:

Temperature=36°C ; pH=7.4; Dissolved Oxygen=20%. Several drops of P-2000 antifoam were added to control foaming. The culture was grown to stationary phase. Cells were harvested by centrifugation at OD650=5.25. A total of 100-300 grams of
10 wet cell paste is typically harvested from ~8.5L of culture.

Partial purification of outer membrane protein fractions from meningococci which elicit heterologous bactericidal antibodies:

100 gms wet weight of cells were suspended, to a volume five times the wet weight, with 10mM HEPES-NaOH, pH 7.4, 1mM Na₂EDTA and lysed by passage
15 through a 110Y microfluidizer equipped with a chamber at ~18,000 psi. The cell lysate was clarified and the cell envelope isolated by centrifugation at 300,000 x g for 1 hour at 10°C. The cell envelopes were washed 2X with the same buffer by suspension with a homogenizer followed by centrifugation as above. The cell envelopes were then extracted with 320mL of 1% (w/v) TritonTM X-100 in 10mM
20 HEPES-NaOH, pH 7.4, 1mM MgCl₂. Referring to Table III below, results from sequential differential detergent extraction using TritonTM X-100 and Zwittergent 3-14 followed by immunization of mice, allowed us to determine that the TritonTM extracts optimally extracted the candidate(s) of interest. This TritonTM X-100 extract, eliciting bactericidal antibody response against 4 out of five strains listed in table III.
25 was then fractionated by preparative isoelectric focusing (IEF) in a BioRad Rotaphor unit. Ampholyte concentrations were 1% pH 3-10 mixed with 1% pH 4-6. As shown in Table III, several fractions were found to elicit a heterologous bactericidal response. The fractions obtained from IEF, which focused in the pH range of 5.5-7.8, elicited a

heterologous response to the most strains as determined by the bactericidal assay.

The pooled IEF fractions were concentrated and the ampholytes removed by ethanol precipitation. A further purification was achieved by adsorbing some of the proteins obtained in the pH range of about 5.5-7.8 on an anion exchange column and

- 5 comparing the bactericidal activity obtained after immunizing mice with the adsorbed and unadsorbed proteins. Referring again to Table II, while many proteins were adsorbed to the anion exchange resin, the proteins which were not adsorbed by the column elicited more heterologous bactericidal antibodies.

TABLE III

Method	Fraction	BC ₅₀ Target Strain				
		H44/76	880049	H355	539*	M982
LOS-depleted	sOMPs	1,000	215	450	NC	50
Detergent Extractions	Cytoplasmic Extract	200	NT	NT	NT	NT
	TX-100	>800	>800	>800	>800	<25
	Zwittergent 3-12	400	>25	100	400	<25
	Zwittergent 3-14	<25	NT	NT	NT	NT
	Zw.3-14 + NaCl	<25	NT	NT	NT	NT
	Sarcosyl	<25	NT	NT	NT	NT
	Zw.3-14 + hcat	<25	NT	NT	NT	NT
Preparative IEF	Fractions 1-3 (pH 2.3-3.9)	50	NT	NT	NT	NT
	Fraction 4 (pH 4.1)	>800	<25	100	<25	NT
	Fraction 5 (pH 4.3)	>800	<25	100	200	NT
	Fraction 6 (pH 4.5)	400	NT	NT	NT	NT
	Fraction 7 (pH 4.8)	<25	NT	NT	NT	NT
	Fractions 8-9 (pH 5.0-5.3)	<25	NT	NT	NT	NT
	Fractions 10-17 (pH 5.5-7.8)	>800	200	<800	<800	NT
Anion Exchange	Adsorbed	400	NT	100	100	NT
	Unadsorbed	>6,400	NT	<800	<800	NT

10 NT: not tested

*Clinical isolate 539 is a homologous strain to 8529, isolated from the same outbreak

As shown in FIG. 1A, two major proteins were present in the unadsorbed fraction as determined by SDS-PAGE. To identify these proteins, two types of analysis were performed. One analysis was to perform limited proteolytic degradation (See FIG. 1A, and FIG. 1B) followed by isolation of peptides and direct protein sequencing. The other analysis was to perform SDS-PAGE followed by gel

15

-74-

excision, proteolytic digestion, and LC-MS/MS (Liquid Chromotography tandem Mass Spectrometry), (see FIG. 3) to obtain mass spectral information on the components of the preparations of interest. (See peptide mapping and sequencing methods described later in this section)

- 5 The *N. meningitidis* A Sanger genomic sequence was analyzed using the methods and algorithms described in Zagursky and Russell, 2001, *BioTechniques*, 31:636-659. This mining analysis yielded over 12,000 possible Open Reading Frames (ORFs). Both the direct sequence data and the mass spectral data described above indicated that the major components of the unadsorbed fraction were the
- 10 products of several ORFs present in an analysis of the Sanger database. The three predominant proteins identified by this methodology correspond to ORFs 4431, 5163 and 2086, (see FIGS. 1B and 3).

- Although ORF 4431 was the most predominant protein identified in the fractions, mouse antibodies to recombinant lipidated 4431 were not bactericidal and
- 15 did not provide a protective response in an animal model. Additional analysis of ORF 5163 is in progress.

 The second most predominant component of the preparations described herein corresponds to the product of ORF 2086.

Immunogenicity Methods:

20 Preparation of antisera:

- Except where noted, protein compositions/vaccines were formulated with 25µg of total protein and were adjuvanted with 20µg QS-21. A 0.2mL dose was administered by subcutaneous (rump) injection to 6-8 week old female Swiss-Webster mice at week 0 and 4. Bleeds were collected at week 0 and 4, and a final
- 25 exsanguination bleed was performed on week 6.

Bactericidal assay:

-75-

Bactericidal assays were performed essentially as described (See Mountzouros and Howell, 2000, *J. Clin. Microbiol.* 38(8):2878-2884). Complement-mediated antibody-dependent bactericidal titers for the SBA were expressed as the reciprocal of the highest dilution of test serum that killed $\geq 50\%$ of the target cells introduced into the assays (BC₅₀ titer).

Methods used to identify 2086 protein:**Cyanogen Bromide cleavage and direct sequencing of fragments:**

Cyanogen Bromide cleavage of Anion Exchange Unadsorbed Fraction (AEUF). The AEUF was precipitated with 90% cold ethanol and was solubilized with 10mg/mL cyanogen bromide in 70% formic acid to a protein concentration of 1mg/mL. The reaction was performed overnight at room temperature in the dark. The cleaved products were dried down by speed vacuum, and the pellet was solubilized with HE/0.1% reduced TX-100. SDS-PAGE followed by N-terminal amino acid sequencing were used to identify the components of this fraction.

Protease digestion/reverse phase/N-terminal sequencing to identify components:

The AEUF was digested with either GluC (V8), LysC or ArgC. The protein to enzyme ratio was 30 μ g protein to 1 μ g enzyme. The digestion was carried out at 37°C overnight. The digested protein mixture (30 μ g) was passed over a seven micron Aquapore RF-300 column and was eluted with a gradient of 10-95% acetonitrile in 0.1% trifluoroacetic acid, and peaks were collected manually. A no protein blank was also run, and the peaks from this were subtracted from the sample chromatogram. Peaks occurring only in the sample run were analyzed by mass spectrometer, and those samples giving a clear mass were analyzed for N-terminal amino acid sequencing.

N-terminal amino acid sequencing:

-76-

For bands excised from a blot, the protein sample is transferred from an SDS gel to a PVDF membrane, stained with Amido Black (10% acetic acid, 0.1% amido black in deionized water) and destained in 10% acetic acid. The desired protein band is then excised from all ten lanes using a methanol cleaned scalpel or mini-Exacto
5 knife and placed in the reaction cartridge of the Applied Biosystems 477A Protein Sequencer. For direct sequencing of samples in solution, the Prosorb cartridge is assembled and the PVDF wetted with 60 μ L of methanol. The PVDF is rinsed with 50 μ L of deionized water and the sample (50 μ L) is loaded to the PVDF. After 50 μ L of deionized water is used to rinse the sample, the Prosorb PVDF is punched out,
10 dried, and placed in the reaction cartridge of the Applied Biosystems 477A Protein Sequencer. For both methods, the Applied Biosystems N-terminal Sequencer is then run under optimal blot conditions for 12 or more cycles (1 cycle Blank, 1 cycle Standard, and 10 or more cycles for desired residue identification) and PTH-amino acid detection is done on the Applied Biosystems 120A PTH Analyzer. The cycles
15 are collected both on an analog chart recorder and digitally via the instrument software. Amino acid assignment is done using the analog and digital data by comparison of a standard set of PTH-amino acids and their respective retention times on the analyzer (cysteine residues are destroyed during conversion and are not detected). Multiple sequence information can be obtained from a single residue and
20 primary versus secondary assignments are made based on signal intensity.

LC-MS/MS

Protein samples purified by IEF were further analyzed by SDS-polyacrylamide gel electrophoresis. Proteins were visualized by Coomassie blue staining, and bands of interest were excised manually, then reduced, alkylated and
25 digested with trypsin (Promega, Madison, WI) *in situ* using an automated in-gel tryptic digestion robot (1). After digestion, peptide extracts were concentrated to a final volume of 10-20 μ L using a Savant Speed Vac Concentrator (ThermoQuest, Holdbrook, NY).

-77-

Peptide extracts were analyzed on an automated microelectrospray reversed phase HPLC. In brief, the microelectrospray interface consisted of a Picofrit fused silica spray needle, 50 cm length by 75 μ m ID, 8 μ m orifice diameter (New Objective, Cambridge MA) packed with 10 μ m C18 reversed-phase beads (YMC, Wilmington, NC) to a length of 10 cm. The Picofrit needle was mounted in a fiber optic holder (Melles Griot, Irvine, CA) held on a home-built base positioned at the front of the mass spectrometer detector. The rear of the column was plumbed through a titanium union to supply an electrical connection for the electrospray interface. The union was connected with a length of fused silica capillary (FSC) tubing to a FAMOS autosampler (LC-Packings, San Francisco, CA) that was connected to an HPLC solvent pump (ABI 140C, Perkin-Elmer, Norwalk, CT). The HPLC solvent pump delivered a flow of 50 μ L /min which was reduced to 250 nL/min using a PEEK microtight splitting tee (Upchurch Scientific, Oak Harbor, WA), and then delivered to the autosampler using an FSC transfer line. The LC pump and autosampler were each controlled using their internal user programs. Samples were inserted into plastic autosampler vials, sealed, and injected using a 5 μ L sample loop.

Microcapillary HPLC-mass spectrometry:

Extracted peptides from in-gel digests were separated by the microelectrospray HPLC system using a 50 minute gradient of 0-50% solvent B (A: 0.1M HoAc, B: 90% MeCN/0.1M HoAc). Peptide analyses were done on a Finnigan LCQ ion trap mass spectrometer (ThermoQuest, San Jose, CA) operating at a spray voltage of 1.5 kV, and using a heated capillary temperature of 150 °C. Data were acquired in automated MS/MS mode using the data acquisition software provided with the instrument. The acquisition method included 1 MS scan (375-1200 m/z) followed by MS/MS scans of the top 3 most abundant ions in the MS scan. The dynamic exclusion and isotope exclusion functions were employed to increase the number of peptide ions that were analyzed (settings: 3 amu = exclusion width, 3 min = exclusion duration, 30 secs = pre-exclusion duration, 3 amu = isotope exclusion width). Automated analysis of MS/MS data was performed using the SEQUEST computer algorithm incorporated into the Finnigan Bioworks data analysis package

-78-

(ThermoQuest, San Jose, CA) using the database of proteins derived from the complete genome of *N. meningitidis* (from Sanger). The results of the study are illustrated in FIG. 3.

5 **Example 2**

Cloning of Recombinant Lipidated P2086 (rLP2086):

A.) Native Leader Sequence:

Source materials:

The ORF 2086 gene was amplified by PCR from a clinical isolate of a serogroup B *Neisseria meningitidis* strain designated 8529. The serogroup, serotype and serosubtype of this strain is shown in parentheses; 8529 (B:15, P1:7b,3). This meningococcal strain was received from The RIVM, Bilthoven, The Netherlands. The mature 2086 protein gene sequence from meningococcal strain 8529 is provided herein as SEQ ID. NO. 212.

15 **PCR Amplification and Cloning Strategy:**

A visual inspection of ORF 2086 indicated that this gene had a potential lipoprotein signal sequence. Additional analysis using a proprietary Hidden Markov Model Lipoprotein algorithm confirmed that ORF 2086 contains a lipoprotein signal sequence. In order to recombinantly express P2086 in a more native-like conformation, oligonucleotide primers were designed to amplify the full length gene with the lipoprotein signal sequence intact and were based on an analysis of the Sanger sequence for *N. meningitidis* A ORF 2086, (5' primer – CT ATT CTG CAT ATG ACT AGG AGC and 3' primer – GCGC GGATCC TTA CTG CTT GGC GGC AAG ACC), which are SEQ ID NO. 304 (Compound No. 4624) and SEQ ID NO. 303 (Compound No. 4623), respectively (See also Table IV herein). The 2086 gene was amplified by polymerase chain reaction (PCR) [ABI 2400 thermal cycler, Applied Biosystems, Foster City, CA] from *N. meningitidis* strain 8529. The correct

-79-

size amplified product was ligated and cloned into pCR2.1-TOPO (Invitrogen). The plasmid DNA was restriction digested with NdeI and BamHI, gel purified and ligated into pET-27b(+) vector (Novagen).

Oligonucleotide primers described herein, were synthesized on a PerSeptive Biosystems oligonucleotide synthesizer, Applied Biosystems, Foster City CA, using β -Cyanoethylphosphoramidite chemistry, Applied Biosystems, Foster City CA. The primers used for PCR amplification of the ORF 2086 gene families are listed in Table IV, which shows non-limiting examples of primers of the present invention.

10 **TABLE IV: PRIMERS**

SEQ ID NO. (Compound No.)	Primer	Sequence	Restriction sites
303 (4623)	Reverse	GCGCGGATCCTTACTGCTTGGCGGCAAGACC	BamHI
304 (4624)	Forward	CTATTCTGCATATGACTAGGAGC	NdeI
305 (4625)	Forward	AGCAGCGGAGGCGGCGGTGTC	
306 (5005)	Forward	TGCCGATGCACTAACCGCACC	
307 (5007)	Reverse	CGTTTCGCAACCATCTTCCCG	
308 (5135)	Reverse	GAGATCTCACTCACTCATTACTGCTTGGCGGCAAGACCGATATG	BglII
309 (5658)	Forward	GCGGATCCAGCGGAGGGGGTGGTGTGCGCC	BamHI
310 (5660)	Reverse	GCGCATGCTTACTGCTTGGCGGCAAGACCGATATG	SphI
311 (6385)	Forward	GCGGATCCAGCGGAGGCGGCGGAAGC	BamHI
312 (6406)	Forward	GCGCAGATCTCATATGAGCAGCGGAGGGGGTGGTGTGCGCCGCGGAYATWGGTGC	BglII and NdeI
313 (6470)	Forward	CTATTCTGCGTATGACTAG	
314 (6472)	Reverse	GTCCGAACGGTAAATTATCGTG	
315 (6473)	Forward	GCGGATCCAGCGGAGGCGGCGGTGTCGCGCC	BamHI
316 (6474)	Forward	GAGATCTCATATGAGCAGCGGAGGCGGCGGAAGC	BglII and NdeI
317 (6495)	Forward	GACAGCCTGATAAACC	
318	Reverse	GATGCCGATTTTCGTGAACC	

-80-

SEQ ID NO. (Compound No.)	Primer	Sequence	Restriction sites
(6496)			
319 (6543)	Reverse	GCGCATGCGTACTGTTTGGCGGCGATG	SphI
320 (6605)	Reverse	GAGATCTCACTCAC TCACTACTGTTTGGC GGCGATGCGCATTTG	BglII
321 (6721)	Forward	GCGCAGATCTCATATGAGCAGCGGAGGCG GCGGAAGCGGAGGCGGCGGTGTCACCGGC GACATAGGCACC	BglII and NdeI

rLP2086 lipoprotein expression utilizing native leader sequence:

Referring to FIG. 5, plasmid pPX7340 was transformed/transfected or infected into BLR(DE3) pLysS host cells (Life Sciences). One transformant was selected and inoculated into 50 mL of Terrific Broth containing 2% glucose, kanamycin (30µg/mL), chloramphenicol (30µg/mL), and tetracycline (12µg/mL). The OD600 for the overnight culture was 6.0. The overnight culture was diluted out in 1 liter of Terrific Broth with 1% glycerol and the same antibiotics. The starting OD600 was 0.4. After 2 hours the OD600 was 1.6 and a pre-induced sample was taken. Cells equivalent to an OD600=1 were centrifuged and the supernatant was removed. The whole cell pellet was resuspended in 150µL Tris-EDTA buffer and 150µL of 2x SDS-PAGE sample buffer. IPTG was added to a final concentration of 1mM. After 3.5 hours a post-induced sample was taken as described and analyzed on SDS-PAGE (See FIG. 4).

Purification of rLP2086:

The rLP2086 was solubilized from *E. coli* following differential detergent extraction. Unlike the P2086 in its native environment, the rLP2086 was not significantly solubilized by TritonTM X-100 or Zwittergent 3-12. The bulk of the rLP2086 was solubilized with sarcosyl, indicating that it interacts with the outer membrane components of *E. coli* differently than it does in *N. meningitidis*. Once solubilized the rLP2086 was purified similarly to the native protein in that many of the contaminating *E. coli* proteins could be removed by adsorption to an anion exchange resin at pH 8. Despite being greater than one half a pH unit above its

-81-

theoretical pI, the rLP2086 remained unadsorbed at pH 8. Further purification was achieved by adsorption of the rLP2086 to a cation exchange resin at pH 4.5.

The homogeneity of the rLP2086 is shown in FIG. 2 following SDS-PAGE. The mass of rLP2086 was determined by MALDI-TOF mass spectral analysis to be 27,836. This mass differs from the theoretical mass of 27,100 by 736, which approximates the mass of the N-terminal lipid modification common to bacterial lipoproteins. Both native and rLP2086 appear to be outer membrane lipoproteins. Attempts with N-terminal sequencing were blocked and this is consistent with the terminal modification.

Purification Methods:

Frozen pellets of BLR DE3 pLysS cells expressing P2086 were resuspended in 10mM HEPES-NaOH/1mM EDTA/1μg/mL Pefabloc SC protease inhibitor (Roche) pH 7.4 (HEP) at 20mL/g wet cell weight and lysed by microfluidizer (Microfluidics Corporation Model 110Y). The cell lysate was centrifuged at 150,000 x g for one hour. The pellet was washed twice with HEP and centrifuged twice, and the resulting membrane pellet was frozen overnight. The pellet was solubilized with 10mM HEPES-NaOH/1mM MgCl₂/1%TX-100 pH 7.4 for 30 minutes, followed by centrifugation at 150,000 x g for 30 minutes. This was repeated three times. The membrane pellet was washed as above twice with 50mM Tris-HCl/5mM EDTA/1% Zwittergent 3-12 pH 8, followed by two washes each of 50mM Tris-HCl/5mM EDTA/1% Zwittergent 3-14 pH 8 and 50mM Tris-HCl/5mM EDTA/1% Zwittergent 3-14/ 0.5M NaCl pH 8.

The rLP2086 was then solubilized with 50mM Tris-HCl/5mM EDTA/ 1% sarcosyl pH 8. This sarcosyl extract was adjusted to 1% Zwittergent 3-14 (Z3-14) and dialyzed twice against a 30 fold excess of 50mM Tris-HCl/5mM EDTA/1% Z3-14. The dialyzed rLP2086 extract was precipitated with 90% ethanol to remove remaining sarcosyl, and solubilized with 50mM Tris-HCl/5mM EDTA/ 1% Z3-14 pH 8 (TEZ).

-82-

Insoluble material was removed by centrifugation, the supernatant was passed over an anion exchange chromatography column, and rLP2086 was collected in the unbound fraction. The unbound material was then dialyzed twice against a 30 fold excess of 25mM NaAc/1% Z3-14 pH 4.5, and passed over a cation exchange chromatography column. The rLP2086 was eluted with a 0-0.3M NaCl gradient and analyzed by SDS-PAGE (Coomassie stain). The rLP2086 pool was determined to be 84% pure by laser densitometry.

Surface Reactivity and Bactericidal Activity of Antisera to rLP2086 Subfamily B.

Referring to Table VII, antisera to purified rLP2086 from the Subfamily B strain 8529, demonstrated surface reactivity to all ten 2086 Subfamily B strains tested by whole cell ELISA. Bactericidal activity was detected against nine of ten 2086 Subfamily B strains expressing heterologous serosubtype antigens, PorAs. These strains are representative of strains causing serogroup B meningococcal disease throughout western Europe, the Americas, Australia, and New Zealand. The only strain which was not killed in the bactericidal assay, 870227, reacted strongly with the anti-rLP2086 (Subfamily B) sera by whole cell ELISA, indicating that this strain expresses a protein with epitopes in common to P2086.

The 2086 Subfamily A strains listed in Table VII, were also tested for surface reactivity by whole cell ELISA. Two out of three of these strains appeared to have a very low level of reactivity, indicating that some 2086 Subfamily A strains may not be cross-reactive with antibodies raised to rLP2086 Subfamily B. The PCR amplification procedure used to identify the 2086 Subfamily B gene from strain 8529 was also performed on strains 870446, NMB and 6557. No 2086 Subfamily B PCR amplified product was detected.

Immunogenicity Methods :

Preparation of antisera:

-83-

Vaccines were formulated as described previously in Example 1. However, a 10 µg dose was used.

Whole cell enzyme-linked immunosorbant assay (ELISA):

N. meningitidis whole cell suspensions were diluted to an optical density of 0.1 at 620nm in sterile 0.01M phosphate, 0.137M NaCl, 0.002M KCl (PBS). From this suspension, 0.1mL were added to each well of Nunc™ Bac T 96 well plates (Cat# 2-69620). Cells were dried on the plates at room temperature for three days, then were covered, inverted and stored at 4°C. Plates were washed three times with wash buffer (0.01M Tris-HCl, 0.139M NaCl/KCl, 0.1% dodecylpoly(oxyethyleneglycolether)_n n=23 (Brij-35®, available from ICI Americas, Inc., Wilmington, Delaware), pH 7.0-7.4). Dilutions of antisera were prepared in PBS, 0.05% Tween™-20/Azide and 0.1mL was transferred to the coated plates. Plates were incubated for two hours at 37°C. Plates were washed three times in wash buffer. Goat-anti-mouse IgG AP (Southern Biotech) was diluted at 1:1500 in PBS/0.05% Tween™-20, 0.1mL was added to each well, and plates were incubated at 37°C for two hours. Plates were washed (as above). Substrate solution was prepared by diluting p-nitrophenyl phosphate (Sigma) in 1M diethanolamine/0.5mM MgCl₂ to 1mg/mL. Substrate was added to the plate at 0.1mL per well and incubated at room temperature for one hour. The reaction was stopped with 50µL /well of 3N NaOH and plates were read at 405nm with 690nm reference.

B.) P4 Leader Sequence:

PCR Amplification and Cloning Strategy:

In order to optimize rLP2086 expression, the 2086 gene was cloned behind the P4 signal sequence of nontypable *Haemophilus influenzae* (Green *et al.*, 1991). Primers utilized for lipoprotein cloning are listed in Table IV and are identified by compound numbers: 5658, 5660, 6473, 6543 and 6385. ORF 2086 was amplified from *N. meningitidis* B strain 8529 using primers with the following compound

-84-

numbers 5658 and 5660. ORF 2086 was amplified from *N. meningitidis* serogroup B strain CDC1573 using primers with the following compound numbers 6385 and 5660. ORF 2086 was amplified from *N. meningitidis* serogroup B strain 2996 using primers with the following compound numbers 6473 and 6543. The N-terminal (5') primers
 5 were designed to be homologous to the mature region of the 2086 gene (starting at the serine residue at amino acid position 3 just downstream of the cysteine). The restriction site BamHI (GGA TTC) was incorporated into the 5' end of each N-terminal primer and resulted in the insertion of a glycine residue in the mature protein at amino acid position 2. The C-terminal (3') primers were designed to be
 10 homologous to the C-terminal end of the 2086 gene and included the Stop codon as well as an SphI site for cloning purposes. The amplified fragment from each *N. meningitidis* B strain was cloned into an intermediate vector and screened by sequence analysis.

Plasmid DNA from correct clones was digested with BamHI and SphI
 15 restriction enzymes (New England Biolabs, (NEB)). A vector designated pLP339 (supplied by applicants' assignee) was chosen as the expression vector. This vector utilizes the pBAD18-Cm backbone and contains the P4 lipoprotein signal sequence and P4 gene of nontypable *Haemophilus influenzae*. The pLP339 vector was partially digested with the restriction enzyme BamHI and then subjected to SphI digestion.
 20 The amplified 2086 fragments (BamHI/SphI) were each ligated separately into the pLP339 vector (partial BamHI/SphI). This cloning strategy places the mature 2086 gene behind the P4 lipoprotein signal sequence. The BamHI site remains in the cloning junction between the P4 signal sequence and the 2086 gene (See the plasmid construct shown in FIG. 7). The following is an example of the sequence at the
 25 BamHI cloning junction:

[P4 signal sequence]— TGT GGA TCC [remaining 2086 mature nucleic acid sequence]

[P4 signal sequence] Cys Gly Ser —[remaining 2086 mature amino acid sequence]

Referring to FIG. 7, each amplified fragment was cloned into a modified pBAD18-Cm vector containing the P4 leader sequence. Fermentation was performed on recombinant *E. coli* BLR pPX7343 which expresses rP4LP2086 (recombinant P4
5 lipidated 2086) to try to increase the cell density by adding additional glucose. The fermentor was filled with 10L complete M9 Minimal medium, according to Sambrook, supplemented with 1% glucose.

The initial concentration of glucose in the fermentor was 45g/L. The fermentor was inoculated to initial OD of ~0.25. At ~OD 25, additional 20g/L glucose
10 was added. The culture was induced with 1% arabinose at glucose depletion at OD 63.4. The fermentation continued until 3 hours after induction. Samples were saved at t=0, 1, 2, 3 post induction and protein quantified using BSA. At t=3, protein yield is ~0.35 g/L, and 7% total cellular protein. A total of 895 grams of wet cell paste was harvested from ~10 L of culture.

15 Purification of the rP4LP2086 was performed using the same methods as described above in Example 2, section A.

Example 3

Development Genetics for Non-lipidated Mature 2086 Protein:

20 To further evaluate the immunogenicity of the 2086 protein, cloning and expression of the non-lipidated form of P2086 were performed.

PCR gene amplification of the ORF 2086:

Oligonucleotides used for PCR amplification of the non-lipidated 2086 gene are listed in the primer table, Table IV. The 2086 gene from strain 8529 can be amplified with primers
25 identified by compound numbers 5135 and 6406 (SEQ ID NOS. 308 and 312, respectively), as indicated in the table. The 2086 gene from strain CDC1573 can be amplified with primers

-86-

identified by compound numbers 5135 and 6474 (SEQ ID NOS. 308 and 316, respectively). The 2086 gene from strain 2996 can be amplified with primers identified by compound numbers 6406 and 6605 (SEQ ID NOS. 312 and 320, respectively).

5 Features of these primers include, a synthetic BglII restriction site in each primer, a synthetic NdeI restriction site in compound numbers 6406 and 6474 and termination codons in all three reading frames are present in compound numbers 5135 and 6605. Primer numbers 6406 and 6474 amplify the 2086 gene with an ATG (Met) fused to the second amino terminal codon (ACG) representing a single amino acid
10 substitution (replaces TGC Cys) of the mature 2086 polypeptide.

The PCR cloning vector was TOPO-PCR2.1, Invitrogen, Valencia, CA.

15 The vector used to express non-lipidated 2086 protein was pET9a from Novagen, Madison, WI.

The *E.coli* cloning strain was Top10, Invitrogen, Carlsbad, CA.

20 The *E. coli* expression strain was BLR(DE3)pLysS, Novagen, Madison, WI.

The culture media for cloning purposes was Terrific Broth liquid or agar, according to Sambrook *et al.*, with 1% sterile glucose substituted for glycerol, and the appropriate antibiotic (ampicillin or kanamycin).

25 Plasmid purification was with Qiagen Spin Miniprep Kit (Valencia, CA).

Preparation of the production strain or cell line for Non-lipidated 2086 expression:

30 The 2086 gene was amplified by polymerase chain reaction (PCR) [AmpliTaq and ABI 2400 thermal cycler, Applied Biosystems, Foster City, CA] from chromosomal DNA derived from meningococcal strain 8529. The PCR amplification of the 2086 gene utilized two oligonucleotide primers in each reaction identified by compound numbers 6474 and 5135 (SEQ ID NOS. 316 and 308, respectively). The amplified 2086 PCR product was cloned

-87-

directly into the TOPO-PCR2.1 cloning vector and selected on Terrific Broth agar supplemented with 100 µg/ml ampicillin and 20 µg/ml X-Gal. White colonies were selected and grown. Plasmid DNA was prepared using a Qiagen miniprep kit and the plasmids were screened for the PCR fragment insert. PCR insert plasmids were subjected to DNA
5 sequencing (Big Dye chemistry on an ABI377 sequencer, Applied Biosystems, Foster City, CA).

Plasmids exhibiting the correct DNA sequence were digested with BglII restriction enzyme and the BglII fragment was gel purified using a GeneClean II purification kit
10 (Bio101, Carlsbad, CA). The purified BglII fragment was cloned into the BamHI site of the expression vector pET9a. The pET9a/2086 clones were selected on Terrific Broth plates supplemented with 30 µg/ml kanamycin. Kanamycin resistant clones were grown and miniprep plasmid DNA was prepared. The plasmids were screened for the appropriate orientation of the 2086 gene in the BamHI site. Correctly oriented plasmids represent a
15 fusion of the T7-antigen to the amino terminus of 2086 gene (rP2086T7). These rP2086T7 gene fusions were transformed into BLR(DE3)pLysS, selected on Terrific Broth/Kan plates, grown in Terrific Broth and induced to express the rP2086T7 fusion protein with 1 mM IPTG (isopropyl β-D-thiogalactopyranoside). The rP2086T7 fusion protein expressed at high
20 levels.

These fusion plasmids were then subjected to a NdeI restriction digest, which deletes the T7-antigen and links the mature 2086 gene directly to the ATG start provided by the vector. These NdeI deleted plasmids were transformed into Top10 cells and selected on Terrific Broth/Kan plates. Candidate clones were grown and
25 miniprep plasmid DNA was prepared. The plasmid DNA was subjected to DNA sequencing to confirm the deletion and the integrity of the 2086 gene sequence. These plasmids are represented by the plasmid map designated pPX7328 (FIG. 6). Plasmids representing the correct DNA sequence were transformed into BLR(DE3)pLysS, selected on Terrific Broth/Kan plates, grown in Terrific Broth and
30 induced to express the 2086 protein with IPTG. The pET9a vector failed to express the mature 2086 protein, in strain BLR(DE3)pLysS, when the T7-Tag was removed.

Production of Non-lipidated 2086 protein:

Purified plasmid DNA was used to transform the expression strain BLR(DE3)pLysS. BLR(DE3)pLysS cells carrying the plasmids are resistant to kanamycin and can be induced
 5 to express high levels of PorA protein by the addition of 1 mM IPTG. The rP2086T7 fusion protein can be expressed as insoluble inclusion bodies in the *E. coli* cell line BLR(DE3)pLysS at ~40% of total protein. This purified fusion protein was used to immunize mice and generated significant levels of bactericidal antibodies against a heterologous meningococcal strain. (See Table V)

10

2086 Non-lipidated gene mutagenesis:

PCR primer mutagenesis was performed on the 5' end of the 2086 gene. Expression studies are under way to determine if the T7-Tag can be removed while
 15 exhibiting the high expression levels of mature rP2086T7.

Purification of non-lipidated rP2086T7:

E. coli BLR(DE3)pLysS cells expressing non-lipidated rP2086T7 were lysed by microfluidizer in 10mM Hepes-NaOH/5mM EDTA/1mM Pefabloc SC pH 7.4.
 20 The cell lysate was then centrifuged at 18,000xg for 30 minutes. The inclusion body pellet was washed three times with 50mM Tris-HCl/5mM EDTA/1% TritonTM X-100 pH 8 followed by centrifugation each time at 24,000xg for 30 min. The inclusion body pellet was then washed twice with 50mM Tris-HCl/5mM EDTA/1% Zwittergent 3-14 pH 8 followed by centrifugation each time at 24,000xg for 15min. The inclusion
 25 body pellet was then solubilized with 50mM Tris-HCl/5mM EDTA/4M Urea pH 8 for two hours followed by centrifugation to remove insoluble material. The supernatant (solubilized rP2086T7) was split into four equal samples. One sample was adjusted to 50mM Tris-HCl/5mM EDTA/250mM NaCl/2M Urea pH8 (no detergent), one was adjusted to 50mM Tris-HCl/5mM EDTA/250mM NaCl/2M Urea/1% hydrogenated
 30 TritonTM X-100 pH8 (TX-100), one was adjusted to 50mM Tris-HCl/5mM EDTA/250mM NaCl/2M Urea/1% Zwittergent 3-12 pH8 (Z3-12), and one was

-89-

adjusted to 50mM Tris-HCl/5mM EDTA/250mM NaCl/2M Urea/1% Zwittergent 3-14 pH8 (Z3-14) using stock solutions. To remove the urea, samples were dialyzed to completion against the respective buffer containing no urea. The samples were then dialyzed to completion against the respective buffer containing no urea and 60mM NaCl to reduce the NaCl concentration. Insoluble material was removed by centrifugation at 2,000xg for 15 minutes, and the resulting supernatant (refolded rP2086T7) was used for further experiments. Homogeneity of rP2086T7 was found to be 91-95% as determined using Coomassie stained SDS-PAGE and laser densitometry.

Immunogenicity Procedure – As described in Example 2

This purified fusion protein was used to immunize mice and generated significant levels of bactericidal antibodies against a heterologous meningococcal strain. (See Table V below):

TABLE V: Bactericidal titers of mouse antibody raised to rP2086T7

MOUSE SERUM	DESCRIPTION	HETEROLOGOUS STRAIN/ H44/76
AF780 week 6	r2086T7 , 10ug	3200
Week 0 pool	Pre-immune serum	10
AE203 week 6	rLP2086, 10 ug (positive control)*	6400

(* positive control sera generated by immunization of mice with rLP2086)

20 **Example 4**

Development of Chimeric Clones of ORF 2086

-90-

The N-terminal region of the 2086 gene from strain CDC-1573 contains a repeated segment not present in the 2086 gene from strains 8529 and 2996 (see FIG. 8). It appears that this repeated segment is responsible for increased levels of recombinant 2086 protein expression from two *E.coli* based expression systems (pET and pBAD). The recombinant protein expression level from the CDC-1573 2086 gene was significantly better in the pET and pBAD expression systems as compared to the recombinant expression levels from the 2086 gene with strains 8529 and 2996 using the same systems. The N-terminal region of the 2086 gene from all three strains is relatively homologous, except for this repeated segment. Therefore, it is reasonable to assume that by fusing the CDC-1573 N-terminus to the 2086 genes from strains 8529 and 2996, that the recombinant 2086 protein levels expressed from these genes will increase when using the pET and pBAD systems.

Materials and Methods:

Chromosomal DNA from strains 8529 and 2996 was purified and used as a template for PCR amplification of the chimeric 2086 gene. PCR primers with the compound numbers 6721 and 5135 (SEQ ID NOS. 321 and 308, respectively) were used to amplify the chimeric 2086 gene from strain 8529 and PCR primers with the compound numbers 6721 and 6605 (SEQ ID NOS. 321 and 320, respectively) were used to amplify the chimeric 2086 gene from strain 2996. The PCR products were cloned directly into the PCR2.1 TOPO vector from Invitrogen and then screened by DNA sequence analysis to identify an intact chimeric 2086 gene. That gene was then cleaved from the PCR2.1 vector with BglII and the BglII fragment was inserted into the BamHI site of the pET9a plasmid. Plasmid inserts were screened for the appropriate orientation and then subjected to a NdeI digestion. The linear NdeI fragments were self-ligated to achieve the deletion of a small NdeI fragment containing the T7-tag sequence contributed by the pET9a vector. This deletion directly links the T7 promoter to the 5' end of the chimeric 2086 gene. The NdeI deleted plasmid was transformed into *E.coli* strain BL21(DE3) and kanamycin resistant colonies were screened for chimeric 2086 protein expression with IPTG induction.

-91-

Initial studies indicate that the chimeric 2086 gene from strain 2996 expresses about twice as much recombinant protein as compared to the native 2996/2086 gene when expressed in the pET9a system. The pBAD system has not been tested yet.

5

Although only one experiment has been performed, the data indicate that there is an enhanced utility from the chimeric 2086 gene. The generation of CDC-1573 N-terminal fusions to the 2086 genes from strains 8529 and 2996 provides enhanced recombinant 2086 protein expression.

10

Example 5

2086 PCR Screening of *N. meningitidis* strains:

In order to determine the conservation of the 2086 gene among clinical isolates, PCR amplification was performed on 88 *N. meningitidis* strains.

15 Initial PCR identification of ORF 2086 utilized primers listed in Table IV (see Example 2 above) identified by compound numbers: 4623, 4624 and 4625 (SEQ ID NOS. 303, 304 and 305, respectively). These primers were designed based on Sanger's *N. meningitidis* serogroup A sequence. To facilitate screening a large number of strains, internal primers were designed for the 2086 gene. A total of 88 *N.*
20 *meningitidis* strains were screened by PCR with the newly designed internal 2086 primers identified by compound numbers 5005 and 5007 (SEQ ID NOS. 306 and 307). With these primers the applicants were able to identify the 2086 gene from 63 of the 88 (~70%) *N. meningitidis* strains, (see Table VI-A).

25 Expanded regions surrounding the 2086 gene in Sanger's *N. meningitidis* serogroup A sequence and TIGR's *N. meningitidis* serogroup B sequence were examined and aligned. Primers were designed to correspond to regions upstream and downstream of the 2086 gene. The purpose was to utilize these primers to amplify greater than full length 2086 genes from a variety of *N. meningitidis* strains for

-92-

sequence comparison. PCR amplification of one strain (6557), using Compound Nos. 6470 and 6472 (SEQ ID NOS: 313 and 314, respectively), resulted in a low yield of product. The strain 6557 amplified product was cloned and plasmid DNA was submitted for sequence analysis. Results indicated a new type of 2086 gene with
 5 greater sequence variability than had previously been seen. The 2086 gene from strain 6557 was ~75% identical at the amino acid level to the other strains sequenced. Interestingly, strain 6557 was one of the 30% of strains that had previously tested negative by 2086 PCR screening described above.

Internal primers specific to the C-terminal variable regions within strain 6557
 10 were designed. These primers were used to screen for the more variable 2086 gene in the ~30% of strains that had previously tested negative by 2086 PCR screening. All available *N. meningitidis* strains (n = 88) were screened by PCR with these newly identified internal 2086 primers (identified by compound numbers 6495 and 6496; SEQ ID NOS. 159 and 160, respectively). Only the ~30% of *N. meningitidis* strains
 15 that had previously tested negative by PCR for 2086 were PCR positive in this screen. The set of genes amplified from the previously PCR negative (~30%) strains should represent a new type of 2086 gene or a second family of 2086 genes and herein are designated 2086 Subfamily A. The set of 2086 genes amplified from the ~70% of strains with the 8529 derived primers are herein designated Subfamily B.

20 Subfamily A of 2086 genes is exemplified by the odd numbered SEQ ID NOS:1-173 without limitation. Subfamily B of 2086 genes is exemplified, without limitation, by the odd numbered SEQ ID NOS: 175-251

N. meningitidis strains used for PCR amplification studies were selected from the following tables, Table VI-A and Table VI-B. The strains listed in the tables are
 25 provided as examples of *N. meningitidis* strains, without limitation. The strains listed in Table VI-A are classified in 2086 protein Subfamily A and the strains listed in Table VI-B are classified in 2086 protein Subfamily B. The strains listed in each table are grouped by serosubtype. The strains are available from the following four sources as indicated in the table: MPHL-Manchester Public Health Laboratory,
 30 Manchester, UK; RIVM, Bilthoven, The Netherlands; University of Iowa, College of

WO 03/063766

PCT/US02/32369

-93-

Medicine, Department of Microbiology, Iowa City, IA; and Walter Reed Army
Institute of Research, Washington, D.C.

TABLE VI-A

Strain	Serosubtype	Source
M97 251854	B:4z, PI:4	MPHL
M98 250622	B:2b, PI:10	MPHL
M98 250572	B:2b, PI:10	MPHL
M98 250771	B:4z, PI:22,14	MPHL
M98 250732	B:4z, PI:22,14a	MPHL
M98 250809	B:15, PI:7,16	MPHL
M97 252697	B:1, PI:6, P1.18,25	MPHL
M97 252988	B:4, PI:6, P1.18,25,6	MPHL
M97 252976	B:4, PI:6, P1.18,25	MPHL
M97 252153	B:4, PI:6, P1.18,25	MPHL
M97 253248	B:15,PI:7, NT, 16	MPHL
CDC1610	P1:NT 4(15), P1.18-7,16-14	CDC
CDC1521	P1.6,3 2b(4)	CDC
CDC1034	P1.7 4(15)	CDC
L8	P1.7,1 15(4)	Walter Reed
CDC1492	P1.7,1 4(15)	CDC
870446	P1.12a,13	RIVM
CDC2369	P1.(9),14	CDC
6557	P1.(9),14, P1.22a,14a	RIVM
2996	P1.5,2, P1.5a,2c	RIVM
NmB	P1.5,2, P1.5a,2c	UIOWA
L3	P1.5,2	Walter Reed
B16B6	P1.5,2	RIVM
CDC1135		CDC
L5	P1.NT, P1.21-6,1	Walter Reed
L4	P1.21,16	Walter Reed
W135		Walter Reed
C11	C:16,P1.7,1	CDC
Y		Walter Reed

WO 03/063766

PCT/US02/32369

-95-

TABLE VI-B

Strain	Serosubtype	Source
M98 250670	B:1, PI:4	MPHL
M98 250024	B:1, PI:4	MPHL
M97 253524	B:1, PI:4	MPHL
M97 252060	B:1, PI:4	MPHL
M97 251870	B:4z, PI:4	MPHL
M97 251836	B:4z, PI:4	MPHL
M97 251830	B:4z, PI:4	MPHL
M97 251905	B:4z, PI:4	MPHL
M97 251898	B:4z, PI:4	MPHL
M97 251885	B:4z, PI:4	MPHL
M97 251876	B:4z, PI:4	MPHL
M97 251994	B:4z, PI:4	MPHL
M97 251985	B:4z, PI:4	MPHL
M97 251957	B:4z, PI:4	MPHL
M97 251926	B:4z, PI:4	MPHL
M97 252045	B:4z, PI:4	MPHL
M97 252038	B:4z, PI:4	MPHL
M97 252026	B:4z, PI:4	MPHL
M97 252010	B:4z, PI:4	MPHL
M97 252098	B:4z, PI:4	MPHL
M97 252083	B:4z, PI:4	MPHL
M97 252078	B:4z, PI:4	MPHL
M98 250735	B:4z, PI:15	MPHL
M98 250797	B:4z, PI:15	MPHL
M98 250768	B:4z, PI:15	MPHL
M98 250716	B:2b, PI:10	MPHL
M98 250699	B:4z, PI:10	MPHL
M98 250393	B:4z, PI:10	MPHL
M98 250173	B:4z, PI:10	MPHL
M97 253462	B:4z, PI:14	MPHL
M98 250762	B:15, PI:7,16	MPHL
M98 250610	B:15, PI:7,16	MPHL
M98 250626	B:15, PI:7,16	MPHL
M97 250571	B:15, PI:16	MPHL
M97 252097	B:15, PI:16, P1.7b,16	MPHL
M97 253092	B:1, PI:6	MPHL
M97 252029	B:15, PI:7, NT	MPHL
M97 251875	B:15, PI:7, NT	MPHL

Strain	Serosubtype	Source
CDC1127	PI.7,16 4(15)	CDC
CDC982	PI.7,16 4(15)	CDC
CDC1359	PI.7,16 4(15)	CDC
CDC798	PI.7,16 15(4)	CDC
CDC1078	PI.7,16 15(4)	CDC
CDC1614	PI.7,16 15(4)	CDC
CDC1658	PI.7,16 15(4)	CDC
H44/76	PI.7,16 15(4)	RIVM
CDC1985	P1.7,13 4(15)	CDC
L6	P1.7,1 ?(4)	Walter Reed
CDC1573	P1.7,1 4(15)	CDC
L7	P1.7,(9),1	Walter Reed
CDC937	P1.7,3, P1.7b,3	CDC
8529	P1.7,3, P1.7b,3	RIVM
880049	P1.7b,4	RIVM
CDC2367	P1.15 4(15)	CDC
H355	P1.19,15	RIVM
CDC1343	P1.14 4(15)	CDC
M982	P1.22,9	RIVM
870227	P1.5c,10	RIVM
B40	P1.5c,10	RIVM
5315	P1.5c,10	RIVM
CDC983	P1.5,2	CDC
CDC852	P1.5,2	CDC
6940	P1.18,25 (6)	RIVM
A4		

Other strains are readily available as isolates from infected individuals.

Example 6

Reactivity of rLP2086 antisera against meningococcal strains:

- 5 The following table, Table VII, shows the cross-reactive and cross protection capacity of the rLP2086 as described above. As indicated in the table, the rLP2086 was processed and analyzed using a variety of techniques including whole cell ELISA (WCE) titers, bactericidal assay (BCA) and Infant Rat (IR) assays to determine the

bacterial cell surface reactivity of a polyclonal antibody raised against the 2086 protein.

TABLE VII

Reactivity of rLP2086-8529 antisera against multiple meningococcal strains

Strain	Serosubtype	WCE	BC	IR
2086 Subfamily A				
870446	P1.12a,13	808,615	>800	
NmB	P1.5a,2c	47,954	<100	
6557	P1.22a,14a	169,479	<25	-
2086 Subfamily B				
880049	P1.7b,4	1,402,767	100	+
H44/76	P1.7,16	8,009,507	>6400	
H355	P1.19,15	10,258,475	3,200	+
6940	P1.18,25(6)	5,625,410	800	
870227	P1.5c,10	4,213,324	<25	+
252097	P1.7b,16	10,354,512	>800	
539/8529	P1.7b,3	11,635,737	3,200	
M982	P1.22,9	1,896,800	800	
CDC-1573	P1.7a,1	208,259	25	
CDC-937	P1.7b,(3)	9,151,863	>800	

+ greater than 10 fold reduction in bacteremia

5 - less than 10 fold reduction in bacteremia

Example 7

Various constructs for expressing ORF2086 protein were prepared. The following table, Table VIII, is an r2086 construct table which is provided for the

purpose of showing examples and illustrating an implementation of the present invention, without limitation thereto.

TABLE VIII
r2086 Construct Summary

Construct	Promoter	Leader	Expression	Extraction	Vector	% total Protein
pPX7340	T7	native	Coomassie	sarcosyl soluble	pET27b	2.5% processed lipoprotein
pPX7341	T7	P4	Coomassie	sarcosyl soluble	pET27b	5% processed lipoprotein
pPX7343	Arabinose	P4	Coomassie	sarcosyl soluble	pBAD18 cm	7-10% processed lipoprotein
pPX7325	T7	T7-tag fusion/mature	Coomassie	inclusion bodies	pET9a	40-50% mature protein
pPX7328	T7	mature	Coomassie	soluble	pET9a	10% mature protein

5 **Example 8**

Further studies with LOS depleted outer membrane proteins identified additional strains producing outer membrane protein(s) other than PorA which were capable of eliciting bactericidal antibodies to strains expressing heterologous serosubtypes. The following describes further studies to identify additional proteins according to one embodiment of the present invention, and specifically outer membrane lipoproteins, which can reduce the number of proteins required in a meningococcal immunogenic composition. These further studies supplement the studies described in the previous examples.

Subcellular fractionation, differential detergent extraction, isoelectric focusing, and ion exchange chromatography were used in conjunction with

-99-

immunization and bactericidal assays against multiple strains to identify small groups of proteins of interest. Direct sequencing of the main components indicated that the N-termini were blocked. Internal protein sequences were obtained by direct sequencing of polypeptides derived from chemical and proteolytic digests. The

5 genomic sequence of a group A meningococcal strain was downloaded from the Sanger Center and analyzed by our Bioinformatics group using existing and proprietary algorithms to create a searchable database. The peptide sequence data indicated that ORF2086 was of interest. Primers based on this orf were used to PCR the P2086 gene from strain 8529. Analysis of the gene sequence, the fact that the N-

10 terminus was blocked, and its subcellular location indicated that P2086 is a lipidated outer membrane protein(LP2086). rLP2086-8529 and variants from other meningococcal strains were recombinantly expressed as lipoproteins in E.coli using the *H.influenzae* P4 signal sequence. These recombinant proteins were isolated from E.coli membranes by differential detergent extraction, purified using ion exchange

15 chromatography, and used to immunize mice. Mouse anti-LP2086 sera were able to facilitate bactericidal activity against several different serosubtype strains of *N.meningitidis*. Further analysis of the P2086 genes from many *N. meningitidis* strains showed that these sequences fell into two groups designated Subfamily A and Subfamily B. (See FIG. 12) The antisera raised against the Subfamily B proteins

20 were bactericidal against nine strains expressing Subfamily B proteins, and one strain expressing a Subfamily A protein. Subfamily A antisera were bactericidal against Subfamily A strains. A mixture of one rPorA and one rLP2086 elicited complementary antibodies extending vaccine coverage beyond that induced by either protein alone.

25 These observations lead to the following conclusions. rLP2086 antigens are capable of eliciting bactericidal antibodies against meningococcal strains expressing heterologous PorAs and heterologous P2086 proteins. The P2086 family of antigens may be a useful vaccine or immunogenic either alone or in combination with other neisserial antigens.

-100-

The following describes the foregoing study in detail. A complex mixture of soluble outer membrane proteins (sOMPs) was found to elicit PorA independent bactericidal antibody against strains expressing heterologous PorA proteins. A process of differential detergent extraction, isoelectric focusing and ion exchange chromatography followed by mouse immunization was used to follow the immunologically active components.

At each step, sera was assayed for surface reactivity and bactericidal activity against several strains containing serosubtype antigens that are representative of the worldwide epidemiology of meningococcal disease.

This process of separation and immunization was used to identify a novel cross-reactive immunogenic candidate for Group B *N. meningitidis*.

Generation of PorA deficient strains - The *porA* chromosomal locus was cloned into plasmid pPX7016 from strain 2996. Within the plasmid the *porA* promoter, the S/D box and the first 38 N-terminal codons have been deleted and replaced with a self contained KanR expressing cassette. The plasmids were linearized with restriction enzymes and naturally transformed into the serosubtype strains PI:5,2; PI:9; PI:7,16; PI:15; PI:4; PI:3 & PI:10. Kanamycin resistant transformants were selected and screened for the loss of PorA by serosubtype specific monoclonals in an ELISA.

Bactericidal Assay : See Mountzourous, K.T. and Howell, A.P. Detection of Complement-Mediated Antibody-Dependent Bactericidal Activity in a Fluorescence-Based Serum Bactericidal Assay for Group B *Neisseria meningitidis*. J Clin Microbiol. 2000;38:2878-2884.

Whole Cell Enzyme Linked Immunosorbant Assay (ELISA): *N. meningitidis* whole cell suspensions were diluted to an optical density of 0.1 at 620nm in sterile 0.01M phosphate, 0.137M NaCl, 0.002M KCl (PBS). From this suspension, 0.1mL were added to each well of Nunc Bac T 96 well plates (Cat# 2-69620). Cells were dried on the plates at 37°C overnight, then were covered, inverted and stored at 4°C.

-101-

Plates were washed three times with wash buffer (0.01M Tris-HCl, 0.139M NaCl/KCl, 0.1% Brij-35, pH 7.0-7.4). Dilutions of antisera were prepared in PBS, 0.05% TweenTM-20/Azide and 0.1mL was transferred to the coated plates and incubated for two hours at 37°C. Plates were washed three times in wash buffer.

- 5 Goat-anti-mouse IgG AP (Southern Biotech) was diluted at 1:1500 in PBS/0.05% TweenTM-20, 0.1mL was added to each well, and plates were incubated at 37°C for two hours. Plates were washed (as above). Substrate solution was prepared by diluting p-nitrophenyl phosphate (Sigma) in diethanolamine at 1mg/ml. Substrate was added to the plate at 0.1mL per well and incubated at room temperature for one
- 10 hour. The reaction was stopped with 50ul/well of 3N NaOH and plates were read at 405nm with 690nm reference.

- Recombinant PorA Induction: The BLR(DE3)/pET9a strains were grown overnight at 37°C in HySoy Broth (Sheffield Products) supplemented with Kan-30 and 2% glucose. In the morning the O/N cultures were diluted 1/20 in HySoy Broth
- 15 Kan-30 and 1% glycerol and grown at 37°C for 1 hour. These cultures were induced by the addition of IPTG to a final concentration of 1mM. The cultures were grown for an additional 2-3 hours and then harvested.

- Recombinant PorA Purification: The rPorA was solubilized from *E. coli* inclusion bodies with 8M Urea, and refolded by dialysis against buffer containing no
- 20 urea. The refolded rPorA was then concentrated by diafiltration and buffer exchanged by G25 column into NaPO₄ pH6. The dialyzed rPorA was then run on a cation exchange column (S FractogelTM) and eluted with 1M NaCl.

-102-

The sOMPs from strain 8529 (P1.7-2,3) elicit PorA independent bactericidal activity in mice against strains expressing heterologous serosubtypes. The following table, Table IX, shows the bactericidal activity in the studied strains.

TABLE IX

Test Strain	Serosubtype	BC ₅₀ Titer ¹
539	P1.7-2,3	1280
539 PorA-	NST ²	1080
H44/76	P1.7,16	3285
H44/76 PorA-	NST	2620
H355	P1.19,15	>1350
H355 PorA-	NST	>1350
880049	P1.7-2,4	290
880049 PorA-	NST	85
M982	P1.22,9	85
M982 PorA-	NST	<50

5

Preparation of sOMPs: *N. meningitidis* membranes were extracted with 1 X-100, Zwittergent 3-14, and Zwittergent 3-14+0.5M NaCl. The sOMPs referred to above were solubilized in the Zwittergent 3-14/0.5M NaCl extract. The extraction is performed using techniques well known to persons skilled in the art, for example, see

10 U.S. Patent No. 6,355,253.

Immunogenicity: Female Swiss-Webster mice were immunized with 25ug total protein adjuvanted with 20ug QS-21 at week 0 and 4. An exsanguination bleed and data analysis were done at week 6.

15 ¹ Bactericidal (BC₅₀) titers represented as the reciprocal of the dilution of anti-sera which reduces viable cell count by 50%. Week 0 normal mouse sera had BC₅₀ titers of <25

² NST = Non Serosubtypable

The following table, Table X, shows the purification and characterization summary for recombinant lipidated P2086 (rLP2086) for both Subfamily A and Subfamily B.

5 Subfamily A rLP2086 Purification

TABLE X

rLP2086 Variant	A.A. Homology (%) ¹	Theoretical pI	Purity (%) ²
870446	75	6.1	80
2996	71	5.9	95
M97 252988	71	6.3	96
C11	68	6.4	82
M98 250771	62	6.1	83

10 Subfamily B rLP2086 Purification

TABLE XI

rLP2086 Variant	A.A. Homology (%) ¹	Theoretical pI	Purity (%) ²
8529	100	7.5	96
M982	94	6.3	96
88049	92	6.2	90
CDC1573	87	5.6	93

Purification Method: All variants were solubilized from *E. coli* membranes with TX-100 (exception rLP2086-8529 which was solubilized with Sarcosyl or Urea).

15 Further purification was achieved with a combination of anion exchange (TMAE), size exclusion and/or cation exchange (S Fractogel) chromatography in a Tris-HCl or NaPO₄ buffer.

1 Amino acid homology as compared to P2086 from strain 8529

2 Purity as determined by SDS-PAGE and laser densitometry of colloidal
20 Coomassie stained band (Simply Blue stain)

Immunogenicity of a Subfamily B member, rLP2086-8529, tested against homologous and heterologous strains

Table XII below shows immunogenicity of a Subfamily B member, rLP2086-8529, tested against homologous and heterologous strains

TABLE XII

Target Strain	P2086 Subfamily	Target Strain Serosubtype	A.A. Homology ^a	Whole Cell ELISA ^b Titer	BC ₅₀ Titer ^c
539	B	P1.7-2,3	100	>1,458,000	3,200
H44/76	B	P1.7,16	100	>1,458,000	3,200
H355	B	P1.19,15	100	>1,458,000	3,200
CDC937	B	P1.7-2,3-4	100	>1,458,000	>800
M97 252097	B	P1.7-2,16	100	>1,458,000	>800
870227	B	P1.5-2,10	100	>1,458,000	<25
6940	B	P1.18,25,6	97	900,162	>800
M982	B	P1.22,9	94	435,909	200
880049	B	P1.7-2,4	92	349,912	400
CDC1573	B	P1.7-1,1	87	102,508	25
870446	A	P1.12-1,13	71	389,829	800
M98 250771	A	P1.22,14	62	139,397	<25
NmB	A	P1.5-1,2-2	71	<2,000	<25

- 5 Vaccination Procedure: 6-8 week old female Swiss-Webster mice were immunized with 10ug rLP2086-8529+20ug QS-21 at week 0 and week 4. Data analysis was performed on the week 6 exsanguination bleed.

aAmino acid homology of P2086 as compared to rLP2086-8529

b Endpoint titers expressed as the reciprocal of the dilution at absorbance = 0.1

- 10 c BC50 titers represented as the reciprocal of the dilution of anti-sera which reduces viable cell count by 50%. Week 0 normal mouse sera had BC50 titers of <10

Table XIII shows immunogenicity of a Subfamily B member, rLP2086-2996, tested against homologous and heterologous strains.

TABLE XIII

Target Strain	P2086 Subfamily	Target Strain Serosubtype	A.A. Homology ^a	Whole Cell ELISA ^b Titer	BC ₅₀ Titer ^c
NmB	A	P1.5-1,2-2	99.6	8,979	<25
870446	A	P1.12-1,13	99	<1,458,000	>800
M97 252697	A	P1.18,25,6	98	320,732	>800
6557	A	P1.22-1,14-1	98	17,319	<25
M98 250732	A	P1.22,14-1	89	241,510	>800
M98 250771	A	P1.22,14	89	447,867	800
H44/76	B	P1.7,16	72	56,386	<25

Vaccination Procedure: 6-8 week old female Swiss-Webster mice were
5 immunized with 10ug rLP2086-2996+20ug QS-21 at week 0 and week 4. Data
analysis was performed on the week 6 exsanguination bleed.

aAmino acid homology of P2086 as compared to rLP2086-2996

b Endpoint titers expressed as the reciprocal of the dilution at absorbance = 0.1

c Bactericidal (BC50) titers represented as the reciprocal of the dilution of
10 anti-sera which reduces viable cell count by 50%. Week 0 normal mouse sera had
BC50 titers of <10.

Table XIV below shows that antisera to rLP2086 and rPorA are
complimentary when mixed and assayed for bactericidal activity.

15

TABLE XIV

Antisera	H44/76 (P1.7,16)	NMB (P1.5-1,2-2)	880049 (P1.7-2,4)	H355 (P1.19,15)	870227 (P1.5-2,10)	6557 (P1.22-1,14-1)
Anti-rLP2086 + three rPorA antisera	>3,200	>800	200	>800	200	200
Controls						
anti-rLP2086	6,400	<25	100	3,200	<25	<25
Corresponding monovalent rPorA antisera	-	1,600	-	-	200	400

WO 03/063766

PCT/US02/32369

-106-

Vaccination Procedure: 6-8 week old female Swiss-Webster mice were immunized with either 10ug rLP2086-8529/20ug QS-21, or 15ug rPorA/100ug MPL at week 0 and week 4. Data analysis was performed on the week 6 exsanguination
5 bleed.

a Bactericidal (BC50) titers represented as the reciprocal of the dilution of anti-sera which reduces viable cell count by 50%. Week 0 normal mouse sera had BC50 titers of <10.

The following table, Table XV, shows that mixtures of rLP2086 Subfamilies
10 and two rPorAs elicit bactericidal antibody in mice.

TABLE XV

	H44/76	6940	880049	M982	M98 250771	M98 250732	M97 252697	870446	NmB	6557
	SfB ^b	SfB	SfB	SfB	SfA ^b	SfA	SfA	SfA	SfA	SfA
	P1.7,16	P1.18 25,6	P1.7- 2,4	P1.22 ,9	P1.22,1 4	P1.22,1 4-1	P1.18,2 5,6	P1.12- 1,13	P1.5- 1,2-2	P1.22 -1,14- 1
Antigen										
rLP2086-8529 + rLP2086-2996	>800	>800	200	400	800	>800	>800	>800	-	<25
rLP2086-8529 + rLP2086-2996 + rP1.5-1,2-2 + rP1.22-1,14-1	>800	800	100	200	400	400	>800	>800	>800	200
Monovalent Controls ^c	>800	>800	200	400	800	>800	>800	>800	>800	800

-107-

Vaccination Procedure: 6-8 week old female Swiss-Webster mice were immunized with 10ug of each protein + 20ug QS-21 at week 0 and week 4. Data analysis was performed on the week 6 exsanguination bleed.

5 a Bactericidal (BC50) titers represented as the reciprocal of the dilution of anti-sera which reduces viable cell count by 50%. Week 0 normal mouse sera had BC50 titers of <10.

bSfA – Subfamily A, SfB – Subfamily B

cRelevant monovalent control: rLP2086-8529, rLP2086-2996, rP1.5-1,2-2 or rP1.22-1,14-1 antisera

10 The following summarizes the results of the above described studies. Anti-rLP2086 antisera is bactericidal against 13/16 test strains. Eleven strains expressing different serosubtypes are killed by anti-P2086 sera. Bactericidal activity of anti-rLP2086 sera is complimentary to anti-rPorA sera. Mixtures of P2086 and PorA elicit complimentary bactericidal antibodies in mice. Differential detergent extraction,
15 purification and immunization in conjunction with a functional antibody assay against many strains can be used to identify new vaccine candidates. P2086 has been identified as a vaccine candidate which elicits bactericidal antibody against strains heterologous in both P2086 and rPorA. Thus, the 2086 family of proteins may be a useful vaccine either alone or in combination with other neisserial antigens.

20

Example 9

In accordance with the previous examples, additional meningococcal strains, of varying serogroups, were screened by PCR for the presence of the ORF 2086 gene. Ultimately, one hundred meningococcal strains were screened. The following describes the
25 study and its overall results. These results supplement the data from the previous examples.

Two sets of internal PCR primers specific to the C-terminal variable regions were utilized to discriminate between Subfamily A and B gene sequences. The presence of a PCR

-108-

amplified product of approximately 350 bp indicated that the 2086 gene sequence was present on the chromosome. All strains yielded a single PCR product of the expected size. The nucleotide sequences of fifty-five full-length ORF 2086 genes were determined, aligned (DNASTar MegAlign) and used to generate a phylogenetic tree. (See FIG. 12).

5

Nine of these 2086 genes were recombinantly expressed as a rLP2086 lipoprotein in a pBAD arabinose inducible promoter system and three of these genes were recombinantly expressed as a rP2086 non-lipidated protein in an IPTG inducible pET system. These recombinant proteins were expressed in E.coli B. The purified recombinant protein was used to immunize mice and the mouse antisera was assayed for its serum IgG titers and its bactericidal activity against a variety of heterologous meningococcal strains.

10

ORF 2086 was amplified by PCR from one of the following, whole meningococcal cells, purified chromosomal DNA or plasmid DNA templates.

15

Nine ORF 2086 genes were cloned into the vector pLP339, which fuses the *Haemophilus* P4 leader sequence to the 5' end of the ORF 2086 genes. E.coli strain BLR was used as the host strain for recombinant expression of the lipidated form of rP2086 from the pBAD/ORF 2086 clones. (See FIG. 10A) The pBAD arabinose inducible promoter drives the expression the P4 signal/ORF 2086 fusion protein to express a lipidated form of rP2086. Three P2086 genes, lacking a signal sequence, were cloned into a pET9a vector behind the highly active T7 phage promoter. E.coli strain BL21(DE3) was used as the host strain for recombinant expression of a non-lipidated form of ORF 2086 from the pET9a/ORF 2086 clones. (See FIG. 10B) The DE3 lysogen in E.coli strain BL21 can be induced to express the T7 RNA polymerase under the control of the lacUV5 promoter by addition of IPTG. See, WCE; FEMS Micro. Lett., 48 (1987) 367-371 and BCA; J. Clin. Microbiol., 38 (2000) 2878-2884.

20

25

The gene, ORF2086, was cloned and sequenced from fifty-five different *N. meningitidis* strains. The nucleotide sequences were aligned (DNASTar MegAlign) and used to generate a phylogenetic tree. (See FIG. 12). This tree reveals two distinct subfamilies of

30

-109-

the ORF 2086 gene nucleotide sequence. The two subfamilies of genes are similar at their 5' ends, but contain considerable variation near their 3' ends. Although there appears to be significant variability, certain key regions of the gene are highly homologous amongst the different strains. These conserved regions may provide functional continuity for the protein and may be indicative of cross-protective epitopes to be exploited as vaccine targets.

The 2086 gene was cloned from several serogroup B meningococcal strains and expressed with and without the lipidation signal sequence. Referring to FIGS. 11A and 11B, gel photographs show the whole cell lysates of E.coli B expressing the r2086 protein. The non-lipidated form fused to the T7-Tag expressed at the highest level. The T7-Tag sequence may provide stability to the mRNA and significantly enhances the level of polypeptide translated. This fusion protein appears to deposit in inclusion bodies and can be purified and refolded readily with known protocols. The lipidated and non-lipidated forms of P2086 are expressed at approximately 5 to 8% of total cellular protein, with the exception of the T7-Tag fusions, which express rP2086 as approximately 50% of total protein. The non-lipidated form of the protein appears to be soluble and localized in the cytoplasm. The lipidated form of the protein appears to be associated with the membrane fractions and is solubilized with detergent.

The recombinant lipidated 2086 protein from *N. meningitidis* B strain 8529 consistently elicits greater serum IgG titers than the non-lipidated form (see Table XVI below), which correlates well with the enhanced level of bactericidal activity against both homologous and heterologous meningococcal strains (see Table XVII below). The protein in its native lipidated form may have superior tertiary structure for antigen presentation and/or the attached lipid may act as an adjuvant stimulating a greater immunogenic response.

TABLE XVI

Immune Response Elicited at Week 6 by WCE using
8529 rP2086 (non-lipidated) vs. 8529 rLP2086 (lipidated)

Mouse Sera		Meningococcal Strains				
Antigen	Adjuvant	H44/76	H355	870227	880049	870446

WO 03/063766

PCT/US02/32369

-110-

(10 ug)	(20 ug)					
rP2088	QS-21	273,238	212,947	102,694	69,124	21,466
rLP2086	QS-21	5,384,306	4,819,061	2,930,946	1,307,091	886,056

TABLE XVII

8529 rP2086 Elicits Weaker Bactericidal Activity than 8529 rLP2086

5

Mouse Sera		Meningococcal Strains			
Antigen (10 ug)	Adjuvant (20 ug)	H44/76	H355	880049	NMB
rP2086	QS-21	200	100	<25	<25
rLP2086	QS-21	6,400	3,200	100	<25
Pre- Immune	-	<10	<10	<10	<10
Positive Control	-	1,600	100	200	1,600

The following is a summary of the results of the study. All *N. meningitidis* B strains tested appear to have one 2086-like gene. At least two families of the 2086 gene are represented:

- 10 Subfamily A - about 30% of strains and Subfamily B - about 70% of strains. The 2086 gene has been cloned and sequenced from 55 *N. meningitidis* strains. Sequences within Subfamily A are ~86-100% identical at the DNA level. Sequence within Subfamily B are ~89.5-100% identical at the DNA level. Sequences within Subfamily A vs. Subfamily B ~60.9%-74% identical at the DNA level. 2086 homologs have been identified by PCR screening in the
- 15 following:

N. meningitidis A, B, C, W135, Y

N. lactamica

N. gonorrhoeae FA1090

20

Several ORF 2086 genes have been cloned and recombinantly expressed

Lipidated versions of P2086 were expressed from nine meningococcal strains.

These recombinant proteins have been purified and used to vaccinate mice.

The resulting antisera is bactericidal.

Non-lipidated versions of P2086 were expressed from three of the above nine strains. rLP2086 consistently elicits a greater immune response than rP2086. rLP2086 also exhibits enhanced bactericidal activity against both homologous and heterologous meningococcal strains.

5

Example 10

The following tables, Tables XVIII and XIX, show the characterization of variants of members of the two subfamilies.

10

TABLE XVIII
Subfamily A rLP2086 Variants -
Characterization

		rLP2086- 252988	rLP2086- 250771	rLP2086- 870446	rLP2086- 2996	rLP2086- C11
Growth Media		HySoy	HySoy	HySoy	HySoy	HySoy
Solubility		rTX-100 ⇒ Z3-12	TX-100	TX-100	rTX-100 ⇒ Z3-12	rTX-100 ⇒ Z3-12
Purification Steps		TMAE S Fractogel SEC	HQ Poros SEC	HQ Poros SEC	TMAE SEC	TMAE S Fractogel
Purity (%)		96	83	80	95	82
Yield (mg/g cell pellet)		0.2	0.7	0.8	0.5 (fermentor)	0.1
Size	SEC (Z3- 12)	134,000	155,000	132,000	163,000	126,000
	MS	27,897 (712 lipid)	-	-	27,878 (750 lipid)	28,139 (682 lipid)
Thermal Denaturation Transition Midpoint (T_M) °C		66 °C	-	NT	65 °C	63 °C
Protein Available		2.7mg	1mg (Z3-12)	5.0mg	44mg	1.1mg

-112-

(mg)					
8529 Sequence Homology (%)	71	62	71	72	68

TABLE XIX
Subfamily B rLP2086 Variants -
Characterization

		rLP2086- 8529	rLP2086- M982	rLP2086- 880049	rLP2086- CDC1573
Growth Media		Apollon (Sanford)	Apollon	HySoy	HySoy
Solubility		4M Urea $\Rightarrow$ Z3-12	rTX-100 $\Rightarrow$ Z3-12	rTX-100 $\Rightarrow$ Z3- 12	rTX-100
Purification Steps		TMAE S Fractogel	TMAE S Fractogel	TMAE S Fractogel	TMAE SEC
Purity (%)		96	96	90	93
Yield (mg/g cell pellet)		0.2 (fermentor)	1.6 (fermentor)	0.4	1.0
Size	SEC (Z3-12)	95,000	110,000 150,000	100,000	120,000
	MS	27,785 (822 lipid)	27,719 (711 lipid)	28,044 (819 lipid)	28,385 (823 lipid)
Thermal Denaturation Transition Midpoint (T_M) °C		70 °C	75 °C	62 °C	NT
Protein Available (mg)		Urea – 34mg Sarc – 36mg	Pool 1 – 47mg Pool 2 – 17mg	3.6mg	4.9mg
8529 Sequence Homology (%)		100	94	92	87

Table XX below provides the results of fluorescent serum bactericidal assays for the 2086 Subfamily A.

TABLE XX

5

Description	250771	870446	6557	NMB	M98 250732	M97 252697
rLP2086-252988, 10µg	>800 (99%)*	>800 (99%)*	<25	-	>800 (99%)*	>800 (93%)*
rLP2086-C11, 10µg	200	>880 (91%)*	<25	-	200	400
rLP2086-250771, 10µg	>800 (92%)*	>800 (99%)*	<25	-	>800 (96%)*	>800 (84%)*
rLP2086-870446, 10µg	400	>800 (99%)*	<25	-	400	400
rLP2086-2996, 10µg	800	>800 (99%)*	<25	-	>800 (93%)*	>800 (72%)*
rLP2086-8529 + rLP2086-2996, 10µg	800	>800 (99%)*	<25	-	>800 (80%)*	>800 (72%)*
rLP2086-8529 + rP1.22a,14a + rP1.5a,2c, 10µg	-	800	200	>800 (98%)*	-	-
rLP2086-8529 + rLP2086-2996 + rP1.22a,14a + rP1.5a,2c, 10µg	400	>800 (99%)*	200	>800 (99%)*	400	>800 (88%)*
NMB/rLP2086- 8529 vesicles, 20µg	-	100	-	400	-	-
rP1.22a,14a, 10µg	25	-	800	-	100	-
rP1.5a,2c, 10µg	-	-	-	>800 (99%)*	-	-
rLP2086-8529, 10µg	-	800	-	-	-	-
rP1.22a,14a, 25µg	200	-	-	-	800	-
rP1.18,25.6, 5µg	-	-	-	-	-	-
nP1.22,9 (M982), 25µg	-	-	100	-	-	-
pre-immune mouse serum (negative control)	<10	<10	<10	<10	<10	<10
	800	400	800	1600	**	**

Notes:

-114-

* Percentage indicates the % BC activity at the 1:800 dilution.

** Positive control not available.

– serum not tested

5

Example 11

The following further demonstrates that P2086 is expressed in neisserial strains and provides additional specific examples of P2086 expression in several strains.

Cell lysates were prepared with cells from plate cultures resuspended in SDS sample buffer and heated at 98°C for four minutes. Samples were loaded at ~30-50ug total protein per well on 10-20% pre-cast gels (ICN) and run at 175V. The gels were transferred to a nitrocellulose membrane, which was then blocked for 30min. with 5% powdered milk in Tris-buffered saline (Blotto). The primary antibody used was a pool of polyclonal antisera raised against individual rLP2086 variants in mice.

Referring to FIGS. 17 and 18, a Western Blot shows the reactivity of rLP2086 mouse antisera to P2086 Subfamily A and B whole cell lysates. For the Subfamily A cell lysate blot, the antisera used were raised against rLP2086-2996, -870446 and -250771 with rLP2086-250771 diluted at 1/500 in Blotto and the others diluted at 1/1000 in Blotto. For the Subfamily B cell lysate blot, the antisera used were raised against rLP2086-8529 (diluted 1/1000 in Blotto), -C1XC1573, -M982 and -880049 (these three diluted 1/500 in Blotto). The primary antisera and blot were incubated at 4°C overnight. The blot was washed, a goat-anti-mouseAP secondary was added at 1/500 in Blotto, and the blot was incubated for 30min. at room temperature. After washing, the blot was developed using the BCIP/NBT Membrane Phosphatase Substrate System (KPL).

30

BIBLIOGRAPHY

References referred to herein above are noted below:

1. 1997. Case definitions for Infectious Conditions Under Public Health Surveillance. CDC.
2. 1995 Sambrook, J. and D. W. Russell. 1995. Current Protocols in Molecular Biology. John Wiley & Sons, Inc., New York.
- 5 3. 1994. Griffin, A. M. and Griffin, H. G., ed., Computer Analysis of Sequence Data, Part I. Humana Press, New Jersey.
4. 1993. Smith, D. W.. ed., Biocomputing: Informatics and Genome Projects. Academic Press, New York
5. 1991. Gribskov, M. and Devereux, J., ed. Sequence Analysis Primer.
10 Stockton Press, New York.
6. 1988. Lesk, A. M., ed. Computational Molecular Biology. Oxford University Press, New York.
7. Abdillahi, H., and J. T. Poolman. 1988. *Neisseria meningitidis* group B serosubtyping using monoclonal antibodies in whole-cell ELISA. *Microbial*
15 *Pathogenesis* 4(1):27-32.
8. Achtman, M. 1995. Epidemic spread and antigenic variability of *Neisseria meningitidis*. *Trends in Microbiology* 3(5):186-92.
9. Alm, R. A., L. S. Ling, D. T. Moir, B. L. King, E. D. Brown, P. C. Doig, D. R. Smith, B. Noonan, B. C. Guild, B. L. deJonge, G. Carmel, P. J. Tummino,
20 A. Caruso, M. Uria-Nickelsen, D. M. Mills, C. Ives, R. Gibson, D. Merberg, S. D. Mills, Q. Jiang, D. E. Taylor, G. F. Vovis, and T. J. Trust. 1999. Genomic-sequence comparison of two unrelated isolates of the human gastric pathogen *Helicobacter pylori* [published erratum appears in Nature 1999 Feb 25;397(6721):719]. *Nature*. 397:176-80.

10. Altschul, S. F., T. L. Madden, A. A. Schaffer, J. Zhang, Z. Zhang, W. Miller, and D. J. Lipman. 1997. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res.* 25:3389-402.
11. Anderson, T. F. 1951. Techniques for the preservation of three-
5 dimensional structure in preparing specimens for the electron microscope. *Trans N Y Acad Sci.* 13:130-134.
12. Ambrosch, F., G. Wiedermann, P. Crooy, and A. M. George. 1983. Immunogenicity and side-effects of a new tetravalent meningococcal polysaccharide vaccine. *Bulletin of the World Health Organization* 61(2):317-23.
- 10 13. Benson, G. 1999. Tandem repeats finder: a program to analyze DNA sequences. *Nucleic Acids Res.* 27:573-80.
14. Carillo, H., D. Lipman, and J. Siam. 1988. *Applied Math* 48:1073.
- 15 15. Chen, C. C., and P. P. Cleary. 1989. Cloning and expression of the streptococcal C5a peptidase gene in *Escherichia coli*: linkage to the type 12 M protein gene. *Infect. Immun.* 57:1740-1745.
16. Chmouryguina, I., A. Suvorov, P. Ferrieri, and P. P. Cleary. 1996. Conservation of the C5a peptidase genes in group A and B streptococci. *Infect. Immun.* 64:2387-2390.
17. Cockerill, F. R., 3rd, R. L. Thompson, J. M. Musser, P. M. Schlievert,
20 J. Talbot, K. E. Holley, W. S. Harmsen, D. M. Ilstrup, P. C. Kohner, M. H. Kim, B. Frankfort, J. M. Manahan, J. M. Steckelberg, F. Roberson, and W. R. Wilson. 1998. Molecular, serological, and clinical features of 16 consecutive cases of invasive streptococcal disease. Southeastern Minnesota Streptococcal Working Group. *Clin Infect Dis.* 26:1448-58.
- 25 18. Courtney, H. S., Y. Li, J. B. Dale, and D. L. Hasty. 1994. Cloning, sequencing, and expression of a fibronectin/fibrinogen-binding protein from group A streptococci. *Infect Immun.* 62:3937-46.

19. Cserzo, M., E. Wallin, I. Simon, G. von Heijne, and A. Elofsson. 1997. Prediction of transmembrane alpha-helices in prokaryotic membrane proteins: the dense alignment surface method. *Protein Engineering*. 10:673-6.
20. Cunningham, M. W., and A. Quinn. 1997. Immunological
5 crossreactivity between the class I epitope of streptococcal M protein and myosin. *Adv Exp Med Biol*. 418:887-92.
21. Dale, J. B., R. W. Baird, H. S. Courtney, D. L. Hasty, and M. S. Bronze. 1994. Passive protection of mice against group A streptococcal pharyngeal infection by lipoteichoic acid. *J Infect Dis*. 169:319-23.
- 10 22. Dale, J. B., M. Simmons, E. C. Chiang, and E. Y. Chiang. 1996. Recombinant, octavalent group A streptococcal M protein vaccine. *Vaccine*. 14:944-8.
23. Dale, J. B., R. G. Washburn, M. B. Marques, and M. R. Wessels. 1996. Hyaluronate capsule and surface M protein in resistance to opsonization of group A
15 streptococci. *Infect Immun*. 64:1495-501.
24. Eddy, S. R. 1996. Hidden Markov models. *Cur Opin Struct Bio*. 6:361-5.
25. Ellen, R. P., and R. J. Gibbons. 1972. M protein-associated adherence of *Streptococcus pyogenes* to epithelial surfaces: prerequisite for virulence. *Infect
20 Immun*. 5:826-830.
26. Eng, J. K., A. L. McCormack, and J. R. Yates, 3rd. 1994. An approach to correlate tandem mass-spectral data of peptides with amino-acid-sequences in a protein database. *Am Soc Mass Spectrometry*. 5:976-89.
27. Fischetti, V. A., V. Pancholi, and O. Schneewind. 1990. Conservation
25 of a hexapeptide sequence in the anchor region of surface proteins from gram-positive cocci. *Mol Microbiol*. 4:1603-5.
28. Fogg, G. C., and M. G. Caparon. 1997. Constitutive expression of fibronectin binding in *Streptococcus pyogenes* as a result of anaerobic activation of *rofA*. *J Bacteriol*. 179:6172-80.

29. Foster, T. J., and M. Hook. 1998. Surface protein adhesins of *Staphylococcus aureus*. *Trends Microbiol.* 6:484-8.
30. Fraser, C. M., S. Casjens, W. M. Huang, G. G. Sutton, R. Clayton, R. Lathigra, O. White, K. A. Ketchum, R. Dodson, E. K. Hickey, M. Gwinn, B. Dougherty, J. F. Tomb, R. D. Fleischmann, D. Richardson, J. Peterson, A. R. Kerlavage, J. Quackenbush, S. Salzberg, M. Hanson, R. van Vugt, N. Palmer, M. D. Adams, J. 31. Gocayne, J. C. Venter, and *et al.* 1997. Genomic sequence of a Lyme disease spirochaete, *Borrelia burgdorferi* [see comments]. *Nature.* 390:580-6.
32. Goldschneider, I., E. C. Gotschlich, and M. S. Artenstein. 1969. Human immunity to the meningococcus. I. The role of humoral antibodies. *Journal of Experimental Medicine* 129(6):1307-26.
33. Goldschneider, I., E. C. Gotschlich, and M. S. Artenstein. 1969. Human immunity to the meningococcus. II. Development of natural immunity. *Journal of Experimental Medicine* 129(6):1327-48.
34. Gotschlich, E. C., I. Goldschneider, and M. S. Artenstein. 1969. Human immunity to the meningococcus. IV. Immunogenicity of group A and group C meningococcal polysaccharides in human volunteers. *Journal of Experimental Medicine* 129(6):1367-84.
35. Gotschlich, E. C., I. Goldschneider, and M. S. Artenstein. 1969. Human immunity to the meningococcus. V. The effect of immunization with meningococcal group C polysaccharide on the carrier state. *Journal of Experimental Medicine* 129(6):1385-95.
36. Hacker, J., G. Blum-Oehler, I. Muhldorfer, and H. Tschape. 1997. Pathogenicity islands of virulent bacteria: structure, function and impact on microbial evolution. *Mol Microbiol.* 23:1089-97.
37. Hanski, E., and M. Caparon. 1992. Protein F, a fibronectin-binding protein, is an adhesion of the group A streptococcus *Streptococcus pyogenes*. *Proc Natl Acad Sci., USA.* 89:6172-76.

38. Hanski, E., P. A. Horwitz, and M. G. Caparon. 1992. Expression of protein F, the fibronectin-binding protein of *Streptococcus pyogenes* JRS4, in heterologous streptococcal and enterococcal strains promotes their adherence to respiratory epithelial cells. *Infect Immun.* 60:5119-5125.
- 5 39. Hernandez-Sanchez, J., J. G. Valadez, J. V. Herrera, C. Ontiveros, and G. Guarneros. 1998. lambda bar minigene-mediated inhibition of protein synthesis involves accumulation of peptidyl-tRNA and starvation for tRNA. *EMBO Journal.* 17:3758-65.
- 10 40. Huang, T. T., H. Malke, and J. J. Ferretti. 1989. The streptokinase gene of group A streptococci: cloning, expression in *Escherichia coli*, and sequence analysis. *Mol Microbiol.* 3:197-205.
41. Hynes, W. L., A. R. Dixon, S. L. Walton, and L. J. Aridgides. 2000. The extracellular hyaluronidase gene (*hyla*) of *Streptococcus pyogenes*. *FEMS Microbiol Lett.* 184:109-12.
- 15 42. Hynes, W. L., L. Hancock, and J. J. Ferretti. 1995. Analysis of a second bacteriophage hyaluronidase gene from *Streptococcus pyogenes*: evidence for a third hyaluronidase involved in extracellular enzymatic activity. *Infect Immun.* 63:3015-20.
- 20 43. Isberg, R. R., and G. Tran Van Nhieu. 1994. Binding and internalization of microorganisms by integrin receptors. *Trends Microbio.* 2:10-4.
44. Jones, K. F., and V. A. Fischetti. 1988. The importance of the location of antibody binding on the M6 protein for opsonization and phagocytosis of group A M6 streptococci. *J Exp Med.* 167:1114-23.
- 25 45. Kihlberg, B. M., M. Collin, A. Olsen, and L. Bjorck. 1999. Protein H, an antiphagocytic surface protein in *Streptococcus pyogenes*. *Infect Immun.* 67:1708-14.
46. Koebnik, R. 1995. Proposal for a peptidoglycan-associating alpha-helical motif in the C-terminal regions of some bacterial cell-surface proteins [letter; comment]. *Molecular Microbiology.* 16:1269-70.

47. Kuipers, O. P., H. J. Boot, and W. M. de Vos. 1991. Improved site-directed mutagenesis method using PCR. *Nucleic Acids Res.* **19**:4558.
48. Kyte, J., and R. F. Doolittle. 1982. A simple method for displaying the hydropathic character of a protein. *Journal of Molecular Biology* **157**:105-132.
- 5 49. Landt, O., H. P. Grunert, and U. Hahn. 1990. A general method for rapid site-directed mutagenesis using the polymerase chain reaction. *Gene* **96**:125-128.
- 10 50. Loessner, M. J., S. Gaeng, and S. Scherer. 1999. Evidence for a holin-like protein gene fully embedded out of frame in the endolysin gene of *Staphylococcus aureus* bacteriophage 187. *J Bacteriol.* **181**:4452-60.
51. Lukashin, A. V., and M. Borodovsky. 1998. GeneMark.hmm: new solutions for gene finding. *Nucleic Acids Res.* **26**:1107-15.
- 15 52. Lukomski, S., C. A. Montgomery, J. Rurangirwa, R. S. Geske, J. P. Barrish, G. J. Adams, and J. M. Musser. 1999. Extracellular cysteine protease produced by *Streptococcus pyogenes* participates in the pathogenesis of invasive skin infection and dissemination in mice. *Infect Immun.* **67**:1779-88.
53. Madore, D. V. 1998. Characterization of immune response as an indicator of *Haemophilus influenzae* type b vaccine efficacy. *Pediatr Infect Dis J.* **17**:S207-10.
- 20 54. Matsuka, Y. V., S. Pillai, S. Gubba, J. M. Musser, and S. B. Olmsted. 1999. Fibrinogen cleavage by the *Streptococcus pyogenes* extracellular cysteine protease and generation of antibodies that inhibit enzyme proteolytic activity. *Infect Immun.* **67**:4326-33.
- 25 55. Mazmanian, S. K., G. Liu, H. Ton-That, and O. Schneewind. 1999. *Staphylococcus aureus* sortase, an enzyme that anchors surface proteins to the cell wall. *Science.* **285**:760-3.
56. McAtee, C. P., K. E. Fry, and D. E. Berg. 1998. Identification of potential diagnostic and vaccine candidates of *Helicobacter pylori* by "proteome" technologies. *Helicobacter.* **3**:163-9.

57. McAtee, C. P., M. Y. Lim, K. Fung, M. Velligan, K. Fry, T. Chow, and D. E. Berg. 1998. Identification of potential diagnostic and vaccine candidates of *Helicobacter pylori* by two-dimensional gel electrophoresis, sequence analysis, and serum profiling. *Clin Diagn Lab Immunol.* 5:537-42.
- 5 58. McAtee, C. P., M. Y. Lim, K. Fung, M. Velligan, K. Fry, T. P. Chow, and D. E. Berg. 1998. Characterization of a *Helicobacter pylori* vaccine candidate by proteome techniques. *J Chromatogr B Biomed Sci Appl.* 714:325-33.
59. Mejlhede, N., J. F. Atkins, and J. Neuhard. 1999. Ribosomal -1 frameshifting during decoding of *Bacillus subtilis* cdd occurs at the sequence CGA
10 AAG. *J. Bacteriol.* 181:2930-7.
60. Molinari, G., S. R. Talay, P. Valentin-Weigand, M. Rohde, and G. S. Chhatwal. 1997. The fibronectin-binding protein of *Streptococcus pyogenes*, SfbI, is involved in the internalization of group A streptococci by epithelial cells. *Infect Immun.* 65:1357-63.
- 15 61. Mountzouros, K. T., and A. P. Howell. 2000. Detection of complement-mediated antibody-dependent bactericidal activity in a fluorescence-based serum bactericidal assay for group B *Neisseria meningitidis*. *J. Clin. Microbiol.* 38(8):2878-2884.
- 20 62. Nakai, K., and M. Kanehisa. 1991. Expert system for predicting protein localization sites in gram-negative bacteria. *Proteins.* 11:95-110.
63. Navarre, W. W., and O. Schneewind. 1999. Surface proteins of gram-positive bacteria and mechanisms of their targeting to the cell wall envelope. *Microbiol Mol Biol Rev.* 63:174-229.
- 25 64. Nielsen, H., J. Engelbrecht, S. Brunak, and G. von Heijne. 1997. Identification of prokaryotic and eukaryotic signal peptides and prediction of their cleavage sites. *Protein Engineering.* 10:1-6.
65. Nizet, V., B. Beall, D. J. Bast, V. Datta, L. Kilburn, D. E. Low, and J. C. De Azavedo. 2000. Genetic locus for streptolysin S production by group A streptococcus. *Infect Immun.* 68:4245-54.

66. Nordstrand, A., W. M. McShan, J. J. Ferretti, S. E. Holm, and M. Norgren. 2000. Allele substitution of the streptokinase gene reduces the nephritogenic capacity of group A streptococcal strain NZ131. *Infect Immun.* 68:1019-25.
67. Olmsted, S. B., S. L. Erlandsen, G. M. Dunny, and C. L. Wells. 1993.
5 High-resolution visualization by field emission scanning electron microscopy of *Enterococcus faecalis* surface proteins encoded by the pheromone-inducible conjugative plasmid pCF10. *J Bacteriol.* 175:6229-37.
68. Park, J., and S. A. Teichmann. 1998. DIVCLUS: an automatic method
10 in the GEANFAMMER package that finds homologous domains in single- and multi-domain proteins. *Bioinformatics.* 14:144-50.
69. Parkhill, J., M. Achtman, K. D. James, S. D. Bentley, C. Churcher, S. R. Klee, G. Morelli, D. Basham, D. Brown, T. Chillingworth, R. M. Davies, P. Davis, K. Devlin, T. Feltwell, N. Hamlin, S. Holroyd, K. Jagels, S. Leather, S. Moule, K. Mungall, M. A. Quail, M. A. Rajandream, K. M. Rutherford, M. Simmonds, J.
15 Skelton, S. Whitehead, B. G. Spratt, and B. G. Barrell. 2000. Complete DNA sequence of a serogroup A strain of *Neisseria meningitidis* Z2491 [see comments]. *Nature.* 404:502-6.
70. Pierschbacher, M. D., and E. Ruoslahti. 1987. Influence of
20 stereochemistry of the sequence Arg-Gly-Asp-Xaa on binding specificity in cell adhesion. *J Biol Chem.* 262:17294-8.
71. Pizza, M., V. Scarlato, V. Maignani, M. M. Giuliani, B. Arico, M. Comanducci, G. T. Jennings, L. Baldi, E. Bartolini, B. Capecchi, C. L. Galeotti, E. Luzzi, R. Manetti, E. Marchetti, M. Mora, S. Nuti, G. Ratti, L. Santini, S. Savino, M. Scarselli, E. Storni, P. Zuo, M. Broecker, E. Hundt, B. Knapp, E. Blair, T. Mason, H.
25 Tettelin, D. W. Hood, A. C. Jeffries, N. J. Saunders, D. M. Granoff, J. C. Venter, E. R. Moxon, G. Grandi, and R. Rappuoli. 2000. Identification of vaccine candidates against serogroup B meningococcus by whole-genome sequencing. *Science* 287(5459):1816-20.

72. Podbielski, A., A. Flosdorff, and J. Weber-Heynemann. 1995. The group A streptococcal virR49 gene controls expression of four structural vir regulon genes. *Infect Immun.* 63:9-20.
73. Poolman, J. T. 1996. Bacterial outer membrane protein vaccines. The
5 meningococcal example. *Advances in Experimental Medicine & Biology* 397:73-7.
74. Proft, T., S. Louise Moffatt, C. J. Berkahn, and J. D. Fraser. 1999. Identification and Characterization of Novel Superantigens from *Streptococcus pyogenes*. *J Exp Med.* 189:89-102.
75. Pugsley, A. P. 1993. The complete general secretory pathway in gram-
10 negative bacteria. *Microbiol Rev.* 57:50-108.
76. Quinn, A., K. Ward, V. A. Fischetti, M. Hemric, and M. W. Cunningham. 1998. Immunological relationship between the class I epitope of streptococcal M protein and myosin. *Infect Immun.* 66:4418-24.
77. Reda, K. B., V. Kapur, D. Goela, J. G. Lamphear, J. M. Musser, and R.
15 R. Rich. 1996. Phylogenetic distribution of streptococcal superantigen SSA allelic variants provides evidence for horizontal transfer of ssa within *Streptococcus pyogenes*. *Infect Immun.* 64:1161-5.
78. Sambrook, J., and D. W. Russell. 2001. *Molecular cloning a laboratory manual*, Third ed, vol. 3. Cold Spring Harbor Laboratory Press, Cold
20 Spring Harbor, New York.
79. Salzberg, S. L., A. L. Delcher, S. Kasif, and O. White. 1998. Microbial gene identification using interpolated Markov models. *Nucleic Acids Res.* 26:544-8.
80. Saukkonen, K., H. Abdillahi, J. T. Poolman, and M. Leinonen. 1987. Protective efficacy of monoclonal antibodies to class 1 and class 3 outer membrane
25 proteins of *Neisseria meningitidis* B:15:P1.16 in infant rat infection model: new prospects for vaccine development. *Microbial Pathogenesis* 3(4):261-7.
81. Sedegah *et al.* 1994. *Immunology.* 91, 9866-9870.

82. Sonnenberg, M. G., and J. T. Belisle. 1997. Definition of *Mycobacterium tuberculosis* culture filtrate proteins by two-dimensional polyacrylamide gel electrophoresis, N-terminal amino acid sequencing, and electrospray mass spectrometry. *Infect Immun.* 65:4515-24.
- 5 83. Sonnhammer, E. L., S. R. Eddy, and R. Durbin. 1997. Pfam: a comprehensive database of protein domain families based on seed alignments. *Proteins.* 28:405-20.
84. Stevens, D. L. 1995. Streptococcal toxic-shock syndrome: spectrum of disease, pathogenesis, and new concepts in treatment. *Emerg Infect Dis.* 1:69-78.
- 10 85. Stockbauer, K. E., L. Magoun, M. Liu, E. H. Burns, Jr., S. Gubba, S. Renish, X. Pan, S. C. Bodary, E. Baker, J. Coburn, J. M. Leong, and J. M. Musser. 1999. A natural variant of the cysteine protease virulence factor of group A streptococcus with an arginine-glycine-aspartic acid (RGD) motif preferentially binds human integrins $\alpha v \beta 3$ and $\alpha IIb \beta 3$ *Proc Natl Acad Sci., USA.* 96:242-7.
- 15 86. Tettelin, H., N. J. Saunders, J. Heidelberg, A. C. Jeffries, K. E. Nelson, J. A. Eisen, K. A. Ketchum, D. W. Hood, J. F. Peden, R. J. Dodson, W. C. Nelson, M. L. Gwinn, R. DeBoy, J. D. Peterson, E. K. Hickey, D. H. Haft, S. L. Salzberg, O. White, R. D. Fleischmann, B. A. Dougherty, T. Mason, A. Ciecko, D. S. Parksey, E. Blair, H. Cittone, E. B. Clark, M. D. Cotton, T. R. Utterback, H. Khouri, H. Qin, J. Vamathevan, J. Gill, V. Scarlato, V. Masignani, M. Pizza, G. Grandi, L. Sun, H. O. Smith, C. M. Fraser, E. R. Moxon, R. Rappuoli, and J. C. Venter. 2000. Complete genome sequence of *Neisseria meningitidis* serogroup B strain MC58. *Science* 287(5459):1809-15.
- 20 87. Ton-That, H., G. Liu, S. K. Mazmanian, K. F. Faull, and O. Schneewind. 1999. Purification and characterization of sortase, the transpeptidase that cleaves surface proteins of *Staphylococcus aureus* at the LPXTG motif. *Proc Natl Acad Sci U S A.* 96:12424-12429.
- 25

-125-

88. von Heinje, G. 1987. Sequence Analysis in Molecular Biology. Academic Press, New York.

89. Weldingh, K., I. Rosenkrands, S. Jacobsen, P. B. Rasmussen, M. J. Elhay, and P. Andersen. 1998. Two-dimensional electrophoresis for analysis of
5 *Mycobacterium tuberculosis* culture filtrate and purification and characterization of six novel proteins. *Infect Immun.* 66:3492-500.

90. Wolff *et al.* 1990. *Science.* 247, 1465-1468.

91. Yutsudo, T., K. Okumura, M. Iwasaki, A. Hara, S. Kamitani, W. Minamide, H. Igarashi, and Y. Hinuma. 1994. The gene encoding a new mitogenic
10 factor in a *Streptococcus pyogenes* strain is distributed only in group A streptococci. *Infection and Immunity.* 62:4000-4004.

92. Zagursky, R.J. and D. Russell. 2001. Bioinformatics: Use in Bacterial Vaccine Discovery. *BioTechniques.* 31:636-659.

-126-

The scope of the claims should not be limited by the preferred embodiments and examples, but should be given the broadest interpretation consistent with the description as a whole.

What is claimed is:

1. A composition comprising a pharmaceutically acceptable carrier, at least one PorA protein and
 - i) at least one isolated and purified polypeptide consisting of an amino acid sequence having greater than 98% sequence identity to the amino acid sequence of SEQ ID NO: 212, 214 or 216;
or
 - ii) a fusion protein comprising said at least one isolated and purified polypeptide, wherein the at least one isolated and purified polypeptide elicits antibodies that specifically bind ORF2086 subfamily B polypeptide in *Neisseria meningitidis* strains of serogroup B in a non-strain specific manner.
2. A composition comprising a pharmaceutically acceptable carrier, at least one PorA protein and;
 - i) at least one isolated and purified polypeptide having greater than 95% amino acid sequence identity to the amino acid sequence of SEQ ID NO: 212 or 216 and having an amino acid sequence length of 255 amino acids or having greater than 95% amino acid sequence identity to the amino acid sequence of SEQ ID NO: 214 and having an amino acid sequence length of 256 amino acids; or
 - ii) a fusion protein comprising said at least one isolated and purified polypeptide; wherein the said at least one isolated and purified polypeptide elicits antibodies that specifically bind ORF2086 subfamily B polypeptide in *Neisseria meningitidis* strains of serogroup B in a non-strain specific manner.
3. The composition of claim 1 or 2, wherein the at least one isolated and purified polypeptide comprises at most 2 amino acid substitutions compared to the amino acid sequence of SEQ ID NO: 212, 214 or 216.
4. The composition of claim 3, wherein the amino acid substitutions are in the N-terminus.
5. The composition of claim 1 or 2, wherein the at least one isolated and purified polypeptide has greater than 98% amino acid sequence identity to the amino acid sequence of SEQ ID NO: 212.
6. The composition of claim 1 or 2, wherein the at least one isolated and purified polypeptide has greater than 98% amino acid sequence identity to the amino acid sequence of SEQ ID NO: 214.

7. The composition of claim 1 or 2, wherein the at least one isolated and purified polypeptide has greater than 98% amino acid sequence identity to the amino acid sequence of SEQ ID NO: 216.
8. The composition of claim 1 or 2, wherein the isolated and purified polypeptide consists of the amino acid sequence of SEQ ID NO: 212.
9. The composition of claim 1 or 2, wherein the at least one isolated and purified polypeptide consists of the amino acid sequence of SEQ ID NO: 214.
10. The composition of claim 1 or 2, wherein the at least one isolated and purified polypeptide consists of the amino acid sequence of SEQ ID NO: 216.
11. A composition comprising at least one isolated and purified polypeptide comprising an amino acid sequence of SEQ ID NO: 214 or 216, at least one PorA protein and a pharmaceutically acceptable carrier, wherein the at least one isolated and purified polypeptide elicits antibodies that specifically bind ORF2086 subfamily B polypeptide in *Neisseria meningitidis* strains of serogroup B in a non-strain specific manner.
12. The composition of claim 11, wherein the at least one isolated and purified polypeptide comprises the amino acid sequence of SEQ ID NO: 214.
13. The composition of claim 11, wherein the at least one isolated and purified polypeptide comprises the amino acid sequence of SEQ ID NO: 216.
14. The composition of any one of claims 1-13, additionally comprising at least one isolated and purified polypeptide having greater than 95% amino acid sequence identity to the amino acid sequence of any of even numbered SEQ ID NOs: 176-210 and 218-246, wherein the at least one isolated and purified polypeptide is immunogenic.
15. The composition of any one of claims 1-13, additionally comprising at least one isolated and purified polypeptide comprising the amino acid sequence of any of even numbered SEQ ID NOs: 176-210 and 218-246.

16. The composition of any one of claims 1-13, additionally comprising at least one isolated and purified polypeptide having greater than 95% amino acid sequence identity to the amino acid sequence of any of even numbered SEQ ID NOs: 2-54 and 62-174, wherein the at least one isolated and purified polypeptide is immunogenic.

17. The composition of any one of claims 1-13, additionally comprising at least one isolated and purified polypeptide comprising the amino acid sequence of any of even numbered SEQ ID NOs: 2-54 and 62-174.

18. The composition of any one of claims 1-13, wherein said at least one isolated and purified polypeptide is nonpathogenic.

19. The composition of any one of claims 1-13, wherein the at least one isolated and purified polypeptide has a molecular weight of about 26,000 to about 30,000 daltons as measured by mass spectroscopy.

20. The composition of claim 19, wherein the at least one isolated and purified polypeptide has a molecular weight of about 28-35 kDa as measured on a 10%-20% SDS polyacrylamide gel.

21. The composition of any one of claims 1-13, wherein said composition additionally comprises a pharmaceutically acceptable buffer, diluent or adjuvant.

22. The composition of any one of claims 1-13, wherein said composition additionally comprises a diluent.

23. The composition of any one of claims 1-13, wherein said composition additionally comprises an adjuvant.

24. The composition of claim 23, wherein said adjuvant is in a liquid state.

25. The composition of claim 23, wherein the adjuvant is aluminum phosphate or aluminum hydroxide.

26. The composition of any one of claims 1-13, wherein the at least one isolated and purified polypeptide is a recombinant protein.

27. The composition of any one of claims 1-13, wherein the at least one isolated and purified polypeptide is isolated from a native *Neisseria* species.
28. The composition of any one of claims 1-13, wherein the at least one isolated and purified polypeptide is a lipoprotein.
29. The composition of any one of claims 1-13, wherein the at least one isolated and purified polypeptide is non-lipidated.
30. The composition of any one of claims 1-13, wherein the composition additionally comprises at least one isolated *Neisseria* species protein PorB, transferrin binding protein, or opacity protein (Opc).
31. The composition of any one of claims 1-13, wherein the composition additionally comprises at least one additional surface antigen of *Neisseria* species, said additional surface antigen being a non-ORF2086 protein.
32. The composition of any one of claims 1-31, wherein said composition additionally comprises a polysaccharide.
33. The composition of claim 1, wherein said composition comprises an additional peptide or protein, wherein the at least one isolated and purified polypeptide and the additional peptide or protein form a conjugate, wherein the at least one isolated and purified polypeptide elicits antibodies in a mammal that specifically bind ORF2086 subfamily B polypeptide in two or more *Neisseria meningitidis* strains of serogroup B.
34. The composition of claim 33, wherein the at least one isolated and purified polypeptide consists of the amino acid sequence of SEQ ID NO: 214.
35. The composition of claim 33, wherein the at least one isolated and purified polypeptide consists of the amino acid sequence of SEQ ID NO: 212 or 216.
36. A composition comprising a pharmaceutically acceptable carrier, at least one PorA protein and one or more of:

- (a) at least one isolated and purified polypeptide consisting of an amino acid sequence having greater than 98% sequence identity to a polypeptide comprising the amino acid sequence of SEQ ID NO: 214 or 216; or
- (b) at least one isolated and purified polypeptide encoded by a polynucleotide having greater than 98% sequence identity with SEQ ID NO: 213 or 215,

wherein the at least one isolated and purified polypeptide elicits antibodies that specifically bind ORF2086 subfamily B polypeptide in *Neisseria meningitidis* strains of serogroup B in a non-strain specific manner.

37. The composition of claim 36, wherein the at least one isolated and purified polypeptide consists of the amino acid sequence of SEQ ID NO: 214 or 216.

38. The composition of claim 36, wherein the at least one isolated and purified polypeptide comprises the amino acid sequence of SEQ ID NO: 214.

39. The composition of claim 36, wherein the at least one isolated and purified polypeptide comprises the amino acid sequence of SEQ ID NO: 216.

40. The composition of any one of claims 36-39, additionally comprising at least one isolated and purified polypeptide having greater than 95% amino acid sequence identity to the amino acid sequence of any of even numbered SEQ ID NOs: 176-210 and 218-246, wherein the at least one isolated and purified polypeptide is immunogenic.

41. The composition of any one of claims 36-39, additionally comprising at least one isolated and purified polypeptide comprising the amino acid sequence of any of even numbered SEQ ID NOs: 176-210 and 218-246.

42. The composition of any one of claims 36-39, additionally comprising at least one isolated and purified polypeptide having greater than 95% amino acid sequence identity to the amino acid sequence of any of even numbered SEQ ID NOs: 2-54 and 62-174, wherein the at least one isolated and purified polypeptide is immunogenic.

43. The composition of any one of claims 36-39, additionally comprising at least one isolated and purified polypeptide comprising the amino acid sequence of any of even numbered SEQ ID NOs: 2-54 and 62-174.
44. The composition of any one of claims 36-39, wherein the composition additionally comprises at least one isolated *Neisseria* species protein PorB, transferrin binding protein, or opacity protein (Opc).
45. The composition of any one of claims 36-39, wherein the composition additionally comprises at least one additional surface antigen of *Neisseria* species, said additional surface antigen being a non-ORF2086 protein.
46. The composition of any one of claims 36-39, wherein the at least one isolated and purified polypeptide has a molecular weight of about 26,000 to about 30,000 Daltons as measured by mass spectroscopy.
47. The composition of claim 46, wherein the at least one isolated and purified polypeptide has a molecular weight of about 28-35 kDa as measured on a 10%-20% SDS polyacrylamide gel.
48. The composition of any one of claims 36-39, wherein said composition additionally comprises a pharmaceutically acceptable buffer, diluent or adjuvant.
49. The composition of any one of claims 36-39, wherein said composition additionally comprises a diluent.
50. The composition of any one of claims 36-39, wherein said composition additionally comprises an adjuvant.
51. The composition of claim 50, wherein said adjuvant is in a liquid state.
52. The composition of claim 50 or 51, wherein said adjuvant comprises aluminum phosphate or aluminum hydroxide.
53. The composition of any one of claims 36-39 and 48-51, wherein the isolated and purified polypeptide is non-lipidated.

54. The composition of any one of claims 36-39 and 48-51, wherein the isolated and purified polypeptide is a recombinant protein.

55. The composition of any one of claims 36-39, wherein the isolated and purified polypeptide is isolated from native *Neisseria* species.

56. The composition of any one of claims 36-39, wherein the isolated and purified polypeptide is a lipoprotein.

57. The composition of any one of claims 36-39, wherein said composition additionally comprises a polysaccharide.

58. The composition of any one of claims 36-39, wherein said at least one isolated and purified polypeptide is comprised in a conjugate, wherein said at least one isolated and purified polypeptide elicits antibodies in a mammal that specifically bind ORF2086 subfamily B polypeptide in two or more strains of *Neisseria meningitidis* of serogroup B.

59. A composition comprising: a pharmaceutically acceptable carrier, at least one PorA protein and at least one isolated and purified polypeptide from a first bacterial strain of *Neisseria* species having greater than 95% amino acid sequence identity to a polypeptide consisting of the amino acid sequence of SEQ ID NO: 212, 214 or 216 that elicits antibodies that specifically bind ORF2086 subfamily B polypeptide in a second strain of *Neisseria meningitidis* of serogroup B.

60. The composition of claim 59, wherein the at least one isolated and purified polypeptide comprises the amino acid sequence of SEQ ID NO: 214.

61. The composition of claim 59, wherein the at least one isolated and purified polypeptide comprises the amino acid sequence of SEQ ID NO: 216.

62. The composition of claim 59, wherein the at least one isolated and purified polypeptide comprises the amino acid sequence of SEQ ID NO: 212.

63. The composition of any one of claims 59-61, wherein the second strain is any of strains: CDC937, M97 252097, 870227, H355, H4476, 8529, 6940, M982, 880049, M97 253524, M97 251885, M97 251926, M98 250670, and CDC1573.

64. The composition of any one of claims 59-61, wherein the first strain is CDC8529.
65. The composition of any one of claims 59-64, wherein the at least one isolated and purified polypeptide is a recombinant protein.
66. The composition of any one of claims 59-64, wherein the at least one isolated and purified polypeptide is isolated from native *Neisseria* species.
67. The composition of any one of claims 59-66, wherein the at least one isolated and purified polypeptide is a lipoprotein.
68. The composition of any one of claims 59-67, wherein said composition additionally comprises a pharmaceutically acceptable buffer, diluent or adjuvant.
69. The composition of any one of claims 59-67, wherein said composition additionally comprises a diluent.
70. The composition of any one of claims 59-67, wherein said composition additionally comprises an adjuvant.
71. The composition of claim 68 or 70, wherein said adjuvant is in a liquid state.
72. The composition of claim 70 or 71, wherein said adjuvant comprises aluminum phosphate or aluminum hydroxide.
73. The composition of any one of claims 59-66, wherein the at least one isolated and purified polypeptide is non-lipidated.
74. The composition of any one of claims 59-73, wherein said composition additionally comprises a polysaccharide.
75. The composition of any one of claims 59-62, wherein said at least one isolated and purified polypeptide is comprised in a conjugate, wherein said at least one isolated and purified polypeptide

elicits antibodies in a mammal that specifically bind ORF2086 subfamily B polypeptide in two or more strains of *Neisseria meningitidis* of serogroup B.

76. The composition of claim 75, wherein the isolated and purified polypeptide consists of the amino acid sequence of SEQ ID NO: 214.

77. The composition of claim 75, wherein the isolated and purified polypeptide consists of the amino acid sequence of SEQ ID NO: 212 or 216.

78. The composition of any one of claims 59-61, additionally comprising at least one isolated and purified immunogenic polypeptide having greater than 95% amino acid sequence identity to the amino acid sequence of any of even numbered SEQ ID NOs: 176-210 and 218-246.

79. The composition of any one of claims 59-61, additionally comprising at least one isolated and purified polypeptide comprising the amino acid sequence of any of even numbered SEQ ID NOs: 176-210 and 218-246.

80. The composition of any one of claims 59-61, additionally comprising at least one isolated and purified polypeptide having greater than 95% amino acid sequence identity to the amino acid sequence of any of even numbered SEQ ID NOs: 2-54 and 62-174, wherein the at least one isolated and purified polypeptide is immunogenic.

81. The composition of any one of claims 59-61, additionally comprising at least one isolated and purified polypeptide comprising the amino acid sequence of any of even numbered SEQ ID NOs: 2-54 and 62-174.

82. A composition comprising a pharmaceutically acceptable carrier, a nucleic acid molecule that encodes at least one PorA protein and a polynucleotide that encodes at least one polypeptide consisting of the amino acid sequence having at least 98% sequence identity to SEQ ID NO: 212, 214 or 216;

wherein the at least one polypeptide elicits antibodies that specifically bind ORF2086 subfamily B polypeptide in *Neisseria meningitidis* strains of serogroup B in a non-strain specific manner.

83. A composition comprising a pharmaceutically acceptable carrier, a nucleic acid molecule that encodes at least one PorA protein and a polynucleotide that encodes at least one polypeptide consisting of the amino acid sequence having at least 99% sequence identity to SEQ ID NO: 214;

wherein the at least one polypeptide elicits antibodies that specifically bind ORF2086 subfamily B polypeptide in *Neisseria meningitidis* strains of serogroup B in a non-strain specific manner.

84. A composition comprising a pharmaceutically acceptable carrier, a nucleic acid molecule that encodes at least one PorA protein and a polynucleotide that encodes at least one polypeptide consisting of the amino acid sequence having at least 99% sequence identity to SEQ ID NO: 216;

wherein the at least one polypeptide elicits antibodies that specifically bind ORF2086 subfamily B polypeptide in *Neisseria meningitidis* strains of serogroup B in a non-strain specific manner.

85. The composition of claim 82, wherein said at least one polynucleotide consists of the nucleic acid sequence of any of the SEQ ID NOs: 211, 213 or 215.

86. The composition of any one of claims 82-85, wherein said composition comprises a vector, wherein said vector comprises the at least one polynucleotide.

87. The composition of any one of claims 82-86, wherein said composition additionally comprises a nucleic acid sequence encoding for an additional peptide, polypeptide or protein.

88. A composition comprising a pharmaceutically acceptable carrier,

a) a first vector comprising a nucleic acid molecule that encodes at least one PorA protein, and a second vector comprising at least one polynucleotide that encodes at least one polypeptide consisting of an amino acid having greater than 98 % amino acid sequence identity to a polypeptide comprising the amino acid sequence of SEQ ID NO: 212, 214 or 216; or

b) a vector comprising a nucleic acid molecule that encodes at least one PorA protein and at least one polynucleotide that encodes at least one polypeptide consisting of an amino acid having greater than 98 % amino acid sequence identity to a polypeptide comprising the amino acid sequence of SEQ ID NO: 212, 214 or 216;

wherein the at least one polypeptide elicits antibodies that specifically bind ORF2086 subfamily B polypeptide in *Neisseria meningitidis* strains of serogroup B in a non-strain specific manner.

89. The composition of claim 88, wherein the vector comprising the at least one polynucleotide comprises the nucleic acid sequence of any of SEQ ID NOs: 213 or 215.

90. The composition of claim 88 or 89, wherein the vector is a plasmid.

91. The composition of claim 88 or 89, wherein the vector is a phage.

92. The composition of claim 88 or 89, wherein the vector is a bacteriophage.

93. The composition of claim 88 or 89, wherein the vector is a moderate phage.

94. A composition comprising:

(a) a pharmaceutically acceptable carrier;

(b) a nucleic acid that encodes at least one PorA protein; and

(c) a polynucleotide prepared by a process comprising:

isolating and purifying from *Neisseria* species of at least one polynucleotide that encodes at least one polypeptide having greater than 95% amino acid sequence identity to a polypeptide comprising the amino acid sequence of SEQ ID NO: 212, 214 or 216;

wherein the at least one polynucleotide encodes at least one polypeptide that elicits antibodies that specifically bind ORF2086 subfamily B polypeptide in *Neisseria meningitidis* strains of serogroup B in a non-strain specific manner.

95. The composition of claim 94, wherein the at least one polynucleotide comprises the nucleic acid sequence of any of SEQ ID NOs: 211, 213 or 215.

96. The composition of claim 94 or 95, wherein the process further comprises introducing a non-native leader sequence to the at least one polynucleotide.

97. The composition of claim 96, wherein the non-native leader sequence is P4 leader sequence SEQ ID NO: 322.

98. A composition comprising:

(a) a pharmaceutically acceptable carrier;

- (b) at least one PorA protein; and
- (c) an isolated and purified polypeptide prepared by a process comprising:
 - i. inserting into a vector
 - at least one polynucleotide that encodes at least one polypeptide consisting of an amino acid sequence having greater than 95% sequence identity to the amino acid sequence of SEQ ID NO: 212, 214 or 216 or that encodes a fusion protein comprising said at least one polypeptide;
 - ii. introducing the vector comprising the polynucleotide in a cell;
 - iii. expressing the at least one polynucleotide that encodes the at least one polypeptide in the cell; and
 - iv. isolating and purifying the at least one polypeptide;

wherein the at least one isolated and purified polypeptide elicits antibodies that specifically bind ORF2086 subfamily B polypeptide in *Neisseria meningitidis* strains of serogroup B in a non-strain specific manner.

99. The composition of claim 98, wherein the at least one polynucleotide encodes at least one polypeptide consisting of an amino acid sequence having greater than 98% amino acid sequence identity to a polypeptide comprising the amino acid sequence of SEQ ID NO: 214.

100. The composition of claim 98, wherein the at least one polynucleotide encodes at least one polypeptide consisting of an amino acid sequence having greater than 98% amino acid sequence identity to a polypeptide comprising the amino acid sequence of SEQ ID NO: 216.

101. The composition of any one of claims 98-100, wherein the vector is a plasmid.

102. The composition of any one of claims 98-100, wherein the vector is a phage.

103. The composition of any one of claims 98-100, wherein the vector is a bacteriophage.

104. The composition of any one of claims 98-100, wherein the vector is a moderate phage.

105. The composition of any one of claims 98-104, wherein the at least one polynucleotide further comprises a non-native leader sequence.

106. The composition of claim 105, wherein the non-native leader sequence is P4 leader sequence SEQ ID NO: 322.
107. Use of:
- i) a nucleic acid comprising a polynucleotide comprising SEQ ID NO: 211, 213 or 215, said polynucleotide encoding a polypeptide that elicits antibodies that specifically bind ORF2086 subfamily B polypeptide in *Neisseria meningitidis* strains of serogroup B in a non-strain specific manner in a subject; and
 - ii) a polynucleotide that encodes at least one PorA protein for preparing a composition for expression in a host cell.
108. Use of:
- i) a nucleic acid comprising a polynucleotide comprising SEQ ID NO: 211, said polynucleotide encoding a polypeptide that elicits antibodies that specifically bind ORF2086 subfamily B polypeptide in *Neisseria meningitidis* strains of serogroup B in a non-strain specific manner in a subject; and
 - ii) a polynucleotide that encodes at least one PorA protein for preparing a composition for expression in a host cell.
109. The use of claim 107 or 108, wherein the nucleic acid is expressed *in vivo*.
110. The use of claim 107 or 108, wherein the nucleic acid is expressed *in vitro*.
111. The use of any one of claims 107-110, wherein the nucleic acid further comprises a P4 leader sequence having the nucleic acid sequence of SEQ ID NO: 322.
112. A method of preparing a composition comprising:
- (a) isolating and purifying from *N. meningitidis* at least one polynucleotide that encodes at least one polypeptide consisting of an amino acid sequence having greater than 95 % sequence identity to a polypeptide comprising the amino acid sequence of SEQ ID NO: 212, 214 or 216; and
 - (b) combining the at least one polynucleotide with a carrier and a polynucleotide that encodes at least one PorA protein;

wherein the at least one polypeptide elicits antibodies that specifically bind ORF2086 subfamily B polypeptide in *Neisseria meningitidis* strains of serogroup B in a non-strain specific manner.

113. A method of preparing a composition comprising:
isolating and purifying from *Neisseria species* any of the polypeptides that elicit antibodies that specifically bind ORF2086 subfamily B and a PorA protein described in any one of claims 1-81 and 98-106, and adding a pharmaceutically acceptable carrier.
114. A method of preparing an antibody composition comprising:
recovering antibodies from an animal after introducing into the animal a composition according to any one of claims 1-81 and 98-106, wherein the antibody specifically binds to a polypeptide consisting of the amino acid sequence of SEQ ID NO: 212, 214 or 216.
115. Use of an effective amount of the composition of any one of claims 1-81 and 98-106 for eliciting in a mammal antibodies that specifically bind ORF2086 subfamily B polypeptide in *Neisseria meningitidis* strains of serogroup B in a non-strain specific manner.
116. The use of claim 115, wherein said composition is suitable for administration parenterally.
117. The use of claim 115, wherein said composition is suitable for administration mucosally.
118. The use of claim 115, wherein the composition is suitable for administration by subcutaneous or intramuscular injection.
119. Use of:
i) a polynucleotide encoding at least one polypeptide comprising the amino acid sequence of SEQ ID NO: 212, 214 or 216; and
ii) a polynucleotide that encodes at least one PorA protein
for preparing a composition for expression in a host cell;
wherein the at least one polypeptide elicits antibodies that specifically bind ORF2086 subfamily B polypeptide in *Neisseria meningitidis* strains of serogroup B in a non-strain specific manner.
120. The use of claim 119, wherein the polynucleotides are expressed *in vivo*.

121. The use of claim 119, wherein the polynucleotides are expressed *in vitro*.
122. The use of any one of claims 119-121, wherein the polynucleotides are expressed from one or more plasmids.
123. The use of any one of claims 119-121, wherein the polynucleotides are expressed from one or more phages.
124. The use of any one of claims 119-123, wherein the polynucleotide encoding the at least one polypeptide further comprises a non-native leader sequence.
125. The use of claim 124, wherein the non-native leader sequence is P4 leader sequence comprising the nucleic acid sequence of SEQ ID NO: 322.
126. A recombinant cell that expresses:
- i) a nucleic acid molecule that:
 - (a) encodes at least one polypeptide having at least 95% sequence identity to SEQ ID NO: 212, 214 or 216;
 - (b) comprises at least one polynucleotide comprising the nucleic acid sequence of SEQ ID NO: 211, 213 or 215 and encoding at least one polypeptide; or
 - (c) encodes a fusion protein comprising the at least one polypeptide encoded by (a) or (b); and
 - ii) a polynucleotide that encodes at least one PorA protein;
- wherein the at least one polypeptide elicits antibodies that specifically bind ORF2086 subfamily B polypeptide in *Neisseria meningitidis* strains of serogroup B in a non-strain specific manner.
127. The recombinant cell of claim 126, wherein the at least one polypeptide has at least 98% sequence identity to SEQ ID NO: 212 or 214.
128. The recombinant cell of claim 126, wherein the at least one polypeptide has at least 98% sequence identity to SEQ ID NO: 216.
129. The recombinant cell of any one of claims 126-128, wherein the recombinant cell is a hybridoma.

130. The recombinant cell of any one of claims 126-128, wherein the recombinant cell is a trioma.
131. A transformed, transfected or infected cell line comprising:
- (I) a recombinant cell that is transformed, transfected or infected to express a polynucleotide that encodes at least one PorA protein and a nucleic acid molecule comprising:
 - (a) at least one polynucleotide that encodes at least one polypeptide comprising the amino acid sequence of any one of SEQ ID NO: 212, 214 or 216; or
 - (b) at least one polynucleotide comprising the nucleic acid sequence of SEQ ID NO: 211, 213 or 215; or
 - (II) a recombinant cell that expresses a polynucleotide that encodes at least one PorA protein and a nucleic acid molecule encoding:
 - i. at least one polypeptide of (a); or
 - ii. at least one polypeptide comprising the amino acid sequence of SEQ ID NO: 214 or 216;
- wherein the at least one polypeptide elicits antibodies that specifically bind ORF2086 subfamily B polypeptide in *Neisseria meningitidis* strains of serogroup B in a non-strain specific manner.
132. The transformed, transfected or infected cell line of claim 131, wherein the recombinant cell is a hybridoma.
133. The transformed, transfected or infected cell line of claim 131, wherein the recombinant cell is a trioma.
134. The composition of claim 82, wherein the polynucleotide comprises SEQ ID NO: 211 or 213.
135. The composition of claim 36 or 82, wherein the polynucleotide comprises SEQ ID NO: 215.
136. The composition of any one of claims 1-77 or 98, wherein the at least one isolated and purified polypeptide is a CDC8529 bacteria strain polypeptide.
137. The composition of any one of claims 82-84, 94-97 and 99-106, wherein the at least one polypeptide is a CDC8529 bacteria strain polypeptide.

138. The composition of any one of claims 82-84, wherein the polynucleotide is a CDC8529 bacterial species polynucleotide.

139. The composition of any one of claims 98-106, wherein the at least one polynucleotide encodes a CDC8529 bacteria strain polypeptide.

140. The use of any one of claims 119-125, wherein the polynucleotide encoding the at least one polypeptide is a CDC8529 bacterial species polynucleotide.

141. The transformed, transfected or infected cell line of any one of claims 131-133, wherein the at least one polypeptide is a CDC8529 bacteria strain polypeptide.

142. The composition of claim 63 wherein the first strain is CDC8529.

143. The use of any one of claims 107-111 and 119-125, wherein the subject is a human or the host cell is a human cell.

144. The use of any one of claims 115-118, wherein the mammal is a human.

145. The composition of any one of claims 1-31, 33-56, 58-73, 75-81, 98-106, 134-135 and 142 further comprising meningococcus serogroup A, C, Y and W 135 polysaccharides and/or polysaccharide conjugates.

146. Use of a composition comprising:

- i) at least one polypeptide having greater than 95% amino acid sequence identity to a polypeptide comprising the amino acid sequence of SEQ ID NO: 212, 214 or 216;
- ii) at least one PorA protein; and
- iii) a pharmaceutically acceptable carrier;

wherein the at least one polypeptide elicits antibodies that specifically bind ORF2086 subfamily B polypeptide in *Neisseria meningitidis* strains of serogroup B in a non-strain specific manner, for inducing a bactericidal immune response against *Neisseria meningitidis* of serogroup B in a mammal.

147. A composition comprising:
- i) at least one polypeptide having greater than 95% amino acid sequence identity to a polypeptide comprising the amino acid sequence of SEQ ID NO: 212, 214 or 216;
 - ii) at least one PorA protein; and
 - iii) a pharmaceutically acceptable carrier;

wherein the at least one polypeptide is immunogenic, for use in inducing a bactericidal immune response against *Neisseria meningitidis* of serogroup B in a mammal.

148. The use of claim 146 or the composition for use of claim 147, wherein the bactericidal immune response is against two or more strains of *Neisseria meningitidis* serogroup B in a mammal.

149. The use or the composition for use of any one of claims 146-148, wherein the mammal is a human.

150. The use or the composition for use of any one of claims 146-149, wherein the at least one polypeptide has greater than 98% amino acid sequence identity to the amino acid sequence of SEQ ID NO: 212.

151. The use or composition for use of any one of claims 146-149, wherein the at least one polypeptide has greater than 98% amino acid sequence identity to the amino acid sequence of SEQ ID NO: 214.

152. The use or composition for use of any one of claim 146-149, wherein the at least one polypeptide has greater than 98% amino acid sequence identity to the amino acid sequence of SEQ ID NO: 216.

153. The use or composition for use of any one of claims 146-149, wherein the at least one polypeptide comprises the amino acid sequence of SEQ ID NO: 212 or 214.

154. The use or composition for use of any one of claims 146-149, wherein the at least one polypeptide comprises the amino acid sequence of SEQ ID NO: 216.

155. The use or composition for use of any one of claims 146-154, additionally comprising at least one protein having greater than 95% amino acid sequence identity to the amino acid sequence of any

of even numbered SEQ ID NOs: 176-210 and 218-246, wherein the at least one protein elicits antibodies that specifically bind ORF2086 subfamily B polypeptide in *Neisseria meningitidis* strains of serogroup B in a non-strain specific manner.

156. The use or composition for use of any one of claims 146-154, additionally comprising at least one polypeptide comprising the amino acid sequence of any of even numbered SEQ ID NOs: 176-210 and 218-246.

157. The use or composition for use of any one of claims 146-154, additionally comprising at least one protein having greater than 95% amino acid sequence identity to the amino acid sequence of any of even numbered SEQ ID NOs: 2-54 and 62-174, wherein the at least one protein elicits antibodies that specifically bind ORF2086 subfamily B polypeptide in *Neisseria meningitidis* strains of serogroup B in a non-strain specific manner.

158. The use or composition for use of any one of claims 146-154, additionally comprising at least one polypeptide comprising the amino acid sequence of any of even numbered SEQ ID NOs: 2-54 and 62-174.

159. The use or composition for use of any one of claims 146-154, wherein said at least one polypeptide is nonpathogenic.

160. The use or composition for use of any one of claims 146-154, wherein the at least one polypeptide has a molecular weight of about 26,000 to about 30,000 daltons as measured by mass spectroscopy.

161. The use or composition for use of claim 160, wherein the at least one polypeptide has a molecular weight of about 28-35 kDa as measured on a 10%-20% SDS polyacrylamide gel.

162. The use or composition for use of any one of claims 146-154, wherein said composition additionally comprises a pharmaceutically acceptable buffer, diluent or adjuvant.

163. The use or composition for use of any one of claims 146-154, wherein said composition additionally comprises a diluent.

164. The use or composition for use of any one of claims 146-154, wherein said composition additionally comprises an adjuvant.

165. The use or composition for use of claim 164, wherein said adjuvant is in a liquid state.

166. The use or composition for use of claim 164, wherein the adjuvant is aluminum phosphate or aluminum hydroxide.

167. The use or composition for use of any one of claims 146-154, wherein the at least one polypeptide is a recombinant protein.

168. The use or composition for use of any one of claims 146-154, wherein the at least one polypeptide is isolated from native *Neisseria* species.

169. The use or composition for use of any one of claims 146-154, wherein the at least one polypeptide is a lipoprotein.

170. The use or composition for use of any one of claims 146-154, wherein the at least one polypeptide is non-lipidated.

171. The use or composition for use of any one of claims 146-154, wherein the composition additionally comprises at least one isolated *Neisseria* species protein PorB, transferrin binding protein, or opacity protein (Opc).

172. The use or composition for use of any one of claims 146-154, wherein the composition additionally comprises at least one additional surface antigen of *Neisseria* species, said additional surface antigen being a non-ORF2086 protein.

173. The use or composition for use of any one of claims 146-154, wherein said composition additionally comprises a polysaccharide.

174. The use or composition for use of any one of claims 146-148, wherein said composition comprises an additional peptide or protein, wherein the polypeptide and the additional peptide or protein form a conjugate and said polypeptide elicits antibodies that specifically bind ORF2086

subfamily B polypeptide in *Neisseria meningitidis* strains of serogroup B in a non-strain specific manner.

175. The use or composition for use of claim 174, wherein the at least one polypeptide comprises the amino acid sequence of SEQ ID NO: 212.

176. The use or composition for use of claim 174, wherein the at least one polypeptide comprises the amino acid sequence of SEQ ID NO: 214.

177. The use or composition for use of claim 174, wherein the at least one polypeptide comprises the amino acid sequence of SEQ ID NO: 216.

178. A composition comprising at least one polypeptide comprising an amino acid sequence having sequence identity greater than 95% to the amino acid sequence of any one of SEQ ID NOs: 212, 214 and 216, wherein the composition additionally comprises at least one PorA protein, wherein the at least one polypeptide elicits antibodies that specifically bind ORF2086 subfamily B polypeptide in *Neisseria meningitidis* strains of serogroup B in a non-strain specific manner.

179. The composition of claim 178, wherein said at least one polypeptide comprises an amino acid sequence having sequence identity greater than 97% to the amino acid sequence of any one of SEQ ID NOs: 212, 214 and 216.

180. The composition of claim 178, wherein said at least one polypeptide comprises the amino acid sequence of any one of SEQ ID NOs: 212, 214 and 216.

181. The composition according to any one of claims 178-180, wherein said at least one polypeptide is lipidated.

182. The composition according to any one of claims 178-180, wherein said at least one polypeptide is non-lipidated.

183. The composition according to any one of claims 178-182, wherein said at least one polypeptide is a recombinant protein.

184. The composition according to any one of claims 178-183, wherein said at least one polypeptide is a fusion protein.

185. The composition according to any one of claims 178-184, wherein said at least one polypeptide is conjugated to a carrier.

186. The composition according to any one of claims 178-185, wherein said composition additionally comprises a pharmaceutically acceptable buffer, diluent or adjuvant.

187. The composition of claim 186, wherein said adjuvant is aluminum hydroxide or aluminum phosphate.

188. The composition according to any one of claims 178-187, wherein the composition additionally comprises at least one PorB, transferrin binding protein or opacity protein (Opc).

189. The composition according to any one of claims 178-187, wherein the composition additionally comprises at least one additional surface antigen of *Neisseria species* said additional surface antigen being a non-ORF2086 protein.

190. The composition according to any one of claims 178-189, further comprising a polysaccharide.

191. The composition according to any one of claims 178-189, further comprising meningococcus serogroup A, C, Y and W 135 polysaccharides and/or polysaccharide conjugates.

192. The composition according to any one of claims 178-191 for use as a medicament against *N. meningitidis* infection.

193. The composition according to any one of claims 178-191 for use as a vaccine.

194. The composition according to any one of claims 178-191 for use in inducing an immune response in a mammal.

195. The composition according to any one of claims 178-191 for use in ameliorating or preventing *N. meningitidis* infection in a human.

196. The composition according to any one of claims 178-191 for parenteral administration.
197. The composition according to any one of claims 178-191 for subcutaneous administration.
198. The composition according to any one of claims 178-191 for intramuscular administration.
199. A composition comprising:
 - i) a pharmaceutically acceptable diluent;
 - ii) at least one PorA protein; and
 - iii) a polypeptide comprising an amino acid sequence having sequence identity greater than 95% to the amino acid sequence of SEQ ID NO: 212;wherein the polypeptide elicits antibodies that specifically bind to an ORF2086 subfamily B polypeptide from a *Neisseria meningitidis* strain of serogroup B in a non-strain specific manner.
200. The composition of claim 199, wherein said polypeptide comprises an amino acid sequence having sequence identity greater than 97% to the amino acid sequence SEQ ID NO: 212.
201. The composition of claim 199, wherein said polypeptide comprises an amino acid sequence having sequence identity greater than 98% to the amino acid sequence SEQ ID NO: 212.
202. The composition of claim 199, wherein said polypeptide comprises an amino acid sequence having sequence identity greater than 99% to the amino acid sequence SEQ ID NO: 212.
203. The composition of claim 199, wherein said polypeptide is lipidated.
204. The composition of claim 199, wherein said polypeptide is non-lipidated.
205. The composition of any one of claims 199 to 204, wherein said polypeptide is a recombinant protein.
206. The composition of any one of claims 199 to 205, wherein said polypeptide is a fusion protein.
207. The composition of any one of claims 199 to 206, wherein said polypeptide is conjugated to a carrier.

208. The composition of any one of claims 199 to 207, wherein said composition additionally comprises a pharmaceutically acceptable buffer.
209. The composition of any one of claims 199 to 208, wherein said composition additionally comprises an adjuvant.
210. The composition of claim 209, wherein said adjuvant is aluminum hydroxide.
211. The composition of claim 209, wherein said adjuvant is aluminum phosphate.
212. The composition of any one of claims 199 to 211, wherein the composition additionally comprises, at least one additional surface antigen of *Neisseria species* said additional surface antigen being a non-ORF2086 protein.
213. The composition of any one of claims 199 to 212, wherein the composition additionally comprises at least one PorB, transferrin binding protein or opacity protein (Opc).
214. The composition of any one of claims 199 to 213, further comprising a polysaccharide.
215. The composition of any one of claims 199 to 213, further comprising meningococcus serogroup A, C, Y and W 135 polysaccharides and/or polysaccharide conjugates.
216. The composition of any one of claims 199 to 215 for use as a medicament against *N. meningitidis* infection.
217. The composition of any one of claims 199 to 215 for use as a vaccine.
218. The composition of any one of claims 199 to 215 for use in inducing an immune response in a mammal.
219. The composition of any one of claims 199 to 215 for use in inducing a bactericidal immune response in a mammal.
220. The composition of any one of claims 199 to 215 for use in ameliorating or preventing *N. meningitidis* infection in a human.
221. The composition according to any one of claims 199 to 220 for parenteral administration.
222. The composition according to any one of claims 199 to 220 for subcutaneous administration.

223. The composition according to any one of claims 199 to 220 for intramuscular administration.

224. The composition according to any one of claims 199 to 220, wherein the polypeptide elicits bactericidal antibodies that specifically bind to an ORF2086 subfamily B polypeptide from a *Neisseria meningitidis* strain of serogroup B in a non-strain specific manner.

225. A composition comprising:

i) a pharmaceutically acceptable diluent;

ii) at least one PorA protein; and

iii) a polypeptide comprising an amino acid sequence having sequence identity greater than 95% to the amino acid sequence of SEQ ID NO: 214;

wherein the polypeptide elicits antibodies that specifically bind to an ORF2086 subfamily B polypeptide from a *Neisseria meningitidis* strain of serogroup B in a non-strain specific manner.

226. The composition of claim 225, wherein said polypeptide comprises an amino acid sequence having sequence identity greater than 97% to the amino acid sequence SEQ ID NO: 214.

227. The composition of claim 225, wherein said polypeptide comprises an amino acid sequence having sequence identity greater than 98% to the amino acid sequence SEQ ID NO: 214.

228. The composition of claim 225, wherein said polypeptide comprises an amino acid sequence having sequence identity greater than 99% to the amino acid sequence SEQ ID NO: 214.

229. The composition of claim 225, wherein said polypeptide is lipidated.

230. The composition of claim 225, wherein said polypeptide is non-lipidated.

231. The composition of any one of claims 225 to 230, wherein said polypeptide is a recombinant protein.

232. The composition of any one of claims 225 to 231, wherein said polypeptide is a fusion protein.

233. The composition of any one of claims 225 to 232, wherein said polypeptide is conjugated to a carrier.

234. The composition of any one of claims 225 to 233, wherein said composition additionally comprises a pharmaceutically acceptable buffer.
235. The composition of any one of claims 225 to 234, wherein said composition additionally comprises an adjuvant.
236. The composition of claim 235, wherein said adjuvant is aluminum hydroxide.
237. The composition of claim 235, wherein said adjuvant is aluminum phosphate.
238. The composition of any one of claims 225 to 237, wherein the composition additionally comprises, at least one additional surface antigen of *Neisseria species* said additional surface antigen being a non-ORF2086 protein.
239. The composition of any one of claims 225 to 238, wherein the composition additionally comprises at least one PorB, transferrin binding protein or opacity protein (Opc).
240. The composition of any one of claims 225 to 239, further comprising a polysaccharide.
241. The composition of any one of claims 225 to 239, further comprising meningococcus serogroup A, C, Y and W 135 polysaccharides and/or polysaccharide conjugates.
242. The composition of any one of claims 225 to 241 for use as a medicament against *N. meningitidis* infection.
243. The composition of any one of claims 225 to 241 for use as a vaccine.
244. The composition of any one of claims 225 to 241 for use in inducing an immune response in a mammal.
245. The composition of any one of claims 225 to 241 for use in inducing a bactericidal immune response in a mammal.
246. The composition of any one of claims 225 to 241 for use in ameliorating or preventing *N. meningitidis* infection in a human.
247. The composition according to any one of claims 225 to 246 for parenteral administration.
248. The composition according to any one of claims 225 to 246 for subcutaneous administration.

249. The composition according to any one of claims 225 to 246 for intramuscular administration.

250. The composition according to any one of claims 225 to 249, wherein the polypeptide elicits bactericidal antibodies that specifically bind to an ORF2086 subfamily B polypeptide from a *Neisseria meningitidis* strain of serogroup B in a non-strain specific manner.

251. A composition comprising:

i) a pharmaceutically acceptable diluent;

ii) at least one PorA protein; and

iii) a polypeptide comprising an amino acid sequence having sequence identity greater than 95% to the amino acid sequence of SEQ ID NO: 216;

wherein the polypeptide elicits antibodies that specifically bind to an ORF2086 subfamily B polypeptide from a *Neisseria meningitidis* strain of serogroup B in a non-strain specific manner.

252. The composition of claim 251, wherein said polypeptide comprises an amino acid sequence having sequence identity greater than 97% to the amino acid sequence SEQ ID NO: 216.

253. The composition of claim 251, wherein said polypeptide comprises an amino acid sequence having sequence identity greater than 98% to the amino acid sequence SEQ ID NO: 216.

254. The composition of claim 251, wherein said polypeptide comprises an amino acid sequence having sequence identity greater than 99% to the amino acid sequence SEQ ID NO: 216.

255. The composition of claim 251, wherein said polypeptide is lipidated.

256. The composition of claim 251, wherein said polypeptide is non-lipidated.

257. The composition of any one of claims 251 to 256, wherein said polypeptide is a recombinant protein.

258. The composition of any one of claims 251 to 257, wherein said polypeptide is a fusion protein.

259. The composition of any one of claims 251 to 258, wherein said polypeptide is conjugated to a carrier.

260. The composition of any one of claims 251 to 259, wherein said composition additionally comprises a pharmaceutically acceptable buffer.
261. The composition of any one of claims 251 to 260, wherein said composition additionally comprises an adjuvant.
262. The composition of claim 261, wherein said adjuvant is aluminum hydroxide.
263. The composition of claim 261, wherein said adjuvant is aluminum phosphate.
264. The composition of any one of claims 251 to 263, wherein the composition additionally comprises at least one additional surface antigen of *Neisseria species* said additional surface antigen being a non-ORF2086 protein.
265. The composition of any one of claims 251 to 264, wherein the composition additionally comprises at least one PorB, transferrin binding protein or opacity protein (Opc).
266. The composition of any one of claims 251 to 265, further comprising a polysaccharide.
267. The composition of any one of claims 251 to 265, further comprising meningococcus serogroup A, C, Y and W 135 polysaccharides and/or polysaccharide conjugates.
268. The composition of any one of claims 251 to 267 for use as a medicament against *N. meningitidis* infection.
269. The composition of any one of claims 251 to 267 for use as a vaccine.
270. The composition of any one of claims 251 to 267 for use in inducing an immune response in a mammal.
271. The composition of any one of claims 251 to 267 for use in inducing a bactericidal immune response in a mammal.
272. The composition of any one of claims 251 to 267 for use in ameliorating or preventing *N. meningitidis* infection in a human.
273. The composition according to any one of claims 251 to 272 for parenteral administration.
274. The composition according to any one of claims 251 to 272 for subcutaneous administration.

275. The composition according to any one of claims 251 to 272 for intramuscular administration.
276. The composition according to any one of claims 251 to 275, wherein the polypeptide elicits bactericidal antibodies that specifically bind to an ORF2086 subfamily B polypeptide from a *Neisseria meningitidis* strain of serogroup B in a non-strain specific manner.
277. The composition of claim 1 or 2, wherein the composition comprises the fusion protein, wherein the at least one isolated and purified polypeptide of the fusion protein elicits antibodies in a mammal that specifically bind ORF2086 subfamily B polypeptide in two or more *Neisseria meningitidis* strains of serogroup B.

FIG. 1A
Identification of Components in the Unadsorbed TMAE
Fraction: SDS-PAGE Isolation of Peptides
(CNBr Cleavage of Unadsorbed TMAE Fraction followed by SDS-PAGE and N-term
sequencing of fragments from PVDF blot)




FIG. 1B
Identification of Components in the Unadsorbed TMAE
Fraction: Reverse Phase Isolation of Peptides

Enzymatic digestion of unadsorbed TMAE fraction followed by
reverse phase chromatography separation of peptides and direct N-
terminal sequencing

Enzymatic Digest	Retention Time of Peptide (min)	Molecular Weight of Peptide (d)	N-term. ID
Gluc (V8)	6.716	2069.7	P5163
LysC	13.800	3351.2	P4431
LysC	13.800	3351.2	P2086
ArgC	6.860	2278.9	P5163

P4431 (SEQ ID NO: 327)
predicted mw 36,775

P2086 (SEQ ID NO: 212)
predicted mw 27,100

P5163 (SEQ ID NO: 328)
predicted mw 7,081

FIG. 2
Purification of rLP2086



FIG. 3
Identification of Components in the Unadsorbed
TMAE Fraction: LC-MS/MS

SDS-PAGE followed by gel excision, proteolytic digestion, and LC-
MS/MS analysis (Liquid Chromatography tandem Mass Spectrometry)




FIG. 4
Expression of rLP2086




FIG. 5




FIG. 6




FIG. 7




FIG. 8

	10	20	30	
1	C S S G G G G	- - - - -	Y A A D I G A G L A D A L T A P L D	8529 nat. pro
1	C S S G G G G	- - - - -	Y A A D I G A G L A D A L T A P L D	2996 nat. pro
1	C S S G G G G	S G G G G	Y T A D I G T C L A D A L T A P L D	1573 nat. pro

5 amino acid repeat

FIG. 9A

Identification of Immunogenic Component in Nm strain 8529





FIG. 9B



Lipidated rP2086
pBAD18-Arabinose promoter




FIG. 10A

Non-Lipidated rP2086
pET9a-T7 promoter
8529, CDC-1573 and 2996




FIG. 10B

FIG. 11B




FIG. 12

Whole Cell ELISA Data for rLP2086 Subfamily A Antisera




FIG. 13

Whole Cell ELISA Data for rLP2086 Subfamily B
Antisera




FIG. 14

rLP2086 Mixing Study – WCE Titers




FIG. 15

rLP2086/rPorA Mixing Study – WCE Titers





FIG. 16

Western Blot Reactivity of rLP2086 Mouse Antisera to P2086 Subfamily B *N. meningitidis* Whole Cell Lysates



1 – Molecular Weight Marker (kDa)

2 – M97 251985

3 – CDC937

4 – 6940

5 – M97 251926

6 – CDC1573

7 – CDC1359

8 – CDC1658

9 – M97 252026


10 – M97 252029

11 – M982

Subfamily B P2086 cell lysates are all Group B *N. meningitidis*

FIG. 17

Western Blot Reactivity of rLP2086 Mouse Antisera to
P2086 Subfamily A *N. meningitidis* and *N. lactamica*
Whole Cell Lysates



1 – Molecular Weight Markers (kDa)
2 – Group A *N. meningitidis* A4 (P2086
Subfamily B)
3 – Group C *N. meningitidis* - C11
4 – Group Y *N. meningitidis* -
ATCC35561
5 – Group W135 *N. meningitidis* -
ATCC35559
6 – *N. lactamica* - UR5

Group B *N. meningitidis*:
7 – CDC1034
8 – M98 250732
9 – NmB
10 – 6557
11 – CDC1521
12 – M97 252153

FIG. 18