

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
21 January 2010 (21.01.2010)

(10) International Publication Number  
WO 2010/009280 A1

(51) International Patent Classification:  
*A61K 31/00* (2006.01)    *A61P 35/02* (2006.01)  
*A61K 31/404* (2006.01)

CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NL, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(21) International Application Number:  
PCT/US2009/050768

(22) International Filing Date:  
16 July 2009 (16.07.2009)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
61/081,805    18 July 2008 (18.07.2008)    US

(71) Applicant (for all designated States except US): NOVARTIS AG [CH/CH]; Lichtstrasse 35, CH-4056 Basel (CH).

(72) Inventor; and

(75) Inventor/Applicant (for US only): ATADJA, Peter, Wisdom [CA/US]; 9 Reeve Street, Acton, MA 01720 (US).

(74) Agent: HOUGHTON, Gregory, C.; Novartis, Patent Department, One Health Plaza, East Hanover, NJ 07936 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ,

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).


Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))

Published:

- with international search report (Art. 21(3))

(54) Title: USE OF HDAC INHIBITORS FOR THE TREATMENT OF HODGKIN'S DISEASE



(57) Abstract: The present invention relates to the use of an HDAC inhibitor, especially an HDAC inhibitor of formula I: [I], wherein the radicals and symbols have the meanings as defined in the specification, for the preparation of a medicament for the treatment of Hodgkin's disease.

## USE OF HDAC INHIBITORS FOR THE TREATMENT OF HODGKIN'S DISEASE

The present invention relates to the use of an HDAC inhibitor for the preparation of a medicament for the treatment of Hodgkins; a method of treating a warm-blooded animal, especially a human, having Hodgkins disease, comprising administering to said animal a therapeutically effective amount of an HDAC inhibitor, especially a compound of formula (I) as defined herein; and to a pharmaceutical composition and a commercial package comprising said combination.

The term "Hodgkins disease", also known as Hodgkin's lymphoma, as used herein is a type of lymphoma first described by Thomas Hodgkin in 1832. Hodgkin's lymphoma is characterized clinically by the orderly spread of disease from one lymph node group to another and by the development of systemic symptoms with advanced disease.


The compounds of formula (I), as defined herein, are histone deacetylase inhibitors (HDAC inhibitors). Reversible acetylation of histones is a major regulator of gene expression that acts by altering accessibility of transcription factors to DNA. In normal cells, histone deacetylase (HDA) and histone acetyltransferase together control the level of acetylation of histones to maintain a balance. Inhibition of HDA results in the accumulation of hyperacetylated histones, which results in a variety of cellular responses.

Surprisingly, it was now found that HDAC inhibitors, especially the compounds of formula (I), as defined herein, are useful in treatment of Hodgkins disease.

Hence, the invention relates to the use of an HDAC inhibitor for the preparation of a medicament for the treatment of Hodgkins disease.

## HDAC Inhibitor Compounds

HDAC inhibitor compounds of particular interest for use in the inventive combination are hydroxamate compounds described by the formula (I):



wherein

R<sub>1</sub> is H; halo; or a straight-chain C<sub>1</sub>-C<sub>6</sub>alkyl, especially methyl, ethyl or *n*-propyl, which methyl, ethyl and *n*-propyl substituents are unsubstituted or substituted by one or more substituents described below for alkyl substituents;

R<sub>2</sub> is selected from H; C<sub>1</sub>-C<sub>10</sub>alkyl, preferably C<sub>1</sub>-C<sub>6</sub>alkyl, e.g., methyl, ethyl or -CH<sub>2</sub>CH<sub>2</sub>-OH; C<sub>4</sub>-C<sub>9</sub>cycloalkyl; C<sub>4</sub>-C<sub>9</sub>heterocycloalkyl; C<sub>4</sub>-C<sub>9</sub>heterocycloalkylalkyl; cycloalkylalkyl, e.g., cyclopropylmethyl; aryl; heteroaryl; arylalkyl, e.g., benzyl; heteroarylalkyl, e.g., pyridylmethyl; -(CH<sub>2</sub>)<sub>n</sub>C(O)R<sub>6</sub>; -(CH<sub>2</sub>)<sub>n</sub>OC(O)R<sub>6</sub>; amino acyl; HON-C(O)-CH=C(R<sub>1</sub>)-aryl-alkyl-; and -(CH<sub>2</sub>)<sub>n</sub>R<sub>7</sub>;

R<sub>3</sub> and R<sub>4</sub> are the same or different and, independently, H; C<sub>1</sub>-C<sub>6</sub>alkyl; acyl; or acylamino; or

R<sub>3</sub> and R<sub>4</sub>, together with the carbon to which they are bound, represent C=O, C=S or C=NR<sub>8</sub>; or

R<sub>2</sub>, together with the nitrogen to which it is bound, and R<sub>3</sub>, together with the carbon to which it is bound, can form a C<sub>4</sub>-C<sub>9</sub>heterocycloalkyl; a heteroaryl; a polyheteroaryl; a non-aromatic polyheterocycle; or a mixed aryl and non-aryl polyheterocycle ring;

R<sub>5</sub> is selected from H; C<sub>1</sub>-C<sub>6</sub>alkyl; C<sub>4</sub>-C<sub>9</sub>cycloalkyl; C<sub>4</sub>-C<sub>9</sub>heterocycloalkyl; acyl; aryl; heteroaryl; arylalkyl, e.g., benzyl; heteroarylalkyl, e.g., pyridylmethyl; aromatic polycycles; non-aromatic polycycles; mixed aryl and non-aryl polycycles; polyheteroaryl; non-aromatic polyheterocycles; and mixed aryl and non-aryl polyheterocycles;

n, n<sub>1</sub>, n<sub>2</sub> and n<sub>3</sub> are the same or different and independently selected from 0-6, when n<sub>1</sub> is 1-6, each carbon atom can be optionally and independently substituted with R<sub>3</sub> and/or R<sub>4</sub>;

X and Y are the same or different and independently selected from H; halo; C<sub>1</sub>-C<sub>4</sub>alkyl, such as CH<sub>3</sub> and CF<sub>3</sub>; NO<sub>2</sub>; C(O)R<sub>1</sub>; OR<sub>9</sub>; SR<sub>9</sub>; CN; and NR<sub>10</sub>R<sub>11</sub>;

R<sub>6</sub> is selected from H; C<sub>1</sub>-C<sub>6</sub>alkyl; C<sub>4</sub>-C<sub>9</sub>cycloalkyl; C<sub>4</sub>-C<sub>9</sub>heterocycloalkyl; cycloalkylalkyl, e.g., cyclopropylmethyl; aryl; heteroaryl; arylalkyl, e.g., benzyl and 2-phenylethethyl; heteroarylalkyl, e.g., pyridylmethyl; OR<sub>12</sub>; and NR<sub>13</sub>R<sub>14</sub>;

R<sub>7</sub> is selected from OR<sub>15</sub>; SR<sub>15</sub>; S(O)R<sub>16</sub>; SO<sub>2</sub>R<sub>17</sub>; NR<sub>13</sub>R<sub>14</sub>; and NR<sub>12</sub>SO<sub>2</sub>R<sub>6</sub>;

R<sub>8</sub> is selected from H; OR<sub>15</sub>; NR<sub>13</sub>R<sub>14</sub>; C<sub>1</sub>-C<sub>6</sub>alkyl; C<sub>4</sub>-C<sub>9</sub>cycloalkyl; C<sub>4</sub>-C<sub>9</sub>heterocycloalkyl; aryl; heteroaryl; arylalkyl, e.g., benzyl; and heteroarylalkyl, e.g., pyridylmethyl;

R<sub>9</sub> is selected from C<sub>1</sub>-C<sub>4</sub>alkyl, e.g., CH<sub>3</sub> and CF<sub>3</sub>; C(O)-alkyl, e.g., C(O)CH<sub>3</sub>; and C(O)CF<sub>3</sub>;

R<sub>10</sub> and R<sub>11</sub> are the same or different and independently selected from H; C<sub>1</sub>-C<sub>4</sub>alkyl; and -C(O)-alkyl;

R<sub>12</sub> is selected from H; C<sub>1</sub>-C<sub>6</sub>alkyl; C<sub>4</sub>-C<sub>9</sub>cycloalkyl; C<sub>4</sub>-C<sub>9</sub>heterocycloalkyl; C<sub>4</sub>-C<sub>9</sub>heterocycloalkylalkyl; aryl; mixed aryl and non-aryl polycycle; heteroaryl; arylalkyl, e.g., benzyl; and heteroarylalkyl, e.g., pyridylmethyl;

R<sub>13</sub> and R<sub>14</sub> are the same or different and independently selected from H; C<sub>1</sub>-C<sub>6</sub>alkyl; C<sub>4</sub>-C<sub>9</sub>cycloalkyl; C<sub>4</sub>-C<sub>9</sub>heterocycloalkyl; aryl; heteroaryl; arylalkyl, e.g., benzyl; heteroarylalkyl, e.g., pyridylmethyl; amino acyl; or

R<sub>13</sub> and R<sub>14</sub>, together with the nitrogen to which they are bound, are C<sub>4</sub>-C<sub>9</sub>heterocycloalkyl; heteroaryl; polyheteroaryl; non-aromatic polyheterocycle; or mixed aryl and non-aryl polyheterocycle;

R<sub>15</sub> is selected from H; C<sub>1</sub>-C<sub>6</sub>alkyl; C<sub>4</sub>-C<sub>9</sub>cycloalkyl; C<sub>4</sub>-C<sub>9</sub>heterocycloalkyl; aryl; heteroaryl; arylalkyl; heteroarylalkyl; and (CH<sub>2</sub>)<sub>m</sub>ZR<sub>12</sub>;

R<sub>16</sub> is selected from C<sub>1</sub>-C<sub>6</sub>alkyl; C<sub>4</sub>-C<sub>9</sub>cycloalkyl; C<sub>4</sub>-C<sub>9</sub>heterocycloalkyl; aryl; heteroaryl; polyheteroaryl; arylalkyl; heteroarylalkyl; and (CH<sub>2</sub>)<sub>m</sub>ZR<sub>12</sub>;

R<sub>17</sub> is selected from C<sub>1</sub>-C<sub>6</sub>alkyl; C<sub>4</sub>-C<sub>9</sub>cycloalkyl; C<sub>4</sub>-C<sub>9</sub>heterocycloalkyl; aryl; aromatic polycycles; heteroaryl; arylalkyl; heteroarylalkyl; polyheteroaryl and NR<sub>13</sub>R<sub>14</sub>;

m is an integer selected from 0-6; and

Z is selected from O; NR<sub>13</sub>; S; and S(O),

or a pharmaceutically acceptable salt thereof.

As appropriate, "unsubstituted" means that there is no substituent or that the only substituents are hydrogen.

Halo substituents are selected from fluoro, chloro, bromo and iodo, preferably fluoro or chloro.

Alkyl substituents include straight- and branched-C<sub>1</sub>-C<sub>6</sub>alkyl, unless otherwise noted. Examples of suitable straight- and branched-C<sub>1</sub>-C<sub>6</sub>alkyl substituents include methyl, ethyl, *n*-propyl, 2-propyl, *n*-butyl, *sec*-butyl, *t*-butyl and the like. Unless otherwise noted, the alkyl substituents include both unsubstituted alkyl groups and alkyl groups that are substituted by one or more suitable substituents, including unsaturation, i.e., there are one or more double or triple C-C bonds; acyl; cycloalkyl; halo; oxyalkyl; alkylamino; aminoalkyl; acylamino; and OR<sub>15</sub>, e.g., alkoxy. Preferred substituents for alkyl groups include halo, hydroxy, alkoxy, oxyalkyl, alkylamino and aminoalkyl.

Cycloalkyl substituents include C<sub>3</sub>-C<sub>9</sub>cycloalkyl groups, such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and the like, unless otherwise specified. Unless otherwise noted, cycloalkyl substituents include both unsubstituted cycloalkyl groups and cycloalkyl groups that are substituted by one or more suitable substituents, including C<sub>1</sub>-C<sub>6</sub>alkyl, halo, hydroxy, aminoalkyl, oxyalkyl, alkylamino and OR<sub>15</sub>, such as alkoxy. Preferred substituents for cycloalkyl groups include halo, hydroxy, alkoxy, oxyalkyl, alkylamino and aminoalkyl.

The above discussion of alkyl and cycloalkyl substituents also applies to the alkyl portions of other substituents, such as, without limitation, alkoxy, alkyl amines, alkyl ketones, arylalkyl, heteroarylalkyl, alkylsulfonyl and alkyl ester substituents and the like.

Heterocycloalkyl substituents include 3- to 9-membered aliphatic rings, such as 4- to 7-membered aliphatic rings, containing from 1-3 heteroatoms selected from nitrogen, sulfur, oxygen. Examples of suitable heterocycloalkyl substituents include pyrrolidyl, tetrahydrofuryl, tetrahydrothiofuranyl, piperidyl, piperazyl, tetrahydropyranyl, morphilino, 1,3-diazepane, 1,4-diazepane, 1,4-oxazepane and 1,4-oxathiapane. Unless otherwise noted, the rings are unsubstituted or substituted on the carbon atoms by one or more suitable substituents, including C<sub>1</sub>-C<sub>6</sub>alkyl; C<sub>4</sub>-C<sub>9</sub>cycloalkyl; aryl; heteroaryl; arylalkyl, e.g., benzyl; heteroarylalkyl, e.g., pyridylmethyl; halo; amino; alkyl amino and OR<sub>15</sub>, e.g., alkoxy. Unless otherwise noted, nitrogen heteroatoms are unsubstituted or substituted by H, C<sub>1</sub>-C<sub>4</sub>alkyl; arylalkyl, e.g., benzyl; heteroarylalkyl, e.g., pyridylmethyl; acyl; aminoacyl; alkylsulfonyl; and arylsulfonyl.

Cycloalkylalkyl substituents include compounds of the formula  $-(CH_2)_{n5}\text{-cycloalkyl}$ , wherein  $n5$  is a number from 1-6. Suitable alkylcycloalkyl substituents include cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl and the like. Such substituents are unsubstituted or substituted in the alkyl portion or in the cycloalkyl portion by a suitable substituent, including those listed above for alkyl and cycloalkyl.

Aryl substituents include unsubstituted phenyl and phenyl substituted by one or more suitable substituents including  $C_1\text{-}C_6$ alkyl; cycloalkylalkyl, e.g., cyclopropylmethyl;  $O(CO)$ alkyl; oxyalkyl; halo; nitro; amino; alkylamino; aminoalkyl; alkyl ketones; nitrile; carboxyalkyl; alkylsulfonyl; aminosulfonyl; arylsulfonyl and  $OR_{15}$ , such as alkoxy. Preferred substituents include including  $C_1\text{-}C_6$ alkyl; cycloalkyl, e.g., cyclopropylmethyl; alkoxy; oxyalkyl; halo; nitro; amino; alkylamino; aminoalkyl; alkyl ketones; nitrile; carboxyalkyl; alkylsulfonyl; arylsulfonyl and aminosulfonyl. Examples of suitable aryl groups include  $C_1\text{-}C_4$ alkylphenyl,  $C_1\text{-}C_4$ alkoxyphenyl, trifluoromethylphenyl, methoxyphenyl, hydroxyethylphenyl, dimethylaminophenyl, aminopropylphenyl, carbethoxyphenyl, methanesulfonylphenyl and tolylsulfonylphenyl.

Aromatic polycycles include naphthyl, and naphthyl substituted by one or more suitable substituents including  $C_1\text{-}C_6$ alkyl; alkylcycloalkyl, e.g., cyclopropylmethyl; oxyalkyl; halo; nitro; amino; alkylamino; aminoalkyl; alkyl ketones; nitrile; carboxyalkyl; alkylsulfonyl; arylsulfonyl; aminosulfonyl and  $OR_{15}$ , such as alkoxy.

Heteroaryl substituents include compounds with a 5- to 7-membered aromatic ring containing one or more heteroatoms, e.g., from 1-4 heteroatoms, selected from N, O and S. Typical heteroaryl substituents include furyl, thienyl, pyrrole, pyrazole, triazole, thiazole, oxazole, pyridine, pyrimidine, isoxazolyl, pyrazine and the like. Unless otherwise noted, heteroaryl substituents are unsubstituted or substituted on a carbon atom by one or more suitable substituents, including alkyl, the alkyl substituents identified above, and another heteroaryl substituent. Nitrogen atoms are unsubstituted or substituted, e.g., by  $R_{13}$ ; especially useful N substituents include H,  $C_1\text{-}C_4$ alkyl, acyl, aminoacyl and sulfonyl.

Arylalkyl substituents include groups of the formula  $-(CH_2)_{n5}\text{-aryl}$ ,  $-(CH_2)_{n5-1}\text{-}(CH\text{-aryl})\text{-}(CH_2)_{n5}\text{-aryl}$  or  $-(CH_2)_{n5-1}CH(\text{aryl})(\text{aryl})$ , wherein aryl and  $n5$  are defined above. Such arylalkyl substituents include benzyl, 2-phenylethyl, 1-phenylethyl, tolyl-3-propyl, 2-phenylpropyl, diphenylmethyl, 2-diphenylethyl, 5,5-dimethyl-3-phenylpentyl and the like.

Arylalkyl substituents are unsubstituted or substituted in the alkyl moiety or the aryl moiety or both as described above for alkyl and aryl substituents.

Heteroarylalkyl substituents include groups of the formula -(CH<sub>2</sub>)<sub>n5</sub>-heteroaryl, wherein heteroaryl and n5 are defined above and the bridging group is linked to a carbon or a nitrogen of the heteroaryl portion, such as 2-, 3- or 4-pyridylmethyl, imidazolylmethyl, quinolylethyl and pyrrolylbutyl. Heteroaryl substituents are unsubstituted or substituted as discussed above for heteroaryl and alkyl substituents.

Amino acyl substituents include groups of the formula -C(O)-(CH<sub>2</sub>)<sub>n</sub>-C(H)(NR<sub>13</sub>R<sub>14</sub>)-(CH<sub>2</sub>)<sub>n</sub>-R<sub>5</sub>, wherein n, R<sub>13</sub>, R<sub>14</sub> and R<sub>5</sub> are described above. Suitable aminoacyl substituents include natural and non-natural amino acids, such as glycanyl, D-tryptophanyl, L-lysanyl, D- or L-homoserinyl, 4-aminobutyric acyl and  $\pm$ -3-amin-4-hexenoyl.

Non-aromatic polycycle substituents include bicyclic and tricyclic fused ring systems where each ring can be 4- to 9-membered and each ring can contain zero, one or more double and/or triple bonds. Suitable examples of non-aromatic polycycles include decalin, octahydroindene, perhydrobenzocycloheptene and perhydrobenzo-[*f*]-azulene. Such substituents are unsubstituted or substituted as described above for cycloalkyl groups.

Mixed aryl and non-aryl polycycle substituents include bicyclic and tricyclic fused ring systems where each ring can be 4- to 9-membered and at least one ring is aromatic. Suitable examples of mixed aryl and non-aryl polycycles include methylenedioxyphenyl, *bis*-methylenedioxyphenyl, 1,2,3,4-tetrahydronaphthalene, dibenzosuberane, dihydroanthracene and 9*H*-fluorene. Such substituents are unsubstituted or substituted by nitro or as described above for cycloalkyl groups.

Polyheteroaryl substituents include bicyclic and tricyclic fused ring systems where each ring can independently be 5- or 6-membered and contain one or more heteroatom, e.g., 1, 2, 3 or 4 heteroatoms, chosen from O, N or S such that the fused ring system is aromatic. Suitable examples of polyheteroaryl ring systems include quinoline, isoquinoline, pyridopyrazine, pyrrolopyridine, furopyridine, indole, benzofuran, benzothifuran, benzindole, benzoxazole, pyrroloquinoline and the like. Unless otherwise noted, polyheteroaryl substituents are unsubstituted or substituted on a carbon atom by one or more suitable substituents, including alkyl, the alkyl substituents identified above and a substituent of the formula -O-(CH<sub>2</sub>CH=CH(CH<sub>3</sub>)(CH<sub>2</sub>))<sub>1-3</sub>H. Nitrogen atoms are unsubstituted or substituted,

e.g., by R<sub>13</sub>, especially useful N substituents include H, C<sub>1</sub>-C<sub>4</sub>alkyl, acyl, aminoacyl and sulfonyl.

Non-aromatic polyheterocyclic substituents include bicyclic and tricyclic fused ring systems where each ring can be 4- to 9-membered, contain one or more heteroatom, e.g., 1, 2, 3 or 4 heteroatoms, chosen from O, N or S and contain zero or one or more C-C double or triple bonds. Suitable examples of non-aromatic polyheterocycles include hexitol, *cis*-perhydro-cyclohepta[b]pyridinyl, decahydro-benzo[f][1,4]oxazepinyl, 2,8-dioxabicyclo[3.3.0]octane, hexahydro-thieno[3,2-*b*]thiophene, perhydropyrrolo[3,2-*b*]pyrrole, perhydronaphthyridine, perhydro-1*H*-dicyclopenta[*b,e*]pyran. Unless otherwise noted, non-aromatic polyheterocyclic substituents are unsubstituted or substituted on a carbon atom by one or more substituents, including alkyl and the alkyl substituents identified above. Nitrogen atoms are unsubstituted or substituted, e.g., by R<sub>13</sub>, especially useful N substituents include H, C<sub>1</sub>-C<sub>4</sub>alkyl, acyl, aminoacyl and sulfonyl.

Mixed aryl and non-aryl polyheterocycles substituents include bicyclic and tricyclic fused ring systems where each ring can be 4- to 9-membered, contain one or more heteroatom chosen from O, N or S, and at least one of the rings must be aromatic. Suitable examples of mixed aryl and non-aryl polyheterocycles include 2,3-dihydroindole, 1,2,3,4-tetrahydroquinoline, 5,11-dihydro-10*H*-dibenz[*b,e*][1,4]diazepine, 5*H*-dibenzo[*b,e*][1,4]diazepine, 1,2-dihydropyrrolo[3,4-*b*][1,5]benzodiazepine, 1,5-dihydro-pyrido[2,3-*b*][1,4]diazepin-4-one, 1,2,3,4,6,11-hexahydro-benzo[*b*]pyrido[2,3-*e*][1,4]diazepin-5-one. Unless otherwise noted, mixed aryl and non-aryl polyheterocyclic substituents are unsubstituted or substituted on a carbon atom by one or more suitable substituents including -N-OH, =N-OH, alkyl and the alkyl substituents identified above. Nitrogen atoms are unsubstituted or substituted, e.g., by R<sub>13</sub>; especially useful N substituents include H, C<sub>1</sub>-C<sub>4</sub>alkyl, acyl, aminoacyl and sulfonyl.


Amino substituents include primary, secondary and tertiary amines and in salt form, quaternary amines. Examples of amino substituents include mono- and di-alkylamino, mono- and di-aryl amino, mono- and di-arylalkyl amino, aryl-arylalkylamino, alkyl-aryl amino, alkyl-arylalkylamino and the like.

Sulfonyl substituents include alkylsulfonyl and arylsulfonyl, e.g., methane sulfonyl, benzene sulfonyl, tosyl and the like.

Acyl substituents include groups of formula -C(O)-W, -OC(O)-W, -C(O)-O-W or -C(O)NR<sub>13</sub>R<sub>14</sub>, where W is R<sub>16</sub>, H or cycloalkylalkyl.

Acylamino substituents include substituents of the formula -N(R<sub>12</sub>)C(O)-W, -N(R<sub>12</sub>)C(O)-O-W and -N(R<sub>12</sub>)C(O)-NHOH and R<sub>12</sub> and W are defined above.

The R<sub>2</sub> substituent HON-C(O)-CH=C(R<sub>1</sub>)-aryl-alkyl- is a group of the formula



Preferences for each of the substituents include the following:

R<sub>1</sub> is H, halo or a straight-chain C<sub>1</sub>-C<sub>4</sub>alkyl;

R<sub>2</sub> is selected from H, C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>4</sub>-C<sub>9</sub>cycloalkyl, C<sub>4</sub>-C<sub>9</sub>heterocycloalkyl, cycloalkylalkyl, aryl, heteroaryl, arylalkyl, heteroarylalkyl, -(CH<sub>2</sub>)<sub>n</sub>C(O)R<sub>6</sub>, amino acyl and -(CH<sub>2</sub>)<sub>n</sub>R<sub>7</sub>;

R<sub>3</sub> and R<sub>4</sub> are the same or different and independently selected from H and C<sub>1</sub>-C<sub>6</sub>alkyl; or

R<sub>3</sub> and R<sub>4</sub>, together with the carbon to which they are bound, represent C=O, C=S or C=NR<sub>8</sub>;

R<sub>5</sub> is selected from H, C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>4</sub>-C<sub>9</sub>cycloalkyl, C<sub>4</sub>-C<sub>9</sub>heterocycloalkyl, aryl, heteroaryl, arylalkyl, heteroarylalkyl, a aromatic polycycle, a non-aromatic polycycle, a mixed aryl and non-aryl polycycle, polyheteroaryl, a non-aromatic polyheterocycle, and a mixed aryl and non-aryl polyheterocycle;

n, n<sub>1</sub>, n<sub>2</sub> and n<sub>3</sub> are the same or different and independently selected from 0-6, when n<sub>1</sub> is 1-6, each carbon atom is unsubstituted or independently substituted with R<sub>3</sub> and/or R<sub>4</sub>;

X and Y are the same or different and independently selected from H, halo, C<sub>1</sub>-C<sub>4</sub>alkyl, CF<sub>3</sub>, NO<sub>2</sub>, C(O)R<sub>1</sub>, OR<sub>9</sub>, SR<sub>9</sub>, CN and NR<sub>10</sub>R<sub>11</sub>;

R<sub>6</sub> is selected from H, C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>4</sub>-C<sub>9</sub>cycloalkyl, C<sub>4</sub>-C<sub>9</sub>heterocycloalkyl, alkylcycloalkyl, aryl, heteroaryl, arylalkyl, heteroarylalkyl, OR<sub>12</sub> and NR<sub>13</sub>R<sub>14</sub>;

R<sub>7</sub> is selected from OR<sub>15</sub>, SR<sub>15</sub>, S(O)R<sub>16</sub>, SO<sub>2</sub>R<sub>17</sub>, NR<sub>13</sub>R<sub>14</sub> and NR<sub>12</sub>SO<sub>2</sub>R<sub>6</sub>;

R<sub>8</sub> is selected from H, OR<sub>15</sub>, NR<sub>13</sub>R<sub>14</sub>, C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>4</sub>-C<sub>9</sub>cycloalkyl, C<sub>4</sub>-C<sub>9</sub>heterocycloalkyl, aryl, heteroaryl, arylalkyl and heteroarylalkyl;

$R_9$  is selected from  $C_1$ - $C_4$ alkyl and  $C(O)$ -alkyl;

$R_{10}$  and  $R_{11}$  are the same or different and independently selected from H,  $C_1$ - $C_4$ alkyl and  $-C(O)$ -alkyl;

$R_{12}$  is selected from H,  $C_1$ - $C_6$ alkyl,  $C_4$ - $C_9$ cycloalkyl,  $C_4$ - $C_9$ heterocycloalkyl, aryl, heteroaryl, arylalkyl and heteroarylalkyl;

$R_{13}$  and  $R_{14}$  are the same or different and independently selected from H,  $C_1$ - $C_6$ alkyl,  $C_4$ - $C_9$ cycloalkyl,  $C_4$ - $C_9$ heterocycloalkyl, aryl, heteroaryl, arylalkyl, heteroarylalkyl and amino acyl;

$R_{15}$  is selected from H,  $C_1$ - $C_6$ alkyl,  $C_4$ - $C_9$ cycloalkyl,  $C_4$ - $C_9$ heterocycloalkyl, aryl, heteroaryl, arylalkyl, heteroarylalkyl and  $(CH_2)_mZR_{12}$ ;

$R_{16}$  is selected from  $C_1$ - $C_6$ alkyl,  $C_4$ - $C_9$ cycloalkyl,  $C_4$ - $C_9$ heterocycloalkyl, aryl, heteroaryl, arylalkyl, heteroarylalkyl and  $(CH_2)_mZR_{12}$ ;

$R_{17}$  is selected from  $C_1$ - $C_6$ alkyl,  $C_4$ - $C_9$ cycloalkyl,  $C_4$ - $C_9$ heterocycloalkyl, aryl, heteroaryl, arylalkyl, heteroarylalkyl and  $NR_{13}R_{14}$ ;


$m$  is an integer selected from 0-6; and

$Z$  is selected from O,  $NR_{13}$ , S and  $S(O)$ ;

or a pharmaceutically acceptable salt thereof.

Useful compounds of the formula (I), include those wherein each of  $R_1$ ,  $X$ ,  $Y$ ,  $R_3$  and  $R_4$  is H, including those wherein one of  $n_2$  and  $n_3$  is 0 and the other is 1, especially those wherein  $R_2$  is H or  $-CH_2-CH_2-OH$ .

One suitable genus of hydroxamate compounds are those of formula (Ia):



wherein


$n_4$  is 0-3;

$R_2$  is selected from H,  $C_1$ - $C_6$ alkyl,  $C_4$ - $C_9$ cycloalkyl,  $C_4$ - $C_9$ heterocycloalkyl, cycloalkylalkyl, aryl, heteroaryl, arylalkyl, heteroarylalkyl,  $-(CH_2)_nC(O)R_6$ , amino acyl and  $-(CH_2)_nR_7$ ; and

$R_5$  is heteroaryl; heteroarylalkyl, e.g., pyridylmethyl; aromatic polycycles; non-aromatic polycycles; mixed aryl and non-aryl polycycles; polyheteroaryl or mixed aryl; and non-aryl polyheterocycles;

or a pharmaceutically acceptable salt thereof.

Another suitable genus of hydroxamate compounds are those of formula (Ia):



wherein


$n_4$  is 0-3;

$R_2$  is selected from H,  $C_1$ - $C_6$ alkyl,  $C_4$ - $C_9$ cycloalkyl,  $C_4$ - $C_9$ heterocycloalkyl, cycloalkylalkyl, aryl, heteroaryl, arylalkyl, heteroarylalkyl,  $-(CH_2)_nC(O)R_6$ , amino acyl and  $-(CH_2)_nR_7$ ;

$R_5$  is aryl; arylalkyl; aromatic polycycles; non-aromatic polycycles and mixed aryl; and non-aryl polycycles, especially aryl, such as *p*-fluorophenyl, *p*-chlorophenyl, *p*- $O$ - $C_1$ - $C_4$ alkylphenyl, such as *p*-methoxyphenyl, and *p*- $C_1$ - $C_4$ alkylphenyl; and arylalkyl, such as benzyl, *ortho*-, *meta*- or *para*-fluorobenzyl, *ortho*-, *meta*- or *para*-chlorobenzyl, *ortho*-, *meta*- or *para*-mono, di- or tri- $O$ - $C_1$ - $C_4$ alkylbenzyl, such as *ortho*-, *meta*- or *para*-methoxybenzyl, *m,p*-diethoxybenzyl, *o,m,p*-triamethoxybenzyl and *ortho*-, *meta*- or *para*-mono, di- or tri- $C_1$ - $C_4$ alkylphenyl, such as *p*-methyl, *m,m*-diethylphenyl;

or a pharmaceutically acceptable salt thereof.

Another interesting genus is the compounds of formula (Ib):




wherein

$R_2'$  is selected from H;  $C_1$ - $C_6$ alkyl;  $C_4$ - $C_6$ cycloalkyl; cycloalkylalkyl, e.g., cyclopropylmethyl;  $(CH_2)_{2-4}OR_{21}$ , where  $R_{21}$  is H, methyl, ethyl, propyl and *i*-propyl; and

$R_5^{\prime \prime}$  is unsubstituted 1*H*-indol-3-yl, benzofuran-3-yl or quinolin-3-yl, or substituted 1*H*-indol-3-yl, such as 5-fluoro-1*H*-indol-3-yl or 5-methoxy-1*H*-indol-3-yl, benzofuran-3-yl or quinolin-3-yl;

or a pharmaceutically acceptable salt thereof.

Another interesting genus of hydroxamate compounds are the compounds of formula (Ic):



wherein

the ring containing  $Z_1$  is aromatic or non-aromatic, which non-aromatic rings are saturated or unsaturated;

$Z_1$  is O, S or N- $R_{20}$ ;

$R_{18}$  is H; halo;  $C_1$ - $C_6$ alkyl (methyl, ethyl, *t*-butyl);  $C_3$ - $C_7$ cycloalkyl; aryl, e.g., unsubstituted phenyl or phenyl substituted by 4-OCH<sub>3</sub> or 4-CF<sub>3</sub>; or heteroaryl, such as 2-furanyl, 2-thiophenyl or 2-, 3- or 4-pyridyl;

$R_{20}$  is H;  $C_1$ - $C_6$ alkyl;  $C_1$ - $C_6$ alkyl- $C_3$ - $C_9$ cycloalkyl, e.g., cyclopropylmethyl; aryl; heteroaryl; arylalkyl, e.g., benzyl; heteroarylalkyl, e.g., pyridylmethyl; acyl, e.g., acetyl, propionyl and benzoyl; or sulfonyl, e.g., methanesulfonyl, ethanesulfonyl, benzenesulfonyl and toluenesulfonyl;

$A_1$  is 1, 2 or 3 substituents which are independently H;  $C_1$ - $C_6$ alkyl; -OR<sub>19</sub>; halo; alkylamino; aminoalkyl; halo; or heteroarylalkyl, e.g., pyridylmethyl;

$R_{19}$  is selected from H;  $C_1$ - $C_6$ alkyl;  $C_4$ - $C_9$ cycloalkyl;  $C_4$ - $C_9$ heterocycloalkyl; aryl; heteroaryl; arylalkyl, e.g., benzyl; heteroarylalkyl, e.g., pyridylmethyl and -(CH<sub>2</sub>CH=CH(CH<sub>3</sub>)(CH<sub>2</sub>))<sub>1-3</sub>H;

$R_2$  is selected from H,  $C_1$ - $C_6$ alkyl,  $C_4$ - $C_9$ cycloalkyl,  $C_4$ - $C_9$ heterocycloalkyl, cycloalkylalkyl, aryl, heteroaryl, arylalkyl, heteroarylalkyl, -(CH<sub>2</sub>)<sub>n</sub>C(O)R<sub>6</sub>, amino acyl and -(CH<sub>2</sub>)<sub>n</sub>R<sub>7</sub>;

$v$  is 0, 1 or 2;

$p$  is 0-3; and


q is 1-5 and r is 0; or

q is 0 and r is 1-5;

or a pharmaceutically acceptable salt thereof. The other variable substituents are as defined above.

Especially useful compounds of formula (Ic), are those wherein  $R_2$  is H, or  $-(CH_2)_pCH_2OH$ , wherein p is 1-3, especially those wherein  $R_1$  is H; such as those wherein  $R_1$  is H and X and Y are each H, and wherein q is 1-3 and r is 0 or wherein q is 0 and r is 1-3, especially those wherein  $Z_1$  is  $N-R_{20}$ . Among these compounds  $R_2$  is preferably H or  $-CH_2-CH_2-OH$  and the sum of q and r is preferably 1.

Another interesting genus of hydroxamate compounds are the compounds of formula (Id):



wherein

$Z_1$  is O, S or  $N-R_{20}$ ;

$R_{18}$  is H; halo;  $C_1-C_6$ alkyl (methyl, ethyl, *t*-butyl);  $C_3-C_7$ cycloalkyl; aryl, e.g., unsubstituted phenyl or phenyl substituted by 4- $OCH_3$  or 4- $CF_3$ ; or heteroaryl;

$R_{20}$  is H;  $C_1-C_6$ alkyl,  $C_1-C_6$ alkyl- $C_3-C_9$ cycloalkyl, e.g., cyclopropylmethyl; aryl; heteroaryl; arylalkyl, e.g., benzyl; heteroarylalkyl, e.g., pyridylmethyl; acyl, e.g., acetyl, propionyl and benzoyl; or sulfonyl, e.g., methanesulfonyl, ethanesulfonyl, benzenesulfonyl, toluenesulfonyl);

$A_1$  is 1, 2 or 3 substituents which are independently H,  $C_1-C_6$ alkyl,  $-OR_{19}$  or halo;

$R_{19}$  is selected from H;  $C_1-C_6$ alkyl;  $C_4-C_9$ cycloalkyl;  $C_4-C_9$ heterocycloalkyl; aryl; heteroaryl; arylalkyl, e.g., benzyl; and heteroarylalkyl, e.g., pyridylmethyl;

p is 0-3; and


q is 1-5 and r is 0; or

q is 0 and r is 1-5;

or a pharmaceutically acceptable salt thereof. The other variable substituents are as defined above.

Especially useful compounds of formula (Id), are those wherein  $R_2$  is H or  $-(CH_2)_pCH_2OH$ , wherein p is 1-3, especially those wherein  $R_1$  is H; such as those wherein  $R_1$  is H and X and Y are each H, and wherein q is 1-3 and r is 0 or wherein q is 0 and r is 1-3. Among these compounds  $R_2$  is preferably H or  $-CH_2-CH_2-OH$  and the sum of q and r is preferably 1.

The present invention further relates to compounds of the formula (Ie):



or a pharmaceutically acceptable salt thereof. The variable substituents are as defined above.

Especially useful compounds of formula (Ie), are those wherein  $R_{18}$  is H, fluoro, chloro, bromo, a  $C_1-C_4$ alkyl group, a substituted  $C_1-C_4$ alkyl group, a  $C_3-C_7$ cycloalkyl group, unsubstituted phenyl, phenyl substituted in the para position, or a heteroaryl, e.g., pyridyl, ring.

Another group of useful compounds of formula (Ie), are those wherein  $R_2$  is H or  $-(CH_2)_pCH_2OH$ , wherein p is 1-3, especially those wherein  $R_1$  is H; such as those wherein  $R_1$  is H and X and Y are each H, and wherein q is 1-3 and r is 0 or wherein q is 0 and r is 1-3. Among these compounds  $R_2$  is preferably H or  $-CH_2-CH_2-OH$  and the sum of q and r is preferably 1. Among these compounds p is preferably 1 and  $R_3$  and  $R_4$  are preferably H.


Another group of useful compounds of formula (Ie), are those wherein  $R_{18}$  is H, methyl, ethyl, *t*-butyl, trifluoromethyl, cyclohexyl, phenyl, 4-methoxyphenyl, 4-trifluoromethylphenyl, 2-furanyl, 2-thiophenyl, or 2-, 3- or 4-pyridyl wherein the 2-furanyl, 2-thiophenyl and 2-, 3- or 4-pyridyl substituents are unsubstituted or substituted as described above for heteroaryl rings;  $R_2$  is H or  $-(CH_2)_pCH_2OH$ , wherein p is 1-3; especially those wherein  $R_1$  is H and X and Y are each H, and wherein q is 1-3 and r is 0 or wherein q is 0 and r is 1-3.

and r is 1-3. Among these compounds R<sub>2</sub> is preferably H or -CH<sub>2</sub>-CH<sub>2</sub>-OH and the sum of q and r is preferably 1.

Those compounds of formula (Ie), wherein R<sub>20</sub> is H or C<sub>1</sub>-C<sub>6</sub>alkyl, especially H, are important members of each of the subgenuses of compounds of formula (Ie) described above.

*N*-hydroxy-3-[4-[[2-hydroxyethyl][2-(1*H*-indol-3-yl)ethyl]-amino]methyl]phenyl]-2*E*-2-propenamide, *N*-hydroxy-3-[4-[[2-(1*H*-indol-3-yl)ethyl]-amino]methyl]phenyl]-2*E*-2-propenamide and *N*-hydroxy-3-[4-[[2-(2-methyl-1*H*-indol-3-yl)-ethyl]-amino]methyl]phenyl]-2*E*-2-propenamide or a pharmaceutically acceptable salt thereof, are important compounds of formula (Ie).

The present invention further relates to the compounds of the formula (If):



or a pharmaceutically acceptable salt thereof. The variable substituents are as defined above.


Useful compounds of formula (If), are include those wherein R<sub>2</sub> is H or -(CH<sub>2</sub>)<sub>p</sub>CH<sub>2</sub>OH, wherein p is 1-3, especially those wherein R<sub>1</sub> is H; such as those wherein R<sub>1</sub> is H and X and Y are each H, and wherein q is 1-3 and r is 0 or wherein q is 0 and r is 1-3. Among these compounds R<sub>2</sub> is preferably H or -CH<sub>2</sub>-CH<sub>2</sub>-OH and the sum of q and r is preferably 1.

*N*-hydroxy-3-[4-[[2-(benzofur-3-yl)-ethyl]-amino]methyl]phenyl]-2*E*-2-propenamide or a pharmaceutically acceptable salt thereof, is an important compound of formula (If).


The compounds described above are often used in the form of a pharmaceutically acceptable salt. Pharmaceutically acceptable salts include, when appropriate, pharmaceutically acceptable base addition salts and acid addition salts, e.g., metal salts, such as alkali and alkaline earth metal salts, ammonium salts, organic amine addition salts

and amino acid addition salts and sulfonate salts. Acid addition salts include inorganic acid addition salts, such as hydrochloride, sulfate and phosphate; and organic acid addition salts, such as alkyl sulfonate, arylsulfonate, acetate, maleate, fumarate, tartrate, citrate and lactate. Examples of metal salts are alkali metal salts, such as lithium salt, sodium salt and potassium salt; alkaline earth metal salts, such as magnesium salt and calcium salt, aluminum salt and zinc salt. Examples of ammonium salts are ammonium salt and tetramethylammonium salt. Examples of organic amine addition salts are salts with morpholine and piperidine. Examples of amino acid addition salts are salts with glycine, phenylalanine, glutamic acid and lysine. Sulfonate salts include mesylate, tosylate and benzene sulfonic acid salts.

Additional HDAC compounds within the scope of formula (I), and their synthesis, are disclosed in WO 02/22577 published March 21, 2002 which is incorporated herein by reference in its entirety. Two preferred compounds within the scope of WO 02/22577 are:



*N*-hydroxy-3-[4-[(2-hydroxyethyl){2-(1*H*-indol-3-yl)ethyl}-amino]methyl]phenyl]-2*E*-2-propenamide, or a pharmaceutically acceptable salt thereof and



*N*-hydroxy-3-[4-[[[2-(2-methyl-1*H*-indol-3-yl)-ethyl]-amino]methyl]phenyl]-2*E*-2-propenamide, or a pharmaceutically acceptable salt thereof.

Furthermore, the invention relates to a method of treating Hodgkins disease, comprising administering a therapeutically effective amount of an HDAC inhibitor to a warm-blooded animal, in particular a human, in need thereof, preferably a therapeutically effective amount of a compound of formula (I), as defined above, or the salt of such compound having

at least one salt-forming group, to a warm-blooded animal, preferably a human, in need thereof.

The term "treatment", as used herein, comprises the treatment of patients having Hodgkins disease or being in a pre-stage of said disease which effects the delay of progression of the disease in said patients.

The present invention provides a method of treating Hodgkins disease comprising administering a HDAC inhibitor in an amount which is therapeutically effective against Hodgkins disease to a warm-blooded animal in need thereof.

The person skilled in the pertinent art is fully enabled to select relevant test models to prove the hereinbefore and hereinafter mentioned beneficial effects on Hodgkins disease of a compound inhibiting the HDAC activity. The pharmacological activity of a compound inhibiting the HDAC activity may, e.g., be demonstrated in a suitable clinical study or by means of the Examples described below.

The present invention also provides the use of a compound of formula (I), as defined herein, and the use of a COMBINATION OF THE INVENTION for the preparation of a medicament for the treatment of Hodgkins disease.

Combination partners include antiproliferative compounds. Such antiproliferative compounds include, but are not limited to aromatase inhibitors; antiestrogens; topoisomerase I inhibitors; topoisomerase II inhibitors; microtubule active compounds; alkylating compounds; histone deacetylase inhibitors; compounds which induce cell differentiation processes; cyclooxygenase inhibitors; MMP inhibitors; mTOR inhibitors; antineoplastic antimetabolites; platin compounds; compounds targeting/decreasing a protein or lipid kinase activity and further anti-angiogenic compounds; compounds which target, decrease or inhibit the activity of a protein or lipid phosphatase; gonadorelin agonists; anti-androgens; methionine aminopeptidase inhibitors; bisphosphonates; biological response modifiers; antiproliferative antibodies; heparanase inhibitors; inhibitors of Ras oncogenic isoforms; telomerase inhibitors; proteasome inhibitors; compounds used in the treatment of hematologic malignancies; compounds which target, decrease or inhibit the activity of Flt-3; Hsp90 inhibitors such as 17-AAG (17-allylaminogeldanamycin, NSC330507), 17-DMAG (17-dimethylaminoethylamino-17-demethoxy-geldanamycin, NSC707545), IPI-504, CNF1010, CNF2024, CNF1010 from Conforma Therapeutics; temozolomide (TEMODAL®); kinesin spindle protein inhibitors, such as SB715992 or SB743921 from GlaxoSmithKline, or

pentamidine/chlorpromazine from CombinatoRx; MEK inhibitors such as ARRY142886 from Array PioPharma, AZD6244 from AstraZeneca, PD181461 from Pfizer and leucovorin. The term "aromatase inhibitor" as used herein relates to a compound which inhibits the estrogen production, i.e. the conversion of the substrates androstenedione and testosterone to estrone and estradiol, respectively. The term includes, but is not limited to steroids, especially atamestane, exemestane and formestane and, in particular, non-steroids, especially aminoglutethimide, roglethimide, pyridoglutethimide, trilostane, testolactone, ketokonazole, vorozole, fadrozole, anastrozole and letrozole. Exemestane can be administered, e.g., in the form as it is marketed, e.g. under the trademark AROMASIN. Formestane can be administered, e.g., in the form as it is marketed, e.g. under the trademark LENTARON. Fadrozole can be administered, e.g., in the form as it is marketed, e.g. under the trademark AFEMA. Anastrozole can be administered, e.g., in the form as it is marketed, e.g. under the trademark ARIMIDEX. Letrozole can be administered, e.g., in the form as it is marketed, e.g. under the trademark FEMARA or FEMAR. Aminoglutethimide can be administered, e.g., in the form as it is marketed, e.g. under the trademark ORIMETEN. A combination of the invention comprising a chemotherapeutic agent which is an aromatase inhibitor is particularly useful for the treatment of hormone receptor positive tumors, e.g. breast tumors.

The term "antiestrogen" as used herein relates to a compound which antagonizes the effect of estrogens at the estrogen receptor level. The term includes, but is not limited to tamoxifen, fulvestrant, raloxifene and raloxifene hydrochloride. Tamoxifen can be administered, e.g., in the form as it is marketed, e.g. under the trademark NOLVADEX. Raloxifene hydrochloride can be administered, e.g., in the form as it is marketed, e.g. under the trademark EVISTA. Fulvestrant can be formulated as disclosed in US 4,659,516 or it can be administered, e.g., in the form as it is marketed, e.g. under the trademark FASLODEX. A combination of the invention comprising a chemotherapeutic agent which is an antiestrogen is particularly useful for the treatment of estrogen receptor positive tumors, e.g. breast tumors.

The term "anti-androgen" as used herein relates to any substance which is capable of inhibiting the biological effects of androgenic hormones and includes, but is not limited to, bicalutamide (CASODEX), which can be formulated, e.g. as disclosed in US 4,636,505. The term "gonadorelin agonist" as used herein includes, but is not limited to abarelix, goserelin and goserelin acetate. Goserelin is disclosed in US 4,100,274 and can be administered, e.g., in

the form as it is marketed, e.g. under the trademark ZOLADEX. Abarelix can be formulated, e.g. as disclosed in US 5,843,901.

The term "topoisomerase I inhibitor" as used herein includes, but is not limited to topotecan, gimatecan, irinotecan, camptothecian and its analogues, 9-nitrocamptothecin and the macromolecular camptothecin conjugate PNU-166148 (compound A1 in WO99/ 17804). Irinotecan can be administered, e.g. in the form as it is marketed, e.g. under the trademark CAMPTOSAR. Topotecan can be administered, e.g., in the form as it is marketed, e.g. under the trademark HYCAMTIN.

The term "topoisomerase II inhibitor" as used herein includes, but is not limited to the anthracyclines such as doxorubicin (including liposomal formulation, e.g. CAELYX), daunorubicin, epirubicin, idarubicin and nemorubicin, the anthraquinones mitoxantrone and losoxantrone, and the podophyllotoxines etoposide and teniposide. Etoposide can be administered, e.g. in the form as it is marketed, e.g. under the trademark ETOPOPHOS. Teniposide can be administered, e.g. in the form as it is marketed, e.g. under the trademark VM 26-BRISTOL. Doxorubicin can be administered, e.g. in the form as it is marketed, e.g. under the trademark ADRIBLASTIN or ADRIAMYCIN. Epirubicin can be administered, e.g. in the form as it is marketed, e.g. under the trademark FARMORUBICIN. Idarubicin can be administered, e.g. in the form as it is marketed, e.g. under the trademark ZAVEDOS. Mitoxantrone can be administered, e.g. in the form as it is marketed, e.g. under the trademark NOVANTRON.

The term "microtubule active agent" relates to microtubule stabilizing, microtubule destabilizing compounds and microtubulin polymerization inhibitors including, but not limited to taxanes, e.g. paclitaxel and docetaxel, vinca alkaloids, e.g., vinblastine, especially vinblastine sulfate, vincristine especially vincristine sulfate, and vinorelbine, discodermolides, cochicine and epothilones and derivatives thereof, e.g. epothilone B or D or derivatives thereof. Paclitaxel may be administered e.g. in the form as it is marketed, e.g. TAXOL. Docetaxel can be administered, e.g., in the form as it is marketed, e.g. under the trademark TAXOTERE. Vinblastine sulfate can be administered, e.g., in the form as it is marketed, e.g. under the trademark VINBLASTIN R.P. Vincristine sulfate can be administered, e.g., in the form as it is marketed, e.g. under the trademark FARMISTIN. Discodermolide can be obtained, e.g., as disclosed in US 5,010,099. Also included are Epothilone derivatives which are disclosed in WO 98/10121, US 6,194,181, WO 98/25929, WO 98/08849, WO 99/43653, WO 98/22461 and WO 00/31247. Especially preferred are Epothilone A and/or B.

The term "alkylating agent" as used herein includes, but is not limited to, cyclophosphamide, ifosfamide, melphalan or nitrosourea (BCNU or Gliadel). Cyclophosphamide can be administered, e.g., in the form as it is marketed, e.g. under the trademark CYCLOSTIN. Ifosfamide can be administered, e.g., in the form as it is marketed, e.g. under the trademark HOLOXAN.

The term "platin compound" as used herein includes, but is not limited to, carboplatin, cisplatin, cisplatinum and oxaliplatin. Carboplatin can be administered, e.g., in the form as it is marketed, e.g. under the trademark CARBOPLAT. Oxaliplatin can be administered, e.g., in the form as it is marketed, e.g. under the trademark ELOXATIN.

The term "compounds targeting/decreasing a protein or lipid kinase activity; or a protein or lipid phosphatase activity; or further anti-angiogenic compounds" as used herein includes, but is not limited to, protein tyrosine kinase and/or serine and/or threonine kinase inhibitors or lipid kinase inhibitors, e.g.,

- a) compounds targeting, decreasing or inhibiting the activity of the platelet-derived growth factor-receptors (PDGFR), such as compounds which target, decrease or inhibit the activity of PDGFR, especially compounds which inhibit the PDGF receptor, e.g. a N-phenyl-2-pyrimidine-amine derivative, e.g. imatinib, SU101, SU6668 and GFB-111;
- b) compounds targeting, decreasing or inhibiting the activity of the fibroblast growth factor-receptors (FGFR);
- c) compounds targeting, decreasing or inhibiting the activity of the insulin-like growth factor receptor I (IGF-IR), such as compounds which target, decrease or inhibit the activity of IGF-IR, especially compounds which inhibit the kinase activity of IGF-I receptor, such as those compounds disclosed in WO 02/092599, or antibodies that target the extracellular domain of IGF-I receptor or its growth factors;
- d) compounds targeting, decreasing or inhibiting the activity of the Trk receptor tyrosine kinase family, or ephrin B4 inhibitors;
- e) compounds targeting, decreasing or inhibiting the activity of the Axl receptor tyrosine kinase family;
- f) compounds targeting, decreasing or inhibiting the activity of the Ret receptor tyrosine kinase;

- g) compounds targeting, decreasing or inhibiting the activity of the Kit/SCFR receptor tyrosine kinase, e.g. imatinib;
- h) compounds targeting, decreasing or inhibiting the activity of the C-kit receptor tyrosine kinases - (part of the PDGFR family), such as compounds which target, decrease or inhibit the activity of the c-Kit receptor tyrosine kinase family, especially compounds which inhibit the c-Kit receptor, e.g. imatinib;
- i) compounds targeting, decreasing or inhibiting the activity of members of the c-Abl family, their gene-fusion products (e.g. BCR-Abl kinase) and mutants, such as compounds which target decrease or inhibit the activity of c-Abl family members and their gene fusion products, e.g. a N-phenyl-2-pyrimidine-amine derivative, e.g. imatinib or nilotinib (AMN107); PD180970; AG957; NSC 680410; PD173955 from ParkeDavis; or dasatinib (BMS-354825);
- j) compounds targeting, decreasing or inhibiting the activity of members of the protein kinase C (PKC) and Raf family of serine/threonine kinases, members of the MEK, SRC, JAK, FAK, PDK1, PKB/Akt, and Ras/MAPK family members, and/or members of the cyclin-dependent kinase family (CDK) and are especially those staurosporine derivatives disclosed in US 5,093,330, e.g. midostaurin; examples of further compounds include e.g. UCN-01, safingol, BAY 43-9006, Bryostatin 1, Perifosine; Ilmofosine; RO 318220 and RO 320432; GO 6976; Isis 3521; LY333531/LY379196; isoquinoline compounds such as those disclosed in WO 00/09495; FTIs; PD184352 or QAN697 (a P13K inhibitor) or AT7519 (CDK inhibitor);
- k) compounds targeting, decreasing or inhibiting the activity of protein-tyrosine kinase inhibitors, such as compounds which target, decrease or inhibit the activity of protein-tyrosine kinase inhibitors include imatinib mesylate (GLEEVEC) or tyrphostin. A tyrphostin is preferably a low molecular weight ( $Mr < 1500$ ) compound, or a pharmaceutically acceptable salt thereof, especially a compound selected from the benzylidenemalonitrile class or the S-arylbenzenemalonitrile or bisubstrate quinoline class of compounds, more especially any compound selected from the group consisting of Tyrphostin A23/RG-50810; AG 99; Tyrphostin AG 213; Tyrphostin AG 1748; Tyrphostin AG 490; Tyrphostin B44; Tyrphostin B44 (+) enantiomer; Tyrphostin AG 555; AG 494; Tyrphostin AG 556, AG957 and adaphostin (4-{[(2,5-dihydroxyphenyl)methyl]amino}-benzoic acid adamanyl ester; NSC 680410, adaphostin);

l) compounds targeting, decreasing or inhibiting the activity of the epidermal growth factor family of receptor tyrosine kinases (EGFR, ErbB2, ErbB3, ErbB4 as homo- or heterodimers) and their mutants, such as compounds which target, decrease or inhibit the activity of the epidermal growth factor receptor family are especially compounds, proteins or antibodies which inhibit members of the EGF receptor tyrosine kinase family, e.g. EGF receptor, ErbB2, ErbB3 and ErbB4 or bind to EGF or EGF related ligands, and are in particular those compounds, proteins or monoclonal antibodies generically and specifically disclosed in WO 97/02266, e.g. the compound of ex. 39, or in EP 0 564 409, WO 99/03854, EP 0520722, EP 0 566 226, EP 0 787 722, EP 0 837 063, US 5,747,498, WO 98/10767, WO 97/30034, WO 97/49688, WO 97/38983 and, especially, WO 96/30347 (e.g. compound known as CP 358774), WO 96/33980 (e.g. compound ZD 1839) and WO 95/03283 (e.g. compound ZM105180); e.g. trastuzumab (Herceptin™), cetuximab (Erbitux™), Iressa, Tarceva, OSI-774, CI-1033, EKB-569, GW-2016, E1.1, E2.4, E2.5, E6.2, E6.4, E2.11, E6.3 or E7.6.3, and 7H-pyrrolo-[2,3-d]pyrimidine derivatives which are disclosed in WO 03/013541; and m) compounds targeting, decreasing or inhibiting the activity of the c-Met receptor, such as compounds which target, decrease or inhibit the activity of c-Met, especially compounds which inhibit the kinase activity of c-Met receptor, or antibodies that target the extracellular domain of c-Met or bind to HGF.

Further anti-angiogenic compounds include compounds having another mechanism for their activity, e.g. unrelated to protein or lipid kinase inhibition e.g. thalidomide (THALOMID) and TNP-470.

Compounds which target, decrease or inhibit the activity of a protein or lipid phosphatase are e.g. inhibitors of phosphatase 1, phosphatase 2A, or CDC25, e.g. okadaic acid or a derivative thereof.

Compounds which induce cell differentiation processes are e.g. retinoic acid,  $\alpha$ -  $\gamma$ - or  $\delta$ -tocopherol or  $\alpha$ -  $\gamma$ - or  $\delta$ -tocotrienol.

The term cyclooxygenase inhibitor as used herein includes, but is not limited to, e.g. Cox-2 inhibitors, 5-alkyl substituted 2-arylamino phenylacetic acid and derivatives, such as celecoxib (CELEBREX), rofecoxib (VIOXX), etoricoxib, valdecoxib or a 5-alkyl-2-

arylaminophenylacetic acid, e.g. 5-methyl-2-(2'-chloro-6'-fluoroanilino)phenyl acetic acid, lumiracoxib.

The term "bisphosphonates" as used herein includes, but is not limited to, etridonic, clodronic, tiludronic, pamidronic, alendronic, ibandronic, risedronic and zoledronic acid. "Etridonic acid" can be administered, e.g., in the form as it is marketed, e.g. under the trademark DIDRONE. "Clodronic acid" can be administered, e.g., in the form as it is marketed, e.g. under the trademark BONEFOS. "Tiludronic acid" can be administered, e.g., in the form as it is marketed, e.g. under the trademark SKELID. "Pamidronic acid" can be administered, e.g. in the form as it is marketed, e.g. under the trademark AREDIA<sup>TM</sup>. "Alendronic acid" can be administered, e.g., in the form as it is marketed, e.g. under the trademark FOSAMAX. "Ibandronic acid" can be administered, e.g., in the form as it is marketed, e.g. under the trademark BONDTRANAT. "Risedronic acid" can be administered, e.g., in the form as it is marketed, e.g. under the trademark ACTONEL. "Zoledronic acid" can be administered, e.g. in the form as it is marketed, e.g. under the trademark ZOMET. The term "mTOR inhibitors" relates to compounds which inhibit the mammalian target of rapamycin (mTOR) and which possess antiproliferative activity such as sirolimus (Rapamune<sup>®</sup>), everolimus (Certican<sup>TM</sup>), CCI-779 and ABT578.

The term "heparanase inhibitor" as used herein refers to compounds which target, decrease or inhibit heparin sulfate degradation. The term includes, but is not limited to, PI-88.

The term " biological response modifier" as used herein refers to a lymphokine or interferons, e.g. interferon  $\gamma$ .

The term "inhibitor of Ras oncogenic isoforms", e.g. H-Ras, K-Ras, or N-Ras, as used herein refers to compounds which target, decrease or inhibit the oncogenic activity of Ras e.g. a "farnesyl transferase inhibitor" e.g. L-744832, DK8G557 or R115777 (Zarnestra).

The term "telomerase inhibitor" as used herein refers to compounds which target, decrease or inhibit the activity of telomerase. Compounds which target, decrease or inhibit the activity of telomerase are especially compounds which inhibit the telomerase receptor, e.g. telomestatin.

The term "methionine aminopeptidase inhibitor" as used herein refers to compounds which target, decrease or inhibit the activity of methionine aminopeptidase. Compounds which target, decrease or inhibit the activity of methionine aminopeptidase are e.g. bengamide or a derivative thereof.

The term "proteasome inhibitor" as used herein refers to compounds which target, decrease or inhibit the activity of the proteasome. Compounds which target, decrease or inhibit the activity of the proteasome include e.g. Bortezomib (Velcade<sup>TM</sup>) and MLN 341. The term "matrix metalloproteinase inhibitor" or ("MMP" inhibitor) as used herein includes, but is not limited to, collagen peptidomimetic and nonpeptidomimetic inhibitors, tetracycline derivatives, e.g. hydroxamate peptidomimetic inhibitor batimastat and its orally bioavailable analogue marimastat (BB-2516), prinomastat (AG3340), metastat (NSC 683551) BMS-279251, BAY 12-9566, TAA211, MMI270B or AAJ996.

The term "compounds used in the treatment of hematologic malignancies" as used herein includes, but is not limited to, FMS-like tyrosine kinase inhibitors e.g. compounds targeting, decreasing or inhibiting the activity of FMS-like tyrosine kinase receptors (Flt-3R); interferon, 1-b-D-arabinofuransylcytosine (ara-c) and bisulfan; and ALK inhibitors e.g. compounds which target, decrease or inhibit anaplastic lymphoma kinase. Compounds which target, decrease or inhibit the activity of FMS-like tyrosine kinase receptors (Flt-3R) are especially compounds, proteins or antibodies which inhibit members of the Flt-3R receptor kinase family, e.g. PKC412, midostaurin, a staurosporine derivative, SU11248 and MLN518.

The term "HSP90 inhibitors" as used herein includes, but is not limited to, compounds targeting, decreasing or inhibiting the intrinsic ATPase activity of HSP90; degrading, targeting, decreasing or inhibiting the HSP90 client proteins via the ubiquitin proteosome pathway. Compounds targeting, decreasing or inhibiting the intrinsic ATPase activity of HSP90 are especially compounds, proteins or antibodies which inhibit the ATPase activity of HSP90 e.g., 17-allylamino,17-demethoxygeldanamycin (17AAG), a geldanamycin derivative; other geldanamycin related compounds; radicicol and HDAC inhibitors.

The term "antiproliferative antibodies" as used herein includes, but is not limited to, trastuzumab (Herceptin<sup>TM</sup>), Trastuzumab-DM1, erbitux, bevacizumab (Avastin<sup>TM</sup>), rituximab

(Rituxan®), PRO64553 (anti-CD40) and 2C4 Antibody. By antibodies is meant e.g. intact monoclonal antibodies, polyclonal antibodies, multispecific antibodies formed from at least 2 intact antibodies, and antibodies fragments so long as they exhibit the desired biological activity.

The term "antileukemic compounds" includes, for example, Ara-C, a pyrimidine analog, which is the 2'-alpha-hydroxy ribose (arabinoside) derivative of deoxycytidine. Also included is the purine analog of hypoxanthine, 6-mercaptopurine (6-MP) and fludarabine phosphate. Somatostatin receptor antagonists as used herein refers to compounds which target, treat or inhibit the somatostatin receptor such as octreotide, and SOM230.

Tumor cell damaging approaches refer to approaches such as ionizing radiation. The term "ionizing radiation" referred to above and hereinafter means ionizing radiation that occurs as either electromagnetic rays (such as X-rays and gamma rays) or particles (such as alpha and beta particles). Ionizing radiation is provided in, but not limited to, radiation therapy and is known in the art. See Hellman, Principles of Radiation Therapy, Cancer, in *Principles and Practice of Oncology*, Devita et al., Eds., 4<sup>th</sup> Edition, Vol. 1, pp. 248-275 (1993).

The term EDG binders as used herein refers a class of immunosuppressants that modulates lymphocyte recirculation, such as FTY720.

The term ribonucleotide reductase inhibitors refers to pyrimidine or purine nucleoside analogs including, but not limited to, fludarabine and/or cytosine arabinoside (ara-C), 6-thioguanine, 5-fluorouracil, cladribine, 6-mercaptopurine (especially in combination with ara-C against ALL) and/or pentostatin. Ribonucleotide reductase inhibitors are especially hydroxyurea or 2-hydroxy-1*H*-isoindole-1,3-dione derivatives, such as PL-1, PL-2, PL-3, PL-4, PL-5, PL-6, PL-7 or PL-8 mentioned in Nandy et al., *Acta Oncologica*, Vol. 33, No. 8, pp. 953-961 (1994).

The term "S-adenosylmethionine decarboxylase inhibitors" as used herein includes, but is not limited to the compounds disclosed in US 5,461,076.

Also included are in particular those compounds, proteins or monoclonal antibodies of VEGF disclosed in WO 98/35958, e.g. 1-(4-chloroanilino)-4-(4-pyridylmethyl)phthalazine or a pharmaceutically acceptable salt thereof, e.g. the succinate, or in WO 00/09495, WO 00/27820, WO 00/59509, WO 98/11223, WO 00/27819 and EP 0 769 947; those as described by Prewett et al, *Cancer Res*, Vol. 59, pp. 5209-5218 (1999); Yuan et al., *Proc Natl Acad Sci U S A*, Vol. 93, pp. 14765-14770 (1996); Zhu et al., *Cancer Res*, Vol. 58, pp. 3209-3214 (1998); and Mordenti et al., *Toxicol Pathol*, Vol. 27, No. 1, pp. 14-21 (1999); in WO 00/37502 and WO 94/10202; ANGIOSTATIN, described by O'Reilly et al., *Cell*, Vol. 79, pp. 315-328 (1994); ENDOSTATIN, described by O'Reilly et al., *Cell*, Vol. 88, pp. 277-285 (1997); anthranilic acid amides; ZD4190; ZD6474; SU5416; SU6668; bevacizumab; or anti-VEGF antibodies or anti-VEGF receptor antibodies, e.g. rhuMAb and RHUFab, VEGF aptamer e.g. Macugon; FLT-4 inhibitors, FLT-3 inhibitors, VEGFR-2 IgG1 antibody, Angiozyme (RPI 4610) and Bevacizumab (Avastin<sup>TM</sup>).

Photodynamic therapy as used herein refers to therapy which uses certain chemicals known as photosensitizing compounds to treat or prevent cancers. Examples of photodynamic therapy includes treatment with compounds, such as e.g. VISUDYNE and porfimer sodium.

Angiostatic steroids as used herein refers to compounds which block or inhibit angiogenesis, such as, e.g., anecortave, triamcinolone, hydrocortisone, 11- $\alpha$ -epihydrocortisol, cortexolone, 17 $\alpha$ -hydroxyprogesterone, corticosterone, desoxycorticosterone, testosterone, estrone and dexamethasone. Implants containing corticosteroids refers to compounds, such as e.g. fluocinolone, dexamethasone.


Other chemotherapeutic compounds include, but are not limited to, plant alkaloids, hormonal compounds and antagonists; biological response modifiers, preferably lymphokines or interferons; antisense oligonucleotides or oligonucleotide derivatives; shRNA or siRNA; or miscellaneous compounds or compounds with other or unknown mechanism of action.

The compounds of the invention are also useful as co-therapeutic compounds for use in combination with other drug substances such as anti-inflammatory, bronchodilatory or

antihistamine drug substances, particularly in the treatment of obstructive or inflammatory airways diseases such as those mentioned hereinbefore, for example as potentiators of therapeutic activity of such drugs or as a means of reducing required dosaging or potential side effects of such drugs. A compound of the invention may be mixed with the other drug substance in a fixed pharmaceutical composition or it may be administered separately, before, simultaneously with or after the other drug substance. Accordingly the invention includes a combination of a compound of the invention as hereinbefore described with an anti-inflammatory, bronchodilatory, antihistamine or anti-tussive drug substance, said compound of the invention and said drug substance being in the same or different pharmaceutical composition.

Suitable anti-inflammatory drugs include steroids, in particular glucocorticosteroids such as budesonide, beclamethasone dipropionate, fluticasone propionate, ciclesonide or mometasone furoate, or steroids described in WO 02/88167, WO 02/12266, WO 02/100879, WO 02/00679 (especially those of Examples 3, 11, 14, 17, 19, 26, 34, 37, 39, 51, 60, 67, 72, 73, 90, 99 and 101), WO 03/035668, WO 03/048181, WO 03/062259, WO 03/064445, WO 03/072592, non-steroidal glucocorticoid receptor agonists such as those described in WO 00/00531, WO 02/10143, WO 03/082280, WO 03/082787, WO 03/104195, WO 04/005229; LTB4 antagonists such LY293111, CGS025019C, CP-195543, SC-53228, BIIL 284, ONO 4057, SB 209247 and those described in US 5451700; LTD4 antagonists such as montelukast and zafirlukast; PDE4 inhibitors such cilomilast (Ariflo® GlaxoSmithKline), Roflumilast (Byk Gulden), V-11294A (Napp), BAY19-8004 (Bayer), SCH-351591 (Schering-Plough), Arofylline (Almirall Prodesfarma), PD189659 / PD168787 (Parke-Davis), AWD-12-281 (Asta Medica), CDC-801 (Celgene), SelCID(TM) CC-10004 (Celgene), VM554/UM565 (Vernalis), T-440 (Tanabe), KW-4490 (Kyowa Hakko Kogyo), and those disclosed in WO 92/19594, WO 93/19749, WO 93/19750, WO 93/19751, WO 98/18796, WO 99/16766, WO 01/13953, WO 03/104204, WO 03/104205, WO 03/39544, WO 04/000814, WO 04/000839, WO 04/005258, WO 04/018450, WO 04/018451, WO 04/018457, WO 04/018465, WO 04/018431, WO 04/018449, WO 04/018450, WO 04/018451, WO 04/018457, WO 04/018465, WO 04/018465, WO 04/019944, WO 04/019945, WO 04/045607 and WO 04/037805; A2a agonists such as those disclosed in EP 409595A2, EP 1052264, EP 1241176, WO 94/17090, WO 96/02543, WO 96/02553, WO 98/28319, WO 99/24449, WO 99/24450, WO 99/24451, WO 99/38877, WO 99/41267, WO 99/67263, WO 99/67264, WO 99/67265, WO 99/67266, WO 00/23457, WO 00/77018, WO 00/78774, WO 01/23399, WO 01/27130, WO

01/27131, WO 01/60835, WO 01/94368, WO 02/00676, WO 02/22630, WO 02/96462, WO 03/086408, WO 04/039762, WO 04/039766, WO 04/045618 and WO 04/046083; A2b antagonists such as those described in WO 02/42298; and beta-2 adrenoceptor agonists such as albuterol (salbutamol), metaproterenol, terbutaline, salmeterol fenoterol, procaterol, and especially, formoterol and pharmaceutically acceptable salts thereof, and compounds (in free or salt or solvate form) of formula I of WO 0075114, which document is incorporated herein by reference, preferably compounds of the Examples thereof, especially a compound of formula



and pharmaceutically acceptable salts thereof, as well as compounds (in free or salt or solvate form) of formula I of WO 04/16601, and also compounds of WO 04/033412.

Suitable bronchodilatory drugs include anticholinergic or antimuscarinic compounds, in particular ipratropium bromide, oxitropium bromide, tiotropium salts and CHF 4226 (Chiesi), and glycopyrrolate, but also those described in WO 01/04118, WO 02/51841, WO 02/53564, WO 03/00840, WO 03/87094, WO 04/05285, WO 02/00652, WO 03/53966, EP 424021, US 5171744, US 3714357, WO 03/33495 and WO 04/018422.

Suitable antihistamine drug substances include cetirizine hydrochloride, acetaminophen, clemastine fumarate, promethazine, loratadine, desloratadine, diphenhydramine and fexofenadine hydrochloride, activastine, astemizole, azelastine, ebastine, epinastine, mizolastine and tefenadine as well as those disclosed in WO 03/099807, WO 04/026841 and JP 2004107299.

Other useful combinations of compounds of the invention with anti-inflammatory drugs are those with antagonists of chemokine receptors, e.g. CCR-1, CCR-2, CCR-3, CCR-4, CCR-5, CCR-6, CCR-7, CCR-8, CCR-9 and CCR10, CXCR1, CXCR2, CXCR3, CXCR4, CXCR5, particularly CCR-5 antagonists such as Schering-Plough antagonists SC-351125, SCH-55700 and SCH-D, Takeda antagonists such as N-[[4-[[[6,7-dihydro-2-(4-methylphenyl)-5H-benzo-cyclohepten-8-yl]carbonyl]amino]phenyl]-methyl]tetrahydro-N,N-dimethyl-2H-pyran-4-aminium chloride (TAK-770), and CCR-5 antagonists described in US

6166037 (particularly claims 18 and 19), WO 00/66558 (particularly claim 8), WO 00/66559 (particularly claim 9), WO 04/018425 and WO 04/026873.

The structure of the active compounds identified by code nos., generic or trade names may be taken from the actual edition of the standard compendium "The Merck Index" or from databases, e.g. Patents International (e.g. IMS World Publications).

The above-mentioned compounds, which can be used in combination with a compound of the formula (I), can be prepared and administered as described in the art, such as in the documents cited above.

A compound of the formula (I) may also be used to advantage in combination with known therapeutic processes, for example, the administration of hormones or especially radiation.

A compound of formula (I) may in particular be used as a radiosensitizer, especially for the treatment of tumors which exhibit poor sensitivity to radiotherapy.

By "combination", there is meant either a fixed combination in one dosage unit form, or a kit of parts for the combined administration where a compound of the formula (I) and a combination partner may be administered independently at the same time or separately within time intervals that especially allow that the combination partners show a cooperative, e.g. synergistic effect.

## EXAMPLES

- The preclinical activity of LBH589 tested against proliferation and viability of 3 Hodgkin lymphoma cell lines, RPMI6666, HD-MY-Z, and L428.
- HL cell lines exhibited extreme sensitivity to LBH589 with IC<sub>50</sub> values ranging from 200 picomolar to 50 nanomolar
- Cell lines of mantle cell and follicular lymphoma demonstrate extreme sensitivity to LBH589
- These concentrations can easily be achieved with doses that can be safely administered in human studies

- CTCL cell lines are extremely sensitive to LBH589 . The pan-DAC inhibitor produces profound tumor regression in a CTCL mouse model.
- LBH589 produces regression of the E<sup>6</sup>-myc mouse lymphoma model and extends survival.
- B-cell lymphoma protein-2 (Bcl2) mediates resistance to LBH589 in E<sup>6</sup>-myc mouse tumors. Bcl2 may be a patient stratification marker

Single agent anti-tumor activity of LBH589 in Hodgkins Lymphoma cell lines in vitro is outlined in table I.

Table I

Cell line	Disease type	IC50	LD50	LD90
RPMI 6666	Hodgkins Lymphoma	0.2	6.7	274.7
HD-MY-Z	Hodgkins Lymphoma	50.8	239.5	>10,000
L-428	Hodgkins Lymphoma	10.5	26.1	57.5


**IC50:** Concentration of LBH589 required to inhibit growth of cell line by 50%

**LD50:** Concentration of LBH589 required to kill a cell line population by 50%

**LD90:** Concentration of LBH589 required to kill a cell line population by 90%

### What is Claimed:

1. The use of an HDAC inhibitor for the preparation of a medicament for the treatment of Hodgkins disease.
  2. Use according to Claim 1, wherein the HDAC inhibitor is a compound of formula (I):



wherein

$R_1$  is H; halo; or a straight-chain C<sub>1</sub>-C<sub>6</sub>alkyl, especially methyl, ethyl or *n*-propyl, which methyl, ethyl and *n*-propyl substituents are unsubstituted or substituted by one or more substituents described below for alkyl substituents;

$R_2$  is selected from H;  $C_1$ - $C_{10}$ alkyl, preferably  $C_1$ - $C_6$ alkyl, e.g., methyl, ethyl or  $-CH_2CH_2-$ OH;  $C_4$ - $C_9$ cycloalkyl;  $C_4$ - $C_9$ heterocycloalkyl;  $C_4$ - $C_9$ heterocycloalkylalkyl; cycloalkylalkyl, e.g., cyclopropylmethyl; aryl; heteroaryl; arylalkyl, e.g., benzyl; heteroarylalkyl, e.g., pyridylmethyl;  $-(CH_2)_nC(O)R_6$ ;  $-(CH_2)_nOC(O)R_6$ ; amino acyl;  $HON-C(O)-CH=C(R_1)-aryl-alkyl-$ ; and  $-(CH_2)_nR_7$ ;

$R_3$  and  $R_4$  are the same or different and, independently, H;  $C_1$ - $C_6$ alkyl; acyl; or acylamino; or

$R_3$  and  $R_4$ , together with the carbon to which they are bound, represent  $C=O$ ,  $C=S$  or  $C=NR_5$ ; or

$R_2$ , together with the nitrogen to which it is bound, and  $R_3$ , together with the carbon to which it is bound, can form a  $C_4$ - $C_9$ heterocycloalkyl; a heteroaryl; a polyheteroaryl; a non-aromatic polyheterocycle; or a mixed aryl and non-aryl polyheterocycle ring;

$R_5$  is selected from H; C<sub>1</sub>-C<sub>6</sub>alkyl; C<sub>4</sub>-C<sub>9</sub>cycloalkyl; C<sub>4</sub>-C<sub>9</sub>heterocycloalkyl; acyl; aryl; heteroaryl; arylalkyl, e.g., benzyl; heteroarylalkyl, e.g., pyridylmethyl; aromatic polycycles; non-aromatic polycycles; mixed aryl and non-aryl polycycles; polyheteroaryl; non-aromatic polyheterocycles; and mixed aryl and non-aryl polyheterocycles;

$n$ ,  $n_1$ ,  $n_2$  and  $n_3$  are the same or different and independently selected from 0-6, when  $n_1$  is 1-6, each carbon atom can be optionally and independently substituted with  $R_3$  and/or  $R_4$ ;

$X$  and  $Y$  are the same or different and independently selected from H; halo;  $C_1$ - $C_4$ alkyl, such as  $CH_3$  and  $CF_3$ ;  $NO_2$ ;  $C(O)R_1$ ;  $OR_9$ ;  $SR_9$ ;  $CN$ ; and  $NR_{10}R_{11}$ ;

$R_6$  is selected from H;  $C_1$ - $C_6$ alkyl;  $C_4$ - $C_9$ cycloalkyl;  $C_4$ - $C_9$ heterocycloalkyl; cycloalkylalkyl, e.g., cyclopropylmethyl; aryl; heteroaryl; arylalkyl, e.g., benzyl and 2-phenylethenyl; heteroarylalkyl, e.g., pyridylmethyl;  $OR_{12}$ ; and  $NR_{13}R_{14}$ ;

$R_7$  is selected from  $OR_{15}$ ;  $SR_{15}$ ;  $S(O)R_{16}$ ;  $SO_2R_{17}$ ;  $NR_{13}R_{14}$ ; and  $NR_{12}SO_2R_6$ ;

$R_8$  is selected from H;  $OR_{15}$ ;  $NR_{13}R_{14}$ ;  $C_1$ - $C_6$ alkyl;  $C_4$ - $C_9$ cycloalkyl;  $C_4$ - $C_9$ heterocycloalkyl; aryl; heteroaryl; arylalkyl, e.g., benzyl; and heteroarylalkyl, e.g., pyridylmethyl;

$R_9$  is selected from  $C_1$ - $C_4$ alkyl, e.g.,  $CH_3$  and  $CF_3$ ;  $C(O)$ -alkyl, e.g.,  $C(O)CH_3$ ; and  $C(O)CF_3$ ;

$R_{10}$  and  $R_{11}$  are the same or different and independently selected from H;  $C_1$ - $C_4$ alkyl; and  $-C(O)$ -alkyl;

$R_{12}$  is selected from H;  $C_1$ - $C_6$ alkyl;  $C_4$ - $C_9$ cycloalkyl;  $C_4$ - $C_9$ heterocycloalkyl;  $C_4$ - $C_9$ heterocycloalkylalkyl; aryl; mixed aryl and non-aryl polycycle; heteroaryl; arylalkyl, e.g., benzyl; and heteroarylalkyl, e.g., pyridylmethyl;

$R_{13}$  and  $R_{14}$  are the same or different and independently selected from H;  $C_1$ - $C_6$ alkyl;  $C_4$ - $C_9$ cycloalkyl;  $C_4$ - $C_9$ heterocycloalkyl; aryl; heteroaryl; arylalkyl, e.g., benzyl; heteroarylalkyl, e.g., pyridylmethyl; amino acyl; or

$R_{13}$  and  $R_{14}$ , together with the nitrogen to which they are bound, are  $C_4$ - $C_9$ heterocycloalkyl; heteroaryl; polyheteroaryl; non-aromatic polyheterocycle; or mixed aryl and non-aryl polyheterocycle;


$R_{15}$  is selected from H;  $C_1$ - $C_6$ alkyl;  $C_4$ - $C_9$ cycloalkyl;  $C_4$ - $C_9$ heterocycloalkyl; aryl; heteroaryl; arylalkyl; heteroarylalkyl; and  $(CH_2)_mZR_{12}$ ;

$R_{16}$  is selected from  $C_1$ - $C_6$ alkyl;  $C_4$ - $C_9$ cycloalkyl;  $C_4$ - $C_9$ heterocycloalkyl; aryl; heteroaryl; polyheteroaryl; arylalkyl; heteroarylalkyl; and  $(CH_2)_mZR_{12}$ ;

$R_{17}$  is selected from  $C_1$ - $C_6$ alkyl;  $C_4$ - $C_9$ cycloalkyl;  $C_4$ - $C_9$ heterocycloalkyl; aryl; aromatic polycycles; heteroaryl; arylalkyl; heteroarylalkyl; polyheteroaryl and  $NR_{13}R_{14}$ ;

m is an integer selected from 0-6; and  
 Z is selected from O; NR<sub>13</sub>; S; and S(O),  
 or a pharmaceutically acceptable salt thereof.

3. Use according to Claim 2, wherein the compound of formula (I) is *N*-hydroxy-3-[4-[[2-(2-methyl-1*H*-indol-3-yl)-ethyl]-amino]methyl]phenyl]-2*E*-2-propenamide having the formula (III):



or a pharmaceutically acceptable salt thereof.


4. Use according to any one of Claims 1 to 3, wherein the Hodgkins disease is a lymphoproliferative disease characterized by characterized clinically by the orderly spread of disease from one lymph node group to another and by the development of systemic symptoms with advanced disease.

5. Use according to any one of Claims 1 to 3, wherein the warm-blooded animal is a human.

6. Use according to any one of Claims 1 to 5, wherein the disease is Hodgkins disease

7. A method of treating Hodgkins disease comprising administering a therapeutically effective amount of an HDAC inhibitor to a warm-blooded animal in need thereof.

8. A method according to Claim 7, comprising administering a therapeutically effective amount of a compound of formula (I):



wherein

R<sub>1</sub> is H; halo; or a straight-chain C<sub>1</sub>-C<sub>6</sub>alkyl, especially methyl, ethyl or n-propyl, which methyl, ethyl and n-propyl substituents are unsubstituted or substituted by one or more substituents described below for alkyl substituents;

R<sub>2</sub> is selected from H; C<sub>1</sub>-C<sub>10</sub>alkyl, preferably C<sub>1</sub>-C<sub>6</sub>alkyl, e.g., methyl, ethyl or -CH<sub>2</sub>CH<sub>2</sub>-OH; C<sub>4</sub>-C<sub>9</sub>cycloalkyl; C<sub>4</sub>-C<sub>9</sub>heterocycloalkyl; C<sub>4</sub>-C<sub>9</sub>heterocycloalkylalkyl; cycloalkylalkyl, e.g., cyclopropylmethyl; aryl; heteroaryl; arylalkyl, e.g., benzyl; heteroarylalkyl, e.g., pyridylmethyl; -(CH<sub>2</sub>)<sub>n</sub>C(O)R<sub>6</sub>; -(CH<sub>2</sub>)<sub>n</sub>OC(O)R<sub>6</sub>; amino acyl; HON-C(O)-CH=C(R<sub>1</sub>)-aryl-alkyl-; and -(CH<sub>2</sub>)<sub>n</sub>R<sub>7</sub>;

R<sub>3</sub> and R<sub>4</sub> are the same or different and, independently, H; C<sub>1</sub>-C<sub>6</sub>alkyl; acyl; or acylamino; or

R<sub>3</sub> and R<sub>4</sub>, together with the carbon to which they are bound, represent C=O, C=S or C=NR<sub>8</sub>; or

R<sub>2</sub>, together with the nitrogen to which it is bound, and R<sub>3</sub>, together with the carbon to which it is bound, can form a C<sub>4</sub>-C<sub>9</sub>heterocycloalkyl; a heteroaryl; a polyheteroaryl; a non-aromatic polyheterocycle; or a mixed aryl and non-aryl polyheterocycle ring;

R<sub>5</sub> is selected from H; C<sub>1</sub>-C<sub>6</sub>alkyl; C<sub>4</sub>-C<sub>9</sub>cycloalkyl; C<sub>4</sub>-C<sub>9</sub>heterocycloalkyl; acyl; aryl; heteroaryl; arylalkyl, e.g., benzyl; heteroarylalkyl, e.g., pyridylmethyl; aromatic polycycles; non-aromatic polycycles; mixed aryl and non-aryl polycycles; polyheteroaryl; non-aromatic polyheterocycles; and mixed aryl and non-aryl polyheterocycles;

n, n<sub>1</sub>, n<sub>2</sub> and n<sub>3</sub> are the same or different and independently selected from 0-6, when n<sub>1</sub> is 1-6, each carbon atom can be optionally and independently substituted with R<sub>3</sub> and/or R<sub>4</sub>;

X and Y are the same or different and independently selected from H; halo; C<sub>1</sub>-C<sub>4</sub>alkyl, such as CH<sub>3</sub> and CF<sub>3</sub>; NO<sub>2</sub>; C(O)R<sub>1</sub>; OR<sub>9</sub>; SR<sub>9</sub>; CN; and NR<sub>10</sub>R<sub>11</sub>;

R<sub>6</sub> is selected from H; C<sub>1</sub>-C<sub>6</sub>alkyl; C<sub>4</sub>-C<sub>9</sub>cycloalkyl; C<sub>4</sub>-C<sub>9</sub>heterocycloalkyl; cycloalkylalkyl, e.g., cyclopropylmethyl; aryl; heteroaryl; arylalkyl, e.g., benzyl and 2-phenylethenyl; heteroarylalkyl, e.g., pyridylmethyl; OR<sub>12</sub>; and NR<sub>13</sub>R<sub>14</sub>;

R<sub>7</sub> is selected from OR<sub>15</sub>; SR<sub>15</sub>; S(O)R<sub>16</sub>; SO<sub>2</sub>R<sub>17</sub>; NR<sub>13</sub>R<sub>14</sub>; and NR<sub>12</sub>SO<sub>2</sub>R<sub>6</sub>;

R<sub>8</sub> is selected from H; OR<sub>15</sub>; NR<sub>13</sub>R<sub>14</sub>; C<sub>1</sub>-C<sub>6</sub>alkyl; C<sub>4</sub>-C<sub>9</sub>cycloalkyl; C<sub>4</sub>-C<sub>9</sub>heterocycloalkyl; aryl; heteroaryl; arylalkyl, e.g., benzyl; and heteroarylalkyl, e.g., pyridylmethyl;

R<sub>9</sub> is selected from C<sub>1</sub>-C<sub>4</sub>alkyl, e.g., CH<sub>3</sub> and CF<sub>3</sub>; C(O)-alkyl, e.g., C(O)CH<sub>3</sub>; and C(O)CF<sub>3</sub>;

R<sub>10</sub> and R<sub>11</sub> are the same or different and independently selected from H; C<sub>1</sub>-C<sub>4</sub>alkyl; and -C(O)-alkyl;

R<sub>12</sub> is selected from H; C<sub>1</sub>-C<sub>6</sub>alkyl; C<sub>4</sub>-C<sub>9</sub>cycloalkyl; C<sub>4</sub>-C<sub>9</sub>heterocycloalkyl; C<sub>4</sub>-C<sub>9</sub>heterocycloalkylalkyl; aryl; mixed aryl and non-aryl polycycle; heteroaryl; arylalkyl, e.g., benzyl; and heteroarylalkyl, e.g., pyridylmethyl;

R<sub>13</sub> and R<sub>14</sub> are the same or different and independently selected from H; C<sub>1</sub>-C<sub>6</sub>alkyl; C<sub>4</sub>-C<sub>9</sub>cycloalkyl; C<sub>4</sub>-C<sub>9</sub>heterocycloalkyl; aryl; heteroaryl; arylalkyl, e.g., benzyl; heteroarylalkyl, e.g., pyridylmethyl; amino acyl; or

R<sub>13</sub> and R<sub>14</sub>, together with the nitrogen to which they are bound, are C<sub>4</sub>-C<sub>9</sub>heterocycloalkyl; heteroaryl; polyheteroaryl; non-aromatic polyheterocycle; or mixed aryl and non-aryl polyheterocycle;

R<sub>15</sub> is selected from H; C<sub>1</sub>-C<sub>6</sub>alkyl; C<sub>4</sub>-C<sub>9</sub>cycloalkyl; C<sub>4</sub>-C<sub>9</sub>heterocycloalkyl; aryl; heteroaryl; arylalkyl; heteroarylalkyl; and (CH<sub>2</sub>)<sub>m</sub>ZR<sub>12</sub>;

R<sub>16</sub> is selected from C<sub>1</sub>-C<sub>6</sub>alkyl; C<sub>4</sub>-C<sub>9</sub>cycloalkyl; C<sub>4</sub>-C<sub>9</sub>heterocycloalkyl; aryl; heteroaryl; polyheteroaryl; arylalkyl; heteroarylalkyl; and (CH<sub>2</sub>)<sub>m</sub>ZR<sub>12</sub>;

R<sub>17</sub> is selected from C<sub>1</sub>-C<sub>6</sub>alkyl; C<sub>4</sub>-C<sub>9</sub>cycloalkyl; C<sub>4</sub>-C<sub>9</sub>heterocycloalkyl; aryl; aromatic polycycles; heteroaryl; arylalkyl; heteroarylalkyl; polyheteroaryl and NR<sub>13</sub>R<sub>14</sub>;

m is an integer selected from 0-6; and

Z is selected from O; NR<sub>13</sub>; S; and S(O),

or a pharmaceutically acceptable salt thereof to a warm-blooded animal in need thereof.

## INTERNATIONAL SEARCH REPORT

International application No  
PCT/US2009/050768

A. CLASSIFICATION OF SUBJECT MATTER  
INV. A61K31/00 A61K31/404 A61P35/02

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, BIOSIS, EMBASE, CHEM ABS Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	REVILL P ET AL: "Panobinostat: Histone deacetylase (HDAC) inhibitor, apoptosis inducer, oncolytic" DRUGS OF THE FUTURE, PROUS SCIENCE, ES, vol. 32, no. 4, 1 January 2007 (2007-01-01), pages 315-322, XP002457526 ISSN: 0377-8282 abstract	1-8
Y	WO 2008/070011 A (NOVARTIS AG [CH]; ATADJA PETER WISDOM [US]) 12 June 2008 (2008-06-12) last two lines of p. 15; p. 16, compound III	1-8

Further documents are listed in the continuation of Box C.

See patent family annex.

## \* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date.
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*&\* document member of the same patent family

Date of the actual completion of the international search

Date of mailing of the international search report

7 October 2009

28/10/2009

Name and mailing address of the ISA/  
European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040,  
Fax: (+31-70) 340-3016

Authorized officer

Dahse, Thomas

## INTERNATIONAL SEARCH REPORT

International application No PCT/US2009/050768
---

## C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	BHALLA K: "ACTIVITY OF THE HISTONE DEACETYLASE INHIBITORS LBH589 AND LAQ824 IN HEMATOLOGIC MALIGNANCIES" HAEMATOLOGICA REPORTS, FONDAZIONE FERRATA STORTI (FERRATA STORTI FOUNDATION), PAVIA, IT, vol. 1, no. 8, 1 January 2005 (2005-01-01), pages 84-88, XP008075104 ISSN: 1824-9337 p. 85, col. 1, second	1-8
X	BHALLA; BALASUBRAMANIAN; DAVID; SIRISAWAD; BUGGY; MAURO; PRACHAND; MILLER; GORDON; EVENS: "The Broad Spectrum HDAC Inhibitor PCI-24781 Induces Caspase- and ROS-Dependent Apoptosis and is Synergistic with Bortezomib in Lymphoma" NATURE PRECEDINGS, [Online] 12 July 2008 (2008-07-12), XP002549131 Retrieved from the Internet: URL: <a href="http://precedings.nature.com/documents/1999/version/1/html">http://precedings.nature.com/documents/1999/version/1/html</a> > [retrieved on 2009-10-07]	1,4-7
Y	p. 12, first sentence of discussion; p. 14, second to p. 15 last	1-8
X	S. VIVIANI, V. BONFANTE, C. FASOLA, P. VALAGUSSA AND A. M. GIANNI: "Phase II study of the histone-deacetylase inhibitor ITF2357 in relapsed/refractory Hodgkin's lymphoma patients" JOURNAL OF CLINICAL ONCOLOGY, [Online] vol. 26, no. 15S, 20 May 2008 (2008-05-20), XP002549132 Retrieved from the Internet: URL: <a href="http://meeting.ascopubs.org/cgi/content/abstract/26/15_suppl/8532">http://meeting.ascopubs.org/cgi/content/abstract/26/15_suppl/8532</a> > [retrieved on 2009-10-07]	1,4-7
Y	abstract, in particular last two sentences	1-8
X	DATABASE BIOSIS [Online] BIOSCIENCES INFORMATION SERVICE, PHILADELPHIA, PA, US; November 2007 (2007-11), BUGLIO DANIELA ET AL: "Vorinostat (SAHA) inhibits STAT6 phosphorylation and transcription, downregulates Bcl-xL, and induces apoptosis in Hodgkin lymphoma" XP002549136 Database accession no. PREV200800215653 abstract	1,4-7
Y		1-8
	-/-	

## INTERNATIONAL SEARCH REPORT

International application No PCT/US2009/050768
---

## C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	O'CONNOR O A: "Clinical experience with the novel histone deacetylase inhibitor vorinostat (suberoylanilide hydroxamic acid) in patients with relapsed lymphoma" BRITISH JOURNAL OF CANCER 200612 GB, vol. 95, no. SUPPL. 1, December 2006 (2006-12), pages S7-S12, XP002549133 ISSN: 0007-0920 1532-1827 p. S9, last of col. 2; p. S10, col. 2, last	1, 4-7
Y		1-8
X	R. G. BOCIEK, J. KURUVILLA, B. PRO, A. WEDGWOOD, Z. LI, M. DROUIN, T. PATTERSON, R. WARD, R. E. MARTELL AND A. YOUNES: "Isotype-selective histone deacetylase (HDAC) inhibitor MGCD0103 demonstrates clinical activity and safety in patients with relapsed/refractory classical Hodgkin Lymphoma (HL)" JOURNAL OF CLINICAL ONCOLOGY, [Online] vol. 26, no. 15S, 20 May 2008 (2008-05-20), XP002549134 Retrieved from the Internet: URL: <a href="http://meeting.ascopubs.org/cgi/content/abstract/26/15_suppl/8507">http://meeting.ascopubs.org/cgi/content/abstract/26/15_suppl/8507</a> [retrieved on 2009-10-07]	1, 4-7
Y	abstract	1-8
X	BHALLA; BALASUBRAMANIAN; DAVID; SIRISAWAD; BUGGY; MAURO; PRACHAND; MILLER; GORDON; EVENS: "The Broad Spectrum HDAC Inhibitor PCI-24781 Induces Caspase- and ROS-Dependent Apoptosis and is Synergistic with Bortezomib in Lymphoma" NATURE PRECEDINGS, [Online] 12 July 2008 (2008-07-12), XP002549135 Retrieved from the Internet: URL: <a href="http://precedings.nature.com/documents/1999/version/1">http://precedings.nature.com/documents/1999/version/1</a> [retrieved on 2009-10-07]	1, 4-7
Y, L	abstract and date	1-8
P, X	DATABASE BIOSIS [Online] BIOSCIENCES INFORMATION SERVICE, PHILADELPHIA, PA, US; November 2008 (2008-11), OTTMANN OLIVER G ET AL: "Phase IA/II Study of Oral Panobinostat (LBH589), a Novel Pan-Deacetylase Inhibitor (Daci) Demonstrating Efficacy in Patients with Advanced Hematologic Malignancies." XP002549137 Database accession no. PREV200900257660 abstract	1-8

**INTERNATIONAL SEARCH REPORT**

Information on patent family members

International application No

PCT/US2009/050768

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2008070011	A 12-06-2008	AR 064072 A1	11-03-2009
		AU 2007328281 A1	12-06-2008
		CA 2670741 A1	12-06-2008
		CL 34672007 A1	11-07-2008
		EC SP099483 A	28-08-2009
		EP 2099451 A2	16-09-2009