




(86) Date de dépôt PCT/PCT Filing Date: 1993/11/04
(87) Date publication PCT/PCT Publication Date: 1994/05/11
(45) Date de délivrance/Issue Date: 2007/04/17
(85) Entrée phase nationale/National Entry: 1995/05/03
(86) N° demande PCT/PCT Application No.: US 1993/010384
(87) N° publication PCT/PCT Publication No.: 1994/010332
(30) Priorité/Priority: 1992/11/04 (GB9223377.4)

(51) Cl.Int./Int.Cl. *C07K 16/46* (2006.01),
C07K 16/28 (2006.01)
(72) Inventeurs/Inventors:
CARR, FRANK J., GB;
HARRIS, WILLIAM J., GB;
TEMPEST, PHILIP R., GB
(73) Propriétaire/Owner:
MEDAREX, INC., US
(74) Agent: BORDEN LADNER GERVAIS LLP

(54) Titre : ANTICORPS HUMANISES CONTRE LES RECEPTEURS Fc DE L'IMMUNOGLOBULINE G DES
PHAGOCYTES MONONUCLEAIRES HUMAINS

(54) Title: HUMANIZED ANTIBODIES TO Fc RECEPTORS FOR IMMUNOGLOBULIN G ON HUMAN MONONUCLEAR
PHAGOCYTES



(57) Abrégé/Abstract:


Humanized antibodies are described which are specific to an Fc receptor (FcR). The humanized antibodies have at least a portion of a complementarity determining region (CDR) derived from a non-human antibody, e.g. murine, with the remaining portions being human in origin. The use of humanized antibodies rather than murine antibodies in human therapy should alleviate some of the problems associated with the use of some murine monoclonal antibodies because only the substituted CDRs will be foreign to a human's immune system. The humanized antibodies can be used in the same manner as their murine counterparts.



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C12P 21/08, C07K 15/28		A1	(11) International Publication Number: WO 94/10332
			(43) International Publication Date: 11 May 1994 (11.05.94)
(21) International Application Number: PCT/US93/10384		(74) Agents: MANDRAGOURAS, Amy, E. et al.; Lahive & Cockfield, 60 State Street, Boston, MA 02109 (US).	
(22) International Filing Date: 4 November 1993 (04.11.93)		(81) Designated States: AU, CA, JP, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).	
(30) Priority data: 9223377.4 4 November 1992 (04.11.92) GB		2148578	
(71) Applicant (for all designated States except US): MEDAREX, INC. [US/US]; 22 Chambers Street, Princeton, NJ 08542 (US).		Published With international search report.	
(72) Inventors; and (75) Inventors/Applicants (for US only) : TEMPEST, Philip, R. [GB/GB]; 63 Brighton Place, Aberdeen AB1 6RT (GB). HARRIS, William, J. [GB/GB]; 3 Caesar Avenue, Carnoustie DD7 6DR (GB). CARR, Frank, J. [GB/GB]; Birchlea, The Holdings, Balmedie, Aberdeenshire AB23 8XU (GB).			

(54) Title: HUMANIZED ANTIBODIES TO Fc RECEPTORS FOR IMMUNOGLOBULIN G ON HUMAN MONONUCLEAR PHAGOCYTES



(57) Abstract

Humanized antibodies are described which are specific to an Fc receptor (FcR). The humanized antibodies have at least a portion of a complementarity determining region (CDR) derived from a non-human antibody, e.g. murine, with the remaining portions being human in origin. The use of humanized antibodies rather than murine antibodies in human therapy should alleviate some of the problems associated with the use of some murine monoclonal antibodies because only the substituted CDRs will be foreign to a human's immune system. The humanized antibodies can be used in the same manner as their murine counterparts.

HUMANIZED ANTIBODIES TO Fc RECEPTORS FOR IMMUNOGLOBULIN G ON HUMAN MONONUCLEAR PHAGOCYTES

5 **Background**

Human Fcγ receptors (FcγR) (reviewed in Fanger, M.W., et al. (1989) Immunology Today 10:92-99), of which there are three structurally and functionally distinct types (i.e., FcγRI, FcγRII and FcγRIII), are well-characterized cell surface glycoproteins that mediate phagocytosis or antibody-dependent cell cytotoxicity (ADCC) of immunoglobulin G
10 (IgG) opsonized targets. Antibodies have been made which are directed towards FcγR for various purposes, e.g., targeting of immunotoxins to a particular target cell type, or radioimaging a particular target cell type. The antibodies typically have been murine antibodies.

Murine monoclonal antibodies are sometimes desirable for human therapeutic
15 applications because the antibodies can be purified in large quantities and are free of contamination by human pathogens such as the hepatitis or human immunodeficiency virus. Murine monoclonal antibodies have been used in some human therapies, however, results have not always been desirable due to the development of an immune response to the "foreign" murine proteins. The immune response has been termed a human anti-mouse
20 antibody or HAMA response (Schroff, R., et al. (1985), Cancer Res., 45, 879-885) and is a condition which causes serum sickness in humans and results in rapid clearance of the murine antibodies from an individual's circulation. The immune response in humans has been shown to be against both the variable and the constant regions of the murine immunoglobulin.

Recombinant DNA technology has provided the ability to alter antibodies by
25 substituting specific immunoglobulin regions from one species with immunoglobulin regions from another species. Neuberger et al. (Patent Cooperation Treaty Patent Application No. PCT/GB85/00392) describes a process whereby the complementary heavy and light chain variable domains of an Ig molecule from one species may be combined with the complementary heavy and light chain Ig constant domains from another species. This
30 process may be used to substitute the murine constant region domains to create a "chimeric" antibody which may be used for human therapy. A chimeric antibody produced as described by Neuberger et al. would have the advantage of having the human Fc region for efficient stimulation of antibody mediated effector functions, such as complement fixation, but would still have the potential to elicit an immune response in humans against the murine ("foreign")
35 variable regions.

Winter (British Patent Application Number GB2188638A) describes a process for altering antibodies by substituting the complementarity determining regions (CDRs) with those from another species. This process may be used to substitute the CDRs from the murine variable region domains of a monoclonal antibody with desirable binding properties

-2-

(for instance to a human pathogen) into human heavy and light chain Ig variable region domains. These altered Ig variable regions may then be combined with human Ig constant regions to create antibodies which are totally human in composition except for the substituted murine CDRs. The "reshaped" or "humanized" antibodies described by Winter elicit a considerably reduced immune response in humans compared to chimeric antibodies because of the considerably less murine components. Further, the half life of the altered antibodies in circulation should approach that of natural human antibodies. However, as stated by Winter, merely replacing the CDRs with complementary CDRs from another antibody which is specific for an antigen such as a viral or bacterial protein, does not always result in an altered antibody which retains the desired binding capacity. In practice, some amino acids in the framework of the antibody variable region interact with the amino acid residues that make up the CDRs so that amino acid substitutions into the human Ig variable regions are likely to be required to restore antigen binding.

15 **Summary of the Invention**

The present invention pertains to humanized antibodies specific to an Fc receptor (FcR). The humanized antibodies have at least a portion of a complementarity determining region (CDR) derived from a non-human antibody, e.g., murine, with the remaining portions being human in origin. The use of humanized antibodies rather than murine antibodies in human therapy should alleviate some of the problems associated with the use of some murine monoclonal antibodies because only the substituted CDRs will be foreign to a human host's immune system.

The present invention further pertains to the use of humanized antibodies specific to an FcR as components in heteroantibodies, bifunctional antibodies, or immunotoxins. The humanized antibody specific to an FcR may be used in the same manner and for the same purpose as its corresponding murine counterpart. For example, the humanized anti-Fc receptor antibody of this invention can be used to treat cancer, allergies, and infectious and autoimmune diseases. Diagnostic applications of the antibodies include their use in assays for FcRI levels and assays for substances that influence FcR levels.

30 **Brief Description of the Drawings**

Figure 1 compares the amino acid sequences of murine 022 VH (SEQ ID NO:3) with the amino acid sequences of humanized NEWM-based VH (022 NMVH) (SEQ ID NO:1) and humanized KOL-based VH (022 KLVH) (SEQ ID NO:2).

35 Murine residues retained in the human portion are indicated by the inverted black triangles.

-3-

Figure 2 compares the amino acid sequences of murine 022 VK (SEQ ID NO:28) with humanized REI-based VK (022 HuVK) (SEQ ID NO:4).

Murine residues retained in the human portion are indicated by the inverted black triangles.

Figure 3 depicts the vector used for expression of the humanized or chimeric
5 022 heavy chain gene.

Figure 4 depicts the vector used for expression of the humanized or chimeric
022 kappa chain gene.

Figure 5 depicts the binding of the test antibodies in the enzyme linked
immunoassay described in the example.

10

Detailed Description

The present invention pertains to a humanized antibody specific for an Fc
receptor. The humanized antibody is made up of a human antibody having at least a portion
of a complementarity determining region (CDR) derived from a non-human antibody. The
15 portion is selected to provide specificity of the humanized antibody for a human Fc receptor.
The humanized antibody has CDR's derived from a non-human antibody and the remaining
portions of the antibody molecule are human.

The antibody may be a complete antibody molecule having full length heavy
and light chains or any fragment thereof, e.g., Fab or (Fab')₂ fragment. The antibody further
20 may be a light chain or heavy chain dimer, or any minimal fragment thereof such as a Fv or a
single chain construct as described in Ladner et al. (U.S. Patent No. 4,946,778, issued August
7, 1990).

The human antibody of the present invention may be any human antibody
capable of retaining non-human CDRs. The preferred human antibody is derived from
25 known proteins NEWM and KOL for heavy chain variable regions (VHs) and REI for Ig
kappa chain, variable regions (VKs). These proteins are described in detail in the examples
below.

"Complementarity determining region" (CDR) is an art recognized
terminology and the technique used for locating the CDRs within the described sequences
30 also is conventional.

The portion of the non-human CDR inserted into the human antibody is
selected to be sufficient for allowing binding of the humanized antibody to the Fc receptor.
A sufficient portion may be selected by inserting a portion of the CDR into the human
antibody and testing the binding capacity of the created humanized antibody using the
35 enzyme linked immunosorbent assay (ELISA) described in the examples below.

All of the CDRs of a particular human antibody may be replaced with at least
a portion of a non-human CDR or only some of the CDRs may be replaced with non-human
CDRs. It is only necessary to replace the number of CDRs required for binding of the

-4-

humanized antibody to the Fc receptor. The exemplified non-human CDR is derived from a murine antibody, particularly the CDR is derived from a monoclonal antibody (mab), mab 22. The mab 22 antibody is specific to the Fc receptor and further is described in WO91/05871 and in Fanger et al. (U.S. Patent No. 4,954,617, issued September 4, 1988).

5

The CDRs are derived from a non-human antibody specific for a human Fc receptor. The CDRs can be derived from known Fc receptor antibodies such as those discussed in the Fanger et al. patent application and issued patent cited above (hereinafter Fanger et al.). The CDR may be derived from an antibody which binds to the Fc receptor at a site which is not blocked by human immunoglobulin G. The antibody also may be specific for the high affinity Fc receptor for human immunoglobulin G. Examples of antibodies from which the non-human CDRs may be derived are mab 32, mab 22, mab 44, mab 62, mab 197 and anti-FcRI antibody 62. A hybridoma cell line producing mAb 22 was deposited with the American Type Culture Collection (ATCC), 12301 Parklawn Drive, Rockville, Md. 20852, on July 9, 1996 and has been assigned ATCC Accession No. HB-12147. A hybridoma cell line producing monoclonal antibody 32.2 was deposited with the ATCC, 12301 Parklawn Drive, Rockville, Md. 20852, on July 1, 1987 and has been assigned ATCC Accession No. HB 9469. The humanized mab 22 antibody producing cell line has been deposited at the American Type Culture Collection (ATCC), 12301 Parklawn Drive, Rockville, Md. 20852 on November 4, 1992 under the designation HA022CL1 and has the Accession No. CRL 11177.

The present invention also pertains to bifunctional antibodies or heteroantibodies having at least one humanized antigen binding region derived from a humanized anti-Fc receptor antibody and at least one antigen binding region specific for a target epitope. The humanized antigen binding region may be derived from a humanized anti-Fc receptor antibody as described above. Bifunctional and heteroantibodies having an antibody portion specific for an Fc receptor are described in detail by Fanger et al.

It should be understood that the humanized antibodies of the present invention may be used in the same manner, e.g., as components of immunotoxins or heteroantibodies, as their corresponding non-humanized counterparts described by Fanger et al. The humanized antibodies further share the same utilities as their non-humanized counterparts. All aspects of the teachings of the Fanger et al. application and patent are incorporated by reference.

The humanized antibody of the present invention may be made by any method capable of replacing at least a portion of a CDR of a human antibody with a CDR derived from a non-human antibody. Winter describes a method which may be used to prepare the humanized antibodies of the present invention (UK Patent Application GB 2188638A, filed

30

4a

on March 26, 1987). The
human CDRs may be replaced with non-human CDRs using oligonucleotide site-directed
35 mutagenesis as described in the examples below.

The humanized antibody of the present invention may be made as described in the brief explanation below. A detailed method for production is set forth in the examples. It should be understood that one of ordinary skill in the art may be able to substitute known

conventional techniques for those described below for the purpose of achieving the same result. The humanized antibodies of the present invention may be produced by the following process:

- 5 (a) constructing, by conventional techniques, an expression vector containing an operon with a DNA sequence encoding an antibody heavy chain in which the CDRs and such minimal portions of the variable domain framework region that are required to retain antibody binding specificity are derived from a non-human immunoglobulin, and the remaining parts of the antibody chain are derived from a human immunoglobulin, thereby producing the vector of the invention;
- 10 (b) constructing, by conventional techniques, an expression vector containing an operon with a DNA sequence encoding a complementary antibody light chain in which the CDRs and such minimal portions of the variable domain framework region that are required to retain donor antibody binding specificity are derived from a non-human immunoglobulin, and the remaining parts of the antibody chain are derived from a human immunoglobulin,
- 15 thereby producing the vector of the invention;
- (c) transfecting the expression vectors into a host cell by conventional techniques to produce the transfected host cell of the invention; and
- (d) culturing the transfected cell by conventional techniques to produce the altered antibody of the invention.

20 The host cell may be cotransfected with the two vectors of the invention, the first vector containing an operon encoding a light chain derived polypeptide and the second vector containing an operon encoding a heavy chain derived polypeptide. The two vectors contain different selectable markers, but otherwise, apart from the antibody heavy and light chain coding sequences, are preferably identical, to ensure, as far as possible, equal

25 expression of the heavy and light chain polypeptides. Alternatively, a single vector may be used, the vector including the sequences encoding both the light and the heavy chain polypeptides. The coding sequences for the light and heavy chains may comprise cDNA or genomic DNA or both.

The host cell used to express the altered antibody of the invention may be

30 either a bacterial cell such as Escherichia coli, or a eukaryotic cell. In particular a mammalian cell of a well defined type for this purpose, such as a myeloma cell or a Chinese hamster ovary cell may be used.

The general methods by which the vectors of the invention may be constructed, transfection methods required to produce the host cell of the invention and

35 culture methods required to produce the antibody of the invention from such host cells are all conventional techniques. Likewise, once produced, the humanized antibodies of the invention may be purified according to standard procedures of the art, including cross-flow

filtration, ammonium sulphate precipitation, affinity column chromatography, gel electrophoresis and the like.

It should be understood that the humanized antibodies of this invention perform in a manner which is the same or similar to that of the non-humanized versions of the same antibodies. It also is noted that the humanized antibodies of this invention may be used for the design and synthesis of either peptide or non-peptide compounds (mimetics) which would be useful for the same therapy as the antibody (Saragobi et al., Science 253:792-795 (1991)).

The following examples are provided as a further illustration of the present invention and should in no way be construed as being limiting.

EXAMPLES

In the following examples all necessary restriction and modification enzymes, plasmids and other reagents and materials were obtained from commercial sources unless otherwise indicated.

In the following examples, unless otherwise indicated, all general recombinant DNA methodology was performed as described in "Molecular Cloning, A Laboratory Manual" (1982) Eds T. Maniatis et al., published by Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York.

In the following examples the following abbreviations were employed:

dCTP	deoxycytidine triphosphate
dATP	deoxyadenosine triphosphate
dGTP	deoxyguanosine triphosphate
dTTP	deoxythymidine triphosphate
DTT	dithiothreitol
C	cytosine
A	adenine
G	guanine
T	thymine
PBS	phosphate buffered saline
PBST	phosphate buffered saline containing 0.05% Tween 20 (pH 7.5)

Example 1 - Production of Humanized Antibodies Specific for an Fc Receptor

The source of the donor CDRs used to prepare the humanized antibody was a murine monoclonal antibody, mab 22, which is specific for the Fc receptor. A mab 22

hybridoma cell line (022WCL-1) was established. Cytoplasmic RNA was prepared from the mab 22 cell line using the method described by Favoloro et al. (Methods in Enzymology 65, 718-749 (1980)). The cDNA

was synthesized using IgG1 and kappa constant region primers. The primer CG1FOR was used for the heavy chain variable (VH) region and the primer CK2FOR was used for the Ig kappa chain variable region (VK). The cDNA synthesis reactions mixtures consisted of 1 µg RNA, 0.5µM CG1FOR or CK2FOR, 250 µM each of dATP, dCTP, dGTP, and dTTP, 50 mM Tris HCl (pH 7.5), 75 mM KCl, 10 mM dithiothreitol, 3 mM MgCl₂ and 20µ RNAGuard* (sold by Pharmacia, Milton Keynes, U.K.) in a total volume of 50 µl. The samples were heated at 72°C for two minutes and slowly cooled to 37°C. Murine moloney leukemia virus reverse transcriptase (100 µl - sold by Life Technologies, Paisley, U.K.) was added to the samples and the transcriptase containing samples were incubated at 42°C for sixty minutes.

VH and VK cDNAs were then amplified using the polymerase chain reaction (PCR) as described by Saiki et al. (Science 239, 487-491 (1988)).

The primers used in the above steps were as follows:

CG1FOR (SEQ ID NO:5) 5' GGAAGCTTAGACAGATGGGGGTGTCGTTTTG 3'

VH1FOR (SEQ ID NO:6) 5' TGAGGAGACGGTGACCGTGGTCCCTTGGCCCCAG 3'

VH1BACK (SEQ ID NO: 7) 5' AGGTSMARCTGCAGSAGTCWGG 3'

SH1BACK (SEQ ID NO:8) 5' TGGAATTCATGGRATGGAGCTGGRTCWTBHTCTT 3'

SH2BACK (SEQ ID NO:9) 5' TGGAATTCATGTRACTTCDGGYTCAACTKRRTTT 3'

CK2FOR (SEQ ID NO:10) 5' GGAAGCTTGAAGATGGATACAGTTGGTGCAGC 3'

VK1BACK (SEQ ID NO:11) 5' GACATTCAGCTGACCCAGTCTCCA 3'

VK5BACK (SEQ ID NO:12) 5' TTGAATTCGGTGCCAGAKCWSAHATYGTKATG 3'

VK6BACK (SEQ ID NO:13) 5' TTGAATTCGGTGGCAGAKCWSAHATYGTKCTC 3'

VK7BACK (SEQ ID NO:14) 5' TTGAATTCGGAGCTGATGGGAACATTGTAATG 3'

Restriction sites incorporated in primers to facilitate cloning are underlined.

The PCR amplification of murine Ig DNA was conducted using the methodology described by Orlandi et al. (Proc. Natl. Acad. Sci USA 86, 3833-3838 (1989)). The DNA/primer mixtures

consisted of RNA/cDNA hybrid (10 µl) and 25pmol each of CG1FOR and SH1BACK or SH2BACK for PCR amplification of VH. The DNA/primer mixtures consisted of RNA/cDNA hybrid (10 µl) and 25pmol each of CK2FOR and VK1BACK, VK5BACK, VK6BACK, VK7BACK for PCR amplification of VK. dATP, dCTP, dGTP and dTTP (250

*Trade-mark

5 μ M each), 10mM Tris HCl (pH 8.3), 60mM KCl, 1.5mM MgCl₂, 0.01% (w/v) gelatin, 0.01% (v/v) Tween 20*, 0.01% (v/v) NP40 and 2.5 μ Amplitaq* (sold by Cetus, Beaconsfield, U.K.) were added to the samples in a final volume of 50 μ l. The samples were subjected to 25-30 thermal cycles of PCR at 94°C for thirty seconds, 55°C for thirty seconds, 72°C for one minute and a final cycle at 72°C for five minutes.

10 The amplified VH and VK DNAs were run on a low melting point agarose gel and purified by Elutip-d* column chromatography (sold by Schleicher and Schueell, Anderman, Walton, U.K.) for cloning and sequencing. The purified VH DNAs were cut with Eco I or Pst I and Hind III and cloned into M13mp18 and mp19 (sold by Pharmacia, Milton Keynes, U.K.). The purified VK DNAs were cut with Pvu II or Eco I and Hind III and cloned into M13mp18 and mp19. For general cloning methodologies see Sambrook et al. (Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY (1989)), the contents of which are expressly incorporated by reference. The resulting collection of clones were sequenced by the dideoxy method using T7 DNA polymerase (sold by Pharmacia, Milton Keynes, U.K.) as described by Sanger et al. (Proc. Natl. Acad. Sci. USA 74, 5463-5467, (1979)).

20 From the sequences of the O22 VH and VR domains the CDR sequences were determined with reference to the database of Kabat et al. ("Sequences of Proteins of Immunological Interest" US Department of Health and Human Services, US Government Printing Office) and utilizing computer assisted alignment with other VH and VK sequences.

25 Transfer of the murine O22 CDRs to human frameworks was achieved by oligonucleotide site-directed mutagenesis as described by Nakamye et al. (Nucleic Acids Res 14, 9679-9687 (1986)). The primers used were as follows:

KL VHCDR1 (SEQ ID NO:15): 5' TGCCTGTCTCACCCAATACATGTAA
TTGTCACTGAAATGAAGCCAGACGMGGAGCGGACAG

30

KL VHCDR2 (SEQ ID NO:16): 5' TGTAATCTTCCCTTCACACTGTCTGGATAGTA
GGTGTAACCTACCACCATCACTAATGGTTGCAACCCACTCAGG

KL VHCDR3 (SEQ ID NO:17): 5' GGGGTCCCTTGGCCCCAGTAGTCCATAGC
35 CCCCTCGTACCTATAGTAGCCTCTTGCAAAAAATAGA

NM VHCDR1 (SEQ ID NO:18): 5' TGGCTGTCTCACCCAATACATGTAAATTGT
CGCTGAAAATGAAGCCAGACACGGTGCAGGTCAGGCTCA

*Trade-mark

NMVHCDR2 (SEQ ID NO:19): 5' TTGCTGGTGTCTCTCAGCATTGTCACTCTC
CCCTTCACACTGTCTGGATAGTAGGTGTA ACTACCA
TCACTAATGGTTCCAATCCACTCAA

5

NMVHCDR3 (SEQ ID NO:20): 5' AGACGGTGACCAAGGACCCTTGGCCCCAG
TAGTCCATAGCCCCCTCGTACCTATAGTAGCCTCTTGACACAATAATAG

10

HuVKCDR1 (SEQ ID NO:21): 5' CTTCTGCTGGTACCAGGCCAAGTAGTTCTTC
TGATTTGAACTGTATAAAACACTTTGACTGGACTTACAGGTGATGGTCAC

HuVKCDR2 (SEQ ID NO:22): 5' GCTTGGCACACCAGATTCCCTAGTGGATG
CCCAGTAGATCAGCAG

15

HuVKCDR3 (SEQ ID NO:23): 5' CCTTGGCCGAACGTCCACGAGGAGAGGTAT
TGATGGCAGTAGTAGGTGG

20 The primer for NMVHCDR1 was extended to include a change of NEWM residues Ser 27
Thr 28 to Phe 27 Ile 28. The primer for NMVHCDR2 was extended to include a change of
NEWM residue Val 71 to Arg 71.

25 The DNA templates used for mutagenesis of VHs comprised human
framework regions from the crystallographically solved protein NEW described by Saul et al.
(J. Biol. Chem. 53, 585-597 (1978)) or KOL described by Schmidt et al. (Z. Physical Chem.
364, 713-747 (1983)). The DNA templates used for mutagenesis of VKs comprised human
framework regions from the crystallographically solved protein REI described by Epp et al.
(Eur. J. Biochem. 45, 513-524 (1974)).

30 M13 based templated M13VHPCR1 (for NEWMVH), M13VHPCR2 (for
KOLVH) and M13VKPCR2 (for REIVK) comprising human frameworks with irrelevant
CDRs were prepared as described by Riechmann et al. (Nature 332, 323-327 (1988)), the
contents of which are expressly incorporated by reference. Oligonucleotide site-directed
mutagenesis was carried out using the following protocol. A 5-fold molar excess of each
phosphorylated mutagenic oligonucleotide was added along with the universal M13
35 sequencing primer (5'- GTAAAACGACGGCCAGT) (SEQ ID NO:24). All of the primers
were annealed in 20ul 0.1M TrisHCl (pH8.0) and 10mM MgCl₂ by heating to 70-85°C for
two minutes and slowly cooling to room temperature. 10 mM DTT, 1 mM ATP, 40 µM each
of dATP, dCTP, dGTP and dTTP, 2.5µ T7 DNA polymerase (sold by United States

Biochemicals) and 0.5µ T4DNA ligase (sold by Life Technologies, Paisley, U.K.) was added to the annealed DNA in a reaction volume of 30µl and incubated at 22° - 37°C for one to two hours. The newly extended and ligated strand was preferentially amplified over the parental strand in a thermostable DNA polymerase directed reaction using the M13 reverse

5 sequencing primer (5' AACAGCTATGACCATG) (SEQ ID NO:25). The reverse sequencing primer is not complementary to the parental strand. The reaction mixture of 50µl contained 1 µl extension/ligation product, 25 pmol M13 reverse sequencing primer, 250µM each of dATP, dCTP, dGTP and dTTP, 1µ Vent DNA polymerase (sold by New England Biolabs, Bishop's Stortford, U.K.) or 2.5µ Amplitaq^{*} (sold by Cetus, Beaconsfield, U.K.) in the
10 appropriate buffer supplied by the enzyme manufacturer and was subjected to thirty thermal cycles of 94°C, 30s, 55°C, 30s, 75° or 72°C, 90s; ending with 5 min at 72°C. A 4µl aliquot of this sample was then amplified by PCR using both M13 universal and reverse sequencing primers in a reaction mixture of 50µl containing 25 pmol of each primer, 250µM each of dATP, dCTP, dGTP and dTTP, 2.5µ Amplitaq^{*} (Cetus) in the buffer supplied by the enzyme
15 manufacturer. Amplified DNAs were digested with HindIII and BamHI and cloned into M13mp19 and sequenced.

Mutagenesis of M13VHPCR2 KOL VH residue Leu71 to Arg71 was by the overlap/extension PCR method of Ho *et al.* (*Gene*, 77, 51-55 (1989)).

The overlapping oligonucleotides used were 5' -

20 TTTACAATATCGAGACAACAGCAA (SEQ ID NO:26) and
5' - TTGCTGTTGTCTCTCGATTGTAAA (SEQ ID NO:27).

The amino acid sequences of the humanized antibodies were compared to the known murine antibodies as shown in Figures 1 and 2. The CDR replaced VH and VK genes were cloned into expression vectors pSVgpt and pSVhyg as shown in Figures 3 and 4 as
25 described by Orlandi *et al.* (cited supra). The CDR replaced NEWMVH and KOLVH genes together with the Ig heavy chain promoter, appropriate splice sites and signal peptide sequences were excised from M13 by digestion with HindIII and BamHI and cloned into the pSVgpt expression vector containing the murine Ig heavy chain enhancer, the gpt gene for selection in mammalian cells and genes for replication and selection in *E. coli*. The plasmid
30 also contains a human IgG1 constant region as described by Takahashi *et al.* (*Cell* 29, 671-675 (1982)). The construction of the kappa chain expression vector was essentially the same except that the gpt gene was replaced by the hygromycin resistance gene and contains a human kappa constant region (Hieter *et al.*, *Cell* 22, 197-207 (1980)).

35 Approximately 5µg of each heavy chain expression vector and 10µg of the kappa chain expression vector were digested with PvuI. The DNAs were mixed together, ethanol precipitated and dissolved in 25µl water. Approximately 5-10 x 10⁶ NSO cells (from European Collection of Animal Cell Cultures, Porton Down, U.K.) were grown to semi-

*Trade-mark

confluency in Dulbecco's modified Eagle's medium (DMEM) plus 10% fetal calf serum (Myoclon^{*} plus, Gibco, Paisley, Scotland), harvested by centrifugation and resuspended in 0.5ml DMEM together with the digested DNA in a cuvette. After five minutes in ice, the cells were given a single pulse of 170V at 960 μ F (Gene-Pulser^{*}, Bio-Rad, Richmond, CA) and left in ice for a further twenty minutes. The cells were then put into 20ml DMEM + supplemented with 10% FCS and allowed to recover for twenty-four to forty-eight hours. After this time, the cells were distributed into a 24-well plate and selective medium was applied (DMEM, 10% FCS, 0.8 μ g/ml mycophenolic acid and 250 μ g/ml xanthine). After three to four days, the medium and dead cells were removed and replaced with fresh selective medium. Transfected clones were visible with the naked eye ten days later.

The presence of human antibody in the medium of wells containing gpt⁺ transfectants was measured using conventional enzyme linked immunosorbent assay (ELISA) techniques. Wells of a microtitre plate (Immolon^{*}, Dynatech, Chantilly, VA) were coated with 100ng goat anti-human IgG antibodies (SeraLab, Crawley Down, U.K.) in 100 μ l 50mM carbonate buffer pH9.6. After washing with PBST (Phosphate buffered saline pH 7.2 containing 0.05% Tween 20) culture medium in 100 μ l PBST (5-50 μ l) was added to each well for one hour at 37°C. The wells were then emptied, washed with PBST and 100 μ l of 1:1000 dilution peroxidase conjugated goat anti-human kappa constant region antibodies (SeraLab, Crawley Down, U.K.) were added for one hour at 37°C. The wells were emptied, washed with PBST and 100 μ l OPD substrate buffer (400 μ g/ml Q-phenylenediamine in 24mM citrate/42mM sodium phosphate pH 5. and 0.0003% (v/v) H₂O₂) was added. The reaction was stopped after a few minutes by the addition of 12.5% H₂SO₄ (25 μ l) and the absorbance at 492nm was measured.

The antibody secreting cells were expanded and antibody was purified from the culture medium by protein A affinity chromatography as described by Harlow and Lane (Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY).

The binding of the antibodies to antigen was measured by ELISA. Wells of a microtitre plate (Immulon 1, Dynatech, Chantilly, VA) were coated with 200ng goat anti-human IgM antibodies (Sera-lab, Crawley Down, U.K.) in 100 μ l 50mM carbonate buffer pH 9.6 at 37°C for at least one hour. Wells were emptied and washed once with PBST and blocked with 1% BSA in PBS at room temperature for thirty minutes. The wells were emptied and washed with PBST and Cos supernatant containing FcRI/IgM fusion protein was added and incubated for one hour at room temperature. Wells were then emptied and washed three times with PBST and test antibodies diluted in 1% BSA/PBS were added and incubated for one hour at room temperature. In addition, each well contained 2 μ g human IgG1, lambda antibody (Sigma, Poole, U.K.) The wells were then emptied, washed three times with PBST and 40ng peroxidase goat anti-human kappa constant region antibodies (Sera-Lab, Crawley

*Trade-mark

Down, U.K.) in 100 μ l 1% BSA/PBS added to each well. After incubation for one hour at room temperature, the wells were emptied, washed three time with PBST and 10 μ l HPD substrate buffer was added. The reaction was stopped by the addition of 25 μ l of 12.5% H₂SO₄ to each well. The absorbance at 492nm was measured and is depicted in Figure 5.

- 5 The test antibodies were the antibody containing irrelevant CDRs (AA), the fully humanized KOL/REI based antibody (KL VHR/HuVK), the mix and match derivatives of the humanized antibody (KL VHR/MuVK and MuVH/HuVK), the humanized NEWM/REI based antibody (NMVK/HuVK) and the chimeric antibody (MuVH/MuVK).

10 **EQUIVALENTS**

Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

-13-

SEQUENCE LISTING

(1) GENERAL INFORMATION:

5

(i) APPLICANT:

- (A) NAME: MEDAREX, INC.
- (B) STREET: 22 Chambers Street
- (C) CITY: Princeton
- 10 (D) STATE: New Jersey
- (E) COUNTRY: USA
- (F) POSTAL CODE (ZIP): 08542
- (G) TELEPHONE: (609) 921-7121
- (H) TELEFAX: (609) 921-7450

15

- (ii) TITLE OF INVENTION: HUMANIZED ANTIBODIES TO Fc RECEPTORS FOR
IMMUNOBLOBULIN G ON HUMAN MONONUCLEAR PHAGOCYTES

- (iii) NUMBER OF SEQUENCES: 28

20

(iv) COMPUTER READABLE FORM:

- (A) MEDIUM TYPE: Floppy disk
- (B) COMPUTER: IBM PC compatible
- (C) OPERATING SYSTEM: PC-DOS/MS-DOS
- 25 (D) SOFTWARE: ASCII text

(v) CURRENT APPLICATION DATA:

- (A) APPLICATION NUMBER:
- (B) FILING DATE:
- 30 (C) CLASSIFICATION:

(vi) PRIOR APPLICATION DATA:

- (A) APPLICATION NUMBER: GB92 23377.4
- (B) FILING DATE: 04-NOV-1992
- 35 (C) CLASSIFICATION:

(vii) ATTORNEY/AGENT INFORMATION:

- (A) NAME: Mandragouras, Amy E.
- (B) REGISTRATION NUMBER: 36.207
- 40 (C) REFERENCE/DOCKET NUMBER: MXI-013PC

(ix) TELECOMMUNICATION INFORMATION:

- (A) TELEPHONE: (617) 227-7400
- (B) TELEFAX: (617) 227-5941

45

(2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 120 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

50

(ii) MOLECULE TYPE: peptide

55

-14-

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

5 Gln Val Gln Leu Gln Glu Ser Gly Pro Gly Leu Val Arg Pro Ser Gln
 1 5 10 15
 Thr Leu Ser Leu Thr Cys Thr Val Ser Gly Phe Ile Phe Ser Asp Asn
 20 25 30
 10 Tyr Met Tyr Trp Val Arg Gln Pro Pro Gly Arg Gly Leu Glu Trp Ile
 35 40 45
 Gly Thr Ile Ser Asp Gly Gly Ser Tyr Thr Tyr Tyr Pro Asp Ser Val
 50 55 60
 15 Lys Gly Arg Val Thr Met Leu Arg Asp Thr Ser Lys Asn Gln Phe Ser
 65 70 75 80
 Leu Arg Leu Ser Ser Val Thr Ala Ala Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 20 Ala Arg Gly Tyr Tyr Arg Tyr Glu Gly Ala Met Asp Tyr Trp Gly Gln
 100 105 110
 25 Gly Ser Leu Val Thr Val Ser Ser
 115 120

(2) INFORMATION FOR SEQ ID NO: 2:

30 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 120 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

35 (ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

40 Glu Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg
 1 5 10 15
 45 Ser Leu Arg Leu Ser Cys Ser Ser Ser Gly Phe Ile Phe Ser Asp Asn
 20 25 30
 Tyr Met Tyr Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45
 50 Ala Thr Ile Ser Asp Gly Gly Ser Tyr Thr Tyr Tyr Pro Asp Ser Val
 50 55 60
 55 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Phe
 65 70 75 80
 Leu Gln Met Asp Ser Leu Arg Pro Glu Asp Thr Gly Val Tyr Phe Cys
 85 90 95

5

55

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

5 Asp Ile Gln Leu Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1 5 10 15
Asp Arg Val Thr Ile Thr Cys Lys Ser Ser Gln Ser Val Leu Tyr Ser
20 25 30
10 Ser Asn Gln Lys Asn Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Lys
35 40 45
Ala Pro Lys Leu Leu Ile Tyr Trp Ala Ser Thr Arg Glu Ser Gly Val
50 55 60
15 Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Phe Thr
65 70 75 80
Ile Ser Ser Leu Gln Pro Glu Asp Ile Ala Thr Tyr Tyr Cys His Gln
85 90 95
20 Tyr Leu Ser Ser Trp Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys
100 105 110

25 (2) INFORMATION FOR SEQ ID NO: 5:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 31 base pairs
(B) TYPE: nucleic acid
30 (C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

35

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

40 GGAAGCTTAG ACAGATGGGG GTGTCGTTTT G

31

(2) INFORMATION FOR SEQ ID NO: 6:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 34 amino acids
45 (B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

50

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

55 Thr Gly Ala Gly Gly Ala Gly Ala Cys Gly Gly Thr Gly Ala Cys Cys
1 5 10 15

-17-

Gly Thr Gly Gly Thr Cys Cys Cys Thr Thr Gly Gly Cys Cys Cys Cys
20 25 30

Ala Gly

5

(2) INFORMATION FOR SEQ ID NO: 7:

(i) SEQUENCE CHARACTERISTICS:

10

- (A) LENGTH: 22 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

15

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

20

AGGTSMARCT GCAGSAGTCW GG

22

(2) INFORMATION FOR SEQ ID NO: 8:

25

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 34 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

30

(ii) MOLECULE TYPE: cDNA

35

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

TGGAATTCAT GGRATGGAGC TGGRTCWTBH TCTT

34

(2) INFORMATION FOR SEQ ID NO: 9:

40

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 33 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

45

(ii) MOLECULE TYPE: cDNA

50

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:

TGGAATTCAT GRACTTCDGG YTCAACTKRR TTT

33

55

-18-

(2) INFORMATION FOR SEQ ID NO: 10:

(i) SEQUENCE CHARACTERISTICS:

5

- (A) LENGTH: 32 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

10

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:

15

GGAAGCTTGA AGATGGATAC AGTTGGTGCA GC

32

(2) INFORMATION FOR SEQ ID NO: 11:

(i) SEQUENCE CHARACTERISTICS:

20

- (A) LENGTH: 24 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

25

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11:

30

GACATTCAGC TGACCCAGTC TCCA

24

(2) INFORMATION FOR SEQ ID NO: 12:

(i) SEQUENCE CHARACTERISTICS:

35

- (A) LENGTH: 32 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

40

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 12:

45

TTGAATTCGG TGCCAGAKCW SAHATYGTKA TG

32

(2) INFORMATION FOR SEQ ID NO: 13:

(i) SEQUENCE CHARACTERISTICS:

50

- (A) LENGTH: 32 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

55

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 13:

TTGAATTCGG TGGCAGAKCW SAHATYGTKC TC

32

5 (2) INFORMATION FOR SEQ ID NO: 14:

(i) SEQUENCE CHARACTERISTICS:

- 10 (A) LENGTH: 32 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

15

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 14:

TTGAATTCGG AGCTGATGGG AACATTGTAA TG

32

20

(2) INFORMATION FOR SEQ ID NO: 15:

(i) SEQUENCE CHARACTERISTICS:

- 25 (A) LENGTH: 61 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

30

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 15:

35 TGCCTGTCTC ACCCAATACA TGTAATTGTC ACTGAAATGA AGCCAGACGM GGAGCGGACA

60

G

61

(2) INFORMATION FOR SEQ ID NO: 16:

40

(i) SEQUENCE CHARACTERISTICS:

- 45 (A) LENGTH: 75 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

50

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 16:

TGTAAATCTT CCCTTCACAC TGTCTGGATA GTAGGTGTAA CTACCACCAT CACTAATGGT

60

55 TGCAACCCAC TCAGG

75

(2) INFORMATION FOR SEQ ID NO: 17:

(i) SEQUENCE CHARACTERISTICS:

5

- (A) LENGTH: 67 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

10

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 17:

15

GGGGTCCCTT GGCCCCAGTA GTCCATAGCC CCCTCGTACC TATAGTAGCC TCTTGCACAA
AAATAGA

60

67

(2) INFORMATION FOR SEQ ID NO: 18:

20

(i) SEQUENCE CHARACTERISTICS:

25

- (A) LENGTH: 68 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

30

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 18:

35

TGGCTGTCTC ACCCAATACA TGTAATTGTC GCTGAAAATG AAGCCAGACA CGGTGCAGGT
CAGGCTCA

60

68

(2) INFORMATION FOR SEQ ID NO: 19:

(i) SEQUENCE CHARACTERISTICS:

40

- (A) LENGTH: 94 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

45

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 19:

50

TTGCTGGTGT CTCTCAGCAT TGTCACCTCTC CCCTTCACAC TGTCTGGATA GTAGGTGTAA
CTACCACCAT CACTAATGGT TCCAATCCAC TCAA

60

94

55

2148578

(2) INFORMATION FOR SEQ ID NO: 20:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 77 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 20:

AGACGGTGAC CAAGGACCCT TGGCCCCAGT AGTCCATAGC CCCCTCGTAC CTATAGTAGC
CTCTTGCACA ATAATAG

(2) INFORMATION FOR SEQ ID NO: 21:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 81 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 21:

CTTCTGCTGG TACCAGGCCA AGTAGTTCTT CTGATTTGAA CTGTATAAAA CACTTTGACT
GGACTTACAG GTGATGGTCA C

(2) INFORMATION FOR SEQ ID NO: 22:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 45 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 22:

GCTTGGCACA CCAGATTCCC TAGTGGATGC CCAGTAGATC AGCAG

-22-

(2) INFORMATION FOR SEQ ID NO: 23:

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 49 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

10

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 23:

15 CCTTGCCCGA ACGTCCACGA GGAGAGGTAT TGATGGCAGT AGTAGGTGG 49

(2) INFORMATION FOR SEQ ID NO: 24:

(i) SEQUENCE CHARACTERISTICS:

- 20 (A) LENGTH: 17 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

25

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 24:

30 GTAAAACGAC GGCCAGT 17

(2) INFORMATION FOR SEQ ID NO: 25:

35

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 16 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- 40 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

45

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 25:

AACAGCTATG ACCATG 16

50 (2) INFORMATION FOR SEQ ID NO: 26:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24 base pairs
- (B) TYPE: nucleic acid
- 55 (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

-23-

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 26:

TTTACAATAT CGAGACAACA GCAA

24

5

(2) INFORMATION FOR SEQ ID NO: 27:

(i) SEQUENCE CHARACTERISTICS:

10

(A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

15

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 27:

20

TTGCTGTTGT CTCTCGATTG TAAA

24

(2) INFORMATION FOR SEQ ID NO: 28:

25

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 112 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

30

(ii) MOLECULE TYPE: peptide

35

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 28:

Asn	Ile	Val	Met	Thr	Gln	Ser	Pro	Ser	Ser	Leu	Ala	Val	Ser	Ala	Gly
1				5				10					15		

40

Glu	Lys	Val	Thr	Met	Ser	Cys	Lys	Ser	Ser	Gln	Ser	Val	Leu	Tyr	Ser
			20				25						30		

Ser	Asn	Gln	Lys	Asn	Tyr	Leu	Ala	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Gln
		35				40						45			

45

Ser	Pro	Lys	Leu	Leu	Ile	Tyr	Trp	Ala	Ser	Thr	Arg	Glu	Ser	Gly	Val
	50				55					60					

50

Pro	Asp	Arg	Phe	Thr	Gly	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Thr
65				70				75						80	

Ile	Ser	Ser	Val	Gln	Ala	Glu	Asp	Leu	Ala	Val	Tyr	Tyr	Cys	His	Gln
				85				90					95		

55

Tyr	Leu	Ser	Ser	Trp	Thr	Phe	Gly	Gly	Gly	Thr	Lys	Leu	Glu	Ile	Lys
				100				105					110		

-24-

CLAIMS:

1. A humanized antibody specific to an Fc receptor Fc γ RI comprising:
a human antibody having at least a portion of a complementarity determining region derived from a non-human antibody, the portion being selected to provide specificity of the humanized antibody for a human Fc receptor, wherein the portion is derived from the humanized antibody produced by the cell line having ATCC Number CRL 11177.
2. The antibody of claim 1 wherein the portion is selected to provide specificity to a human Fc receptor such that the humanized antibody formed binds to the Fc receptor at a site which is not blocked by human immunoglobulin G.
3. The antibody of claim 1 wherein the human antibody is derived from NEWM, KOL, REI protein or a combination thereof.
4. A bifunctional antibody or heteroantibody, comprising:
at least one humanized antigen binding region derived from the humanized anti-Fc receptor antibody of claim 1, and
at least one antigen binding region specific for a target epitope.
5. The antibody of claim 4 wherein the humanized antigen binding region is derived from a humanized anti-Fc receptor antibody selected such that the binding of the humanized antibody to the human Fc receptor is not blocked by human immunoglobulin G.
6. The antibody of claim 4 wherein the target epitope is that of a cancer cell.
7. The antibody of claim 4 wherein the target epitope is that of an infectious agent.

-25-

8. The antibody of claim 4 wherein the target epitope is that of an antibody-producing cell.

9. A humanized antibody comprising:

a portion which binds to the Fc γ RI receptor for human immunoglobulin G, wherein the antigen binding region comprises all complementarity determining regions (CDRs) from the heavy and light chains of a non-human antibody specific for a human Fc γ RI receptor, and

variable domain framework regions from the heavy and light chains of a human antibody, wherein the heavy chain variable region comprises the amino acid sequence of SEQ ID NO: 2, and the light chain variable region comprises the amino acid sequence of SEQ ID NO: 4.

10. The antibody of claim 9 wherein the antigen binding region is selected to provide specificity to a human Fc γ RI receptor such that the humanized antibody formed binds to the Fc γ RI receptor at a site which is not blocked by human immunoglobulin G.

11. The humanized antibody of claim 9 produced by the cell line having ATCC Accession Number CRL 11177.

12. A bifunctional antibody or heteroantibody, comprising:

at least one humanized antigen binding portion derived from a humanized anti-Fc γ RI receptor antibody, wherein said portion is derived from the humanized antibody produced by the cell line having ATCC Number CRL 11177, and

at least one antigen binding region specific for a target epitope.

13. A bifunctional antibody or heteroantibody, comprising:

a portion which binds to the Fc γ RI receptor for human immunoglobulin G, wherein the antigen binding region comprises all complementarity determining regions

-26-

(CDRs) from the heavy and light chains of a non-human antibody specific for a human FcγRI receptor,

variable domain framework regions from the heavy and light chains of a human antibody, wherein the heavy chain variable region comprises the amino acid sequence of SEQ ID NO: 2, and the light chain variable region comprises the amino acid sequence of SEQ ID NO: 4,

and an antigen binding region specific for an antigen present on a target cell.

14. The bifunctional molecule of claim 12 or 13, wherein the target cell is a cancer cell.

15. The bifunctional molecule of claim 12 or 13, wherein the antigen is HER 2/neu.

16. A humanized antibody comprising:

(a) an antigen binding region comprising (i) at least a portion of a complementarity determining region (CDR) from a heavy or a light chain of a murine antibody specific for a human Fc receptor; and (ii) such minimal portions of the variable domain framework region of the murine antibody that are required to retain the binding specificity of the murine antibody, wherein the variable region of said heavy chain comprises the amino acid sequence shown in SEQ ID NO: 1 or 2 and the variable region of said light chain comprises the amino acid sequence shown in SEQ ID NO: 4; and

(b) variable domain framework regions from the heavy and light chains of a human antibody, wherein said humanized antibody induces similar biological effects upon binding to the human Fc receptor as its monoclonal counterpart.

17. The antibody of claim 16 wherein said at least one portion of a complementarity determining region from a heavy or light chain of murine antibody is derived from a non-human antibody.

-27-

18. The antibody of claim 16, which is produced by the hybridoma having ATCC Accession Number CRL 11177.

1 / 6

	10V	20V	30V	40V	50V
022NMVH	QVQLQESGPGGLVRPSQTLTLCTVSGFIFSDNYMYWVRQPPGRGLEWIGTI				
	:VQL ESG GLV:P: :L:L:C..SGFIFSDNYMYWVRQ.P.: LEW::TI				
022MUVH	EVQLVESGGGLVKPGGSLRLSCVASGFIFSDNYMYWVRQTPEKRLEWVATI				
	10 [^]	20 [^]	30 [^]	40 [^]	50 [^]
	60V	70V	80V	90V	100V
022NMVH	SDGGSYTYYPDSVKGRVTMLRDTSKNQFSLRLSSVTAADTAVYYCARGYYR				
	SDGGSYTYYPDSVKGR T: RD.:KN:: L::SS:::DTA:YYCARGYYR				
022MUVH	SDGGSYTYYPDSVKGRFTISRDNAKNNLYLQMSSLKSEDTAIYYCARGYYR				
	60 [^]	70 [^]	80 [^]	90 [^]	100 [^]
	110V 120V				
022NMVH	YEGAMDYWGQGSLVTVSS				
	YEGAMDYWGQG: VTVSS				
022MUVH	YEGAMDYWGQGTSTVTVSS				
	110 [^] 120 [^]				

FIG. 1A

022KLVH	10V	20V	30V	40V	50V
	EVQLVESGGGVVQPGRSLRLSCSSSGFIFSDNYMYWVRQAPGKGLEWVATI				
	EVQLVESGGG:V:PG SLRLSC :SGFIFSDNYMYWVRQ:P.K LEWVATI				
022MUVH					
	EVQLVESGGGLVKPGGSLRLSCVASGFIFSDNYMYWVRQTPEKRLEWVATI				
	10^	20^	30^	40^	50^
	60V	70V	80V	90V	100V
022KLVH					
	SDGGSYTYYPDSVKGRFTISRDN SKNTLFQMDSL RPEDTG VYFCARGYYR				
	SDGGSYTYYPDSVKGRFTISRDN:KN.LI.QM.SLLED TLIYICARGYYR				
022MUVH					
	SDGGSYTYYPDSVKGRFTISRDN AKNNLYLQMSSLKSEDTAIYYCARGYYR				
	60^	70^	80^	90^	100^
	110V	120V			
022KLVH					
	YEGAMDYWGQGTPVTVSS				
	YEGAMDYWGQGT:VTVSS				
022MUVH					
	YEGAMDYWGQGT:VTVSS				
	110^	120^			

FIG. 1B

2148578


3 / 6

022MUVK	10V	20V	30V	40V	50V
	NIVMTQSPSSLAVSAGEKVTMSCKSSQSVLYSSNQKNYLAWYQQKPGQSPK				
	: I : TQSPSSL: .S.G: : VT: : CKSSQSVLYSSNQKNYLAWYQQKPG: : PK				
022HUVK	10^	20^	30^	40^	50^
	60V	70V	80V	90V	100V
	LLIYWASTRESCVPDRFTGSGSGTDFTLTISSVQAEDLAVYYCHQYLSSWT				
	LLIYWASTRESCVP.RF:GSGSGTDFTLTISS:Q:ED:A.YYCHQYLSSWT				
022HUVK	60^	70^	80^	90^	100^
	LLIYWASTRESCVPSRFSGSGSGTDFTFISSLQPEDIAITYYCHQYLSSWT				
	110V				
022MUVK	FGGGTKLEIK				
	FG GTK:EIK				
022HUVK	FGQGTKVEIK				
	110^				

FIG. 2


4 / 6

FIG. 3



5 / 6


FIG. 4




2148578

6 / 6


FIG. 5



CHIMAERIC



RESHAPED



Igh enhancer

Ap^r

pSVgpt

SV40 polyA

colEI
ori

gpt

SV40
promoter