



(86) **Date de dépôt PCT/PCT Filing Date:** 2016/08/03
(87) **Date publication PCT/PCT Publication Date:** 2017/11/23
(45) **Date de délivrance/Issue Date:** 2019/08/27
(85) **Entrée phase nationale/National Entry:** 2018/11/09
(86) **N° demande PCT/PCT Application No.:** IB 2016/054672
(87) **N° publication PCT/PCT Publication No.:** 2017/199070
(30) **Priorité/Priority:** 2016/05/18 (IN201641017204)

(51) **Cl.Int./Int.Cl.** **A61K 31/13** (2006.01),
A61K 31/445 (2006.01), **A61K 31/496** (2006.01),
A61P 25/28 (2006.01)


(72) **Inventeurs/Inventors:**
NIROGI, RAMAKRISHNA, IN;
SHINDE, ANIL KARBHARI, IN;
JAYARAJAN, PRADEEP, IN;
BHYRAPUNENI, GOPINADH, IN;
KAMBHAMPATI, RAMASASTRI, IN;
JASTI, VENKATESWARLU, IN

(73) **Propriétaire/Owner:**
SUVEN LIFE SCIENCES LIMITED, IN

(74) **Agent:** AIRD & MCBURNEY LP

(54) **Titre :** TRIPLE COMBINAISON D'ANTAGONISTES PURS DU RECEPTEURS 5-HT₆, D'INHIBITEURS DE
L'ACÉTYLCHOLINESTERASE ET D'ANTAGONISTES DU RECEPTEUR NMDA

(54) **Title:** TRIPLE COMBINATION OF PURE 5-HT₆ RECEPTOR ANTAGONISTS, ACETYLCHOLINESTERASE INHIBITORS AND
NMDA RECEPTOR ANTAGONIST



Data represents Mean ± SEM of discriminative index. * $p < 0.05$ Vs vehicle (Kruskal-Wallis followed by Dunn's Multiple Comparison Test) N=11-12

(57) **Abrégé/Abstract:**

The present invention relates to a combination of pure 5-HT₆ receptor antagonist, acetylcholinesterase inhibitor and NMDA receptor antagonist. Also, the present invention provides pure 5-HT₆ receptor (5-HT₆R) antagonists, or the pharmaceutically acceptable salt(s) thereof in combination with or as adjunct to acetylcholinesterase inhibitor and N- Methyl-D-aspartate (NMDA) receptor antagonist and their use in the treatment of cognitive disorders. The present invention further provides the pharmaceutical composition containing the said combination.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau(43) International Publication Date
23 November 2017 (23.11.2017)(10) International Publication Number
WO 2017/199070 A1

(51) International Patent Classification:

A61K 31/13 (2006.01) *A61K 31/496* (2006.01)
A61K 31/445 (2006.01) *A61P 25/28* (2006.01)

(21) International Application Number:

PCT/IB2016/054672

(22) International Filing Date:

03 August 2016 (03.08.2016)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:


201641017204 18 May 2016 (18.05.2016) IN

(71) Applicant: **SUVEN LIFE SCIENCES LIMITED**
[IN/IN]; Serene Chambers, Road - 5, Avenue – 7, Banjara Hills, Hyderabad, Telangana 500 034 (IN).(72) Inventors: **NIROGI, Ramakrishna**; Suven Life Sciences Limited, Serene Chambers, Road-5, Avenue – 7, Banjara Hills, Hyderabad, Telangana 500034 (IN). **SHINDE, Anil Karbhari**; Suven Life Sciences Limited, Serene Chambers, Road-5, Avenue – 7, Banjara Hills, Hyderabad,

Telangana, 500034 (IN). **JAYARAJAN, Pradeep**; Suven Life Sciences Limited, Serene Chambers, Road-5, Avenue – 7, Banjara Hills, Hyderabad, Telangana 500034 (IN). **BHYRAPUNENI, Gopinadh**; Suven Life Sciences Limited, Serene Chambers, Road-5, Avenue – 7, Banjara Hills, Hyderabad, Telangana 500034 (IN). **KAMBHAM-PATI, Ramasastri**; Suven Life Sciences Limited, Serene Chambers, Road-5, Avenue – 7, Banjara Hills, Hyderabad, Telangana 500034 (IN). **JASTI, Venkateswarlu**; Suven Life Sciences Limited, Serene Chambers, Road-5, Avenue – 7, Banjara Hills, Hyderabad, Telangana 500034 (IN).

(74) Agent: **GANDHI, Ritu** et al.; Central Square, Suite-328, Plaza III, 20 Manoharlal Khurana Marg, Bara Hindu Rao, (off Rani Jhansi Road), Delhi 110006 (IN).(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN,(54) Title: TRIPLE COMBINATION OF PURE 5-HT₆ RECEPTOR ANTAGONISTS, ACETYLCHOLINESTERASE INHIBITORS AND NMDA RECEPTOR ANTAGONIST

Figure 1a



Data represents Mean ± SEM of discriminative index. * $p < 0.05$ Vs vehicle (Kruskal-Wallis followed by Dunn's Multiple Comparison Test) N=11-12

(57) Abstract: The present invention relates to a combination of pure 5-HT₆ receptor antagonist, acetylcholinesterase inhibitor and NMDA receptor antagonist. Also, the present invention provides pure 5-HT₆ receptor (5-HT₆R) antagonists, or the pharmaceutically acceptable salt(s) thereof in combination with or as adjunct to acetylcholinesterase inhibitor and N-Methyl-D-aspartate (NMDA) receptor antagonist and their use in the treatment of cognitive disorders. The present invention further provides the pharmaceutical composition containing the said combination.

[Continued on next page]

WO 2017/199070 A1

MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States *(unless otherwise indicated, for every kind of regional protection available):* ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— *as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))*

Published:

— *with international search report (Art. 21(3))*

“TRIPLE COMBINATION OF PURE 5-HT₆ RECEPTOR ANTAGONISTS, ACETYLCHOLINESTERASE INHIBITORS AND NMDA RECEPTOR ANTAGONIST”

FIELD OF THE INVENTION

The present invention relates to pure 5-HT₆ receptor (5-HT₆R) antagonists or the pharmaceutically acceptable salt(s) thereof in combination with or as adjunct to acetylcholinesterase inhibitors and N-Methyl-D-aspartate (NMDA) receptor antagonist. The present invention further relates to the use of the combination and the pharmaceutical composition containing the said combination in the treatment of cognitive disorders.

BACKGROUND OF INVENTION

Alzheimer's disease (AD) is the most common cause of dementia worldwide. The exponential rise in the number of cases of AD in the past and the future projection over the next few decades is anticipated to result in great pressure on the social and health-care systems of developed and developing economies alike. AD also imposes tremendous emotional and financial burden to the patient's family and community.

The current list of approved cognitive enhancing drugs for AD is not long and historically been focused on acetylcholinesterase inhibitors (donepezil, rivastigmine and galantamine). These drugs act by inhibiting the hydrolysis of acetylcholine (ACh) into acetate and choline by targeting acetylcholinesterase (AChE) enzyme. Increasing the ACh levels in the synapse can stimulate cholinergic receptors and promote memory function. Although acetylcholinesterase inhibitors (AChEIs) can temporarily delay the progression of cognitive decline in AD, their effects are modest. ACh being present both in the central and peripheral nervous system, AChEIs produce several undesirable side effects such as gastrointestinal disturbances, bradycardia and excessive salivation that are associated with an action on peripheral muscarinic cholinergic receptors (*Expert Opinion on Drug Safety*, 3, 2004, 425–440). The limitation of AChEI class of drugs is that they are poorly tolerated, their efficacy is not sustained and they require constant dose-titration as the disease progresses (*Cochrane Database Systematic Reviews*, 2006, CD005593) which lead to significant patient noncompliance. The incidence and the severity of these side effects increase with the dose amount and in general more pronounced at the initiation of the

treatment or after dose increase. Hence there is an unmet need of alternate therapy for treating cognition disorders.

5-Hydroxytryptamine 6 receptor (5-HT₆R), a member of GPCR family is exclusively expressed in the brain, particularly in areas associated with cognition, such as hippocampus and frontal cortex (*Molecular Pharmacology*, 1993, 43, 320–327). Activation of 5-HT₆R usually represses cholinergic function (*British Journal of Pharmacology*, 1999, 126, 1537–1542), whereas blockade of the receptor improves the cognitive functions. Thus, 5-HT₆R may be a viable target for pharmacologic intervention to improve the cognitive function of patients with AD. As 5-HT₆R is exclusively located centrally, it is believed that 5-HT₆R antagonists would have limited peripheral side effects, including the ones which are commonly associated with cholinesterase inhibitors. Antagonism of this receptor by several investigational compounds has been shown to improve learning and memory in animal models (*CNS & Neurological Disorders - Drug Targets*, 2004, 3, 59–79).

Since blocking 5-HT₆R modulates cholinergic activity, one might expect 5-HT₆R antagonists to complement and/or augment cognitive function through this therapeutic mechanism. This may in turn help to reduce the side effects with better patient compliance and thus can be administered over a longer period.

The glutamatergic system is also involved in learning and memory and is a target for treatment of Alzheimer's disease. Memantine is another approved treatment for Alzheimer's disease, which acts on the glutamatergic system by inhibiting NMDA receptor under conditions of excess stimulation. It may act to protect glutamate neurons from excessive glutamate stimulation, while increasing the signal to noise ratio. It is known that glutamate neurons have synaptic connections on cholinergic neurons in brain areas associated with learning and memory.

The 5-HT₆R antagonists have been shown to increase extracellular glutamate levels in addition to acetylcholine. It is thus possible that the cognitive effects of 5-HT₆R antagonists and other treatments for AD could be the result of interactions with both cholinergic and glutamatergic systems involved in learning and memory.

Since the cause and development of the dementia depend on the different mechanisms, it may be advantageous to use the combination of drugs working through different mechanisms for the treatment of AD. The current approved treatment for AD

includes the combination of acetylcholinesterase inhibitor, donepezil and NMDA receptor antagonist, memantine. However, there remains the need for the new drugs/combination to treat the patients with AD.

5 The compounds of the present invention are pure 5-HT₆R antagonists with high affinity and very high selectivity over closely related serotonin receptor subtypes and improves learning and memory in animals. The 5-HT₆R antagonist compounds mentioned here are described in US7875605. The preparation of these compounds is given in the said patent.

10 As the treatment of AD is chronic in nature, there is a desperate unmet medical need for better and safer treatment options. A therapeutic strategy eagerly sought for AD patients is to target an improvement with an adjunct to existing therapies that would bring additional relief for patients, lower the burden on the caregiver and allow the patient to enjoy a better quality of life without the need for institutional care and/or hospitalization.

15 The instant invention provides pure 5-HT₆R antagonists or the pharmaceutically acceptable salt(s) thereof, which enhances the cognitive function of patients on treatment in combination with AChEIs and NMDA receptor antagonist. The present invention is based on the unusual finding that the combination of compounds with pure 5-HT₆R antagonistic activity, the compounds which act as AChEIs (for example donepezil) and the compounds which act as NMDA receptor antagonists (for example memantine), demonstrate
20 synergistic effect in their pharmacological activity. Memantine acts by blocking the glutamatergic neurotransmissions in the brain. The 5-HT₆R antagonists have been shown to increase extracellular glutamate levels in addition to acetylcholine. Hence it is not anticipated that the combination of a pure 5-HT₆R antagonist + donepezil + memantine would result in synergistic procognitive effects. However surprisingly, the combination of
25 pure 5-HT₆R antagonists + AChEIs + NMDA receptor antagonist (triple combination) showed synergistic effects in animal models, and also increased the levels of acetylcholine, a neurotransmitter that plays a vital role in cognitive improvement. Based on these results one can infer that such combined administration and/or co-treatment of pure 5-HT₆R antagonists + AChEIs + NMDA receptor antagonist, may result in beneficial effect to
30 improve the therapeutic efficacy in humans. Further the pure 5-HT₆R antagonists or the pharmaceutically acceptable salt(s) thereof of the instant invention enhances the effect of the AChEIs and NMDA receptor antagonist in the treatment of cognitive disorders.

SUMMARY OF THE INVENTION

The object of the present invention is to provide an improved combination therapy for the treatment of cognitive disorders, such as Alzheimer's disease, schizophrenia, Parkinson's disease, lewy body dementia, vascular dementia or frontotemporal dementia.

In the first aspect, the present invention relates to a combination of pure 5-HT₆ receptor antagonist, acetylcholinesterase inhibitor and NMDA receptor antagonist.

In yet another aspect, the present invention relates to a combination of pure 5-HT₆ receptor antagonist, acetylcholinesterase inhibitor and NMDA receptor antagonist; wherein the pure 5-HT₆ receptor antagonist is selected from:

1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl) methyl]-1H-indole;

1-[(4-Fluorophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl) methyl]-1H-indole; and

1-[(4-Isopropylphenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl) methyl]-1H-indole; or a pharmaceutically acceptable salt thereof.

In another aspect, the present invention relates to a combination of pure 5-HT₆ receptor antagonist, acetylcholinesterase inhibitor and NMDA receptor antagonist; wherein the pure 5-HT₆ receptor antagonist is 1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole or a pharmaceutically acceptable salt thereof.

In another aspect, the present invention relates to a combination of pure 5-HT₆ receptor antagonist, acetylcholinesterase inhibitor and NMDA receptor antagonist; wherein the pure 5-HT₆ receptor antagonist is 1-[(4-Fluorophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole or a pharmaceutically acceptable salt thereof.

In another aspect, the present invention relates to a combination of pure 5-HT₆ receptor antagonist, acetylcholinesterase inhibitor and NMDA receptor antagonist; wherein the pure 5-HT₆ receptor antagonist is 1-[(4-Isopropylphenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole or a pharmaceutically acceptable salt thereof.

In another aspect, the present invention relates to a combination of pure 5-HT₆ receptor antagonist, acetylcholinesterase inhibitor and NMDA receptor antagonist; wherein the acetylcholinesterase inhibitor is selected from donepezil, galantamine and rivastigmine or a pharmaceutically acceptable salt thereof.

In another aspect, the present invention relates to a combination of pure 5-HT₆ receptor antagonist, acetylcholinesterase inhibitor and NMDA receptor antagonist; wherein the NMDA receptor antagonist is memantine or a pharmaceutically acceptable salt thereof.

In yet another aspect the present invention relates to a combination of 1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole, donepezil and memantine or a pharmaceutically acceptable salt thereof.

In yet another aspect, the present invention relates to the said combination for use in the treatment of cognitive disorders such as Alzheimer's disease, schizophrenia, Parkinson's disease, lewy body dementia, vascular dementia or frontotemporal dementia.

In yet another aspect, the present invention relates to a method of treatment of cognitive disorders such as Alzheimer's disease, schizophrenia, Parkinson's disease, lewy body dementia, vascular dementia or frontotemporal dementia comprising administering to a patient in need thereof a therapeutically effective amount of the said combination.

In yet another aspect, the present invention relates to pure 5-HT₆ receptor antagonist for use in the adjunct treatment of cognitive disorders such as Alzheimer's disease, schizophrenia, Parkinson's disease, lewy body dementia, vascular dementia or frontotemporal dementia in patients on treatment with acetylcholinesterase inhibitor and NMDA receptor antagonist.

In yet another aspect, the present invention relates to the compound, 1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole or a pharmaceutically acceptable salt thereof for use in the adjunct treatment of cognitive disorders such as Alzheimer's disease, schizophrenia, Parkinson's disease, lewy body dementia, vascular dementia or frontotemporal dementia in patients on treatment with donepezil and memantine.

In yet another aspect, the present invention relates to the compound, 1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole or a pharmaceutically acceptable salt thereof for use in combination with or adjunct to an acetylcholinesterase inhibitor and NMDA receptor antagonist for the treatment of cognitive disorders such as Alzheimer's disease, schizophrenia, Parkinson's disease, lewy body dementia, vascular dementia or frontotemporal dementia.

In another aspect, the present invention relates to a method for treatment of cognitive disorders comprising administering to a patient in need thereof a therapeutically

effective amount of 1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole or a pharmaceutically acceptable salt thereof in combination with or as an adjunct to donepezil or a pharmaceutically acceptable salt thereof and memantine or a pharmaceutically acceptable salt thereof.

5 In yet another aspect, the present invention relates to use of a combination of pure 5-HT₆ receptor antagonist, acetylcholinesterase inhibitor and NMDA receptor antagonist for the treatment of cognitive disorders such as Alzheimer's disease, schizophrenia, Parkinson's disease, lewy body dementia, vascular dementia or frontotemporal dementia.

10 In yet another aspect, the present invention relates to use of a combination of 1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole, donepezil and memantine or a pharmaceutically acceptable salt thereof for the treatment of cognitive disorders such as Alzheimer's disease, schizophrenia, Parkinson's disease, lewy body dementia, vascular dementia or frontotemporal dementia.

15 In another aspect, the present invention relates to pharmaceutical composition comprising the 5-HT₆ receptor antagonist, acetylcholinesterase inhibitor and NMDA receptor antagonist and pharmaceutically acceptable excipients or combination thereof.

20 In another aspect, the present invention relates to pharmaceutical composition comprising 1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole, donepezil and memantine or a pharmaceutically acceptable salt thereof and the pharmaceutically acceptable excipients or combination thereof.

25 In another aspect, the present invention relates to pharmaceutical composition comprising the 5-HT₆ receptor antagonist, acetylcholinesterase inhibitor and NMDA receptor antagonist or the pharmaceutically acceptable salt thereof along with the pharmaceutically acceptable excipients or combination thereof for use in the treatment of cognitive disorders such as Alzheimer's disease, schizophrenia, Parkinson's disease, lewy body dementia, vascular dementia or frontotemporal dementia.

30 In another aspect, there is provided a combination comprising pure 5-HT₆ receptor antagonist, acetylcholinesterase inhibitor and NMDA receptor antagonist, wherein the pure 5-HT₆ receptor antagonist is selected from the group consisting of: 1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole; 1-[(4-Fluorophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole; and 1-[(4-Isopropylphenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole; or a pharmaceutically acceptable salt thereof.

In another aspect, there is provided a combination comprising pure 5-HT₆ receptor antagonist, acetylcholinesterase inhibitor and NMDA receptor antagonist, wherein the pure 5-HT₆ receptor antagonist is selected from the group consisting of: 1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole; 1-[(4-Fluorophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole; 1-[(4-Isopropylphenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole; and a pharmaceutically acceptable salt thereof.

In another aspect, there is provided a combination comprising 1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole dimesylate monohydrate, donepezil hydrochloride and memantine hydrochloride.

In yet another aspect, there is provided a compound, 1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole or a pharmaceutically acceptable salt thereof for use in combination with acetylcholinesterase inhibitor and NMDA receptor antagonist for the treatment of Alzheimer's disease in a patient.

In yet another aspect, there is provided a compound, 1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole or a pharmaceutically acceptable salt thereof for use as an adjunct treatment of Alzheimer's disease in a patient on stable treatment with acetylcholinesterase inhibitor and NMDA receptor antagonist.

In still another aspect, there is provided a use of a therapeutically effective amount of 1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole or a pharmaceutically acceptable salt thereof for administration to treat Alzheimer's disease in a patient on stable treatment with acetylcholinesterase inhibitor and NMDA receptor antagonist.

BRIEF DESCRIPTION OF THE DIAGRAMS

Figure 1a depicts the results of the effect of a co-treatment of compound 1 with donepezil and memantine on cognition enhancing properties using object recognition task model.

Figure 1b depicts the results of the effect of a co-treatment of compound 2 with donepezil and memantine on cognition enhancing properties using object recognition task model.

Figure 1c depicts the results of the effect of a co-treatment of compound 3 with donepezil and memantine on cognition enhancing properties using object recognition task model.

Figure 2 depicts the effect of compound 1 alone and in combination with donepezil and memantine on extracellular levels of acetylcholine in ventral hippocampus of male Wistar rats.

Figure 3 depicts the effect of compound 2, donepezil and memantine combination on extracellular levels of acetylcholine in ventral hippocampus of male Wistar rats.

Figure 4 depicts the effect of compound 3, donepezil and memantine combination on extracellular levels of acetylcholine in ventral hippocampus of male Wistar rats.

Figure 5 depicts the effect of compound 1, donepezil and memantine combination on evoked theta modulation in dorsal hippocampus of anesthetized male Wistar rats.

DETAILED DESCRIPTION

Unless otherwise stated, the following terms used in the specification and claims have the meanings given below:

The term, “5-HT₆ receptor antagonist” as used herein refers to a ligand or drug that has affinity towards 5-HT₆ receptor, blocks or inhibits the function / binding of agonist at the 5-HT₆ receptor.

The term, “pure 5-HT₆ receptor antagonist” as used herein refers to 5-HT₆ receptor antagonist which has very high selectivity (>250 fold) over closely related serotonin subtypes like 5-HT_{1A}, 5-HT_{1B}, 5-HT_{1D}, 5-HT_{2A}, 5-HT_{2C}, 5-HT₄, 5-HT_{5A} and 5-HT₇.

Examples of the pure 5-HT₆ receptor antagonists include,

1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole;

1-[(4-Fluorophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole;

and

1-[(4-Isopropylphenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole;

or a pharmaceutically acceptable salt thereof.

Examples of pharmaceutically acceptable salt of the above identified compounds include but not limited to,

1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole dimesylate monohydrate;

1-[(4-Fluorophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole dihydrochloride; and

1-[(4-Isopropylphenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole dihydrochloride.

The term, “acetylcholinesterase inhibitor” as used herein is a chemical or drug that inhibits the acetylcholinesterase enzyme from breaking down acetylcholine, thereby increasing both the level and duration of action of the neurotransmitter acetylcholine. Examples of acetylcholinesterase inhibitor are donepezil, rivastigmine and galantamine. Preferably, the acetylcholinesterase inhibitor is donepezil and rivastigmine. More preferably the acetylcholinesterase inhibitor is donepezil.

Donepezil is a drug approved for treatment of mild, moderate and severe dementia of Alzheimer’s disease. Donepezil is a reversible inhibitor of the enzyme acetylcholinesterase and sold under trade name Aricept[®] as hydrochloride salt.

Rivastigmine is a drug approved for treatment of mild, moderate and severe dementia of Alzheimer’s disease. Rivastigmine is a reversible cholinesterase inhibitor and sold under trade name Exelon[®] and Exelon Patch[®] as tartrate salt.

Galantamine is a drug approved for treatment of mild, moderate and severe dementia of Alzheimer’s disease. Galantamine, a reversible, competitive acetylcholinesterase inhibitor and sold under trade name Razadyne[®] as hydrobromide salt.

The term, “NMDA receptor antagonist” as used herein refers to class of compounds which act on glutamatergic system by inhibiting the NMDA receptor. Example of NMDA receptor antagonist is memantine. Memantine is a drug approved for treatment of moderate to severe dementia of the Alzheimer’s disease. Memantine is NMDA receptor antagonist and sold under trade name Namenda[®] and Namenda XR[®] as hydrochloride salt.

The combination of memantine and donepezil is approved for the treatment of moderate to severe dementia of the Alzheimer's disease and sold under trade name Namzaric[®] as memantine hydrochloride salt and donepezil hydrochloride salt.


The phrase, "therapeutically effective amount" is defined as an amount of a compound of the present invention that (i) treats the particular disease, condition or disorder, (ii) eliminates one or more symptoms of the particular disease, condition or disorder and (iii) delays the onset of one or more symptoms of the particular disease, condition or disorder described herein.

The term, "pharmaceutically acceptable salt" as used herein refers to salts of the active compound and are prepared by reaction with the appropriate organic or inorganic acid or acid derivative, depending on the particular substituents found on the compounds described herein.

The term, "patient" as used herein refers to an animal. Preferably the term "patient" refers to mammal. The term mammal includes animals such as mice, rats, dogs, rabbits, pigs, monkeys, horses and human. More preferably the patient is human.


The term, "Alzheimer's disease" as used herein refers to a dementia that causes problems with memory, thinking and behavior. The Alzheimer's disease can be mild to severe.

The compound 1 as used herein is 1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl) methyl]-1H-indole dimesylate monohydrate which has the chemical structure,




and the process for preparing this compound on a larger scale is described in WO2015083179A1.

The compound 2 as used herein is 1-[(4-Fluorophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole dihydrochloride which has the chemical structure,



The compound 3 as used herein is 1-[(4-Isopropylphenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole dihydrochloride which has the chemical structure,



The term, "treatment" or "treating" as used herein refers to any treatment of a disease in a mammal, including: (a) slowing or arresting the development of clinical symptoms; and/or (b) causing the regression of clinical symptoms.

The term, "compound for use" as used herein embrace any one or more of the following: (1) use of a compound, (2) method of use of a compound, (3) use in the treatment of, (4) the use for the manufacture of pharmaceutical composition / medicament for treatment / treating or (5) method of treatment / treating / preventing / reducing / inhibiting comprising administering an effective amount of the active compound to a subject in need thereof.

The term, "cognitive disorder" as used herein refers to a group of mental health disorders that principally affect learning, memory, perception, and problem solving, and include amnesia, dementia, and delirium. Cognitive disorders can result due to disease, disorder, ailment or toxicity. Example of cognitive disorders includes but not limited to, Alzheimer's disease, schizophrenia, Parkinson's disease, lewy body dementia (LBD), vascular dementia or frontotemporal dementia (FTD). Preferably, the cognitive disorder is Alzheimer's disease.

The term, "adjunct" or "adjunctive treatment" as used herein refers to an additional treatment to a patient who has already received at least one other therapy for cognitive disorders. A drug used as adjunctive therapy is administered to a patient to make that primary treatment works better.

Embodiments

The present invention encompasses all the combinations described herein without limitation, however, preferred aspects and elements of the invention are discussed herein in the form of the following embodiments.

In one embodiment, the present invention relates to the combination of pure 5-HT₆ receptor antagonist, acetylcholinesterase inhibitor and NMDA receptor antagonist; wherein the pure 5-HT₆ receptor antagonist is 1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole dimesylate monohydrate.

In another embodiment, the present invention relates to the combination of pure 5-HT₆ receptor antagonist, acetylcholinesterase inhibitor and NMDA receptor antagonist; wherein the pure 5-HT₆ receptor antagonist is 1-[(4-Fluorophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole dihydrochloride.

In another embodiment, the present invention relates to the combination of pure 5-HT₆ receptor antagonist, acetylcholinesterase inhibitor and NMDA receptor antagonist; wherein the pure 5-HT₆ receptor antagonist is 1-[(4-Isopropylphenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole dihydrochloride.

In another embodiment, the present invention relates to the combination of 1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole, donepezil and memantine or the pharmaceutically acceptable salt thereof.

In another embodiment, the present invention relates to the combination of 1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole, rivastigmine and memantine or the pharmaceutically acceptable salt thereof.

In another embodiment, the present invention relates to the combination of 1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole, galantamine and memantine or the pharmaceutically acceptable salt thereof.

In another embodiment, the present invention relates to the combination of 1-[(4-Fluorophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole, donepezil and memantine or the pharmaceutically acceptable salt thereof.

In another embodiment, the present invention relates to the combination of 1-[(4-Fluorophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole, rivastigmine and memantine or the pharmaceutically acceptable salt thereof.

In another embodiment, the present invention relates to the combination of 1-[(4-Fluorophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole, galantamine and memantine or the pharmaceutically acceptable salt thereof.

In another embodiment, the present invention relates to the combination of 1-[(4-Isopropylphenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole, donepezil and memantine or the pharmaceutically acceptable salt thereof.

In another embodiment, the present invention relates to the combination of 1-[(4-Isopropylphenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole, rivastigmine and memantine or the pharmaceutically acceptable salt thereof.

In another embodiment, the present invention relates to the combination of 1-[(4-Isopropylphenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole, galantamine and memantine or the pharmaceutically acceptable salt thereof.

In another embodiment, the present invention relates to the combination of 1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole dimesylate monohydrate, donepezil hydrochloride and memantine hydrochloride.

In another embodiment, the present invention relates to the combination of 1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole dimesylate monohydrate, rivastigmine tartrate and memantine hydrochloride.

In another embodiment, the present invention relates to the combination of 1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole dimesylate monohydrate, galantamine hydrobromide and memantine hydrochloride.

In another embodiment, the present invention relates to the combination of 1-[(4-Fluorophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole dihydrochloride, donepezil hydrochloride and memantine hydrochloride.

In another embodiment, the present invention relates to the combination of 1-[(4-Fluorophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole dihydrochloride, rivastigmine tartrate and memantine hydrochloride.

In another embodiment, the present invention relates to the combination of 1-[(4-Fluorophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole dihydrochloride, galantamine hydrobromide and memantine hydrochloride.

In another embodiment, the present invention relates to the combination of 1-[(4-Isopropylphenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole dihydrochloride, donepezil hydrochloride and memantine hydrochloride.

In another embodiment, the present invention relates to the combination of 1-[(4-Isopropylphenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole dihydrochloride, rivastigmine tartrate and memantine hydrochloride.

In another embodiment, the present invention relates to the combination of 1-[(4-Isopropylphenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole dihydrochloride, galantamine hydrobromide and memantine hydrochloride.

In another embodiment, the present invention provides the combination of pure 5-HT₆ receptor antagonist, acetylcholinesterase inhibitor and NMDA receptor antagonist which is more effective than the combination of pure 5-HT₆ receptor antagonist and acetylcholinesterase inhibitor, acetylcholinesterase inhibitor and NMDA receptor antagonist or pure 5-HT₆ receptor antagonist and NMDA receptor antagonist.

In another embodiment, the present invention provides the combination of the pure 5-HT₆ receptor antagonist, acetylcholinesterase inhibitor and NMDA receptor antagonist which is more effective than the pure 5-HT₆ receptor antagonist, acetylcholinesterase inhibitor and NMDA receptor antagonist alone.

In another embodiment, the present invention provides the combination of 1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole dimesylate monohydrate, donepezil hydrochloride and memantine hydrochloride which is more effective than the combination of 1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole dimesylate monohydrate and donepezil hydrochloride, donepezil hydrochloride and memantine hydrochloride or 1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole dimesylate monohydrate and memantine hydrochloride.

In another embodiment, the present invention provides the combination of 1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole dimesylate monohydrate, donepezil hydrochloride and memantine hydrochloride which is more effective than 1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole dimesylate monohydrate, donepezil hydrochloride and memantine hydrochloride alone.

In another embodiment the pharmaceutically acceptable salt of pure 5-HT₆ receptor antagonist includes but not limited to dimesylate monohydrate salt, dihydrochloride salt, oxalate salt, tartrate salt and the like. Preferably, the pharmaceutically acceptable salt is dimesylate monohydrate salt and dihydrochloride salt. More preferably, the pharmaceutically acceptable salt is dimesylate monohydrate salt.

In another embodiment, the present invention relates to a method of treating Alzheimer's disease comprising administering to a patient in need thereof a therapeutically effective amount of the said combination.

In another embodiment, the present invention relates to a method of treating Alzheimer's disease comprising administering to a patient in need thereof a therapeutically effective amount of 1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole or a pharmaceutically acceptable salt thereof, acetylcholinesterase inhibitor and NMDA receptor antagonist.

In another embodiment, the present invention relates to a method of treating Alzheimer's disease comprising administering to a patient in need thereof a therapeutically effective amount of 1-[(4-Fluorophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole or a pharmaceutically acceptable salt thereof, acetylcholinesterase inhibitor and NMDA receptor antagonist.

In another embodiment, the present invention relates to a method of treating Alzheimer's disease comprising administering to a patient in need thereof a therapeutically effective amount of 1-[(4-Isopropylphenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole or a pharmaceutically acceptable salt thereof, acetylcholinesterase inhibitor and NMDA receptor antagonist.

In another embodiment, the present invention relates to a method of treating Alzheimer's disease comprising administering to a patient in need thereof a therapeutically effective amount of 1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole or a pharmaceutically acceptable salt thereof in combination with acetylcholinesterase inhibitor and NMDA receptor antagonist.

In another embodiment, the present invention relates to a method of treating Alzheimer's disease comprising administering to a patient in need thereof a therapeutically effective amount of 1-[(4-Fluorophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-

piperazinyl)methyl]-1H-indole or a pharmaceutically acceptable salt thereof in combination with acetylcholinesterase inhibitor and NMDA receptor antagonist.

In another embodiment, the present invention relates to a method of treating Alzheimer's disease comprising administering to a patient in need thereof a therapeutically effective amount of 1-[(4-Isopropylphenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole or a pharmaceutically acceptable salt thereof in combination with acetylcholinesterase inhibitor and NMDA receptor antagonist.

In another embodiment, the present invention relates to a method of treating Alzheimer's disease comprising administering to a patient in need thereof a therapeutically effective amount of 1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole dimesylate monohydrate in combination with acetylcholinesterase inhibitor and NMDA receptor antagonist.

In another embodiment, the present invention relates to a method of treating Alzheimer's disease comprising administering to a patient in need thereof a therapeutically effective amount of 1-[(4-Fluorophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole dihydrochloride in combination with acetylcholinesterase inhibitor and NMDA receptor antagonist.

In another embodiment, the present invention relates to a method of treating Alzheimer's disease comprising administering to a patient in need thereof a therapeutically effective amount of 1-[(4-Isopropylphenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole dihydrochloride in combination with acetylcholinesterase inhibitor and NMDA receptor antagonist.

In another embodiment, the present invention relates to a method of treating Alzheimer's disease comprising administering to a patient in need thereof a therapeutically effective amount of 1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole dimesylate monohydrate in combination with donepezil or a pharmaceutically acceptable salt thereof and memantine or a pharmaceutically acceptable salt thereof.

In another embodiment, the present invention relates to a method of treating Alzheimer's disease comprising administering to a patient in need thereof a therapeutically effective amount of 1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-

piperazinyl)methyl]-1H-indole dimesylate monohydrate in combination with donepezil hydrochloride and memantine hydrochloride.

In another embodiment, the present invention relates to the combination of 1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole dimesylate monohydrate, acetylcholinesterase inhibitor and NMDA receptor antagonist for use in the treatment of Alzheimer's disease.

In yet another aspect, the present invention relates to 1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole or a pharmaceutically acceptable salt thereof for use in the adjunct treatment of Alzheimer's disease in a patient on treatment with acetylcholinesterase inhibitor and NMDA receptor antagonist.

In yet another aspect, the present invention relates to 1-[(4-Fluorophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole or a pharmaceutically acceptable salt thereof for use in the adjunct treatment of Alzheimer's disease in a patient on treatment with acetylcholinesterase inhibitor and NMDA receptor antagonist.

In yet another aspect, the present invention relates to 1-[(4-Isopropylphenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole or a pharmaceutically acceptable salt thereof for use in the adjunct treatment of Alzheimer's disease in a patient on treatment with acetylcholinesterase inhibitor and NMDA receptor antagonist.

In another embodiment, the present invention relates to 1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole dimesylate monohydrate for use in the adjunct treatment of Alzheimer's disease in a patient on treatment with donepezil and memantine or a pharmaceutically acceptable salt thereof.

In another embodiment, the present invention relates to 1-[(4-Fluorophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole dihydrochloride for use in the adjunct treatment of Alzheimer's disease in a patient on treatment with donepezil and memantine or a pharmaceutically acceptable salt thereof.

In another embodiment, the present invention relates to 1-[(4-Isopropylphenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole dihydrochloride for use in the adjunct treatment of Alzheimer's disease in a patient on treatment with donepezil and memantine or a pharmaceutically acceptable salt thereof.

In another embodiment, the present invention relates to use of the combination of pure 5-HT₆ receptor antagonist, acetylcholinesterase inhibitor and NMDA receptor antagonist in the manufacture of a medicament for treatment of Alzheimer's disease.

In another embodiment, the present invention relates to use of pure 5-HT₆ receptor antagonist in the manufacture of a medicament for treatment of Alzheimer's disease in combination with acetylcholinesterase inhibitor and NMDA receptor antagonist.

In another embodiment, the present invention relates to use of pure 5-HT₆ receptor antagonist in the manufacture of a medicament for treatment of Alzheimer's disease as adjunct to acetylcholinesterase inhibitor and NMDA receptor antagonist.

In another embodiment, the present invention relates to use of the 1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole or a pharmaceutically acceptable salt thereof in the manufacture of a medicament for treatment of Alzheimer's disease in combination with donepezil or a pharmaceutically acceptable salt thereof and memantine or a pharmaceutically acceptable salt thereof.

In another embodiment, the present invention relates to use of the 1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole dimesylate monohydrate in the manufacture of a medicament for treatment of Alzheimer's disease in combination with donepezil or a pharmaceutically acceptable salt thereof and memantine or a pharmaceutically acceptable salt thereof.

In another embodiment, the present invention relates to use of the 1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole dimesylate monohydrate in the manufacture of a medicament for treatment of Alzheimer's disease in combination with donepezil hydrochloride and memantine hydrochloride.

In another embodiment, the present invention relates to pure 5-HT₆ receptor antagonist for use in the treatment of Alzheimer's disease in combination with Namzaric[®].

In another embodiment, the present invention relates to a method of treating Alzheimer's disease comprising administering to a patient in need thereof a therapeutically effective amount of the pure 5-HT₆ receptor antagonist in combination with Namzaric[®].

In another embodiment, the present invention relates to the combination for treatment of Alzheimer's disease, wherein Alzheimer's disease is mild Alzheimer's disease.

In another embodiment, the present invention relates to the combination for treatment of Alzheimer's disease, wherein the Alzheimer's disease is moderate Alzheimer's disease.

In another embodiment, the present invention relates to the combination for treatment of Alzheimer's disease, wherein the Alzheimer's disease is severe Alzheimer's disease.

In another embodiment, the present invention relates to the combination wherein the active ingredients can be administered to a patient concurrently or separately.

In yet another aspect, the active ingredients of the combination of the present invention are normally administered by formulating the active ingredients into a pharmaceutical composition in accordance with standard pharmaceutical practice.

In yet another aspect, the active ingredients of the combination of the present invention may be administered by oral, nasal, local, dermal or parenteral routes.

In yet another aspect, the active ingredients of the combination of the present invention can be administered by the same or different route of administration. For instance, the 5-HT₆ receptor antagonist of the instant invention can be administered orally, the acetylcholinesterase inhibitor can be administered transdermally and the NMDA receptor antagonist can be administered locally.

The pharmaceutical compositions of the present invention may be formulated in a conventional manner using one or more pharmaceutically acceptable excipients. The pharmaceutically acceptable excipients are diluents, disintegrants, binders, lubricants, glidants, polymers, coating agents, solvents, cosolvents, preservatives, wetting agents, thickening agents, antifoaming agents, sweetening agents, flavouring agents, antioxidants, colorants, solubilizers, plasticizer, dispersing agents and the like. Excipients are selected from microcrystalline cellulose, mannitol, lactose, pregelatinized starch, sodium starch glycolate, corn starch or derivatives thereof, povidone, crospovidone, calcium stearate, glyceryl monostearate, glyceryl palmitostearate, talc, colloidal silicone dioxide, magnesium stearate, sodium lauryl sulfate, sodium stearyl fumarate, zinc stearate, stearic acid or hydrogenated vegetable oil, gum arabica, magnesia, glucose, fats, waxes, natural or hardened oils, water, physiological sodium chloride solution or alcohols, for example, ethanol, propanol or glycerol, sugar solutions, such as glucose solutions or mannitol solutions and the like or a mixture of the various excipients.

In yet another aspect, the active compounds of the invention may be formulated in the form of pills, tablets, coated tablets, capsules, powder, granules, pellets, patches, implants, films, liquids, semi-solids, gels, aerosols, emulsions, elixirs and the like. Such pharmaceutical compositions and processes for preparing same are well known in the art.

In yet another aspect, the pharmaceutical composition of the instant invention contains 1 to 90 %, 5 to 75 % and 10 to 60 % by weight of the compounds of the instant invention or pharmaceutically acceptable salt thereof. The amount of the active compounds or its pharmaceutically acceptable salt in the pharmaceutical composition(s) can range from about 1 mg to about 500 mg or from about 5 mg to about 400 mg or from about 5 mg to about 250 mg or from about 7 mg to about 150 mg or in any range falling within the broader range of 1 mg to 500 mg.

In yet another aspect, the pharmaceutical composition of the combination of the instant invention can be conventional formulations such as immediate release formulations, modified release formulations such as sustained release formulations, delayed release formulations and extended release formulations or new delivery systems such as oral disintegrating formulations and transdermal patches.

The dose of the active compounds can vary depending on factors such as age and weight of patient, nature, route of administration and severity of the disease to be treated and such other factors. Therefore, any reference regarding pharmacologically effective amount of the compounds 1, 2 and 3 refers to the aforementioned factors.

In yet another aspect, the 5-HT₆ receptor antagonist can be co-administered with acetylcholinesterase inhibitor and NMDA receptor antagonist at a daily dose of 1 mg to 300 mg; such as 1, 5, 10, 20, 25, 30, 50, 75, 100, 150, 200 or 300 mg, preferably at a daily dose of 10, 25, 30, 50, 75, 100, 125 or 150 mg and most preferably at a daily dose of 10, 25, 50, 75, 100 or 125 mg.

In yet another aspect, the acetylcholinesterase inhibitor can be co-administered with 5-HT₆ receptor antagonist and NMDA receptor antagonist at a daily dose of 1 mg to 30 mg; such as 1, 1.5, 2, 3, 4, 4.5, 5, 6, 8, 9.5, 10, 12, 13, 13.3, 15, 16, 23, 24, 25 or 30 mg, preferably at a daily dose of 1, 1.5, 2, 3, 4, 4.5, 5, 6, 8, 9.5, 10, 12, 13, 13.3, 16, 23, 24, or 25 mg and most preferably at a daily dose of 1.5, 3, 4, 4.5, 5, 6, 8, 9.5, 10, 12, 13.3, 16, 23 or 24 mg.

In yet another aspect, the NMDA receptor antagonist, memantine can be co-administered with 5-HT₆ receptor antagonist and acetylcholinesterase inhibitor at a daily dose of 1 mg to 40 mg; such as 5, 10, 14, 20, 28 or 40 mg, preferably at a daily dose of 5, 10, 14, 20 or 28 mg and most preferably at a daily dose of 5, 7, 10, 14, 20 or 28 mg.

In yet another aspect, the acetylcholinesterase inhibitor, donepezil can be co-administered with 5-HT₆ receptor antagonist and NMDA receptor antagonist at a daily dose of 2 mg to 30 mg; such as 2, 5, 10, 15, 23, 25 or 30 mg, preferably at a daily dose of 2, 5, 10, 23 or 25 mg and most preferably at a daily dose of 5, 10 or 23 mg.

In yet another aspect, the acetylcholinesterase inhibitor, rivastigmine can be co-administered with 5-HT₆ receptor antagonist and NMDA receptor antagonist at a daily dose of 0.5 mg to 15 mg; such as 1, 1.5, 3, 4.5, 5, 6, 9.5, 10 or 13.3 mg, preferably at a daily dose of 1, 1.5, 3, 4.5, 5, 6, 9.5 or 13.3 mg and most preferably at a daily dose of 1.5, 3, 4.5, 6, 9.5 and 13.3 mg.

In yet another aspect, the acetylcholinesterase inhibitor, galantamine can be co-administered with 5-HT₆ receptor antagonist and NMDA receptor antagonist at a daily dose of 1 mg to 30 mg; such as 1, 2, 4, 6, 8, 12, 16, 24 and 30 mg, preferably at a daily dose of 2, 4, 6, 8, 12, 16 and 24 mg and most preferably at a daily dose of 4, 8, 12, 16 and 24 mg.

In yet another aspect, the treatment comprises administering to the patient 1 mg to 200 mg of 1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole or a pharmaceutically acceptable salt thereof, per day.

In yet another aspect, the treatment comprises administering to the patient 1 mg to 10 mg of 1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole or a pharmaceutically acceptable salt thereof, per day.

In yet another aspect, the treatment comprises administering to the patient 25 mg to 125 mg of 1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole or a pharmaceutically acceptable salt thereof, per day.

In yet another aspect, the treatment comprises administering to the patient 150 mg to 200 mg of 1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole or a pharmaceutically acceptable salt thereof, per day.

In yet another aspect, the treatment comprises administering to the patient 10 mg to 100 mg of 1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole or a pharmaceutically acceptable salt thereof, per day.

In yet another aspect, the treatment comprises administering to the patient 10 mg to 50 mg of 1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole or a pharmaceutically acceptable salt thereof, per day.

In yet another aspect, the treatment comprises administering to the patient 25 mg to 50 mg of 1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole or a pharmaceutically acceptable salt thereof, per day.

In yet another aspect, the treatment comprises administering to the patient 75 mg to 100 mg of 1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole or a pharmaceutically acceptable salt thereof, per day.

In yet another aspect, the treatment comprises administering to the patient 1 mg to 25 mg of donepezil or a pharmaceutically acceptable salt thereof, per day.

In yet another aspect, the treatment comprises administering to the patient 5 mg to 25 mg of donepezil or a pharmaceutically acceptable salt thereof, per day.

In yet another aspect, the treatment comprises administering to the patient, 5, 10 or 23 mg of donepezil or a pharmaceutically acceptable salt thereof, per day.

In yet another aspect, the treatment comprises administering to the patient 1 mg to 40 mg of memantine or a pharmaceutically acceptable salt thereof, per day.

In yet another aspect, the treatment comprises administering to the patient 5 mg to 30 mg of memantine or a pharmaceutically acceptable salt thereof, per day.

In yet another aspect, the treatment comprises administering to the patient 5, 10, 14, 20 or 28 mg of memantine or a pharmaceutically acceptable salt thereof, per day.

In yet another aspect, the treatment comprises administering the active compounds to the patient one to three times per day, one to three times per week or one to three times per month. Preferably, the treatment comprises administering the compound to a patient once a day, twice a day, or thrice a day. More preferably, the treatment comprises administering the compound to a patient once a day.

Examples

The examples given below are provided by the way of illustration only and therefore should not be construed to limit the scope of the invention.

Abbreviations:

5-HT _{1A}	:	5-Hydroxytryptamine 1A receptor
5-HT _{1B}	:	5-Hydroxytryptamine 1B receptor
5-HT _{1D}	:	5-Hydroxytryptamine 1D receptor
5-HT _{2A}	:	5-Hydroxytryptamine 2A receptor
5-HT _{2C}	:	5-Hydroxytryptamine 2C receptor
5-HT ₄	:	5-Hydroxytryptamine 4 receptor
5-HT _{5A}	:	5-Hydroxytryptamine 5A receptor
5-HT ₆	:	5-Hydroxytryptamine 6 receptor
5-HT ₇	:	5-Hydroxytryptamine 7 receptor
ANOVA	:	Analysis of variance
AP	:	Anterior Posterior
aCSF	:	Artificial Cerebrospinal fluid
cAMP	:	Cyclic adenosine monophosphate
CaCl ₂ . 2H ₂ O	:	Calcium Chloride dehydrate
DV	:	Dorsal Ventral
EC ₅₀	:	Half maximal effective concentration
EDTA	:	Ethylenediaminetetraacetic acid
EEG	:	Electroencephalogram
GPCR	:	G-Protein Coupled Receptor
HCl	:	Hydrochloric acid
h	:	Hour (s)
<i>i.p.</i>	:	Intraperitoneal
<i>i.v.</i>	:	Intravenous
KCl	:	Potassium chloride
K _b	:	Binding constant
K _i	:	Inhibitory constant
LC-MS/MS	:	Liquid chromatography-Mass spectrometry/ Mass spectrometry
mg	:	Milligram

MgCl ₂	:	Magnesium chloride
min	:	Minute (s)
ML	:	Medial Lateral
mM	:	Millimolar
nmol/L	:	Nanomoles per litre
NaCl	:	Sodium chloride
NaH ₂ PO ₄ .2H ₂ O	:	Sodium dihydrogen phosphate dihydrate
Na ₂ HPO ₄ .7H ₂ O	:	Sodium monohydrogen phosphate heptahydrate
NPO	:	Nucleus Pontis Oralis
nM	:	Nanomolar
<i>p.o.</i>	:	Per oral
<i>s.c.</i>	:	Subcutaneous
S.E.M.	:	Standard error of the mean
μM	:	Micromolar
θ	:	Theta

Example 1:

Determination of K_b values at 5-HT₆ receptor:

A stable CHO cell line expressing recombinant human 5-HT₆ receptor and pCRE-Luc reporter system was used for cell-based assay. The assay offers a non-radioactive based approach to determine binding of a compound to GPCRs. In this specific assay, the level of intracellular cAMP which is modulated by activation or inhibition of the receptor is measured. The recombinant cells harbor luciferase reporter gene under the control of cAMP response element.

The above cells were grown in 96 well clear bottom white plates in Hams F12 medium containing 10 % fetal bovine serum (FBS). Prior to the addition of compounds or standard agonist, cells were serum starved overnight. Increasing concentrations of test compound were added along with 10 μM of serotonin in OptiMEM medium to the cells. The incubation was continued at 37°C in CO₂ incubator for 4 hours. Medium was removed and cells were washed with phosphate buffered saline. The cells were lysed and luciferase activity was measured in a Luminometer. Luminescence units were plotted against the compound concentrations using Graphpad software. EC₅₀ values of the compounds were

defined as the concentration required in reducing the luciferase activity by 50 %. The K_b values were calculated by feeding the concentration of agonist used in the assay and its EC_{50} value in the same software.

References: *Molecular Brain Research*, 2001, 90, 110-117 and *British Journal of Pharmacology*, 2006, 148, 1133-1143.

Results:

Compounds 1, 2 and 3 exhibit antagonistic activity in CRE-Luc based reporter gene assay on human recombinant 5-HT₆ receptor with no detectable agonist activity. The K_b values tabulated below are average of three independent experiments.

S. No	Example	K_b (nM)
1	Compound 1	4.2 ± 0.9
2	Compound 2	7.2 ± 1.8
3	Compound 3	1.6 ± 0.3

Example 2:

Determination of K_i value at 5-HT₆ receptor

Compound was tested at MDS pharma services and Novascreen according to the following procedures.

Materials and Methods:

Receptor source: Human recombinant expressed in Hela cells

Radioligand: [³H]-LSD (60-80 Ci/mmol)

Final ligand concentration - [1.5 nM]

Non-Specific Ligand: 5 μ M Serotonin (5-HT)

Reference compound: Methiothepin mesylate

Positive control: Methiothepin mesylate

Incubation conditions: Reactions were carried out in 50 mM Tris-HCl (pH 7.4) containing 10 mM MgCl₂, 0.5 mM EDTA for 60 minutes at 37°C. The reaction was terminated by rapid vacuum filtration onto the glass fiber filters. Radioactivity trapped onto the filters was determined and compared to the control values in order to ascertain any interactions of the test compound(s) with the cloned serotonin 5-HT₆ binding site.

Reference: *Molecular Pharmacology*, 1993, 43, 320-327.

Results:

Compounds 1, 2 and 3 selectively bind to 5-HT₆ receptor when tested by the *in-vitro* radioligand binding technique on human recombinant 5-HT₆ receptor. The K_i values are tabulated below.

S. No	Example	K _i (nM)
1	Compound 1	2.04
2	Compound 2	4.96
3	Compound 3	3.67

Example 3:

Determination of K_i value at 5-HT_{2A} receptor

Compound was tested according to the following procedures.

Materials and Methods:

Receptor source: Recombinant mammalian cells

Radioligand: [³H]-Ketanserine (47.3 Ci/mmol)

Final ligand concentration - [1.75 nM]

Non-Specific Ligand: 0.1 mM 1-Naphthylpiperazine (1-NP)

Reference compound: 1-Naphthylpiperazine (1-NP)

Positive control: 1-Naphthylpiperazine (1-NP)

Incubation conditions: Reactions were carried out in 67 mM Tris-HCl (pH 7.4) for 1 hour at 37°C. The reaction was terminated by rapid vacuum filtration onto the glass fiber filters. Radioactivity trapped onto the filters was determined and compared to the control values in order to ascertain any interactions of the test compound(s) with the cloned serotonin 5-HT_{2A} binding site.

Reference: *Methods in Molecular Biology*, 2002, 190, 31 – 49

Results:

Compounds 1, 2 and 3 bind weakly to 5-HT_{2A} receptor when tested by the in-vitro radioligand binding technique on human recombinant 5-HT_{2A} receptor. The K_i values tabulated below are average of three independent experiments.

S. No	Example	K _i
1	Compound 1	2514 ± 377 nM
2	Compound 2	>10 μM
3	Compound 3	926 ± 317 nM

Example 4:

Test compounds were also evaluated for their 5-HT₆ receptor selectivity over closely related serotonin subtypes like 5-HT_{1A}, 5-HT_{1B}, 5-HT_{1D}, 5-HT_{2A}, 5-HT_{2C}, 5-HT₄, 5-HT_{5A} and 5-HT₇ in commercial panel at Novascreen.

Compounds 1, 2 and 3 have shown selectivity of more than 250-fold over these receptor subtypes.

Example 5:

Object Recognition Task Model

The cognition enhancing properties of compounds of this invention were estimated by using this model.

Male Wistar rats (8-10 weeks old) were used as experimental animals. Four animals were housed in each cage. Animals were kept on 20 % food deprivation from a day prior to experimentation. Water was provided ad libitum throughout the experiment. Animals were maintained on a 12 hours light/dark cycle in temperature and humidity controlled room. The experiment was carried out in an open field made up of acrylic. Rats were habituated to individual arenas (open field) in the absence of any objects on day 1.

Rats received vehicle or donepezil and memantine or test compounds, donepezil and memantine on the day of habituation, before familiar (T₁) and choice (T₂) trials. During the familiarization phase (T₁), the rats were placed individually in the arena for 3 minutes, in which two identical objects (a₁ and a₂) were positioned 10 cm from the wall. 24 hours after T₁, trial for long-term memory test was assessed. The same rats were placed in the same arena as they were placed in T₁ trial. During the choice phase (T₂) rats were allowed to explore the arena for 3 minutes in presence of a copy of familiar object (a₃) and one novel object (b). During the T₁ and T₂ trial, explorations of each object (defined as sniffing, licking, chewing or having moving vibrissae whilst directing the nose towards the object at a distance of less than 1 cm) were recorded using stopwatch.

T₁ is the total time spent exploring the familiar objects (a₁ + a₂).

T₂ is the total time spent exploring the familiar object and novel object (a₃ + b).

Discriminative index is ratio of time spent exploring the novel object divided by sum of time spent exploring the novel object and familiar object in choice trial (T₂).

The object recognition test was performed as described in *Behavioural Brain Research*, 1988, 31, 47-59.

Results:

No significant increase in discriminative index was observed in the group treated with donepezil and memantine when compared to the vehicle treatment. However the group co-treated with test compounds, donepezil and memantine showed significant improvement in the memory end point (discriminative index). The results of this study are provided in figure 1a-1c.

Example 6:**Evaluation of acetylcholine modulation in ventral hippocampus of male Wistar rats****Experimental Procedure**

Male Wistar rats (240-300 g body weight) were stereotaxically implanted with a microdialysis guide cannula in ventral hippocampus (AP: -5.2 mm, ML: + 5.0 mm, DV: - 3.8 mm) under isoflurane anesthesia. Co-ordinates were taken according to atlas for the rat brain (Paxinos and Watson 2004) with reference points taken from bregma and vertical from the skull. The rats were allowed to recover individually for four days in a round bottom Plexiglas bowl with free access to feed and water.

After surgical recovery of 4-5 days, male Wistar rats were connected to dual quartz lined two-channel liquid swivel (Instech, UK) on a counter balance lever arm, which allowed unrestricted movements of the animal. Sixteen hours before start of study, a pre-equilibrated microdialysis probe (4 mm dialysis membrane) was inserted into the ventral hippocampus through the guide cannula. On the day of study, probe was perfused with artificial cerebrospinal fluid (aCSF; NaCl 147 mM, KCl 3 mM, MgCl₂ 1 mM, CaCl₂ 2H₂O 1.3 mM, NaH₂PO₄ 0.2 mM and Na₂HPO₄ 1 mM, pH 7.2) at a flow rate of 1.5 µL/min and a stabilization period of 2 h was maintained. Five basal samples were collected at 20 min intervals prior to the treatment of test compounds (1 or 3 mg/kg, *p.o.*) or vehicle. Donepezil (1 mg/kg, *s.c.*) and memantine (1 mg/kg, *s.c.*) were administered 30 min after administration of test compounds. Dialysate samples were collected for an additional period of 4 hours post treatment of test compounds. Dialysates were stored below -50°C prior to analysis.

Acetylcholine in dialysate was quantified using LC-MS/MS method in the calibration range of 0.103 – 103.491 nmol/L.

All microdialysis data for acetylcholine was plotted as percent change from mean dialysate basal concentrations with 100 % defined as the average of five pre-dose values. The percent change in acetylcholine levels were compared with donepezil and memantine combination using two-way analysis of variance (time and treatment), followed by Bonferroni's posttest. Area under the curve (AUC) values for percent change in acetylcholine levels were calculated and the statistical significance between the mean AUC value were compared against donepezil and memantine treatment using one-way ANOVA followed by Dunnett's test. Statistical significance was considered at a *p* value less than 0.05. Incorrect probe placement was considered as criteria to reject the data from animal.

Reference: Paxinos G. and Watson C. (2004) Rat brain in stereotaxic coordinates. Academic Press, New York

Results:

Compound 1

Treatment with donepezil (1 mg/kg, *s.c.*) and memantine (1 mg/kg, *s.c.*) produced increase in hippocampal acetylcholine levels to the maximum of 1204 ± 106 % of basal levels. Compound 1 produced dose dependent increase of acetylcholine in combination with donepezil (1 mg/kg, *s.c.*) and memantine (1 mg/kg, *s.c.*). The increase in acetylcholine after combination of compound 1, donepezil and memantine was significantly higher compared to donepezil and memantine combination. Mean maximum increase in acetylcholine was observed to be 1383 ± 194 and 2136 ± 288 % of pre-dose levels after 1 and 3 mg/kg, *p.o.*, combinations, respectively (Figure 2 (a)).

Mean area under the curve values (AUC) calculated after treatment of compound 1 (3 mg/kg, *p.o.*), donepezil (1 mg/kg, *s.c.*) and memantine (1 mg/kg, *s.c.*) were significantly higher compared to donepezil (1 mg/kg, *s.c.*) and memantine (1 mg/kg, *s.c.*) combination (Figure 2 (b)).

Compound 2

Treatment with donepezil (1 mg/kg, *s.c.*) and memantine (1 mg/kg, *s.c.*) produced increase in hippocampal acetylcholine levels to the maximum of 691 ± 148 % of basal levels. The increase in acetylcholine after combination of compound 2 (3 mg/kg, *p.o.*), donepezil (1 mg/kg, *s.c.*) and memantine (1 mg/kg, *s.c.*) was significantly higher compared to donepezil and memantine combination with mean maximum increase of 1776 ± 365 % above pre-dose levels (Figure 3 (a)).

Mean area under the curve value (AUC) calculated after treatment of compound 2 (3 mg/kg, *p.o.*), donepezil (1 mg/kg, *s.c.*) and memantine (1 mg/kg, *s.c.*) combination was significantly higher compared to donepezil (1 mg/kg, *s.c.*) and memantine (1 mg/kg, *s.c.*) combination (Figure 3 (b)).

Compound 3

Treatment with donepezil (1 mg/kg, *s.c.*) and memantine (1 mg/kg, *s.c.*) produced increase in hippocampal acetylcholine levels to the maximum of 691 ± 148 % of basal levels. The increase in acetylcholine after combination of compound 3 (3 mg/kg, *p.o.*), donepezil (1 mg/kg, *s.c.*) and memantine (1 mg/kg, *s.c.*) was significantly higher compared to donepezil and memantine combination with mean maximum increase of 1141 ± 258 % above pre-dose levels (Figure 4 (a)).

Mean area under the curve value (AUC) calculated after treatment of compound 3 (3 mg/kg, *p.o.*), donepezil (1 mg/kg, *s.c.*) and memantine (1 mg/kg, *s.c.*) combination was significantly higher compared to donepezil (1 mg/kg, *s.c.*) and memantine (1 mg/kg, *s.c.*) combination (Figure 4 (b)).

Example 7:

Evaluation of theta modulation in dorsal hippocampus of anesthetized male Wistar rats

Synchronous hippocampal EEG activity occurring in θ rhythm (frequency range of 4 to 8 Hz) has been associated with mnemonic processes in vivo.

Experimental Procedure

Male Wistar rats (240-320 g) were anesthetized with 1.2 to 1.5 g/kg urethane intraperitoneally, under anesthesia a catheter was surgically implanted in the left femoral vein for administration of drugs. After cannulation, the animal was placed in a stereotaxic frame for implanting an electrode (stainless steel wire, Plastics One) into the dorsal hippocampus (AP, -3.8 mm; ML, +2.2 mm; DV, -1.5 mm from dura; Paxinos and Watson, 1994) and bipolar stimulating electrode (untwisted stainless steel wires, separated by 0.75–1.0 mm at their tips, Plastics One) was implanted in the NPO (AP, -7.8 mm; ML, ± 1.8 mm; DV, -6.0 mm; Paxinos and Watson, 1994). Additionally one electrode was implanted into the cerebellum which served as a reference. Hippocampal θ rhythm was evoked via a 6-s electrical stimulation train (20-160 μ A, 0.3-ms pulse duration, 250 Hz) delivered to the

NPO at a rate of 0.01 trains/s with a Grass S88 stimulator and PSIU6 stimulus isolation unit (Grass Medical Instruments, Quincy, MA). EEG was recorded at a rate of 1000Hz using Ponemah (Version 5.2) software and stored for off-line analysis using NeuroScore (Version 3.0). Baseline amplitude level was achieved by using the current required to elicit θ rhythm to 50 % of the maximal amplitude under control conditions. After the stabilization period of one hour, Baseline recording was done for 30 min followed by the treatment of vehicle or compound 1 (1 mg/kg, *i.v.*). Donepezil (0.3 mg/kg, *i.v.*) and memantine (0.3 mg/kg, *i.v.*) was administered 30-min after compound 1 treatment and recording was continued for additional 1 hour.

Power in the θ frequency in the stimulation period during the 30 minutes baseline period was calculated and the percent changes in these measures post treatment were calculated. The percent change in relative θ power after combination treatment of compound 1, donepezil and memantine was compared with donepezil and memantine using two-way analysis of variance (time and treatment), followed by Bonferroni's posttest. Statistical significance was considered at a *p* value less than 0.05.

Reference: Paxinos G. and Watson C. (2004) Rat brain in stereotaxic coordinates. Academic Press, New York

Results:

Treatment with donepezil (0.3 mg/kg, *i.v.*) and memantine (0.3 mg/kg, *i.v.*) combination produced about 176 ± 25 % increases in hippocampal θ power. Compound 1 (1 mg/kg, *i.v.*) in combination with donepezil (0.3 mg/kg, *i.v.*) and memantine (0.3 mg/kg, *i.v.*) produced significant increase in θ power levels and peak levels reached up to 221 ± 25 % of pre-dose levels (Figure 5).

We claim:

1. A combination comprising a pure 5-HT₆ receptor antagonist, acetylcholinesterase inhibitor and NMDA receptor antagonist, wherein the pure 5-HT₆ receptor antagonist is
5 selected from the group consisting of:
1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole;
1-[(4-Fluorophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole;
1-[(4-Isopropylphenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole;
and a pharmaceutically acceptable salt thereof.
- 10 2. The combination as claimed in claim 1, wherein the acetylcholinesterase inhibitor is selected from the group consisting of donepezil, rivastigmine, galantamine and a pharmaceutically acceptable salt thereof.
- 15 3. The combination as claimed in claim 1 or 2, wherein the NMDA receptor antagonist is memantine or a pharmaceutically acceptable salt thereof.
4. The combination as claimed in any one of claims 1 to 3, wherein the pure 5-HT₆ receptor antagonist is 1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-
20 1H-indole or a pharmaceutically acceptable salt thereof.
5. The combination as claimed in any one of claims 1 to 4, wherein the pharmaceutically acceptable salt of the pure 5-HT₆ receptor antagonist is selected from the group consisting of:
- 25 1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole dimesylate monohydrate;
1-[(4-Fluorophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole dihydrochloride; and
1-[(4-Isopropylphenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole
30 dihydrochloride.

6. The combination as claimed in any one of claims 1 to 5, wherein the pure 5-HT₆ receptor antagonist is 1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole dimesylate monohydrate.
- 5 7. The combination as claimed in any one of claims 1 to 6, wherein the acetylcholinesterase inhibitor is donepezil or a pharmaceutically acceptable salt thereof.
8. The combination as claimed in any one of claims 1 to 7, wherein the acetylcholinesterase inhibitor is donepezil hydrochloride.
- 10 9. The combination as claimed in any one of claims 1 to 8, wherein the NMDA receptor antagonist is memantine hydrochloride.
- 15 10. A combination comprising 1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole dimesylate monohydrate, donepezil hydrochloride and memantine hydrochloride.
11. The combination as claimed in any one of claims 1 to 10, for use in the treatment of a cognitive disorder in a patient.
- 20 12. The combination as claimed in claim 11, wherein the cognitive disorder is selected from the group consisting of Alzheimer's disease, schizophrenia, Parkinson's disease, Lewy body dementia, vascular dementia, and frontotemporal dementia.
- 25 13. Use of a therapeutically effective amount of the combination as claimed in any one of claims 1 to 10 for administration to a patient to treat a cognitive disorder.
14. The use as claimed in claim 14, wherein the cognitive disorder is selected from the group consisting of Alzheimer's disease, schizophrenia, Parkinson's disease, Lewy body dementia, vascular dementia, and frontotemporal dementia.
- 30

15. A compound, 1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole or a pharmaceutically acceptable salt thereof for use in combination with acetylcholinesterase inhibitor and NMDA receptor antagonist for the treatment of Alzheimer's disease in a patient.
- 5
16. A compound, 1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole or a pharmaceutically acceptable salt thereof for use as an adjunct treatment of Alzheimer's disease in a patient on stable treatment with acetylcholinesterase inhibitor and NMDA receptor antagonist.
- 10
17. The compound as claimed in claim 15 or 16, wherein the acetylcholinesterase inhibitor is selected from the group consisting of donepezil, rivastigmine, galantamine and a pharmaceutically acceptable salt thereof.
- 15
18. The compound as claimed in claim 15 or 16, wherein the NMDA receptor antagonist is memantine or a pharmaceutically acceptable salt thereof.
19. Use of a therapeutically effective amount of 1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole or a pharmaceutically acceptable salt thereof
- 20 for administration to treat Alzheimer's disease in a patient on stable treatment with acetylcholinesterase inhibitor and NMDA receptor antagonist.
20. The use as claimed in claim 19, wherein the acetylcholinesterase inhibitor is selected from the group consisting of donepezil, rivastigmine, galantamine and a pharmaceutically
- 25 acceptable salt thereof.
21. The use as claimed in claim 19, wherein the NMDA receptor antagonist is memantine or a pharmaceutically acceptable salt thereof.
- 30
22. The use as claimed in claim 19, wherein the administration of 1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole or a pharmaceutically acceptable salt thereof is 1 mg to 200 mg per day.

23. The use as claimed in claim 19, wherein the administration of 1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole or a pharmaceutically acceptable salt thereof is 1 mg to 10 mg per day.
- 5
24. The use as claimed in claim 19, wherein the administration of 1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole or a pharmaceutically acceptable salt thereof is 25 mg to 125 mg per day.
- 10
25. The use as claimed in claim 19, wherein the administration of 1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole or a pharmaceutically acceptable salt thereof is 150 mg to 200 mg per day.
- 15
26. The use as claimed in claim 19, wherein the administration of 1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole or a pharmaceutically acceptable salt thereof is 25 mg to 75 mg per day.
- 20
27. The use as claimed in claim 19, wherein the administration of 1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole or a pharmaceutically acceptable salt thereof is 75 mg to 150 mg per day.
28. The use as claimed in claim 20, wherein the administration of donepezil or a pharmaceutically acceptable salt thereof is 1 mg to 30 mg per day.
- 25
29. The use as claimed in claim 21, wherein the administration of memantine or a pharmaceutically acceptable salt thereof is 1 mg to 40 mg per day.
30. A pharmaceutical composition comprising the combination as claimed in any one of claims 1 to 10, and pharmaceutically acceptable excipients.
- 30
31. The pharmaceutical composition as claimed in claim 30, for use in the treatment of cognitive disorders selected from the group consisting of Alzheimer's disease,

schizophrenia, Parkinson's diseases, Lewy body dementia, vascular dementia, and frontotemporal dementia.

32. The pharmaceutical composition as claimed in claim 30 or 31, wherein the 1-[(2-
5 Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole or a
pharmaceutically acceptable salt thereof is present in an amount of 1 mg to 300 mg.

33. The pharmaceutical composition as claimed in claim 30 or 31, wherein the 1-[(2-
Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole or a
10 pharmaceutically acceptable salt thereof is present in an amount of 35 mg to 200 mg.

34. The pharmaceutical composition as claimed in claim 30 or 31, wherein the 1-[(2-
Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole or a
pharmaceutically acceptable salt thereof is present in an amount of 200 mg to 300 mg.

15

35. The pharmaceutical composition as claimed in claim 30 or 31, wherein the 1-[(2-
Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole or a
pharmaceutically acceptable salt thereof is present in an amount of 75 mg or 150 mg.

20 36. The pharmaceutical composition as claimed in any one of claims 30 to 35, wherein the
acetylcholinesterase inhibitor is donepezil or a pharmaceutically acceptable salt thereof and
is present in an amount of 2 mg to 30 mg.

37. The pharmaceutical composition as claimed in any one of claims 30 to 36, wherein the
25 NMDA receptor antagonist is memantine or a pharmaceutically acceptable salt thereof and
is present in an amount of 1 mg to 40 mg.

38. The use as claimed in any one of claims 19 to 29, wherein the administration of 1-[(2-
Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole or a
30 pharmaceutically acceptable salt thereof is by oral, nasal, local, dermal or parenteral routes.

39. The use as claimed in any one of claims 19 to 29, wherein the administration of 1-[(2-Bromophenyl)sulfonyl]-5-methoxy-3-[(4-methyl-1-piperazinyl)methyl]-1H-indole or a pharmaceutically acceptable salt thereof is one to three times per day, one to three times per week or one to three times per month.

5


40. Use of the combination of any one of claims 1 to 10 in the manufacture of a medicament for treatment of a cognitive disorder in a patient.

41. The use of claim 40, wherein the cognitive disorder is selected from the group
10 consisting of Alzheimer's disease, schizophrenia, Parkinson's diseases, Lewy body dementia, vascular dementia, and frontotemporal dementia.


42. Use of the pharmaceutical composition of claim 30 in the manufacture of a medicament for treatment of a cognitive disorder in a patient.

15


43. The use of claim 42, wherein the cognitive disorder is selected from the group consisting of Alzheimer's disease, schizophrenia, Parkinson's diseases, Lewy body dementia, vascular dementia, and frontotemporal dementia.

Figure 1a

Data represents Mean ± SEM of discriminative index. * $p < 0.05$ Vs vehicle (Kruskal-Wallis followed by Dunn's Multiple Comparison Test) N=11-12

Figure 1b


Data represents Mean ± SEM of discriminative index. *** $p < 0.001$ Vs vehicle (Kruskal-Wallis followed by Dunn's Multiple Comparison Test) N=11-14

Figure 1c


Data represents Mean $\pm$ SEM of discriminative index. * $p < 0.05$ Vs vehicle (Kruskal-Wallis followed by Dunn's Multiple Comparison Test) N=17-21

Figure 2

(a)




(b)




(a) Percent change from mean basal levels expressed as Mean $\pm$ SEM. ** $p < 0.01$, *** $p < 0.001$ Vs donepezil + memantine combination (Bonferroni post test). (b) Cumulative changes in acetylcholine levels expressed as mean area under the curve (AUC) $\pm$ S.E.M. ** $p < 0.01$ (Dunnett's Multiple Comparison Test).

Figure 3

(a)




(b)




(a) Percent change from mean basal levels expressed as Mean $\pm$ SEM. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ Vs donepezil + memantine combination (Bonferroni post test). (b) Cumulative changes in acetylcholine levels expressed as mean area under the curve (AUC) $\pm$ S.E.M. * $p < 0.05$ (Dunnett's Multiple Comparison Test).

Figure 4

(a)




(b)




(a) Percent change from mean basal levels expressed as Mean $\pm$ SEM. * $p < 0.05$, ** $p < 0.01$, Vs donepezil + memantine combination (Bonferroni post test). (b) Cumulative changes in acetylcholine levels expressed as mean area under the curve (AUC) $\pm$ S.E.M. * $p < 0.05$ (Dunnett's Multiple Comparison Test).

Figure 5


(a)



(b)



(a) Data expressed as Mean $\pm$ SEM. * p <0.05, ** p <0.01 Vs donepezil and memantine combination (Bonferroni post test) (b) Area under the curve value $\pm$ SEM. * p <0.05 Vs donepezil + memantine combination (Dunnett's post test)



Data represents Mean $\pm$ SEM of discriminative index. * $p < 0.05$ Vs vehicle (Kruskal-Wallis followed by Dunn's Multiple Comparison Test) N=11-12