



(43) International Publication Date
22 September 2016 (22.09.2016)

WIPO | PCT

(10) International Publication Number
WO 2016/149454 A1

- (51) **International Patent Classification:**
C09K 3/32 (2006.01) *A62D 3/30* (2007.01)
- (21) **International Application Number:**
PCT/US2016/022776
- (22) **International Filing Date:**
17 March 2016 (17.03.2016)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**
14/659,919 17 March 2015 (17.03.2015) US
- (71) **Applicant:** **BAKER HUGHES INCORPORATED**
[US/US]; P.o. Box 4740, Houston, Texas 77210-4740 (US).
- (72) **Inventors:** **MAZYAR, Oleg A.**; 2929 Allen Parkway, Suite 2100, Houston, TX 77019-2118 (US). **KHABASHESKU, Valery N.**; 2929 Allen Parkway, Suite 2100, Houston, TX 77019-2118 (US). **CHAKRABORTY, Soma**; 2929 Allen Parkway, Suite 2100, Houston, TX 77019-2118 (US). **AGRAWAL, Gaurav**; 2929 Allen Parkway, Suite 2100, Houston, TX 77019-2118 (US). **HAIN, Toby D.**; 2929 Allen Parkway, Suite 2100, Houston, TX 77019-2118 (US).

(81) **Designated States** (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) **Designated States** (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) **Title:** FUNCTIONALIZED SILICATE NANOPARTICLE COMPOSITION, REMOVING AND EXFOLIATING ASPHALTENES WITH SAME

(57) **Abstract:** Removing an asphaltene particle from a substrate includes contacting a silicate nanoparticle with a chemical group to form a functionalized silicate nanoparticle, the chemical group includes a first portion; and a second portion comprising an aromatic moiety or a nonaromatic moiety, the first portion being bonded to the silicate nanoparticle; contacting the asphaltene particle with the functionalized silicate nanoparticle, the asphaltene particle being disposed on the substrate; interposing the functionalized silicate nanoparticle between the asphaltene particle and the substrate; and separating the asphaltene particle from the substrate with the functionalized silicate nanoparticle to remove the asphaltene particle. A composition includes a functionalized silicate nanoparticle comprising a reaction product of a silicate nanoparticle and a functionalization compound; and a fluid. The functionalization compound includes a chemical group that includes a first portion, the first portion being directly bonded to the silicate nanoparticle in the functionalized silicate nanoparticle; and a second portion including an aromatic moiety or a nonaromatic moiety.



WO 2016/149454 A1

FUNCTIONALIZED SILICATE NANOPARTICLE COMPOSITION, REMOVING AND EXFOLIATING ASPHALTENES WITH SAME

CROSS REFERENCE TO RELATED APPLICATIONS

[0001] This application claims the benefit of U.S. Application No. 14/659919, filed on March 17, 2015, which is incorporated herein by reference in its entirety.

BACKGROUND

[0002] Asphaltenes are a major component in crude oil, and there is general agreement as to the deleterious effects of asphaltenes in the reduction of oil extraction and processing in the petrochemical industry. Asphaltenes can deposit in the pores of formations, blocking the flow of fluids. Additionally, asphaltenes can precipitate from a stream of oil and coat boreholes, production tubing, and transport lines. Moreover, in a processing facility, asphaltenes can foul processing equipment and poison catalysts.

[0003] Asphaltene molecules have been widely reported as having a fused polyaromatic ring system and containing heteroatoms such as sulfur, oxygen, nitrogen, and the like. The heteroatoms may be part of the aromatic ring system or part of other carbocyclic rings, linking groups, or functional groups. Two structural motifs for asphaltene molecules are the so-called continental and archipelago structures. In the continental structure, alkyl chains connect to and branch from a central polyaromatic ring system, which is believed to contain several fused aromatic rings, e.g., 5 or more aromatic rings. In the archipelago structure, multiple polyaromatic ring systems are connected by alkyl chains that may contain a heteroatom, and additional alkyl chains extend freely from the polyaromatic rings. The number of fused aromatic rings in the continental structure can be greater than the number of fused aromatic rings in the archipelago structure.

[0004] In addition to the aromatic regions of the asphaltenes, heteroatoms provide the asphaltenes with polar regions, and the terminal alkyl chains provide hydrophobic regions. Consequently, it is believed that asphaltene molecules aggregate into various micellar structures in oil, with the alkyl chains interacting with the aliphatic oil components. Resin from the oil can insert between aromatic planes of neighboring asphaltene molecules in asphaltene aggregates, aiding in maintaining their micellar structure. Asphaltenes can precipitate from oil in structures where asphaltene molecules form stacked layers having aligned aromatic regions and aligned aliphatic regions.

[0005] Materials and methods for treating and removal of asphaltenes from oil environments such as a reservoir would be well received in the art.

BRIEF DESCRIPTION

[0006] The above and other deficiencies of the prior art are overcome by, in an embodiment, a process for removing an asphaltene particle from a substrate, the process comprising: contacting a silicate nanoparticle with a chemical group to form a functionalized silicate nanoparticle, the chemical group comprising: a first portion; and a second portion comprising an aromatic moiety or a nonaromatic moiety, the first portion being directly bonded to the silicate nanoparticle in the functionalized silicate nanoparticle; and the second portion comprising a moiety of a guanidine, a guanidinium salt, a biguanidine, a biguanidinium salt, or a sulfonium salt; contacting the asphaltene particle with the functionalized silicate nanoparticle, the asphaltene particle being disposed on the substrate; interposing the functionalized silicate nanoparticle between the asphaltene particle and the substrate; and separating the asphaltene particle from the substrate with the functionalized silicate nanoparticle to remove the asphaltene particle from the substrate.

[0007] In another embodiment, a composition comprises: a functionalized silicate nanoparticle comprising a reaction product of a silicate nanoparticle; and a functionalization compound comprising quaternary ammonium salt, quaternary phosphonium salt, alkoxy silane, halide, guanidine, guanidinium salt, biguanidine, biguanidinium salt, sulfonium salt, or a combination thereof; and a fluid, wherein the functionalization compound includes a chemical group comprising: a first portion, the first portion being directly bonded to the silicate nanoparticle in the functionalized silicate nanoparticle; and a second portion comprising: an aromatic moiety or a nonaromatic moiety, and the composition is effective to remove an aromatic compound from a substrate comprising a metal, composite, sand, rock, mineral, glass, formation, downhole element, or a combination thereof.

[0008] A process for removing an asphaltene particle from a substrate comprises: contacting a silicate nanoparticle with a chemical group to form a functionalized silicate nanoparticle, the chemical group comprising: a first portion; and a second portion comprising a nonaromatic moiety, the first portion being directly bonded to the silicate nanoparticle in the functionalized silicate nanoparticle; and the second portion comprising a moiety of a guanidine, a guanidinium salt, a biguanidine, a biguanidinium salt, or a sulfonium salt; contacting the asphaltene particle with the functionalized silicate nanoparticle, the asphaltene particle being disposed on the substrate; interposing the functionalized silicate nanoparticle

between the asphaltene particle and the substrate; and separating the asphaltene particle from the substrate with the functionalized silicate nanoparticle to remove the asphaltene particle from the substrate.

DETAILED DESCRIPTION

[0009] A detailed description of one or more embodiments is presented herein by way of exemplification and not limitation.

[0010] An asphaltene particle includes any collection of asphaltene molecules, for example, a micelle, precipitate, layered asphaltene molecules, aggregate, cluster, and the like. Interactions among the asphaltene molecules in an asphaltene particle can include hydrogen bonding, dipole-dipole interactions, and π - π interactions. Without wishing to be bound by theory, disruption of these interactions can lead to exfoliation of an asphaltene molecule from the asphaltene particle. Since asphaltenes form layered aggregates that resemble the layered sheet structure of graphite, perturbing the layered asphaltene structure allows for asphaltene production from decomposed, e.g., exfoliated asphaltene aggregates. Such deagglomeration is useful for extraction of oil from an oil environment, e.g., a formation, as well as for restoration of the permeability of a plugged or flow-constricted reservoir. The methods and compositions herein are applicable to a multitude of environments such as downhole as well as to a ground environment.





[0011] It has been found that perturbing the internal structure of asphaltene particles, for example, in a micelle or other aggregate, can lead to increased quality of oil containing asphaltenes. Further, degradation of asphaltene aggregates herein enhances production of petroleum fluid in a downhole, subsurface, or ground environment. Furthermore, removal of asphaltene from pores of a rock formation, within a reservoir, or from a sidewall of a tubular, production tubing, borehole, or transportation tube can improve the permeability of such structures, leading to increased quality of oil as well as enhanced oil recovery from, e.g., a reservoir.





[0012] Moreover, without wishing to be bound by theory, it is believed that heteroatoms in the asphaltene structure interact strongly with various materials such as metals, minerals, and polar surfaces. Therefore, asphaltenes coat rock formations, sand, metal, and polymer components such as tubulars, sand screens, or packers. These deleterious adhesions lead to equipment malfunction, failure, or flow blockage. In order to alleviate this issue, a composition herein can separate and remove the asphaltene particles that interact, such as by adsorption or blockage, with such items or materials.

[0013] In an embodiment, a composition includes a functionalized silicate nanoparticle and a fluid. The functionalized silicate nanoparticle is a reaction product of a silicate nanoparticle and a functionalization compound. The functionalization compound includes a chemical group that has a first portion, e.g., a linker, and a second portion (e.g., a tail), which includes an aromatic or a nonaromatic moiety. The first portion is directly bonded to the silicate nanoparticle in the functionalized silicate nanoparticle. The composition is effective to remove an aromatic or olefin compound, e.g., an asphaltene, from a substrate. The substrate can be a metal, composite, sand, rock, mineral, glass, formation, downhole element, or a combination thereof.

[0014] As used herein, the term “aromatic” includes an aryl or heteroaryl group. Thus, an aromatic compound includes an aryl moiety or heteroaryl moiety.

[0015] The functionalization compound can have a structure of formula 1, formula 2, formula 3, formula 4 or a salt thereof, formula 5 or a salt thereof, formula 6 or a salt thereof, or formula 7:





wherein

a is an integer from 1 to 4;

b is an integer from 0 to 3;

the sum of a and b is 4 (i.e., $a + b = 4$) so that the valence of A and Si is completely filled and not exceeded

m is 0 or 1, n is independently an integer from 0 to 20;

Ar is an aromatic moiety or a nonaromatic moiety, wherein each Ar is the same or different, and when Ar is an aromatic moiety, Ar is independently a C₆ to C₃₀ aryl group, C₃ to C₃₀ heteroaryl, or combination thereof;

L is a linker group, wherein each L is the same or different, and L is independently a bond, C₁ to C₃₀ alkylene, C₃ to C₃₀ cycloalkenylene, C₁ to C₃₀ fluoroalkylene, C₃ to C₃₀ cycloalkylene, C₃ to C₃₀ heterocycloalkylene, C₆ to C₃₀ arylene, C₆ to C₄₀ aralkylene, C₆ to C₃₀ aryleneoxy, C₃ to C₃₀ heteroarylene, C₆ to C₄₀ heteroaralkylene, C₂ to C₃₀

alkenylene, C2 to C30 alkynylene, C1 to C30 amide, amine, C1 to C30 oxyalkylene, C1 to C30 oxyarylene, oxygen (O), sulfur (S), or a combination thereof. L can be substituted or unsubstituted (with the exception of O and S). Moreover, L can be linear or branched, with the exception of O and S.

A is nitrogen (N) or phosphorous (P);

X⁻ is an anion of a halogen;

R¹ is a substituent on A or Si, wherein each R¹ is the same or different, and R¹ independently is hydrogen, C1 to C30 alkyl group, C1 to C30 alkenyl group, C1 to C30 alkoxy group, C1 to C30 alkynyl group, C1 to C30 aryloxy, halogen, C6 to C30 aryl group, C1 to C30 amide, amino, C3-C30 cycloalkenyl, C3-C30 cycloalkyl, C3-C30 fluoroalkyl, C1-C30 heteroalkyl, C3-C30 heteroaryl, hydroxy, C1-C30 oxyalkyl, or a combination thereof, and each foregoing group can be substituted or unsubstituted or can be linear or branched;

R₁-R₉ and R₁₁-R₂₄ are independently hydrogen atom, alkyl, heterocyclic, carbonyl, amino, amide, sulfonamide, phosphoramidate, imide moiety, an aromatic moiety, or a combination thereof;

R₁₀ is a bond, alkylene, heterocyclic biradical, carbonyl, amino, amide, sulfonamide, phosphoramidate, imide biradical, an aromatic moiety, or a combination thereof;

R₂₅ is a halogen or an alkoxy group; and

G is of formulas (4)-(7), wherein at least one of R₁-R₉ and R₁₁-R₂₄ is a divalent group attached to oxygen atom or silicon atom.

[0016] As used herein, “alkenyl” refers to a straight or branched chain, monovalent C2-C10 hydrocarbon group having at least one carbon-carbon double bond (e.g., ethenyl (-HC=CH₂)). As used herein, “alkenylene” refers to a straight or branched chain, divalent C2-C30 hydrocarbon group having at least one carbon-carbon double bond (e.g., ethenylene (-HC=CH-)). As used herein, “alkoxy” refers to an alkyl group that is linked via an oxygen (i.e., -O-alkyl). Non-limiting examples of C1 to C30 alkoxy groups include methoxy groups, ethoxy groups, propoxy groups, isobutyloxy groups, sec-butyloxy groups, pentyloxy groups, iso-amtyloxy groups, and hexyloxy groups.

[0017] As used herein, “alkyl” refers to a straight or branched chain saturated aliphatic hydrocarbon having the specified number of carbon atoms, specifically 1 to 12 carbon atoms, more specifically 1 to 6 carbon atoms. Alkyl groups include, for example, groups having from 1 to 50 carbon atoms (C1 to C50 alkyl).

[0018] As used herein, “alkylene” refers to a straight, branched or cyclic divalent aliphatic hydrocarbon group, and can have from 1 to about 18 carbon atoms, more

specifically 2 to about 12 carbons. Exemplary alkylene groups include methylene ($-\text{CH}_2-$), ethylene ($-\text{CH}_2\text{CH}_2-$), propylene ($-(\text{CH}_2)_3-$), cyclohexylene ($-\text{C}_6\text{H}_{10}-$), ethyleneoxy ($-\text{CH}_2\text{CH}_2\text{O}-$), methylenedioxy ($-\text{O}-\text{CH}_2-\text{O}-$), or ethylenedioxy ($-\text{O}-(\text{CH}_2)_2-\text{O}-$).

[0019] As used herein, “alkynyl” refers to a straight or branched chain, monovalent hydrocarbon group having at least one carbon-carbon triple bond (e.g., ethynyl). As used herein, “alkynylene” refers to a straight or branched chain divalent aliphatic hydrocarbon that has one or more unsaturated carbon-carbon bonds, at least one of which is a triple bond (e.g., ethynylene). As used herein, “amide” refers to a group of the formula $-\text{C}(\text{O})-\text{N}(\text{Rx})(\text{Ry})$ or $-\text{N}-\text{C}(\text{O})-\text{Rx}$, wherein Rx is an alkyl, an alkenyl, an alkynyl, a cycloalkyl, an aryl group, or heteroaryl group; and Ry is hydrogen or any of the groups listed for Rx. As used herein, “C1 to C15 amine group” is a group of the formula $-\text{N}(\text{Rw})(\text{Rz})$, wherein Rw is a C1 to C15 alkyl, a C1 to C15 alkenyl, a C1 to C15 alkynyl, a C3 to C15 cycloalkyl, a C6 to C15 aryl, or C3 to C15 heteroaryl group; and Rz is hydrogen or any of the groups listed for Rw.

[0020] As used herein, “aryl” refers to a cyclic moiety in which all ring members are carbon and at least one ring is aromatic, the moiety having the specified number of carbon atoms, specifically 6 to 24 carbon atoms, more specifically 6 to 12 carbon atoms. More than one ring may be present, and any additional rings may be independently aromatic, saturated or partially unsaturated, and may be fused, pendant, spirocyclic or a combination thereof.

[0021] As used herein, “arylalkylene” group is an aryl group linked via an alkylene moiety. The specified number of carbon atoms (e.g., C7 to C30) refers to the total number of carbon atoms present in both the aryl and the alkylene moieties. Representative arylalkyl groups include, for example, benzyl groups.

[0022] As used herein, “arylene” refers to a divalent radical formed by the removal of two hydrogen atoms from one or more rings of an aromatic hydrocarbon, wherein the hydrogen atoms may be removed from the same or different rings (preferably different rings), each of which rings may be aromatic or nonaromatic. As used herein, “aryloxy” refers to an aryl moiety that is linked via an oxygen (i.e., $-\text{O}-\text{aryl}$). As used herein, “cycloalkenyl” refers to a monovalent group having one or more rings and one or more carbon-carbon double bond in the ring, wherein all ring members are carbon (e.g., cyclopentyl and cyclohexyl).

[0023] As used herein, “cycloalkyl” refers to a group that comprises one or more saturated and/or partially saturated rings in which all ring members are carbon, such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, adamantyl and partially saturated variants of the foregoing, such as cycloalkenyl groups (e.g., cyclohexenyl) or cycloalkynyl groups. Cycloalkyl groups do not include an aromatic ring or a heterocyclic

ring. When the numbers of carbon atoms is specified (e.g., C3 to C15 cycloalkyl), the number refers to the number of ring members present in the one or more rings.

[0024] As used herein, “cycloalkenylene” refers to a stable aliphatic 5-15-membered monocyclic or polycyclic, divalent radical having at least one carbon-carbon double bond, which comprises one or more rings connected or bridged together. Unless mentioned otherwise, the cycloalkenylene radical can be linked at any desired carbon atom provided that a stable structure is obtained. If the cycloalkenylene radical is substituted, this may be so at any desired carbon atom, once again provided that a stable structure is obtained. Examples thereof are cyclopentenylene, cyclohexenylene, cycloheptenylene, cyclooctenylene, cyclononenylene, cyclodecenylene, norbornenylene, 2-methylcyclopentenylene, 2-methylcyclooctenylene.

[0025] As used herein, “cycloalkylene” refers to a divalent radical formed by the removal of two hydrogen atoms from one or more rings of a cycloalkyl group (a nonaromatic hydrocarbon that comprises at least one ring).

[0026] As used herein, “fluoroalkyl” refers to an alkyl group in which at least one hydrogen is replaced with fluorine. “Fluoroalkylene” refers to an alkylene group in which at least one hydrogen is replaced with fluorine.

[0027] As used herein, “halogen” refers to one of the elements of group 17 of the periodic table (e.g., fluorine, chlorine, bromine, iodine, and astatine).

[0028] As used herein, the prefix “hetero” means that the compound or group includes a heteroatom (e.g., 1, 2, or 3 heteroatom(s)), wherein the heteroatom(s) is each independently N, O, S, Si, or P.

[0029] As used herein, “heteroalkyl” group is an alkyl group that comprises at least one heteroatom covalently bonded to one or more carbon atoms of the alkyl group. Each heteroatom is independently chosen from nitrogen (N), oxygen (O), sulfur (S), and phosphorus (P).

[0030] As used herein, “heteroaryl” refers to a monovalent carbocyclic ring group that includes one or more aromatic rings, in which at least one ring member (e.g., one, two or three ring members) is a heteroatom. In a C3 to C30 heteroaryl, the total number of ring carbon atoms ranges from 3 to 30, with remaining ring atoms being heteroatoms. Multiple rings, if present, may be pendent, spiro or fused. The heteroatom(s) are generally independently selected from nitrogen (N), oxygen (O), phosphorus (P), and sulfur (S).

[0031] As used herein, “heteroarylene” refers to a divalent radical formed by the removal of two hydrogen atoms from one or more rings of a heteroaryl moiety, wherein the

hydrogen atoms may be removed from the same or different rings (preferably the same ring), each of which rings may be aromatic or nonaromatic.

[0032] As used herein, “oxyalkyl” refers to an alkyl group to which at least one oxygen atom is covalently attached (e.g., via a single bond, forming a hydroxyalkyl or ether group, or double bond, forming a ketone or aldehyde moiety). As used herein, “oxyalkylene” refers to a divalent radical comprising an alkylene group to which at least one oxygen atom is covalently attached (e.g., via a single bond, forming a hydroxyalkylene or an ether group, or double bond, forming a ketone or aldehyde moiety). As used herein, “oxyarylene” moiety is an aromatic group in which all ring members are independently chosen from carbon and oxygen.

[0033] As used herein, “substituted” means that the compound or group is substituted with at least one (e.g., 1, 2, 3, or 4) substituent independently selected from a hydroxyl (-OH), a C1-9 alkoxy, a C1-9 haloalkoxy, an oxo (=O), a nitro (-NO₂), a cyano (-CN), an amino (-NH₂), an azido (-N₃), an amidino (-C(=NH)NH₂), a hydrazino (-NHNH₂), a hydrazono (-C(=NNH₂)-), a carbonyl (-C(=O)-), a carbamoyl group (-C(O)NH₂), a sulfonyl (-S(=O)₂-), a thiol (-SH), a thiocyno (-SCN), a tosyl (CH₃C₆H₄SO₂-), a carboxylic acid (-C(=O)OH), a carboxylic C1 to C6 alkyl ester (-C(=O)OR wherein R is a C1 to C6 alkyl group), a carboxylic acid salt (-C(=O)OM) wherein M is an organic or inorganic anion, a sulfonic acid (-SO₃H₂), a sulfonic mono- or dibasic salt (-SO₃MH or -SO₃M₂ wherein M is an organic or inorganic anion), a phosphoric acid (-PO₃H₂), a phosphoric acid mono- or dibasic salt (-PO₃MH or -PO₃M₂ wherein M is an organic or inorganic anion), a C1 to C12 alkyl, a C3 to C12 cycloalkyl, a C2 to C12 alkenyl, a C5 to C12 cycloalkenyl, a C2 to C12 alkynyl, a C6 to C12 aryl, a C7 to C13 arylalkylene, a C4 to C12 heterocycloalkyl, and a C3 to C12 heteroaryl instead of hydrogen, provided that the substituted atom's normal valence is not exceeded.

[0034] The aromatic moiety of the functionalization compound can include an aryl group or heteroaryl group. Exemplary aryl groups include anthracyl, azulenyl, benzocyclooctenyl, benzocycloheptenyl, biphenyl, chrysenyl, fluorenyl, indanyl, indenyl, naphthyl, pentalenyl, phenalenyl, phenanthrenyl, phenanthryl, phenyl, pyrenyl, tetrahydronaphthyl, a derivative thereof, or a combination thereof.

[0035] Exemplary heteroaryl groups include acridinyl, benzimidazolyl, benzofuranyl, benzofurazanyl, benzothiazolyl, benzothiophenyl, benzoxazolyl, carbazolyl, chromanyl, cinnolinyl, dibenzofuranyl, furazanyl, furopyridinyl, furyl, imidazolyl, indazolyl, indolinyl, indoliziny, indolyl, isochromanyl, isoindolinyl, isoindolyl, isoquinolinyl, isothiazolyl,

isoxazolyl, naphthyridinyl, oxadiazolyl, oxazolyl, perimidinyl, phenanthridinyl, phenanthrolinyl, phenazinyl, phenoxathiinyl, phenothiazinyl, phenoxazinyl, phthalazinyl, pteridinyl, purinyl, pyrazinyl, pyrazolyl, pyridazinyl, pyridinyl, pyrimidinyl, pyrrolyl, quinazolinyl, quinolinyl, quinoxalinyl, tetrazolyl, thiadiazolyl, thiazolyl, thienyl, triazinyl, triazolyl, a derivative thereof, or a combination thereof. The heteroaryl group may be attached at any heteroatom or carbon atom of the ring such that the result is a stable structure. Thus, for example, pyridyl represents 2-, 3-, or 4-pyridyl, thienyl represents 2- or 3-thienyl, and quinolinyl represents 2-, 3-, or 4-quinolinyl, and the like.

[0036] According to an embodiment, a substituent of the aromatic moiety (e.g., the aryl group or heteroaryl group) includes halogen, hydroxy, lower alkyl, lower alkoxy, lower aralkyl, $-NR^2$ (wherein R^2 is a lower alkyl), R^3CONH (wherein R^3 is phenyl or a lower alkyl), and $-OC(O)R^4$ (wherein R^4 is hydrogen, alkyl, or aralkyl). Other substituents for the aromatic moiety can be $-OR^5$, $-OC(O)R^5$, $-NR^5R^6$, $-SR^5$, $-R^5$, $-CN$, $-NO_2$, $-CO_2R^5$, $-CONR^5R^6$, $-C(O)R^5$, $-OC(O)NR^5R^6$, $-NR^6C(O)R^5$, $-NR^6C(O)_2R^5$, $-NR^5-C(O)NR^6R^7$, $-NH-C(NH_2)=NH$, $-NR^5C(NH_2)=NH$, $-NH-C(NH_2)=NR^5$, $-S(O)R^5$, $-S(O)_2R^5$, $-S(O)_2NR^5R^6$, $-N_3$, $-CH(Ph)_2$, perfluoro(C_1 - C_4)alkoxy, and perfluoro(C_1 - C_4)alkyl, in a number ranging from zero to the total number of open valences on the aromatic moiety ring system; and where R^5 , R^6 , and R^7 are independently selected from hydrogen, C_1 - C_8 alkyl or heteroalkyl, unsubstituted aryl and heteroaryl, (unsubstituted aryl)-(C_1 - C_4)alkyl, (unsubstituted aryl)oxy-(C_1 - C_4)alkyl, (unsubstituted heteroaryl)-(C_1 - C_4)alkyl, or (unsubstituted heteroaryl)oxy-(C_1 - C_4)alkyl.

[0037] In a specific embodiment, a substituent for the aromatic moiety of the aromatic compound includes, for example, an alkyl group (having 1 to 20 carbon atoms, specifically 1 to 12 carbon atoms, and more specifically 1 to 8 carbon atoms and including, e.g., methyl, ethyl, iso-propyl, tert-butyl, n-octyl, n-decyl, n-hexadecyl, cyclopropyl, cyclopentyl, cyclohexyl, and the like), an aryl group (having 6 to 30 nuclear carbon atoms, specifically 6 to 20 nuclear carbon atoms and including, for example, phenyl, naphthyl, biphenyl, anthranyl, phenanthryl, pyrenyl, chrysenyl, fluorenyl, and the like), an alkenyl group (having 2 to 20 carbon atoms, specifically 2 to 12 carbon atoms, and more specifically 2 to 8 carbon atoms and including, for example, vinyl, allyl, 2-butenyl, 3-pentenyl, and the like), an alkynyl group (having 2 to 20 carbon atoms, specifically 2 to 12 carbon atoms, and more specifically 2 to 8 carbon atoms and including, for example, propargyl, 3-pentynyl, and the like), an amino group (having 0 to 20 carbon atoms, specifically 0 to 12 carbon atoms, and more specifically 0 to 6 carbon atoms and including, for example, amino, methylamino,

dimethylamino, diethylamino, diphenylamino, dibenzylamino, and the like), an alkoxy group (having 1 to 20 carbon atoms, specifically 1 to 12 carbon atoms, and more specifically 1 to 8 carbon atoms and including, for example, methoxy, ethoxy, butoxy, and the like), an aryloxy group (having 6 to 20 carbon atoms, specifically 6 to 16 carbon atoms, and more specifically 6 to 12 carbon atoms and including, for example, phenoxy, 2-naphthyloxy, and the like), an acyl group (having 1 to 20 carbon atoms, specifically 1 to 16 carbon atoms, and more specifically 1 to 12 carbon atoms and including, for example, acetyl, benzoyl, formyl, pivaloyl, and the like), an alkoxycarbonyl group (having 2 to 20 carbon atoms, specifically 2 to 16 carbon atoms, and more specifically 2 to 12 carbon atoms and including, for example, methoxycarbonyl, ethoxycarbonyl, and the like), an aryloxycarbonyl group (having 7 to 20 carbon atoms, specifically 7 to 16 carbon atoms, and more specifically 7 to 10 carbon atoms and including, for example, phenyloxycarbonyl and the like), an acyloxy group (having 2 to 20 carbon atoms, specifically 2 to 16 carbon atoms, and more specifically 2 to 10 carbon atoms and including, for example, acetoxyl, benzoyloxy, and the like), an acylamino group (having 2 to 20 carbon atoms, specifically 2 to 16 carbon atoms, and more specifically 2 to 10 carbon atoms and including, for example, acetylamino, benzoylamino, and the like), an alkoxycarbonylamino group (having 2 to 20 carbon atoms, specifically 2 to 16 carbon atoms, and more specifically 2 to 12 carbon atoms and including, for example, methoxycarbonylamino and the like), an aryloxycarbonylamino group (having 7 to 20 carbon atoms, specifically 7 to 16 carbon atoms, and more specifically 7 to 12 carbon atoms and including, for example, phenyloxycarbonylamino and the like), a sulfonylamino group (having 1 to 20 carbon atoms, specifically 1 to 16 carbon atoms, and more specifically 1 to 12 carbon atoms and including, for example, methanesulfonylamino, benzenesulfonylamino, and the like), a sulfamoyl group (having 0 to 20 carbon atoms, specifically 0 to 16 carbon atoms and more specifically 0 to 12 carbon atoms and including, for example, sulfamoyl, methylsulfamoyl, dimethylsulfamoyl, phenylsulfamoyl, and the like), a carbamoyl group (having 1 to 20 carbon atoms, specifically 1 to 16 carbon atoms and more specifically 1 to 12 carbon atoms and including, for example, carbamoyl, methylcarbamoyl, diethylcarbamoyl, phenylcarbamoyl, and the like), an alkylthio group (having 1 to 20 carbon atoms, specifically 1 to 12 carbon atoms, and more specifically 1 to 8 carbon atoms and including, for example, methylthio, ethylthio, and the like), an arylthio group (having 6 to 20 carbon atoms, specifically 6 to 16 carbon atoms, and more specifically 6 to 12 carbon atoms and including, for example, phenylthio and the like), a sulfonyl group (having 1 to 20 carbon atoms, specifically 1 to 16 carbon atoms and more specifically 1 to 12 carbon atoms and including,

for example, mesyl, tosyl, and the like), a sulfinyl group (having 1 to 20 carbon atoms, specifically 1 to 16 carbon atoms, and more specifically 1 to 12 carbon atoms and including, for example, methanesulfinyl, benzenesulfinyl, and the like), a ureido group (having 1 to 20 carbon atoms, specifically 1 to 16 carbon atoms, and more specifically 1 to 12 carbon atoms and including, for example, ureido, methylureido, phenylureido, and the like), a phosphoramidate group (having 1 to 20 carbon atoms, specifically 1 to 16 carbon atoms, and more specifically 1 to 12 carbon atoms and including, for example, diethylphosphoramidate, phenylphosphoramidate, and the like), a hydroxy group, a mercapto group, a halogen atom (for example, a fluorine atom, chlorine atom, bromine atom, iodine atom, and the like), a cyano group, a sulfo group, a carboxyl group, a nitro group, a hydroxamic acid, a sulfinio group, a hydrazine group, an imino group, a heterocyclic group (having 1 to 30 carbon atoms, specifically 1 to 12 carbon atoms, including, for example, a nitrogen atom, an oxygen atom and a sulfur atom as a heteroatom and including, for example, imidazolyl, pyridyl, quinolyl, furyl, thienyl, piperidyl, morpholino, benzoxazolyl, benzimidazolyl, benzothiazolyl and carbazolyl), a silyl group (having 3 to 40 carbon atoms, specifically 3 to 30 carbon atoms, and more specifically 3 to 24 carbon atoms and including, for example, trimethylsilyl, triphenylsilyl, and the like), and the like.

[0038] In an embodiment, the R group of formula (1), (2), or (3) is one of the preceding substituents mentioned with the respect to the aromatic moiety.

[0039] In an embodiment, the linking groups of formula (1), (2), or (3) include groups obtained by converting the preceding substituents mentioned above with the respect to the aromatic moiety into divalent groups. Exemplary linking groups are a hetero atom, a substituted or unsubstituted alkylene group having 1 to 6 carbon atoms, a substituted or unsubstituted cycloalkylene group having 3 to 30 carbon atoms, a substituted or unsubstituted arylene group having 6 to 30 nuclear carbon atoms, or a substituted or unsubstituted heteroarylene group having 5 to 30 nuclear carbon atoms. The heteroatom can be, for example, an oxygen atom, a sulfur atom, a nitrogen atom and a silicon atom. The alkylene group can be, for example, methylene, ethylene, propylene, butylene, pentylene, hexylene, heptylene, octylene, dimethylmethylene, diphenylmethylene, and the like. The cycloalkylene group can be, for example, cyclopropylene, cyclobutylene, cyclopentylene and cyclohexylene, 1,1-cyclohexylene, and the like. The arylene group can be, for example, phenylene, biphenylene, terphenylene, naphthylene, anthracenylene, phenathrylene, chrysenylene, pyrenylene, fluorenylene, 2,6-diphenylnaphthalene-4',4''-ene, 2-phenylnaphthalene-2,4'-ene, fluorenylene, and the like. The heteroarylene group can be, for

example, a divalent residue of imidazole, benzimidazole, pyrrole, furan, thiophene, benzothiophene, oxadiazoline, indoline, carbazole, pyridine, quinoline, isoquinoline, benzoquinone, pyrrolidine, imidazolidine, piperidine, pyridylene, and the like.

[0040] According to an embodiment, the functionalization compound can be a quaternary ammonium salt, quaternary phosphonium salt, alkoxy silane, halide, a guanidine, a guanidinium salt, a biguanidine, a biguanidinium salt, a sulfonium salt, or the like.

[0041] Non-limiting examples of the aryl quaternary ammonium salt include benzylammonium chloride, benzyldimethyldodecylammonium chloride, benzyldimethyldodecylammonium chloride, benzyldimethylhexadecylammonium chloride, benzyldimethylhexylammonium chloride, benzyldimethyl(2-hydroxyethyl)ammonium chloride, benzyldimethyl(2-hydroxymethyl)ammonium chloride, benzyldimethyloctylammonium chloride, benzyldimethylstearyl ammonium chloride monohydrate, benzyldimethyltetradecylammonium chloride, benzyldodecyldimethylammonium bromide, benzyltributylammonium bromide, benzyltributylammonium chloride, benzyltributylammonium iodide, benzyltriethylammonium bromide, benzyltriethylammonium chloride, benzyltrimethylammonium bromide, benzyltrimethylammonium dichloriodate, bis(triphenylphosphoranylidene)ammonium chloride, (dodecyldimethyl-2-phenoxyethyl)ammonium bromide, (diisobutylphenoxyethoxyethyl)dimethylbenzylammonium chloride, (4-nitrobenzyl)trimethylammonium chloride, trimethylphenylammonium bromide, trimethylphenylammonium chloride, (vinylbenzyl)trimethylammonium chloride, 3-benzyl-5-(2-hydroxyethyl)-4-methylthiazolium chloride, 1,1-dimethyl-4-phenylpiperazinium iodide, and the like. In an embodiment, the aryl compound is an aryl phosphine such as one of the aforementioned phosphines and the like. In an embodiment, the heteroaryl quaternary ammonium salt includes a heteroaryl group instead of the aryl group in the aryl quaternary ammonium salt here. Exemplary heteroaryl quaternary ammonium salts include pyridyl(methyl)ammonium chloride (bromide, iodide), chinolinyl(methyl)ammonium chloride (bromide, iodide), benzothiophenyl(methyl)ammonium chloride (bromide, iodide), and the like.

[0042] Exemplary aryl quaternary phosphonium salts include benzyltriphenylphosphonium chloride, dimethyldiphenylphosphonium iodide, ethyltriphenylphosphonium iodide, methyltriphenoxyposphonium iodide, tetraphenylphosphonium bromide, and the like. additional quaternary phosphonium salts

includes those derived from commercially available (Sigma Aldrich co.) phosphines such as substituted or unsubstituted triphenylphosphine, naphthylidiphenylphosphine, dinaphthylphenylphosphine, trinaphthylphosphine, 9-anthryldiphenylphosphine, 9-anthryldinaphthylphosphine, diphenylpyrenylphosphine, dinaphthylpyrenylphosphine, bis(pentafluorophenyl)phenylphosphine, (4-bromophenyl)diphenylphosphine, 4-(dimethylamino)phenyldiphenylphosphine, diphenyl(2-methoxyphenyl)phosphine, diphenyl(pentafluorophenyl)phosphine, 2-(diphenylphosphino)benzaldehyde, diphenyl-2-pyridylphosphine, diphenyl(p-tolyl)phosphine, tri-2-furylphosphine, tris(4-chlorophenyl)phosphine, tris(2,6-dimethoxyphenyl)phosphine, tris(4-fluorophenyl)phosphine, tris(3-methoxyphenyl)phosphine, tris(4-methoxyphenyl)phosphine, tris(pentafluoro-phenyl)phosphine, tris(2,4,6-trimethoxyphenyl)phosphine, tris(2,4,6-trimethylphenyl)phosphine, 2-(diphenylphosphino)benzoic acid, 4-(diphenylphosphino)benzoic acid, 4,4'-(phenyl-phosphinidene)bis(benzenesulfonic acid), 3,3',3''-phosphinidynetris(benzenesulfonic acid), tri-m-tolylphosphine, tri-o-tolylphosphine, tri-p-tolylphosphine, (1,2-bis(diphenyl-phosphino)benzene), (2,2'-bis(diphenylphosphino)-1,1'-binaphthyl, and the like. These phosphines can be processed to produce aryl quaternary phosphonium salts as described in U.S. Patent Application No. 10/553,307, the disclosure of which is incorporated herein in its entirety. In an embodiment, the heteroaryl quaternary phosphonium salt includes a heteroaryl group instead of the aryl group in the aryl quaternary phosphonium salts here. Exemplary heteroaryl quaternary phosphonium salts include pyridyl(methyl)phosphonium chloride (bromide, iodide), chinolinyl(methyl)phosphonium chloride (bromide, iodide), benzothiophenyl(methyl)phosphonium chloride (bromide, iodide), and the like.

[0043] The anion X^- of the aromatic compound of formula (1) (e.g., the aryl quaternary ammonium salt, heteroaryl quaternary ammonium salt, aryl quaternary phosphonium salt, or heteroaryl quaternary phosphonium salt) can be a halide, triflate, sulfate, nitrate, hydroxide, carbonate, bicarbonate, acetate, phosphate, oxalate, cyanide, alkylcarboxylate, N-hydroxysuccinimide, N-hydroxybenzotriazole, alkoxide, thioalkoxide, alkane sulfonyloxide, halogenated alkane sulfonyloxide, arylsulfonyloxide, heteroarylsulfonyloxide bisulfate, valerate, oleate, palmitate, stearate, laurate, borate, benzoate, lactate, citrate, maleate, fumarate, succinate, tartrate, naphthylate, mesylate, glucoheptonate, lactobionate, and the like.

[0044] Exemplary aryl alkoxy silanes includes tert-butoxy(chloro)diphenylsilane, aminophenyltrimethoxysilane, 2-(4-pyridylethyl)triethoxysilane, 2-

(trimethoxysilylethyl)pyridine, n-(3-trimethoxysilylpropyl)pyrrole, 3-(m-aminophenoxy)propyltrimethoxy-silane, n-phenylaminopropyltrimethoxy-silane, (phenylaminomethyl)methyl-dimethoxysilane, n-phenylaminomethyltriethoxysilane, 3-(n-styrylmethyl-2-aminoethylamino)-propyltrimethoxysilane hydrochloride, (2-n-benzylaminoethyl)-3-aminopropyl-trimethoxysilane hydrochloride, n-(3-triethoxysilylpropyl)-4,5-dihydroimidazole, 2-(2-pyridylethyl)thiopropyltrimethoxysilane, 2-(4-pyridylethyl)thiopropyltrimethoxysilane, benzoyloxypropyltrimethoxysilane, ((chloromethyl)phenylethyl)-trimethoxysilane, (p-chloromethyl)phenyltrimethoxysilane, ((chloromethyl)phenylethyl)-methyldimethoxysilane, bis(2-diphenylphosphinoethyl)-methylsilylethyltriethoxysilane, diphenylphosphinoethyl-dimethyl-ethoxysilane, 2-(diphenylphosphino)ethyl-triethoxysilane, 2-(2-pyridylethyl)thiopropyltrimethoxysilane, 2-(4-pyridylethyl)thiopropyltri-methoxysilane, 2-(3-trimethoxysilylpropylthio)-thiophene, 3-(n-styrylmethyl-2-aminoethylamino)-propyltrimethoxysilane, (3-cyclopentadienylpropyl)triethoxysilane, styrylethyltrimethoxysilane, 3-(2,4-dinitrophenylamino)propyl-triethoxysilane, 2-hydroxy-4-(3-methyldiethoxysilylpropoxy)diphenylketone, 2-hydroxy-4-(3-triethoxysilylpropoxy)-diphenylketone, o-4-methylcoumarinyl-n-[3-(triethoxy-silyl)propyl]carbamate, 7-triethoxysilylpropoxy-5-hydroxy-flavone, 5-dimethylamino-n-(3-triethoxysilylpropyl)-naphthalene-1-sulfonamide, 2-(2-triethoxysilylpropoxy-5-methyl-phenyl)benzotriazole, 3-(triethoxysilylpropyl)-p-nitrobenzamide, (R)-n-triethoxysilylpropyl-o-quinine-urethane, (R)-n-1-phenylethyl-n'-triethoxysilyl-propylurea, (S)-n-1-phenylethyl-n'-triethoxysilyl-propylurea, and the like available from Gelest Inc, Morrisville, PA. In an embodiment, the heteroaryl alkoxy silanes include a heteroaryl group instead of the aryl group in the aryl alkoxy silane compounds here. Exemplary heteroaryl alkoxy silanes include pyridylmethyltriethoxy silane, furylethyltriethoxysilane, thiophenylethyltriethoxy silane, and the like.

[0045] Exemplary aryl halides include phenyl chloride, 2-chlorotoluene, 2-bromotoluene, 4-chlorotoluene, 4-bromotoluene, 2-chloro-4-methylnaphthalene, 2-bromo-4-methylnaphthalene, 4-chloroanisole, 4-bromoanisole, 2-chlorobenzyl(2-methoxy)ethyl ether, 2-bromobenzyl(2-methoxy)ethyl ether, 2-chlorobenzyl methyl ether, 2-bromobenzyl methyl ether, 2-chlorobenzyl ethyl ether, 2-bromobenzyl ethyl ether, chlorobenzene, bromobenzene, iodobenzene, fluorobenzene, dichlorobenzene, trichlorobenzene, chlorotoluene, 2,4-dichlorotoluene, chloronaphthalene, bromonaphthalene, iodotoluenes, iodonaphthalene, 2-bromo-6-methoxynaphthalene, 4-bromo-isobutylbenzene, triphenylmethane chloride, iodobenzene, bromotoluene, iodonaphthalene, chlorobenzene, phenylphosphine dichloride,

diphenylphosphine mono chloride, (o-chlorophenyl)phosphine dichloride, bis(o-chlorophenyl)phosphine monochloride, 1-naphthylphosphine bromides, chlorotolylphosphine chlorides, dichlorotolylphosphine chlorides, and the like. In an embodiment, the heteroaryl halide includes a heteroaryl group instead of the aryl group in the aryl halide compounds here, e.g., 2,6-dichloropyridine. Exemplary heteroaryl halides include pyridylethyl chloride, furylethyl chloride, thiophenylethyl chloride, and the like.

[0046] In the composition, the functionalized silicate nanoparticle includes a silicate nanoparticle. The silicate nanoparticle contains silicon and oxygen that can be arranged in various structures such as a tetrahedral configuration and can have a shape such as a platelet, sphere, polyhedron, rod, cylinder, a combination thereof, and the like. According to an embodiment, the silicate nanoparticle comprises a silsesquioxane, cyclosilicate, inosilicate, nesosilicate, phyllosilicate, sorosilicate, tectosilicate, or a combination thereof.

[0047] The silicate nanoparticles, from which the composition is formed, are generally particles having an average particle size, in at least one dimension, of less than one micrometer (μm). As used herein “average particle size” refers to the number average particle size based on the largest linear dimension of the particle (sometimes referred to as “diameter”). Particle size, including average, maximum, and minimum particle sizes, may be determined by an appropriate method of sizing particles such as, for example, static or dynamic light scattering (SLS or DLS) using a laser light source. Silicate nanoparticles can include both particles having an average particle size of 250 nm or less, and particles having an average particle size of greater than 250 nm to less than 1 μm (sometimes referred in the art as “sub-micron sized” particles). In an embodiment, a silicate nanoparticle can have an average particle size of about 0.1 nanometers (nm) to about 500 nm, specifically 0.5 nm to 250 nm, more specifically about 1 nm to about 150 nm, more specifically about 1 nm to about 125 nm, and still more specifically about 1 nm to about 75 nm. The silicate nanoparticles may be monodisperse, where all particles are of the same size with little variation, or polydisperse, where the particles have a range of sizes and are averaged. Generally, polydisperse silicate nanoparticles are used. Silicate nanoparticles of different average particle size may be used, and in this way, the particle size distribution of the silicate nanoparticles can be unimodal (exhibiting a single size distribution), bimodal (exhibiting two size distributions), or multi-modal (exhibiting more than one particle size distribution).

[0048] The minimum particle size for the smallest 5 percent of the silicate nanoparticles can be less than 2 nm, specifically less than or equal to 1 nm, and more

specifically less than or equal to 0.5 nm. Similarly, the maximum particle size for 95% of the silicate nanoparticles can be greater than or equal to 900 nm, specifically greater than or equal to 750 nm, and more specifically greater than or equal to 500 nm. The silicate nanoparticles can have a high surface area of greater than 300 m²/g, and in a specific embodiment, 300 m²/g to 1800 m²/g, specifically 500 m²/g to 1500 m²/g. In a particular embodiment, the silsesquioxane has a size from 0.5 nm to 10 nm.

[0049] According to an embodiment, the silicate nanoparticle is a silsesquioxane. Silsesquioxanes, also referred to as polysilsesquioxanes, polyorganosilsesquioxanes, or polyhedral oligomeric silsesquioxanes (POSS), are polyorganosilicon oxide compounds of general formula RSiO_{1.5} (where R is a hydrogen, inorganic group, or organic group) having defined closed or open cage structures (closo or nido structures, which are called respectively completely condensed or incompletely condensed structures). Silsesquioxanes can be prepared by acid and/or base-catalyzed condensation of functionalized silicon-containing monomers such as tetraalkoxysilanes including tetramethoxysilane and tetraethoxysilane, alkyltrialkoxysilanes such as methyltrimethoxysilane and methyltrimethoxysilane, as well as other groups.

[0050] In an embodiment, the silsesquioxane has a closed cage structure, an open cage structure, or a combination thereof. The silsesquioxane can have any shape of cage structure such as cubes, hexagonal prisms, octagonal prisms, decagonal prisms, dodecagonal prisms, and the like. Additionally, the cage structure of the silsesquioxane comprises from 4 to 30 silicon atoms, specifically, 4 to 20 silicon atoms, and more specifically 4 to 16 silicon atoms, with each silicon atom in the cage structure being bonded to oxygen. It should be noted that the term “cage structure” is meant to include the SiO_{1.5} portion of the general silsesquioxane formula RSiO_{1.5}, and not the R-group.

[0051] According to an embodiment, the silsesquioxane comprises a functional group bonded to a silicon atom of the silsesquioxane. In a specific embodiment, the functional group bonded to the silicon atom comprises an alkyl, alkoxy, haloalkyl, cycloalkyl, heterocycloalkyl, cycloalkyloxy, aryl, aralkyl, aryloxy, aralkyloxy, heteroaryl, heteroaralkyl, alkenyl, alkynyl, amine, alkyleneamine, aryleneamine, alkenyleneamine, hydroxy, carboxyl, ether, epoxy, ketone, halogen, hydrogen, or a combination thereof. Thus, the silsesquioxane derivatized with a functional group includes a group such as an alcohol, amine, carboxylic acid, epoxy, ether, fluoroalkyl, halide, imide, ketone, methacrylate, acrylate, silica, nitrile, norbornenyl, olefin, polyethylene glycol (PEG), silane, silanol, sulfonate, thiol, and the like. Furthermore, the silsesquioxane can have from one functional group to as many functional

groups as there are silicon atoms in the cage structure of the silsesquioxane. In a specific embodiment, the silsesquioxane is a derivatized octasilsesquioxane $R_{8-n}H_n(SiO_{1.5})_8$ (where $0 \leq n \leq 8$, and R can be a same or different functional group), and the number of functional groups varies with the number of silicon atoms in the cage structure, i.e., from 0 to 8 functional groups.

[0052] Exemplary silsesquioxanes having a closed cage structure include 1-allyl-3,5,7,9,11,13,15-heptacyclopentylpentacyclo[9.5.1.1(3,9).1(5,15).1(7,13)]octasiloxane; 1-allyl-3,5,7,9,11,13,15-heptaisobutylpentacyclo[9.5.1.1(3,9).1(5,15).1(7,13)]octasiloxane; 1-[3-(2-aminoethyl)amino]propyl-3,5,7,9,11,13,15-isobutylpentacyclo[9.5.1.1(3,9).1(5,15).1(7,13)]octasiloxane; 1-chlorobenzylethyl-3,5,7,9,11,13,15-heptaisobutylpentacyclo[9.5.1.1(3,9).1(5,15).1(7,13)]octasiloxane; 1-(4-chlorobenzyl)-3,5,7,9,11,13,15-heptacyclopentylpentacyclo[9.5.1.1(3,9).1(5,15).1(7,13)]octasiloxane; 1-chloropropyl-3,5,7,9,11,13,15-isobutylpentacyclo[9.5.1.1(3,9).1(5,15).1(7,13)]octasiloxane; (cyanopropyldimethylsilyloxy)heptacyclopentylpentacyclooctasiloxane; 1-(2-trans-cyclohexanediol)ethyl-3,5,7,9,11,13,15-isobutylpentacyclo[9.5.1.1(3,9).1(5,15).1(7,13)]octasiloxane; 1-(3-cyclohexen-1-yl)-3,5,7,9,11,13,15-heptacyclopentylpentacyclo[9.5.1.1(3,9).1(5,15).1(7,13)]octasiloxane; dodecaphenyl-dodecasiloxane; 1-[2-(3,4-epoxycyclohexyl)ethyl]-3,5,7,9,11,13,15-isobutylpentacyclo[9.5.1.1(3,9).1(5,15).1(7,13)]octasiloxane; 1,3,5,7,9,11,13-heptacyclopentyl-15-glycidylpentacyclo[9.5.1.1(3,9).1(5,15).1(7,13)]octasiloxane; 1-(3-glycidyl)propoxy-3,5,7,9,11,13,15-isobutylpentacyclo[9.5.1.1(3,9).1(5,15).1(7,13)]octasiloxane; octakis(tetramethylammonium)pentacyclo[9.5.1.1(3,9).1(5,15).1(7,13)]octasiloxane-1,3,5,7,9,11,13,15-octakis(yloxide)hydrate; 3-hydroxypropylheptaisobutyl-octasiloxane; 1-(3-mercapto)propyl-3,5,7,9,11,13,15-isobutylpentacyclo[9.5.1.1(3,9).1(5,15).1(7,13)]octasiloxane; octacyclohexenylethyldimethylsilyloxy-octasiloxane; 1,3,5,7,9,11,13,15-octacyclohexylpentacyclooctasiloxane; octa[(1,2-epoxy-4-ethylcyclohexyl)dimethylsiloxy]octasiloxane; octa[(3-glycidyloxypropyl)dimethylsiloxy]octasiloxane; octa[(3-hydroxypropyl)dimethylsiloxy]octasiloxane; 1,3,5,7,9,11,13,15-octakis[2-(chlorodimethylsilyl)ethyl]pentacyclo[9.5.1.1(3,9).1(5,15).1(7,13)]octasiloxane; 1,3,5,7,9,11,13,15-

octakis(dimethylsilyloxy)pentacyclo[9.5.1.1(3,9).1(5,15).1(7,13)]octasiloxane;
 1,3,5,7,9,11,13,15-octamethylpentacyclo[9.5.1.1(3,9).1(5,15).1(7,13)]octasiloxane;
 1,3,5,7,9,11,13,15-octaphenylpentacyclo[9.5.1.1(3,9).1(5,15).1(7,13)]octasiloxane;
 1,3,5,7,9,11,13,15-octa(2-trichlorosilyl)ethylpentacyclo[9.5.1.1(3,9).1(5,15).1(7,13)]octasiloxane; 1,3,5,7,9,11,13,15-octavinylpentacyclo[9.5.1.1(3,9).1(5,15).1(7,13)]octasiloxane; 1-(2,3-propanediol)propoxy-3,5,7,9,11,13,15-isobutylpentacyclo[9.5.1.1(3,9).1(5,15).1(7,13)]octasiloxane; 3-(3,5,7,9,11,13,15-heptaisobutylpentacyclo[9.5.1.1(3,9).1(5,15).1(7,13)]octasiloxan-1-yl)propylmethacrylate; (3-tosyloxypropyl)-heptaisobutyl-octasiloxane; 1-(trivinylsilyloxy)-3,5,7,9,11,13,15-heptacyclopentylpentacyclo[9.5.1.1(3,9).1(5,15).1(7,13)]octasiloxane; 1-vinyl-3,5,7,9,11,13,15-isobutylpentacyclo[9.5.1.1(3,9).1(5,15).1(7,13)]octasiloxane, (3-(2,2-bis(hydroxymethyl)butoxy)propyl)dimethylsiloxy-3,5,7,9,11,13,15-isobutylpentacyclo[9.5.1.1(3,9).1(5,15).1(7,13)]octasiloxane; octa(3-hydroxy-3-methylbutyl)dimethylsiloxy)octasiloxane; 1-(3-amino)propyl-3,5,7,9,11,13,15-isobutylpentacyclo[9.5.1.1(3,9).1(5,15).1(7,13)]octasiloxane; 1-(3-amino)propyl-3,5,7,9,11,13,15-isooctylpentacyclo[9.5.1.1(3,9).1(5,15).1(7,13)]octasiloxane; 1,3,5,7,9,11,13,15-octaaminophenylpentacyclo[9.5.1(3,9).1(5,15).1(7,13)]octasiloxane; octa-n-phenylaminopropyl)-octasiloxane; n-methylaminopropyl-heptaisobutyl-octasiloxane; octaethylammoniumoctasiloxane chloride; 1-(4-amino)phenyl-3,5,7,9,11,13,15-cyclohexylpentacyclo[9.5.1.1(3,9).1(5,15).1(7,13)]octasiloxane; 1-(amino)phenyl-3,5,7,9,11,13,15-cyclohexylpentacyclo[9.5.1.1(3,9).1(5,15).1(7,13)]octasiloxane; 1-(4-amino)phenyl-3,5,7,9,11,13,15-heptaisobutylpentacyclo[9.5.1.1(3,9).1(5,15).1(7,13)]octasiloxane; 1-(amino)phenyl-3,5,7,9,11,13,15-heptaisobutylpentacyclo[9.5.1.1(3,9).1(5,15).1(7,13)]octasiloxane; 1-[(3-maleamic acid)propyl]-3,5,7,9,11,13,15-heptacyclohexylpentacyclo[9.5.1.1(3,9).1(5,15).1(7,13)]-octasiloxane; 1-[(3-maleamic acid)propyl]-3,5,7,9,11,13,15-heptaisobutylpentacyclo[9.5.1.1(3,9).1(5,15).1(7,13)]-octasiloxane; octamaleamic acid octasiloxane; trimethoxy-[2-(7-oxabicyclo[4.1.0]hept-3-yl)ethyl]silane, hydrolyzed; 2-[[3-(trimethoxysilyl)propoxy]methyl]-oxirane, hydrolyzed; ethyl 3,5,7,9,11,13,15-heptaethylpentacyclo[9.5.1.1(3,9).1(5,15).1(7,13)]octasiloxane-1-undecanoate; 1-(3-glycidyl)propoxy-3,5,7,9,11,13,15-isooctylpentacyclo[9.5.1.1(3,9).1(5,15).1(7,13)]octasiloxane; 3,7,14-tris{[3-(epoxypropoxy)propyl]dimethylsilyloxy}-1,3,5,7,9,11,14-heptacyclohexyltricyclo[7.3.3.1(5,11)]heptasiloxane; 3,7,14-tris{[3-

(epoxypropoxy)propyl]dimethylsilyloxy}-1,3,5,7,9,11,14-heptaisobutyltricyclo[7.3.3.1(5,11)]heptasiloxane; octafluoropropyl-octasiloxane; endo-3,7,14-trifluoropropyl-1,3,5,7,9,11,14-heptaisobutyltricyclo[7.3.3.1(5,11)]heptasiloxane; 1-chlorobenzyl-3,5,7,9,11,13,15-heptaisobutylpentacyclo[9.5.1.1(3,9).1(5,15).1(7,13)]octasiloxane; 1,3,5,7,9,11,13,15-octakis(1,2-dibromoethyl)-pentacyclo[9.5.1.1(3,9).1(5,15).1(7,13)]octasiloxane; 1-[(3-maleimide)propyl]-3,5,7,9,11,13,15-heptacyclohexylpentacyclo[9.5.1.1(3,9).1(5,15).1(7,13)]octasiloxane; 1-[(3-maleimide)propyl]-3,5,7,9,11,13,15-heptaisobutylpentacyclo[9.5.1.1(3,9).1(5,15).1(7,13)]octasiloxane; 3-(3,5,7,9,11,13,15-heptaisobutylpentacyclo[9.5.1.1(3,9).1(5,15).1(7,13)]octasiloxan-1-yl)propylacrylate; 3-[3,5,7,9,11,13,15-heptacyclohexylpentacyclo[9.5.1.1(3,9).1(5,15).1(7,13)]octasiloxan-1-yl]methylmethacrylate; 3-[3,5,7,9,11,13,15-heptaisobutylpentacyclo[9.5.1.1(3,9).1(5,15).1(7,13)]octasiloxan-1-yl]methylmethacrylate; 3-[3,5,7,9,11,13,15-heptaethylpentacyclo[9.5.1.1(3,9).1(5,15).1(7,13)]octasiloxan-1-yl]methylmethacrylate; 3-[3,5,7,9,11,13,15-heptaethylpentacyclo[9.5.1.1(3,9).1(5,15).1(7,13)]octasiloxan-1-yl]propylmethacrylate; 3-[3,5,7,9,11,13,15-heptaisooctylpentacyclo[9.5.1.1(3,9).1(5,15).1(7,13)]octasiloxan-1-yl]methylmethacrylate; 3-(3,5,7,9,11,13,15-heptaisooctylpentacyclo[9.5.1.1(3,9).1(5,15).1(7,13)]octasiloxan-1-yl)propylmethacrylate; 3-(3,5,7,9,11,13,15-heptaphenylpentacyclo[9.5.1.1(3,9).1(5,15).1(7,13)]octasiloxan-1-yl)propylmethacrylate; octasiloxa-octapropylmethacrylate; octasiloxa-octapropylacrylate; dodecaphenyldecasiloxane; octaisooctyloctasiloxane; phenylheptaisobutyloctasiloxane; phenylheptaisooctyloctasiloxane; isooctylhetpaphenyloctasiloxane; octaisobutyloctasiloxane; octamethyloctasiloxane; octaphenyloctasiloxane; octakis(tetramethylammonium)pentacyclo[9.5.1.1(3,9).1(5,15).1(7,13)]octasiloxane 1,3,5,7,9,11,13,15-octakis(cyloxy)hydrate; octakis(trimethylsiloxy)pentacyclo[9.5.1.1(3,9).1(5,15).1(7,13)]octasiloxane; 3,5,7,9,11,13,15-heptaisobutylpentacyclo[9.5.1.1(3,9).1(5,15).1(7,13)]octasiloxane-1-butyronitrile; 1-[2-(5-norbornen-2-yl)ethyl]-3,5,7,9,11,13,15-heptaethylpentacyclo[9.5.1.1(3,9).1(7,13)]octasiloxane; 1-[2-(5-norbornen-2-yl)ethyl]-3,5,7,9,11,13,15-heptaisobutylpentacyclo[9.5.1.1(3,9).1(7,13)]octasiloxane; 1-allyl-3,5,7,9,11,13,15-heptaisobutylpentacyclo[9.5.1.1(3,9).1(7,13)]octasiloxane; 1,3,5,7,9,11,13-heptaisobutyl-15-vinylpentacyclo[9.5.1.1(3,9).1(5,15).1(7,13)]octasiloxane; 1,3,5,7,9,11,13,15-octa[2-(3-

cyclohexenyl)ethyldimethylsiloxy]pentacyclo[9.5.1.1(3,9).1(5,15).1(7,13)]octasiloxane;
 1,3,5,7,9,11,13,15-octavinylpentacyclo[9.5.1.1(3,9).1(5,15).1(7,13)]octasiloxane;
 1,3,5,7,9,11,13,15-
 octa[vinyldimethylsiloxy]pentacyclo[9.5.1.1(3,9).1(5,15).1(7,13)]octasiloxane;
 1,3,5,7,9,11,13,15-
 octakis(dimethylsilyloxy)pentacyclo[9.5.1.1(3,9).1(5,15).1(7,13)]octasiloxane;
 1,3,5,7,9,11,13,15-octahydropentacyclo[9.5.1.1(3,9).1(5,15).1(7,13)]octasiloxane; 1-(3-
 mercapto)propyl-3,5,7,9,11,13,15-
 isobutylpentacyclo[9.5.1.1(3,9).1(5,15).1(7,13)]octasiloxane; 1-(3-mercapto)propyl-
 3,5,7,9,11,13,15-isooctylpentacyclo[9.5.1.1(3,9).1(5,15).1(7,13)]octasiloxane; and the like.

[0053] Exemplary silsesquioxanes having an open cage structure include
 1,3,5,7,9,11,14-heptacyclohexyltricyclo[7.3.3.1(5,11)]heptasiloxane-endo-3,7,14-triol;
 1,3,5,7,9,11,14-heptacyclopentyltricyclo[7.3.3.1(5,11)]heptasiloxane-endo-3,7,14-triol;
 1,3,5,7,9,11-octaisobutyltetracyclo[7.3.3.1(5,11)]octasiloxane-endo-3,7-diol; 1,3,5,7,9,11,14-
 heptaethyltricyclo[7.3.3.1(5,11)]heptasiloxane-endo-3,7,14-triol; 1,3,5,7,9,11,14-
 heptaisobutyltricyclo[7.3.3.1(5,11)]heptasiloxane-endo-3,7,14-triol; 1,3,5,7,9,11,14-
 heptaisooctyltricyclo[7.3.3.1(5,11)]heptasiloxane-endo-3,7,14-triol; 1,3,5,7,9,11,14-
 heptaphenyltricyclo[7.3.3.1(5,11)]heptasiloxane-endo-3,7,14-triol;
 tricyclo[7.3.3.3(3,7)]octasiloxane-5,11,14,17-tetraol-1,3,5,7,9,11,14,17-octaphenyl; 9-
 {dimethyl[2-(5-norbornen-2-yl)ethyl]silyloxy}-1,3,5,7,9,11,14-heptaisobutyltricyclo[7.3.3.1
 5,11]heptasiloxane-1,5-diol; endo-3,7,14-tris{dimethyl[2-(5-norbornen-2-yl)ethyl]silyloxy}-
 1,3,5,7,9,11,14-heptaisobutyltricyclo[7.3.3.1(5,11)]heptasiloxane;
 [[dimethyl(trifluoromethyl)ethyl]silyloxy]heptacyclopentyltricycloheptasiloxanediol;
 1,3,5,7,9,11,14-heptacyclohexyltricyclo[7.3.3.1(5,11)]heptasiloxane-3,7,14-triol;
 1,3,5,7,9,11,14-heptaisobutyltricyclo[7.3.3.1(5,11)]heptasiloxane-endo-3,7,14-triol;
 1,3,5,7,9,11-octacyclopentyltetracyclo[7.3.3.1(5,11)]octasiloxane-endo-3,7-diol;
 1,3,5,7,9,11,14-hepta-isooctyltricyclo[7.3.3.1(5,11)]heptasiloxane-endo-3,7,14-triol; *endo*-
 3,7,14-trifluoro-1,3,5,7,9,11,14-heptacyclopentyltricyclo[7.3.3.1(5,11)]heptasiloxane; *endo*-
 3,7,14-tris{dimethyl[2-(5-norbornen-2-yl)ethyl]silyloxy}-1,3,5,7,9,11,14-
 heptacyclopentyltricyclo[7.3.3.1(5,11)]heptasiloxane;
 tris((dimethyl(trifluoromethyl)ethyl)silyloxy)heptacyclopentyltricycloheptasiloxane; 3,7,14-
 tris{[3-(epoxypropoxy)propyl]dimethylsilyloxy}-1,3,5,7,9,11,14-
 heptacyclopentyltricyclo[7.3.3.1(5,11)]heptasiloxane, and the like.

[0054] A combination of the silsesquioxanes with an open cage structure or closed cage structure can be used as the silsesquioxane in conjunction with a combination of any of the other silicate nanoparticles.

[0055] In an embodiment, the silicate nanoparticle is a silicate mineral such as cyclosilicate, inosilicate, nesosilicate, phyllosilicate, sorosilicate, tectosilicate, or a combination thereof.

[0056] Cyclosilicates are silicates with tetrahedrons that can link to form rings of three $(\text{Si}_3\text{O}_9)^{-6}$, four $(\text{Si}_4\text{O}_{12})^{-8}$, six $(\text{Si}_6\text{O}_{18})^{-12}$ or nine $(\text{Si}_9\text{O}_{27})^{-18}$ units. Exemplary cyclosilicates include benitoite, axinite, beryl, cordierite, tourmaline, papagoite, eudialyte, milarite, and the like.

[0057] The inosilicate can have a crystalline structure in the form of a chain such as pyroxenes and pyroxenoids (with a crystalline structure of single chains $(\text{SiO}_3)^{-2}$) or amphiboles (with a crystalline structure of double chains $(\text{Si}_4\text{O}_{11})^{-6}$). Non-limiting examples of pyroxenes and pyroxenoids include diopside, spodumene, wollastonite, enstatite, hypersthene, hedenbergite, augite, pectolite, diallage, fassaite, spodumene, jeffersonite, aegirine, omphacite, hiddenite, and the like. Non-limiting examples of amphiboles are calcium amphiboles such as tremolite, actinote, and hornblende; iron-magnesium amphiboles such as grunerite and cummingtonite; and sodium amphiboles such as glaucophane, arfvedsonite and riebeckite; and the like.

[0058] Non-limiting examples of nesosilicates are alite, almandine, andalousite, andalusite, andradite, belite, chloritoid, chondrodite, clinohumite, datolite, dumortierite, fayalite, forsterite, grossular, humite, hydrogrossular, kyanite, norbergite, olivine, phenakite, pyrope, sillimanite, spessartine, staurolite, thaumasite, thorite, titanite, topaz, uvarovite, zircon, and the like.

[0059] The phyllosilicate can be a clay, mica, serpentine, chlorite, or a combination thereof. Exemplary phyllosilicates include antigorite, biotite, chlorite, chrysotile, glauconite, halloysite, illite, kaolinite, lepidolite, lizardite, margarite, montmorillonite, muscovite, palygorskite, phlogopite, pyrophyllite, talc, vermiculite, and the like.

[0060] The sorosilicate can be allanite, clinozoisite, dollaseite, epidote, hemimorphite, ilvaite, lawsonite, prehnite, tanzanite, vesuvianite, zoisite, and the like.

[0061] The tectosilicate can be, for example, albite, alkali-feldspars, analcime, andesine, anorthite, anorthoclase, bytownite, cancrinite, celsiane, chabazite, coesite, cristobalite, feldspar, feldspathoid, hauyne, heulandite, labradorite, lazurite, leucite, marialite, meionite, microcline, mordenite, natrolite, nepheline, nosean, oligoclase, orthoclase, petalite,

plagioclase, quartz, sanidine, scapolite, scolecite, silica, sodalite, stilbite, tridymite, zeolite, and the like.

[0062] Exemplary zeolites include naturally occurring zeolites such as amicitite, analcime, barrerite, bellbergite, bikitaite, boggsite, brewsterite, chabazite, clinoptilolite, cowlesite, dachiardite, edingtonite, epistilbite, erionite, faujasite, ferrierite, garronite, gismondine, gmelinite, gobbinsite, gonnardite, goosecreekite, harmotome, herschelite, heulandite, laumontite, levyne, maricopaite, mazzite, merlinoite, mesolite, montesommaite, mordenite, natrolite, offretite, paranatrolite, paulingite, pentasil, perialite, phillipsite, pollucite, scolecite, sodium dachiardite, stellerite, stilbite, tetranatrolite, thomsonite, tschernichite, wairakite, wellsite, willhendersonite, and yugawaralite. In some embodiments, the zeolite is analcime, chabazite, clinoptilolite, heulandite, natrolite, phillipsite, stilbite, or a combination thereof. A synthetic zeolite also can be used as the tectosilicate of the silicate nanoparticle. The synthetic zeolites can be selected from Zeolite A, Zeolite B, Zeolite F, Zeolite H, Zeolite L, Zeolite T, Zeolite W, Zeolite X, Zeolite Y, Zeolite Omega, Zeolite ZSM-5, Zeolite ZSM-4, Zeolite P, Zeolite N, Zeolite D, Zeolite O, Zeolite S, and Zeolite Z.

[0063] In an embodiment, the silicate nanoparticle can include other elements or components in addition to silicon and oxygen. The silicate nanoparticle can include an oxide, for example, silicon dioxide (SiO_2), aluminum oxide (Al_2O_3), barium oxide (BaO), bismuth trioxide (Bi_2O_3), boron oxide (B_2O_3), calcium oxide (CaO), cesium oxide (CsO), lead oxide (PbO), strontium oxide (SrO), rare earth oxides (e.g., lanthanum oxide (La_2O_3), neodymium oxide (Nd_2O_3), samarium oxide (Sm_2O_3), cerium oxide (CeO_2)), and the like. An exemplary silicate nanoparticle containing SiO_2 includes quartz, cristobalite, tridymite, and the like. The other elements can be, for example, aluminum, antimony, arsenic, barium, beryllium, boron, calcium, cerium, cesium, chromium, cobalt, copper, gallium, gold, iron, lanthanum, lead, lithium, magnesium, manganese, molybdenum, neodymium, nickel, niobium, palladium, phosphorus, platinum, potassium, praseodymium, silver, sodium, tantalum, thorium, titanium, vanadium, zinc, zirconium, and the like. The other elements can occur in the silicate nanoparticle in the form of oxides, carbonates, nitrates, phosphates, sulfates, or halides. Furthermore, the other element can be a dopant in the silicate nanoparticle.

[0064] It is contemplated that the silicate nanoparticle is functionalized with the functionalization compound or a chemical group from the functionalization compound. Functionalization of the silicate nanoparticle to form the functionalized silicate nanoparticle can be achieved by a cation exchange reaction, substitution, condensation, alkoxysilane chemistry, and the like. Without wishing to be bound by the theory, the silicate nanoparticle

can have a group such as a hydroxy group on its surface that interacts with and can react with the functionalization compound or chemical group thereof. As used herein, the bond of the first portion of the chemical group includes covalent bonds as well as ionic bonds.

[0065] As noted above, the functionalization compound includes a chemical group that has a first portion and a second portion, which includes an aromatic moiety or a nonaromatic moiety. The first portion is directly bonded to the silicate nanoparticle in the functionalized silicate nanoparticle. In terms of formulas (1), (2), and (3), the second portion includes the aromatic or nonaromatic moiety Ar. Also, the first portion can include the A group (e.g., nitrogen or phosphorous), Si, oxygen (-O-), or linker group L in formulas (1), (2), and (3). As used herein, the bond of the first portion of the chemical group includes covalent bonds as well as ionic bonds.

[0066] The compounds of formulas (4)-(7) can be attached to the silicate nanoparticles through an ionic bond via "N" or "S." In this instant, the first portion of the chemical group which functionalizes the silicate nanoparticle is an ionic bond and the second portion of the chemical group is a cation of formula (4), formula (5), formula (6) or formula (7). The compounds of formula (8) can be attached to the silicate nanoparticle through Si when R₂₅ is halide or when R₂₅ is an alkoxy group.

[0067] In an embodiment, the functionalization compound is a compound of formula (2) in which the central silicon atom is bonded to the silicate nanoparticle. In an embodiment, the aromatic compound of formula (1) is bonded to the silicate nanoparticle via directly bonding the A group to the silicate nanoparticle.

[0068] In another embodiment, the chemical group is derived from a compound of formula (1), (2), (3), (4), (5), (6), (7), or (8). In a specific embodiment, the chemical group is derived from a quaternary ammonium salt, quaternary phosphonium salt, alkoxy silane, halide, a guanidine, a guanidinium salt, a biguanidinium salt, or a sulfonium salt. Here, the chemical group can be derived from the functionalization compound (e.g., by hydrolysis, photolytic cleavage, thermal decomposition, elimination, substitution, etc.) to produce the chemical group including the aromatic or nonaromatic moiety Ar with or without the linker group L, such that the chemical group is bonded to the silicate nanoparticle. In formula (1), the bond between the A group and linker group, moiety Ar, or R group is broken. In formula (2), the bond between the Si atom and oxygen atom (-O-), linker group L, moiety Ar, or R group is broken. In formula (3), the bond between the halogen X and linker group or moiety Ar is broken. Thus, in an embodiment, the moiety Ar is directly bonded to the silicate nanoparticle. Alternatively, the linker group L can remain attached to the moiety Ar such

that the linker group L is directly bonded to the silicate nanoparticle with the moiety Ar indirectly bonded to the silicate nanoparticle via the linker group L. In an embodiment, the R group is directly bonded to the silicate nanoparticle. Consequently, it is contemplated that the linker group L can be disposed between the silicate nanoparticle and the second portion of the chemical group in the functionalized silicate nanoparticle. Thus, the functionalized silicate nanoparticle can be a reaction product of the silicate nanoparticle with the aromatic or nonaromatic compound, chemical group derived from the aromatic or nonaromatic compound, or a combination thereof.

[0069] In an embodiment, the functionalized silicate nanoparticle can be prepared by contacting the silicate nanoparticle with the functionalization compound or chemical group thereof under conditions effective to functionalize the silicate nanoparticle. The chemical group can be prepared by subjecting the functionalization compound to conditions effective to break bonds within the functionalization compound with a product fragment being the chemical group comprising the first and second portions herein. Conditions include those of temperature, pressure, catalysis (e.g., acid catalysis, metal catalysis, support (e.g., zeolite) promotion, and the like), and the like. After contact of the silicate nanoparticle with the aromatic compound or chemical group, the aromatic compound or chemical group is bonded to the silicate nanoparticle to form the functionalized silicate nanoparticle.

[0070] In a particular embodiment, an aryl alkoxy silane (e.g., trimethyl(2-phenylethoxy)silane) contacts a silicate nanoparticle clay (e.g., montmorillonite), and the silicon of the trimethyl(2-phenylethoxy)silane bonds to the montmorillonite to form a functionalized silicate nanoparticle, having a phenyl ring extending from the surface of the montmorillonite to form a phenyl terminated functionalization. In another embodiment, the trimethyl(2-phenylethoxy)silane is subjected to a lysis condition to produce an aromatic moiety of a (2-phenylethyl)oxidanyl radical, which subsequently bonds to the montmorillonite to form the functionalized silicate nanoparticle, having a phenyl ring extending from the surface of the montmorillonite to form a phenyl terminated functionalization. Here, the linker group is $-\text{CH}_2\text{O}-$, which is directly bonded to the silicate nanoparticle via oxygen (O) and directly bonded to the phenyl ring by the methylene ($-\text{CH}_2-$).

[0071] In another embodiment, a heteroaryl quaternary ammonium salt (e.g., N,N-dimethyl-N-(pyridine-3-ylmethyl)ethanaminium chloride (DMPME)) contacts a silicate nanoparticle silsesquioxane (e.g., 1-allyl-3,5,7,9,11,13,15-heptacyclopentylpentacyclo[9.5.1.1(3,9).1(5,15).1(7,13)]octasiloxane). The resulting functionalized silicate nanoparticle includes the silsesquioxane directly bonded to the amino

nitrogen of the DMPME with the pyridinyl ring extending (via the methylene amino (-CH₂N-) linker group) from the surface of the DMPME to form a pyridinyl terminated functionalization.

[0072] The composition is effective to remove an asphaltene particle from a substrate such as a metal, composite, sand, rock, mineral, glass, formation, downhole element, or a combination thereof. Without wishing to be bound by theory, it is believed that the functionalized silicate nanoparticle is an amphiphile (i.e., having hydrophilic and lipophilic portions) and interacts with both the substrate and the asphaltene particle. It is contemplated that, within the functionalized silicate nanoparticle, the silicate nanoparticle is hydrophilic with a greater affinity for the substrate than the asphaltene particle, and the aromatic or nonaromatic functionalization (either aryl or heteroaryl terminated functionalization) is lipophilic with a greater affinity for the asphaltene particle than the substrate. Moreover, the asphaltene particle has a greater affinity for the aromatic or nonaromatic functionalization of the functionalized silicate nanoparticle than the silicate nanoparticle portion of the functionalized silicate nanoparticle or the substrate.

[0073] In an embodiment, in addition to the functionalized silicate nanoparticle, the composition also includes a fluid. The fluid can be present in an amount to increase a separation and aid removal of the asphaltene particle from the substrate. That is, while the functionalized silicate nanoparticle separates or removes the asphaltene particle from the substrate, the fluid can sweep the removed asphaltene particle (which has been desorbed from the substrate) away from a location proximate to the substrate. In an embodiment, the fluid can also aid in separating the asphaltene particle from the substrate. In an embodiment, the asphaltene particle has a greater affinity for the fluid than the substrate or the silicate nanoparticle portion of the functionalized silicate nanoparticle.

[0074] Exemplary fluids include water (liquid or steam), oil, carbon dioxide, C1-C6 alkane (e.g., gaseous or liquefied at a temperature or pressure of the surrounding environment of the composition), tetrahydrofuran, 1,4-dioxane; diglyme, triglyme, acetonitrile, propionitrile, benzonitrile, N,N-dimethylformamide, N,N-dimethylacetamide, dimethylsulfoxide, nitrobenzene, sulfolane, acetone, butanone, cyclohexanone, diethyl ketone, methyl isobutyl ketone, 2-pentanone, 2-hexanone, 2-heptanone, 3-heptanone, 4-heptanone, 3-pentanone, brine, completion fluid, acid, base, gas, polar solvent, nonpolar solvent, or a combination thereof. Additional exemplary fluids also include those typically encountered downhole, such as hydrocarbons, solvents, or an aqueous environment that includes formation water, seawater, salt (i.e., brine, including formates and inorganic salts,

e.g., NaCl, KCl, CaCl₂, MgCl₂, CaBr₂, ZnBr₂, NaBr, and the like), completion brine, stimulation treatment fluid, remedial cleanup treatment fluid, acidic or corrosive agent such as hydrogen sulfide, hydrochloric acid, or other such corrosive agents, or a combination thereof. Solvents include an inorganic solvent, organic solvent, or a combination thereof. Exemplary solvents include water, alcohols (e.g., methanol, ethanol, and the like), polyhydric alcohols (e.g., diethylene glycol, dipropylene glycol, 1,2-propanediol, 1,4-butanediol, 1,3-butanediol, glycerol, 1,5-pentanediol, 2-ethyl-1-hexanol, and the like), ketones (e.g., acetophenone, methyl-2-hexanone, and the like), ethers (e.g., ethylene glycol monobutyl ether, triethylene glycol monomethyl ether, and the like), carboxylic acid esters (e.g., [2,2-butoxy(ethoxy)]ethyl acetate and the like), esters of carbonic acid (e.g., propylene carbonate and the like), inorganic acids (e.g., hydrofluoric acid, hydrochloric acid, phosphoric acid, sulfuric acid, nitric acid, and the like), organic acids (e.g., those having an C1-C10 alkyl chain, which is a straight or branched chain and can be substituted), or a combination thereof.

[0075] The brine can be, for example, seawater, produced water, completion brine, or a combination thereof. The properties of the brine can depend on the identity and components of the brine. Seawater, as an example, contains numerous constituents such as sulfate, bromine, and trace metals, beyond typical halide-containing salts. On the other hand, produced water can be water extracted from a production reservoir (e.g., hydrocarbon reservoir), produced from the ground. Produced water is also referred to as reservoir brine and often contains many components such as barium, strontium, and heavy metals. In addition to the naturally occurring brines (seawater and produced water), completion brine can be synthesized from fresh water by addition of various salts such as NaCl, CaCl₂, or KCl to increase the density of the brine, such as 10.6 pounds per gallon of CaCl₂ brine. Completion brines typically provide a hydrostatic pressure optimized to counter the reservoir pressures downhole. The above brines can be modified to include an additional salt. In an embodiment, the additional salt included in the brine is NaCl, KCl, NaBr, MgCl₂, CaCl₂, CaBr₂, ZnBr₂, NH₄Cl, sodium formate, cesium formate, and the like. The salt can be present in the brine in an amount from about 0.5 wt.% to about 50 wt.%, specifically about 1 wt.% to about 40 wt.%, and more specifically about 1 wt.% to about 25 wt.%, based on the weight of the composition.

[0076] The density, polarity, hydrophilicity, lipophilicity, and the like of the fluid can be achieved by selection of the foregoing fluids. The selection of the fluid can depend on, for example, a desired density for the composition. In an embodiment, fluid is present in the composition in an amount from about 1 weight percent (wt.%) to about 99 wt.%, specifically

about 10 wt.% to about 90 wt.%, and more specifically about 20 wt.% to about 80 wt.%, based on the weight of the composition.

[0077] The composition can be prepared by combining the fluid with the functionalized silicate nanoparticle. In an embodiment, the fluid is combined with the functionalization compound and silicate nanoparticle with subsequent formation of the functionalized silicate nanoparticle. According to an embodiment, the functionalized silicate nanoparticle can contact an asphaltene particle prior to addition of the fluid. In an embodiment, the silicate nanoparticle is contacted with the chemical group (which includes the aromatic moiety) derived from the functionalization compound, such as by breaking a bond between the chemical group and the rest of the functionalization compound. Consequently, it will be appreciated that the functionalized silicate nanoparticle is a reaction product of the silicate nanoparticle and the functionalization compound or chemical group thereof.

[0078] In an embodiment, a method for making the functionalized silicate nanoparticle includes contacting the silicate nanoparticle with a chemical group to form the functionalized silicate nanoparticle. The chemical group includes a first portion and a second portion comprising an aromatic or nonaromatic moiety. In forming the functionalized silicate nanoparticle, the first portion is directly bonded to the silicate nanoparticle in the functionalized silicate nanoparticle. The aryl or nonaromatic moiety extends from the surface of the functionalized silicate nanoparticle by the first portion.

[0079] The functionalized silicate nanoparticle herein has many uses and beneficial properties. In an embodiment, the functionalized silicate nanoparticle is effective to remove an asphaltene particle or other aromatic species from a substrate comprising a metal, composite, sand, rock, mineral, glass, formation, downhole element, or a combination thereof.

[0080] According to an embodiment, an asphaltene particle, which is disposed on a substrate, can be removed from the substrate by contacting the asphaltene particle with the functionalized silicate nanoparticle. The functionalized silicate nanoparticle can be interposed between the asphaltene particle and the substrate. The asphaltene particle can be separated from the substrate with the functionalized silicate nanoparticle and removed from the substrate.

[0081] In an embodiment, the asphaltene particle can be exfoliated using the functionalized silicate nanoparticle. Without wishing to be bound by theory, it is believed the aromatic moiety of the functionalized silicate nanoparticle can intercalate in a gallery of the

asphaltene particle. The functionalized silicate nanoparticle can be heated to a temperature effective to exfoliate the functionalized silicate nanoparticle, asphaltene particle, or a combination thereof. According to an embodiment, the functionalized silicate nanoparticle expands upon heating. As a result of the expansion, a distance increases between neighboring aromatic moieties that are tethered to the functionalized silicate nanoparticle. The asphaltene particles expand and exfoliate in response to the increased distance between the neighboring aromatic moieties of the functionalized silicate nanoparticle that are intercalated in the gallery of asphaltene molecules of the asphaltene particle. Exfoliation of the asphaltene particle can occur for asphaltene particles attached to the substrate or for those asphaltene particles that are not attached to a substrate.

[0082] The removal of the asphaltene particle from the substrate can also include contacting the asphaltene particle with a fluid. Here, contact of the fluid can increase a distance of separation between the asphaltene particle and the substrate before or after separating the asphaltene particle from the substrate.

[0083] Beneficially, the methods herein, e.g., contacting the silicate nanoparticle with the chemical group to form the functionalized silicate nanoparticle, can be performed in-situ in an environment such as a pipeline, downhole, formation, tubular, frac feature (e.g., a vein or pore), production zone, reservoir, refinery, transport tube, production tube, or a combination thereof. Moreover, the asphaltene particle or exfoliated asphaltene can be removed from the environment after separating the asphaltene particle from the substrate.

[0084] Furthermore, it has been found that perturbing the internal structure of asphaltene particles, for example, in a micelle or other aggregate, can lead to increased quality of oil containing asphaltenes. Additionally, degradation of asphaltene aggregates herein enhances production of petroleum fluid in a downhole, subsurface, or ground environment. Furthermore, removal of asphaltene from pores of a rock formation, within a reservoir, or from a sidewall of a tubular, production tubing, borehole, or transportation tube can improve the permeability of such structures, leading to increased quality of oil as well as increased or prolonged lifetime for oil production.

[0085] In an embodiment, a method for decomposing an asphaltene particle includes contacting the asphaltene particle with the functionalized silicate nanoparticle and causing the intercalating agent to increase a distance between asphaltene molecules in the asphaltene particle to decompose the asphaltene particle. As above, the aromatic moiety or nonaromatic moiety of the functionalized silicate nanoparticle can be disposed in the gallery between adjacent asphaltene molecules or disposed at the periphery of an asphaltene molecule such as

proximate to an edge of an aromatic plane or terminal chain attached to an aromatic portion of an asphaltene molecule in the asphaltene particle. During the exfoliation of the asphaltene particle, the functionalized silicate nanoparticle portion in the gallery forces the adjacent asphaltene molecules away from one another, thereby separating the asphaltene molecules. In this manner, an asphaltene molecule can be exfoliated from the asphaltene particle.

[0086] As used herein, “decomposition” refers to an increased separation distance between asphaltene molecules in an asphaltene particle, expansion of the volume of the asphaltene particle, complete removal of an asphaltene molecule from an asphaltene particle, or a change in the electronic structure or bonding in an asphaltene molecule in an asphaltene particle. Thus, decomposition includes, for example, deagglomeration, exfoliation, disaggregation, and the like. An example of a change in the electronic structure or bonding in an asphaltene molecule in an asphaltene particle includes converting a bond (e.g., converting a π bond to σ bond or vice versa), breaking a bond, or forming a bond.

[0087] Thus, according to an embodiment, the method includes exfoliating an asphaltene particle. In an embodiment, exfoliating includes removing an asphaltene molecule from the asphaltene particle. Exfoliation of an asphaltene particle, in an embodiment, decreases the number of asphaltene molecules in the asphaltene particle. It will be appreciated that exfoliation of asphaltene particles can provide exfoliated asphaltene as a single asphaltene molecule or as a micelle or layered particle containing fewer asphaltene molecules than the non-exfoliated asphaltene particle.

[0088] In a further embodiment, the method includes increasing the temperature of the functionalized silicate nanoparticle, asphaltene particle, or substrate. Increasing the temperature includes techniques that can elevate the temperature to about 60°C to about 1200°C, specifically about 100°C to about 1000°C, and more specifically about 100°C to about 800°C. Such techniques involve, for example, in-situ combustion, steam introduction, heated fluid injection, or a combination comprising at least one of the foregoing. In an embodiment, a downhole environment is heated by introducing steam in an injection well with the steam propagating through the formation and heating the functionalized silicate nanoparticle, asphaltene particle, or substrate. It is contemplated that increasing the temperature can cause reaction, including decomposition of the functionalized silicate nanoparticle, substrate, or asphaltene particle. In addition, the asphaltene particles can be heated to expand, decreasing the mutual attraction among asphaltene molecules therein. Depending on the amount of expansion of the asphaltene particle, asphaltene molecules can exfoliate from the asphaltene particles. In one embodiment, the heating of a functionalized

silicate nanoparticle associated with the asphaltene particle can lead to exfoliation of an asphaltene molecule therefrom.

[0089] Heated fluid injection can include heating a fluid (e.g., a solvent) and subsequently disposing the heated fluid downhole to increase the temperature of the asphaltene particles. In a non-limiting embodiment, in-situ combustion increases the temperature of the functionalized silicate nanoparticle by injecting a gas containing oxygen, for example air, downhole and igniting oil in the reservoir. The combustion releases heat, which can be absorbed by the functionalized silicate nanoparticle or asphaltene particle, in order to exfoliate an asphaltene molecule from the asphaltene particle.

[0090] In certain embodiments, the method further includes applying sonic frequencies to the intercalating agent. The sonic frequencies can be from about 400 hertz (Hz) to about 400 megahertz (MHz), specifically about 800 Hz to about 350 MHz, and more specifically about 1 kilohertz (kHz) to about 300 MHz. A transducer placed near the asphaltene particle can produce the sonic frequency, which can destructively interact with the asphaltene particle or functionalized silicate nanoparticle. Sonic frequencies may induce chemical reactions or expansion of the functionalized silicate nanoparticle and disrupt interparticle bonding in the asphaltene particle, leading to exfoliation of an asphaltene molecule. The sonic frequencies can detach neighboring polyaromatic planes of adjacent asphaltene molecules. Without wishing to be bound by any particular theory, such deterioration of the asphaltene particle may be induced by short-lived, localized disturbances (e.g., a hot spot) produced by the implosion of bubbles in the course of acoustic cavitation.

[0091] In some embodiments, the functionalized silicate nanoparticle is dispersed in a fluid. Such dispersion can occur before or after contacting the asphaltene particle with the functionalized silicate nanoparticle. The fluid can be an organic solvent, inorganic solvent, or a combination comprising at least one of the foregoing. Exemplary fluids are those above and can include CH_3NO_2 , CH_2Cl_2 , CHCl_3 , CCl_4 , $\text{C}_2\text{H}_4\text{Cl}_2$, H_2O , SOCl_2 , SO_2Cl_2 , $\text{S}_3\text{N}_3\text{Cl}_3$, benzene, toluene, o-xylene, dimethyl sulfoxide, furan, tetrahydrofuran, o-dioxane, m-dioxane, p-dioxane, dimethoxyethane, n-methyl-pyrrolidone, n,n-dimethylacetamide, γ -butyrolactone, 1,3-dimethyl-2-imidazolidinone, benzyl benzoate, hexafluorobenzene, octafluorotoluene, pentafluorobenzonitrile, pentafluoropyridine, pyridine, dimethylformamide, hexamethylphosphoramide, nitromethane, benzonitrile, or the like. In an embodiment, the fluid can react with the functionalized silicate nanoparticle to produce product compounds that decompose the asphaltene particle.

[0092] In another embodiment, after contact with the functionalized silicate nanoparticle, the asphaltene particle can be heated. The heat is absorbed by the asphaltene molecule, causing high amplitude vibrational motion of the non-polar groups, e.g., hydrocarbon tails that terminate an asphaltene molecule. In this manner, exfoliation of asphaltene molecules can occur by vibrationally-mediated dissociation or further increased spacing among the asphaltene molecules in the asphaltene particle. Additionally, the heated asphaltene particles can be more miscible with the fluid. Here, the fluid can be as before and can include, for example, an alkane, aromatic solvent, carbon dioxide, carbon disulfide, resin, oil, or a combination thereof. Particular fluids include, 2,2-dimethylpropane, butane, 2,2-dimethylbutane, pentane, hexane, heptane, octane, nonane, decane, undecane, cyclopentane, cyclohexane, benzene, toluene, o-xylene, dimethyl sulfoxide, furan, tetrahydrofuran, o-dioxane, m-dioxane, p-dioxane, dimethoxyethane, n-methyl-pyrrolidone, n,n-dimethylacetamide, γ -butyrolactone, 1,3-dimethyl-2-imidazolidinone, benzyl benzoate, hexafluorobenzene, octafluorotoluene, pentafluorobenzonitrile, pentafluoropyridine, pyridine, dimethylformamide, hexamethylphosphoramide, nitromethane, benzonitrile, and the like.

[0093] In another embodiment, a fluid or surfactant can contact the exfoliated asphaltene particle and allow dispersion of the asphaltene particle, for example, in an oil. Exemplary fluids include solvent such as a polar solvent, aromatic solvent, or a combination comprising at least one of the foregoing. The polar solvent can be an alcohol (e.g., ethanol, propanol, glycol, and the like), amine (e.g., methylamine, diethyl amine, tributyl amine, and the like), amide (e.g., dimethylformamide), ether (e.g., diethyl ether, polyether, tetrahydrofuran, and the like), ester (e.g., ethyl acetate, methyl butyrate, and the like), ketone (e.g., acetone), acetonitrile, dimethylsulfoxide, propylene carbonate, and the like. The aromatic solvent can be, for example, benzene, toluene, xylene, pyridine, hexafluorobenzene, octafluorotoluene, pentafluoropyridine, and the like.

[0094] The methods and materials herein can be used to enhance oil recovery in a reservoir, borehole, downhole, production zone, formation, or a combination thereof. Additionally, the methods and materials can be used to increase flow velocity of oil in a processing facility, refinery, pre-refinery facility, tubular, reactor, or a combination thereof. Removal of the asphaltene molecules from the substrate by the functionalized silicate nanoparticle herein can be used to extract asphaltene deposits that constrict flow in, for example, a tubular, and can restore flow in a plugged reservoir. Additionally, exfoliation of asphaltenes can increase permeability in porous media (e.g., a sand screen that can be deformable such as a polymeric open-cell foam) and flow channels (e.g., a crack in a

formation filled with proppant such as obtained in a fracking process). As a result of exfoliation to decrease the number of asphaltene molecules in an asphaltene particle, oil viscosity also decreases. Lowering the viscosity of the oil improves production efficiency. Additionally, the detrimental effects of asphaltene can be diminished or eliminated, including alleviation of flocculates of asphaltenes that can plug a reservoir or production tubing, restrict flow in a transport line, stabilize water-in-oil emulsions, foul a production facility, alter wettability of porous rock in the reservoir, or poison a refinery catalyst.

[0095] Thus, in an embodiment, a method for producing decomposed asphaltene includes disposing a functionalized silicate nanoparticle in an oil environment and contacting an asphaltene particle in the oil environment with the functionalized silicate nanoparticle. The embodiment also includes decomposing the asphaltene particle to produce decomposed asphaltene. In a certain embodiment, the method also includes breaking a water-in-oil emulsion in response to decomposing the asphaltene particle. Here the oil-in-water emulsion can be a Pickering emulsion that is stabilized by asphaltene particles at the water-oil interface. Upon decomposing the asphaltene particles, the emulsion is destabilized and thus broken.

[0096] In addition, water can be introduced by methods such as hot water injection, steam stimulation, or a combination comprising at least one of the foregoing. It is believed that, in this way, the asphaltene particles decompose as exfoliation of asphaltene molecules in the asphaltene particles occurs. As a result, the viscosity of oil in the oil environment is reduced. Moreover, increasing the mobility of the asphaltene particles by removing them from the substrate is advantageous as noted above. Therefore, the method can be used to enhance oil recovery. In a further embodiment, the method includes increasing a permeability of a reservoir of the oil environment. According to another embodiment, the method further includes producing the oil including the decomposed or removed asphaltene from the oil environment, wherein decomposing the asphaltene particle occurs prior to producing the oil. Alternatively or in addition, decomposing the asphaltene particle can occur subsequent to producing the oil.

[0097] While one or more embodiments have been shown and described, modifications and substitutions may be made thereto without departing from the spirit and scope of the invention. Accordingly, it is to be understood that the present invention has been described by way of illustrations and not limitation. Embodiments herein can be used independently or can be combined.

[0098] All ranges disclosed herein are inclusive of the endpoints, and the endpoints are independently combinable with each other. The ranges are continuous and thus contain every value and subset thereof in the range. The suffix “(s)” as used herein is intended to include both the singular and the plural of the term that it modifies, thereby including at least one of that term (e.g., the colorant(s) includes at least one colorants). “Optional” or “optionally” means that the subsequently described event or circumstance can or cannot occur, and that the description includes instances where the event occurs and instances where it does not. As used herein, “combination” is inclusive of blends, mixtures, alloys, reaction products, and the like.

[0099] As used herein, “a combination thereof” refers to a combination comprising at least one of the named constituents, components, compounds, or elements, optionally together with one or more of the same class of constituents, components, compounds, or elements.

[0100] The use of the terms “a” and “an” and “the” and similar referents in the context of describing the invention (especially in the context of the following claims) are to be construed to cover both the singular and the plural, unless otherwise indicated herein or clearly contradicted by context. “Or” means “and/or.” It should further be noted that the terms “first,” “second,” and the like herein do not denote any order, quantity, or importance, but rather are used to distinguish one element from another. The modifier “about” used in connection with a quantity is inclusive of the stated value and has the meaning dictated by the context (e.g., it includes the degree of error associated with measurement of the particular quantity). The conjunction “or” is used to link objects of a list or alternatives and is not disjunctive; rather the elements can be used separately or can be combined together under appropriate circumstances.

CLAIMS

What is claimed is:

1. A process for removing an asphaltene particle from a substrate, the process characterized by:

contacting a silicate nanoparticle with a chemical group to form a functionalized silicate nanoparticle, the chemical group comprising:

a first portion; and

a second portion comprising an aromatic moiety or a nonaromatic moiety, the first portion being directly bonded to the silicate nanoparticle in the functionalized silicate nanoparticle; and the second portion comprising a moiety of a guanidine, a guanidinium salt, a biguanidine, a biguanidinium salt, or a sulfonium salt;

contacting the asphaltene particle with the functionalized silicate nanoparticle, the asphaltene particle being disposed on the substrate;

interposing the functionalized silicate nanoparticle between the asphaltene particle and the substrate; and

separating the asphaltene particle from the substrate with the functionalized silicate nanoparticle to remove the asphaltene particle from the substrate.

2. The process of claim 1, further characterized by exfoliating the asphaltene particle with the functionalized silicate nanoparticle.

3. The process of claim 2, further characterized by heating the functionalized silicate nanoparticle to a temperature effective to exfoliate the functionalized silicate nanoparticle, asphaltene particle, or a combination thereof.

4. The process of claim 1, further characterized by contacting the asphaltene particle with a fluid to increase a distance of separation between the asphaltene particle and the substrate after separating the asphaltene particle from the substrate, optionally the fluid comprises water, oil, carbon dioxide, a liquefied C1-C6 alkane, completion fluid, brine, acid base, or a combination thereof.

5. The process of any one of claims 1 to 4, wherein contacting the silicate nanoparticle with the group is performed in-situ in an environment comprising, a pipeline, downhole, formation, tubular, frac feature, production zone, reservoir, or a combination thereof.

6. The process of claim 5, further characterized by removing the asphaltene particle from the environment after separating the asphaltene particle from the substrate.

7. The process of any one of claims 1 to 6, wherein the substrate comprises a metal, composite, sand, rock, mineral, glass, formation, downhole element, or a combination thereof.

8. The process of any one of claims 1 to 7, wherein the silicate nanoparticle comprises a silsesquioxane, cyclosilicate, inosilicate, nesosilicate, phyllosilicate, sorosilicate, tectosilicate, or a combination thereof.

9. The process of any one of claims 1 to 8, wherein the aromatic moiety comprises anthracenyl, azulenyl, benzocyclooctenyl, benzocycloheptenyl, biphenyl, chrysenyl, fluorenyl, indanyl, indenyl, naphthyl, pentalenyl, phenalenyl, phenanthrenyl, phenanthryl, phenyl, pyrenyl, tetrahydronaphthyl, a heteroaryl group, a derivative thereof, or a combination thereof.

10. The process of any one of claims 1 to 9, wherein the chemical group is derived from a guanidine, a guanidinium salt, a biguanidine, a biguanidinium salt, a sulfonium salt, or a combination thereof.

11. The process of any one of claims 1 to 10, wherein the first portion of the chemical group comprises a bond, linker group, or a combination thereof; the linker group comprises a bond, C1 to C30 alkylene, C3 to C30 cycloalkenylene, C1 to C30 fluoroalkylene, C3 to C30 cycloalkylene, C3 to C30 heterocycloalkylene, C5 to C30 arylene, C6 to C40 aralkylene, C6 to C30 aryleneoxy, C2 to C30 heteroarylene, C6 to C40 heteroaralkylene, C2 to C30 alkenylene, C2 to C30 alkynylene, C1 to C30 amide, amine, C1 to C30 oxyalkylene, C1 to C30 oxyarylene, oxygen (O), sulfur (S), or a combination thereof, and

the linker group is disposed between the silicate nanoparticle and the second portion of the chemical group in the functionalized silicate nanoparticle.

12. A composition effective to remove an asphaltene particle from a substrate, the composition characterized by

a functionalized silicate nanoparticle comprising a reaction product of

a silicate nanoparticle; and

a functionalization compound comprising quaternary ammonium salt, quaternary phosphonium salt, alkoxy silane, halide, guanidine, guanidinium salt, biguanidine, biguanidinium salt, sulfonium salt, or a combination thereof; and

a fluid,

wherein the functionalization compound includes a chemical group comprising:




a first portion, the first portion being directly bonded to the silicate nanoparticle in the functionalized silicate nanoparticle; and


a second portion comprising an aromatic moiety or a nonaromatic moiety, and the composition is effective to remove an asphaltene particle from a substrate comprising a metal, composite, sand, rock, mineral, glass, formation, downhole element, or a combination thereof.

13. The composition of claim 12, wherein the silicate nanoparticle comprises a silsesquioxane, cyclosilicate, inosilicate, nesosilicate, phyllosilicate, sorosilicate, tectosilicate, or a combination thereof.

14. The composition of claim 12 or claim 13, wherein the second portion comprises an aromatic moiety comprising anthracenyl, azulenyl, benzocyclooctenyl, benzocycloheptenyl, biphenyl, chrysenyl, fluorenyl, indanyl, indenyl, naphthyl, pentalenyl, phenalenyl, phenanthrenyl, phenanthryl, phenyl, pyrenyl, tetrahydronaphthyl, a heteroaryl group, derivatives thereof, or a combination thereof, or

wherein the second portion comprising a moiety of the following formulas or a salt thereof:





$\text{R}_1\text{-R}_9$ and $\text{R}_{11}\text{-R}_{24}$ are independently hydrogen atom, alkyl, heterocyclic, carbonyl, amino, amide, sulfonamide, phosphoramidate, imide moiety, an aromatic moiety, or a combination thereof with the proviso that optionally at least one of $\text{R}_1\text{-R}_9$ and $\text{R}_{11}\text{-R}_{24}$ is a covalent bond or a divalent radical attached to the first portion;

R_{10} is a bond, alkylene, heterocyclic biradical, carbonyl, amino, amide, sulfonamide, phosphoramidate, imide biradical, an aromatic moiety, or a combination thereof.

15. A process for removing an asphaltene particle from a substrate, the process characterized by:

contacting a silicate nanoparticle with a chemical group to form a functionalized silicate nanoparticle, the chemical group comprising:

a first portion; and

a second portion comprising a nonaromatic moiety, the first portion being directly bonded to the silicate nanoparticle in the functionalized silicate nanoparticle; and the second portion comprising a moiety of a guanidine, a guanidinium salt, a biguanidine, a biguanidinium salt, or a sulfonium salt;

contacting the asphaltene particle with the functionalized silicate nanoparticle, the asphaltene particle being disposed on the substrate;

interposing the functionalized silicate nanoparticle between the asphaltene particle and the substrate; and

separating the asphaltene particle from the substrate with the functionalized silicate nanoparticle to remove the asphaltene particle from the substrate,

wherein the chemical group is derived from a quaternary ammonium salt, quaternary phosphonium salt, alkoxy silane, halide, or a combination thereof.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US2016/022776**A. CLASSIFICATION OF SUBJECT MATTER****C09K 3/32(2006.01)i, A62D 3/30(2007.01)i**

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

C09K 3/32; C11D 1/82; C09K 8/524; E21B 43/24; C07C 7/20; C07C 303/02; C10G 1/04; E21B 43/26; C09K 8/584; B01D 17/00; A62D 3/30

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Korean utility models and applications for utility models
Japanese utility models and applications for utility models

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

eKOMPASS(KIPO internal) & Keywords: asphaltene, removing, functionalized silicate nanoparticle, guanidine, guanidinium salt, biguanidine, biguanidinium, sulfonium salt**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2014-106007 A1 (BAKER HUGHES INCORPORATED) 3 July 2014 See paragraphs [0039]–[0041]; claims 1-2, 4-8 and 18-20.	1-6, 12-15
A	US 6048904 A (WIEHE, IRWIN A. et al.) 11 April 2000 See claims 1-2, 7; and column 1, lines 7-10.	1-6, 12-15
A	US 2012-0329887 A1 (PHUKAN, MONJIT et al.) 27 December 2012 See abstract; claim 1; and paragraphs [0001], [0004].	1-6, 12-15
A	US 2010-0096139 A1 (HOLCOMB, DAVID L. et al.) 22 April 2010 See claim 1; and paragraph [0010].	1-6, 12-15
A	US 2012-0255887 A1 (HOLMS, BRADFORD A. et al.) 11 October 2012 See abstract; and claim 1.	1-6, 12-15



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

04 July 2016 (04.07.2016)

Date of mailing of the international search report

04 July 2016 (04.07.2016)

Name and mailing address of the ISA/KR

International Application Division
Korean Intellectual Property Office
189 Cheongsa-ro, Seo-gu, Daejeon, 35208, Republic of Korea

Facsimile No. +82-42-481-8578

Authorized officer

KIM, Dong Seok

Telephone No. +82-42-481-5405



INTERNATIONAL SEARCH REPORTInternational application No.
PCT/US2016/022776**Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)**

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☒ Claims Nos.: 7-11
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fees, this Authority did not invite payment of any additional fees.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- ☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/US2016/022776

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2014-106007 A1	03/07/2014	CA 2896859 A1 US 2014-0187449 A1 US 9012377 B2	03/07/2014 03/07/2014 21/04/2015
US 6048904 A	11/04/2000	AU 2000-19196 A1 AU 2000-19196 B2 CA 2355196 A1 CA 2355196 C EP 1135354 A1 EP 1135354 B1 JP 2002-531591 A WO 00-32546 A1	19/06/2000 23/01/2003 08/06/2000 23/12/2008 26/09/2001 28/04/2004 24/09/2002 08/06/2000
US 2012-0329887 A1	27/12/2012	CA 2838343 A1 CN 103781853 A EP 2723817 A1 MX 2013014563 A US 8779012 B2 WO 2012-177836 A1	27/12/2012 07/05/2014 30/04/2014 21/03/2014 15/07/2014 27/12/2012
US 2010-0096139 A1	22/04/2010	CA 2738929 A1 CA 2739285 A1 EP 2340243 A1 EP 2340243 B1 EP 2350227 A1 EP 2465911 A1 JP 05529876 B2 JP 2012-505891 A MX 2011003953 A US 2011-0275631 A1 US 2012-0168165 A1 US 2012-0175120 A1 US 8785489 B2 WO 2010-045188 A1 WO 2010-045545 A1	22/04/2010 22/04/2010 06/07/2011 08/10/2014 03/08/2011 20/06/2012 25/06/2014 08/03/2012 19/05/2011 10/11/2011 05/07/2012 12/07/2012 22/07/2014 22/04/2010 22/04/2010
US 2012-0255887 A1	11/10/2012	AR 085841 A1 WO 2012-138454 A1	30/10/2013 11/10/2012