

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
19 August 2010 (19.08.2010)

PCT

(10) International Publication Number
WO 2010/092032 A1

(51) International Patent Classification:

A01P 3/00 (2006.01) *A01P 21/00* (2006.01)
A01P 5/00 (2006.01) *A01N 33/12* (2006.01)
A01P 7/02 (2006.01) *A01N 47/02* (2006.01)
A01P 7/04 (2006.01)

(21) International Application Number:

PCT/EP2010/051530

(22) International Filing Date:

9 February 2010 (09.02.2010)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

09152587.3 11 February 2009 (11.02.2009) EP

(71) Applicant (for all designated States except US): **BASF SE** [DE/DE]; 67056 Ludwigshafen (DE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **VOESTE, Dirk** [DE/DE]; Berliner Platz 11, 67117 Limburgerhof (DE).
WILHELM, Ronald [DE/DE]; Platanenweg 11, 65719 Hofheim (DE).

(74) Common Representative: **BASF SE**; 67056 Ludwigshafen (DE).

(81) Designated States (unless otherwise indicated, for every

kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every

kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))



WO 2010/092032 A1

(54) Title: PESTICIDAL MIXTURES

(57) Abstract: The present invention relates to synergistic mixtures of chlormequat chloride with insecticides.

Pesticidal mixtures

Description

5 The present invention relates to synergistic mixtures comprising, as active components,

- 1) chlormequat chloride as compound I; and
- 2) one or more insecticidal or nematicidal compound II selected from the group consisting of

10 IIa) nicotinic receptor agonists/antagonists compounds: clothianidin, dinotefuran, imidacloprid, thiamethoxam, nitenpyram and acetamiprid; and

IIb) macrolide compounds: abamectin, emamectin benzoate and spinosad; and

15 IIc) fipronil or ethiprole
in synergistic effective amounts.

in synergistic effective amounts.

20 The invention furthermore relates to mixtures comprising in addition to the compounds I and II

3 a) a fungicidal compound IIIA selected from the group of strobilurine fungicides IIIA comprising pyraclostrobin, azoxystrobin, dimoxystrobin, enestroburin, fluoxastrobin, kresoxim-methyl, metominostrobin, oryastrobin, picoxystrobin, pyribencarb, trifloxystrobin, 2-(2-(6-(3-chloro-2-methyl-phenoxy)-5-fluoro-pyrimidin-4-yloxy)-phenyl)-2-methoxyimino-N-methyl-acetamide, 3-methoxy-2-(2-(N-(4-methoxy-phenyl)-cyclopropane-carboximidoylsulfanylmethyl)-phenyl)-acrylic acid methyl ester, methyl (2-chloro-5-[1-(3-methylbenzyloxyimino)ethyl]benzyl)carbamate and 2-(2-(3-(2,6-dichlorophenyl)-1-methyl-allylideneaminooxymethyl)-phenyl)-2-methoxyimino-N-methyl-acetamide; and

30 3 b) an fungicidal compound IIIB from the group of carboxamides consisting of N-(3',4',5'-trifluorobiphenyl-2-yl)-3-difluoromethyl-1-methyl-1H-pyrazole-4-carboxamide, N-[2-(4'-trifluoromethylthio)-biphenyl]-3-difluoromethyl-1-methyl-1H-pyrazole-4-carboxamide, bixafen, N-[2-(1,3-dimethylbutyl)-phenyl]-1,3-dimethyl-5-fluoro-1H-pyrazole-4-carboxamide (penflufen), sedaxane, isopyrazam and penthiopyrad);

in synergistic effective amounts.

40 The present invention also relates to the binary mixtures of fipronil and abamectin
in synergistically effective amounts.

These above-referred mixtures are hereinbelow also referred as "inventive mixtures".

5 Moreover, the invention relates to a method for controlling pests, this refers to includes phytopathogenic animal pests and phytopathogenic harmful fungi, using the inventive mixtures and to the use of compound I and compound II (and optionally compound III) for preparing such mixtures, and also to compositions comprising such mixtures.

10 In one embodiment, the present invention provides methods for the control of phytopathogenic animal pests (such as insects, acarids or nematodes) comprising contacting the phytopathogenic animal pest (the insect, acarid or nematode) or their food supply, habitat, breeding grounds or their locus with a pesticidally effective amount of the inventive mixtures.

15 Moreover, in another embodiment the present invention also relates to a method of protecting plants from attack or infestation by phytopathogenic animal pests (insects, acarids or nematodes) comprising contacting the plant, or the soil or water in which the plant is growing, with a pesticidally effective amount of the inventive mixture.

20 Additionally, the present invention also comprises a method for protection of plant propagation material from harmful pests, such as phytopathogenic harmful fungi or phytopathogenic animal pests (such as insects, arachnids or nematodes) comprising contacting the plant propagation materials with an inventive mixture in pesticidally effective amounts

25

The term "plant propagation material" is to be understood to denote all the generative parts of the plant such as seeds and vegetative plant material such as cuttings and tubers (e. g. potatoes), which can be used for the multiplication of the plant. This includes seeds, roots, fruits, tubers, bulbs, rhizomes, shoots, sprouts and other parts of
30 plants, including seedlings and young plants, which are to be transplanted after germination or after emergence from soil. These young plants may also be protected before transplantation by a total or partial treatment by immersion or pouring. In a particular preferred embodiment, the term propagation material denotes seeds.

35 Moreover, the invention relates to a method for controlling phytopathogenic harmful fungi using the inventive mixtures and to the use of the compound I and compound II (and optionally compound III) for preparing such mixtures, and also to compositions comprising such mixtures.

40 The present invention further relates to plant-protecting active ingredient mixtures having synergistically enhanced action of improving the health of plants and to a method of applying such inventive mixtures to the plants.

The compounds I, II, III as well as their pesticidal action and methods for producing them are generally known. For instance, the commercially available compounds may be found in The Pesticide Manual, 14th Edition, British Crop Protection Council (2006) among other publications.

Binary mixtures comprising clothianidin together with a large listing of potential fungicidal mixing partners are disclosed in WO 99/63826.

Combinations generically comprising neonicotinoids and explicitly disclosing imidacloprid together with a large listing of potential fungicidal mixing partners are disclosed in WO 96/3045. WO 06/23899 also discloses binary mixtures of imidacloprid and several other fungicides.

WO 06/069654, WO06/089876 and WO 06/23899 disclose mixtures of neonicotinoids and strobilurins. WO 06/23899 also discloses mixtures of imidacloprid and other fungicides.

WO 08/006541 discloses penthiopyrad mixtures with fungicides, which optionally may comprise an insecticide.

WO 97/22254 discloses mixtures of thiamethoxam with several fungicides mentioning inter alia fludioxonil and metalaxyl. WO 06/128655 disclosed mixtures of neonicotinoids with several azoles as well as mixtures of a huge number of insecticides that can be combined with several fungicides. WO 06/24333 describes a neonicotinoid formulation, which may, as second component comprise at least one further fungicide.

US 2005/0209304 discloses mixtures for seed treatment comprising abamectin and selected fungicides. Azoxystrobin, metalaxyl, fludioxonil and abamectin is disclosed as a specific four way mixture.

However, the specific binary, ternary and quarternary mixtures of the present invention, which are based on compound I and compound II (and optionally compound III) are not disclosed therein, also not the outstanding action, such combinations might have especially regarding seed treatment.

One typical problem arising in the field of pest control lies in the need to reduce the dosage rates of the active ingredient in order to reduce or avoid unfavorable environmental or toxicological effects whilst still allowing effective pest control.

In regard to the instant invention the term "phytopathogenic pests" embrace phytopathogenic animal pests, and phytopathogenic harmful fungi. The term phytopathogenic

animal pests is hereinbelow abbreviated as "animal pest" and the term "phytopathogenic harmful fungi" is hereinbelow abbreviated as "harmful fungi".

5 Another problem encountered concerns the need to have available pest control agents which are effective against a broad spectrum of pests, e.g. both animal pests and harmful fungi.

10 There also exists the need for pest control agents that combine knock-down activity with prolonged control, that is, fast action with long lasting action.

15 Another difficulty in relation to the use of pesticides is that the repeated and exclusive application of an individual pesticidal compound leads in many cases to a rapid selection of pests, that means animal pests, and harmful fungi, which have developed natural or adapted resistance against the active compound in question. Therefore there is a need for pest control agents that help prevent or overcome resistance.

20 Another problem underlying the present invention is the desire for compositions that improve plants, a process which is commonly and hereinafter referred to as "plant health".

25 The term plant health comprises various sorts of improvements of plants that are not connected to the control of pests. For example, advantageous properties that may be mentioned are improved crop characteristics including: emergence, crop yields, protein content, oil content, starch content, more developed root system (improved root growth), improved stress tolerance (e.g. against drought, heat, salt, UV, water, cold), reduced ethylene (reduced production and/or inhibition of reception), tillering increase, increase in plant height, bigger leaf blade, less dead basal leaves, stronger tillers, greener leaf color, pigment content, photosynthetic activity, less input needed (such as fertilizers or water), less seeds needed, more productive tillers, earlier flowering, early grain maturity, less plant verse (lodging), increased shoot growth, enhanced plant vigor, increased plant stand and early and better germination; or any other advantages familiar to a person skilled in the art.

35 In particular, the term plant health relates to improved stress tolerance, preferably against drought, heat, salt, UV, water, cold, more preferably against drought (which correlates to the ability of a plant to have improved abilities to tolerate water deficit).

40 It was therefore an object of the present invention to provide pesticidal mixtures which solve the problems of reducing the dosage rate and / or enhancing the spectrum of activity and / or combining knock-down activity with prolonged control and / or to resistance management and/or promoting the health of plants.

We have found that this object is in part or in whole achieved by the complex mixtures comprising the active compounds defined in the outset.

5 Especially, it has been found that the mixtures as defined in the outset show markedly enhanced action against pests compared to the control rates that are possible with the individual compounds and/or is suitable for improving the health of plants when applied to plants, parts of plants, plant propagation materials (preferably seeds), or at their locus of growth.

10 It has been found that the action of the inventive mixtures goes far beyond the fungicidal and/or insecticidal and/or plant health improving action of the active compounds present in the mixture alone.

15 Moreover, we have found that simultaneous, that is joint or separate, application of the compound I and compound II (and optionally compound III) or successive application of the compound I and compound II (and optionally compound III) allows enhanced control of pests, that means animal pests, and harmful fungi, compared to the control rates that are possible with the individual compounds (synergistic mixtures; synergistic pesti-

20

Moreover, we have found that simultaneous, that is joint or separate, application of the compound I and compound II (and optionally compound III) or successive application of the compound I and compound II (and optionally compound III) provides enhanced plant health effects compared to the plant health effects that are possible with the individual compounds (synergistic mixtures (synergistically enhanced plant health effects)).

25

Preferably, compound II is clothianidin, imidacloprid, thiamethoxam, acetamiprid, abamectin or fipronil.

30 Preferably, compound II is a mixture of compound IIa (which is clothianidin, imidacloprid, thiamethoxam or acetamiprid) and compound IIc (which is preferably fipronil). Preferably, compound II is a mixture of compound IIa (which is clothianidin, imidacloprid, thiamethoxam or acetamiprid) and compound IIb (which is abamectin). Preferably, compound II is a mixture of compound IIa (which is clothianidin, imidaclo-

35

With respect to their intended use, the following mixtures listed in the table 1 below are preferred:

40 In table 1, the following abbreviations are used:
CCC = chlormequatchloride F = Fipronil
AC is Acetamiprid

TMX is thiamethoxam
 IMI is imidacloprid
 C is clothianidin
 AB= Abamectin
 A = Azoxystrobin

Table 1

No	I	II (1)	II (2)	IIIA	IIIB
R-1	CCC	AC	-	-	-
R-2	CCC	TMX	-	-	-
R-3	CCC	IMI	-	-	-
R-4	CCC	C	-	-	-
R-5	CCC	AB	-	-	-
R-6	CCC	F	-	-	-
R-7	CCC	AC	AB	-	-
R-8	CCC	TMX	AB	-	-
R-9	CCC	IMI	AB	-	-
R-10	CCC	C	AB	-	-
R-11	CCC	AC	F	-	-
R-12	CCC	TMX	F	-	-
R-13	CCC	IMI	F	-	-
R-14	CCC	C	F	-	-
R-15	CCC	F	AB	-	-

The present invention also relates to the binary mixtures of
 5 fipronil and abamectin
 in synergistically effective amounts.

The present invention further relates to the binary mixtures comprising synergistically
 effective amounts of the compounds IIA
 10 abamectin and clothianidin; and
 abamectin and imidacloprid; and
 abamectin and acetamiprid; and
 abamectin and thiamethoxam.

In a preferred embodiment, the present invention therefore comprises binary mixtures
 15 of chlormequat-chloride and clothianidin, imidacloprid, thiamethoxam, acetamiprid,
 abamectin or fipronil.

The ratios by weight for the respective binary mixtures comprising compound I and
 20 compound II are from 1:200 to 200:1, preferably from 50:1 to 1:50, more preferably
 from 1:20 to 20:1.

The ratios by weight for the respective binary mixtures comprising two compounds II and compound II are from 1:200 to 200:1, preferably from 50:1 to 1:50, more preferably from 1:20 to 20:1.

- 5 The ratios by weight for the respective ternary mixtures comprising compound and two compounds II are from 200:200:1 to 1:200:200, preferably from 200:1:1 to 1:1:200.

10 Preferably, the ternary and quarternary mixtures according to the present invention comprise as compound IIIA a strobilurine fungicide selected from the group consisting of pyraclostrobin, azoxystrobin, dimoxystrobin, enestroburin, fluoxastrobin, kresoxim-methyl, metominostrobin, orysastrobin, picoxystrobin, pyribencarb and trifloxystrobin, wherein pyraclostrobin, azoxystrobin, orysastrobin and trifloxystrobin are more preferred and pyraclostrobin and orysastrobin are most preferred. Utmost preference is given to pyraclostrobin.

15

In a further preferred embodiment, the ternary and quarternary mixtures according to the present invention comprise a fungicidal compound IIIB from the group of carboxamides consisting of N-(3',4',5'-trifluorobiphenyl-2-yl)- 3-difluoromethyl-1-methyl-1H-pyrazole-4-carboxamide, N-[2-(4'-trifluoromethylthio)-biphenyl]-3-difluoromethyl-1-methyl-1H-pyrazole-4-carboxamide, bixafen, N-[2-(1,3-dimethylbutyl)-phenyl]-1,3-dimethyl-5-fluoro-1H-pyrazole-4-carboxamide (penflufen), sedaxane, isopyrazam and penthiopyrad), wherein N-(3',4',5'-trifluorobiphenyl-2-yl)- 3-difluoromethyl-1-methyl-1H-pyrazole-4-carboxamide, sedaxane, N-[2-(1,3-dimethylbutyl)-phenyl]-1,3-dimethyl-5-fluoro-1H-pyrazole-4-carboxamide (penflufen) and penthiopyrad are more preferred and N-(3',4',5'-trifluorobiphenyl-2-yl)- 3-difluoromethyl-1-methyl-1H-pyrazole-4-carboxamide is most preferred.

20
25

In a further embodiment, the quarternary and fivefold mixtures according to the present invention comprise a mixture of compound IIIA and compound IIIB, wherein for compound IIIA pyraclostrobin, azoxystrobin, orysastrobin and trifloxystrobin are more preferred, pyraclostrobin and orysastrobin are most preferred and pyraclostrobin is utmost preferred;

30

and for compound IIIB N-(3',4',5'-trifluorobiphenyl-2-yl)- 3-difluoromethyl-1-methyl-1H-pyrazole-4-carboxamide, sedaxane, N-[2-(1,3-dimethylbutyl)-phenyl]-1,3-dimethyl-5-fluoro-1H-pyrazole-4-carboxamide (penflufen) and penthiopyrad are more preferred and N-(3',4',5'-trifluorobiphenyl-2-yl)- 3-difluoromethyl-1-methyl-1H-pyrazole-4-carboxamide is most preferred.

35

The ratios by weight for the respective ternary mixtures comprising compound I, the compound II and fungicidal compound III are from 200:200:1 to 1:200:200, preferably from 200:1:1 to 1:1:200.

40

The ratios by weight for the respective quaternary mixtures comprising comprising compound I, the compound II, two compounds III are from are from 1:200:200:200 to 200:1:1:1.

- 5 The ratios by weight for the respective quaternary mixtures comprising comprising compound I, two compounds II and compound III are from are from 1:200:200:200 to 200:1:1:1.

- 10 The ratios by weight for the respective fivefold mixtures comprising comprising compound I, two compounds II and compound IIIA and compound IIIB are from from 1:200:200:200:200 to 200:1:1:1:1

- 15 With respect to their intended use, the following ternary and quaternary and fivefold mixtures of compound I, compound II and compound III (and optionally IV or V) listed in the table 1 below are preferred.

In table 1, the following abbreviations are used:

CCC = chlormequatchloride	F = Fipronil
AC is Acetamidrid	N-(3',4',5'-trifluorobiphenyl-2-yl)- 3-difluoromethyl-1-methyl-1H-pyrazole-4-carboxamide = IIIB1
TMX is thiamethoxam	N-[2-(1,3-dimethylbutyl)-phenyl]-1,3-dimethyl-5-fluoro-1H-pyrazole-4-carboxamide (penflufen) = IIIB2
IMI is imidacloprid	sedaxane = IIIB3
C is clothianidin	penthiopyrad = IIIB4
AB= Abamectin	
P = Pyraclostrobin	
O = Orysastrobin	
T = Trifloxystrobin	
A = Azoxystrobin	

20 Table 1

No	I	II (1)	II (2)	IIIA	IIIB
S-1	CCC	AC	-	P	-
S-2	CCC	TMX	-	P	-
S-3	CCC	IMI	-	P	-
S-4	CCC	C	-	P	-
S-5	CCC	AB	-	P	-
S-6	CCC	F	-	P	-
S-7	CCC	AC	AB	P	-
S-8	CCC	TMX	AB	P	-
S-9	CCC	IMI	AB	P	-
S-10	CCC	C	AB	P	-

No	I	II (1)	II (2)	IIIA	IIIB
S-11	CCC	AC	F	P	-
S-12	CCC	TMX	F	P	-
S-13	CCC	IMI	F	P	-
S-14	CCC	C	F	P	-
S-15	CCC	F	AB	P	-
S-16	CCC	AC	-	O	-
S-17	CCC	TMX	-	O	-
S-18	CCC	IMI	-	O	-
S-19	CCC	C	-	O	-
S-20	CCC	AB	-	O	-
S-21	CCC	F	-	O	-
S-22	CCC	AC	AB	O	-
S-23	CCC	TMX	AB	O	-
S-24	CCC	IMI	AB	O	-
S-25	CCC	C	AB	O	-
S-26	CCC	AC	F	O	-
S-27	CCC	TMX	F	O	-
S-28	CCC	IMI	F	O	-
S-29	CCC	C	F	O	-
S-30	CCC	F	AB	O	-
S-31	CCC	AC	-	T	-
S-32	CCC	TMX	-	T	-
S-33	CCC	IMI	-	T	-
S-34	CCC	C	-	T	-
S-35	CCC	AB	-	T	-
S-36	CCC	F	-	T	-
S-37	CCC	AC	AB	T	-
S-38	CCC	TMX	AB	T	-
S-39	CCC	IMI	AB	T	-
S-40	CCC	C	AB	T	-
S-41	CCC	AC	F	T	-
S-42	CCC	TMX	F	T	-
S-43	CCC	IMI	F	T	-
S-44	CCC	C	F	T	-
S-45	CCC	F	AB	T	-
S-46	CCC	AC	-	A	-
S-47	CCC	TMX	-	A	-
S-48	CCC	IMI	-	A	-
S-49	CCC	C	-	A	-
S-50	CCC	AB	-	A	-

No	I	II (1)	II (2)	IIIA	IIIB
S-51	CCC	F	-	A	-
S-52	CCC	AC	AB	A	-
S-53	CCC	TMX	AB	A	-
S-54	CCC	IMI	AB	A	-
S-55	CCC	C	AB	A	-
S-56	CCC	AC	F	A	-
S-57	CCC	TMX	F	A	-
S-58	CCC	IMI	F	A	-
S-59	CCC	C	F	A	-
S-60	CCC	F	AB	A	-
S-61	CCC	AC	-	-	IIIB1
S-62	CCC	TMX	-	-	IIIB1
S-63	CCC	IMI	-	-	IIIB1
S-64	CCC	C	-	-	IIIB1
S-65	CCC	AB	-	-	IIIB1
S-66	CCC	F	-	-	IIIB1
S-67	CCC	AC	AB	-	IIIB1
S-68	CCC	TMX	AB	-	IIIB1
S-69	CCC	IMI	AB	-	IIIB1
S-70	CCC	C	AB	-	IIIB1
S-71	CCC	AC	F	-	IIIB1
S-72	CCC	TMX	F	-	IIIB1
S-73	CCC	IMI	F	-	IIIB1
S-74	CCC	C	F	-	IIIB1
S-75	CCC	F	AB	-	IIIB1
S-76	CCC	AC	-	-	IIIB2
S-77	CCC	TMX	-	-	IIIB2
S-78	CCC	IMI	-	-	IIIB2
S-79	CCC	C	-	-	IIIB2
S-80	CCC	AB	-	-	IIIB2
S-81	CCC	F	-	-	IIIB2
S-82	CCC	AC	AB	-	IIIB2
S-83	CCC	TMX	AB	-	IIIB2
S-84	CCC	IMI	AB	-	IIIB2
S-85	CCC	C	AB	-	IIIB2
S-86	CCC	AC	F	-	IIIB2
S-87	CCC	TMX	F	-	IIIB2
S-88	CCC	IMI	F	-	IIIB2
S-89	CCC	C	F	-	IIIB2
S-90	CCC	F	AB	-	IIIB2

No	I	II (1)	II (2)	IIIA	IIIB
S-91	CCC	AC	-	-	IIIB3
S-92	CCC	TMX	-	-	IIIB3
S-93	CCC	IMI	-	-	IIIB3
S-94	CCC	C	-	-	IIIB3
S-95	CCC	AB	-	-	IIIB3
S-96	CCC	F	-	-	IIIB3
S-97	CCC	AC	AB	-	IIIB3
S-98	CCC	TMX	AB	-	IIIB3
S-99	CCC	IMI	AB	-	IIIB3
S-100	CCC	C	AB	-	IIIB3
S-101	CCC	AC	F	-	IIIB3
S-102	CCC	TMX	F	-	IIIB3
S-103	CCC	IMI	F	-	IIIB3
S-104	CCC	C	F	-	IIIB3
S-105	CCC	F	AB	-	IIIB3
S-106	CCC	AC	-	-	IIIB4
S-107	CCC	TMX	-	-	IIIB4
S-108	CCC	IMI	-	-	IIIB4
S-109	CCC	C	-	-	IIIB4
S-110	CCC	AB	-	-	IIIB4
S-111	CCC	F	-	-	IIIB4
S-112	CCC	AC	AB	-	IIIB4
S-113	CCC	TMX	AB	-	IIIB4
S-114	CCC	IMI	AB	-	IIIB4
S-115	CCC	C	AB	-	IIIB4
S-116	CCC	AC	F	-	IIIB4
S-117	CCC	TMX	F	-	IIIB4
S-118	CCC	IMI	F	-	IIIB4
S-119	CCC	C	F	-	IIIB4
S-120	CCC	F	AB	-	IIIB4
S-121	CCC	AC	-	P	IIIB1
S-122	CCC	TMX	-	P	IIIB1
S-123	CCC	IMI	-	P	IIIB1
S-124	CCC	C	-	P	IIIB1
S-125	CCC	AB	-	P	IIIB1
S-126	CCC	F	-	P	IIIB1
S-127	CCC	AC	AB	P	IIIB1
S-128	CCC	TMX	AB	P	IIIB1
S-129	CCC	IMI	AB	P	IIIB1
S-130	CCC	C	AB	P	IIIB1

No	I	II (1)	II (2)	IIIA	IIIB
S-131	CCC	AC	F	P	IIIB1
S-132	CCC	TMX	F	P	IIIB1
S-133	CCC	IMI	F	P	IIIB1
S-134	CCC	C	F	P	IIIB1
S-135	CCC	F	AB	P	IIIB1
S-136	CCC	AC	-	P	IIIB2
S-137	CCC	TMX	-	P	IIIB2
S-138	CCC	IMI	-	P	IIIB2
S-139	CCC	C	-	P	IIIB2
S-140	CCC	AB	-	P	IIIB2
S-141	CCC	F	-	P	IIIB2
S-142	CCC	AC	AB	P	IIIB2
S-143	CCC	TMX	AB	P	IIIB2
S-144	CCC	IMI	AB	P	IIIB2
S-145	CCC	C	AB	P	IIIB2
S-146	CCC	AC	F	P	IIIB2
S-147	CCC	TMX	F	P	IIIB2
S-148	CCC	IMI	F	P	IIIB2
S-149	CCC	C	F	P	IIIB2
S-150	CCC	F	AB	P	IIIB2
S-151	CCC	AC	-	P	IIIB3
S-152	CCC	TMX	-	P	IIIB3
S-153	CCC	IMI	-	P	IIIB3
S-154	CCC	C	-	P	IIIB3
S-155	CCC	AB	-	P	IIIB3
S-156	CCC	F	-	P	IIIB3
S-157	CCC	AC	AB	P	IIIB3
S-158	CCC	TMX	AB	P	IIIB3
S-159	CCC	IMI	AB	P	IIIB3
S-160	CCC	C	AB	P	IIIB3
S-161	CCC	AC	F	P	IIIB3
S-162	CCC	TMX	F	P	IIIB3
S-163	CCC	IMI	F	P	IIIB3
S-164	CCC	C	F	P	IIIB3
S-165	CCC	F	AB	P	IIIB3
S-166	CCC	AC	-	P	IIIB4
S-167	CCC	TMX	-	P	IIIB4
S-168	CCC	IMI	-	P	IIIB4
S-169	CCC	C	-	P	IIIB4
S-170	CCC	AB	-	P	IIIB4

No	I	II (1)	II (2)	IIIA	IIIB
S-171	CCC	F	-	P	IIIB4
S-172	CCC	AC	AB	P	IIIB4
S-173	CCC	TMX	AB	P	IIIB4
S-174	CCC	IMI	AB	P	IIIB4
S-175	CCC	C	AB	P	IIIB4
S-176	CCC	AC	F	P	IIIB4
S-177	CCC	TMX	F	P	IIIB4
S-178	CCC	IMI	F	P	IIIB4
S-179	CCC	C	F	P	IIIB4
S-180	CCC	F	AB	P	IIIB4
S-181	CCC	AC	-	O	IIIB1
S-182	CCC	TMX	-	O	IIIB1
S-183	CCC	IMI	-	O	IIIB1
S-184	CCC	C	-	O	IIIB1
S-185	CCC	AB	-	O	IIIB1
S-186	CCC	F	-	O	IIIB1
S-187	CCC	AC	AB	O	IIIB1
S-188	CCC	TMX	AB	O	IIIB1
S-189	CCC	IMI	AB	O	IIIB1
S-190	CCC	C	AB	O	IIIB1
S-191	CCC	AC	F	O	IIIB1
S-192	CCC	TMX	F	O	IIIB1
S-193	CCC	IMI	F	O	IIIB1
S-194	CCC	C	F	O	IIIB1
S-195	CCC	F	AB	O	IIIB1
S-196	CCC	AC	-	O	IIIB2
S-197	CCC	TMX	-	O	IIIB2
S-198	CCC	IMI	-	O	IIIB2
S-199	CCC	C	-	O	IIIB2
S-200	CCC	AB	-	O	IIIB2
S-201	CCC	F	-	O	IIIB2
S-202	CCC	AC	AB	O	IIIB2
S-203	CCC	TMX	AB	O	IIIB2
S-204	CCC	IMI	AB	O	IIIB2
S-205	CCC	C	AB	O	IIIB2
S-206	CCC	AC	F	O	IIIB2
S-207	CCC	TMX	F	O	IIIB2
S-208	CCC	IMI	F	O	IIIB2
S-209	CCC	C	F	O	IIIB2
S-210	CCC	F	AB	O	IIIB2

No	I	II (1)	II (2)	IIIA	IIIB
S-211	CCC	AC	-	O	IIIB3
S-212	CCC	TMX	-	O	IIIB3
S-213	CCC	IMI	-	O	IIIB3
S-214	CCC	C	-	O	IIIB3
S-215	CCC	AB	-	O	IIIB3
S-216	CCC	F	-	O	IIIB3
S-217	CCC	AC	AB	O	IIIB3
S-218	CCC	TMX	AB	O	IIIB3
S-219	CCC	IMI	AB	O	IIIB3
S-220	CCC	C	AB	O	IIIB3
S-221	CCC	AC	F	O	IIIB3
S-222	CCC	TMX	F	O	IIIB3
S-223	CCC	IMI	F	O	IIIB3
S-224	CCC	C	F	O	IIIB3
S-225	CCC	F	AB	O	IIIB3
S-226	CCC	AC	-	O	IIIB4
S-227	CCC	TMX	-	O	IIIB4
S-228	CCC	IMI	-	O	IIIB4
S-229	CCC	C	-	O	IIIB4
S-230	CCC	AB	-	O	IIIB4
S-231	CCC	F	-	O	IIIB4
S-232	CCC	AC	AB	O	IIIB4
S-233	CCC	TMX	AB	O	IIIB4
S-234	CCC	IMI	AB	O	IIIB4
S-235	CCC	C	AB	O	IIIB4
S-236	CCC	AC	F	O	IIIB4
S-237	CCC	TMX	F	O	IIIB4
S-238	CCC	IMI	F	O	IIIB4
S-239	CCC	C	F	O	IIIB4
S-240	CCC	F	AB	O	IIIB4
S-241	CCC	AC	-	A	IIIB1
S-242	CCC	TMX	-	A	IIIB1
S-243	CCC	IMI	-	A	IIIB1
S-244	CCC	C	-	A	IIIB1
S-245	CCC	AB	-	A	IIIB1
S-246	CCC	F	-	A	IIIB1
S-247	CCC	AC	AB	A	IIIB1
S-248	CCC	TMX	AB	A	IIIB1
S-249	CCC	IMI	AB	A	IIIB1
S-250	CCC	C	AB	A	IIIB1

No	I	II (1)	II (2)	IIIA	IIIB
S-251	CCC	AC	F	A	IIIB1
S-252	CCC	TMX	F	A	IIIB1
S-253	CCC	IMI	F	A	IIIB1
S-254	CCC	C	F	A	IIIB1
S-255	CCC	F	AB	A	IIIB1
S-256	CCC	AC	-	A	IIIB2
S-257	CCC	TMX	-	A	IIIB2
S-258	CCC	IMI	-	A	IIIB2
S-259	CCC	C	-	A	IIIB2
S-260	CCC	AB	-	A	IIIB2
S-261	CCC	F	-	A	IIIB2
S-262	CCC	AC	AB	A	IIIB2
S-263	CCC	TMX	AB	A	IIIB2
S-264	CCC	IMI	AB	A	IIIB2
S-265	CCC	C	AB	A	IIIB2
S-266	CCC	AC	F	A	IIIB2
S-267	CCC	TMX	F	A	IIIB2
S-268	CCC	IMI	F	A	IIIB2
S-269	CCC	C	F	A	IIIB2
S-270	CCC	F	AB	A	IIIB2
S-271	CCC	AC	-	A	IIIB3
S-272	CCC	TMX	-	A	IIIB3
S-273	CCC	IMI	-	A	IIIB3
S-274	CCC	C	-	A	IIIB3
S-275	CCC	AB	-	A	IIIB3
S-276	CCC	F	-	A	IIIB3
S-277	CCC	AC	AB	A	IIIB3
S-278	CCC	TMX	AB	A	IIIB3
S-279	CCC	IMI	AB	A	IIIB3
S-280	CCC	C	AB	A	IIIB3
S-281	CCC	AC	F	A	IIIB3
S-282	CCC	TMX	F	A	IIIB3
S-283	CCC	IMI	F	A	IIIB3
S-284	CCC	C	F	A	IIIB3
S-285	CCC	F	AB	A	IIIB3
S-286	CCC	AC	-	A	IIIB4
S-287	CCC	TMX	-	A	IIIB4
S-288	CCC	IMI	-	A	IIIB4
S-289	CCC	C	-	A	IIIB4
S-290	CCC	AB	-	A	IIIB4

No	I	II (1)	II (2)	IIIA	IIIB
S-291	CCC	F	-	A	IIIB4
S-292	CCC	AC	AB	A	IIIB4
S-293	CCC	TMX	AB	A	IIIB4
S-294	CCC	IMI	AB	A	IIIB4
S-295	CCC	C	AB	A	IIIB4
S-296	CCC	AC	F	A	IIIB4
S-297	CCC	TMX	F	A	IIIB4
S-298	CCC	IMI	F	A	IIIB4
S-299	CCC	C	F	A	IIIB4
S-300	CCC	F	AB	A	IIIB4
S-301	CCC	AC	-	T	IIIB1
S-302	CCC	TMX	-	T	IIIB1
S-303	CCC	IMI	-	T	IIIB1
S-304	CCC	C	-	T	IIIB1
S-305	CCC	AB	-	T	IIIB1
S-306	CCC	F	-	T	IIIB1
S-307	CCC	AC	AB	T	IIIB1
S-308	CCC	TMX	AB	T	IIIB1
S-309	CCC	IMI	AB	T	IIIB1
S-310	CCC	C	AB	T	IIIB1
S-311	CCC	AC	F	T	IIIB1
S-312	CCC	TMX	F	T	IIIB1
S-313	CCC	IMI	F	T	IIIB1
S-314	CCC	C	F	T	IIIB1
S-315	CCC	F	AB	T	IIIB1
S-316	CCC	AC	-	T	IIIB2
S-317	CCC	TMX	-	T	IIIB2
S-318	CCC	IMI	-	T	IIIB2
S-319	CCC	C	-	T	IIIB2
S-320	CCC	AB	-	T	IIIB2
S-321	CCC	F	-	T	IIIB2
S-322	CCC	AC	AB	T	IIIB2
S-323	CCC	TMX	AB	T	IIIB2
S-324	CCC	IMI	AB	T	IIIB2
S-325	CCC	C	AB	T	IIIB2
S-326	CCC	AC	F	T	IIIB2
S-327	CCC	TMX	F	T	IIIB2
S-328	CCC	IMI	F	T	IIIB2
S-329	CCC	C	F	T	IIIB2
S-330	CCC	F	AB	T	IIIB2

No	I	II (1)	II (2)	IIIA	IIIB
S-331	CCC	AC	-	T	IIIB3
S-332	CCC	TMX	-	T	IIIB3
S-333	CCC	IMI	-	T	IIIB3
S-334	CCC	C	-	T	IIIB3
S-335	CCC	AB	-	T	IIIB3
S-336	CCC	F	-	T	IIIB3
S-337	CCC	AC	AB	T	IIIB3
S-338	CCC	TMX	AB	T	IIIB3
S-339	CCC	IMI	AB	T	IIIB3
S-340	CCC	C	AB	T	IIIB3
S-341	CCC	AC	F	T	IIIB3
S-342	CCC	TMX	F	T	IIIB3
S-343	CCC	IMI	F	T	IIIB3
S-344	CCC	C	F	T	IIIB3
S-345	CCC	F	AB	T	IIIB3
S-346	CCC	AC	-	T	IIIB4
S-347	CCC	TMX	-	T	IIIB4
S-348	CCC	IMI	-	T	IIIB4
S-349	CCC	C	-	T	IIIB4
S-350	CCC	AB	-	T	IIIB4
S-351	CCC	F	-	T	IIIB4
S-352	CCC	AC	AB	T	IIIB4
S-353	CCC	TMX	AB	T	IIIB4
S-354	CCC	IMI	AB	T	IIIB4
S-355	CCC	C	AB	T	IIIB4
S-356	CCC	AC	F	T	IIIB4
S-357	CCC	TMX	F	T	IIIB4
S-358	CCC	IMI	F	T	IIIB4
S-359	CCC	C	F	T	IIIB4
S-360	CCC	F	AB	T	IIIB4

Within these mixtures, the following are especially preferred: S-1, S-2, S-3, S-4, S-5, S-6, S-7, S-8, S-9, S-10, S-11, S-12, S-13, S-14, S-15, S-61, S-62, S-63, S-64, S-65, S-66, S-67, S-68, S-69, S-70, S-71, S-72, S-73, S-74, S-75, S-121, S-122, S-123, S-124, S-125, S-126, S-127, S-128, S-129, S-130, S-131, S-132, S-133, S-134 and S-135.

The inventive mixtures can further comprise one or more insecticides, fungicides, herbicides.

For use according to the present invention, the mixtures according to the invention can be converted into the customary formulations, for example solutions, emulsions, suspensions, dusts, powders, pastes and granules. The use form depends on the particular intended purpose; in each case, it should ensure a fine and even distribution of the mixtures according to the present invention. The formulations are prepared in a known manner (cf. US 3,060,084, EP-A 707 445 (for liquid concentrates), Browning: "Agglomeration", Chemical Engineering, Dec. 4, 1967, 147-48, Perry's Chemical Engineer's Handbook, 4th Ed., McGraw-Hill, New York, 1963, S. 8-57 und ff. WO 91/13546, US 4,172,714, US 4,144,050, US 3,920,442, US 5,180,587, US 5,232,701, US 5,208,030, GB 2,095,558, US 3,299,566, Klingman: Weed Control as a Science (J. Wiley & Sons, New York, 1961), Hance et al.: Weed Control Handbook (8th Ed., Blackwell Scientific, Oxford, 1989) and Mollet, H. and Grubemann, A.: Formulation technology (Wiley VCH Verlag, Weinheim, 2001).

The agrochemical formulations may also comprise auxiliaries which are customary in agrochemical formulations. The auxiliaries used depend on the particular application form and active substance, respectively.

Examples for suitable auxiliaries are solvents, solid carriers, dispersants or emulsifiers (such as further solubilizers, protective colloids, surfactants and adhesion agents), organic and anorganic thickeners, bactericides, anti-freezing agents, anti-foaming agents, if appropriate colorants and tackifiers or binders (e. g. for seed treatment formulations).

Suitable solvents are water, organic solvents such as mineral oil fractions of medium to high boiling point, such as kerosene or diesel oil, furthermore coal tar oils and oils of vegetable or animal origin, aliphatic, cyclic and aromatic hydrocarbons, e. g. toluene, xylene, paraffin, tetrahydronaphthalene, alkylated naphthalenes or their derivatives, alcohols such as methanol, ethanol, propanol, butanol and cyclohexanol, glycols, ketones such as cyclohexanone and gamma-butyrolactone, fatty acid dimethylamides, fatty acids and fatty acid esters and strongly polar solvents, e. g. amines such as N-methylpyrrolidone.

Solid carriers are mineral earths such as silicates, silica gels, talc, kaolins, limestone, lime, chalk, bole, loess, clays, dolomite, diatomaceous earth, calcium sulfate, magnesium sulfate, magnesium oxide, ground synthetic materials, fertilizers, such as, e. g., ammonium sulfate, ammonium phosphate, ammonium nitrate, ureas, and products of vegetable origin, such as cereal meal, tree bark meal, wood meal and nutshell meal, cellulose powders and other solid carriers.

Suitable surfactants (adjuvants, witters, tackifiers, dispersants or emulsifiers) are alkali metal, alkaline earth metal and ammonium salts of aromatic sulfonic acids, such as ligninsulfonic acid (Borresperse® types, Borregard, Norway) phenolsulfonic acid, naphthalenesulfonic acid (Morwet® types, Akzo Nobel, U.S.A.), dibutyl-naphthalene-sulfonic acid (Nekal® types, BASF, Germany), and fatty acids, alkylsulfonates, alkylarylsulfonates, alkyl sulfates, laurylether sulfates, fatty alcohol sulfates, and sulfated hexa-, hepta- and octadecanolates, sulfated fatty alcohol glycol ethers, furthermore

condensates of naphthalene or of naphthalenesulfonic acid with phenol and formaldehyde, polyoxy-ethylene octylphenyl ether, ethoxylated isooctylphenol, octylphenol, nonylphenol, alkylphenyl polyglycol ethers, tributylphenyl polyglycol ether, tristearylphenyl polyglycol ether, alkylaryl polyether alcohols, alcohol and fatty alcohol/ethylene oxide condensates, ethoxylated castor oil, polyoxyethylene alkyl ethers, ethoxylated polyoxypropylene, lauryl alcohol polyglycol ether acetal, sorbitol esters, lignin-sulfite waste liquors and proteins, denatured proteins, polysaccharides (e. g. methylcellulose), hydrophobically modified starches, polyvinyl alcohols (Mowiol® types, Clariant, Switzerland), polycarboxylates (Sokolan® types, BASF, Germany), polyalkoxylates, polyvinylamines (Lupasol® types, BASF, Germany), polyvinylpyrrolidone and the copolymers thereof.

Examples for thickeners (i. e. compounds that impart a modified flowability to formulations, i. e. high viscosity under static conditions and low viscosity during agitation) are polysaccharides and organic and anorganic clays such as Xanthan gum (Kelzan®, CP Kelco, U.S.A.), Rhodopol® 23 (Rhodia, France), Veegum® (R.T. Vanderbilt, U.S.A.) or Attaclay® (Engelhard Corp., NJ, USA).

Bactericides may be added for preservation and stabilization of the formulation. Examples for suitable bactericides are those based on dichlorophene and benzylalcohol hemi formal (Proxel® from ICI or Acticide® RS from Thor Chemie and Kathon® MK from Rohm & Haas) and isothiazolinone derivatives such as alkylisothiazolinones and benzisothiazolinones (Acticide® MBS from Thor Chemie).

Examples for suitable anti-freezing agents are ethylene glycol, propylene glycol, urea and glycerin.

Examples for anti-foaming agents are silicone emulsions (such as e. g. Silikon® SRE, Wacker, Germany or Rhodorsil®, Rhodia, France), long chain alcohols, fatty acids, salts of fatty acids, fluoroorganic compounds and mixtures thereof.

Suitable colorants are pigments of low water solubility and water-soluble dyes. Examples to be mentioned und the designations rhodamin B, C. I. pigment red 112, C. I. solvent red 1, pigment blue 15:4, pigment blue 15:3, pigment blue 15:2, pigment blue 15:1, pigment blue 80, pigment yellow 1, pigment yellow 13, pigment red 112, pigment red 48:2, pigment red 48:1, pigment red 57:1, pigment red 53:1, pigment orange 43, pigment orange 34, pigment orange 5, pigment green 36, pigment green 7, pigment white 6, pigment brown 25, basic violet 10, basic violet 49, acid red 51, acid red 52, acid red 14, acid blue 9, acid yellow 23, basic red 10, basic red 108.

Examples for tackifiers or binders are polyvinylpyrrolidons, polyvinylacetates, polyvinyl alcohols and cellulose ethers (Tylose®, Shin-Etsu, Japan).

Powders, materials for spreading and dusts can be prepared by mixing or concomitantly grinding the compounds I and/or II and, if appropriate, further active substances, with at least one solid carrier.

Granules, e. g. coated granules, impregnated granules and homogeneous granules, can be prepared by binding the active substances to solid carriers. Examples of solid carriers are mineral earths such as silica gels, silicates, talc, kaolin, attaclay, limestone,

lime, chalk, bole, loess, clay, dolomite, diatomaceous earth, calcium sulfate, magnesium sulfate, magnesium oxide, ground synthetic materials, fertilizers, such as, e. g., ammonium sulfate, ammonium phosphate, ammonium nitrate, ureas, and products of vegetable origin, such as cereal meal, tree bark meal, wood meal and nutshell meal,
5 cellulose powders and other solid carriers.

Examples for formulation types are:

1. Composition types for dilution with water

i) Water-soluble concentrates (SL, LS)

10 10 parts by weight of compounds of the inventive mixtures are dissolved in 90 parts by weight of water or in a water-soluble solvent. As an alternative, wetting agents or other auxiliaries are added. The active substance dissolves upon dilution with water. In this way, a formulation having a content of 10% by weight of active substance is obtained.

ii) Dispersible concentrates (DC)

15 20 parts by weight of compounds of the inventive mixtures are dissolved in 70 parts by weight of cyclohexanone with addition of 10 parts by weight of a dispersant, e. g. polyvinylpyrrolidone. Dilution with water gives a dispersion. The active substance content is 20% by weight.

iii) Emulsifiable concentrates (EC)

20 15 parts by weight of compounds of the inventive mixtures are dissolved in 75 parts by weight of xylene with addition of calcium dodecylbenzenesulfonate and castor oil ethoxylate (in each case 5 parts by weight). Dilution with water gives an emulsion. The composition has an active substance content of 15% by weight.

iv) Emulsions (EW, EO, ES)

25 25 parts by weight of compounds of the inventive mixtures are dissolved in 35 parts by weight of xylene with addition of calcium dodecylbenzenesulfonate and castor oil ethoxylate (in each case 5 parts by weight). This mixture is introduced into 30 parts by weight of water by means of an emulsifying machine (Ultraturrax) and made into a homogeneous emulsion. Dilution with water gives an emulsion. The composition has an
30 active substance content of 25% by weight.

v) Suspensions (SC, OD, FS)

In an agitated ball mill, 20 parts by weight of compounds of the inventive mixtures are comminuted with addition of 10 parts by weight of dispersants and wetting agents and
35 70 parts by weight of water or an organic solvent to give a fine active substance suspension. Dilution with water gives a stable suspension of the active substance. The active substance content in the composition is 20% by weight.

vi) Water-dispersible granules and water-soluble granules (WG, SG)

40 50 parts by weight of compounds of the inventive mixtures are ground finely with addition of 50 parts by weight of dispersants and wetting agents and prepared as water-dispersible or water-soluble granules by means of technical appliances (e. g. extrusion, spray tower, fluidized bed). Dilution with water gives a stable dispersion or solution of

the active substance. The composition has an active substance content of 50% by weight.

vii) Water-dispersible powders and water-soluble powders (WP, SP, SS, WS)

75 parts by weight of compounds of the inventive mixtures are ground in a rotor-stator mill with addition of 25 parts by weight of dispersants, wetting agents and silica gel. Dilution with water gives a stable dispersion or solution of the active substance. The active substance content of the composition is 75% by weight.

viii) Gel (GF)

In an agitated ball mill, 20 parts by weight of compounds of the inventive mixtures are comminuted with addition of 10 parts by weight of dispersants, 1 part by weight of a gelling agent wetters and 70 parts by weight of water or of an organic solvent to give a fine suspension of the active substance. Dilution with water gives a stable suspension of the active substance, whereby a composition with 20% (w/w) of active substance is obtained.

2. Composition types to be applied undiluted

ix) Dustable powders (DP, DS)

5 parts by weight of compounds of the inventive mixtures are ground finely and mixed intimately with 95 parts by weight of finely divided kaolin. This gives a dustable composition having an active substance content of 5% by weight.

x) Granules (GR, FG, GG, MG)

0.5 parts by weight of compounds of the inventive mixtures is ground finely and associated with 99.5 parts by weight of carriers. Current methods are extrusion, spray-drying or the fluidized bed. This gives granules to be applied undiluted having an active substance content of 0.5% by weight.

xi) ULV solutions (UL)

10 parts by weight of compounds of the inventive mixtures are dissolved in 90 parts by weight of an organic solvent, e. g. xylene. This gives a composition to be applied undiluted having an active substance content of 10% by weight.

The agrochemical formulations generally comprise between 0.01 and 95%, preferably between 0.1 and 90%, most preferably between 0.5 and 90%, by weight of active substances. The compounds of the inventive mixtures are employed in a purity of from 90% to 100%, preferably from 95% to 100% (according to NMR spectrum).

The compounds of the inventive mixtures can be used as such or in the form of their compositions, e. g. in the form of directly sprayable solutions, powders, suspensions, dispersions, emulsions, oil dispersions, pastes, dustable products, materials for spreading, or granules, by means of spraying, atomizing, dusting, spreading, brushing, immersing or pouring. The application forms depend entirely on the intended purposes; it is intended to ensure in each case the finest possible distribution of the compounds present in the inventive mixtures.

Aqueous application forms can be prepared from emulsion concentrates, pastes or wettable powders (sprayable powders, oil dispersions) by adding water. To prepare emulsions, pastes or oil dispersions, the substances, as such or dissolved in an oil or solvent, can be homogenized in water by means of a wetter, tackifier, dispersant or emulsifier. Alternatively, it is possible to prepare concentrates composed of active substance, wetter, tackifier, dispersant or emulsifier and, if appropriate, solvent or oil, and such concentrates are suitable for dilution with water.

The active substance concentrations in the ready-to-use preparations can be varied within relatively wide ranges. In general, they are from 0.0001 to 10%, preferably from 0.001 to 1% by weight of compounds of the inventive mixtures .

The compounds of the inventive mixtures may also be used successfully in the ultra-low-volume process (ULV), it being possible to apply compositions comprising over 95% by weight of active substance, or even to apply the active substance without additives.

Various types of oils, wetters, adjuvants, herbicides, fungicides, other pesticides, or bactericides may be added to the active compounds, if appropriate not until immediately prior to use (tank mix). These agents can be admixed with the compounds of the inventive mixtures in a weight ratio of 1:100 to 100:1, preferably 1:10 to 10:1.

Compositions of this invention may also contain fertilizers such as ammonium nitrate, urea, potash, and superphosphate, phytotoxicants and plant growth regulators and safeners. These may be used sequentially or in combination with the above-described compositions, if appropriate also added only immediately prior to use (tank mix). For example, the plant(s) may be sprayed with a composition of this invention either before or after being treated with the fertilizers.

The compounds contained in the mixtures as defined above can be applied simultaneously, that is jointly or separately, or in succession, the sequence, in the case of separate application, generally not having any effect on the result of the control measures.

According to this invention, the compound I and compound II (and optionally compound III) is to be understood to denote, that at least the compound I and compound II (and optionally compound III) occur simultaneously at the site of action (i.e. the pests, such as harmful fungi and animal pests such as insects, arachnids or nematods to be controlled or their habitats such as infected plants, plant propagation materials, particularly seeds, surfaces, materials or the soil as well as plants, plant propagation materials, particularly seeds, soil, surfaces, materials or rooms to be protected from fungal or animal attack) in a effective amount.

This can be obtained by applying the compound I and compound II (and optionally compound III) simultaneously, either jointly (e. g. as tank-mix) or sperately, or in succession, wherein the time interval between the individual applications is selected to ensure that the active substance applied first still occurs at the site of action in a sufficient amount at the time of application of the further active substance(s). The order of application is not essential for working of the present invention.

In binary, ternary and quaternary mixture of the present invention, the weight ratio of the compounds generally depends from the properties of the compounds of the inventive mixtures.

The compounds of the inventive mixtures can be used individually or already partially or completely mixed with one another to prepare the composition according to the invention. It is also possible for them to be packaged and used further as combination composition such as a kit of parts.

In one embodiment of the invention, the kits may include one or more, including all, components that may be used to prepare a subject agrochemical composition. E. g., kits may include the compound I and compound II (and optionally compound III) and/or an adjuvant component and/or a further pesticidal compound (e.g. insecticide or herbicide) and/or a growth regulator component). One or more of the components may already be combined together or pre-formulated. In those embodiments where more than two components are provided in a kit, the components may already be combined together and as such are packaged in a single container such as a vial, bottle, can, pouch, bag or canister. In other embodiments, two or more components of a kit may be packaged separately, i. e., not pre-formulated. As such, kits may include one or more separate containers such as vials, cans, bottles, pouches, bags or canisters, each container containing a separate component for an agrochemical composition. In both forms, a component of the kit may be applied separately from or together with the further components or as a component of a combination composition according to the invention for preparing the composition according to the invention.

The user applies the composition according to the invention usually from a predosage device, a knapsack sprayer, a spray tank or a spray plane. Here, the agrochemical composition is made up with water and/or buffer to the desired application concentration, it being possible, if appropriate, to add further auxiliaries, and the ready-to-use spray liquor or the agrochemical composition according to the invention is thus obtained. Usually, 50 to 500 liters of the ready-to-use spray liquor are applied per hectare of agricultural useful area, preferably 100 to 400 liters.

According to one embodiment, individual compounds of the inventive mixtures formulated as composition (or formulation) such as parts of a kit or parts of a binary or ter-

nary or quaternary mixture may be mixed by the user himself in a spray tank and further auxiliaries may be added, if appropriate (tank mix).

- 5 In a further embodiment, either individual compounds of the inventive mixtures formulated as composition or partially premixed components, e. g. components comprising the compound I and compound II (and optionally compound III) may be mixed by the user in a spray tank and further auxiliaries and additives may be added, if appropriate (tank mix).
- 10 In a further embodiment, either individual components of the composition according to the invention or partially premixed components, e. g. components comprising the compound I and compound II (and optionally compound III and/or [compound IV or V]), can be applied jointly (e. .g. after tankmix) or consecutively.
- 15 As said above, the present invention comprises a method for controlling pests, that means animal pests and harmful fungi, wherein the pest, their habitat, breeding grounds, their locus or the plants to be protected against pest attack, the soil or plant propagation material (preferably seed) are treated with an pesticidally effective amount of a mixture.
- 20 Advantageously, the inventive mixtures are suitable for controlling the following fungal plant diseases (harmful fungi):
Albugo spp. (white rust) on ornamentals, vegetables (e. g. *A. candida*) and sunflowers (e. g. *A. tragopogonis*); *Alternaria* spp. (*Alternaria* leaf spot) on vegetables, rape (*A. brassicola* or *brassicae*), sugar beets (*A. tenuis*), fruits, rice, soybeans, potatoes (e. g. *A. solani* or *A. alternata*), tomatoes (e. g. *A. solani* or *A. alternata*) and wheat; *Aphanomyces* spp. on sugar beets and vegetables; *Ascochyta* spp. on cereals and vegetables, e. g. *A. tritici* (anthracnose) on wheat and *A. hordei* on barley; *Bipolaris* and *Drechslera* spp. (teleomorph: *Cochliobolus* spp.), e. g. Southern leaf blight (*D. maydis*) or Northern
30 leaf blight (*B. zeicola*) on corn, e. g. spot blotch (*B. sorokiniana*) on cereals and e.g. *B. oryzae* on rice and turfs; *Blumeria* (formerly *Erysiphe*) *graminis* (powdery mildew) on cereals (e. g. on wheat or barley); *Botrytis cinerea* (teleomorph: *Botryotinia fuckeliana*: grey mold) on fruits and berries (e. g. strawberries), vegetables (e. g. lettuce, carrots, celery and cabbages), rape, flowers, vines, forestry plants and wheat; *Bremia lactucae*
35 (downy mildew) on lettuce; *Ceratocystis* (syn. *Ophiostoma*) spp. (rot or wilt) on broad-leaved trees and evergreens, e. g. *C. ulmi* (Dutch elm disease) on elms; *Cercospora* spp. (*Cercospora* leaf spots) on corn (e.g. Gray leaf spot: *C. zea-maydis*), rice, sugar beets (e. g. *C. beticola*), sugar cane, vegetables, coffee, soybeans (e. g. *C. sojae* or *C. kikuchii*) and rice; *Cladosporium* spp. on tomatoes (e. g. *C. fulvum*: leaf mold) and cereals, e. g. *C. herbarum* (black ear) on wheat; *Claviceps purpurea* (ergot) on cereals;
40 *Cochliobolus* (anamorph: *Helminthosporium* of *Bipolaris*) spp. (leaf spots) on corn (*C. carbonum*), cereals (e. g. *C. sativus*, anamorph: *B. sorokiniana*) and rice (e. g. *C. miy-*

abeanus, anamorph: *H. oryzae*); *Colletotrichum* (teleomorph: *Glomerella*) spp. (anthracnose) on cotton (e. g. *C. gossypii*), corn (e. g. *C. graminicola*: Anthracnose stalk rot), soft fruits, potatoes (e. g. *C. coccodes*: black dot), beans (e. g. *C. lindemuthianum*) and soybeans (e. g. *C. truncatum* or *C. gloeosporioides*); *Corticium* spp., e. g. *C. sasakii* (sheath blight) on rice; *Corynespora cassiicola* (leaf spots) on soybeans and ornamentals; *Cyloconium* spp., e. g. *C. oleaginum* on olive trees; *Cylindrocarpon* spp. (e. g. fruit tree canker or young vine decline, teleomorph: *Nectria* or *Neonectria* spp.) on fruit trees, vines (e. g. *C. liriodendri*, teleomorph: *Neonectria liriodendri*: Black Foot Disease) and ornamentals; *Dematophora* (teleomorph: *Rosellinia*) *necatrix* (root and stem rot) on soybeans; *Diaporthe* spp., e. g. *D. phaseolorum* (damping off) on soybeans; *Drechslera* (syn. *Helminthosporium*, teleomorph: *Pyrenophora*) spp. on corn, cereals, such as barley (e. g. *D. teres*, net blotch) and wheat (e. g. *D. tritici-repentis*: tan spot), rice and turf; Esca (dieback, apoplexy) on vines, caused by *Formitiporia* (syn. *Phellinus*) *punctata*, *F. mediterranea*, *Phaeomonilla chlamydospora* (earlier *Phaeoacremonium chlamydosporum*), *Phaeoacremonium aleophilum* and/or *Botryosphaeria obtusa*; *Elsinoe* spp. on pome fruits (*E. pyri*), soft fruits (*E. veneta*: anthracnose) and vines (*E. ampelina*: anthracnose); *Entyloma oryzae* (leaf smut) on rice; *Epicoccum* spp. (black mold) on wheat; *Erysiphe* spp. (powdery mildew) on sugar beets (*E. betae*), vegetables (e. g. *E. pisi*), such as cucurbits (e. g. *E. cichoracearum*), cabbages, rape (e. g. *E. cruciferarum*); *Eutypa lata* (*Eutypa* canker or dieback, anamorph: *Cytosporina lata*, syn. *Libertella blepharis*) on fruit trees, vines and ornamental woods; *Exserohilum* (syn. *Helminthosporium*) spp. on corn (e. g. *E. turcicum*); *Fusarium* (teleomorph: *Gibberella*) spp. (wilt, root or stem rot) on various plants, such as *F. graminearum* or *F. culmorum* (root rot, scab or head blight) on cereals (e. g. wheat or barley), *F. oxysporum* on tomatoes, *F. solani* on soybeans and *F. verticillioides* on corn; *Gaeumannomyces graminis* (take-all) on cereals (e. g. wheat or barley) and corn; *Gibberella* spp. on cereals (e. g. *G. zeae*) and rice (e. g. *G. fujikuroi*: Bakanae disease); *Glomerella cingulata* on vines, pome fruits and other plants and *G. gossypii* on cotton; Grain-staining complex on rice; *Guignardia bidwellii* (black rot) on vines; *Gymnosporangium* spp. on rosaceous plants and junipers, e. g. *G. sabinae* (rust) on pears; *Helminthosporium* spp. (syn. *Drechslera*, teleomorph: *Cochliobolus*) on corn, cereals and rice; *Hemileia* spp., e. g. *H. vastatrix* (coffee leaf rust) on coffee; *Isariopsis clavispora* (syn. *Cladosporium vitis*) on vines; *Macrophomina phaseolina* (syn. *phaseoli*) (root and stem rot) on soybeans and cotton; *Microdochium* (syn. *Fusarium*) *nivale* (pink snow mold) on cereals (e. g. wheat or barley); *Microsphaera diffusa* (powdery mildew) on soybeans; *Monilinia* spp., e. g. *M. laxa*, *M. fructicola* and *M. fructigena* (bloom and twig blight, brown rot) on stone fruits and other rosaceous plants; *Mycosphaerella* spp. on cereals, bananas, soft fruits and ground nuts, such as e. g. *M. graminicola* (anamorph: *Septoria tritici*, *Septoria blotch*) on wheat or *M. fijiensis* (black Sigatoka disease) on bananas; *Peronospora* spp. (downy mildew) on cabbage (e. g. *P. brassicae*), rape (e. g. *P. parasitica*), onions (e. g. *P. destructor*), tobacco (*P. tabacina*) and soybeans (e. g. *P. manshurica*); *Phakopsora pachyrhizi* and *P. meibomia* (soybean rust) on soybeans;

Phialophora spp. e. g. on vines (e. g. *P. tracheiphila* and *P. tetraspora*) and soybeans (e. g. *P. gregata*: stem rot); *Phoma lingam* (root and stem rot) on rape and cabbage and *P. betae* (root rot, leaf spot and damping-off) on sugar beets; *Phomopsis* spp. on sunflowers, vines (e. g. *P. viticola*: can and leaf spot) and soybeans (e. g. stem rot: *P.*
5 *phaseoli*, teleomorph: *Diaporthe phaseolorum*); *Physoderma maydis* (brown spots) on corn; *Phytophthora* spp. (wilt, root, leaf, fruit and stem rot) on various plants, such as paprika and cucurbits (e. g. *P. capsici*), soybeans (e. g. *P. megasperma*, syn. *P. sojae*), potatoes and tomatoes (e. g. *P. infestans*: late blight) and broad-leaved trees (e. g. *P.*
10 *ramorum*: sudden oak death); *Plasmiodiophora brassicae* (club root) on cabbage, rape, radish and other plants; *Plasmopara* spp., e. g. *P. viticola* (grapevine downy mildew) on vines and *P. halstedii* on sunflowers; *Podosphaera* spp. (powdery mildew) on rosa-
ceous plants, hop, pome and soft fruits, e. g. *P. leucotricha* on apples; *Polymyxa* spp.,
e. g. on cereals, such as barley and wheat (*P. graminis*) and sugar beets (*P. betae*)
and thereby transmitted viral diseases; *Pseudocercospora herpotrichoides* (eyespot,
15 teleomorph: *Tapesia yallundae*) on cereals, e. g. wheat or barley; *Pseudoperonospora*
(downy mildew) on various plants, e. g. *P. cubensis* on cucurbits or *P. humili* on hop; *Pseudopezicula tracheiphila* (red fire disease or 'rotbrenner', anamorph: *Phialophora*)
on vines; *Puccinia* spp. (rusts) on various plants, e. g. *P. triticina* (brown or leaf rust), *P.*
20 *striiformis* (stripe or yellow rust), *P. hordei* (dwarf rust), *P. graminis* (stem or black rust)
or *P. recondita* (brown or leaf rust) on cereals, such as e. g. wheat, barley or rye, and
asparagus (e. g. *P. asparagi*); *Pyrenophora* (anamorph: *Drechslera*) *tritici-repentis* (tan
spot) on wheat or *P. teres* (net blotch) on barley; *Pyricularia* spp., e. g. *P. oryzae*
(teleomorph: *Magnaporthe grisea*, rice blast) on rice and *P. grisea* on turf and cereals;
25 *Pythium* spp. (damping-off) on turf, rice, corn, wheat, cotton, rape, sunflowers, soy-
beans, sugar beets, vegetables and various other plants (e. g. *P. ultimum* or *P. aphanidermatum*); *Ramularia* spp., e. g. *R. collo-cygni* (*Ramularia* leaf spots, Physiological
leaf spots) on barley and *R. beticola* on sugar beets; *Rhizoctonia* spp. on cotton, rice,
potatoes, turf, corn, rape, potatoes, sugar beets, vegetables and various other plants,
e. g. *R. solani* (root and stem rot) on soybeans, *R. solani* (sheath blight) on rice or *R.*
30 *cerealis* (*Rhizoctonia* spring blight) on wheat or barley; *Rhizopus stolonifer* (black mold,
soft rot) on strawberries, carrots, cabbage, vines and tomatoes; *Rhynchosporium se-
calis* (scald) on barley, rye and triticale; *Sarocladium oryzae* and *S. attenuatum* (sheath
rot) on rice; *Sclerotinia* spp. (stem rot or white mold) on vegetables and field crops,
such as rape, sunflowers (e. g. *S. sclerotiorum*) and soybeans (e. g. *S. rolfsii* or *S. scler-
35*
rotiorum); *Septoria* spp. on various plants, e. g. *S. glycines* (brown spot) on soybeans,
S. tritici (*Septoria* blotch) on wheat and *S.* (syn. *Stagonospora*) *nodorum* (*Stagono-
spora* blotch) on cereals; *Uncinula* (syn. *Erysiphe*) *necator* (powdery mildew, ana-
morph: *Oidium tuckeri*) on vines; *Setosphaeria* spp. (leaf blight) on corn (e. g. *S.*
40 *turcicum*, syn. *Helminthosporium turcicum*) and turf; *Sphacelotheca* spp. (smut) on
corn, (e. g. *S. reiliana*: head smut), sorghum und sugar cane; *Sphaerotheca fuliginea*
(powdery mildew) on cucurbits; *Spongospora subterranea* (powdery scab) on potatoes
and thereby transmitted viral diseases; *Stagonospora* spp. on cereals, e. g. *S. nodorum*

(Stagonospora blotch, teleomorph: Leptosphaeria [syn. Phaeosphaeria] nodorum) on wheat; Synchytrium endobioticum on potatoes (potato wart disease); Taphrina spp., e. g. T. deformans (leaf curl disease) on peaches and T. pruni (plum pocket) on plums; Thielaviopsis spp. (black root rot) on tobacco, pome fruits, vegetables, soybeans and cotton, e. g. T. basicola (syn. Chalara elegans); Tilletia spp. (common bunt or stinking smut) on cereals, such as e. g. T. tritici (syn. T. caries, wheat bunt) and T. controversa (dwarf bunt) on wheat; Typhula incarnata (grey snow mold) on barley or wheat; Urocystis spp., e. g. U. occulta (stem smut) on rye; Uromyces spp. (rust) on vegetables, such as beans (e. g. U. appendiculatus, syn. U. phaseoli) and sugar beets (e. g. U. betae); Ustilago spp. (loose smut) on cereals (e. g. U. nuda and U. avenae), corn (e. g. U. maydis: corn smut) and sugar cane; Venturia spp. (scab) on apples (e. g. V. inaequalis) and pears; and Verticillium spp. (wilt) on various plants, such as fruits and ornamentals, vines, soft fruits, vegetables and field crops, e. g. V. dahliae on strawberries, rape, potatoes and tomatoes.

15

The inventive mixtures are also suitable for controlling harmful fungi and fungal diseases relevant in the protection of stored products or harvest and in the protection of materials. The term "protection of materials" is to be understood to denote the protection of technical and non-living materials, such as adhesives, glues, wood, paper and paperboard, textiles, leather, paint dispersions, plastics, colling lubricants, fiber or fabrics, against the infestation and destruction by harmful microorganisms, such as fungi and bacteria. As to the protection of wood and other materials, the particular attention is paid to the following harmful fungi: Ascomycetes such as Ophiostoma spp., Ceratocystis spp., Aureobasidium pullulans, Sclerophoma spp., Chaetomium spp., Humicola spp., Petriella spp., Trichurus spp.; Basidiomycetes such as Coniophora spp., Coriolus spp., Gloeophyllum spp., Lentinus spp., Pleurotus spp., Poria spp., Serpula spp. and Tyromyces spp., Deuteromycetes such as Aspergillus spp., Cladosporium spp., Penicillium spp., Trichormia spp., Alternaria spp., Paecilomyces spp. and Zygomycetes such as Mucor spp., and in addition in the protection of stored products and harvest the following yeast fungi are worthy of note: Candida spp. and Saccharomyces cerevisiae.

30

They are particularly important for controlling a multitude of harmful fungi on various cultivated plants, such as bananas, cotton, vegetable species (for example cucumbers, beans and cucurbits), barley, grass, oats, coffee, potatoes, corn, fruit species, rice, rye, soya, tomatoes, grapevines, wheat, ornamental plants, sugar cane and also on a large number of plant propagation materials (preferably seeds).

35

The inventive mixtures exhibit also outstanding action against animal pests from the following orders:

40

insects from the order of the lepidopterans (Lepidoptera), for example Agrotis ypsilon, Agrotis segetum, Alabama argillacea, Anticarsia gemmatalis, Argyresthia conjugella,

Autographa gamma, Bupalus piniarius, Cacoecia murinana, Capua reticulana, Cheimantobia brumata, Choristoneura fumiferana, Choristoneura occidentalis, Cirphis unipuncta, Cydia pomonella, Dendrolimus pini, Diaphania nitidalis, Diatraea grandiosella, Earias insulana, Elasmopalpus lignosellus, Eupoecilia ambiguella, Evetria bouliana, Feltia subterranea, Galleria mellonella, Grapholitha funebrana, Grapholitha molesta, Heliothis armigera, Heliothis virescens, Heliothis zea, Hellula undalis, Hibernia defoliaria, Hyphantria cunea, Hyponomeuta malinellus, Keiferia lycopersicella, Lambdina fiscellaria, Laphygma exigua, Leucoptera coffeella, Leucoptera scitella, Lithocolletis blancardella, Lobesia botrana, Loxostege sticticalis, Lymantria dispar, Lymantria monacha, Lyonetia clerkella, Malacosoma neustria, Mamestra brassicae, Orgyia pseudotsugata, Ostrinia nubilalis, Panolis flammea, Pectinophora gossypiella, Peridroma saucia, Phalera bucephala, Phthorimaea operculella, Phyllocnistis citrella, Pieris brassicae, Plathypena scabra, Plutella xylostella, Pseudoplusia includens, Rhyacionia frugiperda, Scrobipalpus absoluta, Sitotroga cerealella, Sparganothis pilleriana, Spodoptera frugiperda, Spodoptera littoralis, Spodoptera litura, Thaumtopoea pityocampa, Tortrix viridana, Trichoplusia ni and Zeiraphera canadensis,

beetles (Coleoptera), for example Agrilus sinuatus, Agriotes lineatus, Agriotes obscurus, Amphimallus solstitialis, Anisandrus dispar, Anthonomus grandis, Anthonomus pomorum, Apthona euphoridae, Athous haemorrhoidalis, Atomaria linearis, Blastophagus piniperda, Blitophaga undata, Bruchus rufimanus, Bruchus pisorum, Bruchus lentis, Byctiscus betulae, Cassida nebulosa, Cerotoma trifurcata, Cetonia aurata, Ceuthorrhynchus assimilis, Ceuthorrhynchus napi, Chaetocnema tibialis, Conoderus vespertinus, Crioceris asparagi, Ctenicera ssp., Diabrotica longicornis, Diabrotica semipunctata, Diabrotica 12-punctata Diabrotica speciosa, Diabrotica virgifera, Epilachna varivestis, Epitrix hirtipennis, Eutinobothrus brasiliensis, Hylobius abietis, Hypera brunneipennis, Hypera postica, Ips typographus, Lema bilineata, Lema melanopus, Leptinotarsa decemlineata, Limonius californicus, Lissorhoptrus oryzophilus, Melanotus communis, Meligethes aeneus, Melolontha hippocastani, Melolontha melolontha, Oulema oryzae, Ortiorrhynchus sulcatus, Otiorrhynchus ovatus, Phaedon cochleariae, Phyllobius pyri, Phyllostreta chrysocephala, Phyllophaga sp., Phyllopertha horticola, Phyllostreta nemorum, Phyllostreta striolata, Popillia japonica, Sitona lineatus and Sitophilus granaria,

flies, mosquitoes (Diptera), e.g. Aedes aegypti, Aedes albopictus, Aedes vexans, Anastrepha ludens, Anopheles maculipennis, Anopheles crucians, Anopheles albimanus, Anopheles gambiae, Anopheles freeborni, Anopheles leucosphyrus, Anopheles minimus, Anopheles quadrimaculatus, Calliphora vicina, Ceratitis capitata, Chrysomya bezziana, Chrysomya hominivorax, Chrysomya macellaria, Chrysops discalis, Chrysops silacea, Chrysops atlanticus, Cochliomyia hominivorax, Contarinia sorghicola, Cordylobia anthropophaga, Culicoides furens, Culex pipiens, Culex nigripalpus, Culex quinquefasciatus, Culex tarsalis, Culiseta inornata, Culiseta melanura, Dacus cucurbi-

- tae, *Dacus oleae*, *Dasineura brassicae*, *Delia antique*, *Delia coarctata*, *Delia platura*, *Delia radicum*, *Dermatobia hominis*, *Fannia canicularis*, *Geomyza Tripunctata*, *Gasterophilus intestinalis*, *Glossina morsitans*, *Glossina palpalis*, *Glossina fuscipes*, *Glossina tachinoides*, *Haematobia irritans*, *Haplodiplosis equestris*, *Hippelates* spp., *Hylemyia*
- 5 *platura*, *Hypoderma lineata*, *Leptoconops torrens*, *Liriomyza sativae*, *Liriomyza trifolii*, *Lucilia caprina*, *Lucilia cuprina*, *Lucilia sericata*, *Lycoria pectoralis*, *Mansonia titillanus*, *Mayetiola destructor*, *Musca domestica*, *Muscina stabulans*, *Oestrus ovis*, *Opomyza florum*, *Oscinella frit*, *Pegomya hysocyami*, *Phorbia antiqua*, *Phorbia brassicae*, *Phorbia coarctata*, *Phlebotomus argentipes*, *Psorophora columbiae*, *Psila rosae*, *Psorophora*
- 10 *discolor*, *Prosimulium mixtum*, *Rhagoletis cerasi*, *Rhagoletis pomonella*, *Sarcophaga haemorrhoidalis*, *Sarcophaga* sp., *Simulium vittatum*, *Stomoxys calcitrans*, *Tabanus bovinus*, *Tabanus atratus*, *Tabanus lineola*, and *Tabanus similis*, *Tipula ol-eracea*, and *Tipula paludosa*
- 15 *thrips* (Thysanoptera), e.g. *Dichromothrips corbetti*, *Dichromothrips* ssp , *Frankliniella fusca*, *Frankliniella occidentalis*, *Frankliniella tritici*, *Scirtothrips citri*, *Thrips oryzae*, *Thrips palmi* and *Thrips tabaci*,
- 20 *termites* (Isoptera), e.g. *Calotermes flavicollis*, *Leucotermes flavipes*, *Heterotermes aureus*, *Reticulitermes flavipes*, *Reticulitermes virginicus*, *Reticulitermes lucifugus*, *Termes natalensis*, and *Coptotermes formosanus*,
- 25 *cockroaches* (Blattaria - Blattodea), e.g. *Blattella germanica*, *Blattella asahinae*, *Periplaneta americana*, *Periplaneta japonica*, *Periplaneta brunnea*, *Periplaneta fuliginosa*, *Periplaneta australasiae*, and *Blatta orientalis*,
- 30 *true bugs* (Hemiptera), e.g. *Acrosternum hilare*, *Blissus leucopterus*, *Cyrtopeltis notatus*, *Dysdercus cingulatus*, *Dysdercus intermedius*, *Eurygaster integriceps*, *Euschistus impictiventris*, *Leptoglossus phyllopus*, *Lygus lineolaris*, *Lygus pratensis*, *Nezara viridula*, *Piesma quadrata*, *Solubea insularis* , *Thyanta perditor*, *Acyrtosiphon onobrychis*, *Adelges laricis*, *Aphidula nasturtii*, *Aphis fabae*, *Aphis forbesi*, *Aphis pomi*, *Aphis gossypii*, *Aphis grossulariae*, *Aphis schneideri*, *Aphis spiraeicola*, *Aphis sambuci*, *Acyrtosiphon pisum*, *Aulacorthum solani*, *Bemisia argentifolii*, *Brachycaudus cardui*, *Brachycaudus helichrysi*, *Brachycaudus persicae*, *Brachycaudus prunicola*, *Brevicoryne brassicae*, *Capitophorus horni*, *Cerosipha gossypii*, *Chaetosiphon fragaefolii*, *Cryptomyzus ribis*, *Dreyfusia nordmanniana*, *Dreyfusia piceae*, *Dysaphis radicola*, *Dysaulacorthum pseudosolani*, *Dysaphis plantaginea*, *Dysaphis pyri*, *Empoasca fabae*, *Hyalopterus pruni*, *Hyperomyzus lactucae*, *Macrosiphum avenae*, *Macrosiphum euphorbiae*, *Macrosiphum rosae*, *Megoura viciae*, *Melanaphis pyriarius*, *Metopolophium*
- 40 *dirhodum*, *Myzus persicae*, *Myzus ascalonicus*, *Myzus cerasi*, *Myzus varians*, *Nasonovia ribis-nigri*, *Nilaparvata lugens*, *Pemphigus bursarius*, *Perkinsiella saccharicida*, *Phorodon humuli*, *Psylla mali*, *Psylla piri*, *Rhopalomyzus ascalonicus*, *Rhopalosiphum*

maidis, *Rhopalosiphum padi*, *Rhopalosiphum insertum*, *Sappaphis mala*, *Sappaphis mali*, *Schizaphis graminum*, *Schizoneura lanuginosa*, *Sitobion avenae*, *Trialeurodes vaporariorum*, *Toxoptera aurantiiand*, *Viteus vitifolii*, *Cimex lectularius*, *Cimex hemipterus*, *Reduvius senilis*, *Triatoma* spp., and *Arilus critatus*.

5

ants, bees, wasps, sawflies (Hymenoptera), e.g. *Athalia rosae*, *Atta cephalotes*, *Atta capiguara*, *Atta cephalotes*, *Atta laevigata*, *Atta robusta*, *Atta sexdens*, *Atta texana*, *Crematogaster* spp., *Hoplocampa minuta*, *Hoplocampa testudinea*, *Monomorium pharaonis*, *Solenopsis geminata*, *Solenopsis invicta*, *Solenopsis richteri*, *Solenopsis xyloni*,
10 *Pogonomyrmex barbatus*, *Pogonomyrmex californicus*, *Pheidole megacephala*, *Dasy-
mutilla occidentalis*, *Bombus* spp. *Vespula squamosa*, *Paravespula vulgaris*, *Paraves-
pula pennsylvanica*, *Paravespula germanica*, *Dolichovespula maculata*, *Vespa crabro*,
Polistes rubiginosa, *Camponotus floridanus*, and *Linepithema humile*,

15

crickets, grasshoppers, locusts (Orthoptera), e.g. *Acheta domestica*, *Gryllotalpa gryllo-
talpa*, *Locusta migratoria*, *Melanoplus bivittatus*, *Melanoplus femurrubrum*, *Melanoplus
mexicanus*, *Melanoplus sanguinipes*, *Melanoplus spretus*, *Nomadacris septemfasciata*,
Schistocerca americana, *Schistocerca gregaria*, *Dociostaurus maroccanus*, *Tachycines
asynamorus*, *Oedaleus senegalensis*, *Zonozerus variegatus*, *Hieroglyphus daganensis*,
20 *Kraussaria angulifera*, *Calliptamus italicus*, *Chortoicetes terminifera*, and *Locustana
pardalina*,

25

Arachnoidea, such as arachnids (Acarina), e.g. of the families *Argasidae*, *Ixodidae* and
Sarcoptidae, such as *Amblyomma americanum*, *Amblyomma variegatum*, *Ambryomma
maculatum*, *Argas persicus*, *Boophilus annulatus*, *Boophilus decoloratus*, *Boophilus
microplus*, *Dermacentor silvarum*, *Dermacentor andersoni*, *Dermacentor variabilis*,
Hyalomma truncatum, *Ixodes ricinus*, *Ixodes rubicundus*, *Ixodes scapularis*, *Ixodes
holocyclus*, *Ixodes pacificus*, *Ornithodoros moubata*, *Ornithodoros hermsi*, *Orni-
thodoros turicata*, *Ornithonyssus bacoti*, *Otobius megnini*, *Dermanyssus gallinae*, *Pso-
roptes ovis*, *Rhipicephalus sanguineus*, *Rhipicephalus appendiculatus*, *Rhipicephalus
evertsi*, *Sarcoptes scabiei*, and *Eriophyidae* spp. such as *Aculus schlechtendali*, *Phyl-
locoptrata oleivora* and *Eriophyes sheldoni*; *Tarsonemidae* spp. such as *Phytonemus
pallidus* and *Polyphagotarsonemus latus*; *Tenuipalpidae* spp. such as *Brevipalpus
phoenicis*; *Tetranychidae* spp. such as *Tetranychus cinnabarinus*, *Tetranychus kan-
zawai*, *Tetranychus pacificus*, *Tetranychus telarius* and *Tetranychus urticae*, *Panony-
chus ulmi*, *Panonychus citri*, and *Oligonychus pratensis*; *Araneida*, e.g. *Latrodectus
mactans*, and *Loxosceles reclusa*,

35
40

fleas (Siphonaptera), e.g. *Ctenocephalides felis*, *Ctenocephalides canis*, *Xenopsylla
cheopis*, *Pulex irritans*, *Tunga penetrans*, and *Nosopsyllus fasciatus*,

silverfish, firebrat (*Thysanura*), e.g. *Lepisma saccharina* and *Thermobia domestica*,

centipedes (Chilopoda), e.g. *Scutigera coleoptrata*,

millipedes (Diplopoda), e.g. *Narceus* spp.,

5

Earwigs (Dermaptera), e.g. *forficula auricularia*,

lice (Phthiraptera), e.g. *Pediculus humanus capitis*, *Pediculus humanus corporis*, *Pthirus pubis*, *Haematopinus eurysternus*, *Haematopinus suis*, *Linognathus vituli*, *Bovicola bovis*, *Menopon gallinae*, *Menacanthus stramineus* and *Solenopotes capillatus*,

10

plant parasitic nematodes such as root-knot nematodes, *Meloidogyne arenaria*, *Meloidogyne chitwoodi*, *Meloidogyne exigua*, *Meloidogyne hapla*, *Meloidogyne incognita*, *Meloidogyne javanica* and other *Meloidogyne* species; cyst nematodes, *Globodera rostochiensis*, *Globodera pallida*, *Globodera tabacum* and other *Globodera* species, *Heterodera avenae*, *Heterodera glycines*, *Heterodera schachtii*, *Heterodera trifolii*, and other *Heterodera* species; seed gall nematodes, *Anguina funesta*, *Anguina tritici* and other *Anguina* species; stem and foliar nematodes, *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzemabosi* and other *Aphelenchoides* species;

20 sting nematodes, *Belonolaimus longicaudatus* and other *Belonolaimus* species; pine nematodes, *Bursaphelenchus xylophilus* and other *Bursaphelenchus* species; ring nematodes, *Criconema* species, *Criconemella* species, *Criconemoides* species, and *Mesocriconema* species; stem and bulb nematodes, *Ditylenchus destructor*, *Ditylenchus dipsaci*, *Ditylenchus myceliophagus* and other *Ditylenchus* species; awl nematodes,

25 *Dolichodorus* species; spiral nematodes, *Helicotylenchus dihystera*, *Helicotylenchus multicinctus* and other *Helicotylenchus* species, *Rotylenchus robustus* and other *Rotylenchus* species; sheath nematodes, *Hemicycliophora* species and *Hemicriconemoides* species; *Hirshmanniella* species; lance nematodes, *Hoplolaimus columbus*, *Hoplolaimus galeatus* and other *Hoplolaimus* species; false root-knot nematodes, *Nacobbus aberrans* and other *Nacobbus* species; needle nematodes, *Longidorus elongatus* and other *Longidorus* species; pin nematodes, *Pratylenchus* species; lesion nematodes, *Pratylenchus brachyurus*, *Pratylenchus coffeae*, *Pratylenchus curvatus*, *Pratylenchus goodeyi*, *Pratylenchus neglectus*, *Pratylenchus penetrans*, *Pratylenchus scribneri*, *Pratylenchus vulnus*, *Pratylenchus zeae* and other *Pratylenchus* species;

35 *Radinaphelenchus cocophilus* and other *Radinaphelenchus* species; burrowing nematodes, *Radopholus similis* and other *Radopholus* species; reniform nematodes, *Rotylenchulus reniformis* and other *Rotylenchulus* species; *Scutellonema* species; stubby root nematodes, *Trichodorus primitivus* and other *Trichodorus* species; *Paratrichodorus minor* and other *Paratrichodorus* species; stunt nematodes, *Tylenchorhynchus claytoni*,

40 *Tylenchorhynchus dubius* and other *Tylenchorhynchus* species and *Merlinius* species; citrus nematodes, *Tylenchulus semipenetrans* and other *Tylenchulus* species; dagger

nematodes, *Xiphinema americanum*, *Xiphinema index*, *Xiphinema diversicaudatum* and other *Xiphinema* species; and other plant parasitic nematode species.

5 The mixtures according to the invention can be applied to any and all developmental stages of pests, such as egg, larva, pupa, and adult. The pests may be controlled by contacting the target pest, its food supply, habitat, breeding ground or its locus with a pesticidally effective amount of the inventive mixtures or of compositions comprising the mixtures.

10 "Locus" means a plant, plant propagation material (preferably seed), soil, area, material or environment in which a pest is growing or may grow.

In general, "pesticidally effective amount" means the amount of the inventive mixtures or of compositions comprising the mixtures needed to achieve an observable effect on growth, including the effects of necrosis, death, retardation, prevention, and removal, 15 destruction, or otherwise diminishing the occurrence and activity of the target organism. The pesticidally effective amount can vary for the various mixtures / compositions used in the invention. A pesticidally effective amount of the mixtures / compositions will also vary according to the prevailing conditions such as desired pesticidal effect and duration, weather, target species, locus, mode of application, and the like. 20

As said above, the present invention comprises a method for improving the health of plants, wherein the plant, the locus where the plant is growing or is expected to grow or plant propagation material, from which the plant grows, is treated with an plant health 25 effective amount of an inventive mixture.

The term "plant effective amount" denotes an amount of the inventive mixtures, which is sufficient for achieving plant health effects as defined hereinbelow. More exemplary information about amounts, ways of application and suitable ratios to be used is given 30 below. Anyway, the skilled artisan is well aware of the fact that such an amount can vary in a broad range and is dependent on various factors, e.g. the treated cultivated plant or material and the climatic conditions.

The term "effective amount" comprises the terms "plant health effective amount" and/or 35 "pesticidally effective amount" as the case may be.

When preparing the mixtures, it is preferred to employ the pure active compounds, to which further active compounds against pests, such as insecticides, herbicides, fungicides or else herbicidal or growth-regulating active compounds or fertilizers can be 40 added as further active components according to need.

The inventive mixtures are employed by treating the harmful fungi or the plants, plant propagation materials (preferably seeds), materials or soil to be protected from fungal attack with a pesticidally effective amount of the active compounds. The application can be carried out both before and after the infection of the materials, plants or plant propagation materials (preferably seeds) by the pests.

In the method of combating harmful fungi depending on the type of compound and the desired effect, the application rates of the mixtures according to the invention are from 0,3 g/ha to 2000 g/ha, preferably 5 g/ha to 2000 g/ha, more preferably from 50 to 900 g/ha, in particular from 50 to 750 g/ha.

In the method of combating animal pests (insects, acarids or nematodes) depending on the type of compound and the desired effect, the application rates of the mixtures according to the invention are from 0,3 g/ha to 2000 g/ha, preferably 5 g/ha to 2000 g/ha, more preferably from 50 to 900 g/ha, in particular from 50 to 750 g/ha.

The inventive mixtures or compositions of these mixtures can also be employed for protecting plants from attack or infestation by animal pests (insects, acarids or nematodes) comprising contacting a plant, or soil or water in which the plant is growing.

In the context of the present invention, the term plant refers to an entire plant, a part of the plant or the propagation material of the plant.

Plants and as well as the propagation material of said plants, which can be treated with the inventive mixtures include all genetically modified plants or transgenic plants, e.g. crops which tolerate the action of herbicides or fungicides or insecticides owing to breeding, including genetic engineering methods, or plants which have modified characteristics in comparison with existing plants, which can be generated for example by traditional breeding methods and/or the generation of mutants, or by recombinant procedures.

For example, mixtures according to the present invention can be applied (as seed treatment, spray treatment, in furrow or by any other means) also to plants which have been modified by breeding, mutagenesis or genetic engineering including but not limiting to agricultural biotech products on the market or in development (cf. http://www.bio.org/speeches/pubs/er/agri_products.asp). Genetically modified plants are plants, which genetic material has been so modified by the use of recombinant DNA techniques that under natural circumstances cannot readily be obtained by cross breeding, mutations or natural recombination. Typically, one or more genes have been integrated into the genetic material of a genetically modified plant in order to improve certain properties of the plant. Such genetic modifications also include but are not lim-

ited to targeted post-translational modification of protein(s), oligo- or polypeptides e. g. by glycosylation or polymer additions such as prenylated, acetylated or farnesylated moieties or PEG moieties.

5 Plants that have been modified by breeding, mutagenesis or genetic engineering, e. g. have been rendered tolerant to applications of specific classes of herbicides, such as hydroxyphenylpyruvate dioxygenase (HPPD) inhibitors; acetolactate synthase (ALS) inhibitors, such as sulfonyl ureas (see e. g. US 6,222,100, WO 01/82685, WO 00/26390, WO 97/41218, WO 98/02526, WO 98/02527, WO 04/106529, WO 05/20673, WO 03/14357, WO 03/13225, WO 03/14356, WO 04/16073) or imidazolinones (see e. g. US 6,222,100, WO 01/82685, WO 00/26390, WO 97/41218, WO 98/02526, WO 98/02527, WO 04/106529, WO 05/20673, WO 03/14357, WO 03/13225, WO 03/14356, WO 04/16073); enolpyruvylshikimate-3-phosphate synthase (EPSPS) inhibitors, such as glyphosate (see e. g. WO 92/00377); glutamine synthetase (GS) inhibitors, such as glufosinate (see e.g. EP-A 242 236, EP-A 242 246) or oxynil herbicides (see e. g. US 5,559,024) as a result of conventional methods of breeding or genetic engineering. Several cultivated plants have been rendered tolerant to herbicides by conventional methods of breeding (mutagenesis), e. g. Clearfield® summer rape (Canola, BASF SE, Germany) being tolerant to imidazolinones, e. g. imazamox. Genetic engineering methods have been used to render cultivated plants such as soybean, cotton, corn, beets and rape, tolerant to herbicides such as glyphosate and glufosinate, some of which are commercially available under the trade names RoundupReady® (glyphosate-tolerant, Monsanto, U.S.A.) and LibertyLink® (glufosinate-tolerant, Bayer CropScience, Germany).

25 Furthermore, plants are also covered that are by the use of recombinant DNA techniques capable to synthesize one or more insecticidal proteins, especially those known from the bacterial genus *Bacillus*, particularly from *Bacillus thuringiensis*, such as δ -endotoxins, e. g. CryIA(b), CryIA(c), CryIF, CryIF(a2), CryIIA(b), CryIIIA, CryIIIB(b1) or Cry9c; vegetative insecticidal proteins (VIP), e. g. VIP1, VIP2, VIP3 or VIP3A; insecticidal proteins of bacteria colonizing nematodes, e. g. *Photorhabdus* spp. or *Xenorhabdus* spp.; toxins produced by animals, such as scorpion toxins, arachnid toxins, wasp toxins, or other insect-specific neurotoxins; toxins produced by fungi, such as *Streptomyces* toxins, plant lectins, such as pea or barley lectins; agglutinins; proteinase inhibitors, such as trypsin inhibitors, serine protease inhibitors, patatin, cystatin or papain inhibitors; ribosome-inactivating proteins (RIP), such as ricin, maize-RIP, abrin, luffin, saporin or bryodin; steroid metabolism enzymes, such as 3-hydroxysteroid oxidase, ecdysteroid-IDP-glycosyl-transferase, cholesterol oxidases, ecdysone inhibitors or HMG-CoA-reductase; ion channel blockers, such as blockers of sodium or calcium channels; juvenile hormone esterase; diuretic hormone receptors (helicokinin receptors); stilben synthase, bibenzyl synthase, chitinases or glucanases. In the context of the present invention these insecticidal proteins or toxins are to be understood ex-

pressly also as pre-toxins, hybrid proteins, truncated or otherwise modified proteins. Hybrid proteins are characterized by a new combination of protein domains, (see, e. g. WO 02/015701). Further examples of such toxins or genetically modified plants capable of synthesizing such toxins are disclosed, e. g., in EP-A 374 753, WO 93/007278, 5 WO 95/34656, EP-A 427 529, EP-A 451 878, WO 03/18810 und WO 03/52073. The methods for producing such genetically modified plants are generally known to the person skilled in the art and are described, e. g. in the publications mentioned above. These insecticidal proteins contained in the genetically modified plants impart to the plants producing these proteins tolerance to harmful pests from all taxonomic groups of 10 athropods, especially to beetles (Coeloptera), two-winged insects (Diptera), and moths (Lepidoptera) and to nematodes (Nematoda). Genetically modified plants capable to synthesize one or more insecticidal proteins are, e. g., described in the publications mentioned above, and some of which are commercially available such as YieldGard® (corn cultivars producing the Cry1Ab toxin), YieldGard® Plus (corn cultivars producing 15 Cry1Ab and Cry3Bb1 toxins), Starlink® (corn cultivars producing the Cry9c toxin), Herculex® RW (corn cultivars producing Cry34Ab1, Cry35Ab1 and the enzyme Phosphinothricin-N-Acetyltransferase [PAT]); NuCOTN® 33B (cotton cultivars producing the Cry1Ac toxin), Bollgard® I (cotton cultivars producing the Cry1Ac toxin), Bollgard® II (cotton cultivars producing Cry1Ac and Cry2Ab2 toxins); VIPCOT® (cotton cultivars 20 producing a VIP-toxin); NewLeaf® (potato cultivars producing the Cry3A toxin); Bt-Xtra®, NatureGard®, KnockOut®, BiteGard®, Protecta®, Bt11 (e. g. Agrisure® CB) and Bt176 from Syngenta Seeds SAS, France, (corn cultivars producing the Cry1Ab toxin and PAT enzyme), MIR604 from Syngenta Seeds SAS, France (corn cultivars producing a modified version of the Cry3A toxin, c.f. WO 03/018810), MON 863 from 25 Monsanto Europe S.A., Belgium (corn cultivars producing the Cry3Bb1 toxin), IPC 531 from Monsanto Europe S.A., Belgium (cotton cultivars producing a modified version of the Cry1Ac toxin) and 1507 from Pioneer Overseas Corporation, Belgium (corn cultivars producing the Cry1F toxin and PAT enzyme).

Furthermore, plants are also covered that are by the use of recombinant DNA techniques capable to synthesize one or more proteins to increase the resistance or tolerance of those plants to bacterial, viral or fungal pathogens. Examples of such proteins are the so-called "pathogenesis-related proteins" (PR proteins, see, e. g. EP-A 392 225), plant disease resistance genes (e. g. potato cultivars, which express resistance genes acting against *Phytophthora infestans* derived from the mexican wild 35 potato *Solanum bulbocastanum*) or T4-lysozym (e. g. potato cultivars capable of synthesizing these proteins with increased resistance against bacteria such as *Erwinia amylovora*). The methods for producing such genetically modified plants are generally known to the person skilled in the art and are described, e. g. in the publications mentioned above.

40 Furthermore, plants are also covered that are by the use of recombinant DNA techniques capable to synthesize one or more proteins to increase the productivity (e. g. bio mass production, grain yield, starch content, oil content or protein content), toler-

ance to drought, salinity or other growth-limiting environmental factors or tolerance to pests and fungal, bacterial or viral pathogens of those plants.

Furthermore, plants are also covered that contain by the use of recombinant DNA techniques a modified amount of substances of content or new substances of content,
5 specifically to improve human or animal nutrition, e. g. oil crops that produce health-promoting long-chain omega-3 fatty acids or unsaturated omega-9 fatty acids (e. g. Nexera® rape, DOW Agro Sciences, Canada).

Furthermore, plants are also covered that contain by the use of recombinant DNA techniques a modified amount of substances of content or new substances of content,
10 specifically to improve raw material production, e. g. potatoes that produce increased amounts of amylopectin (e. g. Amflora® potato, BASF SE, Germany).

Water-soluble concentrates (LS), flowable concentrates (FS), powders for dry treatment (DS), water-dispersible powders for slurry treatment (WS), water-soluble powders
15 (SS), emulsions (ES) emulsifiable concentrates (EC) and gels (GF) are usually employed for the purposes of treatment of plant propagation materials, particularly seeds.

In a preferred embodiment of the invention, the inventive mixtures are used for the protection of the seed and the seedlings' roots and shoots, preferably the seeds.
20

Seed treatment can be made into the seedbox before planting into the field.

For seed treatment purposes, the weight ration in the binary, ternary and quaternary mixtures of the present invention generally depends from the properties of the compounds of the inventive mixtures.
25

Compositions, which are especially useful for seed treatment are e.g.:

- A Soluble concentrates (SL, LS)
- 30 D Emulsions (EW, EO, ES)
- E Suspensions (SC, OD, FS)
- F Water-dispersible granules and water-soluble granules (WG, SG)
- G Water-dispersible powders and water-soluble powders (WP, SP, WS)
- H Gel-Formulations (GF)
- 35 I Dustable powders (DP, DS)

These compositions can be applied to plant propagation materials, particularly seeds, diluted or undiluted. These compositions can be applied to plant propagation materials, particularly seeds, diluted or undiluted. The compositions in question give, after two-to-
40 tenfold dilution, active substance concentrations of from 0.01 to 60% by weight, preferably from 0.1 to 40% by weight, in the ready-to-use preparations. Application can be carried out before or during sowing. Methods for applying or treating agrochemical

compounds and compositions thereof, respectively, on to plant propagation material, especially seeds, are known in the art, and include dressing, coating, pelleting, dusting and soaking application methods of the propagation material (and also in furrow treatment). In a preferred embodiment, the compounds or the compositions thereof, respectively, are applied on to the plant propagation material by a method such that germination is not induced, e. g. by seed dressing, pelleting, coating and dusting.

In the treatment of plant propagation material (preferably seed), the application rates of the inventive mixture are generally for the formulated product (which usually comprises from 10 to 750 g/l of the active(s)) .

The invention also relates to the propagation products of plants, and especially the seed comprising, that is, coated with and/or containing, a mixture as defined above or a composition containing the mixture of two or more active ingredients or a mixture of two or more compositions each providing one of the active ingredients. The plant propagation material (preferably seed) comprises the inventive mixtures in an amount of from 0.1 g to 10 kg per 100 kg of plant propagation material (preferably seed), preferably 0.1 g to 1 kg per 100 kg of plant propagation material (preferably seed).

For example, the ratio by weight of compound IIa is herein preferably between 0,1 – 200 g/100kg plant propagation material (preferably seed), more preferred 1 to 200 g/100kg plant propagation material (preferably seed) and most preferred 1 to 100 g/100kg plant propagation material (preferably seed).

For example, the ratio by weight for compound IIIB is herein preferably between 1 - 200 g/100kg plant propagation material (preferably seed), more preferred 5 to 200 g/100kg plant propagation material (preferably seed), and most preferred 5 to 100g/100kg plant propagation material (preferably seed).

For example, the ratio by weight for the compound IIIA as compound III is herein preferably between 1 - 200 g/100kg plant propagation material (preferably seed), more preferred 1 to 50 g/100kg plant propagation material (preferably seed) and most preferred 1 to 20 g/100kg plant propagation material (preferably seed).

For example, the ratio by weight for compound IIc is herein preferably between 0,1 - 200 g/100kg plant propagation material (preferably seed), more preferred 1 to 200 g/100kg plant propagation material (preferably seed) and most preferred 1 to 50 g/100kg plant propagation material (preferably seed).

For example, the ratio by weight for compound IIb is herein preferably between 1 - 200 g/100kg plant propagation material (preferably seed), more preferred 25 to 200 g/100kg

plant propagation material (preferably seed) and most preferred 50 to 100 g/100kg plant propagation material (preferably seed).

5 The separate or joint application of the compounds of the inventive mixtures is carried out by spraying or dusting the seeds, the seedlings, the plants or the soils before or after sowing of the plants or before or after emergence of the plants.

10 The inventive mixtures are effective through both contact (via soil, glass, wall, bed net, carpet, plant parts or animal parts), and ingestion (bait, or plant part) and through trophallaxis and transfer.

15 Preferred application methods are into water bodies, via soil, cracks and crevices, pastures, manure piles, sewers, into water, on floor, wall, or by perimeter spray application and bait.

20 According to another preferred embodiment of the invention, for use against non crop pests such as ants, termites, wasps, flies, mosquitoes, crickets, locusts, or cockroaches the inventive mixtures are prepared into a bait preparation.

25 The bait can be a liquid, a solid or a semisolid preparation (e.g. a gel). The bait employed in the composition is a product which is sufficiently attractive to incite insects such as ants, termites, wasps, flies, mosquitoes, crickets etc. or cockroaches to eat it. This attractant may be chosen from feeding stimulants or para and / or sex pheromones readily known in the art.

30 Methods to control infectious, non-phytopathogenic diseases transmitted by insects (e.g. malaria, dengue and yellow fever, lymphatic filariasis, and leishmaniasis) with the inventive mixtures and their respective compositions also comprise treating surfaces of huts and houses, air spraying and impregnation of curtains, tents, clothing items, bed nets, tsetse-fly trap or the like. Insecticidal compositions for application to fibers, fabric, knitgoods, non-wovens, netting material or foils and tarpaulins preferably comprise a composition including the inventive mixtures, optionally a repellent and at least one binder.

35 The inventive mixtures and the compositions comprising them can be used for protecting wooden materials such as trees, board fences, sleepers, etc. and buildings such as houses, outhouses, factories, but also construction materials, furniture, leathers, fibers, vinyl articles, electric wires and cables etc. from ants and/or termites, and for controlling ants and termites from doing harm to crops or human being (e.g. when the pests
40 invade into houses and public facilities).

In the case of soil treatment or of application to the pests dwelling place or nest, the quantity of active ingredient ranges from 0.0001 to 500 g per 100 m², preferably from 0.001 to 20 g per 100 m².

- 5 Customary application rates in the protection of materials are, for example, from 0.01 g to 1000 g of active compound per m² treated material, desirably from 0.1 g to 50 g per m².

- 10 Insecticidal compositions for use in the impregnation of materials typically contain from 0.001 to 95 weight %, preferably from 0.1 to 45 weight %, and more preferably from 1 to 25 weight % of at least one repellent and / or insecticide.

- 15 For use in bait compositions, the typical content of active ingredient is from 0.0001 weight % to 15 weight %, desirably from 0.001 weight % to 5% weight % of active compound. The composition used may also comprise other additives such as a solvent of the active material, a flavoring agent, a preserving agent, a dye or a bitter agent. Its attractiveness may also be enhanced by a special color, shape or texture.

- 20 For use in spray compositions, the content of the mixture of the active ingredients is from 0.001 to 80 weights %, preferably from 0.01 to 50 weight % and most preferably from 0.01 to 15 weight %.

Claims

1. Mixtures comprising, as active components,
 - 5 1) chlormequat chloride as compound I; and
 - 2) one or more insecticidal or nematocidal compound II selected from the group consisting of
 - 10 IIa) nicotinic receptor agonists/antagonists compounds: clothianidin, dinotefuran, imidacloprid, thiamethoxam, nitenpyram and acetamiprid; and
 - IIb) macrolide compounds: abamectin, emamectin benzoate and spinosad; and
 - IIc) fipronil or ethiprolein synergistic effective amounts.
15
2. The mixture according to claim 1, wherein one or more insecticidal compound II is selected from
 - 20 IIa) clothianidin, imidacloprid, thiamethoxam and acetamiprid; and
 - IIb) abamectin; and
 - IIc) fipronil.
3. The mixture according to claim 1, wherein the insecticidal compound II is selected from clothianidin, imidacloprid, thiamethoxam, acetamiprid, abamectin; and
25 fipronil.
4. The mixture according to any of claims 1 to 2, additionally comprising one or more fungicidal compound III selected from
 - 30 3 a) a fungicidal compound IIIA selected from the group of strobilurine fungicides IIIA comprising pyraclostrobin, azoxystrobin, dimoxystrobin, enestroburin, fluoxastrobin, kresoxim-methyl, metominostrobin, orysastrobin, picoxystrobin, pyribencarb, trifloxystrobin, 2-(2-(6-(3-chloro-2-methyl-phenoxy)-5-fluoro-pyrimidin-4-yloxy)-phenyl)-2-methoxyimino-N-methyl-acetamide, 3-methoxy-2-(2-(N-(4-methoxy-phenyl)-cyclopropane-
35 carboximidoylsulfanylmethyl)-phenyl)-acrylic acid methyl ester, methyl (2-chloro-5-[1-(3-methylbenzyloxyimino)ethyl]benzyl)carbamate and 2-(2-(3-(2,6-dichlorophenyl)-1-methyl-allylideneaminooxymethyl)-phenyl)-2-methoxyimino-N-methyl-acetamide; and
 - 40 3 b) an fungicidal compound IIIB from the group of carboxamides consisting of N-(3',4',5'-trifluorobiphenyl-2-yl)- 3-difluoromethyl-1-methyl-1H-pyrazole-4-carboxamide, N-[2-(4'-trifluoromethylthio)-biphenyl]-3-difluoromethyl-1-methyl-1H-pyrazole-4-carboxamide, bixafen, N-[2-(1,3-dimethylbutyl)-

phenyl]-1,3-dimethyl-5-fluoro-1H-pyrazole-4-carboxamide (penflufen), sedaxane, isopyrazam and penthiopyrad).

- 5
5. The mixture according to claim 4, wherein one or more fungicidal compound III is selected from
- IIIb) compound IIIB: N-(3',4',5'-trifluorobiphenyl-2-yl)- 3-difluoromethyl-1-methyl-1H-pyrazole-4-carboxamide, sedaxane, N-[2-(1,3-dimethylbutyl)-phenyl]-1,3-dimethyl-5-fluoro-1H-pyrazole-4-carboxamide (penflufen) and penthiopyrad;
- 10
6. The mixture according to claim 4, wherein fungicidal compound III is selected from
- IIIA) compound IIIA: pyraclostrobin, azoxystrobin, oryastrobin, picoxystrobin, trifloxystrobin.
- 15
7. The mixture according to claim 4, wherein fungicidal compound III is pyraclostrobin.
8. The mixture according to claim 4, wherein fungicidal compound III is a binary mixture of compound IIIA and compound IIIB.
- 20
9. A mixture comprising fipronil and abamectin in synergistically effective amounts.
10. A pesticidal composition, comprising a liquid or solid carrier and a mixture as defined in any of claims 1 to 9.
- 25
11. A method for controlling pests and/or improving the health of plants, wherein
- (a) the pest, their habitat, breeding grounds, their locus or the plants to be protected against pest attack, the soil or plant propagation material; or
- (b) the plant, the locus where the plant is growing or is expected to grow or plant propagation material from which the plant grows;
- 30
- are treated with an effective amount of a mixture as defined in any of claims 1 to 9.
12. A method for protection of plant propagation material from pests comprising contacting the plant propagation materials with a mixture as defined in any of claims 1 to 9 in pesticidally effective amounts.
- 35
13. A method as claimed in claim 12, wherein the mixture as defined in any of claims 1 to 9 is applied in an amount of from 0.01 g to 10 kg per 100 kg of plant propagation materials.
- 40

14. A method as claimed in claims 11 to 13, wherein the compounds as defined in any of claims 1 to 9 are applied simultaneously, that is jointly or separately, or in succession.
- 5 15. Plant propagation material, comprising the mixture as defined in any of claims 1 to 9 in an amount of from 0.01 g to 10 kg per 100 kg of plant propagation materials.

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2010/051530

A. CLASSIFICATION OF SUBJECT MATTER
 INV. A01P3/00 A01P5/00 A01P7/02 A01P7/04 A01P21/00
 A01N33/12 A01N47/02

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED
 Minimum documentation searched (classification system followed by classification symbols)
A01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)
EPO-Internal, WPI Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2007/068355 A (BAYER CROPS SCIENCE AG [DE]; JESCHKE PETER [DE]; NAUEN RALF [DE]; PONTZE) 21 June 2007 (2007-06-21) claims 1-4,6,7,17,18,23 page 61, paragraph 1	1-8, 10-15
A	DATABASE CAPLUS [Online] CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; 1989, NEIGEBBAUR, E. F. ET AL: "Use of liquid nitrogen fertilizers, pesticides, and retardants" XP002534249 retrieved from STN Database accession no. 1989:510996 abstract -/--	1-8, 10-15

Further documents are listed in the continuation of Box C. See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"E" earlier document but published on or after the international filing date	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
"O" document referring to an oral disclosure, use, exhibition or other means	"&" document member of the same patent family
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search 17 May 2010	Date of mailing of the international search report 30/06/2010
---	---

Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Authorized officer Marie, Gérald
--	--

INTERNATIONAL SEARCH REPORT

International application No

PCT/EP2010/051530

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	& KHIMIZATSIYA SEL'SKOGO KHOZYAISTVA (1988-1992) , (6), 69-71 CODEN: KSKHE7; ISSN: 0235-2516, 1989, ----- DATABASE CAPLUS [Online] CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; 1983, DUNAEVSKII, A. G. ET AL: "Use of a tricomponent mixture" XP002534250 retrieved from STN Database accession no. 1983:448944 abstract	1-8, 10-15
A	& ZASHCHITA RASTENII (MOSCOW) , (5), 29-30 CODEN: ZSRBXX; ISSN: 0044-1864, 1983, ----- DATABASE CAPLUS [Online] CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; 1997, VOLKMAR, CHRISTA ET AL: "Spider populations on a typical field site in central Germany and special influences of various plant protection intensities" XP002534251 retrieved from STN Database accession no. 1997:475785 abstract	1-8, 10-15
A	& ARCHIVES OF PHYTOPATHOLOGY AND PLANT PROTECTION , 30(6), 533-546 CODEN: APPPER; ISSN: 0323-5408, 1997, ----- DATABASE CAPLUS [Online] CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; 1998, SOKOLOWSKI, ANDRZEJ: "Pesticides and epigeal arthropods of agrocenoses" XP002534252 retrieved from STN Database accession no. 1998:722736 abstract	1-8, 10-15
A	& PESTYCYDY (WARSAW) , (3-4), 55-62 CODEN: PSTYDL; ISSN: 0208-8703, 1997, ----- DATABASE CAPLUS [Online] CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; 1971, ACKERMANN, HEINZ ET AL: "Effect of chlorocholine chloride on the toxicity of cholinesterase inhibitors" XP002534253 retrieved from STN Database accession no. 1971:404397 abstract -/--	1-8, 10-15

INTERNATIONAL SEARCH REPORT

 International application No
 PCT/EP2010/051530

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	& ARCHIV FUER EXPERIMENTELLE VETERINAERMEDIZIN , 24(4), 1045-7 CODEN: AXVMAW; ISSN: 0003-9055, 1970, ----- FR 2 600 494 A (KAO CORP [JP]) 31 December 1987 (1987-12-31) claims 1-5	1-8, 10-15
A	----- WO 95/12314 A1 (RHONE POULENC AGROCHIMIE [FR]; COLLIOT FRANCOIS [FR]; GOUOT JEAN MARIE) 11 May 1995 (1995-05-11) claims 1,2,4-7 examples 1-8 -----	1-8, 10-15

INTERNATIONAL SEARCH REPORT

International application No.
PCT/EP2010/051530

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.

2. As all searchable claims could be searched without effort justifying an additional fees, this Authority did not invite payment of additional fees.

3. As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
1-8(completely); 10-15(partially)

Remark on Protest

- The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. claims: 1-8(completely); 10-15(partially)

A mixture according to independent claim 1;
a composition according to independent claim 10 comprising said mixture;
methods according to independent claims 11 and 12 using said mixture;
a plant propagation material according to independent claim 15 comprising said mixture,

2. claims: 9(completely); 10-15(partially)

A mixture according to independent claim 9;
a composition according to independent claim 10 comprising said mixture;
methods according to independent claims 11 and 12 using said mixture;
a plant propagation material according to independent claim 15 comprising said mixture,

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2010/051530

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 2007068355	A	21-06-2007	AR 058339 A1	30-01-2008
			AU 2006326728 A1	21-06-2007
			CA 2632904 A1	21-06-2007
			CN 101330829 A	24-12-2008
			DE 102005059468 A1	14-06-2007
			EP 1962606 A1	03-09-2008
			JP 2009519258 T	14-05-2009
			KR 20080078871 A	28-08-2008
			US 2009105235 A1	23-04-2009
			ZA 200805039 A	30-12-2009
			FR 2600494	A
AU 7477387 A	07-01-1988			
JP 1773007 C	14-07-1993			
JP 4059281 B	21-09-1992			
JP 63008302 A	14-01-1988			
US 4888049 A	19-12-1989			
WO 9512314	A1	11-05-1995	AT 160672 T	15-12-1997
			AU 690160 B2	23-04-1998
			AU 8109494 A	23-05-1995
			BR 9408163 A	28-10-1997
			CA 2175818 A1	11-05-1995
			CN 1108043 A	13-09-1995
			CN 1140976 A	22-01-1997
			CO 4340720 A1	30-07-1996
			DE 69407198 D1	15-01-1998
			DE 69407198 T2	14-05-1998
			DK 726709 T3	06-04-1998
			EG 20772 A	29-02-2000
			EP 0726709 A1	21-08-1996
			ES 2110308 T3	01-02-1998
			FR 2712144 A1	19-05-1995
			GR 3025592 T3	31-03-1998
			JP 9504538 T	06-05-1997
			MA 23364 A1	01-07-1995
			PL 314183 A1	02-09-1996
			RO 115930 B1	30-08-2000