



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁴ : A61K 31/40, 37/02, 31/425 A61K 31/445, 31/44, 31/47	A2	(11) International Publication Number: WO 87/ 02585 (43) International Publication Date: 7 May 1987 (07.05.87)
(21) International Application Number: PCT/US86/02290 (22) International Filing Date: 31 October 1986 (31.10.86) (31) Priority Application Number: 793,929 (32) Priority Date: 1 November 1985 (01.11.85) (33) Priority Country: US (71) Applicant: SCHERING CORPORATION [US/US]; 2000 Galloping Hill Road, Kenilworth, NJ 07033 (US). (72) Inventor: WATKINS, Robert, W. ; Box 510B, R.R. #1, Great Meadows, NJ 07838 (US). (74) Agents: KANSTAD, Steinar, V. et al.; Schering Plough Corporation, One Giralda Farms, Madison, NJ 07940 (US).		(81) Designated States: AT (European patent), BE (European patent), CH (European patent), DE (European patent), FR (European patent), GB (European patent), IT (European patent), JP, LU (European patent), NL (European patent), SE (European patent). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: TREATING INCREASED INTRAOCULAR PRESSURE CAUSED BY STEROIDS (57) Abstract The rise in intraocular pressure associated with the use of steroidal anti-inflammatory drugs is controlled by use of angiotensin converting enzyme inhibitors. For the purposes of the invention, the preferred ACE inhibitor is 7-[N-(1(S)-carboxy-3-phenylpropyl)-(S)-alanyl]-1,4-dithia-7-azaspiro[4.4]nonane-8(S)-carboxylic acid.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	ML	Mali
AU	Australia	GA	Gabon	MR	Mauritania
BB	Barbados	GB	United Kingdom	MW	Malawi
BE	Belgium	HU	Hungary	NL	Netherlands
BG	Bulgaria	IT	Italy	NO	Norway
BJ	Benin	JP	Japan	RO	Romania
BR	Brazil	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	LI	Liechtenstein	SN	Senegal
CH	Switzerland	LK	Sri Lanka	SU	Soviet Union
CM	Cameroon	LU	Luxembourg	TD	Chad
DE	Germany, Federal Republic of	MC	Monaco	TG	Togo
DK	Denmark	MG	Madagascar	US	United States of America
FI	Finland				

-1-

TREATING INCREASED INTRAOCULAR
PRESSURE CAUSED BY STEROIDS

The present invention relates to the use of angiotensin converting enzyme (ACE) inhibitors for preparing medicaments useful in treating the rise in intraocular pressure (IOP) associated with ophthalmic and systemic use of steroidal anti-inflammatory drugs.

A second aspect of the invention is a pharmaceutical composition comprising an ACE inhibitor and an anti-inflammatory steroid, wherein the ACE inhibitor suppresses the effect of the steroid-induced rise in IOP.

Steroidal anti-inflammatory drugs are commonly prescribed for inflammatory conditions such as arthritis, and for treatment of ocular inflammation due to mechanical or chemical irritants or immunological reactions resulting from such conditions as allergic conjunctivitis, herpes zoster keratitis, iritis, and corneal injury from chemical or thermal burns or penetration of foreign bodies. Anti-inflammatory steroids, however, are known to cause an elevation in IOP resulting in the condition known as ocular hypertension. While the term "ocular hypertension" indicates elevated IOP without optic nerve head damage or loss of vision, the condition is believed by the majority of

-2-

ophthalmologists to represent the earliest phase in the onset of glaucoma, which is characterized by optic nerve damage and loss of vision, and which may lead to blindness. Patients with pre-existing glaucoma are particularly vulnerable to additional elevations in IOP caused by steroidal anti-inflammatory therapy. Also, in some susceptible individuals, topical steroids can induce an intractable glaucoma.

The increase in IOP often associated with the topical optical administration of steroidal anti-inflammatory agents can also occur with other modes of administration of the drugs, including systemic (usually oral), local injection (e.g., depot injection), topical, aerosol or intravitreal administration.

Anti-inflammatory steroids include hydrocortisone, cortisone, prednisolone, prednisone, dexamethasone, methylprednisolone, triamcinolone, betamethasone, alclometasone, flunisolide, beclomethasone, clorocortolone, diflorasone, halcinonide, fluocinonide, flucinolone, desoximetasone, medrysone, paramethasone, 9,21-dichloro-17-[(2-furanylcarbonyl)oxy]-11-hydroxy-16 α -methyl-pregna-1,4-diene-3,20-dione, and fluorometholone, and their pharmaceutically acceptable salts and esters.

The steroids contemplated for use in this invention are exemplified by, but are not limited to, those listed above. Preferred steroids are hydrocortisone, prednisolone, dexamethasone, betamethasone, beclomethasone medrysone and fluorometholone, and their pharmaceutically acceptable salts and esters.

Drugs currently employed to treat glaucoma, e.g., epinephrine and pilocarpine, are not particularly effective in lowering steroid-induced increases in IOP, and often produce serious side effects.

-3-

The elevation in IOP associated with the clinical ophthalmic and systemic use of anti-inflammatory steroids, in particular corticosteroids, can be reduced by the administration of an angiotensin converting enzyme (ACE) inhibitor. The ACE inhibitor may be administered separately from the steroid treatment in order to lower elevated IOP, or may be co-administered with the steroid to suppress the IOP-raising effect of the steroid while not interfering with the anti-inflammatory activity of the steroid.

The present invention therefore relates to a method for reducing and controlling the elevated IOP associated with the ophthalmic and systemic use of steroidal anti-inflammatory agents, which method comprises administering to a mammal in need of such treatment an effective amount of an ACE inhibitor, preferably in an ophthalmologically acceptable topical carrier.


In addition, the present invention relates to a method for suppressing the elevation in IOP associated with the ophthalmic and systemic use of steroidal anti-inflammatory agents, which method comprises administering to a mammal in need of such treatment a combination of an anti-inflammatory effective amount of a steroid and an effective amount of an ACE inhibitor, either separately or in the same pharmaceutical composition.

Any possible combination of dosage forms may be used to administer the combination, e.g., oral steroid/topical ACE inhibitor, topical steroid/oral ACE inhibitor, oral steroid/oral ACE inhibitor, topical steroid/topical ACE inhibitor, and locally injected steroid/topical ACE inhibitor. A preferred combination comprises a steroid and a topical ACE inhibitor. For ophthalmic use, a combination of a topical steroid and a topical ACE inhibitor is preferred.


-4-

Examples of angiotensin converting enzyme (ACE) inhibitors are those disclosed in the following patents and published patent applications:

(a) U.S. Patent No. 4,105,776 to Ondetti et al. discloses proline derivatives which are angiotensin converting enzyme (ACE) inhibitors and have the general formula



wherein R^a is hydroxy, NH_2 or lower alkoxy;
 R_1^a and R_4^a each is hydrogen, lower alkyl, phenyl or phenyl-lower alkyl;
 R_2^a is hydrogen, lower alkyl, phenyl, substituted phenyl wherein the phenyl substituent is halo, lower alkyl or lower alkoxy, phenyl-lower alkyl, diphenyl-lower alkyl, triphenyl-lower alkyl, lower alkylthiomethyl, phenyl-lower alkylthiomethyl, lower alkanoyl-amidomethyl,




$R_6^a - S -$, or R_7^a ;

R_3^a is hydrogen, hydroxy or lower alkyl;

R_5^a is lower alkyl, phenyl or phenyl-lower alkyl;

-5-

R_6^a is lower alkyl, phenyl, substituted phenyl (wherein the phenyl substituent is halo, lower alkyl or lower alkoxy), hydroxy-lower alkyl or amino (carboxy)-lower alkyl;




M^a is O or S;

m^a is 1 to 3;

n^a and p^a each is 0 to 2.

(b) European Patent Application 97050 published December 28, 1983 discloses phosphinylalkanoyl substituted prolines which have the formula




R_1^b is alkyl, aryl, aralkyl, cycloalkyl or cycloalkylalkyl;

R_2^b is cycloalkyl, 3-cyclohexenyl or 2-alkyl-3-cyclohexenyl;


-6-

R_3^b is alkyl, cycloalkyl or phenyl;
 R_4^b is H or alkyl;
 one of R_5^b and R_6^b is H and the other is alkyl- X^b -,
 phenyl- X^b -, alkoxy, phenoxy, phenyl, cycloalkyl,
 alkyl or phenylalkyl; or
 R_5^b and R_6^b together form $-X^bCH_2CH_2X^b-$, X^b is S,
 SO or SO_2 ;
 R_7^b is H or $CH(R_2^b)OCOR_3^b$;
 n^b is 0 or 1;
 'aryl' is phenyl opt. substituted by halo,
 alkyl, alkoxy, alkylthio, OH, alkanoyl, NO_2 ,
 amino, dialkylamino and/or CF_3 ;
 alkyl has 1-10C, cycloalkyl 3-7C, alkoxy 1-8C
 and alkanoyl 2-9C.

(c) European Patent Application 83172
 published July 6, 1983 discloses N-substituted acetyl-L-
 proline derivatives which are said to be angiotensin
 converting enzyme inhibitors and have the formula







R_1^C and R_2^C are H, 1-7C alkyl, 1-7C haloalkyl,
 $(CH_2)_m^C-D^C$, 1-7C aminoalkyl or a group of
 formula:



-7-

D^C is cycloalkyl, furyl, thienyl or pyridinyl;
 A^C is $(CH_2)_n-CHR_{21}^C$, $NR_{22}^C-CHR_{23}^C$ or $O-CHR_{23}^C$;
 n^C is 0 or 1;
 q^C is 0-7;
 R_{21}^C is H, 1-7C alkyl, 1-7C haloalkyl, benzyl or phenethyl;
 R_{22}^C is H or 1-7C alkyl;
 R_{23}^C is H, 1-7C alkyl, 1-7C haloalkyl or $(CH_2)_rCD_1^C$;
 D_1^C is phenyl, 4-hydroxyphenyl, 3,4-dihydroxyphenyl, 3-indolyl, 4-imidazolyl, NH_2 , SH, 1-7C alkylthio, guanidino or $CONH_2$;
 X^C is a group of formula $(II)^C$ -(V)^C, or $NR_4^C-CHR_5^C-COOR_6^C$;



where

R_9^C and R_8^C are H and R_7^C is as defined below;
 R_9^C and R_7^C are H and R_8^C is as defined below;
 R_7^C and R_8^C are H and R_9^C is as defined below;
 R_9^C and R_8^C are H and $-CHR_7^C-$ is replaced by $-CR_{10}^CR_{10}^C$;
 R_9^C is H and $R_7^C + R_8^C$ form a double bond;




-8-

R_9^C is H and $R_7^C + R_8^C$ complete a fused benzene ring; or

R_8^C is H and $R_9^C + R_7^C$ complete a fused benzene ring;

E'^C and E''^C are H or $E'^C + E''^C$ complete a fused benzene ring;

R_7^C is H, 1-7C alkyl, halogen, keto, OH, 2-8C alkanoyl-amino, N_3 , NH_2 , $NR_{19}^C R_{20}^C$, $(CH_2)_m^C - D^C$, $O - CONR_{15}^C R_{15}^C$, 1-7C alkoxy, 1-7C alkylthio, or a group of formula $(VI)^C$, $(VII)^C$ or $(VIII)^C$:




B'^C is a bond or it is O or S; R_8^C is keto, halogen, $O - CONR_{15}^C R_{15}^C$, 1-7C alkoxy, 1-7C alkylthio, or a group $(VII)^C$ or $(VIII)^C$ in which B'^C is O or S;

R_9^C is keto or a group $(VII)^C$ in which B'^C is a bond;

R_{10}^C is halogen or $Y^C R_{16}^C$;

R_{11}^C , R'_{11}^C , R_{12}^C and R'_{12}^C are H or 1-7C alkyl, or R_{11}^C is a group of formula $(IX)^C$ and the other 3 are H;




R_{13}^C is H, 1-4C alkyl, 1-4C alkoxy, 1-4C alkylthio, Cl, Br, F, CF_3 , OH, Ph, PhO, PhS or $PhCH_2$;

-9-

R_{14}^C is H, 1-4C alkyl, 1-4C alkoxy, 1-4C alkylthio, Cl, Br, F, CF_3 or OH;
 m^C is 0-3;
 p^C is 1-3 but it is 2 or 3 only when R_{13}^C or R_{14}^C is H, Me, MeO, Cl or F;
 R_{15}^C is H or 1-4C alkyl;
 Y^C is O or S;
 R_{16}^C is 1-4C alkyl or group (VII)^C in which B^C is a bond; or the two R_{16}^C groups complete a 5- or 6-membered ring in which the C atoms may each bear 1 or 2 alkyls having 1-4C;
 R_4^C is H, 1-7C alkyl, cycloalkyl or $(CH_2)_r^C Ph$;
 R_5^C is H, 1-7C alkyl or $(CH_2)_r^C - D_1^C$;
 r^C is 1-4;
 R_3^C and R_6^C are H, 1-7C alkyl, $PhCH_2$, $PhCH$ or $CHR_{17}^C - O - COR_{18}^C$;
 R_{17}^C is H, 1-7C alkyl or Ph;
 R_{18}^C is H, 1-7C alkyl, 1-7C alkoxy or Ph; or $R_{17}^C + R_{18}^C$ form $(CH_2)_2$, $(CH_2)_3$, $-CH=CH$ or o-phenylene;
 R_{19}^C is 1-7C alkyl, benzyl or phenethyl; and
 R_{20}^C is H or chosen as for R_{19}^C .

(d) European Patent Application 0 012 401 published June 25, 1980 discloses carboxyalkyl dipeptide derivatives which are said to be angiotensin converting enzyme inhibitors and have the formula



-10-

wherein


R^d and R^{6d} are the same or different and are hydroxy, lower alkoxy, lower alkenoxy, dilower alkylamino lower alkoxy (dimethylaminoethoxy), acylamino lower alkoxy (acetylaminomethoxy), acyloxy lower alkoxy (pivaloyloxymethoxy), aryloxy, such as phenoxy, arloweralkoxy, such as benzyloxy, substituted aryloxy or substituted arloweralkoxy wherein the substituent is methyl, halo or methoxy, amino, loweralkylamino, diloweralkylamino, hydroxyamino, arloweralkylamino such as benzylamino; R^{1d} is hydrogen, alkyl of from 1 to 20 carbon atoms which include branched and cyclic and unsaturated (such as allyl) alkyl groups, substituted loweralkyl wherein the substituent can be halo, hydroxy, lower alkoxy, aryloxy such as phenoxy, amino, diloweralkylamino, acylamino, such as acetamido and benzamido, arylamino, guanidino, imidazolyl, indolyl, mercapto, loweralkylthio, arylthio such as phenylthio, carboxy or carboxamido, carboloweralkoxy, aryl such as phenyl or naphthyl, substituted aryl such as phenyl wherein the substituent is lower alkyl, lower alkoxy or halo, arloweralkyl, arloweralkenyl, heteroarlower alkyl or heteroarlower alkenyl such as benzyl, styryl or indolyl ethyl, substituted arloweralkyl, substituted arloweralkenyl, substituted heteroarlower alkyl, or substituted heteroarlower alkenyl, wherein the substituent(s) is halo, dihalo, lower alkyl, hydroxy, lower alkoxy, amino, aminomethyl, acylamino (acetylaminomethyl or benzoylamino) diloweralkylamino, loweralkylamino, carboxyl,

-11-




haloloweralkyl, cyano or sulfonamido;
arloweralkyl or heteroarloweralkyl substituted
on the alkyl portion by amino or acylamino
(acetylamino or benzoylamino);
 R^{2d} and R^{7d} are the same or different and are
hydrogen or lower alkyl;
 R^{3d} is hydrogen, lower alkyl, phenyl lower
alkyl, aminomethyl phenyl lower alkyl, hydroxy
phenyl lower alkyl, hydroxy lower alkyl,
acylamino lower alkyl (such as benzoylamino
lower alkyl, acetylamino lower alkyl), amino
lower alkyl, dimethylamino lower alkyl, halo
lower alkyl, guanidino lower alkyl, imidazolyl
lower alkyl, indolyl lower alkyl, mercapto lower
alkyl, lower alkyl thio lower alkyl;
 R^{4d} is hydrogen or lower alkyl;
 R^{5d} is hydrogen, lower alkyl, phenyl, phenyl
lower alkyl, hydroxy phenyl lower alky, hydroxy
lower alkyl, amino lower alkyl, guanidino lower
alkyl, imidazolyl lower alkyl, indolyl lower
alkyl, mercapto lower alkyl or lower alkylthio
lower alkyl;
 R^{4d} and R^{5d} may be connected together to form an
alkylene bridge of from 2 to 4 carbon atoms, an
alkylene bridge of from 2 to 3 carbon atoms and
one sulfur atom, an alkylene bridge of from 3 to
4 carbon atoms containing a double bond or an
alkylene bridge as above substituted with
hydroxy, loweralkoxy, lower alkyl or dilower
alkyl;
and the pharmaceutically acceptable salts
thereof.

-12-



(e) U.S. Patent No. 4,462,943 to Petrillo et al. discloses carboxyalkyl amino acid derivatives of substituted prolines which are angiotensin converting enzyme inhibitors and have the formula




R_4^e is a substituted proline of the formula





R_6^e is halogen, keto, azido,




-13-




a 1- or 2-naphthyl of the formula



a 1- or 2-naphthyloxy of the formula




-S-lower alkyl,




-14-


or a 1- or 2-naphthylthio of the formula




R_7^e is keto, halogen,



or 1- or 2-naphthloxy of the formula




-S-lower alkyl,




or a 1- or 2-naphthylthio of the formula

-15-



R_8^e is keto or



R_9^e is halogen or $-Y^e-R_{13}^e$.

m^e is zero, one, two, or three.

R_{10}^e is hydrogen, lower alkyl of 1 to 4 carbons, lower alkoxy of 1 to 4 carbons, lower alkylthio of 1 to 4 carbons, chloro, bromo, fluoro, trifluoromethyl, hydroxy, phenyl, phenoxy, phenylthio, or phenylmethyl,


R_{11}^e is hydrogen, lower alkyl of 1 to 4 carbons, lower alkoxy of 1 to 4 carbons, lower alkylthio of 1 to 4 carbons, bromo, fluoro, trifluoromethyl, or hydroxy.

n^e is one, two or three provided that n^e is more than one only if R_{10}^e or R_{11}^e is hydrogen, methyl, methoxy, chloro, or fluoro.

R_{12}^e is hydrogen or lower alkyl of 1 to 4 carbons,


Y^e is oxygen or sulfur,

R_{13}^e is lower alkyl of 1 to 4 carbons.




-16-



or the R_{13}^e group join to complete an unsubstituted 5- or 6-membered ring or said ring in which one or more of the carbons has a lower alkyl of 1 to 4 carbons, or a di(lower alkyl of 1 to 4 carbons) substituent, R^e and R_5^e are independently selected from hydroxy, lower alkoxy, di(lower alkyl)-amino-lower alkoxy, such as dimethylaminoethoxy, lower alkyl-carbonyl-amino-lower alkoxy, such as acetylaminoethoxy, lower alkyl-carbonyloxy-lower alkoxy, such as pivaloyloxymethoxy,











wherein m^e , n^e and R_{10}^e are as defined above, amino, lower alkyl-amino, di(lower alkyl)-amino, hydroxyamino, benzylamino, or phenethylamino, R_1^e is hydrogen, lower alkyl,



halo substituted lower alkyl, hydroxy substituted lower alky,
 $-(CH_2)_q^e$ -cycloalkyl, $-(CH_2)_q^e$ carboxy, $-(CH_2)_q^e$ -S-lower alkyl,




-17-



wherein m^e , n^e , R_{10}^e and R_{11}^e are as defined above, R_{14}^e is lower alkyl, cycloalkyl, or

-18-







and q^{e} is an integer from 1 to 4,

R_2^{e} is hydrogen or lower alkyl,

R_3^{e} is hydrogen, lower alkyl, halo substituted lower alkyl, hydroxy substituted lower alkyl,

$\text{-(CH}_2\text{)}_{q^{\text{e}}}\text{-NH}_2$, $\text{-(CH}_2\text{)}_{q^{\text{e}}}\text{-N-(lower alkyl)}_2$,


$\text{-(CH}_2\text{)}_{q^{\text{e}}}$ guanidinyll, $\text{-(CH}_2\text{)}_{q^{\text{e}}}\text{-SH}$, $\text{-(CH}_2\text{)}_{q^{\text{e}}}\text{-S-}$ lower alkyl,













wherein R_{10}^{e} , n^{e} and q^{e} are as defined above.

(f) U.S. Patent No. 4,470,973 to Natarajan et al. discloses substituted peptides which are angiotensin converting enzyme inhibitors and have the formula


-19-




X^f is an amino or imino acid of the formula





-20-






R_7^f is hydrogen, lower alkyl, halogen, keto, hydroxy






azido, amino,




-21-



a 1- or 2-naphthyl of the formula




a 1- or 2-naphthyloxy of the formula




-S-lower alkyl,



-22-




or a 1- or 2-naphthylthio of the formula



R_8^f is keto, halogen.




-O-lower alkyl, a 1- or 2-naphthyloxy of the formula




-S-lower alkyl


-23-





or a 1- or 2-naphthylthio of the formula



-S-lower alkyl,




or a 1- or 2-naphthylthio of the formula

 R_9^f is keto or

R_{10}^f is halogen or $-Y^f-R_{16}^f$.
 R_{11}^f , R_{11}^f , R_{12}^f and R_{12}^f are independently
 selected from hydrogen and lower alkyl or R_{11}^f ,
 R_{12}^f and R_{12}^f are hydrogen and R_{11}^f is

-24-



R_{13}^f is hydrogen, lower alkyl of 1 to 4 carbons, lower alkoxy of 1 to 4 carbons, lower alkylthio of 1 to 4 carbons, chloro, bromo, fluoro, trifluoromethyl, hydroxy, phenyl, phenoxy, phenylthio, or phenylmethyl.

R_{14}^f is hydrogen, lower alkyl of 1 to 4 carbons, lower alkoxy of 1 to 4 carbons, lower alkylthio of 1 to 4 carbons, chloro, bromo, fluoro, trifluoromethyl, or hydroxy.


m^f is zero, one, two, three, or four.

p^f is one, two or three provided that p is more than one only if R_{13}^f or R_{14}^f is hydrogen, methyl, methoxy, chloro, or fluoro.

R_{15}^f is hydrogen or lower alkyl of 1 to 4 carbons.


Y^f is oxygen or sulfur.

R_{16}^f is lower alkyl of 1 to 4 carbons.





or the R_{16}^f groups join to complete an unsubstituted 5- or 6-membered ring or said ring in which one or more of the carbons has a lower alkyl of 1 to 4 carbons or a di(lower alkyl of 1 to 4 carbons) substituent.






R_4^f is hydrogen, lower alkyl.



-25-



R_5^f is hydrogen, lower alkyl,




r^f is an integer from 1 to 4.

R_{19}^f is lower alkyl, benzyl, or phenethyl.

R₂₀^f is hydrogen, lower alkyl, benzyl or phenethyl.






Rf is hydrogen, lower alkyl, cycloalkyl,

-26-











$-(\text{CH}_2)_2\text{-NH}_2$, $-(\text{CH}_2)_3\text{-NH}_2$, $-(\text{CH}_2)_4\text{-NH}_2$,
 $-(\text{CH}_2)_2\text{-OH}$, $-(\text{CH}_2)_3\text{-OH}$, $-(\text{CH}_2)_4\text{-OH}$,
 $-(\text{CH}_2)_2\text{-SH}$, $-(\text{CH}_2)_3\text{-SH}$, or $-(\text{CH}_2)_4\text{-SH}$.

R_1^f is hydrogen, lower alkyl, halo substituted lower alkyl,



provided that R_1^f is hydrogen only if R^f is other than hydrogen.



R_2^f is



-28-


wherein m^f , R_{14}^f , p^f and r^f are as defined above.

R_6^f is hydrogen, lower alkyl, benzyl, benzhydryl,



R_{17}^f is hydrogen, lower alkyl, cycloalkyl, or phenyl.



R_{18}^f is hydrogen, lower alkyl, lower alkoxy, or phenyl or R_{17}^f and R_{18}^f taken together are $-(\text{CH}_2)_2-$, $-(\text{CH}_2)_3-$, $-\text{CH}=\text{CH}-$, or



R_{21}^f and R_{22}^f are independently selected from hydrogen and lower alkyl.


R_{23}^f is lower alkyl.

R_{24}^f is hydrogen, lower alkyl,



-29-

(g) British Specification No. 2095682 published October 6, 1982 discloses N-substituted-N-carboxyalkyl aminocarbonyl alkyl glycine derivatives which are said to be angiotensin converting enzyme inhibitors and have the formula



either

(A) R^G and R_9^G are OH, 1-6C alkoxy, 2-6C alkenyloxy, di-(1-6C alkyl)amino-(1-6C) alkoxy, 1-6C hydroxyalkoxy, acylamino-(1-6C)alkoxy, acyloxy-(1-6C)alkoxy, aryloxy, aryloxy-(1-6C)alkoxy, NH_2 , mono- or di-(1-6C alkyl)amino, hydroxyamino or aryl-(1-6C)alkylamino; R_1^G - R_5^G , R_7^G and R_8^G are 1-20C alkyl, 2-20C alkenyl, 2-20C alkynyl, aryl, aryl-(1-6C) alkyl having 7-12C or heterocyclyl-(1-6C)alkyl having 7-12C;

R_6^G is cycloalkyl, polycycloalkyl, partly saturated cycloalkyl or polycycloalkyl, cycloalkyl-(1-6C)alkyl having 3-20C, 6-10C aryl, aryl-(1-6C)alkyl, aryl-(2-6C)alkenyl or aryl-(2-6C) alkynyl; or

R_2^G and R_3^G together with the C and N atoms to which they are attached or R_3^G and R_5^G together with the N and C atoms to which they are attached form an N-heterocycle containing 3-5C or 2-4C and a S atom;

-30-

all alkyl, alkenyl and alkynyl are optionally substituted by OH, 1-6C alkoxy, thio(sic), 1-6C alkylthio, NH_2 , mono- or di-(1-6C alkyl)amino, halogen or NO_2 ;


all 'cycloalkyl' groups (including poly and partially unsaturated) are optionally substituted by halogen, 1-6C hydroxyalkyl, 1-6C alkoxy, amino-(1-6C alkyl)amino, di-(1-6C alkyl)amino, SH, 1-6C alkylthio, NO_2 or CF_3 ; and aryl groups are optionally substituted by OH, 1-6C alkoxy, NH_2 , mono- or di-(1-6C alkyl) amino, SH, 1-6C alkylthio, 1-6C hydroxyalkyl, 1-6C aminoalkyl, 1-6C thioalkyl, NO_2 , halogen, CF_3 , OCH_2O , ureido or guanidino; or

(B) R^g and R_9^g are H or 1-6C alkoxy;
 R_1^g and R_2^g are H, 1-6C alkyl, aryl-(1-6C) alkyl having 7-12C or heterocyclyl-(1-6C) alkyl having 6-12C;
 R_3^g - R_5^g , R_7^g and R_8^g are H or 1-6C alkyl;
 R_6^g is cycloalkyl, polycycloalkyl, partly saturated cycloalkyl or polycycloalkyl, cycloalkyl-(1-6C) alkyl having 3-20C, aryl or aryl-(1-6C) alkyl; and

aryl has 6-10C and is optionally substituted by 1-6C alkyl, 2-6C alkenyl, 2-6C alkynyl, OH, 1-6C alkoxy, NH_2 , mono- or di-(1-6C alkyl) amino, SH, 1-6C alkylthio, 1-6C hydroxyalkyl, 1-6C aminoalkyl, 1-6C thioalkyl, NO_2 , halogen, CF_3 , OCH_2O , ureido or guanidino.

(h) U.S. Patent No. 4,470,972 to Gold et al discloses 7-carboxyalkyl-aminoacyl-1,4-dithia-7-azaspiro[4.4]nonane-8-carboxylic acids which are angiotensin converting enzyme inhibitors and have the formula

-31-




wherein:

R^h is lower alkyl, benzyl, benzylthio, benzyloxy, phenylthio or phenoxy;

R_1^h is hydroxy or lower alkoxy;

R_2^h is hydrogen, lower alkyl or amino lower alkyl; and the pharmaceutically acceptable salts thereof.


(i) European Patent Application 0 050 800 published May 5, 1982 discloses carboxyalkyl dipeptides derivatives which are said to be angiotensin converting enzyme inhibitors and have the formula



or a pharmaceutically acceptable salt thereof, wherein R^i and R^{6i} are the same or different and are hydroxy, lower alkoxy, lower alkenyloxy, dilower alkylamino lower alkoxy, acylamino lower alkoxy, acyloxy lower alkoxy, aryloxy, aryllower alkoxy, amino, lower alkylamino, dilower alkylamino, hydroxyamino, aryllower alkylamino, or substituted aryloxy or substituted aryllower alkoxy wherein the substituent is methyl, halo or methoxy; R^{1i} is hydrogen, alkyl of from 1 to 10 carbon atoms, substituted lower alkyl wherein the substituent is hydroxy, lower alkoxy, aryloxy, substituted aryloxy, heteroaryloxy, substituted heteroaryloxy, amino, lower alkylamino, diloweralkylamino, acylamino, arylamino, substituted arylamino, guanidino, imidazolyl, indolyl, lower alkylthio, arylthio, substituted arylthio, carboxy, carbamoyl, lower alkoxy carbonyl, aryl, substituted aryl,

-32-


aralkyloxy, substituted aralkyloxy, aralkylthio or substituted aralkylthio, wherein the aryl or heteroaryl portion of said substituted aryloxy, heteroaryloxy, arylamino, arylthio, aryl, aralkyloxy, aralkylthio group is substituted with a group selected from halo, lower alkyl, hydroxy, lower alkoxy, amino, aminomethyl, carboxyl, cyano, or sulfamoyl; R^{2i} and R^{7i} are the same or different and are hydrogen or lower alkyl; R^{3i} is hydrogen, lower alkyl, phenyl lower alkyl, aminomethylphenyl lower alkyl, hydroxyphenyl lower alkyl, hydroxy lower alkyl, acylamino lower alkyl, amino lower alkyl, dimethylamino lower alkyl, guanidino lower alkyl, imidazolyl lower alkyl, indolyl lower alkyl, or lower alkyl thio lower alkyl; R^{4i} and R^{5i} are the same or different and are hydrogen, lower alkyl or Z^i , or R^{4i} and R^{5i} taken together form a group represented by Q^i , U^i , V^i , Y^i , D^i or E^i , wherein; Z^i is



wherein X^{1i} and X^{2i} independent of each other are O, S or CH_2 , R^{8i} and R^{9i} independent of each other are lower alkyl, lower alkenyl, lower alkynyl, cycloalkyl having 3 to 8 carbon atoms, hydroxy lower alkyl, or $-(CH_2)_{n^i} Ar^i$, wherein n^i is 0, 1, 2 or 3 and Ar^i is unsubstituted or substituted phenyl, furyl, thienyl or pyridyl, wherein said substituted phenyl, furyl, thienyl or pyridyl groups are substituted with at least one group that is independently selected from C_1 to C_4 alkyl, lower alkoxy, lower alkylthio, halo, CF_3 and hydroxy, or R^{8i} and R^{9i} taken together form a bridge W^i , wherein W^i is a single bond or a methylene bridge or a substituted methylene


-33-

bridge when at least one of x^{1i} and x^{2i} is methylene, or w^i is an alkylene or substituted alkylene bridge having 2 or 3 carbon atoms, said substituted methylene bridge or said substituted alkylene bridge having one or two substituents selected from lower alkyl, aryl and aryl lower alkyl groups, and p^i is 0, 1 or 2; with the proviso that at least one of R^{4i} and R^{5i} is z^i , with the proviso that if R^{4i} is z^i and p^i is 0 then x^{1i} and x^{2i} must both be methylene, and with the proviso that if x^{1i} and x^{2i} are both methylene then R^{8i} and R^{9i} must form an alkylene bridge w^i ;
 q^i is




wherein R^{8i} , R^{9i} , x^{1i} and x^{2i} are as defined above, p^i is 0, 1 or 2, q^i is 0, 1 or 2, with the proviso that the sum of p^i and q^i must be 1, 2 or 3, with the proviso that if p^i is 0 then x^{1i} and x^{2i} must be methylene, and with the proviso that if x^{1i} and x^{2i} are methylene then R^{8i} and R^{9i} taken together form a bridge w^i , wherein w^i is as defined above;

v^i is




-34-

wherein R^{8i} , R^{9i} , X^{1i} and X^{2i} are as defined above, p^i is 0, 1 or 2 and q^i is 0, 1 or 2, with the proviso that the sum of p^i and q^i is 1, 2 or 3, with the proviso that if X^{1i} and X^{2i} are CH_2 then R^{8i} and R^{9i} taken together form a bridge w^i , wherein w^i is as defined above; U^i is




wherein W^i is as defined above (except that W^i may also be a methylene bridge when X^{1i} and X^{2i} are oxygen or sulfur), X^{1i} and X^{2i} are as defined above, p^i is 0, 1 or 2, q^i is 0, 1 or 2, with the proviso that the sum of p^i and q^i is 1 or 2, and with the proviso that if p^i is 0, X^{1i} must be CH_2 ; Y^i is




wherein G^i is oxygen, sulfur or CH_2 , a^i is 2, 3 or 4 and b^i is 1, 2, 3, 4 or 5, with the proviso that the sum of a^i and b^i is 5, 6 or 7 or G^i is CH_2 , a^i is 0, 1, 2 or 3, b^i is 0, 1, 2 or 3 with the proviso that the sum of a^i and b^i is 1, 2 or 3, with the proviso that the sum of a^i and b^i may be 1, 2 or 3 only if R^{li} is lower alkyl substituted with aralkylthio or aralkyloxy;

-35-


 D^i is

wherein F^i is O or S, j^i is 0, 1 or 2 and k^i is 0, 1 or 2, with the proviso that the sum of j^i and k^i must be 1, 2 or 3, and m^i is 1, 2 or 3 and t^i is 1, 2 or 3, with the proviso that the sum of m^i and t^i must be 2, 3 or 4;
 E^i is




wherein L^i is O or S, u^i is 0, 1 or 2 and v^i is 0, 1 or 2, with the proviso that the sum of u^i and v^i must be 1 or 2, and h^i is 1 or 2 and s^i is 1 or 2, with the proviso that the sum of h^i and s^i must be 2 or 3.

(j) European Patent Application 0 037 231 published October 7, 1981 discloses acyl derivatives of octahydro-1H-indole-2-carboxylic acids which are said to be angiotensin converting enzyme inhibitors and have the formula



-36-


wherein R^j is hydrogen or lower alkyl; R_1^j is hydrogen,
 lower alkyl, or benzyl; R_2^j is hydrogen or $\overset{\text{O}}{\parallel} R_3^j - C$ -wherein
 R_3^j is lower alkyl, heteroaryl containing 4 to 9 carbon
 atoms and one or two nitrogen, oxygen or sulfur atoms;
 phenyl, substituted phenyl having 1 or 2 substituents
 selected from the group consisting of fluorine, chlorine,
 bromine, lower alkyl or alkoxy; and n is 0 or 1; wherein
 lower alkyl and lower alkoxy include straight or branched
 groups containing 1 to 4 carbon atoms, and
 pharmaceutically acceptable salts of the compounds when
 R^j is hydrogen and when R_3^j is heteroaryl containing 1 or
 2 nitrogen atoms, are disclosed. Also disclosed are
 substituted acyl compounds of octahydro-1H-indole-2-
 carboxylic acid having the formula



wherein R_4^k is hydrogen or lower alkyl; R_5^k is hydrogen,
 lower alkyl or benzyl; R_6^k is hydrogen or lower alkyl;
 Ar^k is phenyl or phenyl substituted with 1 or 2
 substituents selected from the group consisting of
 fluorine, chlorine, bromine, lower alkyl, lower alkoxy,
 hydroxy or amino; and m is 0 to 3; wherein lower alkyl
 and lower alkoxy contain 1 to 4 straight or branched
 carbon atoms; and the pharmaceutically acceptable salts
 thereof.

(L) European Patent Application 0 079 522
 published May 25, 1983 discloses N-carboxymethyl(amino)
 lysyl-proline compounds which are said to be angiotensin
 converting enzyme inhibitors and have the formula where




-37-




wherein:

R^{L} and $\text{R}^{2\text{L}}$ are independently hydrogen; loweralkyl; aralkyl; or aryl;
 $\text{R}^{1\text{L}}$ is hydrogen; branched or straight chain C_1 - 12 alkyl and alkenyl; C_3 - C_9 cycloalkyl and benzofused alkyl; substituted loweralkyl where the substituents are halo, hydroxy loweralkoxy, aryloxy, amino, mono- or diloweralkylamino, acylamino, arylamino, guanidino, mercapto, loweralkylthio, arylthio, carboxy, carboxamido, or loweralkoxycarbonyl; aryl; substituted aryl where the substituents are loweralkyl, loweralkoxy, or halo; arloweralkyl; arloweralkenyl; heteroarloweralkyl; heteroarloweralkenyl; substituted arloweralkyl, substituted arloweralkenyl, substituted heteroarloweralkyl, or substituted heteroarloweralkenyl where the aryl and heteroaryl substituents are halo, dihalo, loweralkyl, hydroxy, loweralkoxy, amino, aminoloweralkyl, acylamino, mono- or diloweralkylamino, carboxyl, haloloweralkyl, nitro, cyano, or sulfonamido, and where the loweralkyl portion of arloweralkyl may be substituted by amino, acylamino, or hydroxyl;

-38-





or



where:

X^{L} and Y^{L} taken together are $-\text{CH}_2-\text{CH}_2-$;



R^{4L} is hydrogen; loweralkyl; aryl; substituted aryl;
 R^{5L} is hydrogen; loweralkyl; aryl or substituted aryl;
 n^{L} is 1 to 3;

W^{L} is absent; $-\text{CH}_2-$; or $-\text{C}(=\text{O})-$;
 Z^{L} is $-(\text{CH}_2)_m^{\text{L}}$, where m is 0 to 2, provided that m^L may not be 0 and w^L may not be absent at the same time; and

-39-

R^{6L} is hydrogen; loweralkyl; halo; or OR^{4L} ;

R^{2L} is $\text{---} (CH_2)_r^L \text{---} B^L \text{---} (CH_2)_s^L \text{---} NR^{7L}R^{15L}$

where

r^L and s^L are independently 0 to 3;

B^L is absent; -O-; -S-; or $-NR^{8L}$;

where R^{8L} is hydrogen; loweralkyl; alkanoyl; or aroyl; and

R^{7L} is $\begin{array}{c} NR^{11L} \\ || \\ -C-R^{9L} \end{array}$; $\begin{array}{c} NR^{11L} \\ || \\ -C-NHR^{10L} \end{array}$; or $\begin{array}{c} N-J^L \\ || \\ -C \begin{array}{c} | \\ K^L \end{array} \end{array} R^{12L}$

where

R^{9L} is loweralkyl; aralkyl; aryl; heteroaryl; or heteroarloweralkyl and these groups substituted by hydroxy, lower alkoxy or halo; carboxyl; carboxamido; nitromethenyl.

R^{10L} is hydrogen; loweralkyl; aryl; or amidino;

R^{11L} is hydrogen; loweralkyl; cyano; amidino; aryl; aroyl; loweralkanoyl; $-C-NHR^{13L}$;

$\begin{array}{c} O \\ || \\ -C-OR^{13L} \end{array}$; $-NO_2$; $-SO_2NH_2$;

or

SO_2R^{13L} ;

R^{12L} is hydrogen; loweralkyl; halo; aralkyl; amino; cyano; mono- or diloweralkylamino; or OR^{4L} ;

R^{13L} is hydrogen; loweralkyl; or aryl;

R^{15L} is hydrogen; lower alkyl; aralkyl; or aryl.


$\begin{array}{c} N-J^L \\ || \\ -C \begin{array}{c} | \\ K^L \end{array} \end{array} R^{12L}$

constitute a basic heterocycle of 5 or 6 atoms or benzofused analogs thereof and optionally containing 1-3 N atoms, an oxygen, a sulfur, an

-40-

- S=O, or an SO₂ group optionally substituted by amino, lower alkyl amino, diloweralkyl amino, lower alkoxy, or aralkyl groups;
- R^{3L} is C₃₋₈ cycloalkyl and benzofused C₃₋₈ cycloalkyl; perhydrobenzofused C₃₋₈ cycloalkyl; aryl; substituted aryl; heteraryl; substituted heteroaryl;
- R^{14L} is hydrogen or loweralkyl; and, a pharmaceutically acceptable salt thereof.

(m) U.S. Patent No. 4,256,761 to Suh et al discloses amides which are angiotensin converting enzyme inhibitors and have the general formula



wherein


R₁^m, R₂^m, R₃^m, R₄^m, R₅^m and R₆^m are hydrogen, alkyl, alkenyl, alkynyl, phenyl-alkyl, and cycloalkyl, and may be the same or different, n^m is an integer from 0 to 4 inclusive, M^m is alkenyl, alkynyl, cycloalkyl, cycloalkyl-alkyl, polycycloalkyl, polycycloalkyl-alkyl, aryl, aralkyl, heteroaryl, heteroaryl-alkyl, hetero-cycloalkyl, hetero-cycloalkyl-alkyl, fused aryl-cycloalkyl, fused aryl-cycloalkyl-alkyl, fused heteroaryl-cycloalkyl, fused heteroaryl-cycloalkyl-alkyl, alkoxyalkyl, alkylthioalkyl, alkylamino-alkyl, or dialkylaminoalkyl, Y^m is hydroxy, alkoxy, amino, or substituted amino, aminoalkanoyl, aryloxy, aminoalkoxy, or hydroxyalkoxy, and

-41-

R_7^m is hydrogen, alkanoyl, carboxyalkanoyl, hydroxyalkanoyl, aminoalkanoyl, cyano, amidino, carbalkoxy, Z^mS or




wherein Z^m is hydrogen, alkyl, hydroxyalkyl, or the radical



wherein R_1^m , R_2^m , R_3^m , R_4^m , R_5^m , R_6^m , n^m , M^m and Y^m are as described above; and where Y is hydroxy, their non-toxic, pharmaceutically acceptable alkali metal, alkaline earth metal, and amine salts.


(n) U.S. Patent No. 4,344,949 to Hoefle et al discloses acyl derivatives of 1,2,3,4-tetrahydroisoquinoline-3-carboxylic acid compounds which are angiotensin converting enzyme inhibitors and have the formula



-42-

wherein R^n is hydrogen, lower alkyl or aralkyl; R_1^n is hydrogen, lower alkyl, or benzyl; R_2^n is hydrogen or lower alkyl, and Ar^n is phenyl or phenyl substituted with 1 or 2 substituents selected from the group consisting of fluorine, chlorine, bromine, lower alkyl, lower alkoxy, hydroxy or amino; X^n and Y^n are independently hydrogen, lower alkyl, lower alkoxy, lower alkylthio, lower alkylsulfinyl, lower alkylsulfonyl, hydroxy, or X^n and Y^n together are methylenedioxy; m^n is 0 to 3; and the pharmaceutically acceptable acid salts thereof.

(b) European Patent 79022 published May 18, 1983 discloses N-amino acyl-azabicyclooctane carboxylic acid derivatives which have the formula



hydrogen atoms at ring positions 1 and 5 are cis to each other and the 3-carboxy group has the endo orientation;

R_1^O is H, allyl, vinyl or the side chain of an optionally protected naturally occurring α -amino acid;


R_2^O is H, 1-6C alkyl, 2-6C alkenyl or aryl(1-4C alkyl);

Y^O is H or OH and Z^O is H, or Y^O and Z^O together are oxygen;

X^O is 1-6C alkyl, 2-6C alkenyl, 5-9C cycloalkyl, 6-12C aryl (optionally substituted by one to three 1-4C alkyl or alkoxy, OH, halo, nitro, amino (optionally substituted by one or two 1-4C alkyl), or methylenedioxy) or indol-3-yl).

-43-

(p) European Patent 46953 published March 10, 1982 discloses N-amino acyl-indoline and tetrahydro isoquinoline carboxylic acids which are angiotensin converting enzyme inhibitors and have the formula



n^{P} is 0 or 1;



is a benzene or cyclohexane ring:

R_1^{P} and R_2^{P} are each 1-6C alkyl, 2-6C alkenyl, 5-7C cycloalkyl, 5-7C cycloalkenyl, 7-12C cycloalkylalkyl, optionally partially hydrogenated 6-10C aryl, 7-14C aralkyl or 5-7 membered monocyclic or 8-10 membered bicyclic heterocyclyl containing 1 or 2 S or O and/or 1-4N atoms; all R_1^{P} and R_2^{P} groups are optionally substituted,
 R_3^{P} is H, 1-6C alkyl, 2-6C alkenyl or 7-14C aralkyl.

Preferred R_1^{P} is Me and R_2^{P} is phenethyl optionally substituted by halogen, Me or OMe.

The above descriptions of suitable classes of ACE inhibitors for use in the present invention were taken from the noted patents and publications or abstracts thereof. Reference should be made to such patents and publications themselves for their full disclosures of such classes and specific compounds within such classes. In describing such suitable ACE inhibitors

-44-

the superscript letters a-p were included to distinguish among the various classes of compounds and the variable substituent groups thereof.

For the purposes of the invention, the preferred ACE inhibitor is 7-[N-(1(S)-carboxy-3-phenylpropyl)-(S)-alanyl]-1,4-dithia-7-azaspiro[4.4]nonane-8(S)-carboxylic acid.

It is believed that any ACE inhibitor will possess the novel utility described herein; however, for purposes of the present invention the preferred ACE inhibitors are compounds which are capable of inhibiting the action of ACE by at least 50% at a concentration of 1 μ M or less when tested by the following standard method:

ACE activity is determined by spectrophotometric assay of the product of the hydrolysis of the synthetic substrate, hippuryl histidyl leucine (HHL), as described by Cushman and Cheung, *Biochem. Pharmacol.*, 20, 1637 (1971). The ACE used is prepared in a manner similar to that of Cheung and Cushman, *Biochem. Biophys. Acta.*, 293, 451 (1973). Incubation for ACE assays is carried out at 37°C. Each 0.25 ml assay mixture contains the following components: 100 mM potassium phosphate buffer containing 300 mM sodium chloride, 5 mM HHL and 1.87 mU enzyme at pH 8.3 and various concentrations of inhibitors. The enzyme reaction is terminated after 60 minutes by the addition of 0.25 ml of 1N HCl. Inhibitors are dissolved in appropriate solvents. Hippuric acid solution for a standard curve is prepared in a similar manner.

Each experiment involves replicate incubations for each condition to be studied. IC_{50} values (the concentration required for the 50% inhibition of ACE activity) are derived from calculated regression lines. Each experiment utilizes multiple concentrations of inhibitor.

-45-

The ACE inhibitors of the invention may exist in isomeric form. The invention contemplates all such isomers both in pure form and admixture, including racemic mixtures.

Many ACE inhibitors are known in the art and may be prepared by known methods or by variations thereof. The following example illustrates the preparation of the preferred ACE inhibitor of the invention..

-46-

Example 17-[N-(1(S)-Carboxy-3-phenylpropyl)-(S)-alanyl]-1,4-dithia-7-azaspiro[4.4]nonane-8(S)-carboxylic acid

A. Dissolve 1-benzyloxycarbonyl-4-keto-(S)-proline (15.7 g, 60 mmol) in acetic acid (45 ml) and ethyl acetate (45 ml), add 1,2-ethanedithiol (6.2 g, 66 mmol) and stir under nitrogen. Add boron trifluoride etherate (redistilled) and stir 4 hours. Pour the reaction mixture into a mixture of water (175 ml) ethyl acetate (120 ml) and ether (60 ml). Wash the organic layer with water (3 x 175 ml) then extract twice with 1.0 N sodium bicarbonate (150 ml, then 50 ml). Acidify the aqueous layer to pH 1 with concentrated hydrochloric acid, extract twice with ether (150 ml, then 50 ml). Wash the organic layer with brine, dry over anhydrous magnesium sulfate and concentrate in vacuo to obtain crude 7-benzyloxycarbonyl-1,4-dithia-7-azaspiro[4.4]-nonane-8(S)-carboxylic acid as an oil (16.1 g).

B. Dissolve the product of Step A in acetic acid (30 ml) by stirring under nitrogen. Add 23% hydrobromic acid in acetic acid (50 ml) stir 3 hours and dilute over 5 minutes with ether (320 ml). Filter the solid, wash twice with ether (2 x 150 ml) and air dry to obtain a purple powder.

Dissolve the crude product in boiling ethanol (120 ml), add Darco grade 6-60 (6 g), and filter hot, washing with 2 x 10 ml hot ethanol. Dilute the filtrate with ether (300 ml) and allow to stand one hour. Filter the resultant precipitate, wash with ether (2 x 50 ml), and air dry to give 1,4-dithia-7-azaspiro[4.4]nonane-8(S)-carboxylic acid hydrobromide as beige crystals.

C. (1)a. To a 5L flask equipped with a magnetic stirrer, dropping funnel and nitrogen inlet tube, add a solution of 190 g (0.92 mole) of S-alanine

-47-

benzyl ester p-toluenesulfonate and 258 g (0.734 mole) of ethyl 2-oxo-4-phenyl butanoate in 1.4L of ethanol. Stir for 2 hours under nitrogen. Add a solution of 17.7 g (0.282 mole) of sodium cyanoborohydride in 550 mL of ethanol, with stirring, over 90 minutes. Stir the solution overnight and concentrate to dryness in vacuo at room temperature. Partition the residue between 500 mL of H₂O and 2L of ether. Dry the ether layer over MgSO₄ and filter. Add 1.3M ethereal HCl to pH 4. Remove the ether and excess HCl in vacuo at room temperature. Slurry the residue in 250 mL of ether and dilute with 750 mL of hexane. Decant the supernatant from the resulting precipitate and wash with two 300 mL portions of ether. Triturate the residue with 300 mL of ether and filter under N₂ to give 134 g of a white solid. Slurry the solid in ether and make basic with saturated aqueous NaHCO₃. Dry the organic layer over MgSO₄, filter and concentrate in vacuo at room temperature to give 135 g of N-(1-carboethoxy-3-phenylpropyl)-(S)-alanine benzyl ester as an amber oil.

Dissolve 135 g of the resultant amber oil in 510 ml of ethyl acetate and add a hot solution of 40.5 g maleic acid in 895 ml of ethyl acetate. Cool to room temperature, filter the resulting precipitate, and recrystallize the precipitate from ethyl acetate to give N-(1(S)-carboethoxy-3-phenylpropyl)-(S)alanine benzyl ester hemimaleate as a solid, m.p. 127-128°C, $[\alpha]_{D26} = 0^\circ(\text{H}_2\text{O})$.

Slurry 7.0 g (0.015 mole) of the above product in ethyl acetate and adjust to pH 8 with saturated aqueous sodium bicarbonate. Wash the organic layer with saturated aqueous sodium chloride solution, dry over magnesium sulfate, filter, and concentrate in vacuo at room temperature to a colorless oil. Dissolve the resultant oil in 100 ml ethanol containing 0.7 g of 10%

-48-

Pd/C and hydrogenate at 60 psi at room temperature for 2 hours. Filter and evaporate the solvent under vacuum at room temperature to obtain 4.0 g of N-(1(S)-carboethoxy-3-phenylpropyl)-(S)-alanine as a white solid, m.p. 147-148°C, $[\alpha]_D^{26} + 24.8^\circ$ (methanol).

(1)b. To a solution of the product of Step (1)a (8.93 g, 32 mmol) and N-hydroxysuccinimide (4.42 g, 38.4 mmol) in dry dimethylformamide (64 ml), add 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (7.3 g, 38.4 mmol) and stir for 12 hours. Pour the reaction mixture into ethyl acetate (320 ml), wash with water (100 ml), brine (100 ml), and dry the organic layer over anhydrous magnesium sulfate. Concentrate in vacuo to obtain N-[1(S)-ethoxycarbonyl-3-phenylpropyl]-(S)-alanine succinimidyl ester as an oil.

(2). Dissolve the product of Step (1)b and 1,4-dithia-7-azaspiro-[4.4]nonane-8(S)-carboxylic acid hydrobromide (Step B) (9.16 g, 32 mmol) in dimethylformamide (155 ml) and cool to 0-5°C. Add triethylamine (8.9 ml, 64 mmol) dropwise over 5 minutes, and rinse the funnel with dimethylformamide (5 ml). Stir at room temperature for 16 hours, concentrate in vacuo to 50 ml and dilute with water (150 ml). Adjust the solution to pH 4.0 with 1N hydrochloric acid. Extract the resultant brown gum with ethyl acetate (4 x 100 ml), combine the extracts, wash with water (20 ml) then brine (20 ml) and dry the organic phase over anhydrous magnesium sulfate. Evaporate the solvent in vacuo to obtain 7-[N-(1(S)-carboethoxy-3-phenylpropyl)-(S)-alanyl]-1,4-dithia-7-azaspiro[4.4]nonane-8(S)-carboxylic acid as a brown oil.

To purify, dissolve the crude product (3.3 g) in methanol (10 ml), place on Sephadex LH-20 (2.5 x 55 cm column) and elute with methanol. Combine the desired fractions and evaporate the solvent in vacuo to give

-49-

a solid white foam. Analytically pure material has an $[\alpha]_{D26} = -29.5^\circ$ (ethanol).

D. Hydrolyze the product of Step C(2) (0.3 g) in methanol (30 ml) with 2.5 N sodium hydroxide (2.0 ml) for 20 hours at room temperature. Concentrate the reaction mixture in vacuo at room temperature. Dissolve the resultant residue in water and place on Bio Rad AG-50W-X2 (100-200 mesh) resin in the hydrogen form. Elute with water (300 ml), then with 2% pyridine in water. Concentrate the desired eluant fractions to obtain the title compound, m.p. 115-117°C, $[\alpha]_{D26} = -22.4^\circ$ (H₂O); $+1.9^\circ$ (ethanol).

The IOP lowering effects of the compositions of the invention may be measured by the procedure described in Watkins et al., J. Ocular. Pharmacol., 1 (2):161 (1985).

Preferred pharmaceutical compositions of the invention comprise the combination of an ACE inhibitor and a steroid in an ophthalmologically acceptable carrier adapted for topical administration to the eye, such as solutions, suspensions, ointments and solid inserts. For solutions and suspensions, those skilled in the art will know that a particular dosage of active ingredient may be calculated if one assumes that one drop of solution is being administered and if one knows the concentration (w/v) of the particular solution that is being administered. Thus, one drop (1/20 ml) of a 0.25% solution (contains 2.5 mg of active per ml) is known to contain .125 mg or 125 μ g of active.

Topical ophthalmic formulations of the invention may combine the following amounts of each constituent:

ACE inhibitor from 0.00001 to 1.0% (w/v) and especially 0.001 to 0.01%. An amount of ACE inhibitor from between 0.005 μ g to 500 μ g, preferably 0.005 μ g to

-50-

50 μ g, and especially 0.005 μ g to 5 μ g of the active composition is applied to the human eye once or twice daily as needed. Individual dosage requirements, i.e., the amount of each dose and the frequency of administration, will depend on the potency of the particular ACE inhibitor, the severity of the increase in IOP and the response of the patient.

Steroid concentrations are preferably from 0.05 to 1.50% (w/v). An amount of steroid from between 20 μ g to 600 μ g of the active composition is applied to the human eye one or more times a day as needed. Individual dosage requirements, i.e. the amount of each dose and the frequency of administration, will depend on the potency of the particular steroid, the severity of the disease and the response of the patient. Appropriate dosage ranges for each steroid are well known to those skilled in the art. The particular steroid used will determine the ACE inhibitor chosen and the dosage of ACE inhibitor administered.

Each component may be administered separately, and separate administration is especially appropriate when the steroid is administered systemically or topically other than in an ophthalmological composition, since topical ophthalmological application of the ACE inhibitor is preferred. The concentration of the steroid in such systemic or topical compositions and the unit dosage weights thereof vary considerably, depending as above on such factors as the potency of the steroid, the severity of the disease and the response of the patient. Appropriate dosage ranges for systemic or topical administration of each steroid are well known in the art.

Since the present invention relates to treatment with a combination of an ACE inhibitor and a steroidal anti-inflammatory agent wherein the ACE

-51-

inhibitor and steroid are administered separately, the invention also relates to combining separate pharmaceutical compositions in kit form, that is, combining two separate units, an ACE inhibitor pharmaceutical composition and a steroid pharmaceutical composition, in a single package. Preferred components of the kit comprise a topical ophthalmological ACE inhibitor pharmaceutical composition and a pharmaceutically acceptable steroid composition. More preferred components of the kit are a topical ophthalmological ACE inhibitor pharmaceutical composition and a topical ophthalmological steroid pharmaceutical composition. A particular advantage of the more preferred embodiment of the kit resides in the ability to provide a combination of an ACE inhibitor composition which can be administered once or twice a day and a steroid composition which may be administered as frequently as once each hour.

Where utilized herein, the terms "controlling" and "suppressing" the elevated intraocular pressure mean the regulation, attenuation and modulation of increased intraocular tension. The term also means that the diminution in the otherwise elevated intraocular pressure obtained by the practice of the invention is maintained for a significant period of time as for example, between consecutive doses of the composition of the invention.

To prepare suitable dosage forms, the active compositions may be conveniently admixed with a non-toxic pharmaceutically acceptable carrier suitable for topical ophthalmologic administration. Typical of such pharmaceutically acceptable carriers are, for example, water, mixtures of water and watermiscible solvents such as lower alkanols or vegetable oils, petroleum based jelly, and including also from 0.5 to 5% by weight of hydroxyethyl cellulose, ethyl oleate, carboxymethyl

-52-

cellulose, polyvinylpyrrolidone, and other water soluble ophthalmologically acceptable non-toxic polymers, for example, cellulose derivatives such as methyl cellulose. The pharmaceutical preparation may also contain non-toxic auxiliary substances such as emulsifying, preserving, wetting, bodying agents and the like, as for example, polyethylene glycols 200, 300, 400 and 600, carbowaxes 1,000, 1,500, 4,000, 6,000 and 10,000, antibacterial components such as quarternary ammonium compounds; phenylmercuric salts known to have cold sterilizing properties and which are non-injurious in use; thimerosal, methyl and propyl paraben, benzyl alcohol, phenyl ethanol; buffering ingredients; antioxidants; and other conventional ingredients.

Additionally, suitable ophthalmic vehicles can be used as carrier media for the present purpose including conventional phosphate buffer vehicle systems, isotonic boric acid vehicles, isotonic alkali chloride vehicles, tris and the like.

The pharmaceutical preparation may also be in the form of a solid insert. For example, one may use a solid water soluble polymer as the carrier for the active ingredients. Inserts that are known in the art that are suitable for this use include those described in United States Patents 3,993,071; 3,986,510; 3,868,445; and 3,867,510. Solid water insoluble inserts, such as those prepared from ethylene vinyl acetate copolymer, may also be utilized.

The compositions of the invention may include additional therapeutic agents in addition to the ACE inhibitor and steroid. For example antibiotics, anesthetics as well as other IOP lowering agents may be present.

-53-

I CLAIM:

1. The use of one or more angiotensin converting enzyme inhibitors for the manufacture of a medicament for treating the rise in intraocular pressure associated with the use of steroidal anti-inflammatory drugs.
2. The use of claim 1 wherein the angiotensin converting enzyme inhibitor is 7-[N-(1(S)-carboxy-3-phenylpropyl)-(S)-alanyl]-1,4-dithia-7-azaspiro[4.4]nonane-8(S)-carboxylic acid.
3. A pharmaceutical composition for treating the rise in intraocular pressure associated with the use of steroidal anti-inflammatory drugs comprising an angiotensin converting enzyme inhibitor together with a pharmaceutically acceptable carrier.
4. The composition of claim 3 wherein the angiotensin converting enzyme inhibitor is 7-[N-(1(S)-carboxy-3-phenylpropyl)-(S)-alanyl]-1,4-dithia-7-azaspiro[4.4]nonane-8(S)-carboxylic acid.
5. The composition of claims 3 or 4 further comprising a steroidal anti-inflammatory drug.
6. A kit comprising in separate containers pharmaceutical composition for controlling the rise in intraocular pressure associated with the use of steroidal anti-inflammatory drugs which comprises in one container an angiotensin converting enzyme inhibitor in a pharmaceutically acceptable carrier and, in a second container, a steroidal pharmaceutical composition in a pharmaceutically acceptable carrier.

-54-

7. The kit of claim 6 further characterized by the angiotensin converting enzyme comprising a topical ophthalmologically acceptable angiotensin converting enzyme.

8. The kit of either of claims 6 or 7 further characterized by the steroidal pharmaceutical composition comprising a topical ophthalmological steroid pharmaceutical composition.

9. The kit of any of claims 6-8 above further characterized by the angiotensin converting enzyme comprising 7[N-(1(S)-carboxy-3-phenylpropyl)-(S)-alanyl]-1,4-dithia-7-azaspiro[4.4]nonane-8(S)-carboxylic acid.