

(19) **DANMARK**

(10) **DK/EP 2651437 T3**



Patent- og
Varemærkestyrelsen

(12) Oversættelse af
europæisk patentskrift

-
- (51) Int.Cl.: **A 61 K 39/04 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2016-05-02**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2016-02-10**
- (86) Europæisk ansøgning nr.: **11799673.6**
- (86) Europæisk indleveringsdag: **2011-12-14**
- (87) Den europæiske ansøgnings publiceringsdag: **2013-10-23**
- (86) International ansøgning nr.: **EP2011072816**
- (87) Internationalt publikationsnr.: **WO2012080369**
- (30) Prioritet: **2010-12-14 US 422723 P**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **GlaxoSmithKline Biologicals S.A., rue de l'Institut, 89, 1330 Rixensart, Belgien**
- (72) Opfinder: **GODART, Stephane Andre Georges, GlaxoSmithKline Biologicals s.a., rue de l'Institut 89, B-1330 Rixensart, Belgien**
LAANAN, Amina, GlaxoSmithKline Biologicals s.a., rue de l'Institut 89, B-1330 Rixensart, Belgien
LEMOINE, Dominique Ingrid, GlaxoSmithKline Biologicals s.a., rue de l'Institut 89, B-1330 Rixensart, Belgien
- (74) Fuldmægtig i Danmark: **Awapatent A/S, Rigensgade 11, 1316 København K, Danmark**
- (54) Benævnelse: **MYCOBAKTERIE-ANTIGEN SAMMENSÆTNING**
- (56) Fremdragne publikationer:
WO-A1-2010/142685
WO-A1-2010/142686
WO-A2-2006/117240
WO-A2-2009/088255
US-A1- 2009 291 062
US-B1- 6 869 607
LEROUX-ROELS ISABEL ET AL: "Evaluation of the Safety and Immunogenicity of Two Antigen Concentrations of the Mtb72F/AS02(A) Candidate Tuberculosis Vaccine in Purified Protein Derivative-Negative Adults", CLINICAL AND VACCINE IMMUNOLOGY, vol. 17, no. 11, November 2010 (2010-11), pages 1763-1771, XP002669941,

DESCRIPTION

Field of the Invention

[0001] The present invention relates to immunogenic compositions comprising an M72 related antigen and having a low ionic strength. The present invention also relates to such immunogenic compositions which further comprise one or more immunostimulants.

Background of the Invention

[0002] Tuberculosis (TB) is a chronic infectious disease caused by infection with *Mycobacterium tuberculosis* and other *Mycobacterium* species. It is a major disease in developing countries, as well as an increasing problem in developed areas of the world. More than 2 billion people are believed to be infected with TB bacilli, with about 9.4 million new cases of TB and 1.7 million deaths each year. 10% of those infected with TB bacilli will develop active TB, each person with active TB infecting an average of 10 to 15 others per year. While annual incidence rates have peaked globally, the number of deaths and cases is still rising due to population growth (World Health Organisation Tuberculosis Facts 2010).

[0003] The protein antigens Mtb72f and M72 (described, for example, in international patent application WO2006/117240) or fragments or derivatives thereof are protein antigens of potential benefit for the treatment or prevention of tuberculosis.

[0004] The formulation of protein antigens is extremely important in order to ensure immunogenicity is maintained. Immunostimulants are sometimes used to improve the immune response raised to any given antigen. However, the inclusion of adjuvants into an immunogenic composition increases the complexity of preparation of the components as well as the complexity of distribution and formulation of the composition. The preparation of each of the adjuvant components as well as the antigenic component must be considered by formulators. In particular, the compatibility of the antigenic component with the adjuvant component should be considered. This is particularly the case where lyophilised antigens or antigenic preparations are intended to be reconstituted with an adjuvant preparation. In such a circumstance, it is important that the buffer of the adjuvant preparation is suitable for the antigen and that immunogenicity or solubility of the antigen is not affected by the adjuvant.

[0005] WO2006/117240 discloses methods of preventing reactivation of *M. tuberculosis* infections using a pharmaceutical composition comprising M72 or Mtb72f fusion proteins.

Summary of the invention

[0006] The present inventors have identified for the first time that M72 related antigens are particularly sensitive to the presence of salts. Without being limited by theory, it is believed M72 related antigens are detrimentally impacted by a phenomenon known as "salting out" which may be defined as the precipitation of a protein from its solution by interaction with salts, such as sodium chloride. The present inventors have found that these antigens aggregate and precipitate at a concentration of sodium chloride as low as 150 mM. Consequently, the stability of immunogenic compositions comprising M72 related antigens can surprisingly be improved by a reduction in the concentration of sodium chloride.

[0007] Accordingly, the present invention provides an immunogenic composition comprising an M72 related antigen, wherein the M72 related antigen comprises a sequence having at least 90% identity to SEQ ID No: 1, wherein the conductivity of the composition is 5 mS/cm or lower and the pH of the composition is in the range 7.0 to 9.0.

Brief description of the drawings

[0008]

Figure 1.

QS21 lytic activity curve

Figure 2.

Percentage of each 3D-MPL congener in the different ASA formulations

Figure 3.

DLS of immunogenic compositions with varied pH and NaCl concentrations after storage

Figure 4.

Nephelometry of immunogenic compositions with varied pH and NaCl concentrations after storage

Figure 5.

Antigenic stability of immunogenic compositions with varied pH and NaCl concentrations following after storage

Figures 6a-6d.

SEC-HPLC analysis of immunogenic compositions with varied pH and NaCl concentrations after storage

Figure 7.

Antigenicity of immunogenic compositions with varied pH and NaCl concentrations after storage

Figure 8.

Conductivity of NaCl standard solutions

Figure 9.

Induction of CD4 T cell responses in mice using immunogenic compositions of the invention

Figure 10.

Induction of CD8 T cell responses in mice using immunogenic compositions of the invention

Figure 11.

Nephelometry of immunogenic compositions with varied pH and NaCl concentrations after storage

Figure 12.

DLS of immunogenic compositions with varied pH and NaCl concentrations after storage

Figure 13.

Antigenicity of immunogenic compositions with varied NaCl concentrations after storage

Brief description of sequence identifiers

[0009]

SEQ ID No: 1 Amino acid sequence for the M72 protein

SEQ ID No: 2 Nucleotide sequence encoding the M72 protein

SEQ ID No: 3 Amino acid sequence for the M72 protein with two N-terminal His residues

SEQ ID No: 4 Nucleotide sequence encoding the M72 protein with two N-terminal His residues

SEQ ID No: 5 Amino acid sequence for the Mtb72f protein

SEQ ID No: 6 Nucleotide sequence encoding the Mtb72f protein

SEQ ID No: 7 Amino acid sequence for the Mtb72f protein with six N-terminal His residues

SEQ ID No: 8 Nucleotide sequence encoding the Mtb72f protein with six N-terminal His residues

SEQ ID No: 9 Nucleotide sequence for CpG Oligo 1 (CpG 1826)

SEQ ID No: 10 Nucleotide sequence for CpG Oligo 2 (CpG 1758)

SEQ ID No: 11 Nucleotide sequence for CpG Oligo 3

SEQ ID No: 12 Nucleotide sequence for CpG Oligo 4 (CpG 2006)

SEQ ID No: 13 Nucleotide sequence for CpG Oligo 5 (CpG 1686)

Detailed description of the invention

[0010] In a first aspect, the present invention provides an immunogenic composition comprising an M72 related antigen, wherein the M72 related antigen comprises a sequence having at least 90% identity to SEQ ID No: 1, the pH of the composition is in the range 7.0 to 9.0 and wherein the conductivity of the composition is 5 mS/cm or lower, 4 mS/cm or lower, or 3 mS/cm or lower. In a particular embodiment the conductivity of the immunogenic composition is 2.5 mS/cm or lower, such as 2.25 mS/cm or lower, or 2.0 mS/cm or lower. In a further specific embodiment the conductivity of the immunogenic composition is 1.5 to 2.5 mS/cm.

[0011] In some embodiments, the present invention provides an immunogenic composition wherein the concentration of salts in said composition is 40 mM or lower. In a particular embodiment the concentration of salts in said composition is 35 mM or lower, such as 30 mM or lower, or 25 mM or lower. In a further specific embodiment the concentration of salts in said composition is 20 to 40 mM, such as 25 to 35 mM.

[0012] In some embodiments the concentration of sodium chloride is 40 mM or lower, 30 mM or lower, 20 mM or lower or 15 mM or lower. In a particular embodiment the concentration of sodium chloride in the immunogenic composition is 10 mM or lower, such as 7.5 mM or lower. Suitably the concentration of sodium chloride in the immunogenic composition is at or below 5 mM. In a further specific embodiment, the immunogenic composition is essentially free of sodium chloride. By essentially free is meant that the concentration of sodium chloride is at or very near to zero mM (such as 3 mM or less, 2 mM or less or 1 mM or less).

[0013] Suitably, the concentration of CaCl_2 in the immunogenic compositions will be 30 mM or lower, 20 mM or lower, 15 mM or lower or 10 mM or lower.

[0014] Suitably, the concentration of MgSO_4 in the immunogenic compositions will be 60 mM or lower, 40 mM or lower, 30 mM or lower, 20 mM or lower or 10 mM or lower.

[0015] Suitably, the total concentration of NH_4^+ , Mg^{2+} and Ca^{2+} ions in the immunogenic compositions will be 60 mM or lower, 40 mM or lower, 30 mM or lower, 20 mM or lower or 10 mM or lower.

[0016] The immunogenic compositions of the invention will be aqueous preparations.

[0017] The conductivity of an immunogenic composition of the invention can be measured using techniques known in the art, for example using a dedicated conductivity meter or other instrument with the capability to measure conductivity. One suitable instrument is the Zetasizer Nano ZS from Malvern Instruments (UK).

[0018] The skilled person can readily test for the concentration of both sodium (Na^+) and chloride (Cl^-) ions using known techniques and kits. For example, sodium can be determined using a kit such as the Sodium Enzymatic Assay Kit (Catalogue Number: BQ011EAE) from Biosupply. Chloride can be determined using a kit such as Chloride Enzymatic Assay Kit (Catalogue Number: BQ006EAE) from Biosupply.

[0019] Tuberculosis (TB) is a chronic infectious disease caused by infection with *Mycobacterium tuberculosis* and other *Mycobacterium* species. It is a major disease in developing countries, as well as an increasing problem in developed areas of the world. More than 2 billion people are believed to be infected with TB bacilli, with about 9.4 million new cases of TB and 1.7 million deaths each year. 10% of those infected with TB bacilli will develop active TB, each person with active TB infecting an average of 10 to 15 others per year. While annual incidence rates have peaked globally, the number of deaths and cases is still rising due to population growth (World Health Organisation *Tuberculosis Facts* 2010).

[0020] *Mycobacterium tuberculosis* infects individuals through the respiratory route. Alveolar macrophages engulf the bacterium, but it is able to survive and proliferate by inhibiting phagosome fusion with acidic lysosomes. A complex immune response involving CD4^+ and CD8^+ T cells ensues, ultimately resulting in the formation of a granuloma. Central to the success of *Mycobacterium tuberculosis* as a pathogen is the fact that the isolated, but not eradicated, bacterium may persist for long periods, leaving an individual vulnerable to the later development of active TB.

[0021] Fewer than 5% of infected individuals develop active TB in the first years after infection. The granuloma can persist for decades and is believed to contain live *Mycobacterium tuberculosis* in a state of dormancy, deprived of oxygen and nutrients. However, recently it has been suggested that the majority of the bacteria in the dormancy state are located in non-macrophage cell types spread throughout the body (Locht et al, Expert Opin. Biol. Ther. 2007 7(11):1665-1677). The development of active TB occurs when the balance between the host's natural immunity and the pathogen changes, for example as a result of an immunosuppressive event (Anderson P Trends in Microbiology 2007 15(1):7-13; Ehlers S Infection 2009 37(2):87-95).

[0022] A dynamic hypothesis describing the balance between latent TB and active TB has also been proposed (Cardana P-J Inflammation & Allergy - Drug Targets 2006 6:27-39; Cardana P-J Infection 2009 37(2):80-86).

[0023] Although an infection may be asymptomatic for a considerable period of time, the active disease is most commonly manifested as an acute inflammation of the lungs, resulting in tiredness, weight loss, fever and a persistent cough. If untreated, serious complications and death typically result.

[0024] Tuberculosis can generally be controlled using extended antibiotic therapy, although such treatment is not sufficient to prevent the spread of the disease. Actively infected individuals may be largely asymptomatic, but contagious, for some time. In addition, although compliance with the treatment regimen is critical, patient behaviour is difficult to monitor. Some patients do not complete the course of treatment, which can lead to ineffective treatment and the development of drug resistance.

[0025] Multidrug-resistant TB (MDR-TB) is a form which fails to respond to first line medications. 3.3% of all TB cases are MDR-TB, with an estimated 440,000 new MDR-TB cases occurring each year. Extensively drug-resistant TB (XDR-TB) occurs when resistance to second line medications develops on top of resistance to first line medications. The virtually untreatable XDR-TB has been confirmed in 58 countries (World Health Organisation Tuberculosis Facts 2010).

[0026] Even if a full course of antibiotic treatment is completed, infection with *M. tuberculosis* may not be eradicated from the infected individual and may remain as a latent infection that can be reactivated. In order to control the spread of tuberculosis, an effective vaccination programme and accurate early diagnosis of the disease are of utmost importance.

[0027] Currently, vaccination with live bacteria is the most widely used method for inducing protective immunity. The most common *Mycobacterium* employed for this purpose is *Bacillus Calmette-Guerin* (BCG), an avirulent strain of *M. bovis* which was first developed over 60 years ago. However, the safety and efficacy of BCG is a source of controversy - while protecting against severe disease manifestation in children, BCG does not prevent the establishment of latent TB or the reactivation of pulmonary disease in adult life. Additionally, some countries, such as the United States, do not vaccinate the general public with this agent.

[0028] Several of the proteins which are strongly expressed during the early stages of *Mycobacterium* infection have been shown to provide protective efficacy in animal vaccination models. However, vaccination with antigens which are highly expressed during the early stages of infection may not provide an optimal immune response for dealing with later stages of infection. Adequate control during latent infection may require T cells which are specific for the particular antigens which are expressed at that time. Post-exposure vaccines which directly target the dormant persistent bacteria may aid in protecting against TB reactivation, thereby enhancing TB control, or even enabling clearance of the infection. A vaccine targeting latent TB could therefore significantly and economically reduce global TB infection rates.

[0029] Subunit vaccines based on late stage antigens could also be utilised in combination with early stage antigens to provide a multiphase vaccine. Alternatively, early and/or late stage antigens could be used to complement and improve BCG vaccination (either by boosting the BCG response or through the development of advanced recombinant BCG strains).

[0030] The protein antigens Mtb72f and M72 are protein antigens of potential benefit for the treatment or prevention of tuberculosis. Mtb72f has been shown to provide protection in a number of animal models (see, for example: Brandt et al Infect. Immun. 2004 72(11):6622-6632; Skeiky et al J. Immunol. 2004 172:7618-7628; Tsenova et al Infect. Immun. 2006 74(4):2392-2401; Reed et al PNAS 2009 106(7):2301-2306). Mtb72f has also been the subject of clinical investigations (Von Eschen et al 2009 Human Vaccines 5(7):475-482). M72 is an improved antigen which incorporates a single serine to alanine mutation relative to Mtb72f, resulting in improved stability characteristics. M72 related antigens have also been shown to be of value in a latent TB model (international patent application WO2006/117240).

[0031] As used herein the term 'M72 related antigen' refers to the M72 protein provided in SEQ ID No: 1 or an immunogenic derivative thereof. As used herein the term "derivative" refers to an antigen that is modified relative to the reference sequence. Immunogenic derivatives are sufficiently similar to the reference sequence to retain the immunogenic properties of the reference sequence and remain capable of allowing an immune response to be raised against the reference sequence. A derivative may, for example, comprise a modified version of the reference sequence or alternatively may consist of a modified version of the reference sequence.

[0032] The M72 related antigen may for example contain fewer than 1500 amino acid residues, such as fewer than 1200 amino acid residues, in particular less than 1000 amino acid residues, especially fewer than 800 amino acid residues.

[0033] T cell epitopes are short contiguous stretches of amino acids which are recognised by T cells (e.g. CD4+ or CD8+ T

cells). Identification of T cell epitopes may be achieved through epitope mapping experiments which are known to the person skilled in the art (see, for example, Paul, *Fundamental Immunology*, 3rd ed., 243-247 (1993); Beißbarth et al *Bioinformatics* 2005 21(Suppl. 1):i29-i37). In a diverse out-bred population, such as humans, different HLA types mean that particular epitopes may not be recognised by all members of the population. As a result of the crucial involvement of the T cell response in tuberculosis, to maximise the level of recognition and scale of immune response, an immunogenic derivative of M72 is desirably one which contains the majority (or suitably all) T cell epitopes intact.

[0034] The skilled person will recognise that individual substitutions, deletions or additions to the M72 protein which alters, adds or deletes a single amino acid or a small percentage of amino acids is an "immunogenic derivative" where the alteration(s) results in the substitution of an amino acid with a functionally similar amino acid or the substitution/deletion/addition of residues which do not substantially impact the immunogenic function.

[0035] Conservative substitution tables providing functionally similar amino acids are well known in the art. In general, such conservative substitutions will fall within one of the amino-acid groupings specified below, though in some circumstances other substitutions may be possible without substantially affecting the immunogenic properties of the antigen. The following eight groups each contain amino acids that are typically conservative substitutions for one another:

1. 1) Alanine (A), Glycine (G);
 2. 2) Aspartic acid (D), Glutamic acid (E);
 3. 3) Asparagine (N), Glutamine (Q);
 4. 4) Arginine (R), Lysine (K);
 5. 5) Isoleucine (I), Leucine (L), Methionine (M), Valine (V);
 6. 6) Phenylalanine (F), Tyrosine (Y), Tryptophan (W);
 7. 7) Serine (S), Threonine (T); and
 8. 8) Cysteine (C), Methionine (M)
- (see, e.g., Creighton, *Proteins* 1984).

[0036] Suitably such substitutions do not occur in the region of an epitope, and do not therefore have a significant impact on the immunogenic properties of the antigen.

[0037] Immunogenic derivatives may also include those wherein additional amino acids are inserted compared to the reference sequence. Suitably such insertions do not occur in the region of an epitope, and do not therefore have a significant impact on the immunogenic properties of the antigen. One example of insertions includes a short stretch of histidine residues (e.g. 2-6 residues) to aid expression and/or purification of the antigen in question.

[0038] Immunogenic derivatives include those wherein amino acids have been deleted compared to the reference sequence. Suitably such deletions do not occur in the region of an epitope, and do not therefore have a significant impact on the immunogenic properties of the antigen.

[0039] The skilled person will recognise that a particular immunogenic derivative may comprise substitutions, deletions and additions (or any combination thereof).

[0040] The terms "identical" or percentage "identity," in the context of two or more polypeptide sequences, refer to two or more sequences or sub-sequences that are the same or have a specified percentage of amino acid residues that are the same (i.e., 70% identity, optionally 75%, 80%, 85%, 90%, 95%, 98% or 99% identity over a specified region), when compared and aligned for maximum correspondence over a comparison window, or designated region as measured using one of the following sequence comparison algorithms or by manual alignment and visual inspection. This definition also refers to the complement of a test sequence. Optionally, the identity exists over a region that is at least 500 amino acids in length, such as at least 600 amino acids or at least 700 amino acids. Suitably, the comparison is performed over a window corresponding to the entire length of the reference sequence (as opposed to the derivative sequence).

[0041] For sequence comparison, one sequence acts as the reference sequence, to which the test sequences are compared. When using a sequence comparison algorithm, test and reference sequences are entered into a computer, subsequence coordinates are designated, if necessary, and sequence algorithm program parameters are designated. Default program parameters can be used, or alternative parameters can be designated. The sequence comparison algorithm then calculates the percentage sequence identities for the test sequences relative to the reference sequence, based on the program parameters.

[0042] A "comparison window", as used herein, refers to a segment in which a sequence may be compared to a reference sequence of the same number of contiguous positions after the two sequences are optimally aligned. Methods of alignment of sequences for comparison are well-known in the art. Optimal alignment of sequences for comparison can be conducted, e.g., by the local homology algorithm of Smith & Waterman, *Adv. Appl. Math.* 2:482 (1981), by the homology alignment algorithm of Needleman & Wunsch, *J. Mol. Biol.* 48:443 (1970), by the search for similarity method of Pearson & Lipman, *Proc. Natl. Acad. Sci. USA* 85:2444 (1988), by computerised implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Dr., Madison, WI), or by manual alignment and visual inspection (see, e.g., *Current Protocols in Molecular Biology* (Ausubel et al., eds. 1995 supplement)).

[0043] One example of a useful algorithm is PILEUP. PILEUP creates a multiple sequence alignment from a group of related sequences using progressive, pairwise alignments to show relationship and percent sequence identity. It also plots a tree or dendrogram showing the clustering relationships used to create the alignment. PILEUP uses a simplification of the progressive alignment method of Feng & Doolittle, *J. Mol. Evol.* 35:351-360 (1987). The method used is similar to the method described by Higgins & Sharp, *CABIOS* 5:151-153 (1989). The program can align up to 300 sequences, each of a maximum length of 5,000 nucleotides or amino acids. The multiple alignment procedure begins with the pairwise alignment of the two most similar sequences, producing a cluster of two aligned sequences. This cluster is then aligned to the next most related sequence or cluster of aligned sequences. Two clusters of sequences are aligned by a simple extension of the pairwise alignment of two individual sequences. The final alignment is achieved by a series of progressive, pairwise alignments. The program is run by designating specific sequences and their amino acid coordinates for regions of sequence comparison and by designating the program parameters. Using PILEUP, a reference sequence is compared to other test sequences to determine the percent sequence identity relationship using the following parameters: default gap weight (3.00), default gap length weight (0.10), and weighted end gaps. PILEUP can be obtained from the GCG sequence analysis software package, e.g., version 7.0 (Devereaux et al., *Nuc. Acids Res.* 12:387-395 (1984)).

[0044] Another example of algorithm that is suitable for determining percent sequence identity and sequence similarity are the BLAST and BLAST 2.0 algorithms, which are described in Altschul et al., *Nuc. Acids Res.* 25:3389-3402 (1977) and Altschul et al., *J. Mol. Biol.* 215:403-410 (1990), respectively. Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information (website at www.ncbi.nlm.nih.gov/). This algorithm involves first identifying high scoring sequence pairs (HSPs) by identifying short words of length W in the query sequence, which either match or satisfy some positive-valued threshold score T when aligned with a word of the same length in a database sequence. T is referred to as the neighbourhood word score threshold (Altschul *et al.*, *supra*). These initial neighbourhood word hits act as seeds for initiating searches to find longer HSPs containing them. The word hits are extended in both directions along each sequence for as far as the cumulative alignment score can be increased. Cumulative scores are calculated using, for nucleotide sequences, the parameters M (reward score for a pair of matching residues; always > 0) and N (penalty score for mismatching residues; always < 0). For amino acid sequences, a scoring matrix is used to calculate the cumulative score. Extension of the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters W, T, and X determine the sensitivity and speed of the alignment. The BLASTN program (for nucleotide sequences) uses as defaults a word length (W) of 11, an expectation (E) of 10, M=5, N=-4 and a comparison of both strands. For amino acid sequences, the BLASTP program uses as defaults a wordlength of 3, and expectation (E) of 10, and the BLOSUM62 scoring matrix (see Henikoff & Henikoff, *Proc. Natl. Acad. Sci. USA* 89:10915 (1989)) alignments (B) of 50, expectation (E) of 10, M=5, N=-4, and a comparison of both strands.

[0045] The BLAST algorithm also performs a statistical analysis of the similarity between two sequences (see, e.g., Karlin & Altschul, *Proc. Natl. Acad. Sci. USA* 90:5873-5787 (1993)). One measure of similarity provided by the BLAST algorithm is the smallest sum probability (P(N)), which provides an indication of the probability by which a match between two nucleotide or amino acid sequences would occur by chance. For example, a nucleic acid is considered similar to a reference sequence if the smallest sum probability in a comparison of the test nucleic acid to the reference nucleic acid is less than about 0.2, more preferably less than about 0.01, and most preferably less than about 0.001.

[0046] In any event, immunogenic derivatives of a polypeptide sequence will have essentially the same activity as the reference sequence. By essentially the same activity is meant at least 50%, suitably at least 75% and especially at least 90% activity of the reference sequence in an *in vitro* restimulation assay of PBMC or whole blood with specific antigens (e.g. restimulation for a period of between several hours to up to two weeks, such as up to one day, 1 day to 1 week or 1 to 2 weeks) that measures the activation of the cells via lymphoproliferation, production of cytokines in the supernatant of culture (measured by ELISA, CBA etc) or characterisation of T and B cell responses by intra and extracellular staining (e.g. using antibodies specific to immune markers, such as CD3, CD4, CD8, IL2, TNF-alpha, IFN-gamma, CD40L, CD69 etc) followed by analysis with a flowcytometer. Suitably, by

essentially the same activity is meant at least 50%, suitably at least 75% and especially at least 90% activity of the reference sequence in a T cell proliferation and/or IFN-gamma production assay.

[0047] Particular derivatives of the M72 protein include those with additional His residues at the N-terminus (e.g. two His residues, as provided in SEQ ID No: 3; or a polyhistadine tag of five or particularly six His residues, which may be used for nickel affinity purification). Mtb72f (SEQ ID No: 5) which contains the original serine residue that has been mutated in M72, is a further derivative of M72, as are Mtb72f proteins with additional His residues at the N-terminus (e.g. two His residues; or a polyhistadine tag of five or particularly six His residues, which may be used for nickel affinity purification).

[0048] Suitably an M72 related antigen will comprise, such as consist of, a sequence having at least 95%, for example at least 99% to SEQ ID No. 1. Optionally, an M72 related antigen will comprise, such as consist of, a sequence having at least 98% identity to SEQ ID No. 1.

[0049] Typical M72 related antigens will comprise, such as consist of, an immunogenic derivative of SEQ ID No: 1 or 3 having a small number of deletions insertions and/or substitutions. Examples are those having deletions of up to 5 residues at 0-5 locations, insertions of up to 5 residues at 0-5 five locations and substitutions of up to 20 residues.

[0050] Other immunogenic derivatives of M72 are those comprising, such as consisting of, a fragment of SEQ ID No: 1 or 3 which is at least 500 amino acids in length, such as at least 600 amino acids in length or at least 700 amino acids in length.

[0051] M72 related antigens may be prepared by methods previously described (WO2006/117240), those provided in the Examples, or methods analogous thereto.

[0052] The immunogenic compositions may comprise one or more further antigenic components. Such additional antigenic components need not themselves be sensitive to the presence of salts in the composition.

[0053] Additional antigenic components may be intended to strengthen or complement the immune responses solicited by the M72 related antigen in the field of tuberculosis prevention and therapy or additional antigens could be associated with other pathogens and are intended for administration with the M72 related antigen for reasons of convenience. Where a number of antigenic components are present within the formulation, these may be provided in the form of individual polypeptides or fusion proteins. In some circumstances additional antigenic components may be provided as a polynucleotide (or polynucleotides).

[0054] It is well known that for parenteral administration solutions should have a pharmaceutically acceptable osmolality to avoid cell distortion or lysis. A pharmaceutically acceptable osmolality will generally mean that solutions will have an osmolality which is approximately isotonic or mildly hypertonic. Suitably the immunogenic compositions of the present invention will have an osmolality in the range of 250 to 750 mOsm/kg, for example, the osmolality may be in the range of 250 to 550 mOsm/kg, such as in the range of 280 to 500 mOsm/kg.

[0055] Osmolality may be measured according to techniques known in the art, such as by the use of a commercially available osmometer, for example the Advanced® Model 2020 available from Advanced Instruments Inc. (USA).

[0056] An "isotonicity agent" is a compound that is physiologically tolerated and imparts a suitable tonicity to a formulation to prevent the net flow of water across cell membranes that are in contact with the formulation.

[0057] Generally, sodium chloride (NaCl) is used as a tonicity agent. The present inventors have shown for the first time that that M72 related antigens are particularly sensitive to "salting out", a process whereby the proteins in solution aggregate or coagulate when in solutions containing high concentrations of salt. Consequently, alternative means are provided for ensuring the immunogenic compositions of the invention have a pharmaceutically acceptable osmolality.

[0058] In a particular embodiment there are provided immunogenic compositions further comprising a non-ionic tonicity agent. A non-ionic tonicity agent for use in an immunogenic composition will itself need to be pharmaceutically acceptable, e.g. suitable for use in humans, as well as being compatible with the M72 related antigen and further compatible with other components such as the immunostimulant(s).

[0059] In one embodiment of the present invention, suitable non-ionic tonicity agents are polyols, sugars (in particular sucrose, fructose, dextrose or glucose) or amino acids such as glycine. In one embodiment the polyol is a sugar alcohol, especially a C3-6 sugar alcohol. Exemplary sugar alcohols include glycerol, erythritol, threitol, arabitol, xylitol, ribitol, sorbitol, mannitol, dulcitol and

iditol. In a specific example of this embodiment, a suitable non-ionic tonicity agent is sorbitol. The skilled person will recognise that an appropriate osmolality may be attained through the use of a mixture of different tonicity agents. In a particular embodiment of the invention the non-ionic tonicity agent in the compositions of the invention incorporates sucrose and/or sorbitol.

[0060] In one embodiment, a suitable concentration of polyol within the immunogenic composition is between about 2.5 and about 15% (w/v), in particular between about 2.5 and about 10% (w/v) for example between about 3 and about 7% (w/v), such as between about 4 and about 6% (w/v). In a specific example of this embodiment, the polyol is sorbitol.

[0061] In another embodiment, the immunogenic composition comprises sucrose and sorbitol. In such circumstances the immunogenic composition may suitably contain between about 2.5 and about 15% (w/v) of sucrose and between about 2.5 and about 15% (w/v) of sorbitol, in particular between about 2.5 and about 10% (w/v) of sucrose and between about 2.5 and about 10% (w/v) of sorbitol, for example, between about 3 and about 7% (w/v) of sucrose and between about 3 and about 7% (w/v) of sorbitol, such as between about 4 and about 6% (w/v) of sucrose and between about 4 and about 6% (w/v) of sorbitol.

[0062] The pH of the immunogenic compositions should be suitable for parenteral administration. Suitably the pH will be in the range 7.25 to 8.75, such as 7.5 to 8.5, in particular pH 7.75 to 8.25. A pH of about 8.0 is of particular interest.

[0063] The pH may be controlled by the use of buffers, including for example Tris or phosphate buffers.

[0064] In a particular embodiment of the invention, the immunogenic composition comprises one or more immunostimulants.

[0065] In one embodiment, the immunostimulant may be a saponin. A particularly suitable saponin for use in the present invention is Quil A and its derivatives. Quil A is a saponin preparation isolated from the South American tree *Quillaja saponaria* Molina and was first described by Dalsgaard et al. in 1974 ("Saponin adjuvants", Archiv. für die gesamte Virusforschung, Vol. 44, Springer Verlag, Berlin, p243-254) to have adjuvant activity. Purified fractions of Quil A have been isolated by HPLC which retain adjuvant activity without the toxicity associated with Quil A (WO88/09336), for example QS7 and QS21 (also known as QA7 and QA21). QS21 is a natural saponin derived from the bark of *Quillaja saponaria* Molina, which induces CD8⁺ cytotoxic T cells (CTLs), Th1 cells and a predominant IgG2a antibody response. QS21 is a preferred saponin in the context of the present invention.

[0066] In a suitable form of the present invention, the saponin adjuvant within the immunogenic composition is a derivative of *saponaria* Molina Quil A, in particular an immunologically active fraction of Quil A, such as QS17 or QS21, suitably QS21.

[0067] Desirably, QS21 is provided in a less reactogenic composition where it is quenched with an exogenous sterol, such as cholesterol for example. Several particular forms of less reactogenic compositions wherein QS21 is quenched with cholesterol exist. In a specific embodiment, the saponin/sterol is in the form of a liposome structure (such as described in WO96/33739, Example 1). In this embodiment the liposomes suitably contain a neutral lipid, for example phosphatidylcholine, which is suitably non-crystalline at room temperature, for example egg yolk phosphatidylcholine, dioleoyl phosphatidylcholine (DOPC) or dilauryl phosphatidylcholine. The liposomes may also contain a charged lipid which increases the stability of the liposome-QS21 structure for liposomes composed of saturated lipids. In these cases the amount of charged lipid is suitably 1-20% w/w, such as 5-10%. The ratio of sterol to phospholipid is 1-50% (mol/mol), suitably 20-25%.

[0068] Suitable sterols include β -sitosterol, stigmasterol, ergosterol, ergocalciferol and cholesterol. In one particular embodiment, the immunogenic composition comprises cholesterol as sterol. These sterols are well known in the art, for example cholesterol is disclosed in the Merck Index, 11th Edn., page 341, as a naturally occurring sterol found in animal fat.

[0069] Where the active saponin fraction is QS21, the ratio of QS21:sterol will typically be in the order of 1:100 to 1:1 (w/w), suitably between 1:10 to 1:1 (w/w), and especially 1:5 to 1:1 (w/w). Suitably excess sterol is present, the ratio of QS21:sterol being at least 1:2 (w/w). In one embodiment, the ratio of QS21:sterol is 1:5 (w/w). The sterol is suitably cholesterol.

[0070] In another embodiment, the immunogenic composition comprises an immunostimulant which is a Toll-like receptor 4 (TLR4) agonist. By "TLR agonist" it is meant a component which is capable of causing a signaling response through a TLR signaling pathway, either as a direct ligand or indirectly through generation of endogenous or exogenous ligand (Sabroe et al, J Immunol 2003 p1630-5). A TLR4 agonist is capable of causing a signaling response through a TLR-4 signaling pathway. A suitable example of a TLR4 agonist is a lipopolysaccharide, suitably a non-toxic derivative of lipid A, particularly monophosphoryl lipid A or more particularly 3-de-O-acylated monophosphoryl lipid A (3D-MPL).

[0071] 3D-MPL is sold under the name MPL by GlaxoSmithKline Biologicals N.A. and is referred throughout the document as

MPL or 3D-MPL see, for example, US Patent Nos. 4,436,727; 4,877,611; 4,866,034 and 4,912,094. 3D-MPL primarily promotes CD4+ T cell responses with an IFN-gamma (Th1) phenotype. 3D-MPL can be produced according to the methods disclosed in GB2220211A. Chemically it is a mixture of 3-de-O-acylated monophosphoryl lipid A with 3, 4, 5 or 6 acylated chains. In the compositions of the present invention small particle 3D-MPL may be used to prepare the immunogenic composition. Small particle 3D-MPL has a particle size such that it may be sterile-filtered through a 0.22 um filter. Such preparations are described in WO94/21292. Suitably, powdered 3D-MPL is used to prepare the immunogenic compositions of the present invention.

[0072] Other TLR4 agonists which can be used are alkyl glucosaminide phosphates (AGPs) such as those disclosed in WO98/50399 or US Patent No. 6,303,347 (processes for preparation of AGPs are also disclosed), suitably RC527 or RC529 or pharmaceutically acceptable salts of AGPs as disclosed in US Patent No. 6,764,840. Some AGPs are TLR4 agonists, and some are TLR4 antagonists.

[0073] Other suitable TLR4 agonists are as described in WO2003/011223 and in WO2003/099195, such as compound I, compound II and compound III disclosed on pages 4-5 of WO2003/011223 or on pages 3-4 of WO2003/099195 and in particular those compounds disclosed in WO2003/011223 as ER803022, ER803058, ER803732, ER804053, ER804057m ER804058, ER804059, ER804442, ER804680 and ER804764. For example, one suitable TLR-4 agonist is ER804057.

[0074] In a particular embodiment, the immunogenic composition comprises both a saponin and a TLR4 agonist. In a specific example, the immunogenic composition comprises QS21 and 3D-MPL.

[0075] A TLR-4 agonist, such as a lipopolysaccharide, such as 3D-MPL, can be used at amounts between 1 and 100 ug per human dose of the immunogenic composition. 3D-MPL may be used at a level of about 50 ug, for example between 40 to 60 ug, suitably between 45 to 55 ug or between 49 and 51 ug or 50 ug. In a further embodiment, the human dose of the immunogenic composition comprises 3D-MPL at a level of about 25 ug, for example between 20 to 30 ug, suitably between 21 to 29 ug or between 22 to 28 ug or between 23 and 27 ug or between 24 and 26 ug, or 25 ug.

[0076] A saponin, such as QS21, can be used at amounts between 1 and 100 ug per human dose of the immunogenic composition. QS21 may be used at a level of about 50 ug, for example between 40 to 60 ug, suitably between 45 to 55 ug or between 49 and 51 ug or 50 ug. In a further embodiment, the human dose of the immunogenic composition comprises QS21 at a level of about 25 ug, for example between 20 to 30 ug, suitably between 21 to 29 ug or between 22 to 28 ug or between 23 and 27 ug or between 24 and 26 ug, or 25 ug.

[0077] Where both TLR4 agonist and saponin are present in the immunogenic composition, then the weight ratio of TLR4 agonist to saponin is suitably between 1:5 to 5:1, suitably between 1:2 to 2:1, such as about 1:1. For example, where 3D-MPL is present at an amount of 50 ug or 25 ug, then suitably QS21 may also be present at an amount of 50 ug or 25 ug, respectively, per human dose of the immunogenic composition. Certain immunogenic compositions of the present invention comprise QS21 and 3D-MPL, at an amount of between 1 and 100 ug of each per human dose, such as at an amount of between 10 and 75 ug of each per human dose. Immunogenic compositions of the present invention may suitably comprise QS21 and 3D-MPL, at an amount of between 15 and 35 ug of each per human dose, such as at an amount of between 20 and 30 ug of each per human dose.

[0078] In one embodiment, the immunostimulant is a TLR9 agonist, for example as set out in WO2008/142133. In a specific example, said TLR9 agonist is an immunostimulatory oligonucleotide, in particular an oligonucleotide containing an unmethylated CpG motif. Such oligonucleotides are well known and are described, for example, in WO96/02555, WO99/33488 and US5,865,462. Suitable TLR9 agonists for use in the immunogenic compositions described herein are CpG containing oligonucleotides, optionally containing two or more dinucleotide CpG motifs separated by at least three, suitably at least six or more nucleotides. A CpG motif is a cytosine nucleotide followed by a guanine nucleotide.

[0079] In one embodiment the internucleotide bond in the oligonucleotide is phosphorodithioate, or possibly a phosphorothioate bond, although phosphodiester and other internucleotide bonds could also be used, including oligonucleotides with mixed internucleotide linkages. Methods for producing phosphorothioate oligonucleotides or phosphorodithioate are described in US5,666,153, US5,278,302 and WO95/26204. Oligonucleotides comprising different internucleotide linkages are contemplated, e.g. mixed phosphorothioate phosphodiesters. Other internucleotide bonds which stabilise the oligonucleotide may be used.

[0080] Examples of CpG oligonucleotides suitable for inclusion in the immunogenic compositions described herein have the following sequences. In one embodiment, these sequences contain phosphorothioate modified internucleotide linkages.

OLIGO 1 (SEQ ID No: 9): TCC ATG ACG TTC CTG ACG TT (CpG 1826)

OLIGO 2 (SEQ ID No: 10): TCT CCC AGC GTG CGC CAT (CpG 1758)

OLIGO 3 (SEQ ID No: 11): ACC GAT GAC GTC GCC GGT GAC GGC ACC ACG

OLIGO 4 (SEQ ID No: 12): TCG TCG TTT TGT CGT TTT GTC GTT (CpG 2006)

OLIGO 5 (SEQ ID No: 13): TCC ATG ACG TTC CTG ATG CT (CpG 1668)

[0081] Alternative CpG oligonucleotides may comprise the sequences above in that they have inconsequential deletions or additions thereto.

[0082] In one embodiment the immunostimulant is a tocol. Tocols are well known in the art and are described in EP0382271. In a particular embodiment, the tocol is alpha-tocopherol or a derivative thereof such as alpha-tocopherol succinate (also known as vitamin E succinate).

[0083] The immunogenic compositions according to the invention may be used in medicine, in particular for the prophylaxis, treatment or amelioration of infection by mycobacteria, such as infection by *Mycobacterium tuberculosis*. The immunogenic compositions will generally be provided for administration to humans, though they may also be of value in veterinary medicine such as for administration to bovines.

[0084] There is provided the use of an immunogenic composition according to the invention in the manufacture of a medicament, in particular a medicament for the prophylaxis, treatment or amelioration of infection by mycobacteria, such as infection by *Mycobacterium tuberculosis*.

[0085] The immunogenic composition may be provided for the purpose of:

- treating active tuberculosis;
- prophylaxis of active tuberculosis, such as by administering to a subject who is uninfected, or alternatively a subject who has latent infection;
- treating latent tuberculosis;
- prophylaxis of latent tuberculosis, such as by administering to a subject who is uninfected; or
- preventing or delaying reactivation of tuberculosis, especially the delay of TB reactivation, for example by a period of months, years or even indefinitely.

[0086] The term "active infection" refers to an infection, e.g. infection by *M. tuberculosis*, with manifested disease symptoms and/or lesions, suitably with manifested disease symptoms.

[0087] The terms "inactive infection", "dormant infection" or "latent infection" refer to an infection, e.g. infection by *M. tuberculosis*, without manifested disease symptoms and/or lesions, suitably without manifested disease symptoms. A subject with latent infection will suitably be one which tests positive for infection, e.g. by PPD or T cell based assays, but which has not demonstrated the disease symptoms and/or lesions which are associated with an active infection.

[0088] The term "primary tuberculosis" refers to clinical illness, e.g., manifestation of disease symptoms, directly following infection, e.g. infection by *M. tuberculosis*. See, Harrison's Principles of Internal Medicine, Chapter 150, pp. 953-966 (16th ed., Braunwald, et al., eds., 2005).

[0089] The terms "secondary tuberculosis" or "postprimary tuberculosis" refer to the reactivation of a dormant, inactive or latent infection, e.g. infection by *M. tuberculosis*. See, Harrison's Principles of Internal Medicine, Chapter 150, pp. 953-966 (16th ed., Braunwald, et al., eds., 2005).

[0090] The term "tuberculosis reactivation" refers to the later manifestation of disease symptoms in an individual that tests positive for infection (e.g. in a tuberculin skin test, suitably in an *in vitro* T cell based assay) test but does not have apparent disease symptoms. Suitably the individual will not have been re-exposed to infection. The positive diagnostic test indicates that the individual is infected, however, the individual may or may not have previously manifested active disease symptoms that had been treated sufficiently to bring the tuberculosis into an inactive or latent state.

[0091] Suitability an immunogenic composition is administered to a subject who is uninfected or who has a latent infection by mycobacteria, such as infection by *Mycobacterium tuberculosis*.

[0092] The volume of immunogenic composition administered may vary depending upon a number of other factors, such as the specific delivery route, e.g. intramuscular, subcutaneous or intradermal. Typically, the volume administered in a single injection (the unit dose) for a human will be in the range of 50 ul to 1 ml, such as 100 ul to 750 ul, especially 400 to 600 ul, for example about 500 ul.

[0093] The quantity of M72 related antigen contained within a single dose is dependent upon clinical needs but a single human dose will typically be in the range of 1 to 100 ug, such as 5 to 50 ug, for example 5 to 20 ug. A single human dose may contain about 10 ug of M72 related antigen.

[0094] Suitably, compositions of the invention will be stable, in which is meant that during storage at 25 °C for a period of 24 hours antigenicity as measured by the techniques described herein remains at least 80% of the antigenicity before storage. Desirably, antigenicity will remain at least 85%, such as at least 90% and in particular at least 95% after storage at 25 °C for a period of 24 hours. For compositions of particular interest, at least 80% of the antigenicity of the composition, such as at least 85%, at least 90% and especially at least 95% remains after storage at 30 °C for a period of 24 hours.

[0095] The present invention will now be further described by means of the following non-limiting examples.

EXAMPLES

Example 1: Preparation of adjuvant composition ASA (sorbitol)

[0096] An adjuvant composition was prepared which comprised 3-de-O-acylated monophosphoryl lipid A and QS21 in a liposomal formulation using sorbitol as a tonicity agent. This was prepared as follows:

A. Method of preparation of liposomes:

[0097] A mixture of lipid (DOPC), cholesterol and 3-de-O-acylated monophosphoryl lipid A in organic solvent was dried down under vacuum. An aqueous solution (phosphate buffered saline [100 mM NaCl, 50 mM Phosphate pH 6.1]) was then added and the vessel agitated until all the lipid was in suspension. This suspension was then prehomogenised with high shear mixer and then high pressure homogenised until the liposome size was reduced to around 90 nm ± 10 nm measured by DLS. Liposomes were then sterile filtered.

B. ASA formulation:

Step 1: Dilution of concentrated liposomes

[0098] Na₂/K Phosphate buffer 100 mM pH 6.1 when diluted 10 times was added to water for injection to reach a 10 mM phosphate buffer concentration in the final formulation. A 30% (w/v) sorbitol solution in water for injection (WFI) was then added to reach a concentration of 4.7% in the final formulation - this was stirred for 15 to 45 minutes at room temperature.

[0099] Concentrated liposomes (made of DOPC, cholesterol and 3D-MPL at 40 mg/ml, 10 mg/ml and 2 mg/ml respectively) were then added to the mix to reach a concentration of 100 ug/ml of 3D-MPL in the final formulation.

[0100] The mixture was subsequently stirred for 15 to 45 minutes at room temperature.

Step 2: QS21 addition

[0101] Using a peristaltic pump, QS21 bulk stock was added to the diluted liposomes under magnetic stirring to reach a 100 ug/ml concentration in the final formulation. The mix was stirred for 15 to 45 minutes.

[0102] Final ASA(sorbitol) formulation contained 2 mg DOPC, 500 ug cholesterol, 100 ug 3D-MPL/ml and 100 ug QS21/ml, 4.7% sorbitol and 5 mM sodium chloride and 10 mM phosphate.

Step 3: pH was checked to be 6.1±0.1

Step 4: Sterile filtration

[0103] Sterile filtration was performed using a polyethersulfone (PES) filter from PALL Corporation.

Step 5: Storage at +2°C to +8°C.

[0104] The adjuvant composition obtained, which comprised 3-de-O-acylated MPL and QS21 in a liposomal formulation and containing sorbitol as a tonicity agent (designated ASA (sorbitol)), was then stored at 4°C.

Example 2: Preparation of adjuvant composition ASA (150 mM NaCl)

[0105] An adjuvant composition was prepared which comprised 3-de-O-acylated monophosphoryl lipid A and QS21 in a liposomal formulation using sodium chloride as a tonicity agent.

A. Method of preparation of liposomes:

[0106] A mixture of lipid (DOPC), cholesterol and 3-de-O-acylated monophosphoryl lipid A (3D-MPL) in organic solvent was dried down under vacuum. Phosphate buffered saline (100 mM NaCl, 50 mM Phosphate pH 6.1) was then added and the vessel agitated until all the lipid was in suspension. This suspension was then prehomogenised with high shear mixer and then high pressure homogenised until the liposomes size was reduced to around 90 nm ± 10 nm measured by DLS. Liposomes were then sterile filtered on 0.22 um PES membrane.

B. ASA formulation:

Step 1: Dilution of concentrated liposomes

[0107] Na₂/K Phosphate buffer 100 mM pH 6.45 when diluted 10 times and NaCl 1.5 M were added to water for injection to reach respectively 10 mM phosphate and NaCl 150 mM concentrations in the final formulation. This mixture was stirred for 5 minutes at room temperature. Concentrated liposomes (made of DOPC, cholesterol and 3D-MPL at 40 mg/ml, 10 mg/ml and 2 mg/ml respectively) were then added to the mix to reach a concentration of 100 ug/ml of 3D-MPL in the final formulation. The mixture was subsequently stirred for 5 to 15 minutes at room temperature.

Step 2: QS21 addition

[0108] QS21 bulk stock was added to the diluted liposomes under magnetic stirring to reach a 100 ug/ml concentration in the final formulation. The mix was stirred at room temperature.

Step 3: pH was checked so as to be 6.1±0.1.

Step 4: Sterile filtration

[0109] Sterile filtration was performed using a polyethersulfone (PES) filter from PALL Corporation.

Step 5: Storage at +2°C to +8°C

[0110] Final composition of ASA(150 mM NaCl) was 2 mg DOPC, 500 ug cholesterol, 100 ug 3-de-O-acylated MPL, 100 ug QS21 per 1 ml, with 10 mM phosphate and 150 mM NaCl.

Example 3: QS21 Lytic Activity

[0111] QS21 is known to lyse red blood cells (RBC). The ASA (sorbitol) adjuvant composition prepared as in Example 1 was tested to ensure that QS21 lytic activity was quenched in the same way as was seen with the equivalent adjuvant composition comprising 150 mM NaCl (ASA (150 mM NaCl)).

[0112] QS21 lytic activity was measured by a haemolysis assay using chicken Red Blood cells (RBC). RBC were centrifuged at 550 g at 4°C. Supernatant was discarded. The pellet was carefully resuspended in PBS buffer to reach the initial volume and the same operation was repeated until supernatant was no longer red (generally 3 times). The pellet was stored at 4°C for 3 to 4 days maximum if not used directly (and washed again the day it is used) or was diluted around 10 times in buffer if used the same day.

[0113] A QS21 dose range curve was prepared in ASA buffer (in salt or in sorbitol buffer following the ASA sample tested) extemporaneously and the adjuvant samples (containing a 50 ug or 90 ug equivalent of QS21 meaning the equivalent of 500 ul or 900 ul ASA) were prepared. Final volume was adjusted to 900 ul in standards and samples with adequate buffer (containing or not sorbitol as a function of the buffer of the sample tested). Due to its opalescence, ASA interferes with optical density (OD). ASA "blanks" were thus prepared and their OD was subtracted from the OD of ASA tested samples. Those blanks corresponded to the same ASA volume as the volume tested in samples, but adjusted to 1 ml with buffer. No RBC were added to these blanks. Standards and samples were then incubated with RBC (100 ul of diluted RBC added to 900 ul of standards and samples) for 30 minutes at room temperature (RT). Samples were then centrifuged 5 minutes at 900 g. Optical density at 540 nm was measured after centrifugation.

[0114] Determination of lytic activity was carried out by a limit test.

1. Limit of detection (LOD) was defined as the lowest concentration of QS21 leading to an OD:
 - Higher than the base level (OD>0.1)
 - Around three times higher than OD's buffer (the "0 ug" QS21)
 - In the ascendant part of the curve
 - Determined for each test.
2. QS21 lytic activity was held to be positive in the adjuvant samples if the OD for the adjuvant sample was greater than the OD_{LOD}.

Example QS21 curve

ug QS21	OD	QS21 quenched
0	0.029	NA
0.5	0.052	< LOD
0.6	0.073	< LOD
0.7	0.091	< LOD
0.8	0.096	< LOD
0.9	0.12	> 98.2%
1	0.195	> 98%

ug QS21	OD	QS21 quenched
1.1	0.212	> 97.8%
1.2	0.348	> 97.6%
1.3	0.479	> 97.4%
1.4	0.612	> 97.2%
1.5	0.669	> 97%
2	1.139	> 96%
2.5	1.294	> 95%
3	1.391	> 94%
5	1.416	> 90%
Adjuvant *	0.03	> 98.2 %
*50ug QS21 equivalent tested. 150 mM sodium chloride buffer.		

[0115] The above data is shown graphically in Figure 1.

[0116] The Limit of Detection in this assay is at 0.9 ug QS21, and OD of 0.12

[0117] The QS21 quenching in an adjuvant composition comprising 150 mM sodium chloride was estimated to be more than 98.2% for the equivalent of 50 ug QS21 tested. In the case of an equivalent of 90 ug tested, conclusion is more than 99%.

[0118] QS21 quenching was then compared with an equivalent adjuvant composition comprising sorbitol and only 5 mM sodium chloride. Data were generated after storage of the ASA at 4°C or after accelerated stability (7 days at 37°C). For the ASA in sorbitol, the QS21 standard curve was realised in a sorbitol containing buffer.

Sample	Timepoint	LOD	QS21 quenched
Adjuvant composition (ASA) 150 mM NaCl	T0	< 1.4	> 97.2%
	7 days 37°C	< 0.9	> 98.2%
Adjuvant Composition (ASA) sorbitol, 5 mM NaCl	T0	< 2	> 97.8%
	7 days 37°C	<1	> 96%
	11 months 4°C	<2	> 97.8%*
Equivalent of 50 ug QS21 tested except * equivalent of 90 ug QS21 tested.			

[0119] It was concluded that QS21 was adequately quenched in a low sodium chloride buffer.

Example 4: MPL congeners.

[0120] Chemically, 3D-MPL is a mixture of 3-de-O-acylated monophosphoryl lipid A with mainly 4, 5 or 6 acylated chains. Each separate 3D-MPL molecule is called a congener. It is important that the congener composition remain constant, with no shift between the proportion of congeners. It is also important that any buffer used enables the congener composition to be the same as in the concentrated liposomes used to make the adjuvant composition.

[0121] As shown on Figure 2, the congener composition was examined in 3D-MPL concentrated liposomes (Conc. Liposomes LIP07-217, first column of Figure 2), an adjuvant composition comprising 3D-MPL liposomes and QS21 in a 150 mM NaCl buffer (Adjuvant 150 mM NaCl, or ASA (150 mM NaCl), second column), and an adjuvant composition comprising 3D-MPL liposomes and QS21 in a sorbitol and 5 mM NaCl buffer (Adjuvant Sorbitol, or ASA (sorbitol), columns 3-7).

[0122] The congener composition was also examined in two lots of ASA (sorbitol) adjuvant at day 0 and 7 days after preparation and maintenance at 37 °C to ensure that there was no evolution over time (see final four columns of Figure 2).

[0123] Relative distribution of tetra-, penta- and hexa-acylated congeners of MPL in concentrated liposomes or ASA (sorbitol) samples was determined by IP-HPLC-Fluo detection (ARD). Both standards and samples were derivatised with dansylhydrazine, which introduces a Fluo-active chromophore on the dissacharide backbone. The derivatised samples were analysed on a C18 reverse phase column using tetrabutylammonium hydroxide (TBAOH) as an ion pair reagent. Congeners containing the same numbers of fatty acyl groups were eluted in distinct groups (tetraacyl, pentaacyl, and hexaacyl). Distribution of congeners is deduced by comparing the peak area of each group to the total peak area of all MPL congeners.

[0124] Figure 2 shows the percentage of each congener. No significant difference in congener composition was found between adjuvant buffers, and the congener composition was consistent over time in the sorbitol buffer.

Example 5: Preparation of adjuvant composition ASA (sorbitol - 2)

[0125] An adjuvant composition was prepared which comprised 3-de-O-acylated monophosphoryl lipid A and QS21, at a reduced level relative to Example 1, in a liposomal formulation using sorbitol as a tonicity agent. This was prepared as follows:

The adjuvant was prepared by 1:1 dilution of ASA(sorbitol), prepared according to Example 1, with a solution containing 10 mM phosphate, 5 mM NaCl, 4.7% sorbitol at pH 6.1.

Final ASA(sorbitol - 2) formulation contained 1 mg DOPC, 250 ug cholesterol, 50 ug 3D-MPL/ml and 50 ug QS21/ml, 4.7% sorbitol, 5 mM sodium chloride and 10 mM phosphate.

Example 6: Preparation of adjuvant composition ASA (sorbitol - 3)

[0126] An adjuvant composition was prepared which comprised 3-de-O-acylated monophosphoryl lipid A and QS21, at a reduced level relative to Example 1, in a liposomal formulation using sorbitol as a tonicity agent. This was prepared as follows:

A. Method of preparation of liposomes:

[0127] A mixture of lipid (DOPC), cholesterol and 3-de-O-acylated monophosphoryl lipid A in organic solvent was dried down under vacuum. An aqueous solution (phosphate buffered saline [100 mM NaCl, 50 mM Phosphate pH 6.1]) was then added and the vessel agitated until all the lipid was in suspension. This suspension was then prehomogenised with high shear mixer and then high pressure homogenised until the liposomes size was reduced to around 90 nm $\pm$ 10 nm measured by DLS. Liposomes were then sterile filtered.

B. ASA formulation:

Step 1: Dilution of concentrated liposomes

[0128] Na₂/K Phosphate buffer 100 mM pH 6.1 when diluted 10 times was added to water for injection to reach a 10 mM phosphate buffer concentration in the final formulation. A 30% (w/v) sorbitol solution in water for injection (WFI) was then added to reach a concentration of 4.7% in the final formulation - this was stirred for 15 to 45 minutes at room temperature.

[0129] Concentrated liposomes (made of DOPC, cholesterol and 3D-MPL at 40 mg/ml, 10 mg/ml and 2 mg/ml respectively) were then added to the mix to reach a concentration of 50 ug/ml of 3D-MPL in the final formulation.

[0130] The mixture was subsequently stirred for 15 to 45 minutes at room temperature.

Step 2: QS21 addition

[0131] Using a peristaltic pump, QS21 bulk stock was added to the diluted liposomes under magnetic stirring to reach a 50 ug/ml concentration in the final formulation. The mix was stirred for 15 minutes.

[0132] Final ASA formulation contained 1 mg DOPC, 250 ug cholesterol, 50 ug 3D-MPL/ml and 50 ug QS21/ml, 4.7% sorbitol and 2.5 mM sodium chloride, 10 mM phosphate.

Step 3: pH was checked to be 6.1±0.1

Step 4: Sterile filtration

[0133] Sterile filtration was performed using a polyethersulfone (PES) filter from PALL Corporation.

Step 5: Storage at +2°C to +8°C.

[0134] The adjuvant composition obtained, which comprised 3-de-O-acylated MPL and QS21 in a liposomal formulation and containing sorbitol as a tonicity agent (designated ASA (sorbitol-3)), was then stored at 4°C.

Example 7: Preparation of adjuvant composition ASA (150 mM NaCl - 2)

[0135] An adjuvant composition was prepared which comprised 3-de-O-acylated monophosphoryl lipid A and QS21, at a reduced level relative to Example 2, in a liposomal formulation using sodium chloride as a tonicity agent. This was prepared as follows:

A. Method of preparation of liposomes:

[0136] A mixture of lipid (DOPC), cholesterol and 3-de-O-acylated monophosphoryl lipid A (3D-MPL) in organic solvent was dried down under vacuum. Phosphate buffered saline (100 mM NaCl, 50 mM Phosphate pH 6.1) was then added and the vessel agitated until all the lipid was in suspension. This suspension was then prehomogenised with high shear mixer and then high pressure homogenised until the liposomes size was reduced to around 90 nm ± 10 nm measured by DLS. Liposomes were then sterile filtered on 0.22 um PES membrane.

B. ASA formulation:

Step 1: Dilution of concentrated liposomes

[0137] Na₂/K Phosphate buffer 100mM pH 6.45 when diluted 10 times and NaCl 1.5 M were added to water for injection to reach respectively 10 mM phosphate and NaCl 150 mM concentrations in the final formulation. This mixture was stirred for 5 minutes at room temperature. Concentrated liposomes (made of DOPC, cholesterol and 3D-MPL at 40 mg/ml, 10 mg/ml and 2 mg/ml respectively) were then added to the mix to reach a concentration of 50 ug/ml of 3D-MPL in the final formulation. The mixture was subsequently stirred for 5 to 15 minutes at room temperature.

Step 2: QS21 addition

[0138] QS21 bulk stock was added to the diluted liposomes under magnetic stirring to reach a 50 ug/ml concentration in the final formulation. The mix was stirred at room temperature.

Step 3: pH was checked so as to be 6.1±0.1.

Step 4: Sterile filtration

[0139] Sterile filtration was performed using a polyethersulfone (PES) filter from PALL Corporation.

Step 5: Storage at +2°C to +8°C

[0140] Final composition of ASA(150 mM NaCl - 2) was 1 mg DOPC, 250 ug cholesterol, 50 ug 3-de-O-acylated MPL, 50 ug QS21 per 1 ml, 10 mM phosphate and 150 mM NaCl.

Example 8: Preparation of Protein Antigens**M72 with two N-terminal His residues (SEQ ID No: 3)****Construction of the M72 expression Vector**

[0141] A plasmid coding for the amino acid sequence of Mtb72f with an additional 6-His tag at the N-terminus was generated by the sequential linkage in tandem of the open reading frames (ORFs) encoding the C terminal fragment of Mtb32a to the full length ORF of Mtb39a followed at the C terminus with the N terminal portion of Mtb32a. This was accomplished by using sequence-specific oligonucleotides containing unique restriction sites (EcoRI and EcoRV) and devoid of the stop codons at the C terminal ends (in the case of the C terminal fragment of Mtb32a and Mtb39a) for polymerase chain reaction (PCR) of genomic DNA from the *M. tuberculosis* strain H37Rv. Using this vector as template, a mutation of Ser706 to Ala was performed by site-directed mutagenesis. The proper orientation of inserts as well as the mutation Ser706Ala was verified by DNA sequencing.

[0142] In order to obtain the vector coding for M72, which just has 2 His residues at the N terminus, four His were deleted making use of a commercial site-directed mutagenesis system. After sequence verification, the M72 coding sequence was excised from the plasmid by enzymatic reaction, gel purified and ligated into a pET vector. The recombinant plasmid was then sequence verified. This plasmid codes for M72 under the control of a T7 promoter. Expression of T7 RNA polymerase is driven from a genomic integrant in the expression host and is induced using a lac operon-based system (lacI) and an IPTG chemical induction signal. The expression plasmid is provided with kanamycin resistance.

[0143] The plasmid coding for the M72 fusion protein under the control of a T7 promoter was transformed into the HMS174 (DE3) strain of *E.coli*, using an electroporation method. The coding sequence of the M72 insert and the flanking regions were sequenced on both strands and were found to be identical to the sequence determined from the original plasmid construct.

Fermentation

[0144] A vial of pelleted working seed was thawed at room temperature. A pre-dilution was prepared by mixing the working seed with 4.9 ml of pre-culture medium. 1 ml of the pre-dilution was used to inoculate the liquid pre-culture which consists of 400 ml of pre-culture medium supplemented with 50 mg/l kanamycin sulfate and 10 g/l glucose.

Pre-culture medium composition

Ingredient	Concentration		
KH ₂ PO ₄	14.83 g/l		
K ₂ HPO ₄	1.65 g/l		
(NH ₄) ₂ SO ₄	5.82 g/l		
Yeast extract	6.21 g/l		

Ingredient	Concentration		
Glycerol 87% (w/w)	14.54 ml/l		
Metal and salt solution (1):	9.7 ml/l		
-FeCl ₃ 6H ₂ O		3.3 g/l	
- MgSO ₄ 7H ₂ O		58 g/l	
- Micro element solution (2):		116 ml/l	
- ZnSO ₄ 7H ₂ O			7.65 g/l
- MnSO ₄ H ₂ O			5.28 g/l
- CuSO ₄ 5H ₂ O			1.1 g/l
- CoCl ₂ 6H ₂ O			1.1 g/l
- H ₃ BO ₃			0.3 g/l
- Na ₂ MoO ₄ 2H ₂ O			2.64 g/l
- HCl 4N			6.2 ml/l
Biotine and CaCl ₂ solution (2):	0.97 ml/l		
- Biotine		0.05 g/l	
- CaCl ₂ 2 H ₂ O		61.7 g/l	
pH of the medium is adjusted to 6.5 with NaOH (25%) solution			
The medium is filtered through 0.22 um			
(1) pH adjusted to 1.50 with HCl (37%) solution; the solution is filtered through 0.22 um			
(2) The solution is filtered through 0.22 um			

[0145] The pre-culture was incubated in a 2 litre shake flask at 30°C under agitation (200 RPM) until the OD_{650nm} reached a value between 2 and 4 (approximate incubation time: 16 hours). At that stage, a 72 litre (total volume) fermenter containing 45 litres of culture medium supplemented with 34 mg/l kanamycin sulfate was inoculated with 52 ml liquid pre-culture.

Culture medium composition

Ingredient	Concentration	
MgSO ₄ 7H ₂ O	0.63 g/l	
FeCl ₃ 6H ₂ O	0.056 g/l	
Micro element solution (1):	1.91 ml/l	
ZnSO ₄ 7H ₂ O		7.65 g/l
- MnSO ₄ H ₂ O		5.28 g/l
- CuSO ₄ 5H ₂ O		1.1 g/l
- CoCl ₂ 6H ₂ O		1.1 g/l
- H ₃ BO ₃		0.3 g/l
- Na ₂ MoO ₄ 2H ₂ O		2.64 g/l
- HCl 4N		6.2 ml/l
HCl 37%	0.40 mL/L	
Yeast extract	35 g/L	
(NH ₄) ₂ SO ₄	2.10 g/l	
KH ₂ PO ₄	18.70 g/l	
Sodium glutamate	2.5 g/l	
Glycerol 87%	0.276 ml/l	

Ingredient	Concentration	
Glucose	20 g/l	
Biotine solution (2):	0.22 ml/l	
- Biotine		1 g/l
CaCl ₂ 2 H ₂ O	0.21 g/l	
The solution is filtered through 0.22 µm		
(1) The solution is filtered through 0.22 µm		
(2) pH adjusted to 11.0 with NaOH (25%) solution; the solution is filtered through 0.22 µm		

[0146] During the growth phase, pH was maintained at 6.8 ± 0.2 by periodic addition of 25% (v/v) NH₄OH and 25% (v/v) H₃PO₄. After incubation for 16 hours at 30°C, fed-batch was started with feed medium.

Feed medium composition

Ingredient	Concentration	
MgSO ₄ 7H ₂ O	1.98 g/l	

Ingredient	Concentration	
FeCl ₃ 6H ₂ O	0.178 g/l	
Micro element solution (1):	6.02 ml/l	
- ZnSO ₄ 7H ₂ O		7.65 g/l
- MnSO ₄ H ₂ O		5.28 g/l
- CuSO ₄ 5H ₂ O		1.1 g/l
- CoCl ₂ 6H ₂ O		1.1 g/l
- H ₃ BO ₃		0.3 g/l
- Na ₂ MoO ₄ 2H ₂ O		2.64 g/l
- HCl 4N		6.2 ml/l
HCl 37%	1.24 ml/l	
Sodium glutamate	5 g/l	
Yeast extract	40 g/l	
Glycerol 87%	590 ml/l	
Biotine solution (2):	2 ml/l	
- Biotine		1 g/l
CaCl ₂ 2 H ₂ O	0.66 g/l	
The solution is filtered through 0.22 µm		
(1) The solution is filtered through 0.22 µm		
(2) pH adjusted to 11.0 with NaOH (25%) solution; the solution is filtered through 0.22 µm		

[0147] The temperature was maintained at 30°C for a further 2 hours, then raised to 37°C until the end of fermentation. The air flow was constantly set to 75 l/min and the dissolved oxygen kept at 17% saturation by feedback control of the agitation and pressure. Small quantities of antifoam solution were added on demand automatically. By the time the OD_{650nm} reached a value of 50 (± 5), 1 mM Isopropyl-beta-D-thiogalactopyranoside (IPTG) was added in order to induce the expression of M72. Fermentation ended after 5 hours from the time point induction was started. The cell culture was cooled down to 15°C under slight agitation and centrifuged (at 4°C) to obtain cell pellets which were thereafter stored at -20°C in aliquots.

Isolation of Inclusion Bodies

[0148] The cell pellets collected from the harvest were thawed at room temperature and disrupted in lysis buffer (10 mM Tris, 50 mM NaCl, pH 8.0) with a high pressure homogenizer. Thereafter the cell lysate was centrifuged and the resulting cell pellets (or inclusion bodies, IBs) were washed with wash buffer containing urea, Tris and NaCl. The IBs were solubilised with solubilisation buffer containing 8 M urea and filtered through a 0.2 µm membrane. This filtered solution was first purified by anion exchange chromatography using a Q Sepharose Fast Flow (QSFF) column. The elution of M72 takes place with a 6 M urea, 20 mM bis-Tris propane, 90 mM NaCl, pH 7.0 solution.

[0149] M72 collected was further purified by Hydroxyapatite chromatography (HA), from which it is eluted with a 6 M urea, 20 mM bis-Tris propane, 250 mM NaCl, pH 7.0 solution. The collected fraction was concentrated with a 30 kDa membrane cassette and diafiltered against 20 mM Tris, pH 7.5. M72 was then sterilised through a 0.22 µm filter. The purified bulk was then aliquoted and stored at -70°C.

Example 9: Investigation of "salting out" in compositions comprising M72 with different salt concentrations at pH 6.1, 7.5 and 8.5.

[0150] The impact of sodium chloride concentration and pH on M72 antigen stability, as assessed by size and antigenicity, was investigated.

Method

[0151] Purified bulk antigen (M72 with two N-terminal His residues, SEQ ID No: 3, as prepared in Example 8) was diluted to a concentration of 100 µg/ml in three different buffers (10mM phosphate buffer at pH 6.1, 20 mM Tris buffer at pH 7.5 and 20 mM Tris buffer at pH 8.5) containing final sodium chloride concentrations of 0, 50, 150, 300 and 450 mM.

[0152] Samples were analysed immediately (TO), stored overnight at 4°C before analysis (TO O/N) or stored at 25°C for 24 hours before analysis (T24h25°C).

[0153] DLS was performed using a Malvern Zetasizer Nano ZS from Malvern Instruments (UK). The instrument was operated using a laser wavelength of 633 nm and power of 4 mW. Scattered light was detected at 173° at a temperature of 22°C. The Z-average diameter (Zav) and polydispersity index (pl) are calculated by the instrument software.

[0154] Nephelometry was performed using a Nepheloskan® Ascent, available from Thermo Fischer Scientific. Analysis was performed in UV transparent Costar® micro-plates available from Corning Inc (USA).

[0155] Antigenicity was quantified by a sandwich ELISA in which the antigen is captured by a M72-specific rabbit polyclonal antibody and subsequently revealed by a M72(Mtb39)-specific mouse monoclonal antibody. All measured values are presented relative to the expected antigenicity based on the purified bulk protein used to prepare the tested formulations.

Results

[0156] The findings of this experiment are presented in Figures 3 to 5.

[0157] The results demonstrate for the first time that the stability of solutions containing an M72 related antigen is sensitive to both pH and sodium chloride concentration. The impact of sodium chloride on antigen size and antigenicity is all the more notable as the pH is lower.

[0158] Antigen size and antigenicity are not stable at pH 6.1 even in the absence of sodium chloride. The addition of 50 mM sodium chloride at pH 6.1 led to a size increase from 35 nm (0 mM sodium chloride at TO) up to 58 nm (TO) or 79 nm after 24 hours at 25°C.

[0159] Antigen size and antigenicity are relatively stable over 24 hours at 25°C at pH 7.5 or 8.5, particularly in the absence of sodium chloride or at a sodium chloride concentration of 50 mM. Nevertheless, increasing the concentration of sodium chloride to 150 mM or greater results in a clear increase in antigen size and reduction in antigenicity.

Example 10: Prevention of "salting out" in compositions comprising M72, immunostimulants and using sorbitol as a tonicity agent

[0160] In order to compare the stability of immunogenic compositions containing 150 mM NaCl with compositions using sorbitol as a tonicity agent, a number of samples were monitored using SEC-HPLC and ELISA.

Method

[0161] Three different lyophilisation cakes were prepared, such that when combined with the appropriate adjuvant formulations from Examples 5 and 7 the desired pH would be obtained:

(a) M72 with two N-terminal His residues - target pH 8.5 in reconstituted vaccine

15.75% (w/v) sucrose solution (prepared in water for injection) was added to water for injection to reach a sucrose concentration of 6.3%. 3% (w/v) Tween80 solution (prepared in water for injection) was then added to reach a concentration of 0.025%. Tris-HCl buffer 1 M pH 8.8 was then added to reach a 50 mM Tris buffer concentration. The mixture was magnetically stirred for 5 minutes at room temperature. Purified bulk antigen (M72 with two N-terminal His residues, SEQ ID No: 3, as prepared in Example 8) was then added to reach a protein concentration of 25 ug/ml. The mixture was magnetically stirred for 10 minutes at room temperature. The pH was checked and found to be 8.8.

0.5 ml of the mixture obtained was filled in 3 ml glass vials then freeze dried.

(b) M72 with two N-terminal His residues - target pH 8.0 in reconstituted vaccine

15.75% (w/v) sucrose solution (prepared in water for injection) was added to water for injection to reach a sucrose concentration of 6.3%. 3% (w/v) Tween80 solution (prepared in water for injection) was then added to reach a concentration of 0.025%. Tris-HCl buffer 1 M pH 8.8 was then added to reach a 20 mM Tris buffer concentration. The mixture was magnetically stirred for 5 minutes at room temperature. Purified bulk antigen (M72 with two N-terminal His residues, SEQ ID No: 3, as prepared in Example 8) was then added to reach a protein concentration of 25 ug/ml. The mixture was magnetically stirred for 10 minutes at room temperature. The pH was checked and found to be 8.8.

0.5 ml of the mixture obtained was filled in 3 ml glass vials then freeze dried.

(c) M72 with two N-terminal His residues - target pH 7.5 in reconstituted vaccine

15.75% (w/v) sucrose solution (prepared in water for injection) was added to water for injection to reach a sucrose concentration of 6.3%. 3% (w/v) Tween80 solution (prepared in water for injection) was then added to reach a concentration of 0.025%. Tris-HCl buffer 1 M pH 8.8 was then added to reach a 12.5 mM Tris buffer concentration. The mixture was magnetically stirred for 5 minutes at room temperature. Purified bulk antigen (M72 with two N-terminal His residues, SEQ ID No: 3, as prepared in Example 8) was then added to reach a protein concentration of 25 ug/ml. The mixture was magnetically stirred for 10 minutes at room temperature. The pH was checked and found to be 8.8.

0.5 ml of the mixture obtained was filled in 3 ml glass vials then freeze dried.

[0162] The lyophilisation cakes described above were reconstituted with 625 ul of the adjuvant solutions prepared in Examples 5 and 7. Upon reconstitution with adjuvant solution, the following immunogenic compositions were obtained:

1. (i) M72 with two N-terminal His residues - ASA(150 mM NaCl - 2) pH 8.5

10 ug antigen (20 ug/ml)

5% w/v sucrose

40 mM Tris

0.02% w/v Tween80

500 ug DOPC

125 ug cholesterol

25 ug 3D-MPL

25 ug QS21

150 mM NaCl

10 mM phosphate

pH 8.5

2. (ii) M72 with two N-terminal His residues - ASA(150 mM NaCl - 2) pH 8.0

10 ug antigen (20 ug/ml)

5% w/v sucrose

16 mM Tris

0.02% w/v Tween80

500 ug DOPC

125 ug cholesterol

25 ug 3D-MPL

25 ug QS21

150 mM NaCl

10 mM phosphate

pH 8.0

3. (iii) M72 with two N-terminal His residues - ASA(150 mM NaCl - 2) pH 7.5

10 ug antigen (20 ug/ml)

5% w/v sucrose

12.5 mM Tris

0.02% w/v Tween80

500 ug DOPC

125 ug cholesterol

25 ug 3D-MPL

25 ug QS21

150 mM NaCl

10 mM phosphate

pH 7.5

4. (iv) M72 with two N-terminal His residues - ASA(sorbitol - 2) pH 8.5

10 ug antigen (20 ug/ml)

5% w/v sucrose

40 mM Tris

0.02% w/v Tween80

500 ug DOPC

125 ug cholesterol

25 ug 3D-MPL

25 ug QS21

5 mM NaCl

4.7% w/v sorbitol

10 mM phosphate

pH 8.5

5. (v) M72 with two N-terminal His residues - ASA(sorbitol - 2) pH 8.0

10 ug antigen (20 ug/ml)

5% w/v sucrose

16 mM Tris

0.02% w/v Tween80

500 ug DOPC

125 ug cholesterol

25 ug 3D-MPL

25 ug QS21

5 mM NaCl

4.7% w/v sorbitol

10 mM phosphate

pH 8.0

6. (vi) M72 with two N-terminal His residues - ASA(sorbitol - 2) pH 7.5

10 ug antigen (20 ug/ml)

5% w/v sucrose

12.5 mM Tris

0.02% w/v Tween80

500 ug DOPC

125 ug cholesterol

25 ug 3D-MPL

25 ug QS21

5 mM NaCl

4.7% w/v sorbitol

10 mM phosphate

pH 7.5

Sample Analysis

[0163] The reconstituted immunogenic compositions described above were characterised after storage at 25°C or 30°C (T0, T6h and T24h).

[0164] SEC-HPLC analysis was performed by injection on a TOSOH TSK-Gel5000Pw λ l (ID 7.8 mm x 30 cm) equilibrated in 20 mM Tris buffer pH 8.5, detection by UV at 210 nm and flow rate 0.5 ml/min.

[0165] Antigenicity was quantified by a sandwich ELISA in which the antigen is captured by a M72-specific rabbit polyclonal antibody and subsequently revealed by a M72(Mtb39)-specific mouse monoclonal antibody. All measured values are presented relative to the expected antigenicity based on the purified bulk protein used to prepare the tested formulations.

Results

[0166] The results are shown in Figures 6a-6d and 7.

[0167] SEC-HPLC profiles are stable after reconstitution in low salt compositions using sorbitol as a tonicity agent at each pH (i.e. pH 7.5, 8.0 and 8.5). This may be contrasted with the SEC-HPLC profiles for immunogenic compositions containing 150 mM NaCl, which show clear changes between the initial profile obtained and those following storage at 25°C or 30°C. This evolution becomes more intense when the pH of the 150 mM NaCl composition is lowered.

[0168] The same conclusions can be drawn in terms of antigenicity, with recoveries remaining largely stable after reconstitution in low salt compositions using sorbitol as a tonicity agent at each pH (i.e. pH 7.5, 8.0 and 8.5) up to 24h at 30°C.

Example 11: Conductivity determination for Immunogenic compositions of the invention.

[0169] The conductivity of a range of immunogenic compositions according to the present invention was measured and compared to the conductivity of control sodium chloride solutions and with an immunogenic composition containing a conventional quantity of sodium chloride.

Method

[0170] A range of standards having sodium chloride concentrations of 0, 75, 100, 150, 250 and 300 mM were prepared from a stock solution of 1500 mM sodium chloride by dilution in water for injection.

[0171] Immunogenic compositions were prepared using M72 with two N-terminal His residues according to the procedures provided in Example 8. To investigate the contribution from the antigen itself and any residual materials in the purified bulk, placebo lyophilisation cakes were also prepared by excluding the antigen component.

[0172] Using a Malvern Zetasizer Nano and 1.5 ml of each sample in folded capillary cells, a voltage of 30 to 150 V (determined automatically by the instrument) was applied and the conductivity determined.

Results

[0173]

Conductivity of sodium chloride standard solutions

Sodium chloride concentration	Conductivity
mM	mS/cm
0	0.0
75	8.2
100	10.7

Sodium chloride concentration	Conductivity
mM	mS/cm
150	15.6
250	23.9
300	30.0

[0174] A standard curve, based on this data, is provided in Figure 8.

Conductivity of test solutions

Description	Sodium chloride concentration mM	Conductivity mS/cm	Equivalent sodium chloride concentration mM
ASA(sorbitol - 2)	5	1.46	9
Placebo pH 8.0 / ASA(sorbitol - 2)	5	1.95	14
M72 pH 8.0 / ASA(sorbitol - 2)	5	1.96	14
Placebo pH 8.5 / ASA(sorbitol - 2)	5	2.36	18
M72 pH 8.5 / ASA(sorbitol - 2)	5	2.28	17
ASA(150 mM NaCl - 2)	150	16	159
Placebo pH 8.5 / ASA(150 mM NaCl - 2)	150	14.8	147
M72 pH 8.5 / ASA(150 mM NaCl - 2)	150	15.3	152

[0175] As can be seen from the data above, the conductivity of solutions which utilise 150 mM NaCl is significantly greater than that of solutions which make minimal use of NaCl.

[0176] The impact of the antigen and any components in the purified bulk is minimal, as placebo preparations have comparable conductivity to their M72 related antigen containing counterparts.

Example 12: Immunogenicity testing of immunogenic compositions of the invention.

[0177] The aim of the this Example was to determine whether or not formulation changes to reduce the quantity of salt in immunogenic compositions of the invention, with a view to improving protein stability, had an impact on *in vivo* immunogenicity.

Method

[0178] Four immunogenic compositions were evaluated:

1. M72 with two N-terminal His residues pH 8.5/ASA (150 mM NaCl - 2)
2. M72 with two N-terminal His residues pH 8.5/ASA (sorbitol - 2)
3. M72 with two N-terminal His residues pH 8/ASA (sorbitol - 2)
4. M72 with two N-terminal His residues pH 7.5/ASA (sorbitol - 2)

[0179] The immunogenicity of these antigen containing compositions was evaluated in C57BL/6 mice. For each of the four compositions, 30 C57BL/6 mice were injected 3 times intramuscularly, on days 0, 14 and 28 with 1 ug of antigen in 50 ul of adjuvant solution (prepared by the procedure provided in Example 10). The elicited M72 specific T cell responses (both CD4 &

CD8) were measured 6 days post last immunisation (6dPIII).

[0180] For the determination of M72-specific cellular responses, peripheral blood lymphocytes from 30 mice/group were collected and pooled (six pools of five mice/group). A red blood cells lysis was performed before plating the cells *in vitro*. The cells were restimulated *in vitro* with a pool of overlapping peptides (15-mer peptides with an 11 amino acid overlap, at 1 ug/ml/peptide) covering the M72 sequence (without the N-terminal His residues). Cells remaining in medium (no peptide stimulation) were used to determine the background responses. Two hours after the co-culture with the peptide pool, brefeldin A was added to the wells (to inhibit cytokine excretion) and the cells were stored overnight at 4 °C. The cells were subsequently stained for the following markers: CD4, CD8, IL-2, IFN-gamma and TNF-alpha.

Results

[0181] Each datapoint in Figures 9 and 10 represents the background subtracted M72-specific CD4 or CD8 T cell response, respectively, of a pool of peripheral blood lymphocytes from five mice six days after the third immunisation. The response is expressed as the percentage of CD4 T cells producing IFN-gamma and/or IL-2 and/or TNF-alpha in response to stimulation with the M72 peptide pool. The bar represents the median of the responses for each group.

[0182] The results in Figure 9 and 10 show that comparable CD4 and CD8 T cell responses are induced following three immunisations with each of the test formulations. Consequently, it may be concluded that a reduction in the quantity of salts present in the immunogenic compositions of the present invention does not lead to a compromise in the induced T cell responses.

Example 13: Investigation of "salting out" in compositions comprising M72 with CaCl₂ or MgSO₄ at pH 6.1 and 8.0.

[0183] To investigate impact of other salts on M72 antigen stability, solutions were prepared with a range of concentrations of CaCl₂ or MgSO₄ and at different pH levels. Visual inspection was used as a readout of stability.

Method

[0184] Purified bulk antigen (M72 with two N-terminal His residues, SEQ ID No: 3) was diluted to a concentration of 100 ug/ml in two different buffers (10mM succinate buffer at pH 6.1 and 10 mM Tris buffer at pH 8.0) containing specified quantities of salts (0 mM; 150 mM or 300 mM NaCl; 40 mM, 80 mM or 160 mM CaCl₂; 87.5 mM, 175 mM or 430 mM MgSO₄).

[0185] Samples were analysed directly after preparation.

[0186] Using a Mettler Toledo conductivity meter and 6 ml of each sample in an unsiliconised glass vial, the conductivity was determined.

Results

[0187]

<u>Group</u>	<u>Salt</u>	<u>Buffer</u>	<u>pH</u> <u>(theoretical)</u>	<u>Conductivity (mS/cm)</u> <u>(measured)</u>	<u>pH</u> <u>(measured)</u>	<u>Visual</u> <u>Observation</u>
A	0 mM	Succinate 10mM	6.1	1.1	6.3	Clear
B	NaCl 150 mM	Succinate 10mM	6.1	13.4	6.1	Clear
C	NaCl 300 mM	Succinate 10mM	6.1	20.0	6.1	Clear
D	CaCl ₂ 40 mM	Succinate 10mM	6.1	8.0	6.1	Opalescent

Group	Salt	Buffer	pH (theoretical)	Conductivity (mS/cm) (measured)	pH (measured)	Visual Observation
E	CaCl ₂ 80 mM	Succinate 10mM	6.1	11.2	5.8	Opalescent + large particles
F	CaCl ₂ 160 mM	Succinate 10mM	6.1	20.2	5.8	Opalescent + large particles
G	MgSO ₄ 87.5 mM	Succinate 10mM	6.1	7.7	6.1	Opalescent
H	MgSO ₄ 175 mM	Succinate 10mM	6.1	12.4	5.9	Opalescent+ very large particles
I	MgSO ₄ 430 mM	Succinate 10mM	6.1	20.4	5.9	Opalescent+ very large particles
J	0 mM	Tris 10mM	8.0	0.463	8.0	Clear
K	NaCl 150 mM	Tris 10mM	8.0	12.13	8.0	Clear
L	NaCl 300 mM	Tris 10mM	8.0	21.1	8.0	Clear
M	CaCl ₂ 40 mM	Tris 10mM	8.0	6.7	8.1	Large particles
N	CaCl ₂ 80 mM	Tris 10mM	8.0	10.8	8.0	Opalescent+ large particles
O	CaCl ₂ 160 mM	Tris 10mM	8.0	19.7	8.0	Opalescent+ large particles
P	MgSO ₄ 87.5 mM	Tris 10mM	8.0	7.5	8.0	Large particles
Q	MgSO ₄ 175 mM	Tris 10mM	8.0	10.9	8.2	Opalescent+ very large particles
R	MgSO ₄ 430 mM	Tris 10mM	8.0	21.7	8.1	Opalescent+ very large particles

[0188] The results demonstrate that solutions containing an M72 related antigen can be sensitive to salts other than sodium chloride. The impact of CaCl₂ or MgSO₄ appears to be more pronounced than for sodium chloride at comparable concentrations or conductivity.

Example 14: Investigation of "salting out" in compositions comprising Mtb72f with different salt concentrations at pH 6.1, 7.5 and 8.5.

[0189] The impact of sodium chloride concentration and pH on Mtb72f antigen stability, as assessed by size, was investigated.

Method

[0190] Purified bulk antigen (Mtb72f with 6 his residues, SEQ ID No: 7) was diluted to a concentration of 100 ug/ml in three different buffers (10mM phosphate buffer at pH 6.1, 20 mM Tris buffer at pH 7.5 and 20 mM Tris buffer at pH 8.5) containing final sodium chloride concentrations of 0, 150 and 450 mM.

[0191] Samples were stored for 24 hours at 4°C or 25°C before analysis.

[0192] Nephelometry was performed using a Nepheloskan® Ascent, available from Thermo Fischer Scientific. Analysis was performed in UV transparent Costar® micro-plates available from Corning Inc (USA).

[0193] DLS was performed using a Dynapro Plate Reader from Wyatt Instruments. The instrument was operated using a laser wavelength of 830 nm and power of 50 mW. Scattered light was detected at 150° at a temperature of 22°C. The mean hydrodynamic diameter and polydispersity index (pl) are calculated by the instrument software.

Results

[0194] The findings of this experiment are presented in Figures 11 and 12.

[0195] Both DLS and nephelometry demonstrate a general trend that Mtb72f is sensitive to salt concentration and pH, in a similar manner to M72 as shown in previous examples. Consequently, the benefits of the present invention apply to M72 related antigens and not just to the M72 sequence itself.

[0196] In the case of a number of DLS samples, instrumentation was unable to determine a specific particle size (shown as NV in Figure 12).

Example 15: Prevention of "salting out" in compositions comprising M72, immunostimulants and using sorbitol as a tonicity agent

[0197] In order to compare the stability of immunogenic compositions containing 150 mM NaCl with compositions using sorbitol as a tonicity agent, samples were monitored using an alternative ELISA.

Method

[0198] Lyophilisation cake was prepared as described in Example 10 (specifically method (a)) such that when combined with the appropriate adjuvant formulations from Example 7 a pH of 8.5 would be obtained.

[0199] The lyophilisation cake describe above were reconstituted with 625 ul of the adjuvant solutions prepared in Example 7. Upon reconstitution with adjuvant solution, the following immunogenic compositions were obtained:

1. (i) M72 with two N-terminal His residues - ASA(150 mM NaCl - 2) pH 8.5

10 ug antigen (20 ug/ml)

5% w/v sucrose

40 mM Tris

0.02% w/v Tween80

500 ug DOPC

125 ug cholesterol

25 ug 3D-MPL

25 ug QS21

150 mM NaCl

10 mM phosphate

pH 8.5

2. (ii) M72 with two N-terminal His residues - ASA(sorbitol - 2) pH 8.5

10 ug antigen (20 ug/ml)

5% w/v sucrose

40 mM Tris
 0.02% w/v Tween80
 500 ug DOPC
 125 ug cholesterol
 25 ug 3D-MPL
 25 ug QS21
 5 mM NaCl
 4.7% w/v sorbitol
 10 mM phosphate
 pH 8.5

[0200] The reconstituted immunogenic compositions described above were characterised after storage at 30°C (T24h) and compared with an extemporaneously prepared sample (T0).

[0201] Antigenicity was quantified by an indirect sandwich ELISA in which the antigen is captured by a M72-specific rabbit polyclonal antibody and subsequently revealed by a M72(Mtb39)-specific mouse monoclonal antibody. Briefly, the plate is coated with anti-M72 rabbit polyclonal antibody at the dilution of 1/8000 in Dulbecco's Phosphate Buffered Saline overnight at 4°C and after four washes the plates were blocked for 1h at 37°C with saturation buffer (PBS, 0.1 % Tween 20, 1% BSA). After the washing step, protein standard (M72 purified bulk: 1950 ug/ml), internal control (M72: 1768 µg/ml) and samples are loaded in wells from the first column of the plate at a concentration of approximately 0.25µg/ml and then a 2-fold serial dilution is performed in the saturation buffer (PBS, 0.025% Tween 20) from well 1 to 12 and incubated 1h30 at 37°C. After the washing step, the immune complex is then incubated 1h at 37°C with anti-M72 mouse monoclonal antibody at a dilution of 1/1000 in saturation buffer (PBS, 0.025% Tween 20). After four washes, a biotinylated rabbit anti-mouse polyclonal antibody was added at a dilution of 1/1000 in saturation buffer (PBS, 0.025% Tween 20). After four washes, the signal was amplified by adding Streptavidin-Horseradish Peroxidase diluted 1/4000 in saturation buffer (PBS, 0.025% Tween 20). After four washes, the signal was revealed by ortho phenylene diamine dihydrochlorid (OPDA) for 15min at RT and the reaction is stopped by addition of HCl 1M. The coloration is proportional to the quantity of bound anti-M72 antibody, and is measured at 490 nm and 620 nm. All washing steps were performed using PBS, 0.025% Tween 20.

[0202] All measured values are presented relative to the expected antigenicity based on the purified bulk protein used to prepare the tested formulations.

Results

[0203] The results are shown in Figure 13. Diamonds indicating the specific measurements for each of the three test samples, with a line indicating the average value.

[0204] Antigen recovery is largely stable after reconstitution in low salt compositions using sorbitol as a tonicity agent at pH 8.5 up to 24h at 30°C. Recovery in ASA(sorbitol-2) was 83.5% after 24 hours (T0 87.1%, meaning 95.9% of the relative antigenicity was maintained), whereas recovery in ASA(NaCl-2) was 54.5% after 24 hours (T0 81.0%, meaning only 67.3% of the relative antigenicity was maintained after storage).

[0205] In summary, Examples 9, 10 and 15 demonstrate for the first time the detrimental impact resulting from pH and NaCl concentration on the stability of immunogenic compositions containing an M72 related antigen. Example 13 extends this work to show that other salts may also have a detrimental impact on the stability of immunogenic compositions containing an M72 related antigen, with Example 14 demonstrating that the effect is also applicable to M72 related sequences.

[0206] Reformulation of the immunogenic compositions with a non-ionic tonicity agent addresses the antigen stability problems.

Additionally, Examples 3, 4 and 12 demonstrate the removal of substantially all NaCl from the immunogenic formulation and its replacement with sorbitol as a tonicity agent does not have a detrimental impact on the stimulation of T cell responses.

[0207] Stability of immunogenic compositions is key and may be particularly challenging when in isolated locations where refrigeration may not be readily accessible. By reducing the presence of salts in the immunogenic compositions, the present inventors have been able to reduce the extent of changes observed when the immunogenic compositions are stored.

SEQUENCE LISTING

[0208]

<110> GlaxoSmithKline Biologicals sa
Godart, Stephane
Laanan, Amina
Lemoine, Dominique

<120> Novel compositions

<130> VB64442

<150> US61/422723

<151> 2010-12-14

<160> 13

<170> PatentIn version 3.5

<210> 1

<211> 723

<212> PRT

<213> Artificial Sequence

<220>

<223> M72 fusion protein

<400> 1

```
Met Thr Ala Ala Ser Asp Asn Phe Gln Leu Ser Gln Gly Gly Gln Gly
1      5      10      15

Phe Ala Ile Pro Ile Gly Gln Ala Met Ala Ile Ala Gly Gln Ile Arg
20     25     30

Ser Gly Gly Gly Ser Pro Thr Val His Ile Gly Pro Thr Ala Phe Leu
35     40     45

Gly Leu Gly Val Val Asp Asn Asn Gly Asn Gly Ala Arg Val Gln Arg
50     55     60

Val Val Gly Ser Ala Pro Ala Ala Ser Leu Gly Ile Ser Thr Gly Asp
65     70     75     80

Val Ile Thr Ala Val Asp Gly Ala Pro Ile Asn Ser Ala Thr Ala Met
85     90     95

Ala Asp Ala Leu Asn Gly His His Pro Gly Asp Val Ile Ser Val Thr
100    105    110

Trp Gln Thr Lys Ser Gly Gly Thr Arg Thr Gly Asn Val Thr Leu Ala
115    120    125

Glu Gly Pro Pro Ala Glu Phe Met Val Asp Phe Gly Ala Leu Pro Pro
130    135    140
```

Glu Ile Asn Ser Ala Arg Met Tyr Ala Gly Pro Gly Ser Ala Ser Leu
 145 150 155 160
 Val Ala Ala Ala Gln Met Trp Asp Ser Val Ala Ser Asp Leu Phe Ser
 165 170 175
 Ala Ala Ser Ala Phe Gln Ser Val Val Trp Gly Leu Thr Val Gly Ser
 180 185 190
 Trp Ile Gly Ser Ser Ala Gly Leu Met Val Ala Ala Ala Ser Pro Tyr
 195 200 205
 Val Ala Trp Met Ser Val Thr Ala Gly Gln Ala Glu Leu Thr Ala Ala
 210 215 220
 Gln Val Arg Val Ala Ala Ala Tyr Glu Thr Ala Tyr Gly Leu Thr
 225 230 235 240
 Val Pro Pro Pro Val Ile Ala Glu Asn Arg Ala Glu Leu Met Ile Leu
 245 250 255
 Ile Ala Thr Asn Leu Leu Gly Gln Asn Thr Pro Ala Ile Ala Val Asn
 260 265 270
 Glu Ala Glu Tyr Gly Glu Met Trp Ala Gln Asp Ala Ala Ala Met Phe
 275 280 285
 Gly Tyr Ala Ala Ala Thr Ala Thr Ala Thr Ala Thr Leu Leu Pro Phe
 290 295 300
 Glu Glu Ala Pro Glu Met Thr Ser Ala Gly Gly Leu Leu Glu Gln Ala
 305 310 315 320
 Ala Ala Val Glu Glu Ala Ser Asp Thr Ala Ala Ala Asn Gln Leu Met
 325 330 335
 Asn Asn Val Pro Gln Ala Leu Gln Gln Leu Ala Gln Pro Thr Gln Gly
 340 345 350
 Thr Thr Pro Ser Ser Lys Leu Gly Gly Leu Trp Lys Thr Val Ser Pro
 355 360 365
 His Arg Ser Pro Ile Ser Asn Met Val Ser Met Ala Asn Asn His Met
 370 375 380
 Ser Met Thr Asn Ser Gly Val Ser Met Thr Asn Thr Leu Ser Ser Met
 385 390 395 400

Leu Lys Gly Phe Ala Pro Ala Ala Ala Ala Gln Ala Val Gln Thr Ala
 405 410 415
 Ala Gln Asn Gly Val Arg Ala Met Ser Ser Leu Gly Ser Ser Leu Gly
 420 425 430
 Ser Ser Gly Leu Gly Gly Gly Val Ala Ala Asn Leu Gly Arg Ala Ala
 435 440 445
 Ser Val Gly Ser Leu Ser Val Pro Gln Ala Trp Ala Ala Ala Asn Gln
 450 455 460
 Ala Val Thr Pro Ala Ala Arg Ala Leu Pro Leu Thr Ser Leu Thr Ser
 465 470 475 480
 Ala Ala Glu Arg Gly Pro Gly Gln Met Leu Gly Gly Leu Pro Val Gly
 485 490 495
 Gln Met Gly Ala Arg Ala Gly Gly Gly Leu Ser Gly Val Leu Arg Val
 500 505 510
 Pro Pro Arg Pro Tyr Val Met Pro His Ser Pro Ala Ala Gly Asp Ile
 515 520 525
 Ala Pro Pro Ala Leu Ser Gln Asp Arg Phe Ala Asp Phe Pro Ala Leu
 530 535 540
 Pro Leu Asp Pro Ser Ala Met Val Ala Gln Val Gly Pro Gln Val Val
 545 550 555 560
 Asn Ile Asn Thr Lys Leu Gly Tyr Asn Asn Ala Val Gly Ala Gly Thr
 565 570 575
 Gly Ile Val Ile Asp Pro Asn Gly Val Val Leu Thr Asn Asn His Val
 580 585 590
 Ile Ala Gly Ala Thr Asp Ile Asn Ala Phe Ser Val Gly Ser Gly Gln
 595 600 605
 Thr Tyr Gly Val Asp Val Val Gly Tyr Asp Arg Thr Gln Asp Val Ala
 610 615 620
 Val Leu Gln Leu Arg Gly Ala Gly Gly Leu Pro Ser Ala Ala Ile Gly
 625 630 635 640
 Gly Gly Val Ala Val Gly Glu Pro Val Val Ala Met Gly Asn Ser Gly
 645 650 655
 Gly Gln Gly Gly Thr Pro Arg Ala Val Pro Gly Arg Val Val Ala Leu
 660 665 670
 Gly Gln Thr Val Gln Ala Ser Asp Ser Leu Thr Gly Ala Glu Glu Thr
 675 680 685
 Leu Asn Gly Leu Ile Gln Phe Asp Ala Ala Ile Gln Pro Gly Asp Ala
 690 695 700
 Gly Gly Pro Val Val Asn Gly Leu Gly Gln Val Val Gly Met Asn Thr
 705 710 715 720
 Ala Ala Ser
 <210> 2
 <211> 2172
 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> Coding sequence for M72 fusion
 <400> 2

```

atgacggccg cgtccgataa cttccagctg tcccaggggtg ggcagggatt cgccattccg      60
atcggggcagg cgatggcgat cgcgggccag atccgatcgg gtgggggggtc acccaacggt      120
catatcgggc ctaccgcctt cctcggcttg ggtgttgtcg acaacaacgg caacggcgca      180
cgagtccaac cgctggctcg gagcgctccg gcggcaagtc tcggcatctc caccggcgac      240
gtgatcaccg cggtcgacgg cgtccgatc aactcggcca ccgcatggo ggaacgcgtt      300
aacgggcatc atcccgggta cgtcatctcg gtgacctggc aaaccaagtc gggcggcacg      360
cgtacaggga acgtgacatt ggccgaggga ccccggccg aattcatggt ggatttcggg      420
gcgttaccac cggagatcaa ctccgcgagg atgtacgccg gcccggggtc ggcctcgtg      480
gtggcccgcg ctcatagtgt ggacagcgtg gcgagtgacc tgttttcggc cgcgtcggcg      540
tttcagtcgg tggctctggg tctgacgggt ggtcgtgga taggttcgtc ggcgggtctg      600
atggtggcgg cggcctcgcc gtatgtggcg tggatgagcg tcaccgcggg gcaggccgag      660
ctgaccgcgg cccaggtccg ggttgctgcg gcggcctacg agacggcgta tgggctgacg      720
gtgccccgcg cggtgatcgc cgagaacggt gctgaactga tgattctgat agcgaccaac      780
ctcttggggc aaaaaccccc ggcgatcgcg gtcaacgagg ccgaatacgg cgagatgtgg      840
gcccaagacg ccgcgcgat gtttggtac gccgcggcga cggcgacggc gacggcgacg      900
ttgctgccgt tcgaggagcg gccggagatg accagcgcgg gtgggtcctc cgagcaggcc      960

gccgcggtcg aggaggcctc cgacaccgcc gcggcgaaac agttgatgaa caatgtgcc      1020
caggcgctgc aacagctggc ccagcccacg cagggcacca cgcctctctc caagctgggt      1080
ggcctgtgga agacgggtct gccgcgatcg tcgccgatca gcaacatggt gtcgatggcc      1140
aacaaccaca tgtcgatgac caactcgggt gtgtcgatga ccaacacctt gagctcgatg      1200
ttgaagggtt ttgctccggc gcggccgcc caggccgtgc aaaccgcggc gcaaaacggg      1260
gtcggggcga tgagctcgct gggcagctcg ctgggttctt cgggtctggg cgggtgggtg      1320
gccgccaaact tgggtcgggc ggcctcggtc ggttcgttgt cggtgccgca ggcctgggcc      1380
gcggccaacc aggcagtcac ccgcgcggcg cgggcgctgc cgtgaccag cctgaccagc      1440
gcccgggaaa gagggcccg gcagatgctg ggcgggctgc cggtggggca gatgggcgcc      1500
agggccggtg gtgggctcag tgggtgtctg cgtgttccgc cgcgacctc tgtgatgccg      1560
cattctccgg cagccggcga tatcggccc cggccttgt cgcaggaccg gttcgccgac      1620
ttccccgcgc tgcccctcga ccgctccgcg atggtcgccc aagtggggcc acaggtggtc      1680
aacatcaaca ccaaactggg ctacaacaac gccgtggggc cggggaccgg catcgtcatc      1740
gatcccaacg gtgtcgtgct gaccaacaac cacgtgatcg cgggcgcac cgacatcaat      1800
gcgttcagcg tcggctccgg ccaaacctac gccgtcgatg tggtcgggta tgaccgcacc      1860
caggatgtcg cgggtgtgca gctgcgcggg gccgggtggc tgccgtcggc ggcgatcggg      1920
ggcgcgctcg cggttgtgta gcccgctgtc gcgatgggca acagcggtgg gcaggcgga      1980
acgccccgtg cggtgccctg cagggtggtc gcgctcggcc aaaccgtgca ggcgtcggat      2040
tcgctgaccg gtgcogaaga gacattgaac gggttgatcc agttcgatgc cgcgatccag      2100
cccgtgatg cgggcggggc cgtcgtcaac ggcctaggac aggtggtcgg tatgaacacg      2160
gccgcgtcct ag                                         2172

```

<210> 3

<211> 725

<212> PRT

<213> Artificial Sequence

<220>

<223> M72 fusion with 2 additional his residues

<400> 3

Met His His Thr Ala Ala Ser Asp Asn Phe Gln Leu Ser Gln Gly Gly
 1 5 10 15
 Gln Gly Phe Ala Ile Pro Ile Gly Gln Ala Met Ala Ile Ala Gly Gln
 20 25 30
 Ile Arg Ser Gly Gly Gly Ser Pro Thr Val His Ile Gly Pro Thr Ala
 35 40 45
 Phe Leu Gly Leu Gly Val Val Asp Asn Asn Gly Asn Gly Ala Arg Val
 50 55 60
 Gln Arg Val Val Gly Ser Ala Pro Ala Ala Ser Leu Gly Ile Ser Thr
 65 70 75 80
 Gly Asp Val Ile Thr Ala Val Asp Gly Ala Pro Ile Asn Ser Ala Thr
 85 90 95
 Ala Met Ala Asp Ala Leu Asn Gly His His Pro Gly Asp Val Ile Ser
 100 105 110
 Val Thr Trp Gln Thr Lys Ser Gly Gly Thr Arg Thr Gly Asn Val Thr
 115 120 125
 Leu Ala Glu Gly Pro Pro Ala Glu Phe Met Val Asp Phe Gly Ala Leu
 130 135 140
 Pro Pro Glu Ile Asn Ser Ala Arg Met Tyr Ala Gly Pro Gly Ser Ala
 145 150 155 160
 Ser Leu Val Ala Ala Ala Gln Met Trp Asp Ser Val Ala Ser Asp Leu
 165 170 175
 Phe Ser Ala Ala Ser Ala Phe Gln Ser Val Val Trp Gly Leu Thr Val
 180 185 190
 Gly Ser Trp Ile Gly Ser Ser Ala Gly Leu Met Val Ala Ala Ala Ser
 195 200 205
 Pro Tyr Val Ala Trp Met Ser Val Thr Ala Gly Gln Ala Glu Leu Thr
 210 215 220
 Ala Ala Gln Val Arg Val Ala Ala Ala Ala Tyr Glu Thr Ala Tyr Gly
 225 230 235 240
 Leu Thr Val Pro Pro Pro Val Ile Ala Glu Asn Arg Ala Glu Leu Met
 245 250 255
 Ile Leu Ile Ala Thr Asn Leu Leu Gly Gln Asn Thr Pro Ala Ile Ala
 260 265 270
 Val Asn Glu Ala Glu Tyr Gly Glu Met Trp Ala Gln Asp Ala Ala Ala
 275 280 285

Met Phe Gly Tyr Ala Ala Ala Thr Ala Thr Ala Thr Ala Thr Leu Leu
 290 295 300

Pro Phe Glu Glu Ala Pro Glu Met Thr Ser Ala Gly Gly Leu Leu Glu
 305 310 315 320

Gln Ala Ala Ala Val Glu Glu Ala Ser Asp Thr Ala Ala Ala Asn Gln
 325 330 335

Leu Met Asn Asn Val Pro Gln Ala Leu Gln Gln Leu Ala Gln Pro Thr
 340 345 350

Gln Gly Thr Thr Pro Ser Ser Lys Leu Gly Gly Leu Trp Lys Thr Val
 355 360 365

Ser Pro His Arg Ser Pro Ile Ser Asn Met Val Ser Met Ala Asn Asn
 370 375 380

His Met Ser Met Thr Asn Ser Gly Val Ser Met Thr Asn Thr Leu Ser
 385 390 395 400

Ser Met Leu Lys Gly Phe Ala Pro Ala Ala Ala Ala Gln Ala Val Gln
 405 410 415

Thr Ala Ala Gln Asn Gly Val Arg Ala Met Ser Ser Leu Gly Ser Ser
 420 425 430

Leu Gly Ser Ser Gly Leu Gly Gly Gly Val Ala Ala Asn Leu Gly Arg
 435 440 445

Ala Ala Ser Val Gly Ser Leu Ser Val Pro Gln Ala Trp Ala Ala Ala
 450 455 460

Asn Gln Ala Val Thr Pro Ala Ala Arg Ala Leu Pro Leu Thr Ser Leu
 465 470 475 480

Thr Ser Ala Ala Glu Arg Gly Pro Gly Gln Met Leu Gly Gly Leu Pro
 485 490 495

Val Gly Gln Met Gly Ala Arg Ala Gly Gly Gly Leu Ser Gly Val Leu
 500 505 510

Arg Val Pro Pro Arg Pro Tyr Val Met Pro His Ser Pro Ala Ala Gly
 515 520 525

Asp Ile Ala Pro Pro Ala Leu Ser Gln Asp Arg Phe Ala Asp Phe Pro
 530 535 540

Ala Leu Pro Leu Asp Pro Ser Ala Met Val Ala Gln Val Gly Pro Gln
545 550 555 560

Val Val Asn Ile Asn Thr Lys Leu Gly Tyr Asn Asn Ala Val Gly Ala
565 570 575

Gly Thr Gly Ile Val Ile Asp Pro Asn Gly Val Val Leu Thr Asn Asn
580 585 590

His Val Ile Ala Gly Ala Thr Asp Ile Asn Ala Phe Ser Val Gly Ser
595 600 605

Gly Gln Thr Tyr Gly Val Asp Val Val Gly Tyr Asp Arg Thr Gln Asp
610 615 620

Val Ala Val Leu Gln Leu Arg Gly Ala Gly Gly Leu Pro Ser Ala Ala
625 630 635 640

Ile Gly Gly Gly Val Ala Val Gly Glu Pro Val Val Ala Met Gly Asn
645 650 655

Ser Gly Gly Gln Gly Gly Thr Pro Arg Ala Val Pro Gly Arg Val Val
660 665 670

Ala Leu Gly Gln Thr Val Gln Ala Ser Asp Ser Leu Thr Gly Ala Glu
675 680 685

Glu Thr Leu Asn Gly Leu Ile Gln Phe Asp Ala Ala Ile Gln Pro Gly
690 695 700

Asp Ala Gly Gly Pro Val Val Asn Gly Leu Gly Gln Val Val Gly Met
705 710 715 720

Asn Thr Ala Ala Ser
725

<210> 4

<211> 2178

<212> DNA

<213> Artificial Sequence

<220>

<223> Coding sequence for M72 fusion with 2 additional his residues

<400> 4

atgcatacaca cggccgcgctc cgataacttc cagctgtccc aggggtgggca gggattcgcc	60
attccgatcg ggcaggcgat ggcatcgcg gccagatcc gatcgggtgg ggggtcaccc	120
accgttcata tcgggcctac cgccttctc ggcttgggtg ttgtcgacaa caacggcaac	180

```

ggcgacagag tccaacgcgt ggtcgggagc gtcocggcgg caagtctcgg catctccacc 240
ggcgacgtga tcaccgcggt cgacggcgct cagatcaact cggccaccgc gatgcgggac 300
gcgcttaacg ggcacatccc cggtgacgtc atctcgggtga cctggcaaac caagtcgggc 360
ggcacgcgta cagggaaacgt gacattggcc gagggacccc cggccgaatt catggtggat 420
ttcggggcgt taccaccgga gatcaactcc gcgaggatgt acgcggcccc gggttcggcc 480
tcgctgggtg cgcgggtca gatgtgggac agcgtggcga gtgacctgtt ttcggccgcg 540
tcggcggttc agtcgggtgt ctggggctct acggtggggt cgtggatagg ttcgtcggcg 600
ggctctgatg tggcggcgcc ctcgccttat gtggcgtgga tgagcgtcac cgcggggcag 660
gccgagctga ccgcgcacca ggtccgggtt gctcggcgg cctacgagac ggcgtatggg 720
ctgacgggtc cccgcgcggt gatcgccgag aaccgtgctg aactgatgat tctgatagcg 780
accaacctct tggggcaaaa caccgcggcg atcgcgtca acgaggccga atacggcgag 840
atgtggggcc aagacgcgcg cgcgatgttt ggctacgccg cggcgacggc gacggcgacg 900
gcgacgttgc tgcctgttca ggaggcgccg gagatgacca gcgcgggtgg gtcctcagag 960
caggcccgcg cggctcagga ggcctccgac accgcgcgg cgaaccagtt gatgaacaat 1020
gtgccccagg cgctgaaca gctggcccag cccacgcagg gcaccacgcc ttcttccaag 1080
ctgggtggcc tgtggaagac ggtctcgcg catcggtcgc cagtcagcaa catggtgtcg 1140
atggccaaca accacatgtc gatgaccaac tcgggtgtgt cagtgaccaa caccctgagc 1200
tcgatgttga agggctttgc tccggcggcg gccgccagg ccgtgcaaac cgcggcgcaa 1260
aacgggggtc gggcgatgag ctcgtgggc agctcgtgg gttctcggg tctggcggt 1320
ggggtggcgg ccaacttggg tcggggcgcc tcggtcggtt cgttgcggt gccgcaggcc 1380
tgggcccggg ccaaccaggc agtcacccc gcggcgcggg cgctgcgct gaccagcctg 1440
accagcgccg cggaaaagag gcccgggcag atgctggcg ggcctgcggt ggggcagatg 1500
ggcgccaggg ccgtgtgtg gctcagtgt gtgctgcgtg tccgcgcg accctatgtg 1560
atgcgcatt ctcggcgagc cggcgatata gcccgccgg ccttgctgca ggaccggttc 1620
gccgacttcc cgcgcgtgcc cctcgaccg tcgcgatgg tcgccaaagt ggggccacag 1680
gtggtcaaca tcaacaccaa actgggctac aacaacgccg tggcgccgg gaccggcatc 1740
gtcatcgatc ccaacggtgt cgtgctgacc aacaaccacg tgatcgcggg cgccaccgac 1800
atcaatgcgt tcagcgtcgg ctcggcccaa acctacggcg tcgatgtggt cgggtatgac 1860
cgcacccagg atgtcgcggt gctcagctg cgcggtgccg gtggcctgcc gtcggcgcg 1920
atcggtggcg gcgtcgcggt tggtagccc gtgctcgca tgggcaacag cggtaggcag 1980
ggcggaacgc ccgctgcggt gcctggcagg gtggtcgcgc tcggccaaac cgtgcaggcg 2040

tcggattcgc tgaccggtgc cgaagagaca ttgaacgggt tgatccagtt cgatgccgcg 2100
atccagcccg gtgatgcggg cgggcccgtc gtcaacggcc taggacaggt ggtcggtatg 2160
aacacggccg cgtcctag 2178

```

<210> 5

<211> 723

<212> PRT

<213> Artificial Sequence

<220>

<223> Mtb72f fusion

<400> 5

Met Thr Ala Ala Ser Asp Asn Phe Gln Leu Ser Gln Gly Gly Gln Gly
 1 5 10 15
 Phe Ala Ile Pro Ile Gly Gln Ala Met Ala Ile Ala Gly Gln Ile Arg
 20 25 30
 Ser Gly Gly Gly Ser Pro Thr Val His Ile Gly Pro Thr Ala Phe Leu
 35 40 45
 Gly Leu Gly Val Val Asp Asn Asn Gly Asn Gly Ala Arg Val Gln Arg
 50 55 60
 Val Val Gly Ser Ala Pro Ala Ala Ser Leu Gly Ile Ser Thr Gly Asp
 65 70 75 80
 Val Ile Thr Ala Val Asp Gly Ala Pro Ile Asn Ser Ala Thr Ala Met
 85 90 95
 Ala Asp Ala Leu Asn Gly His His Pro Gly Asp Val Ile Ser Val Thr
 100 105 110
 Trp Gln Thr Lys Ser Gly Gly Thr Arg Thr Gly Asn Val Thr Leu Ala
 115 120 125
 Glu Gly Pro Pro Ala Glu Phe Met Val Asp Phe Gly Ala Leu Pro Pro
 130 135 140
 Glu Ile Asn Ser Ala Arg Met Tyr Ala Gly Pro Gly Ser Ala Ser Leu
 145 150 155 160
 Val Ala Ala Ala Gln Met Trp Asp Ser Val Ala Ser Asp Leu Phe Ser
 165 170 175
 Ala Ala Ser Ala Phe Gln Ser Val Val Trp Gly Leu Thr Val Gly Ser
 180 185 190

Trp Ile Gly Ser Ser Ala Gly Leu Met Val Ala Ala Ala Ser Pro Tyr
 195 200 205
 Val Ala Trp Met Ser Val Thr Ala Gly Gln Ala Glu Leu Thr Ala Ala
 210 215 220
 Gln Val Arg Val Ala Ala Ala Tyr Glu Thr Ala Tyr Gly Leu Thr
 225 230 235 240
 Val Pro Pro Pro Val Ile Ala Glu Asn Arg Ala Glu Leu Met Ile Leu
 245 250 255
 Ile Ala Thr Asn Leu Leu Gly Gln Asn Thr Pro Ala Ile Ala Val Asn
 260 265 270
 Glu Ala Glu Tyr Gly Glu Met Trp Ala Gln Asp Ala Ala Met Phe
 275 280 285
 Gly Tyr Ala Ala Ala Thr Ala Thr Ala Thr Ala Thr Leu Leu Pro Phe
 290 295 300
 Glu Glu Ala Pro Glu Met Thr Ser Ala Gly Gly Leu Leu Glu Gln Ala
 305 310 315 320
 Ala Ala Val Glu Glu Ala Ser Asp Thr Ala Ala Ala Asn Gln Leu Met
 325 330 335
 Asn Asn Val Pro Gln Ala Leu Gln Gln Leu Ala Gln Pro Thr Gln Gly
 340 345 350
 Thr Thr Pro Ser Ser Lys Leu Gly Gly Leu Trp Lys Thr Val Ser Pro
 355 360 365
 His Arg Ser Pro Ile Ser Asn Met Val Ser Met Ala Asn Asn His Met
 370 375 380
 Ser Met Thr Asn Ser Gly Val Ser Met Thr Asn Thr Leu Ser Ser Met
 385 390 395 400
 Leu Lys Gly Phe Ala Pro Ala Ala Ala Ala Gln Ala Val Gln Thr Ala
 405 410 415
 Ala Gln Asn Gly Val Arg Ala Met Ser Ser Leu Gly Ser Ser Leu Gly
 420 425 430
 Ser Ser Gly Leu Gly Gly Gly Val Ala Ala Asn Leu Gly Arg Ala Ala

435 440 445
 Ser Val Gly Ser Leu Ser Val Pro Gln Ala Trp Ala Ala Ala Asn Gln
 450 455 460
 Ala Val Thr Pro Ala Ala Arg Ala Leu Pro Leu Thr Ser Leu Thr Ser
 465 470 475 480
 Ala Ala Glu Arg Gly Pro Gly Gln Met Leu Gly Gly Leu Pro Val Gly
 485 490 495
 Gln Met Gly Ala Arg Ala Gly Gly Gly Leu Ser Gly Val Leu Arg Val
 500 505 510
 Pro Pro Arg Pro Tyr Val Met Pro His Ser Pro Ala Ala Gly Asp Ile
 515 520 525
 Ala Pro Pro Ala Leu Ser Gln Asp Arg Phe Ala Asp Phe Pro Ala Leu
 530 535 540
 Pro Leu Asp Pro Ser Ala Met Val Ala Gln Val Gly Pro Gln Val Val
 545 550 555 560
 Asn Ile Asn Thr Lys Leu Gly Tyr Asn Asn Ala Val Gly Ala Gly Thr
 565 570 575
 Gly Ile Val Ile Asp Pro Asn Gly Val Val Leu Thr Asn Asn His Val
 580 585 590
 Ile Ala Gly Ala Thr Asp Ile Asn Ala Phe Ser Val Gly Ser Gly Gln
 595 600 605
 Thr Tyr Gly Val Asp Val Val Gly Tyr Asp Arg Thr Gln Asp Val Ala
 610 615 620
 Val Leu Gln Leu Arg Gly Ala Gly Gly Leu Pro Ser Ala Ala Ile Gly
 625 630 635 640
 Gly Gly Val Ala Val Gly Glu Pro Val Val Ala Met Gly Asn Ser Gly
 645 650 655
 Gly Gln Gly Gly Thr Pro Arg Ala Val Pro Gly Arg Val Val Ala Leu
 660 665 670
 Gly Gln Thr Val Gln Ala Ser Asp Ser Leu Thr Gly Ala Glu Glu Thr
 675 680 685
 Leu Asn Gly Leu Ile Gln Phe Asp Ala Ala Ile Gln Pro Gly Asp Ser
 690 695 700
 Gly Gly Pro Val Val Asn Gly Leu Gly Gln Val Val Gly Met Asn Thr
 705 710 715 720

Ala Ala Ser

<210> 6

<211> 2172

<212> DNA

<213> Artificial Sequence

<220>

<223> Coding sequence for Mtb72f fusion

<400> 6

```

atgacggccg cgtccgataa ctccagctg tcccaggtg ggcaggatt cgccattccg      60
atcgggcagg cgatggcgat cgcgggccag atccgatcgg gtggggggtc acccaccgtt    120
cataatcgggc ctaccgcctt cctcggcttg ggtgttgctg acaacaacgg caacggcgca    180
cgagtcacaac cgtcggctcg gagcgctccg gcggcaagtc tcggcatctc caccggcgac    240
gtgatcaccg cggtcgacgg cgctccgac aactcggcca ccgcatggg ggaacgcgtt    300
aacgggcacg atcccgggtg cgtcatctcg gtgacctggc aaaccaagtc gggcggcacg    360
cgtacaggga acgtgacatt ggccgaggga ccccggccg aattcatggt ggatttcggg    420
gcgttaccac cggagatcaa ctccgcgagg atgtacggc gcccgggttc ggcctcgtg      480
gtggccggcg ctcagatgtg ggacagcgtg gcgagtgacc tgttttcggc cgcgtcggcg    540
tttcagtcgg tggtctgggg tctgacggtg ggtctgtgga taggttcgtc ggcgggtctg    600
atggtggcgg cggcctcgcc gtatgtggcg tggatgagcg tcaccgggg gcaggccgag    660
ctgaccgccg cccaggtccg ggttgctcgg gcggcctacg agacggcgta tgggctgacg    720
gtgccccgcg cggtgatcgc cgagaaccgt gctgaactga tgattctgat agcgaccaac    780
ctcttggggc aaaaaccccc ggcatcgcg gtcaacgagg ccgaatacgg cgagatgtgg    840
gcccaagacg ccgcccgcat gtttggtac gcccgggcga cggcgacggc gacggcgacg    900
ttgctgccgt tcgaggaggc gccggagatg accagcgcg gtgggctcct cgagcaggcc    960
gcccggtcgg aggaggccct cgacaccgcc gcggcgaaac agttgatgaa caatgtgcc    1020
caggcgctgc aacagctggc ccagcccacg cagggcacca cgccttcttc caagctgggt    1080
ggcctgtgga agacggctc gccgcacgg tcgccgatca gcaacatggt gtcgatggcc    1140
aacaaccaca tgtcgatgac caactcgggt gtgtcgatga ccaacacct gagctcgatg    1200
ttgaagggtc ttgctccggc ggccggccgc caggccgtgc aaaccggcg gcaaaacggg    1260

gtccgggcga tgagctcgt gggcagctcg ctgggttctt cgggtctggg cggtggggtg    1320
gccgccaaat tgggtcgggc gccctcggtc ggttcgttgt cggtgccgca ggcctgggac    1380
gcggccaacc aggcagtcac ccggcgggcg cggcgctgc cgctgaccag cctgaccagc    1440
gcccgggaaa gagggcccg gcagatgctg ggccggctgc cggtggggca gatggcgcc    1500
agggccggtg gtgggctcag tgggtgctg cgtgttcgcc cgcgacctc tgtgatgccg    1560
cattctcggc cagccggcga tatcgcccc cgggccttgt ccgaggaccg gttcggcgac    1620
ttccccgcgc tgccccctga cccgtccggc atggtcgccc aagtggggcc acagggtggtc    1680
aacatcaaca ccaaactggg ctacaacaac gccgtggggc cggggaccgg catcgtcatc    1740
gatcccaacg gtgtcgtgct gaccaacaac cacgtgatcg cggcgccac cgacatcaat    1800
gcgttcagcg tcggctccgg ccaaacctac gccgtcgatg tggtcgggta tgaccgcacc    1860
caggatgtcg cgggtcgtga gctgcgcgg gccgggtggc tgccgtcggc ggcgatcggg    1920
ggcgggcgtc cggttggtga gccctcgtc gcgatgggca acagcgggtg gcaggcgga    1980
acgccccgtg cggtgccctg cagggtggtc gcgctcggc aaaccgtgca ggcgtcggat    2040
tcgctgaccg gtgccgaaga gacattgaac gggttgatcc agttcgatgc cgcgatccag    2100
cccggtgatt cggcggggcc cgtcgtcaac ggcctaggac aggtggtcgg tatgaacacg    2160
gccgcgtcct ag                                     2172

```

<210> 7

<211> 729

<212> PRT

<213> Artificial Sequence

<220>

<223> Mtb72f fusion with 6 additional his residues

<400> 7

Met His His His His His His Thr Ala Ala Ser Asp Asn Phe Gln Leu
 1 5 10 15
 Ser Gln Gly Gly Gln Gly Phe Ala Ile Pro Ile Gly Gln Ala Met Ala
 20 25 30
 Ile Ala Gly Gln Ile Arg Ser Gly Gly Gly Ser Pro Thr Val His Ile
 35 40 45
 Gly Pro Thr Ala Phe Leu Gly Leu Gly Val Val Asp Asn Asn Gly Asn
 50 55 60
 Gly Ala Arg Val Gln Arg Val Val Gly Ser Ala Pro Ala Ala Ser Leu
 65 70 75 80
 Gly Ile Ser Thr Gly Asp Val Ile Thr Ala Val Asp Gly Ala Pro Ile
 85 90 95
 Asn Ser Ala Thr Ala Met Ala Asp Ala Leu Asn Gly His His Pro Gly
 100 105 110
 Asp Val Ile Ser Val Thr Trp Gln Thr Lys Ser Gly Gly Thr Arg Thr
 115 120 125
 Gly Asn Val Thr Leu Ala Glu Gly Pro Pro Ala Glu Phe Met Val Asp
 130 135 140
 Phe Gly Ala Leu Pro Pro Glu Ile Asn Ser Ala Arg Met Tyr Ala Gly
 145 150 155 160
 Pro Gly Ser Ala Ser Leu Val Ala Ala Ala Gln Met Trp Asp Ser Val
 165 170 175
 Ala Ser Asp Leu Phe Ser Ala Ala Ser Ala Phe Gln Ser Val Val Trp
 180 185 190
 Gly Leu Thr Val Gly Ser Trp Ile Gly Ser Ser Ala Gly Leu Met Val
 195 200 205
 Ala Ala Ala Ser Pro Tyr Val Ala Trp Met Ser Val Thr Ala Gly Gln
 210 215 220
 Ala Glu Leu Thr Ala Ala Gln Val Arg Val Ala Ala Ala Tyr Glu
 225 230 235 240
 Thr Ala Tyr Gly Leu Thr Val Pro Pro Pro Val Ile Ala Glu Asn Arg
 245 250 255
 Ala Glu Leu Met Ile Leu Ile Ala Thr Asn Leu Leu Gly Gln Asn Thr
 260 265 270
 Pro Ala Ile Ala Val Asn Glu Ala Glu Tyr Gly Glu Met Trp Ala Gln
 275 280 285
 Asp Ala Ala Ala Met Phe Gly Tyr Ala Ala Ala Thr Ala Thr Ala Thr
 290 295 300
 Ala Thr Leu Leu Pro Phe Glu Glu Ala Pro Glu Met Thr Ser Ala Gly
 305 310 315 320
 Gly Leu Leu Glu Gln Ala Ala Ala Val Glu Glu Ala Ser Asp Thr Ala
 325 330 335

Ala Ala Asn Gln Leu Met Asn Asn Val Pro Gln Ala Leu Gln Gln Leu
 340 345 350
 Ala Gln Pro Thr Gln Gly Thr Thr Pro Ser Ser Lys Leu Gly Gly Leu
 355 360 365
 Trp Lys Thr Val Ser Pro His Arg Ser Pro Ile Ser Asn Met Val Ser
 370 375 380
 Met Ala Asn Asn His Met Ser Met Thr Asn Ser Gly Val Ser Met Thr
 385 390 395 400
 Asn Thr Leu Ser Ser Met Leu Lys Gly Phe Ala Pro Ala Ala Ala Ala
 405 410 415
 Gln Ala Val Gln Thr Ala Ala Gln Asn Gly Val Arg Ala Met Ser Ser
 420 425 430
 Leu Gly Ser Ser Leu Gly Ser Ser Gly Leu Gly Gly Gly Val Ala Ala
 435 440 445
 Asn Leu Gly Arg Ala Ala Ser Val Gly Ser Leu Ser Val Pro Gln Ala
 450 455 460
 Trp Ala Ala Ala Asn Gln Ala Val Thr Pro Ala Ala Arg Ala Leu Pro
 465 470 475 480
 Leu Thr Ser Leu Thr Ser Ala Ala Glu Arg Gly Pro Gly Gln Met Leu
 485 490 495
 Gly Gly Leu Pro Val Gly Gln Met Gly Ala Arg Ala Gly Gly Gly Leu
 500 505 510
 Ser Gly Val Leu Arg Val Pro Pro Arg Pro Tyr Val Met Pro His Ser
 515 520 525
 Pro Ala Ala Gly Asp Ile Ala Pro Pro Ala Leu Ser Gln Asp Arg Phe
 530 535 540
 Ala Asp Phe Pro Ala Leu Pro Leu Asp Pro Ser Ala Met Val Ala Gln
 545 550 555 560
 Val Gly Pro Gln Val Val Asn Ile Asn Thr Lys Leu Gly Tyr Asn Asn
 565 570 575
 Ala Val Gly Ala Gly Thr Gly Ile Val Ile Asp Pro Asn Gly Val Val
 580 585 590
 Leu Thr Asn Asn His Val Ile Ala Gly Ala Thr Asp Ile Asn Ala Phe
 595 600 605
 Ser Val Gly Ser Gly Gln Thr Tyr Gly Val Asp Val Val Gly Tyr Asp
 610 615 620
 Arg Thr Gln Asp Val Ala Val Leu Gln Leu Arg Gly Ala Gly Gly Leu
 625 630 635 640
 Pro Ser Ala Ala Ile Gly Gly Gly Val Ala Val Gly Glu Pro Val Val
 645 650 655
 Ala Met Gly Asn Ser Gly Gly Gln Gly Gly Thr Pro Arg Ala Val Pro
 660 665 670
 Gly Arg Val Val Ala Leu Gly Gln Thr Val Gln Ala Ser Asp Ser Leu
 675 680 685
 Thr Gly Ala Glu Glu Thr Leu Asn Gly Leu Ile Gln Phe Asp Ala Ala
 690 695 700
 Ile Gln Pro Gly Asp Ser Gly Gly Pro Val Val Asn Gly Leu Gly Gln
 705 710 715 720
 Val Val Gly Met Asn Thr Ala Ala Ser
 725

<210> 8

<211> 2190

<212> DNA

<213> Artificial Sequence

<220>

<223> Coding sequence for M72 fusion with 6 additional his residues

<400> 8

```

atgcatcacc atcaaccatca cacggccgcg tccgataact tccagctgtc ccagggtggg      60
cagggattcg ccattccgat cgggcaggcg atggcgatcg cgggccagat ccgatcgggt      120
gggggggtcac ccaccgttca tatcgggcct accgccttcc tcggcttggg tgttgtcgac      180
aacaacggca acggcgacag agtccaacgc gtggtcggga gcgctcgggc ggcaagtctc      240
ggcatctcca ccggcgacgt gatcaccgcg gtcgacggcg ctccgatcaa ctccggccacc      300
gcgatggcgg acgcgccttaa cgggcatcat cccggtgacg tcatctcggg gacctggcaa      360
accaagtccg gcggcacgcg tacagggaac gtgacattgg ccgagggacc cccggccgaa      420
ttcatggtgg atttcggggc gttaccaccg gagatcaact ccgcgaggat gtacgccggc      480

ccgggttcgg cctcgctggg gccgcgggt cagatgtggg acagcgtggc gagtgaacctg      540
ttttcggccg cgtcggcggt tcagtcgggt gtctggggtc tgacgggtgg gtctgtgata      600
ggttcgtcgg cgggtctgat ggtggcggcg gcctcggcgt atgtggcgtg gatgagcgtc      660
accgcggggc agggcgagct gaccgccgcc caggctcggg ttgctcgggc ggcctacgag      720
acggcgatat ggctgacggg gcccgcggcg gtgatcgccg agaaccgtgc tgaactgatg      780
attctgatat cgaccaacct cttggggcaa aacaccccg cgcctcgggt caacgaggcc      840
gaatacggcg agatgtgggc ccaagacgcc gccgcgatgt ttggctacgc cgcggcgacg      900
gcgacggcga cggcgacggt gctgccgttc gaggaggcgc cggagatgac cagcgcgggt      960
gggctcctcg agcaggccgc cgcggtcgag gaggcctccg acaccgccgc ggcgaaccag     1020
ttgatgaaca atgtgccccg gccgctgcaa cagctggccc agccacgca gggcaccacg     1080
cctctctcca agctgggtgg cctgtggaag acggtctcgc cgcctcgggt gccgatcagc     1140
aacatggtgt cgatggccaa caaccacatg tcgatgacca actcgggtgt gtcgatgacc     1200
aacaccttga gctcgatggt gaagggcttt gctccggcgg cggccgccca ggcgctgcaa     1260
accgcggcgc aaaacggggt ccgggcgatg agctcgtcgg gcagctcgtc gggttcttcg     1320
gggtcgggcg gtgggggtgc cgcacaactg ggtcgggcgg cctcggtcgg ttcgttgtcg     1380
gtgccgcagg cctggggcgc ggccaaccag gcagtcaccc cggcggcgcg ggcgtgccg      1440
ctgaccagcc tgaccagcgc cgcggaaga gggcccgggc agatgctggg cgggctgccg      1500
gtggggcaga tgggcgccag ggccgggtgg gggctcagtg gtgtgctcgc tgttccgccg      1560
cgaccctatg tgatgccgca ttctccgca gccggcgata tcgccccgcc ggccttgtcg      1620
caggaccggg tcgcgcactt cccgcgctg cccctcgacc cgtcccgat ggtcgcccaa      1680
gtggggccac agtggtgcaa catcaacacc aaactgggct acaacaacgc cgtgggcgcc      1740
gggaccggca tcgtcatcga tcccaacggt gtcgtgctga ccaacaacca cgtgatcgcg      1800
ggcgcacccg acatcaatgc gttcagcgtc ggctccggcc aaacctacgg cgtcgatgtg      1860
gtcgggtatg accgcaccca ggatgtcgcg gtgctgcagc tcgcgggtgc cggtggcctg      1920
ccgtcggcgg cgatcgggtg cggcgtcgcg gttggtgagc ccgtcgtcgc gatgggcaac      1980
agcgggtggc agggcggaac gcccgtgcg gtgcctgca ggggtggtcgc gctcggccaa      2040
accgtgcagg cgtcggattc gctgaccggt gccgaagaga cattgaacgg gttgatccag      2100
ttcgatgccg cgatccagcc cgggtattcg ggcgggcccg tcgtcaacgg cctaggacag      2160
gtggtcggta tgaacacggc cgcgtcctag                                     2190

```

<210> 9

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Cpg Oligo 1 - CpG 1826

<400> 9

tccatgacgt tctgacgtt 20

<210> 10

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<223> CpG Oligo 2 - CpG 1758

<400> 10

tctcccagcg tgcgcat 18

<210> 11

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> CpG Oligo 3

<400> 11

accgatgacg tcgccgtga cggcaccacg 30

<210> 12

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> CpG Oligo 4 - CpG 2006

<400> 12

tcgtcgtttt gtcgtttgt cggt 24

<210> 13

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> CpG Oligo 5 - CpG 1686

<400> 13

tccatgacgt tctgatgct 20

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- [WO2006117240A \[0003\] \[0005\] \[0030\] \[0051\]](#)
- [WO8809336A \[0065\]](#)
- [WO9633739A \[0067\]](#)
- [US4436727A \[0071\]](#)
- [US4877611A \[0071\]](#)
- [US4866034A \[0071\]](#)
- [US4912094A \[0071\]](#)
- [GB2220211A \[0071\]](#)
- [WO9421292A \[0071\]](#)
- [WO9850399A \[0072\]](#)
- [US6303347B \[0072\]](#)
- [US6764840B \[0072\]](#)
- [WO2003011223A \[0073\] \[0073\] \[0073\]](#)
- [WO2003099195A \[0073\] \[0073\]](#)
- [WO803022A \[0073\]](#)
- [WO803058A \[0073\]](#)
- [WO803732A \[0073\]](#)
- [WO804053A \[0073\]](#)
- [WO804057A \[0073\]](#)
- [WO804058A \[0073\]](#)
- [WO804059A \[0073\]](#)
- [WO804442A \[0073\]](#)
- [WO804680A \[0073\]](#)
- [WO804764A \[0073\]](#)
- [WO2008142133A \[0078\]](#)
- [WO9602555A \[0078\]](#)
- [WO9933488A \[0078\]](#)
- [US5865462A \[0078\]](#)
- [US5666153A \[0079\]](#)
- [US5278302A \[0079\]](#)
- [WO9526204A \[0079\]](#)
- [EP0362271A \[0082\]](#)
- [US61422723B \[0208\]](#)

Non-patent literature cited in the description

- World Health Organisation Tuberculosis Facts, 2010, [0002] [0025]
- **LOCHT et al.** Expert Opin. Biol. Ther., 2007, vol. 7, 11665-1677 [0021]
- **ANDERSON P** Trends in Microbiology, 2007, vol. 15, 17-13 [0021]
- **EHLERS S** Infection, 2009, vol. 37, 287-95 [0021]
- **CARDANA P-J** Inflammation & Allergy - Drug Targets, 2006, vol. 6, 27-39 [0022]
- **CARDANA P-J** Infection, 2009, vol. 37, 280-86 [0022]
- **BRANDT et al.** Infect. Immun., 2004, vol. 72, 116622-6632 [0030]
- **SKEIKY et al.** J. Immunol., 2004, vol. 172, 7618-7628 [0030]
- **TSENOVA et al.** Infect. Immun., 2006, vol. 74, 42392-2401 [0030]
- **REED et al.** PNAS, 2009, vol. 106, 72301-2306 [0030]
- **VON ESCHEN et al.** Human Vaccines, 2009, vol. 5, 7475-482 [0030]
- **PAUL** Fundamental Immunology 1993 0000243-247 [0033]
- **BEIßBARTH et al.** Bioinformatics, 2005, vol. 21, i29-i37 [0033]
- **CREIGHTON** Proteins, 1984, [0035]
- **SMITHWATERMAN** Adv. Appl. Math., 1981, vol. 2, 482- [0042]
- **NEEDLEMAN WUNSCH** J. Mol. Biol., 1970, vol. 48, 443- [0042]

- **PEARSONLIPMAN**Proc. Natl. Acad. Sci. USA, 1988, vol. 85, 2444- [0042]
- Current Protocols in Molecular Biology19950000 [0042]
- **FENGDOOLITTLE**J. Mol. Evol., 1987, vol. 35, 351-360 [0043]
- **HIGGINSSHARPCABIOS**, 1989, vol. 5, 151-153 [0043]
- **DEVEREAUX et al.**Nuc. Acids Res., 1984, vol. 12, 387-395 [0043]
- **ALTSCHUL et al.**Nuc. Acids Res., 1977, vol. 25, 3389-3402 [0044]
- **ALTSCHUL et al.**J. Mol. Biol., 1990, vol. 215, 403-410 [0044]
- **HENIKOFFHENIKOFF**Proc. Natl. Acad. Sci. USA, 1989, vol. 89, 10915- [0044]
- **KARLINALTSCHUL**Proc. Natl. Acad. Sci. USA, 1993, vol. 90, 5873-5787 [0045]
- Archiv. für die gesamte Virusforschung**DALSGAARD et al.**Saponin adjuvantsSpringer Verlag19740000vol. 44, 243-254 [0065]
- **SABROE et al.**J Immunol, 2003, 1630-5 [0070]
- Harrison's Principles of Internal Medicine20050000953-966 [0088] [0089]

P A T E N T K R A V

1. Immunogen sammensætning omfattende et M72-relateret antigen, hvor det M72-relaterede antigen omfatter en sekvens, der har mindst 90% identitet med SEQ ID No: 1, pH af sammensætningen er i området 7,0 til 9,0 og ledningsevnen af sammensætningen er 5 mS/cm eller lavere.
2. Immunogen sammensætning ifølge krav 1, hvor ledningsevnen af sammensætningen er 4 mS/cm eller lavere.
3. Immunogen sammensætning ifølge krav 2, hvor ledningsevnen af sammensætningen er 3 mS/cm eller lavere.
4. Immunogen sammensætning ifølge et hvilket som helst af krav 1 til 3, hvor koncentrationen af salte i sammensætningen er 40 mM eller lavere.
5. Immunogen sammensætning ifølge et hvilket som helst af krav 1 til 4, yderligere omfattende et ikke-ionisk tonicitetsmiddel.
6. Immunogen sammensætning ifølge krav 5, hvor det ikke-ioniske tonicitetsmiddel er en polyol.
7. Immunogen sammensætning ifølge krav 6, hvor polyolen er sorbitol, og hvor koncentrationen af sorbitol er mellem omkring 4 og omkring 6% (vægt/volumen).
8. Immunogen sammensætning ifølge et hvilket som helst af krav 1 til 7, hvor koncentrationen af sucrose er mellem omkring 4 og omkring 6% (vægt/volumen).
9. Immunogen sammensætning ifølge et hvilket som helst af krav 1 til 8, yderligere omfattende ét eller flere immunstimulerende midler.
10. Immunogen sammensætning ifølge krav 9, hvor den immunogene sammensætning omfatter QS21.
11. Immunogen sammensætning ifølge enten krav 9 eller 10, hvor den immunogene sammensætning omfatter 3-de-O-acyleret monophosphoryl-lipid-A.
12. Immunogen sammensætning ifølge et hvilket som helst af krav 1 til 11, hvor osmolaliteten er 250 til 750 mOsm/kg.
13. Immunogen sammensætning ifølge et hvilket som helst af krav 1 til 12, hvor sammensætningen er tilvejebragt som en enhedsdosis af mellem 50 µl og 1 ml.
14. Immunogen sammensætning ifølge krav 13, hvor enhedsdosen indeholder 5 til 50 µg af M72-relateret protein.
15. Immunogen sammensætning ifølge krav 13 eller 14, hvor enhedsdosen er en human dosis og indeholder mellem 1 og 100 µg af 3D-MPL.
16. Immunogen sammensætning ifølge et hvilket som helst af krav 13 til 15, hvor enhedsdosen er en human dosis og indeholder mellem 1 og 100 µg af QS21.
17. Immunogen sammensætning ifølge et hvilket som helst af krav 1 til 16, hvor pH af sammensætningen er i området 7,5 til 8,5.
18. Immunogen sammensætning ifølge et hvilket som helst af krav 1 til 17, hvor det M72-relaterede antigen omfatter aminosyresekvensen ifølge SEQ ID No: 3.

19. Immunogen sammensætning ifølge et hvilket som helst af krav 1 til 17, hvor det M72-relaterede antigen består af en sekvens, der har mindst 90% identitet med SEQ ID No: 1.

20. Immunogen sammensætning ifølge krav 19, hvor det M72-relaterede antigen består af aminosyresekvensen ifølge SEQ ID No: 3.

21. Immunogen sammensætning ifølge et hvilket som helst af krav 1 til 20, til anvendelse i medicin.

22. Immunogen sammensætning ifølge et hvilket som helst af krav 1 til 21, til anvendelse i en fremgangsmåde til forebyggelse, behandling eller bedring af infektion af mycobakterier.

23. Immunogen sammensætning til anvendelse ifølge enten krav 21 eller 22, til indgivelse til et menneske.

DRAWINGS

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5





Figure 6a






Figure 6b

Figure 6c




Figure 6d




Figure 7




Figure 8




Figure 9




Figure 10




Figure 11




Figure 12




Figure 13

