



US 20090069419A1

(19) **United States**(12) **Patent Application Publication**
Jandeleit et al.(10) **Pub. No.: US 2009/0069419 A1**(43) **Pub. Date: Mar. 12, 2009**(54) **MASKED CARBOXYLATE NEOPENTYL
SULFONYL ESTER CYCLIZATION RELEASE
PRODRUGS OF ACAMPROSATE,
COMPOSITIONS THEREOF, AND METHODS
OF USE**(76) Inventors: **Bernd Jandeleit**, Menlo Park, CA
(US); **Yunxiao Li**, Sunnyvale, CA
(US); **Mark A. Gallop**, Santa Clara,
CA (US); **Noa Zerangue**, Belmont,
CA (US); **Peter A. Virsik**, Portola
Valley, CA (US); **Wolf-Nicolas
Fischer**, Sunnyvale, CA (US)

Correspondence Address:

DORSEY & WHITNEY, LLP
INTELLECTUAL PROPERTY DEPARTMENT
370 SEVENTEENTH STREET, SUITE 4700
DENVER, CO 80202-5647 (US)(21) Appl. No.: **12/204,961**(22) Filed: **Sep. 5, 2008****Related U.S. Application Data**(60) Provisional application No. 60/970,924, filed on Sep.
7, 2007, provisional application No. 61/061,059, filed
on Jun. 12, 2008.**Publication Classification**(51) **Int. Cl.****A61K 31/255** (2006.01)**C07C 233/01** (2006.01)**A61K 31/164** (2006.01)**A61P 25/28** (2006.01)**A61P 25/22** (2006.01)**A61P 25/18** (2006.01)**C07C 303/02** (2006.01)(52) **U.S. Cl. 514/517; 564/192; 558/49; 514/625**

(57)

ABSTRACT

Masked carboxylate neopentyl sulfonyl ester prodrugs of acamprosate, pharmaceutical compositions comprising such prodrugs, and methods of using such prodrugs and compositions thereof for treating diseases are disclosed. In particular, acamprosate prodrugs exhibiting enhanced oral bioavailability and methods of using acamprosate prodrugs to treat neurodegenerative disorders, psychotic disorders, mood disorders, anxiety disorders, somatoform disorders, movement disorders, substance abuse disorders, binge eating disorder, cortical spreading depression related disorders, tinnitus, sleeping disorders, multiple sclerosis, and pain are disclosed.

**MASKED CARBOXYLATE NEOPENTYL
SULFONYL ESTER CYCLIZATION RELEASE
PRODRUGS OF ACAMPROSATE,
COMPOSITIONS THEREOF, AND METHODS
OF USE**

[0001] This application claims the benefit under 35 U.S.C. §119(e) of U.S. Provisional Application Ser. Nos. 60/970,924 filed Sep. 7, 2007, and 61/061,059 filed Jun. 12, 2008, each of which is incorporated by reference in its entirety.

FIELD


[0002] Disclosed herein are masked carboxylate neopentyl sulfonyl ester prodrugs of acamprosate that exhibit enhanced oral bioavailability, pharmaceutical compositions comprising such prodrugs, and methods of using such prodrugs and compositions thereof for treating diseases. In particular, acamprosate prodrugs exhibiting enhanced oral bioavailability and methods of using acamprosate prodrugs to treat neurodegenerative disorders, psychotic disorders, mood disorders, anxiety disorders, somatoform disorders, movement disorders, substance abuse disorders, binge eating disorder, cortical spreading depression related disorders, sleeping disorders, tinnitus, multiple sclerosis, and pain.

BACKGROUND

[0003] Prodrugs are derivatized forms of drugs that following administration are converted or metabolized to an active form of the drug in vivo. Prodrugs are used to modify one or more aspects of the pharmacokinetics of a drug in a manner that enhances the therapeutic efficacy of a drug. For example, prodrugs are often used to enhance the oral bioavailability of a drug. To be therapeutically effective, drugs exhibiting poor oral bioavailability may require frequent dosing, large administered doses, or may need to be administered by other than oral routes, such as intravenously. In particular, many drugs with sulfonic acid groups exhibit poor oral bioavailability.

[0004] Intramolecular cyclization prodrug strategies have been used to modify the pharmacokinetics of drugs (Bundgaard in "A Textbook of Drug Design and Development," Krogsgaard-Larsen and Bundgaard Eds., Harwood Academic, Philadelphia, 1991, pp. 113-192; Bungaard and Nielsen, U.S. Pat. No. 5,073,641; Santos et al., *Bionorganic & Medicinal Chemistry Letters*, 2005, 15, 1595-1598; Papot et al., *Curr Med Chem—Anti-Cancer Agents*, 2002, 2, 155-185; and Shan et al., *J Pharm Sciences* 1997, 86(7), 765-767). Intramolecular cyclization release prodrug strategies have been applied to drugs containing sulfonic acid functional groups. Prodrugs comprising a substituted neopentyl sulfonate ester derivative in which the neopentyl group is removed in vivo by unmasking a nucleophilic heteroatom bonded to a substituted neopentyl moiety followed by intramolecular cyclization to generate the parent drug in the sulfonic acid or sulfonic acid salt form have been described (Roberts and Patch, U.S. Pat. No. 5,596,095; and Roberts et al., *Tetrahedron Lett* 1997, 38(3), 355-358). In such prodrugs the nucleophilic heteroatom can be nitrogen or oxygen and the nitrogen or oxygen nucleophile can be masked with any amine or alcohol protecting group, respectively, capable of being deprotected in vivo. Roberts and Patch also disclose that the masked nucleophilic group can be a carboxylic ester, e.g., —OCOR where R can be aryl, substituted aryl, heteroaryl, C₁₋₈ alkyl, arylalkyl, or heteroarylalkyl. However, Roberts and Patch do not provide biological or pharmacological data to indicate which if any of the substituted neopentyl sulfonate esters release the prodrug in vivo and would therefore be useful for enhancing the oral bioavailability of the corresponding drug.

[0005] 3-(Acetylamino)propylsulfonic acid (also referred to as N-acetylhomotaurine), acamprosate,



is a derivative of homotaurine, a naturally occurring structural analog of γ -aminobutyric acid (GABA) that appears to affect multiple receptors in the central nervous system (CNS). As an antiglutamatergic agent, acamprosate is believed to exert a neuropharmacological effect as an antagonist of N-methyl-D-aspartate (NMDA) receptors. The mechanism of action is believed to include blocking of the Ca²⁺ channel to slow Ca²⁺ influx and reduce the expression of c-fos, leading to changes in messenger RNA transcription and the concomitant modification to the subunit composition of NMDA receptors in selected brain regions (Zornoza et al., *CNS Drug Reviews*, 2003, 9(4), 359-374; and Rammes et al., *Neuropharmacology* 2001, 40, 749-760). In addition, acamprosate may block GABA_B receptors (Daost, et al., *Pharmacol Biochem Behav*, 1992, 41, 669-74; and Johnson et al., *Psychopharmacology* 2000, 149, 327-344). Similar mechanisms are believed to be associated with the activity of other glutamate modulators such as riluzole, N-acetylcysteine, β -lactams, amantadine, lamictal, memantine, neramexane, remacemide, ifenprodil, and dextromethorphan.

[0006] Other diseases or disorders known to be associated with modulation of NMDA activity and for which modulators of NMDA receptor activity are clinically useful include psychotic disorders such as schizophrenia and schizoaffective disorder; mood disorders such as anxiety disorders including posttraumatic stress disorder and obsessive-compulsive disorder, depression, mania, bipolar disorder; and somatoform disorders such as somatization disorder, conversion disorder, hypochondriasis, and body dysmorphic disorder; movement disorders such as Tourette's syndrome, focal dystonia, Huntington's disease, Parkinson's disease, Sydenham's chorea, systemic lupus erythematosus, drug-induced movement disorders, tardive dyskinesia, blepharospasm, tic disorder, and spasticity; substance abuse disorders such as alcohol abuse disorders, narcotic abuse disorders, and nicotine abuse disorders; cortical spreading depression related disorders such as migraine, cerebral damage, epilepsy, and cardiovascular; sleeping disorders such as sleep apnea; multiple sclerosis; and neurodegenerative disorders such as Parkinson's disease, Huntington's disease, Alzheimer's disease, and amyotrophic lateral sclerosis. Recently, acamprosate has been found to be effective in treating tinnitus, or noise originating in the ear, a common disorder (de Azevedo et al., 109th Meeting and OTO EXPO of the Am. Acad. Otolaryngology—Head and Neck Foundation, Los Angeles, Calif., Sep. 25-28, 2005; Azevedo et al, *Rev. Bras. Otorrinolaringol. Engl. Ed.*, 2005, 71, 618-623; and Azevedo et al., WO 2007/082561 A2). Acamprosate analogs (Berthelon et al., U.S. Pat. No. 6,265,437) and salt

forms of acamprosate analogs (Durlach, U.S. Pat. No. 4,355, 043) are also reported to have therapeutic potential.


[0007] There is also evidence that acamprosate may interact with excitatory glutamatergic neurotransmission in general and as an antagonist of the metabotropic glutamate receptor subtype 5 (mGluR5) in particular (De Witte et al., *CNS Drugs* 2005, 19(6), 517-37). The glutamatergic mechanism of action of acamprosate may explain the effects of acamprosate on alcohol dependence and suggests other activities such as in neuroprotection. Dysregulation of the mGluR5 receptor has been implicated in a number of diseases and mGluR5 antagonists have been shown to be effective in treating depression, pain, anxiety disorders, alcohol abuse disorders, drug abuse disorders, nicotine abuse disorders, neurodegenerative disorders such as Parkinson's disease, diabetes, schizophrenia, and gastrointestinal reflux disease.

[0008] Acamprosate is a polar molecule that lacks the requisite physicochemical characteristics for effective passive permeability across cellular membranes. Intestinal absorption of acamprosate is mainly by passive diffusion and to a lesser extent by an active transport mechanism such as via an amino acid transporter (Más-Serrano et al., *Alcohol* 2000, 4(3); and 324-330; Saivin et al., *Clin Pharmacokinet* 1998, 35, 331-345). As a consequence, the oral bioavailability of acamprosate in humans is only about 11%. The mean elimination half-life of acamprosate following intravenous infusion (15 min) is 3.2 ± 0.2 h. Efforts to enhance the gastrointestinal absorption and oral bioavailability of acamprosate include co-administrating the drug with polyglycolysed glycerides (Saslawski et al., U.S. Pat. No. 6,514,524). Acamprosate prodrugs exhibiting enhanced absorption from the lower gastrointestinal tract have the potential to increase the oral bioavailability of the drug and to facilitate administration of acamprosate using sustained release oral dosage forms.

SUMMARY

[0009] Thus, there is a need for new prodrugs of acamprosate with demonstrated enhanced oral bioavailability. In particular, masked carboxylate neopentylsulfonate ester prodrugs of acamprosate that exhibit enhanced absorption throughout the gastrointestinal tract and especially in the large intestine/colon and hence that are suitable for sustained release oral formulations, can enhance the convenience (by reducing the dose and dosing frequency), efficacy, and side effect profile of acamprosate.

[0010] In a first aspect, compounds of Formula (I) are provided:



[0011] or a pharmaceutically acceptable salt thereof; wherein:

[0012] n is chosen from 0, 1, 2, and 3;

[0013] R¹ is chosen from C₁₋₈ alkyl, substituted C₁₋₈ alkyl, C₆₋₁₀ aryl, substituted C₆₋₁₀ aryl, C₃₋₈ cycloalkyl, substituted C₃₋₈ cycloalkyl, C₇₋₁₈ arylalkyl, substituted C₇₋₁₈ arylalkyl, C₄₋₁₆ cycloalkylalkyl, substituted C₄₋₁₆ cycloalkylalkyl, C₁₋₈ heteroalkyl, substituted C₁₋₈ het-

eroalkyl, C₅₋₁₀ heteroaryl, substituted C₅₋₁₀ heteroaryl, C₃₋₈ heterocycloalkyl, substituted C₃₋₈ heterocycloalkyl, C₆₋₁₈ heteroarylalkyl, substituted C₆₋₁₈ heteroarylalkyl, C₄₋₁₆ heterocycloalkylalkyl, substituted C₄₋₁₆ heterocycloalkylalkyl, -C(R⁶)₂OC(O)R⁵, -C(R⁶)₂OC(O)OR⁵, (CH₂)_rN(R⁶)₂, and -(CH₂)_rC(O)N(R⁶)₂, wherein:

[0014] r is chosen from 1, 2 and 3;

[0015] each R⁵ is independently chosen from C₁₋₆ alkyl, substituted C₁₋₆ alkyl, C₆₋₈ aryl, substituted C₆₋₈ aryl, C₃₋₈ cycloalkyl, substituted C₃₋₈ cycloalkyl, C₁₋₆ heteroalkyl, substituted C₁₋₆ heteroalkyl, C₅₋₈ heteroaryl, substituted C₅₋₈ heteroaryl, C₃₋₈ heterocycloalkyl, and substituted C₃₋₈ heterocycloalkyl; and

[0016] each R⁶ is independently chosen from hydrogen, C₁₋₈ alkyl, substituted C₁₋₈ alkyl, C₆₋₈ aryl, substituted C₆₋₈ aryl, C₃₋₈ cycloalkyl, substituted C₃₋₈ cycloalkyl, and -COOR³³; wherein R³³ is chosen from hydrogen and C₁₋₈ alkyl;

[0017] R² and R³ are independently chosen from C₁₋₄ alkyl, substituted C₁₋₄ alkyl, C₁₋₄ alkoxy, substituted C₁₋₄ alkoxy, -NH₂, -NHC(O)R³⁴, -NHR³⁴, and -N(R³⁴)₂; or R² and R³ together with the carbon to which they are bonded form a ring chosen from a C₃₋₈ cycloalkyl, substituted C₃₋₈ cycloalkyl, C₃₋₈ heterocycloalkyl, and substituted C₃₋₈ heterocycloalkyl ring; and

[0018] each R⁴ is independently chosen from hydrogen, halogen, -OR⁷, -CN, -CF₃, =O, -NO₂, -COOR³³, C₁₋₈ alkyl, substituted C₁₋₈ alkyl, C₆₋₁₀ aryl, substituted C₆₋₁₀ aryl, C₃₋₈ cycloalkyl, substituted C₃₋₈ cycloalkyl, C₇₋₁₈ arylalkyl, substituted C₇₋₁₈ arylalkyl, C₄₋₁₆ cycloalkylalkyl, substituted C₄₋₁₆ cycloalkylalkyl, C₁₋₈ heteroalkyl, substituted C₁₋₈ heteroalkyl, C₅₋₁₀ heteroaryl, substituted C₅₋₁₀ heteroaryl, C₃₋₈ heterocycloalkyl, substituted C₃₋₈ heterocycloalkyl, C₆₋₁₈ heteroarylalkyl, substituted C₆₋₁₈ heteroarylalkyl, C₄₋₁₆ heterocycloalkylalkyl, substituted C₄₋₁₆ heterocycloalkylalkyl, -NH₂, -NHC(O)R³⁴, -NHR³⁴, and -NR³⁴₂; wherein

[0019] R⁷ is chosen from hydrogen, C₁₋₈ alkyl, substituted C₁₋₈ alkyl, C₆₋₁₀ aryl, substituted C₆₋₁₀ aryl, C₃₋₈ cycloalkyl, substituted C₃₋₈ cycloalkyl, -C(O)R⁸, -C(O)OR⁸, -C(O)N(R⁹)₂, -C(O)SR⁸, -C(R⁹)₂OC(O)R⁸, -C(R⁹)₂OC(O)OR⁸, -P(O)(OH)₂, and -SO₂OH; wherein:

[0020] each R⁸ is independently chosen from C₁₋₆ alkyl, substituted C₁₋₆ alkyl, C₆₋₈ aryl, substituted C₆₋₈ aryl, C₃₋₈ cycloalkyl, substituted C₃₋₈ cycloalkyl, C₁₋₆ heteroalkyl, substituted C₁₋₆ heteroalkyl, C₅₋₈ heteroaryl, substituted C₅₋₈ heteroaryl, C₃₋₈ heterocycloalkyl, and substituted C₃₋₈ heterocycloalkyl; and

[0021] each R⁹ is independently chosen from hydrogen, C₁₋₈ alkyl, substituted C₁₋₈ alkyl, C₆₋₈ aryl, substituted C₆₋₈ aryl, C₃₋₈ cycloalkyl, substituted C₃₋₈ cycloalkyl, and -COOR³⁵; wherein R³⁵ is chosen from hydrogen and C₁₋₈ alkyl;


[0022] R³³ is chosen from hydrogen and C₁₋₄ alkyl;

[0023] each R³⁴ is independently chosen from hydrogen, C₁₋₈ alkyl, substituted C₁₋₈ alkyl, C₁₋₈ alkoxy, substituted C₁₋₈ alkoxy, C₆₋₁₀ aryl, substituted C₆₋₁₀ aryl, C₆₋₁₀ aryloxy, substituted C₆₋₁₀ aryloxy, C₃₋₈ cycloalkyl, substituted C₃₋₈ cycloalkyl, C₃₋₈ cycloalkoxy, substituted C₃₋₈ cycloalkoxy, C₇₋₁₈ ary-

lalkyl, substituted C₇₋₁₈ arylalkyl, C₄₋₁₆ cycloalkylalkyl, substituted C₄₋₁₆ cycloalkylalkyl, C₁₋₈ heteroalkyl, substituted C₁₋₈ heteroalkyl, C₅₋₁₀ heteroaryl, substituted C₅₋₁₀ heteroaryl, C₃₋₈ heterocycloalkyl, substituted C₃₋₈ heterocycloalkyl, C₆₋₁₈ heteroarylalkyl, substituted C₆₋₁₈ heteroarylalkyl, C₄₋₁₆ heterocycloalkylalkyl, and substituted C₄₋₁₆ heterocycloalkylalkyl; and

[0024] with the proviso that when n is 1, and each of R² and R³ is methyl; then R⁴ is not chosen from —OR⁷, —COOR³³, and substituted heteroalkyl.

[0025] In a second aspect, compounds of Formula (II) are provided.



[0026] or a pharmaceutically acceptable salt thereof; wherein:

[0027] q is chosen from 0, 1, 2, and 3;

[0028] R²² and R²³ are independently chosen from C₁₋₄ alkyl, substituted C₁₋₄ alkyl, C₁₋₄ alkoxy, and substituted C₁₋₄ alkoxy; or R²² and R²³ together with the carbon to which they are bonded form a ring chosen from a C₃₋₈ cycloalkyl, substituted C₃₋₈ cycloalkyl, C₃₋₈ heterocycloalkyl, and substituted C₃₋₈ heterocycloalkyl ring; and

[0029] each R²⁴ is independently chosen from hydrogen, halogen, —OR⁷, —CN, —CF₃, =O, —NO₂, —COOR³³, C₁₋₈ alkyl, substituted C₁₋₈ alkyl, C₆₋₁₀ aryl, substituted C₆₋₁₀ aryl, C₃₋₈ cycloalkyl, substituted C₃₋₈ cycloalkyl, C₇₋₁₈ arylalkyl, substituted C₇₋₁₈ arylalkyl, C₄₋₁₆ cycloalkylalkyl, substituted C₄₋₁₆ cycloalkylalkyl, C₁₋₈ heteroalkyl, substituted C₁₋₈ heteroalkyl, C₅₋₁₀ heteroaryl, substituted C₅₋₁₀ heteroaryl, C₃₋₈ heterocycloalkyl, substituted C₃₋₈ heterocycloalkyl, C₆₋₁₈ heteroarylalkyl, substituted C₆₋₁₈ heteroarylalkyl, C₄₋₁₆ heterocycloalkylalkyl, substituted C₄₋₁₆ heterocycloalkylalkyl, —NH₂, —NHC(O)R³⁴, and —N(R³⁴)₂; wherein:

[0030] R⁷ is chosen from hydrogen, C₁₋₈ alkyl, substituted C₁₋₈ alkyl, C₆₋₁₀ aryl, substituted C₆₋₁₀ aryl, C₃₋₈ cycloalkyl, substituted C₃₋₈ cycloalkyl, —C(O)R⁸, —C(O)OR⁸, —C(O)N(R⁹)₂, —C(O)SR⁸, —C(R⁹)₂OC(O)R⁸, and —C(R⁹)₂OC(O)OR⁸; wherein:

[0031] each R⁸ is independently chosen from C₁₋₆ alkyl, substituted C₁₋₆ alkyl, C₆₋₈ aryl, substituted C₆₋₈ aryl, C₃₋₈ cycloalkyl, substituted C₃₋₈ cycloalkyl, C₁₋₆ heteroalkyl, substituted C₁₋₆ heteroalkyl, C₅₋₈ heteroaryl, substituted C₅₋₈ heteroaryl, C₃₋₈ heterocycloalkyl, and substituted C₃₋₈ heterocycloalkyl;

[0032] each R⁹ is independently chosen from hydrogen, C₁₋₈ alkyl, substituted C₁₋₈ alkyl, C₆₋₈ aryl, substituted C₆₋₈ aryl, C₃₋₈ cycloalkyl, substituted C₃₋₈ cycloalkyl, and —COOR³⁵; wherein R³⁵ is chosen from hydrogen and C₁₋₈ alkyl;

[0033] R³³ is chosen from hydrogen and C₁₋₄ alkyl;

[0034] each R³⁴ is independently chosen from hydrogen, C₁₋₈ alkyl, substituted C₁₋₈ alkyl, C₁₋₈ alkoxy,

substituted C₁₋₈ alkoxy, C₆₋₁₀ aryl, substituted C₆₋₁₀ aryl, C₆₋₁₀ aryloxy, substituted C₆₋₁₀ aryloxy, C₃₋₈ cycloalkyl, substituted C₃₋₈ cycloalkyl, C₃₋₈ cycloalkoxy, substituted C₃₋₈ cycloalkoxy, C₇₋₁₈ arylalkyl, substituted C₇₋₁₈ arylalkyl, C₄₋₁₆ cycloalkylalkyl, substituted C₄₋₁₆ cycloalkylalkyl, C₁₋₈ heteroalkyl, substituted C₁₋₈ heteroalkyl, C₅₋₁₀ heteroaryl, substituted C₅₋₁₀ heteroaryl, C₃₋₈ heterocycloalkyl, substituted C₃₋₈ heterocycloalkyl, C₆₋₁₈ heteroarylalkyl, substituted C₆₋₁₈ heteroarylalkyl, C₄₋₁₆ heterocycloalkylalkyl, and substituted C₄₋₁₆ heterocycloalkylalkyl; and

[0035] with the proviso that when n is 1, and each of R²² and R²³ is methyl; then R²⁴ is not —OH.

[0036] In a third aspect, pharmaceutical compositions are provided comprising at least one pharmaceutically acceptable excipient and at least one compound of Formula (I) or a pharmaceutically acceptable salt thereof.

[0037] In a fourth aspect, methods of treating a disease in a patient are provided comprising administering to a patient in need of such treatment a therapeutically effective amount of a compound of Formula (I) or a pharmaceutically acceptable salt thereof. In certain embodiments, the disease is chosen from a neurodegenerative disorder, a psychotic disorder, a mood disorder, an anxiety disorder, a somatoform disorder, a movement disorder, a substance abuse disorder, binge eating disorder, a cortical spreading depression related disorder, tinnitus, a sleeping disorder, multiple sclerosis, and pain.

DETAILED DESCRIPTION

Definitions

[0038] A dash (“—”) that is not between two letters or symbols is used to indicate a point of bonding to a moiety or substituent. For example, —CONH₂ is attached through the carbon atom.

[0039] “Alkyl” by itself or as part of another substituent refers to a saturated or unsaturated, branched, or straight-chain, monovalent hydrocarbon radical derived by the removal of one hydrogen atom from a single carbon atom of a parent alkane, alkene, or alkyne. Examples of alkyl groups include, but are not limited to, methyl; ethyls such as ethanyl, ethenyl, and ethynyl; propyls such as propan-1-yl, propan-2-yl, prop-1-en-1-yl, prop-1-en-2-yl, prop-2-en-1-yl (allyl), prop-1-yn-1-yl, prop-2-yn-1-yl, etc.; butyls such as butan-1-yl, butan-2-yl, 2-methyl-propan-1-yl, 2-methyl-propan-2-yl, but-1-en-1-yl, but-1-en-2-yl, 2-methyl-prop-1-en-1-yl, but-2-en-1-yl, but-2-en-2-yl, buta-1,3-dien-1-yl, buta-1,3-dien-2-yl, but-1-yn-1-yl, but-1-yn-3-yl, but-3-yn-1-yl, etc.; and the like.

[0040] The term “alkyl” is specifically intended to include groups having any degree or level of saturation, i.e., groups having exclusively single carbon-carbon bonds, groups having one or more double carbon-carbon bonds, groups having one or more triple carbon-carbon bonds, and groups having mixtures of single, double, and triple carbon-carbon bonds. Where a specific level of saturation is intended, the terms “alkanyl,” “alkenyl,” and “alkynyl” are used. In certain embodiments, an alkyl group can have from 1 to 20 carbon atoms, in certain embodiments, from 1 to 10 carbon atoms, in certain embodiments from 1 to 8 carbon atoms, in certain embodiments, from 1 to 6 carbon atoms, in certain embodiments from 1 to 4 carbon atoms, and in certain embodiments, from 1 to 3 carbon atoms.

[0041] “Alkoxy” by itself or as part of another substituent refers to a radical —OR^{31} where R^{31} is chosen from alkyl, heteroalkyl, cycloalkyl, heterocycloalkyl, cycloalkylalkyl, heterocycloalkylalkyl, aryl, heteroaryl, arylalkyl, and heteroarylalkyl, as defined herein. Examples of alkoxy groups include, but are not limited to, methoxy, ethoxy, propoxy, butoxy, cyclohexyloxy, and the like. In certain embodiments, an alkoxy group is C_{1-18} alkoxy, in certain embodiments, C_{1-12} alkoxy, in certain embodiments, C_{1-8} alkoxy, in certain embodiments, C_{1-6} alkoxy, in certain embodiments, C_{1-4} alkoxy, and in certain embodiments, C_{1-3} alkoxy.

[0042] “Aryl” by itself or as part of another substituent refers to a monovalent aromatic hydrocarbon radical derived by the removal of one hydrogen atom from a single carbon atom of a parent aromatic ring system. Aryl encompasses 5- and 6-membered carbocyclic aromatic rings, for example, benzene; bicyclic ring systems wherein at least one ring is carbocyclic and aromatic, for example, naphthalene, indane, and tetralin; and tricyclic ring systems wherein at least one ring is carbocyclic and aromatic, for example, fluorene. Aryl encompasses multiple ring systems having at least one carbocyclic aromatic ring fused to at least one carbocyclic aromatic ring, cycloalkyl ring, or heterocycloalkyl ring. For example, aryl includes 5- and 6-membered carbocyclic aromatic rings fused to a 5- to 7-membered heterocycloalkyl ring containing one or more heteroatoms chosen from N, O, and S. For such fused, bicyclic ring systems wherein only one of the rings is a carbocyclic aromatic ring, the point of attachment may be at the carbocyclic aromatic ring or the heterocycloalkyl ring. Examples of aryl groups include, but are not limited to, groups derived from aceanthrylene, acenaphthylene, acephenanthrylene, anthracene, azulene, benzene, chrysene, coronene, fluoranthene, fluorene, hexacene, hexaphene, hexalene, as-indacene, s-indacene, indane, indene, naphthalene, octacene, octaphene, octalene, ovalene, penta-2,4-diene, pentacene, pentalene, pentaphene, perylene, phenalene, phenanthrene, picene, pleiadene, pyrene, pyranthrene, rubicene, triphenylene, trinaphthalene, and the like. In certain embodiments, an aryl group can have from 6 to 20 carbon atoms (C_{6-20}), from 6 to 12 carbon atoms (C_{6-12}), and in certain embodiments, from 6 to 10 carbon atoms (C_{6-10}).

[0043] “Arylalkyl” by itself or as part of another substituent refers to an acyclic alkyl radical in which one of the hydrogen atoms bonded to a carbon atom, typically a terminal or sp^3 carbon atom, is replaced with an aryl group. Examples of arylalkyl groups include, but are not limited to, benzyl, 2-phenylethan-1-yl, 2-phenyl ethen-1-yl, naphthylmethyl, 2-naphthylethan-1-yl, 2-naphthylethen-1-yl, naphthobenzyl, 2-naphthophenylethan-1-yl and the like. Where specific alkyl moieties are intended, the nomenclature arylalkanyl, arylalkenyl, or arylalkynyl is used. In certain embodiments, an arylalkyl group is C_{7-30} arylalkyl, e.g., the alkanyl, alkenyl or alkynyl moiety of the arylalkyl group is C_{1-10} and the aryl moiety is C_{7-20} , in certain embodiments, an arylalkyl group is C_{6-18} arylalkyl, e.g., the alkanyl, alkenyl or alkynyl moiety of the arylalkyl group is C_{1-8} and the aryl moiety is C_{6-10} .

[0044] “AUC” is the area under a curve representing the concentration of a compound or metabolite thereof in a biological fluid in a patient as a function of time following administration of the compound to the patient. In certain embodiments provided by the present disclosure, the compound is a prodrug of Formula (I) the drug is acamprosate. Examples of biological fluids include plasma, blood, and cerebrospinal fluid. The AUC may be determined by measur-

ing the concentration of a compound or metabolite thereof in a biological fluid such as the plasma or blood using methods such as liquid chromatography-tandem mass spectrometry (LC/MS/MS), at various time intervals, and calculating the area under the plasma concentration-versus-time curve. Suitable methods for calculating the AUC from a drug concentration-versus-time curve are well known in the art. As relevant to the present disclosure, an AUC for acamprosate or metabolite thereof may be determined by measuring over time the concentration of acamprosate or metabolite thereof in the plasma, blood, or other biological fluid or tissue of a patient following administration of a corresponding prodrug of Formula (I) to the patient.

[0045] “Bioavailability” refers to the rate and amount of a drug that reaches the systemic circulation of a patient following administration of the drug or prodrug thereof to the patient and can be determined by evaluating, for example, the plasma or blood concentration-versus-time profile for a drug. Parameters useful in characterizing a plasma or blood concentration-versus-time curve include the area under the curve (AUC), the time to maximum concentration (T_{max}), and the maximum drug concentration (C_{max}), where C_{max} is the maximum concentration of a drug in the plasma or blood of a patient following administration of a dose of the drug or form of drug to the patient, and T_{max} is the time to the maximum concentration (C_{max}) of a drug in the plasma or blood of a patient following administration of a dose of the drug or form of drug to the patient.

[0046] “ C_{max} ” is the maximum concentration of a drug in the plasma or blood of a patient following administration of a dose of the drug or prodrug to the patient.

[0047] “ T_{max} ” is the time to the maximum (peak) concentration (C_{max}) of a drug in the plasma or blood of a patient following administration of a dose of the drug or prodrug to the patient.

[0048] “Compounds” of Formula (I)-(II) disclosed herein include any specific compounds within these formulae. Compounds may be identified either by their chemical structure and/or chemical name. When the chemical structure and chemical name conflict, the chemical structure is determinative of the identity of the compound. The compounds described herein may comprise one or more chiral centers and/or double bonds and therefore may exist as stereoisomers such as double-bond isomers (i.e., geometric isomers), enantiomers, or diastereomers. Accordingly, any chemical structures within the scope of the specification depicted, in whole or in part, with a relative configuration encompass all possible enantiomers and stereoisomers of the illustrated compounds including the stereoisomerically pure form (e.g., geometrically pure, enantiomerically pure, or diastereomerically pure) and enantiomeric and stereoisomeric mixtures. Enantiomeric and stereoisomeric mixtures may be resolved into their component enantiomers or stereoisomers using separation techniques or chiral synthesis techniques well known to those skilled in the art.

[0049] Compounds of Formula (I)-(II) include optical isomers of compounds of Formula (I)-(II), racemates thereof, and other mixtures thereof. In such embodiments, the single enantiomers or diastereomers, i.e., optically active forms, can be obtained by asymmetric synthesis or by resolution of the racemates. Resolution of the racemates may be accomplished, for example, by conventional methods such as crystallization in the presence of a resolving agent, or chromatography, using, for example, a chiral high-pressure liquid

chromatography (HPLC) column. In addition, compounds of Formula (I)-(II) include Z- and E-forms (or cis- and trans-forms) of compounds with double bonds.

[0050] Compounds of Formula (I)-(II) may also exist in several tautomeric forms including the enol form, the keto form, and mixtures thereof. Accordingly, the chemical structures depicted herein encompass all possible tautomeric forms of the illustrated compounds. Compounds of Formula (I)-(II) also include isotopically labeled compounds where one or more atoms have an atomic mass different from the atomic mass conventionally found in nature. Examples of isotopes that may be incorporated into the compounds disclosed herein include, but are not limited to, ^2H , ^3H , ^{11}C , ^{13}C , ^{14}C , ^{15}N , ^{18}O , ^{17}O , etc. Compounds as referred to herein may be free acids, salts, hydrates, solvates, or N-oxides. Thus, when reference is made to compounds of the present disclosure, such as compounds of Formula (I)-(II), it is understood that a compound also implicitly refers to salts, solvates, hydrates, and combinations of any of the foregoing. Certain compounds may exist in multiple crystalline, cocrystalline, or amorphous forms. Compounds of Formula (I)-(II) include pharmaceutically acceptable solvates of a free acid or salt form of any of the foregoing, hydrates of a free acid or salt form of any of the foregoing, as well as crystalline forms of any of the foregoing.

[0051] Compounds of Formula (I)-(II) also include solvates. The term “solvate” refers to a molecular complex of a compound with one or more solvent molecules in a stoichiometric or non-stoichiometric amount. Such solvent molecules are those commonly used in the pharmaceutical art, which are known to be innocuous to a patient, e.g., water, ethanol, and the like. A molecular complex of a compound or moiety of a compound and a solvent can be stabilized by non-covalent intra-molecular forces such as, for example, electrostatic forces, van der Waals forces, or hydrogen bonds. The term “hydrate” refers to a solvate in which the one or more solvent molecules is water.

[0052] Further, when partial structures of the compounds are illustrated, an asterisk (*) indicates the point of bonding of the partial structure to the rest of the molecule.

[0053] “Cycloalkyl” by itself or as part of another substituent refers to a saturated or partially unsaturated cyclic alkyl radical. Where a specific level of saturation is intended, the nomenclature “cycloalkanyl” or “cycloalkenyl” is used. Examples of cycloalkyl groups include groups derived from cyclopropane, cyclobutane, cyclopentane, cyclohexane, and the like. In certain embodiments, a cycloalkyl group is C_{3-15} cycloalkyl, C_{3-12} cycloalkyl, C_{3-10} cycloalkyl or in certain embodiments, C_{3-8} cycloalkyl. Cycloalkyl includes nonaromatic fused ring systems.

[0054] “Cycloalkylalkyl” by itself or as part of another substituent refers to an acyclic alkyl radical in which one of the hydrogen atoms bonded to a carbon atom, typically a terminal or sp^3 carbon atom, is replaced with a cycloalkyl group. Where specific alkyl moieties are intended, the nomenclature cycloalkylalkanyl, cycloalkylalkenyl, or cycloalkylalkynyl is used. In certain embodiments, a cycloalkylalkyl group is C_{7-30} cycloalkylalkyl, e.g., the alkanyl, alkenyl, or alkynyl moiety of the cycloalkylalkyl group is C_{1-10} and the cycloalkyl moiety is C_{6-20} , and in certain embodiments, a cycloalkylalkyl group is C_{7-20} cycloalkylalkyl, e.g., the alkanyl, alkenyl, or alkynyl moiety of the cycloalkylalkyl group is C_{1-8} and the cycloalkyl moiety is

C_{4-20} or C_{6-12} . In certain embodiments, a cycloalkylalkyl group is C_{4-18} cycloalkylalkyl.

[0055] “Disease” refers to a disease, disorder, condition, or symptom of any of the foregoing.

[0056] “Drug” as defined under 21 U.S.C. §321 (g)(1) means “(A) articles recognized in the official United States Pharmacopoeia, official Homeopathic Pharmacopoeia of the United States, or official National Formulary, or any supplement to any of them; and (B) articles intended for use in the diagnosis, cure, mitigation, treatment, or prevention of disease in man or other animals; and (C) articles (other than food) intended to affect the structure or any function of the body of man or other animals . . .”

[0057] “Halogen” refers to a fluoro, chloro, bromo, or iodo group. In certain embodiments, halogen is fluoro, and in certain embodiments, halogen is chloro.

[0058] “Heteroalkyl” by itself or as part of another substituent refer to an alkyl group in which one or more of the carbon atoms (and certain associated hydrogen atoms) are independently replaced with the same or different heteroatomic groups. Examples of heteroatomic groups include, but are not limited to, $-\text{O}-$, $-\text{S}-$, $-\text{O}-\text{O}-$, $-\text{S}-\text{S}-$, $-\text{O}-\text{S}-$, $-\text{NR}^{37}$, $-\text{N}-\text{N}-$, $-\text{N}=\text{N}-$, $-\text{N}=\text{N}-\text{NR}^{37}$, $-\text{PR}^{37}$, $-\text{P}(\text{O})_2-$, $-\text{POR}^{37}$, $-\text{O}-\text{P}(\text{O})_2-$, $-\text{SO}-$, $-\text{SO}_2-$, $-\text{Sn}(\text{R}^{37})_2-$, and the like, where each R^{37} is independently chosen from hydrogen, C_{1-6} alkyl, substituted C_{1-6} alkyl, C_{6-12} aryl, substituted C_{6-12} aryl, C_{7-18} arylalkyl, substituted C_{7-18} arylalkyl, C_{3-7} cycloalkyl, substituted C_{3-7} cycloalkyl, C_{3-7} heterocycloalkyl, substituted C_{3-7} heterocycloalkyl, C_{1-6} heteroalkyl, substituted C_{1-6} heteroalkyl, C_{5-12} heteroaryl, substituted C_{5-12} heteroaryl, C_{6-18} heteroarylalkyl, or substituted C_{6-18} heteroarylalkyl. Reference to, for example, a C_{1-6} heteroalkyl, means a C_{1-6} alkyl group in which at least one of the carbon atoms (and certain associated hydrogen atoms) is replaced with a heteroatom. For example C_{1-6} heteroalkyl includes groups having five carbon atoms and one heteroatom, groups having four carbon atoms and two heteroatoms, etc. In certain embodiments, each R^{37} is independently chosen from hydrogen and C_{1-3} alkyl. In certain embodiments, a heteroatomic group is chosen from $-\text{O}-$, $-\text{S}-$, $-\text{NH}-$, $-\text{N}(\text{CH}_3)-$, and $-\text{SO}_2-$.

[0059] “Heteroaryl” by itself or as part of another substituent refers to a monovalent heteroaromatic radical derived by the removal of one hydrogen atom from a single atom of a parent heteroaromatic ring system. Heteroaryl encompasses multiple ring systems having at least one heteroaromatic ring fused to at least one other ring, which can be aromatic or non-aromatic. Heteroaryl encompasses 5- to 7-membered aromatic, monocyclic rings containing one or more, for example, from 1 to 4, or in certain embodiments, from 1 to 3, heteroatoms chosen from N, O, and S, with the remaining ring atoms being carbon; and 5- to 14-membered bicyclic rings containing one or more, for example, from 1 to 4, or in certain embodiments, from 1 to 3, heteroatoms chosen from N, O, and S, with the remaining ring atoms being carbon, wherein at least one of the rings is an aromatic ring, and wherein at least one heteroatom is present in the at least one aromatic ring. For example, heteroaryl includes a 5- to 7-membered heteroaromatic ring fused to a 5- to 7-membered cycloalkyl ring. For such fused, bicyclic heteroaryl ring systems wherein only one of the rings contains one or more heteroatoms, the point of attachment may be at the heteroaromatic ring or the cycloalkyl ring. In certain embodiments, when the total num-

ber of N, S, and O atoms in the heteroaryl group exceeds one, the heteroatoms are not adjacent to one another. In certain embodiments, the total number of N, S, and O atoms in the heteroaryl group is not more than two. In certain embodiments, the total number of N, S, and O atoms in the aromatic heterocycle is not more than one. In certain embodiments, a heteroaryl group is C₅₋₁₂ heteroaryl, C₅₋₁₀ heteroaryl, and in certain embodiments, C₅₋₆ heteroaryl. The ring of a C₅₋₁₀ heteroaryl has from 4 to 9 carbon atoms, with the remainder of the atoms in the ring being heteroatoms.

[0060] Examples of heteroaryl groups include, but are not limited to, groups derived from acridine, arsinole, carbazole, β -carboline, chromane, chromene, quinoline, furan, imidazole, indazole, indole, indoline, indolizine, isobenzofuran, isochromene, isoindole, isoindoline, isoquinoline, isothiazole, isoxazole, naphthyridine, oxadiazole, oxazole, perimidine, phenanthridine, phenanthroline, phenazine, phthalazine, pteridine, purine, pyran, pyrazine, pyrazole, pyridazine, pyridine, pyrimidine, pyrrole, pyrrolizine, quinazoline, quinoline, quinolizine, quinoxaline, tetrazole, thiadiazole, thiazole, thiophene, triazole, xanthene, and the like. In certain embodiments, a heteroaryl group is from 5- to 20-membered heteroaryl, in certain embodiments from 5- to 10-membered heteroaryl, and in certain embodiments from 5- to 8- heteroaryl. In certain embodiments heteroaryl groups are those derived from thiophene, pyrrole, benzothiophene, benzofuran, indole, pyridine, quinoline, imidazole, oxazole, or pyrazine.

[0061] “Heteroarylalkyl” by itself or as part of another substituent refers to an acyclic alkyl radical in which one of the hydrogen atoms bonded to a carbon atom, is replaced with a heteroaryl group. Typically a terminal or sp³ carbon atom is the atom replaced with the heteroaryl group. Were specific alkyl moieties are intended, the nomenclature “heteroarylalkanyl,” “heteroarylalkenyl,” and “heteroarylalkynyl” is used. In certain embodiments, a heteroarylalkyl group is a 6- to 20-membered heteroarylalkyl, e.g., the alkanyl, alkenyl, or alkynyl moiety of the heteroarylalkyl is 1- to 8-membered and the heteroaryl moiety is a 5- to 12-membered heteroaryl, and in certain embodiments, 6- to 14-membered heteroarylalkyl, e.g., the alkanyl, alkenyl, or alkynyl moiety of the heteroarylalkyl is 1- to 4-membered and the heteroaryl moiety is a 5- to 12-membered heteroaryl. In certain embodiments, a heteroarylalkyl group is C₆₋₁₈ heteroarylalkyl and in certain embodiments, C₆₋₁₀ heteroarylalkyl.


[0062] “Heterocycloalkyl” by itself or as part of another substituent refers to a saturated or partially unsaturated cyclic alkyl radical in which one or more carbon atoms (and any associated hydrogen atoms) are independently replaced with the same or different heteroatom. Typical heteroatoms to replace the carbon atom(s) include, but are not limited to, N, P, O, S, Si, etc. Where a specific level of saturation is intended, the nomenclature “heterocycloalkanyl” or “heterocycloalkenyl” is used. Examples of heterocycloalkyl groups include, but are not limited to, groups derived from epoxides, azirines, thiiranes, imidazolidine, morpholine, piperazine, piperidine, pyrazolidine, pyrrolidine, quinuclidine, and the like. Heterocycloalkyl includes nonaromatic heterocycloalkyl fused ring systems. In certain embodiments, a heterocycloalkyl group is a C₃₋₁₂ heterocycloalkylalkyl, C₃₋₁₀ heterocycloalkylalkyl, and in certain embodiments C₃₋₈ heterocycloalkylalkyl.

[0063] “Heterocycloalkylalkyl” by itself or as part of another substituent refers to an acyclic alkyl radical in which one of the hydrogen atoms bonded to a carbon atom, is

replaced with a heterocycloalkyl group as defined herein. In certain embodiments, a heterocycloalkylalkyl group is a C₄₋₁₈ heterocycloalkylalkyl, C₄₋₁₂ heterocycloalkylalkyl, and in certain embodiments C₄₋₁₀ heterocycloalkylalkyl.

[0064] “Metabolic intermediate” refers to a compound that is formed in vivo by metabolism of a parent compound and that further undergoes reaction in vivo to release an active agent. Compounds of Formula (I) are protected carboxylate nucleophile prodrugs of acamprosate that are metabolized in vivo to provide the corresponding metabolic intermediates of Formula (II). Metabolic intermediates of Formula (II) undergo nucleophilic cyclization to release acamprosate and one or more reaction products. It is desirable that the reaction products or metabolites thereof not be toxic.

[0065] “Neopentyl” refers to a radical in which a methylene carbon is bonded to a carbon atom, which is bonded to three non-hydrogen substituents. Examples of non-hydrogen substituents include carbon, oxygen, nitrogen, and sulfur. In certain embodiments, each of the three non-hydrogen substituents is carbon. In certain embodiments, two of the three non-hydrogen substituents is carbon, and the third non-hydrogen substituent is chosen from oxygen and nitrogen. In certain embodiments, a neopentyl group has the structure:



where R^a and R^b are independently chosen from C₁₋₄ alkyl, substituted C₁₋₄ alkyl, C₁₋₄ alkoxy, and substituted C₁₋₄ alkoxy; or R³ and R⁴ together with the carbon to which they are bonded form a ring chosen from a C₃₋₁₀ cycloalkyl, substituted C₃₋₁₀ cycloalkyl, C₃₋₁₀ heterocycloalkyl, and substituted C₃₋₁₀ heterocycloalkyl ring; and R^c is chosen from carbon, nitrogen, and oxygen. In certain embodiments, each of R^a and R^b is methyl; and R^c is chosen from carbon, nitrogen, and oxygen. In certain embodiments, each of R^a and R^b is methyl; and R^c is carbon; in certain embodiments, nitrogen; and in certain embodiments, oxygen.

[0066] “Parent aromatic ring system” refers to an unsaturated cyclic or polycyclic ring system having a conjugated π (pi) electron system. Included within the definition of “parent aromatic ring system” are fused ring systems in which one or more of the rings are aromatic and one or more of the rings are saturated or unsaturated, such as, for example, fluorene, indane, indene, phenalene, etc. Examples of parent aromatic ring systems include, but are not limited to, acenaphthylene, acenaphthylene, acephenanthrylene, anthracene, azulene, benzene, chrysene, coronene, fluoranthene, fluorene, hexacene, hexaphene, hexalene, as-indacene, s-indacene, indane, indene, naphthalene, octacene, octaphene, octalene, ovalene, penta-2,4-diene, pentacene, pentalene, pentaphene, perylene, phenalene, phenanthrene, picene, pleiadene, pyrene, pyranthrene, rubicene, triphenylene, trinaphthalene, and the like.

[0067] “Parent heteroaromatic ring system” refers to an aromatic ring system in which one or more carbon atoms (and any associated hydrogen atoms) are independently replaced with the same or different heteroatom in such a way as to maintain the continuous π (pi)-electron system characteristic of aromatic systems and a number or out-of-plane π (pi)-electrons corresponding to the Hückel rule (4n+1). Examples of heteroatoms to replace the carbon atoms include, but are

not limited to, N, P, O, S, and Si, etc. In certain embodiments, a heteroatom is chosen from N, O, and S. Specifically included within the definition of “parent heteroaromatic ring systems” are fused ring systems in which one or more of the rings are aromatic and one or more of the rings are saturated or unsaturated, such as, for example, arsindeole, benzodioxan, benzofuran, chromane, chromene, indole, indoline, xanthene, etc. Examples of parent heteroaromatic ring systems include, but are not limited to, arsindeole, carbazole, β -carboline, chromane, chromene, cinnoline, furan, imidazole, indazole, indole, indoline, indolizine, isobenzofuran, isochromene, isoindole, isoindoline, isoquinoline, isothiazole, isoxazole, naphthyridine, oxadiazole, oxazole, perimidine, phenanthridine, phenanthroline, phenazine, phthalazine, pteridine, purine, pyran, pyrazine, pyrazole, pyridazine, pyridine, pyrimidine, pyrrole, pyrrolizine, quinazoline, quinoxaline, quinolizine, quinoxaline, tetrazole, thiadiazole, thiazole, thiophene, triazole, xanthene, and the like.

[0068] “Patient” refers to a mammal, for example, a human.

[0069] “Pharmaceutically acceptable” refers to approved or approvable by a regulatory agency of the Federal or a state government or listed in the U.S. Pharmacopoeia or other generally recognized pharmacopoeia for use in animals, and more particularly in humans.

[0070] “Pharmaceutically acceptable salt” refers to a salt of a compound, which possesses the desired pharmacological activity of the parent compound. Such salts include acid addition salts, formed with inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, and the like; or formed with organic acids such as acetic acid, propionic acid, hexanoic acid, cyclopentanepropionic acid, glycolic acid, pyruvic acid, lactic acid, malonic acid, succinic acid, malic acid, maleic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, 3-(4-hydroxybenzoyl)benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, 1,2-ethane-disulfonic acid, 2-hydroxyethanesulfonic acid, benzenesulfonic acid, 4-chlorobenzenesulfonic acid, 2-naphthalenesulfonic acid, 4-toluenesulfonic acid, camphorsulfonic acid, 4-methylbicyclo[2.2.2]-oct-2-ene-1-carboxylic acid, glucoheptonic acid, 3-phenylpropionic acid, trimethylacetic acid, tertiary butylacetic acid, lauryl sulfuric acid, gluconic acid, glutamic acid, 2-hydroxynaphthoic acid, salicylic acid, stearic acid, muconic acid, and the like; and salts formed when an acidic proton present in the parent compound is replaced by a metal ion, e.g., an alkali metal ion, an alkaline earth ion, or an aluminum ion; or coordinates with an organic base such as ethanolamine, diethanolamine, triethanolamine, N-methylglucamine, and the like. In certain embodiments, pharmaceutically acceptable addition salts include metal salts such as sodium, potassium, aluminum, calcium, magnesium and zinc salts, and ammonium salts such as isopropylamine, diethylamine, and diethanolamine salts. In certain embodiments, a pharmaceutically acceptable salt is the hydrochloride salt. In certain embodiments, a pharmaceutically acceptable salt is the sodium salt. Pharmaceutically acceptable salts may be prepared by the skilled chemist, by treating, for example, a compound of Formula (I) with an appropriate base in a suitable solvent, followed by crystallization and filtration.


[0071] “Pharmaceutically acceptable vehicle” refers to a pharmaceutically acceptable diluent, a pharmaceutically acceptable adjuvant, a pharmaceutically acceptable excipient, a pharmaceutically acceptable carrier, or a combination of any of the foregoing with which a compound provided by

the present disclosure may be administered to a patient and which does not destroy the pharmacological activity thereof and which is non-toxic when administered in doses sufficient to provide a therapeutically effective amount of the compound.

[0072] “Pharmaceutical composition” refers to at least one compound of Formula (I) and at least one pharmaceutically acceptable vehicle with which the at least one compound of Formula (I) is administered to a patient.

[0073] “Prodrug” refers to a derivative of a drug molecule that requires a transformation within the body to release the active drug. Prodrugs are frequently, although not necessarily, pharmacologically inactive until converted to the parent drug. Prodrugs may be obtained by bonding a promoiety (defined herein) typically via a functional group, to a drug. For example, referring to compounds of Formula (I), the promoiety is bonded to the drug, acamprosate, via the sulfonic acid functional group of acamprosate. Compounds of Formula (I) are prodrugs of acamprosate that can be metabolized within a patient’s body to release acamprosate.

[0074] “Promoiety” refers to a group bonded to a drug, typically to a functional group of the drug, via bond(s) that are cleavable under specified conditions of use. The bond(s) between the drug and promoiety may be cleaved by enzymatic or non-enzymatic means. Under the conditions of use, for example following administration to a patient, the bond(s) between the drug and promoiety may be cleaved to release the parent drug. The cleavage of the promoiety may proceed spontaneously, such as via a hydrolysis reaction, or it may be catalyzed or induced by another agent, such as by an enzyme, by light, by acid, or by a change of or exposure to a physical or environmental parameter, such as a change of temperature, pH, etc. The agent may be endogenous to the conditions of use, such as an enzyme present in the systemic circulation of a patient to which the prodrug is administered or the acidic conditions of the stomach or the agent may be supplied exogenously. For example, for a prodrug of Formula (I), the drug is acamprosate (1) and the promoiety has the structure:



where n, R¹, R², R³, and R⁴ are defined herein.

[0075] “Protecting group” refers to a grouping of atoms, which when attached to a reactive group in a molecule masks, reduces, or prevents that reactivity. Examples of amino protecting groups include, but are not limited to, formyl, acetyl, trifluoroacetyl, benzyl, benzyloxycarbonyl (CBZ), tert-butoxycarbonyl (Boc), trimethylsilyl (TMS), 2-trimethylsilyl-ethanesulfonyl (SES), trityl and substituted trityl groups, allyloxycarbonyl, 9-fluorenylmethyloxycarbonyl (FMOC), nitro-veratryloxycarbonyl (NVOC), and the like. Examples of hydroxy protecting groups include, but are not limited to, those in which the hydroxy group is either acylated or alkylated such as benzyl, and trityl ethers as well as alkyl ethers, tetrahydropyranyl ethers, trialkylsilyl ethers, and allyl ethers.

[0076] “Salt” refers to a chemical compound consisting of an assembly of cations and anions. Salts of a compound of the present disclosure include stoichiometric and non-stoichiometric forms of the salt. In certain embodiments, because of

its potential use in medicine, salts of a compound of Formula (I) are pharmaceutically acceptable salts.

[0077] “Substituted” refers to a group in which one or more hydrogen atoms are independently replaced with the same or different substituent group(s). Examples of substituent groups include, but are not limited to, $-M$, $-R^{60}$, $-O^-$, $=O$, $-OR^{60}$, $-SR^{60}$, $-S^-$, $=S$, $-NR^{60}R^{61}$, $=NR^{60}$, $-CF_3$, $-CN$, $-OCN$, $-SCN$, $-NO$, $-NO_2$, $=N_2$, $-N_3$, $-S(O)_2O^-$, $-S(O)_2OH$, $-S(O)_2R^{60}$, $-OS(O)_2O^-$, $-OS(O)_2R^{60}$, $-P(O)(O^{60})_2$, $-P(O)(OR^{60})(O^-)$, $-OP(O)(OR^{60})(OR^{61})$, $-C(O)R^{60}$, $-C(S)R^{60}$, $-C(O)OR^{60}$, $-C(O)NR^{60}R^{61}$, $-C(O)O^-$, $-C(S)OR^{60}$, $-NR^{62}C(O)NR^{60}R^{61}$, $-NR^{62}C(S)NR^{60}R^{61}$, $-NR^{62}C(NR^{63})NR^{60}R^{61}$, and $-C(NR^{62})NR^{60}R^{61}$ where M is halogen; R^{60} , R^{61} , R^{62} , and R^{63} are independently chosen from hydrogen, alkyl, alkoxy, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl, or R^{60} and R^{61} together with the nitrogen atom to which they are bonded form a ring chosen from a heterocycloalkyl ring. In certain embodiments, R^{60} , R^{61} , R^{62} , and R^{63} are independently chosen from hydrogen, C_{1-6} alkyl, C_{1-6} alkoxy, C_{3-12} cycloalkyl, C_{3-12} heterocycloalkyl, C_{6-12} aryl, and C_{6-12} heteroaryl. In certain embodiments, each substituent group is independently chosen from halogen, $-OH$, $-CN$, $-CF_3$, $=O$, $-NO_2$, C_{1-3} alkoxy, C_{1-3} alkyl, $-COOR^{64}$ wherein R^{64} is chosen from hydrogen and C_{1-3} alkyl, and $-N(R^{65})_2$ wherein each R^{65} is independently chosen from hydrogen and C_{1-3} alkyl. In certain embodiments, each substituent group is independently chosen from halogen, $-OH$, $-CN$, $-CF_3$, $-OCF_3$, $=O$, $-NO_2$, C_{1-6} alkoxy, C_{1-6} alkyl, $-COOR^{62}$, $-N(R^{27})_2$, and $-CON(R^{28})_2$; wherein each of R^{26} , R^{27} , and R^{28} is independently chosen from hydrogen and C_{1-6} alkyl.

[0078] In certain embodiments, each substituent group is independently chosen from halogen, $-OH$, $-CN$, $-CF_3$, $=O$, $-NO_2$, C_{1-3} alkoxy, C_{1-3} alkyl, $-COOR^{12}$ wherein R^{12} is chosen from hydrogen and C_{1-3} alkyl, and $-N(R^{12})_2$ wherein each R^{12} is independently chosen from hydrogen and C_{1-3} alkyl. In certain embodiments, each substituent group is independently chosen from halogen, $-OH$, $-CN$, $-CF_3$, $-OCF_3$, $=O$, $-NO_2$, C_{1-6} alkoxy, C_{1-6} alkyl, $-COOR^{12}$, $-N(R^{12})_2$, and $-CONR^{12}_2$; wherein each R^{12} is independently chosen from hydrogen and C_{1-6} alkyl. In certain embodiments, each substituent group is chosen from C_{1-4} alkyl, $-OH$, and $-NH_2$.

[0079] “Sustained release” refers to release of a compound from a dosage form of a pharmaceutical composition at a rate effective to achieve a therapeutic or prophylactic concentration of the compound or active metabolite thereof, in the systemic circulation of a patient over a prolonged period of time relative to that achieved by administration of an immediate release formulation of the same compound by the same route of administration. In some embodiments, release of a compound occurs over a time period of at least about 4 hours, such as at least about 8 hours, at least about 12 hours, at least about 16 hours, at least about 20 hours, and in some embodiments, at least about 24 hours.

[0080] “Treating” or “treatment” of any disease refers to arresting or ameliorating a disease or at least one of the clinical symptoms of a disease or disorder, reducing the risk of acquiring a disease or at least one of the clinical symptoms of a disease, reducing the development of a disease or at least one of the clinical symptoms of the disease or reducing the risk of developing a disease or at least one of the clinical symptoms of a disease. “Treating” or “treatment” also refers to inhibiting the disease, either physically, (e.g., stabilization

of a discernible symptom), physiologically, (e.g., stabilization of a physical parameter), or both, and to inhibiting at least one physical parameter that may or may not be discernible to the patient. In certain embodiments, “treating” or “treatment” refers to delaying the onset of the disease or at least one or more symptoms thereof in a patient which may be exposed to or predisposed to a disease or disorder even though that patient does not yet experience or display symptoms of the disease.


[0081] “Therapeutically effective amount” refers to the amount of a compound that, when administered to a subject for treating a disease, or at least one of the clinical symptoms of a disease, is sufficient to affect such treatment of the disease or symptom thereof. The “therapeutically effective amount” may vary depending, for example, on the compound, the disease and/or symptoms of the disease, severity of the disease and/or symptoms of the disease or disorder, the age, weight, and/or health of the patient to be treated, and the judgment of the prescribing physician. An appropriate amount in any given instance may be ascertained by those skilled in the art or capable of determination by routine experimentation.

[0082] “Therapeutically effective dose” refers to a dose that provides effective treatment of a disease or disorder in a patient. A therapeutically effective dose may vary from compound to compound, and from patient to patient, and may depend upon factors such as the condition of the patient and the route of delivery. A therapeutically effective dose may be determined in accordance with routine pharmacological procedures known to those skilled in the art.

[0083] Reference is now made in detail to certain embodiments of compounds, compositions, and methods. The disclosed embodiments are not intended to be limiting of the claims. To the contrary, the claims are intended to cover all alternatives, modifications, and equivalents.

Compounds

[0084] Certain embodiments provide a compound of Formula (I):



[0085] or a pharmaceutically acceptable salt thereof; wherein:

[0086] n is chosen from 0, 1, 2, and 3;

[0087] R^1 is chosen from C_{1-8} alkyl, substituted C_{1-8} alkyl, C_{6-10} aryl, substituted C_{6-10} aryl, C_{3-8} cycloalkyl, substituted C_{3-8} cycloalkyl, C_{7-18} arylalkyl, substituted C_{7-18} arylalkyl, C_{4-16} cycloalkylalkyl, substituted C_{4-16} cycloalkylalkyl, C_{1-8} heteroalkyl, substituted C_{1-8} heteroalkyl, C_{5-10} heteroaryl, substituted C_{5-10} heteroaryl, C_{3-8} heterocycloalkyl, substituted C_{3-8} heterocycloalkyl, C_{6-18} heteroarylalkyl, substituted C_{6-18} heteroarylalkyl, C_{4-16} heterocycloalkylalkyl, substituted C_{4-16} heterocycloalkylalkyl, $-C(R^6)_2OC(O)R^5$, $-C(R^6)_2OC(O)OR^5$, $-(CH_2)_nN(R^6)_2$, and $-(CH_2)_nC(O)N(R^6)_2$, wherein:

[0088] r is chosen from 1, 2 and 3;

[0089] each R^5 is independently chosen from C_{1-6} alkyl, substituted C_{1-6} alkyl, C_{6-8} aryl, substituted C_{6-8} aryl, C_{3-8} cycloalkyl, substituted C_{3-8} cycloalkyl, C_{1-6} heteroalkyl, substituted C_{1-6} heteroalkyl, C_{5-8} heteroaryl, substituted C_{5-8} heteroaryl, C_{3-8} heterocycloalkyl, and substituted C_{3-8} heterocycloalkyl; and

[0090] each R is independently chosen from hydrogen, C_{1-8} alkyl, substituted C_{1-8} alkyl, C_{6-8} aryl, substituted C_{6-8} aryl, C_{3-8} cycloalkyl, substituted C_{3-8} cycloalkyl, and $-\text{COOR}^{33}$; wherein R^{33} is chosen from hydrogen and C_{1-8} alkyl;

[0091] R^2 and R^3 are independently chosen from C_{1-4} alkyl, substituted C_{1-4} alkyl, C_{1-4} alkoxy, substituted C_{1-4} alkoxy, $-\text{NH}_2$, $-\text{NHC(O)}R^{34}$, $-\text{NHR}^{34}$, and $-\text{N(R}^{34})_2$; or R^2 and R^3 together with the carbon to which they are bonded form a ring chosen from a C_{3-8} cycloalkyl, substituted C_{3-8} cycloalkyl, C_{3-8} heterocycloalkyl, and substituted C_{3-8} heterocycloalkyl ring; and

[0092] each R^4 is independently chosen from hydrogen, halogen, $-\text{OR}^7$, $-\text{CN}$, $-\text{CF}_3$, $=\text{O}$, $-\text{NO}_2$, $-\text{COOR}^{33}$, C_{1-8} alkyl, substituted C_{1-8} alkyl, C_{6-10} aryl, substituted C_{6-10} aryl, C_{3-8} cycloalkyl, substituted C_{3-8} cycloalkyl, C_{7-18} arylalkyl, substituted C_{7-18} arylalkyl, C_{4-16} cycloalkylalkyl, substituted C_{4-16} cycloalkylalkyl, C_{1-8} heteroalkyl, substituted C_{1-8} heteroalkyl, C_{5-10} heteroaryl, substituted C_{5-10} heteroaryl, C_{3-8} heterocycloalkyl, substituted C_{3-8} heterocycloalkyl, C_{6-18} heteroarylalkyl, substituted C_{6-18} heteroarylalkyl, C_{4-16} heterocycloalkylalkyl, substituted C_{4-16} heterocycloalkylalkyl, $-\text{NH}_2$, $-\text{NHC(O)}R^{34}$, and $-\text{N(R}^{34})_2$; wherein:

[0093] R^7 is chosen from hydrogen, C_{1-8} alkyl, substituted C_{1-8} alkyl, C_{6-10} aryl, C_{6-10} substituted aryl, C_{3-8} cycloalkyl, substituted C_{3-8} cycloalkyl, $-\text{C(O)}R^8$, $-\text{C(O)}\text{OR}^8$, $-\text{C(O)}\text{N(R}^8)_2$, $-\text{C(O)}\text{SR}^8$, $-\text{C(R}^9)_2\text{OC(O)}R^8$, $-\text{C(R}^9)_2\text{OC(O)}\text{OR}^8$, $-\text{P(O)}(\text{OH})_2$, and $-\text{SO}_2\text{OH}$; wherein:

[0094] each R^8 is independently chosen from C_{1-6} alkyl, substituted C_{1-6} alkyl, C_{6-8} aryl, substituted C_{6-8} aryl, C_{3-8} cycloalkyl, substituted C_{3-8} cycloalkyl, C_{1-6} heteroalkyl, substituted C_{1-6} heteroalkyl, C_{5-8} heteroaryl, substituted C_{5-8} heteroaryl, C_{3-8} heterocycloalkyl, and substituted C_{3-8} heterocycloalkyl; and

[0095] each R^9 is independently chosen from hydrogen, C_{1-8} alkyl, substituted C_{1-8} alkyl, C_{6-8} aryl, substituted C_{6-8} aryl, C_{3-8} cycloalkyl, substituted C_{3-8} cycloalkyl, and $-\text{COOR}^{35}$; wherein R^{35} is chosen from hydrogen and C_{1-8} alkyl;

[0096] R^{33} is chosen from hydrogen and C_{1-4} alkyl;

[0097] each R^{34} is independently chosen from hydrogen, C_{1-8} alkyl, substituted C_{1-8} alkyl, C_{1-8} alkoxy, substituted C_{1-8} alkoxy, C_{6-10} aryl, substituted C_{6-10} aryl, C_{6-10} aryloxy, substituted C_{6-10} aryloxy, C_{3-8} cycloalkyl, substituted C_{3-8} cycloalkyl, C_{3-8} cycloalkoxy, substituted C_{3-8} cycloalkoxy, C_{7-18} arylalkyl, substituted C_{7-18} arylalkyl, C_{4-16} cycloalkylalkyl, substituted C_{4-16} cycloalkylalkyl, C_{1-8} heteroalkyl, substituted C_{1-8} heteroalkyl, C_{5-10} heteroaryl, substituted C_{5-10} heteroaryl, C_{3-8} heterocycloalkyl, substituted C_{3-8} heterocycloalkyl, C_{6-18} heteroarylalkyl, substituted C_{6-18} heteroarylalkyl, C_{4-16} heterocycloalkylalkyl, and substituted C_{4-16} heterocycloalkylalkyl; and

[0098] with the proviso that when n is 1, and each of R^2 and R^3 is methyl; then R^4 is not chosen from $-\text{OR}^7$, $-\text{COOR}^{33}$, and substituted heteroalkyl.

[0099] In certain embodiments of a compound of Formula (I), each substituent is independently chosen from halogen, $-\text{OH}$, $-\text{CN}$, $-\text{CF}_3$, $-\text{OCF}_3$, $=\text{O}$, $-\text{NO}_2$, C_{1-6} alkoxy, C_{1-6} alkyl, $-\text{COOR}^{30}$, $-\text{N(R}^{31})_2$, and $-\text{CON(R}^{32})_2$; wherein each R^{30} , R^{31} , and R^{32} is independently chosen from hydrogen and C_{1-6} alkyl. In certain embodiments of a compound of Formula (I), each substituent is independently chosen from C_{1-4} alkyl, C_{1-4} alkoxy, $-\text{OH}$, and $-\text{NH}_2$.

[0100] In certain embodiments of a compound of Formula (I), n is chosen from 0, 1, and 2. In certain embodiments of a compound of Formula (I), n is 0, n is 1, n is 2, and in certain embodiments, n is 3.

[0101] In certain embodiments of a compound of Formula (I), R^1 is chosen from C_{1-6} alkyl, substituted C_{1-6} alkyl, C_{3-8} cycloalkyl, substituted C_{3-8} cycloalkyl, C_{6-10} aryl, substituted C_{6-10} aryl, C_{7-12} arylalkyl, substituted C_{7-12} arylalkyl, C_{1-6} heteroalkyl, substituted C_{1-6} heteroalkyl, C_{3-8} heterocycloalkyl, substituted C_{3-8} heterocycloalkyl, C_{5-10} heteroaryl, substituted C_{5-10} heteroaryl, C_{6-12} heteroarylalkyl, and substituted C_{6-12} heteroarylalkyl. In certain embodiments of a compound of Formula (I), R^1 is chosen from C_{1-6} alkyl, C_{3-8} cycloalkyl, C_{6-10} aryl, C_{7-12} arylalkyl, C_{1-6} heteroalkyl, C_{3-8} heterocycloalkyl, C_{5-10} heteroaryl, and C_{6-12} heteroarylalkyl. In certain embodiments of a compound of Formula (I), R^1 is chosen from C_{1-6} alkyl, C_{3-8} cycloalkyl, C_{6-10} aryl, and C_{7-12} arylalkyl.

[0102] In certain embodiments of a compound of Formula (I), R^1 is chosen from $-\text{C(R}^6)_2\text{OC(O)}R^5$ and $-\text{C(R}^6)_2\text{OC(O)}\text{OR}^5$.

[0103] In certain embodiments of a compound of Formula (I), R^1 is chosen from $-\text{C(R}^6)_2\text{OC(O)}R^5$ and $-\text{C(R}^6)_2\text{OC(O)}\text{OR}^5$; R^5 is chosen from C_{1-6} alkyl, substituted C_{1-6} alkyl, C_{6-8} aryl, substituted C_{6-8} aryl, C_{3-8} cycloalkyl, substituted C_{3-8} cycloalkyl, C_{1-6} heteroalkyl, substituted C_{1-6} heteroalkyl, C_{5-8} heteroaryl, substituted C_{5-8} heteroaryl, C_{3-8} heterocycloalkyl, and substituted C_{3-8} heterocycloalkyl; and each R^6 is independently chosen from hydrogen, C_{1-6} alkyl, substituted C_{1-6} alkyl, C_{6-8} aryl, substituted C_{6-8} aryl, C_{3-8} cycloalkyl, and substituted C_{3-8} cycloalkyl.

[0104] In certain embodiments of a compound of Formula (I), R^1 is chosen from $-\text{C(R}^6)_2\text{OC(O)}R^5$ and $-\text{C(R}^6)_2\text{OC(O)}\text{OR}^5$; R^5 is chosen from C_{1-6} alkyl and C_{6-8} aryl; and each R^6 is independently chosen from hydrogen, C_{1-6} alkyl, and C_{6-8} aryl.

[0105] In certain embodiments of a compound of Formula (I) wherein R^1 is chosen from $-\text{C(R}^6)_2\text{OC(O)}R^5$ and $-\text{C(R}^6)_2\text{OC(O)}\text{OR}^5$; R^5 is chosen from methyl, ethyl, n -propyl, isopropyl, n -butyl, isobutyl, $tert$ -butyl, pentyl, cyclohexyl, phenyl, and 3-pyridyl; and one of R^6 is hydrogen and the other of R^6 is chosen from hydrogen, methyl, ethyl, n -propyl, isopropyl, cyclohexyl, and phenyl; or each of R^6 is methyl. In certain of the immediately preceding embodiments, each of R^2 and R^3 is methyl. In certain of the immediately preceding embodiments, each R^4 is hydrogen.

[0106] In certain embodiments of a compound of Formula (I), R^1 is chosen from C_{1-6} alkyl, benzyl, substituted benzyl, morpholinyl- N -ethyl, pyridyl-3-methyl, ethoxycarbonyloxyethyl, and methylbenzyl.

[0107] In certain embodiments of a compound of Formula (I), R^2 and R^3 are independently chosen from C_{1-4} alkyl, substituted C_{1-4} alkyl, C_{1-4} alkoxy, substituted C_{1-4} alkoxy, and $-NH_2$.

[0108] In certain embodiments of a compound of Formula (I), each of R^2 and R^3 is methyl.

[0109] In certain embodiments of a compound of Formula (I), R^2 is methyl; and R^3 is chosen from $-OSO_2(CH_2)_3NHC(O)CH_3$, $-NH_3^+X^-$, and $-CH_2OH$, where X is halogen.

[0110] In certain embodiments of a compound of Formula (I), each R^4 is independently chosen from hydrogen, halogen, $-CN$, $-CF_3$, $=O$, $-NO_2$, C_{1-6} alkoxy, C_{1-6} alkyl, $-COOR^{33}$, and $-N(R^{34})_2$; wherein R^{33} is chosen from hydrogen and C_{1-3} alkyl; and each R^{34} is independently chosen from hydrogen and C_{1-3} alkyl.

[0111] In certain embodiments of a compound of Formula (I), each R^4 is independently chosen from hydrogen and C_{1-4} alkyl. In certain embodiments of a compound of Formula (I), each R^4 is hydrogen.

[0112] In certain embodiments of a compound of Formula (I), R^1 is chosen from C_{1-6} alkyl, morpholinyl-N-ethyl, N,N-dimethylaminoethyl, benzyl, and methylbenzyl. In certain embodiments, R^1 is chosen from methyl, ethyl, n-propyl, isopropyl, 1-butyl, sec-butyl, tert-butyl, isobutyl, pentyl, cyclohexyl, phenyl, and benzyl.

[0113] In certain embodiments of a compound of Formula (I), R^1 is chosen from C_{1-4} alkyl, cyclohexyl, phenyl, benzyl, and cyclohexylmethyl; n is chosen from 0, 1, and 2; each R^4 is hydrogen; and each of R^2 and R^3 is methyl.

[0114] In certain embodiments of a compound of Formula (I), the stereochemistry of the carbon atom to which R^4 is bonded is of the (S)-configuration.

[0115] In certain embodiments of a compound of Formula (I), the stereochemistry of the carbon atom to which R^4 is bonded is of the (R)-configuration.

[0116] In certain embodiments, R^1 is chosen from a carboxyl-protecting group that is cleavable in vivo by enzymatic or chemical means to provide the corresponding amide metabolic intermediate. When cleaved in vivo, the amide-protecting group provides a non-toxic metabolite. Examples of carboxyl-protecting groups include 2-N-(morpholino)ethyl, choline, methyl, methoxyethyl, 9-fluorenylmethyl, methoxymethyl, methylthiomethyl, tetrahydropyranyl, tetrahydrofuranyl, methoxyethoxymethyl, 2-(trimethylsilyl)ethoxymethyl, benzyloxymethyl, pivaloyloxymethyl, phenylacetoxymethyl, triisopropylsilylmethyl, cyanomethyl, acetol, p-bromophenacyl, α -methylphenacyl, p-methoxyphenacyl, desyl, carboxamidomethyl, p-azobenzenecarboxamido-methyl, N-phthalimidomethyl, (methoxyethoxy)ethyl, 2,2,2-trichloroethyl, 2-fluoroethyl, 2-chloroethyl, 2-bromoethyl, 2-iodoethyl, 4-chlorobutyl, 5-chloropentyl, 2-(trimethylsilyl)ethyl, 2-methylthioethyl, 1,3-dithianyl-2-methyl, 2-(p-nitrophenylsulfenyl)ethyl, 2-(p-toluenesulfonyl)ethyl, 2-(2'-pyridyl)ethyl, 2-(p-methoxyphenyl)ethyl, 2-(diphenylphosphino)ethyl, 1-methyl-1-phenylethyl, 2-(4-acetyl-2-nitrophenyl)ethyl, 2-cyanoethyl, heptyl, tert-butyl, 3-methyl-3-pentyl, dicyclopropylmethyl, 2,4-dimethyl-3-pentyl, cyclopentyl, cyclohexyl, allyl, methallyl, 2-methylbut-3-en-2-yl, 3-methylbut-2-(prenyl), 3-buten-1-yl, 4-(trimethylsilyl)-2-buten-1-yl, cinnamyl, α -methylcinnamyl, propargyl, phenyl, 2,6-dimethylphenyl, 2,6-diisopropylphenyl, 2,6-di-tert-butyl-4-methylphenyl, 2,6-di-tert-butyl-4-methoxyphenyl, p-(methylthio)phenyl, pentafluorophenyl, benzyl, triphenylmethyl, diphenylmethyl, bis(o-nitrophenyl)

methyl, 9-anthrylmethyl, 2-(9,10-dioxo)anthrylmethyl, 5-dibenzosuberyl, 1-pyrenylmethyl, 2-(trifluoromethyl)-6-chromonylmethyl, 2,4,6-trimethylbenzyl, p-bromobenzyl, o-nitrobenzyl, p-nitrobenzyl, p-methoxybenzyl, 2,6-dimethoxybenzyl, 4-(methylsulfinyl)benzyl, 4-sulfobenzyl, 4-azidomethoxybenzyl, 4-{a/[1-(4,4-dimethyl-2,6-dioxo-cyclohexylidene)-3-methylbutyl]amino}benzyl, piperonyl, 4-picoyl, trimethylsilyl, triethylsilyl, tert-butyl dimethylsilyl, isopropyl dimethylsilyl, phenyl dimethylsilyl, di-tert-butyl methylsilyl, triisopropylsilyl, and the like. Other carboxyl-protecting groups are known in the art.

[0117] In certain embodiments of a compound of Formula (I), the compound is an embodiment of Formula (I) described herein with the proviso that when n is 1 and each of R^2 and R^3 is C_{1-4} alkyl, then R^4 is not chosen from $-OR^7$, $-COOR^{33}$, and substituted heteroalkyl.

[0118] In certain embodiments of a compound of Formula (I), the compound is chosen from:

- [0119]** methyl 3-{{[3-(acetylamino)propyl]sulfonyloxy}-2,2-dimethylpropanoate;
- [0120]** phenylmethyl 3-{{[3-(acetylamino)propyl]sulfonyloxy}-2,2-dimethylpropanoate;
- [0121]** phenylmethyl 5-{{[3-(acetylamino)propyl]sulfonyloxy}-4,4-dimethylpentanoate;
- [0122]** phenylmethyl 4-{{[3-(acetylamino)propyl]sulfonyloxy}-3,3-dimethylbutanoate;
- [0123]** phenylmethyl 3-{{[3-(acetylamino)propyl]sulfonyloxy}-2-({[3-(acetylamino)propyl]sulfonyloxy)methyl}-2-methylpropanoate;
- [0124]** (4-methoxyphenyl)methyl 3-{{[3-(acetylamino)propyl]sulfonyloxy}-2,2-dimethylpropanoate;
- [0125]** ethyl 3-{{[3-(acetylamino)propyl]sulfonyloxy}-2,2-dimethylpropanoate;
- [0126]** methylethyl 3-{{[3-(acetylamino)propyl]sulfonyloxy}-2,2-dimethylpropanoate;
- [0127]** 2'-morpholinylethyl 3-{{[3-(acetylamino)propyl]sulfonyloxy}-2,2-dimethylpropanoate hydrochloride;
- [0128]** pyridinylmethyl 3-{{[3-(acetylamino)propyl]sulfonyloxy}-2,2-dimethylpropanoate hydrochloride;
- [0129]** ethoxycarbonyloxyethyl 3-{{[3-(acetylamino)propyl]sulfonyloxy}-2,2-dimethylpropanoate;
- [0130]** phenylmethyl 2-({[3-(acetylamino)propyl]sulfonyloxy)methyl}-3-hydroxy-2-methylpropanoate;
- [0131]** methyl 4-{{[3-(acetylamino)propyl]sulfonyloxy}-3,3-dimethylbutanoate;
- [0132]** 3-{{[3-(acetylamino)propyl]sulfonyloxy}-2,2-dimethylpropanoic acid;
- [0133]** 5-{{[3-(acetylamino)propyl]sulfonyloxy}-4,4-dimethylpentanoic acid;
- [0134]** methylethoxycarbonyloxyethyl 3-{{[3-(acetylamino)propyl]sulfonyloxy}-2,2-dimethylpropanoate;
- [0135]** 4-{{[3-(acetylamino)propyl]sulfonyloxy}-3,3-dimethylbutanoic acid;
- [0136]** methyl 3-{{[3-(acetylamino)propyl]sulfonyloxy}-2-[(tert-butoxy)carbonylamino]-2-methylpropanoate;
- [0137]** methyl 3-{{[3-(acetylamino)propyl]sulfonyloxy}-2-amino-2-methyl propanoate;
- [0138]** ethyl 3-{{[3-(acetylamino)propyl]sulfonyloxy}-2,2-diethoxypropanoate;
- [0139]** benzyl 3-{{[3-(acetylamino)propyl]sulfonyloxy}-2,2-diethoxypropanoate; and
- [0140]** a pharmaceutically acceptable salt of any of the foregoing.


[0141] In certain embodiments of a compound of Formula (I), the compound is chosen from:

[0142] phenylmethyl 3-[[3-(acetyl amino)propyl]sulfonyloxy]-2,2-dimethylpropanoate;

[0143] ethyl 3-[[3-(acetylamino)propyl]sulfonyloxy]-2,2-dimethylpropanoate; and

[0144] a pharmaceutically acceptable salt of any of the foregoing.

[0145] Certain embodiments provide a compound of Formula (II):



[0146] or a pharmaceutically acceptable salt thereof; wherein:

[0147] q is chosen from 0, 1, 2, and 3;

[0148] R²² and R²³ are independently chosen from C₁₋₄ alkyl, substituted C₁₋₄ alkyl, C₁₋₄ alkoxy, substituted C₁₋₄ alkoxy, —NH₂, —NHC(O)R³⁴, and —N(R³⁴)₂; or R²² and R²³ together with the carbon to which they are bonded form a ring chosen from a C₃₋₈ cycloalkyl, substituted C₃₋₈ cycloalkyl, C₃₋₈ heterocycloalkyl, and substituted C₃₋₈ heterocycloalkyl ring; and

[0149] each R²⁴ is independently chosen from hydrogen, halogen, —OR⁷, —CN, —CF₃, =O, —NO₂, —COOR³³, C₁₋₈ alkyl, substituted C₁₋₈ alkyl, C₆₋₁₀ aryl, substituted C₆₋₁₀ aryl, C₃₋₈ cycloalkyl, substituted C₃₋₈ cycloalkyl, C₇₋₁₈ arylalkyl, substituted C₇₋₁₈ arylalkyl, C₄₋₁₆ cycloalkylalkyl, substituted C₄₋₁₆ cycloalkylalkyl, C₁₋₈ heteroalkyl, substituted C₁₋₈ heteroalkyl, C₅₋₁₀ heteroaryl, substituted C₅₋₁₀ heteroaryl, C₃₋₈ heterocycloalkyl, substituted C₃₋₈ heterocycloalkyl, C₆₋₁₈ heteroarylalkyl, substituted C₆₋₁₈ heteroarylalkyl, C₄₋₁₆ heterocycloalkylalkyl, substituted C₄₋₁₆ heterocycloalkylalkyl, —NH₂, —NHC(O)R³⁴, —NHR³⁴, and —N(R³⁴)₂; wherein:

[0150] R⁷ is chosen from hydrogen, C₁₋₈ alkyl, substituted C₁₋₈ alkyl, C₆₋₁₀ aryl, substituted C₆₋₁₀ aryl, C₃₋₈ cycloalkyl, substituted C₃₋₈ cycloalkyl, —C(O)R⁸, —C(O)OR⁸, —C(O)N(R⁸)₂, —C(O)SR⁸, —C(R⁹)₂OC(O)R⁸, and —C(R⁹)₂OC(O)OR⁸; wherein:

[0151] each R⁸ is independently chosen from C₁₋₆ alkyl, substituted C₁₋₆ alkyl, C₆₋₈ aryl, substituted C₆₋₈ aryl, C₃₋₈ cycloalkyl, substituted C₃₋₈ cycloalkyl, C₁₋₆ heteroalkyl, substituted C₁₋₆ heteroalkyl, C₅₋₈ heteroaryl, substituted C₅₋₈ heteroaryl, C₃₋₈ heterocycloalkyl, and substituted C₃₋₈ heterocycloalkyl;

[0152] each R⁹ is independently chosen from hydrogen, C₁₋₈ alkyl, substituted C₁₋₈ alkyl, C₆₋₈ aryl, substituted C₆₋₈ aryl, C₃₋₈ cycloalkyl, substituted C₃₋₈ cycloalkyl, and —COOR³⁵; wherein R³⁵ is chosen from hydrogen and C₁₋₈ alkyl;

[0153] R³³ is chosen from hydrogen and C₁₋₄ alkyl;

[0154] each R³⁴ is independently chosen from hydrogen, C₁₋₈ alkyl, substituted C₁₋₈ alkyl, C₁₋₈ alkoxy, substituted C₁₋₈ alkoxy, C₆₋₁₀ aryl, substituted C₆₋₁₀ aryl, C₆₋₁₀ aryloxy, substituted C₆₋₁₀ aryloxy, C₃₋₈

cycloalkyl, substituted C₃₋₈ cycloalkyl, C₃₋₈ cycloalkoxy, substituted C₃₋₈ cycloalkoxy, C₇₋₁₈ arylalkyl, substituted C₇₋₁₈ arylalkyl, C₄₋₁₆ cycloalkylalkyl, substituted C₄₋₁₆ cycloalkylalkyl, C₁₋₈ heteroalkyl, substituted C₁₋₈ heteroalkyl, C₅₋₁₀ heteroaryl, substituted C₅₋₁₀ heteroaryl, C₃₋₈ heterocycloalkyl, substituted C₃₋₈ heterocycloalkyl, C₆₋₁₈ heteroarylalkyl, substituted C₆₋₁₈ heteroarylalkyl, C₄₋₁₆ heterocycloalkylalkyl, and substituted C₄₋₁₆ heterocycloalkylalkyl; and

[0155] with the proviso that when n is 1 and each of R²² and R²³ is methyl; then R²⁴ is not chosen from OR⁷, —COOR³³, and substituted heteroalkyl.

[0156] In certain embodiments of a compound of Formula (II), each substituent is independently chosen from halogen, —OH, —CN, —CF₃, —OCF₃, =O, —NO₂, C₁₋₄ alkoxy, C₁₋₄ alkyl, —COOR³⁰, —N(R³¹)₂, and —CON(R³²)₂; wherein each R³⁰, R³¹, and R³² is independently chosen from hydrogen and C₁₋₄ alkyl. In certain embodiments of a compound of Formula (II), each substituent is chosen from C₁₋₄ alkyl, C₁₋₄ alkoxy, —OH, and —NH₂.

[0157] In certain embodiments of a compound of Formula (II), q is chosen from 0, 1, and 2. In certain embodiments of a compound of Formula (II), q is 0, q is 1, q is 2, and in certain embodiments, q is 3.

[0158] In certain embodiments of a compound of Formula (II), R²² and R²³ are independently chosen from C₁₋₄ alkyl and C₁₋₄ alkoxy.

[0159] In certain embodiments of a compound of Formula (II), each of R²² and R²³ is methyl.

[0160] In certain embodiments of a compound of Formula (II), each R²⁴ is independently chosen from hydrogen, halogen, —OH, —CN, —CF₃, =O, —NO₂, C₁₋₆ alkoxy, C₁₋₆ alkyl, —COOR³³, and —N(R³⁴)₂; wherein R³³ is chosen from hydrogen and C₁₋₃ alkyl; and each R³⁴ is independently chosen from hydrogen and C₁₋₃ alkyl.

[0161] In certain embodiments of a compound of Formula (II), each R²⁴ is chosen from hydrogen, C₁₋₄ alkyl, and C₁₋₄ alkoxy. In certain embodiments of a compound of Formula (II), each R²⁴ is hydrogen.

[0162] In certain embodiments of a compound of Formula (II), the stereochemistry of the carbon atom to which R²⁴ is bonded is of the (S)-configuration.

[0163] In certain embodiments of a compound of Formula (II), the stereochemistry of the carbon atom to which R²⁴ is bonded is of the (R)-configuration.

[0164] In certain embodiments of a compound of Formula (II), q is chosen from 0, 1, and 2; each R²⁴ is chosen from hydrogen and C₁₋₄ alkyl; and each of R²² and R²³ is methyl. In certain embodiments of a compound of Formula (II), the compound is an embodiment of Formula (II) described herein with the proviso that when n is 1 and each of R²² and R²³ is C₁₋₄ alkyl, then R²⁴ is not chosen from OR⁷, —COOR³³, and substituted heteroalkyl.

[0165] In certain embodiments of a compound of Formula (II), the compound is chosen from:

[0166] 3-[[3-(acetylamino)propyl]sulfonyloxy]-2,2-dimethylpropanoic acid;

[0167] 4-[[3-(acetylamino)propyl]sulfonyloxy]-3,3-dimethylbutanoic acid;

[0168] 5-[[3-(acetylamino)propyl]sulfonyloxy]-4,4-dimethylpentanoic acid; and

[0169] a pharmaceutically acceptable salt of any of the foregoing.

[0170] In certain embodiments of compounds of Formula (I)-(II), a pharmaceutically acceptable salt is chosen from a sodium salt, a potassium salt, a lithium salt, an ammonium salt, a calcium salt, a zinc salt, and a magnesium salt. In certain embodiments of compounds of Formula (I)-(II), a pharmaceutically acceptable salt is the hydrochloride salt, and in certain embodiments, the sodium salt.


[0171] In certain embodiments, compounds of Formula (I)-(II) are free acids.

Synthesis


[0172] Compounds disclosed herein may be obtained via the synthetic methods illustrated in Schemes 1-24. Those of ordinary skill in the art will appreciate that a useful synthetic route to the disclosed compounds comprises bonding a substituted neopentyl alcohol or appropriate intermediate thereof bearing a suitable functional group at the neopentyl position of the promoity to acamprosate, i.e. sulfonyl chloride, of acamprosate to form a substituted neopentyl sulfonyl ester moiety.

[0173] General synthetic methods useful in the synthesis of compounds described herein are available in the art. Starting materials useful for preparing compounds and intermediates thereof are commercially available or can be prepared by well-known synthetic methods. Other methods for the synthesis of compounds provided by the present disclosure are either described in the art or will be readily apparent to the skilled artisan in view of the references provided herein and may be used to synthesize the compounds provided by the present disclosure. Accordingly, the methods presented in the schemes are illustrative rather than comprehensive or limiting.


[0174] In certain embodiments, and referring to Scheme 1, commercially available homotaurine 1 can be converted to the corresponding 3-(N-acetyl) homotaurinate derivative 2 using methods or variations thereof disclosed in Durlach, et al., U.S. Pat. No. 4,355,043, DE 30 19 350 C2, or Berthelon, et al., U.S. Pat. No. 6,265,437 B1.



[0175] In certain embodiments, and referring to Scheme 2, commercially available potassium phthalimide 3 can be reacted with 3-propanesultone 4 in a solvent such as ethanol (EtOH) at a temperature from about 25° C. to about 80° C. to provide the corresponding potassium 3-(N-phthalimido) propylsulfonate 5, using methods or variations thereof described by Shue, et al., *Bioorg. Med. Chem. Lett.* 1996, 6, 1709.



[0176] Referring to Scheme 3 (where Q is NHAc, phthalimido, or other useful amine precursor, M is a metal salt, and X is halogen), drugs or suitable precursors of drugs having at least one sulfonic acid group 6, a suitable sulfonic acid derivative thereof such as a tetraalkylammonium salt 7, or certain metal salts of sulfonic acids 8 can be reacted with activation agents to provide the corresponding activated sulfonic acid derivatives, i.e. sulfonyl chlorides. Useful methods are described in Shue, et al., *Bioorg. Med. Chem. Lett.* 1996, 6, 1709; and Korolev, et. al., *Synthesis* 2003, 3, 383-388. For example, activation of sulfonic acid 6, the corresponding tetraalkylammonium salt 7, such as the tetramethylammonium salt of a sulfonic acid derivative, or the corresponding alkali metal salt 8 (n is 1) can be accomplished by reaction with an appropriate chlorination agent such as phosphorous pentachloride (PCl₅), or, alternatively, thionyl chloride (SOCl₂), sulfuryl chloride (SO₂Cl₂), or cyanuric chloride (CICN); in a solvent such as the chlorination agent itself, dichloromethane (DCM), and the like, optionally in the presence of a catalyst such as N,N-dimethylformamide (DMF); and at a temperature from about 0° C. to about 60° C.; to provide the corresponding sulfonic acid chlorides or sulfonyl chlorides 9 such as 3-(N-acetyl)propylsulfonyl chloride (acamprosate chloride) or 3-phthalimido propylsulfonyl chloride.




[0177] In certain embodiments, and referring to Scheme 3, certain activated precursors of drugs having at least one sulfonic acid group 6, for example, where Q is chlorine and X is chlorine, i.e. 3-chloropropylsulfonyl chloride, are commercially available and can be used directly as coupling partners for the synthesis of functionalized prodrug intermediates.

[0178] Masked carboxylate neopentyl sulfonic acid prodrugs, intermediates, and precursors of any of the foregoing can be prepared according to general synthetic Schemes 4-24. In general, activated sulfonic acid intermediates such as sulfonyl chlorides can be coupled with a functionalized neopentyl alcohol in the presence of a base and/or a catalyst at a temperature from about -78°C . to about 65°C . to provide neopentyl sulfonyl ester prodrugs, intermediates, or precursors. Depending on the nature of the functional groups of the sulfonyl moiety and/or the neopentyl alcohol, the intermediates or precursors may be further derivatized or interconverted to provide the desired prodrugs.

[0179] Examples for preparing functionalized neopentyl promoieties and appropriately functionalized neopentyl alcohols such as functionalized 2,2-bis-substituted 3-hydroxy propanoic acid derivatives, that are useful as coupling partners are shown in the following schemes.


[0180] Referring to Scheme 4, 2,2-bis-substituted 3-hydroxy propanoic acid derivative 11 (corresponding to n is 0 in Formula (I)) as a functionalized neopentyl promoiety is provided, where R^1 , R^2 , and R^3 are as defined herein. In certain embodiments, each of R^2 and R^3 is methyl, and R^1 is alkyl or substituted alkyl, and the starting material is 2,2-dimethyl 3-hydroxypropanoic acid (hydroxypivalic acid) 10. In certain embodiments, where R^2 and R^3 are independently chosen from methyl and hydroxymethyl, and R^1 is alkyl or substituted alkyl, the starting material is 2,2-(bis-hydroxymethyl) propionic acid.




[0181] Using known synthetic methods, 2,2-bis-substituted 3-hydroxy propanoic acid derivatives such as 2,2-dimethyl 3-hydroxy propanoic acid (hydroxypivalic acid), 2,2-(bis-hydroxymethyl)propionic acid, and the like, can be converted to the corresponding ester derivative 11 in the presence of an inorganic base such as an alkali carbonate (e.g., Cs_2CO_3 or K_2CO_3), and an alkyl halide reagent such as alkyl or benzylic halides (e.g., ethyl iodide (EtI), isopropyl bro-

mid (iPrBr), or benzyl bromide (BnBr)), in an inert solvent such as N,N-dimethylformamide (DMF), N-methylpyrrolidone (NMP), N,N-dimethylacetamide (DMA, DMAc), dimethylsulfoxide (DMSO), or tetrahydrofuran (THF), at a temperature from about 0°C . to about 100°C .

[0182] Methods for preparing acyloxyalkyl ester derivatives, and alkoxy- or aryloxy-carbonyloxy ester derivatives 14 of 2,2-bis-substituted 3-hydroxy propanoic acid derivatives 12 are shown in Scheme 5 where R^2 , R^3 , R^5 , and R^6 are as defined herein, X is halogen, and Y is oxygen or a bond. In certain embodiments, where each of R^2 and R^3 is methyl, R^5 is ethyl, and R^6 is methyl; the starting materials are 2,2-dimethyl 3-hydroxy propanoic acid (hydroxypivalic acid) and rac-1-chloroethyl ethyl carbonate. Unsubstituted and substituted 1-halogenoalkyl carboxylates, or 1-halogenoalkyl alkyl- or aryl-carbonates are either commercially available or can be prepared from commercially available starting materials adapting procedures or variations thereof according to Harada, et al., *Synth. Commun.* 1994, 24, 767-772; Davidsen, et al., *J. Med. Chem.* 1994, 37, 4423-4429; Jasys, EP 0 061 274 B1; and Wheeler, et al., *J. Med. Chem.* 1979, 22, 657-661, or other methods known in the art. Acyloxyalkyl ester derivatives or alkoxy- and aryloxy-carbonyloxy ester derivative 14 can be obtained by reacting 2,2-bis-substituted 3-hydroxypropanoic acid derivative 12 with a substituted 1-halogenoalkyl carboxylate or a 1-halogenoalkyl alkyl- or aryl-carbonate 13 in the presence of a tertiary organic base such as triethylamine (Et_3N , TEA), diethylisopropylamine (DIEA, Hünigs-base), or NMM (N-methylmorpholine); or an amidine base such as 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU) or 1,5-diazabicyclo(4.3.0)non-5-ene (DBN); either in neat form or in an organic solvent such as 1,2-dichloroethane (DCE), at a temperature from about 0°C . to about 100°C .



[0183] Alternatively, 2,2-bis-substituted 3-hydroxy propanoic acid derivatives can be prepared according to Scheme 6, where R^1 , R^2 , and R^3 are defined herein and Y is oxygen or a bond. In certain embodiments, each of R^2 and R^3 is methyl; R^1 is 3-pyridylmethyl (nicotinyl) or 2-(morpholin-4-yl)ethyl (mofetil); the starting material is 2,2-dimethyl 3-hydroxy propanoic acid (hydroxypivalic acid) 15; and Y is oxygen.




[0184] For example, referring to Scheme 6, protection of hydroxypropanoic acid 15 with mixed trialkyl- or mixed trialkylarylchlorosilanes such as tert-butyl dimethylsilylchlorosilane (TBDMSCI), triisopropylchlorosilane (TIPSCI), tert-butyldiphenylsilyl chlorosilane (TBDPSCI), and the like, in an inert solvent such as dichloromethane (DCM), tetrahydrofuran (THF), or N,N-dimethylformamide (DMF), in the presence of an organic base such as imidazole or triethylamine (Et₃N, TEA), and optionally a catalytic amount of a nucleophilic catalyst such as 4-(N,N-dimethyl)aminopyridine (DMAP), at a temperature from about 0° C. to about 60° C., provides the corresponding 3-trialkyl- or mixed 3-alkylarylsiloxy 2,2-dimethyl propanoic acid intermediates. Other methods for the selective introduction and removal of protecting groups and alternative protection strategies are known in the art.

[0185] Functionalized carboxylic acid derivatives such as carboxylic acid esters or carboxamides of protected 3-trialkyl- or mixed 3-alkylarylsiloxy 2,2-bis-substituted propanoic acids can be obtained through an activation/coupling sequence. For example, 3-trialkyl- or mixed 3-alkylarylsiloxy 2,2-bis-substituted propanoic acids such as 2,2-dimethyl 3-(tert-butyldimethylsilyloxy) propanoic acid can be contacted with an activation agent such as a dehydration agent, e.g., N,N'-dicyclohexylcarbodiimide (DCC); in an inert solvent such as dichloromethane (DCM), acetonitrile (MeCN), and the like; in the presence of an additive such as a nucleophilic acylation catalyst, e.g. 4-(N,N-dimethylamino)pyridine (DMAP); at a temperature from about 0° C. to about 60° C. The activated intermediate of the 3-trialkyl- or mixed 3-alkylarylsiloxy 2,2-dimethyl propanoic acid, such as 2,2-dimethyl 3-(tert-butyldimethylsilyloxy)propanoic acid can then be reacted in the same solvent with a functionalized alcohol such as 2-(morpholin-4-yl)ethanol or 3-pyridylmethanol, to provide the corresponding alkyl-, aryl-, 3-trialkyl-, or mixed 3-alkylarylsiloxy 2,2-bis-substituted propanoate.


[0186] Reaction of alkyl or aryl 2,2-dialkyl 3-trialkylsiloxy propanoates with reagents capable of selectively cleaving the 3-trialkyl or mixed alkylarylsilyl protecting group provide alkyl- or aryl-2,2-dialkyl 3-hydroxy propanoate 16 that is useful neopentyl alcohol promoieties and/or coupling partners. For example, trialkylsilyl or mixed alkylarylsilyl-protected derivatives can be selectively cleaved using fluoride-containing agents such as tetrabutylammonium fluoride (TBAF), potassium fluoride (KF), ammonium fluoride (H₄NF), and hydrogen fluoride (HF); or using hydrogen fluoride complexes with organic bases such as triethylamine tri-

hydrofluoride (Et₃N·3HF) or pyridinium hydrofluoride; in an inert solvent such as tetrahydrofuran (THF); at a temperature from about 0° C. to about 100° C. to provide the corresponding desilylated alkyl 2,2-dialkyl 3-hydroxy propanoate 16.

[0187] As shown in Scheme 7, heteroatom-protected intermediate 18 can be synthesized from an appropriately functionalized 3-hydroxy propanoic acid derivative such as 2-amino-3-hydroxy-2-methylpropanoic acid 17. Standard esterification methods, e.g., anhydrous methanol (MeOH) in the presence of a catalytic amount of an acidic catalyst such as thionyl chloride (SOCl₂), sulfuryl chloride (SO₂Cl₂), concentrated sulfuric acid (H₂SO₄), or trimethylsilyl chloride (TMSCl); or a sulfonic acid derivative such as para-toluene sulfonic acid (TsOH) or camphor sulfonic acid (CSA); at a temperature from about 0° C. to about 100° C. can be used to provide the corresponding protected methyl ester. As shown in Scheme 7, Step 2, methyl 2-amino-3-hydroxy-2-methylpropanoate can be reacted with di-tert-butylpyrocarbonate (Boc₂O) in the presence of a base to provide the corresponding N-Boc protected methyl 2-amino-3-hydroxy-2-methylpropanoate 18. Examples of useful solvents for the reaction shown in Scheme 7, step 2, include a mixture of a 1N aqueous solution of sodium hydroxide (NaOH) and 1,4-dioxane, a saturated aqueous solution of sodium bicarbonate (NaHCO₃) with acetonitrile as a co-solvent, dichloromethane (DCM), and a tertiary organic base, optionally in the presence of a catalyst. Examples of useful tertiary organic bases include triethylamine (TEA) and a catalytic amount of 4-(N,N-dimethylamino)pyridine (DMAP).



[0188] Scheme 8 shows the synthesis of alkyl- or aryl-2,2-dialkoxy 3-hydroxy propanoate 20 as a functionalized neopentyl promoieties where R¹ is as defined herein and R^c and R^d are independently alkyl or R^c and R^d are linked by an alkyl to form a heteroalkyl ring. In certain embodiments where each of R^c and R^b is ethyl (Et), and R¹ is ethyl (Et) or benzyl (Bn), the starting material is the corresponding acrylic acid derivative 19. The carbon-carbon double bond of an acrylic acid ester such as an ethyl acrylate or benzyl acrylate can be dihydroxylated to obtain the corresponding alkyl glycerate using oxidation agents such as potassium permanganate (KMnO₄) in a solvent such as acetone and water, at a temperature from about -78° C. to about 60° C. Methods for the oxidative transformation of alkenes into vicinal diols are known in the art.




[0189] The primary hydroxyl group of an alkyl glycerate can be selectively protected by reacting the alkyl glycerate with a bulky trialkyl chlorosilane such as tert-butyl dimethylsilylchlorosilane (TBDMSCl), triisopropylchlorosilane (TIPSCl), tert-butyl diphenylsilylchlorosilane (TBDPSCl), in an inert solvent such as dichloromethane (DCM), tetrahydrofuran (THF), or N,N-dimethylformamide (DMF), in the presence of an organic base such as imidazole or triethylamine (Et₃N, TEA), optionally in the presence of a catalytic amount of a nucleophilic catalyst such as 4-(N,N-dimethyl)aminopyridine (DMAP) at a temperature from about 0° C. to about 60° C.

[0190] Methods for oxidizing secondary hydroxyl groups to oxo groups, i.e. ketones, are well known. For example, the 2-hydroxyl group of tris alkylsilyl- or mixed alkyl-arylsilyl-protected alkyl glycerate can be oxidized to provide the corresponding alkyl 2-oxo 3-silyloxy propanoate using 1,1,1-tris (acetyloxy)-1,1-dihydro-1,2-benziodoxol-3-(1H)-one (Dess-Martin periodinane) in an inert solvent such as dichloromethane (DCM) at a temperature from about -20° C. to about 25° C.

[0191] In certain embodiments of Scheme 8, alkyl 2,2-dialkoxy 3-trialkylsilyloxy propanoates are provided. Formation of ketals from oxo-compounds, i.e. ketones, is well known. For example, an alkyl 2-oxo 3-trialkylsilyloxy propanoate can be reacted with an excess of alcohol such as ethanol (EtOH) or an appropriately functionalized diol such as ethylene glycol; or with a suitable transacetalization reagent such as a trisalkyl orthoformate, i.e. triethyl orthoformate, either in the neat form or in the presence of an inert solvent and a catalyst such as concentrated sulfuric acid (H₂SO₄), pyridinium para-toluene sulfonate (PPTS), para-toluene sulfonate (TsOH), or camphorsulfonic acid (CSA); at a temperature from about -20° C. to about 100° C. Alternatively, when alcohols are used, the reaction can be carried out by azeotropic removal of water generated during the reaction.


[0192] Reaction of an alkyl 2,2-dialkoxy 3-silyloxy propanoate with reagents capable of selectively cleaving the 3-silyl protecting group provide alkyl 2,2-dialkoxy 3-hydroxy propanoate 20 that is a useful neopentyl alcohol promoiety or coupling partner. For example, reacting an alkyl 2,2-dialkoxy 3-trialkylsilyloxy propanoate with an acid in a solvent at a temperature from about 0° C. to about 100° C. provides the corresponding desilylated alkyl 2,2-dialkoxy 3-hydroxy propanoate 20. Examples of useful acid and solvent mixtures for the reaction include mixtures of acetic acid (HOAc), water, and tetrahydrofuran (THF); and concentrated hydrochloric acid (HCl) in ethanol (EtOH). Alternatively, fluoride-containing agents can be used.

[0193] As shown in Scheme 9 (wherein n, R², R³, and R⁴ are as defined herein; X is halogen such as chloro; Y is hydrogen, alkoxy,; and Q is a NHAc or a precursor to an amine), an activated sulfonic acid derivative such as a sulfonyl chloride of a drug having at least one sulfonic acid group 21, e.g., acamprosate chloride, or alternatively, a similarly activated sulfonic acid derivative of a precursor of a drug having at least one sulfonic acid group can be reacted with a functionalized and externally masked neopentyl alcohol 22 to provide externally masked neopentyl acamprosate prodrugs (neopentyl sulfonyl esters) or precursors or intermediates to such prodrugs 23. Examples of externally masked nucleophile 22 include functionalized 2,2-bis-substituted 3-hydroxy propanoic acid derivatives, such as esters.




[0194] Referring to Scheme 9, when n is 0, a neopentyl promoiety 22 can be a functionalized 2,2-bis-substituted 3-hydroxy propanoic acid ester where Y is —OR¹, X is chlorine, Q is N-acetylamino (NHAc), and R², R³, and R⁴ are defined herein, and the activated sulfonic acid derivative is 3-(N-acetyl)propylsulfonyl chloride (acamprosate chloride) 21. Neopentyl alcohol 22 can be reacted with 3-(N-acetyl)propylsulfonyl chloride 21 in a solvent such as dichloromethane (DCM) in the presence of a base such as triethylamine (Et₃N, TEA), pyridine, or diisopropyl ethyl amine (iPr₂EtN, DIEA); and a nucleophilic catalyst such as 4-(N,N-dimethyl)pyridine (DMAP); at a temperature from about -20° C. to about 25° C. to provide the corresponding internally masked neopentyl sulfonyl ester 23.

[0195] When Q is N-acetylamino (NHAc), R² and R³ are independently chosen from methyl and tert-butoxycarbonylamino (NBoc), Y is methoxy, and n is 0 in Scheme 9, as shown in Scheme 10 the corresponding N-unprotected derivative of neopentyl sulfonylester acamprosate prodrug 25 can be prepared by reacting a N-Boc-protected neopentyl sulfonyl ester derivative 24 with a strong acid in an inert solvent such as trifluoroacetic acid in dichloromethane (DCM) or hydrogen chloride (HCl) in 1,4-dioxane. In the reaction the tert-butoxycarbonyl (Boc) protecting group can be cleaved to provide the corresponding unprotected species in either free amine or in N-protonated form, i.e. ammonium salt, where G is chosen from NH₂, NH₃⁺Cl⁻, and NH₃⁺F₃CCO₂⁻.




[0196] As shown in Scheme 11, and also referring to Scheme 9, when Q is N-acetylamino (NHAc), each of R^2 and R^3 is methyl, Y is benzoyloxy, i.e., phenylmethoxy, and n is 0; the free acid 27 of the corresponding benzyl hydroxypivalic acid conjugate 26 can be obtained by reacting the conjugate with hydrogen in the presence of a heterogeneous catalyst such as palladium on activated carbon, in a solvent such as methanol (MeOH), ethanol (EtOH), or ethyl acetate (EtOAc), at a temperature from about 0° C. to about 50° C. and under a pressure of about 15 psi to about 60 psi.



[0197] Examples of methods for synthesizing functionalized neopentyl promoieties, appropriately functionalized neopentyl alcohols, precursors, or derivatives thereof, such as suitably functionalized 2,2-bis-substituted ω -unsaturated alcohols that are useful as coupling partners are shown in the following schemes.

[0198] Scheme 12 (where R^2 , R^3 , and R^4 are as defined herein; R^a is hydrogen; R^b is hydrogen, alkyl, alkoxy, amide, substituted carbonyl or aryl; and A is hydrogen, hydroxyl, or alkoxy) shows the synthesis of 3,3-bis-substituted 4-hydroxy butanoic acid (corresponding to n is 1 in compound 20) and 4,4-bis-substituted 5-hydroxy pentanoic acid promoieties (corresponding to n is 2 in compound 20). The synthetic method illustrated in Scheme 12 is extendable to homologs such as 5,5-bis-substituted 6-hydroxy hexanoic acids. Conjugates based on 3,3-bis-substituted 4-hydroxy butanoic acid and 4,4-bis-substituted 5-hydroxy pentanoic acid precursor 29 can be derivatized after coupling to a sulfonyl group of a drug or a precursor of a drug having at least one sulfonic acid

group to provide the corresponding neopentyl sulfonyl ester prodrug. In certain embodiments, where each of R^2 and R^3 is methyl, R^4 is hydrogen, and n is either 1 or 2, the starting material 28 can be chosen from 2,2-dimethyl-4-pentenoic acid, methyl 2,2-dimethyl-4-pentenoic acid, 2,2-dimethyl-4-pentenal, and 2,2-dimethyl-5-hexenoic acid.




[0199] For example, referring to Scheme 12, in certain embodiments, A is lower alkoxy such as methoxy, hydroxyl, or hydrogen; each of R^2 and R^3 is methyl, R^4 , each of R^a and R^b is hydrogen; and n is 1 or 2. Using standard synthetic methods, 2,2-dimethyl-4-pentenoic acid, its lower alkyl ester (A is OH or O-lower alkyl), or 2,2-dimethyl-5-hexenoic acid can be converted to the corresponding alcohol 29 by reaction with reducing agents such as lithium aluminum hydride (Li-AlH₄, LAH); in an anhydrous inert solvent such as tetrahydrofuran (THF) or diethyl ether (Et₂O); at a temperature from about -78° C. to about 65° C. Alternatively, the reaction can be carried out using LiBH₄ in an inert solvent such as tetrahydrofuran, at a temperature from about 0° C. to about 25° C. Aldehydes 28, e.g., A is H, and n is 1, can be reduced to the corresponding alcohol 29 using boron hydride reagents such as alkali borohydrides, e.g., NaBH₄, in an alcohol solvent such as methanol (MeOH) or ethanol (EtOH).

[0200] Activated sulfonic acid derivatives such as a sulfonyl chloride of a drug or precursor of a drug having at least one sulfonic acid group 30, e.g. acamprosate chloride, can be reacted with a functionalized coupling partner to provide useful intermediates for preparing externally masked neopentyl sulfonyl ester prodrugs 32 as disclosed herein. Examples of coupling partners include functionalized 2,2-bis-substituted ω -unsaturated alcohols 31.


[0201] Referring to Scheme 13 (where n, R^2 , R^3 , and R^4 are as defined herein, R^a is hydrogen, R^b is hydroxy, alkyl, or an aromatic, X is halogen, and Q is an amine precursor), in certain embodiments where n is 1 or 2, and the corresponding neopentyl promoieties are functionalized 2,2-bis-substituted ω -unsaturated alcohols 31 wherein X is chlorine, and Q is N-acetylamino (NHAc), and the activated sulfonic acid derivative is 3-(N-acetyl)propylsulfonyl chloride (acamprosate chloride) 30. Functionalized 2,2-bis-substituted ω -unsaturated alcohol 31 can be reacted with 3-(N-acetyl)propylsulfonyl chloride (acamprosate chloride) 30 in a solvent such as dichloromethane (DCM) in the presence of a base such as triethylamine (Et₃N, TEA), pyridine, or diisopropyl ethyl amine (iPr₂EtN, DIEA) and a nucleophilic catalyst such as 4-(N,N-dimethyl)pyridine (DMAP) at a temperature from about -20° C. to about 25° C. to provide precursor or intermediate 32 to externally masked neopentyl sulfonyl esters.

Scheme 13



[0202] As shown in Scheme 14, terminally unsaturated sulfonate ester coupling intermediates 33 can then be converted to carboxylic acid ester sulfonate ester intermediates 35 such as ester or amide derivatives. Scheme 14 shows a method for converting the terminal carbon-carbon double bond to a one carbon shortened ester derivative. Other methods for the oxidative transformation of alkenes into aldehydes or carboxylic acid derivatives are described in the art.

Scheme 14




[0203] In certain embodiments of Scheme 14, Q is N-acetyl-amino (NHAc), each of R² and R³ is methyl, R⁴ is hydrogen, each of R^a and R^b is hydrogen, and n is 1 or 2. Oxidative cleavage of the terminal carbon-carbon double bond of alkene 33 by ozonolysis using a gaseous mixture of oxygen and ozone (O₂/O₃) in a solvent such as dichloromethane (DCM) or DCM/alcohol mixtures, i.e., DCM/MeOH=9:1–5:1, at a temperature of about –78° C. followed by reductive decomposition of the intermediate ozonide with a reducing agent


such as dimethyl sulfide (Me₂S), triphenylphosphine (Ph₃P), or tributylphosphine (Bu₃P), provides intermediate one-carbon shortened aldehyde derivative 34. Alternatively, aldehyde derivative 34 can be prepared by oxidation methods using sodium meta-periodate (NaIO₄)/catalytic osmium tetroxide (OsO₄) (Lemieux-Johnson reagent) in a mixture of solvents such as tetrahydrofuran (THF) and water at a temperature from about 0° C. to about 40° C. Other oxidation methods such as the ruthenium-catalyzed oxidative cleavage of olefins to aldehydes can also be useful in preparing aldehyde precursors from unsaturated compounds 33 (Yang, et al., *J. Org. Chem.* 2001, 66, 4814).

[0204] Aldehyde 34 can be converted to the corresponding carboxylic acid ester under oxidative reaction conditions to provide externally masked neopentyl prodrugs that are based on functionalized (n-1), (n-1)-bis-substituted n-hydroxy alkanic acids such as 3,3 bis-substituted 4-hydroxy butanoic acid or 4,4-bis-substituted 5-hydroxy pentanoic acid promoiety. For example, contacting aldehyde 34 with an oxidant such as N-iodosuccinimide (NIS); in the presence of an inorganic base such as alkali carbonate, e.g., K₂CO₃; in a solvent such as methanol or acetonitrile containing an excess of an alcohol; at a temperature from about 0° C. to about 40° C. and in the dark, provides the corresponding carboxylic acid ester as acamprosate prodrug 35 in a single step. Other oxidant systems that can be used in this transformation include Oxone® in an alcohol solvent, bromine or iodine (Br₂, I₂), N-bromosuccinimide (NBS)/2,2'-azobis(2-methylpropionitrile (AIBN), pyridinium dichromate (PDC), and manganese dioxide (MnO₂)/hydrogen cyanide (HCN).

[0205] When P is oxygen in Scheme 15 (where n, R¹, R², R³, and R⁴ are as defined herein, Q is NHAc or an amine precursor, R^a is hydrogen (see Scheme 14), aldehyde derivative 36 can be oxidized to the corresponding free carboxylic acid derivative 37 using standard procedures. For example, Jones-oxidation of aldehyde 36 with excess chromic acid (H₂CrO₄) in a solvent such as acetone or a water/acetone mixture at a temperature from about –10° C. to about 40° C. provides the corresponding carboxylic acid 37. The oxidant Oxone® can also be used to oxidize aldehydes to carboxylic acids in solvents such as N,N-dimethylformamide (DMF) and at a temperature from about 0° C. to about 25° C. Other oxidation systems, for example transition metal-based systems comprising a co-oxidant and an oxidation catalyst can also be used and are well known in the art.

Scheme 15





[0206] When P in Scheme 15 is =CHR^b where R^b is chosen from hydrogen,; and n is 0, 1, or 2, using methods described by Henry, et al., *J. Org. Chem.* 1993, 58, 4745; and Travis, et al., *Org. Lett.* 2003, 5, 1031, the corresponding carboxylic acid intermediate 37 can be prepared by direct oxidation of unsaturated precursor 36 with oxidation mixtures such as chromic acid (H_2CrO_4 , Jones-reagent) or Oxone® ($2\text{HKSO}_5 \cdot \text{KHSO}_4 \cdot \text{K}_2\text{SO}_4$) in the presence of a catalytic amount of osmiumtetroxide (OsO_4) in a solvent such as acetone or N,N-dimethylformamide (DMF) at a temperature from about 0°C . to about 40°C . Other methods for effecting this transformation use systems comprising transition metal oxidation catalysts based on ruthenium, chromium, or tungsten in the presence of co-oxidants such as bleach (NaOCl) or sodium periodate (NaIO_4).


[0207] Carboxylic acids are useful precursors for preparing the corresponding carboxylic acid esters or amides of externally masked neopentyl prodrugs based on suitably functionalized 3,3-bis-substituted 4-hydroxy butanoic acid or 4,4-bis-substituted 5-hydroxy pentanoic acids. For example, referring to Scheme 15, carboxylic acids can be activated in situ to provide the corresponding acid chloride by reacting carboxylic acid 37 with an activating agent such as oxalyl chloride ($(\text{COCl})_2$) or thionyl chloride (SOCl_2) in a solvent such as the chlorination agent itself (neat) or an inert solvent such as dichloromethane (DCM); optionally in the presence of a catalyst such as a catalytic amount of N,N-dimethylformamide (DMF) at a temperature from about 0°C . to about 25°C . The acid chloride can then be quenched with an excess of a functionalized alcohol such as methanol (MeOH) or benzylic alcohol (BnOH) or other suitable alcohol or amine in the presence of a base such as pyridine, triethylamine (Et_3N , TEA), or diisopropylethylamine (iPr_2EtN , DIEA), in an inert solvent such as dichloromethane (DCM); at a temperature from about 0°C . to about 25°C . to provide the corresponding externally masked neopentyl prodrug 38, e.g., an acamprostate prodrug based on a functionalized 3,3-bis-substituted 4-hydroxy butanoic acid or 4,4-bis-substituted 5-hydroxy pentanoic acid.

[0208] Carboxylic acid derivatives may be activated using, for example, any of the activation agents described herein, and the activated intermediates can subsequently be coupled to an alcohol or other functionalized substrate.


[0209] Methods for preparing functionalized 2,2-bis-substituted ω -unsaturated alcohols (functionalized neopentyl alcohols), or derivatives thereof, useful as coupling partners with an activated sulfonic acid, such as sulfonyl chlorides of a drug or a precursor of a drug having at least one sulfonic acid group are provided in Scheme 16 (where n, R^1 , R^2 , and R^3 are as defined herein, R^b is an aromatic, X is hydrogen or alkoxy, and B is hydroxyl and hydrogen or oxygen). For example, in certain embodiments of Scheme 16, wherein each of R^1 and R^2 is methyl; n is 0; R^3 is absent; B is hydroxyl and hydrogen, or oxygen; the starting materials are either pantolactone or

dihydro-4,4-dimethyl-2,3-furandione; R^b is hydrogen; and X is hydrogen or methoxy, 3-O-protected 2,2-dimethylpent-4-en-1-ol 43 can be prepared from the starting materials using protocols or variations thereof according to Blakemore, et al., *J. Org. Chem.* 2005, 70, 5449; Mandel, et al., *Org. Lett.* 2004, 6, 4801; Shiina, et al., *Bull. Chem. Soc. Jpn.* 2001, 74, 113; Lavallée, et al., *Tetrahedron Lett.* 1986, 27, 679; or Ito et al., *Synthesis* 1993, 137.

Scheme 16



-continued



43

[0210] For example, pantolactone 39 where B is hydroxyl and hydrogen or dihydro-4,4-dimethyl-2,3-furandione 39 where B is oxygen can be converted to the corresponding 3,3-dimethylbutan-1,2,4-triol 40 by reaction with a reducing agent such as lithium aluminum hydride (LiAlH_4 , LAH) in an anhydrous inert solvent such as tetrahydrofuran (THF) or diethyl ether (Et_2O), at a temperature from about -78°C . to about 65°C . Alternatively, excess borane dimethyl sulfide complex ($\text{BH}_3\cdot\text{Me}_2\text{S}$) in the presence of a catalytic amount of sodium borohydride (NaBH_4) in an anhydrous inert solvent such as tetrahydrofuran (THF), at a temperature from about 0°C . to about 65°C . can be used for the reaction (Lavallée et al., *Tetrahedron Lett.* 1986, 27, 679-682; and Saito et al., *Chem. Lett.* 1984, 1389-1392). The reaction proceeds with conservation of stereochemical integrity (without racemization) when enantiomerically pure starting materials such as D-pantolactone are used.

[0211] Using methods described in the literature, triol 40 can be converted regioselectively to the corresponding 6-membered ring acetal (benzyliden-type acetal) under thermodynamic conditions using a suitable aldehyde derivative such as benzaldehyde (X is hydrogen), anisaldehyde (X is methoxy), benzaldehyde dimethyl acetal (X is hydrogen), or para-methoxybenzaldehyde dimethyl acetal (X is methoxy) in a solvent such as dichloromethane (DCM) or toluene; and in the presence of a catalyst such as phosphoryl chloride (POCl_3), camphorsulfonic acid (CSA), para-toluenesulfonic acid (TsOH), or pyridinium para-toluenesulfonate (PPTS), at a temperature from about 25°C . to about 110°C . The free hydroxyl group of the cyclic 1,3-acetal protected triol 41 can be oxidized to the corresponding aldehyde derivative using standard Swern-oxidation conditions such as dimethylsulfoxide (DMSO) in the presence of oxalyl chloride ($(\text{COCl})_2$) and triethylamine (Et_3N , TEA) in dichloromethane (DCM) at a temperature of about -78°C . Alternatively, useful oxidants such a sulfur trioxide-pyridine complex ($\text{SO}_3\cdot\text{Py}$) or pyridinium dichromate (PDC) (Comforth reagent) in an inert solvent such as dichloromethane (DCM) at a temperature from about 0°C . to about 25°C . can be used. Another useful oxidant for this transformation is 1,1,1-tris(acetyloxy)-1,1-dihydro-1,2-benziodoxol-3-(1H)-one (Dess-Martin periodinane) in an inert solvent such as dichloromethane (DCM) at a temperature from about -20°C . to about 25°C .


[0212] Wittig-olefination or methylenation of the aldehyde derivative with an appropriately functionalized triphenylphosphoylide or methylenetriphenylphosphorane provides the corresponding alkene derivative 42. The function-


alized phosphoylide or methylenetriphenylphosphorane can be generated in situ from the corresponding methyltriphenylphosphonium halide such as bromide, in a solvent such as tetrahydrofuran (THF) using a base such n-butyllithium (n-BuLi) or potassium tert-butoxide (KOtBu) at a temperature from about -78°C . to about 25°C . Aldehydes and phosphoylides or phosphoranes are reacted in the same solvent at temperatures from about -78°C . to about 65°C . Mild and non-basic reaction conditions such as using alternative methylenation reagents in mixtures of solvents such as dichloromethane (DCM) and tetrahydrofuran (THF) at a temperature from about 0°C . to about 25°C . can also be used to convert aldehyde 41 to alkene 42. For example, methylenation reagents can be generated in situ from zinc dust (Zn), titanium(IV) halides such as titanium(IV) chloride (TiCl_4), and dihalomethanes such as dibromomethane (CH_2Br_2) in an inert organic solvents such tetrahydrofuran (THF) at temperatures from about -78°C . to about 0°C .

[0213] Using methods known to those skilled in the art, regioselective reductive ring opening of 1,3-benzylidene acetal 42 with dialkylaluminum hydride reagents such as diisobutylaluminum hydride (iBu_2AlH , DIBAL(H)) in an inert solvent such as dichloromethane (DCM) at temperatures from about -78°C . to about 25°C . provides the corresponding 3-O-protected 2,2-dimethylpent-4-en-1-ol derivative 43. Alternatively, regioselective reductive acetal ring opening can be accomplished using reducing agents generated from lithium aluminum hydride (LiAlH_4 , LAH) and aluminum (III) chloride (AlCl_3) in an inert solvent such as diethyl ether (Et_2O), at temperature from about 0°C . to about 25°C .

[0214] Alternatively, as shown in Scheme 17 (where PG is a protecting group and R^b is hydrogen, alkoxycarbonyl, or aryl) and using procedures or a variations thereof according to Mandel, et al., *Org. Lett.* 2004, 6, 4801; and Miyoka, et al., *Tetrahedron: Asymmetry* 1995, 6, 587, 3-O-benzylic-protected 2,2-dimethylpent-4-en-1-ol derivatives or 3-O-tert-butyl dimethylsilyl protected 2,2-dimethylpent-4-en-1-ol derivatives can also be prepared from pantolactone 44 using a three step procedure. In certain embodiments of Scheme 17, PG is a protecting group such as benzyl, para-methoxybenzyl, or tert-butyl dimethylsilyl (TBDMS), and R^b is hydrogen or methoxycarbonyl.

Scheme 17





[0215] In certain embodiments, n is 0 and R^1 absent, each of R^2 and R^3 are methyl, B is hydroxyl and hydrogen and the starting material 44 is pantolactone. For example, under mildly basic conditions, the reaction of pantolactone 44 with benzyl bromide (BnBr) in the presence of silver(I) oxide (Ag_2O) in a solvent such as N,N -dimethylformamide (DMF) provides the corresponding O-benzyl-protected pantolactone 45. Other methods for introducing protecting groups into pantolactone are well known in the art and include basic conditions prone to partial racemization if the basicity of the reaction system is not controlled sufficiently. Examples include cesium carbonate promoted O-benylation with benzyl chloride/cesium carbonate in dimethylformamide (DMF) at room temperature (Dueno et al., *Tetrahedron Lett.* 1999, 40, 1843); allylation with silver oxide/allyliodide in diethyl ether (Et_2O) (Aurich et al., *Liebigs Ann. Chem. Recueil* 1997, 2, 423); and alkali metal hydrides, i.e., NaH or organic bases, i.e., diisopropylethyl amine [$(iPr)_2EtN$, Hünigs-Base] with various alkylation agents (Pirrung et al., *Synthesis* 1995, 4, 458; Hart et al., *Heterocycles* 2000, 52(3), 1025; and Gimalova et al., *Russ. J. Org. Chem.* 2005, 41(8), 1183). The formation of the tetrahydropyranyl ether (Klar et al., *Synthesis* 2005, 2, 301) is acid catalyzed

[0216] Using protocols or variations thereof according to O'Brien et al., *Tetrahedron Lett.* 2002, 43, 5491-5494; Weinges et al., *Chem. Ber.* 1994, 127, 1305-1309; Johnston et al., *J. Chem. Soc. Perkin Trans. I*, 2000, 5, 681-695; Iversen et al., *J. Chem. Soc. Chem. Commun.* 1981, 1240-1241; Wessel et al., *J. Chem. Soc. Perkin Trans. I*, 1985, 2247-2250; Enders et al., *Org. Syntheses* 2002, 78, 177-183; and Rai et al., *Tetrahedron Lett.* 2003, 44, 2267-2269, O-benzyl or O-para-methoxybenzyl protecting groups can be added to pantolactone 44 using functionalized acetimidates such as O-benzyl- or O-para-methoxybenzyl 2,2,2-trichloro acetimidates in the presence of a catalyst such triflic acid (F_3CSO_3H), a rare earth triflate such as scandium(III) triflate ($Sc(OTf)_3$), or others in solvent mixtures such as cyclohexane and dichloromethane (DCM), toluene, or acetonitrile (MeCN) at temperatures from about $0^\circ C$. to about $40^\circ C$. Other useful catalyst systems include Brønsted-acids such as para-toluenesulfonic acid (TsOH), camphorsulphonic acid (CSA), trifluoroacetic acid (TFA), and Lewis-acids such as trifluoroborane diethyl ether complex ($BF_3 \cdot Et_2O$), trityl tetrafluoroborate ($TrBF_4$), trityl perchlorate ($TrClO_4$), trimethylsilyl trifluoromethanesulfonate (TMSOTf), or tin triflates ($Sn(OTf)_2$), under similar reaction conditions. When enantiopure starting material such

as D-pantolactone is used, the reaction proceeds with conservation of stereochemical integrity (without racemization) of the stereogenic center.

[0217] Silicon-based protecting groups, such as mixed tris-alkyl or alkylaryl silyl-based protecting groups, i.e. tert-butyldimethyl silyl ($tBuMe_2Si$, TBDMS), or tert-butyldiphenyl silyl ($tBuPh_2Si$, TBDPS), also provide robust protection of the pantolactone hydroxyl group. For example, the free hydroxyl group of pantolactone 44 can be protected with tert-butyldimethyl chlorosilane ($tBuMe_2SiCl$, TBDMSCl) using an inert solvent such as dichloromethane (DCM) or N,N -dimethylformamide (DMF) and an organic base such as imidazole or triethylamine (Et_3N , TEA), optionally in the presence of additives such as catalytic amounts of nucleophilic alkylation catalysts such as 4-(N,N -dimethyl)aminopyridine (DMAP) at a temperature from about $0^\circ C$. to about $60^\circ C$. to provide the corresponding silyl protected pantolactone 45. When enantiopure starting material such as D-pantolactone is used, the reaction proceeds with conservation of stereochemical integrity (without racemization) of the stereogenic center.

[0218] Reduction of O-benzyl- or O-silyl-protected lactone 45 with dialkylaluminum hydride reducing reagents such as diisobutylaluminum hydride [iBu_2AlH , DIBAL(H)] in an inert solvent such as tetrahydrofuran (THF) at temperatures from about $-78^\circ C$. to about $0^\circ C$. provides intermediate lactol or hemiacetal 46 as a mixture of diastereomers or anomers.


[0219] Wittig-olefination of lactol or hemiacetal diastereomer 46 with functionalized triphenylphosphorylides or methylenetriphenylphosphorane (R^b is H) or methyl (triphenylphosphoranylidene)acetate (R^b is CO_2Me) provides the corresponding (substituted) alkene derivative 47. Functionalized phosphorylides can be generated in situ from the corresponding methyltriphenylphosphonium bromide or (carbomethoxymethyl)triphenylphosphonium bromide in a solvent such as tetrahydrofuran (THF) in the presence of a base such n -butyllithium ($nBuLi$) or potassium tert-butoxide ($KOtBu$) at temperatures from about $-78^\circ C$. to about $25^\circ C$. Lactols 46, phosphorylides, or phosphoranes can then be reacted either in the same solvent or in a separate solvent such as 1,2-dichloroethane (DCE) at temperatures from about $0^\circ C$. to about $70^\circ C$. to provide the corresponding 3-O-benzyl or 3-O-tert-butyldimethylsilyl protected 2,2-dimethylpent-4-en-1-ol compound 47. Olefination or methylenation of the lactol 46 can also be accomplished under a variety of Horner-Wadsworth-Emmons (HWE)-conditions with trimethyl phosphonoacetate (i.e., Banwell et al., *J. Chem. Soc., Perkin Trans. 1*, 2002, 22) and an excess of Tebbe's reagent [$Cp_2TiCH_2ClAl(CH_3)_2$] (Martin et al., *Tetrahedron Lett.* 2001, 42, 8373).

[0220] Determination of enantiomeric excess (e.e.) of intermediate 47 can be accomplished by ^1H NMR spectroscopy in the presence of the diamagnetic enantiomerically pure chiral co-solvent (R)-(-)-2,2,2-trifluoro-1-(9-anthryl) ethanol (Pirkle-alcohol) and in comparison with ^1H NMR spectra of the corresponding racemic samples under similar conditions or using other analytical method such as HPLC using chiral stationary phases or enantiopure covalent derivatization agents such as Mosher's chloride.

[0221] Referring to Scheme 18 (where R^1 , R^2 , and R^3 are as defined herein, R^b is hydrogen, alkyl, alkoxycarbonyl, or an aromatic, PG is a protecting group, and X is halogen), activated sulfonic acid derivatives such as a sulfonyl chlorides of a drug having at least one sulfonic acid group 48, e.g. 3-chlo-

propylsulfonyl chloride, can be reacted with a functionalized protected 2,2-disubstituted pent-4-en-ol derivative 49 to provide masked acamprosate neopentyl sulfonyl ester intermediates. The intermediates can be converted to the desired 2,4-dihydroxy 3,3-dimethyl butanoic acid- or pantoic acid-based prodrug. Examples of functionalized protected 2,2-disubstituted pent-4-en-ol derivative 49 include 3-O-protected 2,2-dimethylpent-4-en-1-ol or derivative thereof.

Scheme 18



[0222] In certain embodiments of Scheme 18, the protected neopentyl alcohol 49 is a 3-O-protected 2,2-dimethylpent-4-en-1-ol or a derivative thereof, wherein each of R² and R³ is methyl, R^b is hydrogen or methoxycarbonyl, PG is benzyl, para-methoxybenzyl, or tert-butyl dimethylsilyl (TBDMS), X is chlorine, Q is NHAc, chlorine or 1,3-dioxobenzo[c]azolin-2-yl (phthalyl)-and the activated sulfonic acid derivative 48 is either 3-chloropropylsulfonyl chloride, 2-[3-(chlorosulfonyl)propyl]benzo[c]azolin-1,3-dione, or N-[3-(chlorosulfonyl)propyl]acetamide. Functionalized 3-O-protected 2,2-dimethylpent-4-en-1-ol 49 or derivative thereof can be reacted with 3-chloropropylsulfonyl chloride or 2-[3-(chlorosulfonyl)propyl]benzo[c]azolin-1,3-dione in an inert solvent such as dichloromethane (DCM) in the presence of a base such as triethylamine (Et₃N, TEA), pyridine, or diiso-


propylethylamine (iPr₂EtN, DIEA) and a nucleophilic catalyst such as 4-(N,N-dimethyl)pyridine (DMAP) at a temperature from about -20° C. to about 25° C. to provide the corresponding neopentyl sulfonyl ester intermediate 50 as a precursor or intermediate to externally masked neopentyl sulfonyl ester prodrug 52.

[0223] The terminal alkene group of unsaturated neopentyl sulfonyl ester intermediate or coupling product 50 can be converted to a carboxylic acid derivative, such as an ester or amide derivative, to provide the corresponding functionalized externally masked neopentyl sulfonyl ester prodrug based on the pantoic acid scaffold. Methods for converting terminal carbon-carbon double bonds to ester derivatives are disclosed herein and shown in Scheme 18. Methods for the oxidative transformation of alkenes to aldehydes or carboxylic acid derivatives are also described in the art.

[0224] In certain embodiments of Scheme 18, each of R² and R³ is methyl, R^b is hydrogen or methoxycarbonyl, PG is benzyl, para-methoxybenzyl, or tert-butyl dimethylsilyl (TBDMS), X is chlorine, and Q is NHAc, chlorine, or 1,3-dioxobenzo[c]azolin-2-yl (phthalyl). Using synthetic methods and reaction conditions as previously described the terminal carbon-carbon double bond of alkene intermediate 50 can be oxidatively cleaved to provide the corresponding aldehyde. Using previously described methods and reaction conditions, aldehyde derivatives can be converted to the corresponding free carboxylic acid derivative 51 and subsequently converted to the corresponding carboxylic ester neopentyl intermediate 52.

[0225] Carboxylic acids are useful precursors for preparing the corresponding carboxylic acid esters of externally masked neopentyl sulfonyl ester intermediates or prodrugs based on functionalized (n-1), (n-1)-bis-substituted n-hydroxy alkanolic acids. Methods for preparing acyloxyalkyl ester derivatives, and alkoxy- or aryloxy-carbonyloxy ester derivatives 55 of (n-1), (n-1)-bis-substituted n-hydroxy alkanolic acid derivatives are shown in Scheme 19 where R², R³, R⁴, R⁵, R⁶, and PG are as defined herein, X is halogen such as chlorine or iodine, Y is oxygen or a bond, and Q is NHAc or an amine precursor. In certain embodiments each of R² and R³ is methyl, R⁴ is —OPG with PG is chosen from benzyl, para-methoxybenzyl, or tert-butyl dimethylsilyl (TBDMS), Y is oxygen, R⁵ is isopropyl, ethyl, cyclohexyl, and R⁶ is hydrogen, methyl or isopropyl.



Scheme 19



[0226] For example, chloromethyl carboxylates can be prepared from carboxylic acids using commercially available chloromethyl chlorosulfate in a biphasic mixture of dichloromethane (DCM) and aqueous sodium bicarbonate in the presence of a phase transfer catalyst such as tetra-n-butylammonium hydrogensulfate ($n\text{Bu}_4\text{NHSO}_4$) at a temperature from about -20°C . to about 25°C . Alternative reaction conditions employ chloriodomethane in the presence of a base such as triethylamine (Et_3N , TEA) in a suitable solvent such as N,N-dimethylformamide. Higher substituted 1-halogenoalkyl carboxylates can also be obtained from appropriately functionalized carboxylic acid chlorides and aldehydes in the presence of zinc(II) chloride either neat or in an inert solvent such as dichloromethane (DCM) at a temperature from about -20°C . to about 40°C . Furthermore, 1-halogenoalkyl alkyl- or aryl-carbonates 54 can be prepared from 1-halogenoalkyl chloroformates such as 1-chloroethyl chloroformate, and an alcohol in the presence of a base such as pyridine or triethylamine (Et_3N , TEA) in an inert solvent such as dichloromethane (DCM) at a temperature from about -20°C . to about 25°C .

[0227] Acyloxyalkyl ester derivatives or alkoxy- and aryloxy-carbonyloxy ester derivative 55 can be obtained by reacting carboxylic acid derivative 53 with a substituted 1-halogenoalkyl carboxylate, or a 1-halogenoalkyl alkyl- or aryl-carbonate 54 in the presence of a mild base and halide scavenger such silver(I) carbonate (Ag_2CO_3), silver(I) oxide (Ag_2O), mercury(II) oxide (HgO), or others, either in an inert organic solvent such as toluene, or optionally, neat at a temperature from about 0°C . to about 100°C .

[0228] Pantoic acid derived functionalized neopentyl sulfonyl ester intermediate 56 can be further modified to provide externally masked neopentyl sulfonyl ester prodrugs. Scheme 20 shows methods for converting the functional group Q of an acamprosate precursor to form a N-acetylamino (NHAc) functionality, where R^1 is as described herein, PG is a protecting group, and Q is an amine precursor. Depending on the nature of functional group Q, different synthetic methods can be used.



[0229] In certain embodiments of Scheme 20, R^1 is as defined herein, PG is benzyl or para-methoxybenzyl, and Q is chlorine or 1,3-dioxobenzoc[j]azolin-2-yl (phthalyl). For example, when functional group Q of intermediate 56 is 1,3-dioxobenzoc[j]azolin-2-yl (phthalyl), common synthetic protocols such as the Ing-Manske exchange procedure can be used to liberate the free amino group. When Q is 1,3-dioxobenzoc[j]azolin-2-yl, externally masked neopentyl sulfonyl ester intermediate 56 can be reacted with an excess of hydrazine (H_2NNH_2) in a solvent such as ethyl acetate (EtOAc) and ethanol (EtOH) or mixtures thereof at temperatures from about 0°C . to about 60°C . to provide the corresponding free amine 57. Liberation of the free amine from phthalimide derivative 57 can also be accomplished by reacting with sodium sulfide (Na_2S) in aqueous tetrahydrofuran (THF) or acetone, sodium borohydride (NaBH_4)/acetic acid, or using acid or base catalyzed hydrolysis. Alternative useful and commonly used agents to deprotect phthalimide protecting groups include methylamine (MeNH_2) in ethanol (EtOH) or methanol (MeOH) at room temperature (Motawia et al., *Synthesis* 1989, 384; and Smith et al., *J. Am. Chem. Soc.* 1992, 114, 3134-3136), and n-butylamine ($n\text{BuNH}_2$) in methanol (MeOH) at reflux (Durette et al, *Tetrahedron Lett.* 1979, 42, 4013-4016).

[0230] Free amine 57 can be converted to the corresponding protected intermediate 58 by reaction with an acetylation agent such as acetic anhydride (Ac_2O) or acetyl chloride (AcCl) in a solvent such as dichloromethane (DCM), tetrahydrofuran (THF), or pyridine, optionally in the presence of a base such as pyridine, triethylamine (Et_3N , TEA), or diisopropylethylamine ($i\text{Pr}_2\text{EtN}$, DIEA) and/or a nucleophilic acylation catalyst such 4-(N,N-dimethylaminopyridine) (DMAP) at temperature from about -20°C . to about 40°C . to provide corresponding protected sulfonyl ester intermediate 58.

[0231] In certain embodiments of Scheme 20 where the functional group Q of intermediate 56 is chlorine, the chloro substituent can be converted to an N-acetyl functionality (Q is NHAc) using methods known in the art. For example, intermediate 56 can be reacted with a reagent capable of providing an azide-nucleophile such as sodium azide (NaN_3) or tetrabutylammonium azide ($n\text{Bu}_4\text{NN}_3$), in a polar non-protic solvent such as anhydrous dimethyl sulfoxide (DMSO), anhydrous N,N-dimethylformamide (DMF), acetonitrile (H_3CCN), or mixtures thereof, at a temperature from about 0°C . to about 100°C ., to provide the organic primary azide that can be isolated in pure form. Optionally, the azide can be used directly in the next step following aqueous work-up.

[0232] Primary azides (Q is N_3) can be converted to free amine intermediates that can be isolated in pure form as salts of mineral acids such as hydrogen chloride (HCl) or organic acids such as acetic acid ($\text{H}_3\text{CCO}_2\text{H}$) or trifluoroacetic acid ($\text{F}_3\text{CCO}_2\text{H}$), and the like. Appropriate reagents and reaction conditions for orthogonal and selective deprotection sequences and functional group interconversions can depend

on the nature of the substituents and determined by those skilled in the art. In certain embodiments, where Q is azido, an azide-containing intermediate can be reacted with azide-reducing agents commonly used in similar chemical transformations. For example, azide reducing agents such as hydrogen (H_2) in the presence of a catalyst such as palladium on activated carbon, and a solvent such as methanol (MeOH), ethanol (EtOH), ethyl acetate (EtOAc), or mixtures of any of the foregoing under a pressure from about atmospheric pressure to about 100 psi at a temperature from about 0° C. to about 100° C. can be used.

[0233] Alternatively, the azide functionality can be reduced using metal salts such as stannous chloride ($SnCl_2$) in a protic solvent such as methanol (MeOH), at a temperature from about 0° C. to about 60° C., or, using aryl- or alkylphosphines such as triphenylphosphine (Ph_3P) or tributylphosphine (nBu_3P) in a solvent mixture such as tetrahydrofuran (THF) and water, at a temperature from about 0° C. to about 60° C. The corresponding amine intermediates 57 are provided in either free amine (Q is NH_2), or N-protonated, i.e. ammonium form (Q is NH_3^+), where the counter ion is Cl^- , $H_3CCO_2^-$, $F_3CCO_2^-$, and the like.

[0234] Free amine 57 obtained from reduction of the azide may be directly converted to the corresponding N-acetyl derivative 58 by reacting with an acetylation agent such as acetic anhydride (Ac_2O) or acetyl chloride ($AcCl$) in a solvent such as dichloromethane (DCM), tetrahydrofuran (THF), or pyridine, optionally in the presence of a base such as pyridine, triethylamine (Et_3N , TEA), or diisopropylethylamine (iPr_2EtN , DIEA), and/or a nucleophilic acylation catalyst such as 4-(N,N-dimethylaminopyridine) (DMAP) at a temperature from about -20° C. to about 40° C. to provide the corresponding N-acetylated O-protected neopentyl sulfonyl ester prodrug 58 (Q is $NHAc$). In certain embodiments, acetylation agents, bases, and/or catalysts may be added during or immediately following the azide-reducing step.

[0235] In certain embodiments where PG is benzyl or para-methoxybenzyl and using hydrogenolysis conditions similar to those described herein, O-protecting groups can be cleaved simultaneously with the reduction of the azide-functionality to the corresponding amine to provide the free hydroxyl derivative. In embodiments where the O-deprotection is complete, selective N-acetylation with any of the aforementioned acetylation agents such as acetic anhydride (Ac_2O), acetyl chloride ($AcCl$) using similar reagents such as bases and/or catalysts as described herein may directly provide externally masked neopentyl sulfonyl ester acamprosate prodrugs 59 that incorporate the pantoic acid scaffold as a promoiety.

[0236] In certain embodiments of Scheme 20, where PG is benzyl or para-methoxybenzyl, O-protected externally masked neopentyl sulfonyl ester derivative 58 can be O-deprotected to generate the corresponding externally masked neopentyl sulfonyl ester acamprosate prodrug that incorporates the pantoic acid scaffold as a promoiety 59. The choice of reagents and reaction conditions will depend on the nature of the substituents. For example, in embodiments where PG is benzyl or para-methoxybenzyl, the reducing agent can be hydrogen (H_2), the catalyst can be palladium on activated carbon, and the solvent can be methanol (MeOH), ethanol (EtOH), or ethyl acetate (EtOAc), and reacted under a pressure from about atmospheric pressure to about 100 psi at a temperature from about 0° C. to about 100° C. In certain embodiments, the addition of a catalytic amount of an organic acid, i.e. acetic acid ($HOAc$), i.e. mineral acids (HCl), or other


acidic reagents may activate the catalyst system and facilitate the conversion rate for the transformation.

[0237] In certain embodiments of Scheme 20 where PG is para-methoxybenzyl, O-protected externally masked neopentyl sulfonyl ester derivative 58 may be O-deprotected using orthogonally applicable reagents and reaction conditions to provide the corresponding externally masked neopentyl sulfonyl ester prodrug 59. For example, reacting an O-protected externally masked neopentyl sulfonyl ester acamprosate prodrugs that incorporate the pantoic acid scaffold with an excess of 2,3-dichloro-5,6-dicyano-1,4-benzoquinone (DDQ) in a mixture of solvents such as dichloromethane (DCM) at a temperature from about 0° C. to about 40° C. provides the corresponding externally masked neopentyl sulfonyl ester acamprosate prodrug containing the pantoic acid scaffold as a promoiety 58. Examples of additional reagents and reaction conditions for this transformation include ceric ammonium nitrate ($(H_4N)_2Ce(NO_3)_6$, CAN) and solvents such as water, dichloromethane, or acetonitrile.


[0238] Referring to Scheme 20, selective de-silylation of O-silyl-protected acamprosate prodrug 58 can be accomplished with methods well known in the art to provide acamprosate prodrug 59. For example, trialkylsilyl or mixed alkylarylsilyl-protected derivative 58 can be selectively deprotected using fluoride-containing agents such as tetrabutylammonium fluoride (TBAF), potassium fluoride (KF), ammonium fluoride (H_4NF), or hydrogen fluoride (HF); or using hydrogen fluoride complexes with organic bases such as triethylamine trihydrofluoride ($Et_3N \cdot 3HF$) or pyridinium hydrofluoride; in an inert solvent such as tetrahydrofuran (THF); at a temperature from about 0° C. to about 100° C. to provide the corresponding desilylated acamprosate prodrug 59.

[0239] Alternative routes for preparing sulfonyl ester prodrugs, such as acamprosate prodrugs, from activated sulfonic acid intermediates are shown in Scheme 21. Functionalized 2,2-bis-substituted □-O-protected diol (functionalized neopentyl alcohols) 62, as functionalized coupling partners to activated sulfonic acid derivatives provide useful functionalized and protected neopentyl sulfonyl ester intermediates. The intermediates can be further modified to provide externally masked neopentyl sulfonyl ester prodrugs bearing functionalized (n-1),(n-1)-bis-substituted n-hydroxy alkananoic acid scaffolds in the promoiety and the parent alkananoic acid on which the prodrugs are based is either propanoic acid, butanoic acid, or pentanoic acid. R^1 , R^2 , R^3 , R^4 , Q, X, and PG are defined therein.


Scheme 21



-continued



-continued




[0240] Referring to Scheme 22 (where n , R^2 , R^3 , R^4 , Z , and PG are as defined herein), potentially useful starting materials and methods for preparing functionalized 2,2-bis-substituted ω -O-protected diol (functionalized neopentyl alcohol) 65 are useful as coupling partners with sulfonyl chlorides are either described in the art or will be readily apparent to those skilled in the art. Examples of useful protecting groups (PG) for functionalized 2,2-bis-substituted ω -O-protected diol 65 include benzyl or tris-alkylsilyls and mixed tris-alkyl-arylsilyls.


Scheme 22



[0241] Activated sulfonic acid derivatives can be reacted with functionalized 2,2-bis-substituted ω -O-protected diol 65 to provide useful neopentyl sulfonyl ester intermediates. Referring to Scheme 23, in certain embodiments where each of R^2 and R^3 is methyl, R^4 is hydrogen, each of X and Q is chlorine, PG is benzyl, functionalized 2,2-bis-substituted ω -O-protected diol 68 is 2,2-dimethyl-5-benzylpentan-1-ol, and the activated sulfonic acid derivative 67 is 3-chloropropylsulfonyl chloride. Functionalized 2,2-bis-substituted ω -O-protected diol 68 can be reacted with 3-chloropropylsulfonyl chloride 67 using reaction conditions as described herein. The reaction can be carried out in an inert solvent such as dichloromethane (DCM) in the presence of a base such as triethylamine (Et_3N , TEA), pyridine, or diisopropylethylamine (iPr_2EtN , DIEA) and a nucleophilic catalyst such as 4-(N,N -dimethyl)pyridine (DMAP) at a temperature from about -20°C . to about 25°C . to provide the corresponding protected neopentyl sulfonyl ester intermediate 69.

Scheme 23





[0242] Protected neopentyl sulfonyl ester intermediate 69 can be derivatized after coupling to provide the externally masked neopentyl sulfonyl ester intermediate 70 or if a pro-drug then 73. The chlorine group of intermediate 69 can be converted to the N-acetylamino (NHAc) using methods described herein followed by conversion of the protected hydroxyl group in the ω -position of the promoity precursor to the corresponding carboxylic acid derivative 71 using methods described herein or known in the art.

[0243] In certain embodiments of Scheme 23, each of R^2 and R^3 is methyl, R^4 is hydrogen, m is 2, PG is benzyl, Q is chlorine, and neopentyl alcohol 68 can be derived from 2,2-dimethylglutaric anhydride. Briefly, neopentyl sulfonyl ester intermediate 69, where Q is chlorine, can be reacted with an azide-nucleophile such as sodium azide (NaN_3) or tetrabutylammonium azide (nBu_4NN_3), in a solvent such as dimethyl sulfoxide (DMSO), N,N-dimethylformamide (DMF), or acetonitrile (H_3CCN), at a temperature from about 0°C . to about 100°C ., to provide the organic primary azide. Primary azides (Q is N_3) can be converted to the corresponding free amine intermediate that can be isolated in pure form as a salt of a mineral acid such as hydrogen chloride (HCl) or an organic acid such as acetic acid ($\text{H}_3\text{CCO}_2\text{H}$), trifluoroacetic acid ($\text{F}_3\text{CCO}_2\text{H}$), and the like. Reactants can include a reducing agent such as hydrogen (H_2), a catalyst such as palladium on activated carbon, and a solvent such as methanol (MeOH), ethanol (EtOH), or ethyl acetate (EtOAc) under a pressure from about atmospheric pressure to about 100 psi and at a temperature from about 0°C . to about 100°C . Other useful reducing agents and conditions include metal salts such as

stannous chloride (SnCl_2) in a protic solvent such as methanol (MeOH), at a temperature from about 0°C . to about 60°C ., or, alternatively, aryl- or alkyl-phosphines such as triphenylphosphine (Ph_3P) or tributylphosphine ($\text{n-Bu}_3\text{P}$) in a solvent mixture such as tetrahydrofuran (THF) and water, at a temperature from about 0°C . to about 60°C .

[0244] Free amines or salts thereof can be directly converted to the corresponding N-acetyl derivative 70 by reacting with an acetylation agent such as acetic anhydride (Ac_2O) or acetyl chloride (AcCl) in a solvent such as dichloromethane (DCM), tetrahydrofuran (THF), or pyridine, optionally in the presence of a base such as pyridine, triethylamine (Et_3N , TEA), or diisopropylethylamine (iPr_2EtN , DIEA), and/or a nucleophilic acylation catalyst such as 4-(N,N-dimethylamino)pyridine (DMAP) at a temperature from about -20°C . to about 40°C . to provide corresponding N-acetylated O-protected neopentyl sulfonyl ester intermediate 70 (Q is NHAc). In certain embodiments, acetylation agents, bases, and/or catalysts may be added directly during or immediately following the azido reducing step to provide corresponding N-acetylated O-protected neopentyl sulfonyl ester intermediate 70 (Q is NHAc).

[0245] In certain embodiments, where PG is benzyl or other benzylic protecting group, under hydrogenolysis conditions as described herein, O-protecting groups may be cleaved simultaneously with the reduction of the azide-functionality to provide the corresponding free hydroxyl derivative. In embodiments when O-deprotection is complete, selective N-acetylation with an acetylation agent provides the N-acetylated O-deprotected neopentyl sulfonyl ester intermediate 71

that can be further modified to provide neopentyl sulfonyl ester acamprosate prodrugs that incorporate the (n-1), (n-1)-bis-substituted n-hydroxy alkanic acid scaffold as a promoiety.

[0246] In certain embodiments of Scheme 23, where PG is benzyl or other benzylic protecting group, N-acetylated O-deprotected neopentyl sulfonyl ester intermediate 71 can be prepared via hydrogenolysis of the protecting group using hydrogen (H_2) in the presence of a catalyst such as palladium on activated carbon in a solvent such as methanol (MeOH), ethanol (EtOH), or ethyl acetate (EtOAc) under a pressure from about atmospheric pressure to about 100 psi and at a temperature from about 0° C. to about 100° C. In certain embodiments, the addition of a catalytic amount of an organic acid such as acetic acid (HOAc), mineral acid such as hydrochloric acid (HCl), or other acid can be used to activate the catalyst system and facilitate the conversion rate for the transformation. In embodiments where PG is a tris-alkyl or mixed tris alkyl-arylsilyl protecting group, intermediate 70 can be contacted with a fluoride-containing desilylation agent such as tetrabutylammonium fluoride (TBAF), potassium fluoride (KF), ammonium fluoride (H_4NF), hydrogen fluoride (HF), or hydrogen fluoride complex in an organic base such as triethylamine trihydrofluoride ($Et_3N \cdot 3HF$) or pyridinium hydrofluoride, in an inert solvent such as tetrahydrofuran (THF) at a temperature from about 0° C. to about 100° C. to provide corresponding desilylated intermediate 71. Depending on the nature of PG, other reagents and reactions conditions can be used for the transformation.

[0247] Terminal hydroxyl groups of neopentyl sulfonyl ester intermediate 71 can be converted to aldehydes or carboxylic acids as functionalized building blocks for the preparation of externally masked neopentyl sulfonyl esters prodrugs based on (n-1),(n-1)-bis-substituted n-hydroxy alkanic acid scaffolds. Methods for the oxidative transformation of primary alcohols into aldehydes or carboxylic acids are known to those skilled in the art.

[0248] In certain embodiments of Scheme 23, N-acetyl O-deprotected neopentyl sulfonyl ester intermediate 71 can be oxidized to the corresponding aldehyde 72 or directly to the corresponding carboxylic acid derivative 73. Methods for preparing aldehyde 72 include, for example, reaction with 1,1-tris(acetyloxy)-1,1-dihydro-1,2-benziodoxol-3-(1H)-one (Dess-Martin periodinane) in an inert solvent such as dichloromethane (DCM) at a temperature from about -20° C. to about 25° C. Standard Swern-oxidation conditions such as dimethylsulfoxide (DMSO) in the presence of oxalyl chloride ($(COCl)_2$) and triethylamine (Et_3N , TEA) in dichloromethane (DCM) at a temperature of about -78° C. can also be used. Alternatively, oxidants such as sulfur trioxide-pyridine complex ($SO_3 \cdot Py$) or pyridinium dichromate (PDC) (Cornforth reagent) in an inert solvent such as dichloromethane (DCM) at a temperature from about 0° C. to about 25° C. can be used. Methods for preparing free carboxylic acids include oxidation systems comprising transition metal oxidation catalysts based on ruthenium such as $RuCl_3$ in the presence of the co-oxidant sodium periodate ($NaIO_4$) in a mixture of acetonitrile/carbon tetrachloride/water at a temperature from about 0° C. to about 40° C. Other useful ruthenium compounds include RuO_2 and RuO_4 under similar reaction conditions. In addition, high-valency chromium compounds such as chromium(VI), manganese(VII) com-


pounds, or peroxytungsten compounds, optionally in the presence of a co-oxidant such as bleach ($NaOCl$) or others, can be used.

[0249] Phosphatases are important metabolic enzymes and are classified as phosphoric monoester hydrolases (phosphatases, E.C. 3.1.3.), phosphoric diester hydrolases (phosphodiesterases, E.C. 3.1.4.), triphosphoric monoester hydrolases (E.C. 3.1.5.), diphosphoric monoester hydrolases (pyrophosphatases, E.C. 3.1.7.), and phosphoric ester triester hydrolases (E.C. 3.1.8). Some of these enzymes are active in the de-phosphorylation reaction of xenobiotic organophosphorus compounds, including, for example, alkaline phosphatase, E.C. 3.1.3.1., and others. Phosphate conjugates of pharmaceutical interest are often monoesters. Enzymes believed to be able to dephosphorylate phosphate conjugates via hydrolysis or transphosphorylation include alkaline phosphatases and acid phosphatases. The de-phosphorylation reactions often proceed with high catalytic efficiency in vitro and also in vivo to provide the parent drug. Alkaline phosphatase is a ubiquitous, extracellularly bound to membranes, and widely expressed non-specific esterase of phosphoric monoesters in mammals with an optimal pH for catalysis at about 8.0 and above. The enzyme is found particularly in the gastrointestinal tract, pancreas, liver, bile, placenta, and osteoplasts.

[0250] Incorporation of a phosphate group to has successfully overcome numerous drug delivery problems. Phosphate moieties can either be directly incorporated via a covalent bond to an alcoholic or phenolic hydroxyl group of a parent drug or prodrug (in form of a monoester) or via a chemical linker. Generally these phosphomonoester prodrugs are chemically very stable with long attractive shelf-lives. Introduction of a phosphate group into a xenobiotic often constructively influences a multitude of physiochemical and biological/pharmacokinetic parameters of a drug or prodrug including increasing the aqueous solubility, improving the parenteral dosing regime, and modulating specific pharmacokinetic parameters, i.e., half life (t_{max}), peak concentrations (c_{max}), or concentration time curves (AUCs).

[0251] Examples for phosphate ester prodrugs of alcohols and phenols include the gram-positive antibiotic clindamycin phosphate, the broad-spectrum antifungal fosfluconazole, the orally active human immunodeficiency virus (HIV) inhibitor fosamprenavir, the antineoplastic etoposide phosphate, and double prodrugs such as GPI 15715.

[0252] Methods for preparing phosphate conjugates of acamprosate prodrugs or derivatives thereof, with useful physiochemical, biological, or pharmacokinetic properties are provided in Scheme 24, where R^1 , R^2 are defined herein, Y is oxygen or bond. For example and referring to Scheme 24, using methods well known in the art, the secondary hydroxyl group of acamprosate prodrugs 74, or any other hydroxyl group optionally incorporated in the R^1 and R^2 groups, can be phosphorylated with activated bis-O-protected phosphoric acid diester derivatives to provide the corresponding phosphorylated intermediate 75. The protecting groups of the phosphoric acid moiety are chosen in a way that the protecting groups in compound 75 can be removed orthogonally without affecting any of the additional functional groups to provide the free phosphate conjugate 76.



[0253] In certain embodiments of Scheme 24, where R^1 and R^2 are defined herein, Y is oxygen or bond, the free secondary hydroxyl group of acamprosate prodrug 74 can be reacted with protected and activated phosphorous acid derivatives such as commercially available diphenyl chlorophosphate $(PhO)_2POCl$, dibenzyl chlorophosphate $(BnO)_2POCl$, or di-tert-butyl chlorophosphate $(tBuO)_2POCl$ in a suitable solvent such as dichloromethane (DCM) and in the presence of suitable bases such as triethylamine (Et_3N , TEA), diisopropylethylamine $[(iPr)_2EtN]$, DIEA, Hünig's base, pyridine, and optionally in the presence of catalysts such as 4-(N,N-dimethylamino)pyridine (DMAP) at a temperature from about $0^\circ C$. to about $40^\circ C$.

[0254] Referring to Scheme 24, intermediate 75 can be deprotected and converted to the free phosphate monoesters of acamprosate prodrug 76 with methods known in the art where R^1 and R^2 are defined herein, Y is oxygen or bond. In certain embodiments, PG is phenyl and the phenyl groups are removed via hydrogenolysis under a hydrogen atmosphere format a pressure from about 1 atm to about 100 psi and at a temperature from about $0^\circ C$. to about $60^\circ C$. employing platinum(IV)-based heterogeneous catalysts such as platinum (IV) oxide $Pt(IV)O_2$ or hydrates thereof, i.e. $Pt(IV)O_2 \cdot xH_2O$, $x \sim 1$) (Adam's catalysts) in suitable solvents such as methanol (MeOH), ethanol (EtOH), water, or mixtures thereof optionally in the presence of a trace amount of an acidic additive

such as an organic acid, i.e. acetic acid (HOAc) or diluted mineral acid such as one molar (1.0 M) hydrochloric acid (HCl). If PG in compound 75 is chosen from Bn, then the heterogeneous catalyst can also be palladium on activated carbon in a solvent such as methanol (MeOH), ethyl acetate (EtOAc), mixtures thereof, or others (Scheme 24). If PG in compound 75 is optionally chosen from tBu then the protecting groups can be removed by contacting compounds 75 at temperatures from about $0^\circ C$. to about $40^\circ C$. with strong acids such as trifluoroacetic acid (TFA) in a suitable solvent such as dichloromethane (DCM) or, optionally, with hydrogen chloride (HCl) in 1,4-dioxane or diethyl ether (Et_2O).

Pharmaceutical Compositions

[0255] Pharmaceutical compositions provided by the present disclosure comprise a compound of Formula (I) together with a suitable amount of one or more pharmaceutically acceptable vehicles so as to provide a composition for proper administration to a patient. Examples of suitable pharmaceutical vehicles are known in the art.

[0256] Pharmaceutical compositions comprising a compound of Formula (I) may be manufactured by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping, or lyophilizing processes. Pharmaceutical compositions may be formulated in a conventional manner using one or more physiologically acceptable carriers, diluents, excipients, or auxiliaries, which facilitate processing of compounds of Formula (I) or crystalline form thereof and one or more pharmaceutically acceptable vehicles into formulations that can be used pharmaceutically. Proper formulation is dependent upon the route of administration chosen. In certain embodiments, a pharmaceutical composition comprising a compound of Formula (I) or crystalline form thereof may be formulated for oral administration, and in certain embodiments for sustained release oral administration. Pharmaceutical compositions provided by the present disclosure may take the form of solutions, suspensions, emulsion, tablets, pills, pellets, capsules, capsules containing liquids, powders, sustained-release formulations, suppositories, emulsions, aerosols, sprays, suspensions, or any other form suitable for administration to a patient.

[0257] Pharmaceutical compositions provided by the present disclosure may be formulated in a unit dosage form. A unit dosage form refers to a physically discrete unit suitable as a unitary dose for patients undergoing treatment, with each unit containing a predetermined quantity of at least one compound of Formula (I) calculated to produce an intended therapeutic effect. A unit dosage form may be for a single daily dose, for administration 2 times per day, or one of multiple daily doses, e.g., 3 or more times per day. When multiple daily doses are used, a unit dosage may be the same or different for each dose. One or more dosage forms may comprise a dose, which may be administered to a patient at a single point in time or during a time interval.

[0258] In certain embodiments, a compound of Formula (I) may be incorporated into pharmaceutical compositions to be administered orally. Oral administration of such pharmaceutical compositions may result in uptake of a compound of Formula (I) throughout the intestine and entry into the systemic circulation. Such oral compositions may be prepared in a manner known in the pharmaceutical art and comprise at least one compound of Formula (I) and at least one pharmaceutically acceptable vehicle. Oral pharmaceutical composi-

tions may include a therapeutically effective amount of at least one compound of Formula (I) and a suitable amount of a pharmaceutically acceptable vehicle, so as to provide an appropriate form for administration to a patient.

[0259] Pharmaceutical compositions comprising at least one compound of Formula (I) may be formulated for immediate release for parenteral administration, oral administration, or for any other appropriate route of administration.

[0260] Controlled drug delivery systems may be designed to deliver a drug in such a way that the drug level is maintained within a therapeutically effective window and effective and safe blood levels are maintained for a period as long as the system continues to deliver the drug at a particular rate. Controlled drug delivery may produce substantially constant blood levels of a drug over a period of time as compared to fluctuations observed with immediate release dosage forms. For some drugs, maintaining a constant blood and tissue concentration throughout the course of therapy is the most desirable mode of treatment. Immediate release of drugs may cause blood levels to peak above the level required to elicit a desired response, which may waste the drug and may cause or exacerbate toxic side effects. Controlled drug delivery can result in optimum therapy, and not only can reduce the frequency of dosing, but may also reduce the severity of side effects. Examples of controlled release dosage forms include dissolution controlled systems, diffusion controlled systems, ion exchange resins, osmotically controlled systems, erodible matrix systems, pH independent formulations, gastric retention systems, and the like.

[0261] In certain embodiments, an oral dosage form provided by the present disclosure may be a controlled release dosage form. Controlled delivery technologies can improve the absorption of a drug in a particular region or regions of the gastrointestinal tract.

[0262] The appropriate oral dosage form for a particular pharmaceutical composition provided by the present disclosure may depend, at least in part, on the gastrointestinal absorption properties of a compound of Formula (I), the stability of a compound of Formula (I) in the gastrointestinal tract, the pharmacokinetics of a compound of Formula (I), and the intended therapeutic profile. An appropriate controlled release oral dosage form may be selected for a particular compound of Formula (I). For example, gastric retention oral dosage forms may be appropriate for compounds absorbed primarily from the upper gastrointestinal tract, and sustained release oral dosage forms may be appropriate for compounds absorbed primarily from the lower gastrointestinal tract. Certain compounds are absorbed primarily from the small intestine. In general, compounds traverse the length of the small intestine in about 3 to 5 hours. For compounds that are not easily absorbed by the small intestine or that do not dissolve readily, the window for active agent absorption in the small intestine may be too short to provide a desired therapeutic effect.

[0263] Gastric retention dosage forms, i.e., dosage forms that are designed to be retained in the stomach for a prolonged period of time, may increase the bioavailability of drugs that are most readily absorbed by the upper gastrointestinal tract. For example, certain compounds of Formula (I) may exhibit limited colonic absorption, and be absorbed primarily from the upper gastrointestinal tract. Thus, dosage forms that release a compound of Formula (I) in the upper gastrointestinal tract and/or retard transit of the dosage form through the upper gastrointestinal tract will tend to enhance the oral bio-

availability of the compound of Formula (I). The residence time of a conventional dosage form in the stomach is about 1 to about 3 hours. After transiting the stomach, there is approximately a 3 to 5 hour window of bioavailability before the dosage form reaches the colon. However, if the dosage form is retained in the stomach, the drug may be released before it reaches the small intestine and will enter the intestine in solution in a state in which it can be more readily absorbed. Another use of gastric retention dosage forms is to improve the bioavailability of a drug that is unstable to the basic conditions of the intestine.

[0264] In certain embodiments, pharmaceutical compositions provided by the present disclosure may be practiced with dosage forms adapted to provide sustained release of a compound of Formula (I) upon oral administration. Sustained release oral dosage forms may be used to release drugs over a prolonged time period and are useful when it is desired that a drug or drug form be delivered to the lower gastrointestinal tract. Sustained release oral dosage forms include any oral dosage form that maintains therapeutic concentrations of a drug in a biological fluid such as the plasma, blood, cerebrospinal fluid, or in a tissue or organ for a prolonged time period. Sustained release oral dosage forms include diffusion-controlled systems such as reservoir devices and matrix devices, dissolution-controlled systems, osmotic systems, and erosion-controlled systems. Sustained release oral dosage forms and methods of preparing the same are well known in the art.

[0265] Sustained release oral dosage forms may be in any appropriate form for oral administration, such as, for example, in the form of tablets, pills, or granules. Granules can be filled into capsules, compressed into tablets, or included in a liquid suspension. Sustained release oral dosage forms may additionally include an exterior coating to provide, for example, acid protection, ease of swallowing, flavor, identification, and the like.

[0266] In certain embodiments, sustained release oral dosage forms may comprise a therapeutically effective amount of a compound of Formula (I) and at least one pharmaceutically acceptable vehicle. In certain embodiments, a sustained release oral dosage form may comprise less than a therapeutically effective amount of a compound of Formula (I) and a pharmaceutically effective vehicle. Multiple sustained release oral dosage forms, each dosage form comprising less than a therapeutically effective amount of a compound of Formula (I) may be administered at a single time or over a period of time to provide a therapeutically effective dose or regimen for treating a disease in a patient. In certain embodiments, a sustained release oral dosage form comprises more than one compound of Formula (I).

[0267] Sustained release oral dosage forms provided by the present disclosure can release a compound of Formula (I) from the dosage form to facilitate the ability of the compound of Formula (I) to be absorbed from an appropriate region of the gastrointestinal tract, for example, in the small intestine or in the colon. In certain embodiments, sustained release oral dosage forms may release a compound of Formula (I) from the dosage form over a period of at least about 4 hours, at least about 8 hours, at least about 12 hours, at least about 16 hours, at least about 20 hours, and in certain embodiments, at least about 24 hours. In certain embodiments, sustained release oral dosage forms may release a compound of Formula (I) from the dosage form in a delivery pattern corresponding to about 0 wt % to about 20 wt % in about 0 to about 4 hours;

about 20 wt % to about 50 wt % in about 0 to about 8 hours; about 55 wt % to about 85 wt % in about 0 to about 14 hours; and about 80 wt % to about 100 wt % in about 0 to about 24 hours; where wt % refers to the percent of the total weight of the compound in the dosage form. In certain embodiments, sustained release oral dosage forms may release a compound of Formula (I) from the dosage form in a delivery pattern corresponding to about 0 wt % to about 20 wt % in about 0 to about 4 hours; about 20 wt % to about 50 wt % in about 0 to about 8 hours; about 55 wt % to about 85 wt % in about 0 to about 14 hours; and about 80 wt % to about 100 wt % in about 0 to about 20 hours. In certain embodiments, sustained release oral dosage forms may release a compound of Formula (I) from the dosage form in a delivery pattern corresponding to about 0 wt % to about 20 wt % in about 0 to about 2 hours; about 20 wt % to about 50 wt % in about 0 to about 4 hours; about 55 wt % to about 85 wt % in about 0 to about 7 hours; and about 80 wt % to about 100 wt % in about 0 to about 8 hours.

[0268] Sustained release oral dosage forms comprising a compound of Formula (I) may provide a concentration of the corresponding drug in the plasma, blood, cerebrospinal fluid, or tissue of a patient over time, following oral administration to the patient. The concentration profile of the drug may exhibit an AUC that is proportional to the dose of the corresponding compound of Formula (I).

[0269] Regardless of the specific type of controlled release oral dosage form used, a compound of Formula (I) may be released from an orally administered dosage form over a sufficient period of time to provide prolonged therapeutic concentrations of the compound of Formula (I) in the plasma and/or blood of a patient. Following oral administration, a dosage form comprising a compound of Formula (I) may provide a therapeutically effective concentration of the corresponding drug in the plasma and/or blood of a patient for a continuous time period of at least about 4 hours, of at least about 8 hours, for at least about 12 hours, for at least about 16 hours, and in certain embodiments, for at least about 20 hours following oral administration of the dosage form to the patient. The continuous time periods during which a therapeutically effective concentration of the drug is maintained may be the same or different. The continuous period of time during which a therapeutically effective plasma concentration of the drug is maintained may begin shortly after oral administration or following a time interval.

[0270] An appropriate dosage of a compound of Formula (I) or pharmaceutical composition comprising a compound of Formula (I) may be determined according to any one of several well-established protocols. For example, animal studies such as studies using mice, rats, dogs, and/or monkeys may be used to determine an appropriate dose of a pharmaceutical compound. Results from animal studies may be extrapolated to determine doses for use in other species, such as for example, humans.

Uses

[0271] Compounds of Formula (I) are prodrugs of acamprosate. Thus, compounds of Formula (I) may be administered to a patient suffering from any disease including a disorder, condition, or symptom for which acamprosate is known or hereafter discovered to be therapeutically effective. Methods for treating a disease in a patient provided by the present

disclosure comprise administering to a patient in need of such treatment a therapeutically effective amount of at least one compound of Formula (I).

[0272] Compounds of Formula (I) or pharmaceutical compositions thereof may provide therapeutic or prophylactic plasma and/or blood concentrations of the corresponding drug following oral administration to a patient. The promoiety(ies) of compounds of Formula (I) may be cleaved in vivo either chemically and/or enzymatically to release the parent drug. One or more enzymes present in the intestinal lumen, intestinal tissue, blood, liver, brain, or any other suitable tissue of a patient may enzymatically cleave the promoiety of the administered compounds. For example, a promoiety of a compound of Formula (I) may be cleaved following absorption of the compound from the gastrointestinal tract (e.g., in intestinal tissue, blood, liver, or other suitable tissue of a mammal). For compounds of Formula (I) the masking promoiety is first cleaved enzymatically, chemically, or by both mechanisms to provide a neopentyl promoiety terminated with a nitrogen or oxygen nucleophile. The structures of the oxygen nucleophile metabolic intermediates have the structures of Formula (II). The nucleophilic group can then internally cyclize to release acamprosate.

[0273] In certain embodiments, compounds of Formula (I) may be actively transported across the intestinal endothelium by transporters expressed in the gastrointestinal tract including the small intestine and colon. The drug, e.g., acamprosate, may remain conjugated to the promoiety during transit across the intestinal mucosal barrier to prevent or minimize presystemic metabolism. In certain embodiments, a compound of Formula (I) is essentially not metabolized to acamprosate within gastrointestinal enterocytes, but is metabolized to release acamprosate within the systemic circulation, for example in the intestinal tissue, blood/plasma, liver, or other suitable tissue of a mammal. In such embodiments, compounds of Formula (I) may be absorbed into the systemic circulation from the small and large intestines either by active transport, passive diffusion, or by both active and passive processes. For example, a promoiety may be cleaved after absorption from the gastrointestinal tract, for example, in intestinal tissue, blood, liver, or other suitable tissue of a mammal.

[0274] Compounds of Formula (I) may be administered in similar equivalent amounts of acamprosate and using a similar dosing schedule as described in the art for treatment of a particular disease. For example, in a human subject weighing about 70 kg, compounds of Formula (I) may be administered at a dose over time having an equivalent weight of acamprosate from about 10 mg to about 10 g per day, and in certain embodiments, an equivalent weight of acamprosate from about 1 mg to about 3 g per day. A dose of a compound of Formula (I) taken at any one time can have an equivalent weight of acamprosate from about 1 mg to about 3 g. An acamprosate dose may be determined based on several factors, including, for example, the body weight and/or condition of the patient being treated, the severity of the disease being treated, the incidence of side effects, the manner of administration, and the judgment of the prescribing physician. Dosage ranges may be determined by methods known to one skilled in the art. In certain embodiments, compounds of Formula (I) provide a higher oral bioavailability of acamprosate compared to the oral bioavailability of acamprosate itself when orally administered at an equivalent dose and in an equivalent dosage form. Consequently, a lesser equivalent

amount of acamprosate derived from a compound of Formula (I) may be orally administered to achieve the same therapeutic effect as that achieved when acamprosate itself is orally administered.

[0275] Compounds of Formula (I) may be assayed in vitro and in vivo for the desired therapeutic or prophylactic activity prior to use in humans. For example, in vitro assays may be used to determine whether administration of a compound of Formula (I) is a substrate of a transporter protein, including transporters expressed in the gastrointestinal tract. Examples of certain assay methods applicable to analyzing the ability of compounds of Formula (I) to act as substrates for one or more transporter proteins are disclosed in Zerangue et al., US 2003/0158254. In vivo assays, for example using appropriate animal models, may also be used to determine whether administration of a compound of Formula (I) is therapeutically effective. Compounds of Formula (I) may also be demonstrated to be therapeutically effective and safe using animal model systems.

[0276] In certain embodiments, a therapeutically effective dose of a compound of Formula (I) may provide therapeutic benefit without causing substantial toxicity. Toxicity of compounds of Formula (I) prodrugs, and/or metabolites thereof may be determined using standard pharmaceutical procedures and may be ascertained by one skilled in the art. The dose ratio between toxic and therapeutic effect is the therapeutic index. A dose of a compound of Formula (I) may be within a range capable of establishing and maintaining a therapeutically effective circulating plasma and/or blood concentration of a compound of Formula (I) or acamprosate that exhibits little or no toxicity.

[0277] Compounds of Formula (I) may be used to treat diseases, disorders, conditions, and symptoms of any of the foregoing for which acamprosate is shown to provide therapeutic benefit. Hence, compounds of Formula (I) be used to treat neurodegenerative disorders, psychotic disorders, mood disorders, anxiety disorders, somatoform disorders, movement disorders, substance abuse disorders, binge eating disorder, cortical spreading depression related disorders, tinnitus, sleeping disorders, multiple sclerosis, and pain. The underlying etiology of any of the foregoing diseases being so treated may have a multiplicity of origins.

[0278] Further, in certain embodiments, a therapeutically effective amount of one or more compounds of Formula (I) may be administered to a patient, such as a human, as a preventative measure against various diseases or disorders. Thus, a therapeutically effective amount of one or more compounds of Formula (I) be administered as a preventative measure to a patient having a predisposition for a neurodegenerative disorder, a psychotic disorder, a mood disorder, an anxiety disorder, a somatoform disorder, a movement disorder, a substance abuse disorder, binge eating disorder, a cortical spreading depression related disorder, tinnitus, a sleeping disorder, multiple sclerosis, or pain.

[0279] Substance abuse disorders refer to disorders related to taking a drug of abuse, to the side effects of a medication, and to toxin exposure. Drugs of abuse include alcohol, amphetamines, caffeine, cannabis, cocaine, hallucinogens, inhalants, nicotine, opioids, phencyclidine, or similarly acting arylcyclohexylamines, sedatives, hypnotics, and anxiolytics.

[0280] Alcoholism or alcohol dependence is a chronic disorder with genetic, psychosocial, and environmental causes. Alcoholism refers to "... maladaptive alcohol use with clini-

cally significant impairment as manifested by at least three of the following within any one-year period: tolerance; withdrawal; taken in greater amounts or over longer time course than intended; desire or unsuccessful attempts to cut down or control use; great deal of time spent obtaining, using, or recovering from use; social, occupational, or recreational activities given up or reduced; continued use despite knowledge of physical or psychological sequelae." (Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition, Text Revision, Washington D.C., American Psychiatric Association, 2000 (DSM-IV)). Alcohol use disorders include alcohol dependence and alcohol abuse. Screening tests useful for identifying alcoholism include the Alcohol Dependence Data Questionnaire, the Michigan Alcohol Screening Test, the Alcohol Use Disorders Identification Test, and the Paddington Alcohol Test, and other generally recognized tests for diagnosing alcohol dependence.

[0281] Treatment for alcoholism generally includes psychological, social, and pharmacotherapeutic interventions aimed at reducing alcohol-associated problems and usually involves detoxification and rehabilitation phases. Medications useful in the pharmacologic treatment of alcohol dependence include disulfiram and naltrexone.

[0282] Studies suggest that modulation of mGluR⁵ receptors play a role in substance abuse disorders and that mGluR⁵ receptor antagonists such as MPEP may be useful in treating such conditions including drug abuse disorders.

[0283] Acamprosate has been shown to be effective for maintaining abstinence from alcohol in patients with alcohol dependence that are abstinent at the initiation of acamprosate treatment (Scott et al., *CNS Drugs* 2005, 19(5), 445-464; and Heilig and Egli, *Pharmacology & Therapeutics* 2006, 11, 855-876) and as such is marketed in the United States for the treatment of alcohol abstinence as Camprale (Forest Laboratories and Merck KGaA). Typical acamprosate doses range from about 1-2 gm per day to achieve a steady-state plasma concentration of about 370-640 ng/mL, which occurs at about 3-8 hours post-dose (Overman et al., *Annals Pharmacotherapy* 2003, 37, 1090-1099; Paille et al., *Alcohol* 1995, 30, 239-47; and Pelc et al., *Br. Psychiatry* 1997, 171, 73-77) with a recommended dose of Campral® being two to three 333 mg tablets taken three times daily.

[0284] The efficacy of compounds of Formula (I) and compositions thereof for treating alcohol dependency may be assessed using animal models of alcoholism and using clinical studies. Animal models of alcoholism are known. Clinical protocols for assessing the efficacy of a compound of Formula (I) for treating alcoholism are known.

[0285] The effect of acamprosate on relapse in other substances of abuse has not been extensively studied; however administration of 100 mg/kg acamprosate for 3 days attenuated relapse-like behavior in cocaine conditioned mice (McGeehan and Olive, *Behav Pharmacol* 2006, 17(4), 363-7). Studies suggest that modulation of mGluR⁵ receptors play a role in substance abuse disorders and that mGluR⁵ receptor antagonists such as MPEP may be useful in treating such conditions including drug abuse disorders and nicotine abuse disorders. Therefore, acamprosate may have applicability in treating other substance abuse disorders, including narcotic abuse disorders and nicotine abuse disorders.

[0286] Binge eating disorder is characterized by recurrent episodes of distressing, uncontrollable eating of excessively large amounts of food without the inappropriate compensatory weight loss behaviors of bulimia nervosa or anorexia

nervosa (DSM-IV, Fourth Ed., Text Revision, Washington D.C., American Psychiatric Assoc., 2000). The pathophysiology of binge eating disorders is unknown. Binge eating disorder is associated with psychopathology such as compulsive, impulsive, and affective disorders, medical comorbidity, especially obesity, impaired quality of life, and disability. Emotional cues such as anger, sadness, boredom, and anxiety can trigger binge eating. Impulsive behavior and certain other emotional problems can be more common in people with binge eating disorder. Antidepressant medications, including tricyclic antidepressants, selective serotonin re-uptake inhibitors, as well as some of certain antidepressants, have shown evidence of some therapeutic value in binge eating disorder (Bello and Jainal, *Brain Res Bulletin* 2006, 70, 422-429; Buda-Levin et al., *Physiology & Behavior* 2005, 86, 176-184; and Han et al., *Drug Alcohol Dependence* 2007, prepublication no. DAD-3137, 5 pages).

[0287] The efficacy of compounds of acamprosate prodrugs and compositions for treating binge eating may be assessed using animal models of binge eating and using clinical studies. Animal models of binge eating are known. Clinical protocols useful for assessing the efficacy of an acamprosate prodrug for treating binge eating are also known.

[0288] In certain embodiments, compounds of Formula (I) can be used to treat tinnitus. Tinnitus is the perception of sound in the absence of acoustic stimulation and often involves sound sensations such as ringing, buzzing, roaring, whistling, or hissing that cannot be attributed to an external sound source. Tinnitus is a symptom associated with many forms of hearing loss and can also be a symptom of other health problems.

[0289] Tinnitus can be caused by hearing loss, loud noise, medicine, and other health problems such as allergies, head or neck tumors, cardiovascular disorders such as atherosclerosis, high blood pressure, turbulent blood flow, malformation of capillaries, trauma such as excessive exposure to loud noise, long-term use of certain medications such as salicylates, quinine, cisplatin and certain types of antibiotics, changes to ear bones such as otosclerosis, and jaw and neck injuries. In general, insults or damage to the auditory and somatosensory systems can create an imbalance between inhibitory and excitatory transmitter actions in the midbrain, auditory cortex, and brain stem. This imbalance can cause hyperexcitability of auditory neurons that can lead to the perception of phantom sounds. For acute tinnitus such as tinnitus induced by drugs or loud noises, increased spontaneous firing rates in the auditory nerve fibers have been attributed to reduced levels of central inhibition, presumably by GABA, in central auditory structures leading to neural hyperactivity in the inferior colliculus. Although chronic tinnitus may have a different cause than acute tinnitus, reduced GABA levels have also been implicated.

[0290] A recent clinical trial suggests that acamprosate may be effective in treating tinnitus (Azevedo and Figueiredo, *Rev Bras Otorrinolaringol* 2005, 71(5), 618-23).

[0291] Acamprosate prodrugs of Formula (I) can be used to treat tinnitus, including preventing, reducing, or eliminating tinnitus and/or the accompanying symptoms of tinnitus in a patient. Treating tinnitus refers to any indicia of success in prevention, reduction, elimination, or amelioration of tinnitus, including any objective or subjective parameter such as abatement, remission, diminishing of symptoms, prevention, or lessening of tinnitus symptoms or making the condition

more tolerable to the patient, making the tinnitus less debilitating, or improving a patient's physical or mental well-being.

[0292] The efficacy of an acamprosate prodrug of Formula (I) for treating tinnitus can be assessed using animal models of tinnitus and in clinical results. Methods of evaluating tinnitus in animals and humans are known. The ability of a compound of Formula (I) to treat tinnitus in human patients may be assessed using objective and subjective tests such as those described in Bauer and Brozoski, *Laryngoscope* 2006, 116(5), 675-681. An example of a test used in a clinical context to assess tinnitus treatment outcomes is the Tinnitus Handicap Inventory.

[0293] Neurodegenerative diseases are characterized by progressive dysfunction and neuronal death. Neurodegenerative diseases featuring cell death can be categorized as acute, i.e., stroke, traumatic brain injury, spinal cord injury, and chronic, i.e., amyotrophic lateral sclerosis, Huntington's disease, Parkinson's disease, and Alzheimer's disease. Although these diseases have different causes and affect different neuronal populations, they share similar impairment in intracellular energy metabolism NMDA receptor and non-NMDA receptor mediated excitotoxic injury results in neurodegeneration leading to necrotic or apoptotic cell death. Studies also suggest that mGluR⁵ receptor activity is involved in the etiology of neurodegenerative disorders and that mGluR⁵ modulators can be useful in treating movement and cognitive dysfunction associated with neurodegenerative disorders, as well as exhibit neuroprotective effects.

[0294] Parkinson's disease is a slowly progressive degenerative disorder of the nervous system characterized by tremor when muscles are at rest (resting tremor), slowness of voluntary movements, and increased muscle tone (rigidity). In Parkinson's disease, nerve cells in the basal ganglia, e.g., substantia nigra, degenerate, and thereby reduce the production of dopamine and the number of connections between nerve cells in the basal ganglia. As a result, the basal ganglia are unable to smooth muscle movements and coordinate changes in posture as normal, leading to tremor, incoordination, and slowed, reduced movement (bradykinesia).

[0295] Modulators of NMDA receptor activity have shown therapeutic potential in the management of Parkinson's disease, as well as have mGluR⁵ receptor antagonists. Accordingly, acamprosate may be useful in treating Parkinson's disease.

[0296] Studies suggest that agents that NMDA receptor antagonists or mGluR⁵ receptor antagonists are potentially useful for treating levodopa-induced dyskinesias such as levodopa-induced dyskinesias in Parkinson's disease Fabbrini et al., *Movement Disorders* 2007, 22(10), 1379-1389; and Mela et al., *J Neurochemistry* 2007, 101, 483-497). Accordingly, acamprosate prodrugs provided by the present disclosure may be useful in treating a movement disorder such as levodopa-induced dyskinesias in Parkinson's disease.

[0297] The efficacy of a compound of Formula (I) for treating Parkinson's disease may be assessed using animal models of Parkinson's disease and in clinical studies. Animal models of Parkinson's disease are known. The ability of acamprosate prodrugs to mitigate against L-dopa induced dyskinesias can be assessed using animal models and in clinical trials.

[0298] Alzheimer's disease is a progressive loss of mental function characterized by degeneration of brain tissue. In Alzheimer's disease, parts of the brain degenerate, destroying nerve cells and reducing the responsiveness of the maintaining neurons to neurotransmitters. Abnormalities in brain tis-

sue consist of senile or neuritic plaques, e.g., clumps of dead nerve cells containing an abnormal, insoluble protein called amyloid, and neurofibrillary tangles, twisted strands of insoluble proteins in the nerve cell.

[0299] Excitotoxic cell death is thought to contribute to neuronal cell injury and death in Alzheimer's diseases and other neurodegenerative disorders. Excitotoxicity is due, at least in part, to excessive acylation of NMDA-type glutamate receptors and the concomitant excessive Ca^{2+} influx through the receptor's associated ion channel. NMDA receptor antagonists have shown neuroprotective effects in Alzheimer's disease (Lipton, *NeuroRx* 2004, 1(1), 101-110). As a modulator of the NMDA receptor, acamprosate may have similar effects.

[0300] The efficacy of administering a compound of Formula (I) for treating Alzheimer's disease may be assessed using animal models of Alzheimer's disease and in clinical studies. Useful animal models for assessing the efficacy of compounds for treating Alzheimer's disease are known.

[0301] Huntington's disease is an autosomal dominant neurodegenerative disorder in which specific cell death occurs in the neostriatum and cortex. Onset usually occurs during the fourth or fifth decade of life, with a mean survival at age of onset of 14 to 20 years. Huntington's disease is universally fatal, and there is no effective treatment. Symptoms include a characteristic movement disorder (Huntington's chorea), cognitive dysfunction, and psychiatric symptoms. The disease is caused by a mutation encoding an abnormal expansion of CAG-encoded polyglutamine repeats in the protein, huntingtin.

[0302] Neuroprotective effects of NMDA antagonists such as memantine and ketamine in Huntington's disease have been proposed (Murman et al., *Neurology* 1997, 49(1), 153-161; and Kozachuk, US 2004/0102525).

[0303] The efficacy of administering a compound of Formula (I) for treating Huntington's disease may be assessed using animal models of Huntington's disease and in clinical studies. Animal models of Huntington's disease are known.

[0304] Amyotrophic lateral sclerosis (ALS) is a progressive neurodegenerative disorder characterized by the progressive and specific loss of motor neurons in the brain, brain stem, and spinal cord. ALS begins with weakness, often in the hands and less frequently in the feet that generally progresses up an arm or leg. Over time, weakness increases and spasticity develops characterized by muscle twitching and tightening, followed by muscle spasms and possibly tremors.

[0305] A possible cause of ALS is constitutive opening of the calcium pore in glutamate responsive AMPA channels secondary to a failure of RNA editing. Recent work has shown that endogenous polyamines can block the vulnerability of motor neurons to cell death due to calcium influx through Ca^{2+} -permeable AMP receptors. Acamprosate is believed to have an action at AMPA receptors similar to that of endogenous polyamines. Accordingly, it has been proposed that acamprosate may be useful in treating ALS (Kast and Altschuler, *Med Hypotheses* 2007, 69(4), 836-837).

[0306] The efficacy of a compound of Formula (I) for treating ALS may be assessed using animal models of ALS and in clinical studies. Natural disease models of ALS include mouse models (motor neuron degeneration, progressive motor neuropathy, and wobbler) and the hereditary canine spinal muscular atrophy canine model. Experimentally produced and genetically engineered animal models of ALS can also be useful in assessing therapeutic efficacy. Specifically, the

SODI-G93A mouse model is a recognized model for ALS. Examples of clinical trial protocols useful in assessing treatment of ALS are known.

[0307] Multiple sclerosis (MS) is an inflammatory autoimmune disease of the central nervous system caused by an autoimmune attack against the isolating axonal myelin sheets of the central nervous system. Demyelination leads to the breakdown of conduction and to severe disease with destruction of local axons and irreversible neuronal cell death. The symptoms of MS are highly varied with each patient exhibiting a particular pattern of motor, sensory, and sensory disturbances. MS is typified pathologically by multiple inflammatory foci, plaques of demyelination, gliosis, and axonal pathology within the brain and spinal cord, all of which contribute to the clinical manifestations of neurological disability. Although the causal events that precipitate MS are not fully understood, evidence implicates an autoimmune etiology together with environmental factors and specific genetic predispositions. Functional impairment, disability, and handicap are expressed as paralysis, sensory and occlusive disturbances, spasticity, tremor, lack of coordination, and visual impairment. These symptoms significantly impact the quality of life of the individual.

[0308] Involvement of ionotropic glutamate receptor function including the NMDA receptor, AMPA receptor, and kainite receptor are implicated in the pathology of MS. Compounds that modulate the NMDA and AMPA/kainite family of glutamate receptors have shown neuroprotective effects in multiple sclerosis (Killestein et al., *J Neurol Sci* 2005, 233, 113-115). As a mediator of ionotropic glutamate receptors, acamprosate is potentially useful in treating MS.

[0309] Assessment of MS treatment efficacy in clinical trials can be accomplished using tools such as the Expanded Disability Status Scale and the MS Functional Composite as well as magnetic resonance imaging lesion load, biomarkers, and self-reported quality of life. Animal models of MS shown to be useful to identify and validate potential MS therapeutics include experimental autoimmune/allergic encephalomyelitis (EAE) rodent models that simulate the clinical and pathological manifestations of MS.

[0310] In certain embodiments, compounds of Formula (I) or pharmaceutical compositions thereof can be used to treat a psychotic disorder such as, for example, schizophrenia. Other psychotic disorders for which acamprosate prodrugs provided by the present disclosure may be useful include brief psychotic disorder, delusional disorder, schizoaffective disorder, and schizophreniform disorder.

[0311] Schizophrenia is a chronic, severe, and disabling brain disorder that affects about one percent of people worldwide, including 3.2 million Americans. Schizophrenia encompasses a group of psychotic disorders characterized by dysfunctions of the thinking process, such as delusions, hallucinations, and extensive withdrawal of the patient's interests from other people. Schizophrenia includes the subtypes of paranoid schizophrenia characterized by a preoccupation with delusions or auditory hallucinations, hebephrenic or disorganized schizophrenia characterized by disorganized speech, disorganized behavior, and flat or inappropriate emotions; catatonic schizophrenia dominated by physical symptoms such as immobility, excessive motor activity, or the assumption of bizarre postures; undifferentiated schizophrenia characterized by a combination of symptoms characteristic of the other subtypes; and residual schizophrenia in which a person is not currently suffering from positive symp-

toms but manifests negative and/or cognitive symptoms of schizophrenia (DSM-IV-TR classifications 295.30 (Paranoid Type), 295.10 (Disorganized Type), 295.20 (Catatonic Type), 295.90 (Undifferentiated Type), and 295.60 (Residual Type) (Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition, Text Revision, American Psychiatric Association, 2000). Schizophrenia includes these and other closely associated psychotic disorders such as schizophreniform disorder, schizoaffective disorder, delusional disorder, brief psychotic disorder, shared psychotic disorder, psychotic disorder due to a general medical condition, substance-induced psychotic disorder, and unspecified psychotic disorders (DSM-IV-TR). Schizoaffective disorder is characterized by symptoms of schizophrenia as well as mood disorders such as major depression, bipolar mania, or mixed mania, is included as a subtype of schizophrenia.

[0312] Symptoms of schizophrenia can be classified as positive, negative, or cognitive. Positive symptoms of schizophrenia include delusion and hallucination, which can be measured using, for example, using the Positive and Negative Syndrome Scale (PANSS). Negative symptoms of schizophrenia include affect blunting, anergia, alogia and social withdrawal, can be measured for example, using the Scales for the Assessment of Negative Symptoms (SANS) (Andreasen, 1983, *Scales for the Assessment of Negative Symptoms* (SANS), Iowa City, Iowa). Cognitive symptoms of schizophrenia include impairment in obtaining, organizing, and using intellectual knowledge, which can be measured using the Positive and Negative Syndrome Scale-cognitive subscale (PANSS-cognitive subscale) or by assessing the ability to perform cognitive tasks such as, for example, using the Wisconsin Card Sorting Test.

[0313] The glutamatergic system has been implicated in the etiology and pathophysiology of schizophrenia and modulators of NMDA receptor activity and mGluR⁵ receptor activity such as acamprosate have been proposed as potential therapeutic agents for schizophrenia Paz et al., *Eur Neuropsychopharmacology* 2008, prepublication no. NEUPSY-10085, 14 pages). Accordingly, acamprosate and acamprosate prodrugs provided by the present disclosure may have efficacy in treating the positive, negative, and/or cognitive symptoms of schizophrenia (Kozachuk, US 2004/0102525; and Fogel, U.S. Pat. No. 6,689,816).

[0314] The efficacy of compounds of Formula (I) and pharmaceutical compositions of any of the foregoing for treating schizophrenia may be determined by methods known to those skilled in the art. For example, negative, positive, and/or cognitive symptom(s) of schizophrenia may be measured before, during, and/or after treating the patient. Reduction in such symptom(s) indicates that a patient's condition has improved. Improvement in the symptoms of schizophrenia may be assessed using, for example, the Scale for Assessment of Negative Symptoms (SANS), Positive and Negative Symptoms Scale (PANSS) and using Cognitive Deficits tests such as the Wisconsin Card Sorting Test (WCST).

[0315] The efficacy of a compound of Formula (I) and pharmaceutical compositions of any of the foregoing may be evaluated using animal models of schizophrenic disorders. For example, conditioned avoidance response behavior (CAR) and catalepsy tests in rats are shown to be useful in predicting antipsychotic activity and EPS effect liability.

[0316] In certain embodiments, compounds of Formula (I) or pharmaceutical compositions thereof can be used to treat a mood disorder such as, for example, a bipolar disorder and a depressive disorder.

Bipolar Disorder

[0317] Bipolar disorder is a psychiatric condition characterized by periods of extreme mood. The moods can occur on a spectrum ranging from depression (e.g., persistent feelings of sadness, anxiety, guilt, anger, isolation, and/or hopelessness, disturbances in sleep and appetite, fatigue and loss of interest in usually enjoyed activities, problems concentrating, loneliness, self-loathing, apathy or indifference, depersonalization, loss of interest in sexual activity, shyness or social anxiety, irritability, chronic pain, lack of motivation, and morbid/suicidal ideation) to mania (e.g., elation, euphoria, irritation, and/or suspicious). Bipolar disorder is defined and classified in DSM-IV-TR. Bipolar disorder includes bipolar I disorder, bipolar II disorder, cyclothymia, and bipolar disorder not otherwise specified. Patients afflicted with this disorder typically alternate between episodes of depression (depressed mood, hopelessness, anhedonia, varying sleep disturbances, difficulty in concentration, psychomotor retardation and often, suicidal ideation) and episodes of mania (grandiosity, euphoria, racing thoughts, decreased need for sleep, increased energy, risk taking behavior).

[0318] Inhibitors of glutamate release such as lamotrigine and riluzole, and NMDA antagonists such as memantine and ketamine are being investigated for treating bipolar disorder (Zarate et al., *Am J Psychiatry* 2004, 161, 171-174; Zarate et al., *Biol Psychiatry* 2005, 57, 430-432; and Teng and Demetrio, *Rev Bras Psiquiatr* 2006, 28(3), 251-6).

[0319] Treatment of bipolar disorder can be assessed in clinical trials using rating scales such as the Montgomery-Asberg Depression Rating Scale, the Hamilton Depression Scale, the Raskin Depression Scale, Feiglmer criteria, and/or Clinical Global Impression Scale Score).

[0320] Depressive disorders include major depressive disorder, dysthymic disorder, premenstrual dysphoric disorder, minor depressive disorder, recurrent brief depressive disorder, and postpsychotic depressive disorder of schizophrenia (DSM IV).

[0321] Studies support the involvement of the glutamatergic system in the pathophysiology of depression. NMDA receptor antagonists have shown antidepressant effects in animal models and in clinical studies. Modulators of mGluR⁵ activity have also shown potential efficacy as antidepressants.

[0322] The efficacy of compounds provided by the present disclosure for treating depression can be evaluated in animal models of depression such as the forced swim test, the tail suspension test and others, and in clinical trials.

[0323] Anxiety is defined and classified in DSM-IV-TR. Anxiety disorders include panic attack, agoraphobia, panic disorder without agoraphobia, agoraphobia without history of panic disorder, specific phobia, social phobia, obsessive-compulsive disorder, posttraumatic stress disorder, acute stress disorder, generalized anxiety disorder, anxiety disorder due to a general medical condition, substance-induced anxiety disorder, and anxiety disorder not otherwise specified.

[0324] Neurochemical investigations have linked anxiety to dysfunction in central GABAergic, serotonergic, and noradrenergic systems. Modulators of mGluR⁵ receptors such as the selective antagonist 2-methyl-6-(phenylethynyl)-pyridine have been shown to be effective in treating anxiety

disorders (Lea and Faden, *CNS Drug Rev* 2006, 12(2), 149-66; and Molina-Hernandez et al., *Prog Neuro-Psychopharmacology Biolog Psychiatry* 2006, 30, 1129-1135). In particular, acamprosate has been proposed for the treatment of anxiety disorders (Fogel, U.S. Pat. No. 6,689,816).

[0325] Useful animal models for assessing treatment of anxiety include fear-potentiated startle, elevated plus-maze, X-maze test of anxiety, and the rat social interaction test. Genetic animal models of anxiety are also known as are other animal models sensitive to anti-anxiety agents.

[0326] In clinical trials, efficacy can be evaluated using psychological procedures for inducing experimental anxiety applied to healthy volunteers and patients with anxiety disorders or by selecting patients based on the Structured Clinical interview for DSM-IV Axis I Disorders. One or more scales can be used to evaluate anxiety and the efficacy of treatment including, for example, the Penn State Worry Questionnaire, the Hamilton Anxiety and Depression Scales, the Spielberger State-Trait Anxiety Inventory, and the Liebowitz Social Anxiety Scale.

[0327] In certain embodiments, acamprosate prodrugs provided by the present disclosure may be useful in treating somatoform disorders such as somatization disorder, conversion disorder, hypochondriasis, and body dysmorphic disorder.

[0328] In certain embodiments, movement disorders include myoclonus, tremor, tics, tardive dyskinesia, movement disorders associated with Parkinson's disease and Huntington's disease, progressive supranuclear palsy, Shy-Drager syndrome, tics, Tourette's syndrome, chorea and athetosis, spasmodic torticollis, ataxia, restless legs syndrome, and dystonias. Also included in movement disorders is spasticity.

[0329] Tardive dyskinesia is a neurological disorder caused by the long-term or high-dose use of dopamine antagonists such as antipsychotics. Tardive dyskinesia is characterized by repetitive, involuntary, purposeless movements such as grimacing, tongue protrusion, lip smacking, puckering and pursing of the lips, and rapid eye blinking, and can also involve rapid movements of the arms, legs, and trunk.

[0330] Studies suggest that NMDA receptors are involved in the dyskinesia observed in animal models of tardive dyskinesia and NMDA receptor modulators have to some extent been shown to reverse the effects of neuroleptic induced vacuous chewing movements, an animal model of tardive dyskinesia. Accordingly, acamprosate has been proposed for treating tardive dyskinesia and other movement disorders including tics, Tourette's syndrome, focal dystonias, blepharospasm, and Meige Syndrome (Fogel, U.S. Pat. No. 5,952,389, US 2002/0013366, and US 2006/1028802), and in studies on individual patients has been shown effective in treating tardive dyskinesia, dystonia, and tic at acamprosate doses from about 1,000 mg/day to about 2,000 mg/day.

[0331] Efficacy of tardive dyskinesia treatment can be assessed using animal models.

Spasticity

[0332] Spasticity is an involuntary, velocity-dependent, increased resistance to stretch. Spasticity is characterized by muscle hypertonia and displays increased resistance to externally imposed movement with increasing speed of stretch. Spasticity can be caused by lack of oxygen to the brain before, during, or after birth (cerebral palsy); physical trauma (brain or spinal cord injury); blockage of or bleeding from a blood vessel in the brain (stroke); certain metabolic diseases; adrenoleukodystrophy; phenylketonuria; neurodegenerative dis-

eases such as Parkinson's disease and amyotrophic lateral sclerosis; and neurological disorders such as multiple sclerosis. Spasticity is associated with damage to the corticospinal tract and is a common complication of neurological disease. Diseases and conditions in which spasticity may be a prominent symptom include cerebral palsy, multiple sclerosis, stroke, head and spinal cord injuries, traumatic brain injury, anoxia, and neurodegenerative diseases. Patients with spasticity complain of stiffness, involuntary spasm, and pain. These painful spasms may be spontaneous or triggered by a minor sensory stimulus, such as touching the patient.

[0333] Symptoms of spasticity can include hypertonia (increased muscle tone), clonus (a series of rapid muscle contractions), exaggerated deep tendon reflexes, muscle spasms, scissoring (involuntary crossing of the legs), deformities with fixed joints, stiffness, and/or fatigue caused by trying to force the limbs to move normally. Other complications include urinary tract infections, chronic constipation, fever or other systemic illnesses, and/or pressure sores. The degree of spasticity varies from mild muscle stiffness to severe, painful, and uncontrollable muscle spasms. Spasticity may coexist with other conditions but is distinguished from rigidity (involuntary bidirectional non-velocity-dependent resistance to movement), clonus (self-sustaining oscillating movements secondary to hypertonicity), dystonia (involuntary sustained contractions resulting in twisting abnormal postures), athetoid movement (involuntary irregular confluent writhing movements), chorea (involuntary, abrupt, rapid, irregular, and unsustained movements), ballisms (involuntary flinging movements of the limbs or body), and tremor (involuntary rhythmic repetitive oscillations, not self-sustaining). Spasticity can lead to orthopedic deformity such as hip dislocation, contractures, or scoliosis; impairment of daily living activities such as dressing, bathing, and toileting; impairment of mobility such as inability to walk, roll, or sit; skin breakdown secondary to positioning difficulties and shearing pressure; pain or abnormal sensory feedback; poor weight gain secondary to high caloric expenditure; sleep disturbance; and/or depression secondary to lack of functional independence.

[0334] Treatment of spasticity includes physical and occupational therapy such as functional based therapies, rehabilitation, facilitation such as neurodevelopmental therapy, proprioceptive neuromuscular facilitation, and sensory integration; biofeedback; electrical stimulation; and orthoses. Oral medications useful in treating spasticity include baclofen, benzodiazepines such as diazepam, dantrolene sodium; imidazolines such as clonidine and tizanidine; and gabapentin. Intrathecal medications useful in treating spasticity include baclofen. Chemodenervation with local anesthetics such as lidocaine and xylocaine; type A botulinum toxin and type B botulinum toxin; phenol and alcohol injection can also be useful in treating spasticity. Surgical treatments useful in treating spasticity include neurosurgery such as selective dorsal rhizotomy; and orthopedic operations such as contracture release, tendon or muscle lengthening, tendon transfer, osteotomy, and arthrodesis.

[0335] Studies suggest that NMDA receptor may play a role in the activity of muscle relaxants and that NMDA receptor antagonists may have therapeutic potential in spasticity (Kornhuber and Quack, *Neurosci Lett* 1995, 195, 137-139).

[0336] The efficacy of a compound of Formula (I) for the treatment of spasticity can be assessed using animal models of spasticity and in clinically relevant studies of spasticity of different etiologies. The therapeutic activity may be deter-

mined without determining a specific mechanism of action. Animal models of spasticity are known. For example, animal models of spasticity include the mutant spastic mouse; the acute/chronic spinally transected rat and the acute decerebrate rat; primary observation Irwin Test in the rat; and Rotarod Test in the rat and mouse. Other animal models include spasticity induced in rats following transient spinal cord ischemia (; spasticity in mouse models of multiple sclerosis; and spasticity in rat models of cerebral palsy.

[0337] The efficacy of compounds of Formula (I) may also be assessed in humans using double blind placebo-controlled clinical trials. Clinical trial outcome measures for spasticity include the Ashworth Scale, the modified Ashworth Scale, muscle stretch reflexes, presence of clonus and reflex response to noxious stimuli. Spasticity can be assessed using methods and procedures known in the art such as a combination of clinical examination, rating scales such as the Ashworth Scale, the modified Ashworth scale the spasm frequency scale and the reflex score, biomechanical studies such as the pendulum test, electrophysiologic studies including electromyography, and functional measurements such as the Fugl-Meyer Assessment of Sensorimotor Impairment scale. Other measures can be used to assess spasticity associated with a specific disorder such as the Multiple Sclerosis Spasticity Scale.

[0338] Cortical spreading depression (CSD) is a phenomena believed to be involved in the pathogenesis of migraine. During the early phase of CSD, a slow-propagating wave of hyper- then hypo-activity spreads through the cortex, resulting in hyper- then hypo-vascularization. This is followed by a prolonged period of neuronal depression, which is associated with disturbances in nerve cell metabolism and regional reductions in blood flow. CSD may also activate trigeminal nerve axons, which then release neuropeptides, such as substance P, neurokinin A, and CGRP from axon terminals near the meningeal and other blood vessels that produce an inflammatory response in the area around the innervated blood vessels. CSD is also implicated in progressive neuronal injury following stroke and head trauma; and cerebrovascular disease. Glutamate release and subsequent NMDA receptor activation have been implicated in the spread of CSD. NMDA antagonists such as ifenprodil have been shown effective in preventing CSD in the mouse entorhinal cortex and the NMDA receptor antagonist MK-801 was effective in blocking CSD caused by traumatic injury in rat neocortical brain slices. Accordingly, NMDA receptor antagonists that inhibit the release of glutamate in the neuron can potentially prevent CSD and its consequences. For example, (7-chloro-4-hydroxy-3-(3-phenoxy)phenyl-2-(1H)-quinolone, a high affinity antagonist at the glycine site of the NMDA receptor inhibits the initiation and propagation of spreading depression. Other selective NMDA antagonists and an uncompetitive NMDA receptor blocker have shown potential for treating cortical spreading depression migraine (Menniti et al., *Neuropharmacology* 2000, 39, 1147-1155; and Peeters et al., *J Pharmacology and Experimental Therapeutics* 2007, 321(2), 564-572). Accordingly, acamprosate prodrugs may be useful in treating cortical spreading depression related disorders such as migraine, cerebral injury, epilepsy, and cardiovascular disease.

[0339] Efficacy of acamprosate prodrugs provided by the present disclosure for treating cortical spreading depression can be assessed using animal models of cortical spreading depression.

[0340] Migraine is a neurological disorder that is characterized by recurrent attacks of headache, with pain most often occurring on one side of the head, accompanied by various combinations of symptoms such as nausea, vomiting, and sensitivity to light, sound, and odors. The exact mechanism of migraine initiation and progress is not known. Migraine can occur at any time of day or night, but occurs most frequently on arising in the morning. Migraine can be triggered by various factors, such as hormonal changes, stress, foods, lack of sleep, excessive sleep, or visual, auditory, olfactory, or somatosensory stimulation. In general, there are four phases to a migraine: the prodrome, auras, the attack phase, and postdrome. The prodrome phase is a group of vague symptoms that may precede a migraine attack by several hours, or even a few days before a migraine episode. Prodrome symptoms can include sensitivity to light and sound, changes in appetite, fatigue and yawning, malaise, mood changes, and food cravings. Auras are sensory disturbances that occur before the migraine attack in one in five patients. Positive auras include bright or shimmering light or shapes at the edge of the field of vision. Other positive aura experiences are zigzag lines or stars. Negative auras are dark holes, blind spots, or tunnel vision. Patients may have mixed positive and negative auras. Other neurologic symptoms that may occur at the same time as the aura include speech disturbances, tingling, numbness, or weakness in an arm or leg, perceptual disturbances such as space or size distortions, and confusion. A migraine attack usually lasts from 4 to 72 hours and typically produces throbbing pain on one side of the head, pain worsened by physical activity, nausea, visual symptoms, facial tingling or numbness, extreme sensitivity to light and noise, looking pale and feeling cold, and less commonly tearing and redness in one eye, swelling of the eyelid, and nasal congestion. During the attack the pain may migrate from one part of the head to another, and may radiate down the neck into the shoulder. Scalp tenderness occurs in the majority of patients during or after an attack. After a migraine attack, there is usually a postdrome phase, in which patients may feel exhausted, irritable, and/or be unable to concentrate. Other types of migraine include menstrual migraines, ophthalmologic migraine, retinal migraine, basilar migraine, familial hemiplegic migraine, and status migrainosus.

[0341] It is theorized that persons prone to migraine have a reduced threshold for neuronal excitability, possibly due to reduced activity of the inhibitory neurotransmitter γ -aminobutyric acid (GABA). GABA normally inhibits the activity of the neurotransmitters serotonin (5-HT) and glutamate, both of which appear to be involved in migraine attacks. The excitatory neurotransmitter glutamate is implicated in an electrical phenomenon called cortical spreading depression, which can initiate a migraine attack, while serotonin is implicated in vascular changes that occur as the migraine progresses.

[0342] Acamprosate prodrugs provided by the present disclosure or pharmaceutical composition thereof may be administered to a patient after initiation of the migraine. For example, a patient may be in the headache phase of the migraine or the postdrome phase before the prodrug or pharmaceutical composition is administered. Alternatively, acamprosate prodrugs provided by the present disclosure or pharmaceutical composition thereof may be administered to the patient before the migraine starts, such as once the patient senses that a migraine is developing or when the early symptoms of the migraine have begun. Acamprosate prodrugs

provided by the present disclosure may also be administered to a patient on an ongoing or chronic basis to treat recurrent or frequent occurrences of migraine episodes.

[0343] Migraine may be diagnosed by determining whether some of a person's recurrent headaches meet migraine criteria as disclosed in, for example, see The International Classification of Headache Disorders, 2nd edition, Headache Classification Committee of the International Headache Society, *Cephalalgia* 2004, 24 (suppl 1), 8-160.

[0344] The efficacy of administering at least one compound of Formula (I) for treating migraine can be assessed using animal models of migraine and clinical studies. Animal models of migraine are known. For example, to delineate and assess the effectiveness of an acamprosate prodrug provided by the present disclosure, the frequency of migraine attacks, their severity and their accompanying symptoms may be recorded and measured at baseline, and at 3 months, and 6 months, etc., following initiation of treatment. Anti-migraine and cortical-spreading depression activity of compounds provided by the present disclosure may be determined using methods known in the art.

[0345] Therapeutic efficacy of a compound of Formula (I) or pharmaceutical composition of any of the foregoing for treating migraine may also be determined in various animal models of neuropathic pain or in clinically relevant studies of different types of neuropathic pain. The therapeutic activity may be determined without determining a specific mechanism of action. Animal models for neuropathic pain are known in the art and include, but are not limited to, animal models that determine analgesic activity or compounds that act on the CNS to reduce the phenomenon of central sensitization that results in pain from non-painful or non-noxious stimuli. Other animal models are known in the art, such as hot plate tests, model acute pain and are useful for determining analgesic properties of compounds that are effective when painful or noxious stimuli are present. The progression of migraine is believed to be similar to the progress of epilepsy because an episodic phenomenon underlies the initiation of the epileptic episode and, as such, it is believed that epilepsy animal models may be useful in determining a component of the therapeutic activity of the pharmaceutical composition.

[0346] Sleeping disorders include primary sleep disorders such as dysomnias characterized by abnormalities in the amount, quality, or timing of sleep and parasomnias characterized by abnormal behavioral or physiological events occurring in association with sleep, specific sleep stages, or sleep-wake transitions; sleep disorders related to another mental disorder, sleep disorders due to a general medical condition; and substance-induced sleep disorder (DSM-IV). Dysomnias include breathing-related sleep disorders such as obstructive sleep apnea syndrome characterized by repeated episodes of upper-airway obstruction during sleep; central sleep apnea syndrome characterized by episodic cessation of ventilation during sleep without airway obstruction; and central alveolar hypoventilation syndrome characterized by impairment in ventilatory control that results in abnormally low arterial oxygen levels further worsened by sleep.

[0347] Sleep apnea is a sleep disorder characterized by pauses in breathing during sleep. Clinically significant levels of sleep apnea are defined as five or more events of any type per hour of sleep time. Sleep apnea can be characterized as central, obstructive, and mixed. In central sleep apnea, breathing is interrupted by the lack of effort. In obstructive sleep apnea, a physical block to airflow despite effort results

in interrupted breathing. In mixed sleep apnea, there is a transition from central to obstructive features during the events. Sleep apnea leads to interrupted, poor-quality sleep, nocturnal oxygen desaturation, and a reduction or absence of REM sleep. Sleep apnea may exacerbate or contribute to cardiovascular disease including coronary heart disease, hypertension, ventricular hypertrophy and dysfunction, cardiac arrhythmias, and stroke, by mechanisms such as endothelial damage and dysfunction, increases in inflammatory mediators, increases in prothrombotic factors, increased sympathetic activity, hypoxemia, impaired vagal activity and insulin resistance. Sleep apnea may also contribute to cognitive impairment.

[0348] Acamprosate has been shown to improve sleep in patients being treated for alcohol withdrawal (Staner et al., *Alcohol Clin Exp Res* 2006, 30(9), 1492-9) and preliminary studies suggest that acamprosate at doses of about 1,000 mg/day (333 mg three times per day) may be effective in treating central and obstructive sleep apnea (Hedner et al., WO 2007/032720).

[0349] Sleep apnea can be clinically evaluated using polysomnography or oximetry, and/or using tools such as the Epworth Sleepiness Scale and the Sleep Apnea Clinical Score and/or using polysomnographic recording. Animal models of sleep apnea are known and can be useful in assessing the efficacy of acamprosate prodrugs for treating sleep apnea.

[0350] Pain includes nociceptive pain caused by injury to bodily tissues and neuropathic pain caused by abnormalities in nerves, spinal cord, and/or brain. Pain includes mechanical allodynia, thermal allodynia, hyperplasia, central pain, peripheral neuropathic pain, diabetic neuropathy, breakthrough pain, cancer pain, deafferentation pain, dysesthesia, fibromyalgia syndrome, hyperpathia, incident pain, movement-related pain, myofascial pain, and paresthesia. Pain can be acute or chronic.

[0351] Studies demonstrate the involvement of mGluR⁵ receptors in nociceptive processes and that modulation of mGluR⁵ receptor activity can be useful in treating various pain states such as acute pain, persistent and chronic pain, inflammatory pain, visceral pain, neuropathic pain, non-nociceptive pain, and post-operative pain. NMDA receptor antagonists have also been shown to attenuate central sensitization and hyperplasia in animals and humans.

[0352] Neuropathic pain involves an abnormal processing of sensory input usually occurring after direct injury or damage to nerve tissue. Neuropathic pain is a collection of disorders characterized by different etiologies including infection, inflammation, disease such as diabetes and multiple sclerosis, trauma or compression to major peripheral nerves, and chemical or irradiation-induced nerve damage. Neuropathic pain typically persists long after tissue injury has resolved.

[0353] An essential part of neuropathic pain is a loss (partial or complete) of afferent sensory function and the paradoxical presence of certain hyperphenomena in the painful area. The nerve tissue lesion may be found in the brain, spinal cord, or the peripheral nervous system. Symptoms vary depending on the condition but are usually the manifestations hyperalgesia (the lowering of pain threshold and an increased response to noxious stimuli), allodynia (the evocation of pain by non-noxious stimuli such as cold, warmth, or touch), hyperpathia (an explosive pain response that is suddenly evoked from cutaneous areas with increased sensory detection threshold when the stimulus intensity exceeds sensory threshold), paroxysms (a type of evoked pain characterized

by shooting, electric, shock like or stabbing pain that occurs spontaneously, or following stimulation by an innocuous tactile stimulus or by a blunt pressure), paraesthesia (abnormal but non-painful sensations, which can be spontaneous or evoked, often described as pins and needles), dysesthesia (abnormal unpleasant but not necessarily painful sensation, which can be spontaneous or provoked by external stimuli), referred pain and abnormal pain radiation (abnormal spread of pain), and wind-up like pain and after sensations (the persistence of pain long after termination of a painful stimulus). Patients with neuropathic pain typically describe burning, lancinating, stabbing, cramping, aching and sometimes vice-like pain. The pain can be paroxysmal or constant. Pathological changes to the peripheral nerve(s), spinal cord, and brain have been implicated in the induction and maintenance of chronic pain. Patients suffering from neuropathic pain typically endure chronic, debilitating episodes that are refractory to current pharmacotherapies and profoundly affect their quality of life. Currently available treatments for neuropathic pain, including tricyclic antidepressants and gabapentin, typically show limited efficacy in the majority of patients (Sindrup and Jensen, *Pain* 1999, 83, 389-400).

[0354] There are several types of neuropathic pain. A classification that relates to the type of damage or related pathophysiology causing a painful neuropathy includes neuropathies associated with mechanical nerve injury such as carpal tunnel syndrome, vertebral disk herniation, entrapment neuropathies, ulnar neuropathy, and neurogenic thoracic outlet syndrome; metabolic disease associated neuropathies such as diabetic polyneuropathy; neuropathies associated with neurotropic viral disease such as herpes zoster and human immunodeficiency virus (HIV) disease; neuropathies associated with neurotoxicity such as chemotherapy of cancer or tuberculosis, radiation therapy, drug-induced neuropathy, and alcoholic neuropathy; neuropathies associated with inflammatory and/or immunologic mechanisms such as multiple sclerosis, anti-sulfatide antibody neuropathies, neuropathy associated with monoclonal gammopathy, Sjogren's disease, lupus, vasculitic neuropathy, polyclonal inflammatory neuropathies, Guillain-Barre syndrome, chronic inflammatory demyelinating neuropathy, multifocal motor neuropathy, paraneoplastic autonomic neuropathy, ganglionic acetylcholine receptor antibody autonomic neuropathy, Lambert-Eaton myasthenic syndrome and myasthenia gravis; neuropathies associated with nervous system focal ischemia such as thalamic syndrome (anesthesia dolorosa); neuropathies associated with multiple neurotransmitter system dysfunction such as complex regional pain syndrome (CRPS); neuropathies associated with chronic/neuropathic pain such as osteoarthritis, lower back pain, fibromyalgia, cancer bone pain, chronic stump pain, phantom limb pain, and paraneoplastic neuropathies; neuropathies associated with neuropathic pain including peripheral neuropathies such as postherpetic neuralgia, toxic neuropathies (e.g., exposure to chemicals such as exposure to acrylamide, 3-chlorophene, carbamates, carbon disulfide, ethylene oxide, n-hexane, methyl n-butylketone, methyl bromide, organophosphates, polychlorinated biphenyls, pyriminil, trichlorethylene, or dichloroacetylene), focal traumatic neuropathies, phantom and stump pain, monoradiculopathy, and trigeminal neuralgia; and central neuropathies including ischemic cerebrovascular injury (stroke), multiple sclerosis, spinal cord injury, Parkinson's disease, amyotrophic lateral sclerosis, syringomyelia, neoplasms, arachnoiditis, and post-operative pain; mixed neuropathies

such as diabetic neuropathies (including symmetric polyneuropathies such as sensory or sensorimotor polyneuropathy, selective small-fiber polyneuropathy, and autonomic neuropathy; focal and multifocal neuropathies such as cranial neuropathy, limb mononeuropathy, trunk mononeuropathy, mononeuropathy multiplex, and asymmetric lower limb motor neuropathy) and sympathetically maintained pain. Other neuropathies include focal neuropathy, glossopharyngeal neuralgia, ischemic pain, trigeminal neuralgia, atypical facial pain associated with Fabry's disease, Celiac disease, hereditary sensory neuropathy, or B₁₂-deficiency; mononeuropathies, polyneuropathies, hereditary peripheral neuropathies such as Carcot-Marie-Tooth disease, Refsum's disease, Strumpell-Lorrain disease, and retinitis pigmentosa; acute polyradiculoneuropathy; and chronic polyradiculoneuropathy. Paraneoplastic neuropathies include paraneoplastic subacute sensory neuronopathy, paraneoplastic motor neuron disease, paraneoplastic neuromyotonia, paraneoplastic demyelinating neuropathies, paraneoplastic vasculitic neuropathy, and paraneoplastic autonomic insufficiency.

[0355] The important role of N-methyl-D-aspartate (NMDA) receptors in the development and maintenance of chronic pain associated with central and peripheral nerve injury is well documented. Consequently, NMDA antagonists have been proposed as potential therapeutics for neuropathic pain. NMDA antagonists of different classes have shown efficacy in preclinical models as well as in patients with chronic pain, including neuropathic pain. Several clinical studies have observed a long-lasting relief in some neuropathic pain patients treated with NMDA antagonists (Pud et al., *Pain* 1998, 75(2-3), 349-54; Eisenberg et al., *J Pain* 2007, 8(3), 223-9; Rabben et al., *J Pharmacol Exp Ther* 1999, 289(2), 1060-1066; Correll et al., *Pain Med* 2004, 5(3), 263-75; and Harbut et al., US 2005/0148673).

[0356] Other diseases or disorders for which NMDA antagonists and mGluR⁵ antagonists such as acamprosate may be therapeutically useful include neuroprotection in epilepsy (Chapman et al., *Neuropharmacol* 2000, 39, 1567-1574), cognitive dysfunction (Riedel et al., *Neuropharmacol* 2000, 39, 1943-1951), Down's syndrome, normal cognitive senescence, meningitis, sepsis and septic encephalopathy, CNS vasculitides, leukodystrophies and X-ADL, childbirth and surgical anesthesia, spinal cord injury, hypoglycemia, encephalopathy, tumors and malignancies, cerebellar degenerations, ataxias, bowel syndromes, metabolic bone disease and osteoporosis, obesity, diabetes and pre-diabetic syndromes (Storto et al., *Molecular Pharmacology* 2006, 69(4), 1234-1241), and gastroesophageal reflux disease (Jensen et al., *Eur J Pharmacology* 2005, 519, 154-157).

Administration

[0357] Prodrugs of Formula (I), pharmaceutically acceptable salts of any of the foregoing, and/or pharmaceutical compositions thereof may be administered orally. Prodrugs of Formula (I) and/or pharmaceutical compositions thereof may also be administered by any other convenient route, for example, by infusion or bolus injection, by absorption through epithelial or mucocutaneous linings (e.g., oral mucosa, rectal, and intestinal mucosa, etc.). Administration may be systemic or local. Various delivery systems are known, (e.g., encapsulation in liposomes, microparticles, microcapsules, capsules, etc.) that may be used to administer a compound and/or pharmaceutical composition. Prodrugs of Formula (I) a pharmaceutically acceptable salt of any of the

foregoing, or a pharmaceutical composition thereof may be administered by any appropriate route. Examples of suitable routes of administration include, but are not limited to, intradermal, intramuscular, intraperitoneal, intravenous, subcutaneous, intranasal, epidural, oral, sublingual, intracerebral, intravaginal, transdermal, rectal, inhalation, or topical.

[0358] In certain embodiments, it may be desirable to introduce prodrugs of Formula (I) and/or pharmaceutical compositions thereof into the central nervous system, which may be by any suitable route, including intraventricular, intrathecal, and epidural injection. Intraventricular injection may be facilitated using an intraventricular catheter attached to a reservoir such as an Ommaya reservoir.

[0359] The amount of a prodrug of Formula (I) that will be effective in the treatment of a disease in a patient will depend, in part, on the nature of the condition and can be determined by standard clinical techniques known in the art. In addition, *in vitro* or *in vivo* assays may be employed to help identify optimal dosage ranges. A therapeutically effective amount of prodrug of Formula (I) to be administered may also depend on, among other factors, the subject being treated, the weight of the subject, the severity of the disease, the manner of administration, and the judgment of the prescribing physician.

[0360] For systemic administration, a therapeutically effective dose may be estimated initially from *in vitro* assays. For example, a dose may be formulated in animal models to achieve a beneficial circulating composition concentration range. Initial doses may also be estimated from *in vivo* data, e.g., animal models, using techniques that are known in the art. Such information may be used to more accurately determine useful doses in humans. One having ordinary skill in the art may optimize administration to humans based on animal data.

[0361] A dose may be administered in a single dosage form or in multiple dosage forms. When multiple dosage forms are used the amount of compound contained within each dosage form may be the same or different. The amount of a compound of Formula (I) contained in a dose may depend on the route of administration and whether the disease in a patient is effectively treated by acute, chronic, or a combination of acute and chronic administration.

[0362] In certain embodiments an administered dose is less than a toxic dose. Toxicity of the compositions described herein may be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g., by determining the LD₅₀ (the dose lethal to 50% of the population) or the LD₁₀₀ (the dose lethal to 100% of the population). The dose ratio between toxic and therapeutic effect is the therapeutic index. In certain embodiments, an acamprostate prodrug may exhibit a high therapeutic index. The data obtained from these cell culture assays and animal studies may be used in formulating a dosage range that is not toxic for use in humans. A dose of an acamprostate prodrug provided by the present disclosure may be within a range of circulating concentrations in for example the blood, plasma, or central nervous system, that include the effective dose and that exhibits little or no toxicity. A dose may vary within this range depending upon the dosage form employed and the route of administration utilized. In certain embodiments, an escalating dose may be administered.

Combination Therapy

[0363] In certain embodiments, prodrugs of Formula (I) or pharmaceutically acceptable salts of any of the foregoing can

be used in combination therapy with at least one other therapeutic agent. Prodrugs of Formula (I) and the at least one other therapeutic agent(s) may act additively or, in certain embodiments, synergistically. In certain embodiments, prodrugs of Formula (I) be administered concurrently with the administration of another therapeutic agent. In certain embodiments, prodrugs of Formula (I) or pharmaceutically acceptable salts of any of the foregoing may be administered prior or subsequent to administration of another therapeutic agent. The at least one other therapeutic agent may be effective for treating the same or different disease or disorder.

[0364] When used to treat a disease or disorder a therapeutically effective amount of one or more compounds of Formula (I) may be administered singly, or in combination with other agents including pharmaceutically acceptable vehicles and/or pharmaceutically active agents for treating a disease or disorder, which may be the same or different disease or disorder as the disease or disorder being treated by the one or more compounds of Formula (I). A therapeutically effective amount of one or more compounds of Formula (I) may be delivered together with a compound disclosed herein or combination with another pharmaceutically active agent.

[0365] Methods of the present disclosure include administration of one or more compounds of Formula (I), or pharmaceutical compositions thereof and another therapeutic agent provided the other therapeutic agent does not inhibit the therapeutic efficacy of the one or more compounds of Formula (I) and/or does not produce adverse combination effects.

[0366] In certain embodiments, compositions provided by the present disclosure may be administered concurrently with the administration of another therapeutic agent, which can be part of the same pharmaceutical composition as, or in a different composition than that containing the compound provided by the present disclosure. In certain embodiments, a compound of Formula (I) may be administered prior or subsequent to administration of another therapeutic agent. In certain embodiments of combination therapy, the combination therapy comprises alternating between administering a composition of Formula (I) and a composition comprising another therapeutic agent, e.g., to minimize adverse side effects associated with a particular drug. When a compound of Formula (I) is administered concurrently with another therapeutic agent that may produce adverse side effects including, but not limited to, toxicity, the other therapeutic agent may be administered at a dose that falls below the threshold at which the adverse side effect is elicited.

[0367] In certain embodiments, a pharmaceutical composition may further comprise substances to enhance, modulate and/or control release, bioavailability, therapeutic efficacy, therapeutic potency, stability, and the like. For example, to enhance therapeutic efficacy a compound of Formula (I) may be co-administered with one or more active agents to increase the absorption or diffusion of the compound from the gastrointestinal tract or to inhibit degradation of the drug in the systemic circulation. In certain embodiments, a compound of Formula (I) may be co-administered with active agents having a pharmacological effect that enhance the therapeutic efficacy of the drug.

[0368] In certain embodiments, compounds of Formula (I) or pharmaceutical compositions thereof include, or may be administered to a patient together with, another compound for treating a neurodegenerative disorder, a psychotic disorder, a mood disorder, an anxiety disorder, a somatoform disorder,

movement disorder, a substance abuse disorder, binge eating disorder, a cortical spreading depression related disorder, tinnitus, a sleeping disorder, multiple sclerosis, or pain.

[0369] In certain embodiments, acamprosate prodrugs provided by the present disclosure and pharmaceutical compositions thereof may be administered to a patient for treating a neurodegenerative disorder in combination with a therapy or another therapeutic agent known or believed to be effective in treating a neurodegenerative disorder. In certain embodiments, a neurodegenerative disorder is chosen from Alzheimer's disease, Parkinson's disease, Huntington's disease, and amyotrophic lateral sclerosis.

[0370] Therapeutic agents useful for treating Parkinson's disease include dopamine precursors such as levodopa, dopamine agonists such as bromocriptine, pergolide, pramipexole, and ropinirole, MAO-B inhibitors such as selegiline, anticholinergic drugs such as benztropine, trihexyphenidyl, tricyclic antidepressants such as amitriptyline, amoxapine, clomipramine, desipramine, doxepin, imipramine, maprotiline, nortriptyline, protriptyline, amantadine, and trimipramine, some antihistamines such as diphenhydramine; antiviral drugs such as amantadine; and β -blockers such as propranolol.

[0371] Useful drugs for treating Alzheimer's disease include rosiglitazone, roloxifene, vitamin E, donepezil, tacrine, rivastigmine, galantamine, and memantine.

[0372] Useful drugs for treating symptoms of Huntington's disease include antipsychotics such as haloperidol, chlorpromazine and olanzapine to control hallucinations, delusions and violent outbursts; antidepressants such as fluoxetine, sertraline, and nortriptyline to control depression and obsessive-compulsive behavior; tranquilizers such as benzodiazepines, paroxetine, venlafaxine and beta-blockers to control anxiety and chorea; mood stabilizers such as lithium, valproate, and carbamazepine to control mania and bipolar disorder; and botulinum toxin to control dystonia and jaw clenching. Useful drugs for treating symptoms of Huntington's disease further include selective serotonin reuptake inhibitors (SSRI) such as fluoxetine, paroxetine, sertraline, escitalopram, citalopram, fluvoxamine; norepinephrine and serotonin reuptake inhibitors (NSRI) such as venlafaxine and duloxetine, benzodiazepines such as clonazepam, alprazolam, diazepam, and lorazepam, tricyclic antidepressants such as amitriptyline, nortriptyline, and imipramine; and atypical antidepressants such as buspirone, bupropion, and mirtazapine for treating the symptoms of anxiety and depression; atomoxetine, dextroamphetamine, and modafinil for treating apathy symptoms; amantadine, memantine, and tetrabenazine for treating chorea symptoms; citalopram, atomoxetine, memantine, rivastigmine, and donepezil for treating cognitive symptoms; lorazepam and trazodone for treating insomnia; valproate, carbamazepine and lamotrigine for treating symptoms of irritability; SSRI antidepressants such as fluoxetine, paroxetine, sertraline, and fluvoxamine, NSRI antidepressants such as venlafaxine, and others such as mirtazapine, clomipramine, lomotriline, gabapentin, valproate, carbamazepine, olanzapine, risperidone, and quetiapine for treating symptoms of obsessive-compulsive disorder; haloperidol, quetiapine, clozapine, risperidone, olanzapine, ziprasidone, and aripiprazole for treating psychosis; and pramipexole, levodopa and amantadine for treating rigidity.

[0373] Useful drugs for treating ALS include riluzole. Other drugs of potential use in treating ALS include memantine, tamoxifen, thalidomide, ceftriaxone, sodium phenyl

butyrate, celecoxib, glatiramer acetate, buspirone, creatine, minocycline, coenzyme Q10, oxandrolone, IGF-1, topiramate, xaliproden, and indinavir. Drugs such as baclofen and diazepam can be useful in treating spasticity associated with ALS.

[0374] In certain embodiments, acamprosate prodrugs provided by the present disclosure and pharmaceutical compositions thereof may be administered to a patient for treating a psychotic disorder in combination with a therapy or another therapeutic agent known or believed to be effective in treating a psychotic disorder. In certain embodiments a psychotic disorder is schizophrenia.

[0375] Examples of antipsychotic agents useful in treating positive symptoms of schizophrenia include, but are not limited to, acetophenazine, alseroxylon, amitriptyline, aripiprazole, astemizole, benzquinamide, carphenazine, chlormezanone, chlorpromazine, chlorprothixene, clozapine, desipramine, droperidol, aloperidol, fluphenazine, flupenthixol, glycine, oxapine, mesoridazine, molindone, olanzapine, ondansetron, perphenazine, pimozide, prochlorperazine, procyclidine, promazine, propiomazine, quetiapine, remoxipride, reserpine, risperidone, sertindole, sulpiride, terfenadine, thiethylperazine, thioridazine, thiothixene, trifluoperazine, triflupromazine, trimeprazine, and ziprasidone. Examples of typical antipsychotic agents useful for treating positive symptoms of schizophrenia include acetophenazine, chlorpromazine, chlorprothixene, droperidol, fluphenazine, haloperidol, loxapine, mesoridazine, methotrimeprazine, molindone, perphenazine, pimozide, raclopride, remoxipride, thioridazine, thiothixene, and trifluoperazine. Examples of atypical antipsychotic agents useful for treating positive symptoms of schizophrenia include aripiprazole, clozapine, olanzapine, quetiapine, risperidone, sertindole, and ziprasidone.

[0376] Other antipsychotic agents useful for treating positive symptoms of schizophrenia include amisulpride, balaperidone, blonanserin, butaperazine, carphenazine, eplavanserin, iloperidone, lamictal, osanetant, paliperidone, perospirone, piperacetazine, raclopride, remoxipride, sarizotan, sonepiprazole, sulpiride, ziprasidone, and zotepine; serotonin and dopamine (5HT/D2) agonists such as asenapine and bifeprunox; neurokinin 3 antagonists such as talnetant and osanetant; AMPA/kines such as CX-516, galantamine, memantine, modafinil, ocaperidone, and tolcapone; and α -amino acids such as D-serine, D-alanine, D-cycloserine, and N-methylglycine. Thus, antipsychotic agents include typical antipsychotic agents, atypical antipsychotic agents, and other compounds useful for treating schizophrenia in a patient, and particularly useful for treating the positive symptoms of schizophrenia.

[0377] Examples of agents useful for treating cognitive and/or negative symptoms of schizophrenia include aripiprazole, clozapine, olanzapine, quetiapine, risperidone, sertindole, ziprasidone, asenapine, bifeprunox, iloperidone, lamictal, galantamine, memantine, modafinil, acaperidone, NK3 antagonists such as talnetant and osanetant, AMPA/kines, tolcapone, amisulpride, mirtazapine, lamotrigine, idazoxan, nebogamine, sabcomeline, ispronicline, sarcosine, preclamol, L-camosine, nicotine, raloxifene, pramipexol, escitalopram, estradiol, riluzole, creatine, entacapone, L-threonine, atomoxetine, divalproex sodium, pimozide, provastatin, duloxetine; and NMDA receptor modulators such as glycine, D-serine, and D-cycloserine.

[0378] In certain embodiments, pharmaceutical compositions provided by the present disclosure may be co-administered with another drug useful for treating a symptom of schizophrenia or a disease, disorder, or condition associated with schizophrenia and that is not an antipsychotic agent. For example, acamprosate prodrugs may be co-administered with an antidepressant, such as, but not limited to alprazolam, amitriptyline, amoxapine, bupropion, citalopram, clomipramine, desipramine, doxepin, escitalopram, fluoxetine, fluvoxamine, imipramine, maprotiline, methylphenidate, mirtazapine, nefazodone, nortriptyline, paroxetine, phenelzine, protriptyline, sertraline, tranylcypromine, trazodone, trimipramine, venlafaxine, and combinations of any of the foregoing.

[0379] For example, in certain embodiments, an acamprosate prodrug provided by the present disclosure, or pharmaceutical compositions thereof may be administered to a patient for the treatment of schizophrenia in conjunction with a social therapy for treating schizophrenia such as rehabilitation, community support activities, cognitive behavioral therapy, training in illness management skills, participation in self-help groups, and/or psychotherapy. Examples of psychotherapies useful for treating schizophrenia include Alderian therapy, behavior therapy, existential therapy, Gestalt therapy, person-centered therapy, psychoanalytic therapy, rational-emotive and cognitive-behavioral therapy, reality therapy, and transactional analysis.

[0380] Other examples of drugs useful for treating psychotic disorders include aripiprazole, loxapine, mesoridazine, quetiapine, reserpine, thioridazine, trifluoperazine, and ziprasidone, chlorpromazine, clozapine, fluphenazine, haloperidol, olanzapine, perphenazine, prochlorperazine, risperidone, and thiothixene.

[0381] In certain embodiments, acamprosate prodrugs provided by the present disclosure and pharmaceutical compositions thereof may be administered to a patient for treating a mood disorder in combination with a therapy or another therapeutic agent known or believed to be effective in treating a mood disorder. In certain embodiments, a mood disorder is chosen from a bipolar disorder and a depressive disorder.

[0382] Examples of drugs useful for treating bipolar disorder include aripiprazole, verapamil, carbamazepine, clonidine, clonazepam, lamotrigine, olanzapine, quetiapine, fluoxetine, and ziprasidone.

[0383] In certain embodiments, acamprosate prodrugs provided by the present disclosure and pharmaceutical compositions thereof may be administered to a patient for treating depression in combination with a therapy or another therapeutic agent known or believed to be effective in treating depression.

[0384] Examples of drugs useful for treating depression include tricyclics such as amitriptyline, amoxapine, clomipramine, desipramine, doxepin, imipramine, maprotiline, nortriptyline, protriptyline, and trimipramine; tetracyclics such as maprotiline and mirtazapine; selective serotonin reuptake inhibitors (SSRI) such as citalopram, escitalopram, fluoxetine, fluvoxamine, paroxetine, and sertraline; serotonin and norepinephrine reuptake inhibitors (SNRI) such as venlafaxine and duloxetine; monoamine oxidase inhibitors such as isocarboxazid, phenelzine, selegiline, and tranylcypromine; psychostimulants such as dextroamphetamine and methylphenidate; and other drugs such as bupropion, mirtazapine, nefazodone, trazodone, lithium, and venlafaxine.

[0385] In certain embodiments, acamprosate prodrugs provided by the present disclosure and pharmaceutical compositions thereof may be administered to a patient for treating an anxiety disorder in combination with a therapy or another therapeutic agent known or believed to be effective in treating an anxiety disorder.

[0386] Examples of drugs for useful treating anxiety disorders include alprazolam, atenolol, busiprone, chlordiazepoxide, clonidine, clorazepate, diazepam, doxepin, escitalopram, halazepam, hydroxyzine, lorazepam, nadolol, oxazepam, paroxetine, prochlorperazine, trifluoperazine, venlafaxine, amitriptyline, sertraline, citalopram, clomipramine, fluoxetine, fluvoxamine, and paroxetine.

[0387] In certain embodiments, acamprosate prodrugs provided by the present disclosure and pharmaceutical compositions thereof may be administered to a patient for treating a somatoform disorder in combination with a therapy or another therapeutic agent known or believed to be effective in treating a somatoform disorder.

[0388] Examples of drugs useful for treating somatoform disorders include tricyclic antidepressants such as amitriptyline, and serotonin reuptake inhibitors.

[0389] In certain embodiments, acamprosate prodrugs provided by the present disclosure and pharmaceutical compositions thereof may be administered to a patient for treating a movement disorder in combination with a therapy or another therapeutic agent known or believed to be effective in treating a movement disorder. In certain embodiments, a movement disorder is selected from tardive dyskinesia and spasticity.

[0390] Examples of drugs useful for treating movement disorders include levodopa, mild sedatives such as benzodiazepines including alprazolam, chlordiazepoxide, clonazepam, clorazepate, diazepam, lorazepam, and oxazepam; muscle relaxants such as baclofen, anticholinergic drugs such as trihexyphenidyl and diphenhydramine; antipsychotics such as chlorpromazine, fluphenazine, haloperidol, loxapine, mesoridazine, molindone, perphenazine, pimozide, thioridazine, thiothixene, trifluoperazine, aripiprazole, clozapine, olanzapine, quetiapine, risperidone, and ziprasidone; and antidepressants such as amitriptyline.

[0391] Examples of drugs useful for treating tardive dyskinesia include vitamin E, dizocilpine, memantine, clzapine, lorazepam, diazepam, clonazepam, glycine, D-cycloserine valproic acid, amantadine, ifenprodil, and tetrabenazine.

[0392] Examples of drugs useful for treating spasticity include baclofen, R-baclofen, diazepam, tizanidine, clonidine, dantrolene, 4-aminopyridine, cyclobenzaprine, ketazolam, tiagabine, and botulinum A toxin. Compounds having activity as $\alpha_2\delta$ subunit calcium channel modulators such as gabapentin and pregabalin are believed to be useful as antispasticity agents.

[0393] In certain embodiments, acamprosate prodrugs provided by the present disclosure and pharmaceutical compositions thereof may be administered to a patient for treating a substance abuse disorder in combination with a therapy or another therapeutic agent known or believed to be effective in treating a substance abuse disorder. In certain embodiments, a substance abuse disorder is chosen from an alcohol abuse disorder, a narcotic abuse disorder, and a nicotine abuse disorder.

[0394] Examples of drugs useful for treating alcohol dependency or alcohol abuse disorders include disulfiram, naltrexone, acamprosate, ondansetron, atenolol, chlordiazep-

oxide, clonidine, clorazepate, diazepam, oxazepam, methadone, topiramate, 1-alpha-acetylmethadol, buprenorphine, bupropion, and baclofen.

[0395] Examples of drugs useful for treating opioid abuse disorders include buprenorphine, naloxone, tramadol, methadone, and naltrexone.

[0396] Examples of drugs useful for treating cocaine abuse disorders include disulfiram, modafinil, propranolol, baclofen, vigabatrin, and topiramate.

[0397] Examples of drugs useful for treating nicotine abuse disorders include bupropion, clonidine, rimonabant, varenicline, and nicotine.

[0398] In certain embodiments, acamprosate prodrugs provided by the present disclosure and pharmaceutical compositions thereof may be administered to a patient for treating a cortical spreading depression related disorder in combination with a therapy or another therapeutic agent known or believed to be effective in treating a cortical spreading depression related disorder. In certain embodiments, a cortical spreading depression related disorder is selected from migraine, cerebral injury, epilepsy, and cardiovascular disease.

[0399] Drugs useful for treating migraine can prevent a migraine from occurring, abort a migraine that is beginning, or relieve pain during the migraine episode.

[0400] Prophylactic migraine treatments reduce the frequency of migraines and include non-steroidal anti-inflammatory agents (NSAIDs), adrenergic beta-blockers, calcium channel blockers, tricyclic antidepressants, selective serotonin reuptake inhibitors, anticonvulsants, NMDA receptor antagonists, angiotensin converting enzyme (ACE) inhibitors, angiotensin-receptor blockers (ARBs), leukotriene-antagonists, dopamine agonists, selective 5HT-1D agonists, selective 5HT-1F agonists, AMPA/KA antagonists, CGRP (calcitonin gene related peptide) antagonists, NOS (nitric oxide synthase) inhibitors, blockers of spreading cortical depression, and other therapy. Examples of NSAIDs useful for preventing migraine include aspirin, ibuprofen, fenoprofen, flurbiprofen, ketoprofen, mefenamic acid, and naproxen. Examples of adrenergic beta-blockers useful for preventing migraine include acebutolol, atenolol, imilol, metoprolol, nadolol, pindolol, propranolol, and timolol. Examples of calcium channel blockers useful for preventing migraine include amlodipine, diltiazem, dotarizine, felodipine, flunarizine, nicardipine, nifedipine, nimodipine, nisoldipine, and verapamil. Examples of tricyclic antidepressants useful for preventing migraine include amitriptyline, desipramine, doxepin, imipramine, nortriptyline, and protriptyline. Examples of selective serotonin reuptake inhibitors (SSRIs) useful for preventing migraine include fluoxetine, methysergide, nefazodone, paroxetine, sertraline, and venlafaxine. Examples of other antidepressants useful for preventing migraine include bupropion, nefazodone, norepinephrine, and trazodone.

[0401] Examples of anticonvulsants (antiepileptics) useful for preventing migraine include divalproex sodium, felbamate, gabapentin, lamotrigine, levetiracetam, oxcarbazepine, tiagabine, topiramate, valproate, and zonisamide. Examples of NMDA receptor antagonists useful for preventing migraine include dextromethorphan, magnesium, and ketamine. Examples of angiotensin converting enzyme (ACE) inhibitors useful for preventing migraine include lisinopril. Examples of angiotensin-receptor blockers (ARBs) useful for preventing migraine include candesartan. Examples of leukotriene-antagonists useful for preventing migraine include zileuton, zafirlukast, montelukast, and pranlukast. Examples

of dopamine agonists useful for preventing migraine include α -dihydroergocryptine. Examples of other therapy useful for preventing migraine include botulinum toxin, magnesium, hormone therapies, riboflavin, methylergonovine, cyproheptadine, and phenelzine, and complementary therapies such as counseling/psychotherapy, relaxation training, progressive muscle relaxation, guided imagery, diaphragmatic breathing, biofeedback, acupuncture, and physical and massage therapy.

[0402] Acute migraine treatments intended to eliminate or reduce the severity of the headache and any associated symptoms after a migraine has begun include serotonin receptor agonists, such as triptans (5-hydroxytryptophan (5-HT) agonists) including almotriptan, eletriptan, frovatriptan, naratriptan, rizatriptan, sumatriptan, imotriptan, and zolmitriptan; ergotamine-based compounds such as dihydroergotamine and ergotamine; antiemetics such as metoclopramide and prochlorperazine; and compounds that provide analgesic effects.

[0403] Other examples of drugs used to treat migraine once started include, acetaminophen-aspirin, caffeine, cyproheptadine, methysergide, valproic acid, NSAIDs such as diclofenac, flurbiprofen, ketaprofen, ketorolac, ibuprofen, indomethacin, meclofenamate, and naproxen sodium, opioids such as codeine, meperidine, and oxycodone, and glucocorticoids including dexamethasone, prednisone and methylprednisolone.

[0404] GABA analog prodrugs provided by the present disclosure may also be administered in conjunction with drugs that are useful for treating symptoms associated with migraine such as nausea and vomiting, and depression. Examples of useful therapeutic agents for treating or preventing vomiting include, but are not limited to, 5-HT₃ receptor antagonists such as ondansetron, dolasetron, granisetron, and tropisetron; dopamine receptor antagonists such as prochlorperazine, thiethylperazine, chlorpromazine, metoclopramide, and domperidone; glucocorticoids such as dexamethasone; and benzodiazepines such as lorazepam and alprazolam. Examples of useful therapeutic agents for treating or preventing depression include, but are not limited to, tricyclic antidepressants such as amitriptyline, amoxapine, bupropion, clomipramine, desipramine, doxepin, imipramine, maprotiline, nefazodone, nortriptyline, protriptyline, trazodone, trimipramine, and venlafaxine; selective serotonin reuptake inhibitors such as fluoxetine, fluvoxamine, paroxetine, and sertraline; monoamine oxidase inhibitors such as isocarboxazid, pargyline, phenazine, and tranylcypromine; and psychostimulants such as dextroamphetamine and methylphenidate.

[0405] Useful drugs for treating cerebral trauma include corticosteroids such as betamethasone, budesonide, cortisone, dexamethasone, hydrocortisone, methylprednisolone, prednisolone, prednisone, and triamcinolone, and antithrombotics such as ticlopidine.

[0406] Useful drugs for treating epilepsy include acetazolamide, carbamazepine, gabapentin, mephobarbital, felbamate, fosphenytoin, phenytoin, pregabalin, and valproic acid.

[0407] In certain embodiments, acamprosate prodrugs provided by the present disclosure and pharmaceutical compositions thereof may be administered to a patient for treating tinnitus in combination with a therapy or another therapeutic agent known or believed to be effective in treating tinnitus.

[0408] A second therapeutic agent for treating or preventing tinnitus can have one or more of analgesic, anesthetic, sodium channel blocker, antiemetic, analgesic, and anti-

pyretic properties. Analgesics include, for example, steroidal anti-inflammatory agents, non-steroidal anti-inflammatory agents, selective COX-2 inhibitors, and narcotics. Examples of analgesics include, for example, acetaminophen, amitriptyline, aspirin, buprenorphine, celecoxib, clonidine, codeine, diclofenac, diflunisal, etodolac, fenoprofen, fentanyl, flurbiprofen, hydromorphone, hydroxyzine, ibuprofen, imipramine, indomethacin, ketoprofen, ketorolac, levorphanol, meperidine, methadone, morphine, naproxen, oxycodone, piroxicam, propoxyphene, rofecoxib, sulindac, tolmetin, tramadol, valdecoxib, and combinations of any of the foregoing.

[0409] In certain embodiments, a compound of the present disclosure or pharmaceutical composition thereof can be administered with a N-methyl-D-aspartate (NMDA) receptor antagonist that binds to the NMDA receptor at the competitive NMDA antagonist binding site, the non-competitive NMDA antagonist binding site within the ion channel, or to the glycine site. Examples of NMDA receptor antagonists include amantadine, D-2-amino-5-phosphonopentanoic acid (D-AP5), 3-((±)2-carboxypiperazin-4-yl)-propyl-1-phosphonic acid (CCP), conantokins, 7-chlorokynurenate (7-CK), dextromethorphan, ifenprodil, ketamine, memantine, dizocilpine, gacyclidine, licostinel, phencyclidine, riluzole, traxoprodil, and combinations of any of the foregoing (Sands, U.S. Pat. No. 5,716,961 and Guitton et al., US 2006/0063802). Other drugs that may be useful in treating tinnitus include baclofen, caroverine, piribedil, nimodipine, clonazepam, and trimetazidine.

[0410] An acamprosate prodrug of Formula (I) or pharmaceutical composition thereof can also be used in conjunction with non-pharmacological tinnitus therapies such as, for example, avoidance of ototoxic medications, reduced consumption of alcohol, caffeine and nicotine, reduced stress, the use of background noises or maskers, behavioral therapies such as hypnosis, cognitive therapy, biofeedback, tinnitus retraining therapy.

[0411] In certain embodiments, acamprosate prodrugs provided by the present disclosure and pharmaceutical compositions thereof may be administered to a patient for treating a sleeping disorder in combination with a therapy or another therapeutic agent known or believed to be effective in treating a sleeping disorder.

[0412] Examples of drugs useful for treating sleep apnea include mirtazapine and modafinil.

[0413] In certain embodiments, acamprosate prodrugs provided by the present disclosure and pharmaceutical compositions thereof may be administered to a patient for treating multiple sclerosis in combination with a therapy or another therapeutic agent known or believed to be effective in treating multiple sclerosis.

[0414] Examples of drugs useful for treating MS include corticosteroids such as methylprednisolone; IFN- β such as IFN- β 1a and IFN- β 1b; glatiramer acetate; monoclonal antibodies that bind to the very late antigen-4 (VLA-4) integrin such as natalizumab; immunomodulatory agents such as FTY 720 sphingosine-1 phosphate modulator and COX-2 inhibitors such as BW755c, piroxicam, and phenidone; and neuroprotective treatments including inhibitors of glutamate excitotoxicity and iNOS, free-radical scavengers, and cationic channel blockers; memantine; AMPA antagonists such as topiramate; and glycine-site NMDA antagonists.

[0415] In certain embodiments, acamprosate prodrugs provided by the present disclosure and pharmaceutical compositions thereof may be administered to a patient for treating

pain in combination with a therapy or another therapeutic agent known or believed to be effective in treating pain. In certain embodiments, the pain is neuropathic pain.

[0416] Examples of drugs useful for treating pain include opioid analgesics such as morphine, codeine, fentanyl, meperidine, methadone, propoxyphene, levorphanol, hydromorphone, oxycodone, oxymorphone, and pentazocine; non-opioid analgesics such as aspirin, ibuprofen, ketoprofen, naproxen, and acetaminophen; nonsteroidal anti-inflammatory drugs such as aspirin, choline magnesium trisalicylate, diflunisal, salsalate, celecoxib, rofecoxib, valdecoxib, diclofenac, etodolac, fenoprofen, flubiprofen, ibuprofen, indomethacin, ketoprofen, ketorolac, meclofanamate, mefenamic acid, meloxicam, nabumetone, naproxen, oxaprozin, piroxicam, sulindac, and tometin; and other drugs such as amitriptyline, desipramine, gabapentin, carbamazepine, phenytoin, clonazepam, divalproex, lamotrigine, topiramate, oxcarbazepine, divalproex, butorphanol, tramadol, valdecoxib, vicoprofen, pentazocine, propoxyphene, fenoprofen, piroxicam, indomethacin, hydroxyzine, buprenorphine, benzocaine, clonidine, flurbiprofen, and meperidine.

[0417] The weight ratio of compounds of Formula (I) to a second therapeutic agent may be varied and may depend upon the effective dose of each agent. A therapeutically effective dose of each compound will be used. Thus, for example, when a compound of Formula (I) is combined with another therapeutic agent, the weight ratio of the compound provided by the present disclosure to the second therapeutic agent can be from about 1000:1 to about 1:1000, and in certain embodiments, from about 200:1 to about 1:200.

[0418] Combinations of compounds of Formula (I) and a second therapeutic agent may also be within the aforementioned range, but in each case, an effective dose of each active compound can be used. In such combinations a compound of Formula (I) and second therapeutic agent may be administered separately or in conjunction. In addition, administration of one agent may be prior to, concurrent with, or subsequent to the administration of another therapeutic agent(s). Accordingly, compounds of Formula (I) may be used alone or in combination with other therapeutic agents that are known to be beneficial in treating the same disease being treated with the compound of Formula (I) or other therapeutic agents that affect receptors or enzymes that either increase the efficacy, safety, convenience, or reduce unwanted side effects or toxicity of the compound of Formula (I). Compounds of Formula (I) and the other therapeutic agent may be co-administered, either in concomitant therapy or in a fixed combination. The additional therapeutic agent may be administered by the same or different route than the route used to administer a compound of Formula (I) or pharmaceutical composition of any of the foregoing.

EXAMPLES

[0419] The following examples describe in detail synthesis of compounds of Formula (I)-(II), properties of compounds of Formula (I)-(II), and uses of compounds of Formula (I)-(II). It will be apparent to those skilled in the art that many modifications, both to materials and methods, may be practiced without departing from the scope of the disclosure.

Description 1

General Experimental Protocols

[0420] All reagents and solvents were purchased from commercial suppliers and used without further purification or manipulation.

[0421] Proton NMR spectra (400 MHz) were recorded on a Varian AS 400 NMR spectrometer equipped with an autosampler and data processing computation. CDCl_3 (99.8% D), DMSO-d_6 (99.9% D), or MeOH-d_4 (99.8+% D) were used as solvents unless otherwise noted. The CHCl_3 , DMSO-d_6 , or MeOH-d_3 solvent signals were used for calibration of the individual spectra. Determination of enantiomeric excess (e.e.) of intermediates was accomplished by ^1H NMR-spectroscopy in the presence of the diamagnetic enantiomerically pure chiral co-solvent (R)-(-)-2,2,2-trifluoro-1-(9-anthryl) ethanol (Pirkle-alcohol) and in comparison NMR-spectra of the corresponding racemic samples under similar conditions (Parker, *Chem. Rev.*, 1991, 91, 1441-1457). Analytical thin layer chromatography (TLC) was performed using Whatman, Schleicher & Schuell TLC MK6F silica gel plates (2.5×7.5 cm, 250 μm layer thickness). Dyeing or staining reagents for TLC detection and visualization were prepared according methods known in the art. Ozonolysis reactions were performed using a Welsbach Standard T-series ozone generator. Analytical LC/MS was performed on a Waters 2790 separation module equipped with a Waters Micromass QZ mass spectrometer, a Waters 996 photodiode detector, and a Merck Chromolith UM2072-027 or Phenomenex Luna C-18 analytical column. Mass-guided preparative HPLC purification of final compounds was performed on an instrument equipped with a Waters 600 controller, ZMD Micromass spectrometer, a Waters 2996 photodiode array detector, and a Waters 2700 Sample Manager. Acetonitrile/water gradients containing 0.05% formic acid were used as eluent in both analytical and preparative HPLC experiments. Compound isolation from aqueous solvent mixtures, e.g., acetonitrile/water/0.05% formic acid, was accomplished by primary lyophilization (freeze drying) of the frozen solutions under reduced pressure at room temperature using manifold freeze dryers such as Heto Drywinner DW 6-85-1, Heto FD4, or VIRTIS Freezemobile 25 ES equipped with a high vacuum pump. Optionally, and if the isolated compound had ionizable functional groups such as an amino group or a carboxylic acid, the lyophilization process was conducted in the presence of a slight excess of one molar (1.0 M) hydrochloric acid to yield the purified compounds as the corresponding hydrochloride salt (HCl-salt) or the corresponding protonated free carboxylic acid.

[0422] Chemical names were generated with Chemistry 4-D Draw Pro Version 7.01c (Draw Chemical Structures Intelligently© 1993-2002) from ChemInnovation Software, Inc., San Diego, USA).

Example 1

Synthesis of N-[3-(Chlorosulfonyl)propyl]acetamide (1)

Step A: Tetramethylammonium N-acetylhomotaurate (1a)

[0423] Tetramethylammonium N-acetylhomotaurate (1a) was synthesized adapting procedures disclosed in Durlach, U.S. Pat. No. 4,355,043, DE 3019350, and U.S. Pat. No. 4,199,601. A 250 mL round bottomed flask equipped with a magnetic stir bar was charged with 3-amino-1-propanesulfonic acid (5.0 g, 36.0 mmol) and 20 mL of water. To the stirred solution, 13.0 g (36.0 mmol) of tetramethylammonium hydroxide (25 w-% in water) was added. The solution was stirred at room temperature for 1 hour and acetic anhydride 4.1 mL (4.39 g, 43.0 mmol) was added. The mixture was

stirred overnight at ca. 40° C. (oil bath) to ensure complete conversion. The resulting solution was extracted twice with 30 mL of diethyl ether or tert-butyl methyl ether (MTBE), and residual methanol in the aqueous phase was removed under reduced pressure using a rotary evaporator. The extract was isolated from the residual water in the solution by lyophilization to yield 9.1 g (quant.) of the title compound (1a) as a colorless powder that was used without further purification after additional thorough drying under high vacuum. ^1H NMR (400 MHz, D_2O): δ =1.88-1.95 (m, 2H), 1.97 (s, 3H), 2.88-2.92 (m, 2H), 3.16 (s, 12H), 3.27 (in, 2H) ppm. MS (ESI) m/z 180.04 (M-H^-).

Step B: N-[3-(Chlorosulfonyl)propyl]acetamide (1)

[0424] Adapting a procedure or a variation thereof according to Shue et al., *Bioorg. Med. Chem.* 1996, 6, 1709-1714, and Korolev et al., *Synthesis* 2003, 383-388, a 500 mL round bottomed flask equipped with a magnetic stir bar was charged with tetramethylammonium N-acetylhomotaurate (1a) (9.1 g, 36 mmol), phosphorus pentachloride (PCl_5) (7.9 g, 37 mmol), and 200 mL of anhydrous dichloromethane (200 mL). The solution was heated to reflux overnight. The resulting mixture was washed twice with water (100 mL) and brine (100 mL). The organic layer was dried over magnesium sulfate (MgSO_4), filtered, and the solvents removed by evaporation under reduced pressure using a rotary evaporator to provide 4.6 g (65% yield) of the title compound (1) as a light yellow, viscous liquid. The crude material was of sufficient purity to be used in the next steps. ^1H NMR (400 MHz, CDCl_3): δ =2.07 (s, 3H), 2.23-2.32 (m, 2H), 3.46-3.51 (m, 2H), 3.76-3.80 (m, 2H) ppm. MS (ESI) m/z 200.01 (M+H^+). The material contained various amounts of the cyclization product 2-acetyl-1,1-dioxo-1,2-thiazolidine as determined by ^1H NMR spectroscopy and mass spectroscopy. ^1H NMR (400 MHz, CDCl_3): δ =2.37-2.44 (m, 5H), 3.41 (t, J =7.2 Hz, 2H), 3.84 (t, J =7.2 Hz, 2H) ppm. MS (ESI) m/z 164.01 (M+H^+), 186.00 (M+Na^+).

Description 2

General Procedure for the Preparation of Hydroxypivalic Acid Esters (HPA-Esters) through Reaction of HPA with Alkylating Agents

Method A

[0425] A dry 500 mL round-bottomed flask equipped with a magnetic stirring bar, a reflux condenser, and a rubber septum was charged under an atmosphere of nitrogen with 7.09 g (60.0 mmol) of 3-hydroxy-2,2-dimethylpropanoic acid (hydroxypivalic acid, HPA) or 3-hydroxy-2-(hydroxymethyl)-2-methylpropanoic acid (bis-hydroxypivalic acid, BHPA) and 19.55 g (60.0 mmol) of dried and freshly powdered cesium carbonate (Cs_2CO_3). 150 mL of anhydrous dimethylformamide (DMF) was added followed by 0.7-1.0 equivalents (40.0 to 60.0 mmol) of an appropriately substituted alkyl halide. The reaction mixture was heated under an atmosphere of nitrogen for overnight to ca. 50-70° C. (oil bath) (depending on the boiling point of the halide). The reaction mixture was diluted with ethyl acetate, precipitates were filtered off, and the filtrate was carefully diluted with a one molar (1.0 M) of hydrochloric acid (HCl). The aqueous phase was extracted several times with ethyl acetate and the combined organic extracts were washed with a saturated aqueous solution of sodium hydrogen carbonate (NaHCO_3).

The organic solvents were removed under reduced pressure using a rotary evaporator and the residue was diluted in methyl tert-butyl ether (MTBE). The organic solution was washed five times with water (to remove residual DMF), then brine, dried over anhydrous magnesium sulfate (MgSO_4), filtered and the residual solvents removed under reduced pressure using a rotary evaporator. Generally, the products were of sufficient purity to be used directly in the next step without further purification or, optionally, were purified by fractional distillation under reduced pressure or silica gel column chromatography using ethyl acetate/hexane mixtures as eluent.

Method B

[0426] rac-1-Acyloxyalkyl halides, i.e., chlorides, were either commercially available or were prepared from the corresponding aldehydes and carboxylic acid chlorides in the presence of catalytic amounts of anhydrous zinc chloride (ZnCl_2) adapting literature known methods or variations thereof. As a typical example, a 250 mL round bottomed flask equipped with a magnetic stirring bar, an addition funnel, and a rubber septum was charged under a nitrogen atmosphere with 100.0 mmol of the corresponding aldehyde, a catalytic amount of anhydrous zinc chloride (ZnCl_2) (ca. 5-10 mol-%), and 100-300 mL of an inert solvent such as dichloromethane (DCM). The solution was cooled to ca. 0° C. (ice bath) and to 50.0 mmol of the corresponding carboxylic acid chloride as a solution in DCM was added while stirring. The reaction mixture was stirred at 0° C. for approximately three hours with gradual darkening of the mixture. (Note: If the reaction is run for longer periods and with gradual warming to room temperature, product yields and integrity will become significantly lower after workup.). The solvents and excess of volatile reagents were carefully removed under reduced pressure using a rotary evaporator. The resulting crude reaction mixture was directly pre-purified through a short plug filtration column using n-pentane or hexanes as eluent followed by careful removal of the solvents under slightly reduced pressure using a rotary evaporator. Optionally, the crude reaction mixture was diluted with diethyl ether (Et_2O) and the solution washed with a saturated aqueous solution of sodium hydrogen carbonate (NaHCO_3), water, and brine, dried over anhydrous magnesium sulfate (MgSO_4), filtered and carefully evaporated under reduced pressure using a rotary evaporator to provide the desired product. Occasionally, the products obtained by either pre-purification method were of sufficient purity to be used directly in the next step without further purification or, optionally, were purified further by fractional distillation under reduced pressure or silica gel column chromatography using diethyl ether (Et_2O) or methyl tert-butyl ether (MTBE) mixtures with pentane, hexanes, or heptane as eluent (depending on the volatility of the title compound).

[0427] rac-1-Alkoxy- and aryloxycarbonyloxyalkyl halides, i.e., chlorides, were either commercially available or were prepared from commercially available 1-chloroalkyl chloroformates and the corresponding alcohols or phenols adapting methods known in the literature or variations thereof. As a typical example, a 250 mL round bottomed flask equipped with a magnetic stirring bar and a rubber septum was charged under a nitrogen atmosphere with 10.0 mmol of the corresponding alcohol of phenol, 890 μL (870 mg, 11.0 mmol) of anhydrous pyridine (Pyr) in 10-30 mL of an inert solvent such as dichloromethane (DCM). The solution was cooled to ca. 0° C. (ice bath) and 12.0 mmol of the corre-

sponding rac-1-chloroalkyl chloroformate either in neat form or as a concentrated solution in DCM was added while stirring. The reaction mixture was stirred overnight with gradual warming to room temperature. The solvents and volatile reagents were carefully removed under reduced pressure using a rotary evaporator. The residue was diluted with diethyl ether (Et_2O) or methyl tert-butyl ether (MTBE), washed with 0.1 molar hydrochloric acid (HCl), water, and brine, and then dried over anhydrous magnesium sulfate (MgSO_4). After filtration and evaporation of the solvents under reduced pressure using a rotary evaporator, the desired compound was obtained. Generally, the products were of sufficient purity to be used directly in the next step without further purification or, optionally, were purified by fractional distillation under reduced pressure or silica gel column chromatography using diethyl ether (Et_2O) or methyl tert-butyl ether (MTBE) mixtures with pentane, hexanes, or heptane as eluent (depending on the volatility of the title compound).

[0428] For the synthesis of acyloxyalkyl ester derivatives or alkoxy- and aryloxycarbonyloxyalkyl ester derivatives, a 100 mL round bottomed flask equipped with a magnetic stirring bar and a rubber septum was charged under a nitrogen atmosphere with 709 mg (6.0 mmol) of 3-hydroxy-2,2-dimethylpropanoic acid (hydroxypivalic acid, HPA) or 3-hydroxy-2-(hydroxymethyl)-2-methylpropanoic acid (bis-hydroxypivalic acid, BHPA), and 2.0 equivalents (12.0 mmol) of an appropriately substituted 1-chloro- or 1-iodoalkyl carboxylate or an appropriately substituted 1-chloro- or 1-iodoalkyl alkyl- or arylcarbonate. 2.09 mL (1.55 g, 12.0 mmol) of diethylisopropylamine (DIEA, Hünigs base) were added neat. Optionally, 2 mL of anhydrous 1,2-dichloroethane was added to dissolve the reactants and to facilitate stirring. The reaction mixture was heated to ca. 70° C. and reacted overnight. Volatile reactants or solvents were removed under reduced pressure using a rotary evaporator. The residue was diluted with ethyl acetate (EtOAc) and water. After extraction, the combined extracts were washed with brine and dried over magnesium sulfate (MgSO_4), filtered and the solvents removed under reduced pressure using a rotary evaporator. Optionally, the residues were used in the next step without further purification or were purified using silica gel column chromatography with ethyl acetate/hexane mixtures as eluent.

Example 2

Ethyl 3-hydroxy-2,2-dimethylpropanoate (2)

[0429] Following the general procedure for the synthesis of hydroxypivalic ester derivatives of Description 2, 4.73 g (40.0 mmol) of 3-hydroxy-2,2-dimethylpropanoic acid (hydroxypivalic acid, HPA) was reacted with 3.23 mL (6.24 g, 40.0 mmol) of iodoethane in 80 mL of DMF in the presence of 13.03 g (40.0 mmol) of Cs_2CO_3 . After work-up, 3.40 g (68% yield) of the title compound (2) was obtained as a yellow liquid. The material was of sufficient purity to be used in the next step without further purification. ^1H NMR (400 MHz, CDCl_3): δ =1.20 (s, 6H), 1.28 (t, J =7.2 Hz, 3H), 2.50-2.55 (br. m, 1H), 3.52-3.57 (br. m, 2H), 4.16 (q, J =7.2 Hz, 2H) ppm.

Example 3

Methylethyl 3-hydroxy-2,2-dimethylpropanoate (3)

[0430] Following the general procedure for the synthesis of hydroxypivalic ester derivatives of Description 2, 4.73 g (40.0

mmol) of 3-hydroxy-2,2-dimethylpropanoic acid (hydroxypivalic acid, HPA) was reacted with 3.75 mL (4.92 g, 40.0 mmol) of 2-bromopropane in 80 mL of DMF in the presence of 13.03 g (40.0 mmol) of Cs_2CO_3 . After work-up, 4.40 g (69% yield) of the title compound (3) was obtained as a pale-yellow liquid. The material was of sufficient purity to be used in the next step without further purification. ^1H NMR (400 MHz, CDCl_3): δ =1.18 (s, 6H), 1.25 (d, J =6.4 Hz, 6H), 2.35-2.55 (br. m, 1H), 3.50-3.56 (br. m, 2H), 5.02 (heptett, J =6.0 Hz, 1H) ppm.

Example 4

Phenylmethyl 3-hydroxy-2,2-dimethylpropanoate (4)

[0431] Following the general procedure for the synthesis of hydroxypivalic ester derivatives of Description 2, 7.09 g (60.0 mmol) of 3-hydroxy-2,2-dimethylpropanoic acid (hydroxypivalic acid, HPA) was reacted with 5.94 mL (8.55 g, 50.0 mmol) of benzyl bromide in 150 mL of DMF in the presence of 16.29 g (50.0 mmol) of Cs_2CO_3 . After work-up, 9.60 g (92% yield) of the title compound (4) was obtained as a pale-yellow liquid. The material was of sufficient purity to be used in the next step without further purification. R_f =0.44 (EtOAc/Hxn=1:2). ^1H NMR (400 MHz, CDCl_3): δ =1.21 (s, 6H), 2.34-2.41 (br. m, 1H), 3.55-3.59 (br. m, 2H), 3.82 (s, 3H), 5.09 (s, 2H), 6.86-6.90 (m, 2H), 7.24-7.32 (m, 2H) ppm.

Example 5

(4-Methoxyphenyl)methyl 3-hydroxy-2,2-dimethylpropanoate (5)

[0432] Adapting a procedure or a variation thereof according to Zhou, et al., WO 99/51613, 20.0 g (143 mmol) of sodium 3-hydroxy-2,2-dimethylpropanoate was reacted with 19.7 mL (22.7 g, 145 mmol) of 4-methoxybenzyl chloride in anhydrous N-methyl pyrrolidinone (NMP) as described in the general procedure of Description 2. After work-up, the crude material was purified by silica gel column chromatography using a mixture of ethyl acetate (EtOAc) and hexane (Hxn) (EtOAc/Hxn=1:3) as eluent to provide 11.30 g (32% yield) of the title compound (5) as a pale-yellow liquid. R_f =0.38 (EtOAc/Hxn=1:2). ^1H NMR (400 MHz, CDCl_3): δ =1.21 (s, 6H), 2.34-2.41 (br. m, 1H), 3.55-3.59 (br. m, 2H), 3.82 (s, 3H), 5.09 (s, 2H), 6.86-6.90 (m, 2H), 7.24-7.32 (m, 2H) ppm.

Example 6

Phenylmethyl 3-hydroxy-2-(hydroxymethyl)-2-methylpropanoate (6)

[0433] Following the general procedure for the synthesis of hydroxypivalic ester derivatives of Description 2, 3.35 g (25.0 mmol) of 3-hydroxy-2-(hydroxymethyl)-2-methylpropanoic acid (bis-hydroxypivalic acid, BHPA) was reacted with 1.78 mL (2.57 g, 15.0 mmol) of benzylbromide in 50 mL of DMF in the presence of 6.52 g (15.0 mmol) of Cs_2CO_3 . After work-up, 3.13 g (93% yield) of the title compound (6) was obtained as colorless crystals. The material was of sufficient purity to be used in the next step without further purification. M.p.=71.6-72.8° C. ^1H NMR (400 MHz, CDCl_3): δ =1.10 (s, 3H), 2.77-2.86 (br. m, 2H), 3.75 (d, J =11.6 Hz, 2H), 3.95 (d, J =11.2 Hz, 2H), 5.22 (s, 2H), 7.31-7.41 (m, 5H) ppm.

Example 7

rac-Ethoxycarbonyloxyethyl 3-hydroxy-2,2-dimethylpropanoate (7)

[0434] Following the general procedure for the synthesis of acyloxyalkyl ester derivatives or alkoxy- and aryloxy-carbo-nyloxyalkyl derivatives of Description 2, 709 mg (6.0 mmol) of 3-hydroxy-2,2-dimethylpropanoic acid (hydroxypivalic acid, HPA) was reacted with 1.83 g (12.0 mmol) of 1-chloroethyl ethyl carbonate in the presence of 2.09 mL (1.55 g, 12.0 mmol) of diethylisopropylamine (DIEA, Hünigs base). After work-up and isolation, ca. 1.4 g (quant.) of the title compound (7) was obtained as a colorless liquid. The material was of sufficient purity to be used in the next step without further purification. R_f =0.44 (EtOAc/Hxn=1:2). ^1H NMR (400 MHz, CDCl_3): δ =1.18 (s, 3H), 1.21 (s, 3H), 1.32 (t, J =6.8 Hz, 3H), 1.53 (d, J =5.2 Hz, 3H), 2.35-2.52 (br. m, 1H), 3.48 (d, J =11.2 Hz, 1H), 3.60 (d, J =11.2 Hz, 1H), 4.22 (q, J =6.8 Hz, 2H), 6.70 (q, J =5.2 Hz, 1H) ppm. MS (ESI) m/z 257.00 (M +Na) $^+$.

Description 3

General Procedure for Synthesis of Acamprosate Neopentyl Sulfonyl ester Prodrugs or Intermediates to Prodrugs

Method A

[0435] A 100 mL round-bottomed flask equipped with a magnetic stir bar was charged at ca. 0° C. (icebath) with N-[3-(chlorosulfonyl)propyl]acetamide (1) (ca. 300 mg, 1.5 mmol), the corresponding functionalized neopentyl alcohol (3.0 mmol), and 10 mL of dichloromethane (DCM). To the solution was added 209 μL of triethylamine (TEA, Et_3N) (152 mg, 2.0 mmol) and 4-(N,N-dimethyl)aminopyridine (DMAP) (183 mg, 1.5 mmol). The reaction mixture was stirred overnight at room temperature. Upon completion of the reaction, dichloromethane was evaporated and the residue was diluted with ethyl acetate (EtOAc) and water. The aqueous layer was acidified with a one molar (1.0 M) hydrochloric acid (HCl). After vigorous mixing followed by phase separation, the aqueous layer was extracted twice more with ethyl acetate. The combined organic extracts were successively washed with a saturated aqueous sodium hydrogencarbonate (NaHCO_3) solution, brine, and then dried over MgSO_4 . After filtration, the solvent was evaporated under reduced pressure using a rotary evaporator. The residue was dissolved in a mixture of ca. 60% (v/v) acetonitrile/water and the solution filtered through a 0.2- μm nylon syringe filter and purified by mass-guided preparative HPLC. After lyophilization of the solvents, the corresponding acamprosate neopentyl sulfonyl ester prodrug was obtained as a colorless oil or solid. Alternatively, acamprosate neopentyl sulfonyl ester prodrugs were purified by silica gel chromatography using ethyl acetate/hexane or ethyl acetate/methanol mixtures as eluent followed by removal of the solvents under reduced pressure using a rotary evaporator.

Method B

[0436] Operational (synthesis and work-up) procedures in Method B were identical to the ones described in Method A. The molar ratios of the reactants were similar but N-[3-(chlorosulfonyl)propyl]acetamide (1) the commercially available 3-chloropropylsulfonyl chloride, a precursor to acamprosate

prodrugs, was employed for the synthesis of suitable neopentyl sulfonyl ester intermediates. Acamprosate neopentyl sulfonylester intermediates were generally purified by silica gel chromatography using ethyl acetate (EtOAc) or methyl tert-butyl ether (MTBE) and hexane or n-heptane mixtures as eluent followed by removal of the solvents under reduced pressure using a rotary evaporator.

Example 8

Methyl 3-[[3-(acetylamino)propyl]sulfonyloxy]-2,2-dimethylpropanoate (8)

[0437] Following the general procedure for the synthesis of acamprosate neopentyl sulfonylester prodrugs of Description 3 (Method A), N-[3-(chlorosulfonyl)propyl]acetamide (1) (ca. 1.00 g, 5.0 mmol), dissolved in 8 mL of dichloromethane, was reacted with 397 mg (3.0 mmol) of commercially available methyl 3-hydroxy-2,2-dimethylpropanoate (2) in the presence of 697 μ L of triethylamine (506 mg, 5.0 mmol) and 611 mg (5.0 mmol) of DMAP. After aqueous work-up and purification by mass-guided preparative HPLC, 205 mg (23% yield) of the title compound (8) was obtained as a colorless, viscous oil. ¹H NMR (400 MHz, DMSO-d₆): δ =1.18 (s, 6H), 1.73-1.82 (m, 5H), 3.12 (q, J=5.6 Hz, 2H), 3.29-3.36 (m, 2H), 3.64 (s, 3H), 4.18 (s, 2H), 7.90 (br. t, J=5.2 Hz, 1H) ppm. MS (ESI) m/z 296.08 (M+H)⁺, 318.09 (M+Na)⁺.

Example 9

Ethyl 3-[[3-(acetylamino)propyl]sulfonyloxy]-2,2-dimethylpropanoate (9)

[0438] Following the general procedure for the synthesis of acamprosate neopentyl sulfonylester prodrugs of Description 3 (Method A), N-[3-(chlorosulfonyl)propyl]acetamide (1) (ca. 500 mg, 2.5 mmol) dissolved in 8 mL of dichloromethane was reacted with 585 mg (4.0 mmol) of ethyl 3-hydroxy-2,2-dimethylpropanoate (2) in the presence of 348 μ L of triethylamine (253 mg, 2.5 mmol) and 305 mg (2.5 mmol) of DMAP. Following aqueous work-up and purification by mass-guided preparative HPLC, 208 mg (27% yield) of the title compound (9) was obtained as a colorless, viscous oil. ¹H NMR (400 MHz, CDCl₃): δ =1.28 (s, 6H), 1.29 (t, J=7.2 Hz, 3H), 2.01 (s, 3H), 2.04-2.11 (m, 2H), 3.17-3.22 (m, 2H), 3.40 (q, J=6.4 Hz, 2H), 4.17 (q, J=7.2 Hz, 2H), 4.20 (s, 2H), 5.97-6.04 (br. m, 1H) ppm. MS (ESI) m/z 310.10 (M+H)⁺, 332.04 (M+Na)⁺.

Example 10

Methylethyl 3-[[3-(acetylamino)propyl]sulfonyloxy]-2,2-dimethylpropanoate (10)

[0439] Following the general procedure for the synthesis of acamprosate neopentyl sulfonylester prodrugs of Description 3 (Method A), N-[3-(chlorosulfonyl)propyl]acetamide (1) (ca. 500 mg, 2.5 mmol), dissolved in 8 mL of dichloromethane, was reacted with 585 mg (4.0 mmol) of methylethyl 3-hydroxy-2,2-dimethylpropanoate (3) in the presence of 348 μ L of triethylamine (253 mg, 2.5 mmol) and 305 mg (2.5 mmol) of DMAP. After aqueous work-up and purification by mass-guided preparative HPLC, 192 mg (24% yield) of the title compound (10) was obtained as a colorless, viscous oil. ¹H NMR (400 MHz, CDCl₃): δ =1.25-1.28 (m, 12H), 2.01 (s, 3H), 2.04-2.12 (m, 2H), 3.17-3.22 (m, 2H), 3.41 (q,

J=6.8 Hz, 2H), 4.19 (s, 2H), 5.01 (heptett, J=6.4 Hz, 1H), 5.94-6.02 (br. m, 1H) ppm. MS (ESI) m/z 324.12 (M+H)⁺, 346.06 (M+Na)⁺.

Example 11

Phenylmethyl 3-[[3-(acetylamino)propyl]sulfonyloxy]-2,2-dimethylpropanoate (11)

[0440] Following the general procedure for the synthesis of acamprosate neopentyl sulfonylester prodrugs of Description 3 (Method A), N-[3-(chlorosulfonyl)propyl]acetamide (1) (ca. 8.98 g, 45.0 mmol) dissolved in 150 mL of dichloromethane was reacted with 10.41 g (50.0 mmol) of phenylmethyl 3-hydroxy-2,2-dimethylpropanoate (4) in the presence of 6.97 mL of triethylamine (5.06 g, 50 mmol) and 6.11 g (50 mmol) of DMAP. After aqueous work-up, the crude material was purified by silica gel column chromatography using mixtures of ethyl acetate (EtOAc) and hexane (Hxn) as eluent (EtOAc/Hxn=6:1→9:1→14:1→100% EtOAc) to provide 5.50 g (33% yield) of the title compound (11) as a colorless, viscous oil. R_f=0.59 (EtOAc/MeOH=19:1). ¹H NMR (400 MHz, CDCl₃): δ =1.31 (s, 6H), 1.92-2.01 (m, 5H), 3.06-3.12 (m, 2H), 3.32 (q, J=6.4 Hz, 2H), 4.21 (s, 2H), 5.16 (s, 2H), 5.78-5.84 (br. m, 1H), 7.31-7.41 (m, 5H) ppm. MS (ESI) m/z 372.13 (M+H)⁺, 394.08 (M+Na)⁺.

Example 12

3-[[3-(Acetylamino)propyl]sulfonyloxy]-2,2-dimethylpropanoic acid (12)

[0441] A 100 mL round-bottomed flask equipped with a magnetic stirring bar and a three-way adapter connected to a hydrogen filled balloon at ca. 15 psi was charged with 505 mg (1.36 mmol) of phenylmethyl 3-[[3-(acetylamino)propyl]sulfonyloxy]-2,2-dimethylpropanoate (11) and 400 mg of 10 wt-% palladium on activated carbon. Twenty-five (25) mL of anhydrous ethanol (EtOH) was added and the atmosphere was exchanged to hydrogen with five alternating evacuation/refill cycles. The reaction mixture was stirred for two hours at room temperature and then filtered through a short plug of Celite®. The filter medium was washed several times with EtOH. After partial evaporation of the solvent under reduced pressure using a rotary evaporator, the residual solution was filtered through a 0.2 μ m nylon syringe filter and evaporated under reduced pressure to provide 383 mg (quant.) of the title compound (12) as a colorless oil. ¹H NMR (400 MHz, DMSO-d₆): δ =1.15 (s, 6H), 1.74-1.82 (m, 5H), 3.08-3.16 (m, 2H), 3.29-3.37 (m, 2H), 4.15 (s, 2H), 7.90 (t, J=5.6 Hz, 1H), 12.63 (br. s, 1H) ppm. MS (ESI) m/z 282.13 (M+H)⁺, 304.00 (M+Na)⁺, 280.03 (M-H)⁻.

Example 13

(4-Methoxyphenyl)methyl 3-[[3-(acetylamino)propyl]sulfonyloxy]-2,2-dimethylpropanoate (13)

[0442] Following the general procedure for the synthesis of acamprosate neopentyl sulfonylester prodrugs of Description 3 (Method A), N-[3-(chlorosulfonyl)propyl]acetamide (1) (ca. 500 mg, 2.5 mmol) dissolved in 8 mL of dichloromethane was reacted with 715 mg (3.0 mmol) of (4-methoxyphenyl)methyl 3-hydroxy-2,2-dimethylpropanoate (5) in the presence of 348 μ L of triethylamine (253 mg, 2.5 mmol) and 305 mg (2.5 mmol) of DMAP. After aqueous work-up, the crude product was purified by silica gel column chromatography

using a mixture of ethyl acetate (EtOAc) and methanol (MeOH) as eluent (EtOAc/MeOH=19:1) to provide 150 mg (15% yield) of the title compound (13) as a colorless, viscous oil. R_f =0.32 (EtOAc). ^1H NMR (400 MHz, CDCl_3): δ =1.28 (s, 6H), 1.93-2.02 (m, 5H), 3.06-3.12 (m, 2H), 3.34 (q, J =6.4 Hz, 2H), 3.82 (s, 3H), 4.19 (s, 2H), 5.09 (s, 2H), 5.81-5.88 (br. m, 1H), 6.87-6.92 (m, 2H), 7.26-7.30 (m, 2H) ppm. MS (ESI) m/z 402.14 (M+H) $^+$, 424.09 (M+Na) $^+$.

Example 14

Phenylmethyl 3- $\{[3-(\text{acetylamino})\text{propyl}]\text{sulfonyloxy}\}$ -2- $\{[3-(\text{acetylamino})\text{propyl}]\text{sulfonyloxy}\}$ methyl-2-methylpropanoate (14)

[0443] Following the general procedure for the synthesis of acamprostate neopentyl sulfonylester prodrugs of Description 3 (Method A), N-[3-(chlorosulfonyl)propyl]acetamide (1) (ca. 1.00 g, 5.0 mmol) dissolved in 15 mL of dichloromethane was reacted with 300 mg (1.34 mmol) of phenylmethyl 3-hydroxy-2-(hydroxymethyl)-2-methylpropanoate (6) in the presence of 697 μL of triethylamine (506 mg, 5.0 mmol) and 611 mg (5.0 mmol) of DMAP. After aqueous work-up and purification by mass-guided preparative HPLC, 262 mg (36% yield) of the title compound (14) was obtained as a colorless, viscous oil. ^1H NMR (400 MHz, CDCl_3): δ =1.33 (s, 3H), 1.97-2.06 (m, 10H), 3.14-3.19 (m, 4H), 3.32-3.39 (m, 4H), 4.30 (d, J =9.6 Hz, 2H), 4.42 (d, J =10.0 Hz, 2H), 5.20 (s, 2H), 6.20 (br. t, J =6.0 Hz, 2H), 7.32-7.41 (m, 5H) ppm. MS (ESI) m/z 551.24 (M+H) $^+$, 574.09 (M+Na) $^+$.

Example 15

rac-Phenylmethyl 2- $\{[3-(\text{acetylamino})\text{propyl}]\text{sulfonyloxy}\}$ -methyl-3-hydroxy-2-methylpropanoate (15)

[0444] The corresponding mono-sulfonylester derivative (15) was also isolated from the crude reaction product of Example 14 by mass-guided preparative HPLC and was obtained as a colorless oil. ^1H NMR (400 MHz, CDCl_3): δ =1.27 (s, 3H), 1.97-2.08 (m, 5H), 3.11-3.20 (m, 2H), 3.31-3.38 (m, 2H), 3.72 (d, J =10.8 Hz, 1H), 3.87 (d, J =11.2 Hz, 1H), 4.34 (d, J =9.6 Hz, 1H), 4.47 (d, J =10.0 Hz, 1H), 5.19 (s, 2H), 5.83-5.90 (br. m, 1H), 7.31-7.41 (m, 5H) ppm. MS (ESI) m/z 388.10 (M+H) $^+$, 410.00 (M+Na) $^+$.

Example 16

rac-Ethoxycarbonyloxyethyl 3- $\{[3-(\text{acetylamino})\text{propyl}]\text{sulfonyloxy}\}$ -2,2-dimethylpropanoate (16)

[0445] Following the general procedure for the synthesis of acamprostate neopentyl sulfonylester prodrugs of Description 3 (Method A), N-[3-(chlorosulfonyl)propyl]acetamide (1) (ca. 1.00 g, 5.0 mmol) dissolved in 15 mL of dichloromethane was reacted with ca. 1.4 g (6.0 mmol) of rac-ethoxycarbonyloxyethyl 3-hydroxy-2,2-dimethylpropanoate (7) in the presence of 697 μL of triethylamine (506 mg, 5.0 mmol) and 611 mg (5.0 mmol) of DMAP. After aqueous work-up, the crude material was purified by silica gel column chromatography using mixtures of ethyl acetate (EtOAc) and hexane (Hxn) as eluent (EtOAc/Hxn=4:1 $\rightarrow$ 9:1 $\rightarrow$ 100% EtOAc) to provide 434 mg (22% yield) of the title compound (16) as a colorless, viscous oil. R_f =0.24 (EtOAc/Hxn=4:1). ^1H NMR (400 MHz, CDCl_3): δ =1.29 (s, 6H), 1.34 (t, J =7.2 Hz, 3H), 1.55 (d, J =5.6 Hz, 3H), 2.01 (s, 3H), 2.03-2.11 (m, 2H), 3.16-3.24 (m, 2H),

3.35-3.46 (m, 2H), 4.15-4.28 (m, 4H), 5.40-6.40 (br. m, 1H), 6.76 (q, J =5.6 Hz, 1H) ppm. MS (ESI) m/z 398.01 (M+H) $^+$, 419.95 (M+Na) $^+$.

Example 17

rac-Methylethoxycarbonyloxyethyl 3- $\{[3-(\text{acetylamino})\text{propyl}]\text{sulfonyloxy}\}$ -2,2-dimethylpropanoate (17)

[0446] The title compound (17), an oxidative degradation product produced during preparation of pantoic acid based neopentyl sulfonylester acamprostate prodrugs, was isolated following mass-guided preparative HPLC purification as a colorless, viscous oil (57 mg). R_f =0.15 (EtOAc/Hxn=4:1). ^1H NMR (400 MHz, CDCl_3): δ =1.29 (s, 6H), 1.327/1.330 (2d, superimposed, J =6.4 Hz, 6H), 1.55 (d, J =5.6 Hz, 3H), 2.07 (s, 3H), 2.03-2.11 (m, 2H), 3.18-3.23 (m, 2H), 3.34-3.47 (m, 2H), 4.16 (d, J =9.6 Hz, 1H), 4.21 (d, J =9.2 Hz, 1H), 4.84-4.94 (m, 1H), 5.95-6.01 (br. m, 1H), 6.75 (q, J =5.6 Hz, 1H) ppm. MS (ESI) m/z 412.0 (M+H) $^+$, 434.0 (M+Na) $^+$.

Example 18

2-Morpholin-4-ylethyl 3- $\{[3-(\text{acetylamino})\text{propyl}]\text{sulfonyloxy}\}$ -2,2-dimethylpropanoate Hydrochloride (18)

Step A: 2,2-Dimethyl-3-(1,1,2,2-tetramethyl-1-silapropoxy)propanoic acid (18a)

[0447] A 2 L round bottomed flask equipped with a magnetic stir bar was charged with 11.8 g (100 mmol) of 3-hydroxy-2,2-dimethylpropanoic acid and the acid dissolved in 1 L of anhydrous dichloromethane (DCM). Tert-butyl dimethylchlorosilane (31.7 g, 210 mmol) and imidazole (14.3 g, 210 mmol) was added while stirring. The reaction was monitored by TLC. After the starting material was completely consumed the white precipitate was filtered off and the solvent removed under reduced pressure using a rotary evaporator. The residue was dissolved in 500 mL of diethyl ether (Et_2O). The mixture was stirred for two hours and the colorless precipitate filtered off. After removing diethyl ether under reduced pressure using a rotary evaporator, the residue was dissolved in 600 mL of a mixture of water and acetonitrile (1:1). 8.0 g (200 mmol) of sodium hydroxide was added to the solution and the mixture was stirred overnight at room temperature. The reaction was acidified with a one normal (1N) aqueous solution of hydrogen chloride (HCl) and extracted with ethyl acetate. The combined organic extracts were washed with water and brine, dried over magnesium sulfate (MgSO_4), filtered, and the solvents removed under reduced pressure using a rotary evaporator. The residue was purified by silica gel chromatography using a mixture of ethyl acetate (EtOAc) and hexane (Hxn) (EtOAc/Hxn=1:4) as eluent to provide 23 g (quant.) of the title compound (18a) as a colorless liquid. R_f =0.58 (EtOAc/Hexane=1:4). ^1H NMR (400 MHz, CDCl_3): δ =0.07 (s, 6H), 0.90 (s, 9H), 1.19 (s, 6H), 3.60 (s, 2H) ppm. MS (ESI) m/z 233.01 (M+H) $^+$.

Step B: 2-Morpholin-4-ylethyl 2,2-dimethyl-3-(1,1,2,2-tetramethyl-1-silapropoxy)propanoate (18b)

[0448] To a stirred solution of 2,2-dimethyl-3-(1,1,2,2-tetramethyl-1-silapropoxy)propanoic acid (18a) (2.3 g, 10.0 mmol) in 100 mL of anhydrous dichloromethane (DCM) in a 250 mL round-bottomed flask equipped with a magnetic stir

bar was added N,N'-dicyclohexylcarbodiimide (2.3 g, 11.0 mmol), N-(2-hydroxyethyl)morpholine (1.2 g, 9.0 mmol) and 0.15 g (1.1 mmol) of 4-N,N-dimethylaminopyridine (DMAP). The reaction was stirred overnight at room temperature. The colorless precipitate was filtered off and the solvent removed under reduced pressure using a rotary evaporator. The residue was purified by silica gel chromatography using a mixture of ethyl acetate (EtOAc) and hexane (Hxn) (EtOAc/Hxn=1:1) as eluent to provide 0.53 g (17% yield) of the title compound (18b) as a colorless solid. R_f =0.4 (EtOAc/Hxn=1:1). ^1H NMR (400 MHz, CDCl_3): δ =0.04 (s, 6H), 0.88 (s, 9H), 1.16 (s, 6H), 2.50-2.54 (m, 4H), 2.63 (t, J =6.0 Hz, 2H), 3.57 (s, 2H), 3.68-3.73 (m, 4H), 4.21 (t, J =6.0 Hz, 2H) ppm. MS (ESI) m/z 346.26 (M+H) $^+$.

Step C: 2-Morpholin-4-ylethyl
3-hydroxy-2,2-dimethylpropanoate (18c)

[0449] Adapting a procedure, or a variation thereof according to Pirrung et al., *Bioorg. Med. Chem. Lett.*, 1994, 4, 1345, 0.73 g of triethylamine trihydrofluoride (4.5 mmol) was added to a solution of 2-morpholin-4-ylethyl 2,2-dimethyl-3-(1,1,2,2-tetramethyl-1-silapropoxy)propanoate (18b) (0.52 g, 1.5 mmol) in 10 mL of anhydrous tetrahydrofuran (THF) while stirring. The reaction mixture was stirred overnight at 50° C. After the starting material was completely consumed, the solvent was removed under reduced pressure using a rotary evaporator. The residue was purified by silica gel chromatography using a mixture of ethyl acetate (EtOAc) and methanol (MeOH) (EtOAc/MeOH=4:1) as eluent to provide 330 mg (95% yield) of the title compound (18c). R_f =0.29 (EtOAc). ^1H NMR (400 MHz, CDCl_3): δ =1.21 (s, 6H), 2.47-2.54 (m, 4H), 2.58-2.63 (m, 2H), 3.54 (s, 2H), 3.69-3.73 (m, 4H), 4.33-4.37 (m, 2H) ppm. MS (ESI) m/z 232.10 (M+H) $^+$.

Step D: 2-Morpholin-4-ylethyl 3-[[3-(acetylamino)
propyl]sulfonyloxy]-2,2-dimethylpropanoate Hydro-
chloride (18)

[0450] Following the general procedure for the synthesis of acamprostate neopentyl sulfonylester prodrugs of Description 3 (Method A), N-[3-(chlorosulfonyl)propyl]acetamide (1) (0.43 g, 2.1 mmol) dissolved in 10 mL of dichloromethane (DCM) was reacted with 2-morpholin-4-ylethyl 3-hydroxy-2,2-dimethylpropanoate (18c) (0.33 g, 1.43 mmol) in the presence of 0.30 mL of triethylamine (0.22 g, 2.1 mmol) and 24 mg (0.2 mmol) of DMAP. After mass-guided preparative HPLC purification and lyophilization in the presence of a slight excess of a one molar (1.0 M) hydrochloric acid (HCl), 60 mg (10% yield) of the title compound (18) was obtained as a colorless solid. ^1H NMR (400 MHz, CDCl_3): δ =1.29 (s, 6H), 2.01 (s, 3H), 2.02-2.10 (m, 2H), 2.62-2.66 (m, 4H), 2.76 (t, J =6.0 Hz, 2H), 3.17-3.22 (m, 2H), 3.37-3.43 (m, 2H), 3.72-3.77 (m, 4H), 4.20 (s, 2H), 4.30 (t, J =6.0 Hz, 2H), 6.12-6.20 (m, 1 H) ppm. MS (ESI) m/z 395.10 (M+H) $^+$.

Example 19

3-Pyridylmethyl 3-[[3-(acetylamino)propyl]sulfonyloxy]-2,2-dimethylpropanoate Hydrochloride (19)

Step A: 2-Pyridylmethyl 2,2-dimethyl-3-(1,1,2,2-tetramethyl-1-silapropoxy)propanoate (19a)

[0451] One (1.0) mL of 3-pyridinemethanol (1.2 g, 11.0 mmol) and 4-N,N-dimethylaminopyridine (DMAP) (0.15 g, 1.1 mmol) were added to a solution of 2,2-dimethyl-3-(1,1,

2,2-tetramethyl-1-silapropoxy)propanoic acid (18a) (Example 18, Step A) (2.3 g, 10.0 mmol) in 100 mL of anhydrous dichloromethane (DCM) in a 250 mL round-bottomed flask equipped with a magnetic stir bar was added N,N'-dicyclohexylcarbodiimide (2.3 g, 11.0 mmol) while stirring. The reaction mixture was stirred overnight at room temperature. The colorless precipitate was then filtered off and the solvent removed under reduced pressure using a rotary evaporator. The residue was purified by silica gel chromatography using a mixture of ethyl acetate (EtOAc) and hexanes (Hxn) (EtOAc/Hxn=1:2) as eluent to provide 1.2 g (37% yield) of the title compound (19a) as colorless liquid. R_f =0.43 (EtOAc/Hxn=1:2). ^1H NMR (400 MHz, CDCl_3): δ =0.01 (s, 6H), 0.86 (s, 9H), 1.19 (s, 6H), 3.60 (s, 2H), 5.13 (s, 2H), 7.24-7.30 (m, 1H), 7.65-7.69 (m, 1H), 8.55-8.57 (m, 1H), 8.59-8.61 (m, 1H) ppm. MS (ESI) m/z 324.16 (M+H) $^+$.

Step B: 3-Pyridylmethyl
3-hydroxy-2,2-dimethylpropanoate (19b)

[0452] Adapting a procedure, or a variation thereof according to Pirrung et al., *Bioorg. Med. Chem. Lett.*, 1994, 4, 1345, 1.77 g (11.0 mmol) of triethylamine trihydrofluoride was added to a stirred solution of 3-pyridylmethyl 2,2-dimethyl-3-(1,1,2,2-tetramethyl-1-silapropoxy)propanoate (19a) (1.2 g, 3.7 mmol) in 30 mL of anhydrous tetrahydrofuran (THF). The reaction mixture was stirred overnight at 50° C. After the starting material was completely consumed, the solvent was removed under reduced pressure using a rotary evaporator. The residue was purified by silica gel chromatography using a mixture of ethyl acetate (EtOAc) and methanol (MeOH) (EtOAc/MeOH=10:1) as an eluent to provide 800 mg (quant.) of the title compound (19b) as a colorless solid. R_f =0.34 (EtOAc). ^1H NMR (400 MHz, CDCl_3): δ =1.23 (s, 6H), 2.38-2.52 (br. m, 1H), 3.60 (s, 2H), 5.17 (s, 2H), 7.25-7.33 (m, 1H), 7.65-7.69 (m, 1H), 8.56-8.59 (m, 1 H), 8.60-8.62 (m, 1 H) ppm. MS (ESI) m/z 209.99 (M+H) $^+$.

Step C: 3-Pyridylmethyl 3-[[3-(acetylamino)propyl]
sulfonyloxy]-2,2-dimethylpropanoate Hydrochloride
(19)

[0453] Following the general procedure for the synthesis of acamprostate neopentyl sulfonylester prodrugs of Description 3 (Method A), N-[3-(chlorosulfonyl)propyl]acetamide (1) (0.76 g, 3.8 mmol) dissolved in 20 mL of dichloromethane was reacted with 3-pyridylmethyl 3-hydroxy-2,2-dimethylpropanoate (19b) (0.4 g, 1.9 mmol) in the presence of 0.53 mL of triethylamine (0.38 g, 3.8 mmol) and 50 mg (0.4 mmol) of DMAP. After mass-guided preparative HPLC purification and lyophilization in the presence of a slight excess of a one molar (1.0 M) hydrochloric acid (HCl), 370 mg (47% yield) of the title compound (19) was obtained. ^1H NMR (400 MHz, $\text{MeOH}-d_4$): δ =1.31 (s, 6H), 1.97-2.05 (m, 2H), 2.15 (s, 3H), 3.29-3.34 (m, 2H), 4.41 (t, J =6.8 Hz, 2H), 4.30 (s, 2H), 5.42 (s, 2H), 8.12-8.16 (m, 1H), 8.69-8.73 (m, 1H), 8.85-8.87 (m, 1H), 8.96-8.97 (m, 1H) ppm. MS (ESI) m/z 373.08 (M+H) $^+$.

Example 20

rac-Methyl 3-[[3-(acetylamino)propyl]sulfonyloxy]-
2-[(tert-butoxy)carbonylamino]-2-methylpropanoate
(20)

Step A: rac-Methyl 2-amino-3-hydroxy-2-methyl-
propanoate Hydrochloride (20a)

[0454] Adapting a procedure or variation thereof according to Clemens et al., *Bioorg. Med. Chem. Lett.* 2004, 14, 4903-

4906, a 500 mL round-bottomed flask equipped with a magnetic stirring bar and a reflux condenser was charged with commercially available 2-amino-3-hydroxy-2-methylpropanoic acid (1.67 g, 14.0 mmol) and 200 mL of anhydrous methanol (MeOH). To the stirred solution was added 110 μ L (0.18 g, 1.5 mmol) of neat thionyl chloride (SOCl_2) and the reaction mixture was heated to reflux overnight. The solvent was then removed under reduced pressure using a rotary evaporator. The residue was purified by mass-guided preparative HPLC to provide 550 mg (30% yield) of the title compound (20a) as a colorless solid. ^1H NMR (400 MHz, MeOH-d^4): δ =1.52 (s, 3H), 3.38 (s, 3H), 3.68 (d, J_{gem} =11.2 Hz, 1H), 3.97 (d, J_{gem} =11.2 Hz, 1H) ppm. MS (ESI) m/z 133.94 ($\text{M}+\text{H}$) $^+$. The analytical data for the compound was consistent with that given in then literature.

Step B: rac-Methyl 2-[(tert-butoxy)carbonylamino]-3-hydroxy-2-methylpropanoate (20b)

[0455] Adapting a procedure or variation thereof according to Clemens et al., *Bioorg. Med. Chem. Lett.* 2004, 14, 4903-4906, a 500 mL round-bottomed flask equipped with a magnetic stir bar was charged with (2.5 g, 15.0 mmol) of methyl 2-amino-3-hydroxy-2-methylpropanoate hydrochloride (20a) and 200 mL of a mixture of acetonitrile (H_3CCN) and water ($v/v=1:1$). To the stirred solution was added sodium hydrogen carbonate (NaHCO_3) (2.8 g, 45.0 mmol) and 6.5 g (30.0 mmol) of di-tert-butyl dicarbonate (di-tert-butyl pyrocarbonate, Boc_2O) and the reaction mixture was stirred overnight at room temperature. After the starting material was completely consumed, the reaction was diluted with water and the aqueous phase extracted with ethyl acetate (EtOAc). The organic extracts were washed in brine, dried over anhydrous magnesium sulfate (MgSO_4), filtered, and the solvents removed under reduced pressure using a rotary evaporator. The residue was purified by silica gel chromatography using ethyl acetate (EtOAc) as eluent to provide 0.8 g (24% yield) of the title compound (20b) as a colorless oil. R_f =0.51 (EtOAc). ^1H NMR (400 MHz, CDCl_3): δ =1.46 (s, 9H), 1.49 (s, 3H), 3.77-3.79 (m, 4H), 4.00 (d, J =11.6 Hz, 1H), 5.29 (br. s, 1H) ppm. MS (ESI) m/z 234.06 ($\text{M}+\text{H}$) $^+$. The analytical data for the compound was consistent with that given in the literature.

Step C: Methyl 3-[[3-(Acetylamino)propyl]sulfonyloxy]-2-[(tert-butoxy)carbonylamino]-2-methylpropanoate (20)

[0456] Following the general procedure for the synthesis of acamprostate neopentyl sulfonylester prodrugs of Description 3 (Method A), N-[3-(chlorosulfonyl)propyl]acetamide (1) (ca. 0.4 g, 2.0 mmol) dissolved in 20 mL of dichloromethane was reacted with methyl 2-[(tert-butoxy)carbonylamino]-3-hydroxy-2-methylpropanoate (20b) (0.25 g, 1.1 mmol) in the presence of 0.28 mL of triethylamine (0.20 g, 2.0 mmol) and 24 mg (0.2 mmol) of DMAP. After mass-guided preparative HPLC purification, 210 mg (49% yield) of the title compound (20) was obtained as a clear oil. ^1H NMR (400 MHz, CDCl_3): δ =1.45 (s, 9H), 1.54 (s, 3H), 2.01 (s, 3H), 2.04-2.09 (m, 2H), 3.18 (t, J =7.2 Hz, 2H), 3.39 (q, J =5.6 Hz, 2H), 3.80 (s, 3H), 4.57 (d, J =10.4 Hz, 1H), 4.68 (br. d, J =10.4 Hz, 1H), 5.40 (br. s, 1H), 5.94-6.00 (br. m, 1H) ppm. MS (ESI) m/z 397.03 ($\text{M}+\text{H}$) $^+$.

Example 21

Methyl 3-[[3-(acetylamino)propyl]sulfonyloxy]-2-amino-2-methylpropanoate Hydrochloride (21)

[0457] To a stirred solution of methyl 3-[[3-(acetylamino)propyl]sulfonyloxy]-2-[(tert-butoxy)carbonylamino]-2-me-

thylpropanoate (20) (0.13 g, 0.33 mmol) in 5 mL of dichloromethane (DCM) was added 5 mL of trifluoroacetic acid ($\text{F}_3\text{CO}_2\text{H}$, TFA). The reaction was stirred at room temperature for two hours. The solvent was then removed under reduced pressure using a rotary evaporator. The residue was further purified using mass-guided preparative HPLC. Removal of the solvent by lyophilization in the presence of a slight excess of an one molar (1.0 M) hydrochloric acid (HCl) afforded 60 mg (55% yield) of the title compound (21). ^1H NMR (400 MHz, MeOH-d^4): δ =1.66 (s, 3H), 2.05-2.13 (m, 2H), 2.14 (s, 3H), 3.42-3.48 (m, 4H), 3.93 (s, 3H), 4.47 (d, J =10.8 Hz, 2H), 4.70 (d, J =11.2 Hz, 2H) ppm. MS (ESI) m/z 297.03 ($\text{M}+\text{H}$) $^+$.

Example 22

Ethyl 3-[[3-(acetylamino)propyl]sulfonyloxy]-2,2-diethoxypropanoate (22)

Step A : rac-Ethyl glycerate (22a)

[0458] Adapting a procedure according to Choi et al., *Bioorg. Med. Chem.* 1996, 4, 2105-2114, a 1,000 mL round bottomed flask equipped with a magnetic stirring bar was charged with 15.65 g (99.0 mmol) of potassium permanganate (KMnO_4). The oxidant was dissolved in 150 mL of water and 300 mL of acetone and the reaction mixture was cooled to ca. -78°C . using a dry ice/acetone bath. 9.75 mL of ethyl acrylate (9.01 g, 90.0 mmol) was slowly added while stirring at -78°C . The reaction mixture was slowly warmed to 0°C . The inorganic salts were removed by filtration and the filter residue was washed with 150 mL of acetone. The combined filtrates were concentrated under reduced pressure at temperatures below 40°C . using a rotary evaporator. The product was extracted with ethyl acetate (3 $\times$ 200 mL) and the combined extracts were dried over anhydrous sodium sulfate (Na_2SO_4). After filtration, the solvent was removed under reduced pressure using a rotary evaporator to provide 4.2 g (35% yield) of the title compound (22a) as a clear, colorless oil. The material was of sufficient purity to be used in the next step without further purification. R_f =0.60 ($\text{MeOH}/\text{CHCl}_3=3:7$). ^1H NMR (CDCl_3 , 400 MHz): δ =1.32 (t, J =7.2 Hz, 3H), 2.30-2.70 (br. m, 1H), 3.30-3.60 (br. m, 1H), 3.85 (dd, J =11.6, 3.6 Hz, 1H), 3.91 (dd, J =11.2, 2.8 Hz, 1H), 4.23-4.32 (m, 3H) ppm. MS (ESI) m/z 135.07 ($\text{M}+\text{H}$) $^+$. The analytical data for the compound was consistent with the data given in the literature.

Step B: rac-Ethyl 2-hydroxy-3-(1,1,2,2-tetramethyl-1-silapropoxy)propanoate (22b)

[0459] Adapting procedures or variations thereof according to Shin et al., *J. Org. Chem.* 2000, 65, 7667-7675 and Parkkari et al., *Bioorganic Med. Chem. Lett.* 2004, 14, 3231-3234, a 250 mL round-bottomed flask equipped with a magnetic stirring bar was charged with 1.0 g (7.45 mmol) of rac-ethyl glycerate (22a) and the material dissolved in 20 mL of anhydrous dichloromethane (DCM). Imidazole (0.51 g, 7.45 mmol), 7.45 mL of a one molar (1.0 M solution of tert-butyl dimethylchlorosilane in DCM, and 0.4 g, 2.98 mmol of 4-(N,N-dimethylamino)pyridine (DMAP) were added. The reaction mixture was stirred overnight at room temperature. The reaction mixture was diluted with 100 mL of DCM and washed twice with water and brine. The combined organic extracts were dried over anhydrous magnesium sulfate (MgSO_4). After filtration, the solvent was removed

under reduced pressure using a rotary evaporator to provide 1.8 g (97% yield) of the title compound (22b) as a clear oil that was of sufficient purity to be used in the next step without further purification. $R_f=0.50$ (EtOAc/Hxn=1:1). ^1H NMR (CDCl_3 , 400 MHz): $\delta=0.05$ (s, 3H), 0.06 (s, 3H), 0.87 (s, 9H), 1.30 (t, $J=7.2$ Hz, 3H), 2.98-3.08 (br. m, 1H), 3.85 (dd, $J=10.0$, 3.2 Hz, 1H), 3.92 (dd, $J=10.4$, 3.2 Hz, 1H), 4.17-4.20 (m, 1H), 4.24 (q, $J=7.6$ Hz, 2H) ppm. MS (ESI) $m/z=249.07$ ($\text{M}+\text{H}$) $^+$. The analytical data for the compound was consistent with the proposed structure of the compound as described in Carreiro, et al., ES 2006959.

Step C: Ethyl 2-oxo-3-(1,1,2,2-tetramethyl-1-silapropoxy)propanoate (22c)

[0460] Adapting procedures or variations thereof according to Dess et al., *J. Am. Chem. Soc.* 1991, 113, 7277-7287, 1.6 g (3.77 mmol) of 1,1,1-tris(acetyloxy)-1,1-dihydro-1,2-benziodoxol-3-(1H)-one (Dess-Martin periodinane) was added to a solution of 0.8 g (3.22 mmol) of ethyl 2-hydroxy-3-(1,1,2,2-tetramethyl-1-silapropoxy)propanoate (22b) in 15 mL of anhydrous dichloromethane while stirring at ca. 0° C. (ice bath). The reaction mixture was stirred for six hours at room temperature. The solvent was removed under reduced pressure using a rotary evaporator and the residue diluted with 100 mL of diethyl ether and washed twice with a saturated sodium bicarbonate (NaHCO_3) solution and subsequently with an aqueous 10 wt-% solution of sodium thiosulfate ($\text{Na}_2\text{S}_2\text{O}_3$). The combined organic extracts were dried over anhydrous magnesium sulfate (MgSO_4). The solvents were removed under reduced pressure using a rotary evaporator. The residue was purified by silica gel chromatography using a mixture of ethyl acetate (EtOAc) and hexane (Hxn) as eluent (EtOAc/Hxn=1:4) to provide 0.470 g, (60% yield) of the title compound (22c) as a colorless oil. $R_f=0.50$ (EtOAc/Hxn=1:4). ^1H NMR (CDCl_3 , 400 MHz): $\delta=0.000$ (s, 6H), 0.81 (s, 9H), 1.26 (t, $J=7.2$ Hz, 3H), 4.22 (q, $J=7.2$ Hz, 2H), 4.62 (s, 2H) ppm. MS (ESI) m/z 247.07 ($\text{M}+\text{H}$) $^+$.

Step D: Ethyl 2,2-diethoxy-3-(1,1,2,2-tetramethyl-1-silapropoxy) propanoate (22d)

[0461] To a stirred solution of ethyl 2-oxo-3-(1,1,2,2-tetramethyl-1-silapropoxy)propanoate (22c), 0.362 g (1.469 mmol) in 20 mL of triethylorthoformate at ca. 0° C. (ice bath) was added a catalytic amount (ca. 0.5 mL) of concentrated sulfuric acid (H_2SO_4). The reaction mixture was stirred at 0° C.-10° C. for three to four hours. The reaction mixture was then diluted with dichloromethane (DCM) and washed with water and brine. The organic extract was dried over anhydrous magnesium sulfate (MgSO_4), filtered, and the solvents removed under reduced pressure using a rotary evaporator to provide 0.2 g (43% yield) of the title compound (22d) as a colorless oil that was of sufficient purity to be used in the next step without further purification. $R_f=0.20$ (EtOAc/Hxn=1:4). ^1H NMR (CDCl_3 , 400 MHz): $\delta=0.000$ (s, 6H), 0.83 (s, 9H), 1.19 (t, $J=6.8$ Hz, 3H), 1.27 (t, $J=7.2$ Hz, 6H), 3.39-3.48 (m, 2H), 3.54-3.64 (m, 2H), 3.81 (s, 2H), 4.21 (q, $J=7.2$ Hz, 2H) ppm.

Step E: Ethyl 2,2-diethoxy-3-hydroxy propanoate (22e)

[0462] Adapting procedures or variations thereof according to Bhat et al., *Bioorg. & Med. Chem. Lett.* 1998, 8, 3181-3186, a 100 mL round-bottomed flask equipped with a

magnetic stirring bar was charged with 200 mg (0.63 mmol) of ethyl 2,2-diethoxy-3-(1,1,2,2-tetramethyl-1-silapropoxy)propanoate (22d). Fifteen (15) mL of a mixture of ethanol (EtOH) and concentrated hydrochloric acid (99:1 v/v) were added and the reaction mixture was stirred overnight at room temperature. The solvents were evaporated under reduced pressure using a rotary evaporator and a high vacuum pump to provide 120 mg (93% yield) of the title compound (22e) as a colorless oil that was of sufficient purity to be used directly in the next step without further purification and characterization.

Step F: Ethyl 3-[[3-(acetylamino)propyl]sulfonyloxy]-2,2-diethoxypropanoate (22)

[0463] Following the general procedure for the synthesis of acamprostate neopentyl sulfonyl ester prodrugs of Description 3 (Method A), N-[3-(chlorosulfonyl)propyl]acetamide (1) (ca. 350 mg, 1.75 mmol) dissolved in 10 mL of dichloromethane was reacted with ethyl 2,2-diethoxy-3-hydroxypropanoate (22e) (200 mg, 0.969 mmol) in the presence of 243 μL of triethylamine (176 mg, 1.74 mmol) and 211 mg (1.73 mmol) of DMAP. After mass-guided preparative HPLC purification, 10 mg (3% yield) of the title compound (22) was obtained as a colorless oil. ^1H NMR (400 MHz, $\text{MeOH}-d_4$): $\delta=1.23$ (t, $J=7.2$ Hz, 6H), 1.33 (t, $J=7.6$ Hz, 3H), 1.94-2.01 (m, 5H), 3.25-3.31 (m, 4H), 3.49-3.57 (m, 2H), 3.60-3.69 (m, 2H), 4.28 (q, $J=7.2$ Hz, 2H), 4.42 (s, 2H) ppm. MS (ESI) m/z 370.15 ($\text{M}+\text{H}$) $^+$; 392.14 ($\text{M}+\text{Na}$) $^+$.

Example 23

Benzyl 3-[[3-(acetylamino)propyl]sulfonyloxy]-2,2-diethoxypropanoate (23)

Step A: rac-Benzyl glycerate (23a)

[0464] Adapting a procedure according to Choi et al., *Bioorg. Med. Chem.* 1996, 4, 2105-2114, a 1,000 mL round-bottomed flask equipped with a magnetic stirring bar was charged with 16.0 g (101.7 mmol) of potassium permanganate (KMnO_4). The oxidant was dissolved in 150 mL of water and 300 mL of acetone and the reaction mixture was cooled to ca. -78° C. using a dry ice/acetone bath. 15.0 g (92.5 mmol) of benzyl acrylate was slowly added while stirring at -78° C., and the reaction mixture then slowly warmed to 0° C. The inorganic salts were removed by filtration and the filter residue washed with 150 mL of acetone. The combined filtrates were concentrated under reduced pressure at temperatures below 40° C. using a rotary evaporator. The product was extracted with ethyl acetate (3x200 mL) and the combined extracts were dried over anhydrous sodium sulfate (Na_2SO_4). After filtration, the solvent was removed under reduced pressure using a rotary evaporator. The crude material was purified by silica gel chromatography using a mixture of ethyl acetate (EtOAc) and hexane (Hxn) as eluent (EtOAc/Hxn=1:4) to provide 6.0 g (33% yield) of the title compound (23a) as a clear, colorless oil. $R_f=0.50$ ($\text{CHCl}_3/\text{Acetone}=9:1$). ^1H NMR (CDCl_3 , 400 MHz): $\delta=2.61$ -2.83 (br. m, 2H), 3.87 (dd, $J=12.0$, 3.6 Hz, 1H), 3.92 (dd, $J=11.6$, 3.2 Hz, 1H), 3.12 (br. t, $J=3.6$ Hz, 1H), 5.24 (d, $J=12.4$ Hz, 1H), 5.27 (d, $J=12.4$ Hz, 1H), 7.32-7.40 (m, 5H) ppm. MS (ESI) m/z 197.06 ($\text{M}+\text{H}$) $^+$. The analytical data was consistent with that given in Shin, et al., *J. Org. Chem.*, 2000, 65, 7667-7675.

Step B: rac-Benzyl 2-hydroxy-3-(1,1,2,2-tetramethyl-1-silapropoxy)propanoate (23b)

[0465] Adapting procedures or variations thereof according to Shin et al., *J. Org. Chem.* 2000, 65, 7667-7675 and Parkkari et al., *Bioorganic Med. Chem. Lett.* 2004, 14, 3231-3234, a 250 mL round-bottomed flask equipped with a magnetic stirring bar was charged with 2.0 g (10.2 mmol) of rac-benzyl glycerate (23a) and the material dissolved in 40 mL of anhydrous dichloromethane (DCM). Imidazole (0.69 g, 10.2 mmol), 10.2 mL of a one molar (1.0 M) solution of tert-butyl dimethylchlorosilane in DCM, and 0.50 g, (4.1 mmol) of 4-(N,N-dimethylamino)pyridine (DMAP) were added. The reaction mixture was stirred overnight at room temperature. The reaction mixture was then diluted with 100 mL of DCM and washed twice with water and brine. The combined organic extracts were dried over anhydrous magnesium sulfate (MgSO₄). After filtration, the solvent was removed under reduced pressure and purified by silica gel chromatography using a mixture of ethyl acetate (EtOAc) and hexane (Hxn) as eluent (EtOAc/Hxn=95:5) to provide 2.0 g (75% yield) of the title compound (23b) as a clear oil. *R*_f=0.60 (EtOAc/Hxn=1:4). ¹H NMR (400 MHz, CDCl₃): δ=0.03 (s, 3H), 0.06 (s, 3H), 0.87 (s, 9H), 3.06 (d, J=7.6 Hz, 1H), 3.87 (dd, J=10.4, 3.2 Hz, 1H), 3.97 (dd, J=10.4, 3.2 Hz, 1H), 4.26 (br. dt, J=8.4, 3.2 Hz, 1H), 5.22 (s, 2H), 7.33-7.38 (m, 5H), ppm. MS (ESI) *m/z*=311.19 (M+H)⁺. The analytical data was consistent with that given in Shin, et al., *J. Org. Chem.*, 2000, 65, 7667-7675; and Carreiro, et al., *ES* 2006959 (1989).

Step C: Benzyl 2-oxo-3-(1,1,2,2-tetramethyl-1-silapropoxy)propanoate (23c)

[0466] Adapting procedures or variations thereof according to Dess et al., *J. Am. Chem. Soc.* 1991, 113, 7277-7287, to a stirred solution of benzyl 2-oxo-3-(1,1,2,2-tetramethyl-1-silapropoxy)propanoate (23b), 1.8 g (5.8 mmol) of 1,1,1-tris(acetyloxy)-1,1-dihydro-1,2-benziodoxol-3-(1H)-one (Dess-Martin periodinane) in 10 mL of anhydrous dichloromethane was added at ca. 0° C. (ice bath) 3.0 g (7.0 mmol). The reaction mixture was stirred for six hours at room temperature. The solvent was then removed under reduced pressure using a rotary evaporator. The residue was diluted with 100 mL of diethyl ether and washed twice with a saturated sodium bicarbonate (NaHCO₃) solution and subsequently with a 10 wt-% aqueous solution of sodium thiosulfate (Na₂S₂O₃). The combined organic extracts were dried over anhydrous magnesium sulfate (MgSO₄). The solvents were removed under reduced pressure using a rotary evaporator to provide 1.7 g (95% yield) of the title compound (23c) as a colorless oil. The material was of sufficient purity to be used in the next step without further purification or isolation. *R*_f=0.50 (EtOAc/Hxn=1:4). ¹H NMR (400 MHz, CDCl₃): δ=0.10 (s, 6H), 0.92 (s, 9H), 4.74 (s, 2H), 5.29 (s, 2H), 7.36-7.39 (m, 5H) ppm. MS (ESI) *m/z* 307.12 (M+H)⁺.

Step D: Benzyl 2,2-diethoxy-3-(1,1,2,2-tetramethyl-1-silapropoxy)propanoate (23d)

[0467] A catalytic amount (ca. 2.1 mL) of concentrated sulfuric acid (H₂SO₄) was added to a solution of benzyl 2-oxo-3-(1,1,2,2-tetramethyl-1-silapropoxy)propanoate (23c), 1.7 g, (5.5 mmol) in ca. 90 mL (80.2 g, 0.54 mol) of triethylorthoformate while stirring at ca. 0° C. (ice bath). The reaction mixture was stirred at 0° C.-10° C. for three to four hours. The reaction mixture was then diluted with dichloromethane (DCM) and washed with water and brine. The

organic extract was dried over anhydrous magnesium sulfate (MgSO₄), filtered, and the solvent removed under reduced pressure using a rotary evaporator to provide 2.0 g (95% yield) of the title compound (23d) as a colorless oil that was of sufficient purity to be used in the next step without further purification. *R*_f=0.60 (EtOAc/Hxn=1:4). ¹H NMR (400 MHz, CDCl₃): δ=0.003 (s, 6H), 0.85 (s, 9H), 1.23 (t, J=7.2 Hz, 6H), 3.40-3.64 (m, 4H), 3.84 (s, 2H), 5.26 (s, 2H), 7.30-7.40 (m, 5H) ppm. MS (ESI) *m/z* 383.19 (M+H)⁺, 405.15 (M+Na)⁺.

Step E: Benzyl 2,2-diethoxy-3-hydroxypropanoate (23e)

[0468] Adapting a procedure or a variation thereof according to Bhat et al., *Bioorg. Med. Chem. Lett.* 1998, 8, 3181-3186, a 100 mL round-bottomed flask equipped with a magnetic stirring bar was charged with 2.0 g (5.23 mmol) of benzyl 2,2-diethoxy-3-(1,1,2,2-tetramethyl-1-silapropoxy)propanoate (23d). Fifteen (15) mL of a mixture of ethanol (EtOH) and concentrated hydrochloric acid (95:5 v/v) was added and the reaction mixture stirred overnight at room temperature. The solvents were evaporated under reduced pressure using a rotary evaporator and a high vacuum pump to provide 1.4 g (99% yield) of the title compound (23e) as a colorless oil that was of sufficient purity to be used directly in the next step without further purification or isolation. MS (ESI) *m/z*: 291.1(M+Na)⁺.

Step F: Benzyl 3-{[3-(acetylamino)propyl]sulfonyloxy}-2,2-diethoxypropanoate (23)

[0469] Following the general procedure for the synthesis of acamprostate neopentyl sulfonylester prodrugs of Description 3 (Method A), N-[3-(chlorosulfonyl)propyl]acetamide (1) (ca. 225 mg, 1.11 mmol) dissolved in 10 mL of dichloromethane was reacted with benzyl 2,2-diethoxy-3-hydroxypropanoate (23e) (250 mg, 0.932 mmol) in the presence of 155 μL of triethylamine (113 mg, 1.12 mmol) and 137 mg (1.12 mmol) of DMAP. After mass-guided preparative HPLC purification, 32 mg (8% yield) of the title compound (23) was obtained as a colorless oil. ¹H NMR (400 MHz, DMSO-d₆): δ=1.23 (t, J=7.2 Hz, 6H), 1.70-1.78 (m, 2H), 1.80 (s, 3H), 3.07-3.13 (m, 2H), 3.31-3.37 (m, 2H), 3.37-3.46 (m, 2H), 3.50-3.59 (m, 2H), 4.38 (s, 2H), 5.21 (s, 2H), 7.31-7.40 (m, 5H), 7.89 (br. t, J=5.2 Hz, 1H) ppm. MS (ESI) *m/z* 432.1 (M+H)⁺; 454.1 (M+Na)⁺.

Example 24

Methyl 4-{[3-(acetylamino)propyl]sulfonyloxy}-3,3-dimethylbutanoate (24)

Step A: 2,2-Dimethyl-pent-4-en-1-ol (24a)

[0470] Adapting a procedure or a variation thereof according to Chen et al., *J. Am. Chem. Soc.* 2003, 125, 6697-6704, an oven dried 500 mL round-bottomed flask equipped with a magnetic stirring bar and a pressure equalized addition funnel was charged under a nitrogen atmosphere with 5.00 g (39.0 mmol) of commercially available 2,2-dimethyl-4-pentenoic acid. The acid was dissolved in 100 mL of anhydrous tetrahydrofuran (THF) and cooled to ca. 0° C. (ice bath). Forty (40) mL of a one molar (1.0 M) solution of lithium aluminum hydride (LAH) in THF was slowly added at this temperature and the reaction mixture then stirred overnight with slow warming to room temperature followed by subsequent cool-

ing to ca. 0° C. (ice bath). Subsequent and careful addition of 2.6 mL of water, 5.2 mL of a 10 wt-% aqueous solution of sodium hydroxide (NaOH), and 2.6 mL of water resulted in a colorless precipitate that was filtered off. The filter residue was washed with ethyl acetate (EtOAc) and the combined filtrates were dried over anhydrous magnesium sulfate (MgSO₄) and filtered. Solvents were evaporated under reduced pressure using a rotary evaporator to provide 4.05 g (91% yield) of the title compound (24a) as a colorless oil. R_f =0.30 (EtOAc/H₂O=1:9). ¹H NMR (400 MHz, CHCl₃): δ=0.91 (s, 6H), 2.04 (d, J=8.0 Hz, 2H), 3.34 (s, 2H), 5.01-5.10 (m, 2H), 5.79-5.91 (m, 1H) ppm. MS (ESI) m/z 114.86 (M+H)⁺. The analytical data for the compound was consistent with the structure and the data reported in the literature.

[0471] Alternatively, the title compound (24a) was prepared in comparable yield starting from the corresponding alkyl esters of acid employing comparable synthetic methods, or from commercially available 2,2-dimethyl-4-pentenal through reduction of the aldehyde functionality with sodium borohydride (NaBH₄) in ethanol (EtOH) at room temperature.

Step B: 2,2-Dimethylpent-4-enyl [3-(acetylamino)propyl]sulfonate (24b)

[0472] Following the general procedure for the synthesis of acamprostate neopentyl sulfonylester prodrugs of Description 3 (Method A), N-[3-(chlorosulfonyl)propyl]acetamide (1) (2.6 g, 13 mmol) dissolved in 50 mL of dichloromethane was reacted with 2,2-dimethylpent-4-en-1-ol (24a) (1.0 g, 8.8 mmol) in the presence of 1.9 mL of triethylamine (1.4 g, 14.0 mmol) and 170 mg (1.4 mmol) of DMAP. After aqueous work-up, the residue was purified by silica gel chromatography using ethyl acetate (EtOAc) as eluent to provide 1.4 g (57% yield) of the title compound (24b) as a colorless oil. R_f =0.33 (EtOAc). ¹H NMR (400 MHz, CDCl₃): δ=0.98 (s, 6H), 2.01 (s, 3H), 2.06-2.12 (m, 4H), 3.15-3.19 (m, 2H), 3.43 (dd, J=12.8, 6.8 Hz, 2H), 3.90 (s, 2H), 5.04-5.12 (m, 2H), 5.72-5.81 (m, 2H), 5.87 (br. m, 1H) ppm. MS (ESI) m/z 278.12 (M+H)⁺.

Step C: 2,2-Dimethyl-4-oxobutyl [3-(acetylamino)propyl]sulfonate (24c)

[0473] Adapting a procedure or a variation thereof according to Pappo et al., *J. Org. Chem.* 1956, 21, 478-479, a stirred solution of 2,2-dimethylpent-4-enyl [3-(acetylamino)propyl] sulfonate (24b) (0.1 g, 0.36 mmol) in 3 mL of a mixture of water and tetrahydrofuran (THF) (1:1 v/v) was reacted with 0.36 mL of a 2.5 wt-% solution of osmium tetroxide (OsO₄) in tert-butanol and in the presence of 0.15 g (0.72 mmol) of sodium metaperiodate (NaIO₄). The reaction was monitored by analytical LC/MS. After the starting material was completely consumed, the reaction was quenched by adding a 10 wt-% aqueous sodium hydrogensulfite solution (NaHSO₃). The reaction mixture was then extracted twice with ethyl acetate. The combined organic extracts were subsequently washed with a 10 wt-% aqueous NaHSO₃ solution and brine, dried over magnesium sulfate (MgSO₄), filtered, and the solvents removed under reduced pressure using a rotary evaporator to provide ca. 0.1 g (95% yield) of the title compound (24c) as a colorless oil. The resulting crude material was of sufficient purity for use in the next step without further purification. R_f =0.15 (EtOAc). ¹H NMR (400 MHz, CDCl₃): δ=1.14 (s, 6H), 2.00 (s, 3H), 2.02-2.09 (m, 2H), 2.45 (d, J=2.0

Hz, 2H), 3.15-3.19 (m, 2H), 3.37-3.42 (m, 2H), 4.03 (s, 2H), 5.98-6.05 (br. m, 1H), 9.79 (t, J=2.0 Hz, 1H) ppm. MS (ESI) m/z 280.08 (M+H)⁺.

Step D: Methyl 4-{[3-(acetylamino)propyl]sulfonyloxy}-3,3-dimethylbutanoate (24)

[0474] Adapting a procedure or a variation thereof according to McDonald et al., *J. Org. Chem.* 1989, 54, 1213-1215, to a stirred solution of 2,2-dimethyl-5-oxobutyl [3-(acetylamino)propyl]sulfonate (24c) (0.2 g, 0.68 mmol) and methanol (0.23 mL, 0.24 g, 2.2 mmol) in 10 mL of acetonitrile were added 0.31 g (1.4 mmol) N-iodosuccinimide (NIS) and potassium carbonate (K₂CO₃) (0.28 g, 2.0 mmol). The reaction mixture was stirred overnight at room temperature. The reaction mixture was then diluted with water and extracted twice with ethyl acetate. The combined organic extracts were washed with a 10 wt-% aqueous sodium thiosulfate solution (Na₂S₂O₃) and brine, dried over magnesium sulfate (MgSO₄), filtered, and the solvents removed under reduced pressure using a rotary evaporator. The residue was purified by silica gel chromatography using a mixture of ethyl acetate (EtOAc) and methanol (MeOH) as eluent (EtOAc/MeOH=4:1) to provide 13 mg (12% yield) of the title compound (24) as a colorless oil. ¹H NMR (400 MHz, CHCl₃): δ=1.07 (s, 6H), 2.00 (s, 3H), 2.03-2.10 (m, 2H), 2.34 (s, 2H), 3.17-3.21 (m, 2H), 3.37-3.43 (m, 2H), 3.67 (s, 3H), 4.06 (s, 2H), 6.10-6.20 (br. m, 1H) ppm. MS (ESI) m/z 310.15 (M+H)⁺.

Example 25

Phenylmethyl 4-{[3-(acetylamino)propyl]sulfonyloxy}-3,3-dimethylbutanoate (25)

[0475] Adapting a procedure or a variation thereof according to McDonald et al., *J. Org. Chem.* 1989, 54, 1213-1215, N-iodosuccinimide (NIS) (0.63 g, 2.8 mmol) and potassium carbonate K₂CO₃ (0.39 g, 2.8 mmol) were added to a solution of 2,2-dimethyl-4-oxobutyl [3-(acetylamino)propyl]sulfonate (24c) (0.26 g, 0.93 mmol) and benzyl alcohol (0.29 mL, 0.30 g, 2.8 mmol) in 5 mL acetonitrile while stirring. The mixture was stirred overnight at room temperature. The reaction mixture was then diluted with water and extracted twice with ethyl acetate. The combined organic extracts were washed with a 10 wt-% aqueous sodium thiosulfate solution (Na₂S₂O₃) and brine, dried over magnesium sulfate (MgSO₄), filtered, and the solvents removed under reduced pressure using a rotary evaporator. The residue was purified by mass-guided preparative HPLC to provide 17 mg (5% yield) of the title compound (25) as a colorless oil. ¹H NMR (400 MHz, CHCl₃): δ=1.08 (s, 6H), 2.00 (s, 3H), 2.02-2.09 (m, 2H), 2.40 (s, 2H), 3.12-3.17 (m, 2H), 3.38 (q, J=6.4 Hz, 2H), 4.07 (s, 2H), 5.12 (s, 2H), 5.80-5.88 (br. m, 1H), 7.33-7.40 (m, 5H) ppm. MS (ESI) m/z 386.02 (M+H)⁺; m/z 408.03 (M+Na)⁺.

Alternative Synthesis of Phenylmethyl 4-{[3-(acetylamino)propyl]sulfonyloxy}-3,3-dimethylbutanoate (25)

Step A: 2,2-Dimethyl-4-oxobutyl [3-(acetylamino)propyl]sulfonate (25a)

[0476] A 100 mL round bottomed flask equipped with a magnetic stirring bar and a perforated polyethylene stopper with a 14 gauge-needle connected via silicon tubing to a Welsbach Standard T-Series ozone generator was charged

with 550 mg (1.99 mmol) of 2,2-dimethylpent-4-enyl [3-(acetylamino)propyl]sulfonate (24b) and dissolved in 15 mL of anhydrous dichloromethane. The solution was cooled to ca. -78°C . (dry ice/acetone) and an ozone/oxygen gas mixture was purged through the solution until an intensely blue color indicated an excess of dissolved ozone in the reaction mixture. Excess ozone was removed by successive purges with oxygen and nitrogen gas. Seven-hundred thirty-four (734) μL (621 mg, 10.0 mmol) of dimethyl sulfide was added and the reaction mixture stirred overnight with gradual warming to room temperature. The solvents were then removed under reduced pressure using a rotary evaporator. The residue was purified by silica gel column chromatography using mixtures of ethyl acetate (EtOAc) and methanol (MeOH) as eluent (100% EtOAc $\rightarrow$ EtOAc/2.5 vol-% MeOH $\rightarrow$ EtOAc/5 vol-% MeOH $\rightarrow$ EtOAc/10 vol-% MeOH) to provide 385 mg (69% yield) of the title compound (25a) as a colorless oil. $R_f=0.40$ (EtOAc/MeOH=95:5). The analytical data for the compound was consistent with the proposed structure and data obtained from material prepared using the $\text{OsO}_4/\text{NaIO}_4$ -route.

Step B: 4-{[3-(Acetylamino)propyl]sulfonyloxy}-3,3-dimethylbutanoic Acid (25b)

[0477] Adapting a procedure or a variation thereof according to Fillmore et al., *Encyclopedia of Reagents for Organic Synthesis*, John Wiley & Sons, Ltd., 2001, Jones reagent (chromic acid) was freshly prepared prior to use from 2.00 g (20.0 mmol) of chromium trioxide and 1.333 mL (2.45 g, 25.0 mmol) of concentrated sulfuric acid (H_2SO_4). The chemicals were carefully mixed at 0°C . and the resulting mixture was diluted at the same temperature with water to a total volume of 10 mL to provide an aqueous chromic acid preparation of approximately 2.0 M. The reagent was used without further purification or isolation in the next step.

[0478] Adapting a procedure or variation thereof according to Petrini, et al., *Tetrahedron*, 1986, 42, 151-154, a 4 mL vial equipped with a magnetic stirring bar and screw cap was charged with 55 mg (0.197 mmol) of 2,2-dimethyl-4-oxobutyl [3-(acetylamino)propyl]sulfonate (26a) and 1.6 mL of acetone. To the cooled reaction mixture (ca. 0°C ., ice bath), 111 μL (ca. 0.222 mmol) of the freshly prepared Jones-reagent was added and the reaction mixture stirred for two hours with gradual warming to room temperature. Excess Jones-reagent was consumed by adding 100 μL of isopropanol (i-PrOH) and stirring for 30 minutes at room temperature. The reaction mixture was then diluted with water (pH<1) and extracted with ethyl acetate (EtOAc). The combined organic extracts were washed with brine, dried over magnesium sulfate (MgSO_4), filtered, and evaporated under reduced pressure using a rotary evaporator to provide ca. 50 mg (86% yield) of the title compound (25b) as a colorless oil that was of sufficient purity to be used in the next step without further purification or isolation. ^1H NMR (400 MHz, CDCl_3): $\delta=1.13$ (s, 6H), 2.00-2.12 (m, 5H), 2.38 (s, 2H), 3.14-3.22 (m, 2H), 3.36-3.44 (m, 2H), 4.12 (s, 2H), 6.14-6.22 (m, 1H) ppm. MS (ESI) m/z 296.01 (M+H) $^{+}$; m/z 317.95 (M+Na) $^{+}$; m/z 293.98 (M-H) $^{-}$.

[0479] Alternatively, the title compound (25b) was prepared under mild conditions starting from the unsaturated precursor compound 2,2-dimethylpent-4-enyl [3-(acetylamino)propyl]sulfonate (24c) according to a method described by Henry et al., *J. Org. Chem.* 1993, 58, 4745. In one synthesis, 100 mg (0.36 mmol) of 2,2-dimethylpent-4-

enyl [3-(acetylamino)propyl]sulfonate (24c) dissolved in 3 mL of acetone was reacted with 360 μL (0.04 mmol, ca. 10 mol-%) of a 2.5 wt-% solution of osmiumtetroxide (OsO_4) in tert-butanol (t-BuOH) and 650 μL (ca. 1.30 mmol) of freshly prepared Jones-reagent (ca. 2 M). The reaction mixture was stirred overnight with gradual warming to room temperature. Excess Jones-reagent was then consumed by adding 100 μL of isopropanol (i-PrOH) and stirring for 30 minutes at room temperature followed by the addition of 250 μL of an aqueous 10 wt-% solution of sodium hydrogensulfite (NaHSO_3). The reaction mixture was diluted with water (pH<1) and extracted with dichloromethane (DCM). The combined organic extracts were washed with an aqueous one molar solution of hydrogen chloride (HCl) and brine, dried over magnesium sulfate (MgSO_4), filtered, and evaporated under reduced pressure using a rotary evaporator to provide 100 mg (95% yield) of the title compound (25b) as a colorless oil that was of sufficient purity to be used in the next step without further purification or isolation. The analytical data for the compound was consistent with the proposed structure and the data obtained from material prepared using the sequential route.

Step C: 3-(Chlorocarbonyl)-2,2-dimethylpropyl[3-(acetylamino)propyl]sulfonate (25c)

[0480] An oven dried 100 mL round bottomed flask equipped with a magnetic stirring bar and a rubber septum was charged with ca. 50 mg (0.17 mmol) of 4-{[3-(acetylamino)propyl]sulfonyloxy}-3,3-dimethylbutanoic acid (25b). The material was dissolved in 5 mL of anhydrous dichloromethane (DCM) and two small drops of anhydrous N,N-dimethylformamide (DMF). The solution was cooled to ca. 0°C . (ice bath) and 150 μL (0.3 mmol) of a two molar solution (2 M) of oxalyl chloride in DCM was added. The reaction mixture was stirred with gradual warming to room temperature for three hours and the solvents then removed under reduced pressure on a rotary evaporator to provide the title compound (25c) as a clear, oily semisolid that was used directly in the next step without further purification or characterization.

Step D: Phenylmethyl 4-{[3-(acetylamino)propyl]sulfonyl-oxy}-3,3-dimethylbutanoate (25)

[0481] An oven dried 100 mL round bottomed flask equipped with a magnetic stirring bar and a rubber septum was charged with freshly prepared acid chloride 3-(chlorocarbonyl)-2,2-dimethylpropyl [3-(acetylamino)propyl]sulfonate (25c). The material was dissolved in 3 mL of anhydrous dichloromethane (DCM). One-hundred three (103) μL (108 mg, 1.0 mmol) of anhydrous benzyl alcohol and 61 mg (0.5 mmol) of 4-(N,N-dimethylamino)-pyridine and 40 μL (40 mg, 0.5 mmol) of anhydrous pyridine were added and the reaction mixture stirred overnight at room temperature. The solvents were then removed under reduced pressure using a rotary evaporator and the residue diluted with ethyl acetate (EtOAc). The solution was washed with a one molar aqueous solution of hydrogen chloride (HCl), a saturated aqueous solution of sodium hydrogencarbonate (NaHCO_3) and brine, dried over anhydrous magnesium sulfate (MgSO_4), filtered, and the solvent removed under reduced pressure using a rotary evaporator. The residue was purified by silica gel column chromatography using mixtures of EtOAc and methanol (MeOH) as eluent (100% EtOAc $\rightarrow$ EtOAc/MeOH=95:5) to provide 36 mg (47% yield over three steps) of the title com-

pound (25) as a colorless oil. The analytical data for the compound was consistent with the proposed structure and the data obtained from material prepared using the two step route, e.g., (i) $\text{OsO}_4/\text{NaIO}_4$ and (ii) NIS/ROH .

Example 26

2,2-Dimethyl-5-oxopentyl [3-(acetylamino)propyl] sulfonate (26)

Step A: 2,2-Dimethylhex-5-en-1-ol (26a)

[0482] Adapting procedures or a variations thereof according to Juaristi et al., *J. Org. Chem.* 1991, 56, 1623-1630; Ashby et al., *J. Org. Chem.* 1984, 49, 3545-3556; or Beckwith et al., *J. Chem. Soc., Perkin Trans. 2: Phys. Org. Chem.* 1979, 1535-1539, an oven-dried 100 mL round-bottomed flask equipped with a magnetic stirring bar and a pressure equalized addition funnel was charged under a nitrogen atmosphere with 1.04 g (7.33 mmol) of commercially available 2,2-dimethyl-5-hexenoic acid. The acid was dissolved in 20 mL of anhydrous tetrahydrofuran (THF) and cooled to ca. 0° C. (ice bath). Eight (8.0) mL of a one molar solution (1.0 M) of lithium aluminum hydride (LAH) in THF was slowly added and the reaction mixture stirred for three hours with slow warming to room temperature followed by cooling to ca. 0° C. (ice bath). Careful addition of 0.49 mL of water, 0.98 mL of a 10 wt-% aqueous solution of sodium hydroxide (NaOH), and 0.49 mL of water resulted in a colorless precipitate that was filtered off. The filter residue was washed with ethyl acetate (EtOAc) and the combined filtrates were dried over anhydrous magnesium sulfate (MgSO_4), filtered, and the solvents evaporated under reduced pressure using a rotary evaporator. The residue was purified by silica gel chromatography using a mixture of ethyl acetate (EtOAc) and hexane (Hxn) as eluent (EtOAc/Hxn=1:3) to provide 548 mg (58% yield) of the title compound (26a) as a colorless oil. R_f =0.42 (EtOAc/Hxn=1:4). ^1H NMR (400 MHz, CHCl_3): δ =0.91 (s, 6H), 1.30-1.40 (m, 2H), 2.00-2.08 (m, 2H), 3.35 (s, 2H), 4.91-5.06 (m, 2H), 5.78-5.90 (m, 1H) ppm. MS (ESI) m/z 129.12 ($\text{M}+\text{H}^+$). The analytical data for the compound was consistent with the proposed structure and the data reported in the literature.

Step B: 2,2-Dimethylhex-5-enyl [3-(acetylamino)propyl]sulfonate (26b)

[0483] Following the general procedure for the synthesis of acamprostate neopentyl sulfonylester prodrugs or intermediates of Description 3 (Method A), N-[3-(chlorosulfonyl)propyl]acetamide (1) (1.7 g, 8.6 mmol), dissolved in 50 mL of dichloromethane, was reacted with 2,2-dimethylhex-5-en-1-ol (26a) (0.55 g, 4.3 mmol) in the presence of 1.2 mL of triethylamine (0.87 g, 8.6 mmol) and 100 mg (0.8 mmol) of DMAP. After aqueous work-up, the residue was purified by silica gel chromatography using a mixture of ethyl acetate (EtOAc) and methanol (MeOH) as eluent (EtOAc/MeOH=4:1) to provide 500 mg (40% yield) of the title compound (26b) as a colorless oil. ^1H NMR (400 MHz, CHCl_3): δ =0.98 (s, 6H), 1.38-1.41 (m, 2H), 2.06 (s, 3H), 2.06-2.12 (m, 4H), 3.15-3.19 (m, 2H), 3.43 (dd, J =12.8, 6.4 Hz, 2H), 3.92 (s, 2H), 4.94-5.05 (m, 2H), 5.75-5.85 (m, 1H) ppm. MS (ESI) m/z 292.02 ($\text{M}+\text{H}^+$).

Step C: 2,2-Dimethyl-5-oxopentyl [3-(acetylamino)propyl]sulfonate (26)

[0484] Adapting a procedure or a variation thereof according to Pappo et al., *J. Org. Chem.* 1956, 21, 478-479, a stirred

solution of 2,2-dimethylhex-5-enyl [3-(acetylamino)propyl] sulfonate (26b) (0.5 g, 1.7 mmol) in 15 mL of a mixture of water and tetrahydrofuran (THF) (1:1 v/v) was reacted with 2.0 mL of a 2.5 wt-% solution of osmiumtetroxide (OsO_4) in tert-butanol in the presence of 0.75 g (3.45 mmol) of sodium metaperiodate (NaIO_4). The reaction was monitored using analytical LC/MS. After the starting material was completely consumed, the reaction was quenched by adding a 10 wt-% aqueous sodium hydrogensulfite solution (NaHSO_3). The reaction mixture was then extracted twice with ethyl acetate. The combined organic extracts were subsequently washed with a 10 wt-% aqueous NaHSO_3 solution and brine, dried over magnesium sulfate (MgSO_4), filtered, and the solvents removed under reduced pressure using a rotary evaporator. The residue was purified by silica gel chromatography using a mixture of ethyl acetate (EtOAc) and methanol (MeOH) (EtOAc/MeOH=4:1) as eluent to provide 200 mg (45% yield) of the title compound (26) as a colorless oil. R_f =0.66 (EtOAc/MeOH=4:1). ^1H NMR (400 MHz, CHCl_3): δ =0.96 (s, 6H), 1.62-1.66 (m, 2H), 2.03 (s, 3H), 2.01-2.08 (m, 2H), 2.43-2.47 (m, 2H), 3.14-3.18 (m, 2H), 3.35-3.40 (m, 2H), 3.88 (s, 2H), 6.24-6.28 (br. m, 1H), 9.75 (t, J =2.0 Hz, 1 H) ppm. MS (ESI) m/z 293.98 ($\text{M}+\text{H}^+$).

Alternative Synthesis of 2,2-Dimethyl-5-oxopentyl [3-(acetylamino)propyl]sulfonate (26)

[0485] In one qualitative experiment, a 100 mL round bottomed flask equipped with a magnetic stirring bar and a perforated polyethylene stopper with a 14 gauge-needle connected via silicon tubing to a Welsbach Standard T-Series ozone generator was charged with 2,2-dimethylhex-5-enyl [3-(acetylamino)propyl]sulfonate (26b) and dissolved in anhydrous dichloromethane (DCM). The solution was cooled to ca. -78° C. (dry ice/acetone) and an ozone/oxygen gas mixture was purged through the solution until an intensely blue color indicated an excess of dissolved ozone in the reaction mixture. Excess ozone was removed by successive purges with oxygen and nitrogen gas. An excess of dimethyl sulfide was added and the reaction mixture was stirred overnight with gradual warming to room temperature. The solvents were then removed under reduced pressure using a rotary evaporator. The residue was purified by silica gel column chromatography using mixtures of ethyl acetate (EtOAc) and methanol (MeOH) as eluent to provide the title compound (26) as a colorless oil. R_f =0.40 (EtOAc/MeOH=95:5). The analytical data for the compound are was consistent with the proposed structure and the data obtained from material prepared using the $\text{OsO}_4/\text{NaIO}_4$ -route.

Example 27

Phenylmethyl 5-([3-(acetylamino)propyl]sulfonyloxy)-4,4-dimethylpentanoate (27)

[0486] Adapting a procedure or a variation thereof according to McDonald, et al., *J. Org. Chem.* 1989, 54, 1213-1215, to a stirred solution of 2,2-dimethyl-5-oxopentyl [3-(acetylamino)propyl]sulfonate (26) (0.2 g, 0.68 mmol) and benzyl alcohol (0.23 mL, 0.24 g, 2.2 mmol) in 10 mL of acetonitrile were added 0.31 g (1.4 mmol) N-iodosuccinimide (NIS) and potassium carbonate (K_2CO_3) (0.28 g, 2.0 mmol). The reaction mixture was stirred overnight at room temperature. The reaction mixture was diluted with water and extracted twice with ethyl acetate. The combined organic extracts were washed with a 10 wt-% aqueous sodium thiosulfate solution

($\text{Na}_2\text{S}_2\text{O}_3$) and brine, dried over magnesium sulfate (MgSO_4), filtered, and the solvents removed under reduced pressure using a rotary evaporator. The residue was purified by mass-guided preparative HPLC to provide 43 mg (16% yield) of the title compound (27) as a colorless oil. ^1H NMR (400 MHz, CDCl_3): δ =0.97 (s, 6H), 1.68-1.72 (m, 2H), 1.99 (s, 3H), 2.03-2.10 (m, 2H), 2.35-2.39 (m, 2H), 3.15-3.18 (m, 2H), 3.40 (dd, J =12.8, 6.4 Hz, 2H), 3.90 (s, 2H), 5.12 (s, 2H), 5.98-6.02 (br. m, 1H), 7.32-7.37 (m, 5H) ppm. MS (ESI) m/z 400.06 ($\text{M}+\text{H}$) $^+$.

Example 28

5-[[3-(Acetylamino)propyl]sulfonyloxy]-4,4-dimethylpentanoic Acid (28)

[0487] Caution: Chromium(VI) oxide is a highly toxic cancer suspect agent. All chromium(VI) reagents must be handled with care. The mutagenicity of Cr(VI) compounds, i.e. chromic acid (Jones-Reagent), is well documented. Special care must always be exercised in adding CrO_3 to organic solvents. This reagent must be handled in a fume hood.

[0488] Adapting a procedure or a variation thereof according to Fillmore et al., *Encyclopedia of Reagents for Organic Synthesis*, John Wiley & Sons, Ltd., 2001, Jones reagent (chromic acid) was freshly prepared prior to use from 2.00 g (20.0 mmol) of chromium trioxide and 1.333 mL (2.45 g, 25.0 mmol) of concentrated sulfuric acid (H_2SO_4). The chemicals were carefully mixed at 0° C. and the resulting mixture was carefully diluted at the same temperature with water to total volume of 10 mL to yield an aqueous chromic acid preparation of approximately 2.0 M. The reagent was used in the next step without further purification or isolation.

[0489] Adapting a procedure or variation thereof according to Petrini et al., *Tetrahedron*, 1986, 42, 151-154, a 100 mL round-bottomed flask equipped with a magnetic stirring bar was charged with 2,2-dimethyl-5-oxopentyl [3-(acetylamino)propyl]sulfonate (26) (580 mg, 1.98 mmol). The compound was dissolved in 20 mL of acetone, cooled to ca. 0° C. (ice bath), and 1.19 mL (2.38 mmol) of Jones-reagent (2.0 M in water) was added while stirring. The reaction mixture turned from brown to green in about 30 min. After the starting material was completely consumed, 1 mL of isopropanol (iPrOH) was added to destroy excess oxidant. The reaction mixture was diluted with water and extracted with ethyl acetate. The combined organic extracts were washed with brine, dried over anhydrous magnesium sulfate (MgSO_4), filtered, and the solvents removed under reduced pressure using a rotary evaporator. After purification by mass-guided preparative HPLC, 153 mg (25% yield) of the title compound (28) was obtained as a colorless solid. M.p.: 105.3-124.9° C. ^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ =0.89 (s, 6H), 1.49-1.54 (m, 2H), 1.76-1.83 (m, 5H), 2.17-2.22 (m, 2H), 3.10-3.16 (m, 2H), 3.31-3.35 (m, 2H, superimposed with H_2O signal), 3.88 (s, 2H), 7.90 (br. t, J =5.6 Hz, 1H), 12.06 (br. s, 1H) ppm. MS (ESI) m/z 310.0 ($\text{M}+\text{H}$) $^+$, 332.0 ($\text{M}+\text{Na}$) $^+$, 308.0 ($\text{M}-\text{H}$) $^-$.

Example 29

Bioavailability of Acamprosate Following Administration of Acamprosate Prodrugs to Rats

[0490] Rats were obtained commercially and were precannulated in the jugular vein. Animals were conscious at the time of the experiment. All animals were fasted overnight and until 4 hours post-dosing of a prodrug of Formula (I).

[0491] Rat blood samples (0.3 mL/sample) were collected from all animals prior to dosing and at different time-points up to 24 h post-dose into tubes containing EDTA. Two aliquots (100 μL each) were quenched with 300 μL methanol and stored at -20° C. prior to analysis.

[0492] To prepare analysis standards, 90 μL of rat blood was quenched with 300 μL methanol followed by 10 μL of spiking standard and/or 20 μL of internal standard. The sample tubes were vortexed for at least 2 min and then centrifuged at 3400 rpm for 20 min. The supernatant was then transferred to an injection vial or plate for analysis by LC-MS-MS.

[0493] To prepare samples for analysis, 20 μL of internal standard was added to each quenched sample tube. The sample tubes were vortexed for at least 2 min and then centrifuged at 3400 rpm for 20 min. The supernatant was then transferred to an injection vial or plate for analysis by LC-MS-MS.

[0494] LC-MS-MS analysis was performed using an API 4000 equipped with Agilent 1100 HPLC and a Leap Technologies autosampler. The following HPLC column conditions were used: HPLC column: Thermal-Hypersil-Keystone C18, 4.6 $\times$ 100 mm, 5 μm ; mobile phase A: 0.1% formic acid in water; mobile phase B: 0.1% formic acid in acetonitrile; flow rate: 1.2 mL/min; gradient: 99%A/1%B at 0.0 min; 99%A/1%B at 0.5 min; 5%A/95%B at 1.8 min; 5%A/95%B at 3.5 min; 99%A/1%B at 3.6 min; and 99%A/1%B at 9.0 min. Acamprosate was monitored in negative ion mode. The LOQ was 0.004 $\mu\text{g/mL}$. The standard curve range was 0.004 to 10 $\mu\text{g/mL}$. Prodrug was monitored in positive ion mode. The LOQ and standard curve range was the same as for acamprosate.

[0495] Non-compartmental analysis was performed using WinNonlin software (v.3.1 Professional Version, Pharsight Corporation, Mountain View, Calif.) on individual animal profiles. Summary statistics on major parameter estimates was performed for C_{max} (peak observed concentration following dosing), T_{max} (time to maximum concentration is the time at which the peak concentration was observed), $\text{AUC}_{(0-t)}$ (area under the plasma concentration-time curve from time zero to last collection time, estimated using the log-linear trapezoidal method), $\text{AUC}_{(0-\infty)}$ (area under the plasma concentration time curve from time zero to infinity, estimated using the log-linear trapezoidal method to the last collection time with extrapolation to infinity), and $t_{1/2,z}$ (terminal half-life).

[0496] Acamprosate or acamprosate prodrug was administered by oral gavage to groups of four to six adult male Sprague-Dawley rats (about 250 g). Animals were conscious at the time of the experiment. Acamprosate or acamprosate prodrug was orally or colonically administered in 3.4% Phosal at a dose of 70 mg-equivalents acamprosate per kg body weight.

[0497] The percent relative bioavailability ($F\%$) of acamprosate was determined by comparing the area under the acamprosate concentration vs time curve (AUC) following oral or colonic administration of an acamprosate prodrug or acamprosate with the AUC of the acamprosate concentration vs time curve following intravenous administration of acamprosate on a dose normalized basis. Compounds (9) and (11) exhibited an acamprosate oral bioavailability at least about 5 times greater than the acamprosate oral bioavailability of an equivalent dose of acamprosate itself. Compound (9) exhibited an acamprosate colonic bioavailability at least about 5 times greater than the acamprosate colonic bioavailability of an equivalent dose of acamprosate itself.

Description 4

Use of Clinical Trials to Assess the Efficacy of
Acamprosate Prodrugs for Maintaining Abstinence
from Alcohol

[0498] The efficacy of an acamprosate prodrug for treating alcoholism can be assessed using a randomized, double-blind, double-dummy, placebo-controlled trial. Patients aged 18 to 65 years meeting DSM IV criteria for alcohol dependence and having a history of alcohol dependence for at least 12 months are selected for the study. Patients are required to have undergone detoxification and have had five or more days of abstinence from alcohol before commencing treatment. Patients having a body weight of less than 60 kg receive an equivalent of 1332 mg/day (two 333 mg tablets in the morning and one at midday and in the evening) or placebo, and patients having a bodyweight of greater than 50 kg receive an acamprosate equivalent of 1998 mg/day (two 333 mg tablets in the morning, midday and evening) or placebo. Other acamprosate equivalent doses may be appropriate depending upon the pharmacokinetics of a particular acamprosate prodrug.

[0499] Primary and secondary outcome measures include commonly accepted subjective measures (based mainly on self-reported data) of continuous abstinence rate (CAR, i.e., the percentage of patients completely abstinent throughout the entire treatment and/or follow-up period), cumulative abstinence duration (CAD), the proportion of the total time that CAD represented (CADP, i.e. CAD as a proportion of the total treatment duration) and/or time to first drink (TFD). Surrogate biological markers of relapse such as γ -glutamyl transferase, carbohydrate-deficient transferrin, AST and ALT levels, and mean corpuscular volume can also be determined. Efficacy of acamprosate prodrugs in the maintenance of abstinence in patients with alcohol dependence is reflected in an increased CAR, CADP, and TFD compared to patients receiving placebo.

Description 5

Use of Animal Models to Assess the Efficacy of
Acamprosate Prodrugs for Treating Alcohol With-
drawal

[0500] Withdrawal Seizure-Prone (WSP) and Withdrawal Seizure-Resistant (WSR) mice are used to assess the efficacy of acamprosate prodrugs for treating alcohol withdrawal. Mice are made dependent on ethanol via 72 h of chronic ethanol vapor inhalation. On day 1, mice are weighted, injected with a loading dose of ethanol and pyrazole HCl (Pyr), an alcohol dehydrogenase inhibitor, and placed into ethanol vapor chambers. Controls are placed into air chambers and receive Pyr only. At 24 and 48 h, Pyr boosters are administered to both the experimental and control groups. Blood ethanol concentrations (BECs) for ethanol groups are measured and the ethanol vapor concentrations adjusted to equate ethanol exposure between lines. Mean BECs are maintained between approximately 1.0-2.0 mg/mL, depending upon the effects of the test compound being studied. After 72 h, all mice are removed from the chambers to initiate withdrawal, and ethanol treated mice have blood samples drawn for BEC determinations.

[0501] Following removal from the ethanol or air chambers, mice are scored hourly for handling-induced convulsion (HIC). Scoring is initiated 1 h after removal from ethanol and hourly over the next 12-15 h and again at 24 h. If animals do

not return to baseline HIC levels by 25 h, an additional score is obtained at 48 h. The scale such as the following is used (0—no convulsion after a gently 180° spin; 1—only facial grimace after gentle 180° spin; 2—tonic convulsion elicited by gently 180° spin; 3—tonic-clonic convulsion after 180° spin; 4—tonic convulsion when lifted by tail, no spin; 5—tonic-clonic convulsion when lifted by tail, no spin; 6—severe tonic-clonic convulsion when lifted by tail, no spin; and 7—severe tonic-clonic convulsion elicited before lifting by the tail). The area under the curve is calculated and used to quantitatively evaluate withdrawal severity. An additional index of withdrawal severity is the peak HIC score, calculated by identifying the highest HIC for each individual mouse and averaging this score with the two adjacent scores. Data are analyzed by appropriate statistical methods.

Description 6

Animal Model for Assessing Therapeutic Efficacy of
Acamprosate Prodrugs for Treating Tinnitus

Unilateral Noise Trauma

[0502] The efficacy of acamprosate prodrugs of Formula (I) for treating tinnitus can be assessed using animal models of tinnitus in which unilateral noise trauma is used to induce tinnitus (Bauer and Brozoski, *J Assoc Res Otolaryngology* 2001, 2(1), 54-64; and Guitton et al., US 2006/0063802). Long-Evans rats are first behaviorally acclimated to lever-press for food pellets and then conditioned to respond in a distinctive and standard way to auditory test stimuli. After conditioning, the animals are separated into groups and exposed to unilateral noise trauma for 0, 1, or 2 hours. Animals are anesthetized, placed in a stereotaxic head frame, and unilaterally exposed once to narrowband noise with a peak intensity of 105 dB centered at 16 kHz for 0, 1, or 2 hours before or after behavioral training and testing. The animals are then administered an acamprosate prodrug and suppression of the conditioned response determined and compared to a control group not exposed to noise trauma.

Sodium Salicylate-Induced Sound Experience

[0503] An animal model developed for short-term, acute induced phantom auditory sensations in rats can be used to evaluate acamprosate prodrugs for treating tinnitus. Salicylate-induced animal models of tinnitus are known.

[0504] Female albino rats (Wistar, aged 8-20 weeks) are trained and tested on five consecutive days per week. Training and testing takes place in a commercial conditioning chamber (rat shuttle box, TSE) adapted for the study. Electrical stimuli (0.1-0.5 mA, 100 V, 0.5 s) can be supplied via a shockable floor ground. A resting platform with a mechanical sensor was mounted on one side of the cage, covering the shockable floor and serving as a resting location for the animal. The cage is separated by a wall into two short hallways. At both ends of the hallways, within a recess, small amounts of fluid can be given to an animal, gravity-advanced and controlled by flow resistance- and vibration-muted magnetic shutter valves. A typical open time is 0.5 s, resulting in a reward drop of ca. 20 μ L, supplied to an animal via a curved metal drinking cannula. Reward drops not taken up by the animal are drained off into a reservoir unreachable by the rat. Photo sensors registered the visits of an animal at the feeder recesses. All sensors

are monitored on a computer screen and a top-mounted USB camera provided pictures of the entire floor dimensions of the cage interior.

[0505] Auditory stimuli are generated and presented over three broadened speakers mounted vertically in the cage. A continuous white noise can be played on the central loudspeaker switched off and on with a 100 ms ramp. In parallel to the white noise sound, a pure tone (cue tone, 8 kHz, 70 dB SPL, 200 ms length, 25 ms ramp, repeated five times with 300 ms pause) could be presented over loudspeakers mounted directly over the left and right feeder recesses.

[0506] Animals are trained on auditory stimuli for 30-60 min/day for 5 days/week. Training session length is adapted to the animal's activity. Always 15-18 h prior to behavioral testing (experimental session), the drinking water is withdrawn. The conditioned rats are divided into two groups (one animal per cage for either group). Animals from the first group receive an intraperitoneal injection of sodium salicylate (350 mg/kg bw) while animals from the second group receive an intraperitoneal injection of an equivalent volume of saline. Animals from either group are tested on the same day in a semi-random order exactly 3 h after injection. During the experimental session electrical stimuli are omitted. Four minutes after the start of a session the sugar water reward is stopped and the behavioral performances are recorded from 12-16 min and subsequently analyzed. Within the next 2-5 days rats receive the same training as before the experiment. On the next experimental day animals from the group previously treated with salicylate are injected with saline or test compound and tested again.

[0507] Frequencies of feeder access action of a rat are calculated for periods of sound and periods of silence separately (accesses/min) and normalized (SA activity ratio). The difference of silence activity ratios (ASA ratio) is determined as the silence activity ratio of an animal tested after salicylate injection less the silence activity ratio of the same animal after saline injection. Data is analyzed using appropriate statistical methods.

[0508] During the training procedure, animals are conditioned to discriminate between periods of sound and periods of silence using auditory cues.

[0509] To induce phantom auditory sensations, animals are injected with salicylate (350 mg/kg bw) or an equivalent volume of saline and tested 3 h later. The SA ratio of animals treated with salicylate is significantly higher than the SA ratio for animals treated with saline.

[0510] Test compounds can be administered and their ability to reverse the effects of the salicylate induced phantom auditory sensations determined. Compounds that reduce the increase in the SA ratio following in the salicylate treated animals can have potential in treating tinnitus.

Description 7

Method for Assessing Therapeutic Efficacy of Acamprosate Prodrugs for Treating Tinnitus in Humans

[0511] The efficacy of acamprosate prodrugs of Formula (I) for treating tinnitus in humans can be assessed using methods known in the art.

[0512] Patients are screened using pre-established inclusion and exclusion criteria and selected for their ability to perform a psychophysical loudness matching task using pure tones and broad-band noise (BBN). Examples of inclusion criteria include, for example, age, type of tinnitus, e.g., con-

tinuous or pulsed, duration of tinnitus, Tinnitus Handicap Questionnaire (THQ) score >30, Beck Depression Index (BDI) <13, and criterion performance on loudness matching task using a 1 KHz standard.

[0513] Following screening, selection and enrollment, tinnitus is evaluated before and after an acamprosate prodrug is administered to a patient. Hearing thresholds are evaluated using an objective stimulus loudness match and a tinnitus loudness matching procedure.

[0514] Prior to enrollment, subjects are screened for proficiency in a psychophysical matching task. In the objective stimulus loudness matching procedure, subjects match a binaural 1 KHz standard tone at 20 dB sensation levels to each of five binaural comparison stimuli (BBN, 0.5, 1, 2, and 4 KHz). The loudness match is obtained using a forced two-choice procedure. Each trial begins with the simultaneous presentation of a visual cue and the 1 KHz standard followed by the presentation of the second visual cue and the comparison stimulus. Subjects are instructed to indicate whether the standard and comparison stimuli sound the "same" or "different" in loudness by clicking an on-screen button. An ascending-descending method of limits procedure is used. Subjects are screened using this loudness-matching test and are required to meet inclusion criteria of efficiency (completion time ≤ 1 h) and reliability (standard deviation of match levels ≤ 5 dB).

[0515] The tinnitus loudness matching procedure differs from the objective stimulus loudness matching procedure in that the initial presentation on each trial is a null presentation during which an on-screen message instructs subjects to listen closely to their tinnitus. During this initial 1-sec cue subjects are instructed to use their perception of tinnitus as the standard stimulus. Subjects are instructed to click a "same loudness" button when the loudness of the comparison stimulus matches the loudness of their tinnitus. The presentation order of the comparison stimuli (BBN, 0.5, 1, 2, and 4 KHz) is randomized, and each ascending and descending stimulus series is repeated once, for a total of four tinnitus loudness matches at each of the five comparison stimuli. The intensities of the loudness-match points are recorded and converted to sensation levels of tinnitus loudness using the hearing threshold determined in each session for the comparison stimuli. Psychoacoustically determined tinnitus loudness is reported as dB HL of the maximum sensation-level match obtained within a session.

[0516] Assessment sessions are performed at the initiation of the study and at intervals during the study. Subjects can be given placebo only, an acamprosate prodrug only, a variable including escalating or deescalating dose of an acamprosate prodrug, or a combination of placebo and acamprosate prodrug during the course of a study. The duration of the study can be a few hours, days, weeks, months, or years.

[0517] Primary outcome measures are psychoacoustically determined tinnitus loudness and perceived tinnitus handicap. Tinnitus handicap can be determined using the Tinnitus Handicap Questionnaire, which provides a global score and subscores related to emotional, functional, and cognitive aspects of tinnitus. Secondary outcome measures include general health and quality of life factors determined using, for example, the General Health Survey Short form (RAND 36-Item Health Survey, 1.0, Rand Health, Santa Monica, Calif.) and the Tinnitus Experience Questionnaire, a set of seven scaled questions that evaluate the experiential sensory features of tinnitus. Other questionnaires for assessing tinnitus can be used.

Description 8

Animal Model for Assessing Therapeutic Efficacy of
Acamprosate Prodrugs for Treating Sleep Apnea

[0518] Sprague-Dawley rats are anesthetized and a surgical incision of the scalp is made to allow bilateral implantation of stainless steel screws into the frontal and parietal bones of the skull for electroencephalogram (EEG) recording. Bilateral wire electrodes are placed into the nuchal muscles for electromyogram (EMG) recording. The skin is then sutured and the animals allowed at least 7 days for recovery. Respirations are recorded by placing each rat inside a single chamber plethysmograph. The plethysmograph chamber is flushed with room air at a constant regulated flow rate of 2 L/min. EEG, EMG and respirations are continuously recorded. Sleep apneas are defined as cessation of respiratory effort for at least 2.5 s. The effects of recording hour, sleep state, and acamprosate prodrug administration are analyzed using appropriate statistical methods.

Description 9

Study for Assessing the Therapeutic Efficacy of
Acamprosate Prodrugs for Treating Sleep Apnea in
Humans

[0519] Inclusion criteria are an apnea-hypopnea index (AHI) exceeding 20 based on self-rated sleep duration at previous unattended ventilatory screening or an AHI exceeding 25 in a previous polysomnographic (PSG) recording. A double blind, randomized, placebo-controlled cross-over study comparing the effects of an acamprosate prodrug and placebo is used. Each patient undergoes a complete PSG recording for habituation at night 1. Patients are randomized to receive acamprosate prodrug on night 2 and placebo on night 3, or vice versa. Night 2 is scheduled within 1-21 days after night 1 and night 3 within 7-28 days after night 1 to provide a minimum of 7 days between night 2 and night 3 washout. A complete PSG recording, physical examination, and recording of ECG is performed in an identical manner at all study nights. Blood samples are obtained in the morning after study nights for hematology and clinical chemistry. Adverse events are determined by active questioning. AHI, the number of obstructive apneic/hyponeic events per time, is the primary efficacy variable. Secondary efficacy variables are REM AHI, non-REM AHI, apnea index (AI), hypopnea index (HI), oxygen desaturation index (ODI), minimum overnight oxygen saturation, sleep stage distribution arousal index, REM sleep and slow wave sleep latency, safety and tolerability. An obstructive apnea is defined as loss of nasal pressure accompanied by paradoxical respiratory movements for >10 s. An obstructive hypopnea is defined as a >50% reduction of the nasal pressure signal, but accompanied by chest wall paradoxical motion through most of inspiration for >10 s. Events without respiratory movements are classified as central apneas.

Description 10

Animal Models for Assessing Therapeutic Efficacy
of Acamprosate Prodrugs for Treating Parkinson's
Disease

MPTP Induced Neurotoxicity

[0520] MPTP, or 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine is a neurotoxin that produces a Parkinsonian syndrome in both man and experimental animals. Studies of the mecha-

nism of MPTP neurotoxicity show that it involves the generation of a major metabolite, MPP⁺, formed by the activity of monoamine oxidase on MPTP. Inhibitors of monoamine oxidase block the neurotoxicity of MPTP in both mice and primates. The specificity of the neurotoxic effects of MPP⁺ for dopaminergic neurons appears to be due to the uptake of MPP⁺ by the synaptic dopamine transporter. Blockers of this transporter prevent MPP⁺ neurotoxicity. MPP⁺ has been shown to be a relatively specific inhibitor of mitochondrial complex I activity, binding to complex I at the rotenone binding site and impairing oxidative phosphorylation. In vivo studies have shown that MPTP can deplete striatal ATP concentrations in mice. It has been demonstrated that MPP⁺ administered intrastratially to rats produces significant depletion of ATP as well as increased lactate concentration confined to the striatum at the site of the injections. Compounds that enhance ATP production can protect against MPTP toxicity in mice.

[0521] A prodrug of Formula (I) is administered to mice or rats for three weeks before treatment with MPTP. MPTP is administered at an appropriate dose, dosing interval, and mode of administration for 1 week before sacrifice. Control groups receive either normal saline or MPTP hydrochloride alone. Following sacrifice the two striate are rapidly dissected and placed in chilled 0.1 M perchloric acid. Tissue is subsequently sonicated and aliquots analyzed for protein content using a fluorometer assay. Dopamine, 3,4-dihydroxyphenylacetic acid (DOPAC), and homovanillic acid (HVA) are also quantified. Concentrations of dopamine and metabolites are expressed as nmol/mg protein.

[0522] Prodrugs of Formula (I) that protect against DOPAC depletion induced by MPTP, HVA, and/or dopamine depletion are neuroprotective and therefore can be useful for the treatment of Parkinson's disease.

Haloperidol-Induced Hypolocomotion

[0523] The ability of a compound to reverse the behavioral depressant effects of dopamine antagonists such as haloperidol in rodents and is considered a valid method for screening drugs with potential antiparkinsonian effects. Hence, the ability of prodrugs of Formula (I) to block haloperidol-induced deficits in locomotor activity in mice can be used to assess both in vivo and potential anti-Parkinsonian efficacy.

[0524] Mice used in the experiments are housed in a controlled environment and allowed to acclimatize before experimental use. 1.5 h before testing, mice are administered 0.2 mg/kg haloperidol, a dose that reduces baseline locomotor activity by at least 50%. A test compound is administered 5-60 min prior to testing. The animals are then placed individually into clean, clear polycarbonate cages with a flat perforated lid. Horizontal locomotor activity is determined by placing the cages within a frame containing a 3x6 array of photocells interfaced to a computer to tabulate beam interrupts. Mice are left undisturbed to explore for 1 h, and the number of beam interruptions made during this period serves as an indicator of locomotor activity, which is compared with data for control animals for statistically significant differences.

6-Hydroxydopamine Animal Model

[0525] The neurochemical deficits seen in Parkinson's disease can be reproduced by local injection of the dopaminergic neurotoxin, 6-hydroxydopamine (6-OHDA) into brain

regions containing either the cell bodies or axonal fibers of the nigrostriatal neurons. By unilaterally lesioning the nigrostriatal pathway on only one-side of the brain, a behavioral asymmetry in movement inhibition is observed. Although unilaterally-lesioned animals are still mobile and capable of self maintenance, the remaining dopamine-sensitive neurons on the lesioned side become supersensitive to stimulation. This is demonstrated by the observation that following systemic administration of dopamine agonists, such as apomorphine, animals show a pronounced rotation in a direction contralateral to the side of lesioning. The ability of compounds to induce contralateral rotations in 6-OHDA lesioned rats has been shown to be a sensitive model to predict drug efficacy in the treatment of Parkinson's disease.

[0526] A 2 cm long incision is made along the midline of the scalp and the skin retracted and clipped back to expose the skull. A small hole is then drilled through the skull above the injection site. In order to lesion the nigrostriatal pathway, the injection cannula is slowly lowered to position above the right medial forebrain bundle at -3.2 mm anterior posterior, -1.5 mm medial lateral from the bregma, and to a depth of 7.2 mm below the duramater. Two minutes after lowering the cannula, 6-OHDA is infused at a rate of 0.5 μ L/min over 4 min, to provide a final dose of 8 μ g. The cannula is left in place for an additional 5 min to facilitate diffusion before being slowly withdrawn. The skin is closed, the animal removed from the stereotaxic frame, and returned to its housing. The rats are allowed to recover from surgery for two weeks before behavioral testing.

[0527] Rotational behavior is measured using a rotameter system having stainless steel bowls (45 cm dia $\times$ 15 cm high) enclosed in a transparent Plexiglas cover around the edge of the bowl and extending to a height of 29 cm. To assess rotation, rats are placed in a cloth jacket attached to a spring tether connected to an optical rotameter positioned above the bowl, which assesses movement to the left or right either as partial (45°) or full (360°) rotations.

[0528] To reduce stress during administration of a test compound, rats are initially habituated to the apparatus for 15 min on four consecutive days. On the test day, rats are given a test compound, e.g., a prodrug of Formula (I). Immediately prior to testing, animals are given a subcutaneous injection of a subthreshold dose of apomorphine, and then placed in the harness and the number of rotations recorded for one hour. The total number of full contralateral rotations during the hour test period serves as an index of antiparkinsonian drug efficacy.

L-Dopa Induced Dyskinesia

[0529] The ability of acamprosate prodrugs to mitigate the effects of L-dopa induced dyskinesia can be assessed using animal models.

[0530] Male, Sprague-Dawley rats (250-300 g) are housed and maintained under standard conditions.

[0531] Reserpine (4 mg/kg) is administered under light isoflurane anesthesia. Eighteen hours following reserpine administration, the animals are placed into observation cages. Behavior is assessed using an automated movement detection system that includes dual layers of rectangular grids of sensors containing an array of 24 infrared beams surrounding the cage. Each beam break is registered as an activity count and contributes to the assessment of a variety of different behavioral parameters depending on the location of the event and the timing of successive beam breaks. These parameters

include: (1) horizontal activity, a measure of the number of beams broken on the lower level; and (2) vertical activity, a measure of beams broken on the upper level.

[0532] In one experiment, immediately prior to commencing behavioral assessments, rats are injected with a combination of L-dopa methyl ester and carbidopa (or benserazide). In another study, to assess the effects of acamprosate prodrugs on L-dopa induced activity, animals are randomly assigned to groups. In each group, immediately following L-dopa/carbidopa administration, vehicle or acamprosate prodrug is administered. The behavior of normal, non-reserpine-treated, animals is also assessed. Behavior of the animals in the different groups is monitored for at least 4 hours. Acamprosate prodrugs that reduce the L-dopa-induced locomotion in the reserpine-treated rats are potentially useful in treating Parkinson's disease and/or the symptoms associated with Parkinson's disease.

Description 11

Use of Clinical Trials to Assess the Efficacy of Acamprosate Prodrugs for Treating Parkinson's Disease

[0533] The following clinical study may be used to assess the efficacy of a compound in treating Parkinson's disease.

[0534] Patients with idiopathic PD fulfilling the Queen Square Brain Bank criteria with motor fluctuations and a defined short duration GABA analog response (1.5-4 hours) are eligible for inclusion. Clinically relevant peak dose dyskinesias following each morning dose of their current medication are a further pre-requisite. Patients are also required to have been stable on a fixed dose of treatment for a period of at least one month prior to starting the study. Patients are excluded if their current drug regime includes slow-release formulations of L-Dopa, COMT inhibitors, selegiline, anticholinergic drugs, or other drugs that could potentially interfere with gastric absorption (e.g. antacids). Other exclusion criteria include patients with psychotic symptoms or those on antipsychotic treatment, patients with clinically relevant cognitive impairment, defined as MMS (Mini Mental State) score of less than 24, risk of pregnancy, Hoehn & Yahr stage 5 in off-status, severe, unstable diabetes mellitus, and medical conditions such as unstable cardiovascular disease or moderate to severe renal or hepatic impairment. Full blood count, liver, and renal function blood tests are taken at baseline and after completion of the study.

[0535] A randomized, double blind, and cross-over study design is used. Each patient is randomized to the order in which either L-dopa or one of the two dosages of test compound, e.g., an acamprosate prodrug, is administered in a single-dose challenge in double-dummy fashion in three consecutive sessions. Randomization is by computer generation of a treatment number, allocated to each patient according to the order of entry into the study. All patients give informed consent.

[0536] Patients are admitted to a hospital for an overnight stay prior to administration of test compound the next morning on three separate occasions at weekly intervals. After withdrawal of all antiparkinsonian medication from midnight the previous day, test compound is administered at exactly the same time in the morning in each patient under fasting conditions.

[0537] Patients are randomized to the order of the days on which they receive placebo or test compound. The pharma-

cokinetics of a test compound can be assessed by monitoring plasma acamprosate concentration over time. Prior to administration, a 22 G intravenous catheter is inserted in a patient's forearm. Blood samples of 5 ml each are taken at baseline and 15, 30, 45, 60, 75, 90, 105, 120, 140, 160, 180, 210, and 240 minutes after administering a test compound or until a full off state has been reached if this occurs earlier than 240 minutes after drug ingestion. Samples are centrifuged immediately at the end of each assessment and stored deep frozen until assayed. Plasma acamprosate levels are determined by high-pressure liquid chromatography (HPLC). On the last assessment additional blood may be drawn for routine hematology, blood sugar, liver, and renal function.

[0538] For clinical assessment, motor function is assessed using the United Parkinson's Disease Rating Scale motor score and BrainTest, which is a tapping test performed with a patient's more affected hand on the keyboard of a laptop computer. These tests are carried out at baseline and then immediately following each blood sample until patients reach their full on-stage, and thereafter at 3 intervals of 20 min, and 30 min intervals until patients reach their baseline off-status. Once patients reach their full on-state, video recordings are performed three times at 20 min intervals. The following mental and motor tasks, which have been shown to increase dyskinesia, are monitored during each video session: (1) sitting still for 1 minute; (2) performing mental calculations; (3) putting on and buttoning a coat; (4) picking up and drinking from a cup of water; and (5) walking. Videotapes are scored using, for example, versions of the Goetz Rating Scale and the Abnormal Involuntary Movements Scale to document a possible increase in test compound induced dyskinesia.

[0539] Occurrence and severity of dyskinesia is measured with a Dyskinesia Monitor. The device is taped to a patient's shoulder on their more affected side. The monitor records during the entire time of a challenging session and provides a measure of the frequency and severity of occurring dyskinesias.

[0540] Results can be analyzed using appropriate statistical methods.

Description 12

Use of Clinical Trials to Assess the Efficacy of Acamprosate Prodrugs for Treating Levodopa-Induced Dyskinesias in Parkinson's Disease

[0541] A double-blind placebo-controlled clinical trial such as that described by Goetz et al., *Movement Disorders* 2007, 22(2), 179-186 can be used to assess the efficacy of an acamprosate prodrug for treating levodopa-induced dyskinesias in Parkinson's disease.

[0542] Patients are 30 years of age or older with Parkinson's disease and received levodopa treatment at a stable (at least 4 weeks) and optimized dose. Following enrollment, patients are randomized and receive either placebo or an appropriate dose and regimen of acamprosate prodrug. Levodopa doses are maintained at the baseline level. At appropriate intervals during the study, patients are evaluated for periods during the day characterized by sleep, off, on-without dyskinesias, on-with non-troublesome dyskinesias, and on-with troublesome dyskinesia. The primary outcome is change from baseline in on-time without dyskinesia. Various dyskinesia rating scales such as, for example, the Abnormal Involuntary Movement Scale, Unified Parkinson's Disease Rating Scale (UPDRS) Motor examination (Part III), or

UPDRS Activities of Daily Living assessment (Part III) can also be used. Measures of safety such as frequency and severity of reported adverse events, changes in vital signs, laboratory test results, including ACTH-suppression testing of cortisol levels and electrocardiogram can also be determined.

Description 13

Animal Model for Assessing Therapeutic Efficacy of Acamprosate Prodrugs for Treating Alzheimer's Disease

[0543] Heterozygous transgenic mice expressing the Swedish AD mutant gene, hAPPK670N, M671L (Tg2576) are used as an animal model of Alzheimer's disease. Beginning at 9 months of age, mice are divided into two groups. The first two groups of animals receive increasing doses of an acamprosate prodrug, over six weeks. The remaining control group receives daily saline injections for six weeks.

[0544] Behavioral testing is performed at each drug dose using the same sequence over two weeks in all experimental groups: (1) spatial reversal learning, (2) locomotion, (3) fear conditioning, and (4) shock sensitivity. This order is selected to minimize interference among testing paradigms.

[0545] Acquisition of the spatial learning paradigm and reversal learning are tested during the first five days of test compound administration using a water T-maze. Mice are habituated to the water T-maze during days 1-3, and task acquisition begins on day 4. On day 4, mice are trained to find the escape platform in one choice arm of the maze until 6 to 8 correct choices are made on consecutive trials. The reversal learning phase is then conducted on day 5. During the reversal learning phase, mice are trained to find the escape platform in the choice arm opposite from the location of the escape platform on day 4. The same performance criterion and inter-trial interval are used as during task acquisition.

[0546] Large ambulatory movements are assessed to determine that the results of the spatial reversal learning paradigm are not influenced by the capacity for ambulation. After a rest period of two days, horizontal ambulatory movements, excluding vertical and fine motor movements, are assessed in a chamber equipped with a grid of motion-sensitive detectors on day 8. The number of movements accompanied by simultaneous blocking and unblocking of a detector in the horizontal dimension are measured during a one-hour period.

[0547] The capacity of an animal for contextual and cued memory is tested using a fear conditioning paradigm beginning on day 9. Testing takes place in a chamber that contains a piece of absorbent cotton soaked in an odor-emitting solution such as mint extract placed below the grid floor. A 5-min, 3 trial 80 db, 2800 Hz tone-foot shock sequence is administered to train the animals on day 9. On day 10, memory for context is tested by returning each mouse to the chamber without exposure to the tone and foot shock, and recording the presence or absence of freezing behavior every 10 seconds for 8 minutes. Freezing is defined as no movement, such as ambulation, sniffing or stereotypy, other than respiration.

[0548] On day 11, the response of the animal to an alternate context and to the auditory cue is tested. Coconut extract is placed in a cup and the 80 dB tone is presented, but no foot shock is delivered. The presence or absence of freezing in response to the alternate context is then determined during the first 2 minutes of the trial. The tone is then presented continuously for the remaining 8 minutes of the trial, and the presence or absence of freezing in response to the tone is determined.

On day 12, the animals are tested to assess their sensitivity to the conditioning stimulus, i.e., foot shock. Following the last day of behavioral testing, animals are anesthetized and the brains removed, post-fixed overnight, and sections cut through the hippocampus. The sections are stained to image β -amyloid plaques.

[0549] Data are analyzed using appropriate statistical methods.

Description 14

Animal Model for Assessing Therapeutic Efficacy of Acamprosate Prodrugs for Treating Huntington's Disease

Neuroprotective Effects in a Transgenic Mouse Model of Huntington's Disease

[0550] Transgenic HD mice of the N171-82Q strain and non-transgenic littermates are treated with a prodrug of Formula (I) or a vehicle from 10 weeks of age. The mice are placed on a rotating rod ("rotarod"). The length of time at which a mouse falls from the rotarod is recorded as a measure of motor coordination. The total distance traveled by a mouse is also recorded as a measure of overall locomotion. Mice administered prodrugs of Formula (I) that are neuroprotective in the N 171-82Q transgenic HD mouse model remain on the rotarod for a longer period of time and travel further than mice administered vehicle.

Malonate Model of Huntington's Disease

[0551] A series of reversible and irreversible inhibitors of enzymes involved in energy generating pathways has been used to generate animal models for neurodegenerative diseases such as Parkinson's and Huntington's diseases. In particular, inhibitors of succinate dehydrogenase, an enzyme that impacts cellular energy homeostasis, has been used to generate a model for Huntington's disease. The enzyme succinate dehydrogenase plays a central role in both the tricarboxylic acid cycle as well as the electron transport chain in mitochondria. Malonate is a reversible inhibitor of succinate dehydrogenase. Intrastriatal injections of malonate in rats have been shown to produce dose dependent striatal excitotoxic lesions that are attenuated by both competitive and noncompetitive NMDA antagonists. For example, the glutamate release inhibitor, lamotrigine, also attenuates the lesions. Co-injection with succinate blocks the lesions, consistent with an effect on succinate dehydrogenase. The lesions are accompanied by a significant reduction in ATP levels as well as a significant increase in lactate levels in vivo as shown by chemical shift resonance imaging. The lesions produce the same pattern of cellular sparing, which is seen in Huntington's disease, supporting malonate challenge as a useful model for the neuropathologic and neurochemical features of Huntington's disease.

[0552] To evaluate the effect of acamprosate prodrugs of Formula (I) in this malonate model for Huntington's disease, a prodrug of Formula (I) is administered at an appropriate dose, dosing interval, and route, to male Sprague-Dawley rats. A prodrug is administered for two weeks prior to the administration of malonate and then for an additional week prior to sacrifice. Malonate is dissolved in distilled deionized water and the pH adjusted to 7.4 with 0.1 M HCl. Intrastriatal injections of 1.5 μ L of 3 μ mol malonate are made into the left striatum at the level of the Bregma 2.4 mm lateral to the

midline and 4.5 mm ventral to the dura. Animals are sacrificed at 7 days by decapitation and the brains quickly removed and placed in ice cold 0.9% saline solution. Brains are sectioned at 2 mm intervals in a brain mold. Slices are then placed posterior side down in 2% 2,3,5-tiphenyltetrazolium chloride. Slices are stained in the dark at room temperature for 30 min and then removed and placed in 4% paraformaldehyde pH 7.3. Lesions, noted by pale staining, are evaluated on the posterior surface of each section. The measurements are validated by comparison with measurements obtained on adjacent Nissl stain sections. Compounds exhibiting a neuroprotective effect and therefore potentially useful in treating Huntington's disease show a reduction in malonate-induced lesions.

Description 15

Animal Model for Assessing Therapeutic Efficacy of Acamprosate Prodrugs for Treating Amyotrophic Lateral Sclerosis

[0553] A murine model of SOD1 mutation-associated ALS has been developed in which mice express the human superoxide dismutase (SOD) mutation glycine $\rightarrow$ alanine at residue 93 (SOD1). These SOD1 mice exhibit a dominant gain of the adverse property of SOD, and develop motor neuron degeneration and dysfunction similar to that of human ALS. The SOD1 transgenic mice show signs of posterior limb weakness at about 3 months of age and die at 4 months. Features common to human ALS include astrogliosis, microgliosis, oxidative stress, increased levels of cyclooxygenase/prostaglandin, and, as the disease progresses, motor neuron loss.

[0554] Studies are performed on transgenic mice overexpressing human Cu/Zn—SOD G93A mutations ((B6SJL-TgN (SOD1-G93A) 1 Gur)) and non-transgenic B6/SJL mice and their wild litter mates. Mice are housed on a 12-hr day/light cycle and (beginning at 45 d of age) allowed ad libitum access to either test compound-supplemented chow, or, as a control, regular formula cold press chow processed into identical pellets. Genotyping can be conducted at 21 days of age. The SOD1 mice are separated into groups and treated with a test compound, e.g., an acamprosate prodrug, or serve as controls.

[0555] The mice are observed daily and weighed weekly. To assess health status mice are weighed weekly and examined for changes in lacrimation/salivation, palpebral closure, ear twitch and pupillary responses, whisker orienting, postural and righting reflexes and overall body condition score. A general pathological examination is conducted at the time of sacrifice.

[0556] Motor coordination performance of the animals can be assessed by one or more methods known to those skilled in the art. For example, motor coordination can be assessed using a neurological scoring method. In neurological scoring, the neurological score of each limb is monitored and recorded according to a defined 4-point scale: 0—normal reflex on the hind limbs (animal will splay its hind limbs when lifted by its tail); 1—abnormal reflex of hind limbs (lack of splaying of hind limbs when animal is lifted by the tail); 2—abnormal reflex of limbs and evidence of paralysis; 3—lack of reflex and complete paralysis; and 4—inability to right when placed on the side in 30 seconds or found dead. The primary end point is survival with secondary end points of neurological score and body weight. Neurological score observations and

body weight are made and recorded five days per week. Data analysis is performed using appropriate statistical methods.

[0557] The rotarod test evaluates the ability of an animal to stay on a rotating dowel allowing evaluation of motor coordination and proprioceptive sensitivity. The apparatus is a 3 cm diameter automated rod turning at, for example, 12 rounds per min. The rotarod test measures how long the mouse can maintain itself on the rod without falling. The test can be stopped after an arbitrary limit of 120 sec. Should the animal fall down before 120 sec, the performance is recorded and two additional trials are performed. The mean time of 3 trials is calculated. A motor deficit is indicated by a decrease of walking time.

[0558] In the grid test, mice are placed on a grid (length: 37 cm, width: 10.5 cm, mesh size: 1×1 cm²) situated above a plane support. The number of times the mice put their paws through the grid is counted and serves as a measure for motor coordination.

[0559] The hanging test evaluates the ability of an animal to hang on a wire. The apparatus is a wire stretched horizontally 40 cm above a table. The animal is attached to the wire by its forepaws. The time needed by the animal to catch the string with its hind paws is recorded (60 sec max) during three consecutive trials.

[0560] Electrophysiological measurements (EMG) can also be used to assess motor activity condition. Electromyographic recordings are performed using an electromyography apparatus. During EMG monitoring mice are anesthetized. The measured parameters are the amplitude and the latency of the compound muscle action potential (CMAP). CMAP is measured in gastrocnemius muscle after stimulation of the sciatic nerve. A reference electrode is inserted near the Achilles tendon and an active needle placed at the base of the tail. A ground needle is inserted on the lower back of the mice. The sciatic nerve is stimulated with a single 0.2 msec pulse at supramaximal intensity (12.9 mA). The amplitude (mV) and the latency of the response (ms) are measured. The amplitude is indicative of the number of active motor units, while distal latency reflects motor nerve conduction velocity.

[0561] The efficacy of test compounds can also be evaluated using biomarker analysis. To assess the regulation of protein biomarkers in SOD1 mice during the onset of motor impairment, samples of lumbar spinal cord (protein extracts) are applied to ProteinChip Arrays with varying surface chemical/biochemical properties and analyzed, for example, by surface enhanced laser desorption ionization time of flight mass spectrometry. Then, using integrated protein mass profile analysis methods, data is used to compare protein expression profiles of the various treatment groups. Analysis can be performed using appropriate statistical methods.

Description 16

Assessing Therapeutic Efficacy of Acamprosate Prodrugs for Treating Cortical Spreading Depression

[0562] It has been hypothesized that cortical spreading depression emanating from a site of injury causes secondary damage in the "penumbra" by disrupting ion homeostasis and producing demands on neurons already in a compromised state. Focal CNS injury or ischemia also results in an induction of the immediate early gene c-fos. c-fos induction spreads throughout the injured hemisphere by a process that appears to be dependent on cortical spreading depression. c-fos induction is inhibited by NMDA receptor antagonists.

Thus, both cortical spreading depression and c-fos induction are NMDA receptor activated processes associated with CNS injury and may be components of the cascade leading to neuron death.

[0563] NMDA-induced increase in fos immunoreactivity in mice is determined according to the following protocol. Male CF-1 mice (20 to 25 g) are administered varying doses of an acamprosate prodrug or vehicle. Thirty min later, animals receive intraperitoneal administration of NMDA (75 mg/kg) or vehicle. Sixty min later, animals are terminally anesthetized, brains are removed to ice and immersed for 1 h in 2% paraformaldehyde in phosphate buffered saline, and transferred to 15% sucrose in phosphate buffered saline, incubated overnight, and then frozen at -80° C. Coronal sections through the hippocampal region are taken, washed, and incubated with a sheep anti-fos polyclonal antibody (OA-11-824) for 18 h at 4° C. Sections are washed with phosphate buffered saline and then incubated with biotinylated rabbit anti-sheep antibody for 2 h. After 3 washes in phosphate buffered saline, sections are incubated in Vector ABC solution for 1 h at 25° C., washed 3 times, stained for glucose oxidase, and mounted. Each section is photographed and the intensity of fos-like immunoreactivity in the dentate gyrus is analyzed.

[0564] CNS trauma-induced c-fos mRNA induction in rats is determined according to the following protocol. Male Sprague-Dawley rats (200-250 g) are administered different doses of an acamprosate prodrug or vehicle. After 30 min, animals are anesthetized and a burr hole drilled over the right frontal parietal cortex 3 mm anterior and 3 mm lateral to bregma. An 18-gauge needle is inserted through the hole for 2 min to a depth of about 3 mm into the cortex. After a 60 min recovery animals are sacrificed, the brains removed, and cortices dissected and frozen in liquid nitrogen. Changes in c-fos mRNA expression following needle injury are quantified using procedures known in the art.

[0565] To assess the effects of acamprosate prodrugs on electrically-induced cortical spreading depression in rats, male Sprague-Dawley rats (275-325 g) are anaesthetized. The spontaneously breathing animals are fixed in a stereotaxic frame, a craniotomy is drilled over the parietal cortex, and the dura is removed. Two saline-filled glass recording microelectrodes each containing a Ag/AgCl wire are inserted into the parietal cortex at a depth of about 1 mm and 1.5-2.0 mm apart along the sagittal plane using a micromanipulator. Two saline filled cannulae each containing a Ag/AgCl wire are inserted under the skin of the animal to serve as reference electrodes. Cortical spreading depression is induced in the parietal cortex using a bipolar stimulating electrode placed at 90° to the frontal recording electrode and positioned so that the electrode visibly touches but does not depress the cortex. Electroconvulsive stimulation consists of a train of 5 ms pulses at 40 Hz lasting for 2 s. The threshold stimulation for cortical spreading depression determined by varying the current. Once the threshold current has been determined, the current is increased by 20% for experimental measurements. DC potentials are recorded at 10 min intervals for four control stimulations. An acamprosate prodrug is then administered. DC potentials are again recorded at 10 min intervals. The speed of cortical spreading depression expansion is calculated from the latency difference of the negative DC shift at the rostral and caudal electrodes.

Description 17

Animal Models to Assess the Efficacy of Acamprosate Prodrugs for Treating Migraine

[0566] Therapeutic activity of acamprosate prodrugs provided by the present disclosure may be determined in various

animal models of neuropathic pain or in clinically relevant studies of different types of neuropathic pain. Animal models for neuropathic pain are known in the art and include animal models that determine analgesic activity or compounds that act on the CNS to reduce the phenomenon of central sensitization that results in pain from nonpainful or nonnoxious stimuli. Other animal models that are known in the art, such as hot plate tests, model acute pain, are useful for determining analgesic properties of compounds that are effective when painful or noxious stimuli are present. The progression of migraines is believed to be similar to the progression of epilepsy (because an episodic phenomenon underlies the initiation of the epileptic episode) and, as such, it is believed that epilepsy animal models may be useful in determining efficacy in treating migraine. Analgesic Activity

[0567] The following test can be used to evaluate the analgesic activity of an acamprosate prodrug. Test compound is administered orally to mice. Morphine is administered as a reference substance at 64 mg/kg to mice under the same experimental conditions. A vehicle is administered to mice as a control substance under the same experimental conditions. Test compound, morphine, or vehicle is administered to the mice in a blind study. Sixty minutes after the test compound, morphine, or vehicle is administered, the mice are placed onto a hot metal plate maintained at 54° C. and surrounded by a Plexiglass cylinder. The time taken for the mice to lick their feet is an index of analgesic activity. Effective analgesics increase the latency or amount of time to licking. Latency to the first foot lick is measured, up to a maximum time of 30 sec to prevent tissue damage to the mice.

Hyperreflexia and Flexor Reflex Tests

[0568] Assessment of hyperreflexia, pain, and muscle tone in chronic spinally transected rats is performed using male albino Holtzman-derived rats weighing 270-530 gm. The rats are housed independently and have continuous access to food and water throughout the experiments. Animals are anesthetized. Rats are placed in a stereotaxic frame and anesthesia is maintained. An incision is made so that the paraspinal muscles can be retracted and a laminectomy performed between T6-T9. A one- to two-millimeter portion of the spinal cord is removed by evacuation and replaced with gel foam to reduce bleeding, after which the incision is closed in layers.

[0569] Following the transection, rats are placed in a room in which the ambient temperature is raised to about 27° C. to maintain body temperature. On the following morning post-surgery, the hindquarters of the spinalized rats are bathed and their urine expressed manually by applying pressure to their bladders. Experiments are conducted between 21 and 28 days after surgery. For the first two weeks post-surgery, 0.25 mL of an antibiotic is administered to the rats to prevent bladder infection. A topical antibiotic is applied to any part of the skin that shows signs of decubitus lesions. Within approximately two weeks, all animals regain bladder control and are no longer given antibiotic treatment. Assessment of hyperreflexia and flexor reflex is performed before and after treatment with test compound so that each animal serves as its own control.

[0570] Initial assessment of hyperreflexia is performed by rating the hyperreflexia response elicited with an innocuous stimulus, such as a metal probe. A metal probe is pressed against the lower abdomen at four specific sites. The response is evaluated for each of four trials using a scale ranging from zero (no response in all four trials) to four (a maximum,

tonic-clonic reaction elicited in all four trials). All scores, pre- and post-treatment, are transformed to indicate the percent of hyperreflexia, pain, or muscle tone. The data is analyzed using appropriate statistical methods.

[0571] After determining hyperreflexia before drug treatment, test compound is administered to the rats.

[0572] Polysynaptic flexor-reflex responses, elicited by stimuli that activate high-threshold afferents, are recorded as EMG activity from the ipsilateral hamstring muscle. Supramaximal electric shocks are applied to the hindpaw and recording electrodes are placed in the biceps femoris semitendinosus muscle. Five sets of stimuli are made at each time point. The flexor reflex is recorded, in periods with and without test compound, every 30 min once a stable baseline response is achieved. The data at time zero represent pretreatment control values. The responses are determined in spinalized rats by observing the flexor-reflex response before treatment and at each of 30, 60, 90, and 120 min following administration of test compound, baclofen (10 mg/kg sc), or vehicle (water, 12 ml/kg po). Efficacy is indicated when a test compound is shown to reduce the magnitude of the flexor-reflex responses in a chronic spinalized rat at all time points with similar efficacy to baclofen, the positive control.

Cutaneous Hypersensitivity Test

[0573] The effects of a test compound on nociceptive activation of the trigeminovascular system is determined using an animal model of migraine. A pharmaceutical composition comprising a test compound is administered to cats. To serve as positive and negative controls, a vehicle control is administered to the cats. Efficacy is indicated for compounds that inhibit trigeminovascular activation compared to the trigeminovascular activation in the cats that receive the vehicle.

Yawning

[0574] Yawning is a behavior that has been linked to activation of dopaminergic neurotransmission. Yawning is part of a behavioral syndrome occurring in most patients during a migraine attack. Blockage of quinipirole-induced yawning in rats has been used as an animal model to study the potential antagonism of migraine symptoms.

[0575] Male Sprague Dawley rats are acclimatized for 12 days before testing and at the time of the study. The rats are housed in standard size steel cages with four animals per cage and are maintained on a 12 hour light/dark schedule. Test compound or vehicle is administered 15 min before the dopamine D2 agonist quinipirole in vehicle or the vehicle alone is administered to the animals. The animals are then placed individually in 6 in x 6 in plexiglass observation cages and the number of yawns is counted for the subsequent 30 min. The data is analyzed by an appropriate statistical method.

[0576] The dopamine D2 agonist quinipirole can produce an average of 13-15 yawns per 30 minutes while no yawning behavior is typically observed in vehicle treated animals. Compounds that inhibit quinipirole-induced yawning may be efficacious in treating migraine.

Animal Model of Dural Protein Extravasation

[0577] The following animal model can be employed to determine the ability of an acamprosate prodrug to inhibit protein extravasation, an exemplary functional assay of the neuronal mechanism of migraine.

[0578] Rats or guinea pigs are anesthetized and placed in a stereotaxic frame with the incisor bar set at -3.5 mm for rats or -4.0 mm for guinea pigs. Following a midline sagittal scalp incision, two pairs of bilateral holes are drilled through the skull (6 mm posteriorly, 2.0 and 4.0 mm laterally in rats; 4 mm posteriorly and 3.2 and 5.2 mm laterally in guinea pigs, with all coordinates referenced to bregma). Pairs of stainless steel stimulating electrodes, insulated except at the tips are lowered through the holes in both hemispheres to a depth of 9 mm (rats) or 10.5 mm (guinea pigs) from dura.

[0579] Test compound is administered. About 7 min later a fluorescent dye (e.g., Evans Blue) is administered. The fluorescent dye complexes with proteins in the blood and functions as a marker for protein extravasation. Ten (10) min post-injection of the test compound, the left trigeminal ganglion is stimulated for 3 minutes at a current intensity of 1.0 mA (5 Hz, 4 msec duration) with a potentiostat/galvanostat. Fifteen minutes following stimulation, the animals are killed and exsanguinated with 20 mL of saline. The top of the skull is removed to facilitate collection of the dural membranes. Dural membrane samples are removed from both hemispheres, rinsed with water, and spread flat on microscopic slides. Once dried, the tissues are coverslipped with a 70% glycerol/water solution. A fluorescence microscope equipped with a grating monochromator and a spectrophotometer is used to quantify the amount of fluorescent dye in each sample.

[0580] The extravasation induced by the electrical stimulation of the trigeminal ganglion is an ipsilateral effect (i.e. occurs only on the side of the dura in which the trigeminal ganglion is stimulated). This allows the other (unstimulated) half of the dura to be used as a control. The ratio of the amount of extravasation in the dura from the stimulated side, over the amount of extravasation in the unstimulated side, is calculated. Control animals dosed with only saline, yield, for example, a ratio of about 2.0 in rats and about 1.8 in guinea pigs. In contrast, a compound that effectively prevents the extravasation in the dura from the stimulated side yields a ratio of about 1.0. Dose-response curves can be generated for a test compound and the dose that inhibits the extravasation by 50% (ID_{50}) or 100% (ID_{100}) can be determined.

Amygdala Kindling Model

[0581] A relationship has been reported between migraine, affective illness and epilepsy. Although the three disorders are distinct, they all are paroxysmal dysregulations of the nervous system that partially overlap in their pharmacology. The kindling model for complex-partial seizures is based on the progressive development of seizures combined with electroencephalographic (EEG) paroxysmal patterns induced by repeated initially subconvulsive electrical stimulation of limbic structures, e.g., the basolateral nucleus of the amygdala. Once established, the phenomenon persists for months. Since the amygdala-kindled seizures in animals share numerous characteristics with complex-partial seizures in humans, it is a useful animal model of complex partial seizures. An advantage of using the amygdala kindling model is that both behavioral and EEG parameters of the partial and generalized seizures can be measured. Furthermore, the amygdala kindling model is reported to be appropriate for studying diseases such as migraine, affective illness, and epilepsy which increase in severity over time and in a manner which is related to the number of symptomatic episodes.

[0582] Rats are obtained at an age of 11-12 weeks (body weight 180-200 gm). Rats are maintained separately in plastic cages at controlled temperature (23°C .) and humidity (about 50% RH) with a 12-h light cycle. The rats receive standard diet and tap water ad libitum.

[0583] For implantation of stimulation and recording electrodes, rats are anesthetized and receive stereotaxic implantation of one bipolar electrode in the right basolateral amygdala. Coordinates for electrode implantation are AP-2.2 mm, L-4.8 mm, V-8.5 mm. All coordinates are measured from bregma. Skull screws serve as the reference electrode. The electrode assembly is attached to the skull by dental acrylic cement. After a postoperative period of 2 weeks, constant current stimulations ($500\text{ }\mu\text{A}$, 1 ms, monophasic square-wave pulses, 50/sec for 1 sec) are delivered to the amygdala at intervals of 1/day until ten stage 5 seizures are elicited. The electrical susceptibility of the stimulated region (threshold for induction of afterdischarges) is recorded on the first day of the experiment (initial afterdischarge threshold) as well as after kindling acquisition (with an interval of at least 4 days after the tenth stage 5 seizure) using an ascending staircase procedure. The initial current intensity is $1\text{ }\mu\text{A}$, and the current intensity is increased in steps of about 20% of the previous current at intervals of 1 min until an afterdischarge of at least 3 sec duration is elicited. In addition to afterdischarge threshold, the following parameters of kindled seizures are measured in fully-kindled rats after stimulation with the afterdischarge threshold current: seizure severity is classified as follows: 1—immobility, eye closure, twitching of vibrissae, sniffing, facial clonus; 2—head nodding associated with more severe facial clonus; 3—clonus of one forelimb; 4—rearing, often accompanied by bilateral forelimb clonus; and 5—rearing with loss of balance and falling accompanied by generalized clonic seizures. Seizure duration 1 is the duration of limbic (stage 1-2) and/or motor seizures (stage 3-5). Seizure duration 2 includes the time of limbic and/or motor seizures plus the adjacent time of immobility. Afterdischarge duration 1 (ADD 1) is the time of spikes in the EEG recorded from the site of stimulation with a frequency of at least 1/sec. Afterdischarge duration 2 (ADD 2) is the total time of spikes occurring in the EEG including those, which followed the ADD 1 with lower frequency and amplitude.

[0584] Test compound is administered to the prepared animals. Control experiments are performed 2-3 days before each test compound experiment. For control determinations, rats receive vehicle (e.g., saline) with the pretreatment time of the respective test compound experiment. For all test compound experiments, at least 4 days are interposed between successive administrations in order to avoid alterations in drug potency due to cumulation or tolerance. Data is analyzed using appropriate statistical methods.

[0585] In addition to recordings of anticonvulsant parameters, kindled rats can be observed for adverse effects in order to estimate a therapeutic index. Tests include open field observations, rotarod test, and body temperature. Tests used to evaluate adverse effects are performed in the same manner in control and test compound experiments at two different times, immediately before application of a test compound or vehicle and 13 min after application.

[0586] The rotarod test is carried out with a rod of 6 cm diameter and rotation speed of 8 rpm. Neurological deficit is indicated by inability of the animals to maintain their equilibrium for at least 1 min on the rotating rod. Rats are trained prior to the rotarod evaluation to maintain their balance on the

rod. After treatment with a test compound or vehicle, rats that are not able to maintain their equilibrium on the rod for three subsequent 1 min attempts are considered to exhibit neurological deficit.

[0587] In addition to these quantitative estimations of neurological deficit, behavioral alterations after administration of test compound are noted in the cage and after placing the animals in an open field of 90-100 cm diameter. Muscle tone is estimated by palpation of the abdomen. The extent of deficits in behavior after administration of a test compound is determined by a rating system. Animals are taken out of the cage, placed in an open field, observed for about 1 minute and rated separately for ataxia, abducted hindlimbs, reduced righting, flat body posture, circling, Straub tail, piloerection, hypolocomotion and hyperlocomotion (abdominal muscle tone is evaluated by palpation at the end of the period of observation). All other parameters except ataxia are scored from 0 to 3: 0—absent; 1—equivocal; 2—present; 3—intense. For ataxia: 1—slight ataxia in hind-legs (tottering of the hind quarters); 2—more pronounced ataxia with dragging of hind legs; 3—further increase of ataxia and more pronounced dragging of hind legs; 4—marked ataxia, animals lose balance during forward locomotion; 5—very marked ataxia with frequent loss of balance during forward locomotion; and 6—permanent loss of righting reflexes, but animal still attempts to move forward. Rectal body temperature is measured. Body weight of the animals is recorded once daily before a test compound is administered. Data is analyzed by an appropriate statistical method. The ability of a test compound to increase the electrical threshold for induction of afterdischarges, decrease the severity of seizures, reduce seizure duration, and reduce total afterdischarge duration suggests efficacy in treating migraine.

Description 18

Use of Clinical Trials to Assess the Efficacy of Acamprosate Prodrugs for Treating Migraine

[0588] The efficacy of a compound of Formula (I) in treating migraine may be assessed using a randomized, double blind, placebo-controlled, parallel group, clinical trial. The primary objective of the study is to evaluate the safety and efficacy of a test compound vs placebo in the treatment of recurrent episodes of migraine based on change from the baseline phase to the double-blind phase in the monthly (28 days) migraine episode rate. The secondary objectives are to evaluate the effect of treatment with a test compound versus placebo in migraine patients on percentage of subjects responding to treatment (50% or more reduction in monthly migraine episode rate) and change from the baseline phase to the double-blind phase in migraine days per month, average migraine duration, rescue medication use, average severity of migraine headache, average severity of migraine associated symptoms (nausea, vomiting, photophobia, phonophobia), to provide safety and efficacy data for the comparison a dose of a test compound in the treatment of migraine, and to evaluate the effect of treatment with a dose of a test compound versus placebo in migraine patients on migraine-specific measures of health-related quality of life (HRQL) and SF-36 quality-of-life measures, as well as the correlation between HRQL and migraine frequency.

[0589] The clinical trial is a randomized, double blind, placebo controlled, parallel-group, multicenter study to evaluate the efficacy and safety of one or more doses of a test

compound versus placebo in migraine prophylaxis. Patients are randomized into treatment groups. The patients must have been diagnosed with migraine for at least twelve months, with or without aura, as defined by the International Headache Society (IHS). The IHS diagnostic criteria differ from the definition of a migraine period utilized in this study for evaluation of efficacy. For the purposes of this study a migraine period is defined as the twenty-four hour period starting with the onset of painful migraine symptoms, or aura with successful abortive/rescue treatment. Any recurrence during the twenty-four hour period is considered part of the initial episode. If the migraine pain persists beyond the twenty-four hour period, for the purposes of this study, this is considered a new episode.

[0590] There are four phases in the clinical trial: Baseline, Core Double-Blind, Blinded Extension, and Taper/Exit. The Baseline Phase lasts up to 42 days and includes two periods: Washout and Prospective Baseline. At Baseline Visit 1 (screening), patients are evaluated to ensure that they meet inclusion/exclusion criteria. In addition, a three-month retrospective headache history is recorded. During each of the three months prior to Visit 1, patients should have had no more than 8 migraines and no more than 15 total headache days (migraine plus other headache types). Eligible patients then undergo other study procedures and are given a headache/rescue medication record. Patients maintain this record from Visit I throughout their participation in the clinical trial, documenting the occurrence of any headaches, or auras, as well as the duration, severity, and symptomatology of any migraine attacks. Patients also record the use of any abortive/rescue medication taken for the relief of migraine pain and associated symptoms, or during an aura to prevent migraine pain or relieve symptoms. In addition, for each migraine attack, patients answer the questions on the headache record regarding work loss and productivity. If at the start of the trial, eligible patients are on any prophylactic medication to treat their migraines, they enter a Washout Period of up to 14 days to taper from these medications. This washout is concluded by the time the patient enters the Prospective Baseline Period, 28 days prior to Visit 2 (randomization).

[0591] At Baseline Visit 2 (Day 1), headache/rescue medication record information is reviewed. To be eligible for randomization into the trial a patient must have had 3 to 12 migraine episodes but no greater than 15 (migraine and non-migraine), headache days during the 28 days prior to Visit 2.

[0592] In the Core Double-Blind Phase, patients who complete the Baseline Phase and meet the entry criteria (including Prospective Baseline Period migraine/headache rate) are randomized into treatment groups representing one or more doses of test compound or placebo. The Core Double-Blind Phase has two periods: Titration and Maintenance.

[0593] The Titration Period immediately follows the Baseline Phase and extends for eight weeks (56 days). During this period, patients randomized to test compound are started at an initial dose and the daily dose is increased weekly until the assigned dose is achieved (or maximum tolerated dose, whichever is less). From the third week of Titration until the end of the Maintenance Period, a maximum of two dose level reductions are permitted for unacceptable tolerability problems. If a patient is still in the Titration Period, after a dose reduction, rechallenge is attempted to approach the patient's assigned dose, and, if unsuccessful, the dose is reduced again to the original reduced dose. Patients who have already had their study medication dose decreased by two levels, and are

still experiencing unacceptable tolerability problems, which warrant additional dose reductions, exit the study, or enter the Open Label Extension Phase, where their dose is further adjusted. Clinic visits occur on, for example, Day 29 (Visit 3) and Day 57 (Visit 4/End of Titration).

[0594] During the 18-week Maintenance Period, patients remain on the dose of test compound reached at the end of the Titration Period (the assigned dose or the maximum tolerated dose). If a patient experiences unacceptable tolerability problems, the dose is reduced, but only to the point that there are no more than two dose reductions for the entire Core Phase (Titration plus Maintenance). No rechallenge is permitted during the Maintenance Period, so a patient continues on the reduced dose for the remainder of the period. Patients who have already had their study medication dose decreased by two levels, and are still experiencing unacceptable tolerability problems, which would warrant additional dose reductions, exit the study. Clinic visits occur, for example, on Day 83 (Visit 5), Day 113 (Visit 6), Day 141 (Visit 7) and Day 183 (Visit 8/Core Double-Blind Final Visit or Early Withdrawal).

[0595] Patients are considered to have completed the Core Double-Blind Phase if they complete all 26 weeks of the Phase (8 weeks of Titration and 18 weeks of Maintenance) without prematurely discontinuing study medication. Only patients who complete all 26 weeks of the Core Phase have the option of entering the Blinded Extension Phase.

[0596] During the Blinded Extension Phase, patients remain on test compound at the same dose they achieve during the Core Phase for six months, or until they withdraw. During this phase, patients are not permitted to adjust the dose of test compound. Patients are seen quarterly during this phase (Visits 10 and 11/Blinded Extension Final Visit). Patients are considered to have completed the Blinded Extension Phase if they complete all six months of the Phase without prematurely discontinuing the test compound.

[0597] In the Taper/Exit Phase, patients exiting the study are tapered from study medication. If a patient exits the study during the Core Double-Blind Phase (Titration or Maintenance Period), he or she is tapered from study medication in a blinded fashion. The length of the taper is as long as seven weeks, but varied according to the dose the patient achieves. Patients who exit the study during the Blinded Extension Phase are tapered from their medication following the recommended taper schedule.

[0598] Physical examinations (including height) and neurologic examinations are performed at the beginning and end of the study. A baseline electrocardiogram is performed at the beginning of the study. Vital signs and weight are recorded at each clinic visit. Adverse events are recorded. Quality of Life assessments are performed at intervals, for example, Visits 2 (Day 1), 4 (Day 57/Exit from Titration), 6 (Day 113) and 8 (Day 183/Core Double-Blind Final Visit/Early Withdrawal). Health Care Resource Use information is recorded at intervals, for example, Visits 3 through 8. The occurrence of any headaches or auras, severity and symptomatology of any migraine headaches, and the use of rescue medication is transcribed from a patient's headache record to their case record form at each visit.

[0599] Efficacy evaluations are based on information recorded on the subject's headache/rescue medication record and Health-Related Quality of Life assessments. On the headache/rescue medication record the patients documented the following throughout his/her study participation: occurrence and duration of headaches (and auras if no headache pain

develops), severity of migraine pain and associated symptoms, as well as the use of medication taken to relieve migraine pain or symptoms (or taken during an aura to relieve symptoms or prevent migraine pain). Health-Related Quality of Life (HRQL) assessments are completed at specified intervals throughout the study. The Migraine-Specific Quality of Life questionnaire (MSQ), and the Medical Outcomes Study Short Form-36 (SF-36) can be used to assess HRQL.

[0600] The primary efficacy criterion is the reduction in migraine episodes per month (28 days) during the Core Double-Blind Phase compared to the 28 day Prospective Baseline Period. Secondary efficacy criteria include the percentage of patients responding to treatment (50% or more reduction in the monthly (28 day) migraine episode rate) and reduction from the Prospective Baseline Period to the Core Double-Blind Phase in migraine days per month, monthly rate of all types of headaches, average migraine duration, rescue medication use, average severity of migraine headache, and average severity of migraine-associated symptoms (nausea, vomiting, photophobia, phonophobia). Also included in the secondary efficacy criteria is the effect of treatment with test compound versus placebo on migraine-specific measures of health-related quality of life (HRQL) and SF-36 quality-of-life measures, as well as the correlation between HRQL and migraine frequency. The Medical Outcomes Study Short Form-36 (SF-36) is the most frequently used generic measure of HRQL in migraine patients and has been used in studies of migraine. The SF-36 is a 36-item questionnaire measuring eight domains. The SF-36 has been shown to be reliable and valid in a wide variety of patient populations as well as for migraine patients. The migraine specific quality of life questionnaire (MSQ) can also be administered. The MSQ is a disease-specific instrument developed to assess quality of life relating to migraine. The MSQ has been used in published clinical trials of migraine therapy and has demonstrated evidence of reliability, validity, and responsiveness.

Description 19

Animal Model for Assessing Therapeutic Efficacy of Acamprosate Prodrugs for Treating Schizophrenia

Morris Water Maze

[0601] The Morris Water Maze (MWM) is used as a well-validated hippocampus dependent test of visual-spatial memory. The MWM tests the ability of an animal to locate a hidden platform submerged under water by using extra-maze cues from the test environment. Rats are trained in a pool 1.8 m in diameter and 0.6 µm high, containing water at about 26° C. A 10 cm square transparent platform is hidden in a constant position 1 cm below the water level in the pool. Only distal visuo-spatial cues are available to the rats for location of the submerged platform. The rats are given trials to find the hidden platform. The escape latency, i.e., the time required by the rats to find and climb onto the platform, is recorded for up to 120 s. Each rat is allowed to remain on the platform for 30 s, after which it is removed to its home cage. If the rat did not find the platform within 120 s, it is manually placed on the platform and returned to its home cage after 30 s.

[0602] Male Sprague-Dawley rats weighing 150-200 g are used. Ten days before the beginning of the experiments, the rats are handled once daily to reduce experimental stress. Acamprosate prodrug or control is administered to the rats for three consecutive days before behavioral testing. On each day of behavioral testing the rats are injected with either haloperidol or saline 30 min before behavioral assessment.

PCP-Induced Hyperactivity Model

[0603] Male C57Bl/6J mice are used. Mice are received at 6-weeks of age. Upon receipt, mice are assigned unique identification numbers (tail marked) and are group housed with 4 mice/cage in OPTI mouse ventilated cages. All animals remain housed in groups of four during the study. All mice are acclimated to the colony room for at least two weeks prior to testing and are subsequently tested at an average age of 8 weeks of age. During the period of acclimation, mice and rats are examined on a regular basis, handled, and weighed to assure adequate health and suitability.

[0604] Test compounds are prepared and administered according to the following procedures. An acamprosate prodrug is dissolved in sterile injectable water and administered i.p. at a dose volume of 10 mL/kg at 60 min prior to PCP injection. The amount of acamprosate prodrug administered can range, for example, from 0.01 mg/kg to 100 mg/kg. As a positive control, clozapine (1 mg/kg) is dissolved in 10% DMSO and administered i.p. at a dose volume of 10 mL/kg at 30 min prior to PCP injection. PCP (5 mg/kg) is dissolved in sterile injectable water and administered i.p. at a dose volume of 10 mL/kg.

[0605] The Open Field (OF) test is used to assess both anxiety and locomotor behavior. The open field chambers are Plexiglas square chambers (27.3×27.3×20.3 cm) surrounded by infrared photobeams (16×16×16) to measure horizontal and vertical activity. The analysis is configured to divide the open field into a center and periphery zone. Distance traveled is measured from horizontal beam breaks as a mouse moves, and rearing activity is measured from vertical beam breaks.

[0606] Mice are acclimated to the activity experimental room for at least 1 h prior to testing. Eight animals are tested in each run. Mice are injected with water or acamprosate prodrug, placed in holding cages for 30 min, and then in the OF chamber for 30 min, removed from the OF chamber and injected with either water or PCP and returned to the OF chambers for a 60-minute session. A different group of mice are injected with either 10% DMSO or clozapine and placed in the OF chamber for 30 min, removed from the OF chamber and injected with PCP (5 mg/kg), and returned to the OF chambers for a 60-minute session.

[0607] Data is analyzed by analysis of variance (ANOVA) followed by post-hoc comparisons with Fisher Tests when appropriate. Baseline activity is measured during the first 30 min of the test prior to PCP injection. PCP-induced activity is measured during the 60 min following PCP injection. Statistical outliers that fall above or below 2 standard deviations from the mean are removed from the final analysis. An effect is considered significant if $p < 0.05$.

Auditory Startle and Prepulse Inhibition of Startle (PPI)

[0608] Young, adult male C57Bl/6J mice are used in this study. Mice are received at 6-weeks of age. Upon receipt, mice are assigned unique identification numbers (tail marked) and are group housed in standard mouse cages. For testing, animals are randomly assigned across treatment groups and balanced by PPI chamber.

[0609] Acoustic startle measures an unconditioned reflex response to external auditory stimulation. PPI consisting of an inhibited startle response (reduction in amplitude) to an auditory stimulation following the presentation of a weak auditory stimulus or prepulse, has been used as a tool for the assessment of deficiencies in sensory-motor gating, such as

those seen in schizophrenia. Mice are placed in the PPI chamber (Med Associates) for a 5 min session of white noise (70 dB) habituation. A test session begins immediately after the 5 min acclimation period. The session starts with a habituation block of 6 presentations of the startle stimulus alone, followed by 10 PPI blocks of 6 different types of trials. Trial types are: null (no stimuli), startle (120 dB), startle plus prepulse (4, 8 and 12 dB over background noise i.e., 74, 78 or 82 dB) and prepulse alone (82 dB). Trial types are presented at random within each block. Each trial begins with a 50 ms null period during which baseline movements are recorded. There is a subsequent 20 ms period during which prepulse stimuli are presented and responses to the prepulse measured. Following a 100 ms pause, the startle stimuli are presented for 40 ms and responses are recorded for 100 ms from startle onset. Responses are sampled every ms. The inter-trial interval is variable with an average of 15 s (range from 10 to 20 s). In startle alone trials the basic auditory startle is measured and in prepulse plus startle trials the amount of inhibition of the normal startle is determined and expressed as a percentage of the basic startle response (from startle alone trials), excluding the startle response of the first habituation block.

[0610] For the normal mouse-PPI portion of the study, C57Bl/6J mice are treated with vehicle, haloperidol or acamprosate prodrug and placed back in their holding cages. Thirty min following administration of vehicle or haloperidol and 60 min following injection of vehicle or acamprosate prodrug, normal mouse-PPI testing commences.

[0611] For the PCP-PPI portion of the study, C57Bl/6J mice are treated with vehicle, clozapine, or acamprosate prodrug and returned to their holding cages. Thirty min later, all treatment groups are injected with vehicle or PCP. Thirty min following vehicle or PCP injection, PPI testing commences.

Description 20

Animal Model for Assessing Therapeutic Efficacy of Acamprosate Prodrugs for Treating Anxiety

[0612] The elevated plus-maze test can be used to assess the effects of test compounds on anxiety. A plus-maze consists of two open arms (50×10 cm) and two closed arms (50×10×40 cm). The arms extend from a central platform (10×10 cm) and are raised 50 cm. Each mouse is placed at the center of the maze facing a closed arm and is allowed to explore the maze for 5 min. The time spent in the open arms and the time spent in the closed arms is monitored, and the percent of time spent in the open arms determined. Increased time spent in the open arms indicates an anxiolytic effect for the test condition. A test that measures spontaneous locomotor activity such as measurement in an activity cage can be used to determine whether the test compound also affects locomotor activity. It is desirable that a compound exhibiting an anxiolytic effect not decrease locomotor activity.

Description 21

Animal Models of Depression

Forced Swim Test in Rats

[0613] Male Wistar rats weighting 230-270 g are acclimated to the colony room for a minimum of 1 week, handled daily for at least 4 days and habituated to saline injections for 2 days before the experiments.

[0614] Two glass cylinders (20 cm dia×40 cm height) are separated by black opaque partitions and filled with water at

about 24° C. to a depth of 30 cm. At this depth a rat cannot stand on the cylinder bottom. The water level is 10 cm from the top. Water is changed before each animal is placed into the water tank. An experimental session consists of two trials. During the conditioning trial, rats are gently placed into the cylinders for 15 min. After the trial, rats are dried and placed into a warm cage with the paper towels for 10-15 min before being returned to their home cages. Twenty-four hours later, for the test trial, animals are placed again into the cylinders for a 5-min test session. Tests are video taped for subsequent quantitative behavioral analysis. The frequency and/or total duration are calculated for each of the following categories: passive/immobile behavior (floating is scored when an animal remains in the water with all four limbs motionless, except for occasional alternate movements of paws and tail necessary to prevent sinking and to keep head/nose above the water); active/mobile behaviors (swimming characterized by rigorous movements with all four legs; paddling characterized by floating with rhythmical simultaneous kicks and occasional pushes off the wall to give speed and direction to the drift), including escape-oriented behaviors (climbing characterized by intense movements with all four limbs, with the two forepaws breaking the surface of the water and being directed against the walls of the cylinder; diving characterized by movements towards the bottom of the cylinder with the head of the rat below its hind limbs), and self-directed behaviors (headshakes, vigorous headshakes to get water off the snout and eyes; wiping, rubbing water away from the snout). In addition, at the end of each test trial, fecal boli are counted. A test compound, control, or positive control (e.g., imipramine) is administered prior to the test.

Tail Suspension Test in Mice

[0615] Mice are housed in standard laboratory cages and acclimated. Mice are moved from the housing room to the testing area in their home cages and allowed to adapt to the new environment for at least 1 h before testing. Immobility is induced by tail suspension. Mice are hung individually on a paper adhesive tape, 65 cm above a tabletop. Tape is placed approximately 1 cm from the tip of the tail. Animals are allowed to hang for 6 min and the duration of immobility is recorded. Mice are considered immobile only when hanging passively and completely motionless. Mice from these experiments are used one week later in locomotor activity studies. A test compound, control, or positive control (e.g., imipramine) is administered prior to the test.

Locomotor Activity

[0616] The spontaneous locomotor activity of mice is measured in photoresistor actometers (circular cages, 25 cm in dia, 15 cm high, two light sources, two photoresistors), in which the animals are placed individually 1 h after administration of a test compound. The number of crossings of light beams is measured during the first 30 min of the experimental session. The first measurement is performed 6 min after placing an animal into the actometer.

Description 22

Animal Model for Assessing Therapeutic Efficacy of Acamprosate Prodrugs for Treating Tardive Dyskinesia

[0617] Vacuous chewing movements (VCM) are a rodent model of TD. In this model, animals are treated chronically

with antipsychotics and their vacuous chewing motions are assessed by observation. The model has been shown to be sensitive to differential effects of typical and atypical antipsychotics and potential anti-dyskinetic agents.

[0618] Rats are housed in a controlled environment and allowed to acclimatize prior to testing. In order to limit neuroleptic-induced weight gain, food consumption is restricted to 15 g per animal per day. Rats are weighed biweekly throughout the study.

[0619] For two weeks prior to administration of test compound, animals are handled daily and habituated to the animal colony and the procedures related to drug administration and video recording. Subsequently (week 0), rats undergo a behavior video recording session following which they are randomized to a haloperidol treatment and a control group. The rats in the treatment group receive an intramuscular injection in the thigh muscles with haloperidol decanoate. The control rats are similarly injected with an equal volume of phosphate buffered saline (PBS). The haloperidol decanoate and saline injections are repeated every four weeks, for 20 weeks. Additional behavior video recording sessions are performed at weeks 12, 20 and 24 (i.e., 4 weeks after the last (fifth) injection). During the injection procedures, rats are handled with minimal restraint.

[0620] On the basis of the results of the behavior assessment performed 24 weeks after the first haloperidol injection (i.e., baseline day), the haloperidol-treated rats are assigned to 10 subject-each treatment groups having an equal mean frequency of observed VCM episodes. One week later (i.e., test day), the groups are randomized to receive either 0.5 mL PBS (vehicle) or acamprosate prodrug in 0.5 mL PBS. Rats undergo a 30-150 min video recorded behavior assessment session following administration. Two weeks after the test day (i.e., post-test day), the video recorded behavior assessment session is repeated to investigate longer-term effects of the experimental treatments.

[0621] The videotapes are scored. A VCM episode is defined as a bout of vertical deflections of the lower jaw, which may be accompanied by contractions of the jaw musculature.

Description 23

Animal Model for Assessing Therapeutic Efficacy of Acamprosate Prodrugs for Treating Spasticity

[0622] The mutant spastic mouse is a homozygous mouse that carries an autosomal recessive trait of genetic spasticity characterized by a deficit of glycine receptors throughout the central nervous system. The mouse is normal at birth and subsequently develops a coarse tremor, abnormal gait, skeletal muscle rigidity, and abnormal righting reflexes at two to three weeks of age. Assessment of spasticity in the mutant spastic mouse can be performed using electrophysiological measurements or by measuring the righting reflex (any righting reflex over one second is considered abnormal), tremor (holding mice by their tails and subjectively rating tremor), and flexibility.

[0623] Models of acute spasticity including the acute decerebrate rat, the acute or chronic spinally transected rat, and the chronically spinal cord-lesioned rat. The acute models, although valuable in elucidating the mechanisms involved in the development of spasticity, have come under criticism due to the fact that they are acute. The animals usually die or have total recovery from spasticity. The spas-

ticity develops immediately upon intervention, unlike the spasticity that evolves in the human condition of spasticity, which most often initially manifests itself as a flaccid paralysis. Only after weeks and months does spasticity develop in humans. Some of the more chronic-lesioned or spinally transected models of spasticity do postoperatively show flaccid paralysis. At approximately four weeks post-lesion/transection, the flaccidity changes to spasticity of variable severity. Although all of these models have their own particular disadvantages and lack of true representation of the human spastic condition, they are shown useful in developing treatments for spasticity in humans. Many of these models have also made use of different species, such as cats, dogs, and primates. Baclofen, diazepam, and tizanidine, effective antispastic agents in humans, are effective on different parameters of electrophysiologic assessment of muscle tone in these models.

[0624] The Irwin Test is used to detect physiological, behavioral, and toxic effects of a test substance, and indicates a range of dosages that can be used for later experiments. Typically, rats (three per group) are administered the test substance and are then observed in comparison with a control group given vehicle. Behavioral modifications, symptoms of neurotoxicity, pupil diameter, and rectal temperature are recorded according to a standardized observation grid derived from that of Irwin. The grid contains the following items: mortality, sedation, excitation, aggressiveness, Straub tail; writhes, convulsions, tremor, exophthalmos, salivation, lacrimation, piloerection, defecation, fear, traction, reactivity to touch, loss of righting reflexes, sleep, motor incoordination, muscle tone, stereotypes, head-weaving, catalepsy, grasping, ptosis, respiration, corneal reflex, analgesia, abnormal gait, forepaw treading, loss of balance, head twitches, rectal temperature, and pupil diameter. Observations are performed at 15, 30, 60, 120, and 180 minutes following administration of a test compound, and also 24 hours later.

[0625] In the Rotarod Test rats or mice are placed on a rod rotating at a speed of eight turns per minute. The number of animals that drop from the rod before three minutes is counted and the drop-off times are recorded (maximum: 180 sec). Diazepam, a benzodiazepine, can be administered at 8 mg/kg, i.p., as a reference substance.

Description 24

Animal Model for Assessing Therapeutic Efficacy of Acamprosate Prodrugs for Treating Multiple Sclerosis

[0626] Experiments are conducted on female C57BL/6 mice aged 4-6 weeks weighing 17-20 g. Experimental autoimmune encephalomyelitis (EAE) is actively induced using $\geq 95\%$ pure synthetic myelin oligodendrocyte glycoprotein peptide 35-55 (MOG35-55, MEVGWYRSPFSRV-VHLYRNGK). Each mouse is anesthetized and receives 200 μ g of MOG peptide and 15 μ g of Saponin extract from Quilija bark emulsified in 100 μ L of phosphate-buffered saline. A 25 μ L volume is injected subcutaneously over four flank areas. Mice are also intraperitoneally injected with 200 ng of pertussis toxin in 200 μ L of PBS. A second, identical injection of pertussis toxin is given after 48 h.

[0627] An acamprosate prodrug is administered at varying doses. Control animals receive 25 μ L of DMSO. Daily treatment extends from day 26 to day 36 post-immunization. Clinical scores are obtained daily from day 0 post-immunization

until day 60. Clinical signs are scored using the following protocol: 0, no detectable signs; 0.5, distal tail limpness, hunched appearance and quiet demeanor; 1, completely limp tail; 1.5, limp tail and hindlimb weakness (unsteady gait and poor grip with hindlimbs); 2, unilateral partial hindlimb paralysis; 2.5, bilateral hindlimb paralysis; 3, complete bilateral hindlimb paralysis; 3.5, complete hindlimb paralysis and unilateral forelimb paralysis; 4, total paralysis of hindlimbs and forelimbs.

[0628] Inflammation and demyelination are assessed by histology on sections from the CNS of EAE mice. Mice are sacrificed after 30 or 60 days and whole spinal cords are removed and placed in 0.32 M sucrose solution at 4° C. overnight. Tissues are prepared and sectioned. Luxol fast blue stain is used to observe areas of demyelination. Haematoxylin and eosin staining is used to highlight areas of inflammation by darkly staining the nuclei of mononuclear cells. Immune cells stained with H&E are counted in a blinded manner under a light microscope. Sections are separated into gray and white matter and each sector is counted manually before being combined to give a total for the section. T cells are immunolabelled with anti-CD3+ monoclonal antibody. After washing, sections are incubated with goat anti-rat HRP secondary antibody. Sections are then washed and counterstained with methyl green. Spenocytes isolated from mice at 30 and 60 days post-immunization are treated with lysis buffer to remove red blood cells. Cells are then resuspended in PBS and counted. Cells at a density of about 3×10^6 cells/mL are incubated overnight with 20 μ g/mL of MOG₃₅₋₅₅ peptide. Supernatants from stimulated cells are assayed for IFN- γ protein levels using an appropriate mouse IFN- γ immunoassay system.

Description 25

Animal Models of Pain

[0629] Inflammatory Pain—Formalin test

[0630] A formalin assessment test is used. Fifty μ L of a 5% formalin solution is injected subcutaneously into the dorsal aspect of the right hind paw and the rats are then individually placed into clear observation cages. Rats are observed for a continuous period of 60 min or for periods of time corresponding to phase I (from 0 to 10 min following formalin injection) and phase II (from 30 to 50 min following formalin injection) of the formalin test (Abbott et al., *Pain* 1995, 60, 91-102). The number of flinching behaviors of the injected paw is recorded using a sampling technique in which each animal is observed for one 60-sec period during each 5-min interval. Test compound is administered 30 min or other appropriate interval prior to formalin injection.

Inflammatory Pain—Carrageenan-Induced Acute Thermal Hyperalgesia and Edema

[0631] Paw edema and acute thermal hyperalgesia are induced by injecting 100 μ L of a 1% solution of λ -carrageenan in physiological saline into the plantar surface of the right hind paw. Thermal hyperalgesia is determined 2 h following carrageenan injection, using a thermal paw stimulator. Rats are placed into plastic cubicles mounted on a glass surface maintained at 30° C. and a thermal stimulus in the form of radiant heat emitted from a focused projection bulb is then applied to the plantar surface of each hind paw. The stimulus current is maintained at 4.50 ± 0.05 amp, and the maximum time of exposure is set at 20.48 sec to limit possible

tissue damage. The elapsed time until a brisk withdrawal of the hind paw from the thermal stimulus is recorded automatically using photodiode motion sensors. The right and left hind paw of each rat is tested in three sequential trials at about 5-min intervals. Carrageenan-induced thermal hyperalgesia of paw withdrawal latency ($PWL_{thermal}$) is calculated as the mean of the two shortest latencies. Test compound is administered 30 min before assessment of thermal hyperalgesia.

[0632] The volume of paw edema is measured using water displacement with a plethysmometer 2 h following carrageenan injection by submerging the paw up to the ankle hair-line (approx. 1.5 cm). The displacement of the volume is measured by a transducer and recorded. Test compound is administered at an appropriate time following carrageenan injection, such as for example, 30 min or 90 min.

Visceral Pain

[0633] Thirty min following administration of test compound, mice receive an injection of 0.6% acetic acid in sterile water (10 mL/kg, i.p.). Mice are then placed in table-top Plexiglass observation cylinders (60 cm high×40 cm diameter) and the number of constrictions/writhes (a wave of mild constriction and elongation passing caudally along the abdominal wall, accompanied by a slight twisting of the trunk and followed by bilateral extension of the hind limbs) is recorded during the 5-20 min following acetic acid injection for a continuous observation period of 15 min.

Neuropathic Pain—Spinal Nerve Ligation

[0634] Rats receive unilateral ligation of the lumbar 5 (L5) and lumbar 6 (L6) spinal nerves. The left L5 and L6 spinal nerves of the rat are isolated adjacent to the vertebral column and tightly ligated with a 5-0 silk suture distal to the dorsal root ganglia, and care is taken to avoid injury of the lumbar 4 (L4) spinal nerve. Control rats undergo the same procedure but without nerve ligation. All animals are allowed to recover for at least 1 week and not more than 3 weeks prior to assessment of mechanical allodynia. Mechanical allodynia is measured using calibrated von Frey filaments. Rats are placed into inverted plastic containers (20×12.5×20 cm) on top of a suspended wire mesh grid and acclimated to the test chamber for 20 min. The von Frey filaments are presented perpendicularly to the plantar surface of the selected hind paw, and then held in this position for approximately 8 s with enough force to cause a slight bend in the filament. Positive responses include an abrupt withdrawal of the hind paw from the stimulus or flinching behavior immediately following removal of the stimulus. A 50% paw withdrawal threshold (PWT) is determined. Rats with a $PWT \leq 5.0$ g are considered allodynic and utilized to test the analgesic activity of a test compound. The test compound can be administered 30 min prior to the assessment of mechanical allodynia.

Neuropathic Pain—Chronic Constriction Injury of the Sciatic Nerve

[0635] A model of chronic constriction injury of the sciatic nerve-induced neuropathic pain is used. The right common sciatic nerve is isolated at mid-thigh level and loosely ligated by four chromic gut (4-0) ties separated by an interval of 1 mm. Control rats undergo the same procedure but without sciatic nerve constriction. All animals are allowed to recover for at least 2 weeks and for no more than 5 weeks prior to testing of mechanical allodynia. Allodynic PWT is assessed

in the animals as described for animals with spinal nerve ligation. Only rats with a $PWT \leq 5.0$ g are considered allodynic and utilized to evaluate the analgesic activity of a test compound. Test compound is administered 30 min or other appropriate time prior to the assessment of mechanical allodynia.

Neuropathic Pain—Vincristine-induced Mechanical Allodynia

[0636] A model of chemotherapy-induced neuropathic pain is produced by continuous intravenous vincristine infusion (Nozaki-Taguchi et al., *Pain* 2001, 93, 69-76). Anesthetized rats undergo a surgical procedure in which the jugular vein is catheterized and a vincristine-primed pump is implanted subcutaneously. Fourteen days of intravenous infusion of vincristine (30 μ g/kg/day) results in systemic neuropathic pain of the animal. Control animals undergo the same surgical procedure, with physiological saline infusion. PWT of the left paw is assessed in the animals 14 days post-implantation as described for the spinal nerve ligation model. Test compound is administered 30 min prior to the test for mechanical allodynia in rats with $PWT \leq 5.00$ g before treatment.

Post-Operative Pain

[0637] A model of post-operative pain is performed in rats. The plantar aspect of the left hind paw is exposed through a hole in a sterile plastic drape, and a 1-cm longitudinal incision is made through the skin and fascia, starting 0.5 cm from the proximal edge of the heel and extending towards the toes. The plantaris muscle is elevated and incised longitudinally leaving the muscle origin and insertion points intact. After hemostasis by application of gently pressure, the skin is apposed with two mattress sutures using 5-0 nylon. Animals are then allowed to recover for 2 h following surgery, at which time mechanical allodynia and thermal hyperalgesia are assessed. **[0638]** Effects of test compound on mechanical allodynia are assessed 30 min following administration, with PWT being examined in these animals for both the injured and non-injured paw as described for the spinal nerve ligation model with the von Frey filament systematically pointing towards the medial side of the incision. In a separate experiment, the effects of test compound on thermal hyperalgesia are assessed 30 min following administration of test compound, with $PWL_{thermal}$ being determined as described for the carrageenan-induced thermal hyperalgesia model with the thermal stimulus applied to the center of the incision of the paw planter aspect.

Description 26

Animal Model for Assessing Therapeutic Efficacy of Acamprostate Prodrugs for Treating Binge Eating

[0639] Thirty 2-month old male Sprague Dawley rats are individually housed in a temperature- and humidity-controlled vivarium under a 12:12 light:dark cycle. Three days after being introduced into the vivarium, rats are given overnight access to a bowl of vegetable shortening. The rats are then divided into three groups of ten matched for two-day average chow intake, overnight shortening intake, and body weight.

[0640] The groups and different test phases are designed to test the effects of acamprostate prodrug under different short-

ening access conditions. In phase 1, rats maintained on a feeding protocol that promotes infrequent, large binges (B group) are compared to rats maintained on feeding protocols that promote no binges (FM and C groups). In phase 2, rats maintained for an extended period of time on the infrequent, large binge protocol (B group) are compared to rats that have just started the same binge protocol (FM and C groups). In phase 3, rats maintained on the feeding protocol that promotes infrequent, large binges (B group) are compared to rats on a feeding protocol that promotes more frequent, smaller binges (FM and C groups).

[0641] The three groups are maintained as follows: Binge (B): The (B) rats have continuous access to chow and water. In addition, they are given 2-h access to a separate bowl of vegetable shortening every Monday, Wednesday, and Friday (MWF), during the 2 h prior to no light. During the 2-h shortening access period, the chow and water remain available. This protocol results in infrequent, large episodes of binge-type eating in male rats. This protocol models the intermittent excessive eating behavior that characterizes binge eating. The B rats are maintained on this protocol throughout all phases of the study. Fat-Matched (FM): The rats in group FM are given the same proportions of chow and shortening that the Binge (B) groups consume except that the shortening is mixed into the chow, which is provided continuously. The proportions of chow and shortening consumed by the Binge group each week are determined, and the FM group is provided with a fat-matched chow mixed to that proportion the following week. The FM group has free access to the FM chow and water. The FM group is included to control for possible neural or behavioral effects of dietary fat. The FM group is maintained on the FM chow throughout all phases of the study. During phase 1, the FM group only has access to the FM chow. During phase 2, the FM group has access to a separate bowl of vegetable shortening for 2-h on MWF each week, in addition to the continuously available FM chow. During phase 3, the FM group has 2-h access to the vegetable shortening every day, in addition to the continuously available FM chow. This daily protocol results in more frequent, smaller episodes of binge-type eating. Chow/change (C): The rats in group C have continuous access to the regular chow and water through all phases of the study. During the first phase, the C group only has access to the regular chow diet. During the second phase, the C group has access to a separate bowl of vegetable shortening for 2-h on MWF each week in addition to the continuously available regular chow. During the third phase, the C group has 2-h access to the vegetable shortening every day in addition to the continuously available regular chow.

[0642] The effects of acamprosate prodrugs effects are determined during each of the three phases of the study. In phase 1, the effects of acamprosate prodrug are determined on binge-type consumption of vegetable shortening and on consumption of the regular and FM chow diets. Rats are on their respective diets for about 6 weeks prior to the initiation of acamprosate prodrug testing. In phase 2, the effects of acamprosate prodrug are assessed in rats that are bingeing for a relatively long (B group: three months) or short (FM and C groups: 1 day) period of time (all groups have MWF 2-h access to shortening in addition to their assigned regular or FM chow). In phase 3, the effects of acamprosate prodrug are assessed under conditions of infrequent (B: 2-h MWF) and more frequent (FM and C groups: 2-h daily) shortening access. The FM and C rats are on the daily shortening access


schedule for ten days before the first acamprosate prodrug administration in phase 3. Acamprosate prodrug is not tested in rats with continuous access to a bowl of shortening due to the low 2-h intakes that are generated on that protocol under non-food-deprived conditions. A dose and regimen of acamprosate prodrug is administered as appropriate for the objectives of the study.

[0643] Acamprosate prodrug is administered at an appropriate time prior to the shortening access period. Chow is removed during the 30-min pretreatment period. Shortening and/or chow are weighted and placed into the cage at the beginning of the test period, e.g., 2-h, and then re-weighted at the end of the test period. The data is analyzed using appropriate statistical methods.

[0644] Finally, it should be noted that there are alternative ways of implementing the embodiments disclosed herein. Accordingly, the present embodiments are to be considered as illustrative and not restrictive, and the claims are not to be limited to the details given herein, but may be modified within the scope and equivalents thereof.

What is claimed is:

1. A compound of Formula (I):



or a pharmaceutically acceptable salt thereof, wherein:

n is chosen from 0, 1, 2, and 3;

R¹ is chosen from C₁₋₈ alkyl, substituted C₁₋₈ alkyl, C₆₋₁₀ aryl, substituted C₆₋₁₀ aryl, C₃₋₈ cycloalkyl, substituted C₃₋₈ cycloalkyl, C₇₋₁₈ arylalkyl, substituted C₇₋₁₈ arylalkyl, C₄₋₁₆ cycloalkylalkyl, substituted C₄₋₁₆ cycloalkylalkyl, C₁₋₈ heteroalkyl, substituted C₁₋₈ heteroalkyl, C₅₋₁₀ heteroaryl, substituted C₅₋₁₀ heteroaryl, C₃₋₈ heterocycloalkyl, substituted C₃₋₈ heterocycloalkyl, C₆₋₁₈ heteroarylalkyl, substituted C₆₋₁₈ heteroarylalkyl, C₄₋₁₆ heterocycloalkylalkyl, substituted C₄₋₁₆ heterocycloalkylalkyl, -C(R⁶)₂OC(O)R⁵, -C(R⁶)₂OC(O)OR⁵, -(CH)_rN(R⁶)₂, and (CH₂)_rC(O)N(R⁶)₂, wherein: r is chosen from 1, 2 and 3;

each R⁵ is independently chosen from C₁₋₆ alkyl, substituted C₁₋₆ alkyl, C₆₋₈ aryl, substituted C₆₋₈ aryl, C₃₋₈ cycloalkyl, substituted C₃₋₈ cycloalkyl, C₁₋₆ heteroalkyl, substituted C₁₋₆ heteroalkyl, C₅₋₈ heteroaryl, substituted C₅₋₈ heteroaryl, C₃₋₈ heterocycloalkyl, and substituted C₃₋₈ heterocycloalkyl; and

each R⁶ is independently chosen from hydrogen, C₁₋₈ alkyl, substituted C₁₋₈ alkyl, C₆₋₈ aryl, substituted C₆₋₈ aryl, C₃₋₈ cycloalkyl, substituted C₃₋₈ cycloalkyl, and -COOR³³; wherein R³³ is chosen from hydrogen and C₁₋₈ alkyl;

R² and R³ are independently chosen from C₁₋₄ alkyl, substituted C₁₋₄ alkyl, C₁₋₄ alkoxy, substituted C₁₋₄ alkoxy, -NH₂, -NHC(O)R³⁴, -NHR³⁴ and -N(R³⁴)₂; or R² and R³ together with the carbon to which they are bonded form a ring chosen from a C₃₋₈ cycloalkyl, substituted C₃₋₈ cycloalkyl, C₃₋₈ heterocycloalkyl, and substituted C₃₋₈ heterocycloalkyl ring; and

- each R^4 is independently chosen from hydrogen, halogen, $-\text{OR}^7$, $-\text{CN}$, $-\text{CF}_3$, $=\text{O}$, $-\text{NO}_2$, $-\text{COOR}^{33}$, C_{1-8} alkyl, substituted C_{1-8} alkyl, C_{6-10} aryl, substituted C_{6-10} aryl, C_{3-8} cycloalkyl, substituted C_{3-8} cycloalkyl, C_{7-18} arylalkyl, substituted C_{7-18} arylalkyl, C_{4-16} cycloalkylalkyl, substituted C_{4-16} cycloalkylalkyl, C_{1-8} heteroalkyl, substituted C_{1-8} heteroalkyl, C_{5-10} heteroaryl, substituted C_{5-10} heteroaryl, C_{3-8} heterocycloalkyl, substituted C_{3-8} heterocycloalkyl, C_{6-18} heteroarylalkyl, substituted C_{6-18} heteroarylalkyl, C_{4-16} heterocycloalkylalkyl, substituted C_{4-16} heterocycloalkylalkyl, $-\text{NH}_2$, $-\text{NHC}(\text{O})\text{R}^{34}$, $-\text{NHR}^{34}$, and $-\text{N}(\text{R}^{34})_2$; wherein:
- R^7 is chosen from hydrogen, C_{1-8} alkyl, substituted C_{1-8} alkyl, C_{6-10} aryl, substituted C_{6-10} aryl, C_{3-8} cycloalkyl, substituted C_{3-8} cycloalkyl, $-\text{C}(\text{O})\text{R}^8$, $-\text{C}(\text{O})\text{OR}^8$, $-\text{C}(\text{O})\text{N}(\text{R}^9)_2$, $-\text{C}(\text{O})\text{SR}^8$, $-\text{C}(\text{R}^9)_2\text{OC}(\text{O})\text{R}^8$, $-\text{C}(\text{R}^9)_2\text{OC}(\text{O})\text{OR}^8$, $-\text{P}(\text{O})(\text{OH})_2$, and $-\text{SO}_2\text{OH}$; wherein:
- each R^8 is independently chosen from C_{1-6} alkyl, substituted C_{1-6} alkyl, C_{6-8} aryl, substituted C_{6-8} aryl, C_{3-8} cycloalkyl, substituted C_{3-8} cycloalkyl, C_{1-6} heteroalkyl, substituted C_{1-6} heteroalkyl, C_{5-8} heteroaryl, substituted C_{5-8} heteroaryl, C_{3-8} heterocycloalkyl, and substituted C_{3-8} heterocycloalkyl; and
- each R^9 is independently chosen from hydrogen, C_{1-8} alkyl, substituted C_{1-8} alkyl, C_{6-8} aryl, substituted C_{6-8} aryl, C_{3-8} cycloalkyl, substituted C_{3-8} cycloalkyl, and $-\text{COOR}^{35}$; wherein R^{35} is chosen from hydrogen and C_{1-8} alkyl;
- R^{33} is chosen from hydrogen and C_{1-4} alkyl;
- each R^{34} is independently chosen from hydrogen, C_{1-8} alkyl, substituted C_{1-8} alkyl, C_{1-8} alkoxy, substituted C_{1-8} alkoxy, C_{6-10} aryl, substituted C_{6-10} aryl, C_{6-10} aryloxy, substituted C_{6-10} aryloxy, C_{3-8} cycloalkyl, substituted C_{3-8} cycloalkyl, C_{3-8} cycloalkoxy, substituted C_{3-8} cycloalkoxy, C_{7-18} arylalkyl, substituted C_{7-18} arylalkyl, C_{4-16} cycloalkylalkyl, substituted C_{4-16} cycloalkylalkyl, C_{1-8} heteroalkyl, substituted C_{1-8} heteroalkyl, C_{5-10} heteroaryl, substituted C_{5-10} heteroaryl, C_{3-8} heterocycloalkyl, substituted C_{3-8} heterocycloalkyl, C_{6-18} heteroarylalkyl, substituted C_{6-18} heteroarylalkyl, C_{4-16} heterocycloalkylalkyl, and substituted C_{4-16} heterocycloalkylalkyl; and
- with the proviso that when n is 1, and each of R^2 and R^3 is methyl; then R^4 is not chosen from $-\text{OR}^7$, $-\text{COOR}^{33}$, and substituted heteroalkyl.
- The compound of claim 1, wherein each substituent is independently chosen from halogen, $-\text{OH}$, $-\text{CN}$, $-\text{CF}_3$, $-\text{OCF}_3$, $=\text{O}$, $-\text{NO}_2$, C_{1-4} alkoxy, C_{1-4} alkyl, $-\text{COOR}^{30}$, $-\text{NR}^{31}_2$, and $-\text{CONR}^{32}_2$; wherein: each R^{30} , R^{31} , and R^{32} is independently chosen from hydrogen and C_{1-4} alkyl.
 - The compound of claim 1, wherein n is chosen from 0, 1, and 2.
 - The compound of claim 1, wherein R^1 is chosen from C_{1-6} alkyl, substituted C_{1-6} alkyl, C_{3-8} cycloalkyl, substituted C_{3-8} cycloalkyl, C_{7-12} arylalkyl, substituted C_{7-12} arylalkyl, C_{1-6} heteroalkyl, substituted C_{1-6} heteroalkyl, C_{3-8} heterocycloalkyl, substituted C_{3-8} heterocycloalkyl, C_{6-8} heteroarylalkyl, and substituted C_{6-8} heteroarylalkyl.
 - The compound of claim 4, wherein R^1 is chosen from C_{1-6} alkyl, benzyl, substituted benzyl, morpholinyl- N -ethyl, pyridyl-3-methyl, ethoxycarbonyloxyethyl, and methylbenzyl.
 - The compound of claim 1, wherein R^1 is chosen from $-\text{C}(\text{R}^6)_2\text{OC}(\text{O})\text{R}^5$ and $-\text{C}(\text{R}^6)_2\text{OC}(\text{O})\text{OR}^5$.
 - The compound of claim 6, wherein: R^5 is chosen from methyl, ethyl, n -propyl, isopropyl, n -butyl, isobutyl, tert-butyl, sec-butyl, pentyl, cyclohexyl, phenyl, and 3-pyridyl; and one of R^6 is hydrogen and the other of R^6 is chosen from hydrogen, methyl, ethyl, n -propyl, isopropyl, cyclohexyl, and phenyl; or each of R^6 is methyl.
 - The compound of claim 7, wherein: each of R^2 and R^3 is methyl; and each R^4 is hydrogen.
 - The compound of claim 1, wherein R^2 and R^3 are independently chosen from C_{1-4} alkyl, substituted C_{1-4} alkyl, C_{1-4} alkoxy, substituted C_{1-4} alkoxy, and $-\text{NH}_2$.
 - The compound of claim 9, wherein each of R^2 and R^3 is methyl.
 - The compound of claim 1, wherein each R^4 is hydrogen.
 - The compound of claim 1, wherein the compound is chosen from:
 - methyl 3- $\{[3-(\text{acetylamino})\text{propyl}]\text{sulfonyloxy}\}$ -2,2-dimethylpropanoate;
 - phenylmethyl 3- $\{[3-(\text{acetylamino})\text{propyl}]\text{sulfonyloxy}\}$ -2,2-dimethylpropanoate;
 - phenylmethyl 5- $\{[3-(\text{acetylamino})\text{propyl}]\text{sulfonyloxy}\}$ -4,4-dimethylpentanoate;
 - phenylmethyl 4- $\{[3-(\text{acetylamino})\text{propyl}]\text{sulfonyloxy}\}$ -3,3-dimethylbutanoate;
 - phenylmethyl 3- $\{[3-(\text{acetylamino})\text{propyl}]\text{sulfonyloxy}\}$ -2- $\{[3-(\text{acetylamino})\text{propyl}]\text{sulfonyloxy}\}$ -methyl-2-methylpropanoate;
 - (4-methoxyphenyl)methyl-3- $\{[3-(\text{acetylamino})\text{propyl}]\text{sulfonyloxy}\}$ -2,2-dimethylpropanoate;
 - ethyl 3- $\{[3-(\text{acetylamino})\text{propyl}]\text{sulfonyloxy}\}$ -2,2-dimethylpropanoate;
 - methylethyl 3- $\{[3-(\text{acetylamino})\text{propyl}]\text{sulfonyloxy}\}$ -2,2-dimethylpropanoate;
 - 2'-morpholinylethyl 3- $\{[3-(\text{acetylamino})\text{propyl}]\text{sulfonyloxy}\}$ -2,2-dimethylpropanoate hydrochloride;
 - pyridinylmethyl 3- $\{[3-(\text{acetylamino})\text{propyl}]\text{sulfonyloxy}\}$ -2,2-dimethylpropanoate hydrochloride;
 - ethoxycarbonyloxyethyl 3- $\{[3-(\text{acetylamino})\text{propyl}]\text{sulfonyloxy}\}$ -2,2-dimethylpropanoate;
 - phenylmethyl 2- $\{[3-(\text{acetylamino})\text{propyl}]\text{sulfonyloxy}\}$ -methyl-3-hydroxy-2-methylpropanoate;
 - methyl 4- $\{[3-(\text{acetylamino})\text{propyl}]\text{sulfonyloxy}\}$ -3,3-dimethylbutanoate;
 - 3- $\{[3-(\text{acetylamino})\text{propyl}]\text{sulfonyloxy}\}$ -2,2-dimethylpropanoic acid;
 - 5- $\{[3-(\text{acetylamino})\text{propyl}]\text{sulfonyloxy}\}$ -4,4-dimethylpentanoic acid;
 - methylethoxycarbonyloxyethyl 3- $\{[3-(\text{acetylamino})\text{propyl}]\text{sulfonyloxy}\}$ -2,2-dimethylpropanoate;
 - 4- $\{[3-(\text{acetylamino})\text{propyl}]\text{sulfonyloxy}\}$ -3,3-dimethylbutanoic acid;
 - methyl 3- $\{[3-(\text{acetylamino})\text{propyl}]\text{sulfonyloxy}\}$ -2- $\{[3-(\text{acetylamino})\text{propyl}]\text{sulfonyloxy}\}$ -methyl-2-methylpropanoate;
 - methyl 3- $\{[3-(\text{acetylamino})\text{propyl}]\text{sulfonyloxy}\}$ -2-amino-2-methylpropanoate;
 - ethyl 3- $\{[3-(\text{acetylamino})\text{propyl}]\text{sulfonyloxy}\}$ -2,2-diethoxypropanoate;
 - benzyl 3- $\{[3-(\text{acetylamino})\text{propyl}]\text{sulfonyloxy}\}$ -2,2-diethoxypropanoate; and
 - a pharmaceutically acceptable salt of any of the foregoing.


13. The compound of claim 1, wherein the compound is chosen from:

phenylmethyl 3-{[3-(acetylamino)propyl]sulfonyloxy}-2,2-dimethylpropanoate;

ethyl 3-{[3-(acetylamino)propyl]sulfonyloxy}-2,2-dimethylpropanoate; and

a pharmaceutically acceptable salt of any of the foregoing.

14. A compound of Formula (II):



or a pharmaceutically acceptable salt thereof; wherein:

q is chosen from 0, 1, 2, and 3;

R^{22} and R^{23} are independently chosen from C_{1-4} alkyl, substituted C_{1-4} alkyl, C_{1-4} alkoxy, substituted C_{1-4} alkoxy, $-NH_2$, $-NHC(O)R^{34}$, $-NHR^{34}$, and $-NR^{34}_2$; or R^{22} and R^{23} together with the carbon to which they are bonded form a ring chosen from a C_{3-8} cycloalkyl, substituted C_{3-8} cycloalkyl, C_{3-8} heterocycloalkyl, and substituted C_{3-8} heterocycloalkyl ring; and each R^{24} is independently chosen from hydrogen, halogen, $-OR^7$, $-CN$, $-CF_3$, $=O$, $-NO_2$, $-COOR^{33}$, C_{1-8} alkyl, substituted C_{1-8} alkyl, C_{6-10} aryl, substituted C_{6-10} aryl, C_{3-8} cycloalkyl, substituted C_{3-8} cycloalkyl, C_{7-18} arylalkyl, substituted C_{7-18} arylalkyl, C_{4-16} cycloalkylalkyl, substituted C_{4-16} cycloalkylalkyl, C_{1-8} heteroalkyl, substituted C_{1-8} heteroalkyl, C_{5-10} heteroaryl, substituted C_{5-10} heteroaryl, C_{3-8} heterocycloalkyl, substituted C_{3-8} heterocycloalkyl, C_{6-18} heteroarylalkyl, substituted C_{6-18} heteroarylalkyl, C_{4-16} heterocycloalkylalkyl, substituted C_{4-16} heterocycloalkylalkyl, $-NH_2$, $-NHC(O)R^{34}$, $-NHR^{34}$, and $-N(R^{34})_2$; wherein:

R^7 is chosen from hydrogen, C_{1-8} alkyl, substituted C_{1-8} alkyl, C_{6-10} aryl, substituted C_{6-10} aryl, C_{3-8} cycloalkyl, substituted C_{3-8} cycloalkyl, $-C(O)R^8$, $-C(O)OR^8$, $-C(O)N(R^9)_2$, $-C(O)SR^8$, $-C(R^9)_2OC(O)R^8$, and $-C(R^9)_2OC(O)OR^8$; wherein:

each R^8 is independently chosen from C_{1-6} alkyl, substituted C_{1-6} alkyl, C_{6-8} aryl, substituted C_{6-8} aryl, C_{3-8} cycloalkyl, substituted C_{3-8} cycloalkyl, C_{1-6} heteroalkyl, substituted C_{1-6} heteroalkyl, C_{5-8} heteroaryl, substituted C_{5-8} heteroaryl, C_{3-8} heterocycloalkyl, and substituted C_{3-8} heterocycloalkyl;

each R^9 is independently chosen from hydrogen, C_{1-8} alkyl, substituted C_{1-8} alkyl, C_{6-8} aryl, substituted C_{6-8} aryl, C_{3-8} cycloalkyl, substituted C_{3-8} cycloalkyl, and $-COOR^{35}$; wherein R^{35} is chosen from hydrogen and C_{1-8} alkyl;

R^{33} is chosen from hydrogen and C_{1-4} alkyl;

each R^{34} is independently chosen from hydrogen, C_{1-8} alkyl, substituted C_{1-8} alkyl, C_{1-8} alkoxy, substituted C_{1-8} alkoxy, C_{6-10} aryl, substituted C_{6-10} aryl, C_{6-10} aryloxy, substituted C_{6-10} aryloxy, C_{3-8} cycloalkyl, substituted C_{3-8} cycloalkyl, C_{3-8} cycloalkoxy, substituted C_{3-8} cycloalkoxy, C_{7-18} arylalkyl, substituted C_{7-18} arylalkyl, C_{4-16} cycloalkylalkyl, substituted C_{4-16} cycloalkylalkyl, C_{1-8} heteroalkyl, substituted

C_{1-8} heteroalkyl, C_{5-10} heteroaryl, substituted C_{5-10} heteroaryl, C_{3-8} heterocycloalkyl, substituted C_{3-8} heterocycloalkyl, C_{6-18} heteroarylalkyl, substituted C_{6-18} heteroarylalkyl, C_{4-16} heterocycloalkylalkyl, and substituted C_{4-16} heterocycloalkylalkyl; and with the proviso that when n is 1, and each of R^{22} and R^{23} is methyl; then R^{24} is not chosen from OR^7 , $-COOR^{33}$, and substituted heteroalkyl.

15. The compound of claim 14, wherein each substituent is independently chosen from halogen, $-OH$, $-CN$, $-CF_3$, $-OCF_3$, $=O$, $-NO_2$, C_{1-4} alkoxy, C_{1-4} alkyl, $-COOR^{30}$, $-NR^{31}_2$, and $-CONR^{32}_2$; wherein: each R^{30} , R^{31} , and R^{32} is independently chosen from hydrogen and C_{1-4} alkyl.

16. The compound of claim 14, wherein q is chosen from 0, 1, and 2.

17. The compound of claim 14, wherein R^{22} and R^{23} are independently chosen from C_{1-4} alkyl and C_{1-4} alkoxy.

18. The compound of claim 17, wherein each of R^{22} and R^{23} is methyl.

19. The compound of any one of claims 14 and 18, wherein each R^{24} is hydrogen.

20. The compound of claim 14, wherein the compound is chosen from:

3-{[3-(acetylamino)propyl]sulfonyloxy}-2,2-dimethylpropanoic acid;

4-{[3-(acetylamino)propyl]sulfonyloxy}-3,3-dimethylbutanoic acid;

5-{[3-(acetylamino)propyl]sulfonyloxy}-4,4-dimethylpentanoic acid; and

a pharmaceutically acceptable salt of any of the foregoing.

21. A pharmaceutical composition comprising a compound of claim 1 and at least one pharmaceutically acceptable vehicle.

22. The pharmaceutical composition of claim 21, comprising an amount of the compound effective for the treatment of a disease chosen from a neurodegenerative disorder, a psychotic disorder, a mood disorder, an anxiety disorder, a somatoform disorder, a movement disorder, a substance abuse disorder, binge eating disorder, a cortical spreading depression related disorder, a sleeping disorder, tinnitus, multiple sclerosis, and pain.

23. The pharmaceutical composition of claim 21, wherein the pharmaceutical composition is a sustained release oral dosage formulation.

24. A method of treating a disease in a patient comprising administering to a patient in need of such treatment the pharmaceutical composition of claim 21, wherein the disease is chosen from a neurodegenerative disorder, a psychotic disorder, a mood disorder, an anxiety disorder, a somatoform disorder, a movement disorder, a substance abuse disorder, binge eating disorder, a cortical spreading depression related disorder, a sleeping disorder, tinnitus, multiple sclerosis, and pain.

25. A method of treating a disease in a patient comprising administering to a patient in need of such treatment the compound of claim 1, wherein the disease is chosen from a neurodegenerative disorder, a psychotic disorder, a mood disorder, an anxiety disorder, a somatoform disorder, a movement disorder, a substance abuse disorder, binge eating disorder, a cortical spreading depression related disorder, a sleeping disorder, tinnitus, multiple sclerosis, and pain.

* * * * *