

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
9 November 2006 (09.11.2006)

PCT

(10) International Publication Number
WO 2006/117664 A1

(51) International Patent Classification:

C11C 3/00 (2006.01) *A61P 3/10* (2006.01)
A61K 31/202 (2006.01) *A61P 9/10* (2006.01)
A61K 31/232 (2006.01) *C07C 57/03* (2006.01)
A61K 31/421 (2006.01) *C07C 69/587* (2006.01)
A61P 3/04 (2006.01) *C07D 263/24* (2006.01)

(21) International Application Number:

PCT/IB2006/001155

(22) International Filing Date: 4 May 2006 (04.05.2006)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/677,350 4 May 2005 (04.05.2005) US
0501044-2 4 May 2005 (04.05.2005) SE
0501045-9 4 May 2005 (04.05.2005) SE
60/677,351 4 May 2005 (04.05.2005) US

(71) Applicant (for all designated States except US):
PRONOVA BIOCARE AS [NO/NO]; P.O. Box 420,
N-1327 Lysaker (NO).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **BRYHN,**
Morten [NO/NO]; Storgaten 43, N-3060 Svelvik (NO).
HOLMEIDE, Anne, Kristin [NO/NO]; Orionveien 12,
N-0489 Oslo (NO). **KOPECKY, Jan** [CZ/CZ]; Rozsirena
23, 182 00 Praha 8 (CZ).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declaration under Rule 4.17:


— as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: NEW DHA DERIVATIVES AND THEIR USE AS MEDICAMENTS



(I)


(57) Abstract: Compounds of formula (I); wherein - R₁ and R₂ are the same or different and may be selected from the group consisting of a hydrogen atom, a hydroxy group, an alkyl group, a halogen atom, an alkoxy group, an acyloxy group, an acyl group, an alkenyl group, an alkynyl group, an aryl group, an alkylthio group, an alkoxycarbonyl group, an alkylsulfmlyl group, an alkylsulfonyl group, an amino group, and an alkylamino group; and - X represents a carboxylic acid group, a carboxylate group, or a carboxamide group; or any pharmaceutically acceptable salt, solvate, complex or

pro-drug thereof, with the provisos that the compound of formula (I) is not (all-Z)-4,7,10,13,16,19- docosahexaenoic acid (DHA), alpha-methyl DHA, alpha-methyl DHA methyl ester, alpha-methyl DHA ethyl ester or alpha-hydroxy DHA ethyl ester, are disclosed. A fatty acid composition and a pharmaceutical composition comprising such compounds are also disclosed. The use of such compounds as medicaments, in particular for the treatment of diabetes type 2, is also disclosed.

New DHA derivatives and their use as medicaments

Technical field

The present invention relates to compounds of the general formula (I):



- 5 and their use as medicaments, in particular for the treatment of diabetes mellitus, type 2, and pre-stages thereof. It also relates to a pharmaceutical composition comprising compounds of formula (I), as well as to a fatty acid composition comprising compounds of formula (I).

10 **Background of the invention**

- The increasing incidence of type 2 diabetes mellitus worldwide poses an immense public health and medical challenge for the implementation of successful preventive and treatment strategies. The concurrent rise in overweight and obesity, which is tightly correlated to type 2 diabetes, interferes with diabetes treatment and increases the likelihood of hypertension, dyslipidemia, and atherosclerosis related diseases.

- The pathophysiologic condition precluding the development of type 2 diabetes is related to reduced effects of insulin on peripheral tissues, called insulin resistance. These tissues are mainly muscle, fat and liver. Muscle tissue is the main tissue concerned by insulin resistance in type 2 diabetes. The syndrome characterised by insulin resistance, hypertension, dyslipidemia and a systemic proinflammatory state, is referred to as metabolic syndrome. The prevalence of metabolic syndrome in the adult population in developed countries is 22-39% (Meighs 2003)

- Currently the most promising approach to mitigate and deter the metabolic syndrome is lifestyle intervention with weight reduction, decreased consumption of saturated fat, increased physical activity in combination with appropriate pharmacotherapy. Healthy diets that avoid excess energy intake encompass substitution of mono and polyunsaturated fatty acids in exchange for saturated fat. In particular the long-chain omega-3 fatty acids from fatty fish, namely

eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA) have proven beneficial in prevention of type 2 diabetes.

EPA and DHA have effects on diverse physiological processes impacting normal health and chronic disease, such as the regulation of plasma lipid levels, cardiovascular and immune function, insulin action and neural development and visual function. Firm evidence exist for their beneficial role in the prevention and management of coronary heart disease, dyslipidemias, type 2 diabetes, insulin resistance, and hypertension (Simonopoulos 1999; Geleijnse 2002; Storlien 1998).

Recent studies suggest that omega-3 fatty acids serve as important mediators of gene expression, working via nuclear receptors like the peroxisome proliferator-activated receptors (PPARs) controlling the expression of the genes involved in the lipid and glucose metabolism and adipogenesis (Jump 2002). PPARs are nuclear fatty acid receptors that have been implicated to play an important role in obesity-related metabolic diseases such as hyperlipidemia, insulin resistance, and coronary heart disease.

The three subtypes, α , γ , and δ , have distinct expression pattern and evolved to sense components of different lipoproteins and regulate lipid homeostasis based on the need of a specific tissue. PPAR α potentiates fatty acid catabolism in the liver and is the molecular target of the lipid-lowering fibrates. PPAR γ on the other hand is essential for adipocyte differentiation and mediates the activity of the insulin-sensitizing thiazolidinediones (the glitazones) through mechanisms not fully understood. (Chih-Hao 2003; Yki-Järvinen 2004)

Recently, pharmaceuticals acting as ligands to the PPAR γ receptor have been introduced as treatment of type 2 diabetes (Yki-Järvinen 2004). These compounds called thiazolidinediones or glitazones are drugs that reverse insulin resistance which is the pathophysiologic basis for development of the metabolic syndrome and type 2 diabetes. These compounds, of which rosiglitazone and pioglitazone have been launched as pharmaceuticals, lower fasting and postprandial glucose concentrations (which is being manifest as a pathologic glucose tolerance test), plasma insulin as well as free fatty acid concentrations. In this respect the glitazones act as insulin sensitizers.

However, these improvements are generally accompanied by weight gain and an increase in the subcutaneous adipose-tissue mass (Adams 1997). The use of thiazolidinediones is not only associated with weight gain but a subgroup of patients also have fluid retention and plasma volume expansion, leading to peripheral oedema. The increase in body weight and oedema has been associated with an increase in the incidence of heart failure, which is the reason why the Food and Drug Administration

has included a warning in the prescription information for rosiglitazone (provided by Avandia) and pioglitazone (provided by Takeda). These adverse effects restrict the use of the glitazones especially in patients with coronary heart conditions. Clearly there is a potential for new drugs with positive effects on insulin resistance but with weight reduction activity and no fluid retention tendency.

The effect of the poly-unsaturated fatty acids (PUFAs) on PPARs are not only a result of fatty acid structure and affinity to the receptor. Factors contributing to the composition of the intracellular non-esterified fatty acids (NEFA) levels are also important. This NEFA pool is affected by the concentration of exogenous fatty acids entering the cell and the amount of endogenous synthesised fatty acids, their removal via incorporation into lipids as well as their oxidation pathways. (Pawar 2003)

Although omega-3 fatty acids are weak agonists of PPARs, when compared with pharmacological agonists like the thiolglitazones, these fatty acids have demonstrated improvement in glucose uptake and insulin sensitivity (Storlien 1987). It has been reported that adipocytes were more insulin sensitive and transported more glucose when the polyunsaturated to saturated fatty acid ratio in the diet was increased (Field 1990). Collectively, these data indicate that the 20- and 22-carbon fatty acids, namely EPA and DHA could play a preventive role in the development of insulin resistance.

Due to their limited stability in vivo and their lack of biological specificity, PUFAs have not achieved widespread use as therapeutic agents. Chemical modifications of the n-3 polyunsaturated fatty acids have been performed by several research groups in order to change or increase their metabolic effects.


For example, the hypolipidemic effects of EPA was potentiated by introducing methyl or ethyl in α - or β -position of EPA. (Vaagenes 1999). The compounds also reduced plasma free fatty acid while EPA EE had no effect.

In a recent work published by L. Larsen (Larsen 2005) the authors show that the α -methyl derivatives of EPA and DHA increased the activation of the nuclear receptor PPAR α and thereby the expression of L-FABP compared to EPA/DHA. EPA with an ethyl group in the α -position activated PPAR α with equal strength as α -methyl EPA. The authors suggest that delayed catabolism of these α -methyl FA may contribute to their increased effects due to decreased β -oxidation in mitochondria leading to peroxisomal oxidation.

Alpha-methyl EPA has been shown to be a stronger inhibitor of platelet aggregation than EPA, both in vitro (Larsen 1998) and in vivo (Willumsen 1998).

Patent Abstract of Japan, publication number 05-00974 discloses DHA substituted in alpha-position with an OH-group, however only as an intermediate. No examination as to possible pharmaceutical effects of this compound is disclosed.

Laxdale Limited has also described the use of alpha substituted derivatives of EPA in the treatment of psychiatric or central nervous disorders (US6689812).



(A) α -methyl EPA

None of these modified fatty acids have, however, shown satisfactory pharmaceutical activity, and none of them has reached the pharmaceutical market.

Summary of the invention


The aim of the present invention is to provide new DHA-derivatives having therapeutical activity.

Based on the present invention a number of aspects are presented in the appended claims. Some of these aspects are;

1. Novel compounds, i.e. certain α -substituted polyunsaturated fatty acid derivatives.
2. The novel compounds for use as a medicament and for use in therapy.
3. A fatty acid composition or a pharmaceutical composition comprising the novel compounds.
4. A fatty acid composition comprising the novel compounds for use as a medicament and for use in therapy.
5. Use of the novel compounds for the production of a medicament for the prevention and/or treatment of diabetes in humans or an animal.
6. Use of the novel compounds for the production of a medicament for the treatment and/or the prevention of obesity or an overweight condition.
7. Use of the novel compounds for the production of a medicament for controlling body weight reduction and/or for preventing body weight gain.
8. Use of the novel compounds for the production of a medicament for the treatment and/or prevention of amyloidosis-related diseases.
9. Use of the novel compounds for the production of a medicament for the treatment or prophylaxis of multiple risk factors or cardiovascular diseases.

10. Use of the novel compounds for the production of a medicament for the prevention of stroke, cerebral or transient ischaemic attacks related to atherosclerosis of several arteries.
11. A method for specific treatment of a diabetic condition, preferably type 2 diabetes.
12. A method for controlling body weight reduction, for preventing body weight gain and/or for the treatment and/or the prevention of obesity or an overweight condition.
13. A method for the treatment and/or prevention of amyloidosis-related diseases.
14. A method for the treatment or prophylaxis of multiple risk factors for cardiovascular diseases.
15. A method for the prevention of stroke, cerebral or transient ischaemic attacks related to atherosclerosis of several arteries.
16. Processes for preparing novel fatty acid analogues according to the invention.

The present invention relates to a compound of formula (I):



wherein

- R_1 and R_2 are the same or different and may be selected from the group consisting of a hydrogen atom, a hydroxy group, an alkyl group, a halogen atom, an alkoxy group, an acyloxy group, an acyl group, an alkenyl group, an alkynyl group, an aryl group, an alkylthio group, an alkoxycarbonyl group, an alkylsulfinyl group, an alkylsulfonyl group, an amino group, and an alkylamino group; and
- X represents a carboxylic acid group, a carboxylate group, or a carboxamide group,

or any pharmaceutically acceptable salt, solvate, complex or pro-drug thereof, with the provisos that:

- the compound of formula (I) is not (all-Z)-4,7,10,13,16,19-docosahexaenoic acid (DHA), alpha-methyl DHA, alpha-methyl DHA methyl ester, alpha-methyl DHA ethyl ester, or alpha-hydroxy DHA ethyl ester.

The provisos correspond to the following cases:

- when R_1 is a hydrogen atom, then R_2 is not a hydrogen atom;

- when R₂ is a hydrogen atom, then R₁ is not a hydrogen atom;
- when R₁ is a methyl group, then R₂ is not a hydrogen atom, and X is not a carboxylic acid group, a methylcarboxylate, or an ethylcarboxylate;
- 5 ▪ when R₂ is a methyl group, then R₁ is not a hydrogen atom, and X is not a carboxylic acid group, a methylcarboxylate, or an ethylcarboxylate;
- when R₁ is a hydroxy group, then R₂ is not a hydrogen atom, and X is not an ethylcarboxylate; and
- when R₂ is a hydroxy group, then R₁ is not a hydrogen atom, and X is not an ethylcarboxylate.

10 In a compound according to the invention, said alkyl group may be selected from the group consisting of methyl, ethyl, n-propyl, isopropyl, n-butyl, sec.-butyl, n-hexyl, and benzyl; said halogen atom may be selected from the group consisting of fluorine, chlorine, bromine, and iodine; said alkoxy group may be selected from the group consisting of methoxy, ethoxy, propoxy, isopropoxy, sec.-butoxy, phenoxy,
15 benzyloxy, OCH₂CF₃, and OCH₂CH₂OCH₃; said acyloxy group may be selected from acetoxymethyl, propionoxymethyl, and butyroxymethyl; said alkenyl group may be selected from the group consisting of allyl, 2-butenyl, and 3-hexenyl; said alkynyl group may be selected from the group consisting of propargyl, 2-butyne, and 3-hexyne; said aryl group is a phenyl group; said alkylthio group may be selected from the group
20 consisting of methylthio, ethylthio, isopropylthio, and phenylthio; said alkoxy carbonyl group may be selected from the group consisting of methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, and butoxycarbonyl; said alkylsulfinyl group may be selected from the group consisting of methanesulfinyl, ethanesulfinyl, and isopropanesulfinyl; said alkylsulfonyl group may be selected from
25 the group consisting of methanesulfonyl, ethanesulfonyl, and isopropanesulfonyl; said alkylamino group may be selected from the group consisting of methylamino, dimethylamino, ethylamino, and diethylamino; said carboxylate group may be selected from the group consisting of ethyl carboxylate, methyl carboxylate, n-propyl carboxylate, isopropyl carboxylate, n-butyl carboxylate, sec.-butyl carboxylate, and
30 n-hexyl carboxylate; said carboxamide group may be selected from the group consisting of primary carboxamide, N-methyl carboxamide, N,N-dimethyl carboxamide, N-ethyl carboxamide, and N,N-diethyl carboxamide.

35 In one embodiment of the invention, R₁ and R₂ are selected from the group consisting of a hydrogen atom, a hydroxy group, an alkyl group, a halogen atom, an alkoxy group, an alkylthio group, an alkylsulfinyl group, an alkylsulfonyl group, an amino group, and an alkylamino group.

In another embodiment of the invention, R₁ and R₂ are selected from the group consisting of a hydrogen atom, a hydroxy group, a C₁-C₇ alkyl group, a halogen atom, a C₁-C₇ alkoxy group, a C₁-C₇ alkylthio group, a C₁-C₇ alkylsulfinyl group, a C₁-C₇ alkylsulfonyl group, an amino group, and a C₁-C₇ alkylamino group.

5 Then, said C₁-C₇ alkyl group may be methyl, ethyl, or benzyl; said halogen atom may be fluorine or iodine; said C₁-C₇ alkoxy group may be methoxy or ethoxy; said C₁-C₇ alkylthio group may be methylthio, ethylthio or phenylthio; said C₁-C₇ alkylsulfinyl group may be ethanesulfinyl; said C₁-C₇ alkylsulfonyl group may be ethanesulfonyl; said C₁-C₇ alkylamino group may be ethylamino or diethylamino; and X may
10 represent an ethylcarboxylate or a carboxamide group.

In another embodiment of the invention, R₁ and R₂ are selected from the group consisting of a hydrogen atom, a C₂-C₇ alkyl group, a halogen atom, a C₁-C₇ alkoxy group, a C₁-C₇ alkylthio group, a C₁-C₇ alkylsulfinyl group, a C₁-C₇ alkylsulfonyl group, an amino group, and a C₁-C₇ alkylamino group; and

15 X represents a carboxylate. Then, said C₂-C₇ alkyl group may be ethyl, or benzyl; said halogen atom may be fluorine or iodine; said C₁-C₇ alkoxy group may be methoxy or ethoxy; said C₁-C₇ alkylthio group may be methylthio, ethylthio or phenylthio; said C₁-C₇ alkylsulfinyl group may be ethanesulfinyl; said C₁-C₇ alkylsulfonyl group may be ethanesulfonyl; said C₁-C₇ alkylamino group may be
20 ethylamino or diethylamino; and X represents a an ethylcarboxylate.

In the compound according to formula (I) of the present invention, R₁ and R₂ may be the same or different. When they are different, the compounds of formula (I) are capable of existing in stereoisomeric forms. It will be understood that the invention encompasses all optical isomers of the compounds of formula (I) and
25 mixtures thereof including racemates.

Therefore, the present invention includes, where R₁ is different from R₂, compounds of formula (I) that are racemic or enantiomerically pure, either as the (S) or (R) enantiomer. Therefore, the present invention includes, where R₁ is different from R₂, compounds of formula (I) that are racemic or enantiomeric pure, either as
30 the (S) or (R) stereoisomer.

Within the scope of the invention are enantiomers of the compounds of the formula (I), as hereinbefore defined. Moreover, the enantiomers of the DHA derivatives according to the invention might be in the form of a carboxylic acid, or a pharmaceutically acceptable salt thereof, any ester, anhydride or amide (primary,
35 secondary, tertiary). The acid derivative might be in the form of a phospholipid or a tri- di- or monoglyceride.

In one embodiment of a compound of formula (I) according to the invention, one of R_1 and R_2 represents a C_2 - C_7 alkyl group, e.g. ethyl or benzyl, and the other one represents a hydrogen atom. Preferably, the alkylgroup is ethyl.

In another embodiment of a compound of formula (I) according to the invention, one of R_1 and R_2 represents an alkoxy group, e.g. ethoxy or methoxy, and the other one represents a hydrogen atom.

In another embodiment of a compound of formula (I) according to the invention, one of R_1 and R_2 represents a halogen atom, e.g. fluorine or iodine, and the other one represents a hydrogen atom.

In another embodiment of a compound of formula (I) according to the invention, one of R_1 and R_2 represents an alkylthio group, e.g. ethylthio, methylthio or phenylthio, and the other one represents a hydrogen atom. Preferably, the alkylthiogroup is ethylthio.

In another embodiment of a compound of formula (I) according to the invention, one of R_1 and R_2 represents an alkylsulfonyl group, e.g. ethylsulfonyl, and the other one represents a hydrogen atom.

In another embodiment of a compound of formula (I) according to the invention, one of R_1 and R_2 represents an amino group, and the other one represents a hydrogen atom.

In another embodiment of a compound of formula (I) according to the invention, one of R_1 and R_2 represents an alkyl-amino group, e.g. ethyl-amino or diethyl-amino, and the other one represents a hydrogen atom.

In a further embodiment of a compound of formula (I) according to the invention, R_1 and R_2 are the same and represent C_1 - C_7 -alkyl groups, preferably methyl groups or ethyl groups.

In preferred embodiments of the compound of formula (I), X is a carboxylate, e.g. ethyl carboxylate.

The compound according to the invention may exist in the form of a phospholipid, a tri-, di- or monoglyceride, or in the form of a free acid.

The alpha-substituted DHA-derivatives according to the invention have very surprisingly shown excellent results with regard to pharmaceutical activity. In particular, the fatty acid derivatives according to the present invention possess a huge potential to be used in the treatment and/or prevention of diabetes and pre-stages thereof.

Another aspect of the present invention relates to a compound of formula (I) for use as a medicament.

The invention also relates to a process for the manufacture of a compound of formula (I). For example, a compound of formula (I) may be prepared from (all-Z)-4,7,10,13,16,19-docosahexaenoic acid (DHA). The DHA may e.g. originate from a vegetable, a microbial and/or an animal source, such as a marine fish oil. Another
5 important advantage with compounds of formula (I) is that the fatty acid analogues can be prepared directly from (all-Z)-4,7,10,13,16,19-docosahexaenoic acid (DHA).

In a preferred embodiment of the invention, the fatty acid analogues of formula (I) are prepared from DHA, wherein said DHA is obtained from at least one of vegetable, microbial and animal origins, or combinations thereof. The invention
10 includes therefore derivatives prepared from DHA-containing oil from microbial origin. Suitable, said DHA is produced from a marine oil, such as a fish oil.

Another aspect of the present invention relates to a pharmaceutical composition comprising a compound of formula (I) as an active ingredient. The pharmaceutical composition may further comprise a pharmaceutically acceptable
15 carrier. Suitably, a pharmaceutical composition according to the invention is formulated for oral administration, e.g. in the form of a capsule or a sachet. A suitable daily dosage of a compound of formula (I) according to the present invention is 10 mg to 10 g, in particular 100 mg to 1 g of said compound.

In addition, the present invention relates to a fatty acid composition
20 comprising a compound of formula (I). At least 60%, or at least 90% by weight of the fatty acid composition may be comprised of said compound. The fatty acid composition may further comprise (all-Z)-5,8,11,14,17-eicosapentaenoic acid (EPA), (all-Z)-4,7,10,13,16,19-docosahexaenoic acid (DHA), (all-Z)-6,9,12,15,18-heneicosapentaenoic acid (HPA), and/or (all-Z)-7,10,13,16,19-docosapentaenoic acid
25 (DPA). The fatty acids may be present in the form of derivatives. A fatty acid composition according to the present invention may further comprise a pharmaceutically acceptable antioxidant, e.g. tocopherol. Within the scope of the present invention is also a fatty acid composition described above, for use as a medicament.

30 In a further aspect, the present invention relates to the use of a compound according to formula (I) for the manufacture of a medicament for controlling body weight reduction and/or for preventing body weight gain; for the manufacture of a medicament for the treatment and/or the prevention of obesity or an overweight condition; for the manufacture of a medicament for the prevention and/or treatment of
35 diabetes in an animal, in particular type 2 diabetes; for the manufacture of a medicament for the treatment and/or prevention of amyloidosis-related diseases; for the manufacture of a medicament for the treatment or prophylaxis of multiple risk factors

for cardiovascular diseases, preferably for the treatment of elevated blood lipids for the manufacture of a medicament for prevention of stroke, cerebral or transient ischaemic attacks related to atherosclerosis of several arteries.

In addition, the present invention relates to a method for controlling body weight reduction and/or for preventing body weight gain; a method for the treatment and/or the prevention of obesity or an overweight condition; a method for the prevention and/or treatment of diabetes, in particular type 2 diabetes; a method for the treatment and/or prevention of amyloidosis-related diseases; a method for the treatment or prophylaxis of multiple risk factors for cardiovascular diseases; a method for the prevention of stroke, cerebral or transient ischaemic attacks related to atherosclerosis of several arteries, wherein a pharmaceutically effective amount of a compound of formula (I) is administered to a human or an animal. Suitably, the compound of formula (I) is administered orally to a human or an animal.

15 **Brief description of the drawings**

Fig. 1 is a schematic overview of the free fatty acid pool theory.

Fig. 2 shows an overview of the models and methods used in the present invention for demonstrating effects on the metabolic syndrome and type 2 diabetes

Fig. 3 depicts the free fatty acid concentrations of different compounds according to the invention in liver tissue from animals given these compounds in a concentration of 1.5% of total fat content.

Fig. 4 depicts the intracellular concentrations of DHA in liver tissue from animals given different compounds according to the invention in a concentration of 1.5% of total fat content.

Fig. 5 depicts the binding affinities for the PPAR γ receptor of different compounds according to the invention.

Fig. 6 depicts the binding affinities to the nuclear receptor PPAR α of different compounds according to the invention.

Fig. 7 depicts the binding affinities to the nuclear receptor RXR α of different compounds according to the invention.

Fig. 8 depicts the release of luciferase from transfected cells treated with different compounds according to the invention.

Fig. 9 shows the study design of the experiment of block 4.

Fig. 10 shows the change of body weight during 2 weeks of diet intervention after 8 weeks of HF diet.

Fig. 11 shows the results from luciferase activity, i.e. endogenous PPAR γ -activity).

Fig. 12 shows the endogenous luciferase activity in different compounds according to the invention compared to DHA.

Fig. 13 shows a typical blood glucose elimination curve before and after animals with insulin resistance are given a compound with insulin resistance reducing effect.

Figs. 14, 15 and 16 show different effects of DHA derivatives according to the invention on metabolic syndrome and insulin resistance.

Detailed description of the invention

10 In the research work leading to the present invention, novel DHA-derivatives were prepared, which showed excellent pharmaceutical activity.

Fatty acids enter cells passively or through G-protein coupled transporter systems, such as fatty acid transport proteins. Well inside the cells they are temporarily bound by binding proteins (Fatty acid binding proteins, FABP), which
15 play an important role in directing fatty acids to various intracellular compartments for metabolism and gene expression (Pawar & Jump 2003). (Fig. 1 liver cell).

Esterification of fatty acids into triglycerides, polar lipids, and cholesterol esters and their beta-oxidation (mitochondrial and peroxisomal) requires conversion of fatty acids to acyl CoA thioesters. Other pathways, like microsomal NADPH-
20 dependent mono-oxidation and eicosanoids synthesis, utilise non-esterified fatty acids as substrates. All these reactions are likely to influence cellular levels of free fatty acids (non-esterified) and thereby the amount and type of fatty acids which could be used as ligands to nuclear receptors. Because PPARs are known to bind non-esterified fatty acids it is reasonable to expect that the composition of the free fatty acid pool is
25 an important determinant in the control of PPAR activity.

The composition of the free fatty acid pool is affected by the concentration of exogenous fatty acids entering the cells, and their rate of removal via pathways listed above. Since short and medium chain fatty acids are effectively recruited to these pathways, in practice only the long-chain polyunsaturated fatty acids will be available
30 for liganding to nuclear receptors. In addition, fatty acid structure may also be an important determinant. Even if a series of mono and polyunsaturated fatty acids demonstrated affinity to the PPAR α receptor, EPA and DHA demonstrated the highest binding capacity in experiments with rat liver cells (Pawar & Jump 2003).

Searching for fatty acid candidates available for genetic modification of
35 proteins by interaction with nuclear receptors like the PPARs, it is important to verify that the respective fatty acids will be enriched in the free fatty acid pool.





DHA which enter cells are rapidly converted to fatty acyl-CoA thioesters and incorporated into phospholipids and due to this, the intracellular DHA level is relatively low. These DHA-CoA are also substrate for β -oxidation primarily in the peroxisomes that lead to retroconversion of DHA to EPA, see Fig. 1. Because of the rapid incorporation into neutral lipids and the oxidation pathway DHA will not stay long in the free fatty acid pool. Due to this the effect of DHA on gene expression is probably limited.





The present invention aims at achieving an accumulation of fatty acid derivatives in the free fatty acid pool, rather than incorporation into phospholipids. The present inventors have surprisingly found that the introduction of at least one substituent in the α -position of DHA will lead to a slower oxidation rate in addition to less incorporation into neutral lipids. This will lead to an increased effect on gene expression, since the DHA derivatives will accumulate in the tissue particular within liver, muscle, and adipose cells and trigger local nuclear receptor activity to a greater extent than DHA.

The different substituents according to the invention will give variable affinities of the derivatives to fatty acids binding receptors. It is also possible that changes in affinity to fatty acids binding proteins lead to changes in the biological activity of these α -substituted DHA derivatives of formula (I). Altogether these changes lead to an increased therapeutic effect of the DHA derivatives according to the invention compared to DHA.


EPA (all-Z)-5,8,11,14,17-eicosapentaenoic acid) has earlier been alkylated in α - and β -position to inhibit mitochondrial β -oxidation. DHA is not oxidised in the mitochondria, but rather incorporated into phospholipids. In the peroxisomes though some DHA is retroconverted to EPA. A substituent in the α -position of EPA and DHA will due to this affect different metabolic pathways. It has earlier been shown that α -methyl EPA and β -methyl EPA is incorporated into phospholipids and triglycerids while α -ethyl EPA is not (Larsen 1998). In this study the derivatives were tested as substrates and/or inhibitors of enzymes involved in the eicosanoid cascade. Since most of the substrates for these enzymes are fatty acids liberated from phospholipids it was desired that the derivatives were incorporated into phospholipids. In contrast to this, as mentioned before, we want derivatives that will not incorporate into lipids, but rather accumulate in the NEFA pool.

Throughout this description, the abbreviation "PRB-x", where x is an integer, will be used when describing specific compounds according to the invention. Below, the structural formulas and trivial names for each of these compounds are listed:


PRB-1 α -methyl docosahexaenoic acid ethyl esterPRB-2 α -ethyl docosahexaenoic acid ethyl esterPRB-3 α -ethoxy docosahexaenoic acid ethyl esterPRB-4 α -fluoro docosahexaenoic acid ethyl ester

PRB-5 α,α di-methyl docosaheptaenoic acid ethyl esterPRB-6 α -tiomethyl docosaheptaenoic acid ethyl esterPRB-7 α -tioethyl docosaheptaenoic acid ethyl esterPRB-8 α,α di-ethyl docosaheptaenoic acid ethyl ester


15




PRB-9 α -benzyl docosahexaenoic acid ethyl ester




PRB-10 α -ethanesulfinyl docosahexaenoic acid ethyl ester




PRB-11 α -thiophenyl docosahexaenoic acid ethyl ester




PRB-12 α -hydroxy docosahexaenoic acid ethyl ester





PRB-13 α -methyl docosahexaenoic acid
amide





PRB-14 α -methoxy docosahexaenoic acid
ethyl ester




PRB-15 α -iodo docosahexaenoic acid ethyl
ester


PRB-17 α -amino docosahexanoic acid ethyl ester

PRB-18 (4R,5S)-3-docosa-2,4,6,8,10,12,14,16-octenyl-4-methyl-5-phenyl-oxazolidin-2-one


PRB-19 (4R,5S)-3-[(S)- α -ethyl docosa-2,4,6,8,10,12,14,16-octenyl]-4-methyl-5-phenyl-oxazolidin-2-onePRB-20 (S)-(+)- α -ethyl docosa-2,4,6,8,10,12,14,16-octanoic acid ethyl ester




PRB-21 (4S,5R)-3-docosa-6,8,10,12,14,16,18,20-octanoic acid
4-methyl-5-phenyl-oxazolidin-2-one



PRB-22 (4S,5R)-3-[(R)-α-ethyl docosahexaenoyl]-
4-methyl-5-phenyl-oxazolidin-2-one




PRB-23 (R)-(-)-α-ethyl docosahexanoic acid ethyl ester




PRB-24

2-(1,3-Dioxo-1,3-dihydro-isoindol-2-yl)-docosahexanoic acid ethyl ester



PRB-25 α -ethyl-amino
docosahexanoic acid ethyl
ester



PRB-26 α -diethyl-amino
docosahexanoic acid ethyl
ester

PRB-1 corresponds to a compound of formula (I) in which R_1 or R_2 is methyl, and the other one is hydrogen, and X is ethyl carboxylate.

PRB-2 corresponds to a compound of formula (I) in which R_1 or R_2 is ethyl, and the other one is hydrogen, and X is ethyl carboxylate.

5 PRB-3 corresponds to a compound of formula (I) in which R_1 or R_2 is ethoxy, and the other one is hydrogen, and X is ethyl carboxylate.

PRB-4 corresponds to a compound of formula (I) in which R_1 or R_2 is fluorine, and the other one is hydrogen, and X is ethyl carboxylate.

10 PRB-5 corresponds to a compound of formula (I) in which R_1 and R_2 is methyl, and X is ethyl carboxylate.

PRB-6 corresponds to a compound of formula (I) in which R_1 or R_2 is methylthio, and X is ethyl carboxylate.

PRB-7 corresponds to a compound of formula (I) in which R_1 or R_2 is ethylthio, and the other one is hydrogen, and X is ethyl carboxylate.

15 PRB-8 corresponds to a compound of formula (I) in which R_1 and R_2 is ethyl, and the other one is hydrogen, and X is ethyl carboxylate.

PRB-9 corresponds to a compound of formula (I) in which R_1 or R_2 is benzyl, and the other one is hydrogen, and X is ethyl carboxylate.

20 PRB-10 corresponds to a compound of formula (I) in which R_1 or R_2 is ethanesulfinyl, and the other one is hydrogen, and X is ethyl carboxylate.

PRB-11 corresponds to a compound of formula (I) in which R_1 or R_2 is phenylthio, and the other one is hydrogen, and X is ethyl carboxylate.

PRB-12 corresponds to a compound of formula (I) in which R_1 or R_2 is hydroxy, and the other one is hydrogen, and X is ethyl carboxylate.

25 PRB-13 corresponds to a compound of formula (I) in which R_1 or R_2 is methyl, and the other one is hydrogen, and X is primary carboxamide.

PRB-14 corresponds to a compound of formula (I) in which R_1 or R_2 is methoxy, and the other one is hydrogen, and X is ethyl carboxylate.

30 PRB-15 corresponds to a compound of formula (I) in which R_1 or R_2 is iodine, and the other one is hydrogen, and X is ethyl carboxylate.

PRB-17 corresponds to a compound of formula (I) in which R_1 or R_2 is amino, and the other one is hydrogen, and X is ethyl carboxylate.

PRB-20 corresponds to the (S) stereoisomer of a compound of formula (I) in which R_1 or R_2 is ethyl, and the other one is hydrogen, and X is ethyl carboxylate.

35 PRB-23 corresponds to the (R) stereoisomer of a compound of formula (I) in which R_1 or R_2 is ethyl, and the other one is hydrogen, and X is ethyl carboxylate.

PRB-24 corresponds to a compound of formula (I) in which R₁ or R₂ is N-phtalimide, and the other one is hydrogen, and X is ethyl carboxylate.

PRB-25 corresponds to a compound of formula (I) in which R₁ or R₂ is ethyl-amino, and the other one is hydrogen, and X is primary carboxamide .

5 PRB-26 corresponds to a compound of formula (I) in which R₁ or R₂ is diethyl-amino, and the other one is hydrogen, and X is ethyl carboxylate.

PRB-2 is the most preferred compound according to the present invention. Other preferred compounds according to the invention are PRB-5, PRB-7, and PRB-8.

10 It is to be understood that the present invention encompasses any possible pharmaceutically acceptable salts, solvates, complexes or prodrgs of the compounds of formula (I).

"Prodrugs" are entities which may or may not possess pharmacological activity as such, but may be administered (such as orally or parenterally) and
15 thereafter subjected to bioactivation (for example metabolized) in the body to form the agent of the present invention which is pharmacologically active.

Where X is a carboxylic acid, the present invention also includes salts of the carboxylic acids. Suitable pharmaceutically acceptable salts of carboxy groups includes metal salts, such as for example aluminium, alkali metal salts such as
20 lithium, sodium or potassium, alkaline metal salts such as calcium or magnesium and ammonium or substituted ammonium salts.

A "therapeutically effective amount" refers to the amount of the therapeutic agent which is effective to achieve its intended purpose. While individual patient needs may vary, determination of optimal ranges for effective amounts of each nitric
25 oxide adduct is within the skill of the art. Generally the dosage regimen for treating a condition with the compounds and/or compositions of this invention is selected in accordance with a variety of factors, including the type, age, weight, sex, diet and medical condition of the patient.

By "a medicament" is meant a compound according to formula (I), in any
30 form suitable to be used for a medical purpose, e.g. in the form of a medicinal product, a pharmaceutical preparation or product, a dietary product, a food stuff or a food supplement.

In the context of the present specification, the term "therapy" also includes "prophylaxis" unless there are specific indications to the contrary. The terms
35 "therapeutic" and "therapeutically" should be constructed accordingly.

Treatment includes any therapeutic application that can benefit a human or non-human animal. The treatment of mammals is particularly preferred. Both human

and veterinary treatments are within the scope of the present invention. Treatment may be in respect of an existing condition or it may be prophylactic. It may be of an adult, a juvenile, an infant, a foetus, or a part of any of the aforesaid (e.g. an organ, tissue, cell, or nucleic acid molecule). By "chronic treatment" is meant treatment that
5 continues for some weeks or years.

"A therapeutically or a pharmaceutically active amount" relates to an amount that will lead to the desired pharmacological and/or therapeutic effects.

A compound according to the present invention may for example be included in a food stuff, a food supplement, a nutritional supplement, or a dietary product

10 Alpha-substituted DHA derivatives and EPA (or DHA for that matter) can be bound together and combined on triglyceride form by an esterification process between a mixture of alpha-derivatives, EPA and glycerol catalysed by Novozym 435 (a commercially available lipase from *Candida antarctica* on immobilised form).

The compounds of formula (I) have activity as pharmaceuticals, in particular
15 as triggers of nuclear receptor activity. Thus, the present invention also relates to compounds of formula (I), pharmaceutically acceptable salts, solvates, complexes or pro-drugs thereof, as hereinbefore defined, for use as a medicament and/or for use in therapy. Preferably, the novel compounds, or pharmaceutically acceptable salts, solvates, complexes or pro-drugs thereof, of the invention may be used:

- 20 - for the prevention and/or treatment of diabetes mellitus in humans or animals;
- for controlling body weight reduction and/or for preventing body weight gain;
- for the prevention and/or treatment of obesity or an overweight condition in humans or in an animal;
- for the treatment and/or prevention of amyloidosis-related diseases;
- 25 - for the treatment or prophylaxis of multiple risk factors for cardiovascular diseases;
- for the prevention of stroke, cerebral or transient ischaemic attacks related to atherosclerosis of several arteries.
- for the treatment of TBC or HIV.

30 There are two major forms of diabetes mellitus. One is type 1 diabetes, which is known as insulin-dependent diabetes mellitus (IDDM), and the other one is type 2 diabetes, which is also known as non-insulin-dependent diabetes mellitus (NIDDM). Type 2 diabetes is related to obesity/overweight and lack of exercise, often of gradual onset, usually in adults, and caused by reduced insulin sensitivity, so called peripheral
35 insulin resistance. This leads to a compensatory increase in insulin production. This stage before developing full fetched type 2 diabetes is called the metabolic syndrome and characterized by hyperinsulinemia, insulin resistance, obesity, glucose

intolerance, hypertension, abnormal blood lipids, hypercoagulopathia, dyslipidemia and inflammation, often leading to atherosclerosis of the arteries. Later when insulin production ceases, type 2 diabetes mellitus develops.

In a preferred embodiment, the compounds according to formula (I) may be used for the treatment of type 2 diabetes. The compounds according to formula (I) may also be used for the treatment of other types of diabetes selected from the group consisting of metabolic syndrome, secondary diabetes, such as pancreatic, extrapancreatic/endocrine or drug-induced diabetes, or exceptional forms of diabetes, such as lipoatrophic, myotonic or a disease caused by disturbance of the insulin receptors. The invention also includes treatment of type 2 diabetes. Suitably, compounds of formula (I), as hereinbefore defined, may activate nuclear receptors, preferably PPAR (peroxisome proliferator-activated receptor) α and/or γ .

The compounds of formula (I) may also be used for the treatment and/or prevention of obesity. Obesity is usually linked to an increased insulin resistance and obese people run a high risk of developing type 2 diabetes which is a major risk factor for development of cardiovascular diseases. Obesity is a chronic disease that afflicts an increasing proportion of the population in Western societies and is associated, not only with a social stigma, but also with decreasing life span and numerous problems, for instance diabetes mellitus, insulin resistance and hypertension. The present invention thus fulfils a long-felt need for a drug that will reduce total body weight, or the amount of adipose tissue, of preferably obese humans, towards their ideal body weight without significant adverse side effects.

The compounds according to formula (I) may also be used for the prevention and/or treatment of amyloidosis-related diseases. Amyloidosis-related conditions or diseases associated with deposition of amyloid, preferably as a consequence of fibril or plaque formation, includes Alzheimer's disease or dementia, Parkinson's disease, amyotrophic lateral sclerosis, the spongiform encephalopathies, such as Creutzfeldt-jacob disease, cystic fibrosis, primary or secondary renal amyloidosis, IgA nephropathy, and amyloid deposition in arteries, myocardium and neural tissue. These diseases can be sporadic, inherited or even related to infections such as TBC or HIV, and are often manifested only late in life even if inherited forms may appear much earlier. Each disease is associated with a particular protein or aggregates of these proteins are thought to be the direct origin of the pathological conditions associated with the disease. The treatment of an amyloidosis-related disease can be made either acutely or chronically.

The compounds of formula (I) may also be used for the treatment due to reduction of amyloid aggregates, prevention of misfolding of proteins that may lead

to formation of so called fibrils or plaque, treatment due to decreasing of the production of precursor protein such as A β -protein (amyloid beta protein), and prevention and/or treatment due to inhibiting or slow down the formation of protein fibrils, aggregates, or plaque. Prevention of fibril accumulation, or formation, by
5 administering compounds of formula (I), as hereinbefore defined, is also included herein. In one embodiment, the novel compounds, pharmaceutically acceptable salts, solvates, complexes or pro-drugs thereof, as hereinbefore defined, are used for the treatment of TBC (tuberculosis) or HIV (human immunodeficiency virus).

Further, the compounds of formula (I) may be administered to patients with
10 symptoms of atherosclerosis of arteries supplying the brain, for instance a stroke or transient ischaemic attack, in order to reduce the risk of a further, possible fatal, attack.

The compounds of formula (I) may also be used for the treatment of elevated blood lipids in humans.

15 Additionally, the compounds of formula (I), as hereinbefore defined, are valuable for the treatment and prophylaxis of multiple risk factors known for cardiovascular diseases, such as hypertension, hypertriglyceridemia and high coagulation factor VII phospholipid complex activity. Preferably, the compounds of formula (I) is used for the treatment of elevated blood lipids in humans.

20 The compounds of formula (I) and pharmaceutically acceptable salts, solvates, pro-drugs or complexes thereof may be used on their own but will generally be administered in the form of a pharmaceutical composition in which the compounds of formula (I) (the active ingredient) are in association with a pharmaceutically acceptable adjuvant, diluent or carrier.

25 The present invention thus also provides a pharmaceutical composition comprising a therapeutically effective amount of the compound of formula (I) of the present invention and a pharmaceutically acceptable carrier, diluent or excipients (including combinations thereof).

This is a composition that comprises or consists of a therapeutically
30 effective amount of a pharmaceutically active agent. It preferably includes a pharmaceutically acceptable carrier, diluent or excipients (including combinations thereof). Acceptable carriers or diluents for therapeutic use are well known in the pharmaceutical art. The choice of pharmaceutical carrier, excipient or diluent can be selected with regard to the intended route of
35 administration and standard pharmaceutical practice. The pharmaceutical compositions may comprise as - or in addition to - the carrier, excipient or

diluent any suitable binder(s), lubricant(s), suspending agent(s), coating agent(s), solubilising agent(s).

Pharmaceutical compositions within the scope of the present invention may include one or more of the following: preserving agents, solubilising agents, stabilising agents, s wetting agents, emulsifiers, sweeteners, colourants, flavouring agents, odourants, salts compounds of the present invention may themselves be provided in the form of a pharmaceutically acceptable salt), buffers, coating agents, antioxidants, suspending agents, adjuvants, excipients and diluents.

A pharmaceutical composition according to the invention is preferably formulated for oral administration to a human or an animal. The pharmaceutical composition may also be formulated for administration through any other route where the active ingredients may be efficiently absorbed and utilized, e.g. intravenously, subcutaneously, intramuscularly, intranasally, rectally, vaginally or topically.

In a specific embodiment of the invention, the pharmaceutical composition is shaped in form of a capsule, which could also be microcapsules generating a powder or a sachet. The capsule may be flavoured. This embodiment also includes a capsule wherein both the capsule and the encapsulated fatty acid composition according to the invention is flavoured. By flavouring the capsule it becomes more attractive to the user. For the above-mentioned therapeutic uses the dosage administered will, of course, vary with the compound employed, the mode of administration, the treatment desired and the disorder indicated.

The pharmaceutical composition may be formulated to provide a daily dosage of 10 mg to 10 g. Preferably, the pharmaceutical composition is formulated to provide a daily dosage between 50 mg and 5 g of said composition. Most preferably, the pharmaceutical composition is formulated to provide a daily dosage between 100 mg and 1 g of said composition. By a daily dosage is meant the dosage per 24 hours. The dosage administered will, of course, vary with the compound employed, the mode of administration, the treatment desired and the disorder indicated. Typically, a physician will determine the actual dosage which will be most suitable for an individual subject. The specific dose level and frequency of dosage for any particular patient may be varied and will depend upon a variety of factors including the activity of the specific compound employed, the metabolic stability and length of action of that compound, the age, body weight, general health, sex, diet, mode and time of administration, rate of excretion, drug combination, the severity of the particular condition, and the individual undergoing therapy. The agent and/or the pharmaceutical composition of the present invention may be administered in accordance with a regimen of from 1 to 10 times per day, such as once or twice per

day. For oral and parenteral administration to human patients, the daily dosage level of the agent may be in single or divided doses.

A further aspect of the present invention relates to a fatty acid composition comprising compounds of formula (I). A fatty acid composition comprising
5 compounds of formula (I) increases the natural biological effects of DHA that are a result of regulation of gene expression, and the derivatives according to the present invention will accumulate in the free fatty acid pool.

The fatty acid composition may comprise in the range of 60 to 100 % by weight of the compounds of formula (I), all percentages by weight being based on the
10 total weight of the fatty acid composition. In a preferred embodiment of the invention, at least 80% by weight of the fatty acid composition is comprised of compounds of formula (I). More preferably, the compounds of formula (I) constitute at least 90% by weight of the fatty acid composition. Most preferably, the compounds of formula (I) constitutes more than 95% by weight of the fatty acid composition.

The fatty acid composition may further comprise at least one of the fatty acids
15 (all-Z)-5,8,11,14,17-eicosapentaenoic acid (EPA), (all-Z)-4,7,10,13,16,19-docosahexaenoic acid (DHA), (all-Z)-6,9,12,15,18-heneicosapentaenoic acid (HPA), and (all-Z)-7,10,13,16,19-docosapentaenoic acid (DPAn-3), (all-Z)-8,11,14,17-eicosatetraenoic acid (ETAn-3), or combinations thereof. Further, the fatty acid
20 composition may comprise (all-Z)-4,7,10,13,16-Docosapentaenoic acid (DPAn-6) and/or (all-Z)-5,8,11,14-eicosatetraenoic acid (ARA), or derivatives thereof. The fatty acid composition may also comprise at least these fatty acids, or combinations thereof, in the form of derivatives. The derivatives are suitably substituted in the same way as the DHA derivatives of formula (I), as hereinbefore defined.

The fatty acid composition according to the invention may comprise (all-Z
25 omega-3)-6,9,12,15,18-heneicosapentaenoic acid (HPA), or derivatives thereof, in an amount of at least 1% by weight, or in an amount of 1 to 4% by weight.

Further, the fatty acid composition according to the invention may comprise omega-3 fatty acids other than EPA and DHA that have 20, 21, or 22 carbon atoms,
30 or derivatives thereof, in an amount of at least 1.5% by weight, or in an amount of at least 3% by weight.

In specific embodiments of the invention, the fatty acid composition is a pharmaceutical composition, a nutritional composition or a dietary composition. The fatty acid composition may further comprise an effective amount of a
35 pharmaceutically acceptable antioxidant. Preferably, the antioxidant is tocopherol or a mixture of tocopherols. In a preferred embodiment the fatty acid composition further comprises tocopherol, or a mixture of tocopherols, in an amount of up to 4 mg per g

of the total weight of the fatty acid composition. Preferably, the fatty acid composition comprises an amount of 0.2 to 0.4 mg per g of tocopherols, based on the total weight of the composition.

Another aspect of the invention provides a fatty acid composition, or any
5 pharmaceutically acceptable salt, solvate, pro-drug or complex thereof, comprising compounds of formula (I), as hereinbefore defined, for use as a medicament and/or in therapy. Such a fatty acid composition may be used to prevent and/or treat the same conditions as outlined for the compounds of formula (I) above.

When the fatty acid composition is used as a medicament, it will be
10 administered in a therapeutically or a pharmaceutically active amount.

In a preferred embodiment, the fatty acid composition is administered orally to a human or an animal.

The present invention also provides the use of a compound of formula (I), or a pharmaceutically acceptable salt, solvate, pro-drug or complex thereof, as
15 hereinbefore defined, for the manufacture of a medicament for controlling body weight reduction and/or for preventing body weight gain; for the manufacture of a medicament for the treatment and/or the prevention of obesity or an overweight condition; for the manufacture of a medicament for the prevention and/or treatment of diabetes in a human or animal; for the manufacture of a medicament for the treatment and/or prevention of amyloidosis-related diseases; for the manufacture of a
20 medicament for the treatment and prophylaxis of multiple risk factors known for cardiovascular diseases, such as hypertension, hypertriglyceridemia and high coagulation factor VII phospholipid complex activity; for the manufacture of a medicament for the treatment of TBC or HIV; for the manufacture of a medicament
25 for prevention of stroke, cerebral or transient ischaemic attacks related to atherosclerosis of several arteries; for the manufacturing of a medicament for lowering triglycerides in the blood of mammals and/or elevating the HDL cholesterol levels in the serum of a human patients; or for the manufacturing of a medicament for the treatment and/or prevention of the multi metabolic syndrome termed "metabolic
30 syndrome". All these embodiments also include the use of a fatty acid composition, as hereinbefore defined, comprising compounds of formula (I) for the manufacture of medicaments as outlined above. The present invention further relates to the use of alpha-hydroxy-DHA for the manufacture of medicaments as outlined above.

The present invention also relates to a method for controlling body weight
35 reduction and for preventing body weight gain, wherein a fatty acid composition comprising at least a compound of formula (I), as hereinbefore defined, is administered to a human or an animal.

Further, the invention relates to a method for the treatment and/or the prevention of obesity or an overweight condition, wherein a fatty acid composition comprising at least a compound of formula (I), as hereinbefore defined, is administered to a human or an animal.

5 In a preferred embodiment of the invention, the present invention relates to a method for the prevention and/or treatment of diabetes mellitus, wherein a fatty acid composition comprising at least a compound of formula (I), as hereinbefore defined, is administered to a human or an animal. Preferably, diabetes mellitus is a type 2 diabetes.

10 Other aspects of the present invention relate to;

- a method for the treatment and/or prevention of amyloidosis-related diseases;
- a method for the treatment or prophylaxis of multiple risk factors for cardiovascular diseases;
- a method for prevention of stroke, cerebral or transient ischaemic attacks

15 related to atherosclerosis of several arteries;

wherein a fatty acid composition comprising at least a compound of formula (I), as hereinbefore defined, is administered to a human or an animal.

The fatty acid derivatives of formula (I) may be prepared most effectively from DHA. If the start material is not pure DHA (i.e. not 100% DHA) the final fatty acid composition will contain a mixture of DHA derivatives, as hereinbefore defined, and an amount of other fatty acids than DHA, wherein these fatty acids are substituted in the same way as the novel fatty acid analogous of formula (I). Such embodiments are also included herein.

20 In another embodiment of the invention, the compounds of formula (I) are prepared from (all-Z)-4,7,10,13,16,19-docosahexaenoic acid (DHA), wherein said DHA is obtained from a vegetable, a microbial and/or an animal source, or combinations thereof. Preferably, said DHA is obtained from a marine oil, such as a fish oil.

30 The fatty acids in the composition may also be obtained from a vegetable, a microbial or an animal source, or combinations thereof. Thus, the invention also includes a fatty acid composition prepared from a microbial oil.

The present invention provides processes for preparing novel fatty acid analogous of formula (I), as hereinbefore defined.

35 DHA is produced from biological sources like marine, microbial or vegetable fats. All possible raw materials are mixtures of fatty acids on triglyceride form where DHA constitutes only a fraction of the fatty acids. Typical DHA concentrations are 40% in microbial fats and 10-25% in marine fats. DHA-containing vegetable fats are

during development and fats with high DHA concentrations are expected in the future.

The first process step will always be conversion of the triglycerides to free fatty acids or monoesters. Preferable esters are methyl or ethyl esters, but other esters are possible. In this way the fatty acids bound together three by three on triglycerides are separated from each other and thereby making separation possible. Several methods of separating DHA from other fatty acids are available, the most common ones being short path distillation separating the fatty acids by volatility, and urea precipitation separating the fatty acids by degree of unsaturation. Other methods reported are silver nitrate complexation also separating the fatty acids on degree on unsaturation, esterification reactions catalysed by fatty acid selective lipases in combination with short path distillation and countercurrent extraction with supercritical carbon dioxide.

The most important challenges connected to production of pure DHA is to separate it from the other C20-22 highly unsaturated fatty acids present in all available sources. These fatty acids have properties so similar to DHA that none of the methods mentioned above provide sufficient degree of separation. For some microbial high DHA fats, which have very low levels of C20-22 highly unsaturated fatty acids, short path distillation alone or in combination of other methods mentioned may provide more than 90% purity.

Most DHA containing fats also contain considerable amounts of C20-22 highly unsaturated fatty acids, e.g. EPA (20:5n-3), n-3DPA (22:5n-3), HPA (21:5n-3) and others. The only available method for separating DHA from such fatty acids is preparative High Performance Liquid Chromatography, the stationary phase being silica gel or silver nitrate impregnated silica gel, the mobile phase being selected organic solvents or supercritical carbon dioxide. With this method DHA with more than 97% purity is available. However, it has to be noted that the production costs increases strongly with concentration, as an example is production cost for 97% DHA more 5 times higher than for 90% DHA.

DHA having a purity of 90, 95 eller 97% contains small amounts of other fatty acids. As an example, DHA having a purity of 97% contains n-3DPA (22:5n-3), but also long chain fatty acids, e.g. EPA (20:5n-3), HPA (21:5n-3), and others. However, the other fatty acids will react in a way similar to DHA and provide alpha-substituted derivatives.


Organic synthesis may provide a purification method since DHA and n-6DPA (and 22:5n-6 which normally is present in very low concentrations) are the only known fatty acids that can provide gamma-lactones by cyclisation with the first

double bond. Lactonisation followed by purification and hydrolysis back to DHA may be a possibility, but it is expected that this pathway is even more expensive than HPLC.

In one embodiment, the compounds of formula (I) where R_1 (or R_2) is a hydrogen are prepared through the following processes (Scheme 1). Suitably adapted, these processes can also be used for preparing compounds represented by the general formula (I) where both R_1 and R_2 are e.g. a C_1 - C_7 alkyl group, a benzyl, a halogen, a benzyl, an alkenyl, or an alkynyl.

Compounds represented by the general formula (I) where R_1 is a hydrogen and R_2 denotes a C_1 - C_7 alkyl group, a benzyl, a halogen, a benzyl, an alkenyl, an alkynyl are prepared by reacting a DHA ester with a strong non-nucleophilic base like lithium diisopropylamine or potassium/sodium hexamethyldisilazide in a solvent such as tetrahydrofuran, diethylether at temperatures of -60 to -78 °C, to provide the ester enolate (process 1).


(Scheme 1)



This ester enolate is reacted with an electrophilic reagent like an alkylhalide exemplified by ethyl iodide, benzyl chloride, an acyl halide exemplified by acetyl chloride, benzoyl bromide, a carboxylic anhydride exemplified by acetic anhydride or a electrophilic halogenation reagent exemplified by N-fluorobenzene sulfonimide (NFSI), etc. to provide the monosubstituted derivative (process 2). The ester is further hydrolysed in a solvent like ethanol or methanol to the carboxylic acid derivative by addition of a base like lithium/sodium hydroxide in water at temperatures between 15 - 40 °C.

Claisen condensation of the DHA EE occurs during the treatment of DHA EE with a strong base. This condensation product might possess interesting biological activity. Thus, in one embodiment of the invention the condensation (intermediate) product mentioned above, as well as the use of this product for treatment and/or prevention of diseases according to the present invention, are disclosed.

In a further embodiment, compounds represented by the general formula (I) are synthesised through following processes (Scheme 2).
(Scheme 2)



- Compounds represented by the general formula (I) where R_1 is a hydrogen and R_2 denotes a hydroxy, an alkoxy group, an acyloxy are prepared by reacting a DHA ester with a strong non-nucleophilic base like lithium diisopropylamine or potassium/sodium hexamethyldisilazideane in a solvent such as tetrahydrofuran, diethylether at temperatures of -60 to -78 $^{\circ}\text{C}$, to provide the ester enolate (process 4).
- This ester enolate is reacted with an oxygen source like dimethyldioxirane, 2-(phenylsulfonyl)-3-phenyloxaziridine, molecular oxygen with different additives like trimethylphosphite or different catalysts like a Ni(II) complex to provide α -hydroxy DHA ester(process 5). Reaction of the secondary alcohol with a base like sodiumhydride in a solvent like THF or DMF generates an alkoxide that is reacted

with different electrophilic reagents as alkyl iodide for example; methyl iodide, ethyl iodide, benzyl bromide or an acyl halide, for example; acetyl chloride, benzoyl bromide (process 6). The ester is hydrolysed in a solvent like ethanol or methanol to the carboxylic acid derivative by addition of a base like lithium/sodium hydroxide in
5 water at temperatures between 15-40 °C. (process 7).

The hydroxy-DHA ester is a useful intermediate for the introduction of other functional groups in the α -position according to the invention. The hydroxyl function can be activated by conversion to a halide or tosylate prior to reaction with different nucleophiles like ammonia, amines, thiols, etc. The Mitsunobu reaction is also useful
10 for the conversion of a hydroxyl group into other functional groups. (Mitsunobu, O, Synthesis, 1981, 1).

Compounds represented by the general formula (I), as hereinbefore defined, can also be synthesised by combinations of the different processes previously described. The present invention includes the processes mentioned above.

15 The invention further provides a process for the preparation of a pharmaceutical composition of the invention, which comprises mixing of at least a compound of formula (I), or a pharmaceutically acceptable salt, solvate, complex or pro-drug thereof, as hereinbefore defined, with a pharmaceutically acceptable adjuvant, diluent or a carrier.

20 The enantiomeric pure compounds can be prepared by resolving a racemic compound of formula (I), as hereinbefore defined. The resolution of a compound of formula (I) may be carried out using known resolution procedures, for example by reacting the compound of formula (I) with an enantiomerically pure auxiliary to provide a mixture of diastereomers that can be separated by chromatography.

25 Thereafter the two enantiomers of compound (I) may be regenerated from the separated diastereomers by conventional means, such as hydrolysis.

There is also a possibility to use stoichiometric chiral auxiliaries to effect an asymmetric introduction of the substituents, as hereinbefore defined, in the α -position of DHA. The use of chiral oxazolidin-2-ones has proved to be a particularly effective
30 methodology. The enolates derived from chiral *N*-acyloxazolidines can be quenched with a variety of electrophiles in a highly stereoregulated manner (Ager, Prakash, Schaad 1996).

Examples

35 The invention will now be described in more detail by the following examples, which are not to be constructed as limiting the invention. In the examples the structures were verified by Mass Spectrometry (MS). It should be pointed out that

the fatty acid derivatives may also be produced from low and medium DHA-containing starting material (i.e. about 40-60 w% DHA).

Synthesis protocols

5

Preparation of α -methyl DHA EE (PRB-1)

Butyllithium (228 ml, 0.37 mol, 1.6 M in hexane) was added dropwise to a stirred solution of diisopropylamine (59.5 ml, 0.42 mol) in dry THF (800 ml) under N₂ at 0 °C. The resulting solution was stirred at 0 °C for 30 min., cooled to -78 °C and stirred an additional 30 min. before dropwise addition of DHA EE (100 g, 0.28 mol) in dry THF (500 ml) during 2 h. The dark-green solution was stirred at -78 °C for 30 min. before MeI (28 ml, 0.45 mol) was added. The solution was allowed to reach -20 °C during 1.5 h, then poured into water (1.5 l) and extracted with heptane (2 x 800 ml). The combined organic phases were washed with 1 M HCl (1 l), dried (Na₂SO₄), filtered and evaporated *in vacuo*. The product was purified by dry flash chromatography on silica gel eluting with heptane/EtOAc (99:1) to give 50 g (48%) of the titled compound as a slightly yellow oil;

¹H-NMR (200MHz, CDCl₃) δ 1.02 (t, *J* 7.5 Hz, 3H), 1.20 (d, *J* 6.8 Hz, 3H), 1.29 (t, *J* 7.1 Hz, 3H), 2.0-2.6 (m, 5H), 2.8-3.0 (m, 10H), 4.17 (t, *J* 7.1 Hz, 2H), 5.3-5.5 (m, 12H);

MS (electrospray); 393 [M+Na].

Preparation of α -ethyl DHA EE (PRB-2)


Butyllithium (440 ml, 0.67 mol, 1.6 M in hexane) was added dropwise to a stirred solution of diisopropylamine (111 ml, 0.78 mol) in dry THF (750 ml) under N₂ at 0 °C. The resulting solution was stirred at -78 °C for 45 min. before dropwise addition of DHA EE (200 g, 0.56 mol) in dry THF (1.6 l). The addition of the ester was complete in 4 hours. The dark-green solution was stirred at -78 °C for 30 min. before EtI (65 ml, 0.81 mol) was added. The solution was allowed to reach -40 °C before an additional amount of EtI (5 ml, 0.06 mol) was added, and finally reach -15 °C (during 3 hours from -78 °C) before the mixture was poured into water and extracted with hexane (2x). The combined organic phases were washed with 1 M HCl, water, dried (Na₂SO₄), filtered and evaporated *in vacuo*. The product was purified by flash chromatography on silica gel eluting with heptane/EtOAc (99:1 followed by 50:1) to give 42.2 g (20%) of the titled compound as a yellow oil;

^1H -NMR (200 MHz; CDCl_3) δ 0.8-1.0 (m, 6H), 1.2-1.4 (m, 4H), 1.5-1.7 (m, 2H), 2.12 (m, 2H), 2.3-2.5 (m, 2H), 2.8-3.0 (m, 10H), 4.18 (t, J 7.1 Hz, 2H), 5.3-5.6 (m, 12H);

MS (electrospray); 407 $[\text{M}+\text{Na}]$.

5

Preparation of α -ethoxy-DHA ethylester (PRB-3)



10 To a suspension of 60 % NaH (84.1 mg, 2.1 mmol) in THF, 5 mL, at -78°C under N_2 -atmosphere was added drop wise a solution of α -hydroxy-DHA ethyl ester (**PRB-12**) (372 mg, 1.00 mmol) in THF, 5 mL, the resulting mixture was stirred at -78°C for 20 minutes before ethyl iodide (0.24 mL, 3.01 mmol) was added drop wise. The reaction mixture was gradually warmed to room temperature over night.

15 Saturated aqueous NH_4Cl , 15 mL, was added and the mixture was extracted with diethyl ether, 25 mL x 2, the organic phase was washed with brine, 25 mL, dried (Na_2SO_4) filtered, evaporated *in vacuo* and subjected to flash chromatography on silica gel eluting with heptane/EtOAc (95:5) to yield 68 mg (17 %) of the product as a yellow liquid.

20 ^1H NMR (200 MHz, CDCl_3) δ 0.94 (t, J =7.5 Hz, 3H), 1.16-1.29 (m, 6H), 2.05 (quint, J =7.2 Hz, 2H), 2.50 (m, 2H), 2.76-2.84 (m, 10H), 3.33-3.48 (m, 1H), 3.53-3.71 (m, 1H), 3.83 (dd, J =6.8 Hz, J =6.2 Hz, 1H), 4.18 (q, J =7.1 Hz, 2H), 5.31-5.45 (m, 12 H)

^{13}C NMR (50 MHz, CDCl_3) δ 14.2, 15.1, 20.5, 25.5, 25.6, 25.7, 31.0, 60.8, 25 66.0, 78.7, 124.1, 127.0, 127.8, 127.9, 128.0 (2 signals), 128.2 (2 signals), 128.5, 130.7, 132.0, 172.5 (3 signals hidden)

MS (electrospray); 423 $[\text{M}+\text{Na}]^+$

Preparation of α -fluoro DHA EE (PRB-4)

30 LDA (2.1 mL, 4.2 mmol, 2 M in THF/heptane/ethylbenzene) in dry THF (10 mL) under N_2 at -78°C was dropwise added DHA EE (1 g, 2.8 mmol) in dry THF (30 mL) during 15 min. NFSi (1.06 g, 3.4 mmol) was then added. The solution was allowed to reach RT and stirred for 70 hours. The mixture was poured into water and extracted with hexane (2x). The combined organic phases were washed with 1 M HCl, water, 35 dried (Na_2SO_4), filtered and evaporated *in vacuo*; MS (electrospray); 397 $[\text{M}+\text{Na}]$.

Preparation of α,α -dimethyl DHA EE (PRB-5)

Butyllithium (100 ml, 0.17 mol, 1.6 M in hexane) was added dropwise to a stirred solution of diisopropylamine (28 ml, 0.20 mol) in dry THF (100 ml) under N₂ at 0 °C. The resulting solution was stirred at 0 °C for 30 min., cooled to -78 °C and dropwise added a solution of DHA EE (50 g, 0.14 mol) in dry THF (200 ml). The resulting dark-green solution was stirred at -78 °C for 30 min. before MeI (17 ml, 0.28 mol) was added. The solution was allowed to reach -10 °C, then poured into water and extracted with hexane (2x). The combined organic phases were washed with 1 M HCl, dried (Na₂SO₄), filtered and evaporated *in vacuo*.

The procedure was repeated, but the crude product of α -methyl DHA EE was used instead of DHA EE. The product was purified by dry flash chromatography on silica gel eluting with heptane/EtOAc (99:1 followed by 98:2) to give 31.6 g (59%) of the titled compound as a slightly yellow oil;

¹H-NMR (200 MHz; CDCl₃) δ 1.01 (t, *J* 7.5 Hz, 3H), 1.21 (s, 6H), 1.28 (t, *J* 7.1 Hz, 3H), 2.08 (m, 2H), 2.34 (d, *J* 6.8 Hz, 2H), 2.8-3.0 (m, 10H), 4.15 (q, *J* 7.5 Hz, 2H), 5.3-5.6 (m, 12H);

¹³C-NMR (50 MHz; CDCl₃) δ 14.7, 21.0, 25.3, 26.0, 26.1, 38.3, 42.8, 60.7, 125.8, 127.4, 128.3, 128.5, 128.6, 128.7, 129.0, 130.7, 132.4, 177.9;

MS (electrospray); 385 [M+H].

Preparation of α -thiomethyl DHA (PRB-6)

α -Iodo DHA EE (0.5 g, 1.04 mmol) dissolved in 20 mL THF at 0°C under N₂. MeSNa (80 mg, 1.14 mmol) was added the reaction and the mixture was allowed to stir for a few minutes before it was diluted with heptane. The organic phase was washed with water (2x) dried (Na₂SO₄) and evaporated *in vacuo*. The desired product was isolated by flash chromatography Heptan/EtOAc (30:1) to give α -thiomethyl DHA EE as a pale yellow oil. The α -thiomethyl DHA EE was dissolved in 10 mL EtOH and 10 mL THF. The solution was added LiOH (0.39 g, 9.2 mmol) dissolved in 5 mL water. The reaction mixture was allowed to stir overnight at RT, before diluting with water and heptane. The organic fraction was extracted with 1M LiOH (2x) and the combined aqueous phases was acidified with 5M HCl and extracted with diethyl ether (2x). The combined organic phases was washed with brine, water, dried (Na₂SO₄) and evaporated *in vacuo* to give 183 mg (47 %) of the title compound as a pale yellow oil;

¹H-NMR (200MHz, CDCl₃) δ 0.98 (t, *J* 6.6Hz, 3H), 1.95-2.65 (m, 7H), 2.72-3.05 (m, 10H), 3.12-3.43 (m, 1H), 5.20-5.70 (m, 12H), 10.65 (br s, 1H);

$^{13}\text{H-NMR}$ (50MHz, CDCl_3) δ 14.7, 21.0, 25.9, 26.0, 26.2, 28.8, 125.4, 127.4, 128.1, 128.3, 128.4, 128.7, 128.9, 129.0, 131.6, 132.4, 177.0.

Preparation of α -thioethyl DHA EE (PRB-7)

- 5 α -Iodo DHA EE (11g, 23 mmol) dissolved in 100 mL THF under N_2 at 0°C . EtSNa (2.1 g, 25 mmol) was added the solution and was allowed to stir for 1 hour at 0°C . The reaction was quenched with 1M HCl and diluted with Heptan. The organic phase was washed with water (2x), dried (Na_2SO_4) and evaporated *in vacuo*. The desired product was isolated by flash chromatography Heptan/EtOAc (30:1) to give
- 10 7.3 g (76 %) of the title compound as a pale yellow oil;
- $^1\text{H-NMR}$ (200MHz, CDCl_3) δ 1.1-1.3 (m, 9H), 2.05 (m, 2H), 2.3-2.7 (m, 4H), 2.7-2.9 (m, 10H), 3.25 (m, 1H), 4.17 (q, J 7.1 Hz, 2H), 5.3-5.5 (m, 12H);
- MS (electrospray): 439 [M+Na].

15 **Preparation of α,α -diethyl DHA EE (PRB-8):**

- Butyllithium (38.6 ml, 0.62 mol, 1.6 M in hexane) was added dropwise to a stirred solution of diisopropylamine (9.1 ml, 0.65 mol) in dry THF (200 ml) under N_2 at 0°C . The resulting solution was stirred at 0°C for 30 min., cooled to -78°C and dropwise added a solution of DHA EE (20.0 g, 0.56 mol) in dry THF (100 ml). The
- 20 resulting dark-green solution was stirred at -78°C for 30 min., before EtI (6.8 ml, 0.84 mol) was added. The solution was allowed to reach -10°C , then poured into water and extracted with hexane (2x). The combined organic phases were washed with 1 M HCl, dried (Na_2SO_4), filtered and evaporated *in vacuo*.

- The procedure was repeated, but the crude product of α -ethyl DHA EE was
- 25 used instead of DHA EE. The reaction mixture after addition of EtI was allowed to reach ambient temperature and was stirred over night. The product was purified by dry flash chromatography on silica gel eluting with heptane/EtOAc (99:1 followed by 98:2) to give 10.0 g (43%) of the titled compound as a slightly yellow oil;

- $^1\text{H-NMR}$ (200 MHz; CDCl_3) δ 0.83 (t, J 7.4 Hz, 6H), 0.94 (t, J 5.8 Hz, 3H),
- 30 1.28 (t, J 7.1 Hz, 3H), 1.63 (q, J 7.4 Hz, 4H), 2.10 (m, 2H), 2.34 (d, J 6.9 Hz, 2H), 2.8-3.0 (m, 10H), 4.15 (q, J 7.5 Hz, 2H), 5.3-5.6 (m, 12H);

$^{13}\text{C-NMR}$ (50 MHz; CDCl_3) δ 8.9, 14.7, 21.0, 23.1, 25.9, 26.0, 26.2, 27.4, 31.2, 50.1, 60.6, 125.5, 127.4, 128.3, 128.6, 128.9, 130.5, 132.4, 177.1;

MS (electrospray); 413.3 [M+H], 435.3 [M+Na].

Preparation of α -benzyl DHA EE (PRB-9):

To a stirred solution of diisopropyl amine (0.91 mL, 6.46 mmol) in dry THF (20 mL) under inert atmosphere held at 0°C was added drop wise n-BuLi (1.6 M in hexanes, 3.86 mL, 6.18 mmol). The mixture was stirred at 0°C for 30 minutes, given -
5 78°C and stirred at this temperature for five minutes. DHA EE (2.0 g, 5.62 mmol) in dry THF (10 mL) was added drop wise and the mixture was stirred at -78°C for 20 minutes, then benzyl bromide (0.80 mL, 6.74 mmol) was added. The resulting solution was allowed to reach 0°C over three hours, portioned between water (100 mL) and heptane (100 mL). The aqueous layer was extracted with heptane (50 mL)
10 and the combined organic layer was washed with 1M HCl and dried (Na₂SO₄). Concentration under reduced pressure and purification by flash chromatography (Heptane : EtOAc 99:1) afforded 1.05 g (42%) of the title compound as a colorless oil;

¹H-NMR (200 MHz, CDCl₃): δ 0.99 (t, 3H), 1.18 (t, 3H), 2.08-2.16 (m, 2H),
15 2.35-2.42 (m, 2H), 2.74-2.98 (m, 13H), 4.09 (q, 4H), 5.38-5.50 (m, 10H), 7.19-7.36 (m, 5H);

¹³C-NMR (50 MHz, CDCl₃): δ 14.61, 14.71, 20.99, 25.98, 26.07, 30.07, 38.32, 48.02, 60.88, 126.75, 126.83, 127.46, 128.31, 128.45, 128.53, 128.58, 128.86, 128.77, 129.01, 129.35, 130.55, 132.46, 138.89, 175.39.

20 MS (electrospray): 447.3 [M+H], 469.3 [M+Na].


Preparation of α -ethanesulfinyl DHA EE (PRB-10)

To a solution of α -thioethyl DHA EE (0.5 g, 1.3 mmol) in 15 mL CHCl₃ held at -20 °C under inert atmosphere was added a solution of MCPBA (0.22 g, 1.3 mmol)
25 in 10 mL CHCl₃. The reaction mixture was stirred for 2h at this temperature, filtered and washed with a saturated aqueous solution of NaHCO₃. The aqueous phase was extracted twice with CHCl₃ and the combined organic phase was washed with water and brine, dried with Na₂SO₄, filtered and concentrated. The product was isolated from residual material after flash chromatography using hexane: EtOAc 8:2 to afford
30 0.35 g (70%) of the title compound.

¹H NMR (200 MHz, CDCl₃): δ 0.99 (t, 3H), 1.27-1.45 (m, 6H), 2.09 (m, 2H), 2.79-2.94 (m, 14H), 3.55 (m, 1H), 4.25 (q, 2H), 5.37-5.59 (m, 12H).

¹³C NMR (50 MHz, CDCl₃): δ 7.97, 14.58, 14.68, 20.95, 23.68, 25.17, 25.93, 26.04, 44.20, 45.15, 62.30, 64.08, 123.91, 124.47, 127.41, 127.86, 128.26, 128.40,
35 128.44, 128.72, 128.72, 128.96, 129.12, 132.42, 132.47, 174.55.

MS (electrospray): 455.3 [M+Na].

Preparation of α -thiophenyl-DHA ethylester (PRB-11)


To a solution of α -iodo-DHA ethylester (PRB-15) (3.40 g, 7.05 mmol) in acetone, 20 mL, a solution of sodium phenyl sulfide (1.039 g, 7.86 mmol) in acetone, 110 mL, was added drop wise. The resulting mixture was stirred at ambient temperature for 1 ½ hrs, evaporated *in vacuo* and subjected to flash chromatography on silica gel eluting with heptane/EtOAc 200:1 - 95:5 to yield 2.35 g (72 %) of the product as a yellow liquid.

^1H NMR (200 MHz, CDCl_3) δ 0.97 (t, $J=7.5$ Hz, 3H), 1.18 (t, $J=7.1$ Hz, 3H), 2.09 (quint, $J=7.1$ Hz, 2H), 2.54-2.66 (m, 2H), 2.83-2.86 (m, 10 H), 3.67 (dd, $J=6.8$ Hz, $J=8.3$ Hz, 1 H), 4.12 (q, $J=7.1$ Hz, 2H), 5.24-5.49 (m, 12 H), 7.28-7.33 (m, 3H), 7.46-7.50 (m, 2H)

^{13}C NMR (50 MHz, CDCl_3) δ 14.0, 14.2, 20.5, 25.5, 25.6, 25.7, 29.4, 50.6, 61.1, 125.1, 127.0, 127.7, 127.9, 128.0, 128.3, 128.42, 128.45, 128.9, 131.2, 132.0, 133.0, 133.2, 174.1 (5 signals hidden)

MS (electrospray); 465 $[\text{M}+\text{H}]^+$, 487 $[\text{M}+\text{Na}]^+$

HRMS (EI) calculated for $\text{C}_{30}\text{H}_{40}\text{O}_2\text{S}$: 464.2749, found: 464.2741

Preparation of α -hydroxy-DHA ethylester (PRB-12)

To a solution of diisopropyl amine (19.76 mL, 140 mmol) in dry THF, 40 mL, under N_2 -atmosphere at -78°C was added drop wise 1.6 M BuLi in hexane (87.5 mL, 140 mmol). The resulting mixture was stirred at -78°C for 15 minutes before a solution of DHA ethylester (24.99 g, 70.1 mmol) in THF, 80 mL, was added drop wise. The resulting dark green reaction mixture was stirred for 1 hour at -78°C before triethylphosphite (12.2 mL, 70.1 mmol) was added drop wise and then O_2 was bubbled through the reaction mixture over night while the reaction mixture was kept at -78°C for 5 hrs and then slowly warmed to room temperature. Saturated aqueous NaHCO_3 , 100 mL, was added and the mixture was extracted with diethyl ether, 200

mL x 2. The organic phase was dried (Na_2SO_4), filtered and evaporated *in vacuo* and subjected to flash chromatography on silica gel eluting with heptane/EtOAc 99:1 - 95:5 to yield 4.52 g (17 %) of the product as a yellow liquid.


^1H NMR (200 MHz, CDCl_3) δ 0.92 (t, $J=7.5$ Hz, 3H), 1.24 (t, $J=7.1$ Hz, 3H),
 5 2.02 (quint, $J=7.1$ Hz, 2H), 2.44-2.54 (m, 2H), 2.74-2.87 (m, 10 H), 4.13-4.24 (m, 3H), 5.25-5.94 (m, 12H)

^{13}C NMR (50 MHz, CDCl_3) δ 14.0, 14.1, 20.4, 25.4, 25.5, 25.6, 32.0, 61.5, 69.9, 123.3, 126.9, 127.7, 127.9, 128.08, 128.1, 128.2, 128.4, 131.3, 131.8, 174.4 (4 signals hidden)

10 MS (electrospray); 395 $[\text{M}+\text{Na}]^+$

HRMS (ES) calculated for $\text{C}_{24}\text{H}_{36}\text{O}_3\text{Na}$: 395.2556, found: 395.2543

Preparation of α -methyl-DHA amide (PRB-13)




A solution of α -methyl-DHA (PRB-1 FA) (3.13 g, 9.1 mmol) and oxalyl chloride (8.0 mL, 94.5 mmol) in toluene, 90 mL, was added DMF, 0.1 mL, and the resulting mixture was stirred at ambient temperature under N_2 -atmosphere for 15 ½
 20 hours. The mixture was then evaporated *in vacuo* and the residue was dissolved in THF, 100 mL, cooled to 0°C and aqueous NH_3 (20 mL) was added drop wise. The ice-bath was removed and the mixture was stirred at ambient temperature for 4 hours, water, 50 mL, was added and the aqueous phase was extracted with diethyl ether, 2x100 mL. The organic phase was washed with saturated aqueous NH_4Cl , 50 mL,
 25 dried (Na_2SO_4), filtered and evaporated *in vacuo* and subjected to flash chromatography on silica gel eluting with $\text{CH}_2\text{Cl}_2/2\text{M NH}_3$ in MeOH 97.5:2.5 to yield 2.51 g (80 %) of the product as a yellow liquid.

^1H NMR (200 MHz, CDCl_3) δ 0.91 (t, $J=7.5$ Hz, 3H), 1.10 (d, $J=9.8$ Hz, 3H), 1.94-2.11 (m, 3H), 2.19-2.35 (m, 2H), 2.76-2.77 (m, 10 H), 5.18-5.45 (m, 12 H), 6.03
 30 (s, 1H), 6.72 (s, 1H)

^{13}C NMR (50 MHz, CDCl_3) δ 14.6, 17.6, 20.8, 25.8, 25.9, 32.0, 41.0, 127.3, 128.1, 128.4, 128.6, 128.8, 130.1, 132.2, 179.6 (8 signals hidden)

MS (electrospray); 342 $[\text{M}+\text{H}]^+$, 364 $[\text{M}+\text{Na}]^+$

HRMS (EI) calculated for $\text{C}_{23}\text{H}_{35}\text{NO}$: 341.2719, found: 341.2707

Preparation of α -methoxy-DHA ethylester (PRB-14)

5

To a suspension of 60 % NaH (61.1 mg, 1.53 mmol) in THF, 5 mL, at -78°C under N₂-atmosphere was added drop wise a solution of α -hydroxy-DHA ethyl ester (PRB-12) (373 mg, 1.00 mmol) in THF, 5 mL, the resulting mixture was stirred at -78°C for 20 minutes before methyl iodide (0.13 mL, 2.09 mmol) was added drop wise. The reaction mixture was gradually warmed to room temperature for 5 hrs. Saturated aqueous NH₄Cl, 15 mL, was added and the mixture was extracted with diethyl ether, 25 mL x 2, the organic phase was washed with brine, 25 mL, dried (Na₂SO₄) filtered, evaporated *in vacuo* and subjected to flash chromatography on silica gel eluting with heptane/EtOAc 99:1 – 4:1 to yield 136 mg (35 %) of the product as a yellow liquid.

15

¹H NMR (200 MHz, CDCl₃) δ 0.92 (t, *J*=7.5 Hz, 3H), 1.24 (t, *J*=7.1 Hz, 3H), 2.03 (quint, *J*=7.3 Hz, 2 H), 2.48 (t, *J*=5.7 Hz, 2H), 2.73-2.82 (m, 10 H), 3.34 (s, 3H), 3.74 (t, *J*=6.2Hz, 1H), 4.17 (q, *J*=7.1 Hz, 2 H), 5.24-5.43 (m, 12H)

20

¹³C NMR (50 MHz, CDCl₃) δ 14.1, 20.4, 25.4, 25.5, 25.7, 30.6, 57.9, 60.9, 80.8, 123.7, 126.9, 127.71, 127.73, 127.92, 127.94, 128.07, 128.1, 128.2, 128.4, 130.7, 131.8, 171.9 (3 signals hidden)


MS (electrospray); 409 [M+Na]⁺

HRMS (ES) calculated for C₂₅H₃₈O₃Na: 409.2713, found: 409.2711

Preparation of α -iodo DHA EE (PRB-15)

Diisopropylamine (20 mL, 140 mmol) was dissolved in 150 mL THF under N₂ at -20°C. *n*-BuLi (88mL, 140 mmol, 1.6 M) was added dropwise to the mixture before the solution was cooled to -78°C. DHA EE (50 g, 140 mmol) in 250 mL THF was added dropwise to the solution and the reaction mixture was stirred for 30 min at RT. The resulting mixture was added dropwise to a solution of I₂ (42.8 g, 169 mmol) in 400 mL THF under N₂ at -78°C. The reaction was quenched with 1M HCl and diluted with Heptan. The organic phase was washed with 10% Na₂S₂O₃ (2x), dried (Na₂SO₄), filtered and evaporated *in vacuo*. The desired product was isolated by flash chromatography Heptan/EtOAc (100:1) to give 11.0 g (16 %) of the title compound as a pale yellow oil; MS (Electrospray): 505 [M+Na].

35


Preparation of α -iodo-DHA ethylester (PRB-15)

To a solution of diisopropyl amine (42 mL, 298 mmol) in dry THF, 150 mL, under N₂-atmosphere at -78°C was added drop wise 1.6 M BuLi in hexane (158 mL, 253 mmol). The resulting mixture was stirred at -78°C for 35 minutes before a solution of DHA ethylester (75.05 g, 210 mmol) in THF, 300 mL, was added drop wise. The resulting dark green reaction mixture was stirred for 30 minutes at -78°C before a solution of I₂ (91.06 g, 359 mmol) in THF, 200 mL was added drop wise. The reaction mixture was stirred at -78°C for 20 minutes before it was quenched with water, 200 mL, and extracted with heptane, 300 mL. The organic phase was washed with 1 M HCl, 150 mL, and water, 200 mL, dried (Na₂SO₄), filtered and evaporated *in vacuo*. The resulting crude product was subjected to flash chromatography on silica gel eluting with heptane/EtOAc (100:1) yielding 26.14 g (26%) of the product as a yellow liquid.

¹H NMR (200 MHz, CDCl₃) δ 0.94 (t, *J*=7.5 Hz, 3H), 1.24 (t, *J*=7.1 Hz, 3H), 2.04 (quint, *J*=7.1 Hz, 2H), 2.69-2.84 (m, 12 H), 4.17 (q, *J*=7.1 Hz, 2H), 4.22 (t, *J*=7.9 Hz, 1H), 5.24-5.49 (m, 12 H)

¹³C NMR (50 MHz, CDCl₃) δ 13.7, 14.2, 25.5, 26.0 (2 signals), 25.8, 34.0, 61.7, 126.1, 127.0, 127.4, 127.8, 127.9, 128.0, 128.2, 128.5, 128.5, 131.6, 131.9, 170.9 (4 signals hidden)

MS (electrospray); 505 [M+Na]⁺

Preparation of α -amino-DHA etylester (PRB-17)

A solution of α -phtalimide-DHA ethylester (313.5 mg, 0.62 mmol) in EtOH, 5 mL, was added hydrazine hydrate (46 μ L, 0.95 mmol) and the resulting mixture was

refluxed under N₂-atmosphere for 15 ½ hrs followed by evaporation *in vacuo* and flash chromatography on silica gel eluting with CH₂Cl₂:7M NH₃ in MeOH (99:1-95:1) to yield 149 mg (64 %) of the product as a yellow liquid.


¹H NMR (200 MHz, CDCl₃) δ 0.91 (t, *J*=7.5 Hz, 3H), 1.22 (t, *J*=7.1 Hz, 3H),
 5 1.72 (bs, 2H), 2.02 (quint., *J*=7.2 Hz, 2H), 2.39-2.46 (m, 2H), 2.73-2.82 (m, 10 H),
 3.47 (bs, 1H), 4.13 (q, 2H), 5.23-5.56 (m, 12 H)

¹³C NMR (50 MHz, CDCl₃) δ 14.1, 20.4, 25.4, 25.5, 25.6, 54.1, 60.8, 124.4,
 126.9, 127.7 (2 signals), 127.9, 128.2, 128.3, 128.4, 131.4, 131.9, 189.3 (6 signals
 hidden)

10 MS (electrospray); 372 [M+H]⁺

Preparation of (S)-(+)-α-ethyl DHA EE (PRB-20):


Synthesis of intermediate PRB-18:



DHA (3.00g, 18.3 mmol) was dissolved in dry CH₂Cl₂ (120 mL) held at 0°C under inert atmosphere and added DMAP (2.45 g, 20.1 mmol) and DCC (3.96 g, 19.2 mmol). The mixture was stirred at 0°C for 20 minutes, added (4R,5S)-(+)-4-methyl-5-phenyl-2-oxazolidinone (3.24 g, 18.3 mmol) and stirred at ambient temperature for 20 hours. Filtration and purification by flash chromatography (heptane : EtOAc 6:1) afforded 3.00 g (34%) of intermediate PRB-18 as a colorless oil.

¹H-NMR (200 MHz, CDCl₃): δ 0.93-1.05 (t+d, 6H), 2.11 (m, 2H), 2.51 (m, 2H), 2.80-3.00 (m, 10H), 3.05 (m, 2H), 4.77 (m, 1H), 5.34-5.68 (m, 12H), 5.70 (d, 1H), 7.28-7.32 (m, 2H), 7.37-7.47 (m, 3H).

Synthesis of intermediate PRB-19:




PRB-18 (1.80 g, 3.70 mmol) in dry THF (10 mL) was added drop wise to a solution of LiHMDS (1M in THF, 4.00 mL, 4.00 mmol) in dry THF (15 mL) held at -78°C under inert atmosphere. The mixture was stirred at -78°C for 30 minutes, added EtI (0.89 mL, 11.1 mmol) and slowly given 0°C over one hour. The mixture was then stirred at 0°C for 18 hours and portioned between saturated NH₄Cl (50 mL) and diethyl ether (50 mL). The aqueous layer was extracted with diethyl ether (50 mL) and the combined organic layer was washed with 0.1 M HCl (50 mL) and brine (50 mL). Drying (Na₂SO₄) and purification by flash chromatography (heptane : EtOAc 95:5) afforded 0.52 g (27 %) of intermediate PRB-19 as a colorless oil.

¹H-NMR (200 MHz, CDCl₃): δ 0.88-1.01 (m, 9H), 1.64-1.78 (m, 2H), 2.08 (m, 2H), 2.31 (m, 1H), 2.48 (m, 1H), 2.87 (m, 10H), 3.87 (m, 1H), 4.75 (m, 1H), 5.32 (m, 12H), 5.63 (d, *J* 7.1 Hz, 1H), 7.32 (m, 2H), 7.42 (m, 3H).

¹³C-NMR (50 MHz, CDCl₃): δ 7.26, 11.75, 14.67, 14.98, 20.95, 25.57, 25.93, 26.04, 29.93, 44.59, 55.31, 79.10, 125.21, 126.01, 127.17, 127.42, 128.27, 128.50, 128.55, 128.67, 128.95, 129.09, 130.35, 132.42, 133.80, 153.18, 176.25.

MS (electrospray): 538.2 [M+Na]



PRB-19 (0.25 g, 0.485 mmol) was dissolved in abs EtOH (5 mL) and given 0°C under inert atmosphere. NaOEt (1M in EtOH, 0.54 mL, 0.54 mmol) was added and the mixture was stirred at 0°C for 30 minutes and portioned between water and heptane. The aqueous layer was extracted with heptane and the combined organic layer was washed with 0.1 M HCl and dried. Purification by flash chromatography afforded 0.025 g (13 %) of the title compound PRB-20 as a colorless oil.

44

$^1\text{H-NMR}$ (200 MHz; CDCl_3) δ 0.8-1.0 (m, 6H), 1.2-1.4 (m, 4H), 1.5-1.7 (m, 2H), 2.12 (m, 2H), 2.3-2.5 (m, 2H), 2.8-3.0 (m, 10H), 4.18 (t, 2H), 5.3-5.6 (m, 12H).


MS (electrospray); 407 $[\text{M}+\text{Na}]$.

$[\alpha]_{\text{D}} +1.7^\circ$ ($c=1.5$, ethanol).

5

Preparation of (R)-(-)- α -ethyl DHA EE (PRB-23):

Synthesis of intermediate PRB-21:



10


DHA (1.00g, 3.05 mmol) was dissolved in dry CH_2Cl_2 (20 mL) held at 0°C under inert atmosphere and added DMAP (0.41 g, 3.35 mmol) and DCC (0.66g, 3.20 mmol). The mixture was stirred at 0°C for 20 minutes, added (4S,5R)-(-)-4-methyl-5-phenyl-2-oxazolidinone (0.54 g, 3.05 mmol) and stirred at ambient temperature for 20 hours. Filtration and purification by flash chromatography (heptane : EtOAc 6:1) afforded 1.08 g (73%) of intermediate PRB-21 as a colorless oil.

15

$^1\text{H-NMR}$ (200 MHz, CDCl_3): δ 0.93-1.05 (t+d, 6H), 2.11 (m, 2H), 2.51 (m, 2H), 2.80-3.00 (m, 10H), 3.05 (m, 2H), 4.77 (m, 1H), 5.34-5.68 (m, 12H), 5.70 (d, 1H), 7.28-7.32 (m, 2H), 7.37-7.47 (m, 3H).

20

Synthesis of intermediate PRB-22:



25


PRB-21 (3.25g, 6.67 mmol) in dry THF (15 mL) was added drop wise to a solution of LiHMDS (1M in THF, 7.34 mL, 7.34 mmol) in dry THF (35 mL) held at -

78°C under inert atmosphere. The mixture was stirred at -78°C for 30 minutes, added EtI (1.6 mL, 20.0 mmol) and slowly given 0°C over one hour. The mixture was then stirred at 0°C for 18 hours and portioned between saturated NH₄Cl (50 mL) and diethyl ether (50 mL). The aqueous layer was extracted with diethyl ether (50 mL) and the combined organic layer was washed with 0.1 M HCl (50 mL) and brine (50 mL). Drying (Na₂SO₄) and purification by flash chromatography (heptane : EtOAc 95:5) afforded 1.50 g (44 %) of intermediate PRB-22 as a colorless oil.

¹H-NMR (200 MHz, CDCl₃): δ 0.88-1.01 (m, 9H), 1.64-1.78 (m, 2H), 2.08 (m, 2H), 2.31 (m, 1H), 2.48 (m, 1H), 2.87 (m, 10H), 3.87 (m, 1H), 4.75 (m, 1H), 5.32 (m, 12H), 5.63 (d, *J* 7.1 Hz, 1H), 7.32 (m, 2H), 7.42 (m, 3H).

¹³C-NMR (50 MHz, CDCl₃): δ 7.26, 11.75, 14.67, 14.98, 20.95, 25.57, 25.93, 26.04, 29.93, 44.59, 55.31, 79.10, 125.21, 126.01, 127.17, 127.42, 128.27, 128.50, 128.55, 128.67, 128.95, 129.09, 130.35, 132.42, 133.80, 153.18, 176.25.

MS (electrospray): 538.2 [M+Na]




PRB-22 (0.25 g, 0.485 mmol) was dissolved in abs EtOH (5 mL) and given 0°C under inert atmosphere. NaOEt (1M in EtOH, 0.54 mL, 0.54 mmol) was added and the mixture was stirred at 0°C for 30 minutes and portioned between water and heptane. The aqueous layer was extracted with heptane and the combined organic layer was washed with 0.1 M HCl and dried. Purification by flash chromatography afforded 0.025 g (13 %) of the title compound PRB-23 as a colorless oil.

¹H-NMR (200 MHz; CDCl₃) δ 0.8-1.0 (m, 6H), 1.2-1.4 (m, 4H), 1.5-1.7 (m, 2H), 2.12 (m, 2H), 2.3-2.5 (m, 2H), 2.8-3.0 (m, 10H), 4.18 (t, 2H), 5.3-5.6 (m, 12H);

MS (electrospray); 407 [M+Na].

[α]_D -1.3° (c=1.00, ethanol).

Preparation of α -phthalimide-DHA ethylester (PRB-24)


5 A mixture of the α -hydroxy-DHA ethyl ester (**PRB-12**) (373.5 mg, 1.00 mmol), phthalimide (178 mg, 1.21 mmol) and triphenyl phosphine (313.9 mg, 1.20 mmol) in THF, 10 mL, was cooled to 0°C under N_2 -atmosphere before diisopropyl azodicarboxylate (0.24 mL, 1.24 mmol) was added drop wise. The ice-bath was removed and the reaction mixture was stirred at ambient temperature for 18 hrs, whereupon it was evaporated *in vacuo* and subjected to flash chromatography on silica gel eluting with heptane/EtOAc (99:1-95:1) to yield 323 mg (64 %) of the product as a yellow liquid.

^1H NMR (200 MHz, CDCl_3) δ 0.95 (t, $J=7.5$ Hz, 3H), 1.22 (t, $J=7.1$ Hz, 3H), 2.05 (m, 2H), 2.72-2.84 (m, 11H), 3.02-3.22 (1H), 4.20 (q, $J=7.1$ Hz, 2H), 4.87 (dd, $J=11$ Hz, $J=4.9$ Hz, 1H), 5.17-5.40 (m, 12H), 7.68-7.75 (m, 2H), 7.79-7.85 (m, 2H)

^{13}C NMR (50 MHz, CDCl_3) δ 14.0, 14.1, 20.4, 25.4, 25.4, 25.5, 27.0, 51.8, 61.7, 123.8, 124.3, 126.9, 127.5, 127.7, 127.9, 127.9, 128.1, 128.1, 128.3, 128.4, 131.6, 131.8, 131.8, 134.0, 167.3, 168.7 (2 signals hidden)

MS (electrospray); 502 $[\text{M}+\text{H}]^+$, 524 $[\text{M}+\text{Na}]^+$

20

Preparation of α -ethylamino-DHA ethylester (PRB-25) and α -diethylamino-DHA ethylester (PRB-26)

25

A mixture of the α -amino-DHA ethylester (PRB-17) (746.5 mg, 2.01 mmol), $\text{LiOH}\cdot\text{H}_2\text{O}$ (171.6 mg, 4.09 mmol) and molsieve 4Å (599 mg) in DMF, 4 mL, was added ethylbromide (3.0 mL, 40.2 mmol) and the resulting mixture was stirred at ambient temperature for 71 hrs. The mixture was diluted with diethyl ether, 100 mL, and filtered. The organic phase was washed with 1 M NaOH, 20 mL, and brine, 20 mL, dried (Na_2SO_4), filtered and evaporated *in vacuo* and subjected to flash chromatography on silica gel eluting with heptane:EtOAc (95:5) - CH_2Cl_2 :2M NH_3 in

30

MeOH (99:1) to yield 458 mg (53 %) of PRB-26 as a yellow liquid and 152 mg (19%) of PRB-25 as a yellow liquid.

PRB-26:

¹H NMR (200 MHz, CDCl₃) δ 0.89 (t, *J*=7.5 Hz, 3H), 1.03 (t, 3H), 1.24 (t, *J*=7.1 Hz, 6H), 2.05 (quint, *J*=7.1 Hz, 2H), 2.52 (m, 4H), 2.76-2.85 (m, 12 H), 3.35 (t, 1H), 4.13 (q, *J*=7.1 Hz, 2 H), 5.28-5.44 (m, 12 H)

¹³C NMR (75 MHz, CDCl₃) δ 14.1, 14.3, 14.4, 20.5, 22.6, 25.5, 25.6, 25.7, 31.9, 44.4, 60.1, 62.9, 127.0, 127.8, 128.05, 128.13, 128.17, 128.22, 128.5, 132.0, 173.3 (5 signals hidden)

10

Examples

An overview of the models and methods used in the present invention for demonstrating effects on the metabolic syndrome and type 2 diabetes are presented in Fig. 2: Five blocks of experiments have been performed in order to elucidate the effects of DHA derivatives for reduction of insulin resistance and/or alleviating the metabolic syndrom. The invention shall not be limited to the shown embodiments and examples.

Example 1. Analysis of intracellular free fatty acids (non-esterified fatty acids) in liver cells (block 1 in Fig. 2)

20

Background

In the first block of experiments (see Fig. 2) liver tissue from animals fed PRB-1,2,5, and 7 was analysed with respect to free unesterified fatty acids. The animals were recruited from the fifth block of experiments (pharmacodynamic effects of DHA derivatives in an animal model of metabolic syndrome). The animals had been given DHA (15% of fat content of the diet) or the DHA-derivatives (1,5% of the fat content in their diet) for 8 weeks and were supposed to be in a steady-state situation with stable levels of DHA and the DHA-derivatives intracellularly. Liver tissue was chosen due to the fact that the metabolism rate is very high in liver.

30

Method

The liver samples were homogenized in cold PBS buffer, and extracted immediately with chloroform: methanol (2:1) containing 0.2mM butylated hydroxytoluene (BHT) using cis-10-heptadecenoic acid as internal standard. The organic phases were dried under nitrogen, re-dissolved in acetonitrile with 0.1%

35

acetic acid and 10 μ M BHT for RP-HPLC MS/MS analysis. Total protein content was measured using Bio-Rad method after homogenization.

Agilent 1100 system was used for reverse phase column (Supelco Ascentis C₁₈ column, 25cm x 4.6mm, i.d. 5 μ m) separation of DHA and its PRB derivatives within 22min. The flow phase was iso-gradient acetonitrile-H₂O (87+13, v/v) containing 0.1% acetic acid. The column oven temperature was set at 35°C. The column elute was identified and quantified in the negative electrospray ionisation applying multiple reaction monitoring mode by triple tandem quadrupole mass/mass (ABI Qtrap-4000). The parent-daughter ion pairs were 327.3/327.3 (DHA), 341.3/341.3 (PRB-1), 355.3/355.3 (PRB-2 and PRB-5), 387.3 /387.3 (PRB-7), 267.2/267.2 (I.S. FA 17:1) respectively under unit resolution. The signal collection dwell time was all 100 msec except for FA 17:1 which was set at 200msec. Accurate verification of isomeric PRB compounds was done by combination of the retention time and characteristic mass/charge ratio. The quadratic regression standard curve was used for quantification after internal standard calibration.

Results

Concentration of the different DHA-derivatives and the concentrations of DHA was given as μ g per g of total amount of protein in the liver cells. Fig. 3 depicts the concentrations of the different PRBs from animals given PRB-1, 2, 5 and 7 in a concentration of 1.5% of total fat content in the high fat diet.

The highest intracellular concentration of the PRBs was obtained for PRB-2. Also PRB-5 was enriched intracellularly, although not to the same extent as PRB-2. This finding is unexpected.

Fig. 4 depicts the intracellular concentrations of DHA in liver tissue from animals given the different PRBs. DHA reached a significantly higher level in the animals given PRB-7 compared to the other three DHA-derivatives. Animals given PRB-2 had the lowest concentration of DHA. It seems that PRB-7 is to some extent converted back to DHA.

PRB-2 reached the highest intracellular concentration. This means that PRB-2 will be more available as a ligand to nuclear receptors, a pattern which could be translated into therapeutic effects in handling of blood glucose and blood lipids.

Example 2**Computer based affinity testing (block 2 in Fig. 2)****Background**

5 Nuclear receptors have been sequenced and the amino acid sequence is known for the PPARs and other relevant receptors engaged in the genetic control of glucose and fat. X-ray crystallography and NMR spectroscopy of the PPAR receptors are available and computerised affinity testing of fatty acids liganding to the receptors can be used to estimate binding kinetics. The binding geometrics, often called binding
10 modes or poses, include both positioning of the ligand relative to the receptor and the conformational state of the ligand and the receptor. Effective ligand docking can therefore be analysed.

Affinity of the ligand to the receptor is defined by two different parameters: docking of the ligand (DHA derivative) into the binding site of the receptor and
15 electrostatic bonding between certain amino acids of the receptor and the carboxyl group or side chains in the head of the fatty acid. (Krumrine).

As previously known, the PPAR α receptor is more promiscuous compared to PPAR γ , meaning that PPAR α will accept more fatty acids as ligands compared to PPAR γ . However, since patients with metabolic syndrome or type 2 diabetes are
20 usually obese or overweight and have pathologic blood lipids, mainly elevated triglycerides and low High-Density Cholesterol (HDL-chol) activation of the PPAR α receptor is important. An ideal drug for treatment of metabolic syndrome or type 2 diabetes should act as ligand to both these receptors, preferably with the highest affinity to the PPAR γ receptor.

25

Method

Ranking of the different DHA-derivatives according to their binding affinity was calculated and given as lowest binding affinity (LBA) and average binding affinity (ABE).

30 A total of 15 DHA derivatives (PRB-1 through PRB-15) were tested with the computerized docking method. Some of the derivatives, such as PRB-1, PRB-2, PRB-7, PRB-9, PRB-10, PRB-11, PRB-12, PRB-13, PRB-14 and PRB-15, are presented as r and s enantiomeres and in this case both were tested. The PPAR γ ligands rosiglitazone and pioglitazone, both in the r and s form, were also tested for
35 comparison. These compounds are registered pharmaceuticals for treatment of diabetes.

Results

The results are shown in Table 1, presenting the parameters Lowest binding energy of single confirmation (LBE), average binding energy (ABE) of the correctly posed confirmation and fraction of correctly posed confirmation of the ICM-saved 20 lowest energy confirmation (fbound) of the compounds tested. Affinity to the RXR α was tested in the same setting. The RXR α receptor interacts with the PPAR receptor forming a heterodimer by liganding of a fatty acid.

Fig. 5 depicts the binding affinities for the PPAR γ receptor, which is mainly engaged in the transcription of proteins engaged in handling of blood glucose. Clearly PRB-2 both in the r and the s stereoisomer forms had a good affinity to the PPAR γ receptor. PRB-5 scored somewhat poorer while PRB-8 had the highest ABE score. These findings are highly unsuspected and could be translated into a more effective transcription of the respective PPAR γ activated gene responsible for handling of blood glucose.

Fig. 6 depicts the binding affinities to the nuclear receptor PPAR α which is mainly responsible for metabolisation of fat, blood lipids, fat tissue biology and weight control. Several DHA-derivatives had high binding affinity but PRB8 had the highest score. This is also highly unsuspected.

Fig. 7 depicts the binding affinities to the nuclear receptor RXR α . The physiologic consequence of binding to the RXR α receptor has not been firmly established. It is known that RXR binds to the PPAR receptors thereby forming a heterodimer which then, subsequently, initiates transcription of the defined gene.

Table 1

	PPAR α			PPAR γ			RXR α		
	LBE	ABE	f_{bound}	LBE	ABE	f_{bound}	LBE	ABE	f_{bound}
DHA	-16.14	-13.29 (0.47)	0.60	-11.34	-10.51 (0.21)	0.35	-12.15	-10.72 (0.29)	0.40
cr-PRB1	-16.29	-14.25 (0.53)	0.50	-12.96	-11.82 (0.38)	0.30	-15.68	-14.25 (0.35)	0.30
cs-PRB1	-15.97	-14.01 (0.30)	0.80	-12.74	-10.24 (0.48)	0.65	-17.17	-14.48 (0.44)	0.50
cr-PRB2	-16.00	-14.02 (0.54)	0.50	-13.72	-12.17 (0.54)	0.25	-14.81	-12.80 (1.56)	0.20
cs-PRB2	-16.86	-14.48 (0.27)	0.85	-13.34	-12.05 (0.30)	0.60	-15.57	-12.39 (2.20)	0.20
PRB5	-16.54	-14.37 (0.40)	0.65	-13.16	-11.88 (0.30)	0.50	-18.21	-15.028 (1.03)	0.30
cr-PRB7	-17.06	-15.09 (0.32)	0.80	-12.52	-8.34 (2.30)	0.50	-16.35	-13.72 (0.96)	0.30
cs-PRB7	-16.31	-14.72 (0.37)	0.65	-14.00	-10.84 (0.40)	0.55	-14.63	-12.52 (0.84)	0.30
PRB8	-18.45	-16.41 (0.57)	0.45	-13.39	-13.04 (0.35)	0.10	-17.31	-15.79 (0.57)	0.30
sROSI				-9.47	-9.01 (0.17)	0.20			
sROSI				-10.05	-7.89 (0.91)	0.20			
rPIO				ND					
sPIO				-7.59	-7.59	0.05			
cr-PRB9	-17.15	-15.12	0.35	-15.14	-12.84	0.25	-17.56	-14.30	0.15
cs-PRB9	-15.66	-14.06	0.45	-13.50	-13.50	0.05	-16.63	-15.20	0.15
cr-PRB10	-14.88	-13.46	0.25	-7.31	-7.31	0.05	-15.59	-15.58	0.10
cs-PRB10	-15.17	-12.90	0.50	-11.78	-9.64	0.20	-14.75	-13.05	0.30
crPRB11	-16.50	-12.76	0.30	ND	ND	ND	-15.26	-13.56	0.10
cs-PRB11	-17.77	-13.66	0.20	ND	ND	ND	-14.48	-13.48	0.20
cr-PRB12	-13.23	-11.61	0.65	-11.11	-8.61	0.30	-14.73	-13.13	0.35
cs-PRB12	-15.10	-10.60	0.55	-9.64	-8.20	0.45	-14.08	-8.67	0.60
cr-PRB13	ND	ND	-	ND	ND	ND	-13.20	-10.95	0.25
cs-PRB13	-6.89	-6.89	0.05	ND	ND	ND	-11.84	-13.10	0.50
cr-PRB14	-14.96	-11.99	0.60	-9.50	-8.72	0.20	-15.84	-14.77	0.20
cs-PRB14	-15.42	-11.89	0.40	-11.29	-10.87	0.20	-15.22	-12.77	0.30
cr-PRB15	-14.62	-11.88	0.45	-12.49	-11.05	0.45	-17.01	-13.73	0.45
cs-PRB15	-15.90	-12.90	0.40	-12.15	-10.55	0.50	-16.93	-13.27	0.60

5

ND= Not docked, c = the double bonds in all-cis form. r= R enantioisomer, s= S enantioisomer. ROSI= Rosiglitazone, PIO= Pioglitazone

10

Several of the PRBs have a high LBE and ABE score for the PPAR α and PPAR γ receptors even compared to the mother compound DHA but also to the PPAR γ ligands rosiglitazone and pioglitazone, both in the r and s form. This is an interesting observation indicating that several of the PRBs could be promising competitors to the established anti-diabetics rosiglitazone and pioglitazone.

15

Ethyl derivatives in alfa position of the same fatty acids, both the r and the s form, did not improve affinity. This was especially true for the PPAR γ receptor. As mentioned previously the PPAR α receptor is more promiscuous binding a long series of fatty acids.

In conclusion, many of the DHA-derivatives tested demonstrated interesting affinities to the PPAR α and PPAR γ receptors with binding affinities better than rosiglitazone and pioglitazone.

5 **Example 3**

Affinity testing in transfected cells (block 3 in Fig. 2)

Background

Release of luciferase is correlated to transcription of genes. Binding of a ligand to a nuclear receptor such as PPAR γ induces transcription of the respective
10 gene thereby releasing luciferase. This technique therefore provides a measure of ligand affinity to the receptor as well as activation of the responsible gene.

Method

Transient transfection of COS-1 cells was performed in 6-well plates as
15 described by Graham and van der Eb (Graham). For full length PPAR transfection studies, each well received 5 μ g reporter construct, 2.5 μ g pSV- β -galactosidase as an internal control, 0.4 μ g pSG5-PPAR γ 2. The cells were harvested after 72 h, and the luciferase activity was measured according to the protocol (Promega). The luciferase activity was normalised against β -galactosidase activity. The adipocytes were
20 transfected at D11 of differentiation using 16 μ l LipofectaminPlus reagent, 4 μ l Lipofectamine (Life Technologies Inc.), 0.2 μ g pSG5-PPAR γ , and 100 ng pTK Renilla luciferase as control of transfection efficiency. Three hours after transfection, cells were cultured in serum containing medium and incubated for 48 hours in the same medium containing appropriate agents. The luciferase activities were measured
25 as recommended by the manufacturer (Dual Luciferase assay, Promega). All transfections were performed in triplicate.

Fatty acids (BRL or DHA) and PRBs (stock solutions) were solubilized to 0.1 M final concentration in DMSO. Then, Fatty solubilized to 10mM in DMSO and stored in 1.5 ml tubes (homopolymer, plastic tubes) flushed with argon and stored at -
30 20°C. 10 μ M of PRBs or fatty acids and DMSO (control) was added to the media 5h after transfection. Transfected cells were maintained for 24h before lysis by reporter lysis buffer. Binding of PRBs or fatty acids to the LBD of PPAR activates GAL4 binding to UAS, which in turn stimulates the tk promoter to drive luciferase expression. Luciferase activity was measured using a luminometer (TD-20/20
35 luminometer; Turner Designs, Sunnyvale, CA) and normalized against protein content.

Results

Fig. 8 depicts the release of luciferase from transfected cells treated with different PRBs. The results indicate that PRB-1,2,6,7 and 14 have a significantly higher release of luciferase compared to PRB-3,5,9,10,11,12, and 16.

5

Example 4

Affinity testing in adipose prone animals with metabolic syndrome (block 4 in Fig. 2)

10 Background

An animal model of the metabolic syndrome using the adipose prone mice of the C57BL/6J strain was used to test affinity of PRB-2,5, and 8 compared to 97% DHA and the antidiabetic compound rosiglitazone to PPAR γ , by measuring the release of luciferase from adipose cells taken from these animals. The animals (n=8
15 in each group) were fed high fat diet (fat constituting 60% of total calories, the same diet as used in Block 5) for 8 weeks. Thereafter they were given the PRBs in a dose of 1.5% of the fat content of the diet for another two weeks. The rosiglitazone group was given an amount of 100 mg/KG diet. The control groups continued with either only high fat diet or standard chow. Fig. 9 shows the study design.

20

Method

After sacrifice adipose tissue (epididymal and subcutaneous) was cleared from other structures and cut into millimeter-size pieces. Fat tissue was rinsed in 0,9% NaCl and digested in 5 mL of Krebs-Ringer solution containing Hepes, fatty-acid free
25 bovine serum albumin, 200nM of adenosin, 2 nM of glucose, and 260 U/mL of collagenase for 1,5 h at 37 degrees C in a shaking water bath. After collagenase digestion, adipocytes were separated from tissue debris by filtering. Cells were then washed in Krebs-Ringer solution containing Hepes, fatty-acid free bovine serum albumin, 200nM of adenosin, 2 nM of glucose and kept in a shaking water bath at 37
30 degrees for a maximum of 30 min until electroporation.

Isolated primary adipocytes were transfected by electroporation to measure the specific PPAR gamma response element (PPRE) activity. In this case we incorporated a plasmid encoding firefly luciferase cDNA under control of a PPRE from the acyl-CoA-oxidase gene. The cells were also co-transfected with a plasmid
35 containing cDNA for Renilla luciferase controlled by a constitutively active promoter. The PPRE inducible firefly luciferase activity was normalised according to Renilla luciferase, thus correcting for potential differences in the amount of

transfected cells. To measure luciferase signal we used the Dual-Luciferase® Reporter assay System (Promega, USA).

Pooled epididymal fat tissue was enough to isolate adipocytes for running duplicates. Each of groups was sacrificed in two separated days, and 4 independent
5 transfections for each dietetic group were obtained.

Results

During first 8 weeks of feeding with HF diet (33,7% of fat, w/w), there was a gradual increase of body weight in comparison to control mice fed with chow diet
10 (4,5% w/w). During last 2 weeks of feeding with experimental diets high fat diet animals and animals given high fat diet in combination with Rosiglitazon continued gaining weight, approximately with the same rate as before. In case of PRB-8 and PRB-5 enriched HF diet the weight gain was reduced. However, in case of PRB-2 and DHA (5% w/w) the diet completely stopped the weight gain and even led to reduction
15 of body weight (Fig. 10). Food consumption was recorded occasionally (4x). There were no differences between the HF and the intervention groups.

In case of high fat in combination with Rosiglitazon, the endogenous activity of PPAR γ was approximately 2-fold higher than in all the others diet groups (Fig. 11). Furthermore, these fat cells became more sensitive to additional in vitro stimulation
20 with 5uM Rosiglitazon (5, 12- fold stimulation) in comparison to i.e. HF diet itself (1,5-fold stimulation). This rosiglitazon – sensitizing effect was also recorded in the PRB-2 and the PRB-5 diet group (2,6-fold stimulation).

Data from this study clearly demonstrates acitivity on the nuclear PPAR receptors, in particular with the effects on weight which was most prominent for the
25 groups given PRB-2. Even animals given PRB-5 and PRB-8 did not increase in weight as did the high fat diet group. Interestingly animals given rosiglitazone increaed in weight to the same extent as animals given only the high fat diet. This clearly demonstrates the negative effects of giving only a PPAR γ ligand, like the glitazones, with the risk of weight increase even if insulin resistance is reduced.
30 However, when it comes to PPAR γ activation measured as luciferase activity in this experiment, rosiglitazone scores higher compared to any of the PRBs. Within the PRB groups PRB-2 and PRB-5 had a higher score compared to PRB-8 and DHA only (Fig. 12).

35 Example 5

Pharmacodynamic effects of DHA derivatives in an animal model of metabolic syndrome (block 5 in Fig. 2)

Background

An animal model of the metabolic syndrome using the adipose prone mice of the C57BL/6J strain was used to document effects on typical laboratory and pathological anatomical features common for the metabolic syndrome. When given a high fat diet containing about 60% of fat, the animals are getting obese developing high insulin plasma levels, pathological glucose tolerance test, elevated serum triglycerides and non-esterified fatty acids, and fat liver.

Example 5a

Effect of DHA derivatives in adipose prone mice during 4 months of dietary interventions

Method

All experiments were performed on male C57BL/6 mice, either a substrain C57BL/N (supplier: Charles River, Germany, n = 160, experiments A-C, see below), or a substrain C57BL/6J (supplier: the Jackson laboratory, Bar Harbor, ME, USA, n = 32, experiment D). Total numbers of animals used were higher (n = 170 and 36, respectively), because of culling. In the latter case, animals were bred for several generations (< 20) at the Institute of Physiology. At the beginning of the treatment, animals were 14-week-old and their body weight range was 23.6 – 27.1g. One week before the study start, animals were sorted according to their body weight and assigned to subgroups (n = 8) of similar mean body weight. This method allowed for culling of about 5 – 10 % of animals showing the lowest and highest body weight, respectively. The animals eliminated from the study at this stage were sacrificed by cervical dislocation. Complete health check of mice was performed by the supplier Charles River and at the start of study serological tests were performed by ANLAB (Prague, Czech Republic). In addition, regular health checks were performed in the animal house in 3-mo-intervals using sentinel mice and serological examinations (ANLAB). In all the tests, the animals were free of specific pathogens.

Diets

Animals were fed 3 types of experimental diets:

- (i) Chow diet (ssniff R/M-H from SSNIFF Spezialdiäten GmbH, Soest, Germany; see also <http://ssniff.de>) with protein, fat and carbohydrate forming 33, 9, and 58 energy %, respectively

- (ii) High-fat diet prepared in the laboratory (cHF diet) with protein, fat and carbohydrate forming 15, 59, and 26 energy %, respectively, and well characterized fatty acid composition (with most of the lipids coming from corn oil; see Ruzickova 2004)
- 5 (iii) cHF diets in which 0.15, 0.5, and 1.5 % of fat (specifically the corn oil constituent) was replaced by various PRB-compounds, namely PRB1, PRB2, PRB5, PRB7, and PRB8, or by DHA. All these compounds were in the form of ethyl esters, provided by Pronova Biocare a.s. in sealed containers. Chemical composition of the PRB-compounds was unknown to
- 10 the laboratory performing the experiments (Institute of Physiology, Academy of Sciences Prague, Czech Republic).

After arrival, the PRB-compounds were stored in a refrigerator in original containers. The containers were opened just before preparation of the experimental

15 diets. Diets were kept in plastic bags flushed by nitrogen and stored at -70°C in small aliquots sufficient for feeding animals for one week. Fresh ratios were given in 2-day intervals or daily.

Outline of the study

20 The study was based on 4 individual experiments. In each of the experiments, different PRB-compounds (or DHA, respectively) admixed to cHF diet in three different concentrations (0.15, 0.5, and 1.5 % of the fat content) were tested. In each experiment, a subgroup of plain cHF diet-fed mice was included and served as a control. Mice were caged in groups of 4 and fed standard chow diet until 3 mo of age,

25 when animals (n = 8-13) were randomly assigned to the different test diets. After 2 mo on this new diet (at 5 mo of age), animals were fasted overnight and in the morning, intraperitoneal Glucose Tolerance Test (GTT) was performed. Animals were sacrificed after 4 months on the experimental diets, at 7 mo of age, and the end-point analysis were performed.

30

Study parameters.

The parameters in the study were: Body weight gain (grams), area under the curve (AUC) from intraperitoneal glucose tolerance tests (mMol x 180 min), plasma insulin (ng/ml), serum triglycerides (TAGs, mmol/l), and non-esterified fatty acids

35 (NEFA, mmol/l).

Fig. 13 shows a typical blood glucose elimination curve before and after animals with insulin resistance are given a compound with insulin resistance reducing

effect. Reduction of the area under the curve means that blood glucose is eliminated more effectively due to reduced insulin resistance.

Results

5 The results are shown in the following tables 2, 3 and 4. (* = significant differences compared to cHF diets ($P < 0.05$).)

10 Table 2 shows the effects in animals given 1.5% concentration of the PRB test compounds compared to animals given standard chow (STD), composite high fat diet (cHF) or 97% DHA. Body weight gain was significantly reduced in animals given PRB-2 compared to animals given high fat diet (cHF). Food intake was somewhat lower in this group. The most pronounced reduction in AUC from glucose tolerance tests was seen in the same group and even in animals given PRB-1. Plasma insulin was significantly lower in the PRB-2 group compared to the cHF controls even if the PRB-1 and PRB-5 treated animals showed some effect on this parameter too. The PRB-2 group showed the biggest reduction in triglycerides (TAGs) and non-esterified fatty acids (NEFA).

15 Table 3 shows the effects in animals given a lower concentration, 0.5%, of the PRB test compounds compared to animals given standard chow (STD), composite high fat diet (cHF) or 97% DHA. Body weight gain was somewhat lower in animals given PRB-2 and PRB-5. AUC from the glucose tolerance test as well as plasma insulin, however, was significantly lower only in the PRB-2 group.

20 Table 4 shows the results from the lowest PRB concentration given, 0.15%. Here, the differences were small. Weight gain was somewhat lower in the PRB-1 and PRB-2 groups while AUC was significantly lower only in the PRB-2 group. Plasma insulin was lower in PRB-1, 2 and 7.

25

Table 2 The effect of PRB derivatives after 4 months of treatment with 1.5% concentration

Parameter	STD	cHF	PRB-1	PRB-2	PRB-5	PRB-7	DHA
Body weight (grams)	32.4 ± 0.7	49.6 ± 0.6	44.0 ± 1.5*	30.1 ± 1.1*	46.3 ± 1.6	45.9 ± 1.1*	47.1 ± 0.7*
Body wt. gain (grams)	7.8 ± 0.4	25.2 ± 0.5	20.2 ± 1.3*	6.4 ± 0.8*	22.4 ± 1.4	21.7 ± 0.9*	23.0 ± 0.8*
Food intake (grams/mouse/day)	3.64 ± 0.04	2.70 ± 0.02	2.64 ± 0.03	2.38 ± 0.05*	2.62 ± 0.02	2.68 ± 0.03	2.63 ± 0.02
AUCglucose (mM x 180min)	1124 ± 57	1625 ± 151	913 ± 68*	982 ± 80*	1264 ± 192	1122 ± 73	2132 ± 288*
Fasted glucose (mg/dL)	77 ± 3	145 ± 7	130 ± 14	95 ± 6*	136 ± 12	120 ± 9	138 ± 7
Insulin (ng/mL)	1.03 ± 0.09	5.35 ± 0.36	2.73 ± 0.33	0.60 ± 0.18*	2.47 ± 0.19*	4.42 ± 0.87	6.55 ± 0.31
TAGs (mmol/L)	1.41 ± 0.09	1.45 ± 0.07	1.58 ± 0.08	0.71 ± 0.01*	1.19 ± 0.07	1.15 ± 0.08	1.91 ± 0.26*
NEFA (mmol/L)	0.57 ± 0.05	0.61 ± 0.04	0.63 ± 0.03*	0.54 ± 0.03*	0.72 ± 0.05	0.82 ± 0.06	0.98 ± 0.07

Table 3 The effect of PRB derivatives after 4 months of dietary interventions: 0.5% concentration.

Parameter	STD	CHF	PRB-1	PRB-2	PRB-5	PRB-7	DHA
Body weight (grams)	32.4 ± 0.7	49.6 ± 0.6	47.4 ± 0.6	45.8 ± 1.7	45.7 ± 1.5	48.8 ± 0.9	46.9 ± 0.7*
Body wt. gain (grams)	7.8 ± 0.4	25.2 ± 0.5	23.8 ± 0.5	21.9 ± 0.6	22.0 ± 1.4	24.8 ± 0.8	22.9 ± 0.7*
Food intake (grams/mouse/day)	3.64 ± 0.04	2.70 ± 0.02	2.67 ± 0.04	2.69 ± 0.04	2.63 ± 0.02	2.69 ± 0.03	2.70 ± 0.03
AUCglucose (mM x 180min)	1124 ± 57	1625 ± 151	1596 ± 205	1224 ± 72*	1581 ± 231	1674 ± 203	1816 ± 182
Fasted glucose (mg/dL)	77 ± 3	145 ± 7	131 ± 7	136 ± 7	130 ± 7	152 ± 6	136 ± 8
Insulin (ng/mL)	1.03 ± 0.08	5.35 ± 0.36	3.93 ± 0.59	2.75 ± 0.21*	5.12 ± 0.93	4.10 ± 0.57*	5.82 ± 0.47
TAGs (mmol/L)	1.41 ± 0.09	1.45 ± 0.07	2.03 ± 0.22	1.29 ± 0.08	1.46 ± 0.17	1.42 ± 0.08	1.78 ± 0.08*
NEFA (mmol/L)	0.57 ± 0.05	0.61 ± 0.04	0.73 ± 0.04*	0.75 ± 0.04	0.77 ± 0.03*	0.87 ± 0.04	0.89 ± 0.03

Table 4. The effect of PRB derivatives after 4 months of dietary interventions: 0.15% concentration.

Parameter	STD	CHF	PRB-1	PRB-2	PRB-5	PRB-7	DHA
Body weight (grams)	32.4 ± 0.7	49.6 ± 0.6	47.2 ± 1.3	46.7 ± 1.1	48.0 ± 0.8	47.4 ± 0.8*	48.3 ± 0.6
Body wt. Gain (grams)	7.8 ± 0.4	25.2 ± 0.5	22.9 ± 1.1	22.8 ± 0.9	24.2 ± 0.5	23.2 ± 0.7*	24.3 ± 0.8
Food intake (grams/mouse/day)	3.64 ± 0.04	2.70 ± 0.02	2.63 ± 0.04	2.57 ± 0.03*	2.66 ± 0.02	2.59 ± 0.02	2.79 ± 0.03
AUCglucose (mM x 180min)	1124 ± 57	1625 ± 151	1291 ± 172	1071 ± 148*	1443 ± 70	1425 ± 97	1477 ± 214
Fasted glucose (mg/dL)	77 ± 3	145 ± 7	126 ± 15	132 ± 5	151 ± 5	141 ± 9	141 ± 10
Insulin (ng/mL)	1.03 ± 0.08	5.35 ± 0.36	3.50 ± 0.29	4.00 ± 0.64	6.21 ± 0.45	3.76 ± 0.72*	4.31 ± 0.39*
TAGs (mmol/L)	1.41 ± 0.09	1.45 ± 0.07	1.75 ± 0.08	1.42 ± 0.07	1.64 ± 0.28	1.41 ± 0.11	1.50 ± 0.13
NEFA (mmol/L)	0.57 ± 0.05	0.61 ± 0.04	0.62 ± 0.04*	0.78 ± 0.04*	0.71 ± 0.09	0.85 ± 0.06	0.96 ± 0.07

In conclusion, testing of PRB-1,2,5, and 7 during 4 months in adipose prone animals with insulin resistance and metabolic syndrome demonstrated a clear and unsuspected effect of the PRBs tested, in particular the DHA-derivative PRB-2, on insulin resistance and symptoms of the metabolic syndrome such as weight reduction, reduced AUC in the intraperitoneal glucose tolerance test, lower insulin/plasma levels as well as reduced triglyceride and non-esterified free fatty acids. Effects were observed in the dose of 1.5% as well as in the 0.5% group. Some effects were even noticed in the lowest concentration group of 0.15%.

Testing of the PRB-8 compound was started later, therefore only data from 2 months intervention in three dose groups (1.5%, 0.5% and 0.15%) are given. In the group given 1.5%, body weight (BW) was 28.0 ± 0.7 grams compared to controls 29.6 ± 0.9 , AUC 1031 ± 104 compared to 1074 ± 91 . These differences are small but the trend is interesting. There were no differences between intervention and controls for the lower doses of 0.5% and 0.15%. The data regarding PRB-8 data from 2 months medication showing a trend towards weight reduction and AUC.

Example 5b

Effect of DHA derivatives on established metabolic syndrome and insulin resistance

Method

In another experiment, PRB-2, PRB-5, and PRB-7 were tested in the same breed of animals. In this experiment, animals were initially fed high fat diet (the same as in the previous experiment 5a) for 8 weeks developing insulin resistance and the metabolic syndrome, and then given the PRBs. The start dose was to substitute 15% of the fat content with the PRBs but the animals did not tolerate this dose. After a period of another two weeks the animals were given 1.5% of PRB-2, 5% and 1.5% of PRB-5, and 1.5% and 0.5% of PRB-7.

Results

Weight reduction was very good in the animals given PRB-2. Even the animals given PRB-5 showed some weight reduction but in the higher dose of 5%. Triglycerides were reduced with all derivatives tested compared to the control animals fed composite High Fat diet. Reduction of non-esterified fatty acids was most pronounced with PRB-2 and PRB-5, however in different doses. (See Fig. 14)

Blood cholesterol was reduced in animals given PRB-2 and PRB-5. Blood glucose was not affected due to the fact that these animals are in a pre-diabetic state with normal glucose due to a high insulin production. However, more importantly, plasma insulin was significantly reduced in the PRB-2 group in a much lower concentration compared to the second best DHA-derivative PRB-5. Even PRB-7 showed some effects on the insulin concentration. (See Fig. 15)

PRB-2 showed a statistically significant reduction of the AUC blood glucose at all time points of the curve compared to the baseline values. This means that blood glucose was more effectively removed after treatment of 1,5% of PRB-2. PBR-5 and PBR-7 showed some effect but not to the same extent. (See Fig. 16)

These effects are highly unsuspected and very relevant for a positive effect in metabolic syndrome and type 2 diabetes. These patients are almost exclusively overweight or obese and a weight reductive effect of a drug is highly positive. The mostly used remedies used for treatment of type 2 diabetes today, the thiazolidinedions, which are potent PPAR γ ligands thereby reducing insulin resistance, often result in weight increase which is highly negative for these subset of patients (Yki-Järvinen 2004).

Reduction of serum triglycerides is another very important effect that was demonstrated in the experiments. Patients with metabolic syndrome and type 2 diabetes usually have elevated triglycerides. The triglyceride lowering effects of the DHA-derivatives is a positive finding and again PRB-2 demonstrated the most potent effect with the lowest dose given. The very positive effects on plasma insulin and glucose tolerance test are very promising and highly unsuspected. Taken together the effects obtained with the DHA-derivatives in particular PRB-2 are very promising forming a good basis for development of an antidiabetic drug.

Example 5c

Testing of DHA derivatives on liver fat

Method

Tissue samples from animals in the experiments with DHA derivatives was histologically analysed. After paraffination, tissue samples from liver, adipose tissue, skeletal muscle, pancreas, and kidney were stained with eosin-hematoxylin.

Results

There were no pathological findings in the tissues examined with exception from liver. Control animals fed high fat diet had developed fat liver (liver steatosis).

Fat droplets in the liver can easily be distinguished from normal liver cells. Animals treated with PRB-1, 5, and 7 had low degree of fat liver. However, animals treated with 1.5% of PRB-2 had completely normal liver cells with no trace of steatosis.

5 This is an extremely important finding and very relevant for treatment of patients with insulin resistance, obesity and type 2 diabetes. Liver steatosis is a common finding in these patients which is usually related to an overload of fatty acids and triglycerides, biological markers present in the development of insulin resistance and the metabolic syndrome. DHA-derivatives reduce liver steatosis, and PRB-2 was the most efficient compound showing this effect.

10

Discussion and conclusions

The present application clearly identifies a new group of compounds which are activating nuclear receptors, especially PPAR γ and PPAR α , thereby offering a series of therapeutic effects in the treatment of insulin resistance, the metabolic syndrome, type 2 diabetes, cardiovascular disease and other atherosclerotic related diseases.

15 Members of this group are DHA derivatives with side chains of different kind in the alfa position of the molecule. A large number of alfa-substituted DHA derivatives have been tested and compared with controls as well as pure DHA and EPA. Several of the compounds tested have demonstrated interesting biological effects very relevant for a potential anti-diabetic drug.

20 Interestingly, and not conceivable on beforehand, alfa-ethyl DHA ethyl ester (PRB-2) was significantly more effective in the battery of tests used to demonstrate effects related to insulin resistance and thereby diseases mainly caused by this pathophysiologic condition such as the metabolic syndrome, type 2 diabetes, cardiovascular disease and other atherosclerotic related diseases. Alfa-ethyl DHA ethyl ester was enriched in liver tissue from animals given the different DHA derivatives tested (Block 1) indicating that this compound was not utilised for synthesis of triglycerides, eicosanoids or other metabolic intermediates. Indirectly this would mean that alfa-ethyl DHA would be available for liganding to nuclear receptors like the PPARs.

25 In testing of affinity to PPAR γ and PPAR α using computerized docking technology a large number of the DHA-derivatives showed affinities to both receptors, not least PPAR γ which probably is the most important nuclear receptor engaged in the activation of genes responsible for metabolism of blood glucose. In particular alfa-ethyl DHA (PRB-2) as well as alfa-diethyl DHA (PBR-8) possessed excellent affinity to these nuclear receptors. Compared to alfa-diethyl DHA alfa-ethyl

30

35

DHA has two stereoisomers, the r and the s form. Using the docking technology both stereoisomers possessed about the same affinity to PPAR γ and PPAR α meaning that neither the r or the s form should have advantages compared to the racemic form. In fact the racemic form may have advantages over each one of the stereoisomers.

5 When affinity was tested in transfected cells carrying the nuclear receptor and the subsequent DNA response element, several of the PRBs demonstrated good affinity measured as release of luciferase. Alfa-ethyl DHA (PRB-2) together with PRB-6,7 and 14 demonstrated the best effects.

10 Five of the DHA derivatives have been extensively tested in the C57BL/6 mouse model developing insulin resistance and the metabolic syndrome when fed high fat diet. Alfa-ethyl DHA (PRB-2) has been tested in three individual experiments while PRB-1,5, and 7 were tested in two and alfa-diethyl DHA (PRB-8) was tested in one experiment. All derivatives demonstrated significant biological effects. However, alfa-ethyl DHA (PRB-2) showed the most promising effects with a
15 consistent reduction in body weight, AUC from intraperitoneal glucose tolerance testing, plasma insulin as well as serum triglycerides and non-esterified fatty acids. The effects were obtained on the doses 1.5% and 0.5%. The lowest tested dose 0.15% did not perform convincingly. Alfa-ethyl DHA (PRB-2) in a dose of 1.5% has also demonstrated a normalisation of fat liver, an important pathological finding in
20 patients and animals with insulin resistance and metabolic syndrome.

 Comparing with pure DHA, alfa-ethyl DHA (PRB-2) seems to be 10-30 times as potent as DHA. All in all these findings and the potency compared to the mother molecule DHA are not predictable and highly unexpected.

25 Since alfa-ethyl DHA (PRB-2) seems to work by simultaneous liganding to the nuclear receptors PPAR α and PPAR γ the compound would not only possess therapeutic interesting effects on glucose and lipid metabolism, not least in patients with insulin resistance, metabolic syndrome and type 2 diabetes but also have weight reduction as well as a general anti-inflammatory effect. Directly or through positive intervention on risk factors alfa-ethyl DHA (PRB-2) would have a preventive effect
30 on the development of cardiovascular disease such as myocardial infarction and cerebral stroke as well as having a preventive effect on cardio-vascular mortality.

 Pharmaceuticals acting as PPAR γ ligands are already on the market but even if these compounds are having positive effects on glucose metabolism, they are hampered by adverse effects such as elevated triglycerides, weight increase and
35 oedema. The alfa-substituted DHA derivatives presented in this application are having a combined PPAR γ and PPAR α effect which is probably both relevant and advantageous for patients with insulin resistance, metabolic syndrome and type 2

diabetes. Furthermore, these combinative actions should have important effects also on blood lipids, inflammatory events, atherosclerosis, and thereby cardiovascular disease.

The invention shall not be limited to the shown embodiments and examples.

5


References

- Simonopoulos AP. Essential fatty acids in health and chronic disease. Am J Clin Nutr 1999;70 (Suppl):560S-569S
- 10 Geleijnse JM, Giltay EJ, Grobbee DE, et al. Blood pressure response to fish oil supplementation: metaregression analysis of randomized trials. J Hypertension 2002;20:1493-1499
- 15 Storlien LH, Hulbert AJ, and Else PL. Polyunsaturated fatty acids, membrane function and metabolic diseases such as diabetes and obesity. Curr Opin Clin Nutr Metab Care 1998;1:559-563
- Jump DB. The biochemistry of n-3 polyunsaturated fatty acids. J Biol Chem 20 2002;277:8755-8758
- Pawar A and Jump D. Unsaturated fatty acid regulation of peroxisomes proliferator-activated receptor alfa activity in rat primary hepatocytes. J Biol Chem 2003;278:35931-35939
- 25 Meigs JB, Wilson PWF, Nathan DM, et al. Prevalence and characteristics of the metabolic syndrome in the San Antonio Heart and Framingham offspring studies. Diabetes 2003;52:2160-2167
- 30 Storlien LH, Kraegen WE, Chisholm DJ, et al. Fish oil prevents insulin resistance induced by high fat feeding in rats. Science 1987;237:885-888
- Field CJ, Ryan EA, Thomson ABR, et al. Diet fat composition alters membrane phospholipid composition, insulin binding and glucose metabolism in adipocytes from control and diabetic animals. J Biol Chemistry 1990;265:11143-11150
- 35 Yki-Järvinen, H. Thiazolidinediones. NEJM 2004;351:1106-1118

- Adams M, Montague CT, Prins JB, et al. Activators of peroxisomes proliferator-activated receptor gamma have depot-specific effects on human preadipocyte differentiation. *J Clin Invest* 1997;100:3149-3153
- 5 Ruzickovaj, Rossmeisl M, Prazak T, et al. Omega-3 PUFA of marine origin limit diet-induced obesity in mice by reducing cellularity of adipose tissue, *Lipids* 2004; 39: 1177-1185
- 10 Vaagenes H, Madsen L, Dyrøy E, et al. The hypolipidaemic effect of EPA is potentiated by 2- and 3-methylation. *Biochim Pharmacol* 1999;58:1133-1143
- Larsen L, Granslund L, Holmeide AK, et al. Sulfur-substituted and α -methylated fatty acids as peroxisome proliferator-activated receptor activators. *Lipids* 2005;40:49-57
- 15 Larsen L, Hørvik K, Sørensen HIN, et al. Polyunsaturated thia- and oxa-fatty acids: incorporation into cell-lipids and their effects on arachidonic acid- and eicosanoids synthesis. *Biochim et Biophys Acta* 1997;1348:346-354
- 20 Larsen, et.al *Biochemical Pharmacology* 1998; 55, 405
- Chih-Hao L, Olson P, and Evans RM. Lipid metabolism, metabolic diseases, and peroxisome proliferator-activated receptors. *Endocrinology* 2003;144:2201-2207
- 25 Willumsen N, Waagenes H, Holmsen H, et al. On the effect of 2-deuterium- and 2-methyl-eicosapentaenoic acid derivatives on triglycerides, peroxisomal beta-oxidation and platelet aggregation in rats. *Biochim Biophys Acta* 1998;1369:193-203
- Mitsunobu O, *Synthesis* 1981;1
- 30 Ager DJ, Prakash I, and Schaad DR. 1,2-amino alcohols and their heterocyclic derivatives as chiral auxiliaries in asymmetric synthesis *Chem Rev* 1996;96:835-876

Claims

1. A compound of formula (I):



5 wherein

- R₁ and R₂ are the same or different and may be selected from the group consisting of a hydrogen atom, a hydroxy group, an alkyl group, a halogen atom, an alkoxy group, an acyloxy group, an acyl group, an alkenyl group, an alkynyl group, an aryl group, an alkylthio group, an alkoxycarbonyl group, an alkylsulfinyl group, 10 an alkylsulfonyl group, an amino group, and an alkylamino group; and

- X represents a carboxylic acid group, a carboxylate group, or a carboxamide group,

15 or any pharmaceutically acceptable salt, solvate, complex or pro-drug thereof;

with the provisos that:

▪ the compound of formula (I) is not (all-Z)-4,7,10,13,16,19-docosahexaenoic acid (DHA), alpha-methyl DHA, alpha-methyl DHA methyl ester, alpha-methyl DHA ethyl ester or alpha-hydroxy DHA ethyl ester. 20

2. A compound according to claim 1, wherein said alkyl group is selected from the group consisting of methyl, ethyl, n-propyl, isopropyl, n-butyl, iso-buthyl, 25 sec.-butyl, n-hexyl, and benzyl.

3. A compound according to claim 1, wherein said halogen atom is selected from the group consisting of fluorine, chlorine, bromine, and iodine.

30 4. A compound according to claim 1, wherein said alkoxy group is selected from the group consisting of methoxy, ethoxy, propoxy, isopropoxy, sec.-butoxy, phenoxy, benzyloxy, OCH₂CF₃, and OCH₂CH₂OCH₃.

5. A compound according to claim 1, wherein said alkenyl group is selected from the group consisting of allyl, 2-butenyl, and 3-hexenyl.

6. A compound according to claim 1, wherein said alkynyl group is selected from the group consisting of propargyl, 2-butyne, and 3-hexyne.

7. A compound according to claim 1, wherein said aryl group is a phenyl group.

8. A compound according to claim 1, wherein said alkylthio group is selected from the group consisting of methylthio, ethylthio, isopropylthio, and phenylthio.

9. A compound according to claim 1, wherein said alkoxy carbonyl group is selected from the group consisting of methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, and butoxycarbonyl.

10. A compound according to claim 1, wherein said alkylsulfinyl group is selected from the group consisting of methanesulfinyl, ethanesulfinyl, and isopropanesulfinyl.

11. A compound according to claim 1, wherein said alkylsulfonyl group is selected from the group consisting of methanesulfonyl, ethanesulfonyl, and isopropanesulfonyl.

12. A compound according to claim 1, wherein said alkylamino group is selected from the group consisting of methylamino, dimethylamino, ethylamino, and diethylamino.

13. A compound according to claim 1, wherein said carboxylate group is selected from the group consisting of ethyl carboxylate, methyl carboxylate, n-propyl carboxylate, isopropyl carboxylate, n-butyl carboxylate, sec.-butyl carboxylate, and n-hexyl carboxylate.

14. A compound according to claim 1, wherein said carboxamide group is selected from the group consisting of primary carboxamide, N-methyl carboxamide, N,N-dimethyl carboxamide, N-ethyl carboxamide, and N,N-diethyl carboxamide.

15. A compound according to claim 1, wherein
R₁ and R₂ are selected from the group consisting of a hydrogen atom, a hydroxy
group, an alkyl group, a halogen atom, an alkoxy group, an alkylthio group, an
alkylsulfinyl group, an alkylsulfonyl group, an amino group, and an alkylamino
5 group.

16. A compound according to claim 1, wherein
R₁ and R₂ are selected from the group consisting of a hydrogen atom, a hydroxy
group, a C₁-C₇ alkyl group, a halogen atom, a C₁-C₇ alkoxy group, a C₁-C₇ alkylthio
10 group, a C₁-C₇ alkylsulfinyl group, a C₁-C₇ alkylsulfonyl group, an amino group, and
a C₁-C₇ alkylamino group.

17. A compound according to claim 16, wherein
said C₁-C₇ alkyl group is methyl, ethyl, or benzyl;
15 said halogen atom is fluorine or iodine;
said C₁-C₇ alkoxy group is methoxy or ethoxy;
said C₁-C₇ alkylthio group is methylthio, ethylthio or phenylthio;
said C₁-C₇ alkylsulfinyl group is ethanesulfinyl;
said C₁-C₇ alkylsulfonyl group is ethanesulfonyl;
20 said C₁-C₇ alkylamino group is ethylamino or diethylamino; and
X represents a an ethylcarboxylate or a carboxamide group.

18. A compound according to claim 1, wherein
R₁ and R₂ are selected from the group consisting of a hydrogen atom, a C₂-C₇ alkyl
25 group, a halogen atom, a C₁-C₇ alkoxy group, a C₁-C₇ alkylthio group, a C₁-C₇
alkylsulfinyl group, a C₁-C₇ alkylsulfonyl group, an amino group, and a C₁-C₇
alkylamino group; and
X represents a carboxylate.

30 19. A compound according to claim 18, wherein
said C₂-C₇ alkyl group is ethyl, or benzyl;
said halogen atom is fluorine or iodine;
said C₁-C₇ alkoxy group is methoxy or ethoxy;
said C₁-C₇ alkylthio group is methylthio, ethylthio or phenylthio;
35 said C₁-C₇ alkylsulfinyl group is ethanesulfinyl;
said C₁-C₇ alkylsulfonyl group is ethanesulfonyl;
said C₁-C₇ alkylamino group is ethylamino or diethylamino;

and

X represents a an ethylcarboxylate.

- 5
20. A compound according to claim 1, wherein R_1 and R_2 are different.
21. A compound according to claim 20 in racemic form.
22. A compound according to claim 20 in the form of its R stereoisomer.
- 10 23. A compound according to claim 20 in the form of its S stereoisomer.
24. A compound according to claim 1, wherein one of R_1 and R_2 represents a C_2 - C_7 alkyl group, and the other one represents a hydrogen atom.
- 15 25. A compound according to claim 24, wherein said alkyl group is ethyl.
26. A compound according to claim 25 in racemic form.
27. A compound according to claim 25 in the form of its R stereoisomer.
- 20 28. A compound according to claim 25 in the form of its S stereoisomer.
29. A compound according to claim 24, wherein said alkyl group is benzyl.
- 25 30. A compound according to claim 1, wherein one of R_1 and R_2 represents an alkoxy group, and the other one represents a hydrogen atom.
31. A compound according to claim 30, wherein said alkoxy group is ethoxy or methoxy.
- 30 32. A compound according to claim 1, wherein one of R_1 and R_2 represents a halogen atom, and the other one represents a hydrogen atom.
33. A compound according to claim 32, wherein said halogen atom is fluorine
- 35 or iodine.

34. A compound according to claim 1, wherein one of R₁ and R₂ represents a alkylthio group, and the other one represents a hydrogen atom.

5 35. A compound according to claim 34, wherein said alkylthiogroup is ethylthio

36. A compound according to claim 34, wherein said alkylthiogroup is methylthio or phenylthio.

10 37. A compound according to claim 1, wherein one of R₁ and R₂ represents an alkylsulfonyl group, and the other one represents a hydrogen atom.

15 38. A compound according to claim 37, wherein said alkylsulfonyl group is ethanesulfonyl.

39. A compound according to claim 1, wherein one of R₁ and R₂ represents an amino group, and the other one represents a hydrogen atom.

20 40. A compound according to claim 1, wherein one of R₁ and R₂ represents an alkyl-amino group, and the other one represents a hydrogen atom.

41. A compound according to claim 40, wherein said alkyl-amino group is ethyl-amino or diethyl-amino.

25 42. A compound according to claim 1, wherein R₁ and R₂ represent C₁-C₇-alkyl groups.

30 43. A compound according to claim 42, wherein said alkyl groups are methyl groups.

44. A compound according to claim 42, wherein said alkyl groups are ethyl groups.

35 45. A compound according to any one of the claims 20 to 44, wherein X is ethyl carboxylate.

46. A compound according to any one of the claims 1 to 45, in the form of a phospholipid, a tri-, di- or monoglyceride, or in the form of a free acid.

5 47. A compound according to any one of the claims 1 to 46 for use as a medicament.

48. A process for the manufacture of a compound according to any one of the claims 1 to 46.

10 49. A process according to claim 48, wherein said compound is prepared from (all-Z)-4,7,10,13,16,19-docosahexaenoic acid (DHA).

50. A process according to claim 49, wherein said DHA is prepared from a vegetable, a microbial and/or an animal source.

15 51. A process according to claim 50, wherein said DHA is prepared from a marine oil.

52. A process according to claim 51, wherein said marine oil is fish oil.

20 53. A pharmaceutical composition comprising a compound according to any one of the claims 1 to 46 as an active ingredient.

54. A pharmaceutical composition according to claim 53, further comprising a
25 pharmaceutically acceptable carrier.

55. A pharmaceutical composition according to claim 54, formulated for oral administration.

30 56. A pharmaceutical composition according to claim 55, in the form of a capsule or a sachet.

57. A pharmaceutical composition according to any one of the claims 53 to 56, formulated to provide a daily dosage of 10 mg to 10 g of said compound.

35 58. A pharmaceutical composition according to claim 57, formulated to provide a daily dosage of 100 mg to 1 g of said compound.

59. A fatty acid composition comprising a compound according to any one of the claims 1 to 46.

5 60. A fatty acid composition according to claim 59, wherein at least 60% by weight of the fatty acid composition is comprised of said compound.

61. A fatty acid composition according to claim 60, wherein at least 90% by weight of the fatty acid composition is comprised of said compound.

10

62. A fatty acid composition according to any one of the claims 59 to 61, further comprising fatty acids selected from (all-Z)-5,8,11,14,17-eicosapentaenoic acid (EPA), (all-Z)-4,7,10,13,16,19-docosahexaenoic acid (DHA), (all-Z)-6,9,12,15,18-heneicosapentaenoic acid (HPA), and/or (all-Z)-7,10,13,16,19-docosapentaenoic acid (DPA).

15

63. A fatty acid composition according to claim 62, wherein said fatty acids are present in the form of derivatives.

20 64. A fatty acid composition according to any one of claims 59 to 63, further comprising a pharmaceutically acceptable antioxidant.

65. A fatty acid composition according to claim 64, wherein said antioxidant is tocopherol.

25

66. A fatty acid composition according to any one of the claims 59 to 65, for use as a medicament.

30 67. Use of a compound according to any one of the claims 1 to 46 for the manufacture of a medicament for controlling body weight reduction and/or for preventing body weight gain.

68. Use of a compound according to any one of the claims 1 to 46 for the manufacture of a medicament for the treatment and/or the prevention of obesity or an
35 overweight condition.

69. Use of a compound according to any one of the claims 1 to 46 for the manufacture of a medicament for the prevention and/or treatment of diabetes in an animal.

5 70. Use according to claim 69, wherein the diabetes is type 2 diabetes.

71. Use of a compound according to any one of the claims 1 to 46 for the manufacture of a medicament for the treatment and/or prevention of amyloidosis-related diseases.

10

72. Use of a compound according to any one of the claims 1 to 46 for the manufacture of a medicament for the treatment or prophylaxis of multiple risk factors for cardiovascular diseases, preferably for the treatment of elevated blood lipids.

15 73. Use of a compound according to any one of the claims 1 to 46 for the manufacture of a medicament for prevention of stroke, cerebral or transient ischaemic attacks related to atherosclerosis of several arteries.

20 74. A method for controlling body weight reduction and/or for preventing body weight gain, wherein a pharmaceutically effective amount of a compound according to any one of the claims 1 to 46 is administered to a human or an animal.

25 75. A method for the treatment and/or the prevention of obesity or an overweight condition, wherein a pharmaceutically effective amount of a compound of formula (I) according to any one of the claims 1 to 46 is administered to a human or an animal.

30 76. A method for the prevention and/or treatment of diabetes, wherein a pharmaceutically effective amount of a compound of formula (I) according to any one of the claims 1 to 46 is administered to a human or an animal.

77. A method according to claim 76, wherein the diabetes is type 2 diabetes.

35 78. A method for the treatment and/or prevention of amyloidosis-related diseases, wherein a pharmaceutically effective amount of a compound of formula (I) according to any one of claims 1 to 46 is administered to a human or an animal.

79. A method for the treatment or prophylaxis of multiple risk factors for cardiovascular diseases, wherein a pharmaceutically effective amount of a compound of formula (I) according to any one of the claims 1 to 46 is administered to a human or an animal.


5

80. A method for the prevention of stroke, cerebral or transient ischaemic attacks related to atherosclerosis of several arteries, wherein a pharmaceutically effective amount of a compound of formula (I) according to any one of the claims 1 to 47 is administered to a human or an animal.

10

81. A method according to any one of the claims 74 to 80, wherein the compound of formula (I) is administered orally to a human or an animal.

1/13

*Fig. 1*






Fig. 2

3/13

Free fatty acid concentrations of the different PRBs in liver tissue from animals given 1,5% of PRB-1,2,5, and 7

*Fig. 3*

DHA concentrations in liver tissue from animals given 1,5% of PRB -1,2,5, and 7

*Fig. 4*

4/13




Fig. 5

5/13




Fig. 6




Fig. 7

7/13




Fig. 8




Fig. 9

8/13

B6, change of body weight during 2 weeks diet intervention after 8 weeks of HF diet




Fig. 10

Results from Luciferase assay (PPAR γ activity) including data for Rosiglitazon




Fig. 11

9/13

*Endogenous Luciferase activity in the PRBs
compared to DHA*




Fig. 12

10/13

*Glucose tolerance test before and after
giving a compound reducing insulin resistance*





**Fig. 13**


Fig. 14

Food consumption [grams/mouse/day]						
cHF	PRB2-1.5%	PRB5-5%	PRB5-1.5%	PRB7-1.5%	PRB7-0.5%	
3.19±0.05	2.89±0.07	2.77±0.11	3.13±0.10	3.34±0.09	3.27±0.12	



12/13

Fig. 15



	cHF	PRB2-1.5%	PRB5-5%	PRB5-1.5%	PRB7-1.5%	PRB7-0.5%
Fasting glucose	12	-20*	-13	-8	31	-28*
15 min	-39	-73*	-29	-45*	1	-30
30 min	-18	-125*	-15	-36	-47	-19
60 min	-35	-130*	-28	-51	2	-41
120 min	-49	-108*	-7	-8	36	-16
180 min	-43	-75*	-14	-18	24	-7
AUC	-486	-838*	-40	-194	-175	42




Fig. 16

INTERNATIONAL SEARCH REPORT

International application No.

PCT/IB2006/001155

A. CLASSIFICATION OF SUBJECT MATTER

IPC: see extra sheet

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC: C07D, A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-INTERNAL, WPI DATA, PAJ, CHEM. ABS DATA

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	EP 1310249 A1 (QUATEX N.V.), 14 May 2003 (14.05.2003), claims 1,13-14,18-19, paragraph [0021], abstract --	2-14,17, 19-73
Y	WO 2004078166 A2 (SOLVAY PHARMACEUTICAL GMBH), 16 Sept 2004 (16.09.2004), claims 8-17, abstract --	2-14,17, 19-73
Y	US 5656667 A (BREIVIK ET AL), 12 August 1997 (12.08.1997), claims 14,24,49,52, abstract --	2-14,17, 19-73

☒ Further documents are listed in the continuation of Box C.☒ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

8 August 2006

Date of mailing of the international search report

10 -08- 2006

Name and mailing address of the ISA/
Swedish Patent Office
Box 5055, S-102 42 STOCKHOLM
Facsimile No. +46 8 666 02 86

Authorized officer

Helena Melander/ELY
Telephone No. +46 8 782 25 00

INTERNATIONAL SEARCH REPORT

International application No.

PCT/IB2006/001155

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,Y	EP 1544281 A1 (KAO CORPORATION), 22 June 2005 (22.06.2005), claims 1-3,6, paragraph [0065], abstract --	2-14,17, 19-73
Y	LAILA N LARSEN et al, "Sulfur-Substituted and alpha-Methylated Fatty Acids as Peroxisome Proliferator-Activated Receptor Activators", Lipids, 2005, vol. 40, no. 1, pages 49-57, fig. 1 --	2-14,17, 19-73
A	HEGE VAAGENES et al, "Methylated Eicosapentaenoic Acid and Tetradecylthioacetic Acid: Effects on Fatty Acid Metabolism", Biochemical Pharmacology, 1999, vol. 58, pages 1133-1143, fig. 1-3 --	1-81
A	DATABAS WPI Week 199306 Derwent Publications Ltd., London, GB; Class C07C, AN 1993-233465, & JP 500974 A (SAGAMI CHEM RES CENTRE), 8 January 1993 (1993-01-08) abstract, STN-Caplus, RN 147544-49-8 --	1-81
A	US 20040162348 A1 (PEET ET AL), 19 August 2004 (19.08.2004), claims 1,13, abstract --	1-81
A	EP 1466597 A1 (CLINIGENETICS), 13 October 2004 (13.10.2004) -- -----	1-81

International patent classification (IPC)

C11C 3/00 (2006.01)
A61K 31/202 (2006.01)
A61K 31/232 (2006.01)
A61K 31/421 (2006.01)
A61P 3/04 (2006.01)
A61P 3/10 (2006.01)
A61P 9/10 (2006.01)
C07C 57/03 (2006.01)
C07C 69/587 (2006.01)
C07D 263/24 (2006.01)

Download your patent documents at www.prv.se

The cited patent documents can be downloaded at www.prv.se by following the links:

- In English/Searches and advisory services/Cited documents (service in English) or
- e-tjänster/anförda dokument (service in Swedish).

Use the application number as username.

The password is **QEWFWCSCK**.

Paper copies can be ordered at a cost of 50 SEK per copy from PRV InterPat (telephone number 08-782 28 85).

Cited literature, if any, will be enclosed in paper form.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/IB2006/001155

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 74-81
because they relate to subject matter not required to be searched by this Authority, namely:
Claims 74-81 relate to a method of treatment of the human or animal body by surgery or by therapy, as well as diagnostic
.../...
2. ☒ Claims Nos.: 1, 15-16, 18
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
The initial phase of the search revealed a very large number of documents relevant to the issue of novelty. So many
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- ☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/IB2006/001155

Box II.1

methods /Rule 39.1(iv). Nevertheless, a search has been executed for these claims. The search has been based on the alleged effects of the compounds.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/IB2006/001155

Box II.2

documents were retrieved that it is impossible to determine which parts of the claims may be said to define subject-matter for which protection might legitimately be sought (Article 6 PCT). For these reasons, a meaningful search over the whole breadth of the claims is impossible. Consequently, the search has been restricted to:

The compounds defined in any of the claims 2-14, 17 or 19-45.

INTERNATIONAL SEARCH REPORT
Information on patent family members

04/03/2006

International application No.
PCT/IB2006/001155

EP	1310249	A1	14/05/2003	IT	MI20012384 A	12/05/2003
				JP	2003146874 A	21/05/2003
				US	20030100610 A	29/05/2003
				US	20040162349 A	19/08/2004
WO	2004078166	A2	16/09/2004	AU	2004216856 A	16/09/2004
				BR	PI0408006 A	14/02/2006
				CA	2515328 A	16/09/2004
				EP	1603551 A	14/12/2005
				MX	PA05009432 A	23/11/2005
				US	20040235948 A	25/11/2004
US	5656667	A	12/08/1997	US	5698594 A	16/12/1997
				AT	191889 A	15/06/1994
				AT	398779 B	25/01/1995
				AU	616784 B	07/11/1991
				AU	3896789 A	15/02/1990
				BE	1002547 A	19/03/1991
				CA	1337548 A,C	14/11/1995
				CH	680789 A	13/11/1992
				DE	3926658 A,C	15/02/1990
				DK	173371 B	04/09/2000
				DK	392089 A	12/02/1990
				ES	2018384 A	01/04/1991
				FI	893805 A	12/02/1990
				FR	2635263 A,B	16/02/1990
				GB	2221843 A,B	21/02/1990
				GB	8819110 D	00/00/0000
				GB	8917711 D	00/00/0000
				GR	1000567 B	26/08/1992
				GR	89100507 A	22/08/1990
				HK	84196 A	24/05/1996
				IE	64524 B	09/08/1995
				IE	892317 L	11/02/1990
				IL	91275 A	19/01/1996
				IT	1235879 B	23/11/1992
				IT	8921521 D	00/00/0000
				JP	2104522 A	17/04/1990
				JP	2810916 B	15/10/1998
				KR	126286 B	26/12/1997
				LU	87570 A	08/01/1990
				NL	193459 B,C	01/07/1999
				NL	8902020 A	01/03/1990
				NZ	230022 A	26/10/1990
				SE	504742 C	14/04/1997
				SE	8902701 A	12/02/1990
				US	5502077 A	26/03/1996
				ZA	8905512 A	25/07/1990
EP	1544281	A1	22/06/2005	AU	2003252324 A	11/03/2004
				US	20050129830 A	16/06/2005
				CN	1675344 A	28/09/2005
				JP	2004083902 A	18/03/2004
				WO	2004018598 A	04/03/2004

INTERNATIONAL SEARCH REPORT
Information on patent family members

04/03/2006

International application No.
PCT/IB2006/001155

US	20040162348	A1	19/08/2004	AT	300296	T	15/08/2005
				AU	3065700	A	18/08/2000
				AU	4378099	A	05/01/2000
				BR	0007743	A	27/11/2001
				CA	2360776	A	03/08/2000
				CN	1338929	A,T	06/03/2002
				CN	1739496	A	01/03/2006
				CZ	20012695	A	13/02/2002
				DE	60021525	D,T	01/06/2006
				DK	1148873	T	28/11/2005
				EE	4629	B	15/06/2006
				EE	200100387	A	17/02/2003
				EP	1087715	A	04/04/2001
				EP	1148873	A,B	31/10/2001
				SE	1148873	T3	
				EP	1417963	A	12/05/2004
				ES	2246825	T	01/03/2006
				GB	9901809	D	00/00/0000
				HK	1039564	A	09/12/2005
				HR	20010558	A	31/08/2002
				HU	0105215	A	29/05/2002
				IL	144502	D	00/00/0000
				JP	2002535355	T	22/10/2002
				MX	PA01007662	A	24/06/2003
				NO	20013546	A	25/09/2001
				NZ	513172	A	31/10/2003
				PT	1148873	T	30/11/2005
				RU	2260423	C	20/09/2005
				SI	1148873	T	31/12/2005
				SK	10392001	A	05/02/2002
				TR	200102170	T	00/00/0000
				TW	228041	B	21/02/2005
				US	6384077	B	07/05/2002
				US	6689812	B	10/02/2004
				US	6786906	B	07/09/2004
				US	20020077361	A	20/06/2002
				US	20020183389	A	05/12/2002
				US	20020193439	A	19/12/2002
				WO	0044361	A	03/08/2000
				ZA	200106105	A	03/03/2003

EP	1466597	A1	13/10/2004	WO	2004091603	A	28/10/2004
----	---------	----	------------	----	------------	---	------------
