



US011186592B2

(12) **United States Patent**
Tanzer et al.

(10) **Patent No.:** **US 11,186,592 B2**

(45) **Date of Patent:** **Nov. 30, 2021**

(54) **THIAZOLOPYRIDINE DERIVATIVES AS
ADENOSINE RECEPTOR ANTAGONISTS**

(71) Applicant: **MERCK PATENT GMBH**, Darmstadt
(DE)

(72) Inventors: **Eva-Maria Tanzer**, Darmstadt (DE);
Kai Schiemann, Seeheim-Jugenheim
(DE); **Markus Klein**, Darmstadt (DE)

(73) Assignee: **Merck Patent GmbH**, Darmstadt (DE)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **16/635,356**

(22) PCT Filed: **Jul. 2, 2018**

(86) PCT No.: **PCT/EP2018/067797**

§ 371 (c)(1),

(2) Date: **Jan. 30, 2020**

(87) PCT Pub. No.: **WO2019/025099**

PCT Pub. Date: **Feb. 7, 2019**

(65) **Prior Publication Data**

US 2020/0181171 A1 Jun. 11, 2020

(30) **Foreign Application Priority Data**

Aug. 1, 2017 (EP) 17184272

(51) **Int. Cl.**

C07D 513/04 (2006.01)

C07D 519/00 (2006.01)

(52) **U.S. Cl.**

CPC **C07D 513/04** (2013.01); **C07D 519/00**
(2013.01)

(58) **Field of Classification Search**

CPC C07D 519/00; C07D 513/04

USPC 514/210.16

See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

2004/0235915 A1 11/2004 Flohr et al.

2005/0065151 A1 3/2005 Norcross

OTHER PUBLICATIONS


International Search Report in PCT/EP2018/067797 dated Aug. 21,
2018 (pp. 1-3).

Primary Examiner — Kabsay Habte

(74) *Attorney, Agent, or Firm* — Millen, White, Zelano &
Branigan; Harry B. Shubin

(57) **ABSTRACT**

The invention relates to thiazolopyridine derivatives of the
general formula I,



I


and the use of the compounds of the present invention for the
treatment and/or prevention of hyperproliferative or infec-
tious diseases and disorders in mammals, especially humans,
and pharmaceutical compositions containing such com-
pound.

23 Claims, No Drawings

1

THIAZOLOPYRIDINE DERIVATIVES AS ADENOSINE RECEPTOR ANTAGONISTS

The invention relates to thiazolopyridine derivatives of the general formula I,



and the use of the compounds of the present invention for the treatment and/or prevention of hyperproliferative or infectious diseases and disorders in mammals, especially humans, and pharmaceutical compositions containing such compounds.

BACKGROUND OF THE INVENTION

Adenosine is an ubiquitous modulator of numerous physiological activities, particularly within the cardiovascular, nervous and immune systems. Adenosine is related both structurally and metabolically to the bioactive nucleotides adenosine triphosphate (ATP), adenosine diphosphate (ADP), adenosine monophosphate (AMP) and cyclic adenosine monophosphate (cAMP), to the biochemical methylating agent S-adenosyl-L-methione (SAM) and structurally to the coenzymes NAD, FAD and coenzyme A and to RNA.

Via cell surface receptors, adenosine modulates diverse physiological functions including induction of sedation, vasodilatation, suppression of cardiac rate and contractility, inhibition of platelet aggregability, stimulation of gluconeogenesis and inhibition of lipolysis. Studies show that adenosine is able to activate adenylate cyclases, open potassium channels, reduce flux through calcium channels, and inhibit or stimulate phosphoinositide turnover through receptor-mediated mechanisms (Muller C. E. and Stein B., *Current Pharmaceutical Design*, 2: 501, 1996; Muller C. E., *Exp. Opin. Ther. Patents*, 7(5): 419, 1997).

Adenosine receptors belong to the superfamily of G-protein-coupled receptors (GPCRs). Four major subtypes of adenosine receptors have been pharmacologically, structurally and functionally characterized (Fredholm et al., *Pharm. Rev.*, 46: 143-156, 1994) and referred to as A₁, A_{2A}, A_{2B} and A₃. Though the same adenosine receptor can couple to different G-proteins, adenosine A₁ and A₃ receptors usually couple to inhibitory G-proteins referred to as G_i and G_o which inhibit adenylate cyclase and down-regulate cellular cAMP levels. In contrast, the adenosine A_{2A} and A_{2B} receptors couple to stimulatory G-proteins referred to as G_s, that activate adenylate cyclase and increase intracellular levels of cAMP (Linden J., *Annu. Rev. Pharmacol. Toxicol.*, 41: 775-87 2001).

According to the invention, "adenosine-receptor-selective ligands" are substances which bind selectively to one or more subtypes of the adenosine receptors, thus either mimicking the action of adenosine (adenosine agonists) or blocking its action (adenosine antagonists). According to their receptor selectivity, adenosine-receptor-selective ligands can be divided into different categories, for example ligands

2

which bind selectively to the A₁ or A₂ receptors and in the case of the latter also, for example, those which bind selectively to the A_{2A} or the A_{2B} receptors. Also possible are adenosine receptor ligands which bind selectively to a plurality of subtypes of the adenosine receptors, for example ligands which bind selectively to the A₁ and the A₂, but not to the A₃ receptors. The abovementioned receptor selectivity can be determined by the effect of the substances on cell lines which, after stable transfection with the corresponding cDNA, express the receptor subtypes in question (Olah, M. E. et al., *J. Biol. Chem.*, 267: 10764-10770, 1992). The effect of the substances on such cell lines can be monitored by biochemical measurement of the intracellular messenger cAMP (Klotz, K. N. et al., *Naunyn Schmiedebergs Arch. Pharmacol.* 357: 1-9, 1998).

It is known that the A₁ receptor system include the activation of phospholipase C and modulation of both potassium and calcium ion channels. The A₃ subtype, in addition to its association with adenylate cyclase, also stimulates phospholipase C and so activates calcium ion channels.

The A₁ receptor (326-328 amino acids) was cloned from various species (canine, human, rat, dog, chick, bovine, guinea-pig) with 90-95% sequence identity among the mammalian species. The A_{2A} receptor (409-412 amino acids) was cloned from canine, rat, human, guinea pig and mouse. The A_{2B} receptor (332 amino acids) was cloned from human and mouse with 45% homology of human A_{2B} with human A₁ and A_{2A} receptors. The A₃ receptor (317-320 amino acids) was cloned from human, rat, dog, rabbit and sheep.

The A₁ and A_{2A} receptor subtypes are proposed to play complementary roles in adenosine's regulation of the energy supply. Adenosine, which is a metabolic product of ATP, diffuses from the cell and acts locally to activate adenosine receptors to decrease the oxygen demand (A₁ and A₃) or increase the oxygen supply (A_{2A}) and so reinstate the balance of energy supply/demand within the tissue. The actions of both subtype is to increase the amount of available oxygen to tissue and to protect cells against damage caused by a short term imbalance of oxygen. One of the important functions of endogenous adenosine is preventing damage during traumas such as hypoxia, ischaemia, hypotension and seizure activity. Furthermore, it is known that the binding of the adenosine receptor agonist to mast cells expressing the rat A₃ receptor resulted in increased inositol triphosphate and intracellular calcium concentrations, which potentiated antigen induced secretion of inflammatory mediators. Therefore, the A₃ receptor plays a role in mediating asthmatic attacks and other allergic responses.

These adenosine receptors are encoded by distinct genes and are classified according to their affinities for adenosine analogues and methylxanthine antagonists (Klinger et al., *Cell Signal.*, 14 (2): 99-108, 2002).

Concerning the role of adenosine on the nervous system, the first observations were made on the effects of the most widely used of all psychoactive drugs being caffeine. Actually, caffeine is a well-known adenosine receptor antagonist that is able to enhance the awareness and learning abilities of mammals. The adenosine A_{2A} receptor pathway is responsible for these effects (Fredholm et al., *Pharmacol. Rev.*, 51 (1): 83-133, 1999; Huang et al., *Nat Neurosci.*, 8 (7): 858-9, 2005), and the effects of caffeine on the adenosine A_{2A} receptor signaling pathway encouraged the research of highly specific and potent adenosine A_{2A} antagonists.

In mammals, adenosine A_{2A} receptors have a limited distribution in the brain and are found in the striatum, olfactory tubercle and nucleus acumbens (Dixon et al., *Br. J. Pharmacol.*, 118 (6): 1461-8, 1996). High and intermediate

levels of expression can be observed in immune cells, heart, lung and blood vessels. In the peripheral system, G_3 seems to be the major G-protein associated with adenosine A_{2A} receptor but in the striatum, it has been shown that striatal adenosine A_{2A} receptors mediate their effects through activation of a G-protein referred to as G_o if (Kull et al., *Mol. Pharmacol.*, 58 (4): 772-7, 2000), which is similar to G_3 and also couples to adenylate cyclase.

To date, studies on genetically modified mice and pharmacological analysis suggest that A_{2A} receptor is a promising therapeutic target for the treatment of central nervous system (CNS) disorders and diseases such as Parkinson's disease, Huntington's disease, attention deficit hyperactivity disorders (ADHD), stroke (ischemic brain injury), and Alzheimer's disease (Fredholm et al., *Annu. Rev. Pharmacol. Toxicol.*, 45: 385-412, 2005; Higgins et al.; *Behav. Brain Res.* 185: 32-42, 2007; Dall'Igna et al., *Exp. Neurol.*, 203 (1): 241-5, 2007; Arendash et al., *Neuroscience*, 142 (4): 941-52, 2006; *Trends in Neurosci.*, 29 (11), 647-654, 2006; *Expert Opinion Ther. Patents*, 17, 979-991, 2007; *Exp. Neurol.*, 184 (1), 285-284, 2003; *Prog. Brain Res.* 183, 183-208, 2010; *J. Alzheimer Dis.*, Suppl 1, 1 17-126, 2010; *J. Neurosci.*, 29 (47), 14741-14751, 2009; *Neuroscience*, 166 (2), 590-603, 2010; *J. Pharmacol. Exp. Ther.*, 330 (1), 294-303, 2009; *Frontiers Biosci.*, 13, 2614-2632, 2008) but also for various psychoses of organic origin (Weiss et al., *Neurology*, 61 (11 Suppl 6): 88-93, 2003).

The use of adenosine A_{2A} receptor knockout mice has shown that adenosine A_{2A} receptor inactivation protects against neuronal cell death induced by ischemia (Chen et al., *J. Neurosci.*, 19 (21): 9192-200, 1999 and Monopoli et al., *Neuroreport*, 9 (17): 3955-9, 1998) and the mitochondrial toxin 3-NP (Blum et al., *J. Neurosci.*, 23 (12): 5361-9, 2003). Those results provided a basis for treating ischemia and Huntington's disease with adenosine A_{2A} antagonists. The blockade of adenosine A_{2A} receptors has also an antidepressant effect (El Yacoubi et al., *Neuropharmacology*, 40 (3): 424-32, 2001). Finally, this blockade prevents memory dysfunction (Cunha et al., *Exp. Neurol.*, 210 (2): 776-81, 2008; Takahashi et al., *Front. Biosci.*, 13: 2614-32, 2008) and this could be a promising therapeutic route for the treatment and/or prevention of Alzheimer's disease.

For reviews concerning A_{2A} adenosine receptors see e.g. Moreau et al. (*Brain Res. Reviews* 31: 65-82, 1999) and Svenningsson et al. (*Progress in Neurobiology* 59: 355-396, 1999).

To date, several adenosine A_{2A} receptor antagonists have shown promising potential for treatment of Parkinson's disease. As an example, KW-6002 (Istradefylline) completed a phase III clinical trial in the USA after studies demonstrated its efficacy in alleviation of symptoms of the disease (Bara-Himenez et al., *Neurology*, 61 (3): 293-6, 2003 and Hauser et al., *Neurology*, 61 (3): 297-303, 2003). SCH420814 (Preladenant), which is now in phase II clinical trial in the USA and produces an improvement in motor function in animal models of Parkinson's disease (Neustadt et al., *Bioorg. Med. Chem. Lett.*, 17 (5): 1376-80, 2001) and also in human patients (Hunter J. C., poster Boston 2006—<http://www.a2apd.org/Speakerabstracts/Hunter.pdf>).

Besides the welcome utility of A_{2A} receptor antagonists to treat neurodegenerative diseases, those compounds have been considered for complementary symptomatic indications. These are based on the evidence that A_{2A} receptor activation may contribute to the pathophysiology of a range of neuropsychiatric disorders and dysfunctions such as depression, excessive daytime sleepiness, restless legs syndrome, attention deficit hyperactivity disorder, and cognitive

fatigue (*Neurology*, 61 (Suppl 6), 82-87, 2003; *Behav. Pharmacol.*, 20 (2), 134-145, 2009; *CNS Drug Discov.*, 2 (1), 1-21, 2007).

Some authors suggest the application of A_{2A} antagonists for the treatment of diabetes (WO1999035147; WO2001002400). Other studies suggest the involvement of A_{2A} adenosine receptors in wound healing or atrial fibrillation (*Am. J. Path.*, 6, 1774-1778, 2007; *Arthritis & Rheumatism*, 54 (8), 2632-2642, 2006).

Some of the potent adenosine A_{2A} antagonists discovered in the past by the pharmaceutical companies, have advanced into clinical trials showing positive results and demonstrating the potential of this compound class for the treatment of neurodegenerative disorders like Parkinson's, Huntington's or Alzheimer's disease, but also in other CNS related diseases like depression, restless syndrome, sleep and anxiety disorders (*Clin. Neuropharmacol.*, 33, 55-60, 2010; *J. Neurosci.*, 30 (48), 2010), 16284-16292; *Parkinson Relat. Disord.*, 16 (6), 423-426, 2010; *Expert Opinion Ther. Patents*, 20(8), 987-1005, 2010; *Current Opinion in Drug Discovery & Development*, 13 (4), 466-480, 2010 and references therein; *Mov. Disorders*, 25 (2), S305, 2010).

Known A_{2A} inhibitors are Istradefylline (KW-6002), Preladenant (SCH420814), SCH58261, CGS15943, Tozadenant, Vipadenant (V-2006), V-81444 (CPI-444, HTL-1071, PBF-509, Medi-9447, PNQ-370, ZM-241385, ASO-5854, ST-1535, ST-4206, DT1133 and DT-0926, which are in most cases developed for Parkinson's disease.

Adenosine A_{2B} receptors were cloned from rat hypothalamus (Rivkees and Reppert, 1992), human hippocampus (Pierce et al., 1992), and mouse mast cells (Marquardt et al., 1994), employing standard polymerase chain reaction techniques with degenerate oligonucleotide primers designed to recognize conserved regions of most G protein-coupled receptors. The human A_{2B} receptor shares 86 to 87% amino acid sequence homology with the rat and mouse A_{2B} receptors (Rivkees and Reppert, 1992; Pierce et al., 1992; Marquardt et al., 1994) and 45% amino acid sequence homology with human A_1 and A_{2A} receptors. As expected for closely related species, the rat and mouse A_{2B} receptors share 96% amino acid sequence homology. By comparison, the overall amino acid identity between A_1 receptors from various species is 87% (Palmer and Stiles, 1995). A_{2A} receptors share 90% of homology between species (Ongini and Fredholm, 1996), with most differences occurring in the 2nd extracellular loop and the long C-terminal domain (Palmer and Stiles, 1995). The lowest (72%) degree of identity between species is observed for A_3 receptor sequences (Palmer and Stiles, 1995).

The adenosine analog NECA remains the most potent A_{2B} agonist (Bruns, 1981; Feoktistov and Biaggioni, 1993, 1997; Brackett and Daly, 1994), with a concentration producing a half-maximal effect (EC_{50}) for stimulation of adenylyl cyclase of approximately 2 μ M. It is, however, nonselective and activates other adenosine receptors with even greater affinity, with an EC_{50} in the low nanomolar (A_1 and A_{2A}) or high nanomolar (A_3) range. The characterization of A_{2B} receptors, therefore, often relies on the lack of effectiveness of compounds that are potent and selective agonists of other receptor types. A_{2B} receptors have been characterized by a method of exclusion, i.e., by the lack of efficacy of agonists that are specific for other receptors. The A_{2A} selective agonist CGS-21680 (Webb et al., 1992), for example, has been useful in differentiating between A_{2A} and A_{2B} adenosine receptors (Hide et al., 1992; Chern et al., 1993; Feoktistov and Biaggioni, 1995; van der Ploeg et al., 1996). Both receptors are positively coupled to adenylyl cyclase and are

activated by the nonselective agonist NECA. CGS-21680 is virtually ineffective on A_{2B} receptors but is as potent as NECA in activating A_{2A} receptors, with an EC_{50} in the low nanomolar range for both agonists (Jarvis et al., 1989; Nakane and Chiba, 1990; Webb et al., 1992; Hide et al., 1992; Feoktistov and Biaggioni, 1993; Alexander et al., 1996). A_{2B} receptors have also a very low affinity for the A_1 selective agonist R-PIA (Feoktistov and Biaggioni, 1993; Brackett and Daly, 1994) as well as for the A_3 selective agonist N^6 -(3-iodobenzyl)- N -methyl-5'-carbamoyladen-
 5 osine (IB-MECA) (Feoktistov and Biaggioni, 1997). The agonist profile $NECA > R-PIA = IB-MECA > CGS-21680$ was determined in human erythroleukemia (HEL) cells for A_{2B} -mediated cAMP accumulation. The difference between EC_{50} for NECA and the rest of the agonists is approximately 2
 10 orders of magnitude. Therefore, responses elicited by NECA at concentrations in the low micromolar range (1-10 μ M), but not by R-PIA, IB-MECA or CGS-21680, are character-
 15 istic of A_{2B} receptors.

Whereas A_{2B} receptors have, in general, a lower affinity for agonists compared to other receptor subtypes, this is not true for antagonists. The structure activity relationship of adenosine antagonists on A_{2B} receptors has not been fully
 20 characterized, but at least some xanthines are as or more potent antagonists of A_{2B} receptor subtypes than of other subtypes. In particular, DPSPX (1,3-dipropyl-8-sulphophe-
 25 nylxanthine), DPCPX (1,3-dipropyl-8c-yclopentylxanthine), DPX (1,3 diethylphenylxanthine), the antiasthmatic drug enprofylline (3-n-propylxanthine) and the non-xanthine
 30 compound 2,4-dioxobenzopteridine (alloxazine) have affinities in the mid to high nM range.

Other known A_{2B} inhibitors are ATL801, PSB-605, PSB-1115, ISAM-140, GS6201, MRS1706 and MRS1754.

It is disclosed herein that adenosine receptors play a
 35 non-redundant role in down-regulation of inflammation in vivo by acting as a physiological "STOP" (a termination mechanism) that can limit the immune response and thereby
 40 protect normal tissues from excessive immune damage during pathogenesis of different diseases.

A_{2A} receptor antagonists provide long term enhancement of immune responses by reducing T-cell mediated tolerance to antigenic stimuli, enhancing the induction of memory T cells and enhancing the efficacy of passive antibody admin-
 45 istration for the treatment of cancer and infectious diseases while A_{2A} receptor agonists provide long term reduction of immune responses by enhancing T-cell mediated tolerance to antigenic stimuli, in particular to reduce use of immuno-
 50 suppressive agents in certain conditions.

Immune modulation is a critical aspect of the treatment of
 55 a number of diseases and disorders. T cells in particular play a vital role in fighting infections and have the capability to recognize and destroy cancer cells. Enhancing T cell mediated responses is a key component to enhancing responses to therapeutic agents. However, it is critical in
 60 immune modulation that any enhancement of an immune response is balanced against the need to prevent autoimmunity as well as chronic inflammation. Chronic inflammation and self-recognition by T cells is a major cause for the pathogenesis of systemic disorders such as rheumatoid
 65 arthritis, multiple sclerosis and systemic lupus erythematosus. Furthermore, long term immunosuppression is required in preventing rejection of transplanted organs or grafts.

Tumor-induced immunosuppression is a major hurdle to the efficacy of current cancer therapies. Because of their
 70 remarkable clinical efficacy against a broader range of cancers, recent successes with immune checkpoint blockade

inhibitors such as anti-CTLA-4 and anti-PD-1/PDL1 are revolutionizing cancer treatment.

Adenosine is one of the new promising immunosuppressive targets revealed in preclinical studies. This metabolite is produced by the ectoenzyme—CD73 expressed on host
 5 suppressor cells and tumor cells. Increased expression of CD73 correlates with poor prognosis in patients with a number of cancers, including colorectal cancer (Liu et al., J. Surgical Oncol, 2012), gastric cancer (Lu et al., World J. Gastroenterol., 2013), gallbladder cancer (Xiong et al., Cell and Tissue Res., 2014). Preclinical studies demonstrated that
 10 protumor effects of CD73 can be driven (at least in part) by adenosine-mediated immunosuppression. As disclosed above, adenosine binds to four known receptors A_1 , A_{2A} ,
 15 A_{2B} , and A_3 , with the activation of A_{2A} and A_{2B} receptors known to suppress the effector functions of many immune cells, i.e. A_{2A} and A_{2B} receptors induce adenylylate-cyclase-
 20 dependent accumulation of cAMP leading to immunosuppression. Since antagonizing A_1 and A_3 would counteract the desired effect and A_1 and A_3 agonists serve as potential cardioprotective agents, selectivity towards A_1 and A_3 needs to be achieved (Antonioli et al., Nat. rev. Cancer, 2013, Thiel et al., Microbes and Infection, 2003). In the microenvironment of the tumor, both A_{2A} and A_{2B} receptor activation has
 25 been demonstrated to suppress antitumor immunity and increase the spread of CD73 tumors. In addition, either A_{2A} or A_{2B} blockade with small molecule antagonists can reduce tumor metastasis. It has been found that blocking of A_{2A}
 30 receptor can overcome tumor escape mechanisms including both anergy and regulatory T cell induction caused by tumor cells and cause long-term tumor susceptibility to treatment. Ohta et al. demonstrated rejection of approximately 60% of established CL8-1 melanoma tumors in A_{2A} receptor-deficient mice compared to no rejection in normal mice (Ohta, et al.; PNAS 103 (35): 13132-7, 2006). In agreement, the investigators also showed improved inhibition of tumor growth, destruction of metastases and prevention of neovas-
 35 cularization by anti-tumor T cells after treatment with an A_{2A} receptor antagonist.

Tumors have been shown to evade immune destruction by impeding T cell activation through inhibition of co-stimulatory factors in the B7-CD28 and TNF families, as well as by attracting regulatory T cells, which inhibit anti-tumor T cell responses (Wang, Cancer. Semin. Cancer. Biol. 16: 73-79, 2006; Greenwald, et al., Ann. Rev. Immunol. 23: 515-48, 2005; Watts, Ann. Rev. Immunol. 23: 23-68, 2005; Sadum et al., Clin. Cane. Res. 13 (13): 4016-4025, 2007). Because A_{2A} receptor expression is increased in lymphocytes following activation, therapies that liberate lymphocyte effector responses, such as anti-CTLA-4 and anti-PD-1, may also increase the effects of A_{2A} -mediated immunosuppression. Immune checkpoint blockade in combination with
 45 A_{2A} or dual $A_{2A/2B}$ antagonists increase the magnitude of immune responses to tumors and metastasis. Accordingly, combination of A_{2A} inhibition with anti-PD-1 therapy enhances IFN- γ production by T-cells in a co-culture with MC38 tumor cells, improves mouse survival in 4T1 mammary tumor model and decreases tumor growth in
 50 AT-3ova^{dim} CD73+ tumors (Beavis et al., Cancer Immunol. Res., 2015; Mittal et al., Cancer Res., 2014).

Furthermore, preclinical studies demonstrated that A_{2B} inhibition leads to decreased tumor growth and extended survival of mice in Lewis lung carcinoma, MB49 bladder carcinoma, ortho 4T1 mammary carcinoma models (Ryzhov et al., 2009, Cekic et al., 2012) and the combination of A_{2B} inhibition with anti-PD-1 therapy reduces lung metastases of

B16-F10 melanoma tumors and improves mouse survival in the 4T1 mammary tumor model.

WO 03/050241 describes the methods to increase an immune response to an antigen, increasing vaccine efficacy or increasing an immune response to a tumor antigen or immune cell-mediated tumor destruction by administering an agent that inhibits extracellular adenosine or inhibits adenosine receptors.

WO 2004/089942, WO 2005/000842 and WO 2006/008041 disclose benzothiazole derivatives, including Tozadenant, as A_{2A} inhibitors for the treatment of Parkinson's disease. WO 2004/092171 and WO 2005/028484 disclose similar thiazolopyridine and pyrazolopyrimidine derivatives also as A_{2A} inhibitors for the treatment of Parkinson's disease. However, these compounds do not show significant A_{2B} inhibitory activity and do only show good pharmacokinetic properties in the rat, the Parkinson's disease animal model but not in the mouse, the cancer animal model. Furthermore, the compounds do not show that they are able to prevent immunosuppression and thus are able to support anti-tumor T cell induced inhibition of tumor growth, reduction or destruction of metastases and prevention of neovascularization.

Thus, there remains a need for therapies that provide long term enhancement of immune responses to specific antigens, particularly for the treatment and prevention of hyperproliferative and infectious diseases and disorders and thus the object of the present invention was to provide methods of treatment that allow simplified treatment protocols and enhance immune responses against certain antigens. It was a specific object of the invention to provide improved methods of preventing or treating hyperproliferative and infectious diseases and disorders in a host, especially to provide effective A_{2A} or dual $A_{2A/2B}$ antagonists for the treatment and prevention of such diseases.

SUMMARY OF THE INVENTION

Surprisingly, it has been found that the thiazolopyridine derivatives according to the invention are highly effective inhibitors of the A_{2A} adenosine receptor or both the A_{2A} and A_{2B} adenosine receptors and at the same time have high selectivity over the A_1 and A_3 adenosine receptors, and thus the compounds of the present invention can be used for the treatment of hyperproliferative diseases and disorders such as cancer and infectious diseases and disorders.


Particularly, in contrast to the known adenosine A_{2A} receptor antagonist Tozadenant and similar benzothiazole derivatives, the compounds of the present invention surprisingly show an A_{2A}/A_{2B} dual activity which is preferred for the treatment and/or prevention of hyperproliferative and infectious diseases and disorders as it is disclosed above or the compounds of the present invention show at least a high A_{2A} inhibitory activity together with the other surprising advantages disclosed herein leading to a high efficacy in the treatment and/or prevention of hyperproliferative and infectious diseases and disorders.

Additionally, in comparison with the known adenosine A_{2A} receptor antagonist Tozadenant and similar benzothiazole derivatives, the compounds of the present invention surprisingly show better pharmacokinetic properties in mouse as the animal model relevant for cancer, which is preferred for the treatment and/or prevention of hyperproliferative and infectious diseases and disorders as it is disclosed above.

Furthermore, as discussed above, adenosine in tumor microenvironment can inhibit T cell activity by signaling

through A_{2A} receptors and suppress cytokine secretion by T cells. A_{2A} specific agonists like CGS-21680, similar to adenosine, inhibit T cell cytokine secretion in vitro and in vivo. In contrast, potential A_{2A} antagonists or A_{2A}/A_{2B} dual antagonists can rescue T cells from this inhibition. In contrast to the known adenosine A_{2A} receptor antagonist Tozadenant, the compounds of the present invention show that they are able to rescue T cells from inhibition and are able to prevent the suppression of cytokine secretion as induced by adenosine or A_{2A} specific agonists like CGS-2168, which is preferred for the treatment and/or prevention of hyperproliferative and infectious diseases and disorders as it is disclosed above. Therefore, the compounds of the present invention surprisingly are able to prevent immunosuppression and thus are able to support anti-tumor T cell induced inhibition of tumor growth, reduction or destruction of metastases and prevention of neovascularization.

The invention relates to thiazolopyridine derivatives of the general formula I,



wherein

R^1 is linear or branched alkyl having 1-10 C atoms which is unsubstituted or mono-, di- or trisubstituted by R^5 and in which 1-4 C atoms may be replaced, independently of one another, by O, S, SO, SO_2 , NH, NCH_3 , $-OCO-$, $-NHCONH-$, $-NHCO-$, $-NR^6SO_2R^7-$, $-COO-$, $-CONH-$, $-NCH_3CO-$, $-CONCH_3-$, $-C\equiv C-$ groups and/or $-CH=CH-$ groups, and/or, in addition, 1-10H atoms may be replaced by F and/or Cl, or mono- or bicyclic cyclic alkyl having 3-7 C atoms which is unsubstituted or mono-, di- or trisubstituted by R^5 and in which 1-4 C atoms may be replaced, independently of one another, by O, S, SO, SO_2 , NH, NCH_3 , $-OCO-$, $-NHCONH-$, $-NHCO-$, $-NR^6SO_2R^7-$, $-COO-$, $-CONH-$, $-NCH_3CO-$, $-CONCH_3-$, $-C\equiv C-$ groups and/or by $-CH=CH-$ groups and/or, in addition, 1-10H atoms may be replaced by F and/or Cl, or mono- or bicyclic heteroaryl, heterocyclyl, aryl or cyclic alkylaryl, containing 3 to 14 carbon atoms and 0-4 heteroatoms, independently selected from N, O and S, which is unsubstituted or mono-, di- or trisubstituted by R^5 ,

R^2 is linear or branched alkyl having 1-10 C atoms which is unsubstituted or mono-, di- or trisubstituted by R^5 and in which 1-4 C atoms may be replaced, independently of one another, by O, S, SO, SO_2 , NH, NCH_3 , $-OCO-$, $-NHCONH-$, $-NHCO-$, $-NR^6SO_2R^7-$, $-COO-$, $-CONH-$, $-NCH_3CO-$, $-CONCH_3-$, $-C\equiv C-$ groups and/or $-CH=CH-$ groups, and/or, in addition, 1-10H atoms may be replaced by F and/or Cl, or cyclic alkyl having 3-7 C atoms which is unsubstituted or mono-, di- or trisubstituted by R^5 and in which 1-4 C atoms may be replaced, independently of one another, by O, S, SO, SO_2 , NH, NCH_3 , $-OCO-$, $-NHCONH-$, $-NHCO-$, $-NR^6SO_2R^7-$, $-COO-$, $-CONH-$, $-NCH_3CO-$, $-CONCH_3-$, $-C\equiv C-$ groups and/or

9

by $-\text{CH}=\text{CH}-$ groups and/or, in addition, 1-11H atoms may be replaced by F and/or Cl, or mono- or bicyclic heteroaryl, heterocyclyl, aryl or cyclic alkylaryl, containing 3 to 14 carbon atoms and 0-4 heteroatoms, independently selected from N, O and S, which is unsubstituted or mono-, di- or trisubstituted by R^5 ,

R^3 is linear or branched alkyl or O-alkyl having 1-6 C atoms or cyclic alkyl having 3-6 C atoms, which is unsubstituted or mono-, di- or trisubstituted by H, $=\text{S}$, $=\text{NH}$, $=\text{O}$, OH, cyclic alkyl having 3-6 C atoms, COOH , Hal, NH_2 , SO_2CH_3 , SO_2NH_2 , CN, CONH_2 , NHCOCH_3 , NHCONH_2 or NO_2 ,

R^4 is H, D, linear or branched alkyl having 1-6 C atoms or Hal,

R^5 is H, R^6 , $=\text{S}$, $=\text{NR}^6$, $=\text{O}$, OH, COOH , Hal, NH_2 , SO_2CH_3 , SO_2NH_2 , CN, CONH_2 , NHCOCH_3 , NHCONH_2 , NO_2 , or linear or branched alkyl having 1-10 C atoms which is unsubstituted or mono-, di- or trisubstituted by R^6 and in which 1-4 C atoms may be replaced, independently of one another, by O, S, SO, SO_2 , NH, NCH_3 , $-\text{OCO}-$, $-\text{NHCONH}-$, $-\text{NHCO}-$, $-\text{NR}^6\text{SO}_2\text{R}^7-$, $-\text{COO}-$, $-\text{CONH}-$, $-\text{NCH}_3\text{CO}-$, $-\text{CONCH}_3-$, $-\text{C}=\text{C}-$ groups and/or $-\text{CH}=\text{CH}-$ groups, and/or, in addition, 1-10H atoms may be replaced by F and/or Cl, or mono- or bicyclic cyclic alkyl having 3-7 C atoms which is unsubstituted or mono-, di- or trisubstituted by R^6 and in which 1-4 C atoms may be replaced, independently of one another, by O, S, SO, SO_2 , NH, NCH_3 , $-\text{OCO}-$, $-\text{NHCONH}-$, $-\text{NHCO}-$, $-\text{NR}^6\text{SO}_2\text{R}^7-$, $-\text{COO}-$, $-\text{CONH}-$, $-\text{NCH}_3\text{CO}-$, $-\text{CONCH}_3-$, $-\text{C}=\text{C}-$ groups and/or by $-\text{CH}=\text{CH}-$ groups and/or, in addition, 1-10H atoms may be replaced by F and/or Cl, or mono- or bicyclic heteroaryl, heterocyclyl, aryl or cyclic alkylaryl, containing 3 to 14 carbon atoms and 0-4 heteroatoms, independently selected from N, O and S, which is unsubstituted or mono-, di- or trisubstituted by R^6 ,

R^6 , R^7 are independently of one another selected from the group consisting of H, $=\text{S}$, $=\text{NH}$, $=\text{O}$, OH, COOH , Hal, NH_2 , SO_2CH_3 , SO_2NH_2 , CN, CONH_2 , NHCOCH_3 , NHCONH_2 , NO_2 and linear or branched alkyl having 1-10 C atoms in which 1-4 C atoms may be replaced, independently of one another, by O, S, SO, SO_2 , NH, NCH_3 , $-\text{OCO}-$, $-\text{NHCONH}-$, $-\text{NHCO}-$, $-\text{COO}-$, $-\text{CONH}-$, $-\text{NCH}_3\text{CO}-$, $-\text{CONCH}_3-$, $-\text{C}=\text{C}-$ groups and/or $-\text{CH}=\text{CH}-$ groups, and/or, in addition, 1-10H atoms may be replaced by F and/or Cl,


Hal is F, Cl, Br, or I,

D is deuterium

and physiologically acceptable salts, derivatives, solvates, prodrugs and stereoisomers thereof, including mixtures thereof in all ratios.


The invention preferably relates to a compound of formula I, wherein R^1 is linear or branched alkyl having 1-10 C atoms which is unsubstituted or mono-, di- or trisubstituted by R^4 and in which 1-4 C atoms may be replaced, independently of one another, by O, S, SO, SO_2 , NH, NCH_3 , $-\text{OCO}-$, $-\text{NHCONH}-$, $-\text{NHCO}-$, $-\text{NR}^5\text{SO}_2\text{R}^6-$, $-\text{COO}-$, $-\text{CONH}-$, $-\text{NCH}_3\text{CO}-$, $-\text{CONCH}_3-$, $-\text{C}=\text{C}-$ groups and/or $-\text{CH}=\text{CH}-$ groups, and/or, in addition, 1-10H atoms may be replaced by F and/or Cl, or one of the following structures:

10




which is unsubstituted or mono-, di- or trisubstituted with R^5 and wherein R^2 , R^3 , R^4 , R^5 , R^6 and R^7 have the meanings as disclosed above.

The invention particularly preferably relates to a compound of formula I, wherein R^1 is one of the following structures:



11


-continued



and wherein R², R³, R⁴, R⁵, R⁶ and R⁷ have the meanings as disclosed above.


The invention preferably relates to a compound of formula I, wherein R¹ is phenyl, methylpyrazole or dihydropyran and R², R³, R⁴, R⁵, R⁶ and R⁷ have the meanings as disclosed above.

The invention particularly preferably relates to a compound of formula I, wherein R² is one of the following structures:



12

-continued




which is unsubstituted or mono-, di- or trisubstituted with R⁶

and wherein R¹, R³, R⁴, R⁵, R⁶ and R⁷ have the meanings as disclosed above.


The invention particularly preferably relates to a compound of formula I, wherein

R² is one of the following structures:




13

-continued




14

-continued




15



and wherein R^1 , R^3 , R^4 , R^5 , R^6 and R^7 have the meanings as disclosed above.

The invention preferably relates to a compound of formula I, wherein

R^3 one of the following structures



and R^1 , R^2 , R^4 , R^5 , R^6 and R^7 have the meanings as disclosed above.

The invention preferably relates to a compound of formula I, wherein

R^3 is O-alkyl having 1-6 C atoms, which is unsubstituted or mono-, di- or trisubstituted with F and R^1 , R^2 , R^4 , R^5 , R^6 and R^7 have the meanings as disclosed above.

The invention preferably relates to a compound of formula I, wherein

R^3 is OMe

and R^1 , R^2 , R^4 , R^5 , R^6 and R^7 have the meanings as disclosed above.

The invention particularly preferably relates to a compound of formula I, wherein

16

R^1 is phenyl, methylpyrazole or dihydropyran,

R^3 is OMe

and R^2 , R^4 , R^5 , R^6 and R^7 have the meanings as disclosed above.

The invention preferably relates to a compound of formula I, wherein

R^4 is H, D, methyl, ethyl, F, Br or Cl

and wherein R^1 , R^2 , R^3 , R^5 , R^6 and R^7 have the meanings as disclosed above.

The invention preferably relates to a compound of formula I, wherein

R^4 is H, D, methyl, F, Br or Cl

and wherein R^1 , R^2 , R^3 , R^5 , R^6 and R^7 have the meanings as disclosed above.

The invention preferably relates to a compound of formula I, wherein

R^4 is H

and wherein R^1 , R^2 , R^3 , R^5 , R^6 and R^7 have the meanings as disclosed above.

The invention particularly preferably relates to a compound selected from the group consisting of:

- 1 4-Hydroxy-4-methyl-piperidine-1-carboxylic acid (4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide
- 2 4-Dimethylaminomethyl-N-(4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-benzamide
- 3 4-Methoxymethyl-N-(4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-benzamide
- 4 1-(2-Methoxy-ethyl)-1H-pyrazole-4-carboxylic acid (4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide
- 5 2-Dimethylaminomethyl-N-(4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-isonicotinamide
- 6 2-Methoxymethyl-N-(4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-isonicotinamide
- 7 4-benzyl-4-hydroxy-N-[4-methoxy-7-(morpholin-4-yl)-[1,3]thiazolo[4,5-c]pyridin-2-yl]piperidine-1-carboxamide
- 8 N-[4-methoxy-7-(morpholin-4-yl)-[1,3]thiazolo[4,5-c]pyridin-2-yl]-5-(2-methoxyethoxy)pyrazine-2-carboxamide
- 9 2-Oxa-7-aza-spiro[4.4]nonane-7-carboxylic acid (4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide
- 10 3-Hydroxy-3-methyl-pyrrolidine-1-carboxylic acid (4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide
- 11 8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid (4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide
- 12 4-Hydroxy-4-prop-2-ynyl-piperidine-1-carboxylic acid(4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide
- 13 1-(2-Methoxy-ethyl)-1H-pyrazole-4-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 14 4-Methoxymethyl-N-[4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-benzamide
- 15 N-(4-Methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-N',N'-dimethyl-terephthalamide
- 16 4-Hydroxymethyl-4-methyl-piperidine-1-carboxylic acid (4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide
- 17 (5S)-N-[4-methoxy-7-(morpholin-4-yl)-[1,3]thiazolo[4,5-c]pyridin-2-yl]-2-oxa-7-azaspiro[4.4]nonane-7-carboxamide
- 18 (5R)-N-[4-methoxy-7-(morpholin-4-yl)-[1,3]thiazolo[4,5-c]pyridin-2-yl]-2-oxa-7-azaspiro[4.4]nonane-7-carboxamide
- 19 N-[4-Methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-2-benzamide
- 20 4-Methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-ylamine
- 21 4-Hydroxy-4-methyl-piperidine-1-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 22 4-Dimethylaminomethyl-N-[4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-benzamide
- 23 4-Methoxymethyl-N-[4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-benzamide
- 24 2-Dimethylaminomethyl-N-[4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-isonicotinamide
- 25 2-Dimethylaminomethyl-N-[4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-isonicotinamide
- 26 2-Methoxymethyl-N-[4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]isonicotinamide

27 4-Dimethylaminomethyl-N-[4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-benzamide

28 2-Methoxymethyl-N-[4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-isonicotinamide

29 1-(2-Methoxy-ethyl)-1H-pyrazole-4-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-2-amide

30 4-Hydroxy-4-methyl-piperidine-1-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide

31 Isoxazole-4-carboxylic acid (4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide

32 4-Hydroxy-4-methyl-piperidine-1-carboxylic acid (4-fluoromethoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide

33 4-Imidazol-1-ylmethyl-N-(4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-benzamide

34 7-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid (4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide

35 1-(2-Methoxy-ethyl)-1H-pyrazole-4-carboxylic acid (4-methoxy-7-pyridin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide

36 1-(2-Methoxy-ethyl)-1H-pyrazole-4-carboxylic acid [4-methoxy-7-(6-methyl-pyridazin-3-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide

37 4-Hydroxy-4-methyl-piperidine-1-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

38 4-Difluoromethyl-4-hydroxy-piperidine-1-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

39 4-Hydroxymethyl-4-methyl-piperidine-1-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

40 4-Fluoromethyl-4-hydroxy-piperidine-1-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

41 3-Oxa-9-aza-spiro[5.5]undecane-9-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

42 4-Methyl-piperidine-1-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

43 4-Benzyl-4-hydroxy-piperidine-1-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

44 2-Oxo-1-oxa-3,8-diaza-spiro[4.5]decane-8-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

45 3-Oxo-2,8-diaza-spiro[4.5]decane-8-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

46 4-Dimethylaminomethyl-N-(4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-benzamide

47 4-Methoxymethyl-N-(4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-benzamide

48 2,4-Dioxo-1,3,8-triaza-spiro[4.5]decane-8-carboxylic acid(4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

49 4'-Hydroxy-3',4',5',6'-tetrahydro-2'H-[3,4]bipyridinyl-1'-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

50 4-Oxo-piperidine-1-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

51 1-Oxo-2,8-diaza-spiro[4.5]decane-8-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

52 7-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

53 8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

54 2-Oxa-7-aza-spiro[4.4]nonane-7-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

55 4-Imidazol-1-ylmethyl-N-(4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-benzamide

56 Isoxazole-3-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

57 4-Hydroxy-4-prop-2-ynyl-piperidine-1-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

58 N-(4-Methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-N',N'-dimethyl-terephthalamide

59 N-(4-Methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-4-trifluoromethoxy-benzamide

60 2-Methyl-oxazole-4-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

61 Benzooxazole-5-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

62 N-(4-Methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-4-(2-oxo-pyrrolidin-1-ylmethyl)-benzamide

63 2,3-Dihydro-benzofuran-5-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

64 3-Methoxymethyl-pyrrolidine-1-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

65 Piperidine-1,4-dicarboxylic acid 4-amide 1-[(4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide]

66 4-Diethylamino-N-(4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-benzamide

5 67 4-Difluoromethyl-4-hydroxy-piperidine-1-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide

68 4-Hydroxymethyl-4-methyl-piperidine-1-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide

69 4-Fluoromethyl-4-hydroxy-piperidine-1-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide

10 70 3-Oxa-9-aza-spiro[5.5]undecane-9-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide

71 4-Methyl-piperidine-1-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide

72 4-Benzyl-4-hydroxy-piperidine-1-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide

15 73 2-Oxo-1-oxa-3,8-diaza-spiro[4.5]decane-8-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide

74 3-Oxo-2,8-diaza-spiro[4.5]decane-8-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide

20 75 4-Dimethylaminomethyl-N-[4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-benzamide

76 2,4-Dioxo-1,3,8-triaza-spiro+4.5]decane-8-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide

77 4'-Hydroxy-3',4',5',6'-tetrahydro-2'H-[3,4]bipyridinyl-1'-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide

25 78 4-Oxo-piperidine-1-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]amide

79 1-Oxo-2,8-diaza-spiro[4.5]decane-8-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide

30 80 7-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide

81 8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide

82 2-Oxa-7-aza-spiro[4.4]nonane-7-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide

35 83 4-Imidazol-1-ylmethyl-N-[4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-benzamide

84 Isoxazole-3-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide

85 4-Hydroxy-4-prop-2-ynyl-piperidine-1-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide

40 86 N-[4-Methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-N',N'-dimethyl-terephthalamide

87 N-[4-Methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-4-trifluoromethoxy-benzamide

88 2-Methyl-oxazole-4-carboxylic acid[4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide

89 Benzooxazole-5-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide

45 90 N-[4-Methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-4-(2-oxo-pyrrolidin-1-ylmethyl)-benzamide

91 2,3-Dihydro-benzofuran-5-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide

92 3-Methoxymethyl-pyrrolidine-1-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide

93 Piperidine-1,4-dicarboxylic acid 4-amide 1-[[4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide]

94 4-Diethylamino-N-[4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-benzamide

95 4-Difluoromethyl-4-hydroxy-piperidine-1-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide

55 96 4-Hydroxymethyl-4-methyl-piperidine-1-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide

97 4-Fluoromethyl-4-hydroxy-piperidine-1-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide

60 98 3-Oxa-9-aza-spiro[5.5]undecane-9-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide

99 4-Methyl-piperidine-1-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide

65 100 4-Benzyl-4-hydroxy-piperidine-1-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide

- 101 2-Oxo-1-oxa-3,8-diaza-spiro[4.5]decane-8-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 102 3-Oxo-2,8-diaza-spiro[4.5]decane-8-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 103 2,4-Dioxo-1,3,8-triaza-spiro[4.5]decane-8-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 104 4'-Hydroxy-3',4',5',6'-tetrahydro-2H-[3,4']bipyridinyl-1'-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 105 4-Oxo-piperidine-1-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 106 1-Oxo-2,8-diaza-spiro[4.5]decane-8-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 107 7-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 108 8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 109 2-Oxa-7-aza-spiro[4.4]nonane-7-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 110 4-Imidazol-1-ylmethyl-N-[4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-benzamide
- 111 Isoxazole-3-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]amide
- 112 4-Hydroxy-4-prop-2-ynyl-piperidine-1-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 113 N-[4-Methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-N',N'-dimethyl-terephthalamide
- 114 N-[4-Methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-4-trifluoromethoxy-benzamide
- 115 2-Methyl-oxazole-4-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 116 Benzoxazole-5-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 117 N-[4-Methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-4-(2-oxo-pyrrolidin-1-ylmethyl)-benzamide
- 118 2,3-Dihydro-benzofuran-5-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 119 3-Methoxymethyl-pyrrolidine-1-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 120 Piperidine-1,4-dicarboxylic acid 4-amide 1-{[4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide}
- 121 4-Diethylamino-N-[4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-benzamide
- 122 {1-[4-Methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]carbamoyl}-piperidin-3-yl]-acetic acid
- 123 Pyridine-2,5-dicarboxylic acid 2-dimethylamide 5-([4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide)
- 124 1-Methyl-1H-pyrazole-4-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 125 5-Methyl-isoxazole-4-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 126 1-(2-Methoxy-ethyl)-1H-[1,2,3]triazole-4-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 127 1-Methyl-1H-[1,2,3]triazole-4-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 128 1-Cyano-cyclopropanecarboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 129 3-Cyano-N-[4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-propionamide
- 130 2-Methyl-oxazole-5-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 131 2-Methyl-thiazole-5-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 132 4-Hydroxy-pent-2-ynoic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 133 (S)-3-Methanesulfonyl-pyrrolidine-1-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 134 (S)-3-Fluoro-pyrrolidine-1-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 135 (S)-3-Cyano-pyrrolidine-1-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 136 (R)-3-Dimethylaminomethyl-pyrrolidine-1-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 137 5-Methyl-isoxazole-4-carboxylic acid (4-methoxy-7-morpholin-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide

- 138 1-(2-Methoxy-ethyl)-1H-[1,2,3]triazole-4-carboxylic acid (4-methoxy-7-morpholin-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 139 1-Methyl-1H-[1,2,3]triazole-4-carboxylic acid (4-methoxy-7-morpholin-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 140 Pyridine-2,5-dicarboxylic acid 2-dimethylamide 5-([4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide)
- 141 Pyridine-2,5-dicarboxylic acid 2-dimethylamide 5-([4-methoxy-7-morpholin-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 142 1-Methyl-1H-pyrazole-4-carboxylic acid [7-methoxy-4-(tetrahydro-pyran-4-yl)-1H-benzoimidazol-2-yl]-amide
- 143 5-Methyl-isoxazole-4-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 144 1-(2-Methoxy-ethyl)-1H-[1,2,3]triazole-4-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 145 1-Methyl-1H-[1,2,3]triazole-4-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 146 1-Cyano-cyclopropanecarboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 147 2-Methyl-oxazole-5-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 148 2-Methyl-thiazole-5-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 149 (S)-3-Methanesulfonyl-pyrrolidine-1-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 150 (S)-3-Fluoro-pyrrolidine-1-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 151 (S)-3-Cyano-pyrrolidine-1-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 152 (R)-3-Dimethylaminomethyl-pyrrolidine-1-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 153 1-[4-Methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-3-thiazol-2-ylmethyl-urea
- 154 1-(2-Methoxy-ethyl)-1H-pyrazole-4-carboxylic acid [4-methoxy-7-(2-oxa-7-aza-spiro[4.4]non-7-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 155 1-Methyl-1H-pyrazole-4-carboxylic acid {4-methoxy-7-[(2-methoxy-propyl)-methyl-amino]-thiazolo[4,5-c]pyridin-2-yl]-amide
- 156 1-Methyl-1H-pyrazole-4-carboxylic acid [4-methoxy-7-(5-oxa-2-aza-spiro[3.4]oct-2-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 157 1-Methyl-1H-pyrazole-4-carboxylic acid (4-methoxy-7-piperidin-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 158 1-Methyl-1H-pyrazole-4-carboxylic acid (4-methoxy-7-piperidin-3-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 159 1-Methyl-1H-pyrazole-4-carboxylic acid [7-(carbamoylmethyl-methyl-amino)-4-methoxy-thiazolo[4,5-c]pyridin-2-yl]-amide
- 160 1-Methyl-1H-pyrazole-4-carboxylic acid [4-methoxy-7-(2,2,2-trifluoro-ethoxy)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 161 4-Hydroxy-4-methyl-piperidine-1-carboxylic acid (4-fluoromethoxy-7-morpholin-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 162 4-Hydroxy-4-methyl-piperidine-1-carboxylic acid (4-difluoromethoxy-7-morpholin-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 163 N-(4-methoxy-6-methyl-7-morpholino-thiazolo[4,5-c]pyridin-2-yl)-1-methyl-pyrazole-4-carboxamide
- 164 N-(6-bromo-4-methoxy-7-morpholino-thiazolo[4,5-c]pyridin-2-yl)-1-methyl-pyrazole-4-carboxamide
- 165 1-Methyl-1H-pyrazole-4-carboxylic acid (6-fluoro-4-methoxy-7-morpholin-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 166 N-(6-chloro-4-methoxy-7-morpholino-thiazolo[4,5-c]pyridin-2-yl)-1-methyl-pyrazole-4-carboxamide
- 167 Cyclopropanecarboxylic acid (4-methoxy-7-morpholin-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 168 (S)-7-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid (4-methoxy-7-morpholin-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 169 (R)-7-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid (4-methoxy-7-morpholin-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 170 N-(4-Methoxy-7-morpholin-4-yl)-thiazolo[4,5-c]pyridin-2-yl)-4-(2-oxo-pyrrolidin-1-ylmethyl)-benzamide
- 171 1-Methyl-1H-pyrazole-4-carboxylic acid (4-methoxy-7-morpholin-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 172 (R)-2-Oxa-7-aza-spiro[4.4]nonane-7-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 173 (S)-2-Oxa-7-aza-spiro[4.4]nonane-7-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 174 3-cyano-N-[4-methoxy-7-(morpholin-4-yl)-[1,3]thiazolo[4,5-c]pyridin-2-yl]propanamide

-continued

175 1-cyano-N-[4-methoxy-7-(morpholin-4-yl)-[1,3]thiazolo[4,5-c]pyridin-2-yl]cyclopropane-1-carboxamide

176 (S)-7-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide

177 (R)-7-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide

178 1-Methyl-1H-pyrazole-4-carboxylic acid (6-bromo-4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide

179 Cyclopropanecarboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide

180 N-[4-Methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-4-(2-oxo-pyrrolidin-1-ylmethyl)-benzamide

181 2-Methyl-thiazole-5-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide

182 2-Oxa-7-aza-spiro[4.4]nonane-7-carboxylic acid [7-(3-ethoxy-3-methyl-azetidin-1-yl)-4-methoxy-thiazolo[4,5-c]pyridin-2-yl]-amide

183 2-Oxa-7-aza-spiro[4.4]nonane-7-carboxylic acid (4-methoxy-7-piperidin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide

184 2-Oxa-7-aza-spiro[4.4]nonane-7-carboxylic acid [4-methoxy-7-(5-oxa-2-aza-spiro[3.4]oct-2-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide

185 2-Oxa-7-aza-spiro[4.4]nonane-7-carboxylic acid [4-methoxy-7-(3-methoxy-3-methyl-azetidin-1-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide

186 2-Methyl-oxazole-5-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

187 Cyclopropanecarboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

188 1-Methyl-1H-pyrazole-4-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

189 N-[4-methoxy-7-(morpholin-4-yl)-[1,3]thiazolo[4,5-c]pyridin-2-yl]-2-methyl-1,3-oxazole-5-carboxamide

190 2-Methyl-thiazole-5-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

191 8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid [4-methoxy-7-(5-oxa-2-aza-spiro[3.4]oct-2-yl)-thiazolo[4,5-c]pyridin-2-yl]amide

192 8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid (4-methoxy-7-piperidin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide

193 8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid [7-(4-fluoro-phenyl)-4-methoxy-thiazolo[4,5-c]pyridin-2-yl]-amide

194 8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid [7-(3-fluoro-phenyl)-4-methoxy-thiazolo[4,5-c]pyridin-2-yl]-amide

195 8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid [7-(2-fluoro-phenyl)-4-methoxy-thiazolo[4,5-c]pyridin-2-yl]-amide

196 1-Dimethylaminomethyl-cyclopropanecarboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

197 Cyclopropane-1,1-dicarboxylic acid dimethylamide (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

198 1-Imidazol-1-ylmethyl-cyclopropanecarboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

199 1-Methyl-1H-pyrazole-4-carboxylic acid (4-hydroxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide

200 8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid (4-methoxy-7-piperidin-3-yl-thiazolo[4,5-c]pyridin-2-yl)-amide

201 1-amino-N-[4-methoxy-7-phenyl-[1,3]thiazolo[4,5-c]pyridin-2-yl]-8-azaspiro[4.5]decane-8-carboxamide

202 (1S,2S)-2-methoxy-N-[4-methoxy-7-phenyl-[1,3]thiazolo[4,5-c]pyridin-2-yl]cyclopropane-1-carboxamide

203 1-Amino-cyclopropanecarboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

204 (S)-7-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

205 (R)-7-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

206 8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid (4-methoxy-7-pyridin-3-yl-thiazolo[4,5-c]pyridin-2-yl)-amide

207 (S)-2-Oxa-7-aza-spiro[4.4]nonane-7-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

208 (R)-2-Oxa-7-aza-spiro[4.4]nonane-7-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

209 8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid [7-(2,3-difluoro-phenyl)-4-methoxy-thiazolo[4,5-c]pyridin-2-yl]-amide

210 8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid [7-(2,5-difluoro-phenyl)-4-methoxy-thiazolo[4,5-c]pyridin-2-yl]-amide

211 8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid (4-methoxy-7-pyridin-2-yl-thiazolo[4,5-c]pyridin-2-yl)-amide

212 8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid (4-methoxy-7-pyridin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide

-continued

213 4-(2,5-Dioxo-pyrrolidin-1-yl)-N-(4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-benzamide

214 4-(2,5-Dioxo-pyrrolidin-1-yl)-N-(4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-benzamide

215 (R)-1-Amino-8-aza-spiro[4.5]decane-8-carboxylic acid(4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

216 (S)-1-Amino-8-aza-spiro[4.5]decane-8-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

217 1-Methyl-1H-pyrazole-4-carboxylic acid (6-cyano-4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide

218 N-(4-Methoxy-7-pyridin-3-yl-thiazolo[4,5-c]pyridin-2-yl)-N',N'-dimethyl-terephthalamide

219 N-(4-Methoxy-7-pyridin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-N',N'-dimethyl-terephthalamide

220 8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid (6-chloro-4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

221 1-Methyl-1H-pyrazole-4-carboxylic acid (6-chloro-4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide

222 8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid (6-cyano-4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

223 8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid (4-methoxy-6-methyl-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

224 N-{4-Methoxy-7-[3-(2-methoxy-ethoxy)-phenyl]-thiazolo[4,5-c]pyridin-2-yl}-N',N'-dimethyl-terephthalamide

225 8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid (4-methoxy-7-[3-(2-methoxy-ethoxy)-phenyl]-thiazolo[4,5-c]pyridin-2-yl)-amide

226 4-(2,5-Dioxo-pyrrolidin-1-yl)-N-(6-fluoro-4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-benzamide

227 N-(2-Hydroxy-ethyl)-N'-(4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-N-methyl-terephthalamide

228 N-(2-Hydroxy-ethyl)-N'-(4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-terephthalamide

229 8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid (6-fluoro-4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

230 N-(2-Dimethylamino-ethyl)-N'-(4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-terephthalamide

231 8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid (4,6-dimethyl-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

232 8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide

233 N-{4-methoxy-7-[3-(2-methoxyethoxy)phenyl]thiazolo[4,5-c]pyridin-2-yl}-1-(2-methoxyethyl)-1H-pyrazole-4-carboxamide

234 N-{6-fluoro-4-methoxy-7-[3-(2-methoxyethoxy)phenyl]-[1,3]thiazolo[4,5-c]pyridin-2-yl}-8-oxa-2-azaspiro[4.5]decane-2-carboxamide

235 N-[4-methoxy-7-(oxan-4-yl)-[1,3]thiazolo[4,5-c]pyridin-2-yl]-1-methyl-1H-pyrazole-4-carboxamide

236 N-[4-methoxy-7-(2-methoxyphenyl)-[1,3]thiazolo[4,5-c]pyridin-2-yl]-8-oxa-2-azaspiro[4.5]decane-2-carboxamide

237 8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid [7-(3,6-dihydro-2H-pyran-4-yl)-4-methoxy-thiazolo[4,5-c]pyridin-2-yl]-amide

238 N-[7-(3,6-dihydro-2H-pyran-4-yl)-4-methoxy-[1,3]thiazolo[4,5-c]pyridin-2-yl]-1-methyl-1H-pyrazole-4-carboxamide

239 N-[7-(2,6-dimethoxy-pyridin-3-yl)-4-methoxy-[1,3]thiazolo[4,5-c]pyridin-2-yl]-8-oxa-2-azaspiro[4.5]decane-2-carboxamide

240 8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid (6-fluoro-4-hydroxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

241 2-amino-N-[4-methoxy-7-(oxan-4-yl)-[1,3]thiazolo[4,5-c]pyridin-2-yl]-1,3-thiazole-5-carboxamide

242 3-Amino-pyrrolidine-1-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

243 (R)-3-Amino-pyrrolidine-1-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

244 (S)-3-Amino-pyrrolidine-1-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

245 8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid (7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

246 1-(2-Methoxy-ethyl)-1H-pyrazole-4-carboxylic acid (6-fluoro-4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide

247 Bicyclo[1.1.1]pentane-1,3-dicarboxylic acid dimethylamide (4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide

248 N-(2-Hydroxy-ethyl)-N'-(4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl)-N-methyl-terephthalamide

and physiologically acceptable salts, derivatives, solvates, prodrugs and stereoisomers thereof, including mixtures thereof in all ratios.

All above-mentioned preferred, particularly preferred and very particularly preferred meanings of the above radicals of the compounds of the formula I should be understood in such a way that these preferred particularly preferred and very particularly preferred meanings or embodiments can be combined with one another in any possible combination to give compounds of the formula I and preferred, particularly preferred and very particularly preferred compounds of the formula I of this type are likewise explicitly disclosed hereby.

Hal denotes fluorine, chlorine, bromine or iodine, in particular fluorine, bromine or chlorine.

D or ^2H denotes deuterium.

—(C=O)— or =O denotes carbonyl oxygen and stands for



or oxygen atom bonded to a carbon atom by means of a double bond.

Alkyl is a saturated unbranched (linear) or branched hydrocarbon chain and has 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 C atoms. Alkyl preferably denotes alkenyl methyl, furthermore ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl or tert-butyl, furthermore also pentyl, 1-, 2- or 3-methylbutyl, 1,1-, 1,2- or 2,2-dimethylpropyl, 1-ethylpropyl, hexyl, 1-, 2-, 3- or 4-methylpentyl, 1,1-, 1,2-, 1,3-, 2,2-, 2,3- or 3,3-dimethylbutyl, 1- or 2-ethylbutyl, 1-ethyl-1-methylpropyl, 1-ethyl-2-methylpropyl, 1,1,2- or 1,2,2-trimethylpropyl, linear or branched heptyl, octyl, nonyl or decyl, further preferably, for example, trifluoromethyl.

Cyclic alkyl or cycloalkyl is a saturated cyclic hydrocarbon chain and has 3-10, preferably 3-7 C atoms and preferably denotes cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl or cycloheptyl. Cycloalkyl also denotes a partially unsaturated cyclic alkyl, such as, for example, cyclohexenyl or cyclohexynyl.

Alkenyl denotes an unsaturated unbranched (linear) or branched hydrocarbon chain and has 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 C atoms.

O-alkyl or OA denotes linear or branched alkoxy having 1-6 C atoms, and is preferably methoxyl, furthermore also e.g. ethoxyl, n-propoxyl, isopropoxyl, n-butoxyl, isobutoxyl, sec-butoxyl or tert-butoxyl.

Alkyloxycarbonyl refers to straight or branched chain esters of a carboxylic acid derivative of the present invention, i.e. methyloxycarbonyl (MeOCO—), ethyloxycarbonyl, or butyloxycarbonyl.

Alkylcarbonyl refers to straight or branched chain alkyl and a carboxylic acid group.

Aryl, Ar or aromatic ring denotes a mono- or polycyclic aromatic or fully unsaturated cyclic hydrocarbon chain, for example unsubstituted phenyl, naphthyl or biphenyl, furthermore preferably phenyl, naphthyl or biphenyl, each of which is mono-, di- or trisubstituted, for example, by A, fluorine, chlorine, bromine, iodine, hydroxyl, methoxy, ethoxy, propoxy, butoxy, pentyloxy, hexyloxy, nitro, cyano, formyl, acetyl, propionyl, trifluoromethyl, amino, methylamino, ethylamino, dimethylamino, diethylamino, benzyl, sulfonylamino, methylsulfonylamino, ethylsulfonylamino, propylsulfonylamino, butylsulfonylamino, dimethylsulfonylamino, phenylsulfonylamino, carboxyl, methoxycarbonyl, ethoxycarbonyl, aminocarbonyl.

Heterocycle and heterocyclyl refer to saturated or unsaturated non-aromatic rings or ring systems containing at least one heteroatom selected from O, S and N, further including the oxidized forms of sulfur, namely SO and SO_2 . Examples of heterocycles include tetrahydrofuran (THF), dihydrofuran, 1,4-dioxane, morpholine, 1,4-dithiane, piperazine, piperidine, 1,3-dioxolane, imidazolidine, imidazoline, pyrrolidine, pyrrolidine, tetrahydropyran, dihydropyran, oxathiolane, dithiolane, 1,3-dioxane, 1,3-dithiane, oxathiane, thiomorpholine, and the like.

Heteroaryl means an aromatic or partially aromatic heterocycle that contains at least one ring heteroatom selected from O, S and N. Heteroaryls thus includes heteroaryls fused to other kinds of rings, such as aryls, cycloalkyls and heterocycles that are not aromatic. Examples of heteroaryl groups include: pyrrolyl, isoxazolyl, isothiazolyl, pyrazolyl, pyridyl, oxazolyl, oxadiazolyl, thiadiazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, furanyl, triazinyl, thienyl, pyrimidyl, benzisoxazolyl, benzoxazolyl, benzothiazolyl, benzothiadiazolyl, dihydrobenzofuranyl, indolinyl, pyridazinyl, indazolyl, isoxazolyl, isoindolyl, dihydrobenzothienyl, indolizynyl, cinnolinyl, phthalazinyl, quinazolinyl, naphthyridinyl, carbazolyl, benzodioxinyl, benzodioxolyl, quinoxalinyl, purinyl, furazanyl, thiophenyl, isobenzylfuranlyl, benzimidazolyl, benzofuranlyl, benzothienyl, quinolyl, indolyl, isoquinolyl, dibenzofuranlyl, and the like. For heterocyclyl and heteroaryl groups, rings and ring systems containing from 3-15 atoms are included, forming 1-3 rings.

Mono- or bicyclic saturated, unsaturated or aromatic heterocycle preferably denotes unsubstituted or mono-, di- or trisubstituted 2- or 3-furyl, 2- or 3-thienyl, 1-, 2- or 3-pyrrolyl, 1-, 2-, 4- or 5-imidazolyl, 1-, 3-, 4- or 5-pyrazolyl, 2-, 4- or 5-oxazolyl, 3-, 4- or 5-isoxazolyl, 2-, 4- or 5-thiazolyl, 3-, 4- or 5-isothiazolyl, 2-, 3- or 4-pyridyl, 2-, 4-, 5- or 6-pyrimidinyl, furthermore preferably 1,2,3-triazol-1-, -4- or -5-yl, 1,2,4-triazol-1-, -3- or 5-yl, 1- or 5-tetrazolyl, 1,2,3-oxadiazol-4- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,2,4-thiadiazol-3- or -5-yl, 1,2,3-thiadiazol-4- or -5-yl, 3- or 4-pyridazinyl, pyrazinyl, 1-, 2-, 3-, 4-, 5-, 6- or 7-indolyl, 4- or 5-isoindolyl, 1-, 2-, 4- or 5-benzimidazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzopyrazolyl, 2-, 4-, 5-, 6- or 7-benzoxazolyl, 3-, 4-, 5-, 6- or 7-benzisoxazolyl, 2-, 4-, 5-, 6- or 7-benzothiazolyl, 2-, 4-, 5-, 6- or 7-benzisothiazolyl, 4-, 5-, 6- or 7-benz-2,1,3-oxadiazolyl, 2-, 3-, 4-, 5-, 6-, 7- or 8-quinolyl, 1-, 3-, 4-, 5-, 6-, 7- or 8-isoquinolyl, 3-, 4-, 5-, 6-, 7- or 8-cinnolinyl, 2-, 4-, 5-, 6-, 7- or 8-quinazolinyl, 5- or 6-quinoxalinyl, 2-, 3-, 5-, 6-, 7- or 8-2H-benzo-1,4-oxazinyl, further preferably 1,3-benzodioxol-5-yl, 1,4-benzodioxan-6-yl, 2,1,3-benzothiadiazol-4- or -5-yl or 2,1,3-benzoxadiazol-5-yl.


The heterocyclic radicals may also be partially or fully hydrogenated and also denote, for example, 2,3-dihydro-2-, -3-, -4- or -5-furyl, 2,5-dihydro-2-, -3-, -4- or 5-furyl, tetrahydro-2- or -3-furyl, 1,3-dioxolan-4-yl, tetrahydro-2- or -3-thienyl, 2,3-dihydro-1-, -2-, -3-, -4- or -5-pyrrolyl, 2,5-dihydro-1-, -2-, -3-, -4- or -5-pyrrolyl, 1-, 2- or 3-pyrrolidinyl, tetrahydro-1-, -2- or -4-imidazolyl, 2,3-dihydro-1-, -2-, -3-, -4- or -5-pyrazolyl, tetrahydro-1-, -3- or -4-pyrazolyl, 1,4-dihydro-1-, -2-, -3- or -4-pyridyl, 1,2,3,4-tetrahydro-1-, -2-, -3-, -4-, -5- or -6-pyridyl, 1-, 2-, 3- or 4-piperidinyl, 2-, 3- or 4-morpholinyl, tetrahydro-2-, -3- or -4-pyranyl, 1,4-dioxanyl, 1,3-dioxan-2-, -4- or -5-yl, hexahydro-1-, -3- or -4-pyridazinyl, hexahydro-1-, -2-, -4- or -5-pyrimidinyl, 1-, 2- or 3-piperazinyl, 1,2,3,4-tetrahydro-1-, -2-, -3-, -4-, -5-, -6-, -7- or -8-quinolyl, 1,2,3,4-tetrahydro-1-, -2-, -3-, -4-, -5-, -6-, -7- or -8-isoquinolyl, 2-, 3-, 5-, 6-, 7- or 8-3,4-

dihydro-2H-benzo-1,4-oxazinyl, further preferably 2,3-methylenedioxyphenyl, 3,4-methylenedioxyphenyl, 2,3-ethylenedioxyphenyl, 3,4-ethylenedioxyphenyl, 3,4-(difluoromethylenedioxy)phenyl, 2,3-dihydrobenzofuran-5- or 6-yl, 2,3-(2-oxomethylenedioxy)phenyl or also 3,4-dihydro-2H-1,5-benzodioxepin-6- or -7-yl, furthermore preferably 2,3-dihydrobenzofuranyl or 2,3-dihydro-2-oxofuranyl.

Heterocycle furthermore denotes, for example, 2-oxopiperidin-1-yl, 2-oxopyrrolidin-1-yl, 2-oxo-1H-pyridin-1-yl, 3-oxomorpholin-4-yl, 4-oxo-1H-pyridin-1-yl, 2,6-dioxopiperidin-1-yl, 2-oxopiperazin-1-yl, 2,6-dioxopiperazin-1-yl, 2,5-dioxopyrrolidin-1-yl, 2-oxo-1,3-oxazolidin-3-yl, 3-oxo-2H-pyridazin-2-yl, 2-caprolactam-1-yl(=2-oxoazepan-1-yl), 2-hydroxy-6-oxopiperazin-1-yl, 2-methoxy-6-oxopiperazin-1-yl or 2-azabicyclo[2.2.2]octan-3-on-2-yl.

Heterocycloalkyl here denotes a fully hydrogenated or saturated heterocycle, heterocycloalkenyl (one or more double bonds) or heterocycloalkynyl (one or more triple bonds) denotes a partially or incompletely hydrogenated or unsaturated heterocycle, heteroaryl denotes an aromatic or fully unsaturated heterocycle.

A cyclic alkylaryl group in connection with the present invention means that and one or two aromatic rings Ar are condensed onto an unsubstituted or a mono- or disubstituted cyclic alkyl, in which one or two CH₂ groups and/or, in addition, 1-11H atoms may be replaced, such as, for example, in the radicals depicted below:



Furthermore, the abbreviations below have the following meanings:

Boc ter-butoxycarbonyl
 CBZ benzylloxycarbonyl
 DNP 2,4-dinitrophenyl
 Fmoc 9-fluorenylmethoxycarbonyl
 imi-DNP 2,4-dinitrophenyl in the 1-position of the imidazole ring
 OMe methyl ester
 POA phenoxyacetyl
 DCCldicyclohexylcarbodiimide
 HOBt1-hydroxybenzotriazole

The invention therefore relates to a pharmaceutical preparation comprising the compound according to the present invention and/or one of its physiologically acceptable salts, derivatives, solvates, prodrugs and stereoisomers, including mixtures thereof in all ratios.

The invention also relates to a pharmaceutical preparation according to the invention of this type, comprising further excipients and/or adjuvants.

In addition, the invention relates to an above pharmaceutical preparation according to the invention, comprising at least one further medicament active compound.

Pharmaceutically or physiologically acceptable derivatives are taken to mean, for example, salts of the compound of the present invention, and also so-called prodrug compounds. Prodrug compounds are taken to mean derivatives of the compound of the present invention which have been modified by means of, for example, alkyl or acyl groups (see also amino- and hydroxyl-protecting groups below), sugars or oligopeptides and which are rapidly cleaved or liberated in the organism to form the effective molecules. These also include biodegradable polymer derivatives of the compound of the present invention, as described, for example, in Int. J. Pharm. 115 (1995), 61-67.

The compound of the present invention can be used in its final non-salt form. On the other hand, the present invention also encompasses the use of pepstatin in the form of its pharmaceutically acceptable salts, which can be derived from various organic and inorganic bases by procedures known in the art. Pharmaceutically acceptable salt forms of pepstatin are for the most part prepared by conventional methods. If the compound of the present invention contains a carboxyl group, one of its suitable salts can be formed by reacting the compound of the present invention with a suitable base to give the corresponding base-addition salt. Such bases are, for example, alkali metal hydroxides, including potassium hydroxide, sodium hydroxide and lithium hydroxide; alkaline-earth metal hydroxides, such as barium hydroxide and calcium hydroxide; alkali metal alkoxides, for example potassium ethoxide and sodium propoxide; and various organic bases, such as piperidine, diethanolamine and N-methylglutamine. The aluminium salts of pepstatin are likewise included.

Furthermore, the base salts of the compound of the present invention include aluminium, ammonium, calcium, copper, iron(III), iron(II), lithium, magnesium, manganese(III), manganese(II), potassium, sodium and zinc salts, but this is not intended to represent a restriction.

Of the above-mentioned salts, preference is given to ammonium; the alkali metal salts sodium and potassium, and the alkaline-earth metal salts calcium and magnesium. Salts of the compound of the present invention which are derived from pharmaceutically acceptable organic non-toxic bases include salts of primary, secondary and tertiary amines, substituted amines, also including naturally occurring substituted amines, cyclic amines, and basic ion exchanger resins, for example arginine, betaine, caffeine, chlorprocaine, choline, N,N'-dibenzylethylenediamine (benzathine), dicyclohexylamine, diethanolamine, diethylamine, 2-diethylamino-ethanol, 2-dimethylaminoethanol, ethanolamine, ethylenediamine, N-ethylmorpholine, N-ethylpiperidine, glucamine, glucosamine, histidine, hydrabamine, isopropylamine, lidocaine, lysine, meglumine, N-methyl-D-glucamine, morpholine, piperazine, piperidine, polyamine resins, procaine, purines, theobromine, triethanolamine, triethylamine, trimethylamine, tripropylamine and tris-(hydroxymethyl)methylamine (tromethamine), but this is not intended to represent a restriction.

As mentioned, the pharmaceutically acceptable base-addition salts of pepstatin are formed with metals or amines, such as alkali metals and alkaline-earth metals or organic amines. Preferred metals are sodium, potassium, magnesium and calcium. Preferred organic amines are N,N'-dibenzyl-

ethylenediamine, chlorprocaine, choline, diethanolamine, ethylenediamine, N-methyl-D-glucamine and procaine.

The base-addition salts of the compound of the present invention are prepared by bringing the free acid form into contact with a sufficient amount of the desired base, causing the formation of the salt in a conventional manner. The free acid can be regenerated by bringing the salt form into contact with an acid and isolating the free acid in a conventional manner. The free acid forms differ in a certain respect from the corresponding salt forms thereof with respect to certain physical properties, such as solubility in polar solvents; for the purposes of the invention, however, the salts otherwise correspond to the respective free acid forms thereof.

In view of that stated above, it can be seen that the term "pharmaceutically acceptable salt" in the present connection is taken to mean an active compound which comprises the compound of the present invention in the form of one of its salts, in particular if this salt form imparts improved pharmacokinetic properties on the active compound compared with the free form of the active compound or any other salt form of the active compound used earlier. The pharmaceutically acceptable salt form of the active compound can also provide this active compound for the first time with a desired pharmacokinetic property which it did not have earlier and can even have a positive influence on the pharmacodynamics of this active compound with respect to its therapeutic efficacy in the body.

Solvates of the compound of the present invention are taken to mean adductions of inert solvent molecules pepsatin which form owing to their mutual attractive force. Solvates are, for example, hydrates, such as monohydrates or dihydrates, or alcoholates, i.e. addition compounds with alcohols, such as, for example, with methanol or ethanol.

All physiologically acceptable salts, derivatives, solvates and stereoisomers of these compounds, including mixtures thereof in all ratios, are also in accordance with the invention.

Compounds of the general formula I may contain one or more centres of chirality, so that all stereoisomers, enantiomers, diastereomers, etc., of the compounds of the general formula I are also claimed in the present invention.

The invention also relates to the optically active forms (stereoisomers), the enantiomers, the racemates, the diastereomers and hydrates and solvates of these compounds.

Compounds of the formula I according to the invention may be chiral owing to their molecular structure and may accordingly occur in various enantiomeric forms. They may therefore be in racemic or optically active form. Since the pharmaceutical efficacy of the racemates or stereoisomers of the compounds according to the invention may differ, it may be desirable to use the enantiomers. In these cases, the end product, but also even the intermediates, may be separated into enantiomeric compounds by chemical or physical measures known to the person skilled in the art or already employed as such in the synthesis.

Pharmaceutically or physiologically acceptable derivatives are taken to mean, for example, salts of the compounds according to the invention and also so-called prodrug compounds. Prodrug compounds are taken to mean compounds of the formula I which have been modified with, for example, alkyl or acyl groups (see also amino- and hydroxyl-protecting groups below), sugars or oligopeptides and which are rapidly cleaved or liberated in the organism to form the effective compounds according to the invention. These also include biodegradable polymer derivatives of the

compounds according to the invention, as described, for example, in Int. J. Pharm. 115 (1995), 61-67.

Suitable acid-addition salts are inorganic or organic salts of all physiologically or pharmacologically acceptable acids, for example halides, in particular hydrochlorides or hydrobromides, lactates, sulfates, citrates, tartrates, maleates, fumarates, oxalates, acetates, phosphates, methylsulfonates or p-toluenesulfonates.

Very particular preference is given to the hydrochlorides, the trifluoroacetates or the bistrifluoroacetates of the compounds according to the invention.

Solvates of the compounds of the formula I are taken to mean adductions of inert solvent molecules onto the compounds of the formula I which form owing to their mutual attractive force. Solvates are, for example, hydrates, such as monohydrates or dihydrates, or alcoholates, i.e. addition compounds with alcohols, such as, for example, with methanol or ethanol.

It is furthermore intended that a compound of the formula I includes isotope-labelled forms thereof. An isotope-labelled form of a compound of the formula I is identical to this compound apart from the fact that one or more atoms of the compound have been replaced by an atom or atoms having an atomic mass or mass number which differs from the atomic mass or mass number of the atom which usually occurs naturally. Examples of isotopes which are readily commercially available and which can be incorporated into a compound of the formula I by well-known methods include isotopes of hydrogen, carbon, nitrogen, oxygen, phosphorus, fluorine and chlorine, for example ^2H , ^3H , ^{13}C , ^{14}C , ^{15}N , ^{18}O , ^{17}O , ^{31}P , ^{32}P , ^{35}S , ^{18}F and ^{36}Cl , respectively.

A compound of the formula I, a prodrug thereof or a pharmaceutically acceptable salt of either which contains one or more of the above-mentioned isotopes and/or other isotopes of other atoms is intended to be part of the present invention. An isotope-labelled compound of the formula I can be used in a number of beneficial ways. For example, an isotope-labelled compound of the formula I into which, for example, a radioisotope, such as ^3H or ^{14}C , has been incorporated is suitable for medicament and/or substrate tissue distribution assays. These radio-isotopes, i.e. tritium (^3H) and carbon-14 (^{14}C), are particularly preferred owing to their simple preparation and excellent detectability. Incorporation of heavier isotopes, for example deuterium (^2H), into a compound of the formula I has therapeutic advantages owing to the higher metabolic stability of this isotope-labelled compound. Higher metabolic stability translates directly into an increased in-vivo half-life or lower dosages, which under most circumstances would represent a preferred embodiment of the present invention. An isotope-labelled compound of the formula I can usually be prepared by carrying out the procedures disclosed in the synthesis schemes and the related description, in the example part and in the preparation part in the present text, replacing a non-isotope-labelled reactant with a readily available isotope-labelled reactant.



In order to manipulate the oxidative metabolism of the compound by way of the primary kinetic isotope effect, deuterium (^2H) can also be incorporated into a compound of the formula I. The primary kinetic isotope effect is a change in the rate of a chemical reaction that results from exchange of isotopic nuclei, which in turn is caused by the change in ground state energies necessary for covalent bond formation after this isotopic exchange. Exchange of a heavier isotope usually results in a lowering of the ground state energy for a chemical bond and thus causes a reduction in the rate in rate-limiting bond breakage. If the bond breakage occurs in

29

or in the vicinity of a saddle-point region along the coordinate of a multi-product reaction, the product distribution ratios can be altered substantially. For explanation: if deuterium is bonded to a carbon atom in a non-exchangeable position, rate differences of $k_M/k_D=2-7$ are typical. If this rate difference is successfully applied to a compound of the formula I that is susceptible to oxidation, the profile of this compound in vivo can thereby be drastically modified and result in improved pharmacokinetic properties.

When discovering and developing therapeutic agents, the person skilled in the art attempts to optimise pharmacokinetic parameters while retaining desirable in-vitro properties. It is reasonable to assume that many compounds with poor pharmacokinetic profiles are susceptible to oxidative metabolism. In-vitro liver microsomal assays currently available provide valuable information on the course of oxidative metabolism of this type, which in turn permits the rational design of deuterated compounds of the formula I with improved stability through resistance to such oxidative metabolism. Significant improvements in the pharmacokinetic profiles of the compounds of the formula I are thereby obtained and can be expressed quantitatively in terms of increases in the in-vivo half-life ($T/2$), concentration at maximum therapeutic effect (C_{max}), area under the dose response curve (AUC), and F; and in terms of reduced clearance, dose and costs of materials.

The following is intended to illustrate the above: a compound of the formula I which has multiple potential sites of attack for oxidative metabolism, for example benzylic hydrogen atoms and hydrogen atoms bonded to a nitrogen atom, is prepared as a series of analogues in which various combinations of hydrogen atoms are replaced by deuterium atoms, so that some, most or all of these hydrogen atoms have been replaced by deuterium atoms. Half-life determinations enable favourable and accurate determination of the extent to which the improvement in resistance to oxidative metabolism has improved. In this way, it is determined that the half-life of the parent compound can be extended by up to 100% as the result of deuterium-hydrogen exchange of this type.



30

The replacement of hydrogen by deuterium in a compound of the formula I can also be used to achieve a favourable modification of the metabolite spectrum of the starting compound in order to diminish or eliminate undesired toxic metabolites. For example, if a toxic metabolite arises through oxidative carbon-hydrogen (C—H) bond cleavage, it can reasonably be assumed that the deuterated analogue will greatly diminish or eliminate production of the undesired metabolite, even if the particular oxidation is not a rate-determining step. Further information on the state of the art with respect to deuterium-hydrogen exchange is given, for example in Hanzlik et al., J. Org. Chem. 55, 3992-3997, 1990, Reider et al., J. Org. Chem. 52, 3326-3334, 1987, Foster, Adv. Drug Res. 14, 1-40, 1985, Gillette et al., Biochemistry 33(10), 2927-2937, 1994, and Jarman et al., Carcinogenesis 16(4), 683-688, 1993.


The invention also relates to mixtures of the compounds of the formula I according to the invention, for example mixtures of two diastereomers, for example in the ratio 1:1, 1:2, 1:3, 1:4, 1:5, 1:10, 1:100 or 1:1000. These are particularly preferably mixtures of two stereoisomeric compounds. However, preference is also given to mixtures of two or more compounds of the formula I.

In addition, the invention relates to a process for the preparation of the compounds of the formula I, characterized in that

- a) a compound of the formula II undergoes a reduction to give a compound of formula III, a compound of formula III is reacted with a compound of formula IV at elevated temperature to give a compound of formula V, a compound of formula V is converted to a compound of the formula VI employing the use of catalyst and base, a compound of formula VI is converted to a compound of the formula VII by bromination, a compound of the formula VII is converted to a compound of the formula VIII under essentially basic conditions and a compound of the formula VIII is reacted with a compound of the formula IX under standard amidation or carbamide formation conditions to give a compound of the formula I,


31

32




b) a compound of the formula V is reacted with a compound of the formula X under Suzuki-type reaction conditions to give a compound of the formula VI, a compound of formula VI is converted to a compound of the formula VII by bromination, a compound of formula VII is converted to a compound of the formula VIII under essentially basic conditions and a compound of the formula VIII is reacted with a compound of the formula IX under standard amidation or carbamide formation conditions to give a compound of the formula I,

c) a compound of the formula XII is iodinated to give a compound of the formula XIII, a compound of formula XIII is converted to a compound of the formula XIV by treatment with base and an electrophile, a compound of formula XIV is converted to a compound of the formula XV by reduction, a compound of formula XV is reacted with a compound of formula IV at elevated temperature to give a compound of the formula XVI, a compound of formula XVI is converted under catalytic conditions to a compound of the formula XVII, a compound of the formula XVII is converted to a compound of the



33

formula VIII under basic conditions and a compound of the formula IX under standard amidation or carbamide formation conditions to give a compound of the formula I,



34

enolysis are those which contain correspondingly protected amino, carboxyl and/or hydroxyl groups instead of one or more free amino, carboxyl and/or hydroxyl groups, preferably those which carry an amino-protecting group instead of an H atom which is connected to an N atom. Preference is

- d) the base of a compound of the formula I is converted into one of its salts by treatment with an acid, or
 e) an acid of a compound of the formula I is converted into one of its salts by treatment with a base.

It is also possible to carry out the reactions stepwise in each case and to modify the sequence of the linking reactions of the building blocks with adaptation of the protecting-group concept.

The starting materials or starting compounds are generally known. If they are novel, they can be prepared by methods known per se.

If desired, the starting materials can also be formed in situ by not isolating them from the reaction mixture, but instead immediately converting them further into the compounds of the formula I.

The compounds of the formula I are preferably obtained by liberating them from their functional derivatives by solvolysis, in particular by hydrolysis, or by hydrogenolysis.

furthermore given to starting materials which carry a hydroxyl-protecting group instead of the H atom of a hydroxyl group. Preference is also given to starting materials which carry a protected carboxyl group instead of a free carboxyl group. It is also possible for a plurality of identical or different protected amino, carboxyl and/or hydroxyl groups to be present in the molecule of the starting material. If the protecting groups present are different from one another, they can in many cases be cleaved off selectively.

The term "amino-protecting group" is generally known and relates to groups which are suitable for protecting (blocking) an amino group against chemical reactions, but which can easily be removed after the desired chemical reaction has been carried out elsewhere in the molecule. Typical of such groups are, in particular, unsubstituted or substituted acyl groups, furthermore unsubstituted or substituted aryl (for example 2,4-dinitrophenyl) or aralkyl groups (for example benzyl, 4-nitrobenzyl, triphenylmethyl). Since the amino-protecting groups are removed after

the desired reaction or reaction sequence, their type and size is, in addition, not crucial, but preference is given to those having 1-20, in particular 1-8, C atoms. The term "acyl group" is to be understood in the broadest sense in connection with the present process. It encompasses acyl groups derived from aliphatic, araliphatic, aromatic or heterocyclic carboxylic acids or sulfonic acids and, in particular, alkoxy-carbonyl, aryloxy-carbonyl and especially aralkoxy-carbonyl groups. Examples of such acyl groups are alkanoyl, such as acetyl, propionyl, butyryl, aralkanoyl, such as phenylacetyl, aroyl, such as benzoyl or toluyl, aryoxylkanoyl, such as phenoxyacetyl, alkoxy-carbonyl, such as methoxycarbonyl, ethoxycarbonyl, 2,2,2-trichloroethoxycarbonyl, BOC, 2-iodoethoxycarbonyl, aralkoxy-carbonyl, such as CBZ, 4-methoxybenzyloxy-carbonyl or Fmoc. Preferred acyl groups are CBZ, Fmoc, benzyl and acetyl.

The term "acid-protecting group" or "carboxyl-protecting group" is likewise generally known and relates to groups which are suitable for protecting a —COOH group against chemical reactions, but which can easily be removed after the desired chemical reaction has been carried out elsewhere in the molecule. The use of esters instead of the free acids, for example of substituted and unsubstituted alkyl esters (such as methyl, ethyl, tert-butyl and substituted derivatives thereof), of substituted and unsubstituted benzyl esters or silyl esters, is typical. The type and size of the acid-protecting groups is not crucial, but preference is given to those having 1-20, in particular 1-10, C atoms.

The term "hydroxyl-protecting group" is likewise generally known and relates to groups which are suitable for protecting a hydroxyl group against chemical reactions, but which can easily be removed after the desired chemical reaction has been carried out elsewhere in the molecule. Typical of such groups are the above-mentioned unsubstituted or substituted aryl, aralkyl or acyl groups, furthermore also alkyl groups. Their type and size of the hydroxyl-protecting groups is not crucial, but preference is given to those having 1-20, in particular 1-10, C atoms. Examples of hydroxyl-protecting groups are, inter alia, benzyl, p-nitrobenzoyl, p-toluenesulfonyl and acetyl, where benzyl and acetyl are preferred.

Further typical examples of amino-, acid- and hydroxyl-protecting groups are found, for example, in "Greene's Protective Groups in Organic Synthesis", fourth edition, Wiley-Interscience, 2007.

The functional derivatives of the compounds of the formula I to be used as starting materials can be prepared by known methods of amino-acid and peptide synthesis, as described, for example, in the said standard works and patent applications.

The compounds of the formula I are liberated from their functional derivatives, depending on the protecting group used, for example, with the aid of strong acids, advantageously using trifluoroacetic acid or perchloric acid, but also using other strong inorganic acids, such as hydrochloric acid or sulfuric acid, strong organic acids, such as trichloroacetic acid, or sulfonic acids, such as benzoyl- or p-toluenesulfonic acid. The presence of an additional inert solvent and/or a catalyst is possible, but is not always necessary.

Depending on the respective synthetic route, the starting materials can optionally be reacted in the presence of an inert solvent.

Suitable inert solvents are, for example, heptane, hexane, petroleum ether, DMSO, benzene, toluene, xylene, trichloroethylene-, 1,2-dichloroethane carbon tetrachloride, chloroform or dichloromethane; alcohols, such as methanol, ethanol, isopropanol, n-propanol, n-butanol or tert-butanol;

ethers, such as diethyl ether, diisopropyl ether (preferably for substitution on the indole nitrogen), tetrahydrofuran (THF) or dioxane; glycol ethers, such as ethylene glycol monomethyl or monoethyl ether, ethylene glycol dimethyl-ether (diglyme); ketones, such as acetone or butanone; amides, such as acetamide, dimethylacetamide, N-methylpyrrolidone (NMP) or dimethylformamide (DMF); nitriles, such as acetonitrile; esters, such as ethyl acetate, carboxylic acids or acid anhydrides, such as, for example, such as acetic acid or acetic anhydride, nitro compounds, such as nitromethane or nitro-benzene, optionally also mixtures of the said solvents with one another or mixtures with water.

The amount of solvent is not crucial; 10 g to 500 g of solvent can preferably be added per g of the compound of the formula I to be reacted.

It may be advantageous to add an acid-binding agent, for example an alkali metal or alkaline-earth metal hydroxide, carbonate or bicarbonate or other alkali or alkaline-earth metal salts of weak acids, preferably a potassium, sodium or calcium salt, or to add an organic base, such as, for example, on triethylamine, dimethylamine, pyridine or quinoline, or an excess of the amine component.

The resultant compounds according to the invention can be separated from the corresponding solution in which they are prepared (for example by centrifugation and washing) and can be stored in another composition after separation, or they can remain directly in the preparation solution. The resultant compounds according to the invention can also be taken up in desired solvents for the particular use.

The reaction duration depends on the reaction conditions selected. In general, the reaction duration is 0.5 hour to 10 days, preferably 1 to 24 hours. On use of a microwave, the reaction time can be reduced to values of 1 to 60 minutes.

The compounds of the formula I and also the starting materials for their preparation are, in addition, prepared by known methods, as described in the literature (for example in standard works, such as Houben-Weyl, Methoden der organischen Chemie [Methods of Organic Chemistry], Georg-Thieme-Verlag, Stuttgart), for example under reaction conditions which are known and suitable for the said reactions. Use can also be made here of variants known per se, which are not described here in greater detail.

Conventional work-up steps, such as, for example, addition of water to the reaction mixture and extraction, enable the compounds to be obtained after removal of the solvent. It may be advantageous, for further purification of the product, to follow this with a distillation or crystallisation or to carry out a chromatographic purification.

An acid of the formula I can be converted into the associated addition salt using a base, for example by reaction of equivalent amounts of the acid and base in an inert solvent, such as ethanol, and inclusive evaporation. Suitable bases for this reaction are, in particular, those which give physiologically acceptable salts. Thus, the acid of the formula I can be converted into the corresponding metal salt, in particular alkali or alkaline-earth metal salt, using a base (for example sodium hydroxide, potassium hydroxide, sodium carbonate or potassium carbonate) or into the corresponding ammonium salt. Organic bases which give physiologically acceptable salts, such as, for example, ethanolamine, are also suitable for this reaction.

On the other hand, a base of the formula I can be converted into the associated acid-addition salt using an acid, for example by reaction of equivalent amounts of the base and acid in an inert solvent, such as ethanol, with subsequent evaporation. Suitable acids for this reaction are,

in particular, those which give physiologically acceptable salts. Thus, it is possible to use inorganic acids, for example sulfuric acid, nitric acid, hydrohalic acids, such as hydrochloric acid or hydrobromic acid, phosphoric acids, such as orthophosphoric acid, sulfamic acid, furthermore organic acids, in particular aliphatic, alicyclic, araliphatic, aromatic or heterocyclic, mono- or polybasic carboxylic, sulfonic or sulfuric acids, for example formic acid, acetic acid, propionic acid, pivalic acid, diethylacetic acid, malonic acid, succinic acid, pimelic acid, fumaric acid, maleic acid, lactic acid, tartaric acid, malic acid, citric acid, gluconic acid, ascorbic acid, nicotinic acid, isonicotinic acid, methane- or ethanesulfonic acid, ethanesulfonic acid, 2-hydroxysulfonic acid, benzenesulfonic acid, p-toluenesulfonic acid, naphthalenemom- and disulfonic acids or laurylsulfuric acid. Salts with physiologically unacceptable acids, for example picrates, can be used for the isolation and/or purification of the compounds of the formula I.

It has been found that the compounds of the formula I are well tolerated and have valuable pharmacological properties.

Since adenosine receptors, such as A_{2A} and A_{2B} , are shown to down-regulate the immune response during inflammation and protect tissues from immune damage, inhibition of signaling through adenosine receptors can be used to intensify and prolong the immune response.

Methods are provided herein to increase an immune response. In one example, the method increases desirable and targeted tissue damage, such as damage of a tumor, for example cancer. Disclosed herein are methods of inhibiting one or more processes conducive to the production of extracellular adenosine and adenosine-triggered signaling through adenosine receptors. For example, enhancement of an immune response, local tissue inflammation, and targeted tissue destruction is accomplished by: inhibiting or reducing the adenosine-producing local tissue hypoxia; by degrading (or rendering inactive) accumulated extracellular adenosine; by preventing or decreasing expression of adenosine receptors on immune cells; and/or by inhibiting/antagonizing signaling by adenosine ligands through adenosine receptors. The results disclosed herein demonstrate that by in vivo administration of agents that disrupt the "hypoxia→adenosine accumulation→immunosuppressive adenosine receptor signaling to immune cells" pathway in subjects suffering from various diseases (e.g. cancer and sepsis) can result in in vivo treatment of tumors or improved immunization.

In one example, the method includes administering one or more inhibitors of extracellular adenosine and/or adenosine receptor inhibitors, such as an adenosine receptor antagonist. To increase the efficacy of a vaccine, one or more adenosine receptor inhibitors and/or inhibitors of extracellular adenosine can be administered in conjunction with the vaccine. In one example, one or more adenosine receptor inhibitors or inhibitors of extracellular adenosine are administered to increase an immune response/inflammation. In another example, a method is provided to achieve targeted tissue damage, such as for tumor destruction.

The invention therefore furthermore relates to the use of compounds according to the invention for the preparation of a medicament for the treatment and/or prophylaxis of diseases which are caused, promoted and/or propagated by adenosine or other A_{2A} and/or A_{2B} receptor agonists.

The invention thus also relates, in particular, to a medicament comprising at least one compound according to the invention and/or one of its physiologically acceptable salts, derivatives, solvates, prodrugs and stereoisomers, including

mixtures thereof in all ratios, for use in the treatment and/or prophylaxis of physiological and/or pathophysiological states.

Particular preference is given, in particular, to physiological and/or pathophysiological states which are connected to adenosine A_{2A} and/or A_{2B} receptors.

Physiological and/or pathophysiological states are taken to mean physiological and/or pathophysiological states which are medically relevant, such as, for example, diseases or illnesses and medical disorders, complaints, symptoms or complications and the like, in particular diseases.

The invention furthermore relates to a medicament comprising at least one compound according to the invention and/or one of its physiologically acceptable salts, derivatives, solvates, prodrugs and stereoisomers, including mixtures thereof in all ratios, for use in the treatment and/or prophylaxis of physiological and/or pathophysiological states selected from the group consisting of hyperproliferative and infectious diseases and disorders.

The invention further relates to a medicament comprising at least one compound according to the invention and/or one of its physiologically acceptable salts, derivatives, solvates, prodrugs and stereoisomers, including mixtures thereof in all ratios, for use in the treatment and/or prophylaxis of physiological and/or pathophysiological states selected from the group consisting of hyperproliferative and infectious diseases and disorders, wherein the hyperproliferative disease or disorder is cancer.

The invention thus particularly preferably relates to a medicament comprising at least one compound according to the invention and/or one of its physiologically acceptable salts, derivatives, solvates, prodrugs and stereoisomers, including mixtures thereof in all ratios, wherein the cancer is selected from the group consisting of acute and chronic lymphocytic leukemia, acute granulocytic leukemia, adrenal cortex cancer, bladder cancer, brain cancer, breast cancer, cervical cancer, cervical hyperplasia, cervical cancer, chorio cancer, chronic granulocytic leukemia, chronic lymphocytic leukemia, colon cancer, endometrial cancer, esophageal cancer, essential thrombocytosis, genitourinary carcinoma, glioma, glioblastoma, hairy cell leukemia, head and neck carcinoma, Hodgkin's disease, Kaposi's sarcoma, lung carcinoma, lymphoma, malignant carcinoid carcinoma, malignant hypercalcemia, malignant melanoma, malignant pancreatic insulinoma, medullary thyroid carcinoma, melanoma, multiple myeloma, mycosis fungoides, myeloid and lymphocytic leukemia, neuroblastoma, non-Hodgkin's lymphoma, non-small cell lung cancer, osteogenic sarcoma, ovarian carcinoma, pancreatic carcinoma, polycythemia vera, primary brain carcinoma, primary macroglobulinemia, prostatic cancer, renal cell cancer, rhabdomyosarcoma, skin cancer, small-cell lung cancer, soft-tissue sarcoma, squamous cell cancer, stomach cancer, testicular cancer, thyroid cancer and Wilms' tumor.

The invention further preferably relates to a medicament comprising at least one compound according to the invention and/or one of its physiologically acceptable salts, derivatives, solvates, prodrugs and stereoisomers, including mixtures thereof in all ratios, for use in the treatment and/or prophylaxis of physiological and/or pathophysiological states selected from the group consisting of hyperproliferative and infectious diseases and disorders, wherein the hyperproliferative disease or disorder is selected from the group consisting of age-related macular degeneration, Crohn's disease, cirrhosis, chronic inflammatory-related disorders, proliferative diabetic retinopathy, proliferative vitreoretinopathy, retinopathy of prematurity, granulomatosis,

immune hyperproliferation associated with organ or tissue transplantation and an immunoproliferative disease or disorder selected from the group consisting of inflammatory bowel disease, psoriasis, rheumatoid arthritis, systemic lupus erythematosus (SLE), vascular hyperproliferation secondary to retinal hypoxia and vasculitis.

The invention further preferably relates to a medicament comprising at least one compound according to the invention and/or one of its physiologically acceptable salts, derivatives, solvates, prodrugs and stereoisomers, including mixtures thereof in all ratios, for use in the treatment and/or prophylaxis of physiological and/or pathophysiological states selected from the group consisting of hyperproliferative and infectious diseases and disorders, wherein the infectious disease or disorder is selected from the group consisting of

- a) virally induced infectious diseases which are caused by retroviruses, hepadnaviruses, herpesviruses, flaviviridae and/or adenoviruses wherein the retroviruses are selected from lentiviruses or oncoretroviruses, wherein the lentivirus is selected from the group consisting of HIV-1, HIV-2, FIV, BIV, SIVs, SHIV, CAEV, VMV and EIAV and the oncoretrovirus is selected from the group consisting of HTLV-I, HTLV-II and BLV, the hepadnavirus is selected from the group consisting of HBV, GSHV and WHV, the herpesvirus is selected from the group from the group consisting of HSV I, HSV II, EBV, VZV, HCMV or HHV 8 and the flaviviridae is selected from the group consisting of HCV, West Nile and Yellow Fever,
- b) bacterial infectious diseases which are caused by Gram-positive bacteria wherein the Gram-positive bacteria are selected from the group consisting of methicillin-susceptible and methicillin-resistant staphylococci (including *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Staphylococcus haemolyticus*, *Staphylococcus hominis*, *Staphylococcus saprophyticus*, and coagulase-negative staphylococci), glycopeptides-intermediate susceptible *Staphylococcus aureus* (GISA), penicillin-susceptible and penicillin-resistant streptococci (including *Streptococcus pneumoniae*, *Streptococcus pyogenes*, *Streptococcus agalactiae*, *Streptococcus avium*, *Streptococcus bovis*, *Streptococcus lactis*, *Streptococcus sanguis* and Streptococci Group C (GCS), Streptococci Group G (GGS) and *viridans* streptococci), enterococci (including vancomycin-susceptible and vancomycin-resistant strains such as *Enterococcus faecalis* and *Enterococcus faecium*), *Clostridium difficile*, *Listeria monocytogenes*, *Corynebacterium jeikeium*, *Chlamydia* spp (including *C. pneumoniae*) and *Mycobacterium tuberculosis*,
- c) bacterial infectious diseases which are caused by Gram-negative bacteria wherein the Gram-negative bacteria are selected from the group consisting of the Genus Enterobacteriaceae, including *Escherichia* spp. (including *Escherichia coli*), *Klebsiella* spp., *Enterobacter* spp., *Citrobacter* spp., *Serratia* spp., *Proteus* spp., *Providencia* spp., *Salmonella* spp., *Shigella* spp., the genus *Pseudomonas* (including *P. aeruginosa*), *Moraxella* spp. (including *M. catarrhalis*), *Haemophilus* spp. and *Neisseria* spp.,
- d) infectious diseases induced by intracellular active parasites selected from the group consisting of phylum Apicomplexa, or Sarcocystidophora (including Trypanosoma, Plasmodia, Leishmania, Babesia or Theileria), Cryptosporidia, Sarcocystida, Amoebia, Coccidia and Trichomonadia.

It is intended that the medicaments disclosed above include a corresponding use of the compounds according to the invention for the preparation of a medicament for the

treatment and/or prophylaxis of the above physiological and/or pathophysiological states.

It is additionally intended that the medicaments disclosed above include a corresponding method for the treatment and/or prophylaxis of the above physiological and/or pathophysiological states in which at least one compound according to the invention is administered to a patient in need of such a treatment.

The compounds according to the invention preferably exhibit an advantageous biological activity which can easily be demonstrated in enzyme assays and animal experiments, as described in the examples. In such enzyme-based assays, the compounds according to the invention preferably exhibit and cause an inhibiting effect, which is usually documented by IC₅₀ values in a suitable range, preferably in the micromolar range and more preferably in the nanomolar range.

The compounds according to the invention can be administered to humans or animals, in particular mammals, such as apes, dogs, cats, rats or mice, and can be used in the therapeutic treatment of the human or animal body and in the combating of the above-mentioned diseases. They can furthermore be used as diagnostic agents or as reagents.

Furthermore, compounds according to the invention can be used for the isolation and investigation of the activity or expression of adenosine A_{2A} and/or A_{2B} receptors. In addition, they are particularly suitable for use in diagnostic methods for diseases in connection with disturbed adenosine A_{2A} and/or A_{2B} receptor activity. The invention therefore furthermore relates to the use of the compounds according to the invention for the isolation and investigation of the activity or expression of adenosine A_{2A} and/or A_{2B} receptors or as binders and inhibitors of adenosine A_{2A} and/or A_{2B} receptors.

For diagnostic purposes, the compounds according to the invention can, for example, be radioactively labelled. Examples of radioactive labels are ³H, ¹⁴C, ²³¹I and ¹²⁵I. A preferred labelling method is the iodogen method (Fraker et al., 1978). In addition, the compounds according to the invention can be labelled by enzymes, fluorophores and chemophores. Examples of enzymes are alkaline phosphatase, β-galactosidase and glucose oxidase, an example of a fluorophore is fluorescein, an example of a chemophore is luminol, and automated detection systems, for example for fluorescent colorations, are described, for example, in U.S. Pat. Nos. 4,125,828 and 4,207,554.

The present invention further relates to pharmaceutical compositions containing the compounds of the present invention and their use for the treatment and/or prophylaxis of diseases and disorders where the partial or total inactivation of adenosine A_{2A} and/or A_{2B} receptors could be beneficial.

The compounds of the formula I can be used for the preparation of pharmaceutical preparations, in particular by non-chemical methods. In this case, they are brought into a suitable dosage form together with at least one solid, liquid and/or semi-liquid excipient or adjuvant and optionally in combination with one or more further active compound(s).

The invention therefore furthermore relates to pharmaceutical preparations comprising at least one compound of the formula I and/or physiologically acceptable salts, derivatives, solvates and stereoisomers thereof, including mixtures thereof in all ratios. In particular, the invention also relates to pharmaceutical preparations which comprise further excipients and/or adjuvants, and also to pharmaceutical preparations which comprise at least one further medicinal active compound.

In particular, the invention also relates to a process for the preparation of a pharmaceutical preparation, characterised in that a compound of the formula I and/or one of its physiologically acceptable salts, derivatives, solvates and stereoisomers, including mixtures thereof in all ratios, is brought into a suitable dosage form together with a solid, liquid or semi-liquid excipient or adjuvant and optionally with a further medicament active compound.

The pharmaceutical preparations according to the invention can be used as medicaments in human or veterinary medicine. The patient or host can belong to any mammal species, for example a primate species, particularly humans; rodents, including mice, rats and hamsters; rabbits; horses, cattle, dogs, cats, etc. Animal models are of interest for experimental investigations, where they provide a model for the treatment of a human disease.

Suitable carrier substances are organic or inorganic substances which are suitable for enteral (for example oral), parenteral or topical administration and do not react with the novel compounds, for example water, vegetable oils (such as sunflower oil or cod-liver oil), benzyl alcohols, polyethylene glycols, gelatine, carbohydrates, such as lactose or starch, magnesium stearate, talc, lanolin or Vaseline. Owing to his expert knowledge, the person skilled in the art is familiar with which adjuvants are suitable for the desired medication formulation. Besides solvents, for example water, physiological saline solution or alcohols, such as, for example, ethanol, propanol or glycerol, sugar solutions, such as glucose or mannitol solutions, or a mixture of the said solvents, gel formers, tablet assistants and other active-ingredient carriers, it is also possible to use, for example, lubricants, stabilisers and/or wetting agents, emulsifiers, salts for influencing the osmotic pressure, antioxidants, dispersants, antifoams, buffer substances, flavours and/or aromas or flavour correctants, preservatives, solubilisers or dyes. If desired, preparations or medicaments according to the invention may comprise one or more further active compounds, for example one or more vitamins.

If desired, preparations or medicaments according to the invention may comprise one or more further active compounds and/or one or more action enhancers (adjuvants).

The terms "pharmaceutical formulation" and "pharmaceutical preparation" are used as synonyms for the purposes of the present invention.

As used here, "pharmaceutically tolerated" relates to medicaments, precipitation reagents, excipients, adjuvants, stabilisers, solvents and other agents which facilitate the administration of the pharmaceutical preparations obtained therefrom to a mammal without undesired physiological side effects, such as, for example, nausea, dizziness, digestion problems or the like.

In pharmaceutical preparations for parenteral administration, there is a requirement for isotonicity, euhydration and tolerability and safety of the formulation (low toxicity), of the adjuvants employed and of the primary packaging. Surprisingly, the compounds according to the invention preferably have the advantage that direct use is possible and further purification steps for the removal of toxicologically unacceptable agents, such as, for example, high concentrations of organic solvents or other toxicologically unacceptable adjuvants, are thus unnecessary before use of the compounds according to the invention in pharmaceutical formulations.

The invention particularly preferably also relates to pharmaceutical preparations comprising at least one compound according to the invention in precipitated non-crystalline, precipitated crystalline or in dissolved or suspended form,

and optionally excipients and/or adjuvants and/or further pharmaceutical active compounds.

The compounds according to the invention preferably enable the preparation of highly concentrated formulations without unfavourable, undesired aggregation of the compounds according to the invention occurring. Thus, ready-to-use solutions having a high active-ingredient content can be prepared with the aid of compounds according to the invention with aqueous solvents or in aqueous media.

The compounds and/or physiologically acceptable salts and solvates thereof can also be lyophilised and the resultant lyophilisates used, for example, for the preparation of injection preparations.

Aqueous preparations can be prepared by dissolving or suspending compounds according to the invention in an aqueous solution and optionally adding adjuvants. To this end, defined volumes of stock solutions comprising the said further adjuvants in defined concentration are advantageously added to a solution or suspension having a defined concentration of compounds according to the invention, and the mixture is optionally diluted with water to the pre-calculated concentration. Alternatively, the adjuvants can be added in solid form. The amounts of stock solutions and/or water which are necessary in each case can subsequently be added to the aqueous solution or suspension obtained. Compounds according to the invention can also advantageously be dissolved or suspended directly in a solution comprising all further adjuvants.

The solutions or suspensions comprising compounds according to the invention and having a pH of 4 to 10, preferably having a pH of 5 to 9, and an osmolality of 250 to 350 mOsmol/kg can advantageously be prepared. The pharmaceutical preparation can thus be administered directly substantially without pain intravenously, intraarterially, intra-articularly, subcutaneously or percutaneously. In addition, the preparation may also be added to infusion solutions, such as, for example, glucose solution, isotonic saline solution or Ringer's solution, which may also contain further active compounds, thus also enabling relatively large amounts of active compound to be administered.

Pharmaceutical preparations according to the invention may also comprise mixtures of a plurality of compounds according to the invention.

The preparations according to the invention are physiologically well tolerated, easy to prepare, can be dispensed precisely and are preferably stable with respect to assay, decomposition products and aggregates throughout storage and transport and during multiple freezing and thawing processes. They can preferably be stored in a stable manner over a period of at least three months to two years at refrigerator temperature (2-8° C.) and at room temperature (23-27° C.) and 60% relative atmospheric humidity (R.H.).

For example, the compounds according to the invention can be stored in a stable manner by drying and when necessary converted into a ready-to-use pharmaceutical preparation by dissolution or suspension. Possible drying methods are, for example, without being restricted to these examples, nitrogen-gas drying, vacuum-oven drying, lyophilisation, washing with organic solvents and subsequent air drying, liquid-bed drying, fluidised-bed drying, spray drying, roller drying, layer drying, air drying at room temperature and further methods.

The term "effective amount" denotes the amount of a medicament or of a pharmaceutical active compound which causes in a tissue, system, animal or human a biological or medical response which is sought or desired, for example, by a researcher or physician.

In addition, the term "therapeutically effective amount" denotes an amount which, compared with a corresponding subject who has not received this amount, has the following consequence: improved treatment, healing, prevention or elimination of a disease, syndrome, disease state, complaint, disorder or prevention of side effects or also a reduction in the progress of a disease, complaint or disorder. The term "therapeutically effective amount" also encompasses the amounts which are effective for increasing normal physiological function.

On use of preparations or medicaments according to the invention, the compounds according to the invention and/or physiologically acceptable salts and solvates thereof are generally used analogously to known, commercially available preparations or preparations, preferably in dosages of between 0.1 and 500 mg, in particular 5 and 300 mg, per use unit. The daily dose is preferably between 0.001 and 250 mg/kg, in particular 0.01 and 100 mg/kg, of body weight. The preparation can be administered one or more times per day, for example two, three or four times per day. However, the individual dose for a patient depends on a large number of individual factors, such as, for example, on the efficacy of the particular compound used, on the age, body weight, general state of health, sex, nutrition, on the time and method of administration, on the excretion rate, on the combination with other medicaments and on the severity and duration of the particular disease.

A measure of the uptake of a medicament active compound in an organism is its bioavailability. If the medicament active compound is delivered to the organism intravenously in the form of an injection solution, its absolute bioavailability, i.e. the proportion of the pharmaceutical which reaches the systemic blood, i.e. the major circulation, in unchanged form, is 100%. In the case of oral administration of a therapeutic active compound, the active compound is generally in the form of a solid in the formulation and must therefore first be dissolved in order that it is able to overcome the entry barriers, for example the gastrointestinal tract, the oral mucous membrane, nasal membranes or the skin, in particular the stratum corneum, or can be absorbed by the body. Data on the pharmacokinetics, i.e. on the bioavailability, can be obtained analogously to the method of J. Shaffer et al., *J. Pharm. Sciences*, 88 (1999), 313-318.

Furthermore, medicaments of this type can be prepared by means of one of the processes generally known in the pharmaceutical art.

Medicaments can be adapted for administration via any desired suitable route, for example by the oral (including buccal or sublingual), rectal, pulmonary, nasal, topical (including buccal, sublingual or transdermal), vaginal or parenteral (including subcutaneous, intramuscular, intravenous, intradermal and in particular intra-articular) routes. Medicaments of this type can be prepared by means of all processes known in the pharmaceutical art by, for example, combining the active compound with the excipient(s) or adjuvant(s).

Parenteral administration is preferably suitable for administration of the medicaments according to the invention. In the case of parenteral administration, intra-articular administration is particularly preferred.

The invention thus preferably also relates to the use of a pharmaceutical preparation according to the invention for intra-articular administration in the treatment and/or prophylaxis of physiological and/or pathophysiological states

selected from the group consisting of osteoarthritis, traumatic cartilage injuries, arthritis, pain, allodynia or hyperalgesia.

Intra-articular administration has the advantage that the compound according to the invention can be administered directly into the synovial fluid in the vicinity of the joint cartilage and is also able to diffuse from there into the cartilage tissue. Pharmaceutical preparations according to the invention can thus also be injected directly into the joint gap and thus develop their action directly at the site of action as intended. The compounds according to the invention are also suitable for the preparation of medicaments to be administered parenterally having slow, sustained and/or controlled release of active compound. They are thus also suitable for the preparation of delayed-release formulations, which are advantageous for the patient since administration is only necessary at relatively large time intervals.

The medicaments adapted to parenteral administration include aqueous and non-aqueous sterile injection solutions comprising antioxidants, buffers, bacteriostatics and solutes, by means of which the formulation is rendered isotonic with the blood or synovial fluid of the recipient to be treated; as well as aqueous and non-aqueous sterile suspensions, which can comprise suspension media and thickeners. The formulations can be delivered in single-dose or multi-dose containers, for example sealed ampoules and vials, and stored in the freeze-dried (lyophilised) state, so that only the addition of the sterile carrier liquid, for example water for injection purposes, immediately before use is necessary. Injection solutions and suspensions prepared in accordance with the formulation can be prepared from sterile powders, granules and tablets.

The compounds according to the invention can also be administered in the form of liposome delivery systems, such as, for example, small unilamellar vesicles, large unilamellar vesicles and multilamellar vesicles. Liposomes can be formed from various phospholipids, such as, for example, cholesterol, stearylamine or phosphatidylcholines.

The compounds according to the invention can also be coupled to soluble polymers as targeted medicament excipients. Such polymers can encompass polyvinylpyrrolidone, pyran copolymer, polyhydroxypropylmethacrylamidophenol, polyhydroxyethylaspartamidophenol or polyethylene oxide polylysine, substituted by palmitoyl radicals. The compounds according to the invention can furthermore be coupled to a class of biodegradable polymers which are suitable for achieving slow release of a medicament, for example polylactic acid, poly-epsilon-caprolactone, polyhydroxybutyric acid, polyorthoesters, polyacetals, polydihydroxypyranes, polycyanoacrylates, polylactic-co-glycolic acid, polymers, such as conjugates between dextran and methacrylates, polyphosphoesters, various polysaccharides and polyamines and poly-epsilon-caprolactone, albumin, chitosan, collagen or modified gelatine and crosslinked or amphiphilic block copolymers of hydrogels.

Suitable for enteral administration (oral or rectal) are, in particular, tablets, dragees, capsules, syrups, drops or suppositories, and suitable for topical use are ointments, creams, pastes, lotions, gels, sprays, foams, aerosols, solutions (for example solutions in alcohols, such as ethanol or isopropanol, acetonitrile, DMF, dimethylacetamide, 1,2-propanediol or mixtures thereof with one another and/or with water) or powders. Also particularly suitable for topical uses are liposomal preparations.

In the case of formulation to give an ointment, the active compound can be employed either with a paraffinic or a water-miscible cream base. Alternatively, the active com-

powder can be formulated to a cream with an oil-in-water cream base or a water-in-oil base.

Medicaments adapted to transdermal administration can be delivered as independent plasters for extended, close contact with the epidermis of the recipient.

Thus, for example, the active compound can be supplied from the plaster by means of iontophoresis, as described in general terms in *Pharmaceutical Research*, 3 (6), 318 (1986).

It goes without saying that, besides the constituents particularly mentioned above, the medicaments according to the invention may also comprise other agents usual in the art with respect to the particular type of pharmaceutical formulation.

The invention also relates to a set (kit) consisting of separate packs of

- a) an effective amount of a compound of the formula I and/or physiologically acceptable salts, derivatives, solvates, prodrugs and stereoisomers thereof, including mixtures thereof in all ratios, and
- b) an effective amount of a further medicament active compound.

The set comprises suitable containers, such as boxes or cartons, individual bottles, bags or ampoules. The set may, for example, comprise separate ampoules each containing an effective amount of a compound of the formula I and/or pharmaceutically acceptable salts, derivatives, solvates, prodrugs and stereoisomers thereof, including mixtures thereof in all ratios, and an effective amount of a further medicament active compound in dissolved or lyophilised form.

Furthermore, the medicaments according to the invention can be used in order to provide additive or synergistic effects in certain known therapies and/or can be used in order to restore the efficacy of certain existing therapies.

Besides the compounds according to the invention, the pharmaceutical preparations according to the invention may also comprise further medicament active compounds, for example for use in the treatment of cancer, other anti-tumor medicaments. For the treatment of the other diseases mentioned, the pharmaceutical preparations according to the invention may also, besides the compounds according to the invention, comprise further medicament active compounds which are known to the person skilled in the art in the treatment thereof.

In one principal embodiment, methods are provided for enhancing an immune response in a host in need thereof. The immune response can be enhanced by reducing T cell tolerance, including by increasing IFN- γ release, by decreasing regulatory T cell production or activation, or by increasing antigen-specific memory T cell production in a host. In one embodiment, the method comprises administering a compound of the present invention to a host in combination or alternation with an antibody. In particular subembodiments, the antibody is a therapeutic antibody. In one particular embodiment, a method of enhancing efficacy of passive antibody therapy is provided comprising administering a compound of the present invention in combination or alternation with one or more passive antibodies. This method can enhance the efficacy of antibody therapy for treatment of abnormal cell proliferative disorders such as cancer, or can enhance the efficacy of therapy in the treatment or prevention of infectious diseases. The compound of the present invention can be administered in combination or alternation with antibodies such as rituximab, herceptin or ebitux, for example.

In another principal embodiment, a method of treating or preventing abnormal cell proliferation is provided compris-

ing administering a compound of the present invention to a host in need thereof substantially in the absence of another anti-cancer agent.

In another principal embodiment, a method of treating or preventing abnormal cell proliferation in a host in need thereof is provided, comprising administering a first compound of the present invention substantially in combination with a first anti-cancer agent to the host and subsequently administering a second A_{2A} and/or A_{2B} receptor antagonist. In one subembodiment, the second antagonist is administered substantially in the absence of another anti-cancer agent. In another principal embodiment, a method of treating or preventing abnormal cell proliferation in a host in need thereof is provided, comprising administering a compound of the present invention substantially in combination with a first anti-cancer agent to the host and subsequently administering a second anti-cancer agent in the absence of the antagonist.

Thus, the cancer treatment disclosed here can be carried out as therapy with a compound of the present invention or in combination with an operation, irradiation or chemotherapy. Chemotherapy of this type can include the use of one or more active compounds of the following categories of antitumor active compounds:

(i) antiproliferative/antineoplastic/DNA-damaging active compounds and combinations thereof, as used in medical oncology, such as alkylating active compounds (for example cis-platin, carboplatin, cyclophosphamide, nitrogen mustard, melphalan, chlorambucil, busulfan and nitrosoureas); antimetabolites (for example antifolates such as fluoropyrimidines such as 5-fluorouracil and tegafur, raltitrexed, methotrexate, cytosine arabinoside, hydroxyurea and gemcitabine); antitumor antibiotics (for example anthracyclines, such as adriamycin, bleomycin, doxorubicin, daunomycin, epirubicin, idarubicin, mitomycin-C, dactinomycin and mithramycin); antimitotic active compounds (for example vinca alkaloids, such as vincristine, vinblastine, vindesine and vinorelbine, and taxoids, such as taxol and taxotere); topoisomerase inhibitors (for example epipodophyllotoxins, such as etoposide and teniposide, amsacrine, topotecan, irinotecan and camptothecin) and cell-differentiating active compounds (for example all-trans-retinoic acid, 13-cis-retinoic acid and fenretinide);

(ii) cytostatic active compounds, such as anti-oestrogens (for example tamoxifen, toremifene, raloxifene, droloxifene and idoxifene), oestrogen receptor regulators (for example fulvestrant), anti-androgens (for example bicalutamide, flutamide, nilutamide and cyproterone acetate), LHRH antagonists or LHRH agonists (for example goserelin, leuprorelin and buserelin), progestones (for example megestrol acetate), aromatase inhibitors (for example anastrozole, letrozole, vorazole and exemestane) and inhibitors of 5 α -reductase, such as finasteride;

(iii) active compounds which inhibit cancer invasion including for example metalloproteinase inhibitors, like marimastat, and inhibitors of urokinase plasminogen activator receptor function;

(iv) inhibitors of growth factor function, for example growth factor antibodies, growth factor receptor antibodies, for example the anti-erbB2 antibody trastuzumab [HerceptinTM] and the anti-erbB1 antibody cetuximab [C225], farnesyl transferase inhibitors, tyrosine kinase inhibitors and serine/threonine kinase inhibitors, for example inhibitors of the epidermal growth factor family (for example EGFR family tyrosine kinase inhibitors, such as N-(3-chloro-4-fluorophenyl)-7-methoxy-6-(3-morpholinopropoxy) quinazolin-4-amine (gefitinib, AZD1839), N-(3-ethynylphe-

nyl)-6,7-bis (2-methoxyethoxy)quinazolin-4-amine (erlotinib, OSI-774) and 6-acrylamido-N-(3-chloro-4-fluorophenyl)-7-(3-morpholinopropoxy)quinazolin-4-amine (CI 1033), for example inhibitors of the platelet-derived growth factor family and, for example, inhibitors of the hepatocyte growth factor family;

(v) anti-angiogenic active compounds, such as bevacizumab, angiostatin, endostatin, linomide, batimastat, captopril, cartilage derived inhibitor, genistein, interleukin 12, lavendustin, medroxyprogesterone acetate, recombinant human platelet factor 4, tecogalan, thrombospondin, TNP-470, anti-VEGF monoclonal antibody, soluble VEGF-receptor chimaeric protein, anti-VEGF receptor antibodies, anti-PDGF receptors, inhibitors of integrins, tyrosine kinase inhibitors, serine/threonine kinase inhibitors, antisense oligonucleotides, antisense oligodeoxynucleotides, siRNAs, anti-VEGF aptamers, pigment epithelium derived factor and compounds which have been published in the international patent applications WO 97/22596, WO 97/30035, WO 97/32856 and WO 98/13354);

(vi) vessel-destroying agents, such as combretastatin A4 and compounds which have been published in the international patent applications WO 99/02166, WO 00/40529, WO 00/41669, WO 01/92224, WO 02/04434 and WO 02/08213;

(vii) antisense therapies, for example those directed to the targets mentioned above, such as ISIS 2503, an anti-Ras antisense;

(viii) gene therapy approaches, including, for example, approaches for replacement of abnormal, modified genes, such as abnormal p53 or abnormal BRCA1 or BRCA2, GDEPT approaches (gene-directed enzyme pro-drug

therapy), such as those which use cytosine deaminase, thymidine kinase or a bacterial nitroreductase enzyme, and approaches which increase the tolerance of a patient to chemotherapy or radiotherapy, such as multi-drug resistance therapy; and

(ix) immunotherapy approaches, including, for example, ex-vivo and in-vivo approaches for increasing the immunogenicity of tumour cells of a patient, such as transfection with cytokines, such as interleukin 2, interleukin 4 or granulocyte macrophage colony stimulating factor, approaches for decreasing T-cell anergy, approaches using transfected immune cells, such as cytokine-transfected dendritic cells, approaches for use of cytokine-transfected tumour cells and approaches for use of anti-idiotypic antibodies

(x) chemotherapeutic agents including for example abarelix, aldesleukin, alemtuzumab, alitretinoin, allopurinol, altretamine, amifostine, anastrozole, arsenic trioxide, asparaginase, BCG live, bevacizumab, bexarotene, bleomycin, bortezomib, busulfan, calusterone, camptothecin, capecitabine, carboplatin, carmustine, celecoxib, cetuximab, chlorambucil, cinacalcet, cisplatin, cladribine, cyclophosphamide, cytarabine, dacarbazine, dactinomycin, darbepoetin alfa, daunorubicin, denileukin difitox, dexrazoxane, docetaxel, doxorubicin, dromostanolone, epirubicin, epoetin alfa, estramustine, etoposide, exemestane, filgrastim, floxuridine, fludarabine, fluorouracil, fulvestrant and gemcitabine.

The medicaments from table 1 can preferably, but not exclusively, be combined with the compounds of the formula I.

TABLE 1

Alkylating active compounds	Cyclophosphamide	Lomustine
	Busulfan	Procarbazine
	Ifosfamide	Altretamine
	Melphalan	Estramustine phosphate
	Hexamethylmelamine	Mechloroethamine
	Thiotepa	Streptozocin
	chlorambucil	Temozolomide
	Dacarbazine	Semustine
	Carmustine	
	Cisplatin	Carboplatin
	Oxaliplatin	ZD-0473 (AnorMED)
Platinum active compounds	Spiroplatin	Lobaplatin (Aetema)
	Carboxyphthalatoplatinum	Satraplatin (Johnson Matthey)
	Tetraplatin	BBR-3464
	Orniplatin	(Hoffmann-La Roche)
	Iproplatin	SM-11355 (Sumitomo)
		AP-5280 (Access)
		Tomudex
		Trimetrexate
		Deoxycofomycin
		Fludarabine
		Pentostatin
Antimetabolites	Azacytidine	Raltitrexed
	Gemcitabine	Hydroxyurea
	Capecitabine	Decitabine (SuperGen)
	5-Fluorouracil	Clofarabine (Bioenvision)
	Floxuridine	Irofulven (MGI Pharma)
	2-Chlorodesoxyadenosine	DMDC (Hoffmann-La Roche)
	6-Mercaptopurine	Ethynylcytidine (Taiho)
	6-Thioguanine	Rubitecan (SuperGen)
	Cytarabine	Exatecan mesylate (Daiichi)
	2-Fluorodesoxycytidine	Quinamed (ChemGenex)
	Methotrexate	Gimatecan (Sigma-Tau)
Topoisomerase inhibitors	Idatrexate	Diflomotecan (Beaufour-Ipsen)
	Amsacrine	TAS-103 (Taiho)
	Epirubicin	Elsamitrucin (Spectrum)
	Etoposide	J-107088 (Merck & Co)
	Teniposide or mitoxantrone	BNP-1350 (BioNumerik)
	Irinotecan (CPT-11)	
	7-ethyl-10-hydroxycamptothecin	
	Topotecan	
	Dexrazoxanet (TopoTarget)	
	Pixantrone (Novuspharma)	

TABLE 1-continued

	Rebeccamycin analogue (Exelixis)	CKD-602 (Chong Kun Dang)
	BBR-3576 (Novuspharma)	KW-2170 (Kyowa Hakko)
Antitumour antibiotics	Dactinomycin (Actinomycin D)	Amonafide
	Doxorubicin (Adriamycin)	Azonafide
	Deoxyrubicin	Anthrapyrazole
	Valrubicin	Oxantrazole
	Daunorubicin (Daunomycin)	Losoxantrone
	Epirubicin	Bleomycin sulfate (Blenoxan)
	Therapeutic	Bleomycinic acid
	Idarubicin	Bleomycin A
	Rubidazon	Bleomycin B
	Plicamycin	Mitomycin C
	Porfiromycin	MEN-10755 (Menarini)
	Cyanomorpholinodoxorubicin	GPX-100 (Gem Pharmaceuticals)
	Mitoxantron (Novantron)	
Antimitotic active compounds	Paclitaxel	SB 408075
	Docetaxel	(GlaxoSmithKline)
	Colchicine	E7010 (Abbott)
	Vinblastine	PG-TXL (Cell Therapeutics)
	Vincristine	IDN 5109 (Bayer)
	Vinorelbine	A 105972 (Abbott)
	Vindesine	A 204197 (Abbott)
	Dolastatin 10 (NCI)	LU 223651 (BASF)
	Rhizoxin (Fujisawa)	D 24851 (ASIA Medica)
	Mivobulin (Warner-Lambert)	ER-86526 (Eisai)
	Cemadotin (BASF)	Combretastatin A4 (BMS)
	RPR 109881A (Aventis)	Isohomohalichondrin-B (PharmaMar)
	TXD 258 (Aventis)	ZD 6126 (AstraZeneca)
	Epothilone B (Novartis)	PEG-Paclitaxel (Enzon)
	T 900607 (Tularik)	AZ10992 (Asahi)
	T 138067 (Tularik)	IDN-5109 (Indena)
	Cryptophycin 52 (Eli Lilly)	AVLB (Prescient NeuroPharma)
	Vinflunine (Fabre)	Azaepothilone B (BMS)
	Auristatin PE (Teikoku Hormone)	BNP-7787 (BioNumerik)
	BMS 247550 (BMS)	CA-4-prodrug (OXIGENE)
	BMS 184476 (BMS)	Dolastatin-10 (Nrh)
	BMS 188797 (BMS)	CA-4 (OXIGENE)
Aromatase inhibitors	Taxoprexin (Protarga)	Exemestan
	Aminoglutethimide	Atamestan (BioMedicines)
	Letrozole	YM-511 (Yamanouchi)
	Anastrozole	
	Formestan	
Thymidylate Synthase inhibitors	Pemetrexed (Eli Lilly)	Nolatrexed (Eximias)
	ZD-9331 (BTG)	CoFactor™ (BioKeys)
DNA antagonists	Trabectedin (PharmaMar)	Mafosfamide (Baxter International)
	Glufosfamide (Baxter International)	Apaziquone (Spectrum Pharmaceuticals)
	Albumin + 32 P (isotope solutions)	O6-benzylguanine (Paligent)
	Thymectacin (NewBiotics)	
	Edotreotide (Novartis)	
Farnesyl transferase inhibitors	Arglabin (NuOncology Labs)	Tipifarnib (Johnson & Johnson)
	Lonafarnib (Schering-Plough)	Perillyl alcohol (DOR BioPharma)
	BAY-43-9006 (Bayer)	Zosuquidar trihydrochloride (Eli Lilly)
Pump inhibitors	CBT-1 (CBA Pharma)	Biricodar dicitrate (Vertex)
	Tariquidar (Xenova)	Pivaloyloxymethyl butyrate (Titan)
	MS-209 (Schering AG)	Depsipeptide (Fujisawa)
Histone acetyl transferase inhibitors	Tacedinaline (Pfizer)	CMT-3 (CollaGenex)
	SAHA (Aton Pharma)	BMS-275291 (Celltech)
	MS-275 (Schering AG)	Tezacitabine (Aventis)
Metalloproteinase inhibitors	Neovastat (Aeterna Laboratories)	Didox (Molecules for Health)
Ribonucleoside reductase inhibitors	Marimastat (British Biotech)	
	Gallium maltolate (Titan)	
	Triapin (Vion)	
TNF-alpha agonists/antagonists	Virulizin (Lorus Therapeutics)	Revimid (Celgene)
	CDC-394	(Celgene)
Endothelin-A receptor antagonists	Atrasentan (Abbot)	YM-598 (Yamanouchi)
	ZD-4054 (AstraZeneca)	
Retinoic acid receptor agonists	Fenretinide (Johnson & Johnson)	Alitretinoin (Ligand)
	LGD-1550 (ligand)	

TABLE 1-continued

Immunomodulators	Interferon	Dexosome therapy (Anosys)
	Oncophage (Antigenics)	Pentrix (Australian Cancer Technology)
	GMK (Progenics)	JSF-154 (Tragen)
	Adenocarcinoma vaccine (Biomira)	Cancer vaccine (Intercell)
	CTP-37 (AVI BioPharma)	Norelin (Biostar)
	JRX-2 (Immuno-Rx)	BLP-25 (Biomira)
	PEP-005 (Peplin Biotech)	MGV (Progenics)
	Synchrovax vaccines (CTL Immuno)	!3-Alethin (Dovetail)
	Melanoma vaccines (CTL Immuno)	CLL-Thera (Vasogen)
	p21-RAS vaccine (GemVax)	
Hormonal and antihormonal active compounds	Oestrogens	Prednisone
	Conjugated oestrogens	Methylprednisolone
	Ethinylloestradiol	Prednisolone
	Chlorotrianisene	Aminoglutethimide
	Idenestrol	Leuprolide
	Hydroxyprogesterone caproate	Goserelin
	Medroxyprogesterone	Leuporelin
	Testosterone	Bicalutamide
	Testosterone propionate	Flutamide
	Fluoxymesterone	Octreotide
	Methyltestosterone	Nilutamide
	Diethylstilbestrol	Mitotan
	Megestrol	P-04 (Novogen)
Tamoxifen	2-Methoxyoestradiol (En_treMed)	
Toremofin	Arzoxifen (Eli Lilly)	
Dexamethasone		
Photodynamic active compounds	Talaporphin (Light Sciences)	Pd bacteriopheophorbide (Yeda)
	Theralux (Theratechnologies)	Lutetium texaphyrin (Pharmacyclics)
	Motexaf in-Gadolinium (Pharmacyclics)	Hypericin
		Kahalide F (PharmaMar)
Tyrosine kinase inhibitors	Imatinib (Novartis)	CEP-701 (Cephalon)
	Leflunomide(Sugen/Pharmacia)	CEP-751 (Cephalon)
	ZDI839 (AstraZeneca)	MLN518 (Millenium)
	Erlotinib (Oncogene Science)	PKC412 (Novartis)
	Canertinib (Pfizer)	Phenoxodiol O
	Squalamine (Genaera)	Trastuzumab (Genentech)
	SU5416 (Pharmacia)	C225 (ImClone)
	SU6668 (Pharmacia)	rhu-Mab (Genentech)
	ZD4190 (AstraZeneca)	MDX-H210 (Medarex)
	ZD6474 (AstraZeneca)	2C4 (Genentech)
	Vatalanib (Novartis)	MDX-447 (Medarex)
	PKI166 (Novartis)	ABX-EGF (Abgenix)
	GW2016 (GlaxoSmithKline)	IMC-1C11 (ImClone)
EKB-509 (Wyeth)		
EKB-569 (Wyeth)		
Various other active compounds	SR-27897 (CCK-A inhibitor, Sanofi-Synthelabo)	BCX-1777 (PNP inhibitor, BioCryst)
	Tocladesine (cyclic AMP agonist, Ribapharm)	Ranpirnase (ribonuclease stimulant, Alfacell)
	Alvocidib (CDK inhibitor, Aventis)	Galarubicin (RNA synthesis inhibitor, Dong-A)
	CV-247 (COX-2 inhibitor, Ivy Medical)	Tirapazamine (reducing agent, SRI International)
	P54 (COX-2 inhibitor, Phytopharm)	N-Acetylcysteine (reducing agent, Zambon)
	CapCell™ (CYP450 stimulant, Bavarian Nordic)	R-Flurbiprofen (NF-kappaB inhibitor, Encore)
	GCS-IOO (gal3 antagonist, GlycoGenesys)	3CPA (NF-kappaB inhibitor, Active Biotech)
	G17DT immunogen (gastrin inhibitor, Aphton)	Seocalcitol (vitamin D receptor agonist, Leo)
	Efaproxiral (oxygenator, Allos Therapeutics)	131-I-TM-601 (DNA antagonist, TransMolecular)
	PI-88 (heparanase inhibitor, Progen)	Eflornithin (ODC inhibitor, ILEX Oncology)
	Tesmilifen (histamine antagonist, YM BioSciences)	Minodronic acid (osteoclast inhibitor, Yamanouchi)
	Histamine (histamine H2 receptor agonist, Maxim)	Indisulam (p53 stimulant, Eisai)
	Tiazofurin (IMPDH inhibitor, Ribapharm)	Aplidin (PPT inhibitor, PharmaMar)
	Cilengitide (integrin antagonist, Merck KGaA)	Rituximab (CD20 antibody, Genentech)
	SR-31747 (IL-1 antagonist, Sanofi-Synthelabo)	

TABLE 1-continued

CCI-779 (mTOR kinase inhibitor, Wyeth)	Gemtuzumab (CD33 antibody, Wyeth Ayerst)
Exisulind (PDE-V inhibitor, Cell Pathways)	PG2 (haematopoiesis promoter, Pharmagenesis)
CP-461 (PDE-V inhibitor, Cell Pathways)	Immunol™ (triclosan mouthwash, Endo)
AG-2037 (GART inhibitor, Pfizer)	Triacetyluridine (uridine prodrug, Wellstat)
WX-UK1 (plasminogen activator inhibitor, Wilex)	SN-4071 (sarcoma agent, Signature BioScience)
PBI-1402 (PMN stimulant, ProMetic LifeSciences)	TransMID-107™ (immunotoxin, KS Biomedix)
Bortezomib (proteasome inhibitor, Millennium)	PCK-3145 (apoptosis promoter, Procyon)
SRL-172 (T-cell stimulant, SR Pharma)	Doranidazole (apoptosis promoter, Pola)
TLK-286 (glutathione-S transferase inhibitor, Telik)	CHS-828 (cytotoxic agent, Leo)
PT-100 (growth factor agonist, Point Therapeutics)	trans-Retinoic acid (differentiator, NIH)
Midostaurin (PKC inhibitor, Novartis)	MX6 (apoptosis promoter, MAXIA)
Bryostatin-1 (PKC stimulant, GPC Biotech)	Apomine (apoptosis promoter, ILEX Oncology)
CDA-II (apoptosis promoter, Everlife)	Urocidin (apoptosis promoter, Bioniche)
SDX-101 (apoptosis promoter, Salmédix)	Ro-31-7453 (apoptosis promoter, La Roche)
Ceflatonin (apoptosis promoter, ChemGenex)	Brostallicin (apoptosis promoter, Pharmacia)

Even without further embodiments, it is assumed that a person skilled in the art will be able to use the above description in the broadest scope. The preferred embodiments should therefore merely be regarded as descriptive disclosure which is absolutely not limiting in any way.

The following examples are thus intended to explain the invention without limiting it. Unless indicated otherwise, percent data denote percent by weight. All temperatures are indicated in degrees Celsius. "Conventional work-up": water is added if necessary, the pH is adjusted, if necessary, to values between 2 and 10, depending on the constitution of the end product, the mixture is extracted with ethyl acetate or dichloromethane, the phases are separated, the organic phase is dried over sodium sulfate, filtered and evaporated, and the product is purified by chromatography on silica gel and/or by crystallisation.

Rf values on silica gel; mass spectrometry: EI (electron impact ionisation): M⁺, FAB (fast atom bombardment): (M+H)⁺, THF (tetrahydrofuran), NMP (N-methylpyrrolidone), DMSO (dimethyl sulfoxide), EA (ethyl acetate), MeOH (methanol), TLC (thin-layer chromatography)

List of Abbreviations

AUC Area under the plasma drug concentration-time curve

C_{max} Maximum plasma concentration

CL Clearance

CV Coefficient of variation

CYP Cytochrome P450

DMSO Dimethyl sulfoxide

F Bioavailability

f_a Fraction absorbed

iv Intravenous

LC-MS/MS Liquid chromatography tandem mass spectrometry

LLOQ Lower limit of quantification

NC Not calculated

ND Not determined

PEG Polyethylene glycol

Pgp Permeability glycoprotein

PK Pharmacokinetic(s)

po Per os (oral)

t_{1/2} Half-life

t_{max} Time at which maximum plasma concentration of drug is reached

UPLC Ultra performance liquid chromatography

V_{ss} Volume of distribution (at steady state)

v/v Volume to volume

EXAMPLE 1

Examples of Compounds of the Present Invention

The invention especially relates to the compounds of table 2 and physiologically acceptable salts, derivatives, solvates, prodrugs and stereoisomers thereof, including mixtures thereof in all ratios.

TABLE 2






examples of compounds of the present invention				
No.	Structure	IUPAC-Name	MW	[M + H] ⁺ 1
1		4-Hydroxy-4-methyl-piperidine-1-carboxylic acid (4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide	407.49	408
2		4-Dimethylaminomethyl-N-(4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-benzamide	427.53	429
3		4-Methoxymethyl-N-(4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-benzamide	414.48	415
4		1-(2-Methoxy-ethyl)-1H-pyrazole-4-carboxylic acid (4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide	418.48	419
5		2-Dimethylaminomethyl-N-(4-methoxy-7-morpholin-4-yl)-isonicotinamide	428.51	430

TABLE 2-continued

examples of compounds of the present invention				
No.	Structure	IUPAC-Name	MW	[M + H] ⁺ 1
6		2-Methoxymethyl-N-(4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-isonicotinamide	415.47	416
7		4-benzyl-4-hydroxy-N-[4-methoxy-7-(morpholin-4-yl)-[1,3]thiazolo[4,5-c]pyridin-2-yl]piperidine-1-carboxamide	483.59	485
8		N-[4-methoxy-7-(morpholin-4-yl)-[1,3]thiazolo[4,5-c]pyridin-2-yl]-5-(2-methoxyethoxy)pyrazine-2-carboxamide	446.49	447
9		2-Oxa-7-aza-spiro[4.4]nonane-7-carboxylic acid (4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide	419.50	421
10		3-Hydroxy-3-methyl-pyrrolidine-1-carboxylic acid (4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide	393.47	394

TABLE 2-continued

examples of compounds of the present invention				
No.	Structure	IUPAC-Name	MW	[M + H] ⁺ 1
11		8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid (4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide	433.53	435
12		4-Hydroxy-4-prop-2-ynyl-piperidine-1-carboxylic acid (4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide	431.51	433
13		1-(2-Methoxy-ethyl)-1H-pyrazole-4-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	413.46	414
14		4-Methoxymethyl-N-[4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-benzamide	409.47	410
15		N-(4-Methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-N',N'-dimethyl-terephthalamide	441.51	443

TABLE 2-continued

examples of compounds of the present invention				
No.	Structure	IUPAC-Name	MW	[M + H] ⁺ 1
16		4-Hydroxymethyl-4-methyl-piperidine-1-carboxylic acid (4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide	421.51	423
17		(5S)-N-[4-methoxy-7-(morpholin-4-yl)-[1,3]thiazolo[4,5-c]pyridin-2-yl]-2-oxa-7-azaspiro[4.4]nonane-7-carboxamide	419.50	421
18		(5R)-N-[4-methoxy-7-(morpholin-4-yl)-[1,3]thiazolo[4,5-c]pyridin-2-yl]-2-oxa-7-azaspiro[4.4]nonane-7-carboxamide	419.5	421
19		N-[4-Methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-benzamide	365.42	366
20		4-Methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-ylamine	261.31	262

TABLE 2-continued

examples of compounds of the present invention				
No.	Structure	IUPAC-Name	MW	[M + H] ⁺ 1
21		4-Hydroxy-4-methylpiperidine-1-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	402.48	403
22		4-Dimethylaminomethyl-N-[4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-benzamide	422.51	424
23		4-Methoxymethyl-N-[4-methoxy-7-(tetrahydropyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-benzamide	413.50	414
24		2-Dimethylaminomethyl-N-[4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-isonicotinamide	423.50	424

TABLE 2-continued






examples of compounds of the present invention				
No.	Structure	IUPAC-Name	MW	[M + H] ⁺ 1
25		2-Dimethylaminomethyl-N-[4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-isonicotinamide	427.53	429
26		2-Methoxymethyl-N-[4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-isonicotinamide	410.46	411
27		4-Dimethylaminomethyl-N-[4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-benzamide	426.54	428
28		2-Methoxymethyl-N-[4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-isonicotinamide	414.48	415
29		1-(2-Methoxy-ethyl)-1H-pyrazole-4-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	417.49	418

TABLE 2-continued

examples of compounds of the present invention				
No.	Structure	IUPAC-Name	MW	[M + H] ⁺ 1
30		4-Hydroxy-4-methyl-piperidine-1-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	406.50	408
31		Isoxazole-4-carboxylic acid (4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide	361.38	362
32		4-Hydroxy-4-methyl-piperidine-1-carboxylic acid (4-fluoromethoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide	425.48	426
33		4-Imidazol-1-ylmethyl-N-(4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-benzamide	450.51	452
34		7-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid (4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide	433.52	435

TABLE 2-continued

examples of compounds of the present invention				
No.	Structure	IUPAC-Name	MW	[M + H] ⁺ 1
35		1-(2-Methoxy-ethyl)-1H-pyrazole-4-carboxylic acid (4-methoxy-7-pyridin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide	410.45	411
36		1-(2-Methoxy-ethyl)-1H-pyrazole-4-carboxylic acid [4-methoxy-7-(6-methyl-pyridazin-3-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	425.46	426
37		4-Hydroxy-4-methyl-piperidine-1-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide	398.48	399
38		4-Difluoromethyl-4-hydroxy-piperidine-1-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide	434.46	435
39		4-Hydroxymethyl-4-methyl-piperidine-1-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide	412.51	414

TABLE 2-continued

examples of compounds of the present invention				
No.	Structure	IUPAC-Name	MW	[M + H] ⁺ 1
40		4-Fluoromethyl-4-hydroxy-piperidine-1-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide	416.47	417
41		3-Oxa-9-aza-spiro[5.5]undecane-9-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide	438.54	440
42		4-Methyl-piperidine-1-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide	382.48	383
43		4-Benzyl-4-hydroxy-piperidine-1-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide	474.58	476
44		2-Oxo-1-oxa-3,8-diazaspiro[4.5]decane-8-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide	439.49	440

TABLE 2-continued

examples of compounds of the present invention				
No.	Structure	IUPAC-Name	MW	[M + H] ⁺ 1
45		3-Oxo-2,8-diazaspiro[4.5]decane-8-carboxylic acid (4-methoxy-7-phenylthiazolo[4,5-c]pyridin-2-yl)-amide	437.52	439
46		4-Dimethylaminomethyl-N-(4-methoxy-7-phenylthiazolo[4,5-c]pyridin-2-yl)-benzamide	418.51	420
47		4-Methoxymethyl-N-(4-methoxy-7-phenylthiazolo[4,5-c]pyridin-2-yl)-benzamide	405.47	406
48		2,4-Dioxo-1,3,8-triazaspiro[4.5]decane-8-carboxylic acid (4-methoxy-7-phenylthiazolo[4,5-c]pyridin-2-yl)-amide	452.49	453
49		4'-Hydroxy-3',4',5',6'-tetrahydro-2H-[3,4]bipyridinyl-1'-carboxylic acid (4-methoxy-7-phenylthiazolo[4,5-c]pyridin-2-yl)-amide	461.54	463

TABLE 2-continued

examples of compounds of the present invention				
No.	Structure	IUPAC-Name	MW	[M + H] ⁺ 1
50		4-Oxo-piperidine-1-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide	382.44	383
51		1-Oxo-2,8-diazaspiro[4.5]decane-8-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide	437.52	439
52		7-Oxa-2-azaspiro[4.5]decane-2-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide	424.52	426
53		8-Oxa-2-azaspiro[4.5]decane-2-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide	424.52	426
54		2-Oxa-7-azaspiro[4.4]nonane-7-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide	410.49	411

TABLE 2-continued






examples of compounds of the present invention				
No.	Structure	IUPAC-Name	MW	[M + H] ⁺ 1
55		4-Imidazol-1-ylmethyl-N-(4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-benzamide	441.51	443
56		Isoxazole-3-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide	352.37	353
57		4-Hydroxy-4-prop-2-ynyl-piperidine-1-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide	422.5	424
58		N-(4-Methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-N',N'-dimethyl-terephthalamide	432.5	434
59		N-(4-Methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-4-trifluoromethoxy-benzamide	445.42	446

TABLE 2-continued

examples of compounds of the present invention				
No.	Structure	IUPAC-Name	MW	[M + H] ⁺ 1
60		2-Methyl-oxazole-4-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide	366.39	367
61		Benzooxazole-5-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide	402.43	403
62		N-(4-Methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-4-(2-oxo-pyrrolidin-1-ylmethyl)-benzamide	458.53	460
63		2,3-Dihydro-benzofuran-5-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide	403.45	404
64		3-Methoxymethyl-pyrrolidine-1-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide	398.48	399

TABLE 2-continued






examples of compounds of the present invention				
No.	Structure	IUPAC-Name	MW	[M + H] ⁺ 1
65		Piperidine-1,4-dicarboxylic acid 4-amide 1-[(4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide]	411.48	412
66		4-Diethylamino-N-(4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-benzamide	432.54	434
67		4-Difluoromethyl-4-hydroxy-piperidine-1-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	438.45	439
68		4-Hydroxymethyl-4-methyl-piperidine-1-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	416.5	418
69		4-Fluoromethyl-4-hydroxy-piperidine-1-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	420.46	421

TABLE 2-continued

examples of compounds of the present invention				
No.	Structure	IUPAC-Name	MW	[M + H] ⁺ 1
70		3-Oxa-9-aza-spiro[5.5]undecane-9-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	442.54	444
71		4-Methyl-piperidine-1-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	386.47	387
72		4-Benzyl-4-hydroxy-piperidine-1-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	478.57	480
73		2-Oxo-1-oxa-3,8-diazaspiro[4.5]decane-8-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	443.48	444
74		3-Oxo-2,8-diazaspiro[4.5]decane-8-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	441.51	443

TABLE 2-continued






examples of compounds of the present invention				
No.	Structure	IUPAC-Name	MW	[M + H] ⁺ 1
75		4-Dimethylaminomethyl-N-[4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-benzamide	422.5	424
76		2,4-Dioxo-1,3,8-triazaspiro[4.5]decane-8-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	456.48	457
77		4'-Hydroxy-3',4',5',6'-tetrahydro-2'H-[3,4]bipyridinyl-1'-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	465.53	467
78		4-Oxo-piperidine-1-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	386.43	387
79		1-Oxo-2,8-diazaspiro[4.5]decane-8-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	441.51	443

TABLE 2-continued

examples of compounds of the present invention				
No.	Structure	IUPAC-Name	MW	[M + H] ⁺ 1
80		7-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	428.51	430
81		8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	428.51	430
82		2-Oxa-7-aza-spiro[4.4]nonane-7-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	414.48	415
83		4-Imidazol-1-ylmethyl-N-[4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-benzamide	445.5	447

TABLE 2-continued

examples of compounds of the present invention				
No.	Structure	IUPAC-Name	MW	[M + H] ⁺ 1
84		Isoxazole-3-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	356.36	357
85		4-Hydroxy-4-prop-2-ynyl-piperidine-1-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	426.49	427
86		N-[4-Methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-N',N'-dimethyl-terephthalamide	436.49	437
87		N-[4-Methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-4-trifluoromethoxy-benzamide	449.41	450
88		2-Methyl-oxazole-4-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	370.39	371

TABLE 2-continued

examples of compounds of the present invention				
No.	Structure	IUPAC-Name	MW	[M + H] ⁺ 1
89		Benzooxazole-5-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	406.42	407
90		N-[4-Methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-4-(2-oxo-pyrrolidin-1-ylmethyl)-benzamide	462.53	464
91		2,3-Dihydro-benzofuran-5-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	407.45	408
92		3-Methoxymethyl-pyrrolidine-1-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	402.47	403
93		Piperidine-1,4-dicarboxylic acid 4-amide 1-{[4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide}	415.47	416

TABLE 2-continued

examples of compounds of the present invention				
No.	Structure	IUPAC-Name	MW	[M + H] ⁺ 1
94		4-Diethylamino-N-[4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-benzamide	436.53	438
95		4-Difluoromethyl-4-hydroxy-piperidine-1-carboxylic acid [4-methoxy-7-(tetrahydropyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	442.48	443
96		4-Hydroxymethyl-4-methyl-piperidine-1-carboxylic acid [4-methoxy-7-(tetrahydropyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	420.53	422
97		4-Fluoromethyl-4-hydroxy-piperidine-1-carboxylic acid [4-methoxy-7-(tetrahydropyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	424.49	425
98		3-Oxa-9-aza-spiro[5.5]undecane-9-carboxylic acid [4-methoxy-7-(tetrahydropyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	446.56	448

TABLE 2-continued

examples of compounds of the present invention				
No.	Structure	IUPAC-Name	MW	[M + H] ⁺ 1
99		4-Methyl-piperidine-1-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	390.5	392
100		4-Benzyl-4-hydroxy-piperidine-1-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	482.6	484
101		2-Oxo-1-oxa-3,8-diazaspiro[4.5]decane-8-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	447.51	449
102		3-Oxo-2,8-diazaspiro[4.5]decane-8-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	445.54	447
103		2,4-Dioxo-1,3,8-triazaspiro[4.5]decane-8-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	460.51	462

TABLE 2-continued






examples of compounds of the present invention				
No.	Structure	IUPAC-Name	MW	[M + H] ⁺ 1
104		4'-Hydroxy-3',4',5',6'-tetrahydro-2'H-[3,4]bipyridinyl-1'-carboxylic acid [4-methoxy-7-(tetrahydropyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	469.56	471
105		4-Oxo-piperidine-1-carboxylic acid [4-methoxy-7-(tetrahydropyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	390.46	391
106		1-Oxo-2,8-diazaspiro[4.5]decane-8-carboxylic acid [4-methoxy-7-(tetrahydropyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	445.54	447
107		7-Oxa-2-azaspiro[4.5]decane-2-carboxylic acid [4-methoxy-7-(tetrahydropyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	432.54	434
108		8-Oxa-2-azaspiro[4.5]decane-2-carboxylic acid [4-methoxy-7-(tetrahydropyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	432.54	434

TABLE 2-continued

examples of compounds of the present invention				
No.	Structure	IUPAC-Name	MW	[M + H] ⁺ 1
109		2-Oxa-7-aza-spiro[4.4]nonane-7-carboxylic acid [4-methoxy-7-(tetrahydropyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	418.51	420
110		4-Imidazol-1-ylmethyl-N-[4-methoxy-7-(tetrahydropyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-benzamide	449.53	451
111		Isoxazole-3-carboxylic acid [4-methoxy-7-(tetrahydropyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	360.39	361
112		4-Hydroxy-4-prop-2-ynyl-piperidine-1-carboxylic acid [4-methoxy-7-(tetrahydropyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	430.52	432
113		N-[4-Methoxy-7-(tetrahydropyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-N,N'-dimethyl-terephthalamide	440.52	442

TABLE 2-continued






examples of compounds of the present invention				
No.	Structure	IUPAC-Name	MW	[M + H] ⁺ 1
114		N-[4-Methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-4-trifluoromethoxy-benzamide	453.44	454
115		2-Methyl-oxazole-4-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	374.42	375
116		Benzooxazole-5-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	410.45	411
117		N-[4-Methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-4-(2-oxo-pyrrolidin-1-ylmethyl)-benzamide	466.55	468
118		2,3-Dihydro-benzofuran-5-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	411.48	412

TABLE 2-continued

examples of compounds of the present invention				
No.	Structure	IUPAC-Name	MW	[M + H] ⁺ 1
119		3-Methoxymethyl-pyrrolidine-1-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	406.5	408
120		Piperidine-1,4-dicarboxylic acid 4-amide 1-{[4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide}	419.5	421
121		4-Diethylamino-N-[4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-benzamide	440.56	442
122		{1-[4-Methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]carbamoyl}-piperidin-3-yl}-acetic acid	430.49	431
123		Pyridine-2,5-dicarboxylic acid 2-dimethylamide 5-{[4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide}	437.48	438

TABLE 2-continued

examples of compounds of the present invention				
No.	Structure	IUPAC-Name	MW	[M + H] ⁺ 1
124		1-Methyl-1H-pyrazole-4-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	369.41	370
125		5-Methyl-isoxazole-4-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	370.39	371
126		1-(2-Methoxy-ethyl)-1H-[1,2,3]triazole-4-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	414.45	415
127		1-Methyl-1H-[1,2,3]triazole-4-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	370.39	371
128		1-Cyano-cyclopropanecarboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	354.39	355

TABLE 2-continued

examples of compounds of the present invention				
No.	Structure	IUPAC-Name	MW	[M + H] ⁺ 1
129		3-Cyano-N-[4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-propionamide	342.38	343
130		2-Methyl-oxazole-5-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	370.39	371
131		2-Methyl-thiazole-5-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	386.46	387
132		4-Hydroxy-pent-2-ynoic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	357.39	358

TABLE 2-continued

examples of compounds of the present invention				
No.	Structure	IUPAC-Name	MW	[M + H] ⁺ 1
133		(S)-3-Methanesulfonyl-pyrrolidine-1-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	436.51	438
134		(S)-3-Fluoro-pyrrolidine-1-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	376.41	377
135		(S)-3-Cyano-pyrrolidine-1-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	383.43	384
136		(R)-3-Dimethylaminomethyl-pyrrolidine-1-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	415.52	417

TABLE 2-continued






examples of compounds of the present invention				
No.	Structure	IUPAC-Name	MW	[M + H] ⁺ 1
137		5-Methyl-isoxazole-4-carboxylic acid (4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide	375.41	376
138		1-(2-Methoxy-ethyl)-1H-[1,2,3]triazole-4-carboxylic acid (4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide	419.46	420
139		1-Methyl-1H-[1,2,3]triazole-4-carboxylic acid (4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide	375.41	376
140		Pyridine-2,5-dicarboxylic acid 2-dimethylamide 5-[(4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl)-amide]	441.51	443
141		Pyridine-2,5-dicarboxylic acid 2-dimethylamide 5-[(4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide]	442.5	444

TABLE 2-continued






examples of compounds of the present invention				
No.	Structure	IUPAC-Name	MW	[M + H] ⁺ 1
142		1-Methyl-1H-pyrazole-4-carboxylic acid [7-methoxy-4-(tetrahydro-pyran-4-yl)-1H-benzimidazol-2-yl]-amide	355.4	356
143		5-Methyl-isoxazole-4-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	374.42	375
144		1-(2-Methoxy-ethyl)-1H-1,2,3-triazole-4-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	418,48	419
145		1-Methyl-1H-1,2,3-triazole-4-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	374,42	375
146		1-Cyano-cyclopropanecarboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	358.42	359

TABLE 2-continued

examples of compounds of the present invention				
No.	Structure	IUPAC-Name	MW	[M + H] ⁺ 1
147		2-Methyl-oxazole-5-carboxylic acid [4-methoxy-7-(tetrahydropyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	374.42	375
148		2-Methyl-thiazole-5-carboxylic acid [4-methoxy-7-(tetrahydropyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	390.48	391
149		(S)-3-Methanesulfonyl-pyrrolidine-1-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	440.54	442
150		(S)-3-Fluoro-pyrrolidine-1-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	380.44	381
151		(S)-3-Cyano-pyrrolidine-1-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	387.46	388

TABLE 2-continued





examples of compounds of the present invention				
No.	Structure	IUPAC-Name	MW	[M + H] ⁺ 1
152		(R)-3-Dimethylaminomethylpyrrolidine-1-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	419.55	421
153		1-[4-Methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-3-thiazol-2-ylmethyl-urea	405.5	407
154		1-(2-Methoxy-ethyl)-1H-pyrazole-4-carboxylic acid [4-methoxy-7-(2-oxa-7-aza-spiro[4.4]non-7-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	458.54	460
155		1-Methyl-1H-pyrazole-4-carboxylic acid [4-methoxy-7-[(2-methoxypropyl)-methyl-amino]-thiazolo[4,5-c]pyridin-2-yl]-amide	390.47	391

TABLE 2-continued

examples of compounds of the present invention				
No.	Structure	IUPAC-Name	MW	[M + H] ⁺ 1
156		1-Methyl-1H-pyrazole-4-carboxylic acid [4-methoxy-7-(5-oxa-2-aza-spiro[3.4]oct-2-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	400.46	401
157		1-Methyl-1H-pyrazole-4-carboxylic acid (4-methoxy-7-piperidin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide	372.45	373
158		1-Methyl-1H-pyrazole-4-carboxylic acid (4-methoxy-7-piperidin-3-yl-thiazolo[4,5-c]pyridin-2-yl)-amide	372.45	373
159		1-Methyl-1H-pyrazole-4-carboxylic acid [7-(carbamoylmethyl-methyl-amino)-4-methoxy-thiazolo[4,5-c]pyridin-2-yl]-amide	375.41	376
160		1-Methyl-1H-pyrazole-4-carboxylic acid [4-methoxy-7-(2,2,2-trifluoro-ethoxy)-thiazolo[4,5-c]pyridin-2-yl]-amide	387.34	388

TABLE 2-continued

examples of compounds of the present invention				
No.	Structure	IUPAC-Name	MW	[M + H] ⁺ 1
161		4-Hydroxy-4-methyl-piperidine-1-carboxylic acid (4-fluoromethoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide	425.48	426
162		4-Hydroxy-4-methyl-piperidine-1-carboxylic acid (4-difluoromethoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide	443.47	444
163		N-(4-methoxy-6-methyl-7-morpholino-thiazolo[4,5-c]pyridin-2-yl)-1-methyl-pyrazole-4-carboxamide	388.44	389
164		N-(6-bromo-4-methoxy-7-morpholino-thiazolo[4,5-c]pyridin-2-yl)-1-methyl-pyrazole-4-carboxamide	453.31	454

TABLE 2-continued

examples of compounds of the present invention				
No.	Structure	IUPAC-Name	MW	[M + H] ⁺ 1
165		1-Methyl-1H-pyrazole-4-carboxylic acid (6-fluoro-4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide	392.41	393
166		N-(6-chloro-4-methoxy-7-morpholino-thiazolo[4,5-c]pyridin-2-yl)-1-methyl-pyrazole-4-carboxamide	408.86	410
167		Cyclopropanecarboxylic acid (4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide	334.40	335
168		(S)-7-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid (4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide	433.53	435
169		(R)-7-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid (4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide	433.53	435

TABLE 2-continued

examples of compounds of the present invention				
No.	Structure	IUPAC-Name	MW	[M + H] ⁺ 1
170		N-(4-Methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-4-(2-oxo-pyrrolidin-1-ylmethyl)-benzamide	467.55	469
171		1-Methyl-1H-pyrazole-4-carboxylic acid (4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide	374.42	375
172		(R)-2-Oxa-7-aza-spiro[4.4]nonane-7-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	414.49	415
173		(S)-2-Oxa-7-aza-spiro[4.4]nonane-7-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	414.49	415
174		3-cyano-N-[4-methoxy-7-(morpholin-4-yl)-[1,3]thiazolo[4,5-c]pyridin-2-yl]propanamide	347.40	348

TABLE 2-continued

examples of compounds of the present invention				
No.	Structure	IUPAC-Name	MW	[M + H] ⁺ 1
175		1-cyano-N-[4-methoxy-7-(morpholin-4-yl)-[1,3]thiazolo[4,5-c]pyridin-2-yl]cyclopropane-1-carboxamide	359.41	360
176		(S)-7-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)thiazolo[4,5-c]pyridin-2-yl]-amide	428.51	430
177		(R)-7-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)thiazolo[4,5-c]pyridin-2-yl]-amide	428.51	430
178		1-Methyl-1H-pyrazole-4-carboxylic acid (6-bromo-4-methoxy-7-morpholin-4-ylthiazolo[4,5-c]pyridin-2-yl)-amide	453.32	454
179		Cyclopropanecarboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)thiazolo[4,5-c]pyridin-2-yl]-amide	329.38	330

TABLE 2-continued

examples of compounds of the present invention				
No.	Structure	IUPAC-Name	MW	[M + H] ⁺ 1
180		N-[4-Methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-4-(2-oxo-pyrrolidin-1-ylmethyl)-benzamide	466.56	468
181		2-Methyl-thiazole-5-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	390.49	391
182		2-Oxa-7-aza-spiro[4.4]nonane-7-carboxylic acid [7-(3-ethoxy-3-methyl-azetidin-1-yl)-4-methoxy-thiazolo[4,5-c]pyridin-2-yl]-amide	447.56	449
183		2-Oxa-7-aza-spiro[4.4]nonane-7-carboxylic acid (4-methoxy-7-piperidin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide	417.53	419

TABLE 2-continued

examples of compounds of the present invention				
No.	Structure	IUPAC-Name	MW	[M + H] ⁺ 1
184		2-Oxa-7-aza-spiro[4.4]nonane-7-carboxylic acid [4-methoxy-7-(5-oxa-2-aza-spiro[3.4]oct-2-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	445.54	447
185		2-Oxa-7-aza-spiro[4.4]nonane-7-carboxylic acid [4-methoxy-7-(3-methoxy-3-methyl-azetidin-1-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	433.53	435
186		2-Methyl-oxazole-5-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide	366.40	367
187		Cyclopropanecarboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide	325.39	326
188		1-Methyl-1H-pyrazole-4-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide	365.42	366

TABLE 2-continued

examples of compounds of the present invention				
No.	Structure	IUPAC-Name	MW	[M + H] ⁺ 1
189		N-[4-methoxy-7-(morpholin-4-yl)-[1,3]thiazolo[4,5-c]pyridin-2-yl]-2-methyl-1,3-oxazole-5-carboxamide	375.41	376
190		2-Methyl-thiazole-5-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide	382.47	383
191		8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid [4-methoxy-7-(5-oxa-2-aza-spiro[3.4]oct-2-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	459.57	461
192		8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid (4-methoxy-7-piperidin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide	431.56	433

TABLE 2-continued

examples of compounds of the present invention				
No.	Structure	IUPAC-Name	MW	[M + H] ⁺ 1
193		8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid [7-(4-fluoro-phenyl)-4-methoxy-thiazolo[4,5-c]pyridin-2-yl]-amide	442.51	444
194		8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid [7-(3-fluoro-phenyl)-4-methoxy-thiazolo[4,5-c]pyridin-2-yl]-amide	442.51	444
195		8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid [7-(2-fluoro-phenyl)-4-methoxy-thiazolo[4,5-c]pyridin-2-yl]-amide	442.51	444
196		1-Dimethylaminomethyl-cyclopropanecarboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide	382.49	383
197		Cyclopropane-1,1-dicarboxylic acid dimethylamide (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide	396.47	397

TABLE 2-continued

examples of compounds of the present invention				
No.	Structure	IUPAC-Name	MW	[M + H] ⁺ 1
198		1-Imidazol-1-ylmethyl-cyclopropanecarboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide	405.48	406
199		1-Methyl-1H-pyrazole-4-carboxylic acid (4-hydroxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide	360.40	361
200		8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid (4-methoxy-7-piperidin-3-yl-thiazolo[4,5-c]pyridin-2-yl)-amide	431.56	433
201		1-amino-N-{4-methoxy-7-phenyl-[1,3]thiazolo[4,5-c]pyridin-2-yl}-8-azaspiro[4.5]decane-8-carboxamide	437.57	439
202		(1S,2S)-2-methoxy-N-{4-methoxy-7-phenyl-[1,3]thiazolo[4,5-c]pyridin-2-yl}-8-cyclopropane-1-carboxamide	355.42	356

TABLE 2-continued

examples of compounds of the present invention				
No.	Structure	IUPAC-Name	MW	[M + H] ⁺ 1
203		1-Amino-cyclopropanecarboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide	340.41	341
204		(S)-7-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide	424.52	426
205		(R)-7-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide	424.52	426
206		8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid (4-methoxy-7-pyridin-3-yl-thiazolo[4,5-c]pyridin-2-yl)-amide	425.51	427
207		(S)-2-Oxa-7-aza-spiro[4.4]nonane-7-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide	410.50	411

TABLE 2-continued






examples of compounds of the present invention				
No.	Structure	IUPAC-Name	MW	[M + H] ⁺ 1
208		(R)-2-Oxa-7-aza-spiro[4.4]nonane-7-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide	410.50	411
209		8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid [7-(2,3-difluoro-phenyl)-4-methoxy-thiazolo[4,5-c]pyridin-2-yl]-amide	460.50	462
210		8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid [7-(2,5-difluoro-phenyl)-4-methoxy-thiazolo[4,5-c]pyridin-2-yl]-amide	460.50	462
211		8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid (4-methoxy-7-pyridin-2-yl-thiazolo[4,5-c]pyridin-2-yl)-amide	425.51	427
212		8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid (4-methoxy-7-pyridin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide	425.51	427

TABLE 2-continued

examples of compounds of the present invention					
No.	Structure	IUPAC-Name	MW	[M + H] ⁺	
213		4-(2,5-Dioxo-pyrrolidin-1-yl)-N-(4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-benzamide	458.50	459	
214		4-(2,5-Dioxo-pyrrolidin-1-yl)-N-(4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-benzamide	467.50	469	
215		(R)-1-Amino-8-azaspiro[4.5]decane-8-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide	437.57	439	
216		(S)-1-Amino-8-azaspiro[4.5]decane-8-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide	437.57	439	
217		1-Methyl-1H-pyrazole-4-carboxylic acid (6-cyano-4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide	399.43	400	

TABLE 2-continued

examples of compounds of the present invention				
No.	Structure	IUPAC-Name	MW	[M + H] ⁺ 1
218		N-(4-Methoxy-7-pyridin-3-yl-thiazolo[4,5-c]pyridin-2-yl)-N',N'-dimethyl-terephthalamide	433.49	434
219		N-(4-Methoxy-7-pyridin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-N',N'-dimethyl-terephthalamide	433.49	434
220		8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid (6-chloro-4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide	458.97	460
221		1-Methyl-1H-pyrazole-4-carboxylic acid (6-chloro-4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide	408.87	410
222		8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid (6-cyano-4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide	449.53	451

TABLE 2-continued

examples of compounds of the present invention				
No.	Structure	IUPAC-Name	MW	[M + H] ⁺ 1
223		8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid (4-methoxy-6-methyl-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide	438.55	440
224		N-{4-Methoxy-7-[3-(2-methoxy-ethoxy)-phenyl]-thiazolo[4,5-c]pyridin-2-yl}-N',N'-dimethyl-terephthalamide	506.58	508
225		8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid {4-methoxy-743-(2-methoxy-ethoxy)-phenyl]-thiazolo[4,5-c]pyridin-2-yl}-amide	498.60	500
226		4-(2,5-Dioxo-pyrrolidin-1-yl)-N-(6-fluoro-4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-benzamide	485.49	486
227		N-(2-Hydroxy-ethyl)-N'-(4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-N-methyl-terephthalamide	462.53	464

TABLE 2-continued

examples of compounds of the present invention				
No.	Structure	IUPAC-Name	MW	[M + H] ⁺ 1
228		N-(2-Hydroxy-ethyl)-N'-(4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-terephthalamide	448.50	450
229		8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid (6-fluoro-4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide	442.51	444
230		N-(2-Dimethylamino-ethyl)-N'-(4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-terephthalamide	475.57	477
231		8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid (4,6-dimethyl-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide	422.55	424
232		8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid [4-methoxy-7-(tetrahydropyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide	432.54	434

TABLE 2-continued

examples of compounds of the present invention				
No.	Structure	IUPAC-Name	MW	[M + H] ⁺ 1
233		N-{4-methoxy-7-[3-(2-methoxyethoxy)phenyl]-[1,3]thiazolo[4,5-c]pyridin-2-yl]-1-(2-methoxyethyl)-1H-pyrazole-4-carboxamide	483.55	485
234		N-{6-fluoro-4-methoxy-7-[3-(2-methoxyethoxy)phenyl]-[1,3]thiazolo[4,5-c]pyridin-2-yl]-8-oxa-2-azaspiro[4.5]decane-2-carboxamide	516.59	518
235		N-[4-methoxy-7-(oxan-4-yl)-[1,3]thiazolo[4,5-c]pyridin-2-yl]-1-methyl-1H-pyrazole-4-carboxamide	373.44	374
236		N-[4-methoxy-7-(2-methoxyphenyl)-[1,3]thiazolo[4,5-c]pyridin-2-yl]-8-oxa-2-azaspiro[4.5]decane-2-carboxamide	454.55	456
237		8-Oxa-2-azaspiro[4.5]decane-2-carboxylic acid [7-(3,6-dihydro-2H-pyran-4-yl)-4-methoxy-thiazolo[4,5-c]pyridin-2-yl]-amide	430.53	432

TABLE 2-continued






examples of compounds of the present invention				
No.	Structure	IUPAC-Name	MW	[M + H] ⁺ 1
238		N-[7-(3,6-dihydro-2H-pyran-4-yl)-4-methoxy-[1,3]thiazolo[4,5-c]pyridin-2-yl]-1-methyl-1H-pyrazole-4-carboxamide	371.42	372
239		N-[7-(2,6-dimethoxyphenyl)-4-methoxy-[1,3]thiazolo[4,5-c]pyridin-2-yl]-8-oxa-2-azaspiro[4.5]decane-2-carboxamide	485.56	487
240		8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid (6-fluoro-4-hydroxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide	428.49	429
241		2-amino-N-[4-methoxy-7-(oxan-4-yl)-[1,3]thiazolo[4,5-c]pyridin-2-yl]-1,3-thiazole-5-carboxamide	391.47	392
242		3-Amino-pyrrolidine-1-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide	369.45	370

TABLE 2-continued

examples of compounds of the present invention				
No.	Structure	IUPAC-Name	MW	[M + H] ⁺ 1
243		(R)-3-Amino-pyrrolidine-1-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide	369.45	370
244		(S)-3-Amino-pyrrolidine-1-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide	369.45	370
245		8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid (7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide	394.5	395
246		1-(2-Methoxy-ethyl)-1H-pyrazole-4-carboxylic acid (6-fluoro-4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide	463.47	437
247		Bicyclo[1.1.1]pentane-1,3-dicarboxylic acid dimethylamide (4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide	431.51	433

TABLE 2-continued

examples of compounds of the present invention				
No.	Structure	IUPAC-Name	MW	[M + H] ⁺ 1
248		N-(2-Hydroxy-ethyl)-N-[4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-N-methyl-terephthalamide	470.55	472

TABLE 3

NMR profiles of the compounds of the present invention
The Nos. recited herein corresponds to the numbering
of the compounds, disclosed in table 2.

No. NMR

- 1H NMR (300 MHz, d6-DMSO) ppm = 11.41 (s, 1H), 7.57 (s, 1H), 4.41 (s, 1H), 3.94 (s, 3H), 3.86 – 3.83 (m, 2H), 3.79 – 3.77 (m, 4H), 3.34 – 3.25 (m, 2H), 3.08 – 3.06 (m, 4H), 1.50 – 1.40 (m, 4H), 1.14 (s, 3H).
- 1H NMR (300 MHz, d6-DMSO) ppm = 13.10 (s, 1H), 8.14 (d, J = 8.1, 2H), 7.67 (s, 1H), 7.50 (d, J = 8.4, 2H), 3.98(s, 3H), 3.82 – 3.79 (m, 4H), 3.49 (s, 2H), 3.13 – 3.10 (m, 4H), 2.18 (s, 6H).
- 1H NMR (300 MHz, d6-DMSO) ppm = 13.23(s, 1H), 8.16(d, J = 6, 2H), 7.68(s, 1H), 7.52(d, J = 6, 2H), 4.52 – 4.50(m, 2H), 3.99(s, 3H), 3.83 – 3.80(m, 4H), 3.34 – 3.31(m, 3H), 3.13 – 3.11(m, 4H).
- 1H NMR (400 MHz, d6-DMSO) ppm = 8.48(s, 1H), 8.18(s, 1H), 7.62(s, 1H), 4.35(t, J = 5.2, 2H), 3.97(s, 3H), 3.81 – 3.79(m, 4H), 3.72(t, J = 5.2, 2H), 3.25(s, 3H), 3.11 – 3.09(m, 4H).
- 1H NMR (400 MHz, d6-DMSO) ppm = 12.85(s, 1H), 8.74(d, J = 5.2, 1H), 8.09(s, 1H), 7.94 – 7.93(m, 1H), 7.64(s, 1H), 3.97(s, 3H), 3.82 – 3.80(m, 6H), 3.12 – 3.10(m, 4H), 2.36(s, 6H).
- 1H NMR (400 MHz, d6-DMSO) ppm = 13.6(s, 1H), 8.78(d, J = 5.2, 1H), 8.07(s, 1H), 7.96 – 7.94(m, 1H), 7.70(s, 1H), 4.61(s, 2H), 3.99(s, 3H), 3.82 – 3.80(m, 4H), 3.43(s, 3H), 3.13 – 3.11(m, 4H).
- 1H NMR (500 MHz, DMSO-d6) d 11.54 – 11.25 (m, 1H), 7.56 (s, 1H), 7.28 – 7.17 (m, 5H), 3.97 – 3.90 (m, 5H), 3.79 – 3.75 (m, 4H), 3.23 – 3.13 (m, 2H), 3.09 – 3.04 (m, 4H), 2.71 (s, 2H), 1.49 – 1.37 (m, 4H).
- 1H NMR (500 MHz, DMSO-d6) d 12.82 – 12.74 (m, 1H), 8.95 (d, J = 1.3 Hz, 1H), 8.47 (d, J = 1.3 Hz, 1H), 8.43 – 8.28 (m, 1H), 7.70 (s, 1H), 4.58 – 4.54 (m, 2H), 3.99 (s, 3H), 3.83 – 3.79 (m, 4H), 3.74 – 3.72 (m, 2H), 3.32 (s, 3H), 3.14 – 3.10 (m, 4H).
- 1H NMR (400 MHz, DMSO-d6) ppm = 10.4 (s, 1H), 7.54 (s, 1H), 3.93 (s, 3H), 3.82 – 3.74 (m, 6H), 3.57 – 3.46 (m, 6H), 3.07 (t, J = 4.4, 4H), 1.91 – 1.81 (m, 4H).
- 1H NMR (400 MHz, DMSO-d6) ppm = 11.2 (s, 1H), 7.57(s, 1H), 4.88(s, 1H), 3.94 (s, 3H), 3.79 – 3.77 (m, 4H), 3.58 – 3.21 (m, 4H), 3.08 (t, J = 4.8, 4H), 1.86 – 1.82 (m, 2H), 1.30(s, 3H).
- 1H NMR (400 MHz, DMSO-d6) ppm = 11.3 (s, 1H), 7.57 (s, 1H), 3.94(s, 3H), 3.79 – 3.77 (m, 4H), 3.65 – 3.50 (m, 8H), 3.08 (t, J = 4.8, 4H), 1.85 – 1.83 (m, 2H), 1.52 – 1.49(m, 4H).
- 1H NMR (400 MHz, DMSO-d6) 11.50 (s, 1H), 7.58 (s, 1H), 4.71 (s, 1H), 3.99 (d, J = 13.7 Hz, 2H), 3.95 (s, 3H), 3.81 – 3.75 (m, 4H), 3.19 (t, J = 12.1 Hz, 2H), 3.10 – 3.05 (m, 4H), 2.84 (t, J = 2.6 Hz, 1H), 2.32 (d, J = 2.7 Hz, 2H), 1.71 – 1.45 (m, 4H).
- 1H NMR (400 MHz, DMSO-d6) ppm = 8.49 (s, 1H), 8.19 – 8.16 (m, 3H), 7.91 (s, 1H), 4.36 (t, J = 5.2, 2H), 4.02 (s, 3H), 3.95 (s, 3H), 3.72 (t, J = 5.2, 2H), 3.25 (s, 3H).
- 1H NMR (400 MHz, DMSO-d6) ppm = 13.2 (s, 1H), 8.2 – 8.1 (m, 4H), 7.94 (s, 1H), 7.51(t, J = 8.0, 2H), 4.52 (s, 2H), 4.04 (s, 3H), 3.97 (s, 3H), 3.35 (s, 3H).
- 1H NMR (400 MHz, DMSO, ppm): 13.26(s, 1H), 8.20(d, J = 8.0, 2H), 7.62(s, 1H), 7.56(d, J = 8.0, 2H), 3.97(s, 3H), 3.82 – 3.80(m, 4H), 3.12 – 3.10(m, 4H), 3.01 – 2.87(m, 6H).
- 1H NMR (400 MHz, DMSO, ppm): 11(s, 1H), 7.57(s, 1H), 4.58(t, J = 5.2, 1H), 3.94(s, 3H), 3.79 – 3.77(m, 6H), 3.30 – 3.26(m, 2H), 3.18 – 3.16(m, 2H), 3.08 – 3.06(m, 4H), 1.48 – 1.41(m, 2H), 1.23 – 1.19(m, 2H), 0.91(s, 3H).
- 1H NMR (400 MHz, DMSO, ppm): 11.3(s, 1H), 7.57(s, 1H), 3.94(s, 3H), 3.80 – 3.78(m, 6H), 3.58 – 3.52(m, 6H), 3.08(t, J = 4.4, 4H), 1.91 – 1.83(m, 4H).

TABLE 3-continued

NMR profiles of the compounds of the present invention
The Nos. recited herein corresponds to the numbering
of the compounds disclosed in table 2.

No. NMR

- 18 ¹H NMR (400 MHz, DMSO, ppm): 11.3(s, 1H), 7.57(s, 1H), 3.94(s, 3H), 3.80 – 3.76(m, 6H), 3.58 – 3.52(m, 6H), 3.08(t, J = 4.4, 4H), 1.91 – 1.83(m, 4H).
- 21 ¹H NMR (400 MHz, DMSO-d₆) 11.44(s, 1H), 8.14(s, 1H), 8.11(s, 1H), 7.88 (s, 1H), 4.42(s, 1H), 3.99(s, 3H), 3.94(s, 3H), 3.91 – 3.79 (m, 2H), 3.31 – 3.19 (m, 2H), 1.60 – 1.34 (m, 4H), 1.14 (s, 3H).
- 22 ¹H NMR (400 MHz, DMSO-d₆) 13.17 (s, 1H), 8.23 (s, 1H), 8.22 (s, 1H), 8.14 (d, J = 8.2 Hz, 3H), 7.95 (s, 1H), 7.49 (d, J = 8.0 Hz, 2H), 4.04 (s, 3H), 3.97 (s, 3H), 3.50 (s, 2H), 2.18 (s, 6H).
- 23 ¹H NMR (400 MHz, DMSO-d₆) δ 13.20 (s, 1H), 8.15 (d, J = 8.0 Hz, 2H), 7.91 (s, 1H), 7.51 (d, J = 8.0 Hz, 2H), 4.52 (s, 2H), 4.10 – 3.97 (m, 4H), 3.59 – 3.49 (m, 2H), 3.31 (s, 3H), 3.08 – 2.97 (m, 1H), 2.04 – 1.76 (m, 4H).
- 27 ¹H NMR (400 MHz, DMSO-d₆) δ 8.13 (d, J = 8.2 Hz, 2H), 7.91 (s, 1H), 7.50 (d, J = 8.2 Hz, 2H), 4.06 – 3.96 (m, 5H), 3.59 – 3.49 (m, 4H), 3.08 – 2.98 (m, 1H), 2.20 (s, 6H), 2.00 – 1.79 (m, 4H).
- 29 ¹H NMR (300 MHz, DMSO, ppm): 12.75(s, 1H), 8.54(s, 1H), 8.23(s, 1H), 7.88(s, 1H), 4.37 – 4.34(m, 2H), 4.02 – 3.98(m, 5H), 3.73 – 3.70(m, 2H), 3.56 – 3.49(m, 2H), 3.25 – 3.22(m, 3H), 3.00 – 2.94(m, 1H), 1.98 – 1.93(m, 4H).
- 30 ¹H NMR (400 MHz, DMSO-d₆) 11.45 (s, 1H), 7.79 (s, 1H), 4.41 (s, 1H), 4.02 – 3.92 (m, 5H), 3.84 (d, J = 13.2 Hz, 2H), 3.54 – 3.44 (m, 2H), 3.29 – 3.21 (m, 2H), 3.03 – 2.84 (m, 1H), 1.93 – 1.68 (m, 4H), 1.56 – 1.36 (m, 4H), 1.14 (s, 3H).
- 33 ¹H NMR (400 MHz, DMSO, ppm): 13.20(s, 1H), 8.14(d, J = 8.0, 2H), 7.79(s, 1H), 7.66(s, 1H), 7.40(d, J = 8.0, 2H), 7.23(s, 1H), 6.94(s, 1H), 5.31(s, 2H), 3.98(s, 3H), 3.81 – 3.79(m, 4H), 3.12 – 3.10(m, 4H).
- 34 ¹H NMR (300 MHz, DMSO, ppm): 11.3 (s, 1H), 7.56(s, 1H), 3.93 (s, 3H), 3.79 – 3.76(m, 4H), 3.60 – 3.38(m, 7H), 3.29 – 3.05(m, 5H), 1.63 – 1.53(m, 6H).
- 52 ¹H NMR (400 MHz, DMSO-d₆) 11.39 (s, 1H), 8.03 (s, 1H), 7.69 (d, J = 7.3 Hz, 2H), 7.55 (t, J = 7.7 Hz, 2H), 7.45 (t, J = 7.6 Hz, 1H), 4.03 (s, 3H), 3.70 – 3.38 (m, 6H), 3.30 (m, J = 7.8 Hz, 2H), 1.53 (m, 6H).
- 53 ¹H NMR (400 MHz, DMSO-d₆) 11.38 (s, 1H), 8.03 (d, J = 1.1 Hz, 1H), 7.72 – 7.63 (m, 2H), 7.58 – 7.51 (m, 2H), 7.49 – 7.42 (m, 1H), 4.04 (s, 3H), 3.67 – 3.43 (m, 7H), 3.31 – 3.29 (m, 1H), 1.92 – 1.70 (m, 2H), 1.52 – 1.40 (m, 4H).
- 54 ¹H NMR (400 MHz, DMSO-d₆) 11.42 (s, 1H), 8.03 (s, 1H), 7.70 (m, 2H), 7.55 (t, J = 7.6 Hz, 2H), 7.46 (t, J = 7.3 Hz, 1H), 4.04 (s, 3H), 3.85 – 3.72 (m, 2H), 3.55 (m, 6H), 1.88 (m, 4H).
- 58 ¹H NMR (400 MHz, DMSO, ppm): 13.37(s, 1H), 8.20 – 8.13(m, 3H), 7.75 – 7.73(m, 2H), 7.61 – 7.56(m, 4H), 7.50 – 7.46(m, 1H), 4.08(s, 3H), 3.01 – 2.90(m, 6H).
- 62 ¹H NMR (400 MHz, DMSO, ppm): 13.25(s, 1H), 8.14 – 8.12(m, 3H), 7.75 – 7.73(m, 2H), 7.60 – 7.56(m, 2H), 7.50 – 7.46(m, 1H), 7.41 – 7.39(m, 2H), 4.46(s, 2H), 4.08(s, 3H), 3.29 – 3.25(m, 2H), 2.34 – 2.30(m, 2H), 1.99 – 1.92(m, 2H).
- 80 ¹H NMR (400 MHz, DMSO-d₆) 11.38(s, 1H), 8.15(s, 1H), 8.12(s, 1H), 7.88(d, J = 0.8 Hz, 1H), 4.00(s, 3H), 3.94(s, 3H), 3.68 – 3.28(m, 8H), 1.87 – 1.43(m, 6H).
- 81 ¹H NMR (400 MHz, DMSO-d₆) 11.34(s, 1H), 8.14(s, 1H), 8.12(s, 1H), 7.88(s, 1H), 4.00(s, 3H), 3.94(s, 3H), 3.74 – 3.43 (m, 8H), 1.99 – 1.64 (m, 2H), 1.51(s, 4H).
- 82 ¹H NMR (400 MHz, DMSO-d₆) 11.39(s, 1H), 8.13(d, J = 10.0 Hz, 2H), 7.88(s, 1H), 4.00(s, 3H), 3.94(s, 3H), 3.79(td, J = 7.3, 2.6 Hz, 2H), 3.65 – 3.38 (m, 6H), 1.88 (dq, J = 13.4, 7.3, 6.7 Hz, 4H).
- 113 ¹H NMR (400 MHz, DMSO-d₆): 13.31(s, 1H), 8.20(d, J = 8.3, 2H), 7.91(s, 1H), 7.59(d, J = 8.0, 2H), 4.01(s, 5H), 3.54(t, J = 11.4, 2H), 3.0(s, 4H), 2.91(s, 3H), 2.04 – 1.77(m, 4H).
- 130 ¹H NMR (400 MHz, DMSO-d₆) 13.45 (s, 1H), 8.17 (d, 3H), 7.92 (d, J = 0.7 Hz, 1H), 4.03 (s, 3H), 3.95 (s, 3H), 2.55 (s, 3H).
- 131 ¹H NMR (400 MHz, DMSO-d₆) 13.52 (s, 1H), 8.71 (s, 1H), 8.22 (s, 1H), 8.21 (s, 1H), 7.93 (s, 1H), 4.04 (s, 3H), 3.96 (s, 3H), 2.74 (s, 3H).
- 139 ¹H NMR (400 MHz, DMSO, ppm): 13.11(s, 1H), 8.88(m, 1H), 7.68(s, 1H), 4.16(s, 3H), 3.98(s, 3H), 3.82 – 3.79(m, 4H), 3.12 – 3.09(m, 4H).
- 140 ¹H NMR (400 MHz, DMSO-d₆) 13.54 (s, 1H), 9.26 (dd, J = 2.3, 0.9 Hz, 1H), 8.56 (dd, J = 8.2, 2.3 Hz, 1H), 7.91 (s, 1H), 7.73 (d, J = 8.1 Hz, 1H), 4.11 – 3.90 (m, 5H), 3.61 – 3.49 (m, 2H), 3.08 – 3.01 (m, 4H), 2.94 (s, 3H), 2.03 – 1.69 (m, 4H).
- 141 ¹H NMR (400 MHz, DMSO-d₆) 9.25 (d, J = 2.0 Hz, 1H), 8.54 (dd, J = 8.1, 2.2 Hz, 1H), 7.68 (d, J = 8.1 Hz, 1H), 7.59 (s, 1H), 3.96 (s, 3H), 3.83 – 3.78 (m, 4H), 3.15 – 3.08 (m, 4H), 3.04 (s, 3H), 2.94 (s, 3H).
- 147 ¹H NMR (400 MHz, DMSO, ppm): 13.40(s, 1H), 8.20(s, 1H), 7.92(s, 1H), 4.18 – 3.92(m, 5H), 3.55 – 3.50(m, 2H), 3.04 – 2.98(m, 1H), 2.67(s, 3H), 1.95 – 1.87(m, 4H).
- 156 ¹H NMR (400 MHz, DMSO, ppm): 12.87(s, 1H), 8.48(s, 1H), 8.18(s, 1H), 7.17(s, 1H), 4.02 – 4.00(m, 2H), 3.93 – 3.91(m, 6H), 3.87 – 3.85(m, 2H), 3.79 – 3.76(m, 2H), 2.21 – 2.07(m, 2H), 1.94 – 1.88(m, 2H).
- 162 ¹H NMR (400 MHz, DMSO-d₆) 7.92 – 7.53 (m, 2H), 4.47 (d, J = 2.9 Hz, 1H), 3.98 – 3.69 (m, 6H), 3.28 (s, 2H), 3.12 (dd, J = 5.6, 3.5 Hz, 3H), 1.45 (q, J = 14.5, 13.6 Hz, 4H), 1.14 (s, 3H).
- 165 ¹H NMR (400 MHz, DMSO, ppm): 12.86(s, 1H), 8.50(s, 1H), 8.20(s, 1H), 3.97(s, 3H), 3.91(s, 3H), 3.75 – 3.73(m, 4H), 3.10 – 3.07(m, 4H).

TABLE 3-continued

NMR profiles of the compounds of the present invention	
The Nos. recited herein corresponds to the numbering of the compounds disclosed in table 2.	
No. NMR	
167	¹ H NMR (400 MHz, DMSO, ppm): 12.9(s, 1H), 7.63(s, 1H), 3.96(s, 3H), 3.78 – 3.76(m, 4H), 3.08 – 3.05(m, 4H), 1.97 – 1.92(m, 1H), 1.01 – 0.95(m, 4H).
168	¹ H NMR (300 MHz, DMSO, ppm): 11.3(s, 1H), 7.57(s, 1H), 3.93(s, 3H), 3.79 – 3.76(m, 4H), 3.69 – 3.38(m, 7H), 3.29 – 3.18(m, 1H), 3.08 – 3.05(m, 4H), 1.82 – 1.53(m, 6H).
169	¹ H NMR (300 MHz, DMSO, ppm): 11.3(s, 1H), 7.57(s, 1H), 3.94(s, 3H), 3.79 – 3.76(m, 4H), 3.69 – 3.38(m, 7H), 3.29 – 3.12(m, 1H), 3.08 – 3.05(m, 4H), 1.82 – 1.53(m, 6H).
170	¹ H NMR (400 MHz, DMSO, ppm): 13.21(s, 1H), 8.15(d, J = 8.0, 2H), 7.66(s, 1H), 7.41(d, J = 8.4, 2H), 4.46(s, 2H), 3.98(s, 3H), 3.82 – 3.80(m, 4H), 3.31 – 3.26(m, 2H), 3.12 – 3.10(m, 4H), 2.342.30(m, 2H), 1.98 – 1.94(m, 2H). –
171	¹ H NMR (400 MHz, DMSO, ppm): 12.90(s, 1H), 8.49(s, 1H), 8.20(s, 1H), 7.65(s, 1H), 3.97(s, 3H), 3.92 (s, 3H), 3.83 – 3.79(m, 4H), 3.11 – 3.08(m, 4H).
172	¹ H NMR (400 MHz, DMSO-d ₆)11.39(s, 1H), 8.13(d, J = 10.0Hz, 2H), 7.88(s, 1H), 4.00(s, 3H), 3.94(s, 3H), 3.79(td, J = 7.3, 2.6 Hz, 2H), 3.65 – 3.38 (m, 6H), 1.88 (dq, J = 13.4, 7.3, 6.7 Hz, 4H).
173	¹ H NMR (400 MHz, DMSO-d ₆)11.39(s, 1H), 8.13(d, J = 10.0Hz, 2H), 7.88(s, 1H), 4.00(s, 3H), 3.94(s, 3H), 3.79(td, J = 7.3, 2.6 Hz, 2H), 3.65 – 3.38 (m, 6H), 1.88 (dq, J = 13.4, 7.3, 6.7Hz, 4H).
174	¹ H NMR (500 MHz, DMSO-d ₆) d 12.82 (s, 1H), 7.66 (s, 1H), 3.97 (s, 3H), 3.81 – 3.77 (m, 4H), 3.11 – 3.07 (m, 4H), 2.92 – 2.88 (m, 2H), 2.81 – 2.77 (m, 2H).
175	¹ H NMR (700 MHz, DMSO-d ₆) d 13.52 – 12.30 (m, 1H), 7.68 (s, 1H), 3.97 (s, 3H), 3.80 – 3.77 (m, 4H), 3.10 – 3.07 (m, 4H), 1.92 – 1.88 (m, 2H), 1.84 – 1.82 (m, 2H).
176	¹ H NMR (400 MHz, DMSO-d ₆)11.38(s, 1H), 8.15(s, 1H), 8.12(s, 1H), 7.88(d, J = 0.8 Hz, 1H), 4.00(s, 3H), 3.94(s, 3H), 3.68 – 3.28(m, 8H), 1.87 – 1.43(m, 6H).
177	¹ H NMR (400 MHz, DMSO-d ₆) δ 11.39 (s, 1H), 8.17 – 8.10(m, 2H), 7.89 (d, J = 0.9 Hz, 1H), 4.00 (s, 3H), 3.94 (s, 3H), 3.70 – 3.25 (m, 8H), 1.94 – 1.46 (m, 6H).
178	¹ H NMR (400 MHz, DMSO-d ₆)13.01 (s, 1H), 8.48 (s, 1H), 8.19 (s, 1H), 3.99 (s, 3H), 3.92 (s, 3H), 3.82 – 3.74 (m, 4H), 3.19 – 3.06 (m, 4H).
179	¹ H NMR (400 MHz, DMSO-d ₆)12.76 (s, 1H), 8.18 (s, 1H), 8.16 (s, 1H), 7.89 (s, 1H), 4.02 (s, 3H), 3.93 (s, 3H), 2.03 – 1.91 (m, 1H), 1.15 – 0.72 (m, 4H).
180	¹ H NMR (400 MHz, DMSO-d ₆)13.21 (s, 1H), 8.14 (d, J = 8.3 Hz, 2H), 7.91 (s, 1H), 7.41 (d, J = 8.1 Hz, 2H), 4.47 (s, 2H), 4.07 – 3.83 (m, 5H), 3.60 – 3.47 (m, 2H), 3.28 (t, J = 7.0 Hz, 2H), 2.92 – 3.08(m, 1H), 2.33 (t, J = 8.1 Hz, 2H), 2.05 – 1.66 (m, 6H).
181	¹ H NMR (400 MHz, DMSO, ppm): 13.47(s, 1H), 8.71(s, 1 H), 7.91(s, 1H), 4.00 – 3.98(m, 5H), 3.54 – 3.47(m, 2H), 3.03 – 2.97(m, 1H), 2.74(s, 3H), 1.95 – 1.91(m, 2H), 1.89 – 1.86(m, 2H).
182	¹ H NMR (400 MHz, DMSO, ppm): 11.35(s, 1H), 7.11(s, 1 H), 3.89(s, 3H), 3.82 – 3.74(m, 6H), 3.58 – 3.52(m, 4H), 3.45 – 3.40(m, 4H), 1.91 – 1.82(m, 4H), 1.54(s, 3H), 1.13 – 1.10(m, 3H).
183	¹ H NMR (400 MHz, Chloroform-d)7.85 (s, 1H), 4.14 (s, 3H), 4.03 – 3.92 (m, 2H), 3.80 – 3.25 (m, 9H), 2.97 – 2.73 (m, 3H), 2.21 – 1.80 (m, 8H).
184	¹ H NMR (400 MHz, DMSO, ppm): 11.34(s, 1H), 7.11(s, 1 H), 3.98 – 3.89(m, 2H), 3.83(s, 3H), 3.81 – 3.75(m, 6H), 3.58 – 3.40(m, 6H), 2.19 – 2.16(m, 2H), 1.94 – 1.84(m, 6H).
185	¹ H NMR (400 MHz, DMSO, ppm): 11.10(s, 1H), 7.09(s, 1H), 3.89(s, 3H), 3.81 – 3.76(m, 6H), 3.58 – 3.52(m, 3H), 3.40 – 3.37(m, 3H), 3.19(s, 3H), 1.91 – 1.83(m, 4H), 1.52(s, 3H).
186	¹ H NMR (400 MHz, DMSO, ppm): 13.45(s, 1H), 8.12 – 8.10(m, 2H), 7.72 – 7.70(m, 2H), 7.59 – 7.56(m, 2H), 7.49 – 7.45(m, 1H), 4.07(s, 3H), 2.07(s, 3H).
187	¹ H NMR (400 MHz, DMSO, ppm): 12.99(s, 1H), 8.09(s, 1H), 7.70 – 7.67(m, 2H), 7.57 – 7.53(m, 2H), 7.47 – 7.44(m, 1H), 4.06(s, 3H), 2.00 – 1.93(m, 1H), 1.01 – 0.94(m, 4H).
188	¹ H NMR (400 MHz, DMSO, ppm): 12.93(s, 1H), 8.49(s, 1H), 8.20(s, 1H), 8.10(s, 1H), 7.72 – 7.71(m, 2H), 7.59 – 7.55(m, 2H), 7.49 – 7.45(m, 1H), 4.07(s, 3H), 3.91(s, 3H).
189	¹ H NMR (400 MHz, DMSO-d ₆) d 13.56 – 13.05 (m, 1H), 8.19 (s, 1H), 7.69 (s, 1H), 3.99 (s, 3H), 3.82 – 3.78 (m, 4H), 3.12 – 3.08 (m, 4H), 2.55 (s, 3H).
190	¹ H NMR (400 MHz, DMSO, ppm): 13.51(s, 1H), 8.68(s, 1H), 8.12(s, 1H), 7.73 – 7.71 (m, 2H), 7.59 – 7.55(m, 2H), 7.49 – 7.46(m, 1H), 4.07(s, 3H), 2.73(s, 3H).
191	¹ H NMR (300 MHz, DMSO, ppm): 11.28(s, 1H), 7.09(s, 1H), 3.98 – 3.89(m, 2H), 3.83(s, 3H), 3.81 – 3.75(m, 6H), 3.65 – 3.49(m, 6H), 2.20 – 2.15(m, 2H), 1.96 – 1.76(m, 4H), 1.60 – 1.44(m, 4H).
192	¹ H NMR (300 MHz, DMSO-d ₆)7.72 (s, 1H), 3.95 (s, 3H), 3.72 – 3.44 (m, 8H), 3.20 – 3.03 (m, 3H), 2.81 – 2.59 (m, 3H), 1.93 – 1.66 (m, 6H), 1.60 – 1.32 (m, 4H).
193	¹ H NMR (400 MHz, DMSO, ppm): 11.38(s, 1H), 8.01(s, 1H), 7.76 – 7.69(m, 2H), 7.43 – 7.36(m, 2H), 4.03(s, 3H), 3.70 – 3.40(m, 8H), 1.83(s, 2H), 1.50(t, J = 5.4 Hz, 4H).
194	¹ H NMR (400 MHz, DMSO, ppm): 11.40(s, 1H), 8.08(s, 1H), 7.66 – 7.47(m, 3H), 7.30(tt, J = 8.7, 1.8Hz, 1H), 4.04(s, 3H), 3.73 – 3.42(m, 8H), 1.83(s, 2H), 1.49(d, J = 5.5Hz, 4H).

TABLE 3-continued

NMR profiles of the compounds of the present invention
The Nos. recited herein corresponds to the numbering
of the compounds disclosed in table 2.

No. NMR

- 195 ¹H NMR (400 MHz, DMSO, ppm): 11.38(s, 1H), 7.96(d, J = 1.2 Hz, 1H), 7.66 – 7.49(m, 2H), 7.45 – 7.33(m, 2H), 4.05(s, 3H), 3.72 – 3.37(m, 8H), 1.83(s, 2H), 1.49(t, J = 5.4 Hz, 4H).
- 196 ¹H NMR (400 MHz, DMSO-d₆) δ 13.91 (s, 1H), 8.09 (s, 1H), 7.72 – 7.65 (m, 2H), 7.60 – 7.50 (m, 2H), 7.50 – 7.42 (m, 1H), 4.06 (s, 3H), 2.66 (s, 2H), 2.43 (s, 6H), 1.33 – 1.13 (m, 2H), 0.92 – 0.72 (m, 2H)
- 197 ¹H NMR (400 MHz, DMSO-d₆) δ 12.58 (s, 1H), 8.09 (s, 1H), 7.73 – 7.66 (m, 2H), 7.60 – 7.51 (m, 2H), 7.51 – 7.42 (m, 1H), 4.05 (s, 3H), 3.06 – 2.76 (m, 6H), 1.70 – 1.50 (m, 2H), 1.43 – 1.23 (m, 2H)
- 198 ¹H NMR (400 MHz, DMSO-d₆) δ 12.43 (s, 1H), 8.09 (s, 1H), 7.72 – 7.65 (m, 3H), 7.60 – 7.52 (m, 2H), 7.50 – 7.40 (m, 1H), 7.20 (s, 1H), 6.84 (s, 1H), 4.37 (s, 2H), 4.05 (s, 3H), 1.57 (m, 2H), 1.31 – 1.11 (m, 2H).
- 199 ¹H NMR (400 MHz, DMSO-d₆) δ 12.81 (s, 1H), 11.39 (s, 1H), 8.48 (s, 1H), 8.18 (s, 1H), 6.86 (s, 1H), 3.92 (s, 3H), 3.78 – 3.72 (m, 4H), 2.97 – 2.90 (m, 4H)
- 200 ¹H NMR (400 MHz, DMSO-d₆) δ 7.76 (s, 1H), 3.95 (s, 3H), 3.70 – 3.40 (m, 8H), 3.15 – 3.05 (m, 1H), 3.04 – 2.94 (m, 1H), 2.82 – 2.64 (m, 2H), 2.61 – 2.53 (m, 1H), 2.04 – 1.68 (m, 5H), 1.62 – 1.44 (m, 5H)
- 201 ¹H NMR (400 MHz, DMSO-d₆) δ 7.93 (s, 1H), 7.72 – 7.64 (m, 2H), 7.58 – 7.50 (m, 2H), 7.48 – 7.39 (m, 1H), 4.20 – 3.90 (m, 5H), 3.10 – 2.90 (m, 2H), 2.86 – 2.76 (m, 1H), 1.94 – 1.84 (m, 1H), 1.80 – 1.30 (m, 7H), 1.28 – 1.10 (m, 2H).
- 202 ¹H NMR (400 MHz, DMSO-d₆) δ 13.00 (s, 1H), 8.10 (s, 1H), 7.72 – 7.65 (m, 2H), 7.60 – 7.52 (m, 6.9 Hz, 2H), 7.50 – 7.42 (m, 1H), 4.06 (s, 3H), 3.62 – 3.52 (m, 1H), 3.33 (s, 3H), 2.15 – 2.05 (m, 1H), 1.40 – 1.20 (m, 2H).
- 203 ¹H NMR (400 MHz, DMSO-d₆) δ 8.09 (s, 1H), 7.73 – 7.66 (m, 2H), 7.60 – 7.50 (m, 2H), 7.50 – 7.42 (m, 1H), 6.30 – 5.80 (m, 2H), 4.06 (s, 3H), 1.32 – 1.12 (m, 2H), 1.12 – 0.92 (m, 2H)
- 204 ¹H NMR (400 MHz, DMSO-d₆) δ 11.37 (s, 1H), 8.03 (s, 1H), 7.72 – 7.65 (m, 2H), 7.60 – 7.52 (m, 2H), 7.50 – 7.41 (m, 1H), 4.04 (s, 3H), 3.70 – 3.04 (m, 8H), 2.00 – 1.40 (m, 6H)
- 205 ¹H NMR (400 MHz, DMSO-d₆) δ 11.37 (s, 1H), 8.03 (s, 1H), 7.72 – 7.65 (m, 2H), 7.60 – 7.52 (m, 2H), 7.50 – 7.41 (m, 1H), 4.04 (s, 3H), 3.70 – 3.04 (m, 8H), 2.00 – 1.40 (m, 6H)
- 206 ¹H NMR (400 MHz, DMSO-d₆, ppm): 11.42 (s, 1H), 8.90 (dd, J = 2.3, 0.8 Hz, 1H), 8.66 (dd, J = 4.8, 1.6 Hz, 1H), 8.14 – 8.07 (m, 2H), 7.58 (ddd, J = 7.9, 4.8, 0.9 Hz, 1H), 4.05 (s, 3H), 3.70 – 3.37 (m, 8H), 1.83 (s, 2H), 1.50 (m, 4H).
- 207 ¹H NMR (400 MHz, DMSO-d₆) δ 11.40 (s, 1H), 8.03 (s, 1H), 7.72 – 7.65 (m, 2H), 7.59 – 7.51 (m, 2H), 7.50 – 7.41 (m, 1H), 4.04 (s, 3H), 3.84 – 7.72 (m, 2H), 3.68 – 3.32 (m, 6H), 2.03 – 1.75 (m, 4H)
- 208 ¹H NMR (400 MHz, DMSO-d₆) δ 11.40 (s, 1H), 8.03 (s, 1H), 7.72 – 7.65 (m, 2H), 7.59 – 7.51 (m, 2H), 7.50 – 7.41 (m, 1H), 4.04 (s, 3H), 3.84 – 7.72 (m, 2H), 3.68 – 3.32 (m, 6H), 2.03 – 1.75 (m, 4H)
- 209 ¹H NMR (400 MHz, DMSO-d₆) δ 11.42 (s, 1H), 8.00 (s, 1H), 7.64 – 7.50 (m, 1H), 7.50 – 7.32 (m, 2H), 4.05 (s, 3H), 3.68 – 3.28 (m, 8H), 1.90 – 1.70 (m, 2H), 1.57 – 1.37 (m, 4H)
- 210 ¹H NMR (400 MHz, DMSO-d₆) δ 11.39 (s, 1H), 7.98 (s, 1H), 7.56 – 7.02 (m, 3H), 4.04 (s, 3H), 3.70 – 3.20 (m, 8H), 1.90 – 1.68 (m, 2H), 1.54 – 1.44 (m, 4H)
- 211 ¹H NMR (400 MHz, DMSO-d₆) δ 11.13 (s, 1H), 8.82 – 8.70(m, 2H), 8.26 (d, J = 8.2 Hz, 1H), 8.00 – 7.90 (m, 1H), 7.44 – 7.34 (m, 1H), 4.07 (s, 3H), 3.70 – 3.42 (m, 8H), 1.90 – 1.70 (m, 2H), 1.58 – 1.44 (m, 4H)
- 212 ¹H NMR (400 MHz, DMSO-d₆) δ 11.45 (s, 1H), 8.76 – 8.70 (m, 2H), 8.20 (s, 1H), 7.76 – 7.70 (m, 2H), 4.06 (s, 3H), 3.70 – 3.30 (m, 8H), 1.92 – 1.72 (m, 2H), 1.56 – 1.46 (m, 4H)
- 213 ¹H NMR (400 MHz, DMSO-d₆) δ 13.36 (s, 1H), 8.28 – 8.21 (m, 2H), 8.15 (s, 1H), 7.79 – 7.71 (m, 2H), 7.65 – 7.55 (m, 2H), 7.54 – 7.45 (m, 3H), 4.09 (s, 3H), 2.82 (s, 4H)
- 214 ¹H NMR (400 MHz, DMSO-d₆) δ 13.30 (s, 1H), 8.29 – 8.21 (m, 2H), 7.69 (s, 1H), 7.54 – 7.46 (m, 2H), 4.00 (s, 3H), 3.85 – 3.79 (m, 4H), 3.16 – 3.09 (m, 4H), 2.83 (s, 4H)
- 215 ¹H NMR (400 MHz, DMSO-d₆) δ 7.94 (s, 1H), 7.72 – 7.64 (m, 2H), 7.58 – 7.48 (m, 2H), 7.48 – 7.39 (m, 1H), 4.20 – 3.90 (m, 5H), 3.06 – 2.90 (m, 2H), 2.86 – 2.76 (m, 1H), 1.94 – 1.30 (m, 8H), 1.28 – 1.12 (m, 2H)
- 216 ¹H NMR (400 MHz, DMSO-d₆) δ 7.93 (s, 1H), 7.72 – 7.64 (m, 2H), 7.58 – 7.48 (m, 2H), 7.48 – 7.39 (m, 1H), 4.20 – 3.90 (m, 5H), 3.06 – 2.90 (m, 2H), 2.83 – 2.75 (m, 1H), 1.94 – 1.30 (m, 8H), 1.30 – 1.10 (m, 2H).
- 217 ¹H NMR (400 MHz, DMSO-d₆) δ 13.21 (s, 1H), 8.42 (s, 1H), 8.13 (s, 1H), 4.00 (s, 3H), 3.92 (s, 3H), 3.85 – 3.70 (m, 4H), 3.40 – 3.20 (m, 4H)
- 218 ¹H NMR (400 MHz, DMSO-d₆) δ 13.40 (s, 1H), 8.95 (s, 1H), 8.74 – 8.64 (m, 1H), 8.24 – 8.12 (m, 4H), 7.66 – 7.50 (m, 3H), 4.09 (s, 3H), 3.01 (s, 3H), 2.91 (s, 3H)
- 219 ¹H NMR (400 MHz, DMSO-d₆) δ 8.80 – 8.70 (m, 2H), 8.26 – 8.14 (m, 3H), 7.82 – 7.76 (m, 2H), 7.60 – 7.50 (m, 2H), 4.09 (s, 3H), 3.02 (s, 3H), 2.92 (s, 3H)
- 220 ¹H NMR (400 MHz, DMSO-d₆) δ 11.39 (s, 1H), 7.59 – 7.45 (m, 5H), 4.03 (s, 3H), 3.70 – 3.20 (m, 8H), 1.90 – 1.70 (m, 2H), 1.54 – 1.44 (m, 4H)
- 221 ¹H NMR (400 MHz, DMSO-d₆) δ 12.93 (s, 1H), 8.46 (s, 1H), 8.17 (s, 1H), 3.99 (s, 3H), 3.92 (s, 3H), 3.82 – 3.70 (m, 4H), 3.19 – 3.07 (m, 4H)

TABLE 3-continued

NMR profiles of the compounds of the present invention	
The Nos. recited herein corresponds to the numbering of the compounds disclosed in table 2.	
No. NMR	
222	¹ H NMR (400 MHz, DMSO-d ₆) δ 11.75 (s, 1H), 7.71 – 7.56 (m, 5H), 4.08 (s, 3H), 3.72 – 3.42 (m, 8H), 1.96 – 1.68 (m, 2H), 1.54 – 1.44 (m, 4H)
223	¹ H NMR (400 MHz, DMSO-d ₆) δ 11.17 (s, 1H), 7.56 – 7.41 (m, 5H), 4.00 (s, 3H), 3.70 – 3.20 (m, 5.7 Hz, 8H), 2.33 (s, 3H), 1.99 – 1.79 (m, 2H), 1.53 – 1.43 (m, 4H)
224	¹ H NMR (400 MHz, DMSO-d ₆) δ 13.35 (s, 1H), 8.23 – 8.14 (m, 3H), 7.61 – 7.55 (m, 2H), 7.54 – 7.44 (m, 1H), 7.34 – 7.24 (m, 2H), 7.12 – 7.02 (m, 1H), 4.24 – 4.17 (m, 2H), 4.08 (s, 3H), 3.75 – 3.68 (m, 2H), 3.34 (s, 3H), 3.02 (s, 3H), 2.91 (s, 3H)
225	¹ H NMR (400 MHz, DMSO-d ₆) δ 11.35 (s, 1H), 8.05 (s, 1H), 7.51 – 7.41 (m, 1H), 7.28 – 7.18 (m, 2H), 7.07 – 7.00 (m, 1H), 4.21 – 4.14 (m, 2H), 4.03 (s, 3H), 3.74 – 3.30 (m, 12H), 1.90 – 1.70 (m, 2H), 1.55 – 1.45 (m, 4H).
226	¹ H NMR (400 MHz, DMSO-d ₆) δ 13.26 (s, 1H), 8.27 – 8.20 (m, 2H), 7.54 – 7.46 (m, 2H), 3.99 (s, 3H), 3.80 – 3.73 (m, 4H), 3.14 – 3.07 (m, 4H), 2.82 (s, 4H)
227	¹ H NMR (400 MHz, DMSO-d ₆) δ 13.33 (s, 1H), 8.24 – 8.10 (m, 3H), 7.78 – 7.71 (m, 2H), 7.63 – 7.44 (m, 5H), 4.86 – 4.76 (m, 1H), 4.09 (s, 3H), 3.70 – 3.44 (m, 3H), 3.30 – 3.20 (m, 1H), 3.04 – 2.84 (m, 3H)
228	¹ H NMR (400 MHz, DMSO-d ₆) δ 13.37 (s, 1H), 8.66 – 8.56 (m, 1H), 8.26 – 8.18 (m, 2H), 8.13 (s, 1H), 8.05 – 7.95 (m, 2H), 7.78 – 7.70 (m, 2H), 7.64 – 7.54 (m, 2H), 7.53 – 7.45 (m, 1H), 4.78 – 4.68 (m, 1H), 4.09 (s, 3H), 3.58 – 3.48 (m, 2H), 3.41 – 3.31 (m, 2H)
229	¹ H NMR (400 MHz, DMSO-d ₆) δ 11.33 (s, 1H), 7.65 – 7.51 (m, 4H), 7.51 – 7.42 (m, 1H), 4.02 (s, 3H), 3.70 – 3.30 (m, 8H), 1.90 – 1.70 (m, 2H), 1.55 – 1.45 (m, 4H)
230	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.66 – 8.56 (m, 1H), 8.26 – 8.19 (m, 2H), 8.11 (s, 1H), 8.01 – 7.95 (m, 2H), 7.78 – 7.71 (m, 2H), 7.64 – 7.54 (m, 2H), 7.53 – 7.44 (m, 1H), 4.08 (s, 3H), 3.48 – 3.22 (m, 2H), 2.60 – 2.52 (m, 1H), 2.29 (s, 6H)
231	¹ H NMR (400 MHz, DMSO-d ₆) δ 11.09 (s, 1H), 7.60 – 7.40 (m, 5H), 3.70 – 3.20 (m, 8H), 2.73 (s, 3H), 2.36 (s, 3H), 2.00 – 1.80 (m, 2H), 1.53 – 1.43 (m, 4H)
232	¹ H NMR (400 MHz, DMSO-d ₆) δ 11.29 (s, 1H), 7.80 (s, 1H), 4.03 – 3.98 (m, 2H), 3.96 (s, 3H), 3.70 – 3.20 (m, 10H), 2.99 – 2.89 (m, 1H), 1.96 – 1.66 (m, 6H), 1.60 – 1.40 (m, 4H)
233	¹ H NMR (400 MHz, DMSO-d ₆) δ 12.94 (s, 1H), 8.52 (s, 1H), 8.21 (s, 1H), 8.12 (s, 1H), 7.53 – 7.43 (m, 1H), 7.32 – 7.22 (m, 2H), 7.09 – 7.02 (m, 1H), 4.40 – 4.30 (m, 2H), 4.23 – 4.16 (m, 2H), 4.07 (s, 3H), 3.76 – 3.66 (m, 4H), 3.34 (s, 3H), 3.25 (s, 3H)
234	¹ H NMR (400 MHz, DMSO-d ₆) δ 7.50 – 7.32 (m, 2H), 7.20 – 7.10 (m, 2H), 7.08 – 7.00 (m, 1H), 4.17 – 4.10 (m, 2H), 4.01 (s, 3H), 3.74 – 3.30 (m, 13H), 1.90 – 1.70 (m, 2H), 1.56 – 1.40 (m, 4H)
235	¹ H NMR (400 MHz, DMSO-d ₆) δ 12.88 (s, 1H), 8.51 (s, 1H), 8.21 (s, 1H), 7.88 (s, 1H), 4.05 – 3.95 (m, 5H), 3.92 (s, 3H), 3.57 – 3.47 (m, 2H), 3.05 – 2.95 (m, 1H), 1.98 – 1.77 (m, 4H)
236	¹ H NMR (400 MHz, DMSO-d ₆) δ 11.24 (s, 1H), 7.85 (s, 1H), 7.50 – 7.40 (m, 1H), 7.39 – 7.33 (m, 1H), 7.20 – 7.12 (m, 1H), 7.11 – 7.01 (m, 1H), 4.03 (s, 3H), 3.76 (s, 3H), 3.70 – 3.20 (m, 8H), 1.98 – 1.70 (m, 2H), 1.60 – 1.40 (m, 4H)
237	¹ H NMR (400 MHz, DMSO-d ₆) δ 11.32 (s, 1H), 7.94 (s, 1H), 6.28 – 6.20 (m, 1H), 4.34 – 4.24 (m, 2H), 3.99 (s, 3H), 3.92 – 3.82 (m, 2H), 3.74 – 3.34 (m, 8H), 2.58 – 2.53 (m, 2H), 1.99 – 1.70 (m, 2H), 1.60 – 1.40 (m, 4H)
238	¹ H NMR (400 MHz, DMSO-d ₆) δ 12.91 (s, 1H), 8.51 (s, 1H), 8.22 (s, 1H), 8.02 (s, 1H), 6.33 – 6.26 (m, 1H), 4.35 – 4.25 (m, 2H), 4.03 (s, 3H), 3.95 – 3.85 (m, 5H), 2.61 – 2.54 (m, 2H)
239	¹ H NMR (400 MHz, DMSO-d ₆) δ 11.28 (s, 1H), 7.86 (s, 1H), 7.80 – 7.70 (m, 1H), 6.58 – 6.48 (m, 1H), 4.02 (s, 3H), 3.94 (s, 3H), 3.89 (s, 3H), 3.70 – 3.20 (m, 8H), 1.98 – 1.68 (m, 2H), 1.60 – 1.40 (m, 4H)

50

EXAMPLE 2

Preparation of the Compounds of the Present Invention and Analytical Methods

All solvents used were commercially available and used without further purification. Reactions were typically run using anhydrous solvents under an inert atmosphere of nitrogen. Flash column chromatography was generally carried out using Silica gel 60 (0.035-0.070 mm particle size).

All NMR experiments were recorded either on Bruker Mercury Plus 400 NMR Spectrometer equipped with a Bruker 400 BBFO probe at 400 MHz for proton NMR or on Bruker Mercury Plus 300 NMR Spectrometer equipped with a Bruker 300 BBFO probe at 300 MHz for proton NMR. All deuterated solvents contained typically 0.03% to 0.05% v/v


tetramethylsilane, which was used as the reference signal (set at δ=0.00 for both ¹H and ¹³C).

LC-MS analyses were performed on a SHIMADZU LC-MS machine consisting of an UFLC 20-AD system and LCMS 2020 MS detector. The column used was a Shim-pack XR-ODS, 2.2 μm, 3.0×50 mm. A linear gradient was applied, starting at 95% A (A: 0.05% TFA in water) and ending at 100% B (B: 0.05% TFA in acetonitrile) over 2.2 min with a total run time of 3.6 min. The column temperature was at 40° C. with the flow rate at 1.0 mL/min. The Diode Array detector was scanned from 200-400 nm. The mass spectrometer was equipped with an electro spray ion source (ES) operated in a positive or negative mode. The mass spectrometer was scanned between m/z 90-900 with a scan time of 0.6 s.

65

167

1. N-[4-methoxy-7-(morpholin-4-yl)-[1,3]thiazolo[4,5-c]pyridin-2-yl]-1-(2-methoxyethyl)-1H-pyrazole-4-carboxamide, 4



a. Ethyl
1-(2-methoxyethyl)-1H-pyrazole-4-carboxylate

To a solution of ethyl 1H-pyrazole-4-carboxylate (950 mg, 6.78 mmol) in N,N-dimethylformamide (18 ml), was added 1-bromo-2-methoxyethane (1.14 g, 8.20 mmol), potassium carbonate (1.80 g, 13.8 mmol) at room temperature. The reaction mixture was irradiated with microwave radiation for 2 h at 160° C. When the reaction was done, the solids were filtered off. Then the filtrate was concentrated under vacuum to yield ethyl 1-(2-methoxyethyl)-1H-pyrazole-4-carboxylate as a light yellow liquid (1.50 g, crude). MS: $m/z=199.2$ [M+H]⁺.

b. 1-(2-methoxyethyl)-1H-pyrazole-4-carboxylic Acid

To a solution of ethyl 1-(2-methoxyethyl)-1H-pyrazole-4-carboxylate (1.50 g, crude) in tetrahydrofuran (10 ml) was added aq. LiOH (20 ml, 32 mmol, 1.6 M) at room temperature. The resulting mixture was stirred for 3 h at room temperature. When the reaction was done, it was quenched with H₂O (10 ml). The resulting mixture was extracted with dichloromethane (100 ml×3) and the organic phases were combined, washed with brine and dried over anhydrous Na₂SO₄. The solvent was removed under reduced pressure and the residue was purified by flash chromatography eluting with MeOH in DCM (0% to 70% gradient) to yield 1-(2-methoxyethyl)-1H-pyrazole-4-carboxylic acid as a white solid (600 mg, 52% for 2 steps). MS: $m/z=171.2$ [M+H]⁺.

c. 5-Bromo-4-chloropyridin-3-amine

To a solution of 3-bromo-4-chloro-5-nitropyridine (9.50 g, 40.01 mmol) in EtOH (500 mL) was added NH₄Cl (13.27

168

g, 248 mmol), water (50 ml), and Fe (22.39 g, 401 mmol) at room temperature. The resulting mixture was stirred for 2 h at 80° C. When the reaction was done, the solids were

35 filtered off. The resulting mixture was concentrated under reduced pressure and the residue was purified by flash chromatography eluting with 0% to 60% EtOAc in petrol ether to yield 5-bromo-4-chloropyridin-3-amine as a yellow oil (5.94 g, 72%). MS: $m/z=209.2$ [M+H]⁺.

40 d. N-[7-bromo-[1,3]thiazolo[4,5-c]pyridin-2-yl]benzamide

To a solution of 5-bromo-4-chloropyridin-3-amine (2.97 g, 14.3 mmol) in acetone (64 mL) was added benzoyl isothiocyanate (4.47 g, 27.4 mmol) at room temperature. The resulting mixture was stirred for 16 h at 50° C. When the reaction was done, the solids were collected by filtration to yield N-[7-bromo-[1,3]thiazolo[4,5-c]pyridin-2-yl]benzamide as a yellow solid (3.36 g, 70%). MS: $m/z=333.8$ [M+H]⁺.

55 e. N-[7-(morpholin-4-yl)-[1,3]thiazolo[4,5-c]pyridin-2-yl]benzamide

To a solution of N-[7-bromo-[1,3]thiazolo[4,5-c]pyridin-2-yl]benzamide (336 mg, 1.01 mmol) in dioxane (12 ml) was added morpholine (7 ml), 2nd Generation RuPhos precatalyst (88 mg, 0.11 mmol), RuPhos (104 mg, 0.22 mmol), t-BuOK (380 mg, 3.39 mmol) at room temperature. After three cycles of vacuum/nitrogen flash the reaction mixture was irradiated with microwave radiation for 2 h at 160° C. The solids were filtered off. The filtrate was concentrated under reduced pressure and the residue was purified by flash chromatography eluting with EtOAc in petrol ether (0% to 66% gradient) to yield N-[7-(morpholin-4-yl)-[1,3]thiazolo[4,5-c]pyridin-2-yl]benzamide as a white solid (99 mg, 29%). MS: $m/z=341.0$ [M+H]⁺.

169

f. N-[4-bromo-7-(morpholin-4-yl)-[1,3]thiazolo[4,5-c]pyridin-2-yl]benzamide

At 70° C., to a solution of N-[7-(morpholin-4-yl)-[1,3]thiazolo[4,5-c]pyridin-2-yl]benzamide (653 mg, 1.92 mmol) in N,N-dimethylformamide (55 ml) was added a solution of NBS (410 mg, 2.30 mmol) in N,N-dimethylformamide (15 ml) dropwise in a period of 4 h with stirring. When the addition finished, the resulting mixture was stirred for 10 min at 70° C. The resulting mixture was concentrated under vacuum to yield crude product that was washed with the hot EA (100 ml×3) to yield N-[4-bromo-7-(morpholin-4-yl)-[1,3]thiazolo[4,5-c]pyridin-2-yl]benzamide as a light yellow solid (437 mg, 53%). MS: $m/z=418.8$ [M+H]⁺. ¹H NMR (400 MHz, DMSO-d₆) δ 8.23-8.15 (m, 2H), 7.72 (s, 1H), 7.60-7.46 (m, 3H), 3.86-3.78 (m, 4H), 3.25-3.18 (m, 4H).


g. 4-Methoxy-7-(morpholin-4-yl)-[1,3]thiazolo[4,5-c]pyridin-2-amine

To a solution of N-[4-bromo-7-(morpholin-4-yl)-[1,3]thiazolo[4,5-c]pyridin-2-yl]benzamide (388 mg, 0.94 mmol) in methanol (20 ml) was added MeONa in MeOH (7.9 mL, 42 mmol, 5.4 M) at room temperature. The reaction mixture was irradiated with microwave radiation for 2 h at 140° C. When the reaction was done, it was quenched with H₂O (6 ml). The resulting mixture was extracted with dichloromethane (50 ml×3) and the organic phases were combined, washed with brine and dried over Na₂SO₄. The solvent was

170

in THF (10 ml) was added HATU (353 mg, 0.93 mmol), 4-methylmorpholine (170 mg, 1.68 mmol) under N₂ atmosphere. The reaction was stirred for 16 h at 50° C., and then 4-methoxy-7-(morpholin-4-yl)-[1,3]thiazolo[4,5-c]pyridin-2-amine (75 mg, 0.28 mmol) was added. The resulting solution was stirred for another 16 h at 90° C. in an oil bath. When the reaction was done, the reaction was quenched by the addition of water (10 ml). The resulting solution was extracted with DCM (30 ml×3). The organic phases were combined, washed with brine and dried over Na₂SO₄. The solvent was removed under reduced pressure and the residue was purified by prep-HPLC under the following conditions: Column, XBridge Prep C18 OBD Column, 19×150 mm 5 μm; MeCN in water (with 10 mmol/L NH₄HCO₃), 22% to 40% gradient in 8 min; Detector, UV 254/220 nm to afford N-[4-methoxy-7-(morpholin-4-yl)-[1,3]thiazolo[4,5-c]pyridin-2-yl]-1-(2-methoxyethyl)-1H-pyrazole-4-carboxamide as a white solid (54 mg, 45%). HPLC: 98.8% purity, RT=4.216 min. MS: $m/z=419.1$ [M+H]⁺. ¹H NMR (400 MHz, DMSO-d₆) δ 12.87 (s, 1H), 8.52 (s, 1H), 8.21 (s, 1H), 7.64 (s, 1H), 4.35 (t, J=5.1 Hz, 2H), 3.98 (s, 3H), 3.84-3.76 (m, 4H), 3.71 (t, J=5.1 Hz, 2H), 3.25 (s, 3H), 3.14-3.06 (m, 4H).

2. N-[4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-[1,3]thiazolo[4,5-c]pyridin-2-yl]-4-(methoxymethyl)benzamide, 14



55

i. N-[7-(1-methyl-1H-pyrazol-4-yl)-[1,3]thiazolo[4,5-c]pyridin-2-yl]benzamide

removed under reduced pressure and the residue was purified by flash chromatography eluting with MeOH in DCM (0% to 5% gradient) to yield 4-methoxy-7-(morpholin-4-yl)-[1,3]thiazolo[4,5-c]pyridin-2-amine as a light yellow solid (150 mg, 60%). MS: $m/z=267.0$ [M+H]⁺.

h. N-[4-methoxy-7-(morpholin-4-yl)-[1,3]thiazolo[4,5-c]pyridin-2-yl]-1-(2-methoxyethyl)-1H-pyrazole-4-carboxamide

In a 30-mL sealed tube, to a solution of 1-(2-methoxyethyl)-1H-pyrazole-4-carboxylic acid (143 mg, 0.84 mmol)

To a solution of N-[7-bromo-[1,3]thiazolo[4,5-c]pyridin-2-yl]benzamide (2.94 g, 8.80 mmol) in dioxane (112 mL) was added 1-methyl-4-(tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole (2.19 g, 10.6 mmol), Pd(dppf)Cl₂CH₂Cl₂ (359 mg, 0.44 mmol), sodium carbonate (3.27 g, 30.8 mmol), water (23 ml) at room temperature. After three cycles of vacuum/nitrogen flash, the resulting mixture was stirred for 16 h at 80° C. When the reaction was done, the

65

171

solids were filtered off. The filtrate was concentrated under reduced pressure and the residue was purified by flash chromatography eluting with EtOAc in petrol ether (0% to 100% gradient) to yield N-[7-(1-methyl-1H-pyrazol-4-yl)-[1,3]thiazolo[4,5-c]pyridin-2-yl]benzamide as an off-white solid (2.54 g, 86%). MS: $m/z=336.2$ [M+H]⁺.

j. N-[4-bromo-7-(1-methyl-1H-pyrazol-4-yl)-[1,3]thiazolo[4,5-c]pyridin-2-yl]benzamide

At 50° C., to a solution of N-[7-(1-methyl-1H-pyrazol-4-yl)-[1,3]thiazolo[4,5-c]pyridin-2-yl]benzamide (960 mg, 2.86 mmol) in N,N-dimethylformamide (150 ml) was added NBS (662 mg, 3.72 mmol) in N,N-dimethylformamide (30 mL) dropwise in a period of 1 h with stirring. Then the resulting solution was stirred for 1 h at 50° C. When the reaction was done, it was quenched with H₂O (100 ml). The resulting mixture was extracted with dichloromethane (150 ml×3) and the organic phases were combined, washed with brine and dried over anhydrous Na₂SO₄. The solvent was removed under reduced pressure and the residue was purified by flash chromatography eluting with EtOAc in petrol ether (0% to 60% gradient) to yield N-[4-bromo-7-(1-methyl-1H-pyrazol-4-yl)-[1,3]thiazolo[4,5-c]pyridin-2-yl]benzamide as a white solid (675 mg, 57%). MS: $m/z=414.1$ [M+H]⁺. ¹H NMR (400 MHz, Chloroform-d) δ 9.92 (s, 1H), 8.41 (s, 1H), 8.03 (d, J=7.7 Hz, 2H), 7.98 (s, 1H), 7.92 (s, 1H), 7.76-7.68 (m, 1H), 7.66-7.58 (m, 2H), 4.07 (s, 3H).

k. 4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)thiazolo[4,5-c]pyridin-2-amine

To a solution of N-[4-bromo-7-(1-methyl-1H-pyrazol-4-yl)-[1,3]thiazolo[4,5-c]pyridin-2-yl]benzamide (338 mg, 0.81 mmol) in methanol (12 ml) was added MeONa in MeOH (5.5 ml, 29.7 mmol, 5.4 M) at room temperature. The reaction mixture was irradiated with microwave radiation for 2 h at 140° C. When the reaction was done, it was quenched with ice water (20 mL). The resulting mixture was extracted with dichloromethane (50 ml×3) and the organic


172


phases were combined, washed with brine and dried over Na₂SO₄. The solvent was removed under reduced pressure and the residue was purified by flash chromatography eluting with MeOH in DCM (0% to 9% gradient) to yield 4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)thiazolo[4,5-c]pyridin-2-amine as a light yellow solid (33 mg, 11%). MS: $m/z=382.2$ [M+H]⁺.

l. N-[4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-[1,3]thiazolo[4,5-c]pyridin-2-yl]-4-(methoxymethyl)benzamide, 14

To a solution of 4-(methoxymethyl)benzoic acid (53 mg, 0.32 mmol) in tetrahydrofuran (5 ml) was added 4-methylmorpholine (65 mg, 0.64 mmol), HATU (135 mg, 0.35 mmol) at room temperature. The resulting mixture was stirred for 8 h at 50° C. To above mentioned reaction mixture was added 4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-[1,3]thiazolo[4,5-c]pyridin-2-amine (28 mg, 0.11 mmol) and the resulting mixture was stirred for 16 h at 90° C. When the reaction was done, it was quenched by the addition of water (10 ml). The resulting mixture was extracted with DCM (30 ml×3). The organic phases were combined, washed with brine and dried over Na₂SO₄. The solvent was removed under reduced pressure and the residue was purified by prep-HPLC under the following conditions: Column, XBridge Shield RP18 OBD Column, 19×150 mm 5 μ m; MeCN in water (with 10 mmol/l NH₄HCO₃+0.1% NH₃·H₂O), 30% to 52% gradient in 7 min; Detector, UV 254/220 nm. N-[4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-[1,3]thiazolo[4,5-c]pyridin-2-yl]-4-(methoxymethyl)benzamide as a white solid (18 mg, 40%). HPLC: 99.4% purity, RT=5.40 min. MS: $m/z=410.1$ [M+H]⁺. ¹H NMR (400 MHz, DMSO-d₆) δ 13.24 (s, 1H), 8.25-8.13 (m, 4H), 7.94 (s, 1H), 7.50 (d, J=8.0 Hz, 2H), 4.53 (s, 2H), 4.05 (s, 3H), 3.97 (s, 3H), 3.35 (s, 3H).

3. N-[4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-[1,3]thiazolo[4,5-c]pyridin-2-yl]-4-(methoxymethyl)benzamide, 30





m. 4-Chloro-5-iodo-3-nitropyridin-2-ol

To a solution of 4-chloro-3-nitropyridin-2-ol (40.00 g, 229.22 mmol) in CH₃CN (500 ml) was added NIS (56.72 g, 252.14 mmol) at room temperature. The resulting mixture was stirred for 12 h at 80° C. When the reaction was completed, the solvent was removed under reduced pressure. The residue was purified by flash chromatography eluting with EtOAc in PE (0% to 100% gradient) to yield 4-chloro-5-iodo-3-nitropyridin-2-ol as a yellow solid (46.43 g, 63%). MS: m/z=301.0 [M+H]⁺.

n. 4-Chloro-5-iodo-2-methoxy-3-nitropyridine

To a solution of 4-chloro-5-iodo-3-nitropyridin-2-ol (34.00 g, 112.00 mmol) in toluene (700 ml) was added Ag₂CO₃ (30.88 g, 112.0 mmol), CH₃I (31.81 g, 224.2 mmol) at room temperature. The resulting mixture was stirred for 16 h at 80° C. When the reaction was done, the solids were filtered out. The filtrate was concentrated under reduced pressure and the residue was purified by flash chromatography eluting with EtOAc in PE (0% to 20% gradient) to yield 4-chloro-5-iodo-2-methoxy-3-nitropyridine as a light yellow solid (19.50 g, 91%). ¹H NMR (400 MHz, DMSO-d₆) δ 8.82 (s, 1H), 4.00 (s, 3H).

o. 4-chloro-5-iodo-2-methoxypyridin-3-amine

To a solution of 4-chloro-5-iodo-2-methoxy-3-nitropyridine (20.00 g, 63.59 mmol) in EtOH (1 l) was added NH₄Cl (61.26 g, 1145 mmol), water (180 ml), Fe (53.42 g, 953.85 mmol) at room temperature. The resulting mixture was stirred for 10 h at 80° C. When the reaction was done, the solids were filtered out. The filtrate was concentrated under reduced pressure and the residue was diluted in H₂O (1.5 L). The resulting solution was extracted with dichloromethane (1.5 l×3) and the combined organic layer was washed with brine, dried over anhydrous Na₂SO₄ and filtered. The filtrate was concentrated under reduced pressure and the residue was purified by flash chromatography eluting with EtOAc in PE (0% to 100% gradient) to yield 4-chloro-5-iodo-2-methoxypyridin-3-amine as a light yellow solid (17.36 g, 96%). MS: m/z=284.8 [M+H]⁺.

p. N-(7-iodo-4-methoxythiazolo[4,5-c]pyridin-2-yl)benzamide

To a solution of 4-chloro-5-iodo-2-methoxypyridin-3-amine (11.13 g, 39.13 mmol) in acetone (293 ml) was added benzoyl isothiocyanate (9.58 g, 58.70 mmol). The resulting mixture was stirred for 16 h at 50° C. When the reaction was done, the solvent was removed under reduced pressure and the residue was purified by flash chromatography eluting with EtOAc in PE (0% to 100% gradient) to yield N-(7-

iodo-4-methoxythiazolo[4,5-c]pyridin-2-yl)benzamide as a light yellow solid (16.00 g, 99%). MS: m/z=412.0 [M+H]⁺.

q. N-(7-(3,6-dihydro-2H-pyran-4-yl)-4-methoxythiazolo[4,5-c]pyridin-2-yl)benzamide

To a solution of N-[7-iodo-4-methoxy-[1,3]thiazolo[4,5-c]pyridin-2-yl]benzamide (2.85 g, 6.93 mmol) in dioxane (120 ml) was added 2-(3,6-dihydro-2H-pyran-4-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (2.18 g, 10.40 mmol), Pd(dppf)Cl₂CH₂Cl₂ (566 mg, 0.69 mmol), sodium hydroxide (1.05 g, 26.34 mmol) and water (30 ml) at room temperature. After three cycles of vacuum/nitrogen flash, the resulting mixture was stirred for 16 h at 100° C. When the reaction was done, the solids were filtered out. The filtrate was concentrated under reduced pressure and the residue was purified by flash chromatography eluting with MeOH in DCM (0% to 10% gradient) to yield N-(7-(3,6-dihydro-2H-pyran-4-yl)-4-methoxythiazolo[4,5-c]pyridin-2-yl)benzamide as an off-white solid (702 mg, 28%). MS: m/z=368.2 [M+H]⁺.

r. N-(4-methoxy-7-(tetrahydro-2H-pyran-4-yl)thiazolo[4,5-c]pyridin-2-yl)benzamide

To a solution of N-[7-(3,6-dihydro-2H-pyran-4-yl)-4-methoxy-[1,3]thiazolo[4,5-c]pyridin-2-yl]benzamide (702 mg, 1.91 mmol) in MeOH (30 ml) was added Pd/C (10%, 500 mg) under nitrogen atmosphere. After three cycles of vacuum/hydrogen flash, the reaction mixture was hydrogenated at 50° C. for 16 h under hydrogen atmosphere using a hydrogen balloon. When the reaction was done, the resulting mixture was filtered through a celite pad. The filtrate was concentrated under reduced pressure and the residue was purified by flash chromatography eluting with MeOH in DCM (0% to 10% gradient) to yield N-(4-methoxy-7-(tetrahydro-2H-pyran-4-yl)thiazolo[4,5-c]pyridin-2-yl)benzamide as a white solid (330 mg, 47%). MS: m/z=370.3 [M+H]⁺.

s. 4-methoxy-7-(tetrahydro-2H-pyran-4-yl)thiazolo[4,5-c]pyridin-2-amine

To a solution of N-[4-methoxy-7-(oxan-4-yl)-[1,3]thiazolo[4,5-c]pyridin-2-yl]benzamide (330 mg, 0.89 mmol) in methanol (10 ml) was added water (10 ml), sodium hydroxide (357 mg, 8.93 mmol) at room temperature. The resulting mixture was stirred for 16 h at 90° C. When the reaction was done, methanol was removed under reduced pressure. Water (30 ml) was added to the residue and extracted with dichloromethane (30 ml×3). The combined organic phase was washed with brine, dried over Na₂SO₄ and filtered. The filtrate was concentrated under reduced pressure and the residue was purified by flash chromatography eluting with

175

MeOH in DCM (0% to 10% gradient) to yield 4-methoxy-7-(tetrahydro-2H-pyran-4-yl)thiazolo[4,5-c]pyridin-2-amine as a white solid (106 mg, 42%). MS: $m/z=266.0$ $[M+H]^+$.

t. Phenyl (4-methoxy-7-(tetrahydro-2H-pyran-4-yl)thiazolo[4,5-c]pyridin-2-yl)carbamate

To a solution of 4-methoxy-7-(oxan-4-yl)-[1,3]thiazolo[4,5-c]pyridin-2-amine (87 mg, 0.33 mmol) in THF (20 ml) was added potassium carbonate (136 mg, 0.98 mmol), phenyl chloroformate (308 mg, 1.97 mmol) and pyridine (78 mg, 0.98 mmol) at room temperature. The resulting mixture was stirred for 8 h at 50° C. When the reaction was done, the solvent was removed under reduced pressure. The resulting mixture was diluted with water (30 ml) and extracted with dichloromethane (30 mL×3). The combined organic phases was washed with brine, dried over Na₂SO₄ and filtered. The filtrate was concentrated under reduced pressure to yield phenyl (4-methoxy-7-(tetrahydro-2H-pyran-4-yl)thiazolo[4,5-c]pyridin-2-yl)carbamate as a orange oil (88 mg, 70%), which was used in the next step without further purification. MS: $m/z=386.3$ $[M+H]^+$.

u. N-[4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-[1,3]thiazolo[4,5-c]pyridin-2-yl]-4-(methoxymethyl)benzamide, 30

To a solution of phenyl N-[4-methoxy-7-(oxan-4-yl)-[1,3]thiazolo[4,5-c]pyridin-2-yl]carbamate (88 mg, 0.23 mmol) and 4-methylpiperidin-4-ol (79 mg, 0.68 mmol) in THF (10 ml) was added DIPEA (177 mg, 1.37 mmol) at room temperature. The resulting mixture was stirred for 16 h at 50° C. When the reaction was done, the solvent was removed under reduced pressure and the residue was purified by prep-HPLC under the following conditions: column, XBridge Prep OBD C18 Column, 30×150 mm 5 μm; mobile phase, water (with 10 mmol/l NH₄HCO₃+0.1% NH₃·H₂O) and ACN (13.0% ACN up to 40.0% in 8 min); detector, UV 220 nm. The title compound 4-hydroxy-N-[4-methoxy-7-(oxan-4-yl)-[1,3]thiazolo[4,5-c]pyridin-2-yl]-4-methylpiperidine-1-carboxamide was obtained as a white solid (30 mg, 32%). HPLC: 99.7% purity, RT=3.13 min. MS: $m/z=407.3$ $[M+H]^+$. ¹H NMR (400 MHz, DMSO-d₆) δ 11.45 (s, 1H), 7.79 (s, 1H), 4.41 (s, 1H), 4.02-3.92 (m, 5H), 3.84 (d, J=13.2 Hz, 2H), 3.54-3.44 (m, 2H), 3.29-3.21 (m, 2H), 3.03-2.84 (m, 1H), 1.93-1.68 (m, 4H), 1.56-1.36 (m, 4H), 1.14 (s, 3H).

EXAMPLE 3

Testing Compounds of the Present Invention for Inhibitory Activities Against Human Adenosine Receptors in Recombinant Cells

The functional activities of human A_{2A}, A_{2B}, A₁ and A₃ receptors were determined by quantification of cAMP, being the second messenger for adenosine receptors. For this purpose recombinant HEK293 cells, expressing either human A_{2A} or A_{2B} receptors (both Gs coupled were seeded into 394-well microtiter plates, test compounds and agonist (NECA) were added. After a 15 min incubation, HTRF reagents (cAMP dynamic 2, Cis Bio) were added and the cellular cAMP levels were determined using the ENVISION (Perkin Elmer) plate reader.

176

For human A₁ and A₃ receptors, recombinant CHO cells, expressing either A₁ or A₃-receptor, were used. As both receptors couple to Gi proteins, the assay protocol was adapted:

5 Cells were seeded into 384-well plates, forskolin, test compounds and agonists (CPA for A₁- and IB-MECA for A₃-receptor) were added. After 30 min incubation, HTRF reagents (cAMP dynamic 2, Cis Bio) were added and the cellular cAMP levels were determined using the ENVISION (Perkin Elmer) plate reader. Obtained raw data were normalized against the inhibitor control and the neural control (DMSO) and the normalized data were fitted using Gene-Data software.

The compounds of the present invention show a high selectivity for adenosine A_{2A} and A_{2B} receptors over adenosine A₁ and A₃ receptors (see e.g. the data of some examples of the compounds of the present invention in table 4)

Particularly, in contrast to the known adenosine A_{2A} receptor antagonist Tozadenant and similar benzothiazole derivatives, the compounds of the present invention surprisingly show an A_{2A}/A_{2B} dual activity (see table 4) which is preferred for the treatment and/or prevention of hyperproliferative and infectious diseases and disorders as it is disclosed above or the compounds of the present invention show at least a high A_{2A} inhibitory activity together with the other surprising advantages disclosed herein leading to a high efficacy in the treatment and/or prevention of hyperproliferative and infectious diseases and disorders.

TABLE 4

No.	Functional A2A receptor activity, HEK293, cAMP, IC50 [μM]	Functional A2B receptor activity, HEK293, cAMP, IC50 [μM]	Functional A1 receptor activity, CHO, cAMP, IC50 [μM]	Functional A3 receptor activity, CHO, cAMP, IC50 [μM]
Tozadenant	B	D	D	
3	A	C	D	C
4	A	C	D	D
7	B	B	C	D
9	A	C	D	D
13	A	B	B	D
14	A	C	C	C
15	A	B	D	D
17	A	C	D	D
18	A	C	D	D
21	A	B	C	C
22	A	B	D	
23	A	B	C	D
27	A	B	D	
29	A	B	C	C
30	A	B	C	D
33	A	B	D	
34	A	B		
52	A	B	C	D
53	A	B	D	D
54	A	B	C	D
58	A	A	C	D
62	A	B	C	
80	A	B	D	
81	A	B	D	
82	A	B	D	D
113	A	B	D	D
130	A	B	C	D
131	A	B		D
140	B	B	D	
147	A	B	D	
156	B	B	D	C
165	A	A		C
167	A	B	D	D
168	A	B		
170	A	B		

TABLE 4-continued

No.	Functional A2A receptor activity, HEK293, cAMP, IC50 [μ M]	Functional A2B receptor activity, HEK293, cAMP, IC50 [μ M]	Functional A1 receptor activity, CHO, cAMP, IC50 [μ M]	Functional A3 receptor activity, CHO, cAMP, IC50 [μ M]
171	A	B	C	C
172	A	B	D	
173	A	B	D	D
176	A	B	D	
177	A	B	D	
178	A	B		C
179	A	B	D	C
180	A	B	C	
181	A	B	C	
186	A	B	C	C
187	A	B	C	C
188	A	A		C
190	A	B	C	C
193	A	B	D	
194	A	B	C	D
195	A	B	D	D
201	A	B		
202	B	B	C	C
203	A	B	C	C
204	A	B	C	D
205	A	B	D	D
206	B	B	C	
207	A	B	C	D
208	A	B	D	D
209	A	B	D	D
210	A	B	D	D
212	A	B	D	
213	A	B	C	D
214	A	B	D	
217	A	B		C
218	A	B		D
219	A	B	D	
220	A	B	D	D
221	A	B		C
223	A	B		D
224	A	B		C
225	A	B		D
226	A	A		D
227	A	A		D
228	A	B		C
229	A	A		
230	A	B		C
232	A	B	D	
233	A	A	C	C
234	A	A	B	B
235	A	A	B	C
236	B	B	D	D
237	A	A	C	
238	A	A	A	C
239	B	D	D	D
241	A	A	C	C

A means IC₅₀ value is < 10 nM, B means IC₅₀ value is < 100 nM, C means IC₅₀ value is < 1 μ M, D means IC₅₀ value is > 1 μ M.

EXAMPLE 4

Testing the Effects of the Compounds of the Present Invention Against Endogenous Human A_{2A} Receptor

The endogenous functional activity of the Gs-coupled human A_{2A} receptor was measured in T cells, where this receptor is highly expressed. Determination of receptor activity was done by quantification of cAMP, which is a second messenger for adenosine receptors.

In short, human pan T cells were isolated from human PBMC (MACS Pan T Cell Isolation Kit, Miltenyi Biotec) that have been derived from fresh whole blood. The T cells

were seeded in 384-well microtiter plates and treated with test compounds. After 10 min incubation at room temperature, the A_{2A} adenosine receptor agonist CGS-21680 was added, and the plates were incubated for another 45 min. Finally, HTRF reagents (cAMP Femto Kit, CisBio) were added to the wells, and after 1 h cellular cAMP levels were determined using the ENVISION (Perkin Elmer) plate reader.

The obtained raw data were normalized against the inhibitor control and the neutral control (DMSO) and the normalized data were fitted using Genedata Screener software.

The compounds of the present invention show that they are able to inhibit the A_{2A} receptor expressed in human T cells which incubated with the A_{2A} adenosine receptor agonist CGS-21680 (as measured by quantification of cAMP), which is preferred for the treatment and/or prevention of hyperproliferative and infectious diseases and disorders as it is disclosed above. Therefore, the compounds of the present invention surprisingly are able to prevent immunosuppression and thus are able to support anti-tumor T cell induced inhibition of tumor growth, reduction or destruction of metastases and prevention of neovascularization.

EXAMPLE 5

Testing the Pharmacokinetic Properties of the Compounds of the Present Invention in Rat and Mouse

The objective of the study was to obtain information on the pharmacokinetic properties of the compounds of the present invention in female Wistar rats/mice following single intravenous and oral administration.

Material and Methods:

Animal Experiments (In-Life Phase)

Female Wistar rats/mice (n=6) received either a single intravenous (bolus) injection or an oral administration (by gavage) of the tested compound. Doses of 0.2 and 10 mg/kg (per compound) were given intravenously and per os, respectively, as a solution in DMSO (0.2%)/PEG 200 (40%)/water for iv administration and as a suspension in Methocel (0.5%)/Tween 20 (0.25%) in water for oral dosing. Consecutive blood samples were taken sub-lingually under isoflurane inhalation from 3 animals per route of administration after 0.1 (only iv), 0.25 (only po), 0.5, 1, 2, 4, 6 and 24 h and were further processed to obtain plasma. Also, urine and feces samples of 3 rats per route of administration were collected over the time interval from 0-24 h and were pooled for analysis.

Bioanalytics:

The concentrations of the compounds in plasma, feces were quantified using an UPLC method with tandem mass spectrometric detection (LC-MS/MS) previously developed at the 'Institute of Drug Metabolism and Pharmacokinetics'. The LC-MS/MS system consisted of a Waters Acquity UPLC coupled to an AB Sciex mass spectrometer API 5500 Q-trap. The UPLC separation was carried out on a reversed phase column (HSS T3, 1.8 μ M, 2.1x50 mm) using a mobile phase gradient with 0.1% formic acid and acetonitrile as eluents. The detection of the compounds was performed using multiple reaction monitoring in the positive ionization mode. Plasma samples were spiked with internal standard (20 μ l) and the analyte was extracted from the matrix using tertiary-butyl methyl ether (tBME). The organic phase was evaporated to dryness under a stream of nitrogen. The residue was dissolved in acetonitrile/0.1% formic acid for LC-MS/MS analysis. Feces samples were homogenized

179

with 4-times their volume of an ethanol/water mixture (4:1, v/v). Aliquots of the aqueous-ethanolic extracts were spiked with internal standard, diluted with acetonitrile/water (1:1, v/v) and directly injected into the LC-MS/MS system.

Pharmacokinetic Evaluation:





Pharmacokinetic parameters C_{max} and t_{max} were taken from the observed data. Area under the curve (AUC), clearance (CL), volume (V), half-life ($t_{1/2}$), F and all dose-normalized values were calculated using the custom-made software 'DDS-TOX'. 'DDS-TOX' values were evaluated for several compounds and shown comparable to the values given by the validated software WinNonLin. AUC values were calculated by non-compartmental analysis using the

180

linear up/log down method. Numerical data for mean plasma concentrations and derived pharmacokinetic parameters were rounded to 3 significant digits for presentation. Oral bioavailability and excretion data—expressed as % of dose—are displayed using 2 significant digits.

In comparison with the known adenosine A_{2A} receptor antagonist Tozadenant and similar benzothiazole derivatives, the compounds of the present invention surprisingly show better pharmacokinetic properties in mouse as the animal model relevant for cancer (see table 6), which is preferred for the treatment and/or prevention of hyperproliferative and infectious diseases and disorders as it is disclosed above.

TABLE 6

CHEMISTRY	Name	CL [L/h/kg]	t 1/2 [h]	Vss [L/kg]	Feces iv [%]	CMax (iv) @ 1 mg/kg [ng/ml]
	Tozadenant	8,68	0,184	2,03	23@0.2	337
	No. 1: analogue to Tozadenant	6,86	0,172	1,45	Not determined	449
	Benzothiazole analogue to No. 4	2,18	0,331	0,946	2.6@0.2	940
	No. 4	1.58; 1.28	0.575; 0.817	0.899; 0.929	5.7@0.2; 6.4@0.2	1260; 1390

Testing the Effect of the Compounds of the Present Invention on Mouse T Cells

Background:

Adenosine (Ado) in tumor microenvironment can inhibit T cell activity by signaling through A_{2A} receptors and suppress cytokine secretion by T cells. A_{2A} specific agonists like CGS-21680 does similar job of inhibition of T cell cytokine secretion in vitro and in vivo. Potential A_{2A} antagonists or A_{2A}/A_{2B} dual antagonists can rescue T cells from this inhibition. Herein, we describe the in vitro system we established using Pan T cells from mouse spleens to screen potential A_{2A} antagonists or A_{2A}/A_{2B} dual antagonists for their activity. The method described involves the use of CD3/CD28 pre-coated beads to stimulate Pan T cells purified from mouse splenocytes, combined with the addition of A_{2A} agonist along with potential A_{2A} or A_{2A}/A_{2B} dual antagonists to evaluate potentiation of T cell cytokine production.

Assay Description:

Briefly, mouse Pan T cells are purified from spleens of BALB/c mice using Pan T cell isolation kit Mouse II (MACS Miltenyi biotech Cat #Order no. 130-095-130) according to manufacturer's protocol. The purified T cells are seeded in Nunc™ 96-Well Polystyrene Round Bottom Microwell Plates in RPMI medium with 10% heat inactivated fetal bovine serum. The cells are rested at 37° C. for

1 h before activating with CD3/CD28 pre-coated beads (Dynabeads™ Mouse T-Activator CD3/CD28; Cat #11456D). After 30 min the cells are treated with varying doses of test antagonist(s). The cells are incubated for additional 30 min at 37° C. before treating with A_{2A} agonist CGS-21680 (1 μM) or neutral control (DMSO). After 24 h incubation IL-2 levels in the supernatants and after 48 h incubation IFN-γ levels in the supernatants are measured by ELISAs according to manufacturer's protocol (R&D systems Cat #DY402 (IL-2); DY485 (IFN-γ)). Once the concentrations are calculated, the difference of cytokine concentration of DMSO control and agonist alone control is calculated (called Δ) and the percentage of rescue by each concentration of antagonist is calculated by using Microsoft Excel. These percentages of cytokine rescue in a dose dependent manner of antagonist is plotted in GraphPad Prism software and IC_{50} is calculated.

In contrast to the known adenosine A_{2A} receptor antagonist Tozadenant, the compounds of the present invention show that they are able to rescue T cells from inhibition and are able to prevent the suppression of cytokine secretion as induced by adenosine or A_{2A} specific agonists like CGS-2168 (see table 7), which is preferred for the treatment and/or prevention of hyperproliferative and infectious diseases and disorders as it is disclosed above. Therefore, the compounds of the present invention surprisingly are able to prevent immunosuppression and thus are able to support anti-tumor T cell induced inhibition of tumor growth, reduction or destruction of metastases and prevention of neovascularization.

TABLE 7



No. Name	Structure	Mouse T-Cell IL-2	Mouse IFN-γ
Tozadenant		NA (<50% rescue)	NA (<50% rescue)
1 Tozadenant-analogue: 4-Hydroxy-4-methyl-piperidine-1-carboxylic acid (4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide		500 nM	700 nM

TABLE 7-continued

No.	Name	Structure	Mouse	Mouse
			T-Cell IL-2	IFN- γ
4	1-(2-Methoxy-ethyl)-1H-pyrazole-4-carboxylic acid (4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide		37 nM	130 nM
13	1-(2-Methoxy-ethyl)-1H-pyrazole-4-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide		45 nM	30 nM

30

EXAMPLE 7

Injection Vials

A solution of 100 g of a compound of the present invention and 5 g of disodium hydrogenphosphate in 3 l of bidistilled water is adjusted to pH 6.5 using 2 N hydrochloric acid, filtered under sterile conditions, transferred into injection vials, lyophilised under sterile conditions and sealed under sterile conditions. Each injection vial contains 5 mg of a compound of the present invention.

35

EXAMPLE 8

Solution

A solution is prepared from 1 g of a compound of the present invention, 9.38 g of $\text{NaH}_2\text{PO}_4 \cdot 2\text{H}_2\text{O}$, 28.48 g of $\text{Na}_2\text{HPO}_4 \cdot 12\text{H}_2\text{O}$ and 0.1 g of benzalkonium chloride in 940 ml of bidistilled water. The pH is adjusted to 6.8, and the solution is made up to 1 l and sterilised by irradiation.

45

EXAMPLE 9

Ampoules


A solution of 1 kg of a compound of the present invention in 60 l of bidistilled water is filtered under sterile conditions, transferred into ampoules, lyophilised under sterile conditions and sealed under sterile conditions. Each ampoule contains 10 mg of a compound of the present invention.

60

65

The invention claimed is:

1. A compound of formula I,



I

wherein

R^1 is linear or branched alkyl having 1-10 C atoms which is unsubstituted or mono-, di- or trisubstituted by R^5 and in which 1-4 C atoms may be replaced, independently of one another, by O, S, SO, SO_2 , NH, NCH_3 , $-\text{OCO}-$, $-\text{NHCONH}-$, $-\text{NHCO}-$, $-\text{NR}^6\text{SO}_2\text{R}^7-$, $-\text{COO}-$, $-\text{CONH}-$, $-\text{NCH}_3\text{CO}-$, $-\text{CONCH}_3-$, $-\text{C}\equiv\text{C}-$ groups and/or $-\text{CH}=\text{CH}-$ groups, and/or, in addition, 1-10 H atoms may be replaced by F and/or Cl, or mono- or bicyclic cyclic alkyl having 3-7 C atoms which is unsubstituted or mono-, di- or trisubstituted by R^5 and in which 1-4 C atoms may be replaced, independently of one another, by O, S, SO, SO_2 , NH, NCH_3 , $-\text{OCO}-$, $-\text{NHCONH}-$, $-\text{NHCO}-$, $-\text{NR}^6\text{SO}_2\text{R}^7-$, $-\text{COO}-$, $-\text{CONH}-$, $-\text{NCH}_3\text{CO}-$, $-\text{CONCH}_3-$, $-\text{C}\equiv\text{C}-$ groups and/or by $-\text{CH}=\text{CH}-$ groups and/or, in addition, 1-10 H atoms may be replaced by F and/or Cl, or mono- or bicyclic heteroaryl, heterocyclyl, aryl or cyclic alkylaryl, containing 3 to 14 carbon atoms and 0-4 heteroatoms, independently selected from N, O and S, which is unsubstituted or mono-, di- or trisubstituted by R^5 ,

185

R^2 is linear or branched alkyl having 1-10 C atoms which is unsubstituted or mono-, di- or trisubstituted by R^5 and in which 1-4 C atoms may be replaced, independently of one another, by O, S, SO, SO₂, NH, NCH₃, —OCO—, —NHCONH—, —NHCO—, —NR⁶SO₂R⁷—, —COO—, —CONH—, —NCH₃CO—, —CONCH₃—, —C≡C— groups and/or —CH=CH— groups, and/or, in addition, 1-10 H atoms may be replaced by F and/or Cl, or cyclic alkyl having 3-7 C atoms which is unsubstituted or mono-, di- or trisubstituted by R^5 and in which 1-4 C atoms may be replaced, independently of one another, by O, S, SO, SO₂, NH, NCH₃, —OCO—, —NHCONH—, —NHCO—, —NR⁶SO₂R⁷—, —COO—, —CONH—, —NCH₃CO—, —CONCH₃—, —C≡C— groups and/or by —CH=CH— groups and/or, in addition, 1-11 H atoms may be replaced by F and/or Cl, or mono- or bicyclic heteroaryl, heterocyclyl, aryl or cyclic alkyl, containing 3 to 14 carbon atoms and 0-4 heteroatoms, independently selected from N, O and S, which is unsubstituted or mono-, di- or trisubstituted by R^5 .

R^3 is linear or branched alkyl or O-alkyl having 1-6 C atoms or cyclic alkyl having 3-6 C atoms, which is unsubstituted or mono-, di- or trisubstituted by H, =S, =NH, =O, OH, cyclic alkyl having 3-6 C atoms, COOH, Hal, NH₂, SO₂CH₃, SO₂NH₂, CN, CONH₂, NHCOCH₃, NHCONH₂ or NO₂.

R^4 is H, D, linear or branched alkyl having 1-6 C atoms or Hal.

R^5 is H, R⁶, =S, =NR⁶, =O, OH, COOH, Hal, NH₂, SO₂CH₃, SO₂NH₂, CN, CONH₂, NHCOCH₃, NHCONH₂, NO₂, or linear or branched alkyl having 1-10 C atoms which is unsubstituted or mono-, di- or trisubstituted by R^6 and in which 1-4 C atoms may be replaced, independently of one another, by O, S, SO, SO₂, NH, NCH₃, —OCO—, —NHCONH—, —NHCO—, —NR⁶SO₂R⁷—, —COO—, —CONH—, —NCH₃CO—, —CONCH₃—, —C≡C— groups and/or —CH=CH— groups, and/or, in addition, 1-10 H atoms may be replaced by F and/or Cl, or mono- or bicyclic cyclic alkyl having 3-7 C atoms which is unsubstituted or mono-, di- or trisubstituted by R^6 and in which 1-4 C atoms may be replaced, independently of one another, by O, S, SO, SO₂, NH, NCH₃, —OCO—, —NHCONH—, —NHCO—, —NR⁶SO₂R⁷—, —COO—, —CONH—, —NCH₃CO—, —CONCH₃—, —C≡C— groups and/or by —CH=CH— groups and/or, in addition, 1-10 H atoms may be replaced by F and/or Cl, or mono- or bicyclic heteroaryl, heterocyclyl, aryl or cyclic alkyl, containing 3 to 14 carbon atoms and 0-4 heteroatoms, independently selected from N, O and S, which is unsubstituted or mono-, di- or trisubstituted by R^6 .

R^6 , R^7 are independently of one another selected from the group consisting of H, =S, =NH, =O, OH, COOH, Hal, NH₂, SO₂CH₃, SO₂NH₂, CN, CONH₂, NHCOCH₃, NHCONH₂, NO₂ and linear or branched alkyl having 1-10 C atoms in which 1-4 C atoms may be replaced, independently of one another, by O, S, SO, SO₂, NH, NCH₃, —OCO—, —NHCONH—, —NHCO—, —COO—, —CONH—, —NCH₃CO—, —CONCH₃—, —C≡C— groups and/or —CH=CH— groups, and/or, in addition, 1-10 H atoms may be replaced by F and/or Cl,


Hal is F, Cl, Br, or I,

D is deuterium

or physiologically acceptable salts, solvates, prodrugs or stereoisomers thereof, including mixtures thereof in all ratios.

186


2. The compound according to claim 1, wherein R^1 is linear or branched alkyl having 1-10 C atoms which is unsubstituted or mono-, di- or trisubstituted by R^4 and in which 1-4 C atoms may be replaced, independently of one another, by O, S, SO, SO₂, NH, NCH₃, —OCO—, —NHCONH—, —NHCO—, —NR⁵SO₂R⁶—, —COO—, —CONH—, —NCH₃CO—, —CONCH₃—, —C≡C— groups and/or —CH=CH— groups, and/or, in addition, 1-10 H atoms may be replaced by F and/or Cl, or one of the following structures:




which is unsubstituted or mono-, di- or trisubstituted with R^5

and wherein R^2 , R^3 , R^4 , R^5 , R^6 and R^7 have the meanings as disclosed in claim 1, or physiologically acceptable salts, solvates, prodrugs or stereoisomers thereof, including mixtures thereof in all ratios.

3. The compound according to claim 1, wherein R^1 is one of the following structures:



187



and wherein R², R³, R⁴, R⁵, R⁶ and R⁷ have the meanings as disclosed in claim 1, or physiologically acceptable salts, solvates, prodrugs or stereoisomers thereof, including mixtures thereof in all ratios.


4. The compound according to claim 1, wherein

R¹ is phenyl, methylpyrazole or dihydropyran


and R², R³, R⁴, R⁵, R⁶ and R⁷ have the meanings as in claim 1, or physiologically acceptable salts, solvates, prodrugs or stereoisomers thereof, including mixtures thereof in all ratios.

5. The compound according to claim 1, wherein

R² is one of the following structures:



188




which is unsubstituted or mono-, di- or trisubstituted with R⁵

and wherein R¹, R³, R⁴, R⁵, R⁶ and R⁷ have the meanings as disclosed in claim 1, and physiologi-

189


cally acceptable salts, derivatives, solvates, prodrugs and stereoisomers thereof, including mixtures thereof in all ratios.

6. The compound according to claim 1, wherein R² is one of the following structures:




190

-continued




191



and wherein R^1 , R^3 , R^4 , R^5 , R^6 and R^7 have the meanings as disclosed in claim 1, or physiologically acceptable salts, solvates, prodrugs or stereoisomers thereof, including mixtures thereof in all ratios.

7. The compound according to claim 1, wherein R^3 one of the following structures



and R^1 , R^2 , R^4 , R^5 , R^6 and R^7 have the meanings as disclosed in claim 1, or physiologically acceptable salts, solvates, prodrugs or stereoisomers thereof, including mixtures thereof in all ratios.

192

8. The compound according to claim 1, wherein R^3 is OMe

and R^1 , R^2 , R^4 , R^5 , R^6 and R^7 have the meanings as disclosed in claim 1, or physiologically acceptable salts, solvates, prodrugs or stereoisomers thereof, including mixtures thereof in all ratios.

9. The compound according to claim 1, wherein R^1 is phenyl, methylpyrazole or dihydropyran, R^3 is OMe

and R^2 , R^4 , R^5 , R^6 and R^7 have the meanings as disclosed in claim 1, or physiologically acceptable salts, solvates, prodrugs or stereoisomers thereof, including mixtures thereof in all ratios.

10. The compound according to claim 1, wherein

R^4 is H, D, methyl, ethyl, F, Br or Cl

and wherein R^1 , R^2 , R^3 , R^5 , R^6 and R^7 have the meanings as disclosed in claim 1, or physiologically acceptable salts, solvates, prodrugs or stereoisomers thereof, including mixtures thereof in all ratios.

11. The compound according to claim 1, wherein

R^4 is H,

and wherein R^1 , R^2 , R^3 , R^5 , R^6 and R^7 have the meanings as disclosed in claim 1, or physiologically acceptable salts, solvates, prodrugs or stereoisomers thereof, including mixtures thereof in all ratios.

12. A compound selected from the group consisting of:

- 1 4-Hydroxy-4-methyl-piperidine-1-carboxylic acid (4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide
- 2 4-Dimethylaminomethyl-N-(4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-benzamide
- 3 4-Methoxymethyl-N-(4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-benzamide
- 4 1-(2-Methoxy-ethyl)-1H-pyrazole-4-carboxylic acid (4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide
- 5 2-Dimethylaminomethyl-N-(4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-isonicotinamide
- 6 2-Methoxymethyl-N-(4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-isonicotinamide
- 7 4-benzyl-4-hydroxy-N-[4-methoxy-7-(morpholin-4-yl)-[1,3]thiazolo[4,5-c]pyridin-2-yl]piperidine-1-carboxamide
- 8 N-[4-methoxy-7-(morpholin-4-yl)-[1,3]thiazolo[4,5-c]pyridin-2-yl]-5-(2-methoxyethoxy)pyrazine-2-carboxamide
- 9 2-Oxa-7-aza-spiro[4.4]nonane-7-carboxylic acid (4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide
- 10 3-Hydroxy-3-methyl-pyrrolidine-1-carboxylic acid (4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide
- 11 8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid (4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide
- 12 4-Hydroxy-4-prop-2-ynyl-piperidine-1-carboxylic acid (4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide
- 13 1-(2-Methoxy-ethyl)-1H-pyrazole-4-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 14 4-Methoxymethyl-N-[4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-benzamide
- 15 N-(4-Methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-N',N'-dimethyl-terephthalamide
- 16 4-Hydroxymethyl-4-methyl-piperidine-1-carboxylic acid (4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide
- 17 (5S)-N-[4-methoxy-7-(morpholin-4-yl)-[1,3]thiazolo[4,5-c]pyridin-2-yl]-2-oxa-7-azaspiro[4.4]nonane-7-carboxamide
- 18 (5R)-N-[4-methoxy-7-(morpholin-4-yl)-[1,3]thiazolo[4,5-c]pyridin-2-yl]-2-oxa-7-azaspiro[4.4]nonane-7-carboxamide
- 19 N-[4-Methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-benzamide
- 20 4-Methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-ylamine
- 21 4-Hydroxy-4-methyl-piperidine-1-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 22 4-Dimethylaminomethyl-N-[4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-benzamide
- 23 4-Methoxymethyl-N-[4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-benzamide

- 24 2-Dimethylaminomethyl-N-[4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-isonicotinamide
- 25 2-Dimethylaminomethyl-N-[4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-isonicotinamide
- 26 2-Methoxymethyl-N-[4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-isonicotinamide
- 27 4-Dimethylaminomethyl-N-[4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-benzamide
- 28 2-Methoxymethyl-N-[4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-isonicotinamide
- 29 1-(2-Methoxy-ethyl)-1H-pyrazole-4-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 30 4-Hydroxy-4-methyl-piperidine-1-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 31 Isoxazole-4-carboxylic acid (4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide
- 32 4-Hydroxy-4-methyl-piperidine-1-carboxylic acid (4-fluoromethoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide
- 33 4-Imidazol-1-ylmethyl-N-(4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-benzamide
- 34 7-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid (4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide
- 35 1-(2-Methoxy-ethyl)-1H-pyrazole-4-carboxylic acid (4-methoxy-7-pyridin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide
- 36 1-(2-Methoxy-ethyl)-1H-pyrazole-4-carboxylic acid [4-methoxy-7-(6-methyl-pyridazin-3-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 37 4-Hydroxy-4-methyl-piperidine-1-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide
- 38 4-Difluoromethyl-4-hydroxy-piperidine-1-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide
- 39 4-Hydroxymethyl-4-methyl-piperidine-1-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide
- 40 4-Fluoromethyl-4-hydroxy-piperidine-1-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide
- 41 3-Oxa-9-aza-spiro[5.5]undecane-9-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide
- 42 4-Methyl-piperidine-1-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide
- 43 4-Benzyl-4-hydroxy-piperidine-1-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide
- 44 2-Oxo-1-oxa-3,8-diaza-spiro[4.5]decane-8-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide
- 45 3-Oxo-2,8-diaza-spiro[4.5]decane-8-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide
- 46 4-Dimethylaminomethyl-N-(4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-benzamide
- 47 4-Methoxymethyl-N-(4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-benzamide
- 48 2,4-Dioxo-1,3,8-triazaspiro[4.5]decane-8-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide
- 49 4'-Hydroxy-3',4',5',6'-tetrahydro-2'H-[3,4']bipyridinyl-1'-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide
- 50 4-Oxo-piperidine-1-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide
- 51 1-Oxo-2,8-diaza-spiro[4.5]decane-8-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide
- 52 7-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide
- 53 8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide
- 54 2-Oxa-7-aza-spiro[4.5]nonane-7-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide
- 55 4-Imidazol-1-ylmethyl-N-(4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-benzamide
- 56 Isoxazole-3-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide
- 57 4-Hydroxy-4-prop-2-ynyl-piperidine-1-carboxylic acid(4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide
- 58 N-(4-Methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-N',N'-dimethyl-terephthalamide
- 59 N-(4-Methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-4-trifluoromethoxy-benzamide
- 60 2-Methyl-oxazole-4-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide
- 61 Benzooxazole-5-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

- 62 N-(4-Methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-4-(2-oxo-pyrrolidin-1-ylmethyl)-benzamide
- 5 63 2,3-Dihydro-benzofuran-5-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide
- 64 3-Methoxymethyl-pyrrolidine-1-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide
- 65 Piperidine-1,4-dicarboxylic acid 4-amide 1-[(4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide]
- 10 66 4-Diethylamino-N-(4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-benzamide
- 67 4-Difluoromethyl-4-hydroxy-piperidine-1-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 68 4-Hydroxymethyl-4-methyl-piperidine-1-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 15 69 4-Fluoromethyl-4-hydroxy-piperidine-1-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 70 3-Oxa-9-aza-spiro[5.5]undecane-9-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 20 71 4-Methyl-piperidine-1-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 72 4-Benzyl-4-hydroxy-piperidine-1-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 25 73 2-Oxo-1-oxa-3,8-diaza-spiro[4.5]decane-8-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 74 3-Oxo-2,8-diaza-spiro[4.5]decane-8-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 75 4-Dimethylaminomethyl-N-[4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-benzamide
- 30 76 2,4-Dioxo-1,3,8-triazaspiro[4.5]decane-8-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 77 4'-Hydroxy-3',4',5',6'-tetrahydro-2'H-[3,4']bipyridinyl-1'-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 35 78 4-Oxo-piperidine-1-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 79 1-Oxo-2,8-diaza-spiro[4.5]decane-8-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 80 7-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 40 81 8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 82 2-Oxa-7-aza-spiro[4.4]nonane-7-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo [4,5-c]pyridin-2-yl]-amide
- 83 4-Imidazol-1-ylmethyl-N-[4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-benzamide
- 45 84 Isoxazole-3-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 85 4-Hydroxy-4-prop-2-ynyl-piperidine-1-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo [4,5-c]pyridin-2-yl]-amide
- 86 N-[4-Methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-N',N'-dimethyl-terephthalamide
- 87 N-[4-Methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-4-trifluoromethoxy-benzamide
- 50 88 2-Methyl-oxazole-4-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 89 Benzooxazole-5-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 90 N-[4-Methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-4-(2-oxo-pyrrolidin-1-ylmethyl)-benzamide
- 55 91 2,3-Dihydro-benzofuran-5-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 92 3-Methoxymethyl-pyrrolidine-1-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 93 Piperidine-1,4-dicarboxylic acid 4-amide 1-[(4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl)-amide]
- 60 94 4-Diethylamino-N-[4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-benzamide
- 95 4-Difluoromethyl-4-hydroxy-piperidine-1-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 65 96 4-Hydroxymethyl-4-methyl-piperidine-1-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide

- 97 4-Fluoromethyl-4-hydroxy-piperidine-1-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 98 3-Oxa-9-aza-spiro[5.5]undecane-9-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 99 4-Methyl-piperidine-1-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 100 4-Benzyl-4-hydroxy-piperidine-1-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 101 2-Oxo-1-oxa-3,8-diaza-spiro[4.5]decane-8-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 102 3-Oxa-2,8-diaza-spiro[4.5]decane-8-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 103 2,4-Dioxo-1,3,8-triaza-spiro[4.5]decane-8-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 104 4'-Hydroxy-3',4',5',6'-tetrahydro-2'H-[3,4']bipyridinyl-1'-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 105 4-Oxo-piperidine-1-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 106 1-Oxo-2,8-diaza-spiro[4.5]decane-8-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 107 7-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 108 8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 109 2-Oxa-7-aza-spiro[4.4]nonane-7-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 110 4-Imidazol-1-ylmethyl-N-[4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-benzamide
- 111 Isoxazole-3-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 112 4-Hydroxy-4-prop-2-ynyl-piperidine-1-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 113 N-[4-Methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-N',N'-dimethyl-terephthalamide
- 114 N-[4-Methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-4-trifluoromethoxy-benzamide
- 115 2-Methyl-oxazole-4-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 116 Benzooxazole-5-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 117 N-[4-Methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-4-(2-oxo-pyrrolidin-1-ylmethyl)-benzamide
- 118 2,3-Dihydro-benzofuran-5-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 119 3-Methoxymethyl-pyrrolidine-1-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 120 Piperidine-1,4-dicarboxylic acid 4-amide 1-{[4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo [4,5-c]pyridin-2-yl]-amide}
- 121 4-Diethylamino-N-[4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-benzamide
- 122 {1-[4-Methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]carbamoyl}-piperidin-3-yl}-acetic acid
- 123 Pyridine-2,5-dicarboxylic acid 2-dimethylamide 5-{{[4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide}}
- 124 1-Methyl-1H-pyrazole-4-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 125 5-Methyl-isoxazole-4-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 126 1-(2-Methoxy-ethyl)-1H-[1,2,3]triazole-4-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo [4,5-c]pyridin-2-yl]-amide
- 127 1-Methyl-1H-[1,2,3]triazole-4-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 128 1-Cyano-cyclopropanecarboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 129 3-Cyano-N-[4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-propionamide
- 130 2-Methyl-oxazole-5-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 131 2-Methyl-thiazole-5-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 132 4-Hydroxy-pent-2-ynoic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide

- 133 (S)-3-Methanesulfonyl-pyrrolidine-1-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo [4,5-c]pyridin-2-yl]-amide
- 134 (S)-3-Fluoro-pyrrolidine-1-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 135 (S)-3-Cyano-pyrrolidine-1-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 136 (R)-3-Dimethylaminomethyl-pyrrolidine-1-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo [4,5-c]pyridin-2-yl]-amide
- 10 137 5-Methyl-isoxazole-4-carboxylic acid (4-methoxy-7-morpholin-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 138 1-(2-Methoxy-ethyl)-1H-[1,2,3]triazole-4-carboxylic acid (4-methoxy-7-morpholin-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 139 1-Methyl-1H-[1,2,3]triazole-4-carboxylic acid (4-methoxy-7-morpholin-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 15 140 Pyridine-2,5-dicarboxylic acid 2-dimethylamide 5-{{[4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo [4,5-c]pyridin-2-yl]-amide}}
- 141 Pyridine-2,5-dicarboxylic acid 2-dimethylamide 5-[[4-methoxy-7-morpholin-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide]
- 142 1-Methyl-1H-pyrazole-4-carboxylic acid [7-methoxy-4-(tetrahydro-pyran-4-yl)-1H-benzimidazol-2-yl]-amide
- 143 5-Methyl-isoxazole-4-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 20 144 1-(2-Methoxy-ethyl)-1H-[1,2,3]triazole-4-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 145 1-Methyl-1H-[1,2,3]triazole-4-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 25 146 1-Cyano-cyclopropanecarboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 147 2-Methyl-oxazole-5-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 148 2-Methyl-thiazole-5-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 30 149 (S)-3-Methanesulfonyl-pyrrolidine-1-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 150 (S)-3-Fluoro-pyrrolidine-1-carboxylic acid[4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 151 (S)-3-Cyano-pyrrolidine-1-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 35 152 (R)-3-Dimethylaminomethyl-pyrrolidine-1-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 153 1-[4-Methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-3-thiazol-2-ylmethyl-urea
- 154 1-(2-Methoxy-ethyl)-1H-pyrazole-4-carboxylic acid [4-methoxy-7-(2-oxa-7-aza-spiro[4.4]non-7-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 40 155 1-Methyl-1H-pyrazole-4-carboxylic acid {4-methoxy-7-[(2-methoxy-propyl)-methyl-amino]-thiazolo [4,5-c]pyridin-2-yl]-amide
- 156 1-Methyl-1H-pyrazole-4-carboxylic acid [4-methoxy-7-(5-oxa-2-aza-spiro[3.4]oct-2-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 45 157 1-Methyl-1H-pyrazole-4-carboxylic acid (4-methoxy-7-piperidin-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 158 1-Methyl-1H-pyrazole-4-carboxylic acid (4-methoxy-7-piperidin-3-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 159 1-Methyl-1H-pyrazole-4-carboxylic acid [7-(carbamoylmethyl-methyl-amino)-4-methoxy-thiazolo[4,5-c]pyridin-2-yl]-amide
- 50 160 1-Methyl-1H-pyrazole-4-carboxylic acid [4-methoxy-7-(2,2,2-trifluoro-ethoxy)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 161 4-Hydroxy-4-methyl-piperidine-1-carboxylic acid (4-fluoromethoxy-7-morpholin-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide
- 162 4-Hydroxy-4-methyl-piperidine-1-carboxylic acid (4-difluoromethoxy-7-morpholin-4-yl)-thiazolo[4,5-c]pyridin-2-yl)-amide
- 55 163 N-(4-methoxy-6-methyl-7-morpholino-thiazolo[4,5-c]pyridin-2-yl)-1-methyl-pyrazole-4-carboxamide
- 164 N-(6-bromo-4-methoxy-7-morpholino-thiazolo[4,5-c]pyridin-2-yl)-1-methyl-pyrazole-4-carboxamide
- 165 1-Methyl-1H-pyrazole-4-carboxylic acid (6-fluoro-4-methoxy-7-morpholin-4-yl)-thiazolo[4,5-c]pyridin-2-yl)-amide
- 60 166 N-(6-chloro-4-methoxy-7-morpholino-thiazolo[4,5-c]pyridin-2-yl)-1-methyl-pyrazole-4-carboxamide
- 167 Cyclopropanecarboxylic acid (4-methoxy-7-morpholin-4-yl)-thiazolo[4,5-c]pyridin-2-yl)-amide
- 168 (S)-7-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid (4-methoxy-7-morpholin-4-yl)-thiazolo[4,5-c]pyridin-2-yl)-amide
- 65 169 (R)-7-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid (4-methoxy-7-morpholin-4-yl)-thiazolo[4,5-c]pyridin-2-yl)-amide

170 N-(4-Methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-4-(2-oxo-pyrrolidin-1-ylmethyl)-benzamide

171 1-Methyl-1H-pyrazole-4-carboxylic acid (4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide

172 (R)-2-Oxa-7-aza-spiro[4.4]nonane-7-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo [4,5-c]pyridin-2-yl]-amide

173 (S)-2-Oxa-7-aza-Spiro[4.4]nonane-7-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide

174 3-cyano-N-[4-methoxy-7-(morpholin-4-yl)-[1,3]thiazolo[4,5-c]pyridin-2-yl]propanamide

175 1-cyano-N-[4-methoxy-7-(morpholin-4-yl)-[1,3]thiazolo[4,5-c]pyridin-2-yl]cyclopropane-1-carboxamide

176 (S)-7-Oxa-2-aza-Spiro[4.5]decane-2-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide

177 (R)-7-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide

178 1-Methyl-1H-pyrazole-4-carboxylic acid (6-bromo-4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide

179 Cyclopropanecarboxylic acid [4-methoxy-7-(1-methyl-1H-pyrazol-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide

180 N-[4-Methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-4-(2-oxo-pyrrolidin-1-ylmethyl)-benzamide

181 2-Methyl-thiazole-5-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide

182 2-Oxa-7-aza-spiro[4.4]nonane-7-carboxylic acid [7-(3-ethoxy-3-methyl-azetid-1-yl)-4-methoxy-thiazolo[4,5-c]pyridin-2-yl]-amide

183 2-Oxa-7-aza-spiro[4.4]nonane-7-carboxylic acid (4-methoxy-7-piperidin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide

184 2-Oxa-7-aza-spiro[4.4]nonane-7-carboxylic acid [4-methoxy-7-(5-oxa-2-aza-Spiro[3.4]oct-2-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide

185 2-Oxa-7-aza-spiro[4.4]nonane-7-carboxylic acid [4-methoxy-7-(3-methoxy-3-methyl-azetid-1-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide

186 2-Methyl-oxazole-5-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

187 Cyclopropanecarboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

188 1-Methyl-1H-pyrazole-4-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

189 N-[4-methoxy-7-(morpholin-4-yl)-[1,3]thiazolo[4,5-c]pyridin-2-yl]-2-methyl-1,3-oxazole-5-carboxamide

190 2-Methyl-thiazole-5-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

191 8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid [4-methoxy-7-(5-oxa-2-aza-Spiro[3.4]oct-2-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide

192 8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid (4-methoxy-7-pipendin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide

193 8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid [7-(4-fluoro-phenyl)-4-methoxy-thiazolo[4,5-c]pyridin-2-yl]-amide

194 8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid [7-(3-fluoro-phenyl)-4-methoxy-thiazolo[4,5-c]pyridin-2-yl]-amide

195 8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid [7-(2-fluoro-phenyl)-4-methoxy-thiazolo[4,5-c]pyridin-2-yl]-amide

196 1-Dimethylaminomethyl-cyclopropanecarboxylic acid(4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

197 Cyclopropane-1,1-dicarboxylic acid dimethylamide (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

198 1-Imidazol-1-ylmethyl-cyclopropanecarboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

199 1-Methyl-1H-pyrazole-4-carboxylic acid (4-hydroxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide

200 8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid (4-methoxy-7-piperidin-3-yl-thiazolo[4,5-c]pyridin-2-yl)-amide

201 1-amino-N-{4-methoxy-7-phenyl-[1,3]thiazolo[4,5-c]pyridin-2-yl}-8-azaspiro[4.5]decane-8-carboxamide

202 (1 S, 2 S)-2-methoxy-N-[4-methoxy-7-phenyl-[1,3]thiazolo[4,5-c]pyridin-2-yl]cyclopropane-1-carboxamide

203 1-Amino-cyclopropanecarboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

204 (S)-7-Oxa-2-aza-Spiro[4.5]decane-2-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

205 (R)-7-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

206 8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid (4-methoxy-7-pyridin-3-yl-thiazolo[4,5-c]pyridin-2-yl)-amide

207 (S)-2-Oxa-7-aza-Spiro[4.4]nonane-7-carboxylic acid(4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

208 (R)-2-Oxa-7-aza-spiro[4.4]nonane-7-carboxylic acid(4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

209 8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid [7-(2,3-difluoro-phenyl)-4-methoxy-thiazolo[4,5-c]pyridin-2-yl]-amide

210 8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid [7-(2,5-difluoro-phenyl)-4-methoxy-thiazolo[4,5-c]pyridin-2-yl]-amide

211 8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid (4-methoxy-7-pyridin-2-yl-thiazolo[4,5-c]pyridin-2-yl)-amide

212 8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid (4-methoxy-7-pyridin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide

213 4-(2,5-Dioxo-pyrrolidin-1-yl)-N-(4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-benzamide

214 4-(2,5-Dioxo-pyrrolidin-1-yl)-N-(4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-benzamide

215 (R)-1-Amino-8-aza-spiro[4.5]decane-8-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

216 (S)-1-Amino-8-aza-spiro[4.5]decane-8-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

217 1-Methyl-1H-pyrazole-4-carboxylic acid (6-cyano-4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide

218 N-(4-Methoxy-7-pyridin-3-yl-thiazolo[4,5-c]pyridin-2-yl)-N',N'-dimethyl-terephthalamide

219 N-(4-Methoxy-7-pyridin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-N',N'-dimethyl-terephthalamide

220 8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid (6-chloro-4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

221 1-Methyl-1H-pyrazole-4-carboxylic acid (6-chloro-4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide

222 8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid (6-cyano-4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

223 8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid (4-methoxy-6-methyl-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

224 N-{4-Methoxy-7-[3-(2-methoxy-ethoxy)-phenyl]-thiazolo[4,5-c]pyridin-2-yl}-N',N'-dimethyl-terephthalamide

225 8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid {4-methoxy-7-[3-(2-methoxy-ethoxy)-phenyl]-thiazolo[4,5-c]pyridin-2-yl}-amide

226 4-(2,5-Dioxo-pyrrolidin-1-yl)-N-(6-fluoro-4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-benzamide

227 N-(2-Hydroxy-ethyl)-N'-(4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-N-methyl-terephthalamide

228 N-(2-Hydroxy-ethyl)-N'-(4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-terephthalamide

229 8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid (6-fluoro-4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

230 N-(2-Dimethylamino-ethyl)-N'-(4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-terephthalamide

231 8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid (4,6-dimethyl-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

232 8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid [4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-amide

233 N-{4-methoxy-7-[3-(2-methoxyethoxy)phenyl]-[1,3]thiazolo[4,5-c]pyridin-2-yl}-1-(2-methoxyethyl)-1H-pyrazole-4-carboxamide

234 N-{6-fluoro-4-methoxy-7-[3-(2-methoxyethoxy)phenyl]-[1,3]thiazolo[4,5-c]pyridin-2-yl}-8-oxa-2-azaspiro[4.5]decane-2-carboxamide

235 N-[4-methoxy-7-(oxan-4-yl)-[1,3]thiazolo[4,5-c]pyridin-2-yl]-1-methyl-1H-pyrazole-4-carboxamide

236 N-[4-methoxy-7-(2-methoxyphenyl)-[1,3]thiazolo[4,5-c]pyridin-2-yl]-8-oxa-2-azaspiro[4.5]decane-2-carboxamide

237 8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid [7-(3,6-dihydro-2H-pyran-4-yl)-4-methoxy-thiazolo[4,5-c]pyridin-2-yl]-amide

238 N-[7-(3,6-dihydro-2H-pyran-4-yl)-4-methoxy-[1,3]thiazolo[4,5-c]pyridin-2-yl]-1-methyl-1H-pyrazole-4-carboxamide

239 N-[7-(2,6-dimethoxy-pyridin-3-yl)-4-methoxy-[1,3]thiazolo[4,5-c]pyridin-2-yl]-8-oxa-2-azaspiro[4.5]decane-2-carboxamide

240 8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid (6-fluoro-4-hydroxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

241 2-amino-N-[4-methoxy-7-(oxan-4-yl)-[1,3]thiazolo[4,5-c]pyridin-2-yl]-1,3-thiazole-5-carboxamide

242 3-Amino-pyrrolidine-1-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

243 (R)-3-Amino-pyrrolidine-1-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide









244 (S)-3-Amino-pyrrolidine-1-carboxylic acid (4-methoxy-7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

245 8-Oxa-2-aza-spiro[4.5]decane-2-carboxylic acid (7-phenyl-thiazolo[4,5-c]pyridin-2-yl)-amide

199

-continued

- 246 1-(2-Methoxy-ethyl)-1H-pyrazole-4-carboxylic acid (6-fluoro-4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide
 247 Bicyclo[1.1.1]pentane-1,3-dicarboxylic acid dimethylamide (4-methoxy-7-morpholin-4-yl-thiazolo[4,5-c]pyridin-2-yl)-amide
 248 N-(2-Hydroxy-ethyl)-N'-[4-methoxy-7-(tetrahydro-pyran-4-yl)-thiazolo[4,5-c]pyridin-2-yl]-N-methyl-terephthalamide



I

200

of the formula VI employing the use of catalyst and base, a compound of formula VI is converted to a compound of the formula VII by bromination, a compound of the formula VII is converted to a compound of the formula VIII under essentially basic conditions and a compound of the formula VIII is reacted with a compound of the formula IX under standard amidation or carbamide formation conditions to give a compound of the formula I,

or physiologically acceptable salts, solvates, prodrugs or stereoisomers thereof, including mixtures thereof in all ratios.


13. A process for the preparation of a compound of the formula I, according to claim 1, comprising

- a) a compound of the formula II undergoes a reduction to give a compound of formula III, a compound of formula III is reacted with a compound of formula IV at elevated temperature to give a compound of formula V, a compound of formula V is converted to a compound

- b) a compound of the formula V is reacted with a compound of the formula X under Suzuki-type reaction conditions to give a compound of the formula VI, a compound of formula VI is converted to a compound of the formula VII by bromination, a compound of formula VII is converted to a compound of the formula VIII under essentially basic conditions and a compound of the formula VIII is reacted with a compound of the formula IX under standard amidation or carbamide formation conditions to give a compound of the formula I,


201

202




c) a compound of the formula XII is iodinated to give a compound of the formula XIII, a compound of formula XIII is converted to a compound of the formula XIV by treatment with base and an electrophile, a compound of formula XIV is converted to a compound of the formula XV by reduction, a compound of formula XV is reacted with a compound of formula IV at elevated temperature to give a compound of the formula XVI, a compound

of formula XVI is converted under catalytic conditions to a compound of the formula XVII, a compound of the formula XVII is converted to a compound of the formula VIII under basic conditions and a compound of formula VIII is reacted with a compound of the formula IX under standard amidation or carbamide formation conditions to give a compound of the formula I,



203

204



d) the base of a compound of the formula I is converted into one of its salts by treatment with an acid, or
 e) an acid of a compound of the formula I is converted into one of its salts by treatment with a base.

14. The compound according to claim 1 or physiologically acceptable salts, solvates, prodrugs or stereoisomers thereof, including mixtures thereof in all ratios, as adenosine A_{2A} and/or A_{2B} receptor inhibitor.

15. A pharmaceutical composition comprising at least one compound according to claim 1 and/or physiologically acceptable salts, solvates, prodrugs or stereoisomers thereof, including mixtures thereof in all ratios, and a pharmaceutically acceptable carrier.

16. The pharmaceutical composition according to claim 15, comprising further excipients and/or adjuvants.

17. A process for the preparation of a pharmaceutical preparation, wherein a compound according to claim 1 and/or one of its physiologically acceptable salts, solvates, prodrugs or stereoisomers, including mixtures thereof in all ratios, is brought into a suitable dosage form together with a solid, liquid or semi-liquid excipient or adjuvant.

18. A method of treating a cancer that is acute and chronic lymphocytic leukemia, acute granulocytic leukemia, adrenal cortex cancer, bladder cancer, brain cancer, breast cancer, cervical cancer, cervical hyperplasia, cervical cancer, chorio carcinoma, chronic granulocytic leukemia, chronic lymphocytic leukemia, colon cancer, endometrial cancer, esophageal cancer, essential thrombocytosis, genitourinary carcinoma, glioma, glioblastoma, hairy cell leukemia, head and neck carcinoma, Hodgkin's disease, Kaposi's sarcoma, lung carcinoma, lymphoma, malignant carcinoid carcinoma, malignant hypercalcemia, malignant melanoma, malignant pancreatic insulinoma, medullary thyroid carcinoma, melanoma, multiple myeloma, mycosis fungoides, myeloid and lymphocytic leukemia, neuroblastoma, non-Hodgkin's lymphoma, non-small cell lung cancer, osteogenic sarcoma, ovarian carcinoma, pancreatic carcinoma, polycythemia vera, primary brain carcinoma, primary macroglobulinemia, prostatic cancer, renal cell cancer, rhabdomyosarcoma, skin cancer, small-cell lung cancer, soft-tissue sarcoma, squamous cell cancer, stomach cancer, testicular cancer, thyroid cancer Wilms' tumor in a host in need thereof, comprising administering to said host an effective amount of a compound of claim 1 and/or one of its physiologically accept-

able salts, derivatives, solvates, prodrugs and/or stereoisomers, including mixtures thereof in all ratios.

19. A method of treating an infectious disease or disorder in a host in need thereof, comprising administering to said host an effective amount of a compound according to claim 1 and/or one of its physiologically acceptable salts, derivatives, solvates, prodrugs and/or stereoisomers, including mixtures thereof in all ratios, wherein the hyperproliferative disease or disorder is age-related macular degeneration, Crohn's disease, cirrhosis, chronic inflammatory-related disorders, proliferative diabetic retinopathy, proliferative vitreoretinopathy, retinopathy of prematurity, granulomatosis, immune hyperproliferation associated with organ or tissue transplantation or an immunoproliferative disease or disorder that is inflammatory bowel disease, psoriasis, rheumatoid arthritis, systemic lupus erythematosus (SLE), vascular hyperproliferation secondary to retinal hypoxia or vasculitis.

20. The method according to claim 19, wherein the infectious disease or disorder is

- a) virally induced infectious diseases which are caused by retroviruses, hepadnaviruses, herpesviruses, flaviviridae and/or adenoviruses wherein the retroviruses are lentiviruses or oncoretroviruses, wherein the lentivirus is HIV-1, HIV-2, FIV, BIV, SIVs, SHIV, CAEV, VMV or EIAV and the oncoretrovirus is HTLV-I, HTLV-II or BLV, the hepadnavirus is HBV, GSHV or WHV, the herpesvirus is HSV I, HSV II, EBV, VZV, HCMV or HHV 8 and the flaviviridae is HCV, West Nile or Yellow Fever,
- b) bacterial infectious diseases which are caused by Gram-positive bacteria wherein the Gram-positive bacteria are methicillin-susceptible or methicillin-resistant staphylococci, glycopeptides-intermediate susceptible *Staphylococcus aureus* (GISA), penicillin-susceptible and penicillin-resistant streptococci, enterococci, *Clostridium difficile*, *Listeria monocytogenes*, *Corynebacterium jeikeium*, *Chlamydia* spp or *Mycobacterium tuberculosis*,
- c) bacterial infectious diseases which are caused by Gram-negative bacteria wherein the Gram-negative bacteria are of the Genus Enterobacteriaceae, *Klebsiella* spp., *Enterobacter* spp., *Citrobacter* spp., *Serratia* spp., *Proteus* spp., *Providencia* spp., *Salmonella* spp., *Shi-*

- gella* spp., the genus *Pseudomonas*, *Moraxella* spp., *Haemophilus* spp. or *Neisseria* spp.,
- d) infectious diseases induced by intracellular active parasites that are phylum Apicomplexa, or Sarcosporidia, Cryptosporidia, Sarcocystida, Amoebiasis, Coccidia and Trichomonads.
21. Set (kit) consisting of separate packs of
- a) an effective amount of a compound according to claim 1 and/or physiologically acceptable salts, derivatives, solvates, prodrugs and stereoisomers thereof, including mixtures thereof in all ratios, and
- b) an effective amount of a further medicament active compound.

22. The method according to claim 20, wherein the methicillin-susceptible and/or methicillin-resistant staphylococci are *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Staphylococcus haemolyticus*, *Staphylococcus hominis*, *Staphylococcus saprophyticus*, or coagulase-negative staphylococci, the penicillin-susceptible and penicillin-resistant streptococci are *Streptococcus pneumoniae*, *Streptococcus pyogenes*, *Streptococcus agalactiae*, *Streptococcus avium*, *Streptococcus bovis*, *Streptococcus lactis*, *Streptococcus sanguis* or Streptococci Group C (GCS), Streptococci Group G (GGS) or *viridans* streptococci, the enterococci are vancomycin-susceptible or vancomycin-resistant *Enterococcus faecalis* or *Enterococcus faecium*, the *Chlamydia* spp. is *C. pneumoniae*, the Enterobacteriaceae are *Escherichia*, the *Pseudomonas* are *P. aeruginosa*, and the *Moraxella* spp. are *M. catarrhalis*.

23. The method according to claim 20, wherein the Sarcosporidia is *Trypanosoma*, *Plasmodia*, *Leishmania*, *Babesia* or *Theileria*.

* * * * *