

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
3 April 2003 (03.04.2003)

PCT

(10) International Publication Number
WO 03/027075 A2

(51) International Patent Classification⁷: **C07D 231/56**,
A61K 31/415, A61P 29/00, 43/00

MO 63017 (US). **VAZQUEZ, Michael, L.** [US/US];
614 Castle Meadows Court, Ballwin, MO 63021 (US).
WEIER, Richard, M. [US/US]; 240 Hickory Court, Lake
Bluff, IL 60044 (US).

(21) International Application Number: PCT/US02/29673

(74) Agents: **BAUER, S., Christopher** et al.; Pharmacia Corporation, Corporate Patent Department, 800 North Lindbergh Blvd., Mail Zone O4E, St. Louis, MO 63167 (US).

(22) International Filing Date:
19 September 2002 (19.09.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/323,230 19 September 2001 (19.09.2001) US

(71) Applicant (for all designated States except US): **PHARMACIA CORPORATION** [US/US]; Corporate Patent Department, 800 North Lindberg Blvd., Mail Zone O4E, St. Louis, MI 63167 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **GENG, Lifeng** [CN/US]; 5300 Davis Street, Skokie, IL 60077 (US). **CLARE, Michael** [US/US]; 5154 W. Brown Street, Skokie, IL 60077 (US). **HANSON, Gunnar, J.** [US/US]; 7410 Keystone Avenue, Skokie, IL 60077 (US). **HUANG, He** [CN/US]; 1121 Whitfield Road, Northbrook, IL 60062 (US). **IULA, Donna, M.** [US/US]; 773 Independence Drive, Apartment 5, Palatine, IL 60074 (US). **LIAO, Shuyuan** [US/US]; 136 Walters Avenue, Northbrook, IL 60062 (US). **STEALEY, Michael, A.** [US/US]; 502 Juniper Parkway, Libertyville, IL 60048 (US). **METZ, Suzanne** [US/US]; 525 Westernmill Drive, Chesterfield,

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 03/027075 A2

(54) Title: SUBSTITUTED PYRAZOLYL BENZENESULFAMIDE COMPOUNDS FOR THE TREATMENT OF INFLAMMATION

(57) Abstract: The present invention relates to substituted pyrazolyl derivatives, compositions comprising such, intermediates, methods of making substituted pyrazolyl derivatives, and methods for treating cancer, inflammation, and inflammation-associated disorders, such as arthritis.

SUBSTITUTED PYRAZOLYL BENZENESULFAMIDE COMPOUNDS FOR THE TREATMENT OF INFLAMMATION

5

FIELD OF THE INVENTION

[001] The present invention in general is in the field of anti-inflammatory pharmaceutical agents and specifically relates to substituted pyrazolyl benzenesulfamide derivatives, compositions comprising such, and methods for 10 treating cancer, inflammation, and inflammation-associated disorders, such as arthritis.

BACKGROUND OF THE INVENTION

15 [002] The following description of the background of the invention is provided to aid in the understanding the invention, but is not admitted to be or describe prior art to the invention.

20 [003] NF- κ B is a ubiquitous transcription factor that plays a prominent role in the activation of the immune system and in stress responses by regulating the transcription of many early, inducible genes including proinflammatory cytokines, adhesion molecules, growth factors, enzymes, and receptors (Ghosh S., May, M. J., and Kopp. E (1998) *Annu. Rev. Immunol.* **16**, 115-260; Zandi, E., and Karin, M. (1999) *Mol. Cell. Biol.* **19**, 4547-4551; Karin, M. (1999) *J. 25 Biol. Chem.* **274**, 27339-27342). Specificity of gene expression is determined at a cellular level by a diverse array of external stimuli such as bacterial products including LPS, as well as cytokines, most importantly tumor necrosis factor- α (TNF α) and interleukin- β (IL1 β). Through the synergistic interaction with other transcription factors, further specificity can be achieved while maintaining 30 enormous potential to coordinately induce a large number of functionally related

genes. NF- κ B is composed of homo and heterodimers of the Rel protein family and is sequestered in an inactive form in the cytoplasm by members of the I κ B family of inhibitory proteins (Ghosh S., May, M. J., and Kopp. E (1998) *Annu. Rev. Immunol.* **16**, 115-260; Zandi, E., and Karin, M. (1999) *Mol. Cell. Biol.* **19**, 4547-4551; Karin, M. (1999) *J. Biol. Chem.* **274**, 27339-27342). I κ Bs mask the nuclear localization signal on NF- κ B, preventing nuclear translocation and hence DNA binding to the promoter regions of responsive genes. Stimulation of cells with an agonist that activates NF- κ B leads to a series of biochemical signals, ultimately resulting in the phosphorylation, ubiquitinylation, and 5 degradation of I κ Bs, thereby releasing NF- κ B for nuclear translocation (Ghosh S., May, M. J., and Kopp. E (1998) *Annu. Rev. Immunol.* **16**, 115-260; Zandi, E., and Karin, M. (1999) *Mol. Cell. Biol.* **19**, 4547-4551; Karin, M. (1999) *J. Biol. Chem.* **274**, 27339-27342). Recently, two I κ B kinases (IKK1 or IKK α and IKK2 or IKK β), which phosphorylate I κ Bs and thereby initiate their 10 degradation, have been cloned and characterized by a number of laboratories (Ghosh S., May, M. J., and Kopp. E (1998) *Annu. Rev. Immunol.* **16**, 115-260; Zandi, E., and Karin, M. (1999) *Mol. Cell. Biol.* **19**, 4547-4551; Karin, M. (1999) *J. Biol. Chem.* **274**, 27339-27342). The catalytic subunits, IKK1 and 15 IKK2, are similar structurally as well as enzymatically and exist as a heterodimer in a large protein complex referred to as the IKK signalsome 20 (Regnier, C., Song, H., Gao, X., Goeddel, D., Cao, Z. and Rothe, M. (1997) *Cell* **90**, 373-383; DiDonato, J.A., Hayakawa, M., Rothwarf, D.M., Zandi, E. and Karin, M. (1997) *Nature* **388**, 548-554; Mercurio, F., Zhu, H., Murray, B.W., Shevchenko, A., Bennett, B.L., Li, J.W., Young, D.B., Barbosa, M., Mann, M., 25 Manning, A. and Roa, A. (1997) *Science* **278**, 860-866; Zandi, E. Rothwarf, D.M., Delhase, M., Hayadawa, M and Karin, M. (1997) *Cell* **91**, 243-252; Woronicz, J.D., Gao, X., Cao, Z., Rothe, M. And Goeddel, D.V. (1997) *Science* **278**, 866-869). A third protein, NEMO (IKK γ , IKKAP1), is a regulatory 30 adapter protein necessary for IKK activation and kinase activity (Yamaoka, S., Courtois, G., Bessia, C., Whiteside, S. T., Weil, R., Agou, F., Kirk, H. E., Kay, R. J., and Ireal, A. (1998) *Cell* **93**, 1231-1240; Rothwarf, D. M., Zandi, E.,

Natoli, G., Karin, M. (1998) *Nature* **395**, 297; Mercurio, F., Murray, B. W., Shevchenko, A., Bennet, B. L., Young, D. B., Li, J. W., Pascual, G., Motiwala, A., Zhu, H., Mann, M and Manning, A. M. (1999) *Mol. Cell. Biol.* **2**, 1526-1538). IKK1 and IKK2 are co-expressed in most human adult tissues as well as 5 in different developmental stages of mouse embryos (Regnier, C., Song, H., Gao, X., Goeddel, D., Cao, Z. and Rothe, M. (1997) *Cell* **90**, 373-383; DiDonato, J.A., Hayakawa, M., Rothwarf, D.M., Zandi, E. and Karin, M. (1997) *Nature* **388**, 548-554; Mercurio, F., Zhu, H., Murray, B.W., Shevchenko, A., Bennett, B.L., Li, J.W., Young, D.B., Barbosa, M., Mann, M., Manning, A. and 10 Roa, A. (1997) *Science* **278**, 860-866; Zandi, E. Rothwarf, D.M., Delhase, M., Hayadawa, M and Karin, M. (1997) *Cell* **91**, 243-252; Woronicz, J.D., Gao, X., Cao, Z., Rothe, M. and Goeddel, D.V. (1997) *Science* **278**, 866-869; Hu, M. C. T., and Wang, Y. (1998) *Gene* **222**, 31-40). This kinase complex appears to 15 represent a critical, common denominator in the activation of NF- κ B in a number of signal transduction pathways stimulated by a variety of agonists including cytokines, such as TNF α and IL1 β , microbial products such as LPS and viral proteins such as TAX, as well as phorbol esters, oxidizing agents and serine/tyrosine phosphatases (Ghosh S., May, M. J., and Kopp. E (1998) *Annu. Rev. Immunol.* **16**, 115-260; Zandi, E., and Karin, M. (1999) *Mol. Cell. Biol.* **19**, 20 4547-4551; Karin, M. (1999) *J. Biol. Chem.* **274**, 27339-27342).

[004] IKK1 (also termed IKK α , Regnier, C., Song, H., Gao, X., Goeddel, D., Cao, Z. and Rothe, M. (1997) *Cell* **90**, 373-383; DiDonato, J.A., Hayakawa, M., Rothwarf, D.M., Zandi, E. and Karin, M. (1997) *Nature* **388**, 548-554; 25 Mercurio, F., Zhu, H., Murray, B.W., Shevchenko, A., Bennett, B.L., Li, J.W., Young, D.B., Barbosa, M., Mann, M., Manning, A. And Roa, A. (1997) *Science* **278**, 860-866) was cloned simultaneously by standard biochemical purification of the I κ B kinase activity from TNF α stimulated HeLa S3 cells and by its interaction with the MAP3K, NF- κ B inducing kinase (NIK), in a yeast two- 30 hybrid screen. IKK1 was identified as the previously cloned serine-threonine kinase, CHUK (Connelly, M. and Marcu, K. (1995) *Cell. Mol. Biol. Res.* **41**,

537-549). IKK1 (also termed IKK α) is an 85 kDa, 745 amino acid protein that contains an N-terminal serine/threonine kinase catalytic domain, a leucine zipper-like amphipathic helix, and a C-terminal helix-loop-helix domain. IKK2 (also termed IKK β) was also cloned by standard biochemical purification, 5 copurifying with IKK1 from TNF α stimulated HeLa S3 cells as well as by being identified in the public database from an EST clone with sequence homology to IKK1 (Mercurio, F., Zhu, H., Murray, B.W., Shevchenko, A., Bennett, B.L., Li, J.W., Young, D.B., Barbosa, M., Mann, M., Manning, A. and Roa, A. (1997) *Science* **278**, 860-866; Zandi, E. Rothwarf, D.M., Delhase, M., Hayadawa, M 10 and Karin, M. (1997) *Cell* **91**, 243-252; Woronicz, J.D., Gao, X., Cao, Z., Rothe, M. And Goeddel, D.V. (1997) *Science* **278**, 866-869). IKK2 is an 87 kDa, 756 amino acid protein with the same over all topology as IKK1 except for the addition of an 11 amino acid extension at the C-terminus. IKK1 and IKK2 are 52% identical overall with 65% identity in the kinase domain and 44% 15 identity in the protein interaction domains in the C-terminus. Data obtained using transient mammalian expression analysis, by *in vitro* translation experiments and by coexpression in a baculoviral system reveals that IKK1 and IKK2 associate preferentially as a heterodimer through their leucine zipper motifs. Although homodimers have also been described in these systems, the 20 heterodimer is thought to be the physiologic form of the kinase in mammalian cells (Zandi, E. Rothwarf, D.M., Delhase, M., Hayadawa, M and Karin, M. (1997) *Cell* **91**, 243-252; Li, J., Peet, G.W., Pullen, S.S., Schembri-King, J., Warren, T.C., Marcu, K.B., Kehry, M.R., Barton, R. and Jakes, S. (1998) *J. Biol. Chem.* **273**, 30736-30741). Finally, NEMO (also termed IKK γ) contains 25 three α -helical regions including a leucine zipper, interacts preferentially with IKK2 and is required for activation of the heterodimeric kinase complex perhaps by bringing other proteins into the signalsome complex (Yamaoka, S., Courtois, G., Bessia, C., Whiteside, S. T., Weil, R., Agou, F., Kirk, H. E., Kay, R. J., and Ireal, A. (1998) *Cell* **93**, 1231-1240; Rothwarf, D. M., Zandi, E., Natoli, G., 30 Karin, M. (1998) *Nature* **395**, 297; Mercurio, F., Murray, B. W., Shevchenko, A., Bennet, B. L., Young, D. B., Li, J. W., Pascual, G., Motiwala, A., Zhu, H.,

Mann, M and Manning, A. M. (1999) *Mol. Cell. Biol.* **2**, 1526-1538).

[005] The kinase activities of IKK1 and IKK2 are regulated by phosphorylation and require an intact leucine zipper (LZ) for dimerization as well as an intact helix-loop-helix (HLH) domain, which can exert a positive regulatory effect on kinase activity even when it is expressed in trans with the remainder of the IKK protein (Regnier, C., Song, H., Gao, X., Goeddel, D., Cao, Z. and Rothe, M. (1997) *Cell* **90**, 373-383; DiDonato, J.A., Hayakawa, M., Rothwarf, D.M., Zandi, E. and Karin, M. (1997) *Nature* **388**, 548-554; 5 Mercurio, F., Zhu, H., Murray, B.W., Shevchenko, A., Bennett, B.L., Li, J.W., Young, D.B., Barbosa, M., Mann, M., Manning, A. and Roa, A. (1997) *Science* **278**, 860-866; Zandi, E. Rothwarf, D.M., Delhase, M., Hayadawa, M and Karin, M. (1997) *Cell* **91**, 243-252; Woronicz, J.D., Gao, X., Cao, Z., Rothe, M. and Goeddel, D.V. (1997) *Science* **278**, 866-869; Dehase, M., Hayakawa, M., Chen, 10 Y., and Karin, M. (1999) *Science* **284**, 309-313). Both IKK subunits contain a canonical MAPKK activation loop motif near the N- terminus which is the target for phosphorylation and activation of kinase activity by MAP3Ks such as NIK and MEKK1, although the physiologic regulation by these two upstream kinases awaits further characterization (Zandi, E., and Karin, M. (1999) *Mol. Cell. Biol.* **19**, 4547-4551; Karin, M. (1999) *J. Biol. Chem.* **274**, 27339-27342; 15 Karin, M., and Delhase, M. (1998) *Proc. Natl. Acad. Sci. USA* **95**, 9067-9069). Finally, phosphorylation of serines in the C-terminus of IKK2 results in a decrease in IKK activity and it is postulated to be responsible for the transient kinase activity seen after stimulation of cells with an agonist (Dehase, M., 20 Hayakawa, M., Chen, Y., and Karin, M. (1999) *Science* **284**, 309-313).

[006] IKK2 demonstrates a more potent kinase activity compared to IKK1 using I κ B α or I κ B β as a substrate (Mercurio, F., Zhu, H., Murray, B.W., Shevchenko, A., Bennett, B.L., Li, J.W., Young, D.B., Barbosa, M., Mann, M., 25 Manning, A. and Roa, A. (1997) *Science* **278**, 860-866; Zandi, E. Rothwarf, D.M., Delhase, M., Hayadawa, M and Karin, M. (1997) *Cell* **91**, 243-252;

Woronicz, J.D., Gao, X., Cao, Z., Rothe, M. and Goeddel, D.V. (1997) *Science* **278**, 866-869; Dehase, M., Hayakawa, M., Chen, Y., and Karin, M. (1999) *Science* **284**, 309-313). Mutations of the phospho-acceptor serine residues within the MAPKK activation loop alters IKK2 kinase activity; the serine to alanine substitutions result in decreased kinase activity whereas the serine to glutamic acid substitutions result in a constitutively active kinase. Similar alanine mutations in IKK1 do not result in a decreased stimulation of total IKK activity in response to TNF α or IL1 β (Dehase, M., Hayakawa, M., Chen, Y., and Karin, M. (1999) *Science* **284**, 309-313). IKK2 being the dominant kinase activity within the IKK complex is further supported by the analysis of fibroblasts from mice deficient in IKK1 or IKK2. Fibroblasts lacking IKK1 retain full IKK activity in response to cytokines and could activate NF- κ B. In contrast, fibroblasts lacking IKK2 do not exhibit IKK activity when stimulated with cytokines nor do they activate NF- κ B. Furthermore, the phenotypes of each IKK knock out is unique with IKK1 deficiency resulting in skin and skeletal defects and IKK2 knock out being embryonic lethal due to hepatocyte apoptosis (Li, Q., Antwerp, D. V., Mercurio, F., Lee, K., and Verma, I. M. (1999) *Science* **284**, 321-325; Takeda, K., Tekeuchi, O., Tsujimura, T., Itami, S., Adachi, O., Kawai, T., Sanjo, H., Yoshikawa, K., Terada, N, and Akira, S. (1999) *Science* **284**, 313-316; Hu, Y., Baud, V., Delhase, M., Zhang, P., Deerinck, T., Ellisman, M., Johnson, R., and Karin, M. (1999) *Science* **284**, 315-320; Li, Q., Lu, Q., Hwang, J. Y., Buscher, D., Lee, K., Izpisua-Belmonte, J. C., and Verma, I. M. (1999) *Gene and Development* **13**, 1322-1328; Tanaka, M., Fuentes, M. E., Yamaguchi, K., Durnin, M. H., Dalrymple, S. A., Hardy, K. L., and Goeddel, D. V. (1999) *Immunity* **10**, 421-429).

[007] It is well-known that NF-KB plays a key role in the regulated expression of a large number of pro-inflammatory mediators including cytokines such as IL-6 and IL-8, cell adhesion molecules, such as ICAM and VCAM, and inducible nitric oxide synthase (iNOS). Such mediators are known to play a role in the recruitment of leukocytes at sites of inflammation and in the case of

iNOS, may lead to organ destruction in some inflammatory and autoimmune diseases. The importance of NF- κ B in inflammatory disorders is further strengthened by studies of airway inflammation including asthma in which NF- κ B has been shown to be activated. This activation may underlie the increased 5 cytokine production and leukocyte infiltration characteristic of these disorders. In addition, inhaled steroids are known to reduce airway hyper responsiveness and suppress the inflammatory response in asthmatic airways. In light of the recent findings with regard to glucocorticoid inhibition of NF- κ B, one may speculate that these effects are mediated through an inhibition of NF- κ B.

10 Further evidence for a role of NF- κ B in inflammatory disorders comes from studies of rheumatoid synovium. Although NF- κ B is normally present as an inactive cytoplasmic complex, recent immunohistochemical studies have indicated that NF- κ B is present in the nuclei, and hence active, in the cells comprising rheumatoid synovium. Furthermore, NF- κ B has been shown to be 15 activated in human synovial cells in response to stimulation with TNF- α . Such a distribution may be the underlying mechanism for the increased cytokine and eicosanoid production characteristic of this tissue. See Roshak, A. K., et al., *J. Biol. Chem.*, 271, 31496-31501 (1996). The NF- κ B/Rel and I κ B proteins are also likely to play a key role in neoplastic transformation. Family members are 20 associated with cell transformation in vitro and in vivo because of overexpression, gene amplification, gene rearrangements, or translocations. In addition, rearrangement and/or amplification of the genes encoding these proteins are seen in 20-25% of certain human lymphoid tumors. In addition, a role for NF- κ B in the regulation of apoptosis has been reported strengthening 25 the role of this transcription factor in the control of cell proliferation.

[008] The NF- κ B/Rel and I κ B proteins are also likely to play a key role in neoplastic transformation. Family members are associated with cell transformation in vitro and in vivo because of overexpression, gene amplification, gene rearrangements, or translocations (Gillmore TD, *Trends Genet* 7:318-322, 1991; Gillmore TD, *Oncogene* 18:6925-6937, 1999; Rayet B.

et al., *Oncogene* **18**: 6938-6947, 1991). In addition, rearrangement and/or amplification of the genes encoding these proteins are seen in 20-25% of certain human lymphoid tumors. In addition, a role for NF- κ B in the regulation of apoptosis, cell cycle progression, invasion, and metastasis has been reported 5 (Bours V. et al., *Biochemical Pharmacology* **60**:1085-1090, 2000) strengthening the role of this transcription factor in the control of cell proliferation. The inhibition of NF- κ B has been shown to potentiate TNF- and cancer therapy through increased apoptosis (Wang C-Y et al., *Science* **274**:784-787, 1996; Wang C-Y et al., *Nat Med* **5**:412-417, 1999). It has also been shown that human 10 T-cell leukemia virus type 1 (HTLV1) infected cells (the etiological agent of an aggressive malignancy of activated CD4 $^{+}$ T lymphocytes), IKK α and IKK β are expressed constitutively, which normally function in a transient manner (Chu Z-L et al., *J of Biological Chemistry* **273**:15891-15894, 1998). The HTLV1 transforming and transactivating protein (Tax) has been shown to bind MEKK1 15 and increases the activity of IKK β to enhance phosphorylation of serine residues in I κ B α that lead to its degradation.

[009] Pyrazoles have been described for use in the treatment of inflammation. U.S. Patent No. 5,134,142 to Matsuo et al describes 1,5-diaryl pyrazoles, and 20 specifically, 1-(4-fluorophenyl)-5-[4-(methylsulfonyl)phenyl]-3-trifluoromethyl pyrazole, as having anti-inflammatory activity.

[0010] U.S. Patent No. 3,940,418 to R. Hamilton describes tricyclic 4,5-dihydrobenz[g]indazoles as antiinflammatory agents. In addition, R. Hamilton 25 [J. *Heterocyclic Chem.*, **13**, 545 (1976)] describes tricyclic 4,5-dihydrobenz[g]indazoles as antiinflammatory agents. U.S. Patent No. 5,134,155 describes fused tricyclic pyrazoles having a saturated ring bridging the pyrazole and a phenyl radical as HMG-CoA reductase inhibitors. European publication EP 477,049, published Mar. 25, 1992, describes [4,5-dihydro-1-phenyl-1H-benz[g]indazol-3-yl]amides as having antipsychotic activity. European 30 publication EP 347,773, published Dec. 27, 1989, describes [4,5-dihydro-1-

phenyl-1H-benz[g]indazol-3-yl]propanamides as immunostimulants. M. Hashem et al [*J. Med. Chem.*, **19**, 229 (1976)] describes fused tricyclic pyrazoles, having a saturated ring bridging the pyrazole and a phenyl radical, as antibiotics.

5

[0011] Certain substituted pyrazolyl-benzenesulfonamides have been described in the literature as synthetic intermediates. Specifically, 4-[5-(4-chlorophenyl)-3-phenyl-1H-pyrazol-1-yl]benzenesulfonamide has been prepared from a pyrazoline compound as an intermediate for compounds having hypoglycemic activity [R. Soliman et al, *J. Pharm. Sci.*, **76**, 626 (1987)]. 4-[5-[2-(4-Bromophenyl)-2H-1,2,3-triazol-4-yl]-3-methyl-1H-pyrazol-1-yl]benzenesulfonamide has been prepared from a pyrazoline compound and described as potentially having hypoglycemic activity [H. Mokhtar, *Pak. J. Sci. Ind. Res.*, **31**, 762 (1988)]. Similarly, 4-[4-bromo-5-[2-(4-chlorophenyl)-2H-1,2,3-triazol-4-yl]-3-methyl-1H-pyrazol-1-yl]benzenesulfonamide has been prepared [H. Mokhtar et al, *Pak. J. Sci. Ind. Res.*, **34**, 9 (1991)].

[0012] The phytotoxicity of pyrazole derivatives is described [M. Cocco et al, *Il. Farmaco-Ed. Sci.*, **40**, 272 (1985)], specifically for 1-[4-(aminosulfonyl)phenyl]-5-phenyl-1H-pyrazole-3,4-dicarboxylic acid.


[0013] The use of styryl pyrazole esters for antidiabetes drugs is described [H. Mokhtar et al, *Pharmazie*, **33**, 649-651 (1978)]. The use of styryl pyrazole carboxylic acids for antidiabetes drugs is described [R. Soliman et al, *Pharmazie*, **33**, 184-5 (1978)]. The use of 4-[3,4,5-trisubstituted-pyrazol-1-yl]benzenesulfonamides as intermediates for sulfonylurea anti-diabetes agents is described, and specifically, 1-[4-(aminosulfonyl)phenyl]-3-methyl-5-phenyl-1H-pyrazole-4-carboxylic acid [R. Soliman et al, *J. Pharm. Sci.*, **72**, 1004 (1983)]. A series of 4-[3-substituted methyl-5-phenyl-1H-pyrazol-1-yl]benzenesulfonamides has been prepared as intermediates for anti-diabetes agents, and more specifically, 4-[3-methyl-5-phenyl-1H-pyrazol-1-

yl]benzenesulfonamide [H. Feid-Allah, *Pharmazie*, **36**, 754 (1981)]. In addition, 1-(4-[aminosulfonyl]phenyl)-5-phenylpyrazole-3-carboxylic acid has been prepared from the above described 4-[3-methyl-5-phenyl-1*H*-pyrazol-1-yl]benzenesulfonamide compound [R. Soliman et al, *J. Pharm. Sci.*, **70**, 602 (1981)].

[0014] WO 00/27822 discloses tricyclic pyrazole derivatives, WO 00/59901 discloses dihydroindeno pyrazoles, WO 95/15315 discloses diphenyl pyrazole compounds, WO 95/15317 discloses triphenyl pyrazole compounds, WO 10 95/15318 discloses tri-substituted pyrazole compounds, and WO 96/09293 discloses benz[g]indazolyl derivatives. WO 95/15316 discloses substituted pyrazolyl benzenesulfamide derivatives.

DETAILED DESCRIPTION OF THE INVENTION

[0015] A class of compounds, which are useful in treating cancer, inflammation, and inflammation related disorders, is defined by Formula I:



wherein

A is $(CH_2)_m-CR^{15}=CR^{16}-(CH_2)_n$

m is independently selected from 0, 1, or 2;

B is a 5 or 6 membered heteroaryl, aryl, saturated or unsaturated heterocyclic wherein said aryl, heteroaryl, or heterocyclic are optionally substituted with R^1 , R^2 , and R^{12} ;

X is selected from the group consisting of: N and C;

5 **Y** and **Z** are independently selected from the group consisting of: N, CH, CR^3 , S, and O;

R¹ is selected from the group consisting of: hydrido, halogen, alkyl, aryl, heteroaryl, alkenyl, alkynyl, haloalkyl, CN, NO_2 , OR^5 , $OCOOR^5$, CO_2R^7 , $CON(R^6)R^7$, COR^6 , SR^6 , SOR^6 , SO_2R^6 , NR^6R^7 , NR^6COR^7 , NR^6CONHR^7 , $NR^6SO_2R^7$, $NR^6SO_2NHR^7$, and $SO_2N(R^6)R^7$ wherein R^6 and R^7 may be taken together to form a 3-7 membered carbocyclic ring having 1 to 3 substituted or unsubstituted heteroatoms selected from the group consisting of: S, SO, SO_2 , O, and NR^6 ; wherein said alkenyl, alkynyl, alkyl, aryl, heteroaryl or OR^5 are optional substituted with, hydrido, halogen, alkyl, hydroxyalkyl, aryl, heteroaryl, haloalkyl, $COCF_3$, CN, NO_2 , OR^5 , $OCOOR^5$, CO_2R^7 , $CON(R^6)R^7$, COR^6 , SR^6 , SOR^6 , SO_2R^6 , NR^6R^7 , NR^6COR^7 , NR^6CONHR^7 , $NR^6SO_2R^7$, $NR^6SO_2NHR^7$, and $SO_2N(R^6)R^7$ wherein R^6 and R^7 may be taken together to form a 3-7 membered carbocyclic ring having 1 to 3 substituted or unsubstituted heteroatoms selected from the group consisting of: S, SO, SO_2 , O, and NR^6 ;

10 **R²** is selected from the group consisting of: halogen, hydrido, hydroxyalkyl, alkyl, OR^6 , CN, NO_2 , SR^6 , NHR^6 , $CON(R^6)R^7$, $NHCONHR^6$, CO_2H , and haloalkyl;

15 **R¹** and **R²** may be taken together to form a 5 to 7 membered saturated or unsaturated carbocyclic ring optionally containing 0 to 3 heteroatoms selected from the group consisting of N, O, or S, and wherein said ring is optionally substituted with R^1 ;

20

25

R³ is selected from the group consisting of: substituted or unsubstituted amidine, alkylamino, aminoalkyl, CONHR⁷, NH₂, NHCOR⁶, and CH₂NHCOR⁶;

R⁴ is selected from the group consisting of: halogen, alkylsulfinyl, alkylsulfonyl, cyano, alkoxy carbonyl, alkyl, haloalkyl, hydrido, hydroxyalkyl, haloalkoxy, heterocyclic, nitro, acylamino, aryl, heteroaryl, and alkenyl, OR¹³, SR⁸, SO₂N(R⁸)R⁸, NHR⁹, NHCOR⁹, NR⁹COR⁹, NHCO(OR⁹), NR⁹CO(OR⁹), NR⁸SO₂R¹⁰, NHSO₂N(R¹⁰)R¹⁰, NR⁶CON(R¹⁰)R¹⁰, COR⁹, CO₂R⁸, CON(R⁸)R⁸, wherein R⁸ and R^{8'} may be taken together to form a 3-7 membered carbocyclic ring having 1 to 3 substituted or unsubstituted heteroatoms selected from S, SO, SO₂, O, N, and NR⁶, and wherein R¹⁰ and R^{10'} may be taken together to form a 3-7 membered carbocyclic ring having 1 to 3 substituted or unsubstituted heteroatoms selected from S, SO, SO₂, O, N, and NR⁶ wherein said aryl, heterocyclic, heteroaryl, or alkenyl are optionally substituted with R⁹;

R⁵ is selected from the group consisting of: hydrido, alkyl, aryl, arylalkyl, heteroaryl, heterocyclicalkyl, and heteroarylalkyl, wherein aryl, alkyl, arylalkyl, heteroaryl, heterocyclicalkyl, or heteroarylalkyl are optionally substituted with one or more radicals selected from the group consisting of OR¹⁴, N(R¹⁴)R¹⁴, and glycols;

R⁶ is independently selected from the group consisting of: hydrido, aryl, heteroaryl, lower alkyl, haloalkyl, alkenyl, alkynyl, hydroxyalkyl, aminoalkyl, alkylaminoalkyl, alkoxy, alkoxyalkyl, heterocyclicalkyl, and heterocyclic;

R⁷ is independently selected from the group consisting of: hydrido, aryl, heteroaryl, lower alkyl, haloalkyl, alkenyl, alkynyl,

hydroxyalkyl, aminoalkyl, alkylaminoalkyl, alkoxy, alkoxyalkyl, heterocyclicalkyl, and heterocyclic;

5 **R⁸** is independently selected from the group consisting of:
hydrido, aryl, heteroaryl, arylalkyl, heterocyclic, haloalkyl, arylalkylamino, alkylaminoalkyl, dialkylaminoalkyl, alkyl, alkenyl, alkynyl, heteroarylalkyl, and heterocyclicalkyl;

10 **R^{8'}** is independently selected from the group consisting of:
hydrido, aryl, heteroaryl, arylalkyl, heterocyclic, haloalkyl, arylalkylamino, alkylaminoalkyl, dialkylaminoalkyl, alkyl, alkenyl, alkynyl, heteroarylalkyl, and heterocyclicalkyl;

15 **R⁹** is independently selected from the group consisting of:
hydrido, lower alkyl, aryl, heteroaryl, arylalkyl, heterocyclic, cycloalkyl, heterocyclicalkyl, haloalkyl, arylalkylamino, amino, aminoalkyl, aminoacyl, nitro, azido, and heteroarylalkyl, wherein alkyl, aryl, heteroaryl, aminoalkyl, or arylalkyl are optionally substituted with one or more radical selected from the group consisting of: alkylsulfonamide, sulfamyl, alkyl, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, aminoalkyl, alkylaminoalkyl, alkoxy, halogen, acyloxy, oxy, formyl, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, nitro, azido, benzyloxy, dialkylaminoacyl, thioalkyl, aminoacyloxy, thiocyanate, isothiocyanate, alkyldioxy, hydroxyalkyl, alkylamino, alkyloxycarbonyl, alkoxyalkyl, alkenylamino, alkynylamino, alkenyl, alkynyl, dialkylaminoalkyloxy, and heterocyclic optionally substituted with alkyl, alkylamino, aminoalkyl, hydroxyalkyl, and alkylaminoalkyl;

20 **R¹⁰** is independently selected from the group consisting of:
hydrido, lower alkyl, heteroaryl, heterocyclic, haloalkyl, arylalkylamino, heteroarylalkyl, aryl, and arylalkyl, wherein aryl, heteroaryl, heterocyclic, or arylalkyl are optionally substituted

25

30

with one or more radical selected from alkyl, alkoxy, halogen, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, and heterocyclic;

5 **R¹⁰** is independently selected from the group consisting of: hydrido, lower alkyl, heteroaryl, heterocyclic, haloalkyl, arylalkylamino, heteroarylalkyl, aryl, and arylalkyl, wherein aryl, heteroaryl, heterocyclic, or arylalkyl are optionally substituted with one or more radical selected from alkyl, alkoxy, halogen, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, and heterocyclic; and

10 **R¹¹** is selected from the group consisting of: hydrido, halogen, haloalkyl, CN, CO₂R⁵, lower alkyl, lower alkenyl, lower alkynyl, alkoxy, and CONH₂;

15 **R¹²** is selected from the group consisting of: hydrido, halogen, alkyl, and alkoxy;

20 **R¹³** is selected from the group consisting of: hydrido, alkyl, aryl, arylalkyl, heteroaryl, heterocyclicalkyl, and heteroarylalkyl, wherein aryl, alkyl, arylalkyl, heteroaryl, heterocyclicalkyl, or heteroarylalkyl are optionally substituted with one or more radicals selected from the group consisting of OR¹⁴, N(R¹⁴)R¹⁴', and glycols;

25 **R¹⁴** is independently selected from the group consisting of hydrido, and lower alkyl; and

R¹⁴' is independently selected from the group consisting of hydrido, and lower alkyl;

30 **R¹⁵** is selected from the group consisting of: hydrido, halogen, alkyl, cycloalkyl, aryl, haloalkyl, heteroaryl, heterocyclic, alkylalkene, alkylalkyne, hydroxy, hydroxyalkyl, alkylhydroxy, amino, aminoalkyl, alkylamino, alkylaminoalkyl,


alkylhydroxyalkyl, nitro, cyano, alkylthio, alkylsulfinyl, alkylsulfonyl; wherein aryl or arylalkyl are optionally substituted with one or more radical selected from alkyl, alkoxy, halo, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, heterocyclic; and

10 R^{16} is selected from the group consisting of: hydrido, halogen, alkyl, cycloalkyl, aryl, haloalkyl, heteroaryl, heterocyclic, alkylalkene, alkylalkyne, hydroxy, hydroxyalkyl, alkylhydroxy, amino, aminoalkyl, alkylamino, alkylaminoalkyl, alkylhydroxyalkyl, nitro, cyano, alkylthio, alkylsulfinyl, alkylsulfonyl; wherein aryl or arylalkyl are optionally substituted with one or more radical selected from alkyl, alkoxy, halo, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, heterocyclic; wherein R^{15} and R^{16} may be taken together to form a **15** 5 to 7 membered saturated or unsaturated carbocyclic ring optionally containing 0 to 3 heteroatoms selected from the group consisting of N, O, or S, and wherein said ring is optionally substituted with R^1 ;

20

or isomers, tautomers, carriers, esters, prodrugs, pharmaceutically acceptable salts thereof.

[0016] Another class of compounds is defined by formula II



5

wherein

A is $(CH_2)_m-CR^{15}=CR^{16}-(CH_2)_m$

m is independently selected from 0, 1, or 2;

B is a 5 or 6 membered heteroaryl, aryl, saturated or unsaturated heterocyclic wherein said aryl, heteroaryl, or heterocyclic are optionally substituted with **R¹**, **R²**, and **R¹²**;

R¹ is selected from the group consisting of: hydrido, halogen, alkyl, aryl, heteroaryl, alkenyl, alkynyl, haloalkyl, CN, NO₂, OR⁵, OCOOR⁵, CO₂R⁷, CON(R⁶)R⁷, COR⁶, SR⁶, SOR⁶, SO₂R⁶, NR⁶R⁷, NR⁶COR⁷, NR⁶CONHR⁷, NR⁶SO₂R⁷, NR⁶SO₂NHR⁷, and SO₂N(R⁶)R⁷ wherein R⁶ and R⁷ may be taken together to form a 3-7 membered carbocyclic ring having 1 to 3 substituted or unsubstituted heteroatoms selected from the group consisting of: S, SO, SO₂, O, and NR⁶; wherein said alkenyl, alkynyl, alkyl, aryl, heteroaryl or OR⁵ are optional substituted with, hydrido, halogen, alkyl, hydroxyalkyl, aryl, heteroaryl, haloalkyl, COCF₃, CN, NO₂, OR⁵, OCOOR⁵, CO₂R⁷, CON(R⁶)R⁷, COR⁶, SR⁶, SOR⁶, SO₂R⁶, NR⁶R⁷, NR⁶COR⁷, NR⁶CONHR⁷, NR⁶SO₂R⁷, NR⁶SO₂NHR⁷, and SO₂N(R⁶)R⁷ wherein R⁶ and R⁷ may be taken together to form a 3-7 membered carbocyclic ring having 1 to 3

substituted or unsubstituted heteroatoms selected from the group consisting of: S, SO, SO₂, O, and NR⁶;

5 **R²** is selected from the group consisting of: halogen, hydrido, hydroxyalkyl, alkyl, OR⁶, CN, NO₂, SR⁶, NHR⁶, CON(R⁶)R⁷, NHCONHR⁶, CO₂H, and haloalkyl;

R¹ and R² may be taken together to form a 5 to 7 membered saturated or unsaturated carbocyclic ring optionally containing 0 to 3 heteroatoms selected from the group consisting of N, O, or S, and wherein said ring is optionally substituted with R¹;

10 **R³** is selected from the group consisting of: substituted or unsubstituted amidine, alkylamino, aminoalkyl, CONHR⁷, NH₂, NHCOR⁶, and CH₂NHCOR⁶;

15 **R⁴** is selected from the group consisting of: halogen, alkylsulfinyl, alkylsulfonyl, cyano, alkoxycarbonyl, alkyl, haloalkyl, hydrido, hydroxyalkyl, haloalkoxy, heterocyclic, nitro, acylamino, aryl, heteroaryl, and alkenyl, OR¹³, SR⁸, SO₂N(R⁸)R⁸, NHR⁹, NHCOR⁹, NR⁹COR⁹, NHCO(OR⁹), NR⁹CO(OR⁹), NR⁸SO₂R¹⁰, NHSO₂N(R¹⁰)R¹⁰, NR⁶CON(R¹⁰)R¹⁰, COR⁹, CO₂R⁸, CON(R⁸)R⁸, wherein R⁸ and R^{8'} may be taken together to form a 3-7 membered carbocyclic ring having 1 to 3 substituted or unsubstituted heteroatoms selected from S, SO, SO₂, O, N, and NR⁶, and wherein R¹⁰ and R^{10'} may be taken together to form a 3-7 membered carbocyclic ring having 1 to 3 substituted or unsubstituted heteroatoms selected from S, SO, SO₂, O, N, and NR⁶ wherein said aryl, heterocyclic, heteroaryl, or alkenyl are optionally substituted with R⁹;

20 **R⁵** is selected from the group consisting of: hydrido, alkyl, aryl, arylalkyl, heteroaryl, heterocyclicalkyl, and heteroarylalkyl, wherein aryl, alkyl, arylalkyl, heteroaryl, heterocyclicalkyl, or heteroarylalkyl are optionally substituted with one or more

25

30

radicals selected from the group consisting of OR^{14} , $N(R^{14})R^{14'}$, and glycols;

5 R^6 is independently selected from the group consisting of: hydrido, aryl, heteroaryl, lower alkyl, haloalkyl, alkenyl, alkynyl, hydroxyalkyl, aminoalkyl, alkylaminoalkyl, alkoxy, alkoxyalkyl, heterocyclicalkyl, and heterocyclic;

10 R^7 is independently selected from the group consisting of: hydrido, aryl, heteroaryl, lower alkyl, haloalkyl, alkenyl, alkynyl, hydroxyalkyl, aminoalkyl, alkylaminoalkyl, alkoxy, alkoxyalkyl, heterocyclicalkyl, and heterocyclic;

15 R^8 is independently selected from the group consisting of: hydrido, aryl, heteroaryl, arylalkyl, heterocyclic, haloalkyl, arylalkylamino, alkylaminoalkyl, dialkylaminoalkyl, alkyl, alkenyl, alkynyl, heteroarylalkyl, and heterocyclicalkyl;

20 R^8' is independently selected from the group consisting of: hydrido, aryl, heteroaryl, arylalkyl, heterocyclic, haloalkyl, arylalkylamino, alkylaminoalkyl, dialkylaminoalkyl, alkyl, alkenyl, alkynyl, heteroarylalkyl, and heterocyclicalkyl;

25 R^9 is independently selected from the group consisting of: hydrido, lower alkyl, aryl, heteroaryl, arylalkyl, heterocyclic, cycloalkyl, heterocyclicalkyl, haloalkyl, arylalkylamino, amino, aminoalkyl, aminoacyl, nitro, azido, and heteroarylalkyl, wherein alkyl, aryl, heteroaryl, aminoalkyl, or arylalkyl are optionally substituted with one or more radical selected from the group consisting of: alkylsulfonamide, sulfamyl, alkyl, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, aminoalkyl, alkylaminoalkyl, alkoxy, halogen, acyloxy, oxy, formyl, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, nitro, azido, benzyloxy, dialkylaminoacyl, thioalkyl, aminoacyloxy, thiocyanate, isothiocyanate, alkyldioxy, hydroxyalkyl, alkylamino,

alkyloxycarbonyl, alkoxyalkyl, alkenylamino, alkynylamino, alkenyl, alkynyl, dialkylaminoalkyloxy, and heterocyclic optionally substituted with alkyl, alkylamino, aminoalkyl, hydroxyalkyl, and alkylaminoalkyl;

5 **R¹⁰** is independently selected from the group consisting of: hydrido, lower alkyl, heteroaryl, heterocyclic, haloalkyl, arylalkylamino, heteroarylalkyl, aryl, and arylalkyl, wherein aryl, heteroaryl, heterocyclic, or arylalkyl are optionally substituted with one or more radical selected from alkyl, alkoxy, halogen, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, and heterocyclic;

10 **R^{10'}** is independently selected from the group consisting of: hydrido, lower alkyl, heteroaryl, heterocyclic, haloalkyl, arylalkylamino, heteroarylalkyl, aryl, and arylalkyl, wherein aryl, heteroaryl, heterocyclic, or arylalkyl are optionally substituted with one or more radical selected from alkyl, alkoxy, halogen, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, and heterocyclic;

15 **R¹¹** is selected from the group consisting of: hydrido, halogen, haloalkyl, CN, CO₂R⁵, lower alkyl, lower alkenyl, lower alkynyl, alkoxy, and CONH₂;

20 **R¹²** is selected from the group consisting of: hydrido, halogen, alkyl, and alkoxy;

25 **R¹³** is selected from the group consisting of: hydrido, alkyl, aryl, arylalkyl, heteroaryl, heterocyclicalkyl, and heteroarylalkyl, wherein aryl, alkyl, arylalkyl, heteroaryl, heterocyclicalkyl, or heteroarylalkyl are optionally substituted with one or more radicals selected from the group consisting of OR¹⁴, N(R¹⁴)R^{14'}, 30 and glycols;

R¹⁴ is independently selected from the group consisting of hydrido, and lower alkyl; and

R^{14'} is independently selected from the group consisting of hydrido, and lower alkyl;

5 **R¹⁵** is selected from the group consisting of: hydrido, halogen, alkyl, cycloalkyl, aryl, haloalkyl, heteroaryl, heterocyclic, alkylalkene, alkylalkyne, hydroxy, hydroxyalkyl, alkylhydroxy, amino, aminoalkyl, alkylamino, alkylaminoalkyl, alkylhydroxyalkyl, nitro, cyano, alkylthio, alkylsulfinyl, alkylsulfonyl; wherein aryl or arylalkyl are optionally substituted with one or more radical selected from alkyl, alkoxy, halo, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, heterocyclic; and

10 **R¹⁶** is selected from the group consisting of: hydrido, halogen, alkyl, cycloalkyl, aryl, haloalkyl, heteroaryl, heterocyclic, alkylalkene, alkylalkyne, hydroxy, hydroxyalkyl, alkylhydroxy, amino, aminoalkyl, alkylamino, alkylaminoalkyl, alkylhydroxyalkyl, nitro, cyano, alkylthio, alkylsulfinyl, alkylsulfonyl; wherein aryl or arylalkyl are optionally substituted with one or more radical selected from alkyl, alkoxy, halo, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, heterocyclic; wherein **R¹⁵** and **R¹⁶** may be taken together to form a

15 20 25 5 to 7 membered saturated or unsaturated carbocyclic ring optionally containing 0 to 3 heteroatoms selected from the group consisting of N, O, or S, and wherein said ring is optionally substituted with **R¹**;

30 or isomers, tautomers, carriers, esters, prodrugs, pharmaceutically acceptable salts thereof.

Definitions

[0017] The present invention includes the use of all hydrates, solvates, 5 complexes and prodrugs of the compounds of this invention. Prodrugs are any covalently bonded compounds, which release the active parent drug according to Formula I or II in vivo. If a chiral center or another form of an isomeric center is present in a compound of the present invention all forms of such isomer or isomers, including enantiomers and diastereomers, are intended to be covered 10 herein. Compounds containing a chiral center may be used as a racemic mixture, an enantiornerically enriched mixture, or the racemic mixture may be separated using well-known techniques and an individual enantiomer may be used alone. In cases in which compounds have unsaturated carbon-carbon double bonds, both the cis (Z) and trans (E) isomers are within the scope of this invention. In 15 cases wherein compounds may exist in tautomeric forms, such as keto-enol tautomers, each tautomeric form is contemplated as being included within this invention whether existing in equilibrium or predominantly in one form.

[0018] The meaning of any substituent at any one occurrence in Formula I and 20 II or any sub-formula thereof is independent of its meaning, or any other substituents meaning, at any other occurrence, unless specified otherwise.

[0019] The term "alkyl" is used, either alone or within other terms such as "haloalkyl" and "alkylsulfonyl"; it embraces linear or branched radicals having 25 one to about twenty carbon atoms or, preferably, one to about twelve carbon atoms. More preferred alkyl radicals are "lower alkyl" radicals having one to about ten carbon atoms. Most preferred are lower alkyl radicals having one to about five carbon atoms. Examples of such radicals include methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, pentyl, isoamyl, hexyl, 30 octyl and the, like. The term "hydrido" denotes a single hydrogen atom (H). This hydrido radical may be attached, for example, to an oxygen atom to form a

hydroxyl radical or two hydrido radicals may be attached to a carbon atom to form a methylene (-CH₂-) radical. The term "halo" means halogens such as fluorine, chlorine, and bromine or iodine atoms. The term "haloalkyl" embraces radicals wherein any one or more of the alkyl carbon atoms is substituted with 5 halo as defined above. Specifically embraced are monohaloalkyl, dihaloalkyl, and polyhaloalkyl radicals. A monohaloalkyl radical, for one example, may have a bromo, chloro, or a fluoro atom within the radical. Dihalo radicals may have two or more of the same halo atoms or a combination of different halo radicals and polyhaloalkyl radicals may have more than two of the same halo atoms or a 10 combination of different halo radicals. The term "hydroxyalkyl" embraces linear or branched alkyl radicals having one to about ten carbon atoms any one of which may be substituted with one or more hydroxyl radicals. The terms "alkoxy" and "alkoxyalkyl" embrace linear or branched oxy-containing radicals each having alkyl portions of one to about ten carbon atoms, such as methoxy 15 radical. The term "alkoxyalkyl" also embraces alkyl radicals having two or more alkoxy radicals attached to the alkyl radical, that is, to form monoalkoxyalkyl and dialkoxyalkyl radicals. The "alkoxy" or "alkoxyalkyl" radicals may be further substituted with one or more halo atoms, such as fluoro, chloro, or bromo, to provide "haloalkoxy" or "haloalkoxyalkyl" radicals. Examples of 20 "alkoxy" radicals include methoxy, butoxy, and trifluoromethoxy. The term "aryl", alone or in combination, means a carbocyclic aromatic system containing one, two, or three rings wherein such rings may be attached together in a pendent manner or may be fused. The term "aryl" embraces aromatic radicals such as phenyl, naphthyl, tetrahydronaphthyl, indane, and biphenyl. The term 25 "heterocyclic" embraces saturated, partially saturated, and unsaturated heteroatom-containing ring-shaped radicals, where the heteroatoms may be selected from nitrogen, sulfur and oxygen. Examples of saturated heterocyclic radicals include pyrrolidyl and morpholinyl. The term "heteroaryl" embraces unsaturated heterocyclic radicals. Examples of unsaturated heterocyclic radicals, 30 also termed "heteroaryl" radicals include thienyl, pyrrolyl, furyl, pyridyl, pyrimidyl, pyrazinyl, pyrazolyl, oxazolyl, isoxazolyl, imidazolyl, thiazolyl, and

tetrazolyl. The term also embraces radicals where heterocyclic radicals are fused with aryl radicals. Examples of such fused bicyclic radicals include benzofuran, benzothiophene, and the like. The term "heterocyclic alkyl" embraces alkyl attached to the heterocyclic. The term "sulfonyl", whether used alone or linked to other terms such as alkylsulfonyl, denotes respectively divalent radicals – SO_2- . "Alkylsulfonyl", embraces alkyl radicals attached to a sulfonyl radical, where alkyl is defined as above. The term "arylsulfonyl" embraces sulfonyl radicals substituted with an aryl radical. The terms "sulfamyl" or "sulfonamidyl", whether alone or used with terms such as "N-alkylsulfamyl", "N-arylsulfamyl", "N,N-dialkylsulfamyl" and "N-alkyl-N-arylsulfamyl", denotes a sulfonyl radical substituted with an amine radical, forming a sulfonamide ($-\text{SO}_2-\text{NH}_2$). The terms "N-alkylsulfamyl" and "N,N-dialkylsulfamyl" denote sulfamyl radicals substituted, respectively, with one alkyl radical, a cycloalkyl ring, or two alkyl radicals. The terms "N-arylsulfamyl" and "N-alkyl-N-arylsulfamyl" denote sulfamyl radicals substituted, respectively, with one aryl radical, and one alkyl and one aryl radical. The terms "carboxy" or "carboxyl", whether used alone or with other terms, such as "carboxyalkyl", denotes $-\text{CO}_2\text{H}$. The term "carboxyalkyl" embraces radicals having a carboxy radical as defined above, attached to an alkyl radical. The term "carbonyl", whether used alone or with other terms, such as "alkylcarbonyl", denotes $-(\text{C}=\text{O})-$. The term "alkylcarbonyl" embraces radicals having a carbonyl radical substituted with an alkyl radical. An example of an "alkylcarbonyl" radical is $\text{CH}_3-(\text{C}=\text{O})-$. The term "alkylcarbonylalkyl" denotes an alkyl radical substituted with an "alkylcarbonyl" radical. The term "alkoxycarbonyl" means a radical containing an alkoxy radical, as defined above, attached via an oxygen atom to a carbonyl ($\text{C}=\text{O}$) radical. Examples of such "alkoxycarbonyl" radicals include $(\text{CH}_3)_3\text{CO}-\text{C}=\text{O}-$ and $-(\text{O}=\text{C})-\text{OCH}_3$. The term "alkoxycarbonylalkyl" embraces radicals having "alkoxycarbonyl", as defined above substituted to an alkyl radical. Examples of such "alkoxycarbonylalkyl" radicals include $(\text{CH}_3)_3\text{COC}(\text{=O})-(\text{CH}_2)_2-$ and $-(\text{CH}_2)_2(\text{O}=\text{C})-\text{COCH}_3$. The term "amido" when used by itself or with other terms such as "amidoalkyl", "N-monoalkylamido", "N-monoaryl amido",

"N,N-dialkylamido", "N-alkyl-N-arylamido", "N-alkyl-N-hydroxyamido" and "N-alkyl-N-hydroxyamidoalkyl", embraces a carbonyl radical substituted with an amino radical. The terms "N-alkylamido" and "N,N-dialkylamido" denote amido groups which have been substituted with one alkyl radical and with two alkyl radicals, respectively. The terms "N-monoarylamido" and "N-alkyl-N-arylamido" denote amido radicals substituted, respectively, with one aryl radical, and one alkyl and one aryl radical. The term "N-alkyl-N-hydroxyamido" embraces amido radicals substituted with a hydroxyl radical and with an alkyl radical. The term "N-alkyl-N-hydroxyamidoalkyl" embraces alkyl radicals substituted with an N-alkyl-N-hydroxyamido radical. The term "amidoalkyl" embraces alkyl radicals substituted with amido radicals. The term "aminoalkyl" embraces alkyl radicals substituted with amino radicals. The term "alkylaminoalkyl" embraces aminoalkyl radicals having the nitrogen atom substituted with an alkyl radical. The term "amidino" denotes an $-\text{C}(\text{=NH})-\text{NH}_2$ radical. The term "cyanoamidino" denotes an $-\text{C}(\text{=N}-\text{CN})-\text{NH}_2$ radical. The term "heterocycloalkyl" embraces heterocyclic-substituted alkyl radicals such as pyridylmethyl and thienylmethyl. The term "aralkyl" embraces aryl-substituted alkyl radicals such as benzyl, diphenylmethyl, triphenylmethyl, phenethyl, and diphenethyl. The terms benzyl and phenylmethyl are interchangeable. The term "cycloalkyl" embraces radicals having three to ten carbon atoms, such as cyclopropyl cyclobutyl, cyclopentyl, cyclohexyl, and cycloheptyl. The term "cycloalkenyl" embraces unsaturated radicals having three to ten carbon atoms, such as cylopropenyl, cyclobutenyl, cyclopentenyl, cyclohexenyl, and cycloheptenyl. The term "alkylthio" embraces radicals containing a linear or branched alkyl radical, of one to ten carbon atoms, attached to a divalent sulfur atom. An example of "alkylthio" is methylthio, $(\text{CH}_3-\text{S}-)$. The term "alkylsulfinyl" embraces radicals containing a linear or branched alkyl radical, of one to ten carbon atoms, attached to a divalent $-\text{S}(\text{=O})-$ atom. The terms "N-alkylamino" and "N, N-dialkylamino" denote amino groups which have been substituted with one alkyl radical and with two alkyl radicals, respectively. The term "acyl", whether used alone, or within a

term such as "acylamino", denotes a radical provided by the residue after removal of hydroxyl from an organic acid. The term "acylamino" embraces an amino radical substituted with an acyl group. An examples of an "acylamino" radical is acetyl amino ($\text{CH}_3\text{C}(=\text{O})-\text{NH}-$).

5

[0020] Compounds of Formula I and II would be useful for, but not limited to, the treatment of inflammation in a subject, and for treatment of other inflammation-associated disorders, such as, as an analgesic in the treatment of pain and headaches, or as an antipyretic for the treatment of fever. For example, 10 compounds of Formula I and II would be useful to treat arthritis, including but not limited to rheumatoid arthritis, spondylo arthropathies, gouty arthritis, osteoarthritis, systemic lupus erythematosus, and juvenile arthritis. Such compounds of Formula I and II would be useful in the treatment of asthma, bronchitis, menstrual cramps, tendinitis, bursitis, and skin related conditions 15 such as psoriasis, eczema, burns, and dermatitis. Compounds of Formula I and II also would be useful to treat gastrointestinal conditions such as inflammatory bowel disease, Crohn's disease, gastritis, irritable bowel syndrome, and ulcerative colitis and for the prevention of colorectal cancer. Compounds of Formula I and II would be useful in treating inflammation in such diseases as 20 vascular diseases, migraine headaches, periarteritis nodosa, thyroiditis, aplastic anemia, Hodgkin's disease, sclerodoma, rheumatic fever, type I diabetes, myasthenia gravis, sarcoidosis, nephrotic syndrome, Behcet's syndrome, polymyositis, gingivitis, hypersensitivity, conjunctivitis, swelling occurring after injury, myocardial ischemia, and the like. The compounds are useful as 25 antiinflammatory agents, such as for the treatment of arthritis, with the additional benefit of having significantly less harmful side effects. The compounds of formula I or II are useful as agents for treating cancer or as an anticancer agents. The compounds of formula I or II may be proapoptotic, antiapoptotic, anticell cycle progressive, antiinvasive, and antimetastatic. More 30 specifically, the compounds of this invention are useful in the treatment of a variety of cancers including, but not limited to: carcinoma such as bladder,

breast, colon, kidney, liver, lung, including small cell lung cancer, esophagus, gall-bladder, ovary, pancreas, stomach, cervix, thyroid, prostate, and skin, including squamous cell carcinoma; hematopoietic tumors of lymphoid lineage, including leukemia, acute lymphocytic leukemia, acute lymphoblastic leukemia,

5 B-cell lymphoma, T-cell-lymphoma, Hodgkin's lymphoma, non-Hodgkin's lymphoma, hairy cell lymphoma and Burkett's lymphoma; hematopoietic tumors of myeloid lineage, including acute and chronic myelogenous leukemias, myelodysplastic syndrome and promyelocytic leukemia; tumors of mesenchymal origin, including fibrosarcoma and rhabdomyosarcoma; tumors of

10 the central and peripheral nervous system, including astrocytoma, neuroblastoma, glioma and schwannomas; other tumors, including melanoma, seminoma, teratocarcinoma, osteosarcoma, xeroderma pigmentosum, keratoanthoma, thyroid follicular cancer and Kaposi's sarcoma. Due to the key role of protein kinases in the regulation of cellular proliferation, these

15 compounds are also useful in the treatment of a variety of cell proliferative disorders such as, for instance, benign prostate hyperplasia, familial adenomatosis, polyposis, neuro-fibromatosis, psoriasis, vascular smooth cell proliferation associated with atherosclerosis, pulmonary fibrosis, arthritis glomerulonephritis and post-surgical stenosis and restenosis. The compounds of

20 formula I or II may be used as an antiviral agent. The compounds of this invention are useful as inhibitors of protein kinases. The compounds of this invention are useful as inhibitors of IKK1 and/or IKK2, IKK α /IKK β heterodimer, TBK or IKKi. The compounds of the invention may also be useful as inhibitors of other protein kinases such as, for instance, protein kinase C in

25 different isoforms, cyclin dependent kinase (cdk), Met, PAK-4, PAK-5, ZC-1, STLK-2, DDR-2, Aurora 1, Aurora 2, Bub-1, PLK, Chk1, Chk2, HER2, raf1, MEK1, MAPK, EGF-R, PDGF-R, FGF-R, IGF-R, VEGF-R, PI3K, weel kinase, Src, Abl, Akt, ILK, MK-2, IKK-2, Cdc7, Nek, and thus be effective in the treatment of diseases associated with other protein kinases. The present

30 invention preferably includes compounds, which selectively inhibit IKK2 over IKK1. Preferably, the compounds have an IKK2 IC50 of less than 1 μ M, and

have a selectivity ratio of IKK2 inhibition over IKK1 inhibition of at least 50, and more preferably of at least 100. Even more preferably, the compounds have an IKK1 IC₅₀ of greater than 10 μ M, and more preferably of greater than 100 μ M. The compounds of formula may also be used to treat angiogenesis 5 associated cardiovascular, ophthalmology and osteoporosis disorders. The compounds of the present invention may also be used for treatment of knee injury such as sport injuries.

[0021] While it is possible for an active ingredient to be administered alone as 10 the raw chemical, it is preferable to present it as a pharmaceutical formulation. The present invention comprises a pharmaceutical composition comprising a therapeutically effective amount of a compound of the present invention in association with at least one pharmaceutically acceptable carrier, adjuvant, or diluent. The present invention also comprises a method of treating inflammation 15 or inflammation associated disorders in a subject, the method comprising administering to the subject having such inflammation or disorders a therapeutically effective amount of a compound of the present invention. Also included in the family of compounds of the present invention are the pharmaceutically acceptable salts thereof. The term "pharmaceutically 20 acceptable salts" embraces salts commonly used to form alkali metal salts and to form addition salts of free acids or free bases. The nature of the salt is not critical, provided that it is pharmaceutically acceptable. Suitable pharmaceutically acceptable acid addition salts of compounds of the present invention may be prepared from an inorganic acid or from an organic acid. 25 Examples of such inorganic acids are hydrochloric, hydrobromic, hydroiodic, nitric, carbonic, sulfuric, and phosphoric acid. Appropriate organic acids may be selected from aliphatic, cycloaliphatic, aromatic, araliphatic, heterocyclic, carboxylic and sulfonic classes of organic acids, examples of which are formic, acetic, propionic, succinic, glycolic, gluconic, lactic, malic, tartaric, citric, 30 ascorbic, glucuronic, maleic, fumaric, pyruvic, aspartic, glutamic, benzoic, anthranilic, mesylic, salicyclic, salicyclic, hydroxybenzoic, phenylacetic,

mandelic, embonic (pamoic), methanesulfonic, ethanesulfonic, benzenesulfonic, pantothenic, toluenesulfonic, 2-hydroxyethanesulfonic, sulfanilic, stearic, cyclohexylaminosulfonic, algenic, β -hydroxybutyric, salicyclic, galactaric and galacturonic acid. Suitable pharmaceutically acceptable base addition salts of 5 compounds of the present invention include metallic salts made from aluminum, calcium, lithium, magnesium, potassium, sodium and zinc or organic salts made from N,N'-dibenzylethylenediamine, chloroprocaine, choline, diethanolamine, ethylenediamine, meglumine (N-methyl-glucamine) and procaine. All of these salts may be prepared by conventional means from the corresponding compound 10 of the present invention by reacting, for example, the appropriate acid or base with the compound of the present invention.

[0022] Also embraced within this invention are pharmaceutical compositions comprising one or more compounds of the present invention in association with 15 one or more non-toxic, pharmaceutically acceptable carriers and/or diluents and/or adjuvants and/or excipient (collectively referred to herein as "carrier" materials) and, if desired, other active ingredients. Accordingly, the compounds of the present invention may be used in the manufacture of a medicament.

Pharmaceutical compositions of the compounds of the present invention 20 prepared as herein before described may be formulated as solutions or lyophilized powders for parenteral administration. Powders may be reconstituted by addition of a suitable diluent or other pharmaceutically acceptable carrier prior to use. The liquid formulation may be a buffered, isotonic aqueous solution. The compounds of the present invention may be 25 administered by any suitable route, preferably in the form of a pharmaceutical composition adapted to such a route, and in a dose effective for the treatment intended. The compounds and composition may, for example, be administered intravascularly, intraperitoneally, intravenously, subcutaneously, intramuscularly, intramedullary, orally, or topically. For oral administration, the 30 pharmaceutical composition may be in the form of, for example, a tablet, capsule, suspension, or liquid. The active ingredient may also be administered

by injection as a composition wherein, for example, normal isotonic saline solution, standard 5% dextrose in water or buffered sodium or ammonium acetate solution may be used as a suitable carrier. Such formulation is especially suitable for parenteral administration, but may also be used for oral 5 administration or contained in a metered dose inhaler or nebulizer for insufflation. It may be desirable to add excipients such as polyvinylpyrrolidone, gelatin, hydroxy cellulose, acacia, polyethylene glycol, mannitol, sodium chloride, or sodium citrate. The pharmaceutical composition is preferably made in the form of a dosage unit containing a particular amount of the active 10 ingredient. Examples of such dosage units are tablets or capsules. The amount of therapeutically active compound that is administered and the dosage regimen for treating a disease condition with the compounds and/or compositions of this invention depends on a variety of factors, including the age, weight, sex and medical condition of the subject, the severity of the disease, the route and 15 frequency of administration, and the particular compound employed, and thus may vary widely. The pharmaceutical compositions may contain active ingredient in the range of about 0.1 to 2000 mg, preferably in the range of about 0.5 to 500 mg and most preferably between about 1 and 100 mg. A daily dose of about 0.01 to 100 mg/kg bodyweight, preferably between about 0.1 and about 20 50 mg/kg body weight and most preferably between about 1 to 20 mg/kg bodyweight, may be appropriate. The daily dose can be administered in one to four doses per day. For therapeutic purposes, the compounds of this invention are ordinarily combined with one or more adjuvants appropriate to the indicated route of administration. If administered orally, the compounds may be admixed 25 with lactose, sucrose, starch powder, cellulose esters of alkanoic acids, cellulose alkyl esters, talc, stearic acid, magnesium stearate, magnesium oxide, sodium and calcium salts of phosphoric and sulfuric acids, gelatin, acacia gum, sodium alginate, polyvinylpyrrolidone, and/or polyvinyl alcohol, and then tableted or encapsulated for convenient administration. Such capsules or tablets may 30 contain a controlled release formulation as may be provided in a dispersion of active compound in a sustained release material such as glyceryl monostearate,

glyceryl distearate, hydroxypropylmethyl cellulose alone or with a wax. Formulations for parenteral administration may be in the form of aqueous or non-aqueous isotonic sterile injection solutions or suspensions. These solutions and suspensions may be prepared from sterile powders or granules having one or more of the carriers or diluents mentioned for use in the formulations for oral administration. The compounds may be dissolved in water, polyethylene glycol, propylene glycol, ethanol, corn oil, cottonseed oil, peanut oil, sesame oil, benzyl alcohol, sodium chloride, and/or various buffers. The pharmaceutical preparations are made following the conventional techniques of pharmacy involving milling, mixing, granulating, and compressing, when necessary, for tablet forms; or milling, mixing and filling for hard gelatin capsule forms. When a liquid carrier is used, the preparation will be in the form of a syrup, elixir, emulsion, or an aqueous or non-aqueous suspension. Such a liquid formulation may be administered orally or filled into a soft gelatin capsule. For rectal administration, the compounds of the present invention may also be combined with excipients such as cocoa butter, glycerin, gelatin, or polyethylene glycols and molded into a suppository. The methods of the present invention include topical administration of the compounds of the present invention. By topical administration is meant non-systemic administration, including the application of a compound of the invention externally to the epidermis, to the buccal cavity and instillation of such a compound into the ear, eye, and nose, wherein the compound does not significantly enter the blood stream. By systemic administration is meant oral, intravenous, intraperitoneal, and intramuscular administration. The amount of a compound of the present invention (hereinafter referred to as the active ingredient) required for therapeutic or prophylactic effect upon topical administration will, of course, vary with the compound chosen, the nature and severity of the condition being treated and the animal undergoing treatment, and is ultimately at the discretion of the physician.

30 [0023] The topical formulations of the present invention, both for veterinary and for human medical use, comprise an active ingredient together with one or

more acceptable carriers therefore, and optionally any other therapeutic ingredients. The carrier must be "acceptable" in the sense of being compatible with the other ingredients of the formulation and not deleterious to the recipient thereof. Formulations suitable for topical administration include liquid or semi-

5 liquid preparations suitable for penetration through the skin to the site of where treatment is required such as: liniments, lotions, creams, ointments or pastes, and drops suitable for administration to the eye, ear or nose. The active ingredient may comprise, for topical administration, from 0.01 to 5.0 wt% of the formulation.

10

[0024] Drops according to the present invention may comprise sterile aqueous or oily solutions or suspensions and may be prepared by dissolving the active ingredient in a suitable aqueous solution of a bactericidal and/or fungicidal agent and/or any other suitable preservative, and preferably including a surface active agent. The resulting solution may then be clarified by filtration, transferred to a suitable container, which is then sealed and sterilized by autoclaving, or maintaining at 90-100° C for half an hour. Alternatively, the solution may be sterilized by filtration and transferred to the container by an aseptic technique. Examples of bactericidal and fungicidal agents suitable for 15 inclusion in the drops are phenylmercuric nitrate or acetate (0.00217c), benzalkonium chloride (0.0 1%) and chlorhexidine acetate (0.0 1%). Suitable solvents for the preparation of an oily solution include glycerol, diluted alcohol, and propylene glycol.

20 25 [0025] Lotions according to the present invention include those suitable for application to the skin or eye. An eye lotion may comprise a sterile aqueous solution optionally containing a bactericide and may be prepared by methods similar to those for the preparation of drops. Lotions or liniments for application to the skin may also include an agent to hasten drying and to cool the skin, such as an alcohol or acetone, and/or a moisturizer such as glycerol or an oil such as 30 castor oil or arachis oil. Creams, ointments, or pastes according to the present

invention are semi-solid formulations of the active ingredient for external application. They may be made by mixing the active ingredient in finely divided or powdered form, alone or in solution or suspension in an aqueous or non-aqueous fluid, with the aid of suitable machinery, with a greasy or non-greasy 5 basis. The basis may comprise hydrocarbons such as hard, soft or liquid paraffin, glycerol, beeswax, a metallic soap; a mucilage; an oil of natural origin such as almond, corn, arachis, castor or olive oil; wool fat or its derivatives, or a fatty acid such as stearic or oleic acid together with an alcohol such as propylene glycol or macrogols. The formulation may incorporate any suitable 10 surface-active agent such as an anionic, cationic, or non-ionic surface-active agent such as sorbitan esters or polyoxyethylene derivatives thereof. Suspending agents such as natural gums, cellulose derivatives or inorganic materials such as silicaceous silicas, and other ingredients such as lanolin may also be included. Other adjuvants and modes of administration are well and widely known in the 15 pharmaceutical art. Although this invention has been described with respect to specific embodiments, the details of these embodiments are not to be construed as limitations.

[0026] Another aspect of the present invention is chemical intermediates 20 useful in the synthesis of the compounds of Formula I and II.

[0027] Another aspect of the present invention is methods of syntheses of the compounds of Formula I and II.


25 GENERAL SYNTHETIC PROCEDURES

[0028] The starting materials used herein are commercially available or are prepared by routine methods well known to those of ordinary skill in the art and can be found in standard reference books, such as the COMPENDIUM OF 30 ORGANIC SYNTHETIC METHODS, Vol. I-VI (published by Wiley-Interscience).

[0029] The compounds of the invention can be synthesized according to the following procedures of Schemes I-X, wherein the R1-R16 substituents, linker A, are as defined for Formula I and II, above, except where further noted.


5

10 [0030] SCHEME I




Synthetic Scheme I illustrates the procedure used to prepare the antiinflammatory pyrazoles of the present invention. 1,3-Dicarbonyl compounds such as 1, or the shown enol form which is in equilibrium with the 1,3-diketone, are allowed to react with a substituted hydrazine hydrochloride 2 in warm methanol or ethanol or acetic acid to provide the pyrazoles 3 via a condensation reaction. When A = -CH₂CH₂-, the central ring may be aromatized to provide A = -CH=CH-, by using an oxidant such as DDQ, Pd or Pt on carbon with cyclooctadiene or other H₂ acceptor, or sulfur in an appropriate solvent.

[0031] SCHEME II



Synthetic Scheme II illustrates the procedure for the preparation of substituted diketones **1**. An appropriately substituted ketone **4**, including, but not limited to; 1-indanones, 1-tetralones, and 1-benzosuberones, is first treated with base, such as sodium methoxide, lithium bistrimethylsilylamine or lithium 5 diisopropylamide (LDA), followed by condensation with a suitable acylating agent, such as, dimethyl or diethyl oxalate, in an appropriate solvent, such as methanol, diethyl ether or tetrahydrofuran, to provide 1,3-dicarbonyl compounds **1** which are suitable for conversion into antiinflammatory pyrazoles as illustrated in Scheme I. Alternatively, the dicarbonyl compounds **1** can be 10 directly prepared from commercially available cyclic ketones **4**.

[0032] SCHEME III




15

Synthetic Scheme III illustrates a three-step procedure used for the preparation of substituted 1-tetralones. In step one, an appropriate substituted benzene **5** is 20 condensed with succinic anhydride and a catalyst such as aluminum chloride into the corresponding 4-phenyl-4-ketobutanoic acid derivatives **6**. In step two, the keto group of the 4-phenyl-4-ketobutanoic acids **6** is reduced using catalytic hydrogenation or Wolff-Kishner type reductions, thus providing 4-

phenylbutanoic acids **7**. In addition, ketone reductions can be carried out using metal amalgams. In step three, the 4-phenylbutanoic acids are treated with a mixture of trifluoroacetic anhydride, and trifluoroacetic acid to effect intramolecular Friedel-Crafts acylation affording selected tetralones **8**.


5 Alternatively, the Friedel-Crafts acylation can be affected with other strong acids such as polyphosphoric acid, sulfuric acid or aluminum chloride.

[0033] SCHEME IV



Synthetic Scheme IV describes an alternate synthetic route to 1-tetralones **8**. In step one, addition of allylmagnesium bromide in a suitable solvent such as, THF or diethyl ether, to an appropriately substituted benzoate **9** affords the 1-phenylbut-3-ene-1-ones **10**. In step two, the 1-phenylbut-3-ene-1-ones **10** can be cyclized under Friedel-Crafts alkylation conditions, provided R4 is a ring activating substituent, using catalysts such as aluminum chloride to produce 1-tetralones **8**.


20 [0034] SCHEME V



25 Scheme V describes the direct modification of 1-tetralone to substituted tetralones. Commercially available 1-tetralone may be treated with a variety of

electrophilic reagents such as bromine, ammonium nitrite or vinylsilanes, represented by E^+ , with or without a catalyst to generate directly a substituted tetralone **8**, containing bromo, nitro or vinyl groups. Such tetralones **8** can be further embellished to provide the desired substitution patterns. Mixtures may 5 be readily separated using chromatographic techniques.


[0035] SCHEME VI



10

An alternate to Scheme V is Scheme VI wherein an appropriately substituted decalin is subjected to electrophilic addition to generate substituted decalins **11**. Substituted decalins may also be prepared by Friedel-Crafts alkylation of substituted benzenes. Substituted decalins **11** can then be oxidized to the 15 tetralones **8** using oxidants such as $KMnO_4$ or SeO_2 .

[0036] SCHEME VII




20

Scheme VII describes the modification of existing tetralones into analogs containing differing functional groups that can also be further modified. By example, hydroxy tetralone (**8a** where $R_4 = OH$) can be converted to the triflate **8b** by treatment with trifluoromethane sulfonic anhydride. Triflate **8b** can the 25 be subjected to $Pd(OAc)_2$ an appropriate phosphine and CO in the presence of methanol to generate tetralone **12** containing a carboxy methyl group. Triflates

can be used in a variety of palladium coupling reactions to introduce additional functional groups.

[0037] SCHEME VIII


5



Synthetic Scheme VIII illustrates a three step procedure used for the preparation of substituted 1-indanones **16**. In step one, an appropriate substituted 10 benzaldehyde **13** is condensed with methyl acetate and a catalyst such as triethylamine into the corresponding methyl cinnamate derivatives **14**. Additionally, commercially available cinnamates may be used in the following steps. In step two the olefin group of the cinnamate **14** is reduced using catalytic hydrogenation and the ester hydrolyzed with base, such as NaOH, thus 15 providing 3-phenylpropanoic acids **15**. In step three, the 3-phenylpropanoic acids are treated with a mixture of trifluoroacetic anhydride and trifluoroacetic acid to effect intramolecular Friedel-Crafts acylation affording selected 1-indanones **16**. Alternatively, the Friedel-Crafts acylation can be effected with other strong acids such as sulfuric acid or aluminum chloride.


20

[0038] SCHEME IX



Synthetic Scheme IX illustrates a two-step route for the preparation of substituted 1-indanones **16**. Commercially available methyl benzoates **9**, or other alkyl esters, may be treated with a vinyl lithium reagent to afford phenylvinyl ketones **17**. Alternatively, dimethylamides or N-methyl-O-methylhydroxamides may be used in place of the esters. Also, other vinyl metals, such as; vinylmagnesium bromide may be used in place of the vinyl lithium reagent. The resulting phenylvinyl ketones may be cyclized using Friedel-Crafts alkylating catalysts, such as aluminum chloride.

[0039] SCHEME X




15

Synthetic Scheme X illustrates a three step procedure used for the preparation of substituted 1-benzosuberones **20**. In step one, an appropriate substituted benzene **5** is condensed with glutaric anhydride and a catalyst such as aluminum chloride into the corresponding 5-phenyl-5-ketopentanoic acid derivatives **18**. In step two, the keto group of the 5-phenyl-5-ketopentanoic acids **18** is reduced

using catalytic hydrogenation or Wolff-Kishner type reductions, thus providing 5-phenylpentanoic acids **19**. In addition, ketone reductions can also be carried out using metal amalgams. In step three, the 5-phenylpentanoic acids are treated with a mixture of trifluoroacetic anhydride, and trifluoroacetic acid to effect 5 intramolecular Friedel-Crafts acylation affording selected benzosuberones **20**. Alternatively, the Friedel-Crafts acylation can be affected with other strong acids such as polyphosphoric acid, H_2SO_4 or $AlCl_3$. Alternatively, 5-phenyl-5-ketopentanoic acids **18**, can be prepared from glutaric acid and a phenyllithium or a phenyl Grignard reagent appropriately substituted and compatible with 10 reaction conditions.

[0040] SCHEME XI



Scheme XI describes the general synthesis of pyrazolo[4,3-c]quinoline analogs of this present invention. In step 1, aniline was heated with acrylic acid in a solvent such toluene to give N-phenyl- β -aniline. In step two, the acid was treated with PPA at 100°C to give dihydroquinolinone. In step three, the ketone was first treated with a base such as LiHMDS, followed by diethyl oxalate to afford the intermediate. This intermediate was then reacted with a substituted phenylhydrazine to form pyrazole in step four; substituents can be sulfonamide and sulfone (aromatization occurred in this reaction presumably by β -elimination). In step five; the aromatized pyrazole was converted to amide by using ammonia in ethanol.


[0041] The complete content of all publications, patents, and patent applications cited in this disclosure are herein incorporated by reference as if each individual publication, patent, or patent application were specifically and

individually indicated to incorporated by reference. Although the foregoing invention has been described in some detail by way of illustration and example for the purposes of clarity of understanding, it will be readily apparent to one skilled in the art in light of the teachings of this invention that changes and 5 modifications can be made without departing from the spirit and scope of the present invention. The following examples are provided for exemplification purposes only and are not intended to limit the scope of the invention, which has been described in broad terms above.

10

EXAMPLES

[0042] Example 1




15

To 7-nitro-1-tetralone (4.6 g, 0.024 mol) and ethyl oxalate (3.5 mL, 0.026 mol) in ether (100 mL), was added dropwise lithium bis(trimethylsilyl)amide (1M in THF, 26 mL). The slurry was stirred overnight and filtered to give the product as an olive green solid, 6.2 g (87% yield). ¹H NMR (DMSO-*d*₆ / 300 MHz) 8.45 20 (d, 1H); 8.05 (d of d, 1H); 7.42 (d, 1H); 4.08 (q, 2H); 2.82-2.72 (m, 2H); 2.51- 2.43 (m, 2H); 1.21 (t, 3H).

[0043] Example 2

Ethyl 1-{4-[(aminothio)peroxy]phenyl}-8-nitro-4,5-dihydro-1H- 25 benzo[g]indazole-3-carboxylate




The material of Example 1 (6.2 g, 0.021 mol) and 4-sulfonamidophenylhydrazine hydrochloride (5.1 g, 0.023 mol) were stirred in methanol (100 mL) overnight. Conc. HCl (2 mL) was added to the thick slurry and the contents were heated on a steam bath for 1 hour. Contents were allowed 5 to cool and filtered to give an off-white solid, 6.9 g. NMR and LC/MS analysis show the solid to contain two components, the desired and the hydrated pyrazole. TFA (60 mL) and TFAA (20 mL) were added to the solid and heated on a steam bath for 1 hour. Contents were concentrated *in vacuo* leaving the product as a solid, 6.4 g (69% yield). FABHRMS m/z 443.1020 (M+H, 10 $C_{20}H_{19}N_4O_6S$ requires 443.1025). 1H NMR (DMSO-*d*₆ / 300 MHz) 8.10 (d of d, 1H); 8.03 (d, 2H); 7.82 (d, 2H); 7.70 (d, 1H); 7.62 (s, 1H); 7.50 (d, 1H); 4.33 (q, 2H); 3.20-2.95 (m, 4H); 1.33 (t, 3H).

Anal. Calcd for $C_{20}H_{18}N_4O_6S$: C, 54.29; H, 4.10; N, 12.66. Found: C, 54.49; H, 15 4.00; N, 12.52.

[0044] Example 3
1-{4-[(aminothio)peroxy]phenyl}-8-nitro-4,5-dihydro-1H-benzo[g]indazole-3-carboxamide

20




The material of Example 2 (718 mg, 0.0016 mol), conc. ammonium hydroxide (30 mL), and methanol (15 mL) were stirred in a stoppered flask for 72 hours. 25 Contents were filtered to give a light amber solid (606 mg). The solid was recrystallized from acetonitrile to give the product as a light amber solid, 450 mg (68% yield). FABHRMS m/z 414.0902 (M+H, $C_{18}H_{16}N_5O_5S$ requires 414.0872). 1H NMR (DMSO-*d*₆ / 300 MHz) 8.15 - 7.95 (m, 3H); 7.83 (d, 2H); 7.80-7.40 (m, 6H); 3.20-2.95 (m, 4H).

Anal. Calcd for $C_{18}H_{15}N_5O_5S$: C, 52.30; H, 3.66; N, 16.94. Found: C, 52.04; H, 3.64; N, 16.61.

5 [0045] Example 4a

ethyl 8-amino-1-{4-[(aminothio)peroxy]phenyl}-4,5-dihydro-1H-benzo[g]indazole-3-carboxylate




10 The product of Example 2 (2.0 g) and 10% Pd/C (350 mg) in DMF (20 mL) were shaken at 55 psi hydrogen for 3 hours. The catalyst was removed by filtration and the filtrate was concentrated under vacuum to yield a yellow wax. The wax was triturated with methanol and the resulting amber solid was collected by filtration to afford the title compound (1.6 g).

15

[0046] Example 4b

8-amino-1-{4-[(aminothio)peroxy]phenyl}-4,5-dihydro-1H-benzo[g]indazole-3-carboxamide



20 The product of Example 4a (968 mg), conc. ammonium hydroxide (10 mL), and methanol (5 mL) were heated at 95°C in a sealed pressure bottle for 16 h. The mixture was allowed to cool and let stand for 3 h. The resulting amber solid was collected by filtration to yield the title compound (630 mg).


25 FABHRMS m/z 384.1136 (M+H, $C_{18}H_{18}N_5O_3S$ requires 384.1130). 1H NMR (DMSO- d_6 / 300 MHz) 7.95 (d, 2H); 7.75 (d, 2H); 7.53 (br s, 1H); 7.43 (br s,

1H); 7.32 (br s, 1H); 7.01 (d, 1H); 6.44 (d of d, 1H); 6.03 (s, 1H); 4.81 (s, 2H); 2.93-2.65 (m, 4H).

5 Anal. Calcd for $C_{18}H_{17}N_5O_3S$: C, 56.38; H, 4.47; N, 18.27. Found: C, 56.31; H, 4.42; N, 18.31.

[0047] Example 4c

8-amino-1-{4-[(aminothio)peroxy]phenyl}-1H-benzo[g]indazole-3-carboxamide




The product of Example 4b (1 g) and 5% Pd/C (300 mg) suspended in cumene (50 mL) and NMP (5 mL) was heated at reflux for 16 h. The mixture was then filtered and concentrated. The residue was diluted with methanol (10 mL) and added to water, after 1 h the resulting precipitate was collected by filtration and 15 dried to yield the title compound (0.9 g).

[0048] Example 5

8-(acetylamino)-1-{4-[(aminothio)peroxy]phenyl}-4,5-dihydro-1H-benzo[g]indazole-3-carboxamide

20




To the material of Example 4b (1.0 g, 0.0026 mol) in DMF (15 mL) was added dropwise a mixture of acetic anhydride (0.283 mL, 0.003 mol) and pyridine 25 (0.243 mL, 0.003 mol) in DMF (5 mL). Contents were stirred overnight, diluted with water (75 mL), and filtered to give the desired as a white solid, 1.0 g (90% yield). FABHRMS m/z 426.1235 ($M+H$, $C_{20}H_{20}N_5O_4S$ requires 426.1236). 1H

NMR (DMSO-*d*₆ / 300 MHz) 9.80 (s, 1H); 8.00 (d, 2H); 7.75 (d, 2H); 7.60 (s, 1H); 7.48 (s, 2H); 7.39 (s, 1H); 7.30 (d, 1H); 7.15 (s, 1H); 2.90 (s, 4H); 1.92 (s, 3H).

5 Anal. Calcd for C₂₀H₁₉N₅O₄S (1H₂O): C, 54.17; H, 4.77; N, 15.79. Found: C, 54.20; H, 4.97; N, 15.77.

[0049] Example 6

10 1-{4-[(aminothio)peroxy]phenyl}-8-{[(methylthio)peroxy]amino}-4,5-dihydro-1H-benzo[g]indazole-3-carboxamide




To the material of Example 4 (1.2 g, 0.003 mol) and triethylamine (0.278 mL, 0.0035 mol) in DMF (10 mL) at 0°C, was added dropwise methanesulfonyl chloride (0.278 mL, 0.0035 mol) in CH₂Cl₂ (2 mL). Contents were stirred overnight, slowly coming to room temperature. Contents were diluted with water (50 mL) and filtered to give the product as an off-white solid, 524 mg (37% yield). FABHRMS m/z 462.0917 (M+H, C₁₉H₂₀N₅O₅S₂ requires 462.0906). ¹H NMR (DMSO-*d*₆ / 300 MHz) 9.60 (s, 1H); 7.98 (d, 2H); 7.80 (d, 2H); 7.60 (s, 1H); 7.50 (s, 2H); 7.40 (s, 1H); 7.37 (d, 1H); 7.02 (s, 1H); 6.75 (s, 1H); 2.93 (s, 4H); 2.75 (s, 3H).

25 Anal. Calcd for C₁₉H₁₉N₅O₅S₂: C, 49.45; H, 4.15; N, 15.17. Found: C, 49.19; H, 3.77; N, 15.53.

[0050] Examples 7-34

Synthesis of the sulfonamide/amide/urea library

SCHEME XII



The sulfonamides, amides, and urea were synthesized in a library format by 5 using a Bohdan reaction block. The starting materials are 8-amino-1-{4-[(aminothio)peroxy]phenyl}-4,5-dihydro-1H-benzo[g]indazole-3-carboxamide (Example 4) and appropriate sulfonyl chlorides, acyl chlorides and isocyanates. Twenty-eight reactions were done in this library.

10 The general procedure is as follows: 48 mg of 8-amino-1-{4-[(aminothio)peroxy]phenyl}-1H-benzo[g]indazole-3-carboxamide in 1 mL pyridine was placed in each reaction vessel, then 1.2 eq. of a sulfonyl chloride was added, and the mixture was shaken overnight. Then 3 mL methylene chloride and 300 mg of resin PS-trisamine were added, and then shaken over 15 night. After filtration and washing with 2 mL methanol twice, the filtrates were combined and solvents evaporated. The residue was dissolved in 2 mL dimethylformamide, and MS-TsOH resin (450 mg) was added and shaken for 48 hours. After filtration and washing with 2 mL DMF, the combined filtrate was analyzed by LC-MS and LC. Then the filtrate was evaporated on a 20 SpeedVac and the residue were suspended in 2 mL of $\text{H}_2\text{O}/\text{tBuOH}$, and lyophilized for 2 days. All compounds were obtained in solid form, and the majority of the compounds have about 90% purity. Table 1 shows the compound identification, structure, IKK heterodimer assay values (expressed as IC50), and weight (determined by Mass Spectroscopy) for the compounds from 25 the sulfonamide library.

[0051] Table 1 shows the compound identification, compound, IKK heterodimer assay values expressed as IC₅₀, formula weight, and mass spectroscopy characterization for the compounds from the library.

TABLE 1

COMPOUND	STRUCTURE	EXAMPLE	HetD	Formula Weight	Mass Spec
8-amino-1-{4-[(aminothio)peroxy]phenyl}-1H-benzo[g]indazole-3-carboxamide		Example 4c	≤ 1 μM	381.41	382
8-amino-1-{4-[(diallylamino)sulfonyl]phenyl}-1H-benzo[g]indazole-3-carboxamide		Example 7	≤ 1 μM	461.54	462
1-[4-(aminosulfonyl)phenyl]-8-[(3-methoxybenzoyl)amino]-1H-benzo[g]indazole-3-carboxamide		Example 8	≤ 1 μM	515.55	516
1-[4-(aminosulfonyl)phenyl]-8-[(3-chlorobenzoyl)amino]-1H-benzo[g]indazole-3-carboxamide		Example 9	≤ 1 μM	519.97	520
1-[4-(aminosulfonyl)phenyl]-8-[(3-(trifluoromethyl)benzoyl)amino]-1H-benzo[g]indazole-3-carboxamide		Example 10	1 ≤ 10 μM	553.52	554
1-[4-(aminosulfonyl)phenyl]-8-[(3-methylbenzoyl)amino]-1H-benzo[g]indazole-3-carboxamide		Example 11	≤ 1 μM	499.55	500
1-[4-(aminosulfonyl)phenyl]-8-[(3-bromobenzoyl)amino]-1H-benzo[g]indazole-3-carboxamide		Example 12	≤ 1 μM	564.42	565

TABLE 1 cont

COMPOUND	STRUCTURE	EXAMPLE	HetD	Formula Weight	Mass Spec
1-[4-(aminosulfonyl)phenyl]-8-[(pyridin-3-ylcarbonyl)amino]-1H-benzo[g]indazole-3-carboxamide		Example 13	≤ 1 μM	486.51	487
1-[4-(aminosulfonyl)phenyl]-8-[(2-chlorobenzoyl)amino]-1H-benzo[g]indazole-3-carboxamide		Example 14	≤ 1 μM	519.97	520
1-[4-(aminosulfonyl)phenyl]-8-[(3-bromophenyl)sulfonyl]amino-1H-benzo[g]indazole-3-carboxamide		Example 15	1 ≤ 10 μM	600.47	601
1-[4-(aminosulfonyl)phenyl]-8-[(3-chlorophenyl)sulfonyl]amino-1H-benzo[g]indazole-3-carboxamide		Example 16	1 ≤ 10 μM	556.02	557
1-[4-(aminosulfonyl)phenyl]-8-[(3-cyanobenzoyl)amino]-1H-benzo[g]indazole-3-carboxamide		Example 17	≤ 1 μM	510.53	511
1-[4-(aminosulfonyl)phenyl]-8-[(3-methylphenyl)sulfonyl]amino-1H-benzo[g]indazole-3-carboxamide		Example 18	10 ≤ 100 μM	535.6	536

TABLE 1 cont

COMPOUND	STRUCTURE	EXAMPLE	HetD	Formula Weight	Mass Spec
1-[4-(aminosulfonyl)phenyl]-8-({[3-(trifluoromethyl)phenyl]sulfonyl}amino)-1H-benzo[g]indazole-3-carboxamide		Example 19	10 ≤ 100 μM	589.57	590
8-(acetylamino)-1-[4-(aminosulfonyl)phenyl]-1H-benzo[g]indazole-3-carboxamide		Example 20	≤ 1 μM	423.45	424
1-[4-(aminosulfonyl)phenyl]-8-[(methylsulfonyl)amino]-1H-benzo[g]indazole-3-carboxamide		Example 21	≤ 1 μM	459.51	460
1-[4-(aminosulfonyl)phenyl]-8-{{[3,4-dichlorophenyl]sulfonyl}amino}-1H-benzo[g]indazole-3-carboxamide		Example 22	1 ≤ 10 μM	590.47	591
1-[4-(aminosulfonyl)phenyl]-8-{{[(2,4,5-trichlorophenyl)sulfonyl]amino}-1H-benzo[g]indazole-3-carboxamide		Example 23	1 ≤ 10 μM	624.91	625
1-[4-(aminosulfonyl)phenyl]-8-{{[3,5-bis(trifluoromethyl)phenyl]sulfonyl}amino}-1H-benzo[g]indazole-3-carboxamide		Example 24	1 ≤ 10 μM	657.57	658

TABLE 1 cont

COMPOUND	STRUCTURE	EXAMPLE	HetD	Formula Weight	Mass Spec
1-[4-(aminosulfonyl)phenyl]-8-[(2-methoxybenzoyl)amino]-1H-benzo[g]indazole-3-carboxamide		Example 25	1 ≤ 10 μM	515.55	516
1-[4-(aminosulfonyl)phenyl]-8-[(2-(trifluoromethyl)benzoyl)amino]-1H-benzo[g]indazole-3-carboxamide		Example 26	≤ 1 μM	553.52	554
1-[4-(aminosulfonyl)phenyl]-8-[(2-methylbenzoyl)amino]-1H-benzo[g]indazole-3-carboxamide		Example 27	≤ 1 μM	499.55	500
1-[4-(aminosulfonyl)phenyl]-8-[(2,6-dichlorobenzoyl)amino]-1H-benzo[g]indazole-3-carboxamide		Example 28	≤ 1 μM	554.41	555
1-[4-(aminosulfonyl)phenyl]-8-[(2-(trifluoromethoxy)benzoyl)amino]-1H-benzo[g]indazole-3-carboxamide		Example 29	≤ 1 μM	569.52	570
1-[4-(aminosulfonyl)phenyl]-8-[(2,3-dichlorobenzoyl)amino]-1H-benzo[g]indazole-3-carboxamide		Example 30	≤ 1 μM	554.41	555


TABLE 1 cont

COMPOUND	STRUCTURE	EXAMPLE	HetD	Formula Weight	Mass Spec
1-[4-(aminosulfonyl)phenyl]-8-[(2-fluorobenzoyl)amino]-1H-benzo[g]indazole-3-carboxamide		Example 31	≤ 1 μM	503.51	504
1-[4-(aminosulfonyl)phenyl]-8-[(2-chloropyridin-3-yl)carbonyl]amino]-1H-benzo[g]indazole-3-carboxamide		Example 32	≤ 1 μM	520.96	521
1-[4-(aminosulfonyl)phenyl]-8-[(2-chlorophenyl)sulfonyl]amino]-1H-benzo[g]indazole-3-carboxamide		Example 33	1 ≤ 10 μM	556.02	557
1-[4-(aminosulfonyl)phenyl]-8-(isonicotinoylamino)-1H-benzo[g]indazole-3-carboxamide		Example 34	≤ 1 μM	486.51	487

[0052] Example 35

5 1-[3-(aminosulfonyl)phenyl]-8-[(2-chlorobenzoyl)amino]-1H-benzo[g]indazole-3-carboxamide

SCHEME XIII



[0053] Step 1

ethyl 1-[3-(aminosulfonyl)phenyl]-8-nitro-4,5-dihydro-1H-benzo[g]indazole-3-

10 carboxylate

This compound was prepared using the same procedures for Ethyl 1-{4-
 15 [(aminothio)peroxy]phenyl}-8-nitro-4,5-dihydro-1H-benzo[g]indazole-3-
 carboxylate of Example 2, where 3-(aminosulfonamido)- phenylhydrazine was
 20 used in place of 4-(aminosulfonamido)- phenylhydrazine. Its structure was
 confirmed by ¹H NMR and MS (443, M+1). C₂₀H₁₈N₄O₆S, Calc.: C: 54.29, H:
 4.10, N: 12.66; Found, C: 54.27, H: 4.09, N: 12.55.

[0054] Step 2

20 8-amino-1-[3-(aminosulfonyl)phenyl]-1H-benzo[g]indazole-3-carboxamide

A sealed solution of ethyl 1-[3-(aminosulfonyl)phenyl]-8-nitro-4,5-dihydro-1H-
 benzo[g]indazole-3-carboxylat from step 1 (2.20 g, 4.98 mmol) in excess

NH₃/MeOH was heated at 110 °C for 10 h. The mixture was then concentrated to 10 mL, cooled to RT and the solid product collected by filtration. The solid was dissolved in HOAc, exposed to H₂ (5 psi) at RT in the presence of Pd-C (5%) for 2 h. The mixture was filtered through celite pad, concentrated, taken 5 into MeOH, filtered again through celite. The clear solution was concentrated, recrystallized in EtOH to give a grey solid (450 mg). Its structure was confirmed by ¹H NMR and MS (382, M+1). C₁₈H₁₅N₅O₃S.H₂O.(Et₂O)_{0.2}, Calc.: C: 54.51, H: 4.62, N: 16.91; Found, C: 54.79, H: 4.06, N: 16.72.


10 [0055] Step 3

1-[3-(aminosulfonyl)phenyl]-8-[(2-chlorobenzoyl)amino]-1H-benzo[g]indazole-3-carboxamide

To a stirred solution of 8-amino-1-[3-(aminosulfonyl)phenyl]-1H-15 benzo[g]indazole-3-carboxamide from step 2 (76.2 mg, 0.20 mmol) in pyridine (25 mL) at RT was added 2-chlorobenzoic chloride (53 mg, 0.3 mmol). After 14 h, trisamine was added and the mixture was stirred for 2 h. The mixture was filtered through a silica gel pad with EtOAc and concentrated. The resulting title product was a solid (42 mg, 40%). Its structure was confirmed by ¹H NMR and 20 MS (521, M+1). C₂₅H₁₈ClN₅O₄S, Calc.: C: 57.75, H: 3.49, N: 13.47; Found, C: 57.06, H: 3.65, N: 13.51.


[0056] Example 36

25 8-[(2-chlorobenzoyl)amino]-1-[4-(methylsulfonyl)phenyl]-1H-benzo[g]indazole-3-carboxamide



[0057] Step 1

ethyl 1-[4-(methylsulfonyl)phenyl]-8-nitro-4,5-dihydro-1H-benzo[g]indazole-3-carboxylate




5 The material of Example 1 (1.88g) was suspended in 50 ml EtOH in a round bottom flask, and then 4-methylethylsulfonylphenylhydrazine (1.2g) was added. The mixture was refluxed overnight, then the solvent evaporated, and the residue was purified by column chromatography (ethyl acetate/methylene chloride, 1/1) to yield the titled compound (0.9g, characterized by NMR, MS and HPLC).

10

[0058] Step 2

1-[4-(methylsulfonyl)phenyl]-8-nitro-4,5-dihydro-1H-benzo[g]indazole-3-carboxamide


15



The material from step 1 (0.9g) was subjected to aminolysis by using EtOH as solvent and ammonia at 110°C under 400 psi for 20 hours. After the solvent was 20 evaporated, the crude material, which was impure, was used for next step without purification or analysis.

[0059] Step 3

8-amino-1-[4-(methylsulfonyl)phenyl]-1H-benzo[g]indazole-3-carboxamide




The material from the step 2 (0.9g) was subjected to hydrogenation by using 5 DMF-AcOH as solvent, 20% Pd(OH)₂/C as catalyst under 5 psi pressure for 1 hour. The solvent was evaporated, and 1 mL conc. HCl and 30 mL water were added, and the solution was filtered. The filtrate was concentrated to about 10 mL, then purified by reverse phase HPLC (5% CH₃CN to 95% CH₃CN in water with TFA in 30 minutes) to give the titled compound (180mg), which was 10 analyzed by NMR, MS, and microanalysis confirming the structure.

[0060] Step 4

8-[(2-chlorobenzoyl)amino]-1-[4-(methylsulfonyl)phenyl]-1H-benzo[g]indazole-3-carboxamide

15



The material from step 3 (102mg) was dissolved in 10 mL pyridine, then 2-chlorobenzoyl chloride (61mg) in 2 mL methylene chloride was added. The 20 mixture was stirred at room temperature for 2 hours, then MeOH (3 mL) and PS-trisamine resin (1g) were added, and stirred overnight. After filtration and solvent evaporation, the residue obtained was suspended in tBuOH and water,

and then lyophilized. The title compound (110 mg), analyzed by NMR, MS, and microanalysis, was obtained.

[0061] Examples 37-42

5


[0062] The compounds of Examples 37-42 shown in Table 2 were synthesized in a similar manner as described in Examples 35 and 36.

Table 2

Compound No., Structure	Mol. Wt.	Compound Name(s)	IKK2 Resin IC 50	LCMS (M+H)	Example
	518.98	8-[(2-chlorobenzoyl)amino]-1-[4-(methylsulfonyl)phenyl]-1H-benzo[g]indazole-3-carboxamide	$\leq 1 \mu\text{M}$	519	37
	498.56	8-[(2-methylbenzoyl)amino]-1-[4-(methylsulfonyl)phenyl]-1H-benzo[g]indazole-3-carboxamide	$\leq 1 \mu\text{M}$	499	38
	552.53	1-[4-(methylsulfonyl)phenyl]-8-[(2-(trifluoromethyl)benzoyl)amino]-1H-benzo[g]indazole-3-carboxamide	$1 \leq 10 \mu\text{M}$	553	39
	553.42	8-[(2,3-dichlorobenzoyl)amino]-1-[4-(methylsulfonyl)phenyl]-1H-benzo[g]indazole-3-carboxamide	$\leq 1 \mu\text{M}$	554	40
	519.97	8-[(2-chloropyridin-3-yl)carbonyl]amino]-1-[4-(methylsulfonyl)phenyl]-1H-benzo[g]indazole-3-carboxamide	$\leq 1 \mu\text{M}$	520	41

Compound No., Structure	Mol. Wt.	Compound Name(s)	IKK2 Resin IC 50	LCMS (M+H)	Example
	502.52	8-[(2-fluorobenzoyl)amino]-1-[4-(methylsulfonyl)phenyl]-1H-benzo[g]indazole-3-carboxamide	1 ≤ 10 μM	503	42

[0063] Example 43



5 [0064] Step 1


ethyl 1-(4-fluorophenyl)-8-nitro-4,5-dihydro-1H-benzo[g]indazole-3-carboxylate



10 [0065] This compound was synthesized in a similar manner as described in Example 36 step 1 using 4-fluorophenylhydrazine and the material of Example 1.

[0066] Step 2

15 ethyl 8-amino-1-(4-fluorophenyl)-4,5-dihydro-1H-benzo[g]indazole-3-carboxylate



20 This compound was synthesized in a similar manner as described in Example 4a.

[0067] Step 3

8-amino-1-(4-fluorophenyl)-4,5-dihydro-1H-benzo[g]indazole-3-carboxamide




5

This compound was synthesized in a similar manner as described in Example 4b.

[0068] Step 4

10 8-amino-1-(4-fluorophenyl)-1H-benzo[g]indazole-3-carboxamide




The product of step 3 (2g) was stirred in cumene (150 mls) and warmed to 75 degrees to effect solution. Pd/C (5%; 1g) was added and the contents were 15 refluxed overnight under Ar. The reaction mixture was filtered hot thru a fritted filter and the Pd/C was washed with hot ethyl acetate. The filtrate was evaporated and the residue was triturated with toluene, filtered, and air dried to give 1.9 g (90%) of titled material. LCMS and NMR are consistent for the titled material.

20

[0069] Step 5


8-[(3-chloroisonicotinoyl)amino]-1-(4-fluorophenyl)-1H-benzo[g]indazole-3-carboxamide



3-chloro-4-pyridylcarboxylic acid (125 mg; .00078 moles) was dissolved in DMF (6 mls) and diisopropylethylamine (110 mg; 1.2 equivalents) and HATU (330 mg; 1.1 equivalents) were added and the contents were stirred at room temperature under N₂ for 30 minutes. The material of step 4 of Example 43 (250 mg; .00078 moles) was added and the mixture was stirred overnight at room temperature. Contents of the reaction mixture were poured into water (30 mls) with stirring, the precipitate was filtered, and air dried. The crude product was recrystallized from EtOH/water to give the purified product (300 mg; 84%). LCMS and NMR were consistent for the titled material. C₂₄H₁₅FCIN₅O₂. MW=459.87. Calc:C, 62.68; H, 3.29; N, 15.23. Found: C, 62.29; H, 3.51; N, 15.01.

15 [0070] Synthesis of Examples 44 and 45

SCHEME XIV



[0071] Example 44

The title compound of Example 44 (460 mg) is suspended in 5ml of DMF, then bromine (340 mg) in 15 ml of acetic acid was added in two portions, and the reaction mixture was stirred at room temperature for 15 minutes. After 5 evaporation of the solvent, the residue was suspended in aq. NaHCO₃, then filtered and washed with water, ether, acetonitrile. After drying under vacuum, 0.37g of desired compound was obtained and characterized by HPLC, LC-MS, ¹HMR. NOE study confirmed the substitution position.

[0072] Example 45

The title compound of Example 44 (0.27g) is suspended in DME (5ml) and 5 water (1ml). Under N₂, Pd(PPh₃)₄ (50mg), Na₂CO₃ (426mg) and 3-pyridylboronic acid (130mg) are added. The reaction mixture was placed in CEM Discover microwave reactor at 120C for 60 minutes with 150W power. After the reaction was completed, and solvent evaporated, the residue was dissolved in 30 ml MeOH and 5 drops of conc. HCl, and purified by reversed 10 phase HPLC(20% CH₃CH to 80% CH₃CN in 30 minutes). 80 mg of desired compound was obtained and characterized by LC-MS, ¹H NMR, CHN analysis, and HPLC purity analysis.

[0073] Examples 46-53

15

The compounds of Examples 46-53 shown in Table 3 were synthesized as in Examples 44 and 45 using the appropriate aryl, substituted aryl, heteroaryl, substituted heteroaryl, substituted arylalkyl, substituted heteroarylalkyl, or cycloalkyl.

20

[0074] The bioactivity in the IKK2 Resin assay for the compounds of Examples 43-53 is shown in Table 3.

Table 3

Compound No., Structure	Mol. Wt.	Compound Name(s)	IKK2 Resin IC 50	LCMS (M+H)	Example
	459.87	8-[(3-chloroisonicotinoyl)amino]-1-(4-fluorophenyl)-1H-benzo[g]indazole-3-carboxamide	≤1 μM	460	43


Compound No., Structure	Mol. Wt.	Compound Name(s)	IKK2 Resin IC 50	LCMS (M+H)	Example
	538.77	5-bromo-8-[(3-chloroisonicotinoyl)amino]-1-(4-fluorophenyl)-1H-benzo[g]indazole-3-carboxamide	$\leq 1 \mu\text{M}$	539	44
	609.88	8-[(3-chloroisonicotinoyl)amino]-1-(4-fluorophenyl)-5-pyridin-3-yl-1H-benzo[g]indazole-3-carboxamide dihydrochloride	$1 \leq 10 \mu\text{M}$	610	45
	320.33	8-amino-1-(4-fluorophenyl)-1H-benzo[g]indazole-3-carboxamide	$10 \leq 100 \mu\text{M}$	321	46
	458.88	8-[(2-chlorobenzoyl)amino]-1-(4-fluorophenyl)-1H-benzo[g]indazole-3-carboxamide	$1 \leq 10 \mu\text{M}$	459	47
	320.33	8-amino-1-(3-fluorophenyl)-1H-benzo[g]indazole-3-carboxamide	$\leq 20 \mu\text{M}$	321	48
	458.88	8-[(2-chlorobenzoyl)amino]-1-(3-fluorophenyl)-1H-benzo[g]indazole-3-carboxamide	$\leq 50 \mu\text{M}$	459	49
	398.42	1-(4-fluorophenyl)-8-[(methylsulfonyl)amino]-1H-benzo[g]indazole-3-carboxamide	$\leq 1 \mu\text{M}$	399	50
	459.87	8-{[(2-chloropyridin-3-yl)carbonyl]amino}-1-(4-fluorophenyl)-1H-benzo[g]indazole-3-carboxamide	$\leq 1 \mu\text{M}$	460	51
	477.86	8-{[(2-chloropyridin-3-yl)carbonyl]amino}-1-(2,4-difluorophenyl)-1H-benzo[g]indazole-3-carboxamide	$\leq 1 \mu\text{M}$	478	52

Compound No., Structure	Mol. Wt.	Compound Name(s)	IKK2 Resin IC 50	LCMS (M+H)	Example
	446.49	1-(4-fluorophenyl)-8-[(4-methylpiperazin-1-yl)carbonyl]amino]-1H-benzo[g]indazole-3-carboxamide	≤ 20	447	53

[0075] Example 54


1-(1,3-benzodioxol-5-yl)-8-[(2-chlorobenzoyl)amino]-1H-benzo[g]indazole-3-carboxamide

5



[0076] Step 1


10 ethyl 1-(1,3-benzodioxol-5-yl)-8-nitro-4,5-dihydro-1H-benzo[g]indazole-3-carboxylate



This compound was synthesized in a similar manner as described in Examples 36 step 1 using 3,4-methylenedioxophenylhydrazine and the material of 15 Example 1.

[0077] Step 2


ethyl 8-amino-1-(1,3-benzodioxol-5-yl)-4,5-dihydro-1H-benzo[g]indazole-3-carboxylate



5 This compound was synthesized in a similar manner as described in Example
4a.

[0078] Step 3

8-amino-1-(1,3-benzodioxol-5-yl)-4,5-dihydro-1H-benzo[g]indazole-3-
10 carboxamide




This compound was synthesized in a similar manner as described in Example
4b.

15

[0079] Step 4


8-amino-1-(1,3-benzodioxol-5-yl)-1H-benzo[g]indazole-3-carboxamide



[0080] This compound was synthesized in a similar manner as described in Examples 36 step 4.

5 [0081] Step 5

1-(1,3-benzodioxol-5-yl)-8-[(3-chloroisocitonoyl)amino]-1H-benzo[g]indazole-3-carboxamide



10

This compound was synthesized in a similar manner as described in Example 36 step 5.

[0082] The compounds of Examples 54-58 shown in Table 4 were 15 synthesized in a similar manner as described in Examples 35, 36 and 44.

Table 4

Compound No., Structure	Mol. Wt.	Compound Name(s)	IKK2 Resin IC 50	LCMS (M+H)	Example
	484.90	1-(1,3-benzodioxol-5-yl)-8-[(2-chlorobenzoyl)amino]-1H-benzo[g]indazole-3-carboxamide	≤1 μM	485	54
	485.89	1-(1,3-benzodioxol-5-yl)-8-[(2-chloropyridin-3-yl)carbonyl]amino]-1H-benzo[g]indazole-3-carboxamide	≤1 μM	486	55

Compound No., Structure	Mol. Wt.	Compound Name(s)	IKK2 Resin IC 50	LCMS (M+H)	Example
	346.35	8-amino-1-(1,3-benzodioxol-5-yl)-1H-benzo[g]indazole-3-carboxamide	1 ≤ 10 μM	347	56
	564.79	1-(1,3-benzodioxol-5-yl)-5-bromo-8-[(2-chloropyridin-3-yl)carbonyl]amino-1H-benzo[g]indazole-3-carboxamide	≤1 μM	565	57
	485.89	1-(1,3-benzodioxol-5-yl)-8-[(3-chloroissonicotinoyl)amino]-1H-benzo[g]indazole-3-carboxamide	≤1 μM	486	58

BIOLOGICAL EVALUATION

[0083] *Materials*

5

SAM² TM 96 Biotin capture plates were from Promega. Anti-FLAG affinity resin, FLAG-peptide, NP-40 (Nonidet P-40), BSA, ATP, ADP, AMP, LPS (*E. coli* serotype 0111:B4), and dithiothreitol were obtained from Sigma Chemicals. Antibodies specific for NEMO (IKK γ) (FL-419), IKK1(H-744), IKK2(H-470) 10 and IkB α (C-21) were purchased from Santa Cruz Biotechnology. Ni-NTA resin was purchased from Qiagen. Peptides were purchased from American Peptide Company. Protease inhibitor cocktail tablets were from Boehringer Mannheim. Sephadryl S-300 column was from Pharmacia LKB Biotechnology. Centriprep-10 concentrators with a molecular weight cutoff of 10 kDa and 15 membranes with molecular weight cut-off of 30 kDa were obtained from Amicon. [γ -³³P] ATP (2500 Ci/mmol) and [γ -³²P] ATP (6000 Ci/mmol) were purchased from Amersham. The other reagents used were of the highest grade commercially available.

20 [0084] *Cloning and Expression*

cDNAs of human IKK1 and IKK2 were amplified by reverse transcriptase-polymerase chain reaction from human placental RNA (Clonetech). hIKK1 was subcloned into pFastBac HTa (Life Technologies) and expressed as N-terminal His₆-tagged fusion protein. The hIKK2 cDNA was amplified using a reverse oligonucleotide primer which incorporated the peptide sequence for a FLAG-epitope tag at the C-terminus of the IKK2 coding region (DYKDDDDKD). The hIKK2:FLAG cDNA was subcloned into the baculovirus vector pFastBac. The rhIKK2 (S177S, E177E) mutant was 25 constructed in the same vector used for wild type rhIKK2 using a QuikChangeTM mutagenesis kit (Stratagene). Viral stocks of each construct 30

were used to infect insect cells grown in 40L suspension culture. The cells were lysed at a time that maximal expression and rhIKK activity were demonstrated. Cell lysates were stored at -80 °C until purification of the recombinant proteins was undertaken as described below.

5

[0085] *Enzyme Isolation*

All purification procedures were carried out at 4 °C unless otherwise noted. Buffers used are: buffer A: 20 mM Tris-HCl, pH 7.6, containing 50 mM NaCl, 10 20 mM NaF, 20 mM β -Glycerophosphate, 500 uM sodiumortho-vanadate, 2.5 mM metabisulfite, 5 mM benzamidine, 1 mM EDTA, 0.5 mM EGTA, 10% glycerol, 1 mM DTT, 1X Complete™ protease inhibitors; buffer B: same as buffer A, except 150 mM NaCl, and buffer C: same as buffer A, except 500 mM NaCl.

15

[0086] *Isolation of rhIKK1 homodimer*

Cells from an 8 liter fermentation of baculovirus-expressed IKK1 tagged with His peptide were centrifuged and the cell pellet (MOI 0.1, I=72 hr) was re- 20 suspended in 100 ml of buffer C. The cells were microfluidized and centrifuged at 100,000 X g for 45 min. The supernatant was collected, imidazole added to the final concentration of 10 mM and incubated with 25 ml of Ni-NTA resin for 2 hrs. The suspension was poured into a 25 ml column and washed with 250 ml of buffer C and then with 125 ml of 50 mM imidazole in buffer C. rhIKK1 25 homodimer was eluted using 300 mM imidazole in buffer C. BSA and NP-40 were added to the enzyme fractions to the final concentration of 0.1 %. The enzyme was dialyzed against buffer B, aliquoted and stored at -80 °C.

[0087] *Isolation of rhIKK2 homodimer*

30

A 10 liter culture of baculovirus-expressing IKK2 tagged with FLAG peptide was centrifuged and the cell pellet (MOI=0.1 and I=72 hrs) was re-suspended in buffer A. These cells were microfluidized, and centrifuged at 100,000 X g for 45 min. Supernatant was passed over a G-25 column equilibrated with Buffer 5 A. Protein peak was collected and incubated with anti-FLAG affinity resin on a rotator overnight in buffer B. The resin was washed in batch with 10-15 bed volumes of buffer C. Washed resin was poured into a column and rhIKK2 homodimer was eluted using 5 bed volumes of buffer B containing FLAG peptide. 5 mM DTT, 0.1% NP-40 and BSA (concentrated to 0.1% in final 10 amount) was added to the eluted enzyme before concentrating in using an Amicon membrane with a molecular weight cut-off of 30 kDa. Enzyme was aliquoted and stored at -80 °C.

[0088] *Isolation of rhIKK1/IKK2 heterodimer*

15

The heterodimer enzyme was produced by coinfection in a baculovirus system (FLAG IKK2/IKK1 His; MOI=0.1 and I=72 hrs). Infected cells were centrifuged and the cell pellet (10.0 g) was suspended in 50 ml of buffer A. The protein suspension was microfluidized and centrifuged at 100,000 X g for 45 20 min. Imidazole was added to the supernatant to a final concentration of 10 mM. The protein was allowed to bind 25 ml of Ni-NTA resin by mixing for 2 hrs. The protein-resin slurry was poured into a 25 ml column and washed with 250 ml of buffer A containing 10 mM imidazole followed by 125 ml of buffer A containing 50 mM imidazole. Buffer A, containing 300 mM imidazole, was 25 then used to elute the protein. A 75 ml pool was collected and NP-40 was added to a final concentration of 0.1%. The protein solution was then dialyzed against buffer B. The dialyzed heterodimer enzyme was then allowed to bind to 25 ml of anti-FLAG M2 agarose affinity gel overnight with constant mixing. The protein-resin slurry was then centrifuged for 5 min at 2,000 rpm. The 30 supernatant was collected and the resin re-suspended in 100 ml of buffer C containing 0.1% NP-40. The resin was washed with 375 ml of buffer C

containing 0.1 % NP-40. The protein-resin was poured into a 25 ml column and the enzyme eluted using buffer B containing FLAG peptide. Enzyme fractions (100 ml) were collected and concentrated to 20 ml using an Amicon membrane with molecular weight cut-off of 30 kDa. Bovine serum albumin was added to 5 the concentrated enzyme to final concentration of 0.1 %. The enzyme was then aliquoted and stored at -80 °C.

[0089] *Cell Culture*

10 The wild type (wt) human pre-B cell line, 70Z/3, and its mutant, 1.3E2, were generously provided by Dr. Carol Sibley. Wt 70Z/3 and 1.3E2 cells were grown in RPMI 1640 (Gibco) supplemented with 7 % defined bovine serum (Hyclone) and 50 µM 2-mercaptoethanol. Human monocytic leukemia THP-1 cells, obtained from ATCC, were cultured in RPMI 1640 supplemented with 10% 15 defined bovine serum, 10 mM HEPES, 1.0 mM sodium pyruvate and 50 µM 2-mercaptoethanol. For experiments, cells were plated in 6 well plates at 1x10⁶ cells/ml in fresh media. Pre-B cells were stimulated by the addition of 10 µg/ml LPS for varying lengths of time ranging from 0-4 hr. THP-1 cells were stimulated by the addition of 1 µg/ml LPS for 45 minutes. Cells were pelleted, 20 washed with cold 50 mM sodium phosphate buffer, pH 7.4 containing 0.15 M NaCl and lysed at 4 °C in 20 mM Hepes buffer, pH 7.6 containing 50 mM NaCl, 1 mM EDTA, 1 mM EGTA, 1 mM sodium orthovanadate, 10 mM β-glycerophosphate, 1 mM NaF, 1 mM PMSF, 1 mM DTT and 0.5 % NP40 (lysis buffer). The cytosolic fractions obtained following centrifugation at 10,000 X g 25 were stored at -80° C until used.

[0090] *Immunoprecipitation and Western Blotting*

30 SF9 cells paste containing rhIKKs were centrifuged (100,000 X g, 10 min) to remove debris. rhIKKs were immunoprecipitated (100 µg of cell paste) from the cell supernatant using 3 µg of anti-NEMO antibody (FL-419), followed by

coupling to protein A sepharose beads. rhIKKs were also immunoprecipitated from affinity chromatography purified protein preparations (1 μ g) using anti-FLAG, anti-His or anti-NEMO antibodies (1-4 μ g) followed by protein A sepharose coupling. The native, human IKK complex was immunoprecipitated 5 from THP-1 cell homogenates (300 μ g/condition) using the anti-NEMO antibody. Immune complexes were pelleted and washed 3 times with 1 ml cold lysis buffer. Immunoprecipitated rhIKKs were chromatographed by SDS-PAGE (8% Tris-glycine) and transferred to nitrocellulose membranes (Novex) and detected by chemiluminescence (SuperSignal) using specific anti-IKK 10 antibodies (IKK2 H-470, IKK1 H-744). Native IKK2, I κ B α and NEMO proteins from cytosolic lysates (20-80 μ g) were separated by SDS-PAGE and visualized by chemiluminescence using specific antibodies.

[0091] *Phosphatase Treatment*

15 Immunoprecipitated rhIKKs were washed 2 times in 50 mM Tris-HCl, pH 8.2 containing 0.1 mM EDTA, 1 mM DTT, 1 mM PMSF and 2 mM MnCl₂ and resuspended in 50 μ l. Phosphatase (λ PPase, 1000 U) was pre-diluted in the same buffer and added to the IKK samples. Following an incubation at room 20 temperature for 30 minutes with intermittent mixing, cold lysis buffer was added to the tubes to stop the reaction. After several washes, 10 % of the beads were removed for Western analysis, and the remaining material was pelleted and resuspended in 100 μ l of the buffer used for the *in vitro* kinase assay.

25 **[0092] *IKK α SAM Enzyme Assay***

IKK α kinase activity was measured using a biotinylated I κ B α peptide (Gly-Leu-Lys-Lys-Glu-Arg-Leu-Leu-Asp-Asp-Arg-His-Asp-Ser₃₂-Gly-Leu-Asp-Ser₃₆-Met-Lys-Asp-Glu-Glu), a SAM² TM 96 Biotin capture plate, and a vacuum 30 system. The standard reaction mixture contained 5 μ M biotinylated I κ B α peptide, 1 μ M [γ -³³P] ATP (about 1 X 10⁵ cpm), 1 mM DTT, 50 mM KCl, 2

mM MgCl₂, 2 mM MnCl₂, 10 mM NaF, 25 mM Hepes buffer, pH. 7.6 and enzyme solution (1-10 μ l) in a final volume of 50 μ l. After incubation at 25 °C for 30 min, 25 μ l of the reaction mixture was withdrawn and added to a SAM²™ 96 Biotin capture 96-well plate. Each well was then washed successively with 800 μ l 2 M NaCl, 1.2 ml of NaCl containing 1% H₃PO₄, 400 μ l H₂O, and 200 μ l 95% ethanol. The plate was allowed to dry in a hood at 25 °C for 1 hr and then 25 μ l of scintillation fluid (Microscint 20) was added to each well. Incorporation of [γ -³³P] ATP was measured using a Top-Count NXT (Packard). Under each assay condition, the degree of phosphorylation of I κ B α peptide substrate was linear with time and concentration for all purified enzymes. Results from the biotinylated peptide assay were confirmed by SDS-PAGE analysis of kinase reaction utilizing a GST-I κ B α ₁₋₅₄ and [γ -³²P] ATP. The resulting radiolabeled substrate was quantitated by Phosphoimager (Molecular Dynamics). An ion exchange resin assay was also employed using [γ -³³P] ATP and GST-I κ B α ₁₋₅₄ fusion protein as the substrates. Each assay system yielded consistent results in regard to K_m and specific activities for each of the purified kinase isoforms. One unit of enzyme activity was defined as the amount required to catalyze the transfer of 1 nmole of phosphate from ATP to I κ B α peptide per min. Specific activity was expressed as units per mg of protein. For experiments related to K_m determination of purified enzymes, various concentrations of ATP or I κ B α peptide were used in the assay at either a fixed I κ B α or ATP concentration. For I κ B α peptide K_m, assays were carried out with 0.1 μ g of enzyme, 5 μ M ATP and I κ B α peptide from 0.5 to 20 μ M. For ATP K_m, assays were carried out with 0.1 μ g of enzyme, 10 μ M I κ B α peptide and ATP from 0.1 to 10 μ M. For K_m determination of rhIKK1 homodimer, due to its low activity and higher K_m for I κ B α peptide, rhIKK1 homodimer (0.3 μ g) was assayed with 125 μ M I κ B α peptide and a 5-fold higher specific activity of ATP (from 0.1 to 10 μ M) for ATP K_m experiments and a 5-fold higher specific activity of 5 μ M ATP and I κ B α peptide (from 5 to 200 μ M) for I κ B α peptide K_m experiments.

[0093] IKK β Resin Enzyme Assay

IKK β kinase activity was measured using a biotinylated I κ B α peptide (Gly-Leu-
5 Lys-Lys-Glu-Arg-Leu-Leu-Asp-Asp-Arg-His-Asp-Ser₃₂-Gly-Leu-Asp-Ser₃₆-
Met-Lys-Asp-Glu-Glu) (American Peptide Co.). 20 μ l of the standard reaction
mixture contained 5 μ M biotinylated I κ B α peptide, 0.1 μ Ci/reaction [γ -³³P]
ATP (Amersham) (about 1 X 10⁵ cpm), 1 μ M ATP (Sigma), 1 mM DTT
(Sigma), 2 mM MgCl₂ (Sigma), 2 mM MnCl₂ (Sigma), 10 mM NaF (Sigma), 25
10 mM Hepes (Sigma) buffer, pH 7.6 and 20 μ l enzyme solution and 10 μ l
inhibitor in a final volume of 50 μ l. After incubation at 25 °C for 30 min, 150
 μ l resin (Dowex anion-exchange resin AG1X8 200-400 mesh) in 900 mM
formate, pH 3.0 was added to each well to stop the reaction. Resin was allowed
to settle for one hour and 50 μ l of supernatant was removed to a Micolite-2 flat
15 bottom plate (Dynex). 150 μ l of scintillation fluid (Microscint 40) (Packard)
was added to each well. Incorporation of [γ -³³P] ATP was measured using a
Top-Count NXT (Packard).


[0094] IKK heterodimer Resin Enzyme Assay

20 IKK heterodimer kinase activity was measured using a biotinylated I κ B α
peptide (Gly-Leu-Lys-Lys-Glu-Arg-Leu-Leu-Asp-Asp-Arg-His-Asp-Ser₃₂-Gly-
Leu-Asp-Ser₃₆-Met-Lys-Asp-Glu-Glu) (American Peptide Co.). 20 μ l of the
standard reaction mixture contained 5 μ M biotinylated I κ B α peptide, 0.1
25 μ Ci/reaction [γ -³³P] ATP (Amersham) (about 1 X 10⁵ cpm), 1 μ M ATP
(Sigma), 1 mM DTT (Sigma), 2 mM MgCl₂ (Sigma), 2 mM MnCl₂ (Sigma), 10
mM NaF (Sigma), 25 mM Hepes (Sigma) buffer, pH 7.6 and 20 μ l enzyme
solution and 10 μ l inhibitor in a final volume of 50 μ l. After incubation at 25
°C for 30 min, 150 μ l resin (Dowex anion-exchange resin AG1X8 200-400
30 mesh) in 900 mM formate, pH 3.0 was added to each well to stop the reaction.

Resin was allowed to settle for one hour and 50 μ l of supernatant was removed to a Micolite-2 flat bottom plate (Dynex). 150 μ l of scintillation fluid (Microscint 40) (Packard) was added to each well. Incorporation of [γ - 33 P] ATP was measured using a Top-Count NXT (Packard).

WHAT IS CLAIMED IS:

1. A compound of formula I



5

wherein

A is $(CH_2)_m-CR^{15}=CR^{16}-(CH_2)_m$ **m** is independently selected from 0, 1, or 2;

10 **B** is a 5 or 6 membered heteroaryl, aryl, saturated or unsaturated heterocyclic wherein said aryl, heteroaryl, or heterocyclic are optionally substituted with **R**¹, **R**², and **R**¹²;

X is selected from the group consisting of: N and C;

15 **Y** and **Z** are independently selected from the group consisting of: N, CH, CR³, S, and O;

R¹ is selected from the group consisting of: hydrido, halogen, alkyl, aryl, heteroaryl, alkenyl, alkynyl, haloalkyl, CN, NO₂, OR⁵, OCOOR⁵, CO₂R⁷, CON(R⁶)R⁷, COR⁶, SR⁶, SOR⁶, SO₂R⁶, NR⁶R⁷, NR⁶COR⁷, NR⁶CONHR⁷, NR⁶SO₂R⁷, NR⁶SO₂NHR⁷, and SO₂N(R⁶)R⁷ wherein R⁶

20 and R⁷ may be taken together to form a 3-7 membered carbocyclic ring having 1 to 3 substituted or unsubstituted heteroatoms selected from the group consisting of: S, SO, SO₂, O, and NR⁶; wherein said alkenyl, alkynyl, alkyl, aryl, heteroaryl or OR⁵ are optional substituted with, hydrido, halogen, alkyl, hydroxyalkyl, aryl, heteroaryl, haloalkyl, COCF₃, CN, NO₂, OR⁵, OCOOR⁵, CO₂R⁷, CON(R⁶)R⁷, COR⁶, SR⁶, SOR⁶, SO₂R⁶, NR⁶R⁷, NR⁶COR⁷, NR⁶CONHR⁷, NR⁶SO₂R⁷,

25

NR⁶SO₂NHR⁷, and SO₂N(R⁶)R⁷ wherein R⁶ and R⁷ may be taken together to form a 3-7 membered carbocyclic ring having 1 to 3 substituted or unsubstituted heteroatoms selected from the group consisting of: S, SO, SO₂, O, and NR⁶;

5 R² is selected from the group consisting of: halogen, hydrido, hydroxyalkyl, alkyl, OR⁶, CN, NO₂, SR⁶, NHR⁶, CON(R⁶)R⁷, NHCONHR⁶, CO₂H, and haloalkyl;

R¹ and R² may be taken together to form a 5 to 7 membered saturated or unsaturated carbocyclic ring optionally containing 0 to 3 heteroatoms selected from the group consisting of N, O, or S, and wherein said ring is optionally substituted with R¹;

10 R³ is selected from the group consisting of: substituted or unsubstituted amidine, alkylamino, aminoalkyl, CONHR⁷, NH₂, NHCOR⁶, and CH₂NHCOR⁶;

15 R⁴ is selected from the group consisting of: halogen, alkylsulfinyl, alkylsulfonyl, cyano, alkoxy carbonyl, alkyl, haloalkyl, hydrido, hydroxyalkyl, haloalkoxy, heterocyclic, nitro, acylamino, aryl, heteroaryl, and alkenyl, OR¹³, SR⁸, SO₂N(R⁸)R^{8'}, NHR⁹, NHCOR⁹, NR⁹COR⁹, NHCO(OR⁹), NR⁹CO(OR⁹), NR⁸SO₂R¹⁰, NHSO₂N(R¹⁰)R^{10'}, NR⁶CON(R¹⁰)R^{10'}, COR⁹, CO₂R⁸, CON(R⁸)R^{8'}, wherein R⁸ and R^{8'} may be taken together to form a 3-7 membered carbocyclic ring having 1 to 3 substituted or unsubstituted heteroatoms selected from S, SO, SO₂, O, N, and NR⁶, and wherein R¹⁰ and R^{10'} may be taken together to form a 3-7 membered carbocyclic ring having 1 to 3 substituted or unsubstituted heteroatoms selected from S, SO, SO₂, O, N, and NR⁶ wherein said aryl, heterocyclic, heteroaryl, or alkenyl are optionally substituted with R⁹;

20 R⁵ is selected from the group consisting of: hydrido, alkyl, aryl, arylalkyl, heteroaryl, heterocyclicalkyl, and heteroarylalkyl, wherein aryl, alkyl, arylalkyl, heteroaryl, heterocyclicalkyl, or heteroarylalkyl are optionally substituted with one or more radicals selected from the group consisting of OR¹⁴, N(R¹⁴)R^{14'}, and glycols;

25

30

R⁶ is independently selected from the group consisting of: hydrido, aryl, heteroaryl, lower alkyl, haloalkyl, alkenyl, alkynyl, hydroxyalkyl, aminoalkyl, alkylaminoalkyl, alkoxy, alkoxyalkyl, heterocyclicalkyl, and heterocyclic;

5 **R⁷** is independently selected from the group consisting of: hydrido, aryl, heteroaryl, lower alkyl, haloalkyl, alkenyl, alkynyl, hydroxyalkyl, aminoalkyl, alkylaminoalkyl, alkoxy, alkoxyalkyl, heterocyclicalkyl, and heterocyclic;

10 **R⁸** is independently selected from the group consisting of: hydrido, aryl, heteroaryl, arylalkyl, heterocyclic, haloalkyl, arylalkylamino, alkylaminoalkyl, dialkylaminoalkyl, alkyl, alkenyl, alkynyl, heteroarylalkyl, and heterocyclicalkyl;

15 **R^{8'}** is independently selected from the group consisting of: hydrido, aryl, heteroaryl, arylalkyl, heterocyclic, haloalkyl, arylalkylamino, alkylaminoalkyl, dialkylaminoalkyl, alkyl, alkenyl, alkynyl, heteroarylalkyl, and heterocyclicalkyl;

20 **R⁹** is independently selected from the group consisting of: hydrido, lower alkyl, aryl, heteroaryl, arylalkyl, heterocyclic, cycloalkyl, heterocyclicalkyl, haloalkyl, arylalkylamino, amino, aminoalkyl, aminoacyl, nitro, azido, and heteroarylalkyl, wherein alkyl, aryl, heteroaryl, aminoalkyl, or arylalkyl are optionally substituted with one or more radical selected from the group consisting of: alkylsulfonamide, sulfamyl, alkyl, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, aminoalkyl, alkylaminoalkyl, alkoxy, halogen, acyloxy, oxy, formyl, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, nitro, azido, benzyloxy, dialkylaminoacyl, thioalkyl, aminoacyloxy, thiocyanate, isothiocyanate, alkyldioxy, hydroxyalkyl, alkylamino, alkyloxycarbonyl, alkoxyalkyl, alkenylamino, alkynylamino, alkenyl, alkynyl, dialkylaminoalkyloxy, and heterocyclic optionally substituted with alkyl, alkylamino, aminoalkyl, hydroxyalkyl, and alkylaminoalkyl;

25

30

R¹⁰ is independently selected from the group consisting of: hydrido, lower alkyl, heteroaryl, heterocyclic, haloalkyl, arylalkylamino, heteroarylalkyl, aryl, and arylalkyl, wherein aryl, heteroaryl, heterocyclic, or arylalkyl are optionally substituted with one or more radical selected from alkyl, alkoxy, halogen, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, and heterocyclic;

5 **R^{10'}** is independently selected from the group consisting of: hydrido, lower alkyl, heteroaryl, heterocyclic, haloalkyl, arylalkylamino, heteroarylalkyl, aryl, and arylalkyl, wherein aryl, heteroaryl, heterocyclic, or arylalkyl are optionally substituted with one or more radical selected from alkyl, alkoxy, halogen, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, and heterocyclic;

10 **R¹¹** is selected from the group consisting of: hydrido, halogen, haloalkyl, CN, CO₂R⁵, lower alkyl, lower alkenyl, lower alkynyl, alkoxy, and CONH₂;

15 **R¹²** is selected from the group consisting of: hydrido, halogen, alkyl, and alkoxy;

20 **R¹³** is selected from the group consisting of: hydrido, alkyl, aryl, arylalkyl, heteroaryl, heterocyclicalkyl, and heteroarylalkyl, wherein aryl, alkyl, arylalkyl, heteroaryl, heterocyclicalkyl, or heteroarylalkyl are optionally substituted with one or more radicals selected from the group consisting of OR¹⁴, N(R¹⁴)R^{14'}, and glycols;

25 **R¹⁴** is independently selected from the group consisting of hydrido, and lower alkyl; and

R^{14'} is independently selected from the group consisting of hydrido, and lower alkyl;

30 **R¹⁵** is selected from the group consisting of: hydrido, halogen, alkyl, cycloalkyl, aryl, haloalkyl, heteroaryl, heterocyclic, alkylalkene, alkylalkyne, hydroxy, hydroxyalkyl, alkylhydroxy, amino, aminoalkyl,


alkylamino, alkylaminoalkyl, alkylhydroxyalkyl, nitro, cyano, alkylthio, alkylsulfinyl, alkylsulfonyl; wherein aryl or arylalkyl are optionally substituted with one or more radical selected from alkyl, alkoxy, halo, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, heterocyclic, and

5 R^{16} is selected from the group consisting of: hydrido, halogen, alkyl, cycloalkyl, aryl, haloalkyl, heteroaryl, heterocyclic, alkylalkene, alkylalkyne, hydroxy, hydroxyalkyl, alkylhydroxy, amino, aminoalkyl, alkylamino, alkylaminoalkyl, alkylhydroxyalkyl, nitro, cyano, alkylthio, alkylsulfinyl, alkylsulfonyl; wherein aryl or arylalkyl are optionally substituted with one or more radical selected from alkyl, alkoxy, halo, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, heterocyclic, wherein

10 R^{15} and R^{16} may be taken together to form a 5 to 7 membered saturated or unsaturated carbocyclic ring optionally containing 0 to 3 heteroatoms selected from the group consisting of N, O, or S, and wherein said ring is optionally substituted with R^1 ;

15 or isomers, tautomers, carriers, esters, prodrugs, pharmaceutically acceptable salts thereof.

20 2. A compound of formula II



wherein

A is $(\text{CH}_2)_m\text{-CR}^{15}=\text{CR}^{16}\text{-}(\text{CH}_2)_m$

m is independently selected from 0, 1, or 2;

B is a 5 or 6 membered heteroaryl, aryl, saturated or unsaturated

5 heterocyclic wherein said aryl, heteroaryl, or heterocyclic are optionally substituted with R^1 , R^2 , and R^{12} ;

10 R^1 is selected from the group consisting of: hydrido, halogen, alkyl, aryl, heteroaryl, alkenyl, alkynyl, haloalkyl, CN, NO_2 , OR^5 , OCOOR^5 , CO_2R^7 , $\text{CON}(\text{R}^6)\text{R}^7$, COR^6 , SR^6 , SOR^6 , SO_2R^6 , NR^6R^7 , NR^6COR^7 , $\text{NR}^6\text{CONHR}^7$, $\text{NR}^6\text{SO}_2\text{R}^7$, $\text{NR}^6\text{SO}_2\text{NHR}^7$, and $\text{SO}_2\text{N}(\text{R}^6)\text{R}^7$ wherein R^6 and R^7 may be taken together to form a 3-7 membered carbocyclic ring having 1 to 3 substituted or unsubstituted heteroatoms selected from the group consisting of: S, SO, SO_2 , O, and NR^6 ; wherein said alkenyl, alkynyl, alkyl, aryl, heteroaryl or OR^5 are optional substituted with, hydrido, halogen, alkyl, hydroxyalkyl, aryl, heteroaryl, haloalkyl, COCF_3 , CN, NO_2 , OR^5 , OCOOR^5 , CO_2R^7 , $\text{CON}(\text{R}^6)\text{R}^7$, COR^6 , SR^6 , SOR^6 , SO_2R^6 , NR^6R^7 , NR^6COR^7 , $\text{NR}^6\text{CONHR}^7$, $\text{NR}^6\text{SO}_2\text{R}^7$, $\text{NR}^6\text{SO}_2\text{NHR}^7$, and $\text{SO}_2\text{N}(\text{R}^6)\text{R}^7$ wherein R^6 and R^7 may be taken together to form a 3-7 membered carbocyclic ring having 1 to 3 substituted or unsubstituted heteroatoms selected from the group consisting of: S, SO, SO_2 , O, and NR^6 ;

15 R^2 is selected from the group consisting of: halogen, hydrido, hydroxyalkyl, alkyl, OR^6 , CN, NO_2 , SR^6 , NHR^6 , $\text{CON}(\text{R}^6)\text{R}^7$, NHCONHR^6 , CO_2H , and haloalkyl;

20 R^1 and R^2 may be taken together to form a 5 to 7 membered saturated or unsaturated carbocyclic ring optionally containing 0 to 3 heteroatoms selected from the group consisting of N, O, or S, and wherein said ring is optionally substituted with R^1 ;

25 R^3 is selected from the group consisting of: substituted or unsubstituted amidine, alkylamino, aminoalkyl, CONHR^7 , NH_2 , NHCOR^6 , and $\text{CH}_2\text{NHCOR}^6$;

R⁴ is selected from the group consisting of: halogen, alkylsulfinyl, alkylsulfonyl, cyano, alkoxycarbonyl, alkyl, haloalkyl, hydrido, hydroxyalkyl, haloalkoxy, heterocyclic, nitro, acylamino, aryl, heteroaryl, and alkenyl, OR¹³, SR⁸, SO₂N(R⁸)R^{8'}, NHR⁹, NHCOR⁹, NR⁹COR⁹, NHCO(OR⁹), NR⁹CO(OR⁹), NR⁸SO₂R¹⁰, NHSO₂N(R¹⁰)R^{10'}, NR⁶CON(R¹⁰)R^{10'}, COR⁹, CO₂R⁸, CON(R⁸)R^{8'}, wherein R⁸ and R^{8'} may be taken together to form a 3-7 membered carbocyclic ring having 1 to 3 substituted or unsubstituted heteroatoms selected from S, SO, SO₂, O, N, and NR⁶, and wherein R¹⁰ and R^{10'} may be taken together to form a 3-7 membered carbocyclic ring having 1 to 3 substituted or unsubstituted heteroatoms selected from S, SO, SO₂, O, N, and NR⁶ wherein said aryl, heterocyclic, heteroaryl, or alkenyl are optionally substituted with R⁹;

R⁵ is selected from the group consisting of: hydrido, alkyl, aryl, arylalkyl, heteroaryl, heterocyclicalkyl, and heteroarylalkyl, wherein aryl, alkyl, arylalkyl, heteroaryl, heterocyclicalkyl, or heteroarylalkyl are optionally substituted with one or more radicals selected from the group consisting of OR¹⁴, N(R¹⁴)R^{14'}, and glycols;

R⁶ is independently selected from the group consisting of: hydrido, aryl, heteroaryl, lower alkyl, haloalkyl, alkenyl, alkynyl, hydroxyalkyl, aminoalkyl, alkylaminoalkyl, alkoxy, alkoxyalkyl, heterocyclicalkyl, and heterocyclic;

R⁷ is independently selected from the group consisting of: hydrido, aryl, heteroaryl, lower alkyl, haloalkyl, alkenyl, alkynyl, hydroxyalkyl, aminoalkyl, alkylaminoalkyl, alkoxy, alkoxyalkyl, heterocyclicalkyl, and heterocyclic;

R⁸ is independently selected from the group consisting of: hydrido, aryl, heteroaryl, arylalkyl, heterocyclic, haloalkyl, arylalkylamino, alkylaminoalkyl, dialkylaminoalkyl, alkyl, alkenyl, alkynyl, heteroarylalkyl, and heterocyclicalkyl;

R^{8'} is independently selected from the group consisting of: hydrido, aryl, heteroaryl, arylalkyl, heterocyclic, haloalkyl, arylalkylamino,

alkylaminoalkyl, dialkylaminoalkyl, alkyl, alkenyl, alkynyl, heteroarylalkyl, and heterocyclicalkyl;

5 R^9 is independently selected from the group consisting of: hydrido, lower alkyl, aryl, heteroaryl, arylalkyl, heterocyclic, cycloalkyl, heterocyclicalkyl, haloalkyl, arylalkylamino, amino, aminoalkyl, aminoacyl, nitro, azido, and heteroarylalkyl, wherein alkyl, aryl, heteroaryl, aminoalkyl, or arylalkyl are optionally substituted with one or more radical selected from the group consisting of: alkylsulfonamide, sulfamyl, alkyl, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, aminoalkyl, alkylaminoalkyl, alkoxy, halogen, acyloxy, oxy, formyl, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, nitro, azido, benzyloxy, dialkylaminoacyl, thioalkyl, aminoacyloxy, thiocyanate, isothiocyanate, alkyldioxy, hydroxyalkyl, alkylamino, alkyloxycarbonyl, alkoxyalkyl, alkenylamino, alkynylamino, alkenyl, alkynyl, dialkylaminoalkyloxy, and heterocyclic optionally substituted with alkyl, alkylamino, aminoalkyl, hydroxyalkyl, and alkylaminoalkyl;

10 R^{10} is independently selected from the group consisting of: hydrido, lower alkyl, heteroaryl, heterocyclic, haloalkyl, arylalkylamino, heteroarylalkyl, aryl, and arylalkyl, wherein aryl, heteroaryl, heterocyclic, or arylalkyl are optionally substituted with one or more radical selected from alkyl, alkoxy, halogen, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, and heterocyclic;

15 $R^{10'}$ is independently selected from the group consisting of: hydrido, lower alkyl, heteroaryl, heterocyclic, haloalkyl, arylalkylamino, heteroarylalkyl, aryl, and arylalkyl, wherein aryl, heteroaryl, heterocyclic, or arylalkyl are optionally substituted with one or more radical selected from alkyl, alkoxy, halogen, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, and heterocyclic;

20 $R^{10''}$ is independently selected from the group consisting of: hydrido, lower alkyl, heteroaryl, heterocyclic, haloalkyl, arylalkylamino, heteroarylalkyl, aryl, and arylalkyl, wherein aryl, heteroaryl, heterocyclic, or arylalkyl are optionally substituted with one or more radical selected from alkyl, alkoxy, halogen, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, and heterocyclic;

25 $R^{10'}$ is independently selected from the group consisting of: hydrido, lower alkyl, heteroaryl, heterocyclic, haloalkyl, arylalkylamino, heteroarylalkyl, aryl, and arylalkyl, wherein aryl, heteroaryl, heterocyclic, or arylalkyl are optionally substituted with one or more radical selected from alkyl, alkoxy, halogen, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, and heterocyclic;

30 $R^{10''}$ is independently selected from the group consisting of: hydrido, lower alkyl, heteroaryl, heterocyclic, haloalkyl, arylalkylamino, heteroarylalkyl, aryl, and arylalkyl, wherein aryl, heteroaryl, heterocyclic, or arylalkyl are optionally substituted with one or more radical selected from alkyl, alkoxy, halogen, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, and heterocyclic;

R¹¹ is selected from the group consisting of: hydrido, halogen, haloalkyl, CN, CO₂R⁵, lower alkyl, lower alkenyl, lower alkynyl, alkoxy, and CONH₂;

5 **R¹²** is selected from the group consisting of: hydrido, halogen, alkyl, and alkoxy;

10 **R¹³** is selected from the group consisting of: hydrido, alkyl, aryl, arylalkyl, heteroaryl, heterocyclicalkyl, and heteroarylalkyl, wherein aryl, alkyl, arylalkyl, heteroaryl, heterocyclicalkyl, or heteroarylalkyl are optionally substituted with one or more radicals selected from the group consisting of OR¹⁴, N(R¹⁴)R^{14'}, and glycols;

15 **R¹⁴** is independently selected from the group consisting of hydrido, and lower alkyl; and

20 **R^{14'}** is independently selected from the group consisting of hydrido, and lower alkyl;

25 **R¹⁵** is selected from the group consisting of: hydrido, halogen, alkyl, cycloalkyl, aryl, haloalkyl, heteroaryl, heterocyclic, alkylalkene, alkylalkyne, hydroxy, hydroxyalkyl, alkylhydroxy, amino, aminoalkyl, alkylamino, alkylaminoalkyl, alkylhydroxyalkyl, nitro, cyano, alkylthio, alkylsulfinyl, alkylsulfonyl; wherein aryl or arylalkyl are optionally substituted with one or more radical selected from alkyl, alkoxy, halo, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, heterocyclic; and

30 **R¹⁶** is selected from the group consisting of: hydrido, halogen, alkyl, cycloalkyl, aryl, haloalkyl, heteroaryl, heterocyclic, alkylalkene, alkylalkyne, hydroxy, hydroxyalkyl, alkylhydroxy, amino, aminoalkyl, alkylamino, alkylaminoalkyl, alkylhydroxyalkyl, nitro, cyano, alkylthio, alkylsulfinyl, alkylsulfonyl; wherein aryl or arylalkyl are optionally substituted with one or more radical selected from alkyl, alkoxy, halo, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, heterocyclic, wherein R¹⁵ and R¹⁶ may be taken together to form a 5 to 7 membered saturated or


unsaturated carbocyclic ring optionally containing 0 to 3 heteroatoms selected from the group consisting of N, O, or S, and wherein said ring is optionally substituted with R¹;

5

or isomers, tautomers, carriers, esters, prodrugs, pharmaceutically acceptable salts thereof.

3. The compound of claim 2 of formula II

10



wherein

A is (CH₂)_m-CR¹⁵=CR¹⁶-(CH₂)_m

15 m is independently selected from 0, 1, or 2;

B is a 5 or 6 membered heteroaryl, aryl, saturated or unsaturated heterocyclic wherein said aryl, heteroaryl, or heterocyclic are optionally substituted with R¹, R², and R¹²;

20 R¹ is selected from the group consisting of: hydrido, halogen, alkyl, aryl, heteroaryl, alkenyl, alkynyl, haloalkyl, CN, NO₂, OR⁵, OCOOR⁵, CO₂R⁷, CON(R⁶)R⁷, COR⁶, SR⁶, SOR⁶, SO₂R⁶, NR⁶R⁷, NR⁶COR⁷, NR⁶CONHR⁷, NR⁶SO₂R⁷, NR⁶SO₂NHR⁷, and SO₂N(R⁶)R⁷ wherein R⁶ and R⁷ may be taken together to form a 3-7 membered carbocyclic ring having 1 to 3 substituted or unsubstituted heteroatoms selected from the group consisting of: S, SO, SO₂, O, and NR⁶; wherein said alkenyl,

25

alkynyl, alkyl, aryl, heteroaryl or OR^5 are optional substituted with, hydrido, halogen, alkyl, hydroxyalkyl, aryl, heteroaryl, haloalkyl, $COCF_3$, CN , NO_2 , OR^5 , $OCOOR^5$, CO_2R^7 , $CON(R^6)R^7$, COR^6 , SR^6 , SOR^6 , SO_2R^6 , NR^6R^7 , NR^6COR^7 , NR^6CONHR^7 , $NR^6SO_2R^7$, $NR^6SO_2NHR^7$, and $SO_2N(R^6)R^7$ wherein R^6 and R^7 may be taken together to form a 3-7 membered carbocyclic ring having 1 to 3 substituted or unsubstituted heteroatoms selected from the group consisting of: S, SO, SO_2 , O, and NR^6 ;

5 R^2 is hydrido;

10 R^3 is selected from the group consisting of: substituted or unsubstituted amidine, alkylamino, aminoalkyl, $CONHR^7$, NH_2 , $NHCOR^6$, and CH_2NHCOR^6 ;

15 R^4 is selected from the group consisting of: halogen, alkylsulfinyl, alkylsulfonyl, cyano, alkoxy carbonyl, alkyl, haloalkyl, hydrido, hydroxyalkyl, haloalkoxy, heterocyclic, nitro, acylamino, aryl, heteroaryl, and alkenyl, OR^{13} , SR^8 , $SO_2N(R^8)R^{8'}$, NHR^9 , $NHCOR^9$, NR^9COR^9 , $NHCO(OR^9)$, $NR^9CO(OR^9)$, $NR^8SO_2R^{10}$, $NHSO_2N(R^{10})R^{10'}$, $NR^6CON(R^{10})R^{10'}$, COR^9 , CO_2R^8 , $CON(R^8)R^{8'}$, wherein R^8 and $R^{8'}$ may be taken together to form a 3-7 membered carbocyclic ring having 1 to 3 substituted or unsubstituted heteroatoms selected from S, SO, SO_2 , O, N, and NR^6 , and wherein R^{10} and $R^{10'}$ may be taken together to form a 3-7 membered carbocyclic ring having 1 to 3 substituted or unsubstituted heteroatoms selected from S, SO, SO_2 , O, N, and NR^6 wherein said aryl, heterocyclic, heteroaryl, or alkenyl are optionally substituted with R^9 ;

20 R^5 is selected from the group consisting of: hydrido, alkyl, aryl, arylalkyl, heteroaryl, heterocyclic alkyl, and heteroarylalkyl, wherein aryl, alkyl, arylalkyl, heteroaryl, heterocyclic alkyl, or heteroarylalkyl are optionally substituted with one or more radicals selected from the group consisting of OR^{14} , $N(R^{14})R^{14'}$, and glycols;

25 R^6 is independently selected from the group consisting of: hydrido, aryl, heteroaryl, lower alkyl, haloalkyl, alkenyl, alkynyl, hydroxyalkyl,

30

aminoalkyl, alkylaminoalkyl, alkoxy, alkoxyalkyl, heterocyclicalkyl, and heterocyclic;

5 **R**⁷ is independently selected from the group consisting of: hydrido, aryl, heteroaryl, lower alkyl, haloalkyl, alkenyl, alkynyl, hydroxyalkyl, aminoalkyl, alkylaminoalkyl, alkoxy, alkoxyalkyl, heterocyclicalkyl, and heterocyclic;

10 **R**⁸ is independently selected from the group consisting of: hydrido, aryl, heteroaryl, arylalkyl, heterocyclic, haloalkyl, arylalkylamino, alkylaminoalkyl, dialkylaminoalkyl, alkyl, alkenyl, alkynyl, heteroarylalkyl, and heterocyclicalkyl;

15 **R**^{8'} is independently selected from the group consisting of: hydrido, aryl, heteroaryl, arylalkyl, heterocyclic, haloalkyl, arylalkylamino, alkylaminoalkyl, dialkylaminoalkyl, alkyl, alkenyl, alkynyl, heteroarylalkyl, and heterocyclicalkyl;

20 **R**⁹ is independently selected from the group consisting of: hydrido, lower alkyl, aryl, heteroaryl, arylalkyl, heterocyclic, cycloalkyl, heterocyclicalkyl, haloalkyl, arylalkylamino, amino, aminoalkyl, aminoacyl, nitro, azido, and heteroarylalkyl, wherein alkyl, aryl, heteroaryl, aminoalkyl, or arylalkyl are optionally substituted with one or more radical selected from the group consisting of: alkylsulfonamide, sulfamyl, alkyl, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, aminoalkyl, alkylaminoalkyl, alkoxy, halogen, acyloxy, oxy, formyl, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, nitro, azido, benzyloxy, dialkylaminoacyl, thioalkyl, aminoacyloxy, thiocyanate, isothiocyanate, alkyldioxy, hydroxyalkyl, alkylamino, alkyloxycarbonyl, alkoxyalkyl, alkenylamino, alkynylamino, alkenyl, alkynyl, dialkylaminoalkyloxy, and heterocyclic optionally substituted with alkyl, alkylamino, aminoalkyl, hydroxyalkyl, and alkylaminoalkyl;

25 **R**¹⁰ is independently selected from the group consisting of: hydrido, lower alkyl, heteroaryl, heterocyclic, haloalkyl, arylalkylamino,

heteroarylalkyl, aryl, and arylalkyl, wherein aryl, heteroaryl, heterocyclic, or arylalkyl are optionally substituted with one or more radical selected from alkyl, alkoxy, halogen, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, and heterocyclic;

5 **R¹⁰** is independently selected from the group consisting of: hydrido, lower alkyl, heteroaryl, heterocyclic, haloalkyl, arylalkylamino, heteroarylalkyl, aryl, and arylalkyl, wherein aryl, heteroaryl, heterocyclic, or arylalkyl are optionally substituted with one or more radical selected from alkyl, alkoxy, halogen, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, and heterocyclic;

10 **R¹¹** is selected from the group consisting of: hydrido, halogen, haloalkyl, CN, CO₂R⁵, lower alkyl, lower alkenyl, lower alkynyl, alkoxy, and CONH₂;

15 **R¹²** is selected from the group consisting of: hydrido, halogen, alkyl, and alkoxy;

R¹³ is selected from the group consisting of: hydrido, alkyl, aryl, arylalkyl, heteroaryl, heterocyclicalkyl, and heteroarylalkyl, wherein aryl, alkyl, arylalkyl, heteroaryl, heterocyclicalkyl, or heteroarylalkyl are optionally substituted with one or more radicals selected from the group consisting of OR¹⁴, N(R¹⁴)R^{14'}, and glycols;

20 **R¹⁴** is independently selected from the group consisting of hydrido, and lower alkyl; and

25 **R^{14'}** is independently selected from the group consisting of hydrido, and lower alkyl;

R¹⁵ is selected from the group consisting of: hydrido, halogen, alkyl, cycloalkyl, aryl, haloalkyl, heteroaryl, heterocyclic, alkylalkene, alkylalkyne, hydroxy, hydroxyalkyl, alkylhydroxy, amino, aminoalkyl, alkylamino, alkylaminoalkyl, alkylhydroxyalkyl, nitro, cyano, alkylthio, alkylsulfinyl, alkylsulfonyl; wherein aryl or arylalkyl are optionally

substituted with one or more radical selected from alkyl, alkoxy, halo, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, heterocyclic; and


5 R^{16} is selected from the group consisting of: hydrido, halogen, alkyl, cycloalkyl, aryl, haloalkyl, heteroaryl, heterocyclic, alkylalkene, alkylalkyne, hydroxy, hydroxyalkyl, alkylhydroxy, amino, aminoalkyl, alkylamino, alkylaminoalkyl, alkylhydroxyalkyl, nitro, cyano, alkylthio, alkylsulfinyl, alkylsulfonyl; wherein aryl or arylalkyl are optionally substituted with one or more radical selected from alkyl, alkoxy, halo, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, heterocyclic; wherein

10 R^{15} and R^{16} may be taken together to form a 5 to 7 membered saturated or unsaturated carbocyclic ring optionally containing 0 to 3 heteroatoms selected from the group consisting of N, O, or S, and wherein said ring is

15 optionally substituted with R^1 ;

or isomers, tautomers, carriers, esters, prodrugs, pharmaceutically acceptable salts thereof.

20 4. The compound of claim 2 of formula II



wherein

A is $(CH_2)_m-CR^{15}=CR^{16}-(CH_2)_n$

25 m is independently selected from 0, 1, or 2;

B is a 5 or 6 membered heteroaryl, aryl, saturated or unsaturated heterocyclic wherein said aryl, heteroaryl, or heterocyclic are optionally substituted with R¹, R², and R¹²;

R¹ is selected from the group consisting of: hydrido, halogen, alkyl, aryl, heteroaryl, alkenyl, alkynyl, haloalkyl, CN, NO₂, OR⁵, OCOOR⁵, CO₂R⁷, CON(R⁶)R⁷, COR⁶, SR⁶, SOR⁶, SO₂R⁶, NR⁶R⁷, NR⁶COR⁷, NR⁶CONHR⁷, NR⁶SO₂R⁷, NR⁶SO₂NHR⁷, and SO₂N(R⁶)R⁷ wherein R⁶ and R⁷ may be taken together to form a 3-7 membered carbocyclic ring having 1 to 3 substituted or unsubstituted heteroatoms selected from the group consisting of: S, SO, SO₂, O, and NR⁶; wherein said alkenyl, alkynyl, alkyl, aryl, heteroaryl or OR⁵ are optional substituted with, hydrido, halogen, alkyl, hydroxyalkyl, aryl, heteroaryl, haloalkyl, COCF₃, CN, NO₂, OR⁵, OCOOR⁵, CO₂R⁷, CON(R⁶)R⁷, COR⁶, SR⁶, SOR⁶, SO₂R⁶, NR⁶R⁷, NR⁶COR⁷, NR⁶CONHR⁷, NR⁶SO₂R⁷, NR⁶SO₂NHR⁷, and SO₂N(R⁶)R⁷ wherein R⁶ and R⁷ may be taken together to form a 3-7 membered carbocyclic ring having 1 to 3 substituted or unsubstituted heteroatoms selected from the group consisting of: S, SO, SO₂, O, and NR⁶;

R² is selected from the group consisting of: halogen, hydrido, hydroxyalkyl, alkyl, OR⁶, CN, NO₂, SR⁶, NHR⁶, CON(R⁶)R⁷, NHCONHR⁶, CO₂H, and haloalkyl;

R¹ and R² may be taken together to form a 5 to 7 membered saturated or unsaturated carbocyclic ring optionally containing 0 to 3 heteroatoms selected from the group consisting of N, O, or S, and wherein said ring is optionally substituted with R¹;

R³ is selected from the group consisting of: substituted or unsubstituted amidine, alkylamino, aminoalkyl, CONHR⁷, NH₂, NHCOR⁶, and CH₂NHCOR⁶;

R⁴ is selected from the group consisting of: halogen, alkylsulfinyl, alkylsulfonyl, cyano, alkoxy carbonyl, alkyl, haloalkyl, hydrido, hydroxyalkyl, haloalkoxy, heterocyclic, nitro, acylamino, aryl,

heteroaryl, and alkenyl, OR^{13} , SR^8 , $SO_2N(R^8)R^{8'}$, NHR^9 , $NHCOR^9$, NR^9COR^9 , $NHCO(OR^9)$, $NR^9CO(OR^9)$, $NR^8SO_2R^{10}$, $NHSO_2N(R^{10})R^{10'}$, $NR^6CON(R^{10})R^{10'}$, COR^9 , CO_2R^8 , $CON(R^8)R^{8'}$, wherein R^8 and $R^{8'}$ may be taken together to form a 3-7 membered carbocyclic ring having 1 to 3 substituted or unsubstituted heteroatoms selected from S, SO, SO_2 , O, N, and NR^6 , and wherein R^{10} and $R^{10'}$ may be taken together to form a 3-7 membered carbocyclic ring having 1 to 3 substituted or unsubstituted heteroatoms selected from S, SO, SO_2 , O, N, and NR^6 wherein said aryl, heterocyclic, heteroaryl, or alkenyl are optionally substituted with R^9 ;

5 **R⁵** is selected from the group consisting of: hydrido, alkyl, aryl, arylalkyl, heteroaryl, heterocyclicalkyl, and heteroarylalkyl, wherein aryl, alkyl, arylalkyl, heteroaryl, heterocyclicalkyl, or heteroarylalkyl are optionally substituted with one or more radicals selected from the group consisting of OR^{14} , $N(R^{14})R^{14'}$, and glycols;

10 **R⁶** is independently selected from the group consisting of: hydrido, aryl, heteroaryl, lower alkyl, haloalkyl, alkenyl, alkynyl, hydroxyalkyl, aminoalkyl, alkylaminoalkyl, alkoxy, alkoxyalkyl, heterocyclicalkyl, and heterocyclic;

15 **R⁷** is independently selected from the group consisting of: hydrido, aryl, heteroaryl, lower alkyl, haloalkyl, alkenyl, alkynyl, hydroxyalkyl, aminoalkyl, alkylaminoalkyl, alkoxy, alkoxyalkyl, heterocyclicalkyl, and heterocyclic;

20 **R⁸** is independently selected from the group consisting of: hydrido, aryl, heteroaryl, arylalkyl, heterocyclic, haloalkyl, arylalkylamino, alkylaminoalkyl, dialkylaminoalkyl, alkyl, alkenyl, alkynyl, heteroarylalkyl, and heterocyclicalkyl;

25 **R^{8'}** is independently selected from the group consisting of: hydrido, aryl, heteroaryl, arylalkyl, heterocyclic, haloalkyl, arylalkylamino, alkylaminoalkyl, dialkylaminoalkyl, alkyl, alkenyl, alkynyl, heteroarylalkyl, and heterocyclicalkyl;

30 **R⁹** is independently selected from the group consisting of: hydrido, aryl, heteroaryl, arylalkyl, heterocyclic, haloalkyl, arylalkylamino, alkylaminoalkyl, dialkylaminoalkyl, alkyl, alkenyl, alkynyl, heteroarylalkyl, and heterocyclicalkyl;

R⁹ is independently selected from the group consisting of: hydrido, lower alkyl, aryl, heteroaryl, arylalkyl, heterocyclic, cycloalkyl, heterocyclicalkyl, haloalkyl, arylalkylamino, amino, aminoalkyl, aminoacyl, nitro, azido, and heteroarylalkyl, wherein alkyl, aryl, heteroaryl, aminoalkyl, or arylalkyl are optionally substituted with one or more radical selected from the group consisting of: alkylsulfonamide, sulfamyl, alkyl, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, aminoalkyl, alkylaminoalkyl, alkoxy, halogen, acyloxy, oxy, formyl, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, nitro, azido, benzyloxy, dialkylaminoacyl, thioalkyl, aminoacyloxy, thiocyanate, isothiocyanate, alkyldioxy, hydroxyalkyl, alkylamino, alkyloxycarbonyl, alkoxyalkyl, alkenylamino, alkynylamino, alkenyl, alkynyl, dialkylaminoalkyloxy, and heterocyclic optionally substituted with alkyl, alkylamino, aminoalkyl, hydroxyalkyl, and alkylaminoalkyl;

R¹⁰ is independently selected from the group consisting of: hydrido, lower alkyl, heteroaryl, heterocyclic, haloalkyl, arylalkylamino, heteroarylalkyl, aryl, and arylalkyl, wherein aryl, heteroaryl, heterocyclic, or arylalkyl are optionally substituted with one or more radical selected from alkyl, alkoxy, halogen, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, and heterocyclic;

R^{10'} is independently selected from the group consisting of: hydrido, lower alkyl, heteroaryl, heterocyclic, haloalkyl, arylalkylamino, heteroarylalkyl, aryl, and arylalkyl, wherein aryl, heteroaryl, heterocyclic, or arylalkyl are optionally substituted with one or more radical selected from alkyl, alkoxy, halogen, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, and heterocyclic;

R¹¹ is hydrido;

\mathbf{R}^{12} is selected from the group consisting of: hydrido, halogen, alkyl, and alkoxy;

\mathbf{R}^{13} is selected from the group consisting of: hydrido, alkyl, aryl, arylalkyl, heteroaryl, heterocyclicalkyl, and heteroarylalkyl, wherein aryl, alkyl, arylalkyl, heteroaryl, heterocyclicalkyl, or heteroarylalkyl are optionally substituted with one or more radicals selected from the group consisting of OR^{14} , $\text{N}(\text{R}^{14})\text{R}^{14'}$, and glycols;

\mathbf{R}^{14} is independently selected from the group consisting of hydrido, and lower alkyl; and

10 $\mathbf{R}^{14'}$ is independently selected from the group consisting of hydrido, and lower alkyl;

\mathbf{R}^{15} is selected from the group consisting of: hydrido, halogen, alkyl, cycloalkyl, aryl, haloalkyl, heteroaryl, heterocyclic, alkylalkene, alkylalkyne, hydroxy, hydroxyalkyl, alkylhydroxy, amino, aminoalkyl, alkylamino, alkylaminoalkyl, alkylhydroxyalkyl, nitro, cyano, alkylthio, alkylsulfinyl, alkylsulfonyl; wherein aryl or arylalkyl are optionally substituted with one or more radical selected from alkyl, alkoxy, halo, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, heterocyclic; and

15 \mathbf{R}^{16} is selected from the group consisting of: hydrido, halogen, alkyl, cycloalkyl, aryl, haloalkyl, heteroaryl, heterocyclic, alkylalkene, alkylalkyne, hydroxy, hydroxyalkyl, alkylhydroxy, amino, aminoalkyl, alkylamino, alkylaminoalkyl, alkylhydroxyalkyl, nitro, cyano, alkylthio, alkylsulfinyl, alkylsulfonyl; wherein aryl or arylalkyl are optionally substituted with one or more radical selected from alkyl, alkoxy, halo, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, heterocyclic; wherein \mathbf{R}^{15} and \mathbf{R}^{16} may be taken together to form a 5 to 7 membered saturated or unsaturated carbocyclic ring optionally containing 0 to 3 heteroatoms


20 selected from the group consisting of N, O, or S, and wherein said ring is optionally substituted with \mathbf{R}^1 ;

25

30

or isomers, tautomers, carriers, esters, prodrugs, pharmaceutically acceptable salts thereof.

5 5. The compound of claim 2 of formula II



wherein

10 **A** is $(CH_2)_m-CR^{15}=CR^{16}-(CH_2)_m$
m is independently selected from 0, 1, or 2;

B is a 5 or 6 membered heteroaryl, aryl, saturated or unsaturated heterocyclic wherein said aryl, heteroaryl, or heterocyclic are optionally substituted with **R¹**, **R²**, and **R¹²**;

15 **R¹** is selected from the group consisting of: hydrido, halogen, alkyl, aryl, heteroaryl, alkenyl, alkynyl, haloalkyl, CN, NO₂, OR⁵, OCOOR⁵, CO₂R⁷, CON(R⁶)R⁷, COR⁶, SR⁶, SOR⁶, SO₂R⁶, NR⁶R⁷, NR⁶COR⁷, NR⁶CONHR⁷, NR⁶SO₂R⁷, NR⁶SO₂NHR⁷, and SO₂N(R⁶)R⁷ wherein R⁶ and R⁷ may be taken together to form a 3-7 membered carbocyclic ring having 1 to 3 substituted or unsubstituted heteroatoms selected from the group consisting of: S, SO, SO₂, O, and NR⁶; wherein said alkenyl, alkynyl, alkyl, aryl, heteroaryl or OR⁵ are optional substituted with, hydrido, halogen, alkyl, hydroxyalkyl, aryl, heteroaryl, haloalkyl, COCF₃, CN, NO₂, OR⁵, OCOOR⁵, CO₂R⁷, CON(R⁶)R⁷, COR⁶, SR⁶, SOR⁶, SO₂R⁶, NR⁶R⁷, NR⁶COR⁷, NR⁶CONHR⁷, NR⁶SO₂R⁷,

20 25

NR⁶SO₂NHR⁷, and SO₂N(R⁶)R⁷ wherein R⁶ and R⁷ may be taken together to form a 3-7 membered carbocyclic ring having 1 to 3 substituted or unsubstituted heteroatoms selected from the group consisting of: S, SO, SO₂, O, and NR⁶;

5 R² is selected from the group consisting of: halogen, hydrido, hydroxyalkyl, alkyl, OR⁶, CN, NO₂, SR⁶, NHR⁶, CON(R⁶)R⁷, NHCONHR⁶, CO₂H, and haloalkyl;

R¹ and R² may be taken together to form a 5 to 7 membered saturated or unsaturated carbocyclic ring optionally containing 0 to 3 heteroatoms selected from the group consisting of N, O, or S, and wherein said ring is optionally substituted with R¹;

10 R³ is selected from the group consisting of: substituted or unsubstituted amidine, alkylamino, aminoalkyl, CONHR⁷, NH₂, NHCOR⁶, and CH₂NHCOR⁶;

15 R⁴ is selected from the group consisting of: halogen, alkylsulfinyl, alkylsulfonyl, cyano, alkoxy carbonyl, alkyl, haloalkyl, hydrido, hydroxyalkyl, haloalkoxy, heterocyclic, nitro, acylamino, aryl, heteroaryl, and alkenyl, OR¹³, SR⁸, SO₂N(R⁸)R^{8'}, NHR⁹, NHCOR⁹, NR⁹COR⁹, NHCO(OR⁹), NR⁹CO(OR⁹), NR⁸SO₂R¹⁰, NHSO₂N(R¹⁰)R^{10'}, NR⁶CON(R¹⁰)R^{10'}, COR⁹, CO₂R⁸, CON(R⁸)R^{8'}, wherein R⁸ and R^{8'} may be taken together to form a 3-7 membered carbocyclic ring having 1 to 3 substituted or unsubstituted heteroatoms selected from S, SO, SO₂, O, N, and NR⁶, and wherein R¹⁰ and R^{10'} may be taken together to form a 3-7 membered carbocyclic ring having 1 to 3 substituted or unsubstituted heteroatoms selected from S, SO, SO₂, O, N, and NR⁶ wherein said aryl, heterocyclic, heteroaryl, or alkenyl are optionally substituted with R⁹;

20 R⁵ is selected from the group consisting of: hydrido, alkyl, aryl, arylalkyl, heteroaryl, heterocyclicalkyl, and heteroarylalkyl, wherein aryl, alkyl, arylalkyl, heteroaryl, heterocyclicalkyl, or heteroarylalkyl are optionally substituted with one or more radicals selected from the group consisting of OR¹⁴, N(R¹⁴)R^{14'}, and glycols;

25

30

R⁶ is independently selected from the group consisting of: hydrido, aryl, heteroaryl, lower alkyl, haloalkyl, alkenyl, alkynyl, hydroxyalkyl, aminoalkyl, alkylaminoalkyl, alkoxy, alkoxyalkyl, heterocyclicalkyl, and heterocyclic;

5 **R⁷** is independently selected from the group consisting of: hydrido, aryl, heteroaryl, lower alkyl, haloalkyl, alkenyl, alkynyl, hydroxyalkyl, aminoalkyl, alkylaminoalkyl, alkoxy, alkoxyalkyl, heterocyclicalkyl, and heterocyclic;

10 **R⁸** is independently selected from the group consisting of: hydrido, aryl, heteroaryl, arylalkyl, heterocyclic, haloalkyl, arylalkylamino, alkylaminoalkyl, dialkylaminoalkyl, alkyl, alkenyl, alkynyl, heteroarylalkyl, and heterocyclicalkyl;

15 **R^{8'}** is independently selected from the group consisting of: hydrido, aryl, heteroaryl, arylalkyl, heterocyclic, haloalkyl, arylalkylamino, alkylaminoalkyl, dialkylaminoalkyl, alkyl, alkenyl, alkynyl, heteroarylalkyl, and heterocyclicalkyl;

20 **R⁹** is independently selected from the group consisting of: hydrido, lower alkyl, aryl, heteroaryl, arylalkyl, heterocyclic, cycloalkyl, heterocyclicalkyl, haloalkyl, arylalkylamino, amino, aminoalkyl, aminoacyl, nitro, azido, and heteroarylalkyl, wherein alkyl, aryl, heteroaryl, aminoalkyl, or arylalkyl are optionally substituted with one or more radical selected from the group consisting of: alkylsulfonamide, sulfamyl, alkyl, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, aminoalkyl, alkylaminoalkyl, alkoxy, halogen, acyloxy, oxy, formyl, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, nitro, azido, benzyloxy, dialkylaminoacyl, thioalkyl, aminoacyloxy, thiocyanate, isothiocyanate, alkyldioxy, hydroxyalkyl, alkylamino, alkyloxycarbonyl, alkoxyalkyl, alkenylamino, alkynylamino, alkenyl, alkynyl, dialkylaminoalkyloxy, and heterocyclic optionally substituted with alkyl, alkylamino, aminoalkyl, hydroxyalkyl, and alkylaminoalkyl;

25 **R¹⁰**

30 **R¹¹**

R¹⁰ is independently selected from the group consisting of: hydrido, lower alkyl, heteroaryl, heterocyclic, haloalkyl, arylalkylamino, heteroarylalkyl, aryl, and arylalkyl, wherein aryl, heteroaryl, heterocyclic, or arylalkyl are optionally substituted with one or more radical selected from alkyl, alkoxy, halogen, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, and heterocyclic;

5 **R^{10'}** is independently selected from the group consisting of: hydrido, lower alkyl, heteroaryl, heterocyclic, haloalkyl, arylalkylamino, heteroarylalkyl, aryl, and arylalkyl, wherein aryl, heteroaryl, heterocyclic, or arylalkyl are optionally substituted with one or more radical selected from alkyl, alkoxy, halogen, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, and heterocyclic;

10 **R¹¹** is hydrido;

15 **R¹²** is hydrido;

20 **R¹³** is selected from the group consisting of: hydrido, alkyl, aryl, arylalkyl, heteroaryl, heterocyclicalkyl, and heteroarylalkyl, wherein aryl, alkyl, arylalkyl, heteroaryl, heterocyclicalkyl, or heteroarylalkyl are optionally substituted with one or more radicals selected from the group consisting of OR¹⁴, N(R¹⁴)R^{14'}, and glycols;

25 **R¹⁴** is independently selected from the group consisting of hydrido, and lower alkyl; and

R^{14'} is independently selected from the group consisting of hydrido, and lower alkyl;

30 **R¹⁵** is selected from the group consisting of: hydrido, halogen, alkyl, cycloalkyl, aryl, haloalkyl, heteroaryl, heterocyclic, alkylalkene, alkylalkyne, hydroxy, hydroxyalkyl, alkylhydroxy, amino, aminoalkyl, alkylamino, alkylaminoalkyl, alkylhydroxyalkyl, nitro, cyano, alkylthio, alkylsulfinyl, alkylsulfonyl; wherein aryl or arylalkyl are optionally substituted with one or more radical selected from alkyl, alkoxy, halo,

haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, heterocyclic; and

5 **R¹⁶** is selected from the group consisting of: hydrido, halogen, alkyl, cycloalkyl, aryl, haloalkyl, heteroaryl, heterocyclic, alkylalkene, alkylalkyne, hydroxy, hydroxyalkyl, alkylhydroxy, amino, aminoalkyl, alkylamino, alkylaminoalkyl, alkylhydroxyalkyl, nitro, cyano, alkylthio, alkylsulfinyl, alkylsulfonyl; wherein aryl or arylalkyl are optionally substituted with one or more radical selected from alkyl, alkoxy, halo, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, heterocyclic, wherein


10 **R¹⁵** and **R¹⁶** may be taken together to form a 5 to 7 membered saturated or unsaturated carbocyclic ring optionally containing 0 to 3 heteroatoms selected from the group consisting of N, O, or S, and wherein said ring is optionally substituted with **R¹**;

15

or isomers, tautomers, carriers, esters, prodrugs, pharmaceutically acceptable salts thereof.

6. The compound of claim 2 of formula II

20



A is $(CH_2)_m-CR^{15}=CR^{16}-(CH_2)_m$

m is independently selected from 0, 1, or 2;

B is a 5 or 6 membered heteroaryl, aryl, saturated or unsaturated heterocyclic wherein said aryl, heteroaryl, or heterocyclic are optionally substituted with R^1 , R^2 , and R^{12} ;

5 R^1 is selected from the group consisting of: hydrido, halogen, alkyl, aryl, heteroaryl, alkenyl, alkynyl, haloalkyl, CN, NO_2 , OR^5 , $OCOOR^5$, CO_2R^7 , $CON(R^6)R^7$, COR^6 , SR^6 , SOR^6 , SO_2R^6 , NR^6R^7 , NR^6COR^7 , NR^6CONHR^7 , $NR^6SO_2R^7$, $NR^6SO_2NHR^7$, and $SO_2N(R^6)R^7$ wherein R^6 and R^7 may be taken together to form a 3-7 membered carbocyclic ring having 1 to 3 substituted or unsubstituted heteroatoms selected from the group consisting of: S, SO, SO_2 , O, and NR^6 ; wherein said alkenyl, alkynyl, alkyl, aryl, heteroaryl or OR^5 are optional substituted with, hydrido, halogen, alkyl, hydroxyalkyl, aryl, heteroaryl, haloalkyl, $COCF_3$, CN, NO_2 , OR^5 , $OCOOR^5$, CO_2R^7 , $CON(R^6)R^7$, COR^6 , SR^6 , SOR^6 , SO_2R^6 , NR^6R^7 , NR^6COR^7 , NR^6CONHR^7 , $NR^6SO_2R^7$, $NR^6SO_2NHR^7$, and $SO_2N(R^6)R^7$ wherein R^6 and R^7 may be taken together to form a 3-7 membered carbocyclic ring having 1 to 3 substituted or unsubstituted heteroatoms selected from the group consisting of: S, SO, SO_2 , O, and NR^6 ;

10 R^2 is hydrido;

20 R^3 is selected from the group consisting of: substituted or unsubstituted amidine, alkylamino, aminoalkyl, $CONHR^7$, NH_2 , $NHCOR^6$, and CH_2NHCOR^6 ;

25 R^4 is selected from the group consisting of: halogen, alkylsulfinyl, alkylsulfonyl, cyano, alkoxy carbonyl, alkyl, haloalkyl, hydrido, hydroxyalkyl, haloalkoxy, heterocyclic, nitro, acylamino, aryl, heteroaryl, and alkenyl, OR^{13} , SR^8 , $SO_2N(R^8)R^{8'}$, NHR^9 , $NHCOR^9$, NR^9COR^9 , $NHCO(OR^9)$, $NR^9CO(OR^9)$, $NR^8SO_2R^{10'}$, $NHSO_2N(R^{10'})R^{10'}$, $NR^6CON(R^{10'})R^{10'}$, COR^9 , CO_2R^8 , $CON(R^8)R^{8'}$, wherein R^8 and $R^{8'}$ may be taken together to form a 3-7 membered carbocyclic ring having 1 to 3 substituted or unsubstituted heteroatoms selected from S, SO, SO_2 , O, N, and NR^6 , and wherein $R^{10'}$ and $R^{10'}$ may be taken together to form a 3-7

membered carbocyclic ring having 1 to 3 substituted or unsubstituted heteroatoms selected from S, SO, SO₂, O, N, and NR⁶ wherein said aryl, heterocyclic, heteroaryl, or alkenyl are optionally substituted with R⁹;

5 R⁵ is selected from the group consisting of: hydrido, alkyl, aryl, arylalkyl, heteroaryl, heterocyclicalkyl, and heteroarylalkyl, wherein aryl, alkyl, arylalkyl, heteroaryl, heterocyclicalkyl, or heteroarylalkyl are optionally substituted with one or more radicals selected from the group consisting of OR¹⁴, N(R¹⁴)R¹⁴, and glycols;

10 R⁶ is independently selected from the group consisting of: hydrido, aryl, heteroaryl, lower alkyl, haloalkyl, alkenyl, alkynyl, hydroxyalkyl, aminoalkyl, alkylaminoalkyl, alkoxy, alkoxyalkyl, heterocyclicalkyl, and heterocyclic;

15 R⁷ is independently selected from the group consisting of: hydrido, aryl, heteroaryl, lower alkyl, haloalkyl, alkenyl, alkynyl, hydroxyalkyl, aminoalkyl, alkylaminoalkyl, alkoxy, alkoxyalkyl, heterocyclicalkyl, and heterocyclic;

20 R⁸ is independently selected from the group consisting of: hydrido, aryl, heteroaryl, arylalkyl, heterocyclic, haloalkyl, arylalkylamino, alkylaminoalkyl, dialkylaminoalkyl, alkyl, alkenyl, alkynyl, heteroarylalkyl, and heterocyclicalkyl;

25 R^{8'} is independently selected from the group consisting of: hydrido, aryl, heteroaryl, arylalkyl, heterocyclic, haloalkyl, arylalkylamino, alkylaminoalkyl, dialkylaminoalkyl, alkyl, alkenyl, alkynyl, heteroarylalkyl, and heterocyclicalkyl;

30 R⁹ is independently selected from the group consisting of: hydrido, lower alkyl, aryl, heteroaryl, arylalkyl, heterocyclic, cycloalkyl, heterocyclicalkyl, haloalkyl, arylalkylamino, amino, aminoalkyl, aminoacyl, nitro, azido, and heteroarylalkyl, wherein alkyl, aryl, heteroaryl, aminoalkyl, or arylalkyl are optionally substituted with one or more radical selected from the group consisting of: alkylsulfonamide, sulfamyl, alkyl, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino,

aminoalkyl, alkylaminoalkyl, alkoxy, halogen, acyloxy, oxy, formyl, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, nitro, azido, benzyloxy, dialkylaminoacyl, thioalkyl, aminoacyloxy, thiocyanate, isothiocyanate, alkyldioxy, hydroxyalkyl, alkylamino, alkyloxycarbonyl, alkoxyalkyl, alkenylamino, alkynylamino, alkenyl, alkynyl, dialkylaminoalkyloxy, and heterocyclic optionally substituted with alkyl, alkylamino, aminoalkyl, hydroxyalkyl, and alkylaminoalkyl;

5 **R¹⁰** is independently selected from the group consisting of: hydrido, lower alkyl, heteroaryl, heterocyclic, haloalkyl, arylalkylamino, heteroarylalkyl, aryl, and arylalkyl, wherein aryl, heteroaryl, heterocyclic, or arylalkyl are optionally substituted with one or more radical selected from alkyl, alkoxy, halogen, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, and heterocyclic;

10 **R^{10'}** is independently selected from the group consisting of: hydrido, lower alkyl, heteroaryl, heterocyclic, haloalkyl, arylalkylamino, heteroarylalkyl, aryl, and arylalkyl, wherein aryl, heteroaryl, heterocyclic, or arylalkyl are optionally substituted with one or more radical selected from alkyl, alkoxy, halogen, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, and heterocyclic;

15 **R^{10'}** is independently selected from the group consisting of: hydrido, lower alkyl, heteroaryl, heterocyclic, haloalkyl, arylalkylamino, heteroarylalkyl, aryl, and arylalkyl, wherein aryl, heteroaryl, heterocyclic, or arylalkyl are optionally substituted with one or more radical selected from alkyl, alkoxy, halogen, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, and heterocyclic;

20 **R¹¹** is hydrido;

25 **R¹²** is hydrido;

25 **R¹³** is selected from the group consisting of: hydrido, alkyl, aryl, arylalkyl, heteroaryl, heterocyclicalkyl, and heteroarylalkyl, wherein aryl, alkyl, arylalkyl, heteroaryl, heterocyclicalkyl, or heteroarylalkyl are optionally substituted with one or more radicals selected from the group consisting of OR¹⁴, N(R¹⁴)R^{14'}, and glycols;

30 **R¹⁴** is independently selected from the group consisting of hydrido, and lower alkyl; and


R^{14}' is independently selected from the group consisting of hydrido, and lower alkyl;

R^{15} is selected from the group consisting of: hydrido, halogen, alkyl, cycloalkyl, aryl, haloalkyl, heteroaryl, heterocyclic, alkylalkene, 5 alkylalkyne, hydroxy, hydroxyalkyl, alkylhydroxy, amino, aminoalkyl, alkylamino, alkylaminoalkyl, alkylhydroxyalkyl, nitro, cyano, alkylthio, alkylsulfinyl, alkylsulfonyl; wherein aryl or arylalkyl are optionally substituted with one or more radical selected from alkyl, alkoxy, halo, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, 10 phenoxy, benzyloxy, dialkylaminoalkyloxy, heterocyclic; and

R^{16} is selected from the group consisting of: hydrido, halogen, alkyl, cycloalkyl, aryl, haloalkyl, heteroaryl, heterocyclic, alkylalkene, alkylalkyne, hydroxy, hydroxyalkyl, alkylhydroxy, amino, aminoalkyl, alkylamino, alkylaminoalkyl, alkylhydroxyalkyl, nitro, cyano, alkylthio, 15 alkylsulfinyl, alkylsulfonyl; wherein aryl or arylalkyl are optionally substituted with one or more radical selected from alkyl, alkoxy, halo, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, heterocyclic; wherein R^{15} and R^{16} may be taken together to form a 5 to 7 membered saturated or 20 unsaturated carbocyclic ring optionally containing 0 to 3 heteroatoms selected from the group consisting of N, O, or S, and wherein said ring is optionally substituted with R^1 ;

25 or isomers, tautomers, carriers, esters, prodrugs, pharmaceutically acceptable salts thereof.

7. The compound of claim 2 of the formula



wherein

B is a 5 or 6 membered heteroaryl, aryl, saturated or unsaturated

5 heterocyclic wherein said aryl, heteroaryl, or heterocyclic are optionally substituted with R¹, R², and R¹²;

R¹ is selected from the group consisting of: hydrido, halogen, alkyl, aryl, heteroaryl, alkenyl, alkynyl, haloalkyl, CN, NO₂, OR⁵, OCOOR⁵,

CO₂R⁷, CON(R⁶)R⁷, COR⁶, SR⁶, SOR⁶, SO₂R⁶, NR⁶R⁷, NR⁶COR⁷,

10 NR⁶CONHR⁷, NR⁶SO₂R⁷, NR⁶SO₂NHR⁷, and SO₂N(R⁶)R⁷ wherein R⁶ and R⁷ may be taken together to form a 3-7 membered carbocyclic ring

having 1 to 3 substituted or unsubstituted heteroatoms selected from the group consisting of: S, SO, SO₂, O, and NR⁶; wherein said alkenyl, alkynyl, alkyl, aryl, heteroaryl or OR⁵ are optional substituted with,

15 hydrido, halogen, alkyl, hydroxyalkyl, aryl, heteroaryl, haloalkyl, COCF₃, CN, NO₂, OR⁵, OCOOR⁵, CO₂R⁷, CON(R⁶)R⁷, COR⁶, SR⁶,

SOR⁶, SO₂R⁶, NR⁶R⁷, NR⁶COR⁷, NR⁶CONHR⁷, NR⁶SO₂R⁷,

NR⁶SO₂NHR⁷, and SO₂N(R⁶)R⁷ wherein R⁶ and R⁷ may be taken

together to form a 3-7 membered carbocyclic ring having 1 to 3

20 substituted or unsubstituted heteroatoms selected from the group consisting of: S, SO, SO₂, O, and NR⁶;

R² is selected from the group consisting of: halogen, hydrido,

hydroxyalkyl, alkyl, OR⁶, CN, NO₂, SR⁶, NHR⁶, CON(R⁶)R⁷,

NHCONHR⁶, CO₂H, and haloalkyl;

R^1 and R^2 may be taken together to form a 5 to 7 membered saturated or unsaturated carbocyclic ring optionally containing 0 to 3 heteroatoms selected from the group consisting of N, O, or S, and wherein said ring is optionally substituted with R^1 ;

5 R^3 is selected from the group consisting of: substituted or unsubstituted amidine, alkylamino, aminoalkyl, $CONHR^7$, NH_2 , $NHCOR^6$, and CH_2NHCOR^6 ;

R^4 is selected from the group consisting of: halogen, alkylsulfinyl, alkylsulfonyl, cyano, alkoxycarbonyl, alkyl, haloalkyl, hydrido, hydroxyalkyl, haloalkoxy, heterocyclic, nitro, acylamino, aryl, heteroaryl, and alkenyl, OR^{13} , SR^8 , $SO_2N(R^8)R^{8'}$, NHR^9 , $NHCOR^9$, NR^9COR^9 , $NHCO(OR^9)$, $NR^9CO(OR^9)$, $NR^8SO_2R^{10}$, $NHSO_2N(R^{10})R^{10'}$, $NR^6CON(R^{10})R^{10'}$, COR^9 , CO_2R^8 , $CON(R^8)R^{8'}$, wherein R^8 and $R^{8'}$ may be taken together to form a 3-7 membered carbocyclic ring having 1 to 3 substituted or unsubstituted heteroatoms selected from S, SO, SO_2 , O, N, and NR^6 , and wherein R^{10} and $R^{10'}$ may be taken together to form a 3-7 membered carbocyclic ring having 1 to 3 substituted or unsubstituted heteroatoms selected from S, SO, SO_2 , O, N, and NR^6 wherein said aryl, heterocyclic, heteroaryl, or alkenyl are optionally substituted with R^9 ;

10 R^5 is selected from the group consisting of: hydrido, alkyl, aryl, arylalkyl, heteroaryl, heterocyclicalkyl, and heteroarylalkyl, wherein aryl, alkyl, arylalkyl, heteroaryl, heterocyclicalkyl, or heteroarylalkyl are optionally substituted with one or more radicals selected from the group consisting of OR^{14} , $N(R^{14})R^{14'}$, and glycols;

15 R^6 is independently selected from the group consisting of: hydrido, aryl, heteroaryl, lower alkyl, haloalkyl, alkenyl, alkynyl, hydroxyalkyl, aminoalkyl, alkylaminoalkyl, alkoxy, alkoxyalkyl, heterocyclicalkyl, and heterocyclic;

20 R^7 is independently selected from the group consisting of: hydrido, aryl, heteroaryl, lower alkyl, haloalkyl, alkenyl, alkynyl, hydroxyalkyl,

25 R^8 and $R^{8'}$ are taken together to form a 3-7 membered carbocyclic ring having 1 to 3 substituted or unsubstituted heteroatoms selected from S, SO, SO_2 , O, N, and NR^6 wherein said aryl, heterocyclic, heteroaryl, or alkenyl are optionally substituted with R^9 ;

30 R^9 is independently selected from the group consisting of: hydrido, aryl, heteroaryl, lower alkyl, haloalkyl, alkenyl, alkynyl, hydroxyalkyl,

aminoalkyl, alkylaminoalkyl, alkoxy, alkoxyalkyl, heterocyclicalkyl, and heterocyclic;

5 **R⁸** is independently selected from the group consisting of: hydrido, aryl, heteroaryl, arylalkyl, heterocyclic, haloalkyl, arylalkylamino, alkylaminoalkyl, dialkylaminoalkyl, alkyl, alkenyl, alkynyl, heteroarylalkyl, and heterocyclicalkyl;

10 **R^{8'}** is independently selected from the group consisting of: hydrido, aryl, heteroaryl, arylalkyl, heterocyclic, haloalkyl, arylalkylamino, alkylaminoalkyl, dialkylaminoalkyl, alkyl, alkenyl, alkynyl, heteroarylalkyl, and heterocyclicalkyl;

15 **R⁹** is independently selected from the group consisting of: hydrido, lower alkyl, aryl, heteroaryl, arylalkyl, heterocyclic, cycloalkyl, heterocyclicalkyl, haloalkyl, arylalkylamino, amino, aminoalkyl, aminoacyl, nitro, azido, and heteroarylalkyl, wherein alkyl, aryl, heteroaryl, aminoalkyl, or arylalkyl are optionally substituted with one or more radical selected from the group consisting of: alkylsulfonamide, sulfamyl, alkyl, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, aminoalkyl, alkylaminoalkyl, alkoxy, halogen, acyloxy, oxy, formyl, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, nitro, azido, benzyloxy, dialkylaminoacyl, thioalkyl, aminoacyloxy, thiocyanate, isothiocyanate, alkyldioxy, hydroxyalkyl, alkylamino, alkyloxycarbonyl, alkoxyalkyl, alkenylamino, alkynylamino, alkenyl, alkynyl, dialkylaminoalkyloxy, and heterocyclic optionally substituted with alkyl, alkylamino, aminoalkyl, hydroxyalkyl, and alkylaminoalkyl;

20 **R¹⁰** is independently selected from the group consisting of: hydrido, lower alkyl, heteroaryl, heterocyclic, haloalkyl, arylalkylamino, heteroarylalkyl, aryl, and arylalkyl, wherein aryl, heteroaryl, heterocyclic, or arylalkyl are optionally substituted with one or more radical selected from alkyl, alkoxy, halogen, haloalkyl, cyano,

haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, and heterocyclic;

5 $R^{10'}$ is independently selected from the group consisting of: hydrido, lower alkyl, heteroaryl, heterocyclic, haloalkyl, arylalkylamino, heteroarylalkyl, aryl, and arylalkyl, wherein aryl, heteroaryl, heterocyclic, or arylalkyl are optionally substituted with one or more radical selected from alkyl, alkoxy, halogen, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, and heterocyclic;

10 R^{11} is selected from the group consisting of: hydrido, halogen, haloalkyl, CN, CO_2R^5 , lower alkyl, lower alkenyl, lower alkynyl, alkoxy, and $CONH_2$;

R^{12} is selected from the group consisting of: hydrido, halogen, alkyl, and alkoxy;

15 R^{13} is selected from the group consisting of: hydrido, alkyl, aryl, arylalkyl, heteroaryl, heterocyclicalkyl, and heteroarylalkyl, wherein aryl, alkyl, arylalkyl, heteroaryl, heterocyclicalkyl, or heteroarylalkyl are optionally substituted with one or more radicals selected from the group consisting of OR^{14} , $N(R^{14})R^{14'}$, and glycols;

20 R^{14} is independently selected from the group consisting of hydrido, and lower alkyl;

$R^{14'}$ is independently selected from the group consisting of hydrido, and lower alkyl;

25 R^{15} is selected from the group consisting of: hydrido, halogen, alkyl, cycloalkyl, aryl, haloalkyl, heteroaryl, heterocyclic, alkylalkene, alkylalkyne, hydroxy, hydroxyalkyl, alkylhydroxy, amino, aminoalkyl, alkylamino, alkylaminoalkyl, alkylhydroxyalkyl, nitro, cyano, alkylthio, alkylsulfinyl, alkylsulfonyl; wherein aryl or arylalkyl are optionally substituted with one or more radical selected from alkyl, alkoxy, halo, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, heterocyclic; and

R¹⁶ is selected from the group consisting of: hydrido, halogen, alkyl, cycloalkyl, aryl, haloalkyl, heteroaryl, heterocyclic, alkylalkene, alkylalkyne, hydroxy, hydroxyalkyl, alkylhydroxy, amino, aminoalkyl, alkylamino, alkylaminoalkyl, alkylhydroxyalkyl, nitro, cyano, alkylthio, alkylsulfinyl, alkylsulfonyl; wherein aryl or arylalkyl are optionally substituted with one or more radical selected from alkyl, alkoxy, halo, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, heterocyclic; wherein R¹⁵ and R¹⁶ may be taken together to form a 5 to 7 membered saturated or unsaturated carbocyclic ring optionally containing 0 to 3 heteroatoms selected from the group consisting of N, O, or S, and wherein said ring is optionally substituted with R¹;

or isomers, tautomers, carriers, esters, prodrugs, pharmaceutically acceptable salts thereof.

15

8. The compound of claim 2

wherein

20 A is (CH₂)_m-CR¹⁵=CR¹⁶-(CH₂)_m
m is independently selected from 0, 1, or 2;

25 B is a 5 or 6 membered heteroaryl, aryl, saturated or unsaturated heterocyclic wherein said aryl, heteroaryl, or heterocyclic are optionally substituted with R¹, R², and R¹²;

R¹ is selected from the group consisting of: SOR⁶, SO₂R⁶, NR⁶R⁷, NR⁶COR⁷, NR⁶SO₂R⁷, and SO₂N(R⁶)R⁷ wherein R⁶ and R⁷ may be taken together to form a 3-7 membered carbocyclic ring having 1 to 3 substituted or unsubstituted heteroatoms selected from the group consisting of: S, SO, SO₂, O, and NR⁶;

30 R² is hydrido;

R³ is selected from the group consisting of: CONHR⁷, NH₂, NHCOR⁶, and CH₂NHCOR⁶;

R⁴ is selected from the group consisting of: halogen, alkylsulfinyl, alkylsulfonyl, cyano, alkoxycarbonyl, alkyl, haloalkyl, hydrido, hydroxyalkyl, haloalkoxy, heterocyclic, nitro, acylamino, aryl, heteroaryl, and alkenyl, OR¹³, SR⁸, SO₂N(R⁸)R^{8'}, NHR⁹, NHCOR⁹, NR⁹COR⁹, NHCO(OR⁹), NR⁹CO(OR⁹), NR⁸SO₂R¹⁰, NHSO₂N(R¹⁰)R^{10'}, NR⁶CON(R¹⁰)R^{10'}, COR⁹, CO₂R⁸, CON(R⁸)R^{8'}, wherein R⁸ and R^{8'} may be taken together to form a 3-7 membered carbocyclic ring having 1 to 3 substituted or unsubstituted heteroatoms selected from S, SO, SO₂, O, N, and NR⁶, and wherein R¹⁰ and R^{10'} may be taken together to form a 3-7 membered carbocyclic ring having 1 to 3 substituted or unsubstituted heteroatoms selected from S, SO, SO₂, O, N, and NR⁶ wherein said aryl, heterocyclic, heteroaryl, or alkenyl are optionally substituted with R⁹;

R⁵ is selected from the group consisting of: hydrido, alkyl, aryl, arylalkyl, heteroaryl, heterocyclicalkyl, and heteroarylalkyl, wherein aryl, alkyl, arylalkyl, heteroaryl, heterocyclicalkyl, or heteroarylalkyl are optionally substituted with one or more radicals selected from the group consisting of OR¹⁴, N(R¹⁴)R^{14'}, and glycols;

R⁶ is independently selected from the group consisting of: hydrido, aryl, heteroaryl, lower alkyl, haloalkyl, alkenyl, alkynyl, hydroxyalkyl, aminoalkyl, alkylaminoalkyl, alkoxy, alkoxyalkyl, heterocyclicalkyl, and heterocyclic;

R⁷ is independently selected from the group consisting of: hydrido, aryl, heteroaryl, lower alkyl, haloalkyl, alkenyl, alkynyl, hydroxyalkyl, aminoalkyl, alkylaminoalkyl, alkoxy, alkoxyalkyl, heterocyclicalkyl, and heterocyclic;

R⁸ is independently selected from the group consisting of: hydrido, aryl, heteroaryl, arylalkyl, heterocyclic, haloalkyl, arylalkylamino, alkylaminoalkyl, dialkylaminoalkyl, alkyl, alkenyl, alkynyl, heteroarylalkyl, and heterocyclicalkyl;

R^{8'} is independently selected from the group consisting of: hydrido, aryl, heteroaryl, arylalkyl, heterocyclic, haloalkyl, arylalkylamino, alkylaminoalkyl, dialkylaminoalkyl, alkyl, alkenyl, alkynyl, heteroarylalkyl, and heterocyclicalkyl;

5 **R⁹** is independently selected from the group consisting of: hydrido, lower alkyl, aryl, heteroaryl, arylalkyl, heterocyclic, cycloalkyl, heterocyclicalkyl, haloalkyl, arylalkylamino, amino, aminoalkyl, aminoacyl, nitro, azido, and heteroarylalkyl, wherein alkyl, aryl, heteroaryl, aminoalkyl, or arylalkyl are optionally substituted with one or

10 more radical selected from the group consisting of: alkylsulfonamide, sulfamyl, alkyl, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, aminoalkyl, alkylaminoalkyl, alkoxy, halogen, acyloxy, oxy, formyl, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, nitro, azido, benzyloxy, dialkylaminoacyl, thioalkyl, aminoacyloxy, thiocyanate, isothiocyanate, alkyldioxy, hydroxyalkyl, alkylamino, alkyloxycarbonyl, alkoxyalkyl, alkenylamino, alkynylamino, alkenyl, alkynyl, dialkylaminoalkyloxy, and heterocyclic optionally substituted with alkyl, alkylamino, aminoalkyl, hydroxyalkyl, and alkylaminoalkyl;

15 **R¹⁰** is independently selected from the group consisting of: hydrido, lower alkyl, heteroaryl, heterocyclic, haloalkyl, arylalkylamino, heteroarylalkyl, aryl, and arylalkyl, wherein aryl, heteroaryl, heterocyclic, or arylalkyl are optionally substituted with one or more radical selected from alkyl, alkoxy, halogen, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, and heterocyclic;

20 **R^{10'}** is independently selected from the group consisting of: hydrido, lower alkyl, heteroaryl, heterocyclic, haloalkyl, arylalkylamino, heteroarylalkyl, aryl, and arylalkyl, wherein aryl, heteroaryl, heterocyclic, or arylalkyl are optionally substituted with one or more radical selected from alkyl, alkoxy, halogen, haloalkyl, cyano,

25 haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, and heterocyclic;

30 **R^{10''}** is independently selected from the group consisting of: hydrido, lower alkyl, heteroaryl, heterocyclic, haloalkyl, arylalkylamino, heteroarylalkyl, aryl, and arylalkyl, wherein aryl, heteroaryl, heterocyclic, or arylalkyl are optionally substituted with one or more radical selected from alkyl, alkoxy, halogen, haloalkyl, cyano,

haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, and heterocyclic;

5 **R¹¹** is hydrido, halogen, haloalkyl, CN, CO₂R⁵, lower alkyl, lower alkenyl, lower alkynyl, alkoxy, and CONH₂;

R¹² is hydrido;

10 **R¹³** is selected from the group consisting of: hydrido, alkyl, aryl, arylalkyl, heteroaryl, heterocyclicalkyl, and heteroarylalkyl, wherein aryl, alkyl, arylalkyl, heteroaryl, heterocyclicalkyl, or heteroarylalkyl are optionally substituted with one or more radicals selected from the group consisting of OR¹⁴, N(R¹⁴)R¹⁴, and glycols;

R¹⁴ is independently selected from the group consisting of hydrido, and lower alkyl;

15 **R^{14'}** is independently selected from the group consisting of hydrido, and lower alkyl;

R¹⁵ is selected from the group consisting of: hydrido, halogen, alkyl, cycloalkyl, aryl, haloalkyl, heteroaryl, heterocyclic, alkylalkene, alkylalkyne, hydroxy, hydroxyalkyl, alkylhydroxy, amino, aminoalkyl, alkylamino, alkylaminoalkyl, alkylhydroxyalkyl, nitro, cyano, alkylthio, alkylsulfinyl, alkylsulfonyl; wherein aryl or arylalkyl are optionally substituted with one or more radical selected from alkyl, alkoxy, halo, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, heterocyclic; and

20 **R¹⁶** is selected from the group consisting of: hydrido, halogen, alkyl, cycloalkyl, aryl, haloalkyl, heteroaryl, heterocyclic, alkylalkene, alkylalkyne, hydroxy, hydroxyalkyl, alkylhydroxy, amino, aminoalkyl, alkylamino, alkylaminoalkyl, alkylhydroxyalkyl, nitro, cyano, alkylthio, alkylsulfinyl, alkylsulfonyl; wherein aryl or arylalkyl are optionally substituted with one or more radical selected from alkyl, alkoxy, halo, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, heterocyclic; wherein

25 **R¹⁵** and **R¹⁶** may be taken together to form a 5 to 7 membered saturated or

unsaturated carbocyclic ring optionally containing 0 to 3 heteroatoms selected from the group consisting of N, O, or S, and wherein said ring is optionally substituted with R¹;

5 or isomers, tautomers, carriers, esters, prodrugs, pharmaceutically acceptable salts thereof.

9. The compound of claim 2

10

wherein

A is (CH₂)_m-CR¹⁵=CR¹⁶-(CH₂)_m

m is independently selected from 0, 1, or 2;

15 **B** is a 5 or 6 membered heteroaryl, aryl, saturated or unsaturated heterocyclic wherein said aryl, heteroaryl, or heterocyclic are optionally substituted with R¹, R², and R¹²;

R¹ is selected from the group consisting of: SOR⁶, SO₂R⁶, and SO₂N(R⁶)R⁷ wherein R⁶ and R⁷ may be taken together to form a 3-7 membered carbocyclic ring having 1 to 3 substituted or unsubstituted

20 heteroatoms selected from the group consisting of: S, SO, SO₂, O, and NR⁶; wherein said alkenyl, alkynyl, alkyl, aryl, heteroaryl or OR⁵ are optional substituted with, hydrido, halogen, alkyl, hydroxyalkyl, aryl, heteroaryl, haloalkyl, COCF₃, CN, NO₂, OR⁵, OCOOR⁵, CO₂R⁷, CON(R⁶)R⁷, COR⁶, SR⁶, SOR⁶, SO₂R⁶, NR⁶R⁷, NR⁶COR⁷,

25 NR⁶CONHR⁷, NR⁶SO₂R⁷, NR⁶SO₂NHR⁷, and SO₂N(R⁶)R⁷ wherein R⁶ and R⁷ may be taken together to form a 3-7 membered carbocyclic ring having 1 to 3 substituted or unsubstituted heteroatoms selected from the group consisting of: S, SO, SO₂, O, and NR⁶;

R² is hydrido;

30 **R³** is CONHR⁷;

R⁴ is selected from the group consisting of: halogen, alkylsulfinyl, alkylsulfonyl, cyano, alkoxycarbonyl, alkyl, haloalkyl, hydrido, hydroxyalkyl, haloalkoxy, heterocyclic, nitro, acylamino, aryl, heteroaryl, and alkenyl, OR¹³, SR⁸, SO₂N(R⁸)R^{8'}, NHR⁹, NHCOR⁹, NR⁹COR⁹, NHCO(OR⁹), NR⁹CO(OR⁹), NR⁸SO₂R¹⁰, NHSO₂N(R¹⁰)R^{10'}, NR⁶CON(R¹⁰)R^{10'}, COR⁹, CO₂R⁸, CON(R⁸)R^{8'}, wherein R⁸ and R^{8'} may be taken together to form a 3-7 membered carbocyclic ring having 1 to 3 substituted or unsubstituted heteroatoms selected from S, SO, SO₂, O, N, and NR⁶, and wherein R¹⁰ and R^{10'} may be taken together to form a 3-7 membered carbocyclic ring having 1 to 3 substituted or unsubstituted heteroatoms selected from S, SO, SO₂, O, N, and NR⁶ wherein said aryl, heterocyclic, heteroaryl, or alkenyl are optionally substituted with R⁹;

R⁵ is selected from the group consisting of: hydrido, alkyl, aryl, arylalkyl, heteroaryl, heterocyclicalkyl, and heteroarylalkyl, wherein aryl, alkyl, arylalkyl, heteroaryl, heterocyclicalkyl, or heteroarylalkyl are optionally substituted with one or more radicals selected from the group consisting of OR¹⁴, N(R¹⁴)R^{14'}, and glycols;

R⁶ is independently selected from the group consisting of: hydrido, aryl, heteroaryl, lower alkyl, haloalkyl, alkenyl, alkynyl, hydroxyalkyl, aminoalkyl, alkylaminoalkyl, alkoxy, alkoxyalkyl, heterocyclicalkyl, and heterocyclic;

R⁷ is independently selected from the group consisting of: hydrido, aryl, heteroaryl, lower alkyl, haloalkyl, alkenyl, alkynyl, hydroxyalkyl, aminoalkyl, alkylaminoalkyl, alkoxy, alkoxyalkyl, heterocyclicalkyl, and heterocyclic;

R⁸ is independently selected from the group consisting of: hydrido, aryl, heteroaryl, arylalkyl, heterocyclic, haloalkyl, arylalkylamino, alkylaminoalkyl, dialkylaminoalkyl, alkyl, alkenyl, alkynyl, heteroarylalkyl, and heterocyclicalkyl;

R^{8'} is independently selected from the group consisting of: hydrido, aryl, heteroaryl, arylalkyl, heterocyclic, haloalkyl, arylalkylamino,

alkylaminoalkyl, dialkylaminoalkyl, alkyl, alkenyl, alkynyl, heteroarylalkyl, and heterocyclicalkyl;

5 R^9 is independently selected from the group consisting of: hydrido, lower alkyl, aryl, heteroaryl, arylalkyl, heterocyclic, cycloalkyl, heterocyclicalkyl, haloalkyl, arylalkylamino, amino, aminoalkyl, aminoacyl, nitro, azido, and heteroarylalkyl, wherein alkyl, aryl, heteroaryl, aminoalkyl, or arylalkyl are optionally substituted with one or more radical selected from the group consisting of: alkylsulfonamide, sulfamyl, alkyl, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, aminoalkyl, alkylaminoalkyl, alkoxy, halogen, acyloxy, oxy, formyl, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, nitro, azido, benzyloxy, dialkylaminoacyl, thioalkyl, aminoacyloxy, thiocyanate, isothiocyanate, alkyldioxy, hydroxyalkyl, alkylamino, alkyloxycarbonyl, alkoxyalkyl, alkenylamino, alkynylamino, alkenyl, alkynyl, dialkylaminoalkyloxy, and heterocyclic optionally substituted with alkyl, alkylamino, aminoalkyl, hydroxyalkyl, and alkylaminoalkyl;

10 R^{10} is independently selected from the group consisting of: hydrido, lower alkyl, heteroaryl, heterocyclic, haloalkyl, arylalkylamino, heteroarylalkyl, aryl, and arylalkyl, wherein aryl, heteroaryl, heterocyclic, or arylalkyl are optionally substituted with one or more radical selected from alkyl, alkoxy, halogen, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, and heterocyclic;

15 $R^{10'}$ is independently selected from the group consisting of: hydrido, lower alkyl, heteroaryl, heterocyclic, haloalkyl, arylalkylamino, heteroarylalkyl, aryl, and arylalkyl, wherein aryl, heteroaryl, heterocyclic, or arylalkyl are optionally substituted with one or more radical selected from alkyl, alkoxy, halogen, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, and heterocyclic;

R¹¹ is selected from the group consisting of: hydrido, halogen, haloalkyl, CN, CO₂R⁵, lower alkyl, lower alkenyl, lower alkynyl, alkoxy, and CONH₂;

5 **R¹²** is selected from the group consisting of: hydrido, halogen, alkyl, and alkoxy;

R¹³ is selected from the group consisting of: hydrido, alkyl, aryl, arylalkyl, heteroaryl, heterocyclicalkyl, and heteroarylalkyl, wherein aryl, alkyl, arylalkyl, heteroaryl, heterocyclicalkyl, or heteroarylalkyl are optionally substituted with one or more radicals selected from the group consisting of OR¹⁴, N(R¹⁴)R^{14'}, and glycols;

10 **R¹⁴** is independently selected from the group consisting of hydrido, and lower alkyl; and

R^{14'} is independently selected from the group consisting of hydrido, and lower alkyl;

15 **R¹⁵** is selected from the group consisting of: hydrido, halogen, alkyl, cycloalkyl, aryl, haloalkyl, heteroaryl, heterocyclic, alkylalkene, alkylalkyne, hydroxy, hydroxyalkyl, alkylhydroxy, amino, aminoalkyl, alkylamino, alkylaminoalkyl, alkylhydroxyalkyl, nitro, cyano, alkylthio, alkylsulfinyl, alkylsulfonyl; wherein aryl or arylalkyl are optionally substituted with one or more radical selected from alkyl, alkoxy, halo, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, heterocyclic; and


20 **R¹⁶** is selected from the group consisting of: hydrido, halogen, alkyl, cycloalkyl, aryl, haloalkyl, heteroaryl, heterocyclic, alkylalkene, alkylalkyne, hydroxy, hydroxyalkyl, alkylhydroxy, amino, aminoalkyl, alkylamino, alkylaminoalkyl, alkylhydroxyalkyl, nitro, cyano, alkylthio, alkylsulfinyl, alkylsulfonyl; wherein aryl or arylalkyl are optionally substituted with one or more radical selected from alkyl, alkoxy, halo, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, heterocyclic; wherein

25 **R¹⁵** and **R¹⁶** may be taken together to form a 5 to 7 membered saturated or

unsaturated carbocyclic ring optionally containing 0 to 3 heteroatoms selected from the group consisting of N, O, or S, and wherein said ring is optionally substituted with R¹;

5

10. The compound of claim 6 of the formula



10 R⁶ is independently selected from the group consisting of: hydrido, aryl, heteroaryl, lower alkyl, haloalkyl, alkenyl, alkynyl, hydroxyalkyl, aminoalkyl, alkylaminoalkyl, alkoxy, alkoxyalkyl, heterocyclicalkyl, and heterocyclic;

15 R⁷ is independently selected from the group consisting of: hydrido, aryl, heteroaryl, lower alkyl, haloalkyl, alkenyl, alkynyl, hydroxyalkyl, aminoalkyl, alkylaminoalkyl, alkoxy, alkoxyalkyl, heterocyclicalkyl, and heterocyclic wherein R⁶ and R⁷ may be taken together to form a 3-7 membered carbocyclic ring having 1 to 3 substituted or unsubstituted heteroatoms selected from the group consisting of: S, SO, SO₂, O, and NR⁶;

20 R¹⁰ is independently selected from the group consisting of: hydrido, lower alkyl, heteroaryl, heterocyclic, haloalkyl, arylalkylamino, heteroarylalkyl, aryl, and arylalkyl, wherein aryl, heteroaryl, heterocyclic, or arylalkyl are optionally substituted with one or more radical selected from alkyl, alkoxy, halogen, haloalkyl, cyano,

25

haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, and heterocyclic;

5 **R¹⁵** is selected from the group consisting of: hydrido, halogen, alkyl, cycloalkyl, aryl, haloalkyl, heteroaryl, heterocyclic, alkylalkene, alkylalkyne, hydroxy, hydroxyalkyl, alkylhydroxy, amino, aminoalkyl, alkylamino, alkylaminoalkyl, alkylhydroxyalkyl, nitro, cyano, alkylthio, alkylsulfinyl, alkylsulfonyl; wherein aryl or arylalkyl are optionally substituted with one or more radical selected from alkyl, alkoxy, halo, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, heterocyclic; and


10 **R¹⁶** is selected from the group consisting of: hydrido, halogen, alkyl, cycloalkyl, aryl, haloalkyl, heteroaryl, heterocyclic, alkylalkene, alkylalkyne, hydroxy, hydroxyalkyl, alkylhydroxy, amino, aminoalkyl, alkylamino, alkylaminoalkyl, alkylhydroxyalkyl, nitro, cyano, alkylthio, alkylsulfinyl, alkylsulfonyl; wherein aryl or arylalkyl are optionally substituted with one or more radical selected from alkyl, alkoxy, halo, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, heterocyclic; wherein

15 **R¹⁵** and **R¹⁶** may be taken together to form a 5 to 7 membered saturated or unsaturated carbocyclic ring optionally containing 0 to 3 heteroatoms selected from the group consisting of N, O, or S, and wherein said ring is optionally substituted with **R¹**;

20 or isomers, tautomers, carriers, esters, prodrugs, pharmaceutically acceptable salts thereof.

25

11. The compound of claim 6 of the formula



5 \mathbf{R}^6 is independently selected from the group consisting of: hydrido, aryl, heteroaryl, lower alkyl, haloalkyl, alkenyl, alkynyl, hydroxyalkyl, aminoalkyl, alkylaminoalkyl, alkoxy, alkoxyalkyl, heterocyclicalkyl, and heterocyclic;

10 \mathbf{R}^7 is independently selected from the group consisting of: hydrido, aryl, heteroaryl, lower alkyl, haloalkyl, alkenyl, alkynyl, hydroxyalkyl, aminoalkyl, alkylaminoalkyl, alkoxy, alkoxyalkyl, heterocyclicalkyl, and heterocyclic wherein \mathbf{R}^6 and \mathbf{R}^7 may be taken together to form a 3-7 membered carbocyclic ring having 1 to 3 substituted or unsubstituted heteroatoms selected from the group consisting of: S, SO, SO₂, O, and NR⁶;

15 \mathbf{R}^9 is independently selected from the group consisting of: hydrido, lower alkyl, aryl, heteroaryl, arylalkyl, heterocyclic, cycloalkyl, heterocyclicalkyl, haloalkyl, arylalkylamino, amino, aminoalkyl, aminoacyl, nitro, azido, and heteroarylalkyl, wherein alkyl, aryl, heteroaryl, aminoalkyl, or arylalkyl are optionally substituted with one or more radical selected from the group consisting of: alkylsulfonamide, sulfamyl, alkyl, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, aminoalkyl, alkylaminoalkyl, alkoxy, halogen, acyloxy, oxy, formyl, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, nitro, azido, benzyloxy, dialkylaminoacyl, thioalkyl, aminoacyloxy, thiocyanate, isothiocyanate, alkyldioxy, hydroxyalkyl, alkylamino, alkyloxycarbonyl, alkoxyalkyl, alkenylamino, alkynylamino, alkenyl, alkynyl, dialkylaminoalkyloxy, and heterocyclic optionally


substituted with alkyl, alkylamino, aminoalkyl, hydroxyalkyl, and alkylaminoalkyl;

5 **R¹⁵** is selected from the group consisting of: hydrido, halogen, alkyl, cycloalkyl, aryl, haloalkyl, heteroaryl, heterocyclic, alkylalkene, alkylalkyne, hydroxy, hydroxyalkyl, alkylhydroxy, amino, aminoalkyl, alkylamino, alkylaminoalkyl, alkylhydroxyalkyl, nitro, cyano, alkylthio, alkylsulfinyl, alkylsulfonyl; wherein aryl or arylalkyl are optionally substituted with one or more radical selected from alkyl, alkoxy, halo, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, heterocyclic; and

10 **R¹⁶** is selected from the group consisting of: hydrido, halogen, alkyl, cycloalkyl, aryl, haloalkyl, heteroaryl, heterocyclic, alkylalkene, alkylalkyne, hydroxy, hydroxyalkyl, alkylhydroxy, amino, aminoalkyl, alkylamino, alkylaminoalkyl, alkylhydroxyalkyl, nitro, cyano, alkylthio, alkylsulfinyl, alkylsulfonyl; wherein aryl or arylalkyl are optionally substituted with one or more radical selected from alkyl, alkoxy, halo, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, heterocyclic; wherein
15 **R¹⁵** and **R¹⁶** may be taken together to form a 5 to 7 membered saturated or unsaturated carbocyclic ring optionally containing 0 to 3 heteroatoms selected from the group consisting of N, O, or S, and wherein said ring is optionally substituted with **R¹**;

20 or isomers, tautomers, carriers, esters, prodrugs, pharmaceutically acceptable salts thereof.

25 12. The compound of claim 6 of the formula



5 \mathbf{R}^6 is independently selected from the group consisting of: hydrido, aryl, heteroaryl, lower alkyl, haloalkyl, alkenyl, alkynyl, hydroxyalkyl, aminoalkyl, alkylaminoalkyl, alkoxy, alkoxyalkyl, heterocyclicalkyl, and heterocyclic;

10 \mathbf{R}^7 is independently selected from the group consisting of: hydrido, aryl, heteroaryl, lower alkyl, haloalkyl, alkenyl, alkynyl, hydroxyalkyl, aminoalkyl, alkylaminoalkyl, alkoxy, alkoxyalkyl, heterocyclicalkyl, and heterocyclic wherein \mathbf{R}^6 and \mathbf{R}^7 may be taken together to form a 3-7 membered carbocyclic ring having 1 to 3 substituted or unsubstituted heteroatoms selected from the group consisting of: S, SO, SO₂, O, and NR⁶;

15 \mathbf{R}^9 is independently selected from the group consisting of: hydrido, lower alkyl, aryl, heteroaryl, arylalkyl, heterocyclic, cycloalkyl, heterocyclicalkyl, haloalkyl, arylalkylamino, amino, aminoalkyl, aminoacyl, nitro, azido, and heteroarylalkyl, wherein alkyl, aryl, heteroaryl, aminoalkyl, or arylalkyl are optionally substituted with one or more radical selected from the group consisting of: alkylsulfonamide, sulfamyl, alkyl, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, aminoalkyl, alkylaminoalkyl, alkoxy, halogen, acyloxy, oxy, formyl, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, nitro, azido, benzyloxy, dialkylaminoacyl, thioalkyl, aminoacyloxy, thiocyanate, isothiocyanate, alkyldioxy, hydroxyalkyl, alkylamino, alkyloxycarbonyl, alkoxyalkyl, alkenylamino, alkynylamino, alkenyl, alkynyl, dialkylaminoalkyloxy, and heterocyclic optionally

substituted with alkyl, alkylamino, aminoalkyl, hydroxyalkyl, and alkylaminoalkyl;

5 R^{15} is selected from the group consisting of: hydrido, halogen, alkyl, cycloalkyl, aryl, haloalkyl, heteroaryl, heterocyclic, alkylalkene, alkylalkyne, hydroxy, hydroxyalkyl, alkylhydroxy, amino, aminoalkyl, alkylamino, alkylaminoalkyl, alkylhydroxyalkyl, nitro, cyano, alkylthio, alkylsulfinyl, alkylsulfonyl; wherein aryl or arylalkyl are optionally substituted with one or more radical selected from alkyl, alkoxy, halo, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, heterocyclic; and

10 R^{16} is selected from the group consisting of: hydrido, halogen, alkyl, cycloalkyl, aryl, haloalkyl, heteroaryl, heterocyclic, alkylalkene, alkylalkyne, hydroxy, hydroxyalkyl, alkylhydroxy, amino, aminoalkyl, alkylamino, alkylaminoalkyl, alkylhydroxyalkyl, nitro, cyano, alkylthio, alkylsulfinyl, alkylsulfonyl; wherein aryl or arylalkyl are optionally substituted with one or more radical selected from alkyl, alkoxy, halo, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, heterocyclic; wherein
15 R^{15} and R^{16} may be taken together to form a 5 to 7 membered saturated or unsaturated carbocyclic ring optionally containing 0 to 3 heteroatoms selected from the group consisting of N, O, or S, and wherein said ring is optionally substituted with R^1 ;

20 or isomers, tautomers, carriers, esters, prodrugs, pharmaceutically acceptable salts thereof.


25 13. The compound of claim 6 selected from the group consisting of:
30 8-amino-1-{4-[(diallylamino)sulfonyl]phenyl}-1H-benzo[g]indazole-3-carboxamide,

1-[4-(aminosulfonyl)phenyl]-8-[(3-methoxybenzoyl)amino]-1H-
benzo[g]indazole-3-carboxamide,
1-[4-(aminosulfonyl)phenyl]-8-[(3-chlorobenzoyl)amino]-1H-
benzo[g]indazole-3-carboxamide,
5 1-[4-(aminosulfonyl)phenyl]-8-[(3-(trifluoromethyl)benzoyl)amino]-
1H-benzo[g]indazole-3-carboxamide,
1-[4-(aminosulfonyl)phenyl]-8-[(3-methylbenzoyl)amino]-1H-
benzo[g]indazole-3-carboxamide,
1-[4-(aminosulfonyl)phenyl]-8-[(3-bromobenzoyl)amino]-1H-
10 benzo[g]indazole-3-carboxamide,
1-[4-(aminosulfonyl)phenyl]-8-[(pyridin-3-ylcarbonyl)amino]-1H-
benzo[g]indazole-3-carboxamide,
1-[4-(aminosulfonyl)phenyl]-8-[(2-chlorobenzoyl)amino]-1H-
benzo[g]indazole-3-carboxamide,
15 1-[4-(aminosulfonyl)phenyl]-8-[(3-bromophenyl)sulfonyl]amino]-1H-
benzo[g]indazole-3-carboxamide,
1-[4-(aminosulfonyl)phenyl]-8-[(3-chlorophenyl)sulfonyl]amino]-1H-
benzo[g]indazole-3-carboxamide,
1-[4-(aminosulfonyl)phenyl]-8-[(3-cyanobenzoyl)amino]-1H-
20 benzo[g]indazole-3-carboxamide,
1-[4-(aminosulfonyl)phenyl]-8-[(3-methylphenyl)sulfonyl]amino]-1H-
benzo[g]indazole-3-carboxamide,
1-[4-(aminosulfonyl)phenyl]-8-({[3-
25 (trifluoromethyl)phenyl]sulfonyl}amino)-1H-benzo[g]indazole-3-
carboxamide,
8-(acetylamino)-1-[4-(aminosulfonyl)phenyl]-1H-benzo[g]indazole-3-
carboxamide,
1-[4-(aminosulfonyl)phenyl]-8-[(methylsulfonyl)amino]-1H-
benzo[g]indazole-3-carboxamide,
30 1-[4-(aminosulfonyl)phenyl]-8-[(3,4-dichlorophenyl)sulfonyl]amino)-
1H-benzo[g]indazole-3-carboxamide,

1-[4-(aminosulfonyl)phenyl]-8-{{(2,4,5-trichlorophenyl)sulfonyl}amino}-1H-benzo[g]indazole-3-carboxamide,
1-[4-(aminosulfonyl)phenyl]-8-{{[3,5-bis(trifluoromethyl)phenyl}sulfonyl}amino)-1H-benzo[g]indazole-3-carboxamide,
5
1-[4-(aminosulfonyl)phenyl]-8-[(2-methoxybenzoyl)amino]-1H-benzo[g]indazole-3-carboxamide,
1-[4-(aminosulfonyl)phenyl]-8-{{[2-(trifluoromethyl)benzoyl}amino}-1H-benzo[g]indazole-3-carboxamide,
10
1-[4-(aminosulfonyl)phenyl]-8-[(2-methylbenzoyl)amino]-1H-benzo[g]indazole-3-carboxamide,
1-[4-(aminosulfonyl)phenyl]-8-[(2,6-dichlorobenzoyl)amino]-1H-benzo[g]indazole-3-carboxamide,
1-[4-(aminosulfonyl)phenyl]-8-{{[2-(trifluoromethoxy)benzoyl}amino}-1H-benzo[g]indazole-3-carboxamide,
15
1-[4-(aminosulfonyl)phenyl]-8-[(2,3-dichlorobenzoyl)amino]-1H-benzo[g]indazole-3-carboxamide,
1-[4-(aminosulfonyl)phenyl]-8-[(2-fluorobenzoyl)amino]-1H-benzo[g]indazole-3-carboxamide,
20
1-[4-(aminosulfonyl)phenyl]-8-{{(2-chloropyridin-3-yl)carbonyl}amino}-1H-benzo[g]indazole-3-carboxamide,
1-[4-(aminosulfonyl)phenyl]-8-{{(2-chlorophenyl)sulfonyl}amino}-1H-benzo[g]indazole-3-carboxamide,
1-[4-(aminosulfonyl)phenyl]-8-(isonicotinoylamino)-1H-benzo[g]indazole-3-carboxamide,
25
1-[3-(aminosulfonyl)phenyl]-8-[(2-chlorobenzoyl)amino]-1H-benzo[g]indazole-3-carboxamide,
8-[(2-chlorobenzoyl)amino]-1-[4-(methylsulfonyl)phenyl]-1H-benzo[g]indazole-3-carboxamide,
30
8-[(2-methylbenzoyl)amino]-1-[4-(methylsulfonyl)phenyl]-1H-benzo[g]indazole-3-carboxamide,

1-[4-(methylsulfonyl)phenyl]-8-{[2-(trifluoromethyl)benzoyl]amino}-1H-benzo[g]indazole-3-carboxamide,
 8-[(2,3-dichlorobenzoyl)amino]-1-[4-(methylsulfonyl)phenyl]-1H-benzo[g]indazole-3-carboxamide,
 5 8-{[(2-chloropyridin-3-yl)carbonyl]amino}-1-[4-(methylsulfonyl)phenyl]-1H-benzo[g]indazole-3-carboxamide, and
 8-[(2-fluorobenzoyl)amino]-1-[4-(methylsulfonyl)phenyl]-1H-benzo[g]indazole-3-carboxamide.

10 14. The compound of claim 2 of the formula



wherein

15 \mathbf{A} is $(\text{CH}_2)_m\text{-CR}^{15}=\text{CR}^{16}\text{-}(\text{CH}_2)_m$
 \mathbf{m} is independently selected from 0, 1, or 2;
 \mathbf{R}^4 is selected from the group consisting of: halogen, alkylsulfinyl, alkylsulfonyl, cyano, alkoxy carbonyl, alkyl, haloalkyl, hydrido, hydroxyalkyl, haloalkoxy, heterocyclic, nitro, acylamino, aryl, heteroaryl, and alkenyl, OR^{13} , SR^8 , $\text{SO}_2\text{N}(\text{R}^8)\text{R}^{8'}$, NHR^9 , NHCOR^9 , NR^9COR^9 , $\text{NHCO}(\text{OR}^9)$, $\text{NR}^9\text{CO}(\text{OR}^9)$, $\text{NR}^8\text{SO}_2\text{R}^{10'}$, $\text{NHSO}_2\text{N}(\text{R}^{10'})\text{R}^{10'}$, $\text{NR}^6\text{CON}(\text{R}^{10'})\text{R}^{10'}$, COR^9 , CO_2R^8 , $\text{CON}(\text{R}^8)\text{R}^{8'}$, wherein R^8 and $\text{R}^{8'}$ may be taken together to form a 3-7 membered carbocyclic ring having 1 to 3 substituted or unsubstituted heteroatoms selected from S, SO, SO_2 , O, N, and NR^6 , and wherein R^{10} and $\text{R}^{10'}$ may be taken together to form a 3-7 membered carbocyclic

20

25

ring having 1 to 3 substituted or unsubstituted heteroatoms selected from S, SO, SO₂, O, N, and NR⁶ wherein said aryl, heterocyclic, heteroaryl, or alkenyl are optionally substituted with R⁹;

5 R⁶ is independently selected from the group consisting of: hydrido, aryl, heteroaryl, lower alkyl, haloalkyl, alkenyl, alkynyl, hydroxyalkyl, aminoalkyl, alkylaminoalkyl, alkoxy, alkoxyalkyl, heterocyclicalkyl, and heterocyclic;

10 R⁸ is independently selected from the group consisting of: hydrido, aryl, heteroaryl, arylalkyl, heterocyclic, haloalkyl, arylalkylamino, alkylaminoalkyl, dialkylaminoalkyl, alkyl, alkenyl, alkynyl, heteroarylalkyl, and heterocyclicalkyl;

15 R^{8'} is independently selected from the group consisting of: hydrido, aryl, heteroaryl, arylalkyl, heterocyclic, haloalkyl, arylalkylamino, alkylaminoalkyl, dialkylaminoalkyl, alkyl, alkenyl, alkynyl, heteroarylalkyl, and heterocyclicalkyl;

20 R⁹ is independently selected from the group consisting of: hydrido, lower alkyl, aryl, heteroaryl, arylalkyl, heterocyclic, cycloalkyl, heterocyclicalkyl, haloalkyl, arylalkylamino, amino, aminoalkyl, aminoacyl, nitro, azido, and heteroarylalkyl, wherein alkyl, aryl, heteroaryl, aminoalkyl, or arylalkyl are optionally substituted with one or more radical selected from the group consisting of: alkylsulfonamide, sulfamyl, alkyl, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, aminoalkyl, alkylaminoalkyl, alkoxy, halogen, acyloxy, oxy, formyl, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, nitro, azido, benzyloxy, dialkylaminoacyl, thioalkyl, aminoacyloxy, thiocyanate, isothiocyanate, alkyldioxy, hydroxyalkyl, alkylamino, alkyloxycarbonyl, alkoxyalkyl, alkenylamino, alkynylamino, alkenyl, alkynyl, dialkylaminoalkyloxy, and heterocyclic

25

30

optionally substituted with alkyl, alkylamino, aminoalkyl, hydroxyalkyl, and alkylaminoalkyl;

5 **R¹⁰** is independently selected from the group consisting of: hydrido, lower alkyl, heteroaryl, heterocyclic, haloalkyl, arylalkylamino, heteroarylalkyl, aryl, and arylalkyl, wherein aryl, heteroaryl, heterocyclic, or arylalkyl are optionally substituted with one or more radical selected from alkyl, alkoxy, halogen, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, and heterocyclic;

10 **R^{10'}** is independently selected from the group consisting of: hydrido, lower alkyl, heteroaryl, heterocyclic, haloalkyl, arylalkylamino, heteroarylalkyl, aryl, and arylalkyl, wherein aryl, heteroaryl, heterocyclic, or arylalkyl are optionally substituted with one or more radical selected from alkyl, alkoxy, halogen, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, and heterocyclic;

15 **R¹³** is selected from the group consisting of: hydrido, alkyl, aryl, arylalkyl, heteroaryl, heterocyclicalkyl, and heteroarylalkyl, wherein aryl, alkyl, arylalkyl, heteroaryl, heterocyclicalkyl, or heteroarylalkyl are optionally substituted with one or more radicals selected from the group consisting of OR¹⁴, N(R¹⁴)R^{14'}, and glycols;

20 **R¹⁴** is independently selected from the group consisting of hydrido, and lower alkyl; and


25 **R^{14'}** is independently selected from the group consisting of hydrido, and lower alkyl;

30 **R¹⁵** is selected from the group consisting of: hydrido, halogen, alkyl, cycloalkyl, aryl, haloalkyl, heteroaryl, heterocyclic, alkylalkene, alkylalkyne, hydroxy, hydroxyalkyl, alkylhydroxy,

amino, aminoalkyl, alkylamino, alkylaminoalkyl,
alkylhydroxyalkyl, nitro, cyano, alkylthio, alkylsulfinyl,
alkylsulfonyl; wherein aryl or arylalkyl are optionally substituted
with one or more radical selected from alkyl, alkoxy, halo,
5
haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy,
hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy,
heterocyclic; and
R¹⁶ is selected from the group consisting of: hydrido, halogen,
alkyl, cycloalkyl, aryl, haloalkyl, heteroaryl, heterocyclic,
10
alkylalkene, alkylalkyne, hydroxy, hydroxyalkyl, alkylhydroxy,
amino, aminoalkyl, alkylamino, alkylaminoalkyl,
alkylhydroxyalkyl, nitro, cyano, alkylthio, alkylsulfinyl,
alkylsulfonyl; wherein aryl or arylalkyl are optionally substituted
with one or more radical selected from alkyl, alkoxy, halo,
15
haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy,
hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy,
heterocyclic; wherein R¹⁵ and R¹⁶ may be taken together to form a
5 to 7 membered saturated or unsaturated carbocyclic ring
optionally containing 0 to 3 heteroatoms selected from the group
20
consisting of N, O, or S, and wherein said ring is optionally
substituted with R¹;
or isomers, tautomers, carriers, esters, prodrugs,
pharmaceutically acceptable salts thereof.

25

15. The compound of claim 2 of the formula



wherein

A is $(CH_2)_m-CR^{15}=CR^{16}-(CH_2)_m$

m is independently selected from 0, 1, or 2;

5 R^9 is independently selected from the group consisting of:
hydrido, lower alkyl, aryl, heteroaryl, arylalkyl, heterocyclic,
cycloalkyl, heterocyclicalkyl, haloalkyl, arylalkylamino, amino,
aminoalkyl, aminoacyl, nitro, azido, and heteroarylalkyl, wherein
alkyl, aryl, heteroaryl, aminoalkyl, or arylalkyl are optionally
10 substituted with one or more radical selected from the group
consisting of: alkylsulfonamide, sulfamyl, alkyl, alkylthio,
alkylsulfinyl, alkylsulfonyl, alkylamino, aminoalkyl,
alkylaminoalkyl, alkoxy, halogen, acyloxy, oxy, formyl,
haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy,
15 hydroxyalkyloxy, phenoxy, nitro, azido, benzyloxy,
dialkylaminoacyl, thioalkyl, aminoacyloxy, thiocyanate,
isothiocyanate, alkyldioxy, hydroxyalkyl, alkylamino,
alkyloxycarbonyl, alkoxyalkyl, alkenylamino, alkynylamino,
20 alkenyl, alkynyl, dialkylaminoalkyloxy, and heterocyclic
optionally substituted with alkyl, alkylamino, aminoalkyl,
hydroxyalkyl, and alkylaminoalkyl;

R^{15} is selected from the group consisting of: hydrido, halogen, alkyl, cycloalkyl, aryl, haloalkyl, heteroaryl, heterocyclic, alkylalkene, alkylalkyne, hydroxy, hydroxyalkyl, alkylhydroxy, amino, aminoalkyl, alkylamino, alkylaminoalkyl, alkylhydroxyalkyl, nitro, cyano, alkylthio, alkylsulfinyl, alkylsulfonyl,; wherein aryl or arylalkyl are optionally substituted

with one or more radical selected from alkyl, alkoxy, halo, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, heterocyclic; and

5 **R¹⁶** is selected from the group consisting of: hydrido, halogen, alkyl, cycloalkyl, aryl, haloalkyl, heteroaryl, heterocyclic, alkylalkene, alkylalkyne, hydroxy, hydroxyalkyl, alkylhydroxy, amino, aminoalkyl, alkylamino, alkylaminoalkyl, alkylhydroxyalkyl, nitro, cyano, alkylthio, alkylsulfinyl, alkylsulfonyl,; wherein aryl or arylalkyl are optionally substituted with one or more radical selected from alkyl, alkoxy, halo, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, heterocyclic; wherein R¹⁵ and R¹⁶ may be taken together to form a

10 5 to 7 membered saturated or unsaturated carbocyclic ring optionally containing 0 to 3 heteroatoms selected from the group consisting of N, O, or S, and wherein said ring is optionally substituted with R¹;

15

20 or isomers, tautomers, carriers, esters, prodrugs, pharmaceutically acceptable salts thereof.

16. The compound of claim 15 selected from the group selected from the group consisting of:

25 1-(1,3-benzodioxol-5-yl)-8-[(2-chlorobenzoyl)amino]-1H-
benzo[g]indazole-3-carboxamide,
1-(1,3-benzodioxol-5-yl)-8-{[(2-chloropyridin-3-yl)carbonyl]amino}-
1H-benzo[g]indazole-3-carboxamide,
8-amino-1-(1,3-benzodioxol-5-yl)-1H-benzo[g]indazole-3-carboxamide,
30 1-(1,3-benzodioxol-5-yl)-5-bromo-8-{[(2-chloropyridin-3-
yl)carbonyl]amino}-1H-benzo[g]indazole-3-carboxamide, and

1-(1,3-benzodioxol-5-yl)-8-[(3-chloroisonicotinoyl)amino]-1H-benzo[g]indazole-3-carboxamide.

17. The compound of claim 2 of the formula

5



wherein

A is $(CH_2)_m-CR^{15}=CR^{16}-(CH_2)_m$

10 **m** is independently selected from 0, 1, or 2;

R⁴ is selected from the group consisting of: halogen, alkylsulfinyl, alkylsulfonyl, cyano, alkoxy carbonyl, alkyl, haloalkyl, hydrido, hydroxyalkyl, haloalkoxy, heterocyclic, nitro, acylamino, aryl, heteroaryl, and alkenyl, OR^{13} , SR^8 , $SO_2N(R^8)R^{8'}$, NHR^9 , $NHCOR^9$, NR^9COR^9 , $NHCO(OR^9)$, $NR^9CO(OR^9)$, $NR^8SO_2R^{10}$, $NHSO_2N(R^{10})R^{10'}$, $NR^6CON(R^{10})R^{10'}$, COR^9 , CO_2R^8 , $CON(R^8)R^{8'}$, wherein R^8 and $R^{8'}$ may be taken together to form a 3-7 membered carbocyclic ring having 1 to 3 substituted or unsubstituted heteroatoms selected from S, SO, SO₂, O, N, and NR^6 , and wherein R^{10} and $R^{10'}$ may be taken together to form a 3-7 membered carbocyclic ring having 1 to 3 substituted or unsubstituted heteroatoms selected from S, SO, SO₂, O, N, and NR^6 wherein said aryl, heterocyclic, heteroaryl, or alkenyl are optionally substituted with R^9 ;

20 **R**⁶ is independently selected from the group consisting of: hydrido, aryl, heteroaryl, lower alkyl, haloalkyl, alkenyl, alkynyl,

25

hydroxyalkyl, aminoalkyl, alkylaminoalkyl, alkoxy, alkoxyalkyl, heterocyclicalkyl, and heterocyclic;

5 **R⁸** is independently selected from the group consisting of: hydrido, aryl, heteroaryl, arylalkyl, heterocyclic, haloalkyl, arylalkylamino, alkylaminoalkyl, dialkylaminoalkyl, alkyl, alkenyl, alkynyl, heteroarylalkyl, and heterocyclicalkyl;

10 **R^{8'}** is independently selected from the group consisting of: hydrido, aryl, heteroaryl, arylalkyl, heterocyclic, haloalkyl, arylalkylamino, alkylaminoalkyl, dialkylaminoalkyl, alkyl, alkenyl, alkynyl, heteroarylalkyl, and heterocyclicalkyl;

15 **R⁹** is independently selected from the group consisting of: hydrido, lower alkyl, aryl, heteroaryl, arylalkyl, heterocyclic, cycloalkyl, heterocyclicalkyl, haloalkyl, arylalkylamino, amino, aminoalkyl, aminoacyl, nitro, azido, and heteroarylalkyl, wherein alkyl, aryl, heteroaryl, aminoalkyl, or arylalkyl are optionally substituted with one or more radical selected from the group consisting of: alkylsulfonamide, sulfamyl, alkyl, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, aminoalkyl, alkylaminoalkyl, alkoxy, halogen, acyloxy, oxy, formyl, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, nitro, azido, benzyloxy, dialkylaminoacyl, thioalkyl, aminoacyloxy, thiocyanate, isothiocyanate, alkyldioxy, hydroxyalkyl, alkylamino, alkyloxycarbonyl, alkoxyalkyl, alkenylamino, alkynylamino, alkenyl, alkynyl, dialkylaminoalkyloxy, and heterocyclic optionally substituted with alkyl, alkylamino, aminoalkyl, hydroxyalkyl, and alkylaminoalkyl;

20 **R¹⁰** is independently selected from the group consisting of: hydrido, lower alkyl, heteroaryl, heterocyclic, haloalkyl, arylalkylamino, heteroarylalkyl, aryl, and arylalkyl, wherein aryl, heteroaryl, heterocyclic, or arylalkyl are optionally substituted

with one or more radical selected from alkyl, alkoxy, halogen, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, and heterocyclic;

5 **R¹⁰** is independently selected from the group consisting of: hydrido, lower alkyl, heteroaryl, heterocyclic, haloalkyl, arylalkylamino, heteroarylalkyl, aryl, and arylalkyl, wherein aryl, heteroaryl, heterocyclic, or arylalkyl are optionally substituted with one or more radical selected from alkyl, alkoxy, halogen, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, and heterocyclic;

10 **R¹³** is selected from the group consisting of: hydrido, alkyl, aryl, arylalkyl, heteroaryl, heterocyclicalkyl, and heteroarylalkyl, wherein aryl, alkyl, arylalkyl, heteroaryl, heterocyclicalkyl, or heteroarylalkyl are optionally substituted with one or more radicals selected from the group consisting of OR¹⁴, N(R¹⁴)R¹⁴, and glycols;

15 **R¹⁴** is independently selected from the group consisting of hydrido, and lower alkyl; and

20 **R¹⁴** is independently selected from the group consisting of hydrido, and lower alkyl;

25 **R¹⁵** is selected from the group consisting of: hydrido, halogen, alkyl, cycloalkyl, aryl, haloalkyl, heteroaryl, heterocyclic, alkylalkene, alkylalkyne, hydroxy, hydroxyalkyl, alkylhydroxy, amino, aminoalkyl, alkylamino, alkylaminoalkyl, alkylhydroxyalkyl, nitro, cyano, alkylthio, alkylsulfinyl, alkylsulfonyl; wherein aryl or arylalkyl are optionally substituted with one or more radical selected from alkyl, alkoxy, halo, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy,

30

hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, heterocyclic; and

R¹⁶ is selected from the group consisting of: hydrido, halogen, alkyl, cycloalkyl, aryl, haloalkyl, heteroaryl, heterocyclic, alkylalkene, alkylalkyne, hydroxy, hydroxyalkyl, alkylhydroxy, amino, aminoalkyl, alkylamino, alkylaminoalkyl, alkylhydroxyalkyl, nitro, cyano, alkylthio, alkylsulfinyl, alkylsulfonyl; wherein aryl or arylalkyl are optionally substituted with one or more radical selected from alkyl, alkoxy, halo, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, heterocyclic; wherein R¹⁵ and R¹⁶ may be taken together to form a 5 to 7 membered saturated or unsaturated carbocyclic ring optionally containing 0 to 3 heteroatoms selected from the group consisting of N, O, or S, and wherein said ring is optionally substituted with R¹;

or isomers, tautomers, carriers, esters, prodrugs, pharmaceutically acceptable salts thereof.

20

18. The compound of claim 16 selected from the group consisting of:

8-[(3-chloroisonicotinoyl)amino]-1-(4-fluorophenyl)-1H-benzo[g]indazole-3-carboxamide,

25

benzo[g]indazole-3-carboxamide.

8-[(3-chloroisonicotinoyl)amino]-

benzo[g]indazole-3-carboxamide dihydrochloride,
8-amino-1-(4-fluorophenyl)-1H-benzo[g]indazole-3-carboxamide

8 [(2-chlorobenzoyl)amino]-1-(4-fluorophenyl)-1H-benzo[4,5]azepine

8-[(2-chlorobenzoyl)amino]-1-(4-fluorophenyl)-1H-benzo[g]indazole-3-carboxamide,

8-amino-1-(3-fluorophenyl)-1H-benzo[g]indazole-3-carboxamide,
8-[(2-chlorobenzoyl)amino]-1-(3-fluorophenyl)-1H-benzo[g]indazole-3-
carboxamide,
1-(4-fluorophenyl)-8-[(methylsulfonyl)amino]-1H-benzo[g]indazole-3-
carboxamide,
5 8-[(2-chloropyridin-3-yl)carbonyl]amino]-1-(4-fluorophenyl)-1H-
benzo[g]indazole-3-carboxamide,
8-[(2-chloropyridin-3-yl)carbonyl]amino]-1-(2,4-difluorophenyl)-1H-
benzo[g]indazole-3-carboxamide, and
10 1-(4-fluorophenyl)-8-[(4-methylpiperazin-1-yl)carbonyl]amino]-1H-
benzo[g]indazole-3-carboxamide.

15 19. A composition comprising the compound of claim 1, 2, 3, 4, 5, 6,
7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, or 18 and at least one
pharmaceutically acceptable carrier.

20 20. A method of treating cancer, inflammation or an inflammation
associated disorder in a subject, said method comprising administering
to the subject having or susceptible to such cancer, inflammation or
inflammation associated disorder, a therapeutically-effective amount of a
compound of claim 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12 13, 14, 15, 16, 17,
or 18.

25 21. The method of claim 20 for use in the treatment of cancer.

22. The method of claim 20 for use in the treatment of inflammation.

23. The method of claim 20 for use in the treatment of an
inflammation-associated disorder.

24. The method of claim 23 wherein the inflammation-associated disorder is arthritis.

5 25. The method of claim 23 wherein the inflammation-associated disorder is pain

26. The method of claim 23 wherein the inflammation-associated disorder is fever.