

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
20 May 2010 (20.05.2010)

PCT

(10) International Publication Number
WO 2010/056233 A1

(51) International Patent Classification:
A61K 8/30 (2006.01) *A61Q 19/10* (2006.01)

(21) International Application Number:
PCT/US2008/083087

(22) International Filing Date:
11 November 2008 (11.11.2008)

(25) Filing Language: English

(26) Publication Language: English

(71) Applicant (for all designated States except US): **COLGATE-PALMOLIVE COMPANY** [US/US]; 300 Park Avenue, New York, NY 10022 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **BOYKE, Christine** [US/US]; 378 Inwood Court, Somerset, NJ 08873 (US). **HOLERCA, Marian, N.** [RO/US]; 90 Rodney Avenue, Somerset, NJ 08873 (US).

(74) Agent: **MORGAN, Michael, F.**; Colgate- Palmolive Company, 909 River Road, Po Box 1343, Piscataway, NJ 08854 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))



WO 2010/056233 A1

(54) Title: COMPOSITION WITH A COLOR TO INDICATE COVERAGE

(57) Abstract: A composition comprising at least one indicator, at least one light stabilizer, and at least one antioxidant. The composition is useful as a cleanser that provides a usage timer based on color change to indicate to a person that sufficient time has elapsed for cleansing.

COMPOSITION WITH A COLOR TO INDICATE COVERAGE

BACKGROUND OF THE INVENTION

[0001] When applying compositions to a substrate, it may not be able to be determined if the desired area of the substrate has been covered by the composition. While a composition may be colored to assist in visually identifying coverage, it may not be desired that the color remains on the substrate.

SUMMARY OF THE INVENTION

[0002] A method comprising applying an indicator composition to a desired area of a substrate, wherein the applying is conducted until the desired area is covered as indicated by a color of the composition, and wherein the composition comprises:

- a. at least one indicator;
- b. at least one light stabilizer; and
- c. at least one antioxidant.

[0003] The present invention provides a color changing composition that remains stable over time on the shelf, but will change color during use.

DETAILED DESCRIPTION

[0004] As used throughout, ranges are used as a shorthand for describing each and every value that is within the range. Any value within the range can be selected as the terminus of the range. In addition, all references cited herein are hereby incorporated by reference in their entireties. In the event of a conflict in a definition in the present disclosure and that of a cited reference, the present disclosure controls.

[0005] Unless otherwise specified, all percentages and amounts expressed herein and elsewhere in the specification should be understood to refer to percentages by weight. The amounts given are based on the active weight of the material.

[0006] The method of the present invention is suitable for providing a color change during use of a composition to initially indicate to a person that a desired area of a substrate has been covered by the composition. After a period of time, the color of the composition can change to no longer be observable to a person. The substrate to be covered can be coated with the composition to any desired depth to indicate coverage of the substrate.

[0007] In certain embodiments, the substrate is skin or hair, and the composition is a personal care composition. The composition can be applied to skin (or hair). The composition will have a color associated with it. The coverage on the skin can be determined by observing the areas of the skin that have the colored composition. The composition can be applied until the

desired area is covered to assure that the benefit of the composition is obtained in the desired area.

[0008] Examples of compositions include, but are not limited to, sunscreens, lotions, creams, hand sanitizers, skin cleansers, body washes, shower gels, liquid hand soaps, bar soaps, shampoos, conditioners, antiperspirants, or deodorants. In other embodiments, the composition can be any desired application that can be applied to any desired type of substrate.

[0009] The timing of the color change can be adjusted to be observable by a person. By observable it is meant that the color change can be seen by a non-color blind person with an unaided eye at 20/20 or corrected to 20/20 with glasses or contact lenses at a distance of 30 cm from the composition under incandescent light, florescent light, or sunlight. In certain embodiments, the color change is observed by a person between 1 second to 30 minutes, before about 5 minutes, about 2 minutes, about 30 seconds, about 25 seconds, about 20 seconds, about 15 seconds, about 10 seconds, about 5 seconds, 5 to 30 seconds, or 10 to 30 seconds. The amount of time for color change can be adjusted by changing the concentration of the indicator and/or changing the starting pH. Time is also proportional to the concentration of the indicator. The more indicator that is present, the longer the time. Also, the further the pH is away from neutral pH (7), the longer the time to color change will be. For example, a composition can be made more basic, which would then take longer to color change as the pH is changed in the system to become more basic.

[0010] To provide the color change, an indicator is included. An indicator can be included in any amount to provide the color change. As more indicator is included in the composition, the color intensity increases to become more distinctly observable to a person. In one embodiment, the amount of indicator is 0.01 to 10% by weight. In one embodiment, the amount of indicator is 0.01 to 1% by weight. In one embodiment, the amount of indicator is about 0.5% by weight.

[0011] Examples of indicators that can be used include, but are not limited to, litmus, carmine, carminic acid, curry powder, thymol blue, pentamethoxy red, tropaeolin O, tropaeolin OO, tropaeolin OOO, 2, 4-dinitrophenol, methyl yellow, methyl orange, bromophenol blue, tetrabromophenol blue, alizarin sodium sulfonate, alpha-naphthyl red, para-ethoxychrysoidine, bromocresol green, methyl red, bromocresol purple, chlorophenol red, bromothymol blue, para-nitrophenol, azolitmin, phenol red, neutral red, rosolic acid, cresol red, naphtholphthalein, phenolphthalein, naphtholbenzein, thymolphthalein, Nile blue, alizarin yellow, salicyl yellow, diazo violet, nitramine, Poirrier's blue, trinitrobenzoic acid,

and extracts from beets, blackberries, blueberries, carrots, cherries, delphinium petals, geranium petals, grapes, grape seeds, horse chestnut leaves, morning glories, pansy petals, petunia petals, primrose, poppy petals, purple peonies, red radish, red cabbage, rhubarb, rose petals, strawberries, tea, turmeric, tulip petals, thyme, violet petals, and vanilla. Multiple indicators can be used in combination. In some embodiments, the indicator is selected from litmus and carminic acid. In one embodiment, the indicator is litmus. Litmus can be provided by lichen extract. Litmus changes blue under basic conditions and red under acidic conditions.

[0012] The composition can be designed to color change from a first pH to a higher pH or from a first pH to a lower pH. The upper and lower limits for pH are any pH that can be tolerated by skin. In one embodiment, the pH is 2 to 12. The pH of the composition can be adjusted to a desired starting pH by using any desired acid or base. In one embodiment, sodium hydroxide (NaOH) can be selected as the base and citric acid as the acid.

[0013] The pH change is provided by the substrate and/or the person applying the composition to a substrate. The pH of skin is about 5.5. When the composition is basic, the composition becomes more acidic during use. When the composition is acidic, the pH moves to become less acidic.

[0014] In one embodiment using litmus as the indicator, the pH can be adjusted to about 9.5 using sodium hydroxide. If the pH is too high, the composition can be adjusted back to about 9.5 using citric acid. As the composition is used, the blue litmus changes to red when contacted with skin.

[0015] Indicators may degrade upon exposure to light in solution. For a commercial product that will sit on store shelves and in locations where the product is used, the product will be exposed to light sources. To overcome the exposure to light, a light stabilizer is included to allow the product to remain stable over the targeted life of the product. Additionally, indicators can be oxidized in solution. For a product containing an indicator that contains materials or air that oxidize the indicator, an antioxidant is included to allow the product to remain stable over the targeted life of the product. Light degradation and oxidation is particularly noticeable in litmus and carminic acid.

[0016] A light stabilizer is included in the composition. A light stabilizer is any material that stabilizes indicators against degradation from exposure to light. In one embodiment, the light stabilizer is an ascorbate salt. In one embodiment, the light stabilizer is sodium ascorbate. In this embodiment, the sodium ascorbate is particularly useful for litmus and/or carminic acid.

[0017] The light stabilizer can be present in any amount that stabilizes the composition. In one embodiment, the amount is 0.01 to 1% by weight. In another embodiment, the amount is 0.05 to 0.5% by weight. In one embodiment, the amount is about 0.25% by weight.

[0018] An antioxidant is included in the composition. An antioxidant is any material that stabilizes an indicator against degradation from oxidation. In one embodiment, the light stabilizer is benzophenone-4. In this embodiment, the benzophenone-4 is particularly useful for litmus and/or carminic acid.

[0019] The antioxidant can be present in any amount that stabilizes the composition. In one embodiment, the antioxidant is present in an amount of 0.01 to 0.1% by weight. In one embodiment, the amount is about 0.05% by weight.

[0020] The light stabilizer and the antioxidant can both independently be selected from the CTFA (Cosmetic, Toiletries, and Fragrance Association) list of antioxidants. See the list that is available circa July 2008 at ctfa-online.org. Examples include, but are not limited to, Acacia Catechu Wood Extract, Acacia Victoriae Fruit Extract, Acanthopanax Senticosus (Eleuthero) Root Water, Acer Palmatum Leaf Extract, Acetamidocaproic Acid, Acetyl Benzoyloxy Prasterone, Acetyl Cysteine, Acetyl Hexapeptide-22, 2-Acetylhydroquinone, Achillea Millefolium Flower Extract, Adamantanylcarboxamido Hydroxylbenzamide, Adamantanylcarboxamido Methylhydroxylbenzamide, Adamantanyl Dihydrocaffeamide, Adamantanyl Dihydroxybenzamide, Adamantanyl Hydroxybenzamide, Adamantanyl Hydroxylterephthalamide, Adamantanyl Methylhydroxylterephthalamide, Adamantanyl Trihydroxybenzamide, Aesculus Hippocastanum (Horse Chestnut) Extract, Aframomum Angustifolium Leaf Oil, Agastache Rugosa Extract, Agrimonia Eupatoria Root Extract, Ajuga Reptans Cell Culture Extract, Ajuga Reptans Leaf Extract, Alchemilla Vulgaris Leaf Extract, Allium Cepa (Onion) Root Extract, Allium Fistulosum Root Extract, Allium Odorum Seed Extract, Alpinia Uraiensis Stalk/Leaf Water, Amaranthus Hypochondriacus Seed Extract, Aminoethanesulfinic Acid, Aminopropyl Ascorbyl Phosphate, Aminopropyl Methylenedioxyphenyl Phosphate, Aminopropyl Tocopheryl Phosphate, Angelica Furfijuga Flower/Leaf/Stem Extract, Angelica Keiskei Extract, Angoroside C, Anserine, Apigenin, Apium Graveolens (Celery) Seed Extract, Arabidopsis Thaliana Extract, Arbutin, Alpha-Arbutin, Arbutus Unedo Fruit Extract, Arctium Lappa Fruit Extract, Argon, Artemisia Maritima Extract, Asarum Heterotropoides Extract, Asarum Heterotropoides Rhizome Extract, Ascorbic Acid, Ascorbic Acid Polypeptide, Ascorbyl Dipalmitate, Ascorbyl Glucoside, Ascorbyl Linoleate, Ascorbyl Methylsilanol Pectinate, Ascorbyl Palmitate, Ascorbyl Phosphate Succinoyl Pentapeptide-12, Ascorbyl Stearate, Ascorbyl

Tetraisopalmitate, Ascorbyl Tocopheryl Maleate, Asiaticoside, Avena Sativa (Oat) Kernel Extract, Bacillus/Rice Bran Extract/Soybean Extract Ferment Filtrate, Bambusa Vulgaris Shoot Extract, Benzoguanamine, Beta Vulgaris (Beet) Root Powder, Betula Ermanii Stem Extract, BHA, BHT, Bis-demethoxycurcumin, Bis-Hydroxyethyl Tocopherylsuccinoylamido Hydroxypropane, Bis-(Tetramethyl Hydroxypiperidinyl) Sebacate, Boesenbergia Pandurata Rhizome Extract, Borago Officinalis Extract, Botrytis Ferment Extract Filtrate, Buddleja Axillaris Leaf Extract, Butylated Xylenol, t-Butylbenzamido Hydroxylbenzamide, t-Butylbenzamido Hydroxylphenylacetamide, t-Butylbenzamido Methylhydroxylbenzamide, t-Butylphenyl Imidazolylphenyl Sulfonamide, 4-Butylresorcinol, Caffeic Acid, Calcium Ascorbate, Calophyllum Inophyllum Seed Oil, Camellia Sinensis Catechins, Camellia Sinensis Leaf Extract, Camellia Sinensis Leaf Oil, Campsis Grandiflora Flower Extract, Capparis Moonii Fruit Extract, Capsella Bursa-Pastoris Sprout Water, Capsicum Annuum Fruit Extract, Carnosic Acid, Carotenoids, Carpinus Laxiflora Stem Extract, Carpinus Tschonoskii Stem Extract, Castanopsis Cuspidata Stem Extract, Cayaponia Tayuya Root Extract, Celosia Cristata Extract, Cercis Chinensis Flower/Leaf/Stem Extract, Cereus Grandiflorus (Cactus) Extract, Chaenomeles Sinensis Fruit Extract, Chitosan Ascorbate, Chitosan Glycolate, Chitosan Salicylate, Chlorogenic Acids, Chrysanthellum Indicum Flower Water, Chrysanthemum Boreale Flower Extract, Chrysanthemum Indicum Flower/Leaf/Stem Extract, Cimicifuga Dahurica Root Extract, Cimicifuga Racemosa Root Extract, Cinnamomum Zeylanicum Bark Extract, Citrus Junos Seed Extract, Citrus Medica Vulgaris Fruit Extract, Citrus Unshiu Fruit Powder, Cobalt DNA, Codonopsis Lanceolata Extract, Coffea Arabica Fruit Extract, Colloidal Platinum, Commelina Communis Leaf/Flower/Stem Extract, Copper Adenosine Triphosphate, Copper Pyridoxal 5-Phosphate, Coptis Chinensis Root Extract, Cordyceps Sinensis Extract, Cornus Controversa Leaf Extract, Crotonaldehyde, Cucumaria Frondosa Extract, Curcumin, Cyamopsis Tetragonoloba (Guar) Symbiosome Extract, Cyclopia Genistoides Flower/Leaf/Stem Extract, Cyclopia Genistoides Leaf Extract, Cyclopia Subternata Flower/Leaf/Stem Extract, Cynoches Cooperi (Orchid) Extract, Cymbidium Goeringii Extract, Cynara Cardunculus Seed Oil, Cysteine, Cysteine HCl, Davidsonia Pruriens Fruit Extract, Decapeptide-6, Decapeptide-7, Decursinol, Decyl Mercaptomethylimidazole, Demethoxycurcumin, Diacetylcurcumin, Diamylhydroquinone, Di-t-Butylhydroquinone, Dicetyl Thiodipropionate, Dictyophora Indusiata (Mushroom) Extract, Dictyopteris Membranacea Extract, Dicyclopentadiene/t-Butylcresol Copolymer, Digalloyl Trioleate, Dihydroquercetin, Dihydroxy Methylchromone, Diisooctyl Sebacate, Dilauryl Thiodipropionate, Dimethoxybenzamido Phenylhydroxylacetamide, Dimethoxy Di-

p-Cresol, Dimethylmethoxy Chromanol, Dimyristyl Thiodipropionate, Dioleoyl Tocopheryl Methylsilanol, Dioscorea Batatas Extract, Diosmine, Diospyros Kaki Calyx Extract, Diretinyl Ether, Disodium Ascorbyl Sulfate, Disodium Isostearyl Ascorbyl Phosphate, Disodium Rutinyl Disulfate, Disodium Salicylphosphate, Disodium Ubiquinone, Distearyl Thiodipropionate, Ditridecyl Thiodipropionate, Dodecyl Gallate, Dunaliella Bardawil Powder, Echinops Sphaerocephalus Seed Oil, Ellagic Acid, Empetrum Nigrum Flower/Fruit/Leaf Extract, Epigallocatechin Gallate, Epimedium Sagittatum Leaf/Stem Extract, Ergothioneine, Erigeron Canadensis Flower Extract, Eriobotrya Japonica Leaf Protoplasts, Eriocaulon Buergerianum Flower/Stem Extract, Erythorbic Acid, Ethylbisiminomethylguaiacol Manganese Chloride, Ethyl Ferulate, Ethylhexyl Ferulate, Ethylhexyl Gallate, Eucommia Ulmoides Bark Extract, Euphorbia Jolkini Extract, Euphorbia Supina Stem Extract, Excoecaria Bicolor Leaf Extract, Ferulic Acid, Feruloyl Soy Glycerides, Foeniculum Vulgare (Fennel) Seed Extract, Forsythia Suspensa Fruit Extract, Fragaria Ananassa (Strawberry) Seed Oil, Fragaria Vesca (Strawberry) Flower Extract, Fragilaria Pinnata Extract, Furfuryl Palmitate, Galla Rhois Gallnut Extract, Ganoderma Japonicum Mycelium Ferment Extract Filtrate, Genistein Glucoside, Ginkgo Leaf Terpenoids, Glucosylrutin, Glyceryl Chromonyl Ether, Glyceryl Diferulate, Glycine Max (Soybean) Symbiosome Extract, Glycine Soja (Soybean) Oil, Glycyrrhiza Glabra (Licorice) Root Extract, Glycyrrhiza Glabra (Licorice) Root Water, Grifola Frondosa Fruiting Body Extract, Grifola Frondosa (Maitake) Mycelium Ferment Filtrate Extract, Gynostemma Pentaphyllum Extract, Gynostemma Pentaphyllum Leaf Extract, Gynostemma Pentaphyllum Leaf/Stem Extract, Haematococcus Pluvialis Extract, Haematococcus Pluvialis Oil, Haematococcus Pluvialis Powder, Hedera Helix (Ivy) Extract, Hesperetin, Hesperetin Laurate, Hesperidin Methyl Chalcone, Hexyloxy Trimethylphenol, Honokiol, Human Oligopeptide-9 Hexapeptide-25, Hydrolyzed Anona Cherimolia Fruit Extract, Hydrolyzed Aspergillus/Ginseng Extract Ferment, Hydrolyzed Gardenia Florida Extract, Hydrolyzed Grape Skin, Hydrolyzed Lycium Barbarum Fruit Extract, Hydrolyzed Olive Fruit, Hydrolyzed Olive Fruit Extract, Hydrolyzed Proanthocyanidin, Hydrolyzed Rice Leaf Extract, Hydrolyzed Saccharomyces/Sodium Selenate Ferment Extract, Hydrolyzed Sophora Japonica Fruit Extract, Hydrolyzed Soy Extract, Hydrolyzed Typha Angustifolia Pollen/Root Extract, Hydrolyzed Wheat Bran, Hydroquinone, p-Hydroxyanisole, Hydroxydecyl Ubiquinone, Hydroxylamine HCl, Hydroxylamine Sulfate, Hydroxyphenyl Dihydroxybenzamide, Illicium Religiosum Branch/Leaf Extract, Inositol Hexaniacinate Hexaascorbate, Ipomoea Batatas Tuber Extract, Isooctyl Caprylate/Caprates, Isooctyl

Thioglycolate, Isoquercitrin, Juglans Regia (Walnut) Seedcoat Extract, Juniperus Communis Sprout Extract, Kaempferol, Kojic Acid, Kojyl Glucoside, Kojyl Methylene dioxy cinnamate, Kou-Cha Ekisu, Lactobacillus/Portulaca Oleracea Ferment Extract, Lactobacillus/Rice Bran/Saccharomyces/Camellia Sinensis Leaf Extract Ferment, Lactobacillus/Saccharomyces/Pichia Anomala/Camellia Sinensis Leaf Extract/Artemisia Princeps Leaf Extract/Honey Ferment Filtrate, Lactobacillus/Saccharomyces/Pichia Anomala/Camellia Sinensis Leaf Extract/Honey Ferment Extract Filtrate, Lactobacillus/Saccharomyces/Pichia Anomala/Camellia Sinensis Leaf Extract/Honey Ferment Filtrate, Lactobacillus/Wasabia Japonica Root Ferment Extract, Larix Sibirica Wood Extract, Lawsonia Inermis (Henna) Extract, Lens Culinaris (Lentil) Symbiosome Extract, Lespedeza Cuneata Extract, Ligusticum Striatum Root Extract, Lilium Candidum Flower Extract, Lindera Strychnifolia Leaf Extract, Linoleyl Gallate, Linseed Oil Ascorbate Esters, Liriodendron Tulipifera Leaf Water, Lotus Japonicus Symbiosome Extract, Lupinus Subcarnosus Symbiosome Extract, Lycium Chinense Fruit Extract, Lycopene, Maackia Fauriei Stem Extract, Madecassoside, Magnesium Ascorbate, Magnesium Ascorbate/PCA, Magnesium Ascorbylborate, Magnesium Ascorbyl Phosphate, Magnolia Grandiflora Bark Extract, Magnolol, Malachite Extract, Mallotus Japonicus Leaf Extract, Malpighia Punicifolia (Acerola) Fruit Extract, Malus Domestica Fruit Extract, Manganese Adenosine Triphosphate, Manganese Dioxide, Manganese Fructose Diphosphate, Manganese Pyridoxal 5-Phosphate, Matrine, Matteuccia Struthiopteris Extract, Medicago Sativa (Alfalfa) Symbiosome Extract, Melaleuca Alternifolia (Tea Tree) Leaf Oil, Melatonin, Melia Azadirachta Conditioned Media/Culture, Methoxy PEG-7 Ascorbic Acid, Methoxy-PEG-7 Rutinyl Succinate, Methoxytrimethylphenyl Dihydroxyphenyl Propanol, Methyl Di-t-butyl Hydroxyhydrocinnamate, Methylene Di-t-Butylcresol, 3,4-Methylene dioxybenzyl Methylsulfone, Methyl Methacrylate/Trimethoxysilylpropyl Methacrylate Crosspolymer, Methyl Myristic Acid, Methylsilanol Ascorbate, Mirabilis Jalapa Flower Extract, Momordica Grosvenori Fruit Extract, Monascus/Rice Ferment, Morus Alba Fruit Extract, Morus Alba Leaf Extract, Morus Alba Root Extract, Morus Bombycis Extract, Morus Bombycis Stem Extract, Morus Bombycis Wood Extract, Murdannia Keisak Extract, Murraya Exotica Leaf Extract, Musa Balbisiana Fruit Extract, Narcissus Tazetta Bulb Extract, Nelumbo Nucifera Seed Extract, Niacinamide Hydroxybenzoate, Nictoflorin, Nordihydroguaiaretic Acid, Octadecyl Di-t-butyl-4-hydroxyhydrocinnamate, Octanicotinoyl Epigallocatechin Gallate, Octapeptide-4, Olea Europaea (Olive) Bud Extract, Olea Europaea (Olive) Flower Water, Olea Europaea (Olive) Fruit Unsaponifiables, Oligopeptide-28, Oligopeptide-29,

Oligopeptide-30, Oligopeptide-31, Orostachys Japonica Extract, Oryza Sativa (Rice) Seed Water, Oxycoccus Palustris Seed Oil, Paecilomyces Japonica/Grape/Cucumber Juice Extract Ferment Filtrate, Paeonol, Palmatine, Palmitoyl Camellia Sinensis Extract, Palmitoyl Coffee Bean Extract, Palmitoyl Grape Seed Extract, Palmitoyl Grapevine Shoot Extract, Palmitoyl Olive Leaf Extract, Pearl Extract, PEG/PPG-2/5 Tocopheryl Ether, PEG/PPG-5/10 Tocopheryl Ether, PEG/PPG-5/20 Tocopheryl Ether, PEG/PPG-5/30 Tocopheryl Ether, PEG/PPG-30/10 Tocopheryl Ether, PEG/PPG-50/20 Tocopheryl Ether, PEG/PPG-70/30 Tocopheryl Ether, PEG/PPG-100/70 Tocopheryl Ether, Pelvetia Siliquosa Extract, Pentaerythrityl Tetra-di-t-butyl Hydroxyhydrocinnamate, Perilla Ocymoides Seed Extract, Perillyl Alcohol, Peucedanum Japonicum Branch Extract, Phaseolus Angularis Seed Extract, Phaseolus Vulgaris (Kidney Bean) Sprout Extract, Phellinus Linteus Extract, Phenethyl Caffate, Phenylethyl Resorcinol, Phenylthioglycolic Acid, Phloretin, Phloroglucinol, Physocarpus Amurensis Stem Extract, Phytosteryl Linoleate, Phytosteryl Linoleate/Linolenate, Piceatannol, Pikea Robusta Extract, Pimpinella Brachycarpa Extract, Pinus Densiflora Extract, Pinus Densiflora Leaf Extract, Pinus Nigra Bud/Needle Extract, Pinus Pinaster Bark/Bud Extract, Pinus Radiata Bark Extract, Pinus Sylvestris Bud Extract, Piperlonguminine, Piper Nigrum (Pepper) Seed, Piper Umbellatum Root Extract, Pisum Sativum Symbiosome Extract, Platycarya Strobilacea Branch Extract, Platycodon Grandiflorum Root Extract, Pogostemon Cablin Leaf/Stem Extract, Polygonum Cuspidatum Root Extract, Polyquaternium-82, Porphyridium Polysaccharide, Porphyridium Polysaccharide Hydroxypropyltrimonium Chloride, Portulaca Oleracea Flower/Leaf/Stem Extract, Potassium Ascorbyl Tocopheryl Phosphate, Potassium Sulfite, Pothomorphe Umbellata Leaf Extract, PPG-2 Tocophereth-5, PPG-5 Tocophereth-2, PPG-10 Tocophereth-30, PPG-20 Tocophereth-50, PPG-30 Tocophereth-70, PPG-70 Tocophereth-100, PPG-5 Tocopheryl Ether, Propyl Gallate, Protocatechuicaldehyde, Prunella Vulgaris Extract, Prunus Cerasus (Bitter Cherry) Extract, Prunus Cerasus (Bitter Cherry) Fruit Extract, Prunus Cerasus (Bitter Cherry) Leaf Extract, Prunus Mume Flower, Prunus Mume Flower Extract, Prunus Mume Flower Powder, Prunus Salicina Fruit Extract, Prunus Spinosa Wood Extract, Prunus Tomentosa Fruit Extract, Pueraria Lobata Symbiosome Extract, Puerarin, Punica Granatum Bark/Fruit Extract, Pyridoxine Hydroxybenzoate, Pyridoxine Hydroxycitrate, Pyridoxine Salicylate, Pyridoxine Serinate, Pyridyloxide t-Butylnitron, Pyrus Malus (Apple) Peel Extract, Quercetin, Quercetin Caprylate, Quercus Infectoria Fruit Extract, Quercus Salicina Stem Extract, Resacetophenone, Resorcinol, Resveratrol, Retinyl Formyl Aspartamate, Rhodochrosite Extract, Rhododendron Mucronulatum Flower Extract, Rhodymenia Palmata

Extract, Rhus Javanica Extract, Rhus Semialata Leaf Extract, Rosa Davurica Bud Extract, Rosa Indica Flower Extract, Rosa Wichuraiana Stem Extract, Rosmarinic Acid, Rosmarinus Officinalis (Rosemary) Flower Extract, Rosmarinus Officinalis (Rosemary) Leaf Extract, Rubus Arcticus Fruit Extract, Rubus Chamaemorus Fruit Extract, Rutin, Rutinyl Succinate, Ryoku-Cha Ekisu, Saccharomyces/Carrageenan Extract/Sarcoditheca Gaudichaudii Extract Ferment, Saccharomyces/Pueraria Lobata Symbiosome Extract Ferment, Saccharomyces/Rhodobacter/Lactobacillus/Leuconostoc/Streptomyces Griseus/Aspergillus/Bacillus Ferment Filtrate, Salnacedin, Sambucus Nigra Bud Extract, Sanguisorba Minor Extract, Santalum Acuminatum Fruit Extract, Sarcoditheca Gaudichaudii Extract, Sargassum Pallidum Extract, Sarothamnus Scoparius Extract, Sasa Senanensis Leaf Extract, Saururus Chinensis Flower Extract, Saururus Chinensis Leaf/Root Extract, Schinopsis Quebracho-Colorado Wood Extract, Schizandra Sphenanthera Fruit Extract, Sedum Rosea Root Extract, Selenium/Glycine Soja (Soybean) Sprout Extract, Sideritis Syriaca Extract, Smilax China Extract, Smilax China Fruit Extract, Smilax China Root Extract, Smithsonite Extract, Sodium Ascorbate, Sodium Ascorbyl/Cholesteryl Phosphate, Sodium Ascorbyl Phosphate, Sodium Bisulfite, Sodium Erythorbate, Sodium Grape Seed Extract Phosphate, Sodium Metabisulfite, Sodium Phosphono-Pyridoxylidenerhodanine, Sodium Sulfite, Sodium Thioglycolate, Sodium Tocopheryl Phosphate, Sodium Zinc Histidine Dithiooctanamide, Solanum Lycopersicum (Tomato) Fruit Extract, Sophora Flavescens Root Extract, Sorbityl Furfural, Spirulina Maxima Powder, Sponge Extract, Stearyl Gallate, Stephanandra Incisa Stem Extract, Streptococcus Thermophilus/Lactobacillus/Bifidobacterium/Licorice Root Extract Ferment Filtrate, Succinoyl Ascorbate Pentapeptide-6, Syzygium Leuhmanii Fruit Extract, Tangeritin, TBHQ, Terminalia Ferdinandiana Fruit Extract, Tetrabutyl Ethylidinebisphenol, Tetrahexyldecyl Ascorbate, Tetrahydrobisdemethoxydiferuloylmethane, Tetrahydrocurcumin Diacetate, Tetrahydrodemethoxydiferuloylmethane, Tetrahydrodiferuloylmethane, Tetramethylbutyl Dihydroxybenzamide, Tetramethylchromanol Glucoside, Teupolioside, Theobroma Cacao (Cocoa) Husk Extract, Theobroma Cacao (Cocoa) Seed Extract, Thioctic Acid, Thiodiglycol, Thiodiglycolamide, Thiodiglycolic Acid, Thioglycolic Acid, Thiolactic Acid, Thiosalicylic Acid, Thiotaurine, Thuja Orientalis Extract, Thuja Orientalis Leaf Extract, Thymol Trimethoxycinnamate, Tococysteamide, Tocophereth-5, Tocophereth-10, Tocophereth-12, Tocophereth-18, Tocophereth-50, Tocopherol, Tocophersolan, Tocopheryl Acetate, Tocopheryl Dimethylglycinate, Tocopheryl Dimethylglycinate HCl, Tocopheryl Linoleate, Tocopheryl Linoleate/Oleate, Tocopheryl Nicotinate, Tocopheryl Retinoate, Tocopheryl

Succinate, Tocoquinone, Toluene, o-Tolyl Biguanide, Torilis Japonica Extract, Totarol, Trapa Bicornis Nut Extract, Tremella Fuciformis Sporocarp Extract, Tricholoma Magnivelare Extract, Trifolium Pratense (Clover) Seed Extract, Trifolium Pratense (Clover) Symbiosome Extract, Trigonella Foenum Symbiosome Extract, Tripropylene Glycol, Tris-BHT Mesitylene, Tris(Nonylphenyl)Phosphite, Trisodium Ascorbyl Isopalmitate Phosphate, Trisodium Ascorbyl Palmitate Phosphate, Trisodium Fructose Diphosphate, Trisodium Resveratrol Triphosphate, Triticum Aestivum (Wheat) Leaf Extract, Tyrosyl Histidine HCl, Ubiquinol, Ubiquinone, Ulva Lactuca Powder, Uron-Cha Ekisu, Vaccinium Macrocarpon (Cranberry) Fruit Powder, Vaccinium Myrtillus Bud Extract, Vaccinium Myrtillus Stem Extract, Vaccinium Vitis-Idaea Fruit Extract, Vaccinium Vitis-Idaea Leaf Protoplasts, Vaccinium Vitis-Idaea Seed Oil, Vanda Coerulea Extract, Viburnum Awabuki Leaf Extract, Vicia Sativa Symbiosome Extract, Viola Mandshurica Flower Extract, Vitis Vinifera (Grape) Fruit Powder, Vitis Vinifera (Grape) Juice, Vitis Vinifera (Grape) Juice Extract, Vitis Vinifera (Grape) Seed Extract, Vitis Vinifera (Grape) Skin Extract, Vitis Vinifera (Grape) Skin Powder, Volcanic Rock, Wine Extract, Xylyl Dibutylbenzofuranone, Zinc Adenosine Triphosphate, Zinc Ascorbate, Zinc Dibutyldithiocarbamate, Zinc Fructose Diphosphate, and Zinc Pyridoxal 5-Phosphate. In one embodiment, the light stabilizer and the antioxidant are different materials.

[0021] In one embodiment, a combination of sodium ascorbate and benzophenone-4 are selected as the light stabilizer and antioxidant, respectively. This combination provides stability to light and oxidation. In one embodiment, the amount of sodium ascorbate is about 0.25% by weight, and the amount of benzophenone-4 is about 0.05% by weight.

[0022] Stability is measured up to 13 weeks for compositions placed in sunlight for light stability and for oxidation at 25°C and 49°C. Light stability is measured by color fading using UV-Vis spectroscopy. Oxidation is measured by color fading using UV-Vis spectroscopy and with pH measurements. This test can be conducted by taking 1g of sample and diluting with 9g of water to make a dilute solution. The dilute solution is placed into a UV-Vis spectrometer. The UV-Vis spectrometer settings are: wavelength set to scan 400 nm to 800 nm, scan speed is set to fast, cycle time is 60 seconds, and results are recorded in absorbance units with a lower limit of 0.0 and an upper limit of 2.0.

[0023] In one embodiment, the composition has a second light absorption value after 13 weeks at 49°C that is no more than 20% different from a first light absorption value that is measured initially according to a UV-Vis spectrometer test. The wavelength that is selected for the measurement is based on the peak absorption for the indicator that is used. For litmus,

the absorption is measured at about 590 nm. When there is more than one indicator used, the absorption measurements are taken at each of the peak absorptions for each material. In other embodiments, the value is not more than 15%, 10%, or 5% different. In one embodiment, the wavelength used for measuring is 590 nm. In one embodiment, the pH does not change more than +/- 0.5 units.

[0024] When the composition is a cleanser, the composition contains at least one surfactant. The surfactant may be one or more anionic surfactants, amphoteric surfactants, nonionic surfactants, cationic surfactants, and combinations thereof. Those of ordinary skill in the art will be aware of suitable surfactants and other additives readily identifiable from the *International Cosmetic Ingredient Dictionary and Handbook*, 10th ed., (2004). Surfactants can be included in any desired amount. In one embodiment, surfactants are present in the composition in an amount of 0 to 40% by weight. In one embodiment, the surfactants are present in an amount of 1 to 40% by weight. In one embodiment, surfactants are present in the composition in an amount of 5 to 40% by weight. In one embodiment, the surfactants are present in an amount of 1 to 10% by weight.

[0025] A variety of anionic surfactants can be utilized in the moisturizing body wash composition including, for example, long chain alkyl (C₆-C₂₂) materials such as long chain alkyl sulfates, long chain alkyl sulfonates, long chain alkyl phosphates, long chain alkyl ether sulfates, long chain alkyl alpha olefin sulfonates, long chain alkyl taurates, long chain alkyl isethionates (SCI), long chain alkyl glyceryl ether sulfonates (AGES), sulfosuccinates and the like. These anionic surfactants can be alkoxyated, for example, ethoxyated, although alkoxylation is not required. These surfactants are typically highly water soluble as their sodium, potassium, alkyl and ammonium or alkanol ammonium containing salt form and can provide high foaming cleansing power. Other equivalent anionic surfactants may be used. In one embodiment, the anionic surfactant comprises sodium laureth sulfate, sodium pareth sulfate, and combinations thereof. Anionic surfactants can be included in any desired amount. In one embodiment, anionic surfactants are present in the composition in an amount of 0 to 15% by weight. In one embodiment, anionic surfactants are present in an amount of 4 to 12 % by weight.

[0026] Amphoteric surfactants may also be included in the composition. These surfactants are typically characterized by a combination of high surfactant activity, lather forming and mildness. Amphoteric surfactants include, but are not limited to, derivatives of aliphatic secondary and tertiary amines in which the aliphatic radical can be straight chain or branched and wherein one of the aliphatic substituents contains 8 to 18 carbon atoms and one contains

an anionic water solubilizing group, *e.g.*, carboxy, sulfonate, sulfate, phosphate, or phosphonate. Examples of such compounds include sodium 3-dodecylaminopropionate, sodium 3-dodecylaminopropane sulfonate, N-alkyl taurines and N-higher alkyl aspartic acids. Other equivalent amphoteric surfactants may be used. Examples of amphoteric surfactants include, but are not limited to, a range of betaines including, for example, high alkyl betaines, such as coco dimethyl carboxymethyl betaine, lauryl dimethyl carboxy-methyl betaine, lauryl dimethyl alpha-carboxyethyl betaine, cetyl dimethyl carboxymethyl betaine, lauryl bis-(2-hydroxyethyl)carboxy methyl betaine, stearyl bis-(2-hydroxypropyl)carboxymethyl betaine, oleyl dimethyl gamma-carboxypropyl betaine, and lauryl bis-(2-hydroxypropyl)alpha-carboxyethyl betaine, sulfobetaines such as coco dimethyl sulfopropyl betaine, stearyl dimethyl sulfopropyl betaine, amido betaines, amidosulfobetaines and the like. Betaines having a long chain alkyl group, particularly coco, may be particularly useful as are those that include an amido groups such as the cocamidopropyl and cocoamidoethyl betaines. Amphoteric surfactants can be included in any desired amount. In one embodiment, amphoteric surfactants are present in the composition in an amount of 0 to 15% by weight. In one embodiment, the amphoteric surfactants are present in the composition in an amount of 1 to 6% by weight.

[0027] Examples of nonionic surfactants include, but are not limited to, polysorbate 20, long chain alkyl glucosides having C₈-C₂₂ alkyl groups; coconut fatty acid monoethanolamides such as cocamide MEA; coconut fatty acid diethanolamides, fatty alcohol ethoxylates (alkylpolyethylene glycols); alkylphenol polyethylene glycols; alkyl mercaptan polyethylene glycols; fatty amine ethoxylates (alkylaminopolyethylene glycols); fatty acid ethoxylates (acylpolyethylene glycols); polypropylene glycol ethoxylates (for example the PLURONIC™ block copolymers commercially available from BASF); fatty acid alkylolamides, (fatty acid amide polyethylene glycols); N-alkyl-, N-alkoxypolyhydroxy fatty acid amides; sucrose esters; sorbitol esters; polyglycol ethers; and combinations thereof. Nonionic surfactants can be included in any desired amount. In one embodiment, nonionic surfactants are present in the composition in an amount of 0 to 3% by weight. In one embodiment, nonionic surfactants are present in the composition in an amount of 0.5 to 1.5% by weight.

[0028] Cationic surfactants can also be included in the composition. Examples of cationic surfactants include, but are not limited to any quaternium or polyquaternium compound. Cationic surfactants can be included at any desired level. In one embodiment, cationic surfactants are present in the composition in an amount of 0 to 2 % by weight. In one

embodiment, cationic surfactants are present in the composition in an amount of 0.1 to 0.3 % by weight.

[0029] Many additional surfactants are described in McCutcheon's Detergents and Emulsifiers (1989) and other reference materials that are well known to those of ordinary skill in the art.

[0030] In one embodiment, the surfactant is a combination of an anionic surfactant and an amphoteric surfactant. In one embodiment, the surfactant comprises a C10-C16 alcohol ethoxylate and cocoamidopropyl betaine. In one embodiment, the alcohol ethoxylate is sodium C10-C16 alcohol ethoxylate.

[0031] When the composition is a sanitizer, the composition can contain an alcohol, such as ethanol. Sanitizers can be thickened with any material to provide a desired viscosity to the sanitizer composition. Examples of thickening agents include, but are not limited to, synthetic thickening agents, polymeric gums, polysaccharides, pectin, alginate, arabinogalactan, carrageen, gellan gum, xanthum gum, guar gum, rhamsan gum, furcellaran gum, and other natural gum. A synthetic thickening agent in one embodiment is a polyacrylate. One acrylate aqueous solution is manufactured by Lubrizol as CARBOPOL™ resins, also known as CARBOMER™, which are hydrophilic high molecular weight, crosslinked acrylic acid polymers. In one embodiment, the polymer is CARBOPOL™ Aqua SF-1. Other polymers that can be used include, but are not limited to, CARBOPOL™ Aqua 30, CARBOPOL™ 940 with a molecular weight of approximately 4,000,000, and CARBOPOL™ 934 with a molecular weight of approximately 3,000,000. The thickening agents can be used alone or in combination. The amount of thickening agent can be any amount that provides for a desired level of thickening. In one embodiment, the thickening agent is present in an amount of 0.01 to 15% by weight of the composition. In other embodiments, the amount of thickening agent is 1% to 10%.

[0032] The composition can also contain glycerin. The glycerin can be included in any desired amount. In one embodiment, the amount of glycerin is up to 5% by weight. Glycerin can change the feel of skin after washing. At higher pH, skin can feel tight after washing.

[0033] In other embodiments, the composition can contain particulate materials. An example of a particulate material is shea butter beads, such as Butyrospermum Parkii beads from ISP Corp. available as CAPTIVATES™ 2485.

[0034] The composition can additionally contain a colorant to adjust the color of the composition to any desired color. The colorant can work in combination with the indicator to

provide color to the composition. Even with the colorant present, the indicator still works to change the color of the composition when the pH changes.

[0035] Examples of colorants include, but are not limited to, annatto, caramel, carmine, β -carotene, bismuth citrate, disodium EDTA-copper, potassium sodium copper chlorophyllin (chlorophyllin copper-complex), dihydroxyacetone, bismuth oxychloride, guaiazulene, henna, iron oxides, ferric ammonium ferrocyanide, ferric ferrocyanide, chromium hydroxide green, chromium oxide greens, guanine, lead acetate, pyrophyllite, mica, silver, titanium dioxide, aluminum powder, bronze powder, copper powder, ultramarines, manganese violet, zinc oxide, luminescent zinc sulfide, D&C Black No. 2, D&C Black No. 3, FD&C Blue No. 1, D&C Blue No. 4, D&C Brown No. 1, FD&C Green No. 3, D&C Green No. 5, D&C Green No. 6, D&C Green No. 8, D&C Orange No. 4, D&C Orange No. 5, D&C Orange No. 10, D&C Orange No. 11, FD&C Red No. 4, D&C Red No. 6, D&C Red No. 7, D&C Red No. 17, D&C Red No. 21, D&C Red No. 22, D&C Red No. 27, D&C Red No. 28, D&C Red No. 30, D&C Red No. 31, D&C Red No. 33, D&C Red No. 34, D&C Red No. 36, FD&C Red No. 40, D&C Violet No. 2, Ext. D&C Violet No. 2, FD&C Yellow No. 5, FD&C Yellow No. 6, D&C Yellow No. 7, Ext. D&C Yellow No. 7, D&C Yellow No. 8, D&C Yellow No. 10, and D&C Yellow No. 11. In one embodiment, FD&C Blue #1 is used as a second colorant in compositions containing litmus or carminic acid as the indicator. In this embodiment, the Blue #1 is present at about 0.003 weight %.

[0036] The composition can be formulated to have any desired viscosity. In some embodiments, the composition has Brookfield viscosity of greater than 0 to 35,000 mPas (cps). In one embodiment, the viscosity is less than 14,000 mPas (cps) so that it is pumpable. In other embodiments, the viscosity is greater than 2500 to 14,000 mPas (cps). Brookfield viscosity is measured on a DVII viscometer with spindle 5 at 20 rpm for 1 minute.

[0037] In some embodiments, the composition is formulated as a foaming hand cleanser that has a low viscosity. The viscosity range for foaming liquid hand soap is 25 ± 15 mPas (cps). Brookfield viscosity is measured on a DVII viscometer with spindle 3 at 100 rpm for 1 minute.

WHAT IS CLAIMED IS:

1. A method comprising applying an indicator composition to a desired area of a substrate, wherein the applying is conducted until the desired area is covered as detected by a color of the composition covering the desired area, and wherein the composition comprises:
 - a. at least one indicator;
 - b. at least one light stabilizer; and
 - c. at least one antioxidant.
2. The method of claim 1, wherein the composition has a second light absorption value after 13 weeks at 49°C that is no more than 20% different from a first light absorption value that is measured initially according to a UV-Vis spectrometer test for a peak absorption for the indicator.
3. The method of claim 1, wherein the indicator is present in an amount of 0.01 to 10% by weight in the composition.
4. The method of claim 1, wherein the light stabilizer is present in an amount of 0.01 to 1% by weight in the composition.
5. The method of claim 1, wherein the antioxidant is present in an amount of 0.01 to 0.1% by weight in the composition.
6. The method of claim 1, wherein the indicator is at least one indicator selected from litmus, carmine, carminic acid, curry powder, thymol blue, pentamethoxy red, tropaeolin O, tropaeolin OO, tropaeolin OOO, 2, 4-dinitrophenol, methyl yellow, methyl orange, bromophenol blue, tetrabromophenol blue, alizarin sodium sulfonate, alpha-naphthyl red, para-ethoxychrysoidine, bromocresol green, methyl red, bromocresol purple, chlorophenol red, bromothymol blue, para-nitrophenol, azolitmin, phenol red, neutral red, rosolic acid, cresol red, naphtholphthalein, phenolphthalein, naphtholbenzein, thymolphthalein, Nile blue, alizarin yellow, salicyl yellow, diazo violet, nitramine, Poirrier's blue, trinitrobenzoic acid, extracts from: beets, blackberries, blueberries, carrots, cherries, delphinium petals, geranium petals, grapes, grape seeds, horse chestnut leaves, morning glories, pansy petals, petunia petals, primrose, poppy petals, purple peonies, red radish, red cabbage, rhubarb, rose petals, strawberries, tea, turmeric, tulip petals, thyme, violet petals, and vanilla.
7. The method of claim 1, wherein the indicator is at least one indicator selected from litmus and carminic acid.

8. The method of claim 1, wherein the light stabilizer comprises sodium ascorbate and the antioxidant comprises benzophenone-4.
9. The method of claim 1, wherein the composition further comprises a surfactant.
10. The method of claim 1, wherein the composition comprises:
 - a. 0.01 to 1% by weight litmus,
 - b. 0.05 to 1% by weight sodium ascorbate, and
 - c. 0.05 to 0.1% by weight benzophenone-4.
11. The method of claim 1, wherein the composition comprises:
 - a. about 0.5 weight % litmus,
 - b. about 0.25 weight % sodium ascorbate, and
 - c. about 0.05 weight % benzophenone-4.
12. The method of claim 1, wherein the composition further comprises glycerin.
13. The method of claim 1, wherein the composition comprises:
 - a. 0.01 to 1% by weight litmus,
 - b. 0.05 to 1% by weight sodium ascorbate,
 - c. 0.05 to 0.1% by weight benzophenone-4;
 - d. glycerin in an amount up to 5% by weight; and
 - e. a surfactant.
14. The method of claim 1, wherein a color change is observed before about 2 minutes.
15. The method of claim 1, wherein a color change is observed at a time between 1 second to 30 minutes.

A. CLASSIFICATION OF SUBJECT MATTER*A61K 8/30(2006.01)i, A61Q 19/10(2006.01)i*

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC8: A61K8, A61K7, A61K49, A61Q, A61L, C11D3

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

eKOMPASS(KIPO internal), FPD, PAJ, PUBMED

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 2005/0049157 A1 (KIMBERLY-CLARK WORLDWIDE, INC.) 03 March 2005 See claims and paragraph [0055].	1-15
A	US 2008/0081020 A1 (Yeong H. Huang et al.) 03 April 2008 See claims.	1-15
A	JP 2002226898 A (NISSHIN KAGAKU KK et al.) 14 August 2002 See whole document.	1-15
A	JP 06285140 A (OGURA NOBUO) 11 October 1994 See whole document.	1-15

 Further documents are listed in the continuation of Box C. See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

10 AUGUST 2009 (10.08.2009)

Date of mailing of the international search report

11 AUGUST 2009 (11.08.2009)

Name and mailing address of the ISA/KR

Korean Intellectual Property Office
Government Complex-Daejeon, 139 Seonsa-ro, Seo-gu,
Daejeon 302-701, Republic of Korea

Facsimile No. 82-42-472-7140

Authorized officer

SONG, Keon Hyoung

Telephone No. 82-42-481-5607



INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/US2008/083087

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 2005/0049157 A1	03.03.2005	WO 05023972 A1 MX 6002287 A KR 6125691 A JP 2007504294 T2 EP 1697490 A1 DE 602004012892 C0	17.03.2005 17.05.2006 06.12.2006 01.03.2007 06.09.2006 15.05.2008
US 2008/0081020 A1	03.04.2008	WO 2008060355 A2 CA 2665313 AA AU 7320012 AA	22.05.2008 22.05.2008 22.05.2008
JP 2002226898 A	14.08.2002	WO 0042163 A1 AU 2957999 A1	20.07.2000 01.08.2000
JP 06285140 A	11.10.1994	None	