



US00PP13592P3

(12) **United States Plant Patent**
Olesen et al.

(10) **Patent No.:** **US PP13,592 P3**

(45) **Date of Patent:** **Feb. 25, 2003**

(54) **MINIATURE ROSE VARIETY 'POULGO004'**

(52) **U.S. Cl.** **Plt./121**

(76) Inventors: **L. Pernille Olesen**, Hillerødvejen 49, DK-3480 Fredensborg (DK); **Mogens N. Olesen**, Hillerødvejen 49, DK-3480 Fredensborg (DK)

(58) **Field of Search** **Plt./118, 121, 117, Plt./116, 119, 122**

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 21 days.

Primary Examiner—Bruce R. Campell
Assistant Examiner—Annette H. Para

(21) Appl. No.: **09/776,154**

(22) Filed: **Feb. 2, 2001**

(65) **Prior Publication Data**

US 2002/0148011 P1 Oct. 10, 2002

(57) **ABSTRACT**

A new miniature rose plant which has abundant, light red flowers and attractive foliage. This new and distinct variety has shown to be uniform and stable in the resulting generations from asexual propagation.

(51) **Int. Cl.**⁷ **A01H 5/00**

1 Drawing Sheet

1

2

SUMMARY OF THE INVENTION

Classification:

Botanical.—*Rosa hybrida* 'POULgo004'.

Commercial.—Miniature.

The present invention constitutes a new and distinct variety of miniature rose plant which originated from a controlled crossing between an unnamed, unpatented seedling and 'POULrijk', described and illustrated under U.S. Plant patent application Ser. No. 09/261,445, dated Mar. 3, 1999. The two parents were crossed during the summer of 1994 and the resulting seeds were planted in a controlled environment in Fredensborg, Denmark. The new variety is named 'POULgo004'.

The new rose may be distinguished from its seed parent, an unnamed seedling, by the following combination of characteristics:

1. The seed parent exhibits pink blooms, where 'POULgo004' exhibits red blooms.

The new variety may be distinguished from its pollen parent, 'POULrijk' by the following combination of characteristics:

1. 'POULrijk' exhibits pink blooms, where those of 'POULgo004' are red.

2. 'POULrijk' is a shrub rose with a spreading, low-growing habit, while 'POULgo004' is more procumbent and suitable for use in hanging baskets.

The objective of the hybridization of this rose variety was to create a new and distinct variety for use in hanging baskets with unique qualities, such as:

1. Uniform and abundant flowers;
2. Vigorous, but compact growth when propagated by cuttings;
3. Disease resistance;
4. A miniature rose suitable for use in hanging basket containers in the nursery industry.

This combination of qualities is not present in previously available commercial cultivars of this type and distinguish 'POULgo004' from all other varieties of which we are aware.

As part of their rose development program, L. Pernille Olesen and Mogens N. Olesen germinated the seeds from

the aforementioned hybridization during winter 1994 and conducted evaluations on the resulting seedlings in a controlled environment in Fredensborg, Denmark.

'POULgo004' was selected in the spring 1995 by the inventors as a single plant from the progeny of the aforementioned hybridization.

Asexual reproduction of 'POULgo004' by traditional budding and rooted cuttings was first done by L. Pernille and Mogens N. Olesen in their nursery in Fredensborg, Denmark in July 1995. This initial and other subsequent asexual propagations conducted in controlled environments have demonstrated that the characteristics of 'POULgo004' are true to type and are transmitted from one generation to the next.

BRIEF DESCRIPTION OF THE DRAWING

The accompanying color illustration shows as true as is reasonably possible to obtain in color photographs of this type the variety rose 'POULgo004'.

Specifically illustrated in SHEET 1 is a plant of 'POULgo004' growing in a 20-cm hanging basket.

DETAILED DESCRIPTION OF THE VARIETY

The following is a description of 'POULgo004', as observed in its growth in Jackson County, Oreg., on plants aged 18 months. Color references are made using The Royal Horticultural Society (London, England) Colour Chart, 1995, except where common terms of color are used.

For a comparison, several physical characteristics of the rose variety 'POULrijk', a rose variety from the same inventors described and illustrated in U.S. Plant patent application Ser. No. 09/261,445, filed Mar. 3, 1999 are compared to 'POULgo004' in Chart 1.

CHART 1

	'POULgo004'	'POULrijk'
Color of open flower, outer side, middle zone.	Red Group 53D to Red-Purple Group 57A	Red-Purple Group 62B.

CHART 1-continued

	'POULgo004'	'POULrijk'
Petalage	Single	Single
Upon opening, color basal petal spot, outer side	Red Group 53C	Green-White Group 157D.

Parents:

Seed parent.—An unnamed seedling.

Pollen parent.—'POULrijk'.

FLOWER AND FLOWER BUD

Blooming habit: Continuous.

Flower bud:

Size.—18 mm in length from base of receptacle to end of bud.

Bud form.—Long, pointed ovoid.

Bud color.—As sepals unfold, Red Group 53D and Red Group 57A.

Sepals.—Shape: Sepal apex is cirrose; base is flat at union with peduncle. Length: 15 to 30 mm. Width: 10 mm. Color: Upper surface is Yellow-Green Group 144 B. Lower surface is Yellow-Green Group 144 B. Appendages: Weak foliaceous appendages on 4 of the five sepals. Pubescence: Moderate. Stipitate Glands: Stipitate glands present on sepal margins.

Receptacle.—Surface: Glaucous. Shape: Funnel-shaped. Size: 5 mm (h)×6 mm (w). Color: Green Group 144B, with some anthocyanin intonation of Greyed-Purple Group 183D.

Peduncle.—Surface: Many stipitate glands and fine white hairs present. Length: 28 to 32 mm. Color: Yellow-Green Group 144B. Strength: Somewhat strong. Anthocyanin: Greyed-Purple Group 183D.

Borne.—2 to 3 buds per stem.

Flower bloom:

Fragrance.—Light, fresh scent.

Duration.—As a pot plant, flowers last from 12 to 15 days.

Size.—Small. Average flower diameter is between 40 to 45 mm when open.

Form.—Shape of flower when viewed from the side: Upon opening, upper part: Flattened convex. Upon opening, lower part: Flattened convex. After opening, upper part: Flattened convex. After opening, lower part: Concave. Petalage: Very double. Average range: 85 to 90 petals, with 10 to 15 petaloids.

Color:

Upon opening, petals.—Outermost petals: Upper surface: Red Group 53C. Lower surface: Red Group 53B. Innermost petals: Upper surface: Red Group 53C. Lower surface: Red Group 53B.

Upon opening, basal petal spots.—Upper surface: Yellow Group 4D. Lower surface: Yellow Group 4D.

After opening, petals.—Outermost petals: Upper surface: Red Group 53C. Lower surface: Red Group 53B. Innermost petals: Upper surface: Red Group 53C. Lower surface: Red Group 53B.

After opening, basal petal spots.—Upper surface: Yellow Group 4D. Lower surface: Yellow Group 4D.

General tonality: On open flower, Red Group 53C. No change in the general tonality at the end of the 7th day. Afterwards, general tonality is Red Group 53C and Red Group 57A.

Petals:

Petal reflex.—Slight.

Shape.—Obovate.

Petaloids.—3 to 4. Petaloids are 5 to 7 mm long and 3 to 4 mm wide. Petaloid coloration is Red Group 43B on both upper and lower surfaces.

Thickness.—Average.

Arrangement.—Imbricated.

Reproductive organs:

Filaments.—Color: Yellow-Green Group 154D. Length: 4 mm.

Pistils.—Length: 6 mm. Quantity: 30.

Pollen.—Color: Yellow Group 12B. Quantity: Average.

Anthers.—Length: 1 mm. Color: Brown Group 200C. Quantity: 20 to 24.

Stigmas.—Height: Stigmas are even in height relative to anthers. Color: Greyed-Green Group 196D.

Styles.—Color: Greyed-Green Group 192A. Length: 4 mm.

Hips.—None observed.

PLANT

Plant growth: Compact, spreading growth. Average height of the plant is 36 cm; the average width is 30 cm.

Stems:

Color.—Young wood: Yellow-Green Group 144B. Older wood: Yellow-Green Group 144A and B.

Thorns.—Incidence: 10 to 12 thorns per 10 cm of stem.

Size.—Average length is 3 mm. Color: Greyed-Yellow Group 160D. Shape: Linear.

Surface.—Older wood: Smooth. Young wood: Smooth.

Plant foliage:

Number of leaflets on normal leaves in middle of a stem.—5.

Leaf size.—60 mm (l) by 40 mm (w).

Quantity.—Average.

Glossiness.—Upper side weak glossiness.

Color.—Mature foliage: Upper Leaflet Surface: Yellow-Green Group 146A. Lower Leaflet Surface: Yellow-Green Group 146C. Juvenile foliage: Upper Leaflet Surface: Yellow-Green Group 146A. Lower Leaflet Surface: Yellow-Green Group 146C. Anthocyanin intonation: Greyed-Red Group 185A on margins of upper leaflet surface, all of lower surface, petiole, rachis, and stipule.

Plant leaflets:

Stipule.—Size: 10 mm (l)×4 mm (w). Color: Yellow-Green Group 144B and C. Anthocyanin: Greyed-Red Group 183C. Stipitate Glands: On edges of stipules.

Petiole.—Length: 7 mm. Color: Yellow-Green Group 144B. Underneath: Yellow-Green Group 144B. Stipitate glands and occasionally a single, small thorn. Margins: Yellow-Green Group 144A. Anthocyanin: Greyed-Red Group 183C.

Rachis.—Color: Yellow-Green Group 144B. Underneath: Yellow-Green Group 144B. Stipitate glands and occasionally a single, small thorn. Margins: Yellow-Green Group 144A. Stipitate glands present. Anthocyanin: Greyed-Red Group 183C.

Leaflet.—Edge: Dentate. Shape: Ovate. Surface: Moderately glossy. Texture: Leathery. Arrangement: Odd pinnate. Ventation: Reticulate.

Disease resistance: Above average resistance to mildew, rust, black spot, and Botrytis under normal growing conditions in Jackson County, Ore.

US PP13,592 P3

5

Cold hardiness: The variety 'POULgo004' has been found to be resistant to damage from cold in USDA Zone 7.

We claim:

1. A new and distinct variety of rose plant of the miniature rose class, substantially as herein illustrated and described as

6

a distinct and novel rose variety due to its abundant red flowers, procumbent to arching habit, disease resistance, and extended period of bloom.

* * * * *

