

(12) PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 200050958 B2**
(10) Patent No. **778910**

(54) Title
Methods and compositions for the manufacture of C-3' and C-4' anthracycline antibiotics

(51)⁷ International Patent Classification(s)
C07H 001/00 C07H 015/24

(21) Application No: **200050958**

(22) Application Date: **2000.03.15**

(87) WIPO No: **WO00/56267**

(30) Priority Data

(31) Number	(32) Date	(33) Country
60/125505	1999.03.19	US

(43) Publication Date : **2000.10.09**

(43) Publication Journal Date : **2000.12.07**

(44) Accepted Journal Date : **2004.12.23**

(71) Applicant(s)
Waldemar Priebe; Izabela Fokt; Teresa Przewloka; Marta Krawczyk; Piotr Skibicki; Grzegorz Grynkiewicz; Roman Perez-Soler

(72) Inventor(s)
Waldemar Priebe; Izabela Fokt; Teresa Przewloka; Marta Krawczyk; Piotr Skibicki; Grzegorz Grynkiewicz; Roman Perez-Soler

(74) Agent/Attorney
F.B. Rice and Co., 139 Rathdowne Street, CARLTON VIC 3053

(56) Related Art
WO 1994/020114
AU 56888/90



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A61K	A2	(11) International Publication Number: WO 00/56267 (43) International Publication Date: 28 September 2000 (28.09.00)
(21) International Application Number: PCT/IB00/00527 (22) International Filing Date: 15 March 2000 (15.03.00) (30) Priority Data: 60/125,505 19 March 1999 (19.03.99) US (63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application US 60/125,505 (CON) Filed on 19 March 1999 (19.03.99) (71)(72) Applicants and Inventors: PRIEBE, Waldemar [US/US]; 4239 Emory Avenue, Houston, TX 77005 (US). FOKT, Izabela [PL/US]; 1908 Nursery Road, The Woodlands, TX 77380 (US). PRZEWLOKA, Teresa [PL/US]; Apartment No.1406, 1200 Sawmill Road, The Woodlands, TX 77380 (US). KRAWCZYK, Marta [PL/US]; Apartment #2308, 175 N. Locust Hill Drive, Lexington, KY 40503 (US). SKIBICKI, Piotr [PL/PL]; Apartment 24, Waszyngton Street 39, PL-04-015 Warsaw (PL). GRYNKIEWICZ, Gregorz [PL/PL]; u. Zielona 16 B/2, PL-05-092 Lominaki (PL). PEREZ-SOLER, Roman [PL]; - (PL). <i>564 1st Ave. #20T New York NY 10016 United States of America</i>	(74) Agent: BOWLES, Teresa, J.; Fulbright & Jaworski, L.L.P., Suite 2400, 600 Congress Avenue, Austin, TX 78701 (US). (81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published Without international search report and to be republished upon receipt of that report. <i>→ 4210 S. Ravinia Dr. Apt. No. 114 Greenfield WI 53221 United States of America</i>	
(54) Title: METHODS AND COMPOSITIONS FOR THE MANUFACTURE OF C-3' AND C-4' ANTHRACYCLINE ANTIBIOTICS (57) Abstract The present invention discloses new and novel substituted anthracyclines with modified alkyl-aromatic ring substitutions on the C-3' of the sugar moiety or modified or unmodified alkyl-aromatic ring substitutions at the C-4' of the sugar moiety. It also discloses novel methods for the preparation of sugar substrates and methods for the preparation of anthracycline antibiotics. These anthracycline analogs show high cytotoxicity <i>in vitro</i> against several tumor cell lines.		

DESCRIPTION

METHODS AND COMPOSITIONS FOR THE MANUFACTURE OF C-3' AND C-4' ANTHRACYCLINE ANTIBIOTICS

BACKGROUND OF THE INVENTION

5 The U.S. Government may own rights in the present invention pursuant to National Institute of Health grants numbered C3A55270 and CA50320.

1. Field of the Invention

10 The present invention relates generally to the treatment of cancer. More particularly, it concerns novel compounds useful for chemotherapy, methods of synthesis of these compounds and methods of treatment employing these compounds. These novel drugs comprise two main classes of compounds; one bearing modified substituents at the C-3' sugar moiety and the other bearing modifications at the C-4' sugar moiety. In addition, some of these analogs might also be modified at the aglycon and/or sugar moiety. These novel anthracycline analogs display high anti-tumor activity and can be used as potent drugs active against multi-drug resistant tumors. These compounds are related to other anti-tumor anthracyclines such as daunorubicin, idarubicin, epirubicin, and doxorubicin. The cytotoxic potency of these new compounds is significantly higher when compared to doxorubicin.

20

2. Description of Related Art

25 Resistance of tumor cells to the killing effects of chemotherapy is one of the central problems in the management of cancer. It is now apparent that at diagnosis many human tumors already contain cancer cells that are resistant to standard chemotherapeutic agents. Spontaneous mutation toward drug resistance is estimated to occur in one of every 10^6 to 10^7 cancer cells. This mutation rate appears to be independent of any selective pressure from drug therapy, although radiation therapy and chemotherapy may give rise to additional mutations and contribute to tumor progression within cancer cell populations (Goldie *et al.*, 1979; Goldie *et al.*, 1984;

Nowell, 1986). The cancer cell burden at diagnosis is therefore of paramount importance because even tumors as small as 1 cm (10^9 cells) could contain as many as 100 to 1,000 drug-resistant cells prior to the start of therapy.

5 Selective killing of only the tumor cells sensitive to the drugs leads to an overgrowth of tumor cells that are resistant to the chemotherapy. Mechanisms of drug resistance include decreased drug accumulation (particularly in multi-drug resistance), accelerated metabolism of the drug and other alterations of drug metabolism, and an increase in the ability of the cell to repair drug-induced damage (Curt *et al.*, 1984; and
10 Kolate, 1986). The cells that overgrow the tumor population not only are resistant to the agents used but also tend to be resistant to other drugs, many of which have dissimilar mechanisms of action. This phenomenon, called pleiotropic drug resistance or multi-drug resistance (MDR), may account for much of the drug resistance that occurs in previously treated cancer patients. The development of drug resistance is
15 one of the major obstacles in the management of cancer. One of the traditional ways to attempt to circumvent this problem of drug resistance has been combination chemotherapy.

 Combination drug therapy is the basis for most chemotherapy employed to
20 treat breast, lung, and ovarian cancers as well as Hodgkin's disease, non-Hodgkin's lymphomas, acute leukemias, and carcinoma of the testes. Combination chemotherapy uses the differing mechanisms of action and cytotoxic potentials of multiple drugs.

25 Although combination chemotherapy has been successful in many cases, the need still exists for new anti-cancer drugs. These new drugs could be such that they are useful in conjunction with standard combination chemotherapy, or these new drugs could attack drug resistant tumors by having the ability to kill cells of multiple resistance phenotypes.

A drug that exhibits the ability to overcome multiple drug resistance could be employed as a chemotherapeutic agent either alone or in combination with other drugs. The potential advantages of using such a drug in combination with chemotherapy would be the need to employ fewer toxic compounds in the combination, cost savings, and a synergistic effect leading to a treatment regime involving fewer treatments.

The commonly used chemotherapeutic agents are classified by their mode of action, origin, or structure, although some drugs do not fit clearly into any single group. The categories include alkylating agents, anti-metabolites, antibiotics, alkaloids, and miscellaneous agents (including hormones). Agents in the different categories have different sites of action.

Antibiotics are biologic products of bacteria or fungi. They do not share a single mechanism of action. The anthracyclines daunorubicin and doxorubicin (DOX) are some of the more commonly used chemotherapeutic antibiotics. The anthracyclines achieve their cytotoxic effect by several mechanisms, including inhibition of topoisomerase II; intercalation between DNA strands, thereby interfering with DNA and RNA synthesis; production of free radicals that react with and damage intracellular proteins and nucleic acids; chelation of divalent cations; and reaction with cell membranes. The wide range of potential sites of action may account for the broad efficacy as well as the toxicity of the anthracyclines (Young *et al.*, 1985).

The anthracycline antibiotics are produced by the fungus *Streptomyces peuceitius* var. *caesius*. Although they differ only slightly in chemical structure, daunorubicin has been used primarily in the acute leukemias, whereas doxorubicin displays broader activity against human neoplasms, including a variety of solid tumors. The clinical value of both agents is limited by an unusual cardiomyopathy, the occurrence of which is related to the total dose of the drug; it is often irreversible. In a search for agents with high anti-tumor activity but reduced cardiac toxicity, anthracycline derivatives and related compounds have been prepared. Several of

these have shown promise in the early stages of clinical study, and some, like epirubicin and idarubicin, are used as drugs. Epirubicin outsells doxorubicin in Europe and Japan, but it is not sold in the U.S.

5 The anthracycline antibiotics have tetracycline ring structures with an unusual sugar, daunosamine, attached by glycosidic linkage. Cytotoxic agents of this class all have quinone and hydroquinone moieties on adjacent rings that permit them to function as electron-accepting and donating agents. Although there are marked differences in the clinical use of daunorubicin and doxorubicin, their chemical
10 structures differ only by a single hydroxyl group on C14. The chemical structures of daunorubicin and doxorubicin are shown in FIG. 1.

 Doxorubicin's broad spectrum of activity against most hematological malignancies as well as carcinomas of the lung, breast, and ovary has made it a
15 leading agent in the treatment of neoplastic disease (Arcamone, 1981; Lown, 1988; Priebe, 1995). Since the discovery of daunorubicin and doxorubicin (FIG. 1), the mechanistic details of the anti-tumor activity of anthracycline antibiotics have been actively investigated (Priebe, 1995a; Priebe, 1995b; Booser, 1994).

20 Unfortunately, concomitant with its anti-tumor activity, DOX can produce adverse systemic effects, including acute myelosuppression, cumulative cardiotoxicity, and gastrointestinal toxicity (Young *et al.*, 1985). At the cellular level, in both cultured mammalian cells and primary tumor cells, DOX can select for multiple mechanisms of drug resistance that decrease its chemotherapeutic efficacy.
25 These mechanisms include P-gp-mediated MDR and MPR-mediated MDR, characterized by the energy-dependent transport of drugs from the cell (Bradley *et al.*, 1988), and resistance conferred by decreased topoisomerase II activity, resulting in the decreased anthracycline-induced DNA strand scission (Danks *et al.*, 1987; Pommier *et al.*, 1986; Moscow *et al.*, 1988).

Among the potential avenues of circumvention of systemic toxicity and cellular drug resistance of the natural anthracyclines is the development of semisynthetic anthracycline analogues which demonstrate greater tumor-specific toxicity and less susceptibility to various forms of resistance.

5

SUMMARY OF THE INVENTION

It is desirable to provide compositions of agents that display increased cytotoxicity when compared with doxorubicin and can prevent and/or overcome multi-drug resistance and exhibit reduced cardiotoxicity. This invention involves novel
10 compounds that have utility as anti-tumor and/or chemotherapeutic drugs, methods of synthesizing these compounds and methods of using these compounds to treat patients with cancer. The invention is based on the discovery that anthracycline derivatives with substitutions attached to their C-3 or C-4 carbons in the sugar moiety have a surprisingly strong ability to kill multi-drug resistant tumor cells.

15

New anthracycline-based agents designed to interact and crosslink with DNA have been synthesized. These analogs contain substitutions at the C-3 or C-4 sugar moiety. Synthesized compounds displayed activity significantly higher than that of parental daunorubicin or doxorubicin. In brief, in vitro the compounds WP755,
20 WP756, WP757, WP758, WP775, WP778, WP784, WP786, WP790, WP791 modified at the C-3 and WP744, WP783 and WP750 modified at the C-4 were significantly more effective as measured by resistance index (RI) (Table 2). The RI values for the 3-O-substituted analogs vary from 1.2 to 36 and are low when compared to the RI value of 253 and > 200 for DOX, wherein, a higher RI value indicates that a compound is less
25 effective against MDR. Similarly, RI values were very low (1.4-D8.8) for 4-O-alkylated analogs whereas RI values for DOX varied from > 42.6 to > 200 for MDR1 type of resistance and from 10 to 16 for the MRP form of resistance. Lower RI values indicate greater efficacy of the drug against MDR tumors. The inventors also designed and synthesized other analogs. The observed

activity and high potency against multi-drug resistant tumors indicate that these analogs are different from the parental drugs like doxorubicin and daunorubicin.


The inventors synthesized a series of analogs substituted at the aromatic ring
5 of the C-3'-substituent which were then combined with modifications at the aglycon moiety. The inventors discovered that substitution at the aromatic ring increased the potency and alter the mechanism of action of the drugs making them significantly more active than doxorubicin. The mechanism of action of this class of drugs might involve direct interaction of the aromatic ring with cellular targets like DNA,
10 topoisomerase II and topoisomerase I. Substitution of a benzyl ring at C-3' modifies drug interaction with P-glycoprotein, and subsequently the resistance index, thereby makes these compounds more effective against MDR tumors in comparison to the parent drug. *In vitro* evaluation identified the following C-3' substituted anthracycline analogs: WP831, WP791, WP790, WP786, WP785, WP784, WP780,
15 WP778, WP775, WP774, WP765, WP758, WP757, WP756 and WP755 as unusually effective cytotoxic agents when compared to DOX.

The inventors also synthesized anthracycline analogs with substitutions at the C-4' sugar and demonstrated that these analogs overcome both, (a) multi-drug
20 resistance (MDR) caused by overexpression of the *MDR1* gene and (b) MDR-associated protein (MRP)-related resistance caused by overexpression of the *MRP* gene. Such modifications also contribute to the drugs ability to circumvent others forms of drug resistance and increased bioavailability. An increased steric hindrance at C-4' in doxorubicin might reduce drug interaction with P-glycoprotein and MRP and in combination, the increased lipophilicity caused by introduction of
25 aromatic ring further contributes to increase intracellular drug concentration in MDR cells. Such modifications alter cellular uptake and retention of the drugs without affecting interaction with cellular targets, which results in cytotoxic effects. *In vitro* evaluation identified the following C-4' substituted anthracycline analogs: WP799,
30 WP797, WP794, WP787, WP783, WP750, WP744, WP727, WP764 and WP571 as unusually potent cytotoxic agents when compared to DOX.

The anthracycline compounds bearing the C-3' substitutions have among others, O-Benzyl, N-Benzyl or S-Benzyl substitutions where the phenyl, an aromatic ring of the benzyl group is substituted. The compounds bearing C-4' substitutions at the sugar have among others, O-Benzyl, N-Benzyl or S-Benzyl substitutions where the benzyl group is a substituted or an unsubstituted benzene group. These compounds are depicted in FIGs. 2-25. These compounds exhibit cytotoxic activity substantially different from the activities of doxorubicin or daunorubicin. These compounds are active against doxorubicin resistant tumors and/or are usually similar or more cytotoxic than doxorubicin against sensitive tumors.

10

According to the invention there is provided a substituted anthracycline having the formula:



15 wherein, R^1 denotes an alkyl chain, a $(-COCH_2R^{13})$ group or a $(C(OH)-CH_2R^{13})$;

wherein, R^{13} is a hydrogen (-H) group, a hydroxyl group (-OH); a methoxy group (-OCH₃), an alkoxy group having 1-20 carbon atoms, an alkyl group having 1-20 carbon atoms, an aryl group having 1-20 carbon atoms, a fatty acyl group having the general structure $-O-CO(CH_2)_nCH_3$, wherein n = an integer from 1 to about 20, a fatty acyl group having the general structure $-O-CO(CH_2)_l(CH=CH)_m(CH_2)_nCH_3$, wherein l

20

is an integer between 1 to 3, m is an integer between 1 and 6, and n is an integer between 1 and 9, a $-\text{OCO}-(\text{CH}_2)_n-\text{CH}_2\text{NH}_2$, or $\text{OCO}-(\text{CH}_2)_n-\text{CO}_2\text{H}$:

each of R^2 and R^3 is, independently of the other, a hydrogen (-H), a hydroxyl group (-OH) or a methoxy group ($-\text{OCH}_3$);

R^4 is a hydrogen (-H) group, a methoxy group ($-\text{OCH}_3$), a hydroxyl group (-OH) or a halide;

each of Y^1 and Y^2 is, independently of the other, a double bonded oxygen, sulphur, or nitrogen atom;

Z is a -H, -OH, a $-\text{CO}_2\text{H}$ group or a $-\text{CO}_2\text{R}$ group;

R^7 , R^8 , are, independently, -H, -OH, a halide, $-\text{OR}^{19}$, -SH, $-\text{SR}^{19}$, $-\text{NH}_2$, $-\text{NHR}^{19}$, $-\text{N}(\text{R}^{19})_2$ or $-\text{CH}_3$, and R^7 can additionally be a saccharide, wherein R^{19} is an alkyl chain, an alkylating moiety, a cycloalkyl chain, a cyclic ring, or a hydrogen;


R^9 is an -H, $-\text{CH}_3$, alkyl, aryl, CH_2OH , or a CH_2F group;

R^{10} , R^{11} and R^{12} are, independently, -H, -OH, a halide, -OR, -SH, -SR, $-\text{NH}_2$, -NHR, $-\text{N}(\text{R})_2$, or a $-\text{CH}_3$;

one of R^5 and R^6 is a -H;

one of R^5 and R^6 is a X-alkyl-aromatic-ring (-XAAR) substituent, wherein, A is an alkyl group and wherein, AR is a substituted phenyl ring, or a substituted five-member ring, a heteroatomic five-member ring, or a heteroatomic six-member ring, of the form:

7b



;

wherein, $R^{14} - R^{18}$ are independently a (-H) group, a hydroxyl group (-OH), a methoxy group (-OCH₃), a nitro group (-NO₂), an amine group (-NH₂), a halide, an alkoxy group having 1-20 carbon atoms, an alkyl group having 1-20 carbon atoms, an aryl group having 1-20 carbon atoms, an alkyl-amino group, an alkyl-thio group, a cyano group (CN, SCN), an -CO₂H group or an -CO₂R group; and


the aromatic ring is unsubstituted, disubstituted, trisubstituted, tetrasubstituted or pentasubstituted; and

X is a -N, a -S, a -SO, or -SO₂ group; and

A is (CH₂)_n where n = 0-10;

wherein, if R⁵ is a XAAR substituent R⁶ is not and if R⁶ is a XAAR substituent R⁵ is not.

According to the invention there is provided a substituted anthracycline having the formula:



wherein, R^1 denotes an alkyl chain, a $(-\text{COCH}_2\text{R}^{13})$ group, or a $(\text{C}(\text{OH})-\text{CH}_2\text{R}^{13})$ group;

wherein, R^{13} is a hydrogen $(-\text{H})$ group, a hydroxyl group $(-\text{OH})$, a methoxy group $(-\text{OCH}_3)$, an alkoxy group having 1-20 carbon atoms, an alkyl group having 1-20 carbon atoms, an aryl group having 1-20 carbon atoms, a fatty acyl group having the general structure $-\text{O}-\text{CO}(\text{CH}_2)_n\text{CH}_3$, wherein n = an integer from 1 to about 20, a fatty acyl group having the general structure $-\text{O}-\text{CO}(\text{CH}_2)_l(\text{CH}=\text{CH})_m(\text{CH}_2)_n\text{CH}_3$, wherein l is an integer between 1 to 3, m is an integer between 1 and about 6, and n is an integer between 1 to about 9, $-\text{OCO}-(\text{CH}_2)_n-\text{CH}_2\text{NH}_2$ or $\text{OCO}-(\text{CH}_2)_n-\text{CO}_2\text{H}$;

each of R^2 and R^3 is, independently of the other, a hydrogen $(-\text{H})$, a hydroxyl group $(-\text{OH})$ or a methoxy group $(-\text{OCH}_3)$;

R^4 is a hydrogen $(-\text{H})$ group, a methoxy group $(-\text{OCH}_3)$, a hydroxyl group $(-\text{OH})$ or a halide;

each of Y^1 and Y^2 is, independently of the other, a double bonded oxygen, sulphur or nitrogen atom;

5 Z is a $-H$, $-OH$, a $-CO_2H$ group or a $-CO_2R$ group;

R^5 and R^6 , are, independently, $-H$, $-OH$, a halide, $-OR^{19}$, $-SH$, $-SR^{19}$, $-NH_2$, $-NHR^{19}$, $-N(R^{19})_2$, or $-CH_3$, and R^6 can additionally be an alkylating moiety, wherein R^{19} is an alkyl chain, an alkylating moiety, a cycloalkyl chain, a cyclic ring or a hydrogen,

10


R^9 is an $-H$, $-CH_3$, alkyl, aryl, CH_2OH or a CH_2F group;

R^{10} , R^{11} and R^{12} are, independently, $-H$, $-OH$, a halide, $-OR$, $-SH$, $-SR$, $-NH_2$, $-NHR$, $-N(R)_2$ or $-CH_3$;

15

one of R^7 and R^8 is a $-H$;

one of R^7 and R^8 is a X-alkyl aromatic-ring ($-XAAR$) substituent, wherein, A is an alkyl group and wherein, AR is an unsubstituted phenyl ring, a substituted phenyl ring,
20 a substituted five-member ring or a heteroatomic five-member ring, of the general form;



;

wherein, R^{14} - R^{18} are independently a $(-H)$ group, a hydroxyl group $(-OH)$, a methoxy group $(-OCH_3)$, a nitro group $(-NO_2)$, an amine group $(-NH_2)$, a halide, an alkoxy group
25

having 1-20 carbon atoms, an alkyl group having 1-20 carbon atoms, an aryl group having 1-20 carbon atoms, an alkyl-amino group, an alkyl-thio group, a cyano group (CN, SCN), an $\text{-CO}_2\text{H}$ group or an $\text{-CO}_2\text{R}$ group; and

- 5 the aromatic ring is unsubstituted, disubstituted, trisubstituted, tetrasubstituted or pentasubstituted; and

X is a -O , -N , -S , -SO or a -SO_2 group; and

A is $(\text{CH}_2)_n$, where $n = 0\text{-}10$;

- 10 wherein if R^7 is a XAAR substituent R^8 is not and if R^8 is a XAAR substituent R^7 is not.

According to the invention there is provided a method for the synthesis of amine containing anthracyclines comprising:


- 15 (a) obtaining a substituted sugar azide, wherein the azido substitution is at the 1', 2', 3', 4' or 5' position on the sugar, said azide serving as a masked and neutral form of amine substituent;

(b) subjecting said azide to a coupling reaction with an anthracyclinone; and

- (c) subjecting said azide to a reduction reaction;


20 wherein an amine containing anthracycline is produced

In some specific embodiments, the C-3 -substituted anthracycline compounds of the present invention have the general formula:



wherein, R' denotes any suitable group or combination of groups that form but are not limited to a nucleic acid intercalator or binding compound; a topoisomerase inhibitor, including but not limited to, an alkyl chain; a (-COCH₂R¹³) group; or a (-C(OH)-CH₂R¹³); wherein, R¹³ is a hydrogen (-H) group or a hydroxyl group (-OH); a methoxy group (-OCH₃); an alkoxy group having 1-20 carbon atoms; an alkyl group having 1-20 carbon atoms; an aryl group having 1-20 carbon atoms; a fatty acyl group


having the general structure $(-O-CO(CH_2)_nCH_3)$ wherein $n =$ an integer from 1 to about 20; or a fatty acyl group having the general structure $(-O-CO(CH_2)_l(CH=CH)_m(CH_2)_nCH_3)$ wherein l is an integer between 1 to 3, m is an integer between 1 and about 6, and n is an integer between 1 to about 9; or a chain(R) such as $-OCO-(CH_2)_n-CH_2NH_2$; or $OCO-(CH_2)_n-CO_2H$ and its salts; each of R^2 and R^3 is, independently of the other, a hydrogen (-H), a hydroxyl group (-OH); a methoxy group (-OCH₃); R^4 is a hydrogen (-H) group; a methoxy group (-OCH₃); a hydroxyl group (-OH); or a halide; each of Y^1 and Y^2 is, independently of the other, a double bonded oxygen, sulphur, or nitrogen atom; Z is a -H; -OH; a -CO₂H group; or a -CO₂R group; R^7 , R^8 , are, independently, -H; -OH; a halide; -OR¹⁹; -SH; -SR¹⁹; -NH₂; -NHR¹⁹; -N(R¹⁹)₂; -CH₃; and R^7 can additionally be a saccharide; wherein R^{19} is an alkyl chain; an alkylating moiety; a cycloalkyl chain; a cyclic ring; or a hydrogen; R^9 can be -H; -CH₃; alkyl; aryl; CH₂OH, CH₂F; R^{10} , R^{11} and R^{12} are, independently, -H; -OH; a halide; -OR; -SH; -SR; -NH₂; -NHR; -N(R)₂; -CH₃; one of R^5 and R^6 is a -H; one of R^5 and R^6 is a X-alkyl-aromatic-ring (AAR) substituent such as -XAAR, wherein, A is an alkyl group and wherein, AR is an substituted phenyl ring; or a substituted five-member ring; or a heteroatomic five-member ring; or a heteroatomic six-member ring such as a pyridine ring; of the form;



; wherein, $R^{14}-R^{18}$ are independently a (-H) group; a hydroxyl group (-OH); a methoxy group (-OCH₃); a nitro group (-NO₂), an amine group (-NH₂), a halide; an alkoxy group having 1-20 carbon atoms; an alkyl group having 1-20 carbon atoms; an aryl group having 1-20 carbon atoms; an alkyl-amino group; an alkyl-thio group; a cyano group (CN, SCN); an -CO₂H group; an -CO₂R group; and the aromatic ring may be disubstituted, trisubstituted, tetrasubstituted or pentasubstituted; and X is a -O,


-N or -S, or -SO, or -SO₂ group; and A is (CH₂)_n where n = 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, wherein, if R⁵ is a XAAR substituent R⁶ is not and if R⁶ is a XAAR substituent R⁵ is not.

- 5 In other specific embodiments, the C-4' substituted anthracycline compounds of the present invention have the general formula:



- wherein, R¹ denotes any suitable group or combination of groups that form but are not limited to a nucleic acid intercalator or binding compound; a topoisomerase inhibitor, including but not limited to, an alkyl chain; a (-COCH₂R¹³) group; or a (C(OH)-CH₂R¹³) wherein, R¹³ is a hydrogen (-H) group or a hydroxyl group (-OH); a methoxy group (-OCH₃); an alkoxy group having 1-20 carbon atoms; an alkyl group having 1-20 carbon atoms; an aryl group having 1-20 carbon atoms; a fatty acyl group having the general structure (-O-CO(CH₂)_nCH₃) wherein n = an integer from 1 to about 20; or a fatty acyl group having the general structure (-O-CO(CH₂)_l(CH=CH)_m(CH₂)_nCH₃), wherein l is an integer between 1 to 3, m is an integer between 1 and about 6, and n is an integer between 1 to about 9; or a chain(R) such as -OCO-(CH₂)_n-CH₂NH₂; or OCO-(CH₂)_n-CO₂H and its salts; each of R² and R³ is, independently of the other, a hydrogen (-H), a hydroxyl group (-OH); a methoxy group (-OCH₃); R⁴ is a hydrogen (-H) group; a methoxy group (-OCH₃); a hydroxyl

group (-OH); or a halide; each of Y^1 and Y^2 is, independently of the other, a double bonded oxygen, sulphur, or nitrogen atom; Z is a -H; -OH; a -CO₂H group; or a -CO₂R group; R^5 , R^6 , are, independently, -H; -OH; a halide; -OR¹⁹; -SH; -SR¹⁹; -NH₂; -NHR¹⁹; -N(R¹⁹)₂; -CH₃; wherein R¹⁹ is an alkyl chain; an alkylating moiety; a cycloalkyl chain; a cyclic ring; or a hydrogen; and R⁶ can additionally be a an alkylating moiety; R⁹ can be -H; -CH₃; alkyl; aryl; CH₂OH, CH₂F; R¹⁰, R¹¹ and R¹² are, independently, -H; -OH; a halide; -OR; -SH; -SR; -NH₂; -NHR; -N(R)₂; -CH₃; one of R⁷ and R⁸ is a -H; one of R⁷ and R⁸ is a X-alkyl aromatic-ring (AAR) substituent such as -XAAR, wherein, A is an alkyl group and wherein, AR is an unsubstituted phenyl ring; or a substituted phenyl ring; or a substituted five-member ring such as a pyridine ring; or a heteroatomic five-member ring, of the general form;



; wherein, R¹⁴-R¹⁸ are independently a (-H) group; a hydroxyl group (-OH); a methoxy group (-OCH₃); a nitro group (-NO₂), an amine group (-NH₂), a halide; an alkoxy group having 1-20 carbon atoms; an alkyl group having 1-20 carbon atoms; an aryl group having 1-20 carbon atoms; an alkyl-amino group; an alkyl-thio group; a cyano group (CN, SCN); an -CO₂H group; an -CO₂R group; and the aromatic ring may be disubstituted, trisubstituted, tetrasubstituted or pentasubstituted; and X is a -O, -N or -S, or -SO, or -SO₂ group; and A is (CH₂)_n where n = 0, 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10; wherein if R⁷ is a XAAR substituent R⁸ is not and if R⁸ is a XAAR substituent R⁷ is not.

Certain specific embodiments of the anthracyclines of the invention are shown in FIGs. 2-25.

The present application also comprises methods of preparing novel substituted sugar substrates and their use in the synthesis of the novel anthracycline analogs described in this invention. In certain embodiments the method for the synthesis of 4'-O-benzylated sugars is described. These 4-O-benzylated sugars may comprise one of two main classes; glycal sugars or 1-O-silylated sugars. Examples of the benzylated sugars encompassed by this invention are WP567, WP735, WP736, WP819, WO820, WP821, WP822, WP823, WP824 and WP825.

Related embodiments describe the method for synthesizing glycals using various bases including but not limited to NaH. Other related embodiments describe the synthesis of glycals using various solvents including but not limited to DMF. Yet other related embodiments describe the synthesis of glycals using various alkylating agents including but not limited to benzyl chloride and benzyl bromide.

In certain embodiments, the method for the synthesis of amine containing analogs of anthracyclines is described. Further embodiments describe the use of substituted sugar azides for the synthesis of said amine containing anthracyclines wherein the azido substitution can be at the 1', 2', 3', 4' or 5' position on the sugar. The azide group serves as a masked or neutral form of amine substituent allowing for a coupling reaction (explained in Example 2, Procedure A). This allows the generation of selective conditions to reduce azides (explained in Example 2, Procedure B). In one example the amine containing anthracyclines synthesized by this procedure are 14-OH analogs similar to doxorubicin, epirubicin or daunorubicin such as WP744 and WP769. The azido sugar used in the preparation of this compound is WP819 (explained in Example 1, Procedure A and B).

This procedure also allows the use of 14-O-blocked aglycons as these blocked groups survive the steps in which azides are reduced and can be selectively removed at a later stage (Example 2, Procedure B).

The present application comprises methods of preparing substituted anthracyclines and the preparation of important sugar substrates. In devising the synthetic schemes and compounds of the present invention, the inventors have created a variety of novel compounds. These compounds and their methods of synthesis are
5 described elsewhere in the specification, examples and figures and are given "WP" numbers. The structure of a compound designated with a "WP" number is ascertainable by reviewing the specification and figures. Exemplary specific anthracycline compounds that are encompassed by the invention are WP831, WP791, WP790, WP787, WP786, WP785, WP784, WP780, WP778, WP775, WP774,
10 WP765, WP764, WP758, WP757, WP756, WP755, WP799, WP797, WP794, WP783, WP750, WP744, WP727 and WP571. Exemplary specific sugar substrates that are encompassed by the invention are WP567, WP735, WP736, WP819, WP820, WP821, WP822, WP823, WP824, WP825.

15 The invention also considers methods of treating a patient with cancer, comprising administering to the patient a therapeutically effective amount of the contemplated substituted anthracycline compounds and therapeutic kits comprising, in suitable container means, a pharmaceutically acceptable composition comprising the contemplated substituted anthracycline compounds.

20 Throughout this specification the word "comprise", or variations such as "comprises" or "comprising", will be understood to imply the inclusion of a stated element, integer or step, or group of elements, integers or steps, but not the exclusion of any other element, integer or step, or group of elements, integers or steps.

25 Any discussion of documents, acts, materials, devices, articles or the like which has been included in the present specification is solely for the purpose of providing a context for the present invention. It is not to be taken as an admission that any or all of these matters form part of the prior art base or were common general knowledge in the
30 field relevant to the present invention as it existed before the priority date of each claim of this application.

BRIEF DESCRIPTION OF THE DRAWINGS

35 The following drawings form part of the present specification and are included to further demonstrate certain aspects of the present invention. The invention may be

better understood by reference to one or more of these drawings in combination with the detailed description of specific embodiments presented herein.

- 5 FIG. 1. Structure of Doxorubicin and Daunorubicin
 FIG. 2. Structure of WP831
 FIG. 3. Structure of WP791
 FIG. 4. Structure of WP790
 FIG. 5. Structure of WP787
 FIG. 6. Structure of WP786




	FIG. 7.	Structure of WP785
	FIG. 8.	Structure of WP784
	FIG. 9.	Structure of WP780
	FIG. 10.	Structure of WP778
5	FIG. 11.	Structure of WP775
	FIG. 12.	Structure of WP774
	FIG. 13.	Structure of WP758
	FIG. 14.	Structure of WP757
	FIG. 15.	Structure of WP756
10	FIG. 16.	Structure of WP755
	FIG. 17.	Structure of WP799
	FIG. 18.	Structure of WP797
	FIG. 19.	Structure of WP794
	FIG. 20.	Structure of WP783
15	FIG. 21.	Structure of WP750
	FIG. 22.	Structure of WP744
	FIG. 23.	Structure of WP727
	FIG. 24.	Structure of WP567
	FIG. 25.	Structure of WP793
20	FIG. 26.	Structure of WP819
	FIG. 27.	Structure of WP820
	FIG. 28.	Structure of WP821
	FIG. 29.	Structure of WP822
	FIG. 30.	Structure of WP823
25	FIG. 31.	Structure of WP824
	FIG. 32.	Structure of WP825
	FIG. 33.	Structure of WP764
	FIG. 34.	Structure of WP765
	FIG. 35.	Synthesis of 4'-O-Benzylated Anthracyclines from 4-O-
30	Benzylated	glycals
	FIG. 36.	Synthesis of 4'-O-Benzyl-3'-Amino-Anthracyclines

- FIG. 37. Synthesis of 4-O-Benzyl-Glycals and its 3-O-Derivatives
- FIG. 38. Synthesis of 4'-O-Benzyl-3'-Deamino-Anthracyclines from 1-O-silylated 4-O-Benzyl-Hexopyranoses
- FIG. 39. Examples of Selective Alkylation of Acylated Glycals
- 5 FIG. 40. Synthesis of 3'-Azido-Sugars and its 4-O-Benzylated Derivatives
- FIG. 41. Alternative Synthesis of 3-Azido-4-O-Benzyl-Daunosamine from 3-Azidoacosamine
- FIG. 42. Structure of WP735
- 10 FIG. 43. Structure of WP736

DESCRIPTION OF ILLUSTRATIVE EMBODIMENTS

The present invention provides new and novel DNA intercalating agents.

- 15 These agents are substituted anthracyclines. These compounds show high activity against resistant tumors and cells. A novel approach of the invention produces compounds that are as active or more so than the parent compounds. Furthermore, the inventors' discovery is also for the design of effective DNA-binding substituted anthracyclines.

20

- The anthracycline compounds have a tetracycline ring structure with sugars attached by a glycosidic linkage. Cytotoxic agents of this class have quinone and hydroquinone moieties that permit them to function as electron-accepting and electron donating agents. Doxorubicin and daunorubicin are examples of compounds of this class (FIG. 1). These compounds act by intercalating with DNA. Examples of exemplary anthracyclinones and anthracyclines are given in Table I.
- 25

Table 1. List of Exemplary anthracyclonones and anthracyclines.**Anthracyclonones**

Rhodomycinone
Isorhodomycinone
Pyrromycinone
4-Demethoxydaunomycinone
4-Demethoxyadriamycinone
Daunomycinone
Adriamycinone

Anthracyclines

Daunorubicin
Doxorubicin
Epirubicin
Idarubicin
Pyrromycin
Aclacinamycine
Isorhodomycline
Carminomycine
Doxorubicine 14-esters:
Doxorubicin 14-acetate
Doxorubicin 14-propionate
Doxorubicin 14-octanoate
Doxorubicine 14-benzoate
Doxorubicine 14-phenylacetate
4'-Epidaunorubicin

TABLE 1 (CONT'D)

4'-Epidoxorubicin
4'-Iododaunorubicin
4'-Iododoxorubicin
4'-Deoxydaunorubicin
4'-Deoxydoxorubicin
3'-Hydroxydaunorubicin
3'-Hydroxydoxorubicin
4-Demethoxydaunorubicin
4-Demethoxydoxorubicin
4'-Epi-4-demethoxydaunorubicin
4'-Epi-4-demethoxydoxorubicin

Following long-standing patent law convention, the words "a" and "an", when used in the specification including the claims, denotes one or more.

5 "Aryl" may be a phenyl or alkyl group, unsubstituted or substituted with an amine, alkylamine, nitro, carboxy, sulfonic acid, hydroxy, oxyalkyl, or halide.

The term "saccharide" includes oxidized, reduced or substituted saccharides. Saccharides of this invention include, but are not limited to, ribose, arabinose, xylose, lyxose, allose, altrose, glucose, mannose, fructose, glucose, idose, galactose, talose, 10 ribulose, sorbose, tagatose, gluconic acid, glucuronic acid, glucaric acididuronic acid rhamnose, fucose, N-acetyl glucosamine, N-acetyl galactosamine, N-acetyl neuraminic acid, sialic acid, derivatives of saccharides such as acetals, amines, and phosphorylated sugars, oligosaccharides, as well as open chain forms of various sugars, and the like.

15

An individual skilled in the art of organic synthesis in light of the present disclosure is able to prepare a large variety of substituted sugars and substituted anthracyclines having C-3'-substituents or having C-4'-substitutents which are

expected to have chemotherapeutic activities and may be used in the treatment of cancer and/or other diseases. Exemplary substituted anthracyclines having C-3' alkylated anthracyclines modified at benzyl ring of the present invention are WP831, WP791, WP790, WP787, WP786, WP785, WP784, WP780, WP778, WP775, 5 WP774, WP758, WP757, WP756, WP755 (FIGs. 2-16). . Exemplary substituted anthracyclines having 4'-substituted-benzylated anthracyclines of the present invention are WP799, WP797, WP794, WP783, WP750, WP744, WP727 and WP571 (FIGs. 17-25). Such specific anthracyclines having C-3' substituted moieties or having C-4'-substituted moieties have been synthesized by the inventors and have 10 been analyzed and the structure confirmed by n.m.r and elemental analysis. The methods of the present application enable one of skill in the art to synthesize these compounds and many other related compounds without undue experimentation.

The present discoveries may be utilized in conjunction with certain techniques 15 that are well-known in the biological arts and that are further described in the following sections.

PHARMACEUTICAL COMPOSITIONS

20 The anti-tumor compounds of this invention can be administered to kill tumor cells by any method that allows contact of the active ingredient with the agent's site of action in the tumor. They can be administered by any conventional methods available for use in conjunction with pharmaceuticals, either as individual therapeutically active ingredients or in a combination of therapeutically active ingredients. They can be 25 administered alone but are generally administered with a pharmaceutically acceptable carrier selected on the basis of the chosen route of administration and standard pharmaceutical practice.

Aqueous compositions of the present invention will have an effective amount 30 of anthracycline to kill or slow the growth of cancer cells. Further the potential recognition of genes can be accomplished by the synthesis of substituted

anthracyclines having C-3' alkylated anthracyclines modified at benzyl ring and having 4'-substituted-benzylated anthracyclines with specific structures that allow for the recognition of specific parts of DNA. Such compositions will generally be dissolved or dispersed in a pharmaceutically acceptable carrier or aqueous medium.

5

The phrases "pharmaceutically or pharmacologically acceptable" refer to molecular entities and compositions that do not produce an adverse, allergic or other untoward reaction when administered to an animal, or human, as appropriate. As used herein, "pharmaceutically acceptable carrier" includes any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents and the like. The use of such media and agents for pharmaceutical active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active ingredients, its use in the therapeutic compositions is contemplated. Supplementary active ingredients, such as other anti-cancer agents, can also be incorporated into the compositions.

10

15

In addition to the compounds formulated for parenteral administration, such as intravenous or intramuscular injection, other pharmaceutically acceptable forms include, e.g., tablets or other solids for oral administration; time release capsules; and any other form currently used, including cremes, lotions, mouthwashes, inhalants and the like.

20

A. Parenteral Administration

The active compounds will often be formulated for parenteral administration, e.g., formulated for injection via the intravenous, intramuscular, subcutaneous, or even intraperitoneal routes. The preparation of an aqueous composition that contains an anthracycline of the present invention as an active ingredient will be known to those of skill in the art in light of the present disclosure. Typically, such compositions can be prepared as injectables, either as liquid solutions or suspensions; solid forms suitable for using to prepare solutions or suspensions upon the addition of a liquid prior to injection can also be prepared; and the preparations can also be emulsified.

25

30

Solutions of the active compounds as free base or pharmacologically acceptable salts can be prepared in water suitably mixed with a surfactant, such as hydroxypropylcellulose. Dispersions can also be prepared in glycerol, liquid
5 polyethylene glycols, and mixtures thereof and in oils. Under ordinary conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms.

In some forms, it will be desirable to formulate the novel compounds in salt
10 form, generally to improve the solubility and bioavailability and to provide an active drug form more readily assimilated. As used herein, the term "pharmaceutically acceptable salt" refers to compounds which are formed from acidifying a substituted anthracycline solution with suitable physiologically tolerated acids. Suitable
15 physiologically tolerated acids are organic and inorganic acids, such as hydrochloric acid, sulfuric acid, phosphoric acid, acetic acid, citric acid, oxalic acid, malonic acid, salicylic acid, maleic acid, methane sulfonic acid, isothionic acid, lactic acid, gluconic acid, glucuronic acid, amidosulfuric acid, benzoic acid, tartaric acid and pamoic acid. Typically, such salt forms of the active compound will be provided or mixed prior to
20 use.

The pharmaceutical forms suitable for injectable use include sterile aqueous solutions or dispersions; formulations including sesame oil, peanut oil or aqueous propylene glycol; and sterile powders for the extemporaneous preparation of sterile
25 injectable solutions or dispersions. In all cases the form must be sterile and must be fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms, such as bacteria and fungi.

The active compounds may be formulated into a composition in a neutral or
30 salt form. Pharmaceutically acceptable salts, include the acid addition salts and which

are formed with inorganic acids such as, for example, hydrochloric or phosphoric acids, or such organic acids as acetic, oxalic, tartaric, mandelic, and the like.

The carrier can also be a solvent or dispersion medium containing, for
5 example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), suitable mixtures thereof, and vegetable oils. The proper fluidity can be maintained, for example, by the use of a coating, such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. The prevention of the action of microorganisms can be
10 brought about by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars or sodium chloride. Prolonged absorption of the injectable compositions can be brought about by the use in the compositions of agents delaying absorption, for example, aluminum
15 monostearate and gelatin.

Sterile injectable solutions are prepared by incorporating the active compounds in the required amount in the appropriate solvent with various of the other ingredients enumerated above, as required, followed by filtered sterilization.
20 Generally, dispersions are prepared by incorporating the various sterilized active ingredients into a sterile vehicle which contains the basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum-drying and freeze-drying techniques which yield a powder of
25 the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

In certain cases, the therapeutic formulations of the invention could also be prepared in forms suitable for topical administration, such as in creams and lotions.
30 These forms may be used for treating skin-associated diseases, such as various sarcomas.

Upon formulation, solutions will be administered in a manner compatible with the dosage formulation and in such amount as is therapeutically effective. The formulations are easily administered in a variety of dosage forms, such as the type of injectable solutions described above, with even drug release capsules and the like being employable.

For parenteral administration in an aqueous solution, for example, the solution should be suitably buffered if necessary and the liquid diluent first rendered isotonic with sufficient saline or glucose. These particular aqueous solutions are especially suitable for intravenous, intramuscular, subcutaneous and intraperitoneal administration. In this connection, sterile aqueous media which can be employed will be known to those of skill in the art in light of the present disclosure. For example, one dosage could be dissolved in 1 mL of isotonic NaCl solution and either added to 1000 mL of hypodermoclysis fluid or injected at the proposed site of infusion, (see for example, "Remington's Pharmaceutical Sciences" 15th Edition, pages 1035-1038 and 1570-1580). Some variation in dosage will necessarily occur depending on the condition of the subject being treated. The person responsible for administration will, in any event, determine the appropriate dose for the individual subject.

B. Oral Administration

In certain embodiments, active compounds may be administered orally. This is contemplated for agents which are generally resistant, or have been rendered resistant, to proteolysis by digestive enzymes. Such compounds are contemplated to include all those compounds, or drugs, that are available in tablet form from the manufacturer and derivatives and analogues thereof.

For oral administration, the active compounds may be administered, for example, with an inert diluent or with an assimilable edible carrier, or they may be enclosed in hard or soft shell gelatin capsule, or compressed into tablets, or incorporated directly with the food of the diet. For oral therapeutic administration, the

active compounds may be incorporated with excipients and used in the form of ingestible tablets, buccal tables, troches, capsules, elixirs, suspensions, syrups, wafers, and the like. Such compositions and preparations should contain at least 0.1% of active compound. The percentage of the compositions and preparations may, of course, be varied and may conveniently be between about 2 to about 60% of the weight of the unit. The amount of active compounds in such therapeutically useful compositions is such that a suitable dosage will be obtained.

The anthracycline analogs described in the present invention may be administered alone or with cyclodextrins, or substituted cyclodextrin formulations.

The tablets, troches, pills, capsules and the like may also contain the following: a binder, as gum tragacanth, acacia, cornstarch, or gelatin; excipients, such as dicalcium phosphate; a disintegrating agent, such as corn starch, potato starch, alginic acid and the like; a lubricant, such as magnesium stearate; and a sweetening agent, such as sucrose, lactose or saccharin may be added or a flavoring agent, such as peppermint, oil of wintergreen, or cherry flavoring. When the dosage unit form is a capsule, it may contain, in addition to materials of the above type, a liquid carrier. Various other materials may be present as coatings or to otherwise modify the physical form of the dosage unit. For instance, tablets, pills, or capsules may be coated with shellac, sugar or both. A syrup or elixir may contain the active compounds sucrose as a sweetening agent methyl and propylparabens as preservatives, a dye and flavoring, such as cherry or orange flavor. Of course, any material used in preparing any dosage unit form should be pharmaceutically pure and substantially non-toxic in the amounts employed. In addition, the active compounds may be incorporated into sustained-release preparation and formulations.

Upon formulation, the compounds will be administered in a manner compatible with the dosage formulation and in such amount as is therapeutically effective. The formulations are easily administered in a variety of dosage forms, such as those described below in specific examples.

THERAPIES

One of the major challenges in oncology today is the effective treatment of a given tumor. Tumors are often resistant to traditional therapies. Thus, a great deal of effort is being directed at finding efficacious treatment of cancer. One way of achieving this is by combining new drugs with the traditional therapies and is discussed below. In the context of the present invention, it is contemplated that therapies using the anthracycline analogs could be used in conjunction with surgery, chemotherapy, radiotherapy and indeed gene therapeutic intervention. It also may prove effective to combine the anthracycline analog based chemotherapy with antisense or immunotherapies directed at tumor marker.

"Effective amounts" are those amounts of a candidate substance effective to reproducibly inhibit decrease, reduce, inhibit or otherwise abrogate the growth of a cancer cell in an assay in comparison to levels in untreated cells.

A. Standard Therapies

a. **Chemotherapy:** A variety of chemical compounds, also described as "chemotherapeutic agents", function to induce DNA damage, are used to treat tumors. Chemotherapeutic agents contemplated to be of use, include, adriamycin, 5-fluorouracil (5FU), etoposide (VP-16), camptothecin, actinomycin-D, mitomycin, cisplatin (CDDP), hydrogen peroxide, carboplatin, procarbazine, mechlorethamine, cyclophosphamide, ifosfamide, melphalan, chlorambucil, bisulfan, nitrosurea, dactinomycin, daunorubicin, doxorubicin, bleomycin, plicomycin, tamoxifen, taxol, transplatinum, vincristin, vinblastin and methotrexate to mention a few.

Agents that damage DNA include compounds that interfere with DNA replication, mitosis and chromosomal segregation. Such chemotherapeutic compounds include adriamycin, also known as doxorubicin, etoposide, verapamil, podophyllotoxin, and the like. Widely used in a clinical setting for the treatment of neoplasms, these

compounds are administered through bolus injections intravenously at doses ranging from 25-75 mg/m² at 21 day intervals for adriamycin, to 35-50 mg/m² for etoposide intravenously or double the intravenous dose orally.

- 5 Agents that disrupt the synthesis and fidelity of nucleic acid precursors and subunits also lead to DNA damage. A number of such agents have been developed, particularly useful are agents that have undergone extensive testing and are readily available. 5-fluorouracil (5-FU), is one such agent that is preferentially used by neoplastic tissue, making it particularly useful for targeting neoplastic cells. Thus, 10 although quite toxic, 5-FU, is applicable with a wide range of carriers, including topical and even intravenous administrations with doses ranging from 3 to 15 mg/kg/day.

- Agents that directly cross-link nucleic acids, specifically DNA, are envisaged to facilitate DNA damage leading to a useful antineoplastic treatment. For example, 15 cisplatin, and other DNA alkylating agents may be used. Cisplatin has been widely used to treat cancer, with efficacious doses used in clinical applications of 20 mg/m² for 5 days every three weeks for a total of three courses. Cisplatin is not absorbed orally and must therefore be delivered *via* injection intravenously, subcutaneously, intratumorally or intraperitoneally.

20

- b. Radiotherapy:** Radiotherapeutic agents and factors include radiation and waves that induce DNA damage for example, γ -irradiation, X-rays, UV-irradiation, microwaves, electronic emissions, radioisotopes, and the like. Therapy may be achieved by irradiating the localized tumor site with the above described forms of radiations. It is 25 most likely that all of these factors effect a broad range of damage DNA, on the precursors of DNA, the replication and repair of DNA, and the assembly and maintenance of chromosomes.

- Dosage ranges for X-rays range from daily doses of 50 to 200 roentgens for 30 prolonged periods of time (3 to 4 weeks), to single doses of 2000 to 6000 roentgens.

Dosage ranges for radioisotopes vary widely, and depend on the half-life of the isotope, the strength and type of radiation emitted, and the uptake by the neoplastic cells.

c. **Surgery:** Surgical treatment for removal of the cancerous growth is generally a standard procedure for the treatment of tumors and cancers. This attempts to remove the entire cancerous growth. However, surgery is generally combined with chemotherapy and/or radiotherapy to ensure the destruction of any remaining neoplastic or malignant cells.

B. Combination Therapies

To kill cells, inhibit cell growth, inhibit metastasis, inhibit angiogenesis or otherwise reverse or reduce the malignant phenotype of tumor cells, the methods of standard therapy discussed above are generally insufficient as tumors are often resistant to several of these agents. Often combining a host of different treatment methods prove most effective in cancer therapy. Further, several AIDS afflicted patients have a higher risk of developing cancers. Combination therapy in these cases is required to treat AIDS as well as the cancer. Using the methods and compounds developed in the present invention, one would generally contact a "target" cell with an anthracycline analog synthesized in the present invention and at least one other agent. These compositions would be provided in a combined amount effective to kill or inhibit proliferation of the cell. This process may involve contacting the cells with the an anthracycline analog synthesized in the present invention and the other agent(s) or factor(s) at the same time. This may also be achieved by contacting the cell with a single composition or pharmacological formulation that includes both agents, or by contacting the cell with two distinct compositions or formulations, at the same time, wherein one composition includes the anthracycline analogs synthesized herein and the other includes the agent.

Alternatively, the anthracycline analog-based treatment may precede or follow the other agent treatment by intervals ranging from minutes to weeks. In embodiments where the other agent and anthracycline analog-based therapy are applied separately to

the cell, one would generally ensure that a significant period of time did not expire between the time of each delivery, such that the agent and anthracycline analog-based treatment would still be able to exert an advantageously combined effect on the cell. In such instances, it is contemplated that one would contact the cell with both modalities within about 12-24 h of each other and, more preferably, within about 6-12 h of each other, with a delay time of only about 12 h being most preferred. In some situations, it may be desirable to extend the time period for treatment significantly, however, where several days (2, 3, 4, 5, 6 or 7) to several weeks (1, 2, 3, 4, 5, 6, 7 or 8) lapse between the respective administrations.

It also is conceivable that more than one administration of either anthracycline analog-based treatment or the other agent will be desired. Various combinations may be employed, where anthracycline analog-based treatment is "A" and the other agent is "B", as exemplified below:

A/B/A B/A/B B/B/A A/A/B B/A/A A/B/B B/B/B/A B/B/A/B

A/A/B/B A/B/A/B A/B/B/A B/B/A/A B/A/B/A B/A/A/B B/B/B/A

A/A/A/B B/A/A/A A/B/A/A A/A/B/A A/B/B/B B/A/B/B B/B/A/B

Other combinations are also contemplated. Again, to achieve cell killing, both agents are delivered to a cell in a combined amount effective to kill the cell.

The invention also encompasses the use of a combination of one or more DNA damaging agents, whether chemotherapeutic compounds or radiotherapeutics as described in the section above, together with the anthracycline analog. The invention also contemplates the use of the anthracycline analog in combination with surgical removal of tumors to treat any remaining neoplastic or metastasized cells. Further, immunotherapy may be directed at tumor antigen markers that are found on the surface of tumor cells. The invention also contemplates the use of C-3' and C-4' substituted

anthracycline analog-based treatments in combination with gene therapy, directed toward a variety of oncogenes, such as, tumor markers, cell cycle controlling genes. For example, combining the anthracycline analog-based treatment and gene therapy towards oncogenes such as p53, p16, p21, Rb, APC, DCC, NF-1, NF-2, BCRA2, p16, FHIT, WT-1, MEN-I, MEN-II, BRCA1, VHL, FCC, MCC, *ras*, *myc*, *neu*, *raf*, *erb*, *src*, *fms*, *jun*, *trk*, *ret*, *gsp*, *hst*, *bcl* and *abl* mutations.

The other agent may be prepared and used as a combined therapeutic composition, or kit, by combining it with the anthracycline analog-based treatment, as described above. The skilled artisan is directed to "Remington's Pharmaceutical Sciences" 15th Edition, chapter 33, in particular pages 624-652. Some variation in dosage will necessarily occur depending on the condition of the subject being treated. The person responsible for administration will, in any event, determine the appropriate dose for the individual subject. Moreover, for human administration, preparations should meet sterility, pyrogenicity, general safety and purity standards as required by FDA Office of Biologics standards.

It is proposed that the regional delivery of anthracycline analog-based drugs described in the present invention to patients with tumors will be a very efficient method for delivering a therapeutically effective chemical to counteract the clinical disease. Similarly, other chemotherapeutics, radiotherapeutics, gene therapeutic agents may be directed to a particular, affected region of the subjects body. Alternatively, systemic delivery of anthracycline analog-based treatment and/or the agent may be appropriate in certain circumstances, for example, where extensive metastasis has occurred.

It also should be pointed out that any of the standard or other therapies may prove useful by themselves in treating a cancer. In this regard, reference to chemotherapeutics and anthracycline analog-based treatment in combination should also be read as a contemplation that these approaches may be employed separately.

When such combination therapy is employed for the treatment of a tumor, the cytotoxic agent may be administered at a dosage known in the art to be effective for treating the tumor. However, the anthracycline analog-based compounds may produce an additive or synergistic effect with a cytotoxic agent against a particular tumor.

5 Thus, when such combination therapy is used, the dosage of anthracycline analog-based drugs administered may be less than that administered when the cytotoxic agent is used alone. Similarly, for patients afflicted by AIDS, AZT/protease inhibitors will be used with anthracycline analog compounds, or other herein mentioned therapeutic agent(s). Again the dosage of anthracycline analog-based compounds or other
10 conjunctively utilized agent, may be altered to suit the AIDS treatment.

Preferably, the patient is treated with anthracycline analog-based compounds for about 1 to 14 days, preferably 4 to 14 days, prior to the beginning of therapy with a cytotoxic agent, and thereafter, on a daily basis during the course of such therapy.

15 Daily treatment with the anthracycline analogs can be continued for a period of, for example, 1 to 365 days after the last dose of the cytotoxic agent is administered.

* * *

The following examples are included to demonstrate preferred embodiments
20 of the invention. It should be appreciated by those of skill in the art that the techniques disclosed in the examples that follow represent techniques discovered by the inventor to function well in the practice of the invention, and thus can be considered to constitute preferred modes for its practice. However, those of skill in the art should, in light of the present disclosure, appreciate that many changes can be
25 made in the specific embodiments that are disclosed and still obtain a like or similar result without departing from the spirit and scope of the invention.

EXAMPLE 1**Synthesis of Exemplary Substituted Anthracyclines having C-3' Alkylated Anthracyclines Modified at Benzyl Ring****5 General Procedure for Making C-3' Alkylated Anthracyclines Modified at Benzyl Ring****A. Synthesis of 3'-N-nitro benzyl derivatives of daunorubicin and 4-demethoxy-daunorubicin:****10 1. Procedure A**

Daunorubicin (1 mmol), 4-nitrobenzyl bromide (1 mmol), and sodium carbonate (250 g) were dissolved in DMF (10 ml). Dichloromethane (10 ml) was added to this solution, and obtained mixture was stirred at room temperature until all substrate was converted into the product. (monitored by TLC). After reaction was completed the reaction mixture was diluted with dichloromethane (100 ml), and washed with water until neutral. The organic solution was dried over sodium sulfate, then drying agent was removed and solvent evaporated under diminished pressure, and product was purified by column chromatography (Silica Gel 60 Merck), using chloroform, chloroform/methanol 98:2, 95:5 as an eluent.

20

1.1 The following compounds were obtained from daunorubicin according the above procedure (Procedure A):

(i) WP 756 FIG.15

25 ¹H-n.m.r (CDCl₃) δ: 13.99, 13.40 (2s, 1H ea, 6,11-OH), 8.05 (d, 1H, J= 7.5 Hz, H-1), 7.93 (d, 1H, J = 8.1 Hz, H-2), 7.80 (dd, 1H, J = 8.2 Hz, H-3), 7.60-7.52 (m, 2H, H-aromatic), 7.44-7.40 (m, 2H, H-aromatic), 5.54 (d, J = 3.2 Hz, H-1'), 5.32 (, 1H, H-7), 4.67 (s, 1H, 9-OH), 4.13 - 4.06 (m, 4H, OMe and H-5'), 4.04 (d, 1H, J = 13.6 Hz, Ch2Ph), 3.95 (d, 1H, J = 13.6 Hz, Ch2Ph), 3.72 (, 1H, H-4'), 3.25 (dd, 1H, J = 18.1 Hz, J = 1.1 Hz, H=10), 2.97 (d, 1H, J = 18.9 Hz, H-10), 3.00 - 2.92 (m, 1H, H-

30

3'), 2.44 (s, 3 H, 14-CH₃), 2.38 (d, 1H, J = 14.82 Hz, H-8), 2.12 (dd, 1H, J = 14.8 Hz, J = 4.0 Hz, H-8), 1.86 -1.69 (m, 2H, H-2'a,e), 1.4 (d, 3H, J = 6.6 Hz, H-6').

Anal. Elem. Calcd. for: C₃₄H₃₅O₁₂N₂Cl x 1.5 H₂O : C: 56.24; H:5.27; N:3.86; Cl:4.88.

5 Found: C: 56.24; H:5.27; N: 3.86; Cl: 4.88.

(ii) WP 784 FIG.8

¹H-n.m.r (CDCl₃) δ: 13.98, 13.29 (2s, 1H ea, 6,11-OH), 8.14 (s, 1H, H-aromatic), 8.08 (d, 1H, J = 8.1 Hz, H-arom), 8.03 (d, 1H, J = 7.6 Hz, H-1), 7.78 (dd, 10 1H, J = J = 8.15 Hz H-2), 7.62 (d, 1H, J = 7.6 Hz, H-aromatic), 7.46 (dd, 1H, J = J = 7.9 Hz, H-aromatic), 7.39 (d, 1H, J = 8.4 Hz, H-aromatic) 5.53 (d, J = 3.0 Hz, H-1'), 5.3 (, 1H, H-7), 4.62 (s, 1H, 9-OH), 4.10 - 4.08 (m, 1H, H-5'), 4.08 (s, 3H, OMe), 3.9 (d, 1H, J = 13.6 Hz, Ch2Ph), 3.81 (d, 1H, J = 13.6 Hz, Ch2Ph), 3.67 (, 1H, H-4'), 3.22 (dd, 1H, J = 18.8 Hz, J = 1.8 Hz, H=10), 2.96 (d, 1H, J = 18.8 Hz, H-10), 2.96 - 15 2.92 (m, 1H, H-3'), 2.41 (s, 3 H, 14-CH₃), 2.36 (d, 1H, J = 14.8 Hz, H-8), 2.11 (dd, 1H, J = 14.8 Hz, J = 2.1 Hz, H-8), 1.82 -1.75 (m, 2H, H-2'a,e), 1.37 (d, 3H, J = 6.5 Hz, H-6').

Anal. Elem. Calcd. for: C₃₄H₃₅O₁₂N₂Cl x 1.5 H₂O : C: 56.24; H:5.27; N:3.86; Cl:4.88.

20 Found: C: 55.94; H:5.27; N: 3.78; Cl: 4.92.

(iii) WP 755 FIG.16

¹H-n.m.r (CDCl₃) δ: 14.02, 13.30 (2s, 1H ea, 6,11-OH), 8.17 (d, 1H, J = 8.6 Hz, H-aromatic), 8.05 (d, 1H, J = 7.8 Hz, H-1), 7.81 (dd, 1H, J = J = 7.8 Hz, H-2), 25 7.49 - 7.41 (m, 4H, H-3 and H-aromatic), 5.54 (s, 1H, H-1'), 5.32 (, 1H, H-7), 4.64 (s, 1H, 9-OH), 4.11 (s, 3H, OMe), 4.09 (q, 1H, J = 6.5 Hz, H-5'), 3.93 (d, 1H, J = 14 Hz, Ch2Ph), 3.83 (d, 1H, J = 14 Hz, Ch2Ph), 3.67 (, 1H, H-4'), 3.25 (d, 1H, J = 18.9 Hz, H=10), 2.98 (d, 1H, J = 18.9 Hz, H-10), 2.96 - 2.92 (m, 1H, H-3'), 2.43 (s, 3 H, 14-CH₃), 2.38 (d, 1H, J = 15.2 Hz, H-8), 2.13 (dd, 1H, J = 15.2 Hz, J = 4.13 Hz, H-8), 30 1.82 -1.78 (m, 2H, H-2'a,e), 1.38 (d, 3H, J = 6.6 Hz, H-6').

Anal. Elem. Calcd. for: $C_{34}H_{35}O_{12}N_2Cl \times H_2O$: C: 56.95; H:5.2; N:3.91;
Cl:4.94.

Found: C: 56.99; H:5.19; N: 3.81; Cl: 4.89.

5 **1.2 The following compounds were obtained from 4-demethoxy-daunorubicin using Procedure A:**

(iv) **WP 790 FIG.4**

1H -n.m.r ($CDCl_3$) δ : 13.58, 13.34 (2s, 1H ea, 6,11-OH), 8.36 - 8.32 (m, 2H, H-1,4), 8.90 (d, 1H, J = 8.2 Hz, H-arom), 7.85 - 7.80 (m, 2H, H-2,3), 7.57 (dd, 1H, J = J = 7.5 Hz, H-aromatic), 7.51 (d, 1H, J = 6.5 Hz, H-aromatic), 7.40 (dd, 1H, J = J = 7.1 Hz, H-aromatic), 5.50 (d, 1H, J = 3.6 Hz, H-1'), 5.28 (, 1H, H-7), 4.66 (s, 1H, 9-OH), 4.10 (m, 1H, J = 6.6 Hz, H-5'), 4.03 (d, 1H, J = 14.7 Hz, Ch2Ph), 3.94 (d, 1H, J = 14.7 Hz, Ch2Ph), 3.70 (, 1H, H-4'), 3.25 (dd, 1H, J = 19 Hz, J = 1.7 Hz, H=10),
15 3.00 (d, 1H, J = 19 Hz, H-10), 2.96 - 2.91 (m, 1H, H-3'), 2.40 (s, 3 H, 14-CH₃), 2.38 (d, 1H, J = 14.9 Hz, H-8), 2.10 (dd, 1H, J = 14.9 Hz, J = 4.1 Hz, H-8), 1.82 (ddd, 1H, J = 13.3 Hz, J = 4.2 Hz, H-2'a), 1.74 (dd, 1H, J = 13.3 Hz, J = 4.9 Hz, H-2'e), 1.38 (d, 3H, J = 6.6 Hz, H-6').

Anal. Elem. Calcd. for: $C_{33}H_{33}O_{11}N_2Cl \times 1.5 H_2O$: C: 56.94; H:5.21; N:4.02;
20 Cl:5.09.

Found: C: 57.00; H:5.13; N: 3.98; Cl: 5.16.

(v) **WP 774 FIG.12**

1H -n.m.r ($CDCl_3$) δ : 13.59, 13.35 (2s, 1H ea, 6,11-OH), 8.37 - 8.32 (m, 2H, H-1,4), 8.13 (d, 1H, J = 8.2 Hz, H-arom), 7.86 - 7.82 (m, 2H, H-2,3), 7.44 (d, 2H, H-aromatic), 5.50 (s, 1H, H-1'), 5.28 (, 1H, H-7), 4.62 (s, 1H, 9-OH), 4.09 (m, 1H, J = 6.4 Hz, H-5'), 3.91 (d, 1H, J = 14.7 Hz, Ch2Ph), 3.83 (d, 1H, J = 14.7 Hz, Ch2Ph), 3.66 (, 1H, H-4'), 3.25 (dd, 1H, J = 19 Hz, J = 1.8 Hz, H=10), 2.98 (d, 1H, J = 19 Hz, H-10), 2.96 - 2.93 (m, 1H, H-3'), 2.40 (s, 3 H, 14-CH₃), 2.37 (d, 1H, J = 13.0 Hz, H-8), 2.11 (dd, 1H, J = 13.0 Hz, J = 4.0 Hz, H-8), 1.84- 1.76 (m, 2H, H-2'a,e), 1.28 (d, 3H, J = 6.4 Hz, H-6').
30

Anal. Elem. Calcd. for: $C_{33}H_{33}O_{11}N_2Cl \times 0.5 H_2O$: C: 57.15; H:4.87; N:4.04; Cl:5.11.

Found: C: 57.15; H:5.08; N: 4.01; Cl: 5.11.

- 5 **1.3 The following compound WP787 was obtained when 4-amino-daunorubicin using benzyl bromide according to procedure A:**

(vi) WP 787 FIG.5

1H -n.m.r ($CDCl_3$) δ : 13.98, 13.29 (2s, 1H ea, 6,11-OH), 8.02 (d, 1H, J = 7.3
10 Hz, H-1), 7.81 (dd, 1H, J = 7.8 Hz, H-2), 7.40 (d, 1H, J = 8.1 Hz, H-3), 7.35 (s, 4H, H-aromatic), 7.31 - 7.26 (m, 1H, H-aromatic), 5.42 (d, 1H, J = 4.0 Hz, H-1'), 5.25 (, 1H, H-7), 4.64 (s, 1H, 9-OH), 4.14 (q, 1H, J = 6.6 Hz, H-5'), 4.11 (d, 1H, J = 12.7 Hz, Ch2Ph), 4.07 (s, 3H, OMe), 3.77 (d, 1H, J = 12.7 Hz, Ch2Ph), 3.72 - 3.68 (m, 1H, H-3'), 3.22 (dd, 1H, J = 18.8 Hz, J = 1.8 Hz, H=10), 2.94 (d, 1H, J = 18.8 Hz, H-10),
15 2.76 (d, 1H, J = 3.9 Hz, H-4'), 2.41 (s, 3 H, 14-CH₃), 2.29 (d, 1H, J = 14.7 Hz, H-8).

Anal. Elem. Calcd. for: $C_{34}H_{36}O_{10}NCl$: C: 62.43; H:5.55; N:2.14; Cl:5.42.

(vii) WP 831 FIG.2 The following compound was obtained from daunorubicin using 4-picolyl chloride according to Procedure A:

1H -n.m.r ($CDCl_3$) δ : 13.97, 13.37 (2s, 1H ea, 6,11-OH), 8.50 (d, 2H, H-arom), 8.02 (d, 1H, J = 7.5Hz, H-1), 7.78 (dd, 1H, J = 7.8 Hz, H-2), 7.39 (d, 1H, H-3), 7.18 (m, 2H, H-aromatic), 5.51 (d, 1H, J = 3.8 Hz, H-1'), 5.28 (, 1H, H-7), 4.62 (s, 1H, 9-OH), 4.08 (m, 4H, H-5' and OMe), 3.81 (d, 1H, J = 14.2 Hz, Ch2Ph), 3.72 (d, 1H, J = 14.2 Hz, Ch2Ph), 3.64 (, 1H, H-4'), 3.37 (dd, 1H, J = 19 Hz, J = 1.5 Hz,
25 H=10), 2.94 (d, 1H, J = 19 Hz, H-10), 2.96 - 2.93 (m, 1H, H-3'), 2.41 (s, 3 H, 14-CH₃), 2.36 (d, 1H, J = 14.8 Hz, H-8), 2.10 (dd, 1H, J = 14.8 Hz, J = 4.1 Hz, H-8), 1.83-1.75 (m, 2H, H-2'a,e), 1.36 (d, 3H, J = 6.4 Hz, H-6').

Anal. Elem. Calcd. for: $C_{33}H_{36}O_{10}N_2Cl_2 \times 3 H_2O$: C: 53.16; H:5.68; N:3.76; Cl:9.51.

30 Found: C: 53.70; H:5.59; N: 3.67; Cl: 9.36.

1.4 The following compound WP 786 was obtained according procedure A using 4-fluorobenzyl bromide as alkylating agent:

(viii) WP 786 FIG.6

5 ¹H-n.m.r (CDCl₃) δ: 13.98, 13.30 (2s, 1H ea, 6,11-OH), 8.03 (d, 1H, J = 7.6 Hz, H-1), 7.78 (dd, 1H, J = 7.6 Hz, H-2), 7.39 (d, 1H, J = 7.8 Hz, H-3), 7.23 - 7.20 (m, 2H, H-aromatic), 7.00 - 6.95 (m, 2H, H-aromatic), 5.51 (d, 1H, J = 3.7 Hz, H-1'), 5.3 (, 1H, H-7), 4.69 (s, 1H, 9-OH), 4.08 (s, 3H, OMe), 4.06 (q, 1H, J = 6.6 Hz, H-5'), 3.96 (d, 1H, J = 12.7 Hz, Ch2Ph), 3.64 (d, 1H, J = 12.7 Hz, Ch2Ph), 3.64 (, 1H, H-4'), 3.23 (dd, 1H, J = 18.9 Hz, J = 1.9 Hz, H=10), 2.96 (d, 1H, J = 18.9 Hz, H-10), 2.96 - 2.93 (m, 1H, H-3'), 2.42 (s, 3 H, 14-CH₃), 2.37 (d, 1H, J = 13.8 Hz, H-8), 2.09 (dd, 1H, J = 13.8 Hz, J = 4.1 Hz, H-8), 1.78 (ddd, 1H, J = 13.1 Hz, J = 4.0 Hz, H-2'a), 1.67 (dd, 1H, J = 13.1 Hz, J = 4.9 Hz, H-2'e), 1.37 (d, 3H, J = 6.6 Hz, H-6').

10

15 **B. Synthesis of 3'-N-nitro benzyl derivatives of doxorubicin and 4-demethoxydoxorubicin:**

2. Procedure B

Doxorubicin having hydroxyl at C-14 protected by silylation (1 mmol), 4-nitrobenzyl bromide (1 mmol), and sodium carbonate (250 mg) were dissolved in
20 DMF (10 ml). Dichloromethane (10 ml) was added to this solution, and obtained mixture was stirred at room temperature until all substrate was converted into the product. (monitored by TLC control). After reaction was completed the reaction mixture was diluted with di chloromethane (100 ml), and washed with water until neutral. The organic solution was dried over sodium sulfate, then drying agent and
25 solvent were removed, and product was purified by column chromatography (Silica Gel 60 Merck), using chloroform, chloroform/methanol 98:2, as eluent.

Obtained compounds was dissolved in THF (6 ml) and to this 1 N water solution of HCl (9 ml) was added. The mixture was stirred at room temp. overnight , then it was diluted with chloroform, and neutralized with 10 % water solution of
30 potassium bicarbonate. Organic solution was dried over sodium sulfate. Drying agent was filtered off, and solvent was evaporated to dryness. Crude product was purified

by column chromatography (Silica Gel 60 Merck), using chloroform/methanol 98:2, 95:5 as eluent.

The following compounds were obtained according to this Procedure B:

5

(ix) WP 780 FIG. 9

¹H-n.m.r (CDCl₃) δ: 13.96, 13.26 (2s, 1H ea, 6,11-OH), 8.03 (d, 1H, J = 7.7 Hz, H-aromatic), 7.91 (d, 1H, J = 8.1 Hz, H-1), 7.79 (dd, 1H, J = 8.1 Hz, H-2), 7.56 (dd, 1H, J = J = 7.8 Hz H-aromatic), 7.50 (d, 1H, J = 6.8 Hz, H-aromatic), 7.40-7.30 (m, 2H, H-3 and H-aromatic), 5.52 (d, 1H, J = 3.6 Hz, H-1'), 5.3 (, 1H, H-7), 4.77 (s, 10 2H, 14-CH₂), 4.75 (s, 1H, 9-OH), 4.08 (s, 3H, OMe), 4.03 (d, 1H, J = 13.6 Hz, Ch2Ph), 3.99 (q, 1H, J = 6.6 Hz, H-5') 3.93 (d, 1H, J = 13.6 Hz, Ch2Ph), 3.70 (, 1H, H-4'), 3.27 (dd, 1H, J = 18.9 Hz, J = 1.7 Hz, H=10), 3.03 (d, 1H, J = 18.9 Hz, H-10), 2.89 (m, 1H, H-3'), 2.36 (d, 1H, J = 14.8 Hz, H-8), 2.16 (dd, 1H, J = 14.8 Hz, J = 2.9 15 Hz, H-8), 1.84 (ddd, 1H, J = 13.1 Hz, J = 4.0 Hz, H-2'a), 1.68 (dd, 1H, J = 13.1 Hz, J = 4.6 Hz, H-2'e), 1.39 (d, 3H, J = 6.6 Hz, H-6').

Anal. Elem. Calcd. for: C₃₄H₃₅O₁₃N₂Cl x 2 H₂O: C: 54.37; H:5.23; N:3.73; Cl:4.72.

Found: C: 54.49; H:5.13; N: 3.70; Cl: 4.76.

20

(x) WP 765 FIG.34

¹H-n.m.r (CDCl₃) δ: 13.96, 13.24 (2s, 1H ea, 6,11-OH), 8.13 (d, 1H, J = 7.6 Hz, H-1), 8.01 (d, 1H, J = 7.5 Hz, H-2), 7.71 (d, 1H, J = 7.8 Hz, H-3), 7.45 - 7.40 (m, 4H, H-aromatic), 5.52 (, 1H, H-1'), 5.31 (, 1H, H-7), 4.75 (s, 2H, 14-CH₂), 4.70 (s, 25 1H, 9-OH), 4.00 (s, 3H, OMe), 3.99 (q, 1H, J = 6.6 Hz, H-5') 3.91 (d, 1H, J = 14.0 Hz, Ch2Ph), 3.82 (d, 1H, J = 14.0 Hz, Ch2Ph), 3.66 (, 1H, H-4'), 3.24 (d, 1H, J = 18.8 Hz, H=10), 3.00 (d, 1H, J = 18.8 Hz, H-10), 2.90-2.88 (m, 1H, H-3'), 2.37 (d, 1H, J = 14.8 Hz, H-8), 2.15 (dd, 1H, J = 14.8 Hz, J = 4.0 Hz, H-8), 1.8 - 1.6 (m, 2H, H-2'a,e), 1.39 (d, 3H, J = 6.5 Hz, H-6').

30

Anal. Elem. Calcd. for: C₃₄H₃₅O₁₃N₂Cl x 0.5 H₂O: C: 59.39; H:5.13; N:4.07.

Found: C: 59.00; H:5.15; N: 4.07

2.1 The following compound WP778 was obtained from 14-O-silylated 4-demethoxydoxorubicin following compound was prepared according

Procedure B:

(xi) WP 778 FIG.10

¹H-n.m.r (CDCl₃) δ: 13.57, 13.30 (2s, 1H ea, 6,11-OH), 8.36 - 8.32 (m, 2H, H-1,4), 8.13 (d, 2H, J = 8.7 Hz, H-arom), 7.88 - 7.83 (m, 2H, H-2,3), 7.44 (d, 2H, J = 8.7 Hz, H-aromatic), 5.52 (d, 1H, J = 3.3 Hz, H-1'), 5.30 (, 1H, H-7), 4.76 (, 2H, 14-CH₂), 4.73 (s, 1H, 9-OH), 4.01 (m, 1H, J = 6.6 Hz, H-5'), 3.93 (d, 1H, J = 14.1 Hz, Ch2Ph), 3.83 (d, 1H, J = 14.1 Hz, Ch2Ph), 3.68 (, 1H, H-4'), 3.29 (dd, 1H, J = 19 Hz, J = 1.8 Hz, H=10), 3.06 (d, 1H, J = 19 Hz, H-10), 2.96 - 2.88 (m, 1H, H-3'), 2.38 (d, 1H, J = 14.7 Hz, H-8), 2.15 (dd, 1H, J = 14.7 Hz, J = 4.0 Hz, H-8), 1.88 - 1.74 (m, 1H, H-2'a,e), 1.29 (d, 3H, J = 6.6 Hz, H-6').

C. Synthesis of 3'-N-amino-substituted benzyl derivatives of daunorubicin and 4-methoxy-daunorubicin:

3. Procedure C

Products obtained according to the procedure A and B (0.1 mmol) were dissolved in the mixture of dichloromethane and methyl alcohol (1:1 v/v) (10 ml), then stannous chloride (II) (1.1 g) was added and the mixture was stirred at room temp. until all substrate disappeared (TLC control). The reaction mixture was then diluted with chloroform, and saturated solution of sodium bicarbonate (100 ml) was added. The reaction mixture was stirred at room temp. for 3 hr, then the inorganic salts were filtered off, and washed with chloroform. Organic solution was washed with water until neutral, then dried over Na₂SO₄. Drying agent and solvent were removed, and product was purified by column chromatography (Silica Gel 60 Merck), using chloroform/methanol 98:2, 95:5, 9:1 as an eluent.

Following compounds were prepared according to the Procedure C:

(xii) WP 758 FIG. 13

¹H-n.m.r (CDCl₃) δ: 13.98, 13.28 (2s, 1H ea, 6,11-OH), 8.04 (d, 1H, J= 8.0 Hz, H-1), 7.78 (dd, 1H, J = 8.0 Hz, H-2), 7.38 (d, 1H, J = 8.0 Hz, H-3), 7.05 (td, 1H, J = 7.6 Hz, J=1.3 Hz, H-aromatic), 6.97 (dd, 2H, J = 7.8 Hz, J=1.4 Hz, H-aromatic),
5 6.63 (m, 2H, H-aromatic), 5.51 (, 1H, H-1'), 5.28 (, 1H, H-7), 4.08 (s, 3H, OMe), 4.13 - 4.08 (q, 1H, J = 6.5 Hz, H-5'), 3.82 (d, 1H, J = 12.5 Hz, Ch2Ph), 3.73 (, 1H, H-4'), 3.71 (d, 1H, J = 12.5 Hz, Ch2Ph), 3.23 (dd, 1H, J = 18.9 Hz, J = 1.7 Hz, H=10), 2.95 (d, 1H, J = 18.9 Hz, H-10), 2.99 - 2.93 (m, 1H, H-3'), 2.44 (s, 3H, 14-CH₃), 2.7 (, 2H, NH, OH), 2.34 (d, 1H, J = 14.9 Hz, H-8), 2.10 (dd, 1H, J = 14.9 Hz,
10 J = 4.1 Hz, H-8), 1.81 - 1.74 (m, 2H, H-2'a,e), 1.37 (d, 3H, J = 6.6 Hz, H-6').

Anal. Elem. Calcd. for: C₃₄H₃₈O₁₀N₂Cl₂ : C: 57.88; H:5.43; N:3.97; Cl:10.05.

Found: C: 57.85; H:5.44; N: 3.93; Cl: 9.96

(xiii) WP 757 FIG.14

¹H-n.m.r (CDCl₃) δ: 14.00 (, 1H, H-6 or H-11), 8.06 (d, 1H, J= 7.7 Hz, H-1), 7.80 (dd, 1H, J = 8.1 Hz, H-2), 7.41 (d, 1H, J = 7.7 Hz, H-3), 7.05 (d, 2H, J = 8.3 Hz, H-aromatic), 6.62 (d, 2H, J = 8.3 Hz, H-aromatic), 5.53 (d, 1H, J= 3.4 Hz, H-1'), 5.31 (, 1H, H-7), 4.75 (s, 1H, 9-OH), 4.10 (s, 3H, OMe), 4.10 (q, 1H, J = 6.5 Hz, H-5'), 3.71 (d, 1H, J = 15.0 Hz, Ch2Ph), 3.7 (, 1H, H-4'), 3.58 (d, 1H, J = 15.0 Hz, Ch2Ph), 3.28 (dd, 1H, J = 19.0 Hz, J = 1.0 Hz, H=10), 3.0 (d, 1H, J = 19.0 Hz, H-10),
20 3.02 - 2.92 (m, 1H, H-3'), 2.45 (s, 3H, 14-CH₃), 2.40 (d, 1H, J = 14.9 Hz, H-8), 2.11 (dd, 1H, J = 14.4 Hz, J = 4.0 Hz, H-8), 1.78 (ddd, 1H, J = 12.5 Hz, J = 8.6 Hz, J = 3.8 Hz, H-2'a), 1.63 (dd, 1H, J = 12.5 Hz, J = 5.2 Hz, H-2'e) 1.40 (d, 3H, J = 6.5 Hz, H-6').

25 Anal. Elem. Calcd. for: C₃₄H₃₆O₁₀N₃ : C: 64.55; H:5.74; N: 4.43.

Found: C: 64.27; H:5.90; N: 4.28

(xiv) WP 791 FIG.3

¹H-n.m.r (CDCl₃) δ: 8.37 (m, 2H, H-1,4), 7.87 - 7.83 (m, 2H, H-2,3), 7.04 (dd, 1H, J = 8.0 Hz, H-aromatic), 6.97 (d, 2H, J = 7.1 Hz, H-aromatic), 6.66 - 6.62 (m, 2H, H-aromatic), 5.51 (d, 1H, J= 3.5 Hz, H-1'), 5.29 (, 1H, H-7), 4.67 (s, 1H, 9-OH),
30

4.11 (q, 1H, $J = 6.5$ Hz, H-5'), 3.80 (d, 1H, $J = 12.3$ Hz, Ch2Ph), 3.7 (, 1H, H-4'),
3.68 (d, 1H, $J = 12.3$ Hz, Ch2Ph), 3.27 (dd, 1H, $J = 19.0$ Hz, $J = 1.4$ Hz, H=10), 3.0
(d, 1H, $J = 19.0$ Hz, H-10), 2.97 - 2.88 (m, 1H, H-3'), 2.40 (s, 3 H, 14-CH₃), 2.37
(d, 1H, $J = 15.2$ Hz, H-8), 2.11 (dd, 1H, $J = 15.2$ Hz, $J = 4.0$ Hz, H-8), 1.89 - 1.73 (m,
5 2H, H-2'a,e), 1.35 (d, 3H, $J = 6.5$ Hz, H-6').

(xv) WP 764 FIG.33

¹H-n.m.r (DMSO-d₆) δ : 7.91-7.89 (m, 2H, H-1,2), 7.62 -7.60 (m, 1H, H-3),
7.00 (d, 2H, $J = 8.2$ Hz, H-arom), 6.5 (d, 2H, $J = 8.2$ Hz, H-aromatic), 5.19 (d, 1H, $J =$
10 3.4 Hz, H-1'), 4.89 (, 1H, H-7), 4.38 (s, 1H, 9-OH), 4.14 (q, 1H, $J = 6.5$ Hz, H-5'),
3.98 (s, 3H, OMe), 3.81 (m, 1H, H-4'), 3.78 (s, 2H, H-aromatic), 2.96 (d, 1H, $J =$
18.0 Hz, H=10), 2.88 (d, 1H, $J = 19.0$ Hz, H-10), 2.57 (m, 1H, H-3'), 2.25 (s, 3 H,
14-CH₃), 2.14 (dd, 1H, $J = 14.3$ Hz, $J = 3.0$ Hz, H-8), 2.07 (dd, 1H, $J = 14.3$ Hz, $J =$
5.0 Hz, H-8), 1.66 (ddd, 1H, $J = 14.2$ Hz, $J = 13.0$ Hz, $J = 3.5$ Hz, H-2'a), 1.55 (dd,
15 1H, $J = 13.0$ Hz, $J = 4.4$ Hz, H-2'e) 1.15 (d, 3H, $J = 6.5$ Hz, H-6').

EXAMPLE 2**Synthesis of Exemplary C-4'-Alkyl-Aromatic Ring Anthracyclines**

5 General procedures for making anthracycline analogs with C-4'-substituted sugars.

A. Synthesis of 4'-O-benzylated anthracyclines

1. Procedure A: Coupling and deacetylation.

10 In the one flask a mixture of aglycon (1 mmol), HgBr₂ (0.7 g), HgO (2.8 g) and molecular sieves 4 Å (1 g) in dichloromethane (50 ml) was prepared and stirred at room temperature. At the same time, in the second flask, a trimethylsilyl bromide (3 mmol) was added to the solution of 1-O-silylated 3-azido-sugar (2 mmol) in dichloromethane (8 mL). Such reaction mixture was stirred at RT and its progress was monitored by TLC. (Toluene/ acetone = 8:1). After disappearance of starting
15 sugar, solvent was removed under diminished pressure. Then hexanes (10 mL) were added to the residue and the mixture was evaporated. Such addition and removal of hexanes were repeated three times. Subsequently, the resulting dry residue was dissolved in the dichloromethane (10 mL). This solution was added in three portions to the first flask containing earlier prepared solution of aglycone and mercuric salts.
20 The final reaction mixture was stirred at RT for 10 min. (progress of the reaction was monitored by TLC), then it was diluted with dichloromethane (50 mL), and filtered through Celite. Filtrate was washed with 10% water soln. of KI, then twice with water, and it was dried over Na₂SO₄. The crude product was purified by column chromatography and subsequently deacetylated in the mixture of methylene chloride
25 and methanol (2:1, v/v) (50 mL), using solid potassium carbonate (2 g). The reaction mixture was stirred at RT until all substrate disappeared. (progress of the reaction was monitored by TLC). The mixture was diluted with dichloromethane (100 mL), then 1N water solution of HCl (15 mL), and water (50 mL) were added. The layers were separated. The organic layer was washed with water until neutral, then with
30 brine, and it was dried over Na₂SO₄. The product was purified by column chromatography.

2. Procedure B: Reduction and desilylation.

The coupling product containing azido group at the C-3' position (1 mmol) was dissolved in THF (8 mL). Triphenyl phosphine (2 mmol) was added to this solution, and reaction mixture was stirred at room temperature overnight. Progress of the reaction was monitored by TLC until disappearance of substrate. Then the reaction mixture was diluted with chloroform (50 mL) and 2N NH_3 in methanol (2 mL) was added to this solution. Reaction mixture was stirred for 1 h, then the solution was washed with water until neutral, then with brine, and dried over Na_2SO_4 . Obtained crude product was purified by column chromatography. Anthracycline antibiotic with hydroxyl protected at 14 (1 mmol) was dissolved in THF (10 mL), then 1N HCl (15 mL) was added and the mixture was stirred at room temperature until all substrate disappeared as judge by TLC. The mixture was diluted with chloroform (20 mL). The layers were separated and the water solution was washed with chloroform until organic layer was colorless, and then saturated solution of NaHCO_3 was added to adjust pH to 8-9. Basic solution was extracted with chloroform contained 5% of methanol. Combined extracts were washed with brine and dried over dry Na_2SO_4 . The solvent was removed under diminished pressure and dry product was dissolved in methanol (4 mL). 1N solution of HCl in methanol (1.5 mL) followed by diethyl ether were added to this solution. Obtained solid hydrochloride was filtered off and washed with diethyl ether to remove excess of acid. When the filtrate was neutral, solid hydrochloride was precipitated from methanol/diethyl ether to give analytically pure product.

(i) WP 750A FIG. 36

3'-Deamino-3'-azido-

4'-O-benzyl-14-O-tert-butyl-diphenylsilyl-4'-epi doxorubicin.

Product was obtained according to the A. Yield 85%.

$\text{nmr}(\text{CDCl}_3)$ δ : 14.00, 13.19 (2s, 1H each, 6-OH, 11-OH), 8.02 (dd, 1H, $J = 8.47$ Hz, $J = 0.98$ Hz, H-1), 7.77 (dd, 1H, $J = J = 8.39$ Hz, H-2), 7.70-7.76 (m, 4H), 7.45-7.30 (m, 12H, H-3), 5.39 (d, 1H, $J = 3.72$ Hz, H-1'), 5.17 (bs, 1H, H-7), 4.89 (s,

2H, 14-CH₂), 4.81 (d, 1H, $J = 10.60$ Hz, CH₂Ph), 4.60 (d, 1H, $J = 10.60$ Hz, CH₂Ph), 4.16 (s, 1H, 9-OH), 4.07 (s, 3H, OMe), 3.71 (dq, 1H, $J = 9.4$ Hz, $J = 6.2$ Hz, H-5'), 3.56 (ddd, 1H, $J = 12.53$ Hz, $J = 9.38$ Hz, $J = 4.91$ Hz, H-3'), 2.97 (dd, 1H, $J = 9.4$ Hz, H-4'), 2.97 (dd, 1H, $J = 18.94$ Hz, $J = 1.24$ Hz, H-10), 2.76 (d, 1H, $J = 18.94$ Hz, H-10), 2.1-2.0 (m, 3H, H-8a,e, and H-2'e), 1.64 (ddd, 1H, $J = 13.28$ Hz, $J = 13.28$ Hz, $J = 4.07$ Hz, H-2'a), 1.18 (d, 3H, $J = 6.2$ Hz, H-6'), 1.12 (s, 9H, t-Bu).

(ii) WP 750 FIG.21

4'-O-Benzyl-4'epi-doxorubicin

10 **3'-Deamino-3'-azido-4'-O-benzyl-14-O-tert butyldiphenylsilyl-doxorubicin.**

Product was obtained according to the Procedure (A), yield 90%.

nmr(CDC13) δ : 13.90, 13.18 (2s, 1H each, 6-OH, 11-H), 8.01 (dd, 1H, $J = 7.49$ Hz, $J = 0.9$ Hz, H-1), 7.76 (dd, 1H, $J = 8.11$ Hz, H-2), 7.72-7.69 (m, 4H), 15 7.47-7.33 (m, 12H, H-3 and H-aromatic), 5.49 (d, 1H, $J = 3.26$ Hz, H-1'), 5.16 (, 1H, H-7), 4.83 (ABq, 2H, 14-CH₂), 4.83 (d, 1H, $J = 11.14$ Hz, CH₂Ph), 4.57 (d, 1H, $J = 11.28$ Hz, CH₂Ph), 4.27 (s, 1H, 9-OH), 4.05 (s, 3H, OMe), 3.75 (q, 1H $J = 6.4$ Hz, H-5'), 3.42-3.36 (m, 2H, H-4', H-3), 3.00 (d, 1H, $J = 19.09$ Hz, H-10), 2.80 (d, 1H, $J = 19.03$ Hz, H-10), 2.17 (ddd, 1H, $J = 13.06$ Hz, $J = 3.86$ Hz, H-2'a), 2.1-1.95 (m, 20 2H, H-8a,e), 1.81 (dd, 1H, $J = 13.06$ Hz, $J = 3.81$ Hz, H-2'e), 1.09 (s, 9H, t-Bu), 0.975 (d, 3H, $J = 6.4$ Hz, H-6').

(iii) WP 744 FIG.22

4'-O-Benzyl-doxorubicin

25 Product was obtained according to the procedure B., yield 40

n.m.r (DMSO) δ : 14.00, 13.22 (2s, 1H each, 6-OH, 11-OH), 8.15 (, 2H, NH₂), 7.91-7.89 (m, 2H, H-1,2), 7.65-7.63 (m, 2H, H-3), 7.47-7.30 (m, 5H, H-aromatic), 5.45 (s, 1H, 14-OH), 5.40 (d, 1H, $J = 2.48$ Hz, H-1'), 4.9 (, 1H, H-7), 4.85 (, 1H, 9-OH), 4.79 (d, 1H, $J = 11.13$ Hz, Ch₂Ph), 4.57 (s, 2H, 14-CH₂), 4.56 (d, 1H, $J = 11.13$ Hz, Ch₂Ph), 4.23 (q, 1H, $J = 6.51$ Hz, H-5'), 3.97 (s, 3H, OMe), 3.72 (, 1H, H-4'), 3.52-3.45 (m, 1H, H-3'), 2.97 (d, 1H, $J = 18.24$ Hz, H-10), 2.85 (d, 1H, $J = 18.24$

Hz, H-10), 2.14-2.05 (m, 2H, H-8), 1.87 (ddd, 1H, $J = 12.63$ Hz, $J = 12.63$ Hz, $J = 3.5$ Hz, H-2'a), 1.71 (dd, 1H, $J = 12.63$ Hz, $J = 4.38$ Hz, H-2'e), 1.12 (d, 3H, $J = 6.48$ Hz, H-6').

5 (iv) WP 783 FIG.20

4'-O-Benzyl-4-demethoxydaunorubicin

The mixture of aglycone (1 mmol) HgBr₂ (0.7 g), HgO (2.8 g) and molecular sieves 4 Å (1 g) in methylene chloride (50 mL) was prepared and stirred at room temperature. Trimethylsilyl bromide (3 mmol) was added to the solution of sugar
10 (2 mmol) in dichloromethane (8 mL). The obtained mixture was stirred at room temp., progress of the reaction was monitored by TLC (Toluene/acetone = 8:1), until all substrate disappeared from reaction mixture, the solvent was removed under diminished pressure. Hexanes (10 mL) were added to the residue. Solvent was removed again, and the second portion of hexanes was added and then removed from
15 the mixture. The addition and removal of hexanes was repeated three times then dry residue was dissolved in dichloromethane (10 mL). Obtained solution was added in three portions to the earlier prepared solution of aglycone and mercuric salts. The mixture was stirred at room temperature for 10 min. (reaction was controlled by TLC, then it was diluted with methylene chloride (50 mL), and filtered through Celite.
20 Filtrate was washed with 10% water soln. of KI, then twice with water, and it was dried over Na₂SO₄. The crude product was purified by column chromatography (Eluent: toluene, toluene:acetone = 100:1, 98:2). Product contained azido group in position 3' (1 mmol) was dissolved in THF (8 mL). Triphenyl phosphine (2 mmol) was added to this solution, and obtained mixture was stirred at room temperature
25 overnight. Progress of the reaction was controlled by TLC. The reaction mixture was then diluted with chloroform, then 2N NH₃ in methanol (2 mL) was added to this solution. Everything was stirred together for 1 hr, then the solution was washed with water until neutral, then with brine, and dried over Na₂SO₄. Crude product was purified by column chromatography (eluent: chloroform, chloroform:methanol = 98:2,
30 95:5). Pure free amine was dissolved in methanol (4 mL), 1N solution of HCl in methanol (1.5 mL) followed by diethyl ether were added to this solution. Obtained

solid hydrochloride was filtered off and washed with diethyl ether to remove excess of acid. When the filtrate was neutral, solid hydrochloride was precipitated from methanol/diethyl ether, gave analytically pure product. (Yield 40%).

5 ¹H n.m.r (CDCl₃) δ: 8.35-8.33 (m, 2H, H-1,4), 7.85-7.80 (m, 2H, H-2,3),
7.42-7.29 (m, 5H, aromatic), 5.50 (d, 1H, J = 3.55 Hz, H-1'), 5.28 (, 1H, H-7), 4.82 (d,
1H, J = 11.52 Hz, aromatic), 4.76 (, 2H, NH₂), 4.67 (d, 1H, J = 11.51 Hz, aromatic),
4.10 (q, 1H, J = 6.58 Hz, H-5'), 3.44 (, 1H, H-4'), 3.24 (dd, 1H, J = 19.0 Hz, J = 1.59
10 Hz, H-10), 3.05-2.95 (m, 1H, H-3'), 3.00 (d, 1H, J = 19.0 Hz, H-10), 2.34 (d, 1H, J =
14.77 Hz, H-8), 2.07 (dd, 1H, J = 14.74 Hz, J = 4.0 Hz, H-8), 1.85 (ddd, 1H, J = J =
13.01 Hz, J = 3.1 Hz, H-2'a), 1.70 (dd, 1H, J = 13.09 Hz, J = 4.53 Hz), 1.32 (d, 3H, J
= 6.57 Hz, H-6').

(v) WP 799 FIG. 17

15 **4'-(2,6-difluoro-benzyloxy)-doxorubicine**

Product was obtained according to the procedure A and B, yield 45 % ¹H
n.m.r (CDCl₃) δ: 7.9 (d, 1H, J = 7.45 Hz, H-1), 7.72 (dd, 1H, J = 7.9 Hz, J = 7.9 Hz,
H-2), 7.23 (d, 1H, J = 7.8 Hz, H-3), 7.28-7.24 (m, 2H, aromatic), 6.89-6.86 (m, 2H,
2,6 aromatic) 5.44 (d, 1H, J = 2.5 Hz, H-1'), 5.26 (, 1H, H-7), 4.76 (ABq, 2H,
20 aromatic), 4.71 (s, 4H, aromatic and 9-OH) 4.10-3.98 (m, 1H, H-5'), 4.02 (s, 3H,
OMe) 3.42 (, 1H, H-4'), 3.20 (d, 1H, J = 19 Hz, H-10), 2.97 (d, 1H, J = 19 Hz, H-10),
3.01-2.90 (m, 1H, H-3'), 2.27 (d, 1H, J = 14.8 Hz, H-8), 2.07 (dd, 1H, J = 14 Hz, J =
3.7 Hz, H-8), 1.75-1.64 (m, 2H, H-2'a, H-2'e), 1.29 (d, 3H, J = 6.5 Hz, H-6').

25

(vi) WP 797 FIG.18

4'-(4-Fluoro-benzyloxy)-4-demethoxydaunorubicine

Product was obtained according to the procedure A and B, yield 60% ¹H n.m.r
30 (CDCl₃) δ: 8.3-8.27 (m, 2H H-1, H-4), 7.81-7.76 (m, 2H, H-2,3), 7.36-7.29 (m, 2H,
4F-Ph-CH₂), 7.02-6.96 (m, 2H, 4F-Ph-CH₂), 5.46 (d, 1H, J = 3.5 Hz, H-1'), 5.24 (,

1H, H-7), 4.66 (ABq, 2H, 4F-Ph-CH₂), 4.07 (q, 1H, J = 6.59 Hz, H-5'), 3.39 (, 1H, H-4'), 3.20 (dd, 1H, J = 19 Hz, J = 1.9 Hz, H-10), 2.95 (d, 1H, J = 19 Hz, H-10), 3.0-2.9 (m, 1H, H-3'), 2.37 (s, 3H, OMe), 2.30 (d, 1H, J = 14 Hz, H-8), 2.03 (dd, 1H, J = 14 Hz, J = 4.2 Hz, H-8), 1.88 (ddd, 1H, J = 13 Hz, J = 13 Hz, J = 3.93 Hz, H-2'a),
5 1.67 (dd, 1H, J = 13 Hz, J = 4.6 Hz, H-2'e), 1.26 (d, 3H, J = 6.6 Hz, H-6').

B. Synthesis of 3-azido-2,3,6-trideoxyhexopyranoses (3-azido-daunosamine and 3 azidoacosamine), sugars used in the synthesis of 4'-O-benzylated analogs of 3' amino-anthracyclines.

(vii) WP 219 FIG. 40

4-O-acetyl-3-azido-1-O-t-butyldimethylsilyl-2,3,6-trideoxy-b-L-arabino hexapyranose

The mixture of 3,4-di-O-acetyl-L-rhamnal (1 mol) and water (2700 mL) was
15 heated to 70 °C, and it was stirred at this temperature until substrate is hydrolyzed (TLC). After substrate disappearance the mixture was cooled down to 0 °C, then sodium azide (2.2 mol) followed by acetic acid (85 mL) were added to the mixture, and it was stirred at 0 °C for 2 hr. Second portion of sodium azide (2.2 mol) and acetic acid (85 mL), and methylene chloride (1500 mL) were added to the reaction mixture,
20 and it was stirred at room temperature overnight. The organic layer was separated, the water layer was extracted with dichloromethane (3 x 500 mL). Combined organic solutions were washed with water until neutral, then dried over Na₂SO₄. The drying agent and solvent were removed, and obtained product, light yellow oil was dried, gave crystals (0.9 mol). Obtained mixture of azides was dissolved in *N,N*
25 dimethylformamide (300 ml), then imidazole (256 g), and tertbutyldimethylsilyl chloride (185 g) were added. The mixture was stirred at room temperature overnight, then it was diluted with water (600 mL) and extracted with hexanes (4 x 500 mL). The combined organic extracts were washed with water until neutral, then with brine, and it was dried over Na₂SO₄. The drying agent and solvent were removed and crude
30 product - light yellow oil was dried with vacuum pump overnight. Product after silylation was dissolved in methanol (1400 mL), and cooled down to 0 °C then

sodium methanolate (1M solution in methanol) (14 ml) was added. The reaction mixture was stirred at 0 °C for 3 hr., then 1 N water solution of HCl (14 ml) was added. The mixture was diluted with water (1400 mL), and extracted with hexanes (3 x 500 mL), then with dichloromethane (3 x 500 mL). The hexanes extracts were combined and washed with water until neutral, then with brine, and it was dried over Na₂SO₄. Crude product was purified by column chromatography, using as an eluent: hexanes, hexanes: ethyl acetate = 98:2, gave colorless oil 4-O-acetyl-3-azido-1-O-t-butylidimethylsilyl-2,3,6-trideoxy-β-L-arabino hexopyranose (158 g; 0.48 mol). [α]_D 1.46° (c=1.045, chloroform).

n.m.r (CDCl₃) d: 4.82 (dd, 1 H, *J* = 9.3 Hz, *J* = 1.9 Hz, H-1), 4.67 (t, 1H, *J* = *J* = 9.8 Hz, H-4), 3.50 (ddd, 1H, *J* = 4.9 Hz, *J* = 12.8 Hz, *J* = 9.8 Hz, H-3), 3.44 (qd, 1H, *J* = 6.3 Hz, *J* = 9.3 Hz, H-5), 2.20 (ddd, 1H, *J* = 12.8 Hz, *J* = 4.9 Hz, *J* = 1.9 Hz, H-2e), 2.16 (s, 3H, OAc), 1.69 (ddd, 1H, *J* = 12.8 Hz, *J* = 12.8 Hz, *J* = 9.3 Hz, H-2a), 1.21 (d, 3H, *J* = 6.3 Hz, H-6), 0.90 (s, 9H, t-BuSi), 0.13, 0.11 (s, 3H, Me₂Si).

Anal elem calc for C₁₄H₂₇N₃O₄Si : C 51.04, H 8.26, N 12.75.

Found: C 51.3, H 8.33, N 12.47.

(viii) **WP 460 FIG. 40**

3-azido-1-O-t-butylidimethylsilyl-2,3,6-trideoxy-β-L-arabino-hexapyranose

Obtained in previous step sugar (100 g) was dissolved in methanol (500 mL), then solid potassium carbonate (100 g) was added to this solution. The mixture was stirred at room temperature. After 0.5 hr reaction was completed, solid salts were filtered off. Organic solution was diluted with water (300 mL), and extracted with hexanes (3 x 300 mL). Combined extracts were washed with water until neutral, dried over sodium sulfate. The drying agent and solvents were removed gave pure product (white crystals), 82.8 g, yield 95 %; [α]_D+28.9° (c=1.3, chloroform). ¹H n.m.r (CDCl₃), d: 4.83 (dd, 1H, *J* = 9.3 Hz, *J* = 2.0 Hz, H-1), 3.42 (q, 1H, *J* = 4.8 Hz, *J* = 12.6 Hz, *J* = 9.6 Hz, H-3), 3.35 (dq, 1H, *J* = 6.1 Hz, *J* = 9.0 Hz, H-5), 3.17 (td, 1H, *J*

= 9.6 Hz, $J = 9.0$ Hz, $J = 3.6$ Hz, H-4), 2.30 (d, 1H, $J = 3.6$ Hz, 4-OH), 2.21 (ddd, 1H, $J = 12.6$ Hz, $J = 4.8$ Hz, $J = 2.0$ Hz, H-2e), 1.68 (ddd, 1H, $J = 12.6$ Hz, $J = 12.6$ Hz, $J = 9.3$ Hz, H-2a), 1.34 (d, 3H, $J = 6.1$ Hz, H-6), 0.92 (s, 9H, t-Bu), 0.14, 0.13 (2s, 1H each, Me₂Si).

5 Anal elem calc for C₁₂H₂₅N₃O₃Si: C 50.14, H 8.77, N 14.62.

Found: C 50.20, H 8.78, N 14.54.

C. Alternative method for resolution of mixture 3-azido-L-arabino- and 3-azido L-ribo-hexapyranoses.

10 3-Azido-2,3,6-trideoxy-L-arabino- and 3-azido-2,3,6-trideoxy-L-ribo hexapyranoses obtained by hydrolysis of 3,4-do-O-acetyl-L-rhamnal and subsequent Michael addition of HN₃, were separated by crystallization.

(ix) WP 417 FIG. 40

15 **Synthesis of 3-azido-4-O-benzyl-2,3,6-trideoxy-L-lyxohexapyranoside**

t-Butyldimethylsilyl 3-azido-2,3,6-trideoxy-L-arabinohexopyranose (WP460) (18.0 g, 62.7 mmol) was added to a solution of dry pyridine (38 ml) dichloromethane (1200 ml). The solution was purged with argon and cooled to -40 °C. To this vigorous stirred mixture, solution of trifluoromethanesulfonic anhydride (38 ml, 63.4 g, 225 mmol) in methylene chloride (100 ml) was added dropwise through septum, during 1.5 h. The reaction mixture was allowed to warm up to the room temperature, then it was diluted with methylene chloride (300 ml) and washed with sodium acetate (10% aqueous solution) and phosphate buffer (pH 8) (2 x 200 ml). The solvent was evaporated twice with toluene (200 ml). The oily residue was dissolved in dry
20 dimethylformamide (25 ml) and tetrabutylammonium acetate (22.0 g, 71 mmol) was added to this solution. The mixture was stirred for 1h at room temperature. After the complete disappearance of the triflate (TLC) the mixture was diluted with water (300 ml), and extracted with ethyl acetate (3 x 250 ml). Combined extracts were washed with 5% NaHCO₃, and water until neutral. The organic layer was then dried over
25 anhydrous sodium sulfate. Crude product was purified by column chromatography gave pure WP417 (15.5 g, 47.1 mmol). Yield: 75.1%; [α]_D + 5.3° (c=1.3, methylene
30

chloride); ^1H n.m.r (CDCl₃) δ : 5.06 (d, 1H, J = 3.1 Hz, H-4), 4.78 (dd, 1H, J = 8.4 Hz, J = 2.8 Hz, H-1), 3.60 (q, 1H, J = 6.44 Hz, H-5), 3.42 (ddd, 1H, J = 12.1 Hz, J = 5.3 Hz, J = 3.2 Hz, H-3), 2.18 (s, 3H, OAc), 2.00-1.86 (m, 2H, H-2a, H-2e), 1.18 (d, 3H, J = 6.5 Hz, H-6), 0.91 (s, 9H, t-Bu), 0.14, 0.12 (2s, 3H each, Me₂Si).

5 Anal Calc for: C₁₄H₂₇N₃O₄Si: H 8.26, C 51.04, N 12.75.

Found: H 8.31, C 51.22, N 12.68

(x) **WP 418 FIG. 40**

t-Butyldimethylsilyl 3-azido- -2,3,6-trideoxy- β -L-lyxo hexapyranose

10 Obtained in previous step sugar WP471 (15.5 g) was dissolved in methanol (100 ml), then solid potassium carbonate (15.5 g) was added to this solution. The mixture was stirred at room temperature. Progress of the reaction was monitored by T.L.C. After 0.5 hr reaction was completed, solid salts were filtered off. Organic solution was diluted with water (100 ml), and extracted with hexanes (3 x 200 ml),
15 Combined extracts were washed with water until neutral and dried over sodium sulfate. The drying agent and solvents were removed gave pure product (white crystals), 13.25 g Yield : 98%

mp 74-76 °C, $[\alpha]_D$ -4.1° (c=1.8, dichloromethane) ^1H n.m.r (CDCl₃) δ : 4.74 (dd, 1H, J = 9.1 Hz, J = 2.4 Hz, H-1), 3.60 (dd, 1H, J = 2.7 Hz, J = 8.4 Hz, H-4),
20 3.50 (q, 1H, J = 6.5 Hz, H-5), 3.30 (ddd, 1H, J = 2.9 Hz, J = 12.7 Hz, J = 2.7 Hz, H-3), 2.03 (d, 1H, J = 8.4 Hz, 4-OH), 1.99 (ddd, 1H, J = 12.7 Hz, J = 2.9 Hz, J = 2.4 Hz, H-2e), 1.84 (td, 1H, J = 12.7 Hz, J = 12.7 Hz, J = 9.1 Hz, H-2a), 1.29 (d, 3H, J = 6.5 Hz, H-6), 0.90 (s, 9H, t-Bu), 0.13, 0.11 (s, 3H each, Me₂Si).

Anal Calc for: C₁₂H₂₅N₃O₃Si: H 8.77, C 50.14, N 14.62.

25 Found: H 8.80, C 50.11, N 14.52

(xi) **WP 819 FIG. 26**

1-O-t-butyldimethylsilyl-3-Azido-4-O-benzylo-2,3,6-trideoxy- β -L-lyxo hexapyranose

30 Sodium hydride (1 g of 60 % suspension in mineral oil) was added to a cooled to 0 °C solution of WP 418 (3.9 g, 13.5 mmol) in dry N, N dimethylformamide (40

ml). The reaction mixture was stirred for 15 min, then solution of benzyl bromide (2.97 ml, 4.27 g, 25 mmol) was added. The stirring continued for 0.5 h at 0 °C, then mixture was allowed to warm up to room temperature. The reaction mixture was poured into the ice/water and extracted with ethyl acetate (3 x 100 ml). The combined
5 extracts were washed with water, then with saturated solution of sodium bicarbonate and dried over magnesium sulfate. The pure product (3.145 g, 8.34 mmol) was separated by column chromatography. Yield : 61.8%. ¹H n.m.r (CDCl₃) δ: 7.4-7.2 (m, 5H, H-aromatic), 4.84 (d, 1H, *J* = 11.4 Hz, H-aromatic), 4.70 (dd, 1H, *J* = 9 Hz, *J* = 2.19 Hz, H-1), 4.61 (d, 1H, *J* = 11.4 Hz, Ch₂Ph), 3.39 (q, 1H, *J* = 6.1 Hz, H-5),
10 3.33 (d, 1H, *J* = 2.67 Hz, H-4), 3.25 (ddd, 1H, *J* = 4.46 Hz, *J* = 12.8 Hz, *J* = 2.67 Hz, H-3), 2.03 (ddd, 1H, *J* = 12.3 Hz, *J* = 12.8 Hz, *J* = 9 Hz, H-2a), 1.90 (ddd, 1H, *J* = 12.3 Hz, *J* = 4.46 Hz, *J* = 2.19 Hz, H-2e), 1.15 (d, 3H, *J* = 6.1 Hz, H-6), 0.90 (s, 9H, t-Bu), 0.13, 0.11 (s, 3H each, Me₂Si).

15 **D. Direct synthesis of 3-azido-daunosamine.**

(xii) **WP 417 FIG. 40**

**4-O-acetyl-3-Azido-1-O-t-butyldimethylsilyl-2,3,6-trideoxy-β-L-lyxo
hexapyranose**

The mixture of 3,4-di-O-acetyl-fucal (1 mol) and water (2700 mL) was heated
20 to 80 °C, and it was stirred at this temperature until all 3,4-di-O-acetyl-fucal hydrolysed (TLC). After all substrate disappeared from reaction mixture, the mixture was cooled down to 0°C, then sodium azide (2.2 mol) followed by acetic acid (85 mL) were added to the mixture, and it was stirred at 0 °C for 2 hr. Second portion of sodium azide (2.2 mol) and acetic acid (85 mL), and dichloromethane (1500 mL)
25 were added to the reaction mixture, and it was stirred at room temperature overnight. The organic layer was separated, the water layer was extracted with dichloromethane (3 x 500 mL). Combined organic solutions were washed with water until neutral, then dried over Na₂SO₄. The drying agent and solvent were removed, and obtained product mixture, a light yellow oil was dried, to gave crystals (0.89 mol). Obtained
30 mixture of azides was dissolved in dimethylformamide (300 mL), then imidazole (256 g), and tert butyldimethylsilyl chloride (185 g) were added. The mixture was stirred at

room temperature overnight, then it was diluted with water (600 mL) and extracted with hexanes (4 x 500 mL). The combined organic extracts were washed with water until neutral, then with brine, and it was dried over Na₂SO₄. The drying agent and solvent were removed and crude product, light yellow oil was dried overnight.

- 5 Product after silylation was dissolved in methanol (1400 mL), and cooled down to 0 °C then sodium methanolate (1 M solution in methanol) (14 mL) was added. The reaction mixture was stirred at 0°C for 3 hr., then 1 N water solution of HCl (14 mL) was added. The mixture was diluted with water (1400 mL), and extracted with hexanes (3 x 500 mL), then with methylene chloride (3 x 500 mL). The hexanes
10 extracts were combined and washed with water until neutral, then with brine, and it was dried over Na₂SO₄. Crude product was purified by column chromatography, using hexanes, hexanes: ethyl acetate = 98:2 as eluent, gave analytically pure product (48.5 g; 0.168 mol); [α]_D +5.3° (c=1.3, chloroform) 1H n.m.r (CDCl₃) δ: 5.06 (d, 1H, *J* = 3.1 Hz, H-4), 4.78 (dd, 1H, *J* = 8.4 Hz, *J* = 2.8 Hz, H-1), 3.60 (q, 1H, *J* = 6.44
15 Hz, H-5), 3.42 (ddd, 1H, *J* = 12.1 Hz, *J* = 5.3 Hz, *J* = 3.2 Hz, H-3), 2.18 (s, 3H, OAc), 2.00-1.86 (m, 2H, H 2a, H-2e), 1.18 (d, 3H, *J* = 6.5 Hz, H-6), 0.91 (s, 9H, t-Bu), 0.14, 0.12 (2s, 3H each, Me₂Si).

Anal elem calc for: C₁₄H₂₇N₃O₄Si : C 51.04, H 8.26, N 12.75.

Found: C 51.22, H 8.31, N 12.68.

20

E. 4'-O-benzylated-3'-hydroxy-doxorubicin analogs:

(xiii) WP 568 FIG.35

25 Synthesis of 3 -Oacetyl-4 -Obenzyl-3 -deamino-14-Ot-butyl-diphenylsilyl-doxorubicin.

A mixture of 14-O-t-butyl-diphenylsilyl-adriamycinone (0.84 g, 1.29 mmol), 3-O-acetyl-4-O-benzyl-L-fucal (0.84 g, 3.2 mmol), triphenylphosphine hydrobromide (0.055 g, 0.16 mmol) in dichloromethane (18 mL) was stirred for 48 hr. During this time additional portion of 3-O-acetyl-4-O-benzyl-L-fucal (0.175 g, 0.7 mmol) was
30 added and stirred for 24 hours. The reaction mixture was diluted with dichloromethane (200 mL), washed with water and dried with anhydrous Na₂SO₄.

The product was purified by chromatography on silicagel column using dichloromethane : acetone (95 : 5 vol.) to give 3 -O-acetyl-4 -O-benzyl-3 -deamino-14-O-t-butyldiphenylsilyl-doxorubicin (WP 568) (1.0 g, 1.09 mmol). Yield: 84%. [a]_D 116.84° (c=0.0354, chloroform : methanol 1 : 1). ¹H n.m.r (CDCl₃) δ: 13.94 (s, 1H, OH-6), 13.24 (s, 1H, OH-11), 8.1 (d, 1H, J = 7.8 Hz, H-1), 7.68-7.85 (m, 5H, J = 8.1 Hz, from silyl, H-2), 7.5-7.2 (m, 12H, J = 7.3 Hz, from silyl, Ch₂Ph, H-3), 5.49 (d, 1H, J = 3.9 Hz, H-1), 5.19 (bs, 1H, H-7), 5.0-4.78 (m, 3H, J = 19.8 Hz, H-14, H-3), 4.7 (d, 1H, J = 11.6 Hz, Ch₂Ph), 4.59 (d, 1H, J = 11.8 Hz, Ch₂Ph), 4.29 (s, 1H, OH-9), 4.07 (s, 3H, CH₃O), 3.90 (q, 1H, J = 6.5 Hz, H-5), 3.60 (bs, 1H, H-4), 3.02 (d, 1H, J = 19.3 Hz, H-10eq), 2.85 (d, 1H, J = 19.1 Hz, H-10ax), 2.3-1.95 (m, 3H, H-8, H-2a), 1.93 (s, 3H, CH₃CO), 1.8 (dd, 1H, J = 3.8 Hz, J = 12.7 Hz, H-2e), 1.43 (s, 9H, tBu), 1.07 (d, 3H, J = 6.5 Hz, H-6).

(xiv) WP 569 FIG.35

Synthesis of 4 -Obenzyl-3 -deamino-3 -hydroxy-14-Ot-butyldiphenylsilyl-doxorubicin.

3 -O-Acetyl-4 -O-benzyl-3 -deamino-14-O-t-butyldiphenylsilyl-doxorubicin (WP 568) (1.0 g, 1.09 mmol) was dissolved in mixture of tetrahydrofuran (20 mL) and methanol (80 mL), and then anhydrous potassium carbonate (1.0 g, 7.2 mmol) was added. The reaction mixture was stirred vigorously at room temperature for 2 h, until substrate disappeared. Then reaction mixture was diluted with dichloromethane (300 mL) and poured into 0.05 N HCl (100 mL), extracted with dichloromethane, washed with water, dried with anhydrous Na₂SO₄. The solvent was evaporated and the residue was chromatographed using dichloromethane : acetone (9 : 1 vol.) to give 4 -O-benzyl-3 -deamino-3 -hydroxy-14-O-t-butyldiphenylsilyl-doxorubicin (WP 569) (0.6 g, 0.687 mmol). Yield: 63%. [a]_D 52.94° (c=0.022, chloroform : methanol 1 : 1). ¹H n.m.r (CDCl₃) δ: 13.93 (s, 1H, OH-6), 13.23 (bs, 1H, OH-11), 8.02 (d, 1H, J = 7.6 Hz, H-1), 7.82-7.67 (m, 5H, from silyl, J = 8.2 Hz, H-2), 7.45-7.25 (m, 12H, J = 8.3 Hz, from silyl, Ch₂Ph, H-3), 5.47 (d, 1H, J = 3 Hz, H-1), 5.19 (bs, 1H, H-7), 4.95 (d, 1H, J = 19.8 Hz, H-14), 4.84 (d, 1H, J = 19.8 Hz, H-14), 4.80 (d, 1H, J = 11.6 Hz, Ch₂Ph), 4.64 (d, 1H, J = 11.6 Hz, Ch₂Ph), 4.43 (s, 1H, OH-9), 4.07 (s, 3H, CH₃O),

3.92 (q, 1H, J = 6.5 Hz, H-5), 3.70 (bs, 1H, H-3), 3.45 (d, 1H, J = 2.3 Hz, H-4), 3.0 (d, 1H, J = 19 Hz, H-10eq), 2.83 (d, 1H, J = 19 Hz, H-10ax), 2.35 (s, 1H, OH), 2.12-1.95 (m, 2H, H-8), 1.92-1.7 (m, J = 3.8 Hz, H-2) 1.13 (s, 12H, tBu), 1.11 (s, 3H, H-6).

5

(xv) Synthesis of 4 -Obenzyl-3 -Deamino-3 -Hydroxy-Doxorubicin - WP 727 FIG. 23

4 -O-Benzyl-3 -deamino-3 -hydroxy-14-O-t-butylidiphenylsilyl-doxorubicin (WP 569) (0.6 g, 0.687 mmol) was dissolved in THF (33 mL), and a reagent (25 mL) / made from THF (24 mL), dichloromethane (12 mL), pyridine (0.66 mL), 1.0M Bu₄NF (0.6 mL)/ was added, and stirred vigorously until substrate disappear. Then the reaction mixture was poured into solution containing ice (50 g), 0.1 N HCl (100 mL), dichloromethane (100 mL). Organic layer was separated and water layer was extracted with dichloromethane (3 x 50 mL). The combined organic layers were washed with water, dried with anhydrous Na₂SO₄, evaporated, and chromatographed on silicagel column using dichloromethane : acetone (7 : 3 vol.) to give pure product 4 -O-benzyl-3 -deamino-3 -hydroxy-doxorubicin (WP 570) (0.34 g, 0.536 mmol). Yield: 78%. [α]_D²⁰ 101.00° (c=0.029, chloroform : methanol 1 : 1). ¹H n.m.r (CDCl₃) δ: 13.93 (s, 1H, OH-6), 13.22 (s, 1H, OH-11), 8.02 (d, 1H, J = 8 Hz, H-1), 7.77 (t, 1H, J = 8 Hz, H-2), 7.38 (m, 6H, H-aromatic, H-3), 5.55 (d, 1H, J = 3.2 Hz, H-1), 5.26 (s, 1H, H-7), 4.83 (dd, 1H, J = 11. 8 Hz, H-14), 4.7-4.65 (m, 4H, J = 2.75 Hz, H-14, H-aromatic, OH-9), 4.07 (s, 3H, CH₃O), 4.03 (q, 1H, J = 6.6 Hz, H-5), 3.9-3.78 (m, 1H, H-3), 3.57 (d, 1H, J = 2.52 Hz, H-4), 3.25 (dd, 2H, J = 1.5 Hz, J = 18.7 Hz, H-10eq), 3.01 (d, bs, 3H, J = 18.8 Hz, H-10ax, OH-3), 2.32 (bd, 1H, J_{8e-8a} = 14.8 Hz, H-8eq), 2.13 (dd, 1H, J = 3.9 Hz, J_{8a-8e} = 14.7 Hz, H-8ax), 1.95 (dt, 1H, J = 4 Hz, J = 12.8 Hz, H-2a), 1.84 (dd, 1H, J = 4.8 Hz, J = 12.8 Hz, H-2e), 1.33 (d, 3H, J = 6.5 Hz, H-6).

(xvi) WP 571 FIG. 38**Synthesis of 4 -Obenzyl-3 -deamino-3 -Otriethylsilyl-14-Opentanoyl-doxorubicin.**

A mixture of 14-O-pentanoyl-adriamycinone (0.6 g, 1.2 mmol), 4-O-benzyl-3-O-triethylsilyl-L-fucal (1.1 g, 3.3 mmol), sieves 4 (0.5 g), TPHB (0.06 g, 0.18 mmol) in dichloromethane (20 mL) was stirred overnight at room temperature. Then the reaction mixture was diluted with dichloromethane (100 mL) and washed with water, dried with anhydrous Na₂SO₄; the solvent was evaporated, and the residue was chromatographed on silicagel column using dichloromethane : acetone (98 : 2 vol.) to give 4 -O-benzyl-3 -deamino-3 -O-triethylsilyl-14-O-pentanoyl-doxorubicin (0.89 g, 1.07 mmol). Yield: 89%. ¹H n.m.r (CDCl₃) δ: 13.94 (s, 1H, OH-6), 13.25 (s, 1H, OH-11), 8.03 (d, 1H, J = 7.8 Hz, H-1), 7.77 (t, 1H, J = 8 Hz, H-2), 7.43-7.26 (m, 6H, J = 7.2 Hz, H-aromatic, H-3), 5.55 (d, 1H, J = 3.7 Hz, H-1), 5.31 (bs, 1H, H-7), 5.32 (d, 1H, J = 18.2 Hz, H-14), 5.12 (d, 1H, J = 18.2 Hz, H-14), 4.98 (d, 1H, J = 12.6 Hz, CH₂Ph), 4.92 (s, 1H, OH-9), 4.67 (d, 1H, J = 12.7 Hz, CH₂Ph), 4.08 (s, 3H, CH₃O), 4.02-3.90 (q, m, 2H, J = 6.5 Hz, H-5, H-3), 3.45 (bs, 1H, H-4), 3.28 (d, 1H, J = 19 Hz, H-10eq), 3.03 (d, 1H, J = 19 Hz, H-10ax), 2.50-2.38 (m, 4H, J = 7.4 Hz, CH₂ from pentanoyl, H-8), 2.22 (dt, 1H, J = 3.1 Hz, J = 12.7 Hz, H-2a), 2.06 (dd, 1H, J = 3.8 Hz, J = 12.7 Hz, H-2e), 1.77-1.62 (m, 2H, J = 7.4 Hz, CH₂ from pentanoyl), 1.48 - 1.35 (m, 2H, J = 7.5 Hz, CH₂ from pentanoyl), 1.23 (d, 3H, J = 6.5 Hz, H-6), 0.95 (t, t, 12H, J = 7.8 Hz, CH₃ from pentanoyl, (Et)₃Si), 0.58 (q, 6H, J = 8 Hz, (Et)₃Si).

(xvii) WP 571 FIG. 38**Synthesis of 4 -Obenzyl-3 -Deamino-3 -Hydroxy-14-Opentanoyl-Doxorubicin**

4 -O-Benzyl-3 -deamino-3 -O-triethylsilyl-14-O-pentanoyl-doxorubicin (0.715 g, 0.86 mmol) was dissolved in THF (15 mL), and then 0.1 N HCl (10 mL) was added. Reaction mixture was stirred at room temperature for 1 h until substrate disappeared. Then the reaction mixture was diluted with water (30 mL); a precipitate was filtered, washed with water and dried at 100 °C under diminished pressure overnight to give pure 3 -deamino-4 -O-benzyl-3 -O-hydroxy-14-O-pentanoyl-

doxorubicin (WP 571) (0.56 g, 0.78 mmol). Yield: 91%. ¹H n.m.r (CDCl₃) δ: 13.91 (s, 1H, OH-6), 13.18 (s, 1H, OH-11), 7.99 (d, 1H, J = 7.7 Hz, H-1), 7.75 (t, 1H, J = 8 Hz, H-2), 7.46-7.29 (m, 6H, H-aromatic, H-3), 5.54 (d, 1H, J = 3.2 Hz, H-1), 5.32 (d, 1H, J = 18.1 Hz, H-14), 5.25 (bs, 1H, H-7), 5.10 (d, 1H, J = 18.2 Hz, H-14), 4.85 (d, 1H, J = 11.7 Hz, CH₂Ph), 4.74 (s, 1H, OH-9), 4.69 (d, 1H, J = 11.7 Hz, CH₂Ph), 4.12-4.00 (q, s, 4H, H-5, CH₃O), 3.90-3.70 (m, 1H, H-3), 3.58 (d, 1H, J = 2 Hz, H-4), 3.25 (d, 1H, J = 18.7 Hz, H-10eq), 2.97 (d, 1H, J = 18.7 Hz, H-10ax), 2.52-2.38 (m, 3H, J = 7.4 Hz, CH₂ from pentanoyl, H-8), 2.08 (dd, 1H, J = 3.8 Hz, J = 14.9 Hz, H-8), 1.95 (dt, J = 4 Hz, J = 12.9 Hz, H-2a), 1.89-1.82 (dd, 1H, J = 5 Hz, J = 12.9 Hz, H-2e), 1.75 (d, 1H, J = 9.4 Hz, OH-3), 1.75-1.65 (m, 2H, J = 7.4 Hz, CH₂ from pentanoyl), 1.48-1.3 (m, 2H, J = 7.4 Hz, CH₂ from pentanoyl), 1.36 (d, 3H, J = 6.5 Hz, H-6), 0.95 (t, 3H, J = 7.4 Hz, CH₃ from pentanoyl).

EXAMPLE 3

Novel Method for the Synthesis of 4-O-Benzylated Sugars

Described herein is a method for the preparation of the 4-O-benzylated sugars, used as precursors in the synthesis of 4-O-benzylated-3-hydroxy-anthracyclines like WP727, WP571, WP794, or WP793. This is a novel method superior to other methods described earlier. A direct selective alkylation of acetylated glycols is used which gives almost exclusively products alkylated at C-4. The inventors approach is different from other known methods in the following ways: (1) a one step approach wherein a fully blocked compound is alkylated allowing for an easier and simpler synthesis versus the standard method of alkylating free hydroxyl groups, (2) because the process is selective the major products are 4-O-alkylated compounds. In this method, the first step of the reaction is a selective deblocking (deacylation) of the hydroxyl at C-4 followed by a rapid alkylation of this free hydroxyl group. In contrast, alkylation of unblocked (hydroxy compounds) glycols gives mixtures of

mono-benzylated and dibenzylated products and 4-O-benzylated-glycals are obtained in lower yields.

Such 4-O-benzylated glycals can be used directly towards synthesis of the 4-O-benzylated anthracycline analogs described in the previous example, using electrophilic addition to glycals in the presence of triphenylphosphine hydrobromide (TPHB), for example the synthesis of WP727.

The inventors method works well not only for the 6-deoxy-glycals like fucal and rhamnal but to also for glycals like D-glucal or D-galactal having hydroxyl groups at C-3, 4, and 6 positions. Direct alkylation of 3,4,6-tri-O-acetylated-D-glucal or -D-galactal give 4-O-benzylated-glycals as a the main product. Therefore, this method is of general use.

More importantly, the inventors found that these reactions can be continued in the same vessel to the next step to synthesize 3 -hydroxy-4-O-benzylated glycals. For this the other acetyl groups are removed by the addition of methanol to the reaction mixture. Methanol forms *in situ* sodium methoxide, which in turn deacetylates all hydroxyl groups. Described below are the experimental procedures and data for the 4-O-benzylated substrates which are either acetylated or bear free hydroxyl groups, and data for other 4-O-benzylated sugars to prove the general nature of the inventors approach.

This method can be used to synthesize 4-O-benzylated products and also other 4-O-alkylated products. The 4-O-benzylated glycals can then be used directly in the coupling reactions with proper aglycon like daunomycinone or can be first transformed into 4-O-alkylated 1-O-silylated hexopyranoses which can further be coupled using a variety of other coupling methods as for example in the synthesis of WP793 and WP794.

A. Synthesis of 4-O-Alkylated-Glycals : Fig 39**1. Examples of Selective Benzylation of Glycals at the C-4 Position****(i) Selected procedure for 4-O-benzyl-L-rhamnal exemplifying**

general procedure. To a solution of NaH (2.5 g, 0.063 mmol) in DMF (30 mL) cooled to ?C 3,4-di-O-acetyl-L-rhamnal (6.5 g, 0.03 mmol) was added. The reaction mixture was stirred for 15 min, and benzyl chloride (6 mL, 0.046 mmol) was added. Then the reaction was allowed to warm up to room temperature and stirred until TLC showed complete consumption of starting material. The reaction was cooled to 0°C methanol (10 ml) was added, with continuous stirring for 10 min. The reaction mixture was then poured into a solution containing 1 N HCl (10 mL), ice (20 g), and ethyl acetate (50 mL). The organic phase was separated and aqueous phase was extracted with ethyl acetate (3 x 50 mL). The combined organic solution was washed with water, dried with anhydrous Na₂SO₄, and concentrated to give an oily-solid product that was crystallized from dichloromethane/hexane.

(ii) 4-OBenzyl-L-rhamnal Yield: 73%, mp 110-110.5 C. [a] D -44.12° (c=1.05, ethyl acetate). ¹H n.m.r (CDCl₃) δ: 7.6-7.0 (m, 5H, H-aromatic), 6.32 (dd, 1H, J = 5.7 Hz, H-1), 4.90-4.72 (dd, 2H, J = 11.8 Hz, CH₂Ph), 4.69 (dd, 1H, J_{1,2} = 5.7 Hz, J_{2,3} = 2 Hz, H-2), 4.34 (m, 1H, H-3), 3.97-3.85 (dq, 1H, J_{5,4} = 9.4 Hz, J_{5,6} = 6.3 Hz, H-5), 3.32-3.22 (dd, 1H, J_{4,5} = 9.4 Hz, H-4), 1.70 (d, 1H, J = 5.8 Hz, OH), 1.40 (d, 3H, J_{6,5} = 6.3 Hz, H-6).

(iii) 4-OBenzyl-L-fucal Yield: 63%, mp 77.0-77.5 C. [a] D 13.3° (c=1.18, chloroform). ¹H n.m.r (CDCl₃) δ: 7.55-7.30 (m, 5H, H-aromatic), 6.36 (dd, 1H, J_{1,2} = 6 Hz, J_{1,3} = 0.9 Hz, H-1), 4.8 (m, 2H, H-aromatic), 5.75-5.68 (m, 1H, H-2), 4.45-4.35 (m, 1H, H-3), 4.08 (q, 1H, J_{5,6} = 6.6 Hz, H-5), 3.68 (d, 1H, J = 5.1 Hz, H-4), 2.23 (d, 1H, J = 10 Hz, OH), 1.35 (d, 1H, J_{6,5} = 6.6 Hz, H-6).

(iv) 4-OBenzyl-D-glucal Yield: 72%, mp 98.5-100 C. [a] D 10.37° (c=1.4, chloroform). ¹H n.m.r (CDCl₃) δ: 7.5-7.25 (m, 5H, H-aromatic), 6.37 (dd,

1H, J_{1,2} = 5.9 Hz, J_{1,3} = 1.1 Hz, H-1), 4.95-4.78 (dd, 2H, H-aromatic), 4.75 (dd, 1H, J_{2,1} = 5.9 Hz, J_{2,3} = 2.3 Hz, H-2), 4.41 (m, 1H, H-3), 4.02-3.82 (m, 3H, H-6, H-5), 3.65 (t, J = 6.9 Hz, H-4), 1.90 (d, J = 12.7 Hz, 1H, OH).

5 (v) **4-O-Benzyl-D-galactal** Yield: 55%, mp 98.5-99° C.

2. **Selected procedure for preparation of 3-Oacetyl-4-Obenzyl-L-rhamnal exemplifying general procedure.**

10 To a solution of NaH (2.5 g, 0.063 mmol) in DMF (30 mL) cooled to 0 C 3,4-di-O-acetyl-L-rhamnal was added (6.5 g, 0.03 mmol). The reaction was stirred for 15 min, and BnCl (6 ml, 0.046 mmol) was added. Then the reaction was allowed to warm to room temperature and stirred vigorously until TLC showed complete consumption of starting material. After 1.5 h reaction mixture was poured into solution containing 1 N HCl (10 mL), ice (20 g), and ethyl acetate (50 mL). The
15 organic phase was separated and aqueous phase was extracted with ethyl acetate (3 x 50 ml). The combined organic solution was washed with water, dried with anhydrous Na₂SO₄, and concentrated to give an oil that was purified by column chromatography on silicagel column using hexane : ethyl acetate 95 : 5 as eluents to give pure 3-O-acetyl-4-O-benzyl-L-rhamnal.

20

(vi) **3-OAcetyl-4-Obenzyl-L-rhamnal** Yield: 50%, bp 200 C/0.04 mm Hg. [α]_D 55.77° (c=1.15, ethyl acetate). 1H n.m.r (CDCl₃) δ: 7.42-7.25 (m, 5H, H-aromatic), 6.4 (d, 1H, J = 6.5 Hz, H-1), 5.43-5.36 (m, 1H, H-3), 4.78 (dd, dd, 3H, J_{2,3} = 2.5 Hz, J_{2,1} = 5.8 Hz, J = 11.5 Hz, CH₂Ph, H-2), 4.1-4.0 (m, 1H, J_{5,4} = 8.4 Hz, J_{5,6} = 6.5 Hz, H-5), 3.57-3.49 (dd, 1H, J_{4,5} = 8.2 Hz, H-4), 2.02 (s, 3H, CH₃CO),
25 1.38 (d, 3H, J_{6,5} = 6.7 Hz, H-6).

(vii) **4-OBenzyl-3,6-di-Oacetyl-D-glucal** Yield: 50%, mp 52-55.5 C. [α]_D -7.34° (c=1.4, ethyl acetate). 1H n.m.r (CDCl₃) δ: 7.42-7.18 (m, 5H, H-aromatic),
30 6.42 (d, 1H, J = 6.3 Hz, H-1), 5.43-5.38 (m, 1H, J = 5.3 Hz, J = 3.2 Hz, H-3), 4.83-4.78 (dd, 1H, J = 5.9 Hz, J = 3.1 Hz, H-2), 4.72 (d, 1H, J = 11.6 Hz, CH₂Ph), 4.64 (d,

1H, J = 11.6 Hz, Ch2Ph), 4.42-4.29 (m, 2H, H-6), 4.2-4.12 (m, 1H, H-5), 3.85-3.78 (dd, 1H, J_{4,5} = 8 Hz, H-4), 2.07 (s, 3H, CH₃CO), 2.04 (s, 3H, CH₃CO).

(viii) **4-OBenzyl-3,6-di-Oacetyl-D-galactal** Yield: 59%. ¹H n.m.r (CDCl₃) d: 7.42-7.18 (m, 5H, H-aromatic), 6.4 (d, 1H, J = 6.2 Hz, H-1), 5.53-5.4 (m, 1H, J = 4.4 Hz, J = 4 Hz, H-3), 4.85 - 4.77 (dd, 1H, J_{2,1} = 6.1 Hz, J_{2,3} = 4.3 Hz, H-2), 4.74 (d, 1H, J = 11.8 Hz, Ch2Ph), 4.55 (d, 1H, J = 11.9 Hz, Ch2Ph), 4.52-4.43 (dd, 1H, J = 11.7 Hz, J = 8.5 Hz, H-5), 4.34-4.2 (m, 2H, J = 11.7 Hz, H-6), 3.99 (t, 1H, J_{4,3} = 4 Hz, H-4), 2.09 (s, 3H, CH₃CO), 2.06 (s, 3H, CH₃CO).

(ix) **3-OAcetyl-4-Obenzyl-L-fucal** Yield: 58% bp 190 C/0.02 mm Hg. [α]_D 61.4 (c=0.6, chloroform).

(x) **3-OAcetyl-4-Obenzyl-L-arabinal** Yield: 59%.

(xi) **3-OAcetyl-4-Obenzyl-L-xylal** Yield: 39%. ¹H n.m.r (CDCl₃) d: 7.4-7.25 (m, 5H, H-aromatic), 6.01 (d, 1H, J_{1,2} = 6.1 Hz, H-1), 5.9 (t, 1H, J_{2,1} = 6.3 Hz, H-2), 5.4-5.25 (m, 1H, H-3), 4.75 - 4.62 (dd, 2H, J = 12.2 Hz, Ch2Ph, H-2), 4.15 (m, 1H, J_{5,5} = 11.6 Hz, H-5), 3.9 (m, 1H, J_{5,5} = 11.7 Hz, H-5), 3.65-3.57 (m, 1H, H-4), 2.02 (s, 3H, CH₃CO).

3. General procedure for preparation of 3-O-silylated-4-O-benzyl-L-glycals.

4-O-Benzyl-L-fucal (1.5 g, 10.4 mmol) was silylated with Et₃SiCl (1.1 ml, 10.4 mmol) in presence of imidazole (1.1 g, 17 mmol) in DMF (3 mL). Reaction was finished after 2 h, then water (2 mL) was added, and stirring was continued for 5 min. Then the reaction mixture was extracted with ethyl ether (3 x 10 ml). The combined organic solution was washed with water, dried with anhydrous Na₂SO₄, and concentrated to give an oil that was distilled on Kugelrohr 150 C/0.03 mm Hg to give pure 4-O-benzyl-3-O-triethylsilyl-L-fucal (2.2 g, 6.6 mmol). Yield: 97%.

(xii) **4-O-Benzyl-3-O-t-butyldimethylsilyl-L-rhamnal**. Yield: 98%.

EXAMPLE 4

Assessment of Anti-tumor Activity *In Vitro*

5

Compounds synthesized using the methods described above were tested using a standard MTT assay (Green *et al.*, 1984) against human carcinoma sensitive (KB) and multi-drug-resistant (KBV1) cells and MCF-7 and MCF-7/VP-16 resistant cells characterized as having the MRP (multi-drug resistant associated protein) phenotype.

10 The use of an MTT assay using these cells is recognized as an accepted assay for anti-tumor activity by those in the field.

Methods

In vitro Cytotoxicity against MCF-7, MCF-7/VP-16, and MCF-7/DOX cell

15 lines. *In vitro* drug cytotoxicities against human breast carcinoma wild-type MCF-7 and MRP-resistant MCF-7/VP-16 cells were assessed by using the MTT reduction assay, as previously reported (Green *et al.*, 1984). The MTT dye was obtained from Sigma Chemical Co. (St. Louis, MO). Cells were plated in 96-well microassay culture plates (10^4 cells/well) and grown overnight at 37°C in a 5% CO₂ incubator.

20 Drugs were then added to the wells to achieve a final drug concentration ranging from 0.1 to 50 µg/mL. Four wells were used for each concentration. Control wells were prepared by adding appropriate volumes of calcium- and magnesium-free PBS (pH 7.4). Wells containing culture medium without cells were used as blanks. The plates were incubated at 37°C in a 5% CO₂ incubator for 72 hours. Upon completion of the

25 incubation, 20 µL of stock MTT dye solution (5 mg/mL) was added to each well. After a 4-hour incubation, 100 µL of buffer containing 50% *N,N*-dimethylformamide and 20% SDS was added to solubilize the MTT formazan. Complete solubilization was achieved by placing the plate in a mechanical shaker for 30 minutes at room temperature. The optical density of each well was then measured with a microplate

spectrophotometer at a wavelength of 570 nm. The percent cell viability was calculated by the following equation:

$$\% \text{ cell viability} = (\text{OD treated wells} / \text{OD control wells}) \times 100$$

where OD is the mean optical density from four determinations. The percent cell viability values were plotted against the drug concentrations used, and the ID_{50} was calculated from the curve. Cytotoxicity experiments were repeated at least three times.

Results and Discussion

Drug resistance, both *de novo* and acquired, by human tumors is currently a major factor limiting the effectiveness of chemotherapy. Thus, for the *in vitro* evaluation of substituted anthracyclines having C-3' alkylated anthracyclines modified at benzyl ring and having 4'-substituted-benzylated anthracyclines, the inventors selected two sensitive cell lines: a human carcinoma KB and MCF-7 human breast cancer, the multi-drug-resistant (MDR) counterpart of KB (KBV1 carcinoma), which overexpresses *MDR1* gene that encodes a membrane transport glycoprotein (P-gp), the MCF-7/VP-16 cell line that overexpresses the multi-drug-resistant associated protein (MRP), and the MCF-7/dox cell line which overexpresses *MDR1* gene. Using this system, the inventors evaluate a drug's cytotoxic potential against human tumors and at the same time identify compounds that might have unique activity against MDR tumors (Priebe *et al.*, 1993).

Table 2 shows the *in vitro* evaluation of cytotoxic properties of WP791, WP790, WP786, WP785, WP784, WP778, WP775, WP774, WP758, WP757, WP756, WP755 and doxorubicin (DOX) in KB and KB-V1 cells.

TABLE 2.

***In Vitro* Cytotoxicity against Sensitive And MDR
Tumor Cell Lines¹**

5

Compound	KB	KB-V1	RI ²
	$\mu\text{g/ml}$	$\mu\text{g/ml}$	
WP786	0.60 $\pm$ 0.20	4.9 $\pm$ 0.2	8.1
WP784	2.9 $\pm$ 0.8	0.8 Only one test	-
WP785	0.43 $\pm$ 0.05	49 $\pm$ 12	114
WP790	3.93 $\pm$ 0.81	4.6 $\pm$ 2.3	1.2
WP791	0.67 $\pm$ 0.25	4.6 $\pm$ 3.2	7
WP774	4.5 $\pm$ 1.3	6.33.5	1.4
WP775	0.4 $\pm$ 0.2	4.1 $\pm$ 3.5	10
DOX	0.50 $\pm$ 0.0	>100	>200
WP755	1.50 $\pm$ 0.80	1.80 $\pm$ 0.80	1.20
WP756	2.60 $\pm$ 0.30	2.80 $\pm$ 0.30	1.08
WP757	0.67 $\pm$ 0.15	24.00	36
WP758	0.67 $\pm$ 0.06	2.77 $\pm$ 0.25	4.13
WP778	0.67 $\pm$ 0.41	3.83 $\pm$ 2.75	5.7
DOX	0.43 $\pm$ 0.19	109 $\pm$ 28	253

¹ mean of at least 4 experiments; MTT assay

10

Table 3 shows *in vitro* cytotoxic properties of WP783, WP750, WP744 and doxorubicin (DOX) in KB, KB-V1, MCF-7 and MCF-7/VP-16 cells

TABLE 3.***In Vitro* Cytotoxicity against Sensitive and Typical MDR and MRP Tumor Cell Lines¹**

	KB	KBV1	RI ²	MCF-7	MCF-7/VP-16	RI ³	MCF-7/DOX	RI ⁴
	EXP. 1							
WP744	1.44±0.4 5	2.76±0.5 2	1.84	1.63±0.5	2.3±1.9	1.41	5.25	3.22
DOX	2.35	>100	>42.6	0.3	3.0	10	>100	>333
	EXP. 2							
WP744	1.4±0.45	2.1±0.5	1.5	1.6±0.5	2.3±0.9	1.4		
DOX	2.1±0.8	174±3 2	82.9	0.50±0.26	8.3±0.5	16.6		
WP744	0.48±0.0 3	4.07±1.8 5	8.5	0.37±0.11	0.60±0.10	1.6	0.89±0.10	2.41
DOX	0.60±0.1 3	>100	>167	0.53±0.25	6.87±0.58	13	>100	>189
WP783	0.43±0.2 1	1.1±0.8	2.6					
WP750	6.4	56±29	8.8					
DOX	0.50±0.0	>100	>200					

Table 4 shows data for fragmented DNA expressed as percentage of total DNA determined by quantitative apoptotic fragmentation assay in CEM leukemic cells incubated with drugs for 24 h. Results are from two independent studies carried out in duplicate.

TABLE 4.**Apoptotic Fragmentation by WP744 in Comparison to Doxorubicin.**

Concentration	WP744		DOX	
μM	% Total DNA	SE	% Total DNA	SE
0	0		0	
0.05	17.8	6.8		
0.1	31.3	12.9	-0.2	0.3
0.5	46.7	9.8	27.1	12.7
1			28.7	9
2			38.3	7.4

Direct comparisons of the cytotoxicity of WP791, WP790, WP786, WP785, WP784, WP783, WP778, WP775, WP774, WP758, WP757, WP756, WP755, WP750 with DOX indicated that the C-3' and C-4' substitutions drastically increase the potency of those compounds. Compound WP785 displayed similar profile of activity to doxorubicin and its RI was of 114 was clearly higher than that of other analogs tested.

EXAMPLE 5**Treatment of Tumors with Analogs of
Anthracyclines with Substitutions at C-3' or C-4' Sugars**

5 Treatment with the substituted anthracyclines having C-3'-alkylated anthracyclines modified at benzyl ring or having C-4'-substituted-benzylated anthracyclines of the present invention is similar to the treatment regimes of other anthracyclines and their derivatives, although some modifications to dosage may be warranted. For example, standard treatment with doxorubicin is described in
10 *Remington's Pharmaceutical Sciences* as follows.

Doxorubicin is administered intravenously to adults at 60 to 75 mg/m² at 21-day intervals or 25 to 30 mg/m² on each of 2 or 3 successive days repeated at 3- or 4-week intervals or 20 mg/m² once a week. The lowest dose should be used in elderly
15 patients, when there is prior chemotherapy or neoplastic marrow invasion or when the drug is combined with other myelopoietic suppressant drugs. The dose should be reduced by 50% if the serum bilirubin lies between 1.2 and 3 mg/dL and by 75% if above 3 mg/dL. The lifetime total dose should not exceed 550 mg/m² in patients with normal heart function and 400 mg/m² in patients with abnormal heart function and
20 400 mg/m² on each of 3 consecutive days, repeated every 4 weeks. Prescribing limits are as with adults. It has been reported that a 96-hour continuous infusion is as effective as and much less toxic than the same dose given by bolus injections.

Of course, modifications of the treatment regimes due to the unique nature of
25 the substituted anthracyclines having C-3'-alkylated anthracyclines modified at benzyl ring and having C-4'-substituted anthracyclines of the present invention are possible and well within the ability of one skilled in the art. Appropriate modifications may be ascertained by following the protocols in the following examples for *in vivo* testing and developments of human protocols.

EXAMPLE 6***In Vivo* Prevention of Tumor Development Using Analogs of Anthracyclines with Substitutions at C-3' or C-4' Sugars**

5 In an initial round of *in vivo* trials, a mouse model of human cancer with the histologic features and metastatic potential resembling tumors seen in humans (Katsumata *et al.*, 1995) is used. The animals are treated with analogs of anthracyclines having C-3'-substitutions as described or anthracycline analogs with
10 C-4'-substitutions synthesized in the present invention to determine the suppression of tumor development.

These novel analogs are tested *in vivo* for anti-tumor activity against murine leukemia L1210, P388, and P388 resistant to doxorubicin. In conjunction with these
15 studies, the acute and sub-acute toxicity is studied in mice (LD10, LD50, LD90). In a more advanced phase of testing, the anti-tumor activity of substituted anthracyclines synthesized in the present invention, are tested against human xenografts is assessed and cardiotoxicity studies performed is done in a rat or rabbit model.

20 These studies are based on the discovery that these analogs of anthracyclines have anti-cancer activity for MDR cancer cells. The invention provides a useful preventive and therapeutic regimen for patients with MDR tumors.

Two groups of mice of a suitable cancer model are treated with doses of
25 substituted anthracyclines with substitutions at the C-3'-sugar or having C-4'-substituted sugar moiety. Several combinations and concentrations of the substituted anthracyclines are tested. Control mice are treated with buffer only.

The effect of substituted anthracyclines on the development of breast tumors is
30 compared with the control group by examination of tumor size and histopathologic examination (breast tissue is cut and stained with hematoxylin and eosin) of breast tissue. With the chemopreventive potential of WP831, WP791, WP790, WP787,

WP786, WP785, WP784, WP780, WP778, WP775, WP774, WP758, WP757, WP756, WP755, WP799, WP797, WP794, WP783, WP750, WP744, WP727, WP571, WP418 and WP417, it is predicted that, unlike the control group of mice that develop tumors, the testing group of mice is resistant to tumor development.

5

EXAMPLE 7

Human Treatment with Analogs of Anthracyclines with Substitutions at C-3' or C-4' Sugars

10

This example describes a protocol to facilitate the treatment of cancer using substituted anthracyclines having C-3' alkylated anthracyclines modified at benzyl ring and having 4'-substituted-benzylated anthracyclines .

15

A cancer patient presenting, for example, an MDR cancer is treated using the following protocol. Patients may, but need not, have received previous chemo- radio- or gene therapeutic treatments. Optimally, the patient exhibits adequate bone marrow function (defined as peripheral absolute granulocyte count of $> 2,000/\text{mm}^3$ and platelet count of $100,000/\text{mm}^3$, adequate liver function (bilirubin 1.5mg/dl) and adequate renal function (creatinine 1.5mg/dl)).

20

Exemplary Protocol for the Treatment of Multi-Drug Resistant Cancer

25

A composition of the present invention is typically administered orally or parenterally in dosage unit formulations containing standard, well known, non-toxic physiologically acceptable carriers, adjuvants, and vehicles as desired. The term parenteral as used herein includes subcutaneous injections, intravenous, intramuscular, intra-arterial injection, or infusion techniques. The analogs of anthracyclines synthesized in the present invention may be delivered to the patient before, after, or concurrently with the other anti-cancer agents.

30

A typical treatment course may comprise about six doses delivered over a 7- to 21-day period. Upon election by the clinician, the regimen may be continued six

doses every three weeks or on a less frequent (monthly, bimonthly, quarterly etc.) basis. Of course, these are only exemplary times for treatment, and the skilled practitioner will readily recognize that many other time-courses are possible.

5 A major challenge in clinical oncology is that many cancers are multi-drug resistant. One goal of the inventors' efforts has been to find ways to improve the efficacy of chemotherapy. In the context of the present invention, the substituted anthracyclines have a surprising cytotoxicity against such cancers.

10 To kill MDR cancer cells using the methods and compositions described in the present invention, one will generally contact a target cell with a bisanthracycline of the present invention. These compositions are provided in an amount effective to kill or inhibit the proliferation of the cell.

15 In certain embodiments, it is contemplated that one would contact the cell with agent(s) of the present invention about every 6 hours to about every one week. In some situations, however, it may be desirable to extend the time period for treatment significantly where several days (2, 3, 4, 5, 6, 7 or more) to several weeks (1, 2, 3, 4, 5, 6, 7, or more) lapse between respective administrations.

20

 Regional delivery of anthracycline analogs is an efficient method for delivering a therapeutically effective dose to counteract the clinical disease. Likewise, the chemotherapy may be directed to a particular affected region. Alternatively, systemic delivery of active agents may be appropriate.

25

 The therapeutic composition of the present invention is administered to the patient directly at the site of the tumor. This is in essence a topical treatment of the surface of the cancer. The volume of the composition should usually be sufficient to ensure that the tumor is contacted by the substituted anthracyclines.

30

In one embodiment, administration simply entails injection of the therapeutic composition into the tumor. In another embodiment, a catheter is inserted into the site of the tumor, and the cavity may be continuously perfused for a desired period of time.

5

Clinical responses may be defined by acceptable measure. For example, a complete response may be defined by the disappearance of all measurable disease for at least a month, whereas a partial response may be defined by a 50% or greater reduction of the sum of the products of perpendicular diameters of all evaluable tumor nodules or at least one month with no tumor sites showing enlargement. Similarly, a mixed response may be defined by a reduction of the product of perpendicular diameters of all measurable lesions by 50% or greater, with progression in one or more sites.

10

15

Of course, the above-described treatment regimes may be altered in accordance with the knowledge gained from clinical trials such as those described in Example 7. Those of skill in the art are able to take the information disclosed in this specification and optimize treatment regimes based on the clinical trials described in the specification.

20

EXAMPLE 8

Clinical Trials with Analogs of Anthracyclines with Substitutions at C-3' or C-4' on the Sugar Moiety

25

This example is concerned with the development of human treatment protocols using the substituted anthracyclines. These compounds are of use in the clinical treatment of various MDR cancers in which transformed or cancerous cells play a role. Such treatment is a particularly useful tool in anti-tumor therapy, for example, in treating patients with ovarian, breast and lung cancers that are resistant to conventional chemotherapeutic regimens.

30

The various elements of conducting a clinical trial, including patient treatment and monitoring, is known to those of skill in the art in light of the present disclosure. The following information is being presented as a general guideline for use in establishing substituted anthracyclines drugs made by the use of this invention, in
5 clinical trials.

Patients with human metastatic breast and/or epithelial ovarian carcinoma, colon cancer leukemia, or sarcoma are chosen for clinical study. Measurable disease is not required, however the patient must have easily accessible pleural effusion and/or
10 ascites. Further the patients must carry tumors that express MDR phenotype. In an exemplary clinical protocol, patients may undergo placement of a Tenckhoff catheter, or other suitable device, in the pleural or peritoneal cavity and undergo serial sampling of pleural/peritoneal effusion. Typically, one will wish to determine the absence of known loculation of the pleural or peritoneal cavity, creatinine levels that are below 2
15 mg/dl, and bilirubin levels that are below 2 mg/dl. The patient should exhibit a normal coagulation profile.

In regard to the substituted anthracyclines drug administration, a Tenckhoff catheter, or alternative device, may be placed in the pleural cavity or in the peritoneal
20 cavity, unless such a device is already in place from prior surgery. A sample of pleural or peritoneal fluid can be obtained, so that baseline cellularity, cytology, LDH, and appropriate markers in the fluid (CEA, CA15-3, CA 125, p185) and in the cells (E1A, p185) may be assessed and recorded.

In the same procedure, substituted anthracyclines may be administered. The administration may be in the pleural/peritoneal cavity, directly into the tumor, or in a systemic manner. The starting dose may be 0.5mg/kg body weight. Three patients may be treated at each dose level in the absence of grade > 3 toxicity. Dose escalation may be done by 100% increments (0.5mg, 1mg, 2mg, 4mg) until drug related Grade II
25 30 toxicity is detected. Thereafter, dose escalation may proceed by 25% increments. The administered dose may be fractionated equally into two infusions, separated by 6 hours

if the combined endotoxin levels determined for the lot of bisanthracycline exceed 5EU/kg for any given patient.

The substituted anthracyclines may be administered over a short infusion time or at a steady rate of infusion over a 7- to 21-day period. The bisanthracycline infusion may be administered alone or in combination with the anti-cancer drug. The infusion given at any dose level is dependent upon the toxicity achieved after each. Hence, if Grade II toxicity was reached after any single infusion, or at a particular period of time for a steady rate infusion, further doses should be withheld or the steady rate infusion stopped unless toxicity improves. Increasing doses of substituted anthracyclines in combination with an anti-cancer drug is administered to groups of patients until approximately 60% of patients show unacceptable Grade III or IV toxicity in any category. Doses that are 2/3 of this value could be defined as the safe dose.

Physical examination, tumor measurements, and laboratory tests should, of course, be performed before treatment and at intervals of about 3-4 weeks later. Laboratory studies should include CBC, differential and platelet count, urinalysis, SMA-12-100 (liver and renal function tests), coagulation profile, and any other appropriate chemistry studies to determine the extent of disease, or determine the cause of existing symptoms. Also appropriate biological markers in serum should be monitored, e.g. CEA, CA 15-3, p185 for breast cancer, and CA 125, p185 for ovarian cancer.

To monitor disease course and evaluate the anti-tumor responses, it is contemplated that the patients should be examined for appropriate tumor markers every 4 weeks, if initially abnormal, with twice weekly CBC, differential and platelet count for the 4 weeks; then, if no myelosuppression has been observed, weekly. If any patient has prolonged myelosuppression, a bone marrow examination is advised to rule out the possibility of tumor invasion of the marrow as the cause of pancytopenia. Coagulation profile shall be obtained every 4 weeks. An SMA-12-100 shall be

performed weekly. Pleural/peritoneal effusion may be sampled 72 hours after the first dose, weekly thereafter for the first two courses, then every 4 weeks until progression or off study. Cellularity, cytology, LDH, and appropriate markers in the fluid (CEA, CA15-3, CA 125, p185) and in the cells (p185) may be assessed. For an example of an evaluation profile, see Table 3. When measurable disease is present, tumor measurements are to be recorded every 4 weeks. Appropriate radiological studies should be repeated every 8 weeks to evaluate tumor response. Spirometry and DLCO may be repeated 4 and 8 weeks after initiation of therapy and at the time study participation ends. A urinalysis may be performed every 4 weeks.

Clinical responses may be defined by acceptable measure. For example, a complete response may be defined by the disappearance of all measurable disease for at least a month. Whereas a partial response may be defined by a 50% or greater reduction of the sum of the products of perpendicular diameters of all evaluable tumor nodules or at least one month with no tumor sites showing enlargement. Similarly, a mixed response may be defined by a reduction of the product of perpendicular diameters of all measurable lesions by 50% or greater with progression in one or more sites.

Table 5. EVALUATIONS BEFORE AND DURING THERAPY

EVALUATIONS	PRE-STUDY	TWICE WEEKLY	WEEKLY	EVERY 4 WEEKS	EVERY 8 WEEKS
History	X			X	
Physical	X			X	
Tumor Measurements	X			X	
CBC	X	X ¹	X		
Differential	X	X ¹	X		
Platelet Count	X	X ¹	X		
SMA12-100 (SGPT, Alkaline Phosphatase, Bilirubin, Alb/Total Protein)	X		X		
Coagulation Profile	X			X	
Serum Tumor markers (CEA, CA15-3, CA-125, Her-2/neu)	X			X ³	
Urinalysis	X			X	
X-rays:					
chest	X		X ⁴		
others	X				X
Pleural/Peritoneal Fluids: (cellularity, cytology, LDH, tumor markers, E1A, HER-2/neu)	X		X ⁵	X	
Spirometry and DLCO	X			X ⁶	X ⁶

¹ For the first 4 weeks, then weekly, if no myelosuppression is observed.

² As indicated by the patient's condition.

³ Repeated every 4 weeks if initially abnormal.

⁴ For patients with pleural effusion, chest X-rays may be performed at 72 hours after first dose, then prior to each treatment administration.

⁵ Fluids may be assessed 72 hours after the first dose, weekly for the first two courses and then every 4 weeks thereafter.

⁶ Four and eight weeks after initiation of therapy.

5

10

All of the compositions and methods disclosed and claimed herein can be made and executed without undue experimentation in light of the present disclosure. While the compositions and methods of this invention have been described in terms of preferred embodiments, it will be apparent to those of skill in the art that variations may be applied to the compositions and methods and in the steps or in the sequence of steps of the method described herein without departing from the concept, spirit and scope of the invention. More specifically, it will be apparent that certain agents which are both chemically and physiologically related may be substituted for the agents described herein while the same or similar results would be achieved. All such similar substitutes and modifications apparent to those skilled in the art are deemed to be within the spirit, scope and concept of the invention as defined by the appended claims.

REFERENCES

The following references, to the extent that they provide exemplary procedural or other details supplementary to those set forth herein, are specifically incorporated herein by reference.

- 5 Arcamone F., Doxorubicin. Anti-Cancer Antibiotics. New York: Academic Press, 1981.
- Bell *et al.*, *J. Clin. Oncol.*, 3:311, 1985
- Bertino, *J. Clin. Oncol.*, 3:293, 1985
- Bodley, *et al.*, *Cancer Res.*, 49:5969-5978, 1989.
- 10 Bodley *et al.*, *Cancer Res.*, 49:5969.
- Booser DJ and Hortobagyi GN., "Anthracycline Antibiotics in Cancer Therapy. Focus on Drug Resistance", *Drugs*, 47:223-258, 1994.
- Bradley *et al.*, *Biochem. Biophys. Acta.*, 948:87, 1988.
- Capranico *et al.*, "Sequence-Selective Topoisomerase II Inhibition by Anthracycline
- 15 Derivatives in SV40 DNA: Relationship with DNA Binding Affinity and Cytotoxicity", *Biochem*, 29:562-569, 1990.
- Curt *et al.*, *Cancer Treat. Rep.*, 68:87, 1984.
- Danks *et al.*, *Cancer Res.*, 47:1297, 1987.
- Denny *et al.*, "Potential Anti-tumor Agents. 39. Anilino Ring Geometry of
- 20 Amsacrine and Derivatives: Relationship to DNA binding and Anti-tumor Activity", *J. Med. Chem.*, 26(11):1625-1630, 1983.
- Dervan P. *Science*, 232:464-471, 1986.
- Fojo *et al.*, *P.N.A.S.*, 84:265, 1987.
- Ganapath, *et al.*, *Br. J. Cancer*, 60:819, 1989.
- 25 Goldie *et al.*, *Cancer Res.*, 44:3643, 1984.
- Goldie *et al.*, *Cancer Treat. Rep.*, 63:1727, 1979.
- Green *et al.*, *J. Immunol. Methods*, 70:257-268, 1984.
- Gros *et al.*, *Nature*, 323:728, 1986
- Israel *et al.*, 1987.
- 30 Israel *et al.*, *Cancer Chemother. Pharmacol.*, 25:177, 1989.

- Katsumata *et al.*, "Prevention of Breast Tumor Development *In Vivo* by Down-Regulation of the p185^{neu} Receptor", *Nature Med.*, 1:644-648, 1995.
- Kolate, *Science*, 231:220, 1986.
- Lown JW, "Targeting the DNA Minor Groove for Control of Biological Function:
5 Progress, Challenges and Prospects", *Chemtracts - Org. Chem.*, 6:205-237, 1993.
- Lown JW, "Anthracycline and Anthracenedione-Based Anticancer Agents, Bioactive Molecules, Vol. 6, Amsterdam: Elsevier, 1988.
- Norris *et al.*, "Expression of the gene for multi-drug-resistance-associated protein and
10 outcome in patients with neuroblastoma", *N. Engl. J. Med.*, 334:231-238, 1996.
- Nowell, *Cancer Res.*, 46:2203, 1986.
- Pommier *et al.*, *Cancer Res.*
- Priebe *et al.*, "3'-Hydroxy-esorubicin substituted at C- 2'", *J. Antibiot.*, 45:386-393, 1992.
- 15 Priebe *et al.*, "Removal of the basic center from doxorubicin partially overcomes multi-drug resistance and decreases cardiotoxicity", *Anti-Cancer Drugs*, 4:37-48, 1993.
- Priebe W, "Anthracycline Antibiotics. Novel Analogues, Methods of Delivery, and Mechanisms of Action", Washington, DC: American Chemical Society, 1995.
- 20 Priebe W. "Mechanism of Action-Governed Design of Anthracycline Antibiotics: A "Turn-Off/Turn-On" Approach", *Current Pharmaceutical Design*, 1:51-68, 1995.
- Rubinstein *et al.*, "Comparison of *In Vitro* Anti-Cancer-Drug-Screening Data Generated with A Tetrazolium Assay Versus a Protein Assay Against a
25 Diverse Panel of Human Tumor Cell Lines", *J. Nat'l. Cancer Inst.*, 82:1113-1120, 1990.
- Stryer, "Biochemistry", Freeman and Co., 1981.
- Sweatman *et al.*, *J. Cell. Pharmacol.*, 1:95-102.
- Traganos *et al.*, *Cancer Res.*, 45:6273, 1985.
- 30 Wakelin LPG, "Medicinal Research Rev.", 6:275-340, 1986.
- Young *et al.*, *N. Engl. J. Med.*, 312:692, 1985.


U. S. Patent No. 4,263,428 April 21, 1981.

U.S. Patent No. 4,345,070, August 17, 1982.

U.S. Patent No. 4,438,105, March 20, 1984.

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. A substituted anthracycline having the formula:



5

wherein, R^1 denotes an alkyl chain, a $(-COCH_2R^{13})$ group or a $(C(OH)-CH_2R^{13})$;

wherein, R^{13} is a hydrogen $(-H)$ group, a hydroxyl group $(-OH)$; a methoxy group $(-OCH_3)$,
 an alkoxy group having 1-20 carbon atoms, an alkyl group having 1-20 carbon atoms, an
 10 aryl group having 1-20 carbon atoms, a fatty acyl group having the general structure
 $-O-CO(CH_2)_nCH_3$, wherein n = an integer from 1 to about 20, a fatty acyl group having the
 general structure $-O-CO(CH_2)_l(CH=CH)_m(CH_2)_nCH_3$, wherein l is an integer between 1 to
 3, m is an integer between 1 and 6, and n is an integer between 1 and 9, a $-OCO-(CH_2)_n-$
 CH_2NH_2 , or $OCO-(CH_2)_n-CO_2H$:

15

each of R^2 and R^3 is, independently of the other, a hydrogen $(-H)$, a hydroxyl group $(-OH)$
 or a methoxy group $(-OCH_3)$;

R^4 is a hydrogen (-H) group, a methoxy group (-OCH₃), a hydroxyl group (-OH) or a halide;

each of Y^1 and Y^2 is, independently of the other, a double bonded oxygen, sulphur, or
5 nitrogen atom;

Z is a -H, -OH, a -CO₂H group or a -CO₂R group;


R^7 , R^8 , are, independently, -H, -OH, a halide, -OR¹⁹, -SH, -SR¹⁹, -NH₂, -NHR¹⁹, -N(R¹⁹)₂
10 or -CH₃, and R^7 can additionally be a saccharide, wherein R^{19} is an alkyl chain, an alkylating moiety, a cycloalkyl chain, a cyclic ring, or a hydrogen;

R^9 is an -H, -CH₃, alkyl, aryl, CH₂OH, or a CH₂F group;

15 R^{10} , R^{11} and R^{12} are, independently, -H, -OH, a halide, -OR, -SH, -SR, -NH₂, -NHR, -N(R)₂, or a -CH₃;

one of R^5 and R^6 is a -H;

20 one of R^5 and R^6 is a X-alkyl-aromatic-ring (-XAAR) substituent, wherein, A is an alkyl group and wherein, AR is an substituted phenyl ring, or a substituted five-member ring, a heteroatomic five-member ring, or a heteroatomic six-member ring, of the form:



;

wherein, $R^{14} - R^{18}$ are independently a (-H) group, a hydroxyl group (-OH), a methoxy group (-OCH₃), a nitro group (-NO₂), an amine group (-NH₂), a halide, an alkoxy group having 1-20 carbon atoms, an alkyl group having 1-20 carbon atoms, an aryl group having 1-20 carbon atoms, an alkyl-amino group, an alkyl-thio group, a cyano group (CN, SCN), an -CO₂H group or an -CO₂R group; and


the aromatic ring is unsubstituted, disubstituted, trisubstituted, tetrasubstituted or pentasubstituted; and

X is a -N, a -S, a -SO, or -SO₂ group; and


A is (CH₂)_n where n = 0-10;

wherein, if R⁵ is a XAAR substituent R⁶ is not and if R⁶ is a XAAR substituent R⁵ is not.


2. The substituted anthracycline of claim 1 having the formula:




3. The substituted anthracycline of claim 1 having the formula:




4. The substituted anthracycline of claim 1 having the formula:




5. The substituted anthracycline of claim 1 having the formula:




6. The substituted anthracycline of claim 1 having the formula:




7. The substituted anthracycline of claim 1 having the formula:




- 5 8. The substituted anthracycline of claim 1 having the formula:




9. The substituted anthracycline of claim 1 having the formula:




10. The substituted anthracycline of claim 1 having the formula:




11. The substituted anthracycline of claim 1 having the formula:




12. The substituted anthracycline of claim 1 having the formula:




13. The substituted anthracycline of claim 1 having the formula:



14. The substituted anthracycline of claim 1 having the formula:




15. The substituted anthracycline of claim 1 having the formula:




WP778

16. The substituted anthracycline of claim 1 having the formula:



WP780

17. A substituted anthracycline having the formula:



5 wherein, R^1 denotes an alkyl chain, a $(-COCH_2R^{13})$ group, or a $(C(OH)-CH_2R^{13})$ group;

wherein, R^{13} is a hydrogen (-H) group, a hydroxyl group (-OH), a methoxy group (-OCH₃), an alkoxy group having 1-20 carbon atoms, an alkyl group having 1-20 carbon atoms, an aryl group having 1-20 carbon atoms, a fatty acyl group having the general structure
 10 $-O-CO(CH_2)_nCH_3$, wherein n = an integer from 1 to about 20, a fatty acyl group having the general structure $-O-CO(CH_2)_l(CH=CH)_m(CH_2)_nCH_3$, wherein l is an integer between 1 to 3, m is an integer between 1 and about 6, and n is an integer between 1 to about 9, $-OCO-(CH_2)_n-CH_2NH_2$ or $OCO-(CH_2)_n-CO_2H$;

15 each of R^2 and R^3 is, independently of the other, a hydrogen (-H), a hydroxyl group (-OH) or a methoxy group (-OCH₃);

R^4 is a hydrogen (-H) group, a methoxy group (-OCH₃), a hydroxyl group (-OH) or a halide;

each of Y^1 and Y^2 is, independently of the other, a double bonded oxygen, sulphur or nitrogen atom;

5 Z is a $-H$, $-OH$, a $-CO_2H$ group or a $-CO_2R$ group;

R^5 and R^6 , are, independently, $-H$, $-OH$, a halide, $-OR^{19}$, $-SH$, $-SR^{19}$, $-NH_2$, $-NHR^{19}$, $-N(R^{19})_2$, or $-CH_3$, and R^6 can additionally be an alkylating moiety, wherein R^{19} is an alkyl chain, an alkylating moiety, a cycloalkyl chain, a cyclic ring or a hydrogen,

10

R^9 is an $-H$, $-CH_3$, alkyl, aryl, CH_2OH or a CH_2F group;


R^{10} , R^{11} and R^{12} are, independently, $-H$, $-OH$, a halide, $-OR$, $-SH$, $-SR$, $-NH_2$, $-NHR$, $-N(R)_2$ or $-CH_3$;

15

one of R^7 and R^8 is a $-H$;

one of R^7 and R^8 is a X-alkyl aromatic-ring (-XAAR) substituent, wherein, A is an alkyl group and wherein, AR is an unsubstituted phenyl ring, a substituted phenyl ring, a substituted five-member ring or a heteroatomic five-member ring, of the general form;

20



;

wherein, R^{14} - R^{18} are independently a (-H) group, a hydroxyl group (-OH), a methoxy group (-OCH₃), a nitro group (-NO₂), an amine group (-NH₂), a halide, an alkoxy group having 1-20 carbon atoms, an alkyl group having 1-20 carbon atoms, an aryl group having 1-20 carbon atoms, an alkyl-amino group, an alkyl-thio group, a cyano group (CN, SCN),
 5 an -CO₂H group or an -CO₂R group; and

the aromatic ring is unsubstituted, disubstituted, trisubstituted, tetrasubstituted or pentasubstituted; and

X is a -O, -N, -S, -SO or a -SO₂ group; and


10

A is (CH₂)_n, where n = 0-10;


wherein if R^7 is a XAAR substituent R^8 is not and if R^8 is a XAAR substituent R^7 is not.

18. The substituted anthracycline of claim 17 having the formula:

15




19. The substituted anthracycline of claim 17 having the formula:




10

20. The substituted anthracycline of claim 17 having the formula:




20

21. The substituted anthracycline of claim 17 having the formula:



WP787

- 5 22. The substituted anthracycline of claim 17 having the formula:




10

WP750


23. The substituted anthracycline of claim 17 having the formula:

15




WP783

24. The substituted anthracycline of claim 17 having the formula:




10

25. The substituted anthracycline of claim 17 having the formula:




26. The substituted anthracycline of claim 17 having the formula:



27. The substituted anthracycline of claim 17 having the formula:

25



WP764

- 5 28. A method for the synthesis of amine containing anthracyclines comprising:
- (a) obtaining a substituted sugar azide, wherein the azido substitution is at the 1', 2', 3', 4' or 5' position on the sugar, said azide serving as a masked and neutral form of amine substituent;
 - (b) subjecting said azide to a coupling reaction with an anthracyclinone; and
 - (c) subjecting said azide to a reduction reaction;
- 10 wherein an amine containing anthracycline is produced.
29. The method of claim 28 wherein the amine containing anthracycline is a 14-hydroxy analog of anthracyclines.
- 15 30. The method of claim 28 wherein the amine containing anthracycline is an analog of doxorubicin.
31. The method of claim 28 wherein the amine containing anthracycline is an analog of
- 20 daunorubicin.
32. The method of claim 28 wherein the amine containing anthracycline is WP744.

33. The method of claim 28 wherein the amine containing anthracycline is WP764.

34. A substituted anthracycline according to either claim 1 or claim 17, substantially as
5 herein described with reference to any one or more of the Examples and/or accompanying
Figures.

35. A method for the synthesis of amine containing anthracyclines according to claim
28, substantially as herein described with reference to any one or more of the Examples
10 and/or accompanying Figures.


Dated this thirteenth day of October 2004

Waldemar Priebe, Izabella Fokt, Marta
Krawczyk, Roman Perez-Soler, Teresa
Przewłoka, Piotr Skibicki, Grzegorz
Grynkiewicz


Patent Attorneys for the Applicant:

F B RICE & CO


1 / 22



DOXORUBICIN



DAUNORUBICIN

FIG. 1. Structure of Doxorubicin and Daunorubicin

WP831

FIG. 2 Structure of WP831




FIG. 3 Structure of WP791




FIG. 4 Structure of WP790

3 / 22




FIG. 5 Structure of WP787






FIG. 6 Structure of WP786

4 / 22

**FIG. 7** Structure of WP785**FIG. 8** Structure of WP784




FIG. 9 Structure of WP780




FIG. 10 Structure of WP778




FIG. 11 Structure of WP775




FIG. 12 Structure of WP774




FIG. 13 Structure of WP758




FIG. 14 Structure of WP757




FIG. 15 Structure WP756




FIG. 16 Structure WP755




FIG. 17 Structure of WP799




FIG. 18 Structure of WP797




FIG. 19 Structure of WP794




FIG. 20 Structure of WP783






FIG. 21 Structure of WP750




WP744

FIG. 22 Structure of WP744



WP727

FIG. 23 Structure of WP727



WP567

FIG. 24 Structure of WP567




FIG. 25 Structure of WP793




FIG. 26 Structure of WP819




FIG. 27 Structure of WP820




FIG. 28 WP821




FIG. 29 Structure of WP822




FIG. 30 Structure of WP823




FIG. 31 Structure of WP824




FIG. 32 Structure of WP825





FIG. 33 Structure of WP764



WP765

FIG. 34 Structure of WP765

FIG. 35

SYNTHESIS OF 4'-O-BENZYLATED ANTHRACYCLINES FROM 4-O-BENZYLATED GLYCALS




FIG. 36

SYNTHESIS OF 4'-O-BENZYL-3'-AMINO-ANTHRACYCLINES





FIG. 37

SYNTHESIS OF 4-O-BENZYL-GLYCALS AND ITS
3-O-DERIVATIVES

The same procedure was used for the synthesis of other O-benzylated glycals.

FIG. 38

SYNTHESIS OF 4'-O-BENZYL-3'-DEAMINO-ANTHRACYCLINES
FROM 1-O-SILYLATED 4-O-BENZYL-HEXOPYRANOSES





FIG. 39

Examples of Selective Alkylation of Acylated Glycals




L-Rhamnal: WP- $R^1=OAc$, $R^2=H$, $R^3=OAc$, $R^4=Me$, $R^5=OH$, $R^6=H$, $R^7=OBn$

L-Rhamnal: WP- $R^1=OAc$, $R^2=H$, $R^3=OAc$, $R^4=Me$, $R^5=OAc$, $R^6=H$, $R^7=OBn$

L-Fucal: WP- $R^1=OAc$, $R^2=OAc$, $R^3=H$, $R^4=Me$, $R^5=OH$, $R^6=OBn$, $R^7=H$

L-Fucal: WP- $R^1=OAc$, $R^2=OAc$, $R^3=H$, $R^4=Me$, $R^5=OAc$, $R^6=OBn$, $R^7=H$

L-Xylal: WP- $R^1=OAc$, $R^2=H$, $R^3=OAc$, $R^4=H$, $R^5=OAc$, $R^6=H$, $R^7=OBn$



D-Glucal: WP- $R^1=OAc$, $R^2=OAc$, $R^3=H$, $R^4=CH_2OAc$, $R^5=OH$, $R^6=OBn$, $R^7=H$, $R^8=CH_2OH$

D-Glucal: WP- $R^1=OAc$, $R^2=OAc$, $R^3=H$, $R^4=CH_2OAc$, $R^5=OAc$, $R^6=OBn$, $R^7=H$, $R^8=CH_2OAc$

D-Galactal: WP- $R^1=OAc$, $R^2=H$, $R^3=OAc$, $R^4=CH_2OAc$, $R^5=OH$, $R^6=H$, $R^7=OBn$, $R^8=CH_2OH$

D-Galactal: WP- $R^1=OAc$, $R^2=H$, $R^3=OAc$, $R^4=CH_2OAc$, $R^5=Ac$, $R^6=H$, $R^7=OBn$, $R^8=CH_2OA$

D-Arabinol: WP- $R^1=OAc$, $R^2=H$, $R^3=OAc$, $R^4=H$, $R^5=OH$, $R^6=H$, $R^7=OH$, $R^8=H$

FIG. 40

Synthesis of 3'-azido-sugars and its 4-O-benzylated derivatives




FIG. 41

ALTERNATIVE SYNTHESIS OF 3-AZIDO-4-O-BENZYL-DAUNOSAMINE FROM 3-AZIDOACOSAMINE





FIG. 42 Structure of WP735




FIG. 43 Structure of WP736