

A&A Ref: 151302

PUBLICATION PARTICULARS AND ABSTRACT
(Section 32(3)(a) - Regulations 22(1)(g) and 31)

21	01	PATENT APPLICATION NO	22	LODGING DATE	43	ACCEPTANCE DATE
----	----	-----------------------	----	--------------	----	-----------------

2004/8664

26 October 2004

5.2.07

51	INTERNATIONAL CLASSIFICATION	NOT FOR PUBLICATION
----	------------------------------	---------------------

A61K

CLASSIFIED BY: ISA

71	FULL NAME(S) OF APPLICANT(S)
----	------------------------------

Endocyte, Inc.

72	FULL NAME(S) OF INVENTOR(S)
----	-----------------------------

LEAMON, Christopher Paul

PARKER, Matthew A.

EARLIEST PRIORITY CLAIMED	COUNTRY		NUMBER		DATE	
	33	US	31	60/378,571	32	6 May 2002

NOTE: The country must be indicated by its International Abbreviation - see schedule 4 of the Regulations

54	TITLE OF INVENTION
----	--------------------

Vitamin-targeted imaging agents

57	ABSTRACT (NOT MORE THAN 150 WORDS)
----	------------------------------------

NUMBER OF SHEETS	79
------------------	----

One sheet(s) containing the abstract is/are attached.

If no classification is furnished, Form P.9 should accompany this form.
The figure of the drawing to which the abstract refers is attached.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
13 November 2003 (13.11.2003)

PCT

(10) International Publication Number
WO 03/092742 A1

(51) International Patent Classification: A61K 49/00 (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BC, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, IL, IN, IS, JP, KE, KG, KP, KR, KZ, L, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(21) International Application Number: PCT/US03/14379 (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, T, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IL, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(22) International Filing Date: 6 May 2003 (06.05.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data: 60/378,571 6 May 2002 (06.05.2002) US

(71) Applicant (for all designated States except US): ENDOCYTIC, INC. [US/US]; 1205 Kent Avenue, West Lafayette, IN 47906 (US)

(72) Inventors; and

(75) Inventors/Applicants (for US only): LEAMON, Christopher, Paul [US/US]; 5830 Farm Ridge Road, West Lafayette, IN 47906 (US). PARKER, Matthew, A. [US/US]; 2350 Sixth Avenue, Apt. 5F, San Diego, CA 92101 (US).



(74) Agent: LAMMERT, Steven, R.; Barnes & Thornburg, 11 South Meridian Street, Indianapolis, IN 46204 (US).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: VITAMIN-TARGETED IMAGING AGENTS



(57) Abstract: The invention relates to compounds and methods for targeting radionuclide-based imaging agents to cells having receptors for a vitamin, or vitamin receptor binding derivative or analog thereof, by using such a vitamin as the targeting ligand for the imaging agent. The invention provides a compound of the formula (I) or (II) for use in such methods. In the compound, V is a vitamin that is a substrate for receptor-mediated transmembrane transport *in vivo*, or a vitamin receptor binding derivative or analog thereof, L is a divalent linker, R is a side chain of an amino acid of the formula H₂NCHR₁COOH, M is a cation of a radionuclide, n is 1 or 0, K is 1 or 0, and the compound can be in a pharmaceutically acceptable carrier therefor. The vitamin-based compounds can be used to target radionuclides to cells, such as a variety of tumor cell types, for use in diagnostic imaging of the targeted cells.

WO 03/092742 A1

-1-

VITAMIN-TARGETED IMAGING AGENTS

FIELD OF THE INVENTION

The invention relates to compounds and methods for targeting an imaging agent to cells of an animal. More particularly, radionuclide-based imaging agents are targeted to cells having receptors for a vitamin by using such a vitamin, or a vitamin receptor binding derivative or an analog thereof, as the targeting ligand for the imaging agent.

BACKGROUND AND SUMMARY OF THE INVENTION

Transmembrane transport is a critical cellular function. Because practitioners have recognized the importance of transmembrane transport to many areas of medical and biological science, including drug therapy and gene transfer, there have been significant research efforts directed to the understanding and application of such processes. Thus, for example, transmembrane delivery of nucleic acids has been attempted through the use of protein carriers, antibody carriers, liposomal delivery systems, electroporation, direct injection, cell fusion, viral carriers, osmotic shock, and calcium-phosphate mediated transformation. However, many of those techniques are limited both by the types of cells in which transmembrane transport occurs and by the conditions required for successful transmembrane transport of exogenous molecules. Furthermore, many of these techniques are limited by the type and size of the exogenous molecule that can be transported across the cell membrane without loss of bioactivity.

One mechanism for transmembrane transport of exogenous molecules having wide applicability is receptor-mediated endocytosis. Advantageously, receptor-mediated endocytosis occurs both *in vivo* and *in vitro*. Receptor-mediated endocytosis involves the movement of ligands bound to membrane receptors into the interior of an area bounded by the membrane through invagination of the membrane. The process is initiated or activated by the binding of a receptor-specific ligand to the receptor. Many receptor-mediated endocytic systems have been characterized, including those resulting in internalization of galactose, mannose, mannose 6-phosphate, transferrin, asialoglycoprotein, folate, transcobalamin (vitamin B₁₂), α -2

-2-

macroglobulins, insulin, and other peptide growth factors such as epidermal growth factor (EGF).

Receptor mediated endocytosis has been utilized for delivering exogenous molecules such as proteins and nucleic acids to cells. Generally, a specific ligand is chemically conjugated by covalent, ionic, or hydrogen bonding to an exogenous molecule of interest, forming a conjugate molecule having a moiety (the ligand portion) that is still recognized in the conjugate by a target receptor. Using this technique the phototoxic protein psoralen has been conjugated to insulin and internalized by the insulin receptor endocytotic pathway (Gasparro, Biochem. Biophys. Res. Comm. 141(2), pp. 502-509, Dec. 15, 1986); the hepatocyte specific receptor for galactose terminal asialoglycoproteins has been utilized for the hepatocyte-specific transmembrane delivery of sialoorosomucoid-poly-L-lysine non-covalently complexed to a plasmid (Wu, G. Y., J. Biol. Chem., 262(10), pp. 4429-4432, 1987); the cell receptor for EGF has been utilized to deliver poly-nucleotides covalently linked to EGF to the cell interior (Mayers, European Patent Application 86810614.7, published Jun. 6, 1988); the intestinal situated cellular receptor for the organometallic vitamin B₁₂-intrinsic factor complex has been used to mediate delivery of a drug, a hormone, a bioactive peptide and an immunogen complexed with vitamin B₁₂ to the circulatory system after oral administration (Russell-Jones et al., European Patent Application 86307849.9, published Apr. 29, 1987); the mannose-6-phosphate receptor has been used to deliver low density lipoproteins to cells (Murray, G. J. and Neville, D. M., Jr., J. Biol. Chem., Vol. 255 (24), pp. 1194-11948, 1980); the cholera toxin binding subunit receptor has been used to deliver insulin to cells lacking insulin receptors (Roth and Maddox, J. Cell. Phys. Vol. 115, p. 151, 1983); and the human chorionic gonadotropin receptor has been employed to deliver a ricin α -chain coupled to HCG to cells with the appropriate HCG receptor (Oeltmann and Heath, J. Biol. Chem., vol. 254, p. 1028 (1979)).

In one embodiment the present invention involves the transmembrane transport of a radionuclide-based imaging agent across a membrane having receptors for a vitamin, or a vitamin receptor binding derivative or analog thereof. A cell membrane bearing vitamin receptors, or receptors for vitamin derivatives or analogs, is contacted with a vitamin-imaging agent conjugate for a time sufficient to initiate

-3-

and permit transmembrane transport of the conjugate, and the biodistribution of the vitamin-imaging agent conjugate in the animal is monitored. In another embodiment, the vitamin/vitamin derivative or analog targeting moiety simply binds to a cell surface vitamin receptor to concentrate the chelated radionuclide on the cell surface.

5 The invention takes advantage of (1) the location of vitamin receptors and (2) the associated receptor-mediated endocytic processes. For example, the invention takes advantage of the unique expression, overexpression, or preferential expression of vitamin receptors, transporters, or other surface-presented proteins that specifically bind vitamins, or derivatives or analogs thereof, on tumor cells or other 10 cell types which overexpress such receptors. Accordingly, the invention can be used to detect cells, such as tumor cells or other cell types, which overexpress vitamin receptors, or receptors for vitamin derivatives or analogs, by taking advantage of the receptor-mediated endocytic processes that occur when such cells are contacted with the vitamin-imaging agent conjugate.

15 Vitamin receptors, such as the high-affinity folate receptor (FR) is expressed at high levels, for example, on cancer cells. Epithelial cancers of the ovary, mammary gland, colon, lung, nose, throat, and brain have all been reported to express elevated levels of the FR. In fact, greater than 90% of all human ovarian tumors are known to express large amounts of this receptor. Thus, the present invention can be 20 used for the diagnostic imaging of a variety of tumor types, and of other cell types involved in disease states.

25 Radionuclide chelators complexed to ligands have been used as non-invasive probes for diagnostic imaging purposes. For example, vasoactive intestinal peptide, somatostatin analogs, and monoclonal antibodies have been used as ligands to localize radionuclides to cells, such as tumor cells. Monoclonal antibodies, and various fragments thereof, initially received the most attention because it was believed that precise tumor-specific targeting might be achieved using monoclonal antibodies as targeting ligands. Unfortunately, this approach was problematic because 30 i) antibodies have prolonged circulation times due to their large size which is unfavorable for imaging purposes, ii) antibodies are expensive to produce, iii) antibodies can be immunogenic, and, accordingly, must be humanized when multiple doses are used, and iv) tumor to non-target tissue ratios (⁹⁹T/NT) of antibody-

-4-

linked radionuclides are sub-optimal. Thus, the focus has recently been directed to the use of smaller tumor-specific ligands that do not have such limitations.

Vitamins, such as folic acid, have been used for the targeting of imaging agents to tumor cells, and are advantageous because of their small size. The 5 first folic acid-based targeting complex described for *in vivo* tumor imaging was a histamine derivative containing ^{125}I odine. This complex was not considered a relevant clinical candidate because of the long-lived ^{125}I radionuclide component. Subsequently, a deferoxamine-folate conjugate for tumor imaging was developed 10 (deferoxamine chelates ^{67}Ga , a gamma-emitting radionuclide that has a 78 hour half-life). Hepatobiliary clearance was noted with this conjugate and, thus, preclinical development was stopped due to anticipated problems in accurately imaging regional abdominal locations. This obstacle was overcome, however, by replacing the deferoxamine chelator with diethylenetriamine pentaacetic acid (DTPA), an efficient chelator of ^{111}In (68 hour half life). The primary route of elimination of ^{111}In -DTPA-folate was confirmed to be through the kidneys.

More recently, $^{99\text{m}}\text{Tc}$ has been adopted as the preferred radionuclide for diagnostic imaging, because i) the radionuclide is easily obtained from commercially available ^{99}Mo - $^{99\text{m}}\text{Tc}$ generators, ii) the cost of producing large amounts of $^{99\text{m}}\text{Tc}$ is insignificant compared to the cost of producing ^{111}In , and iii) $^{99\text{m}}\text{Tc}$ has a much shorter (6 hour) half life, which allows higher radionuclide doses to be administered, yielding higher resolution images without the risk of hazardous radiation exposure to vital organs.


Several folate-based $^{99\text{m}}\text{Tc}$ conjugates have been developed. For example, folate conjugates of $^{99\text{m}}\text{Tc}$ -6-hydrazinonicotinamido-hydrazido (HYNIC; Guo, et al., *J. Nucl. Med.*, 40(9): 1563-1569 (1999)), $^{99\text{m}}\text{Tc}$ -12-amino-3,3,9,9-tetramethyl-5-oxa-4,8 diaza-2,10-decanedione dioxime (OXA) (Linder, et al., *Soc. Nucl. Med., Proc. 47th Annual Meeting*, 2000, 41(5): 119P), $^{99\text{m}}\text{Tc}$ -ethylenedicycysteine (Egan, et al., *Cancer Biother. & Radiopharm.*, 13(6): 427-435 (1998)), and $^{99\text{m}}\text{Tc}$ -DTPA-folate (Mathias, et al., *Bioconjug. Chem.*, 11(2): 253-257 (2000)) have shown promising *in vivo* tumor uptake qualities. However, there is a need for alternative vitamin-based $^{99\text{m}}\text{Tc}$ conjugates, or vitamin-based conjugates employing other radionuclides, that exhibit optimal tumor to non-target tissue ratios (T/NT) and are

-5-

eliminated through the kidneys. Such vitamin-based conjugates should be suitable for clinical development as tumor imaging agents, and for the diagnosis of other disease states.


In one embodiment is provided a compound of the formula

5



wherein V is a vitamin, or a vitamin receptor binding derivative or analog thereof, L is a divalent linker, R is a side chain of an amino acid of the formula $H_2N-CH(R)COOH$, M is a cation of a radionuclide, n is 1 or 0, and k is 1 or 0. The vitamin is a substrate for receptor-mediated transmembrane transport in vivo.


10 In another embodiment is provided a composition for diagnostic imaging comprising a compound of the formula



15 wherein V is a vitamin, or a vitamin receptor binding derivative or analog thereof, L is a divalent linker, R is a side chain of an amino acid of the formula $H_2N-CH(R)COOH$, M is a cation of a radionuclide, n is 1 or 0, and a pharmaceutically acceptable carrier therefor. The vitamin is a substrate for receptor-mediated transmembrane transport in vivo.


20 In yet another embodiment a method is provided of imaging a population of cells in an animal, wherein the cells are characterized by a vitamin receptor on the surface of the cells. The method comprises the steps of administering to the animal an effective amount of a composition comprising a compound of the formula

-6-




wherein V is a vitamin, or a receptor binding derivative or analog thereof, specific for the cell surface vitamin receptor, L is a divalent linker, R is a side chain of an amino acid of the formula $H_2N\text{CHRCOOH}$, M is a cation of a radionuclide, n is 1 or 0, and a pharmaceutically acceptable carrier therefor, and monitoring the biodistribution of the compound in the animal.

In another embodiment a compound is provided of the formula



wherein V is a vitamin that is a substrate for receptor-mediated transmembrane transport in vivo, or a vitamin receptor binding derivative or analog thereof, L is a divalent linker, R is a side chain of an amino acid of the formula $H_2N\text{CHRCOOH}$, M is a cation of a radionuclide, n is 1 or 0, and k is 1 or 0.


In still another embodiment, a composition for diagnostic imaging is provided comprising a compound of the formula



wherein V is a vitamin that is a substrate for receptor-mediated transmembrane transport in vivo, or a vitamin receptor binding derivative or analog thereof, L is a divalent linker, R is a side chain of an amino acid of the formula $H_2N\text{CHRCOOH}$, M is a cation of a radionuclide, n is 1 or 0, and a pharmaceutically acceptable carrier therefor.

In yet another embodiment, a method of imaging a population of cells in an animal is provided wherein the cells are characterized by a vitamin receptor on

the surface of the cells. The method comprises the steps of administering to the animal an effective amount of a composition comprising a compound of the formula



5 wherein V is the vitamin, or a receptor binding derivative or analog thereof, specific for the cell surface vitamin receptor, L is a divalent linker, R is a side chain of an amino acid of the formula $\text{H}_2\text{N-CHRCOOH}$, M is a cation of a radionuclide, n is 1 or 0, and a pharmaceutically acceptable carrier therefor, and monitoring the biodistribution of the compound in the animal.

10 In any of these embodiments, V in the compound can be, for example, a vitamin selected from the group consisting of folate, riboflavin, thiamine, vitamin B₁₂, and biotin, or a derivative or analog thereof. In any of these embodiments, the radionuclide in the compound can be selected, for example, from the group consisting of radioisotopes of gallium, indium, copper, technetium, and rhodium.

15

BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1. Structure of EC20, an exemplary compound used as an imaging agent in accordance with the invention.

Fig. 2. HPLC radiochromatogram of ^{99m}Tc-EC20. Samples of ^{99m}Tc-EC20 were eluted isocratically on a Waters Nova-Pak C18 (3.9 x 150 mm) column using an aqueous mobile phase containing 20% methanol and 0.2% trifluoroacetic acid at a flow rate of 1 mL/min. The HPLC analysis was monitored with both the UV detector (280 nm) and a Bioscan FC-3200 radiodetector. Peak A, free ^{99m}Tc; Peak B, a folate-containing chelate of unknown structure; Peaks C and D, diastereomers possessing either a *syn* or *anti* configuration of the technetium-oxygen bond in the Dap-Asp-Cys chelating ring of EC20.

Fig. 3. Structures of Re-EC20 and ^{99m}Tc-EC20 isomers (*syn* or *anti* position of metal-oxo bond).

Fig. 4. Blocking of ³H-folic acid binding to KB cells with various folate-containing competitors. KB cells were incubated for 15 min on ice with 100

-8-

nM 3 H-folic acid in the presence and absence of increasing competitor concentrations. (•) Folic acid; (■) EC20; (▲) EC20:Re isomer A; (▼) EC20:Re isomer B; (□) DTPA-Folate. Error bars represent 1 standard deviation (n = 3).

Fig. 5. Time-dependent association of 99m Tc-EC20. KB cells were incubated with 10 nM 99m Tc-EC20 for increasing periods of time at 37°C. Following multiple washes, cells were harvested and counted for associated radioactivity. Error bars represent 1 standard deviation (n = 3).

Fig. 6. Concentration-dependent association of 99m Tc-EC20. KB cells were incubated for 2 hr at 37°C in the presence of increasing concentrations of 99m Tc-EC20. Following multiple washes, cells were harvested and counted for associated radioactivity. Error bars represent 1 standard deviation (n = 3).

Fig. 7. Concentration-dependent association of 99m Tc-EC20 "peak B." KB cells were incubated for 2 hr at 37°C in the presence of increasing concentrations of "Peak B" that was chromatographically isolated from the 99m Tc-EC20 formulation. Following multiple washes, cells were harvested and counted for associated radioactivity. Error bars represent 1 standard deviation (n = 3). (•), Peak B; (○), Peak B plus 1 mM folic acid.

Fig. 8. Blood clearance of 99m Tc-EC20 in Balb/c mice. Each animal received an intravenous dose of 50 μ g/kg EC20 (67 nmol/kg) in approximately 0.1 mL during brief diethyl ether anesthesia. At the designated times post-injection, each animal was euthanized by CO₂ asphyxiation, blood was collected and counted for associated radioactivity. Error bars represent 1 standard deviation (n = 3 animals).

Fig. 9. Whole-body gamma images (ventral view). Images were obtained 4 hr following intravenous administration of 99m Tc-EC20 to a Balb/c mouse bearing a subcutaneous folate receptor-positive M109 tumor. Only the kidneys (K) and tumor (T) exhibit significant accumulation of this radiotracer.

Fig. 10. Structures of EC11, EC13, EC14, EC15, EC19, EC20, EC31, and EC53.

Fig. 11. Tissue distribution of 99m Tc-EC20 in Balb/c mice bearing FR-positive M109 tumors and FR-negative 4T1 tumors.

Fig. 12. HPLC analysis of EC11.

Fig. 13. Mass spectroscopy analysis of EC11.

-9-

Fig. 14. NMR analysis of EC11.

Fig. 15. HPLC analysis of EC13.

Fig. 16. NMR analysis of EC14.

Fig. 17. Mass spectroscopy analysis of EC15.

Fig. 18. HPLC analysis of EC19.

Fig. 18. Mass spectroscopy analysis of EC19.

Fig. 3Q HPLC analysis of EC31.

Fig. 21 HPLC analysis of EC53.


Fig. 32. Mass spectroscopy analysis of EC53.

Fig. 22. Mass spectroscopy analysis of EC53.


DETAILED DESCRIPTION OF THE INVENTION

In accordance with the invention, compounds and methods are provided for targeting radionuclide-based imaging agents to cell populations that uniquely express, overexpress, or preferentially express vitamin receptors. Accordingly, a vitamin, or a receptor binding derivative or analog thereof, is used as the targeting ligand for the imaging agent. The vitamin-imaging agent conjugate can be used to target radionuclides to cells and to concentrate the radionuclides in a cell population, such as a tumor cell population, for use in diagnostic imaging.

20 The invention provides a composition for diagnostic imaging comprising a compound of the formula




81



-10-


for use in such methods. In the compound, V is a vitamin, or a vitamin receptor binding derivative or analog thereof, L is a divalent linker, R is a side chain of an amino acid of the formula $H_2N-CH_2-RCOOH$, M is a cation of a radionuclide, and n is 1 or 0. The vitamin, or vitamin receptor binding derivative or analog thereof, is a substrate for receptor-mediated transmembrane transport in vivo.

5 The invention also provides compounds of the formulas



and

10



wherein V is a vitamin, or a vitamin receptor binding derivative or analog thereof, L is a divalent linker, R is a side chain of an amino acid of the formula $H_2N-CH_2-RCOOH$, M is a cation of a radionuclide, n is 1 or 0, and k is 1 or 0. The vitamin is a substrate for receptor-mediated transmembrane transport in vivo.

15 Exemplary of these compounds is a compound referred to as EC20 depicted in Fig. 1. Exemplary of other compounds for use in accordance with this invention are compounds denominated as EC11, EC13, EC14, EC15, EC19, EC31, and EC53 (see Fig. 10). The vitamin moiety (e.g., the folic acid moiety in EC20) provides high affinity binding to cellular receptors. The compounds also contain a bifunctional peptide-based chelator, which provides the site for chelation of the radionuclide, for example, ^{99m}Tc (see Fig. 1), and the compounds can, optionally, contain a linker through which the vitamin moiety is covalently bonded to the chelating moiety.

20 In accordance with the invention, the vitamin moiety of the compounds is a vitamin that is a substrate for receptor-mediated transmembrane transport in vivo, or a vitamin receptor binding derivative or analog thereof. The vitamin is linked,

- 11 -

optionally, through a linker (L) to the chelator portion of the compounds. The chelator portion comprises an α , β -diaminopropionic acid moiety linked to a cysteine group through a third amino acid residue. The chelator portion of the compound is adapted to bind a radionuclide cation (M) (where $k = 1$).

5 In accordance with the invention, the compounds with bound radionuclides are referred to as "vitamin-imaging agent conjugates."

The structure of the linker, if present, is not critical to the invention. Thus, for example, it can be any biocompatible divalent linker. Typically, the linker comprises about 1 to about 30 carbon atoms, more typically about 2 to about 20 10 carbon atoms. Lower molecular weight linkers (i.e., those having an approximate molecular weight of about 30 to about 300) are typically employed. Furthermore, the vitamin moiety may be a vitamin, or a derivative or analog thereof. For example, folate contains one glutamic acid in the L configuration linked to pteroic acid. As shown in Fig. 1, EC20 comprises a folic acid analog linked to the chelator moiety because EC20 has the glutamic acid in the D configuration. EC11 and EC14 contain 15 two glutamic acid residues and, thus, these compounds can also, for example, be considered derivatives of folic acid (Fig. 10).

Among vitamins believed to trigger receptor-mediated endocytosis, and having application in accordance with the presently disclosed method are niacin, 20 pantothenic acid, folic acid, riboflavin, thiamine, biotin, vitamin B₁₂, and the lipid soluble vitamins A, D, E and K. These vitamins, and their analogs and derivatives, constitute vitamins that can be coupled with imaging agents to form the vitamin-chelator conjugates for use in accordance with the invention. Preferred vitamin 25 moieties include folic acid, biotin, riboflavin, thiamine, vitamin B₁₂, and analogs and derivatives of these vitamin molecules, and other related vitamin receptor-binding molecules.

Folic acid, folinic acid, pteroic acid, pteropolyglutamic acid, and folate receptor-binding pteridines such as tetrahydropterins, dihydrofolates, 30 tetrahydrofolates, and their deaza and di-deaza analogs can be used in accordance with the invention. The terms "deaza" and "di-deaza" analogs refers to the art-recognized folate analogs having a carbon atom substituted for one or two nitrogen atoms in the naturally occurring folic acid structure. For example, the deaza analogs include the 1-

-12-

deaza, 3-deaza, 5-deaza, 8-deaza, and 10-deaza analogs. The dideaza analogs include, for example, 1,5-dideaza, 5,10-dideaza, 8,10-dideaza, and 5,8-dideaza analogs. The foregoing are folate analogs or derivatives and can bind to folate receptors. Other folate analogs or derivatives useful in accordance with the invention are the folate receptor-binding analogs aminopterin, amethopterin (methotrexate), N^{10} -methylfolate, 2-deaminohydroxyfolate, deaza analogs such as 1-deazamethopterin or 3-deazamethopterin, and 3'5'-dichloro-4-amino-4-deoxy- N^{10} -methylpteroylglutamic acid (dichloromethotrexate).

The vitamin, or derivative or analog thereof, can be capable of selectively binding to the population of cells to be visualized due to preferential expression on the targeted cells of a receptor for the vitamin, or derivative or analog, wherein the receptor is accessible for binding. The binding site for the vitamin can include receptors for any vitamin molecule capable of specifically binding to a receptor wherein the receptor or other protein is uniquely expressed, overexpressed, or preferentially expressed by the population of cells to be visualized. A surface-presented protein uniquely expressed, overexpressed, or preferentially expressed by the cells to be visualized is a receptor not present or present at lower amounts on other cells providing a means for selective, rapid, and sensitive visualization of the cells targeted for diagnostic imaging using the vitamin-imaging agent conjugates of the present invention.

In accordance with the invention the vitamin-imaging agent conjugates are capable of high affinity binding to receptors on cancer cells or other cells to be visualized. The high affinity binding can be inherent to the vitamin moiety or the binding affinity can be enhanced by the use of a chemically modified vitamin (i.e., an analog or a derivative) or by the particular chemical linkage between the vitamin and the chelator moiety that is present in the conjugate.

In accordance with the invention, the chelator can be conjugated with multiple, different vitamins, or vitamin receptor binding derivatives or analogs, to enhance the opportunity for binding to the respective cell membrane receptors. Alternatively, independent portions of the dose of a vitamin-imaging agent conjugate can constitute different vitamin-imaging agent conjugates to enhance the opportunity for binding to the respective cell membrane receptors.

-13-

Generally, any manner of forming a complex between the chelator and the vitamin, or vitamin receptor binding derivative or analog, can be utilized in accordance with the present invention. The chelator can form a complex with the vitamin, or vitamin receptor binding derivative or analog, by direct conjugation of the chelator and the vitamin by using a divalent linker. Alternatively, the vitamin and the chelator may be conjugated without employing a linker. If a linker is used, the linker can directly conjugate the vitamin, or vitamin receptor binding derivative or analog, and the chelator through a hydrogen, ionic, or covalent bond. Also, in accordance with this invention the divalent linker can comprise an indirect means for associating the chelator with the vitamin, or vitamin receptor binding derivative or analog, such as by connection through intermediary linkers, spacer arms, or bridging molecules. Both direct and indirect means for association must not prevent the binding of the vitamin, or vitamin receptor binding derivative or analog, to the vitamin receptor on the cell membrane for operation of the method of the present invention.

Covalent bonding of the vitamin, or vitamin receptor binding derivative or analog, and the chelator can occur, whether or not a linker is employed, through the formation of amide, ester or imine bonds between acid, aldehyde, hydroxy, amino, or hydrazo groups. For example, a carboxylic acid on the vitamin moiety or on the chelator can be activated using carbonyldiimidazole or standard carbodiimide coupling reagents such as 1-ethyl-3-(3-dimethylaminopropyl)-carbodiimide (EDC) and thereafter reacted with the other component of the conjugate, or with a linker, having at least one nucleophilic group, viz hydroxy, amino, hydrazo, or thiol, to form the vitamin-chelator conjugate coupled, with or without a linker, through ester, amide, or thioester bonds.

The radionuclides suitable for diagnostic imaging include radioisotopes of gallium, indium, copper, technetium and rhenium, including isotopes ¹¹¹In, ^{99m}Tc, ⁶⁴Cu, ⁶⁷Ga or ⁶⁸Ga. These radionuclides are cationic and are complexed with the chelator through the chelating group of the conjugate to form the vitamin-imaging agent conjugate.

The vitamin-imaging agent conjugates in accordance with the invention are utilized to selectively visualize, using scintigraphic imaging techniques, a population of cells in an animal wherein the population of cells uniquely expresses,

- 14 -

overexpresses, or preferentially expresses receptors for a vitamin, or a vitamin receptor binding derivative or analog thereof. The vitamin-imaging agent conjugates can be used to visualize populations of pathogenic cells, as long as the cells uniquely or preferentially express or overexpress vitamin receptors or receptors that bind

5 vitamin derivatives or analogs.

The invention is applicable to populations of pathogenic cells that cause a variety of pathologies including cancer, and diseases mediated by any other type of pathogenic cells that overexpress vitamin receptors, or receptors capable of binding vitamin derivatives or analogs. Thus, the population of pathogenic cells can be tumorigenic, including benign tumors and malignant tumors, or it can be non-tumorigenic. If the cell population is a cancer cell population, the cancer cells can arise spontaneously or by such processes as mutations present in the germline of the host animal or somatic mutations, or the cancer can be chemically-, virally-, or radiation-induced. The invention can be utilized for diagnostic imaging of such cancers as carcinomas, sarcomas, lymphomas, Hodgkin's disease, melanomas, mesotheliomas, Burkitt's lymphoma, nasopharyngeal carcinomas, and myelomas. The cancer cell population can include, but is not limited to, oral, nasopharyngeal, thyroid, endocrine, skin, gastric, esophageal, laryngeal, throat, pancreatic, colon, bladder, bone, ovarian, cervical, uterine, breast, testicular, prostate, rectal, kidney, liver, lung, and brain cancers. In embodiments where the cell population is a cancer cell population, tumor cells, including cells of the primary tumor or cells that have metastasized or are in the process of dissociating from the primary tumor, can be visualized using the vitamin-imaging agent conjugate.

The vitamin-imaging agent conjugates of the present invention can be used to diagnose a disease state or to monitor the progression of disease. For example, the diagnostic imaging method in accordance with the invention can be used to monitor the progression of cancer in combination with prophylactic treatments to prevent return of a tumor after its removal by any therapeutic approach including surgical removal of the tumor, radiation therapy, chemotherapy, or biological therapy.

30 The compositions and methods of the present invention can be used for both human clinical medicine and veterinary applications. Thus, the animal harboring the population of cells that are visualized can be human or, in the case of veterinary

-15-

applications, can be a laboratory, agricultural, domestic, or wild animal. The present invention can be applied to animals including, but not limited to, humans, laboratory animals such as rodents (e.g., mice, rats, hamsters, etc.), rabbits, monkeys, chimpanzees, domestic animals such as dogs, cats, and rabbits, agricultural animals such as cows, 5 horses, pigs, sheep, goats, and wild animals in captivity such as bears, pandas, lions, tigers, leopards, elephants, zebras, giraffes, gorillas, dolphins, and whales.

The compositions for diagnostic imaging comprise an amount of the vitamin-imaging agent conjugate effective to visualize the cells targeted for diagnostic imaging in an animal when administered in one or more doses. The diagnostic imaging composition containing the vitamin-imaging agent conjugate is preferably administered to the animal parenterally, e.g., intradermally, subcutaneously, 10 intramuscularly, intraperitoneally, intravenously, or intrathecally. Alternatively, the composition containing the vitamin-imaging agent conjugate can be administered to the animal by other medically useful processes, and any effective dose and suitable dosage form can be used, including oral and inhalation dosage forms.

Examples of parenteral dosage forms include aqueous solutions of the vitamin-imaging agent conjugate, in isotonic saline, 5% glucose or other well-known pharmaceutically acceptable liquid carriers such as liquid alcohols, glycols, esters, and amides. The parenteral dosage form in accordance with this invention can be in 20 the form of a reconstitutable lyophilizate comprising the dose of the vitamin-imaging agent conjugate.

The dosage of the vitamin-imaging agent conjugate in the diagnostic imaging composition can vary significantly depending on the size of the animal, the cell population targeted for diagnostic imaging, the specific vitamin-imaging agent conjugate being used, and the route of administration of the conjugate. The effective amount to be administered to the animal is based on body surface area, weight, and physician assessment of the condition of the animal. An effective dose can range from about 1 ng/kg to about 1 mg/kg, more preferably from about 100 ng/kg to about 500 μ g/kg, and most preferably from about 100 ng/kg to about 25 μ g/kg.

Any effective regimen for administering the diagnostic imaging composition containing the vitamin-imaging agent conjugate can be used. For example, the diagnostic imaging composition can be administered as a single dose, or


-16-

it can be administered in multiple doses, if necessary, to achieve visualization of the targeted cell population. Additional injections of the diagnostic imaging composition containing the vitamin-imaging agent conjugate can be administered to the animal at an interval of days or months after the initial injections(s), and the additional 5 injections can be useful for monitoring the progress of the disease state. The diagnostic imaging composition containing the vitamin-imaging agent conjugate can also be administered in combination with unlabeled vitamin. "In combination with" means that the unlabeled vitamin can be either co-administered with the imaging agent or the unlabeled vitamin can be preinjected before administration of the imaging agent to improve image quality. For example, the imaging agent can be administered in combination with about 0.5 ng/kg to about 100 mg/kg, or about 1 μ g/kg to about 100 mg/kg, or about 100 μ g/kg to about 100 mg/kg of the unlabeled vitamin.

10 The diagnostic imaging composition is typically formulated for parenteral administration and is administered to the animal in an amount effective to enable imaging of the targeted cell population. Typically, the diagnostic imaging composition containing the vitamin-targeted imaging agent is administered to the animal, and following a period of time to allow delivery and concentration of the vitamin-imaging agent conjugate in the targeted cell population, the animal is subjected to the imaging procedure and imaging is enabled by the vitamin-imaging agent conjugate. When used for monitoring the progression of disease or diagnosis, 15 imaging procedures are typically carried out about 1 to about 6 hours post administration of the diagnostic imaging composition containing the vitamin-imaging agent conjugate.

20 The invention also provides a method of imaging a population of cells in an animal wherein the cells are characterized by a vitamin receptor on the surface of the cells. The method comprises the steps of administering to the animal an effective amount of a composition comprising a compound of the formula

-17-



EXAMPLE 1

Materials

N^{10} -trifluoroacetylpteroic acid was purchased from Eprova AG, 25 Schaffhausen, Switzerland. Peptide synthesis reagents were purchased from NovaBiochem and Bachem. ^{99m}Tc Sodium Pertechnetate was supplied by Syncor. $[ReO_2(en)_2]Cl$ was prepared according to Rouschias (Rouschias, G., Chem. Rev., 74:

-18-

531 (1974)). Cellulose plates and DEAE ion exchange plates were purchased from J.T. Baker.

EXAMPLE 2

Synthesis, Purification, and Analytical Characterization of EC20

5 EC20 was prepared by a polymer-supported sequential approach using the Fmoc-strategy (see Scheme 1 below; Fmoc = 9-fluorenylmethoxycarbonyl; Boc = tert-butyloxycarbonyl; Dap = diaminopropionic acid; DMF = dimethylformamide; DIP-EA = diisopropylethylamine). EC20 was synthesized on an acid-sensitive Wang resin loaded with Fmoc-L-Cys(Trt)-OH. Benzotriazole-1-yl-oxy-tris-pyrrolidinomophosphoniumhexafluorophosphate (PyBOP) was applied as the activating reagent to ensure efficient coupling using low equivalents of amino acids. Fmoc protecting groups were removed after every coupling step under standard conditions (20% piperidine in DMF). After the last assembly step the peptide was cleaved from the polymeric support by treatment with 92.5% trifluoroacetic acid containing 2.5% ethanedithiol, 2.5% triisopropylsilane and 2.5% deionized water. This reaction also resulted in simultaneous removal of the t-Bu, Boc and trityl protecting groups. Finally, the trifluoroacetyl moiety was removed in aqueous ammonium hydroxide to give EC20.


20 The crude EC20 product was purified by HPLC using an Xterra RP18 30 x 300 mm, 7 μ m column (Waters); mobile phase 32 mM HCl (A), MeOH (B); gradient conditions starting with 99% A and 1% B, and reaching 89% A and 11% B in 37 min by a flow rate of 20 mL/min. Under these conditions, EC20 monomer typically eluted at 14.38 min, whereas EC20 disulfide dimer (minor contaminant) eluted at 16.83 min. All other compounds shown in Fig. 10 can be prepared using a similar synthesis scheme except for EC15 which is synthesized as shown in Scheme 2 below.

25 Two milligrams of HPLC-purified EC20 were dissolved in 0.62 mL of D₂O, and a 500 MHz ¹H-NMR spectrum was collected. Table 1 (see below) lists the chemical shifts, signal shapes, and J values for all non-exchangeable protons in EC20 30 molecule.

-19-

■EC20 was also analyzed by elec-trospray-mass spectrometry. Major positive ion peaks (m/z, relative intensity): 745.1, 100; 747.1, 44; 556.8, 32; 570.8, 16.


-20-

Scheme 1^a

^aReagents and conditions: i) 20% Piperidine, DMF; ii) Fmoc-Asp(OtBu)-OH, PyBop, DIPEA, DMF; iii) Boc-Dap(Fmoc)-OH, PyBop, DIPEA, DMF; iv) Fmoc-D-Glu-OtBu, PyBop, DIPEA, DMF; v) N¹⁰-TFA-Pte-OH, DIPEA, DMSO; vi) F₃CCO₂H, HSCH₂CH₂SH, iPr₂SiH; vii) H₄NOH, pH = 10.3.

Table 1. ¹H-NMR data for EC20. EC20 was dissolved in D₂O and a 500 MHz spectrum was collected. Chemical shifts (δ) are in ppm. The signal for HOD at δ = 4.80 ppm was used as the reference. pD = 4.78; s = singlet; d = doublet; m = multiplet.

Residue	Protons observed	Chemical Shift (δ)	Signals	J values
Pte	H-7	8.76	s	
	2 x H-9	4.64	s	
	H-12 a. H-16	7.68	d	³ J _{12,13} = ³ J _{15,16} =8.8 Hz
	H-13 a. H-15	6.8	d	
D-Glu	H-2	4.41	dd	³ J _{2,3} =9.1 Hz; ³ J _{2,3b} =4.5 Hz
	H-3a	2.08	m	² J _{3a,3b} =14.2 Hz
	H-3B	2.27	m	
	2 x H-4	2.44	dd	³ J _{3a,4} = ³ J _{4b,4} =5.6 Hz
Dpr	H-2	4.1	dd; X of ABX System	³ J _{2,3A} =6.6 Hz; ³ J _{2,3B} =4.7 Hz
	H-3A	3.52	dd; A of ABX System	² J _{A,B} =14.7 Hz
	H-3B	3.72	dd; B of ABX System	
Asp	H-2	4.71	dd; X of ABX System	
	H-3A	2.62	dd; A of ABX System	³ J _{2,3A} =9.5 Hz; ³ J _{2,3B} =4.3 Hz
	H-3B	2.81	dd; B of ABX System	² J _{A,B} =16.1 Hz
Cys	H-2	4.3	dd; X of ABX System	
	H-3A	2.85	dd; A of ABX System	³ J _{2,3A} =5.5 Hz; ³ J _{2,3B} =4.4 Hz
	H-3B	2.89	dd; B of ABX System	² J _{A,B} =14.1 Hz

Scheme 2^a

^aReagents and conditions: i) 20% Piperidine, DMF; ii) Fmoc-Asp(OtBu)-OH, PyBop, DIPEA, DMF; iii) Fmoc-Cys(Trt)-OH, PyBop, DIPEA, DMF; iv) Fmoc-D-Glu(OtBu), PyBop, DIPEA, DMF; v) N¹⁰-TFA-Pte-OH, DIPEA, DMSO; vi) TFAA, HS-CH₂CH₂SH, iPr₂SiH; vii) H₄NOH, pH = 10.3

5

EXAMPLE 3

Preparation of the Non-Radioactive Reagent vial and of ^{99m}Tc-EC20

EC20 kits were used for preparation of the ^{99m}Tc-EC20 radioactive drug substance. Each kit contained a sterile, non-pyrogenic lyophilized mixture of 0.1 mg EC20, 80 mg sodium α -D-glucosidate, 80 mg tin (II) chloride dihydrate, and sufficient sodium hydroxide or hydrochloric acid to adjust the pH to 6–8 ± 0.2 prior to lyophilization. The lyophilized powder was sealed in a 5 mL vial under an argon atmosphere. The kits were then stored frozen at -20°C until use or expiration (current shelf life is > 2 years). Importantly, the tin (II) chloride component is required to reduce the added ^{99m}Tc-pertechnetate, while the sodium α -D-glucosidate component is necessary to stabilize the newly reduced ^{99m}Tc prior to its final chelation to the EC20 compound.

^{99m}Tc-EC20 was prepared as follows (i.e., chelation of ^{99m}Tc to EC20). First, a boiling water bath containing a partially submerged lead vial shield was prepared. The top of an EC20 vial was swabbed with 70% ethanol to sanitize the surface and the vial was placed in a suitable shielding container. Using a shielded syringe with 27-gauge needle, 1 mL of sterile Sodium Pertechnetate ^{99m}Tc Injection

-22-

(15 to 20 mCi) in 0.9% sodium chloride was injected into the shielded vial. Before removal of the syringe from the vial, a volume of gas from the vial equal to the volume of pertechnetate added was withdrawn in order to normalize the pressure inside the vial. The vial was gently swirled for 30 seconds to ensure complete dissolution of the lyophilized powder. The vial was then placed into the lead shield that was standing in the boiling water bath. The solution was heated for ~18 minutes and then cooled to room temperature for a minimum of 15 min. This solution can be stored at room temperature (15-25°C) protected from light, but it should be used within 6 hours of preparation.

10 The radiochemical stability of the radioactive drug substance was determined by HPLC after storing at room temperature protected from light for up to 24 hours. Samples of the ^{99m}Tc-EC20 solution (20 µL) were analyzed using an HPLC system consisting of a Waters 600E Multisolvent Delivery System and 490 UV detector, a Bioscan EC-3200 radiodetector, Lambda v1.5 radiochromatogram software, and a Waters Nova-Pak C18 (3.9 x 150 mm) column. Injected samples were eluted isocratically using an aqueous mobile phase containing 20% methanol and 0.1% trifluoroacetic acid at a flow rate of 1 mL/min. The HPLC analysis was monitored with both the UV detector (280 nm) and the gamma radiodetector. Notably, the radiochemical purity of ^{99m}Tc-EC20 remained greater than 90% for at least 24 hours in all cases.

EXAMPLE 4

Determination of Radiochemical purity of ^{99m}Tc-EC20 by TLC

The major radiochemical impurities in the preparation of ^{99m}Tc-EC20 will be 1) ^{99m}Tc pertechnetate, 2) ^{99m}Tc-glucoheptonate (ligand exchange precursor), 3) non-specific binding ^{99m}Tc (^{99m}Tc bound at a site other than the expected Dap-Asp-Cys chelating moiety of the EC20 molecule), and 4) hydrolyzed ^{99m}Tc. Since ^{99m}Tc-EC20 was being tested for possible clinical use, a three-TLC-based method was developed to determine the amounts of each impurity and to estimate the overall radiochemical purity of ^{99m}Tc-EC20.

In the first system a cellulose plate was developed with deionized water. ^{99m}Tc-EC20, ^{99m}Tc-glucoheptonate, non-specific binding ^{99m}Tc and ^{99m}Tc

-23-

Pertechnetate move to the solvent front ($R_f = 1.0$), while hydrolyzed ^{99m}Tc remains at the origin ($R_f = 0.0$). The cellulose plate was cut into two pieces at $R_f = 0.3$ (1.5 cm from origin) and each piece was counted using a dose calibrator. The percent of hydrolyzed ^{99m}Tc was calculated as follows: $A = \% \text{ Hydrolyzed} = \frac{(\mu\text{Ci in bottom piece}/\mu\text{Ci in both pieces})}{2} \times 100$.

5 In the second system, a cellulose plate was developed with acetone and 0.9% NaCl (7:3, v/v). ^{99m}Tc -pertechnetate moves with $R_f = 0.9$, while $^{99m}\text{Tc-EC20}$, $^{99m}\text{Tc-glucoheptonate}$, non-specific binding ^{99m}Tc and hydrolyzed ^{99m}Tc remain at the origin ($R_f = 0.0$). The cellulose/acetone-saline plate was cut into two pieces at $R_f = 0.6$ (3.0 cm from the origin) and each piece was counted using a dose calibrator. The percent of ^{99m}Tc -pertechnetate was calculated as follows: $B = \% \text{ }^{99m}\text{Tc-pertechnetate} = \frac{(\mu\text{Ci in top piece}/\mu\text{Ci in both pieces})}{2} \times 100$.

10 Finally, in the third system a DEAE ion exchange plate was developed with 0.3 M Na_2SO_4 . $^{99m}\text{Tc-glucoheptonate}$ moves to the solvent front ($R_f = 1.0$), nonspecific binding ^{99m}Tc moves with $R_f = 0.6$, and $^{99m}\text{Tc-EC20}$, hydrolyzed ^{99m}Tc and ^{99m}Tc -pertechnetate remain near the origin ($^{99m}\text{Tc-EC20}$, $R_f = 0.1$; hydrolyzed ^{99m}Tc : $R_f = 0.0$; ^{99m}Tc pertechnetate: $R_f = 0.3$). The cellulose/ Na_2SO_4 plate was cut into two pieces at 2.5 cm from the origin and each piece was counted using a dose calibrator. The percent of $^{99m}\text{Tc-glucoheptonate}$ and non-specific binding ^{99m}Tc were calculated as follows: $C = \% \text{ }^{99m}\text{Tc-Glucoheptonate} + \% \text{ non-specific binding }^{99m}\text{Tc} = (\mu\text{Ci in top piece}/\mu\text{Ci in both pieces}) \times 100$. The overall radiochemical purity of $^{99m}\text{Tc-EC20}$ was then calculated as follows: Radiochemical purity = $100 - (A + B + C)$.

15 As shown in Fig. 2, HPLC analysis of the $^{99m}\text{Tc-EC20}$ formulation shows four radiochemical components, designated as Peaks A through D. Peak A was confirmed to be free ^{99m}Tc and this by-product is reproducibly present at < 2%. Peak B, which was different from that of $^{99m}\text{Tc-glucoheptonate}$ (data not shown) eluted with a retention time of 2.8 min. This component represented about 3% of the mixture and was thought to result from ^{99m}Tc chelating at some other site on the EC20 molecule besides the expected Dap-Asp-Cys moiety. Peaks C and D (retention times of 4.8 minutes and 13.2 minutes, respectively), account for the majority of the formulated radiochemical activity.

-24-

EXAMPLE 5

Synthesis of Re-EC20

Fifty-two mg (0.010 mmol) of EC20 and $[\text{ReO}_2(\text{en})_2]\text{Cl}$ (52 mg, 0.014 mmol) were dissolved in 6 mL and 1 mL argon-purged phosphate buffer (0.05 M, pH 5.8), respectively. The two solutions were combined and heated under an argon atmosphere in a boiling water bath for 2 hours. The reaction mixture was frozen and lyophilized overnight. The crude product was purified by HPLC (Xterra RPI8 column, 1.9 x 150 mm, 10 mM $\text{NH}_4\text{OAc}/\text{CH}_3\text{CN}$, flow rate 10 mL/mm; gradient 1% to 8%). The fractions were collected, lyophilized and stored at -20°C until use.

Because no mass spectral facilities were available for analysis of radioactive materials, the non-radioactive rhenium analog, Re-EC20, was analyzed. Both rhenium and technetium are Group VIIA metals that have significant similarity in physical and chemical properties. They also form similar complexes with organic ligands. This analogous chemical behavior has been frequently used in structure elucidation of new classes of technetium radiopharmaceuticals based on non-radioactive rhenium analogues. Interestingly, HPLC analysis of Re-EC20 also showed two major peaks eluting at 5 and 14.2 minutes, respectively, similar to Peaks C and D for ^{99m}Tc -EC20 (chromatogram not shown). Mass spectral analysis confirmed that these two components were isomers corresponding to the Re-EC20 complex ($m/z = 945$). In fact, these species were likely diastereomers possessing either a *syn* or *anti* configuration of the technetium-oxygen bond in the Dap-Asp-Cys chelating ring, as depicted in Fig. 3. Because i) the two peaks in the Re-EC20 chromatogram represent isomeric complexes, and ii) reports of similar isomerism in technetium complexes exist, it is likely that components C and D in the ^{99m}Tc -EC20 radiochromatogram are also isomers.

EXAMPLE 6

Cell Culture

Cells were grown continuously as a monolayer using folate-free RPMI medium (FFRPMI) containing 10% heat-inactivated fetal calf serum (HIFCS) at 37°C in a 5% CO_2 /95% air-humidified atmosphere with no antibiotics. The HIFCS contained its normal complement of endogenous folates which enabled the cells to

-25-

sustain growth in this more physiologically-relevant medium. All cell experiments were performed using FFRPMI containing 10% HIFCS (FFRPMI/HIFCS) as the growth medium, except where indicated.

5

EXAMPLE 7

Relative Affinity Assay

The relative affinity of various folate derivatives was determined according to the method described by Westerhoff et al. (Mol. Pharm., 48: 459-471 (1995)) with slight modification. Briefly, FR-positive KB cells were gently trypsinized in 0.25% trypsin/PBS at room temperature for 3 minutes and then diluted in FFRPMI/HIFCS. Following a 5 min 800 x g spin and one PBS wash, the final cell pellet was suspended in FFRPMI 16-40 (no serum). Cells were incubated for 15 min on ice with 100 nM of ³H-folic acid in the absence and presence of increasing concentrations of folate-containing test articles. Samples were centrifuged at 10,000 x g for 5 min, cell pellets were suspended in buffer, were transferred to individual vials containing 5 mL of scintillation cocktail, and were then counted for radioactivity. Negative control tubes contained only the ³H-folic acid in FFRPMI (no competitor). Positive control tubes contained a final concentration of 1 mM folic acid, and CPMs measured in these samples (representing non-specific binding of label) were subtracted from all samples. Notably, relative affinities were defined as the inverse molar ratio of compound required to displace 50% of ³H-folic acid bound to KB-FR, and the relative affinity of folic acid for the FR was set to 1.

The capacity of EC20 to directly compete with folic acid for binding to cell surface FRs was measured using this assay. Importantly, a relative affinity value of 1.0 implies that the test article ligand has an affinity for the FR equal to folic acid. Likewise, values lower than unity reflect weaker affinity, and values higher than unity reflect stronger affinity.

Cultured KB cells were incubated with 100 nM ³H-folic acid in the presence of increasing concentrations of non-radioactive folic acid, EC20, Rhenium-EC20 (isomer A; Peak C), Rhenium-EC20 (isomer B; peak 0), or a related folate-based radiopharmaceutical, DTPA-folate. Following a 15-minute incubation at 4°C, cells were rinsed free of unbound material and counted for residual cell-associated

-26-

radioactivity. The quantity of bound radioactivity was plotted against the concentration of unlabeled ligand, and IC₅₀ values (concentration of ligand required to block 50% of ³H-folic acid binding) were estimated. As shown in Fig. 4 and Table 2 (below), EC20 was determined to have an affinity of 0.92 relative to that of folic acid. 5 for human FRs. Both isomers of Rhenium-EC20 displayed relative affinity values that were very similar to, if not better than, the parent EC20 molecule (1.42 and 1.37 for Re-EC20 isomer A and isomer B, respectively). DTPA-folate, an ¹¹¹In-chelating folate radiopharmaceutical agent, displayed a relative affinity of 0.87 for the folate receptor. Thus, chemical modification of folate with various metal chelating motifs 10 did not disturb the vitamin's intrinsic affinity for the FR.

-27-

Table 2. Relative Affinity Estimations. Relative affinities (RA) were defined as the inverse molar ratio of compound required to displace 50% of ^3H -folic acid bound to FR-positive KB cells. The relative affinity of folic acid was set to 1. Each test article was evaluated in triplicate.

5

Test Article	IC ₅₀ (nM)	S.D.	RA	S.D.
Folic Acid	118	± 19	1.00	
EC20	128	± 25	0.92	± 0.23
EC20:Re isomer 1	83	± 16	1.42	± 0.36
EC20:Re isomer 2	86	± 3	1.37	± 0.23
DTPA-Folate	136	± 12	0.87	± 0.16

EXAMPLE 8

Time-Dependent Cell Uptake

KB cells were seeded in 12-well Falcon plates and allowed to form sub-confluent monolayers overnight. Following one rinse with 1 mL of fresh FFRPMI/HIFCS, each well received 1 mL of FFRPMI/HIFCS containing 10 nM $^{99\text{m}}\text{Tc}$ -EC20. Cells were incubated for predetermined times at 37°C and then rinsed four times with 1 mL of ice-cold PBS, pH 7.4. The cell monolayers were dissolved in 0.5 mL of PBS, pH 7.4 containing 1% sodium dodecyl sulfate for 15 min at room temperature and then counted for radioactivity using a Packard gamma counter. The protein in each sample was quantitated using a BioRad DC Protein Assay kit, and cellular protein values were converted to cell number using the conversion factor of 2.23×10^{-7} mg protein per cell. Final tabulated values were expressed in terms of molecules of EC20 per cell.

The kinetics of $^{99\text{m}}\text{Tc}$ -EC20 uptake into FR-positive KB cells was quantitatively measured using this protocol. As shown in Fig. 5, steady-state uptake was reached within two hours at 37°C, where approximately 3.2 million molecules of EC20 were cell-associated, whereas half-maximal cell association occurred 9 minutes after mixing 1.0 nM of this radiopharmaceutical with the cells. Interestingly, the half-maximal saturation point was reached in only 37 seconds when cells were incubated

-28-

with a 10-fold higher concentration of 99m Tc-EC20 (100 nM; data not shown).

EXAMPLE 9

Concentration-Dependent Cell Uptake

KB cells were seeded in 12-well Falcon plates and allowed to form sub-confluent monolayers overnight. Following one rinse with 1 mL of fresh FERPMI/HIFCS, each well received 1 mL of FERPMI/HIFCS containing increasing concentrations of 99m Tc-EC20. Cells were incubated for 2 hours at 37°C and then rinsed four times with 1 mL of ice-cold PBS, pH 7.4. The monolayers were dissolved in 0.5 mL of PBS, pH 7.4 containing 1% sodium dodecyl sulfate for 15 min at room temperature and then counted for radioactivity using a Packard gamma counter. Protein content was determined as described above, and final tabulated values were expressed in terms of molecules of EC20 per cell.

As shown in Fig. 6, the cell uptake of 99m Tc-EC20 was found to be dependent on the extracellular concentration. The particular KB cells used were determined to bind a maximum of four million molecules of the folate radiopharmaceutical per cell. Scatchard analysis of the data estimated the K_D of binding to be 3.2 nM, a value comparable with the K_D observed for the vitamin folate binding to these same cells.

Although the full identity of the Peak B component was not established, UV absorption analysis indicated that it contained a folate moiety (i.e., the absorption spectrum contained folate's signature secondary absorption peak at 363 nm). This HPLC-purified radiolabeled material (Peak B material) was collected and then added to cultured KB cells. As shown in Fig. 7, the cell uptake of the 99m Tc-labeled Peak B component was also found to be dependent on the extracellular concentration. Scatchard analysis of the data estimated the K_D of binding to be 1.1 nM. Interestingly, the cell association of Peak B was completely blocked in the presence of excess folic acid, indicating that this minor formulation by-product is also capable of targeting FR-positive cells for radiodiagnostic purposes.

-29-

EXAMPLE 10

Blood Clearance

Animals used for this study were maintained on a folate-free diet (Harlan #TD-90261) for approximately three weeks prior to dose administration.

5 Accumulation to this special diet is essential because regular rodent diets contain large amounts of folic acid (6 mg/kg chow) and promote high serum folate levels in mice. Furthermore, previous studies have shown that mice placed on a folate-free diet for 3 weeks had maintained a safe serum folate level of $25 \pm 7 \text{ nM}$, which is slightly higher than the 9-14 nM concentration measurable in human serum.

10 The $^{99m}\text{Tc-EC20}$ solution was prepared on the day of use and had initially contained 100 μg of EC20 per milliliter. The solution was further diluted with sterile saline to prepare working stock solutions. The radiochemical purity of the product was estimated to be ~94% by TLC. Each animal received a dose of 50 $\mu\text{g/kg}$ EC20 (67 nmol/kg) in approximately 0.1 mL volume i.v. via the tail vein during brief diethyl ether anesthesia. At the designated times (see Fig. 8) post-injection, each animal was euthanized by CO_2 asphyxiation, and blood was immediately collected by cardiac puncture.

15 As shown in Fig. 8, $^{99m}\text{Tc-EC20}$ was rapidly removed from circulation in the Balb/c mouse. The plasma half life of this radiopharmaceutical was estimated to be ~4 minutes, and less than 0.2% of the injected $^{99m}\text{Tc-EC20}$ dose remained in circulation after four hours (assuming that blood represents 5.5% of the total body mass). This data indicates that folate conjugates are rapidly removed from circulation following intravenous administration, and that valuable tissue biodistribution data can be obtained after only a few hours post-injection without the concern for non-specific tissue uptake due to blood-borne radioactivity.

EXAMPLE 11

Tissue Distribution Studies

20 The ability of $^{99m}\text{Tc-EC20}$ to target tumors *in vivo* was assessed using a FR-positive M109 model. These tumor cells are syngeneic for the Balb/c mouse, and they reproducibly form subcutaneous solid tumors within two weeks post inoculation. $^{99m}\text{Tc-EC14}$, which is structurally similar to $^{99m}\text{Tc-EC20}$ except it

-30-

contains one additional D-Glu residue (i.e., Pte-D-Glu-D-Glu- β Dpr-Asp-Cys), 99m Tc-EC 28 (a non-pterinate containing control consisting of benzoyl-D-Glu- β -Glu- β Dpr-Asp-Cys), and the previously reported 111 In-DTPA-folate radiopharmaceutical were also evaluated in this bioassay. Importantly, the 99m Tc-EC 28 control agent will not bind to cell surface FRs because it lacks an essential pteridine ring moiety.

Four to five week-old mice (Balb/c strain) were purchased from Harlan Sprague Dawley, Inc. (Indianapolis, IN) and were maintained on a folate-free diet for a total of three weeks prior to the experiment. Syngeneic FR-positive M 109 tumor cells (1×10^6 per animal) were inoculated in the subcutis of the right axilla two weeks prior to the experiment. All mice were females, and the tumor weights were 54.2 ± 2.8 mg on the day of this experiment. A stock 99m Tc-EC20 solution containing 100 μ g of agent per milliliter was prepared on the day of use, and its radiochemical purity was > 96%. The two additional 99m Tc-chelating agents, 99m Tc-EC14 and 99m Tc-EC28 as well as 111 In-DTPA-folate were also prepared to > 90% radiochemical purity. All solutions were diluted with either saline alone or a saline solution containing 100 equivalents of folic acid (for competition) such that the final radiopharmaceutical concentration was 10 μ mol/mL.

Animals received an approximate 40 μ mol/kg i.v. dose of test article in 1.00 μ L volume via a lateral tail vein during brief diethyl ether anesthesia. Four hours post-injection, animals were sacrificed by CO₂ asphyxiation, and dissected. Selected tissues were removed, weighed, and counted to determine 99m Tc distribution. CPM values were decay-corrected, and results were tabulated as % injected dose per gram of wet weight tissue.

As shown in Table 3 (below), the three "folate" containing radiopharmaceuticals, 99m Tc-EC14, 99m Tc-EC20 and 111 In-DTPA-Folate, predominantly accumulated in the FR-positive tumor and kidneys, however the kidneys concentrated a higher percent injected dose per gram of tissue (%ID/g) than did the tumor. Interestingly, the net tumor accumulation of 111 In-DTPA-Folate and 99m Tc-EC20 was nearly the same (19 and 17% ID/g, respectively), whereas the tumor uptake of 99m Tc-EC14 was somewhat less at ~ 10% ID/g. Nonetheless, all three agents displayed high tumor to blood ratios (> 30 to 1).

-31-

Table 3. Biodistribution of Folate Radiopharmaceuticals in Balb/c Mice Bearing Subcutaneous M109 Tumors.

	% Injected Dose per Gram Tissue (4 hr post i.v. injection)*						
	^{99m} Tc-EC14 + Folic acid	^{99m} Tc-EC14 + Folic acid	^{99m} Tc-EC20	^{99m} Tc-EC20 + Folic acid	¹¹¹ In-DTPA- Folate	¹¹¹ In-DTPA- Folate + Folic acid	^{99m} Tc-EC28
Blood	0.31 ± 0.14	0.19 ± 0.07	0.34 ± 0.03	0.09 ± 0.02	0.21 ± 0.10	0.09 ± 0.04	0.06 ± 0.04
Heart	2.39 ± 0.64	0.08 ± 0.01	1.57 ± 0.26	0.08 ± 0.01	2.57 ± 0.82	0.06 ± 0.02	0.03 ± 0.01
Lung	2.08 ± 0.40	0.15 ± 0.04	2.22 ± 0.63	0.31 ± 0.26	1.72 ± 0.61	0.09 ± 0.2	0.05 ± 0.01
Liver	3.44 ± 2.19	1.37 ± 0.98	3.56 ± 0.25	1.15 ± 0.22	5.21 ± 2.63	0.81 ± 0.03	0.50 ± 0.26
Spleen	2.68 ± 2.49	2.99 ± 1.43	0.95 ± 0.15	0.38 ± 0.33	3.30 ± 2.33	1.46 ± 0.73	0.60 ± 0.38
Intestine	1.70 ± 0.55	0.32 ± 0.11	2.56 ± 0.61	2.93 ± 1.49	1.87 ± 0.69	0.82 ± 0.14	0.47 ± 0.19
Kidney	98.0 ± 40.7	5.94 ± 0.52	138 ± 12.4	5.64 ± 2.13	191 ± 79.2	3.14 ± 1.96	0.62 ± 0.14
Muscle	0.99 ± 0.28	0.09 ± 0.11	0.67 ± 0.20	0.06 ± 0.02	1.19 ± 0.48	0.05 ± 0.04	0.02 ± 0.01
Stomach	1.47 ± 0.58	0.10 ± 0.03	1.45 ± 0.55	3.35 ± 5.19	1.62 ± 0.65	0.25 ± 0.20	0.21 ± 0.19
Tumor	9.83 ± 2.77	0.43 ± 0.52	17.2 ± 1.02	0.45 ± 0.18	19.3 ± 5.86	0.46 ± 0.42	0.11 ± 0.06
Tumor/Blood	34.1 ± 7.41	2.00 ± 2.00	51.0 ± 8.20	4.70 ± 1.30	102 ± 43.4	5.00 ± 4.60	2.00 ± 0.50

* Values shown represent the mean ± s.d. of data from 3 animals.

5

Folate-specific targeting was further demonstrated by two distinct methods. First, the accumulation of ^{99m}Tc-EC14, ^{99m}Tc-EC20 and ¹¹¹In-DTPA-folate in the FR-positive tumor and kidneys was effectively blocked (> 94%) when these agents were co-administered with a 100-fold excess of folic acid. Second, the ^{99m}Tc-EC28 control agent failed to appreciably accumulate in the kidneys and tumor. Both observations show that an intact "folate-like" (or pteroylate) moiety is required to afford targeted uptake and retention of these radiopharmaceutical agents into FR-positive tissues.

15

EXAMPLE 12

Gamma Scintigraphy

M109 tumor cells (1×10^6 per animal¹⁷) were inoculated in the subcutis of the right axilla of Balb/c mice two weeks prior to the experiment. Animals received an approximate 50 $\mu\text{mol}/\text{kg}$ i.v. dose of tracer article in 100 μL volume via a lateral tail vein during brief diethyl ether anesthesia. Four hours post-injection, animals were sacrificed by CO₂ asphyxiation and then placed on top of an image acquisition surface. Whole body image acquisition was performed for 1 minute at a count rate of 50-75,000 counts per minute using a T-technicare Omega 500 Sigma 410 Radioisotope Gamma Camera. All data were analyzed using a Medasys MIS-DOS-based computer equipped with Medasys Pinnacle software.

-32-

Uptake of 99m Tc-EC20 by the FR-positive M109 tumors and kidneys was demonstrated using this gamma scintigraphy protocol. As shown in Fig. 9, a ventral image of a mouse injected with 99m Tc-EC20 as described above localizes the gamma radiation to the two kidneys (K) and the M109 tumor mass (T; shoulder region). No appreciable radiotracer was observed in other body tissues. A similar image profile has been reported for the 111 In-DTPA-Folate radiopharmaceutical.

EXAMPLE 13

Urinary Excretion and Metabolism

10 The urinary HPLC speciation profile of 99m Tc-EC20 was obtained using Balb/c mice. Mice (~20 g each) were injected with 1 mCi (6.7 nmol) of 99m Tc-EC20 via a lateral tail vein. Following a 1, 4, or 6 hour time period, groups of two mice were euthanized by CO₂ asphyxiation and urine was collected. After filtration through a GV 13 Millex filter, the radiochemical speciation was assessed using an 15 HPLC system equipped with a Nova-Pak C18 3.9 x 150 mm column and a radiochemical detector. The system was isocratically eluted with 20% methanol containing 0.1% TFA at a flow rate of 1 mL/minute.

It was previously determined that the primary elimination route for 111 In-DTPA-Folate was via the urine. Similar to the HPLC profile shown in Fig. 2, 20 both the 99m Tc-EC20 standard and the urine samples exhibited four radioactive peaks. As shown in Table 4 (below), the radiochemical purity of the standard (sum of peaks C and D presumably corresponding to the *syn* and *anti* 99m Tc-EC20) remained constant at ~ 93% over the 6 hr duration of this experiment. The amount of free 99m Tc in the standard (peak A) was ~ 2%. Importantly, peak B within this radiochemical 25 profile is believed to be EC20 chelated with 99m Tc at an unconventional, less stable position, however the radioactivity measured in this fraction was not included in the overall radiochemical purity estimation for 99m Tc-EC20. This data collectively indicates that the formulation remained stable in saline solution throughout this 6 hr investigation.

30 After 1 and 4 hours post-injection into Balb/C mice, the radiochemical speciation profile of 99m Tc-EC20 in the mouse urine did not change. The radioactivity present in the urine at 6 hours post-injection, however, was too low to accurately

-33-

assay by HPLC. The proportion of parent drug among radioactive species recovered in urine remained relatively constant at approximately 90% throughout the four hours during which it could be quantitated. This value is very similar to the 93% purity of the standard indicating that 99m Tc-EC20 is predominately excreted into the urine in an unmodified form.

10 **Table 4. Excretion and Metabolism of 99m Tc-EC20 from the Balb/c Mouse.** Mice were injected with 1 mCi (6.7 nmol) of 99m Tc-EC20 via a lateral tail vein. At the indicated times, groups of two mice were euthanized and urine was collected. The radiochemical speciation was then determined by HPLC. The area percent sum of peaks C and D (syn and anti isomers) is used to calculate the overall purity of intact 99m Tc-EC20.

Peak	RT (min)	Area Percent							
		99m Tc-EC20 Standard			Urine Samples (two mice/timepoint)				
		0 hr	1 hr	6 hr	1 hr		4 hr		
A (pertechnetate)	1.4	2	2.1	1.8	8.3	6.3	9.4	10.2	
B (unknown)	3.4	4.5	4.5	4.8	2.5	2.6	5.4	0	
C (isomer 1)	5.5	15.5	15.7	15.9	20.4	18.1	7.3	11.1	
D (isomer 2)	18.5	78	77.7	77.5	68.8	73	77.9	78.7	
Sum C and D		93.5	93.4	93.4	89.2	91.1	85.2	89.8	

15

EXAMPLE 14

Serum Protein Binding

Fresh rat serum, and commercial male human serum (type AB donors, Sigma Chemical Co.) were used to evaluate *in vitro* binding of 99m Tc-EC20 to serum proteins. One minute after 99m Tc-EC20 was mixed with 1 mL of serum at room temperature, 0.3 mL of the serum solution was transferred to a clean Amicon Centrifree® ultrafiltration device (30,000 NMWL) in triplicate. Within one minute of loading the centrifuge with the serum solution, the device was spun at 1000 x g for 20 minutes at 20°C. 50 μ L samples of the original solution, and off the filtrate from each device, was transferred to a clean tube and counted in an automatic gamma counter. A control solution of 99m Tc-EC20 mixed with 1 mL of normal saline was ultrafiltered in an identical fashion. The percentage of free 99m Tc was calculated for each of the three samples.

-34-

While 99m Tc-EC20 exhibited only a minor level of non-specific binding to the ultrafiltration device (~ 5%), approximately 70% of it was found to predominantly associate with the > 30kDa serum protein fraction in solutions of rat or human serum (69% and 72%, respectively). Importantly, since 99m Tc-EC20 does effectively and preferentially accumulate within FR-positive tissues (see Table 2 and Fig. 8), its apparent affinity for serum proteins does not appear to affect this radiotracer's ability to target FRs *in vivo*.

EXAMPLE 15

Tissue Distribution Studies

The protocols used in this example are similar to those described in Example 11. The ability of 99m Tc-EC20 to target tumors *in vivo* was further assessed using FR-positive M109 and FR-negative 4T1 tumor models. Six week-old female Balb/c mice (n = 3/dose group) were purchased from Harlan Sprague Dawley, Inc. (Indianapolis, IN) and were maintained on a folate-free diet (Harlan TEKLAD) for a total of seven days prior to tumor cell inoculation.

Syngeneic, FR-positive M109 tumor cells (2×10^6 P_0 per animal) or FR-negative 4T1 cells (5×10^5 P_0 per animal) were inoculated subcutaneously in 100 μl of folate-free RPMI-1640 containing 1% syngeneic mouse serum. A stock 99m Tc-EC20 solution containing 100 μg of agent per milliliter was prepared on the day of use as described above.

Sixteen days after tumor cell inoculation, the animals were injected intravenously with 500 or 1800 nmoles/kg of EC20 for M109 tumor-bearing animals and 500 nmoles/kg of EC20 for 4T1 tumor-bearing animals (3 mice per dose group). All injections were in 100 μl volumes. Four hours post-injection, animals were sacrificed by CO_2 asphyxiation, and blood was collected by cardiac puncture and the animals were dissected. Selected tissues (heart, lungs, liver, spleen, kidney, intestines, stomach, muscle, and tumor) were removed, weighed, and counted in an automatic gamma counter to determine 99m Tc distribution. Uptake of the radiopharmaceutical in terms of percentage injected dose of wet weight tissue (% ID/g) was calculated by reference to standards prepared from dilutions of the injected preparation.

-35-

As shown in Fig. 11, folate receptor-specific targeting was demonstrated because 99m Tc-EC20 predominantly accumulated in the FR-positive M109 tumors and kidneys, and not in the FR-negative 4T1 tumors. Uptake in the FR-negative 4T1 tumors was 7.6-fold lower than in the FR-positive M109 tumors.

5 Uptake of 99m Tc-EC20 in normal tissues, except kidney as expected, was low. These results show that 99m Tc-EC20 targeting is FR-specific.

EXAMPLE 16

Tissue Distribution Studies

10 The protocols used in this example are similar to those described in Example 11. The ability of 99m Tc-EC11 (peptide -A₁), 99m Tc-EC13 (peptide-A₃), and 99m Tc-EC14 (peptide-A₂) to target tumors *in vivo* was assessed using the FR-positive KB tumor model. Four week-old male nude mice (n = 4/group) were maintained on a folate-free diet for a total of ten days prior to tumor cell inoculation.

15 FR-positive KB tumor cells (0.25 x 10⁶ per animal) were inoculated subcutaneously in the intracapsular region. Fourteen days after tumor cell inoculation, the animals (n = 4/group) were injected intravenously with 99m Tc-EC11, 99m Tc-EC13, or 99m Tc-EC14 at the doses (about 12 μ g/kg) of the conjugates shown in Table 5 below. Stocks of 99m Tc-EC11, 99m Tc-EC13, and 99m Tc-EC14 solutions were prepared on the day of use as described above. About a 20-fold excess of free folate (about 200 μ g/kg) was co-administered to control animals (n = 4/group). Four hours post-injection, animals were sacrificed by CO₂ asphyxiation, and blood was collected by cardiac puncture and the animals were dissected. Selected tissues were removed, weighed, and counted in an automatic gamma counter to determine 99m Tc distribution.

20 25 Uptake of the radiopharmaceutical in terms of percentage injected dose of wet weight tissue (% ID/g) was calculated by reference to standards prepared from dilutions of the injected preparation.

As shown in Table 5, folate receptor-specific targeting was demonstrated because 99m Tc-EC11, 99m Tc-EC13, and 99m Tc-EC14 predominantly accumulated in the FR-positive KB tumors and kidneys. The accumulation was blocked by co-administration of free folate. These results show that 99m Tc-EC11, 99m Tc-EC13, and 99m Tc-EC14 can target tumors *in vivo* in a FR-specific manner.

-36-

Similar results (see Table 6 below) were obtained with ^{99m}Tc-EC 53 (the all D-enantiomer of EC20) using similar protocols except that the dose of ^{99m}Tc-EC53 was about 50 µg/kg and about a 100-fold excess of free folate or cold EC53 was used. As shown in Table 6, folate receptor-specific targeting was demonstrated because ^{99m}Tc-EC53 predominantly accumulated in the FR-positive KB tumors and kidneys. The accumulation was blocked by co-administration of free folate. These results show that ^{99m}Tc-EC53 can target tumors *in vivo* in a FR-specific manner.

5

- 37 -

TABLE 5

		Percentage of Injected 99m Tc Dose per Gram (Tissue Wet Mass)					
		Peptide A-Folate (EC11)	Peptide A ₂ -Folate (EC14)	Peptide A ₂ -Folate (EC13)	Peptide A ₂ -Folate (EC13)	HYNIC-Folate	
<i>Tumor mass (g):</i>	0.112 \pm 0.027	0.125 \pm 0.032	0.160 \pm 0.037	0.171 \pm 0.044	0.213 \pm 0.067	0.0773 \pm 0.041	0.179 \pm 0.050
<i>Animal Mass (g):</i>	28.9 \pm 1.3	27.1 \pm 1.6	27.6 \pm 0.6	27.3 \pm 2.7	30.0 \pm 1.3	27.5 \pm 1.4	29.0 \pm 2.0
<i>Animal Quantity & Gender:</i>	4M	4M	4M	4M	4M	4M	4M
<i>Folate-Conjugate Dose (μg/kg):</i>	11.9 \pm 0.5	13.04 \pm 1.12	12.6 \pm 0.4	12.87 \pm 1.53	11.7 \pm 0.9	13.03 \pm 1.05	1.58 \pm 0.03
<i>Folic Acid Dihydrate Dose† (μg/kg):</i>	0	195.1 \pm 17.9	0	192.6 \pm 22.9	0	191.7 \pm 15.5	0
<i>Blood:</i>	0	0.41 \pm 0.04	0	0.40 \pm 0.05	0	0.40 \pm 0.03	0
<i>Heart:</i>	0.21 \pm 0.01	0.25 \pm 0.01	0.19 \pm 0.02	0.12 \pm 0.02	0.25 \pm 0.03	0.16 \pm 0.02	0.31 \pm 0.06
<i>Lungs:</i>	2.5 \pm 0.3	0.36 \pm 0.03	1.0 \pm 0.5	0.24 \pm 0.01	1.6 \pm 0.1	0.20 \pm 0.03	3.4 \pm 0.3
<i>Liver & Gall Bladder:</i>	1.2 \pm 0.2	0.35 \pm 0.03	1.6 \pm 0.3	0.24 \pm 0.02	1.1 \pm 0.1	0.23 \pm 0.02	1.2 \pm 0.2
<i>Spine:</i>	5.4 \pm 1.4	1.6 \pm 0.1	4.5 \pm 1.0	0.68 \pm 0.07	3.5 \pm 0.7	0.65 \pm 0.04	3.9 \pm 0.9
<i>Kidney (one):</i>	0.38 \pm 0.03	0.23 \pm 0.01	0.41 \pm 0.06	0.15 \pm 0.01	0.39 \pm 0.06	0.15 \pm 0.02	0.39 \pm 0.12
<i>Stomach, Intestines & Contents:</i>	67.8 \pm 6.9	55.5 \pm 2.3	44.2 \pm 6.4	20.6 \pm 2.4	54.5 \pm 2.4	29.1 \pm 2.2	41.2 \pm 8.4
<i>Muscle:</i>	1.4 \pm 0.1	1.1 \pm 0.3	1.4 \pm 0.1	0.50 \pm 0.10	2.5 \pm 0.2	3.7 \pm 0.7	1.5 \pm 0.2
<i>Tumor:</i>	2.95 \pm 0.57	1.47 \pm 0.24	5.57 \pm 0.76	2.0 \pm 0.5	1.7 \pm 0.2	0.21 \pm 0.03	2.2 \pm 0.3
<i>Tumor/Blood:</i>	13.7 \pm 2.1	5.9 \pm 0.9	29.3 \pm 5.2	17.0 \pm 4.5	22.5 \pm 1.4	10.2 \pm 1.5	15.1 \pm 2.1
<i>Tumor/Liver:</i>	0.57 \pm 0.14	0.94 \pm 0.09	1.3 \pm 0.3	3.0 \pm 0.6	1.6 \pm 0.2	2.6 \pm 0.3	1.3 \pm 0.3
<i>Tumor/Kidney:</i>	0.043 \pm 0.005	0.027 \pm 0.005	0.13 \pm 0.03	0.10 \pm 0.01	0.10 \pm 0.01	0.058 \pm 0.010	0.11 \pm 0.01
<i>Tumor/Muscle:</i>	1.6 \pm 0.3	3.9 \pm 0.08	2.4 \pm 0.6	7.7 \pm 1.7	3.3 \pm 0.3	8.2 \pm 2.0	2.1 \pm 0.4

Values shown represent the mean \pm standard deviation. Blood was assumed to account for 11% of total body mass. Tumor/Background Tissue ratios based on corresponding % Injected Dose per Gram data.

† Folic Acid dose co-injected with the 99m Tc-Folate-Conjugate dose.
 n = 3

- 38 -

TABLE 6

	^{99m} Tc-ECS3		^{99m} Tc-ECS3 plus Folic Acid		^{99m} Tc-ECS3 plus EC53	
	Average	STD	Average	STD	Average	STD
Test Article Dose (ug/se) (mmol/kg)	50.0 67.0		50.0 67.0		50.0 67.0	
Co-dosed competitor (nmol/kg)			6700.0		6700.0	
Tumor mass (g)	0.2	0.17	0.20	0.14	0.19	0.15
Animals	3F		3F		3F	
Blood	0.38	0.03	0.244	0.028	0.24	0.06
Heart	1.09	0.28	0.15	0.03	0.19	0.03
Lung	0.89	0.30	0.23	0.04	0.26	0.07
Liver	3.86	0.96	4.40	1.15	3.77	0.48
Intestine	3.53	0.86	4.33	3.67	3.96	1.86
Kidney	77.99	6.19	10.12	6.91	7.97	1.52
Muscle	0.76	0.31	0.11	0.04	0.12	0.06
Spleen	0.67	0.22	0.27	0.06	0.41	0.11
Stomach	1.04	0.36	0.30	0.15	0.18	0.01
Tumor	11.77	4.26	0.53	0.11	1.88	1.56
Tumor/Kidney	31.4	13.7	2.2	0.6	4.5	4.8
Tumor/liver	3.4	2.2	0.1	0.0	0.3	0.1
Tumor/Muscle	17.1	7.0	5.2	1.4	8.5	8.0
Tumor/Kidney	0.2	0.07	0.07	0.05	0.14	0.16

DISCUSSION

The invention provides a conjugate of a vitamin and a radionuclide chelator for clinical development as an imaging agent. Exemplary of such an imaging agent is the newly designed, synthesized, and radiochemically characterized folate-based radionuclide chelator, ^{99m}Tc -EC20.

^{99m}Tc -EC20, a small molecular weight peptide derivative of folate that contains a D- γ -Glu peptide linkage (see Fig. 1), was synthesized using an efficient solid-phase synthetic procedure. In its natural form, folate (or pterooyl-glutamate) has a single glutamyl residue present in an L configuration. However, a D-Glu enantiomer residue was incorporated into the EC20 molecule. Importantly, similar to EC20, substitution of the L-Glu residue in folic acid with a D-Glu residue does not alter the ability of folic acid to bind to the high affinity FR.

EC20 was found to efficiently chelate ^{99m}Tc when in the presence of α -D-glucoheptonate and tin (II) chloride. When analyzed by radiochemical HPLC, > 95% of the resulting ^{99m}Tc -EC20 formulation consisted of a mixture of *syn* and *anti* stereoisomers, each equally capable of binding to FR with high affinity (see Fig. 3). Approximately 3% of the ^{99m}Tc in the formulation was chelated to EC20 at some other site on the EC20 molecule than the expected Dap-Asp-Cys moiety. Although this component was not isolated in sufficient quantity for optimal characterization, it was shown to bind to FR with high affinity (see Fig. 6). Finally, the remaining 2% of the radioactivity in the ^{99m}Tc -EC20 formulation was attributed to free ^{99m}Tc .

^{99m}Tc -EC20 demonstrated both time- and concentration-dependent association with FR-positive cells. ^{99m}Tc -EC20 was rapidly cleared from the blood ($t_{1/2} \sim 4$ min), which is important for diagnostic imaging agents, and ^{99m}Tc -EC20 preferentially accumulated in large amounts within FR-positive tumors.

The performance of ^{99m}Tc -EC20 was directly compared to that of a similar FR targeting agent, ^{111}In -DTPA-Folate, using two different methods. First, both folate-based radiopharmaceuticals were found to equally compete with folic acid for binding to KB FRs (see Fig. 3 and Table 1). Second, the biodistribution of each agent in tumor-bearing mice was nearly identical (see Table 2). High tumor uptake and tumor-to-blood ratios were measured for ^{99m}Tc -EC20. Taken together these

-40-

results suggest that like ^{111}In -DTPA-folate, $^{99\text{m}}\text{Tc}$ -EC20 will effectively localize in FR-positive tumors when clinically administered to patients.

Several folate-based $^{99\text{m}}\text{Tc}$ conjugates have previously been described. Limited biodistribution data is available on a $^{99\text{m}}\text{Tc}$ -12-amino-3,3,9,9-tetramethyl-5-oxa-4,8 diaza-2,1 0-dodecanedioxime (OX-A) folate conjugate, however moderate levels (~ 7% ID/g) of tracer uptake in a KB tumor was reported. Studies involving the biodistribution of a $^{99\text{m}}\text{Tc}$ -ethylene-dicysteine-folate conjugate in mammary tumor-bearing rats were also reported. The rats in that study were fed a folate-rich diet. Thus, low tumor uptake and low tumor-to-blood ratios were obtained. Lastly, a $^{99\text{m}}\text{Tc}$ -6-hydrazinonicotinamido-hydrazido (HYNIC) folate derivative (HYNIC-folate) was shown to accumulate in large amounts within 24JK-FBP tumors. Interestingly, $^{99\text{m}}\text{Tc}$ -EC20 accumulated within M109 tumors to nearly identical levels as that of HYNIC-folate in 241JK-FBP tumors (~ 17% ID/g) (Table 2). These two agents also displayed roughly 50:1 tumor-to-blood ratios at 4 hours post intravenous injection.

In summary, a new peptide derivative of folate was created to efficiently chelate $^{99\text{m}}\text{Tc}$. This new compound, $^{99\text{m}}\text{Tc}$ -EC20, avidly binds to FR-positive tumor cells *in vitro* and *in vivo*. EC20 was found to bind cultured folate receptor (FR)-positive tumor cells in both a time- and concentration-dependent manner with very high affinity ($K_D \sim 3 \text{ nM}$). Using an *in vitro* relative affinity assay, EC20 was also found to effectively compete with ^3H -folic acid for cell binding when presented either alone or as a formulated metal chelate. Following intravenous injection into Balb/c mice, $^{99\text{m}}\text{Tc}$ -EC20 was rapidly removed from circulation (plasma $t_{1/2} \sim 4 \text{ min}$) and excreted into the urine in a non-metabolized form. Data from gamma scintigraphic and quantitative biodistribution studies performed in M109 tumor-bearing Balb/c mice confirmed that $^{99\text{m}}\text{Tc}$ -EC20 predominantly accumulates in FR-positive tumor and kidney tissues. These results show that $^{99\text{m}}\text{Tc}$ -EC20 is an effective, non-invasive radiodiagnostic imaging agent for the detection of FR-positive tumors. Other EC20-related imaging agents were also shown to be effective, including EC11, EC13, EC14, and EC53.


Each year ~ 26,000 women in the United States are diagnosed with ovarian cancer, and less than 50% of those women survive more than five years. One

-41-

reason for the low survival rate is the difficulty in diagnosing this form of cancer. Because of the fear of rupturing an unidentified abdominal mass and the potential for spreading cancer throughout the abdominal cavity, fine needle biopsy is not often performed. Rather, the diagnosis and staging of suspicious ovarian masses is typically done through surgical laparotomy, which is an invasive and expensive procedure. Since ^{99m}Tc-EC20 binds tightly to FR present in large amounts on ovarian cancers (among others), this radiopharmaceutical provides an inexpensive, non-invasive but reliable method for the early diagnosis of malignant ovarian cancer. Importantly, ^{99m}Tc-EC20 may also help guide the clinical decision process by making possible more definitive and earlier diagnosis of recurrent or residual disease.

CLAIMS:

1. A compound of the formula



wherein V is a vitamin that is a substrate for receptor-mediated transmembrane transport in vivo, or a vitamin receptor binding derivative or analog thereof;

L is a divalent linker;

R is a side chain of an amino acid;

M is a cation of a radionuclide;

n is 1 or 0; and

k is 1 or 0.


2. The compound of claim 1 wherein V is a vitamin selected from the group consisting of folate, riboflavin, thiamine, vitamin B₁₂, and biotin, or a vitamin receptor binding derivative or analog thereof.

3. The compound of claim 1 wherein the radionuclide is selected from the group consisting of isotopes of gallium, indium, copper, technetium, and rhenium.

4. The compound of claim 3 wherein the radionuclide is an isotope of technetium.


5. The compound of claim 1 wherein V is folate, or a folate receptor binding analog or derivative thereof.

6. The compound of claim 1 having the formula




- 43 -


7. The compound of claim 1 having the formula



8. The compound of claim 1 having the formula




9. The compound of claim 1 having the formula




10. The compound of claim 1 having the formula

AMENDED SHEET



- 44 -



11. The compound of claim 1 having the formula



12. The compound of claim 1 having the formula

13. A composition for diagnostic imaging comprising a compound of the formula $\text{V}-(\text{L})_n-\text{NH}-\text{CH}(\text{H}_2\text{N})-\text{CH}(\text{CO}-\text{NH}-\text{CH}(\text{R})\text{CO}-\text{NH}-\text{CH}(\text{COOH})\text{CH}_2-\text{SH}$ 

AMENDED SHEET

1/23




FIG. 1




FIG. 2

SUBSTITUTE SHEET (RULE 26)

2/23




FIG. 3




FIG. 4

SUBSTITUTE SHEET (RULE 26)

3/23




FIG. 5




FIG. 6

SUBSTITUTE SHEET (RULE 26)

4/23





FIG. 7

FIG. 8
SUBSTITUTE SHEET (RULE 26)

5/23




FIG. 9

SUBSTITUTE SHEET (RULE 26)

6/23






FIG. 10A

SUBSTITUTE SHEET (RULE 26)

7/23








FIG. 20B

SUBSTITUTE SHEET (RULE 26)

8/23



EC31 (Pte-L-βDpr-L-Asp-L-Cys)



EC53 (Pte-D-γGlu-D-βDpr-D-Asp-D-Cys)

FIG. 10C

SUBSTITUTE SHEET (RULE 26)

9/23




FIG. 21

SUBSTITUTE SHEET (RULE 26)

10/23




FIG. 12

SUBSTITUTE SHEET (RULE 26)

11/23





FIG. 13

SUBSTITUTE SHEET (RULE 26)

12/23



13/23





FIG. 15

SUBSTITUTE SHEET (RULE 26)


14/23

FIG. 16



SUBSTITUTE SHEET (RULE 26)

15/23




SUBSTITUTE SHEET (RULE 26)

16/23

MOBILE PHASE A=0.1%TFA IN H₂O
 B=0.1%TFA IN CH₃CN
 COLUMN: SYMMETRY C18 (4.6X150 mm). INJECTION: 20 μ L.
 GRADIENT 0% B - 100% B 30MIN. FLOW RATE: 1 mL/min.
 CHROMATOGRAM AT 280 nm.
 BY LE-CUN XU, Ph.D. ENDOCYTE, INC.

SAMPLE NAME	A5(OLD SAMPLE)	SAMPLE TYPE	UN-KNOWN
VIAL	1	DATE ACQUIRED	
INJECTION	1	ACQ METHOD SET	60% B 100% B 30MIN 1MLMIN
INJECTION VOLUME	20.00 μ L	PROCESSING METHOD	LC DEFAULT PROCESSING
CHANNEL	996	DATE PROCESSED	
RUN TIME	20.0 MINUTES		

AUTO-SCALED CHROMATOGRAM



SAMPLE NAME	A5(REF-PURIFIED SAMPLE)	SAMPLE TYPE	UNKNOWN
VIAL	1	DATE ACQUIRED	
INJECTION	1	ACQ METHOD SET	60% B 100% B 30MIN 1MLMIN
INJECTION VOLUME	20.00 μ L	PROCESSING METHOD	LC DEFAULT PROCESSING
CHANNEL	996	DATE PROCESSED	
RUN TIME	30.0 MINUTES		

AUTO-SCALED CHROMATOGRAM





FIG. 18
 SUBSTITUTE SHEET (RULE 26)


17/23



SUBSTITUTE SHEET (RULE 26)

18/23

PROJECT NAME: EC20
 SAMPLE NAME: Pro- β Dop-Asp-Cys FRACTION #2
 INJECTION VOLUME: 20.00 μ l
 DATE ACQUIRED:
 AQC METHOD: SET: g1% B20M IN 20% B 1_5 min AND Eq
 PROCESSING METHOD: 15
 CURRENT DATE: DATE PROCESSED:




PEAK RESULTS

	NAME	RT	AREA	HEIGHT	%AREA
1	19.433	976094	7436	1.37	

FIG. 20

SUBSTITUTE SHEET (RULE 26)

19/23



NAME	RT	AREA	HEIGHT	% AREA
1	13.102	98094	7436	1.37
2	15.325	131550	9546	1.88
3	15.795	5974085	505938	85.48
4	16.360	390992	26982	5.59
5	16.877	322515	21694	4.61
6	17.189	79995	7670	1.06

FIG. 21
 SUBSTITUTE SHEET (RULE 26)

20/23

#1 STATUS LOG TIME: 0.01

37 STATES API SOURCE

ALI SOURCE	:	3.52
SOURCE VOLTAGE (V)	:	0.24
SOURCE CURRENT (mA)	:	NO
VAPORIZER THERMOCOUPLE OK:	:	-0.00
VAPORIZER TEMP (C)	:	58.98
SHEATH GAS FLOW RATE (l)	:	3.17
AUX GAS FLOW RATE (l)	:	YES
CAPILLARY RTD OK	:	20.64
CAPILLARY VOLTAGE (V)	:	224.60
CAPILLARY TEMP (C)	:	NO
B. HV SUPPLY AT LIMIT	:	

VACUUM	:	YES
VACUUM OK	:	YES
ION GAUGE PRESSURE OK	:	YES
ION GAUGE STATUS	:	ON
ION GAUGE (X1 0-5 TORR)	:	1.52
CONVECTION PRESSURE OK	:	YES
CONVECTION GAUGE (TORR)	:	0.84
TURBO PUMP		
STATUS	:	RUNNING
LIFE (HOURS)	:	16359.00
SPEED (RPM)	:	60000.00
POWER (WATTS)	:	39.20
TEMPERATURE (C)	:	40.00

ION OPTICS






FIG. 22A

SUBSTITUTE SHEET (RULE 26)

21/23



SUBSTITUTE SHEET (RULE 26)

FIG. 22B

22/23

#1 STATUS LOG TIME: 0.003

API SOURCE

SOURCE VOLTAGE (KV)	:	4.52
SOURCE CURRENT (mA)	:	19.48
VAPORIZER THERMOCOUPLE OK:	:	NO
VAPORIZER TEMP (C)	:	-0.00
SHEATH GAS FLOW RATE (L)	:	58.93
AUX GAS FLOW RATE (L)	:	3.06
CAPILLARY RTD OK	:	YES
CAPILLARY VOLTAGE (V)	:	-10.10
CAPILLARY TEMP (C)	:	225.60
8 KV SUPPLY AT LIMIT	:	NO

VACUUM	:	
VACUUM OK	:	YES
ION GAUGE PRESSURE OK	:	YES
ION GAUGE STATUS	:	ON
ION GAUGE (x10 ⁻⁵ TORR)	:	1.52
CONVECTION PRESSURE OK	:	YES
CONVECTION GAUGE (TORR)	:	0.84

TURBO PUMP

STATUS	:	RUNNING
LIFE (HOURS)	:	16359.00
SPEED (RPM)	:	60000.00
POWER (WATTS)	:	39.20
TEMPERATURE (C)	:	40.00

ION OPTICS




FIG. 23A

SUBSTITUTE SHEET (RULE 26)

23/23





FIG. 23B

SUBSTITUTE SHEET (RULE 26)