



(86) Date de dépôt PCT/PCT Filing Date: 2019/03/05
(87) Date publication PCT/PCT Publication Date: 2019/09/12
(45) Date de délivrance/Issue Date: 2022/11/29
(85) Entrée phase nationale/National Entry: 2020/09/08
(86) N° demande PCT/PCT Application No.: CN 2019/077028
(87) N° publication PCT/PCT Publication No.: 2019/170088
(30) Priorité/Priority: 2018/03/06 (CN201810182708.1)

(51) Cl.Int./Int.Cl. *C07D 498/04* (2006.01),
A61K 31/5383 (2006.01), *A61P 35/00* (2006.01)

(72) Inventeurs/Inventors:
ZHANG, QIANG, CN;
YU, SHANNAN, CN;
WANG, ZHONGXIANG, CN;
FENG, SHOUYE, CN;
ZHENG, NANQIAO, CN;
YANG, HAILONG, CN;


...

(73) Propriétaire/Owner:
BEIJING SCITECH-MQ PHARMACEUTICALS LIMITED,
CN

(74) Agent: BLANEY MCMURTRY LLP

(54) Titre : COMPOSE DE TYPES OXAZINO-QUINAZOLINE ET OXAZINO-QUINOLINE, METHODE DE PREPARATION ET UTILISATIONS CONNEXES

(54) Title: OXAZINO-QUINAZOLINE AND OXAZINO-QUINOLINE TYPE COMPOUND, PREPARATION METHOD AND USES THEREOF



(57) Abrégé/Abstract:

Provided are an oxazino-quinazoline and oxazino-quinoline type compound, a preparation method, and uses thereof. More particularly provided is a compound shown in formula (I), an isomer, a hydrate, a solvate, a pharmaceutically acceptable salt, a

(72) Inventeurs(suite)/Inventors(continued): YANG, LEIFU, CN; ZHANG, HONGBO, CN; ZHOU, LIKAI, CN;
XU, ZHANQIANG, CN


(57) Abrégé(suite)/Abstract(continued):

prodrug thereof, a preparation method, and uses thereof in preparing a drug acting as a kinase inhibitor.

(see formula I)

ABSTRACT

Provided are an oxazino-quinazoline and oxazino-quinoline type compound, a preparation method, and uses thereof. More particularly provided is a compound shown in formula (I), an isomer, a hydrate, a solvate, a pharmaceutically acceptable salt, a prodrug thereof, a preparation method, and uses thereof in preparing a drug acting as a kinase inhibitor.



Oxazino-quinazoline and Oxazino-quinoline Type Compound, Preparation Method and Uses Thereof

TECHNICAL FIELD


5 [0001] The present disclosure belongs to the technical field of medicinal chemistry and specifically relates to an oxazino-quinazoline and oxazino-quinoline compound, preparation method thereof and application thereof.

BACKGROUND OF THE INVENTION

10 [0002] Protein kinase is an important signal messenger of cell life activity, which can catalyze the transfer of γ -phosphate group at the terminal of the ATP to the hydroxyl receptor in the amino acid residues (serine, threonine, tyrosine) of the substrate, thereby activating the target protein (Johnson L. N., and Lewis R. J., (2001) Structural basis for control by phosphorylation. *Cheminform.* 101, 2209). Protein kinases are involved in many physiological processes,
15 including cell proliferation, survival, apoptosis, metabolism, transcription, and differentiation (Adams J. A., (2001) Kinetic and catalytic mechanisms of protein kinases. *Chemical reviews.* 101, 2271). Among the existing drug targets in the human body, members of the protein kinase family account for up to 10% (Santos R., Ursu O., Gaulton A., et al. (2017) A comprehensive map of molecular drug targets. *Nature Reviews Drug Discovery.* 16, 19).

20 [0003] Epidermal growth factor receptor (ErbB) tyrosine kinase can regulate cell proliferation, migration, differentiation, apoptosis, and cell movement through a variety of pathways. In many forms of malignant tumors, members of the ErbB family and part of their ligands are often overexpressed, amplified, or mutated, which makes them become important therapeutic targets. The family of protein kinases includes: ErbB1/EGFR/HER1, ErbB2/HER2, ErbB3/HER3 and
25 ErbB4/HER4. Wherein, EGFR and HER2 are important targets for the development of non-small cell lung cancer and breast cancer drugs (Dienstmann R., et. al., (2001) Personalizing Therapy with Targeted Agents in Non-Small Cell Lung Cancer. *ONCOTARGET.* 2(3), 165.; Mitri Z., et. al. (2012) The HER2 Receptor in Breast Cancer: Pathophysiology, Clinical Use, and New Advances in Therapy., *Chemotherapy Research & Practice.*, Volum 2012 (23),
30 743193). In addition, in terms of structural characteristics, EGFR and HER2 are highly conserved in the intracellular tyrosine kinase structural region (ATP pocket) containing catalytically active sites. Therefore, EGFR and HER2 kinase inhibitors that are already marketed and under research often have similar chemical structures.


[0004] The kinase inhibitors Gefitinib, Erlotinib, and Icotinib target EGFR for the treatment of non-small cell carcinoma. Afatinib, Lapatinib and Neratinib target HER2 and EGFR, wherein Afatinib is used to treat non-small cell carcinoma, Lapatinib and Neratinib are used to treat breast cancer. These kinase inhibitors all contain a quinazoline or quinoline core, and a hydrophobic aromatic substituent is introduced at the 4-position connected through the heteroatoms.



[0005] The above-mentioned marketed kinase inhibitor drugs generally have shortcomings such as drug resistance and severe toxic side effects. In particular, Lapatinib and Neratinib produce serious gastrointestinal side effects after administration, including emesis and diarrhea. Therefore, there is urgent need to develop new HER2 and EGFR kinase inhibitor drugs. Structure-based drug design strategies can discover new active molecules with better efficacy, drug metabolism and drug toxicology, and the development of new active molecules with novel core structures can often facilitate the discovery of a whole series of new drug molecules.

SUMMARY OF THE INVENTION

[0006] The present disclosure provides a compound represented by formula (I), or isomers, hydrates, solvates, pharmaceutically acceptable salts or prodrugs thereof,




Formula (I)

[0007] in the formula (I),

[0008] X is O, or NH;

[0009] Y is N or C-Z, wherein, Z is -H or -CN;



5 [0011] R³ is -H, halogen, C₁-C₃ alkyl, C₁-C₃ alkoxy, C₁-C₃ alkyl substituted by halogen, or C₁-C₃ alkoxy substituted by halogen;




[0013] M is O or S;

[0014] T is linear C₁-C₃ alkyl, or linear C₁-C₃ alkyl independently substituted by R⁵ and R⁶,
10 respectively;

[0015] R⁵ and R⁶ are independently -H, halogen, C₁-C₃ alkyl, or C₁-C₃ alkyl substituted by halogen;

[0016] R⁴ is aryl, 5- to 6-membered heteroaryl, aryl substituted by 1-3 identical or different R⁷, or 5- to 6-membered heteroaryl substituted by 1-3 identical or different R⁷, wherein the
15 heteroaryl group is a heteroaryl group containing 1-3 heteroatoms selected from N, O or S;

[0017] R⁷ is -H, halogen, amino, hydroxy, cyano, C₁-C₃ alkylthio, amino substituted with mono- or di-C₁-C₃ alkyl, C₃-C₄ cycloalkyl, unsubstituted or substituted C₁-C₆ alkyl, or unsubstituted or substituted C₁-C₆ alkoxy, wherein the substituent of the substituted C₁-C₆ alkyl is halogen, hydroxy, amino substituted with mono- or di-C₁-C₃ alkyl, or C₁-C₃ alkoxy, and
20 wherein the substituent of the substituted C₁-C₆ alkoxy is halogen, C₁-C₃ alkoxy, or amino substituted with mono- or di-C₁-C₃ alkyl;



[0020] T₁ is linear C₁-C₈ alkyl, or linear C₁-C₈ alkyl independently substituted by R⁹ and R¹⁰,
25 respectively;

[0021] R⁹ and R¹⁰ are each independently -H, or C₁-C₃ alkyl;


[0022] R⁸ is -H, hydroxy, C₁-C₃ alkyl, C₃-C₇ cycloalkyl, C₁-C₃ alkoxy, C₁-C₃ alkylthio, 4- to 7-membered heterocyclyl or -NR¹¹R¹²;

[0023] R¹¹ and R¹² are each independently -H, C₁-C₆ alkyl, C₃-C₆ cycloalkyl, C₁-C₆ alkyl
30 substituted by hydroxy, or C₁-C₆ alkyl substituted by C₁-C₃ alkoxy;

[0024] the 4- to 7-membered heterocyclyl is a heterocyclyl containing 1-2 heteroatoms

selected from N, O or S, the heterocyclcyl is unsubstituted or substituted by one or two of the group consisting of: C₁-C₃ alkyl, aldehyde group, C₁-C₄ alkylacyl, aminoacyl, aminoacyl wherein the amino is substituted with mono- or di-C₁-C₃ alkyl, C₁-C₃ alkylsulfonyl, and C₁-C₃ alkylsulfinyl, or the sulfur in the heterocycle is oxidized by one to two oxygen atoms.

5 [0025] In one alternative embodiment,



[0027] R³ is -H, fluorine, chlorine, bromine, methyl, ethyl, methoxy, ethoxy, trifluoromethyl or trifluoromethoxy;



10 [0029] M is O or S;


[0030] T is linear C₁-C₂ alkyl, or linear C₁-C₂ alkyl independently substituted by R⁵ and R⁶, respectively;

[0031] R⁵ and R⁶ are each independently is -H, -F, methyl, ethyl or trifluoromethyl;

[0032] R⁴ is aryl, 5- to 6-membered heteroaryl, aryl substituted by 1-2 identical or different R⁷, or 5- to 6-membered heteroaryl substituted by 1-2 identical or different R⁷, wherein the aryl or heteroaryl group is selected from the group consisting of: phenyl, pyridyl, pyrimidinyl, thiazolyl, thienyl, pyrrolyl, thiadiazolyl, furyl, oxazolyl or isoxazolyl;

[0033] R⁷ is -H, -F, -Cl, -Br, -CF₃, -OCF₃, amino, hydroxy, cyano, methylthio, ethylthio, propylthio, isopropylthio, methylamino, ethylamino, dimethylamino, diethylamino, cyclopropyl, cyclobutyl, methyl, ethyl, propyl, isopropyl, methoxy, ethoxy, propoxy, isopropoxy, hydroxymethyl, hydroxyethyl, hydroxypropyl, methylaminomethyl, methylaminoethyl, methylaminopropyl, dimethylaminomethyl, dimethylaminoethyl, dimethylaminopropyl, methoxymethyl, methoxyethyl, methoxypropyl, ethoxymethyl, ethoxyethyl, ethoxypropyl, methoxyethoxy, methoxypropoxy, ethoxyethoxy, ethoxypropoxy, methylaminoethoxy, methylaminopropoxy, ethylaminoethoxy, ethylaminopropoxy, dimethylaminoethoxy, dimethylaminopropoxy, diethylaminoethoxy, or diethylaminopropoxy.

[0034] In another alternative embodiment,



30 [0037] T₁ is linear C₁-C₆ alkyl, or linear C₁-C₆ alkyl independently substituted by R⁹ and R¹⁰, respectively;


[0038] R^9 and R^{10} are independently -H or methyl;

[0039] R^8 is -H, hydroxy, methyl, ethyl, propyl, isopropyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, methoxy, ethoxy, propoxy, isopropoxy, methylthio, ethylthio, propylthio, isopropylthio, 5- to 6-membered heterocyclyl or $-NR^{11}R^{12}$,

5 [0040] R^{11} and R^{12} are each independently -H, methyl, ethyl, propyl, butyl, pentyl, hexyl, isopropyl, sec-butyl, 1-ethylpropyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, hydroxyethyl, hydroxypropyl, hydroxybutyl, hydroxypentyl, hydroxyhexyl, methoxyethyl, methoxypropyl, methoxybutyl, methoxypentyl, methoxyhexyl, ethoxyethyl, ethoxypropyl, ethoxybutyl, ethoxypentyl, ethoxyhexyl, propoxyethyl, propoxypropyl, propoxybutyl, propoxypentyl, propoxyhexyl, isopropoxyethyl, isopropoxypropyl, isopropoxybutyl, isopropoxypentyl or isopropoxyhexyl;

[0041] the 5- to 6-membered heterocyclyl is a heterocyclyl containing 1-2 heteroatoms selected from N, O or S, the 5- to 6-membered heterocyclyl is unsubstituted or substituted by one or two of the group consisting of: methyl, ethyl, propyl, isopropyl, aldehyde group, formyl, acetyl, propionyl, butyryl, isobutyryl, aminoacyl, methylaminoacyl, dimethylaminoacyl, methylsulfonyl, ethylsulfonyl, isopropylsulfonyl, methylsulfinyl, ethylsulfinyl, and isopropylsulfinyl, or sulfur in the heterocycle is oxidized by one to two oxygen atoms;

[0042] the 5- to 6-membered heterocycle is selected from:



20 [0043] Alternatively,

[0044] R^2 is or ,

[0045] L_1 is selected from: or ,

[0046] T_1 is linear C_1 - C_6 alkyl, or linear C_1 - C_6 alkyl independently substituted by R^9 and R^{10} , respectively;

25 [0047] R^9 and R^{10} are each independently -H or methyl;


[0048] R^8 is -H, hydroxy, methyl, ethyl, propyl, isopropyl, methoxy, ethoxy, propoxy, isopropoxy, methylthio, ethylthio, propylthio, isopropylthio, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, 5- to 6-membered heterocyclyl or $-NR^{11}R^{12}$,

[0049] R^{11} and R^{12} are each independently -H, methyl, ethyl, propyl, butyl, pentyl, hexyl, isopropyl, sec-butyl, 1-ethylpropyl, cyclopropyl, cyclobutyl, cyclopentyl, hydroxyethyl, hydroxypropyl, hydroxybutyl, hydroxypentyl, methoxyethyl, methoxypropyl, methoxybutyl, methoxypentyl, ethoxyethyl, ethoxypropyl, ethoxybutyl, ethoxypentyl, propoxyethyl,

propoxypropyl, propoxybutyl, propoxypentyl, isopropoxyethyl, isopropoxypropyl, isopropoxybutyl or isopropoxypentyl;


[0050] the 5- to 6-membered heterocyclyl is a heterocyclyl containing 1-2 heteroatoms selected from N, O or S, the 5- to 6-membered heterocyclyl is unsubstituted or substituted by one or two of the group consisting of: methyl, ethyl, propyl, isopropyl, aldehyde group, formyl, acetyl, propionyl, butyryl, isobutyryl, aminoacyl, methylaminoacyl, dimethylaminoacyl, methylsulfonyl, ethylsulfonyl, isopropylsulfonyl, methylsulfinyl, ethylsulfinyl, and isopropylsulfinyl, or the sulfur in the heterocycle is oxidized by one to two oxygen atoms;

[0051] the 5- to 6-membered heterocyclyl is selected from:



[0052] R^{13} is -H, amino, methylamino, dimethylamino, methyl, ethyl, propyl, or isopropyl.

[0053] According to another aspect of the present disclosure, a compound of formula (I), or isomers, hydrates, solvates, pharmaceutically acceptable salts and prodrugs thereof are provided,




Formula (I)

[0054] in the formula (I),

[0055] X is O, or NH;

20 [0056] Y is N or C-Z, wherein Z is -H or -CN, alternatively, Y is N;



[0058] R^3 is -H, halogen, C_1 - C_3 alkyl, C_1 - C_3 alkoxy, C_1 - C_3 alkyl substituted by halogen or C_1 - C_3 alkoxy substituted by halogen; alternatively, R^3 is -H, fluorine, chlorine, bromine, methyl, ethyl, methoxy, ethoxy, trifluoromethyl or trifluoromethoxy;

25 [0059] L is or ,


[0060] M is O or S;

[0061] T is linear C_1 - C_3 alkyl, or linear C_1 - C_3 alkyl independently substituted by R^5 and R^6 ,


respectively; alternatively, T is linear C₁-C₂ alkyl, or linear C₁-C₂ alkyl independently substituted by R⁵, and R⁶, respectively;

[0062] R⁵ and R⁶ are each -H, halogen, C₁-C₃ alkyl, or C₁-C₃ alkyl substituted by halogen; alternatively, R⁵ and R⁶ are each -H, -F, methyl, ethyl or trifluoromethyl;

- 5 [0063] R⁴ is substituted or unsubstituted fused heteroaryl, wherein the substituted fused heteroaryl is substituted by 1-3 identical or different R⁷, the fused ring heteroaryl group is a heteroaryl group containing 1-3 heteroatoms selected from N, O or S; alternatively, R⁴ is



- [0064] R⁷ is -H, halogen, amino, hydroxy, cyano, C₁-C₃ alkylthio, amino substituted with
10 mono- or di-C₁-C₃ alkyl, C₃-C₄ cycloalkyl, unsubstituted or substituted C₁-C₆ alkyl, or
unsubstituted or substituted C₁-C₆ alkoxy, wherein the substituent of the substituted C₁-C₆ alkyl
is halogen, hydroxy, amino substituted with mono- or di-C₁-C₃ alkyl, or C₁-C₃ alkoxy, and
wherein the substituent of the substituted C₁-C₆ alkoxy is halogen, C₁-C₃ alkoxy, or amino
substituted with mono- or di-C₁-C₃ alkyl; alternatively, R⁷ is -H, -F, -Cl, -Br, -CF₃, -OCF₃,
15 amino, hydroxy, cyano, methylthio, ethylthio, propylthio, isopropylthio, methylamino,
ethylamino, dimethylamino, diethylamino, cyclopropyl, cyclobutyl, methyl, ethyl, propyl,
isopropyl, methoxy, ethoxy, propoxy, isopropoxy, hydroxymethyl, hydroxyethyl,
hydroxypropyl, methylaminomethyl, methylaminoethyl, methylaminopropyl,
dimethylaminomethyl, dimethylaminoethyl, dimethylaminopropyl, methoxymethyl,
20 methoxyethyl, methoxypropyl, ethoxymethyl, ethoxyethyl, ethoxypropyl, methoxyethoxy,
methoxypropoxy, ethoxyethoxy, ethoxypropoxy, methylaminoethoxy, methylaminopropoxy,
ethylaminoethoxy, ethylaminopropoxy, dimethylaminoethoxy, dimethylaminopropoxy,
diethylaminoethoxy, or diethylaminopropoxy;




- 25 [0066] R³ is -H, fluorine, chlorine, bromine, methyl, ethyl, methoxy, ethoxy, trifluoromethyl
or trifluoromethoxy;





[0068] M is O or S;



- [0069] T is linear C₁-C₂ alkyl, or linear C₁-C₂ alkyl independently substituted by R⁵ and R⁶,
30 respectively;

[0070] R⁵ and R⁶ are each independently -H, -F, methyl, ethyl or trifluoromethyl;

[0071] R^4 is unsubstituted or substituted , wherein the group is substituted by 1-3 identical or different R^7 ,

[0072] R^7 is -H, -F, -Cl, -Br, -CF₃, -OCF₃, amino, hydroxy, cyano, methylthio, ethylthio, propylthio, isopropylthio, methylamino, ethylamino, dimethylamino, diethylamino, cyclopropyl, cyclobutyl, methyl, ethyl, propyl, isopropyl, methoxy, ethoxy, propoxy, isopropoxy, hydroxymethyl, hydroxyethyl, hydroxypropyl, methylaminomethyl, methylaminoethyl, methylaminopropyl, dimethylaminomethyl, dimethylaminoethyl, dimethylaminopropyl, methoxymethyl, methoxyethyl, methoxypropyl, ethoxymethyl, ethoxyethyl, ethoxypropyl, methoxyethoxy, methoxypropoxy, ethoxyethoxy, ethoxypropoxy, methylaminoethoxy, methylaminopropoxy, ethylaminoethoxy, ethylaminopropoxy, dimethylaminoethoxy, dimethylaminopropoxy, diethylaminoethoxy, or diethylaminopropoxy.

[0073] R^2 is  or ,

[0074] L_1 is selected from:  or .



[0075] T_1 is linear C₁-C₈ alkyl, or linear C₁-C₈ alkyl independently substituted by R^9 and R^{10} , respectively;



[0076] R^9 and R^{10} are each independently -H, or C₁-C₃ alkyl;

[0077] R^8 is -H, hydroxy, C₁-C₃ alkyl, C₃-C₇ cycloalkyl, C₁-C₃ alkoxy, C₁-C₃ alkylthio, 4- to 7-membered heterocyclyl or -NR¹¹R¹²;

[0078] R^{11} and R^{12} are each independently -H, C₁-C₆ alkyl, C₃-C₆ cycloalkyl, C₁-C₆ alkyl substituted by hydroxy or C₁-C₆ alkyl substituted by C₁-C₃ alkoxy;

[0079] the 4- to 7-membered heterocyclyl is a heterocyclyl containing 1-2 heteroatoms selected from N, O or S, the heterocyclyl is unsubstituted or substituted by any one or two of the group consisting of: C₁-C₃ alkyl, aldehyde group, C₁-C₄ alkylacyl, aminoacyl, aminoacyl wherein the amino is substituted with mono- or di-C₁-C₃ alkyl, C₁-C₃ alkylsulfonyl, C₁-C₃ alkylsulfinyl, hydroxy, halogen, C₁-C₃ hydroxyalkyl, and C₁-C₃ haloalkyl, or the sulfur in the heterocycle is oxidized by one to two oxygen atoms.

[0080] alternatively, R^2 is  or ,

[0081] L_1 is selected from:  or .

[0082] T_1 is linear C₁-C₆ alkyl, or linear C₁-C₆ alkyl independently substituted by R^9 and R^{10} , respectively;


[0083] R^9 and R^{10} are each independently -H or methyl;



[0084] R^8 is -H, hydroxy, methyl, ethyl, propyl, isopropyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, methoxy, ethoxy, propoxy, isopropoxy, methylthio, ethylthio, propylthio, isopropylthio, 5- to 6-membered heterocyclyl or $-NR^{11}R^{12}$,



[0085] R^{11} and R^{12} are each independently -H, methyl, ethyl, propyl, butyl, pentyl, hexyl, isopropyl, sec-butyl, 1-ethylpropyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, hydroxyethyl, hydroxypropyl, hydroxybutyl, hydroxypentyl, hydroxyhexyl, methoxyethyl, methoxypropyl, methoxybutyl, methoxypentyl, methoxyhexyl, ethoxyethyl, ethoxypropyl, ethoxybutyl, ethoxypentyl, ethoxyhexyl, propoxyethyl, propoxypropyl, propoxybutyl, propoxypentyl, propoxyhexyl, isopropoxyethyl, isopropoxypropyl, isopropoxybutyl, isopropoxypentyl or isopropoxyhexyl;

[0086] the 5- to 6-membered heterocyclyl is a heterocyclyl containing 1-2 heteroatoms selected from N, O or S, the 5- to 6-membered heterocyclyl is unsubstituted or substituted by any one or two from the group consisting of: methyl, ethyl, propyl, isopropyl, aldehyde group, formyl, acetyl, propionyl, butyryl, isobutyryl, aminoacyl, methylaminoacyl, dimethylaminoacyl, methylsulfonyl, ethylsulfonyl, isopropylsulfonyl, methylsulfinyl, ethylsulfinyl, isopropylsulfinyl, hydroxy, fluorine, chlorine, hydroxymethyl, hydroxyethyl, hydroxypropyl, and trifluoromethyl, or the sulfur in the heterocycle is oxidized by one to two oxygen atoms;

[0087] the 5- to 6-membered heterocycle is selected from:



[0088] Alternatively, R^2 is  or .

[0089] L_1 is selected from:  or .

[0090] T_1 is linear C_1 - C_6 alkyl, or linear C_1 - C_6 alkyl independently substituted by R^9 and R^{10} , respectively;

25 [0091] R^9 and R^{10} are each independently -H or methyl;


[0092] R^8 is -H, hydroxy, methyl, ethyl, propyl, isopropyl, methoxy, ethoxy, propoxy, isopropoxy, methylthio, ethylthio, propylthio, isopropylthio, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, 5- to 6-membered heterocyclyl or $-NR^{11}R^{12}$,

[0093] R^{11} and R^{12} are each independently -H, methyl, ethyl, propyl, butyl, pentyl, hexyl, isopropyl, sec-butyl, 1-ethylpropyl, cyclopropyl, cyclobutyl, cyclopentyl, hydroxyethyl, hydroxypropyl, hydroxybutyl, hydroxypentyl, methoxyethyl, methoxypropyl, methoxybutyl, methoxypentyl, ethoxyethyl, ethoxypropyl, ethoxybutyl, ethoxypentyl, propoxyethyl,

propoxypropyl, propoxybutyl, propoxypentyl, isopropoxyethyl, isopropoxypropyl, isopropoxybutyl or isopropoxypentyl;


[0094] the 5- to 6-membered heterocyclyl is a heterocyclyl containing 1-2 heteroatoms selected from N, O or S, wherein the 5- to 6-membered heterocyclyl is unsubstituted or substituted by any one or two of the group consisting of: methyl, ethyl, propyl, isopropyl, aldehyde group, formyl, acetyl, propionyl, butyryl, isobutyryl, aminoacyl, methylaminoacyl, dimethylaminoacyl, methylsulfonyl, ethylsulfonyl, isopropylsulfonyl, methylsulfinyl, ethylsulfinyl, isopropylsulfinyl, hydroxy, fluorine, chlorine, hydroxymethyl, hydroxyethyl, and trifluoromethyl, or the sulfur in the heterocycle is oxidized by one to two oxygen atoms;

10 [0095] the 5- to 6-membered heterocyclyl is selected from:



[0096] R^{13} is -H, amino, methylamino, dimethylamino, methyl, ethyl, propyl or isopropyl.

[0097] According to yet another aspect of the present invention, a compound of formula (I),
15 or isomers, hydrates, solvates, pharmaceutically acceptable salts and prodrugs thereof are provided.




Formula (I)

[0098] in the formula (I),

20 [0099] X is O, or NH;

[0100] Y is N or C-Z, wherein Z is -H or -CN;



[0102] R^3 is -H, halogen, C_1 - C_3 alkyl, C_1 - C_3 alkoxy, C_1 - C_3 alkyl substituted by halogen or C_1 - C_3 alkoxy substituted by halogen;

25 [0103] L is

[0104] M is O or S;



[0105] T is linear C_1 - C_3 alkyl, or linear C_1 - C_3 alkyl independently substituted by R^5 and R^6 ,



respectively;

[00106] R^5 and R^6 are each independently -H, halogen, C₁-C₃ alkyl, or C₁-C₃ alkyl substituted by halogen;

[00107] R^4 is aryl, 5- to 6-membered heteroaryl, aryl substituted by 1-3 identical or different R^7 , or 5- to 6-membered heteroaryl substituted by 1-3 identical or different R^7 , wherein the heteroaryl group is a heteroaryl group containing 1-3 heteroatoms selected from N, O or S;

[00108] R^7 is -H, halogen, amino, hydroxy, cyano, C₁-C₃ alkylthio, amino substituted with mono- or di-C₁-C₃ alkyl, C₃-C₄ cycloalkyl, unsubstituted or substituted C₁-C₆ alkyl, or unsubstituted or substituted C₁-C₆ alkoxy, wherein the substituent of the substituted C₁-C₆ alkyl is halogen, hydroxy, amino substituted with mono- or di-C₁-C₃ alkyl, or C₁-C₃ alkoxy, and wherein the substituent of the substituted C₁-C₆ alkoxy is halogen, C₁-C₃ alkoxy, or amino substituted with mono- or di-C₁-C₃ alkyl;

[00109] R^2 is  or .

[00110] L_1 is selected from:  or .

[00111] T_1 is linear C₁-C₈ alkyl, or linear C₁-C₈ alkyl independently substituted by R^9 and R^{10} , respectively;


[00112] R^9 and R^{10} are each independently -H, or C₁-C₃ alkyl;

[00113] R^8 is -H, hydroxy, C₁-C₃ alkyl, C₃-C₇ cycloalkyl, C₁-C₃ alkoxy, C₁-C₃ alkylthio, 4- to 7-membered heterocyclyl or -NR¹¹R¹²;



[00114] R^{11} and R^{12} are each independently -H, C₁-C₆ alkyl, C₃-C₆ cycloalkyl, C₁-C₆ alkyl substituted by hydroxyl or C₁-C₆ alkyl substituted by C₁-C₃ alkoxy;

[00115] the 4- to 7-membered heterocyclyl is a heterocyclyl containing 1-2 heteroatoms selected from N, O or S, the heterocyclyl is unsubstituted or substituted by any one or two of the group consisting of: C₁-C₃ alkyl, hydroxy, halogen, C₁-C₃ hydroxyalkyl, and C₁-C₃ haloalkyl.

[00116] In one alternative embodiment, X is O, or NH, Y is N;

[00117] R^1 is .

[00118] R^3 is -H, fluorine, chlorine, bromine, methyl, ethyl, methoxy, ethoxy, trifluoromethyl or trifluoromethoxy;

[00119] L is  or .

[00120] M is O or S;



[00121] T is linear C₁-C₂ alkyl, or linear C₁-C₂ alkyl independently substituted by R⁵ and R⁶, respectively;



[00122] R⁵ and R⁶ are each independently -H, -F, methyl, ethyl or trifluoromethyl;

[00123] R⁴ is aryl, 5- to 6-membered heteroaryl, aryl substituted by 1-2 identical or different R⁷, or 5- to 6-membered heteroaryl substituted by 1-2 identical or different R⁷, wherein the aryl or heteroaryl group is selected from the group consisting of: phenyl, pyridyl, pyrimidinyl, thiazolyl, thienyl, pyrrolyl, thiadiazolyl, furyl, oxazolyl or isoxazolyl;

[00124] R⁷ is -H, -F, -Cl, -Br, -CF₃, -OCF₃, amino, hydroxy, cyano, methylthio, ethylthio, propylthio, isopropylthio, methylamino, ethylamino, dimethylamino, diethylamino, cyclopropyl, cyclobutyl, methyl, ethyl, propyl, isopropyl, methoxy, ethoxy, propoxy, isopropoxy, hydroxymethyl, hydroxyethyl, hydroxypropyl, methylaminomethyl, methylaminoethyl, methylaminopropyl, dimethylaminomethyl, dimethylaminoethyl, dimethylaminopropyl, methoxymethyl, methoxyethyl, methoxypropyl, ethoxymethyl, ethoxyethyl, ethoxypropyl, methoxyethoxy, methoxypropoxy, ethoxyethoxy, ethoxypropoxy, methylaminoethoxy, methylaminopropoxy, ethylaminoethoxy, ethylaminopropoxy, dimethylaminoethoxy, dimethylaminopropoxy, diethylaminoethoxy, or diethylaminopropoxy.

[00125] In another alternative embodiment,

[00126] R² is  or ,

[00127] L₁ is selected from:  or .

[00128] T₁ is linear C₁-C₆ alkyl, or linear C₁-C₆ alkyl independently substituted by R⁹ and R¹⁰, respectively;

[00129] R⁹ and R¹⁰ are each independently -H or methyl;


[00130] R⁸ is -H, hydroxy, methyl, ethyl, propyl, isopropyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, methoxy, ethoxy, propoxy, isopropoxy, methylthio, ethylthio, propylthio, isopropylthio, 5- to 6-membered heterocyclyl or -NR¹¹R¹²,

[00131] R¹¹ and R¹² are each independently -H, methyl, ethyl, propyl, butyl, pentyl, hexyl, isopropyl, sec-butyl, 1-ethylpropyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, hydroxyethyl, hydroxypropyl, hydroxybutyl, hydroxypentyl, hydroxyhexyl, methoxyethyl, methoxypropyl, methoxybutyl, methoxypentyl, methoxyhexyl, ethoxyethyl, ethoxypropyl, ethoxybutyl, ethoxypentyl, ethoxyhexyl, propoxyethyl, propoxypropyl, propoxybutyl, propoxypentyl, propoxyhexyl, isopropoxyethyl, isopropoxypropyl, isopropoxybutyl, isopropoxypentyl or isopropoxyhexyl;



[00132] the 5- to 6-membered heterocyclyl is a heterocyclyl containing 1-2 heteroatoms



selected from N, O or S, the 5- to 6-membered heterocyclyl is unsubstituted or substituted by any one or two of the group consisting of: methyl, ethyl, propyl, isopropyl, hydroxy, fluorine, chlorine, hydroxymethyl, hydroxyethyl, hydroxypropyl, or trifluoromethyl;

[00133] the 5- to 6-membered heterocycle is selected from:



[00134] Alternatively,

[00135] R^2 is  or ,

[00136] L_1 is selected from:  or .

[00137] T_1 is linear C_1 - C_6 alkyl, or linear C_1 - C_6 alkyl independently substituted by R^9 and R^{10} , respectively;


[00138] R^9 and R^{10} are each independently -H or methyl;

[00139] R^8 is -H, hydroxy, methyl, ethyl, propyl, isopropyl, methoxy, ethoxy, propoxy, isopropoxy, methylthio, ethylthio, propylthio, isopropylthio, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, 5- to 6-membered heterocyclyl or $-NR^{11}R^{12}$,

15 [00140] R^{11} and R^{12} are each independently is -H, methyl, ethyl, propyl, butyl, pentyl, hexyl, isopropyl, sec-butyl, 1-ethylpropyl, cyclopropyl, cyclobutyl, cyclopentyl, hydroxyethyl, hydroxypropyl, hydroxybutyl, hydroxypentyl, methoxyethyl, methoxypropyl, methoxybutyl, methoxypentyl, ethoxyethyl, ethoxypropyl, ethoxybutyl, ethoxypentyl, propoxyethyl, propoxypropyl, propoxybutyl, propoxypentyl, isopropoxyethyl, isopropoxypropyl, 20 isopropoxybutyl or isopropoxypentyl;

[00141] the 5- to 6-membered heterocyclyl is a heterocyclyl containing 1-2 heteroatoms selected from N, O or S, the 5- to 6-membered heterocyclyl is unsubstituted or substituted by one or two of the group consisting of: methyl, ethyl, propyl, isopropyl, hydroxy, fluorine, chlorine, hydroxymethyl, hydroxyethyl, and trifluoromethyl;

25 [00142] the 5- to 6-membered heterocyclyl is selected from:




[00143] R^{13} is -H, amino, methylamino, dimethylamino, methyl, ethyl, propyl, or isopropyl.

[00144] The present disclosure also relates to a method for treating diseases or disorders


mediated by kinases such as EGFR, HER2, HER3 and HER4, which includes administering a therapeutically effective amount of compounds of formula (I) or salts thereof to a patient in need (human or other mammals, especially human), and the diseases or disorders mediated by EGFR, HER2, HER3 and HER4 kinase include those mentioned above.

- 5 **[00145]** The present disclosure provides a method for preparing the above compound or a pharmaceutically acceptable salt, isomer, hydrate, solvate, or prodrug thereof, which comprises the following steps:


[00146] the compound of formula (I) is prepared by the reaction of $R^2C(O)Cl$ with the compound of formula (VIII), or by the chlorination reaction of R^2COOH followed by the
10 reaction with the compound of formula (VIII);



[00147] or, when R^2 is $\text{---}L_1\text{---}T_1\text{---}R^8$, wherein L_1 is $\text{---}CH=CH\text{---}$, and R^8 is $-NR^{11}R^{12}$, the compound of formula (I) is prepared by the reaction of compound of formula (VIII) with





followed by the reaction with $HNR^{11}R^{12}$,



15


[00148] or, when R^2 is $\text{---}L_1\text{---}R^8$ or $\text{---}L_1\text{---}T_1\text{---}R^8$, wherein L_1 is $\text{---}CH=CH\text{---}$, the compound of formula (I) is prepared by the reaction of compound of formula (IX) with



20 **[00149]** $R^1, R^2, R^8, R^9, R^{10}, R^{11}, R^{12}, X, Y, L_1$ and T_1 are as defined above.


[00150] The present disclosure also provides a compound represented by formula (VIII),

wherein R¹ and Y are as defined above,




[00151] The present disclosure also provides a method for preparing the compound represented by formula (VIII), R¹ is as defined above, and Y is N. The preparation method includes the following steps:

[00152] Method A:




[00153] Compound of formula (VIII) is prepared by fully contacting 2,3,4,9-tetrahydro-10H-[1,4]oxazino[2,3-f]quinazolin-10-one as represented by formula (V) with R¹XH, followed by

[00154] or method B:



[00155] 2,3,4,9-tetrahydro-10H-[1,4]oxazino[2,3-f]quinazolin-10-one as represented by formula (V) is fully contacted with Castro's reagent to afford the compound of formula (VI), and compound (VI) is further reacted with R¹XH to afford the compound represented by formula (VIII);

[00156] or method C:



[00157] The compound represented by the formula (II) is reacted with a chlorinating reagent

and then reacted with R^1XH to afford the compound represented by the formula (VII), and the compound represented by the formula (VII) is further subjected to a reduction and cyclization reaction to afford the compound represented by the formula (VIII).

5 DETAILED DESCRIPTION

[00158] The term "substituted" as used herein, includes multiple substituents (e.g., phenyl, aryl, heteroalkyl, heteroaryl), preferably 1 to 5 substituents, more preferably 1 to 3 substituents, most preferably 1 or 2 substituents, independently selected from the list of substituents.

[00159] Unless otherwise specified, alkyl includes saturated linear and branched hydrocarbon group, C_1 - C_8 represents the number of carbon atoms of an alkyl is 1-8. Similarly, for example, C_1 - C_3 represents the number of carbon atoms of an alkyl is 1-3, e.g., C_1 - C_6 alkyl includes methyl, ethyl, propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, n-pentyl, 3-(2-methyl)butyl, 2-pentyl, 2-methylbutyl, neopentyl, n-hexyl, 2-hexyl, and 2-methylpentyl. Alkoxy is an ether consisting of a linear or branched alkyl as previously described. Similarly, alkenyl and alkynyl groups include linear or branched alkenyl or alkynyl groups.

[00160] Cycloalkyl refers to a cyclic group formed by carbon atoms. For example, C_3 - C_7 represents a cycloalkyl group having 3 to 7 carbon atoms, including cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, and cycloheptyl. Similarly, cyclic alkenyl group is also included herein.

[00161] The term "aryl" as used herein, unless otherwise specified, refers to an unsubstituted or substituted aromatic group, such as phenyl, naphthyl, anthracenyl. The term "arylacyl" refers to $-C(O)$ -aryl.

[00162] "Oxidized by one or two oxygen atoms" refers to a sulfur atom oxidized by one oxygen atom to form a double bond between the sulfur and oxygen, or oxidized by two oxygen atoms to form double bonds between the sulfur and two oxygen atoms.

[00163] The term "heterocyclyl" as used herein, unless otherwise specified, represents an unsubstituted or substituted stable 3 to 8 membered monocyclic saturated ring system consisting of carbon atoms and 1 to 3 heteroatoms selected from N, O, and S, wherein the N, S heteroatoms can be optionally oxidized, and the N heteroatoms can also be optionally quaternized. The heterocycle can be attached through any heteroatom or carbon atom to form a stable structure. Examples of such heterocyclyl rings include, but are not limited to, azetidiny, pyrrolidiny, tetrahydrofurany, tetrahydrothiazoly, tetrahydropyrany, morpholinyl, thiomorpholinyl, piperidiny, piperazinyl, oxopiperazinyl, oxopiperidiny, dioxolany, dioxany, tetrahydroimidazoly, tetrahydrooxazoly, thiomorpholine oxide, thiomorpholine dioxide and

oxadiazolyl.

[00164] The term "heteroaryl" as used herein, unless otherwise specified, represents an unsubstituted or substituted stable 5 or 6 membered monocyclic aromatic ring system, and may also represent unsubstituted or substituted 9 or 10-membered benzo-fused heteroaromatic ring system or a bicyclic heteroaromatic ring system consisting of carbon atoms and one to three heteroatoms selected from N, O, S, wherein the N, S heteroatoms may optionally be oxidized, and N heteroatoms may optionally be quaternized. Heteroaryl can be attached at any heteroatom or carbon atom to form a stable structure. Heteroaryl includes but is not limited to thienyl, furyl, imidazolyl, pyrrolyl, thiazolyl, oxazolyl, isoxazolyl, pyranyl, pyridinyl, piperazinyl, pyrimidinyl, pyrazinyl, pyridazinyl, pyrazolyl, thiadiazolyl, triazolyl, indolyl, azaindolyl, indazolyl, azaindazolyl, benzimidazolyl, benzofuryl, benzothienyl, benzoisoxazolyl, benzoxazolyl, benzopyrazolyl, benzothiazolyl, benzothiadiazolyl, benzotriazolyl, adeninyl, quinolinyl, or isoquinolinyl.


[00165] The term "carbonyl" refers to a C(O) group.

[00166] Whenever the terms "alkyl" or "aryl" or either of their prefix roots appear in the name of a substituent (eg, aralkyl, dialkylamino), it shall be interpreted to contain those limitations given for the above "alkyl" and "aryl". Designated numbers of carbon atoms (e.g., C₁-C₆) shall independently represent the number of carbon atoms in an alkyl moiety or an alkyl moiety in a larger substituent (wherein the alkyl group is the prefix root).

[00167] The present disclosure also provides methods for preparing the corresponding compounds, wherein the compounds disclosed herein can be prepared using a variety of synthetic methods, including the methods described below. The compounds of the present disclosure, or pharmaceutically acceptable salts, isomers or hydrates thereof could be synthesized using the following methods and the known synthetic methods in the art of organic synthesis, or by variations of those methods as understood by those skilled in the art. The preferred methods include, but are not limited to, the methods described below.

[00168] The synthetic route of the compound of formula (I) is illustrated by the formula (I) wherein Y is N. The present disclosure are mainly illustrated by the following three preparation schemes:

[00169] **Preparation route I of the compound represented by formula (I):** wherein, R¹, R², R⁸, R⁹, R¹⁰, R¹¹, R¹², X, L₁ and T₁ are as defined above.



[00170] Reaction step a): 5-chloro-6-nitroquinazolin-4(3H)-one is fully contacted with ethylene glycol and sodium hydride to afford the compound represented by formula (II) of 5-(2-hydroxyethoxy)-6-nitroquinazolin-4(3H)-one.

[00171] Reaction step b): 5-(2-hydroxyethoxy)-6-nitroquinazolin-4(3H)-one represented by formula (II) is fully contacted with a chlorinating reagent, followed by adding water to afford 5-(2-chloroethoxy)-6-nitroquinazolin-4(3H)-one represented by formula (III). The chlorinating reagent includes, but is not limited to, any one or the combination of two or more of phosphorus oxychloride, sulfoxide chloride, phosphorus trichloride, phosphorus pentachloride and chlorine gas.

[00172] Reaction step c): 5-(2-chloroethoxy)-6-nitroquinazolin-4(3H)-one represented by formula (III) is subjected to a reduction reaction to afford 5-(2-chloroethoxy)-6-aminoquinazolin-4(3H)-one represented by formula (IV). The conditions of the reduction reaction include, but are not limited to, hydrogen and raney nickel, hydrogen and palladium on carbon, iron powder, zinc powder and stannous chloride.

[00173] Reaction step d): 5-(2-chloroethoxy)-6-aminoquinazolin-4(3H)-one represented by formula (IV) is dissolved in a solvent, and heated to afford 2,3,4,9-tetrahydro-10H-

[1,4]oxazino[2,3-f]quinazolin-10-one represented by formula (V); the solvent is preferably selected from any one or the combination of two or more of the following: methanol, ethanol, isopropanol, tetrahydrofuran, N,N-dimethylformamide (DMF), N,N-dimethylacetamide (DMA), N-methylpyrrolidone (NMP), dioxane, and dichloroethane;

5 [00174] alternatively, the reaction can be carried out under base-catalyzed conditions, the base includes, but is not limited to, any one or the combination of two or more of the following: triethylamine, diisopropylethylamine, pyridine, 4-dimethylaminopyridine, 1,8-diazabicycloundec-7-ene, N-methylmorpholine, sodium carbonate, potassium carbonate and cesium carbonate.

10 [00175] Reaction step e): 2,3,4,9-tetrahydro-10H-[1,4]oxazino[2,3-f]quinazolin-10-one represented by formula (V) is fully contacted with R^1XH and Castros reagent to afford an oxazino-quinazoline compound represented by formula (VII).


[00176] Alternatively, Castros reagent is selected from any one or the combination of two of the following: (benzotriazol-1-yloxy)tris(dimethylamino)phosphonium hexafluorophosphate (BOP) or (benzotriazol-1-yloxy)tripyrrolidinophosphonium hexafluorophosphate (PyBOP); the reaction can be carried out under base-catalyzed conditions, the base includes, but is not limited to, any one or the combination of two or more of the following: triethylamine, diisopropylethylamine, triethylenediamine (DABCO), 1,8-diazabicycloundec-7-ene (DBU), pyridine, N-methylmorpholine, 4-dimethylaminopyridine, sodium carbonate, potassium carbonate and cesium carbonate.

20 [00177] Reaction step f-A): the compound represented by formula (VIII) is condensed with $R^2C(O)Cl$, or the product of the reaction of R^2COOH and a chlorinating reagent, to afford the compound represented by formula (I);




[00178] the chlorinating agent is preferably selected from any one or the combination of two or more of the following: phosphorus oxychloride, thionyl chloride, oxalyl chloride, phosphorus trichloride and phosphorus pentachloride;

[00179] alternatively, the reaction can be carried out under base-catalyzed conditions, the base includes, but is not limited to, any one or the combination of two or more of the following: triethylamine, diisopropylethylamine, pyridine, 4-dimethylaminopyridine, 1,8-diazabicycloundec-7-ene, N-methylmorpholine, sodium carbonate, potassium carbonate and cesium carbonate.

30 [00180] Reaction step f-B): or, when R^2 is , and R^8 is $HNR^{11}R^{12}$, and L_1 is



followed by adding amine having R^{11} and R^{12} ($R^{11}-\overset{\text{H}}{\text{N}}-R^{12}$) to afford the compound represented by formula (I); alternatively, the above reaction can be carried out in an organic solvent, the organic solvent includes, but is not limited to, any one or the combination of two or more of the following: tetrahydrofuran (THF), N,N-dimethylformamide (DMF), N,N-dimethylacetamide (DMA), N-methylpyrrolidone (NMP), dioxane, and dichloroethane;

[00181] Reaction step f-C): or, when R^2 is  or , and L_1 is , the compound of formula (IX) is obtained from the reaction of step f-C1 by the reaction of the compound of formula (VIII) and 2-(diethoxyphosphoryl)acetic acid under the action of a condensing agent, and the compound of formula (IX) is further reacted with

10 $\text{OHC}-R^8$ or $\text{OHC}-T_1-R^8$ in the reaction step f-C2 to afford compound of formula (I).

[00182] Alternatively, the condensing agent includes, but is not limited to, any one or two or more of the following: carbodiimide type condensing agent, onium salt-based condensing agent, organophosphorus condensing agent and other types of condensing agent, alternatively, any one or the combination of two or more of the following: N,N'-carbonyldiimidazole (CDI), N,N'-dicyclohexylcarbodiimide (DCC), N,N'-diisopropylcarbodiimide (DIC), 1-hydroxybenzotriazole (HOBt), N,N'-diisopropylethylamine (DIEA), 1-hydroxy-7-azabenzotriazole (HOAt), *O*-(benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium tetrafluoroborate (TBTU), (benzotriazol-1-yloxy)tris(dimethylamino)phosphonium hexafluorophosphate (BOP), *O*-(benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HBTU), *O*-(6-chlorobenzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (HCTU), *O*-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HATU), propylphosphonic anhydride (T3P), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDCI), 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (EDC), (benzotriazol-1-yloxy)tripyrrolidinophosphonium hexafluorophosphate (PyBOP), and (3H-1,2,3-triazolo[4,5-b]pyridin-3-yloxy)tripyrrolidinophosphonium hexafluorophosphate (PyAOP);


[00183] alternatively, this step can be carried out in an organic base, the organic base includes, but is not limited to, any one or the combination of two or more of the following: triethylamine, diisopropylethylamine (DIEA), pyridine, 4-dimethylaminopyridine (DMAP), 2,6-dimethylpyridine (Lutidine), 1,8-diazabicycloundec-7-ene (DBU) or N-methylmorpholine.

[00184] Alternatively, step f-C2 can be carried out in an aprotic solvent under the action of an base. The aprotic solvent includes, but is not limited to, any one or the combination of two or

more of the following: tetrahydrofuran (THF), N,N-dimethylformamide (DMF), N,N-dimethylacetamide (DMA), N-methylpyrrolidone (NMP), and dioxane; the base includes, but is not limited to, one of sodium hydride, and lithium bistrimethylsilylamide, or both.

[00185] The above steps f-A), f-B) and f-C) are parallel steps, that is, the compound represented by formula (I) can be prepared by the compound represented by formula (VIII) through one of steps f-A), f-B) and f-C), that is, the compound represented by formula (I) can be prepared by the compound represented by formula (VIII) through step f-A), or the compound represented by formula (I) can be prepared by the compound represented by formula (VIII) through step f-B), or the compound represented by formula (I) can be prepared by the compound represented by formula (VIII) through step f-C).

[00186] **Preparation route II of the compound represented by formula (I)**, wherein, R^1 , R^2 , R^8 , R^9 , R^{10} , R^{11} , R^{12} , X, L_1 and T_1 are as defined above.



[00187] Reaction step a): 2,3,4,9-tetrahydro-10H-[1,4]oxazino[2,3-f]quinazolin-10-one represented by formula (V) is fully contacted with Castros reagent to afford the compound represented by formula (VI);

[00188] alternatively, Castros reagent is selected from any one of (benzotriazol-1-yloxy)tris(dimethylamino)phosphonium hexafluorophosphate (BOP) or (benzotriazol-1-yloxy)tripyrrolidinophosphonium hexafluorophosphate (PyBOP), or both;

[00189] alternatively, the above reaction can be carried out under basic conditions, the base includes, but is not limited to, any one or the combination of two or more of the following: triethylamine, diisopropylethylamine, triethylenediamine, 1,8-diazabicycloundec-7-ene (DBU), pyridine, N-methylmorpholine, 4-dimethylaminopyridine, sodium carbonate, potassium carbonate and cesium carbonate.

[00190] Reaction step b): 10-((1H-benzo[d][1,2,3]triazol-1-yl)oxy)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazoline represented by formula (VI) and R^1XH are fully contacted in an organic solvent to afford the compound represented by formula (VIII);

[00191] alternatively, the organic solvent is selected from any one or the combination of two or more of the following: methanol, ethanol, isopropanol, tetrahydrofuran, N,N-dimethylformamide (DMF), N,N-dimethylacetamide (DMA), N-methylpyrrolidone (NMP), dioxane, and dichloroethane;


[00192] alternatively, the reaction can be carried out under base-catalyzed conditions, the base includes, but is not limited to, any one or the combination of two or more of the following: triethylamine, diisopropylethylamine, pyridine, 4-dimethylaminopyridine, 1,8-diazabicycloundec-7-ene, N-methylmorpholine, sodium carbonate, potassium carbonate, cesium carbonate, and sodium hydride;



[00193] alternatively, the reaction can be carried out under acid-catalyzed conditions, the acid includes, but is not limited to, any one or the combination of two or more of the following: methanesulfonic acid, p-toluenesulfonic acid, benzenesulfonic acid, trifluoroacetic acid, and hydrochloric acid.

[00194] Reaction step c-A): the compound represented by formula (VIII) is subjected to a condensation reaction with $R^2C(O)Cl$ or the product of R^2COOH and chlorinating agent to afford the compound represented by formula (I);

[00195] the chlorinating agent is selected from any one or the combination of two or more of phosphorus oxychloride, thionyl chloride, oxalyl chloride, phosphorus trichloride and phosphorus pentachloride;




[00196] alternatively, the above reaction can be carried out under basic conditions, and the base includes, but is not limited to, any one or the combination of two or more of the following: triethylamine, diisopropylethylamine, pyridine, 4-dimethylaminopyridine, 1,8-diazabicycloundec-7-ene, N-methylmorpholine, sodium carbonate, potassium carbonate, and cesium carbonate.

[00197] Reaction step c-B): or, when R^2 is , and R^8 is $HNR^{11}R^{12}$, and L_1 is

, the compound represented by formula (VIII) is reacted with 

followed by adding amine having R^{11} and R^{12} ($R^{11}-\text{N}(H)-R^{12}$) to afford the compound represented by formula (I).

[00198] Alternatively, the above reaction can be carried out in an organic solvent, which includes, but is not limited to, any one or the combination of two or more of the following: tetrahydrofuran (THF), N,N-dimethylformamide (DMF), N,N-dimethylacetamide (DMA), N-methylpyrrolidone (NMP), dioxane and dichloroethane;

[00199] Reaction step c-C): or, when R^2 is  or , and L_1 is , the compound of formula (IX) is prepared by the compound of formula (VIII) and 2-(diethoxyphosphoryl)acetic acid through the reaction in step c-C1 under the action of a

condensing agent, the compound represented by formula (IX) is further reacted with $\text{OHC}-R^8$ or $\text{OHC}-T_1-R^8$ through the reaction in step c-C2 to afford the compound represented by formula (I).

[00200] Alternatively, the condensing agent includes, but is not limited to, any one or more of the following: carbodiimide type condensing agent, onium salt-based condensing agent, organophosphorus condensing agent and other types of condensing agent, alternatively, one or the combination of two or more of the following: N,N'-carbonyldiimidazole (CDI), N,N'-dicyclohexylcarbodiimide (DCC), N,N'-diisopropylcarbodiimide (DIC), 1-hydroxybenzotriazole (HOBt), N,N-diisopropylethylamine (DIEA), 1-hydroxy-7-azabenzotriazole (HOAt), *O*-(benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium tetrafluoroborate (TBTU), (benzotriazol-1-yloxy)tris(dimethylamino)phosphonium hexafluorophosphate (BOP), *O*-(benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HBTU), *O*-(6-chlorobenzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (HCTU), *O*-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HATU), propylphosphonic anhydride (T3P), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDCI), 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (EDC), (benzotriazol-1-yloxy)tripyrrolidinophosphonium hexafluorophosphate (PyBOP), and (3H-1,2,3-triazolo[4,5-b]pyridin-3-yloxy)tripyrrolidinophosphonium hexafluorophosphate (PyAOP).


[00201] Alternatively, this step can be carried out in an organic base, the organic base includes,

but is not limited to, any one or the combination of two or more of the following: triethylamine, diisopropylethylamine (DIEA), pyridine, 4-dimethylaminopyridine (DMAP), 2,6-dimethylpyridine (Lutidine), 1,8-diazabicycloundec-7-ene (DBU) and N-methylmorpholine.

[00202] Alternatively, step c-C2 can be carried out in an aprotic solvent under the action of a base. The aprotic solvent includes, but is not limited to, any one or the combination of two or more of the following: tetrahydrofuran (THF), N,N-dimethylformamide (DMF), N,N-dimethylacetamide (DMA), N-methylpyrrolidone (NMP), and dioxane; the alkali includes, but is not limited to, any one of sodium hydride, and lithium bistrimethylsilylamide, or both.

[00203] The above steps c-A), c-B) and c-C) are parallel steps, that is, the compound represented by formula (I) can be prepared by the compound represented by formula (VIII) through one of steps c-A), c-B) and c-C), that is, the compound represented by formula (I) can be prepared by the compound represented by formula (VIII) through step c-A), or the compound represented by formula (I) can be prepared by the compound represented by formula (VIII) through step c-B), or the compound represented by formula (I) can be prepared by the compound represented by formula (VIII) through step c-C).

[00204] Preparation route III of the compound represented by formula (I), wherein, R^1 , R^2 , R^8 , R^9 , R^{10} , R^{11} , R^{12} , X, L_1 and T_1 are as defined above.



[00205] Reaction step a): the compound represented by formula (II), 5-(2-hydroxyethoxy)-6-

nitroquinazolin-4(3H)-one, is reacted with a chlorinating reagent followed by contacting with R^1XH to afford quinazoline compounds represented by formula (VII);


[00206] alternatively, the chlorinating reagent include, but are not limited to, any one or the combination of two or more of the following: phosphorus oxychloride, thionyl chloride, phosphorus trichloride, phosphorus pentachloride and chlorine gas;



[00207] Reaction step b): the quinazoline compounds represented by formula (VII) are subjected to reducing conditions to afford oxazino-quinazoline compounds represented by formula (VIII),

[00208] the reduction conditions include, but are not limited to, hydrogen and raney nickel, hydrogen and palladium on carbon, iron powder, zinc powder, and stannous chloride.

[00209] Reaction step c-A): the compound represented by formula (VIII) is subjected to a condensation reaction with $R^2C(O)Cl$ or the product of R^2COOH and a chlorinating reagent to obtain a compound represented by formula (I);




[00210] the chlorinating agent is selected from any one or a combination of two or more of the following: phosphorus oxychloride, thionyl chloride, oxalyl chloride, phosphorus trichloride, and phosphorus pentachloride; alternatively, the above reaction can be carried out under basic conditions, the base includes, but is not limited to, triethylamine, diisopropylethylamine, pyridine, 4-dimethylaminopyridine, 1,8-diazabicycloundec-7-ene, N-methylmorpholine, sodium carbonate, potassium carbonate and cesium carbonate

[00211] Reaction step c-B): or, when R^2 is , and R^8 is $HNR^{11}R^{12}$, and L_1 is

, the compound represented by formula (VIII) is reacted with 

followed by adding amine having R^{11} and R^{12} ($R^{11}-N(R^{12})$) to afford the compound represented by formula (I).

[00212] Alternatively, the above reaction can be carried out in an organic solvent, the organic solvent includes, but is not limited to, any one or a combination of two or more of the following: tetrahydrofuran (THF), N,N-dimethylformamide (DMF), N,N-dimethylacetamide (DMA), N-methylpyrrolidone (NMP), dioxane, and dichloroethane;

[00213] Reaction step c-C): or, when R^2 is  or , and L_1 is , the compound of formula (IX) is prepared by the compound of formula (VIII) and

2-(diethoxyphosphoryl)acetic acid through the reaction in step c-C1 under the action of a

condensing agent, the compound represented by formula (IX) is further reacted with $\text{OHC}-\text{R}^8$ or $\text{OHC}-\text{T}_1-\text{R}^8$ through reaction of step c-C2 to afford the compound represented by formula (I);

[00214] alternatively, the condensing agent includes, but is not limited to, any one or more of the following: carbodiimide type condensing agent, onium salt-based condensing agent, organophosphorus condensing agent and other types of condensing agent, alternatively, one or the combination of two or more of the following: N,N'-carbonyldiimidazole (CDI), N,N'-dicyclohexylcarbodiimide (DCC), N,N'-diisopropylcarbodiimide (DIC), 1-hydroxybenzotriazole (HOBt), N,N-diisopropylethylamine (DIEA), 1-hydroxy-7-azabenzotriazole (HOAt), *O*-(benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium tetrafluoroborate (TBTU), (benzotriazol-1-yloxy)tris(dimethylamino)phosphonium hexafluorophosphate (BOP), *O*-(benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HBTU), *O*-(6-chlorobenzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (HCTU), *O*-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HATU), propylphosphonic anhydride (T3P), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDCI), 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (EDC), (benzotriazol-1-yloxy)tripyrrolidinophosphonium hexafluorophosphate (PyBOP), and (3H-1,2,3-triazolo[4,5-b]pyridin-3-yloxy)tripyrrolidinophosphonium hexafluorophosphate (PyAOP);

[00215] alternatively, this step can be carried out in an organic base, the organic base includes, but is not limited to, any one or the combination of two or more of the following: triethylamine, diisopropylethylamine (DIEA), pyridine, 4-dimethylaminopyridine (DMAP), 2,6-dimethylpyridine (Lutidine), 1,8-diazabicycloundec-7-ene (DBU) and N-methylmorpholine.

[00216] Alternatively, step c-C2 can be carried out in an aprotic solvent under the action of a base. The aprotic solvent includes, but is not limited to, any one or the combination of two or more of the following: tetrahydrofuran (THF), N,N-dimethylformamide (DMF), N,N-dimethylacetamide (DMA), N-methylpyrrolidone (NMP), and dioxane; the alkali includes, but is not limited to, any one of sodium hydride, and lithium bistrimethylsilylamide, or both.

[00217] The above steps c-A), c-B) and c-C) are parallel selection steps, that is, the compound represented by formula (I) can be prepared by the compound represented by formula (VIII) through one of steps c-A), c-B) and c-C), that is, the compound represented by formula (I) can be prepared by the compound represented by formula (VIII) through step c-A), or the compound represented by formula (I) can be prepared by the compound represented by formula (VIII)

through step c-B), or the compound represented by formula (I) can be prepared by the compound represented by formula (VIII) through step c-C).

[00218] It is apparent that the compounds of Formula (I), the isomers, crystalline forms or prodrugs, and pharmaceutically acceptable salts thereof, may exist in both solvated and unsolvated forms. For example, the solvated form can be a hydrate form. The disclosure includes both solvated and unsolvated forms.

[00219] The compounds of the present disclosure may have asymmetric carbon atoms. Such diastereomeric mixtures can be separated into their individual diastereomers on the basis of their physical chemical differences by methods known to those skilled in the art, for example, by chromatography or fractional crystallization. Enantiomers can be separated by converting the enantiomeric mixtures into a diastereomeric mixture by reaction with an appropriate optically active compound, separating the diastereomers and converting (e.g., hydrolyzing) the individual diastereomers to the corresponding pure enantiomers. All such isomers, including diastereomer mixtures and pure enantiomers are considered as part of the disclosure.

[00220] The compound of the present disclosure as an active ingredient, and the method of preparing the same, are both included in the present disclosure. Moreover, the crystalline form of some of the compounds may exist as polymorphs, and such forms may also be included in the present disclosure. Additionally, some of the compounds may form solvates with water (i.e., hydrates) or common organic solvents, and such solvates are also included within the scope of the disclosure.

[00221] The compounds of the disclosure may be used in the free form for treatment or, when appropriate, in the form of a pharmaceutically acceptable salt or other derivative for treatment. As used herein, the term "pharmaceutically acceptable salt" refers to organic and inorganic salts of the compounds of the present disclosure which are suitable for use in human and lower animals without undue toxicity, irritation, allergic response, etc., and have reasonable benefit/risk ratio. Pharmaceutically acceptable salts of amines, carboxylic acids, phosphonates, and other types of compounds are well known in the art. The salt can be formed by reacting a compound of the disclosure with a suitable free base or acid, including, but not limited to, salts with inorganic acids such as hydrochloric acid, hydrobromic acid, phosphoric acid, sulfuric acid, perchloric acid or with organic acids such as acetic acid, oxalic acid, maleic acid, tartaric acid, citric acid, succinic acid, malonic acid. Or the salts may be obtained by methods well known in the art, such as ion exchange. Other pharmaceutically acceptable salts include adipate, alginate, ascorbate, aspartate, besylate, benzoate, bisulfate, borate, butyrate, camphorate, camphorsulfonate, citrate, digluconate, lauryl sulfate, ethanesulfonate, formate, fumarate,

glucoheptonate, glycerol phosphate, glyconate, hemisulfate, hexanoate, hydroiodide, 2-hydroxyethanesulfonate, lactobionate, lactate, laurate, lauryl sulfate, malate, maleate, mesylate, 2-naphthalenesulfonate, nicotinate, nitrate, oleate, palmitate, pamoate, pectate, persulphate, per-3-phenylpropionate, phosphate, picrate, propionate, stearate, sulfate, thiocyanate, p-toluenesulfonate, undecanoate, and the like. Representative alkali or alkaline earth metal salts include salts of sodium, lithium, potassium, calcium, magnesium, and the like. Other pharmaceutically acceptable salts include suitable non-toxic salts of ammonium, quaternary ammonium, and amine cations formed from halides, hydroxides, carboxylates, sulfates, phosphates, nitrates, lower alkyl sulfonates and aryl sulfonates.

10 **[00222]** Further, the term "prodrug" as used herein means that a compound can be converted into a compound of Formula (I) of the present disclosure *in vivo*. Such transformation is affected by hydrolysis of the prodrug in the blood or enzymatic conversion to the parent compound in the blood or tissue.

15 **[00223]** Pharmaceutical compositions of this disclosure comprise a compound of the formulas described herein or a pharmaceutically acceptable salt thereof; an additional agent selected from a kinase inhibitory agent (small molecule, polypeptide, antibody, etc.), an immunosuppressant, an anticancer agent, an anti-viral agent, anti-inflammatory agent, antifungal agent, antibiotic, or an anti-vascular hyper proliferation compound; and any pharmaceutically acceptable carrier, adjuvant or vehicle.

20 **[00224]** The compounds of the present disclosure may be used alone or in combination with one or more of other compounds of the present disclosure or with one or more of other agents. When administered in combination, the therapeutic agents can be formulated for simultaneous or sequential administration at different times, or the therapeutic agents can be administered as a single composition. By "combination therapy", it refers to the use of a compound of the disclosure in combination with another agent in the form of co-administration of each agent or sequential administration of each agent, in either case, for the purpose of achieving the optimal results. Co-administration includes dosage form for simultaneous delivery, as well as separate dosage forms for each compound. Thus, administration of the compounds of the disclosure can be combined with other therapies known in the art, for example, radiation therapy or cytostatic agents, cytotoxic agents, other anticancer agents, and the like as used in the treatment of cancer, in order to improve the symptoms of cancer. The administration sequence is not limited in the present disclosure. The compounds of the present disclosure may be administered before, simultaneously, or after other anticancer or cytotoxic agents.

[00225] To prepare the pharmaceutical ingredient of the present disclosure, one or more

compounds of Formula (I) or salts thereof as an active ingredient can be intimately mixed with a pharmaceutical carrier, which is carried out according to a conventional pharmaceutical Formulation technique. The carrier can be used in a wide variety of forms depending on the form of preparation which is designed for different administration modes (for example, oral or parenteral administration). Suitable pharmaceutically acceptable carriers are well known in the art. A description of some of these pharmaceutically acceptable carriers can be found in the Handbook of Pharmaceutical Excipients, published jointly by the American Pharmaceutical Association and the Pharmaceutical Society of Great Britain.

[00226] The pharmaceutical composition of the present disclosure may have the following forms, for example, those suitable for oral administration, such as tablets, capsules, pills, powders, sustained release forms, solutions or suspensions; those for parenteral injections such as clear solutions, suspensions, emulsion; or those for topical use such as ointments, creams; or as a suppository for rectal administration. The pharmaceutical ingredients may also be presented in unit dosage form for single administration in a precise dosage. The pharmaceutical ingredient will include a conventional pharmaceutical carrier or excipient and a compound as an active ingredient prepared according to the present disclosure, and may also include other medical or pharmaceutical preparations, carriers, adjuvants, and the like.

[00227] Therapeutic compounds can also be administered to mammals other than humans. The drug dosage for a mammal will depend on the species of the animal and its disease condition or its disordered condition. The therapeutic compound can be administered to the animal in the form of a capsule, a bolus, or a tablet or liquid drench. The therapeutic compound can also be introduced into the animal by injection or infusion. These drug forms are prepared in a traditional manner complying with standard veterinary practice. As an alternative, the therapeutic compounds can be mixed with the animal feed and fed to the animal, so that the concentrated feed additive or premix can be prepared by mixing ordinary animal feed.

[00228] It is a further object of the present disclosure to provide a method for treating cancer in a subject in need thereof, including a method for administering to the subject a therapeutically effective amount of a composition containing the compound of the present disclosure.

[00229] The present disclosure also includes the use of the compounds of the present disclosure or pharmaceutically acceptable derivatives thereof in the manufacture of drugs for the treatment of cancer and autoimmune diseases associated with tyrosine kinases EGFR, HER2, HER3 and HER4, wherein the diseases include, but are not limited to, cancer (including non-solid tumors, solid tumors, primary or metastatic cancer, as indicated elsewhere herein and including one or more of other therapies to which the cancer is resistant or refractory), as well as other diseases


(including, but not limited to, ocular fundus diseases, psoriasis, atheroma, pulmonary fibrosis, liver fibrosis, myelofibrosis, and the like). The cancer includes, but is not limited to any one of non-small cell lung cancer, small cell lung cancer, breast cancer, pancreatic cancer, glioma, glioblastoma, ovarian cancer, cervical cancer, colorectal cancer, melanoma, endometrial cancer, prostate cancer, bladder cancer, leukemia, gastric cancer, liver cancer, gastrointestinal stromal tumor, thyroid cancer, chronic granulocytic leukemia, acute myeloid leukemia, non-Hodgkin's lymphoma, nasopharyngeal carcinoma, esophageal cancer, brain tumor, B-cell and T-cell lymphoma, lymphoma, multiple myeloma, biliary cancer and sarcoma, and cholangiocarcinoma.

[00230] The present disclosure is better illustrated by referring to the examples provided below, wherein all temperatures are in degrees Celsius unless otherwise stated.


Detailed Embodiments

[00231] Synthesis of the compound represented by formula VIII (intermediate of compounds of the present disclosure)

[00232] Preparation of N-(4-phenoxyphenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine (Intermediate No. VIII-1)



[00233] Step 1) Preparation of 5-(2-hydroxyethoxy)-6-nitroquinazolin-4(3H)-one (II)




20

[00234] Ethylene glycol (352.7 g, 5.7 mol) was dissolved in 1L of DMF, cooled in an ice bath, to which sodium hydride was added (68.2 g, 2.8 mol), and stirred for 0.5 h. 5-chloro-6-nitroquinazolin-4(3H)-one (128 g, 0.57 mol) was added, and the reaction was slowly warmed to room temperature and stirred until the reaction was completed. Ethyl acetate was added until a large amount of solid was precipitated out, which was filtered with suction, and the resulting solid was slurried with water. The slurry was adjusted to weak acidic with hydrochloric acid and filtered with suction to afford 129.7g of a white solid with a yield of 91%. ¹H NMR (DMSO-d₆, 400 MHz) δ 12.55 (1H, s), 8.13-8.28 (2H, m), 7.52 (1H, d, J=8.9 Hz), 4.76 (1H, brs), 4.04-

4.32 (2H, m), 3.60-3.84 (2H, m); MS: 252[M+H]⁺.

[00235] Step 2) Preparation of 5-(2-chloroethoxy)-6-nitro-N-(4-phenoxyphenyl)quinazolin-4-amine (VII-1)




5 **[00236]** 5-(2-hydroxyethoxy)-6-nitroquinazolin-4(3H)-one (3 g, 11.94 mmol) was added to a round-bottomed flask, to which thionyl chloride was added and stirred to dissolve it, catalytic amount of dimethylformamide was added dropwise, the reaction solution was heated to reflux until the raw materials were completely reacted, the reaction solution was evaporated to dryness under reduced pressure to afford a yellow solid, which was directly dissolved in
10 dichloromethane, then 4-phenoxyaniline (2.2 g, 11.94 mmol) in ethanol was added and stirred until the reaction was completed. N-hexane was added and stirred until a large amount of solid was precipitated out, which was filtered with suction, washed with petroleum ether, and dried in air to afford 3.9 g of yellow solid with a yield of 88%. MS: 437[M+H]⁺.

[00237] Step 3) Preparation of N-(4-phenoxyphenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine (Intermediate No. VIII-1)
15

[00238] 5-(2-chloroethoxy)-6-nitro-N-(4-phenoxyphenyl)quinazolin-4-amine (VII-1) (3.9 g, 8.7 mmol) was added to a round-bottomed flask, to which a mixed solvent of ethanol and water was added. Then iron powder (1.3 g, 22.7 mmol) and acetic acid (1.85 mL, 32.27 mmol) were added in sequence, and the reaction solution was heated and stirred until the reaction was
20 completed. The solvent was distilled off, and extracted with ethyl acetate, concentrated, and subjected to column chromatography to afford 2.2 g of yellow solid with a yield of 65%. ¹H NMR (300 MHz, DMSO-d₆) δ 9.92 (s, 1H), 8.47 (s, 1H), 7.79 (d, J = 8.9 Hz, 2H), 7.44 – 7.37 (m, 3H), 7.35 – 7.28 (m, 1H), 7.16 – 7.13 (m, 1H), 7.09 – 7.06 (m, 2H), 7.04 – 7.00 (m, 2H), 6.82 – 6.74 (m, 1H), 4.71 – 4.66 (m, 2H), 4.13 – 4.08 (m, 2H); MS: 371[M+H]⁺.


25

[00239] Preparation of intermediate N-(4-(3-methylphenoxy)phenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine (Intermediate No. VIII-2)




[00240] Step 1) is the same as the step 1) of the synthetic route of Intermediate No. VIII-1.

[00241] Step 2) Preparation of 5-(2-chloroethoxy)-6-nitroquinazolin-4(3H)-one (III)




5 [00242] 5-(2-hydroxyethoxy)-6-nitroquinazolin-4(3H)-one (129.7 g, 0.52 mol) was placed in a flask, to which 200 ml of phosphorus oxychloride was added and heated to reflux until the reaction was completed. Phosphorus oxychloride was evaporated off, and slurried with water until a large amount of solid was precipitated out, which was filtered with suction to afford 120.8g of white solid with a yield of 87%. ¹H NMR (400 MHz, DMSO-d₆) δ 12.55 (s, 1H), 8.24 – 8.21 (m, 2H), 7.56 (d, J = 9.0 Hz, 1H), 4.41 (t, J = 5.4 Hz, 2H), 3.96 (t, J = 5.4 Hz, 2H); MS: 270[M+H]⁺.

[00243] Step 3) Preparation of 5-(2-chloroethoxy)-6-aminoquinazolin-4(3H)-one (IV)



15 [00244] 5-(2-chloroethoxy)-6-nitroquinazolin-4(3H)-one (120.8 g, 0.45 mol) was dissolved in a mixed solvent of methanol and tetrahydrofuran, to which 40 g raney nickel was added, and stirred at room temperature under hydrogen atmosphere until the reaction was completed. The reaction was filtered with suction, and concentrated to afford 107.4 g of a yellow solid with a yield of 100%. ¹H NMR (400 MHz, DMSO-d₆) δ 11.76 (s, 1H), 7.74 (s, 1H), 7.24 (d, J = 2.2 Hz, 2H), 5.32 (s, 2H), 4.15 (t, J = 5.6 Hz, 2H), 3.97 (t, J = 5.6 Hz, 2H); MS: 240[M+H]⁺.

20 [00245] Step 4) Preparation of 2,3,4,9-tetrahydro-10H-[1,4]oxazino[2,3-f]quinazolin-10-one (V)




[00246] 5-(2-chloroethoxy)-6-aminoquinazolin-4(3H)-one (107.4 g, 0.45 mol) was dissolved in 1L of DMF, to which triethylamine (94 mL, 0.68 mol) was added, heated until the reaction was completed, and then DMF was distilled off. Dichloromethane was added and stirred until

a large amount of solid was precipitated out, which was filtered with suction to afford 80g of a white solid with a yield of 88%. ¹H NMR (DMSO-d₆, 300 MHz) δ 11.66 (1H, s), 7.67 (1H, s), 6.96 – 7.03 (2H, m), 6.11 (1H, s), 4.13 – 4.21 (2H, m), 3.25 – 3.33 (2H, m); MS: 204[M+H]⁺.

[00247] Step 5) Preparation of N-(4-(3-methylphenoxy)phenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine (Intermediate No. VIII-2)


[00248] 2,3,4,9-tetrahydro-10H-[1,4]oxazino[2,3-f]quinazolin-10-one (0.5 g, 2.46 mmol), 4-(m-tolyloxy)aniline (979 mg, 4.92 mmol), (benzotriazol-1-yl)tris(dimethylamino)phosphonium hexafluorophosphate (BOP)(1.4 g, 3.20 mmol) were placed in a round-bottomed flask, to which 5 ml of acetonitrile was added, and 1,8-diazabicycloundec-7-ene(DBU)(0.56 g, 3.69 mmol) was added after stirring well, and then was stirred under room temperature until the reaction was completed. The solvent was distilled off, and the resulting mixture was purified using silica gel column chromatography to afford 710 mg of brown solid with a yield of 75%. MS: 385[M+H]⁺.

[00249] Preparation of intermediate N-(3-trifluoromethyl-4-(3-fluorobenzoyloxy)phenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine (Intermediate No. VIII-3)



[00250] Step 1) to Step 4) are the same as Step 1) to Step 4) in the preparation method of Intermediate No. VIII-2.

[00251] Step 5) Preparation of 10-((1H-benzo[d][1,2,3]triazol-1-yl)oxy)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazoline (VI)




[00252] 2,3,4,9-tetrahydro-10H-[1,4]oxazino[2,3-f]quinazolin-10-one (20.3g, 100 mmol) and (benzotriazol-1-yloxy)tris(dimethylamino)phosphonium hexafluorophosphate (BOP) (44.2 g, 100 mmol) were placed in a round-bottomed flask, to which acetonitrile was added and stirred well, and 1,8-diazabicycloundec-7-ene (DBU)(15.2 g, 100 mmol) was then added, and stirred at room temperature until the reaction was completed. Water was added and stirred until a large amount of solid was precipitated out, which was filtered with suction to afford 28g of yellow solid with a yield of 87%. ¹H NMR (400 MHz, DMSO-d₆) δ 8.26 (s, 1H), 8.19 (d, J = 8.4 Hz, 1H), 7.78 (d, J = 8.3 Hz, 1H), 7.69-7.58 (m, 1H), 7.58 – 7.47 (m, 3H), 6.70 (s, 1H), 4.41 (t, J = 4.2 Hz, 2H), 3.58 – 3.47 (m, 2H); MS: 321[M+H]⁺.

[00253] Step 6) Preparation of N-(3-trifluoromethyl-4-(3-fluorobenzyloxy)phenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine(Intermediate No. VIII-3)

[00254] 10-((1H-benzo[d][1,2,3]triazol-1-yl)oxy)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazoline (320 mg, 1 mmol) and 4-((3-fluorobenzyl)oxy)-3-(trifluoromethyl)aniline (285 mg, 1 mmol), p-toluenesulfonic acid monohydrate (17 mg, 0.1 mmol) were dissolved in isopropanol, and stirred at room temperature until the reaction was completed. Water was added and stirred, and filtered with suction to afford 424 mg of yellow solid product with a yield of 90%. MS: 471[M+H]⁺.













[00255] Preparation of Intermediate No. VIII-4 to Intermediate No. VIII-63






















[00256] Step 1) to Step 4) are the same as Step 1) to Step 4) in the synthetic method of Intermediate No. VIII-2.








[00257] Step 5): See Step 5) in the synthetic route of Intermediate No. VIII-2, wherein the same operation was used, and the method was carried out using 2,3,4,9-tetrahydro-10H-[1,4]oxazino[2,3-f]quinazolin-10-one (V) as the starting material, and replacing 4-m-tolyloxyaniline with an equal molar equivalent of R¹XH in the table below. Specific compounds are as follows:








Intermediate No.	R ¹ X	Compound name	LCMS m/z = (M+H) +
------------------	------------------	---------------	-----------------------------








VIII-4		N-(3-chloro-4-(3-fluorobenzoyloxy)phenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine	437
VIII-5		N-(3-chloro-4-(pyridin-2-ylmethoxy)phenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine	420
VIII-6		N-(2-fluoro-4-phenoxyphenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine	389
VIII-7		N-(3-fluoro-4-phenoxyphenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine	389
VIII-8		N-(4-(2-fluorophenoxy)phenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine	389
VIII-9		N-(4-(3-(trifluoromethyl)phenoxy)phenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine	439
VIII-10		N-(4-(3-chlorophenoxy)phenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine	405
VIII-11		N-(4-(3-fluorophenoxy)phenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine	389
VIII-12		N-(4-(4-fluorophenoxy)phenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine	389
VIII-13		N-(4-(4-chlorophenoxy)phenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine	405
VIII-14		N-(4-(2-methoxyphenoxy)phenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine	401
VIII-15		N-(4-(2-pyridyloxy)phenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine	372








VIII-16		N-(4-(3-pyridyloxy)phenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine	372
VIII-17		N-(4-(3-methoxyphenoxy)phenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine	401
VIII-18		N-(4-(thiazol-2-yloxy)phenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine	378
VIII-19		N-(3-chloro-4-phenoxyphenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine	405
VIII-20		N-(4-(4-fluorobenzoyloxy)phenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine	403
VIII-21		N-(4-(3-fluorobenzoyloxy)phenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine	403
VIII-22		N-(4-(3-trifluoromethylbenzyloxy)phenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine	453
VIII-23		N-(4-(pyridin-2-ylmethoxy)phenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine	386
VIII-24		N-(4-(thiophen-2-ylmethoxy)phenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine	391
VIII-25		N-(4-(thiazol-2-ylmethoxy)phenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine	392
VIII-26		N-(4-(benzylthio)phenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine	401


VIII-27		N-(3-methoxy-4-(3-fluorobenzoyloxy)phenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine	433
VIII-28		N-(3-fluoro-4-(3-fluorobenzoyloxy)phenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine	421
VIII-29		N-(4-(4-fluorophenylthio)phenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine	405
VIII-30		N-(4-(2-fluoro-5-methylphenoxy)phenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine	403
VIII-31		N-(4-(2-fluoro-5-chlorophenoxy)phenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine	423
VIII-32		N-(4-(2,5-difluorophenoxy)phenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine	407
VIII-33		N-(4-(1-(3-fluorophenyl)ethoxy)phenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine	417
VIII-34		N-(4-(1-(pyridin-2-yl)ethoxy)phenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine	400

VIII-35		N-(4-(pyridin-3-ylmethoxy)phenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine	386
VIII-36		N-(4-(pyridin-4-ylmethoxy)phenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine	386
VIII-37		N-(4-(2-(2-(dimethylamino)ethoxy)phenoxy)phenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine	458
VIII-38		N-(4-(2-(3-(dimethylamino)propoxy)phenoxy)phenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine	472
VIII-39		N-(4-(2-(2-methoxyethoxy)phenoxy)phenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine	445
VIII-40		N-(4-phenoxyphenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine	371
VIII-41		N-(4-(benzyloxy)phenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine	385

VIII-42		N-(4-(2-fluorobenzyloxy)phenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine	403
VIII-43		N-(4-(2-chlorobenzyloxy)phenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine	419
VIII-44		N-(4-(2-methylbenzyloxy)phenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine	399
VIII-45		N-(4-(2-methoxybenzyloxy)phenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine	415
VIII-46		N-(4-(3-chlorobenzyloxy)phenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine	419
VIII-47		N-(4-(3-methylbenzyloxy)phenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine	399
VIII-48		N-(4-(3-cyanobenzyloxy)phenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine	410

VIII-49		N-(4-(3-methoxybenzyloxy)phenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine	415
VIII-50		N-(4-(4-chlorobenzyloxy)phenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine	419
VIII-51		N-(4-(4-methylbenzyloxy)phenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine	399
VIII-52		N-(4-(2,5-difluorobenzyloxy)phenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine	421
VIII-53		N-(4-(2-chloro-5-fluorobenzyloxy)phenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine	437
VIII-54		10-(3-chloro-4-(pyridin-2-ylmethoxy)phenoxy)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazoline	421
VIII-55		10-(3-chloro-4-(3-fluorobenzyloxy)phenoxy)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazoline	438


VIII-56		10-(4-phenoxyphenoxy)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazoline	372
VIII-57		10-(4-(4-chlorophenoxy)phenoxy)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazoline	406
VIII-58		10-(4-(4-fluorophenoxy)phenoxy)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazoline	390
VIII-59		10-(4-(2,5-dichlorophenoxy)phenoxy)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazoline	440
VIII-60		10-(4-(pyridin-2-ylmethoxy)phenoxy)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazoline	387
VIII-61		10-(4-(2-fluorobenzoyloxy)phenoxy)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazoline	404
VIII-62		10-(4-(3-fluorobenzoyloxy)phenoxy)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazoline	404

VIII-63		10-(4-(4-fluorobenzoyloxy)phenoxy)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazoline	404
---------	---	--	-----

[00258] Preparation of example compounds


[00259] Example 1

[00260] Preparation of 1-(10-((4-(3-(trifluoromethyl)phenoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one



[00261] N-(4-(3-trifluoromethylphenoxy)phenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine (Intermediate No. VIII-9)(219 mg, 0.5 mmol) was dissolved in tetrahydrofuran, to which acryloyl chloride (45.3 mg, 0.5 mmol) was added, and stirred at room temperature until the reaction was completed. The reaction was quenched by adding potassium carbonate aqueous solution, extracted with ethyl acetate, and the organic phase was concentrated and purified by silica gel column chromatography to afford 196 mg of an off-white solid, with a yield of 80%. ¹H NMR (DMSO-d₆, 300 MHz) δ 9.92 (s, 1H), 8.47 (s, 1H), 8.03 – 7.71 (m, 3H), 7.69–7.58 (m, 1H), 7.48 (d, J=7.8 Hz, 1H), 7.38 – 7.23 (m, 3H), 7.24 – 7.03 (m, 2H), 6.92 – 6.69 (m, 1H), 6.43 – 6.18 (m, 1H), 6.00 – 5.76 (m, 1H), 4.68 (t, J=4.5 Hz, 2H), 4.07 (t, J=4.5 Hz, 2H); ¹³CNMR (101 MHz, DMSO-d₆) δ 158.57, 157.66, 154.56, 151.72, 149.22, 144.48, 142.80, 135.76, 131.85, 130.33, 125.24, 123.57, 122.09, 121.93, 120.45, 119.89, 119.85, 119.21, 114.25, 114.21, 106.39, 68.71, 60.34, 45.62; MS: 493[M+H]⁺.





[00262] Examples 2-42












[00263] With reference to the preparation method of Example 1, wherein exactly the same operations were used, and N-(4-(3-trifluoromethylphenoxy)phenyl)-3,4-dihydro-2H-





[1,4]oxazino[2,3-f]quinazolin-10-amine was replaced with the same molar equivalent of intermediate represented by formula (VIII) wherein R¹X is the substituent in the table below.






The specific example compounds are shown in the table below:





Ex am ple No	Starti ng Inter medi ate No.	R ¹ X	Example compounds		
			Name	LCMS m/z = (M+H) +	HNMR
2	VIII-1		1-(10-((4-phenoxyphenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one	425	(400 M, DMSO-d ₆) δ9.88(s, 1H), 8.46(s, 1H), 7.81-7.79(m, 3H), 7.42-7.38(m, 2H), 7.32-7.30(m, 1H), 7.15-7.07(m, 5H), 7.03-7.01(m, 1H), 6.35-6.30(m, 1H), 5.90-5.87(m, 1H), 4.68(m, 2H), 4.07(m, 2H).
3	VIII-4		1-(10-((3-chloro-4-((3-fluorobenzoyloxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one	491	(400 M, DMSO-d ₆) δ9.83(s, 1H), 8.47(s, 1H), 7.99(s, 1H), 7.68-7.66(m, 1H), 7.48-7.45(m, 1H), 7.34-7.25(m, 6H), 7.21-7.17(m, 1H), 6.75-6.70(m, 1H), 5.80-5.76(m, 1H), 5.28(s, 2H), 4.74 – 4.66 (m, 2H), 4.06 – 3.96 (m, 2H).
4	VIII-5		1-(10-((3-chloro-4-(pyridin-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one	474	(400 M, DMSO-d ₆) δ9.82(s, 1H), 8.61(s, 1H), 8.46(s, 1H), 8.01(s, 1H), 7.89-7.86(m, 1H), 7.68-7.57(m, 2H), 7.39-7.36(m, 1H), 7.32-7.25(m, 2H), 7.21-7.17(m, 1H), 6.81-6.75(m, 1H), 6.34-6.30(m, 1H), 5.90-5.87(m, 1H), 5.31(s, 2H), 4.74 – 4.66 (m, 2H), 4.06 – 3.96 (m, 2H).
5	VIII-2		1-(10-((4-(m-tolyloxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,	439	(300 MHz, DMSO-d ₆) δ 9.87 (s, 1H), 8.45 (s, 1H), 7.82 – 7.70 (m, 3H), 7.39 – 7.20 (m, 2H), 7.06 (d, J = 8.9 Hz, 2H), 6.94





			3-f[quinazolin-4-yl]prop-2-en-1-one		(d, J = 7.5 Hz, 1H), 6.87 – 6.65 (m, 3H), 6.39 – 6.22 (m, 1H), 6.05 – 5.69 (m, 1H), 4.67 (t, J = 4.6 Hz, 2H), 4.07 (t, J = 4.6 Hz, 2H), 2.29 (s, 3H).
6	VIII-10		1-(10-((4-(3-chlorophenoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one	459	(300 MHz, DMSO-d ₆) δ 9.90 (s, 1H), 8.46 (s, 1H), 8.04 – 7.68 (m, 3H), 7.49 – 7.37 (m, 1H), 7.32 (d, J = 9.0 Hz, 1H), 7.24 – 7.10 (m, 3H), 7.09 – 6.93 (m, 2H), 6.88 – 6.62 (m, 1H), 6.49 – 6.14 (m, 1H), 5.96 – 5.75 (m, 1H), 4.68 (t, J = 4.6 Hz, 2H), 4.07 (t, J = 4.5 Hz, 2H).
7	VIII-11		1-(10-((4-(3-fluorophenoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one	443	(300 MHz, DMSO-d ₆) δ 9.90 (s, 1H), 8.47 (s, 1H), 7.94 – 7.74 (m, 3H), 7.51 – 7.38 (m, 1H), 7.32 (d, J = 9.0 Hz, 1H), 7.15 (d, J = 8.8 Hz, 2H), 7.04 – 6.90 (m, 1H), 6.90 – 6.71 (m, 3H), 6.38 – 6.24 (m, 1H), 6.01 – 5.73 (m, 1H), 4.75 – 4.60 (m, 2H), 4.14 – 4.00 (m, 2H).
8	VIII-8		1-(10-((4-(2-fluorophenoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one	443	(300 MHz, DMSO-d ₆) δ 9.85 (s, 1H), 8.43 (s, 1H), 7.91 – 7.70 (m, 3H), 7.40 (s, 1H), 7.30 (d, J = 9.0 Hz, 1H), 7.27 – 7.11 (m, 3H), 7.04 (d, J = 8.8 Hz, 2H), 6.87 – 6.69 (m, 1H), 6.40 – 6.25 (m, 1H), 5.93 – 5.82 (m, 1H), 4.66 (t, J = 4.6 Hz, 2H), 4.06 (t, J = 4.6 Hz, 2H).
9	VIII-12		1-(10-((4-(4-fluorophenoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one	443	(300 MHz, DMSO-d ₆) δ 9.85 (s, 1H), 8.44 (s, 1H), 7.82 – 7.73 (m, 3H), 7.35 – 7.16 (m, 3H), 7.14 – 7.00 (m, 4H), 6.86 – 6.71 (m, 1H), 6.38 – 6.25 (m, 1H), 5.93 – 5.82 (m, 1H), 4.67 (t, J = 4.6 Hz, 2H), 4.06 (t, J = 4.6 Hz, 2H).





10	VIII-13		1-(10-((4-(4-chlorophenoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one	459	(300 MHz, DMSO-d ₆) δ 9.89 (s, 1H), 8.56 – 8.36 (m, 1H), 8.03 – 7.62 (m, 3H), 7.44 (d, J = 8.4 Hz, 2H), 7.31 (d, J = 8.8 Hz, 1H), 7.16 – 6.97 (m, 4H), 6.78 (s, 1H), 6.32 (d, J = 16.7 Hz, 1H), 5.88 (d, J = 10.4 Hz, 1H), 4.77 – 4.57 (m, 2H), 4.17 – 3.95 (m, 2H).
11	VIII-14		1-(10-((4-(2-methoxyphenoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one	455	(400 MHz, DMSO-d ₆) δ 9.79 (s, 1H), 8.41 (s, 1H), 7.82 (s, 1H), 7.73 – 7.61 (m, 2H), 7.29 (d, J = 9.0 Hz, 1H), 7.24 – 7.16 (m, 2H), 7.10 – 6.97 (m, 2H), 6.93 – 6.87 (m, 2H), 6.86 – 6.71 (m, 1H), 6.39 – 6.27 (m, 1H), 5.96 – 5.83 (m, 1H), 4.66 (t, J = 4.6 Hz, 2H), 4.06 (t, J = 4.6 Hz, 2H), 3.77 (s, 3H).
12	VIII-15		1-(10-((4-(pyridin-2-yloxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one	426	(300 MHz, DMSO-d ₆) δ 9.89 (s, 1H), 8.49 – 8.32 (m, 3H), 7.89 – 7.79 (m, 3H), 7.47 – 7.36 (m, 2H), 7.32 (d, J = 9.0 Hz, 1H), 7.15 (d, J = 8.8 Hz, 2H), 6.87 – 6.71 (m, 1H), 6.38 – 6.26 (m, 1H), 5.93 – 5.83 (m, 1H), 4.68 (t, J = 4.6 Hz, 2H), 4.07 (t, J = 4.5 Hz, 2H).
13	VIII-16		1-(10-((4-(pyridin-3-yloxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one	426	(300 MHz, DMSO-d ₆) δ 9.89 (s, 1H), 8.46 (s, 1H), 8.16 (d, J = 4.6 Hz, 1H), 7.94 – 7.71 (m, 4H), 7.32 (d, J = 9.1 Hz, 1H), 7.22 – 7.10 (m, 3H), 7.04 (d, J = 8.3 Hz, 1H), 6.91 – 6.62 (1H, m), 6.41 – 6.26 (m, 1H), 5.94 – 5.82 (m, 1H), 4.74 – 4.61 (m, 2H), 4.13 – 4.01 (m, 2H).
14	VIII-17		1-(10-((4-(3-methoxyphenoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one	455	(300 MHz, DMSO-d ₆) δ 9.88 (s, 1H), 8.45 (s, 1H), 7.99 – 7.69 (m, 3H), 7.41 – 7.19 (m, 2H), 7.09 (d, J = 8.8 Hz, 2H), 6.89 – 6.67 (m, 2H), 6.65 –

			3-f[quinazolin-4-yl]prop-2-en-1-one		6.49 (m, 2H), 6.39 – 6.23 (m, 1H), 5.99 – 5.83 (m, 1H), 4.76 – 4.61 (m, 2H), 4.14 – 3.98 (m, 2H), 3.75-3.72 (m, 3H).
15	VIII-18		1-(10-((4-(thiazol-2-yloxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one	432	(300 MHz, DMSO-d ₆) δ 9.95 (s, 1H), 8.49 (s, 1H), 8.03 – 7.72 (m, 3H), 7.47 – 7.28 (m, 4H), 7.22 (d, J = 3.8 Hz, 1H), 6.91 – 6.65 (m, 1H), 6.40 – 6.21 (m, 1H), 5.97 – 5.81 (m, 1H), 4.69 (t, J = 4.6 Hz, 2H), 4.07 (t, J = 4.6 Hz, 2H).
16	VIII-7		1-(10-((3-fluoro-4-phenoxyphenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one	443	(400 MHz, DMSO-d ₆) δ 9.99 (s, 1H), 8.55 (s, 1H), 8.27 – 8.03 (m, 1H), 7.88 (s, 1H), 7.69 – 7.56 (m, 1H), 7.52 – 7.30 (m, 3H), 7.30 – 7.19 (m, 1H), 7.17 – 7.07 (m, 1H), 7.04 – 6.91 (m, 2H), 6.87 – 6.68 (m, 1H), 6.50 – 6.24 (m, 1H), 6.02 – 5.81 (m, 1H), 4.75 – 4.57 (m, 2H), 4.08 (t, J = 4.7 Hz, 2H).
17	VIII-19		1-(10-((3-chloro-4-phenoxyphenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one	459	(300 MHz, DMSO-d ₆) δ 9.96 (s, 1H), 8.50 (s, 1H), 8.06 – 7.69 (m, 3H), 7.51 – 7.31 (m, 5H), 7.29-7.18 (m, 2H), 6.78 (s, 1H), 6.32 (d, J = 16.8 Hz, 1H), 5.88 (d, J = 10.7 Hz, 1H), 4.77 – 4.59 (m, 2H), 4.18 – 4.00 (m, 2H).
18	VIII-20		1-(10-((4-(4-fluorobenzyl)oxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one	457	(300 MHz, DMSO-d ₆) δ 9.76 (s, 1H), 8.40 (s, 1H), 7.80 (s, 1H), 7.72 – 7.61 (m, 2H), 7.59 – 7.46 (m, 2H), 7.36 – 7.15 (m, 3H), 7.04 (d, J = 8.8 Hz, 2H), 6.88 – 6.69 (m, 1H), 6.41 – 6.24 (m, 1H), 5.98 – 5.77 (m, 1H), 5.11 (s, 2H), 4.66 (t, J = 4.7 Hz, 2H), 4.06 (t, J = 4.6 Hz, 2H).

19	VIII-21		1-(10-((4-((3-fluorobenzyl)oxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one	457	(300 MHz, DMSO-d ₆) δ 9.78 (s, 1H), 8.40 (s, 1H), 7.81 (s, 1H), 7.65 (d, J = 8.8 Hz, 2H), 7.55 – 7.40 (m, 1H), 7.38 – 7.24 (m, 3H), 7.20 – 7.11 (m, 1H), 7.05 (d, J = 8.9 Hz, 2H), 6.78 (s, 1H), 6.41 – 6.23 (m, 1H), 5.95 – 5.80 (m, 1H), 5.17 (s, 2H), 4.74 – 4.60 (m, 2H), 4.17 – 4.00 (m, 2H).
20	VIII-22		1-(10-((4-((3-trifluoromethylbenzyl)oxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one	507	(300 MHz, DMSO-d ₆) δ 9.77 (s, 1H), 8.40 (s, 1H), 7.87 – 7.59 (m, 7H), 7.28 (d, J = 9.0 Hz, 1H), 7.07 (d, J = 8.8 Hz, 2H), 6.86 – 6.70 (m, 1H), 6.38 – 6.25 (m, 1H), 5.93 – 5.82 (m, 1H), 5.25 (s, 2H), 4.66 (t, J = 4.6 Hz, 2H), 4.06 (t, J = 4.7 Hz, 2H).
21	VIII-23		1-(10-((4-(pyridin-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one	440	(300 MHz, DMSO-d ₆) δ 9.78 (s, 1H), 8.59 (s, 1H), 7.91 – 7.79 (m, 2H), 7.65 (d, J = 9.9 Hz, 2H), 7.58 – 7.44 (m, 2H), 7.37 – 7.25 (m, 2H), 7.08 – 7.01 (m, 2H), 6.77 (s, 1H), 6.40 – 6.22 (m, 1H), 5.96 – 5.80 (m, 1H), 5.21 (s, 2H), 4.79 – 4.61 (m, 2H), 4.15 – 3.96 (m, 2H).
22	VIII-24		1-(10-((4-(thiophen-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one	445	(300 MHz, DMSO-d ₆) δ 9.77 (s, 1H), 8.40 (s, 1H), 7.80 (s, 1H), 7.66 (d, J = 8.8 Hz, 2H), 7.56 (d, J = 5.2 Hz, 1H), 7.36 – 7.18 (m, 2H), 7.13 – 6.98 (m, 3H), 6.88 – 6.69 (m, 1H), 6.41 – 6.26 (m, 1H), 5.97 – 5.82 (m, 1H), 5.32 (s, 2H), 4.70 – 4.61 (m, 2H), 4.12 – 3.97 (m, 2H).
23	VIII-25		1-(10-((4-(thiazol-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one	446	(400 MHz, DMSO-d ₆) δ 9.78 (s, 1H), 8.41 (s, 1H), 7.87 – 7.79 (m, 3H), 7.73 – 7.65 (m, 2H), 7.29 (d, J = 9.0 Hz, 1H), 7.14

			[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one		– 7.07 (m, 2H), 6.85 – 6.73 (m, 1H), 6.38 – 6.27 (m, 1H), 5.92 – 5.83 (m, 1H), 5.47 (s, 2H), 4.67 (t, J = 4.6 Hz, 2H), 4.06 (t, J = 4.7 Hz, 2H).
24	VIII-26		1-(10-((4-(benzylthio)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one	455	(300 MHz, DMSO-d ₆) δ 9.89 (s, 1H), 8.48 (s, 1H), 7.87 – 7.74 (m, 3H), 7.37 (d, J = 2.7 Hz, 1H), 7.39 – 7.17 (m, 7H), 6.86 – 6.70 (m, 1H), 6.38 – 6.25 (m, 1H), 5.93 – 5.82 (m, 1H), 4.67 (t, J = 4.6 Hz, 2H), 4.22 (s, 2H), 4.06 (t, J = 4.7 Hz, 2H).
25	VIII-3		1-(10-((4-((3-fluorobenzyl)oxy)-3-(trifluoromethyl)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one	525	(300 MHz, DMSO-d ₆) δ 9.87 (s, 1H), 8.44 (s, 1H), 8.07 (d, J = 2.6 Hz, 1H), 8.02 – 7.91 (m, 1H), 7.84 (s, 1H), 7.55 – 7.41 (m, 1H), 7.37 (d, J = 9.1 Hz, 1H), 7.34 – 7.23 (m, 3H), 7.23 – 7.14 (m, 1H), 6.86 – 6.71 (m, 1H), 6.39 – 6.26 (m, 1H), 5.94 – 5.84 (m, 1H), 5.34 (s, 2H), 4.66 (t, J = 4.5 Hz, 2H), 4.06 (t, J = 4.5 Hz, 2H).
26	VIII-27		1-(10-((4-((3-fluorobenzyl)oxy)-3-(methoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one	487	(300 MHz, DMSO-d ₆) δ 9.76 (s, 1H), 8.43 (s, 1H), 7.81 – 7.79 (m, 1H), 7.59 – 7.37 (m, 2H), 7.36 – 7.25 (m, 4H), 7.16 (s, 1H), 7.04 (d, J = 8.0 Hz, 1H), 6.78 – 6.75 (m, 1H), 6.32 (d, J = 16.7 Hz, 1H), 5.88 (d, J = 10.2 Hz, 1H), 5.14 (s, 2H), 4.74 – 4.59 (m, 2H), 4.11 – 3.99 (m, 2H), 3.82 (s, 3H).
27	VIII-28		1-(10-((4-((3-fluorobenzyl)oxy)-3-fluorophenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one	475	(300 MHz, DMSO-d ₆) δ 9.87 (s, 1H), 8.45 (s, 1H), 7.88 – 7.73 (m, 2H), 7.46 – 7.40 (m, 2H), 7.30 – 7.26 (m, 4H), 7.16 – 7.00 (m, 1H), 6.79 – 6.77 (m, 1H), 6.35 – 6.31 (m, 1H), 5.88 – 5.86 (m, 1H), 5.23 (s, 2H), 4.71 – 4.58

			1-one		(m, 2H), 4.08 – 4.01 (m, 2H).
28	VIII-29		1-(10-((4-((4-fluorophenyl)thio)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one	459	(300 MHz, DMSO-d ₆) δ 9.96 (s, 1H), 8.53 (s, 1H), 8.21 (d, J = 2.6 Hz, 1H), 7.96 – 7.77 (m, 2H), 7.45 – 7.30 (m, 3H), 7.20 (d, J = 8.8 Hz, 1H), 7.16 – 7.09 (m, 1H), 7.01 – 6.91 (m, 2H), 6.87 – 6.70 (m, 1H), 6.33 (d, J = 16.4 Hz, 1H), 5.89 (d, J = 10.7 Hz, 1H), 4.72 – 4.64 (m, 2H), 4.13 – 4.03 (m, 2H).
29	VIII-6		1-(10-((2-fluoro-4-phenoxyphenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one	443	(400 MHz, DMSO-d ₆) δ 10.34 (s, 1H), 8.86 – 8.71 (m, 1H), 8.60 (s, 1H), 7.88 (s, 1H), 7.52 – 7.44 (m, 2H), 7.39 (d, J = 9.1 Hz, 1H), 7.28 – 7.16 (m, 4H), 7.00 – 6.93 (m, 1H), 6.84 – 6.72 (m, 1H), 6.40 – 6.28 (m, 1H), 5.95 – 5.87 (m, 1H), 4.43 (t, J = 4.7 Hz, 2H), 4.04 (t, J = 4.7 Hz, 2H).
30	VIII-37		1-(10-((4-(2-(2-(dimethylamino)ethoxy)phenoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one	512	(400 MHz, Methanol-d ₄) δ 8.44 – 8.22 (m, 1H), 7.98 – 7.49 (m, 3H), 7.48 – 7.21 (m, 2H), 7.19 – 6.97 (m, 4H), 6.92 – 6.65 (m, 2H), 6.47 – 6.31 (m, 1H), 5.95 – 5.77 (m, 1H), 5.06 – 5.00 (m, 2H), 4.13 – 4.02 (m, 2H), 3.30 (s, 6H), 2.35 – 2.19 (m, 4H).
31	VIII-38		1-(10-((4-(2-(3-(dimethylamino)propoxy)phenoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one	526	(400 MHz, Methanol-d ₄) δ 8.44 – 8.28 (m, 1H), 7.89 – 7.57 (m, 3H), 7.27 (d, J = 9.1 Hz, 1H), 7.23 – 7.11 (m, 2H), 7.08 – 7.01 (m, 2H), 6.96 (d, J = 7.7 Hz, 1H), 6.90 – 6.80 (m, 2H), 6.79 – 6.69 (m, 1H), 6.46 – 6.30 (m, 1H), 5.98 – 5.78 (m, 1H), 4.15 – 4.06 (m, 2H), 4.03 – 3.91 (m, 2H), 3.31 (s, 6H), 2.55 – 2.45 (m, 2H), 2.36 – 2.28 (m, 4H).


32	VIII-39		1-(10-((4-(2-(2-methoxyethoxy)phenoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one	499	(400 MHz, DMSO-d ₆) δ 9.90 – 9.77 (m, 1H), 8.55 – 8.31 (m, 1H), 7.86 – 7.64 (m, 3H), 7.40 – 7.24 (m, 2H), 7.17 (d, J = 6.6 Hz, 2H), 7.03 – 6.95 (m, 2H), 6.93 (d, J = 8.5 Hz, 1H), 6.78 (s, 1H), 6.32 (d, J = 16.8 Hz, 1H), 5.88 (d, J = 10.8 Hz, 1H), 4.66 (t, J = 4.8 Hz, 2H), 4.12 (t, J = 4.8 Hz, 2H), 4.06 (t, J = 4.5 Hz, 2H), 3.56 (t, J = 4.7 Hz, 2H), 3.23 (s, 3H).
33	VIII-56		1-(10-(4-phenoxyphenoxy)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one	426	(300 MHz, DMSO-d ₆) δ 8.60 (s, 1H), 8.03 (s, 1H), 7.53 – 7.39 (m, 3H), 7.28 (d, J = 8.6 Hz, 2H), 7.20 – 7.04 (m, 5H), 6.80 (s, 1H), 6.33 (d, J = 17.0 Hz, 1H), 5.89 (d, J = 10.7 Hz, 1H), 4.64 – 4.51 (m, 2H), 4.14 – 3.98 (m, 2H).
34	VIII-54		1-(10-(3-chloro-4-(pyridin-2-ylmethoxy)phenoxy)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one	475	(400 MHz, DMSO-d ₆) δ 8.68 – 8.53 (m, 2H), 8.13 – 7.95 (m, 1H), 7.94 – 7.85 (m, 1H), 7.61 (d, J = 7.8 Hz, 1H), 7.57 – 7.45 (m, 2H), 7.43 – 7.35 (m, 1H), 7.32 (d, J = 8.9 Hz, 1H), 7.26 – 7.19 (m, 1H), 6.87 – 6.74 (m, 1H), 6.39 – 6.28 (m, 1H), 5.93 – 5.86 (m, 1H), 5.33 (s, 2H), 4.54 (t, J = 4.5 Hz, 2H), 4.05 (t, J = 4.5 Hz, 2H).
35	VIII-55		1-(10-(3-chloro-4-((3-fluorobenzyl)oxy)phenoxy)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one	492	(400 MHz, DMSO-d ₆) δ 8.58 (s, 1H), 8.02 (s, 1H), 7.56 – 7.43 (m, 3H), 7.41 – 7.27 (m, 3H), 7.27 – 7.14 (m, 2H), 6.84 – 6.73 (m, 1H), 6.39 – 6.24 (m, 1H), 5.97 – 5.80 (m, 1H), 5.29 (s, 2H), 4.53 (t, J = 4.6 Hz, 2H), 4.05 (t, J = 4.6 Hz, 2H).

36	VIII-57		1-(10-(4-(4-chlorophenoxy)phenoxy)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one	460	(400 MHz, DMSO-d ₆) δ 8.59 (s, 1H), 8.02 (s, 1H), 7.51 – 7.45 (m, 3H), 7.32 – 7.27 (m, 2H), 7.17 – 7.12 (m, 2H), 7.11 – 7.07 (m, 2H), 6.85 – 6.74 (m, 1H), 6.37 – 6.29 (m, 1H), 5.92 – 5.88 (m, 1H), 4.54 (t, J = 4.4 Hz, 2H), 4.05 (t, J = 4.4 Hz, 2H).
37	VIII-58		1-(10-(4-(4-fluorophenoxy)phenoxy)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one	444	(400 MHz, DMSO-d ₆) δ 8.59 (s, 1H), 8.02 (s, 1H), 7.49 (d, J = 9.1 Hz, 1H), 7.43 – 7.19 (m, 4H), 7.19 – 7.01 (m, 4H), 6.85 – 6.76 (m, 1H), 6.41 – 6.26 (m, 1H), 5.95 – 5.81 (m, 1H), 4.54 (t, J = 4.6 Hz, 2H), 4.05 (t, J = 4.7 Hz, 2H).
38	VIII-59		1-(10-(4-(2,5-dichlorophenoxy)phenoxy)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one	494	(400 MHz, DMSO-d ₆) δ 8.59 (s, 1H), 8.01 (s, 1H), 7.82 – 7.78 (m, 1H), 7.51 – 7.46 (m, 2H), 7.31 – 7.27 (m, 2H), 7.20 – 7.16 (m, 1H), 7.11 – 7.07 (m, 2H), 6.84 – 6.75 (m, 1H), 6.36 – 6.29 (m, 1H), 5.92 – 5.86 (m, 1H), 4.54 (t, J = 4.4 Hz, 2H), 4.05 (t, J = 4.4 Hz, 2H).
39	VIII-60		1-(10-(4-(pyridin-2-ylmethoxy)phenoxy)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one	441	(400 MHz, DMSO-d ₆) δ 8.61 – 8.59 (m, 1H), 8.55 (s, 1H), 8.08 – 7.93 (m, 1H), 7.89 – 7.84 (m, 1H), 7.58 – 7.54 (m, 1H), 7.50 – 7.46 (m, 1H), 7.39 – 7.34 (m, 1H), 7.19 – 7.16 (m, 2H), 7.13 – 7.08 (m, 2H), 6.85 – 6.74 (m, 1H), 6.33 (d, J = 16.4 Hz, 1H), 5.91 – 5.86 (m, 1H), 5.22 (s, 2H), 4.53 (t, J = 4.8 Hz, 2H), 4.05 (t, J = 4.8 Hz, 2H).

40	VIII-61		1-(10-(4-((2-fluorobenzyl)oxy)phenoxy)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one	458	(400 MHz, DMSO-d ₆) δ 8.56 (s, 1H), 8.01 (s, 1H), 7.63 – 7.57 (m, 1H), 7.50 – 7.46 (m, 1H), 7.46 – 7.41 (m, 1H), 7.31 – 7.24 (m, 2H), 7.21 – 7.17 (m, 2H), 7.14 – 7.09 (m, 2H), 6.86 – 6.74 (m, 1H), 6.38 – 6.29 (m, 1H), 5.92 – 5.86 (m, 1H), 5.18 (s, 2H), 4.58 – 4.51 (m, 2H), 4.09 – 4.01 (m, 2H).
41	VIII-62		1-(10-(4-((3-fluorobenzyl)oxy)phenoxy)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one	458	(400 MHz, DMSO-d ₆) δ 8.55 (s, 1H), 7.99 (s, 1H), 7.50 – 7.42 (m, 2H), 7.35 – 7.29 (m, 2H), 7.20 – 7.14 (m, 3H), 7.11 – 7.07 (m, 2H), 6.86 – 6.71 (m, 1H), 6.37 – 6.27 (m, 1H), 5.92 – 5.84 (m, 1H), 5.18 (s, 2H), 4.56 – 4.49 (m, 2H), 4.08 – 4.00 (m, 2H).
42	VIII-63		1-(10-(4-((4-fluorobenzyl)oxy)phenoxy)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one	458	(400 MHz, DMSO-d ₆) δ 8.55 (s, 1H), 8.00 (s, 1H), 7.56 – 7.51 (m, 2H), 7.50 – 7.45 (m, 1H), 7.27 – 7.20 (m, 2H), 7.19 – 7.14 (m, 2H), 7.11 – 7.05 (m, 2H), 6.85 – 6.72 (m, 1H), 6.36 – 6.27 (m, 1H), 5.93 – 5.84 (m, 1H), 5.13 (s, 2H), 4.53 (t, J = 4.4 Hz, 2H), 4.05 (t, J = 4.4 Hz, 2H).

[00264] Example 43


[00265] Preparation of (E)-4-(dimethylamino)-1-(10-((4-(3-(trifluoromethyl)phenoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one



[00266] N-(4-(3-trifluoromethylphenoxy)phenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-




f]quinazolin-10-amine (VIII-9)(219 mg, 0.5 mmol) was dissolved in a mixed solvent of tetrahydrofuran and dimethylformamide, and trans-4-dimethylaminocrotonic acid hydrochloride (92 mg, 0.5 mmol) was added, and stirred at room temperature until the reaction was completed. The reaction was quenched by adding potassium carbonate aqueous solution, extracted with ethyl acetate, the organic phase was concentrated and purified by silica gel column chromatography to afford 208mg of an off-white solid with a yield of 76%. ¹H NMR (400 MHz, DMSO-d₆) δ 9.92 (s, 1H), 8.48 (s, 1H), 7.93 – 7.84 (m, 3H), 7.66 – 7.58 (m, 1H), 7.48 (d, J = 7.8 Hz, 1H), 7.35 – 7.25 (m, 3H), 7.23 – 7.14 (m, 2H), 6.88 – 6.76 (m, 1H), 6.58 (d, J = 15.2 Hz, 1H), 4.68 (t, J = 4.6 Hz, 2H), 4.06 (t, J = 4.6 Hz, 2H), 3.12 – 3.05 (m, 2H), 2.18 (s, 6H); ¹³CNMR (101 MHz, DMSO-d₆) δ 158.57, 157.66, 154.56, 151.72, 149.22, 144.48, 142.80, 135.76, 131.85, 130.33, 125.24, 123.57, 122.09, 121.93, 120.45, 119.89, 119.85, 119.21, 114.25, 114.21, 106.39, 68.71, 60.34, 45.62; MS: 550[M+H]⁺.




[00267] Examples 44-90









[00268] With reference to the preparation method of Example 43, wherein exactly the same operations were used, and N-(4-(3-trifluoromethylphenoxy)phenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine (VIII-9) was replaced with the same molar equivalent of intermediate represented by formula (VIII) wherein R¹X is the substituent in the table below. The specific example compounds are shown in the table below:




Example No.	Starting Intermediate No.	R ¹ X	Example compounds		
			Name	LC MS m/z = (M+H) ⁺	HNMR
44	VIII-1		(E)-4-(dimethylamino)-1-(10-((4-phenoxyphenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one	482	(400 M, CDCl ₃) δ9.56(s, 1H), 8.62(s, 1H), 7.71-7.68(m, 2H), 7.61(brs, 1H), 7.47-7.45(m, 1H), 7.38-7.34(m, 2H), 7.14-7.05(m, 6H), 6.72-6.64(m, 1H), 4.72-4.66(m, 2H),




					4.24-4.18(m, 2H), 3.38-3.31(m, 2H), 2.46(s, 6H)
45	VIII-4		(E)-1-(10-((3-chloro-4-((3-fluorobenzyl)oxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(dimethylamino)but-2-en-1-one	548	(400 MHz, DMSO-d ₆) δ 9.82 (s, 1H), 8.46 (s, 1H), 8.01 (d, J = 2.5 Hz, 1H), 7.83 (s, 1H), 7.71 – 7.63 (m, 1H), 7.53 – 7.43 (m, 1H), 7.37 – 7.22 (m, 4H), 7.21 – 7.12 (m, 1H), 6.89 – 6.74 (m, 1H), 6.58 (d, J = 15.3 Hz, 1H), 5.27 (s, 2H), 4.66 (t, J = 4.6 Hz, 2H), 4.05 (t, J = 4.6 Hz, 2H), 3.10 (d, J = 5.8 Hz, 2H), 2.19 (s, 6H).
46	VIII-5		(E)-1-(10-((3-chloro-4-(pyridin-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(dimethylamino)but-2-en-1-one	531	(400 MHz, DMSO-d ₆) δ 9.81 (s, 1H), 8.64 – 8.58 (m, 1H), 8.46 (s, 1H), 8.01 (d, J = 2.6 Hz, 1H), 7.92 – 7.75 (m, 2H), 7.71 – 7.63 (m, 1H), 7.58 (d, J = 7.9 Hz, 1H), 7.42 – 7.34 (m, 1H), 7.34 – 7.22 (m, 2H), 6.87 – 6.76 (m, 1H), 6.58 (d, J = 15.2 Hz, 1H), 5.31 (s, 2H), 4.66 (t, J = 4.6 Hz, 2H), 4.05 (t, J = 4.6 Hz, 2H), 3.09 (d, J = 5.8 Hz, 2H), 2.19 (s, 6H).
47	VIII-2		(E)-4-(dimethylamino)-1-(10-((4-(m-tolyl)oxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one	496	(400 MHz, DMSO-d ₆) δ 9.87 (s, 1H), 8.45 (s, 1H), 7.89 – 7.76 (m, 3H), 7.34 – 7.23 (m, 2H), 7.11 – 7.04 (m, 2H), 6.98 – 6.92 (m, 1H), 6.86 – 6.77 (m, 3H), 6.58 (d, J = 15.4 Hz,




					1H), 4.72 – 4.66 (m, 2H), 4.12 – 4.02 (m, 2H), 3.12 – 3.04 (m, 2H), 2.30 (s, 3H), 2.18 (s, 6H).
48	VIII-10		(E)-4-(dimethylamino)-1-(10-((4-(3-chlorophenoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one	516	(400 MHz, DMSO-d ₆) δ 9.90 (s, 1H), 8.47 (s, 1H), 7.90 – 7.82 (m, 3H), 7.42 (t, J = 8.1 Hz, 1H), 7.31 (d, J = 9.0 Hz, 1H), 7.22 – 7.11 (m, 3H), 7.07 – 6.94 (m, 2H), 6.88 – 6.76 (m, 1H), 6.59 (d, J = 15.3 Hz, 1H), 4.67 (t, J = 4.6 Hz, 2H), 4.06 (t, J = 4.6 Hz, 2H), 3.11 (d, J = 5.7 Hz, 2H), 2.20 (s, 6H).
49	VIII-11		(E)-4-(dimethylamino)-1-(10-((4-(3-fluorophenoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one	500	(400 MHz, DMSO-d ₆) δ 9.90 (s, 1H), 8.47 (s, 1H), 7.93 – 7.74 (m, 3H), 7.46 – 7.38 (m, 1H), 7.31 (d, J = 9.0 Hz, 1H), 7.18 – 7.12 (m, 2H), 6.99 – 6.93 (m, 1H), 6.89 – 6.77 (m, 3H), 6.59 (d, J = 15.3 Hz, 1H), 4.72 – 4.64 (m, 2H), 4.06 (t, J = 4.7 Hz, 2H), 3.13 – 3.05 (m, 2H), 2.18 (s, 6H).
50	VIII-8		(E)-4-(dimethylamino)-1-(10-((4-(2-fluorophenoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one	500	(400 MHz, DMSO-d ₆) δ 9.85 (s, 1H), 8.44 (s, 1H), 7.77 (d, J = 8.9 Hz, 3H), 7.46 – 7.36 (m, 1H), 7.30 (d, J = 9.0 Hz, 1H), 7.27 – 7.21 (m, 2H), 7.20 – 7.13 (m, 1H), 7.04 (d, J = 8.8 Hz, 2H), 6.87 – 6.77 (m, 1H), 6.58 (d, J = 15.3 Hz, 1H),

					4.66 (t, J = 4.6 Hz, 2H), 4.05 (t, J = 4.6 Hz, 2H), 3.10 – 3.04 (m, 2H), 2.17 (s, 6H).
51	VIII-12		(E)-4-(dimethylamino)-1-(10-((4-(4-fluorophenoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one	500	(400 MHz, DMSO-d ₆) δ 9.86 (s, 1H), 8.45 (s, 1H), 7.89 – 7.76 (m, 3H), 7.30 (d, J = 9.0 Hz, 1H), 7.28 – 7.19 (m, 2H), 7.13 – 7.01 (m, 4H), 6.89 – 6.77 (m, 1H), 6.58 (d, J = 15.3 Hz, 1H), 4.67 (t, J = 4.6 Hz, 2H), 4.05 (t, J = 4.6 Hz, 2H), 3.08 (d, J = 5.8 Hz, 2H), 2.18 (s, 6H).
52	VIII-13		(E)-4-(dimethylamino)-1-(10-((4-(4-chlorophenoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one	516	(400 MHz, DMSO-d ₆) δ 9.88 (s, 1H), 8.46 (s, 1H), 7.89 – 7.77 (m, 3H), 7.48 – 7.40 (m, 2H), 7.31 (d, J = 9.0 Hz, 1H), 7.15 – 7.08 (m, 2H), 7.08 – 7.01 (m, 2H), 6.86 – 6.77 (m, 1H), 6.58 (d, J = 15.2 Hz, 1H), 4.67 (t, J = 4.6 Hz, 2H), 4.06 (t, J = 4.7 Hz, 2H), 3.12 – 3.04 (m, 2H), 2.18 (s, 6H).
53	VIII-14		(E)-4-(dimethylamino)-1-(10-((4-(2-methoxyphenoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one	512	(400 MHz, DMSO-d ₆) δ 9.79 (s, 1H), 8.41 (s, 1H), 7.83 (s, 1H), 7.72 – 7.63 (m, 2H), 7.29 (d, J = 9.0 Hz, 1H), 7.25 – 7.16 (m, 2H), 7.08 – 7.03 (m, 1H), 7.03 – 6.95 (m, 1H), 6.95 – 6.87 (m, 2H), 6.87 – 6.76 (m, 1H), 6.57 (d, J = 15.3 Hz, 1H), 4.69 – 4.61 (m, 2H), 4.05




					(t, J = 4.7 Hz, 2H), 3.77 (s, 3H), 3.11 – 3.04 (m, 2H), 2.17 (s, 6H).
54	VIII-15		(E)-4-(dimethylamino)-1-(10-((4-(pyridin-2-yloxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one	483	(400 MHz, DMSO-d ₆) δ 9.90 (s, 1H), 8.47 (s, 1H), 8.45 (t, J = 1.8Hz, 1H), 8.36 (t, J = 3.0Hz, 1H), 7.85 (d, J = 8.8 Hz, 3H), 7.43 (t, J = 2.3Hz, 2H), 7.31 (d, J = 9.0 Hz, 1H), 7.15 (d, J = 8.8 Hz, 2H), 6.88 – 6.76 (m, 1H), 6.58 (d, J = 15.3 Hz, 1H), 4.67 (t, J = 4.7 Hz, 2H), 4.06 (t, J = 4.7 Hz, 2H), 3.08 (d, J = 5.9 Hz, 2H), 2.18 (s, 6H).
55	VIII-16		(E)-4-(dimethylamino)-1-(10-((4-(pyridin-3-yloxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one	483	(400 MHz, DMSO-d ₆) δ 9.89 (s, 1H), 8.47 (s, 1H), 8.19 – 8.13 (m, 1H), 7.90 – 7.83 (m, 2H), 7.83 – 7.78 (m, 2H), 7.31 (d, J = 9.0 Hz, 1H), 7.19 – 7.15 (m, 2H), 7.15 – 7.11 (m, 1H), 7.04 (d, J = 8.3 Hz, 1H), 6.86 – 6.78 (m, 1H), 6.58 (d, J = 15.3 Hz, 1H), 4.68 (t, J = 4.6 Hz, 2H), 4.06 (t, J = 4.6 Hz, 2H), 3.08 (d, J = 5.8 Hz, 2H), 2.18 (s, 6H).
56	VIII-17		(E)-4-(dimethylamino)-1-(10-((4-(3-methoxyphenoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one	512	(400 MHz, DMSO-d ₆) δ 9.87 (s, 1H), 8.46 (s, 1H), 7.96 – 7.74 (m, 3H), 7.35 – 7.24 (m, 2H), 7.15 – 7.03 (m, 2H), 6.89 – 6.75 (m, 1H), 6.74 – 6.69 (m, 1H), 6.63 – 6.52 (m, 3H),




					4.72 – 4.66 (m, 2H), 4.06 (t, J = 4.6 Hz, 2H), 3.75 (s, 3H), 3.12 – 3.04 (m, 2H), 2.18 (s, 6H).
57	VIII-18		(E)-4-(dimethylamino)-1-(10-((4-(thiazol-2-yloxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one	489	(400 MHz, DMSO-d ₆) δ 9.95 (s, 1H), 8.49 (s, 1H), 7.96 – 7.82 (m, 3H), 7.40 (d, J = 8.9 Hz, 2H), 7.36 – 7.28 (m, 2H), 7.23 (d, J = 3.8 Hz, 1H), 6.88 – 6.76 (m, 1H), 6.64 – 6.54 (m, 1H), 4.68 (t, J = 4.6 Hz, 2H), 4.06 (t, J = 4.6 Hz, 2H), 3.08 (d, J = 5.8 Hz, 2H), 2.18 (s, 6H).
58	VIII-7		(E)-4-(dimethylamino)-1-(10-((3-fluoro-4-phenoxyphenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one	500	(400 MHz, Methanol-d ₄) δ 8.51 (s, 1H), 8.32 (s, 1H), 8.05 – 8.01 (m, 1H), 7.88 – 7.82 (m, 1H), 7.75 – 7.72 (m, 1H), 7.51 – 7.45 (m, 1H), 7.27 – 7.21 (m, 2H), 7.18 – 7.14 (m, 2H), 6.94 – 6.87 (m, 2H), 6.70 – 6.64 (m, 1H), 4.74 (d, J = 4.8 Hz, 2H), 4.17 (t, J = 4.8 Hz, 2H), 3.23 – 3.19 (m, 2H), 2.33 (s, 6H).
59	VIII-19		(E)-4-(dimethylamino)-1-(10-((3-chloro-4-phenoxyphenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one	516	(400 MHz, DMSO-d ₆) δ 9.95 (s, 1H), 8.53 (s, 1H), 8.21 (d, J=2.6 Hz, 1H), 7.94 – 7.76 (m, 2H), 7.45 – 7.36 (m, 2H), 7.33 (d, J=9.0 Hz, 1H), 7.20 (d, J=8.8 Hz, 1H), 7.12 (t, J=7.4 Hz, 1H), 7.00 – 6.91 (m, 2H), 6.86 –



					6.80 (m, 1H), 6.62 – 6.56 (m, 1H), 4.68 (t, J=4.5 Hz, 2H), 4.06 (t, J=4.6 Hz, 2H), 3.14 – 3.06 (m, 2H), 2.18 (s, 6H).
60	VIII-20		(E)-4-(dimethylamino)-1-(10-((4-((4-fluorobenzyl)oxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one	514	(400 MHz, DMSO-d ₆) δ 9.76 (s, 1H), 8.41 (s, 1H), 7.90 – 7.72 (m, 1H), 7.70 – 7.62 (m, 2H), 7.57 – 7.49 (m, 2H), 7.32 – 7.20 (m, 3H), 7.10 – 7.02 (m, 2H), 6.87 – 6.76 (m, 1H), 6.57 (d, J = 15.3 Hz, 1H), 5.12 (s, 2H), 4.66 (t, J = 4.7 Hz, 2H), 4.05 (t, J = 4.6 Hz, 2H), 3.11 – 3.04 (m, 2H), 2.18 (s, 6H).
61	VIII-21		(E)-4-(dimethylamino)-1-(10-((4-((3-fluorobenzyl)oxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one	514	(400 MHz, DMSO-d ₆) δ 9.76 (s, 1H), 8.40 (s, 1H), 7.80 (s, 1H), 7.70 – 7.63 (m, 2H), 7.51 – 7.40 (m, 1H), 7.35 – 7.24 (m, 3H), 7.22 – 7.12 (m, 1H), 7.06 (d, J = 8.9 Hz, 2H), 6.87 – 6.75 (m, 1H), 6.57 (d, J = 15.2 Hz, 1H), 5.17 (s, 2H), 4.65 (t, J = 4.6 Hz, 2H), 4.04 (t, J = 4.6 Hz, 2H), 3.11 – 3.04 (m, 2H), 2.17 (s, 6H).
62	VIII-22		(E)-4-(dimethylamino)-1-(10-((4-((3-(trifluoromethyl)benzyl)oxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one	564	(400 MHz, DMSO-d ₆) δ 9.77 (s, 1H), 8.41 (s, 1H), 7.86 – 7.76 (m, 3H), 7.75 – 7.61 (m, 4H), 7.28 (d, J = 9.0 Hz, 1H), 7.08 (d, J = 8.9 Hz, 2H), 6.87 – 6.75 (m, 1H), 6.57

					(d, J = 15.3 Hz, 1H), 5.25 (s, 2H), 4.65 (t, J = 4.6 Hz, 2H), 4.05 (t, J = 4.6 Hz, 2H), 3.08 (d, J = 5.8 Hz, 2H), 2.18 (s, 6H).
63	VIII-23		(E)-4-(dimethylamino)-1-(10-((4-(pyridin-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one	497	(400 MHz, DMSO-d ₆) δ 9.76 (s, 1H), 8.63 – 8.56 (m, 1H), 8.40 (s, 1H), 7.90 – 7.80 (m, 1H), 7.80 (s, 1H), 7.66 (d, J = 8.9 Hz, 2H), 7.54 (d, J = 7.9 Hz, 1H), 7.40 – 7.32 (m, 1H), 7.28 (d, J = 8.9 Hz, 1H), 7.07 (d, J = 8.9 Hz, 2H), 6.87 – 6.75 (m, 1H), 6.57 (d, J = 15.3 Hz, 1H), 5.21 (s, 2H), 4.65 (t, J = 4.7 Hz, 2H), 4.04 (t, J = 4.6 Hz, 2H), 3.08 (d, J = 5.8 Hz, 2H), 2.17 (s, 6H).
64	VIII-24		(E)-4-(dimethylamino)-1-(10-((4-(thiophen-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one	502	(400 MHz, DMSO-d ₆) δ 9.77 (s, 1H), 8.41 (s, 1H), 7.80 (s, 1H), 7.72 – 7.64 (m, 2H), 7.59 – 7.53 (m, 1H), 7.28 (d, J = 9.0 Hz, 1H), 7.26 – 7.21 (m, 1H), 7.11 – 7.01 (m, 3H), 6.86 – 6.76 (m, 1H), 6.57 (d, J = 15.1 Hz, 1H), 5.32 (s, 2H), 4.69 – 4.62 (m, 2H), 4.09 – 4.02 (m, 2H), 3.11 – 3.04 (m, 2H), 2.17 (s, 6H).
65	VIII-25		(E)-4-(dimethylamino)-1-(10-((4-(thiazol-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one	503	(400 MHz, DMSO-d ₆) δ 9.78 (s, 1H), 8.41 (s, 1H), 7.86 – 7.79 (m, 3H), 7.74 – 7.65 (m, 2H),




			f]quinazolin-4-yl)but-2-en-1-one		7.28 (d, J = 9.0 Hz, 1H), 7.14 – 7.07 (m, 2H), 6.87 – 6.75 (m, 1H), 6.57 (d, J = 15.3 Hz, 1H), 5.47 (s, 2H), 4.66 (t, J = 4.7 Hz, 2H), 4.05 (t, J = 4.7 Hz, 2H), 3.07 (d, J = 5.5 Hz, 2H), 2.17 (s, 6H).
66	VIII-26		(E)-4-(dimethylamino)-1-(10-((4-(benzylthio)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one	512	(400 MHz, DMSO-d ₆) δ 9.89 (s, 1H), 8.48 (s, 1H), 7.91 – 7.76 (m, 3H), 7.42 – 7.20 (m, 8H), 6.89 – 6.75 (m, 1H), 6.58 (d, J = 15.3 Hz, 1H), 4.67 (t, J = 4.6 Hz, 2H), 4.23 (s, 2H), 4.05 (t, J = 4.6 Hz, 2H), 3.12 – 3.00 (m, 2H), 2.18 (s, 6H).
67	VIII-3		(E)-4-(dimethylamino)-1-(10-((4-((3-(trifluoromethyl)phenoxy)methyl)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one	582	(400 MHz, DMSO-d ₆) δ 9.87 (s, 1H), 8.44 (s, 1H), 8.08 (d, J = 2.6 Hz, 1H), 7.99 – 7.93 (m, 1H), 7.84 (s, 1H), 7.54 – 7.45 (m, 1H), 7.37 (d, J = 9.1 Hz, 1H), 7.34 – 7.24 (m, 3H), 7.22 – 7.15 (m, 1H), 6.86 – 6.78 (m, 1H), 6.58 (d, J = 15.2 Hz, 1H), 5.34 (s, 2H), 4.69 – 4.62 (m, 2H), 4.09 – 4.02 (m, 2H), 3.13 – 3.04 (m, 2H), 2.18 (s, 6H).
68	VIII-27		(E)-4-(dimethylamino)-1-(10-((4-((3-(fluorobenzyl)oxy)-3-methoxyphenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one	544	(400 MHz, DMSO-d ₆) δ 9.76 (s, 1H), 8.43 (s, 1H), 7.81 (s, 1H), 7.50 – 7.40 (m, 2H), 7.35 – 7.25 (m, 4H), 7.21 – 7.11 (m, 1H), 7.04 (d, J = 8.8 Hz,



					1H), 6.87 – 6.77 (m, 1H), 6.58 (d, J = 15.3 Hz, 1H), 5.14 (s, 2H), 4.67 (t, J = 4.5 Hz, 2H), 4.09 – 4.02 (m, 2H), 3.83 (s, 3H), 3.15 – 3.03 (m, 2H), 2.18 (s, 6H).
69	VIII-28		(E)-4-(dimethylamino)-1-(10-((4-((3-fluorobenzyl)oxy)-3-fluorophenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one	532	(400 MHz, DMSO-d ₆) δ 9.83 (s, 1H), 8.48 (s, 1H), 7.96 – 7.77 (m, 2H), 7.54 – 7.40 (m, 2H), 7.37 – 7.15 (m, 5H), 6.89 – 6.77 (m, 1H), 6.58 (d, J = 15.4 Hz, 1H), 5.23 (s, 2H), 4.73 – 4.61 (m, 2H), 4.10 – 4.00 (m, 2H), 3.14 – 3.00 (m, 2H), 2.17 (s, 6H).
70	VIII-29		(E)-4-(dimethylamino)-1-(10-((4-((4-fluorophenyl)thio)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one	516	(400 MHz, DMSO-d ₆) δ 9.97 (s, 1H), 8.51 (s, 1H), 7.96 – 7.82 (m, 3H), 7.43 – 7.32 (m, 5H), 7.27 – 7.20 (m, 2H), 6.86 – 6.78 (m, 1H), 6.58 (d, J = 15.2 Hz, 1H), 4.71 – 4.64 (m, 2H), 4.09 – 4.04 (m, 2H), 3.12 – 3.06 (m, 2H), 2.18 (s, 6H).
71	VIII-30		(E)-4-(dimethylamino)-1-(10-((4-(2-fluoro-5-methylphenoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one	514	(400 MHz, DMSO-d ₆) δ 9.96 – 9.72 (m, 1H), 8.43 (s, 1H), 7.98 – 7.63 (m, 3H), 7.37 – 7.15 (m, 2H), 7.10 – 6.92 (m, 4H), 6.88 – 6.70 (m, 1H), 6.67 – 6.47 (m, 1H), 4.83 – 4.47 (m, 2H), 4.14 – 3.87 (m, 2H), 3.15 – 2.94 (m, 2H), 2.35 – 2.24




					(m, 3H), 2.22 – 2.00 (s, 6H).
72	VIII-31		(E)-4-(dimethylamino)-1-(10-((4-(5-chloro-2-fluorophenoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one	534	(400 MHz, DMSO-d ₆) δ 9.88 (s, 1H), 8.45 (s, 1H), 7.96 – 7.67 (m, 3H), 7.47 (dd, J = 10.8, 8.8 Hz, 1H), 7.33 – 7.24 (m, 2H), 7.22 – 7.05 (m, 3H), 6.87 – 6.73 (m, 1H), 6.58 (d, J = 15.2 Hz, 1H), 4.66 (t, J = 4.6 Hz, 2H), 4.05 (t, J = 4.9 Hz, 2H), 3.07 (d, J = 5.8 Hz, 2H), 2.17 (s, 6H).
73	VIII-32		(E)-4-(dimethylamino)-1-(10-((4-(2,5-difluorophenoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one	518	(400 MHz, DMSO-d ₆) δ 9.83 (s, 1H), 8.43 (s, 1H), 8.02 – 7.61 (m, 3H), 7.57 – 7.41 (m, 1H), 7.38 – 7.21 (m, 2H), 7.18 – 7.08 (m, 1H), 7.02 (d, J = 8.6 Hz, 2H), 6.87 – 6.72 (m, 1H), 6.57 (d, J = 15.4 Hz, 1H), 4.65 (t, J = 4.6 Hz, 2H), 4.04 (t, J = 4.7 Hz, 2H), 3.07 (d, J = 5.8 Hz, 2H), 2.17 (s, 6H).
74	VIII-33		(E)-4-(dimethylamino)-1-(10-((4-(1-(3-fluorophenyl)ethoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one	528	(400 MHz, DMSO-d ₆) δ 9.70 (s, 1H), 8.37 (s, 1H), 7.77 (s, 1H), 7.55 (d, J = 8.6 Hz, 2H), 7.47 – 7.34 (m, 1H), 7.30 – 7.14 (m, 3H), 7.14 – 7.00 (m, 1H), 6.94 (d, J = 8.6 Hz, 2H), 6.85 – 6.71 (m, 1H), 6.56 (d, J = 15.4 Hz, 1H), 5.61 – 5.39 (m, 1H), 4.62 (t, J = 4.7 Hz, 2H), 4.02 (t, J = 4.7 Hz, 2H), 3.07 (d, J = 5.8 Hz,

					2H), 2.16 (s, 6H), 1.56 (d, J = 6.3 Hz, 3H).
75	VIII-34		(E)-4-(dimethylamino)-1-(10-((4-(1-(pyridin-2-yl)ethoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one	511	(400 MHz, DMSO-d ₆) δ 9.73 (d, J = 19.8 Hz, 1H), 8.66 – 8.51 (m, 1H), 8.44 – 8.29 (m, 1H), 7.88 – 7.74 (m, 2H), 7.63 – 7.51 (m, 2H), 7.45 (d, J = 7.4 Hz, 1H), 7.40 – 7.22 (m, 2H), 7.01 – 6.88 (m, 2H), 6.88 – 6.70 (m, 1H), 6.55 (d, J = 18.1 Hz, 1H), 5.62 – 5.35 (m, 1H), 4.73 – 4.54 (m, 2H), 4.11 – 3.92 (m, 2H), 3.41 – 3.33 (m, 3H), 3.06 (d, J = 5.8 Hz, 2H), 2.16 (s, 6H).
76	VIII-35		(E)-4-(dimethylamino)-1-(10-((4-(pyridin-3-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one	497	(400 MHz, DMSO-d ₆) δ 9.76 (s, 1H), 8.69 (d, J = 2.5 Hz, 1H), 8.59 – 8.51 (m, 1H), 8.40 (s, 1H), 7.93 – 7.86 (m, 1H), 7.84 – 7.70 (m, 1H), 7.70 – 7.60 (m, 2H), 7.47 – 7.38 (m, 1H), 7.32 – 7.22 (m, 1H), 7.10 – 7.00 (m, 2H), 6.84 – 6.74 (m, 1H), 6.57 (d, J = 15.2 Hz, 1H), 5.18 (s, 2H), 4.90 – 4.42 (m, 2H), 4.04 (t, J = 4.7 Hz, 2H), 3.09 (d, J = 5.8 Hz, 2H), 2.23 – 2.02 (s, 6H).

77	VIII-36		(E)-4-(dimethylamino)-1-(10-((4-(pyridin-4-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one	497	(400 MHz, DMSO-d ₆) δ 9.76 (s, 1H), 8.59 (d, J = 5.1 Hz, 2H), 8.40 (s, 1H), 7.80 (s, 1H), 7.67 (d, J = 8.6 Hz, 2H), 7.46 (d, J = 5.0 Hz, 2H), 7.27 (d, J = 9.0 Hz, 1H), 7.06 (d, J = 8.6 Hz, 2H), 6.90 – 6.73 (m, 1H), 6.57 (d, J = 15.3 Hz, 1H), 5.22 (s, 2H), 4.65 (t, J = 4.6 Hz, 2H), 4.04 (t, J = 4.6 Hz, 2H), 3.08 (d, J = 6.1 Hz, 2H), 2.17 (s, 6H).
78	VIII-41		(E)-1-(10-((4-(benzyloxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(dimethylamino)but-2-en-1-one	496	(400 MHz, DMSO-d ₆) δ 9.75 (s, 1H), 8.40 (s, 1H), 7.80 (s, 1H), 7.67 – 7.63 (m, 2H), 7.49 – 7.45 (m, 2H), 7.43 – 7.38 (m, 2H), 7.36 – 7.33 (m, 1H), 7.27 (d, J = 9.2 Hz, 1H), 7.07 – 7.03 (m, 2H), 6.84 – 6.77 (m, 1H), 6.62 – 6.53 (m, 1H), 5.13 (s, 2H), 4.65 (t, J = 4.4 Hz, 2H), 4.04 (t, J = 4.4 Hz, 2H), 3.09 (d, J = 5.6 Hz, 2H), 2.18 (s, 6H).
79	VIII-42		(E)-4-(dimethylamino)-1-(10-((4-((2-fluorobenzyl)oxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one	514	(400 MHz, DMSO-d ₆) δ 9.77 (s, 1H), 8.40 (s, 1H), 7.80 (s, 1H), 7.69 – 7.65 (m, 2H), 7.60 – 7.56 (m, 1H), 7.46 – 7.40 (m, 1H), 7.29 – 7.26 (m, 2H), 7.25 – 7.23 (m, 1H), 7.08 – 7.05 (m, 2H), 6.85 – 6.77 (m, 1H), 6.60 – 6.53 (m, 1H), 5.17 (s, 2H),

					4.65 (t, J = 4.4 Hz, 2H), 4.04 (t, J = 4.4 Hz, 2H), 3.07 (dd, J = 6.0, 1.6 Hz, 2H), 2.17 (s, 6H).
80	VIII-43		(E)-4-(dimethylamino)-1-(10-((4-((2-chlorobenzyl)oxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one	530	(400 MHz, DMSO-d ₆) δ 9.77 (s, 1H), 8.40 (s, 1H), 7.80 (s, 1H), 7.68 – 7.66 (m, 2H), 7.64 – 7.60 (m, 1H), 7.54 – 7.51 (m, 1H), 7.42 – 7.39 (m, 2H), 7.28 (d, J = 9.0 Hz, 1H), 7.08 – 7.05 (m, 2H), 6.85 – 6.77 (m, 1H), 6.57 (d, J = 15.2 Hz, 1H), 5.19 (s, 2H), 4.65 (t, J = 4.6 Hz, 2H), 4.04 (t, J = 4.6 Hz, 2H), 3.07 (dd, J = 6.0, 1.6 Hz, 2H), 2.17 (s, 6H).
81	VIII-44		(E)-4-(dimethylamino)-1-(10-((4-((2-methylbenzyl)oxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one	510	(400 MHz, DMSO-d ₆) δ 9.76 (s, 1H), 8.40 (s, 1H), 7.79 (s, 1H), 7.67 – 7.63 (m, 2H), 7.43 (d, J = 7.2 Hz, 1H), 7.29 – 7.21 (m, 4H), 7.09 – 7.05 (m, 2H), 6.84 – 6.77 (m, 1H), 6.57 (d, J = 15.2 Hz, 1H), 5.11 (s, 2H), 4.65 (t, J = 4.4 Hz, 2H), 4.04 (t, J = 4.4 Hz, 2H), 3.07 (dd, J = 6.0, 1.6 Hz, 2H), 2.35 (s, 3H), 2.17 (s, 6H).
82	VIII-45		(E)-4-(dimethylamino)-1-(10-((4-((2-methoxybenzyl)oxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one	526	(400 MHz, DMSO-d ₆) δ 9.76 (s, 1H), 8.40 (s, 1H), 7.80 (s, 1H), 7.66 – 7.63 (m, 2H), 7.41 (dd, J = 7.2, 1.6 Hz, 1H), 7.37 – 7.32 (m, 1H), 7.27 (d, J = 9.0 Hz, 1H), 7.08 –


					7.05 (m, 1H), 7.04 – 7.01 (m, 2H), 7.00 – 6.95 (m, 1H), 6.85 – 6.77 (m, 1H), 6.57 (d, J = 15.2 Hz, 1H), 5.08 (s, 2H), 4.65 (t, J = 4.6 Hz, 2H), 4.04 (t, J = 4.6 Hz, 2H), 3.84 (s, 3H), 3.07 (dd, J = 6.0, 1.6 Hz, 2H), 2.17 (s, 6H).
83	VIII-46		(E)-1-(10-((4-((3-chlorobenzyl)oxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(dimethylamino)but-2-en-1-one	530	(400 MHz, DMSO-d ₆) δ 9.76 (s, 1H), 8.40 (s, 1H), 7.79 (s, 1H), 7.66 (d, J = 9.0 Hz, 2H), 7.57 – 7.48 (m, 1H), 7.48 – 7.34 (m, 3H), 7.27 (d, J = 9.0 Hz, 1H), 7.07 – 6.99 (m, 2H), 6.88 – 6.70 (m, 1H), 6.57 (d, J = 15.2 Hz, 1H), 5.16 (s, 2H), 4.65 (t, J = 4.7 Hz, 2H), 4.04 (t, J = 4.6 Hz, 2H), 3.14 – 2.97 (m, 2H), 2.17 (s, 6H).
84	VIII-47		(E)-4-(dimethylamino)-1-(10-((4-((3-methylbenzyl)oxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one	510	(400 MHz, DMSO-d ₆) δ 9.75 (s, 1H), 8.40 (s, 1H), 7.80 (s, 1H), 7.66 – 7.63 (m, 2H), 7.29 – 7.25 (m, 4H), 7.16 – 7.13 (m, 1H), 7.05 – 7.02 (m, 2H), 6.84 – 6.77 (m, 1H), 6.57 (d, J = 15.2 Hz, 1H), 5.09 (s, 2H), 4.65 (t, J = 4.4 Hz, 2H), 4.04 (t, J = 4.4 Hz, 2H), 3.07 (dd, J = 6.0, 1.6 Hz, 2H), 2.33 (s, 3H), 2.17 (s, 6H).

85	VIII-48		(E)-4-(dimethylamino)-1-(10-((4-((3-cyanobenzyl)oxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one	521	(400 MHz, DMSO-d ₆) δ 9.76 (s, 1H), 8.40 (s, 1H), 7.94 – 7.92 (m, 1H), 7.86 – 7.78 (m, 3H), 7.68 – 7.65 (m, 2H), 7.65 – 7.61 (m, 1H), 7.28 (d, J = 9.0 Hz, 1H), 7.08 – 7.05 (m, 2H), 6.85 – 6.78 (m, 1H), 6.57 (d, J = 15.2 Hz, 1H), 5.20 (s, 2H), 4.65 (t, J = 4.6 Hz, 2H), 4.04 (t, J = 4.6 Hz, 2H), 3.07 (dd, J = 6.0, 1.6 Hz, 2H), 2.17 (s, 6H).
86	VIII-49		(E)-4-(dimethylamino)-1-(10-((4-((3-methoxybenzyl)oxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one	526	(400 MHz, DMSO-d ₆) δ 9.75 (s, 1H), 8.40 (s, 1H), 7.79 (s, 1H), 7.67 – 7.57 (m, 2H), 7.38 – 7.19 (m, 2H), 7.10 – 6.95 (m, 4H), 6.94 – 6.74 (m, 2H), 6.57 (d, J = 15.3 Hz, 1H), 5.11 (s, 2H), 4.68 – 4.57 (m, 2H), 4.04 (t, J = 4.7 Hz, 2H), 3.77 (s, 3H), 3.10 – 2.99 (m, 2H), 2.17 (s, 6H).
87	VIII-50		(E)-4-(dimethylamino)-1-(10-((4-((4-chlorobenzyl)oxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one	530	(400 MHz, DMSO-d ₆) δ 9.76 (s, 1H), 8.40 (s, 1H), 7.80 (s, 1H), 7.67 – 7.64 (m, 2H), 7.51 – 7.49 (m, 2H), 7.47 – 7.45 (m, 2H), 7.27 (d, J = 9.0 Hz, 1H), 7.06 – 7.02 (m, 2H), 6.84 – 6.77 (m, 1H), 6.57 (d, J = 15.2 Hz, 1H), 5.14 (s, 2H), 4.65 (t, J = 4.6 Hz, 2H), 4.04 (t, J = 4.6 Hz, 2H), 3.07 (dd, J

					= 6.0, 1.6 Hz, 2H), 2.17 (s, 6H).
88	VIII-51		(E)-4-(dimethylamino)-1-(10-((4-((4-methylbenzyl)oxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one	510	(400 MHz, DMSO-d ₆) δ 9.75 (s, 1H), 8.39 (s, 1H), 7.79 (s, 1H), 7.65 – 7.62 (m, 2H), 7.35 (d, J = 7.6 Hz, 2H), 7.27 (d, J = 9.2 Hz, 1H), 7.20 (d, J = 7.6 Hz, 2H), 7.04 – 7.01 (m, 2H), 6.84 – 6.77 (m, 1H), 6.57 (d, J = 15.2 Hz, 1H), 5.08 (s, 2H), 4.65 (t, J = 4.4 Hz, 2H), 4.04 (t, J = 4.4 Hz, 2H), 3.07 (dd, J = 6.0, 1.6 Hz, 2H), 2.31 (s, 3H), 2.17 (s, 6H).
89	VIII-52		(E)-1-(10-((4-((2,5-difluorobenzyl)oxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(dimethylamino)but-2-en-1-one	532	(400 MHz, DMSO-d ₆) δ 9.78 (s, 1H), 8.41 (s, 1H), 7.80 (s, 1H), 7.72 – 7.64 (m, 2H), 7.47 – 7.37 (m, 1H), 7.38 – 7.20 (m, 3H), 7.12 – 7.02 (m, 2H), 6.88 – 6.75 (m, 1H), 6.57 (d, J = 15.3 Hz, 1H), 5.16 (s, 2H), 4.65 (t, J = 4.7 Hz, 2H), 4.04 (t, J = 4.7 Hz, 2H), 3.13 – 3.02 (m, 2H), 2.17 (s, 6H).
90	VIII-53		(E)-1-(10-((4-((2-chloro-5-fluorobenzyl)oxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(dimethylamino)but-2-en-1-one	548	(400 MHz, DMSO-d ₆) δ 9.78 (s, 1H), 8.41 (s, 1H), 7.80 (s, 1H), 7.73 – 7.62 (m, 2H), 7.63 – 7.53 (m, 1H), 7.53 – 7.41 (m, 1H), 7.36 – 7.17 (m, 2H), 7.14 – 7.02 (m, 2H), 6.88 – 6.73 (m, 1H), 6.57 (d, J = 15.3 Hz, 1H), 5.18 (s, 2H),


					4.65 (t, J = 4.7 Hz, 2H), 4.04 (t, J = 4.7 Hz, 2H), 3.13 – 3.00 (m, 2H), 2.18 (s, 6H).
--	--	--	--	--	--

[00269] Example 91: Preparation of (E)-1-(10-((4-phenoxyphenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(pyrrolidin-1-yl)but-2-en-1-one



[00270] N-(4-phenoxyphenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine (VIII-1) (185 mg, 0.5 mmol) was dissolved in a mixed solvent of dichloromethane and dimethylformamide, to which 4-bromocrotonyl chloride (91 mg, 0.5 mmol) was added, and stirred at room temperature until the reaction was completed. The reaction was quenched with water, extracted with ethyl acetate, the organic phase was concentrated and directly dissolved in acetonitrile, to which diisopropylethylamine (129 mg, 1 mmol) and pyrrolidine (67 mg, 1 mmol) were added, and stirred at room temperature until the reaction was completed. The reaction was quenched by adding water, extracted with ethyl acetate, and the organic phase was concentrated and purified by silica gel column chromatography to afford 63 mg of a off-white solid with a yield of 25%. ¹H NMR (400 MHz, DMSO-d₆) δ 9.87 (s, 1H), 8.46 (s, 1H), 7.84 – 7.78 (m, 3H), 7.44 – 7.37 (m, 2H), 7.32 – 7.28 (m, 1H), 7.16 – 7.11 (m, 1H), 7.10 – 7.06 (m, 2H), 7.05 – 7.00 (m, 2H), 6.90 – 6.82 (m, 1H), 6.62 – 6.55 (m, 1H), 4.67 (t, J = 4.4 Hz, 2H), 4.05 (t, J = 4.4 Hz, 2H), 3.27 – 3.23 (m, 2H), 2.49 – 2.45 (m, 4H), 1.72 – 1.67 (m, 4H); MS: 508[M+H]⁺.

[00271] Examples 92-111














[00272] With reference to the preparation method of Example 91, wherein exactly the same operations were used, and N-(4-phenoxyphenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-







f]quinazolin-10-amine (VIII-1) was replaced with the same molar equivalent of intermediate represented by formula (VIII) wherein R¹X is the substituent in the table below and pyrrolidine





was replaced with equivalent molar equivalent of HN-R^{12}
 R^{11} . The specific example compounds are shown in the table below:




5





Example No.	Starting Intermediate No.	R ₁ X		Example compound		
				Name	LC MS m/z = (M + H) ⁺	HNMR
92	VIII-1			(E)-4-(diethylamino)-1-(10-((4-phenoxyphenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one	510	(400 M, CDCl ₃) δ 9.55(s, 1H), 8.62(s, 1H), 7.70-7.68(m, 2H), 7.61(brs, 1H), 7.47-7.45(m, 1H), 7.38-7.35(m, 2H), 7.15-7.05(m, 6H), 6.72-6.64(m, 1H), 4.70(s, 2H), 4.22(s, 2H), 3.64(br, 2H), 3.02-2.98(m, 2H), 1.66(br, 2H), 1.31-1.27(m, 6H)
93	VIII-1			(E)-1-(10-((4-phenoxyphenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(piperidin-1-yl)but-2-en-1-one	522	(400 M, DMSO-d ₆) δ 9.87(s, 1H), 8.45(s, 1H), 7.81-7.79(m, 3H), 7.42-7.38(m, 2H), 7.30-7.28(m, 1H), 7.15-7.01(m, 5H), 6.84-6.80(m, 1H), 6.60-6.56(m, 1H), 4.69-4.63(m, 2H), 4.09-4.03(m, 2H), 3.14-3.08(m, 2H), 2.39-2.33(m, 4H), 1.49-1.47(m, 4H), 1.42-1.36(m, 2H)
94	VIII-1			(E)-4-morpholino-1-(10-((4-phenoxyphenyl)amino)-2,3-	524	(300 MHz, DMSO-d ₆) δ 9.87(s, 1H), 8.45(s, 1H), 7.88 – 7.73(m, 3H), 7.44 –





				dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one		7.36 (m, 2H), 7.30 (d, J = 9.0 Hz, 1H), 7.16 – 7.00 (m, 5H), 6.79 (s, 1H), 6.68 – 6.53 (m, 1H), 4.71 – 4.58 (m, 2H), 4.09 – 4.03 (m, 2H), 3.57 (t, J = 4.6 Hz, 4H), 3.14 (d, J = 5.5 Hz, 2H), 2.43 – 2.34 (m, 4H).
95	VIII-1			(E)-4-(4-methylpiperazin-1-yl)-1-(10-((4-phenoxyphenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one	537	(400 MHz, DMSO-d₆) δ 9.87 (s, 1H), 8.46 (s, 1H), 7.86 – 7.72 (m, 3H), 7.42 – 7.38 (m, 2H), 7.30 (d, J = 8.9 Hz, 1H), 7.13 (t, J = 7.4 Hz, 1H), 7.10 – 7.06 (m, 2H), 7.05 – 7.00 (m, 2H), 6.85 – 6.76 (m, 1H), 6.59 (d, J = 15.2 Hz, 1H), 4.67 (t, J = 4.6 Hz, 2H), 4.05 (t, J = 4.6 Hz, 2H), 3.14 (d, J = 5.7 Hz, 2H), 2.49 – 2.22 (m, 8H), 2.17 (s, 3H).
96	VIII-4			(E)-1-(10-((3-chloro-4-((3-fluorobenzyl)oxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-morpholinobut-2-en-1-one	590	(400 MHz, DMSO-d₆) δ 9.82 (s, 1H), 8.47 (s, 1H), 8.00 (d, J = 2.4 Hz, 1H), 7.90 – 7.74 (m, 1H), 7.70 – 7.65 (m, 1H), 7.52 – 7.44 (m, 1H), 7.35 – 7.24 (m, 4H), 7.22 – 7.15 (m, 1H), 6.85 – 6.77 (m, 1H), 6.66 – 6.56 (m, 1H), 5.27 (s, 2H), 4.66 (t, J = 4.4 Hz, 2H), 4.04 (t, J = 4.4 Hz, 2H), 3.61 – 3.55 (m, 4H), 3.17 – 3.11 (m, 2H),





						2.44 – 2.36 (m, 4H)
97	VIII-6			(E)-1-(10-((2-fluoro-4-phenoxyphenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(piperidin-1-yl)but-2-en-1-one	540	(400 MHz, Methanol-d ₄) δ 8.77 – 8.65 (m, 1H), 8.48 – 8.41 (m, 1H), 7.69 – 7.47 (m, 1H), 7.38 – 7.29 (m, 2H), 7.28 – 7.13 (m, 2H), 7.06 – 6.95 (m, 2H), 6.94 – 6.86 (m, 1H), 6.86 – 6.71 (m, 2H), 6.68 – 6.61 (m, 1H), 5.39 (s, 2H), 4.38 – 4.25 (m, 2H), 3.76 – 3.67 (m, 2H), 3.08 – 2.91 (m, 4H), 1.64 – 1.45 (m, 6H).
98	VIII-7			(E)-1-(10-((3-fluoro-4-phenoxyphenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(piperidin-1-yl)but-2-en-1-one	540	(400 MHz, Methanol-d ₄) δ 8.51 (s, 1H), 8.08 – 7.99 (m, 1H), 7.93 – 7.79 (m, 1H), 7.51 – 7.45 (m, 1H), 7.40 – 7.30 (m, 3H), 7.17 – 6.97 (m, 4H), 6.94 – 6.87 (m, 1H), 6.66 (d, J = 15.2 Hz, 1H), 4.76 – 4.65 (m, 2H), 4.21 – 4.13 (m, 2H), 3.25 – 3.19 (m, 2H), 2.58 – 2.42 (m, 4H), 1.68 – 1.60 (m, 4H), 1.55 – 1.47 (m, 2H).
99	VIII-8			(E)-1-(10-((4-fluorophenoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-((4-methoxybutyl)amino)but-2-en-1-one	558	(400 MHz, Methanol-d ₄) δ 8.45 – 8.40 (m, 1H), 7.77 – 7.69 (m, 3H), 7.40 – 7.34 (m, 2H), 7.30 – 7.26 (m, 1H), 7.23 – 7.19 (m, 2H), 7.07 – 7.02 (m, 2H), 7.02 – 6.94 (m, 1H), 6.82 – 6.74 (m, 1H),



						4.84 – 4.69 (m, 2H), 4.19 – 4.14 (m, 2H), 3.70 (s, 3H), 3.19 – 3.12 (m, 2H), 2.94 – 2.83 (m, 2H), 2.32 – 2.14 (m, 2H), 1.69 – 1.66 (m, 4H).
100	VIII-5			(E)-1-(10-((3-chloro-4-(pyridin-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(cyclopropyl(methyl)amino)but-2-en-1-one	557	(400 MHz, DMSO-d ₆) δ 9.82 (s, 1H), 8.64 – 8.55 (m, 1H), 8.46 (s, 1H), 8.01 (d, J = 2.6 Hz, 1H), 7.95 – 7.73 (m, 2H), 7.70 – 7.62 (m, 1H), 7.58 (d, J = 7.8 Hz, 1H), 7.42 – 7.34 (m, 1H), 7.32 – 7.16 (m, 2H), 6.89 – 6.78 (m, 1H), 6.55 (d, J = 15.2 Hz, 1H), 5.31 (s, 2H), 4.65 (t, J = 4.6 Hz, 2H), 4.04 (t, J = 4.7 Hz, 2H), 3.32 – 3.29 (m, 2H), 2.27 (s, 3H), 1.78 – 1.72 (m, 1H), 0.49 – 0.23 (m, 4H).
101	VIII-5			(E)-1-(10-((3-chloro-4-(pyridin-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(cyclopropylamino)but-2-en-1-one	543	(400 MHz, DMSO-d ₆) δ 9.82 (s, 1H), 8.60 (d, J = 4.8 Hz, 1H), 8.46 (s, 1H), 8.01 (d, J = 2.6 Hz, 1H), 7.90 – 7.87 (m, 1H), 7.66 (d, J = 6.2 Hz, 1H), 7.58 (d, J = 7.8 Hz, 1H), 7.50 – 7.48 (m, 1H), 7.39 – 7.36 (m, 1H), 7.31 – 7.28 (m, 1H), 7.27 – 7.25 (m, 1H), 7.08 (d, J = 7.8 Hz, 1H), 6.57 (d, J = 14.8 Hz, 1H), 5.31 (s, 2H), 4.64 (t, J = 4.6 Hz, 2H), 4.05 (t, J = 4.1

						Hz, 2H), 3.51 – 3.50 (m, 1H), 3.42 – 3.41 (m, 2H), 2.68 – 2.66 (m, 1H), 0.41 – 0.37 (m, 2H), 0.27 – 0.23 (m, 2H).
102	VIII-5			(E)-1-(10-((3-chloro-4-(pyridin-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(isopropylamino)butan-2-one	545	(400 MHz, DMSO-d ₆) δ 9.83 (s, 1H), 8.60 (d, J = 4.8 Hz, 1H), 8.47 (s, 1H), 8.01 (d, J = 2.8 Hz, 1H), 7.92 – 7.80 (m, 2H), 7.67 (dd, J = 8.8, 2.8 Hz, 1H), 7.58 (d, J = 7.8 Hz, 1H), 7.39 – 7.35 (m, 1H), 7.33 – 7.25 (m, 2H), 6.89 – 6.80 (m, 1H), 6.80 – 6.68 (m, 1H), 5.31 (s, 2H), 4.67 (t, J = 4.6 Hz, 2H), 4.07 (t, J = 4.6 Hz, 2H), 3.68 – 3.62 (m, 2H), 3.31 – 3.31 (m, 1H), 2.68 – 2.66 (m, 1H), 1.14 (d, J = 6.4 Hz, 6H).
103	VIII-5			(E)-1-(10-((3-chloro-4-(pyridin-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-morpholinobutan-2-one	573	(400 MHz, DMSO-d ₆) δ 9.83 (s, 1H), 8.62 – 8.58 (m, 1H), 8.47 (s, 1H), 8.00 (d, J = 2.6 Hz, 1H), 7.93 – 7.78 (m, 2H), 7.68 – 7.63 (m, 1H), 7.61 – 7.56 (m, 1H), 7.41 – 7.35 (m, 1H), 7.33 – 7.24 (m, 2H), 6.86 – 6.77 (m, 1H), 6.72 – 6.60 (m, 1H), 5.31 (s, 2H), 4.66 (t, J = 4.6 Hz, 2H), 4.05 (t, J = 4.6 Hz, 2H), 3.70 – 3.60 (m, 4H), 3.18 – 3.12 (m, 2H), 2.66 – 2.51 (m,

						4H).
104	VIII-5			(E)-1-(10-((3-chloro-4-(pyridin-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(4-methylpiperazin-1-yl)but-2-en-1-one	586	(400 MHz, DMSO-d ₆) δ 9.82 (s, 1H), 8.64 – 8.56 (m, 1H), 8.46 (s, 1H), 8.01 (d, J = 2.6 Hz, 1H), 7.93 – 7.86 (m, 1H), 7.86 – 7.68 (m, 1H), 7.68 – 7.63 (m, 1H), 7.58 (d, J = 7.8 Hz, 1H), 7.43 – 7.33 (m, 1H), 7.33 – 7.23 (m, 2H), 6.87 – 6.75 (m, 1H), 6.70 – 6.52 (m, 1H), 5.31 (s, 2H), 4.66 (t, J = 4.5 Hz, 2H), 4.04 (t, J = 4.6 Hz, 2H), 3.14 (d, J = 5.7 Hz, 2H), 2.52 – 2.30 (m, 8H), 2.22 (s, 3H).
105	VIII-5			(E)-1-(10-((3-chloro-4-(pyridin-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(diethylamino)but-2-en-1-one	559	(400 MHz, DMSO-d ₆) δ 9.84 (s, 1H), 8.63 – 8.59 (m, 1H), 8.47 (s, 1H), 8.01 (d, J = 2.6 Hz, 1H), 7.93 – 7.86 (m, 1H), 7.79 – 7.61 (m, 2H), 7.58 (d, J = 7.8 Hz, 1H), 7.41 – 7.35 (m, 1H), 7.34 – 7.24 (m, 2H), 6.90 – 6.61 (m, 2H), 5.32 (s, 2H), 4.68 (t, J = 4.7 Hz, 2H), 4.06 (t, J = 4.7 Hz, 2H), 3.39 – 3.31 (m, 6H), 1.18 – 0.97 (m, 6H).
106	VIII-5			(E)-1-(10-((3-chloro-4-(pyridin-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-	557	(400 MHz, DMSO-d ₆) δ 9.83 (s, 1H), 8.61 (d, J = 4.8 Hz, 1H), 8.47 (s, 1H), 8.03 – 8.00 (m, 1H), 7.92 – 7.84 (m, 2H), 7.69 – 7.65 (m, 1H), 7.60 – 7.57 (m,


				4-yl)-4-(pyrrolidin-1-yl)but-2-en-1-one		1H), 7.39 – 7.35 (m, 1H), 7.32 – 7.25 (m, 2H), 6.88 – 6.82 (m, 1H), 6.61 – 6.56 (m, 1H), 5.32 (s, 2H), 4.66 (t, J = 4.7 Hz, 2H), 4.05 (t, 2H), 3.36 – 3.30 (m, 2H), 2.61 – 2.53 (m, 4H), 1.78 – 1.68 (m, 4H).
107	VIII-5			(E)-1-(10-((3-chloro-4-(pyridin-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(piperidin-1-yl)but-2-en-1-one	571	(400 MHz, DMSO-d ₆) δ 9.83 (s, 1H), 8.63 – 8.56 (m, 1H), 8.47 (s, 1H), 8.01 (d, J = 2.6 Hz, 1H), 7.92 – 7.71 (m, 2H), 7.72 – 7.65 (m, 1H), 7.59 (d, J = 7.8 Hz, 1H), 7.42 – 7.34 (m, 1H), 7.34 – 7.23 (m, 2H), 6.92 – 6.61 (m, 2H), 5.31 (s, 2H), 4.68 (t, J = 4.7 Hz, 2H), 4.07 (t, J = 4.6 Hz, 2H), 3.36 – 3.30 (m, 4H), 1.86 – 1.30 (m, 8H).
108	VIII-5			(E)-4-(4-acetylpiperazin-1-yl)-1-(10-((3-chloro-4-(pyridin-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl) but-2-en-1-one	614	(400 MHz, DMSO-d ₆) δ 9.86 (s, 1H), 8.65 – 8.56 (m, 1H), 8.47 (s, 1H), 8.00 (s, 1H), 7.96 – 7.72 (m, 2H), 7.71 – 7.63 (m, 1H), 7.58 (d, J = 7.8 Hz, 1H), 7.41 – 7.35 (m, 1H), 7.35 – 7.23 (m, 2H), 6.88 – 6.77 (m, 1H), 6.75 – 6.58 (m, 1H), 5.31 (s, 2H), 4.67 (t, J = 4.7 Hz, 2H), 4.05 (t, J = 4.6 Hz, 2H), 3.52 – 3.35 (m, 6H), 2.79 – 2.51 (m, 4H), 2.00 (s,

						3H).
109	VIII-5			(E)-1-(10-((3-chloro-4-(pyridin-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(cyclobutylamino)but-2-en-1-one	557	(400 MHz, DMSO-d ₆) δ 9.83 (s, 1H), 8.63 – 8.59 (m, 1H), 8.50 – 8.45 (m, 1H), 8.01 (d, J = 2.6 Hz, 1H), 7.93 – 7.82 (m, 2H), 7.71 – 7.65 (m, 1H), 7.58 (d, J = 7.8 Hz, 1H), 7.41 – 7.35 (m, 1H), 7.33 – 7.24 (m, 2H), 6.91 – 6.83 (m, 1H), 6.59 (d, J = 13.9 Hz, 1H), 5.32 (s, 2H), 4.65 (t, J = 4.6 Hz, 2H), 4.10 – 4.00 (m, 2H), 3.35 – 3.34 (m, 1H), 3.32 (d, J = 6.3 Hz, 2H), 3.24 – 3.20 (m, 1H), 2.13 – 2.04 (m, 2H), 1.70 (d, J = 9.1 Hz, 2H), 1.65 – 1.55 (m, 2H).
110	VIII-5			(E)-1-(10-((3-chloro-4-(pyridin-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(cyclobutyl(methyl)amino)but-2-en-1-one	571	(400 MHz, DMSO-d ₆) δ 9.83 (s, 1H), 8.64 – 8.59 (m, 1H), 8.46 (s, 1H), 8.01 (d, J = 2.5 Hz, 1H), 7.94 – 7.77 (m, 2H), 7.70 – 7.63 (m, 1H), 7.58 (d, J = 7.8 Hz, 1H), 7.41 – 7.34 (m, 1H), 7.31 – 7.24 (m, 2H), 6.87 – 6.77 (m, 1H), 6.62 – 6.52 (m, 1H), 5.31 (s, 2H), 4.66 (t, J = 4.6 Hz, 2H), 4.04 (t, J = 4.6 Hz, 2H), 3.03 (d, J = 5.9 Hz, 2H), 2.90 – 2.80 (m, 1H), 2.04 (s, 3H), 2.01 – 1.92 (m, 2H), 1.82 – 1.71 (m, 2H), 1.64 – 1.53 (m,

111	VIII-5			(E)-1-(10-((3-chloro-4-(pyridin-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(isopropyl(methyl)amino)but-2-en-1-one	559	2H). (400 MHz, DMSO-d ₆) δ 9.83 (s, 1H), 8.63 – 8.58 (m, 1H), 8.46 (s, 1H), 8.01 (d, J = 2.6 Hz, 1H), 7.92 – 7.86 (m, 1H), 7.80 (s, 1H), 7.69 – 7.64 (m, 1H), 7.58 (d, J = 7.8 Hz, 1H), 7.41 – 7.35 (m, 1H), 7.31 – 7.25 (m, 2H), 6.86 – 6.77 (m, 1H), 6.64 – 6.55 (m, 1H), 5.31 (s, 2H), 4.66 (t, J = 4.6 Hz, 2H), 4.04 (t, J = 4.7 Hz, 2H), 3.22 – 3.18 (m, 2H), 2.87 – 2.76 (m, 1H), 2.13 (s, 3H), 0.96 (d, J = 6.5 Hz, 6H).
-----	--------	---	---	---	-----	--


[00273] Example 112

[00274] Preparation of (S,E)-1-(10-((3-chloro-4-(pyridin-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-3-(1-methylpyrrolidin-2-yl)prop-2-en-1-one



5

[00275] Step 1) Preparation of diethyl (2-(10-((3-chloro-4-(pyridin-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-2-oxoethyl)phosphate



10 [00276] N-(3-chloro-4-(pyridin-2-ylmethoxy)phenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-


f]quinazolin-10-amine (VIII-5)(210 mg, 0.5 mmol), 2-(diethoxyphosphoryl)acetic acid (0.5 mmol) were dissolved in tetrahydrofuran, to which N,N'-carbonyldiimidazole (81 mg, 0.5 mmol) was added, stirred at room temperature until the reaction was completed. Water and ethyl acetate were added for extraction, the organic phase was concentrated and purified using column chromatography to afford 253 mg of yellow solid with a yield of 85%. MS: 598[M+H]⁺.

[00277] Step 2) Preparation of (S,E)-1-(10-((3-chloro-4-(pyridin-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-3-(1-methylpyrrolidin-2-yl)prop-2-en-1-one

[00278] diethyl (2-(10-((3-chloro-4-(pyridin-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-2-oxoethyl)phosphate (253 mg, 0.43 mmol) was dissolved in tetrahydrofuran, which was cooled to -78°C, and 1 mol/l of toluene solution of bis-trimethylsilylamide lithium (0.83 mL, 0.83 mmol) was added dropwise, and stirred until the all of the materials were disappeared. (S)-1-methylpyrrolidinyl-2-carbaldehyde (48.6 mg, 0.43 mmol) was added, the reaction was warmed to room temperature and stirred until the reaction was completed. Water and ethyl acetate were added for extraction, the organic phase was concentrated and purified using column chromatography to afford 167 mg of off-white solid with a yield of 70%. ¹H NMR (300 MHz, DMSO-d₆) δ 9.82 (s, 1H), 8.65 – 8.56 (m, 1H), 8.46 (s, 1H), 8.01 (d, J = 2.6 Hz, 1H), 7.95 – 7.72 (m, 2H), 7.71 – 7.61 (m, 1H), 7.58 (d, J = 7.9 Hz, 1H), 7.40 – 7.32 (m, 1H), 7.32 – 7.26 (m, 2H), 6.77 – 6.64 (m, 1H), 6.64 – 6.48 (m, 1H), 5.31 (s, 2H), 4.66 (s, 2H), 4.16 – 3.92 (m, 2H), 3.09 – 2.94 (m, 1H), 2.91 – 2.74 (m, 1H), 2.29 – 2.11 (m, 4H), 2.06 – 1.91 (m, 1H), 1.82 – 1.64 (m, 2H), 1.62 – 1.44 (m, 1H); MS: 557[M+H]⁺.

[00279] Example 113

[00280] Preparation of 1-(10-((3-chloro-4-(pyridin-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-yne-1-one




[00281] But-2-ynoic acid (42 mg, 0.5 mmol) was dissolved in dichloromethane, and 0.05 mL of dimethylformamide was added dropwise, which was cooled in an ice bath and then oxalyl chloride (32 mg, 0.25 mmol) was added. After stirring for 0.5h, N-(3-chloro-4-(pyridin-2-ylmethoxy)phenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine (VIII-5)(210 mg, 0.5 mmol) was added, and the reaction was warmed to room temperature and stirred until the

reaction was completed. The reaction was quenched by potassium carbonate aqueous solution, extracted with ethyl acetate, and the organic phase was concentrated and purified using column chromatography to afford 75 mg of off-white solid product, with a yield of 31%. ¹H NMR (400 MHz, DMSO-d₆) δ 9.82 (s, 1H), 8.63 – 8.59 (m, 1H), 8.46 (s, 1H), 8.40 – 8.34 (m, 1H), 8.01 – 7.97 (m, 1H), 7.92 – 7.86 (m, 1H), 7.69 – 7.63 (m, 1H), 7.60 – 7.57 (m, 1H), 7.40 – 7.36 (m, 1H), 7.33 – 7.25 (m, 2H), 5.34 – 5.28 (m, 2H), 4.76 – 4.68 (m, 2H), 4.28 (s, 2H), 2.14 (s, 3H). MS: 486[M+H]⁺.

[00282] Example 114: Preparation of (R,E)-1-(10-(3-chloro-4-(pyridin-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-3-(1-methylpyrrolidin-2-yl)prop-2-en-1-one

[00283] Step 1): Preparation of diethyl (2-(10-((3-chloro-4-(pyridin-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-2-oxoethyl)phosphonate




15

[00284] N-(3-chloro-4-(pyridin-2-ylmethoxy)phenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine (210 mg, 0.5 mmol), 2-(diethoxyphosphoryl)acetic acid (98 mg, 0.5 mmol) were dissolved in tetrahydrofuran, to which N,N'-carbonyldiimidazole (81 mg, 0.5 mmol) was added, the solution was stirred in a 40°C oil bath until the reaction was completed. Water and ethyl acetate were added for extraction, the organic phase was concentrated and purified using column chromatography to afford 254 mg of yellow solid with a yield of 85%. MS: 598[M+H]⁺.

20

[00285] Step 2) Preparation of (R,E)-1-(10-(3-chloro-4-(pyridin-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-3-(1-methylpyrrolidin-2-yl)prop-2-en-1-one


25



[00286] diethyl (2-(10-((3-chloro-4-(pyridin-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-2-oxoethyl)phosphonate (254 mg, 0.43 mmol) was

dissolved in tetrahydrofuran, cooled to -78°C , and 1 mol/l of toluene solution of bis-trimethylsilylamide lithium (0.65 mL, 0.65 mmol) was added dropwise, which was stirred until the all of the materials were disappeared. (R)-1-methylpyrrolidinyl-2-carbaldehyde (48.6 mg, 0.43 mmol) was added, and the reaction was warmed to room temperature and stirred until the reaction was completed. Water and ethyl acetate were added for extraction, the organic phase was concentrated and separated by high performance liquid chromatography to afford 154 mg of white solid with a yield of 65%. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 9.82 (s, 1H), 8.60 (d, $J = 4.9$ Hz, 1H), 8.46 (s, 1H), 8.01 (d, $J = 2.5$ Hz, 1H), 7.92-7.74 (m, 2H), 7.69-7.64 (m, 1H), 7.58 (d, $J = 7.8$ Hz, 1H), 7.40-7.34 (m, 1H), 7.31-7.24 (m, 2H), 6.73-6.66 (m, 1H), 6.55 (d, $J = 15.4$ Hz, 1H), 5.31 (s, 2H), 4.70-4.63 (m, 2H), 4.11-3.97 (m, 2H), 2.99 (d, $J = 8.2$ Hz, 1H), 2.82-2.74 (m, 1H), 2.20 (s, 3H), 2.16 (d, $J = 8.7$ Hz, 1H), 2.06-1.92 (m, 1H), 1.71 (d, $J = 9.4$ Hz, 2H), 1.63-1.49 (m, 1H); MS: 557[M+H] $^{+}$.

[00287] Example 115: Preparation of (E)-1-(10-((3-chloro-4-(pyridin-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(4-hydroxypiperidin-1-yl)but-2-en-1-one




[00288] N-(3-chloro-4-(pyridin-2-ylmethoxy)phenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine (210 mg, 0.5 mmol) was dissolved in dimethylformamide, to which 4-bromocrotonyl chloride (91 mg, 0.5 mmol) was added, and stirred at room temperature until the reaction was completed. The reaction was quenched by adding water, extracted with ethyl acetate, the organic phase was concentrated and directly dissolved in acetonitrile, to which diisopropylethylamine (129 mg, 1 mmol) and piperidin-4-ol (101 mg, 1 mmol) were added, and stirred at room temperature until the reaction was completed. The reaction was quenched by the addition of water, extracted with ethyl acetate, and the organic phase was concentrated and purified by high performance liquid chromatography to afford 97 mg of white solid with a yield of 33%. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 9.81 (s, 1H), 8.60 (d, $J = 4.8$ Hz, 1H), 8.46 (s, 1H), 8.01 (d, $J = 2.6$ Hz, 1H), 7.92-7.85 (m, 1H), 7.81 (s, 1H), 7.71-7.63 (m, 1H), 7.58 (d, $J = 7.9$ Hz, 1H), 7.41-7.33 (m, 1H), 7.28 (t, $J = 9.4$ Hz, 2H), 6.86-6.76 (m, 1H), 6.57 (d, $J = 15.4$ Hz, 1H), 5.31 (s, 2H), 4.66 (t, $J = 4.6$ Hz, 2H), 4.50 (d, $J = 4.1$ Hz, 1H), 4.04 (t, $J = 4.7$ Hz, 2H), 3.44 (s, 1H), 3.11 (d, $J = 5.6$ Hz, 2H), 2.69 (d, $J = 12.0$ Hz, 2H), 2.07 (t, $J = 10.7$ Hz, 2H), 1.70

(d, $J = 12.4$ Hz, 2H), 1.37 (d, $J = 11.1$ Hz, 2H); MS: 587[M+H]⁺.


[00289] Example 116: Preparation of (E)-1-(10-((3-chloro-4-(pyridin-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(4-

5 hydroxyl-4-methylpiperidin-1-yl)but-2-en-1-one



[00290] It was prepared by a method similar to that of Example 115, except that piperidin-4-ol was replaced by the same molar equivalent of 4-methylpiperidin-4-ol. ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.81 (s, 1H), 8.60 (d, $J = 4.9$ Hz, 1H), 8.45 (s, 1H), 8.00 (s, 1H), 7.94-7.72 (m, 2H), 7.69-7.61 (m, 1H), 7.58 (d, $J = 7.8$ Hz, 1H), 7.41-7.32 (m, 1H), 7.26 (t, $J = 8.7$ Hz, 2H), 6.86-6.74 (m, 1H), 6.62-6.46 (m, 1H), 5.30 (s, 2H), 4.65 (t, $J = 4.7$ Hz, 2H), 4.10 (s, 1H), 4.03 (t, $J = 4.8$ Hz, 2H), 3.19-3.06 (m, 2H), 2.46-2.28 (m, 4H), 1.51-1.26 (m, 4H), 1.08 (s, 3H); MS: 601[M+H]⁺.


[00291] Example 117: Preparation of (E)-1-(10-((3-chloro-4-(pyridin-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(4-(hydroxymethyl)piperidin-1-yl)but-2-en-1-one



[00292] It was prepared by a method similar to that of Example 115, except that piperidin-4-ol was replaced by the same molar equivalent of piperidin-4-ylmethanol. ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.81 (s, 1H), 8.60 (d, $J = 4.8$ Hz, 1H), 8.46 (s, 1H), 8.01 (d, $J = 2.5$ Hz, 1H), 7.88 (t, $J = 8.0$ Hz, 2H), 7.70-7.62 (m, 1H), 7.58 (d, $J = 7.9$ Hz, 1H), 7.40-7.33 (m, 1H), 7.28 (t, $J = 9.4$ Hz, 2H), 6.87-6.75 (m, 1H), 6.57 (d, $J = 15.3$ Hz, 1H), 5.31 (s, 2H), 4.66 (d, $J = 5.0$ Hz, 2H), 4.37 (t, $J = 5.3$ Hz, 1H), 4.04 (s, 2H), 3.23 (t, $J = 5.9$ Hz, 2H), 3.12 (d, $J = 5.7$ Hz, 2H), 2.83 (d, $J = 11.1$ Hz, 2H), 1.92 (t, $J = 11.4$ Hz, 2H), 1.63 (d, $J = 12.6$ Hz, 2H), 1.37-1.25 (m, 1H), 1.18-1.04 (m, 2H); MS: 601[M+H]⁺.


[00293] Example 118: Preparation of (E)-1-(10-((3-chloro-4-(pyridin-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(4-

fluoropiperidin-1-yl)but-2-en-1-one




[00294] It was prepared by a method similar to that of Example 115, except that piperidin-4-ol was replaced by the same molar equivalent of 4-fluoropiperidine. ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.81 (s, 1H), 8.63-8.57 (m, 1H), 8.46 (s, 1H), 8.01 (d, *J* = 2.5 Hz, 1H), 7.93-7.76 (m, 2H), 7.70-7.62 (m, 1H), 7.58 (d, *J* = 7.9 Hz, 1H), 7.40-7.34 (m, 1H), 7.32-7.22 (m, 2H), 6.87-6.77 (m, 1H), 6.59 (d, *J* = 15.2 Hz, 1H), 5.31 (s, 2H), 4.66 (t, *J* = 4.7 Hz, 3H), 4.04 (t, *J* = 4.4 Hz, 2H), 3.15 (d, *J* = 5.6 Hz, 2H), 2.55 (d, *J* = 6.9 Hz, 2H), 2.34 (s, 2H), 1.91-1.77 (m, 2H), 1.70 (s, 2H); MS: 589[M+H]⁺.

[00295] Example 119: Preparation of (E)-1-(10-((3-chloro-4-(pyridin-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-((2-methoxyethyl)(methyl)amino)but-2-en-1-one




[00296] It was prepared by a method similar to that of Example 115, except that piperidin-4-ol was replaced by the same molar equivalent of N-(2-methoxyethyl)methylamine. ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.81 (s, 1H), 8.60 (d, *J* = 4.8 Hz, 1H), 8.46 (s, 1H), 8.01 (d, *J* = 2.6 Hz, 1H), 7.93-7.73 (m, 2H), 7.70-7.61 (m, 1H), 7.58 (d, *J* = 7.8 Hz, 1H), 7.42-7.31 (m, 1H), 7.31-7.18 (m, 2H), 6.87-6.73 (m, 1H), 6.64-6.51 (m, 1H), 5.30 (s, 2H), 4.65 (t, *J* = 4.7 Hz, 2H), 4.04 (t, *J* = 4.7 Hz, 2H), 3.40 (t, *J* = 5.8 Hz, 2H), 3.24-3.14 (m, 5H), 2.55-2.51 (m, 2H), 2.21 (s, 3H); MS: 575[M+H]⁺.

[00297] Example 120: Preparation of (E)-1-(10-((3-chloro-4-(pyridin-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-((3-hydroxypropyl)(methyl)amino)but-2-en-1-one




[00298] It was prepared by a method similar to that of Example 115, except that piperidin-4-ol was replaced by the same molar equivalent of 3-(methylamino)-1-propanol. ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.81 (s, 1H), 8.60 (d, *J* = 4.8 Hz, 1H), 8.46 (s, 1H), 8.01 (d, *J* = 2.6 Hz, 1H), 7.92-7.76 (m, 2H), 7.71-7.63 (m, 1H), 7.58 (d, *J* = 7.9 Hz, 1H), 7.37 (t, *J* = 6.3 Hz, 1H), 7.32-7.23 (m, 2H), 6.88-6.77 (m, 1H), 6.58 (d, *J* = 15.2 Hz, 1H), 5.31 (s, 2H), 4.65 (d, *J* = 4.8 Hz, 2H), 4.39 (s, 1H), 4.04 (t, *J* = 4.6 Hz, 2H), 3.44 (d, *J* = 5.8 Hz, 2H), 3.15 (d, *J* = 5.7 Hz, 2H), 2.39 (t, *J* = 7.3 Hz, 2H), 2.17 (s, 3H), 1.62-1.52 (m, 2H); MS: 575[M+H]⁺.

[00299] Example 121: Preparation of (S,E)-1-(10-((3-chloro-4-(pyridin-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(2,4-dimethylpiperazin-1-yl)but-2-en-1-one




[00300] It was prepared by a method similar to that of Example 115, except that piperidin-4-ol was replaced by the same molar equivalent of (S)-1,3-dimethylpiperazine. ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.80 (s, 1H), 8.59 (d, *J* = 4.7 Hz, 1H), 8.45 (s, 1H), 8.00 (d, *J* = 2.6 Hz, 1H), 7.87 (t, *J* = 7.7, 1.9 Hz, 1H), 7.84-7.69 (m, 1H), 7.69-7.62 (m, 1H), 7.57 (d, *J* = 7.8 Hz, 1H), 7.41-7.32 (m, 1H), 7.26 (t, *J* = 9.3 Hz, 2H), 6.89-6.78 (m, 1H), 6.65-6.51 (m, 1H), 5.29 (s, 2H), 4.72-4.56 (m, 2H), 4.13-3.90 (m, 2H), 3.55-3.41 (m, 1H), 3.05-2.94 (m, 1H), 2.72-2.62 (m, 1H), 2.58-2.52 (m, 1H), 2.47-2.30 (m, 2H), 2.28-2.20 (m, 1H), 2.10 (s, 3H), 2.04-1.96 (m, 1H), 1.74 (t, *J* = 10.1 Hz, 1H), 0.96 (d, *J* = 6.2 Hz, 3H); MS: 600[M+H]⁺.

[00301] Example 122: Preparation of (E)-1-(10-((3-chloro-4-(pyridin-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-((2-hydroxyethyl)(methyl)amino)but-2-en-1-one




[00302] It was prepared by a method similar to that of Example 115, except that piperidin-4-ol was replaced by the same molar equivalent of 2-methylaminoethanol. ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.81 (s, 1H), 8.60 (d, *J* = 4.8 Hz, 1H), 8.46 (s, 1H), 8.01 (d, *J* = 2.6 Hz, 1H), 7.91-7.79 (m, 2H), 7.70-7.63 (m, 1H), 7.58 (d, *J* = 7.8 Hz, 1H), 7.37 (t, *J* = 6.3 Hz, 1H), 7.34-7.23 (m, 2H), 6.89-6.78 (m, 1H), 6.61 (d, *J* = 15.5 Hz, 1H), 5.31 (s, 2H), 4.66 (t, *J* = 4.5 Hz, 2H), 4.38 (t, *J* = 5.4 Hz, 1H), 4.05 (t, *J* = 4.7 Hz, 2H), 3.52-3.44 (m, 2H), 3.20 (d, *J* = 5.7 Hz, 2H), 2.44 (t, *J* = 6.3 Hz, 2H), 2.22 (s, 3H); MS: 561[M+H]⁺.

[00303] Example 123: Preparation of (R,E)-1-(10-((3-chloro-4-(pyridin-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(2,4-dimethylpiperazin-1-yl)but-2-en-1-one



[00304] It was prepared by a method similar to that of Example 115, except that piperidin-4-ol was replaced by the same molar equivalent of (R)-1,3-dimethylpiperazine. ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.82 (s, 1H), 8.60 (d, *J* = 4.8 Hz, 1H), 8.46 (s, 1H), 8.01 (d, *J* = 2.6 Hz, 1H), 7.92-7.72 (m, 2H), 7.71-7.63 (m, 1H), 7.58 (d, *J* = 7.8 Hz, 1H), 7.37 (t, *J* = 6.3 Hz, 1H), 7.28 (t, *J* = 8.8 Hz, 2H), 6.90-6.80 (m, 1H), 6.66-6.51 (m, 1H), 5.31 (s, 2H), 4.72-4.59 (m, 2H), 4.13-3.95 (m, 2H), 3.50 (d, *J* = 15.7 Hz, 1H), 3.13-2.96 (m, 2H), 2.70 (d, *J* = 13.2 Hz, 1H), 2.54 (s, 1H), 2.41 (s, 1H), 2.26 (t, *J* = 10.6 Hz, 1H), 2.11 (s, 3H), 2.04 (d, *J* = 11.1 Hz, 1H), 1.76 (s, 1H), 0.97 (d, *J* = 6.2 Hz, 3H); MS: 600[M+H]⁺.


[00305] Example 124: Preparation of (E)-1-(10-((4-([1,2,4]triazolo[1,5-a]pyridin-7-yloxy)-3-methylphenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(dimethylamino)but-2-en-1-one



[00306] N-(4-([1,2,4]triazolo[1,5-a]pyridin-7-yloxy)-3-methylphenyl)-3,4-dihydro-2H-

[1,4]oxazino[2,3-f]quinazolin-10-amine (213 mg, 0.5 mmol) was dissolved in dimethylformamide, to which (E)-4-(dimethylamino)but-2-enoyl chloride hydrochloride (137 mg, 0.75 mmol) was added at room temperature, and stirred at room temperature until the reaction was completed. The reaction was quenched by the addition of potassium carbonate aqueous solution, extracted with ethyl acetate, and the organic phase was concentrated and purified by high performance liquid chromatography to afford 200 mg of white solid with a yield of 75%. ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.93 (s, 1H), 8.94 (d, *J* = 7.5 Hz, 1H), 8.50 (s, 1H), 8.38 (s, 1H), 7.97-7.69 (m, 3H), 7.32 (d, *J* = 9.0 Hz, 1H), 7.22 (d, *J* = 8.6 Hz, 1H), 7.08-6.98 (m, 1H), 6.87-6.72 (m, 2H), 6.64-6.52 (m, 1H), 4.69 (t, *J* = 4.6 Hz, 2H), 4.06 (t, *J* = 4.5 Hz, 2H), 3.14-3.01 (m, 2H), 2.33-2.03 (m, 9H); MS: 537[M+H]⁺.

[00307] Example 125: Preparation of (E)-1-(10-((4-([1,2,4]triazolo[1,5-a]pyridin-7-yloxy)-3-chlorophenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(dimethylamino)but-2-en-1-one




15

[00308] N-(4-([1,2,4]triazolo[1,5-a]pyridin-7-yloxy)-3-chlorophenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine (223 mg, 0.5 mmol) was dissolved in dimethylformamide, to which (E)-4-(dimethylamino)but-2-enoyl chloride hydrochloride (137 mg, 0.75 mmol) was added at room temperature, and stirred at room temperature until the reaction was completed. The reaction was quenched by the addition of potassium carbonate aqueous solution, extracted with ethyl acetate, and the organic phase was concentrated and purified by high performance liquid chromatography to afford 195 mg of white solid with a yield of 70%. ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.03 (s, 1H), 8.97 (d, *J* = 7.3 Hz, 1H), 8.56 (s, 1H), 8.41 (s, 1H), 8.32 (d, *J* = 2.5 Hz, 1H), 7.98-7.93 (m, 1H), 7.93-7.73 (m, 1H), 7.47 (d, *J* = 8.8 Hz, 1H), 7.34 (d, *J* = 9.1 Hz, 1H), 7.11-7.02 (m, 1H), 6.92 (d, *J* = 2.6 Hz, 1H), 6.87-6.77 (m, 1H), 6.65-6.52 (m, 1H), 4.69 (t, 2H), 4.06 (t, *J* = 4.7 Hz, 2H), 3.11-3.02 (m, 2H), 2.17 (s, 6H); MS: 557[M+H]⁺.

25

[00309] Example 126: Preparation of (E)-1-(10-((4-([1,2,4]triazolo[1,5-a]pyridin-7-yloxy)-3-


chlorophenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(4-methylpiperazin-1-yl)but-2-en-1-one



[00310] N-(4-([1,2,4]triazolo[1,5-a]pyridin-7-yloxy)-3-chlorophenyl)-3,4-dihydro-2H-

5 [1,4]oxazino[2,3-f]quinazolin-10-amine (223 mg, 0.5 mmol) was dissolved in dimethylformamide, to which 4-bromocrotonyl chloride (91 mg, 0.5 mmol) was added, and stirred at room temperature until the reaction was completed. The reaction was quenched by adding water, extracted with ethyl acetate, the organic phase was concentrated and directly dissolved in acetonitrile, to which diisopropylethylamine (129 mg, 1 mmol) and 1-methylpiperazine (100 mg, 1 mmol) were added, and stirred at room temperature until the
10 reaction was completed. The reaction was quenched by the addition of water, extracted with ethyl acetate, the organic phase was concentrated and purified by high-performance liquid chromatography to afford a white solid 125 mg with a yield of 41%. ¹H NMR (400 MHz, MeOH-*d*₄) δ 8.66 (d, *J* = 7.5 Hz, 1H), 8.46-8.38 (m, 2H), 8.21-8.18 (m, 2H), 7.73 (d, *J* = 2.6 Hz, 1H), 7.71 (d, *J* = 2.6 Hz, 1H), 7.29-7.24 (m, 2H), 7.00-6.96 (m, 1H), 6.91-6.83 (m, 1H),
15 6.74 (d, *J* = 2.5 Hz, 1H), 6.58 (d, *J* = 15.3 Hz, 1H), 4.64 (t, *J* = 4.7 Hz, 2H), 4.06 (t, 2H), 3.18 (dd, *J* = 6.1, 1.6 Hz, 2H), 2.76-2.62 (m, 4H), 2.61-2.45 (m, 4H), 2.40 (s, 3H); MS: 612[M+H]⁺.

[00311] Example 127: Preparation of (E)-1-(10-((4-([1,2,4]triazolo[1,5-a]pyridin-7-yloxy)-3-chlorophenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(isopropyl(methyl)amino)but-2-en-1-one
20




[00312] It was prepared by a method similar to that of Example 126, except that 1-methylpiperazine was replaced by the same molar equivalent of N-isopropylmethylamine. ¹H

NMR (400 MHz, MeOH- d_4) δ 8.65 (d, J = 7.5 Hz, 1H), 8.40 (d, J = 1.7 Hz, 2H), 8.22-8.16 (m, 2H), 7.79-7.68 (m, 2H), 7.29-7.22 (m, 2H), 6.99-6.95 (m, 1H), 6.92-6.85 (m, 1H), 6.73 (d, J = 2.6 Hz, 1H), 6.68-6.62 (m, 1H), 4.64 (t, J = 4.7 Hz, 2H), 4.06 (t, J = 4.7 Hz, 2H), 3.41 (dd, J = 6.6, 1.4 Hz, 2H), 3.06-2.98 (m, 1H), 2.30 (s, 3H), 1.06 (d, J = 6.6 Hz, 6H); MS: 585[M+H]⁺.

5


[00313] Example 128: Preparation of (E)-1-(10-((4-([1,2,4]triazolo[1,5-a]pyridin-7-yloxy)-3-chlorophenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(piperidin-1-yl)but-2-en-1-one



[00314] It was prepared by a method similar to that of Example 126, except that 1-methylpiperazine was replaced by the same molar equivalent of piperidine. ¹H NMR (400 MHz, MeOH- d_4) δ 8.67 (d, J = 7.5 Hz, 1H), 8.43 (s, 2H), 8.23-8.18 (m, 2H), 7.73 (dd, J = 8.8, 2.6 Hz, 2H), 7.30-7.25 (m, 2H), 6.99 (dd, J = 7.5, 2.6 Hz, 1H), 6.93-6.84 (m, 1H), 6.75 (d, J = 2.6 Hz, 1H), 6.63 (d, J = 15.3 Hz, 1H), 4.65 (t, J = 4.7 Hz, 2H), 4.07 (t, J = 4.7 Hz, 2H), 3.38-3.31 (m, 2H), 2.62 (s, 4H), 1.64-1.57 (m, 4H), 1.48-1.40 (m, 2H); MS: 597[M+H]⁺.

15

[00315] Example 129: Preparation of (E)-1-(10-((4-([1,2,4]triazolo[1,5-a]pyridin-7-yloxy)-3-chlorophenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(cyclobutyl(methyl)amino)but-2-en-1-one




20

[00316] It was prepared by a method similar to that of Example 126, except that 1-methylpiperazine was replaced by the same molar equivalent of N-methylcyclobutylamine hydrochloride. ¹H NMR (400 MHz, MeOH- d_4) δ 8.66 (dd, J = 7.4, 0.6 Hz, 1H), 8.42 (s, 1H),

8.40 (s, 1H), 8.24-8.17 (m, 2H), 7.72 (dd, $J = 8.8, 2.6$ Hz, 2H), 7.30-7.24 (m, 2H), 7.00-6.96 (m, 1H), 6.93-6.85 (m, 1H), 6.74 (d, $J = 2.6$ Hz, 1H), 6.58 (d, $J = 15.2$ Hz, 1H), 4.65 (t, $J = 4.6$ Hz, 2H), 4.06 (t, 2H), 3.16 (dd, $J = 6.7, 1.4$ Hz, 2H), 3.00-2.91 (m, 1H), 2.14 (s, 3H), 2.06-1.98 (m, 2H), 1.90-1.79 (m, 2H), 1.69-1.57 (m, 2H); MS: 597[M+H]⁺.

5


[00317] Example 130: Preparation of (E)-1-(10-((4-([1,2,4]triazolo[1,5-a]pyridin-7-yloxy)-3-chlorophenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-((2-methoxyethyl)(methyl)amino)but-2-en-1-one



10 **[00318]** It was prepared by a method similar to that of Example 126, except that 1-methylpiperazine was replaced by the same molar equivalent of N-(2-methoxyethyl)methylamine. ¹H NMR (400 MHz, MeOH-*d*₄) δ 8.79 (dd, $J = 7.5, 0.7$ Hz, 1H), 8.56 (s, 1H), 8.34-8.32 (m, 2H), 7.94-7.89 (m, 1H), 7.88-7.87 (m, 1H), 7.86-7.85 (m, 1H), 7.41 (dd, $J = 8.9, 4.0$ Hz, 2H), 7.12 (dd, $J = 7.5, 2.6$ Hz, 1H), 7.07-6.99 (m, 1H), 6.89-6.87 (m, 1H),
15 6.71 (d, $J = 15.3$ Hz, 1H), 4.77 (t, $J = 4.8$ Hz, 2H), 4.19 (t, $J = 4.7$ Hz, 2H), 3.55 (t, $J = 5.5$ Hz, 2H), 3.35 (t, 2H), 3.35 (s, 3H), 2.68 (t, $J = 5.5$ Hz, 2H), 2.37 (s, 3H); MS: 601 [M+H]⁺.

[00319] Example 131: Preparation of (E)-1-(10-((4-([1,2,4]triazolo[1,5-a]pyridin-7-yloxy)-3-chlorophenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(diethylamino)but-2-en-1-one


20



[00320] It was prepared by a method similar to that of Example 126, except that 1-methylpiperazine was replaced by the same molar equivalent of diethylamine. ¹H NMR (400


MHz, MeOH-*d*₄) δ 8.79 (dd, *J* = 7.5, 0.7 Hz, 1H), 8.56 (s, 1H), 8.33 (s, 1H), 8.32 (d, *J* = 2.5 Hz, 1H), 7.91-7.88 (m, 1H), 7.88-7.86 (m, 1H), 7.86 (d, *J* = 2.6 Hz, 1H), 7.43-7.39 (m, 2H), 7.13-7.10 (m, 1H), 7.09-7.01 (m, 1H), 6.88-6.87 (m, 1H), 6.70 (d, *J* = 15.3 Hz, 1H), 4.77 (t, *J* = 4.7 Hz, 2H), 4.20-4.16 (m, 2H), 3.39 (dd, *J* = 6.3, 1.5 Hz, 2H), 2.64 (q, *J* = 7.2 Hz, 4H), 1.12 (t, *J* = 7.2 Hz, 6H); MS: 585 [M+H]⁺.

[00321] Example 132: Preparation of (E)-1-(10-((4-([1,2,4]triazolo[1,5-a]pyridin-7-yloxy)-3-methylphenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(4-methylpiperazin-1-yl)but-2-en-1-one




[00322] N-(4-([1,2,4]triazolo[1,5-a]pyridin-7-yloxy)-3-methylphenyl)-3,4-dihydro-2H-[1,4]oxazino[2,3-f]quinazolin-10-amine (212 mg, 0.5 mmol) was dissolved in dimethylformamide, to which 4-bromocrotonyl chloride (91 mg, 0.5 mmol) was added, and stirred at room temperature until the reaction was completed. The reaction was quenched by adding water, extracted with ethyl acetate, the organic phase was concentrated and directly dissolved in acetonitrile, to which diisopropylethylamine (129 mg, 1 mmol) and 1-methylpiperazine (100 mg, 1 mmol) were added, and stirred at room temperature until the reaction was completed. The reaction was quenched by the addition of water, extracted with ethyl acetate, and the organic phase was concentrated and purified by high performance liquid chromatography to afford 110 mg of a white solid with a yield of 37%. ¹H NMR (400 MHz, MeOH-*d*₄) δ 8.77 (d, *J* = 7.5 Hz, 1H), 8.49 (s, 1H), 8.31 (s, 1H), 7.88-7.83 (m, 1H), 7.83-7.80 (m, 2H), 7.79-7.77 (m, 1H), 7.38 (d, *J* = 9.1 Hz, 1H), 7.21 (d, *J* = 8.5 Hz, 1H), 7.12-7.09 (m, 1H), 7.03-6.96 (m, 1H), 6.84 (d, *J* = 2.5 Hz, 1H), 6.70 (d, *J* = 15.3 Hz, 1H), 4.76 (t, *J* = 4.7 Hz, 2H), 4.18 (t, *J* = 4.7 Hz, 2H), 3.30-3.28 (m, 2H), 2.78-2.55 (m, 8H), 2.45 (s, 3H), 2.27 (s, 3H); MS: 592 [M+H]⁺.

[00323] Example 133: Preparation of (E)-1-(10-((4-([1,2,4]triazolo[1,5-a]pyridin-7-yloxy)-3-methylphenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(isopropyl(methyl)amino)but-2-en-1-one



[00324] It was prepared by a method similar to that of Example 132, except that 1-methylpiperazine was replaced by the same molar equivalent of N-isopropylmethylamine. ¹H NMR (400 MHz, MeOH-*d*₄) δ 8.76 (d, *J* = 7.5 Hz, 1H), 8.48 (s, 1H), 8.31 (s, 1H), 7.89-7.83 (m, 1H), 7.81 (t, *J* = 2.4 Hz, 2H), 7.78 (d, *J* = 2.7 Hz, 1H), 7.40-7.36 (m, 1H), 7.21 (d, *J* = 8.5 Hz, 1H), 7.12-7.08 (m, 1H), 7.06-6.98 (m, 1H), 6.84 (d, *J* = 2.6 Hz, 1H), 6.71 (d, *J* = 15.2 Hz, 1H), 4.76 (t, *J* = 4.7 Hz, 2H), 4.18 (t, *J* = 4.7 Hz, 2H), 3.41-3.39 (m, 2H), 3.03-2.96 (m, 1H), 2.32 (s, 3H), 2.27 (s, 3H), 1.13 (s, 3H), 1.12 (s, 3H); MS: 565 [M+H]⁺.


- [00325]** Example 134: Preparation of (E)-1-(10-((4-([1,2,4]triazolo[1,5-a]pyridin-7-yloxy)-3-methylphenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(piperidin-1-yl)but-2-en-1-one



- [00326]** It was prepared by a method similar to that of Example 132, except that 1-methylpiperazine was replaced by the same molar equivalent of piperidine. ¹H NMR (400 MHz, MeOH-*d*₄) δ 8.64 (d, *J* = 7.5 Hz, 1H), 8.37 (s, 1H), 8.19 (s, 1H), 7.79-7.75 (m, 1H), 7.70-7.67 (m, 2H), 7.66 (d, *J* = 2.7 Hz, 1H), 7.26 (d, *J* = 9.0 Hz, 1H), 7.09 (d, *J* = 8.4 Hz, 1H), 7.00-6.96 (m, 1H), 6.89-6.82 (m, 1H), 6.75 (s, 1H), 6.71 (d, *J* = 2.6 Hz, 1H), 4.66 (d, *J* = 4.7 Hz, 2H), 4.10-4.05 (m, 2H), 3.63 (d, *J* = 6.6 Hz, 2H), 2.97-2.86 (m, 4H), 2.15 (s, 3H), 1.75-1.67 (m, 4H), 1.57-1.50 (m, 2H); MS: 577 [M+H]⁺.

[00327] Example 135: Preparation of (E)-1-(10-((4-([1,2,4]triazolo[1,5-a]pyridin-7-yloxy)-3-methylphenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-


(cyclobutyl(methyl)amino)but-2-en-1-one



[00328] It was prepared by a method similar to that of Example 132, except that 1-methylpiperazine was replaced by the same molar equivalent of N-methylcyclobutylamine hydrochloride. ¹H NMR (400 MHz, MeOH-*d*₄) δ 8.74 (d, *J* = 7.5 Hz, 1H), 8.50 (s, 1H), 8.45 (s, 1H), 8.29 (s, 1H), 7.88-7.81 (m, 1H), 7.80-7.75 (m, 2H), 7.34 (d, *J* = 9.1 Hz, 1H), 7.19-7.16 (m, 1H), 7.09-7.05 (m, 1H), 7.02-6.94 (m, 1H), 6.84-6.75 (m, 2H), 4.75 (t, *J* = 4.6 Hz, 2H), 4.18 (t, *J* = 4.6 Hz, 2H), 3.56 (d, *J* = 6.8 Hz, 2H), 3.42-3.37 (m, 1H), 2.66 (d, *J* = 9.4 Hz, 2H), 2.49 (s, 3H), 2.24 (s, 3H), 2.17-2.09 (m, 2H), 1.85-1.75 (m, 2H); MS: 577 [M+H]⁺.

10


[00329] Example 136: Preparation of (E)-1-(10-((4-([1,2,4]triazolo[1,5-a]pyridin-7-yloxy)-3-methylphenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-((2-methoxyethyl)(methyl)amino)but-2-en-1-one



[00330] It was prepared by a method similar to that of Example 132, except that 1-methylpiperazine was replaced by the same molar equivalent of N-(2-methoxyethyl)methylamine. ¹H NMR (400 MHz, MeOH-*d*₄) δ 8.76 (dd, *J* = 7.6, 0.7 Hz, 1H), 8.48 (s, 1H), 8.31 (s, 1H), 7.93-7.83 (m, 1H), 7.83-7.80 (m, 2H), 7.79-7.77 (m, 1H), 7.38 (d, *J* = 9.1 Hz, 1H), 7.21 (d, *J* = 8.5 Hz, 1H), 7.12-7.08 (m, 1H), 7.07-6.99 (m, 1H), 6.85-6.83 (m, 1H), 6.70 (d, *J* = 15.3 Hz, 1H), 4.76 (t, *J* = 4.7 Hz, 2H), 4.18 (t, 2H), 3.57-3.52 (m, 2H), 3.35-3.34 (m, 5H), 2.67 (t, *J* = 5.5 Hz, 2H), 2.35 (s, 3H), 2.27 (s, 3H); MS: 581 [M+H]⁺.

20

[00331] Example 137: Preparation of (E)-1-(10-((4-([1,2,4]triazolo[1,5-a]pyridin-7-yloxy)-3-methylphenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(diethylamino)but-2-en-1-one



5 [00332] It was prepared by a method similar to that of Example 132, except that 1-methylpiperazine was replaced by the same molar equivalent of diethylamine. ¹H NMR (400 MHz, MeOH-*d*₄) δ 8.76 (dd, *J* = 7.5, 0.7 Hz, 1H), 8.48 (s, 1H), 8.31 (s, 1H), 7.84-7.80 (m, 2H), 7.78 (d, *J* = 2.7 Hz, 1H), 7.38 (d, *J* = 9.1 Hz, 1H), 7.21 (d, *J* = 8.5 Hz, 1H), 7.12-7.10 (m, 1H), 7.10-7.08 (m, 1H), 7.07-7.01 (m, 1H), 6.84 (dd, *J* = 2.7, 0.7 Hz, 1H), 6.70 (d, *J* = 15.3 Hz, 1H),
10 4.77 (t, *J* = 4.7 Hz, 2H), 4.19 (d, *J* = 4.7 Hz, 2H), 3.41-3.39 (m, 2H), 2.67-2.63 (m, 4H), 2.27 (s, 3H), 1.12 (t, *J* = 7.1 Hz, 6H); MS: 565 [M+H]⁺.

[00333] Assay Example 1.

[00334] The assay for the inhibition of EGFR and HER2 kinase activity by small molecular
15 compounds was carried out using the method as follows:

[00335] 1) Dilution of the compounds

[00336] In a 96-well plate a, the compounds were diluted with DMSO using a 3-fold
gradient dilution to form 11 concentrations, the 12th concentration is pure DMSO (as a
positive control); and in a new 96-well plate b the above solutions were diluted 25 times with
20 ultrapure water (DMSO concentration is 4%).

[00337] 2) Transferring the compounds to 384-well plate

[00338] The compound solutions diluted with ultrapure water in the 96-well plate b
above was transferred to the corresponding wells of a 384-well plate in duplicate.

[00339] 3) Addition of 4×kinase solution: 2.5μl of the above 4× kinase solution was
25 taken using multichannel pipette and added to the corresponding reaction wells of the 384-
well plate, mixed well and pre-reacted at room temperature for 5 minutes.

[00340] 4) Addition of 2×substrate/ATP mixed solution: 5μl of the above
2×substrate/ATP mixed solution was taken using multichannel pipette and added to the
corresponding reaction wells of the 384-well plate.

- [00341] 5) Negative control: negative control wells were set in the 384-well plate, and 2.5µl 4×substrate, 2.5µl 4×enzyme solution, 2.5µl 1×Kinase Assay Buffer and 2.5µl ultrapure water containing 4% DMSO were added to each well.
- [00342] 6) Mixed by centrifugation and kept at room temperature for 2 hours in the dark.
- [00343] 7) Termination of the enzymatic reaction:
- [00344] 5 µl of the above 4× stop solution was pipetted to the corresponding wells of the 384-well plate, centrifuged and mixed, and reacted at room temperature for 5 minutes.
- [00345] 8) Development reaction:
- [00346] 5µl of the above 4x detection solution was pipetted into the corresponding wells of the 384-well plate, centrifuged and mixed, and reacted at room temperature for 1 hour.
- [00347] 9) The 384-well plate was placed into a microplate reader and the signal was detected using the corresponding program.
- [00348] 10) IC₅₀ analysis:
- [00349] Well reading value=10000*EU665 value/ EU615 value
- [00350] Inhibition rate =(reading value of positive control well - reading value of experimental well)/(reading value of positive control well - reading value of negative control well)*100%
- Corresponding IC₅₀s can be calculated by entering the drug concentrations and the corresponding inhibition rates into GraphPad Prism 5.
- [00351] Conditions of experiment for screening EGFR kinase inhibitory molecules:
- [00352] The final concentration of EGFR kinase in the reaction system is 0.35 nM, and the final concentration of ATP is 150 µM, the final concentration of substrate *ULight*TM-labeled JAK-1 (Tyr1023) Peptide is 100 nM, and the enzymatic reaction time is 2 hours.
- [00353] The maximum final concentration of the compound in the reaction system is 2.5 µM, 11 concentrations were made with a 3-fold gradient dilution, and the minimal final concentration is 0.042 nM. The final concentration of DMSO is 1%.
- [00354] Conditions of experiment for screening HER2 kinase inhibitory molecules:
- [00355] The final concentration of HER2 kinase in the reaction system is 10 nM, the final concentration of ATP is 10 µM, the final concentration of substrate *ULight*TM-labeled PolyGT is 100 nM, and the enzymatic reaction time is 2 hours.
- [00356] The maximum final concentration of the compound in the reaction system is 2.5 µM, 11 concentrations were made with a 3-fold gradient dilution, and the minimal final

concentration is 0.042 nM. The final concentration of DMSO is 1%.

[00357] Assay results of the inhibitory activity of some compounds disclosed herein on tyrosine kinases were listed in Table (1), wherein A means IC₅₀ is less than or equal to 50 nM, B means IC₅₀ is greater than 50 nM but less than or equal to 500 nM, C means IC₅₀ is greater than 500 nM but less than or equal to 5000 nM, D means IC₅₀ is greater than 5000 nM, and NT means that the compound was not tested for the corresponding kinase.

[00358] Table (1), assay results of the inhibitory activity of the compounds disclosed herein on EGFR and HER2 kinases

Example	HER2	EGFR	Example	HER2	EGFR
No.	IC ₅₀ nM	IC ₅₀ nM	No.	IC ₅₀ nM	IC ₅₀ nM
1	A	NT	70	A	A
2	A	A	71	A	A
3	A	A	72	A	A
4	A	A	73	A	B
5	A	NT	74	A	B
6	A	NT	75	A	B
7	A	NT	76	A	C
8	A	NT	77	A	C
9	A	NT	78	A	NT
10	A	NT	79	A	NT
11	A	NT	80	A	NT
12	A	NT	81	A	NT
13	A	NT	82	A	NT
14	A	NT	83	A	A
15	A	NT	84	A	NT
16	A	NT	85	A	NT
17	A	NT	86	A	NT
18	A	NT	87	A	NT
19	A	NT	88	A	NT
20	A	NT	89	A	NT
21	A	NT	90	A	NT
22	A	NT	91	A	A
23	A	NT	92	A	A
24	A	NT	93	A	A
25	A	NT	94	A	A
26	A	NT	95	A	A
27	A	NT	96	A	A
28	A	NT	97	A	NT
29	A	NT	98	A	NT
30	A	NT	99	A	B
31	B	NT	100	A	NT
32	A	NT	101	B	NT
33	A	A	102	A	NT
34	A	A	103	A	NT
35	A	A	104	A	A

36	A	NT	105	A	A
37	A	A	106	A	A
38	A	NT	107	A	A
39	A	A	108	A	A
40	A	A	109	A	NT
41	A	A	110	A	A
42	A	A	111	A	A
43	A	A	112	A	NT
44	A	A	113	A	NT
45	A	A	114	A	NT
46	A	A	115	A	NT
47	A	A	116	A	NT
48	A	A	117	A	NT
49	A	A	118	A	NT
50	A	A	119	A	NT
51	A	A	120	A	NT
52	A	A	121	A	NT
53	A	A	122	A	NT
54	A	C	123	A	NT
55	A	C	124	A	B
56	A	A	125	A	B
57	A	B	126	A	A
58	A	NT	127	A	B
59	A	A	128	A	B
60	A	A	129	A	B
61	A	A	130	A	B
62	A	A	131	A	B
63	A	B	132	A	B
64	A	B	133	A	C
65	A	A	134	A	B
66	A	A	135	A	B
67	A	A	136	A	B
68	A	A	137	A	C
69	A	A			

[00359] Assay Example 2.

[00360] The assay for the inhibition of cell proliferation by small molecular compounds was carried out using the method as follows:

5 **[00361]** 1. 600 μ L pancreatin was added to a T75 cell culture flask, which was digested in a 37°C incubator for about 1 min before 5 mL of DMEM complete culture solution was added, blew evenly, transferred to a 15 mL centrifuge tube, and centrifuged at 1000 rpm for 4 min;

[00362] 2. The supernatant was removed and 5mL DMEM complete culture solution was added, blew evenly, and 10 μ L cell suspension was taken and mixed with 10 μ L 0.4% Trypan Blue, and counted using a cell counter;

10

[00363] 3. BT474 and HCC827 cell lines were seeded in 96-well plates at a cell density of

10,000 and 3000 cells/well/80 μ L, respectively, and cultured overnight. Only sterile water was added to the 36 wells on the periphery of the 96-well plate without adding cells, and only the 60 wells in the center of the 96-well plate were used for cell assays and controls;

[00364] 4. Dilution of compounds: the compounds were diluted with a 3-fold dilution to make

5 10 concentrations in total with an initial concentration of 10mM;

[00365] 5. 20 μ L of different compounds with different concentrations were added to each well, and 20 μ L of complete culture solution was added to the remaining wells. The final concentration of DMSO in each well is 0.25%;

[00366] 6. After 72 h of incubation, 10 μ L CCK-8 reagent was added to each well, and
10 incubated at 37°C for 1-2 h; and the OD value was read at 450 nm;

[00367] 7. Cell survival rate (%)=[(As-Ab)/(Ac-Ab)]*100%

[00368] As: Assay well (medium containing cell, CCK-8, compound)

[00369] Ac: Control well (medium containing cell, CCK-8)

[00370] Ab: Blank well (CCK-8, medium without cell and compound)

15 [00371] 8. The value was imported into Graphpad Prism5 software for the calculation of IC₅₀s (compound concentration at which 50% maximum survival rate is observed).

[00372] Table (2) lists the assay results of representative compounds disclosed herein on the viability of BT474 and HCC827 cancer cells. Wherein A means IC₅₀ is less than or equal to 50 nM, B means IC₅₀ is greater than 50 nM but less than or equal to 500 nM, C means IC₅₀ is
20 greater than 500 nM but less than or equal to 5000 nM, D means IC₅₀ is greater than 5000 nM, and NT means that the compound was not tested for the corresponding cell.

[00373] Table (2) Assay results of representative compounds disclosed herein on cell viability

Example	BT474	HCC827	Example	BT474	HCC827
No.	IC ₅₀	IC ₅₀	No.	IC ₅₀	IC ₅₀
44	A	C	92	A	C
46	A	B	93	A	B
47	A	C	94	A	C
48	B	C	95	A	B
49	A	B	96	A	B
50	A	C	100	A	B
51	A	C	101	NT	C
53	A	C	102	A	B
54	A	C	103	A	B
55	A	C	104	A	B
56	A	C	105	A	B
57	A	C	106	A	B
58	A	B	107	A	C
59	A	C	108	A	B
60	A	C	109	A	B
61	A	D	110	A	B


62	B	D	111	A	C
63	A	C	112	A	B
64	A	B	113	B	C
65	A	B	114	A	NT
66	A	C	115	A	NT
68	A	C	116	A	B
69	A	C	117	A	NT
70	A	NT	118	A	NT
71	A	C	119	A	B
72	A	B	120	A	NT
73	A	C	121	A	NT
74	A	C	122	A	NT
76	A	B	123	A	NT
78	A	C	124	A	NT
79	A	C	125	A	NT
80	A	C	126	A	NT
81	A	C	127	A	NT
82	A	C	128	A	NT
83	A	C	129	A	NT
84	A	B	130	A	NT
85	A	C	131	A	NT
86	A	C	132	A	NT
87	B	C	133	A	NT
88	A	C	134	A	NT
89	A	C	135	A	NT
90	A	C	136	A	NT
91	A	C	137	A	NT

[00374] The biological data provided by the present disclosure indicates that the compounds of the present disclosure have extremely strong inhibitory ability on HER2 kinase and HER2 expressing cells, which are beneficial to the treatment or prevention of diseases caused by abnormality of HER2 kinase. Another feature of the compound of the present disclosure is that its ability to inhibit cells with EGFR expression is relatively weak, thereby greatly reducing side effects caused by excessive inhibition of EGFR.

[00375] The above is a preferred embodiment of the present disclosure, and it should be noted that those skilled in the art can make various improvements and modifications to the embodiments of the present disclosure without departing from the principles of the present disclosure. These improvements and modifications are also considered to be within the scope of the disclosure.

CLAIMS

1. A compound of Formula (I), or a stereoisomer, hydrate, solvate, or pharmaceutically acceptable salt thereof, wherein:




Formula (I)


in the Formula (I),

X is O, or NH;

Y is N or C-Z, wherein Z is -H or -CN;



R³ is -H, halogen, C₁-C₃ alkyl, C₁-C₃ alkoxy, C₁-C₃ alkyl substituted by halogen, or C₁-C₃ alkoxy substituted by halogen;





M is O or S;

T is linear C₁-C₃ alkyl, or linear C₁-C₃ alkyl independently substituted by R⁵ and R⁶, respectively;

R⁵ and R⁶ are independently -H, halogen, C₁-C₃ alkyl, or C₁-C₃ alkyl substituted by halogen;

R⁴ is aryl, 5- to 6-membered heteroaryl, aryl substituted by 1-3 identical or different R⁷, or 5- to 6-membered heteroaryl substituted by 1-3 identical or different R⁷, wherein the heteroaryl group is a heteroaryl group containing 1-3 heteroatoms selected from N, O or S;

R⁷ is -H, halogen, amino, hydroxy, cyano, C₁-C₃ alkylthio, amino substituted with mono- or di-C₁-C₃ alkyl, C₃-C₄ cycloalkyl, unsubstituted or substituted C₁-C₆ alkyl, or unsubstituted or substituted C₁-C₆ alkoxy, wherein the substituent of the substituted C₁-C₆ alkyl is halogen, hydroxy, amino substituted with mono- or di-C₁-C₃ alkyl, or C₁-C₃ alkoxy, and wherein the substituent of the substituted C₁-C₆ alkoxy is halogen, C₁-C₃ alkoxy, or amino substituted with mono- or di-C₁-C₃ alkyl;



T₁ is linear C₁-C₈ alkyl, linear C₁-C₈ alkyl independently substituted by R⁹ and R¹⁰, respectively;

R⁹ and R¹⁰ are each independently -H, or C₁-C₃ alkyl;

R⁸ is -H, hydroxy, C₁-C₃ alkyl, C₃-C₇ cycloalkyl, C₁-C₃ alkoxy, C₁-C₃ alkylthio, 4- to 7-membered heterocyclyl or -NR¹¹R¹²;

R¹¹ and R¹² are each independently -H, C₁-C₆ alkyl, C₃-C₆ cycloalkyl, C₁-C₆ alkyl substituted by hydroxy, or C₁-C₆ alkyl substituted by C₁-C₃ alkoxy;


the 4- to 7-membered heterocyclyl is a heterocyclyl containing 1-2 heteroatoms selected from N, O or S, the heterocyclyl is unsubstituted or substituted by one or two of: C₁-C₃ alkyl, aldehyde group, C₁-C₄ alkylacyl, aminoacyl, aminoacyl wherein the amino is substituted with mono- or di-C₁-C₃ alkyl, C₁-C₃ alkylsulfonyl, and C₁-C₃ alkylsulfinyl, or the sulfur in the heterocycle is oxidized by one to two oxygen atoms.

2. The compound of Formula (I), or the stereoisomer, hydrate, solvate, or pharmaceutically acceptable salt thereof according to claim 1 wherein:


in the Formula (I),

X is NH;

Y is N;



R³ is -H, halogen, C₁-C₃ alkyl, or C₁-C₃ alkyl substituted by halogen;





M is O or S;

T is linear C₁-C₃ alkyl;

R⁴ is aryl, 5- to 6-membered heteroaryl, or aryl substituted by 1-3 identical or different R⁷, wherein the heteroaryl group is a heteroaryl group containing 1-3 heteroatoms selected from N, O or S;

R⁷ is -H, or halogen;




T₁ is linear C₁-C₈ alkyl;


R⁸ is -NR¹¹R¹²; and

R¹¹ and R¹² are each independently -H, C₁-C₆ alkyl, or C₃-C₆ cycloalkyl.

3. The compound, or stereoisomer, hydrate, solvate, or pharmaceutically acceptable salt thereof according to claim 1, wherein:



R³ is -H, fluorine, chlorine, bromine, methyl, ethyl, methoxy, ethoxy, trifluoromethyl or trifluoromethoxy;



M is O or S;



10 T is linear C₁-C₂ alkyl, or linear C₁-C₂ alkyl independently substituted by R⁵ and R⁶, respectively;

R⁵ and R⁶ are independently is -H, -F, methyl, ethyl or trifluoromethyl;

15 R⁴ is aryl, 5- to 6-membered heteroaryl, aryl substituted by 1-2 identical or different R⁷, or 5- to 6-membered heteroaryl substituted by 1-2 identical or different R⁷, wherein the aryl or heteroaryl group is phenyl, pyridyl, pyrimidinyl, thiazolyl, thienyl, pyrrolyl, thiadiazolyl, furyl, oxazolyl or isoxazolyl; and

20 R⁷ is -H, -F, -Cl, -Br, -CF₃, -OCF₃, amino, hydroxy, cyano, methylthio, ethylthio, propylthio, isopropylthio, methylamino, ethylamino, dimethylamino, diethylamino, cyclopropyl, cyclobutyl, methyl, ethyl, propyl, isopropyl, methoxy, ethoxy, propoxy, isopropoxy, hydroxymethyl, hydroxyethyl, hydroxypropyl, methylaminomethyl, methylaminoethyl, methylaminopropyl, dimethylaminomethyl, dimethylaminoethyl, dimethylaminopropyl, methoxymethyl, methoxyethyl, methoxypropyl, ethoxymethyl, ethoxyethyl, ethoxypropyl, methoxyethoxy, methoxypropoxy, ethoxyethoxy, ethoxypropoxy, methylaminoethoxy, methylaminopropoxy, ethylaminoethoxy, ethylaminopropoxy, 25 dimethylaminoethoxy, dimethylaminopropoxy, diethylaminoethoxy, or diethylaminopropoxy.

4. The compound, or stereoisomer, hydrate, solvate, or pharmaceutically acceptable salt thereof according to claim 1 or 3, wherein:



T₁ is linear C₁-C₆ alkyl, or linear C₁-C₆ alkyl independently substituted by R⁹ and R¹⁰,

respectively;


R⁹ and R¹⁰ are independently -H or methyl;

R⁸ is -H, hydroxy, methyl, ethyl, propyl, isopropyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, methoxy, ethoxy, propoxy, isopropoxy, methylthio, ethylthio, propylthio, isopropylthio, 5- to 6-membered heterocyclyl or -NR¹¹R¹²;

R¹¹ and R¹² are each independently -H, methyl, ethyl, propyl, butyl, pentyl, hexyl, isopropyl, sec-butyl, 1-ethylpropyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, hydroxyethyl, hydroxypropyl, hydroxybutyl, hydroxypentyl, hydroxyhexyl, methoxyethyl, methoxypropyl, methoxybutyl, methoxypentyl, methoxyhexyl, ethoxyethyl, ethoxypropyl, ethoxybutyl, ethoxypentyl, ethoxyhexyl, propoxyethyl, propoxypropyl, propoxybutyl, propoxypentyl, propoxyhexyl, isopropoxyethyl, isopropoxypropyl, isopropoxybutyl, isopropoxypentyl or isopropoxyhexyl;

the 5- to 6-membered heterocyclyl is a heterocyclyl containing 1-2 heteroatoms selected from N, O or S, the 5- to 6-membered heterocyclyl is unsubstituted or substituted by one or two of: methyl, ethyl, propyl, isopropyl, aldehyde group, formyl, acetyl, propionyl, butyryl, isobutyryl, aminoacyl, methylaminoacyl, dimethylaminoacyl, methylsulfonyl, ethylsulfonyl, isopropylsulfonyl, methylsulfinyl, ethylsulfinyl, or isopropylsulfinyl, or sulfur in the heterocycle is oxidized by one to two oxygen atoms;

the 5- to 6-membered heterocycle is



5. The compound, stereoisomer, hydrate, solvate, or pharmaceutically acceptable salt thereof according to any one of claims 1, 3 and 4, wherein:

R² is or ;

L₁ is or ;

T₁ is linear C₁-C₆ alkyl, or linear C₁-C₆ alkyl independently substituted by R⁹ and R¹⁰, respectively;

R⁹ and R¹⁰ are each independently -H or methyl;


R⁸ is -H, hydroxy, methyl, ethyl, propyl, isopropyl, methoxy, ethoxy, propoxy, isopropoxy, methylthio, ethylthio, propylthio, isopropylthio, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, 5- to 6-membered heterocyclyl or -NR¹¹R¹²;

R¹¹ and R¹² are each independently -H, methyl, ethyl, propyl, butyl, pentyl, hexyl,

isopropyl, sec-butyl, 1-ethylpropyl, cyclopropyl, cyclobutyl, cyclopentyl, hydroxyethyl, hydroxypropyl, hydroxybutyl, hydroxypentyl, methoxyethyl, methoxypropyl, methoxybutyl, methoxypentyl, ethoxyethyl, ethoxypropyl, ethoxybutyl, ethoxypentyl, propoxyethyl, propoxypropyl, propoxybutyl, propoxypentyl, isopropoxyethyl, isopropoxypropyl, isopropoxybutyl or isopropoxypentyl;


the 5- to 6-membered heterocyclyl is a heterocyclyl containing 1-2 heteroatoms selected from N, O or S, the 5- to 6-membered heterocyclyl is unsubstituted or substituted by one or two of: methyl, ethyl, propyl, isopropyl, aldehyde group, formyl, acetyl, propionyl, butyryl, isobutyryl, aminoacyl, methylaminoacyl, dimethylaminoacyl, methylsulfonyl, ethylsulfonyl, isopropylsulfonyl, methylsulfinyl, ethylsulfinyl, or isopropylsulfinyl, or the sulfur in the heterocycle is oxidized by one to two oxygen atoms;

the 5- to 6-membered heterocyclyl is



R^{13} is -H, amino, methylamino, dimethylamino, methyl, ethyl, propyl, or isopropyl.

6. A compound of Formula (I), or a stereoisomer, hydrate, solvate, or pharmaceutically acceptable salt thereof, wherein:




Formula (I)


in the Formula (I),

X is O, or NH;

Y is N or C-Z, wherein Z is -H or -CN;



R^3 is -H, halogen, C_1 - C_3 alkyl, C_1 - C_3 alkoxy, C_1 - C_3 alkyl substituted by halogen or C_1 - C_3 alkoxy substituted by halogen;





M is O or S;



T is linear C₁-C₃ alkyl, or linear C₁-C₃ alkyl independently substituted by R⁵ and R⁶, respectively;

R⁵ and R⁶ are each -H, halogen, C₁-C₃ alkyl, or C₁-C₃ alkyl substituted by halogen;

5 R⁴ is substituted or unsubstituted fused heteroaryl, the substituted fused heteroaryl is substituted by 1-3 identical or different R⁷, the fused ring heteroaryl group is a heteroaryl group containing 1-3 heteroatoms selected from N, O or S;

10 R⁷ is -H, halogen, amino, hydroxy, cyano, C₁-C₃ alkylthio, amino substituted with mono- or di-C₁-C₃ alkyl, C₃-C₄ cycloalkyl, unsubstituted or substituted C₁-C₆ alkyl, or unsubstituted or substituted C₁-C₆ alkoxy, wherein the substituent of the substituted C₁-C₆ alkyl is halogen, hydroxy, amino substituted with mono- or di-C₁-C₃ alkyl, or C₁-C₃ alkoxy, and wherein the substituent of the substituted C₁-C₆ alkoxy is halogen, C₁-C₃ alkoxy, or amino substituted with mono- or di-C₁-C₃ alkyl;

R² is  or 

15 L₁ is  or 

T₁ is linear C₁-C₈ alkyl, or linear C₁-C₈ alkyl independently substituted by R⁹ and R¹⁰, respectively;

R⁹ and R¹⁰ are each independently -H, or C₁-C₃ alkyl;

20 R⁸ is -H, hydroxy, C₁-C₃ alkyl, C₃-C₇ cycloalkyl, C₁-C₃ alkoxy, C₁-C₃ alkylthio, 4- to 7-membered heterocyclyl or -NR¹¹R¹²;

R¹¹ and R¹² are each independently -H, C₁-C₆ alkyl, C₃-C₆ cycloalkyl, C₁-C₆ alkyl substituted by hydroxy or C₁-C₆ alkyl substituted by C₁-C₃ alkoxy; and


25 the 4- to 7-membered heterocyclyl is a heterocyclyl containing 1-2 heteroatoms selected from N, O or S, the heterocyclyl is unsubstituted or substituted by any one or two of: C₁-C₃ alkyl, aldehyde group, C₁-C₄ alkylacyl, aminoacyl, aminoacyl wherein the amino is substituted with mono- or di-C₁-C₃ alkyl, C₁-C₃ alkylsulfonyl, C₁-C₃ alkylsulfinyl, hydroxy, halogen, C₁-C₃ hydroxyalkyl, or C₁-C₃ haloalkyl, or the sulfur in the heterocycle is oxidized by one to two oxygen atoms.

30 7. The compound of Formula (I), or the stereoisomer, hydrate, solvate, or pharmaceutically acceptable salt thereof according to claim 6 wherein:


in the Formula (I),

X is NH;

Y is N;





R³ is -H, halogen, C₁-C₃ alkyl, or C₁-C₃ alkyl substituted by halogen;



5 M is O or S;

T is linear C₁-C₃ alkyl;

R⁴ is substituted or unsubstituted fused heteroaryl, wherein the fused ring heteroaryl group is a heteroaryl group containing 1-3 heteroatoms selected from N, O or S;




T₁ is linear C₁-C₈ alkyl;

R⁸ is -NR¹¹R¹²; and

R¹¹ and R¹² are each independently -H, or C₁-C₆ alkyl.

15 8. The compound, or stereoisomer, hydrate, solvate, or pharmaceutically acceptable salt thereof according to claim 6, wherein:




R³ is -H, fluorine, chlorine, bromine, methyl, ethyl, methoxy, ethoxy, trifluoromethyl or trifluoromethoxy;



M is O or S;

T is linear C₁-C₂ alkyl, or linear C₁-C₂ alkyl independently substituted by R⁵ and R⁶, respectively;


R⁵ and R⁶ are each independently -H, -F, methyl, ethyl or trifluoromethyl;



R⁷ is -H, -F, -Cl, -Br, -CF₃, -OCF₃, amino, hydroxy, cyano, methylthio, ethylthio,

propylthio, isopropylthio, methylamino, ethylamino, dimethylamino, diethylamino, cyclopropyl, cyclobutyl, methyl, ethyl, propyl, isopropyl, methoxy, ethoxy, propoxy, isopropoxy, hydroxymethyl, hydroxyethyl, hydroxypropyl, methylaminomethyl, methylaminoethyl, methylaminopropyl, dimethylaminomethyl, dimethylaminoethyl, dimethylaminopropyl, methoxymethyl, methoxyethyl, methoxypropyl, ethoxymethyl, ethoxyethyl, ethoxypropyl, methoxyethoxy, methoxypropoxy, ethoxyethoxy, ethoxypropoxy, methylaminoethoxy, methylaminopropoxy, ethylaminoethoxy, ethylaminopropoxy, dimethylaminoethoxy, dimethylaminopropoxy, diethylaminoethoxy, or diethylaminopropoxy.

9. A compound of Formula (I), or a stereoisomer, hydrate, solvate, or pharmaceutically acceptable salt thereof, wherein:




Formula (I)


in the Formula (I),

X is O, or NH;

Y is N or C-Z, wherein Z is -H or -CN;



R³ is -H, halogen, C₁-C₃ alkyl, C₁-C₃ alkoxy, C₁-C₃ alkyl substituted by halogen or C₁-C₃ alkoxy substituted by halogen;



M is O or S;


T is linear C₁-C₃ alkyl, or linear C₁-C₃ alkyl independently substituted by R⁵ and R⁶, respectively;

R⁵ and R⁶ are each independently -H, halogen, C₁-C₃ alkyl, or C₁-C₃ alkyl substituted by halogen;

R⁴ is aryl, 5- to 6-membered heteroaryl, aryl substituted by 1-3 identical or different R⁷, or 5- to 6-membered heteroaryl substituted by 1-3 identical or different R⁷, wherein the heteroaryl group is a heteroaryl group containing 1-3 heteroatoms selected from N, O or S;

R⁷ is -H, halogen, amino, hydroxy, cyano, C₁-C₃ alkylthio, amino substituted with mono- or di-C₁-C₃ alkyl, C₃-C₄ cycloalkyl, unsubstituted or substituted C₁-C₆ alkyl, or unsubstituted

or substituted C₁-C₆ alkoxy, wherein the substituent of the substituted C₁-C₆ alkyl is halogen, hydroxy, amino substituted with mono- or di-C₁-C₃ alkyl, or C₁-C₃ alkoxy, and wherein the substituent of the substituted C₁-C₆ alkoxy is halogen, C₁-C₃ alkoxy, or amino substituted with mono- or di-C₁-C₃ alkyl;



T_1 is linear C₁-C₈ alkyl, or linear C₁-C₈ alkyl independently substituted by R^9 and R^{10} , respectively;

R^9 and R^{10} are each independently -H, or C₁-C₃ alkyl;

10 R^8 is -H, hydroxy, C₁-C₃ alkyl, C₃-C₇ cycloalkyl, C₁-C₃ alkoxy, C₁-C₃ alkylthio, 4- to 7-membered heterocyclyl or -NR¹¹R¹²;

R^{11} and R^{12} are each independently -H, C₁-C₆ alkyl, C₃-C₆ cycloalkyl, C₁-C₆ alkyl substituted by hydroxyl or C₁-C₆ alkyl substituted by C₁-C₃ alkoxy; and

15 the 4- to 7-membered heterocyclyl is a heterocyclyl containing 1-2 heteroatoms selected from N, O or S, the heterocyclyl is unsubstituted or substituted by any one or two of: C₁-C₃ alkyl, hydroxy, halogen, C₁-C₃ hydroxyalkyl, or C₁-C₃ haloalkyl.

10. The compound, or stereoisomer, hydrate, solvate, or pharmaceutically acceptable salt thereof according to any one of claims 1-9, wherein the compound is

20 1-(10-((4-(3-(trifluoromethyl)phenoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one;

1-(10-((4-phenoxyphenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one;

25 1-(10-((3-chloro-4-((3-fluorobenzyloxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one;

1-(10-((3-chloro-4-(pyridin-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one;

1-(10-((4-(m-tolyl)oxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one;

30 1-(10-((4-(3-chlorophenoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one;

1-(10-((4-(3-fluorophenoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one;

1-(10-((4-(2-fluorophenoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one;

1-(10-((4-(4-fluorophenoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one;

5 1-(10-((4-(4-chlorophenoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one;

1-(10-((4-(2-methoxyphenoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one;

1-(10-((4-(pyridin-2-yloxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one;

10 1-(10-((4-(pyridin-3-yloxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one;

1-(10-((4-(3-methoxyphenoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one;

15 1-(10-((4-(thiazol-2-yloxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one;

1-(10-((3-fluoro-4-phenoxyphenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one;

1-(10-((3-chloro-4-phenoxyphenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one;

20 1-(10-((4-((4-fluorobenzyl)oxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one;

1-(10-((4-((3-fluorobenzyl)oxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one;

25 1-(10-((4-((3-trifluoromethylbenzyl)oxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one;

1-(10-((4-(pyridin-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one;

1-(10-((4-(thiophen-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one;

30 1-(10-((4-(thiazol-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one;

1-(10-((4-(benzylthio)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one;

1-(10-((4-((3-fluorobenzyl)oxy)-3-(trifluoromethyl)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one;

1-(10-((4-((3-fluorobenzyl)oxy)-3-(methoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one;

5 1-(10-((4-((3-fluorobenzyl)oxy)-3-fluorophenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one;

1-(10-((4-((4-fluorophenyl)thio)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one;

1-(10-((2-fluoro-4-phenoxyphenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one;

10 1-(10-((4-(2-(2-(dimethylamino)ethoxy)phenoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one;

1-(10-((4-(2-(3-(dimethylamino)propoxy)phenoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one;

15 1-(10-((4-(2-(2-methoxyethoxy)phenoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one;

1-(10-(4-phenoxyphenoxy)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one;

1-(10-(3-chloro-4-(pyridin-2-ylmethoxy)phenoxy)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one;

20 1-(10-(3-chloro-4-((3-fluorobenzyl)oxy)phenoxy)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one;

1-(10-(4-(4-chlorophenoxy)phenoxy)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one;

25 1-(10-(4-(4-fluorophenoxy)phenoxy)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one;

1-(10-(4-(2,5-dichlorophenoxy)phenoxy)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one;

1-(10-(4-(pyridin-2-ylmethoxy)phenoxy)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one;

30 1-(10-(4-((2-fluorobenzyl)oxy)phenoxy)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one;

1-(10-(4-((3-fluorobenzyl)oxy)phenoxy)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one;

- 1-(10-(4-((4-fluorobenzyl)oxy)phenoxy)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)prop-2-en-1-one;
- (E)-4-(dimethylamino)-1-(10-((4-(3-(trifluoromethyl)phenoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one;
- 5 (E)-4-(dimethylamino)-1-(10-((4-phenoxyphenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one;
- (E)-1-(10-((3-chloro-4-((3-fluorobenzyl)oxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(dimethylamino)but-2-en-1-one;
- (E)-1-(10-((3-chloro-4-(pyridin-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-10 [1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(dimethylamino)but-2-en-1-one;
- (E)-4-(dimethylamino)-1-(10-((4-(m-tolyl)oxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one;
- (E)-4-(dimethylamino)-1-(10-((4-(3-chlorophenoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one;
- 15 (E)-4-(dimethylamino)-1-(10-((4-(3-fluorophenoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one;
- (E)-4-(dimethylamino)-1-(10-((4-(2-fluorophenoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one;
- (E)-4-(dimethylamino)-1-(10-((4-(4-fluorophenoxy)phenyl)amino)-2,3-dihydro-4H-20 [1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one;
- (E)-4-(dimethylamino)-1-(10-((4-(4-chlorophenoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one;
- (E)-4-(dimethylamino)-1-(10-((4-(2-methoxyphenoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one;
- 25 (E)-4-(dimethylamino)-1-(10-((4-(pyridin-2-yloxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one;
- (E)-4-(dimethylamino)-1-(10-((4-(pyridin-3-yloxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one;
- (E)-4-(dimethylamino)-1-(10-((4-(3-methoxyphenoxy)phenyl)amino)-2,3-dihydro-4H-30 [1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one;
- (E)-4-(dimethylamino)-1-(10-((4-(thiazol-2-yloxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one;
- (E)-4-(dimethylamino)-1-(10-((3-fluoro-4-phenoxyphenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one;

- (E)-4-(dimethylamino)-1-(10-((3-chloro-4-phenoxyphenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one;
- (E)-4-(dimethylamino)-1-(10-((4-((4-fluorobenzyl)oxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one;
- 5 (E)-4-(dimethylamino)-1-(10-((4-((3-fluorobenzyl)oxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one;
- (E)-4-(dimethylamino)-1-(10-((4-((3-(trifluoromethyl)benzyl)oxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one;
- (E)-4-(dimethylamino)-1-(10-((4-(pyridin-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one;
- 10 (E)-4-(dimethylamino)-1-(10-((4-(thiophen-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one;
- (E)-4-(dimethylamino)-1-(10-((4-(thiazol-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one;
- 15 (E)-4-(dimethylamino)-1-(10-((4-(benzylthio)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one;
- (E)-4-(dimethylamino)-1-(10-((4-((3-fluorobenzyl)oxy)-3-(trifluoromethyl)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one;
- 20 (E)-4-(dimethylamino)-1-(10-((4-((3-fluorobenzyl)oxy)-3-methoxyphenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one;
- (E)-4-(dimethylamino)-1-(10-((4-((3-fluorobenzyl)oxy)-3-fluorophenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one;
- (E)-4-(dimethylamino)-1-(10-((4-((4-fluorophenyl)thio)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one;
- 25 (E)-4-(dimethylamino)-1-(10-((4-(2-fluoro-5-methylphenoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one;
- (E)-4-(dimethylamino)-1-(10-((4-(5-chloro-2-fluorophenoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one;
- 30 (E)-4-(dimethylamino)-1-(10-((4-(2,5-difluorophenoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one;
- (E)-4-(dimethylamino)-1-(10-((4-(1-(3-fluorophenyl)ethoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one;
- (E)-4-(dimethylamino)-1-(10-((4-(1-(pyridin-2-yl)ethoxy)phenyl)amino)-2,3-dihydro-

4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one;
 (E)-4-(dimethylamino)-1-(10-((4-(pyridin-3-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one;
 (E)-4-(dimethylamino)-1-(10-((4-(pyridin-4-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-
 5 [1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one;
 (E)-1-(10-((4-(benzyloxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(dimethylamino)but-2-en-1-one;
 (E)-4-(dimethylamino)-1-(10-((4-((2-fluorobenzyl)oxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one;
 10 (E)-4-(dimethylamino)-1-(10-((4-((2-chlorobenzyl)oxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one;
 (E)-4-(dimethylamino)-1-(10-((4-((2-methylbenzyl)oxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one;
 (E)-4-(dimethylamino)-1-(10-((4-((2-methoxybenzyl)oxy)phenyl)amino)-2,3-dihydro-
 15 4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one;
 (E)-1-(10-((4-((3-chlorobenzyl)oxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(dimethylamino)but-2-en-1-one;
 (E)-4-(dimethylamino)-1-(10-((4-((3-methylbenzyl)oxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one;
 20 (E)-4-(dimethylamino)-1-(10-((4-((3-cyanobenzyl)oxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one;
 (E)-4-(dimethylamino)-1-(10-((4-((3-methoxybenzyl)oxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one;
 (E)-4-(dimethylamino)-1-(10-((4-((4-chlorobenzyl)oxy)phenyl)amino)-2,3-dihydro-4H-
 25 [1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one;
 (E)-4-(dimethylamino)-1-(10-((4-((4-methylbenzyl)oxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one;
 (E)-1-(10-((4-((2,5-difluorobenzyl)oxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(dimethylamino)but-2-en-1-one;
 30 (E)-1-(10-((4-((2-chloro-5-fluorobenzyl)oxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(dimethylamino)but-2-en-1-one;
 (E)-1-(10-((4-phenoxyphenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(pyrrolidin-1-yl)but-2-en-1-one;
 (E)-4-(diethylamino)-1-(10-((4-phenoxyphenyl)amino)-2,3-dihydro-4H-


[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one;
 (E)-1-(10-((4-phenoxyphenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(piperidin-1-yl)but-2-en-1-one;
 (E)-4-morpholino-1-(10-((4-phenoxyphenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one;
 5 (E)-4-(4-methylpiperazin-1-yl)-1-(10-((4-phenoxyphenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one;
 (E)-1-(10-((3-chloro-4-((3-fluorobenzyl)oxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-morpholinobut-2-en-1-one;
 10 (E)-1-(10-((2-fluoro-4-phenoxyphenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(piperidin-1-yl)but-2-en-1-one;
 (E)-1-(10-((3-fluoro-4-phenoxyphenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(piperidin-1-yl)but-2-en-1-one;
 (E)-1-(10-((4-(2-fluorophenoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-((4-methoxybutyl)amino)but-2-en-1-one;
 15 (E)-1-(10-((3-chloro-4-(pyridin-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(cyclopropyl(methyl)amino)but-2-en-1-one;
 (E)-1-(10-((3-chloro-4-(pyridin-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(cyclopropylamino)but-2-en-1-one;
 20 (E)-1-(10-((3-chloro-4-(pyridin-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(isopropylamino)but-2-en-1-one;
 (E)-1-(10-((3-chloro-4-(pyridin-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-morpholinobut-2-en-1-one;
 (E)-1-(10-((3-chloro-4-(pyridin-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(4-methylpiperazin-1-yl)but-2-en-1-one;
 25 (E)-1-(10-((3-chloro-4-(pyridin-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(diethylamino)but-2-en-1-one;
 (E)-1-(10-((3-chloro-4-(pyridin-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(pyrrolidin-1-yl)but-2-en-1-one;
 30 (E)-1-(10-((3-chloro-4-(pyridin-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(piperidin-1-yl)but-2-en-1-one;
 (E)-4-(4-acetylpiperazin-1-yl)-1-(10-((3-chloro-4-(pyridin-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-en-1-one;
 (E)-1-(10-((3-chloro-4-(pyridin-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-

[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(cyclobutylamino)but-2-en-1-one;
 (E)-1-(10-((3-chloro-4-(pyridin-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-
 [1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(cyclobutyl(methyl)amino)but-2-en-1-one;
 (E)-1-(10-((3-chloro-4-(pyridin-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-
 5 [1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(isopropyl(methyl)amino)but-2-en-1-one;
 (S,E)-1-(10-((3-chloro-4-(pyridin-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-
 [1,4]oxazino[2,3-f]quinazolin-4-yl)-3-(1-methylpyrrolidin-2-yl)prop-2-en-1-one;
 1-(10-((3-chloro-4-(pyridin-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-
 [1,4]oxazino[2,3-f]quinazolin-4-yl)but-2-yn-1-one;
 10 (R,E)-1-(10-(3-chloro-4-(pyridin-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-
 [1,4]oxazino[2,3-f]quinazolin-4-yl)-3-(1-methylpyrrolidin-2-yl)prop-2-en-1-one;
 (E)-1-(10-((3-chloro-4-(pyridin-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-
 [1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(4-hydroxypiperidin-1-yl)but-2-en-1-one;
 (E)-1-(10-((3-chloro-4-(pyridin-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-
 15 [1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(4-hydroxyl-4-methylpiperidin-1-yl)but-2-en-1-one;
 (E)-1-(10-((3-chloro-4-(pyridin-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-
 [1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(4-(hydroxymethyl)piperidin-1-yl)but-2-en-1-one;
 (E)-1-(10-((3-chloro-4-(pyridin-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-
 [1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(4-fluoropiperidin-1-yl)but-2-en-1-one;
 20 (E)-1-(10-((3-chloro-4-(pyridin-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-
 [1,4]oxazino[2,3-f]quinazolin-4-yl)-4-((2-methoxyethyl)(methyl)amino)but-2-en-1-one;
 (E)-1-(10-((3-chloro-4-(pyridin-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-
 [1,4]oxazino[2,3-f]quinazolin-4-yl)-4-((3-hydroxypropyl)(methyl)amino)but-2-en-1-one;
 (S,E)-1-(10-((3-chloro-4-(pyridin-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-
 25 [1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(2,4-dimethylpiperazin-1-yl)but-2-en-1-one;
 (E)-1-(10-((3-chloro-4-(pyridin-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-
 [1,4]oxazino[2,3-f]quinazolin-4-yl)-4-((2-hydroxyethyl)(methyl)amino)but-2-en-1-one;
 (R,E)-1-(10-((3-chloro-4-(pyridin-2-ylmethoxy)phenyl)amino)-2,3-dihydro-4H-
 [1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(2,4-dimethylpiperazin-1-yl)but-2-en-1-one;
 30 (E)-1-(10-((4-([1,2,4]triazolo[1,5-a]pyridin-7-yloxy)-3-methylphenyl)amino)-2,3-
 dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(dimethylamino)but-2-en-1-one;
 (E)-1-(10-((4-([1,2,4]triazolo[1,5-a]pyridin-7-yloxy)-3-chlorophenyl)amino)-2,3-
 dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(dimethylamino)but-2-en-1-one;
 (E)-1-(10-((4-([1,2,4]triazolo[1,5-a]pyridin-7-yloxy)-3-chlorophenyl)amino)-2,3-

dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(4-methylpiperazin-1-yl)but-2-en-1-one;
 (E)-1-(10-((4-([1,2,4]triazolo[1,5-a]pyridin-7-yloxy)-3-chlorophenyl)amino)-2,3-
 dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(isopropyl(methyl)amino)but-2-en-1-one;
 (E)-1-(10-((4-([1,2,4]triazolo[1,5-a]pyridin-7-yloxy)-3-chlorophenyl)amino)-2,3-
 5 dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(piperidin-1-yl)but-2-en-1-one;
 (E)-1-(10-((4-([1,2,4]triazolo[1,5-a]pyridin-7-yloxy)-3-chlorophenyl)amino)-2,3-
 dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(cyclobutyl(methyl)amino)but-2-en-1-one;
 (E)-1-(10-((4-([1,2,4]triazolo[1,5-a]pyridin-7-yloxy)-3-chlorophenyl)amino)-2,3-
 dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-((2-methoxyethyl)(methyl)amino)but-2-en-
 10 1-one;
 (E)-1-(10-((4-([1,2,4]triazolo[1,5-a]pyridin-7-yloxy)-3-chlorophenyl)amino)-2,3-
 dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(diethylamino)but-2-en-1-one;
 (E)-1-(10-((4-([1,2,4]triazolo[1,5-a]pyridin-7-yloxy)-3-methylphenyl)amino)-2,3-
 dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(4-methylpiperazin-1-yl)but-2-en-1-one;
 15 (E)-1-(10-((4-([1,2,4]triazolo[1,5-a]pyridin-7-yloxy)-3-methylphenyl)amino)-2,3-
 dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(isopropyl(methyl)amino)but-2-en-1-one;
 (E)-1-(10-((4-([1,2,4]triazolo[1,5-a]pyridin-7-yloxy)-3-methylphenyl)amino)-2,3-
 dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(piperidin-1-yl)but-2-en-1-one;
 (E)-1-(10-((4-([1,2,4]triazolo[1,5-a]pyridin-7-yloxy)-3-methylphenyl)amino)-2,3-
 20 dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(cyclobutyl(methyl)amino)but-2-en-1-one;
 (E)-1-(10-((4-([1,2,4]triazolo[1,5-a]pyridin-7-yloxy)-3-methylphenyl)amino)-2,3-
 dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-((2-methoxyethyl)(methyl)amino)but-2-en-
 1-one; or
 (E)-1-(10-((4-([1,2,4]triazolo[1,5-a]pyridin-7-yloxy)-3-methylphenyl)amino)-2,3-
 25 dihydro-4H-[1,4]oxazino[2,3-f]quinazolin-4-yl)-4-(diethylamino)but-2-en-1-one.


11. A method for preparing the compound as defined in any one of claims 1-10, characterized in that it comprises the following steps:

the compound represented by Formula (I) is prepared by the reaction of $R^2C(O)Cl$ with
 30 the compound represented by Formula (VIII) or by the chlorinating reaction of R^2COOH
 followed by the reaction with the compound represented by Formula (VIII)




when R^2 is $\text{---}L_1-T_1-R^8$, wherein L_1 is $\text{---}\text{CH=CH}\text{---}$, and R^8 is $-NR^{11}R^{12}$, the compound represented by Formula (I) is prepared by reacting the compound represented by

Formula (VIII) with $\text{O=C(Cl)-CH=CH-T}_1\text{Br}$ and $HNR^{11}R^{12}$




or when R^2 is $\text{---}L_1-R^8$ or $\text{---}L_1-T_1-R^8$, wherein L_1 is $\text{---}\text{CH=CH}\text{---}$, the compound represented by Formula (I) is prepared by reacting the compound represented by

Formula (IX) with $OHC-R^8$ or $OHC-T_1-R^8$



$R^1, R^2, R^8, R^9, R^{10}, R^{11}, R^{12}, X, Y, L_1$ and T_1 are as defined in any one of claims 1-9.


12. A compound represented by Formula (VIII),



wherein R^1, X and Y are as defined in any one of claims 1-9.

13. A method for preparing the compound of claim 12, characterized in that X and R^1 are as defined in any one of claims 1-9, and Y is N , and the preparation method thereof comprises the following steps:


method A:



2,3,4,9-tetrahydro-10H-[1,4]oxazino[2,3-f]quinazolin-10-one represented by Formula (V), and R¹XH are fully contacted with Castros reagent to afford the compound represented by

5 Formula (VIII);


method B:



2,3,4,9-tetrahydro-10H-[1,4]oxazino[2,3-f]quinazolin-10-one represented by Formula (V) is fully contacted with Castros reagent to afford the compound represented by Formula (VI), and the compound of Formula (VI) is further reacted with R¹XH to afford the compound represented by Formula (VIII);

10

or method C:



the compound represented by Formula (II) is reacted with a chlorinating reagent and then reacted with R¹XH to afford the compound represented by Formula (VII), and the compound represented by Formula (VII) is further subjected to reduction cyclization reaction to afford the compound represented by Formula (VIII).

15

14. The compound of Formula (I) according to any one of claims 1-10, or the pharmaceutically acceptable salt, stereoisomer, hydrate or solvate thereof, wherein, the pharmaceutically acceptable salt of the compound is selected from one or more of the following salts of the said compound: hydrochloride salt, hydrobromide salt, hydroiodide salt, perchlorate


20

salt, sulfate salt, nitrate salt, phosphate salt, formate salt, acetate salt, propionate salt, glycolate salt, lactate salt, succinate salt, maleate salt, tartrate salt, malate salt, citrate salt, fumarate salt, gluconate salt, benzoate salt, mandelate salt, methanesulfonate salt, isethionate salt, benzenesulfonate salt, oxalate salt, palmitate salt, 2-naphthalenesulfonate salt, p-toluenesulfonate salt, cyclohexylsulfamate salt, salicylate salt, hexonate salt, trifluoroacetate salt, aluminum salt, calcium salt, chloroprocaine salt, choline salt, diethanolamine salt, ethylenediamine salt, lithium salt, magnesium salt, potassium salt, sodium salt and zinc salt.


15. A pharmaceutical composition for treating one or more diseases related to tyrosine
10 kinases EGFR, HER2, HER3 and HER4, which comprises the compound of Formula (I) or the pharmaceutically acceptable salt, stereoisomer, hydrate or solvate thereof as defined in any one of claims 1-10 and one or more pharmaceutically acceptable carriers or excipients, wherein the one or more diseases are ocular fundus disease, xerophthalmia, psoriasis, leucoderma, dermatitis, alopecia areata, rheumatoid arthritis, colitis, multiple sclerosis, systemic lupus
15 erythematosus, Crohn's disease, atherosclerosis, pulmonary fibrosis, liver fibrosis, myelofibrosis, non-small cell lung cancer, small cell lung cancer, breast cancer, pancreatic cancer, glioma, glioblastoma, ovarian cancer, cervical cancer, colorectal cancer, melanoma, endometrial cancer, prostate cancer, bladder cancer, leukemia, gastric cancer, liver cancer, gastrointestinal stromal tumor, thyroid cancer, chronic myeloid leukemia, acute myeloid
20 leukemia, non-Hodgkin's lymphoma, nasopharyngeal cancer, esophageal cancer, brain tumor, B-cell and T-cell lymphoma, lymphoma, multiple myeloma, biliary cancer, sarcoma, or cholangiocarcinoma.

16. A pharmaceutical composition comprising the compound of formula (I) or the
25 pharmaceutically acceptable salt, stereoisomer, hydrate, or solvate thereof as defined in any one of claims 1-10, one or more other therapeutic agents, and one or more pharmaceutically acceptable carriers or excipients, wherein the one or more other therapeutic agents comprise a kinase inhibitory agent, an immunosuppressant, an anticancer agent, an anti-viral agent, anti-inflammatory agent, antifungal agent, antibiotic, or an anti-vascular hyper proliferation
30 compound.

17. A compound having the following structure:




18. A method of preparing the compound of claim 17, which comprises the steps of:



5 wherein,


in said step b), 5-(2-hydroxyethoxy)-6-nitroquinazolin-4(3H)-one represented by Formula (II) is contacted with a chlorinating reagent to afford 5-(2-chloroethoxy)-6-nitroquinazolin-4(3H)-one represented by Formula (III).

10 19. The compound according to claim 10 having the following structure:




or the pharmaceutically acceptable salt thereof.

20. The compound according to claim 10 having the following structure:




15 or the pharmaceutically acceptable salt thereof.

21. The compound according to claim 10 having the following structure:




or the pharmaceutically acceptable salt thereof.

22. The compound according to claim 10 having the following structure:



5 or the pharmaceutically acceptable salt thereof.

23. The compound according to claim 10 having the following structure:



or the pharmaceutically acceptable salt thereof.

10 24. A pharmaceutical composition comprising the compound of Formula (I) or the pharmaceutically acceptable salt, stereoisomer, hydrate, or solvate thereof as defined in any one of claims 1-10, and one or more pharmaceutically acceptable carriers or excipients.

15 25. A pharmaceutical composition comprising the compound or the pharmaceutically acceptable salt thereof as defined in any one of claims 19-23, and one or more pharmaceutically acceptable carriers or excipients.

20 26. A pharmaceutical composition comprising the compound or the pharmaceutically acceptable salt thereof as defined in any one of claims 19-23, one or more other therapeutic agents, and one or more pharmaceutically acceptable carriers or excipients, wherein the one or more other therapeutic agents comprise a kinase inhibitory agent, an immunosuppressant, an anticancer agent, an anti-viral agent, anti-inflammatory agent, antifungal agent, antibiotic, or an

anti-vascular hyper proliferation compound.

27. A pharmaceutical composition for treating one or more diseases related to tyrosine kinases EGFR, HER2, HER3 and HER4, which comprises the compound or the pharmaceutically acceptable salt thereof as defined in any one of claims 19-23 and one or more pharmaceutically acceptable carriers or excipients, wherein the one or more diseases are ocular fundus disease, xerophthalmia, psoriasis, leucoderma, dermatitis, alopecia areata, rheumatoid arthritis, colitis, multiple sclerosis, systemic lupus erythematosus, Crohn's disease, atherosclerosis, pulmonary fibrosis, liver fibrosis, myelofibrosis, non-small cell lung cancer, small cell lung cancer, breast cancer, pancreatic cancer, glioma, glioblastoma, ovarian cancer, cervical cancer, colorectal cancer, melanoma, endometrial cancer, prostate cancer, bladder cancer, leukemia, gastric cancer, liver cancer, gastrointestinal stromal tumor, thyroid cancer, chronic myeloid leukemia, acute myeloid leukemia, non-Hodgkin's lymphoma, nasopharyngeal cancer, esophageal cancer, brain tumor, B-cell and T-cell lymphoma, lymphoma, multiple myeloma, biliary cancer, sarcoma, or cholangiocarcinoma.

28. Use of the compound of Formula (I), or the stereoisomer, hydrate, solvate, or pharmaceutically acceptable salt thereof as defined in any one of claims 1-10 to treat one or more diseases related to tyrosine kinases EGFR, HER2, HER3 and HER4, wherein the one or more diseases are ocular fundus disease, xerophthalmia, psoriasis, leucoderma, dermatitis, alopecia areata, rheumatoid arthritis, colitis, multiple sclerosis, systemic lupus erythematosus, Crohn's disease, atherosclerosis, pulmonary fibrosis, liver fibrosis, myelofibrosis, non-small cell lung cancer, small cell lung cancer, breast cancer, pancreatic cancer, glioma, glioblastoma, ovarian cancer, cervical cancer, colorectal cancer, melanoma, endometrial cancer, prostate cancer, bladder cancer, leukemia, gastric cancer, liver cancer, gastrointestinal stromal tumor, thyroid cancer, chronic myeloid leukemia, acute myeloid leukemia, non-Hodgkin's lymphoma, nasopharyngeal cancer, esophageal cancer, brain tumor, B-cell and T-cell lymphoma, lymphoma, multiple myeloma, biliary cancer, sarcoma, or cholangiocarcinoma.

29. Use of the compound, or the pharmaceutically acceptable salt thereof as defined in any one of claims 19-23 to treat one or more diseases related to tyrosine kinases EGFR, HER2, HER3 and HER4, wherein the one or more diseases are ocular fundus disease, xerophthalmia, psoriasis, leucoderma, dermatitis, alopecia areata, rheumatoid arthritis, colitis, multiple sclerosis, systemic lupus erythematosus, Crohn's disease, atherosclerosis, pulmonary fibrosis,

liver fibrosis, myelofibrosis, non-small cell lung cancer, small cell lung cancer, breast cancer, pancreatic cancer, glioma, glioblastoma, ovarian cancer, cervical cancer, colorectal cancer, melanoma, endometrial cancer, prostate cancer, bladder cancer, leukemia, gastric cancer, liver cancer, gastrointestinal stromal tumor, thyroid cancer, chronic myeloid leukemia, acute myeloid leukemia, non-Hodgkin's lymphoma, nasopharyngeal cancer, esophageal cancer, brain tumor, B-cell and T-cell lymphoma, lymphoma, multiple myeloma, biliary cancer, sarcoma, or cholangiocarcinoma.

30. Use of the compound of Formula (I), or the stereoisomer, hydrate, solvate, or pharmaceutically acceptable salt thereof as defined in any one of claims 1-10 in the manufacture of a medicament for the treatment of one or more diseases related to tyrosine kinases EGFR, HER2, HER3 and HER4, wherein the one or more diseases are ocular fundus disease, xerophthalmia, psoriasis, leucoderma, dermatitis, alopecia areata, rheumatoid arthritis, colitis, multiple sclerosis, systemic lupus erythematosus, Crohn's disease, atherosclerosis, pulmonary fibrosis, liver fibrosis, myelofibrosis, non-small cell lung cancer, small cell lung cancer, breast cancer, pancreatic cancer, glioma, glioblastoma, ovarian cancer, cervical cancer, colorectal cancer, melanoma, endometrial cancer, prostate cancer, bladder cancer, leukemia, gastric cancer, liver cancer, gastrointestinal stromal tumor, thyroid cancer, chronic myeloid leukemia, acute myeloid leukemia, non-Hodgkin's lymphoma, nasopharyngeal cancer, esophageal cancer, brain tumor, B-cell and T-cell lymphoma, lymphoma, multiple myeloma, biliary cancer, sarcoma, or cholangiocarcinoma.

31. Use of the compound or the pharmaceutically acceptable salt thereof as defined in any one of claims 19-23 in the manufacture of a medicament for the treatment of one or more diseases related to tyrosine kinases EGFR, HER2, HER3 and HER4, wherein the one or more diseases are ocular fundus disease, xerophthalmia, psoriasis, leucoderma, dermatitis, alopecia areata, rheumatoid arthritis, colitis, multiple sclerosis, systemic lupus erythematosus, Crohn's disease, atherosclerosis, pulmonary fibrosis, liver fibrosis, myelofibrosis, non-small cell lung cancer, small cell lung cancer, breast cancer, pancreatic cancer, glioma, glioblastoma, ovarian cancer, cervical cancer, colorectal cancer, melanoma, endometrial cancer, prostate cancer, bladder cancer, leukemia, gastric cancer, liver cancer, gastrointestinal stromal tumor, thyroid cancer, chronic myeloid leukemia, acute myeloid leukemia, non-Hodgkin's lymphoma, nasopharyngeal cancer, esophageal cancer, brain tumor, B-cell and T-cell lymphoma, lymphoma, multiple myeloma, biliary cancer, sarcoma, or cholangiocarcinoma.

