



(86) Date de dépôt PCT/PCT Filing Date: 2002/06/14
(87) Date publication PCT/PCT Publication Date: 2002/12/27
(45) Date de délivrance/Issue Date: 2008/12/16
(85) Entrée phase nationale/National Entry: 2003/11/27
(86) N° demande PCT/PCT Application No.: NL 2002/000391
(87) N° publication PCT/PCT Publication No.: 2002/103012
(30) Priorités/Priorities: 2001/06/15 (US09/882,621);
2001/06/15 (EP01202304.0)

(51) Cl.Int./Int.Cl. *C12N 7/00* (2006.01),
C07K 14/005 (2006.01), *C12N 15/10* (2006.01),
C12N 15/63 (2006.01), *C40B 30/04* (2006.01),
C40B 40/02 (2006.01), *C40B 50/06* (2006.01),
C07K 16/00 (2006.01), *C07K 19/00* (2006.01)

(72) Inventeurs/Inventors:
HOUTZAGER, ERWIN, NL;
LOGTENBERG, TON, NL;
DE KRUIF, CORNELIS ADRIAAN, NL

(73) Propriétaire/Owner:
CRUCCELL HOLLAND B.V., NL

(74) Agent: BORDEN LADNER GERVAIS LLP

(54) Titre : PHAGES CHIMERIQUES
(54) Title: CHIMAERIC PHAGES

(57) **Abrégé/Abstract:**

The invention relates to the field of generating helper phages and phage display libraries for the identification of binding molecules. The invention provides chimaeric phages having a coat comprising a protein mixture comprising a fusion protein wherein a proteinaceous molecule is fused to a functional form of a phage coat protein and a mutant form of the phage coat protein, the mutant form being impaired in binding to a host cell receptor. The invention further provides new phage collections, novel helper phages and methods and means for producing chimaeric phages, infectious phages and helper phages according to the invention.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
27 December 2002 (27.12.2002)

PCT

(10) International Publication Number
WO 02/103012 A1

(51) International Patent Classification⁷: **C12N 15/10**,
G01N 33/68, C12N 7/00

(21) International Application Number: PCT/NL02/00391

(22) International Filing Date: 14 June 2002 (14.06.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
01202304.0 15 June 2001 (15.06.2001) EP
09/882,621 15 June 2001 (15.06.2001) US

(71) Applicant (*for all designated States except US*): **CRU-CELL HOLLAND B.V.** [NL/NL]; Archimedesweg 4, NL-2333 CN Leiden (NL).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **HOUTZAGER, Erwin** [NL/NL]; Koenestraat 13, NL-3958 XD Amerongen (NL). **LOGTENBERG, Ton** [NL/NL]; Dorpsstraat 19, NL-3433 CH Nieuwegein (NL). **DE KRUIF, Cornelis, Adriaan** [NL/NL]; Zilvergeldstraat 11, NL-3513 VN Utrecht (NL).

(74) Agent: **PRINS, A., W.**; Nieuwe Parklaan 97, NL-2587 BN Den Haag (NL).

(81) Designated States (*national*): AE, AG, AL, AM, AT (utility model), AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ (utility model), CZ, DE (utility model), DE, DK (utility model), DK, DM, DZ, EC, EE (utility model), EE, ES, FI (utility model), FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK (utility model), SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: CHIMAERIC PHAGES

(57) Abstract: The invention relates to the field of generating helper phages and phage display libraries for the identification of binding molecules. The invention provides chimaeric phages having a coat comprising a protein mixture comprising a fusion protein wherein a proteinaceous molecule is fused to a functional form of a phage coat protein and a mutant form of the phage coat protein, the mutant form being impaired in binding to a host cell receptor. The invention further provides new phage collections, novel helper phages and methods and means for producing chimaeric phages, infectious phages and helper phages according to the invention.



WO 02/103012 A1

Title: Chimaeric phages

FIELD OF THE INVENTION

5 The present invention relates to the fields of molecular biology and immunology. The invention in particular relates to the field of generating helper phages and phage display libraries for the identification of binding molecules.

10

BACKGROUND OF THE INVENTION

 An individual needs to have a dynamic immune system that is able to adapt rapidly to respond adequately to potentially harmful microorganisms and to respond to the exposure of a highly diverse and continuously changing environment. Higher organisms have evolved specialized molecular mechanisms to ensure the implementation of clonally-distributed, highly diverse repertoires of antigen-receptor molecules expressed by cells of the immune system: immunoglobulin (Ig) molecules on B lymphocytes and T cell receptors on T lymphocytes. A primary repertoire of (generally low affinity) Ig receptors is established during B cell differentiation in the bone marrow as a result of rearrangement of germ line-encoded gene segments. Further refinement of Ig receptor specificity and affinity occurs in peripheral lymphoid organs where antigen-stimulated B lymphocytes activate a somatic hypermutation machinery that specifically targets the immunoglobulin variable (V) regions. During this process, B cell clones with mutant Ig receptors of higher affinity for the inciting antigen are stimulated into clonal proliferation and maturation into antibody-secreting plasma cells (reviewed in Berek and Milstein, 1987).

15
20
25
30
35

Recombinant DNA technology has been used to mimic many aspects of the processes that govern the generation and selection of natural human antibody repertoires (reviewed in Winter and Milstein. 1991; Vaughan et al. 1998). The construction of large repertoires of antibody fragments (such as Fab fragments or single chain Fv fragments, scFv's) expressed on the surface of filamentous phage particles and the selection of such phages by "panning" on antigens has been developed as a versatile and rapid method to obtain antibodies of desired specificities (reviewed in Burton and Barbas. 1994). A subsequent optimization of the affinity of individual phage antibodies was achieved by creating mutant antibody repertoires of the selected phages and sampled for higher affinity descendants by selection for binding to antigen under more stringent conditions (reviewed in Hoogenboom. 1994).

M13 and M13-derived phages (sometimes also called viruses) are filamentous phages that can selectively infect F-pili bearing (F^+) *Escherichia coli* (*E.coli*) cells. The phage genome encodes 11 proteins, while the phage coat itself consists of 5 of these proteins: gene3, -6, -7, -8 and -9 (g3, g6, g7, g8 and g9) proteins that are bound to and that protect the (circular) single stranded DNA (ssDNA) of the viral genome. The life cycle of the virus can be subdivided into different phases. The g3 protein (g3p) of M13 phages and M13-derivatives comprises three functional domains: D1, D2 and D3, linked by two glycin-rich linkers. An alternative nomenclature for g3p domains has also been generally accepted, in which D1, D2 and D3 are named N1, N2 and CT, respectively. The N-terminal D1/D2 regions interact with the C-terminal D3 region as has been found by Chatellier et al. (1999) using several deletion mutants of g3p. Considering that functionality of a D3 domain of the protein is required for assembly of stable phages,

a less-, or non-infectious mutant of said phage coat protein preferably comprises a D3 region of said g3p, or comprises a functional part, derivative and/or analogue of said D3 region. The D3 domain is thought to bind to DNA
5 inside the viral particle. Loss of the D3 domain functionally results in rare phage-like particles that are very long and very fragile (Pratt et al. 1969; Crissman and Smith. 1984; Rakonjac and Model. 1998). The D1 and D2 domain are thought to interact with each other until the
10 phage binds to the bacteria, while D1/D2 also interact with D3 at certain stages (Chatellier et al. 1999). The linkers present in g3p between D1, D2 and D3 apparently also play a role in infectivity of the phage particle (Nilsson et al. 2000). Studies in which a protease
15 cleavage site was introduced between D1 and D2 showed that after cleavage the phage particle became non-infectious (Kristensen and Winter. 1998). Functional analysis of g3p showed that of the g3p N-terminal regions, the D1 domain is essential for infection. Loss of this domain results in
20 phages that cannot infect bacteria (Lubkowski et al. 1998; Nelson et al. 1981; Deng et al. 1999; Riechmann and Holliger. 1997; Holliger and Riechmann. 1997). It has been shown that the D2 domain interacts with the D1 domain of g3p on the phage (Figure 1). Due to competition of
25 proteins located on the F-pilus (on F⁺ bacteria) that have higher affinity for D2 than for D1, the D1 and D2 domain of the g3p dissociate from each other. The binding of D2 to the F-pilus results in a process that leads to retraction of the F-pilus towards the *E.coli* cell
30 membrane. Due to this process the phage particle comes in close contact with the bacterial membrane. Now the dissociated D1 domain can interact with bacterial proteins such as the TolA receptor, leading to the introduction of the phage DNA into the *E.coli* cell (Lubkowski et al.
35 1999). The fact that removal of the D2 domain does not prevent infection but enables phages to infect *E.coli*

lacking F-pili (Riechmann and Holliger. 1997; Deng et al. 1999) shows that the presence of the D2 domain increases specificity and that D2 has an important role in preventing F-pili independent infections. The binding of
5 D1 to the specific receptors on the surface of the *E.coli* cell (a feature that is not F⁺-specific) is represented in Figure 2. This process triggers the injection of the viral genome into the bacterium (as depicted in Figure 3).
Although loss of the D2 domain results in the formation of
10 phage particles that can infect *E.coli* in a somewhat reduced specific manner, it appears that the level of infections from such a population of phages is significantly reduced. After infection of an *E.coli* by a phage particle, the ssDNA of the virus becomes double
15 stranded due to the action of a number of bacterial enzymes. The double stranded phage genome now serves as a template for the transcription and translation of all 11 genes located on the phage genome. Besides these protein-encoding regions, the phage genome contains an intergenic
20 region: the F1-origin of replication initiation (F1-ORI). The DNA sequence of this F1-ORI can be divided in 2 separate subregions. One subregion is responsible for the initiation and termination of the synthesis of ssDNA via the so-called 'rolling circle mechanism' and the other
25 subregion is responsible for the packaging initiation of the formed circular ssDNA leading to the formation and release of new virus particles.

It has been shown that polypeptides, such as stretches of amino acids, protein parts or even entire
30 proteins can be added by means of molecular genetics to the terminal ends of a number of particle coat proteins, without disturbing the functionality of these proteins in the phage life cycle (Smith. 1985; Cwirla et al. 1990; Devlin et al. 1990; Bass et al. 1990; Felici et al. 1993;
35 Luzzago et al. 1993).

This feature enables investigators to display peptides or proteins on phages, resulting in the generation of peptide- or protein expression phage display libraries. One of the proteins that has been used in the art to fuse with polypeptides for phage display purposes, is the g3 protein (g3p), which is a coat protein that is required for an efficient and effective infectivity and subsequent entry of the viral genome into the *E.coli* cell.

For the production of phages that display polypeptides fused to the g3p coat protein, investigators in the art introduced a plasmid together with the phage genome in *E.coli* cells. This plasmid contains an active promoter upstream of an in-frame fusion between the g3 encoding gene and a gene of interest (X) encoding, for instance, polypeptides such as proteins such as antibodies or fragments such as Fab fragments or scFv's. The introduction of this plasmid together with the genome of the helper phage in an *E.coli* cell results in the generation of phages that contain on their coat either the wild type g3p from the viral genome, the fusion product g3p-X from the plasmid or a mixture of the two, since one phage particle carries five g3p's on its surface. The process of g3p or g3p-X incorporation is generally random. The presence of an F1-ORI sequence in the g3p-X expression vector (plasmid) misleads the phage synthesis machinery in such a way that two types of circular ssDNA are formed: one is derived from the genome of the phage and the other is derived from the expression vector. During the synthesis of new phages the machinery is unable to distinguish the difference between these two forms of ssDNA resulting in the synthesis of a mixed population of phages, one part containing the phage genome and one part harbouring the vector DNA. Due to these processes the mixture contains at least some phages in which the phenotypic information on the outside (the g3p-X fusion protein) is conserved within the genotypic information

inside the particle (the g3p-X expression vector). An infectious wild type phage and a phage carrying a fusion protein attached to g3p are depicted in Figure 4. The art teaches that there are several problems that concern the
5 use of these basic set-ups.

The high level of genotypic wild type phages in phage populations grown in bacteria that contain both the phage genome and the expression vector compelled investigators to design mutant Fl-ORI sequences in M13 genomes. Such
10 mutant M13-strains are less effective in incorporating their genome in phage particles during phage assembly, resulting in an increased percentage of phages containing vector sequences when co-expressed. These mutant phages, such as the commercially available strains R408, VCSM13
15 and M13K07, are called "helper phages". The genome of these helper phages may contain genes required to assemble new (helper-) phages in *E.coli* and to subsequently infect new F-pili expressing *E.coli*. Both VCSM13 and M13K07 were provided with an origin of replication initiation (ORI) of
20 the P15A type that results in the multiplication of the viral genome in *E.coli*. Moreover, the ORI introduction ensures that after cell division the old and newly formed *E.coli* contains at least one copy of the viral genome.

It was suggested and finally proven by several
25 investigators that the introduction of plasmids containing a g3p-scFv fusion product together with the genome of the helper phages in *E.coli* cells results in approximately 99% of newly formed phages that harbor the g3p-scFv fusion protein expression plasmid but nevertheless lack the g3p-scFv fusion on its surface (Beekwilder et al. 1999). The
30 absence of g3p-X is a significant disadvantage in the use of display libraries for the identification of specific proteins or peptides such as scFv's that bind to a target of interest (such as tumor antigens). It implies that in
35 the case of phage display libraries at least a 100-fold excess of produced phages must be used in an experiment in

order to perform a selection with all possible fusion proteins present. The art teaches that this overload of relatively useless phages in an experiment leads to (too) many false positives. For instance, at least 10^{12} phages should be added to a panning experiment in order to have 1 copy of each possible fusion present in the experiment, since such a library contains approximately 10^{10} different g3p-scFv fusions (1%). The phages in this approximate 1% express generally only 1 g3p-scFv fusion on their coat together with four normal g3p's (no fusions), while the rest of the helper phages (approximately 99%) express five g3p's and no g3p-scFv fusions. To ensure, theoretically, the presence of 100 copies of each separate fusion protein in a panning experiment, one needs to use approximately 10^{14} phages in such an experiment. Persons skilled in the art generally attempt to use an excess of at least 100-fold of each single unique fusion protein, to ensure the presence of sufficient numbers of each separate fusion and not to lose relevant binders too quickly in first panning rounds. That number of phages (10^{14}) is more or less the maximum of phage particles that a milliliter (ml) can hold. The viscosity of such a solution is extremely high and therefore relatively useless. Especially when ELISA panning strategies are used (in which the volume of one well is only 200 μ l) such libraries cannot be used.

In addition to these problems, it is assumed that generally, depending on the antigen and the stringency of washing procedures, an average of 1 in every 10^7 phages will bind to the antigen due to a-specific binding. Thus, for the application of 10^{12} scFv expressing input phages (1%) to a panning procedure, one has to add approximately 10^{14} phages (99% of which do not express a scFv fragment). It is generally assumed that from these 10^{12} phages approximately 10^4 particles might be putatively interesting phages. However, depending on the washing conditions, the number of calculated background phages

that are normally found by using libraries present in the art after one round of panning, was approximately 10^6 - 10^7 while only a few of these phages appear to be relevant binders. This is one of the most significant problems recognized in the art: too many background phages show up as initial binders in the phage mix after the first round of panning, while only a few significant and interesting binders are present in this mix. So, the absolute number of isolated phages after one round of panning is clearly too high (10^6 - 10^7). Moreover, also in subsequent rounds of panning non-specific background phages remain present. In libraries used in the art, most of these non-specific binders will amplify on bacteria upon amplification to continue appearing in a second round of panning.

Therefore, the art teaches that the background level of a-specifically binding phages and the total number of phages per ml in these types of libraries is unacceptably high and remains high during subsequent rounds of panning.

A possibility that was suggested by investigators in the art as a solution to the problem of obtaining too many background phages that lack a g3p-X fusion was to remove the g3p-encoding gene entirely from the helper phage genome. In principle, this system ensures that during phage synthesis in an *E.coli* cell (that received the g3-less phage genome and a g3p-X fusion protein expression vector) only g3p-X proteins are incorporated in the newly formed phage coat. By doing so, each synthesized phage will express five copies of the g3p-X fusion product and hardly any phages are synthesized that express the g3p alone or that express less than five g3p-X fusions. R408-d3 and M13MDΔD3 are two examples of g3-minus helper phages (Dueñas and Borrebaeck. 1995; Rakonjac et al. 1997). Because the genome of these phages is not capable of supporting g3p synthesis, phage particles that carry less than five g3p-X fusion proteins can hardly be formed, or, if formed, are found to be non-infectious due to

instability, since the art teaches that five g3p's are necessary to ensure a stable phage particle.

To produce helper phages that do not contain the g3 gene, but that are nevertheless infectious and that can be used to generate libraries of phages that carry five g3p-X fusion proteins, and that lack phages with less than five g3p-X fusions, it was recognized in the art that an external source for g3p was required. Such a source can be a vector without F1-ORI but that nevertheless contains an active promoter upstream of the full open reading frame (ORF) of g3. One major problem that is recognized by persons skilled in the art is that after the generation step of producing newly formed helper phages lacking a g3 gene, the yield is dramatically low. In fact, the yield of all described systems is below 10^{10} phages per liter, meaning that for a library of 10^{10} individual clones at least 100 liter of helper phage culture is necessary (NB: the helper phages need to be purified) in order to grow the library once. The art thus teaches that phage libraries generated with such low titers of helper phages are not useful for phage display purposes and that therefore these libraries cannot be used for panning experiments. One method for complementation of g3p deletion phages was presented recently, in which wild-type g3p was provided by a nucleic acid encoding the wild-type g3p, wherein the nucleic acid was stably incorporated into the host cell genome (Rondot et al. 2001).

Phages that express deleted g3p's fused to heterologous proteins have been generated. For the construction of most conventional Fab libraries and some scFv phage display libraries the D1 domain and parts of the D2 domain were removed to ensure a shorter fusion protein, which was considered in the art as a product that could be translated easier than a full length g3p linked to a full length Fab fragment. The shorter g3p part would not prevent the generation of a viable and useful helper

phage. Of course, such phages still depend on full length g3p's that are present on their surface next to the deleted g3p fusion with the Fab fragment for functional infectivity of *E.coli* cells. Also, phages that express
5 deleted g3p's fused to ligand-binding proteins have been generated that depend on their infectious abilities on antigens that were fused to the parts of g3p that were missing from the non-infectious phage particle (Krebber et al. 1997; Spada et al. 1997). These particles depend for
10 their infectivity on an interaction between the ligand-binding protein, such as an antibody or a fragment thereof and their respective ligand (or antigen). However, this interaction-dependency reduces the efficiency of infection, due to elimination of a direct linkage between
15 the g3p domains, and a general inhibitory effect of the soluble N-terminal part of g3p coupled to the antigen.

The g3-minus helper phages R408-d3 and M13MDΔD3 mentioned above, lack in their genome a bacterial ORI and a selection marker. The absence of a selection marker in
20 the g3-minus genomes has a significant effect on the production scale of helper phages, because it results in an overgrowth of bacteria that do not contain the helper phage genome. It is known that bacteria grow slower when infected with the helper phage or virus. Therefore,
25 bacteria that lack the phage genome quickly overgrow the other bacteria that do contain the genome. Another effect of the lack of an ORI or a selection marker is that g3-minus phage genomes cannot be kept in dividing bacteria during the production and expansion of phage display
30 libraries. This is a very important negative feature because overgrowth of bacteria that lost the phage genome or that did never receive one, appear to have a growth advantage over bacteria that do contain the phage genome. In addition of course, such 'empty' bacteria are not
35 capable of producing any phage and as a result of this the phage display vectors including fusion protein fragments

in such helper phages lacking-bacteria are lost permanently.

As mentioned, the g3p's are thought to be essential for the assembly of stable M13-like phages and because of their crucial role in infection, g3p's should be provided otherwise when g3-minus helper phages are to be generated. There is a prejudice in the art against making phage display libraries that lack g3p's because phages lacking g3p's are not stable. Rakonjac et al. (1997) constructed a VCSM13 g3-minus helper phage in parallel to a R408 g3-minus helper phage and used helper plasmids with either the psp or the lac promoter upstream of a full length g3 sequence to substitute g3 during helper phage synthesis (Model et al. 1997). However, the art teaches that the lac promoter has the disadvantage that it cannot be shut off completely, even not in the presence of high concentrations of glucose (3-5%) in the medium (Rakonjac and Model. 1998). An additional problem that is well known in the art is that even very low levels of g3p in *E.coli* can block infection of M13-like phages. Moreover, it has been shown that co-encapsidation of plasmids together with the phage genome can occur (Russel and Model. 1989; Krebber et al. 1995; Rakonjac et al. 1997). If co-encapsidation occurs with the lac driven helper plasmid it will compete with the lac driven vectors used in the phage display resulting in the efficient production of infectious phage particles that will not contain the g3p-X fusion product. Together, the art thus teaches that the lac promoter is not the best candidate promoter in the helper plasmid system. The psp promoter has the advantage to be relatively silent in *E.coli* until infection (Rakonjac et al. 1997). Upon M13-class phage infection the psp promoter becomes activated and now the helper plasmid will produce g3 proteins. However, the disadvantage of this promoter is that the level of RNA production cannot be regulated with external factors, but has to be

regulated by either mutating (and change the activity of) the promotor, changing the ribosomal binding site (RBS) or other elements that influence the promotor activity. To figure out the ideal level of promotor activity in a specific *E.coli* strain can be time consuming and needs to be optimized for each *E.coli* strain separately. The art also teaches that the psp promotor system is not very attractive for large-scale helper phage production due to the inflexibility of *E.coli* strains, the time consuming optimization and the significant low level of helper phage production.

A significant problematic feature of all helper phage systems described is the occurrence of unwanted recombination events between the helper genome and the (helper-) plasmids. The problem that confronts investigators in the art is the fact that the g3 DNA sequences in the helper phages are homologous to the g3 sequences in the phage display vector and/or the helper phage plasmid. This results in many cases in recombination between the two DNA strains and therefore loss of functionality of the library as a whole.

It is an object of the present invention to deal with some of the problems and drawbacks known from the art as described above, concerning the generation of phage particles and helper phages, the use of helper phages in the production of phage display libraries, and the problems and drawbacks known for the identification of relevant binding molecules using such libraries.

SHORT DESCRIPTION OF THE FIGURES

- Figure 1.** Schematic representation of the g3 protein (g3p) present in the coat of M13 phages. The D3 domain of g3p is attached to the single stranded DNA inside the particle via the g8 protein (g8p), while the D1 and D2 domains interact with each other outside the particle and can be used for fusion with for example scFv.
- Figure 2.** Schematic representation of the interaction between the D2 domain of g3p with the F-pilus on the surface of *E.coli*, with a subsequent interaction of the D1 domain with other components of the bacterial surface.
- Figure 3.** Schematic representation of the interaction of the D2 domain of g3p with the F-pilus (left) and the D1 domain of g3p with the TolA receptor (see Figure 2) and the subsequent entry (right) of the phage genome into the cytoplasm of the bacterium.
- Figure 4.** Schematic representation of a wild type phage expressing five g3p's on its infectious end (left) and a recombinant phage expressing four wild type g3p's and one g3p-X fusion protein on its infectious end (right). The recombinant phage also harbors the genetic information of the fusion protein present on the surface.
- Figure 5.** Schematic representation of the pBAD/gIII-g3 helper vector harboring the full-length g3 gene under the control of the AraC/BAD promoter and further harboring an ampicillin resistance and a ColE1 origin of replication (ORI).
- Figure 6.** (A) Schematic representation of the VCSM13 helper phage genome. The 11 genes are indicated, as well as the kanamycin resistance gene (Kan R), the packaging

signal (PS) and the large and small fragments of the original intergenic region (IG). (B) Sequence of the VCSM13 genome shown in (A). The translation of the g3 gene is given in one letter code. (C) Schematic representation of the VCSM13-derived g3-minus helper phage genome deleted for the open reading frame (ORF) of the g3 gene. (D) Sequence of the part of the helper phage genome that surrounds the position of the g3 deletion depicted in (C). The HindIII site at position 3431 is underlined, followed by the 6 codons in bold face upstream of the TAA stop codon.

Figure 7. (A) Schematic representation of the VCSM13-derived D3 helper phage genome deleted for the D1 and D2 domains of the g3 gene. The D3 part of the g3 gene encodes the carboxy-terminal part of the g3 protein enabling the generation of stable, but essentially non-infectious helper phages. The D3 helper phage genome is identical to the CT helper phage genome. (B) Sequence of the D3 domain of g3 present in the nucleic acid shown in (A). The BamHI site at position 3488 as well as the GTG start and TAA stop codon are underlined.

Figure 8. Codon usage adaptation in the g3 gene and the D3 domain to prevent homologous recombination during helper phage production and during phage display library amplification. The left panel shows the obtained one-letter coded amino acids and the right panel shows the optimal codons.

30

Figure 9. Schematic representation of the pUC-g3 helper plasmid harboring the full length g3 gene under the control of the lac promoter and further harboring an ampicillin resistance gene (Amp R) and a ColE1 origin of replication (ORI).

35

Figure 10. Schematic representation of the VCSM13-derived N2CT helper phage genome deleted for the N1 (D1) domain of the g3 gene that confers infectivity to the phage.

5 **Figure 11.** Schematic representation of the VCSM13-derived p3-minus helper phage genome expressing the g3 leader followed by only seven amino acids. At the DNA level, a large part of the N2 (D2) domain and the entire CT (D3) domain are still present. The former C-terminal region of
10 gene III is present but does not encode functional protein through frame shift and therefore includes many in-frame stop codons.

SUMMARY OF THE INVENTION

The current invention provides methods and means that substantially lack the drawbacks outlined above and that
5 are characterized in providing novel phage particles such as chimaeric phages, novel helper phages, libraries comprising said chimaeric phages and methods and means to produce said chimaeric phages and said helper phages.

The invention provides a chimaeric phage having a
10 coat comprising a mixture of proteins, said mixture comprising a fusion protein wherein a proteinaceous molecule is fused to a functional form of a phage coat protein, said mixture further comprising a mutant form of said phage coat protein, said mutant form being impaired
15 in binding to a host cell receptor. The invention also provides a chimaeric phage having a coat comprising a mixture of proteins, said mixture comprising a fusion protein wherein a proteinaceous molecule is fused to a phage coat protein, or to a fragment or derivative
20 thereof, and wherein said fusion protein is functional so as to render the chimaeric phage infectious, said mixture further comprising a mutant form of said phage coat protein, said mutant form being impaired in binding to a host cell receptor.

25 In one embodiment, the invention provides a helper phage comprising nucleic acid encoding phage proteins or functional equivalents thereof that are essential for the assembly of said helper phage, said nucleic acid further encoding a mutant form of a phage coat protein, said
30 mutant form being impaired in binding to a host cell receptor, and wherein said helper phage does not comprise nucleic acid encoding a functional form of said phage coat protein.

The invention further provides methods and means for
35 producing phage particles, chimaeric phages, infectious phages and helper phage according to the invention. The

invention also provides phage collections such as phage display libraries comprising chimaeric phages and/or infectious phages according to the invention.

According to one aspect of the present invention there is provided a helper phage displaying a mixture of proteins on its surface, characterized in that said mixture comprises a) a g3 phage coat protein or part, derivative or analogue thereof for mediating infection of a host by the helper phage, and b) a mutant form of the g3 phage coat protein, wherein the D1 region, D2 region or both regions of the g3 protein are deleted.

According to a further aspect of the present invention there is provided a method of producing a helper phage displaying a mixture of proteins on its surface, said mixture comprising a g3 phage coat protein or part, derivative or analogue thereof for mediating infection of a host by the helper phage and a mutant form of the g3 phage coat protein, wherein the D1 region, D2 region or both regions of the g3 protein are deleted, the method comprising the steps of: a) providing a host cell with a first nucleic acid encoding the g3 phage coat protein or part, derivative or analogue thereof for mediating infection of a host by the helper phage; b) providing the host cell with a second nucleic acid encoding the mutant form of the g3 phage coat protein, said second nucleic acid further comprising all other proteins that are essential for the assembly of the helper phage in the host cell; and c) culturing the host cell to allow assembly of the helper phage.

According to another aspect of the present invention there is provided a method of producing a chimaeric phage particle, the method comprising the steps of: a) providing a host cell with a first nucleic acid encoding a fusion protein, wherein the fusion protein comprises a proteinaceous molecule fused to a g3 phage coat protein or part, derivative or analogue thereof for mediating infection of a host by the chimaeric phage particle; b) providing the host cell with a second nucleic acid encoding a mutant form of the g3 phage coat protein, wherein the mutant form comprises a deletion of the D1 region, D2 region or both regions of the g3 phage coat protein, said host cell further comprising a nucleic acid encoding all other proteins that are essential for the assembly of the chimaeric phage particle in the host cell; and c) culturing the host cell to allow assembly of the chimaeric phage particle, the chimaeric phage particle displaying a mixture of proteins on its surface, the mixture comprising the mutant form of the g3 phage coat protein and the fusion protein, wherein the fusion protein is attached to the chimaeric filamentous phage particle via the g3 phage coat protein or part, derivative or analogue thereof for mediating infection of a host by the chimaeric phage particle.

According to a still further aspect of the present invention there is provided a method of producing a chimaeric phage particle, the method comprising the steps of: a) providing a host cell with a first nucleic acid encoding a fusion protein, wherein the fusion protein comprises a proteinaceous molecule fused to a g3 phage

coat protein or part, derivative or analogue thereof for mediating infection of a host by the chimaeric phage particle; b) providing the host cell with a second nucleic acid encoding a mutant form of the g3 phage coat protein, wherein the mutant form comprises a deletion of the D1 region, D2 region or both regions of the g3 phage coat protein, the second nucleic acid further encoding all other proteins that are essential for the assembly of the chimaeric phage particle in the host cell; and c) culturing the host cell to allow assembly of the chimaeric phage particle, the chimaeric phage particle displaying a mixture of proteins on its surface, the mixture comprising the mutant form of the g3 phage coat protein and the fusion protein, wherein the fusion protein is attached to the chimaeric phage particle via the g3 phage coat protein or part, derivative or analogue thereof for mediating infection of a host by the chimaeric phage particle.

DETAILED DESCRIPTION OF THE INVENTION

The present invention provides a chimaeric phage having a coat comprising a mixture of proteins, said mixture comprising a fusion protein wherein a proteinaceous molecule is fused to a functional form of a phage coat protein, said mixture further comprising a mutant form of said phage coat protein, wherein said mutant form is characterized in that a phage comprising no wild type phage coat protein from which said mutant form is derived and having a coat comprising said mutant form and no copies of said functional form, is less infectious than a phage comprising no wild type phage coat protein from which said mutant form is derived and having a coat comprising said mutant form and at least one copy of said functional form. In one embodiment of the invention, said mutant form is characterized in that a phage comprising no wild type phage coat protein from which said mutant form is derived and carrying said mutant form and no copies of said fusion protein is less infectious than a phage comprising no wild type phage coat protein from which said mutant form is derived and carrying in addition to said mutant form at least one copy of said fusion protein. Preferably, said mutant form is characterized in that a phage comprising no wild type phage coat protein from which said mutant form is derived and carrying said mutant form and no copies of said fusion protein or said functional form is non-infectious. More preferably, said mutant form is further characterized in that a phage having a coat comprising said mutant form in the presence or absence of copies of said functional form, is stable. Stable as used herein means that the part of g3p that is still present in said mutant form (and that is also present in said functional form) ensures features such as DNA binding and rigidity of the phage, but does not contribute to infectiousness of said phage as do the

domains in said functional form that are not present in said mutant form. The invention also provides an infectious phage containing at least one copy of a mutant form of a phage coat protein, wherein said mutant form has
5 lost the ability to mediate infection of a natural host by said infectious phage.

In a preferred embodiment, said phage coat protein is the g3 protein (g3p) present in the coat of phages such as M13 and R408. More preferably, said mutant form comprises
10 a mutation in the D1 and/or the D2 region of g3p. In another preferred aspect of the invention said chimaeric phage and/or said infectious phage are part of a phage collection such as a phage display library. In a more preferred aspect of the invention such a phage collection
15 consists essentially of chimaeric phages or infectious phages provided by the invention. Also preferred are chimaeric phages or infectious phages according to the invention that comprise binding moieties, such as antibodies or fragments thereof as part of said fusion
20 protein.

In another embodiment the invention provides a method for producing a phage particle comprising the steps of: providing a host cell with a first nucleic acid encoding a fusion protein, said fusion protein comprising a
25 proteinaceous molecule fused to a functional form of a phage coat protein; providing said host cell with a second nucleic acid encoding a mutant form of said phage coat protein, said mutant form being characterized in that a phage comprising no wild type phage coat protein from
30 which said mutant form is derived and having a coat comprising said mutant form and no copies of said functional form is less infectious than a phage comprising no wild type phage coat protein from which said mutant form is derived and having a coat comprising at least one
35 copy of said functional form, and wherein said host cell comprises an additional nucleic acid sequence encoding at

least all other proteins or functional equivalents thereof, that are essential for the assembly of said phage particle in said host cell; and culturing said host cell to allow assembly of said phage particle. The invention
5 provides a method for producing a phage particle, said method comprising the steps of: providing a host cell with a first nucleic acid encoding a fusion protein, said fusion protein comprising a proteinaceous molecule fused to a functional form of a phage coat protein; providing
10 said host cell with a second nucleic acid encoding a mutant form of said phage coat protein, said mutant form being impaired in binding to a host cell receptor; and culturing said host cell to allow assembly of said phage particle. Preferably, these methods according to the
15 invention for producing a phage particle are applied for producing said chimaeric phage and/or said infectious phage. More preferably said method is applied for producing a phage particle such as a chimaeric phage or an infectious phage provided by the invention that comprise
20 nucleic acid encoding said mutant form under the control of a controllable promoter such as the AraC/BAD promoter, the lac promoter or the psp promoter.

In another embodiment the invention provides a helper phage comprising nucleic acid encoding phage proteins or
25 functional equivalents thereof that are essential for the assembly of said helper phage, said nucleic acid further encoding a mutant form of a phage coat protein, wherein said mutant form is characterized in that a phage comprising no wild type phage coat protein from which said
30 mutant form is derived and having a coat comprising said mutant form and no copies of a functional form of said phage coat protein, is less infectious than a phage comprising no wild type phage coat protein from which said mutant form is derived and having a coat comprising at
35 least one copy of said functional form, wherein said functional form is characterized in that it renders a

phage particle carrying said functional form in its coat infectious, and wherein said helper phage does not comprise expressible nucleic acid encoding said functional form. The invention provides a helper phage comprising
5 nucleic acid encoding phage proteins or functional equivalents thereof that are essential for the assembly of said helper phage, said nucleic acid further encoding a mutant form of a phage coat protein, said mutant form being impaired in binding to a host cell receptor, and
10 wherein said helper phage does not comprise nucleic acid encoding a functional form of said phage coat protein.

In yet another embodiment the invention provides a method for producing a helper phage comprising the steps of: providing a host cell with a first nucleic acid
15 encoding a functional form of a phage coat protein; providing said host cell with a second nucleic acid encoding a mutant form of said phage coat protein, wherein said mutant form is characterized in that a phage comprising no wild type phage coat protein from which said
20 mutant form is derived and having a coat comprising said mutant form, is less infectious than a phage comprising no wild type phage coat protein from which said mutant form is derived and having a coat comprising at least one copy of said functional form, wherein said host cell comprises
25 an additional nucleic acid sequence encoding at least all other proteins or functional equivalents thereof that are essential for the assembly of said helper phage in said host cell; and culturing said host cell to allow assembly of said helper phage. The invention also provides a method
30 for producing a helper phage comprising the steps of: providing a host cell with a first nucleic acid encoding a functional form of a phage coat protein; providing said host cell with a second nucleic acid encoding a mutant form of a phage coat protein, said mutant form being
35 impaired in binding to a host cell receptor, wherein said host cell comprises an additional nucleic acid sequence

encoding at least all other proteins or functional equivalents thereof that are essential for the assembly of said helper phage in said host cell; and culturing said host cell to allow assembly of said helper phage.

5 In another aspect the invention provides methods and means for producing a phage particle such as a chimaeric phage, an infectious phage or a helper phage according to the invention, wherein separate nucleic acids encoding either (1) a functional form of said phage coat protein
10 alone or fused to a proteinaceous molecule or (2) a mutant form of said phage coat protein, each comprise codons in the overlapping regions between the protein encoding parts, that essentially do not render a homologous recombination event between said separate nucleic acids.
15 In a preferred aspect said separate nucleic acids each comprises non-interfering origins of replication and unique selection markers.

In another embodiment the invention provides a method for the enrichment of a first binding pair member in a
20 repertoire of first binding pair members selected from the group consisting of an antibody, an antibody fragment, a single chain Fv fragment, a Fab fragment, a variable region, a CDR region, an immunoglobulin or a functional part thereof, said first binding pair member being
25 specific for a second binding pair member, said method comprising the steps of: contacting a phage collection comprising chimaeric or infectious phages according to the invention, with material comprising said second binding pair member under conditions allowing specific binding;
30 removing non-specific binders; and recovering specific binders, said specific binders comprising said first binding pair member. Said material may comprise second binding pair members such as purified proteins, recombinant proteins and/or proteins present in or on
35 cells. In a preferred embodiment, the invention provides a method for the enrichment of a first binding pair member

that furthermore comprise the steps of: recovering from a phage a DNA sequence encoding said first specific binding pair member; sub-cloning said DNA sequence in a suitable expression vector; expressing said DNA sequence in a
5 suitable host; and culturing said suitable host under conditions whereby said first specific binding pair member is produced. A suitable expression vector may be a plasmid vector comprising an active promoter that regulates the expression of said first specific binding pair member in
10 suitable hosts like eukaryotic cells such as yeast cells or mammalian cells.

In another aspect, the invention provides a nucleic acid molecule comprising a sequence encoding a mutant form of a phage coat protein, said mutant form being
15 characterized in that a phage, comprising no wild type phage coat protein from which said mutant form is derived and having a coat comprising said mutant form and no functional form of said phage coat protein, is less infectious than a phage comprising no wild type phage coat
20 protein from which said mutant form is derived and having a coat comprising said mutant form and at least one copy of said functional form of said phage coat protein, wherein said functional form is characterized in that it renders a phage carrying said functional form in its coat
25 infectious. Said nucleic acid molecule may furthermore comprise all relevant nucleic acid encoding proteins that are required for assembly of a phage in a host cell.

The invention provides a chimaeric phage having a
30 coat comprising a mixture of proteins, said mixture comprising a fusion protein wherein a proteinaceous molecule is fused to a functional form of a phage coat protein, said mixture further comprising a mutant form of said phage coat protein, wherein said mutant form is
35 characterized in that a phage comprising no wild type phage coat protein from which said mutant form is derived

and having a coat comprising a mutant form of said phage coat protein and no copy or copies of said functional form of said phage coat protein, is less infectious than a phage comprising no wild type phage coat protein from which said mutant form is derived and having a coat comprising at least one functional form of said phage coat protein. 'Functional form' as used herein refers to a phage coat protein that contributes significantly to the infectiousness of the particle to which it is attached.

10 The phage coat protein itself is not infectious, but the functional form renders the phage particle to which it is attached infectious. Besides the contribution to infectivity of the phage particle, the phage coat protein also sustains other functions such as stabilization of the phage particle. A mutant form of a phage coat protein according to the invention may render the phage particle less infectious or non-infectious but should still sustain other functions of said phage coat protein such as stabilization of the phage. A phage that comprises no wild type phage coat protein from which said mutant form is derived and comprises no functional forms of said phage coat protein but does contain only mutant forms of said phage coat protein is less infectious than a phage that comprises no wild type phage coat protein from which said mutant form is derived and comprises one or more functional forms of said phage coat protein next to the mutant forms of said phage coat protein in its coat. 'Less infectious' as used herein may also mean non-infectious. Although a chimaeric phage of the invention comprises at least one copy of the mutant form of said phage coat protein in its coat, the chimaeric phage has infectious capability because it also comprises at least one functional form of said phage coat protein in its coat. A 'functional form of a phage coat protein' as used herein also means a part, derivative and/or an analogue thereof that still harbors functionality in rendering the phage

infectious to which it is attached. 'Mutant form of a phage coat protein' as used herein also means a part, a derivative and/or an analogue of said mutant form, wherein said mutant form is characterized in that a phage,
5 comprising no wild type phage coat protein from which said mutant form is derived and having a coat comprising only mutant forms of said phage coat protein or parts, derivatives and/or analogues thereof, is less- or non-infectious as compared to a phage comprising no wild type
10 phage coat protein from which said mutant form is derived and having a coat comprising at least one functional form of said phage coat protein.

In a preferred embodiment said phage coat protein is the g3 protein (g3p) that in a wild type or functional
15 form renders a phage to which it is attached, infectious. As is outlined in Figure 2 and 3, certain parts of the g3 protein are involved in the recognition of host cell receptors. A mutant form is 'impaired' in receptor binding if an alteration in the g3 protein renders the receptor to
20 be recognized and bound by the g3 protein, or parts thereof, to a lower extend than if no alteration is present. Therefore, 'impaired' as used herein means that the coat protein, such as g3p, binds the host cell receptor less efficiently, or that g3p has completely lost
25 the ability to bind and/or recognize the host cell receptor. In a preferred embodiment the mutant form of said phage coat protein comprises an alteration in the g3 protein consisting of a mutation in either the D1 region, the D2 region or both. An 'alteration' or 'mutation' as
30 used herein means one or multiple point mutations, stretches of mutations, deletions, substitutions, replacements and/or swapping of parts. In a more preferred embodiment said alteration in the g3 protein is a deletion of substantially all of the D1 and/or the D2 region. Said
35 alteration may also mean a substitution of the deleted g3 protein part by a protein or a peptide not contributing to

the infectivity of the helper phage, the chimaeric phage, the infectious phage or the phage particle.

In a more preferred embodiment of the invention a chimaeric phage or an infectious phage according to the invention comprises a nucleic acid encoding a fusion protein, wherein a proteinaceous molecule is fused to a functional form of said phage coat protein and wherein said chimaeric phage of the invention comprises a M13, M13K07, VCSM13 or a R408 strain or a mutant, derivative or analogue strain derived from either one of these strains. A proteinaceous molecule according to the invention is fused to said functional form of said phage coat protein and comprises a protein such as a ligand-binding moiety or an immunoglobulin (such as an antibody). A proteinaceous molecule may also mean a peptide such as a random stretch of amino acids or a non-random stretch of amino acids such as an antibody fragment or derivatives thereof (Fab fragment, a single chain Fv fragment (scFv), a variable region or a CDR region). A proteinaceous molecule can also mean first specific binding pair member or can mean fusions between different kinds of (fragments of) proteins and/or fusions between (fragments of) proteins and (random and non-random) peptides such as antibody-recognized tags.

A chimaeric phage of the invention relies significantly for infection on the functional form of the phage coat protein g3p and on the presence of part(s) of said phage coat protein that contribute to the infectivity of the phage. Said mutant form of said phage coat protein is mutated in the part(s) of said phage coat protein that render the phage infectious. The mutation is exemplified in, but not limited to, deletions, residue- or fragment substitutions, swaps and/or replacements by other protein fragments rendering it less infectious. Said protein fragments may or may not be related to phage coat proteins or fragments thereof, and are essentially not capable of inducing infection of the phage particle into a host cell.

In another embodiment the invention provides a phage collection comprising a chimaeric phage or an infectious phage according to the invention. Phages of the present invention are particularly useful for the generation of phage display libraries. Therefore, in a more preferred embodiment said phage collection is a phage display library. In an even more preferred embodiment said phage collection consists essentially of chimaeric phages or of infectious phages of the invention. A proteinaceous molecule, such as (random or not random) stretches of amino acids, peptides, protein parts or even entire proteins can be fused to the phage coat proteins and can form a first specific binding pair member. This fusion is typically done at the terminal ends of the coat protein. Addition typically does not affect the function of the phage coat protein. Moreover, it also often does not interfere with the function of the added moiety. Thus it is possible to generate libraries that for instance can be used to locate and to clone specific binding molecules. Such libraries can comprise peptides or larger molecules. Preferably, said larger molecules comprise a protein such as an antibody or a functional part, derivative and/or analogue thereof, such as full length heavy and/or light chains from an immunoglobulin molecule, or fragments of immunoglobulins such as Fab fragments, single chain Fv (scFv) fragments, CDR regions, sole variable regions and/or combinations of the above.

In another aspect the invention provides a method for making a phage particle which comprises the steps of providing a host cell with a first nucleic acid encoding a fusion protein, said fusion protein comprising a proteinaceous molecule fused to a functional form of a phage coat protein or to a functional part, derivative and/or analogue of said phage coat protein, providing said host cell with a second nucleic acid encoding a mutant form of said phage coat protein, said mutant form being

characterized as described above and wherein less infectious may also mean non-infectious, and culturing said cell to allow assembly of said phage, said host cell otherwise or additionally comprising nucleic acid encoding at least all essential proteins, or functional equivalents of said essential proteins for the assembly of said phage particle. In a preferred embodiment the invention provides a method wherein the nucleic acid encoding at least all other proteins or functional equivalents thereof that are essential for the assembly of said phage particle in said host cell is comprised by a helper phage and said helper phage is used to deliver the nucleic acid to said host cell. In a more preferred embodiment the nucleic acid that is delivered by said helper phage also comprises said second nucleic acid encoding said mutant form of said phage coat protein. In an even more preferred embodiment said first and said second nucleic acid are separate nucleic acids that each may comprise separate unique selection markers to ensure that the host cell comprises at least one copy of each separate nucleic acid and that each comprise separate unique origins of replication to ensure no interferences during replication. In another aspect of the invention the number of possible homologous recombination events between overlapping stretches of nucleic acid sequences between the separate nucleic acids is reduced due to the use of different codons within each nucleic acid.

Using a method of the invention it is possible to generate phage particles that comprise at least two variants derived from the same coat protein. Such phage particles have not been used in the art. The art teaches the use of phage particles either having a coat holding deleted g3 proteins, g3 proteins linked to heterologous proteins, deleted g3 proteins linked to heterologous proteins or limited numbers of g3 proteins, but no mixtures thereof. The present invention provides novel

phage particles (chimaeric phages, infectious phages and helper phages) in which the relative number of the g3p variants (either fused to a heterologous moiety or not) present in the coat can vary. Typically, one wants to influence the relative amount of the various variants in said phage coat. To that end it is preferred that expression of said fusion protein and/or expression of said mutant form of said phage coat protein is regulatable (controllable). Preferably, this is achieved by regulating the expression of the gene encoding said phage coat protein at a transcriptional level. Thus, preferably, expression of said fusion protein and/or said mutant form of said phage coat protein is under the control of a promoter that is well controlled. Examples of such promoters are the lac promoter, the psp promoter and the AraC/BAD promoter, the latter being influenced by the concentration of glucose or arabinose in the medium. Particularly advantageous is the AraC/BAD promoter. The AraC/BAD promoter is preferred because it is a promoter that is controlled in a very tight manner. This promoter is for all practical purposes silent in the presence of glucose and only slightly leaky in the absence of glucose. This means that a low concentration or absence of glucose renders the promoter active, resulting in up-regulation of the gene of interest that is under the control of the promoter. If, for example, the deletion mutant of g3 (lacking the D1 and D2 infectious regions) is under the control of such a promoter, the relative number of deletion mutants as compared to full length (or at least functional) g3-fusion proteins would be low when the glucose concentration is relatively high. In principle, this system results in a tight regulation of the number of deletion mutants and functional coat proteins such as g3 on the coat of a phage particle, and therefore the percentage of such coat proteins and deletion mutants can be regulated. In addition, the activity of the AraC/BAD

promoter can be regulated very tightly by the addition of arabinose to the medium. The concentration arabinose used determines the level of protein expressed in *E.coli* cells. Therefore, optimal regulation of phage coat protein content is accomplished by using this AraC/BAD promoter and by altering the culturing conditions of the host cell. The use of a promoter that is dependent on arabinose such as the AraC/BAD promoter, instead of IPTG, such as the lac-operon, prevents possible problems that will occur due to co-encapsidation of the helper plasmid in viral particles during helper phage synthesis. As described above, it has been shown that co-encapsidation of plasmids together with the phage genome does occur (Russel and Model. 1989; Krebber et al. 1995; Rakonjac et al. 1997). If co-encapsidation occurs with a lac driven helper plasmid, it may compete with the lac driven vectors, generally used for the phage display, resulting in the production of infectious phage particles that will not contain the g3p-X fusion product. This problem most likely does not occur when the AraC/BAD promoter is used. However, in the studies presented by the present invention such competition problems with the lac promoter were not observed. Therefore, other regulatable promoters such as the psp and lac promoter can be used to generate phages according to the invention. It is therefore also a part of the invention to use such promoters as an alternative to the AraC/BAD promoter.

A method of the invention for the production of a phage particle is preferably used to produce a chimaeric phage according to the invention. The host can be provided with nucleic acid encoding a phage protein in any suitable way. Preferably, however, said host is provided with a helper phage according to the invention. A helper phage according to the invention comprises nucleic acid encoding other phage proteins or functional equivalents thereof that are essential for the assembly of said helper phage,

said nucleic acid further encoding a mutant form of a phage coat protein, wherein said mutant form is characterized in that a phage comprising no wild type phage coat protein from which said mutant form is derived
5 and having a coat comprising mutant forms of said phage coat protein and no functional forms of said phage coat protein, is less infectious than a phage comprising no wild type phage coat protein from which said mutant form is derived and having a coat comprising at least one
10 functional form of said phage coat protein and wherein said helper phage does not comprise nucleic acid encoding a functional form of said phage coat protein. With other phage proteins is meant other than said functional form of said phage coat protein and said mutant form of said phage
15 coat protein. Nucleic acid encoding the latter may be provided to the host cell in an alternative fashion. However, said helper phage may further comprise nucleic acid encoding said functional form of said phage coat protein or said mutant form or both. Preferably, said
20 helper phage does not comprise nucleic acid encoding said fusion protein. In this way said helper phage is uniform and may be used to produce phages that preferentially comprise nucleic acid encoding said fusion protein in the absence of nucleic acid encoding any other required helper
25 phage protein. Thus preferably said fusion protein and said mutant form of said phage coat protein are encoded by separate nucleic acids. Preferably, each of said separate nucleic acids comprises a unique selection marker. Preferably, said separate nucleic acids comprise non-
30 interfering origins of replication, wherein said origins of replication do not compete with one another resulting in bacterial cells that tolerate the separate nucleic acids for the generation of new phage particles.

The art teaches that it has been very difficult to
35 generate helper phage batches, wherein said helper phages harbor nucleic acid that encodes all essential proteins

required for assembly of a phage particle in a bacterial host cell and wherein said nucleic acid lacks a gene encoding g3p. Subsequently, the art teaches that it is difficult to produce phage libraries using such helper
5 phages. Several difficulties are known in the art that hamper a proper generation of such helper phage batches. The present invention provides methods and means and a good combination of features such as the use of specific origins of replication, selection markers and codons in
10 overlapping stretches of DNA, that enable the production of phage batches containing high titers of useful helper phages, that can subsequently be applied for the generation of chimaeric or infectious phages according to the invention. Therefore, in one embodiment the invention
15 provides methods and means for the production of helper phages that carry functional forms and/or wild type forms of a phage coat protein in their coat, but that nevertheless lack nucleic acid encoding for said functional form and/or said wild type form of said phage
20 coat protein. The invention provides a method for producing a helper phage comprising the steps of: providing a host cell with a first nucleic acid encoding a functional form of a phage coat protein, providing said host cell with a second nucleic acid encoding a mutant
25 form of said phage coat protein, wherein said mutant form is characterized in that a phage comprising no wild type phage coat protein from which said mutant form is derived and having a coat comprising mutant forms of said phage coat protein and no copies of said functional form, is
30 less infectious than a phage comprising no wild type phage coat protein from which said mutant form is derived and having a coat comprising at least one functional form of said phage coat protein, and culturing said host cell to allow assembly of said helper phage, said host cell
35 additionally comprising nucleic acid encoding at least all other proteins or functional equivalents thereof that are

essential for the assembly of said helper phage in said host cell. Preferably said method does not comprise the incorporation of a wild type form of said phage coat protein in said helper phage. Preferably said phage coat protein is the g3 protein present in the coat of most, if not all bacteriophages. Preferably, said other proteins are encoded by said second nucleic acid, wherein expression of said mutant form of said phage coat protein and/or the expression of said functional form is regulated by altering the culturing conditions of the host cell and that is preferably under the control of a controllable promoter such as the AraC/BAD, psp and lac promoter as described above.

A phage coat protein is said to be contributing significantly to the infectiousness of a phage when it allows said phage, upon incorporation in a phage, to recognize, bind and/or infect a bacterial host in a manner comparable to a wild type version of said phage coat protein. As used herein the terms less-infectious or non-infectious refers to a phage, carrying no functional or wild type forms of a phage coat protein, preferably g3p, and that exhibits a significant impaired, diminished infectious capability as compared to the wild type phage (particle) as determinable in for instance plaque assays well known to persons skilled in the art. As used herein, the terms less- and non-infectious may also refer to a decrease in host cell specificity. In general, a non-infectious phage is unable to recognize, bind and/or enter a bacterial cell as outlined for wild type phages in Figure 2 and 3, whereas a less infectious phage is capable of doing so with less efficiency. Preferably, a phage having mutant forms infects the host cell with an efficiency which is less than 50% compared to a phage carrying at least a functional form, more preferably less than 10%, still more preferably less than 1%, under conditions which are otherwise comparable.

In embodiments wherein said phage coat protein comprises a g3p, said mutant form preferably comprises at least a structural part, derivative and/or analogue of the D3 region. Said structural part, derivative and/or
5 analogue comprise at least the functionality of g3p to allow assembly of stable phage particles.

A derivative of a protein as used herein comprises the same activity as the protein the derivative is derived from. When the protein is a functional form of a phage
10 coat protein the derivative comprises the same functionality. When the protein is a mutant form of a phage coat protein the derivative is also a mutant form of said phage coat protein. When the protein is a mutant form of a phage coat protein that renders a phage carrying only
15 mutant forms of said phage coat protein less infectious, the derivative also is a phage coat protein that renders a phage carrying only derivatives and/or mutant forms of said phage coat protein less infectious. Typical derivatives are proteins comprising one or more
20 conservative amino acid substitutions. However, derivatives may also comprise insertions and/or deletions. Furthermore, derivatives may also comprise swaps of amino acids or sequences of amino acids within the same protein or between two or more related and/or unrelated proteins.
25 An analogue of a protein as used herein comprises essentially the same activity as the protein the analogue is analogous to. When the protein is a functional form of a phage coat protein the analogue comprises the same functionality.

30 Prior to the invention it has been difficult to generate sufficient numbers of helper phages that lack a functional g3 gene or a part, derivative and/or analogue thereof in their genome, but that nevertheless are infectious for one round of infection through *E.coli*
35 cells. The present invention succeeds in generating such helper phages efficiently. Such helper phages carry

functional forms and/or wild type g3 proteins (g3p's) on their surface, but nevertheless lack nucleic acid encoding said functional form and/or said wild type g3 protein. The invention further provides methods and means to generate

5 libraries comprising chimaeric phages with the help of such helper phages. Preferably, phage display libraries are generated. More preferably, libraries display a large variety of single chain Fv (scFv) fragments. Using the means and methods of the invention phage libraries can be

10 generated that contain a significant higher number of infectious phages as compared to the number of non- or less-infectious phages than were described and are present in the art. At the same time the helper phages used to produce such libraries become (through an infection round

15 in *E.coli* cells) essentially non-infectious, because the g3 gene is not present in an infectivity-contributing form in said phages. Thus after infection of a bacterial host, said phages cannot spread to other bacterial cells except of course through division of the already infected host

20 into daughter cells. Such libraries are therefore also provided by the invention. The generated libraries are particularly useful for panning experiments because the titers of phages per milliliter are significantly higher than was used in the art until the present invention.

25 Moreover, libraries of the invention display less a-specific stickiness. Thus said libraries display less false positives than libraries in the art. Moreover, after one round of panning only phages that display a functional form on their surface (preferably fused to a proteinaceous

30 molecule) can be amplified in *E.coli* cells, while phages that do not carry any functional forms of said phage coat protein but only carry mutant forms of said phage coat protein are essentially non-infectious and cannot be amplified on *E.coli* cells. Therefore, the number of

35 remaining phages that are used for a second round of panning is significantly decreased. As a result of using

the chimaeric phages of the invention present in the libraries provided by the present invention the number of panning rounds is decreased and the number of relevant binders is obtained in a much more sufficient manner as
5 was possible before the present invention.

In another aspect the invention provides nucleic acids and helper phages comprising said nucleic acids that comprise genomic DNA sequences in which at least the domains of g3p that are responsible and contributing to
10 infection are functionally removed. The invention also provides helper phage genomic DNA's in which the leader and at least the D3 domain are unaffected and fused together. Said nucleic acids are preferably based on VCSM13 and M13K07 genomic sequences. Due to a lack of a
15 functional D1 domain phage particles produced by said nucleic acids are essentially non-infectious.

'Essentially' in this context means that no spread or at least significantly less spread of said phages to other bacterial cells than the production bacterium occurs
20 through g3p provided infectious features. This absence of spread to other bacterial cells is due to the absence of a functional form of g3p. If during production a source for wild type form or functional form of g3p is provided, produced phages can infect a bacterium. However, if said
25 bacterium produces phages as a result of the infection then a resulting phage particle is not capable of infecting another bacterium unless again a source for infectious g3p is provided during production. A chimaeric phage or an infectious phage of the invention preferably
30 comprises a part of g3p that ensures the generation of a stable phage particle after one round of infection in *E.coli* cells. To this end said helper phage preferably comprises a nucleic acid encoding a mutant form of said g3p. Preferably, said mutant form comprises D3 or a
35 functional part, derivative and/or analogue thereof.

A phage display library can for instance be generated by providing a collection of bacteria with a library of nucleic acids encoding g3p fused to a range of different proteinaceous molecules and infecting said bacteria with helper phages of the invention. In a particularly preferred embodiment a library of phage display particles that is produced with these helper phages contains phages that do not carry any infectious parts of g3p on their surface and phages that carry one or two full length g3p-X fusions, next to non-infectious or less-infectious parts of g3p deletion proteins. Phages in these library mixtures that do not express g3p-X fusion proteins cannot infect bacteria anymore, since they were generated in the absence of infectious g3p parts that are not fused to X (X comprises a proteinaceous molecule or a fusion partner of interest such as immunoglobulins or fragments of immunoglobulins such as Fab fragments or scFv fragments).

In one aspect the invention provides helper phages that combine the presence of a selection marker with the presence of a bacterial origin of replication (ORI) to overcome the described problems in the production of g3-minus helper phages and subsequently for the generation of phage display libraries. The presence of such a combination ensures the production of large amounts of helper phages and/or helper genomes. g3-minus helper phages with an ORI and a resistance marker can be made from the helper phages VCSM13 and M13K07. These helper phages, unlike M13 or R408, do contain a kanamycin resistance gene from the Tn903 transposon and a P15A ORI that are both inserted in the intergenic region of the phage genome. Another aspect of the invention is the fact that because of this resistance gene and the presence of this particular ORI, these helper phages can grow easily in large quantities, while empty, or no plasmid- or no genome-containing bacteria are removed under the selection pressure, and that no interferences occur between ORI's

from the phage genome and the helper plasmid or between ORI's from the phage genome and the display library plasmids, when both nucleic acids are present in the same bacterial host cell. VCSM13 and M13K07 contain the P15A
5 ORI. To prevent the disappearance of the helper genome or the helper vector the ORI's should not cause any interferences and therefore P15A derived ORI's are not used in the vector. For the vector applied for the production of helper phages, the ColE1 ORI was chosen.
10 Besides the features and effects mentioned above, it is also important that the helper vector for the generation of helper phages does not carry an F1 ORI to prevent the incorporation of the helper vector in the phage particle instead of the viral genome.

15 The invention also provides vectors enabling a regulated expression of the mutant form and/or the functional form of g3p by the use of a regulatable promoter and that furthermore contain a resistance gene that is different from the kanamycin resistance gene
20 present in the helper genome. This complementary resistance is here provided by the beta-lactamase (ampicillin) gene since its product is relatively stable and ensures complete killing of bacteria that do not express the gene product.

25 The pBAD/gIII vector (Invitrogen) can be used as a backbone vector for the production of helper vectors of the invention. Alternatively, vectors such as pUC19 can be used. Preferably, further features of these basic helper vectors are that regions of sequence homology are
30 minimized which significantly decreases the possibility of homologous recombination.

The invention further provides the use of TOP10, LMG and/or XL1blue bacterial host cells for the production of helper phages that contain a g3-minus genome but are
35 nevertheless infectious due to the g3p present on the phages because it was delivered by the helper vector. The

genotype of the TOP10 and LMG bacteria ensures that they can transport arabinose into the cell but that they cannot metabolize it (genotype: araABCD- and araEFGH+). In addition, the TOP10 bacteria are recA and endA deficient
5 which diminishes the chance of recombination and mutation. Furthermore, the TOP10 bacteria are F-, which makes them resistant to phages that might contaminate phage batches of interest.

The present invention also provides the sequence of
10 the VCSM13 phage genome and use thereof for the purpose of phage display library construction, cloning phage and plasmid mutants useful in general molecular biology methods and means.

The present invention provides further a partially
15 deleted g3 gene that is still present in the helper phage genome to provide stable but essentially non-infectious helper phages that harbor infectious g3p's on their coat. The invention describes this partially deleted g3 gene, that is made synthetically by using synthetic primers in
20 such a way that the functionally deleted g3 gene encodes the same protein on an amino acid level as compared to the other part of the g3 gene that is present in the same bacterium, but the codons that are used do not lead to homologous recombination events. Because the leader
25 sequences in the different settings are very different there is no need to change these regions in the helper genome, phage display vector (with the scFv encoding genes) or helper vector. In principle it does not matter whether the g3 gene in the helper genome, in the phage
30 display vector or helper vector has been changed as long as the two overlapping (in amino acid content) and previously homologous g3 parts that are introduced into one *E.coli* cell do not match.

The present invention describes the use of codon
35 changes in the g3 gene for the production of helper phages that are infectious due to g3p's encoded by the helper

vectors but that lack a wild type or at least an infectious g3 gene in their genome and for the production of chimaeric phages according to the invention. The use of codon changes ensures a diminished chance for homologous recombination effects that might occur during the process of helper phage generation. The invention preferably provides the use of codon changes in the g3 gene or parts thereof for the generation of phage display libraries in which the helper phage genome, that is brought into an *E.coli* cell together with nucleic acids encoding for g3p-X fusion proteins, is not homologous to the g3 gene present in the DNA encoding for the g3p-X fusion protein. These codon changes ensure that the chance for homologous recombination events in the generation of phage display libraries is significantly decreased through which the quality of these libraries and uses thereof are significantly improved.

EXPERIMENTAL PROCEDURES

Primers

The following primers (obtained from Genset or
 5 Invitrogen) were used in the generation of the different
 vectors and helper phage genomic constructs. Most
 restriction enzymes hardly or fail to digest DNA if their
 corresponding palindrome is near the end of the DNA.
 Therefore a stretch of 8 nucleotides was added to the 5'
 10 end of the D3, g3-minus and g3 ORF primers in which this
 stretch is an A/T rich non-hybridising 8-mer.

D3 primers**D3 BamHI Forward**

15 5'-GGATCCTCTGGTTCCGGTGATTTTGATTATG-3'

D3 BamHI Backward

5'-GGATCCAGCGGAGTGAGAATAGAAAGGAAC-3'

g3-minus primers

20 **g3 minus HindIII Forward**

5'AAGCTTCTGCGTAATAAGGAGTCTTAATCATGC-3'

g3 minus HindIII Backward

5'-AAGCTTGTTGAAAATCTCCAAAAAAAAGGC-3'

25 g3 ORF primers

g3 ORF NcoI Forward

5'-CCATGGCTGAAACTGTTGAAAGTTGTTTAGC-3'

g3 ORF XbaI Backward

5'-TCTAGATTAAGACTCCTTATTACGCAGTATG-3'

30

CT and N2CT primers**SnaBIclon**

5'-TTAGGTTGGTGCCTTCGTAG-3'

WO 02/103012

PCT/NL02/00391

Bamlead

5'-GGATCCAGCGGAGTGAGAATAGAAAGG-3'

BglN2

5'-AGATCTGGTACTAAACCTCCTGAGTACGG-3'

5 **BamCT**

5'-GGATCCTCTGGTTCCGGTGATTTTGATTATG-3'

PacIclon

5'-TTGCTTCTGTAAATCGTCGC-3'

10 pUC-g3 primers**H3leadA**5'-CAAATTCTATTTCAAGGAGACAGTCATAATGAAAAATTATTATTC
GCAATTCCTTTAG-3'**H3leadB**

15 5'-GATTACGCCAAGCTTGCATGCAAATTCTATTTCAAGGAGA-3'

p3endEco

5'-GCTAACATACTGCGTAATAAGGAGTCTTAAGAATTCCAGTTCTTT-3'

PCR reactions and product isolation

20 PCR reactions with D3, g3-minus and g3 ORF primers were as a standard (except for the elongation time of the DNA synthesis cycle step) performed using the following 50 μ l hot start PCR scheme and the AmpliTaq®PCR kit from Perkin Elmer: 1 μ l 10mM dNTP (Roche Diagnostics), 4 μ l

25 25mM MgCl₂, 5 μ l 10X PCR buffer supplied with the kit, 5 μ l 2.5 μ M Forward primer, 5 μ l 2.5 μ M Backward primer, 0.3 μ l 5 units/ μ l AmpliTaq®, 10-50 ng template, sterile bidistilled water. All components were kept on ice until placing in the pre-heated PCR block. The standard program

30 was as follows. 12 cycles of 25 sec 94°C, 52°C annealing for 25 sec, 72°C polymerisation ending with one cycle of 72°C for 7 min followed by storage at 4°C. The time of polymerisation for new helper genome synthesis was 12 min, and for g3 amplification and for AraC gene and AraC/BAD

35 promoter amplification this was set at 90 sec. PCR

WO 02/103012

PCT/NL02/00391

reactions with pUC-g3 primers were performed as above, but with Pwo polymerase (Boehringer Mannheim) instead of AmpliTaq®, and with the following program: 30 cycles of 45 sec 94°C, 50°C annealing for 30 sec and 72°C polymerisation for 1 min, ending with one cycle of 68°C for 8 min followed by storage at 4°C. PCR reactions with CT and N2CT primers were performed as above, but with Taq polymerase (Gibco) instead of AmpliTaq®, and with the following program: 25 cycles of 30 sec 96°C, 53°C annealing for 30 sec and 72°C polymerisation for 2 min, ending with one cycle of 72°C for 10 min followed by storage at 4°C. An exception was the PCR reaction with N2CT primers BglN2 and PacIclon, which was performed using Taq DNA polymerase recombinant (Invitrogen) with the following program: 30 cycles of 45 sec 94°C, 55°C annealing for 30 sec and 72°C polymerisation for 2.5 min, ending with one cycle of 72°C for 10 min followed by storage at 4°C.

All PCR products were separated on 0.5%-1% TBE agarose gels containing 100 ng/ml ethidium bromide. After imaging the desired fragments were cut out using sterile disposable surgical knives and isolated with Qiagen's gel purification kit according to the guided protocol.

25 Ligation reactions

Ligation reactions were as a standard performed in the following reaction mixtures:

50 ng vector or helper genome
25 ng insert
30 4 µl 5X ligation buffer (Gibco BRL)
1 µl T4-ligase (Gibco-BRL, 200 units/µl)
sterile bidistilled water to 20 µl

However, for the construction of the pUC-g3 helper plasmid, T4 DNA ligase (Roche, 1 unit/µl) was used, while
35 T4 DNA ligase (NEB, 400 units/µl) was used in the construction of the helper phage genomes CT, N2CT and p3-

minus. The mixtures were incubated overnight at 6-16°C. Then, 30 µl sterile water, 5 µl K-acetate 3M pH 4.8 acidic acid adjusted (KAc), 1 µl glycogen 10 mg/ml and 50 µl isopropanol or 96% ethanol were added and mixed
5 thoroughly. After 15 min of precipitation the tubes were centrifuged at maximum speed at 4°C for 10 min. The pellet was once washed with 1 ml 70% ethanol and after drying dissolved in 10 µl sterile water. Half of this volume was used for electroporation together with 50 µl competent
10 cells.

Sequencing

Sequencing of the clones was performed according to the instruction guide sent along with the ABI PRISM BigDye
15 Terminator Cycle Sequencing Kit (Applied Biosystems) at 50°C annealing temperature. All clones were sequenced in order to verify the correctness of the products. The sequence of the VCSM13 helper phage genome was determined by the primer-walking method on both strands, a method
20 generally known to persons skilled in the art.

Electroporation

All bacterial strains, except those that were used for the helper phage production, were acquired from
25 manufacturers as electroporation competent cells with the highest competence available and transformed according to the manufactures protocol using 0.1 cm cuvettes (BioRad). The production of helper phages, however, is dependent on TOP10 or LMG cells (Stratagene) containing the helper
30 plasmid (pBAD/gIII-g3). These cells were made competent and stored at -80°C until use as follows: One colony of the bacteria was used to inoculate in 10 ml 2X TY with ampicillin (100 µg/ml) and for LMG also with tetracycline (10 µg/ml) and cultured by vigorously shaking at 30°C
35 overnight. Next, the cultures were spun down at 3000 rpm for 5 min. The pellet was resuspended in 500 ml fresh 2X

YT including the antibiotics and cultured until OD 0.5 in a 2 l Erlenmeyer flask on a shaking platform at 37°C. These cells were allowed to cool on ice water for 45 min and centrifuged in pre-cooled buckets and rotor at 3000 rpm at 4°C for 25 min in a Sorvall centrifuge using a GLA-3000 rotor. The supernatant was discarded and the cells were slowly and carefully resuspended in 100 ml ice-cold 10% glycerol. The centrifugation and glycerol steps were repeated twice. The final pellet was taken up very carefully in 5 ml 10% ice-cold glycerol and aliquoted in pre-cooled eppendorf tubes. Next, these tubes were immersed in a mixture of ethanol and dry ice for 5 min to ensure very quick freezing of the cells. The tubes containing the electrocompetent cells were stored at -80°C until use. A comparable procedure was used to make competent cells of XL-1 (Stratagene) containing the helper plasmid pUC-g3.

Phage production

The desired F⁺ *E.coli* strain is inoculated in 2X YT medium containing the required antibiotics and cultured at 37°C at 220 rpm until OD 0.2. The (helper-) phage is added to the culture and incubated for 30-45 min at 37°C in a non-shaking waterbath. Then, kanamycine (50 µg/ml) is added to the cells and cells are further incubated at 220 rpm at 37°C for 30-45 min. Subsequently, this solution is spun at 3500 rpm at room temperature for 15 min. The supernatant is removed carefully and the pellet is brought to the desired volume of 2X YT medium containing all required antibiotics. Cells are cultured overnight at 30°C for a phage display library and at 37°C for regular (helper-) phages on a shaking platform.

Titer determination

One colony of Xl1blue (Stratagene) is inoculated in 5 ml 2X YT containing 10 µg tetracycline per ml (YT-T) in a

50 ml tube (Falcon) and cultured at 37°C at 220 rpm overnight. 200 µl of this culture is added to 5 ml YT-T and cultured until OD 0.2. Then, the phage stock is diluted in YT and a dilution series as required is made to determine the number of plaque forming units. For each dilution step 100 µl of the O.D. 0.2 XL1Blue culture is taken and added to 100 µl of the phages. This mixture is incubated for 25 min at 37°C in a water bath (not shaking). The 200 µl of bacterial cells is pipetted onto a 2X YT-broth plate containing the required antibiotics. The suspension is spread using a sterile glass rod. After drying the plates they are inverted and transferred into a 37°C incubator. After overnight culture the number of colonies are counted. Each colony indicates the presence of 1 infective phage particle in the original phage solution. The number of infectious particles per ml of the analyzed stock is calculated. The phage particles are ELISA tested according to the protocol supplied with anti-M13 and anti-M13-HRP conjugate (Pharmacia).

20

Isolation of DNA from phages

An overnight phage culture is grown as described above. If a large scale isolation of DNA was required the BioRad Plasmid Maxi Prep kit was used according to the manufactures instructions, except for the elution step which is done with 10 mM Tris pH8.5 at 65°C for 10 min. Small or medium scale isolations were performed using Qiagen's mini-prep kit according to the instruction supplied with the kit except for the elution step. The elution step here was performed at 65°C for 10 min.

30

PEG precipitation

The medium containing bacteria and phages is collected in 450 ml buckets. The mixture is spun in a pre-cooled Sorvall centrifuge using a GSA-3000 rotor at 8000 rpm for 20 min. Then, 90 ml 20% PEG/2.5 M NaCl is pipetted

35

into clean 450 ml buckets. 360 ml of the supernatant of the centrifuged medium containing the phages is brought into the PEG containing buckets and mixed well. The mixture is set on ice water for 2 h or overnight in the fridge. The precipitate is pelleted by centrifugation in a pre-cooled Sorvall centrifuge at 8000 rpm for 20 min. The supernatant is decanted and the buckets are left to drip out for 5 min in order to remove as much Precipitation buffer as possible. Subsequently, 32 ml PBS/1% bovine serum albumin (BSA) is added to the 450 ml buckets containing the pelleted phages and buckets are rotated on a bottle roller for 15 min. The solution is transferred to a SS-34 compatible centrifuge tube and spun in a pre-cooled Sorvall centrifuge containing a SS-34 rotor (or equivalent equipment) for 25 min at 13,000 rpm. This step removes all kind of debris and small bacteria. In the meantime, the plunger is removed from a 50 ml syringe and attached to a 0.45 μ M filter (Whatmann). The centrifuged supernatant is transferred into the syringe and pushed through the filter. This step removes all small bacteria and other cells. 8 ml 20% PEG/2.5 M NaCl is added and mixed well. The tubes are set on ice for 1 h. The high-speed centrifugation step is repeated as described above. The supernatant is decanted and the tube is let to drip out on a paper towel for 5 min. The phage pellet is dissolved in 5 ml PBS/1% BSA. Then, 5 ml 100% glycerol is added to the phage solution and mixed well. Phages are stored at -20°C. Typically, the solution contains approximately 2 to 5×10^{13} infectious phage particles per ml.

WO 02/103012

PCT/NL02/00391

EXAMPLES

To illustrate the invention, the following examples are provided, not intended to limit the scope of the invention.

5

Example 1.**Cloning of the pBAD/gIII-g3 helper vector.**

The full Open Reading Frame (ORF) of the g3 gene was generated by using M13KO7 (Gibco-BRL) DNA as a template in
10 a standard PCR reaction together with g3 ORF NcoI Forward and g3 ORF XbaI Backward primers. The purified PCR product and the pBAD/gIII vector were both digested with NcoI (NEB) and XbaI (Roche Diagnostics) simultaneously in buffer H (Roche Diagnostics) for 4 h at 37°C. After
15 ligation, isolation, electroporation in TOP10 (Stratagene) and LMG (Stratagene) cells, two correct clones were selected by sequencing and grown on a large scale followed by the isolation of the DNA. The DNA was reprecipitated with 70% ethanol and in the presence of KAc and the pellet
20 was washed twice with 70% ethanol. After drying the DNA was dissolved in sterile bidistilled water and stored at -20°C until use. The resulting plasmid pBAD/gIII-g3 is depicted in Figure 5. This helper vector contains the full sized g3 gene under the control of the AraC/BAD promoter,
25 ampicillin resistance gene and a ColE1 ORI as most important features.

Example 2.**Cloning of the g3-minus helper phage genome.**

30 VCSM13 (Stratagene) is a widely used helper phage, however, its sequence is not publicly available. The full DNA sequence of the VCSM13 genome was determined and is depicted schematically in Figures 6A and in nucleotide order in Figure 6B. The use of g3 minus HindIII Forward and
35 g3 minus HindIII Backward primers and M13KO7 and VCSM13 as

templates in a standard PCR reaction resulted in the formation of a PCR product that contained HindIII sites at both ends of the DNA. After separation, gel isolation and purification, digestion with HindIII (Roche Diagnostics) and re-purification of the DNA, the product was self-ligated under standard ligation conditions and the resulting helper phage genome, named g3-minus, was electroporated into XL1Blue cells (Stratagene). The transformed cells were resuspended in 5 ml 2TY medium and cultured shaking at 37°C for 1 h. Kanamycin was added to an end-concentration of 50 µg/ml and the cells were allowed to grow at the same conditions for another 5 h. The culture was centrifuged at 3000 rpm for 15 min and the supernatant passed through a 0.22 µm filter to remove bacteria. At the same time, a culture of exponentially growing XL-1 Blue bacteria was prepared. Fractions of the filtrate (50-1000 µl), containing phage particles, were added to 5 ml of XL-1 Blue bacteria and incubated at 37°C for 30 min without shaking. The culture was centrifuged again, the supernatant discarded and the cells were plated on 2X YT-K-T plates and transferred into an incubator at 37°C for overnight growth. Eight correct clones that lack the g3 ORF (checked through the BamHI site) and include the introduced HindIII site were isolated and used for g3-less helper phage production in the presence of the pBAD/gIII-g3 helper plasmid. Only two clones that were able to form phages in the presence of the helper plasmid were kept. From these clones a large quantity of DNA was isolated and stored for further experiments. The obtained VCSM13-derived g3-minus helper phage genome, in which the entire g3 gene except for the last six codons was replaced by a HindIII site, is depicted schematically in Figure 6C, while the g3-minus sequence is depicted in Figure 6D.

Example 3.**Cloning of the D3 helper phage genome with a partially deleted g3 gene.**

The construction of helper phage genome that express
5 only the D3 part of the g3 gene was comparable to the
above described g3-minus helper phages with the exception
that the primers used were D3 BamHI Forward and D3 BamHI
Backward primers in order to generate the new genome. All
other procedures were the same as for the g3-minus
10 procedure, except for the use of BamHI instead of HindIII.
In the end, the DNA of two correct clones was kept and
stored at -20°C. The final VCSM13-derived construct of the
helper phage genome (named D3: it only expresses this part
of the g3 gene that does not contribute to the
15 infectiousness of the phage particle) is depicted in
Figure 7A, while the D3 sequence is depicted in Figure 7B.

Example 4.**Production of infectious helper phages that do not carry a
20 gene encoding the wild type g3p.**

The procedures for generating g3-less and generating
partially deleted g3 (or D3 expressing) helper phages are
identical. Frozen competent TOP10 or LMG cells that
contain the pBAD/gIII-g3 helper plasmid were
25 electroporated using 100 ng helper phage DNA. After
recovery the cells were transferred into 4X 250 ml 2X YT-
K-A medium supplemented with 0.05% arabinose (Sigma).
Phages were produced during overnight culture at 37°C and
vigorously shaking. The next day the phages were purified
30 and stored according to the standard procedures of
precipitation and storage that were described supra. The
number of infectious particles and the number of phages
were determined by titration and ELISA procedures that
were also described supra. For g3-less helper phages
35 approximately 5×10^{11} infectious particles were synthesized

while for D3 approximately 5×10^{13} infectious phages were formed using these procedures.

Example 5.

- 5 **Amplification and harvesting of phage display libraries containing infectious phages carrying g3p-scFv fusions and non-infectious helper phages, using g3 minus and partially g3 deleted helper phages.**

A frozen library was inoculated as follows. In
10 general approximately 5-10 μ l concentrated stocks were inoculated in 25 ml 2X YT containing the required antibiotics and 5% glucose and grown at 37°C with vigorously shaking. At OD 0.3-0.4 (after about 2-3 h), a 500-1000 fold excess of helper phage was added. The medium
15 containing the helper phages and bacteria was incubated in a water bath at 37°C without shaking for 25 min. Removal of dead cells and excess of phage particles was realized after a centrifugation step at 3000 rpm for 15 min. The pellet was resuspended in 250 ml 2X YT with antibiotics
20 but without glucose and grown at 30°C with good aeration overnight. The next day the formed phages were isolated and stored using the standard PEG/NaCl procedure.

Example 6.

- 25 **Codon usage in g3 and the partially deleted g3 gene (D3) to prevent homologous recombination during helper phage production and library amplification.**

In order to prevent possible recombinations between a genomic nucleic acid encoding the helper phage g3 protein
30 region (or a part thereof such as the D3 domain) and other nucleic acids (like the phage display vector and the AraC/BAD helper vector), a series of helper phages are designed that contain changed codons within the g3p region. Newly translated g3p's are identical to the wild
35 type g3 protein or protein part (D3). Due to these

changes, g3-ORF coding DNA domains cannot or barely recombine with the phage display vectors or the AraC/BAD-g3 helper vector. The codons that are used to generate non-homologous g3 genes are depicted in Figure 8 and are
 5 optimal for the *E.coli* transcription machinery.

PCR generation of helper phage genomes (VCSM13, M13K07, D3, g3-minus or AraC/BAD) with g3-leader backward and g3 end forward primers with NotI restriction sites ensure the generation of PCR products containing all helper
 10 phage components and genes, except for the g3 ORF.

New g3 regions are constructed with overlapping primers and are inserted in helper phages. The PCR-generated g3p or parts thereof are digested with NotI and ligated in the NotI digested PCR generated helper phage genome.
 15 After transformation and selection of correct helper genomes (with a new g3 gene), helper phages are grown as described.

Example 7.

20 **Selection of thyroglobulin-interacting phages using a library amplified with helper phages comprising only the D3 part of g3p in their genome.**

In order to validate the D3 helper phages in standard phage selections, a selection was performed
 25 using an antibody phage display library that was amplified using the D3 helper phages as described above. Procedures that were used were essentially as described by De Kruif et al. (1995a). Briefly, thyroglobulin was coated to a plastic tube. The tube was blocked in PBS
 30 containing 2% milk (MPBS) whereafter the antibody phage display library, also blocked in MPBS, was added to the tube. The phages were allowed to bind for 2 h, whereafter non-binding phages were removed by washing the tube in PBS containing 0.1% TweenTM-20 as described by De Kruif et
 35 al. (1995a). The binding phages were eluted in 50 mM Glycin/HCl pH 2.2 (10 min at RT) and used to infect

freshly grown XL-1 Blue bacteria. The bacteria were plated on 2TY agar plates containing the appropriate antibiotics and glucose, incubated overnight at 37°C and used to prepare an enriched phage display library; phage
5 D3 was again used as helper phage. The procedure was repeated once, whereafter individual *E.coli* colonies were used to prepare monoclonal phage antibodies. These monoclonal phage antibodies were tested in ELISA for their ability to bind specifically to the thyroglobulin
10 antigen. Results show that after two rounds of selection 25/46 colonies show positive binding to thyroglobulin. Previously we found that by using a general VCS-M13 as a helper phage, at least 3 rounds of selection were required to obtain specific binders in this selection
15 format.

Example 8.

**Selection of phages interacting with myeloma cells using a library amplified with helper phages comprising only
20 the D3 part of g3p in their genome.**

In order to validate the D3 helper phages in selections on intact cells, a selection was performed using an antibody phage display library that was amplified using the D3 helper phage. Procedures that were
25 used were essentially as described by De Kruif et al. (1995a and 1995b). Briefly, myeloma cells (AML, CD33+, CD34+) were obtained from the blood of a patient undergoing treatment at the Utrecht University Hospital (The Netherlands). 0.5 ml phage library was added to 3 ml
30 RPMI medium containing 10% FCS (RPMIS) and incubated on ice for 15 min. The myeloma cells were added and the cell suspension was rotated at 4°C for 2 h. The cells were washed 5 times in 50 ml ice-cold RPMIS whereafter the binding phages were eluted (in 50 mM Glycin/HCl pH 2.2
35 for 10 min at RT) and used to infect freshly grown XL-1 Blue bacteria. The bacteria were plated on 2TY agar

plates containing the appropriate antibiotics and glucose, incubated overnight at 37°C and used to prepare an enriched phage display library. Again, the D3 expressing helper phages were used as helper. The
5 procedure was repeated once, whereafter individual *E.coli* colonies were used to prepare monoclonal phage antibodies. These monoclonal phage antibodies were tested in FACS procedures for their ability to bind myeloma cells. Results show that 23 out of 41 clones tested bound
10 specifically to epitopes expressed on the myeloma cells. Generally, three or more rounds of selection are required to obtain similar numbers of binding phages using identical procedures, with the exception of using VCS-M13 helper phages instead of the helper phages described by
15 the invention.

Example 9.

Cloning of the pUC-g3 helper plasmid.

As an alternative to the pBAD/gIII-g3 helper plasmid
20 described above, a pUC19-based helper plasmid for the expression of wild-type g3p under the control of a lac promotor was constructed. The use of H3leadA and p3endEco primers and VCSM13 as a template in a PCR reaction detailed above resulted in the formation of a PCR product
25 containing the g3 gene sequence with 28 additional nucleotides upstream of the gene and 15 additional nucleotides including an introduced EcoRI site downstream. The use of H3leadB and p3endEco primers and this PCR product as a template in a PCR reaction detailed
30 above resulted in the formation of an extended PCR product containing the g3 gene sequence, with introduced HindIII and EcoRI sites upstream and downstream of the gene, respectively. The extended PCR product was cloned into plasmid pCR4-BluntII-TOPO using the Zero Blunt TOPO
35 PCR Cloning Kit (Invitrogen) and the sequence of the insert was checked. Plasmid pUC19 (New England Biolabs)

and the pCR4-BluntII-TOPO derivative containing the g3 gene were both digested with HindIII and EcoRI. The fragment containing the g3 gene from the pCR4-TOPO derivative and the vector fragment of pUC19 were purified
5 from gel and ligated. The resulting plasmid, named pUC-g3, was electroporated into XL-1 cells (Stratagene) and a correct clone was selected by sequencing DNA isolated from the transformed cells. Helper vector pUC-g3, which contains the g3 gene under the control of the lac
10 promotor, is depicted schematically in Figure 9.

Example 10.

Alternative cloning of helper phage genomes with a partially deleted g3 gene.

15 In the cloning procedures described in example 2 and 3, a very large region of the phage genome is amplified by PCR, with the risk of introducing artificial mutations due to the limited fidelity of DNA polymerase. An alternative strategy was applied to produce helper
20 genomes with a partially deleted g3 gene. In this strategy, a part of the g3 sequence encoding the g3 leader, and a part of the g3 sequence encoding a C-terminal portion of the g3 protein, were amplified separately. The two fragments were combined in a shuttle
25 vector and subsequently cloned into the phage genome. To distinguish the resulting constructs from the D3 helper phage described in example 3, the generally accepted alternative nomenclature for the g3 domains was used, in which D1, D2 and D3 are named N1, N2 and CT,
30 respectively. The use of SnaBIclon and Bamlead primers and VCSM13 as a template in a PCR reaction detailed above resulted in the formation of a PCR product containing the g3 leader sequence, with a native SnaBI site upstream and an introduced BamHI site downstream. The use of BamCT and
35 PacIclon primers and VCSM13 as a template in a PCR reaction detailed above resulted in the formation of a

PCR product containing the g3 CT domain, with an introduced BamHI site upstream and a native PacI site downstream. The use of BglN2 and PacIclon primers and VCSM13 as a template in a PCR reaction detailed above

5 resulted in the formation of a PCR product containing the g3 N2 and CT domains, with an introduced BglII site upstream and a native PacI site downstream. All PCR products were cloned into plasmid pCR4-TOPO using the TOPO TA Cloning Kit for Sequencing (Invitrogen) and the

10 sequences of the inserts were checked. The pCR4-TOPO derivative containing the leader sequence was digested with NotI and BamHI, and the insert was isolated from agarose gel. The pCR4-TOPO derivative containing the CT domain was also digested with NotI and BamHI, and the

15 vector fragment was isolated. The insert was ligated with the vector fragment, resulting in a pCR4-TOPO derivative containing the g3 gene in which the N1 and N2 domains are deleted. This plasmid and VCSM13 were both digested with SnaBI and PacI. After isolation the plasmid insert was

20 ligated with the VCSM13 vector fragment. The sequence of the insert in the resulting phagemid, named CT and identical to D3 depicted in Figure 7A, was checked. A comparable procedure was used to construct a phagemid lacking only the N1 domain. For this construct, the pCR4-

25 TOPO derivative containing the N2 and CT domains was digested with NotI and BglII, and the vector fragment was isolated. The insert containing the leader sequence was ligated with the vector fragment (the cohesive ends of BglII and BamHI are compatible), resulting in a pCR4-TOPO

30 derivative containing the g3 gene in which the N1 domain is deleted. This gene was sub-cloned to VCSM13 as described above and the sequence of the insert in the resulting phagemid, named N2CT, was checked. The VCSM13-derived helper phage genome N2CT is depicted

35 schematically in Figure 10.

Example 11.**Cloning of the p3-minus helper phage genome.**

In the cloning of the g3-minus helper phage genome described in example 2, almost the entire g3 sequence was
5 deleted. The g3 gene overlaps the ribosome-binding site of the g6 gene and may contain other important features as well. An alternative cloning strategy was applied, in which part of the N2 domain and the entire CT domain are retained at the DNA level, but a frame shift deletes
10 these domains at the protein level. The pCR4-TOPO derivative containing the g3 leader fused to the g3 CT domain (see above) and VCSM13 were both cut with BamHI and SnaBI. After isolation from gel, the insert fragment of the pCR4-TOPO derivative, containing the g3 leader,
15 was ligated with the vector fragment of VCSM13, containing a C-terminal portion of the N2 domain and the entire CT domain of g3. The sequence of the insert in the resulting phagemid, named p3-minus, was checked. In this construct, a frame shift occurs at the BamHI site located
20 between the g3 leader and the C-terminal portion of the N2 domain, resulting in the introduction of over 30 stop codons downstream of the BamHI site. The VCSM13-derived helper phage genome p3-minus is depicted schematically in Figure 11.

25

REFERENCES

- Balint RF and Larrick JW (1993) Antibody engineering by parsimonious mutagenesis. *Gene* 137:109
- 5 Barbas CF, Hu D, Dunlop N, Sawyer L, Cababa D, Hendry RM, Nara PL, Burton DR (1994) In vitro evolution of a neutralizing human antibody to human immunodeficiency virus type 1 to enhance affinity and broaden strain
- 10 cross-reactivity. *Proc Natl Acad Sci USA*. 91:3809
- Bass SH, Greene R and Wells JA (1990) Hormone phage: An enrichment method for variant proteins with altered binding properties. *Proteins* 8:309-314
- 15 Beekwilder J, Rakonjac J, Jongsma M, Bosch D (1999) A phagemid vector using the E.coli phage shock promoter facilitates phage display of toxic proteins. *Gene* 288:23-31
- 20 Berek C, and Milstein C (1987) Mutation drift and repertoire shift in the maturation of the immune response. *Immunol Rev* 96:23
- 25 Burton DR and Barbas CF (1994) Human antibodies from combinatorial libraries. *Adv Immunol* 57:191
- Chatellier J, Hartley O, Griffiths AD, Fersht AR, Winter G, Riechmann L (1999) Interdomain interactions within the
- 30 gene 3 protein of filamentous phage. *FEBS Letters* 463:371-374
- Crissman JW and Smith GP (1984) Gene-III protein of filamentous phage: evidence for a carboxyl-terminal
- 35 domain with a role in morphogenesis. *Virology* 132:445-455

Cwirla SE, Peters EA, Barrett RE and Dower WJ (1990)
Peptides of phage: A vast library of peptides for
identifying ligands. Proc Natl Acad Sci USA 87:6378-638

- 5 De Kruif J, Boel E and Logtenberg T (1995a) Selection and
application of human single chain Fv antibody fragments
from a semi-synthetic phage antibody display library with
designed CDR3 regions. J Mol Biol 248: 97-105
- 10 De Kruif J, Terstappen L, Boel E and Logtenberg T (1995b)
Rapid selection of cell subpopulation-specific human
monoclonal antibodies from a synthetic phage antibody
library. Proc Natl Acad Sci USA 92:3938-3942
- 15 Deng LW, Malik P, Perham RN (1999) Interaction of the
globular domains of pIII protein of filamentous
bacteriophage fd with the F-pilus of Escherichia coli.
Virology 253:271-277
- 20 Devlin JJ, Panganiban LC and Devlin PE (1990) Random
peptide libraries: A source of specific protein binding
molecules. Science 249:404-406
- 25 Dueñas M, Borrebaeck CA (1995) Novel helper phage design:
intergenic region affects the assembly of bacteriophage
and the size of antibody libraries. FEMS microbiology
letters 125:317-322
- 30 Felici F, Luzzago A and Cortese R (1993) Mimicking of
discontinuous epitopes by phage displayed proteins. II.
Selection of clones recognized by a protective monoclonal
antibody against the *Bordetella pertussis* toxin from
phage peptide libraries. Gene 128:21-27

- Hawkins RE, Russel SJ, Winter G (1992) Selection of phage antibodies by binding affinity: mimicking affinity maturation. *J Mol Biol* 226:889
- 5 Holliger P, Riechmann L (1997) A conserved infection pathway for filamentous bacteriophage is suggested by the structure of the membrane penetration domain of the minor coat protein g3p from phage fd. *Structure* 5:265-275
- 10 Hoogenboom HR (1994) Designing and optimizing library selection strategies for generating high-affinity antibodies. *Trends in Biotechnol.* 15:62
- Krebber C, Spada S, Desplancq D, Pluckthun A (1995) Co-
15 selection of cognate antibody-antigen pairs by selectively-infective phage. *FEBS Lett.* 377:227-31
- Krebber C, Spada S, Desplancq D, Krebber A, Ge L, Plückthun (1997) Selectively-infective Phage (SIP): A
20 mechanistic dissection of a novel in vivo selection for protein-ligand interactions. *J Mol Biol* 268:607-618
- Kristensen P, Winter G (1998) Proteolytic selection for protein folding using filamentous bacteriophage. *Folding*
25 *and Design* 3:321-328
- Low NM, Holliger PH, Winter G (1996) Mimicking somatic hypermutation: affinity maturation of antibodies displayed on bacteriophage using a bacterial mutator
30 strain. *J Mol Biol* 260:359
- Lubkowski J, Hennecke F, Pluckthun A, Wlodawer A (1998) The structural basis of phage display elucidated by the crystal structure of the N-terminal domains of g3p. *Nat*
35 *Struct Biol* 5:140-147

- Lubkowski J, Hennecke F, Pluckthun A, Wlodawer A (1999) Filamentous phage infection: crystal structure of g3p in complex with its coreceptor, the C-terminal domain of TolA. *Structure* 7:711-722
- 5 Luzzago A, Felici F, Tramontano A, Pessi A and Cortese R (1993) Mimicking of discontinuous epitopes by phage displayed proteins. I. Epitope mapping of human H ferritin using a phage display library of constrained peptides.
- 10 Gene 128:51-57
- Lopez J and Webster RE (1983) Morphogenesis of filamentous bacteriophage f1: orientation of extrusion and production of polyphage. *Virology* 127:177-193
- 15 Model P, Jovanovic G, Dworkin J. (1997) The Escherichia coli phage shock protein (psp) operon. *Mol Microbiol* 24:255-261
- 20 Nelson FK, Friedman SM and Smith GP (1981) Filamentous phage DNA cloning vectors: A noninfective mutant with a nonpolar deletion in gene III. *Virology* 108:338-350
- 25 Nilsson N, Malmborg A-C, Borrebaeck CA (2000) The phage infection process: a functional role for the distal linker region of bacteriophage protein 3. *J Virol* 74:4229-4235
- 30 Pratt D, Tzagoloff H and Beaudoin J (1969) Conditional lethal mutants of the small filamentous coliphage. II. Two genes for coat proteins. *Virology* 39:42-53
- Rakonjac J, Jovanovic G, Model P. (1997) Filamentous phage infection-mediated gene expression: construction and propagation of the gIII deletion mutant helper phage
- 35 R408D3. *Gene* 198:99-103

Rakonjac J, Model P. (1998) Roles of pIII in filamentous phage assembly. J Mol Biol 282:25-41

5 Riechmann L, Holliger P (1997) The C-terminal domain of TolA is the coreceptor for filamentous phage infection of E. coli. Cell 90:351-360

10 Rondot S, Koch J, Breitling F, Dübel S (2001) A helper phage to improve single-chain antibody presentation in phage display. Nature Biotech 19: 75-78

15 Russel M, Model P (1989) Genetic analysis of the filamentous bacteriophage packaging signal and of the proteins that interact with it. J Virol 63:3284-3295

Smith GP (1985) Filamentous fusion phage: Novel expression vectors that display cloned antigens on the surface of the virion. Science 228:1315-1317

20 Spada S, Krebber C and Pluckthun A (1997) Selectively infective phages (SIP). Biol Chem 378:445-456

Vaughan TJ, Osbourn JK, Tempest PR (1998) Human antibodies by design. Nat Biotechnol 16:535

25 Winter G, Milstein C (1991) Man-made antibodies. Nature 349:293

30 Yang W-P, Green K, Pinz-Sweeney S, Briones AT, Burton DR, Barbas CF (1995) CDR walking mutagenesis for the affinity maturation of a potent human ant-HIV-1 antibody into the picomolar range. J Mol Biol 254:392

SEQUENCE LISTING

<110> Crucell Holland B.V.

<120> Chimaeric phages

<130> PAT 55798W-1

<140> 2,449,071

<141> 2002-06-14

<150> EP 01202304.0

<151> 2001-06-15

<150> US 09/882,621

<151> 2001-06-15

<160> 18

<170> PatentIn version 3.1

<210> 1

<211> 31

<212> DNA

<213> Artificial sequence

<220>

<223> /note= "D3 BamHI Forward primer"

<400> 1

ggatcctctg gttccggtga ttttgattat g

31

<210> 2

<211> 30

<212> DNA

<213> Artificial sequence

<220>

<223> /note= "D3 BamHI Backward primer"

<400> 2

ggatccagcg gagtgagaat agaaaggaac

30

<210> 3

<211> 33

<212> DNA

<213> Artificial sequence

<220>

<223> /note= "g3 minus HindIII Forward primer"

<400> 3

aagcttctgc gtaataagga gtcttaatca tgc 33

<210> 4
 <211> 31
 <212> DNA
 <213> Artificial sequence

<220>
 <223> /note= "g3 minus HindIII Backward primer"

<400> 4
 aagcttggtg aaaatctcca aaaaaaaagg c 31

<210> 5
 <211> 31
 <212> DNA
 <213> Artificial sequence

<220>
 <223> /note= "g3 ORF NcoI Forward primer"

<400> 5
 ccattggctga aactgttgaa agttgttttag c 31

<210> 6
 <211> 31
 <212> DNA
 <213> Artificial sequence

<220>
 <223> /note= "g3 ORF XbaI Backward primer"

<400> 6
 tctagattaa gactccttat tacgcagtat g 31

<210> 7
 <211> 20
 <212> DNA
 <213> Artificial sequence

<220>
 <223> /note= "SnaBIclon primer"

<400> 7
 ttaggttggt gccttcgtag 20

<210> 8
 <211> 27
 <212> DNA
 <213> Artificial sequence

<220>
 <223> /note= "Bamlead primer"

 <400> 8
 ggatccagcg gagtgagaat agaaagg 27

<210> 9
 <211> 29
 <212> DNA
 <213> Artificial sequence

<220>
 <223> /note= "BglN2 primer"

 <400> 9
 agatctggta ctaaacctcc tgagtacgg 29

<210> 10
 <211> 31
 <212> DNA
 <213> Artificial sequence

<220>
 <223> /note= "BamCT primer"

 <400> 10
 ggatcctctg gttccggtga ttttgattat g 31

<210> 11
 <211> 20
 <212> DNA
 <213> Artificial sequence

<220>
 <223> /note= "PacIclon primer"

 <400> 11
 ttgcttctgt aaatcgtcgc 20

<210> 12
 <211> 59
 <212> DNA
 <213> Artificial sequence

<220>
 <223> /note= "H3leadA primer"

 <400> 12
 caaattctat ttcaaggaga cagtcataat gaaaaaatta ttattcgcaa ttccttttag 59

<210> 13
 <211> 40
 <212> DNA
 <213> Artificial sequence

<220>
 <223> /note= "H3leadB primer"

<400> 13
 gattacgccca agcttgcattg caaattctat ttcaaggaga 40

<210> 14
 <211> 45
 <212> DNA
 <213> Artificial sequence

<220>
 <223> /note= "p3endEco primer"

<400> 14
 gctaacatac tgcgtaataa ggagtcttaa gaattccagt tcttt 45

<210> 15
 <211> 8669
 <212> DNA
 <213> Helper phage VCSM13

<220>
 <221> CDS
 <222> (3430)..(4704)
 <223>

<400> 15
 tgcttgactc cccgtcgtgt agataactac gatacgggag ggcttaccat ctggccccag 60
 tgctgcaatg ataccgagag acccacgctc accggctcca gatttatcag caataaacca 120
 gccagccgat tcgagctcgc ccggggatcg accagttggt gatattgaac ttttgctttg 180
 ccacggaacg gtctgcgttg tcgggaagat gcgtgatctg atccttcaac tcagcaaaag 240
 ttcgatttat tcaacaaagc cgccgtcccg tcaagtcagc gtaatgctct gccagtgtta 300
 caaccaatta accaattctg attagaaaaa ctcatcgagc atcaaatgaa actgcaattt 360
 attcatatca ggattatcaa taccatattt ttgaaaaagc cgtttctgta atgaaggaga 420
 aaactcaccg aggcagttcc ataggatggc aagatcctgg tatcggctctg cgattccgac 480
 tcgtccaaca tcaatacaac ctattaattt cccctcgtca aaaataaggt tatcaagtga 540
 gaaatcacca tgagtgcga ctgaatccgg tgagaatggc aaaagcttat gcatttcttt 600

ccagacttgt tcaacaggcc agccattacg ctcgtcatca aaatcactcg catcaaccaa	660
accgttattc attcgtgatt gcgcctgagc gagacgaaat acgcgatcgc tgttaaaagg	720
acaattacaa acaggaatcg aatgcaaccg gcgcaggaac actgccagcg catcaacaat	780
attttcacct gaatcaggat attcttctaa tacctggaat gctgttttcc cggggatcgc	840
agtgggtgagt aaccatgcat catcaggagt acggataaaa tgcttgatgg tcggaagagg	900
cataaattcc gtcagccagt ttagtctgac catctcatct gtaacatcat tggcaacgct	960
acctttgcca tgtttcagaa acaactctgg cgcategggc ttcccataca atcgatagat	1020
tgtegcacct gattgcccga cattatcgcg agccatttta taccatata aatcagcatc	1080
catgttgga ttaaatcgcg gcctcgagca agacgtttcc cgttgaatat ggctcataac	1140
cccccttgta ttactgttta tgtaagcaga cagttttatt gttcatgatg atatattttt	1200
atcttgtgca atgtaacatc agagattttg aaacacaacg tggctttccc ccccccccc	1260
ctgcaggctc cgggctattc ttttgattta taagggattt tgccgatttc ggcctattgg	1320
ttaaaaaatg agctgattta acaaaaattt aacgcgaatt ttaacaaaat attaacgttt	1380
acaatttaaa tatttgctta tacaatcttc ctgtttttgg ggcttttctt attatcaacc	1440
ggggtacata tgattgacat gctagtttta cgattaccgt tcatcgattc tcttgtttgc	1500
tccagactct caggcaatga cctgatagcc tttgtagacc tctcaaaaat agctaccctc	1560
tccggcatga atttatcagc tagaacggtt gaatatcatg ttgatgggtga tttgactgtc	1620
tccggccttt ctcacccttt tgaatcttta cctacacatt actcaggcat tgcatttaaa	1680
atatatgagg gttctaaaaa tttttatcct tgcgttgaaa taaaggcttc tcccgcaaaa	1740
gtattacagg gtcataatgt ttttggtaca accgatttag ctttatgctc tgaggcttta	1800
ttgcttaatt ttgctaattc tttgccttgc ctgtatgatt tattggatgt taacgctact	1860
actattagta gaattgatgc caccttttca gctcgcgccc caaatgaaaa tatagctaaa	1920
caggttattg accatttgcg aaatgtatct aatgggtcaaa ctaaactctac tcgttcgcag	1980
aattgggaat caactgttac atggaatgaa acttccagac accgtacttt agttgcatat	2040
ttaaaacatg ttgagctaca gcaccagatt cagcaattaa gctctaagcc atccgcaaaa	2100
atgacctctt atcaaaagga gcaattaaag gtactctcta atcctgacct gttggagttt	2160
gcttccggtc tggttcgctt tgaagctcga attaaaacgc gatatttgaa gtctttcggg	2220
cttctctta atctttttga tgcaatccgc tttgcttctg actataatag tcagggtaaa	2280

```

gacctgattt ttgatttatg gtcattctcg tttctgaac tgtttaaagc atttgagggg 2340
gattcaatga atatttatga cgattccgca gtattggacg ctatccagtc taaacatttt 2400
actattaccc cctctggcaa aacttctttt gcaaaagcct ctcgctattt tggtttttat 2460
cgtcgtctgg taaacgaggg ttatgatagt gttgctctta ctatgcctcg taattccttt 2520
tggcgttatg tatctgcatt agttgaatgt ggtattccta aatctcaact gatgaatctt 2580
tctacctgta ataatgttgt tccgtaggtt cgttttatta acgtagattt ttcttcccaa 2640
cgtcctgact ggtataatga gccagttctt aaaatcgcat aaggtaattc acaatgatta 2700
aagttgaaat taaaccatct caagcccaat ttactactcg ttctgggtgt tctcgtcagg 2760
gcaagcctta ttcactgaat gagcagcttt gttacgttga tttgggtaat gaatatccgg 2820
ttcttgtcaa gattactctt gatgaaggtc agccagccta tgcgcctggg ctgtacaccg 2880
ttcatctgtc ctctttcaaa gttggtcagt tcggttccct tatgattgac cgtctgcgcc 2940
tcgttccggc taagtaacat ggagcaggtc gcggatttcg acacaattta tcaggcgatg 3000
atacaaactc ccgttgtact ttgtttcgcg cttggtataa tcgctggggg tcaaagatga 3060
gtgttttagt gtattctttc gcctctttcg ttttaggttg gtgccttcgt agtggcatta 3120
cgtattttac ccgtttaatg gaaacttcct catgaaaaag tcttttagtcc tcaaagcctc 3180
tgtagccgtt gctaccctcg ttccgatgct gtctttcgtc gctgaggggtg acgatcccgc 3240
aaaagcggcc ttttaactccc tgcaagcctc agcgaccgaa tatatcggtt atgcgtgggc 3300
gatggttggt gtcattgtcg gcgcaactat cggtatcaag ctgtttaaga aattcacctc 3360
gaaagcaagc tgataaaccg atacaattaa aggctccttt tggagccttt ttttttggag 3420
attttcaac gtg aaa aaa tta tta ttc gca att cct tta gtt gtt cct ttc 3471
      Val Lys Lys Leu Leu Phe Ala Ile Pro Leu Val Val Pro Phe
        1             5             10

tat tct cac tcc gct gaa act gtt gaa agt tgt tta gca aaa ccc cat 3519
Tyr Ser His Ser Ala Glu Thr Val Glu Ser Cys Leu Ala Lys Pro His
15             20             25             30

aca gaa aat tca ttt act aac gtc tgg aaa gac gac aaa act tta gat 3567
Thr Glu Asn Ser Phe Thr Asn Val Trp Lys Asp Asp Lys Thr Leu Asp
              35             40             45

cgt tac gct aac tat gag ggc tgt ctg tgg aat gct aca ggc gtt gta 3615
Arg Tyr Ala Asn Tyr Glu Gly Cys Leu Trp Asn Ala Thr Gly Val Val
              50             55             60

gtt tgt act ggt gac gaa act cag tgt tac ggt aca tgg gtt cct att 3663

```

Val	Cys	Thr	Gly	Asp	Glu	Thr	Gln	Cys	Tyr	Gly	Thr	Trp	Val	Pro	Ile		
		65					70					75					
ggg	ctt	gct	atc	cct	gaa	aat	gag	ggc	ggc	ggc	tct	gag	ggc	ggc	ggc		3711
Gly	Leu	Ala	Ile	Pro	Glu	Asn	Glu	Gly	Gly	Gly	Ser	Glu	Gly	Gly	Gly		
	80					85					90						
tct	gag	ggc	ggc	ggc	tct	gag	ggc	ggc	ggc	act	aaa	cct	cct	gag	tac		3759
Ser	Glu	Gly	Gly	Gly	Ser	Glu	Gly	Gly	Gly	Thr	Lys	Pro	Pro	Glu	Tyr		
95					100					105					110		
ggc	gat	aca	cct	att	ccg	ggc	tat	act	tat	atc	aac	cct	ctc	gac	ggc		3807
Gly	Asp	Thr	Pro	Ile	Pro	Gly	Tyr	Thr	Tyr	Ile	Asn	Pro	Leu	Asp	Gly		
				115						120					125		
act	tat	ccg	cct	ggc	act	gag	caa	aac	ccc	gct	aat	cct	aat	cct	tct		3855
Thr	Tyr	Pro	Pro	Gly	Thr	Glu	Gln	Asn	Pro	Ala	Asn	Pro	Asn	Pro	Ser		
			130					135					140				
ctt	gag	gag	tct	cag	cct	ctt	aat	act	ttc	atg	ttt	cag	aat	aat	agg		3903
Leu	Glu	Glu	Ser	Gln	Pro	Leu	Asn	Thr	Phe	Met	Phe	Gln	Asn	Asn	Arg		
		145					150					155					
ttc	cga	aat	agg	cag	ggc	gca	tta	act	gtt	tat	acg	ggc	act	gtt	act		3951
Phe	Arg	Asn	Arg	Gln	Gly	Ala	Leu	Thr	Val	Tyr	Thr	Gly	Thr	Val	Thr		
	160					165					170						
caa	ggc	act	gac	ccc	gtt	aaa	act	tat	tac	cag	tac	act	cct	gta	tca		3999
Gln	Gly	Thr	Asp	Pro	Val	Lys	Thr	Tyr	Tyr	Gln	Tyr	Thr	Pro	Val	Ser		
175					180					185					190		
tca	aaa	gcc	atg	tat	gac	gct	tac	tgg	aac	ggc	aaa	ttc	aga	gac	tgc		4047
Ser	Lys	Ala	Met	Tyr	Asp	Ala	Tyr	Trp	Asn	Gly	Lys	Phe	Arg	Asp	Cys		
				195				200						205			
gct	ttc	cat	tct	ggc	ttt	aat	gag	gat	cca	ttc	gtt	tgt	gaa	tat	caa		4095
Ala	Phe	His	Ser	Gly	Phe	Asn	Glu	Asp	Pro	Phe	Val	Cys	Glu	Tyr	Gln		
			210					215					220				
ggc	caa	tcg	tct	gac	ctg	cct	caa	cct	cct	gtc	aat	gct	ggc	ggc	ggc		4143
Gly	Gln	Ser	Ser	Asp	Leu	Pro	Gln	Pro	Pro	Val	Asn	Ala	Gly	Gly	Gly		
		225					230					235					
tct	ggc	ggc	ggc	tct	ggc	ggc	ggc	tct	gag	ggc	ggc	ggc	tct	gag	ggc		4191
Ser	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Ser	Glu	Gly	Gly	Gly	Ser	Glu	Gly		
	240					245					250						
ggc	ggc	tct	gag	ggc	ggc	ggc	tct	gag	gga	ggc	ggc	tcc	ggc	ggc	ggc		4239
Gly	Gly	Ser	Glu	Gly	Gly	Gly	Ser	Glu	Gly	Gly	Gly	Ser	Gly	Gly	Gly		
255					260					265					270		
tct	ggc	tcc	ggc	gat	ttt	gat	tat	gaa	aag	atg	gca	aac	gct	aat	aag		4287
Ser	Gly	Ser	Gly	Asp	Phe	Asp	Tyr	Glu	Lys	Met	Ala	Asn	Ala	Asn	Lys		
				275					280					285			

ggg gct atg acc gaa aat gcc gat gaa aac gcg cta cag tct gac gct	4335
Gly Ala Met Thr Glu Asn Ala Asp Glu Asn Ala Leu Gln Ser Asp Ala	
290 295 300	
aaa ggc aaa ctt gat tct gtc gct act gat tac ggt gct gct atc gat	4383
Lys Gly Lys Leu Asp Ser Val Ala Thr Asp Tyr Gly Ala Ala Ile Asp	
305 310 315	
ggt ttc att ggt gac gtt tcc ggc ctt gct aat ggt aat ggt gct act	4431
Gly Phe Ile Gly Asp Val Ser Gly Leu Ala Asn Gly Asn Gly Ala Thr	
320 325 330	
ggt gat ttt gct ggc tct aat tcc caa atg gct caa gtc ggt gac ggt	4479
Gly Asp Phe Ala Gly Ser Asn Ser Gln Met Ala Gln Val Gly Asp Gly	
335 340 345 350	
gat aat tca cct tta atg aat aat ttc cgt caa tat tta cct tcc ctc	4527
Asp Asn Ser Pro Leu Met Asn Asn Phe Arg Gln Tyr Leu Pro Ser Leu	
355 360 365	
cct caa tcg gtt gaa tgt cgc cct ttt gtc ttt ggc gct ggt aaa cca	4575
Pro Gln Ser Val Glu Cys Arg Pro Phe Val Phe Gly Ala Gly Lys Pro	
370 375 380	
tat gaa ttt tct att gat tgt gac aaa ata aac tta ttc cgt ggt gtc	4623
Tyr Glu Phe Ser Ile Asp Cys Asp Lys Ile Asn Leu Phe Arg Gly Val	
385 390 395	
ttt gcg ttt ctt tta tat gtt gcc acc ttt atg tat gta ttt tct acg	4671
Phe Ala Phe Leu Leu Tyr Val Ala Thr Phe Met Tyr Val Phe Ser Thr	
400 405 410	
ttt gct aac ata ctg cgt aat aag gag tct taa tcatgccagt tcttttgggt	4724
Phe Ala Asn Ile Leu Arg Asn Lys Glu Ser	
415 420	
attccggttat tattgcgttt cctcggtttc cttctggtaa ctttgttcgg ctatctgctt	4784
acttttctta aaaagggtt cggttaagata gctattgcta tttcattggt tcttgctctt	4844
attattgggc ttaactcaat tcttggtgggt tatctctctg atattagcgc tcaattaccc	4904
tctgactttg ttcagggtgt tcagttaatt ctcccgtcta atgcgcttcc ctgtttttat	4964
gttattctct ctgtaaaggc tgctattttc atttttgacg ttaaacaaaa aatcgtttct	5024
tatttggtt gggataaata atatggctgt ttattttgta actggcaaata taggctctgg	5084
aaagacgctc gtttagcgttg gtaagattca ggataaaatt gtagctgggt gcaaaatagc	5144
aactaatctt gatttaaggc ttcaaaacct cccgcaagtc gggagggttcg ctaaaacgcc	5204
tcgcgttctt agaataccgg ataagccttc tatactctgat ttgcttgcta ttgggcgcgg	5264
taatgattcc tacgatgaaa ataaaaacgg cttgcttggt ctcgatgagt gcggtacttg	5324

gtttaataacc cgttcttgga atgataagga aagacagccg attattgatt ggtttctaca 5384
 tgctcgtaaa ttaggatggg atattatattt tcttggtcag gacttatcta ttgttgataa 5444
 acaggcgcggt tctgcattag ctgaacatgt tgtttattgt cgtcgtctgg acagaattac 5504
 ttacacctttt gtcggtactt tatattctct tattactggc tcgaaaatgc ctctgcctaa 5564
 attacatggtt ggcgttggtt aatatggcga ttctcaatta agccctactg ttgagcggtg 5624
 gctttatact ggtaagaatt tgtataacgc atatgatact aaacaggcct tttctagtaa 5684
 ttatgattcc ggtgtttatt cttatttaac gccttattta tcacacgggc ggtatttcaa 5744
 accattaaat ttaggtcaga agatgaaatt aactaaaata tatttgaaaa agttttctcg 5804
 cgttctttgt cttgcgattg gatttgcac agcatttaca tatagttata taaccaacc 5864
 taagccggag gttaaaaagg tagtctctca gacctatgat tttgataaat tcactattga 5924
 ctcttctcag cgtcttaatc taagctatcg ctatgttttc aaggattcta agggaaaatt 5984
 aattaatagc gacgatttac agaagcaagg ttattcactc acatatattg atttatgtac 6044
 tgtttccatt aaaaaaggta attcaaataa aattgttaaa tgtaattaat tttgttttct 6104
 tgatgtttgt ttcacatctt tcttttgctc aggttaattga aatgaataat tcgcctctgc 6164
 gcgattttgt aacttggtat tcaaagcaat caggcgaatc cgttattggt tctcccgatg 6224
 taaaagggtac tgttactgta tattcatctg acgttaaacc tgaaaatcta cgcaatttct 6284
 ttatttctgt ttacgtgca aataattttg atatggtagg ttctaaccct tccattattc 6344
 agaagtataa tccaaacaat caggattata ttgatgaatt gccatcatct gataatcagg 6404
 aatatgatga taattccgct ccttctgggtg gtttctttgt tccgcaaaat gataatgtta 6464
 ctcaaacttt taaaattaat aacgttcggg caaaggattt aatacgagtt gtcgaattgt 6524
 ttgtaaagtc taatacttct aaatcctcaa atgtattatc tattgacggc tctaattctat 6584
 tagttgtag tgctcctaaa gatatttttag ataaccttc tcaattcctt tcaactgttg 6644
 atttgccaac tgaccagata ttgattgagg gtttgatatt tgagggtcag caagggtgatg 6704
 ctttagatth ttcatthgct gctggctctc agcgtggcac tgttgcaggc ggtgttaata 6764
 ctgaccgcct cacctctggt ttatcttctg ctggtggttc gttcggtatt tttaatggcg 6824
 atgttttagg gctatcagtt cgcgcattaa agactaatag ccattcaaaa atattgtctg 6884
 tgccacgtat tcttacgctt tcagggtcaga agggttctat ctctgttggc cagaatgtcc 6944
 cttttattac tggtcgtgtg actggtgaat ctgccaatgt aaataatcca tttcagacga 7004

ttgagcgtca	aaatgtaggt	atitccatga	gcgtttttcc	tggtgcaatg	gctggcggta	7064
atattgttct	ggatattacc	agcaaggccg	atagtttgag	ttcttctact	caggcaagtg	7124
atgttattac	taatcaaaga	agtattgcta	caacgggttaa	tttgcgatgat	ggacagactc	7184
ttttactcgg	tggcctcact	gattataaaa	acactttctca	ggattctggc	gtaccgttcc	7244
tgtctaaaat	ccctttaatc	ggcctcctgt	ttagctcccc	ctctgattct	aacgaggaaa	7304
gcacgttata	cgtgctcgtc	aaagcaacca	tagtacgcgc	cctgtagcgg	cgcattaagc	7364
gcggcgggtg	tggtgggttac	gcgcagcgtg	accgctacac	ttgccagcgc	cctagcgcgc	7424
gctcctttcg	ctttcttccc	ttcctttctc	gccacgttcg	ccggctttcc	ccgtcaagct	7484
ctaaatcggg	ggctcccttt	agggttccga	tttagtgctt	tacggcacct	cgaccccaaa	7544
aaacttgatt	tgggtgatgg	ttcacgtagt	gggccatcgc	cctgatagac	ggtttttcgc	7604
cctttgacgt	tggagtccac	gttctttaat	agtggactct	tgttccaaac	tggaacaaca	7664
ctcaacccta	tctcgggacg	gatcgcttca	tgtggcagga	gaaaaaaggc	tgcaccggtg	7724
cgtcagcaga	atatgtgata	caggatatat	tccgcttcc	cgtcactga	ctcgctacgc	7784
tgggtcgttc	gactgcggcg	agcggaaatg	gcttacgaac	ggggcggaga	tttccctggaa	7844
gatgccagga	agatacttaa	cagggaagtg	agagggccgc	ggcaaagccg	tttttccata	7904
ggctccgccc	ccctgacaag	catcacgaaa	tctgacgctc	aatcagtggt	tggcgaaacc	7964
cgacaggact	ataaagatac	caggcgtttc	cccctggcgg	ctccctcgtg	cgtctcctg	8024
ttcctgcctt	tcggtttacc	ggtgtcattc	cgtgttatg	gcgcggtttg	tctcattcca	8084
cgcctgacac	tcagttccgg	gtaggcagtt	cgtccaagc	tggactgtat	gcacgaaccc	8144
cccgttcagt	ccgaccgctg	cgccttatcc	ggtaactatc	gtcttgagtc	caaccgggaa	8204
agacatgcaa	aagcaccact	ggcagcagcc	actggtaatt	gatttagagg	agttagtctt	8264
gaagtcatgc	gccgggttaag	gctaaactga	aaggacaagt	tttggtgact	gcgctcctcc	8324
aagccagtta	cctcgggttca	aagagttggg	agctcagaga	accttcgaaa	aaccgccctg	8384
caaggcgggt	ttttcgtttt	cagagcaaga	gattacgcgc	agaccaaacc	gatctcaaga	8444
agatcatctt	attaaggggt	ctgacgctca	gtggaacgaa	aactcacgtt	aagggatttt	8504
ggtcatgaga	ttatcaaaaa	ggatcttcac	ctagatcctt	ttaaattaaa	aatgaagttt	8564
taaatcaatc	taaagtatat	atgagtaaac	ttgggtctgac	agttaccaat	gcttaatcag	8624
tgaggcacct	atctcagcga	tctgtctatt	tcgttcatcc	atagt		8669

<400> 16

Gly Gly Ser Gly Gly Gly Ser Glu Gly Gly Gly Ser Glu Gly Gly Gly
245 250 255

Ser Glu Gly Gly Gly Ser Glu Gly Gly Gly Ser Gly Gly Gly Ser Gly
 260 265 270
 Ser Gly Asp Phe Asp Tyr Glu Lys Met Ala Asn Ala Asn Lys Gly Ala
 275 280 285
 Met Thr Glu Asn Ala Asp Glu Asn Ala Leu Gln Ser Asp Ala Lys Gly
 290 295 300
 Lys Leu Asp Ser Val Ala Thr Asp Tyr Gly Ala Ala Ile Asp Gly Phe
 305 310 315 320
 Ile Gly Asp Val Ser Gly Leu Ala Asn Gly Asn Gly Ala Thr Gly Asp
 325 330 335
 Phe Ala Gly Ser Asn Ser Gln Met Ala Gln Val Gly Asp Gly Asp Asn
 340 345 350
 Ser Pro Leu Met Asn Asn Phe Arg Gln Tyr Leu Pro Ser Leu Pro Gln
 355 360 365
 Ser Val Glu Cys Arg Pro Phe Val Phe Gly Ala Gly Lys Pro Tyr Glu
 370 375 380
 Phe Ser Ile Asp Cys Asp Lys Ile Asn Leu Phe Arg Gly Val Phe Ala
 385 390 395 400
 Phe Leu Leu Tyr Val Ala Thr Phe Met Tyr Val Phe Ser Thr Phe Ala
 405 410 415
 Asn Ile Leu Arg Asn Lys Glu Ser
 420

<210> 17
 <211> 266
 <212> DNA
 <213> Helper phage VCSM13

<400> 17
 ggttatgcgt gggcgatggg tgttggtcatt gtcggcgcaa ctatcggtat caagctgttt 60
 aagaaattca cctcgaaagc aagctgataa accgatacaa ttaaaggctc cttttggagc 120
 cttttttttt ggagattttc aacaagcttc tgcgtaataa ggagtcttaa tcatgccagt 180
 tcttttgggt attccgttat tattgcgttt cctcggtttc cttctggtaa ctttgttcgg 240
 ctatctgcta acttttctta aaaagg 266

<210> 18
 <211> 832
 <212> DNA
 <213> Helper phage VCSM13

```

<400> 18
ttgactccct gcaagcctca gcgaccgaat atatcggtta tgcgtgggcg atggttgttg      60
tcattgtcgg cgcaactatc ggtatcaagc tgtttaagaa attcacctcg aaagcaagct      120
gataaacgga tacaattaaa ggctcctttt ggagcctttt tttttggaga ttttcaacgt      180
gaaaaaatta ttattcgcaa ttccttttagt tgttcctttc tattctcact ccgctggatc      240
ctctggttcc ggtgattttg attatgaaaa tatggcaaac gctaataagg gggctatgac      300
cgaaaatgcc gatgaaaacg cgctacagtc tgacgctaaa ggcaaacttg attctgtcgc      360
tactgattac ggtgctgcta tcgacggttt cattggtgac gtttccggcc ttgctaattg      420
taatggtgct actggtgatt ttgctggctc taattcccaa atggctcaag tcggtgacgg      480
tgataattca cctttaatga ataatttcog tcaatattta ccttcccttc ctcaatcggc      540
tgaatgtcgc ccttttgtct ttcgcgctgg taaaccatat gaattttcta ttgattgtga      600
caaaataaac ttattccgtg gtgtctttgc gtttctttta tatgttgcca cctttatgta      660
tgtattttcg acgtttgcta acatactgcg taataaggag tcttaatcat gccagttctt      720
ttgggtattc cgttattatt gcgtttcctc ggtttccttc tggtaacttt gttcggctat      780
ctgctaactt ttcttaaaaa gggcttcggc aagatagcta ttgctatttc at              832

```

CLAIMS

1. A helper phage displaying a mixture of proteins on its surface, characterized in that said mixture comprises a) a g3 phage coat protein or a part thereof capable of mediating infection of a host by the helper phage, and b) a mutant form of the g3 phage coat protein, wherein the D1 region, D2 region or both regions of the g3 protein are deleted.
2. A helper phage according to claim 1, characterized in that the genome of the helper phage comprises a nucleic acid encoding the mutant form of the g3 phage coat protein.
3. A helper phage according to claim 1 or 2, characterized in that the genome of the helper phage does not comprise a nucleic acid encoding the g3 phage coat protein or a part thereof capable of mediating infection of a host by the helper phage.
4. A helper phage according to any one of claims 1-3, characterized in that the genome of the helper phage further encodes other proteins that are essential for the assembly of the helper phage.
5. A helper phage according to any one of claims 1-4, characterized in that the genomic DNA sequences of the helper phage are based on the genomic sequences of VCSM13 or M13K07.
6. A helper phage according to any one of claims 1-5, characterized in that the helper phage is infectious.
7. A method of producing a helper phage displaying a mixture of proteins on its surface, said mixture comprising a g3 phage coat protein or a part thereof capable of mediating infection of a host by the helper phage and a mutant form of the g3 phage

coat protein, wherein the D1 region, D2 region or both regions of the g3 protein are deleted, the method comprising the steps of:

- a) providing a host cell with a first nucleic acid encoding the g3 phage coat protein or part thereof capable of mediating infection of a host by the helper phage;
- b) providing the host cell with a second nucleic acid encoding the mutant form of the g3 phage coat protein, said second nucleic acid further comprising all other proteins that are essential for the assembly of the helper phage in the host cell; and
- c) culturing the host cell to allow assembly of the helper phage.

8. A method according to claim 7, characterized in that the expression of the g3 phage coat protein or a part thereof capable of mediating infection of a host by the helper phage, the mutant form of the g3 phage coat protein, or both is regulatable by altering the culturing conditions of the host cell.

9. A method according to claim 7 or 8, characterized in that the expression of the g3 phage coat protein or a part thereof capable of mediating infection of a host by the helper phage, the mutant form of the phage coat protein, or both is under control of a regulatable promoter.

10. A method according to claim 9, characterized in that the regulatable promoter comprises the AraC/BAD promoter, the psp promoter, the lac promoter, or a functional equivalent thereof.

11. A method according to any one of claims 7-10, characterized in that the first nucleic acid and the second nucleic acid each comprise a unique selection marker.

12. A method according to any one of claims 7-11, characterized in that the first nucleic acid and the second nucleic acid each comprises a unique origin of

replication.

13. A method according to any one of claims 7-12, characterized in that the first nucleic acid and the second nucleic acid comprise codons that essentially do not lead to a homologous recombination event between the first nucleic acid and the second nucleic acid.

14. A helper phage obtainable by a method according to any one of claims 7-13.

15. A method of producing a chimaeric phage particle, the method comprising the steps of:

- a) providing a host cell with a first nucleic acid encoding a fusion protein, wherein the fusion protein comprises a proteinaceous molecule fused to a g3 phage coat protein or a part thereof capable of mediating infection of a host by the chimaeric phage particle;
- b) providing the host cell with a second nucleic acid encoding a mutant form of the g3 phage coat protein, wherein the mutant form comprises a deletion of the D1 region, D2 region or both regions of the g3 phage coat protein, said host cell further comprising a nucleic acid encoding all other proteins that are essential for the assembly of the chimaeric phage particle in the host cell; and
- c) culturing the host cell to allow assembly of the chimaeric phage particle, the chimaeric phage particle displaying a mixture of proteins on its surface, the mixture comprising the mutant form of the g3 phage coat protein and the fusion protein, wherein the fusion protein is attached to the chimaeric filamentous phage particle via the g3 phage coat protein or a part thereof capable of mediating infection of a host by the chimaeric phage particle.

16. A method of producing a chimaeric phage particle, the method comprising the steps of:

- a) providing a host cell with a first nucleic acid encoding a fusion protein, wherein the fusion protein comprises a proteinaceous molecule fused to a g3 phage coat protein or a part thereof capable of mediating infection of a host by the chimaeric phage particle;
- b) providing the host cell with a second nucleic acid encoding a mutant form of the g3 phage coat protein, wherein the mutant form comprises a deletion of the D1 region, D2 region or both regions of the g3 phage coat protein, the second nucleic acid further encoding all other proteins that are essential for the assembly of the chimaeric phage particle in the host cell; and
- c) culturing the host cell to allow assembly of the chimaeric phage particle, the chimaeric phage particle displaying a mixture of proteins on its surface, the mixture comprising the mutant form of the g3 phage coat protein and the fusion protein, wherein the fusion protein is attached to the chimaeric phage particle via the g3 phage coat protein or a part thereof capable of mediating infection of a host by the chimaeric phage particle.

17. A method according to claim 16, characterized in that the second nucleic acid is provided by a helper phage to the host cell.

18. A method according to claim 17, characterized in that the helper phage is a helper phage according to one of claims 1-6 or claim 14.

19. A method according to any one of claims 15-18, characterized in that the expression of the fusion protein or a part thereof capable of mediating infection of a host by the chimaeric phage particle, the mutant form of the g3 phage coat protein, or both is regulatable by altering the culturing conditions of the host cell.

20. A method according to any one of claims 15-19, characterized in that the expression of the fusion protein or a part thereof capable of mediating infection of a host by the chimaeric phage particle, the mutant form of the g3 phage coat protein, or

both is under control of a regulatable promoter.

21. A method according to claim 20, characterized in that the regulatable promoter comprises the AraC/BAD promoter, the psp promoter, the lac promoter, or a functional equivalent thereof.

22. A method according to any one of claims 15-21, characterized in that the first nucleic acid and the second nucleic acid each comprise a unique selection marker.

23. A method according to any one of claims 15-22, characterized in that the first nucleic acid and the second nucleic acid each comprise a unique origin of replication.

24. A method according to any one of claims 15-23, characterized in that the first nucleic acid and the second nucleic acid comprise codons that essentially do not lead to homologous recombination between the first nucleic acid and the second nucleic acid.

25. A method according to any one of claims 15-24, characterized in that the proteinaceous molecules comprises an antibody, a Fab fragment, a single chain Fv fragment, a variable region, a CDR region, an immunoglobulin or a functional part thereof.

26. Use of a helper phage according to any one of claims 1-6 or claim 14 for generating a phage particle or a phage display library.

27. A method for enriching a first binding pair member that is specific for a second binding pair member, the method comprising the steps of:

- a) preparing a phage collection comprising chimaeric phages comprising a repertoire of first binding pair members by performing a method according to any one claims 15-25,
- b) contacting the obtained phage collection with material comprising said second binding pair member under conditions allowing specific binding between the first binding pair member and the second binding pair member,
- c) removing chimaeric phages that do not bind to the material,
- d) recovering chimaeric phages binding to the material.

28. A method according to claim 27, the method further comprising the steps of:

- a) recovering from the chimaeric phages binding to the material a DNA sequence encoding the first binding pair member,
- b) subcloning the DNA sequence in a suitable expression vector
- c) expressing the DNA sequence in a suitable host under conditions allowing production of the first binding pair member.

29. A method according to claim 27 or 28, wherein the first binding pair member is selected from the group consisting of an antibody or fragment thereof, a single chain Fv fragment, a Fab fragment, a variable region, a CDR region, and an immunoglobulin.

30. A method according to claim 28 or 29, wherein the host is a mammalian cell.




Fig. 1




Fig. 2

3/17




Fig. 3

4/17

No inserted ORF Inserted ORF




Fig. 4

5/17




Fig. 5

6/17




Fig. 6A

Fig. 6B

1	TGCCTGACTCCCCGTCGTGTAGATAACTACGATACGGGAGGGCTTACCATCTGGCCCCAGTGTGCAATGATACCGCGAGACCCACGCTCACCGGCTCCA
101	GATTTATCAGCAATAAACAGCCAGCCGATTCGAGCTCGCCCGGGATCGACCAGTTGGTGATTTTGAACTTTTGCTTTGCCACGGAACGGTCTGCGTTG
201	TCGGGAAGATCGTGATCTGATCCTTCAACTCAGCAAAAAGTTCGATTTATTCAACAAGCCGCCGTCCTCAAGTCAGCGTAATGCTCTGCCAGTGTTA
301	CAACCAATTAACCAATTCTGATTAGAAAAACTCATCGAGCATCAAAATGAAACTGCAATTTATTATATCAGGATTATCAATACCATATTTTGA AAAAGC
401	CGTTTCTGTAATGAAGGAGAAAACTCACCGAGGCAGTTCCATAGGATGGCAAGATCCTGGTATCGGTCTGCGATTCCGACTCGTCCAACATCAATACAAC
	HindIII
	~~~~~
501	CTATTAATTTCCCCTCGTCAAAAAATAAGGTTATCAAGTGAGAAATCACCATGAGTGACGACTGAATCCGGTGAGATGGCAAAAAGCTTATGCATTTCTTT
601	CCAGACTTGTTCACAGGCCAGCCATTACGCTCGTCATCAAAAATCACTCGCATCAACCAACCGTTATTCAATTCGTGATTGCGCCTGAGCGAGACGAAAT
701	ACGCGATCGCTGTTAAAAAGGACAAATTACAAACAGGAATCGAATGCAACCGCGCAGGAACACTGCCAGCGCATCAACAATATTTTCACCTGAATCAGGAT
801	ATTCTTCTAATACCTGGAATGCTGTTTTCCCGGGGATCGCAGTGGTGAGTAACCATGCATCATCAGGAGTACGGATAAAAATGCTTGATGGTCGGAAGAGG
901	CATAAATTCGGTCAGCCAGTTTAGTCTGACCATCTCATCTGTAAACATCATTTGGCAACGCTACCTTTGCCATGTTTCAGAAAACAACCTCTGGCGCATCGGGC
1001	TTCCCATACAAATCGATAGATTGTGCGACCTGATTGCGGAGCCCATTTATACCCATATAAATCAGCATCCATGTTGGAATTTAATCGCG
1101	GCCTCGAGCAAGACGTTTCCCGTTGAATATGGCTCATAAACCCCTTGTTACTGTTTATGTAAGCAGACAGTTTATTTGTTCAATGATGATATATTTTT
1201	ATCTTGTGCAATGTAACATCAGAGATTTTGAAAACACAACGTGGCTTTCCTCCCCCCCCCTGCAGGTCTCGGGCTATTCTTTTGATTATAAAGGATTT
1301	TGCCGATTTCCGGCCTATTGGTTAAAAATGAGCTGATTAAACAAAAATTAACCGCGAATTTTAAACAAAAATATAACGTTTACAATTTAAAAATATTGCTTA
1401	TACAACTTCCCTGTTTTTGGGGCTTTTCTTATTATCAACCGGGGTACATATGATTGACATGCTAGTTTTTACGATTACCGTTTCATCGATTTCTCTTTGTC
1501	TCCAGACTCTCAGGCAATGACCTGATAGCCTTTGTAGACCTCTCAAAAAATAGTACCCCTCTCCGGCATGAATTTATCAGCTAGAAAGGTTGAAATATCATG
1601	TTGATGGTGATTTGACTGTCTCCGGCCTTTCACCCCTTTTGAATCTTTACCTACACATTACTCAGGCATTGCATTTAAAAATATATGAGGGTTCTAAAAA
1701	TTTTTATCCTTGCGTTGAATAAAGGCTTCTCCCGCAAAAGTATTACAGGGTCATAATGTTTTTGGTACAACCGATTAGCTTTATGCTCTGAGGCTTTA
1801	TTGCTTAAATTTTGCTAAATCTTTTGCCCTTGCCCTGTATGATTTATGGATGTTAACGCTACTACTATTAGTAGAATTGATGCCACCTTTTCAGCTCGCGCCC
1901	CAAAATGAAAAATATAGCTAAACAGGTTATTGACCAATTTGCGAAAAATGTATCTAATGGTCAAACTAAATCTACTCGTTCGCAGAAATTGGGAATCAACTGTTAC
2001	ATGGAATGAAACTTCCAGACACCGTACTTTAGTTGCATATTTAAACACATGTTGAGCTACAGCACCCAGATTCAAGCAATTAAGCTCTAAGCCATCCGC AAAA
2101	ATGACCTCTTATCAAAAGGAGCAATTAAAGGTACTCTCTAATCCTGACCTGTTGGAGTTTGCTTCCGGTCTGGTTCGCTTTGAAGCTCGAATTAAACGC
2201	GATATTTGAAGTCTTTCGGGCTTCCTCTTAATCTTTTTTGATGCAATCCGCTTTGCTTCTGACTATAATAGTCAGGTAAGACCTGATTTTGTGATTTATG
2301	GTCAATCTCGTTTTCTGAACTGTTTAAAGCATTTGAGGGGATTC AATGAATATTTATGACGATTCGCGAGTATTGGACGCTATCCAGTCTAAACAATTTT
2401	ACTATTACCCCTCTGGCAAAACTTCTTTTGCAAAAGCCTCTCGCTATTTTGGTTTTTATCGTCTGTGTAACGAGGGTTATGATAGTGTGCTCTTA
2501	CTATGCCCTCGTAATTCCTTTTGGCGTTATGTATCTGCATTAGTTGAATGTGGTATTCCTAAATCTCAACTGATGAATCTTCTACCTGTAATAATGTTGT
2601	TCCGTTAGTTCGTTTTATTAAACGTAGATTTTTTCTTCCCACGTCCTGACTGGTATAATGAGCCAGTTCTTAAAAATCGCATAAAGTAATTCACAATGATTA
2701	AAGTTGAAATTA AACCATCTCAAGCCCAATTTACTACTCGTTCTGGTGTTTCTCGTCAGGGCAAGCCCTTATTCACTGAATGAGCAGCTTTGTTACGTTGA
2801	TTTGGGTAATGAATATCCGGTTCCTTGTC AAGATTACTCTTGATGAAGGTCAGCCAGCCTATGCGCCTGGTCTGTACACCGTTTCACTGTCTCTTTCAAA
2901	GTTGGTCAGTTCGGTTCCCTTATGATTGACCGTCTGCGCCTCGTTCCGGCTAAGTAACATGGAGCAGGTCGCGGATTTTCGACACAATTTATCAGGCGGATG
3001	ATACAAATCTCCGTTGTACTTTTGTTCGCGCTTGGTATAATCGCTGGGGTCAAAGATGAGTGTTTTAGTGTATTCTTTTCGCCCTCTTTTCGTTTAGGTTG

**SUBSTITUTE SHEET (RULE 26)**

Fig. 6B, contd.

	PacI	
	~~~~~	
5901	TGATTTTGATAAAATTCACTATTGACTCTTCTCAGCGTCTTAATCTAAGCTATCGCTATGTTTCAAGGATTCTAAGGGAAAAATTAATTAATAGCGACGAT	
6001	TTACAGAAAGCAAGGTTATTCACTCACATATATTGATTTATGTACTGTTTCCATTAAAAAAGGTAATTCAAATGAAAATTTGTTAAATGTAATTAATTTTGT	
6101	TTCTTGATGTTTGTTCATCATCTTCTTTTGCTCAGGTAATTGAAATGAATAATTCGCCCTCTGCGCGATTTTGTAACCTTGCTATTCAAAGCAATCAGGCG	
6201	AATCCGTTATTGTTTCTCCGATGTAAAAAGGTACTGTTACTGTATATTCACTGACGTTAAACCTGAAAAATCTACGCAATTTCTTTATTCTGTTTACG	
6301	TGCAAAATAAATTTTGATATGGTAGGTTCTAACCCCTTCAACCTTCCATTATTCAGAAGTATAATCCAAACAATCAGGATTATATGATGAATTGCCATCATCTGATAAT	
6401	CAGGAATATGATGATAAATCCCGCTCCCTTCTGGTGTTTCTTTGTCCGCAAAATGATAATGTTACTCAAACTTTTAAAAATTAATAACGTTTCGGGCAAGG	
6501	ATTTAATACGAGTTGTCGAATTGTTGTAAAGTCTAATACTTCTAAATCCTCAAATGTATTATCTATTGACGGCTCTAATCTAATCTATTAGTTGTAGTCTCC	
6601	TAAAGATAATTTTAGATAAACCTTCCCTCAATTCCTTTCAACTGTGTGATTTGCCAACTGACCAAGATATTGATTGAGGGTTTGATATTTGAGGTTTCAGCAAGGT	
6701	GATGCTTTAGATTTTTCATTTGCTGCTGCTCAGCGTGGCACTGTTGCAGGCGGTGTTAATACTGACCGCCTCACCTCTGTTTATCTTCTGCTGGTG	
6801	GTTCGTTCCGGTATTTTAATGGCGATGTTTTAGGGCTATCAGTTCGCGCATTAAGACTAATAGCCATTCAAAAAATATTGCTGTGTCACGTTATCTTTAC	
6901	GCTTTCAGGTCAGAAGGTTCTATCTGTGTGGCCAGAAATGTCCTTTTATTACTGGTCGTGACTGGTGAATCTGCCAATGTTCAATTAATCCATTTCAG	
7001	ACGATTGAGCGTCAAAATGTAGGTATTTCCATGAGCGTTTTTCCTGTTGCAATGGCTGGCGGTAATATTGTTCTGGATATTACCAGCAAGGCCGATAGTT	
7101	TGAGTCTTCTACTCAGGCAAGTGATGTTATTACTAATCAAAGAAGTATGTACAACGGTTAATTTGCGTGATGGACAGACTCTTTTACTCGGTGGCCT	
7201	CAC TGATTATAAAAAACACTTCTCAGGATTCTGGCGTACCGTTCCTGTCTAAAAATCCCTTTAATCGGCCCTCCTGTTTAGCTCCCGCTCTGATTCTAAACGAG	
7301	GAAAGCACGTTATACGTGCTCGTCAAAGCAACCATAGTACGGGCCCTGTAGCGCGGCATTAAAGCGCGGGGTGGTGGTTACGGCGCAGCGTGACCGCT	
7401	ACACTTGCCAGCGCCCTAGCGCCCGCTCCTTTTCGCTTTCTTCCCTTCTCGCCACGTTCCCGGCTTTCCCGCTCAAAGCTCTAAATCGGGGGCTCC	
7501	CTTTAGGGTTCCGATTTAGTGCTTTACGGCACCTCGACCCCAAAAAACTTGATTGGGTGATGGTTCAAGTAGTGGCCATCGCCCTGATAGACGGTTTTT	
7601	TCGCCCTTTGACGTTGGAGTCCACGTTCTTTAATAGTGGA CTCTGTTCCAAACTGGAAACAACACTCAACCTATCTCGGACGGATCGCTTCATGTGGC	
7701	AGGAGAAAAAGGCTGCACCGGTGCGTCAGCAGAAATATGTCGCTACAGGATATATTCGGCTTCCTCGCTCACTGACTCGCTACGCTCGGTCGTTTCGACTGC	
7801	GGCGAGCGGAAATGGCTTACGAAACGGGGCGGAGATTTCCTGGAAAGATGCCAGGAAGTACCTAACAGGGAAGTGAGAGGCGCGGCAAGCCGTTTTC	
7901	CATAGGCTCCGCCCCCTGACAAGCATCACGAAATCTGACGCTCAAATCAGTGGTGCGAAACCCGACAGGACTATAAAGATACCAGGCGTTTCCCCCTG	
8001	GCGGCTCCCTCGTCGCTCTCCTGTTCCCTGCTTTCGGTTTACCGGTGTCATTCCGCTGTTATGCGCGGTTTGTCTCATTCACCGCTGACACTCAGTT	
8101	CCGGTAGGCAGTTCGCTCCAAGCTGGACTGTATGCACGAACCCCGTTTCAGTCCGACCGCTGCGCCTTATCCGGTAACTATCGTCTTGAGTCCAAACCC	
8201	GGAAAGACATGCAAAAGCACCACTGGCAGCAGCCACTGGTAATTGATTTAGAGGAGTTAGTCTTGAAGTCATCGCCCGGTTAAGGCTAAACTGAAAGGAC	
8301	AAGTTTGGTGACTGCGCTCCTCCAAGCCAGTTACCTCGGTTCAAAGAGTTGGTAGCTCAGAGAACCTTCGAAAAACCCGCCCTGCAAGGCGGTTTTCG	
8401	TTTTCAGAGCAAGAGATTACGCGCAGACCAAAACGATCTCAAGAGAGATCATCTTATTAAAGGGTCTGACGCTCAGTGGAAACGAAAACTCACGTTAAGGGA	
8501	TTTTCGTCATGAGATTATCAAAAAGGATCTTCACCTAGATCCTTTTAAATTA AAAATGAAGTTTAAATCAATCTAAAGTATATATGAGTAAACTTGGTC	
8601	TGACAGTTACCAAATGCTTAATCAGTGAGGCACCTATCTCAGCGATCTGTCTATTTTCGTTTCATCCATAGT	

10/17




Fig. 6C

11/17

G3-minus protein

GGTTATGCGTGGGCGATGGTTGTTGTCATTGTCGGCGCAACTATCGGTATCAAGCTGTTTAAGAAATTCACCTCG
AAAGCAAGCTGATAAACCGATACAATTAAAGGCTCCTTTTGGAGCCTTTTTTTTGGAGATTTTCAACAAGCTTC
TGCGTAATAAGGAGTCTTAATCATGCCAGTTCTTTGGGTATTCCGTTATTATTGCGTTTCCTCGGTTTCCTTCT
GGTAACTTTGTTCCGGCTATCTGCTAACTTTTCTTAAAAAGG

Fig. 6D

12/17




Fig. 7A

13/17

D3 domain of g3 Protein

TTGACTCCCTGCAAGCCTCAGCGACCGAATATATCGGTTATGCGTGGGCGATGGTTGTTGTCATTGTCGGCGCAA
CTATCGGTATCAAGCTGTTTAAGAAATTCACCTCGAAAGCAAGCTGATAAACCGATACAATTAAAGGCTCCTTTT
GGAGCCTTTTTTTTTTGGAGATTTTCAACGTGAAAAAATTATTATTCGCAATTCCTTTAGTTGTTCTTTCTATTC
TCACTCCGCTGGATCCTCTGGTTCCGGTGATTTTGATTATGAAAATATGGCAAACGCTAATAAGGGGGCTATGAC
CGAAAATGCCGATGAAAACGCGCTACAGTCTGACGCTAAAGGCAAACCTTGATTCTGTCGCTACTGATTACGGTGC
TGCTATCGACGGTTTCATTGGTGACGTTTCCGGCCTTGCTAATGGTAATGGTGCTACTGGTGATTTTGCTGGCTC
TAATTCCCAAATGGCTCAAGTCGGTGACGGTGATAATTCACCTTTAATGAATAATTTCCGTCAATATTTACCTTC
CCTTCCTCAATCGGTTGAATGTCGCCCTTTTGTCTTTCGCGCTGGTAAACCATATGAATTTTCTATTGATTGTGA
CAAAATAAACTTATTCCGTGGTGTCTTTGCGTTTCTTTTATATGTTGCCACCTTTATGTATGTATTTTCGACGTT
TGCTAACATACTGCGTAATAAGGAGTCTTAATCATGCCAGTTCTTTTGGGTATTCCGTTATTATTGCGTTTCCTC
GGTTTCCTTCTGGTAACTTTGTTTCGGCTATCTGCTAACTTTTCTTAAAAAGGGCTTCGGTAAGATAGCTATTGCT
ATTTCAT

Fig. 7B

A	GCG GCA GCC GCT
C	TGC TGT
D	GAT GAC
E	GAA GAG
F	TTT TTC
G	GGT GGC
H	CAT CAC
I	ATT ATC ATG
K	AAA
L	CTG
M	ATG
N	AAC AAT
P	CCG
Q	CAG
R	CGT CGC
S	AGC TCT
T	ACC ACG
V	GTG GTT GTC
W	TGG
Y	TAT TAC
STOP	TAA TGA

Fig. 8

15/17




Fig. 9

16/17




Fig. 10

17/17




Fig. 11