



(86) **Date de dépôt PCT/PCT Filing Date:** 2006/04/27
 (87) **Date publication PCT/PCT Publication Date:** 2006/11/09
 (45) **Date de délivrance/Issue Date:** 2016/06/14
 (85) **Entrée phase nationale/National Entry:** 2007/11/29
 (86) **N° demande PCT/PCT Application No.:** US 2006/016050
 (87) **N° publication PCT/PCT Publication No.:** 2006/118954
 (30) **Priorités/Priorities:** 2005/04/29 (US60/676,043);
 2005/09/13 (US60/716,630)

(51) **Cl.Int./Int.Cl. A61K 33/24** (2006.01)

(72) **Inventeurs/Inventors:**
 MCGINNIS, JAMES F., US;
 CHEN, JUNPING, US;
 WONG, LILY, US;
 SEZATE, STEVE, US;
 SEAL, SUDIPTA, US;
 PATIL, SWANAND, US

(73) **Propriétaires/Owners:**
 UNIVERSITY OF CENTRAL FLORIDA RESEARCH
 FOUNDATION, INC., US;
 UNIVERSITY OF OKLAHOMA, CORP., US

(74) **Agent:** GOWLING WLG (CANADA) LLP

(54) **Titre : INHIBITION D'ESPECES REACTIVES DE L'OXYGENE ET PROTECTION DE CELLULES MAMMALIENNES**
 (54) **Title: INHIBITION OF REACTIVE OXYGEN SPECIES AND PROTECTION OF MAMMALIAN CELLS**

(57) **Abrégé/Abstract:**

Methods and compositions useful for neuronal protection in retinal cells in vitro and the protection of mammalian cells from reactive oxygen species in vivo are provided. Ultrafine nano-size cerium oxide particles, less than 10 nanometers in diameter, have been provided to decrease reactive oxygen species (ROS) in retina tissue that generates large amounts of ROS. These reactive oxygen species (ROS) are involved in light-induced retina degeneration and age-related macular degeneration (AMD). Cerium oxide nanoparticles have been used to promote the lifespan of retinal neurons and protect the neurons from apoptosis induced by hydrogen peroxide in vitro and in vivo. The neuronal protection in retinal cells is achieved by decreasing generation of intracellular reactive oxygen species (ROS). Thus, cerium oxide particles are used to promote the longevity of retinal neurons in vitro and mammalian cells in vivo.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

CORRECTED VERSION

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
9 November 2006 (09.11.2006)

PCT

(10) International Publication Number
WO 2006/118954 A3

(51) International Patent Classification:

A61K 31/40 (2006.01) G02B 3/00 (2006.01)
 A61K 31/685 (2006.01) G02B 7/00 (2006.01)
 A61K 31/66 (2006.01) B32B 23/02 (2006.01)
 G02C 7/02 (2006.01)

(21) International Application Number:

PCT/US2006/016050

(22) International Filing Date: 27 April 2006 (27.04.2006)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/676,043 29 April 2005 (29.04.2005) US
 60/716,630 13 September 2005 (13.09.2005) US

(71) Applicant: UNIVERSITY OF CENTRAL FLORIDA
RESEARCH FOUNDATION, INC. [US/US]; 12443 Re-
search Parkway, Suite 302, Orlando, FL 32826-3252 (US).(71) Applicant (for all designated States except US): UNIVER-
SITY OF OKLAHOMA, CORP. [US/US]; 1000 Stanton
L. Young Blvd., Oklahoma City, OK 73104 (US).(72) Inventors: MCGINNIS, James, F.; 708 Foxhill Drive,
Edmond, OK 73034 (US). CHEN, Junping; 14425 No.
Pennsylvania Avenue, Unit #B, Oklahoma City, OK 73134
(US). WONG, Lily; 14200 No. May Avenue, Apt. #224,
Oklahoma City, OK 73134 (US). SEZATE, Steve; 2301
N.W. 118th Street, Oklahoma City, OK 73120 (US).
SEAL, Sudipta; 1637 Riverbirch Avenue, Oviedo, FL
32765 (US). PATIL, Swanand; 11931 Pasteur Drive,
Orlando, FL 32826 (US).(74) Agent: STEINGBERGER, Brian; Law Offices of Brian
S. Steinberger, P.A., 101 Brevard Avenue, Cocoa, FL
32922 (US).(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV,
LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI,
NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG,
SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US,
UZ, VC, VN, YU, ZA, ZM, ZW.(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,
FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT,
RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA,
GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

(88) Date of publication of the international search report:
24 May 2007(48) Date of publication of this corrected version:
23 August 2007(15) Information about Correction:
see PCT Gazette No. 34/2007 of 23 August 2007For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: INHIBITION OF REACTIVE OXYGEN SPECIES AND PROTECTION OF MAMMALIAN CELLS

(57) Abstract: Methods and compositions useful for neuronal protection in retinal cells in vitro and the protection of mammalian cells from reactive oxygen species in vivo are provided. Ultrafine nano-size cerium oxide particles, less than 10 nanometers in diameter, have been provided to decrease reactive oxygen species (ROS) in retina tissue that generates large amounts of ROS. These reactive oxygen species (ROS) are involved in light-induced retina degeneration and age-related macular degeneration (AMD). Cerium oxide nanoparticles have been used to promote the lifespan of retinal neurons and protect the neurons from apoptosis induced by hydrogen peroxide in vitro and in vivo. The neuronal protection in retinal cells is achieved by decreasing generation of intracellular reactive oxygen species (ROS). Thus, cerium oxide particles are used to promote the longevity of retinal neurons in vitro and mammalian cells in vivo.



WO 2006/118954 A3

INHIBITION OF REACTIVE OXYGEN SPECIES AND PROTECTION OF MAMMALIAN CELLS

5

FIELD OF THE INVENTION

This invention relates to biological uses of nanoceria particles, and in particular to methods and compositions useful for neuronal protection in retinal cells *in vitro* and the protection of mammalian cells from reactive oxygen species *in vivo*.

10

BACKGROUND AND PRIOR ART

Cerium is a silvery metallic element, belonging to the lanthanide group. Cerium Oxide (CeO₂) is used in precision polishing and lapping applications. Ultra fine nano-size cerium oxide, less than 10 nanometers, is more efficient for coating purposes. Recently, it was reported by B. Rzigalinski et al. that nanoparticles prolong the life of cortical neurons in culture 4 fold over the cells without treatment; decreased the intracellular Ca²⁺ concentration and prevented UV damage of cortical neurons. Based on its chemical characteristics, this effect is partially due to a decrease of reactive oxygen species (ROS).

15
20

Retina tissue generates a large amount of ROS which are involved in light-induced retina degeneration and age-related macular degeneration (AMD). The present invention tests the hypothesis that nanoparticles can promote the lifespan of retinal neurons in culture and protect them from apoptosis induced by hydrogen peroxide (H₂O₂) *in vitro* by decreasing the intracellular concentration of reactive oxygen species.

In 2004, T. H. Margrain et al. discuss the state of research in the treatment of age-related macular degeneration in *Progress in Retinal and Eye Research*, 2004, 23: 523-531, "Do Blue Light Filters Confer Protection Against Age-Related Macular Degeneration?" The problem of apoptosis in the body is discussed in an article by P. Moongkarndi et al. in "Antiproliferation, Antioxidation and Induction of Apoptosis by Garcinia Mangostana (Mangosteen) on SKBR3 Human Breast Cancer Cell Line", *Jl. of Ethnopharmacology*, 2004, 90: 161-166.

Often persons suffering from light-induced retina degeneration and age-related macular degeneration (AMD) are without satisfactory remedies to prevent the eventual outcome of blindness. There are some proteins available for neuronal protection of retinal cells, however, they are big molecules and over time their effect may fade away.

It is desirable to find reliable solutions to prolong the lifespan of retinal neurons so that blindness is avoided for persons with retina degeneration and AMD.

In addition to diseases of the eye, many human diseases are due to the death of cells in specific tissues or organs. The majority of those diseases are due to accumulation of metabolic insults from reactive oxygen species originating within or outside of the cells. These diseases include all forms of blindness whether hereditary, light-induced, or physical damage such as occurs in retinal detachment. In addition, damage due to

ageing, stroke, cardiac infarction, burns, etc, which proceed through reactive oxygen species, can be addressed with the nanoceria particles of the present invention.

The present invention promotes a longer lifespan for retinal neurons. The greatest benefit of the nanoceria is its ability to get inside the cells and provide protection from reactive oxygen species (ROS); other body systems and tissues can also be protected from damage due to ROS.

SUMMARY OF THE INVENTION

A primary objective of the present invention is to promote the lifespan of retinal neurons in culture.

A secondary objective of the present invention is to protect retinal neurons from apoptosis induced by hydrogen peroxide (H_2O_2) *in vitro*.

A third objective of the present invention is to protect retinal neurons from apoptosis induced by reactive oxygen species *in vivo*.

A fourth objective of the present invention is to inhibit the rise in the intracellular concentration of reactive oxygen species (ROS).

A fifth objective of the present invention is to provide method for inhibiting apoptosis induced by H_2O_2 of retinal neurons *in vitro* in a dose and time dependent manner.

A sixth objective of the present invention is to provide method for inhibiting apoptosis induced by reactive oxygen species in retinal neurons *in vivo* in a dose and time dependent manner.

A seventh objective of the present invention is to provide a method for preventing an increase in the intracellular reactive oxygen species (ROS) in a dose and time dependent manner.

An eighth objective of the present invention is to manufacture and modify cerium oxide (CeO_2) nanoparticles for effective use in neuronal protection in retinal cells.

A ninth objective of the present invention is to manufacture and modify cerium oxide (CeO_2) nanoparticles for effective use in mammalian cells *in vivo* to inhibit damage caused by reactive oxygen species (ROS).

A preferred composition for promoting longevity of retinal neurons includes at least one of $\text{CeO}_{.n1}$ wherein $0 < n1 < 2$, and $0 < n2 < 3$ in the form of ultra-fine particles. The preferred ultra-fine particles have a diameter in a range between approximately 1 nanometer (nm) and approximately 10 nm and the preferred $\text{CeO}_{.n1}$ is further defined as $n1$ equals approximately 2.

A preferred composition for inhibiting apoptosis induced by hydrogen peroxide oxidation of retinal neurons includes at least one of $\text{CeO}_{.n1}$ wherein $0 < n1 < 2$, and $0 < n2 < 3$ in the form of ultra-fine particles. The preferred ultra-fine particles have a diameter in a range between approximately 1 nanometer (nm) and approximately 10 nm and the preferred $\text{CeO}_{.n1}$ is further defined as $n1$ equals approximately 2.

A preferred composition for inhibiting apoptosis of retinal neurons in a dose and time dependent manner includes at least one of $\text{CeO}_{.n1}$ wherein $0 < n1 < 2$, and $0 < n2 < 3$ in the form of ultra-fine particles. The preferred ultra-fine particles have a diameter in a range between approximately 1 nanometer (nm) and approximately 10 nm and the preferred $\text{CeO}_{.n1}$ is further defined as $n1$ equals approximately 2.

A more preferred composition that decreases the concentration of intracellular reactive oxygen species (ROS) includes at least one of $\text{CeO}_{.n1}$ wherein $0 < n1 < 2$, and $0 < n2 < 3$ in the form of ultra-fine particles. The more preferred composition is used in the treatment of diseases of the retina selected from the group consisting of light-induced retina degeneration and age-related macular degeneration, and is also used *in vivo* for the treatment of diseases in mammalian cells to inhibit damage caused by reactive oxygen species (ROS).

The mammalian cells that can be treated by the composition of the present invention, include, but are not limited to, retinal neurons, brain cells, heart cells, skin cells, liver cells, kidney cells and peripheral nervous system cells.

A preferred method for promoting longevity of retinal neurons includes preparing
5 ultra-fine particles of at least one of CeO_2 wherein $0 < n_1 < 2$, and $0 < n_2 < 3$ in a preselected concentration, and adding the preselected concentration of CeO_2 wherein $0 < n_1 < 2$, and $0 < n_2 < 3$ to primary retinal neurons. The preferred ultra-fine particles have a diameter in a range between approximately 1 nanometer (nm) and approximately 10 nm and the CeO_2 is further defined as n_1 equals approximately 2.
10 The preferred preselected concentrations of CeO_2 are in a range between approximately 3 nanomolar (nM) and approximately fifty nanomolar (nM), more preferably in a range between approximately 3 nanomolar (nM) and approximately twenty nanomolar (nM).

It is also preferred that the preselected concentrations of CeO_2 are added to primary retinal neurons *in vitro* and/or administered to mammalian cells *in vivo* to protect
15 the mammalian body system from damage to any tissue due to reactive oxygen species (ROS).

Further objects and advantages of the present invention will be apparent from the following detailed description of a presently preferred embodiment which is illustrated schematically in the accompanying drawings.

20 BRIEF DESCRIPTION OF THE FIGURES

Fig. 1 is a timeline of exposure of a primary retinal neuron culture to treatment with cerium oxide nanoparticles.

Fig. 2 is a graph showing the percentage of apoptotic retinal neurons in culture with and without 5 nanomoles (nM) cerium oxide nanoparticle treatment at different time points.

25 Fig. 3 shows the initial cerium oxide nanoparticle treatments followed by hydrogen peroxide incubation, and time line of exposure.

Fig. 4A is a flow cytometry plot of the control sample of primary retinal neurons with no cerium oxide nanoparticle treatments and no hydrogen peroxide incubation.

Fig. 4B is a flow cytometry plot of retinal neurons in the presence of 100 micromoles (μM) hydrogen peroxide.

5 Fig. 4C is a flow cytometry plot of retinal neurons treated with 1nM cerium oxide in the presence of 100 micromoles (μM) hydrogen peroxide.

Fig. 4D is a flow cytometry plot of retinal neurons treated with 3nM cerium oxide in the presence of 100 micromoles (μM) hydrogen peroxide.

10 Fig. 4E is a flow cytometry plot of retinal neurons treated with 5nM cerium oxide in the presence of 100 micromoles (μM) hydrogen peroxide.

Fig. 4F is a flow cytometry plot of retinal neurons treated with 10nM cerium oxide in the presence of 100 micromoles (μM) hydrogen peroxide.

Fig. 4G is a flow cytometry plot of retinal neurons treated with 20nM cerium oxide in the presence of 100 micromoles (μM) hydrogen peroxide.

15 Fig. 4H is a flow cytometry plot of retinal neurons treated with 20nM cerium oxide.

Fig. 5 shows relative viable retinal neurons with and without incubation of different concentrations of cerium oxide nanoparticles for time periods between approximately 12 hours and approximately 96 hours.

20 Fig. 6 shows the initial cerium oxide nanoparticle treatments, hydrogen peroxide and DCFH-DA incubation, and time line of exposure.

Fig. 7A is a graph of the intracellular level of reactive oxygen species (ROS) of retinal neurons after 30 minutes incubation with cerium oxide nanoparticles.

Fig. 7B is a graph of the intracellular level of reactive oxygen species (ROS) of retinal neurons after 12 hours incubation with cerium oxide nanoparticles.

Fig. 7C is a graph of the intracellular level of reactive oxygen species (ROS) of retinal neurons after 24 hours incubation with cerium oxide nanoparticles.

5 Fig. 7D is a graph of the intracellular level of reactive oxygen species (ROS) of retinal neurons after 96 hours incubation with cerium oxide nanoparticles.

DESCRIPTION OF THE PREFERRED EMBODIMENTS

10

Also, the terminology used herein is for the purpose of description and not of limitation.

The present invention has two embodiments describing in detail the *in vitro* and *in vivo* treatment of mammalian cells with ultra fine nano-size cerium oxide particles, less
15 than 10 nanometers in diameter, to protect the “body system” from damage to any tissue due to reactive oxygen species (ROS).

Hydrogen peroxide (H_2O_2) is one of many reactive oxygen species. In the present invention, H_2O_2 is added directly to cultures for the *in vitro* treatments. H_2O_2 is not added
20 to the live tissue samples, since H_2O_2 is one of the ROS products of light damage. The discussion below confirms that nanoceria particles inhibit all forms of reactive oxygen species (ROS).

For example, the nanoceria particles of the present invention can protect the brain against stroke and reperfusion injury, the heart cells from effects of cardiac infarction, the
25 skin from UV rays and burn injuries. Neurodegeneration (e.g., Alzheimer’s, Parkinson’s,

dementia, amyotrophic lateral sclerosis) and potentially mental retardation (due to loss of brain cells) within the central and peripheral nervous systems can also be inhibited. This protection can extend to diseases which produce chronic problems such as cirrhosis of the liver or kidney or the multi-organ effects of aging itself. The nanoceria can become the
5 universal treatment for all major and minor diseases and events which involve reactive oxygen species.

The nanoceria (CeO₂ nanoparticles) have the ability to destroy toxic products of metabolism known as reactive oxygen species (ROS). It has been shown in one embodiment that the nanoceria particles prevent the ROS induced death of mammalian
10 retinal neurons in culture (*in vitro*) and subsequently prolonged the life of the cells in culture and protected the cells from ROS. In the second embodiment, the ability to provide mammalian cell protection *in vivo* is disclosed.

The first embodiment of the present invention provides a method and composition for promoting the lifespan of retinal neurons and protecting the nerve cells in the eye
15 from apoptosis induced by hydrogen peroxide (H₂O₂) *in vitro* by decreasing generation of intracellular reactive oxygen species. The treatment of the eye with ultra fine nano-size cerium oxide is a significant advance in biological uses of cerium oxide. Persons afflicted with such conditions as, light-induced retina degeneration and age-related macular degeneration (AMD) have hope for brighter, clearer vision.

20 The examples below provide further detail on the preparation and treatment of retinal nerve cells with CeO₂ nanoparticles.

Example 1

A primary retinal neuron culture is obtained from albino rat pups. Retinae of Sprague-Dawley albino rat pups (0-2-day old) were dissected out and mechanically dissociated in 25ml of DMEM/F12 medium. After being filtered through 230 μ m and 140 μ m sieves, the dissociated cells were centrifuged at 1200rpm for 5min. The cell pellets were re-suspended in the medium to 1×10^5 cells/ml. 1ml of the cell suspension was plated in each well pre-treated with 10 μ g/ml of poly-D-lysine. The cells were maintained in the medium until day 7, when different concentrations of CeO₂ nanoparticles were added to the cultures. The timeline for the addition of CeO₂ nanoparticles is shown in Fig. 1. The treated neuronal cells were harvested on day 14, day 19, day 24 and day 29 after the beginning of treatment on day 7. The percentage of apoptotic retinal neurons in the culture with and without 5nM CeO₂ nanoparticle treatment is shown in Fig. 2 at the 12th day, 14th day, 19th day, 24th day and 29th day. Data are shown in $M \pm S.D.$ Statistics were collected by Student t-test ($n=3$, $*p<0.05$, $**p<0.01$). Fig. 2 confirms that at every testing period the control with no CeO₂ nanoparticle treatment had a higher percentage of apoptotic retinal neurons in the culture, in contrast to the decreased percentage of apoptotic retinal neurons in the cells treated with 5nM CeO₂ nanoparticles.

Example 2

The detection of apoptosis by flow cytometry is illustrated in Figures 3, 4A – 4H and 5. After periods of incubation with CeO₂ or H₂O₂, the cells were washed with serum free medium 3 times, followed by treatment of 1ml of 1 \times trypsin for 2min. After

centrifuging, the cell pellet was resuspended in 500 μ l of 1 \times PBS containing 5 μ l of Annexin V-FITC and 25 μ l of Propidium Iodide (PI). The kit used for the analysis is commercially available from Beckman Coulter and is known as the "ANNEXIN V-FITC Kit." The mixture was incubated on ice for 10 minutes. The fluorescent emissions of FITC and PI were detected by flow cytometry (Beckman Coulter) with the excitation filters of 492 nanometers (nm) and 550nm. The FITC fluorescent emissions signals the presence of cells undergoing apoptosis; whereas, the PI signals with an automatic red color fluorescence the binding to DNA fragments identifying cells in a necrotic stage.

Fig. 3 shows a treatment timeline with CeO₂ incubation after the 7th day of treatment at the time intervals of 12 hours, 24 hours, 72 hours and 96 hours with each treated sample subsequently exposed to 12 hours incubation time with H₂O₂.

Figures 4A – 4H are representative flow cytometry plots of retinal neurons with and without incubation with CeO₂ nanoparticles. Measurements were taken after 96 hours. The cytometry plot shows activity of approximately 10,000 cells in four quadrants, as described below.

C1 represents the percentage of 10,000 cells showing Annexin V positive signals, which are interpreted to indicate the percentage of 10,000 cells undergoing apoptosis.

C2 represents the percentage of 10,000 cells showing both Annexin V and PI positive signals, which is interpreted as the percentage of 10,000 cells which are in late apoptotic or necrotic stage. (AnnexinV signals apoptotic stage; PI signals necrotic stage.)

C3 represents the percentage of 10,000 cells showing *neither* Annexin V nor PI positive signals, which is interpreted as the percentage of 10,000 cells *which are still viable*.

C4 represents the percentage of 10,000 cells showing a PI positive signal, which is interpreted as the percentage of 10,000 cells in a necrotic stage.

Quadrants C2 and C4 show the percentage of 10,000 cells committed to die. In quadrant C1 the percentage of 10,000 cells undergoing apoptosis are shown and some
5 may be salvaged. It is important to observe the percentage of 10,000 cells in quadrant C2 for the efficacy of the cerium oxide treatment of the present invention.

Focusing on the viable cells, the control in Fig. 4A has no CeO₂ nanoparticle treatment and 73.9% of the cell population is viable after 96 hours. Fig. 4B is treated with 100 μM H₂O₂, causing a deadly assault and leaving the lowest percentage (57.7%)
10 of viable cells. The addition of gradually increasing concentrations of cerium oxide with 100 μM H₂O₂, are shown in Figures 4C – 4G.

Fig. 4C is treated with 1nM CeO₂ nanoparticles and 100 μM H₂O₂ and 60.6% of the cell population remains viable. Fig. 4D is treated with 3nM CeO₂ nanoparticles and 100 μM H₂O₂ and 67.0% of the cell population is viable. Fig. 4E is treated with 5nM
15 CeO₂ nanoparticles and 100 μM H₂O₂, with 68.9% of the cell population remaining viable. Fig. 4F is treated with 10nM CeO₂ nanoparticles and 100 μM H₂O₂ and 72.6% of the cell population is viable. Fig. 4G is treated with 20nM CeO₂ nanoparticles and 100 μM H₂O₂ and 72.3% of the cell population remains viable.

Fig. 4H is treated with 20nM CeO₂ nanoparticles without the addition of H₂O₂
20 and 73.4% of the cell population remains viable showing that the CeO₂ nanoparticles alone have no negative effect on the cell population.

The data in Figures 4A - 4H can also be summarized from the analysis of cells undergoing apoptosis as shown in quadrant C1. The control, Fig. 4A shows 16.2% of

cells undergoing apoptosis under normal conditions, without any treatment, after 96 hours. Fig. 4B shows 26.1% of cells undergoing apoptosis after the 100 μM H_2O_2 challenge. Figures 4C – 4G show there are 28.3%, 21.9%, 22.2%, 18.5% and 15.8% of cells undergoing apoptosis with 1, 3, 5, 10 and 20nM CeO_2 nanoparticle treatment, respectively. Fig. 4H shows 15.7% of cells undergoing apoptosis with 20nM CeO_2 nanoparticle treatment, which is a slight improvement over no treatment at all as shown by the control in Fig. 4A.

Thus, cerium oxide nanoparticles inhibit apoptosis in retinal neurons *in vitro* in a dose and time dependent manner.

Fig. 5 shows relative viable retinal neurons with and without incubation of different concentration of CeO_2 nanoparticles for different time periods. The concentration of CeO_2 nanoparticles were 1nM, 3nM, 5nM, 10nM, 20nM each with 100 μM H_2O_2 ; one sample was treated with only 100 μM H_2O_2 and another sample was treated with only 20nM CeO_2 nanoparticles. The measurement of relative viable cells for each group was determined after 12 hours, 24 hours, 48 hours, 72 hours and 96 hours. The effects showed dose and time dependency.

5nM cerium oxide nanoparticles started to decrease the apoptosis at 24h of incubation. As incubation time increased, the protective effect from 5nM nanoparticles was more significantly increased. Additionally, 10nM and 20nM nanoparticles began to have effects late with 96h of incubation. (Statistical analysis was done by ANOVA, and Duncan test for post hoc analysis. Data are presented in $M \pm S.D.$ $n \geq 3$, $*p < 0.05$, $**p < 0.01$). Fig. 5 shows that among the various treatment doses, 5nM CeO_2

nanoparticle treatment gave the earliest response with moderate protective effect and 20nM CeO₂ nanoparticle treatment gave the highest protective effect with late response.

Example 3

Intracellular reactive oxygen species (ROS) production was measured in both
5 CeO₂ nanoparticle treated and control cells using 29,79-dichlorofluorescein diacetate (DCFH-DA, Sigma). Briefly, the retinal neurons were exposed to CeO₂ nanoparticles with different concentrations and various incubation times. After incubation, the cells were incubated with 10μM DCFH-DA (dissolved in dimethylsulfoxide (DMSO)) at 37°C for 30 min. The cells then were incubated with 1mM H₂O₂ at 37°C for 30 min after the
10 excess DCFH-DA was washed with PBS. The cells were harvested as described above. The intensity of fluorescence was detected by flow cytometry with the excitation filter of 485nm. The ROS level was calculated as a ratio: ROS = mean intensity of treated cells divided by mean intensity of control cells.

Fig. 6 is an experimental paradigm showing the beginning of the CeO₂
15 nanoparticle treatment on the 7th day and simultaneous incubation with H₂O₂ and DCFH-DA and a timeline of exposure at 30 minutes, 12 hours, 24 hours, and 96 hours. Figures 7A – 7D show how the CeO₂ nanoparticles decreased the generation of ROS in a dose and time dependent manner. The ROS generated in the groups of 5nM, 10nM, and 20nM nanoparticle incubation were statistically significantly less than the group without
20 treatment at the earliest (12h) of the tested points. 3nM nanoparticle incubation had an effect at 24h. However, 1nM nanoparticles did not show any significant decrease within the tested time point. Statistical analysis was done by ANOVA, and Duncan test for post hoc analysis. Data were shown in M±S.D. n=3, * * p<0.01; *p<0.05.

The cerium oxide of the present invention includes those cerium compounds that have reacted with atmospheric oxygen to have stable oxide layers identified as CeO_x , wherein x is less than 1 or equal to 2 ($0 < x < 2$) and ($0 < x < 3$).

The cerium oxide particles of the present invention are characterized as ultra-fine and are preferably in a size range of from approximately 1 nanometer in diameter to approximately 10 nanometers in diameter; more preferably from approximately 1 nm to approximately 7 nm. A judicious selection of particle size is required by someone skilled in the art and is not a limitation of the present invention.

The results of testing in the above examples document the ability of cerium oxide nanoparticles to promote the longevity of retinal neurons *in vitro* and inhibit apoptosis induced by hydrogen peroxide on retinal neurons *in vitro* in a dose and time dependent manner. It has also been determined that cerium oxide nanoparticles decrease generation of intracellular reactive oxygen species in a dose and time dependent manner. Thus, the present invention represents a significant advance in the treatment in degenerative diseases of the retina, such as, but not limited to light-induced retina degeneration and age-related macular degeneration (AMD).

In the second embodiment of the present invention a rat "light damage" model for retinal degeneration was used as the test system. The data demonstrate that the nanocerium prevented the death of retinal neurons when given prior to the "light insult". The nanocerium particles protected the cells at the time of exposure as well as prevented the subsequent death seen days later in the untreated animals. The *in vivo* route of administration, including, direct injection into the eye, intravenous, intraperitoneal, intramuscular, oral or topically on the eye or skin may improve the result. Similarly, the

time of administration of the nanoparticle, both before or after an insult, is important. Thus, the nanoceria will also prevent the death of retinal cells due to glaucoma, diabetic retinopathy, inherited retinal degeneration (for example, Retinitis Pigmentosa), macular degeneration, retinal detachment or any disease or event which proceeds through the
5 production of ROS. These particles should preserve and prolong vision when administered *in vivo*.

Example 4

Rats were injected intravitreally with 2 microliters (μ l) of nanoceria (concentrations from 0.1 to 1.0 micromolar) three days before they were exposed for six
10 hours to a bright light (2700 LUX) in a light box. The animals were returned to normal lighting for 3.5 days, then killed, the eyes enucleated, fixed, processed for paraffin embedding, sectioned, and either stained with H & E or processed for immunocytochemistry.

The number of photoreceptors remaining was determined with the H&E sections
15 by using a microscope connected to a digital camera to record the images and then measuring the thickness of the outer nuclear layer every 240 microns from the optic nerve along both the superior and the inferior retina to the ora seratta. The data was then plotted as retina thickness versus the distance from the optic nerve head. The changes in other layers of retinal cells were recorded with images but not quantified. Cells actively
20 undergoing apoptosis were also visualized using a commercially available "Apoptosis kit" and recording the microscopic images with a fluorescence microscope.

The data demonstrate that the nanoceria at all concentrations tested prevented the immediate death of retinal cells shortly after exposure to light as well as the ongoing

death seen days later in the untreated animals. We therefore have demonstrated *in vivo* that these nanoceria particles can protect cells within the rat retina from light-induced cell death.

Prior to the present invention, it was not known that nanoceria particles could be used *in vivo* for preventing blindness or death of retinal cells, or for preventing the death *in vivo* of any other cell type in any disease. The chemical properties of CeO₂ nanoparticles enable the destruction of reactive oxygen species (ROS) produced by toxins and/or products of oxygen metabolism within cells.

Thus, nanoceria can protect any cell type from ROS induced damages. Diseases which could possibly be prevented, cured or ameliorated would include hereditary blindness, macular degeneration, glaucoma, diabetic retinopathy, retinal detachment, and other blinding diseases which involve ROS would be potentially solved. Similarly, in other cells within the central nervous system (CNS), neuronal death in strokes, degenerative diseases such as Alzheimer's Disease, Parkinson's, Huntington's Disease, and the death of peripheral nerves are preventable or at least the rate of cell death can be decreased and would result in prolonged function of the cells, tissues, organs, and individual.

The scope of the claims should not be limited by the preferred embodiments set forth in the examples, but should be given the broadest interpretation consistent with the description as a whole.

We claim:

1. A composition of matter consisting essentially of a plurality of nanoceria particles in contact with isolated retinal neurons wherein the plurality of nanoceria particles comprises at least one of CeO_2 and Ce_2O_3 in the form of a plurality of ultra-fine particles, wherein each particle has a diameter in a range between approximately 1 nanometer (nm) and approximately 10 nm, and the combination of the plurality of nanoceria particles in contact with isolated retinal neurons inhibits apoptosis of the isolated retinal neurons in a dose and time dependent manner.
2. The plurality of nanoceria particles in contact with isolated retinal neurons of claim 1, wherein the dosage amount is selected from at least one of 3 (nanomole) nM, 5 nM, 10 nM, and 20 nM of nanoceria particles.
3. Use of ultra-fine particles of at least one of CeO_2 and Ce_2O_3 in concentration of approximately 3 nanomolar (nM) to approximately 50 nanomolar (nM) for promoting longevity of retinal neurons, wherein the ultra-fine particles have a diameter in a range between approximately 1 nanometer (nm) and approximately 10 nm.
4. The use of claim 3, wherein the ultra-fine particles are ultra-fine particles of CeO_2 .
5. The use of claim 4, wherein the concentration of CeO_2 is in a range between approximately 3 nanomolar (nM) and approximately 50 nanomolar (nM).
6. Use of ultra-fine particles of at least one of CeO_2 and Ce_2O_3 in concentration of approximately 3 nanomolar (nM) to approximately 50 nanomolar (nM) for protecting mammalian body system from tissue damage due to reactive oxygen species (ROS), wherein the ultra-fine particles have a diameter in a range between approximately 1 nanometer (nm) and approximately 10 nm.
7. The use of claim 6, wherein the ultra-fine particles are ultra-fine particles of CeO_2 .
8. The use of claim 7, wherein the concentration of CeO_2 is in a range between approximately 3 nanomolar (nM) and approximately 50 nanomolar (nM).




Fig. 1




Fig. 2




Fig. 3




Fig. 4A




Fig. 4B




Fig. 4C




Fig. 4D

5nM CeO₂ + 100μM




Fig. 4E

10nM CeO₂ + 100μM H₂O₂




Fig. 4F

20nM CeO₂ + 100μM H₂O₂




Fig. 4G

20nM CeO₂




Fig. 4H




Fig. 5




Fig. 6




Fig. 7A




Fig. 7B




Fig. 7C




Fig. 7D