639712

Form 1 Regulation 9

COMMONWEALTH OF AUSTRALIA

Patents Act 1952

APPLICATION FOR A STANDARD PATENT

We, AECI LIMITED of Carlton Centre, Commissioner Street, Johannesburg, Transvaal 1645, Republic of South Africa, hereby apply for the grant of a standard patent for an invention entitled:

"ACTIVE INGREDIENT DOSAGE DEVICE"

which is described in the accompanying complete specification.

Details of basic application(s):

Number of basic application: 89/6653

Name of Convention country in which basic application was filed: Republic of South Africa

Date of basic application: 30 August 1989

Our address for service is: Industrial Property Section, ICI Australia Operations Proprietary Limited, P.O. Box 4311, 1 Nicholson Street, Melbourne, Victoria, 3001, Australia

Dated this 29th day of August 1990

AECI LIMITED

By its Patent Attorney

Raymond L. Evans

To: The Commissioner of Patents

FORM 8 Case No: AECI 682

Regulation 12(2)

COMMONWEALTH OF AUSTRALIA

PATENTS ACT 1952

DECLARATION IN SUPPORT OF A CONVENTION APPLICATION FOR A PATENT

In support of the Application made by AECI LIMITED for patent for an invention entitled: Active Ingredient Dosage Device

I, Marthinus Jacobus Frederick Potgieter of AECI Limited

do solemnly and sincerely declare as follows:-

1. I am authorized by the applicants for the patent to make this Declaration on its behalf.
2. The basic application, as defined by Section 141 of the Act, was made in Republic of South Africa on the 30th day of August 1989 by AECI Limited.
3. I/We John Hoy and Phillipus Jansen van Rensburg

   of 9 Cheetah Springs, Cnr. Cheetah Crescent and Gemsbok Lane, Rivonia and of 3 Rooiels Road, Sharonlea, Ext 3, Randburg

   ~~is~~/are the actual inventor(s) of the invention and the facts upon which the applicant is entitled to make the application are as follows:-

   Applicant is the assignee of the invention from the actual inventors

4. The basic application referred to in paragraph 2 of this Declaration was the first application made in a Convention country in respect of the invention, the subject of the application.

Declared at Johannesburg, South Africa this 30th day August of 1990.

.....

TO: The Commissioner of Patents
Canberra

AU9062043

(12) PATENT ABRIDGMENT (11) Document No. AU-B-62043/90
(19) AUSTRALIAN PATENT OFFICE (10) Acceptance No. 639712

(54) Title
ACTIVE INGREDIENT DOSAGE DEVICE

(51)[5] International Patent Classification(s)
A01N 025/12 A01N 043/70 A01N 047/34 A01N 053/00
A01N 043/653

(21) Application No. : 62043/90 (22) Application Date : 30.08.90

(30) Priority Data

(31) Number (32) Date (33) Country
89/6653 30.08.89 ZA SOUTH AFRICA

(43) Publication Date : 07.03.91

(44) Publication Date of Accepted Application : 05.08.93

(71) Applicant(s)
AECI LIMITED

(72) Inventor(s)
JOHN HOY; PHILLIPUS JANSEN VAN RENSBURG

(74) Attorney or Agent
ICI AUSTRALIA OPERATIONS PTY LTD , Industrial Property Section, P.O. Box 4311, MELBOURNE VIC 3001

(56) Prior Art Documents
US 4677003
US 4606909

(57) Claim

1. A method of making a dosage device, which method comprises

admixing at least one pesticide as the active ingredient in solid form at 25°C and which has an average particle size of less than 5 microns, with a disintegrating agent, to provide a compressible mix; and

compressing the mix into a unitary dosage device.

14. A dosage device which is in compressed unitary form and which comprises an admixture of a pesticide which is in solid form at 25°C, is sparingly soluble in water, and has an average particle size of less than 5 microns, and a disintegrating agent, with the device being capable of disintegrating in water to form a suspension of the pesticide in the water, when introduced into sufficient water such that the concentration of the pesticide in the water is above the solubility limit of the pesticide in water.

.../2

21. A method of treating an article or locus, which comprises

introducing a dosage device as claimed in any one of Claims 13 to 20 inclusive, into a predetermined volume of water, with the volume of water being such that the concentration of the active ingredient in the water is greater than the solubility limit of the active ingredient in the water;

allowing the dosage device to disintegrate, thereby to form a suspension of the active ingredient in the water; and

applying the suspension to an article or locus to be treated.

639712

P/00/011
Form 10

PATENTS ACT 1952

COMPLETE SPECIFICATION

(ORIGINAL)

FOR OFFICE USE

Short Title:

Int. Cl:

Application Number:
Lodged:

Complete Specification—Lodged:
Accepted:
Lapsed:
Published:

Priority:

Related Art:

TO BE COMPLETED BY APPLICANT

Name of Applicant: AECI LIMITED

Address of Applicant: 16th Floor Office Tower
Carlton Centre
Commissioner Street
Johannesburg 2001 South Africa

Actual Inventor: John HOY and
Phillipus Jansen Van RENSBURG

Address for Service: Industrial Property Section
ICI Australia Operations Proprietary Limited
1 Nicholson Street, P.O. Box 4311
Melbourne 3001, Victoria, Australia.

Complete Specification for the invention entitled: "ACTIVE INGREDIENT DOSAGE DEVICE"

The following statement is a full description of this invention, including the best method of performing it known to me:—*

---

* Note: The description is to be typed in double spacing, pica type face, in an area not exceeding 250 mm in depth and 160 mm in width, on tough white paper of good quality and it is to be inserted inside this form.

14599/78—L Printed by C. J. THOMPSON, Commonwealth Government Printer, Canberra

ABSTRACT

A method of making a dosage device comprises admixing at least one active ingredient which is in solid form at 25°C and which has an average particle size of less than 5 microns, with a disintegrating agent, to provide a compressible mix; and compressing the mix into a unitary dosage device.

THIS INVENTION relates to a dosage device. It relates also to a method of forming a dosage device, and to a method of treating an article or locus.

According to a first aspect of the invention, there is provided a method of making a dosage device, which method comprises

admixing at least one active ingredient which is in solid form at 25°C, and which has an average particle size of less than 5 microns, with a disintegrating agent, to provide a compressible mix; and

compressing the mix into a unitary dosage device.

The method may include comminuting the active ingredient in solid form and having an average particle size greater than 5 microns, to have an average particle size of less than 5 microns. The active ingredient may be comminuted sufficiently to have an average particle size of 1-3 microns. In one embodiment of the invention, the comminution may be effected by dry milling the active ingredient, eg by means of micronization, to the desired particle size. The milled active ingredient may then be admixed with the disintegrating agent in particulate form. In

another embodiment of the invention, the comminution may be effected by forming a suspension concentrate of the active ingredient in a suitable carrier liquid in which the active ingredient is non-soluble or sparingly soluble, and wet milling the suspension concentrate to obtain the desired active ingredient particle size. The supension concentrate may then be admixed with the disintegrating agent by absorbing it into the disintegrating agent. The milled suspension concentrate may, however, be spray dried, if desired, and then admixed with the disintegrating agent.

While the active ingredient can be any suitable active ingredient, such as a therapeutic agent, anthelmintic, a pigment or dye, or the like, the Applicant believes that the method will find particular, but thus not necessarily exclusive, application in making dosage devices in which the active ingredient is a pesticide, eg an insecticide, herbicide, fungicide, or the like.

The pesticide may be one which is sparingly soluble in water, with water hence being the carrier liquid for use in forming the suspension concentrate and the concentration of the pesticide in the water, in the suspension concentrate, thus being greater than the solubility limit of the pesticide in water. The pesticide may have a water solubility of less than 1000 mg/l at 25°C, preferably less than 50 mg/l at 25°C. Preferably, the pesticide should have a melting point exceeding 70°C.

The pesticide may be a herbicide such as atrazine, simazine, cyanazine, terbuthylazine, diuron,

chlorsulphuron, metsulfuron, tralkoxydin, or 2-(2-chloro-4-mesylbenzoyl) cyclohexane-1,3-dione; an insecticide such as deltamethrin, lindane, carbaryl, endosulfan, or carbofuran; a fungicide such as thiophanate methyl, carbendazim, flutriafol, hexaconazole, chlorothalonil, copper oxychloride, captan or thiram; or an acaricide such as hexythiazox, cyhexatin or amitraz.

The dosage device may comprise, on a mass basis

active ingredient	-	5 - 60 parts
self disintegrating agent	-	20 - 70 parts

The disintegrating agent may be capable of disintegrating by effervescing or swelling on contact with water. When it is capable of swelling on contact with water, it may be a cross-linked polyvinyl pyrrolidone which also acts as a binder. For example, the cross-linked polyvinylpyrrolidone may then be that available under the trade name POLYPLASDONE XL from GAF Corp., or that available under the trade name KOLLIDON CL from BASF. However, it can instead be any other suitable disintegrating agent capable of swelling on contact with water such as a modified cellulose gum, for example that available under the trade name AC-DI-SOL from FMC Corporation, or a sodium starch glycolate such as that available under the trade name EXPLOTAB from Protea Chemical Services. When it is capable of effervescing on contact with water, it may be an acid and base combination such as tartaric acid and an alkali metal carbonate or bicarbonate, eg sodium bicarbonate.

The method may include admixing at least one further substance with the active ingredient,

before comminution thereof and/or after comminution thereof. Thus, when the suspension concentrate is formed, the method may include adding at least one dispersing agent to the suspension concentrate before wet milling thereof. When dry milling is effected, the dispersing agent may be added before or after the milling. The dispersing agent may be a surfactant such as that conventionally used in a wettable powder or suspension concentrate formulation, for example sodium lignosulphonate; sodium naphthalene sulphonic acid/formaldehyde condensate; sodium alkyl aryl sulphonate; a nonyl phenol alkylene oxide, such as nonyl phenol ethylene oxide condensate or nonyl phenol ethylene/propylene oxide; a sodium lauryl sulphate which also acts as wetting agent, such as that available under the trade name EMPICOL LZ from Lankro; a sodium diisopropyl naphthalene sulfonate which also acts as wetting agent, such as that available under the trade name AEROSOL OS from Cyanamid; a sodium salt of naphthalene sulfonic acid formaldehyde condensate, such as that available under the trade name TAMOL NNO from BASF; oxyethylated polyarylphenol phosphate, which is a dispensing agent in aqueous media, and an example of which is obtainable under the trade name SOPROPHOR FL from Rhone-Poulenc; or the like.

The further substance may instead or also comprise one or more of the following:

- an absorptive carrier such as a colloidal silica, for example AEROSIL 200 (trade name), diatomaceous earth, or a clay such as attapulgite;
- a binder such as a microfine cellulose, for example that obtainable under the trade name

ELCEMA P100 from Degussa, and which also acts as a filler and disintegrating agent; or lactose monohydrate, for example that obtainable under the trade name LUDIPRESS from BASF, and which is also a direct tabletting auxiliary;

- a lubricant such as magnesium stearate;
- a flow improving agent such as an absorptive silica, for example SIPERNAT 22S (trade name) from Degussa, which is a spray-dried ground silica, and acts as a free flow/anti-caking agent; and
- a water soluble filler such as soluble starch, urea, or sodium chloride.

The dosage device may comprise, by mass

active ingredient	-	5 - 60%
binder	-	10 - 30%
flow improving agent	-	0.5 - 10%
wetting/dispersing agent	-	0.5 - 5%
disintegrating agent	-	3 - 50%
lubricant	-	0,05 - 2%
filler	-	2 - 30%

The invention also extends to a dosage device when made in accordance with the method of the first aspect of the invention.

According to a second aspect of the invention, there is provided a dosage device which is in compressed unitary form and which comprises an admixture of an active ingredient which is in solid form at 25°C and which has an average particle size of less than 5 microns, and a disintegrating agent.

According to a third aspect of the invention, there is provided a method of treating an article or locus, which comprises

introducing a dosage device according to the invention, into a predetermined volume of water, with the volume of water being such that the concentration of the active ingredient in the water is greater than the solubility limit of the active ingredient in the water;

allowing the dosage device to disintegrate, thereby to form a suspension of the active ingredient in the water; and

applying the suspension to an article or locus to be treated.

The invention will now be described by way of the following non-limiting examples (all proportions in parts by mass):

EXAMPLE 1

Step 1

A suspension concentrate having the following composition was made up by admixing the following components:

as active ingredient, thiophanate methyl, technical (93% pure) (equivalent to 40,08 parts active ingredient)	-	43.10
as dispersing agent, TAMOL NNO	-	3.50
as further dispersing agent, TERIC 200 (trade name) which is a nonyl phenol alkylene oxide condensate available from ICI Australia	-	1.50
as antifoaming agent, SILCOLAPSE 5000A (trade name) which is an antifoaming silicone and is obtainable from ICI	-	0.04
water	-	51.86

The suspension concentrate was milled by passing it twice through a horizontal bead mill (laboratory model KDL Dyno-mill) to obtain an average solids particle size of 2 microns.

Step 2

Tablets were formed having the following composition:

suspension concentrate from Step 1 (equivalent to 35.0 parts active ingredient)	87.32
as wetting agent, EMPICOL LZ	1.0
as flow improving agent, SIPERNAT 22S	0.5
as disintegrating agent, POLYPLASDONE XL	40.9
as binder, ELCEMA P100	15.4
as lubricant, magnesium stearate	0.1

To form the tablets, the following steps were followed:

(i) The suspension concentrate was absorbed into the disintegrating agent;

(ii) This mixture was granulated by passing it through a 16 mesh British Standard Sieve;

(iii) The granules were then dried in a vacuum oven at 70°C;

(iv) The remainder of the constituents were admixed with the dried granules; and

(v) The mixture was pressed into 2.5 g tablets at 1 tonne pressure.

Each tablet hence contained 0.875 gms active ingredient. In use, 1 tablet is added to 1 liter of water at ambient temperature, eg at 25°C. The

tablet disintegrates within 30 seconds to form a suspension of the active ingredient in the water. Microscopic examination of the suspension revealed that most of the suspended particles were in the 1-3 micron range.

EXAMPLE 2

Step 1

A suspension concentrate having the composition as set out hereunder was made up and milled as described in EXAMPLE 1.

hexaconazole, technical (87,3% pure) (equivalent to 20 parts active ingredient)	-	22.9
TERIC 200	-	2.6
SILCOLAPSE 5000A	-	0.1
water	-	74.4

Step 2

Tablets were made up to have the composition as set out hereunder, following the procedure of EXAMPLE 1 save that the dispersing agent was added to the milled suspension concentrate before it was absorbed into the disintegrating agent.

suspension concentrate from Step 1 (equivalent to 30 parts active ingredient)	-	150.0
POLYPLASDONE XL (disintegrating agent)	-	58.4
SOPROPHOR FL (dispersing agent)	-	3.1
magnesium stearate	-	0.1

When the resulting tablets were dispersed in standard hard water containing 340 ppm calcium as calcium carbonate, to a concentration of 0.005% active ingredient, a suspensibility of 88% after 30 minutes was obtained.

EXAMPLE 3

Step 1

A suspension concentrate having a composition as set out hereunder was made up and milled as described in EXAMPLE 1.

deltamethrin, technical (98% pure) (equivalent to 64 parts active ingredient)	65.31
TERIC 200	2.73
SILCOLAPSE 5000A	0.08
water	31.47

Step 2

Tablets were made up to have the composition as set out hereunder, following the procedure of EXAMPLE 1, except that the mixture was not granulated but was pressed directly, after drying, using a commercial MANESTY press.

suspension concentrate from Step 1 (equivalent to 40 parts active ingredient)	62.50
KOLLIDON CL (disintegrating agent)	24.25
ELCEMA P100 (binder)	30.60
EMPICOL LZ (wetter)	2.[illegible]
SIPERNAT 22 S (flow improving agent)	0.50
magnesium stearate	0.10

The tablets were evaluated by dispersing them in water at the rate of 1 tablet of 2.5 gm (1 gm active ingredient) in 5 liters of water, ie 0.2 gm active ingredient per liter of water. The solubility of deltamethrin in water at 20°C is less than 2 micrograms/liter of water. The tablets disintegrated within 30 seconds.

Suspensibility tests indicated that 65% of the active ingredient was still in suspension after 30 minuts in standard hard water containing 340 ppm calcium as calcium carbonate.

EXAMPLE 4

Step 1

A suspension concentrate having a composition as set out hereunder was made up and milled as described in EXAMPLE 1:

atrazine, technical (98% pure)	-	50.0
TERIC 200	-	2.5
SILCOLAPSE 5000A	-	0.1
water	-	47.4

Step 2

Tablets were made up to have the composition as set out hereunder, following the procedure set out in EXAMPLE 1:

suspension concentrate from Step 1 (equivalent to 36.45 parts active ingredient)	-	74.4
KOLLIDON CL	-	27.9
ELCEMA P100	-	28.4
EMPICOL LZ	-	4.0
SIPERNAT 22S	-	0.5
magnesium stearate	-	0.1

20 mm diameter tablets weighing 3.5 gms each were pressed at 5 MPa, using a laboratory precision press. The resulting tablets were evaluated for hardness by subjecting single tablets to an increasing mass applied edge-on. The tablets withstood pressures up to 5 kg. Disintegration time was measured by dropping a single tablet into approximately 2 litres of water at ambient

temperature, and agitating lightly. The test tablet disintegrated completely within 30 seconds. Microscopic examination of the resultant suspension revealed that most particles were in the 1-3 micron range.

EXAMPLE 5

2,5 gm tablets were made up by homogeneously admixing the components as set out hereunder, and pressing them into the tablets at 1.5 tonnes pressure.

deltamethrin, technical	-	40.0
EMPICOL LZ (wetting agent)	-	2.0
TAMOL NNO (dispersing agent)	-	5.0
KOLLIDON CL (disintegrating agent)	-	25.0
ELCEMA P100 (binder)	-	27.4
SIPERNAT 22S (flow improving agent)	-	0.5
magnesium stearate (lubricant)	-	0.1

The deltamethrin is dry milled prior to the admixing thereof to have an average particle size less than 5 microns. For example, it may be in micronized form as available from Roussel Uclaf, with 50% (by mass) of its particles having a particle size less than 2 microns. Alternatively, the suspension concentrate of Example 3 can be spray dried, and the resultant powder admixed with the remaining ingredients as set out above.

The tablets, when introduced into water at the rate of one tablet in 5 liters of water, disintegrate within 1 minute. Microscopic examination revealed that most of the particles in the suspension were in the 2 micron range.

The Applicant has found that with active pesticides requiring a very low rate of application, typically in the order of a few grams per hectare, very small pesticide particles, typically having an average particle size less than 5 microns, dispersed in the prescribed carrier liquid, usually water, are highly desirable for effective, accurate and even distribution of the pesticide on application. Furthermore, the smaller the particle size, the greater the surface area thereof, which promotes effective release of the pesticide after application to a locus or substrate. However, it has hitherto been a problem when providing pesticides in tablet form, that if the pesticide particles are too small, unsatisfactory dispersion rates of the tablet in the carrier liquid result. However, it has surprisingly been found that in the method and dosage device of the present invention in which the average pesticide particle size is less than 5 microns, and typically in the order of 1 to 3 microns, rapid disintegration and dispersion rates are achieved. Moreover, the resulting tablets have adequate hardness which permit handling in the field, and the tablets on dispersion have excellent suspension properties.

The dosage devices of the present invention thus provide a good vehicle for such pesticides, since they are compact and hence easily transported and stored, and are also in a form in which they are handled safely. Furthermore, there is not a problem of having to dispose of large used pesticide containers. They are furthermore easy to disperse and apply effectively and accurately as set out hereinbefore, ie with little wastage.

Furthermore, the Applicant was also surprised to find that the resulting average pesticide particle size of the active ingredient, after the tablets had been dispersed in water, was of the desired order of 1-3 microns in spite of the fact that the active ingredient, after having been milled down to less than 5 microns, was then compressed with the disintegrating agent into tablet form during which agglomeration into larger particle sizes would have been expected. However, as stated, it was surprisingly found that the average particle size of the active ingredient, in the resultant suspensions, was still in the range of 1-3 microns.

CLAIMS: The claims defining the invention are as follows:-

1. A method of making a dosage device, which method comprises

admixing at least one pesticide as the active ingredient in solid form at 25°C and which has an average particle size of less than 5 microns, with a disintegrating agent, to provide a compressible mix; and

compressing the mix into a unitary dosage device.

2. A method according to Claim 1, which comprises comminuting the active ingredient in solid form and having an average particle size greater than 5 microns, to have an average particle size of less than 5 microns.

3. A method according to Claim 2, wherein the active ingredient is comminuted to have an average particle size of 1-3 microns.

4. A method according to Claim 2 or Claim 3, wherein the comminution is effected by dry milling the active ingredient to the desired particle size.

5. A method according to Claim 2 or Claim 3, wherein the comminution is effected by forming a suspension concentrate of the active ingredient in a suitable carrier liquid in which the active ingredient is non-soluble or sparingly soluble, and wet milling the suspension concentrate to obtain the desired active ingredient particle size.

AUSTRALIAN PATENT OFFICE LS

6. A method according to Claim 5, wherein the active ingredient is sparingly soluble in water, with water hence being the carrier liquid for use in forming the suspension concentrate and the concentration of the active ingredient in the water, in the suspension concentrate, thus being greater than the solubility limit of the active ingredient in water.

7. A method according to Claim 5 or Claim 6, wherein the milled suspension concentrate is absorbed into the disintegrating agent to effect the admixing thereof with the disintegrating agent.

8. A method according to claim 5 or Claim 6, wherein the milled suspension concentrate is spray dried, and then admixed with the disintegrating agent.

9. A method according to any one of Claims 2 to 8 inclusive, wherein the active ingredient is a pesticide and the disintegrating agent is capable of disintegrating by effervescing.

10. A method acording to any one of Claims 2 to 8 inclusive, wherein the active ingredient is a pesticide and the disintegrating agent is capable of disintegrating by swelling on contact with water.

11. A method according to any one of Claims 2 to 10 inclusive, which includes admixing at least one further substance with the active ingredient, before comminution thereof and/or after comminution thereof.

12. A method according to Claim 11, wherein the further substance comprises one or more of a surfactant, an absorptive carrier, a binder, a lubricant, a flow improving agent, and a filler.

13. A dosage device when made in accordance with the method as claimed in any one of Claims 1 to 12 inclusive.

14. A dosage device which is in compressed unitary form and which comprises an admixture of a pesticide which is in solid form at 25°C, is sparingly soluble in water, and has an average particle size of less than 5 microns, and a disintegrating agent, with the device being capable of disintegrating in water to form a suspension of the pesticide in the water, when introduced into sufficient water such that the concentration of the pesticide in the water is above the solubility limit of the pesticide in water.

15. A dosage device according to Claim 14, wherein the pesticide has a water solubility of less than 1000 mg/l at 25°C, and the average particle size of the active ingredient is from 1 to 3 microns.

16. A dosage device according to Claim 15, wherein the pesticide is selected from the group consisting in an insecticide, herbicide, fungicide and an acaricide, with the disintegrating agent being capable of disintegrating the device by effervescing or swelling on contact with water.

AUSTRALIAN PATENT OFFICE LS

17. A dosage device according to Claim 16, wherein the pesticide is selected from the group consisting in atrazine, simazine, cyanazine, terbuthylazine, diuron, chlorsulphuron, metsulfuron, tralkoxydim, 2-(2-chloro-4-mesylbenzoyl) cyclohexane-1,3-dione, deltamethrin, lindane, carbaryl, endosulfan, carbofuran, thiophanate methyl, carbendazim, flutriafol, hexaconazole, chlorothalonil, copper oxychloride, captan, thiram, hexythiazox, cyhexatin and amitraz, with the disintegrating agent being selected from the group consisting in a cross-linked polyvinyl pyrrolidone, a modified cellulose gum, a sodium starch glycolate, and an acid/base combination.

18. A dosage device according to Claim 14 which is that obtained by forming a suspension concentrate of the pesticide in a suitable carrier liquid in which the pesticide is non-soluble or sparingly soluble, wet milling the suspension concentrate to obtain the pesticide average particle size of less than 5 microns, admixing the suspension concentrate with the disintegrating agent by absorbing it into the disintegrating agent, and compressing the resultant mix into the unitary form.

19. A dosage device according to Claim 14, which comprises as pesticide,

atrazine, simazine, cyanazine, terbuthylazine, diuron, chlorsulphuron, metsulfuron, tralkoxydim, 2-(2-chloro-4-mesylbenzoyl) cyclohexane-1,3-dione, deltamethrin, lindane, carbaryl, endosulfan, carbofuran, thiophanate methyl, carbendazim, flutriafol, hexaconazole, chlorothalonil, copper oxychloride, captan, thiram, hexythiazox, cyhexatin or amitraz - 5-60% (by mass)

AUSTRALIAN PATENT OFFICE

as disintegrating agent,
cross-linked polyvinyl pyrrolidone, a modified cellulose gum, a sodium starch glycolate, or an acid/base combination - 3-50% (by mass)

as binder,
a microfined cellulose or lactose monohydrate - 10-30% (by mass)

as flow improving agent,
an absorptive silica - 0.5-10% (by mass)

as wetting/dispersing agent,
sodium lignosulphonate, sodium naphthalene sulphonic acid/formaldehyde condensate, sodium alkyl aryl sulphonate, a nonyl phenol alkylene oxide, a sodium lauryl sulphate, a sodium diisopropyl naphthalene sulfonate, a sodium salt of naphthalene sulfonic acid formaldehyde condensate, or oxyethylated polyarylphenol phosphate - 0.5-5% (by mass)

as lubricant,
magnesium stearate - 0.05-2% (by mass)

as filler,
soluble starch, urea or sodium chloride - 2-30% (by mass)



20. A dosage device which is in compressed unitary form and which comprises an admixture of a pesticide which is in solid form at 25°C, is sparingly soluble in water, and has an average particle size of 1 to 3 microns, and a disintegrating agent, with the device being capable of disintegrating in water to form a suspension of the pesticide in the water, when introduced into sufficient water such that the concentration of the pesticide in the water is above the solubility limit of the pesticide in water, with the average particle size of the pesticide in the suspension being in the range of 1 to 3 microns.

21. A method of treating an article or locus, which comprises

introducing a dosage device as claimed in any one of Claims 13 to 20 inclusive, into a predetermined volume of water, with the volume of water being such that the concentration of the active ingredient in the water is greater than the solubility limit of the active ingredient in the water;

allowing the dosage device to disintegrate, thereby to form a suspension of the active ingredient in the water; and

applying the suspension to an article or locus to be treated.

22. A novel method of making a dosage device, substantially as described and exemplified herein.

AUSTRALIAN PATENT OFFICE LS

23. A novel dosage device, substantially as described and exemplified herein.

24. A novel method of treating an article, substantially as described and exemplified herein.

DATED this 20th day of May 1993

AECI LIMITED
By its Patent Attorney

.....
David Gibson

AUSTRALIAN PATENT OFFICE LS