

(12) PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 199528043 B2**
(10) Patent No. **707366**

(54) Title
Long-acting oxytetracycline composition

(51)⁶ International Patent Classification(s)
A61K 031/65 A61K 047/10
A61K 047/00 A61K 047/22
A61K 047/02

(21) Application No: **199528043**

(22) Application Date: **1995.07.05**

(87) WIPO No: **WO96/01634**

(30) Priority Data

(31) Number	(32) Date	(33) Country
9413873	1994.07.09	GB

(43) Publication Date : **1996.02.09**

(43) Publication Journal Date : **1996.03.21**

(44) Accepted Journal Date : **1999.07.08**

(71) Applicant(s)
Norbrook Laboratories Limited

(72) Inventor(s)
Alan Patterson; Drew Holmes

(74) Agent/Attorney
DAVIES COLLISON CAVE, 1 Little Collins Street, MELBOURNE VIC 3000

OPI DATE 09/02/96 APPLN. ID 28043/95
AOJP DATE 21/03/96 PCT NUMBER PCT/GB95/01583



AU9528043

PCT)

(51) International Patent Classification ⁶ : A61K 31/65, 47/02, 47/10, 47/22, 47/00		A1	(11) International Publication Number: WO 96/01634
			(43) International Publication Date: 25 January 1996 (25.01.96)
(21) International Application Number: PCT/GB95/01583 (22) International Filing Date: 5 July 1995 (05.07.95) (30) Priority Data: 9413873.2 9 July 1994 (09.07.94) GB (71) Applicant (for all designated States except US): NORBROOK LABORATORIES LIMITED [GB/GB]; Station Works, Camlough Road, Newry BT35 6JP (GB). (72) Inventors; and (75) Inventors/Applicants (for US only): PATTERSON, Alan [GB/GB]; 60 Quarry Road, Belfast BT4 2NQ (GB). HOLMES, Drew [GB/GB]; 9 Village Manor, Bryansford, Newcastle, Co Down BT33 0SL (GB). (74) Agent: FITZPATRICKS; 4 West Regent Street, Glasgow G2 1RS (GB).			(81) Designated States: AU, CA, CN, GB, JP, MX, NZ, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: LONG-ACTING OXYTETRACYCLINE COMPOSITION			
(57) Abstract An injectable composition of higher residual effect with reduced detrimental effects such as pain at injection site, swelling, tissue irritancy or necrosis and containing as active principle a tetracycline compound, either as the free base or a salt thereof with a physiologically acceptable acid, complexed with a substantially equimolar amount of a magnesium compound, is solubilised in a water miscible solvent system comprising, either (i) a) glycerol formal in an amount of from about 10 to about 50 % v/v; with b) polyethylene glycol in an amount of from about 1 to 15 % v/v; or (ii) from about 25 to about 75 % v/v of N-methylpyrrolidone, said composition optionally containing a pH modifier in an amount sufficient to maintain a physiochemically acceptable pH, the balance being made up with water q.s.			

Long-Acting Oxytetracycline Composition

This invention relates to injectable formulations containing tetracyclines, particularly oxytetracycline, which exhibit higher residual effect with less of the known detrimental effects such as pain at injection site, swelling, tissue irritancy or necrosis.

Preparation of pharmaceutical compositions containing tetracyclines and oxytetracycline in particular has always presented a challenge due to aqueous solubility constraints which firstly have impact upon composition stability, and secondly upon parenteral administration.

Prior art oxytetracycline compositions have exhibited relatively high viscosity at low temperatures which makes injection difficult, have shown poor stability and suffered limitations on strength of active principle. Thus considerable research has gone into determining suitable complexing agents and more favourable co-solvents to address these shortcomings. A review of the art suggests that presence of calcium, and especially magnesium in the formulation now appears mandatory as a complexing agent and whereas some improvements have been made in stability and delivery by adopting various co-solvent systems, higher concentration loadings and residual effect remain areas in which improvements are needed. This is especially of interest for veterinary purposes where the need is to deliver high effective doses with minimum effort in animal handling and detrimental effect on the animal requiring treatment.

At the current time prior art so-called "long-acting" oxytetracycline formulations typically contain 200 mg/ml oxytetracycline and are administered at 20 mg/kg body weight, having activity as determined by residual blood levels of oxytetracycline detectable for up to about four days or so.

An object of this invention is to provide a composition of substantially greater long acting effect whilst minimising to the greatest extent possible the defects observed in previously proposed formulations. In particular the invention to be particularly described hereinbelow

provides for administration of an oxtetracycline formulation at dose rates of from 10 to 40 mg/kilogram bodyweight, giving at 30 mg/kg in animals an extended duration of effective plasma levels against susceptible organisms in excess of 9 days which is a surprising achievement in the light of the known prior art.

Solubility of oxytetracycline in non-aqueous solvents was considered by Eugene Gans and Takeru Higuchi, Journal of the American Pharmaceutical Association, 1957, Vol XLVI, pp 587-591.

The patent literature in this area is extensive and one could refer to the following patents which are illustrative of the decades of research carried out on formulation of tetracycline compositions:

GB-A-894 619, GB-A-1 131 007, GB-A-1 250 304, GB-A-1 286 351, GB-A-1 427 882, GB-A-1 494 558, GB-A-1 508 601, GB-A-1 514 838, GB-A-1 520 197, GB-A-1 538 903 GB-A-1 563 478, GB-A-1 592 053, GB-B-2 047 097; EP-B-38 103, EP-B-96 942; US-A-2 516 080, US-A-2 980 584, US-A-2 990 331, US-A-3 062 717, US-A-3 219 529, US-A-3 557 280, US-A-3 712 949, US-A-3 957 972, US-A-4 011 313, US-A-4 018 889, US-A-4 020 162, US-A-4 126 680, US-A-4 386 083, US-A-4 399 127, US-A-4 772 460, US-A-4 957 972, and US-A-5 075 295.

From these documents it is apparent that a variety of water-dispersible complex-stabilisers or water-miscible co-solvents have been proposed including 2-pyrrolidone, polyvinyl pyrrolidone, polyethylene glycols, caprolactam, 2-piperidone, and glycerol formal (a reaction product of glycerol and formaldehyde) in specific formulations. However it is by no means clear that the said co-solvents are equally interchangeable nor can the effect of such a change be entirely predictable for a given formulation.

US-A-4 386 083 proposes use of glycerol formal in conjunction with magnesium acetate and magnesium chloride, whilst US-A-4 772 460 proposes use of N-methylpyrrolidone (1-methyl-2-pyrrolidone) and a soluble magnesium compound.

US-A-5 075 295 is particularly directed to a composition aiming to achieve up to 30% oxytetracycline, which contains polyethylene glycol 400 and magnesium oxide, but examples given only appear to show a capability of achieving up to 25% oxytetracycline and there is to applicant's knowledge no current commercially available product capable of achieving greater than 20%.

Accordingly this invention provides a composition containing as active principle a tetracycline compound, either as the free base or a salt thereof with a physiologically acceptable acid, complexed with a substantially equimolar amount of a magnesium compound, solubilised in a water miscible solvent system comprising, either

- (i) a) glycerol formal in an amount of from about 10 to about 50% v/v; with
- b) polyethylene glycol in an amount of from about 1 to 15% v/v.

said composition optionally containing a pH modifier in an amount sufficient to maintain a physiochemically acceptable pH, the balance being made up with water q.s.

The composition optionally contains a thickener such as polyvinyl pyrrolidone in an amount of up to 10% w/v, and may contain usual formulation aids or auxiliaries typically used for such formulations. Thus the composition may contain antioxidants, e.g. sodium formaldehyde sulfoxylate and pH adjusting agents e.g. monoethanolamine, to provide a preferred pH range of from about 7.5 to about 9.5, more preferably from about 8.5 to about 9.0.

Preferably the composition contains a magnesium compound such as magnesium oxide or a salt e.g magnesium chloride.

The preferred compositions contain oxytetracycline as the base or its hydrochloride in an amount of from about 15 to about 35% w/v, complexed with an equimolar ratio of a magnesium compound, preferably a salt, solubilised in a solvent system comprising polyethylene glycol in an amount



of from about 1 to about 15% v/v and glycerol formal in an amount of from about 10 to about 50% v/v. In particular the most preferred composition contains about 30% w/v oxytetracycline, about 40% glycerol formal, about 10% v/v polyethylene glycol with a magnesium-containing complexing agent or stabiliser, antioxidant and water making up the balance. In that composition magnesium oxide is suitably present in an amount of about 2.7% w/v and, as antioxidant, sodium formaldehyde sulfoxylate in an amount of about 0.4% w/v may be used. Thus according to the present invention there is provided a formulation capable of providing from about 10 to about 40 mg/kg bodyweight consisting of:

Oxytetracycline	300 mg
Magnesium oxide	27 mg
Sodium formaldehyde sulfoxylate	4 mg
Glycerol formal	0.4 ml
Polyethylene glycol	0.1 ml
Monoethanolamine	q.s. pH 8.6 to 8.8
Water for injections	<u>to</u> 1 ml

The invention will now be further described by way of example for the purposes of practical illustration only.

An oxytetracycline formulation was prepared according to the procedure indicated below using the following components:

Active Ingredient -

Oxytetracycline	30% w/v
-----------------	---------

Excipients -

Magnesium oxide	2.7% w/v
Sodium formaldehyde sulfoxylate	0.4% w/v
Glycerol formal	40% w/v
Polyethylene glycol	10% w/v
Monoethanolamine	q.s. pH 8.6 to 8.8
Water for injections	<u>to</u> 100% w/v

A controlled environment having an inert atmosphere was provided within which suitable mixing and temperature controllable heating apparatus was assembled. A nitrogen blanket is considered suitable for this purpose. The above

components of the proposed composition were mixed by initially mixing a proportion of the total water with the selected solvents. The sodium formaldehyde sulfoxylate, magnesium oxide and oxytetra-cycline were added sequentially whilst mixing continuously and maintaining a temperature of approximately 65°C until all the constituents have dissolved. Thereafter, the composition is cooled to below 30°C and the pH is adjusted to lie within the range of 8.0 to 9.0, in this case by adding a sufficient amount of mono-ethanolamine. Finally the volume is made up with water, the pH checked and adjusted if necessary, and the composition is filtered through a 0.2 µm filter and filled into appropriate containers.

15 In alternative embodiments, where use of a thickener such as polyvinyl pyrrolidone is called for then it should preferably be added after the sodium formaldehyde sulfoxylate.

Throughout this specification and the claims which follow, unless the context requires otherwise, the word "comprise", and variations such as "comprises" and "comprising", will be understood to imply the inclusion of a stated integer or step or group of integers or steps but not the exclusion of any other integer or step or group of integers or steps.

The following Tables provide details of Examples 1 to 14 each of which provided compositions showing excellent stability and which achieved the desired dosage levels and long acting effect.



TABLE 1:

INGREDIENTS	EXAMPLE									
	1	2	3	4	5	6	7	8	9	10
Oxytetracycline (% w/v)	30.0	30.0	30.0	30.0	30.0	30.0	30.0	30.0	15.0	35.0
Magnesium Oxide (% w/v)	2.7	2.7	2.7	2.7	2.7	2.7	2.7	13.25*	1.3	3.06
Sodium Formaldehyde Sulphoxylate (% w/v)	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.5	0.4	0.4
Glycerol Formal (% v/v)	30.0	30.0	30.0	35.0	35.0	35.0	40.0	40.0	40.0	40.0
Polyethylene Glycol 200 (% v/v)	10.0	15.0	20.0	10.0	15.0	20.0	10.0	10	10.0	10.0
Polyvinyl Pyrrolidone K12 (% w/v)			3.0							
Water to (% v/v)	100	100	100	100	100	100	100	100	100	100

*Magnesium Chloride

TABLE 2:

INGREDIENTS	EXAMPLE			
	11	12	13	14
Oxytetracycline (% w/v)	30	30	25	35
Magnesium Oxide (% w/v)	2.78	2.78	2.3	3.21
N-Methyl Pyrrolidone (% v/v)	30.0	60.0	60.0	60.0
Sodium Formaldehyde Sulphoxylate (% w/v)	0.4	0.4	0.40	0.4
Water to (% v/v)	100	100	100	100

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS

1. A composition containing as active principle a tetracycline compound, either as the free base or a salt thereof with a physiologically acceptable acid, complexed with a substantially equimolar amount of a magnesium compound, solubilised in a water miscible solvent system comprising,

a) glycerol formal in an amount of from about 10 to about 50% v/v; with

b) polyethylene glycol in an amount of from about 1 to 15% v/v;

said composition optionally containing a pH modifier in an amount sufficient to maintain a physiochemically acceptable pH, the balance being made up with water q.s.

2. A composition according to claim 1 comprising as a thickener polyvinyl pyrrolidone in an amount of up to about 10% w/v

3. A composition according to claim 1 or claim 2 wherein the magnesium compound is magnesium oxide.

4. A composition according to claim 1 or claim 2 wherein the magnesium compound is a magnesium salt.

5. A composition according to claim 4 wherein the magnesium salt is magnesium chloride.

6. A composition according to claim 1 wherein the tetracycline compound is oxytetracycline base or its hydrochloride in an amount of from about 15 to about 35% w/v.



7. A composition according to claim 1 wherein the composition contains about 30% w/v oxytetracycline, about 40% glycerol formal, about 10% v/v poly-ethylene glycol with a magnesium-containing complexing agent or stabiliser, antioxidant and water making up the balance.

8. A composition according to claim 7 wherein magnesium oxide is present in an amount of about 2.7% w/v and, as antioxidant, sodium formaldehyde sulfoxylate in an amount of about 0.4% w/v may be used.

9. An injectable composition for treatment of animals which consists of:

Oxytetracycline	300 mg
Magnesium oxide	27 mg
Sodium formaldehyde sulfoxylate	4 mg
Glycerol formal	0.4 ml
Polyethylene glycol	0.1 ml
Monoethanolamine	q.s. pH 8.6 to 8.8
Water for injections	<u>to</u> 1 ml.

10. A method for administering oxytetracycline to an animal in need thereof comprising the administration of a formulation according to any one of claims 6 to 9 at a dosage rate of from about 10 to about 40 mg of oxytetracycline per kilogram of bodyweight.

11. Use of oxytetracycline in the manufacture of a composition according to any one of claims 6 to 9 wherein said composition is adapted for administration of from about 10 to about 40 mg of oxytetracycline per kilogram of bodyweight.

12. An injectable composition for treatment of animals according to any one of the Examples 1 to 10 hereinbefore.

DATED this 23rd day of April 1999

Norbrook Laboratories Limited.

By its Patent Attorneys

DAVIES COLLISON CAVE

