



(86) Date de dépôt PCT/PCT Filing Date: 2003/12/18
(87) Date publication PCT/PCT Publication Date: 2004/07/15
(45) Date de délivrance/Issue Date: 2012/04/17
(85) Entrée phase nationale/National Entry: 2005/05/19
(86) N° demande PCT/PCT Application No.: US 2003/040937
(87) N° publication PCT/PCT Publication No.: 2004/058178
(30) Priorité/Priority: 2002/12/23 (US60/436,274)

(51) Cl.Int./Int.Cl. *C07K 16/28* (2006.01),
A61K 38/17 (2006.01), *A61K 38/20* (2006.01),
A61K 39/395 (2006.01), *A61K 48/00* (2006.01),
A61P 17/02 (2006.01), *C07K 16/24* (2006.01)

(72) Inventeurs/Inventors:
BOWMAN, EDWARD P., US;
CHEN, SHI-JUAN, US;
CUA, DANIEL J., US;
MOORE, KEVIN W., US;
CHURAKOVA, TATYANA, US;
NGUYEN, HONG-NHUNG Y., US;
CHAN, JASON R., US

(73) Propriétaire/Owner:

(54) Titre : IL-23 DESTINE AU TRAITEMENT DES LESIONS CUTANEEES
(54) Title: IL-23 FOR TREATING CUTANEOUS WOUNDS

(57) **Abrégé/Abstract:**

It is provided methods of treatment for skin disorders. In particular, treatment, the skin disorders are generally inflammatory skin disorders, including improper wound healing. It is further provided the use of an IL-23 complex for treating a cutaneous wound, the IL-23 complex comprising a polypeptide comprising residues 352 to 521 of SEQ ID NO: 8 or a conservatively modified variant thereof comprising an individual amino acid substitution; and a polypeptide comprising residues 25 to 330 of SEQ ID NO: 8 or a conservatively modified variant thereof comprising an individual amino acid substitution. It is also provided the use of an IL-23 fusion protein for treating a cutaneous wound, the IL-23 fusion protein comprising residues 352 to 521 of SEQ ID NO: 8 and residues 25 to 330 of SEQ ID NO: 8.

(73) **Propriétaires(suite)/Owners(continued):**SCHERING CORPORATION, US

(74) **Agent:** NORTON ROSE CANADA S.E.N.C.R.L.,S.R.L./LLP

ABSTRACT :

It is provided methods of treatment for skin disorders. In particular, treatment, the skin disorders are generally inflammatory skin disorders, including improper wound healing. It is further provided the use of an IL-23 complex for treating a cutaneous wound, the IL-23 complex comprising a polypeptide comprising residues 352 to 521 of SEQ ID NO: 8 or a conservatively modified variant thereof comprising an individual amino acid substitution; and a polypeptide comprising residues 25 to 330 of SEQ ID NO: 8 or a conservatively modified variant thereof comprising an individual amino acid substitution. It is also provided the use of an IL-23 fusion protein for treating a cutaneous wound, the IL-23 fusion protein comprising residues 352 to 521 of SEQ ID NO: 8 and residues 25 to 330 of SEQ ID NO: 8.

IL-23 FOR TREATING CUTANEOUS WOUNDS

5

FIELD OF THE INVENTION

The present invention relates generally to uses of mammalian cytokine-like molecules and related reagents. More specifically, the invention relates to identification of mammalian cytokine-like proteins and inhibitors thereof that modulate skin or wound healing, e.g., inflammatory skin
10 conditions.

BACKGROUND OF THE INVENTION

Cytokines are small proteins that mediate signaling and communication between cells of the immune system, e.g., T cells, B cells, dendritic cells, and macrophages. These proteins mediate a
15 number of cellular activities, including proliferation, growth, differentiation, migration, cell activation, and response to infection, foreign antigens, and wounds.

A particularly important family of cytokines is interleukin-6 (IL-6) family. These cytokines exhibit a wide range of often overlapping biological functions that are transmitted via multichain cell surface receptors, which are typically formed by high-affinity, cytokine-specific receptor chains and
20 lower-affinity, signal-transducing chains. Receptor subunits are often shared among members of this cytokine subfamily.

Recently, a novel helical cytokine was identified that has structural homology to the IL-6 family of cytokines. This protein was designated p19, and was shown to be part of a novel composite factor consisting of a disulfide-bridged complex between p19 and the p40 subunit of IL-12. This
25 novel p19p40 complex, also known as IL-23, is naturally expressed by activated mouse and human dendritic cells and has biological activities that are similar to but distinct from IL-12 (see, e.g., Oppmann *et al.* (2000) *Immunity* 13:715-725). The p19 subunit of IL-23 is also known as "IL-23p19."

The present invention identifies and provides IL-23, IL-23 agonists, and variants and
30 derivatives thereof, as modulators of skin disorders, for example, for use in the treatment or diagnosis of skin conditions and disorders or wound healing, see, e.g., Fitzpatrick, *et al.* (eds.) (1993) *Dermatology in General Medicine 4th ed.*, McGraw-Hill, NY; Bos (ed.) (1989) *Skin Immune System*, CRC Press, Boca Raton, FL; Callen (1996) *General Practice Dermatology*, Appleton and Lange, Norwalk, CN; Rook, *et al.* (eds.) (1998) *Textbook of Dermatology*, Blackwell Publ., Malden, MA;
35 Habifor and Habie (1995) *Clinical Dermatology: A Color Guide to Diagnosis and Therapy*, Mosby,

Phila., PA.; Grob (ed.) (1997) *Epidemiology, Causes and Prevention of Skin Diseases*, Blackwell, Malden, MA; Hess and Salcido (2000) *Wound Care*, Springhouse Pub. Co., Springhouse, PA; Mani, et al. (1999) *Chronic Wound Healing: Clinical Measurement and Basic*, Balliere Tindall Ltd., London, UK; Wyngaarden and Smith (eds.) (1985) *Cecil's Textbook of Medicine*, W.B. Saunders Co., Phila., PA; Berkow (ed.) (1982) *The Merck Manual of Diagnosis and Therapy*, Merck Sharp & Dohme Research Laboratories, West Point, PA; Braunwald, et al. (eds.) (1991) *Harrison's Principles of Internal Medicine, 12th Ed.*, McGraw-Hill, Inc., NY,

The present invention provides methods and reagents for the treatment, prevention, and diagnosis of wounds and wound healing, e.g., burns, wounds of cartilage, nerves and spinal cord, muscle, soft tissues, blood vessels and angiogenesis, ulcers and pressure sores, bone fractures and osteoporosis, and for promoting skin growth, e.g., at harvested or donor sites used in skin grafting (see, e.g., Yamaguchi and Yoshikawa (2001) *J. Dermatol.* 28:521-534; Cairns, et al. (1993) *Arch. Surg.* 128:1246-1252; Hom, et al. (2002) *Facial Plast. Surg.* 18:41-52; Hackam and Ford (2002) *Surg. Infect. (Larchmt.)* 3(Suppl. 1):S23-S35; Oshima, et al. (2002) *Hum. Cell.* 15:118-128; Lal, et al. (2000) *Growth Horm. IGF Res.* 10 (Suppl. B):S39-S43; Rose and Herndon (1997) *Burns* 23:S19-S26; Schryvers, et al. (2000) *Arch. Phys. Med. Rehabil.* 81:1556-1562; Hidaka, et al. (2002) *Orthop. Clin. North Am.* 33:439-446; Dagum (1998) *J. Hand Ther.* 11:111-117; Coutts, et al. (2001) *Clin. Orthop.* 391 (Suppl.):S271-S279; Larsson (2002) *Scand. J. Surg.* 91:140-146; Goldstein (2000) *Clin. Orthop.* 379 (Suppl.):S113-119; Lieberman, et al. (2002) *Mol. Therapy* 6:141-147; Tuli, et al. (2003) *Arthritis Res. Ther.* 5:235-238; Li, et al. (2003) *Microsc. Res. Tech.* 60:107-114; van Hinsbergh, et al. (2001) 936:426-437; Conway, et al. (2001) *Cardiovasc. Res.* 49:507-521).

Skin wound healing involves a number of phases: inflammation, first with neutrophil and later monocyte/macrophage inflammation, new tissue formation, including matrix formation and differentiation of a neoepithelium, and finally remodeling and maturation. The initial inflammatory phase allows clot formation, controls infection, and promotes vascularization, and produces growth factors. If not controlled properly, the inflammation can lead to pathological healing, e.g., ulcers or scars.

Fibroblasts deposit provisional matrix or granulation tissue, while the newly formed provisional matrix is later degraded in a tissue remodeling process. Degradation of extracellular matrix is mediated by proteases, such as matrix metalloproteases (MMP), gelatinase, and collagenase, as well as protease inhibitors. An imbalance in matrix formation and degradation leads, at one extreme, to chronic ulcers and, on the other extreme, to fibrosis. For example, keloids, an "overhealed response," are fibrous tissue outgrowths (Michalik, et al. (2001) *J. Cell Biol.* 154:799-814; Okada, et al. (1997) *J. Cell Biol.* 137:67-77; Fedyk, et al. (2001) *J. Immunol.* 166:5749-5755;

Ravanti and Kahari (2000) *Int. J. Mol. Med.* 6:391-407; Peled, *et al.* (2000) *Clin. Past. Surg.* 27:489-500).

Matrix formation and reepithelialization depend on angiogenesis (Montesinos, *et al.* (1997) *J. Exp. Med.* 186:1615-1620; Malinda, *et al.* (1998) *J. Immunol.* 160:1001-1006).

5 Growth factors used in wound healing induce expression of anti-microbial factors, e.g., defensins, cathelicidins, secretory protease inhibitor, and gelatinase-associated lipocalin (from neutrophils) (Sorensen, *et al.* (2003) *J. Immunol.* 170:5583-5589).

10 In wound healing, cells such as platelets, monocyte/macrophages, T cells, and other immune cells, infiltrate the wound and produce factors that regulate growth of tissue. These factors include TGF, tumor necrosis factor (TNF), IL-1, IL-4, IL-6, oncostatin M, GRO-alpha, various angiogenic factors, and chemokines. In turn, these factors stimulate, for expression of, e.g., extracellular matrix and tissue inhibitor of metalloproteases (TIMP). (Ihn and Tamaki (2000) *J. Immunol.* 165:2149-2155; Feugate, *et al.* (2002) *J. Cell. Biol.* 156:161-172). Myofibroblasts, cells that are fibrogenic, are important for wound closure and contraction. Disease states characterized by accumulation of myofibroblasts include pulmonary fibrosis and scleroderma (Feugate, *et al.* (2002) *J. Cell Biol.* 156:161-172).

Wound healing of skin and other tissues is a complex process involving proliferation and migration of immune cells, endothelial cells, fibroblasts, stromal cells, myofibroblasts, smooth muscle cells, pericytes, and keratinocytes.

20 Parameters used to measure healing include rate of healing, breaking strength of healed wounds, degree of epithelialization, thickness of granulation tissue, and density of extracellular matrix (Matsuda, *et al.* (1998) *J. Exp. Med.* 187:297-306).

25 Ischemia or ischemia reperfusion, as occurs with traumatic injury and "muscle unloading" (chronic bedrest), results in neutrophil infiltration, where the neutrophils often produce tissue damage in excess to the damage caused by the ischemia. Consistent with this adverse effect of neutrophils on healing is that administration of cytokine antagonists, including antagonists to IL-1 or TNF, can also improve wound healing under certain conditions, even where the cytokine is ultimately required for normal repair (see, e.g., Ley (2003) *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 285:R718-R719; Graves, *et al.* (2001) *J. Immunol.* 167:5316-5320). The present invention provides methods using an IL-23 antagonist to inhibit neutrophil-induced tissue injury, e.g., after trauma, wounding, or prolonged bedrest.

35 Improper wound healing, e.g., of cutaneous wounds, can result in chronic discomfort or disfigurement, and can lead to further complications, e.g., infections or dehydration. Thus, a need exists for effective treatment, both prophylactic and curative, to alleviate the symptoms of those

conditions. Alternatively, methods of diagnosis, e.g., of abnormal or modified health of those tissues will be useful. The present invention provides both.

SUMMARY OF THE INVENTION

5 The present invention is based, in part, upon the discovery that an IL-23 fusion protein, e.g., a fusion protein comprising the p19 subunit linked to the p40 subunit, enhanced wound healing response in various mouse models.

 The invention provides a method of treating or improving healing comprising administering to a subject an effective amount of an agonist or antagonist of IL-23. Also provided is the above
10 method, wherein the agonist or antagonist comprises a polypeptide of IL-23, or a derivative or variant thereof; a binding composition derived from an antibody that specifically binds to IL-23 or to IL-23R; or a nucleic acid encoding a polypeptide of IL-23, or a derivative or variant thereof. In addition, the invention provides the above method wherein the derivative or variant comprises an IL-23 hyperkine; wherein the agonist comprises a complex of a mature sequence of SEQ ID NO:10; and a
15 mature sequence of SEQ ID NO:12; or the above method wherein the nucleic acid further comprises an expression vector.

 In another aspect, the invention provides a method of treating or improving healing comprising administering to a subject an effective amount of an agonist or antagonist of IL-23, wherein the healing is of a skin or cutaneous wound; of an ulcer or graft; or is improper healing.
20 Also provided is the above method wherein the treating or improving increases a pressure required to break a healed or healing wound; a stiffness of a healed or healing wound; a rate of healing of a wound; a granulation layer thickness of a healed or healing wound; recruitment of a cell to or towards a wound; or antimicrobial activity. In yet another aspect, the invention provides the above method wherein the cell is a CD11b⁺, MHC Class II⁺ cell; a monocyte/macrophage; a CD31⁺ endothelial cell;
25 or an immune cell. Also provided is the above method wherein the recruitment is in or towards a granulation tissue; wherein
the increased wound breaking pressure is about a 15% or about a 20% increase in wound breaking pressure; or the increased stiffness is about a 15% or about a 20% increase in stiffness. In another embodiment, the present invention provides the above method wherein the treating or improving
30 comprises increased angiogenesis; or immune surveillance; or the above method wherein the increased angiogenesis is mediated by ICAM-1 or -2; or the increased immune surveillance is mediated by dendritic cells.

 Yet another aspect of the above invention provides a method of treating or improving healing comprising administering to a subject an effective amount of an agonist or antagonist of IL-23,
35 wherein the treating or improving comprises increased expression of a nucleic acid or protein of a cytokine in addition to IL-23; a signaling molecule; an anti-microbial molecule; a protease or

protease inhibitor; or a molecule of the extracellular matrix; or the above method wherein the cytokine nucleic acid or protein is IL-17, IL-6, IL-19, GRO-alpha, or GM-CSF; or wherein wherein the nucleic acid or protein is lactoferrin; DEC-205; CD50; nitric oxide synthase; or secretory leukoprotease inhibitor; or CD40L.

5 Also provided is the above method, wherein the antagonist comprises a nucleic acid; a blocking antibody to IL-23 or to IL-23R; or a soluble receptor derived from an extracellular part of IL-23R; the above method wherein the nucleic acid comprises an anti-sense nucleic acid; or interference RNA.

10 Yet another aspect of the present invention provides an agonist of IL-23 derived from the binding site of an antibody that specifically binds to an IL-23 receptor; the above agonist that is a polyclonal antibody; a monoclonal antibody; an Fab, Fv, or F(ab')₂ fragment; humanized; a peptide mimetic; or detectably labeled. In another embodiment, the present invention provides the above agonist comprising a complex of a polypeptide of the mature sequence of SEQ ID NO:10 and a polypeptide of the mature sequence of SEQ ID NO:12; the above agonist comprising a complex of
15 two polypeptides of the mature sequence of SEQ ID NO:10 and two polypeptides of the mature sequence of SEQ ID NO:12. Moreover the invention provides the above agonist wherein contact of the agonist to a cell expressing hIL-23R and hIL-12beta1 results in an increase in proliferation of the cell. Also provided is a kit comprising the above agonist and a compartment; or instructions for use or disposal. Also provided is a nucleic acid encoding an agonist of IL-23 derived from the binding
20 site of an antibody that specifically binds to an IL-23 receptor.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

25

As used herein, including the appended claims, the singular forms of words such as "a," "an," and "the," include their corresponding plural references unless the context clearly dictates otherwise.

30

I. General.

Interleukin-23 (IL-23) is a heterodimeric cytokine composed of a novel p19 subunit (a.k.a. IL-B30) and the p40 subunit of IL-12 (Oppmann, *et al.*, *supra*). The p19 subunit was identified during a computational search for members of the IL-6 helical cytokine family characterized by their
35 unique four α -helix bundle. Genetic analysis of the family, of which oncostatin-M, IL-11, cardiotrophin-1, and leukaemia inhibitory factor are members, reveals the closest evolutionary neighbor of p19 to be the p35 subunit of IL-12. Like p35, p19 requires co-expression of p40 for

biological activity (Wiekowski, *et al.* (2001) *J. Immunol.* 166:7563-7570). The IL-23 receptor (IL-23R) comprises a novel receptor subunit (IL-23R), that binds p19, and IL-12R β 1, that binds p40 (Parham, *et al.* (2002) *J. Immunol.* 168:5699-5708). These two receptor subunits form the functional signaling complex and are expressed on CD4⁺CD45Rb^{lo} memory T cells as well as interferon-
5 gamma (IFN γ) activated bone marrow macrophages (Parham, *et al.*, *supra*).

Preliminary characterization of IL-23 suggests that it has potent effects on memory T cells from both humans and mice, as measured by proliferation and IFN γ production. Consistent with the immunostimulatory properties of IL-23, mice in which haematopoietic cells constitutively express transgenic p19 have widespread multi-organ inflammation that results in premature death
10 (Wiekowski, *et al.*, *supra*). The inflammatory disease is characterized by intense macrophage infiltration, neutrophilia, and elevated levels of proinflammatory monokines such as IL-1 and TNF, suggesting that IL-23 may also act on myeloid cells.

Recent studies analyzing the necessity of IL-12 in resistance to infectious diseases have yielded divergent results, depending on whether p35^{-/-} or p40^{-/-} mice are used. The former, which
15 specifically lack IL-12 but express IL-23, are resistant to infection, whereas the latter, unable to express both IL-12 and IL-23, are more susceptible.

Transgenic mice deficient for the p19 subunit of IL-23 (IL-23p19) were resistant to EAE, a CNS autoimmune disease mediated by TH1 cells and inflammatory macrophages, while wild-type and heterozygous p19 control mice were highly susceptible. Mice deficient in the p40 subunit of IL-
20 12 (IL-12p40 deficient mice) were also resistant to EAE, while mice deficient in the p35 subunit of IL-12 (IL-12p35 deficient mice) were highly susceptible to EAE. This is indicative of a role of IL-23 in the induction of EAE.

p19 deficient mice had a notable altered wound healing response following subcutaneous injection of an oil emulsion in mice. The p19 deficient mice were also defective in a variety of
25 mouse disease models that required monocyte/macrophage activation. Monocytes/macrophages are known to stimulate wound repair, see, e.g., Schaffer and Nanney (1996) *Intl. Rev. Cytol* 169:151-181. In particular, it is shown below that delivery of IL-23 polypeptide into mouse skin could attract CD11b⁺/Class II⁺ activated monocyte/macrophage populations.

30 II. Definitions.

“Agonist of IL-23” and “IL-23 agonist” encompasses an agonistic antibody that specifically binds to IL-23 receptor (IL-23R) and increases the signaling properties of IL-23R. Agonist of IL-23 also encompasses an agonistic antibody that specifically binds to the complex of IL-23R and IL-12R β 1.

A "blocking antibody" encompasses, e.g., an antibody that specifically binds to IL-23 and prevents or impairs signaling mediated by IL-23 and IL-23R. A blocking antibody also encompasses an antibody that specifically binds to IL-23R and prevents or impairs signaling mediated by IL-23R, or mediated by IL-23 and IL-23R.

5 "Conservatively modified variants" applies to both amino acid and nucleic acid sequences. With respect to particular nucleic acid sequences, conservatively modified variants refers to those nucleic acids which encode identical or essentially identical amino acid sequences, or where the nucleic acid does not encode an amino acid sequence, to essentially identical nucleic acid sequences. Because of the degeneracy of the genetic code, a large number of functionally identical nucleic acids
10 may encode any given protein.

As to amino acid sequences, one of skill will recognize that an individual substitution to a nucleic acid, peptide, polypeptide, or protein sequence which substitutes an amino acid or a small percentage of amino acids in the encoded sequence for a conserved amino acid is a "conservatively modified variant." Conservative substitution tables providing functionally similar amino acids are
15 well known in the art. An example of a conservative substitution is the exchange of an amino acid in one of the following groups for another amino acid of the same group (U.S. Pat. No. 5,767,063 issued to Lee, *et al.*; Kyte and Doolittle (1982) *J. Mol. Biol.* 157:105-132):

- (1) Hydrophobic: Norleucine, Ile, Val, Leu, Phe, Cys, or Met;
- (2) Neutral hydrophilic: Cys, Ser, Thr;
- 20 (3) Acidic: Asp, Glu;
- (4) Basic: Asn, Gln, His, Lys, Arg;
- (5) Residues that influence chain orientation: Gly, Pro;
- (6) Aromatic: Trp, Tyr, Phe;
- (7) Small amino acids: Gly, Ala, Ser.

25 The phrase "effective amount" means an amount sufficient to ameliorate a symptom or sign of the medical condition. Typical mammalian hosts will include mice, rats, cats, dogs, and primates, including humans. An effective amount for a particular patient may vary depending on factors such as the condition being treated, the overall health of the patient, the method route and dose of administration and the severity of side affects. When in combination, an effective amount is in ratio
30 to a combination of components and the effect is not limited to individual components alone.

An "endpoint" for assessing or diagnosing improved healing, e.g., wound healing, includes, without limitation: wound breaking pressure; stiffness, looseness; immune surveillance; angiogenesis; wound-related anti-microbial activity; inflammation, e.g., by neutrophils; degree of expression of genes or polypeptides indicative of inflammation, angiogenesis, reepithelialization,
35 anti-microbial action, and remodeling, e.g., matrix formation, matrix breakdown, or granulization.

As determined by a suitable endpoint, the present invention provides a method to improve healing by 10% or more, more generally by 15% or more, most generally by 20% or more, typically 25% or more, more typically by 30% or more, most typically by 35% or more, often by 40% or more, more often by 50% or more, most often by 60% or more, usually by 70% or more, more usually by 80% or more, most usually by 90% or more, ideally by 100% (i.e., 2-fold) or more, more ideally by 4-fold or more, and most ideally by 8-fold or more. As determined by a suitable endpoint, the present invention also provides a method to improve healing by about 10%, more generally by about 15%, most generally by about 20%, typically about 25%, more typically by about 30%, most typically by about 35%, often by about 40%, more often by about 50%, most often by about 60%, usually by about 70%, more usually by about 80%, most usually by about 90%, ideally by about 100% (i.e., 2-fold), more ideally by about 4-fold, and most ideally by about 8-fold.

“Improper wound healing” encompasses the absence or abnormally slow progression of the healing of a wound, e.g., delayed re-epithelialization. Improper wound healing can be found, e.g., in diabetic ulcers and abscesses, pressure ulcers, infected wounds, burns, advanced age, inadequate perfusion, and obesity. Improper wound healing also encompasses injuries leading to scars or to persistent infections, see, e.g., Singer and Clark (1999) *New Engl. J. Med.* 341:738-746; Calhoun, *et al.* (2002) *Adv. Skin Wound Care* 15:31-45; Rico, *et al.* (2002) *J. Surg. Res.* 102:193-197; Thomas (2001) *Cleve. Clin. J. Med.* 68:704-722; Ashcroft, *et al.* (2002) *Biogerontology* 3:337-345; Thomason (1999) *Home Care Provid.* 4:156-161; Gallagher (1997) *Ostomy. Wound Manage.* 43:18-27.

Parameters and endpoints used to assess wound healing and response to therapeutic, pharmacological, and diagnostic agents, include a number of histological, physiological, and biochemical parameters, e.g., infiltration, activation, or differentiation of neutrophils, monocytes, and macrophages, e.g., differentiation of monocytes to reparative macrophages, and appearance of new stroma, blood vessels, and nerves. Suitable parameters also include expression levels of signaling agents, e.g., transforming growth factor, interleukin-1, and insulin-like growth factor. Measures of epithelization, e.g., rate and thickness, migration of epidermal cells, granulation thickness, degradation and maturation of extracellular matrix, e.g., provisional matrix versus collagenous matrix, wound strength (breaking strength), and fibroblast proliferation rate and phenotype, are also suitable parameters. Increased granulation tissue thickness can resulting stronger healed wounds (see, e.g., Singer and Clark, *supra*, Werner and Grose (2002) *Physiol. Rev.* 83:835-870; Matsuda, *et al.* (1998) *J. Exp. Med.* 187:297-306; Wankell, *et al.* (2001) *EMBO J.* 20:5361-5372).

A composition that is “labeled” is detectable either directly or indirectly by, e.g., spectroscopic, photochemical, biochemical, metabolic, immunochemical, isotopic, or chemical methods. For example, useful labels include epitope tags, fluorettes, ^{32}P , ^{33}P , ^{35}S , ^{14}C , ^3H , ^{125}I , stable isotopes, fluorescent compounds, electron-dense reagents, substrates, or enzymes, e.g., as used in enzyme-linked immunoassays, see, e.g., Invitrogen (2002) *Catalogue*, Carlsbad, CA; Molecular

Probes (2002) *Catalogue*, Molecular Probes, Eugene, OR; Rozinov and Nolan (1998) *Chem. Biol.* 5:713-728.

Increase in "anti-microbial activity" encompasses an increase in expression, concentration, or level of a mediator of anti-microbial activity, both in the presence and absence of demonstrated reduction in biological activity, concentration, population, or number of a microbe in contact with a human or animal subject or host. The mediator of anti-microbial activity can be, e.g., an immune cell responsive to a bacterial, viral, fungal, or protozoal, or parasitic antigen, or an anti-microbial molecule, such as a defensin. "Anti-microbial activity" embraces, e.g., phagocytosis and any activity that is generally or usually associated with phagocytosis, e.g., exposure of a microbe to toxic oxygen. "Anti-microbial activity" also encompasses an change in expression, concentration, or level of a cell, gene, protein, or small molecule that is generally or usually associated with anti-microbial action, e.g., an increase in expression of a neutrophil gene. "Anti-microbial activity" is not limited to an increase in expression, i.e., it also encompasses a decrease in expression, where that decrease promotes anti-microbial activity.

"Proliferation" or "rate of proliferation" can be measured, e.g., by assessing the increase in cell number over a predetermined period or interval of time, or by the number or proportion of cells in S phase at any given point in time.

III. Agonists and Antagonists.

The present invention provides methods of using IL-23 agonists including the full length cytokine protein (SEQ ID NO: 2 or 4). Also provided is a fusion protein, also known as "IL-23 hyperkine" (SEQ ID NO: 6 or 8), comprising p19 linked to p40 with a FLAG sequence as described for IL-6 in, e.g., Oppmann, *et al.*, *supra*; Fischer, *et al.* (1997) *Nature Biotechnol.* 15:142-145; Rakemann, *et al.* (1999) *J. Biol. Chem.* 274:1257-1266; and Peters, *et al.* (1998) *J. Immunol.* 161:3575-3581, thereof. The invention also provides agonistic anti-IL-23R antibodies that are agonistic to the IL-23 receptor, e.g., antibodies that stimulate the IL-23 receptor in the absence or presence of IL-23.

Peptides of those sequences, or variants thereof, will be used to induce receptor signaling. Also contemplated are small molecules which also induce receptor signaling. Agonists of the present invention will be useful in the treatment of various inflammatory skin disorders, including but not limited to wound healing, skin disorders associated with impaired recruitment of myeloid/monocyte cells.

The invention provides IL-23 antagonists, e.g., a blocking antibody that binds to IL-23, a blocking antibody that binds to IL-23R, a soluble receptor based on the extracellular portion of IL-23R, and nucleic acids. The IL-23 antagonists of the present invention encompass nucleic acids that are anti-sense nucleic acids and RNA interference nucleic acids (see, e.g., Arenz and Schepers (2003)

Naturwissenschaften 90:345-359; Sazani and Kole (2003) *J. Clin. Invest.* 112:481-486; Pirollo, *et al.* (2003) *Pharmacol. Therapeutics* 99:55-77; Wang, *et al.* (2003) *Antisense Nucl. Acid Drug Devel.* 13:169-189).

5 III. Antibodies and Related Reagents.

Antibodies and binding compositions derived from an antigen-binding site of an antibody are provided. These include humanized antibodies, monoclonal antibodies, polyclonal antibodies, and binding fragments, such as Fab, F(ab)₂, and Fv fragments, and engineered versions thereof. The antibody or binding composition may be agonistic or antagonistic. Antibodies that simultaneously
10 bind to a ligand and receptor are contemplated. Monoclonal antibodies will usually bind with at least a K_D of about 1 mM, more usually at least about 300 μM, typically at least about 100 μM, more typically at least about 30 μM, preferably at least about 10 μM, and more preferably at least about 3 μM or better.

Monoclonal, polyclonal, and humanized antibodies can be prepared. See, e.g., Sheperd and
15 Dean (eds.) (2000) *Monoclonal Antibodies*, Oxford Univ. Press, New York, NY; Kontermann and Dubel (eds.) (2001) *Antibody Engineering*, Springer-Verlag, New York; Harlow and Lane (1988) *Antibodies A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, pp. 139-243; Carpenter, *et al.* (2000) *J. Immunol.* 165:6205; He, *et al.* (1998) *J. Immunol.* 160:1029; Tang, *et al.* (1999) *J. Biol. Chem.* 274:27371-27378).

20 Single chain antibodies, single domain antibodies, and bispecific antibodies are described, see, e.g., Malecki, *et al.* (2002) *Proc. Natl. Acad. Sci. USA* 99:213-218; Conrath, *et al.* (2001) *J. Biol. Chem.* 276:7346-7350; Desmyter, *et al.* (2001) *J. Biol. Chem.* 276:26285-26290, Kostelney, *et al.* (1992) *J. Immunol.* 148:1547-1553; U.S. Pat. Nos. 5,932, 448; 5,532,210; 6,129,914; 6,133,426; 4,946,778.

25 The invention also encompasses deamidated binding compositions, e.g., antibodies, and methods of using deamidated binding compositions (see, e.g., Zhang and Czupryn (2003) *J. Pharm. Biomed. Anal.* 30:1479-1490; Perkins, *et al.* (2000) *Pharm. Res.* 17:1110-1117; Lehrman, *et al.* (1992) *J. Protein Chem.* 11:657-663).

Antigen fragments may be joined to other materials, such as fused or covalently joined
30 polypeptides, to be used as immunogens. An antigen and its fragments may be fused or covalently linked to a variety of immunogens, such as keyhole limpet hemocyanin, bovine serum albumin, or ovalbumin (Coligan, *et al.* (1994) *Current Protocols in Immunol.*, Vol. 2, 9.3-9.4, John Wiley and Sons, New York, NY). Peptides of suitable antigenicity can be selected from the polypeptide target, using an algorithm, such as those of Parker, *et al.* (1986) *Biochemistry* 25:5425-5432; Welling, *et al.*
35 (1985) *FEBS Lett.* 188:215-218; Jameson and Wolf (1988) *Cabios* 4:181-186; or Hopp and Woods (1983) *Mol. Immunol.* 20:483-489.

Purification of antigen is not necessary for the generation of antibodies. Immunization can be performed by DNA vector immunization. See, e.g., Wang, *et al.* (1997) *Virology* 228:278-284. Alternatively, animals can be immunized with cells bearing the antigen of interest. Splenocytes can then be isolated from the immunized animals, and the splenocytes can fused with a myeloma cell line to produce a hybridoma. Resultant hybridomas can be screened for production of the desired antibody by functional assays or biological assays, that is, assays not dependent on possession of the purified antigen. Immunization with cells may prove superior for antibody generation than immunization with purified antigen (Meyaard, *et al.* (1997) *Immunity* 7:283-290; Wright, *et al.* (2000) *Immunity* 13:233-242; Preston, *et al.* (1997) *Eur. J. Immunol.* 27:1911-1918; Kaithamana, *et al.* (1999) *J. Immunol.* 163:5157-5164).

Antibody affinity, i.e., antibody to antigen binding properties can be measured, e.g., by surface plasmon resonance or enzyme linked immunosorbent assay (ELISA) (see, e.g., Maynard and Georgiou (2000) *Annu. Rev. Biomed. Eng.* 2:339-376; Karlsson, *et al.* (1991) *J. Immunol. Methods* 145:229-240; Neri, *et al.* (1997) *Nat. Biotechnol.* 15:1271-1275; Jonsson, *et al.* (1991) *Biotechniques* 11:620-627; Friguet, *et al.* (1985) *J. Immunol. Methods* 77:305-319; Hubble (1997) *Immunol. Today* 18:305-306).

Antibodies of the present invention will usually bind with at least a K_D of about 10^{-3} M, more usually at least 10^{-6} M, typically at least 10^{-7} M, more typically at least 10^{-8} M, preferably at least about 10^{-9} M, and more preferably at least 10^{-10} M, and most preferably at least 10^{-11} M (see, e.g., Presta, *et al.* (2001) *Thromb. Haemost.* 85:379-389; Yang, *et al.* (2001) *Crit. Rev. Oncol. Hematol.* 38:17-23; Carnahan, *et al.* (2003) *Clin. Cancer Res.* (Suppl.) 9:3982s-3990s; Wilchek, *et al.* (1984) *Meth. Enzymol.* 104:3-55).

Antibodies to IL-23R, where the anti-IL-23R antibody has substantially the same nucleic acid and amino acid sequence as those recited herein, but possessing substitutions that do not substantially affect the functional aspects of the nucleic acid or amino acid sequence, are within the definition of the contemplated invention. Variants with truncations, deletions, additions, and substitutions of regions which do not substantially change the biological functions of these nucleic acids and polypeptides are also within the definition of the contemplated invention.

A humanized antibody encompasses a human antibody, antibody fragment, single chain antibody, and the like, that has one or more amino acid residues introduced into it from a source which is non-human (import antibody). The amino acids used for grafting may comprise the entire variable domain of the source, one or more of the complementary determining regions (CDRs) of the source, or all six of the CDRs of the source antibody. With grafting of the import amino acids or polypeptide regions on to the host antibody, the corresponding amino acids or regions of the host antibody are generally removed. A humanized antibody will comprise substantially all of at least one, and typically two, variable domains (Fab, Fab', F(ab')₂, Fabc, Fv) in which all or substantially all

of the CDR regions correspond to those of a non-human immunoglobulin and all or substantially all of the framework regions are those of a human immunoglobulin consensus sequence. The framework regions and CDRs are highly conserved in sequence and conformation and can be accurately predicted, e.g., for use in grafting CDRs into an acceptor human antibody framework.

5 CDR regions can be grafted into a naturally occurring human acceptor framework, or in a consensus framework derived from many human antibodies. A number of human variable light (VL) and variable heavy (VH) consensus sequences have been identified. For humanization, a chain of the mouse antibody can be compared with the available human framework chains, where the human chain of closest homology is chosen for grafting (see, e.g., Maynard and Georgiou, *supra*; Li, *et al.* 10 (2002) *Immunol. Revs.* 190:53-68; Co, *et al.* (1991) *Proc. Natl. Acad. Sci. USA* 88:2869-2873; Sims, *et al.* (1993) *J. Immunol.* 151:2296-2308; Sato, *et al.* (1994) *Mol. Immunol.* 31:371-381; Morea, *et al.* (2000) *Methods* 20:267-279; Kabat *et al.* (1991) *Sequences of Proteins of Immunological Interest*, 5th ed., 4 vol., U.S. Department of Health Human Services, NIH, USA; U.S. Pat. No. 6,538,111, issued to Koike, *et al.*; U.S. Pat. No. 6,329,511, issued to Vasquez, *et al.*).

15 The humanized antibody of the present invention also encompasses substitutions, deletions, and/or insertions, using standard techniques of site-directed mutagenesis, e.g., those used for alanine scanning, see, e.g., Jin and Wells (1994) *Protein Sci.* 3:2351-2357; Cunningham and Wells (1997) *Curr. Opin. Struct. Biol.* 7:457-462; Jones, *et al.* (1998) *J. Biol. Chem.* 273:11667-11674; U.S. Pat. No. 4,816,567 issued to Cabilly, *et al.*

20 Embodiments of the present invention encompass fusion proteins, purification tags, and epitope tag, at an N-terminus, C-terminus, or positions within the polypeptide, e.g., FLAG tag and GSH-S transferase fusion protein. Amino acid changes can alter, add, or eliminate post-translational processes of the agonist anti-IL-23R antibody, e.g., sites for O- and N-glycosylation, and positions of cysteine residues used for disulfide formation, see, e.g., Wright and Morrison (1997) *Trends* 25 *Biotechnol.* 15:26-32; Kunkel, *et al.* (2000) *Biotechnol. Prog.* 16:462-470.

Binding properties of the humanized antibody can be improved by the following procedure, e.g., involving site-directed mutagenesis. Computer modeling allows visualization of which mouse framework amino acid residues are likely to interact with mouse CDRs. These "contacting" mouse framework amino acids are then superimposed on the homologous human framework. Where the 30 superimposition indicates that the mouse "contacting" framework amino acid is different from the corresponding human framework amino acid, human amino acid is changed to the corresponding mouse framework amino acid. "Contact" means interchain contact between a light chain and heavy chain, where, e.g., the amino acids are predicted to be within about 3 Angstroms of each other.

Site-directed mutagenesis can also be desirable where the amino acid of the human 35 framework is rare for that position and the corresponding amino acid in the mouse immunoglobulin is common for that position in human immunoglobulin sequences. Here, the human framework amino

acid can be mutated to the corresponding donor framework amino acid, see, e.g., U.S. Pat. No. 6,407,213, issued to Carter *et al.*; U.S. Pat. No. 6,180,370, issued to Queen, *et al.*, Jung, *et al.* (2001) *J. Mol. Biol.* 309:701-716.

5 The humanized antibody can comprise at least a portion of an immunoglobulin constant region (Fc), e.g., of a human immunoglobulin. The antibody can optionally include the CH1, hinge, CH2, CH3, and CH4 regions of the heavy chain. The humanized antibody can be selected from any class of immunoglobulins, including IgM, IgG, IgD, IgA and IgE, and any isotype, including IgG1, IgG2, IgG3 and IgG4. Standard methods can be used to improve, or remove, effector function. Effector function includes binding to FcRn, FcγR, and complement. Half-life can be improved,
10 e.g., by using human IgG2 or IgG4 subclasses or by altering residues in the hinge region (see, e.g., Clark (2000) *Immunol. Today* 21:397-402; Presta, *et al.* (2002) *Biochem. Soc. Trans.* 30:487-490; Morea, *et al.* (2000) *Methods* 20:267-279).

The CDR and framework regions of the humanized antibody need not correspond precisely to the import or host sequences, e.g., these sequences can be mutagenized by substitution, insertion or
15 deletion of at least one residue so that the residue at that site does not correspond to either the consensus or the import antibody. Such mutations, however, will not be extensive. Usually, at least 75% of the humanized antibody residues will correspond to those of the parental FR and CDR sequences, more often 90%, and most preferably greater than 95%.

Ordinarily, amino acid sequence variants of the humanized anti-IL-23R antibody will have an
20 amino acid sequence having at least 75% amino acid sequence identity with the original humanized antibody amino acid sequences of either the heavy or the light chain (e.g. as in SEQ ID NOs:2 and 4), more preferably at least 80%, more preferably at least 85%, more preferably at least 90%, and most preferably at least 95%. Identity or homology with respect to this sequence is defined herein as the percentage of amino acid residues in the candidate sequence that are identical with the humanized
25 anti-IL-23R residues, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity, and not considering any conservative substitutions as part of the sequence identity. None of N-terminal, C-terminal, or internal extensions, deletions, or insertions into the antibody sequence shall be construed as affecting sequence identity or homology.

An alternative to humanization is to use human antibody libraries displayed on phage or
30 human antibody libraries contained in transgenic mice (see, e.g., Vaughan, *et al.* (1996) *Nat. Biotechnol.* 14:309-314; Barbas (1995) *Nature Med.* 1:837-839; de Haard, *et al.* (1999) *J. Biol. Chem.* 274:18218-18230; McCafferty *et al.* (1990) *Nature* 348:552-554; Clackson *et al.* (1991) *Nature* 352:624-628; Marks *et al.* (1991) *J. Mol. Biol.* 222:581-597; Mendez, *et al.* (1997) *Nature Genet.* 15:146-156; Hoogenboom and Chames (2000) *Immunol. Today* 21:371-377; Barbas, *et al.*
35 (2001) *Phage Display: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring

Harbor, New York; Kay, *et al.* (1996) *Phage Display of Peptides and Proteins: A Laboratory Manual*, Academic Press, San Diego, CA; de Bruin, *et al.* (1999) *Nat. Biotechnol.* 17:397-399).

IV. Nucleic Acids, Vectors, and Protein Purification.

5 An "expression vector" is a nucleic acid construct, generated recombinantly or synthetically, with one or more predetermined nucleic acid elements that permit transcription of a particular nucleic acid. Typically, the expression vector includes a nucleic acid to be transcribed operably linked to a promoter.

10 The light chain and heavy chain of the agonistic anti-IL-23R antibody can be encoded by one nucleic acid, where expression of the light chain is operably linked to a first promoter, and where expression of the heavy chain is operably linked to a second promoter. Alternatively, both light and heavy chains can be encoded by one nucleic acid, where expression of both chains is operably linked to one promoter. The nucleic acid or nucleic acids encoding the light chain and the heavy chain can be provided as one or as two vectors. The methods of the present invention encompass incorporation
15 of the one or two vectors into the genome of a host cell (see, e.g., Chadd and Chamow (2001) *Curr. Opin. Biotechnol.* 12:188-194; Houdebine (2000) *Transgenic Res.* 9:305-320; Stoger, *et al.* (2002) *Curr. Opin. Biotechnol.* 13:161-166).

The nucleic acid encoding the light chain can further comprise a first vector, while the nucleic acid encoding the heavy chain can further comprise a second vector. Alternatively, one
20 vector may comprise the nucleic acids encoding the light chain and the heavy chain.

For long-term or scaled-up expression of the agonistic anti-IL-23R antibody, one vector, containing the nucleic acids encoding both the light chain and the heavy chain, can be incorporated into the host genome, e.g., where incorporation is at one point or a plurality of points in the host genome. Coexpression of the light chain and heavy chain in a host cell produces a soluble antibody.
25 The host cell can be, e.g., a mammalian, transformed or immortalized, insect, plant, yeast, or bacterial cell. The host cell may further comprise a transgenic animal. Combinations of the above embodiments are contemplated, e.g., where the light chain is simultaneously expressed by a vector that is incorporated in the host cell's genome and by a vector that is not incorporated in the genome.

Purification of an antibody, or fragments thereof, can involve ion exchange chromatography,
30 immunoprecipitation, epitope tags, affinity chromatography, high pressure liquid chromatography, and use of stabilizing agents, detergents or emulsifiers (Dennison and Lovrien (1997) *Protein Expression Purif.* 11:149-161; Murby, *et al.* (1996) *Protein Expression Purif.* 7:129-136; Ausubel, *et al.* (2001) *Curr. Protocols Mol. Biol.*, Vol. 3, John Wiley and Sons, New York, NY, pp. 17.0.1-17.23.8; Rajan, *et al.* (1998) *Protein Expression Purif.* 13:67-72; Amersham-Pharmacia (2001)
35 *Catalogue*, Amersham-Pharmacia Biotech, Inc., pp. 543-567, 605-654; Gooding and Regnier (2002) *HPLC of Biological Molecules*, 2nd ed., Marcel Dekker, NY).

V. Kits.

This invention contemplates an agonistic anti-IL23R antibody, fragments thereof, nucleic acids encoding an agonistic anti-IL-23R antibody, or fragments thereof, in a diagnostic kit.

5 Encompassed is the use of binding compositions, including antibodies or antibody fragments, for the detection of IL-23R and metabolites and breakdown products thereof, and for the detection of IL-23R-dependent activities, e.g., biochemical or cellular activity. Conjugated antibodies are useful for diagnostic or kit purposes, and include antibodies coupled with a label or polypeptide, e.g., a dye, isotopes, enzyme, or metal, see, e.g., Le Doussal, *et al.* (1991) *J. Immunol.* 146:169-175; Gibellini, *et al.* (1998) *J. Immunol.* 160:3891-3898; Hsing and Bishop (1999) *J. Immunol.* 162:2804-2811; Everts, 10 *et al.* (2002) *J. Immunol.* 168:883-889.

The invention provides a kit, where the kit comprises a compartment containing an agonistic anti-IL-23R antibody, an antigenic fragment thereof, or a nucleic acid encoding an agonistic anti-IL-23R antibody, or a fragment thereof. In another embodiment the kit has a compartment, a nucleic 15 acid, e.g., a probe, primer, or molecular beacon, see, e.g., Zammattéo, *et al.* (2002) *Biotech. Annu. Rev.* 8:85-101; Klein (2002) *Trends Mol. Med.* 8:257-260.

The kit may comprise, e.g., a reagent and a compartment, a reagent and instructions for use, or a reagent with both a compartment and instructions for use. A kit for determining the binding of a test compound, e.g., acquired from a biological sample or from a chemical library, can comprise a 20 control compound, a labeled compound, and a method for separating free labeled compound from bound labeled compound. Diagnostic assays can be used with biological matrices such as live cells, cell extracts and lysates, fixed cells, cell cultures, bodily fluids, or forensic samples. Various assay formats exist, such as radioimmunoassays (RIA), ELISA, and lab on a chip (U.S. Pat. Nos. 6,176,962 and 6,517,234).

25 The method can further comprise contacting a sample from a control subject, normal subject, or normal tissue or fluid from the test subject, with the binding composition. Moreover, the method can additionally comprise comparing the specific binding of the composition to the test subject with the specific binding of the composition to the normal subject, control subject, or normal tissue or fluid from the test subject. Expression or activity of a test sample or test subject can be compared 30 with that from a control sample or control subject. A control sample can comprise, e.g., a sample of non-affected or non-inflamed tissue in a patient suffering from an immune disorder. Expression or activity from a control subject or control sample can be provided as a predetermined value, e.g., acquired from a statistically appropriate group of control subjects.

35 VI. Diagnostic uses; Therapeutic Compositions; Methods.

The present invention provides methods for the treatment and diagnosis of healing, improper healing, wound healing, and improper wound healing, e.g., of the skin. Provided are methods of improving normal wound healing, e.g., by improving the rate of healing, and of treating improper wound healing, e.g., wounds characterized by ulcers or excess fibrosis. Moreover, the invention provides methods of treating and preventing wound-related infections.

Gene therapy of skin disorders may be performed using a variety of methods. Delivery vehicles are well described in the art, see, e.g., Boulikas (1998) *Gene Therapy and Molecular Biology, Vol. 1*, Gene Therapy Press, Palo Alto, CA; Jolly, *et al.* (1994) *Cancer Gene Therapy* 1:51-64; Kimura, *et al.* (1994) *Human Gene Therapy* 5:845-852; and Kaplitt, *et al.* (1994) *Nat. Genetics* 6:148-153.

To prepare pharmaceutical or sterile compositions including a cytokine or a small molecule agonist, the entity is admixed with a pharmaceutically acceptable carrier or excipient which is preferably inert. Preparation of such pharmaceutical compositions is known in the art, see, e.g., *Remington's Pharmaceutical Sciences* and *U.S. Pharmacopeia: National Formulary*, Mack Publishing Company, Easton, PA (1984).

Cytokines are normally administered parentally, preferably intravenously. Since such proteins or peptides may be immunogenic they are preferably administered slowly, either by a conventional IV administration set or from a subcutaneous depot, e.g. as taught by Tomasi, *et al.*, U.S. patent 4,732,863. Means to minimize immunological reactions may be applied. Small molecule entities may be orally active. For treatment of skin disorders, the present invention may also be administered topically, see, e.g., Gilman, *et al.* (eds.) (1990) *Goodman and Gilman's: The Pharmacological Bases of Therapeutics, 8th ed.*, Pergamon Press; and *Remington's Pharmaceutical Sciences, 17th ed.* (1990), Mack Publishing Co., Easton, Penn.

Parenteral therapeutics may be administered in aqueous vehicles such as water, saline, or buffered vehicles with or without various additives and/or diluting agents. Alternatively, a suspension, such as a zinc suspension, can be prepared to include the peptide. Such a suspension can be useful for subcutaneous (SQ) or intramuscular (IM) injection, see, e.g., Avis, *et al.* (eds.) (1993) *Pharmaceutical Dosage Forms: Parenteral Medications 2d ed.*, Dekker, NY; Lieberman, *et al.* (eds. 1990) *Pharmaceutical Dosage Forms: Tablets 2d ed.*, Dekker, NY; Lieberman, *et al.* (eds. 1990) *Pharmaceutical Dosage Forms: Disperse Systems* Dekker, NY; Fodor, *et al.* (1991) *Science* 251:767-773, Coligan (ed.) *Current Protocols in Immunology*; Hood, *et al.* *Immunology Benjamin/Cummings*; Paul (ed.) *Fundamental Immunology*; Academic Press; Parce, *et al.* (1989) *Science* 246:243-247; Owicki, *et al.* (1990) *Proc. Natl. Acad. Sci. USA* 87:4007-4011; and Blundell and Johnson (1976) *Protein Crystallography*, Academic Press, New York.

Selecting an administration regimen for a therapeutic depends on several factors, including the serum or tissue turnover rate of the entity, the level of symptoms, the immunogenicity of the

entity, and the accessibility of the target cells, timing of administration, absorption through epithelial layers, etc. Preferably, an administration regimen maximizes the amount of therapeutic delivered to the patient consistent with an acceptable level of side effects. Accordingly, the amount of biologic delivered depends in part on the particular entity and the severity of the condition being treated.

- 5 Guidance in selecting appropriate doses of cytokine or small molecules are determined using standard methodologies.

Determination of the appropriate dose is made by the clinician, e.g., using parameters or factors known or suspected in the art to affect treatment or predicted to affect treatment. Generally, the dose begins with an amount somewhat less than the optimum dose and it is increased by small
10 increments thereafter until the desired or optimum effect is achieved relative to any negative side effects. Important diagnostic measures include those of symptoms of, e.g., the inflammation or level of inflammatory cytokines produced. Preferably, a biologic that will be used is derived from the same species as the animal targeted for treatment, thereby minimizing a humoral response to the reagent.

- 15 Antibodies, antibody fragments, and cytokines can be provided by continuous infusion, or by doses at intervals of, e.g., one day, one week, or 1-7 times per week. Doses may be provided intravenously, subcutaneously, topically, orally, nasally, rectally, intramuscular, intracerebrally, or by inhalation. A preferred dose protocol is one involving the maximal dose or dose frequency that avoids significant undesirable side effects. A total weekly dose is generally at least 0.05 $\mu\text{g/kg}$ body
20 weight, more generally at least 0.2 $\mu\text{g/kg}$, most generally at least 0.5 $\mu\text{g/kg}$, typically at least 1 $\mu\text{g/kg}$, more typically at least 10 $\mu\text{g/kg}$, most typically at least 100 $\mu\text{g/kg}$, preferably at least 0.2 mg/kg, more preferably at least 1.0 mg/kg, most preferably at least 2.0 mg/kg, optimally at least 10 mg/kg, more optimally at least 25 mg/kg, and most optimally at least 50 mg/kg, see, e.g., Yang, *et al.* (2003) *New Engl. J. Med.* 349:427-434; Herold, *et al.* (2002) *New Engl. J. Med.* 346:1692-1698; Liu, *et al.* (1999)
25 *J. Neurol. Neurosurg. Psych.* 67:451-456; Portielji, *et al.* (2003) *Cancer Immunol. Immunother.* 52:133-144. The desired dose of a small molecule therapeutic, e.g., a peptide mimetic, natural product, or organic chemical, is about the same as for an antibody or polypeptide, on a moles/kg basis.

- The present invention also provides for administration of biologics in combination with
30 known therapies, e.g., steroids, particularly glucocorticoids, which alleviate the symptoms, e.g., associated with inflammation, or antibiotics or anti-infectives. Daily dosages for glucocorticoids will range from at least about 1 mg, generally at least about 2 mg, and preferably at least about 5 mg per day. Generally, the dosage will be less than about 100 mg, typically less than about 50 mg, preferably less than about 20 mg, and more preferably at least about 10 mg per day. In general, the
35 ranges will be from at least about 1 mg to about 100 mg, preferably from about 2 mg to 50 mg per

day. Suitable dose combinations with antibiotics, anti-infectives, or anti-inflammatories are also known.

The present invention provides agonists and antagonists of IL-23 for modulating genes relating to healing, e.g., wound healing. Also provided are methods of diagnosis of healing, e.g.,
5 involving detecting expression or changes in expression of IL-23 modulated genes and gene products. These genes and gene products include, e.g., nitric oxide synthase 2 (NOS2), lactoferrin, IL-19, DEC-205, CD50 (ICAM-2), IL-25, TNFSF7 (CD27L), eosinophilic basic protein, and others.

The invention provides a method to modulate expression of MMP-7, e.g., for the treatment of wound healing. Matrix proteolysis is a hallmark of inflammation. Matrilysin, a metalloprotease, is
10 used in wound repair (see, e.g., Parks, *et al.* (2001) *Chest* 120:36S-41S; Wilson, *et al.* (1999) *Science* 286:113-117).

The present invention provides methods for modulating activities and proteins relating to neutrophils, such as neutrophil chemoattractants and proteins and metabolites expressed by neutrophils. IL-23 stimulates IL-17 expression which, in turn, stimulates production of chemokines
15 that attract neutrophils. Increased expression or activity of neutrophil response, lactoferrin, IL-17, IL-6, and nitric oxide, are found in a number of inflammatory conditions, and can play a role in modulating wound healing (see, e.g., Tsokos, *et al.* (2002) *Virchows Arch.* 441:494-499; Linden (2001) *Int. Arch. Allergy Immunol.* 126:179-184; Sheppard (2002) *Chest* 121:21S-25S; Redington (2000) *Monaldi Arch. Chest Dis.* 55:317-323; Vignola, *et al.* (2001) *Curr. Allergy Asthma Rep.*
20 1:108-115).

Provided are methods of modulating expression of lactoferrin, a protein produced by neutrophils (see, e.g., Boyton, *et al.* (2002) *Brit. Medical Bull.* 61:1-12; Singh, *et al.* (2002) *Nature* 417:552-555; Gomez, *et al.* (2002) *Infect. Immun.* 70:7050-7053).

Provided are methods for modulating expression of neutrophil elastase, e.g., for modulating
25 wound healing (see, e.g., Tkalcevic, *et al.* (2000) *Immunity* 12:201-210; Aprikyan, *et al.* (2001) *Curr. Opinion Immunol.* 13:535-538; Tremblay, *et al.* (2003) *Curr. Opin. Investig. Drugs* 4:556-565; Lee, *et al.* (2001) *Curr. Opinion Crit. Care* 7:1-7; Shapiro (2002) *Am. J. Respir. Cell Mol. Biol.* 26:266-268).

Also provided are methods to modulate neutrophil attractants for promoting wound healing,
30 e.g., IL-17, nitric oxide, and GRO-alpha. IL-17 modulates neutrophil recruitment (see, e.g., Ye, *et al.* (2001) *J. Exp. Med.* 194:519-527; Ye, *et al.* (2001) *Am. J. Respir. Cell Mol. Biol.* 25:335-340). Nitric oxide, synthesized by nitric oxide synthase, can promote wound healing, e.g., by attracting monocytes and neutrophils to the wound, see, e.g., Schwentker, *et al.* (2002) *Nitric Oxide* 7:1-10; MacMicking, *et al.* (1997) *Annu. Rev. Immunol.* 15:323-350. CXCL-1 (a.k.a. GRO-alpha) promotes
35 wound healing, e.g., by attracting neutrophils to wounds and stimulating keratinocyte proliferation

and angiogenesis (see, e.g., Gillitzer, *et al.* (2001) *J. Leukoc. Biol.* 69:513-521; Li and Thornhill (2000) *Cytokine* 12:1409-1413).

IL-6 promotes the healing of injuries, e.g., skin wounds, see, e.g., Gallucci, *et al.* (2001) *J. Interferon Cytokine Res.* 21:603-609; Sugawara, *et al.* (2001) *Cytokine* 15:328-336; Erdag, *et al.* (2002) *Ann. Surg.* 235:113-124; Nadeau, *et al.* (2002) *Microbes Infect.* 4:1379-1387; Imanishi, *et al.* (2000) *Prog. Retin. Eye Res.* 19:113-129; Gregory, *et al.* (1998) *J. Immunol.* 160:6056-6061.

Interferon-gamma (IFNgamma) mediates proper wound healing, e.g., by modulating actin and collagen content, contractile capacity, and scar formation, see, e.g., Moulin, *et al.* (1998) *Exp. Cell Res.* 238:283-293; Ahdieh, *et al.* (2001) *Am. J. Physiol. Cell Physiol.* 281:C2029-C2038; 10 Cornelissen, *et al.* (2000) *J. Dent. Res.* 79:1782-1788; Shtrichman, *et al.* (2001) *Curr. Opin. Microbiol.* 4:251-259; Ikeda, *et al.* (2002) *Cytokine Growth Factor Rev.* 13:95-109; Rottenberg, *et al.* (2002) *Curr. Opin. Immunol.* 14:444-451).

IFNgamma production is stimulated by CD27 (a.k.a. TNFRSF7). CD27 also stimulates cell proliferation and is implicated in the activation and development of T cells, and in T cell-dependent 15 antibody production, including IgE production, by B cells (see, e.g., Takeda, *et al.* (2000) *J. Immunol.* 164:1741-1745; Nagumo, *et al.* (1998) *J. Immunol.* 161:6496-6502; Tomiyama, *et al.* (2002) *J. Immunol.* 168:5538-5550; Busse and Lemanske (2001) *New Engl. J. Med.* 344:350-362).

MUC5ac serves a number of biological functions, including wound healing, see, e.g., Dohrman, *et al.* (1998) *Biochim. Biophys. Acta* 1406:251-259; Rose, *et al.* (2000) *J. Aerosol. Med.* 20 13:245-261; Rogers (2000) *Monaldi Arch. Chest Dis.* 55:324-332; Enss, *et al.* (2000) *Inflamm. Res.* 49:162-169.

The broad scope of this invention is best understood with reference to the following examples, which are not intended to limit the inventions to the specific embodiments.

EXAMPLES

I. General Methods.

Standard methods of biochemistry and molecular biology are described or referenced, e.g., in
 5 Maniatis, *et al.* (1982) *Molecular Cloning, A Laboratory Manual*, Cold Spring Harbor Press, Cold Spring Harbor, NY; Sambrook and Russell (2001) *Molecular Cloning, 3rd ed.*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY; Wu (1993) *Recombinant DNA, Vol. 217*, Academic Press, San Diego, CA; Innis, *et al.* (eds.) (1990) *PCR Protocols: A Guide to Methods and Applications*, Academic Press, N.Y. Standard methods are also found in Ausbel, *et al.* (2001)
 10 *Current Protocols in Molecular Biology, Vols. 1-4*, John Wiley and Sons, Inc. New York, NY, which describes cloning in bacterial cells and DNA mutagenesis (Vol. 1), cloning in mammalian cells and yeast (Vol. 2), glycoconjugates and protein expression (Vol. 3), and bioinformatics (Vol. 4). Methods for producing fusion proteins are described. See, e.g., Invitrogen (2002) *Catalogue*, Carlsbad, CA; Amersham Pharmacia Biotech (2002), *Catalogue*, Piscataway, NJ; Liu, *et al.* (2001)
 15 *Curr. Protein Pept. Sci.* 2:107-121; Graddis, *et al.* (2002) *Curr. Pharm. Biotechnol.* 3:285-297. Standard methods of histology are described (Carson (1997) *Histotechnology: A Self-Instructional Text, 2nd ed.*, Am. Soc. Clin. Pathol. Press, Chicago, IL; Bancroft and Gamble (eds.) (2002) *Theory and Practice of Histological Techniques, 5th ed.*, W.B. Saunders Co., Phila., PA).

Methods for flow cytometry, including fluorescence activated cell sorting (FACS), are
 20 available, see, e.g., Owens, *et al.* (1994) *Flow Cytometry Principles for Clinical Laboratory Practice*, John Wiley and Sons, Hoboken, NJ; Givan (2001) *Flow Cytometry, 2nd ed.*; Wiley-Liss, Hoboken, NJ; Shapiro (2003) *Practical Flow Cytometry*, John Wiley and Sons, Hoboken, NJ.

Standard methods of histology of the immune system are described, see, e.g., Muller-Harmelink (ed.) (1986) *Human Thymus: Histopathology and Pathology*, Springer Verlag, New York,
 25 NY; Hiatt, *et al.* (2000) *Color Atlas of Histology*, Lippincott, Williams, and Wilkins, Phila, PA; Louis, *et al.* (2002) *Basic Histology: Text and Atlas*, McGraw-Hill, New York, NY.

Methods for antibody production and modification are described in, e.g., Coligan, *et al.* (2001) *Current Protocols in Immunology, Vol. 1*, John Wiley and Sons, Inc., New York; Harlow and Lane (1999) *Using Antibodies, Cold Spring Harbor Laboratory Press*, Cold Spring Harbor, NY);
 30 Harlow and Lane (1988) *Antibodies A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY; Einhauer, *et al.* (2001) *J. Biochem. Biophys. Methods* 49:455-465. Methods for adenovirus engineering and transfection, e.g., into cells or mammals, are described (Hurst, *et al.* (2002) *New Engl. J. Med.* 169:443-453; Danthinne and Imperiale (2000) *Gene Ther.* 7:1707-1714; Carlisle (2002) *Curr. Op. Mol. Ther.* 4:306-312.

35 Methods for protein purification such as immunoprecipitation, column chromatography, electrophoresis, isoelectric focusing, centrifugation, and crystallization, are described (Coligan, *et al.*

(2000) *Current Protocols in Protein Science, Vol. 1*, John Wiley and Sons, Inc., New York).

Chemical analysis, chemical modification, post-translational modification, and glycosylation of proteins is described. See, e.g., Coligan, *et al.* (2000) *Current Protocols in Protein Science, Vol. 2*, John Wiley and Sons, Inc., New York; Walker (ed.) (2002) *Protein Protocols Handbook*, Humana Press, Towota, NJ; Lundblad (1995) *Techniques in Protein Modification*, CRC Press, Boca Raton, FL. Techniques for characterizing binding interactions are described (Coligan, *et al.* (2001) *Current Protocols in Immunology, Vol. 4*, John Wiley and Sons, Inc., New York; Parker, *et al.* (2000) *J. Biomol. Screen.* 5: 77-88; Karlsson, *et al.* (1991) *J. Immunol. Methods* 145:229-240; Neri, *et al.* (1997) *Nat. Biotechnol.* 15:1271-1275; Jonsson, *et al.* (1991) *Biotechniques* 11:620-627; Friguet, *et al.* (1985) *J. Immunol. Methods* 77: 305-319; Hubble (1997) *Immunol. Today* 18:305-306; Shen, *et al.* (2001) *J. Biol. Chem.* 276:47311-47319).

Computer analysis is performed using software for determining, e.g., antigenic fragments, signal and leader sequences, protein folding, and functional domains, are available, see, e.g., Vector NTI® Suite (Informax, Inc., Bethesda, MD); GCG Wisconsin Package (Accelrys, Inc., San Diego, CA), and DeCypher® (TimeLogic Corp., Crystal Bay, Nevada); Menne, *et al.* (2000) *Bioinformatics* 16:741-742. Public sequence databases were also used, e.g., from GenBank and others.

II. Wound Generation in p19 Deficient Mice.

Mice deficient in the p19 subunit of IL-23 (p19 knockout mice; p19KO mice) were injected with 1 mg of heat killed *Mycobacterium tuberculosis* (strain H37 RA) emulsified in incomplete Freund's adjuvant (IFA) over four dorsal-flank sites. After 14-18 days post exposure to bacterial antigen, the p19KO mice showed significant hair loss and skin lesions. These results suggest that in the absence of IL-23, mice have an abnormal immune response to bacterial antigen challenge leading to an impaired wound healing response.

III. Incision Wound Generation and Assessment of Histological Responses.

Incision wounding (5 mm) was performed on the dorsal skin of wild-type (WT) and p19KO mice as described (see, e.g., Cohen and Mast (1990) *Advances in Understanding Trauma and Burn Injury* 30:S149-S155). Eighteen hours after linear incision wounding samples from skin surrounding the incision were harvested and subjected to confocal microscopy. Tissue sections were stained with anti IA-FITC and anti-CD11b-APC. p19KO mice showed delayed recruitment of CD11b+, MHC-class II positive cells, i.e., a delay in the recruitment of bone marrow derived monocytes during the wound healing process.

IL-23 was found to induce recruitment of activated monocytes/macrophages. C57BL/6 mice were injected with 10 micrograms of recombinant IL-23 (rIL-23) at a dorsal intradermal site. Skin samples were stained with anti-IA-FITC and anti-CD11b-APC and analyzed by confocal microscopy.

IL-23 induced recruitment of MHC-class II⁺, CD11b⁺ monocytes to the site of cytokine injection. PBS treated skin samples appeared negative for recruitment of these monocytes. Taken together, the altered healing response of p19KO mice to injected *Mycobacterium tuberculosis* and the IL-23 recruitment of activated monocytes/macrophages suggest a role of IL-23 in the wound-healing response.

IL-23 also induced monocyte/macrophage recruitment in wound bed granulation tissue. Ten micrograms of rIL-23 was delivered intradermally to eight sites surrounding two 6 mm full thickness dorsal excisional wounds immediately post-wounding. On day 3, the two wounds and surrounding skin were excised, frozen in OCT, sectioned and stained with anti-CD11b. Counterstaining was done with hematoxylin (blue). IL-23 treated wound tissue exhibited higher monocyte/macrophage infiltrate than buffer control 3 days post-wounding. Similar results were found for CD31⁺ endothelial cell migration.

Recombinant IL-23 was found to augment wound healing. Ten micrograms of rIL-23 was delivered intradermally to eight sites surrounding 6 mm full thickness excisional wounds on the backs of healing competent wild-type Balb/c mice immediately post wounding. On day 3 post wounding the two excisional wounds plus additional surrounding skin were excised. Wounds were fixed in formaldehyde, paraffin embedded and stained with hematoxylin and eosin. Histology slides were assessed for the depth of granulation tissue. IL-23 treated mice exhibited a greater thickness of the granulation tissue layer as compared to buffer control treated mice. Increase in the granulation tissue layer is imperative for late stage wound healing (see, e.g., Schaffer and Nanney, *supra*). Similar results were at 3, 6, and 10 days post wounding.

The same procedure was utilized to determine if IL-23 augments the normal re-epithelialization that occurs in wound healing. IL-23 treated mice exhibited thicker keratinocyte layers than buffer control mice.

IV. Incision Wound Healing and Assessment of Breaking Strength.

Twelve week old male C57Bl/6NT mice were shaved and hair was removed from the dorsum using a depilatory (Nair®, Church and Dwight Co., Princeton, NJ). Two 0.5 cm incisional wounds were created approximately 4 cm down from the nape of the neck and 1 cm away from the midline using surgical scissors. Each mouse received four 20 microliter intradermal injections with either saline or mIL-23 (10 micrograms per mouse) around the periphery of each wound to span the entire length of the incision on both sides. Each wound was closed using a medical adhesive (Mastisol®, Ferndale Labs., Ferndale, MI) and transparent dressing (OpSite 3000®, Smith and Nephew, Largo, FL). Three days later, mice were anesthetized with a ketamine/xylazine cocktail and wounds were analyzed *in vivo* using a biomechanical tissue characterizer (BTC-2000, SRLI Technologies, Nashville, TN), according to manufacturer's instructions. The BTC-2000 uses a high resolution

target laser and vacuum to measure skin deformation over a range of negative pressure and time. Negative pressure was applied at a rate of 10 mm Hg/second until wound rupture occurred.

Data were analyzed by plotting skin deformation as a function of negative pressure to give: 1) the stiffness of the healing wound (mmHg negative pressure required to deform the tissue) and 2) the total strength of the healing wound (i.e. mmHg negative pressure required to break the wound). Tests comparing saline-treated controls with mIL-23-treated experimentals demonstrated that 46% more negative pressure was required to break mIL-23-treated wounds (195.6 mm Hg) than to break to saline-treated wounds (134.1 mm Hg). Moreover, the healing wound following mIL-23 treatment (262.7 mm Hg/mm) was stiffer in comparison to saline-treated controls (198.4 mm Hg/mm).

Administration of an IL-23 agonist is contemplated to result in a desirable increase in stiffness (or reduction in looseness) of the healed wound, beyond which, an undesirable increase in stiffness (or undesirable reduction of looseness) occurs. Hence, an IL-23 antagonist is provided for reducing any undesirable stiffness (or undesired reduction of looseness) of a naturally healing wound or an IL-23 agonist treated wound. Alternatively, for example, an IL-23 agonist is contemplated to result in a decrease in undesirable stiffness, e.g., due to excess fibrosis. Hence, an IL-23 agonist is provided to reduce any undesirable stiffness.

V. IL-23 Treatment Increases Granulation Tissue.

The wound healing response in the mouse back excisional model is mediated by the combination of increased granulation tissue formation, re-epithelialization, and wound contraction. One phase of wound contraction in this model occurs very early and contributes significantly to the overall wound closure. To confirm that IL-23 wound healing promotion activities were also seen in a mouse model that did not have a prominent early wound contraction aspect to the healing response, the mouse head excisional model was chosen.

10 micrograms of recombinant IL-23 was delivered intradermally to 4 sites surrounding one 3 mm full thickness excisional wound on the crown of Balb/c mice's heads immediately post-wounding. On Day 3 post-wounding, the wound and surrounding skin were excised. Wounds were fixed in formaldehyde, paraffin embedded, and stained with hematoxylin and eosin (H/E). Histology slides from the center of the wound were assessed for the depth of granulation tissue.

The data at day 3 post wound infliction indicates that the increased granulation tissue activity resulting from IL-23 treatment was seen prior to the start of later-stage contraction and therefore is independent of early phase wound contraction. Note that a later-wound contraction phase also occurs in this model, but this occurs only after day 4-6.

VI. Gene Therapy.

IL-23 is delivered to the site of a cutaneous wound using gene therapy technology.

Therapeutic delivery to the skin is accomplished by *ex vivo* or *in vivo* methods (see, e.g., Khavari (1997) *Mol. Med. Today* 3:533-538 and Khavari, *et al.* (2002) *J. Int. Med.* 252:1-10).

Recombinant adenoviral preparation were prepared by conventional techniques, see, e.g.,
 5 Srivastava in WO 93/09239, Samulski *et al.* (1989) *J Virol.* 63:3822-3828; Mendelson *et al.* (1988) *Virol.* 166:154-165; and Flotte *et al.* (1993) *Proc. Natl. Acad. Sci. USA* 90:10613-10617). Increasing number of adenoviral particles encoding mouse IL-23 hyperkine or green fluorescent protein (GFP) or saline diluent were delivered intradermally to 8 sites surrounding two 6 mm full thickness excisional wounds on the backs of Balb/c mice immediately post-wounding. On Day 3 post-
 10 wounding, the two wounds and surrounding skin were excised. Wounds were fixed in formaldehyde, paraffin embedded, and stained with hematoxylin and eosin (H/E). Histology slides from the center of the wound were assessed for the depth of granulation tissue. Delivery of mIL-23 hyperkine by this method resulted in an increased wound healing response as compared to vehicle alone.

15 VII. Gene Expression with mIL-23 Hyperkine Treatment.

Excisional wounds were created on the backs of C57BI6/NT mice. Wounds were treated with 10 micrograms of mIL-23 hyperkine or saline control by intradermal (i.d.) injection around the periphery of the wound. Wounds were harvested at Day 1 and at Day 3 post-wounding. Skin samples distal to the wounded site were obtained as "non-wounded" controls. Taqman® real time
 20 PCR analysis (Applied Biosystems, Foster City, CA) was performed on the tissue samples. Expression data were relative to ubiquitin expression, where ubiquitin expression was set to one (1.0). Expression data were then compared according to the indicated pairs of data sets.

Expression was examined with and without IL-23 hyperkine treatment, in the absence of wounding (Table 1) and with and without IL-23 hyperkine with wounding (Table 2).. C57BI6/NT
 25 mice were treated with IL-23 hyperkine or saline, followed by determination of expression of the indicated gene by Taqman® real time PCR analysis (Table 1). Each mouse was injected intradermally, in the back, with either saline or with 10 micrograms IL-23 hyperkine. Tissue samples were taken and extracted at either 1, 3, or 7 days after injection, where the samples from the three dates were pooled, and then used for Taqman® analysis. The ratio of gene expression with and
 30 without IL-23 hyperkine treatment is shown (Table 1). IL-23 hyperkine provoked an increase in expression, of 2-fold or greater, of 15 of the 157 genes tested. These included IL-6 (33-fold), IL-19 (32-fold), and CXCL-1 (GRO-alpha) (11-fold) (Table 1).

Table 1. Ratio of [Gene expression with IL-23] / [Gene expression with saline]. Control (saline) and experimental (IL-23) data were acquired without wounding.

IL-6	33
IL-19	32
CXCL-1 (GRO-alpha)	11
IL-17	9
mMUC-5ac.fcgi	8
secretory leukoprotease inhibitor (SLPI)	5
granulocyte macrophage-colony stimulating factor (GM-CSF)	5
TNFSF5 (CD40L)	3
MAdCAM-1	3
interferon-gamma (IFN-gamma)	3
IL-9	3
12-lipoxygenase	2
tissue inhibitor of metalloproteinases-1 (TIMP-1)	2
IL-1alpha	2
IL-17RC	2

Expression data from Day 1 (1 day after wounding) and from Day 3 (3 days after wounding) are shown (Table 2). IL-23 with wounding stimulated increases in expression over that found with wounding only for a number of genes associated with healing or neutrophil response, including IL-17, nitric oxide synthase, lactoferrin, and matrix metalloproteases (Table 2).

Table 2. Ratio of [Gene expression with IL-23, with wound] / [Gene expression with saline, with wound]

	Day 1	Day 3
IL-17	234	6.7
nitric oxide synthase 2 (NOS2)	17	--
lactoferrin	5.3	8.9
IL-12 (p35 subunit)	3.9	--
IL-12 (p40 subunit)	3.6	--
CD107 (LAMP-2; mac-3)	3.5	--
integrin beta7 chain	3.4	--
TNFSF11 (RANKL)	3.3	--
CD11a (LFA-1a chain)	3.0	--
IL-19	2.9	2.0
endoperoxidase synthase type II (COX-2)	2.8	--

IL-1RA	2.8	--
IL-17RA	2.7	--
CD29 (integrin b1)	2.6	--
leukotriene A4 hydrolase	2.5	--
MMP-13	2.4	--
MMP-8 (collagenase 2)	2.4	--
leukotriene B4 receptor	2.4	--
TNFRSF11a (RANK)	2.4	--
M-CSF1	2.3	--
TIMP-1	2.3	--
CD14	2.3	--
TNF-alpha converting enzyme (TACE)	2.3	--
Mac-1 alpha subunit (CD11b)	2.3	--
DEC-205	2.2	--
CD50 (ICAM-1)	2.2	--
CCL3 MIP-1alpha	2.2	--
CD86 (B7-2)	2.2	--
CD80 (B7-1)	2.1	--
IL-80	2.1	--
fibronectin	2.0	--
TNFRSF6a (FAS)	2.0	--
neutrophil elastase	0.1	13.7
interferon-gamma (IFN-gamma)	0.7	8.7
matrilysin (MMP-7)	0.1	5.3
TNFRSF7u (CD27)	0.8	4.9
IL-13Ra2	1.4	4.8
IL-12Rb1	1.6	3.8
myeloperoxidase	1.1	2.7
eosinophil major basic protein	0.4	2.7
proteinase 3	1.6	2.6
IL-15	1.4	2.5

IL-23 dependent trends in gene expression in common with two strains of mice (C57B1/6NT mice; Balb/c mice) were as follows. A panel of 158 genes was screened to determine which are regulated by IL-23 treatment during wound healing. Only five of these genes were upregulated by at

least 2-fold (Table 3), and only three of the genes were down regulated by 2-fold or greater (Table 3).

IL-17 has been shown to regulate genes important for remodeling in other tissue systems. IL-17 potentiates both MMP-3 and TIMP-1 expression and secretion stimulated by TNF and IL-1 in human colonic subepithelial myofibroblasts (Bamba, *et al.* (2003) *J Gastroenterol.* 38:548-554). In human osteoblasts, IL-17 synergizes with TNF-alpha, TGF-beta, and IFN-gamma to stimulate production of MMP-13 (Rifas and Arackal (2003) *Arthritis Rheum.* 48:993-1001). Given that IL-17 alone has minimal effects in these models, the major role of IL-17 may be to amplify the remodeling response. IL-19 is found in skin and has been associated with psoriasis (Ghoreschik, *et al.* (2003) *Nature Med.* 9:40-46; Gallagher, *et al.* (2000) *Genes Immunity* 1:442-450). Lactoferrin is an iron-binding protein present in mature neutrophil granules with a number of biological functions including anti-microbial properties and may thus protect against bacterial infection in the wound (Masson, *et al.* (1969) *J. Exp. Med.* 130:643-658; Farnaud and Evans (2003) *Mol Immunol.* 40:395-405). ICAM-2 is constitutively expressed on all vascular endothelial cells suggesting that IL-23 may promote angiogenesis in the wound (Yasuda, *et al.* (2002) *Am. J. Physiol.* 282:C917-C925; Sakurai, *et al.* (2003) *Invest. Ophthalmol. Vis. Sci.* 44:2743-2749; Moromizato, *et al.* (2000) *Am. J. Pathol.* 157:1277-1281). DEC-205 is a dendritic cell marker implicated in the dendritic cell maturation (Anjuere, *et al.* (1999) *Blood* 93:590-598; Bonifax, *et al.* (2002) *J. Exp. Med.* 196:1627-1638). Thus IL-23 of the present invention can promote immune surveillance in the skin.

Taken together, IL-23 treatment in wounds may have pleiotropic effects to accelerate the wound healing response. IL-23 treatment also reduced the expression of 3 genes in the wound environment, i.e., IL-25, TNSF7, and eosinophilic basic protein. Infusion of mice with IL-25 induced IL-4, IL-5, and IL-13 gene expression associated with lung remodeling pathology (Fort, *et al.* (2001) *Immunity* 15:985-995). Thus, IL-23 may modulate or reduce excess fibrosis in wound healing.

Relevant methodology was as follows. Hair was shaved and removed with Nair from the dorsum of 9 week old male C57Bl/6NT and 8 week old male Balb/c mice. Two 6 mm diameter excisional wounds were created approximately 3.5 cm down from the nape of the neck and 0.6 cm away from the midline using a punch biopsy tool. Each mouse was given mIL-23 hyperkine (10 micrograms per mouse) or saline in four 20 microliters intradermal injections around the periphery of each wound. A transparent dressing was applied and mice were allowed to recover and heal. One day later, mice were sacrificed and wounds were excised with a margin of approximately 1.5 mm. In addition, a sample of non-wounded skin from each mouse was obtained. All samples were snap frozen in liquid nitrogen and stored at -80 degrees Celsius for further processing. RNA was extracted from each sample, pooled together into appropriate groups, and analyzed by real time PCR for expression of a panel of 158 candidate genes.

Table 3. Modulation of gene expression, by at least 2-fold, in response to IL-23 in both C57B1/6NT mice and Balb/c mice.

Genes up-regulated.	C57B1/6NT mouse	Balb/c mouse
IL-17	234.5	207.9
Lactoferrin	5.3	3.3
IL-19	2.9	2.2
DEC-205	2.3	2.8
CD50 (ICAM-2)	2.3	2.1
Genes down-regulated.	C57B1/6NT mouse	Balb/c mouse
IL-25	5.88	5.56
TNSF7-CD27L	3.45	100
Eosinophilic basic protein	2.13	2.33

VIII. Agonist Anti-IL-23R Antibodies.

Mouse anti-IL-23R antibodies were prepared against human IL-23R using standard methods.

- 5 The resultant anti-IL-23R antibodies were screened for agonist activity using cells transfected with hIL-23R and hIL-12Rbeta1, where agonist activity was determined by increases in cell proliferation. Proliferation of transfectant Ba/F3 cells was measured by colorimetric methods using Alamar Blue, a growth indicator dye.

- 10 Cell proliferation was measured after culture in Roswell Park Memorial Institute (RPMI)-1640 medium, fetal calf serum (10%), 0.05 mM 2-mercaptoethanol, glutamine, penicillin, streptomycin, and mouse interleukin-3 (mIL-3) (10 ng/ml). Cells were incubated in the presence of one concentration of antibody, where the concentrations ranged from 0.01 to 10,000 ng/ml. Baseline cell proliferation was that in absence of antibody. Maximal proliferation with the antibodies tested occurred in the concentration range of 1000 to 10,000 ng/ml. "Detectable stimulation," e.g., of cell activity, cell proliferation, or a predetermined activity, refers, e.g., to a comparison of proliferation in the presence and absence of, e.g., an IL-23 agonist. "Detectable" may be a function of the context, e.g., of the reagents, instrumentation, or biological system.

- 15 Antibodies that stimulated expression included TC48-8B10.D5 (light chain, SEQ ID NOs:9 and 10; heavy chain, SEQ ID NOs:11 and 12), TC48-1H3.G5, TC48-2C9A5, and TC48-5B12.C9. 20 Of these antibodies, TC48-8B10.D5 showed the greatest agonist activity, in terms of antibody concentration provoking ½ maximal increase in cell proliferation. 100% cell proliferation means maximal increase in cell proliferation in response to the antibody, in titration curves (Table 4). The predicted cleavage point for the signal sequence of the light chain of TC48-8B10.D5 is between

amino acids 22 (Ala) and 23 (Glu) of SEQ ID NO:10, while the predicted cleavage point for the heavy chain is between amino acids 19 (Ser) and 20 (Gln) of SEQ ID NO:12.

Table 4. Proliferation of Ba/F3-2.2lo cells in response to agonistic antibodies.

Antibody	Maximal increase cell proliferation in response to antibody (units of OD _{570-600 nm})	Antibody concentration providing ½ maximal increase in cell proliferation.
TC48-8B10.D5 (SEQ ID NOs: 9-12)	0.026	10 ng/ml
TC48-2C9A5	0.021	20 ng/ml
TC48-1H3.G5	0.018	50 ng/ml
TC48-5B12.C9	0.026	100 ng/ml

5 The invention provides an agonistic murine antibody that specifically binds human IL-23R, TC48-8B10.D5. This antibody comprises a nucleic acid (SEQ ID NO:9) and polypeptide (SEQ ID NO:10) of the light chain, and the nucleic acid (SEQ ID NO:11) and polypeptide (SEQ ID NO:12) of the heavy chain of TC48-8B10.D5. Also provided are the nucleic acids and polypeptides comprising the hypervariable regions of SEQ ID NOs:9-12.

10 The hypervariable regions of the light chain are: ITSTDIDDDMI (amino acids 46-56); EGNTLRP (amino acids 72-78); and LQSDNMPLT (amino acids 111-119), of SEQ ID NO:10). The hypervariable regions of the heavy chain are: GYTFTSYWMN (amino acids 45-54); MIDPLDSETHYNQMFKD (amino acids 69-87); and GDNYAMDY (amino acids 118-126), of SEQ ID NO:12.

15

IX. Rat Anti-Mouse IL-23R Antibodies and Wound Healing.

Antibodies against mouse IL-23R were raised in rats using standard methods, resulting in antibodies named 5C10, 29A5, and 10E11 (Table 5). An IL-23 hyperkine comprising a p19 subunit and p40 subunit covalently connected by way of an elasti-linker (InvivoGen, San Diego, CA) was prepared, and named "IL-23 elastikine." The anti-mouse IL-23R antibodies, IL-23 elastikine (control), 36E10 antibody (isotype control), and saline (control) were tested in a wound healing assay. The results demonstrated a stimulation of wound healing with the IL-23 elastikine, and lesser stimulation with the 5C10 antibody (Table 5):

25

Table 5. Wound healing; thickness of granulation tissue (micrometers).

Saline control	170 micrometers
IL-23 elastikine control	300
36E10 isotype control	170
5C10	205
29A5	170
10E11	125

The sequences in the Sequence Listing are summarized (Table 6):

Table6. Sequences in Sequence Listing.

SEQ ID NO:	Nucleic acid or Polypeptide
1	Mouse p19 nucleic acid
2	Mouse p19 polypeptide
3	Human p19 nucleic acid
4	Human p19 polypeptide
5	Mouse hyperkine nucleic acid
6	Mouse hyperkine polypeptide
7	Human hyperkine nucleic acid
8	Human hyperkine polypeptide
9	Mouse anti-human agonist Ab light chain nucleic acid
10	Mouse anti-human agonist Ab light chain polypeptide
11	Mouse anti-human agonist Ab heavy chain nucleic acid
12	Mouse anti-human agonist Ab heavy chain polypeptide

5

Many modifications and variations of this invention can be made without departing from its spirit and scope, as will be apparent to those skilled in the art. The specific embodiments described herein are offered by way of example only, and the invention is to be limited by the terms of the appended claims, along with the full scope of equivalents to which such claims are entitled; and the invention is not to be limited by the specific embodiments that have been presented herein by way of example.

10

30a

SEQUENCE LISTING

<110> Schering Corporation

<120> Uses of mammalian cytokine; related reagents

<130> 3085-1091CA

<140> CA 2,506,672

<141> 2003-12-18

<150> US 60/436,274

<151> 2002-12-23

<160> 12

<170> PatentIn version 3.1

<210> 1

<211> 1203

<212> DNA

<213> Mus musculus

<220>

<221> CDS

<222> (113)..(700)

<220>

<221> mat_peptide

<222> (176)..(700)

<400> 1

cgcttagaag tcggactaca gagttagact cagaaccaaa ggaggtggat aggggggtcca 60

caggcctggg gcagatcaca gagccagcca gatctgagaa gcaggggaaca ag atg ctg 118

Met Leu

-20

gat tgc aga gca gta ata atg cta tgg ctg ttg ccc tgg gtc act cag 166

Asp Cys Arg Ala Val Ile Met Leu Trp Leu Leu Pro Trp Val Thr Gln

-15

-10

-5

ggc ctg gct gtg cct agg agt agc agt cct gac tgg gct cag tgc cag 214

Gly Leu Ala Val Pro Arg Ser Ser Ser Pro Asp Trp Ala Gln Cys Gln

-1 1

5

10

cag ctc tct cgg aat ctc tgc atg cta gcc tgg aac gca cat gca cca 262

Gln Leu Ser Arg Asn Leu Cys Met Leu Ala Trp Asn Ala His Ala Pro

15

20

25

30b

gcg gga cat atg aat cta cta aga gaa gaa gag gat gaa gag act aaa	310
Ala Gly His Met Asn Leu Leu Arg Glu Glu Glu Asp Glu Glu Thr Lys	
30 35 40 45	
aat aat gtg ccc cgt atc cag tgt gaa gat ggt tgt gac cca caa gga	358
Asn Asn Val Pro Arg Ile Gln Cys Glu Asp Gly Cys Asp Pro Gln Gly	
50 55 60	
ctc aag gac aac agc cag ttc tgc ttg caa agg atc cgc caa ggt ctg	406
Leu Lys Asp Asn Ser Gln Phe Cys Leu Gln Arg Ile Arg Gln Gly Leu	
65 70 75	
gct ttt tat aag cac ctg ctt gac tct gac atc ttc aaa ggg gag cct	454
Ala Phe Tyr Lys His Leu Leu Asp Ser Asp Ile Phe Lys Gly Glu Pro	
80 85 90	
gct cta ctc cct gat agc ccc atg gag caa ctt cac acc tcc cta cta	502
Ala Leu Leu Pro Asp Ser Pro Met Glu Gln Leu His Thr Ser Leu Leu	
95 100 105	
gga ctc agc caa ctc ctc cag cca gag gat cac ccc cgg gag acc caa	550
Gly Leu Ser Gln Leu Leu Gln Pro Glu Asp His Pro Arg Glu Thr Gln	
110 115 120 125	
cag atg ccc agc ctg agt tct agt cag cag tgg cag cgc ccc ctt ctc	598
Gln Met Pro Ser Leu Ser Ser Ser Gln Gln Trp Gln Arg Pro Leu Leu	
130 135 140	
cgt tcc aag atc ctt cga agc ctc cag gcc ttt ttg gcc ata gct gcc	646
Arg Ser Lys Ile Leu Arg Ser Leu Gln Ala Phe Leu Ala Ile Ala Ala	
145 150 155	
cgg gtc ttt gcc cac gga gca gca act ctg act gag ccc tta gtg cca	694
Arg Val Phe Ala His Gly Ala Ala Thr Leu Thr Glu Pro Leu Val Pro	
160 165 170	
aca gct taaggatgcc caggttccca tggctaccat gataagacta atctatcagc	750
Thr Ala	
175	
ccagacatct accagttaat taaccatta ggacttgtgc tgttcttgtt tcgtttgttt	810
tgcgtgaagg gcaaggacac cattattaaa gagaaaagaa acaaacccca gagcaggcag	870
ctggctagag aaaggagctg gagaagaaga ataaagtctc gagcccttgg ccttggaagc	930
gggcaagcag ctgcgtggcc tgaggggaag ggggcggtgg catcgagaaa ctgtgagaaa	990
accagagca tcagaaaaag tgagcccagg ctttggccat tatctgtaag aaaaacaaga	1050
aaaggggaac attatacttt cctgggtggc tcagggaat gtgcagatgc acagtactcc	1110
agacagcagc tctgtacctg cctgctctgt ccctcagttc taacagaatc tagtcactaa	1170
gaactaacag gactaccaat acgaactgac aaa	1203

30c

<210> 2
 <211> 196
 <212> PRT
 <213> Mus musculus

<400> 2
 Met Leu Asp Cys Arg Ala Val Ile Met Leu Trp Leu Leu Pro Trp Val
 -20 -15 -10
 Thr Gln Gly Leu Ala Val Pro Arg Ser Ser Ser Pro Asp Trp Ala Gln
 -5 -1 1 5 10
 Cys Gln Gln Leu Ser Arg Asn Leu Cys Met Leu Ala Trp Asn Ala His
 15 20 25
 Ala Pro Ala Gly His Met Asn Leu Leu Arg Glu Glu Glu Asp Glu Glu
 30 35 40
 Thr Lys Asn Asn Val Pro Arg Ile Gln Cys Glu Asp Gly Cys Asp Pro
 45 50 55
 Gln Gly Leu Lys Asp Asn Ser Gln Phe Cys Leu Gln Arg Ile Arg Gln
 60 65 70 75
 Gly Leu Ala Phe Tyr Lys His Leu Leu Asp Ser Asp Ile Phe Lys Gly
 80 85 90
 Glu Pro Ala Leu Leu Pro Asp Ser Pro Met Glu Gln Leu His Thr Ser
 95 100 105
 Leu Leu Gly Leu Ser Gln Leu Leu Gln Pro Glu Asp His Pro Arg Glu
 110 115 120
 Thr Gln Gln Met Pro Ser Leu Ser Ser Ser Gln Gln Trp Gln Arg Pro
 125 130 135
 Leu Leu Arg Ser Lys Ile Leu Arg Ser Leu Gln Ala Phe Leu Ala Ile
 140 145 150 155
 Ala Ala Arg Val Phe Ala His Gly Ala Ala Thr Leu Thr Glu Pro Leu
 160 165 170
 Val Pro Thr Ala
 175

<210> 3
 <211> 570
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (1)..(567)

30d

<220>

<221> mat_peptide

<222> (64)..(567)

<400> 3

atg ctg ggg agc aga gct gta atg ctg ctg ttg ctg ctg ccc tgg aca	48
Met Leu Gly Ser Arg Ala Val Met Leu Leu Leu Leu Leu Pro Trp Thr	
-20 -15 -10	
gct cag ggc aga gct gtg cct ggg ggc agc agc cct gcc tgg act cag	96
Ala Gln Gly Arg Ala Val Pro Gly Gly Ser Ser Pro Ala Trp Thr Gln	
-5 -1 1 5 10	
tgc cag cag ctt tca cag aag ctc tgc aca ctg gcc tgg agt gca cat	144
Cys Gln Gln Leu Ser Gln Lys Leu Cys Thr Leu Ala Trp Ser Ala His	
15 20 25	
cca cta gtg gga cac atg gat cta aga gaa gag gga gat gaa gag act	192
Pro Leu Val Gly His Met Asp Leu Arg Glu Glu Gly Asp Glu Glu Thr	
30 35 40	
aca aat gat gtt ccc cat atc cag tgt gga gat ggc tgt gac ccc caa	240
Thr Asn Asp Val Pro His Ile Gln Cys Gly Asp Gly Cys Asp Pro Gln	
45 50 55	
gga ctc agg gac aac agt cag ttc tgc ttg caa agg atc cac cag ggt	288
Gly Leu Arg Asp Asn Ser Gln Phe Cys Leu Gln Arg Ile His Gln Gly	
60 65 70 75	
ctg att ttt tat gag aag ctg cta gga tcg gat att ttc aca ggg gag	336
Leu Ile Phe Tyr Glu Lys Leu Leu Gly Ser Asp Ile Phe Thr Gly Glu	
80 85 90	
cct tct ctg ctc cct gat agc cct gtg gcg cag ctt cat gcc tcc cta	384
Pro Ser Leu Leu Pro Asp Ser Pro Val Ala Gln Leu His Ala Ser Leu	
95 100 105	
ctg ggc ctc agc caa ctc ctg cag cct gag ggt cac cac tgg gag act	432
Leu Gly Leu Ser Gln Leu Leu Gln Pro Glu Gly His His Trp Glu Thr	
110 115 120	
cag cag att cca agc ctc agt ccc agc cag cca tgg cag cgt ctc ctt	480
Gln Gln Ile Pro Ser Leu Ser Pro Ser Gln Pro Trp Gln Arg Leu Leu	
125 130 135	
ctc cgc ttc aaa atc ctt cgc agc ctc cag gcc ttt gtg gct gta gcc	528
Leu Arg Phe Lys Ile Leu Arg Ser Leu Gln Ala Phe Val Ala Val Ala	
140 145 150 155	
gcc cgg gtc ttt gcc cat gga gca gca acc ctg agt ccc taa	570
Ala Arg Val Phe Ala His Gly Ala Ala Thr Leu Ser Pro	
160 165	

30e

<210> 4
 <211> 189
 <212> PRT
 <213> Homo sapiens

<400> 4
 Met Leu Gly Ser Arg Ala Val Met Leu Leu Leu Leu Leu Pro Trp Thr
 -20 -15 -10
 Ala Gln Gly Arg Ala Val Pro Gly Gly Ser Ser Pro Ala Trp Thr Gln
 -5 -1 1 5 10
 Cys Gln Gln Leu Ser Gln Lys Leu Cys Thr Leu Ala Trp Ser Ala His
 15 20 25
 Pro Leu Val Gly His Met Asp Leu Arg Glu Glu Gly Asp Glu Glu Thr
 30 35 40
 Thr Asn Asp Val Pro His Ile Gln Cys Gly Asp Gly Cys Asp Pro Gln
 45 50 55
 Gly Leu Arg Asp Asn Ser Gln Phe Cys Leu Gln Arg Ile His Gln Gly
 60 65 70 75
 Leu Ile Phe Tyr Glu Lys Leu Leu Gly Ser Asp Ile Phe Thr Gly Glu
 80 85 90
 Pro Ser Leu Leu Pro Asp Ser Pro Val Ala Gln Leu His Ala Ser Leu
 95 100 105
 Leu Gly Leu Ser Gln Leu Leu Gln Pro Glu Gly His His Trp Glu Thr
 110 115 120
 Gln Gln Ile Pro Ser Leu Ser Pro Ser Gln Pro Trp Gln Arg Leu Leu
 125 130 135
 Leu Arg Phe Lys Ile Leu Arg Ser Leu Gln Ala Phe Val Ala Val Ala
 140 145 150 155
 Ala Arg Val Phe Ala His Gly Ala Ala Thr Leu Ser Pro
 160 165

<210> 5
 <211> 1610
 <212> DNA
 <213> Mus musculus

<220>
 <221> CDS
 <222> (7)..(1599)

30f

```

<400> 5
agatct atg tct gca ctt ctg atc cta gct ctt gtt gga gct gca gtt      48
      Met Ser Ala Leu Leu Ile Leu Ala Leu Val Gly Ala Ala Val
        1             5             10

gct gac tac aaa gac gat gac gac aag ctt atg tgg gag ctg gag aaa      96
Ala Asp Tyr Lys Asp Asp Asp Asp Lys Leu Met Trp Glu Leu Glu Lys
15             20             25             30

gac gtt tat gtt gta gag gtg gac tgg act ccc gat gcc cct gga gaa      144
Asp Val Tyr Val Val Glu Val Asp Trp Thr Pro Asp Ala Pro Gly Glu
              35             40             45

aca gtg aac ctc acc tgt gac acg cct gaa gaa gat gac atc acc tgg      192
Thr Val Asn Leu Thr Cys Asp Thr Pro Glu Glu Asp Asp Ile Thr Trp
              50             55             60

acc tca gac cag aga cat gga gtc ata ggc tct gga aag acc ctg acc      240
Thr Ser Asp Gln Arg His Gly Val Ile Gly Ser Gly Lys Thr Leu Thr
              65             70             75

atc act gtc aaa gag ttt ctr gat gct ggc cag tac acc tgc cac aaa      288
Ile Thr Val Lys Glu Phe Leu Asp Ala Gly Gln Tyr Thr Cys His Lys
      80             85             90

gga ggc gag act ctg agc cac tca cat ctg ctg ctc cac aag aag gaa      336
Gly Gly Glu Thr Leu Ser His Ser His Leu Leu Leu His Lys Lys Glu
95             100             105             110

aat gga att tgg tcc act gaa att tta aaa aat ttc aaa aac aag act      384
Asn Gly Ile Trp Ser Thr Glu Ile Leu Lys Asn Phe Lys Asn Lys Thr
              115             120             125

ttc ctg aag tgt gaa gca cca aat tac tcc gga cgg ttc acg tgc tca      432
Phe Leu Lys Cys Glu Ala Pro Asn Tyr Ser Gly Arg Phe Thr Cys Ser
              130             135             140

tgg ctg gtg caa aga aac atg gac ttg aag ttc aac atc aag agc agt      480
Trp Leu Val Gln Arg Asn Met Asp Leu Lys Phe Asn Ile Lys Ser Ser
              145             150             155

agc agt tcc cct gac tct cgg gca gtg aca tgt gga atg gcg tct ctg      528
Ser Ser Ser Pro Asp Ser Arg Ala Val Thr Cys Gly Met Ala Ser Leu
      160             165             170

tct gca gag aag gtc aca ctg gac caa agg gac tat gag aag tat tca      576
Ser Ala Glu Lys Val Thr Leu Asp Gln Arg Asp Tyr Glu Lys Tyr Ser
175             180             185             190

gtg tcc tgc cag gag gat gtc acc tgc cca act gct gag gag acc ctg      624
Val Ser Cys Gln Glu Asp Val Thr Cys Pro Thr Ala Glu Glu Thr Leu
              195             200             205

ccc att gaa ctg gcg ttg gaa gca cgg cag cag aat aaa tat gag aac      672
Pro Ile Glu Leu Ala Leu Glu Ala Arg Gln Gln Asn Lys Tyr Glu Asn
              210             215             220

```

30g

tac agc acc agc ttc ttc atc agg gac atc atc aaa cca gac ccg ccc	720
Tyr Ser Thr Ser Phe Phe Ile Arg Asp Ile Ile Lys Pro Asp Pro Pro	
225 230 235	
aag aac ttg cag atg aag cct ttg aag aac tca cag gtg gag gtc agc	768
Lys Asn Leu Gln Met Lys Pro Leu Lys Asn Ser Gln Val Glu Val Ser	
240 245 250	
tgg gag tac cct gac tcc tgg agc act ccc cat tcc tac ttc tcc ctc	816
Trp Glu Tyr Pro Asp Ser Trp Ser Thr Pro His Ser Tyr Phe Ser Leu	
255 260 265 270	
aag ttc ttt gtt cga atc cag cgc aag aaa gaa aag atg aag gag aca	864
Lys Phe Phe Val Arg Ile Gln Arg Lys Lys Glu Lys Met Lys Glu Thr	
275 280 285	
gag gag ggg tgt aac cag aaa ggt gcg ttc ctc gta gag aag aca tct	912
Glu Glu Gly Cys Asn Gln Lys Gly Ala Phe Leu Val Glu Lys Thr Ser	
290 295 300	
acc gaa gtc caa tgc aaa ggc ggg aat gtc tgc gtg caa gct cag gat	960
Thr Glu Val Gln Cys Lys Gly Gly Asn Val Cys Val Gln Ala Gln Asp	
305 310 315	
cgc tat tac aat tcc tcr tgc agc aag tgg gca tgt gtt ccc tgc agg	1008
Arg Tyr Tyr Asn Ser Ser Cys Ser Lys Trp Ala Cys Val Pro Cys Arg	
320 325 330	
gtc cga tcc tct aga ggt gga tca ggc tcc gga ggt agt gga ggt ggg	1056
Val Arg Ser Ser Arg Gly Gly Ser Gly Ser Gly Gly Ser Gly Gly Gly	
335 340 345 350	
gga tct aag ctt ctg gct gtg cct agg agt agc agt cct gac tgg gct	1104
Gly Ser Lys Leu Leu Ala Val Pro Arg Ser Ser Ser Pro Asp Trp Ala	
355 360 365	
cag tgc cag cag ctc tct cgg aat ctc tgc atg cta gcc tgg aac gca	1152
Gln Cys Gln Gln Leu Ser Arg Asn Leu Cys Met Leu Ala Trp Asn Ala	
370 375 380	
cat gca cca gcg gga cat atg aat cta cta aga gaa gaa gag gat gaa	1200
His Ala Pro Ala Gly His Met Asn Leu Leu Arg Glu Glu Glu Asp Glu	
385 390 395	
gag act aaa aat aat gtg ccc cgt atc cag tgt gaa gat ggt tgt gac	1248
Glu Thr Lys Asn Asn Val Pro Arg Ile Gln Cys Glu Asp Gly Cys Asp	
400 405 410	
cca caa gga ctc aag gac aac agc cag ttc tgc ttg caa agg atc cgc	1296
Pro Gln Gly Leu Lys Asp Asn Ser Gln Phe Cys Leu Gln Arg Ile Arg	
415 420 425 430	
caa ggt ctg gtt ttt tat aag cac ctg ctt gac tct gac atc ttc aaa	1344
Gln Gly Leu Val Phe Tyr Lys His Leu Leu Asp Ser Asp Ile Phe Lys	
435 440 445	

30h

ggg gag cct gct cta ctc cct gat agc ccc atg gag caa ctt cac acc	1392
Gly Glu Pro Ala Leu Leu Pro Asp Ser Pro Met Glu Gln Leu His Thr	
450 455 460	
tcc cta cta gga ctc agc caa ctc ctc cag cca gag gat cac ccc cgg	1440
Ser Leu Leu Gly Leu Ser Gln Leu Leu Gln Pro Glu Asp His Pro Arg	
465 470 475	
gag acc caa cag atg ccc agc ctg agt tct agt cag cag tgg cag cgc	1488
Glu Thr Gln Gln Met Pro Ser Leu Ser Ser Ser Gln Gln Trp Gln Arg	
480 485 490	
ccc ctt ctc cgt tcc aag atc ctt cga agc ctc cag gcc ttt ttg gcc	1536
Pro Leu Leu Arg Ser Lys Ile Leu Arg Ser Leu Gln Ala Phe Leu Ala	
495 500 505 510	
ata gct gcc cgg gtc ttt gcc cac gga gca gca act ctg act gag ccc	1584
Ile Ala Ala Arg Val Phe Ala His Gly Ala Ala Thr Leu Thr Glu Pro	
515 520 525	
tta gtg cca aca gct taagcggccg c	1610
Leu Val Pro Thr Ala	
530	

<210> 6
 <211> 531
 <212> PRT
 <213> Mus musculus

<400> 6
 Met Ser Ala Leu Leu Ile Leu Ala Leu Val Gly Ala Ala Val Ala Asp
 1 5 10 15
 Tyr Lys Asp Asp Asp Asp Lys Leu Met Trp Glu Leu Glu Lys Asp Val
 20 25 30
 Tyr Val Val Glu Val Asp Trp Thr Pro Asp Ala Pro Gly Glu Thr Val
 35 40 45
 Asn Leu Thr Cys Asp Thr Pro Glu Glu Asp Asp Ile Thr Trp Thr Ser
 50 55 60
 Asp Gln Arg His Gly Val Ile Gly Ser Gly Lys Thr Leu Thr Ile Thr
 65 70 75 80
 Val Lys Glu Phe Leu Asp Ala Gly Gln Tyr Thr Cys His Lys Gly Gly
 85 90 95
 Glu Thr Leu Ser His Ser His Leu Leu Leu His Lys Lys Glu Asn Gly
 100 105 110
 Ile Trp Ser Thr Glu Ile Leu Lys Asn Phe Lys Asn Lys Thr Phe Leu
 115 120 125

30i

Lys	Cys	Glu	Ala	Pro	Asn	Tyr	Ser	Gly	Arg	Phe	Thr	Cys	Ser	Trp	Leu	130	135	140	
Val	Gln	Arg	Asn	Met	Asp	Leu	Lys	Phe	Asn	Ile	Lys	Ser	Ser	Ser	Ser	145	150	155	160
Ser	Pro	Asp	Ser	Arg	Ala	Val	Thr	Cys	Gly	Met	Ala	Ser	Leu	Ser	Ala	165	170	175	
Glu	Lys	Val	Thr	Leu	Asp	Gln	Arg	Asp	Tyr	Glu	Lys	Tyr	Ser	Val	Ser	180	185	190	
Cys	Gln	Glu	Asp	Val	Thr	Cys	Pro	Thr	Ala	Glu	Glu	Thr	Leu	Pro	Ile	195	200	205	
Glu	Leu	Ala	Leu	Glu	Ala	Arg	Gln	Gln	Asn	Lys	Tyr	Glu	Asn	Tyr	Ser	210	215	220	
Thr	Ser	Phe	Phe	Ile	Arg	Asp	Ile	Ile	Lys	Pro	Asp	Pro	Pro	Lys	Asn	225	230	235	240
Leu	Gln	Met	Lys	Pro	Leu	Lys	Asn	Ser	Gln	Val	Glu	Val	Ser	Trp	Glu	245	250	255	
Tyr	Pro	Asp	Ser	Trp	Ser	Thr	Pro	His	Ser	Tyr	Phe	Ser	Leu	Lys	Phe	260	265	270	
Phe	Val	Arg	Ile	Gln	Arg	Lys	Lys	Glu	Lys	Met	Lys	Glu	Thr	Glu	Glu	275	280	285	
Gly	Cys	Asn	Gln	Lys	Gly	Ala	Phe	Leu	Val	Glu	Lys	Thr	Ser	Thr	Glu	290	295	300	
Val	Gln	Cys	Lys	Gly	Gly	Asn	Val	Cys	Val	Gln	Ala	Gln	Asp	Arg	Tyr	305	310	315	320
Tyr	Asn	Ser	Ser	Cys	Ser	Lys	Trp	Ala	Cys	Val	Pro	Cys	Arg	Val	Arg	325	330	335	
Ser	Ser	Arg	Gly	Gly	Ser	Gly	Ser	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	340	345	350	
Lys	Leu	Leu	Ala	Val	Pro	Arg	Ser	Ser	Ser	Pro	Asp	Trp	Ala	Gln	Cys	355	360	365	
Gln	Gln	Leu	Ser	Arg	Asn	Leu	Cys	Met	Leu	Ala	Trp	Asn	Ala	His	Ala	370	375	380	
Pro	Ala	Gly	His	Met	Asn	Leu	Leu	Arg	Glu	Glu	Glu	Asp	Glu	Glu	Thr	385	390	395	400
Lys	Asn	Asn	Val	Pro	Arg	Ile	Gln	Cys	Glu	Asp	Gly	Cys	Asp	Pro	Gln	405	410	415	
Gly	Leu	Lys	Asp	Asn	Ser	Gln	Phe	Cys	Leu	Gln	Arg	Ile	Arg	Gln	Gly	420	425	430	

30j

Leu Val Phe Tyr Lys His Leu Leu Asp Ser Asp Ile Phe Lys Gly Glu
 435 440 445
 Pro Ala Leu Leu Pro Asp Ser Pro Met Glu Gln Leu His Thr Ser Leu
 450 455 460
 Leu Gly Leu Ser Gln Leu Leu Gln Pro Glu Asp His Pro Arg Glu Thr
 465 470 475 480
 Gln Gln Met Pro Ser Leu Ser Ser Ser Gln Gln Trp Gln Arg Pro Leu
 485 490 495
 Leu Arg Ser Lys Ile Leu Arg Ser Leu Gln Ala Phe Leu Ala Ile Ala
 500 505 510
 Ala Arg Val Phe Ala His Gly Ala Ala Thr Leu Thr Glu Pro Leu Val
 515 520 525
 Pro Thr Ala
 530

<210> 7
 <211> 1580
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (7)..(1569)

<400> 7
 agatct atg tct gca ctt ctg atc cta gct ctt gtt gga gct gca gtt 48
 Met Ser Ala Leu Leu Ile Leu Ala Leu Val Gly Ala Ala Val
 1 5 10

 gct gac tac aaa gac gat gac gac aag ctt ata tgg gaa ctg aag aaa 96
 Ala Asp Tyr Lys Asp Asp Asp Asp Lys Leu Ile Trp Glu Leu Lys Lys
 15 20 25 30

 gat gtt tat gtc gta gaa ttg gat tgg tat ccg gat gcc cct gga gaa 144
 Asp Val Tyr Val Val Glu Leu Asp Trp Tyr Pro Asp Ala Pro Gly Glu
 35 40 45
 atg gtg gtc ctc acc tgt gac acc cct gaa gaa gat ggt atc acc tgg 192
 Met Val Val Leu Thr Cys Asp Thr Pro Glu Glu Asp Gly Ile Thr Trp
 50 55 60

 acc ttg gac cag agc agt gag gtc tta ggc tct ggc aaa acc ctg acc 240
 Thr Leu Asp Gln Ser Ser Glu Val Leu Gly Ser Gly Lys Thr Leu Thr
 65 70 75

 atc caa gtc aaa gag ttt gga gat gct ggc cag tac acc tgt cac aaa 288
 Ile Gln Val Lys Glu Phe Gly Asp Ala Gly Gln Tyr Thr Cys His Lys
 80 85 90

30k

gga ggc gag gtt cta agc cat tcg ctc ctg ctg ctt cac aaa aag gaa	336
Gly Gly Glu Val Leu Ser His Ser Leu Leu Leu Leu His Lys Lys Glu	
95 100 105 110	
gat gga att tgg tcc act gat att tta aag gac cag aaa gaa ccc aaa	384
Asp Gly Ile Trp Ser Thr Asp Ile Leu Lys Asp Gln Lys Glu Pro Lys	
115 120 125	
aat aag acc ttt cta aga tgc gag gcc aag aat tat tct gga cgt ttc	432
Asn Lys Thr Phe Leu Arg Cys Glu Ala Lys Asn Tyr Ser Gly Arg Phe	
130 135 140	
acc tgc tgg tgg ctg acg aca atc agt act gat ttg aca ttc agt gtc	480
Thr Cys Trp Trp Leu Thr Thr Ile Ser Thr Asp Leu Thr Phe Ser Val	
145 150 155	
aaa agc agc aga ggc tct tct gac ccc caa ggg gtg acg tgc gga gct	528
Lys Ser Ser Arg Gly Ser Ser Asp Pro Gln Gly Val Thr Cys Gly Ala	
160 165 170	
gct aca ctc tct gca gag aga gtc aga ggg gac aac aag gag tat gag	576
Ala Thr Leu Ser Ala Glu Arg Val Arg Gly Asp Asn Lys Glu Tyr Glu	
175 180 185 190	
tac tca gtg gag tgc cag gag gac agt gcc tgc cca gct gct gag gag	624
Tyr Ser Val Glu Cys Gln Glu Asp Ser Ala Cys Pro Ala Ala Glu Glu	
195 200 205	
agt ctg ccc att gag gtc atg gtg gat gcc gtt cac aag ctc aag tat	672
Ser Leu Pro Ile Glu Val Met Val Asp Ala Val His Lys Leu Lys Tyr	
210 215 220	
gaa aac tac acc agc agc ttc ttc atc agg gac atc atc aaa cct gac	720
Glu Asn Tyr Thr Ser Ser Phe Phe Ile Arg Asp Ile Ile Lys Pro Asp	
225 230 235	
cca ccc aac aac ttg cag ctg aag cca tta aag aat tct cgg cag gtg	768
Pro Pro Asn Asn Leu Gln Leu Lys Pro Leu Lys Asn Ser Arg Gln Val	
240 245 250	
gag gtc agc tgg gag tac cct gac acc tgg agt act cca cat tcc tac	816
Glu Val Ser Trp Glu Tyr Pro Asp Thr Trp Ser Thr Pro His Ser Tyr	
255 260 265 270	
ttc tcc ctg aca ttc tgc gtt cag gtc cag ggc aag agc aag aga gaa	864
Phe Ser Leu Thr Phe Cys Val Gln Val Gln Gly Lys Ser Lys Arg Glu	
275 280 285	
aag aaa gat aga gtc ttc acc gac aag acc tca gcc acg gtc atc tgc	912
Lys Lys Asp Arg Val Phe Thr Asp Lys Thr Ser Ala Thr Val Ile Cys	
290 295 300	
cgc aaa aat gcc agc att agc gtg cgg gcc cag gac cgc tac tat agc	960
Arg Lys Asn Ala Ser Ile Ser Val Arg Ala Gln Asp Arg Tyr Tyr Ser	
305 310 315	

301

tca tct tgg agc gaa tgg gca tct gtg ccc tgc agt ggt agc ggc tct Ser Ser Trp Ser Glu Trp Ala Ser Val Pro Cys Ser Gly Ser Gly Ser 320 325 330	1008
tct aga ggt gga tca ggc tcc gga ggt agt gga ggt ggg gga tct aag Ser Arg Gly Gly Ser Gly Ser Gly Gly Ser Gly Gly Gly Ser Lys 335 340 345 350	1056
ctt aga gct gtg cct ggg ggc agc agc cct gcc tgg act cag tgc cag Leu Arg Ala Val Pro Gly Gly Ser Ser Pro Ala Trp Thr Gln Cys Gln 355 360 365	1104
cag ctt tca cag aag ctc tgc aca ctg gcc tgg agt gca cat cca cta Gln Leu Ser Gln Lys Leu Cys Thr Leu Ala Trp Ser Ala His Pro Leu 370 375 380	1152
gtg gga cac atg gat cta aga gaa gag gga gat gaa gag act aca aat Val Gly His Met Asp Leu Arg Glu Glu Gly Asp Glu Glu Thr Thr Asn 385 390 395	1200
gat gtt ccc cat atc cag tgt gga gat ggc tgt gac ccc caa gga ctc Asp Val Pro His Ile Gln Cys Gly Asp Gly Cys Asp Pro Gln Gly Leu 400 405 410	1248
agg gac aac agt cag ttc tgc ttg caa agg atc cac cag ggt ctg att Arg Asp Asn Ser Gln Phe Cys Leu Gln Arg Ile His Gln Gly Leu Ile 415 420 425 430	1296
ttt tat gag aag ctg cta gga tcg gat att ttc aca ggg gag cct tct Phe Tyr Glu Lys Leu Leu Gly Ser Asp Ile Phe Thr Gly Glu Pro Ser 435 440 445	1344
ctg ctc cct gat agc cct gtg gcg cag ctt cat gcc tcc cta ctg ggc Leu Leu Pro Asp Ser Pro Val Ala Gln Leu His Ala Ser Leu Leu Gly 450 455 460	1392
ctc agc caa ctc ctg cag cct gag ggt cac cac tgg gag act cag cag Leu Ser Gln Leu Leu Gln Pro Glu Gly His His Trp Glu Thr Gln Gln 465 470 475	1440
att cca agc ctc agt ccc agc cag cca tgg cag cgt ctc ctt ctc cgc Ile Pro Ser Leu Ser Pro Ser Gln Pro Trp Gln Arg Leu Leu Leu Arg 480 485 490	1488
ttc aaa atc ctt cgc agc ctc cag gcc ttt gtg gct gta gcc gcc cgg Phe Lys Ile Leu Arg Ser Leu Gln Ala Phe Val Ala Val Ala Ala Arg 495 500 505 510	1536
gtc ttt gcc cat gga gca gca acc ctg agt ccc taagcggccg c Val Phe Ala His Gly Ala Ala Thr Leu Ser Pro 515 520	1580

<210> 8
 <211> 521
 <212> PRT

30m

<213> Homo sapiens

<400> 8

Met	Ser	Ala	Leu	Leu	Ile	Leu	Ala	Leu	Val	Gly	Ala	Ala	Val	Ala	Asp	1	5	10	15
Tyr	Lys	Asp	Asp	Asp	Asp	Lys	Leu	Ile	Trp	Glu	Leu	Lys	Lys	Asp	Val	20	25	30	
Tyr	Val	Val	Glu	Leu	Asp	Trp	Tyr	Pro	Asp	Ala	Pro	Gly	Glu	Met	Val	35	40	45	
Val	Leu	Thr	Cys	Asp	Thr	Pro	Glu	Glu	Asp	Gly	Ile	Thr	Trp	Thr	Leu	50	55	60	
Asp	Gln	Ser	Ser	Glu	Val	Leu	Gly	Ser	Gly	Lys	Thr	Leu	Thr	Ile	Gln	65	70	75	80
Val	Lys	Glu	Phe	Gly	Asp	Ala	Gly	Gln	Tyr	Thr	Cys	His	Lys	Gly	Gly	85	90	95	
Glu	Val	Leu	Ser	His	Ser	Leu	Leu	Leu	Leu	His	Lys	Lys	Glu	Asp	Gly	100	105	110	
Ile	Trp	Ser	Thr	Asp	Ile	Leu	Lys	Asp	Gln	Lys	Glu	Pro	Lys	Asn	Lys	115	120	125	
Thr	Phe	Leu	Arg	Cys	Glu	Ala	Lys	Asn	Tyr	Ser	Gly	Arg	Phe	Thr	Cys	130	135	140	
Trp	Trp	Leu	Thr	Thr	Ile	Ser	Thr	Asp	Leu	Thr	Phe	Ser	Val	Lys	Ser	145	150	155	160
Ser	Arg	Gly	Ser	Ser	Asp	Pro	Gln	Gly	Val	Thr	Cys	Gly	Ala	Ala	Thr	165	170	175	
Leu	Ser	Ala	Glu	Arg	Val	Arg	Gly	Asp	Asn	Lys	Glu	Tyr	Glu	Tyr	Ser	180	185	190	
Val	Glu	Cys	Gln	Glu	Asp	Ser	Ala	Cys	Pro	Ala	Ala	Glu	Glu	Ser	Leu	195	200	205	
Pro	Ile	Glu	Val	Met	Val	Asp	Ala	Val	His	Lys	Leu	Lys	Tyr	Glu	Asn	210	215	220	
Tyr	Thr	Ser	Ser	Phe	Phe	Ile	Arg	Asp	Ile	Ile	Lys	Pro	Asp	Pro	Pro	225	230	235	240
Asn	Asn	Leu	Gln	Leu	Lys	Pro	Leu	Lys	Asn	Ser	Arg	Gln	Val	Glu	Val	245	250	255	
Ser	Trp	Glu	Tyr	Pro	Asp	Thr	Trp	Ser	Thr	Pro	His	Ser	Tyr	Phe	Ser	260	265	270	
Leu	Thr	Phe	Cys	Val	Gln	Val	Gln	Gly	Lys	Ser	Lys	Arg	Glu	Lys	Lys	275	280	285	

30n

Asp Arg Val Phe Thr Asp Lys Thr Ser Ala Thr Val Ile Cys Arg Lys
 290 295 300

Asn Ala Ser Ile Ser Val Arg Ala Gln Asp Arg Tyr Tyr Ser Ser Ser
 305 310 315 320

Trp Ser Glu Trp Ala Ser Val Pro Cys Ser Gly Ser Gly Ser Ser Arg
 325 330 335

Gly Gly Ser Gly Ser Gly Gly Ser Gly Gly Gly Gly Ser Lys Leu Arg
 340 345 350

Ala Val Pro Gly Gly Ser Ser Pro Ala Trp Thr Gln Cys Gln Gln Leu
 355 360 365

Ser Gln Lys Leu Cys Thr Leu Ala Trp Ser Ala His Pro Leu Val Gly
 370 375 380

His Met Asp Leu Arg Glu Gly Asp Glu Glu Thr Thr Asn Asp Val
 385 390 395 400

Pro His Ile Gln Cys Gly Asp Gly Cys Asp Pro Gln Gly Leu Arg Asp
 405 410 415

Asn Ser Gln Phe Cys Leu Gln Arg Ile His Gln Gly Leu Ile Phe Tyr
 420 425 430

Glu Lys Leu Leu Gly Ser Asp Ile Phe Thr Gly Glu Pro Ser Leu Leu
 435 440 445

Pro Asp Ser Pro Val Ala Gln Leu His Ala Ser Leu Leu Gly Leu Ser
 450 455 460

Gln Leu Leu Gln Pro Glu Gly His His Trp Glu Thr Gln Gln Ile Pro
 465 470 475 480

Ser Leu Ser Pro Ser Gln Pro Trp Gln Arg Leu Leu Leu Arg Phe Lys
 485 490 495

Ile Leu Arg Ser Leu Gln Ala Phe Val Ala Val Ala Ala Arg Val Phe
 500 505 510

Ala His Gly Ala Ala Thr Leu Ser Pro
 515 520

<210> 9
 <211> 711
 <212> DNA
 <213> Mus musculus

<400> 9
 atgacatgc tctcactagc tcctctcctc agccttcttc tcctctgtgt ctctgattct 60
 agggcagaaa caactgtgac ccagtctcca gcatccctgt ccgtggctac aggagaaaaa 120
 gtcactatca gatgcataac cagcactgat attgatgatg atatgatctg gtaccagcag 180

300

```

aagccagggg aacctcctaa gctccttatt tcagaaggca atactcttcg tcctggagtc 240
ccatcccgcgt tctccagcag tggctatggc acagattttg tttttacaat tgaaaacacg 300
ctctcagaag atgttgcaga ttactactgt ttgcaaagtg ataacatgcc tctcacgttc 360
ggtgctggga ccaaggtgga gctgaaacgg gctgatgctg caccaactgt atccatcttc 420
ccaccatcca tggaacagtt aacatctgga ggtgccacag tcgtgtgctt cgtgaacaac 480
ttctatccca gagacatcag tgtcaagtgg aagattgatg gcagtgaaca acgagatggc 540
gtcctggaca gtgttactga tcaggacagc aaagacagca cgtacagcat gagcagcacc 600
ctctcgttga ccaaggttga atatgaaagg cataacctct atacctgtga gggtgttcat 660
aagacatcat cctcaccgcgt cgtcaagagc ttcaacagga atgagtgtta g 711

```

```

<210> 10
<211> 236
<212> PRT
<213> Mus musculus

```

```

<400> 10
Met Thr Met Leu Ser Leu Ala Pro Leu Leu Ser Leu Leu Leu Leu Cys
1          5          10          15

Val Ser Asp Ser Arg Ala Glu Thr Thr Val Thr Gln Ser Pro Ala Ser
          20          25          30

Leu Ser Val Ala Thr Gly Glu Lys Val Thr Ile Arg Cys Ile Thr Ser
          35          40          45

Thr Asp Ile Asp Asp Asp Met Ile Trp Tyr Gln Gln Lys Pro Gly Glu
          50          55          60

Pro Pro Lys Leu Leu Ile Ser Glu Gly Asn Thr Leu Arg Pro Gly Val
65          70          75          80

Pro Ser Arg Phe Ser Ser Ser Gly Tyr Gly Thr Asp Phe Val Phe Thr
          85          90          95

Ile Glu Asn Thr Leu Ser Glu Asp Val Ala Asp Tyr Tyr Cys Leu Gln
          100          105          110

Ser Asp Asn Met Pro Leu Thr Phe Gly Ala Gly Thr Lys Val Glu Leu
          115          120          125

Lys Arg Ala Asp Ala Ala Pro Thr Val Ser Ile Phe Pro Pro Ser Met
          130          135          140

Glu Gln Leu Thr Ser Gly Gly Ala Thr Val Val Cys Phe Val Asn Asn
145          150          155          160

```

30p

Phe Tyr Pro Arg Asp Ile Ser Val Lys Trp Lys Ile Asp Gly Ser Glu
 165 170 175

Gln Arg Asp Gly Val Leu Asp Ser Val Thr Asp Gln Asp Ser Lys Asp
 180 185 190

Ser Thr Tyr Ser Met Ser Ser Thr Leu Ser Leu Thr Lys Val Glu Tyr
 195 200 205

Glu Arg His Asn Leu Tyr Thr Cys Glu Val Val His Lys Thr Ser Ser
 210 215 220

Ser Pro Val Val Lys Ser Phe Asn Arg Asn Glu Cys
 225 230 235

<210> 11
 <211> 1406
 <212> DNA
 <213> Mus musculus

<400> 11
 atgggatgga gctgtatcat cctcttcttg gtagcaacag ctacaggtgt ccactcccag 60
 gtccaactgc agcagcctgg ggctgagctg gtgaggcctg gggcttcagg gaagctgtcc 120
 tgcaaggctt ctggctacac cttcaccagc tactggatga actgggtgaa gcagaggcct 180
 ggacaaggcc ttgaatggat tggatatgatt gatccttttag acagtgaaac tcactataat 240
 caaatgttca aggacaaggc cacattgact gtagacaaat cctccagcac agcctacatg 300
 cagctcagca gcctgacatc tgaggactat gcggtctatt actgtgcaag aggggataac 360
 tactatgcta tggactactg ggggtcaagga acctcagtca ccgtctcctc agccaaaacg 420
 acacccccat ctgtctatcc actggcccct ggatctgctg cccaaactaa ctccatggtg 480
 accctgggat gcctgggtcaa gggctatttc cctgagccag tgacagtgac ctggaactct 540
 ggatccctgt ccagcgggtgt gcacaccttc ccagctgtcc tgcagtctga cctctacact 600
 ctgagcagct cagtgactgt cccctccagc acctggccca gcgagaccgt cacctgcaac 660
 gttgcccacc cggccagcag caccaagggtg gacaagaaaa ttgtgcccag ggattgtggt 720
 tgtaagcctt gcatatgtac agtcccagaa gtatcatctg tcttcatctt cccccaaag 780
 cccaaggatg tgctcaccat tactctgact cctaagggtca cgtgtgttgt ggtagacatc 840
 agcaaggatg atccccgaggt ccagttcagc tggttttag atgatgtgga ggtgcacaca 900
 gctcagacgc aaccccggga ggagcagttc aacagcactt tccgctcagt cagtgaactt 960
 cccatcatgc accaggactg gctcaatggc aaggagttca aatgcagggt caacagtgca 1020

30q

```

gcttttcctg ccccatcga gaaaaccatc tccaaaacca aaggcagacc gaaggctcca 1080
caggtgtaca ccattccacc tcccaaggag cagatggcca aggataaagt cagtctgacc 1140
tgcatgataa cagacttctt ccctgaagac attactgtgg agtggcagtg gaatgggcag 1200
ccagcggaga actacaagaa cactcagccc atcatggact cagatggctc ttacttcgtc 1260
tacagcaagc tcaatgtgca gaagagcaac tgggaggcag gaaatacttt cacctgctct 1320
gtgttacatg agggcctgca caaccaccat actgagaaga gcctctccca ctctcctggt 1380
aatgatccc agagtccagt ggcccc 1406

```

```

<210> 12
<211> 467
<212> PRT
<213> Mus musculus

```

```

<400> 12
Met Gly Trp Ser Cys Ile Ile Leu Phe Leu Val Ala Thr Ala Thr Gly
1          5          10          15

Val His Ser Gln Val Gln Leu Gln Gln Pro Gly Ala Glu Leu Val Arg
20          25          30

Pro Gly Ala Ser Gly Lys Leu Ser Cys Lys Ala Ser Gly Tyr Thr Phe
35          40          45

Thr Ser Tyr Trp Met Asn Trp Val Lys Gln Arg Pro Gly Gln Gly Leu
50          55          60

Glu Trp Ile Gly Met Ile Asp Pro Leu Asp Ser Glu Thr His Tyr Asn
65          70          75          80

Gln Met Phe Lys Asp Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Ser
85          90          95

Thr Ala Tyr Met Gln Leu Ser Ser Leu Thr Ser Glu Asp Tyr Ala Val
100         105         110

Tyr Tyr Cys Ala Arg Gly Asp Asn Tyr Tyr Ala Met Asp Tyr Trp Gly
115         120         125

Gln Gly Thr Ser Val Thr Val Ser Ser Ala Lys Thr Thr Pro Pro Ser
130         135         140

Val Tyr Pro Leu Ala Pro Gly Ser Ala Ala Gln Thr Asn Ser Met Val
145         150         155         160

Thr Leu Gly Cys Leu Val Lys Gly Tyr Phe Pro Glu Pro Val Thr Val
165         170         175

Thr Trp Asn Ser Gly Ser Leu Ser Ser Gly Val His Thr Phe Pro Ala
180         185         190

```

[illegible]

What is claimed is:

1. Use of an IL-23 complex for treating a cutaneous wound, said IL-23 complex comprising:

a) a polypeptide comprising residues 352 to 521 of SEQ ID NO: 8 or a conservatively modified variant thereof comprising an individual amino acid substitution; and

b) a polypeptide comprising residues 25 to 330 of SEQ ID NO: 8 or a conservatively modified variant thereof comprising an individual amino acid substitution; wherein the IL-23 complex increases the expression of IL-17 at least 2-fold in excisional wounds on the backs of C57Bl/6NT mice.

2. Use of an IL-23 complex in the manufacture of a medicament for treating a cutaneous wound, said IL-23 complex comprising:

a) a polypeptide comprising residues 352 to 521 of SEQ ID NO: 8 or a conservatively modified variant thereof comprising an individual amino acid substitution; and

b) a polypeptide comprising residues 25 to 330 of SEQ ID NO: 8 or a conservatively modified variant thereof comprising an individual amino acid substitution; wherein the IL-23 complex increases the expression of IL-17 at least 2-fold in excisional wounds on the backs of C57Bl/6NT mice.

3. The use of claim 1 or 2, wherein the wound is an incision wound, an excision wound, an ulcer or a graft.

4. The use of any one of claims 1-3, wherein treatment with the IL-23 complex increases:

- a) the pressure required to break a healed or healing wound;
- b) stiffness of a healed or healing wound;
- c) healing rate of the wound;
- d) granulation layer thickness of a healed or healing wound;
- e) antimicrobial activity in the healed or healing wound; or
- f) angiogenesis.

5. The use of any one of claims 1-4, wherein treatment with the IL-23 complex increases recruitment of a cell to the wound, and wherein the cell is:

- a) a CD11b⁺ MHC class II cell;

- b) a monocyte/macrophage; or
- c) a CD31+ endothelial cell.

6. The use of any one of claims 1-5, wherein the polypeptide comprising residues 25 to 330 of SEQ ID NO: 8 has no amino acid substitutions.

7. The use of claim 6, wherein the individual amino acid substitution in the polypeptide comprising residues 352 to 521 of SEQ ID NO: 8 is the exchange of one small amino acid for another small amino acid, wherein the small amino acids are Gly, Ala and Ser.

8. Use of an IL-23 fusion protein for treating a cutaneous wound, said IL-23 fusion protein comprising:

- a) residues 352 to 521 of SEQ ID NO: 8 or a conservatively modified variant thereof comprising an individual amino acid substitution; and
- b) residues 25 to 330 of SEQ ID NO: 8 or a conservatively modified variant thereof comprising an individual amino acid substitution; wherein the IL-23 fusion protein increases the expression of IL-17 at least 2-fold in excisional wounds on the backs of C57Bl/6NT mice.

9. Use of an IL-23 fusion protein in the manufacture of a medicament for treating a cutaneous wound, said IL-23 fusion protein comprising:

- a) residues 352 to 521 of SEQ ID NO: 8 or a conservatively modified variant thereof comprising an individual amino acid substitution; and
- b) residues 25 to 330 of SEQ ID NO: 8 or a conservatively modified variant thereof comprising an individual amino acid substitution; wherein the IL-23 fusion protein increases the expression of IL-17 at least 2-fold in excisional wounds on the backs of C57Bl/6NT mice.

10. The use of claim 8 or 9, wherein the IL-23 fusion protein comprises residues 25 to 521 of SEQ ID NO: 8 or a conservatively modified variant thereof comprising an individual amino acid substitution.

11. The use of any one of claims 8-10, wherein the wound is an incision wound, an excision wound, an ulcer or a graft.

12. The use of any one of claims 8-11, wherein treatment with the IL-23 fusion protein increases:

- a) the pressure required to break a healed or healing wound;
- b) stiffness of a healed or healing wound;
- c) healing rate of the wound;
- d) granulation layer thickness of a healed or healing wound;
- e) antimicrobial activity in the healed or healing wound; or
- f) angiogenesis.

13. The use of any one of claims 8-12, wherein treatment with the IL-23 fusion protein increases the recruitment of a cell to the wound, and wherein the cell is:

- a) a CD11b+ MHC class II cell;
- b) a monocyte/macrophage; or
- c) a CD31+ endothelial cell.

14. The use of any one of claims 8-13, wherein residues 25 to 330 of SEQ ID NO: 8 have no amino acid substitutions.

15. The use of claim 14, wherein the individual amino acid substitution in residues 352 to 521 of SEQ ID NO: 8 is the exchange of one small amino acid for another small amino acid, wherein the small amino acids are Gly, Ala and Ser.

16. Use of an IL-23 complex for treating a cutaneous wound, said IL-23 complex comprising:

- a) a polypeptide comprising residues 352 to 521 of SEQ ID NO: 8; and
- b) a polypeptide comprising residues 25 to 330 of SEQ ID NO: 8.

17. Use of an IL-23 complex in the manufacture of a medicament for treating a cutaneous wound, said IL-23 complex comprising:

- a) a polypeptide comprising residues 352 to 521 of SEQ ID NO: 8; and
- b) a polypeptide comprising residues 25 to 330 of SEQ ID NO: 8.

18. Use of an IL-23 fusion protein for treating a cutaneous wound, said IL-23 fusion protein comprising:

- a) residues 352 to 521 of SEQ ID NO: 8; and
- b) residues 25 to 330 of SEQ ID NO: 8.

19. Use of an IL-23 fusion protein in the manufacture of a medicament for treating a cutaneous wound, said IL-23 fusion protein comprising:

- a) residues 352 to 521 of SEQ ID NO: 8; and
- b) residues 25 to 330 of SEQ ID NO: 8.