



US 20110206776A1

(19) **United States**
(12) **Patent Application Publication**
Tom et al.

(10) **Pub. No.: US 2011/0206776 A1**
(43) **Pub. Date: Aug. 25, 2011**

(54) **METHODS OF MANUFACTURE OF IMMUNOCOMPATIBLE AMNIOTIC MEMBRANE PRODUCTS**

filed on Feb. 18, 2010, provisional application No. 61/369,562, filed on Jul. 30, 2010.

Publication Classification

(76) Inventors: **Samson Tom**, Baltimore, MD (US); **Alla Danilkovitch**, Columbia, MD (US); **Dana Yoo**, Sykesville, MD (US); **Timothy Jansen**, Baltimore, MD (US); **Jin-Qiang Kuang**, Woodstock, MD (US); **Jennifer Michelle Marconi**, Glen Burnie, MD (US)

(51) **Int. Cl.**
A61K 35/50 (2006.01)
A61P 17/02 (2006.01)
A01N 1/02 (2006.01)
(52) **U.S. Cl.** **424/583; 435/1.3**

(21) Appl. No.: **13/030,562**
(22) Filed: **Feb. 18, 2011**

Related U.S. Application Data

(60) Provisional application No. 61/338,464, filed on Feb. 18, 2010, provisional application No. 61/338,489,

(57) **ABSTRACT**
Provided herein is a placental product comprising an immunocompatible amniotic membrane. Such placental products can be cryopreserved and contain viable therapeutic cells after thawing. The placental product of the present invention is useful in treating a patient with a tissue injury (e.g. wound or burn) by applying the placental product to the injury. Similar application is useful with ligament and tendon repair and for engraftment procedures such as bone engraftment.






FIGURE 1A
Freezing - Styrofoam Box vs. Freezer Shelf




FIGURE 1B
Freezing - Styrofoam Box vs. Freezer Shelf





FIGURE 1C
Freezing - Styrofoam Box vs. Freezer Shelf






FIGURE 1E
Freezing - Styrofoam Box vs. Freezer Shelf




FIGURE 1F
Freezing- Styrofoam Box vs. Freezer Shelf




FIGURE 1G

Freezing - Styrofoam Box vs. Freezer Shelf




FIGURE 1H

Freezing - Styrofoam Box vs. Freezer Shelf




FIGURE 2




FIGURE 3




FIGURE 4




FIGURE 5




FIGURE 6




FIGURE 7

Expression of IL-2R on T-cells stimulated by Placental Derived Cells (donor 37 fresh)




FIGURE 8




FIGURE 9A




FIGURE 9B




FIGURE 9C




FIGURE 10A




FIGURE 10B




FIGURE 10C




FIGURE 11




FIGURE 12




FIGURE 13A




FIGURE 13B




FIGURE 13C




FIGURE 14




FIGURE 15A

Expression of bone reparative proteins in amniotic membrane homogenates




FIGURE 15B

Expression of IGF-1 in amniotic membrane homogenates




FIGURE 16




FIGURE 17




FIGURE 19




FIGURE 20A



FIGURE 20B



FIGURE 20C



FIGURE 20D




FIGURE 20E



FIGURE 21A




FIGURE 21B




FIGURE 21C




FIGURE 21D




FIGURE 21E



METHODS OF MANUFACTURE OF IMMUNOCOMPATIBLE AMNIOTIC MEMBRANE PRODUCTS

RELATED APPLICATIONS

[0001] This application claims priority to:

[0002] U.S. Provisional Applications Ser. No. 61/338, 464 entitled "Selectively Immunodepleted Chorionic Membranes", filed on Feb. 18, 2010 bearing Docket No. 22924US01,

[0003] U.S. Provisional Applications Ser. No. 61/338, 489 entitled "Selectively Immunodepleted Amniotic Membranes", filed on Feb. 18, 2010 bearing Docket No. 22925US01, and

[0004] U.S. Provisional Applications Ser. No. 61/369, 562 entitled "Therapeutic Products Comprising Vitalized Placental Dispersions" filed on Jul. 30, 2010 bearing Docket No. 23498US01, the contents of which are hereby incorporated by reference in their entireties.

[0005] This application is being co-filed on Feb. 18, 2011 with, and incorporates by reference, applications entitled:

[0006] "Methods of Manufacture of Immunocompatible Chorionic Membrane Products",

[0007] "Immunocompatible Chorionic Membrane Products",

[0008] "Immunocompatible Amniotic Membrane Products",

[0009] "Therapeutic Products Comprising Vitalized Placental Dispersions", and

[0010] "Methods of Manufacture of Therapeutic Products Comprising Vitalized Placental Dispersions"

FIELD OF THE INVENTION

[0011] The present technology relates to products to facilitate wound healing such as placenta membrane-derived products and biologic skin. The present technology relates to products to protect injured or damaged tissue, or as a covering to prevent adhesions, to exclude bacteria, to inhibit bacterial activity, or to promote healing or growth of tissue. An example of such a placental membrane is an amniotic membrane. The field also relates to methods of manufacturing and methods of use of such membrane-derived products.

BACKGROUND OF THE INVENTION

[0012] Fresh or decellularized placental membranes have been used typically in surgical applications since at least 1910 when Johns Hopkins Hospital reported the use of placental membrane for dermal applications. Subsequently unseparated amnion and chorion were used as skin substitutes to treat burned or ulcerated surfaces. During the 1950's and 60's Troensegaard-Hansen applied boiled amniotic membranes to chronic leg ulcers.

[0013] The human amniotic membrane (AM) is the innermost of the fetal membranes deriving from the amniotic sac and constituting the lining of the amniotic cavity. It is approximately 0.02 to 0.5 mm thick. The AM consists of five layers: a thin layer rests on the basement membrane and contacts the amniotic fluid, an underlying layer of connective tissue attaching the basement membrane that consists of three layers: a compact layer, a layer of fibroblast, and a spongy layer. The spongy layer is adjacent to the cellular layer of the chorion. The amnion is essentially devoid of vasculature.

[0014] Both fresh and frozen AMs have been used for wound healing therapy. When fresh AM is used, there is increased risk of disease transmission. According to published reports, fresh amniotic tissue exhibits cell viability of 100%, however within 28 days of storage above 0° C. diminished cell viability to 15 to 35%. Freezing over a time of 3 weeks reduced cell viability to 13 to 18%, regardless of the temperature or medium.

[0015] Lee and Tseng report the successful cryopreservation of AM in glycerol and Dulbeccos Modified Eagle medium (DMEM) at -80° C., although such cryopreservation dramatically decreases cell viability. The cryopreservation of AM in glycerol and DMEM is recommended by the FDA. According to published reports, glycerol storage of AM resulted in immediate cell death. Glycerol cryopreserved AM (-80° C.) and glycerol-preserved AM (-4° C.) are sufficient to provide a matrix for wound healing, but fail to provide sufficient cell viability to bestow biological effectiveness

[0016] Gajiwala and Gajiwala report the successful preservation of AM by freeze-drying (lyophilisation) and gamma-irradiation. According to this method, AM is pasteurized at 60° C., treated with 70% ethanol, and freeze-dried to remove most of the remaining moisture. Then the AM is sterilized by exposure of 25 kGy gamma-radiation in a Cobalt 60 Gamma chamber unit or at an ISO-certified radiation plant. The sterilized AM can be stored at room temperature for a short period (up to 6 months).

[0017] Gomes reports preservation of AM with lyophilisation followed by sterilization in ethylene oxide.

[0018] Rama et al reported the cryopreservation of AM in 10% dimethyl sulfoxide (DMSO) instead of glycerol and achieved a cell viability of about 40%.

[0019] Two placental tissue graft products containing living cells, Apligraf and Dermagraft, are currently commercially available. Both Apligraf and Dermagraft comprise ex vivo cultured cells. Neither Apligraf nor Dermagraft comprise stem cells. Furthermore, neither Apligraf nor Dermagraft comprise Insulin-like Growth Factor Binding Protein-1 (IGFBP-1) and adiponectin, which are key factors in the natural wound healing process. In addition, neither Apligraf nor Dermagraft exhibit a protease-to-protease inhibitor ratio favorable for wound healing. As wound healing is a multifactorial biological process, many factors are needed to properly treat a wound; products having non-native cellular populations are less capable of healing wounds relative to a product having an optimal population of cells representing the native array. It would represent an advance in the art to provide a chorion-derived biologic skin substitute comprising a population of cells representing the native array of factors, including, for example, growth factors and cytokines.

[0020] Apligraf is a living, bi-layered skin substitute manufactured using neonatal foreskin keratinocytes and fibroblasts with bovine Type I collagen. As used in this application, Apligraf refers to the product available for commercial sale in November 2009.

[0021] Dermagraft is cryopreserved human fibroblasts derived from newborn foreskin tissue seeded on extracellular matrix. According to its product literature, Dermagraft requires three washing steps before use which limits the practical implementation of Dermagraft as a wound dressing relative to products that require less than three washing steps. As used in this application, Dermagraft refers to the product available for commercial sale in November 2009.

[0022] Engineered wound dressings such as Apligraf and Dermagraft do not provide the best potential for wound healing because they comprise sub-optimal cellular compositions and therefore do not provide proper wound healing. For example, neither Apligraf nor Dermagraft comprises stem cells and, as a result, the ratio between different factors secreted by cells does not enable efficient wound healing. Additionally, some factors that are important for wound healing, including EGF, IGFBP1, and adiponectin, are absent from both Apligraf and Dermagraft. Additionally, some factors, including MMPs and TIMPs, are present in proportions that differ greatly from the proportions found in the natural wound healing process; this difference significantly alters, among other things, upstream inflammatory cytokine pathways which in turn allows for sub-optimal micro-environments at the wound site.

[0023] Paquet-Fifield et al. report that mesenchymal stem cells and fibroblasts are important for wound healing (J Clin Invest, 2009, 119: 2795). No product has yet been described that comprise mesenchymal stem cells and fibroblasts.

[0024] Both MMPs and TIMPs are among the factors that are important for wound healing. However, expression of these proteins must be highly regulated and coordinated. Excess of MMPs versus TIMPs is a marker of poor chronic wound healing (Liu et al, Diabetes Care, 2009, 32: 117; Mwaura et al, Eur J Vasc Endovasc Surg, 2006, 31: 306; Trengove et al, Wound Rep Reg, 1999, 7: 442; Vaalamo et al, Hum Pathol, 1999, 30: 795).

[0025] α 2-macroglobulin is known as a plasma protein that inactivates proteinases from all 4 mechanistic classes: serine proteinases, cysteine proteinases, aspartic proteinases, and metalloproteinases (Borth et al., FASEB J, 1992, 6: 3345; Baker et al., J Cell Sci, 2002, 115:3719). Another important function of this protein is to serve as a reservoir for cytokines and growth factors, examples of which include TGF, PDGF, and FGF (Asplin et al, Blood, 2001, 97: 3450; Huang et al, J Biol Chem, 1988; 263: 1535). In chronic wounds like diabetic ulcers or venous ulcers, the presence of high amount of proteases leads to rapid degradation of growth factors and delays in wound healing. Thus, a placental membrane wound dressing comprising α 2-macroglobulin would constitute an advance in the art.

[0026] An in vitro cell migration assay is important for assessing the wound healing potential of a skin substitute. The process of wound healing is highly complex and involves a series of structured events controlled by growth factors (Goldman, Adv Skin Wound Care, 2004, 1:24). These events include increased vascularization, infiltration by inflammatory immune cells, and increases in cell proliferation. The beginning stages of wound healing revolve around the ability of individual cells to polarize towards the wound and migrate into the wounded area in order to close the wound area and rebuild the surrounding tissue. An assay capable of evaluating the wound healing potential of skin substitutes by examining the correlation between cell migration and wound healing would represent an advance in the art.

SUMMARY OF THE INVENTION

[0027] The present invention provides a pharmaceutically acceptable placental product.

[0028] A placental product according to the present invention comprises an immunocompatible amniotic membrane in a cryopreservation medium (optionally cryopreserved) and viable native therapeutic cells and native therapeutic factors.

[0029] In some embodiments, the amniotic membrane of the placental product is selectively devitalized.

[0030] There is now provided a placental product that is selectively depleted of substantially all immunogenic cells.

[0031] There is now provided a placental product that does not contain ex vivo cultured cells.

[0032] In some embodiments, the placental product further comprises a chorionic membrane that is selectively devitalized.

[0033] There is now provided a placental product that comprises at least one of Epidermal Growth Factor, IGFBP1, and Adiponectin.

[0034] Optionally, the therapeutic factors include one or more of IGFBP1, adiponectin, α 2-macroglobulin, bFGF, and EGF. Optionally, the therapeutic factors include MMP-9 and TIMP1, wherein the ratio of MMP-9:TIMP1 is from about 7 to about 10. Optionally, the therapeutic factors include IGFBP1, adiponectin, α 2-macroglobulin, bFGF, EGF, MMP-9, and TIMP1. Optionally, the therapeutic factors include IGFBP1, adiponectin, α 2-macroglobulin, bFGF, MMP-9, and TIMP1, wherein the ratio of MMP-9:TIMP1 is from about 7 to about 10 to one. Optionally, the therapeutic factor is present in a substantial amount in comparison to unprocessed human placental membrane. Optionally, each placental product embodiment optionally is devoid of ex-vivo expanded cultured cells.

[0035] The present invention also provides a method of manufacturing a placental product comprising: obtaining a placenta, wherein the placenta comprises an amniotic membrane, selectively depleted of immunogenicity, and cryopreserving the placenta, thereby providing a placental product. According to the present invention, the step of selective depletion comprises removing immunogenic cells (e.g. CD14+ macrophages and vascularized tissue-derived cells) and/or immunogenic factors (e.g. TNF α). Optionally, the step of selective depletion comprises removing CD14+ macrophages by refrigerating the placental product for a period of time (e.g. about 30-60 mins.) at a temperature above freezing (e.g. at 2-8° C.), and then freezing, whereby CD14+ macrophages are selectively killed relative to therapeutic cells.

[0036] The present invention also provides a method of screening a placental product for therapy comprising assaying the placental product for immunogenicity and/or therapeutic value. Optionally, the step of assaying the placental product for immunogenicity comprises a Mixed Lymphocyte Reaction (MLR) and/or Lipopolysaccharide (LPS)-induced Tumor Necrosis Factor (TNF)- α secretion. Optionally, the step of assaying the placental product for therapeutic value comprises assaying the placental product for cell migration induction.

[0037] The present invention also provides a method of treating a subject comprising administering a placental product to the subject. Optionally, the step of administering comprises applying the placental product to a wound, for example, topically applying the placental product to a skin wound. In one embodiment, a placental product is used in a tendon or ligament surgery to promote healing of a tendon or ligament.

[0038] The present inventors have identified a need for the development of amniotic membrane products comprising at least one of IGFBP1, and adiponectin, providing superior wound healing properties.

BRIEF DESCRIPTION OF THE DRAWINGS

[0039] FIG. 1 depicts freezing rates of various freezing methods.

[0040] FIG. 2 depicts process cell recovery for amniotic membrane as a function of cryo volume.

[0041] FIG. 3 depicts process cell recovery for chorionic membrane as a function of cryo volume.

[0042] FIG. 4 depicts the effects of refrigeration time and freezing parameters on process (cryopreservation) cell recovery for the amniotic membrane.

[0043] FIG. 5 depicts the effects of refrigeration time and freezing parameters on process (cryopreservation) cell recovery for the chorionic membrane.

[0044] FIG. 6 shows representative images of the live/dead staining of the epithelial layer of fresh amniotic membrane.

[0045] FIG. 7 depicts expression of IL-2sR from T-cells stimulated by placental derived cells from various membrane preparations.

[0046] FIG. 8 depicts expression of IL-2sR from T-cells stimulated by placental derived cells from various membrane preparations after cryopreservation.

[0047] FIG. 9 depicts LPS stimulated TNF a released from various membrane preparations.

[0048] FIG. 10 shows images of cultured cells isolated from various membrane preparations.

[0049] FIG. 11 depicts a correlation between IL-2sR release in an MLR assay of various membrane preparations and the number of CD45+ cells.

[0050] FIG. 12 depicts the IL-2sR release in an MLR assay of amniotic membrane preparations and the number of CD45+ cells.

[0051] FIG. 13 depicts expression of EGF (A), IGFBP1 (B), and Adiponectin (C) in amniotic and/or chorionic membranes.

[0052] FIG. 14 depicts expression of IFN-2 α and TGF- β 3 in amniotic membrane homogenates.

[0053] FIG. 15 depicts expression of BMP-2, BMP-4, PLAB, PIGF (A), and IGF-1 (B) in amniotic membrane homogenates.

[0054] FIG. 16 depicts the ratio of MMPs to TIMPs in various membrane products.

[0055] FIG. 17 depicts expression of EGF in chorion and amnion membranes measured by ELISA in two separate placenta donors.

[0056] FIG. 18 depicts the Cell Biolabs 24-well Cytoselect wound healing assay.

[0057] FIG. 19 depicts representative images of HMVECs treated with 5% conditioned media from amniotic, chorionic, or a combination of amniotic/chorionic preparations.

[0058] FIG. 20 depicts the remarkable efficacy of placental products for treating diabetic foot ulcers in patient 1.

[0059] FIG. 21 depicts the remarkable efficacy of placental products for treating diabetic foot ulcers in patient 2.

DETAILED DESCRIPTION OF THE INVENTION

[0060] As used herein, the following definitions apply:

[0061] “Exemplary” (or “e.g.” or “by example”) means a non-limiting example.

[0062] “hCMSCs” means human chorionic membrane stromal cells. hCMSCs are generally positive for CD73?, CD70, CD90, CD105, and CD166; negative for CD45 and CD34. hCMSCs differentiate to mesodermal lineages (osteogenic, chondrogenic, and adipogenic).

[0063] “Selective depletion of immunogenicity” or “selective depletion of immunogenic cells or factors” or “selective depletion” means a placental product that retains live therapeutic cells and/or retains therapeutic efficacy for the treatment of tissue injury yet is free, substantially free, or depleted of at least one of immune cell type (e.g. CD14+ macrophages, trophoblasts, and/or vascular-tissue derived cells) and/or immunogenic factor that are otherwise present in a native placenta or amniotic membrane.

[0064] “MSC” means mesenchymal stem cells and include fetal, neonatal, adult, or post-natal. “MSCs” include amniotic MSCs (AMSCs). MSCs generally express one or more of CD73, CD90, CD105, and CD166.

[0065] “Native cells” means cells that are native, resident, or endogenous to the placental membrane, i.e. cells that are not exogenously added to the placental membrane.

[0066] “Native factors” means placental membrane factors that are native, resident, or endogenous to the placental membrane, i.e. factors that are not exogenously added to the placental membrane.

[0067] “Placental products” means the instant placental products disclosed and claimed herein.

[0068] “Substantial amount” of an element of the present invention, e.g. native factors, native cells, therapeutic factors, or selective depletion, means a value at least about 10% or more in comparison to an unprocessed, not cryopreserved, fresh membrane sample. A substantial amount can optionally be at least about 50%.

[0069] “Therapeutic cells” or “beneficial cells” means stromal cells, MSCs, fibroblasts, and/or epithelial cells.

[0070] “Therapeutic factors” means placenta- or amniotic membrane-derived factors that promote wound healing. Examples include IGFBP1, adiponectin, α 2-macroglobulin, and/or bFGF. Other examples include MMP-9 and TIMP1.

[0071] “Stromal cells” refers to a mixed population of cells present (optionally in native proportions) composed of neonatal mesenchymal stem cells and neonatal fibroblasts. Both neonatal mesenchymal stem cells and neonatal fibroblasts are immunoprivileged; neither express surface proteins present on immunogenic cell types.

[0072] In some embodiments, the present technology discloses placental products for clinical use, including use in wound healing such as diabetic foot ulcers, venous leg ulcers, and burns. The manufacturing process optionally eliminates essentially all potentially immunogenic cells from the placental membrane while preserving of specific cells that play an important role in wound healing.

[0073] In some embodiments, the present technology discloses a placental product that is selectively devitalized. There is now provided a placental product that is selectively depleted of substantially all immunogenic cells. There is now provided a placental product that does not contain ex vivo cultured cells. There is now provided a placental product that comprises at least one of IGFBP1, and adiponectin. There is now provided a placental product that comprises Epidermal Growth Factor, IGFBP1. There is now provided a placental product that comprises adiponectin.

[0074] In some embodiments, the present technology discloses a method of cryopreserving a placental product that preserves the viability of specific beneficial cells that are the primary source of factors for the promotion of healing to the

wound healing process while selectively depleting immunogenic cells (e.g. killing or rendering non-immunogenic).

[0075] In some embodiments, the present technology discloses a bioassay to test immunogenicity of manufactured placental products.

[0076] In some embodiments, the present technology discloses a placental product exhibiting a therapeutic ratio of MMP to TIMP comparable to that exhibited in vivo. The present inventors have identified a need for the development of placental products exhibiting a ratio of MMP-9 and TIMP1 of about 7-10 to one.

[0077] In some embodiments, the present technology discloses a placental product that comprises α 2-macroglobulin.

[0078] There is now provided a placental product that inactivates substantially all serine proteinases, cysteine proteinases, aspartic proteinases, and metalloproteinases present in the amniotic membrane. There is now provided a placental product that inactivates substantially all serine proteinases present in the amniotic membrane. There is now provided a placental product that inactivates substantially all cysteine proteinases present in the amniotic membrane. There is now provided a placental product that inactivates substantially all aspartic proteinases present in the amniotic membrane. There is now provided a placental product that inactivates substantially all metalloproteinases present in the amniotic membrane.

[0079] In some embodiments, the present technology discloses a placental product that comprises bFGF.

[0080] In some embodiments, the present technology discloses a placental product exhibiting a protease-to-protease inhibitor ratio favorable for wound healing.

[0081] In some embodiments, the present technology discloses a cell migration assay capable of evaluating the wound-healing potential of a placental product.

[0082] IGFBP1 and adiponectin are among the factors that are important for wound healing. Evaluation of proteins derived from placental products prepared according to the presently disclosed technology reveal that EGF is one of the major factors secreted in higher quantities by these tissues. Additionally, the importance of EGF for wound healing together with high levels of EGF detected in the presently disclosed amniotic membranes support selection of EGF as a potency marker for evaluation of membrane products manufactured for clinical use pursuant to the present disclosure.

[0083] The present technology discloses a cryopreservation procedure for a placental product that selectively depletes immunogenic cells and preserves the viability of other beneficial cells (including at least one or 2 or all of mesenchymal stem cells, epithelial cells and fibroblasts. In some embodiments, the beneficial cells are the primary source of factors for the promotion of healing.

[0084] Placental products, their usefulness, and their immunocompatibility are surprisingly enhanced by depletion of maternal trophoblast and selective elimination of CD14+ fetal macrophages. Immunocompatibility can be demonstrated by any means commonly known by the skilled artisan, such demonstration can be performed by the mixed Lymphocyte Reaction (MLR) and by lipopolysaccharide (LPS)-induced Tumor Necrosis Factor (TNF)- α secretion.

[0085] The instant placental products contain bFGF optionally at a substantial concentration.

[0086] The instant placental products optionally secrete factors that stimulate cell migration and/or wound healing.

The presence of such factors can be demonstrated by any commonly recognized method,

[0087] For example, commercially available wound healing assays exist (Cell Biolabs) and cell migration can be assessed by cell line HMVEC (Lonza Inc.). In one embodiment, conditioned medium from the present placental products enhance cell migration.

[0088] The placental products disclosed herein are useful in treating a number of wounds including: tendon repair, cartilage repair (e.g. femoral condyle, tibial plateau), ACL replacement at the tunnel/bone interface, dental tissue augmentation, fistulas (e.g. Crohn's disease, G-tube, tracheoesophageal), missing tissue at adhesion barriers (e.g. nasal septum repair, vaginal wall repair, abdominal wall repair, tumor resection), dermal wounds (e.g. partial thickness burns, toxic epidermal necrolysis, epidermolysis bullosa, pyoderma gangrenosum, ulcers e.g. diabetic ulcers (e.g. foot), venous leg ulcers), surgical wounds, hernia repair, tendon repair, bladder repair, periosteum replacement, keloids, organ lacerations, epithelial defects, and repair or replacement of a tympanic membrane.

[0089] The placental products disclosed herein exhibit one or more of the following properties beneficial to the wound healing process:

[0090] a. an epithelial cell layer, wherein the approximate number of cells per cm² of the amniotic membrane is about 10,000 to about 360,000 or about 40,000 to about 90,000.

[0091] b. a thick basement membrane (comprising one or more of Collagen Type I, III, IV, laminin, and fibronectin),

[0092] c. a stromal cell layer;

[0093] d. an amniotic membrane with a thickness of about 20 to about 50 μ m,

[0094] e. high thrombin activity,

[0095] f. low immunogenicity,

[0096] g. cryopreserved/cryopreserveable,

[0097] h. amniotic MSC,

[0098] i. analgesic effect

[0099] j. reduces scarring,

[0100] k. anti-inflammatory proteins such as IL-1a and IL-10,

[0101] l. anti-inflammatory suppression of lymphocyte reactivity in vitro, for example by inhibition of CD8+ and CD4+ proliferation or increased CD4+ Treg cells,

[0102] m. antibacterial proteins such as defensins and allantoin (bacteriolytic proteins),

[0103] n. angiogenic and mitogenic factors that promote re-epithelialization such as EGF, HGF, and VEGF,

[0104] o. cells that are positive for CD70, CD90, CD105, and CD166 and negative for CD45 and CD34,

[0105] p. cells that express HLA-G,

[0106] q. cells that express a placenta specific MHC-1 antigen important to immune tolerance that inhibits both NK cytotoxicity and T-cell-mediated cytotoxicity and activation of immune cells (e.g. interferon- γ secretion),

[0107] r. cells that express IDO and FAS ligand, which likely contribute to immune tolerance,

[0108] s. cells with a capacity to differentiate into 1-Human Amniotic Epithelial Cells (hAECs)

[0109] t. cells with a capacity to differentiate to neural, hepatocyte, and pancreatic cells,

[0110] u. Human Amniotic Membrane Stromal Cells (hAMSCs) differentiate to mesodermal lineages (osteogenic, chondrogenic, and adipogenic) and to all three germ

layers-ectoderm (neural), mesoderm (skeletal muscle, cardiomyocytic, and endothelial), and ectoderm (pancreatic),

[0111] v. Cells that expression CD49d by hAMSCs distinguishes hAMSCs from hAECs,

[0112] w. hAMSCs that positive for the embryonic cytoplasmic marker Oct-4 that plays a role in maintaining pluripotency and self renewal,

[0113] x. hAECs that are positive for SSEA-3, SSEA-4, TRA-1-60, TRA-1-81, and negative for SSEA-4 and non-tumorigenic.

[0114] The present inventors have now identified a need for the development of placental products that do not contain ex vivo cultured cells.

[0115] The present inventors have now identified a need for the development of placental products comprising IGFBP1.

[0116] The present inventors have now identified a need for the development of placental products comprising adiponectin.

[0117] The present inventors have now identified a need for the development of placental products exhibiting a protease-to-protease inhibitor ratio favorable for wound healing.

[0118] The present inventors have now identified a need for the development of a method of cryopreserving placental products that preserves the viability of specific cells that are other beneficial cells that are the primary source of factors for the promotion of healing to the wound healing process while selectively depleting immunogenic cells from chorionic membranes.

[0119] The present inventors have now identified a need for the development of a bioassay to test immunogenicity of manufactured placental products.

[0120] The present inventors have now identified a need for the development of placental products exhibiting a ratio of MMP to TIMP comparable to that exhibited in vivo. The present inventors have now identified a need for the development of placental products exhibiting a ratio of MMP-9 and TIMP1 of about 7-10 to one.

[0121] The present inventors have now identified a need for the development of placental products that comprise α 2-macroglobulin.

[0122] The present inventors have now identified a need for the development of placental products that inactivate serine proteinases, cysteine proteinases, aspartic proteinases, and metalloproteinases. The present inventors have now identified a need for the development of placental products that inactivate serine proteinases. The present inventors have now identified a need for the development of placental products that inactivate cysteine proteinases. The present inventors have now identified a need for the development of placental products that inactivate aspartic proteinases. The present inventors have now identified a need for the development of placental products that inactivate metalloproteinases.

[0123] The present inventors have now identified a need for the development of placental products that comprise bFGF.

[0124] The present inventors have now identified a need for the development of a cell migration assay to evaluate the potential of placental membrane products.

[0125] The present inventors have now identified a need for the development of a placental product for wound healing that comprises mesenchymal stem cells, epithelial cells and fibroblasts.

Placental Product

Overview

[0126] One embodiment of the present invention provides a placental product comprising a cryopreservation medium and

an amniotic membrane, wherein the amniotic membrane comprises viable native therapeutic cells and native therapeutic factors, and wherein the cryopreservation medium comprises a cryopreserving amount of a cryopreservative. According to this embodiment, the amniotic membrane is substantially free of at least one at least one or both of the following immunogenic cell types: CD14+ macrophages and vascularized tissue-derived cells.

[0127] In one embodiment, the amniotic membrane comprises one or more layers which exhibit the architecture of the native amniotic membrane (e.g. has not been homogenized or treated with collagenase).

[0128] In one embodiment, the placental product is suitable for dermal application to a wound.

[0129] With the teachings provided herein, the skilled artisan can now produce the present placental products. The present disclosure provides methods of manufacture that produce the technical features of the present placental products. Accordingly, in one embodiment, the placental product is manufactured by steps taught herein. The present placental products are not limited to products manufactured by the methods taught here. For example, products of the present invention could be produced through methods that rely on screening steps; e.g. steps to screen for preparations with the described technical features and superior properties.

[0130] The present placental product comprises one or more of the following technical features:

[0131] a. the viable therapeutic native cells are capable of differentiating into cells of more than one lineage (e.g. osteogenic, adipogenic and/or chondrogenic lineages),

[0132] b. the native therapeutic factors include IGFBP1,

[0133] c. the native therapeutic factors include adiponectin,

[0134] d. the native therapeutic factors include α 2-macroglobulin,

[0135] e. the native therapeutic factors include bFGF,

[0136] f. the native therapeutic factors include EGF,

[0137] g. the native therapeutic factors include MMP-9 and TIMP1,

[0138] h. the native therapeutic factors include MMP-9 and TIMP1 in a ratio of about 7 to about 10,

[0139] i. the placental product does not comprise ex-vivo cultured cells,

[0140] j. the cryopreservative medium is present in an amount of greater than about 20 ml or greater than about 50 ml,

[0141] k. the cryopreservative comprises DMSO,

[0142] l. cryopreservative comprises DMSO in a majority amount,

[0143] m. the cryopreservation medium further comprises albumin, optionally wherein the albumin is HSA,

[0144] n. the cryopreservative comprises DMSO and albumin (e.g. HSA),

[0145] o. comprises about 5,000 to about 240,000 cells/cm² or about 20,000 to about 60,000,

[0146] p. the amniotic membrane comprises at least: about 2,000, or about 2,400, or about 4,000, or about 6,000, or about 8,000, or about 10,000, or about 10,585, or about 15,000 stromal cells per unit cm² of the amniotic membrane,

[0147] q. the amniotic membrane comprises about 2,000 to about 15,000 of stromal cells per cm² of the amniotic membrane,

[0148] r. comprises stromal cells wherein at least: about 40%, or about 50%, or about 60%, or about 70%, or about

- 74.3%, or about 83.4 or about 90%, or about 92.5% of the stromal cells are viable after a freeze-thaw cycle,
- [0149] s. comprises stromal cells wherein about 40% to about 92.5% of the stromal cells are viable after a freeze-thaw cycle,
- [0150] t. the amniotic membrane has a thickness of about 20 μm to about 50 μm ,
- [0151] u. secretes less than about any of: 420 pg/mL, 350 pg/mL, or 280 pg/mL TNF- α into a tissue culture medium upon placing a 2 cm \times 2 cm piece of the tissue product in a tissue culture medium and exposing the tissue product to a bacterial lipopolysaccharide for about 20 to about 24 hours,
- [0152] v. cryopreservation and thawing, secretes less than about any of: 420 pg/mL, 350 pg/mL, or 280 pg/mL TNF- α into a tissue culture medium upon placing a 2 cm \times 2 cm piece of the tissue product in a tissue culture medium and exposing the tissue product to a bacterial lipopolysaccharide for about 20 to about 24 hours,
- [0153] w. after refrigeration, cryopreservation and thawing, secretes less than about any of: 420 pg/mL, 350 pg/mL, or 280 pg/mL TNF- α into a tissue culture medium upon placing a 2 cm \times 2 cm piece of the tissue product in a tissue culture medium and exposing the tissue product to a bacterial lipopolysaccharide for about 20 to about 24 hours,
- [0154] x. further comprises an chorionic membrane,
- [0155] y. amniotic membrane comprises a layer of amniotic epithelial cells,
- [0156] z. further comprises an chorionic membrane, wherein the amniotic membrane and the chorionic membrane are associated to one another in the native configuration,
- [0157] aa. further comprises an chorionic membrane, wherein the amniotic membrane and the chorionic membrane are not attached to one another in the native configuration,
- [0158] bb. further comprises a chorionic membrane wherein the chorionic membrane comprises about 2 to about 4 times more stromal cells relative to the amniotic membrane,
- [0159] cc. does not comprise a chorionic membrane;
- [0160] dd. comprises chorionic membrane, wherein the chorionic membrane comprises about 2 to about 4 times more stromal cells relative to the amniotic membrane, and
- [0161] ee. is suitable for dermal application to a wound;

Cells

- [0162] According to the present invention, a placental product comprises native therapeutic cells of the amniotic membrane. The cells comprise one or more of stromal cells, MSCs, fibroblasts, and epithelial cells.
- [0163] In one embodiment, the native therapeutic cells comprise viable stromal cells.
- [0164] In one embodiment, the native therapeutic cells comprise viable MSCs.
- [0165] In one embodiment, the native therapeutic cells comprise viable fibroblasts.
- [0166] In one embodiment, the native therapeutic cells comprise viable epithelial cells.
- [0167] In one embodiment, the native therapeutic cells comprise viable MSCs and viable fibroblasts.
- [0168] In one embodiment, the native therapeutic cells comprise viable MSCs, viable fibroblasts, and viable epithelial cells.

[0169] In one embodiment, the native therapeutic cells comprise viable stromal cells and viable epithelial cells.

[0170] In one embodiment, the therapeutic native cells are viable cells capable of differentiating into cells of more than one lineage (e.g. osteogenic, adipogenic and/or chondrogenic lineages).

[0171] In one embodiment, the placental product comprises about 10,000 to about 360,000 or about 40,000 to about 90,000 cells per cm^2 .

[0172] In one embodiment, the placental product comprises at least: about 2,000, or about 2,400, or about 4,000, or about 6,000, or about 8,000, or about 10,000, or about 10,585, or about 15,000 stromal cells per unit cm^2 of the placental product.

[0173] In one embodiment, the placental product comprises about 2,000 to about 15,000 of stromal cells per cm^2 of the placental product.

[0174] In one embodiment, the placental product comprises stromal cells wherein at least: about 40%, or about 50%, or about 60%, or about 70%, or about 74.3%, or about 83.4 or about 90%, or about 92.5% of the stromal cells are viable after a freeze-thaw cycle.

[0175] In one embodiment, the placental product comprises stromal cells wherein about 40% to about 92.5% of the stromal cells are viable after a freeze-thaw cycle.

[0176] In one embodiment, the placental product comprises less than about 1% of CD14+ macrophages per total cells.

[0177] In one embodiment, the amniotic membrane of the placental product comprises about 2 to about 4 times less stromal cells relative to a chorionic membrane of the same area derived from the same placenta.

[0178] In one embodiment, the placental product further comprises membrane chorionic membrane containing about 2 to about 4 times more stromal cells relative to the amniotic membrane.

[0179] In one embodiment, the amniotic membrane of the placental product comprises MSCs in an amount of: at least about 1%, at least about 2%, at least about 3%, at least about 4%, at least about 5%, about 1% to about 10%, or about 3% to about 10%, relative to the total number of cells in the amniotic membrane of the placental product. Optionally, at least: about 40%, about 50%, about 60%, or about 70% of the MSCs are viable after a freeze-thaw cycle.

[0180] In one embodiment, the amniotic membrane of the placental product comprises fibroblasts in an amount of: about 1%, about 20%, about 5% to about 15%, at least about 1%, at least about 2%, at least about 3%, or at least about 4 relative to the total number of cells in the amniotic membrane of the placental product. Optionally, at least: about 40%, about 50%, about 60%, or about 70% of the fibroblasts are viable after a freeze-thaw cycle.

[0181] In one embodiment, the amniotic membrane of the placental product comprises stromal cells in an amount of: about 5% to about 40%, about 5% to about 30%, about 10% to about 30%, about 15% to about 25%, at least about 5%, at least about 10%, or at least about 15%, relative to the total number of cells in the amniotic membrane of the placental product. Optionally, at least: about 40%, about 50%, about 60%, or about 70% of the stromal cells are viable after a freeze-thaw cycle.

[0182] In one embodiment, the amniotic membrane of the placental product comprises epithelial cells in an amount of: about 60% to about 90%, about 70% to about 90%, about 40%

to about 90%, about 50% to about 90%, at least about 40%, at least about 50%, least about 60%, or at least about 70%, relative to the total number of cells in the amniotic membrane of the placental product. Optionally, at least: about 40%, about 50%, about 60%, or about 70% of the epithelial cells are viable after a freeze-thaw cycle.

[0183] In one embodiment, the amniotic membrane of the placental product comprises MSCs and functional macrophages in a ratio of greater than about any of: 5:1, 7:1, or 10:1.

[0184] In one embodiment, the amniotic membrane of the placental product comprises fibroblasts and functional macrophages in a ratio of greater than about any of: 10:1, 15:1, 20:1, or 25:1.

[0185] In one embodiment, the amniotic membrane of the placental product comprises fibroblasts and MSCs in a ratio of: about 4:1 to about 1:1 or about 3:1 to about 3:2, or about 2:1.

[0186] In one embodiment, the amniotic membrane of the placental product comprises MSCs in an amount of: at least about 1,000 cells/cm², at least about 2,000 cells/cm², about 1,000 to about 5,000 cells/cm², or about 2,000 to about 5,000 cells/cm². Optionally, at least: about 40%, about 50%, about 60%, or about 70% of the MSCs are viable after a freeze-thaw cycle.

[0187] In one embodiment, the amniotic membrane of the placental product comprises fibroblasts in an amount of: at least about 2,000 cells/cm², at least about 4,000 cells/cm², about 2,000 to about 9,000 cells/cm², or about 2,000 to about 9,000 cells/cm². Optionally, at least: about 40%, about 50%, about 60%, or about 70% of the fibroblasts are viable after a freeze-thaw cycle.

[0188] In one embodiment, the amniotic membrane of the placental product comprises stromal cells in an amount of: at least about 4,000, at least about 8,000 cells/cm², about 4,000 to about 18,000 cells/cm², or about 4,000 to about 18,000 cells/cm². Optionally, at least: about 40%, about 50%, about 60%, or about 70% of the stromal cells are viable after a freeze-thaw cycle.

[0189] In one embodiment, the amniotic membrane of the placental product comprises epithelial cells in an amount of: at least about 10,000 cells/cm², at least about 20,000 cells/cm², at least about 32,000 cells/cm², about 10,000 to about 72,000 cells/cm², about 20,000 to about 72,000 cells/cm², or about 32,000 to about 72,000 cells/cm². Optionally, at least: about 40%, about 50%, about 60%, or about 70% of the epithelial cells are viable after a freeze-thaw cycle.

[0190] In one embodiment, the amniotic membrane of the placental product comprises functional macrophages in an amount of: less than about 3,000 cells/cm², less than about 1,000 cells/cm², or less than about 500 cells/cm².

[0191] In one embodiment, the placental product comprises a layer of amniotic epithelial cells.

[0192] In one embodiment, the placental product comprises a chorionic membrane but is substantially free of trophoblasts.

[0193] In one embodiment, the placental product is substantially free of functional CD14+ macrophages.

[0194] In one embodiment, the placental product is substantially free of vascularized tissue-derived cells.

[0195] In one embodiment, the placental product is substantially free of trophoblasts, functional CD14+ macrophages, and vascularized tissue-derived cells. Optionally, the placental product comprises viable stromal cells. Optionally, the placental product comprises viable MSCs. Optionally, the

placental product comprises viable fibroblasts. Optionally, the placental product comprises viable epithelial cells. Optionally, the placental product comprises viable MSCs, fibroblasts, and epithelial cells.

[0196] In one embodiment, the placental product comprises a chorionic membrane but is substantially free of maternal decidual cells.

[0197] In one embodiment, the placental product comprises a chorionic membrane but is substantially free of maternal leukocytes and/or trophoblast cells.

[0198] In one embodiment, the placental product is substantially free of ex-vivo cultured cells.

Placental Factors

[0199] According to the present invention, a placental product comprises native therapeutic factors of the amniotic membrane.

[0200] In one embodiment, the factors include one or more of: IGFBP1, adiponectin, α 2-macroglobulin, bFGF, EGF, MMP-9, and TIMP1. Optionally, the factors are present in amounts/cm² that are substantially similar to that of a native amniotic membrane or layer thereof (e.g. \pm 10% or 20%).

[0201] In one embodiment, the factors include IGFBP1, adiponectin, α 2-macroglobulin, bFGF, EGF, MMP-9, and TIMP1. Optionally, the factors are present in ratios that are substantially similar to that of a native amniotic membrane or layer thereof. Optionally, the factors are present in amounts/cm² that are substantially similar to that of a native amniotic membrane or layer thereof (e.g. \pm 10% or 20%).

[0202] In one embodiment, the factors include MMP-9 and TIMP1 in a ratio of about 7 to about 10 (e.g. about 7). Optionally, the factors are present in amounts/cm² that are substantially similar to that of a native amniotic membrane or layer thereof (e.g. \pm 10% or 20%).

[0203] In one embodiment, the factors include one or more (e.g. a majority or all) of the factors listed in Table 16. Optionally, the factors are present in ratios that are substantially similar to that of a native amniotic membrane or layer thereof. Optionally, the factors are present in amounts/cm² that are substantially similar to that of a native amniotic membrane or layer thereof (e.g. \pm 10% or 20%).

[0204] In one embodiment, the placental product or layer thereof comprises substantially less TNF- α /cm² than a native amniotic membrane or layer thereof, respectively.

[0205] In one embodiment, the placental product or layer thereof secretes substantially less TNF- α /cm² than a native placental product or layer thereof, respectively.

[0206] In one embodiment, the placental product secretes less than about any of: 420 pg/mL, 350 pg/mL, or 280 pg/mL TNF- α into a tissue culture medium upon placing a 2 cm \times 2 cm piece of the tissue product in a tissue culture medium and exposing the tissue product to a bacterial lipopolysaccharide for about 20 to about 24 hours.

[0207] In one embodiment, after cryopreservation and thawing, the placental product secretes less than about any of: 420 pg/mL, 350 pg/mL, or 280 pg/mL TNF- α into a tissue culture medium upon placing a 2 cm \times 2 cm piece of the tissue product in a tissue culture medium and exposing the tissue product to a bacterial lipopolysaccharide for about 20 to about 24 hours;

[0208] In one embodiment, after refrigeration, cryopreservation and thawing, the placental product secretes less than about any of: 420 pg/mL, 350 pg/mL, or 280 pg/mL TNF- α into a tissue culture medium upon placing a 2 cm \times 2 cm piece

of the tissue product in a tissue culture medium and exposing the tissue product to a bacterial lipopolysaccharide for about 20 to about 24 hours.

[0209] In one embodiment, the placental product further comprises an exogenously added inhibitor of TNF- α . Optionally, the inhibitor of TNF- α is IL-10.

[0210] In one embodiment, the product has been treated with an antibiotic

Architecture

[0211] A placental product of the present invention comprises one or more layers which exhibit the architecture of the native amniotic membrane. With the teachings provided herein, the skilled artisan will recognize placental layers that exhibit native architecture, for example, layers that have not been homogenized or treated with collagenase or other enzyme that substantially disrupts the layer.

[0212] In one embodiment, the placental product comprises a stromal cell layer with native architecture of the amniotic membrane.

[0213] In one embodiment, the placental product comprises a basement membrane with native architecture of the amniotic membrane.

[0214] In one embodiment, the placental product comprises an epithelial cell layer with native architecture of the amniotic membrane.

[0215] In one embodiment, the placental product comprises a stromal cell layer and a basement layer with native architecture of the amniotic membrane.

[0216] In one embodiment, the placental product comprises a stromal layer, a basement layer, and an epithelial cell layer with native architecture of the amniotic membrane.

[0217] In one embodiment, the placental product or amniotic membrane thereof has a thickness of about 20 μm to about 50 μm .

[0218] In one embodiment, the placental product comprises a chorionic membrane but is substantially free of trophoblasts. In one embodiment, the placental product comprises a basement membrane with native architecture of the chorionic membrane and the chorionic membrane is substantially free of trophoblasts. Optionally, the maternal side of the chorionic membrane comprises fragments of extracellular matrix proteins in a concentration substantially greater than that of a native chorionic membrane. Optionally, the placental product has been treated with Dispase (e.g. Dispase II) and/or a substantial portion of the extracellular matrix protein fragments comprises terminal leucine or phenylalanine.

[0219] In one embodiment, the placental product further comprises a chorionic membrane. Optionally, the amniotic membrane and the chorionic membrane in the placental product are associated to one another in the native configuration. Alternatively, the amniotic membrane and the chorionic membrane are not attached to one another in the native configuration.

[0220] In one embodiment, the placental product does not comprise a chorionic membrane.

Formulation

[0221] According to the present invention, the placental product can be formulated with a cryopreservation medium.

[0222] In one embodiment, the cryopreservation medium comprising one or more cell-permeating cryopreservatives, one or more non cell-permeating cryopreservatives, or a combination thereof.

[0223] Optionally, the cryopreservation medium comprises one or more cell-permeating cryopreservatives selected from DMSO, a glycerol, a glycol, a propylene glycol, an ethylene glycol, or a combination thereof.

[0224] Optionally, the cryopreservation medium comprises one or more non cell-permeating cryopreservatives selected from polyvinylpyrrolidone, a hydroxyethyl starch, a polysaccharide, a monosaccharides, a sugar alcohol, an alginate, a trehalose, a raffinose, a dextran, or a combination thereof.

[0225] Other examples of useful cryopreservatives are described in "Cryopreservation" (BioFiles Volume 5 Number 4—Sigma-Aldrich® datasheet).

[0226] In one embodiment, the cryopreservation medium comprises a cell-permeating cryopreservative, wherein the majority of the cell-permeating cryopreservative is DMSO. Optionally, the cryopreservation medium does not comprise a substantial amount of glycerol.

[0227] In one embodiment, the cryopreservation medium comprises DMSO. Optionally, the cryopreservation medium does not comprise glycerol in a majority amount. Optionally, the cryopreservation medium does not comprise a substantial amount of glycerol.

[0228] In one embodiment, the cryopreservation medium comprises additional components such as albumin (e.g. HSA or BSA), an electrolyte solution (e.g. Plasma-Lyte), or a combination thereof.

[0229] In one embodiment, the cryopreservation medium comprises 1% to about 15% albumin by weight and about 5% to about 20% cryopreservative by volume (e.g. about 10%). Optionally, the cryopreservative comprises DMSO (e.g. in a majority amount).

[0230] In one embodiment, the placental product is formulated in greater than about 20 ml or greater than about 50 ml of cryopreservation medium. Optionally, the cryopreservative comprises DMSO (e.g. in a majority amount). Optionally, the cryopreservation medium does not comprise a substantial amount of glycerol.

[0231] In one embodiment, the placental product is placed on nitrocellulose paper.

[0232] In one embodiment, the placenta is cut into a plurality of sections. Optionally, the sections are less than about 10 cm \times 10 cm. Optionally, the sections are between about 2 cm \times 2 cm and 5 cm \times 5 cm.

Manufacture

Overview

[0233] A placental product of the present invention can be manufactured from a placenta in any suitable manner that provides the technical features taught herein. According to the present invention, a placenta product comprises at least an immunocompatible amniotic membrane.

[0234] In one embodiment, a placental product is manufactured by a method comprising:

- a. obtaining a placenta,
- b. selectively depleting the placenta of immunogenicity; and
- c. cryopreserving the placenta.

[0235] Optionally, the method comprises a step of removing vascularized tissue from the placenta, for example, by lysing red blood cells, by removing blood clots, or a combination thereof.

[0236] Optionally, the method comprises a step of treating the placenta with one or more antibiotics.

[0237] Optionally, the method comprises a step of selectively depleting CD14+ macrophages, optionally as demonstrated by a substantial decrease in LPS stimulation of TNF α release.

[0238] Optionally, the step of cryopreserving the placenta comprises freezing the placenta in a cryopreservation medium which comprises one or more cell-permeating cryopreservatives, one or more non cell-permeating cryopreservatives, or a combination thereof.

[0239] Optionally, the step of cryopreserving the placenta comprises refrigerating for a period of time and then freezing, thereby selectively depleting CD14+ macrophages optionally as demonstrated by a substantial decrease in LPS stimulation of TNF α release.

[0240] Optionally, the method comprises retaining a layer of epithelial cells of the amniotic membrane.

[0241] Optionally, the method comprises a step of removing the chorionic membrane or portion thereof. Optionally, the method comprises removing trophoblasts from the chorionic membrane while retaining the stromal cell layer, reticular layer, and/or basement membrane of the chorionic membrane.

[0242] An exemplary placental product of the present invention can be manufactured or provided with a bandage or wound dressing.

Immunocompatibility and Selective Depletion

[0243] In one embodiment, the invention the placental product is immunocompatible. Immunocompatibility can be accomplished by any selective depletion step that removes immunogenic cells or factors or immunogenicity from the placenta (or amniotic membrane thereof).

[0244] In one embodiment, the placental product is made immunocompatible by selectively depleting it of functional immunogenic cells. A placenta can be made immunocompatible by selectively removing immunogenic cells from the placenta (or amniotic membrane thereof) relative to therapeutic cells. For example, immunogenic cells can be removed by killing the immunogenic cells or by purification of the placenta there from.

[0245] In one embodiment, the placental product is made immunocompatible by selectively depleting trophoblasts, for example, by removal of the trophoblast layer.

[0246] In one embodiment, the placenta is made immunocompatible by selective depletion of functional CD14+ macrophages, optionally as demonstrated by a substantial decrease in LPS stimulation of TNF α release or by MLR assay.

[0247] In one embodiment, the placenta is made immunocompatible by selective depletion of vascularized tissue-derived cells.

[0248] In one embodiment, the placenta is made immunocompatible by selective depletion of functional CD14+ macrophages, trophoblasts, and vascularized tissue-derived cells.

[0249] In one embodiment, the placenta product is made immunocompatible by selective depletion of trophoblasts and/or CD14+ macrophages, optionally as demonstrated by a substantial decrease in LPS stimulation of TNF α release or by MLR assay.

Trophoblast Removal

[0250] In one embodiment, Immunocompatibility (or selective depletion) is accomplished by removal or depletion of trophoblasts from the placental product. Trophoblasts can be removed by removing the chorionic membrane from the placental product or by removing trophoblasts from the chorionic membrane while retaining at least one of the basement layer, reticular layer, or stromal cell layer of the chorionic membrane. Surprisingly, such a placental product has one or more of the following superior features:

- a. is substantially non-immunogenic;
- b. provides remarkable healing time; and
- c. provides enhanced therapeutic efficacy.

[0251] In one embodiment, trophoblasts are removed while retaining the basement layer, reticular layer, and/or stromal cell layer of the chorionic membrane.

[0252] Trophoblasts can be removed in any suitable manner which substantially diminishes the trophoblast content of the placental product. Optionally, the trophoblasts are selectively removed or otherwise removed without eliminating a substantial portion of one or more therapeutic components from the chorionic membrane (e.g. MSCs, placental factors, etc). Optionally, a majority (e.g. substantially all) of the trophoblasts are removed.

[0253] One method of removing trophoblasts comprises treating the placenta (e.g. chorion or amino-chorion) with a digestive enzyme such as dispase (e.g. dispase II) and separating the trophoblasts from the placenta. Optionally, the step of separating comprises mechanical separation such as peeling or scraping. Optionally, scraping comprises scraping with a soft instrument such as a finger.

[0254] One method of removing trophoblasts comprises treating the chorionic membrane with dispase for about 30 to about 45 minutes separating the trophoblasts from the placenta. Optionally, the dispase is provided in a solution of about less than about 1% (e.g. about 0.5%). Optionally, the step of separating comprises mechanical separation such as peeling or scraping. Optionally, scraping comprises scraping with a soft instrument such as a finger.

[0255] Useful methods of removing trophoblasts from a placenta (e.g. chorion) are described by Portmann-Lanz et al. ("Placental mesenchymal stem cells as potential autologous graft for pre- and perinatal neuroregeneration"; American Journal of Obstetrics and Gynecology (2006) 194, 664-73), ("Isolation and characterization of mesenchymal cells from human fetal membranes"; Journal Of Tissue Engineering And Regenerative Medicine 2007; 1: 296-305.), and (Concise Review: Isolation and Characterization of Cells from Human Term Placenta: Outcome of the First International Workshop on Placenta Derived Stem Cells").

[0256] In one embodiment, trophoblasts are removed before cryopreservation.

Macrophage Removal

[0257] In one embodiment, functional macrophages are depleted or removed from the placental product. Surprisingly, such a placental product has one or more of the following superior features:

- a. is substantially non-immunogenic;
- b. provides remarkable healing time; and
- c. provides enhanced therapeutic efficacy.

[0258] Functional macrophages can be removed in any suitable manner which substantially diminishes the macroph-

age content of the placental product. Optionally, the macrophages are selectively removed or otherwise removed without eliminating a substantial portion of one or more therapeutic components from the placenta (e.g. MSCs, placental factors, etc). Optionally, a majority (e.g. substantially all) of the macrophages are removed.

[0259] One method of removing immune cells such as macrophages comprises killing the immune cells by rapid freezing rates such as 60-100° C./min.

[0260] Although immune cells can be eliminated by rapid freezing rates, such a method can also be detrimental to therapeutic cells such as stromal cells (e.g. MSCs).

[0261] The present inventors have discovered a method of selectively killing CD14+ macrophages can be selectively killed by refrigerating the placenta for a period of time (e.g. for at least about 10 min such as for about 30-60 mins) at a temperature above freezing (e.g. incubating at 2-8° C.) and then freezing the placenta (e.g. incubating at -80° C.±5° C.). Optionally, the step of freezing comprises freezing at a rate of less than 10°/min (e.g. less than about 5°/min such as at about 1°/min).

[0262] In one embodiment, the step of refrigerating comprises soaking the placenta in a cryopreservation medium (e.g. containing DMSO) for a period of time sufficient to allow the cryopreservation medium to penetrate (e.g. equilibrate with) the placental tissues. Optionally, the step of freezing comprises reducing the temperature at a rate of about 1°/min. Optionally, the step of freezing comprises freezing at a rate of less than 10°/min (e.g. less than about 5°/min such as at about 1°/min).

[0263] In one embodiment, the step of refrigerating comprises soaking the placenta in a cryopreservation medium (e.g. containing DMSO) at a temperature of about -10-15° C. (e.g. at 2-8° C.) for at least about any of: 10 min, 20 min, 30 min, 40 min, or 50 min.

[0264] In another embodiment the step of refrigerating comprises soaking the placenta in a cryopreservation medium (e.g. containing DMSO) at a temperature of about -10-15° C. (e.g. at 2-8° C.) for about any of: 10-120, 20-90 min, or 30-60 min. Optionally, the step of freezing comprises freezing at a rate of less than 10°/min (e.g. less than about 5°/min such as at about 1°/min).

Removal of Vascularized Tissue-Derived Cells

[0265] In one embodiment, vascularized tissue-derived cells (or vascularized tissue) are depleted or removed from the placental product. Surprisingly, such a placental product has one or more of the following superior features:

- is substantially non-immunogenic;
- provides remarkable healing time; and
- provides enhanced therapeutic efficacy.

[0266] Vascularized tissue-derived cells can be removed in any suitable manner which substantially diminishes such cell content of the placental product. Optionally, the vascularized tissue-derived cells are selectively removed or otherwise removed without eliminating a substantial portion of one or more therapeutic components from the placenta (e.g. MSCs, placental factors, etc).

[0267] In one embodiment, removal of vascularized tissue-derived cells comprises rinsing the amniotic membrane (e.g. with buffer such as PBS) to remove gross blood clots and any excess blood cells.

[0268] In one embodiment, removal of vascularized tissue-derived cells comprises treating the amniotic membrane with an anticoagulant (e.g. citrate dextrose solution).

[0269] In one embodiment, removal of vascularized tissue-derived cells comprises rinsing the amniotic membrane (e.g. with buffer such as PBS) to remove gross blood clots and any excess blood cells, and treating the amniotic membrane with an anticoagulant (e.g. citrate dextrose solution).

[0270] In one embodiment, the chorionic membrane is retained and removal of vascularized tissue-derived cells comprises separating the chorion from the placenta by cutting around the placental skirt on the side opposite of the umbilical cord. The chorion on the umbilical side of the placenta is not removed due to the vascularization on this side.

[0271] In one embodiment, the chorionic membrane is retained and removal of vascularized tissue-derived cells comprises separating the chorion from the placenta by cutting around the placental skirt on the side opposite of the umbilical cord and rinsing the amniotic membrane and chorionic membrane (e.g. with buffer such as PBS) to remove gross blood clots and any excess blood cells.

[0272] In one embodiment, the chorionic membrane is retained and removal of vascularized tissue-derived cells comprises separating the chorion from the placenta by cutting around the placental skirt on the side opposite of the umbilical cord and treating the amniotic membrane and chorionic membrane with an anticoagulant (e.g. citrate dextrose solution).

[0273] In one embodiment, the chorionic membrane is retained and removal of vascularized tissue-derived cells comprises separating the chorion from the placenta by cutting around the placental skirt on the side opposite of the umbilical cord, rinsing the chorionic membrane amniotic membrane (e.g. with buffer such as PBS) to remove gross blood clots and any excess blood cells, and treating the amniotic membrane with an anticoagulant (e.g. citrate dextrose solution).

Selective Depletion of Immunogenicity as Demonstrated by a Substantial Decrease in LPS Stimulation of TNF α Release.

[0274] In one embodiment, the placental product is selectively depleted of immunogenicity as demonstrated by a reduction in LPS stimulated TNF- α release. In one embodiment, the placental product is selectively depleted of macrophages.

[0275] In one embodiment, TNF- α is depleted by killing or removal of macrophages.

[0276] In one embodiment, TNF- α is functionally depleted by treatment with IL-10, which suppresses TNF- α secretion.

Preservation

[0277] A placental product of the present invention may be used fresh or may be preserved for a period of time. Surprisingly, cryopreservation results in immunocompatible placental products.

[0278] In one embodiment, a placental product is cryopreserved. A placental product may be cryopreserved by incubation at freezing temperatures (e.g. a -80° C.±5° C.) in a cryopreservative medium.

[0279] Cryopreservation can comprise, for example, incubating the placental product at 4° C. for 30-60 min, and then incubating at -80° C. until use. The placental product may then be thawed for use. Optionally, the placental product is cryopreserved in a manner such that cell viability is retained surprisingly well after a freeze-thaw cycle.

[0280] In one embodiment, cryopreservation comprises storage in a cryopreservation medium comprising one or more cell-permeating cryopreservatives, one or more non cell-permeating cryopreservatives, or a combination thereof. Optionally, the cryopreservation medium comprises one or more cell-permeating cryopreservatives selected from DMSO, a glycerol, a glycol, a propylene glycol, an ethylene glycol, or a combination thereof. Optionally, the cryopreservation medium comprises one or more non cell-permeating cryopreservatives selected from polyvinylpyrrolidone, a hydroxyethyl starch, a polysaccharide, a monosaccharides, a sugar alcohol, an alginate, a trehalose, a raffinose, a dextran, or a combination thereof. Other examples of useful cryopreservatives are described in “Cryopreservation” (BioFiles Volume 5 Number 4—Sigma-Aldrich® datasheet).

[0281] In one embodiment, the cryopreservation medium comprises a cell-permeating cryopreservative, wherein the majority of the cell-permeating cryopreservative is DMSO. Optionally, the cryopreservation medium does not comprise a substantial amount of glycerol.

[0282] In one embodiment, the cryopreservation medium comprises DMSO. Optionally, the cryopreservation medium does not comprise glycerol in a majority amount. Optionally, the cryopreservation medium does not comprise a substantial amount of glycerol.

[0283] In one embodiment, the cryopreservation medium comprises additional components such as albumin (e.g. HSA or BSA), an electrolyte solution (e.g. Plasma-Lyte), or a combination thereof.

[0284] In one embodiment, the cryopreservation medium comprises 1% to about 15% albumin by weight and about 5% to about 20% cryopreservative by volume (e.g. about 10%). Optionally, the cryopreservative comprises DMSO (e.g. in a majority amount).

[0285] In one embodiment, cryopreservation comprises placing the placenta on nitrocellulose paper.

[0286] In one embodiment, the placenta is cut into a plurality of sections before cryopreservation. Optionally, the sections are placed on nitrocellulose paper before refrigeration.

Methods of Use

[0287] The placental products of the present invention may be used to treat any tissue injury. A method of treatment may be provided, for example, by administering to a subject in need thereof, a placental product of the present invention.

[0288] A typical administration method of the present invention is topical administration. Administering the present invention can optionally involve administration to an internal tissue where access is gained by a surgical procedure.

[0289] Placental products can be administered autologously, allogeneically or xenogeneically.

[0290] In one embodiment, a present placental product is administered to a subject to treat a wound. Optionally, the wound is a laceration, scrape, thermal or chemical burn, incision, puncture, or wound caused by a projectile. Optionally, the wound is an epidermal wound, skin wound, chronic wound, acute wound, external wound, internal wounds, congenital wound, ulcer, or pressure ulcer. Such wounds may be accidental or deliberate, e.g., wounds caused during or as an adjunct to a surgical procedure. Optionally, the wound is closed surgically prior to administration.

[0291] In one embodiment, a present placental product is administered to a subject to treat a burn. Optionally, the burn

is a first-degree burn, second-degree burn (partial thickness burns), third degree burn (full thickness burns), infection of burn wound, infection of excised and unexcised burn wound, loss of epithelium from a previously grafted or healed burn, or burn wound impetigo.

[0292] In one embodiment, a present placental product is administered to a subject to treat an ulcer, for example, a diabetic ulcer (e.g. foot ulcer).

[0293] In one embodiment, a placental product is administered by placing the placental product directly over the skin of the subject, e.g., on the stratum corneum, on the site of the wound, so that the wound is covered, for example, using an adhesive tape. Additionally or alternatively, the placental product may be administered as an implant, e.g., as a subcutaneous implant.

[0294] In one embodiment, a placental product is administered to the epidermis to reduce rhtids or other features of aging skin. Such treatment is also usefully combined with so-called cosmetic surgery (e.g. rhinoplasty, rhytidectomy, etc.).

[0295] In one embodiment, a placental product is administered to the epidermis to accelerate healing associated with a dermal ablation procedure or a dermal abrasion procedure (e.g. including laser ablation, thermal ablation, electric ablation, deep dermal ablation, sub-dermal ablation, fractional ablation, and microdermal abrasion).

[0296] Other pathologies that may be treated with placental products of the present invention include traumatic wounds (e.g. civilian and military wounds), surgical scars and wounds, spinal fusions, spinal cord injury, avascular necrosis, reconstructive surgeries, ablations, and ischemia.

[0297] In one embodiment, a placental product of the present invention is used in a tissue graft procedure. Optionally, the placental product is applied to a portion of the graft which is then attached to a biological substrate (e.g. to promote healing and/or attachment to the substrate). By way of non-limiting example, tissues such as skin, cartilage, ligament, tendon, periosteum, perichondrium, synovium, fascia, mesenter and sinew can be used as tissue graft.

[0298] In one embodiment, a placental product is used in a tendon or ligament surgery to promote healing of a tendon or ligament. Optionally, the placental product is applied to portion of a tendon or ligament which is attached to a bone. The surgery can be any tendon or ligament surgery, including, e.g. knee surgery, shoulder, leg surgery, arm surgery, elbow surgery, finger surgery, hand surgery, wrist surgery, toe surgery, foot surgery, ankle surgery, and the like. For example, the placental product can be applied to a tendon or ligament in a grafting or reconstruction procedure to promote fixation of the tendon or ligament to a bone.

[0299] Through the insight of the inventors, it has surprisingly been discovered that placental products of the present invention provide superior treatment (e.g. healing time and/or healing strength) for tendon and ligament surgeries. Tendon and ligament surgeries can involve the fixation of the tendon or ligament to bone. Without being bound by theory, the present inventors believe that osteogenic and/or chondrogenic potential of MSCs in the present placental products promotes healing process and healing strength of tendons or ligaments. The present inventors believe that the present placental products provide an alternative or adjunctive treatment to periosteum-based therapies. For example, useful periosteum based treatments are described in Chen et al. (“Enveloping the tendon graft with periosteum to enhance tendon-

bone healing in a bone tunnel: A biomechanical and histologic study in rabbits"; *Arthroscopy*. 2003 March; 19(3): 290-6), Chen et al. ("Enveloping of periosteum on the hamstring tendon graft in anterior cruciate ligament reconstruction"; *Arthroscopy*. 2002 May-June; 18(5):27E), Chang et al. ("Rotator cuff repair with periosteum for enhancing tendon—bone healing: a biomechanical and histological study in rabbits"; *Knee Surgery, Sports Traumatology, Arthroscopy* Volume 17, Number 12, 1447-1453), each of which are incorporated by reference.

[0300] As non-limiting example of a method of tendon or ligament surgery, a tendon is sutured to and/or wrapped or enveloped in a placental membrane and the tendon is attached to a bone. Optionally, the tendon is placed into a bone tunnel before attached to the bone.

[0301] In one embodiment, the tendon or ligament surgery is a graft procedure, wherein the placental product is applied to the graft. Optionally, the graft is an allograft, xenograft, or an autologous graft.

[0302] In one embodiment, the tendon or ligament surgery is repair of a torn ligament or tendon, wherein the placental product is applied to the torn ligament or tendon.

[0303] Non-limiting examples of tendons to which a placental product can be applied include a digitorum extensor tendon, a hamstring tendon, a bicep tendon, an Achilles Tendon, an extensor tendon, and a rotator cuff tendon.

[0304] In one embodiment, a placental product of the present invention is used to reduce fibrosis by applying the placental product to a wound site.

[0305] In one embodiment, a placental product of the present invention is used as an anti-adhesion wound barrier, wherein the placental product is applied to a wound site, for example, to reduce fibrosis (e.g. postoperative fibrosis).

[0306] Non-limiting examples of wound sites to which the placental product can be applied include those that are surgically induced or associated with surgery involving the spine, laminectomy, knee, shoulder, or child birth, trauma related wounds or injuries, cardiovascular procedures, angiogenesis stimulation, brain/neurological procedures, burn and wound care, and ophthalmic procedures. For example, optionally, the wound site is associated with surgery of the spine and the stromal side of the placental product is applied to the dura (e.g. the stromal side facing the dura). Direction for such procedures, including the selection of wound sites and/or methodologies, can be found, for example, in WO 2009/132186 and US 2010/0098743, which are hereby incorporated by reference.

[0307] A placental product of the present invention can optionally be used to reduce adhesion or fibrosis of a wound. Postoperative fibrosis is a natural consequence of all surgical wound healing. By example, postoperative peridural adhesion results in tethering, traction, and compression of the thecal sac and nerve roots, which cause a recurrence of hyperesthesia that typically manifests a few months after laminectomy surgery. Repeated surgery for removal of scar tissue is associated with poor outcome and increased risk of injury because of the difficulty of identifying neural structures that are surrounded by scar tissue. Therefore, experimental and clinical studies have primarily focused on preventing the adhesion of scar tissue to the dura matter and nerve roots. Spinal adhesions have been implicated as a major contributing factor in failure of spine surgery. Fibrotic scar tissue can cause compression and tethering of nerve roots, which can be associated with recurrent pain and physical impairment.

[0308] Without being bound by theory, the present inventors believe that placental products taught herein are useful to reduce adhesion or fibrosis of a wound, at least in part, because the placental products can perform the very critical function in-situ of providing a immunoprivileged environment (i.e. relatively high resistance against immune responses) in the human development process. One advantage of the wound dressings and processes of the present invention is that an anti-adhesion barrier is provided which can be used to prevent adhesions following surgery, and in particular following back surgery.

[0309] In the preceding paragraphs, use of the singular may include the plural except where specifically indicated. As used herein, the words "a," "an," and "the" mean "one or more," unless otherwise specified. In addition, where aspects of the present technology are described with reference to lists of alternatives, the technology includes any individual member or subgroup of the list of alternatives and any combinations of one or more thereof.

[0310] The disclosures of all patents and publications, including published patent applications, are hereby incorporated by reference in their entireties to the same extent as if each patent and publication were specifically and individually incorporated by reference.

[0311] It is to be understood that the scope of the present technology is not to be limited to the specific embodiments described above. The present technology may be practiced other than as particularly described and still be within the scope of the accompanying claims.

[0312] Likewise, the following examples are presented in order to more fully illustrate the present technology. They should in no way be construed, however, as limiting the broad scope of the technology disclosed herein.

[0313] The presently described technology and its advantages will be better understood by reference to the following examples. These examples are provided to describe specific embodiments of the present technology. By providing these specific examples, it is not intended limit the scope and spirit of the present technology. It will be understood by those skilled in the art that the full scope of the presently described technology encompasses the subject matter defined by the claims appending this specification, and any alterations, modifications, or equivalents of those claims.

EXAMPLES

[0314] Other features and embodiments of the present technology will become apparent from the following examples which are given for illustration of the present technology rather than for limiting its intended scope.

Example 1

Characterization of Placental Membranes

[0315] Cells in placental membranes were characterized by Fluorescence Activated Cell Sorting (FACS) demonstrated the presence of stromal cells (Mesenchymal Stem Cell-like cells) in addition to fetal epithelial cells and fibroblasts.

[0316] One unique characteristic of the presently disclosed placental products is the presence of MSCs, which have been shown to be one of three types of cells (in addition to epithelial cells and fibroblasts) that are important for wound healing. Placental membranes secrete a variety of factors involved in wound healing such as angiogenic factors, factors supporting proliferation and migration of epithelial cells and fibro-

blasts, factors attracting endothelial stem cells from blood circulation to the wound site, antibacterial factors, and others.

[0317] Evaluation of proteins secreted by exemplary placental products of the invention in comparison to Apligraf and Dermagraft demonstrated a number of growth factors present in the tested products that are involved in wound healing. Examples are Vascular Endothelial Growth Factor (VEGF), Platelet-Derived Growth Factor (PDGF), Transforming Growth Factor (TGF) and others. However, several unique factors including Epidermal Growth Factor (EGF), which is one of the key factors for wound healing, are present in placental membranes and absent in Apligraf and Dermagraft. Also, placental membranes have a favorable protease-to-protease inhibitor ratio for wound healing. In an in vitro model of wound healing (cell migration assay, disclosed herein), the present inventors have demonstrated that placental membranes secrete factors promoting cell migration that will support wound closure.

Example 2

Exemplary Manufacturing Process of a Placental Product

[0318] In one embodiment, the present invention is a method of manufacturing a placental product comprising an amniotic membrane and optionally a chorionic membrane from placenta post partum. One such method is:

- [0319] a. Remove umbilical cord close to placental surface,
- [0320] b. Blunt dissect of the amnion to placental skirt,
- [0321] c. Flip placenta over and completely remove amnion,
- [0322] d. Rinse amnion in PBS to remove red blood cells,
- [0323] e. Rinse amnion once with 11% ACD-A solution to assist in red blood cell removal,
- [0324] f. Rinse amnion with PBS to remove ACD-A solution,
- [0325] g. Use PBS to remove any remaining blood from the amnion,
- [0326] h. Gently remove the connective tissue layer from the amnion,
- [0327] i. Place the amnion in PBS and set aside,
- [0328] j. Place the amnion into a bottle containing antibiotic solution and incubate at 37° C.±2° C. for 24-28 hrs,
- [0329] k. Remove bottle from the incubator and rinse membrane with PBS to remove antibiotic solution,
- [0330] l. Mount amnion (epithelial side up) on reinforced nitrocellulose paper and cut to size,
- [0331] m. Place into an FP-90 cryobag and heat seal,
- [0332] n. Add 50 mL cryopreservation solution to the bag through a syringe and remove any air trapped within the bag with the syringe,
- [0333] o. Tube seal the solution line on the FP-90 bag,
- [0334] p. Place filled bag into secondary bag and heat seal,
- [0335] q. Place unit into packaging carton,
- [0336] r. Refrigerate at 2-8° C. for 30-60 minutes, Freeze at -80° C.±5° C. inside a Styrofoam container.

Example 3

Exemplary Manufacturing Process of a Placental Product Containing an Amniotic Membrane and a Chorionic Membrane

[0337] In one embodiment, the present invention is a method of manufacturing a placental product comprising an

amniotic membrane and optionally a chorionic membranes from placenta post partum. One such method is:

- [0338] a. Remove umbilical cord close to placental surface,
- [0339] b. Blunt dissect of the amnion to placental skirt,
- [0340] c. Flip placenta over and completely remove amnion,
- [0341] d. Remove chorion by cutting around placental skirt,
- [0342] e. Rinse both membranes in PBS to remove red blood cells,
- [0343] f. Rinse both membranes once with 11% ACD-A solution to assist in red blood cell removal,
- [0344] g. Rinse both membranes with PBS to remove ACD-A solution, \
- [0345] h. Treat chorion in 0.5% dispase solution at 37° C.±2° C. for 30-45 minutes, optionally, during dispase incubation period, use PBS to remove any remaining blood from the amnion,
- [0346] i. Gently remove the connective tissue layer from the amnion,
- [0347] j. Place the amnion in PBS and set aside,
- [0348] k. When dispase treatment is complete, rinse chorion with PBS to remove dispase solution,
- [0349] l. Gently remove trophoblast layer from the chorion,
- [0350] m. Place the amnion and chorion each into a bottle containing antibiotic solution and incubate at 37° C.±2° C. for 24-28 hrs,
- [0351] n. Remove bottles from the incubator and rinse each membrane with PBS to remove antibiotic solution,
- [0352] o. Mount amnion (epithelial side up) or chorion on reinforced nitrocellulose paper and cut to size,
- [0353] p. Place each piece into an FP-90 cryobag and heat seal,
- [0354] q. Add 50 mL cryopreservation solution to the bag through a syringe and remove any air trapped within the bag with the syringe,
- [0355] r. Tube seal the solution line on the FP-90 bag,
- [0356] s. Place filled bag into secondary bag and heat seal,
- [0357] t. Place unit into packaging carton,
- [0358] u. Refrigerate at 2-8° C. for 30-60 minutes,
- [0359] v. Freeze at -80° C.±5° C. inside a Styrofoam container.

Example 4

Exemplary Placental Product Manufacturing Process

[0360] One method manufacturing a placental product comprising an amniotic membrane according to the presently disclosed manufacturing procedure is as follows:

[0361] One method of manufacturing a placental product comprising a chorionic membrane product and an amniotic membrane product according to the presently disclosed manufacturing procedure was as follows:

[0362] The placenta was processed inside a biological safety cabinet. The umbilical cord was first removed, and the amniotic membrane was peeled from the underlying chorionic membrane using blunt dissection. The membrane was rinsed with phosphate buffered saline (PBS) (Gibco Invitrogen, Grand Island, N.Y.) to remove gross blood clots and any excess blood cells. The membrane was then washed with 11% anticoagulant citrate dextrose solution (USP) formula A (ACD-A) (Baxter Healthcare Corp., Deerfield, Ill.) in saline (Baxter Healthcare Corp., Deerfield, Ill.) to remove remaining blood cells.

[0363] The stromal side of the amnion was cleaned by gently scraping away any remaining connective tissue.

[0364] The amnion was then each disinfected in 500 mL of antibiotic solution consisting of gentamicin sulfate (50 µg/mL) (Abraxis Pharmaceutical Products, Schaumburg, Ill.), vancomycin HCl (50 µg/mL) (Hospira Inc., Lake Forest, Ill.), and amphotericin B (2.5 µg/mL) (Sigma Aldrich, St. Louis, Mo.) in DMEM at 37° C.±2° C. for 24-28 hours. Vented caps were used with the incubation flasks. After the incubation period, the membrane was washed with PBS to remove any residual antibiotic solution.

[0365] The membrane was mounted on Optitran BA-S 85 reinforced nitrocellulose paper (Whatman, Dassel, Germany) and cut to the appropriate size prior to packaging into an FP-90 cryobag (Charter Medical Ltd., Winston-Salem, N.C.). The stromal side of the amnion was mounted towards the nitrocellulose paper. Once the membrane unit was placed into the FP-90 cryobag and the cryobag was heat sealed, 50 mL of a cryopreservation solution containing 10% dimethyl sulfoxide (DMSO) (Bioniche Teo. Inverin Co., Galway, Ireland) and 5% human serum albumin (HSA) (Baxter, West Lake Village, Calif.) in PlasmaLyte-A (Baxter Healthcare Corp., Deerfield, Ill.) were added through the center tubing line. Any excess air was removed, and the tubing line was subsequently sealed.

[0366] The FP-90 cryobag was placed into a mangar bag (10 in.×6 in.) (Mangar Industries, New Britain, Pa.), which was then heat sealed. The mangar bag was placed into a packaging carton (10.5 in.×6.5 in.×0.6 in.) (Diamond Packaging, Rochester, N.Y.). All cartons were refrigerated at 2-8° C. for 30-60 minutes prior to freezing at -80° C.±5° C. inside a Styrofoam container.

Example 5

Exemplary Manufacturing Process of a Placental Product Comprising Chorionic Membrane and Amniotic Membrane

[0367] One method of manufacturing a placental product comprising a chorionic membrane product and an amniotic membrane product according to the presently disclosed manufacturing procedure was as follows:

[0368] The placenta was processed inside a biological safety cabinet. The umbilical cord was first removed, and the amniotic membrane was peeled from the underlying chorionic membrane using blunt dissection. Subsequently, the chorion was removed by cutting around the placental skirt on the side opposite of the umbilical cord. The chorion on the umbilical side of the placenta was not removed due to the vascularization on this side. Both membranes were rinsed with phosphate buffered saline (PBS) (Gibco Invitrogen, Grand Island, N.Y.) to remove gross blood clots and any excess blood cells. The membranes were then washed with 11% anticoagulant citrate dextrose solution (USP) formula A (ACD-A) (Baxter Healthcare Corp., Deerfield, Ill.) in saline (Baxter Healthcare Corp., Deerfield, Ill.) to remove remaining blood cells.

[0369] The chorion was then incubated in 200 mL of a 0.5% dispase (BD Biosciences, Bedford, Mass.) solution in Dulbecco's modified eagles media (DMEM) (Lonza, Walkersville, Md.) at 37° C.±2° C. for 30-45 minutes to digest the connective tissue layer between the chorion and adjacent trophoblast layer. During this incubation period, the stromal side of the amnion was cleaned by gently scraping away any

remaining connective tissue. Once the chorion incubation period was complete, the chorion was rinsed with PBS to remove the dispase solution. Subsequently, the trophoblast layer was removed by gently peeling or scraping away these maternal decidual cells.

[0370] The amnion and chorion were then each disinfected in 500 mL of antibiotic solution consisting of gentamicin sulfate (50 µg/mL) (Abraxis Pharmaceutical Products, Schaumburg, Ill.), vancomycin HCl (50 µg/mL) (Hospira Inc., Lake Forest, Ill.), and amphotericin B (2.5 µg/mL) (Sigma Aldrich, St. Louis, Mo.) in DMEM at 37° C.±2° C. for 24-28 hours. Vented caps were used with the incubation flasks. After the incubation period, the membranes were washed with PBS to remove any residual antibiotic solution.

[0371] The membranes were mounted on Optitran BA-S 85 reinforced nitrocellulose paper (Whatman, Dassel, Germany) and cut to the appropriate size prior to packaging into an FP-90 cryobag (Charter Medical Ltd., Winston-Salem, N.C.). For the amnion, the stromal side was mounted towards the nitrocellulose paper. Once a membrane unit was placed into the FP-90 cryobag and the cryobag was heat sealed, 50 mL of a cryopreservation solution containing 10% dimethyl sulfoxide (DMSO) (Bioniche Teo. Inverin Co., Galway, Ireland) and 5% human serum albumin (HSA) (Baxter, West Lake Village, Calif.) in PlasmaLyte-A (Baxter Healthcare Corp., Deerfield, Ill.) were added through the center tubing line. Any excess air was removed, and the tubing line was subsequently sealed.

[0372] The FP-90 cryobag was placed into a mangar bag (10 in.×6 in.) (Mangar Industries, New Britain, Pa.), which was then heat sealed. The mangar bag was placed into a packaging carton (10.5 in.×6.5 in.×0.6 in.) (Diamond Packaging, Rochester, N.Y.). All cartons were refrigerated at 2-8° C. for 30-60 minutes prior to freezing at -80° C.±5° C. inside a Styrofoam container.

Example 6

Quantitative Evaluation of Cell Number and Cell Viability after Enzymatic Digestion of Placental Membranes

[0373] Amnion and chorion membranes and present placental products (from above) were evaluated for cell number and cell viability throughout the process. These analyses were performed on fresh placental tissue (prior to the antibiotic treatment step), placental tissue post antibiotic treatment, and product units post thaw. Cells were isolated from the placental membranes using enzymatic digestion. For the frozen product units, the FP-90 cryobags were first removed from the packaging cartons and mangar bags. Then the FP-90 cryobags were thawed for 2-3 minutes in a room temperature water bath. Early experiments involved the use of a 37° C.±2° C. water bath. After thaw, the placental membranes were removed from the FP-90 cryobag and placed into a reservoir containing saline (Baxter Healthcare Corp., Deerfield, Ill.) for a minimum of 1 minute and a maximum of 60 minutes. Each membrane was detached from the reinforced nitrocellulose paper prior to digestion.

[0374] Amniotic membranes were digested with 40 mL of 0.75% collagenase (Worthington Biochemical Corp., Lakewood, N.J.) solution at 37° C.±2° C. for 20-40 minutes on a rocker. After collagenase digestion, the samples were centrifuged at 2000 rpm for 10 minutes. The supernatant was removed, and 40 mL of 0.05% trypsin-EDTA (Lonza, Walk-

ersville, Md.) were added and incubated at 37° C. ±2° C. for an additional 5-15 minutes on a rocker. The trypsin was warmed to 37° C. ±2° C. in a water bath prior to use. After trypsin digestion, the suspension was filtered through a 100 µm cell strainer nylon filter to remove any debris. Centrifugation at 2000 rpm for 10 minutes was performed, and supernatant was removed. Cell pellets were reconstituted with a volume of PlasmaLyte-A that was proportional to the pellet size, and 20 µL of the resuspended cell suspension were mixed with 80 µL of trypan blue (Sigma Aldrich, St. Louis, Mo.) for counting. The cell count sample was placed into a hemocytometer and evaluated using a microscope.

[0375] Chorionic membranes were digested with 25 mL of 0.75% collagenase solution at 37° C. ±2° C. for 20-40 minutes on a rocker. After collagenase digestion, the suspension was filtered through a 100 µm cell strainer nylon filter to remove any debris. Centrifugation at 2000 rpm for 10 minutes was performed, and supernatant was removed. Cell pellets were reconstituted with a volume of PlasmaLyte-A that was proportional to the pellet size, and 20 µL of the resuspended cell suspension were mixed with 80 µL of trypan blue for counting. The cell count sample was placed into a hemocytometer and evaluated using a microscope.

[0376] Placenta membranes were analyzed prior to any processing to determine the initial characteristics of the membranes. Table 1 contains the average cell count per cm² and cell viability values for the amniotic and chorionic membranes from 32 placenta lots.

[0377] The average cell count per cm² for the amniotic membrane was 91,381 cells with a corresponding average cell viability of 84.5%. For the chorionic membrane, the average cell count per cm² was 51,614 cells with a corresponding cell viability of 86.0%.

[0378] As certain methods of manufacture are taught herein to retain amniotic membrane cells, these data further demonstrate that the present methods can produce placental products that comprise an amniotic membrane containing about 10,000 to about 360,000 cells/cm².

[0379] Since the amniotic membrane consists of epithelial cells and stromal cells, experiments were conducted to determine the ratio of epithelial cells to stromal cells. Amniotic membranes from 3 placenta lots were analyzed. First, a 5 cm×5 cm piece of amniotic membrane was digested with approximately 25 mL of 0.05% trypsin-EDTA (Lonza, Walkersville, Md.) at 37° C. ±2° C. in a water bath for 30 minutes. After the incubation step, epithelial cells were removed by gently scraping the cells from the membrane. After rinsing with PBS (Gibco Invitrogen, Grand Island, N.Y.), the membrane was subsequently digested in the same manner as chorionic membrane (described above). In addition, another intact 5 cm×5 cm piece of amniotic membrane was digested using the standard procedure (described above) to determine the total number of cells. The percentage of stromal cells was then determined by dividing the cell count from the amniotic membrane with the epithelial cells removed with the cell count from the intact membrane.

[0380] Results indicate that 19% of the total cells were stromal cells. Therefore, approximately 17,362 stromal cells were present in amniotic membrane with approximately 74,019 epithelial cells. These data indicated that there are approximately 3 times more stromal cells in chorionic membranes as compared to amniotic membranes. As certain methods of manufacture are taught herein to retain stromal cells, this ratio demonstrates that the present methods can produce

placental products that comprise a chorionic membrane and amniotic membrane, wherein the chorionic membrane comprises about 2 to about 4 times more stromal cells relative to the amniotic membrane.

TABLE 1

Cell count per cm ² and cell viability values from fresh placental tissue from 32 donors.			
Membrane	Statistics	Cell Count per cm ²	Cell Viability
Amnion	Average	91,381	84.5%
	SD	49,597	3.7%
Chorion	Average	51,614	86.0%
	SD	25,478	4.7%

[0381] Cell count and cell viability was assessed after the antibiotic treatment step. Table 2 provides the results from these analyses. Cell recoveries from this step for the amniotic membrane and the chorionic membrane were 87.7% and 70.3%, respectively.

TABLE 2

Cell count per cm ² , cell viability, and process (antibiotic treatment) cell recovery values for post antibiotic placental tissue from 28 donors.				
Membrane	Statistics	Cell Count per cm ²	Cell Viability	Process Cell Recovery
Amnion	Average	75,230	84.4%	87.7%
	SD	46,890	4.2%	49.4%
Chorion	Average	33,028	85.6%	70.3%
	SD	18,595	4.4%	31.1%

Example 7

Development of a Placental Product Cryopreservation Procedure

[0382] Cryopreservation is a method that provides a source of tissues and living cells. A main objective of cryopreservation is to minimize damage to biological materials during low temperature freezing and storage. Although general cryopreservation rules are applicable to all cells, tissues, and organs, optimization of the cryopreservation procedure is required for each type of biological material. The present application discloses a cryopreservation procedure for placental membrane products that can selectively deplete immunogenic cells from the placental membranes; and preserve viability of other beneficial cells that are the primary source of factors for the promotion of healing.

[0383] During cryopreservation method development for placental membranes, the present inventors evaluated key parameters of cryopreservation including volume of cryopreservative solution, effect of tissue equilibration prior to freezing, and cooling rates for a freezing procedures.

[0384] Acceptance of tissue allografts in the absence of immunosuppression will depend on the number of satellite immune cells present in the tissue. Cryopreservation is an approach which can be utilized to reduce tissue immunogenicity. This approach is based on differential susceptibility of different cell types to freezing injury in the presence of DMSO; leukocytes are sensitive to fast cooling rates. The freezing rate of 1° C./min is considered optimal for cells and tissues including immune cells. Rapid freezing rates such as

60-100° C./min eliminate immune cells. However, this type of procedure is harmful to other tissue cells, which are desirable for preservation according to the present invention. The developed cryopreservation procedure utilized a cryopreservation medium containing 10% DMSO, which is a key component protecting cells from destruction when water forms crystals at low temperatures. The second step of cryopreservation was full equilibration of placental membrane in the cryopreservation medium, which was achieved by soaking membranes in the cryopreservation medium for 30-60 min at 4° C. This step allowed DMSO to penetrate the placental tissues. Although there are data in the literature showing that tissue equilibration prior to freezing affects survival of immune cells (Taylor & Bank, Cryobiology, 1988, 25:1),

[0385] It was unexpectedly found that 30-60 min placental membrane equilibration in a DMSO-containing solution at 2-8° C. increases sensitivity of immune cells to freezing so that these type of cells cannot withstand freezing even at a 1° C./min freezing rate.

[0386] Temperature mapping experiments were performed to analyze the temperature profiles of potential cryopreservation conditions for the membrane products. These results are illustrated in FIG. 1. Eight (8) FP-90 cryobags were filled with either 20 mL or 50 mL of cryopreservation solution, and temperature probes were placed inside each cryobag. The first set of parameters (conditions 1 through 4 of FIG. 1a through FIG. 1d, respectively) involved a 30-minute refrigeration (2-8° C.) step prior to freezing (-80° C. ± 5° C.). In addition, the analysis involved freezing of the cryobags either inside a Styrofoam container or on the freezer shelf. The second set of parameters (conditions 5 through 8 of FIG. 1e through FIG. 1h, respectively) involved direct freezing (-80° C. ± 5° C.) of the cryobags either inside a Styrofoam container or on the freezer shelf. The results indicated that condition 6 and condition 2 exhibited the most gradual temperature decreases. Gradual temperature decreases are typically desired in order to preserve cell viability. The difference between condition 6 and condition 2 was that condition 2 included a 30-minute refrigeration step. Therefore, the decrease in temperature from the start of freezing to -4° C., where latent heat evolution upon freezing occurs, was examined further. For condition 6, the rate of cooling was approximately -1° C./minute during this period. The rate of cooling for condition 2 was approximately -0.4° C./minute during the same timeframe. Therefore, condition 2 was selected for incorporation into a non-limiting cryopreservation process since slower rates of cooling are generally desired to maintain optimal cell viability.

[0387] FIG. 2 depicts the effects of cryopreservation solution volume on process (cryopreservation) cell recovery for the amniotic membrane. The analysis of the 10 mL cryopreservation solution volume involved 5 placenta lots, and the analysis of the 20 mL cryopreservation solution volume included 3 lots. For the 50 mL cryopreservation solution volume, 14 placenta lots were analyzed.

[0388] As depicted in FIG. 2, the 50 mL volume of cryopreservation solution volume provided superior cell recovery compared to that of the 10 mL and 20 mL. Since the 50 mL cryopreservation solution volume provided the slowest cooling rate, FIG. 2 indicates that a cooling rate of less than about 2° C./minute or less than about 1° C./minute can provide a superior method of manufacture according to the present invention.

[0389] FIG. 3 shows the results of a similar study of analysis of the cryopreservation cell recovery for the chorionic membrane, demonstrating that a cryopreservation solution volume of 50 mL was optimal. The analysis of the 10 mL cryopreservation solution volume involved 5 placenta lots, and the analysis of the 20 mL cryopreservation solution volume included 3 lots. For the 50 mL cryopreservation solution volume, 16 placenta lots were analyzed.

[0390] Experiments were conducted to evaluate different potential freezing conditions to maximize cell recovery after the cryopreservation process. FIG. 4 (cells from amniotic membrane) and FIG. 5 (cells from chorionic membranes) depict these results, showing the effects of refrigeration time and freezing parameters on process (cryopreservation) cell recovery for the chorionic membrane. Three conditions were analyzed. These conditions were also linked to the temperature mapping studies. The first condition involved directly freezing the product unit on a shelf within the freezer (-80° C. ± 5° C.). The second condition also contained a direct freeze, but the product unit was placed into a Styrofoam container within the freezer. The third condition included a refrigeration (2-8° C.) period of 30 minutes prior to the freezing step. For the amniotic membrane, 3 placenta lots were evaluated. Two (2) placenta lots were analyzed for the chorionic membrane. Results indicated that the third condition was optimal for both membrane types.

[0391] All of the cryopreservation parameters that were assessed for the amniotic and chorionic membranes are summarized in Table 3 and Table 4. The evaluation of the cell recoveries and cell viabilities from these experiments resulted in the selection of the final parameters for the manufacturing process. In addition, all average cell viability values were $\geq 70\%$.

TABLE 3

Post thaw cell count per cm ² , cell viability, and process (cryopreservation) cell recovery values for the amniotic membrane.						
Parameter	Condition Tested	Statistics	Cell		Process	
			Count per cm ²	Cell Viability	Cell Recovery	Comments/Conclusions
Refrigerate at 2-8° C. for 30-60 min and freeze at -80° C. ± 10° C.	All conditions	Average	55,709	83.4%	64.2%	Overall assessment.
		SD	45,210	4.4%	22.5%	
		N	32	32	32	

TABLE 3-continued

Post thaw cell count per cm ² , cell viability, and process (cryopreservation) cell recovery values for the amniotic membrane.						
Parameter	Condition Tested	Statistics	Cell Count per cm ²	Cell Viability	Process Cell Recovery	Comments/Conclusions
Refrigeration time interval	30 min	Average	52,173	83.1%	63.7%	No significant difference found in process cell recovery. A 30-60 min range was established.
		SD	39,750	4.5%	21.4%	
		N	26	26	26	
	60 min	Average	71,033	85.0%	66.5%	
		SD	66,525	3.9%	29.3%	
		N	6	6	6	
Thawing temperature	37° C. ± 2° C. water bath	Average	48,524	83.3%	64.0%	No significant difference found in process cell recovery. The room temp condition was selected for logistical reasons.
		SD	27,804	1.7%	34.4%	
		N	7	7	7	
	Room temp water bath	Average	57,721	83.5%	64.3%	
		SD	49,271	4.9%	19.0%	
		N	25	25	25	
Holding period after transfer into saline	1-15 min	Average	50,873	83.1%	65.0%	No significant difference found in process cell recovery. Membranes can be held in saline for up to 1 hr.
		SD	38,969	3.9%	24.2%	
		N	26	26	26	
	1 hr	Average	76,667	85.1%	61.0%	
		SD	66,565	6.2%	14.3%	
		N	6	6	6	
Tissue size	5 cm × 5 cm	Average	58,431	83.3%	62.8%	No decrease in process cell recovery from the 5 cm × 5 cm product to the 2 cm × 2 cm product. Both sizes were acceptable for use.
		SD	47,603	4.5%	21.7%	
		N	28	28	28	
	2 cm × 2 cm	Average	36,656	84.4%	73.9%	
		SD	13,175	3.4%	29.5%	
		N	4	4	4	

TABLE 4

Post thaw cell count per cm ² , cell viability, and process (cryopreservation) cell recovery values for the chorionic membrane.						
Parameter	Condition Tested	Statistics	Cell Count per cm ²	Cell Viability	Process Cell Recovery	
Refrigerate at 2-8° C. for 30-60 min and freeze at -80° C. ± 10° C.	All conditions	Average	23,217	87.3%	102.8%	
		SD	9,155	4.1%	65.5%	
		N	27	27	27	
Dispase treatment	30 min	Average	22,354	85.7%	81.1%	No decrease in process cell recovery for the 45 min treatment. A 30-45 min range was established.
		SD	9,505	5.1%	32.4%	
		N	24	24	24	
	45 min	Average	27,125	90.6%	172.6%	
		SD	7,963	2.2%	101.2%	
		N	6	6	6	

TABLE 4-continued

Post thaw cell count per cm ² , cell viability, and process (cryopreservation) cell recovery values for the chorionic membrane.						
Parameter	Condition Tested	Statistics	Cell Count per cm ²	Cell Viability	Process Cell Recovery	
Refrigeration time interval	30 min	Average	23,815	86.8%	102.2%	The process recovery value was >80% for the 60 min time interval. A 30-60 min range was established.
		SD	9,681	5.2%	68.8%	
		N	25	25	25	
	60 min	Average	20,773	85.8%	84.9%	
		SD	7,356	4.7%	14.4%	
		N	5	5	5	
Thawing temperature	37° C. ± 2° C. water bath	Average	33,360	85.9%	114.7%	No significant difference found in process cell recovery. The room temp condition was selected for logistical reasons.
		SD	8,497	4.0%	38.1%	
		N	5	5	5	
	Room temp water bath	Average	21,298	86.8%	96.3%	
		SD	8,189	5.3%	67.2%	
		N	25	25	25	
Holding period after transfer into saline	1-15 min	Average	23,733	86.6%	100.6%	No significant difference found in process cell recovery. Membranes can be held in saline for up to 1 hr.
		SD	9,674	5.1%	67.0%	
		N	26	26	26	
	1 hr	Average	20,550	87.0%	91.4%	
		SD	6,575	4.8%	32.0%	
		N	4	4	4	
Tissue size	5 cm × 5 cm	Average	23,391	86.1%	99.6%	No decrease in process cell recovery from the 5 cm × 5 cm product to the 2 cm × 2 cm product. Both sizes were acceptable for use.
		SD	8,865	5.0%	58.7%	
		N	23	23	23	
	2 cm × 2 cm	Average	23,036	88.4%	98.7%	
		SD	11,362	5.0%	81.3%	
		N	7	7	7	

Notes:

cm = centimeter; min = minutes; temp = temperature; hr = hour, SD = standard deviation; N = number.

[0392] These data indicate that a step of dispase treatment for about 30-45 mins provides a superior method of manufacture according to the present invention. These data also indicate that a step of refrigeration at 2-8° C. for 30-60 min before freezing provides a superior method of manufacture according to the present invention.

Example 8

Qualitative Evaluation of Cell Viability by Tissue Staining

[0393] The amniotic and chorionic membranes were stained using a LIVE/DEAD® Viability/Cytotoxicity kit (Molecular Probes Inc., Eugene, Oreg.) to qualitatively assess cell viability. Staining was performed as per the manufacturer's protocol. Membrane segments of approximately 0.5 cm×0.5 cm were used. Evaluation of stained membranes was performed using a fluorescent microscope. An intense uniform green fluorescence indicated the presence of live cells, and a bright red fluorescence indicated the presence of dead cells. Images of fresh amniotic and chorionic membranes as well as cryopreserved amniotic and chorionic membranes

demonstrated that the manufacturing process did not alter the phenotypic characteristics of the membranes post thaw.

[0394] FIG. 6 contains representative images of fresh amniotic and chorionic membranes as well as cryopreserved amniotic and chorionic membranes. These images demonstrated that the manufacturing process did not alter the phenotypic characteristics of the membranes and the proportion of viable cell types (epithelial and stromal cells) in the membranes post thaw.

[0395] FIG. 6. Representative images of the live/dead staining of the epithelial layer of fresh amniotic membrane (A); epithelial layer of cryopreserved amniotic membrane (B); stromal layer of fresh amniotic membrane (C); stromal layer of cryopreserved amniotic membrane (D); fresh chorionic membrane (E); and cryopreserved chorionic membrane (F). Live cells are green, and dead cells are red.

Example 9

Placental Tissue Immunogenicity Testing

[0396] One unique feature of the human amnion and chorion is the absence of fetal blood vessels that prevent

mobilization of leukocytes from fetal circulation. On the fetal side, macrophages resident in the chorioamniotic mesodermal layer represent the only population of immune cells. Thus, fetal macrophages present in the chorion and amnion are the major source of tissue immunogenicity. However, the number of macrophages in amnion is significantly lower (Maggatti et al, *Stem Cells*, 2008, 26: 182), and this explains the low immunogenicity of amnion and the ability to use it across HLA barriers without matching between the donor and recipient (Akle et al, *Lancet*, 1981, 8254:1003; Ucakhan et al., *Cornea*, 2002, 21:169). In contrast, the chorion is considered immunogenic. In a study where the amnion was used together with the chorion for plastic repair of conjunctival defects, the success rate was low (De Roth Arch *Ophthalmol*, 1940, 23: 522). Without being bound by theory, the present inventors believe that removal of CD14+ cells from placental membranes eliminates activation of lymphocytes in vitro. In addition to the presence of fetal macrophages, the present inventors believe that immunogenicity of chorion can be mediated by contamination of blood cells coming from the maternal trophoblast, which contains blood vessels. Thus, the processing of placental membrane for clinical use can be enhanced by purification of the amnion and chorion from maternal trophoblasts and selective elimination of all CD14+ fetal macrophages.

[0397] Immunogenicity testing can be used to characterize a placental product as safe clinical therapeutics. For example, two bioassays can be used to test immunogenicity of manufactured placental products: Mixed Lymphocyte Reaction (MLR) and Lipopolysaccharide (LPS)-induced Tumor Necrosis Factor (TNF)- α secretion.

Example 10

Mixed Lymphocyte Reaction (MLR)

[0398] An MLR is a widely used in vitro assay to test cell and tissue immunogenicity. The assay is based on the ability of immune cells (responders) derived from one individual to recognize allogeneic Human Leukocyte Antigen (HLA) and other antigenic molecules expressed on the surface of allogeneic cells and tissues (stimulators) derived from another individual when mixed together in a well of an experimental tissue culture plate. The response of immune cells to stimulation by allogeneic cells and tissues can be measured using a variety of methods such as secretion of particular cytokines (e.g., Interleukin (IL)-2), expression of certain receptors (e.g., IL-2R), or cell proliferation, all of which are characteristics of activated immune cells.

[0399] Placental tissue samples representing different steps of the presently disclosed manufacturing process were used for immunogenicity testing. These samples included amnion with chorion and trophoblast as a starting material and separated choriotrophoblast, chorion, trophoblast, and amnion. Both freshly purified and cryopreserved (final products) tissues were tested.

[0400] For the MLR assay, cells from placental tissues were isolated using 280 U/mL of collagenase type II (Worthington, Cat No. 4202). Tissues were treated with enzyme for 60-90 min at 37° C. \pm 2° C., and the resulting cell suspension was filtered through a 100 μ m filter to remove tissue debris. Single cell suspensions were then centrifuged using a Beckman, TJ-6 at 2000 rpm for 10 min and washed twice with DPBS. Supernatant was discarded after each wash, and cells were resuspended in 2 mL of DMEM (Invitrogen, Cat No. 11885) and evaluated for cell number and cell viability by counting cells in the presence of Trypan blue dye (Invitrogen, Cat No. 15250-061). For the MLR, placental-derived cells were

mixed with allogeneic hPBMCs at a 1:5 ratio in 24-well culture plates in DMEM supplemented with 5% fetal bovine serum (FBS) and incubated for 4 days in the incubator containing 5% CO₂, 95% humidity at 37° C. \pm 2° C. Human Peripheral Blood Mononuclear Cells (hPBMCs) alone were used as a negative control, and a mixture of two sets of hPBMCs derived from two different donors was used as a positive MLR control. After 4 days of incubation, cells were collected from wells, lysed using a lysis buffer (Sigma, Cat No. C2978) supplemented with protease inhibitor cocktail (Roche, Cat No. 11836153001), and IL-2R was measured in cell lysates using the sIL-2R ELISA kit (R&D Systems, Cat No. SR2A00) according to the established protocol (G-SOP-Q088).

[0401] The level of IL-2R is a measure of activation of T-cells in response to immunogenic molecules expressed by allogeneic cells. Results of 2 out of 12 representative experiments are shown in FIG. 7 and FIG. 8. Results presented in these figures demonstrate a method of manufacture of placental membranes, resulting in low immunogenicity of the final products.

[0402] As depicted in FIG. 7: the manufacturing process serially reduces immunogenicity of the placental product. Samples representing different steps of the manufacturing process (Chorion+Trophoblast (CT), Trophoblast (T), Amnion (AM), and Chorion (CM)) were co-cultured with hPBMCs for 4 days. IL-2 α R was measured in cell lysates as a marker of T-cell activation. Negative control shows a basal level of immune cell activation: PBMCs derived from one donor were cultured alone. Positive control: a mixture of PBMCs derived from 2 different donors.

[0403] As depicted in FIG. 8, selective depletion of immunogenicity results from the present cryopreservation process of producing the present placental products, as evidenced by the significant decrease in immunogenicity upon cryopreservation.

Example 11

LPS-Induced TNF- α Secretion by Placental Membrane Cells

[0404] As described herein, fetal macrophages present in the amnion and chorion are a major source of tissue immunogenicity. Without being bound by theory, the present inventors believe that removal of CD14+ cells from placental membrane eliminates activation of lymphocytes and that depletion of allogeneic donor tissue macrophages decreases the level of inflammatory cytokine secretion and tissue immunogenicity. The inventors also believe that reduction of tissue immunogenicity can also be reached by depletion of TNF- α with anti-TNF- α antibodies or suppression of TNF- α secretion by IL-10. Macrophages in fetal placental membranes respond to bacteria by secretion of inflammatory cytokines. The secretion of TNF- α by fresh placental membranes in vitro in response to bacterial LPS is significantly higher in the chorionic membrane. Thus, the present inventors believe that immunogenicity of placental membranes is mediated by macrophages, the amount and/or activity of which is higher in the chorionic membrane.

[0405] According to the present invention, selective depletion of macrophages is an approach to selectively deplete immunogenicity of the amniotic and chorionic membranes, allowing the use of both allogeneic membranes for clinical applications. The assay of functional macrophages in a placental product is used here as an assay for immunogenicity testing (e.g. in production or prior to clinical use) based on the facts that: macrophages are the source of immunogenicity in chorionic membranes. Macrophages in placenta-derived

membranes respond to bacterial LPS by secretion of high levels of TNF- α ; and TNF- α is a critical cytokine involved in immune response and allograft tissue rejection. Therefore, secretion of TNF- α by placenta-derived membranes in response to LPS is used here to characterize tissue immunogenicity and for pre-use screening.

Example 12

Establishment of Allowed LPS-Induced TNF- α Secretion Level by Placental Membranes

[0406] Data from published reports regarding the level of TNF- α , which is associated with the absence or an insignifi-

cant immune response in a variety of experimental systems, are presented in Table 5. These data indicate that a TNF- α level below 100 pg/mL correlates with a low immune response. The ability of amniotic and chorionic membranes to produce TNF- α spontaneously and in response to bacteria or bacterial LPS in vitro has been shown by a number of investigators. Table 6 summarizes such data. The lowest spontaneous TNF- α secretion by amniotic membrane of about 70 pg/cm² of the membrane was reported by Fortunato et al. All reports also showed that fresh placental membranes secrete large amounts of TNF- α in response to bacteria or bacterial LPS, which is attributed to the presence of viable functional macrophages.

TABLE 5

TNF- α levels associated with no or an insignificant immune response.			
Description of experimental system	TNF- α levels associated with the absence/reduction of immune response	Comments	References
IL-10-induced inhibition of MLR in vitro. TNF was measured in tissue culture supernatant by ELISA. MLR using skin tissue explants (0.02 cm ² per well) as stimulators in the presence or absence of IL-10 (skin explant assay). Skin tissue destruction was assessed microscopically, and severity was assigned based on histopathological tissue damage.	Mean 260 pg/mL Mean 100 pg/mL		Wang et al., Transplantation, 2002, 74: 772
Endogenous TNF production in MLR in the presence or absence of anti-TNF antibodies. TNF levels were assessed using the WEHI-164 cytotoxicity assay.	~0.04 U/mL for the negative control and MLR in the presence of anti-TNF antibodies, which correlated with no or significant inhibition of lymphocyte proliferation	TNF activity per mg is not provided.	Shalaby et al, J Immunol, 1988, 141: 499
TNF levels in BAL fluid of lung isografts, unmodified allograft, and alveolar macrophages (AM) depleted allograft in rats.	Isograft: below detection; AM-depleted allograft: ~15 pg/mL of BAL (total 75 pg/5 ml of BAL)	Unmodified allograft: ~45 pg/mL (immunogenic)	Sekine et al, J Immunol, 1997, 159: 4084.
TNF levels in MLR after 48 hours in the presence or absence of advanced glycation end products (MLR inhibitors).	~<200 pg/mL TNF correlated with a complete inhibition of MLR		Ohashi et al, Clin Immunol, 2009, e-pub ahead of print
TNF levels in MLR.	<100 pg/mL TNF in MLR with HLA-matched donors (control, no stimulation)		Toungouz et al, Hum Immunol, 1993, 38: 221
TNF activity in MLR when pieces of cryopreserved skin allografts (~0.2 cm ²) were incubated with hPBMCs for 24 hours. Positive control: hPBMC + LPS; negative - hPBMC alone.	Negative control ~20 U of TNF activity; MLR with skin explants - 0-40 U; Positive control - 600 U	Unit of activity was calculated as TNF in ng/mL divided by OD at 570 nm for the same experimental well	Lomas et al, Cell Tissue Bank, 2004, 5: 23.

TABLE 5-continued

TNF- α levels associated with no or an insignificant immune response.			
Description of experimental system	TNF- α levels associated with the absence/reduction of immune response	Comments	References
Cytokine time course in MLR, including TNF.		Optimal TNF after 24 hours - ~150 pg/mL	Jordan & Ritter, J Immunol Meth, 2002, 260: 1
MLR using skin tissue explants (0.02 cm ² per well) as stimulators in the presence or absence of anti-TNF antibodies (skin explant assay). Skin tissue destruction was assessed microscopically, and severity was assigned based on histopathological tissue damage.	For no skin destruction: 0.5-1.1 pg/mL for HLA compatible responders, and 2.6-1376 pg/mL for unmatched MLR	Recalculation per 1 cm ² of skin tissue: lowest TNF non-immunogenic level is 100 pg/cm ²	Dickinson et al, Cytokine, 1994, 6: 141

TABLE 6

Secretion of TNF in vitro by fresh amniotic and chorionic membranes.			
Description of experimental system	TNF levels secreted by fresh placental membranes in culture	Comments/ recalculations of the lowest TNF levels per cm ²	References
TNF secretion by "fresh" amnion and chorion tissues (1.44 cm ²) incubated for 24 hours in the presence or absence of LPS (500 ng/mL).	Chorion: basal 3.3 \pm 0.46 ng/cm ² , LPS-induced - 150-250 ng/cm ² Amnion: basal 2.5 \pm 1.3 ng/cm ² , LPS-induced - ~50 ng/cm ²	Lowest TNF level for amnion is 1200 pg/cm ²	Zaga et al., Biol Reprod, 2004, 71: 1296
TNF secretion by "fresh" amnion and chorion tissues (1.8 cm diameter disks: 2.5 cm ²) incubated for 24 hours in the presence or absence of <i>E. Coli</i> in 1 mL medium.	Basal ~1-2.5 pg/ μ g total protein in the medium for both amnion and chorion; <i>E. Coli</i> -induced: amnion - 29.2 (14.5-35.3) pg and chorion - 53.15 (40-94.2) pg per μ g total protein	Lowest TNF level for amnion is 800 pg/cm ²	Zaga-Clavellina et al, Reprod Biol Endocrinol, 2007, 5: 46
TNF secretion by "fresh" amnion and chorion tissues (chorion 8-10 mg tissue/mL; amnion 5-7 mg/mL, 0.02-0.04 cm ²) incubated for 20 hours in the presence or absence of LPS (5 μ g/mL).	Basal: ~2-64 U/mL or 8-10 mg chorion; <1 U/mL for 5-7 mg amnion. LPS-induced: >100 U/10 mg for chorion and ~15-17 U/10 mg for amnion	1 unit = ~100-200 pg/mL; Lowest TNF level for amnion is <100 pg/mL corresponding to <2500 pg/cm ²	Paradowska et al, Placenta, 1997, 18: 441
TNF secretion by "fresh" amnion (0.57 cm ²) in 0.8 mL incubated for 24 hours in the presence or absence of LPS (50 ng/mL).	Amnion: Basal - 40 pg/mL, LPS-induced - 410 pg/mL	Lowest TNF level for fresh amnion is ~70 pg/cm ²	Fortunato et al, Am J Obstet Gynecol, 1996, 174: 1855

TABLE 6-continued

Secretion of TNF in vitro by fresh amniotic and chorionic membranes.			
Description of experimental system	TNF levels secreted by fresh placental membranes in culture	Comments/recalculations of the lowest TNF levels per cm ²	References
TNF secretion by "fresh" amnion and chorion tissues (4 cm ²) incubated for 24 hours in the presence or absence of LPS (1-1000 ng/mL)	Basal: Amnion ~7-13 ng/mL/g tissue); Chorion ~18 ng/mL/g tissue LPS-induced (1000 ng/mL): Amnion ~14 ng/mL/g), Chorion ~27 ng/mL/g	Amnion is 5-7 mg corresponds ~0.02-0.04 cm ² ; 1 g is ~6 cm ² ; Lowest TNF level for amnion is ~1000 pg/cm ²	Thiex et al, Reprod Biol Endocrinol, 2009, 7: 117

Example 13

LPS-Induced TNF- α Secretion and CT induced MLR Immunogenicity Assay

[0407] 2 cm \times 2 cm pieces of placental membranes representing intermediates and final products were placed in tissue culture medium and exposed to bacterial LPS (1 μ g/mL) for 20-24 hr. After 24 hours, tissue culture supernatant were collected and tested for the presence of TNF- α using a TNF- α ELISA kit (R&D Systems) according to the manufacturer's protocol. Human hPBMCs (SeraCare) known to contain monocytes responding to LPS by secretion of high levels of TNF- α were used as a positive control in the assay. hPBMCs and placental tissues without LPS were also included as controls in the analysis. In this assay, TNF detected in the culture medium from greater than 70 pg/cm² (corresponding to 280 pg/mL) for both spontaneous and LPS-induced TNF- α secretion was considered immunogenic.

[0408] As depicted in FIG. 9A and FIG. 9B, the manufacturing process serially reduces immunogenicity of the placental product. Samples representing different steps of the manufacturing process (Amnion+Chorion+Trophoblast (ACT), Chorion+Trophoblast (CT), Amnion (AM), and Chorion (CM)) were incubated in the presence of LPS for 24 hr, and after that tissue culture supernatants were tested for the TNF- α by ELISA. Tissues cultured in medium without LPS show the basal level of TNF secretion. PBMCs, which are known to secrete high levels of TNF, were used as a positive control.

[0409] The low levels of TNF- α and the absence of the response to LPS by AM and CM indicates the absence of viable functional macrophages that are the major source of immunogenicity for amniotic and chorionic membranes. Results of this assay showed a correlation with the MLR data: tissues that produce high levels of TNF- α in response to LPS are immunogenic in the MLR assay (FIG. 9A and FIG. 9B, for TNF- α secretion; FIG. 9, C-MLR).

[0410] As depicted in FIG. 9A and FIG. 9B, the manufacturing process serially reduces immunogenicity of the placental product. Samples representing different steps of the manufacturing process (Amnion+Chorion+Trophoblast (ACT), Chorion+Trophoblast (CT), Amnion (AM), and Chorion (CM)) were incubated in the presence of LPS for 24 hr, and after that tissue culture supernatants were tested for the TNF- α by ELISA. Tissues cultured in medium without LPS

show the basal level of TNF secretion. PBMCs, which are known to secrete high levels of TNF, were used as a positive control.

[0411] Choriotrophoblast (CT), which secreted high levels of TNF- α (FIG. 9, B), was tested in MLR against two different PBMC donors. CT cells were co-cultured with PBMCs for 4 days. IL-2 α R was measured in cell lysates as a marker of T-cell activation. Positive control: a mixture of PBMCs derived from 2 different donors.

[0412] FIG. 9C shows that preparations producing high levels of TNF- α are immunogenic. Choriotrophoblast (CT), which secreted high levels of TNF- α (FIG. 9 B), was tested in MLR against two different PBMC donors. CT cells were co-cultured with PBMCs for 4 days. IL-2 α R was measured in cell lysates as a marker of T-cell activation. Positive control: a mixture of PBMCs derived from 2 different donors.

Example 14

Analysis of Placental Cells by FACS

[0413] Knowing the cellular composition of amnion and chorionic membranes is important for developing a thorough understanding of potential functional roles in wound healing and immunogenicity. Previous reports demonstrated that both amnion and chorion contains multiple cell types. Purified amnion has two major cellular layers: epithelial cells and stromal. In addition to epithelial cells and fibroblasts, stromal cells were identified in the amnion and chorion. Although there are no fetal blood vessels within either the amniotic or chorionic membranes, both membranes comprise resident fetal macrophages. The close proximity to maternal blood circulation and decidua provide a potential source of immunogenic cells (maternal leukocytes and trophoblast cells) and therefore are a potential source of immunogenicity. To investigate the cellular composition of the amnion and chorion, FACS analysis was performed.

Example 14.1

FACS Procedure: Single Cell Suspension Preparation

[0414] Purified amnion and chorionic membranes were used for cellular phenotypic analysis via FACS. Cells from amnion and chorion were isolated using 280 U/mL collagenase type II (Worthington, Cat No. 4202). Tissues were treated with enzyme for 60-90 min at 37 $^{\circ}$ C. \pm 2 $^{\circ}$ C., and the

resulting cell suspension was filtered through a 100 µm filter to remove tissue debris. Single cell suspensions were then centrifuged using a Beckman TJ-6 at 2000 rpm for 10 min and washed twice with DPBS. Supernatant was discarded after each wash, and cells were resuspended in 2 mL of FACS staining buffer (DPBS+0.09% NaN₃+1% FBS).

Example 14.2

Immunolabeling Cells for Specific Cellular Markers

[0415] Once the single cell suspension was prepared according to Example 10, a minimum of 1×10⁵ cells in 100 µL of FACS staining buffer was treated with antibodies labeled with fluorescent dye. Table 7 provides descriptions of the antibodies and the amounts used. For cell surface markers, cells were incubated for 30 min at room temperature in the dark with antibodies followed by washing twice with FACS staining buffer by centrifugation at 1300 rpm for 5 min using a Beckman TJ-6 centrifuge. Cells were then resuspended in 400 µL of FACS staining buffer and analyzed using a BD FACSCalibur flow cytometer. To assess cell viability, 10 µL of 7-AAD reagent (BD, Cat No. 559925) was added just after the initial FACS analysis and analyzed again. For intracellular staining, cells were permeabilized and labeled following the manufacturer's recommendations (BD Cytotfix/Cytoperm, Cat No. 554714) and analyzed using a BD FACSCalibur flow cytometer.

TABLE 7

Description of reagents used for placental cell characterization by FACS.				
Cell marker antibody and label type	Cat No.	Volume of antibody solution used	Cell marker type	Cell marker specificity
IgG1 isotype-PE	BD 559320	5 µL	Cell surface	Isotype control
CD105-PE	Caltag MHCD10504	20 µL	Cell surface	MSC marker
CD166-PE	BD 559263	80 µL	Cell surface	MSC marker
CD45-PE	BD 555483	10 µL	Cell surface	Hematopoietic cell marker
IgG2a isotype-PE	BD 555574	2 µL	Cell surface	Isotype control
CD14-PE	BD 555398	20 µL	Cell surface	Monocyte marker
HLA-DR-PE	BD 556644	20 µL	Cell surface	HLA class II specific for antigen-presenting cells
IgG1 isotype-FITC	BD555748	5 µL	Cell surface	Isotype control
CD86-FITC	BD 557343	20 µL	Cell surface	Immune co-stimulatory marker
CD40-FITC	BD 556624	20 µL	Cell surface	Immune co-stimulatory marker
IgG1 isotype-unlabeled	Dako X0931	10 µL	Intracellular	Isotype control
Cytokeratin 7-unlabeled	Dako M7018	2 µL	Intracellular	Trophoblast marker
Rabbit anti-mouse FITC	Dako F0261	5 µL	Intracellular	Secondary antibody

Example 15

Phenotypic Analysis of Placental Cells

[0416] FACS analysis of single cell suspensions of both amnion and chorion membranes demonstrates that both

membranes contain cells expressing markers specific for mesenchymal stem cells (refer to Table 8), implicating the presence of MSCs. In addition, several immunogenic markers, which are more likely expressed on CD14+ placental macrophages, were detected. The % ranges for different markers are wide. It can be explained by: 1) high variability in cell number between placenta donors; and 2) technical issues, which include the presence of high and variable cellular and tissue debris in the cellular suspension. Although debris can be gated out, debris particles that are comparable with cells by size will affect the accuracy of the calculated % for each tested marker. In addition, Table 9 provides a FACS analysis of cells isolated from the amniotic and chorionic membranes that were cultured in 10% FBS in DMEM at 37° C.±2° C. until confluency (passage 0 cells).

[0417] These data demonstrated that cells derived from amniotic and chorionic membranes retained a phenotype similar to MSCs after culturing. In conclusion, the presence of stromal cells in placental tissues was confirmed by FACS analysis.

[0418] As certain methods of manufacture are taught herein to retain amniotic membrane cells, these data indicate that the present methods can produce placental products that comprise a amniotic membrane containing MSC-like cells that express CD105, CD166, C90, and/or CD73 (also referred to herein as AMSCs or MSCs).

TABLE 8

Characterization of the cellular composition of placental membranes based on selective CD markers.			
	Marker	Amnion (% range)	Chorion (% range)
MSC Markers	CD105	72.1-88.2	6.4-78.5
	CD166	17.3-58.0	4.8-51.5
Hematopoietic Cell Markers	CD14	6.93-10.5	0.9-6.1
	CD45	4.4-9.9	4.6-14.7
Immune co-stimulatory markers	HLA-DR	0-5.6	0-14.7
	CD86	24.3-49.6	4.9-22.5
Trophoblast marker	CD40	7.0-68.7	2-5.8
	Cytokeratin-7	1.36-4.66	2.71-23.07

TABLE 9

FACS analysis of cultured cells (passage 0) from placenta lot D16.		
Cell Surface Marker	Amnion (%)	Chorion (%)
CD45	2.18	0.53
CD166	92.77	82.62
CD105	83.02	86.73
CD49a	92.28	92.26
CD73	89.57	94.57
CD41a	-0.03	-0.05
CD34	-0.23	-0.25
HLA-DR	-0.23	-0.19
CD19	-0.19	-0.22
CD14	-0.25	-0.27
CD90	99.12	98.00

Example 16

Aherence of Cells Derived from Placental Products

[0419] Therapeutic cells, in optional embodiment of the present invention, are adherent, express specific cellular

markers such as CD105 and lack expression of other markers such as CD45, and demonstrate the ability to differentiate into adipocytes, osteoblasts, and

[0420] The expression of specific cellular markers has already been described in Example 15. To determine if the cells within the placental product derived from the chorionic membrane can adhere to plastic and differentiate into one of the lineages, cells were isolated from the placental product derived from the amnion as described in this invention and cultured at 37° C. ± 2° C. and expanded.

[0421] FIG. 10 shows representative images of cells isolated and cultured from amniotic (FIG. 10.A) and chorionic (FIG. 10.B) membranes demonstrating plastic adherence, which is a key feature of MSCs. As a comparison, a representative image of MSCs isolated and expanded from human bone marrow aspirate is also provided (FIG. 10.C). Together, these data show that cells derived from amniotic and chorionic membranes retain a phenotype similar to MSCs after culturing as demonstrated by the cellular markers present in addition to the ability of the cells to adhere to plastic. In conclusion, the presence of MSCs in placental tissues was confirmed by FACS analysis and tissue culture

Example 17

Live CD45+ FACS Analysis

[0422] As CD45 is a general marker for hematopoietic cells and therefore a marker for the presence immunogenic cells, the presence of CD45+ cells may correlate well with how immunogenic a tissue may be. An initial study indeed showed a correlation between amount of immunogenicity as measured via an in vitro MLR assay of placental tissue at various stages within the manufacturing process (as described previously), and the amount of CD45+ cells was determined via FACS analysis. As FIG. 11 demonstrates, membranes that trigger the expression of high levels of IL-2sR on hPBMC responders in MLR also contained a high percentage of CD45+ cells, indicating that immunogenicity of placental membranes can be correlated with the number of CD45+ cells. Further studies revealed, however, that quantifying CD45+ cells via FACS alone showed high variability that did not allow for the establishment of a safety threshold for CD45+ cells in placental membranes. Accordingly, the inventors evaluated whether or not viability of CD45+ cells is correlated with immunogenicity.

[0423] To eliminate some of the variability in CD45+ measurements via FACS, viability of CD45+ cells was assessed, as dead CD45+ cells do not contribute to immunogenicity. To ensure an accurate assessment of live CD45+ cells, a pilot experiment was conducted in which a single cell suspension of amnion membrane was spiked in with a known concentration of live CD45+ cells (hPBMCs) ranging from a theoretical 1.25% to 20% (0.75-12%-actual % of the spiked cells) of the total cell concentration in suspension. Cells were stained with CD45-PE antibody at determined concentrations (refer to Table 10), incubated with 7-AAD cell viability test reagent, and analyzed using a BD FACSCalibur. Table 10 demonstrates that recovery of known amounts of CD45+ cells was not correct (4th column in the table). For example, although 12% of PBMCs was spiked into a single-cell suspension of amnion membrane, only 4.26% of CD45+ cells were recovered according to FACS analysis (>60% difference from the actual spike). To correlate with immunogenicity, MLR was also performed in parallel. Briefly, single cell suspensions of

amniotic membrane spiked with various amounts of live hPBMCs were co-cultured with another donor of PBMCs in the MLR. FIG. 12 depicts a correlation between the amount of CD45+ cells present in amnion-derived cell suspensions and immunogenicity in MLR in vitro. Table 10 and FIG. 12 show that the suspensions spiked with higher amounts of live CD45+ cells resulted in higher immunogenicity as measured by IL-2sR expression on the hPBMC responder donor.

TABLE 10

% CD45+ recovery experiments.				
Sample Description (in % of cell types in the mixture)	% CD45+ cells (detected by FACS)	Actual spike (% based on 60% CD45+ cells in this hPBMC batch)	% Difference from actual spike	Cell suspension immunogenicity (tested in MLR and expressed as IL-2R in pg/mL)
100% amnion	0.65	N/A	N/A	20.23
0% PBMC	N/A	N/A	N/A	15.6 (negative control)
100% PBMC	61.51	N/A	N/A	86.31 (positive control)
20% PBMC + 80% Amnion	4.26	12	64.5%	24.38
10% PBMC + 90% Amnion	2.24	6	62.7%	21.17
5% PBMC + 95% Amnion	1.7	3	43.3%	16.75
2.5% PBMC + 97.5% Amnion	1.36	1.5	Not calculated*	15.9
1.25% PBMC + 98.75% Amnion	1.06	0.75	Not calculated*	12.27

Notes:

N/A—not applicable;

*Not calculated - values are close to the method detection limits

Example 18

Protein Array Analyses

[0424] The protein profiles of amniotic and chorionic membranes were investigated using a SearchLight Multiplex chemiluminescent array. The presence of proteins in tissue membrane extracts and secreted by tissues in culture medium was investigated. For comparison, two commercially available products containing living cells, Apligraf and Dermagraft, were assayed.

Example 18.1

Dermagraft

[0425] Dermagraft membrane was thawed and washed according to the manufacturer's instructions. Dermagraft membrane was cut into 7.5 cm² pieces. For tissue lysates, one 7.5 cm² piece of membrane was snap frozen in liquid nitrogen followed by pulverization using a mortar and pestle. Crushed tissue was transferred to a 1.5 mL microcentrifuge tube and 500 μL of Lysis buffer (Cell Signaling Technologies, Cat No. 9803) with protease inhibitor (Roche, Cat No. 11836153001) was added and incubated on ice for 30 min with frequent vortexing. The sample was then centrifuged at 16000 g for 10

min. The supernatant was collected and sent for protein array analysis by Aushon Biosystems. For tissue culture, one 7.5 cm² piece of membrane was plated onto a well of a 12-well dish and 2 mL of DMEM +1% HSA+ antibiotic/antimycotic were added and incubated at 37° C. ±2° C. for 3, 7, or 14 days. After incubation, tissue and culture media were transferred to a 15 mL conical tube and centrifuged at 2000 rpm for 5 min. Culture supernatant was collected and sent for protein array analysis by Aushon Biosystems.

Example 18.2

Apligraf

[0426] Apligraf membrane was cut into 7.3 cm² pieces. For tissue lysates, one 7.3 cm² piece of membrane was snap frozen in liquid nitrogen followed by pulverization using a mortar and pestle. Crushed tissue was transferred to a 1.5 mL microcentrifuge tube and 500 µL of Lysis buffer (Cell Signaling Technologies, Cat No. 9803) with protease inhibitor (Roche, Cat No. 11836153001) was added and incubated on ice for 30 min with frequent vortexing. The sample was then centrifuged at 16000 g for 10 min. The supernatant was collected and sent for protein array analysis by Aushon Biosystems. For tissue culture, one 7.3 cm² piece of membrane was plated onto a well of a 12-well dish and 2 mL of DMEM +1% HSA+ antibiotic/antimycotic were added and incubated at 37° C. ±2° C. for 3, 7, or 14 days. After incubation, tissue and culture media were transferred to a 15 mL conical tube and centrifuged at 2000 rpm for 5 min. Culture supernatant was collected and sent for protein array analysis by Aushon Biosystems.

Example 18.3

Amniotic and Chorionic Membranes

[0427] Amniotic and chorionic membranes were isolated and packaged at -80° C. ±5° C. according to the manufacturing protocols disclosed herein in Example 3. Packaged membranes were then thawed in a 37° C. ±2° C. water bath and washed 3 times with DPBS. Membranes were cut into 8 cm² pieces. For tissue lysates, one 8 cm² piece of membrane was snap frozen in liquid nitrogen followed by pulverization using a mortar and pestle. Crushed tissue was transferred to a 1.5 mL microcentrifuge tube and 500 µL of Lysis buffer (Cell Signaling Technologies, Cat No. 9803) with protease inhibitor (Roche, Cat No. 11836153001) was added and incubated on ice for 30 min with frequent vortexing. Tissue lysate was then centrifuged at 16000 g for 10 min. The supernatant was collected and sent for protein array analysis by Aushon Biosystems. For tissue culture, one 8 cm² piece of membrane was plated onto a well of a 12-well dish and 2 mL of DMEM +1% HSA+ antibiotic/antimycotic were added and incubated at 37° C. ±2° C. for 3, 7, or 14 days. After incubation, tissue and culture media were transferred to a 15 mL conical tube and centrifuged at 2000 rpm for 5 min. Culture supernatant was collected and sent for protein array analysis by Aushon Biosystems.

[0428] Initial testing consisted of an analysis of 36 proteins that are important for wound healing. The list of identified proteins is described in Table 11.

TABLE 11

List of selected proteins for analysis.		
Protein Group Based on Functionality		Comments
Metalloproteases	Matrix Metalloproteinase 1 (MMP1), MMP2,3,7,8,9,10,13	Matrix and growth factor degradation; facilitate cell migration.
MMP Inhibitors	Tissue Inhibitors of MMPs (TIMP1 and 2)	Have angiogenic activity; can be placed in the "angiogenic factors" group.
Angiogenic Factors	Angiotensin-2 (Ang-2); basic Fibroblast Growth Factor basic (bFGF); heparin-bound Epidermal Growth Factor (HB-EGF); EGF; FGF-7 (also known as Keratinocyte Growth Factor-KGF); Platelet derived Growth Factors (PDGF) AA, AB, and BB; Vascular Endothelial Growth Factor (VEGF), VEGF-C and VEGF-D; Neutrophil gelatinase-associated lipocalin (NGAL); Hepatocyte Growth Factor (HGF); Placenta Growth Factor (PlGF); Pigment Epithelium Derived Factor (PEDF); Thrombopoietin (TPO)	Majority of these factors also have growth and migration stimulatory activities and can be placed in a group of growth factors.
Protease Inhibitor/Protein Carrier	Alpha-2-macroglobulin	Inhibit protease activity; regulate growth factor activity.
Growth Factors	See "angiogenic factors" + Transforming Growth Factor alpha (TGF-a)	See "angiogenic factors."

TABLE 11-continued

List of selected proteins for analysis.		
Protein Group Based on Functionality		Comments
Cytokines	Adiponectin (Acrp-30)	Affect keratinocyte functions.
	Granulocyte Colony-Stimulating Factor (G-CSF)	Protection from infections.
	Interleukin1 Receptor Antagonist (IL-1RA)	Regulate activity of inflammatory cytokine IL-1.
	Leukemia Inhibitory Factor (LIF)	Support angiogenic growth factors.
Chemokines	SDF-1beta	Attracts endothelial and other stem cells from circulation to wound site.
Regulators of IGF	Insulin-like growth factor binding protein (IGFBP1,2,3)	Regulate IGF activity.

Example 18

Protein Expression in Present Placental Products

[0429] Preliminary protein array data analyses showed that the majority of selected testing factors (refer to Table 11) were expressed in amniotic membrane, chorionic membrane, Apligraf, and Dermagraft. Three proteins were identified as unique for the amniotic membrane and/or the chorionic membrane which are undetectable in Apligraf and Dermagraft. These proteins are EGF, IGFBP1, and Adiponectin. All three proteins are important for wound healing. FIG. 13 depicts expression of EGF (A), IGFBP1 (B), and Adiponectin (C) in amniotic or chorionic membranes. AM75 and AM 78 are placental products of the present invention (e.g. cryopreserved), CM75 and CM78 are cryopreserved chorionic membrane products. These proteins are believed by the inventors to facilitate the therapeutic efficacy of the present placental products for wound healing.

[0430] These data indicate that the present methods can produce placental products that comprise an amniotic membrane containing EGF, IGFBP1, and/or adiponectin.

Example 19

Wound Healing Proteins are Secreted for a Minimum of 14 Days

[0431] Placental products of the present invention demonstrate a durable effect, desirable for wound healing treatments. The extracellular matrix and presence of viable cells within the amniotic membrane described in this invention allow for a cocktail of proteins that are known to be important for wound healing to be present for at least 14 days. Amniotic membranes were thawed and plated onto tissue culture wells and incubated at 37° C.±2° C. for 3, 7, and 14 days. At each time point, a sample of the culture supernatant was collected and measured through protein array analysis as described in Example 18. Table 12 illustrates the level of various secreted factors in tissue culture supernatants from two donors of amniotic membranes at 3, 7 and 14 days as measured through protein array analysis.

TABLE 12

Levels of proteins secreted in amnion tissue culture supernatants at different time points (pg/ml).			
	Day 3	Day 7	Day 14
hACRP30	548.03	766.73	371.56
hAlpha2Macroglobulin	69687.55	31764.00	48477.62
hANG2	0.00	9.28	1.65
hEGF	3.06	2.51	2.32
hbFGF	40.80	85.46	269.97
hFibronectin	1932101.25	3506662.00	6019286.50
hHBEGF	41.78	80.50	78.09
hHGF	5358.09	9327.67	18081.16
hIGFBP1	2654.57	6396.11	4666.88
hIGFBP2	4379.76	23797.46	21784.21
hIGFBP3	36030.52	107041.71	13350.99
hIL1ra	116593.20	675.09	4927.52
hKGF	7.29	13.86	36.59
hMMP1	323249.53	1727765.60	15272931.52
hMMP10	14804.44	20557.91	16194.56
hMMP13	92.92	408.17	399.01
hMMP2	38420.90	322500.72	3283119.13
hMMP3	66413.54	283513.74	3598175.53
hMMP7	128.51	147.65	4005.14
hMMP8	463.32	2109.21	2331.47
hMMP9	6139.53	25810.38	60483.67
hNGAL	15754.19	70419.63	721923.09
hPDGFAA	18.02	58.69	16.31
hPDGFAB	16.58	58.41	28.30
hPDGFBB	1.94	21.67	5.84
hPEDF	6793.74	21645.90	169990.84
hSDF1b	0.00	24.09	37.12
hTGfα	15.05	14.89	205.90
hTGfβ1	334.07	341.53	680.33
hTGfβ2	119.59	207.79	731.96
hTIMP1	197743.23	437492.21	247661.65
hTIMP2	4724.25	19970.76	189810.51
hTSP1	0.00	0.00	1274.62
hTSP2	13820.61	59695.21	991366.59
hVEGF	44.98	57.45	7.40
hVEGFC	548.03	766.73	371.56

Example 20

Interferon 2α (IFN-2α) and Transforming Growth Factor-β3 (TGF-β3)

[0432] Placental products described in this invention have been analyzed for the presence of IFN-2α and TGF-β3.

Briefly, after thawing, the membranes were homogenized and centrifuged at 16,000 g to collect the resulting supernatants. Supernatants were analyzed on a commercially available ELISA kit from MabTech (IFN-2 α) and R&D Systems (TGF- β 3). FIG. 14 shows significant expression of IFN-2 α (A) and TGF- β 3 (B) in placental product homogenates.

[0433] Without being bound by theory, interferon-2 α and TGF- β 3 may aid in the prevention of scar and contracture formation. IFN-2 α may serve a role to decrease collagen and fibronectin synthesis and fibroblast-mediated wound contracture.

Example 21

Tissue Reparative Proteins in Amniotic Membranes

[0434] Placental product homogenates were analyzed for the presence of proteins that are important in tissue repair.

[0435] Placental product described in this invention have been analyzed for the presence of tissue reparative proteins. Briefly, the thawed products were incubated in DMEM+10% FBS for 72 hrs. The membranes were then homogenized in a bead homogenizer with the culture media. The homogenates were centrifuged, and the supernatants were analyzed on commercially available ELISA kits from R&D Systems. FIG. 15 shows significant expression of BMP-2, BMP-4, PLAB, PIGF, and IGF-1 in several donors of amniotic membranes.

[0436] Without being bound by theory, the inventors believe that efficacy of the present placental products for wound repair are due, in part, to the role of BMPs, IGF-1, and PIGF in the development and homeostasis of various tissues by regulating key cellular processes. BMP-2 and BMP-4 may stimulate differentiation of MSCs to osteoblasts in addition to promote cell growth; placental BMP or PLAB is a novel member of the BMP family that is suggested to mediate embryonic development. Insulin-like growth factor 1 (IGF-1) may promotes proliferation and differentiation of osteoprogenitor cells. Placental derived growth factor (PIGF) may acts as a mitogen for osteoblasts.

Example 22

MMPs and TIMPs

[0437] Both MMPs and TIMPs are among the factors that are important for wound healing. However, expression of these proteins must be highly regulated and coordinated. Excess of MMPs versus TIMPs is a marker of poor chronic wound healing. We investigated expression of MMPs and TIMPs and its ratio in amniotic membrane and chorionic membrane and compared it to the expression profile in Apligraf and Dermagraft.

[0438] Results showed that all membranes express MMPs and TIMPs; however, the ratio in the thawed placental products and chorionic membranes is significantly lower. Therefore, these membranes will be more beneficial for wound healing (FIG. 16).

[0439] Accumulated data indicate that the MMP to TIMP ratio is higher in cases of non-healing wounds. For example, the ratio between MMP-9 and TIMP1 is approximately 7-10 to one for good healing and 18-20 or higher for poor healing. Analysis of the ratio between MMPs and TIMPs secreted by placental tissues, Apligraf, and Dermagraft showed that the amniotic and chorionic membrane products contain MMPs

and TIMPs at an approximate ratio of 7, which is favorable for wound healing. In contrast, Dermagraft had a ratio >20, and Apligraf had a ratio >200.

[0440] These data indicate that the present methods can produce placental products that comprise an amniotic membrane containing MMP-9 and TIMP1 at a ratio of about 7-10 to one.

Example 23

α 2-Macroglobulin

[0441] α 2-macroglobulin is known as a plasma protein that inactivates proteinases from all 4 mechanistic classes, serine proteinases, cysteine proteinases, aspartic proteinases, and metalloproteinases. Another important function of this protein is to serve as a reservoir for cytokines and growth factors, examples of which include TGF, PDGF, and FGF. In the chronic wounds like diabetic ulcers or venous ulcers, the presence of high amount of proteases leads to rapid degradation of growth factors and delays in wound healing. Thus, the presence of α 2-macroglobulin in products designed for chronic wound healing will be beneficial. Results of the protein array analysis showed that amniotic and chorionic membranes contain α 2-macroglobulin (Table 13). Although these preliminary data show high variability between donors, the importance of this protein in wound healing prompted the additional evaluation of α 2-macroglobulin in placental tissues using a single analyte ELISA instead of protein array, which is a useful tool to evaluate the presence of multiple proteins in one sample for profiling.

[0442] These data indicate that the present methods can produce placental products that comprise an amniotic membrane containing α 2-macroglobulin.

TABLE 13

Expression of α 2-macroglobulin in placental tissue protein extracts.	
Sample	α 2-macroglobulin (pg/mL/cm ²)
AM75	7
CM75	790
AM78	53042
CM78	1014

Example 24

Establishment of EGF as a Marker for Amniotic Tissue Potency

[0443] EGF is among the factors that are important for wound healing (Schultz et al., 1991, Komarcevic, 2000, and Hong et al., 2006). The absence or decreased amount of EGF is one characteristic of chronic wounds (Harding et al., 2002). Evaluation of proteins derived from amniotic membrane samples prepared according to the developed manufacturing process disclosed by the present application reveal that EGF is one of the major factors secreted in higher quantities by these tissues. The importance of EGF for wound healing together with high levels of EGF detected in the presently disclosed amniotic membranes support selection of EGF as a potency marker for evaluation of membrane products manufactured for clinical use pursuant to the present disclosure. A commercially available ELISA kit from R&D Systems was selected for evaluation of its suitability to measure EGF

secreted by amniotic membranes. ELISA qualification meets the standards established by the FDA and ICH guidances for bioanalytical assay validation (Validation of Analytical Procedures: Text and Methodology Q2(R1), 1994; ICH Harmonized Tripartite Guideline and Guidance for Industry Bioanalytical Method Validation, 2001). Amniotic membranes evaluated for expression of EGF by this method confirmed protein array data and further demonstrated that EGF was a unique factor expressed at clinically significant levels in these tissues.

Example 25

Amniotic Tissue Expression of EGF

[0444] Protein array analysis provided initial evidence that EGF was uniquely expressed in amniotic membranes but not in chorionic membranes (Table 14). The levels of EGF measured in amniotic membranes were of clinical significance.

TABLE 14

Protein array data showing range of expression of EGF in amniotic and chorionic membranes from multiple donors.		
	Amnion (pg/ml)	Chorion (pg/ml)
EGF	127.3-361.4	0-0.8

[0445] These data indicate that the present methods can produce placental products that comprise an amniotic membrane containing EGF, optionally in substantial amounts.

Homogenate of Placental Products

[0446] Placental products were thawed until no remaining frozen cryomedia was present. Membranes were then removed from bags and cut into 4 cmx2 cm pieces while still adhered to nitrocellulose. Each piece of tissue was then removed from the nitrocellulose and washed twice with PBS. Each tissue was then snap frozen in a homogenization tube using liquid nitrogen. Subsequently, one pre-cooled 5 mm steel bead was added to each tube; samples were then homogenized using a Qiagen Tissue Lyser according to the manufacturer's recommendations in 500 µL homogenization media. Tissue homogenates were stored at -80° C.±5° C. until analyzed by ELISA for EGF expression.

ELISA Procedure and Validation

[0447] Samples were analyzed for the expression of EGF using the Quantikine Human EGF ELISA Kit (R&D Systems) according to the manufacturer's recommendations. Several parameters were tested to establish the test criteria and to show the suitability of this ELISA kit to measure EGF in placental tissue samples. Assay performance was assessed by analyzing linearity, range, lower and upper limits of quantitation (LLOQ and ULOQ), precision, accuracy, and robustness. Experimental data (Table 14) showed that the quantitation range of this assay was 3.9-250 pg/mL EGF. The intra- and inter-assay CVs ranged from 1.22 to 5.80% and 2.73 to 7.53%, respectively. Additionally, sample recovery analysis demonstrated accuracy within 20%. Furthermore, this assay showed dilutional linearity and specificity. Ruggedness was also demonstrated by assay insensitivity to variations introduced by different analysts.

TABLE 15

Established ELISA parameters for measuring EGF in placenta homogenates.	
Calibration Standard Range	3.9-250 pg/mL
Assay Quantitation Range	7.8-250 pg/mL
LLOQ	7.8 pg/mL
LOD	2.18 pg/mL
ULOQ	250 pg/mL

EGF Expression in Amniotic Membranes

[0448] Measurement of EGF in amniotic preparations has proven to be both reliable and reproducible. Measurement of EGF in multiple donors showed that this method of quantification was a valuable means of evaluating potency in tissue prepared pursuant to the present disclosure for use in a clinical setting. FIG. 17 shows representative expression of EGF in a thawed placental product and chorionic membrane prepared and analyzed by the methods described above. Results have been reproduced in multiple tissue preparations.

[0449] These data indicate that the present methods can produce placental products that comprise an amniotic membrane containing EGF.

Example 26

Placental Tissues Enhance Cell Migration and Wound Healing

[0450] The process of wound healing is highly complex and involves a series of structured events controlled by growth factors (Goldman, 2004). These events include increased vascularization, infiltration by inflammatory immune cells, and increases in cell proliferation. The beginning stages of wound healing revolve around the ability of individual cells to polarize towards the wound and migrate into the wounded area in order to close the wound area and rebuild the surrounding tissue. Upon proper stimulation, several different types of cells including epithelial, endothelial, mesenchymal, and fibroblastic cells are implicated in the wound healing process (Pastar et al, 2008 and Bannasch et al., 2000). Specifically, they proliferate and migrate into the wound area to promote healing. Therefore, experiments were conducted to determine if factors secreted from amniotic and chorionic membranes produced pursuant to the present disclosure promote cell migration and wound field closure. To accomplish this, a commercially available wound healing assay (Cell Biolabs) and a highly accepted human microvascular endothelial cell line (HMVEC, Lonza Inc.) were utilized. Results indicated that cell migration was enhanced by treatment with conditioned media from the placental membranes.

In Vitro Cell Migration Assay

[0451] Human microvascular endothelial cells (HMVECs) were grown under normal cell culture conditions in defined complete media (Lonza Inc.). To assess migration and wound field closure, a commercially available wound healing assay was used (Cell Biolab).

[0452] FIG. 18 depicts the Cell Biolabs 24-well Cytoselect wound healing assay. (Figure reproduced from Cell Biolabs).

[0453] Cells were collected via trypsinization, pelleted, and counted before being resuspended in complete media at a density of 2×10^5 cells/mL. 250 μ L (5×10^4 cells) of cell suspension was then pipetted into each side of a well containing a wound healing insert (Cytoselect 24-well Wound Healing Assay Plate, Cell Biolabs). The cells were grown for 24 hours in complete media. After 24 hours, the wound inserts were removed. At the same time, complete media was removed and replaced with experimental media. Complete media and basal media were used as positive and negative controls, respectively. To generate experimental media, placental membranes were incubated for 3 days in DMEM with 1% human serum albumin (HSA) in a tissue culture incubator. The resulting

[0457] These data demonstrate that the methods of manufacture according to the present invention produce placental products with unexpectedly superior levels of factors that promote wound healing.

Example 27

Biochemical Profile of the Supernatants from Exemplary Placental Tissue Products

[0458] Table 16 depicts the biochemical profile of the supernatants of exemplary placental products of the invention (results adjusted per cm^2 after subtraction of the negative background).

TABLE 16

Factors in Placental Tissue Product (pg/cm^2).							
	Units	Apligraf	DermaGraft	AM75	CM75	AM78	CM78
hMMP1	$\text{pg}/\text{ml}/\text{cm}^2$	1964945.37	14818.20	2821.85	3531.81	117326.89	95.46
hMMP7	$\text{pg}/\text{ml}/\text{cm}^2$	911.54	0.00	0.00	0.00	3.96	0.00
hMMP10	$\text{pg}/\text{ml}/\text{cm}^2$	0.00	0.00	113.94	0.00	0.00	0.00
hMMP13	$\text{pg}/\text{ml}/\text{cm}^2$	21.61	0.00	0.00	0.00	0.71	0.00
hMMP3	$\text{pg}/\text{ml}/\text{cm}^2$	208281.70	180721.52	170.26	161.52	8325.17	0.00
hMMP9	$\text{pg}/\text{ml}/\text{cm}^2$	8872.28	19321.39	214.78	1455.11	630.56	57.59
hMMP2	$\text{pg}/\text{ml}/\text{cm}^2$	153341.77	19712.21	287.14	37.93	3823.38	24.44
hMMP8	$\text{pg}/\text{ml}/\text{cm}^2$	36.92	12.19	0.00	0.00	0.00	0.00
hTIMP1	$\text{pg}/\text{ml}/\text{cm}^2$	2487.18	10909.84	569.23	883.05	28743.48	97.94
hTIMP2	$\text{pg}/\text{ml}/\text{cm}^2$	7285.53	1796.56	89.29	13.72	424.06	4.83
MMP/TIMP		239.26	19.72	6.81	6.26	4.50	2.62

tissue and media were then placed in eppendorf tubes and spun at high speed in a microcentrifuge. The supernatants were collected and stored at $-80^\circ \text{C} \pm 2^\circ \text{C}$. until use. For migration and wound healing studies, conditioned media from placental membranes was diluted 1:20 in basal media before being added to experimental wells. After 18 hours, the media was removed, and the cells were fixed for 20 min in 4% paraformaldehyde and stained with crystal violet. The wound field in each well was then photographed. Wound healing was determined by the amount of wound field still visible at the end of the experiment when compared to control pictures taken before conditioned media was added to the wells.

Placental Membrane Conditioned Media Supports Cell Migration and Wound Field Closure

[0454] Conditioned media from amniotic and chorionic membranes was used to assess the potential for these membranes to promote cell migration and wound field closure. Conditioned media from placental amniotic, chorionic, and a combination of amniotic/chorionic membranes supported migration of cells into the experimental wound field.

[0455] FIG. 19 depicts representative images of cells treated with 5% conditioned media from amniotic, chorionic, or a combination of amniotic/chorionic tissue as well as positive and negative controls. Wound field is 0.9 mm in width.

[0456] The ability of factors from placental membranes produced pursuant to the present disclosure to promote HMVEC migration indicated that these tissues have the ability to enhance wound healing. Additionally, based on the insight of the inventors, it has been surprisingly discovered that these tissues also enhance revascularization since the HMVEC cell line is derived from vascular endothelial cells.

Example 28

Biochemical Profile of the Lysates from Exemplary Placental Tissue Products

[0459] Table 17 depicts the biochemical profile of the lysates of exemplary placental tissue products of the invention (results adjusted per cm^2 after subtraction of the negative background).

TABLE 17

	AM75 lysate pg/ml	AM78 lysate pg/ml	CM75 lysate pg/ml	CM78 lysate pg/ml
hACRP30	50.8	1154.6	1213.7	225.3
hAlpha2-Macroglobulin	1910.6	426191.6	8174.4	9968.6
hEGF	127.3	361.4	0.0	0.8
hbFGF	119.1	821.5	375.0	351.3
hGCSF	0.7	3.2	1.2	0.7
hHBEGF	127.5	168.0	15.4	84.5
hHGF	3943.7	15060.0	29979.6	50392.8
hIGFBP1	5065.0	9456.6	934.0	1443.6
hIGFBP2	12460.8	5569.7	135.9	134.6
hIGFBP3	50115.7	41551.4	4571.5	11970.2
hIL1ra	3881.0	32296.9	5168.2	525.5
hKGF	1.4	8.8	3.1	1.5
hLIF	0.0	4.2	0.0	0.0
hMMP1	9144.1	20641.2	2882.9	6582.3
hMMP10	0.0	15.5	79.3	87.5
hMMP2	2067.3	4061.9	949.5	748.8
hMMP3	0.0	36.2	0.0	0.0
hMMP7	5.1	11.4	4.5	9.1
hMMP8	0.0	0.0	0.0	0.0
hMMP9	92.2	2878.1	2676.2	1259.3
hNGAL	6900.1	6175.9	938.5	229.7
hPDGFAA	0.0	12.5	39.8	35.2
hPDGFAB	11.2	31.3	14.4	14.0
hPDGFbb	4.6	13.4	4.0	1.3

TABLE 17-continued

	AM75 lysate pg/ml	AM78 lysate pg/ml	CM75 lysate pg/ml	CM78 lysate pg/ml
hPEDF	0.0	652.6	0.0	0.0
hTIMP1	7958.1	35955.6	50712.3	17419.9
hTIMP2	3821.8	7443.2	640.7	780.0
hVEGF	3.3	11.8	125.2	8.4
hVEGFC	46.5	150.0	123.5	51.7
hVEGFD	25.7	31.0	15.0	20.4

Example 29

Use of Placental Products for Treating Diabetic Foot Ulcers

[0460] Purpose: Despite bioengineered skin substitutes that contain human fibroblasts or a combination of human fibroblasts and keratinocytes, published rates of chronic wound healing remain low, with approximately half of all wounds recalcitrant to even these newer therapies. Morbidity and mortality from diabetic foot ulceration are substantial as the 5 year mortality rate following a lower extremity amputation is between 39% and 68%. (Page J. J of Foot & Ankle Surgery 2002; 41(4):251-259; Isumi Y., et al. Diabetes Res and Clin Practice 2009; 83:126-131).

[0461] A instant membrane product, which provides necessary angiogenic and anti-inflammatory growth factors was introduced in an effort to improve outcomes of patients with chronic skin ulceration at-risk for amputation.

[0462] Objective: Patients with chronic diabetic foot ulceration, unresponsive to available therapy and at-risk for amputation were considered for treatment. All wounds were aggressively debrided prior to graft application. Patients were evaluated regularly and application of a membrane product of the present invention was at the discretion of the treating physician. Offloading was encouraged in both patients.

Introduction

[0463] According to the United States Food and Drug Administration (FDA), a chronic, cutaneous ulcer is defined as a wound that has failed to proceed through an orderly and timely series of events to produce a durable structural, functional and cosmetic closure(2). The most common chronic wounds include pressure ulcers and leg ulcers. The triad of peripheral neuropathy, deformity, and minor trauma has emerged as the most frequent causes of insult that lead to ulcerations. In terms of healing rates, an appropriate benchmark for a chronic wound is a decrease of 10% to 15% in size every week, or 50% decrease in size over a one-month period. A three-year retrospective cohort study performed by Ramsey et al. of 8,905 patients in a large health maintenance organization who have diabetes reported a 5.8% cumulative incidence of ulceration. At the time of diagnosis, 15% of these patients developed osteomyelitis and 16% required partial amputation of a lower limb.

[0464] Approximately 80% to 85% of lower extremity amputations are preceded by foot ulcerations. Morbidity and mortality from diabetic foot ulceration are substantial as the 5 year mortality rate following a lower extremity amputation is between 39% and 68%. (2). These mortality rates are higher than the five-year mortality rates for breast cancer, colon cancer, and prostate cancer.

[0465] Despite all of the advances in bioengineered tissue for the treatment of chronic diabetic ulcerations, there are an abundance of patients whose ulcerations are resistant to therapy, and result in a chronic wound. Because of healing rates that only approach 50% with these newer therapies, the use of stem cells in regenerative medicine has been of particular interest recently. The ultimate aim is to promote restoration of functional skin. A preliminary study was performed by Fiami et al. in which they isolated mesenchymal stem cells from umbilical cord blood and inoculated them onto a piece of de-epithelialized dermis. The results of this preliminary study showed that peripheral stem cells are capable of surviving and expressing neoangiogenesis. In addition to showing promise for tissue repair, mesenchymal stem cells exhibit low immunogenicity and can be transplanted universally without having to undergo compatibility testing between the donor and recipient.

[0466] In this study, clinical evidence of remarkable healing using an instant membrane product for the treatment of two chronic wounds that amputation was considered. The fundamentals of wound management are still the cornerstone of comprehensive wound care in any treatment protocol including adequate debridement, offloading, maintaining a moist environment, and adequate perfusion and infection control.

Materials

[0467] An instant membrane product was made as taught herein, comprising an allograft derived from the aminon comprising a bilayer of native epithelial cells on a basement membrane and a stoma layer consisting of neonatal fibroblasts, extracellular matrix (ECM) and mesenchymal stem cells (MSC).

Limb Salvage: Case One

History and Physical Examination

[0468] A 70 year old male presented to the emergency department with bulla formation on the dorsolateral aspect of his right foot between the fourth and fifth digits, edema and pain, and a small lesion lateral to the fifth digit. The patient reported a history of minor trauma to the area two weeks prior to presentation. The patient had a history of type II diabetes mellitus, hypertension, heart failure, chronic obstructive pulmonary disease, and chronic kidney disease treated with hemodialysis three times a week. The patient had a surgical history of an aorta-venous graft replacement. He denied any history of alcohol, tobacco or drug use. Physical exam revealed no active purulent drainage or malodor, and no tenderness on palpation. The vascular exam revealed non-palpable pulses in the dorsalis pedis and posterior tibial arteries. Doppler exam revealed a monophasic dorsalis pedis pulse with a biphasic posterior tibial artery pulse. The fifth digit had gangrenous changes and was cold on palpation. There were ischemic changes of the fourth digit. Radiographic evaluation revealed scattered air densities indicative of soft tissue gas in the fourth interspace as well as the tip of the fifth digit.

Preoperative Management

[0469] The patient was started on intravenous antibiotics of vancomycin and piperacillin and tazobactam at appropriate renal dosing.

Operative Management

[0470] The patient was taken to the operating room where an incision and drainage of the fourth interspace was per-

formed, and a partial fifth ray amputation to the level of the metatarsal head was performed without complication. The wound was left open and packed with sterile gauze moistened with sterile normal saline, and covered with a sterile compressive dressing. Intraoperative findings revealed liquefactive necrosis of surrounding tissues with purulence and malodor. The patient underwent 2 subsequent surgical debridements, with the second resulting in further removal of the fourth and fifth metatarsal shafts. In a third surgery further debridement of necrotic soft tissue and amputation of the fourth digit was performed. On May 20, 2010 treatment with an instant membrane product was initiated. Prior to the graft placement the patient had undergone successful recanalization of the popliteal artery and the peroneal artery without significant residual stenosis.

Postoperative Course

[0471] The patient followed up with his podiatric surgeon within 2 days of being discharged from the hospital. Upon initial exam, there were no clinical signs of infection, and the proximal dorsal incision appeared coapted. The third digit was dusky and cool in appearance. Radiographs were taken which showed no evidence of soft tissue gas or acute osteomyelitis. A dry sterile dressing was applied. The patient received applications of the instant membrane product at 6 additional visits in an outpatient office. Prior to each application the wound was evaluated for abscess, cellulitis, drainage, hematoma formation, and infection. At each visit, the wound decreased in size and appeared more granular in nature as compared to previous visits. At the time of the third application the wound had decreased in size 50%.

[0472] At 19 weeks the wound was considered closed, and the patient was instructed to remain weight bearing on the affected limb with the use of a surgical shoe only.

[0473] Photographs of the remarkable wound healing mediated by a placental product of the present invention as shown in FIG. 20. Panel A: First application of an instant membrane product; B: 8 weeks post first an instant membrane product application; C: 10½ weeks post first an instant membrane product application; D: 12 weeks post first an instant membrane product application; E: 19 weeks post first an instant membrane product application.

Limb Salvage: Case Two

History and Physical Examination

[0474] A 44-year old male presented to an outpatient office with a large ulceration on the plantar aspect of his left hallux, secondary to a previous trauma a few weeks prior to the visit. The patient had a history of diabetes mellitus for the past five years complicated by peripheral neuropathy, hypertension, dyslipidemia, and osteomyelitis. Past surgical history included abdominal aortic aneurysm repair and circumcision. On physical exam the ulceration measured 4.0 cm×2.0 cm×1.5 cm, probing to the distal phalanx with exposed tendon. There was no ascending cellulitis or lymphangitis, and no increased temperature gradient. Capillary fill time, hair growth, and tissue turgor were all normal. There were palpable pulses in the dorsalis pedis and the posterior tibial artery. Radiographic exam was negative for soft tissue gas. Magnetic resonance imaging revealed osteomyelitis in the distal aspect of the proximal phalanx and the distal phalanx of

the great toe with a small soft tissue abscess in the region of the dorsal soft tissue adjacent to the distal phalanx.

Preoperative Management

[0475] The patient was started on intravenous antibiotics. He was taken to the operating room for excisional debridement of all nonviable tissue and application of the instant membrane product.

Operative Management

[0476] The ulceration was debrided to healthy tissue with utilization of both sharp dissection and Versajet™, leaving the head of the proximal phalanx exposed plantarly. The instant membrane product was then placed over the wound bed and exposed bone. The patient tolerated the procedure without complication. The patient was discharged from the hospital the day after surgery on a five week course of intravenous antibiotic therapy.

Postoperative Course

[0477] The patient was instructed to remain strictly non-weight bearing to the affected limb, and returned for follow-up on post-operative day 6. The dressing was clean, dry and intact. There were no post-operative complications such as abscess, cellulitis, discomfort, or drainage and no clinical signs of infection. The patient received a total of 7 stem cell graft applications over the course of the next 8 weeks. At each visit the wound was inspected for clinical signs of infection. Evaluation at each visit revealed marked development in granulation tissue to the wound base and significant decrease in size. Eight weeks after the initial application of the allograft tissue the wound was closed.

[0478] Photographs of the remarkable wound healing mediated by a placental product of the present invention as shown in FIG. 21. Panel A: Osteomyelitis, tendon exposed, probed to bone. First stem cell graft was applied after surgical debridement; B: Status post 1 application of stem cell graft, wound is granular in nature and no signs of infection; C: 3 weeks post surgical intervention; 2 applications on the instant membrane product, the wound is considerably smaller in circumference and depth; D: 6 weeks post surgical intervention the wound is almost closed; E: 8 weeks and 7 applications of the instant membrane product, the wound is closed.

CONCLUSION

[0479] Despite the tremendous progress in skin tissue engineering in the past few decades, current therapy has limited efficacy in the treatment of chronic diabetic ulceration. As shown in this case report of two patients, the use of advanced therapies containing stem cells may prove useful to ultimately heal these patients in lieu of amputation, reduce mortality rates, and at the same time be a cost effective alternative to standard treatments currently on the market. Both patients highlighted in this case report received 7 applications of a membrane product of the present invention. Complete healing occurred in both patients. There were no reported complications associated with treatment; the instant membrane product was safe and effective in an initial evaluation of two patients with diabetic foot ulceration at-risk for amputation. These results indicate that patients with recalcitrant, chronic wounds should be considered for this novel therapy.

- 1. A method of manufacturing a placental product comprising the steps of:
 - a. obtaining a placenta;
 - b. selectively depleting the amniotic membrane of a substantial portion of one or more types of immunogenic cells, wherein
 - i. the one or more types of immunogenic cells comprise functional CD14+ macrophages; and
 - ii. the step of selective depletion preserves the viability of a substantial portion of native therapeutic cells selected from MSCs, fibroblasts, epithelial cells, and a combination thereof; and
 - c. cryopreserving the amniotic membrane in a cryopreservation medium; and
 - d. optionally, removing substantially all vascularized tissue or vascularized tissue-derived immunogenic cells from the placenta, or amniotic membrane thereof.
- 2. The method of claim 1, wherein said depleting step comprises removing trophoblasts from the chorionic membrane and retaining the chorionic membrane.
- 3. The method of claim 1, wherein the chorionic membrane is not retained.
- 4. The method of claim 1, wherein said depleting step comprises selectively killing or inactivating the functional CD14+ macrophages.
- 5. The method of claim 4, wherein said depleting step comprises refrigerating the placenta for a period of time before the step of cryopreservation.
- 6. The method of claim 5, wherein the step of refrigerating comprises incubating the placenta in the cryopreservation medium for the period of time.
- 7. The method of claim 6, wherein the period of time is a period of time sufficient to allow the placenta or amniotic membrane thereof to equilibrate.
- 8. The method of claim 5, wherein the period of time is at least: about 10 min, about 20 min, about 30 min, about 40 min, or about 50 min.
- 9. The method of claim 5, wherein the period of time is: about 10 min to about 120 min, about 20 min to about 90 min, or 30 min to about 60 min.
- 10. The method of any of claims 5-9, wherein the refrigerating comprises incubating the placenta for the period of time at a temperature of about -10° C. to about 15° C. or about 2°-8° C.
- 11. The method claim 4, wherein said the depleting step comprises treating the placenta with an anti-TNF- α antibody.
- 12. The method claim 4, wherein said depleting step comprises treating the placenta with IL-10.
- 13. The method of any of the preceding claims, wherein the step of cryopreserving comprises reducing the temperature at a rate of less than any of: about 10° C./min, about 5° C./min, about 3° C./min, about 2° C./min, or about 1° C./min.
- 14. The method of claim 13, wherein the cryopreservation medium comprises a volume of at least any of: about 20 ml, about 30 ml, about 40 ml, about 45 ml, or about 50 ml.
- 15. The method of any of claims 1-12, wherein the cryopreservative comprises a cell-permeating cryopreservative, a non-cell-permeating cryopreservative, or a combination thereof, optionally wherein the permeating cryopreservative comprises:
 - a. DMSO; or
 - b. DMSO in a majority amount.

- 16. The method of claim 15, wherein the cell-permeating cryopreservative:
 - a. does not comprise glycerol in a majority amount; or
 - b. does not comprises a substantial amount of glycerol.
- 17. The method of claim 15, wherein the cell-permeating cryopreservative comprises a DMSO in amount of: about 2% to about 20%, about 5% to about 20%, about 5% to about 15%, or about 7% to about 13%.
- 18. The method of claim 15, wherein the cryopreservation medium further comprises albumin, optionally wherein the albumin is HSA.
- 19. The method of claim 17, wherein the cryopreservation medium further comprises albumin, optionally wherein the albumin is HSA.
- 20. The method of claims 15-19, wherein the step of cryopreserving comprises reducing the temperature at a rate of less than any of: about 10° C./min, about 5° C./min, about 3° C./min, about 2° C./min, or about 1° C./min.
- 21. The method of any of claims 1-12, wherein the step of removing substantially all vascularized tissue or vascularized tissue-derived immunogenic cells from the placenta, or amniotic membrane thereof comprises lysing red blood cells, removing blood clots or a combination thereof from the placenta.
- 22. The method of claim 21, wherein said removing substantially all vascularized tissue or vascularized tissue-derived immunogenic cells comprises treating the placenta with an anticoagulant.
- 23. The method of claim 22, wherein the anticoagulant is a citrate.
- 24. The method of any of claims 1-12 further comprising the step of treating the placenta with one or more antibiotics.
- 25. The method of any of claims 1-12, wherein the method comprises retaining the basement layer, the compact layer, the layer of fibroblast, and the spongy layer of the amniotic membrane.
- 26. The method of claim 25 wherein at least about 15.9% to about 100%, or about 25% to about 100%, or about 35% to about 100%, or about 50% to about 100%, or about 75% to about 100%, or about 90% to about 100%, or about 95% to about 100%, or substantially all amniotic epithelial cells are not removed from the placenta.
- 27. A placental product produced by the method of any of claims 1-26.
- 28. The placental product of claim 27, wherein the membrane is dermatologically acceptable after thawing.
- 29. A method of treating a patient comprising administering to an injury a placental product of claim 28 to the injury.
- 30. The method of claim 29, wherein the injury is a surgical procedure.
- 31. The method of claim 30 wherein the surgical procedure is a tendon repair or an engraftment.
- 32. The method of claim 30, wherein the injury is an ulcer, optionally a foot ulcer, optionally a diabetic foot ulcer.
- 33. The method of any of claims 29-32 wherein the patient has diabetes.
- 34. The method of any of claims 29-32 wherein the injury is associated with ischemia.

* * * * *