



(86) Date de dépôt PCT/PCT Filing Date: 2002/11/29
(87) Date publication PCT/PCT Publication Date: 2003/07/24
(45) Date de délivrance/Issue Date: 2012/05/15
(85) Entrée phase nationale/National Entry: 2004/06/28
(86) N° demande PCT/PCT Application No.: JP 2002/012473
(87) N° publication PCT/PCT Publication No.: 2003/059865
(30) Priorités/Priorities: 2001/12/27 (JP2001-396300);
2002/05/23 (JP2002-149069);
2002/05/23 (JP2002-149078);
2002/06/21 (JP2002-182032)

(51) Cl.Int./Int.Cl. *C07C 229/36* (2006.01),
A23L 1/236 (2006.01), *A23L 2/60* (2006.01),
C07C 227/08 (2006.01), *C07C 229/24* (2006.01),
C07C 251/36 (2006.01), *C07C 51/353* (2006.01),
C07C 51/38 (2006.01), *C07C 59/285* (2006.01),
C07C 59/347 (2006.01), *C07C 59/48* (2006.01),
C07D 209/08 (2006.01), *C07D 209/20* (2006.01)

(72) Inventeurs/Inventors:
KAWAHARA, SHIGERU, JP;
AMINO, YUSUKE, JP;
MORI, KENICHI, JP;
FUNAKOSHI, NAO, JP;
TAKEMOTO, TADASHI, JP

(73) Propriétaire/Owner:

(54) Titre : PROCEDES DE PREPARATION DE COMPOSES D'ACIDE GLUTAMIQUE ET D'INTERMEDIAIRES DESDITS
COMPOSES, ET NOUVEAUX INTERMEDIAIRES UTILISES DANS LESDITS PROCEDES
(54) Title: PROCESSES OF PRODUCING GLUTAMIC ACID COMPOUNDS AND PRODUCTION INTERMEDIATES
THEREFOR AND NOVEL INTERMEDIATE FOR THE PROCESSES

(57) **Abrégé/Abstract:**

A process for preparing industrially and efficiently glutamic acid compounds such as monatin useful as sweeteners or intermediates for the production of drugs or the like a process for preparation of intermediates used therein novel intermediates included among them a process for preparation of optically active monatin a process for preparation of intermediates used therein and novel intermediates included among them.

Specifically, a process for the preparation of glutamic acid compounds or salts thereof including monatin, which comprises preparing a ketoglutaric acid compound serving as a precursor of the objective glutamic acid compound by condensing a specific pyruvic acid compound with oxalacetic acid or pyruvic acid through cross-aldol reaction and, if necessary, subjecting the obtained condensate to decarboxylation, and replacing the carbonyl group of the ketoglutaric acid compound by an amino group.

(73) Propriétaires(suite)/Owners(continued):AJINOMOTO CO., INC., JP

(74) Agent: NORTON ROSE CANADA S.E.N.C.R.L.,S.R.L./LLP

Abstract

A process for preparing industrially and efficiently glutamic acid compounds such as monatin useful as sweeteners or intermediates for the production of drugs or the like a process for preparation of intermediates used therein novel intermediates included among them a process for preparation of optically active monatin a process for preparation of intermediates used therein and novel intermediates included among them.

Specifically, a process for the preparation of glutamic acid compounds or salts thereof including monatin, which comprises preparing a ketoglutaric acid compound serving as a precursor of the objective glutamic acid compound by condensing a specific pyruvic acid compound with oxalacetic acid or pyruvic acid through cross-aldol reaction and, if necessary, subjecting the obtained condensate to decarboxylation, and replacing the carbonyl group of the ketoglutaric acid compound by an amino group.

DESCRIPTION

PROCESSES OF PRODUCING GLUTAMIC ACID COMPOUNDS AND PRODUCTION INTERMEDIATES THEREFOR AND NOVEL INTERMEDIATE FOR THE PROCESSES


Technical Field

The present invention relates to processes of producing glutamic acid compounds typically including monatin, which are useful as production intermediates for sweetener or pharmaceutical products, as well as production intermediates therefor, including a novel important intermediate. More specifically, the invention relates to a process of industrially efficiently producing the glutamic acid compounds, a process of producing production intermediates therefor, including a novel intermediate, and a process of producing optically active monatin, a process of producing production intermediates therefor, including a novel intermediate, and the like.

Background Art

Glutamic acid compounds typically including monatin are compounds promising for use as sweetener or production intermediates for pharmaceutical products and the like. For example, it has been known that 4-hydroxy-4-(3-


indolylmethyl)-2-aminoglutaric acid (3-(1-amino-1,3-dicarboxy-3-hydroxybutan-4-yl)indole, sometimes referred to as "monatin" hereinbelow) represented by the following formula (7') in the (2S,4S) form is contained in the root of a plant *Schlerochiton ilicifolius* and has sweetness at a level several hundreds-fold that of sucrose (see JP-A-64-25757 (US 4,975,298)).




In the specification, the term "monatin" is not limited to the (2S, 4S) form naturally occurring but is used as the generic name of 4-hydroxy-4-(3-indolylmethyl)-2-aminoglutaric acid (3-(1-amino-1,3-dicarboxy-3-hydroxybutan-4-yl)indole) including the individual isomers in the forms of (2S, 4S), (2S, 4R), (2R, 4S), and (2R, 4R).

The following reports have been issued about processes of producing monatin (the following examples (2) to (5)) and protected monatin (the following example (1)).


(1) Process described in Tetrahedron Letters, 2001, Vol. 42, No. 39, pp. 6793-6796;




(2) Process described in Organic Letters, 2000, Vol. 2, No. 19, pp. 2967-2970;



(3) Process described in USP 5,994,559;




(4) Process described in Synthetic Communications, 1994, Vol. 24, No. 22, pp. 3197-3211;



and

(5) Process described in Synthetic Communications, 1993, Vol.

23, No. 18, pp. 2511-2526 and US 4,975, 298 and US 5,128,164;



Because any of the processes requires multiple steps, however, the industrial practice of the processes actually involves much difficulty. Some of the references shown above or other references (see T. Kitahara, et al., Japanese Agrochemical Association, the 2000-th Conference, Abstracts of Proceedings, 3B128β (p.221)) describe the examination of processes of producing optically active monatin. However, disadvantageously, the processes require multiple steps and involve steps which are very tough steps to practise industrially. Thus, it has been desired to develop an industrial process of efficiently producing glutamic acid compounds typically including monatin, particularly an industrial process of efficiently producing optically active monatin.

Disclosure of the Invention

(Problem of the Invention)

The problems to be solved by the invention are to provide processes of industrially and efficiently producing glutamic acid compounds typically including monatin and production

intermediates therefor (including salt forms of them) and to provide important intermediates therefor. More specifically, the invention provides a process of industrially efficiently producing glutamic acid compounds, a process of producing production intermediates therefor, including a novel important intermediate, and a process of producing optically active monatin, a process of producing production intermediates therefor and an important novel intermediate therefor.

(Means for Solving the Problems)

The inventors have made investigations so as to solve the problems. The inventors have found that glutamic acid compounds typically including monatin (including salt forms thereof) can be efficiently produced by condensing a specific pyruvic acid compound and oxalacetic acid or pyruvic acid together with cross aldol reaction to produce ketoglutaric acid compounds as precursors of the intended glutamic acid compounds and then converting the carbonyl group in the resulting ketoglutaric acid compounds to amino group.

In case of aldol reaction using carbonyl compounds of different types as in the invention, generally, four types of products are produced in mixture through the self aldol reaction of the same types of compounds and the cross aldol reaction of different types of compounds. Although the self aldol condensation reaction of oxalacetic acid (Journal of Organic Chemistry, 1973, Vol. 38, No. 20, pp.3582-3585) or



pyruvic acid (Journal of American Chemical Society, 1964, Vol. 86, pp. 2805-2810; Analytical Chemistry, 1986, Vol. 58, No. 12, pp. 2504-2510) is already known, as is cross aldol reaction in a system including a carbonyl compound which does not condense with itself, such as glyoxylic acid or oxalacetic acid, so that a single product can relatively readily be obtained (Tetrahedron Letters, 1987, Vol. 28, pp. 1277-1280), there has been no report of any example of selectively obtaining a single cross aldol reaction product between oxalacetic acid or pyruvic acid and pyruvic acid compounds.




Additionally, the inventors have found that an optically active monatin can be obtained by reacting a glutaric acid compound of the following formula (9) with a specific optically active amine to form a diastereomer salt, then crystallizing and separating the resulting diastereomer salt, further dissociating the diastereomer salt or exchanging the diastereomer salt with a different salt to obtain an optically active glutaric acid compound, then converting the alkoxyimino group (or hydroxyimino group) of the diastereomer salt or the optically active glutaric acid compound to an amino group, crystallizing the resulting monatin represented by the following formula (13) (racemate at the 2-position) in a mixed solvent of water and an organic solvent.

Based on so many findings described above, the invention has been achieved finally.

In other words, the invention includes inventions relating to the following production processes [1] through [23] and the novel substance [24] in their individual various modes.

[1] A process of producing a glutamic acid compound represented by the following formula (7) or a salt thereof, including a step of treating a pyruvic acid compound represented by the following formula (1) and oxalacetic acid represented by the following formula (2) by a cross aldol reaction and decarboxylation reaction or of treating the pyruvic acid compound (except for pyruvic acid) and a pyruvic acid represented by the following formula (2') by a cross aldol reaction, to obtain a ketoglutaric acid compound represented by the following formula (4) or a salt thereof, and a step of converting the carbonyl group of the ketoglutaric acid compound or a salt thereof to an amino group, where the pyruvic acid compound, oxalacetic acid and pyruvic acid may individually be in salt forms thereof:






in the above formulas, R^1 represents a group selected from alkyl groups, aryl groups, aralkyl groups and heterocyclic ring-containing hydrocarbon groups; and R^1 may have at least one substituent selected from halogen atoms, hydroxyl group, alkyl groups with one to 3 carbon atoms, alkoxy groups with one to 3 carbon atoms and amino group.

[2] A process as described above in [1], where the step of converting the carbonyl group of the ketoglutaric acid compound represented by the formula (4) or a salt thereof to an amino group includes a step of reacting an amine compound represented by the following formula (5) or a salt thereof with the

ketoglutaric acid or a salt thereof, to obtain a glutaric acid compound represented by the following formula (6) or a salt thereof, and a step of treating the resulting glutaric acid compound or a salt thereof by reducing reaction:







in the above formulas, R^1 represents a group selected from alkyl groups, aryl groups, aralkyl groups and heterocyclic ring-containing hydrocarbon groups; R^2 represents a hydrogen atom or a group selected from alkyl groups, aryl groups and aralkyl groups; and R^1 may have at least one substituent selected from halogen atoms, hydroxyl group, alkyl groups with one to 3 carbon atoms, alkoxy groups with one to 3 carbon atoms and amino group.

[3] A process as described above in [1], where the step of converting the carbonyl group of the ketoglutaric acid represented by the formula (4) or a salt thereof to an amino group includes a step of treating the ketoglutaric acid compound or a salt thereof by reductive amination.

[4] A process as described above in [1] through [3], where the cross aldol reaction is done within a range of pH 10 to 14.

[5] A process of producing a ketoglutaric acid compound represented by the following formula (4) or a salt thereof,

including a step of treating a pyruvic acid compound represented by the following formula (1) and oxalacetic acid represented by the following formula (2) by a cross aldol reaction and decarboxylation reaction, or of treating the pyruvic acid compound (except for pyruvic acid) and a pyruvic acid represented by the following formula (2') by a cross aldol reaction, where the pyruvic acid compound, oxalacetic acid and pyruvic acid may individually be in salt forms thereof:







in the above formulas, R^1 represents a group selected from alkyl

groups, aryl groups, aralkyl groups and heterocyclic ring-containing hydrocarbon groups; and R^1 may have at least one substituent selected from halogen atoms, hydroxyl group, alkyl groups with one to 3 carbon atoms, alkoxy groups with one to 3 carbon atoms and amino group.

[6] A process as described above in [5], where the cross aldol reaction is done within a range of pH 10 to 14.



[7] A process of producing a glutamic acid compound represented by the following formula (7) or a salt thereof, including a step of reacting a ketoglutaric acid compound represented by the following formula (4) or a salt thereof with an amine compound represented by the following formula (5) or a salt thereof, to obtain a glutamic acid compound represented by the following formula (6) or a salt thereof, and a step of treating the resulting glutamic acid compound or a salt thereof by a reducing reaction:










in the above formulas, R^1 represents a group selected from alkyl groups, aryl groups, aralkyl groups and heterocyclic ring-containing hydrocarbon groups; R^2 represents a hydrogen atom or a group selected from alkyl groups, aryl groups and aralkyl groups; and R^1 may have at least one substituent selected from halogen atoms, hydroxyl group, alkyl groups with one to 3 carbon atoms, alkoxy groups with one to 3 carbon atoms and amino group.

[8] A process of producing a glutamic acid compound represented by the following formula (7) or a salt thereof, including a step of treating a ketoglutaric acid compound represented by the following formula (4) or a salt thereof by a reductive amination reaction:




in the above formulas, R^1 represents a group selected from alkyl groups, aryl groups, aralkyl groups and heterocyclic ring-containing hydrocarbon groups; and R^1 may have at least one substituent selected from halogen atoms, hydroxyl group, alkyl groups with one to 3 carbon atoms, alkoxy groups with one to 3 carbon atoms and amino group.

[9] A process of producing monatin represented by the following formula (7') or a salt thereof, including a step of treating indole-3-pyruvic acid represented by the following formula (1') and oxalacetic acid represented by the following formula (2) by an aldol reaction and a decarboxylation reaction, or of treating indole-3-pyruvic acid represented by the following formula (1') and pyruvic acid represented by the following formula (2') by a cross aldol reaction, to obtain 4-hydroxy-4-(3-indolylmethyl)-2-ketoglutaric acid represented by the following formula (4') or a salt thereof, and a step of converting the carbonyl group of the ketoglutaric acid or a salt thereof to an amino group, where indole-3-pyruvic acid, oxalacetic acid and pyruvic acid may individually be in salt forms thereof:



[10] A process as described above in [9], where the step of converting the carbonyl group of 4-hydroxy-4-(3-

indolylmethyl)-2-ketoglutaric acid represented by the formula (4') or a salt thereof to an amino group includes a step of reacting an amine compound represented by the following formula (5) or a salt thereof with the ketoglutaric acid or a salt thereof to obtain a glutaric acid compound represented by the following formula (6') or a salt thereof, and a step of treating the glutaric acid compound or a salt thereof by a reducing reaction:






in the formula, R^2 represents a hydrogen atom, or a substituent selected from alkyl groups, aryl groups and aralkyl groups.


[11] A process as described above in [9], where the step of converting the carbonyl group of 4-hydroxy-4-(3-indolylmethyl)-2-ketoglutaric acid represented by the formula (4') or a salt thereof to amino group includes a step of treating the ketoglutaric acid compound or a salt thereof by a reductive amination reaction.

[12] A process as described above in [9] to [11], where the

cross aldol reaction is done within a range of pH 10 to 14.


[13] A process of producing 4-hydroxy-4-(3-indolylmethyl)-2-ketoglutaric acid represented by the following formula (4') or a salt thereof, including a step of treating indole-3-pyruvic acid represented by the following formula (1') and oxalacetic acid represented by the following formula (2) by a cross aldol reaction and a decarboxylation reaction, or of treating indole-3-pyruvic acid represented by the following formula (1') and pyruvic acid represented by the following formula (2') by aldol reaction, where indole-3-pyruvic acid, oxalacetic acid and pyruvic acid may individually be in salt forms thereof:







[14] A process as described above in [13], where the cross aldol reaction is done within a range of pH 10 to 14.


[15] A process of producing monatin represented by the following formula (7') or a salt thereof, including a step of reacting 4-hydroxy-4-(3-indolylmethyl)-2-ketoglutaric acid represented by the following formula (4') or a salt thereof with an amine compound represented by the following formula (5) or a salt thereof, to obtain a glutaric acid compound represented by the following formula (6') or a salt thereof, and a step of subsequently treating the glutaric acid compound or a salt thereof by a reducing reaction:







in the formula, R^2 represents hydrogen atom and a substituent selected from alkyl groups, aryl groups and aralkyl groups.

[16] A process of producing monatin represented by the following formula (7') or a salt thereof, including a step of treating 4-hydroxy-4-(3-indolylmethyl)-2-ketoglutaric acid represented by the following formula (4') or a salt thereof by a reductive amination reaction:






[17] A process of producing an optically active monatin represented by the following formula (8) or a salt thereof, including the following steps a to c:




[in the formula, * indicates an asymmetric center and independently represents R- or S-configuration]:

step a: a step of obtaining an optically active glutaric acid compound salt represented by the following formula (11)




[in the formula, R², R³, R⁴, R⁵, R⁶ and R⁷ represent the same meanings as described below; in the formula, * indicates an asymmetric center and independently represents R- or S-configuration] by reacting a glutaric acid compound

represented by the following formula (9)




[in the formula, R^2 represents a hydrogen atom, an alkyl group, an aryl group or an aralkyl group; and the bond marked with a wavy line indicates that both the R-configuration and the S-configuration are included] with an optically active amine represented by the following formula (10)




[in the formula, R^3 , R^4 , R^5 , R^6 and R^7 independently represent a hydrogen atom or an alkyl group with one to 3 carbon atoms; * indicates an asymmetric center and represents R-configuration or S-configuration], to form a diastereomer salt, and a step of separating the diastereomer salt by crystallization;

step b: a step of generating monatin represented by the following formula (13) or a salt thereof




[in the formula, * indicates an asymmetric center and represents R- or S-configuration; and the bond marked with a wavy line means that both the R-configuration and the S-configuration are included], by dissociating the optically active glutaric acid compound salt represented by the formula (11) or exchanging the optically active glutaric acid compound salt with a different salt, as necessary, to prepare an optically active glutaric acid compound represented by the following formula (12) or a salt thereof (excluding the optically active glutaric acid compound salt represented by the formula (11))



[in the formula, R^2 represents hydrogen atom, an alkyl group, an aryl group or an aralkyl group; in the formula, * indicates an asymmetric center and represents the R-configuration or the S-configuration], and a step of converting the alkoxyimino group or hydroxyimino group to an amino group;


step c: a step of obtaining an optically active monatin represented by the formula (8) or a salt thereof by crystallizing monatin represented by the formula (13) or a salt thereof using a mixed solvent of water and an organic solvent.

[18] A process of producing an optically active monatin represented by the following formula (8) or a salt thereof, including the following steps b and c:




[in the formula, * indicates an asymmetric center and independently represents the R- or S-configuration]

step b: a step of generating monatin represented by the following formula (13) or a salt thereof




[in the formula, * indicates an asymmetric center and represents R- or S-configuration; and the bond marked with a wavy line means that both the R-configuration and the S-configuration are included], by dissociating an optically active glutaric acid compound salt represented by the following

formula (11)



[in the formula, R^2 represents hydrogen atom, an alkyl group, an aryl group or an aralkyl group; R^3 , R^4 , R^5 , R^6 and R^7 independently represent hydrogen atom or an alkyl group with one to 3 carbon atoms; and * indicates an asymmetric center and independently represents the R- or S-configuration] or exchanging the optically active glutaric acid compound salt with a different salt on a needed basis, to prepare an optically active glutaric acid compound represented by the following formula (12) or a salt thereof (excluding the optically active glutaric acid compound salt represented by the formula (11))



[in the formula, R^2 represents hydrogen atom, an alkyl group, an aryl group or an aralkyl group; and * indicates an asymmetric center and represents the R-configuration or the S-configuration],


and by treating the optically active glutaric acid compound by

a reaction to convert the alkoxyimino group or hydroxyimino group thereof to an amino group


and

step c: a step of obtaining an optically active monatin represented by the formula (8) or a salt thereof by crystallizing the monatin represented by the formula (13) or a salt thereof using a mixed solvent of water and an alcohol.

[19] A process of producing an optically active monatin represented by the following formula (8) or a salt thereof




[in the formula, * indicates an asymmetric center and independently represents the R- or S-configuration], including a step of crystallizing the salt of monatin represented by the following formula (13) using a mixed solvent of water and alcohol:




[in the formula, * indicates an asymmetric center and represents R- or S-configuration; and the bond marked with a

wavy line means both the R-configuration and the S-configuration are included].


[20] A process of producing an optically active glutaric acid compound salt represented by the following formula (11)



[in the formula, R^2 represents a hydrogen atom, an alkyl group, an aryl group or an aralkyl group; R^3 , R^4 , R^5 , R^6 and R^7 independently represent a hydrogen atom or an alkyl group with one to 3 carbon atoms; and * indicates an asymmetric center and independently represents the R- or S-configuration], including a step of reacting a glutaric acid compound represented by the following formula (9)




[in the formula, R^2 represents hydrogen atom, an alkyl group, an aryl group or an aralkyl group; and the bond marked with a wavy line means that both the R-configuration and S-configuration are included] with an optically active amine represented by the following formula (10)



(10)


[in the formula, R^3 , R^4 , R^5 , R^6 and R^7 independently represent a hydrogen atom or an alkyl group with one to 3 carbon atoms; * indicates an asymmetric center and represents the R- or S-configuration], to form a diastereomer salt, and a step of separating the diastereomer salt by crystallization.

[21] A process of producing an optically active glutaric acid compound represented by the following formula (12) or a salt thereof (excluding the optically active glutaric acid compound salt represented by the formula (11))




(12)

[in the formula, R^2 represents a hydrogen atom, an alkyl group, an aryl group or an aralkyl group; * indicates an asymmetric center and represents the R- or S-configuration], including a step of dissociating an optically active glutaric acid compound salt represented by the following formula (11)




[in the formula, R^2 represents a hydrogen atom, an alkyl group, an aryl group or an aralkyl group; R^3 , R^4 , R^5 , R^6 and R^7 independently represent a hydrogen atom or an alkyl group with one to 3 carbon atoms; and * indicates an asymmetric center and independently represents the R- or S-configuration] or exchanging the optically active glutaric acid compound salt with a different salt.


[22] A process of producing monatin represented by the following formula (13) or a salt thereof:



[in the formula, * indicates an asymmetric center and represents the R- or S-configuration; the bond marked with a wavy line indicates that both the R-configuration and the S-configuration are included], including a step of dissociating an optically active glutaric acid compound salt represented by the following formula (11)




[in the formula, R^2 represents a hydrogen atom, an alkyl group, an aryl group or an aralkyl group; R^3 , R^4 , R^5 , R^6 and R^7 independently represent a hydrogen atom or an alkyl group with one to 3 carbon atoms; and * indicates an asymmetric center and independently represents the R- or S-configuration] or exchanging the optically active glutaric acid compound salt with a different salt on a needed basis, to prepare an optically active glutaric acid compound represented by the following formula (12) or a salt thereof (excluding the optically active glutaric acid compound salt represented by the formula (11))




[in the formula, R^2 represents a hydrogen atom, an alkyl group, an aryl group or an aralkyl group; * indicates an asymmetric center and represents the R-configuration or the S-configuration] and a step of treating the resulting optically active glutaric acid compound by a reaction to convert the alkoxyimino group or hydroxyimino group thereof to an amino





group.





[23] A process of producing monatin represented by the following structural formula (7') (including salt forms thereof), the process passing through a process according to any one of claims 1 to 22:



[24] A compound represented by any one of the following formulas: (4'), (6'), (7''), (11), (12), (14), (15), (16) or (17) (including salt forms thereof), where in the formulas, R^2 represents a hydrogen atom, an alkyl group, an aryl group or an aralkyl group; R^3 , R^4 , R^5 , R^6 and R^7 independently represent a hydrogen atom or an alkyl group with one to 3 carbon atoms; and * indicates an asymmetric center and represents the R- or S-configuration:







In the case where a compound is used or prepared in an appropriate salt form in accordance with the invention, there is no specific limitation on the salt form. Such a salt form includes for example sodium salt, potassium salt, lithium salt, magnesium salt, calcium salt, ammonium salt, and






dicyclohexylammonium salt. By a salt formation process, desalting process, salt exchange process and the like as already known for routine use, the intended salt can be produced.

Mode for Carrying out the Invention

The mode for carrying out the invention is now described in detail below.

(Production of a ketoglutaric acid compound by cross aldol reaction and decarboxylation reaction between a pyruvic acid compound and oxalacetic acid and derivative preparation of glutamic acid compound)

A pyruvic acid compound represented by the following formula (1) and oxalacetic acid represented by the following formula (2) are treated by a cross aldol reaction and decarboxylation reaction or a pyruvic acid compound (excluding pyruvic acid) and pyruvic acid represented by the following formula (2') are treated by aldol reaction, to obtain a ketoglutaric acid compound represented by the following formula (4) or a salt thereof, and then, the carbonyl group of the ketoglutaric acid compound or a salt thereof is converted to an amino group, to produce a glutamic acid compound represented by the following formula (7) or a salt thereof. In this case, the pyruvic acid compound, oxalacetic acid and pyruvic acid may individually be in salt forms.



In the formulas, R^1 represents a group selected from alkyl groups, aryl groups, aralkyl groups and heterocyclic ring-containing hydrocarbon groups. These groups may have at least

one substituent selected from halogen atoms (iodine atom, bromine atom, chlorine atom, fluorine atom, etc.), hydroxyl group, alkyl groups with one to 3 carbon atoms, alkoxy groups with one to 3 carbon atoms and amino group.

As R^1 , alkyl groups, aryl groups, aralkyl groups and heterocyclic ring-containing hydrocarbon groups with one to 11 carbon atoms (never including the carbon number in substituents if these groups contain substituents) are preferable. For example, R^1 includes alkyl groups such as isopropyl group, isobutyl group and 1-methylpropyl group, aryl groups such as phenyl group and 3-indolyl group, aralkyl groups such as benzyl group, 2-phenylethyl group and 2-naphthylmethyl group and heterocyclic ring-containing hydrocarbon groups such as 3-indolylmethyl group and 3-(6-methylindolyl)methyl group.

For the aldol reaction between the pyruvic acid compound represented by formula (1) and the pyruvic acid represented by formula (2'), herein, it is never the case that the pyruvic acid compound represented by formula (1) is pyruvic acid, namely the case that R^1 is a methyl group (an alkyl group with one carbon atom).

Examples of R^1 with substituents include R^1 having an aromatic ring or heterocyclic ring, provided that the aromatic ring or heterocyclic ring contains at least one substituent selected from alkyl groups with one to 3 carbon atoms, alkoxy groups with one to 3 carbon atoms, and amino groups. When a

benzyl group or 3-indolylmethyl group for example is selected as R^1 in the formula, specifically, the benzene ring or indole ring contained in the group may contain at least one substituent selected from halogen atoms (iodine atom, bromine atom, chlorine atom, fluorine atom, etc.), hydroxyl group, alkyl groups with one to 3 carbon atoms, alkoxy groups with one to 3 carbon atoms, and amino group.

In the case where R^1 is a 3-indolylmethyl group, in other words where indole-3-pyruvic acid (formula 1') is used as the pyruvic acid compound, 4-hydroxy-4-(3-indolylmethyl)-2-ketoglutaric acid (formula 4') or a salt thereof can be obtained, and is an intermediate important for monatin production. Subsequently, by converting the carbonyl group of the ketoglutaric acid or a salt thereof to an amino group, monatin (formula 7') or a salt thereof can be produced.

(Cross aldol reaction)

The cross aldol reaction is preferably carried out under alkaline conditions. The pyruvic acid compound and oxalacetic acid, or the pyruvic acid compound (excluding pyruvic acid) and pyruvic acid may be present in an appropriate solvent for the reaction.

As the reaction solvent, polar solvents such as water, methanol, ethanol, propanol, acetonitrile and dimethylformamide or mixed solvents thereof are preferable.

Particularly, water and a mixed solvent (hydrous organic solvent) of water and polar solvents are preferable.

The pH of the solvent is within a range of preferably 10 to 14, more preferably 10.5 to 14, still more preferably 11 to 13.

When the pH is too high, the yield is likely to decrease. When the pH is too low, secondary reactions are likely to take place during the cross aldol reaction.

Bases may satisfactorily be used to achieve these pH under alkaline conditions, and include, for example, inorganic bases such as alkali metal salts and alkali earth metal salts including alkali earth metal hydroxides and carbonates, e.g., lithium hydroxide, sodium hydroxide, potassium hydroxide, sodium carbonate, potassium carbonate and calcium carbonate, and organic bases such as triethylamine.

Though the amount of oxalacetic acid or pyruvic acid to be used to the amount of the pyruvic acid compound has no specific limitation and the reaction yield is likely to be improved when pyruvic acid is used in excess, generally one to 10 equivalents and preferably 3 to 6 equivalents are used to one equivalent of the pyruvic acid compound.

The reaction can be carried at a reaction temperature within a range of preferably -10 to 70°C, more preferably 10 to 50°C. When the reaction temperature is too low, the intended reaction progresses so slowly that secondary reactions are

likely to occur. When the reaction temperature is high, the intended ketoglutaric acid compound (or a salt thereof) is likely to be decomposed.

The reaction time has no specific limitation, and is generally one to 72 hours, preferably 3 to 24 hours.

(Decarboxylation reaction)

The reaction employing oxalacetic acid is then progressed on to a decarboxylation reaction, to subsequently obtain the intended ketoglutaric acid compound (or a salt thereof). The decarboxylation reaction of the condensate from the aldol reaction between oxalacetic acid and the pyruvic acid compound can be achieved by a spontaneous decarboxylation reaction. However, the decarboxylation reaction can be effectively carried out by adding an acid or a metal ion or both to the reaction solution. The acid for use then includes for example hydrochloric acid, sulfuric acid, phosphoric acid, acetic acid, p-toluenesulfonic acid, solid acids such as ion exchange resins, while the metal ion includes for example transition metal ions such as nickel ion, copper ion, and iron ion. As the reaction temperature, preferably about -10 to 100°C, more preferably about 0 to 60°C can be selected.

The reaction solution after the cross aldol reaction or after the cross aldol reaction and decarboxylation reaction may satisfactorily be used for the subsequent step as it is. From

the reaction solution the ketoglutaric acid compound (or a salt thereof) represented by the formula (4) is isolated and purified, for use in the subsequent step. When the subsequent amination step is continuously carried out, generally, the ketoglutaric acid compound (or a salt thereof) need not be isolated. After the completion of the reaction, the reaction solution is concentrated or distilled off, if necessary, for the amination step. By using the same solvent as used in the cross aldol reaction step for the amination step, the following step can be carried out with no distillation or substitution of reaction solvent, or the like. In the case where the ketoglutaric acid compound represented by formula (4) is obtained as a salt, the salt is prepared in a free form by a method known to persons skilled in the art, for use in the amination step. However, generally, it is not necessary to do so. The salt can be used in its salt form.

In the cross aldol reaction (and decarboxylation reaction if necessary) in accordance with the invention where R^1 is a 3-indolylmethyl group, i.e. where indole-3-pyruvic acid (formula 1') is used as the pyruvic acid compound, 4-hydroxy-4-(3-indolylmethyl)-2-ketoglutaric acid (formula 4'), an intermediate important for monatin production, or a salt thereof, can be produced.



(Conversion of carbonyl group to amino group)

After the cross aldol reaction is carried out (and

subsequently the decarboxylation reaction is carried out if necessary), the carbonyl group of the ketoglutaric acid compound represented by formula (4) or a salt thereof is converted to an amino group, to produce a glutamic acid compound represented by formula (7). The reaction to convert the carbonyl group to an amino group is with no specific limitation and is carried out for example by the following methods.

(Conversion example 1 of carbonyl group to amino group)

After the cross aldol reaction is carried out (and subsequently the decarboxylation reaction is carried out if necessary), an amine compound (which may be in a salt form) represented by the following formula (5) is reacted with the ketoglutaric acid compound represented by formula (4) or a salt thereof, to produce a glutaric acid compound represented by formula (6) or a salt thereof, which is then treated by a reducing reaction to produce a glutamic acid compound represented by formula (7).



In the formula, R^1 is as described above.

In the case where R^1 is a 3-indolylmethyl group, i.e. in the case where 4-hydroxy-4-(3-indolylmethyl)-2-ketoglutaric acid (formula 4') is used as the ketoglutaric acid represented by formula (4) or a salt thereof, herein, a glutaric acid compound represented by formula (6') or a salt thereof is produced, which is then treated by a reduction reaction to produce monatin represented by formula (7') or a salt thereof.

In the formula, R^2 represents a hydrogen atom or a group selected from alkyl groups, aryl groups and aralkyl groups and the like. R^2 is preferably selected from a hydrogen atom and alkyl groups and aralkyl groups with 7 or less carbon atoms.

Specifically, R^2 is preferably a hydrogen atom, methyl group or benzyl group, particularly preferably a hydrogen atom. In other words, specific examples of the amine compound represented by formula (5) preferably include hydroxylamine, methoxyamine and benzyloxyamine, particularly preferably include hydroxylamine.

The salt of the amine compound represented by formula (5) includes salt forms of the amine compound with organic acids or inorganic acids and specifically includes for example hydroxylamine hydrochloride salt, hydroxylamine sulfate salt and methoxyamine hydrochloride salt.

When hydroxylamine hydrochloride reacts with 4-hydroxy-4-(3-indolylmethyl)-2-ketoglutaric acid represented

by the formula (4'), for example, the corresponding 4-hydroxy-4-(3-indolylmethyl)-2-hydroxyiminoglutaric acid (the compound of formula (6') where R^2 is hydrogen atom) is obtained in good yield.

For the reaction of the ketoglutaric acid compound represented by the formula (4) or a salt thereof with the amine compound represented by the formula (5) or a salt thereof, the reaction temperature can be set at preferably about -10 to 100°C, more preferably about 0 to 60°C. Further, the reaction time can be set at preferably about one to 100 hours, more preferably about 1 to 24 hours.

For the reaction of the amine compound or a salt thereof, the pH of the reaction solution is preferably 2 or more because the reaction progresses slowly when the pH is too low. The reaction can be progressed more preferably at about pH 2 to 13, still more preferably at about pH 4 to 12.

The ratio of the amine compound or a salt thereof to be used with not particularly limited. However, the amine compound (or a salt thereof) is used preferably at about one to 7 moles, more preferably at about one to 2 moles per one mole of the ketoglutaric acid compound (or a salt thereof) represented by formula (4).



As the reaction solvent, polar solvents such as water, methanol, ethanol, propanol, acetonitrile and dimethylformamide, or mixed solvents thereof are preferable.

Particularly, water and a mixed solvent of water with polar solvents (hydrous organic solvents) are preferable.

As the glutaric acid compound (or a salt thereof) represented by the formula (6) thus obtained, the reaction solution may satisfactorily be used as it is for the subsequent step. Otherwise, the glutaric acid compound may be isolated and purified from the reaction solution, prior to use in the next step.

Where isolation and purification is carried out, methods known to a person skilled in the art such as extraction and crystallization can appropriately be used. For example, in the case of the glutaric acid compound represented by formula (6') by acidifying the reaction solution by adjusting the pH of the reaction solution with acids such as hydrochloric acid, extracting the glutaric acid compound in organic solvents such as ethyl acetate, concentrating the resulting organic layer and crystallizing the residue in a mixed solvent of aqueous ammonia and alcohol, the glutaric acid compound represented by formula (6') can be obtained as the diammonium salt thereof in crystal. Using ion exchange resins or adsorption resins or the like, the compound in the free form can be isolated from the reaction solution. The glutaric acid compound represented by formula (6') or a salt thereof as obtained in such manner is generally a racemate, from which the optically active form can be obtained by a method described below.

Then, the glutaric acid compound represented by the formula (6) (or a salt thereof) is treated by a reduction reaction, to produce a glutamic acid compound represented by the following formula (7). Through the reaction, the alkoxyimino group (or hydroxyimino group) at the 2-position in the glutaric acid compound represented by the formula (6) can be converted to amino group.



In the formulas, R^1 and R^2 are as described above.

The reduction of the alkoxyimino group (or hydroxyimino group) to an amino group can preferably be done by a hydrogenation reaction using a catalyst for hydrogenation.

As the catalyst for hydrogenation, there can be used palladium-type catalysts (palladium carbon, etc.), platinum-type catalysts (platinum carbon, etc.), rhodium-type catalysts (rhodium carbon, etc.), ruthenium-type catalysts

(ruthenium carbon, etc.), nickel-type catalysts (Raney-nickel, etc.) and the like.

These catalysts are used within a range of preferably 0.1 to 20 mol%, more preferably 0.5 mol% to 5 mol% of the substrate.

As the reaction solvent, polar solvents such as water, methanol, ethanol, propanol, acetonitrile and dimethylformamide or mixed solvents thereof are preferable. Particularly, water and a mixed solvent (hydrous organic solvent) of water with polar solvents are preferable.

The reaction of this step is preferably done under alkaline conditions, generally within a range of pH 7 to 14, preferably within a range of pH 8 to 12. In the case of using rhodium-type catalysts (rhodium carbon, etc.), in particular, the reaction is done generally within a range of pH 7.5 to 11, preferably within a range of pH 8 to 10. In the case of using nickel-type catalysts (Raney nickel, etc.), however, the reaction preferably progresses under neutral conditions, generally within a range of pH 5 to 9, preferably within a range of pH 6.5 to 7.5. If pH is too high, the reaction is likely to give rise to an increase of by-products. If pH is too low, the reaction is likely to progress slowly. When the reaction is carried out under alkaline conditions, the type of an alkali for use in pH adjustment is not specifically limited. For reduction reactions using rhodium-type catalysts and palladium-type catalysts, aqueous ammonia is used for the


reaction, and is particularly preferable because of the increase of the yield and low by-products.


The hydrogenation reaction is preferably done in a hydrogen atmosphere. As the hydrogen pressure, a range of preferably 0.5 to 100 atmospheres, more preferably 3 to 70 atmospheres is desirable for the reaction.

The reaction temperature is within a range of preferably -20 to 100°C, more preferably 0 to 70°C. The reaction time can be 6 to 24 hours.

(Conversion example 2 of carbonyl group to amino group)

By converting the carbonyl group at the 2-position in the ketoglutaric acid compound represented by the following formula (4) (or a salt thereof) by a reductive amination reaction using amines such as ammonia, benzylamine and 1-phenylethylamine, a glutamic acid compound represented by the following formula (7) can be obtained.





In the formulas, R^1 is as described above.

When R^1 is a 3-indolylmethyl group, i.e. in the case where 4-hydroxy-4-(3-indolylmethyl)-2-ketoglutaric acid (formula 4') is used as the ketoglutaric acid compound represented by formula (4) or a salt thereof, herein, monatin represented by formula (7') or a salt thereof can be produced.

The amine can be used within a range of preferably one to 10 equivalents to the ketoglutaric acid compound (or a salt thereof). When ammonia is to be used as the amine, herein, a large excess of ammonia is preferably used.

As the reducing catalyst, hydride catalysts such as NaBH_4 in addition to the hydrogenation catalysts described above can be used. In the case of a hydride catalyst, the catalyst can be used at an amount generally within a range of 0.5 to 2 equivalents. In the case of a hydrogenation catalyst, the catalyst at an amount similar to the amount used for contact hydrogenation of the glutaric acid compound represented by the formula (6) can be used. The reaction can preferably be progressed at a reaction temperature within a range of preferably 0 to 50°C , more preferably 20 to 35°C . The reaction


time is within a range of preferably 1 to 72 hours. Where a hydrogenation catalyst is used, the reaction can be done at a hydrogen pressure within 1 to 15 atmospheres.

As the reaction solvent, polar solvents such as water, methanol, ethanol, propanol, acetonitrile and dimethylformamide or mixed solvents thereof are preferable. Particularly, water and a mixed solvent (hydrous organic solvent) of water with polar solvents are preferable.

The glutaric acid compound (or a salt thereof) represented by formula (7) as obtained by the processes in the two examples can be isolated and purified, using methods known to a person skilled in the art, such as extraction and crystallization. Where R^1 is 3-indolylmethyl group, i.e. in the case where 4-hydroxy-4-(3-indolylmethyl)-2-ketoglutaric acid (formula 4') is used as the ketoglutaric acid compound represented by formula (4) or a salt thereof, herein, monatin represented by formula (7') or a salt thereof can be produced. Monatin represented by formula (7') or a salt thereof can be isolated and purified by a method for producing optically active monatin as described below, to obtain the optically active form.

(Production of optically active monatin)


Monatin has asymmetric carbon atoms at the 2- and 4-positions, so the following four types of optical isomers exist.




When 4-hydroxy-4-(3-indolylmethyl)-2-ketoglutaric acid (formula 4') is used as the ketoglutaric acid compound represented by formula (4) or a salt thereof as described above, reaction with an amine compound represented by formula (5) or a reagent generating the compound or a reagent being capable of generating the compound can produce the glutaric acid compound represented by the formula (6') or a salt thereof, which is generally a racemate.

The racemate or the glutaric acid compound containing the R-form and S-form at an appropriate ratio (these are included in the glutaric acid compound represented by the formula (9)) is treated by the following steps a to c, to obtain optically active forms of monatin or a salt thereof.


step a: a step of obtaining an optically active glutaric acid compound salt represented by the following formula (11)



[in the formula, R^2 , R^3 , R^4 , R^5 , R^6 and R^7 represent the same meanings as described below; and * indicates an asymmetric center and independently represents R- or S-configuration], by reacting a glutaric acid compound represented by the following formula (9)




[in the formula, R^2 represents hydrogen atom, an alkyl group, an aryl group or an aralkyl group; and the bond marked with a wavy line expresses that both the R-configuration and the S-configuration are included] with an optically active amine represented by the following formula (10)




[in the formula, R^3 , R^4 , R^5 , R^6 and R^7 independently represent

a hydrogen atom or an alkyl group with one to 3 carbon atoms;
 and * indicates an asymmetric center and represents R-configuration or S-configuration], to form a diastereomer salt and then separating the diastereomer salt by crystallization;
 step b: a step of generating monatin represented by the following formula (13) or a salt thereof



[in the formula, * indicates an asymmetric center and represents R- or S-configuration; and the bond marked with a wavy line means that both the R-configuration and the S-configuration are included], by dissociating the optically active glutaric acid compound salt represented by formula (11) or exchanging the optically active glutaric acid compound salt with a different salt as necessary, to prepare an optically active glutaric acid compound represented by the following formula (12) or a salt thereof (excluding the optically active glutaric acid compound salt represented by the formula (11))



[in the formula, R^2 represents a hydrogen atom, an alkyl group, an aryl group or an aralkyl group; and * indicates an asymmetric center and represents the R-configuration or the S-configuration], and subsequently converting the alkoxyimino group or hydroxyimino group to an amino group;

step c: a step of obtaining an optically active monatin represented by formula (8) or a salt thereof by crystallizing monatin represented by formula (13) or a salt thereof using a mixed solvent of water and an organic solvent.

The step a is now described.

So as to allow the glutaric acid compound represented by the formula (9) to form a diastereomer salt together with the optically active amine represented by the formula (10), for example, such compounds or salts thereof are dissolved in a solvent for the reaction. If the glutaric acid compound is in a salt form thereof, the salt is converted to the free form by neutralizing the salt with an acid as necessary and subsequently extracting the free form into an organic solvent, and then, the free form reacts with an optically active amine to form the salt. Additionally, an acid may satisfactorily be added to a solvent containing the glutaric acid compound salt dissolved therein to neutralize the salt, and then, an optically active amine may be added for reaction to form the diastereomer salt. When the glutaric acid compound salt represented by the formula (11) is formed by a salt exchange

reaction with an optically active amine represented by the formula (10) in a solvent, the salt form can even be used as is to react with the optically active amine represented by the formula (10). In this case, the optically active amine represented by the formula (10) is preferably used in a salt form such as the hydrochloride salt or sulfate salt.

Particularly preferable examples of the optically active amine represented by the formula (10) include (R)-(+)-1-phenylethylamine and (S)-(-)-1-phenylethylamine, where R^3 , R^4 , R^5 , R^6 and R^7 in the formula are hydrogen atoms.



The optically active amine is used at an amount of preferably about 0.1 to 1-fold mole, more preferably about 0.3 to 0.6-fold that of the glutaric acid compound.

The reaction temperature is set within a range of preferably about -20 to 100°C, more preferably about 0 to 60°C. The reaction time is not particularly limited but is short enough for rapid formation of the salt.

The reaction solvent includes a single solvent selected from water, methanol, ethanol, acetonitrile, toluene and ethyl acetate, and an appropriate mixed solvent of two or more thereof. Particularly, water or a mixed solvent of water and an organic solvent miscible with water (for example, polar solvents such as methanol, ethanol and acetonitrile) is preferably used. Of these, a single solvent of water is more preferable.

After the completion of the reaction, for example, the

reaction solution is concentrated if necessary, and water is added to crystallize the diastereomer salt. The reaction solution may be cooled if required. Because water is a poor solvent of the resulting diastereomer salt, water or a mixed solvent of water with an organic solvent miscible with water is used as the reaction solvent for forming the diastereomer salt, so that the crystals can be deposited and crystallized, concurrently with the progress of the reaction (salt formation). The crystals obtained by crystallization are separated from the reaction solution by filtration and the like, to obtain the diastereomer salt represented by the formula (11). The diastereomer salt obtained as crystals in the case of using water as the poor solvent varies depending on the steric configuration of the optically active amine used. When using (R) - (+) - 1-phenylethylamine as the optically active amine represented by the formula (10) for the 4-hydroxy-4-(3-indolylmethyl)-2-hydroxyiminoglutaric acid (a compound of the formula (9) where R^2 is hydrogen atom), for example, the diastereomer salt represented by the following formula (14) can be obtained. In case of using (S) - (-) - 1-phenylethylamine, the diastereomer salt represented by the following formula (15) can be obtained.



In the case of using (R) - (+) -1-phenylethylamine, in other words, a crystal of the diastereomer salt can be obtained, where the glutaric acid compound in the S-configuration at the 4-position forms the salt together with (R) - (+) -1-phenylethylamine. In the case of using (S) - (-) -1-phenylethylamine, a crystal of the diastereomer salt can be obtained, where the glutaric acid compound in the R-configuration at the 4-position forms the salt together with (S) - (-) -1-phenylethylamine. A person skilled in the art can select an optically active amine suitable for the intended compound, to form and crystallize such diastereomer salt, so that a diastereomer salt with the desired steric configuration can be obtained.

Further, the mother liquor, after the separation of these

diastereomer salts as crystals, contains, as its main component, a glutaric acid compound in a steric configuration opposite to the glutaric acid compound separated as crystals. Thus, by adding an optically active amine with a steric configuration opposite to that of the optically active amine used to form the salt in the mother liquor, to form and crystallize the diastereomer salt in the same manner as described above, one additional diastereomer salt crystal can be separated from the mother liquor. In other words, the step a in accordance with the invention can be applied to the mother liquor to obtain another diastereomer salt by crystallization as described above.

Step b is now described below.

The optically active glutaric acid compound salt represented by formula (11) as obtained in step a is dissociated to form an optically active glutaric acid compound represented by the formula (12), if necessary. Alternatively, the salt can be replaced by a different salt (salt exchange), if necessary.

Methods known to persons skilled in the art can be used to effect such dissociation and replacement with a different salt (salt exchange). The dissociation method is for example a method including a step of dissolving or suspending the salt in water, alcohol or a mixed solvent thereof, a step of neutralizing the resulting solution or suspension with acids

such as hydrochloric acid or sulfuric acid, and an extraction step into an organic solvent, a method including a step of dissolving the salt in water, and separating the free form represented by formula (12) with ion exchange resins or adsorption resins. For isolation of the free form, the intended isolation can readily be carried out by a known method, for example distillation of a resin eluent solution, or of an extract solution containing the free form, under reduced pressure. In case of substitution with a different salt, for example, the optically active glutaric acid compound salt represented by formula (11) can be dissolved in aqueous alkali metal solutions of sodium hydroxide and potassium hydroxide and aqueous ammonia solution, from which the free optically active amine is extracted in an organic solvent, so that the salt can be exchanged. Using a known method such as distillation under reduced pressure of the aqueous solution after extraction or crystallization, additionally, the salt of the optically active glutaric acid compound represented by the formula (12) (excluding the optically active glutaric acid compound salt represented by formula (11)) can be isolated.

Monatin represented by formula (13) can be produced from the optically active glutaric acid compound represented by formula (12) thus obtained or from a salt thereof by treating the alkoxyimino group (or hydroxyimino group) thereof with a reaction to convert the group to an amino group. Further,

monatin represented by formula (13) can be produced by treating the alkoxyimino group (or hydroxyimino group) of the optically active glutaric acid compound salt represented by formula (11) with a reaction to convert the group to an amino group in the same manner as described above.

As described above, the conversion of the alkoxyimino group (or hydroxyimino group) to amino group can be done by a hydrogenation reaction using a catalytic hydrogenation reagent, under reaction conditions as described above.

After the completion of the reaction, the catalyst is removed by filtration and the like and the filtrate is concentrated, if necessary, from which monatin represented by the formula (13) can be obtained by isolation methods known to a person skilled in the art (for example, crystallization, HPLC, etc.). If the next step c is to be continuously done, additionally, monatin represented by the formula (13) need not be isolated, in general. After the completion of the reaction, the catalyst is removed from the reaction solution by filtration and the like. Then, the reaction solution is concentrated or distilled off, if necessary, in order to carry out the crystallization according to the step c. By using the same solvent as used as the crystallization solvent for the hydrogenation reaction, the next step c can be carried out without distillation of the reaction solvent or solvent replacement or the like. When a base is used for the

hydrogenation reaction, monatin in the reaction solution exists in a salt form thereof. By treating the reaction solution after removal of the catalyst with an ion exchange resin and the like, for example, monatin can be prepared as the free form or can be converted to a different salt (salt exchange including for example the conversion of an ammonium salt to a sodium salt or potassium salt or the like). Then, the resulting product can be crystallized. For carrying out the next step c, preferably, the resulting monatin is used in the salt form thereof as it is.

The step c is now described.

Although the monatin represented by formula (13) or a salt thereof as obtained at the step b can retain the optical activity at the 4-position, the resulting monatin can be recovered as a mixture of the S form and R form at the 2-position. The monatin and a salt thereof can be optically resolved by crystallization according to the step c described next, so that optically active monatin or a salt thereof, optically active at both the 2- and 4-positions, can be obtained.

The monatin represented by formula (13) or a salt thereof is treated by a crystallization step using a mixed solvent of water and an organic solvent, to obtain optically active monatin (crystal) represented by formula (8).


When a base is used for the hydrogenation reaction at the

step (b), generally, monatin can be obtained in a salt form thereof, which is preferably treated at a crystallization step, as it is in the salt form. In this case, water is a good solvent for the monatin salt. The crystallization method is not particularly limited, and includes for example methods known to a person skilled in the art, such as crystallization under cooling and concentration prior to crystallization. For crystallizing the monatin salt, for example, the monatin crystal in the free form can be obtained by adding an acid to an aqueous solution containing the monatin salt dissolved therein to neutralize the solution and by adding an organic solvent to the resulting solution. Because monatin is likely to be decomposed by acid, however, the invention can preferably be used particularly in the case where monatin is to be obtained in a salt form.

As the organic solvent, an organic solvent miscible with water can be used. Particularly, alcohols such as methanol, ethanol, propanol and isopropanol are preferable. A mixed solvent of different two types or more organic solvents may satisfactorily be used as the organic solvent. The ratio of an organic solvent and water in the mixed solvent with water may satisfactorily be set preferably within a range of organic solvent: water = about 1:0.1 to 1:1 in volume ratio, more preferably within a range of organic solvent: water = about 1:0.3 to 1:0.9 in volume ratio. The crystallization

temperature may satisfactorily be set within a range of preferably about -20 to 100°C, more preferably about 0 to 60°C.

As shown below in the following schemes, the steric configuration of the monatin crystal obtained at step c is as follows. In case of using monatin represented by formula (13) in the R form at the 4-position and in the S form at the 4-position, the monatin crystals (2R, 4R) and (2S, 4S) are obtained, respectively. Additionally, the mother liquor after crystal separation individually contain (2S, 4R) monatin and (2R, 4S) monatin as the main component. By treating the mother solutions with adsorption resins and the like, (2S, 4R) monatin or (2R, 4S) monatin can be isolated.



If desired, the optically active monatin obtained in the free form can be prepared as a salt form. By methods known to persons skilled in the art (salt formation), the monatin can be prepared in salt forms, for example as a sodium salt or

potassium salt. Further, even the optically active monatin obtained in a salt form thereof can be obtained similarly in the free form if desired or can be converted to a different salt. By utilizing methods known to a person skilled in the art, for example, the salt can be converted to the free form by a method for converting salts to free forms via dissociation and the like. Alternatively, the salt can be converted to a different salt by exchanging the resulting salt with a different salt (salt exchange).

Preferable embodiments for carrying out the Invention

The invention is now described in detail in the following Examples. However, the invention is not limited to the Examples.

In the Examples, further, the optical purity was assayed by HPLC under the following conditions.

<Column for separating optical isomers>

SUMICHIRAL OA-7100 manufactured by Sumika Chemical Analysis Service

<Eluent>

20 mM phosphate buffer, pH 2.8: acetonitrile = 7:3

<Column temperature>

10°C; and

<Flow rate>

0.6 ml/min.

<Example 1>

Synthesis of 4-hydroxy-4-(3-indolylmethyl)-2-ketoglutaric acid; NO. 1

After 8.28 g of potassium hydroxide (having a purity of 85 % by weight) was dissolved in 27 ml of water, 3.0 g (14.76 mmol) of indole-3-pyruvic acid and 5.85 g (44.29 mmol) of oxalacetic acid were added to the resulting solution, for reaction at ambient temperature for 72 hours (about pH 13 at the start of the reaction). An ion exchange resin (AmberliteTM IR 120B H AG) was added to the reaction solution to adjust the solution to pH 3.0, for extraction into 200 ml of ethyl acetate at 0°C. 100 ml of saturated aqueous sodium bicarbonate was added to the resulting ethyl acetate layer, and ethyl acetate in the ethyl acetate layer was distilled off, and pH of the solution was re-adjusted to 7.9 with an ion exchange resin (IRA400 OH AG manufactured by Organo Corporation). The resulting solution was freeze-dried as it was. 4-Hydroxy-4-(3-indolylmethyl)-2-ketoglutarate sodium salt was obtained as a crude product. Then, 40 ml of water and 200 ml of ethanol were added to the resulting residue, in which the solid was filtered off. The resulting mother solution was concentrated to dryness, to obtain 4-hydroxy-4-(3-indolylmethyl)-2-ketoglutarate sodium salt of 1.5 g as a crude product.

<Example 2>

Synthesis of 4-hydroxy-4-(3-indolylmethyl)-2-ketoglutaric acid; NO. 2

After 18.91 g (286.5 mmol) of potassium hydroxide (at a content of 85 % by weight) was dissolved in 64.45 ml of water, 7.50 g (35.8 mmol at a content of 97.0 % by weight) of indole-3-pyruvic acid and 14.18 g (107.4 mmol) of oxalacetic acid were added to and dissolved in the resulting solution (about pH 13 at the start of the reaction). The dissolved mixture was stirred at 35°C for 24 hours. Afterwards, 40.0 ml of 3N hydrochloric acid was added for neutralization (pH = 7.0), to obtain 153.5 g of a neutralized reaction solution. The neutralized reaction solution contained 5.55 g of 4-hydroxy-4-(3-indolylmethyl)-2-ketoglutaric acid at a yield of 53.5 % (vs. indole-3-pyruvic acid). Water was added to the neutralized reaction solution to make up 168 ml, and the solution passed through a resin column (diameter of 4.8 cm) packed with a synthetic adsorbent (DIAION-SP207TM manufactured by Mitsubishi Chemical Corporation) of 840 ml. Then, pure water was passed through the column at a flow rate of 23.5 ml per minute. 1.73 to 2.55 L/L-R were collected, to obtain an aqueous solution containing 3.04 g of 4-hydroxy-4-(3-indolylmethyl)-2-ketoglutaric acid - a yield of 54.7 % (relative to the amount with which the resin was charged). (NMR measurement)

$^1\text{H-NMR}$ (400MHz, D_2O) : δ 3.03 (d, 1H, $J=14.6$ Hz) , 3.11 (d, 1H, $J=14.6$ Hz) , 3.21 (d, 1H, $J=18.1$ Hz) , 3.40 (d, 1H, $J=18.1$ Hz) , 7.06 - 7.15 (m, 3H) , 7.39 (d, 1H, $J=7.8$ Hz) , 7.66 (d, 1H, $J=7.8$ Hz).

$^{13}\text{C-NMR}$ (400MHz, D_2O) : δ 35.43, 47.91, 77.28, 109.49, 112.05, 119.44, 119.67, 121.91, 125.42, 128.41, 136.21, 169.78, 181.43, 203.58.

(Molecular weight measurement)

Theoretical ESI-MS value $\text{C}_{14}\text{H}_{13}\text{NO}_6 = 291.07$

Analytical value = 290.02 (MH^-)

<Example 3>

Synthesis of 4-hydroxy-4-(3-indolylmethyl)-2-ketoglutaric acid; NO. 3

After 3.70 g (56.0 mmol) of potassium hydroxide (at a content of 85 % by weight) was dissolved in 72.1 ml of water, 0.81 g (4.0 mmol) of indole-3-pyruvic acid and 3.17 g (24.0 mmol) of oxalacetic acid were added to and dissolved in the resulting solution (about pH 13 at the start of the reaction). The dissolved mixture was stirred at 35°C for 24 hours. A part of the reaction solution was treated with hydroxylamine to prepare 4-hydroxy-4-(3-indolylmethyl)-2-hydroxyiminoglutaric acid, which was analyzed by HPLC. Consequently, it was found that 4-hydroxy-4-(3-indolylmethyl)-2-ketoglutaric acid was generated at a yield of 76.6 % (vs. indole-3-pyruvic acid).

<Example 4>

Synthesis of 4-benzyl-4-hydroxy-2-ketoglutaric acid

After 16.23 g of potassium hydroxide (having a purity of 85 % by weight) was dissolved in 48 ml of water, 5.0 g (30.5 mmol) of phenylpyruvic acid and 12.1 g (91.4 mmol) of oxalacetic acid were added to the resulting solution, for reaction at ambient temperature for 72 hours (about pH 13 at the start of the reaction). Using conc. hydrochloric acid, the reaction solution was adjusted to pH 2.2, and extracted in ethyl acetate. The organic layer was rinsed in aqueous saturated sodium chloride, dried over anhydrous magnesium sulfate and concentrated to obtain the residue. The residue was recrystallized in ethyl acetate and toluene, to obtain 4-benzyl-4-hydroxy-2-ketoglutaric acid of 2.8 g (11.3 mmol) in crystal.

(NMR measurement)

^1H NMR (D_2O) δ : 2.48 (d, $J=14.4$ Hz, 0.18H) , 2.60 (d, $J=14.4$ Hz, 0.18H) , 2.85–3.30 (m, 3.64H) , 7.17–7.36 (m, 5H) .

(Molecular weight measurement)

Theoretical ESI-MS value $\text{C}_{12}\text{H}_{12}\text{O}_6 = 252.23$

Analytical value = 251.22 (MH^-)

<Example 5>

Synthesis of 4-hydroxy-4-(3-indolylmethyl)-2-

hydroxyiminoglutaric acid; NO. 1

After 13.8 g of potassium hydroxide (having a purity of 85 % by weight) was dissolved in 50 ml of water, 5.0 g (24.6 mmol) of indole-3-pyruvic acid and 9.8 g (73.8 mmol) of oxalacetic acid were added to the resulting solution, for reaction at ambient temperature for 72 hours (about pH 13 at the start of the reaction). To the reaction solution was added 6.8 g (98.4 mmol) of hydroxylamine hydrochloride salt. Then, the reaction solution was adjusted to pH 7.5 with aqueous 4N sodium hydroxide solution. After the reaction solution had been stirred for 24 hours at ambient temperature, the reaction solution was adjusted to pH 2.6 with 6N hydrochloric acid. After extraction using ethyl acetate, the organic layer was rinsed in aqueous saturated sodium chloride, dried over anhydrous magnesium sulfate and concentrated to dryness. The resulting residue was dissolved in 10 ml of aqueous 14 % ammonia, followed by gradual dropwise addition of 70 ml of ethanol, and stirred at ambient temperature for 3 hours. The resulting slurry was filtered. The resulting crystals were dried to obtain 2.7 g (7.9 mmol) of 4-hydroxy-4-(3-indolylmethyl)-2-hydroxyiminoglutaric acid in the form of an ammonium salt.

(NMR measurement)

^1H NMR (DMSO- d_6) δ : 2.66 (s, 2H) , 2.89 (d, $J=14.4$ Hz, 1H) , 3.04 (d, $J=14.4$ Hz, 1H) , 6.89–6.94 (m, 1H) , 6.97–7.03 (m, 1H) , 7.11 (d, $J=2.8$ Hz, 1H) , 7.27 (d, $J=7.8$ Hz, 1H) , 7.53 (d, $J=7.8$ Hz, 1H) , 10.71 (br s, 1H) .

(Molecular weight measurement)

Theoretical ESI-MS value $\text{C}_{14}\text{H}_{14}\text{N}_2\text{O}_6 = 306.28$

Analytical value = 305.17 (MH^-)

<Example 6>

Synthesis of 4-hydroxy-4-(3-indolylmethyl)-2-methoxyiminoglutaric acid

After 9.12 g (138.1 mmol) of potassium hydroxide (at a content of 85 % by weight) was dissolved in 23 ml of water, 2.55 g (12.2 mmol at a content of 97.0 % by weight) of indole-3-pyruvic acid and 7.46 g (56.5 mmol) of oxaloacetic acid were added to and dissolved in the resulting solution (about pH 13 at the start of the reaction). The solution was stirred at 35°C for 24 hours. To the reaction solution was gradually added 5.76 g (69 mmol) of methoxyamine hydrochloride salt while the reaction solution was adjusted to around pH 10 with aqueous 25 % sodium hydroxide solution. After the reaction had continued at ambient temperature for 14 hours, the reaction solution was adjusted to pH 2.23, using 6N hydrochloric acid, and was then extracted in ethyl acetate. The organic layer was rinsed with aqueous saturated sodium chloride and dried over anhydrous

magnesium sulfate. After the magnesium sulfate had been filtered off, the resulting solution was concentrated to obtain 4.66 g of residue. The resulting residue was approximately purified by silica gel column chromatography, and further coarsely purified by preparative thin layer chromatography (PTLC; ethyl acetate/hexane/acetic acid = 5/5/1), to obtain 0.93 g (2.92 mmol) of the title compound 4-hydroxy-4-(3-indolylmethyl)-2-methoxyiminoglutaric acid; yield 24 % (vs. indole-3-pyruvic acid)).

(NMR measurement)

$^1\text{H-NMR}$ (400MHz, DMSO- d_6) : δ 2.89 (d, $J=14.9$ Hz, 1H) , 3.04 (s, 2H) , 3.15 (d, $J=14.9$ Hz, 1H) , 3.90 (s, 3H) , 6.91 - 6.96 (m, 1H) , 6.98 - 7.04 (m, 1H) , 7.09 - 7.12 (m, 1H) , 7.29 (d, $J=7.4$ Hz, 1H) , 7.50 (d, $J=7.4$ Hz, 1H) , 10.80 (br s, 1H) .

<Example 7>

Synthesis of 4-benzyl-4-hydroxy-2-hydroxyiminoglutaric acid

After 16.23 g (having a purity of 85 % by weight) of potassium hydroxide was dissolved in 45 ml of water, 5.0 g (30.5 mmol) of phenylpyruvic acid and 12.1 g (91.4 mmol) of oxalacetic acid were added to the resulting solution, for reaction at ambient temperature for 24 hours (about pH 13 at the start of the reaction). To the reaction solution was added 8.5 g (121.8 mmol) of hydroxylamine hydrochloride salt, for reaction at ambient temperature for 72 hours. The reaction solution was

adjusted to pH 2.6, using 6N hydrochloric acid, and was then extracted in ethyl acetate. The organic layer was rinsed with aqueous saturated sodium chloride, dried over anhydrous magnesium sulfate and concentrated to dryness. The resulting residue was recrystallized in 20 ml of ethyl acetate and 80 ml of toluene, to obtain 4.0 g (15.1 mmol) of 4-benzyl-4-hydroxy-2-hydroxyiminoglutaric acid.

(NMR measurement)

^1H NMR (DMSO- d_6) δ : 2.80 (d, $J=13.9$ Hz, 1H) , 2.99 (d, $J=12.7$ Hz, 1H) , 3.01 (d, $J=13.9$ Hz, 1H) , 3.03 (d, $J=12.7$ Hz, 1H) , 7.13 – 7.25 (m, 5H).

(Molecular weight measurement)

Theoretical ESI-MS value $\text{C}_{12}\text{H}_{13}\text{NO}_6 = 267.24$

Analytical value = 266.12 (MH^-)

<Example 8>

Synthesis of 4-hydroxy-4-(3-indolylmethyl)-2-aminoglutaric acid (monatin); NO. 1

0.13 g (0.38 mmol) of 4-hydroxy-4-(3-indolylmethyl)-2-hydroxyiminoglutaric acid ammonium salt was dissolved in 5 ml of aqueous 28 % ammonia, to which 0.09 g of 5 % rhodium carbon was added, for reaction at ambient temperature and a hydrogen pressure of 7.5 atmospheres. 14 hours later, the catalyst was filtered off, and the resulting solution was concentrated to dryness, to obtain a mixture of 0.075 g (0.23 mmol) of the ammonium salt of (2S, 4S)/(2R, 4R)-4-hydroxy-4-(3-

indolylmethyl)-2-aminoglutaric acid (monatin) and 0.036 g (0.11 mmol) of the ammonium salt of (2S, 4R)/(2R, 4S)-4-hydroxy-4-(3-indolylmethyl)-2-aminoglutaric acid (monatin). (NMR measurement)

^1H NMR (D_2O) δ : 2.05 (dd, $J=12.2, 15.1\text{Hz}$, 0.67H) , 2.21 (dd, $J= 9.9, 15.6\text{ Hz}$, 0.33H) , 2.48 (dd, $J= 3.2, 15.6\text{ Hz}$, 0.33H) , 2.68 (dd, $J= 2.2, 15.1\text{ Hz}$, 0.67H) , 3.08 (d, $J= 14.4\text{ Hz}$, 0.67H) , 3.17 – 3.25 (m, 0.66H) , 3.28 (d, $J= 14.4\text{ Hz}$, 0.67H) , 3.63 (dd, $J= 2.2, 12.2\text{ Hz}$, 0.67H) , 3.98 (dd, $J=3.2, 9.9\text{Hz}$, 0.33H) , 7.12 – 7.18 (m, 1H) , 7.19 – 7.26 (m, 2H) , 7.45 – 7.51 (m, 1H) , 7.70 – 7.76 (m, 1H) .

(Molecular weight measurement)

Theoretical ESI-MS value $\text{C}_{14}\text{H}_{16}\text{N}_2\text{O}_5 = 292.29$

Analytical value = 291.28 (MH^-)

<Example 9>

Synthesis of 4-hydroxy-4-(3-indolylmethyl)-2-aminoglutaric acid (monatin); NO. 2

0.264 g (0.824 mmol) of 4-hydroxy-4-(3-indolylmethyl)-2-methoxyiminoglutaric acid was dissolved in 10 ml of aqueous 28 % ammonia. 0.18 g of 5 % rhodium carbon (dry product) was added and the mixture stirred at a hydrogen pressure of 7.5 atmospheres pressures for 18 hours. The catalyst was filtered off and the solvent was distilled off under reduced pressure, to obtain the residue. The resulting residue was analyzed by NMR, and a mixture of 0.115 g (0.395

mmol; yield 48 %) of (2S, 4S)/(2R, 4R)-4-hydroxy-4-(3-indolylmethyl)-2-aminoglutaric acid (monatin) and 0.065 g (0.223 mmol; yield 27 %) of (2S, 4R)/(2R, 4S)-4-hydroxy-4-(3-indolylmethyl)-2-aminoglutaric acid was found to have been generated.

<Example 10>

Synthesis of 4-hydroxy-4-(3-indolylmethyl)-2-aminoglutaric acid (monatin); NO. 3

1.0 g (2.94 mmol) of 4-hydroxy-4-(3-indolylmethyl)-2-hydroxyiminoglutarate ammonium salt was dissolved in 10 ml of water, to which 1 ml of a RaneyTM-nickel catalyst (manufactured by Kawaken Fine Chemicals Co., Ltd.; developed nickel catalyst NDHT-90) was added with a syringe, and the mixture stirred at a hydrogen pressure of 20 atmospheres for 10 hours. The catalyst was filtered off, and the resulting solution was concentrated to obtain the residue. The residue was analyzed by NMR. It was shown that 0.29 g (0.89 mmol; yield, 30 %) of (2S, 4S)/(2R, 4R)-4-hydroxy-4-(3-indolylmethyl)-2-aminoglutaric acid (monatin) and 0.29 g (0.89 mmol; yield, 30 %) of (2S, 4R)/(2R, 4S)-4-hydroxy-4-(3-indolylmethyl)-2-aminoglutaric acid were generated.

<Example 11>

Synthesis of 2-amino-4-benzyl-4-hydroxyglutaric acid; NO.1

0.25 g (0.94 mmol) of 4-benzyl-4-hydroxy-2-hydroxyiminoglutaric acid was dissolved in 10 ml of aqueous 50 % methanol solution, to which 0.5 ml of aqueous 28 % ammonia was added. 1.0 g of 5 % palladium-carbon (50 % hydrous product) was added, for reaction at ambient temperature and a hydrogen pressure of 7.7 atmospheres. 72 hours later, the catalyst was filtered off, and the reaction solution was concentrated to dryness, to obtain 0.10 g (0.35 mmol) of the ammonium salt of (2S, 4S)/(2R, 4R)-2-amino-4-benzyl-4-hydroxyglutaric acid and 0.10 g (0.35 mmol) of the ammonium salt of (2R, 4S)/(2S, 4R)-2-amino-4-benzyl-4-hydroxyglutaric acid as a mixture. (NMR measurement)

^1H NMR (D_2O) δ : 1.94 (dd, $J=11.9, 15.3$ Hz, 0.5H) , 2.10 (dd, $J= 10.2, 15.3$ Hz, 0.5H) , 2.36 (dd, $J= 3.1, 15.3$ Hz, 0.5H) , 2.56 (dd, $J= 2.4, 15.3$ Hz, 0.5H) , 2.81 (d, $J= 13.6$ Hz, 0.5H) , 2.94 (d, $J= 13.5$ Hz, 0.5H) , 3.01 (d, $J= 13.5$ Hz, 0.5H) , 3.06 (d, $J= 13.6$ Hz, 0.5H) , 3.55 (dd, $J= 2.4, 11.9$ Hz, 0.5H) , 3.88 (dd, $J= 3.1, 10.2$ Hz, 0.5H), 7.17 – 7.31 (m, 5H) .

(Molecular weight measurement)

Theoretical ESI-MS value $\text{C}_{12}\text{H}_{15}\text{NO}_5 = 253.26$

Analytical value = 252.23 (MH^-)

<Example 12>

Synthesis of 2-amino-4-benzyl-4-hydroxyglutaric acid; NO.2

0.13 g (0.52 mmol) of 4-benzyl-4-hydroxy-2-ketoglutaric acid and 0.11 ml (1.0 mmol) of benzylamine were dissolved in

5 ml of methanol, to which 0.1 g of 5 % palladium carbon (50 % hydrous product) was added, for reaction under a hydrogen atmosphere at ambient temperature and atmospheric pressure. Two days later, the catalyst was filtered off, and the reaction solution was concentrated to dryness, to obtain 0.03 g (0.12 mmol) of (2S, 4S)/(2R, 4R)-2-amino-4-benzyl-4-hydroxyglutaric acid and 0.06 g (0.24 mmol) of (2R, 4S)/(2S, 4R)-2-amino-4-benzyl-4-hydroxyglutaric acid as a mixture.

<Example 13>

Synthesis of 4-hydroxy-4-(4-hydroxyphenylmethyl)-2-hydroxyiminoglutamic acid

To 10 ml of water, in which were dissolved 3.18 g of potassium hydroxide, were added 1.0 g (5.55 mmol) of 4-hydroxyphenylpyruvic acid and 2.2 g (16.7 mmol) of oxalacetic acid, for reaction at ambient temperature for 72 hours (about pH 13 at the start of the reaction). Hydroxylamine hydrochloride salt (1.54 g (22.2 mmol)) was added to the reaction solution, for reaction at ambient temperature for 10 hours. The reaction solution was then adjusted to pH 2.6, using 6N hydrochloric acid, for extraction using ethyl acetate. The organic layer was rinsed with aqueous saturated sodium chloride, dried over anhydrous magnesium sulfate and concentrated to dryness, to obtain 0.7 g (2.47 mmol) of 4-hydroxy-4-(4-hydroxyphenylmethyl)-2-hydroxyiminoglutamic acid as a crude

product. Further, the crude product was recrystallized in methanol and toluene, to obtain 0.22 g (0.78 mmol) of 4-hydroxy-4-(4-hydroxyphenylmethyl)-2-hydroxyiminoglutaric acid in crystals.

(NMR measurement)

^1H NMR (DMSO- d_6) δ : 2.67 (d, $J=13.7$ Hz, 1H), 2.89 (d, $J=13.7$ Hz, 1H), 2.95 (d, $J=12.5$ Hz, 1H), 2.99 (d, $J=12.5$ Hz, 1H), 6.59 (d, $J=8.0$ Hz, 2H), 6.97 (d, $J=8.0$ Hz, 2H), 9.11 (br s, 1H).

(Molecular weight measurement)

Theoretical ESI-MS value $\text{C}_{12}\text{H}_{13}\text{NO}_6 = 283.24$

Analytical value = 281.93 (MH^-)

<Example 14>

Synthesis of 4-hydroxy-4-(4-hydroxyphenylmethyl)-2-hydroxyiminoglutarate acid

0.06 g (0.21 mmol) of 4-hydroxy-4-(4-hydroxyphenylmethyl)-2-hydroxyiminoglutarate acid was dissolved in 2.5 ml of aqueous 28 % ammonia, to which 0.04 g of 5 % rhodium carbon was added for reaction at ambient temperature and a hydrogen pressure of 7.5 atmospheres. 14 hours later, the catalyst was filtered off, and the resulting solution was concentrated to dryness, to obtain a mixture of 0.044 g (0.145 mmol) of (2S, 4S)/(2R, 4R)-4-hydroxy-4-(4-hydroxyphenylmethyl)-2-aminoglutaric acid and 0.021 g (0.069 mmol) of (2S, 4R)/(2R, 4S)-4-hydroxy-4-(4-

hydroxyphenylmethyl)-2-aminoglutaric acid.

^1H NMR (D_2O) δ : 1.89 (dd, $J=11.9, 15.7$ Hz, 0.68H) , 2.06 (dd, $J= 10.2, 15.0$ Hz, 0.32H) , 2.30 (dd, $J= 3.3, 15.0$ Hz, 0.32H) , 2.51 (dd, $J= 2.4, 15.7$ Hz, 0.68H) , 2.70 (d, $J= 13.4$ Hz, 0.68H) , 2.83 (d, $J= 13.4$ Hz , 0.32H) , 2.90 (d , $J= 13.4$ Hz , 0.32H) , 2.96 (d, $J= 13.4$ Hz, 0.68 H) , 3.52 (dd, $J= 2.4, 11.9$ Hz, 0.68H) , 3.84 (dd, $J=3.3, 10.2$ Hz, 0.32H) , 6.71 – 6.77 (m, 2H) , 7.02 – 7.08 (m, 2H).

(Molecular weight measurement)

Theoretical ESI-MS value $\text{C}_{12}\text{H}_{15}\text{NO}_6 = 269.26$

Analytical value = 268.11 (MH^-)

<Example 15>

Synthesis of 4-hydroxy-4-(3-indolylmethyl)-2-ketoglutaric acid; NO. 4

12.30 g (58.7 mmol; at a purity of 97.0 % by weight) of indole pyruvic acid was added to and dissolved in 209 ml of water containing 2.45 g of sodium hydroxide dissolved therein. Over a period of two hours, 47.61 g of an aqueous 25 % by weight sodium hydroxide solution and a mixture of 25.85 g (293.5 mmol) of pyruvic acid and 25.85 g of water were added to the resulting solution under a nitrogen atmosphere at 35°C , while the reaction system was kept at pH 11.0. Subsequently, the reaction system was agitated for 14 hours. In this way, a reaction solution was obtained, which contained 4-hydroxy-4-(3-indolylmethyl)-2-ketoglutaric acid at a yield of 44.1 % (vs.

indolepyruvic acid). 3.60 g of 1N hydrochloric acid was added to the solution for neutralization (pH = 6.91), to obtain 275ml of a neutralized reaction solution.

168 ml of the thus obtained neutralized reaction solution was passed through a resin column (having a diameter of 4.8 cm) packed with 840 ml of a synthetic adsorbent (DIAION-SP207TM manufactured by Mitsubishi Chemical Corporation). Then, pure water was passed through the column at a flow rate of 23.5 ml per minute, to collect 1.7 to 2.9 L/L-R to obtain 4-hydroxy-4-(3-indolylmethyl)-2-ketoglutaric acid at high purity at a yield of 66.3 %.

(NMR spectrum)

¹H-NMR(400MHz, D₂O): δ 3.03 (d, 1H, J = 14.6 Hz), 3.11(d, 1H, J = 14.6 Hz), 3.21(d, 1H, J= 18.1 Hz), 3.40 (d, 1H, J = 18.1 Hz), 7.06-7.15 (m, 3H), 7.39 (d, 1H, J = 7.8 Hz), 7.66 (d, 1H, J = 7.8 Hz).

¹³C-NMR(400MHz, D₂O): δ 35.43, 47.91, 77.28, 109.49, 112.05, 119.44, 119.67, 121.91, 125.42, 128.41, 136.21, 169.78, 181.43, 203.58.

(Mass analysis)

Theoretical ESI-MS value C₁₄H₁₃NO₆ = 291.07

Analytical value = 290.02 (MH⁻)

<Example 16>

Synthesis of 4-hydroxy-4-(3-indolylmethyl)-2-hydroxyiminoglutaric acid; NO. 2

After 1.0 g (4.92 mmol) of indole-3-pyruvic acid was added to and dissolved in 10 ml of aqueous saturated sodium carbonate solution, the resulting solution was adjusted to pH 12.55 using aqueous 25 % sodium hydroxide solution. After 1.3 g (14.8 mmol) of pyruvic acid was added, the resulting solution was adjusted to pH 12.6 using aqueous 25 % sodium hydroxide solution, for reaction at ambient temperature for 2 hours, to obtain a reaction solution containing 4-hydroxy-4-(3-indolylmethyl)-2-ketoglutaric acid. 1.37 g (19.7 mmol) of hydroxylamine hydrochloride salt was added to the reaction solution, while the reaction solution was adjusted to a pH value around neutrality using aqueous 25 % sodium hydroxide solution, and then stirred at ambient temperature for 4 hours. Using conc. hydrochloric acid, the reaction solution was adjusted to an acidic pH value, to extract the organic matter in ethyl acetate. The organic layer was rinsed with aqueous saturated sodium chloride, dried over anhydrous magnesium sulfate and subsequently concentrated, to obtain the residue. The residue was recrystallized in aqueous 28 % ammonia and ethanol, to obtain 0.52 g (1.5 mmol) of 4-hydroxy-4-(3-indolylmethyl)-2-hydroxyiminoglutaric acid in crystals (a yield of 31 % vs. indole-3-pyruvic acid).

(NMR spectrum)

^1H NMR (DMSO- d_6) δ : 2.66 (s, 2H) , 2.89 (d, $J=14.4$ Hz, 1H) , 3.04 (d, $J=14.4$ Hz, 1H) , 6.89 – 6.94 (m, 1H) , 6.97 – 7.03 (m, 1H) , 7.11 (d, $J=2.8$ Hz, 1H) , 7.27 (d, $J=7.8$ Hz, 1H) , 7.53 (d, $J=7.8$ Hz, 1H) , 10.71 (br s, 1H) .

(Mass analysis)

Theoretical ESI-MS value $\text{C}_{14}\text{H}_{14}\text{N}_2\text{O}_6 = 306.28$

Analytical value = 305.17 (MH^-)

<Example 17>

Synthesis of 4-hydroxy-4-(3-indolylmethyl)-2-hydroxyiminoglutaric acid; NO. 3

After 10.0 g (49.2 mmol) of indole-3-pyruvic acid was added to and dissolved in 98 ml of aqueous saturated sodium carbonate solution, the resulting solution was adjusted to pH 12.4 using aqueous 25 % sodium hydroxide solution. After 16.3 g (147.6 mmol) of sodium pyruvate was added for reaction at ambient temperature for 2 hours, a reaction solution containing 4-hydroxy-4-(3-indolylmethyl)-2-ketoglutaric acid was obtained. 13.7 g (197 mmol) of hydroxylamine hydrochloride salt was added to the reaction solution while the reaction solution was adjusted to a pH value around neutrality using aqueous 25 % sodium hydroxide solution, and agitated at ambient temperature for 4 hours. Using conc. hydrochloric acid, the reaction solution was adjusted to an acidic pH value, in order to extract the organic matter in ethyl acetate. The organic layer was rinsed with aqueous saturated sodium chloride, dried

over anhydrous magnesium sulfate and subsequently concentrated, to obtain the residue. The residue was recrystallized in aqueous 28 % ammonia and ethanol, to obtain 5.51 g (16.2 mmol) 4-hydroxy-4-(3-indolylmethyl)-2-hydroxyiminoglutaric acid in crystals (a yield of 32 % vs. indole-3-pyruvic acid).

<Example 18>

After 1.0 g (4.92 mmol) of indole-3-pyruvic acid was added to and dissolved in 10 ml of aqueous saturated sodium carbonate solution in the same manner as in Example 16, the resulting solution was adjusted to pH 12.7 using aqueous 25 % sodium hydroxide solution. After 1.3 g (14.8 mmol) of pyruvic acid was added, the resulting solution was adjusted to pH 10.0 using aqueous 25 % sodium hydroxide solution, for reaction at ambient temperature for 6 hours. 1.37 g (19.7 mmol) of hydroxylamine hydrochloride salt was added to the reaction solution while the reaction solution was adjusted to a pH value around neutrality using aqueous 25 % sodium hydroxide solution, and stirred at ambient temperature for 13 hours. Using conc. hydrochloric acid, the reaction solution was adjusted to an acidic pH value, to extract the organic matter in ethyl acetate. The organic layer was rinsed with aqueous saturated sodium chloride, dried over anhydrous magnesium sulfate and subsequently concentrated, to obtain the residue. The residue was analyzed by HPLC, by which it was shown that 4-hydroxy-

4-(3-indolylmethyl)-2-hydroxyiminoglutaric acid was generated at a yield of about 14 %.

<Example 19>

Synthesis of 4-hydroxy-4-(3-indolylmethyl)-2-hydroxyiminoglutaric acid; NO. 4

73.8 g (352 mmol) of indole-3-pyruvic acid was added to and dissolved in 917 g of aqueous 1.6 wt % sodium hydroxide solution. The resulting solution was adjusted to 35°C, to which 310.2 g (1761 mmol) of aqueous 50 % pyruvic acid solution was added dropwise over 2 hours, while the reaction solution was retained at pH 11.1 using aqueous 30 % sodium hydroxide solution. After reaction for another 4.5-hours, a reaction solution containing 4-hydroxy-4-(3-indolylmethyl)-2-ketoglutaric acid was obtained. 367.2 g (2114 mmol) of aqueous 40 % hydroxylamine hydrochloride salt solution was added to the reaction solution while the reaction solution was kept at pH 7 using aqueous 30 % sodium hydroxide solution, and stirred at 5°C for 17.5 hours. Using conc. hydrochloric acid, the reaction solution was adjusted to pH 2, in order to extract the organic matter in ethyl acetate. The organic layer was rinsed with aqueous saturated sodium chloride and concentrated, to obtain the residue. The residue was recrystallized in 60 ml of aqueous 28 % ammonia and 1350 ml of 2-propanol, to obtain 43.4 g (142 mmol) of the diammonium salt of 4-hydroxy-4-(3-indolylmethyl)-2-

hydroxyiminoglutaric acid in crystals (a yield of 40 % vs. indole-3-pyruvic acid).

<Example 20>

Production of (R)-(+)-1-phenylethylamine salt of (4S)-4-hydroxy-4-(3-indolylmethyl)-2-hydroxyiminoglutaric acid

After 44.7 g (0.131 mol) of the ammonium salt of 4-hydroxy-4-(3-indolylmethyl)-2-hydroxyiminoglutaric acid was dissolved in 500 ml of water at 25°C, the resulting aqueous solution was adjusted to pH 2 using 25.5 g of 36 % hydrochloric acid. The acidic solution was extracted in 1300 ml of ethyl acetate, and the resulting ethyl acetate solution was rinsed with 200 ml of aqueous saturated sodium chloride solution. 500 ml of an aqueous sodium carbonate solution (13.9 g (0.131 mole) of sodium carbonate) was added to the resulting ethyl acetate solution for agitation, to separate the aqueous alkali solution from ethyl acetate. 23.1 g of 36 % hydrochloric acid was added to the resulting aqueous alkali solution, to adjust the solution to pH 2. 6.99 g (57.6 mmol) of (R)-(+)-1-phenylethylamine was added dropwise to the resulting aqueous acidic solution and the solution stirred at 25°C for one hour. The resulting crystals were filtered and dried under reduced pressure, to obtain 21.8 g (47.8 mmol) of the (R)-(+)-1-phenylethylamine salt of (4S)-4-hydroxy-4-(3-indolylmethyl)-2-hydroxyiminoglutaric acid. (Yield, 72.7 %;

optical purity, 87.4 %.) .

$^1\text{H-NMR}$ (400MHz, DMSO-d_6) σ : 1.48 (d, 3H, $J = 6.8$ Hz), 2.63(d, 1H, $J = 14.0$ Hz), 2.70(d, 1H, $J = 14.0$ Hz), 2.90 (d, 1H, $J = 14.1$ Hz), 3.06 (d, 1H, $J = 14.1$ Hz), 4.40 (q, 1H, $J = 6.8$ Hz), 6.91-7.54 (m, 10H).

<Example 21>

Production of (S)-(-)-1-phenylethylamine salt of (4R)-4-hydroxy-4-(3-indolylmethyl)-2-hydroxyiminoglutaric acid

(S)-(-)-1-phenylethylamine of 7.12 g (58.7 mmol) was added dropwise to the solution remaining in Example 20 after the crystals of that example were filtered off, and the mixture stirred at 25°C for one hour. The resulting crystals were filtered off and dried under reduced pressure, to obtain 23.8 g (53.3 mol) of the (S)-(-)-1-phenylethylamine salt of (4R)-4-hydroxy-4-(3-indolylmethyl)-2-hydroxyiminoglutaric acid. (Yield, 81.1 %; optical purity, 92.1 %.)

<Example 22>

Production of the ammonium salt of (4S)-4-hydroxy-4-(3-indolylmethyl)-2-hydroxyiminoglutaric acid

200 ml of water and 18.5 g of aqueous 28 % ammonia were added to 21.8 g (51.0 mmol) of the (R)-(+)-1-phenylethylamine salt of (4S)-4-hydroxy-4-(3-indolylmethyl)-2-hydroxyiminoglutaric acid at 25°C so as to dissolve the salt, followed by further addition of 200 ml of toluene and stirring.

The aqueous layer obtained by the partition of the resulting layers was heated to 60°C. To the resulting aqueous solution was added dropwise 900 ml of 2-propanol over 2 hours. After the aqueous 2-propanol solution was cooled to 10°C over 5 hours, the solution was stirred at 10°C for 10 hours. The resulting crystals were filtered and dried under reduced pressure, to obtain 14.75 g of the ammonium salt of (4S)-4-hydroxy-4-(3-indolylmethyl)-2-hydroxyiminoglutaric acid. (Yield, 85.1 %; optical purity, 99.0 %.)

Melting point; 205°C (decomposed)

Specific rotation $[\alpha]^{20}_D + 13.4$ (c = 1.00, H₂O)

<Example 23>

Production of the ammonium salt of (4R)-4-hydroxy-4-(3-indolylmethyl)-2-hydroxyiminoglutaric acid

In the same manner as in the Example described above, 16.2 g of the ammonium salt of (4R)-4-hydroxy-4-(3-indolylmethyl)-2-hydroxyiminoglutaric acid was recovered from 23.8 g (53.3 mmol) of the (S)-(-)-1-phenylethylamine salt of (4R)-4-hydroxy-4-(3-indolylmethyl)-2-hydroxyiminoglutaric acid. (Yield, 89.3 %; optical purity, 99.9 %.)

Specific rotation $[\alpha]^{20}_D - 13.6$ (c = 1.00, H₂O)

<Example 24>

Production of (2S, 4S) monatin

4.5 g (13.1 mmol) of the ammonium salt of (4S)-4-hydroxy-4-(3-indolylmethyl)-2-hydroxyiminoglutamic acid was dissolved in 100 ml of aqueous 28 % ammonia, followed by addition of 3.4 g of 5 % rhodium carbon (50 % hydrous product), for reaction at ambient temperature and a hydrogen pressure of 10 atmospheres (1 MPa). After 24 hours, the catalyst was filtered off. The filtrate was concentrated. 40 ml of aqueous 90 % ethanol was added to the concentrate, followed by stirring at 25°C for 1.5 hours. The deposited crude crystals were filtered. 40 ml of aqueous 90 % ethanol was added to the crude crystals, followed by stirring at 25°C for 1.5 hours. The deposited purified crystals were filtered off and dried under reduced pressure, to obtain 0.57 g (1.84 mmol) of the ammonium salt of (2S, 4S) monatin. (Yield, 14.1 %; optical purity, 99.5 %).

$^1\text{H NMR}$ (400MHz, D_2O) δ : 2.06 (dd, $J=11.8, 15.3\text{Hz}$, 1H) , 2.67 (dd, $J=2.0, 15.2\text{ Hz}$, 1H) , 3.08 (d, $J=14.4\text{ Hz}$, 1H) , 3.28 (d, $J=14.4\text{ Hz}$, 1H) , 3.63 (dd, $J=2.2, 12.2\text{ Hz}$, 1H) , 7.12 – 7.16 (m, 1H) , 7.20 – 7.24 (m, 2H) , 7.48 – 7.49 (m, 1H) , 7.71 – 7.73 (m, 1H).

Theoretical ESI-MS value $\text{C}_{14}\text{H}_{16}\text{N}_2\text{O}_5 = 292.29$

Analytical value = 291.28 (MH^-)

<Example 25>

Production of ammonium salt of (2S, 4S) monatin

14.0 g (41.1 mmol) of the ammonium salt of (4S)-4-

hydroxy-4-(3-indolylmethyl)-2-hydroxyiminoglutaric acid was dissolved in 120 ml of aqueous 28 % ammonia, followed by addition of 7.38 g of 5 % rhodium carbon (50 % hydrous product), for reaction at 25°C and a hydrogen pressure of 1 MPa. After 24 hours, the catalyst was filtered off. The filtrate was concentrated. 110 ml of aqueous 88 % ethanol was added to 17.68 g of the concentrate, for stirring at 25°C for 19 hours. The resulting crude crystals were filtered off and dissolved in 15 ml of water, followed by addition of 100 ml of ethanol. After stirring at 25°C for 1.5 hours, the deposited purified crystals were filtered and dried under reduced pressure, to obtain 4.94 g (16.0 mmol) of the ammonium salt of (2S, 4S) monatin. (Yield, 39.2 %; optical purity, 99.9 %.)

<Example 26>

Production of free form of (2S, 4S) monatin

2.22 g (7.18 mmol) of the ammonium salt of (2S, 4S) monatin obtained in the above Example was dissolved in a mixed solvent of 4.5 ml of water and 4.2 ml (71.8 mmol) of acetic acid, followed by dropwise addition of 50 ml of ethanol to the resulting solution at 25°C over about 3 hours. After another 0.5-hr stirring, the resulting crystal was filtered and dried under reduced pressure, to obtain 1.93 g (6.62 mmol) of (2S, 4S) monatin. (Yield, 92.2 %; ammonium content, 0.19 wt %.)

<Comparative Example 1>

Using cinchonidine in place of the optically active amine used in Example 20, the same procedures were carried out. The optical purity of the resulting crystals was 0 %.

<Comparative Example 2>

L-Lysine was used in place of the optically active amine used in Example 20. However, no crystals were obtained.

<Comparative Example 3>



L-Arginine was used in place of the optically active amine used in Example 20. However, no crystals were obtained.




Industrial Applicability

In accordance with the invention, glutamic acid compounds typically including monatin useful as sweetener or an intermediate for producing pharmaceutical products can be efficiently produced industrially. In accordance with the invention, further, optically active monatin can be efficiently produced industrially.

CLAIMS

1. A process of producing a glutamic acid compound represented by the following formula (7) or a salt thereof, including a step of treating a pyruvic acid compound represented by the following formula (1) and oxalacetic acid represented by the following formula (2) by a cross aldol reaction and decarboxylation reaction or of treating the pyruvic acid compound (except for pyruvic acid) and a pyruvic acid represented by the following formula (2') by a cross aldol reaction, to obtain a ketoglutaric acid compound represented by the following formula (4) or a salt thereof, and a step of converting the carbonyl group of the ketoglutaric acid compound or a salt thereof to an amino group, in which the pyruvic acid compound, oxalacetic acid and pyruvic acid may individually be in salt forms thereof:






in the above formulas, R^1 represents a group selected from alkyl groups, aryl groups, aralkyl groups and heterocyclic ring-containing hydrocarbon groups; and R^1 may have at least one substituent selected from halogen atoms, hydroxyl group, alkyl groups with one to 3 carbon atoms, alkoxy groups with one to 3 carbon atoms and amino group.

2. A process according to claim 1, wherein the step of converting the carbonyl group of the ketoglutaric acid compound represented by the formula (4) or a salt thereof to an amino group includes a step of reacting an amine compound represented by the following formula (5) or a salt thereof with the

ketoglutaric acid or a salt thereof, to obtain a glutaric acid compound represented by the following formula (6) or a salt thereof, and a step of treating the resulting glutaric acid compound or a salt thereof by a reducing reaction:







in the above formulas, R^1 represents a group selected from alkyl groups, aryl groups, aralkyl groups and heterocyclic ring-containing hydrocarbon groups; R^2 represents a hydrogen atom or a group selected from alkyl groups, aryl groups and aralkyl groups; and R^1 may have at least one substituent selected from halogen atoms, hydroxyl group, alkyl groups with one to 3 carbon atoms, alkoxy groups with one to 3 carbon atoms and amino group.

3. A process according to claim 1, wherein the step of converting the carbonyl group of the ketoglutaric acid compound represented by the formula (4) or a salt thereof to an amino group includes a step of treating the ketoglutaric acid or a salt thereof by a reductive amination reaction.

4. A process according to claim 1, wherein the cross aldol reaction is done within a range of pH 10 to 14.

5. A process of producing a ketoglutaric acid compound represented by the following formula (4) or a salt thereof,


including a step of treating a pyruvic acid compound represented by the following formula (1) and oxalacetic acid represented by the following formula (2) by a cross aldol reaction and decarboxylation reaction, or of treating the pyruvic acid compound (except for pyruvic acid) and a pyruvic acid represented by the following formula (2') by a cross aldol reaction, in which the pyruvic acid compound, oxalacetic acid and pyruvic acid may individually be in salt forms thereof:





in the above formulas, R^1 represents a group selected from alkyl groups, aryl groups, aralkyl groups and heterocyclic ring-containing hydrocarbon groups; and R^1 may have at least one substituent selected from halogen atoms, hydroxyl group, alkyl groups with one to 3 carbon atoms, alkoxy groups with one to 3 carbon atoms and amino group.

6. A process according to claim 5, wherein the cross aldol reaction is done within a range of pH 10 to 14.


7. A process of producing a glutamic acid compound represented by the following formula (7) or a salt thereof, including a step of reacting a ketoglutaric acid compound represented by the following formula (4) or a salt thereof with an amine compound represented by the following formula (5) or a salt thereof, to obtain a glutaric acid compound represented by the following formula (6) or a salt thereof, and a step of treating the resulting glutaric acid compound or a salt thereof by a reducing reaction:










in the above formulas, R^1 represents a group selected from alkyl groups, aryl groups, aralkyl groups and heterocyclic ring-containing hydrocarbon groups; R^2 represents a hydrogen atom or a group selected from alkyl groups, aryl groups and aralkyl groups; and R^1 may have at least one substituent selected from halogen atoms, hydroxyl group, alkyl groups with one to 3 carbon atoms, alkoxy groups with one to 3 carbon atoms and amino group.

8. A process of producing a glutamic acid compound represented by the following formula (7) or a salt thereof, including a step of treating a ketoglutaric acid compound represented by the following formula (4) or a salt thereof by a reductive amination reaction:




in the above formulas, R^1 represents a group selected from alkyl groups, aryl groups, aralkyl groups and heterocyclic ring-containing hydrocarbon groups; and R^1 may have at least one substituent selected from halogen atoms, hydroxyl group, alkyl groups with one to 3 carbon atoms, alkoxy groups with one to 3 carbon atoms and amino group.

9. A process of producing monatin represented by the following formula (7') or a salt thereof, including a step of treating indole-3-pyruvic acid represented by the following formula (1') and oxalacetic acid represented by the following formula (2) by an aldol reaction and a decarboxylation reaction, or of treating indole-3-pyruvic acid represented by the following formula (1') and pyruvic acid represented by the following formula (2') by a cross aldol reaction, to obtain 4-hydroxy-4-(3-indolylmethyl)-2-ketoglutaric acid represented by the following formula (4') or a salt thereof, and a step of converting the carbonyl group of the ketoglutaric acid or a salt thereof to an amino group, in which indole-3-pyruvic acid, oxalacetic acid and pyruvic acid may individually be in salt forms thereof:



10. A process according to claim 9, wherein the step of converting the carbonyl group of 4-hydroxy-4-(3-

indolylmethyl)-2-ketoglutaric acid represented by the formula (4') or a salt thereof to an amino group includes a step of reacting an amine compound represented by the following formula (5) or a salt thereof with the ketoglutaric acid or a salt thereof to obtain a glutaric acid compound represented by the following formula (6') or a salt thereof, and a step of treating the glutaric acid compound or a salt thereof by a reducing reaction:






in the formula, R^2 represents a hydrogen atom, or a substituent selected from alkyl groups, aryl groups and aralkyl groups.


11. A process according to claim 9, wherein the step of converting the carbonyl group of 4-hydroxy-4-(3-indolylmethyl)-2-ketoglutaric acid represented by the formula (4') or a salt thereof to an amino group includes a step of treating the ketoglutaric acid compound or a salt thereof by a reductive amination reaction.

12. A process according to claims 9 to 11, wherein the cross

aldol reaction is done within a range of pH 10 to 14.


13. A process of producing 4-hydroxy-4-(3-indolylmethyl)-2-ketoglutaric acid represented by the following formula (4') or a salt thereof, including a step of treating indole-3-pyruvic acid represented by the following formula (1') and oxalacetic acid represented by the following formula (2) by a cross aldol reaction and a decarboxylation reaction, or of treating indole-3-pyruvic acid represented by the following formula (1') and pyruvic acid represented by the following formula (2') by an aldol reaction, in which indole-3-pyruvic acid, oxalacetic acid and pyruvic acid may individually be in salt forms thereof:







14. A process according to claim 13, wherein the cross aldol reaction is done within a range of pH 10 to 14.


15. A process of producing monatin represented by the following formula (7') or a salt thereof, including a step of reacting 4-hydroxy-4-(3-indolylmethyl)-2-ketoglutaric acid represented by the following formula (4') or a salt thereof with an amine compound represented by the following formula (5) or a salt thereof, to obtain a glutaric acid compound represented by the following formula (6') or a salt thereof, and a step of subsequently treating the glutaric acid compound or a salt thereof by a reducing reaction:






in the formula, R^2 represents a hydrogen atom or a substituent selected from alkyl groups, aryl groups and aralkyl groups.

16. A process of producing monatin represented by the following formula (7') or a salt thereof, including a step of treating 4-hydroxy-4-(3-indolylmethyl)-2-ketoglutaric acid represented by the following formula (4') or a salt thereof by a reductive amination reaction:





17. A compound represented by the following formulas (4') or (6') or a salt thereof, in which in the formulas, R^2 represents a hydrogen atom, an alkyl group, an aryl group or an aralkyl group:

