



(11)

EP 3 312 158 A1

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:
25.04.2018 Bulletin 2018/17

(51) Int Cl.:
C07D 207/46 (2006.01) **C07C 327/20** (2006.01)
C07C 327/32 (2006.01) **C07H 3/06** (2006.01)
C08B 37/00 (2006.01) **C07K 14/31** (2006.01)
A61K 39/08 (2006.01) **A61K 39/085** (2006.01)

(21) Application number: 17198371.1

(22) Date of filing: 21.07.2009

(84) Designated Contracting States:
AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL
PT RO SE SI SK SM TR
Designated Extension States:
AL BA RS

(30) Priority: 21.07.2008 US 135493 P
20.02.2009 US 208155 P

(62) Document number(s) of the earlier application(s) in accordance with Art. 76 EPC:
09800657.0 / 2 315 747

(71) Applicant: **The Brigham and Women's Hospital, Inc.**
Boston, MA 02115 (US)

(72) Inventors:

- **PIER, Gerald, B.**
Brookline, MA Massachusetts 02446 (US)
- **NIFANTIEV, Nikolay**
117593 Moscow (RU)
- **TSVETKOV, Yury**
119421 Moscow (RU)
- **GENING, Marina**
125047 Moscow (RU)

(74) Representative: **Potter Clarkson LLP**
The Belgrave Centre
Talbot Street
Nottingham NG1 5GG (GB)

Remarks:

This application was filed on 25-10-2017 as a divisional application to the application mentioned under INID code 62.

(54) **METHODS AND COMPOSITIONS RELATING TO SYNTHETIC BETA-1,6 GLUCOSAMINE OLIGOSACCHARIDES**

(57) The invention relates to the compositions of synthetic oligo- β -(1 \rightarrow 6)-2-amino-2-deoxy-D-glucopyranosides conjugated to carriers, and methods for making and use same.
ed.

Description**RELATED APPLICATIONS**

5 **[0001]** This application claims priority to U.S. provisional applications 61/135493 and 61/208155, filed on July 21, 2008 and February 20, 2009, respectively, the entire contents of both of which are incorporated by reference herein.

GOVERNMENT SUPPORT

10 **[0002]** The present invention was supported in part by a grant from the United States National Institutes of Health RO1AI046706. The U.S. Government has rights in the invention.

FIELD OF THE INVENTION

15 **[0003]** The invention relates to compositions and methods relating to synthetic oligo- β -(1 \rightarrow 6)-2-amino-2-deoxy-D-glucopyranosides which are referred to herein interchangeably as oligo- β -(1 \rightarrow 6)-D-glucosamines or oligoglucosamines.

BACKGROUND OF THE INVENTION

20 **[0004]** *Staphylococci* are gram-positive bacteria which normally inhabit and colonize the skin and mucus membranes of humans. If the skin or mucus membrane becomes damaged during surgery or other trauma, the *Staphylococci* may gain access to internal tissues causing infection to develop. If the *Staphylococci* proliferate locally or enter the lymphatic or blood system, serious infectious complications such as those associated with *Staphylococcal* bacteremia may result. These complications include septic shock, endocarditis, arthritis, osteomyelitis, pneumonia, and abscesses in various organs.

25 **[0005]** *Staphylococci* include both coagulase-positive organisms that produce a free coagulase and coagulase-negative organisms that do not produce this free coagulase. *Staphylococcus aureus* is the most common coagulase-positive form of *Staphylococci*. *S. aureus* generally causes infection at a local site, either extravascular or intravascular, which ultimately may result in bacteremia. *S. aureus* is also a leading cause of acute osteomyelitis, and causes *Staphylococcal* pneumonia infections. Additionally, *S. aureus* is responsible for approximately 1-9% of the cases of bacterial meningitis and 10-15% of brain abscesses.

30 **[0006]** There are at least thirty-one known species of coagulase-negative *Staphylococci*, including *S. epidermidis*, *S. saprophyticus*, *S. hominis*, *S. warneri*, *S. haemolyticus*, *S. saprophiticus*, *S. cohnii*, *S. xylosus*, *S. simulans*, and *S. capitis*. *S. epidermidis* is the most frequent infection-causing agent associated with intravenous access devices, and the most frequent isolate in primary nosocomial bateremias. *S. epidermidis* is also associated with prosthetic valve endocarditis.

35 **[0007]** *Staphylococcus* is also a common source of bacterial infection in animals. For instance, *Staphylococcal* mastitis is a common problem in ruminants such as cattle, sheep, and goats. The disease is generally treated with antibiotics to reduce the infection but the treatment is a costly procedure and still results in a loss of milk production. The most effective vaccines identified to date are live, intact *S. aureus* vaccines administered subcutaneously. The administration of live vaccines, however, is associated with the risk of infection. For that reason, many researchers have attempted to produce killed *S. aureus* vaccines and/or to isolate capsular polysaccharides or cell wall components which will induce immunity to *S. aureus*.

40 **[0008]** Carrier compounds are sometimes used in vaccines in order to enhance the immune response to the antigen. For example, the carrier in some vaccines is useful for stimulating T cell help in response to the antigenic moiety. Antigenic moieties that are naturally occurring or fragments of naturally occurring substances are sometimes less amenable and less facilely manipulated and/or conjugated to other moieties such as for example carrier compounds, and this can reduce the therapeutic impact of such vaccines.

50 SUMMARY OF THE INVENTION

55 **[0009]** The invention provides novel methods and compounds for generating antigenic compositions such as but not limited to vaccines. In particular, the invention provides novel methods for modifying oligosaccharide antigens. These methods involve the controlled synthesis of oligosaccharide (a) comprised of a predetermined order of monosaccharide monomers and (b) conjugated to a carrier compound. The resulting conjugates are better able to stimulate (whether by induction or enhancement) an immune response to the microbial polysaccharide antigen of interest. As described herein, such immune responses are useful in the treatment and/or prevention of various infections including but not limited to *Staphylococcal* infections.


[0010] Various aspects of the invention relate to particular linkers (or linking agents or spacers or linking reagents, as the terms are used interchangeably herein), and their use in synthesizing conjugates. Of particular interest are conjugates of oligosaccharides and various carrier compounds. These oligosaccharides include oligo- β -(1 \rightarrow 6)-D-glucosamine (linked glucosamine, as referred to herein) which is a comprised of glucosamine monomers attached to each other by a β -(1 \rightarrow 6) linkage. One or more of the various monomers that make up the linked glucosamine may be N-acetylated. Thus, the monomers may be D-glucosamine or N-acetyl-D-glucosamine monomers, and the linked glucosamine may comprise a defined order of one or both types of these monomers (or monosaccharides, as the terms are used interchangeably herein). The oligoglucosamines of the invention also comprise spacers having thiol containing groups on their "reducing" ends. The spacer is used to link the oligoglucosamines to carriers. The carriers may be comprised of amino acids (such as peptides or proteins) although they are not so limited.

[0011] Thus, in one aspect, the invention provides a compound of Formula I



Formula I


wherein X is any atom or group, Y is a sulfur blocking group, and n is greater than 1. In one embodiment, Y is an acyl group. In another embodiment, Y is an acetyl group, and the compound has the structure of Formula II



Formula II

wherein Ac is an acetyl group.


[0012] In still another embodiment, the compound is an activated ester of Formula I. In another embodiment, the compound is a cyano, azido or haloid derivative an activated ester of Formula I. In another embodiment, the compound has the structure of Formula III



Formula III

40

wherein Ac is an acetyl group, and the compound is referred to as N-hydroxysuccinimidyl 4-acetylsulfanyl butyrate. In another embodiment, the compound has the structure of Formula IV




Formula IV

50

wherein Ac is an acetyl group and the compound is referred to as N-nitrophenyl 4-acetylsulfanyl butyrate.


[0013] In another aspect, the invention provides a method for synthesizing N-hydroxysuccinimidyl 4-acetylsulfanyl butyrate (Formula III) comprising reacting 4-acetylsulfanylbutyric acid (Formula V)



Formula V

with N-hydroxysuccinimidyl trifluoroacetate (CF_3COOSu) to yield N-hydroxysuccinimidyl 4-acetylsulfanyl butyrate (Formula III).

10 [0014] In another aspect, the invention provides a method for synthesizing N-nitrophenyl 4-acetylsulfanyl butyrate (IV) comprising reacting 4-acetylsulfanylbutyric acid (Formula V)




Formula V

20 with N-nitrophenyl trifluoroacetate (CF_3COOpNp) to yield N-nitrophenyl 4-acetylsulfanyl butyrate (Formula IV).

[0015] In another aspect, the invention provides a method for synthesizing an oligosaccharide conjugated to a carrier comprising reacting an oligosaccharide (a) first with a compound having a structure of Formula Va or Vb



Formula Va




Formula Vb

30 wherein m is a number selected from 1 to 10, p is a number selected from 1 to 20, and R is H or a alkyl group, (b) second with a compound of Formula I, II, III or IV, and (c) third with a carrier, wherein the oligosaccharide is β -1-6 linked glucosamine.

[0016] In yet another aspect, the invention provides a composition comprising an oligosaccharide bearing an O-linked linker, wherein the linker comprises

35




Formula VI.

[0017] In one embodiment, the oligosaccharide is β -1-6 linked glucosamine. In another embodiment, the oligosaccharide is 2-20 monomers in length. In another embodiment, the oligonucleotide is 5-11 monomers in length. In still another embodiment, the oligosaccharide is 7, 9 or 11 monomers in length.


45 [0018] In one embodiment, the oligosaccharide is 0% acetylated or 100% acetylated. In another embodiment, the oligosaccharide is 0-40% acetylated.

[0019] In still another aspect, the invention provides a composition comprising an oligosaccharide-carrier conjugate comprising an oligosaccharide conjugated to a carrier through a linker that is

50




Formula VIIa




Formula VIIIb

wherein the linker is O-linked to the oligosaccharide and N-linked to the carrier.

10 [0020] In still another aspect, the invention provides a method for synthesizing an oligosaccharide-carrier conjugate comprising reacting an oligosaccharide conjugated to a linker of Formula IXa or IXb




Formula IXa



Formula IXb

with a carrier having modified amino group following reaction with a compound of Formula



Formula X

to yield an oligosaccharide conjugated to a carrier compound through a linker having a structure of Formula VIIa or VIIb.

[0021] In one embodiment, the carrier is a peptide. In another embodiment, the carrier is a protein. An example of a carrier is tetanus toxoid.

[0022] In one embodiment, the oligosaccharide is β -1-6 linked glucosamine.

[0023] In one embodiment, the composition has a oligosaccharide to μ -C-linked glucosamine ratio of 1:1, 2:1, 3:1, 4:1, 5:1, 6:1, 7:1, 8:1, 9:1, 10:1, 20:1, 30:1, 40:1, 50:1, 60:1, 70:1, 80:1, 90:1 or 100:1.

[0024] In some embodiments, the oligosaccharide is 2-20 monomers in length, 5-11 monomers in length, or 5, 7, 9 or 11 monomers in length.

[0025] In one embodiment, the oligosaccharide is 100% acetylated. In another embodiment, the oligosaccharide is 0% acetylated. In still another embodiment, the oligosaccharide is 0-40% acetylated.

[0026] In some embodiments, the composition further comprises a pharmaceutically acceptable carrier, an adjuvant, and/or an anti-bacterial agent.

50 [0027] In another aspect, the invention provides a method for stimulating an immune response in a subject comprising administering to a subject in need thereof an oligosaccharide-carrier conjugate as described above or an oligosaccharide-carrier conjugate synthesized by a method described above in an amount effective to stimulate an immune response in the subject.

[0028] In one embodiment, the subject is a human. In another embodiment, the subject is non-human.

55 [0029] In another embodiment, the method further comprises isolating antibodies or antibody-forming cells from the subject.

[0030] In another embodiment, the method further comprises administering an adjuvant to the subject. In another embodiment, the method further comprises administering an anti-bacterial agent to the subject.

[0031] In still another aspect, the invention provides a method for treating or preventing a infection in a subject comprising administering to a subject having or at risk of developing an infection an effective amount for inducing an immune response of an oligosaccharide-carrier conjugate as described above or an oligosaccharide-carrier conjugate synthesized by a method as described above, wherein the infection is caused by a bacterial species that makes or is capable of making PNAG.

[0032] In one embodiment, the infection is a *Staphylococcus* infection. In a related embodiment, the *Staphylococcus* infection is *Staphylococcus aureus* infection. In another related embodiment, the *Staphylococcus* infection is *Staphylococcus epidermidis*. In one embodiment, the subject is at risk of exposure to *Staphylococcus*. In another embodiment, the subject has been exposed to *Staphylococcus*.

[0033] In other embodiments, the infection is an *E. coli* infection, a *Y. pestis* infection, a *Y. entercolitica* infection, a *Y. pseudotuberculosis* infection, an *Aggregatibacter actinomycetemcomitans* infection, an *Actinobacillus pleuropneumoniae* infection, a *Bordetella pertussis* infection, a *B. parapertussis* infection, a *B. bronchiseptica* infection, an *Acinetobacter* infection, a *Burkholderia* infection such as *Burkholderia cenocepacia*, a *Stenatrophomonas maltophilia* infection, a *Shigella* infection, or a *Klebsiella* infection such as *Klebsiella pneumoniae*.

[0034] In some embodiments, the oligosaccharide-carrier conjugate is administered with an adjuvant and/or an anti-bacterial agent.

[0035] In still another aspect, the invention provides a method for producing antibodies comprising administering to a subject an effective amount, for producing antibodies specific for native PNAG, of an oligosaccharide-carrier conjugate as described above or an oligosaccharide-carrier conjugate synthesized by a method as described above, and isolating antibodies from the subject.

[0036] In another aspect, the invention provides a method for producing monoclonal antibodies comprising administering to a subject an effective amount, for producing antibodies specific for native PNAG, of an oligosaccharide-carrier conjugate as described above or an oligosaccharide-carrier conjugate synthesized by a method as described above, harvesting antibody-producing cells from the subject, fusing the antibody-producing cells from the subject to myeloma cells, and harvesting antibody produced from a fusion subclone.

[0037] In one embodiment, the antibodies are polyclonal antibodies. In one embodiment, the subject is a rabbit. In another embodiment, the subject is a human.

[0038] It should be appreciated that all combinations of the foregoing concepts and additional concepts discussed in greater detail below (provided such concepts are not mutually inconsistent) are contemplated as being part of the inventive subject matter disclosed herein. In particular, all combinations of claimed subject matter appearing at the end of this disclosure are contemplated as being part of the inventive subject matter disclosed herein. It should also be appreciated that terminology explicitly employed herein that also may appear in any disclosure incorporated by reference should be accorded a meaning most consistent with the particular concepts disclosed herein.

35 BRIEF DESCRIPTION OF THE FIGURES

[0039]

40 FIGs. 1A and B are graphs showing binding of antisera raised to non-acetylated nonaglucosamine (9GlcNH₂) to PNAG (A) or dPNAG (B) from *S. aureus*.

FIGs. 2A and B are graphs showing binding of antisera raise to fully acetylated nonaglucosamine (9GlcNAc) to PNAG or dPNAG from *S. aureus*.

FIGs. 3A and B are graphs showing binding of antisera raised to conjugated 9GlcNAc or 9GlcNH₂ to 1 1GlcNAc or 1 1GlcNH₂.

45 FIG. 4 is a graph showing killing of *S. aureus* strain MN8 bacteria using an antisera raised in two rabbits, one received the 9GlcNH₂-TT conjugate and the second received the 9GlcNAc-TT conjugate with the antiserum taken 2 weeks after the last injection of vaccine.

50 FIG. 5 is a graph showing killing of *S. aureus* strain MN8 bacteria using an antisera raised in two rabbits, one of which received the 9GlcNH₂-TT conjugate and the second received the 9GlcNAc-TT conjugate with the antiserum taken 4 weeks after the last injection of vaccine and also shows, for comparison, killing of the same bacteria by an antiserum raised to a conjugate vaccine consisting of the dPNAG molecule of ~ 100 kDa conjugated to tetanus toxoid (TT) and further labeled (051).

FIGs. 6-8 are graphs comparing the killing of *S. aureus* strains (FIGs. 6 and 8, LAC (NT, USA300); FIG. 7, SF8300 (NT, USA300)) by rabbit antisera (bleed 1) to fully acetylated or non-acetylated 9-mer oligoglucosamine (9GlcNH₂) conjugated to TT (9GlcNH₂-TT). For comparison, killing of the same bacteria by an antiserum raised to a conjugate vaccine consisting of the dPNAG molecule of ~ 100 kDa conjugated to tetanus toxoid (TT) and further labeled (051).

55 FIGs. 9-16 are graphs comparing the killing of *S. aureus* strains (FIG. 9, MN8 (capsular polysaccharide (CP) 8); FIG. 10, LAC (non-typable (NT), USA300)); FIG. 11, SF8300 (NT, USA300)); FIG. 12, Newman (CP5); FIG. 13,

PS80; FIG. 14, Reynolds (CP5); FIG. 15, Reynolds (non-typable); FIG. 16, Reynolds (CP8)) by a rabbit antiserum (labeled "bleed 2") to fully acetylated or non-acetylated 9-mer oligoglucosamine conjugated to TT. For comparison, killing of the same bacteria by an antiserum raised to a conjugate vaccine consisting of the dPNAG molecule of ~ 100 kDa conjugated to tetanus toxoid (TT) and further labeled (051).

5 FIG. 16A is a graph showing the killing of two PNAG-positive (*E. coli* J and *E. coli* P) but not PNAG-negative (*E. coli* H) *E. coli* strains by rabbit antisera to 9GlcNH₂-TT obtained 6 weeks after the last immunization.

FIGs. 17, 18 and 19 are graphs showing the results of an in vivo study relating to prevention of *S. aureus* skin abscess infection after immunization with an antiserum raised to the 9-mer nonacetylated oligoglucosamine conjugated to TT (9GlcNH₂-TT, bleed 2) and administered to mice 24 hr prior to infection with 2 x 10⁴ (FIG. 18), 2 x 10⁵ (FIG. 19) or 2 x 10⁶ (FIG. 20) CFU of *S. aureus* strain LAC.

10 FIG. 20 is a summary of the results in FIGs. 17-19.

FIGs. 21, 22 and 23 are graphs showing the results of in vivo studies relating to prevention of *S. aureus* abscess infection after immunization with an antiserum raised to the 9-mer nonacetylated oligoglucosamine and administered to mice prior to infection with 1 x 10⁶ CFU *S. aureus* MN8 (FIG. 21), 4 x 10⁶ CFU *S. aureus* Newman (FIG. 22), and 15 4 x 10⁶ CFU *S. aureus* NewmanΔica and 1.5 x 10⁶ CFU *S. aureus* MN8Δica (FIG. 23).

DETAILED DESCRIPTION OF THE INVENTION

[0040] The invention relates broadly to the synthesis and use of oligosaccharide conjugates. The invention provides 20 de novo synthesis methods and the compositions used therein to generate novel compositions. These synthetic routes facilitate modification of the oligosaccharide or polysaccharide that would not otherwise be possible using naturally occurring polysaccharide antigens.

[0041] The invention provides inter alia methods for the preparation of oligo- β -(1 \rightarrow 6)-D-glucosamine oligosaccharides 25 or polysaccharides having a defined order of monomers, methods for the conjugation of such oligosaccharides or polysaccharides to linkers for subsequent conjugation to carrier compounds, methods for the preparation of the oligosaccharide-carrier conjugates or polysaccharide-carrier conjugates, and compositions of these various compounds. The invention further relates to various novel linkers that are unexpectedly more useful than previously known linkers in these preparation methods. The resultant oligosaccharide-carrier conjugates are useful for stimulating an immune response 30 in vivo in human and non-human subjects, including generation of antibodies to the oligosaccharide themselves and the corresponding naturally occurring PNAG and dPNAG antigens.

[0042] PNAG and dPNAG antigens are described in greater detail in published PCT application WO 2004/043405. Briefly, PNAG refers to poly N-acetyl glucosamine which is a surface polysaccharide made by various bacterial species 35 including but not limited to *Staphylococci*, such as *S. aureus* and *S. epidermidis*. PNAG exists naturally in both highly and poorly acetylated forms. A "highly acetylated" form of PNAG is a PNAG having greater than 50% acetate substitutions. Poorly acetylated forms of PNAG (referred to herein as dPNAG) may have 0-40% acetylation. (See Formula VII where 40 R¹ represents the location of the acetyl group, if present.) Native PNAG is a mixture of PNAG forms with varying degrees of acetylation. Native PNAG may include dPNAG in a mixture with more highly acetylated forms of PNAG. PNAG or dPNAG may be comprised of hundreds or thousands or more glucosamine units (or monomers).

[0043] The oligosaccharides of the invention are intended to mimic regions of PNAG or dPNAG. Thus, when used in 45 vivo the oligosaccharide-carrier conjugates induce immune responses directed to regions of the oligosaccharide that are similar or identical to PNAG and/or dPNAG and therefore such immune responses are useful in targeting bacterial species that make or are capable of making PNAG and/or dPNAG.

[0044] In some aspects of the invention, the oligosaccharides are comprised of only D-glucosamine, or only N-acetyl-D-glucosamine units, or a predetermined ratio and order of both types of these monomers. The ratio and order is intended 50 to mimic in some embodiments the ratios and orders found in native PNAG. The oligosaccharides are manipulated according to the invention to comprise a spacer (or a linker, as the terms are used interchangeably herein) having a thiol group at its terminus (e.g., its reducing end).


[0045] The preparation of oligosaccharides comprising amine groups, such as linked glucosamine oligosaccharides 55 (or oligoglucosamines, as the terms are used interchangeably herein) suitable for conjugation to one or more carriers has proved challenging in the art. This is partly because the stereo-specific synthesis of linked glucosamine requires the use of participating but temporary acyl N-protecting groups (so-called "participating" groups) in the glycosyl donors in order to form the necessary β -glycoside bond between the monomers. N-phthaloyl, N-trichloroethoxycarbonyl and some other moieties are suitable as participating groups. Certain other participating groups however are less suitable including the N-acetyl participating groups that are present in some of the oligosaccharides contemplated by the invention. As an example, N-acetylated glycosyl donors are of low reactivity and give only moderate yields of glycosylation products. In addition, the presence of N-acetyl groups in glycosyl donor complicates the glycosylation reaction due to oxazoline intermediate formation, migration of N-acetyl groups, and other undesirable chemical reactions.

[0046] With regards to linked glucosamines, their structure and more particularly the number of amino groups they

contain necessitates the introduction of the linker before total liberation of amino groups. Removal of the above-mentioned temporary N-protecting group to prepare free oligosaccharide is carried out under basic conditions. The most effective reagent for removal of an N-phthaloyl participating group is hydrazine hydrate in boiling ethanol. This reagent also effectively removes O-acyl protecting groups including acetyl and benzoyl groups which may be contained in the oligosaccharide of interest.

[0047] Commercially available linkers that have been used for the attachment of amino-containing ligands to proteins are pentafluorophenyl S-acetylthioglycolate (chemical structure 8 shown in the Examples) and N-hydroxysuccinimidyl 3-acetylsulfanyl-propionate (chemical structure 12 shown in the Examples). These linking reagents can be used to introduce a thiol moiety into a oligosaccharide; however, as described in greater detail in the Examples, both were unstable under the conditions of removal of phthaloyl groups. Example 5 shows that a linker based on thioglycolic acid undergoes oxidative rearrangement, and Example 6 shows that a derivative of 3-mercaptopropionic acid gives a complex mixture of side products. Both transformations result in loss of the necessary thiol function.

[0048] The invention therefore is based in part on the discovery and use of a class of linkers that is effective and better than previously known linkers for conjugating oligosaccharides (including amine containing versions thereof) to a carrier such as a protein. This class of linkers is defined as derivatives of ω -acetylsulfanyl carbonic acid of Formula I (where $n > 1$).



Formula I

This class of linkers provides effective N-acylation during attachment to an oligosaccharide. The linker may be an activated ester of Formula I or its cyano, azido, or haloid derivative. It may be another derivative of Formula I provided such derivative is active as an acylating agent and thus suitable for attachment to oligosaccharides of the present invention. Y represents a temporary blocking group of sulfur atoms, which are known in the art and which include acyl and acetyl groups. Removal of the Y group liberates an SH group which is needed for attachment of the oligosaccharide to the carrier. X represents any leaving group that provides the necessary acylating ability to the compound of Formula I.


[0049] It is to be understood that any of the linker classes provided by the invention can be used to conjugate oligosaccharides to carrier compounds.

[0050] As an example, one linker class may comprise the structure of Formula II (where $n > 1$):



Formula II.


[0051] As another example, another linker class may comprise N-hydroxysuccinimidyl derivatives. An example of such a linker, as described in Examples 1 and 3, is N-hydroxysuccinimidyl 4-acetylsulfanyl butyrate (compound 2 on Scheme 1, Examples 1 and 3). This linker has the structure of Formula III as follows:



Formula III.

This linker is stable under conditions of total deprotection of carbohydrates such as oligosaccharides and polysaccharides.


[0052] Another example of an activated ester is N-nitrophenyl 4-acetylsulfanyl butyrate (compound 3, Examples 2 and 3). This compound has the structure of Formula IV as follows:



Formula IV

(N-nitrophenyl 4-acetylsulfanyl butyrate).


10 [0053] The invention provides a method for synthesizing N-hydroxysuccinimidyl 4-acetylsulfanyl butyrate (Formula III). This method involves reacting 4-acetylsulfanylbutyric acid (Formula V)



Formula V

20 with N-hydroxysuccinimidyl trifluoroacetate (CF_3COOSu) to yield N-hydroxysuccinimidyl 4-acetylsulfanyl butyrate (Formula III).

[0054] The invention further provides a method for synthesizing N-nitrophenyl 4-acetylsulfanyl butyrate (Formula IV). This method involves reacting 4-acetylsulfanylbutyric acid (Formula V)




Formula V

30 with N-nitrophenyl trifluoroacetate (CF_3COOpNp) to yield N-nitrophenyl 4-acetylsulfanyl butyrate (Formula IV).

[0055] These synthesis methods are described in greater detail in the Examples.


35 [0056] The invention further provides compositions comprising an oligosaccharide comprising a linker conjugated to the O1-atom of the glucosamine unit at the "reducing end" of the oligosaccharide. This O1-conjugated linker is used to attach an SH-containing linker *via* the reaction with the compound of Formula I. The O1-conjugated linker have the structure of Formula Va (below) having an aminoalkyl group and where m may vary from 1 to 10 and R may be H or a simple alkyl group (e.g., a methyl or an ethyl group). Alternatively, an O1-conjugated linker may have the structure of Formula Vb (below) where p may vary from 1 to 20 and R is the same as in Formula Va.



Formula Va

Formula Vb


45 [0057] The oligosaccharides conjugates of the present invention therefore may be synthesized by coupling the compounds of Formula Va or Vb with the compound of Formula I. The transformation and subsequent removal of the temporary S-blocking group (Example 8) or reduction of the corresponding intermediate disulfide (Example 7) results in the generation of a final linking group used in the conjugation of the oligosaccharides with a carrier compound. Such 50 final linking groups are described in Examples 7 and 8 have the structure of Formula VI



Formula VI.

[0058] The oligosaccharide may comprise 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 30, 40, 50, 60, 70, 80, 90 or 100 monosaccharide monomers. In some important embodiments, the oligosaccharide comprises 5 or more monomers including 5, 7, 9 or 11 monomers. In some embodiments, the oligosaccharide comprises 2-20 monomers, or 3-20 monomers, or 4-20 monomers, or 5-20 monomers.

5 [0059] In some important embodiments, the monomer is glucosamine and the oligosaccharide is a linked glucosamine. The structure of a glucosamine monomer (as present in a linked glucosamine) is as follows:




15 Formula VII


where R¹ is H in the case of glucosamine units with a "free" amino group, or R¹ is an acetyl group (COCH₃) in the case of N-acetylated glucosamine units. These units are connected through β -(1 \rightarrow 6)-linkages. Any number of glucosamine units may be linked, including 2, 3, 4, 5, 6, 7, 8, 9, 10, 50, or more units up to 100 units (whether substituted with acetyl groups or un substituted).

20 [0060] The degree of N-acetylation in the compounds of Formula VII may vary. It may range from 0-50% N-acetylation (i.e., 0-50% of R¹ are acetyl groups), including 0-40% N-acetylation. In some embodiments, the β -(1 \rightarrow 6) linked glucosamine is less than 50%, less than 40%, less than 30%, less than 20%, less than 10%, or less than 5% N-acetylated. In some important embodiments, the level of N-acetylation and the position of acetyl groups within an oligosaccharide 25 are known by virtue of the synthesis method. That is, the oligosaccharide may be synthesized from the ordered arrangement of glucosamine units having or lacking N-acetyl groups. The Examples provide methods for producing oligosaccharides having defined and ordered acetyl substitutions. In addition, reference can be made to Gening et al. Carbohydrate Research 2007 342:567-575, Gening et al. Russian J Bioorganic Chem 2006 32(4):389-399, Yang and Du Carbohydrate Research 2003 338:495-502, Yang et al. Carbohydrate Research 2003 338:1313-1318, and Fridman et al. Organic Letters 2002 4(2):281-283.

30 [0061] The invention still further provides a composition comprising an oligosaccharide-carrier conjugate comprising an oligosaccharide conjugated to a carrier compound through a linker of Formula VIIIa or VIIIb.



40 Formula VIIIa




45 Formula VIIIb

50 wherein the linker is O-linked to the oligosaccharide and N-linked to the carrier compound.


[0062] The invention further provides a method for synthesizing an oligosaccharide-carrier conjugate by reacting oligosaccharide conjugates comprising SH-terminated linkers such as those having structures of Formula IXa or IXb

5



Formula IXa

10




15

Formula IXb

(in the case of compounds from Examples 7 and 8)

with a carrier compound in which terminal amino groups are modified by attachment of a compound of Formula X

20



25

Formula X

to yield an oligosaccharide conjugated to a carrier compound through a linker having a structure of Formula VIIa or VIIb.

[0063] In some embodiments, a linker of Formula VIIa or VIIb is linked through its terminal CH₂-group to the O1-atom of glucosamine unit at the "reducing end" of the oligosaccharide. Such a linkage is considered an O-linkage, and the oligosaccharide is referred to as being O-linked to the linker.

[0064] In some embodiments, a linker of Formula VIIa or VIIb is linked through the terminal CO group to an amino group in the carrier compound by an amide bond. Such a linkage is considered an N-linkage, and the carrier is referred to as being N-linked to the linker.

[0065] Preparation of conjugates of amino-group-containing carrier and oligoglucosamines having an SH-terminated linker, as described above and in Examples 8 and 9, may use the reagent SBAP. The invention however contemplates the use of other reagents suitable for linking amino-group-containing carriers and SH-terminated oligosaccharides (particularly as described by G. Hermanson "Bioconjugate Techniques", 2nd Edition, Academic Press, 2008).

Carrier Compound

40

[0066] A "carrier compound" (or carrier, as the terms are used interchangeably herein) as used herein is a compound that is conjugated to the oligosaccharide through the use of a linker of the invention. Typically, the carrier compound is one that enhances the immune response to the oligosaccharide ligand.

[0067] Carrier compounds include but are not limited to proteins, peptides, polysaccharides, nucleic acids, or other polymers, lipids, and small oligomeric molecules, particularly the dendrimers. In some embodiments, the carrier compound may be naturally occurring or may be derived from a naturally occurring entity. Proteins include, for example, plasma proteins such as serum albumin, immunoglobulins, apolipoproteins and transferrin; bacterial polypeptides such as tetanus toxoid (TT), TRPLE, β -galactosidase, polypeptides such as herpes gD protein, allergens, diphtheria toxoid, salmonella flagellin, hemophilus pilin, hemophilus 15kDa, 28-30kDa and 40kDa membrane proteins, *Escherichia coli* heat label enterotoxin Irb, cholera toxin, and viral proteins including rotavirus VP and respiratory syncytial virus f and g proteins. The carriers useful in the invention include any protein that is safe for administration to mammals and optionally that is an immunologically effective carrier protein. Thus, in some embodiments, the carrier compound may itself be immunogenic. Examples include compounds that have been used in or as vaccines against a bacterial species such as but not limited to those listed herein.

[0068] Carrier compounds that are useful particularly for immunization include proteins such as keyhole limpet hemocyanin, serum albumin, bovine thyroglobulin, or soy bean trypsin inhibitor. Any other compound that is immunogenic in the species of animal to be immunized can similarly be used.

[0069] As shown in the Examples, the oligosaccharide to carrier ratio in the oligosaccharide-carrier conjugates of the

invention may vary. For example, an oligosaccharide-carrier conjugate may have an oligosaccharide to carrier ratio of 1:1, at least 2:1, at least 3:1, at least 4:1, at least 5:1, at least 6:1, at least 7:1, at least 8:1, at least 9:1 or at least 10:1. In still other embodiments, the ratio may be at least 20:1, at least 30:1, at least 40:1, at least 50:1, at least 60:1, at least 70:1, at least 80:1, at least 90:1, or greater, up to for example 100:1 or 700:1 depending on the capacity of carrier compound for attachment of oligosaccharide ligands. As an example, a conjugate that has an oligosaccharide to carrier ratio of at least 3:1 is a conjugate that has at least three oligosaccharide moieties attached to a single carrier compound.

Utility

[0070] The compositions of the invention have a number of *in vitro* and *in vivo* uses. For example, the compositions of the invention are useful for producing an antibody response, e.g., as a vaccine for active immunization of humans and animals to prevent infection by species of bacteria that make or are capable of making native PNAG, including but not limited to *Staphylococcus*; as a vaccine for immunization of humans or animals to produce anti-dPNAG antibodies that can be administered to other humans or animals to prevent or treat infections by species of bacteria that make or are capable of making native PNAG, including but not limited to *Staphylococcus*; as an antigen to screen for biological agents such as monoclonal antibodies capable of preventing infection by species of bacteria that make or are capable of making native PNAG, including but not limited to *Staphylococcus*, libraries of genes involved in making antibodies, or peptide mimetics; as a diagnostic reagent for infections by species of bacteria that make or are capable of making native PNAG, including but not limited to *Staphylococcus*; and as a diagnostic reagent for determining the immunologic status of humans or animals in regard to their susceptibility to infections by species of bacteria that make or are capable of making native PNAG, including but not limited to *Staphylococcus*.

Treatment and Prevention of Infections

[0071] The invention provides a method for treating or preventing infection by species of bacteria that make or are capable of making native PNAG, including but not limited to *Staphylococcus*, in a subject comprising administering to a subject having or at risk of developing such an infection an effective amount for inducing an immune response of an oligosaccharide-carrier conjugate of the invention.

[0072] Another aspect of the invention provides a method for evaluating the ability of a conjugates of the invention to protect against infection by species of bacteria that make or are capable of making native PNAG, including but not limited to *Staphylococcus*, in a subject. The method involves administering to the subject an effective amount of the conjugate, wherein the conjugate induces active immunity, exposing the subject to bacterial species that make or are capable of making native PNAG, and testing for the presence of the bacterial species in the subject.

[0073] The subject may be one that is subject is at risk of exposure to the bacterial species, or one that has been exposed to the bacterial species. The conjugate may be administered in a composition together with other agents such as but not limited to one or more adjuvants, and/or one or more anti-bacterial agents, etc.

[0074] The infection may be a *Staphylococcus* infection. The *Staphylococcus* infection may a *Staphylococcus aureus* infection, a *Staphylococcus epidermidis* infection, but it is not so limited. The infection may be an *E. coli* infection, a *Y. pestis* infection, a *Y. enterocolitica* infection, a *Y. pseudotuberculosis* infection, an *Aggregatibacter actinomycetemcomitans* infection, an *Actinobacillus pleuropneumoniae* infection, a *Bordetella pertussis* infection, a *B. parapertussis* infection, a *B. bronchiseptica* infection, an *Acinetobacter* infection including infection by *Acinetobacter* complex organisms, a *Burkholderia* infection including an infection by *Burkholderia* complex organisms, a *Stenatrophomonas maltophilia* infection, a *Shigella* (different species) infection, and a *Klebsiella* (different species) infection.

[0075] The antibodies generated according to the invention may also be used to prevent or treat infection by any infectious or pathogenic microbe that makes a molecule that reacts with the antibodies induced by an immune response to the oligosaccharide-carrier conjugate.

[0076] An "effective amount for inducing an immune response (e.g., an antibody response)" as used herein is an amount which is sufficient to (i) assist the subject in producing its own immune protection by e.g. inducing the production of antibodies in the subject, inducing the production of memory cells, and possibly a cytotoxic lymphocyte reaction etc. and/or (ii) prevent infection from occurring in a subject which is exposed to an infectious or pathogenic microbe that makes or is capable of making PNAG, including but not limited to *Staphylococcus*.

[0077] In some preferred embodiments, the effective amount of a vaccine for stimulating an immune response is an amount of vaccine that is capable of eliciting the production of antibodies that are cross-reactive with at least two species of *Staphylococcus*, e.g., *S. aureus* and *S. epidermidis*.

[0078] One of ordinary skill in the art can assess whether an amount is sufficient to induce active immunity by routine methods known in the art. For instance, the ability of a specific conjugate to produce antibody in a mammal can be assessed by screening for antibodies in a mouse or other subject using conjugates or their corresponding oligosaccharides.

5 [0079] Subjects as used herein include human and non-human subjects. Non-human subjects include but are not limited to companion animals such as dogs, cats, ferrets, birds, and the like, agricultural animals such as cows, pigs, goats, sheep, horses, chickens, and the like, zoo animals such as giraffes, lions, tigers, elephants, bears, and the like, laboratory animals such as mice, rats, rabbits, and the like. The subject may be a human over 60 years of age. The subject may be one that is healthy.

10 [0080] The subjects to be treated according to the invention include hospitalized patients who are at risk of developing *Staphylococcal* infection as a result of exposure to the bacteria in the hospital environment. Particular high risk populations for developing infection by *S. aureus* include, for example, renal disease patients on dialysis, and individuals undergoing high risk surgery. High risk populations for developing infection by *S. epidermidis* also include, for example, patients with indwelling medical devices, such as intravenous lines (e.g., central lines), or prostheses (e.g., hip or knee replacement prostheses), because clinical isolates are often highly adherent to plastic surfaces. In some embodiments, the subject is a subject that has received a medical device implant and in other embodiments, the subject is one that has not received a medical device implant but may be scheduled to receive one. Subjects at a high risk of developing infection by *S. epidermidis* further include, for example, pre-term neonates and patients undergoing chemotherapy. Additional subjects 15 to be treated according to the invention include hospitalized patients or individuals in the community who become ill and who are at risk of developing infections with microbes that make or are capable of making PNAG as a result of exposure to the bacteria in the hospital or community environments.

20 Immune Response Induction and Antibody Generation

25 [0081] The invention further provides methods for stimulating an immune response in a subject comprising administering to a subject in need thereof an oligosaccharide-carrier conjugate of the invention in an amount effective to stimulate an immune response in the subject. The immune response may be an antigen-specific immune response. It may be a cellular and/or a humoral immune response. For example, the immune response may result in the production of antibodies and/or antibody-producing cells.

30 [0082] "Passive immunity" as used herein involves the administration of antibodies to a subject, wherein the antibodies are produced in a different subject (including subjects of the same and different species), such that the antibodies attach to the surface of the bacteria and cause the bacteria to be phagocytosed.

35 [0083] The antibodies generated using the conjugates of the invention may be administered to any subject at risk of developing an infection by a species that makes PNAG or a species that makes another molecule that reacts with the antibodies to induce passive immunity, and in some embodiments may be particularly suited for subjects incapable of inducing active immunity. Since vaccination with the antigen might not be completely effective in high risk immunocompromised subjects, these subjects will benefit from treatment with antibody preparations raised against the oligosaccharide-carrier conjugates of the invention to prevent or treat infections such as those due to *Staphylococcus aureus*. A subject that is incapable of inducing an immune response is an immunocompromised subject (e.g. patient undergoing chemotherapy, AIDS patient, etc.) or a subject that has not yet developed an immune system (e.g. pre-term neonate).

40 [0084] Thus, the invention provides a method for producing antibodies comprising administering to a subject an effective amount for producing antibodies specific for the PNAG molecule expressed by an organism such as *Staphylococcus* using an oligosaccharide-carrier conjugate of the invention, and isolating antibodies from the subject. The antibodies may be polyclonal antibodies. The antibodies may be further modified.

45 [0085] The invention further provides a method for producing monoclonal antibodies comprising administering to a subject an effective amount for producing antibodies specific for the PNAG molecule using an oligosaccharide-carrier conjugate of the invention, harvesting antibody-producing cells from any tissue containing such cells such as spleen or blood from the subject, fusing the antibody-producing cells from the subject to myeloma cells, and harvesting antibody produced from a fusion subclone.

50 [0086] The invention contemplates the generation of a variety of antibodies specific to the oligo- β -(1 \rightarrow 6)-D-glucosamines of present invention. These include chimeric antibodies such as humanized antibodies and antibody fragments, as well as intact monoclonal and polyclonal antibodies. A "humanized monoclonal antibody" as used herein is a human monoclonal antibody or functionally active fragment thereof having at least human constant regions and an antigen binding region (such as 1, 2, 3, 4, 5, or 6 CDRs, or 1 or 2 variable regions, or Fab or F(ab)₂ fragments) from a species other than a human. Humanized monoclonal antibodies, for example, may be constructed by replacing the non-CDR regions of a non-human mammalian antibody with similar regions of human antibodies while retaining the epitopic specificity of the original antibody. For example, non-human CDRs and optionally some of the framework regions may be covalently joined to human FR and/or Fc/pFc' regions to produce a functional antibody. European Patent Application 0239400, the entire contents of which is hereby incorporated by reference, provides an exemplary teaching of the production and use of humanized monoclonal antibodies in which at least the CDR portion of a murine (or other non-human mammal) antibody is included in the humanized antibody. There are entities in the United States which will synthesize humanized antibodies from specific murine antibody regions commercially, such as Protein Design Labs

(Mountain View California), Abgenix, and Medarex.

[0087] An intact humanized monoclonal antibody in an isolated form or in a pharmaceutical preparation is particularly suited to some aspects of the invention. Humanized antibodies have particular clinical utility because they will not evoke an immune response in humans against the antibody itself. In one preferred embodiment, a murine CDR is grafted into the framework region of a human antibody to prepare the "humanized antibody." See, e.g., L. Riechmann et al., *Nature* 332, 323 (1988); M. S. Neuberger et al., *Nature* 314, 268 (1985) and EPA 0 239 400 (published Sep. 30, 1987).

[0088] Human monoclonal antibodies may be made by any of the methods known in the art, such as those disclosed in US Patent No. 5,567,610, issued to Borrebaeck et al., US Patent No. 565,354, issued to Ostberg, US Patent No. 5,571,893, issued to Baker et al, Kozber, *J. Immunol.* 133: 3001 (1984), Brodeur, et al., *Monoclonal Antibody Production Techniques and Applications*, p. 51-63 (Marcel Dekker, Inc, new York, 1987), and Boerner et al., *J. Immunol.*, 147: 86-95 (1991).

[0089] In addition to the conventional methods for preparing human monoclonal antibodies, such antibodies may also be prepared by immunizing transgenic animals that are capable of producing human antibodies (e.g., Jakobovits et al., *PNAS USA*, 90: 2551 (1993), Jakobovits et al., *Nature*, 362: 255-258 (1993), Brugermann et al., *Year in Immunol.*, 7:33 (1993) and US Patent No. 5,569,825 issued to Lonberg).

[0090] Human antibodies may also be obtained by recovering antibody-producing lymphocytes from the blood or other tissues of humans. These lymphocytes can be treated to produce cells that grow on their own in the laboratory under appropriate culture conditions. The cell cultures can be screened for production of antibody to the conjugates of the invention and then cloned. Clonal cultures can be used to produce human monoclonal antibodies, or the genetic elements 20 encoding the variable portions of the heavy and light chain of the antibody can be cloned and inserted into nucleic acid vectors for production of antibody of different types.

[0091] Antibody fragments are also encompassed by the invention. Well-known functionally active antibody fragments include but are not limited to $F(ab')_2$, Fab, Fv and Fd fragments of antibodies. These fragments which lack the Fc fragment of intact antibody, clear more rapidly from the circulation, and may have less non-specific tissue binding than an intact antibody (Wahl et al., *J. Nucl. Med.* 24:316-325 (1983)). For example, single-chain antibodies can be constructed in accordance with the methods described in U.S. Patent No. 4,946,778 to Ladner et al. Such single-chain antibodies include the variable regions of the light and heavy chains joined by a flexible linker moiety. Methods for obtaining a single domain antibody ("Fd") which comprises an isolated variable heavy chain single domain, also have been reported (see, for example, Ward et al., *Nature* 341:644-646 (1989), disclosing a method of screening to identify an antibody heavy 30 chain variable region (V_H single domain antibody) with sufficient affinity for its target epitope to bind thereto in isolated form). Methods for making recombinant Fv fragments based on known antibody heavy chain and light chain variable region sequences are known in the art and have been described, e.g., Moore et al., US Patent No. 4,462,334. Other references describing the use and generation of antibody fragments include e.g., Fab fragments (Tijssen, *Practice and Theory of Enzyme Immunoassays* (Elsevier, Amsterdam, 1985)), Fv fragments (Hochman et al., *Biochemistry* 12: 1130 35 (1973); Sharon et al., *Biochemistry* 15: 1591 (1976); Ehrlich et al., U.S. Patent No. 4,355,023) and portions of antibody molecules (Audilore-Hargreaves, U.S. patent No. 4,470,925). Thus, those skilled in the art may construct antibody fragments from various portions of intact antibodies without destroying the specificity of the antibodies.

Compositions and Pharmaceutical Preparations

[0092] The compositions of the invention, including for example the oligosaccharide-carrier conjugates of the invention, may be formulated together with a pharmaceutically acceptable vehicle. The term "pharmaceutically-acceptable vehicle" as used herein means one or more compatible solid or liquid filler, diluents or encapsulating substances which are suitable for administration to a human or other animal. In the present invention, the term "vehicle" denotes an organic 40 or inorganic ingredient, natural or synthetic, with which the active ingredient is combined to facilitate the application. The components of the pharmaceutical compositions also are capable of being commingled with the conjugates, and with each other, in a manner such that there is no interaction which would substantially impair the desired pharmaceutical efficiency.

[0093] Thus, the composition of present invention may be regarded as a pharmaceutical preparation. It may be used *in vivo*, but its use is not so limited. When used *in vivo*, it may be used in human or non-human subjects, whether for therapeutic, prophylactic or research purposes. As an example, the compositions may be used to generate antibodies and/or antibody producing cells in non-human subjects such as mice, rabbits, and other suitable animal hosts.

[0094] The invention therefore provides pharmaceutical preparations comprising any of the foregoing oligosaccharide-carrier conjugates, which may be used as vaccines. These preparations may comprise the conjugates in an amount effective to stimulate an immune response, such as an antigen-specific immune response. These preparations may comprise other constituents or components such as but not limited to adjuvants and anti-bacterial agents. Such preparations may routinely contain pharmaceutically acceptable concentrations of salt, buffering agents, preservatives, compatible carriers, adjuvants, and optionally other therapeutic ingredients.

[0095] A suitable carrier media for formulating a vaccine includes sodium phosphate-buffered saline (pH 7.4) or 0.125 M aluminum phosphate gel suspended in sodium phosphate-buffered saline at pH 6 and other conventional media. Generally, vaccines contain from about 5 to about 100 μ g, and preferably about 10-50 μ g of the antigen to elicit effective levels of antibody in warm-blooded mammals.

5 [0096] An adjuvant is any substance which is incorporated into or administered (simultaneously or otherwise) with an antigen, which potentiates the immune response to the antigen in the subject. Adjuvants include but are not limited to aluminum compounds, e.g., gels, aluminum hydroxide and aluminum phosphate, and Freund's complete or incomplete adjuvant (e.g., in which the antigen is incorporated in the aqueous phase of a stabilized water in paraffin oil emulsion). The paraffin oil may be replaced with different types of oils, e.g., squalene or peanut oil. Other materials with adjuvant properties include BCG (attenuated *Mycobacterium tuberculosis*), calcium phosphate, levamisole, isoprinosine, poly-anions (e.g., poly A:U), lentinan, pertussis toxin, lipid A, saponins, QS-21 and peptides, e.g. muramyl dipeptide, and immunostimulatory oligonucleotides such as CpG oligonucleotides. Rare earth salts, e.g., lanthanum and cerium, may also be used as adjuvants. The amount of adjuvants depends on the subject and the particular antigen used and can be readily determined by one skilled in the art without undue experimentation.

10 [0097] An anti-bacterial agent is an agent that kills bacteria (e.g., through lysis) or prevents their division. The use of antibiotics in the treatment of bacterial infection is routine. Anti-bacterial agents include penicillin G, penicillin V, ampicillin, amoxicillin, bacampicillin, cyclacillin, epicillin, hetacillin, pivampicillin, methicillin, nafcillin, oxacillin, cloxacillin, dicloxacillin, flucloxacillin, carbenicillin, ticarcillin, avlocillin, mezlocillin, piperacillin, amdinocillin, cephalexin, cephradine, cefadroxil, cefaclor, cefazolin, cefuroxime axetil, cefamandole, cefonicid, cefoxitin, cefotaxime, ceftizoxime, cefmenoxine, ceftriaxone, moxalactam, cefotetan, cefoperazone, ceftazidime, imipenem, clavulanate, timentin, sulbactam, neomycin, erythromycin, metronidazole, chloramphenicol, clindamycin, lincomycin, vancomycin, trimethoprim-sulfamethoxazole, aminoglycosides, quinolones, tetracyclines and rifampin. (See Goodman and Gilman's, Pharmacological Basics of Therapeutics, 8th Ed., 1993, McGraw Hill Inc.)

15 [0098] The conjugates may be attached covalently or non-covalently to other moieties including but not limited to detectable labels such as imaging agents, fluorophores, enzymes, and the like.

20 [0099] Compositions suitable for parenteral administration conveniently comprise a sterile aqueous preparation, which can be isotonic with the blood of the recipient. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution, and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono or di-glycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables. Carrier formulations suitable for subcutaneous, intramuscular, intraperitoneal, intravenous, etc. administrations may be found in Remington's Pharmaceutical Sciences, Mack Publishing Company, Easton, PA.

25 [0100] The preparations of the invention are administered in effective amounts. An effective amount, as discussed above, in some instances is that amount that will alone, or together with further doses, induce active or passive immunity depending on the subject. It is believed that doses ranging from 1 nanogram/kilogram to 100 milligrams/kilogram, depending upon the mode of administration, will be effective. The preferred range is believed to be between 500 nanograms and 500 micrograms/kilogram, and most preferably between 1 microgram and 100 micrograms/kilogram. The absolute amount will depend upon a variety of factors including whether the administration is performed on a high risk subject not yet infected with the bacteria or on a subject already having an infection, the concurrent treatment, the number of doses and the individual patient parameters including age, physical condition, size and weight. These are factors well known to those of ordinary skill in the art and can be addressed with no more than routine experimentation. It is preferred generally that a maximum dose be used, that is, the highest safe dose according to sound medical judgment.

30 [0101] Multiple doses of the compositions of the invention are contemplated. Generally immunization schemes involve the administration of a high dose of an antigen followed by subsequent lower doses of antigen after a waiting period of several weeks. Further doses may be administered as well. The dosage schedule for passive immunization would be quite different with more frequent administration if necessary. Any regimen that results in an enhanced immune response to bacterial infection and/or subsequent protection from infection may be used. Desired time intervals for delivery of multiple doses of a particular conjugate can be determined by one of ordinary skill in the art employing no more than routine experimentation.

35 [0102] A variety of administration routes are available. The particular mode selected will depend, of course, upon the particular conjugate selected, the particular condition being treated and the dosage required for therapeutic efficacy. The methods of this invention, generally speaking, may be practiced using any mode of administration that is medically acceptable, meaning any mode that produces effective levels of an immune response without causing clinically unacceptable adverse effects. Preferred modes of administration are parenteral routes. The term "parenteral" includes subcutaneous, intravenous, intramuscular, intraperitoneal, and intrasternal injection, or infusion techniques. Other routes include but are not limited to oral, nasal, dermal, sublingual, and local.

40 [0103] The following Examples are included for purposes of illustration and are not intended to limit the scope of the invention.

EXAMPLES


[0104] Aspects of the invention are further illustrated by the following non-limiting Examples. These Examples illustrate *inter alia* how to make the oligo- β -(1 \rightarrow 6)-D-glucosamine oligosaccharides and how to conjugate such oligosaccharides to carriers such as protein carriers. Such conjugates may be used, *inter alia*, as vaccines.

Example 1. Synthesis of N-hydroxysuccinimidyl 4-acetylsulfanylbutyrate 2

[0105]

10

15


Scheme 1. Synthesis of the linking reagent 2. a: CF₃COOSu, pyridine.

[0106] To a solution of the acid 1 (Hogg, J. Heather; Ollmann, Ian R.; Wetterholm, Anders; Andberg, Martina Blomster; Haeggstroem, Jesper; et al.; Chem. Europ. J.; EN; 4; 9; 1998; 1698-1713) (237 mg, 1.46 mmol) and N-hydroxysuccinimidyl trifluoroacetate (431 mg, 2.02 mmol) in dichloromethane (4 mL) was added pyridine (355 μ L, 4.4 mmol) and the resulting solution was stirred at room temperature for 2 h. The mixture was diluted with dichloromethane (50 mL), and washed with ice-cold 1 M HCl and water, and concentrated. Silica gel column chromatography (toluene - ethyl acetate 9:1) of the residue gave the active ester 2 (359 mg, 95%) as colorless syrup. ¹H NMR data (300 MHz, CDCl₃) δ 2.00 (m, 2 H, β -CH₂), 2.33 (s, 3 H, CH₃COS), 2.68 (t, 2 H, J 7.4 Hz, γ -CH₂), 2.81 (s, 4 H, 2 CH₂ of succinimide), 2.96 (t, 2 H, J 7.1 Hz, α -CH₂). ¹³C NMR data (62.9 MHz, CDCl₃): δ 24.7 (β -CH₂), 25.6 (CH₂ of succinimide), 27.9 (γ -CH₂), 29.7 (α -CH₂), 30.7 (CH₃COS), 168.0 (COON), 169.2 (CO of succinimide), 195.4 (CH₃COS).

Example 2. Synthesis of 4-nitrophenyl 4-acetylsulfanylbutyrate 3

[0107]

35

Scheme 2. Synthesis of the linking reagent 3. a: CF₃COOpNp, Et₃N.

40


50

[0108] To a solution of the acid 1 (233 mg, 1.44 mmol) (Hogg, J. Heather; Ollmann, Ian R.; Wetterholm, Anders; Andberg, Martina Blomster; Haeggstroem, Jesper; et al.; Chem. Europ. J.; EN; 4; 9; 1998; 1698-1713) and 4-nitrophenyl trifluoroacetate (470 mg, 2 mmol) in dichloromethane was added triethylamine (400 μ L, 2.88 mmol) and the solution was stirred for 2 h at room temperature. The mixture was worked-up as described for 2 and the active ester 3 (385 mg, 94%) was isolated by silica gel column chromatography (toluene - ethyl acetate 92:8). ¹H NMR data (300 MHz, CDCl₃) δ 2.03 (m, 2 H, β -CH₂), 2.36 (s, 3 H, CH₃COS), 2.68 (t, 2 H, J 7.2 Hz, γ -CH₂), 3.01 (t, 2 H, J 7.1 Hz, α -CH₂), 7.28, 8.26 (2 d, 4 H, aromatics). ¹³C NMR data (62.9 MHz, CDCl₃): δ 24.7 (β -CH₂), 28.0 (γ -CH₂), 30.7 (CH₃COS), 32.7 (α -CH₂), 122.5, 125.2, 145.3, 155.3 (aromatic C), 170.4 (COO), 195.5 (CH₃COS).

Example 3. Synthesis of the ligands 6 and 7 using linking reagent 2 or 3

[0109]

55



Scheme 3. Synthesis of the ligands. a: 1) H_2 , $\text{Pd}(\text{OH})_2/\text{C}$, MeOH/THF 2 or 3, Et_3N , CH_2Cl_2 ; b: $\text{N}_2\text{H}_4 \cdot \text{H}_2\text{O}$, EtOH , Δ ; c: DTT , NaHCO_3 , Ac_2O , $\text{MeOH/H}_2\text{O}$.

[0110] Nonasaccharide 4 (M.L. Gening, Y.E. Tsvetkov, G.B. Pier, N.E. Nifantiev, «Synthesis of oligo- β (1 \rightarrow 6)-glucosamines corresponding to the fragments of the surface polysaccharide of *Staphylococcus aureus*» *Carbohydr. Res.* 342 (2007), 567-575) (110 mg, 0.023 mmol) was dissolved in a mixture of MeOH (3 ml), THF (6 ml) and 1M HCl (0.2 ml) and $\text{Pd}(\text{OH})_2/\text{C}$ (110 mg) was added. The resultant mixture was stirred under hydrogen atmosphere for 1 h. The catalyst was filtered off and the solvents were evaporated. The residue was dissolved in a mixture of CH_2Cl_2 (2 ml) and DMF (1 ml) and then linker reagent 2 (12 mg, 0.046 mmol, solution in 1 ml of CH_2Cl_2) and Et_3N (100 μl) were added. After 30 minutes the mixture was diluted with toluene, concentrated and subjected to silica gel column chromatography (toluene/ Me_2CO , 4:1) to give 5 (98 mg, 98%) as a colorless foam. Introduction of the linker was confirmed by the presence of its characteristic signals in NMR spectra of the protected carbohydrate ligand. ^1H NMR (500 MHz, CDCl_3): δ 1.88 (m, 2 H, β - CH_2), 2.21 (t, 2 H, J 7.2 Hz, γ - CH_2), 2.31 (s, 3 H, CH_3COS), 2.88 (t, 2 H, J 7.1 Hz, α - CH_2). ^{13}C NMR data (125 MHz, CDCl_3): δ 25.7 (β - CH_2), 28.5 (α - CH_2), 30.6 (CH_3COS), 35.1 (γ - CH_2).

[0111] A mixture of the protected nonasaccharide 5 (95 mg, 0.02 mmol), EtOH (5 ml) and $\text{N}_2\text{H}_4 \cdot \text{H}_2\text{O}$ (0.5 ml) was stirred under reflux for 1 h and then concentrated and subjected to gel permeation chromatography (TSK gel Toyopearl HW 40S, 2.5 x 40 cm) in 0.1M aqueous AcOH to furnish nonasaccharide 6 (28 mg, 86%). Directly after gel chromatography SH-derivative may be obtained as it was detected by mass-spectrometry, but after storage in solution it converted to

corresponding disulfide as it was confirmed by ^{13}C NMR experiment. ^1H NMR (500 MHz, D_2O): δ 1.95 (m, 2 H, $\beta\text{-CH}_2$), 2.32 (t, 2 H, J 7.2 Hz, $\gamma\text{-CH}_2\text{SH}$), 2.71 (t, 2 H, J 7.1 Hz, $\alpha\text{-CH}_2$). ^{13}C NMR data (125 MHZ, D_2O): δ 29.7 ($\beta\text{-CH}_2$), 35.8 ($\gamma\text{-CH}_2\text{SH}$), 38.5 ($\alpha\text{-CH}_2$), 42.1 ($\gamma\text{-CH}_2\text{SS}$). Mass-spectra: calculated for $\text{C}_{61}\text{H}_{114}\text{N}_{10}\text{O}_{38}\text{S}$ 543.234 [$\text{M}+3\text{H}]^{3+}$, experimental 543.243 [$\text{M}+3\text{H}]^{3+}$.

5 [0112] Nonasaccharide **6** (15 mg) was dissolved in a mixture of water/MeOH (2 ml, 1:1 v/v) and dithiothreitol (15 mg) and NaHCO_3 (20 mg) were added. The resultant mixture was stirred for 5 minutes and then acetic anhydride (100 μl) was added and stirring was continued for 30 minutes before the solvents were evaporated. The product was purified by gel permeation chromatography (TSK gel Toyopearl HW 40S, 2.5 x 40 cm) in 0.1M aqueous AcOH to give acetylated product **7** (15 mg, 95%) ^1H NMR (500 MHz, D_2O): δ 1.75 (m, 2 H, $\beta\text{-CH}_2$), 2.11 (t, 2 H, J 7.2 Hz, $\gamma\text{-CH}_2$), 2.43 (s, 3 H, CH_3COS), 2.75 (t, 2 H, J 7.1 Hz, $\alpha\text{-CH}_2$). ^{13}C NMR data (125 MHZ, D_2O): δ 25.7 ($\beta\text{-CH}_2$), 28.5 ($\alpha\text{-CH}_2$), 30.6 (CH_3COS), 35.1 ($\gamma\text{-CH}_2$). Mass-spectra: calculated for $\text{C}_{81}\text{H}_{134}\text{N}_{10}\text{O}_{48}\text{S}$ 1024.418 [$\text{M}+2\text{H}]^{2+}$, experimental 1024.427 [$\text{M}+2\text{H}]^{2+}$

Example 4. Synthesis of the protected ligand 5 using linking reagent 3

15 [0113] Nonasaccharide **4** (60 mg, 0.013 mmol) was dissolved in a mixture of MeOH (1.5 ml), THF (3ml) and 1M HCl (0.1 ml) and $\text{Pd}(\text{OH})_2/\text{C}$ (60 mg) was added. The resultant mixture was stirred under hydrogen atmosphere for 1 h. The catalyst was filtered off and the solvents were evaporated. The residue was dissolved in a mixture of CH_2Cl_2 (2 ml) and DMF (1 ml) and then linker reagent **3** (20 mg, 0.07 mmol, solution in 1 ml of CH_2Cl_2) and Et_3N (50 μl) were added. After 20 h the mixture was diluted with toluene, concentrated and subjected to silica gel column chromatography (toluene/Me₂CO, 4:1) to give **5** (37 mg, 68%) as a colorless foam.

Example 5. Study of applicability of linking reagent 8 for the preparation of ligands for further conjugation with protein

[0114]

25

30


35

40

45

50

55




Scheme 4. Synthesis of the ligands.. a: 1)H₂, Pd(OH)₂/C, MeOH/THF 2) 8, Et₃N, CH₂Cl₂; b: N₂H₄•H₂O, EtOH, Δ.

[0115] Pentasaccharide **9** (150 mg, 0.026 mmol) (M.L. Gening, Y.E. Tsvetkov, G.B. Pier, N.E. Nifantiev, «Synthesis of oligo- β (1 \rightarrow 6)-glucosamines corresponding to the fragments of the surface polysaccharide of *Staphylococcus aureus*» Carbohydr. Res. 342 (2007), 567-575) was dissolved in a mixture of MeOH (1.5 ml), THF (3ml) and 1M HCl (0.1 ml) and Pd(OH)₂/C (150 mg) was added. The resultant mixture was stirred under hydrogen atmosphere for 1 h. The catalyst was filtered off and the solvents were evaporated. The residue was dissolved in a mixture of CH₂Cl₂(2 ml) and DMF (1 ml) and then linker reagent **8** (30 mg, 0.1 mmol, solution in 0.1 ml of CH₂Cl₂) and Et₃N (50 μ l) were added. After 1 h the mixture was diluted with toluene, concentrated and subjected to silica gel column chromatography (toluene/Me₂CO, 4:1) to give **10** (132 mg, 89%) as a colorless foam. ¹H NMR data (500 MHz, CDCl₃): δ 2.38 (s, 3H, CH₃COS), 3.53 (s, 2H, CH₂S); ¹³C NMR data (125 MHZ, CDCl₃): δ 30.2 (CH₃COS), 32.9 (CH₂S).

[0116] A mixture of the protected pentasaccharide **10** (100 mg, 0.017 mmol), EtOH (5 ml) and N₂H₄•H₂O (0.5 ml) was stirred under reflux for 1 h and then concentrated and subjected to gel permeation chromatography (TSK gel Toyopearl HW 40S, 2.5 x 40 cm) in 0.1M aqueous AcOH to furnish pentasaccharide **11** (32 mg, 93%). ¹H NMR data (500 MHz, D₂O): δ 7.17 (s, 1 H, CH=N); ¹³C NMR data (125 MHZ, D₂O): δ 134.6 (C=N-NH₂), 167.4 (C(O)-CH=N). Mass-spectra: calculated for C₃₅H₆₇N₈O₂₂ 951.438 [M+H]⁺, experimental 951.448 [M+H]⁺.

Example 6. Study of applicability of linking reagent 12 for the preparation of ligands for further conjugation with protein

[0117]




30 Scheme 5. Synthesis of the ligands.. a: 1) H_2 , $\text{Pd}(\text{OH})_2/\text{C}$, MeOH/THF 2) 12, Et_3N , CH_2Cl_2 ;
b: $\text{N}_2\text{H}_4 \cdot \text{H}_2\text{O}$, EtOH , Δ

[0118] Pentasaccharide 9 (100 mg, 0.017 mmol) (M.L. Gening, Y.E. Tsvetkov, G.B. Pier, N.E. Nifantiev, «Synthesis of oligo- β (1 \rightarrow 6)-glucosamines corresponding to the fragments of the surface polysaccharide of *Staphylococcus aureus*» Carbohydr. Res. 342 (2007), 567-575) was dissolved in a mixture of MeOH (1.5 ml), THF (3 ml) and 1M HCl (0.1 ml) and Pd(OH)₂/C (150 mg) was added. The resultant mixture was stirred under hydrogen atmosphere for 1 h. The catalyst was filtered off and the solvents were evaporated. The residue was dissolved in a mixture of CH₂Cl₂ (2 ml) and DMF (1 ml) and then linker reagent **12** (15 mg, 0.061 mmol, solution in 0.1 ml of CH₂Cl₂) and Et₃N (50 μ l) were added. After 1 h the mixture was diluted with toluene, concentrated and subjected to silica gel column chromatography (toluene/Me₂CO, 4:1) to give **13** (87 mg, 85%) as a colorless foam. ¹H NMR data (500 MHz, CDCl₃): δ 2.33 (s, 3H, CH₃COS), 2.52 (m, 2H, COCH₂), 3.15 (t, 2H, J 7.6, CH₂S); ¹³C NMR data (125 MHz, CDCl₃): δ 25.2 (CH₃COS), 29.2 (COCH₂), 35.9 (CH₂S).
[0119] A mixture of the protected pentasaccharide **13** (80 mg, 0.015 mmol), EtOH (5 ml) and N₂H₄ \cdot H₂O (0.5 ml) was stirred under reflux for 1 h and then concentrated and subjected to gel permeation chromatography (TSK gel Toyopearl HW 40S, 2.5 x 40 cm) in 0.1M aqueous AcOH to give complex mixture of nonidentified products.

Example 7. Preparation of the conjugate of tetanus toxoid with ligand 6 - "TT-9NH₂"

[0120]



Conjugate of Tetanus Toxoid with Ligand 6 ("TT-9NH₂")

Scheme 6. Conjugation with Ligand 6.

Step 1. Protein modification. Tetanus toxoid (4 mg in 120 μ l, stock solution) was diluted with 400 μ l of pH 7.2 buffer (0.1 M sodium phosphate, 0.15 M NaCl, 10 mM EDTA), the solution of SBAP (2.6 mg) in DMSO (80 μ l) was added and the mixture was incubated for 2 h at RT. Unreacted SBAP was removed using PD-10 column in pH 8.0 running buffer (0.1 M sodium phosphate, 0.15 M NaCl, 10 mM EDTA) and resultant 3.5 ml solution of modified protein was concentrated to 400 μ l.


Step 2. Disulfide reduction. Immobilized TCEP Disulfide Reducing Gel (200 μ l of 50% slurry in water) was centrifuged, an excess of water was removed and disulfide 6 (1.5 mg in 100 μ l of pH 8.0 buffer(0.1 M sodium phosphate, 0.15 M NaCl, 10 mM EDTA)) was added. After incubation on a rotor rack at room temperature for 45 min, the solution of SH-derivative was separated from the gel by centrifugation and immobilized TCEP was washed with the same pH 8.0 buffer (3 x 100 μ l).

Step 3. Conjugation. The solution of the ligand obtained in Step 2 (400 μ l in pH 8.0 buffer) was immediately combined with modified protein (400 μ l in pH 8.0 buffer, Step 1.) and stirred overnight at room temperature. After this time, the conjugate was separated from uncoupled components by gel filtration on Superose 6 prep-grade column. Fractions, containing TT-9NH₂ conjugate were pooled, concentrated and stored frozen at -20°C.

[0121] Chemical analysis of conjugate. Conjugate was analyzed for its contents of oligosaccharides using hexosamine assay described by Smith and Gilkerson (R.L. Smith and E. Gilkerson. 1979 Analytical Biochem. 98: 478-480) with compounds **6** as a standard and for protein with the Bradford assay (M.M. Bradford, 1976, Analytical Biochem. 72:248 - 254). According to these assays, conjugate TT-9NH₂ contains 74 carbohydrate ligands per protein molecule (x = 74).

50 Example 8. Preparation of the conjugate of tetanus toxoid with ligand 7 - "TT-9NAc"

[0122]



Conjugate of Tetanus Toxoid with Ligand 7 ("TT-9NAc")

Scheme 7. Conjugation with Ligand 7.

Step 1. Protein modification. TT was modified with SBAP as described above for conjugate TT-9NH₂.

Step 2. S-Acetyl deprotection. Nonasaccharide 7 (2.1 mg) was dissolved in 200 μ l of 7% aqueous NH_3 solution, the mixture was kept at room temperature for 1 hour and then lyophilized.

Step 3. Conjugation. Lyophilized oligosaccharide was immediately dissolved in 400 μ l of pH 8.0 buffer (0.1 M sodium phosphate, 0.15 M NaCl, 10 mM EDTA) and mixed with 400 μ l of TT-modified solution in the same buffer.

35 sodium phosphate, 0.15 M NaCl, 10 mM EDTA) and mixed with 400 μ l of TT-modified solution in the same buffer. Reaction mixture was stirred overnight at room temperature. After this time, the conjugate was separated from uncoupled components by gel filtration on Superose 6 prep-grade column. Fractions containing TT-9NAc conjugate were pooled, concentrated and stored frozen at -20°C.

[0123] Chemical analysis of conjugate. The analysis was performed the same way as for conjugate TT-9NH₂ (Example 7) and revealed that conjugate TT-9Nac contains 71 carbohydrate ligands per protein molecule ($\gamma = 71$).

Example 9. Antibody production using oligosaccharide conjugates

[0124] **Methods.** Rabbits were immunized subcutaneously with 10 µg polysaccharide equivalent of nonaglucosamine (i.e., 9 linked monomers) conjugated to tetanus toxoid (TT) twice, one week apart, with an equivalent volume of Specol adjuvant. On the third week, rabbits were immunized three times (i.e., on Monday, Wednesday and Friday) with 10 µg PS-equivalent IV in saline. After the last immunization, rabbits were rested for two weeks and blood was taken every two weeks. Data in this presentation incorporate results from the first (bleed 1) and second (bleed 2) sera obtained post-immunization

50 [0125] **Results.** FIGs. 1A and B show the data relating to binding of antisera raised to non-acetylated nona-glucosamine (9GlcNH_2) to PNAG (A) or dPNAG (B) from *S. aureus*. dPNAG used in this experiment was about 15% acetylated. However, it is to be understood that the level of acetylation may range from 0-40%. The data show that the antisera bound comparably to both PNAG and dPNAG.

[0126] FIGs. 2A and B show the data relating to binding of antisera raise to fully acetylated non-glucosamine (9GlcNAc) to PNAG or dPNAG from *S. aureus*. The data show that the antisera bound better to highly acetylated PNAG than to dPNAG.

[0127] FIGs. 3A and B show the data relating to the binding of antisera raised to TT-conjugated 9GlcNAc or 9GlcNH₂ to unconjugated 11GlcNAc or 11GlcNH₂. The data show that the antiserum raised against acetylated 9GlcNAc bound

better to acetylated 11GlcNAc than did the antiserum raised against non-acetylated 9GlcNH₂. The opposite was true for binding to 9GlcNH₂, as antisera to 9GlcNH₂-TT conjugate was off-scale at serum dilutions of less than 6400.

[0128] FIGs. 4 and 5 show results using antisera from bleed 1 against *S. aureus* MN8 and two USA300 strains. FIG. 4 compares the killing of *S. aureus* MN8 (CP8) by rabbit antisera (referred to as "bleed 1") to fully acetylated or non-acetylated 9-mer oligoglucosamine conjugated to tetanus toxoid (TT). FIG. 5 compares the killing of *S. aureus* MN8 by rabbit antisera (bleed 1) to fully acetylated or non-acetylated 9-mer oligoglucosamine conjugated to TT. Anti-dPNAG-TT conjugate antiserum was used as a comparator. FIG. 6 compares the killing of *S. aureus* LAC (NT, USA300) by rabbit sera (bleed 1) to fully acetylated or non-acetylated 9-mer oligoglucosamine conjugated to TT. FIG. 7 compares the killing of *S. aureus* SF8300 (NT, USA300) by rabbit sera (bleed 1) to fully acetylated or non-acetylated 9-mer oligoglucosamine conjugated to TT. FIG. 8 compares the killing of *S. aureus* LAC (NT, USA300) by rabbit sera (bleed 1) to fully acetylated or non-acetylated 9-mer oligoglucosamine conjugated to TT. In general, serum raised to 9GlcNH₂-TT demonstrated the best overall activity but in most assays was only slightly better than serum raised to 9GlcNAc-TT.

[0129] FIGs. 9-16 show killing data resulting from the use of bleed 2 sera. Rabbit anti dPNAG-TT control used as comparator for strains MN8, SF8300 and LAC. Goat anti-dPNAG-TT was used as a comparator for Newman (CP5), PS80 (CP8) and the isogenic strains Reynolds CP5, Reynolds non-typable and Reynolds CP8.

[0130] FIG. 9 compares the killing of *S. aureus* MN8 (CP8) by rabbit sera (bleed 2) to fully acetylated or non-acetylated 9-mer oligoglucosamine conjugated to TT. FIG. 10 compares the killing of *S. aureus* LAC (NT, USA300) by rabbit sera (bleed 2) to fully acetylated or non-acetylated 9-mer oligoglucosamine conjugated to TT. FIG. 11 compares the killing of *S. aureus* SF8300 (NT, USA300) by rabbit sera (bleed 2) to fully acetylated or non-acetylated 9-mer oligoglucosamine conjugated to TT. FIG. 12 compares the killing of *S. aureus* Newman (CP5) by rabbit sera (bleed 2) to fully acetylated or non-acetylated 9-mer oligoglucosamine conjugated to TT. FIG. 13 compares the killing of *S. aureus* PS80 by rabbit sera (bleed 2) to fully acetylated or non-acetylated 9-mer oligoglucosamine conjugated to TT. FIG. 14 compares the killing of *S. aureus* Reynolds (CP5) by rabbit sera (bleed 2) to fully acetylated or non-acetylated 9-mer oligoglucosamine conjugated to TT. FIG. 15 compares the killing of *S. aureus* Reynolds (non-typable) by rabbit sera (bleed 2) to fully acetylated or non-acetylated 9-mer oligoglucosamine conjugated to TT. FIG. 16 compares the killing of *S. aureus* Reynolds (CP8) by rabbit sera (bleed 2) to fully acetylated or non-acetylated 9-mer oligoglucosamine conjugated to TT.

[0131] Generally, a higher degree of killing was achieved using antisera raised to the oligosaccharide conjugates than with antiserum raised to dPNAG-TT. Killing of strains LAC and SF8300 was better with bleed 2 sera as compared with bleed 1 sera even though the ELISA binding curves are similar. Using bleed 2, a greater difference in killing using the antisera raised to 9GlcNH₂ is seen as compared to lysis using serum raised to 9GlcNAc.

Example 10. Rabbit antisera and opsonic killing of *E. coli*.

[0132] Rabbit antibodies in post-immunization rabbit antisera (as described above) mediated opsonic killing of two *E. coli* strains previously shown to produce PNAG, but not a third strain lacking the pga genes encoding the biosynthetic enzymes for PNAG in *E. coli* (FIG. 16A).

Example 11. Mouse skin abscess models.

[0133] FIGs. 17, 18 and 19 show the results of an in vivo study using a mouse skin abscess model and challenge with *S. aureus* strain LAC (USA300). Antisera raised to 9GlcNH₂ demonstrated protective efficacy against an *S. aureus* LAC (USA300) skin infection. Group 1 (labeled 9GlcNH₂-TT) was administered 0.2 ml of 9GlcNH₂-TT antiserum (bleed 2) intraperitoneally 24 hours prior to infection. Group 2 (labeled NRS) was administered 0.2 ml of normal rabbit serum (NRS) 24 hours prior to infection. Mice were infected with 2 x 10⁴ CFU (FIG. 17), 2 x 10⁵ CFU (FIG. 19), or 2 x 10⁶ CFU (FIG. 20) in a 100 µl subcutaneous injection (per abscess) with microdex beads (10 g/ml). *S. aureus* LAC strain was grown in TSB overnight, then washed and added to the microdex beads prior to administration. After 72 hours, each abscess was excised, resuspended in 1 ml of TSB, homogenized, diluted, and then 100 µl of homogenate were plated along with serial dilutions. The lower limit of detection was 10 CFU/abscess. FIGs. 17, 18 and 19 show that the number of CFU per abscess was greatly reduced in mice administered the 9GlcNH₂-TT antiserum as compared to the normal rabbit serum. FIG. 20 summarizes the results of FIGs. 18-20 showing that at every dose of *S. aureus*, the mice administered 9GlcNH₂ were better protected against the *S. aureus* challenge.

[0134] FIGs. 21 and 22 show the results of two in vivo studies using the same mouse skin abscess model and procedures described in the preceding paragraph but the challenge strains are two additional *S. aureus* strains, MN8 and Newman. Mice were infected with 1 x 10⁶ CFU of strain MN8 (FIG. 21) or 4 x 10⁶ CFU of strain Newman (FIG. 22) in a 100 µl subcutaneous injection (per abscess) with microdex beads (10 g/ml). FIGs. 21 and 22 show that the number of CFU per abscess was significantly reduced in mice administered the 9GlcNH₂-TT antiserum as compared to the normal rabbit serum.

[0135] FIG. 23 shows the inability of the antiserum raised to the 9GlcNH₂-TT conjugate vaccine to significantly (P>0.05)

reduce the CFU/abscess of *S. aureus* strains MN8Δ*ica* and NewmanΔ*ica*. These two strains have had the *ica* genetic locus removed and can no longer synthesize the PNAG surface polysaccharide. In the absence of the PNAG antigen, the antibody to the 9GlcNH₂ oligosaccharide cannot provide any protective immunity to *S. aureus* skin infections.

5 Example 12. Protective efficacy against *E. coli* peritonitis.

[0136] The protective efficacy of antibody to 9GlcNH₂-TT was tested in a lethal peritonitis model of *E. coli* infection. This antibody protected all immunized mice against infection caused by two PNAG-positive *E. coli* isolates (Table 1, UTI strains J and P) whereas all controls receiving NRS did not survive. There was no protection afforded by antibody to 9GlcNH₂-TT antiserum against PNAG-negative *E. coli* strain H.

10 **Table 1. Protective efficacy of antibody to 9GlcNH₂-TT against lethal peritonitis caused by *E. coli*.**

Challenge <i>E. coli</i> strain	Number of survivors out of total mice	P value (Fisher's exact test)	
	Anti-9GlcNH ₂ TT	NRS ^a	
J (PNAG ⁺)	8/8	0/8	0.0002
P (PNAG ⁺)	8/8	0/8	0.0002
H (PNAG ⁻)	0/8	0/8	1.0

15 ^a NRS: normal rabbit serum

20 Discussion of Results

[0137] Using the fully non-acetylated, synthetic GlcNH₂-TT conjugate vaccines, it has been found according to the invention that no acetates are required for generating high levels of opsonic and protective antibodies in animals, that conjugating a molecule as small as five GtcNH₂-monomers in size is sufficient for a robust immune response, and that these antibodies readily bind to highly N-acetylated PNAG, poorly acetylated dPNAG and the non-acetylated oligosaccharides. These protective antibodies will therefore bind to naturally occurring PNAG regardless of its composition, including its level of acetylation.

[0138] The invention further contemplates producing vaccines that comprise multiple antigens including the non-acetylated oligosaccharides of the invention. The synthesis of GlcNH₂-oligomers with a reducing end linker containing a reactive sulfhydryl group suggests that vaccines targeting microbes that make PNAG could be made more effective by conjugating the GlcNH₂-oligosaccharides to microbial proteins that function as virulence factors and vaccine antigens. For example, the LcrV protein of *Y. pestis* is a target for protection against plague (Garmory et al. Vaccine 22:947-57 (2004); Overheim et al. Infect. Immun. 73:5152-9 (2005); Quenee et al. Infect. Immun. 76:2025-2036 (2008)) but serologic variants of this protein are known among strains of *Y. pestis* circulating in central Asia (Anisimov et al. Clin. Microbiol. Rev. 17:434-464 (2004)), making it possible such strains could evade immunity engendered by a single LcrV vaccine component. As PNAG is expressed by *Y. pestis*, conjugating GlcNH₂-oligomers to LcrV might enhance the protective coverage of a plague vaccine. This approach to vaccine production is attractive since the synthetic version of dPNAG oligosaccharides can be produced fairly inexpensively and, importantly, will not have any other microbial contaminants.

[0139] Overall these findings indicate that small sized oligomers of beta(1→6)-linked glucosamine conjugated to a carrier protein can induce high titers of opsonic antibody that is also protective against experimental *S. aureus* skin infection and lethal peritonitis due to *E. coli*.

45 **EQUIVALENTS**

[0140] While several inventive embodiments have been described and illustrated herein, those of ordinary skill in the art will readily envision a variety of other means and/or structures for performing the function and/or obtaining the results and/or one or more of the advantages described herein, and each of such variations and/or modifications is deemed to be within the scope of the inventive embodiments described herein. More generally, those skilled in the art will readily appreciate that all parameters, dimensions, materials, and configurations described herein are meant to be exemplary and that the actual parameters, dimensions, materials, and/or configurations will depend upon the specific application or applications for which the inventive teachings is/are used. Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific inventive embodiments described herein. It is, therefore, to be understood that the foregoing embodiments are presented by way of example only and that, within the scope of the appended claims and equivalents thereto, inventive embodiments may be practiced otherwise than as specifically described and claimed. Inventive embodiments of the present disclosure are directed to each individual feature, system, article, material, kit, and/or method described herein. In addition, any combination of two or more such

features, systems, articles, materials, kits, and/or methods, if such features, systems, articles, materials, kits, and/or methods are not mutually inconsistent, is included within the inventive scope of the present disclosure.

[0141] All definitions, as defined and used herein, should be understood to control over dictionary definitions, definitions in documents incorporated by reference, and/or ordinary meanings of the defined terms.

[0142] All references, patents and patent applications disclosed herein are incorporated by reference with respect to the subject matter for which each is cited, which in some cases may encompass the entirety of the document.

[0143] The indefinite articles "a" and "an," as used herein in the specification and in the claims, unless clearly indicated to the contrary, should be understood to mean "at least one."

[0144] The phrase "and/or," as used herein in the specification and in the claims, should be understood to mean "either or both" of the elements so conjoined, i.e., elements that are conjunctively present in some cases and disjunctively present in other cases. Multiple elements listed with "and/or" should be construed in the same fashion, i.e., "one or more" of the elements so conjoined. Other elements may optionally be present other than the elements specifically identified by the "and/or" clause, whether related or unrelated to those elements specifically identified. Thus, as a non-limiting example, a reference to "A and/or B", when used in conjunction with open-ended language such as "comprising" can refer, in one embodiment, to A only (optionally including elements other than B); in another embodiment, to B only (optionally including elements other than A); in yet another embodiment, to both A and B (optionally including other elements); etc.

[0145] As used herein in the specification and in the claims, "or" should be understood to have the same meaning as "and/or" as defined above. For example, when separating items in a list, "or" or "and/or" shall be interpreted as being inclusive, i.e., the inclusion of at least one, but also including more than one, of a number or list of elements, and, optionally, additional unlisted items. Only terms clearly indicated to the contrary, such as "only one of" or "exactly one of," or, when used in the claims, "consisting of," will refer to the inclusion of exactly one element of a number or list of elements. In general, the term "or" as used herein shall only be interpreted as indicating exclusive alternatives (i.e. "one or the other but not both") when preceded by terms of exclusivity, such as "either," "one of," "only one of," or "exactly one of." "Consisting essentially of," when used in the claims, shall have its ordinary meaning as used in the field of patent law.

[0146] As used herein in the specification and in the claims, the phrase "at least one," in reference to a list of one or more elements, should be understood to mean at least one element selected from any one or more of the elements in the list of elements, but not necessarily including at least one of each and every element specifically listed within the list of elements and not excluding any combinations of elements in the list of elements. This definition also allows that elements may optionally be present other than the elements specifically identified within the list of elements to which the phrase "at least one" refers, whether related or unrelated to those elements specifically identified. Thus, as a non-limiting example, "at least one of A and B" (or, equivalently, "at least one of A or B," or, equivalently "at least one of A and/or B") can refer, in one embodiment, to at least one, optionally including more than one, A, with no B present (and optionally including elements other than B); in another embodiment, to at least one, optionally including more than one, B, with no A present (and optionally including elements other than A); in yet another embodiment, to at least one, optionally including more than one, A, and at least one, optionally including more than one, B (and optionally including other elements); etc.

[0147] It should also be understood that, unless clearly indicated to the contrary, in any methods claimed herein that include more than one step or act, the order of the steps or acts of the method is not necessarily limited to the order in which the steps or acts of the method are recited.

[0148] In the claims, as well as in the specification above, all transitional phrases such as "comprising," "including," "carrying," "having," "containing," "involving," "holding," "composed of," and the like are to be understood to be open-ended, i.e., to mean including but not limited to. Only the transitional phrases "consisting of" and "consisting essentially of" shall be closed or semi-closed transitional phrases, respectively, as set forth in the United States Patent Office Manual of Patent Examining Procedures, Section 2111.03.

[0149] The following represent possible embodiments according to the invention:

Paragraph 1. A compound of Formula I



Formula I

where X is any atom or group, Y is a sulfur blocking group, and n is greater than 1.

Paragraph 2. The compound of paragraph 1, wherein Y is an acyl group.

Paragraph 3. The compound of paragraph 1 or 2, wherein Y is an acetyl group, and the compound has the structure of Formula II

5



10

Formula II

wherein Ac is an acetyl group.


Paragraph 4. The compound of paragraph 1, wherein the compound is an activated ester of Formula I.

15

Paragraph 5. The compound of paragraph 1, wherein the compound is a cyano, azido or haloid derivative an activated ester of Formula I.

Paragraph 6. The compound of paragraph 4, wherein the compound has the structure of Formula III

20



25

Formula III


(N-hydroxysuccinimidyl 4-acetylsulfanyl butyrate)

wherein Ac is an acetyl group.

30

Paragraph 7. The compound of paragraph 4, wherein the compound has the structure of Formula IV

35



Formula IV

40

(N-nitrophenyl 4-acetylsulfanyl butyrate)

wherein Ac is an acetyl group.

45

Paragraph 8. A method for synthesizing N-hydroxysuccinimidyl 4-acetylsulfanyl butyrate (Formula III) comprising reacting 4-acetylsulfanylbutyric acid (Formula V)




50

Formula V

wherein Ac is an acetyl group, with N-hydroxysuccinimidyl trifluoroacetate (CF_3COOSu) to yield N-hydroxysuccinimidyl 4-acetylsulfanyl butyrate (Formula III).

55

Paragraph 9. A method for synthesizing N-nitrophenyl 4-acetylsulfanyl butyrate (IV) comprising reacting 4-acetylsulfanylbutyric acid (Formula V)



5


Formula V

wherein Ac is an acetyl group, with N-nitrophenyl trifluoroacetate (CF_3COOPn) to yield N-nitrophenyl 4-acetylsulfonyl butyrate (Formula IV).


10

Paragraph 10. A method for synthesizing an oligosaccharide conjugated to a carrier comprising reacting an oligosaccharide first with a compound having a structure of Formula Va or Vb

15



Formula Va



Formula Vb


20

wherein m is a number selected from 1 to 10, p is a number selected from 1 to 20, and R is H or an alkyl group, second with a compound of Formula I, II, III or IV, and third with a carrier, wherein the oligosaccharide is β -1-6 linked glucosamine.

25

Paragraph 11. A composition comprising an oligosaccharide bearing an O-linked linker, wherein the linker comprises

30



Formula VI.

35

Paragraph 12. The composition of paragraph 11, wherein the oligosaccharide is β -1-6 linked glucosamine.

Paragraph 13. The composition of paragraph 10 or 11, wherein the oligosaccharide is 2-20 monomers in length.

40

Paragraph 14. The composition of paragraph 10 or 11, wherein the oligosaccharide is 5-11 monomers in length.

45

Paragraph 15. The composition of paragraph 10 or 11, wherein the oligosaccharide is 7, 9 or 11 monomers in length.


Paragraph 16. The composition of paragraph 15, wherein the oligosaccharide is 100% acetylated.

50


Paragraph 17. The composition of paragraph 15, wherein the polysaccharide is less than 50% acetylated.

Paragraph 18. A composition comprising an oligosaccharide-carrier conjugate comprising an oligosaccharide conjugated to a carrier through a linker that is

55



Formula VIIa or



5

Formula VIIIb

wherein the linker is O-linked to the oligosaccharide and N-linked to the carrier.

Paragraph 19. The composition of paragraph 18, wherein the carrier is a peptide.

10

Paragraph 20. The composition of paragraph 18, wherein the carrier is a protein.

Paragraph 21. The composition of paragraph 18, wherein the carrier is tetanus toxoid.

Paragraph 22. The composition of paragraph 18, wherein the oligosaccharide is β -1-6 linked glucosamine.

Paragraph 23. The composition of paragraph 18, wherein the composition has a oligosaccharide to carrier ratio of 1:1, 2:1, 3:1, 4:1, 5:1, 6:1, 7:1, 8:1, 9:1, 10:1, 20:1, 30:1, 40:1, 50:1, 60:1, 70:1, 80:1, 90:1 or 100:1.

Paragraph 24. The composition of paragraph 22, wherein the oligosaccharide is 2-20 monomers in length.

Paragraph 25. The composition of paragraph 22, wherein the oligosaccharide is 5-11 monomers in length.

Paragraph 26. The composition of paragraph 22, wherein the oligosaccharide is 7, 9 or 11 monomers in length.

Paragraph 27. The composition of paragraph 22, wherein the oligosaccharide is 0-40% acetylated.


Paragraph 28. The composition of paragraph 18, further comprising a pharmaceutically acceptable carrier.

Paragraph 29. The composition of paragraph 18, further comprising an adjuvant.

Paragraph 30. The composition of paragraph 18, further comprising an anti-bacterial agent.

Paragraph 31. A method for synthesizing an oligosaccharide-carrier conjugate comprising reacting an oligosaccharide conjugated to a linker of Formula IXa or IXb


40



45


Formula IXa

50



Formula IXb

55 with a carrier having modified amino group following reaction with a compound of Formula



5

Formula X

to yield an oligosaccharide conjugated to a carrier compound through a linker having a structure of Formula VIIa or VIIb.

10

Paragraph 32. The method of paragraph 31, wherein the oligosaccharide-carrier conjugate has a oligosaccharide to carrier ratio of 1:1, 2:1, 3:1, 4:1, 5:1, 6:1, 7:1, 8:1, 9:1, 10:1, 20:1, 30:1, 40:1, 50:1, 60:1, 70:1, 80:1, 90:1 or 100:1.

15

Paragraph 33. The method of paragraph 31, wherein the carrier is a peptide.

15

Paragraph 34. The method of paragraph 31, wherein the carrier is a protein.

20

Paragraph 35. The method of paragraph 31, wherein the carrier is a tetanus toxoid.

20

Paragraph 36. The method of paragraph 31, wherein the oligosaccharide is β -1-6 linked glucosamine.

25

Paragraph 37. The method of paragraph 31, wherein the oligosaccharide is 2-20 monomers in length.

25

Paragraph 38. The method of paragraph 31, wherein the oligosaccharide is 5-11 monomers in length.

25

Paragraph 39. The method of paragraph 31, wherein the oligosaccharide is 7, 9 or 11 monomers in length.

25

Paragraph 40. The method of paragraph 36, wherein the oligosaccharide 0-40% acetylated.

30

Paragraph 41. A method for stimulating an immune response in a subject comprising administering to a subject in need thereof an oligosaccharide-carrier conjugate of any one of paragraphs 18 to 27 or an oligosaccharide-carrier conjugate synthesized by a method of any one of paragraphs 31 to 40 in an amount effective to stimulate an immune response in the subject.

35

Paragraph 42. The method of paragraph 41, wherein the subject is a human.

35

Paragraph 43. The method of paragraph 41, wherein the subject is non-human.

40

Paragraph 44. The method of paragraph 41, 42 or 43, further comprising isolating antibodies or antibody-forming cells from the subject.

45

Paragraph 45. The method of paragraph 41, 42, 43, or 44, further comprising administering an adjuvant to the subject.

45

Paragraph 46. The method of paragraph 41-44 or 45, further comprising administering an anti-bacterial agent to the subject.

50

Paragraph 47. A method for treating or preventing an infection in a subject comprising administering to a subject having or at risk of developing an infection an effective amount for inducing an immune response of an oligosaccharide-carrier conjugate of any one of paragraphs 18 to 27 or an oligosaccharide-carrier conjugate synthesized by a method of any one of paragraphs 31 to 40, wherein the infection is caused by a bacterial species that produces PNAG.

50

Paragraph 48. The method of paragraph 47, wherein the infection is a *Staphylococcus* infection.

55

Paragraph 49. The method of paragraph 47, wherein the *Staphylococcus* infection is *Staphylococcus aureus* infection.

Paragraph 50. The method of paragraph 47, wherein the *Staphylococcus* infection is *Staphylococcus epidermidis*.

Paragraph 51. The method of paragraphs 47-49 or 50, wherein the subject is at risk of exposure *Staphylococcus*.

Paragraph 52. The method of paragraph 47-49 or 50, wherein the subject has been exposed to *Staphylococcus*.

5 Paragraph 53. The method of paragraph 47-51 or 52, wherein the infection is an *E. coli* infection, a *Y. pestis* infection, a *Y. enterocolitica* infection, a *Y. pseudotuberculosis* infection, an *Aggregatibacter actinomycetemcomitans* infection, an *Actinobacillus pleuropneumoniae* infection, a *Bordetella pertussis* infection, a *B. parapertussis* infection, a *B. bronchiseptica* infection, an *Acinetobacter* infection, a *Burkholderia* infection, a *Stenatrophomonas maltophilia* infection, a *Shigella* infection, or a *Klebsiella* infection.

10 Paragraph 54. The method of paragraph 47-52 or 53, wherein the oligosaccharide-carrier conjugate is administered with an adjuvant.

15 Paragraph 55. A method for producing antibodies comprising:

15 administering to a subject an effective amount, for producing antibodies specific for PNAG, of an oligosaccharide-carrier conjugate of any one of paragraphs 18 to 27 or an oligosaccharide-carrier conjugate synthesized by a method of any one of paragraphs 31 to 40, and isolating antibodies from the subject.

20 Paragraph 56. A method for producing monoclonal antibodies comprising:

25 administering to a subject an effective amount, for producing antibodies specific for PNAG, of an oligosaccharide-carrier conjugate of any one of paragraphs 18 to 27 or an oligosaccharide-carrier conjugate synthesized by a method of any one of paragraphs 31 to 40, harvesting antibody-producing cells from the subject, fusing antibody-producing cells from the subject to myeloma cells, and harvesting antibody produced from a fusion subclone.

30 Paragraph 57. The method of paragraph 55, wherein the antibodies are polyclonal antibodies.

Paragraph 58. The method of paragraph 55 or 56, wherein the subject is a rabbit.

35 Paragraph 59. The method of paragraph 55 or 56, wherein the subject is human.

Claims

40 1. A method for synthesizing an oligosaccharide conjugated to a carrier comprising reacting an oligosaccharide first with a compound having a structure of Formula Va or Vb




Formula Va



Formula Vb

50 wherein m is a number selected from 1 to 10, p is a number selected from 1 to 20, and R is H or an alkyl group, second with a compound of Formula I, II, III or IV, and third with a carrier, wherein the oligosaccharide is β -1-6 linked glucosamine.

55 2. A composition comprising an oligosaccharide bearing an O-linked linker, wherein the linker comprises



5

Formula VI.

3. The composition of claim 2, wherein the oligosaccharide is β -1-6 linked glucosamine.
- 10 4. The composition of claim 1 or 2, wherein the oligosaccharide is 2-20 monomers in length.
5. The composition of claim 1 or 2, wherein the oligosaccharide is 5-11 monomers in length.
- 15 6. The composition of claim 1 or 2, wherein the oligosaccharide is 7, 9 or 11 monomers in length.
7. The composition of claim 6, wherein the oligosaccharide is 100% acetylated.
8. The composition of claim 6, wherein the polysaccharide is less than 50% acetylated.

20

25

30

35

40

45

50

55




Fig. 1A




Fig. 1B




Fig. 2A




Fig. 2B





Fig. 3B




Fig. 4




Fig. 5




Fig. 6




Fig. 7




Fig. 8




Fig. 9




Fig. 10




Fig. 11





Fig. 12






Fig. 14




Fig. 15




Fig. 16




Fig. 16A




Fig. 17




Fig. 18





Fig. 19






Fig. 21




Fig. 22




Fig. 23



EUROPEAN SEARCH REPORT

Application Number
EP 17 19 8371

5

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (IPC)
10	X WO 98/47915 A1 (NOVARTIS A.-G., SWITZ.; NOVARTIS-ERFINDUNGEN VERWALTUNGSGESELLSCHAFT M) 29 October 1998 (1998-10-29) * the whole document, in particular compounds A8-A10, A12 * -----	2,4,5	INV. C07D207/46 C07C327/20 C07C327/32 C07H3/06 C08B37/00 C07K14/31 A61K39/08 A61K39/085
15	X WO 97/19105 A1 (NOVARTIS AG, SWITZ.; GLYCOTECH CORP.) 29 May 1997 (1997-05-29) * the whole document *-----	2-8	
20	X THOMA G ET AL: "Novel glycodendrimers self-assemble to nanoparticles which function as polyvalent ligands in vitro and in vivo", ANGEWANDTE CHEMIE. INTERNATIONAL EDITION, VCH VERLAG, WEINHEIM, DE, vol. 41, no. 17, 2 September 2002 (2002-09-02), pages 3195-3198, XP002376807, ISSN: 0570-0833 * scheme 1 *-----	2,4	
25			TECHNICAL FIELDS SEARCHED (IPC)
30	X THOMA G ET AL: "SYNTHESIS OF OLIGOSACCHARIDE-POLYLYSINE CONJUGATES: A WELL CHARACTERIZED SIALYL LEWIS POLYMER FOR ELISA", JOURNAL OF THE AMERICAN CHEMICAL SOCIETY, AMERICAN CHEMICAL SOCIETY, WASHINGTON, DC; US, vol. 119, no. 31, 6 August 1997 (1997-08-06), XP000198261, ISSN: 0002-7863, DOI: DOI:10.1021/JA970657T * scheme 2 *-----	2,4	C07D C07H C08B C07K A61K
35			
40			
45			
4	The present search report has been drawn up for all claims		
50	Place of search Munich	Date of completion of the search 15 March 2018	Examiner Klein, Didier
	CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document	T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	
55			



EUROPEAN SEARCH REPORT

Application Number

EP 17 19 8371

5

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (IPC)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
10	X O. BLIXT ET AL: "Solid-Phase Enzymatic Synthesis of a Lewis a Trisaccharide Using an Acceptor Reversibly Bound to Sepharose", JOURNAL OF CARBOHYDRATE CHEMISTRY, vol. 16, no. 2, 1 February 1997 (1997-02-01), pages 143-154, XP055459203, UK ISSN: 0732-8303, DOI: 10.1080/07328309708006516 * page 146 *	2,4	
15			
20	X WRIGHT K ET AL: "Preparation of synthetic glycoconjugates as potential vaccines against Shigella flexneri serotype 2a disease", ORGANIC & BIOMOLECULAR CHEMISTRY, vol. 2, no. 10, 21 May 2004 (2004-05-21), pages 1518-1527, XP002305375, ISSN: 1477-0520, DOI: 10.1039/B400986J * the whole document *	1,4-8	
25			TECHNICAL FIELDS SEARCHED (IPC)
30			
35			
40			
45			
46	The present search report has been drawn up for all claims		
47	Place of search Munich	Date of completion of the search 15 March 2018	Examiner Klein, Didier
48	CATEGORY OF CITED DOCUMENTS		
49	X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document	T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	
50			
51			
52			
53			
54			
55			

ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.

EP 17 19 8371

5 This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

15-03-2018

10	Patent document cited in search report	Publication date		Patent family member(s)	Publication date
15	WO 9847915 A1 29-10-1998		AT	332918 T	15-08-2006
			AU	733282 B2	10-05-2001
			CA	2284729 A1	29-10-1998
			CN	1252807 A	10-05-2000
			DE	69835201 T2	14-06-2007
			EP	0970114 A1	12-01-2000
			ES	2268776 T3	16-03-2007
			JP	3474583 B2	08-12-2003
			JP	2001500528 A	16-01-2001
			PT	970114 E	29-12-2006
			US	6399071 B1	04-06-2002
			US	2002164347 A1	07-11-2002
			WO	9847915 A1	29-10-1998

25	WO 9719105 A1 29-05-1997		AU	7569796 A	11-06-1997
			EP	0866802 A1	30-09-1998
			JP	3372551 B2	04-02-2003
			JP	H11509861 A	31-08-1999
			US	6677164 B1	13-01-2004
			WO	9719105 A1	29-05-1997
30			-----		

35			-----		

40			-----		

45			-----		

50			-----		

55			-----		

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 61135493 A [0001]
- US 61208155 B [0001]
- WO 2004043405 A [0042]
- EP 0239400 A [0086] [0087]
- US 5567610 A, Borrebaeck [0088]
- US 565354 A, Ostberg [0088]
- US 5571893 A, Baker [0088]
- US 5569825 A, Lonberg [0089]
- US 4946778 A, Ladner [0091]
- US 4462334 A, Moore [0091]
- US 4355023 A, Ehrlich [0091]
- US 4470925 A, Audilore-Hargreaves [0091]

Non-patent literature cited in the description

- **GENING et al.** *Carbohydrate Research*, 2007, vol. 342, 567-575 [0060]
- **GENING et al.** *Russian J Bioorganic Chem*, 2006, vol. 32 (4), 389-399 [0060]
- **YANG ; DU.** *Carbohydrate Research*, 2003, vol. 338, 495-502 [0060]
- **YANG et al.** *Carbohydrate Research*, 2003, vol. 338, 1313-1318 [0060]
- **FRIDMAN et al.** *Organic Letters*, 2002, vol. 4 (2), 281-283 [0060]
- **G. HERMANSON.** *Bioconjugate Techniques*. Academic Press, 2008 [0065]
- **L. RIECHMANN et al.** *Nature*, 1988, vol. 332, 323 [0087]
- **M. S. NEUBERGER et al.** *Nature*, 1985, vol. 314, 268 [0087]
- **KOZBER.** *J. Immunol.*, 1984, vol. 133, 3001 [0088]
- **BRODEUR et al.** *Monoclonal Antibody Production Techniques and Applications*. Marcel Dekker, Inc, 1987, 51-63 [0088]
- **BOERNER et al.** *J. Immunol.*, 1991, vol. 147, 86-95 [0088]
- **JAKOBIVITS et al.** *PNAS USA*, 1993, vol. 90, 2551 [0089]
- **JAKOBIVITS et al.** *Nature*, 1993, vol. 362, 255-258 [0089]
- **BRUGGERMANN et al.** *Year in Immunol.*, 1993, vol. 7, 33 [0089]
- **WAHL et al.** *J. Nucl. Med.*, 1983, vol. 24, 316-325 [0091]
- **WARD et al.** *Nature*, 1989, vol. 341, 644-646 [0091]
- **TIJSSEN.** *Practice and Theory of Enzyme Immunoassays*. Elsevier, 1985 [0091]
- **HOCHMAN et al.** *Biochemistry*, 1973, vol. 12, 1130 [0091]
- **SHARON et al.** *Biochemistry*, 1976, vol. 15, 1591 [0091]
- **GOODMAN ; GILMAN'S.** *Pharmacological Basics of Therapeutics*. McGraw Hill Inc, 1993 [0097]
- *Remington's Pharmaceutical Sciences*. Mack Publishing Company [0099]
- **HOGG, J. HEATHER ; OLLMANN, IAN R. ; WETTERHOLM, ANDERS ; ANDBERG, MARTINA BLOMSTER ; HAEGGSTROEM, JESPER et al.** *Chem. Europ. J.*, 1998, vol. 4 (9), 1698-1713 [0106]
- **HOGG, J. HEATHER ; OLLMANN, IAN R. ; WETTERHOLM, ANDERS ; ANDBERG, MARTINA BLOMSTER ; HAEGGSTROEM, JESPER et al.** *Chem. Europ. J.*, 1998, vol. 4 (9), 1698-1713 [0108]
- **M.L. GENING ; Y.E. TSVETKOV ; G.B. PIER ; N.E. NIFANTIEV.** *Synthesis of oligo- β (1 \rightarrow 6)-glucosamines corresponding to the fragments of the surface polysaccharide of *Staphylococcus aureus**. *Carbohydr. Res.*, 2007, vol. 342, 567-575 [0110] [0115] [0118]
- **R.L. SMITH ; E. GILKERSON.** *Analytical Biochem.*, 1979, vol. 98, 478-480 [0121]
- **M.M. BRADFORD.** *Analytical Biochem.*, 1976, vol. 72, 248-254 [0121]
- **GARMORY et al.** *Vaccine*, 2004, vol. 22, 947-57 [0138]
- **OVERHEIM et al.** *Infect. Immun.*, 2005, vol. 73, 5152-9 [0138]
- **QUENEY et al.** *Infect. Immun.*, 2008, vol. 76, 2025-2036 [0138]
- **ANISIMOV et al.** *Clin. Microbiol. Rev.*, 2004, vol. 17, 434-464 [0138]

摘　　要

本發明涉及與載體綴合的合成的寡- β -(1→6)-2-氨基-2-去氧-D-吡喃葡萄糖苷的組合物，以及其製備方法和用途。