

(19) **DANMARK**



Patent- og
Varemærkestyrelsen

(10) **DK/EP 1692271 T4**

(12) Oversættelse af ændret
europæisk patentskrift

-
- (51) Int.Cl.: **C 12 N 1/21 (2006.01)** **C 12 P 7/46 (2006.01)** **C 12 N 15/63 (2006.01)**
C 12 N 15/74 (2006.01) **C 12 R 1/00 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2022-09-26**
- (80) Dato for Den Europæiske Patentmyndigheds
bekendtgørelse om opretholdelse af patentet i ændret form: **2022-08-03**
- (86) Europæisk ansøgning nr.: **04734158.1**
- (86) Europæisk indleveringsdag: **2004-05-20**
- (87) Den europæiske ansøgnings publiceringsdag: **2006-08-23**
- (86) International ansøgning nr.: **KR2004001210**
- (87) Internationalt publikationsnr.: **WO2005052135**
- (30) Prioritet: **2003-11-27 KR 20030084934** **2004-04-23 KR 20040028105**
- (84) Designerede stater: **AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IT LI LU MC NL PL PT
RO SE SI SK TR**
- (73) Patenthaver: **Korea Advanced Institute of Science and Technology, 373-1, Guseong-dong, Yuseong-gu, 305-701
Daejeon, Sydkorea**
- (72) Opfinder: **LEE, Sang Yup, EXPO Apt. 212-702, Jeonmin-dong, Yuseong-gu, Daejeon 305-761, Sydkorea
LEE, Sang Jun, Daewoo Apt. 305-604, Jijok-dong, Yuseong-gu, Daejeon 305-330, Sydkorea**
- (74) Fuldmægtig i Danmark: **Plougmann Vingtoft A/S, Strandvejen 70, 2900 Hellerup, Danmark**
- (54) Benævnelse: **HIDTIL UKENDTE VOMBAKTERIEVARIANTER OG FREMGANGSMÅDE TIL FREMSTILLING AF
RAVSYRE VED ANVENDELSE DERAFT**
- (56) Fremdragne publikationer:
WO-A-97/16528
WO-A1-02/00846
KR-A- 99 014 331
KR-A- 2002 003 712
US-A- 5 143 833
US-A- 5 521 075
**DIAZ-RICCI J C ET AL: "Effect of alteration of the acetic acid synthesis pathway on the fermentation pattern of
Escherichia coli" BIOTECHNOLOGY AND BIOENGINEERING, WILEY & SONS, HOBOKEN, NJ, US, vol. 38, no.
11, 1991, pages 1318-1324, XP002334441 ISSN: 0006-3592**
**MILLARD CYNTHIA SANVILLE ET AL: "Enhanced production of succinic acid by overexpression of
phosphoenolpyruvate carboxylase in Escherichia coli" APPLIED AND ENVIRONMENTAL MICROBIOLOGY,
WASHINGTON,DC, US, vol. 62, no. 5, May 1996 (1996-05), pages 1808-1810, XP002132795 ISSN: 0099-2240**
**LEE P.C. ET AL: 'Isolation and characterization of new succinic acid-producing bacterium, Mannheimia
succiniciproducents MBEL55E, from bovine rumen' APPL. MICROBIOL. BIOTECHNOL. vol. 58, 08 February
2002, pages 663 - 668**

Fortsættes ...

- LEE P.C. ET AL: 'Batch and continuous cultures of Mannheimia succiniciproducens MBEL55E for the production of succinic acid from whey and corn steep liquor' BIOPROCESS. BIOSYST. ENG vol. 26, 03 October 2003, pages 63 - 67
- ZEIKUS J.G. ET AL: 'Biotechnology of succinic acid production and markets for derived industrial products' APPL MICROBIOL BIOTECHNOL no. 51, 1999, pages 545 - 552
- HYOHAK SONG; SANG YUP LEE: 'Production of succinic acid by bacterial fermentation' ENZYME AND MICROBIAL TECHNOLOGY vol. 39, 2006, pages 352 - 361
- MCKINLAY ET AL APPL MICROBIL BIOTECHNOL vol. 76, 04 July 2007, pages 727 - 740
- SAMUELO ET AL: 'Influence of CO₂-HCO₃⁻ levels and pH on growth, succinate production and enzyme activities of Anaerobiospirillum succiniproducens' APPLIED AND ENVIRONMENTAL MICROBIOLOGY vol. 57, no. 10, 01 October 1991, pages 3013 - 3019
- SONG ET AL: 'modeling of batch fermentation kinetics for succinic acid production by Mannheimia succiniproducens' BIOCHEMICAL ENGINEERING JOURNAL vol. 40, 2008, pages 107 - 115
- SAMUELOV ET AL: 'whey fermentation by anaerobiospirillum succiniproducens for production of a succinate based animal feed additive' APPLIED AND ENVIRONMENTAL MICROBIOLOGY vol. 65, no. 5, 1999, pages 2260 - 2263
- VAN DER WERF ET AL: 'environmental and physiological factors affecting the succinate product ratio during carbohydrate fermentation by actinobacillus sp. 130z' ARCH MICROBIOL vol. 167, 1997, pages 332 - 342

DESCRIPTION

TECHNICAL FIELD

[0001] The present invention relates to a rumen bacterial mutant which produce succinic acid at high concentration while producing little or no other organic acids, as well as a method for producing succinic acid, which is characterized by the culture of such mutants in anaerobic conditions.

BACKGROUND ART

[0002] Various anaerobic microorganisms, including *Succinivibrio dextrinosolvens*, *Fibrobacter succinogenes*, *Ruminococcus flavefaciens* and the like, produce succinic acid as an end product by glucose metabolism (Zeikus, Annu. Rev. Microbiol., 34:423, 1980). Strains that produce succinic acid at industrially useful yield have not yet been reported except for *Anaerobiospirillum succiniciproducens* known to produce succinic acid at high concentration and high yield from glucose upon the presence of excessive CO₂ (David et al., Int. J. Syst. Bacteriol., 26:498, 1976). However, since *Anaerobiospirillum succiniciproducens* is an obligate anaerobic microorganism, a fermentation process of producing succinic acid using this microorganism has a shortcoming that the process itself becomes unstable even upon exposure to a very small amount of oxygen.

[0003] To overcome this shortcoming, *Mannheimia succiniciproducens* 55E was developed that is a strain having not only resistance to oxygen but also high organic acid productivity. However, since this strain produces formic acid, acetic acid and lactic acid in addition to succinic acid, it has shortcomings in that it has low yield and costs a great deal in a purification process of removing other organic acids except succinic acid.

[0004] Recombinant *E. coli* strains for the production of succinic acid have been reported in various literatures. If the *E. coli* strains have disruptions of a gene coding for lactate dehydrogenase and a gene coding for pyruvate formate-lyase, it is hard for them to grow in anaerobic conditions. Furthermore, they have too low yield to apply them to industrial field, since, although lactic acid is not produced as a fermentation product, other metabolites (acetic acid and ethanol) account for about half of the production of succinic acid. In an attempt to overcome such shortcomings, *E. coli* cells were grown in aerobic conditions, and then anaerobic conditions were applied to induce the fermentation of succinic acid. However, this attempt still has low productivity (Vemuri et al., J. Ind. Microbiol. Biotechnol., 28:325, 2002). Also, other examples were reported in which the genes of enzymes, such as pyruvate carboxylase, phosphoenolpyruvate carboxylase, phosphoenolpyruvate carboxykinase, and malic enzyme, that immobilize CO₂ in a metabolic pathway of succinic acid fermentation, are introduced into *E. coli*, thereby increasing the production of succinic acid (Vemuri et al., Appl.

Environ. Microbiol., 68:1715, 2002; Millard et al., Appl. Environ. Microbiol., 62:1808, 1996; Chao and Liao, Appl. Environ. Microbiol., 59:4261, 1993; Stols and Donnelly, Appl. Environ. Microbiol., 63:2695, 1997).

[0005] Meanwhile, it is known that the disruption of *ptsG* in *E. coli* contributes to an improvement of bacterial production and succinic acid production (Chatterjee et al., Appl. Environ. Microbiol., 67:148, 2001), but most of rumen bacteria have no *ptsG*, and thus have an advantage that they do not require a removal process of *ptsG* as in the case of *E. coli*. Recently, an attempt is actively conducted in which the genes of enzymes that immobilize CO₂ in a metabolic pathway of succinic acid fermentation are introduced into rumen bacteria, including genus *Actinobacillus* and genus *Anaerobiospirillum*. However, in this attempt, other organic acids were produced at large amounts or the yield of succinic acid was so low, as a result of that, it did not reach an industrially applicable level.

[0006] WO 02/00846 discloses an organic acid producing micro-organisms and a process for preparing organic acids employing the same.

[0007] WO 97/16528 discloses a method for isolating succinic, acid producing bacteria comprising increasing the biomass of an organism which lacks the ability to catabolize pyruvate, and then subjecting the biomass to glucose-rich medium in an anaerobic environment to enable pyruvate-catabolyzing mutants to grow.

DISCLOSURE OF INVENTION

[0008] Accordingly, during our extensive studies to develop bacterial strains that produce succinic acid at high yield, the present inventors constructed bacterial mutant *Mannheimia* sp. LPK (KCTC 10558BP) by the disruption of a lactate dehydrogenase gene (*ldhA*) and a pyruvate formate-lyase gene (*pfl*) from *Mannheimia succiniciproducens* 55E, which is a kind of rumen bacteria, and constructed bacterial mutants *Mannheimia* sp. LPK7 and LPK4, by the disruption of phosphotransacetylase gene (*pta*) and an acetate kinase gene (α ck4), and a phosphoenolpyruvate carboxylase gene (*ppc*), respectively from the LPK strain, and then confirmed that the culture of such bacterial mutants in anaerobic conditions provides succinic acid at high yield, thereby completing the present invention.

[0009] Therefore, a main object of the present invention is to provide a rumen bacterial mutant that produces succinic acid at high yield while producing no other organic acids, as well as a producing method thereof

[0010] Another object of the present invention is to provide a method of producing succinic acid, which is characterized by the culture of the above bacterial mutants in anaerobic conditions.

[0011] To achieve the above objects, in one aspect, the present invention provides a rumen

bacterial mutant in which a lactate dehydrogenase-encoding gene (*ldhA*) and a pyruvate formate-lyase-encoding gene (*pfl*) have been disrupted, and has the property of producing succinic acid at high concentration while producing little organic acids in anaerobic conditions, wherein the rumen bacterial mutant is *Mannheimia* sp. LPK according to Figure 3.

[0012] In another aspect, the present invention provides a rumen bacterial mutant in which a lactate dehydrogenase-encoding gene (*ldhA*), a pyruvate formate-lyase-encoding gene (*pfl*), a phosphotransacetylase-encoding gene (*pta*) and an acetate kinase-encoding gene (*ackA*) have been disrupted, and has the property of producing succinic acid at high concentration while producing little organic acids in anaerobic conditions, wherein the rumen bacterial mutant is *Mannheimia* sp. LPK7 according to Figure 8.

[0013] In still another aspect, the present invention provides a rumen bacterial mutant in which a lactate dehydrogenase-encoding gene (*ldhA*), a pyruvate formate-lyase-encoding gene (*pfl*), and a phosphoenolpyruvate carboxylase-encoding gene (*ppc*) have been disrupted, and has the property of producing succinic acid at high concentration while producing little organic acids in anaerobic conditions, wherein the rumen bacterial mutant is *Mannheimia* sp. LPK4 according to Figure 9.

[0014] In the present invention, the rumen bacteria according to the invention produce only succinic acid while producing little or no other organic acids.

[0015] In the method for producing the rumen bacterial mutant, the disruptions of the *ldhA* and *pfl* genes are performed by homologous recombination. The homologous recombination is performed using a genetic exchange vector containing a disrupted *ldhA* and a genetic exchange vector containing a disrupted *pfl*. The vector containing a disrupted *ldhA* is pMLKO-sacB, and the vector containing a disrupted *pfl* is pMPKO-sacB. (Embodiments not covered by the claimed invention.)

[0016] The disruptions of the *pta* and *ackA* genes are performed by homologous recombination. The homologous recombination is performed using a genetic exchange vector containing a disrupted *pta* and *ackA*. The genetic exchange vector containing a disrupted *pta* and *ackA* is pPTA-sacB.

[0017] The disruption of the *ppc* gene is performed by homologous recombination. The homologous recombination is performed using a genetic exchange vector containing a disrupted *ppc*. The genetic exchange vector containing a disrupted *ppc* is pPPC-sacB.

[0018] In the present invention, the rumen bacterial mutant having disruptions of a lactate dehydrogenase-encoding gene (*ldhA*) and a pyruvate formate-lyase-encoding gene (*pfl*) is *Mannheimia* sp. LPK (KCTC 10558BP).

[0019] Described outside the scope of the invention is a genetic exchange vector pMLKO-sacB containing a disrupted *ldhA*; a genetic exchange vector pMPKO-sacB containing a disrupted

pfl; a genetic exchange vector pPTA- *sacB* containing a disrupted *pta* and *ackA*; and a genetic exchange vector pPPC-*sacB* containing a disrupted *ppc*.

[0020] In another further aspect, the present invention provides a method for producing succinic acid, the method comprising the steps of: culturing the above rumen bacterial mutants in anaerobic condition; and recovering succinic acid from the culture broth.

[0021] As used herein, the term "disruption" means that the genes encoding the enzymes are modified such that the enzymes cannot be produced.

[0022] In the present invention, each of the lactate dehydrogenase gene (*ldhA*) and the pyruvate formate-lyase gene (*pfl*) was identified from the genomic information of *Mannheimia succiniciproducens* 55E, which is a kind of rumen bacteria, and then, all the two genes were removed from *Mannheimia succiniciproducens* 55E using a vector having disruptions of the genes, thereby constructing the bacterial mutant *Mannheimia* sp. LPK (KCTC 10558BP). Next, each of *pta-ackA* genes and a *ppc* gene was disrupted from the bacterial mutant *Mannheimia* sp. LPK, thereby constructing various bacterial mutants. Then, such bacterial mutants were confirmed to produce succinic acid at high concentration while producing little or no other organic acids.

[0023] The inventive bacterial mutants (*Mannheimia* sp. LPK, LPK4 and LPK7) are facultative anaerobic, gram-negative, non-mobile rods or coccobacilli, do not produce endospores, and can produce succinic acid in anaerobic conditions.

BRIEF DESCRIPTION OF DRAWINGS

[0024]

FIG. 1 shows a process of constructing a vector containing a disrupted *ldhA* (pMLKO-*sacB*).

FIG. 2 shows a process of constructing a vector containing a disrupted *pfl* (pMPKO-*sacB*).

FIG. 3 shows a process of constructing a bacterial mutant (LPK) by disrupting *ldhA* and *pfl* genes from *Mannheimia succiniciproducens* 55E.

FIG. 4 is an electrophoresis photograph showing the disruption of *ldhA* and *pfl* genes from *Mannheimia* sp. LPK (M: lambda *Hind*III size marker; lanes 1-3: PCR product LU1 & KM1 (1.5 kb); lanes 4-6: PCR product LD2 & KM2 (1.7 kb); lanes 7-9: PCR product PU1 & CM1 (2.2 kb); and lanes 10-12: PCR product PD2 & CM2 (1.6 kb)).

FIG. 5 shows the culture characteristics of *Mannheimia* sp. LPK in anaerobic conditions saturated with CO₂.

FIG. 6 shows a process of constructing vector containing a disrupted *pta* and *ackA* (pPTA-*sacB*).

FIG. 7 is a process of constructing a vector containing a disrupted *ppc* (pPPC-sacB).

FIG. 8 shows a process of constructing bacterial mutant LPK7 by disrupting *pta* and *ackA* genes from *Mannheimia* sp. LPK.

FIG. 9 shows a process of constructing bacterial mutant LPK4 by disrupting a *ppc* gene from *Mannheimia* sp. LPK.

FIG. 10 is an electrophoresis photograph showing the disruption of *pta* and *ackA* genes from *Mannheimia* sp. LPK7 (M: 1-kb ladder size marker; lane 1: PCR product P 13 & P 14 (1.1 kb); and lane 2: PCR product P 15 & P16 (1.5 kb)).

FIG. 11 is an electrophoresis photograph showing the disruption of a *ppc* gene from *Mannheimia* sp. LPK4 (M: 1-kb ladder size marker; lane 1: PCR product P 13 & P 17 (1.1 kb); and lane 2: PCR product P 15 & P18 (1.5 kb)).

FIG. 12 shows the cultivation characteristics of *Mannheimia* sp. LPK7 in anaerobic conditions saturated with O_2 .

FIG. 13 shows the cultivation characteristics of *Mannheimia* sp. LPK4 in anaerobic conditions saturated with CO_2 .

DETAILED DESCRIPTION OF THE INVENTION

[0025] The present invention will hereinafter be described in further detail by examples. It will however be obvious to a person skilled in the art that these Examples are given for illustrative purpose only, and the present invention is not limited to or by the examples.

[0026] Particularly, the following examples illustrate only a method comprising disrupting genes from a genus *Mannheimia* strain to obtain bacterial mutants and then producing succinic acid at high concentration by these bacterial mutants.

[0027] Furthermore, the following examples illustrate only a certain medium and culture method. However, the use of other mediums different from, such as whey, corn steep liquor (CSL), as described in literatures (Lee et al., Bioprocess Biosyst Eng., 26:63, 2003; Lee et al, Appl. Microbiol. Biotechnol., 58:663, 2002; Lee et al., Biotechnol. Lett., 25:111, 2003; Lee et al., Appl. Microbiol. Biotechnol., 54:23, 2000; Lee et al., Biotechnol. Bioeng., 72:41, 2001), and the use of various methods, such as fed-batch culture and continuous culture, will also be obvious to a person skilled in the art.

Example 1: Construction of pMLKO-sacB

[0028] In order to disrupt a lactate dehydrogenase gene (*ldhA*) by homologous recombination, a gene exchange vector was constructed in the following manner. First, the *genomic DNA* of *Mannheimia succiniciproducens* 55E QLCTC 0769BP), as a template, was subjected to PCR using primers set forth in SEQ ID NO: 1 and SEQ ID NO: 2 below, and then, the obtained PCR fragment was cut with *SacI* and *PstI* and introduced into pUC18 (New England Biolabs, Inc., Beverly, Mass.), thereby constructing pUC18-L1.

SEQ ID NO: 1: 5'-CAGTGAAGGAGCTCCGTAACGCATCCGCCG (LS1)

SEQ ID NO: 2: 5'-CTTTATCGAATCTGCAGGCGGTTTCCAAAA (LP1)

[0029] Thereafter, the genomic DNA of *Mannheimia succiniciproducens* 55E, as a template, was subjected to PCR using primers set forth in SEQ ID NO: 3 and SEQ ID NO: 4 below, and the resulting PCR fragment was cut with *PstI* and *HindIII* and introduced into the pUC18-L1, thereby constructing pUC18-L1-L2.

SEQ ID NO: 3: 5'-GTACTGTAAACTGCAGCTTTCATAGTTAGC (LP2)

SEQ ID NO: 4: 5'-GCCGAAAGTCAAGCTTGCCGTCGTTTAGTG (LH2)

[0030] pUC4K (Pharmacia, Freiburg, Germany) was cut with *PstI*, and the resulting kanamycin-resistant gene was fused with pUC18-L1-L2 cut with *PstI*, thereby constructing pUC18-L1-KmR-L2. A linker set forth in SEQ ID NO: 5 was inserted into the pUC18-L1-KmR-L2 cut with *SacI*, thereby making a new *XbaI* cutting site.

SEQ ID NO: 5: 5'-TCTAGAAGCT

[0031] PCR on pKmobSacB (Schafer et al., Gene, 145:69, 1994) as a template was performed using primers set forth in SEQ ID NO: 6 and 7 below, and the resulting PCR product was cut with *XbaI* and inserted into the above *XbaI* restriction enzyme site, thereby constructing pMLKO-sacB (FIG. 1).

SEQ ID NO: 6: 5'-GCTCTAGACCTTCTATCGCCTTCTTGACG (SXF)

SEQ ID NO: 7: 5'-GCTCTAGAGGCTACAA.AATCACGGGCGTC (SXR)

Example 2: Construction of pMPKO-sacB

[0032] In order to disrupt a pyruvate formate-lyase gene (*pfl*) by homologous recombination, a genetic exchange vector was constructed in the following manner. A pKmobsacB template containing a *sacB* gene (Genbank 02730) was subjected to PCR using primers set forth in SEQ ID NO: 8 and SEQ ID NO: 9 below. The resulting *sacB* product was cut with *Pst*I and *Bam*HI and inserted into pUC19 (Stratagene Cloning Systems, La Jolla, Calif.), thereby constructing pUC19-sacB.

SEQ ID NO: 8: 5'-AGCGGATCCCCTTCTATCGCCTTCTTGACG (SBG)

SEQ ID NO: 9: 5'-GTCCTGCAGGGCTACAAAATCACGGGCGTC (SPR)

[0033] The genomic DNA of *Mannheimia succiniciproducens* 55E, as a template, was subjected to PCR using primers set forth in SEQ ID NO: 10 and SEQ ID NO: 11 below. The resulting PCR fragment was cut with *Bam*HI and fused with the pUC19-sacB cut with *Bam*HI, thereby constructing pUC19-sacB-pfl.

SEQ ID NO: 10: 5'-CATGGCGGATCCAGGTACGCTGATTTTCGAT (PB1)

SEQ ID NO: 11: 5'-CAAGGATCCAACGGATAAAGCTTTTATTAT (PB2)

[0034] In order to obtain a chloramphenicol-resistant gene, pACYC184 (New England Biolabs, Inc., Beverly, Mass.) as a template was subjected to PCR using primers set forth in SEQ ID NO: 12 and SEQ ID NO: 13 below. The resulting PCR product was cut with *Sma*I and fused with the pUC19-sacB-pfl cut with *Bst*1107I, thereby constructing pMPKO-sacB (FIG. 2).

SEPA ID NO: 12: 5'-CTCGAGCCCCGGGGTTTAAGGGCACCAATAA (CTR)

SEQ ID NO: 13: 5'-CTCGAGCCCCGGGCTTTGCGCCGAATAAAT (CTF)

Example 3: Construction of *Mannheimia* sp. LPK strain

[0035] FIG. 3 shows a process of constructing a mutant strain (LPK) by disrupting *ldhA* and *pfl* genes from *Mannheimia succiniciproducens* 55E. *Mannheimia succiniciproducens* 55E was plated on LB-glucose medium containing 10 g/l of glucose, and cultured at 37°C for 36 hours. The colony formed was inoculated in 10 ml of LB-glucose liquid medium, and cultured for 12 hours. The culture broth which had been sufficiently grown was inoculated by 1% in 100 ml of LB-glucose liquid medium, and cultured in a shaking incubator at 200 rpm and 37°C.

[0036] When the culture broth reached an OD of about 0.2-0.3 after 4~hours, it was

centrifuged at 4°C and 4000 rpm for 10 minutes to collect cells. Then, the cells were resuspended in 200 ml of 10% glycerol solution at 4°C. The suspension was centrifuged at 4°C and 4000 rpm for 10 minutes, and the cells were collected and resuspended in 200 ml of 10% glycerol solution at 4°C, and then centrifuged at 4°C and 4000rpm for 10 minutes to collect the cells. The cells were suspended in glycerol at a volume ratio of 1:1, to obtain cell concentrate.

[0037] The cell concentrate thus obtained was mixed with the genetic exchange vectors pMLKO-sacB and pMPKO-sacB constructed in Examples 1 and 2, and then subjected to electroporation under conditions of 1.8 kV, 25 µF and 200 ohms. 1 ml of LB-glucose liquid medium was added to the electroporated mixture and cultured in a shaking incubator at 37°C and 200rpm for one hour. The culture broth was plated on LB-glucose solid medium containing a suitable antibiotic [Km (final concentration of 25 µg/ml) or Cm (6.8 µg/ml) and cultured at 37°C for 48 hours or more. In order to select a colony where only double crossover occurred, the colonies formed were streaked on LB-sucrose medium (LB medium with 100g/l sucrose) containing Km 25 µg/ml) or Cm (6.8µg/ml). After 24 hours, the formed colonies were streaked again on the same plate.

[0038] The colony (mutant) formed on the plate were cultured in LB-glucose liquid medium containing an antibiotic, and a genomic DNA was isolated from the cultured strain by the method described in Rochelle et al. (FEMS Microbiol. Lett., 100:59, 1992). PCR was performed using the isolated mutant genomic DNA as a template, and the PCR product was electrophoresed to confirm the disruption of *ldhA* and *pfl* genes from the PCR product.

[0039] In order to confirm the disruption of the *ldhA* gene, PCRs were performed twice in the following manners. First, the mutant genomic DNA as a template was subjected to PCR using primers set forth in SEQ ID NO: 14 and SEQ ID NO: 15.

SEQ ID NO: 14: 5'-GACGTTTCCCGTTGAATATGGC (KM1)

SEQ ID NO:15: 5'-CATTGAGGCGTATTATCAGGAAAC (LU1)

[0040] Then, the mutant genomic DNA as a template was subjected to PCR using primers set forth in SEQ ID NO: 16 and SEQ ID NO: 17 below. The products obtained in the two PCRs were subjected to gel electrophoresis to confirm the disruption of *ldhA* by their size (1.5 kb) (FIG. 4).

SEQ ID NO: 16: 5'-GCAGTTTCATTTGATGCTCGATG (KM2)

SEQ ID NO: 17: 5'-CCTCTTACGATGACGCATCTTTCC (LD2)

[0041] In order to confirm the disruption of *pfl*, PCRs were performed twice in the following manner. First, the mutant genomic DNA as a template was subjected to PCR using primers set

forth in SEQ ID NO: 18 and SEQ ID NO: 19 below.

SEQ ID NO: 18: 5'-GGTGGTATATCCAGTGATTTTTTTCTCCAT (CM1)

SEQ ID NO: 19: 5'-CTTTGCAACATTATGGTATGTATTGCCG (PU1)

[0042] Then, the mutant genomic DNA as a template was subjected to PCR using primers set forth in SEQ ID NO: 20 and SEQ ID NO: 21. The products obtained in the two PCRs were subjected to gel electrophoresis to confirm the disruption of *pfl* by their size (1.5kb) (FIG. 4). In FIG. 4, M represents a Lambda HindIII size marker, lanes 1-3 represent the PCR product LU1 & KM1 (1.5kb), lanes 4-6 represent the PCR product LD2 & KM2 (1.7kb), lanes 7-9 represent the PCR product PU1 & CM1 (2.2kb), and lanes 10-12 represent the PCR product PD2 & CM2 (1.6kb).

SEQ ID NO: 20: 5'-TACTGCGATGAGTGGCAGGGCGGGGCGTAA (CM2)

SEQ ID NO: 21: 5'-CCCCAGCATGTGCAAATCTTCGTCAC (PD2)

[0043] The disruption of *ldhA* was confirmed by the fact that the product resulted from the PCR using the primers (LU1 and KM1) of SEQ ID NO: 14 and SEQ ID NO: 15 has a size of 1.5 kb and at the same time the product resulted from the PCR using the primers (LD2 and KM2) of SEQ ID NO: 16 and SEQ ID NO: 17 has a size of 1.7 kb. And, the disruption of *pfl* was confirmed by the fact that the product resulted from the PCR using the primers (PU1 and CM1) of SEQ ID NO: 18 and SEQ ID NO: 19 has a size of 2.2 kb and at the same time the product resulted from the PCR using the primers (PD2 and CM2) of SEQ ID NO: 20 and SEQ ID NO: 21 has a size of 1.6 kb. The position of each primer is shown in FIG. 3. The mutant constructed by the above method, i.e., a bacterial mutant having disruptions of *ldhA* and *pfl*, was named "*Mannheimia* sp. LPK" and deposited under accession number KCTC 10881BP on November 26, 2003 in the Korean Collection for Type Cultures (KCTC), Korean Research Institute of Bioscience and Biotechnology (KRIBB).

Example 4: Fermentation characteristics of *Mannheimia* sp. LPK

[0044] In order to examine the fermentation characteristics of *Mannheimia* sp. LPK constructed in Example 3 above, the mutant was cultured in anaerobic conditions saturated with CO₂, and the resulting reaction product was analyzed. First, carbon dioxide was introduced into 100 ml of preculture medium consisting of 20g/L glucose, 5g/L polypeptone, 5g/L yeast extract, 3g/L K₂HPO₄, 1g/L NaCl, 1g/L (NH₄)₂SO₄, 0.2g/L CaCl₂ - 2H₂O, 0.2g/L MgCl₂ - 6H₂O and 10g/L MgCO₃, and then, *Mannheimia* sp. LPK was inoculated in the preculture medium and precultured at 39°C for 14 hours. Then, 0.9 L of culture medium

consisting of 20g/L glucose, 5g/L polypeptone, 5g/L yeast extract, 3g/L K_2HPO_4 , 1g/L NaCl, 5g/L $(NH_4)_2SO_4$, 0.2g/L $CaCl_2 \cdot 2H_2O$, 0.2g/L $MgCl_2 \cdot 6H_2O$ and 5g/L Na_2CO_3 was put in a 2.5-L culture tank, and 100 ml of the precultured microorganisms were inoculated in the culture medium and batch-cultured at 39°C and pH 6.5 while supplying carbon dioxide at a flow rate of 0.25vvm.

[0045] The concentration of cells in the culture broth was measured with a spectrophotometer (Ultraspec 3000, Pharmacia Biotech., Sweden), and the amounts of succinate, glucose, lactate, acetate and formate were measured by HPLC (Aminex HPX-87H column, Bio-Rad, USA) (FIG. 5). Symbols in FIG. 5, refer to changes in the concentrations of cells (●), succinate (○), glucose (■), formate (◇) and acetate (Δ) with the passage of culture time. As shown in FIG. 5, after 14 hours of culture, the concentration of consumed glucose was 20g/L and the concentration of produced succinate was 17.2g/L, indicating that the yield of succinate (the amount of produced succinate/the amount of consumed glucose) is 81% and the volume productivity of succinate (the concentration of produced succinate/elapsed time) is 1.23 g/L/h. The inventive method of producing succinic acid by culturing *Mannheimia* sp. LPK in anaerobic conditions saturated with CO_2 showed a great increase in yield as compared to that of producing succinic acid by culturing parent strain *Mannheimia succiniciproducens* 55E in anaerobic conditions saturated with CO_2 , and showed a ratio of succinic acid : acetic acid of 40.7:1, indicating that it can produce succinic acid with little or no by-products.

Example 5: Construction of pPTA-sacB

[0046] In order to disrupt a phosphotransacetylase gene (*pta*) and an acetate kinase gene (*ackA*) by homologous recombination, a genetic exchange vector was constructed in the following manner. First, the genomic DNA of *Mannheimia* sp. LPK (KCTC 10558BP), as a template, was amplified by PCR using primers set forth in SEQ ID NO: 22 and SEQ ID NO: 23 below, and the resulting PCR fragment was cut with *Xba*I and *Bam*HI and introduced into pUC19, thereby constructing pUC19-PTA1.

SEQ ID NO: 22: 5'-GCTCTAGATATCCGCAGTATCACTTTCTGCGC

SEQ ID NO:23: 5'-TCCGCAGTCGGATCCGGGTAAACCGCACAG

[0047] Thereafter, the genomic DNA of *Mannheimia* sp. LPK as a template was amplified by PCR using primers set forth in SEQ ID NO: 24 and SEQ ID NO: 25 below, and the resulting PCR fragment was cut with *Xba*I and *Sac*I and introduced into the pUC19-PTA1, thereby constructing pUC19-PTA12.

SEQ ID NO: 24: 5'-GGGGAGCTCGCTAACTTAGCTTCTAAAGGCCATGT TTCC

SEQ ID NO: 25: 5'-GCTCTAGATATCCGGGTCAATATCGCCGCAAC

[0048] As a template, plasmid pIC156 (Steinmetz et al., Gene, 142:79, 1994) containing a spectinomycin-resistant gene (GenBank X02588) was amplified by PCR using primers set forth in SEQ ID NO: 26 and SEQ ID NO: 27 below, and the resulting PCR fragment (spectinomycin-resistant gene) was cut with *EcoRV* and introduced into the pUC19-PTA12, thereby constructing pUC19-PTA1 S2 having the spectinomycin-resistant gene. The constructed pUC19-PTA1 S2 was cut with *SacI* and *BamHI* and introduced into pUC19-SacB (see Example 2), thereby constructing a pPTA-sacB vector (FIG. 6).

SEQ ID NO: 26: 5'-GAATTCGAGCTCGCCCGGGGATCGATCCTC

SEQ ID NO: 27: 5'-CCCGGGCCGACAGGCTTTGAAGCATGCAAATGTCAC

Example 6: Construction of pPPC-sacB

[0049] In order to disrupt a phosphoenolpyruvate carboxylase gene (*ppc*) by homologous recombination, a genetic exchange vector was constructed in the following manner. First, the genomic DNA of *Mannheimia* sp. LPK, as a template, was amplified by PCR using primers set forth in SEQ ID NO: 28 and SEQ ID NO: 29, and the resulting PCR fragment was cut with *XbaI* and *BanHI* and introduced into pUC 19, thereby constructing pUC19-PPC1.

SEQ ID NO: 28: 5'-TACGGATCCCCAGA.AAATCGCCCCCATGCCGA

SEQ ID NO: 29: 5'-GCTCTAGATATCGTTTGATATTGTTCCGCCACATTTG

[0050] Thereafter, the genomic DNA of *Mannheimia* sp. LPK, as a template, was subjected to PCR using primers set forth in SEQ ID NO: 30 and SEQ ID NO: 31, and the resulting PCR fragment was cut with *XbaI* and *SacI* and introduced into the pUC19-PPC1, thereby constructing pUC19-PPC12.

SEQ ID NO: 30: 5'-GCTCTAGATATCCGTCAGGAAAGCACCCGCCATAGC

SEQ ID NO: 31: 5'-GGGGAGCTCGTGTGGCGCTGCGGAAGTAAGGCAAAAATC

[0051] A spectinomycin-resistant gene cut with *EcoRV* (see Example 5) was introduced into the pUC19-PPC12 to construct pUC19-PPC1S2. The pUC19-PPC1S2 was cut with *SacI* and *BamHI* and introduced into the pUC19-SacB, thereby constructing a pPPC-sacB vector (FIG. 7).

Example 7: Construction of *Mannheimia* sp. LPK7 and LPK4 strains

[0052] FIG. 8 and FIG. 9 show processes of constructing mutant strains LPK7 and LPK4 by disrupting *pta-ackA* and *ppc* from *Mannheimia* sp. LPK, respectively. *Mannheimia* sp. LPK was plated on LB-glucose medium containing 10g/l glucose, and cultured at 37°C for 36 hours. The colony formed was inoculated in 10 ml LB-glucose liquid medium and cultured for 12 hours. The culture broth which had been sufficiently grown was inoculated by 1% in 100 ml LB-glucose liquid medium and cultured in a shaking incubator at 37°C.

[0053] Cell concentrate was collected from the resulting culture broth in the same manner as described in Example 3. The collected cell concentrate was mixed with the genetic exchange vectors pPTA-sacB and pPPC-sacB constructed in Examples 5 and 6, and then subjected to electroporation under conditions of 1.8 kV, 25°F and 200 ohms. The electroporated mixture was added with 1 ml of LB-glucose liquid medium and cultured in a shaking incubator at 200 rpm and 37°C for one hour.

[0054] The culture broth was plated on LB-glucose solid medium containing a spectinomycin antibiotic (final concentration: 50 (g/ml), and cultured at 37°C for at least 48 hours. In order to select a colony where double crossover occurred, the colonies formed were streaked on LB-sucrose medium (LB medium containing 100 g/l of sucrose) containing 50 (g/ml of spectinomycin. After 24 hours, the formed colonies were re-streaked on the same plate. The colony (mutant) formed on the plate was cultured in LB-glucose liquid medium containing an antibiotic, and a genomic DNA was isolated from the cultured strain by the method of Rochelle et al. The isolated mutant genomic DNA as a template was amplified by PCR, and the PCR product was electrophoresed to confirm the disruption of each of *pta-ackA* and *ppc*.

[0055] To confirm the disruption of *pta-ackA*, PCRs were performed twice in the following manner. First, the mutant genomic DNA as a template was subjected to PCR using primers set forth in SEQ ID NO: 32 and SEQ ID NO: 33 below. Then, the mutant genomic DNA as a template was subjected to PCR using primers set forth in SEQ ID NO: 34 and SEQ ID NO: 35.

SEQ ID NO: 32: 5'-CCTGCAGGCATGCAAGCTTGGGCTGCAGGTCGACTC

SEQ ID NO: 33: 5'-GCTGCCAAACAACCGAAAATACCGCAATAAACGGC

SEQ ID NO: 34: 5'-GCATGTAACCTTTACTGGATATAGCTAGAAAAGGCATCGGGGAG

SEQ ID NO: 35: 5'-GCAACGCGAGGGTCAATACCGAAGGATTTGCGCG

[0056] The products obtained in the two PCRs were subjected to gel electrophoresis to confirm the disruption of *pta-ackA* by their size (FIG. 10). In FIG. 10, M represents a 1-kb ladder size marker, lane 1 represents the PCR product P13 & P14 (1.1 kb), and lane 2 represents the PCR

product P15 & P16 (1.5 kb). The disruption of *pta-ackA* was confirmed by the fact the product resulted from the

[0057] PCR using the primers of SEQ ID NO: 32 and SEQ ID NO: 33 (P13 & P14) has a size of 1.1 kb at the same time the product resulted from the PCR using the primers of SEQ ID NO: 34 and SEQ ID NO: 35 (P15 & P16) has a size of 1.5 kb. The positions of the primers are shown in FIG. 8. The mutant strain constructed as described above, i.e., a strain resulted from the disruption of *pta-ackA* from *Mannheimia* sp. LPK, was named "*Mannheimia* sp. LPK7" and deposited under accession number "KCTC 10626BP" in KCTC, an international depositary authority.

[0058] Furthermore, to confirm the disruption of *ppc*, PCRs were performed twice in the following manner. First, the mutant genomic DNA as a template was subjected to PCR using primers set forth in SEQ ID NO: 32 and SEQ ID NPO: 36. Then, the mutant genomic DNA as a template was subjected to PCR using primers set forth in SEQ ID NO: 34 and SEQ ID NO: 37.

SEQ ID NO: 36: 5'-GATCCAGGGAATGGCACGCAGGCTTTCAACGCCGCC

SEQ ID NO: 37: 5'-GCAAAGCCAGAGGAATGGATGCCATTAACCAATAGCG

[0059] The products obtained in the two PCRs were subjected to gel electrophoresis to confirm the disruption of *ppc* by their size (FIG. 11). In FIG. 11, M represents a 1-kb ladder size marker, lane 1 is the PCR product P13 & P17 (1.1kb), and lane 2 represents the PCR product P15 & P18 (1.5kb). The disruption of *ppc* was confirmed by the fact that the product resulted from the PCR using the primers of SEQ ID NO: 32 and SEQ ID NO: 36 (P13 & P17) has a size of 1.1 kb at the same time the product resulted from the PCR using the primers of SEQ ID NO: 34 and SEQ ID NO: 37 (P15 & P18) has a size of 1.5 kb. The positions of the primers are shown in FIG. 9. The mutant strain constructed as described above, i.e., a strain resulted from the disruption of *ppc* from *Mannheimia* sp. LPK, was named "*Mannheimia* sp. LPK4".

Example 8: Fermentation characteristics of LPK7 and LPK4

[0060] In order to examine the fermentation characteristics of *Mannheimia* sp. LPK7 and LPK4 constructed in Example 7 above, the mutant strains were cultured in anaerobic conditions saturated with CO₂, and the resulting reaction products were analyzed. First, carbon dioxide was introduced into 200ml of the preculture medium as described in Example 4, and each of *Mannheimia* sp. LPK7 and LPK4 was inoculated in the preculture medium and precultured at 39°C for 24 hours. Next, 1.8 L of a culture medium, which is the same as that in Example 4 except that glucose concentration is 18 g/L (final 100mM), was put in a 6.6 L culture tank, and 100 ml of the precultured microorganisms was inoculated in the culture medium and then batch-cultured at 39°C and pH 6.5 while supplying carbon dioxide at a flow rate of 0.25wm.

[0061] The concentrations of cells, succinate, glucose, lactate, acetate and formate were measured in the same manner as in Example 4 (FIG. 12 and FIG. 13). Symbols in FIG. 12 and FIG. 13 refer to changes in the concentrations of cells (● in upper portion), succinate (● in lower portion), glucose (□), formate (◆) and acetate (▲) with the passage of culture time. As shown in FIG. 12, after 22 hours of the culture of *Mannheimia* sp. LPK7, the concentration of consumed glucose was 100mM and the concentration of produced succinate was 124mM, indicating that the yield of succinate (the amount of produced succinate/the amount of consumed glucose) is 124 mol%. And, the production of acetate was remarkably reduced (Table 1). Also, as shown in FIG. 13, after 22 hours of the culture of *Mannheimia* sp. LPK4, the concentration of consumed glucose was 100mM and the concentration of produced succinate was 123.7mM, indicating that the yield of succinate (the amount of produced succinate/the amount of consumed glucose) is 123.7 mol%. And, the production of acetate was greatly reduced as compared to that in the wild type (Table 1).

[0062] The inventive method of producing succinic acid by culturing *Mannheimia* sp. LPK7 in anaerobic conditions saturated with CO₂ showed a great increase in the yield of succinic acid and also a 9.8 times increase in the ratio of succinic acid: acetic acid, as compared to that of producing succinic acid by culturing parent strain *Mannheimia succiniciproducens* 55E in anaerobic conditions saturated with CO₂, indicating that the inventive method can produce succinic acid with producing little or no byproducts (Table 1).

[0063] As reported by Bulter et al., even if acetate-producing genes in microorganisms known till now are all disrupted, a significant amount of acetate is produced in amino acid and fatty acid metabolisms which are still not established (Bulter et al. PNAS, 101:2299, 2004). Thus, the present invention cut off all acetate production pathways known till now, and achieved succinate fermentation at high yield and concentration.

Table 1: Comparison of products from fermentation of LPK4 and LPK7 and product from fermentation of 55E in anaerobic conditions

Strain	Fermentation products (mM)						S/A ratio (fold)
	Succinate	Acetate	Formate	Lactate	Pyruvate	Ethanol	
55E	99.1	40.6	53.8	8.2	13	<1.0	2.44 (1.0)
LPK4	123.7±6.2	28.1±5.4	ND	ND	12.2±6.3	<1.0	4.40(1.8)
LPK7	124.0±5.2	5.2±0.2	ND	ND	36.36±4.7	<1.0	23.84(9.8)

[0064] While the present invention has been described in detail with reference to the specific features, it will be apparent to persons skilled in the art that this description is only for a preferred embodiment and does not limit the scope of the present invention. Thus, the substantial scope of the present invention will be defined by the appended claims.

INDUSTRIAL APPLICABILITY

[0065] As described and provided above in detail, *Mannheimia* sp. mutant strains (LPK, LPK7 and LPK4) produce succinic acid in anaerobic conditions saturated with CO₂ and are facultative anaerobic strains having high resistance to oxygen. Thus, the production of succinic acid using such mutants can not only eliminate the fermentation process instability caused by oxygen exposure, etc., but also eliminate the production of other organic acids, as compared to the prior method of producing succinic acid using obligate anaerobic strains, thereby making it possible to optimize and maximize a purification process and production yield.

Sequence Listing

[0066]

<110> Korea Advanced Institute of Science and Technology

<120> NOVEL RUMEN BACTERIA VARIANTS AND PROCESS FOR PREPARING SUCCINIC ACID EMPLOYING THE SAME

<130> PP-B0038

<150> KR 10-2003-0084934

<151> 2003-11-27

<150> KR 10-2004-0028105

<151> 2004-04-23

<160> 37

<170> KopatentIn 1.71

<210> 1

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer LS1

<400> 1

cagtgaagga gctccgtaac gcatccgccg 30

<210> 2

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer LP1

<400> 2

ctttatcgaa tctgcaggcg gtttcaaaa 30

<210> 3

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer LP2

<400> 3

gtactgtaaa ctgcagcttt catagttagc 30

<210> 4

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer LH2 .

<400> 4

gccgaaagtc aagcttgccg tcgttagtg 30

<210> 5

<211> 10

<212> DNA

<213> Artificial Sequence

<220>

<223> Linker 1

<400> 5

tctagaagct 10

<210> 6

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer SXF

<400> 6

gctctagacc ttctatgcc ttctgacg 29

<210> 7

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer SXR

<400> 7

gctctagagg ctacaaaatc acgggcgtc 29

<210> 8

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer SBG

<400> 8

agcggatccc cttctatcgc cttcttgacg 30

<210> 9

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer SPR

<400> 9

gtcctgcagg gctacaaaat cacgggcgtc 30

<210> 10

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer PB1

<400> 10

catggcggat ccaggtacgc tgatttcgat 30

<210> 11

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer PB2

<400> 11

caaggatcca acggataaag cttttattat 30

<210> 12

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer CTR

<400> 12

ctcgagcccg gggtttaagg gcaccaataa 30

<210> 13

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer CTF

<400> 13

ctcgagcccc gggctttgcg ccgaataaat 30

<210> 14

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer KM1

<400> 14

gacgtttccc gttgaatatg gc 22

<210> 15

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer LU1

<400> 15

cattgaggcg tattatcagg aaac 24

<210> 16

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer KM2

<400> 16

gcagtttcat ttgatgctcg atg 23

<210> 17

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer LD2

<400> 17

cctcttacga tgacgcatct ttcc 24

<210> 18

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer CM1

<400> 18

ggtggtatat ccagtgattt tttctccat 30

<210> 19

<211> 28

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer PU1

<400> 19

ctttgcaaca ttatggtatg tattgccg 28

<210> 20

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer CM2

<400> 20

tactgcgatg agtggcaggg cggggcgtaa 30

<210> 21

<211> 26

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer PD2

<400> 21

ccccagcatg tgcaaattctt cgtcac 26

<210> 22

<211> 32

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer

<400> 22

gctctagata tccgcagtat cactttctgc gc 32

<210> 23

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer

<400> 23

tccgcagtcg gatccgggtt aaccgcacag 30

<210> 24

<211> 39

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer

<400> 24

ggggagctcg ctaacttagc ttctaaaggc catgtttcc 39

<210> 25

<211> 32

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer

<400> 25

gctctagata tccgggtcaa tatcgccgca ac 32

<210> 26

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer

<400> 26

gaattcgagc tcgcccgggg atcgatcctc 30

<210> 27

<211> 36

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer

<400> 27

cccgggccga caggcttga agcatgcaaa tgtcac 36

<210> 28

<211> 32

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer

<400> 28

tacggatccc cagaaaatcg ccccatgcc ga 32

<210> 29
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer

<400> 29
gctctagata tcgttgata ttgtccgcc acatttg 37

<210> 30
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer

<400> 30
gctctagata tccgtcagga aagcaccgc catagc 36

<210> 31
<211> 39
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer

<400> 31
ggggagctcg tgtggcgctg cggaagtaag gcaaaaatc 39

<210> 32
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer

<400> 32
cctgcaggca tgcaagcttg ggctgcaggt cgactc 36

<210> 33
<211> 35
<212> DNA
<213> Artificial Sequence

<220>

<223> Primer

<400> 33

gctgccaaac aaccgaaaat accgcaataa acggc 35

<210> 34

<211> 43

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer

<400> 34

gcatgtaact ttactggata tagctagaaa aggcacggg gag 43

<210> 35

<211> 34

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer

<400> 35

gcaacgcgag ggtcaatacc gaaggatttc gccg 34

<210> 36

<211> 36

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer

<400> 36

gatccaggga atggcacgca ggctttcaac gccgcc 36

<210> 37

<211> 37

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer

<400> 37

gcaaagccag aggaatggat gccattaacc aatagcg

37

REFERENCES CITED IN THE DESCRIPTION

Cited references

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- [WO0200846A](#) [\[0006\]](#)
- [WO9716528A](#) [\[0007\]](#)
- [KR1020030084934](#) [\[0066\]](#)
- [KR1020040028105](#) [\[0066\]](#)

Non-patent literature cited in the description

- **ZEIKUS** Annu. Rev. Microbiol., 1980, vol. 34, 423- [\[0002\]](#)
- **DAVID et al.** Int. J. Syst. Bacteriol., 1976, vol. 26, 498- [\[0002\]](#)
- **VEMURI et al.** J. Ind. Microbiol. Biotechnol., 2002, vol. 28, 325- [\[0004\]](#)
- **VEMURI** Appl. Environ. Microbiol., 2002, vol. 68, 1715- [\[0004\]](#)
- **MILLARD et al.** Appl. Environ. Microbiol., 1996, vol. 62, 1808- [\[0004\]](#)
- **CHAOLIAO** Appl. Environ. Microbiol., 1993, vol. 59, 4261- [\[0004\]](#)
- **STOLSDONNELLY** Appl. Environ. Microbiol., 1997, vol. 63, 2695- [\[0004\]](#)
- **CHATTERJEE et al.** Appl. Environ. Microbiol., 2001, vol. 67, 148- [\[0005\]](#)
- **LEE et al.** Bioprocess Biosyst Eng., 2003, vol. 26, 63- [\[0027\]](#)
- **LEE et al.** Appl. Microbiol. Biotechnol., 2002, vol. 58, 663- [\[0027\]](#)
- **LEE et al.** Biotechnol. Lett., 2003, vol. 25, 111- [\[0027\]](#)
- **LEE et al.** Appl. Microbiol. Biotechnol., 2000, vol. 54, 23- [\[0027\]](#)
- **LEE et al.** Biotechnol. Bioeng., 2001, vol. 72, 41- [\[0027\]](#)

- **SCHAFER et al.**Gene, 1994, vol. 145, 69- [\[0031\]](#)
- **ROCHELLE et al.**FEMS Microbiol. Lett., 1992, vol. 100, 59- [\[0038\]](#)
- **STEINMETZ et al.**Gene, 1994, vol. 142, 79- [\[0048\]](#)
- **BULTER et al.**PNAS, 2004, vol. 101, 2299- [\[0053\]](#)

Patentkrav

1. Vombakteriemutant, hvor et lactatdehydrogenase-kodende gen (*ldhA*) og et pyruvatformiatlyase-kodende gen (*pfl*) er blevet afbrudt, og som har den egenskab at den kan producere ravsyre med høj koncentration mens den producerer næsten ingen organiske syrer under anaerobe betingelser, hvor
5 vombakteriemutanten er *Mannheimia* sp. LPK ifølge Figur 3.
2. Vombakteriemutant, hvor et lactatdehydrogenase-kodende gen (*ldhA*), et pyruvatformiatlyase-kodende gen (*pfl*), et phosphotransacetylase-kodende gen
10 (*pta*) og et acetatkinase-kodende gen (*ackA*) er blevet afbrudt, og som har den egenskab at den kan producere ravsyre med høj koncentration mens den producerer næsten ingen organiske syrer under anaerobe betingelser, hvor vombakteriemutanten er *Mannheimia* sp. LPK7 ifølge Figur 8.
- 15 3. Vombakteriemutant, hvor et lactatdehydrogenase-kodende gen (*ldhA*), et pyruvatformiatlyase-kodende gen (*pfl*) og et phosphoenolpyruvatcarboxylase-kodende gen (*ppc*) er blevet afbrudt, og som har den egenskab at den kan producere ravsyre med høj koncentration mens den producerer næsten ingen organiske syrer under anaerobe betingelser, hvor vombakteriemutanten er
20 *Mannheimia* sp. LPK4 ifølge Figur 9.
4. Vombakteriemutant ifølge et hvilket som helst af kravene 1-3, hvor vombakterierne er homo-fermentative bakterier.
- 25 5. Vombakteriemutant ifølge krav 1, hvor *Mannheimia* sp. LPK er KCTC 10558BP.
6. Vombakteriemutant ifølge krav 2, hvor *Mannheimia* sp. LPK7 er KCTC 10626BP.
7. Fremgangsmåde til fremstilling af ravsyre, hvilken fremgangsmåde omfatter
30 trinnene: at dyrke vombakteriemutanten ifølge et hvilket som helst af kravene 1-6 under anaerobe betingelser; og at isolere ravsyre fra dyrkningsvæsken.

8. Fremgangsmåde til fremstilling af ravsyre ifølge krav 7, hvor dyrkningstrinnet er homo-fermentering.

DRAWINGS

FIG. 1




FIG. 2




FIG. 3





FIG. 4

M 1 2 3 4 5 6



M 7 8 9 10 11 12



FIG. 5




FIG. 6




FIG. 7




FIG. 8




FIG. 9




FIG. 10




FIG. 11



FIG. 12




FIG. 13

