



US00PP13586P2

(12) **United States Plant Patent**
Dümmen

(10) **Patent No.:** **US PP13,586 P2**

(45) **Date of Patent:** **Feb. 18, 2003**

(54) **NEW GUINEA IMPATIENS PLANT NAMED**
‘DUEPETWHITE’

(75) **Inventor:** **Marga Dümmen**, Rheinberg (DE)

(73) **Assignee:** **Dümmen Jungpflanzen GbR**,
Rheinberg (DE)

(*) **Notice:** Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) **Appl. No.:** **09/996,367**

(22) **Filed:** **Nov. 30, 2001**

(51) **Int. Cl.⁷** **A01H 5/00**

(52) **U.S. Cl.** **Plt./318**

(58) **Field of Search** **Plt./318**

(56) **References Cited**
PUBLICATIONS

UPOV-ROM GTITM Computer Database 2002/03, GTI
Jouve Retrieval Software, Citation for Impatiens ‘Duepet-
white’.*

* cited by examiner

Primary Examiner—Bruce R. Campell

Assistant Examiner—June Hwu

(74) *Attorney, Agent, or Firm*—C. A. Whealy

(57) **ABSTRACT**

A new and distinct cultivar of new Guinea Impatiens plant
named ‘Duepetwhite’, characterized by its upright, rounded
and uniform plant habit; freely branching and freely flow-
ering habit; white-colored flowers that are positioned above
and beyond the leaves; and dark green-colored leaves.

1 Drawing Sheet

1

BOTANICAL CLASSIFICATION/CULTIVAR
DENOMINATION

Impatiens hawkeri cultivar Duepetwhite.

BACKGROUND OF THE INVENTION

The present Invention relates to a new and distinct cul-
tivar of New Guinea Impatiens plant, botanically known as
Impatiens hawkeri, and hereinafter referred to by the name
‘Duepetwhite’.

The new Impatiens is a product of a planned breeding
program conducted by the Inventor in Rheinberg, Germany.
The objective of the breeding program is to develop New
Guinea Impatiens cultivars that have a uniform plant habit
and attractive flower colors.

The new Impatiens originated from a cross made by the
Inventor of a proprietary selection of *Impatiens hawkeri*
identified as code number 94-104-1, not patented, as the
female, or seed parent, with a proprietary selection of
Impatiens hawkeri identified as code number L-94-9, not
patented, as the male, or pollen parent. The cultivar Duepet-
white was discovered and selected by the Inventor as a
flowering plant within the progeny of the stated cross in a
controlled environment in Rheinberg, Germany.

Asexual reproduction of the new cultivar by terminal
cuttings taken in Rheinberg, Germany, has shown that the
unique features of this new Impatiens are stable and repro-
duced true to type in successive generations.

SUMMARY OF THE INVENTION

The following traits have been repeatedly observed and
are determined to be the unique characteristics of ‘Duepet-
white’. These characteristics in combination distinguish
‘Duepetwhite’ as a new and distinct Impatiens cultivar:

1. Upright, rounded and uniform plant habit.
2. Freely branching and freely flowering habit.
3. White-colored flowers that are positioned above and
beyond the leaves.
4. Dark green-colored leaves.

2

Plants of the new Impatiens can be compared to plants of
the female parent, the selection 94-104-1. In side-by-side
comparisons conducted in Rheinberg, Germany, plants of
the new Impatiens were more uniform in plant habit than
plants of the selection 94-104-1.

Plants of the new Impatiens can be compared to plants of
the male parent, the selection L-94-9. In side-by-side com-
parisons conducted in Rheinberg, Germany, plants of the
new Impatiens differed from plants of the selection L-94-9
in the following characteristics:

1. Plants of the new Impatiens were more compact than
plants of the selection L-94-9.
2. Plants of the new Impatiens had larger flowers than
plants of the selection L-94-9.

Plants of the new Impatiens are similar to plants of the
cultivar Moorea, disclosed in U.S. Plant Pat. No. 9,147, in
flower color. However, in side-by-side comparisons con-
ducted in Rheinberg, Germany, plants of the new Impatiens
differed from plants of the cultivar Moorea in the following
characteristics:

1. Plants of the new Impatiens were slightly shorter than
plants of the cultivar Moorea.
2. Plants of the new Impatiens had smaller leaves than
plants of the cultivar Moorea.
3. Plants of the new Impatiens were more freely flowering
than plants of the cultivar Moorea.

BRIEF DESCRIPTION OF THE PHOTOGRAPH

The accompanying colored photograph illustrates the
overall appearance of the new cultivar, showing the colors as
true as it is reasonably possible to obtain in colored repro-
ductions of this type. Colors in the photograph may differ
slightly from the color values cited in the detailed botanical
description which accurately describe the colors of the new
Impatiens. The photograph comprises a side perspective
view of a typical flowering plant of ‘Duepetwhite’ grown in
a container.

DETAILED BOTANICAL DESCRIPTION

The cultivar Duepetwhite has not been observed under all possible environmental conditions. The phenotype may vary somewhat with variations in environment such as temperature and light intensity without, however, any variance in genotype.

The aforementioned photographs, following observations and measurements describe plants grown in Rheinberg, Germany, under commercial practice in a glass-covered greenhouse. Plants were about 16 weeks from cuttings and were grown in 12-cm containers. During the production of the plants, day and night temperatures averaged 18° C. and light levels were about 4,500 lux.

In the following description, color references are made to The Royal Horticultural Society Colour Chart, edition 1995, except where general terms of ordinary dictionary significance are used.

Botanical classification: *Impatiens hawkeri* cultivar Duepetwhite.

Parentage:

Female parent.—Proprietary selection of *Impatiens hawkeri* identified as code number 94-104-1, not patented.

Male parent.—Proprietary selection of *Impatiens hawkeri* identified as code number L-94-9, not patented.

Propagation:

Type cutting.—Terminal cuttings.

Time to initiate roots.—Summer: About 7 days at 22° C.

Winter: About 10 days at 22° C.

Time to produce a rooted cutting.—Summer: About 18 days at 22° C. Winter: About 25 days at 22° C.

Root description.—Fine and freely branching.

Plant description:

General appearance.—Upright, rounded and uniform plant growth habit; freely branching and flowering habit.

Crop time.—From unrooted cuttings, about 16 weeks are required to produce finished flowering plants in 12-cm containers.

Plant height.—About 16 cm.

Plant diameter or spread.—About 41 cm.

Lateral branches.—Quantity per plant: About ten. Length: About 15 cm. Diameter: About 8 mm. Internode length: About 4.25 cm. Color: 144A.

Foliage description.—Arrangement: Opposite or in whorls. Length: About 11 cm. Width: About 3.9 cm. Shape: Elliptic to ovate. Apex: Acuminate to apiculate. Base: Obtuse. Margin: Serrulate with ciliation. Texture: Smooth, glabrous; leathery. Venation pattern: Pinnate. Color: Young foliage, upper surface: 147A. Young foliage, lower surface: 147B. Fully expanded foliage, upper surface: 147A. Fully expanded foliage, lower surface: 147C. Venation,

upper and lower surfaces: 144A. Petiole: Length: About 1.3 cm. Diameter: About 2.5 mm. Color, upper and lower surfaces: 144C.

Flower description:

Flower type and flowering habit.—Single white-colored flowers. Freely and continuously flowering; usually about five to ten flowers and flower buds per lateral branch. Flowers positioned above and beyond the foliage and typically face upward or outward. Petals self-cleaning; gynoecium persistent. Flowers not fragrant.

Flower longevity.—Flowers lasts about one week on the plant.

Flowering season.—Year-round under greenhouse conditions; in the garden, flowering from spring until fall. Plants begin flowering about eight weeks after planting.

Flower buds.—Length: About 1.5 cm. Diameter: About 9 mm. Shape: Ovoid. Color: 145A.

Flower diameter.—About 6.5 cm.

Flower depth.—About 5 mm.

Petals.—Quantity: Five per flower, imbricate. Length: About 3 cm. Width: About 3.5 cm. Shape: Obcordate. Apex: Notched, lobed. Base: Acute. Margin: Entire. Aspect: Mostly flat. Texture: Smooth; satiny. Color: When opening and fully opened, upper surface: 155C; color does not fade with subsequent development. When opening and fully opened, lower surface: 155C.

Spur.—Quantity: One per flower. Length: About 5.1 cm. Diameter: At apex: Less than 1 mm. At flower: About 2.5 mm. Aspect: Curved downward. Color: 154C.

Peduncles.—Length: About 3.8 cm. Diameter: About 1.5 mm. Strength: Moderately strong; flexible. Color: 144C.

Reproductive organs.—Androecium: Stamen quantity/arrangement: Five fused at anthers, hooded; filaments free. Anther length: About 6 mm. Anther shape: Oval. Anther color: 7C. Pollen amount: Abundant. Pollen color: 155B. Gynoecium: Pistil quantity: One per flower. Pistil length: About 6 mm. Stigma color: 145D. Style length: Less than 1 mm. Style color: 150B. Ovary arrangement: Five-celled. Ovary color: 146A.

Seeds/fruits.—Seed and fruit development has not been observed.

Disease/pest resistance: Plants of the new *Impatiens* have not been observed to be resistant to pathogens and pests common to *Impatiens*.

It is claimed:

1. A new and distinct cultivar of New Guinea *Impatiens* plant named 'Duepetwhite', as illustrated and described.

* * * * *

