



Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2681267 C 2013/11/19

(11)(21) **2 681 267**

(12) **BREVET CANADIEN
CANADIAN PATENT**

(13) **C**

(86) Date de dépôt PCT/PCT Filing Date: 2008/04/14
(87) Date publication PCT/PCT Publication Date: 2009/02/19
(45) Date de délivrance/Issue Date: 2013/11/19
(85) Entrée phase nationale/National Entry: 2009/10/08
(86) N° demande PCT/PCT Application No.: US 2008/060186
(87) N° publication PCT/PCT Publication No.: 2009/023299
(30) Priorité/Priority: 2007/04/13 (US60/911,645)

(51) Cl.Int./Int.Cl. *A61K 31/10* (2006.01),
A61P 35/00 (2006.01), *A61P 35/04* (2006.01),
C07C 323/20 (2006.01)

(72) Inventeurs/Inventors:
QU, ZHICAN, US;
ROY, ANSHU MITTAL, US;
ANANTHAN, SUBRAMANIAM, US

(73) Propriétaire/Owner:
SOUTHERN RESEARCH INSTITUTE, US

(74) Agent: MOFFAT & CO.

(54) Titre : AGENTS ANTI-ANGIOGENIQUES ET PROCEDES D'UTILISATION
(54) Title: ANTI-ANGIOGENIC AGENTS AND METHODS OF USE

(57) **Abrégé/Abstract:**

The present disclosure relates generally to treating or preventing diseases associated with angiogenesis by administering to a patient certain compounds found to inhibit or substantially reduce angiogenesis. Compounds employed according to the present disclosure exhibit good anti-angiogenic activity as well as demonstrate a prophylactic effect for preventing and substantially reducing angiogenesis. Examples of such compounds include Ritanserin, Amiodarone, Terfenadine, Perphenazine, Bithionol, and Clomipramine



ABSTRACT

The present disclosure relates generally to treating or preventing diseases associated with angiogenesis by administering to a patient certain compounds found to inhibit or substantially reduce angiogenesis. Compounds employed according to the present disclosure exhibit good anti-angiogenic activity as well as demonstrate a prophylactic effect for preventing and substantially reducing angiogenesis. Examples of such compounds include Ritanserin, Amiodarone, Terfenadine, Perphenazine, Bithionol, and Clomipramine

ANTI-ANGIOGENIC AGENTS AND METHODS OF USE

TECHNICAL FIELD

5 The present disclosure relates generally to treating or preventing diseases associated with angiogenesis by administering to the patient certain compounds found to inhibit or substantially reduce angiogenesis. Compounds employed according to the present disclosure exhibit good anti-angiogenic activity as well as demonstrate a prophylactic effect in preventing and substantially reducing angiogenesis. Examples of such compounds include Ritanserin,
10 Amiodarone, Terfenadine, Perphenazine, Bithionol, and Clomipramine.

BACKGROUND

Angiogenesis is the formation of new blood vessels out of pre-existing capillaries and is a sequence of events that is of key importance in many physiologic and pathologic processes.

15 Normal tissue growth, such as in embryonic development, wound healing, and the menstrual cycle, is characterized by dependence on new vessel formation for the supply of oxygen and nutrients as well as removal of waste products. A large number of different and unrelated diseases are also associated with formation of new vasculature. Among certain pathologies are conditions in which angiogenesis is low, and should be enhanced to improve disease conditions.

20 Pathologies which involve inadequate blood vessel formation include peripheral and coronary ischemia and infarction, chronic wound healing failure, and ulcers. More frequently, however, excessive angiogenesis is an important characteristic of various pathologies including pathologies characterized or associated with an abnormal or uncontrolled proliferation of cells. Pathologies which involve excessive angiogenesis include, for example, cancer (both solid and

25 hematologic tumors), cardiovascular diseases (such as atherosclerosis and restenosis), chronic inflammation (rheumatoid arthritis, Crohn's disease), diabetes (diabetic retinopathy), psoriasis, endometriosis, neovascular glaucoma and adiposity. See Griffioen & Molema, *Angiogenesis: Potentials for Pharmacologic Intervention in the Treatment of Cancer, Cardiovascular Diseases, and Chronic Inflammation*, PHARMACOL. REV. 52, 237-268 (2000).

30 Generally speaking, the angiogenic process entails the proliferation and migration of a normally quiescent endothelium, the controlled proteolysis of the pericellular matrix, and the

synthesis of new extracellular matrix components by developing capillaries. The establishment of new intra- and intercellular contacts and the morphological differentiation of endothelial cells to capillary-like tubular networks provide support for their subsequent maturation, branching, remodeling and selective regression to form a highly organized, functional microvascular network. The autocrine, paracrine and amphicrine interactions of the vascular endothelium with its surrounding stromal components, as well as with the pro-angiogenic and angiostatic cytokines and growth factors orchestrating physiologic angiogenesis, are normally tightly regulated both spatially and temporally. See Gasparini, *The Rationale and Future Potential of Angiogenesis Inhibitors in Neoplasia*, DRUGS, 58(1):17-38 (1999)

The best known anti-angiogenic agents targeting endothelial cell proliferation are Vascular Endothelial Growth Factor ("VEGF") inhibitors. VEGF, a potent angiogenic growth factor, is over expressed in most human solid tumors and in the retina associated eye disorders. The VEGF receptors are mainly enriched in endothelial cells transducing VEGF signaling in many pathological angiogenesis conditions. Growth-stimulated endothelial cells are also sensitive to tyrosine kinase inhibitors targeting VEGF receptors, such as the recent FDA approved anti-cancer drugs Sunitinib (SU11248) and Srafenib (BAY 43-9006). Anti-VEGF and VEGF receptor agents are able to arrest endothelial cell proliferation and prevent new blood vessel growth. In addition to VEGF, many other growth factors such as fibroblast growth factors (FGFs) and platelet derived growth factors (PDGF) also play important roles in endothelial activation. Recently, resistance to anti-angiogenic agents targeting only VEGF signaling is emerging presumably due to other growth factor mediated alternative signaling pathways.

Angiogenesis is crucial to the growth of neoplastic tissues. For more than 100 years, tumors have been observed to be more vascular than normal tissues. Several experimental studies have suggested that both primary tumor growth and metastasis require neovascularization. In contrast to the well orchestrated process described above for normal tissue growth, the pathologic angiogenesis necessary for active tumor growth is generally sustained and persistent, with the initial acquisition of the angiogenic phenotype being a common mechanism for the development of a variety of solid and hematopoietic tumor types. See Folkman, J., *CANCER MEDICINE*, 132-152 (5th Ed., B.C. Decker Inc.) (2000). Tumors that are unable to recruit and sustain a vascular network typically remain dormant as asymptomatic


lesions *in situ*. Metastasis is also angiogenesis-dependent—for a tumor cell to metastasize successfully, it generally must gain access to the vasculature in the primary tumor, survive the circulation, arrest in the microvasculature of the target organ, exit from this vasculature, grow in the target organ, and induce angiogenesis at the target site. Thus, angiogenesis appears to be
 5 necessary at the beginning as well as the completion of the metastatic cascade.

The criticality of angiogenesis to the growth and metastasis of neoplasms thus provides an optimal potential target for chemotherapeutic efforts. Appropriate anti-angiogenic agents may act directly or indirectly to influence tumor-associated angiogenesis either by delaying its onset (i.e., blocking an "angiogenic switch") or by blocking the sustained and focal neovascularization
 10 that is characteristic of many tumor types. Anti-angiogenesis therapies directed against the tumor-associated endothelium and the multiple molecular and cellular processes and targets implicated in sustained pathologic angiogenesis are being actively evaluated for their safety and efficacy in multiple clinical trials. See Deplanque & Harris, Anti-angiogenic Agents: Clinical Trial Design and Therapies in Development, EUR. J. CANCER, 36: 1713-1724 (2000). However,
 15 there has been limited success to date with the discovery and/or identification of safe and/or effective anti-angiogenic agents.

SUMMARY

The present disclosure relates generally to treating or preventing diseases associated with
 20 angiogenesis.

In one embodiment, there are provided methods for inhibiting angiogenesis with a compound of Formula (I)



Formula (I)

wherein:

R is hydrogen, hydroxy or lower alkyloxy;


5 R₁ is a member selected from the group consisting of hydrogen and lower alkyl;

Alk is a lower alkanediyl radical;

X is a member selected from the group consisting of -S-, -CH₂- and -C(R₂)=C(R₃)-, said
R₂ and R₃ being each independently hydrogen or lower alkyl;

A is a bivalent radical having the formula -CH₂CH₂-, -CH₂CH₂CH₂- or

10




wherein R₄ and R₅ are each independently selected from the group consisting of
hydrogen, halo, amino and lower alkyl; and

Ar₁ and Ar₂ are each independently selected from the group consisting of pyridinyl,
thienyl and phenyl, being optionally substituted with halo, hydroxy, lower alkyloxy, lower alkyl
15 and trifluoromethyl; and pharmaceutically acceptable salts, solvates, and prodrugs thereof.
Alternatively, Alk is an 1,2-ethanediyl radical.

In one embodiment, the compound of Formula (I) is Ritanserin.

In another embodiment, there are provided methods for inhibiting angiogenesis with a
compound of Formula (II):

20



Formula (II)

wherein:

R_1 is an alkyl group containing 1 to 6 carbon atoms;

R_2 is selected from the group consisting of hydrogen and methyl;


NR_3 is a radical selected from the group consisting of a dimethylamino, diethylamino, dipropylamino, piperidino, piperazino, pyrrolidino, morpholino, and N-substituted heteroaryl; and

Y and Y_1 are independently selected from the group consisting of hydrogen, fluorine, bromine, chlorine, and iodine; and pharmaceutically acceptable salts, solvates, and prodrugs thereof. Alternatively, Y and Y_1 are identical and are selected from hydrogen, fluorine, bromine, chlorine, and iodine.

In a broad aspect, then, the present invention relates to use of a therapeutically effective amount of bithionol in the manufacture of a medicament for inhibiting the growth or metastasis of a solid tumor in a patient in need thereof.

In one embodiment, the compound of Formula (III) is Terfenadine.

In another embodiment, there are provided methods for inhibiting angiogenesis with a compound of Formula (IV):



Formula (IV)

wherein

X at each occurrence independently represents hydrogen, chlorine or bromine;

A represents OH, OR, NR₁R₂, OC(O)R, OC(O)OR, C(O)OH, C(O)OR, C(O)NR₁R₂, OC(O)N R₁R₂, NHC(O)N R₁R₂ or NHC(NH)N R₁R₂;

R represents alkyl, cycloalkyl, heterocycle, aryl, arylalkyl, or heterocyclalkyl;

R₁ and R₂ each independently represent hydrogen, alkyl, cycloalkyl, heterocycle, aryl, arylalkyl, heterocyclalkyl, or R₁ and R₂ can together form a 3 to 8 membered heterocyclic ring which may be further optionally substituted with one to four (CH₂)_nA substituents;


WO 2009/023299

PCT/US2008/060186

n is a number between 0 and 5 inclusive; and pharmaceutically acceptable salts, solvates, and prodrugs thereof.

In another embodiment, there are provided methods for inhibiting angiogenesis with a compound of Formula (IV-a):

5



Formula (IV-a)

10 wherein

X represents hydrogen, chlorine or bromine;

A represents OH, OR, NR_1R_2 , OC(O)R , OC(O)OR , C(O)OH , C(O)OR , $\text{C(O)NR}_1\text{R}_2$, $\text{OC(O)NR}_1\text{R}_2$, $\text{NHC(O)NR}_1\text{R}_2$ or $\text{NHC(NH)NR}_1\text{R}_2$;

R represents alkyl, cycloalkyl, heterocycle, aryl, arylalkyl, or heterocyclalkyl;

15


R_1 and R_2 each independently represent hydrogen, alkyl, cycloalkyl, heterocycle, aryl, arylalkyl, heterocyclalkyl, or R_1 and R_2 can together form a 3 to 8 membered heterocyclic ring;

n is a number between 0 and 5 inclusive; and pharmaceutically acceptable salts, solvates, and prodrugs thereof.

In one embodiment, the compound of Formula (IV) or Formula (IV-a) is Perphenazine.

20

In another embodiment, there are provided methods for inhibiting angiogenesis with a compound of Formula (V):



Formula (V)


wherein,

Y represents O, S, or S=O;

5 X at each occurrence independently represents halogen or OH; and

n at each occurrence is independently an integer between 0 and 5 inclusive; and
pharmaceutically acceptable salts, solvates, and prodrugs thereof.

Also provided are methods for inhibiting angiogenesis with a compound of Formula (V-a)



Formula (V-a)

wherein


Y represents O, S, or S=O;

X at each occurrence independently represents halogen or OH; and

15 n at each occurrence is independently an integer between 0 and 5 inclusive; and
pharmaceutically acceptable salts, solvates, and prodrugs thereof.

In one embodiment, the compound of Formula (V) or Formula (V-a) is Bithionol.

In another embodiment, there are provided methods for inhibiting angiogenesis with a compound of Formula (VI):



Formula (VI)

wherein

5 X represents a member selected from the group consisting of the ethylene radical -CH₂CH₂- and the vinylene radical -CH=CH-;

R represents a member selected from the group consisting of the methyl radical, the ethyl radical, the propyl radical, chlorine and bromine,

Y represents an alkylene radical with 2-3 carbon atoms, and

10 Am represents a member selected from the group consisting of a lower dialkylamino group, the pyrrolidino, piperidino, piperazino, morpholino and N-methyl-piperidyl(2)-group; and pharmaceutically acceptable salts, solvates, and prodrugs thereof.

In one embodiment, the compound of Formula (VI) is Clomipramine.

15 Also provided are methods of inhibiting the growth or metastasis of an angiogenesis-dependent tumor with compounds of the present disclosure. Another embodiment is drawn to methods for treating a disease or disorder associated with angiogenesis such as neoplastic diseases, restenosis, rheumatoid arthritis, Crohn's disease, diabetic retinopathy, psoriasis, endometriosis, macular degeneration, neovascular glaucoma, and adiposity with compounds of the present disclosure.

20 BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 shows the dose response curve plotted against concentration for cell proliferation assays using Ritanserin.

FIG. 2 shows the results of an endothelial cell migration assay using Ritanserin and plots the percent of migrated cells relative to a control.

FIG. 3. shows the results of an endothelial tube-formation assay using Ritanserin and plots the percentage of tubule formation relative to a control.

FIG. 4 shows the results of an *ex vivo* angiogenesis CAM assay using Ritanserin and plots the vascular density index.

5 FIG. 5. plots the results in microvessel density of an *in vivo* mouse matrigel plug assay using Ritanserin.

FIG. 6 shows the results of a xenograft mouse model assay using Ritanserin and plots the change in tumor volume over time.

10 FIG. 7 shows the dose response curve plotted against concentration for cell proliferation assays using Amiodarone Hydrochloride.

FIG. 8 shows the results of an endothelial cell migration assay using Amiodarone Hydrochloride and plots the percent of migrated cells relative to a control.

FIG. 9 shows the results of an endothelial tube-formation assay using Amiodarone Hydrochloride and plots the percentage of tubule formation relative to a control.

15 FIG. 10 shows the results of an *ex vivo* angiogenesis CAM assay using Amiodarone and plots the vascular density index.

FIG. 11 shows the dose response curve plotted against concentration for cell proliferation assays using Terfenadine.

20 FIG. 12 shows the results of an endothelial cell migration assay using Terfenadine and plots the percent of migrated cells relative to a control.

FIG. 13 shows the results of an endothelial tube-formation assay using Terfenadine and plots the percentage of tubule formation relative to a control.

FIG. 14. shows the results of an *ex vivo* angiogenesis CAM assay using Terfenadine and plots the vascular density index.

25 FIG. 15 shows the dose response curve plotted against concentration for cell proliferation assays using Perphenazine.

FIG. 16 shows the results of an endothelial cell migration assay using Perphenazine and plots the percent of migrated cells relative to a control.

30 FIG. 17 shows the results of an endothelial tube-formation assay using Perphenazine and plots the percentage of tubule formation relative to a control.

FIG. 18 shows the dose response curve plotted against concentration for cell proliferation assays using Bithionol.

FIG. 19 shows the results of an endothelial cell migration assay using Bithionol and plots the percent of migrated cells relative to a control.

5 FIG. 20 shows the results of an endothelial tube-formation assay using Bithionol and plots the percentage of tubule formation relative to a control.

FIG. 21. shows the results of an *ex vivo* angiogenesis CAM assay using Bithionol and plots the vascular density index.

10 FIG. 22 shows the dose response curve plotted against concentration for cell proliferation assays using Clomipramine.

FIG. 23 shows the results of an endothelial cell migration assay using Clomipramine and plots the percent of migrated cells relative to a control.

FIG. 24 shows the results of an endothelial tube-formation assay using Clomipramine and plots the percentage of tubule formation relative to a control.

15 FIG. 25. shows the results of an *ex vivo* angiogenesis CAM assay using Clomipramine and plots the vascular density index.

DETAILED DESCRIPTION OF THE DISCLOSURE

20 The present disclosure relates generally to treating or preventing diseases associated with angiogenesis in a patient in need thereof by administering to the patient certain compounds found to inhibit or substantially reduce angiogenesis.

As employed above and throughout the disclosure, the following terms, unless otherwise indicated, shall be understood to have the following meanings.

25 "Alkyl" refers to an aliphatic hydrocarbon group which may be straight, branched or cyclic having from 1 to about 10 carbon atoms in the chain, and all combinations and subcombinations of ranges therein. "Branched" refers to an alkyl group in which a lower alkyl group, such as methyl, ethyl or propyl, is attached to a linear alkyl chain. In certain embodiments, the alkyl group is a C₁-C₄ alkyl group, i.e., a branched or linear alkyl group having from 1 to about 4 carbons. In other embodiments, the alkyl group is a C₁-C₃ alkyl group, i.e., a

branched or linear alkyl group having from 1 to about 3 carbons. Exemplary alkyl groups include methyl, ethyl, n-propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, hexyl, heptyl, octyl, nonyl and decyl.

5 "Aryl" refers to an aromatic carbocyclic radical containing from about 6 to about 10 carbons, and all combinations and subcombinations of ranges therein. The aryl group may be optionally substituted with one or two or more substituents. Exemplary aryl groups include monocyclic groups, such as phenyl, and bicyclic groups, such as naphthyl.

10 "Heteroaryl" refers to an aromatic carbocyclic radical containing from about 4 to about 10 members, and all combinations and subcombinations of ranges therein, wherein one or more of the members is an element other than carbon, for example, nitrogen, oxygen or sulfur. Exemplary heteroaryl groups include monocyclic groups, such as pyridyl, and bicyclic groups, such as indolyl.

"Arylalkyl" refers to an alkyl substituted with an aryl group. A "C₁-C₄ arylalkyl group", for example, has a C₁-C₄ alkyl group substituted with an aryl group.

15 "Heteroarylalkyl" refers to an alkyl substituted with a heteroaryl group. A "C₁-C₄ heteroarylalkyl group", for example, has a C₁-C₄ alkyl group substituted with an heteroaryl group.

20 The term "cycloalkyl" used alone or as part of a larger moiety shall include cyclic C₃-C₁₀ hydrocarbons which are completely saturated or which contain one or more units of unsaturation, but which are not aromatic. Cycloaliphatic groups are typically C₃-C₁₀, more typically C₃-C₇.

The term "non-aromatic heterocycle", used alone or as part of a larger moiety as in "heterocyclalkyl", refers to non-aromatic ring systems typically having five to fourteen members, preferably five to ten, in which one or more ring carbons, preferably one to four, are each replaced by a heteroatom such as N, O, or S.

25 "Effective amount" refers to an amount of a compound as described herein that may be therapeutically effective to treat a disease or disorder associated with angiogenesis. The precise

amount of these compounds required will vary with the particular compounds or employed, the age and condition of the subject to be treated, and the nature and severity of the condition. However, the effective amount may be determined by one of ordinary skill in the art with only routine experimentation.

5 "Pharmaceutically acceptable" refers to those compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other problem complications commensurate with a reasonable benefit/risk ratio.

"Pharmaceutically acceptable salts" refer to derivatives of the disclosed compounds
 10 wherein the parent compound is modified by making acid or base salts thereof. The compounds of this disclosure form acid and base addition salts with a wide variety of organic and inorganic acids and bases and includes the physiologically acceptable salts which are often used in pharmaceutical chemistry. Such salts are also part of this disclosure. Typical inorganic acids used to form such salts include hydrochloric, hydrobromic, hydroiodic, nitric, sulfuric, phosphoric,
 15 hypophosphoric and the like. Salts derived from organic acids, such as aliphatic mono and dicarboxylic acids, phenyl substituted alkonic acids, hydroxyalkanoic and hydroxyalkandioic acids, aromatic acids, aliphatic and aromatic sulfonic acids, may also be used. Such pharmaceutically acceptable salts thus include acetate, phenylacetate, trifluoroacetate, acrylate, ascorbate, benzoate, chlorobenzoate, dinitrobenzoate, hydroxybenzoate, methoxybenzoate, methylbenzoate, o-
 20 acetoxybenzoate, naphthalene-2-benzoate, bromide, isobutyrate, phenylbutyrate, β -hydroxybutyrate, butyne-1,4-dioate, hexyne-1,4-dioate, cabrate, caprylate, chloride, cinnamate, citrate, formate, fumarate, glycollate, heptanoate, hippurate, lactate, malate, maleate, hydroxymaleate, malonate, mandelate, mesylate, nicotinate, isonicotinate, nitrate, oxalate, phthalate, teraphthalate, phosphate, monohydrogenphosphate, dihydrogenphosphate, metaphosphate, pyrophosphate, propiolate,
 25 propionate, phenylpropionate, salicylate, sebacate, succinate, suberate, sulfate, bisulfate, pyrosulfate, sulfite, bisulfite, sulfonate, benzene-sulfonate, p-bromobenzenesulfonate, chlorobenzenesulfonate, ethanesulfonate, 2-hydroxyethanesulfonate, methanesulfonate, naphthalene-1-sulfonate, naphthalene-2-sulfonate, p-toluenesulfonate, xylenesulfonate, tartarate, and the like.

Bases commonly used for formation of salts include ammonium hydroxide and alkali and alkaline earth metal hydroxides, carbonates, as well as aliphatic and primary, secondary and tertiary amines, aliphatic diamines. Bases especially useful in the preparation of addition salts include sodium hydroxide, potassium hydroxide, ammonium hydroxide, potassium carbonate,
 5 methylamine, diethylamine, and ethylene diamine.

"Patient" refers to animals, including mammals, preferably humans.

"Metabolite" refers to any substance resulting from chemical changes involved in the processes of growth and repair in a living organism, including the anabolic and catabolic processes.

10 A "Prodrug" is a compound that is converted within the body into its active form that has a medical effect. Prodrugs may be useful when the active drug may be too toxic to administer systemically, the active drug is absorbed poorly by the digestive tract, or the body breaks down the active drug before it reaches its target. Methods of making prodrugs are disclosed in Hans Bundgaard, DESIGN OF PRODRUGS (Elsevier Science Publishers B.V. 1985).

15

"Solvates" refers to the compound formed by the interaction of a solvent and a solute and includes hydrates. Solvates are usually crystalline solid adducts containing solvent molecules within the crystal structure, in either stoichiometric or nonstoichiometric proportions.


20 The term "comprising" (and its grammatical variations) as used herein is used in the inclusive sense of "having" or "including" and not in the exclusive sense of "consisting only of." The term "consisting essentially of" as used herein is intended to refer to including that which is explicitly recited along with what does not materially affect the basic and novel characteristics of that recited or specified.

25 The terms "a", "an", and "the" as used herein are understood to encompass the plural as well as the singular unless the context clearly dictates otherwise.

Certain acidic or basic compounds of the present disclosure may exist as zwitterions. All forms of the compounds, including free acid, free base and zwitterions, are contemplated to be within the scope of the present disclosure.

In the formulas described and claimed herein, it is intended that when any symbol appears more than once in a particular formula or substituent, its meaning in each instance is independent of the other.

In one embodiment, there are provided methods for inhibiting angiogenesis with a
5 compound of Formula (I)



Formula (I)

wherein:


10 R is hydrogen, hydroxy or lower alkyloxy;

R₁ is a member selected from the group consisting of hydrogen and lower alkyl;

Alk is a lower alkanediyl radical;

X is a member selected from the group consisting of -S-, -CH₂- and -C(R₂)=C(R₃)-, said
R₂ and R₃ being each independently hydrogen or lower alkyl;

15 A is a bivalent radical having the formula -CH₂CH₂-, -CH₂CH₂CH₂- or




wherein R₄ and R₅ are each independently selected from the group consisting of
hydrogen, halo, amino and lower alkyl; and

Ar₁ and Ar₂ are each independently selected from the group consisting of pyridinyl,
20 thienyl and phenyl, being optionally substituted with halo, hydroxy, lower alkyloxy, lower alkyl
and trifluoromethyl; and pharmaceutically acceptable salts, solvates, and prodrugs thereof.
Alternatively, Alk is an 1,2-ethanediyl radical.

In one embodiment, the compound of Formula (I) is Ritanserin.

Methods of synthesizing compounds of Formula (I) are well known in the art and disclosed in U.S. Patent No. 4, 533, 665 to Kennis *et al.* and U.S. Patent No. 4,485,107 to Kennis *et al.*

In another embodiment, there are provided methods for inhibiting angiogenesis with a
5 compound of Formula (II):



Formula (II)

wherein:

R₁ is an alkyl group containing 1 to 6 carbon atoms;

10 R₂ is selected from the group consisting of hydrogen and methyl;

NR₃ is a radical selected from the group consisting of a dimethylamino, diethylamino, dipropylamino, piperidino, piperazino, pyrrolidino, morpholino, and N-substituted heteroaryl; and


15 Y and Y₁ are independently selected from the group consisting of hydrogen, fluorine, bromine, chlorine, and iodine; and pharmaceutically acceptable salts, solvates, and prodrugs thereof. Alternatively, Y and Y₁ are identical and are selected from hydrogen, fluorine, bromine, chlorine, and iodine.

20 In certain embodiments, NR₃ is selected from the group consisting of dimethylamino, diethylamino, dipropylamino, piperidino, pyrrolidino, and morpholino; Y and Y₁ are the same or different and are selected from the group consisting of hydrogen, iodine, and bromine.

In one embodiment, the compound of Formula (II) is Amiodarone.

Methods of synthesizing compounds of Formula (II) are well known in the art and are disclosed in U.S. Patent No. 3,248,401 to Tondeur *et al.*

In another embodiment, there are provided methods for inhibiting angiogenesis with a
5 compound of Formula (III):



Formula (III)

wherein

R is selected from the group consisting of hydrogen or hydroxy;

10 R₁ is hydrogen; or

R and R₁ taken together form a second bond between the carbon atoms bearing R and R₁;


n is an integer from 1 to 3; and

Z is selected from the group consisting of thienyl, phenyl, or substituted phenyl wherein the substituents on the substituted phenyl may be attached at the ortho, meta, or para positions of
15 the substituted phenyl ring and are selected from the group consisting of a halogen atom, a straight or branched lower alkyl chain of from 1 to 4 carbon atoms, a lower alkoxy group of from 1 to 4 carbon atoms, a di(lower)alkylamino group, or a saturated monocyclic heterocyclic ring selected from the group consisting of pyrrolidino, piperidino, morpholino, or N-(lower)alkylpiperazino; and pharmaceutically acceptable salts, solvates, and prodrugs thereof.

20 In one embodiment, the compound of Formula (III) is Terfenadine.

Methods of synthesizing compounds of Formula (III) are well known in the art and disclosed in U.S. Patent No. 3,878,217 to Carr *et al.* and U.S. Patent No. 4,254,129 to Carr *et al.*

In another embodiment, there are provided methods for inhibiting angiogenesis with a
5 compound of Formula (IV):



10

Formula (IV)

wherein

X at each occurrence independently represents hydrogen, chlorine or bromine;

A represents OH, OR, NR₁R₂, OC(O)R, OC(O)OR, C(O)OH, C(O)OR, C(O)NR₁R₂,
15 OC(O)N R₁R₂, NHC(O)N R₁R₂ or NHC(NH)N R₁R₂;


R represents alkyl, cycloalkyl, heterocycle, aryl, arylalkyl, or heterocyclalkyl;

R₁ and R₂ each independently represent hydrogen, alkyl, cycloalkyl, heterocycle, aryl, arylalkyl, heterocyclalkyl, or R₁ and R₂ can together form a 3 to 8 membered heterocyclic ring which may be further optionally substituted with one to four (CH₂)_nA substituents;

20 n is a number between 0 and 5 inclusive; and pharmaceutically acceptable salts, solvates, and prodrugs thereof.

Methods of synthesizing compounds of Formula (IV) are well known in the art and are disclosed in U.S. Patent No. 2,645,640 to Choisy-le-Roi ,

25 In another embodiment, there are provided methods for inhibiting angiogenesis with a compound of Formula (IV-a):



Formula (IV-a)

wherein

X represents hydrogen, chlorine or bromine;

A represents OH , OR , NR_1R_2 , $OC(O)R$, $OC(O)OR$, $C(O)OH$, $C(O)OR$, $C(O)NR_1R_2$,
 10 $OC(O)N R_1R_2$, $NHC(O)N R_1R_2$ or $NHC(NH)N R_1R_2$;

R represents alkyl, cycloalkyl, heterocycle, aryl, arylalkyl, or heterocyclalkyl;


R_1 and R_2 each independently represent hydrogen, alkyl, cycloalkyl, heterocycle, aryl,
 arylalkyl, heterocyclalkyl, or R_1 and R_2 can together form a 3 to 8 membered heterocyclic ring;

n is a number between 0 and 5 inclusive; and pharmaceutically acceptable salts, solvates,
 15 and prodrugs thereof.

In one embodiment, the compound of Formula (IV) or Formula (IV-a) is Perphenazine.

Methods of synthesizing compounds of Formula (IV-a) are well known in the art and are
 disclosed in U.S. Patent No. 2,860,138 to Sherlock *et al.*

20 In another embodiment, there are provided methods for inhibiting angiogenesis with a
 compound of Formula (V):



Formula (V)

wherein,

Y represents O, S, or S=O;


5 X at each occurrence independently represents halogen or OH; and

n is an integer between 0 and 5 inclusive; and pharmaceutically acceptable salts, solvates, and prodrugs thereof.

In one embodiment, the compound of Formula (V) or Formula (V-a) is Bithionol.

Also provided are methods for inhibiting angiogenesis with a compound of Formula (V-

10 a)



Formula (V-a)

wherein

Y represents O, S, or S=O;


15 X at each occurrence independently represents halogen or OH; and

n at each occurrence is independently an integer between 0 and 5 inclusive; and pharmaceutically acceptable salts, solvates, and prodrugs thereof.

Methods of synthesizing compounds of Formula (V) and Formula (V-a) are well known in the art and disclosed in U.S. Patent No. 3,506,720 to Basel *et al.* and U.S. Patent No.

20 2,849,494 to Cooper *et al.*

In another embodiment, there are provided methods for inhibiting angiogenesis with a compound of Formula (VI):



Formula (VI)

5 wherein

X represents a member selected from the group consisting of the ethylene radical -CH₂CH₂- and the vinylene radical -CH=CH-;

R represents a member selected from the group consisting of the methyl radical, the ethyl radical, the propyl radical, chlorine and bromine,

10 Y represents an alkylene radical with 2-3 carbon atoms, and

Am represents a member selected from the group consisting of a lower dialkylamino group, the pyrrolidino, piperidino, piperazino, morpholino and N-methyl-piperidyl(2)-group; and pharmaceutically acceptable salts, solvates, and prodrugs thereof.

In one embodiment, the compound of Formula (VI) is Clomipramine.

15 Methods of synthesizing compounds of Formula (VI) are well known in the art and disclosed in U.S. Patent No. 3,467,650 to Riechen *et al.*

In another embodiment, there are provided methods for inhibiting angiogenesis with a compound selected from Ritanserin, Amiodarone hydrochloride, Terfenadine, Perphenazine, 20 Bithionol, Bithionol sulfoxide, Clomipramine hydrochloride, Fexofenadine, and combinations thereof.

Also provided are methods of inhibiting the growth or metastasis of an angiogenesis-dependent tumor with compounds of the present disclosure. Another embodiment is drawn to methods for treating a disease or disorder associated with angiogenesis such as neoplastic

diseases, restenosis, rheumatoid arthritis, Crohn's disease, diabetic retinopathy, psoriasis, endometriosis, macular degeneration, neovascular glaucoma, and adiposity with compounds of the present disclosure.

5 In certain embodiments, the present disclosure is directed to methods for inhibiting angiogenesis and/or inhibiting the growth or metastasis of a tumor. As used herein, the term "inhibit" means that the amount of tumor growth or metastasis and/or the occurrence of angiogenesis in patients that have received a compound, as described herein, may be desirably reduced as compared to patients that have not received that compound. Thus, in one form, the inhibitory methods of the present disclosure comprise administering to a patient an effective
10 amount of an anti-angiogenic agent. The term "anti-angiogenic agent", as used herein, refers to compounds which may inhibit angiogenesis.


In other embodiments, the disclosure is directed to methods for treating a disease or disorder associated with angiogenesis. These methods may include the step of identifying a patient having such a disease, including patients who would benefit from the treatment methods
15 described herein. Diseases or disorders associated with angiogenesis include, for example, conditions in which angiogenesis plays a role in the pathology or progression of the condition, such that inhibition of angiogenesis in a patient having such a condition may delay or prevent the further progression of the condition, or lead to remission or regression of the disease state. Such conditions are frequently characterized by or associated with abnormal cellular proliferation and
20 include, for example, neoplastic diseases. As used herein, the term "treating a disease or disorder" refers to the administration of agents intended to limit the extent, progression and/or severity of a condition in a patient, as compared to patients that have not been so treated. As used herein, the term "neoplastic disease" refers to any condition characterized by the presence of an aberrant growth of abnormal cells or tissue, including, but not limited to all cancers and
25 tumors, whether benign or malignant. "Treating neoplastic disease" refers to the administration of a chemotherapeutic agent that will inhibit the further growth or metastasis of any neoplastic tissue that may exist in a patient and/or stimulate regression of such neoplasms, including reducing the size and/or number of such neoplasms and/or inducing the death of neoplastic cells.

Prodrug forms of the compounds bearing various nitrogen functions (amino, hydroxyamino, amide, etc.) may include the following types of derivatives where each R group individually may be hydrogen, substituted or unsubstituted alkyl, aryl, alkenyl, alkynyl, heterocycle, alkylaryl, aralkyl, aralkenyl, aralkynyl, cycloalkyl or cycloalkenyl groups as defined earlier.

- 5 (a) Carboxamides, -NHC(O)R
- (b) Carbamates, -NHC(O)OR
- (c) (Acyloxy)alkyl Carbamates, NHC(O)OROC(O)R
- (d) Enamines, -NHCR(=CHCO₂R) or -NHCR(=CHCONR₂)
- (e) Schiff Bases, -N=CR₂
- 10 (f) Mannich Bases (from carboximide compounds), RCONHCH₂NR₂

Preparations of such prodrug derivatives are discussed in various literature sources (examples are: Alexander *et al.*, J. Med. Chem. 1988, 31, 318; Aligas-Martin *et al.*, PCT WO pp/41531, p.30). The nitrogen function converted in preparing these derivatives is one (or more) of the nitrogen atoms of a compound of the invention.

- 15 Prodrug forms of carboxyl-bearing compounds of the invention include esters (-CO₂R) where the R group corresponds to any alcohol whose release in the body through enzymatic or hydrolytic processes would be at pharmaceutically acceptable levels. Another prodrug derived from a carboxylic acid form of the invention may be a quaternary salt type



- 20 of structure described by Bodor *et al.*, J. Med. Chem. 1980, 23, 469.

It is of course understood that the compounds of the present invention relate to all optical isomers and stereo-isomers at the various possible atoms of the molecule.

- Pharmaceutically acceptable salts of the compounds of the present invention include those derived from pharmaceutically acceptable inorganic or organic acids. Examples of suitable acids
- 25 include hydrochloric, hydrobromic, sulfuric, nitric, perchloric, fumaric, maleic, phosphoric, glycollic, lactic, salicylic, succinic, toluene-p-sulfonic, tartaric, acetic, citric, methanesulfonic, formic, benzoic, malonic, naphthalene-2-sulfonic, trifluoroacetic and benzenesulfonic acids. Salts derived from appropriate bases include alkali such as sodium and ammonia.

- The compounds of the present invention can be synthesized by persons skilled in the art once aware of the present disclosure without undue experimentation. Procedures are available in the chemical literature suitable for preparing the requisite sugars or nucleosides. Along these lines, see Choi, Jong-Ryoo; Kim, Jeong-Min; Roh, Kee-Yoon; Cho, Dong-Gyu; Kim, Jae-Hong;
- 5 Hwang, Jae-Taeg; Cho, Woo-Young; Jang, Hyun-Sook; Lee, Chang-Ho; Choi, Tae-Saeng; Kim, Chung-Mi; Kim, Yong-Zu; Kim, Tae-Kyun; Cho, Seung-Joo; Kim, Gyoung-Won PCT Int. Appl. (2002), 100 pp. WO 0257288 A1 20020725. Holy, Antonin; Votruba, Ivan; Tloustova, Eva; Masojidkova, Milena. Collection of Czechoslovak Chemical Communications (2001), 66(10), 1545-1592. Rejman, Dominik; Masojidkova, Milena; De Clercq, Eric; Rosenberg, Ivan
 - 10 Nucleosides, Nucleotides & Nucleic Acids (2001), 20(8), 1497-1522; Ubasawa, Masaru; Sekiya, Kouichi PCT Int. Appl. (2001), 39 pp WO 0164693 A1 20010907. Otmar, Miroslav; Masojidkova, Milena; Votruba, Ivan; Holy, Antonin. Collection of Czechoslovak Chemical Communications (2001), 66(3), 500-506. Michal; Hocek, Michal; Holy, Antonin. Collection of Czechoslovak Chemical Communications (2000), 65(8), 1357-1373. Jeffery, A.
 - 15 L.; Kim, J.-H.; Wiemer, D. F. Tetrahedron (2000), 56(29), 5077-5083. Holy, Antonin; Guenter, Jaroslav; Dvorakova, Hana; Masojidkova, Milena; Andrei, Graciela; Snoeck, Robert; Balzarini, Jan; De Clercq, Erik. Journal of Medicinal Chemistry (1999), 42(12), 2064-2086. Janeba, Zlatko; Holy, Antonin; Masojidkova, Milena. Collection of Czechoslovak Chemical Communications (2001), 66(9), 1393-1406. Holy, Antonin; Guenter, Jaroslav; Dvorakova,
 - 20 Hana; Masojidkova, Milena; Andrei, Graciela; Snoeck, Robert; Balzarini, Jan; De Clercq, Erik. Journal of Medicinal Chemistry (1999), 42(12), 2064-2086. Dang, Qun; Erion, Mark D.; Reddy, M. Rami; Robinsion, Edward D.; Kasibhatla, Srinivas Rao; Reddy, K. Raja. PCT Int. Appl. (1998), 126 pp WO 9839344 A1 19980911. Arimilli, Murty N.; Cundy, Kenneth C.; Dougherty, Joseph P.; Kim, Choung U.; Oliyai, Reza; Stella, Valentino J. PCT Int. Appl.
 - 25 (1998), 74 pp WO 9804569. Sekiya, Kouichi; Takashima, Hideaki; Ueda, Naoko; Kamiya, Naohiro; Yuasa, Satoshi; Fujimura, Yoshiyuki; Ubasawa, Masaru Journal of Medicinal Chemistry (2002), 45(14), 3138-3142. Ubasawa, Masaru; Sekiya, Kouichi; Takashima, Hideaki; Ueda, Naoko; Yuasa, Satoshi; Kamiya, Naohiro. Eur. Pat. Appl. (1997), 56 pp EP 785208 A1 19970723. Hocek, Michal; Masojidkova, Milena; Holy, Antonin, Collection of
 - 30 Czechoslovak Chemical Communications (1997), 62(1), 136-146. Holy, Antonin; Votruba,

- Ivan; Tloustova, Eva; Masojdkova, Milena. Collection of Czechoslovak Chemical Communications (2001), 66(10), 1545-1592. Holy, Antonin; De Clercq, Erik Desire Alice. PCT Int. Appl. (1996), 57 pp. WO 9633200 A1 19961024. Rejman, Dominik; Rosenberg, Ivan. Collection of Czechoslovak Chemical Communications (1996), 61(Spec. Issue), S122-5 S123. Holy, Antonin; Dvorakova, Hana; Jindrich, Jindrich; Masojdkova, Milena; Budesinsky, Milos; Balzarini, Jan; Andrei, Graciella; De Clercq, Erik. Journal of Medicinal Chemistry (1996), 39(20), 4073-4088. Guanti, Giuseppe; Merlo, Valeria; Narisano, Enrica. Tetrahedron (1995), 51(35), 9737-46. Takashima, Hideaki; Inoue, Naoko; Ubasawa, Masaru; Sekiya, Kouichi; Yabuuchi, Shingo Eur. Pat. Appl. (1995), 88 pp. EP 632048 A1 19950104.
- 10 Alexander, Petr; Holy, Antonin; Masojdkova, Milena, Collection of Czechoslovak Chemical Communications (1994), 59(8), 1853-69. Alexander, Petr; Holy, Antonin; Masojdkova, Milena; Collection of Czechoslovak Chemical Communications (1994), 59(8), 1853-69. Jindrich, Jindrich; Holy, Antonin; Dvorakova, Hana. Collection of Czechoslovak Chemical Communications (1993), 58(7), 1645-67. Holy, Antonin. Collection of Czechoslovak
- 15 Chemical Communications (1993), 58(3), 649-74. Guanti, Giuseppe; Merlo, Valeria; Narisano, Enrica; Tetrahedron (1995), 51(35), 9737-46. Emishetti, Purushotham; Brodfuehrer, Paul R.; Howell, Henry G.; Sapino, Chester, Jr. PCT Int. Appl. (1992), 43 pp. WO 9202511 A1 19920220. Glazier, Arnold. PCT Int. Appl. (1991), 131 pp. WO 9119721. Kim, Choung Un; Luh, Bing Yu; Misco, Peter F.; Bronson, Joanne J.; Hitchcock, Michael J. M.; Ghazzouli, Ismail;
- 20 Martin, John C Journal of Medicinal Chemistry (1990), 33(4), 1207-13. Rosenberg, Ivan; Holy, Antonin; Masojdkova, Milena. Collection of Czechoslovak Chemical Communications (1988), 53(11B), 2753-77. Rosenberg, Ivan; Holy, Antonin; Masojdkova, Milena. Collection of Czechoslovak Chemical Communications (1988), 53(11B), 2753-77.

The following non-limiting examples illustrate and describe aspects of the present

25 disclosure. The examples show and describe only limited embodiments but it is to be understood that the disclosure is capable of use in various other combinations, modifications, and environments and is capable of changes or modifications within the scope of the concept as expressed herein, commensurate with the teachings and/or the skill or knowledge of the relevant art.

Example 1

Cell Proliferation Assay

Human primary cell lines, arterial endothelia, such as HAEC and HPAEC, venous endothelia HUVEC (from Cambrex BioScience Rockland Inc.), and lung fibroblasts LL47 (from the American Type Culture Collection) are cultured by following the instruction and used to evaluate the target compounds for their differential activity against human endothelia verses fibroblasts by CellTiter-Glo Luminescent Cell viability Assay. This assay, generating luminescent signals, is based on quantification of ATP levels in cell cultures. Amount of ATP produced in cell culture reflects the number of viable cells. Hence, this assay is often used to estimate cell proliferation and cytotoxic effects of test compounds. Cells are seeded in 96-well plates with their growth medium, about 5×10^3 cell per well. After 24 hours, various doses of the target compounds are added to the cultures, having four replicates for each dose. After 72 hours of treatment, CellTiter-Glo reagent is added to the cultures following the manufacturer's instruction and luminescence is measured. The control groups are given DMSO vehicle only. IC₅₀s of target compounds for endothelia and fibroblasts proliferation in their growth medium are determined based on the dose response curve and plotted against the range of concentrations.

Example 2

Endothelial Cell Migration Assay

Endothelial cell migration is a key step of the angiogenesis process, which is crucial for on-site recruitment of blood vessel formation. The trans-well filter/inserts chamber of Biocoat Endothelial Cell Migration Angiogenesis System (BD Biosciences) is used for endothelial cell migration assay, which is a 24-Transwell chamber plate containing 3-um pore size inserts coated with human fibronectin. The inserts are incubated at 37°C with 0.1% bovine serum albumin containing endothelial cell basal medium for 1 hour. Endothelial cells (HUVEC) are starved with 0.1% bovine serum albumin in endothelial cell basal medium for 4 to 5 hours before the cell harvest and then seeded (1×10^5 per well) in upper chambers of the Transwell plate with various treatments in 100 ul of 0.1% bovine serum albumin in endothelial cell basal medium. The full growth medium that contains various chemoattractants is added in the lower chambers. The cells are allowed to migrate for 22 ± 1 hours at 37°C. Unmigrated cells at the inside of the inserts are

carefully removed with a Q-tip. Migrated cells at the lower part of the Transwell inserts are then fixed with 4% paraformaldehyde, stained with Hoechst 33342, and photographed under a fluorescent microscope. Three microscopic view fields of each filter/insert are analyzed for the number of migrated cells. Triplicate filter/insert chamber cultures are carried for each test concentration and control. Data are expressed as average number of migrated cells per microscopic view field of 10x magnification and IC₅₀ are calculated based on the dose curve.

10

Example 3

Endothelial Tube Formation Assay

Human endothelial cells are cultured *in vitro* on extracellular matrix, which stimulates the attachment and differentiation of endothelial cells into tubules. The endothelial tube formation assay is based on this phenomenon. Endothelial cells (HUVEC) are seeded (1.5×10^4 per well) in a 96-well plates coated with extracellular matrix and treated with the target compounds at different concentrations with the full growth medium in triplicate. The cells are allowed to form endothelial tubes at 37°C for about 18 hours and photographed under an inverted light microscope. The tubule lengths are quantified using image analysis software, Image-Pro Plus (Media Cybernetics, Inc., Silver Spring, MD). Data are expressed as average tube lengths of three view fields for each well and triplicate wells for each treatment condition. IC₅₀ values are calculated based on the dose curve.

15

20

Example 4

Chick Chorioallantoic Membrane (CAM) Assay

Chicken embryo chorioallantoic membrane provides an ideal *in vivo* model for the physiologic process of angiogenesis. Angiogenic modulators applied on methylcellulose discs placed on top of chicken embryo chorioallantoic membrane are able to alter the development of new blood vasculature. See Staton *et al*, *Current Methods for Assaying Angiogenesis in vitro and in vivo*, INT. J. EXP. PATHOL., 85:233-48 (2004).

The angiogenesis chorioallantoic membrane assay is used for *ex vivo* evaluation of the anti-

25

30

angiogenic potential of synthetic compounds. Fertilized chicken eggs are incubated at 37.5°C in a humidified egg incubator with forced air circulation. On embryo Day 3, eggs are cracked open and embryos are transferred into 100-mm³ Petri plates to continue their development in a cell culture incubator at 37.5°C. Pre-made methylcellulose discs in about 2 mm in diameter are
5 gently implanted on top of embryo chorioallantoic membrane on embryonic Day 5 and then testing compound in solution or control vehicle is applied on top of methylcellulose disc. The embryos are incubated for two more days in the cell culture incubator. The chorioallantoic membranes are examined and quantitatively analyzed for new blood vessel formation at embryonic Day 7. The treatment effects on angiogenesis are evaluated by determining vascular
10 density index (VDI) of each CAM with viable chicken embryos. The VDI represents the number of intersections made by blood vessels with three equidistant concentric circles on the area covered by methylcellulose discs by using *Image Pro Plus* software. The data are expressed as average VDI based on the quantitative analysis for each treatment group ($N \geq 5$) with stander deviation.

15

Example 5

Matrigel Plug Assay in Mouse

The Matrigel Plug Assay is an *in vivo* new blood vessel formation assay, which is widely used for evaluation of anti-angiogenic activities of synthetic compounds and recombinant
20 proteins. Female C57BL/6 mice around 8-10 weeks old and High Concentration Matrigel Matrix is used for the Matrigel Plug experiments. There are four or five mice in each treatment group and two Matrigel plugs in each mouse. Matrigel is mixed with 50ng/ml VEGF, 50ng/ml FGFb, and 3ng/ml heparin as angiogenic stimuli. Target compounds in different dosages are either mixed with Matrigel or given by i.p. or i.v. or by oral. Matrigel at 4°C is subcutaneously
25 injected, 500ul Matrigel for each plug, into each sides of mouse where the hair has been shaved. The injected Matrigel will rapidly form a single solid gel plug. Plugs from each group are collected about two weeks after the Matrigel inoculation. Mice are euthanized by inhalation of CO₂ and mouse-skin is pulled back to expose the plug. The intact plugs are removed and fixed in 10% formalin for histological analysis. Sections (5um thickness) of paraffin-embedded plugs are
30 immunostained with specific antibody against CD31 and counterstained with H&E. CD31-

positive microvessels in an entire cross sectional area of each Matrigel plug are counted. For each group of mice, about six Matrigel plugs are quantitatively analyzed to assess any statistical significant difference of microvessel density between control and compound treated groups.

5

Example 6**Anti-tumor efficacy of Compounds evaluated by Xenograft Nude Mouse Model**

Tumor growth depends on angiogenesis. Inhibition of tumor angiogenesis has become an effective treatment for cancers. Standard Xenograft nude mouse models are used to evaluate potential anti-angiogenic agents for their anti-tumor activity. Human tumor cells or fragments are
 10 implanted into 5 to 7-weeks old nude mice. Target compounds are given to each group of mice by i.p., or i.v., or oral. Tumor size and mouse body weight are monitored twice a week. Average tumor volumes at each time points are expressed with standard deviation against tumor growth time, which is days after xenograft inoculation.

Procedures such as those described in the above examples were employed to give the data
 15 presented in the Figures and in Table I below.

Table I								
Compound	IC₅₀ (μM) Arterial Endothelia Proliferation	IC₅₀ (μM) Venous endothelia Proliferation	IC₅₀ (μM) Fibroblasts Proliferation	Inhibition in Endothelial migration		Inhibition in Endothelia Tube- Formation		Inhibition in CAM Assay (10μg)
				2μM	8μM	2μM	8μM	
Ritanserin	11.0	10.5	>50	55%	65%	11%	33%	21%
Amiodarone hydrochloride	7.5	8.8	20.0	36%	38%	33%	33%	27%
Terfenadine	3.0	5.5	7.0	34%	100%	20%	43%	28%
Clomipramine hydrochloride	10.8	4.5	28.0	58%	63%	12%	26%	19%
Perphenazine	4.2	11.1	10.8	43%	46%	7%	22%	-
Bithionol	2.0	6.0	10.5	5%	100%	19%	38%	30%

Formulations

The compounds of the present disclosure can be administered by any conventional means
5 available for use in conjunction with pharmaceuticals, either as individual therapeutic agents or
in a combination of therapeutic agents. They can be administered alone, but generally
administered with a pharmaceutical carrier selected on the basis of the chosen route of
administration and standard pharmaceutical practice. The compounds can also be administered
in conjunction with other therapeutic agents such as interferon (IFN), interferon α -2a, interferon
10 α -2b, consensus interferon (CIFN), ribavirin, amantadine, remantadine, interleukine-12,
ursodeoxycholic acid (UDCA), and glycyrrhizin.

The pharmaceutically acceptable carriers described herein, for example, vehicles,
adjuvants, excipients, or diluents, are well-known to those who are skilled in the art. Typically,
the pharmaceutically acceptable carrier is chemically inert to the active compounds and has no
15 detrimental side effects or toxicity under the conditions of use. The pharmaceutically acceptable
carriers can include polymers and polymer matrices.

The compounds of this disclosure can be administered by any conventional method
available for use in conjunction with pharmaceuticals, either as individual therapeutic agents or
in a combination of therapeutic agents.

20 The dosage administered will, of course, vary depending upon known factors, such as the
pharmacodynamic characteristics of the particular agent and its mode and route of
administration; the age, health and weight of the recipient; the nature and extent of the
symptoms; the kind of concurrent treatment; the frequency of treatment; and the effect desired.
A daily dosage of active ingredient can be expected to be about 0.001 to 1000 milligrams (mg)
25 per kilogram (kg) of body weight, with the preferred dose being 0.1 to about 60 mg/kg.

Dosage forms (compositions suitable for administration) contain from about 1 mg to
about 500 mg of active ingredient per unit. In these pharmaceutical compositions, the active
ingredient will ordinarily be present in an amount of about 0.5-95% weight based on the total
weight of the composition.

The active ingredient can be administered orally in solid dosage forms, such as capsules, tablets, and powders, or in liquid dosage forms, such as elixirs, syrups and suspensions. It can also be administered parenterally, in sterile liquid dosage forms. The active ingredient can also be administered intranasally (nose drops) or by inhalation of a drug powder mist. Other dosage
5 forms are potentially possible such as administration transdermally, via patch mechanism or ointment.

Formulations suitable for oral administration can consist of (a) liquid solutions, such as an effective amount of the compound dissolved in diluents, such as water, saline, or orange juice; (b) capsules, sachets, tablets, lozenges, and troches, each containing a predetermined amount of
10 the active ingredient, as solids or granules; (c) powders; (d) suspensions in an appropriate liquid; and (e) suitable emulsions. Liquid formulations may include diluents, such as water and alcohols, for example, ethanol, benzyl alcohol, propylene glycol, glycerin, and the polyethylene alcohols, either with or without the addition of a pharmaceutically acceptable surfactant, suspending agent, or emulsifying agent. Capsule forms can be of the ordinary hard- or soft-
15 shelled gelatin type containing, for example, surfactants, lubricants, and inert fillers, such as lactose, sucrose, calcium phosphate, and corn starch. Tablet forms can include one or more of the following: lactose, sucrose, mannitol, corn starch, potato starch, alginic acid, microcrystalline cellulose, acacia, gelatin, guar gum, colloidal silicon dioxide, croscarmellose sodium, talc, magnesium stearate, calcium stearate, zinc stearate, stearic acid, and other excipients, colorants,
20 diluents, buffering agents, disintegrating agents, moistening agents, preservatives, flavoring agents, and pharmacologically compatible carriers. Lozenge forms can comprise the active ingredient in a flavor, usually sucrose and acacia or tragacanth, as well as pastilles comprising the active ingredient in an inert base, such as gelatin and glycerin, or sucrose and acacia, emulsions, and gels containing, in addition to the active ingredient, such carriers as are known in
25 the art.

The compounds of the present disclosure, alone or in combination with other suitable components, can be made into aerosol formulations to be administered via inhalation. These aerosol formulations can be placed into pressurized acceptable propellants, such as dichlorodifluoromethane, propane, and nitrogen. They also may be formulated as
30 pharmaceuticals for non-pressured preparations, such as in a nebulizer or an atomizer.

Formulations suitable for parenteral administration include aqueous and non-aqueous, isotonic sterile injection solutions, which can contain anti-oxidants, buffers, bacteriostats, and solutes that render the formulation isotonic with the blood of the intended recipient, and aqueous and non-aqueous sterile suspensions that can include suspending agents, solubilizers, thickening agents, stabilizers, and preservatives. The compound can be administered in a physiologically acceptable diluent in a pharmaceutical carrier, such as a sterile liquid or mixture of liquids, including water, saline, aqueous dextrose and related sugar solutions, an alcohol, such as ethanol, isopropanol, or hexadecyl alcohol, glycols, such as propylene glycol or polyethylene glycol such as poly(ethyleneglycol) 400, glycerol ketals, such as 2,2-dimethyl-1,3-dioxolane-4-methanol, ethers, an oil, a fatty acid, a fatty acid ester or glyceride, or an acetylated fatty acid glyceride with or without the addition of a pharmaceutically acceptable surfactant, such as a soap or a detergent, suspending agent, such as pectin, carbomers, methylcellulose, hydroxypropylmethylcellulose, or carboxymethylcellulose, or emulsifying agents and other pharmaceutical adjuvants.

Oils, which can be used in parenteral formulations include petroleum, animal, vegetable, or synthetic oils. Specific examples of oils include peanut, soybean, sesame, cottonseed, corn, olive, petrolatum, and mineral. Suitable fatty acids for use in parenteral formulations include oleic acid, stearic acid, and isostearic acid. Ethyl oleate and isopropyl myristate are examples of suitable fatty acid esters. Suitable soaps for use in parenteral formulations include fatty alkali metal, ammonium, and triethanolamine salts, and suitable detergents include (a) cationic detergents such as, for example, dimethyldialkylammonium halides, and alkylpyridinium halides, (b) anionic detergents such as, for example, alkyl, aryl, and olefin sulfonates, alkyl, olefin, ether, and monoglyceride sulfates, and sulfosuccinates, (c) nonionic detergents such as, for example, fatty amine oxides, fatty acid alkanolamides, and polyoxyethylene polypropylene copolymers, (d) amphoteric detergents such as, for example, alkyl β -aminopropionates, and 2-alkylimidazoline quaternary ammonium salts, and (e) mixtures thereof.

The parenteral formulations typically contain from about 0.5% to about 25% by weight of the active ingredient in solution. Suitable preservatives and buffers can be used in such formulations. In order to minimize or eliminate irritation at the site of injection, such compositions may contain one or more nonionic surfactants having a hydrophile-lipophile

balance (HLB) of from about 12 to about 17. The quantity of surfactant in such formulations ranges from about 5% to about 15% by weight. Suitable surfactants include polyethylene sorbitan fatty acid esters, such as sorbitan monooleate and the high molecular weight adducts of ethylene oxide with a hydrophobic base, formed by the condensation of propylene oxide with propylene glycol.

Pharmaceutically acceptable excipients are also well-known to those who are skilled in the art. The choice of excipient will be determined in part by the particular compound, as well as by the particular method used to administer the composition. Accordingly, there is a wide variety of suitable formulations of the pharmaceutical composition of the present disclosure. The following methods and excipients are merely exemplary and are in no way limiting. The pharmaceutically acceptable excipients preferably do not interfere with the action of the active ingredients and do not cause adverse side-effects. Suitable carriers and excipients include solvents such as water, alcohol, and propylene glycol, solid absorbants and diluents, surface active agents, suspending agent, tableting binders, lubricants, flavors, and coloring agents.

The formulations can be presented in unit-dose or multi-dose sealed containers, such as ampules and vials, and can be stored in a freeze-dried (lyophilized) condition requiring only the addition of the sterile liquid excipient, for example, water, for injections, immediately prior to use. Extemporaneous injection solutions and suspensions can be prepared from sterile powders, granules, and tablets. The requirements for effective pharmaceutical carriers for injectable compositions are well known to those of ordinary skill in the art (5, 6). See Banker and Chalmers PHARMACEUTICS AND PHARMACY PRACTICE, 238-250 (J.B. Lippincott Co., Philadelphia, PA Eds. 1982) and Toissel, ASHP HANDBOOK ON INJECTABLE DRUGS, 622-630 (4th ed. 1986).

Formulations suitable for topical administration include lozenges comprising the active ingredient in a flavor, usually sucrose and acacia or tragacanth; pastilles comprising the active ingredient in an inert base, such as gelatin and glycerin, or sucrose and acacia; and mouthwashes comprising the active ingredient in a suitable liquid carrier; as well as creams, emulsions, and gels containing, in addition to the active ingredient, such carriers as are known in the art.

Additionally, formulations suitable for rectal administration may be presented as suppositories by mixing with a variety of bases such as emulsifying bases or water-soluble bases.

Formulations suitable for vaginal administration may be presented as pessaries, tampons, creams, gels, pastes, foams, or spray formulas containing, in addition to the active ingredient, such carriers as are known in the art to be appropriate.

Suitable pharmaceutical carriers are described in Remington's Pharmaceutical Sciences, 5 Mack Publishing Company, a standard reference text in this field.

The dose administered to an animal, particularly a human, in the context of the present disclosure should be sufficient to affect a therapeutic response in the animal over a reasonable time frame. One skilled in the art will recognize that dosage will depend upon a variety of factors including a condition of the animal, the body weight of the animal, as well as the severity 10 and stage of the condition being treated.

A suitable dose is that which will result in a concentration of the active agent in a patient which is known to affect the desired response. The preferred dosage is the amount which results in maximum inhibition of the condition being treated, without unmanageable side effects.

The size of the dose also will be determined by the route, timing and frequency of 15 administration as well as the existence, nature, and extend of any adverse side effects that might accompany the administration of the compound and the desired physiological effect.

Useful pharmaceutical dosage forms for administration of the compounds according to the present disclosure can be illustrated as follows:

Hard Shell Capsules

20 A large number of unit capsules are prepared by filling standard two-piece hard gelatine capsules each with 100 mg of powdered active ingredient, 150 mg of lactose, 50 mg of cellulose and 6 mg of magnesium stearate.

Soft Gelatin Capsules

25 A mixture of active ingredient in a digestible oil such as soybean oil, cottonseed oil or olive oil is prepared and injected by means of a positive displacement pump into molten gelatin to form soft gelatin capsules containing 100 mg of the active ingredient. The capsules are washed and dried. The active ingredient can be dissolved in a mixture of polyethylene glycol, glycerin and sorbitol to prepare a water miscible medicine mix.

Tablets

A large number of tablets are prepared by conventional procedures so that the dosage unit was 100 mg of active ingredient, 0.2 mg. of colloidal silicon dioxide, 5 mg of magnesium stearate, 275 mg of microcrystalline cellulose, 11 mg. of starch, and 98.8 mg of
5 lactose. Appropriate aqueous and non-aqueous coatings may be applied to increase palatability, improve elegance and stability or delay absorption.

Immediate Release Tablets/Capsules

These are solid oral dosage forms made by conventional and novel processes. These
10 units are taken orally without water for immediate dissolution and delivery of the medication. The active ingredient is mixed in a liquid containing ingredient such as sugar, gelatin, pectin and sweeteners. These liquids are solidified into solid tablets or caplets by freeze drying and solid state extraction techniques. The drug compounds may be compressed with viscoelastic and thermoelastic sugars and polymers or effervescent components to produce porous matrices
15 intended for immediate release, without the need of water.

Moreover, the compounds of the present disclosure can be administered in the form of nose drops, or metered dose and a nasal or buccal inhaler. The drug is delivered from a nasal solution as a fine mist or from a powder as an aerosol.

The foregoing description of the disclosure illustrates and describes the present
20 disclosure. Additionally, the disclosure shows and describes only the preferred embodiments but, as mentioned above, it is to be understood that the disclosure is capable of use in various other combinations, modifications, and environments and is capable of changes or modifications within the scope of the concept as expressed herein, commensurate with the above teachings and/or the skill or knowledge of the relevant art.

25 The embodiments described hereinabove are further intended to explain best modes known of practicing it and to enable others skilled in the art to utilize the disclosure in such, or other, embodiments and with the various modifications required by the particular applications or uses. Accordingly, the description is not intended to limit it to the form disclosed herein. Also, it is intended that the appended claims be construed to include alternative embodiments.

CLAIMS

1. Use of a therapeutically effective amount of bithionol in the manufacture of a medicament for inhibiting the growth or metastasis of a solid tumor in a patient in need thereof.
2. The use of claim 2, wherein the medicament is for inhibiting the growth of a solid tumor.
3. The use of claim 2, wherein the medicament is for inhibiting metastasis of a solid tumor.
4. The use according to Claim 1, wherein the tumor is a breast tumor.
5. The use according to Claim 1, wherein the tumor is an ovarian tumor.

Application number / Numéro de demande: 2681267

Figures: 2, 3, 4, 8, 9, 10, 12, 13, 14, 16, 17, 19
20, 21, 23, 24, 25

Pages: _____

Unscannable items received with this application
(Request original documents in File Prep. Section on the 10th floor)

Documents reçu avec cette demande ne pouvant être balayés
(Commander les documents originaux dans la section de la préparation
des dossiers au 10^{ième} étage)

Ritanserin
Cell Proliferation Assays




Figure 1

Ritanserlin
In Vivo Mouse Matrigel Plug Assay




Figure 5

Ritanserin
Xenograft Mouse Model

Animal Group	# of mice	Tumor Line	Treatment (Start on Day 0)	Dose mg/kg (i.p. injection everyday)	Inhibition tumor size
1	10	MX-1	Vehicle	-	-
2	6	MX-1	Ritanserin	1.2 – 2.4 – 4.8	40%
3	10	SKOV-3	Vehicle	-	
4	6	SKOV-3	Ritanserin	1.2 – 2.4 – 4.8	41%




Figure 6

Amiodarone Hydrochloride
Cell Proliferation Assays




Figure 7

Terfenadine
Cell Proliferation Assays

Human Arterial Endothelia (HIAEC)	Human Venous Endothelia (HUVEC)	Human Fibroblasts (LL47)
--------------------------------------	------------------------------------	-----------------------------




Figure 11

Perphenazine
Cell Proliferation Assays




Figure 15

Bithionol
Cell Proliferation Assays




Figure 18

Clomipramine
Cell Proliferation Assays




Figure 22