



HU000032086T2

(19) **HU****MAGYARORSZÁG****Szellemi Tulajdon Nemzeti Hivatala**(11) Lajstromszám: **E 032 086**(13) **T2**

EURÓPAI SZABADALOM SZÖVEGÉNEK FORDÍTÁSA

(21) Magyar ügyszám: **E 12 834840**(22) A bejelentés napja: **2012. 09. 28.**(51) Int. Cl.: **C07D401/04** (2006.01)

(96) Az európai bejelentés bejelentési száma:

EP 20120834840

(86) A nemzetközi (PCT) bejelentési szám:

PCT/JP 12/075084

(97) Az európai bejelentés közzétételi adatai:

EP 2762473 A1 **2013. 04. 04.**

(87) A nemzetközi közzétételi szám:

WO 13047749

(97) Az európai szabadalom megadásának meghirdetési adatai:

EP 2762473 B1 **2016. 08. 31.**

(30) Elsőbbségi adatok:

2011213687 **2011. 09. 29.** **JP**

(73) Jogosult(ak):

Mitsui Chemicals Agro, Inc., Tokyo 103-0027 (JP)

(72) Feltaláló(k):

UMETANI, Hideki, Shiga 520-2362 (JP)
KONDO, Nobuhiro, Chiba 297-0017 (JP)
KAJINO, Fumie, Shiga 520-2362 (JP)

(74) Képviselő:

PINTZ ÉS TÁRSAI Szabadalmi, Védjegy és Jogi Iroda Kft., Budapest(54) **SAIB-tartalmú (szukróz-acetát-izobutirát tartalmú) tartós hatású helyi érzéstelenítő összetétel**

Az európai szabadalom ellen, megadásának az Európai Szabadalmi Közlönyben való meghirdetésétől számított kilenc hónapon belül, felszólalást lehet benyújtani az Európai Szabadalmi Hivatalnál. (Európai Szabadalmi Egyezmény 99. cikk(1))

A fordítást a szabadalmat az 1995. évi XXXIII. törvény 84/H. §-a szerint nyújtotta be. A fordítás tartalmi helyességét a Szellemi Tulajdon Nemzeti Hivatala nem vizsgálta.



(11)

EP 2 762 473 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the grant of the patent:

31.08.2016 Bulletin 2016/35

(21) Application number: **12834840.6**

(22) Date of filing: **28.09.2012**

(51) Int Cl.:

C07D 401/04 ^(2006.01)

(86) International application number:

PCT/JP2012/075084

(87) International publication number:

WO 2013/047749 (04.04.2013 Gazette 2013/14)

(54) **PRODUCTION METHOD FOR 4, 4-DIFLUORO-3,4-DIHYDROISOQUINOLINE DERIVATIVE**

HERSTELLUNGSVERFAHREN FÜR 4, 4-DIFLUOR-3,4-DIHYDROISOCHINOLINDERIVAT

PROCÉDÉ DE FABRICATION DE DÉRIVÉ DE 4,4-DIFLUORO-3,4-DIHYDROISOQUINOLÉINE

(84) Designated Contracting States:

**AL AT BE BG CH CY CZ DE DK EE ES FI FR GB
GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO
PL PT RO RS SE SI SK SM TR**

(30) Priority: **29.09.2011 JP 2011213687**

(43) Date of publication of application:

06.08.2014 Bulletin 2014/32

(73) Proprietor: **Mitsui Chemicals Agro, Inc.**

Tokyo 103-0027 (JP)

(72) Inventors:

- **UMETANI, Hideki**
Shiga 520-2362 (JP)
- **KONDO, Nobuhiro**
Chiba 297-0017 (JP)
- **KAJINO, Fumie**
Shiga 520-2362 (JP)

(74) Representative: **Raynor, Stuart Andrew**

J A Kemp
14 South Square
Gray's Inn
London WC1R 5JJ (GB)

(56) References cited:

WO-A1-2005/070917 WO-A1-2007/011022

- **JIKKEN KAGAKU KOZA 19 YUKIGOSEI-TANKA**
SUIO HALOGEN KAGOBUTSU 1992, pages 378
- 384
- **NOBUO ISHIKAWA ET AL. FUSSO NO**
KAGOBUTSU - SONO KAGAKU TO OYO 1979,
pages 80 - 83, XP008174256
- **HAUFE, G.: 'Triethylamine trishydrofluoride in**
synthesis' JOURNAL FUER PRAKTISCHE
CHEMIE/ CHEMIKER-ZEITUNG vol. 338, no. 2,
1996, pages 99 - 113, XP009013259
- **DAI, X. ET AL.: 'Advances in the application of**
triethylamine tris(hydrofluoride) to organic
synthesis' CHINESE JOURNAL OF ORGANIC
CHEMISTRY 0253-2786 vol. 29, no. 9, 2009, pages
1307 - 16, XP008173621

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 2 762 473 B1

Description


TECHNICAL FIELD

[0001] The present invention relates to a method for producing a 4,4-difluoro-3,4-dihydroisoquinoline derivative.

BACKGROUND ART

[0002] Numerous chemicals have been proposed for the purpose of controlling diseases in agricultural and horticultural crops. For example, Patent Document 1 and Patent Document 2 disclose chemicals containing a 4,4-difluoro-3,4-dihydroisoquinoline derivative represented by general formula (1):


[Chemical Formula 1]



[0003] Wherein R1 and R2 independently represent an optionally substituted alkyl group having 1 to 6 carbon atoms or R1 and R2 together with the carbon atom to which they are bound form an optionally substituted cycloalkyl group having 3 to 10 carbon atoms, X represents a halogen atom, optionally substituted alkyl group having 1 to 6 carbon atoms or optionally substituted alkoxy group having 1 to 6 carbon atoms, n represents an integer of 0 to 4, Y represents a halogen atom, optionally substituted alkyl group having 1 to 6 carbon atoms or optionally substituted alkoxy group having 1 to 6 carbon atoms, and m represents an integer of 0 to 6, and these chemicals are known to be useful as agricultural and horticultural microbicides. Consequently, a method capable of providing a compound represented by general formula (1) on an industrial scale is important.

[0004] However, a specific method for preparing the aforementioned 4,4-difluoro-3,4-dihydroisoquinoline derivative represented by general formula (1) is not described in the aforementioned patent documents. When producing this group of compounds, an efficient method comprises deoxyfluorinating the ketone group of an isoquinolin-4(3H)-one derivative represented by general formula (4) disclosed in Patent Document 1:

[Chemical Formula 2]



wherein R1, R2, X, Y, n and m are the same as previously described. When (diethylamino)sulfur trifluoride as a typical deoxyfluorination reagent (see Non-Patent Document 1) was reacted with a compound represented by general formula (4) as indicated in the comparative examples as described hereinafter, in addition to the reaction progressing slowly, the yield was low at 28.9%. In addition, since (diethylamino)sulfur trifluoride is highly reactive, it has the disadvantage of being difficult to handle during large-scale production.

[0005] With the foregoing background, there has been a fervent desire for the development of a production method that enables 4,4-difluoro-3,4-dihydroisoquinoline derivatives to be synthesized easily and allows them to be produced on an industrial scale.

[0006]

Patent Document 2: International Publication No. WO 2011/77514

Non-Patent Documents

[0007] Non-Patent Document 1: Journal of Organic Chemistry, Vol. 40, pp. 574-578 (1975)

DISCLOSURE OF THE INVENTION

PROBLEMS TO BE SOLVED BY THE INVENTION

[0008] An object of the present invention is to provide a simple and efficient method for producing a 4,4-difluoro-3,4-dihydroisoquinoline derivative.


MEANS FOR SOLVING THE PROBLEMS

[0009] As a result of conducting extensive studies to solve the aforementioned problems, it was found that a target 4,4-difluoro-3,4-dihydroisoquinoline derivative can be produced after reacting a 3,4-dihydroisoquinoline derivative and a brominating agent to convert to a 4,4-dibromo-3,4-dihydroisoquinoline derivative, by reacting the 4,4-dibromo-3,4-dihydroisoquinoline derivative with hydrogen fluoride. This method makes it possible to supply 4,4-difluoro-3,4-dihydroisoquinoline derivatives both easily and efficiently, thereby leading to completion of the present invention.

[0010] Namely, the present invention is:


[1] a method for producing a compound represented by general formula (1):

[Chemical Formula 3]



wherein R1 and R2 independently represent an optionally substituted alkyl group having 1 to 6 carbon atoms or R1 and R2 together with the carbon atom to which they are bound form an optionally substituted cycloalkyl group having 3 to 10 carbon atoms, X represents a halogen atom, optionally substituted alkyl group having 1 to 6 carbon atoms or optionally substituted alkoxy group having 1 to 6 carbon atoms, n represents an integer of 0 to 4, Y represents a halogen atom, optionally substituted alkyl group having 1 to 6 carbon atoms or optionally substituted alkoxy group having 1 to 6 carbon atoms, and m represents an integer of 0 to 6, comprising reacting a compound represented by general formula (2):


[Chemical Formula 4]



wherein R1, R2, X, Y, n and m are the same as previously defined, with hydrogen fluoride;

[2] the method for producing a compound represented by general formula (1) described in [1], wherein the compound represented by general formula (2) is obtained by reacting a compound represented by general formula (3):

[Chemical Formula 5]



wherein R1, R2, X, Y, n and m are the same as in previous [1], with a brominating agent;

[3] the method for producing a compound represented by general formula (1) described in [1], wherein R1 and R2 independently represent an optionally substituted alkyl group having 1 to 6 carbon atoms, n = 0 and m = 0; and,

[4] the method for producing a compound represented by general formula (1) described in [2], wherein R1 and R2 independently represent an optionally substituted alkyl group having 1 to 6 carbon atoms, n = 0 and m = 0.

EFFECTS OF THE INVENTION

[0011] According to the present invention, a method for producing a large amount of 4,4-difluoro-3,4-dihydroisoquinoline derivative can be provided. In addition, the method of the present invention is suitable for an industrial manufacturing method since the target compound can be prepared efficiently by simple operation.

MODE FOR CARRYING OUT THE INVENTION

[0012] The following provides a detailed explanation of embodiments for carrying out the present invention.

[0013] An explanation of general formula (1) is first provided.

[0014] R1 and R2 in general formula (1) are independent and may be the same or different.

[0015] The substituents of the optionally substituted alkyl group having 1 to 6 carbon atoms at R1 and R2 in general formula (1) refer to halogen atoms and alkoxy groups having 1 to 6 carbon atoms. The halogen atom is fluorine, chlorine, bromine or iodine. The alkoxy group having 1 to 6 carbon atoms represents a linear or branched alkoxy group, such as a methoxy group, methoxy group, propoxy group, isopropoxy group, butyloxy group, isobutyloxy group, s-butyloxy group, t-butyloxy group, pentoxy group, isopentoxy group, 2-methylbutyloxy group, neopentoxy group, 1-ethylpropoxy group, hexyloxy group, 4-methylpentoxy group, 3-methylpentoxy group, 2-methylpentoxy group, 1-methylpentoxy group, 3,3-dimethylbutyloxy group, 2,2-dimethylbutyloxy group, 1,1-dimethylbutyloxy group, 1,2-dimethylbutyloxy group, 1,3-dimethylbutyloxy group, 2,3-dimethylbutyloxy group or 2-ethylbutyloxy group. It is preferably an alkoxy group having 1 to 4 carbon atoms and more preferably a methoxy group, ethoxy group, propoxy group or isopropoxy group. There are no particular limitations on the number of substituents and each substituent may be the same or different.

[0016] The alkyl group in the optionally substituted alkyl group having 1 to 6 carbon atoms at R1 and R2 in general formula (1) represents a linear or branched alkyl group, such as a methyl group, ethyl group, propyl group, isopropyl group, butyl group, isobutyl group, s-butyl group, t-butyl group, pentyl group, isopentyl group, 2-methylbutyl group, neopentyl group, 1-ethylpropyl group, hexyl group, 4-methylpentyl group, 3-methylpentyl group, 2-methylpentyl group,

1-methylpentyl group, 3,3-dimethylbutyl group, 2,2-dimethylbutyl group, 1,1-dimethylbutyl group, 1,2-dimethylbutyl group, 1,3-dimethylbutyl group, 2,3-dimethylbutyl group or 2-ethylbutyl group. It is preferably an alkyl group having 1 to 3 carbon atoms and more preferably a methyl group or ethyl group.

[0017] The substituents of the optionally substituted cycloalkyl group having 3 to 10 carbon atoms formed by R1 and R2 together with the carbon atom to which they are bound in general formula (1) have the same meaning as the substituents of the optionally substituted alkyl group having 1 to 6 carbon atoms at R1 and R2 in general formula (1). There are no particular limitations on the number of substituents and each substituent may be the same or different.

[0018] The cycloalkyl group in the optionally substituted cycloalkyl group having 3 to 10 carbon atoms formed by R1 and R2 together with the carbon atom to which they are bound in general formula (1) refers to monocyclic or polycyclic cycloalkyl group having 3 to 10 carbon atoms, such as a cyclobutyl group, cyclopentyl group, cyclohexyl group, cycloheptyl group or norbornyl group. It is preferably a cyclobutyl group, cyclopentyl group, cyclohexyl group or cycloheptyl group, and more preferably a cyclopentyl group.

[0019] The halogen atom at X in general formula (1) refers to fluorine, chlorine, bromine or iodine.

[0020] The optionally substituted alkyl group having 1 to 6 carbon atoms at X in general formula (1) has the same meaning as the optionally substituted alkyl group having 1 to 6 carbon atoms at R1 and R2 in general formula (1).

[0021] The substituents of the optionally substituted alkoxy group having 1 to 6 carbon atoms at X in general formula (1) refer to a halogen atom, that is, fluorine, chlorine, bromine or iodine. There are no particular limitations on the number of substituents and each substituent may be the same or different.

[0022] The alkoxy group of the optionally substituted alkoxy group having 1 to 6 carbon atoms at X in general formula (1) refers to a linear or branched alkoxy group, such as a methoxy group, ethoxy group, propoxy group, isopropoxy group, butyloxy group, isobutyloxy group, s-butyloxy group, t-butyloxy group, pentoxy group, isopentoxy group, 2-methylbutyloxy group, neopentoxy group, 1-ethylpropoxy group, hexyloxy group, 4-methylpentoxy group, 3-methylpentoxy group, 2-methylpentoxy group, 1-methylpentoxy group, 3,3-dimethylbutyloxy group, 2,2-dimethylbutyloxy group, 1,1-dimethylbutyloxy group, 1,2-dimethylbutyloxy group, 1,3-dimethylbutyloxy group, 2,3-dimethylbutyloxy group or 2-ethylbutyloxy group. It is preferably an alkoxy group having 1 to 4 carbon atoms and more preferably a methoxy group, ethoxy group, propoxy group or isopropoxy group.

n in general formula (1) is an integer of 0 to 4.

X may be the same or different when n in general formula (1) is 2 or more.

[0023] The halogen atom at Y in general formula (1) has the same meaning as the halogen atom at X in general formula (1).

[0024] The optionally substituted alkyl group having 1 to 6 carbon atoms at Y in general formula (1) has the same meaning as the optionally substituted alkyl group having 1 to 6 carbon atoms at X in general formula (1).

[0025] The optionally substituted alkoxy group having 1 to 6 carbon atoms at Y in general formula (1) has the same meaning as the optionally substituted alkoxy group having 1 to 6 carbon atoms at X in general formula (1).

m in general formula (1) is an integer of 0 to 6.

Y may be the same or different when m in general formula (1) is 2 or more.

R1, R2, X, Y, n and m in general formula (2) have the same meanings as in general formula (1).

[0026] The following provides an explanation of a method for converting from a compound represented by general formula (2) to a compound represented by general formula (1).

[0027] The hydrogen fluoride used in the reaction may be hydrogen fluoride alone or a reagent that is stabilized by hydrogen bonding, such as triethylamine trihydrofluoride, pyridine hydrofluoride or 1,3-dimethyl-2-imidazolidinone hydrofluoride. There are no particular limitations on the form of the reagent provided the reagent contains hydrogen fluoride and allows the target reaction to proceed.

[0028] Although there are no particular limitations on the amount of hydrogen fluoride used provided it is more than 2 equivalents based on the compound represented by general formula (2), it is preferably 2 equivalents to 20 equivalents from the viewpoint of economy.

[0029] A solvent can be used during the reaction. Although there are no particular limitations on the solvent provided it allows the reaction to proceed, examples of solvents that can be used include benzene-based solvents such as toluene, xylene, benzene, chlorobenzene or dichlorobenzene, nitrile-based solvents such as acetonitrile, ester-based solvents such as ethyl acetate, isopropyl acetate or butyl acetate, amide-based solvents such as N-methylpyrrolidone, N,N-dimethylformamide or N,N-dimethylacetamide, urea-based solvents such as 1,3-dimethyl-2-imidazolidinone, basic solvents such as pyridine, collidine, triethylamine or tributylamine, ether-based solvents such as tetrahydrofuran, diethyl ether, diisopropyl ether or methyl t-butyl ether, chlorine-based solvents such as dichloromethane, dichloroethane, chloroform or carbon tetrachloride, and hydrocarbon-based solvents such as hexane, heptane, cyclohexane or methylcy-

clohexane. In addition, these solvents can be used alone, or two or more types can be mixed at an arbitrary ratio.

[0030] Although there are no particular limitations on the amount of solvent used provided it allows the reaction to proceed, it is preferably 2 times to 30 times the weight of the compound represented by general formula (2) from the viewpoint of economy.

[0031] Although there are no particular limitations on the reaction temperature provided it allows the reaction to proceed, it is higher than 30°C and lower than 120°C or the boiling point of the solvent. The reaction temperature can be suitably set according to the reaction states.

[0032] Method for post-treatment of reaction can consist of mixing the reaction mixture with an aqueous alkaline solution obtained by dissolving potassium hydroxide, sodium hydroxide, sodium carbonate, potassium carbonate, sodium bicarbonate or potassium bicarbonate followed by a liquid separation procedure. At this time, a solvent that is incompatible with water can be added as necessary, examples of which include benzene-based solvents such as toluene, xylene, benzene, chlorobenzene or dichlorobenzene, ester-based solvents such as ethyl acetate, isopropyl acetate or butyl acetate, ether-based solvents such as diethyl ether, diisopropyl ether or methyl t-butyl ether, chlorine-based solvents such as dichloromethane, dichloroethane or chloroform, and hydrocarbon-based solvents such as hexane, heptane, cyclohexane or methylcyclohexane. In addition, these solvents can be used alone, or two or more types can be mixed at an arbitrary ratio. There are no particular limitations on the number of liquid separation procedures, and liquid separation can be carried out corresponding to the target purity and yield.


[0033] Although the moisture in the aforementioned resulting reaction mixture containing compound (1) can be removed with a desiccant such as sodium sulfate or magnesium sulfate, this operation is not essential.

[0034] The aforementioned resulting reaction mixture containing compound (1) can be subjected to distillation under reduced pressure to remove the solvent provided the compound does not decompose.

[0035] The reaction mixture containing compound (1) obtained after distilling off the solvent can be purified by washing, re-precipitating or recrystallizing with a suitable solvent. Examples of solvents used include water, alcohol-based solvents such as methanol, ethanol or isopropyl alcohol, benzene-based solvents such as toluene, xylene, benzene, chlorobenzene or dichlorobenzene, ester-based solvents such as ethyl acetate, isopropyl acetate or butyl acetate, ether-based solvents such as diethyl ether, diisopropyl ether or methyl t-butyl ether, and hydrocarbon-based solvents such as hexane, heptane, cyclohexane or methylcyclohexane. At this time, one type of solvent can be used alone or two or more types mixed at an arbitrary ratio can be used. In addition, the reaction mixture can also be purified by column chromatography. Purification is suitably set according to the target purity.

[0036] Moreover, the reaction mixture containing compound (1) can also be isolated as a salt of a compound represented by general formula (1) such as a compound represented by general formula (4):

[Chemical Formula 6]



wherein R1, R2, X, Y, n and m are the same as in general formula (1), Z represents an acid, and p is from 0.5 to 2.

[0037] The acid at Z in general formula (4) refers to an inorganic acid such as hydrochloric acid, sulfuric acid or phosphoric acid, or an organic acid such as methanesulfonic acid, p-toluenesulfonic acid, oxalic acid or succinic acid.

[0038] The value of p in general formula (4) is 0.5 to 2.

[0039] A compound represented by general formula (4) can be prepared by adding a suitable solvent to the mixture containing a compound represented by general formula (1) followed by adding an acid.

[0040] Examples of the solvents added when preparing a compound represented by general formula (4) include water, alcohol-based solvents such as methanol, ethanol or isopropyl alcohol, benzene-based solvents such as toluene, xylene, benzene, chlorobenzene or dichlorobenzene, ether-based solvents such as tetrahydrofuran, diethyl ether, diisopropyl ether or methyl t-butyl ether, ester-based solvents such as ethyl acetate, isopropyl acetate or butyl acetate, and hydrocarbon-based solvents such as hexane, heptane, cyclohexane or methylcyclohexane. In addition, there are no particular limitations on the form in which solvents are used, and one type of solvent may be used alone or two or more types may be mixed at an arbitrary ratio.

[0041] Although there are no particular limitations on the amount of acid used when preparing a compound represented by general formula (4) provided the amount is more than 1 equivalent, the amount used is 1 equivalent to 15 equivalents from the viewpoint of economy.

[0042] The value of p of the resulting salt is 1 or 2 in the case of a monoacid, and the value of p of the resulting salt is 0.5 or 1 in the case of a diacid. There are no particular limitations on the form of the salt and it may be one salt alone or a mixture of a monoacid salt with a diacid salt.

[0043] A compound represented by general formula (4) can be washed, re-precipitated or recrystallized with a suitable solvent. Examples of solvents used include water, alcohol-based solvents such as methanol, ethanol or isopropyl alcohol, benzene-based solvents such as toluene, xylene, benzene, chlorobenzene or dichlorobenzene, ether-based solvents such as tetrahydrofuran, diethyl ether, diisopropyl ether or methyl t-butyl ether, ester-based solvents such as ethyl acetate, isopropyl acetate or butyl acetate, and hydrocarbon-based solvents such as hexane, heptane, cyclohexane or methylcyclohexane. There are no particular limitations on these solvents provided the target procedure can be carried out, and it may be one type of solvent alone or a mixed solvent of two or more types of solvents.

[0044] A compound represented by general formula (4) can be converted to a compound represented by general formula (1) by a basic substance. The basic substance refers to a substance such as potassium hydroxide, sodium hydroxide, sodium carbonate, potassium carbonate, sodium bicarbonate or potassium bicarbonate, and these can be used while dissolved in water. In addition, extraction can be carried out as necessary with a solvent that is incompatible with water, examples of which include benzene-based solvents such as toluene, xylene, benzene, chlorobenzene or dichlorobenzene, ester-based solvents such as ethyl acetate or butyl acetate, ether-based solvents such as diethyl ether, diisopropyl ether or methyl t-butyl ether, chlorine-based solvents such as dichloromethane, dichloroethane or chloroform, and hydrocarbon-based solvents such as hexane, heptane, cyclohexane or methylcyclohexane. There are no particular limitations on the number of liquid separation procedures, and the number thereof can be set as is suitable. The resulting compound represented by general formula (1) can be purified by washing, re-precipitation, recrystallization or column chromatography and the like using the same procedure as the aforementioned method for post-treatment of the reaction. Method for purification can be suitably set according to the target purity.

[0045] The following provides an explanation of a method for obtaining a compound represented by general formula (2).

[0046] R1, R2, X, Y, n and m in general formula (3) have the same meanings as in general formula (1).

[0047] Examples of brominating agents include 1,3-dibromo-5,5-dimethylhydantoin and N-bromosuccinimide.

[0048] A compound represented by general formula (3) can be prepared with reference to Patent Document 1.

[0049] When converting a compound represented by general formula (3) to a compound represented by general formula (2) with a brominating agent, a radical initiator such as a peracid or azo compound or light irradiation is required.

[0050] Although there are no particular limitations on the radical initiator provided the target bromination is allowed to proceed, a radical initiator having a 10-hour half-life temperature of lower than 90°C is preferable.

[0051] Examples of the peracid as radical initiators include diisobutyl peroxide, cumyl peroxyneodecanoate, di-n-propyl peroxydicarbonate, diisopropyl peroxydicarbonate, di-sec-butyl peroxydicarbonate, 1,1,3,3-tetramethylbutyl peroxyneodecanoate, di(4-t-butylcyclohexyl) peroxydicarbonate, di(2-ethylhexyl) peroxydicarbonate, t-hexyl peroxyneodecanoate, t-butyl peroxyneodecanoate, t-butyl peroxyneohexanoate, t-hexyl peroxyneohexanoate, t-butyl peroxyneopentanoate, t-hexyl peroxyneopentanoate, di(3,5,5-trimethylhexanoyl) peroxide, dilauryl peroxide, 1,1,3,3-tetramethylbutyl peroxy-2-ethylhexanoate, disuccinic acid peroxide, 2,5-dimethyl-2,5-di(2-ethylhexanoylperoxy)hexane, t-hexyl peroxy-2-ethylhexanoate, di(4-methylbenzoyl) peroxide, t-butyl peroxy-2-ethylhexanoate, mixtures of di(3-methylbenzoyl) peroxide, benzoyl(3-methylbenzoyl) peroxide and dibenzoyl peroxide, dibenzoyl peroxide, 1,1-di(t-butylperoxy)-2-methylcyclohexane and 1,1-di(t-hexylperoxy)-3,3,5-trimethylcyclohexane.

[0052] Examples of the azo compound as radical initiators include 2,2'-azobis(isobutyronitrile), 2,2'-azobis(4-methoxy-2,4-dimethylvaleronitrile), 2,2'-azobis(2,4-dimethylvaleronitrile), dimethyl 2,2'-azobis(2-methylpropionate), 2,2'-azobis(2-methylbutyronitrile) and 1,1'-azobis(cyclohexane-1-carbonitrile).

[0053] There are no particular limitations on the amount of radical initiator used provided it allows the target reaction to proceed. The amount used is preferably 0.001 equivalents to 0.30 equivalents from the viewpoint of economy.

[0054] There are no particular limitations on the amount of brominating agent used provided it allows the target reaction to proceed, and it is more than 2 equivalents as bromine atoms. The amount used is preferably 2 equivalents to 4 equivalents as bromine atoms from the viewpoint of economy.

[0055] A solvent can be used when carrying out the reaction. Examples of the solvents include chlorine-based benzene solvents such as chlorobenzene or dichlorobenzene, halogen-based solvents such as carbon tetrachloride, hydrocarbon-based solvents such as hexane, heptane, cyclohexane or methylcyclohexane, and ester-based solvents such as ethyl acetate, isopropyl acetate or butyl acetate.

[0056] Although there are no particular limitations on the amount of solvent used in the reaction provided it allows the reaction to proceed, it is preferably 3 times the weight to 30 times the weight of the compound represented by general formula (3).

[0057] The reaction temperature can be set according to the type of radical initiator, and is higher than 30°C and lower than 150°C or the boiling point of the solvent.

[0058] As for a method for post-treatment of the reaction, by-products can be removed by carrying out a filtration procedure in the case that by-products formed from the brominating agent, such as 5,5-dimethylhydantoin in the case

of 1,3-dibromo-5,5-dimethylhydantoin, have precipitated.

[0059] The reaction mixture of a compound represented by general formula (2) can be washed, re-precipitated or recrystallized with a suitable solvent. Examples of the solvents used at this time include benzene-based solvents such as toluene, xylene, benzene, chlorobenzene or dichlorobenzene, ester-based solvents such as ethyl acetate, isopropyl acetate or butyl acetate, ether-based solvents such as diethyl ether, diisopropyl ether or methyl t-butyl ether, chlorine-based solvents such as dichloromethane, dichloroethane or chloroform, and hydrocarbon-based solvents such as hexane, heptane, cyclohexane or methylcyclohexane. In addition, these solvents can be used alone or as a mixture of two or more types at an arbitrary ratio. In addition, the reaction mixture can also be purified by column chromatography. Purification can be suitably carried out according to the target purity.

[0060] A compound represented by general formula (2) obtained by reacting a compound represented by general formula (3) with a brominating agent can be converted to a compound represented by general formula (1) by reacting with hydrogen fluoride.

[0061] As a result, a 4,4-difluoro-3,4-dihydroisoquinoline derivative can be efficiently produced.

EXAMPLES


[0062] Although the following provides a more detailed description of the present invention by indicating examples thereof, the present invention is not limited to these examples. 3-(3,3-dimethyl-3,4-dihydroisoquinolin-1-yl)quinoline is referred to as Compound (I), 3-(4,4-dibromo-3,3-dimethyl-3,4-dihydroisoquinolin-1-yl) quinoline is referred to as Compound (II), 3-(4,4-difluoro-3,3-dimethyl-3,4-dihydroisoquinolin-1-yl) quinoline is referred to as Compound (III), 1,3-dibromo-5,5-dimethylhydantoin is referred to as DBH, and high-performance liquid chromatography is referred to as

HPLC.

[Comparative Example 1] Synthesis of Compound (III)

[0063] Using 3,3-dimethyl-1-(quinoline-3-yl)isoquinolin-4(3H)-one (referred to as Compound (IV)) as Substrate

[Chemical Formula 7]




[0064] 20 mL of (diethylamino)sulfur trifluoride were added to a mixture of 4.57 g of Compound (IV) and 5 mL of methylene chloride followed by heat refluxing for 13 hours. After cooling in air, the reaction mixture was treated with ice-cooled saturated aqueous sodium bicarbonate followed by extraction with methylene chloride. The resulting methylene chloride layer was washed with saturated brine solution and dried with magnesium sulfate followed by distilling off the solvent under reduced pressure and purifying the resulting residue by chromatography to obtain the target substance (1.42 g, yield: 28.9%). Raw material (2.89 g, recovery rate: 63.2%) was simultaneously recovered.

[Example 1] Synthesis of Compound (II) by DBH

[0065]

[Chemical Formula 8]



[0066] 4.8 g of Compound (I) were dissolved in 48 ml of chlorobenzene followed by raising the temperature to 93°C. 2.64 g of DBH and 0.42 g of 2,2'-azobis(isobutyronitrile) (AIBN) were added and stirred for 5 minutes followed by again adding 2.64 g of DBH and 0.42 g of AIBN and stirring for 2 hours. After cooling to 15°C, the mixture was stirred for 1 hour and then filtered. After distilling the filtrate under reduced pressure to remove the solvent, 5 ml of a mixture of ethyl acetate and hexane (ethyl acetate:hexane = 4:1) were added to the residue followed by stirring at 15°C, further adding 15 ml of hexane and stirring for 1 hour at the same temperature. The precipitate was then filtered out to obtain 6.68 g of Compound (II) as a pale yellow solid. The purity was 94.9%.


Material Data of Compound (II):

[0067] ¹H-NMR (CDCl₃) δ: 9.13 (1H, d, J = 2.0 Hz), 8.38 (1H, d, J = 2.0 Hz), 8.21 (2H, t, J = 8.1 Hz), 7.89 (1H, d, J = 8.3 Hz), 7.82-7.78 (1H, m), 7.62 (2H, td, J = 7.7, 4.1 Hz), 7.45-7.41 (1H, m), 7.24 (1H, d, J = 7.3 Hz), 1.79 (6H, br s).

[Example 2] Synthesis of Compound (III) Using Triethylamine

[0068] Trihydrofluoride

[Chemical Formula 9]



[0069] 5.0 g of Compound (II) obtained in Example 1 and 5.73 g of triethylamine trihydrofluoride were added to 30 ml of xylene and allowed to react for 4 hours at 90°C. Next, 50 g of an 18% aqueous potassium hydroxide solution were dropped in while cooling with ice followed by stirring at room temperature. After separating the liquids of the resulting reaction mixture, the organic layer was concentrated under reduced pressure. 13 ml of methanol were added to the residue, and the resulting solution was added dropwise in 50% aqueous methanol solution. 26 ml of water were additionally added followed by stirring. The resulting precipitate was filtered out to obtain 3.33 g of the title compound as a pale yellow solid. The yield was 88%, thereby demonstrating the present method to be extremely superior to the method of Comparative Example 1. In addition, the ¹H-NMR data of the resulting compound coincided with that described in Patent Document 1.

[Example 3] Synthesis of Compound (III) Using Triethylamine Trihydrofluoride

[0070] 1.47 g of triethylamine trihydrofluoride were added to 7.5 ml of acetonitrile followed by adding 1.21 g of Compound (II) and allowing to react for 4 hours at 90°C. Measurement of the reaction mixture at this time by HPLC indicated that Compound (III) had formed in a reaction yield of 90%. After cooling to room temperature, the reaction mixture was added to an aqueous potassium hydroxide solution. Next, the resulting solution was extracted with ethyl acetate followed by drying with magnesium sulfate. After removing the magnesium sulfate, an aqueous methanol solution was added to the residue followed by stirring and filtering out the precipitate to obtain 0.67 g of Compound (III) as a pale yellow solid. Yield: 80%.

[Example 4] Synthesis of Compound (III) Using Triethylamine Trihydrofluoride

[0071] 0.85 g of triethylamine trihydrofluoride were added to 4 ml of toluene followed by adding 0.70 g of Compound (II) and allowing to react for 4 hours at 90°C. Measurement of the reaction mixture at this time by HPLC indicated that Compound (III) had formed in a reaction yield of 96%. After allowing to cool to room temperature, the reaction mixture was added to a 5% aqueous potassium hydroxide solution. After separating the liquids, the solvent was distilled off under reduced pressure. An aqueous methanol solution was added to the resulting residue and the precipitate was filtered out to obtain 0.43 g of Compound (III) as a pale yellow solid. Yield: 84%.

[Example 5] Synthesis of Compound (III) Using Triethylamine Trihydrofluoride

[0072] 0.80 g of triethylamine trihydrofluoride and 1.0g of Compound (II) were added to 6 ml of heptane and allowed to react for 4 hours at 90°C. Measurement of the resulting reaction mixture by HPLC indicated that Compound (III) had formed in a reaction yield of 93%.

[Example 6] Synthesis of Compound (III) Using Triethylamine Trihydrofluoride

[0073] The reaction was carried out in the same manner as in Example 5 with the exception of using butyl acetate instead of heptane. Measurement of the resulting reaction mixture by HPLC indicated that Compound (III) had formed in a reaction yield of 78%.

[Example 7] Synthesis of Compound (III) Using Triethylamine Trihydrofluoride

[0074] 0.88 g of triethylamine trihydrofluoride were added to 4 ml of triethylamine followed by adding 0.72 g of Compound (II) and allowing to react for 4 hours at 90°C. Measurement of this reaction mixture by HPLC indicated that Compound (III) had formed in a reaction yield of 82%.


[Example 8] Synthesis of Compound (III) Using 70% Pyridine Hydro fluoride

[0075] 0.43 g of 70% pyridine hydrofluoride and 263 mg of pyridine were added to 6 ml of toluene followed by charging with 1.01 g of Compound (II). Next, the reaction mixture was stirred for 4 hours at 85°C. Analysis of the resulting reaction mixture by HPLC indicated that Compound (III) had formed in a reaction yield of 87%.

[Example 9] Synthesis of Compound (II) by N-bromosuccinimide

[0076]

[Chemical Formula 10]




[0077] 10 ml of chlorobenzene added with 1 g of Compound (I) were heated to 93°C. Next, 1.40 g of N-bromosuccinimide and 29 mg of AIBN were added and allowed to react for 2 hours at the same temperature. Measurement of the reaction mixture by HPLC indicated that Compound (II) had formed in a reaction yield of 90%.

[Example 10] Synthesis of Compound (III) from Compound (I)

[0078]

[Chemical Formula 11]



[0079] 26.0 g of DBH and 650.2 mg of di(4-tert-butylcyclohexyl) peroxydicarbonate (purity: 93%) were added to 483.87 g of a chlorobenzene solution containing 21.73 g of Compound (I) followed by heating to 65°C. After stirring for 2.5 hours at 65°C, the reaction mixture was cooled to 45°C and a portion of the chlorobenzene was distilled off under reduced pressure. 213.7 g of the resulting reaction mixture was filtered to obtain 223.4 g of filtrate. The chlorobenzene was further distilled off under reduced pressure to obtain 82.91 g of a chlorobenzene solution of Compound (II) (37.97% by weight, yield: 93.4%).

[0080] 5.10 g of triethylamine trihydrofluoride were added to 82.77 g of the chlorobenzene solution of Compound (II) obtained by the aforementioned reaction followed by heating to 85°C and stirring for 6 hours. After cooling to 60°C, 170.0 g of 20% aqueous potassium hydroxide solution were added followed by cooling to room temperature and stirring for 15 minutes. A liquid separation procedure was then carried out to obtain 90.05 g of an organic layer. As a result of analyzing the organic layer by HPLC, Compound (III) was confirmed to have been formed in a yield of 93.4%. The reaction liquid was concentrated under reduced pressure to obtain 35.21 g of a black solution. 189.11 g of ethanol and 12.94 g of concentrated hydrochloric acid were added to the resulting solution followed by heating to 75°C and stirring for 30 minutes. The solution was then cooled to 2°C and stirred for 3 hours followed by filtering out the precipitate. 21.85 g of the resulting pale yellow solid were a hydrochloride of Compound (III). Purity: 97.4%, Yield: 84%.

Material Data of Hydrochloride of Compound (III):

[0081] ¹H-NMR (DMSO-D₆) δ: 9.32 (1H, d, J = 1.8 Hz), 9.04 (1H, d, J = 1.8 Hz), 8.31 (2H, dd, J = 8.3, 1.8 Hz), 8.06 (1H, dt, J = 10.7, 3.9 Hz), 7.93 (1H, d, J = 7.6 Hz), 7.88-7.82 (2H, m), 7.75 (1H, t, J = 7.5 Hz), 7.57 (1H, d, J = 7.6 Hz), 1.40 (6H, s).

Melting point: 188°C to 191°C


Elementary Analysis: C:66.8%, H:5.0%, N:7.8%, Cl:10%, F:11%

[0082] 105.0 g of methyl t-butyl ether were added to 28.00 g of a 10% aqueous sodium hydroxide solution followed by adding 21.00 g of the aforementioned hydrochloride of Compound (III) while stirring. After stirring for 30 minutes at room temperature, the liquids were separated and the resulting organic layer was washed with 40 g of water. 27.00 g of ethanol were added to the resulting organic layer followed by heating to 59°C and distilling off the methyl t-butyl ether. After cooling the solution to 10°C, 84.0 g of water were added followed by stirring for 1 hour at room temperature. The precipitated solid was filtered and dried to obtain 18.79 g of Compound (III) as a pale yellow solid (purity: 98.1%).

[Example 11] Synthesis of

[0083] 6-bromo-3-(4,4-dibromo-3,3-dimethyl-3,4-dihydroisoquinolin-1-yl)quinoline

[Chemical Formula 12]



[0084] 36.98 g of 6-bromo-3-(3,3-dimethyl-3,4-dihydroisoquinolin-1-yl)quinoline were dissolved in 740 ml of chlorobenzene followed by the addition of 34.74 g of DBH and 4.33 g of di(4-tert-butylcyclohexyl) peroxydicarbonate (purity: 93%)

and heating to 80°C. After stirring for 4 hours at 80°C, the reaction liquid was cooled to 18°C and then filtered. After distilling the filtrate under reduced pressure to remove the solvent, 168 g of chloroform were added to the residue followed by heating to 60°C and stirring for 10 minutes at the same temperature. After cooling to 20°C the reaction mixture was allowed to stand without stirring for 2 hours at the same temperature. The precipitate was then filtered out to obtain 36.03 g of the title compound as a solid. Yield: 68%.


Material Data of Title Compound:

[0085] $^1\text{H-NMR}$ (CDCl_3) δ : 9.13 (1H, d, $J = 2.1$ Hz), 8.27 (1H, d, $J = 2.1$ Hz), 8.22 (1H, dd, $J = 7.8, 1.1$ Hz), 8.05 (2H, dd, $J = 3.1, 1.5$ Hz), 7.85 (1H, dd, $J = 9.2, 2.1$ Hz), 7.64 (1H, td, $J = 7.6, 1.2$ Hz), 7.43 (1H, td, $J = 7.6, 1.2$ Hz), 7.21 (1H, dd, $J = 7.6, 0.9$ Hz), 1.65 (6H, brs).

[Example 12] Synthesis of 6-bromo-3-(4,4-difluoro-3,3-dimethyl-3,4-dihydroisoquinolin-1-yl)quinoline

[0086]

[Chemical Formula 13]



35.93 g of

[0087] 6-bromo-3-(4,4-dibromo-3,3-dimethyl-3,4-dihydroisoquinolin-1-yl)quinoline were dissolved in 216 ml of toluene followed by the addition of 36.54 g of triethylamine trihydrofluoride, heating to 85°C and stirring for 4 hours at the same temperature. After cooling to 30°C, 248.0 g of 20% aqueous potassium hydroxide solution were added followed by stirring for 30 minutes. An organic layer obtained by carrying out a liquid separation procedure was washed with water and the organic layer was dried with sodium sulfate. After filtering out the sodium sulfate, the filtrate was concentrated under reduced pressure to obtain 27.10 g of brown oil. 62.90 g of ethanol were added to the resulting brown oil followed by heating to 70°C and stirring for 10 minutes. After cooling the solution to 2°C and stirring for 2 hours, the precipitate was filtered out. 22.31 g of the resulting white solid were the title compound. Yield: 81%.


Material Data of Title Compound:

[0088] $^1\text{H-NMR}$ (CDCl_3) δ : 9.15 (1H, d, $J = 2.1$ Hz), 8.30 (1H, d, $J = 2.1$ Hz), 8.05-8.04 (2H, m), 7.88 (1H, d, $J = 7.6$ Hz), 7.85 (1H, dd, $J = 9.2, 2.1$ Hz), 7.67 (1H, td, $J = 7.5, 1.0$ Hz), 7.55 (1H, t, $J = 7.6$ Hz), 7.30 (1H, dd, $J = 7.8, 0.8$ Hz), 1.46 (6H, s).

[Example 13] Synthesis of 7-bromo-3-(4,4-difluoro)-3,3-dimethyl-3,4-dihydroisoquinolin-1-yl)quinoline

[0089]

[Chemical Formula 14]



55.7 mg of

[0090] 7-bromo-3-(4,4-dibromo-3,3-dimethyl-3,4-dihydroisoquinolin-1-yl)quinoline, prepared in the same manner as

in Example 11 with the exception of using 7-bromo-3-(3,3-dimethyl-3,4-dihydroisoquinolin-1-yl)quinoline instead of 6-bromo-3-(3,3-dimethyl-3,4-dihydroisoquinolin-1-yl)quinoline, were dissolved in 0.33 ml of toluene followed by the addition of 60 mg of triethylamine trihydrofluoride. The reaction mixture was heated to 95°C and stirred for 4 hours at the same temperature. After cooling to 25°C, 6.0 g of 10% aqueous potassium hydroxide solution were added followed by stirring for 1 hour. After adding 6 ml of toluene, the liquids were separated and the resulting organic layer was washed with 6 g of water followed by drying the organic layer with sodium sulfate. After filtering out the sodium sulfate, the filtrate was concentrated under reduced pressure and the resulting residue was purified by silica gel column chromatography. 26.4 mg of the resulting white solid were the title compound. Yield: 62%.


Material Data of Title Compound:

[0091] $^1\text{H-NMR}$ (CDCl_3) δ : 9.14 (1H, d, $J = 2.1$ Hz), 8.38-8.36 (2H, m), 7.88 (1H, d, $J = 7.6$ Hz), 7.76 (1H, d, $J = 8.6$ Hz), 7.71-7.65 (2H, m), 7.55 (1H, t, $J = 7.6$ Hz), 7.31 (1H, dd, $J = 7.6, 0.6$ Hz), 1.45 (6H, s).

[Example 14] Synthesis of 3-(4,4-dibromo-3,3-dimethyl-3,4-dihydroisoquinolin-1-yl)-7-fluoroquinoline

[0092]

[Chemical Formula 15]



[0093] 103.2 mg of 3-(3,3-dimethyl-3,4-dihydroisoquinolin-1-yl)-7-fluoroquinoline were dissolved in 2 ml of chlorobenzene followed by the addition of 116.3 mg of DBH and 7.3 mg of di(4-tert-butylcyclohexyl) peroxydicarbonate (purity: 93%) and heating to 75°C. After stirring for 3 hours at 75°C, the reaction liquid was cooled to 25°C and filtered. After distilling the filtrate under reduced pressure to remove the solvent, the resulting residue was purified by silica gel column chromatography. 109.6 mg of the resulting solid were the title compound. Yield: 70%.


Material Data of Title Compound:

[0094] $^1\text{H-NMR}$ (CDCl_3) δ : 9.12 (1H, d, $J = 2.1$ Hz), 8.37 (1H, d, $J = 2.1$ Hz), 8.22 (1H, dd, $J = 8.0, 1.2$ Hz), 7.89 (1H, dd, $J = 8.9, 6.1$ Hz), 7.81 (1H, dd, $J = 10.1, 2.4$ Hz), 7.63 (1H, td, $J = 7.6, 1.2$ Hz), 7.45-7.39 (2H, m), 7.23 (1H, dd, $J = 7.6, 1.2$ Hz), 1.68 (6H, br s).

[Example 15] Synthesis of 3-(4,4-difluoro-3,3-dimethyl-3,4-dihydroisoquinolin-1-yl)-7-fluoroquinoline

[0095]

[Chemical Formula 16]



101.0 mg of

[0096] 3-(4,4-dibromo-3,3-dimethyl-3,4-dihydroisoquinolin-1-yl)-7-fluoroquinoline were dissolved in 0.6 ml of toluene followed by the addition of 60 mg of triethylamine trihydrofluoride. The mixture was heated to 90°C and stirred for 4 hours at the same temperature. After cooling to 25°C, 6.0 g of 10% aqueous potassium hydroxide solution were added followed by stirring for 1 hour. After adding 6 ml of toluene, an organic layer obtained by carrying out a liquid separation

procedure was washed with 6 g of water and the organic layer was dried with sodium sulfate. After filtering out the sodium sulfate, the filtrate was concentrated under reduced pressure and the resulting residue was purified by silica gel column chromatography. 57.1 mg of the resulting colorless oil were the title compound. Yield: 77%.


Material Data of Title Compound:

[0097] $^1\text{H-NMR}$ (CDCl_3) δ : 9.15 (1H, d, $J = 2.1$ Hz), 8.40 (1H, d, $J = 2.1$ Hz), 7.91-7.87 (2H, m), 7.81 (1H, dd, $J = 9.8, 2.4$ Hz), 7.67 (1H, t, $J = 7.5$ Hz), 7.55 (1H, t, $J = 7.6$ Hz), 7.41 (1H, td, $J = 8.6, 2.7$ Hz), 7.33 (1H, d, $J = 7.6$ Hz), 1.46 (6H, s).

[Example 16] Synthesis of 3-(4,4-dibromo-3-chloromethyl-3-methyl-3,4-dihydroisoquinolin-1-yl)quinoline

[0098]

[Chemical Formula 17]



651.0 mg of

[0099] 3-(3-chloromethyl-3-methyl-3,4-dihydroisoquinolin-1-yl)quinoline were dissolved in 13.35 g of chlorobenzene followed by the addition of 696.2 mg of DBH and 87.0 mg of di(4-tert-butylcyclohexyl) peroxydicarbonate (purity: 93%) and heating to 65°C. After stirring for 5 hours at 65°C, the reaction liquid was cooled to 25°C and then filtered. After distilling the filtrate to remove the solvent, the residue was purified by silica gel column chromatography. 461.4 mg of the resulting solid were the title compound. Yield: 48%.


Material Data of Title Compound:

[0100] $^1\text{H-NMR}$ (CDCl_3) δ : 9.16 (1H, d, $J = 2.1$ Hz), 8.43 (1H, d, $J = 2.1$ Hz), 8.20 (2H, t, $J = 9.2$ Hz), 7.91 (1H, dd, $J = 8.3, 1.2$ Hz), 7.82 (1H, m), 7.66 (1H, td, $J = 7.6, 1.2$ Hz), 7.62 (1H, m), 7.47 (1H, td, $J = 7.6, 1.2$ Hz), 7.32 (1H, dd, $J = 7.6, 0.9$ Hz), 4.42 (2H, br s), 1.43 (3H, br s).

[Example 17] Synthesis of 3-(3-chloromethyl-4,4-difluoro-3-methyl-3,4-dihydroisoquinolin-1-yl)quinoline

[0101]

[Chemical Formula 18]



461.4 mg of

[0102] 3-(4,4-dibromo-3-chloromethyl-3-methyl-3,4-dihydroisoquinolin-1-yl)quinoline were dissolved in 3 ml of toluene followed by the addition of 520 mg of triethylamine trihydrofluoride. The mixture was then heated to 90°C and stirred for 6 hours at the same temperature. After cooling to 25°C, 7.0 g of 20% aqueous potassium hydroxide solution were added and stirred for 30 minutes. After adding ethyl acetate, an organic layer obtained by carrying out a liquid separation procedure was dried with sodium sulfate. After filtering out the sodium sulfate, the filtrate was concentrated under reduced

pressure and the resulting residue was purified by silica gel column chromatography. 332.2 mg of the resulting solid were the title compound. Yield: 97%.

Material Data of Title Compound:

[0103] $^1\text{H-NMR}$ (CDCl_3) δ : 9.17 (1H, d, $J = 2.1$ Hz), 8.43 (1H, d, $J = 2.1$ Hz), 8.19 (1H, d, $J = 8.6$ Hz), 7.90 (2H, t, $J = 8.6$ Hz), 7.82 (1H, m), 7.69 (1H, td, $J = 7.6, 0.9$ Hz), 7.62 (1H, m), 7.5 (1H, t, $J = 7.6$ Hz), 7.47 (1H, dd, $J = 7.6, 0.9$ Hz), 3.99 (2H, s), 1.48 (3H, s).


INDUSTRIAL APPLICABILITY

[0104] According to the present invention, a 4,4-difluoro-3,4-dihydroisoquinoline derivative can be provided both easily and efficiently. Moreover, the present invention has high value in terms of industrial use since it enables industrial production to be carried out advantageously.

Claims


1. A method for producing a compound represented by general formula (1):

[Chemical Formula 20]



wherein R1 and R2 independently represent an optionally substituted alkyl group having 1 to 6 carbon atoms or R1 and R2 together with the carbon atom to which they are bound form an optionally substituted cycloalkyl group having 3 to 10 carbon atoms, X represents a halogen atom, optionally substituted alkyl group having 1 to 6 carbon atoms or optionally substituted alkoxy group having 1 to 6 carbon atoms, n represents an integer of 0 to 4, Y represents a halogen atom, optionally substituted alkyl group having 1 to 6 carbon atoms or optionally substituted alkoxy group having 1 to 6 carbon atoms, and m represents an integer of 0 to 6, comprising reacting a compound represented by general formula (2):


[Chemical Formula 21]



wherein R1, R2, X, Y, n and m are the same as previously defined, with hydrogen fluoride.

2. The method for producing a compound represented by general formula (1) described in claim 1, wherein the compound represented by general formula (2) is obtained by reacting a compound represented by general formula (3):

[Chemical Formula 22]




10 wherein R1, R2, X, Y, n and m are the same as in claim 1, with a brominating agent.

3. The method for producing a compound represented by general formula (1) described in claim 1, wherein R1 and R2 independently represent an optionally substituted alkyl group having 1 to 6 carbon atoms, n = 0 and m = 0.
- 15 4. The method for producing a compound represented by general formula (1) described in claim 2, wherein R1 and R2 independently represent an optionally substituted alkyl group having 1 to 6 carbon atoms, n = 0 and m = 0.

Patentansprüche


- 20 1. Verfahren zum Herstellen einer Verbindung, die durch die allgemeine Formel (1) dargestellt wird:

[Chemische Formel 20]



35 wobei R1 und R2 unabhängig voneinander eine optional substituierte Alkylgruppe mit 1 bis 6 Kohlenstoffatomen oder R1 und R2 zusammen mit dem Kohlenstoffatom, an das sie gebunden sind, eine optional substituierte Cycloalkylgruppe mit 3 bis 10 Kohlenstoffatomen darstellen, wobei X ein Halogenatom, eine optional substituierte Alkylgruppe mit 1 bis 6 Kohlenstoffatomen oder eine optional substituierte Alkoxygruppe mit 1 bis 6 Kohlenstoffatomen, n eine ganze Zahl von 0 bis 4, Y ein Halogenatom, eine optional substituierte Alkylgruppe mit 1 bis 6 Kohlenstoffatomen oder eine optional substituierte Alkoxygruppe mit 1 bis 6 Kohlenstoffatomen und m eine ganze Zahl von 0 bis 6 darstellt, umfassend das Reagieren einer durch die allgemeine Formel (2) dargestellten Verbindung:

45 [Chemische Formel 21]




55 wobei R1, R2, X, Y, n und m genauso wie oben definiert sind, mit Fluorwasserstoff.

2. Verfahren zum Herstellen einer Verbindung, die durch die in Anspruch 1 beschriebene allgemeine Formel (1) dargestellt wird, wobei die durch die allgemeine Formel (2) dargestellte Verbindung durch das Reagieren einer durch

die allgemeine Formel (3) dargestellten Verbindung erhalten wird:

[Chemische Formel 22]




wobei R₁, R₂, X, Y, n und m genauso wie in Anspruch 1 sind, mit einem Bromierungsmittel.

3. Verfahren zum Herstellen einer Verbindung, die durch die in Anspruch 1 beschriebene allgemeine Formel (1) dargestellt wird, wobei R₁ und R₂ unabhängig voneinander eine optional substituierte Alkylgruppe mit 1 bis 6 Kohlenstoffatomen darstellen, n = 0 und m = 0.
4. Verfahren zum Herstellen einer Verbindung, die durch die in Anspruch 2 beschriebene allgemeine Formel (1) dargestellt wird, wobei R₁ und R₂ unabhängig voneinander eine optional substituierte Alkylgruppe mit 1 bis 6 Kohlenstoffatomen darstellen, n = 0 und m = 0.

Revendications


1. Procédé de fabrication d'un composé représenté par la formule générale (1) :

[Formule chimique 20]



dans laquelle R₁ et R₂ représentent indépendamment un groupe alkyle éventuellement substitué comportant de 1 à 6 atomes de carbone, ou R₁ et R₂ conjointement à l'atome de carbone auquel ils sont liés forment un groupe cycloalkyle éventuellement substitué comportant de 3 à 10 atomes de carbone, X représente un atome de la famille des halogènes, un groupe alkyle éventuellement substitué comportant de 1 à 6 atomes de carbone ou un groupe alcoxy éventuellement substitué comportant de 1 à 6 atomes de carbone, n représente un nombre entier de 0 à 4, Y représente un atome de la famille des halogènes, un groupe alkyle éventuellement substitué comportant de 1 à 6 atomes de carbone ou un groupe alcoxy éventuellement substitué comportant de 1 à 6 atomes de carbone, et m représente un nombre entier de 0 à 6, comprenant une étape consistant à faire réagir un composé représenté par la formule générale (2) :


[Formule chimique 21]



dans laquelle R1, R2, X, Y, n et m sont tels que définis précédemment, avec du fluorure d'hydrogène.

- 15 2. Procédé de fabrication d'un composé représenté par la formule générale (1) décrite dans la revendication 1, dans lequel le composé représenté par la formule générale (2) est obtenu en faisant réagir un composé représenté par la formule générale (3) :

[Formule chimique 22]



dans laquelle R1, R2, X, Y, n et m sont tels que définis dans la revendication 1, avec un agent de bromation.

- 35 3. Procédé de fabrication d'un composé représenté par la formule générale (1) décrite dans la revendication 1, dans laquelle R1 et R2 représentent indépendamment un groupe alkyle éventuellement substitué comportant de 1 à 6 atomes de carbone, n = 0 et m = 0.
- 40 4. Procédé de fabrication d'un composé représenté par la formule générale (1) décrite dans la revendication 2, dans laquelle R1 et R2 représentent indépendamment un groupe alkyle éventuellement substitué comportant de 1 à 6 atomes de carbone, n = 0 et m = 0.

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description


- WO 200570917 A [0006]
- WO 201177514 A [0006]

Non-patent literature cited in the description


- *Journal of Organic Chemistry*, 1975, vol. 40, 574-578
[0007]

MÓDSZER 4,4-DIFLUOR-3,4-DIHDRO-IZOKINOLIN SZÁRMAZÉK GYÁRTÁSÁRA
SZABADALMI IGÉNYPONTOK

1. Módszer az (1)-es általános képlet szerinti vegyület gyártására:
[20. Kémiai képlet]




ahol R1 és R2 jelentése egymástól függetlenül egy opcionálisan szubsztituált, 1-6 szénatommal rendelkező alkilcsoport, vagy R1 és R2 együttesen a szénatommal, amellyek egymáshoz kapcsolódva, egy olyan opcionálisan szubsztituált cikloalkil csoportot alkotnak, amely 3-10 szénatommal rendelkezik, X jelentése halogénatom, opcionálisan szubsztituált, 1-6 szénatommal rendelkező alkilcsoport vagy opcionálisan szubsztituált, 1-6 szénatommal rendelkező alkoxycsoport, n jelentése egész szám 0-tól 4-ig; Y jelentése halogénatom, opcionálisan szubsztituált, 1-6 szénatommal rendelkező alkilcsoport vagy opcionálisan szubsztituált, 1-6 szénatommal rendelkező alkoxycsoport, és m jelentése egész szám 0-tól 6-ig, magában foglalva a (2)-es általános képlet szerinti vegyület reakcióba lépését:



ahol R1, R2, X, Y, n és m az előbb említettek szerint meghatározott, és a reakció hidrogénnel és fluoriddal történik.

2. A módszer az (1)-es általános képlet szerinti vegyület gyártására az 1-es igénypont által leírt módon, ahol a (2)-es általános képlet szerinti vegyületet a (3)-as általános képlet szerinti vegyület reakcióba lépésével nyerik ki:
[22. Kémiai képlet]



ahol R1, R2, X, Y, n és m az 1-es igénypont szerint meghatározott, és a reakció egy brómozószerrel történik.

3. A módszer az (1)-es általános képlet szerinti vegyület gyártására az 1-es igénypont által leírt módon, ahol R1 és R2 jelentése egymástól függetlenül egy opcionálisan szubsztituált, 1-6 szénatommal rendelkező alkilcsoport, továbbá $n=0$ és $m=0$.
4. A módszer az (1)-es általános képlet szerinti vegyület gyártására az 2-es igénypont által leírt módon, ahol R1 és R2 jelentése egymástól függetlenül egy opcionálisan szubsztituált, 1-6 szénatommal rendelkező alkilcsoport, továbbá $n=0$ és $m=0$.



SZTNH-100029657