

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
8 July 2010 (08.07.2010)(10) International Publication Number
WO 2010/077603 A1

(51) International Patent Classification:

A01N 25/08 (2006.01) A61K 31/4192 (2006.01)
C07D 249/04 (2006.01)

(21) International Application Number:

PCT/US2009/066979

(22) International Filing Date:

7 December 2009 (07.12.2009)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/120,661 8 December 2008 (08.12.2008) US
12/426,742 20 April 2009 (20.04.2009) US(71) Applicant (for all designated States except US): **NORTH CAROLINA STATE UNIVERSITY** [US/US]; 1021 Main Campus Drive, Campus Box 8210, Raleigh, NC 27695 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **MELANDER, Christian** [US/US]; 3309 Milton Road, Raleigh, NC 27609 (US). **CAVANAGH, John** [GB/US]; 200 Tattenhall Drive, Cary, NC 27518 (US). **RITCHIE, David, F.** [US/US]; 4804 Marlborough Way, Durham, NC 27713 (US). **ROGERS, Steven, A.** [US/US]; 1900 Trailwood Heights Lane, Apt. 204, Raleigh, NC 27603 (US).**HUIGENS, III, Robert, W.** [US/US]; 245 Old Grove Lane, Apex, NC 27502 (US).(74) Agent: **MYERS BIGEL SIBLEY & SAJOVEC, P.A.**; P.O. Box 37428, Raleigh, NC 27627 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

[Continued on next page]

(54) Title: INHIBITION AND DISPERSION OF BIOFILMS IN PLANTS WITH IMIDAZOLE-TRIAZOLE DERIVATIVES




Figure 1

(57) Abstract: Disclosure is provided for methods of preventing, removing or inhibiting microbial biofilm formation or microbial infection in a plant or plant part thereof, including applying thereto a treatment effective amount of an active compound as described herein, or an agriculturally acceptable salt thereof. Methods of enhancing a microbicide (e.g., including a copper, antibiotic, bacteriophage, etc.) and/or plant defense activator are also provided, including applying an active compound as described herein. Compositions comprising an active compound as described herein in an agriculturally acceptable carrier are also provided, and in some embodiments the compositions further include a microbicide (e.g., including copper, antibiotic, bacteriophage, etc.) and/or plant defense activator.

WO 2010/077603 A1



Published:

— *with international search report (Art. 21(3))*

INHIBITION AND DISPERSION OF BIOFILMS IN PLANTS WITH IMIDAZOLE-TRIAZOLE DERIVATIVES

Christian Melander, John Cavanagh, David F. Ritchie,
Steven A. Rogers and Robert W. Huigens III

Related Applications

This application claims the benefit under 35 U.S.C. § 119(e) of U.S. Application No. 61/120,661, filed December 8, 2008 the disclosure of which is incorporated herein by reference in its entirety. This application is related to U.S. application serial no. 12/426,742, filed April 20, 2009, and published October 22, 2009, as publication no. 2009/0263438, the disclosure of which is incorporated herein by reference in its entirety.

Field of the Invention

This invention relates to compositions and methods useful for controlling biofilms and microorganisms in plants, particularly vascular plants.

Background of the Invention

New approaches are urgently needed to improve agricultural production, given the steadily growing global population that is predicted to reach 6-9 billion persons by mid-century, the continual strain on existing and finite agricultural lands, and the recent diversion of valuable agricultural land from production of crops to production of biomass for fuels. Here we describe new approaches to increase agricultural production by controlling the adverse effects of microorganisms on plants.

The five main crops on which modern societies depend most heavily include corn, cotton, rice, soybeans, and wheat. All of these crops are affected in a deleterious manner by biofilm formation. In addition, other valuable plants such as those producing fruits and vegetables are similarly affected. Plants grown for biomass stand to increase as a valuable crop, albeit not for food, and also can benefit from protection from biofilm formation. Forestry crops and ornamentals also suffer from biofilms.

Summary of the Invention

The present invention is a method of preventing, removing or inhibiting microbial biofilm formation or microbial infection in a plant or plant part thereof, comprising applying to the plant or plant part a treatment effective amount of a compound selected from the group consisting of compounds of **Formula (I)**, **Formula (I)(a)(1)**, **Formula (I)(b)(1)**, **Formula (I)(a)(2)**, **Formula (I)(b)(2)**, **Formula (I)(i)**, **Formula (I)(i)(a)**, **Formula (II)**, **Formula (II)(a)**, **Formula (II)(i)**, **Formula (II)(i)(a)**, **Formula (III)**, **Formula (III)(a)**, **Formula (III)(b)**, **Formula (III)(b)(i)**, **Formula (III)(b)(ii)**, **Formula (IV)**, **Formula (IV)(a)**, **Formula (IV)(i)**,

Formula (IV)(i)(a), Formula (V), Formula (V)(a), Formula (V)(i), Formula (V)(i)(a), Formula (VI), Formula (VI)(a), Formula (VI)(i) and Formula (VI)(i)(a) as described herein, or an agriculturally acceptable salt thereof.

In some embodiments, the plant is a fruit or a vegetable crop plant.

5 In some embodiments, the plant is a citrus tree, and the compound is applied in an amount effective to treat or control a bacterial disease selected from the group consisting of canker, bacterial spot, Black Pit (fruit), Blast, citrus variegated chlorosis, and Citrus Huanglongbing. In some embodiments, the citrus tree is selected from the group consisting of orange, grapefruit, Mandarin, lemon, lime and Kumquat.

10 In some embodiments, the plant is a pome fruit, and the compound is applied in an amount effective to treat or control a bacterial disease selected from the group consisting of Fire Blight, Crown Gall, Blister spot and Hairy root. In some embodiments, the pome fruit is selected from the group consisting of apple, pear, quince, Asian pear, and loquats.

In some embodiments, the plant is a Musa species such as a banana, and the compound is
15 applied in an amount effective to treat or control *Ralstonia solanacearum*.

In some embodiments, the plant is a cole (Brassicaceae) such as cabbage or broccoli, and the compound is applied in an amount effective to treat or control black rot (*Xanthomonas campestris*).

In some embodiments, the plant is a winegrape, and the compound is applied in an
20 amount effective to treat or control for Pierce's disease (*Xylella fastidiosa*) or crown gall (*Agrobacterium vitas*, *A. tumefaciens*).

In some embodiments, the plant is a stone fruit or nut (e.g., peaches, nectarines, plums, almonds, walnuts), and the compound is applied in an amount effective to treat or control bacterial spot and/or blight caused by *Xanthomonas arboricola*; blight caused by *Pseudomonas syringae*; crown gall caused by *Agrobacterium tumefaciens*; phony peach and plum; or almond
25 leaf scorch caused by *Xylella fastidiosa*.

In some embodiments, the plant is a landscape and/or shade tree (e.g., oak, maple, birch, etc.) for bacterial leaf scorch disease (e.g., cause by *Xylella fastidiosa*).

In some embodiments, the plant is a potato, and the compound is applied in an amount
30 effective to treat or control soft rot or black leg (*Erwinia*, *Pectobacterium*).

In some embodiments, the plant is a pepper plant, and the compound is applied in an amount effective to treat or control a bacterial disease selected from the group consisting of Bacterial Spot, Bacterial wilt, Bacterial canker, and Syringae seedling blight and leaf spot.

In some embodiments, the plant is a tomato plant, and the compound is applied in an
35 amount effective to treat or control a bacterial disease selected from the group consisting of:

bacterial canker, bacterial speck, bacterial spot, bacterial stem rot and fruit rot, Bacterial wilt, Pith necrosis, and *Syringae* leaf spot.

In some embodiments, the plant is a soybean plant, and the compound is applied in an amount effective to treat or control a bacterial disease selected from the group consisting of
5 Bacterial blight, Bacterial pustules, Bacterial wilt, Bacterial crinkle leaf, Bacterial tan spot, and Wildfire.

In some embodiments, the plant is corn, and the compound is applied in an amount effective to treat or control a bacterial disease selected from the group consisting of: Bacterial leaf blight, stalk rot, bacterial stripe, chocolate spot, holcus spot all causes by *Pseudomonas*
10 species, Bacterial leaf spot caused by *Xanthomonas* species, Bacterial stalk rot, top rot and Stewart's disease caused by *Erwinia* species, seed rot-seedling blight caused by *Bacillus* species, Purple leaf sheath caused by *Hemiparasitic bacteria*, Corn stunt caused by *Spriroplasma kunkelii*, Goss's bacterial wilt and blight caused by *Clavibacter michiganensis*.

In some embodiments, the plant is cotton, and the compound is applied in an amount
15 effective to treat or control a bacterial disease selected from the group consisting of Bacterial blight caused by *Xanthomonas* species, and Crown gall caused by *Agrobacterium* species and Lint degradation caused by *Erwinia* species.

In some embodiments, the plant is wheat, and the compound is applied in an amount effective to treat or control a bacterial disease selected from the group consisting of Bacterial leaf
20 blight, bacterial sheath rot and Basal glume rot caused by *Pseudomonas* species, Bacterial mosaic and Spike blight caused by *Clavibacter* species, Black chaff caused by *Xanthomonas* species, and Pink seed caused by *Erwinia* (*Pantoea*) species.

In some embodiments, the plant is rice, and the compound is applied in an amount effective to treat or control a bacterial disease selected from the group consisting of bacterial
25 blight and leaf streak caused by *Xanthomonas* species, Foot rot caused by *Erwinia* species, Grain rot caused by *Burkholderia* species, and Sheath brown rot caused by *Pseudomonas* species.

In some embodiments, the plant is pineapple, and the compound is applied in an amount effective to treat or control a bacterial disease selected from the group consisting of bacterial
30 heart rot, fruit collapse, bacterial fruitlet brown rot, marbled fruit, pink fruit and soft rot caused by *Erwinia* species, and Acetic souring caused by Acetic acid bacteria.

In some embodiments, the microbial biofilm formation or microbial infection is caused by a fungi. In some embodiments, the compound is applied to the plant in an amount effective to treat or control a fungal disease selected from the group consisting of rots, leaf molds, blights, wilts, damping-off, spot, root rot, stem rot, mildew, brown spot, gummosis, melanose, post-

bloom fruit drop, scab, alternaria, canker, flyspeck, fruit blotch, dieback, downy mildews, ear rots, anthracnose bunts, smut, rust, eyespot and pecky rice.

In some embodiments, the plant is citrus, and the compound is applied in an amount effective to treat or control a fungal disease selected from the group consisting of: Alternaria brown spot caused by *Alternaria alternaria*, Brown rot caused by *Phytophthora citricola*, Greasy spot and Greasy spot rind blotch caused by *Mycosphaerella citri*, Melanose caused by *Diaporthe citri*, Phytophthora foot rot, gummosis and root rot caused by *Phytophthora citrophthora*, *Phytophthora palmivora*, *Phytophthora syringae* and other *Phytophthora spp.*, Post bloom fruit drop caused by *Colletotrichum acutatum*, and Scab caused by *Elsinoe fawcettii*.

In some embodiments, the plant is Pome fruit, and the compound is applied in an amount effective to treat or control a fungal disease selected from the group consisting of: Apple scab caused by *Venturia inaequalis*, Bitter rot caused by *Colletotrichum gloeosporioides*, Diplodia canker caused by *Diplodia mutila*, Phytophthora crown, collar, root and fruit rot caused by *Phytophthora spp.*, Powdery mildew caused by *Podosphaera leucotricha*, Pacific Coast pear rust, Cedar apple rust, Quince rust caused by *Gymnosporangium spp.*, and Flyspeck caused by *Schizothyrium pomi*.

In some embodiments, the plant is Peppers, and the compound is applied in an amount effective to treat or control a fungal disease selected from the group consisting of: Anthracnose caused by *Colletotrichum spp.*, Damping-off and root rot caused by *Rhizoctonia solani*, *Phytophthora spp.*, *Fusarium spp.*, and *Pythium spp.*, Phytophthora blight caused by *Phytophthora capsici*, and Verticillium wilt caused by *Verticillium albo-atrum*.

In some embodiments, the plant is Tomato, and the compound is applied in an amount effective to treat or control a fungal disease selected from the group consisting of: Alternaria stem canker caused by *Alternaria alternaria*, Anthracnose caused by *Colletotrichum spp.*, Fusarium crown, root rot and wilt caused by *Fusarium oxysporum*, Gray mold caused by *Botrytis cinerea*, Late blight caused by *phytophthora infestans*, Pythium damping-off and fruit rot caused by *Pythium spp.*, Rhizoctonia damping-off and fruit rot caused by *Rhizoctonia solani*, Septoria leaf spot caused by *Septoria lycopersici*, Verticillium wilt caused by *Verticillium albo-atrum*, and White mold caused by *Sclerotinia sclerotiorum*.

In some embodiments, the plant is Soybean, and the compound is applied in an amount effective to treat or control a fungal disease selected from the group consisting of: Phytophthora root and stem rot caused by *Phytophthora sojae*, Pythium root rot, damping-off and seed decay caused by *Pythium spp.*, Brown stem rot caused by *Phialophora gregata*, Rhizoctonia root and stem rot caused by *Rhizoctonia solani*, Stem canker, pod and stem blight caused by *Diaporthe phaseolorum*, Phomopsis seed decay caused by *Phomopsis longicollis*, Charcoal rot caused by

Macrophomina phaseolina, Sclerotinia stem rot caused by *Sclerotinia sclerotiorum*, Sudden death syndrome caused by *Fusarium solani*, and Soybean Rust caused by *Phakopsora pachyrhizi*.

In some embodiments, the plant is Grape, and the compound is applied in an amount effective to treat or control a fungal disease selected from the group consisting of: Alternaria rot caused by *Alternaria alternaria*, Angular leaf spot caused by *Mycosphaerella angulata*, Botrytis bunch rot and blight caused by *Botrytis cinerea*, Diplodia cane dieback and bunch rot caused by *Diplodia natalensis*, Downy mildew caused by *Plasmopara viticola*, Phytophthora crown and root rot caused by *Phytophthora* spp., Powdery mildew caused by *Uncinula necator*, Ripe rot caused by *Glomerella cingulata*, Septoria leaf spot caused by *Septoria ampelopsidis*, and Verticillium wilt caused by *Verticillium dahliae*.

In some embodiments, the plant is Potato, and the compound is applied in an amount effective to treat or control a fungal disease selected from the group consisting of: Brown spot, Black pit and Early blight caused by *Alternaria* spp., Fusarium dry rot and wilt caused by *Fusarium* spp., Gangrene caused by *Phoma* spp., Late blight and Pink rot caused by *Phytophthora* spp., Rhizoctonia canker and black scurf caused by *Rhizoctonia solani*, Rosellinia black rot caused by *Rosellinia* spp., Septoria leaf spot caused by *Septoria lycopersici*, Stem rot caused by *Sclerotium rolfsii*, Verticillium wilt caused by *Verticillium albo-atrum*, and White mold caused by *Sclerotinia sclerotiorum*.

In some embodiments, the plant is Pineapple, and the compound is applied in an amount effective to treat or control a fungal disease selected from the group consisting of: Anthracnose caused by *Colletotrichum ananas*, Butt rot and White leaf spot caused by *Chalara paradoxa*, Leaf spot caused by *Curvularia eragrostidis*, Phytophthora heart rot caused by *Phytophthora cinnamomi* and *Phytophthora parasitica*, Root rot and Seedling blight caused by *Pythium* spp., and Leaking brown ring caused by *Toffliadis dimenationa*.

In some embodiments, the plant is Cotton, and the compound is applied in an amount effective to treat or control a fungal disease selected from the group consisting of: Anthracnose caused by *Glomerella gossypii*, Boll rot caused by *Colletotrichum gossypii*, *Fusarium* spp., *Phytophthora* spp., or *Rhizoctonia solani*, Fusarium wilt caused by *Fusarium oxysporum*, Leaf spot caused by *Alternaria* spp., *Cercospora gossypina*, *Rhizoctonia solani*, and *Stemphylium solani*, Lint contamination caused by *Aspergillus flavus*, Powdery mildew caused by *Leveillula taurica*, Cotton rust caused by *Puccinia schedonnardii*, Southwestern cotton rust caused by *Puccinia cacabata*, Tropical cotton rust caused by *Phakopsora gossypii*, Southern blight caused by *Sclerotium rolfsii*, Seedling disease complex caused by *Colletotrichum gossypii*, *Fusarium*

spp., *Pythium* spp., *Rhizoctonia solani*, or *Thielaviopsis basicola*, Stem canker caused by *Phoma exigua*, and Verticillium wilt caused by *Verticillium dahliae*.


In some embodiments, the plant is Corn, and the compound is applied in an amount effective to treat or control a fungal disease selected from the group consisting of: Anthracnose caused by *Colletotrichum graminicola*, Aspergillus ear and kernel rot caused by *Aspergillus flavus*, Banded leaf, sheath spot, root rot and stalk rot caused by *Rhizoctonia solani*, Brown spot, Black spot and Stalk rot caused by *Physoderma maydis*, Curvularia leaf spot caused by *Curvularia clavata*, Diplodia ear rot, stalk rot, seed rot and seedling blight caused by *Diplodia* spp., Downey mildews caused by *Sclerophthora* spp. or *Peronosclerospora* spp., Ear rots caused by *Alternaria alternaria*, Ergot caused by *Claviceps gigantea*, Fusarium ear, stalk, kernel, root, seed rot, seedling blight caused by *Fusarium* spp., Cercospora leaf spot caused by *Cercospora zeae-maydis*, Helminthosporium ear rot caused by *Helminthosporium carbonum*, Pythium root rot and stalk rot caused by *Pythium* spp., Rhizoctonia ear rot caused by *Rhizoctonia zeae*, Common corn rust and Southern corn rust caused by *Puccinia* spp., Southern blight caused by *Athelia rolfsii*, Common smut caused by *Ustilago zeae*, Southern corn leaf blight and stalk rot caused by *Cochliobolus heterostrophus*, and storage rots caused by *Aspergillus* spp. and *Penicillium* spp.

In some embodiments, the plant is Rice, and the compound is applied in an amount effective to treat or control a fungal disease selected from the group consisting of: Black kernel caused by *Curvularia lunata*, Blast caused by *Pyricularia oryzae*, Brown spot caused by *Cochliobolus miyabeanus*, Downy mildew caused by *Sclerophthora macrospora*, False smut caused by *Ustilaginoidea virens*, Narrow brown leaf spot caused by *Cercospora janseana*, Pecky rice caused by *Fusarium* spp., *Microdochium oryzae*, or *Sarocladium oryzae*, Root rot caused by *Fusarium* spp. or *Pythium* spp., Seedling blight caused by fungi (e.g., *Cochliobolus miyabeanus*, *Curvularia* spp., *Fusarium* spp., *Rhizoctonia solani*, *Sclerotium rolfsii* and *Athelia rolfsii*), Stackburn caused by *Alternaria padwickii*, Stem rot caused by *Magnaporthe salvinii*, Water-mold (seed-rot and seedling disease) caused by *Achlya* spp., *Fusarium* spp., or *Pythium* spp.

In some embodiments, the plant is Wheat, and the compound is applied in an amount effective to treat or control a fungal disease selected from the group consisting of: Alternaria leaf blight caused by *Alternaria trititica*, Anthracnose caused by *Colletotrichum graminicola*, Black head molds caused by *Cladosporium* spp., *Epicoccum* spp., *Sporobolomyces* spp. or *Stemphylium* spp., Common bunt caused by *Tilletia* spp., Crown rot, seedling blight and dryland root rot caused by *Fusarium* spp. or *Gibberella* spp., Downey mildew caused by *Sclerophthora macrospora*, Dwarf bunt caused by *Tilletia controversa*, Ergot caused by *Claviceps purpurea*, Eyespot caused by *Tapesia yallundae*, Leaf rust caused by *Puccinia trititica*, Loose smut caused

by *Ustilago tritici*, Microscopia leaf spot caused by *Phaeosphaeria microscopia*, Phoma spot caused by *Phoma* spp., Powdery mildew caused by *Erysiphe graminis*, Pythium root rot, Snow rot caused by *Pythium* spp., Rhizoctonia root rot caused by *Rhizoctonia solani*, Scab (head blight) caused by *Fusarium* spp. or *Gibberella* spp., Southern blight caused by *Sclerotium rolfsii*,
 5 Speckled snow mold caused by *Typhula* spp., Stem rust caused by *Puccinia graminis*, storage molds caused by *Aspergillus* spp. or *Penicillium* spp., Take-all caused by *Gaeumannomyces graminis*, and Zoosporic root rot caused by *Lagenia radicola*.


A further aspect of the present invention is an agricultural composition comprising: (a) an agriculturally acceptable carrier (e.g., an aqueous carrier or a solid particulate carrier); and (b)
 10 an antimicrobial or biofilm preventing, removing or inhibiting compound selected from the group consisting of compounds of **Formula (I)**, **Formula (I)(a)(1)**, **Formula (I)(b)(1)**, **Formula (I)(a)(2)**, **Formula (I)(b)(2)**, **Formula (I)(i)**, **Formula (I)(i)(a)**, **Formula (II)**, **Formula (II)(a)**, **Formula (II)(i)**, **Formula (II)(i)(a)**, **Formula (III)**, **Formula (III)(a)**, **Formula (III)(b)**, **Formula (III)(b)(i)**, **Formula (III)(b)(ii)**, **Formula (IV)**, **Formula (IV)(a)**, **Formula (IV)(i)**,
 15 **Formula (IV)(i)(a)**, **Formula (V)**, **Formula (V)(a)**, **Formula (V)(i)**, **Formula (V)(i)(a)**, **Formula (VI)**, **Formula (VI)(a)**, **Formula (VI)(i)** and **Formula (VI)(i)(a)** as described herein, or an agriculturally acceptable salt thereof. In some embodiments, the composition further includes a microbicide. In some embodiments, the microbicide comprises copper (e.g., copper hydroxide). In some embodiments, the microbicide comprises an antibiotic or a bacteriophage. In
 20 some embodiments, the composition further includes a plant defense activator. In some embodiments, the composition further includes both a plant defense activator and a microbicide. In some embodiments, the compound is a compound of **Formula (II)(a)(5)(D)**:



or an agriculturally acceptable salt thereof.


Further provided are methods of enhancing the effects of a microbicide comprising
 25 applying an active compound selected from the group consisting of compounds of **Formula (I)**, **Formula (I)(a)(1)**, **Formula (I)(b)(1)**, **Formula (I)(a)(2)**, **Formula (I)(b)(2)**, **Formula (I)(i)**, **Formula (I)(i)(a)**, **Formula (II)**, **Formula (II)(a)**, **Formula (II)(i)**, **Formula (II)(i)(a)**, **Formula (III)**, **Formula (III)(a)**, **Formula (III)(b)**, **Formula (III)(b)(i)**, **Formula (III)(b)(ii)**,
 30 **Formula (IV)**, **Formula (IV)(a)**, **Formula (IV)(i)**, **Formula (IV)(i)(a)**, **Formula (V)**, **Formula (V)(a)**, **Formula (V)(i)**, **Formula (V)(i)(a)**, **Formula (VI)**, **Formula (VI)(a)**, **Formula (VI)(i)** and **Formula (VI)(i)(a)** as described herein, in combination with said microbicide. In some

embodiments, the microbicide comprises copper (e.g., copper hydroxide). In some embodiments, the microbicide is an antibiotic or a bacteriophage. In some embodiments, the applying step is carried out by applying the active compound and the microbicide simultaneously. In some embodiments, the applying step is carried out by applying the active compound and the microbicide sequentially. In some embodiments, the compound is a compound of **Formula (II)(a)(5)(D)**:



or an agriculturally acceptable salt thereof.

Also provided are methods of enhancing the effects of a plant defense activator comprising applying an active compound selected from the group consisting of compounds of **Formula (I)**, **Formula (I)(a)(1)**, **Formula (I)(b)(1)**, **Formula (I)(a)(2)**, **Formula (I)(b)(2)**, **Formula (I)(i)**, **Formula (I)(i)(a)**, **Formula (II)**, **Formula (II)(a)**, **Formula (II)(i)**, **Formula (II)(i)(a)**, **Formula (III)**, **Formula (III)(a)**, **Formula (III)(b)**, **Formula (III)(b)(i)**, **Formula (III)(b)(ii)**, **Formula (IV)**, **Formula (IV)(a)**, **Formula (IV)(i)**, **Formula (IV)(i)(a)**, **Formula (V)**, **Formula (V)(a)**, **Formula (V)(i)**, **Formula (V)(i)(a)**, **Formula (VI)**, **Formula (VI)(a)**, **Formula (VI)(i)** and **Formula (VI)(i)(a)** as described herein, in combination with said plant defense activator. In some embodiments, the applying step is carried out by applying the active compound and the microbicide simultaneously. In some embodiments, the applying step is carried out by applying the active compound and the microbicide sequentially. In some embodiments, the compound is a compound of **Formula (II)(a)(5)(D)**:



or an agriculturally acceptable salt thereof.

A further aspect of the present invention is an antimicrobial or biofilm preventing, removing or inhibiting compound selected from the group consisting of compounds of **Formula (I)**, **Formula (I)(a)(1)**, **Formula (I)(b)(1)**, **Formula (I)(a)(2)**, **Formula (I)(b)(2)**, **Formula (I)(i)**, **Formula (I)(i)(a)**, **Formula (II)**, **Formula (II)(a)**, **Formula (II)(i)**, **Formula (II)(i)(a)**, **Formula (III)**, **Formula (III)(a)**, **Formula (III)(b)**, **Formula (III)(b)(i)**, **Formula (III)(b)(ii)**, **Formula (IV)**, **Formula (IV)(a)**, **Formula (IV)(i)**, **Formula (IV)(i)(a)**, **Formula (V)**, **Formula (V)(a)**, **Formula (V)(i)**, **Formula (V)(i)(a)**, **Formula (VI)**, **Formula (VI)(a)**, **Formula (VI)(i)**

and **Formula (VI)(i)(a)** as described herein, for use in treating or preventing a bacterial or fungal infection in a plant or plant part as described above and below.

Brief Description of the Drawings

Figure 1. Construction of the initial 2-AIT library.

Figure 2. Variation on tether length.

Figure 3. Effects of bacterial spot disease on pepper plants and crops.

Figure 4. Evaluation of foliar disease on inoculated pepper plants.

Figure 5. Pepper fruit yield per inoculated plant.

Figure 6. Evaluation of foliar disease on pepper plants measured by plot average.

Figure 7. Average pepper fruit yield per plant/plot, July 1 and 9 harvests.

Detailed Description of the Preferred Embodiments

The present invention is further described below. All patent references referred to in this patent application are hereby incorporated by reference in their entirety as if set forth fully herein.

A. Definitions.

“Active compound” as used herein refers to the various embodiments of compounds described in Section B (triazole derivatives) set forth below.

“Plant” as used herein includes all members of the plant kingdom, including higher (or “vascular”) plants and lower (“non-vascular”) plants, and particularly including all plants in the divisions Filicinae, Gymnospermae (or “gymnosperm”), and Angiospermae (or “Angiosperm”). Nonvascular plants of the present invention include, but are not limited to, bryophytes.

A plant of the present invention includes, but is not limited to, a crop plant, a turf grass, an ornamental species, a species grown for timber or pulp, a species grown for biofuels or species grown for pharmaceuticals. Additionally, plants of the present invention include, but are not limited to, tobacco, tomato, potato, sugar beet, pea, carrot, cauliflower, broccoli, soybean, canola, sunflower, alfalfa, cotton, rapeseed, Arabidopsis, peach, pepper, apple, chile, peanut, orange, grape, coffee, cassava, spinach, lettuce, cucumber, wheat, maize, rye, rice, turfgrass, oat, barley, sorghum, millet, sugarcane, or banana.

“Angiosperm” as used herein includes, but is not limited to, plants of the sub-classes Monocotyledoneae (or monocots) and Dicotyledoneae (or dicots).

Monocotyledoneae (or monocots) as used herein includes but is not limited to Amaryllidaceae-the Amaryllis Family, Gramineae(Poaceae)-the Grass Family, Liliaceae-the Lily Family, Orchidaceae-the Orchid Family, Palmae(Aracaceae)-the Palm Family; and Lemnace—the duckweed family.

Dicotyledoneae (or dicots) as used herein includes but is not limited to Cactaceae-the Cactus Family, Compositae (Asteraceae)- the Sunflower Family, Cruciferae (Brassicaceae) – the Mustard Family, Cucurbitaceae-the Gourd Family, Ericaceae-the Heath Family, Euphorbiaceae-the Spurge Family, Lauraceae-the Laurel Family, Leguminosae (Fabaceae)-the Pea Family,
 5 Rosaceae-the Rose Family, Rutaceae-the Rue Family, Solanaceae-the Nightshade Family, and Umbelliferae (Apiaceae)-the Carrot family.

Gymnospermae (or “Gymnosperms”) as used herein includes but is not limited to conifers.

“Conifer,” as used herein, refers to a member of the order Coniferae in the sub-phylum
 10 Gymnospermae in the phylum Spermatophyta. Exemplary conifers which may be used in practicing the present invention are the members of the family Pinaceae, which include, for example, loblolly pine (*Pinus taeda*), slash pine (*Pinus elliotii*), longleaf pine (*Pinus palustris*), shortleaf pine (*Pinus echinata*), ponderosa pine (*Pinus ponderosa*), red pine (*Pinus resinosa*), jack pine (*Pinus banksiana*), Eastern white pine (*Pinus strobus*), Western white pine (*Pinus monticola*),
 15 sugar pine (*Pinus lambertiana*), lodgepole pine (*Pinus contorta*), Monterey pine (*Pinus radiata*), Afghan pine (*Pinus eldarica*), Scots pine (*Pinus sylvestris*), and Virginia pine (*Pinus virginiana*); Douglas-fir (*Pseudotsuga menziesii*); Western hemlock (*Tsuga canadensis*); Sitka spruce (*Picea glauca*); redwood (*Sequoia sempervirens*); the true firs including silver fir (*Abies amabilis*), grand fir (*Abies grandis*) noble fir (*Abies procera*), white fir (*Abies concolor*),
 20 balsam fir (*Abies balsamea*); and the cedars which include Western red cedar (*Thuja plicata*), incense cedar (*Libocedrus decurrens*), Port Orford cedar (*Chamaecyparis lawsoniana*), and Alaska yellow-cedar (*Chamaecyparis nootkatensis*); and Western larch (*Larix occidentalis*). *See, e.g.*, US Patent No. 5,122,466 to Stomp et al.

“Duckweed” as used herein includes plants of the genus *Lemna* (*L. aequinoctialis*, *L.*
 25 *disperma*, *L. ecuadoriensis*, *L. gibba*, *L. japonica*, *L. minor*, *L. miniscula*, *L. obscura*, *L. perpusilla*, *L. tenera*, *L. trisulca*, *L. turionifera*, *L. valdiviana*); genus *Spirodela* (*S. intermedia*, *S. polyrrhiza*, *S. punctata*); genus *Wolffia* (*Wa. angusta*, *Wa. arrhiza*, *Wa. australina*, *Wa. borealis*, *Wa. brasiliensis*, *Wa. columbiana*, *Wa. elongata*, *Wa. globosa*, *Wa. microscopica*, *Wa. neglecta*) and genus *Wolffiella* (*Wl. caudata*, *Wl. denticulata*, *Wl. gladiata*, *Wl. hyalina*, *Wl. lingulata*, *Wl.*
 30 *repunda*, *Wl. rotunda*, and *Wl. neotropica*). *See, e.g.*, US Patent No. 7,161,064 to Stomp et al.

Particular examples of plants include but are not limited to all cereal and grain crops, herbs and spices, oil seed crops, sugarcane, vegetable crops, brassica vegetables, bulb vegetables, cucurbit vegetables and fruit, leafy vegetables, fruiting vegetables, legume vegetables, root and tuber vegetables, tree, vine and shrub crops, berry crops, citrus (*e.g.*, orange, grapefruit,
 35 Mandarin (including Tangerine and Satsuma), lemon, lime, and kumquat), pome fruit (*e.g.*,


apple, pear, quince, Asian pear, loquat, etc.), stone fruit (e.g., peach, apricot, prune, plum, cherries, almond, etc.), miscellaneous tree food crops, non-food tree crops, tree nuts, tropical and subtropical trees and fruit, vine crops, pasture grasses, forage legumes, and rangeland, grass seed or sod production, pastures, cotton, corn, soybeans, rice, wheat, greenhouse/shadehouse grown plants, ornamental, plant nurseries, Christmas trees, golf courses and turf, forestry, tobacco, orchids, flowers and roses, foliage crops, algae such as green algae, bryophytes (mosses, liverworts, hornworts), etc. Note that “foliage crops” refers to the types of plants (ferns, etc.) that are typically used in home or commercial settings for decorative purposes; this alone constitutes a very large commercial industry.

“Plant part” as used herein refers to seeds, roots, leaves, shoots, fruits (e.g., apples, pineapples, citrus fruit, etc.), vegetables, tubers, flowers (e.g., cut flowers such as roses, as well as the reproductive parts of plants), petals, stem, trunk, etc., harvested or collected from a plant as described herein. The plant part of a vascular plant may be a non-vascular part, such as a seed or meristem (growing tip of a shoot).


“Applying” as described herein can be carried out directly or indirectly by any suitable technique, including topically applying to the plant or plant part, applying to the media in which the plant or plant part is grown, stored, displayed or maintained (e.g., adding to water in which the stems of cut flowers are placed), etc. Note that the plant may be grown in any suitable media, including but not limited to soil, potting soil, soilless media such as sand and hydroponic media (including solution culture, medium culture, and deep water culture), etc.

“Agricultural composition” as described herein may be in any suitable form, including but not limited to: wettable powders, dry flowables, soluble powders, water dispersibles, liquids, dusts, emulsifiable concentrates, flowables, fumigants, water dispersible granules, liquid concentrates, granules, water soluble packages, wettable powders in water soluble films, emulsions, etc.

“Triazole” refers to the commonly known structures:



“Imidazole” refers to the commonly known structure:



“H” refers to a hydrogen atom. “C” refers to a carbon atom. “N” refers to a nitrogen atom. “O” refers to an oxygen atom. “Halo” refers to F, Cl, Br or I. The term “hydroxy,” as used herein, refers to an -OH moiety. “Br” refers to a bromine atom. “Cl” refers to a chlorine atom. “I” refers to an iodine atom. “F” refers to a fluorine atom.

5 An “acyl group” is intended to mean a group $-C(O)-R$, where R is a suitable substituent (for example, an acetyl group, a propionyl group, a butyryl group, a benzoyl group, or an alkylbenzoyl group).

“Alkyl,” as used herein, refers to a straight or branched chain hydrocarbon containing from 1 or 2 to 10 or 20 or more carbon atoms (e.g., C₂, C₃, C₄, C₅, C₆, C₇, C₈, C₉, C₁₀, C₁₁,
10 C₁₂, C₁₃, C₁₄, C₁₅, etc.). Representative examples of alkyl include, but are not limited to, methyl, ethyl, n-propyl, iso-propyl, n-butyl, sec-butyl, iso-butyl, tert-butyl, n-pentyl, isopentyl, neopentyl, n-hexyl, 3-methylhexyl, 2,2-dimethylpentyl, 2,3-dimethylpentyl, n-heptyl, n-octyl, n-nonyl, n-decyl, and the like. In some embodiments, alkyl groups as described herein are optionally substituted (e.g., from 1 to 3 or 4 times) with independently selected H, halo, hydroxy,
15 acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide.

As generally understood by those of ordinary skill in the art, “saturation” refers to the state in which all available valence bonds of an atom (e.g., carbon) are attached to other atoms.
20 Similarly, “unsaturation” refers to the state in which not all the available valence bonds are attached to other atoms; in such compounds the extra bonds usually take the form of double or triple bonds (usually with carbon). For example, a carbon chain is “saturated” when there are no double or triple bonds present along the chain or directly connected to the chain (e.g., a carbonyl), and is “unsaturated” when at least one double or triple bond is present along the chain or directly
25 connected to the chain (e.g., a carbonyl). Further, the presence or absence of a substituent depending upon chain saturation will be understood by those of ordinary skill in the art to depend upon the valence requirement of the atom or atoms to which the substituent binds (e.g., carbon).

The term “optionally substituted” indicates that the specified group is either unsubstituted, or substituted by one or more suitable substituents. A “substituent” is an atom or
30 atoms substituted in place of a hydrogen atom on the parent chain or cycle of an organic molecule, for example, H, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide.

“Alkenyl,” as used herein, refers to a straight or branched chain hydrocarbon containing
35 from 1 or 2 to 10 or 20 or more carbons, and containing at least one carbon-carbon double bond,

formed structurally, for example, by the replacement of two hydrogens. Representative examples of "alkenyl" include, but are not limited to, ethenyl, 2-propenyl, 2-methyl-2-propenyl, 3-butenyl, 4-pentenyl, 5-hexenyl, 2-heptenyl, 2-methyl-1-heptenyl, 3-decenyl and the like. In some embodiments, alkenyl groups as described herein are optionally substituted (e.g., from 1 to 3 or 4
5 times) with independently selected H, halo, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide.

"Alkynyl," as used herein, refers to a straight or branched chain hydrocarbon group containing from 1 or 2 to 10 or 20 or more carbon atoms, and containing at least one carbon-
10 carbon triple bond. Representative examples of alkynyl include, but are not limited, to acetylenyl, 1-propynyl, 2-propynyl, 3-butyne, 2-pentyne, 1-butyne and the like. In some embodiments, alkynyl groups as described herein are optionally substituted (e.g., from 1 to 3 or 4 times) with independently selected H, halo, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro,
15 carbonyl, carboxy, amino acid sidechain, amino acid and peptide.

The term "cycloalkyl," as used herein, refers to a saturated cyclic hydrocarbon group containing from 3 to 8 carbons or more. Representative examples of cycloalkyl include, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, and cyclooctyl. In some
20 embodiments, cycloalkyl groups as described herein are optionally substituted (e.g., from 1 to 3 or 4 times) with independently selected H, halo, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide.

"Heterocyclo," as used herein, refers to a monocyclic or a bicyclic ring system. Monocyclic heterocycle ring systems are exemplified by any 5 or 6 member ring containing 1, 2,
25 3, or 4 heteroatoms independently selected from the group consisting of: O, N, and S. The 5 member ring has from 0 to 2 double bonds, and the 6 member ring has from 0-3 double bonds. Representative examples of monocyclic ring systems include, but are not limited to, azetidine, azepine, aziridine, diazepine, 1,3-dioxolane, dioxane, dithiane, furan, imidazole, imidazoline, imidazolidine, isothiazole, isothiazoline, isothiazolidine, isoxazole, isoxazoline, isoxazolidine,
30 morpholine, oxadiazole, oxadiazoline, oxadiazolidine, oxazole, oxazoline, oxazolidine, piperazine, piperidine, pyran, pyrazine, pyrazole, pyrazoline, pyrazolidine, pyridine, pyrimidine, pyridazine, pyrrole, pyrroline, pyrrolidine, tetrahydrofuran, tetrahydrothiophene, tetrazine, tetrazole, thiadiazole, thiadiazoline, thiadiazolidine, thiazole, thiazoline, thiazolidine, thiophene, thiomorpholine, thiomorpholine sulfone, sulfoxide, thiopyran, triazine, triazole, trithiane, and the
35 like. Bicyclic ring systems are exemplified by any of the above monocyclic ring systems fused to

an aryl group as defined herein, a cycloalkyl group as defined herein, or another monocyclic ring system as defined herein. Representative examples of bicyclic ring systems include but are not limited to, for example, benzimidazole, benzothiazole, benzothiadiazole, benzothiophene, benzoxadiazole, benzoxazole, benzofuran, benzopyran, benzothiopyran, benzodioxine, 1,3-benzodioxole, cinnoline, indazole, indole, indoline, indolizine, naphthyridine, isobenzofuran, isobenzothiophene, isoindole, isoindoline, isoquinoline, phthalazine, pyranopyridine, quinoline, quinolizine, quinoxaline, quinazoline, tetrahydroisoquinoline, tetrahydroquinoline, thiopyranopyridine, and the like. In some embodiments, heterocyclo groups as described herein are optionally substituted (e.g., from 1 to 3 or 4 times) with independently selected H, halo, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide.

“Aryl” as used herein refers to a fused ring system having one or more aromatic rings. Representative examples of aryl include, azulenyl, indanyl, indenyl, naphthyl, phenyl, tetrahydronaphthyl, and the like. The aryl groups of this invention can be substituted with 1, 2, 3, 4, or 5 substituents independently selected from alkenyl, alkenyloxy, alkoxy, alkoxyalkoxy, alkoxycarbonyl, alkyl, alkylcarbonyl, alkylcarbonyloxy, alkylsulfinyl, alkylsulfonyl, alkylthio, alkynyl, aryl, aryloxy, azido, arylalkoxy, arylalkyl, aryloxy, carboxy, cyano, formyl, halogen, haloalkyl, haloalkoxy, hydroxy, hydroxyalkyl, mercapto, nitro, sulfamyl, sulfo, sulfonate, $-NR'R''$ (wherein, R' and R'' are independently selected from hydrogen, alkyl, alkylcarbonyl, aryl, arylalkyl and formyl), and $-C(O)NR'R''$ (wherein R' and R'' are independently selected from hydrogen, alkyl, alkylcarbonyl, aryl, arylalkyl, and formyl). In some embodiments, aryl groups as described herein are optionally substituted (e.g., from 1 to 3 or 4 times) with independently selected H, halo, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide.


“Heteroaryl” means a cyclic, aromatic hydrocarbon in which one or more carbon atoms have been replaced with heteroatoms. If the heteroaryl group contains more than one heteroatom, the heteroatoms may be the same or different. Examples of heteroaryl groups include pyridyl, pyrimidinyl, imidazolyl, thienyl, furyl, pyrazinyl, pyrrolyl, pyranyl, isobenzofuranyl, chromenyl, xanthenyl, indolyl, isoindolyl, indolizynyl, triazolyl, pyridazinyl, indazolyl, purinyl, quinolizynyl, isoquinolyl, quinolyl, phthalazinyl, naphthyridinyl, quinoxalinyl, isothiazolyl, and benzo[b]thienyl. Preferred heteroaryl groups are five and six membered rings and contain from one to three heteroatoms independently selected from the group consisting of: O, N, and S. The heteroaryl group, including each heteroatom, can be unsubstituted or substituted with from 1 to 4

suitable substituents, as chemically feasible. For example, the heteroatom S may be substituted with one or two oxo groups, which may be shown as =O. In some embodiments, heteroaryl groups as described herein are optionally substituted (e.g., from 1 to 3 or 4 times) with independently selected H, halo, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo,
 5 aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide.

“Alkoxy,” as used herein, refers to an alkyl group, as defined herein, appended to the parent molecular moiety through an oxy group, as defined herein. Representative examples of alkoxy include, but are not limited to, methoxy, ethoxy, propoxy, 2-propoxy, butoxy, tert-butoxy,
 10 pentyloxy, hexyloxy and the like. In some embodiments, alkoxy groups as described herein are optionally substituted (e.g., from 1 to 3 or 4 times) with independently selected H, halo, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide.

An “amine” or “amino” is intended to mean the group -NH_2 . “Optionally substituted” amines refers to -NH_2 groups wherein none, one or two of the hydrogens is replaced by a suitable substituent as described herein, such as alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo,
 15 aryl, heteroaryl, alkoxy, carbonyl, carboxy, etc. In some embodiments, one or two of the hydrogens are optionally substituted with independently selected, halo, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide. Disubstituted amines may have substituents that are bridging, i.e., form a heterocyclic ring structure that includes the amine nitrogen.


An “amide” as used herein refers to an organic functional group having a carbonyl group (C=O) linked to a nitrogen atom (N), or a compound that contains this group, generally depicted
 25 as:



wherein, R and R' can independently be any covalently-linked atom or atoms, for example, H, halo, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy,
 30 amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide.


A “thiol” or “mercapto” refers to an -SH group or to its tautomer =S .

A “sulfone” as used herein refers to a sulfonyl functional group, generally depicted as:



wherein, R can be any covalently-linked atom or atoms, for example, H, halo, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide.

A “sulfoxide” as used herein refers to a sulfinyl functional group, generally depicted as:



wherein, R can be any covalently-linked atom or atoms, for example, H, halohydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide.

The term “oxo,” as used herein, refers to a =O moiety. The term “oxy,” as used herein, refers to a -O- moiety.

“Nitro” refers to the organic compound functional group $-\text{NO}_2$.

“Carbonyl” is a functional group having a carbon atom double-bonded to an oxygen atom ($-\text{C}=\text{O}$). “Carboxy” as used herein refers to a $-\text{COOH}$ functional group, also written as $-(\text{C}=\text{O})-\text{OH}$.

“Amino acid sidechain” as used herein refers to any of the 20 commonly known groups associated with naturally-occurring amino acids, or any natural or synthetic homologue thereof. An “amino acid” includes the sidechain group and the amino group, alpha-carbon atom, and carboxy groups, as commonly described in the art. Examples of amino acids include glycine, and glycine that is substituted with a suitable substituent as described herein, such as alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, carbonyl, carboxy, etc., or an agriculturally acceptable salt thereof. For example, “Histidine” is one of the 20 most commonly known amino acids found naturally in proteins. It contains an imidazole side chain substituent. Other examples of naturally-occurring amino acids include lysine, arginine, aspartic acid, glutamic acid, asparagine, glutamine, serine, threonine, tyrosine, alanine, valine, leucine, isoleucine, phenylalanine, methionine, cryptophan, and cysteine. Also included in the definitions of “amino acid sidechain” and “amino acid” is proline, which is commonly included in the definition of an amino acid, but is technically an imino acid. As used in this application, both the

naturally-occurring L-, and the non-natural D-amino acid enantiomers are included. The single letter code for amino acids is A (Ala), C (Cys), D (Asp), E (Glu), F (Phe), G (Gly), H (His), I (Ile), K (Lys), L (Leu), M (Met), N (Asn), P (Pro), Q (Gln), R (Arg), S (Ser), T (Thr), V (Val), W (Trp), and Y (Tyr). A “peptide” is a linear chain of amino acids covalently linked together, typically through an amide linkage, and contains from 1 or 2 to 10 or 20 or more amino acids, and is also optionally substituted and/or branched.

“Agriculturally acceptable salt” is intended to mean a salt that retains the biological effectiveness of the free acids and bases of a specified compound and that is not biologically or otherwise undesirable. Examples of agriculturally acceptable salts include sulfates, pyrosulfates, bisulfates, sulfites, bisulfites, phosphates, monohydrogenphosphates, dihydrogenphosphates, metaphosphates, pyrophosphates, chlorides, bromides, iodides, acetates, propionates, decanoates, caprylates, acrylates, formates, isobutyrate, caproates, heptanoates, propiolates, oxalates, malonates, succinates, suberates, sebacates, fumarates, maleates, butyne-1,4-dioates, hexyne-1,6-dioates, benzoates, chlorobenzoates, methylbenzoates, dinitrobenzoates, hydroxybenzoates, methoxybenzoates, phthalates, sulfonates, xylenesulfonates, phenylacetates, phenylpropionates, phenylbutyrates, citrates, lactates, γ -hydroxybutyrates, glycollates, tartrates, methane-sulfonates, propanesulfonates, naphthalene-1-sulfonates, naphthalene-2-sulfonates, and mandelates.

The term “optionally substituted” indicates that the specified group is either unsubstituted, or substituted by one or more suitable substituents. A “substituent” is an atom or atoms substituted in place of a hydrogen atom on the parent chain or cycle of an organic molecule.

B. Active compounds.

Active compounds are provided below. In some of the embodiments provided in the present invention, active compounds are derivatives of triazole. In some embodiments, active compounds include imidazole-triazole conjugates. In some embodiments, active compounds include 2-aminoimidazole-triazole conjugates (“2-AIT”). Active compounds as described herein can be prepared as detailed below or in accordance with known procedures or variations thereof that will be apparent to those skilled in the art.

As will be appreciated by those of skill in the art, the active compounds of the various formulas disclosed herein may contain chiral centers, e.g. asymmetric carbon atoms. Thus, the present invention is concerned with the synthesis of both: (i) racemic mixtures of the active compounds, and (ii) enantiomeric forms of the active compounds. The resolution of racemates into enantiomeric forms can be done in accordance with known procedures in the art. For

example, the racemate may be converted with an optically active reagent into a diastereomeric pair, and the diastereomeric pair subsequently separated into the enantiomeric forms.


Geometric isomers of double bonds and the like may also be present in the compounds disclosed herein, and all such stable isomers are included within the present invention unless
 5 otherwise specified. Also included in active compounds of the invention are tautomers (e.g., tautomers of triazole and/or imidazole) and rotamers.

All chains defined by the formulas herein which include three or more carbons may be saturated or unsaturated unless otherwise indicated.


Carbons or other atoms along a chain identified by the Formulas herein may be identified
 10 by number, and when identified by number shall be numbered from left to right. For example:




To illustrate where there are two or more discrete chains, for **Formula (II)(i)(a)**
 15 described below, wherein R^2, R^3, R^4, R^5, R^7 and $R^8 = \text{H}$, and $R^6 = \text{phenyl}$:



the exemplary structure below shows $n=5$, saturated, one of either R^x or $R^y = \text{methyl}$ at C4; $m=3$,
 20 unsaturated, $R^u = \text{methyl}$ at C2 (R^v is absent at C2); and R^x, R^y, R^u and $R^v = \text{H}$ at all other occurrences:



Active compounds for carrying out the present invention include compounds of **Formula (I)**:



(I)

wherein:

R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 and R^8 are each independently selected from the group consisting of: H, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide;

each occurrence of R^x and R^y is present or absent (depending upon chain saturation), and is independently selected from the group consisting of: H, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide; and

A, B, D, E, F, G and H are each independently selected from carbon, N, S and O, wherein at least one of D, E, F, G and H is carbon; and


$n=0$ to 20, saturated or unsaturated;

or an agriculturally acceptable salt thereof.

This formula may be optionally substituted (e.g., from 1 to 3 or 4 times) with independently selected H, halo, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide.

As will be appreciated by those of skill in the art, a given substituent (R^1 - R^8) may be present or absent depending upon the valence requirement of the atom or atoms to which the substituent binds (e.g., carbon versus nitrogen).

In some embodiments of **Formula (I)**, R^1 is a substituted amino, A, B, F, G and H are each N, and D and E are each carbon, generally depicted by **Formulas (I)(a)(1) – (I)(b)(1)**:



(I)(a)(1)

wherein:


R^{1a} , R^{1b} , R^2 , R^3 , R^5 and R^6 are each independently selected from the group consisting of: H, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide; and

each occurrence of R^x and R^y is present or absent (depending upon chain saturation), and is independently selected from the group consisting of: H, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide; and

$n=0$ to 20;

or an agriculturally acceptable salt thereof.

This formula may be optionally substituted (e.g., from 1 to 3 or 4 times) with independently selected H, halo, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide.



(I)(a)(2)

wherein:

R^{1a} , R^{1b} , R^2 , R^3 , R^5 and R^8 are each independently selected from the group consisting of: H, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino,

amide, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide; and


each occurrence of R^x and R^y is present or absent (depending upon chain saturation), and is independently selected from the group consisting of: H, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide; and

$n=0$ to 20;

or an agriculturally acceptable salt thereof.

This formula may be optionally substituted (e.g., from 1 to 3 or 4 times) with independently selected H, halo, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide.

In some embodiments of **Formula (I)**, R^1 is a substituted amino, A, B, F, G and D are each N, and D and E are each carbon, generally depicted by **Formula (I)(b)(2)**:



(I)(b)(2)

wherein:

R^{1a} , R^{1b} , R^2 , R^3 , R^5 and R^8 are each independently selected from the group consisting of: H, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide;

each occurrence of R^x and R^y is present or absent (depending upon chain saturation), and is independently selected from the group consisting of: H, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide; and


$n=0$ to 20;

or an agriculturally acceptable salt thereof.

This formula may be optionally substituted (e.g., from 1 to 3 or 4 times) with independently selected H, halo, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl,

alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide.

Active compounds further include compounds of **Formula (I)(i)**:



(I)(i)

wherein:

$R^1, R^2, R^3, R^4, R^5, R^6, R^7$ and R^8 are each independently selected from the group consisting of: H, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide;

A, B, D, E, F, G and H are each independently selected from carbon, N, S and O, wherein at least one of D, E, F, G and H is carbon; and


$n=0$ to 20, saturated or unsaturated;

or an agriculturally acceptable salt thereof.

This formula may be optionally substituted (e.g., from 1 to 3 or 4 times) with independently selected H, halo, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide.

As will be appreciated by those of skill in the art, a given substituent (R^1 - R^8) may be present or absent depending upon the valence requirement of the atom or atoms to which the substituent binds (e.g., carbon versus nitrogen).

In some embodiments of **Formula (I)(i)**, R^1 is a substituted amino; R^2, R^3, R^4, R^5, R^7 and $R^8=H$; A, B, F, G and H are each N, and D and E are each carbon, generally depicted by **Formula (I)(i)(a)**:



(I)(i)(a)

wherein:


R⁶ is selected from the group consisting of: H, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide; and

n=0 to 20, saturated or unsaturated;

or an agriculturally acceptable salt thereof.

This formula may be optionally substituted (e.g., from 1 to 3 or 4 times) with independently selected H, halo, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide.

Active compounds for carrying out the present invention include compounds of **Formula (II)**:



(II)

wherein:

R¹, R², R³, R⁴, R⁵, R⁶, R⁷ and R⁸ are each independently selected from the group consisting of: H, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide;

each occurrence of R^x, R^y, R^u and R^v is present or absent (depending upon chain saturation), and is independently selected from the group consisting of: H, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide;

A, B, D, E, F, G and H are each independently selected from carbon, N, S and O, wherein at least one of D, E, F, G and H is carbon; and

n=0 to 20; and


m=0 to 20;

or an agriculturally acceptable salt thereof.

This formula may be optionally substituted (e.g., from 1 to 3 or 4 times) with independently selected H, halo, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide.

5 As will be appreciated by those of skill in the art, a given substituent (R^1 - R^8) may be present or absent depending upon the valence requirement of the atom or atoms to which the substituent binds (e.g., carbon versus nitrogen).

In some embodiments of **Formula (II)**, R^1 is a substituted amino, A, B, F, G and H are each N, and D and E are each carbon, generally depicted by **Formula (II)(a)**:



(II)(a)

wherein:

R^{1a} , R^{1b} , R^2 , R^3 , R^5 and R^6 are each independently selected from the group consisting of: H, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide;

each occurrence of R^x , R^y , R^u and R^v is present or absent (depending upon chain saturation), and is independently selected from the group consisting of: H, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide;

$n=0$ to 20; and

$m=0$ to 20;

or an agriculturally acceptable salt thereof.


This formula may be optionally substituted (e.g., from 1 to 3 or 4 times) with independently selected H, halo, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide.

In some embodiments of **Formula (II)(a)**, R^{1a} , R^{1b} , R^2 , R^3 and R^5 are each H and R^6 is phenyl, examples of which include, but are not limited to, the following exemplary **Formulas**. Each occurrence of R^x , R^y , R^u and R^v present is H unless otherwise indicated.


Formulas (II)(a)(1)(A) – (II)(a)(1)(D), wherein $n=1$:

5


$m=1$:




$m=2$:



$m=3$:




$m=3$, R^u =methyl at C2




10

Formulas (II)(a)(2)(A) – (II)(a)(2)(D), wherein $n=2$:


$m=1$:




$m=2$:



$m=3$:




$m=3$, R^u =methyl at C2:




15

Formulas (II)(a)(3)(A) – (II)(a)(3)(D), wherein $n=3$:


$m=1$:




$m=2$:



$m=3$:




$m=3$, R^u =methyl at C2:




20


Formula (II)(a)(4)(D), wherein $n=4$; $m=3$, R^u =methyl at C2:



Formula (II)(a)(5)(D), wherein $n=5$; $m=3$, R^u =methyl at C2:




Formula (II)(a)(6)(D), wherein $n=6$; $m=3$, R^u =methyl at C2:



5

Active compounds further include compounds of **Formula (II)(i)**:



wherein:

10 R^1 and R^6 are each independently selected from the group consisting of: H, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide;

15 A, B, D, E, F, G and H are each independently selected from carbon, N, S and O, wherein at least one of D, E, F, G and H is carbon;


$n=0$ to 20, saturated or unsaturated; and

$m=0$ to 20, saturated or unsaturated;

or an agriculturally acceptable salt thereof.

20 This formula may be optionally substituted (e.g., from 1 to 3 or 4 times) with independently selected H, halo, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide.

In some embodiments of **Formula (II)(i)**, R^1 is a substituted amino, A, B, F, G and H are each N, and D and E are each carbon, generally depicted by **Formula (II)(i)(a)**:



(II)(i)(a)

wherein:

R^{1a} , R^{1b} and R^6 are each independently selected from the group consisting of: H, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide;

$n=0$ to 20, saturated or unsaturated; and

$m=0$ to 20, saturated or unsaturated;


or an agriculturally acceptable salt thereof.

This formula may be optionally substituted (e.g., from 1 to 3 or 4 times) with independently selected H, halo, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide.


In some embodiments of **Formula (II)(i)(a)**, R^{1a} and R^{1b} are each H, and R^6 is heteroaryl, examples of which include, but are not limited to, the following exemplary **Formulas**:

Formulas (II)(i)(a)(1)(E) – (II)(i)(a)(1)(L), wherein $n=1$:



$m=1$; R^4 is thiophenyl:




$m=2$; R^4 is thiophenyl:




$m=1$; R^4 is furyl:





m=1; R⁴ is indolyl:



m=2; R⁴ is indolyl:




m=1; R⁴ is benzimidazolyl:



Further embodiments include **Formulas (II)(i)(a)(2)(E) – (II)(i)(a)(2)(L)**, wherein n=2; **Formulas (II)(i)(a)(3)(E) – (II)(i)(a)(3)(L)**, wherein n=3; **Formulas (II)(i)(a)(4)(E) – (II)(i)(a)(4)(L)**, wherein n=4; **Formulas (II)(i)(a)(5)(E) – (II)(i)(a)(5)(L)**, wherein n=5; **Formulas (II)(i)(a)(6)(E) – (II)(i)(a)(6)(L)**, wherein n=6; and so on.

These formulas may be optionally substituted (e.g., from 1 to 3 or 4 times) with independently selected H, halo, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide.

Also provided are compounds of **Formula (III)**:



(III)

wherein:

R⁵, R⁶, R⁷, R⁸ and R⁹ are each independently selected from the group consisting of: H, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide; and


A, B, D, E and F are each independently selected from carbon, N, S and O, wherein at least one of A, B, D, E and F is carbon;

or an agriculturally acceptable salt thereof.

This formula may be optionally substituted (e.g., from 1 to 3 or 4 times) with independently selected H, halo, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide.

5 As will be appreciated by those of skill in the art, a given substituent (R^5 - R^9) may be present or absent depending upon the valence requirement of the atom or atoms to which the substituent binds (e.g., carbon versus nitrogen).

In some embodiments of **Formula (III)**, B, D and E are each N, and A and F are each carbon, generally depicted as **Formula (III)(a)**:



(III)(a)


wherein:

R^5 , R^6 and R^9 are each independently selected from the group consisting of: H, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, 15 thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide;

or an agriculturally acceptable salt thereof.

This formula may be optionally substituted (e.g., from 1 to 3 or 4 times) with independently selected H, halo, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, 20 alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide.

In some embodiments of **Formula (III)**, A, D and E are each N, and B and F are each carbon, generally depicted as **Formula (III)(b)**:



(III)(b)


wherein:

R^5 , R^6 and R^9 are each independently selected from the group consisting of: H, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide; and

5 or an agriculturally acceptable salt thereof.

This formula may be optionally substituted (e.g., from 1 to 3 or 4 times) with independently selected H, halo, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide.

10 In some embodiments of **Formula (III)(b)**, R^6 and R^9 are each H, generally depicted by **Formula (III)(b)(i)**:



(III)(b)(i)


wherein:

15 R^5 is selected from the group consisting of: H, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide; and

or an agriculturally acceptable salt thereof.

This formula may be optionally substituted (e.g., from 1 to 3 or 4 times) with independently
20 selected H, halo, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide.

In some embodiments of **Formula (III)(b)**, R^5 and R^6 are each H, generally depicted by **Formula (III)(b)(ii)**:



(III)(b)(ii)


wherein:

R^9 is selected from the group consisting of: H, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo,
30 oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide; and

or an agriculturally acceptable salt thereof.

This formula may be optionally substituted (e.g., from 1 to 3 or 4 times) with independently selected H, halo, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide.

Also provided are compounds of **Formula (IV)**:



wherein:

$R^1, R^2, R^3, R^4, R^5, R^6, R^7$ and R^8 are each independently selected from the group consisting of: H, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide;

each occurrence of R^x, R^y, R^u, R^v, R^z and R^w is present or absent (depending upon chain saturation), and is independently selected from the group consisting of: H, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide;

A, B, D, E, F, G and H are each independently selected from carbon, N, S and O, wherein at least one of D, E, F, G and H is carbon; and

$n=0$ to 20;


$m=0$ to 20; and

$p=0$ to 20

or an agriculturally acceptable salt thereof.

This formula may be optionally substituted (e.g., from 1 to 3 or 4 times) with independently selected H, halo, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide.

In some embodiments of **Formula (IV)**, R^1 is a substituted amino, A, B, F, G and H are each N, and D and E are each carbon, generally depicted by **Formula (IV)(a)**:



(IV)(a)

wherein:

R^{1a} , R^{1b} , R^2 , R^3 , R^5 and R^6 are each independently selected from the group consisting of: H, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide;

each occurrence of R^x , R^y , R^u , R^v , R^z and R^w is present or absent (depending upon chain saturation), and is independently selected from the group consisting of: H, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide;

$n=0$ to 20;


$m=0$ to 20; and

$p=0$ to 20

or an agriculturally acceptable salt thereof.

This formula may be optionally substituted (e.g., from 1 to 3 or 4 times) with independently selected H, halo, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide.

In some embodiments, R^6 is a group:




(i)

wherein:

X, Y and Z are each independently selected from the group consisting of: H, methyl, Br and Cl.


In some embodiments, R^6 is a group:



wherein:

- 5 R^{20} , R^{21} , R^{22} , R^{23} and R^{24} are each independently selected from the group consisting of: H, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide.

Further provided are compounds of **Formula (IV)(i)**:



wherein:

- R^1 and R^6 are each independently selected from the group consisting of: H, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide;

A, B, D, E, F, G and H are each independently selected from carbon, N, S and O, wherein at least one of D, E, F, G and H is carbon;

$n=0$ to 20, saturated or unsaturated;


20 $m=0$ to 20, saturated or unsaturated; and

$p=0$ to 20, saturated or unsaturated;

or an agriculturally acceptable salt thereof.

This formula may be optionally substituted (e.g., from 1 to 3 or 4 times) with independently selected H, halo, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide.

In some embodiments of **Formula (IV)(i)**, R^1 is a substituted amino, A, B, F, G and H are each N, and D and E are each carbon, generally depicted by **Formula (IV)(i)(a)**:



(IV)(i)(a)

wherein:

R^{1a} , R^{1b} and R^6 are each independently selected from the group consisting of: H, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide;

$n=0$ to 20, saturated or unsaturated;

$m=0$ to 20, saturated or unsaturated; and


$p=0$ to 20, saturated or unsaturated;

or an agriculturally acceptable salt thereof.

This formula may be optionally substituted (e.g., from 1 to 3 or 4 times) with independently selected H, halo, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide.

In some embodiments of **Formula (IV)(i)(a)**, R^{1a} , R^{1b} and R^6 are each H. In some embodiments of **Formula (IV)(i)(a)**, R^{1a} and R^{1b} are each H, and R^6 is aryl or heteroaryl.

Also provided are compounds of **Formula (V)**:



(V)

wherein:

R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 and R^8 are each independently selected from the group consisting of: H, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide;

each occurrence of R^x , R^y , R^u , R^v , R^z and R^w is present or absent (depending upon chain

saturation), and is independently selected from the group consisting of: H, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide;

A, B, D, E, F, G and H are each independently selected from carbon, N, S and O, wherein
 5 at least one of D, E, F, G and H is carbon; and

$n=0$ to 20;


$m=0$ to 20; and

$p=0$ to 20

or an agriculturally acceptable salt thereof.

10 This formula may be optionally substituted (e.g., from 1 to 3 or 4 times) with independently selected H, halo, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide.

In some embodiments of **Formula (V)**, R^1 is a substituted amino, A, B, F, G and H are
 15 each N, and D and E are each carbon, generally depicted by **Formula (V)(a)**:



(V)(a)

wherein:

R^{1a} , R^{1b} , R^2 , R^3 , R^5 and R^6 are each independently selected from the group consisting of: H,
 20 hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide;

each occurrence of R^x , R^y , R^u , R^v , R^z and R^w is present or absent (depending upon chain saturation), and is independently selected from the group consisting of: H, hydroxy, acyl, alkyl,
 25 alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide;

$n=0$ to 20;


$m=0$ to 20; and

$p=0$ to 20

or an agriculturally acceptable salt thereof.

This formula may be optionally substituted (e.g., from 1 to 3 or 4 times) with independently selected H, halo, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide.

Further provided are compounds of **Formula (V)(i)**:



wherein:

R^1 and R^6 are each independently selected from the group consisting of: H, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide;

A, B, D, E, F, G and H are each independently selected from carbon, N, S and O, wherein at least one of D, E, F, G and H is carbon;

$n=0$ to 20, saturated or unsaturated;


$m=0$ to 20, saturated or unsaturated; and

$p=0$ to 20, saturated or unsaturated;

or an agriculturally acceptable salt thereof.

This formula may be optionally substituted (e.g., from 1 to 3 or 4 times) with independently selected H, halo, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide.

In some embodiments of **Formula (V)(i)**, R^1 is a substituted amino, A, B, F, G and H are each N, and D and E are each carbon, generally depicted by **Formula (V)(i)(a)**:



wherein:


R^{1a} , R^{1b} and R^6 are each independently selected from the group consisting of: H, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide;

- 5 $n=0$ to 20, saturated or unsaturated;
 $m=0$ to 20, saturated or unsaturated; and
 $p=0$ to 20, saturated or unsaturated;
 or an agriculturally acceptable salt thereof.

10 This formula may be optionally substituted (e.g., from 1 to 3 or 4 times) with independently selected H, halo, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide.

15 In some embodiments of **Formula (V)(i)(a)**, R^{1a} , R^{1b} and R^6 are each H, alkyl, cycloalkyl or heterocyclo. In some embodiments of **Formula (V)(i)(a)**, R^{1a} and R^{1b} are each H, and R^6 is aryl.

Also provided are compounds of **Formula (VI)**:



wherein:

- 20 R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 and R^8 are each independently selected from the group consisting of: H, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide;

- 25 each occurrence of R^x , R^y , R^u , R^v , R^z and R^w is present or absent (depending upon chain saturation), and is independently selected from the group consisting of: H, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide;

A, B, D, E, F, G and H are each independently selected from carbon, N, S and O, wherein at least one of D, E, F, G and H is carbon; and

n=0 to 20;


m=0 to 20; and

p=0 to 20;

or an agriculturally acceptable salt thereof.

- 5 This formula may be optionally substituted (e.g., from 1 to 3 or 4 times) with independently selected H, halo, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide.

10 In some embodiments of **Formula (VI)**, R¹ is a substituted amino, A, B, F, G and H are each N, and D and E are each carbon, generally depicted by **Formula (VI)(a)**:



(VI)(a)

wherein:

- 15 R^{1a}, R^{1b}, R², R³, R⁵ and R⁶ are each independently selected from the group consisting of: H, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide;

- 20 each occurrence of R^x, R^y, R^u, R^v, R^z and R^w is present or absent (depending upon chain saturation), and is independently selected from the group consisting of: H, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide;

n=0 to 20;


m=0 to 20; and

p=0 to 20;

- 25 or an agriculturally acceptable salt thereof.

This formula may be optionally substituted (e.g., from 1 to 3 or 4 times) with independently selected H, halo, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide.

Further provided are compounds of **Formula (VI)(i)**:



wherein:

- 5 R^1 and R^6 are each independently selected from the group consisting of: H, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide;

- 10 A, B, D, E, F, G and H are each independently selected from carbon, N, S and O, wherein at least one of D, E, F, G and H is carbon;

$n=0$ to 20, saturated or unsaturated;


$m=0$ to 20, saturated or unsaturated; and

$p=0$ to 20, saturated or unsaturated;

or an agriculturally acceptable salt thereof.

- 15 This formula may be optionally substituted (e.g., from 1 to 3 or 4 times) with independently selected H, halo, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide.

- 20 In some embodiments of **Formula (VI)(i)**, R^1 is a substituted amino, A, B, F, G and H are each N, and D and E are each carbon, generally depicted by **Formula (VI)(i)(a)**:



wherein:

- 25 R^{1a} , R^{1b} and R^6 are each independently selected from the group consisting of: H, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide;

$n=0$ to 20, saturated or unsaturated;

m=0 to 20, saturated or unsaturated; and

p=0 to 20, saturated or unsaturated;

or an agriculturally acceptable salt thereof.

This formula may be optionally substituted (e.g., from 1 to 3 or 4 times) with independently
5 selected H, halo, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl,
alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid
sidechain, amino acid and peptide.

In some embodiments of **Formula (VI)(i)(a)**, R^{1a} and R^{1b} are each H, and R⁶ is aryl or
heteroaryl.

10 **C. Microbicides and plant defense activators.**

In some embodiments, an active compound described herein is applied in combination
with a microbicide. "Microbicide" as used herein refers to a substance with the ability to kill or
to inhibit the growth of microorganisms (e.g., bacteria, fungal cells, protozoa, etc.), which
microbicide is not an active compound in the group herein disclosed of triazole derivatives.

15 Common microbicides used for microbial control in plants include copper compounds. Examples
of copper compounds include, but are not limited to, Bordeaux mixture, copper hydroxide,
copper oxychloride, copper sulfate, cuprous oxide, mancozeb or oxine-copper. However,
microorganisms (e.g., bacteria such as Xanthomonas and Pseudomonas) may become resistant to
treatment with copper.

20 In some embodiments, resistant microorganisms (e.g., copper-resistant bacteria) are
rendered more susceptible to a microbicides and/or the effectiveness of treatment with a
microbicides is enhanced upon application in combination with an active compound described
herein (e.g., fruit or vegetable yield is increased as compared to diseased plant producing the
fruit or vegetable that is untreated or treated only with the microbicide).

25 Other microbicides include, but are not limited to, azoles such as azaconazole, bitertanol,
propiconazole, difenoconazole, diniconazole, cyproconazole, epoxiconazole, fluquinconazole,
flusilazole, flutriafol, hexaconazole, imazalil, imibenconazole, ipconazole, tebuconazole,
tetraconazole, fenbuconazole, metconazole, myclobutanil, perfurazate, penconazole,
bromuconazole, pyrifenoxy, prochloraz, triadimefon, triadimenol, triflumizole or triticonazole;
30 pyrimidinyl carbinols such as ancymidol, fenarimol or nuarimol; 2-amino-pyrimidine such as
bupirimate, dimethirimol or ethirimol; morpholines such as dodemorph, fenpropidin,
fenpropimorph, spiroxamin or tridemorph; anilinopyrimidines such as cyprodinil, pyrimethanil
or mepanipyrim; pyrroles such as fenpiclonil or fludioxonil; phenylamides such as benalaxyl,
furalaxyl, metalaxyl, R-metalaxyl, ofurace or oxadixyl; benzimidazoles such as benomyl,
35 carbendazim, debacarb, fuberidazole or thiabendazole; dicarboximides such as chlozolinate,

dichlozoline, iprodione, myclozoline, procymidone or vinclozolin; carboxamides such as carboxin, fenfuram, flutolanil, mepronil, oxycarboxin or thifluzamide; guanidines such as guazatine, dodine or iminoctadine; strobilurines such as azoxystrobin, kresoxim-methyl, metominostrobin, SSF-129, methyl 2[(2-trifluoromethyl)-pyrid-6-yloxy-methyl]-3-methoxy-
 5 acrylate or 2-[$\{\alpha[(\alpha\text{-methyl-3-trifluoromethyl-benzyl})\text{imino}]\text{-oxy}\}\text{-o-tolyl}\}$]-glyoxylic acid-methylester-O-methyloxime (trifloxystrobin); dithiocarbamates such as ferbam, mancozeb, maneb, metiram, propineb, thiram, zineb or ziram; N-halomethylthio-dicarboximides such as captafol, captan, dichlofluanid, fluoromide, folpet or tolyfluanid; nitrophenol derivatives such as dinocap or nitrothal-isopropyl; organo phosphorous derivatives such as edifenphos, iprobenphos,
 10 isoprothiolane, phosdiphen, pyrazophos or toclofos-methyl; and other compounds of diverse structures such as acibenzolar-S-methyl, harpin, anilazine, blasticidin-S, chinomethionat, chloroneb, chlorothalonil, cymoxanil, dichione, diclomezine, dicloran, diethofencarb, dimethomorph, dithianon, etridiazole, famoxadone, fenamidone, fentin, ferimzone, fluazinam, flusulfamide, fenhexamid, fosetyl-aluminium, hymexazol, kasugamycin, methasulfocarb,
 15 pencycuron, phthalide, polyoxins, probenazole, propamocarb, pyroquilon, quinoxifen, quintozone, sulfur, triazoxide, tricyclazole, triforine, validamycin, (S)-5-methyl-2-methylthio-5-phenyl-3-phenylamino-3,5-di-hydroimidazol-4-o- ne (RPA 407213), 3,5-dichloro-N-(3-chloro-1-ethyl-1-methyl-2-oxopropyl)-4-methylbenzamide (RH-7281), N-allyl-4,5-dimethyl-2-trimethylsilylthiophene-3-carboxamide (MON 65500), 4-chloro-4-cyano-N,N-dimethyl-5-p-
 20 tolylimidazole-1-sulfon-amide (IKF-916), N-(1-cyano-1,2-dimethylpropyl)-2-(2,4-dichlorophenoxy)-propionamide (AC 382042) or iprovalicarb (SZX 722).

An "antibiotic" as used herein is a type of "microbicide." Common antibiotics include aminoglycosides, carbacephems (e.g., loracarbef), carbapenems, cephalosporins, glycopeptides (e.g., teicoplanin and vancomycin), macrolides, monobactams (e.g., aztreonam) penicillins,
 25 polypeptides (e.g., bacitracin, colistin, polymyxin B), quinolones, sulfonamides, tetracyclines, etc. Antibiotics treat infections by either killing or preventing the growth of microorganisms. Many act to inhibit cell wall synthesis or other vital protein synthesis of the microorganisms.

Aminoglycosides are commonly used to treat infections caused by Gram-negative bacteria. Examples of aminoglycosides include, but are not limited to amikacin, gentamicin,
 30 kanamycin, neomycin, netilmicin, streptomycin, tobramycin, and paromomycin.

Carbapenems are broad-spectrum antibiotics, and include, but are not limited to, ertapenem, doripenem, imipenem/cilastatin, and meropenem.

Cephalosporins include, but are not limited to, cefadroxil, cefazolin, cefalotin (cefalothin), cefalexin, cefaclor, cefamandole, cefoxitin, cefprozil, loracarbef, cefuroxime,

cefixime, cefdinir, cefditoren, cefoperazone, cefotaxime, cefpodoxime, ceftazidime, ceftibuten, ceftizoxime, ceftriaxone, cefepime, cefpirome, and ceftobiprole.

Macrolides include, but are not limited to, azithromycin, clarithromycin, dirithromycin, erythromycin, roxithromycin, troleandomycin, telithromycin and spectinomycin.

5 Penicillins include, but are not limited to, amoxicillin, ampicillin, azlocillin, bacampicillin, carbenicillin, cloxacillin, dicloxacillin, flucloxacillin, mezlocillin, meticillin, nafcillin, oxacillin, penicillin, piperacillin and ticarcillin.

10 Quinolones include, but are not limited to, ciprofloxacin, enoxacin, gatifloxacin, gemifloxacin, levofloxacin, lomefloxacin, moxifloxacin, norfloxacin, ofloxacin and trovafloxacin.

Sulfonamides include, but are not limited to, mafenide, prontosil, sulfacetamide, sulfamethizole, sulfanilamide, sulfasalazine, sulfisoxazole, trimethoprim, and co-trimoxazole (trimethoprim-sulfamethoxazole).

15 Tetracyclines include, but are not limited to, demeclocycline, doxycycline, minocycline, oxytetracycline and tetracycline.

Other antibiotics include arsphenamine, chloramphenicol, clindamycin, lincomycin, ethambutol, fosfomycin, fusidic acid, furazolidone, isoniazid, linezolid, metronidazole, mupirocin, nitrofurantoin, platensimycin, pyrazinamide, quinupristin/dalfopristin, rifampin (rifampicin), tinidazole, etc.

20 Other microbicides that may be used in combination with the active compounds of the present invention include bacteriophages (bacterial viruses) such as Bacillus. Examples of bacteriophage microbicides include, but are not limited to, AgriPhageTM (OmniLytics, Inc., Salt Lake City, Utah) and Serenade® (AgraQuest, Davis, California). See, e.g., U.S. Patent Nos. 5,919,447 and 6,077,506 to Marrone et al.; U.S. Patent No. 6,103,228 to Heins et al.; and U.S.
25 Patent Application Publication 20080152684.

In some embodiments, an active compound described herein is applied in combination with a plant defense activator. A “plant defense activator” as used herein is a compound that improves disease resistance by activating a plant’s natural defense mechanisms, e.g., induces the plant to produce disease-fighting compounds. Examples of plant defense activators include, but
30 are not limited to, prohexadione-calcium (Apogee), Cropset (plant booster element complex), probenazole, potassium phosphate (e.g., ProPhyt®, Helena Chemical Company), harpin protein (e.g., Messenger®, Eden Biosciences Ltd, Bothell, WA), acibenzolar or acibenzolar-S-methyl (e.g., ActigardTM, Syngenta Crop Production, Inc, Greensboro, NC), streptomycin sulfate, reynoutria sachalinensis extract (reysa), etc.

D. Agrochemical compositions.

Active compounds of the present invention can be used to prepare agrochemical compositions in like manner as other antimicrobial compounds. See, e.g., U.S. Pat. Application 2006/0094739; see also U.S. Pat. Nos. 6,617,330; 6,616,952; 6,569,875; 6,541,500, and 6,506,794.

Active compounds described herein can be used for protecting plants against diseases that are caused by microorganisms, including biofilm-forming microorganisms. The active compounds can be used in the agricultural sector and related fields as active ingredients for controlling plant pests. The active compounds can be used to inhibit or destroy the pests that occur on plants or parts of plants (fruit, blossoms, leaves, stems, tubers, roots) of different crops of useful plants, optionally while at the same time protecting also those parts of the plants that grow later e.g. from phytopathogenic microorganisms.

Active compounds may be used as dressing agents for the treatment of plant propagation material, in particular of seeds (fruit, tubers, grains) and plant cuttings (e.g. rice), for the protection against fungal infections as well as against phytopathogenic fungi occurring in the soil.

The active compounds can be used in the form of compositions and can be applied to the crop area or plant to be treated, simultaneously or in succession with further compounds. These further compounds can be e.g. fertilizers or micronutrient donors or other preparations which influence the growth of plants. They can also be selective herbicides as well as insecticides, fungicides, bactericides, nematocides, molluscicides, plant growth regulators, plant activators or mixtures of several of these preparations, if desired together with further carriers, surfactants or application promoting adjuvants customarily employed in the art of formulation.

Suitable carriers and adjuvants can be solid or liquid and are substances useful in formulation technology, e.g. natural or regenerated mineral substances, solvents, dispersants, wetting agents, tackifiers, thickeners, binders or fertilizers.

The active compounds are used in unmodified form or, preferably, together with the adjuvants conventionally employed in the art of formulation. To this end they are conveniently formulated in known manner to emulsifiable concentrates, coatable pastes, directly sprayable or dilutable solutions, dilute emulsions, wettable powders, soluble powders, dusts, granulates, and also encapsulations e.g. in polymeric substances. As with the type of the compositions, the methods of application, such as spraying, atomizing, dusting, scattering, coating or pouring, are chosen in accordance with the intended objectives and the prevailing circumstances.

The formulation, i.e. the compositions containing the active compound and, if desired, a solid or liquid adjuvant, are prepared in known manner, typically by intimately mixing and/or

grinding the compound with extenders, e.g. solvents, solid carriers and, optionally, surface active compounds (surfactants).

Suitable carriers and adjuvants may be solid or liquid and correspond to the substances ordinarily employed in formulation technology, such as, e.g. natural or regenerated mineral substances, solvents, dispersants, wetting agents, tackifiers, thickeners, binding agents or fertilizers. Such carriers are for example described in WO 97/33890.

Further surfactants customarily employed in the art of formulation are known to the expert or can be found in the relevant literature.

The agrochemical formulations will usually contain from 0.1 to 99% by weight, preferably from 0.1 to 95% by weight, of a compound described herein, 99.9 to 1% by weight, preferably 99.8 to 5% by weight, of a solid or liquid adjuvant, and from 0 to 25% by weight, preferably from 0.1 to 25% by weight, of a surfactant.

Whereas it is preferred to formulate commercial products as concentrates, the end user will normally use dilute formulations.

The compositions may also contain further adjuvants such as stabilizers, antifoams, viscosity regulators, binders or tackifiers as well as fertilizers, micronutrient donors or other formulations for obtaining special effects.

E. Methods of use.

Target crops or plants to be treated with active compounds and compositions of the invention typically comprise the following species of plants: cereal (wheat, barley, rye, oat, rice, maize, sorghum and related species); beet (sugar beet and fodder beet); pomes, drupes and soft fruit (apples, pears, plums, peaches, almonds, cherries, strawberries, raspberries and blackberries); leguminous plants (beans, lentils, peas, soybeans); oil plants (rape, mustard, poppy, olives, sunflowers, coconut, castor oil plants, cocoa beans, groundnuts); cucumber plants (pumpkins, cucumbers, melons); fiber plants (cotton, flax, hemp, jute); citrus fruit (oranges, lemons, grapefruit, mandarins); vegetables (spinach, lettuce, asparagus, cabbages, carrots, onions, tomatoes, potatoes, paprika); lauraceae (avocado, cinnamon, camphor) or plants such as tobacco, nuts, coffee, eggplants, sugar cane, tea, pepper, vines including grape-bearing vines, hops, bananas, pineapple, turf and natural rubber plants, as well as ornamentals (flowers, shrubs, broad-leaved trees and evergreens, such as conifers). This list does not represent any limitation.

1. Bacterial infections. The methods, active compounds and compositions can be used to treat bacterial infections in a variety of plants, with specific examples including but not limited to those set forth below.

Citrus. In citrus trees (including orange, lemon, lime, and grapefruit) active compounds and compositions as described herein can be used to treat or control a variety of microbial

diseases, including but not limited to canker (caused by *Xanthomonas campestris* or *Xanthomonas axonopodis* infection), bacterial spot (caused by *Xanthomonas campestris* pv. *Citrumelo* infection); Black Pit(fruit) (caused by *Pseudomonas syringae* infection); Blast (caused by *Pseudomonas syringae* infection) citrus variegated chlorosis (caused by *Xylella fastidiosa* infection), and Citrus Huanglongbing (HLB) caused by *Candidatus Liberibacter asiaticus*.

Pome Fruit. In pome fruits (including apple, pear, quince, Asian pear, and loquat), active compounds and compositions as described herein can be used to treat or control a variety of microbial infections, including but not limited to Fire Blight (caused by *Erwinia amylovora* infection), Crown Gall (caused by *Agrobacterium tumefaciens* infection); Blister spot (caused by *Pseudomonas syringae* infection) and Hairy root (caused by *Agrobacterium rhizogenes* infection).

Peppers. In pepper plants, active compounds and compositions as described herein can be used to treat or control a variety of microbial infections, including but not limited to: Bacterial Spot (caused by *Xanthomonas campestris* pv. *vesicatoria* infection); Bacterial wilt (caused by *Ralstonia solanacearum* infection), and Syringae seedling blight and leaf spot (caused by *Pseudomonas syringae* infection).

Tomatoes. In tomato plants, active compounds and compositions as described herein can be used to treat or control a variety of microbial infections, including but not limited to: Bacterial canker (caused by *Clavibacter michiganensis*), Bacterial speck (caused by *Pseudomonas syringae*), Bacterial spot (caused by *Xanthomonas campestris vesicatoria*), Bacterial stem rot and fruit rot (caused by *Erwinia carotovora*), Bacterial wilt (caused by *Ralstonia solanacearum*), Pith necrosis (caused by *Pseudomonas corrugate*), and Syringae leaf spot (caused by *Pseudomonas syringae*).

Soybeans. In soybeans, active compounds and compositions as described herein can be used to treat or control a variety of microbial infections, including but not limited to: Bacterial blight (caused by *Pseudomonas amygdale*), Bacterial pustules (caused by *Xanthomonas axonopodis* pv. *Glycines*), and Bacterial wilt (caused by *Ralstonia solanacearum* or *Curtobacterium flaccumfaciens*).

Corn, Cotton, Wheat and Rice. In corn, cotton, wheat and rice, active compounds and compositions as described herein can be used to treat or control a variety of microbial infections, including but not limited to: bacterial blights, leaf spots and leaf streak caused by *Xanthomonas* species; bacterial sheath rot, stripe and spot caused by *Pseudomonas* species; and to bacterial stalk and top rot, wilt, foot rot, pink seed and lint degradation caused by *Erwinia* species.

Pineapple. In pineapple, active compounds and compositions as described herein can be used to treat or control a variety of microbial infections, including but not limited to: Bacterial heart rot and Fruit collapse (caused by *Erwinia chrysanthemi*), Bacterial fruitlet brown rot (caused by *Erwinia ananas*), Marbled fruit and Pink fruit (caused by *Erwinia herbicola*), Soft rot (caused by *Erwinia carotovora*), and Acetic souring (caused by *Acetic acid bacteria*).

The above listing is but a sampling, and active compounds and compositions as described herein may also be used to treat or control bacteria (some of which are named above) in a variety of plants. For example, the bacteria *Xylella fastidiosa* infects citrus trees as noted above (citrus variegated chlorosis), and also infects grapevines (Pierce's disease). Other plant hosts of *Xylella fastidiosa* include, but are not limited to, ornamentals, oleander (leaf scorch), almond, coffee, maple, mulberry, elm, sycamore, alfalfa, etc. Similarly, *Ralstonia solanacearum* infects soybeans (bacterial wilt) as well as banana (Moko disease), tobacco (Granville wilt), geranium (southern bacterial wilt), potato (brown rot) and a wide variety of other plants, including ginger and mulberry.

2. Fungal infections. In addition to treating or controlling bacterial infections, active compounds and compositions as described herein can be used to treat or control fungal infections such as rots, leaf molds, blights, wilts, damping-off, spot, root rot, stem rot, mildew, brown spot, gummosis, melanose, post-bloom fruit drop, scab, alternaria, canker, flyspeck, fruit blotch, dieback, downy mildews, ear rots, anthracnose bunts, smut, rust, eyespot and pecky rice. Genera of plant-pathogenic fungi that can be treated or controlled by the active compounds, compositions, and methods described herein include but are not limited to: *Pythium* spp., *Fusarium* spp., *Rhizoctonia* spp., *Cercospora* spp., *Alternaria* spp., *Colletotrichum* spp., *Ustilago* spp., *Phoma* spp., *Gibberella* spp., *Penicillium* spp., *Glomerella* spp., *Diplodia* spp., *Curvularia* spp., *Sclerospora* spp., *Peronosclerospora* spp., *Cercospora* spp., *Puccinia* spp., *Ustilago* spp., *Aspergillus* spp., *Phomopsis* spp., *Diaporthe* spp., *Botrytis* spp., *Verticillium* spp., *Phytophthora* spp.

Particular fungal infections that can be treated or controlled by the methods, compounds and compositions described herein, in vegetables and greenhouse crops, include *Phytophthora* blight (caused by *Phytophthora capsici*) and *Pythium* damping-off (caused by *Pythium* spp.).

Note that *Phytophthora* also has adverse effects on crops ranging from pineapples to cotton. It can kill woody citrus seedlings and young citrus trees (oranges, grapefruits, lemons, limes). In the greenhouse, germinating seed and seedlings are very susceptible to damping-off caused by *Phytophthora*, *Pythium*, *Sclerotinia* and *Rhizoctonia* species. The cost to the grower to lose his crop to any of these fungi is substantial. The loss can happen at transplant time or when the crop is ready to be harvested.

The problems of fungi are not restricted to traditional crops but also extend to forestry products and have worldwide scope. *Phytophthora cinnamomi* is a soil-borne water mould that leads to a condition in plants called “root rot” or “dieback.” *P. cinnamomi* causes root rot affecting woody ornamentals including azalea, dogwood, forsythia, Fraser fir, hemlock, Japanese holly, juniper, rhododendron, white pine, and American chestnut. *P. cinnamomi* is responsible for the destruction of the elegant American chestnut tree. In Australia, *P. cinnamomi* has spread through the forests of western Australia, and into coastal forests of Victoria, where entire plant ecosystems are being obliterated. Given that *P. cinnamomi* is a soil-borne water mould that infects the roots, almost the entire action takes place below ground. This problem highlights the importance of developing new compounds to counter fungal infections, even those that directly affect only the roots of the plant rather than the more visible effects on fruits or vegetables.

Active compounds of the invention can be applied to plants or plant loci in accordance with known techniques. The compound(s) can be tank mixed with other agricultural, turf, ornamental nursery, forestry and all other plant-labeled compatible pesticides. The compound(s) can be applied to seed. The compound(s) can be applied to edible and non-edible crops. The compound(s) can be applied to roots and all other parts of all plants. The compound(s) can be applied in greenhouses. The compound(s) can be applied and used in food-processing facilities. The compound(s) can be applied to plastic food bags and containers. The compound(s) can be applied as a solid, as its free base, or as a salt. The salts can include, but are not limited to, HI, HCl, HBr, H₂SO₄, acetic acid, and trifluoroacetic acid. The compound(s) can be applied as a solution from 0.0001% to 99.9%. The compound(s) can be applied as a solid or solution with copper-based fungicidal compounds. The compound(s) can be applied with specific additional active agents, including but not limited to bactericides, fungicides, pesticides, biological insecticides and microbial insecticides.

Application can be carried out with any suitable equipment or technique, such as: Aerial – Fixed wing and Helicopter; Ground Broadcast Spray – Boom or boomless system, pull-type sprayer, floaters, pick-up sprayers, spray coupes, speed sprayers, and other broadcast equipment, water wagons and water bags; Low pressure boom sprayers, High pressure sprayers; Air blast sprayers; Low volume air sprayers (mist blowers); Ultra-low volume sprayers (ULV); Aerosol Generators (foggers); Dusters; Soil Injector; Hand-Held or High-Volume Spray Equipment – knapsack and backpack sprayers, pump-up pressure sprayers, hand guns, motorized spray equipment; Selective Equipment—Recirculating sprayers, shielded and hooded sprayers; Controlled droplet applicator (CDA) hand-held or boom-mounted applicators that produce a spray consisting of a narrow range of droplet size; Any and all greenhouse sprayers; Micro-sprinkler or drip irrigation systems; Chemigation.

One method of applying an active compound of the invention, or an agrochemical composition which contains at least one of said compounds, is foliar application. The frequency of application and the rate of application will depend on the risk of infestation by the corresponding pathogen. However, the active compounds can also penetrate the plant through the roots via the soil (systemic action) by drenching the locus of the plant with a liquid formulation, or by applying the compounds in solid form to the soil, e.g. in granular form (soil application). In crops of water such as rice, such granulates can be applied to the flooded rice field. The active compounds may also be applied to seeds (coating) by impregnating the seeds or tubers either with a liquid formulation of the fungicide or coating them with a solid formulation.

The term locus as used herein is intended to embrace the fields on which the treated crop plants are growing, or where the seeds of cultivated plants are sown, or the place where the seed will be placed into the soil. The term seed is intended to embrace plant propagating material such as cuttings, seedlings, seeds, and germinated or soaked seeds.

Advantageous rates of application are normally from 5 g to 2 kg of active ingredient (a.i.) per hectare (ha), preferably from 10 g to 1 kg a.i./ha, most preferably from 20 g to 600 g a.i./ha. When used as seed drenching agent, convenient dosages are from 10 mg to 1 g of active substance per kg of seeds.

F. Combination treatments.

In some embodiments, methods of enhancing the effects of a microbicide (such as a microbicide comprising copper, e.g., Kocide® 2000 or Kocide® 3000 (DuPont™, with active ingredient copper hydroxide) are disclosed, comprising the step of applying an active compound in combination with a microbicide, the active compound being applied in an amount effective to enhance the effects of the microbicide.

In some embodiments, methods of enhancing the effects of a plant defense activator are disclosed, comprising the step of applying an active compound in combination with a plant defense activator, the active compound being applied in an amount effective to enhance the effects of the plant defense activator.

“Enhancing” the effects of a microbicide by applying an active compound in combination with the microbicide refers to increasing the effectiveness of the microbicide, such that the microorganism killing and/or growth inhibition is higher at a certain concentration of the microbicide applied in combination with the active compound than without. In some embodiments, a bacteria or other microorganism is “sensitized” to the effects of a microbicide, such that the bacteria or other microorganism that was resistant to the microbicide prior to applying the active compound (e.g., little to none, or less than 20, 10, 5 or 1% are killed upon

application) is rendered vulnerable to that microbicide upon or after applying the active compound (e.g., greater than 20, 30, 40, 50, 60, 70, 80, 90, or 95% or more are killed).

Similarly, "enhancing" the effects of a plant defense activator by applying an active compound in combination with the plant defense activator refers to increasing the effectiveness of the plant defense activator, such that the microorganism killing and/or growth inhibition is higher at a certain concentration of the plant defense activator applied in combination with the active compound than without. In some embodiments, a bacteria or other microorganism is "sensitized" to the effects of a plant defense activator, such that the bacteria or other microorganism that was resistant to the effects of the plant defense activator prior to applying the active compound (e.g., little to none, or less than 20, 10, 5 or 1% are killed upon application) is rendered vulnerable to the effects of that plant defense activator upon or after applying the active compound (e.g., greater than 20, 30, 40, 50, 60, 70, 80, 90, or 95% or more are killed).

As used herein, the application of two or more compounds (inclusive of active compounds and microbicides) "in combination" means that the two compounds are applied closely enough in time that the application of or presence of one alters the biological effects of the other. The two compounds may be applied simultaneously (concurrently) or sequentially.

Simultaneous application of the compounds may be carried out by mixing the compounds prior to application, or by applying the compounds at the same point in time but at different sites of the plant or using different types of applications, or applied at times sufficiently close that the results observed are indistinguishable from those achieved when the compounds are applied at the same point in time.


Sequential application of the compounds may be carried out by applying, e.g., an active compound at some point in time prior to application of a microbicide, such that the prior application of active compound enhances the effects of the microbicide (e.g., percentage of microorganisms killed and/or slowing the growth of microorganisms). In some embodiments, an active compound is applied at some point in time prior to the initial application of a microbicide. Alternatively, the microbicide may be applied at some point in time prior to the application of an active compound, and optionally, applied again at some point in time after the application of an active compound.

EXAMPLES

EXAMPLE 1: Synthesis of 2-aminoimidazole-triazole (2-AIT) chemical library.

There is a paucity of reactions that have been reported to be compatible with 2-aminoimidazoles. To test the applicability of the Cu(I)-catalyzed [3+2] alkyne/azide cycloaddition (Click reaction, see Kolb et al., *Angewandte Chemie-International Edition* **2001**, 11, 2004-2021; Rodionov et al., *Angewandte Chemie-International Edition* **2005**, 15, 2210-2215), we synthesized the alkyne


derived 2-aminoimidazole **1** and tested its ability to participate in a Cu(I)-catalyzed [3+2] cycloaddition with benzyl azide.



Scheme 1. Synthesis of alkyne 2-aminoimidazole **1**. a) 5% Na/Hg, H₂O, pH = 1.5-2.0, 0-5 °C. b) cyanamide, H₂O, pH = 4.3, 95 °C.

The alkyne derived 2-aminoimidazole (2-AI) was synthesized as outlined in Scheme 1. Amino acid **2** (Kotha et al., *Tetrahedron* **2002**, 45, 9203-9208) was subjected to small scale Akabori reduction (Akabori, *Berichte Der Deutschen Chemischen Gesellschaft* **1933**, 66, 151-158), which, followed by condensation with cyanamide (Xu Yz et al., *J Org Chem* **1997**, 3, 456-464) delivered the target alkyne 2-AI **1** in 88% yield. With **1** in hand, we explored various conditions to elicit the Cu-catalyzed [3+2] cycloaddition between **1** and benzyl azide (**Table 1**).

Table 1



Scale (mg)	Cu(I) Source ^a	Solvent	Base	Temp	Yield
20	CuI	THF	DIEA	RT	NR
20	CuI	THF	DIEA	40 °C	NR
20	CuSO ₄ /NaAsc	EtOH/H ₂ O (1:1)	—	RT	NR
20	CuSO ₄ /NaAsc	EtOH/H ₂ O (1:1)	—	40 °C	86%
100	CuSO ₄ /NaAsc	EtOH/H ₂ O (1:1)	—	40 °C	Decomp
100	CuSO ₄ /NaAsc	<i>t</i> -BuOH/H ₂ O/CH ₂ Cl ₂ (1:1:1)	—	RT	93%

^a NaAsc = Sodium Ascorbate

Reactions in THF using Cu(I) yielded no reaction and only returned starting material. We then switched to using CuSO₄ and sodium ascorbate in a 1:1 solvent mixture of H₂O/EtOH. Again, no reaction was noted. However, when the reaction was heated to 40 °C we noted clean conversion to the desired 2-AIT conjugate **3** in 86% yield. Unfortunately, when the reaction was

scaled up, we observed a significant amount of decomposition. Room temperature click reactions have been noted when a 1:1:1 solvent mixture of H₂O/EtOH/CH₂Cl₂ (Lee et al., *Tetrahedron Letters* **2006**, 29, 5105-5109) is employed instead of the 1:1 H₂O/EtOH mixture. When these reaction conditions were tested, we observed conversion to **3** in 93% yield.


5 With the methodology established to access 2-AIT conjugates, we employed the synthetic approach outlined in Scheme 1 to synthesize 2-AI alkynes **4** and **5** in which we systematically extended the methylene space between the alkyne and the 2-AI. The click reaction was then performed between each of the 2-AI alkynes and 12 azides to yield an initial 2-AIT library (shown below). Each compound was characterized (¹H NMR, ¹³C NMR, HRMS).

10 In conclusion, we have developed a synthetic approach to access 2-aminoimidazole/triazole (2-AIT) conjugates that is underpinned by the Cu(I)-catalyzed [3+2] alkyne/azide cycloaddition. Using this chemistry we have assembled a focused library of 2-AIT conjugates.


1. Experimental Protocols for 2-AIT Conjugate Synthesis

15 All reagents used for chemical synthesis were purchased from commercially available sources and used without further purification. Chromatography was performed using 60 Å mesh standard grade silica gel from Sorbtech (Sorbent Technologies, Inc., Atlanta, Georgia). NMR solvents were obtained from Cambridge Isotope Laboratories, Inc. (Andover, Massachusetts) and used as received. ¹H NMR (300 MHz or 400 MHz) and ¹³C NMR (75 MHz or 100 MHz) spectra
20 were recorded at 25°C on Varian Mercury spectrometers. Chemical shifts (δ) are given in ppm relative to tetramethylsilane or the respective NMR solvent; coupling constants (*J*) are in hertz (Hz). Abbreviations used are s = singlet, bs = broad singlet, d = doublet, dd = doublet of doublets, t = triplet, dt = doublet of triplets, bt = broad triplet, qt = quartet, m = multiplet, bm =broad multiplet and br = broad. High and low resolution mass spectra were obtained at the
25 North Carolina State Mass Spectrometry Laboratory for Biotechnology. FAB experiments were carried out with a JOEL HX110HF mass spectrometer while ESI experiments were carried out on an Agilent LC-TOF mass spectrometers.

Chemical Library:



Synthesis:




5


To a 50 mL round-bottomed flask equipped with a magnetic stirbar was added 3-furan methanol (1.00 g, 10.2 mmol) and a solution of diphenyl phosphoryl azide (3.37 g, 12.2 mmol) in toluene (30 mL). The stirring solution is allowed to cool to 0°C, in which 1, 8 Diazabicyclo [5. 4. 0.] undec-7-ene (1.86 g, 12.2 mmol) was added dropwise. The reaction is allowed to slowly warm to ambient temperature for an additional 16 hours of stirring. After this period, the reaction mixture is washed with water (2 x 20 mL) and then with 5% HCl (20 mL). Volatiles are evaporated under reduced pressure. The resulting residue is then purified by column chromatography (1:9 ethyl acetate/hexane) providing 3-azidomethyl furan (1.19 g, 95%) as a colorless oil. ¹H NMR (400 MHz, CDCl₃) δ 7.46 (d, 1H), δ 7.44 (s, 1H), δ 6.42 (d, 1H), δ 4.20 (s, 2H). ppm; ¹³C NMR (75 MHz, CDCl₃) δ 144.1, 141.1, 110.4, 92.1, 45.8 ppm; LRMS (EI) calcd

15


for $C_5H_5N_3O$ (M^+) 123, found 123.




Following the same procedure used to synthesize 3-azidomethyl furan, indole-3-methanol (2.00 g, 13.6 mmol) was converted to 3-azidomethyl indole (1.31 g, 56%) as a yellow oil. ^1H NMR (300 MHz, $CDCl_3$) δ 8.19 (bs, 1H), δ 7.71 (d, 1H), δ 7.39 (m, 2H), δ 7.19 (m, 2H), δ 4.54 (s, 2H) ppm; ^{13}C NMR (75 MHz, $CDCl_3$) δ 130.3, 125.9, 125.3, 122.2, 120.3, 120.3, 119.7, 118.7, 111.9 ppm; LRMS (EI) calcd for $C_9H_8N_4$ (M^+) 172, found 172.




Following the same procedure used to synthesize 3-azidomethyl furan, furfuryl alcohol (2.50 g, 25.5 mmol) was converted to 2-azidomethyl furan (2.96 g, 95%) as a colorless oil. ^1H NMR (300 MHz, $CDCl_3$) δ 7.43 (d, 1H), δ 6.36 (m, 2H), δ 4.29 (s, 2H) ppm; ^{13}C NMR (75 MHz, $CDCl_3$) δ 148.2, 110.7, 109.6, 47.2 ppm; LRMS (EI) calcd for $C_5H_5N_3O$ (M^+) 123, found 123.




Following the same procedure used to synthesize 3-azidomethyl furan, thiophene-3-methanol (3.14 g, 27.6 mmol) was converted to 3-azidomethyl thiophene (3.72 g, 97%) as a colorless oil. ^1H NMR (300 MHz, $CDCl_3$) δ 7.38 (d, 1H), δ 7.23 (s, 1H), δ 7.10 (d, 1H), δ 4.36 (s, 2H) ppm; ^{13}C (75 MHz, $CDCl_3$) δ 136.4, 127.6, 127.1, 124.0, 49.9 ppm; LRMS (EI) calcd for $C_5H_5N_3S$ (M^+) 139, found 139.




2-chloroethyl-5,6-dimethyl-1H-benzimidazole was synthesized through the treatment of 4,5-dimethyl-1,2-phenylenediamine to conditions outlined by Hortelano (Raban et al., Org. Chem. **1985**, 50 (13), 2205–2210). The resulting product was transformed to 2-azidomethyl-5,6-dimethyl-1H-benzimidazole following conditions outlined by Hankovszky resulting in a yellow solid (Hiales et al., Synthesis **1978**, 4, 313–315). ^1H NMR (300 MHz, $CDCl_3$) δ 7.26 (s, 2H), δ 4.72 (s, 2H), δ 2.37 (s, 6H) ppm; ^{13}C NMR (75 MHz, DMSO) δ 153.7, 134.3, 130.1, 125.1, 124.6, 117.2, 113.6, 48.0, 20.6, 19.6 ppm; HRMS (FAB) calcd for $C_{10}H_{11}N_5$ (M^+) 201.1014, found 201.1010.




To a 100 mL round-bottomed flask equipped with a magnetic stir was added trans-2-methyl-3-phenyl-2-propen-1-ol (2.00 g, 13.5 mmol) and 75 mL of methylene chloride. The solution was then cooled to 0°C while stirring. Then, triethylamine (2.75 g, 27.0 mmol) is added followed by a dropwise addition of methanesulfonyl chloride (2.34 g, 20.4 mmol) and a two hour stir period. The reaction mixture is washed with water (2 x 75 mL), dried with sodium sulfate and then concentrated de vacuo. The crude mixture is then dissolved in 75 mL of DMF and then stirred via magnetic stir bar. To this mixture, sodium azide (1.76 g, 27.0 mmol) is added. The reaction mixture is then heated to 80°C and allowed to stir for two hours. At this time, volatiles are concentrated de vacuo and the resulting residue is purified via column chromatography (1:9 ethyl acetate/hexane) providing (3-Azido-2-methylpropenyl)-benzene (2.08 g, 89%) as a colorless oil. ¹H NMR (300 MHz, CDCl₃) δ 7.36 – 7.25 (m, 5H), δ 6.53 (s, 1H), δ 3.87 (s, 2H) ppm; ¹³C NMR (75 MHz, CDCl₃) δ 129.4, 129.2, 128.9, 128.6, 128.4, 127.1, 59.9, 52.0, 22.3, 16.5 ppm; LRMS (EI) calcd for C₁₀H₁₁N₃ (M⁺) 173, found 173.




Following the same procedure used to synthesize (3-Azido-2-methylpropenyl) benzene, thiophene-3-ethanol (2.00 g, 15.5 mmol) was converted to 3-azidoethyl-thiophene (2.06 g, 86%) as a colorless oil. ¹H NMR (300 MHz, CDCl₃) δ 7.36 (s, 1H), δ 7.14 (d, 1H), δ 7.04 (d, 1H), δ 3.54 (t, 2H), δ 2.98 (t, 2H) ppm; ¹³C NMR (75 MHz, CDCl₃) δ 138.7, 128.4, 126.3, 122.2, 52.0, 30.1 ppm; LRMS (EI) calcd for C₆H₇N₃S (M⁺) 153, found 153.




2-Amino-hex-5-ynoic acid methyl ester hydrochloride was synthesized using the same methods previously reported for the synthesis of 2-Amino-pent-4-ynoic acid methyl ester hydrochloride (Kotha et al., Tetrahedron **2002**, 58, 9203–9208). ¹H NMR (300 MHz, D₂O) δ 4.16 (t, 1H), δ 3.71 (s, 3H), δ 2.32 (t, 1H), δ 2.29 (m, 2H), δ 2.06 (m, 2H) ppm; ¹³C NMR (75 MHz, D₂O) δ 170.4, 82.4, 71.4, 53.9, 52.1, 28.7, 14.4 ppm; HRMS (ESI) calcd for C₇H₁₁NO₂ (M⁺) 142.0859, found 142.862.




2-Amino-hept-6-ynoic acid methyl ester hydrochloride was synthesized using the same methods previously reported for the synthesis of 2-Amino-pent-4-ynoic acid methyl ester hydrochloride (Kotha et al., Tetrahedron **2002**, 58, 9203–9208). ¹H NMR (300 MHz, DMSO) δ 8.75 (s, 2H), δ 3.99 (m, 1H), δ 3.72 (s, 3H), δ 2.82 (t, 1H), δ 2.17 (m, 2H), δ 1.89 (m, 2H), δ 1.51 (m, 2H) ppm; ¹³C NMR (75 MHz, DMSO) δ 163.7, 83.1, 69.2, 60.9, 30.6, 29.5, 23.9, 17.6 ppm; HRMS (ESI) calcd for C₈H₁₃NO₂ (M⁺) 156.1019, found 156.1017.



2-Amino-oct-7-ynoic acid methyl ester hydrochloride was synthesized using the same methods previously reported for the synthesis of 2-Amino-pent-4-ynoic acid methyl ester hydrochloride (Kotha et al., Tetrahedron **2002**, 58, 9203–9208). ¹H NMR (300 MHz, D₂O) δ 4.16 (t, 1H), δ 3.84 (s, 3H), δ 2.35 (t, 1H), δ 2.24 (m, 2H), δ 1.95 (m, 2H), δ 1.52 (m, 4H) ppm; ¹³C NMR (75 MHz, D₂O) δ 170.9, 85.6, 69.6, 53.6, 52.9, 29.3, 27.0, 23.4, 17.3 ppm; HRMS (ESI) calcd for C₉H₁₅NO₂ (M⁺) 170.1176, found 170.1171.




2-Amino-non-8-ynoic acid methyl ester hydrochloride was synthesized using the same methods previously reported for the synthesis of 2-Amino-pent-4-ynoic acid methyl ester hydrochloride (Kotha et al., Tetrahedron **2002**, 58, 9203–9208). ¹H NMR (300 MHz, D₂O) δ 4.21 (t, 1H), δ 3.90 (s, 3H), δ 2.41 (t, 1H), δ 2.28 (m, 2H), δ 2.01 (m, 2H), δ 1.51 (m, 6H) ppm; ¹³C NMR (75 MHz, D₂O) δ 171.1, 86.4, 69.4, 53.7, 53.1, 29.8, 27.5, 27.4, 23.8, 17.6 ppm; HRMS (ESI) calcd for C₁₀H₁₈NO₂ (M⁺) 184.1332, found 184.1329.




2-Amino-dec-9-ynoic acid methyl ester hydrochloride was synthesized using the same


methods previously reported for the synthesis of 2-Amino-pent-4-ynoic acid methyl ester hydrochloride (Kotha et al., Tetrahedron **2002**, 58, 9203–9208). ^1H NMR (300 MHz, D_2O) δ 4.16 (t, 1H), δ 3.86 (s, 3H), δ 2.36 (t, 1H), δ 2.21 (m, 2H), δ 1.98 (m, 2H), δ 1.55 – 1.39 (m, 8H) ppm; ^{13}C NMR (75 MHz, D_2O) δ 171.2, 86.7, 69.3, 53.7, 53.1, 29.8, 27.7, 27.7, 27.6, 24.1, 17.6 ppm; HRMS (ESI) calcd for $\text{C}_{11}\text{H}_{20}\text{NO}_2$ (M^+) 198.1488, found 198.1488.




2-Amino-pent-4-ynoic acid methyl ester hydrochloride (2.91 g, 17.8 mmol) was treated to an Akabori reduction followed by a cyanamide condensation employing conditions previously reported to produce 4-Prop-2-ynyl-1H-imidazol-2-ylamine hydrochloride (1.65 g, 59%) as a yellow oil (Olofson et al., Journal of Organic Chemistry **1997**, 62, (23), 7918–7919). ^1H NMR (300 MHz, CD_3OD) δ 6.30 (s, 1H), δ 5.02 (d, 2H), δ 2.26 (t, 1H) ppm; ^{13}C NMR (75 MHz, CD_3OD) δ 150.1, 127.2, 109.6, 83.6, 69.8, 15.9 ppm; HRMS (ESI) calcd for $\text{C}_6\text{H}_7\text{N}_3$ (M^+) 122.0712, found 122.0713.




2-Amino-hex-5-ynoic acid methyl ester hydrochloride (2.53 g, 14.2 mmol) was treated to an Akabori reduction followed by a cyanamide condensation employing conditions previously reported to produce 4-But-3-ynyl-1H-imidazol-2-ylamine hydrochloride (1.17 g, 48%) as a pale yellow oil (Olofson et al., Journal of Organic Chemistry **1997**, 62, (23), 7918–7919). ^1H NMR (300 MHz, CD_3OD) δ 6.52 (s, 1H), δ 2.61 (t, 2H), δ 2.42 (m, 2H), δ 2.27 (t, 1H) ppm; ^{13}C NMR (75 MHz, CD_3OD) δ 147.4, 126.2, 109.4, 81.9, 69.9, 23.8, 17.4 ppm; HRMS (ESI) calcd for $\text{C}_7\text{H}_{10}\text{N}_3$ (M^+) 136.0869, found 136.0865.




2-Amino-hept-6-ynoic acid methyl ester hydrochloride (2.00 g, 10.4 mmol) was treated to an Akabori reduction followed by a cyanamide condensation employing conditions previously reported to produce 4-Pent-4-ynyl-1H-imidazol-2-ylamine hydrochloride (1.75 g, 90%) as a pale oil (Olofson et al., Journal of Organic Chemistry **1997**, 62, (23), 7918–7919). ^1H NMR (300 MHz, CDCl_3) δ 6.68 (bs, 2H), δ 6.24 (s, 1H), δ 2.51 (t, 2H), δ 2.17 (t, 1H), δ 1.95 (s, 1H), δ 1.74 (m, 2H) ppm; ^{13}C NMR (75 MHz, CDCl_3) δ 148.3, 132.7, 111.6, 84.4, 69.1, 28.0, 26.0, 18.2 ppm; HRMS (ESI) calcd for $\text{C}_{10}\text{H}_{16}\text{N}_3$ (M^+) 150.1026, found 150.1029.




2-Amino-oct-7-ynoic acid methyl ester hydrochloride (2.90 g, 14.1 mmol) was treated to an Akabori reduction followed by a cyanamide condensation employing conditions previously reported to produce 4-Hex-5-ynyl-1H-imidazol-2-ylamine hydrochloride (2.45 g, 87%) as a pale yellow solid (Olofson et al., Journal of Organic Chemistry **1997**, 62, (23), 7918–7919). ¹H NMR (300 MHz, CD₃OD) δ 6.43 (s, 1H), δ 2.44 (t, 2H), 2.14 (t, 1H), δ 2.12 (m, 2H), δ 1.64 (m, 2H), δ 1.47 (m, 2H) ppm; ¹³C NMR (75 MHz, CD₃OD) δ 147.3, 127.6, 108.5, 83.4, 68.7, 27.7, 27.1, 23.8, 17.5 ppm; HRMS (ESI) calcd for C₉H₁₄N₃ (M⁺) 164.1182, found 164.1182.



2-Amino-non-8-ynoic acid methyl ester hydrochloride (2.02g, 9.20mmol) was treated to an Akabori reduction followed by a cyanamide condensation employing conditions previously reported to produce 4-Hept-6-ynyl-1H-imidazol-2-ylamine hydrochloride (1.04 g, 53%) as a pale yellow solid (Olofson et al., Journal of Organic Chemistry **1997**, 62, (23), 7918–7919). ¹H NMR (300 MHz, CD₃OD) δ 6.17 (s, 1H), δ 2.19 (t, 2H), δ 1.90 (t, 1H), δ 1.86 (m, 2H), δ 1.24 – 1.13 (m, 6H) ppm; ¹³C NMR (75 MHz, CD₃OD) δ 147.5, 128.3, 108.8, 83.8, 65.5, 28.4, 28.3, 28.2, 24.5, 17.8 ppm; HRMS (ESI) calcd for C₁₀H₁₆N₃ (M⁺) 178.1338, found 178.1337.




2-Amino-dec-9-ynoic acid methyl ester hydrochloride (1.50 g, 6.42 mmol) was treated to an Akabori reduction followed by a cyanamide condensation employing conditions previously reported to produce 4-Oct-7-ynyl-1H-imidazol-2-ylamine hydrochloride (0.774 g, 53%) as a pale yellow solid (Olofson et al., Journal of Organic Chemistry **1997**, 62, (23), 7918–7919). ¹H NMR (400 MHz, CD₃OD) δ 6.09 (s, 1H), δ 2.19 (t, 2H), δ 1.95 (t, 1H), 1.93 (m, 2H), δ 1.39 – 1.11 (m, 8H) ppm; ¹³C NMR (75 MHz, CD₃OD) δ 148.5, 131.0, 110.1, 83.9, 68.3, 28.6, 28.5, 28.4, 28.3, 25.7, 17.8 ppm; HRMS (ESI) calcd for C₁₁H₁₈N₃ (M⁺) 192.1495, found 192.1495.




To a 50 mL round-bottomed flask equipped with a magnetic stirbar was added 1-H-1, 2, 3-triazole (0.192 g, 2.78 mmol) and DMF (10 mL) and then cooled to 0°C while stirring. Then,

sodium hydride (60% dispersion in mineral oil) (0.133 g, 3.33 mmol) is added to the reaction mixture and was slowly allowed to warm to ambient temperature. Then, 1-iodo-4-pentyne (0.647 g, 3.33 mmol) was added dropwise. The reaction mixture was then heated to 80°C and allowed to stir for 2.5 hours. Water (20 mL) was then added to the reaction mixture and then extracted with ethyl acetate (2 x 20 mL). The organic phase was dried with sodium sulfate and concentrated *de vacuo* followed by a purification by column chromatography (ethyl acetate/hexane) to produce 1-Pent-4-ynyl-1H-[1,2,3]triazole (0.349 g, 93%) as a colorless oil. ¹H NMR (300 MHz, CDCl₃) δ 7.67 (s, 1H), δ 7.59 (s, 1H), δ 4.53 (t, 2H), δ 2.20 (t, 2H), δ 2.17 (m, 2H), δ 2.04 (s, 1H) ppm; ¹³C NMR (75 MHz, CDCl₃) δ 133.9, 123.9, 82.2, 70.4, 48.7, 28.9, 15.7 ppm; HRMS (ESI) calcd for C₇H₁₀N₃ (M⁺) 136.0869, found 136.0866.

General procedure for click reactions: The terminal alkyne (1.0 equiv.) was dissolved in a 1:1:1 mixture of *tert*-butyl alcohol, water and methylene chloride (ca. 10 mL per 0.300 g of terminal alkyne). To this solution, the appropriate azide (1.2 equiv.) was added while stirring vigorously at room temperature. Copper (II) sulfate pentahydrate (15 mol%) and sodium ascorbate (45 mol%) were then added sequentially to the solution. Reaction mixtures were allowed to stir until completion via TLC analysis (12 – 24 hrs). The solvents were then removed *de vacuo* in which the resulting residue was dissolved in methanol and purified by flash chromatography (10 – 20% ammonia saturated methanol: methylene chloride). The resulting fractions were evaporated under reduced pressure followed by a 24hr high vacuum treatment to remove all ammonia traces. Methanol saturated with HCl is then added to the purified product in which all volatiles are then removed under reduced pressure.




4-Prop-2-ynyl-1H-imidazol-2-ylamine hydrochloride (0.127 g, 0.809 mmol) was reacted with (3-Azido-2-methyl-propenyl)-benzene (0.168 g, 0.971 mmol) following the general procedure for click reactions outlined above to produce 4-[1-(2-Methyl-3-phenyl-allyl)-1H [1,2,3]triazol-4-ylmethyl]-1H-imidazol-2-ylamine hydrochloride (0.244 g, 91%) of a pale yellow solid. ¹H NMR (300 MHz, D₂O) δ 7.94 (s, 1H), δ 7.40 – 7.35 (m, 5H), δ 6.56 (s, 1H), δ 5.10 (s, 2H), δ 3.99 (s, 2H), δ 1.76 (s, 3H) ppm; ¹³C δ 145.8, 138.1, 136.9, 136.5, 136.4, 132.8, 128.3, 128.0, 127.6, 125.3, 123.4, 49.3, 23.8, 23.4; HRMS (ESI) calcd for C₁₆H₁₈N₆ (M⁺) 295.1665, found 295.1665.




4-Prop-2-ynyl-1H-imidazol-2-ylamine hydrochloride (0.095 g, 0.603 mmol) was reacted


with benzyl azide (0.096 g, 0.723 mmol) following the general procedure for click reactions outlined above to produce 4-(1-Benzyl-1H-[1,2,3]triazol-4-ylmethyl)-1H-imidazol-2-ylamine hydrochloride (0.151 g, 86%) of a pale yellow solid. ¹H NMR (300 MHz, D₂O) δ 7.79 (s, 1H), δ 7.42 – 7.35 (m, 5H), δ 6.43 (s, 1H), δ 5.58 (s, 2H), δ 3.86 (s, 2H) ppm; ¹³C δ 145.4, 136.6, 136.4, 128.8, 128.7, 126.9, 126.8, 126.3, 124.1, 121.8, 109.6, 53.3, 24.1 ppm; HRMS (ESI) calcd for C₁₃H₁₆N₆O (M⁺) 254.1352, found 254.1352.



4-Prop-2-ynyl-1H-imidazol-2-ylamine hydrochloride (0.101 g, 0.639 mmol) was reacted with 3-azidomethyl-furan (0.094 g, 0.767 mmol) following the general procedure for click reactions outlined above to produce 4-(1-Furan-3-ylmethyl-1H-[1,2,3]triazol-4-ylmethyl)-1H-imidazol-2-ylamine hydrochloride (0.077 g, 43%) of a pale yellow solid. ¹H NMR (300 MHz, CD₃OD) δ 8.12 (s, 1H), δ 7.60 (s, 1H), δ 7.38 (s, 1H), δ 6.53 (s, 1H), δ 6.37 (s, 1H), δ 5.46 (s, 2H), δ 3.95 (s, 2H) ppm; ¹³C NMR (75 MHz, CD₃OD) δ 146.1, 145.3, 144.1, 141.0, 123.3, 123.1, 110.3, 110.2, 109.9, 51.3, 19.9 ppm; HRMS (ESI) calcd for C₁₁H₁₂N₆O (M⁺) 244.1145, found 244.1145.




4-Prop-2-ynyl-1H-imidazol-2-ylamine hydrochloride (0.096 g, 0.061 mmol) was reacted with 2-azidomethyl-furan (0.089 g, 0.729 mmol) following the general procedure for click reactions outlined above to produce 4-(1-Furan-2-ylmethyl-1H-[1,2,3]triazol-4-ylmethyl)-1H-imidazol-2-ylamine hydrochloride (0.078 g, 46%) as a pale yellow solid. ¹H NMR (300 MHz, CD₃OD) δ 7.93 (s, 1H), δ 7.41 (s, 1H), 6.49 (s, 2H), δ 6.32 (s, 1H), δ 5.56 (s, 2H), δ 3.89 (s, 2H) ppm; ¹³C NMR (75 MHz, CD₃OD) δ 146.2, 146.1, 142.4, 141.5, 122.3, 122.2, 108.9, 109.2, 108.8, 51.6, 18.9 ppm; HRMS (ESI) calcd for C₁₁H₁₂N₆O (M⁺) 245.1145, found 245.1147.




4-Prop-2-ynyl-1H-imidazol-2-ylamine hydrochloride (0.096 (0.061 mmol) was reacted with 3-azidomethyl thiophene (0.102 g, 0.732 mmol) following the general procedure for click reactions outlined above to produce 4-(1-Thiophen-3-ylmethyl-1H-[1,2,3]triazol-4-ylmethyl)-1H-imidazol-2-ylamine hydrochloride (0.079 g, 44%) of a pale yellow solid. ¹H NMR (300 MHz, CD₃OD) δ 8.18 (s, 1H), δ 7.44 (s, 1H), δ 7.30 (d, 1H), δ 6.98 (d, 1H), δ 6.53 (s, 1H), δ 5.57 (s, 2H), δ 3.95 (s, 2H) ppm; ¹³C (75 MHz, CD₃OD) δ 145.7, 145.1, 143.7, 139.7, 129.8, 121.8,


111.5, 109.8, 109.2, 52.7, 21.5 ppm; HRMS (ESI) calcd for $C_{11}H_{12}N_6S$ (M^+) 260.0919, found 260.0919.




4-Prop-2-ynyl-1H-imidazol-2-ylamine hydrochloride (0.101 g, 0.641 mmol) was reacted with 3-(2-Azido-ethyl)-thiophene (0.118 g, 0.769 mmol) following the general procedure for click reactions outlined above to produce 4-[1-(2-Thiophen-3-yl-ethyl)-1H [1,2,3]triazol-4-ylmethyl]-1H-imidazol-2-ylamine hydrochloride (0.119 g, 60%) of a pale yellow solid. ^1H NMR (300 MHz, CD_3OD) δ 7.42 (s, 1H), δ 7.19 (t, 1H), δ 6.90 (s, 1H), δ 6.78 (d, 1H), δ 6.09 (s, 1H), δ 4.47 (t, 2H), δ 3.67 (s, 2H), δ 3.09 (t, 2H) ppm; ^{13}C ; (75 MHz, CD_3OD) δ 151.2, 145.7, 137.7, 130.5, 127.8, 125.8, 122.9, 121.9, 110.3, 50.798, 30.7, 23.3 ppm; HRMS (ESI) calcd for $C_{12}H_{14}N_6S$ (M^+) 274.1001, found 274.1007.




4-Prop-2-ynyl-1H-imidazol-2-ylamine hydrochloride (0.047 g, 0.295 mmol) was reacted with 2-azidomethyl-5,6-dimethyl-1H-benzimidazole (0.071 g, 0.354 mmol) following the general procedure for click reactions outlined above to produce 4-[1-(5,6 Dimethyl-1H-benzoimidazol-2-ylmethyl)-1H-[1,2,3]triazol-4-ylmethyl]-1H-imidazol-2-ylamine dihydrochloride (0.029 g, 25%) of a yellow solid. ^1H NMR (300 MHz, CD_3OD) δ 8.09 (s, 1H), δ 7.45 (s, 2H), δ 6.52 (s, 1H), δ 6.08 (s, 2H), δ 3.93 (s, 2H), δ 2.35 (s, 6H) ppm; ^{13}C (75 MHz, CD_3OD) δ 142.2, 136.9, 136.6, 124.5, 124.3, 113.8, 113.4, 110.0, 100.4, 85.8, 80.6, 75.3, 74.1, 45.0, 20.9, 19.28 ppm; HRMS (ESI) calcd for $C_{16}H_{18}N_8$ (M^+) 323.1727, found 323.1734.




4-Prop-2-ynyl-1H-imidazol-2-ylamine hydrochloride (0.091 g, 0.576 mmol) was reacted with 2-azidomethyl-1H-benzimidazole, which was synthesized using previously reported methods (Hiales et al., Synthesis **1978**, 4, 313-315), (0.120 g, 0.691 mmol) following the general procedure for click reactions outlined above to produce 4-[1-(1H-Benzoimidazol-2-ylmethyl)-1H-[1,2,3]triazol-4-ylmethyl] 1Himidazol-2-ylamine dihydrochloride (0.125 g, 59%) of a yellow solid. ^1H NMR (300 MHz, CD_3OD) δ 8.01 (s, 1H), δ 7.58 (d, 2H), δ 7.31 (t, 2H), δ 6.44 (s, 1H), δ 5.98 (s, 2H), δ 3.84 (s, 2H) ppm; ^{13}C (75 MHz, CD_3OD) δ 135.6, 124.7, 123.7, 123.5, 122.8, 112.8, 112.3, 108.4, 83.6, 78.9, 74.0, 73.2, 44.2, 19.3 ppm; HRMS (ESI) calcd for $C_{14}H_{14}N_8$ (M^+) 295.1414, found 295.1420.




4-But-3-ynyl-1H-imidazol-2-ylamine hydrochloride (0.0735 g, 0.428 mmol) was reacted with (3-azido-propyl) benzene, which was synthesized using previously reported methods (Suenaga et al., Tetrahedron Letters **2003**, 44, 5799–5801), (0.083 g, 0.514 mmol) following the general procedure for click reactions outlined above to produce 4-{2-[1-(3-Phenyl-propyl)-1H-[1,2,3]triazol-4-yl]-ethyl}-1H-imidazol-2-ylamine hydrochloride (0.051 g, 36%) of a pale yellow oil. ¹H NMR (300 MHz, CD₃OD) δ 7.87 (s, 1H), δ 7.32 – 7.22 (m, 5H), δ 6.54 (s, 1H), δ 4.31 (t, 2H), δ 3.05 (t, 2H), δ 2.93 (t, 2H), δ 2.67 (t, 2H), δ 2.23 (m, 2H) ppm; ¹³C (75 MHz CD₃OD) δ 153.3, 146.6, 140.4, 134.3, 128.8, 128.7, 128.6, 126.6, 121.5, 67.5, 53.9, 32.8, 32.9, 29.5 ppm; HRMS (ESI) calcd for C₁₆H₂₀N₆ (M⁺) 296.1822, found 296.1828.




4-But-3-ynyl-1H-imidazol-2-ylamine hydrochloride (0.062 g, 0.362) was reacted with (3-Azido-2-methyl-propenyl)-benzene (0.075 g, 0.434 mmol) following the general procedure for click reactions outlined above to produce 4-{2-[1-(2-Methyl-3-phenyl-allyl)-1H-[1,2,3]triazol-4-yl]-ethyl}-1H-imidazol-2-ylamine hydrochloride (0.054 g, 43%) of a pale yellow oil. ¹H NMR (300 MHz, CD₃OD) δ 8.39 (s, 1H), δ 7.12 – 7.07 (m, 5H), δ 6.54 (s, 1H), δ 6.36 (s, 1H), δ 5.09 (s, 2H), δ 3.01 (t, 2H), δ 2.76 (t, 2H), δ 1.60 (s, 3H) ppm; ¹³C; 147.7, 143.1, 136.3, 132.7, 129.9, 128.9, 128.6, 128.5, 128.2, 127.4, 127.1, 124.9, 109.9, 61.2, 23.0, 22.3, 14.7 ppm; HRMS (ESI) calcd for C₁₆H₂₀N₆ (M⁺) 308.1749, found 308.1742.




4-But-3-ynyl-1H-imidazol-2-ylamine hydrochloride (0.068 g, 0.399 mmol) was reacted with 2-azidomethyl furan (0.059 g, 0.478 mmol) following the general procedure for click reactions outlined above to produce 4-[2-(1-Furan-2-ylmethyl-1H-[1,2,3]triazol-4-yl)-ethyl]-1H-imidazol-2-ylamine hydrochloride (0.085 g, 72%) of a pale yellow solid. ¹H NMR (300 MHz, CD₃OD) δ 8.06 (s, 1H), δ 7.53 (s, 1H), δ 6.59 (t, 1H), δ 6.49 (s, 1H), δ 6.44 (dd, 1H), δ 3.05 (t, 2H), δ 2.89 (t, 2H) ppm; ¹³C (75 MHz, CD₃OD) δ 156.4, 154.9, 147.5, 144.1, 126.0, 125.1, 123.8, 110.7, 109.3, 69.8, 23.8, 17.4 ppm; HRMS (ESI) calcd for C₁₂H₁₄N₆O (M⁺) 259.1301, found 259.1305.




4-But-3-ynyl-1H-imidazol-2-ylamine hydrochloride (0.078 g, 0.45 mmol) was reacted with 3-(2-Azido-ethyl)-thiophene (0.083 g, 0.543 mmol) following the general procedure for click reactions outlined above to produce 4-{2-[1-(2-Thiophen-3-yl-ethyl)-1H-[1,2,3]triazol-4-yl]-ethyl}-1H-imidazol-2-ylamine hydrochloride (0.069 g, 47%) of a pale yellow oil. ¹H NMR (300 MHz, CD₃OD) δ 7.33 (s, 1H), δ 7.19 (dd, 1H), δ 6.85 (d, 1H), δ 6.72 (d, 1H), δ 6.06 (s, 1H), δ 4.45 (t, 2H), δ 3.08 (t, 2H), δ 2.79 (t, 2H), δ 2.59 (t, 2H), ppm; ¹³C (75 MHz, CD₃OD) δ 149.3, 147.0, 137.7, 132.1, 127.7, 125.8, 122.4, 121.9, 110.6, 50.8, 30.7, 26.6, 24.8 ppm; HRMS (ESI) calcd for C₁₃H₁₇N₆S (M⁺) 289.1229, found 289.1231.




4-But-3-ynyl-1H-imidazol-2-ylamine hydrochloride (0.066 g, 0.383 mmol) was reacted with (3-azido-propenyl)-benzene, which was synthesized using previously reported methods (Rad et al., Tetrahedron Letters **2007**, 48, 3445–3449), (0.079 g, 0.460 mmol) following the general procedure for click reactions outlined above to produce 4-{2-[1-(3-Phenyl-allyl)-1H-[1,2,3]triazol-4-yl]-ethyl}-1H-imidazol-2-ylamine hydrochloride (0.049 g, 39%) of a pale yellow oil. ¹H NMR (300 MHz, CD₃OD) δ 8.19 (s, 1H), δ 7.44 (d, 2H), δ 7.29 (m, 3H), δ 6.73 (d, 1H), δ 6.53 (s, 1H), δ 6.42 (m, 2H), δ 5.24 (d, 2H), δ 3.07 (t, 2H), δ 2.88 (t, 2H) ppm; ¹³C 148.6, 145.3, 139.4, 139.2, 129.8, 129.7, 129.6, 128.5, 128.4, 127.6, 54.6, 47.3, 45.9, 26.3, 22.6 ppm; HRMS (ESI) calcd for C₁₆H₁₉N₆ (M⁺) 295.1665, found 295.1670.




4-But-3-ynyl-1H-imidazol-2-ylamine hydrochloride (0.066 g, 0.384 mmol) was reacted with 3-azidomethyl-furan (0.057 g, 0.461 mmol) following the general procedure for click reactions outlined above to produce 4-{2-[1-(Furan-3-ylmethyl)-1H-[1,2,3]triazol-4-yl]-ethyl}-1H-imidazol-2-ylamine hydrochloride (0.061 g, 54%) as a pale yellow solid. ¹H NMR (300 MHz, CD₃OD) δ 7.70 (s, 1H), δ 7.61 (s, 1H), δ 7.46 (s, 1H), δ 6.42 (s, 1H), δ 6.38 (s, 1H), δ 5.41 (s, 2H), δ 2.95 (t, 2H), δ 2.83 (t, 2H) ppm; ¹³C (75 MHz, CD₃OD) δ 146.3, 144.2, 141.6, 141.5, 126.5, 122.3, 120.2, 109.9, 109.1, 44.8, 24.2, 23.9 ppm; HRMS (ESI) calcd for C₁₂H₁₄N₆O (M⁺) 259.1301, found 259.1306.




4-But-3-ynyl-1H-imidazol-2-ylamine hydrochloride (0.073 g, 0.423 mmol) was reacted with benzyl azide (0.068 g, 0.509 mmol) following the general procedure for click reactions outlined above to produce 4-[2-(1-benzyl-1H-[1,2,3]triazol-4-yl)-ethyl]-1H-imidazol-2-ylamine hydrochloride (0.059 g, 46%) as a pale yellow oil. ^1H NMR (300 MHz, CD_3OD) δ 7.83 (d, 1H), δ 7.37 – 7.24 (m, 5H), δ 6.44 (d, 1H), δ 5.54 (d, 2H), δ 2.96 (m, 2H), δ 2.83 (m, 2H) ppm; ^{13}C (75 MHz, CD_3OD) δ 163.7, 147.3, 146.4, 135.7, 128.8, 128.4, 127.9, 127.8, 126.5, 122.6, 109.1, 53.7, 24.1, 23.9 ppm; HRMS (ESI) calcd for $\text{C}_{14}\text{H}_{16}\text{N}_6$ (M^+) 269.1515, found 269.1513.




4-But-3-ynyl-1H-imidazol-2-ylamine hydrochloride (0.073 g, 0.429 mmol) was reacted with 3-azidomethyl-indole (0.089 g, 0.515 mmol) following the general procedure for click reactions outlined above to produce 4-{2-[1-(1H-Indol-3-ylmethyl)-1H-[1,2,3]triazol-4-yl]-ethyl}-1H-imidazol-2-ylamine hydrochloride (0.086 g, 58%) as a yellow oil. ^1H NMR (300 MHz, CD_3OD) δ 8.11 (s, 1H), δ 7.84 (m, 2H), δ 7.41 (m, 2H), δ 7.06 (d, 1H), δ 5.66 (s, 2H), δ 3.04 (t, 2H), δ 2.85 (t, 2H) ppm; ^{13}C (75 MHz, CD_3OD) δ 147.5, 144.9, 144.8, 134.7, 131.9, 127.2, 127.1, 127.1, 125.8, 125.3, 124.4, 109.5, 49.9, 23.6, 23.2 ppm; HRMS (ESI) calcd for $\text{C}_{17}\text{H}_{19}\text{N}_7$ (M^+) 321.1701, found 321.1704.




4-But-3-ynyl-1H-imidazol-2-ylamine hydrochloride (0.061 g, 0.361 mmol) was reacted with 2-azidomethyl-1H-benzimidazole (0.075 g, 0.433 mmol) following the general procedure for click reactions outlined above to produce 4-{2-[1-(1H-Benzoimidazol-2-ylmethyl)-1H-[1,2,3]triazol-4-yl]-ethyl}-1H-imidazol-2-ylamine dihydrochloride (0.066 g, 48%) of a yellow solid. ^1H NMR (300 MHz, CD_3OD) δ 8.05 (s, 1H), δ 7.68 (m, 2H), δ 7.48 (m, 2H), δ 6.38 (s, 1H), δ 6.15 (s, 2H), δ 2.90 (t, 2H), δ 2.76 (t, 2H) ppm; ^{13}C (75 MHz, CD_3OD) δ 147.4, 146.9, 146.8, 131.3, 127.0, 126.4, 124.1, 114.1, 109.2, 44.7, 24.0, 23.9 ppm; HRMS (ESI) calcd for $\text{C}_{15}\text{H}_{17}\text{N}_8$ (M^+) 309.1570, found 309.1572.




4-But-3-ynyl-1H-imidazol-2-ylamine hydrochloride (0.0734 g, 0.432 mmol) was reacted with 2-azidomethyl-5,6-dimethyl-1H-benzimidazole (0.104 g, 0.518 mmol) following the general procedure for click reactions outlined above to produce 4-{2-[1-(5,6-Dimethyl-1H-benzimidazol-2-ylmethyl)-1H-[1,2,3]triazol-4-yl]-ethyl}-1H-imidazol-2-ylamine dihydrochloride (0.104 g, 59%) of a pale yellow solid. ¹H NMR (300 MHz, CD₃OD) δ 7.70 (s, 1H), δ 7.19 (s, 2H), δ 6.23 (s, 1H), δ 5.66 (s, 2H), δ 2.85 (t, 2H), δ 2.68 (t, 2H), □ 2.21 (s, 6H) ppm; ¹³C (75 MHz, CD₃OD) δ 148.2, 147.0, 146.9, 132.3, 128.9, 122.8, 120.0, 115.1, 109.7, 100.4, 25.1, 24.4, 19.2 ppm; HRMS (ESI) calcd for C₁₇H₂₀N₈ (M⁺) 337.1883, found 337.1886.




4-Pent-4-ynyl-1H-imidazol-2-ylamine hydrochloride (0.050 g, 0.269 mmol) was reacted with (3-azido-propyl) benzene (0.044 g, 0.273 mmol) following the general procedure for click reactions outlined above to produce 4-{3-[1-(3-Phenyl-propyl)-1H-[1,2,3]triazol-4-yl]-propyl}-1H-imidazol-2-ylamine hydrochloride (0.0318 g, 34%) as a pale yellow solid. ¹H NMR (300 MHz, DMSO) δ 7.91 (s, 1H), δ 7.29 – 7.17 (m, 5H), δ 6.62 (s, 2H), δ 6.57 (s, 1H), δ 4.29 (t, 2H), δ 2.59 (t, 2H), δ 2.53 (t, 2H), δ 2.43 (t, 2H) δ 2.09 (m, 2H), δ 1.83 (m, 2H) ppm; ¹³C NMR (75 MHz, DMSO) δ 163.6, 156.2, 155.1, 147.4, 146.9, 141.4, 129.1, 127.1, 126.7, 122.6, 109.4, 49.4, 32.6, 32.0, 28.1, 24.9, 24.2 ppm; HRMS (ESI) calcd for C₁₇H₂₂N₆ (M⁺) 310.1978, found 310.1977.




4-Pent-4-ynyl-1H-imidazol-2-ylamine hydrochloride (0.063 g, 0.340 mmol) was reacted with 3-azidomethyl-furan (0.050 g, 0.406 mmol) following the general procedure for click reactions outlined above to produce 4-[3-(1-Furan-3-ylmethyl-1H-[1,2,3]triazol-4-yl)-propyl]-1H-imidazol-2-ylamine hydrochloride (0.061 g, 58%) as a pale yellow solid. ¹H NMR (300 MHz, CD₃OD) δ 7.70 (s, 1H), δ 7.56 (s, 1H), δ 7.41 (s, 1H), δ 6.44 (s, 1H), δ 6.35 (s, 1H), δ 5.35 (s, 2H), δ 2.65 (t, 2H), δ 2.45 (t, 2H), δ 1.86 (m, 2H) ppm; ¹³C NMR (75 MHz, CD₃OD) δ 147.3, 147.3, 144.2, 141.6, 127.2, 122.1, 120.2, 109.9, 108.8, 44.7, 27.8, 24.2, 23.6 ppm; HRMS (ESI) calcd for C₁₃H₁₆N₆O (M⁺) 272.1458, found 272.1462.




4-Pent-4-ynyl-1H-imidazol-2-ylamine hydrochloride (0.093 g, 0.500 mmol) was reacted with 2-azidomethyl-furan (0.074 g, 0.601 mmol) following the general procedure for click reactions outlined above to produce 4-[3-(1-Furan-2-ylmethyl-1H-[1,2,3]triazol-4-yl)-propyl]-1H-imidazol-2-ylamine hydrochloride (0.075, 50%) as a pale yellow solid. ¹H NMR (300 MHz, DMSO) δ 7.85 (s, 1H), δ 7.65 (s, 1H), δ 6.70 (s, 2H), δ 6.64 (s, 1H), δ 6.52 (t, 1H), δ 6.46 (s, 1H), δ 5.58 (s, 2H), δ 2.61 (t, 2H), δ 2.41 (t, 2H), δ 1.82 (m, 2H) ppm; ¹³C NMR (75 MHz, DMSO) δ 163.6, 149.4, 147.9, 147.4, 144.3, 128.5, 122.6, 111.5, 110.3, 109.9, 46.3, 28.4, 24.9 ppm; HRMS (ESI) calcd for C₁₃H₁₆N₆O (M⁺) 272.1458, found 272.1460.



4-Pent-4-ynyl-1H-imidazol-2-ylamine hydrochloride (0.090 g, 0.486 mmol) was reacted with 3-Azidomethyl-thiophene (0.081 g, 0.582 mmol) following the general procedure for click reactions outlined above to produce 4-[3-(1-Thiophen-3-ylmethyl-1H-[1,2,3]triazol-4-yl)-propyl]-1H-imidazol-2-ylamine hydrochloride (0.0727 g, 46%) as a pale yellow solid. ¹H NMR (300 MHz, CD₃OD) δ 7.61 (s, 1H), δ 7.24 (s, 1H), δ 7.23 (d, 1H), δ 6.87 (d, 1H), δ 6.29 (s, 1H), δ 2.54 (t, 2H), 2.33 (t, 2H), δ 1.75 (m, 2H) ppm; ¹³C NMR (75 MHz, CD₃OD) δ 147.6, 147.4, 147.3, 136.1, 127.9, 127.0, 126.9, 124.2, 122.2, 109.1, 53.1, 27.9, 24.3; HRMS (ESI) calcd for C₁₃H₁₆N₆S (M⁺) 289.1229, found 289.1234.



4-Pent-4-ynyl-1H-imidazol-2-ylamine hydrochloride (0.096 g, 0.517 mmol) was reacted with (3-Azido-2-methyl-propenyl)-benzene (0.110 g, 0.635 mmol) following the general procedure for click reactions outlined above to produce 5-[3-[1-(2-Methyl-3-phenyl-allyl)-1H-[1,2,3]triazol-4-yl]-propyl]-1H-imidazol-2-ylamine hydrochloride (0.076, 41%) as a pale yellow solid. ¹H NMR (300 MHz, CD₃OD) δ 7.65 (s, 1H), δ 7.14 – 7.04 (m, 5H), δ 6.35 (s, 1H), δ 6.28 (s, 1H), δ 4.94 (s, 2H), δ 2.57 (t, 2H), δ 2.34 (t, 2H), δ 1.75 (m, 2H), δ 1.57 (s, 3H) ppm; ¹³C NMR (75 MHz, CD₃OD) δ 147.5, 136.9, 132.5, 129.7, 128.8, 128.6, 128.4, 128.3, 128.2, 126.9, 122.5, 109.2, 58.1, 61.9, 28.0, 24.3, 24.3, 24.1, 14.6 ppm; HRMS (ESI) calcd for C₁₈H₂₂N₆ (M⁺) 323.1978, found 323.1984.




4-Pent-4-ynyl-1H-imidazol-2-ylamine hydrochloride (0.081 g, 0.437 mmol) was reacted with benzyl azide (0.071 g, 0.533 mmol) following the general procedure for click reactions outlined above to produce 4-[3-(1-Benzyl-1H-[1,2,3]triazol-4-yl)-propyl]-1H-imidazol-2-


ylamine hydrochloride (0.073 g, 53%) as a pale yellow solid. ^1H NMR (300 MHz, DMSO) δ 7.93 (s, 1H), δ 7.37 – 7.27 (m, 5H), δ 6.63 (s, 2H), δ 6.44 (s, 1H), δ 2.61 (t, 2H), δ 2.40 (t, 2H), δ 1.82 (m, 2H) ppm; ^{13}C NMR (75 MHz, DMSO) δ 163.5, 148.1, 147.4, 135.9, 129.4, 128.7, 128.5, 128.4, 122.8, 122.8, 115.9, 109.9, 53.3, 28.5, 25.0; HRMS (ESI) calcd for $\text{C}_{15}\text{H}_{18}\text{N}_6$ (M^+) 282.1665, found 282.1674.



4-Pent-4-ynyl-1H-imidazol-2-ylamine hydrochloride (0.090 g, 0.485 mmol) was reacted with 3-(2-Azido-ethyl)-thiophene (0.089 g, 0.581 mmol) following the general procedure for click reactions outlined above to produce 4-{3-[1-(2-Thiophen-3-yl-ethyl)-1H-[1,2,3]triazol-4-yl]-propyl}-1H-imidazol-2-ylamine hydrochloride (0.067 g, 41%) as a pale yellow solid. ^1H NMR (300 MHz, CD_3OD) δ 7.42 (s, 1H), δ 7.12 (d, 1H), δ 6.83 (s, 1H), δ 6.70 (d, 1H), δ 6.29 (s, 1H), δ 4.41 (t, 2H), δ 3.04 (t, 2H), δ 2.50 (t, 2H), δ 2.29 (t, 2H), δ 1.72 (m, 2H) ppm; ^{13}C NMR (75 MHz, CD_3OD) δ 146.8, 146.8, 137.7, 127.8, 127.6, 125.7, 122.6, 121.9, 109.0, 50.7, 30.7, 27.9, 23.7 ppm; HRMS (ESI) calcd for $\text{C}_{14}\text{H}_{18}\text{N}_6\text{S}$ (M^+) 302.1313, found 302.1317.




4-Pent-4-ynyl-1H-imidazol-2-ylamine hydrochloride (0.115 g, 0.620 mmol) was reacted with (3-azido-propenyl)-benzene (0.119 g, 0.748 mmol) following the general procedure for click reactions outlined above to produce 4-{3-[1-(3-Phenyl-allyl)-1H-[1,2,3]triazol-4-yl]-propyl}-1H-imidazol-2-ylamine hydrochloride (0.082 g, 43%) as a pale yellow solid. ^1H NMR (300 MHz, CDCl_3) δ 7.34 – 7.24 (m, 6H), δ 6.67 (d, 1H), δ 6.34 (q, 1H), δ 6.25 (s, 1H), δ 5.08 (d, 2H), δ 2.72 (t, 2H), δ 2.47 (t, 2H), δ 1.91 (m, 2H) ppm; ^{13}C (75 MHz, CD_3OD) 149.1, 137.9, 133.5, 129.5, 128.9, 128.7, 128.3, 128.2, 127.7, 126.8, 54.8, 48.9, 31.3, 29.6, 27.5 ppm; HRMS (ESI) calcd for $\text{C}_{17}\text{H}_{21}\text{N}_6$ (M^+) 308.1822, found 308.1821.




4-Hex-5-ynyl-1H-imidazol-2-ylamine hydrochloride (0.089 g, 0.447 mmol) was reacted with (3-Azido-2-methyl-propenyl)-benzene (0.085 g, 0.491 mmol) following the general procedure for click reactions outlined above to produce 4-{4-[1-(2-Methyl-3-phenyl-allyl)-1H-[1,2,3]triazol-4-yl]-butyl}-1H-imidazol-2-ylamine (0.132 g, 79%) as a pale yellow solid. ^1H NMR (300 MHz, CD_3OD) δ 8.36 (s, 1H), 7.03 (m, 5H), 6.49 (s, 1H), 6.25 (s, 1H), 5.05 (s, 2H), δ 2.67 (t, 2H), δ 2.73 (t, 2H), δ 1.56 (s, 3H), δ 1.42 (m, 4H) ppm; ^{13}C (75 MHz, CD_3OD) δ 147.3, 144.7, 136.4, 132.7, 129.9, 128.9, 128.6, 128.5, 128.3, 127.5, 127.2, 126.9, 108.7, 61.2, 27.4,


27.3, 23.8, 22.8, 14.8 ppm; HRMS (ESI) calcd for C₁₉H₂₄N₆ (M⁺) 336.2135, found 336.2134.



4-Hept-6-ynyl-1H-imidazol-2-ylamine hydrochloride (0.060 g, 0.281 mmol) was reacted with (3-Azido-2-methyl-propenyl)-benzene (0.058 g, 0.336 mmol) following the general procedure for click reactions outlined above to produce 4-{5-[1-(2-Methyl-3-phenyl-allyl)-1H-[1,2,3]triazol-4-yl]-pentyl}-1H-imidazol-2-ylamine hydrochloride (0.064 g, 65%) as a pale yellow solid. ¹H NMR (300 MHz, CD₃OD) δ 8.23 (s, 1H), δ 6.93 – 6.83 (m, 5H), δ 6.38 (s, 1H), δ 6.08 (s, 1H), δ 4.93 (s, 2H), δ 2.51 (t, 2H), δ 2.09 (t, 2H), δ 1.43 (s, 3H), δ 1.41 (m, 2H), δ 1.25 (m, 2H), δ 1.05 (m, 2H) ppm; ¹³C (75 MHz, CD₃OD) δ 147.7, 144.8, 136.3, 132.8, 129.8, 128.9, 128.6, 128.5, 128.2, 127.6, 127.5, 127.0, 108.5, 61.3, 28.0, 27.6, 27.5, 24.0, 22.9, 14.7 ppm; HRMS (ESI) calcd for C₂₀H₂₇N₆ (M⁺) 351.2291, found 351.2291.



4-Oct-7-ynyl-1H-imidazol-2-ylamine hydrochloride (0.098 g, 0.568 mmol) was reacted with (3-Azido-2-methyl-propenyl)-benzene (0.118 g, 0.681 mmol) following the general procedure for click reactions outlined above to produce 4-{6-[1-(2-Methyl-3-phenyl-allyl)-1H-[1,2,3]triazol-4-yl]-hexyl}-1H-imidazol-2-ylamine hydrochloride (0.147 g, 85%) as a pale yellow solid. ¹H NMR (300 MHz, CD₃OD) δ 8.42 (s, 1H), δ 7.25 – 7.15 (m, 5H), δ 6.65 (s, 1H), δ 6.36 (s, 1H), δ 5.23 (s, 2H), δ 2.78 (t, 2H), δ 2.38 (t, 2H), δ 1.74 (s, 3H), δ 1.64 (m, 2H), δ 1.51 (m, 2H), δ 1.33 (m, 4H) ppm; ¹³C (75 MHz, CD₃OD) δ 147.5, 136.3, 132.9, 132.8, 129.9, 128.9, 128.6, 128.5, 128.3, 127.7; HRMS (ESI) calcd for C₂₁H₂₉N₆ (M⁺) 365.2448, found 365.2448.



1-Pent-4-ynyl-1H-[1,2,3]triazole (0.100 g, 0.739 mmol) was reacted with (3-Azido-2-methylpropenyl)-benzene (0.154 g, 0.889 mmol) following the general procedure for click reactions outlined above to produce 1-(2-Methyl-3-phenyl-allyl)-4-(3-[1,2,3]triazol-1-yl-propyl)-1H-[1,2,3]triazole (0.227 g, Quantitative) as a white solid. ¹H NMR (300 MHz, DMSO) δ 7.91 (s, 1H), δ 7.77 (s, 2H), δ 7.40 – 7.25 (m, 5H), δ 6.47 (s, 1H), δ 5.05 (s, 2H), δ 4.48 (t, 2H), δ 2.62 (t, 2H), δ 2.50 (t, 2H), δ 2.21 (m, 2H), δ 1.74 (s, 3H) ppm; ¹³C (75 MHz, DMSO) δ 146.4, 137.1, 134.8, 133.6, 132.8, 129.4, 129.3, 129.2, 128.9, 127.6, 123.0, 57.9, 54.0, 32.9, 29.6, 22.7, 16.2 ppm; HRMS (ESI) calcd for C₁₇H₂₁N₆ (M⁺) 309.1822, found 309.1821.

EXAMPLE 2: Activity testing of 2-AIT library members. A standard crystal violet reporter assay is employed to assess the effect of compounds from the 2-amino on the formation of biofilms. Among others, the following strains are tested:

Xanthomonas Xccl

5 *Xanthomonas* Xcv.135

Xanthomonas Xcv5

Xanthomonas Xccp

Xanthomonas Xcp60

Xanthomonas Xcp25

10 *Ralstonia solanacearum* K66




Xanthomonas is a Gram-negative rod-shaped bacterium that is a common plant pathogen. *Xanthomonas* bacteria grow almost exclusively in plants. *Xanthomonas* species testing includes *X. vesicatoria* (crop=tomato), *X. euvesicatoria* (crop=pepper), *X. campestris* (crop=crucifers, particularly cabbage), *X. zinniae* (crop=zinnia), and *X. fragariae* (crop=strawberry). *Ralstonia*
15 *solanacearum* is a Gram-negative bacterium that is found in soil.

Bacteria are allowed to form biofilms in a multi-well plate in the absence or presence of one or more compounds. Planktonic (or free growing) bacteria are then removed, wells washed vigorously, and crystal violet added. Crystal violet stains the remaining bacteria which, following ethanol solubilization, is quantitated by spectrophotometry (A_{540}). Time-dependent
20 and concentration-dependent analysis of each compound are performed.


EXAMPLE 3: Activity testing of 2-AIT library members on *Xanthomonas*. Biofilm formation on PVC microtiter wells was accomplished using *Xanthomonas* strains Xcv 135 (known to infect peppers and tomatoes) and Xcv 5 (known to infect tomatoes but not peppers) as models. The Starting Optical Density (OD at 600 nm) for biofilm attachment assay was 0.55, the
25 temperature for this assay was 28 °C, the duration of incubation was 6 hours under, and the assay was static.

Biofilm inhibition results are as follows for screens with Xcv 135. The Xcv 5 strain is tested in the same manner.

Compound Screened	% Inhibition at 20 μ M (vs. Xcv 135)
30 Formula (II)(a)(5)(D)	86 %
Formula (II)(a)(6)(D)	85 %
Formula (II)(i)(a)(2)(J)	0 %



- 5 **EXAMPLE 4:** Activity testing in pepper plants inoculated with *Xanthomonas euvesicatoria* (bacterial spot). To test the effects of the triazole derivatives for plant biofilm inhibition activity, the compound of Formula (II)(a)(5)(D) was used as an exemplary compound (the “biofilm inhibitor/disperser” or “BFI” in the text hereinafter of Example 4).



- 10 The compound was evaluated under field conditions to determine whether it controls or enhances control of bacterial spot (caused by the bacterium *Xanthomonas euvesicatoria*) of pepper when applied alone or in tank-mixtures with a copper (Kocide 3000), an antibiotic (GWN-9350, gentamicin), or a putative plant defense activator (Prophyt). Tests were performed April – July 2008 at the North Carolina Agricultural Research Service Sandhills Research Station,
- 15 Montgomery County.

- EXPERIMENTAL DESIGN and TREATMENT APPLICATIONS: Each treatment consisted of two 7-plant rows running east-west replicated in a four-block completely randomized design. Each 7-plant row was 10.5 ft x 2 ft. This area was used to calculate the quantity of spray material extrapolated for a per acre basis. Plants of bell pepper cultivar ‘X3R-
- 20 Camelot’ were transplanted to the field on April 24, 2008. On May 6, two plants in the south row of each plot of each treatment were inoculated with a suspension of copper-resistant strains of the bacterial pathogen. These plants were destined to serve as the inoculum source for each plot.

Each of the spray test materials was mixed in 1.5 liters of water and applied using a backpack sprayer by making a single pass on each side of the plant row. Treatments were applied weekly starting immediately after the plants had been inoculated May 6 followed by 7 additional weekly sprays for a total of 8 applications through June 24.

DATA: Ratings of foliar disease were started May 22, when symptoms were observed on the inoculated plants and continued weekly through June 25. Disease was most severe on the inoculated plants and then spread to the non-inoculated plants. Data from these two groups (inoculated plants and the non-inoculated) were evaluated separately. The results are expressed as disease progress over time using the calculated Area Under the Disease Progress Curve (AUDPC). Two fruit harvests were done; July 1 and 8. Because there had been plant loss in some plots, the number of plants per plot was counted and yields (number of fruit and weight) were calculated and are reported on a per plant basis per treatment. Results are presented in **Table 2** below.


TABLE 2.

Treatment & rate/acre (based on 100 gal water/acre)	Mean AUDPC for inoculated plants*	Mean AUDPC for entire plot*	Mean number pepper fruit/plant	Mean fruit yield (lb)/plant
1 – NT Check	162 a**	45 a **	6.0 ab **	1.85 ab **
2 – Kocide 3000 (30% MCE) 1.25 lb	123 b	37 ab	6.5 ab	1.86 ab
3 – BFI 0.467 oz (35 mg/L)	113 b	42 a	4.8 b	1.46 b
4 – GWN-9350 (10%) 3.5 lb + GWN-65 1.0 pt	74 c	15 c	8.6 a	2.52 a
5 – ProPhyt 5.0 pt	114 b	37 ab	6.5 ab	2.06 ab
6 – ProPhyt 5.0 pt + BFI 0.467 oz (35 mg/L)	64 c	25 bc	6.3 ab	1.74 b
7 – GWN-9350 + GWN-65 1.0 pt + BFI 0.467 oz (35 mg/L)	41 c	12 c	8.3 a	2.51 a
8 – Kocide 3000 1.25 lb + BFI 0.467 oz (35 mg/L)	62 c	16 c	8.6 a	2.54 a
L.S.D. $\alpha = 0.05$ and probability of > treatment F value	34 0.0001	16 0.0006	2.6 0.0466	0.76 0.0483

* 0 – 9 rating scale with 0 = no disease, 1 = at least 1 diseased leaf, 2 = 1-5% foliage diseased or defoliated, 3 = 6-10%, 4 = 11-15%, 5 = 16-25%, 6 = 26-50%, 7 = 51-75%, 8 = 76-99%, and 9 = 100% leaves diseased or plant completely defoliated.

** Means within a column followed by the same letter do not differ significantly, L.S.D. $\alpha = 0.05$.


EXAMPLE 5: Activity testing in fungus. The compound of Formula (II)(a)(5)(D) was used as an exemplary compound to evaluate the ability of 2-AIT compounds to inhibit the formation of *Candida albicans* biofilms.



5 It was screened at 100 μ M for its ability to inhibit to the formation of *C. albicans* biofilms using a crystal violet reporter assay. Briefly, biofilms were allowed to form for 24 hours in a 96-well microtiter plate in the absence or presence of 100 μ M of the compound. The wells were subsequently washed thoroughly with water to remove free-floating and loosely adherent fungus, and then treated with crystal violet. Crystal violet stains the remaining surface attached fungus
10 (i.e. the biofilm), which following solubilization, can be quantified by spectrophotometry (A_{540}).

From this initial screen, we determined that the exemplary compound was able to inhibit *C. albicans* formation by 12% at 100 μ M. Follow up growth curves at 100 μ M demonstrated that this anti-biofilm activity was non-fungicidal (data not shown).

15 With this initial success of inhibiting fungal biofilms with a 2-AI derivative, we asked the question whether analogue synthesis could deliver alternative 2-AIT derivatives with enhanced anti-biofilm activity in the context of fungal biofilms. Previous work in our lab has demonstrated that 2-AI-based inhibitors of biofilms can be sub-divided into three separate sections: 1) the 2-AI head, 2) the linker region, and 3) the tail region. Structure activity relationship (SAR) data indicates that selectivity and activity can be tuned/enhanced by modification of the tail region.




20

Based upon this data, we synthesized a new pilot library of 2-AI derivatives for anti-biofilm testing in which diversity could be rapidly assembled via substituents off the triazole ring through commercially available carboxylic acids.

25 The synthetic approach to this library is outlined in **Scheme 2**. In our previous synthesis of 2-AIT conjugates, we employed the alkyne-derived 2-AI as a precursor to the Cu^I-mediated [3+2] alkyne/azide cycloaddition (click reaction). Although this reaction worked well, purification of the resulting product was cumbersome due to the use of copious amounts of ammonia saturated methanol for column chromatography. Therefore, we decided to revise the

route by employing a boc-protected 2-AI alkyne that would allow more traditional means of purification (i.e. methanol/dichloromethane columns). The boc-protected scaffold was synthesized from 7-octynoic acid by treatment with oxalyl chloride followed by diazomethane and quenching the resulting α -diazo ketone with HBr to generate the intermediate α -bromo ketone. Cyclization with boc-guanidine then delivered the target 2-AI alkyne **5**.



Scheme 2. Synthesis of triazole substituted 2-AI derivatives. a) i. (COCl)₂, CH₂Cl₂, DMF (cat.), ii. CH₂N₂, Et₂O-CH₂Cl₂ iii. HBr 82 % b) Boc-guanidine, DMF, rt (59 %) c) Azide, CuSO₄, Na Ascorbate, EtOH, CH₂Cl₂, H₂O d) Trifluoroacetic acid, CH₂Cl₂ (58 – 94 % over two steps).

Once **5** had been synthesized, we assembled a diverse array of azido amides to employ in the click reaction to create our pilot library of 2-AIT conjugates. Briefly, 2-bromo-ethylamine was treated with sodium azide to deliver 2-azido-ethylamine, which following acylation (via the respective acid chloride) generated the azido amides for elaboration into the 2-AIT pilot library. Each azido amide was then subjected to the click reaction with the 2-AI **5**. Boc-deprotection (TFA/CH₂Cl₂) followed by counterion exchange (trifluoroacetate for chloride) delivered the target 2-AIT library for anti-biofilm screening.

Each member of the pilot library was assayed at 100 μ M for its ability to inhibit the formation of *C. albicans* biofilms using the crystal violet reporter assay. From this assay, 2-AIT derivatives **7f** and **7m** were determined to be the most potent. Subsequent dose response studies

revealed that **7f** had an IC₅₀ of 2.9 μM while **7m** had an IC₅₀ of 3.3 μM (**Table 3**). Growth curve and colony count analysis of **7f** and **7m** at respective IC₅₀ values demonstrated their antibiofilm activity to be non-fungicidal (data not shown).

Next, we addressed whether **7f** and **7m** could disperse pre-formed *C. albicans* biofilms. *C. albicans* was allowed to establish biofilms in 96-well microtiter plate for 24 hours. Plates were then washed to remove any free floating or loosely adherent fungus. The appropriate 2-AIT (**7f** and **7m**) was then added to each well at 75 μM and the plate was allowed to incubate at 37 °C for 24 hours. Wells were then washed with water and stained with CV to quantify any remaining biofilms. In comparison to biofilms treated with media only, compound **7f** dispersed 56% while **7m** dispersed 62% of the pre-formed biofilm. Once we had established that both compounds could disperse pre-formed biofilms, we quantified this effect by determining **7f** and **7m**'s EC₅₀ value against pre-formed *C. albicans* biofilms. Here, EC₅₀ is defined as the concentration at which the compound will disperse 50% of a pre-formed biofilm. Dose response studies revealed EC₅₀'s of 37.2 μM and 24.7 μM for **7f** and **7m** respectively (**Table 1**). From a medical perspective, molecules that simply inhibit the formation of a biofilm could be used in a prophylactic sense; however, given that a majority of patients already have an established biofilm infection when they seek medical intervention, molecules that are effective against a pre-formed biofilm are more clinically significant.

Once we had established that these next generation 2-AIT conjugates had the ability to inhibit and disperse *C. albicans* biofilms, we addressed whether members of this library would also inhibit and disperse biofilms from *Cryptococcus neoformans*, an opportunistic fungal strain known to infect immunosuppressed patients, especially those with HIV infections. Initial screening of our library showed that **7e** and **7m** had potent anti-biofilm activity against *C. neoformans*. Follow up dose response studies revealed IC₅₀'s of 1.3 μM and 8.0 μM (**Table 3**). Comparison of fungal growth in the presence or absence of either compound indicated that each compound was not fungicidal. Unfortunately, neither of these compounds was able to disperse pre-formed *C. neoformans* biofilms at the concentrations tested.

Table 3. Fungal biofilm inhibition and dispersion.

Compound	IC ₅₀ (μM)	EC ₅₀ (μM)
<i>C. albicans</i>		
7f	2.9 ± 0.7	37.2 ± 5.7
7m	3.3 ± 1.6	24.7 ± 4.5

<i>S. cerevisiae</i>		
7g	2.7 ± 0.1	207.4 ± 7.3
7h	50.4 ± 2.2	353.7 ± 7.3
7j	130.6 ± 16.9	> 400
<i>C. neoformans</i>		
7e	1.3 ± 0.3	-
7m	8.0 ± 3.4	-

Synthesis: All reagents used for chemical synthesis were purchased from commercially available sources and used without further purification. Chromatography was performed using 60 Å mesh standard grade silica gel from Sorbtech. NMR solvents were obtained from Cambridge Isotope Labs and used as is. ¹H NMR (300 MHz or 400 MHz) and ¹³C NMR (75 MHz or 100 MHz) spectra were recorded at 25°C on Varian Mercury spectrometers. Chemical shifts (δ) are given in ppm relative to tetramethylsilane or the respective NMR solvent; coupling constants (*J*) are in hertz (Hz). Abbreviations used are s = singlet, bs = broad singlet, d = doublet, dd = doublet of doublets, t = triplet, dt = doublet of triplets, bt = broad triplet, qt = quartet, m = multiplet, bm = broad multiplet and br = broad.

N-(2-azidoethyl)tetradecanamide: To a 25 mL round-bottomed flask equipped with a magnetic stir bar was added 2-azidoethanamine (0.102 g, 1.18 mmol), DCM (5 mL) and then triethylamine (0.239 g, 2.37 mmol). To this reaction mixture, tetradecanoyl chloride (0.292 g, 1.18 mmol) was added dropwise and allowed to stir at room temperature for 24 hr. Then, the reaction mixture was concentrated de vacuo and then purified via silica gel column chromatography (100% dichloromethane to 1:40 methanol: dichloromethane) to give N-(2-azidoethyl)tetradecanamide (0.245 g, 79 % yield). ¹H NMR (300 MHz, CDCl₃) δ 5.86 (s, 1H), δ 3.42 (s, 4H), δ 2.18 (t, *J* = 7.8 Hz, 2H), δ 1.62 (m, 2H), δ 1.24 (m, 20H), δ 0.87 (t, *J* = 3.9 Hz, 3H) ppm; ¹³C NMR (75 MHz, CDCl₃) δ 173.7, 51.2, 39.1, 36.9, 32.1, 29.9, 29.8, 29.7, 29.6, 29.5, 25.9, 22.9, 14.4 ppm; HRMS (ESI) calcd for C₁₆H₃₂N₄O (M⁺) 296.2576, found 296.2566.

N-(2-azidoethyl)octadec-9-enamide: As in the synthesis of N-(2-azidoethyl)tetradecanamide, octadec-9-enoyl chloride (0.374 g, 1.24 mmol) was reacted with 2-azidoethanamine (0.107 g, 1.24 mmol) and triethylamine (0.252 g, 2.49 mmol) in dichloromethane (5 mL) to give

N-(2-azidoethyl)octadec-9-enamide (0.309 g, 71 % yield). ¹H NMR (300 MHz, CDCl₃) δ 6.30 (s, 1H), δ 5.29 (m, 2H), δ 3.39 (d, *J* = 2.1 Hz, 2H), δ 3.38 (d, *J* = 2.4 Hz, 2H), δ 2.19 (t, *J* = 7.2 Hz, 2H), δ 1.97 (m, 4H), δ 1.59 (t, *J* = 7.5 Hz, 2H), δ 1.26 (m, 20H), δ 0.84 (t, *J* = 6.6 Hz,

3H) ppm; ^{13}C NMR (75 MHz, CDCl_3) δ 173.9, 130.2, 129.9, 51.1, 39.1, 36.8, 32.1, 30.0, 29.9, 29.7, 29.5, 29.4, 29.3, 27.4, 27.3, 25.9, 22.9, 14.3 ppm; HRMS (ESI) calcd for $\text{C}_{20}\text{H}_{38}\text{N}_4\text{O}$ (M^+) 350.3046, found 350.3039.

N-(2-azidoethyl)thiophene-2-sulfonamide: As in the synthesis of N-(2-

5 azidoethyl)tetradecanamide, thiophene-2-sulfonyl chloride (0.218 g, 1.19 mmol) was reacted with 2-azidoethanamine (0.103 g, 1.19 mmol) and triethylamine (0.241 g, 2.39 mmol) in dichloromethane (5 mL) to give N-(2-azidoethyl)thiophene-2-sulfonamide (0.221 g, 80 % yield). ^1H NMR (300 MHz, CDCl_3) δ 7.62 (d, J = 1.2 Hz, 1H), δ 7.59 (d, J = 1.2 Hz, 1H), 7.08 (t, J = 3.9 Hz, 1H), 5.52 (s, 1H), 3.41 (t, J = 5.4 Hz, 2H), 3.17 (q, J = 5.4, 3.8 Hz, 2H) ppm; ^{13}C NMR (75 MHz, CDCl_3) δ 140.6, 132.7, 132.6, 127.9, 50.9, 42.9 ppm; HRMS (ESI) calcd for $\text{C}_6\text{H}_8\text{N}_4\text{O}_2\text{S}_2$ (M^+) 232.0089, found 232.0084.

N-(2-azidoethyl)-5-(dimethylamino)naphthalene-1-sulfonamide: As in the synthesis

of N-(2-azidoethyl)tetradecanamide, 5-(dimethylamino)naphthalene-1-sulfonyl chloride (0.321 g, 1.19 mmol) was reacted with 2-azidoethanamine (0.103 g, 1.19 mmol) and triethylamine
15 (0.241 g, 2.38 mmol) in dichloromethane (5 mL) to give N-(2-azidoethyl)-5-(dimethylamino)naphthalene-1-sulfonamide (0.328 g, 86 % yield). ^1H NMR (300 MHz, CDCl_3) δ 8.52 (d, J = 8.4 Hz, 1H), δ 8.28 (d, J = 8.4 Hz, 1H), δ 8.22 (d, J = 0.9 Hz, 1H), δ 7.53 (t, J = 8.1 Hz, 1H), δ 7.50 (t, J = 7.5 Hz, 1H), δ 7.17 (d, J = 7.8 Hz, 1H), δ 5.58 (t, J = 1.8 Hz, 1H), δ 3.28 (t, J = 5.7 Hz, 2H), δ 3.03 (q, J = 6.3, 5.7 Hz, 2H), δ 2.85 (s, 6H) ppm; ^{13}C NMR (75 MHz, CDCl_3) δ 152.3, 134.8, 130.9, 130.1, 129.7, 128.9, 123.4, 118.9, 115.6, 51.0, 45.6, 42.6 ppm; HRMS (ESI) calcd for $\text{C}_{14}\text{H}_{17}\text{N}_5\text{O}_2\text{S}$ (M^+) 319.1103, found 319.1104.

N-(2-azidoethyl)-2,3,4,5,6-pentamethylbenzenesulfonamide: As in the synthesis of N-

(2-azidoethyl)tetradecanamide, 2,3,4,5,6-pentamethylbenzene-1-sulfonyl chloride (0.304 g, 1.23 mmol) was reacted with 2-azidoethanamine (0.106 g, 1.23 mmol) and triethylamine (0.249 g,
25 2.46 mmol) in dichloromethane (5 mL) to give N-(2-azidoethyl)-2,3,4,5,6-pentamethylbenzenesulfonamide (0.296 g, 81 % yield). ^1H NMR (300 MHz, CDCl_3) δ 5.12 (t, J = 3.9 Hz, 1H) δ 3.35 (t, J = 5.4 Hz, 2H), δ 3.05 (q, J = 6.0, 5.1 Hz, 2H), 2.59 (s, 6H), δ 2.28 (s, 3H), 2.24 (s, 6H) ppm; ^{13}C NMR (75 MHz, CDCl_3) δ 139.9, 136.1, 135.2, 134.2, 51.1, 42.3, 19.2, 17.9, 17.3 ppm; HRMS (ESI) calcd for $\text{C}_{13}\text{H}_{20}\text{N}_4\text{O}_2\text{S}$ (M^+) 296.1307, found 296.1304.

N-(2-azidoethyl)benzenesulfonamide: As in the synthesis of N-(2-

azidoethyl)tetradecanamide, benzenesulfonyl chloride (0.201 g, 1.16 mmol) was reacted with 2-azidoethanamine (0.100 g, 1.16 mmol) and triethylamine (0.235 g, 2.32 mmol) in dichloromethane (5 mL) to give N-(2-azidoethyl)benzenesulfonamide (0.289 g, 84 % yield). ^1H NMR (300 MHz, CDCl_3) δ 7.90 (d, J = 1.8 Hz, 2H), δ 7.55 (m, 3H), δ 5.47 (s, 1H), δ 3.39 (t, J =

5.1 Hz, 2H), δ 3.12 (t, J = 5.4 Hz, 2H) ppm; ^{13}C NMR (75 MHz, CDCl_3) δ 139.8, 133.2, 129.5, 127.2, 50.9, 42.6 ppm; HRMS (ESI) calcd for $\text{C}_6\text{H}_8\text{N}_4\text{O}_2\text{S}$ (M^+) 226.0524, found 226.0523.

(E)-N-(2-azidoethyl)-4-phenylbut-3-enamide: As in the synthesis of N-(2-azidoethyl)tetradecanamide, (E)-4-phenylbut-3-enoyl chloride (0.198 g, 1.19 mmol) was reacted with 2-azidoethanamine (0.114 g, 1.19 mmol) and triethylamine (0.239 g, 2.37 mmol) in dichloromethane (5 mL) to give (E)-N-(2-azidoethyl)-4-phenylbut-3-enamide (0.168 g, 55 % yield). ^1H NMR (300 MHz, CDCl_3) δ 7.68 (d, J = 15.6 Hz, 1H), δ 7.45 (d, J = 3.3 Hz, 2H), δ 7.29 (m, 3H), δ 7.04 (t, J = 4.1 Hz, 1H), δ 6.59 (d, J = 15.9 Hz, 1H), δ 3.55 (q, J = 6.0, 5.1 Hz, 2H), δ 3.47 (t, J = 1.5 Hz, 2H) ppm; ^{13}C NMR (75 MHz, CDCl_3) δ 166.9, 141.6, 134.9, 130.1, 129.1, 128.1, 120.8, 51.1, 39.4 ppm; HRMS (ESI) calcd for $\text{C}_{11}\text{H}_{12}\text{N}_4\text{O}$ (M^+) 216.1011, found 216.1005.

N-(2-azidoethyl)-2-(phenylthio)acetamide: As in the synthesis of N-(2-azidoethyl)tetradecanamide, 2-(phenylthio)acetyl chloride (0.222 g, 1.19 mmol) was reacted with 2-azidoethanamine (0.102 g, 1.19 mmol) and triethylamine (0.240 g, 2.37 mmol) in dichloromethane (5 mL) to give N-(2-azidoethyl)-2-(phenylthio)acetamide (0.179 g, 64 % yield). ^1H NMR (300 MHz, CDCl_3) δ 7.29 (m, 4H), δ 7.18 (m, 2H), δ 3.60 (s, 2H), 3.34 (m, 4H) ppm; ^{13}C NMR (75 MHz, CDCl_3) δ 168.6, 134.8, 129.5, 128.6, 127.1, 50.8, 39.3, 37.7 ppm; HRMS (ESI) calcd for $\text{C}_{10}\text{H}_{12}\text{N}_4\text{OS}$ (M^+) 236.0732, found 236.0729.

N-(2-azidoethyl)palmitamide: As in the synthesis of N-(2-azidoethyl)tetradecanamide, (0.338 g, 1.23 mmol) was reacted with 2-azidoethanamine (0.106 g, 1.23 mmol) and triethylamine (0.249 g, 2.46 mmol) in dichloromethane (5 mL) to give N-(2-azidoethyl)palmitamide (0.336 g, 84 % yield). ^1H NMR (300 MHz, CDCl_3) δ 6.11 (s, 1H), δ 3.41 (s, 4H), δ 2.17 (t, J = 7.5 Hz, 2H), δ 1.60 (m, 2H), δ 1.23 (m, 24H), δ 0.85 (t, 6.3 Hz, 3H) ppm; ^{13}C NMR (75 MHz, CDCl_3) δ 173.9, 51.1, 39.1, 36.9, 32.1, 29.9, 29.8, 29.8, 29.7, 29.6, 29.5, 25.9, 22.9, 14.3 ppm; HRMS (ESI) calcd for $\text{C}_{18}\text{H}_{36}\text{N}_4\text{O}$ (M^+) 324.2889, found 324.2879.

N-(2-azidoethyl)decanamide: As in the synthesis of N-(2-azidoethyl)tetradecanamide, decanoyl chloride (0.251 g, 1.32 mmol) was reacted with 2-azidoethanamine (0.114 g, 1.32 mmol) and triethylamine (0.267 g, 2.64 mmol) in dichloromethane (5 mL) to give N-(2-azidoethyl)decanamide (0.262 g, 83 % yield). ^1H NMR (300 MHz, CDCl_3) δ 6.49 (s, 1H), δ 3.36 (s, 2H), δ 3.35 (s, 2H), δ 2.14 (t, J = 7.2 Hz, 2H), δ 1.46 (m, 2H), δ 1.21 (m, 12H), δ 0.81 (t, J = 6.0 Hz, 3H) ppm; ^{13}C NMR (75 MHz, CDCl_3) δ 174.1, 50.9, 39.1, 36.8, 32.0, 29.7, 29.6, 29.5, 29.4, 25.9, 22.8, 14.3 ppm; HRMS (ESI) calcd for $\text{C}_{12}\text{H}_{24}\text{N}_4\text{O}$ (M^+) 240.1950, found 240.1947.

N-(2-azidoethyl)-2-iodobenzamide: As in the synthesis of N-(2-azidoethyl)tetradecanamide, 2-iodobenzoyl chloride (0.312 g, 1.17 mmol) was reacted with 2-azidoethanamine (0.101 g, 1.17 mmol) and triethylamine (0.237 g, 2.34 mmol) in

dichloromethane (5 mL) to give N-(2-azidoethyl)-2-iodobenzamide (0.272 g, 74 % yield). ¹H NMR (300 MHz, CDCl₃) δ 7.84 (d, J = 7.2 Hz, 1H), δ 7.33 (m, 2H), δ 7.08 (t, J = 3.3 Hz, 1H), δ 6.51 (s, 1H), δ 3.54 (s, 2H), δ 3.53 (s, 2H) ppm; ¹³C NMR (75 MHz, CDCl₃) δ 169.9, 141.9, 140.0, 131.5, 128.4, 128.4, 92.7, 50.8, 39.6 ppm; HRMS (ESI) calcd for C₉H₉IN₄O (M+) 315.9821, found 315.9820.

N-(2-azidoethyl)-4-tert-butylbenzamide: As in the synthesis of N-(2-azidoethyl)tetradecanamide, 4-tert-butylbenzoyl chloride (0.244 g, 1.23 mmol) was reacted with 2-azidoethanamine (0.106 g, 1.23 mmol) and triethylamine (0.249 g, 2.47 mmol) in dichloromethane (5 mL) to give N-(2-azidoethyl)-4-tert-butylbenzamide (0.221 g, 73 % yield). ¹H NMR (300 MHz, CDCl₃) δ 7.73 (d, J = 6.6 Hz, 2H), δ 7.43 (d, J = 9.0 Hz, 2H), δ 6.95 (t, J = 3.9 Hz, 1H), δ 3.59 (q, J = 5.4, 5.7 Hz, 2H), δ 3.50 (t, J = 5.1 Hz, 2H), δ 1.31 (s, 9H) ppm; ¹³C NMR (75 MHz, CDCl₃) δ 168.2, 155.4, 131.4, 127.2, 125.8, 51.1, 39.7, 35.2, 31.4 ppm; HRMS (ESI) calcd for C₁₃H₁₈N₄O (M+) 246.1480, found 246.1479.

N-(2-azidoethyl)-3,5-difluorobenzamide: As in the synthesis of N-(2-azidoethyl)tetradecanamide, 3,5-difluorobenzoyl chloride (0.219 g, 1.24 mmol) was reacted with 2-azidoethanamine (0.107 g, 1.24 mmol) and triethylamine (0.251 g, 2.48 mmol) in dichloromethane (5 mL) to give N-(2-azidoethyl)-3,5-difluorobenzamide (0.280 g, 59 % yield). ¹H NMR (300 MHz, CDCl₃) δ 7.32 (s, 2H), 7.29 (s, 1H), 6.91 (t, J = 2.4 Hz, 1H), 3.58 (q, J = 5.7, 5.1 Hz, 2H), 3.51 (t, J = 4.8 Hz, 2H) ppm; ¹³C NMR (75 MHz, CDCl₃) δ 165.9, 164.7, 161.5, 161.4, 137.6, 110.7, 106.9, 50.7, 39.9 ppm; HRMS (ESI) calcd for C₉H₈F₂N₄O₂S (M+) 226.0666, found 226.0662.

N-(2-azidoethyl)-2,4,6-trichlorobenzamide: As in the synthesis of N-(2-azidoethyl)tetradecanamide, (0.301 g, 1.23 mmol) was reacted with 2-azidoethanamine (0.106 g, 1.23 mmol) and triethylamine (0.249 g, 2.47 mmol) in dichloromethane (5 mL) to give N-(2-azidoethyl)-2,4,6-trichlorobenzamide (0.317 g, 88 % yield). ¹H NMR (300 MHz, CDCl₃) δ 7.27 (2H, s), δ 6.88 (s, 1H), δ 3.54 (s, 2H), δ 3.53 (s, 2H) ppm; ¹³C NMR (75 MHz, CDCl₃) δ 164.4, 135.9, 134.3, 132.9, 128.2, 50.7, 39.4 ppm; HRMS (ESI) calcd for C₉H₇N₄O (M+) 291.9685, found 291.9681.

N-(2-azidoethyl)-2-naphthamide: As in the synthesis of N-(2-azidoethyl)tetradecanamide, 2-naphthoyl chloride (0.237 g, 1.24 mmol) was reacted with 2-azidoethanamine (0.107 g, 1.24 mmol) and triethylamine (0.252 g, 2.49 mmol) in dichloromethane (5 mL) to give N-(2-azidoethyl)-2-naphthamide (0.234 g, 77 % yield). ¹H NMR (300 MHz, CDCl₃) δ 8.69 (s, 1H), δ 7.84 (m, 4H), δ 7.50 (m, 2H), 6.97 (t, J = 4.2 Hz, 1H), δ 3.68 (q, J = 4.8, 1.2 Hz, 2H), 3.56 (t, J = 3.0 Hz, 2H) ppm; ¹³C NMR (75 MHz, CDCl₃) δ 168.3,

135.1, 132.8, 131.5, 129.2, 128.8, 128.0, 127.9, 127.1, 123.8, 51.2, 39.8 ppm; HRMS (ESI) calcd for $C_{13}H_{12}N_4O$ (M^+) 240.1011, found 240.1007.

N-(2-azidoethyl)-4-heptylbenzamide: As in the synthesis of N-(2-azidoethyl)tetradecanamide, 4-heptylbenzoyl chloride (0.289 g, 1.21 mmol) was reacted with 2-azidoethanamine (0.104 g, 1.21 mmol) and triethylamine (0.245 g, 2.42 mmol) in dichloromethane (5 mL) to give N-(2-azidoethyl)-4-heptylbenzamide (0.260 g, 75 % yield). ^1H NMR (300 MHz, $CDCl_3$) δ 7.73 (d, J = 7.8 Hz, 2H), δ 7.19 (d, J = 7.8 Hz, 3H), δ 3.57 (q, J = 5.7, 5.4 Hz, 2H), δ 3.47 (t, J = 5.7 Hz, 2H), δ 2.61 (t, J = 7.5 Hz, 2H), δ 1.61 (m, 2H), δ 1.26 (m, 8H), δ 0.87 (t, J = 6.0 Hz, 3H) ppm; ^{13}C NMR (75 MHz, $CDCl_3$) δ 168.3, 147.3, 131.7, 128.8, 127.4, 50.9, 39.7, 36.1, 32.0, 31.4, 29.4, 29.3, 22.9, 14.3 ppm; HRMS (ESI) calcd for $C_{16}H_{24}N_4O$ (M^+) 288.1950, found 288.1943.

N-(2-azidoethyl)-4-butylbenzamide: As in the synthesis of N-(2-azidoethyl)tetradecanamide, (0.242 g, 1.22 mmol) was reacted with 2-azidoethanamine (0.105 g, 1.22 mmol) and triethylamine (0.246 g, 2.43 mmol) in dichloromethane (5 mL) to give N-(2-azidoethyl)-4-butylbenzamide (0.224 g, 75 % yield). ^1H NMR (300 MHz, $CDCl_3$) δ 7.73 (d, J = 8.4 Hz, 2H), δ 7.39 (t, J = 1.8 Hz, 1H), δ 7.17 (d, J = 8.1 Hz, 2H), δ 3.55 (q, J = 8.7, 5.7 Hz, 2H), δ 3.45 (t, J = 5.7 Hz, 2H), δ 2.60 (t, J = 7.8 Hz, 2H), δ 1.56 (m, 2H), δ 1.30 (m, 2H), δ 0.90 (t, J = 4.2 Hz, 3H) ppm; ^{13}C NMR (75 MHz, $CDCl_3$) δ 168.5, 147.2, 131.7, 128.8, 127.4, 50.9, 39.7, 35.7, 33.5, 22.5, 14.1 ppm; HRMS (ESI) calcd for $C_{13}H_{18}N_4O$ (M^+) 246.1481, found 246.1476.

N-(2-azidoethyl)-4-hexylbenzamide: As in the synthesis of N-(2-azidoethyl)tetradecanamide, 4-hexylbenzoyl chloride (0.293 g, 1.30 mmol) was reacted with 2-azidoethanamine (0.112 g, 1.30 mmol) and triethylamine (0.264 g, 2.61 mmol) in dichloromethane (5 mL) to give N-(2-azidoethyl)-4-hexylbenzamide (0.299 g, 84 % yield). ^1H NMR (300 MHz, $CDCl_3$) δ 7.75 (d, J = 8.1 Hz, 2H), δ 7.43 (t, J = 3.9 Hz, 1H), δ 7.17 (d, J = 7.8 Hz, 2H), δ 3.55 (q, J = 6.3, 5.7 Hz, 2H), δ 3.46 (t, J = 5.7 Hz, 2H), δ 2.59 (t, J = 7.5 Hz, 2H), δ 1.58 (m, 2H), δ 1.29 (m, 4H), δ 0.87 (t, J = 6.9 Hz, 3H) ppm; ^{13}C NMR (75 MHz, $CDCl_3$) δ 168.6, 147.3, 131.7, 130.3, 128.8, 127.5, 50.9, 39.7, 35.9, 31.6, 31.1, 22.7, 14.2 ppm; HRMS (ESI) calcd for $C_{15}H_{22}N_4O$ (M^+) 274.1794, found 274.1789.

N-(2-azidoethyl)-4-pentylbenzamide: As in the synthesis of N-(2-azidoethyl)tetradecanamide, 4-pentylbenzoyl chloride (0.283 g, 1.35 mmol) was reacted with 2-azidoethanamine (0.116 g, 1.35 mmol) and triethylamine (0.273 g, 2.69 mmol) in dichloromethane (5 mL) to give N-(2-azidoethyl)-4-pentylbenzamide (0.267 g, 76 % yield). ^1H NMR (300 MHz, $CDCl_3$) δ 7.74 (d, J = 8.1 Hz, 2H), δ 7.61 (t, J = 5.1 Hz, 1H), δ 7.15 (d, J = 8.4 Hz, 2H), δ 3.54 (q, J = 5.7, 5.1 Hz, 2H), δ 3.44 (t, J = 5.7 Hz, 2H), δ 2.58 (t, J = 7.5 Hz, 2H), δ 1.57 (m, 2H), δ 1.26 (m, 4H), δ 0.86 (t, J = 6.6 Hz, 3H) ppm; ^{13}C NMR (75 MHz, $CDCl_3$) δ

168.6, 147.2, 131.7, 128.7, 127.5, 50.8, 39.7, 35.9, 31.6, 31.1, 22.7, 14.2 ppm; HRMS (ESI) calcd for $C_{14}H_{20}N_4O$ (M⁺) 260.1637, found 260.1632.

N-(2-azidoethyl)biphenyl-4-carboxamide: As in the synthesis of N-(2-azidoethyl)tetradecanamide, (0.283 g, 1.31 mmol) was reacted with 2-azidoethanamine (0.113 g, 1.31 mmol) and triethylamine (0.264 g, 2.61 mmol) in dichloromethane (5 mL) to give N-(2-azidoethyl)biphenyl-4-carboxamide (0.297 g, 85 % yield). ¹H NMR (300 MHz, CDCl₃) δ 7.88 (d, 2H), δ 7.67 (m, 4H), δ 7.44 (m, 3H), δ 6.63 (s, 1H), δ 3.67 (q, J = 6.0, 1.8 Hz, 2H), δ 3.58 (t, J = 6.0 Hz, 2H) ppm; ¹³C NMR (75 MHz, CDCl₃) δ 167.8, 144.8, 140.1, 132.9, 129.2, 128.3, 127.8, 127.5, 127.4, 51.2, 39.7 ppm; HRMS (ESI) calcd for $C_{15}H_{14}N_4O$ (M⁺) 266.1168, found 266.1162.

N-(2-azidoethyl)-4,5-dibromo-1-methyl-1H-pyrrole-2-carboxamide: As in the synthesis of N-(2-azidoethyl)tetradecanamide, (0.529 g, 1.38 mmol) was reacted with 2-azidoethanamine (0.119 g, 1.38 mmol) and triethylamine (0.279 g, 2.75 mmol) in dichloromethane (5 mL) to give N-(2-azidoethyl)-4,5-dibromo-1-methyl-1H-pyrrole-2-carboxamide (0.331 g, 69 % yield). ¹H NMR (300 MHz, CD₃OD) δ 6.84 (s, 1H), δ 3.91 (s, 3H), δ 3.50 (t, J = 3.6 Hz, 2H), δ 3.25 (t, J = 1.5 Hz, 2H) ppm; ¹³C NMR (75 MHz, CD₃OD) δ 161.7, 127.7, 114.7, 111.4, 97.8, 50.4, 38.9, 35.1 ppm; HRMS (ESI) calcd for $C_8H_9N_5O$ (M⁺) 348.9174, found 348.9183.

N-(2-azidoethyl)-1H-pyrrole-2-carboxamide: As in the synthesis of N-(2-azidoethyl)tetradecanamide, (0.249 g, 1.17 mmol) was reacted with 2-azidoethanamine (0.101 g, 1.17 mmol) and triethylamine (0.237 g, 2.34 mmol) in dichloromethane (5 mL) to give N-(2-azidoethyl)-1H-pyrrole-2-carboxamide (0.137 g, 66 % yield). ¹H NMR (300 MHz, CD₃OD) δ 6.88 (d, J = 1.5 Hz, 1H), δ 6.77 (d, J = 2.4 Hz, 1H), δ 6.14 (t, J = 2.4 Hz, 1H), δ 3.46 (t, J = 5.7 Hz, 2H), δ 3.37 (q, J = 5.7, 14.7 Hz, 2H) ppm; ¹³C NMR (75 MHz, CD₃OD) δ 162.9, 125.6, 122.2, 121.9, 110.9, 109.2, 50.6, 38.9 ppm; HRMS (ESI) calcd for $C_7H_9N_5O$ (M⁺) 179.0807, found 179.0803.

N-(2-azidoethyl)-4-bromo-1H-pyrrole-2-carboxamide: As in the synthesis of N-(2-azidoethyl)tetradecanamide, (0.408 g, 1.34 mmol) was reacted with 2-azidoethanamine (0.115 g, 1.34 mmol) and triethylamine (0.270 g, 2.67 mmol) in dichloromethane (5 mL) to give N-(2-azidoethyl)-4-bromo-1H-pyrrole-2-carboxamide (0.299 g, 82 % yield). ¹H NMR (300 MHz, CD₃OD) δ 6.93 (s, 1H), δ 6.80 (s, 1H), δ 3.49 (t, J = 0.6 Hz, 2H), δ 3.41 (t, J = 0.9 Hz, 2H) ppm; ¹³C NMR (75 MHz, CD₃OD) δ 161.7, 126.1, 122.0, 112.4, 96.4, 50.5, 38.9 ppm; HRMS (ESI) calcd for $C_7H_8BrN_5O$ (M⁺) 256.9912, found 256.9908.

N-(2-azidoethyl)-4-nonylbenzamide: As in the synthesis of N-(2-azidoethyl)tetradecanamide, the acid chloride (0.354 g, 1.40 mmol) was reacted with 2-

azidoethanamine (0.121 g, 1.40 mmol) and triethylamine (0.284 g, 2.80 mmol) in dichloromethane (5 mL) to give N-(2-azidoethyl)-4-nonylbenzamide (0.339 g, 80 % yield). ¹H NMR (300 MHz, CDCl₃) δ 6.21 (s, 1H), δ 3.40 (s, 4H), δ 2.19 (t, J = 7.2 Hz, 2H), δ 1.62 (t, J = 6.0 Hz, 2H), δ 1.22 (bs, 14 H), δ 0.842 (t, J = 6.3 Hz, 3H) ppm; ¹³C NMR (75 MHz, CDCl₃) δ 173.8, 51.1, 30.1, 36.8, 32.1, 29.8, 29.7, 29.6, 29.5, 29.2, 25.9, 22.9, 14.3 ppm; HRMS (ESI) calcd for C₁₄H₂₈N₄O (M⁺) 268.2263, found 268.2259.

N-(2-azidoethyl)-4-nonylbenzamide: As in the synthesis of N-(2-azidoethyl)tetradecanamide, (0.383 g, 1.43 mmol) the acid chloride was reacted with 2-azidoethanamine (0.124 g, 1.43 mmol) and triethylamine (0.290 g, 2.87 mmol) in dichloromethane (5 mL) to give N-(2-azidoethyl)-4-nonylbenzamide (0.441 g, 97 % yield). ¹H NMR (300 MHz, CDCl₃) δ 7.72 (d, 2H), δ 7.26 (t, 1H), δ 7.18 (d, 2H), δ 3.57 (q, 2H), δ 3.45 (t, 2H), δ 2.60 (t, 2H), δ 1.58 (m, 2H), δ 1.27 (m, 12H), δ 0.86 (t, 3H) ppm; ¹³C NMR (75 MHz, CDCl₃) δ 168.5, 147.3, 131.7, 128.8, 127.4, 50.9, 39.7, 36.1, 32.1, 31.5, 29.8, 29.7, 29.6, 29.5, 22.9, 14.4 ppm; HRMS (ESI) calcd for C₁₈H₂₈N₄O (M⁺) 316.2263, found 316.2268.

N-(2-azidoethyl)-4-octylbenzamide: As in the synthesis of N-(2-azidoethyl)tetradecanamide, (0.262 g, 1.19 mmol) the acid chloride was reacted with 2-azidoethanamine (0.103 g, 1.19 mmol) and triethylamine (0.242 g, 2.39 mmol) in dichloromethane (5 mL) to give N-(2-azidoethyl)-4-octylbenzamide (0.220 g, 69 % yield). ¹H NMR (300 MHz, CDCl₃) δ 7.73 (d, 2H), δ 7.37 (t, 1H), δ 7.17 (d, 2H), δ 3.56 (q, 2H), δ 3.44 (t, 2H), δ 2.59 (t, 2H), δ 1.58 (m, 2H), δ 1.27 (m, 10H), δ 0.86 (t, 3H) ppm; ¹³C NMR (75 MHz, CDCl₃) δ 168.4, 147.3, 131.7, 128.8, 127.4, 50.9, 39.7, 36.1, 32.1, 31.5, 29.7, 29.5, 29.5, 22.9, 14.3 ppm; HRMS (ESI) calcd for C₁₇H₂₈N₄O (M⁺) 302.2107, found 302.2104.

(E)-N-(2-azidoethyl)-2-methyl-3-phenylacrylamide: To a 25 mL round-bottomed flask equipped with a magnetic stir bar was added (E)-2-methyl-3-phenylacrylic acid (0.0512 g, 3.15 mmol) and dichloromethane (10 mL). Oxalyl chloride (0.400 g, 3.15 mmol) was added dropwise to the reaction mixture and allowed to stir for one hour. Then, the reaction mixture was concentrated de vacuo. To the crude mixture was then added dichloromethane (10 mL), 2-azidoethanamine (0.299 g, 3.47 mmol) and then triethylamine (0.351 g, 3.47 mmol) and allowed to stir for two hours. The reaction mixture was then concentrated de vacuo and then purified via silica gel column chromatography (100% dichloromethane to 1:40 methanol: dichloromethane) to give (E)-N-(2-azidoethyl)-2-methyl-3-phenylacrylamide (0.698 g, 96 % yield). ¹H NMR (300 MHz, CDCl₃) δ 7.39 (m, 6H), δ 7.01 (s, 1H), δ 3.51 (t, J = 4.5 Hz, 2H), δ 3.45 (t, J = 4.8 Hz, 2H), δ 2.13 (s, 3H) ppm; ¹³C NMR (75 MHz, CDCl₃) δ 170.5, 136.2, 134.5, 132.0, 130.3, 129.6, 128.9, 128.6, 128.3, 50.9, 39.8, 14.5 ppm; HRMS (ESI) calcd for C₁₂H₁₄N₄O (M⁺) 230.1168, found 230.1165.

N-(2-azidoethyl)-2',4'-difluoro-3-hydroxybiphenyl-4-carboxamide: To a 25 mL round-bottomed flask equipped with a magnetic stir bar was added 2',4'-difluoro-3-hydroxybiphenyl-4-carboxylic acid (0.301, 1.20 mmol), *N,N*-Dimethylformamide (5 mL), *N,N'*-Dicyclohexylcarbodiimide (0.248 g, 1.20 mmol) and *n*-methylmorpholine (0.25 mL). The reaction mixture was cooled to 0 °C and allowed to stir. Then, 2-azidoethanamine (0.1037 g, 1.20 mmol) was added dropwise and allowed to slowly warm to room temperature while stirring for 24 hr. The reaction mixture was diluted with water, extracted with dichloromethane, washed with 1N HCl, washed with saturated sodium bicarbonate, washed with brine and then concentrated de vacuo. The resulting residue was then purified via silica gel column chromatography (1:40 methanol: dichloromethane to 1:10 methanol: dichloromethane) to give N-(2-azidoethyl)-2',4'-difluoro-3-hydroxybiphenyl-4-carboxamide (0.232 g, 61 % yield). ¹H NMR (300 MHz, CDCl₃) δ 7.60 (s, 1H), δ 7.47 (d, J = 6.0 Hz, 1H), δ 7.29 (m, 2H), δ 7.02 (d, J = 6.6 Hz, 1H), δ 6.88 (m, 2H) ppm; ¹³C NMR (75 MHz, CDCl₃) δ 170.3, 160.8, 135.1, 131.3, 131.2, 131.3, 131.2, 126.8, 118.8, 114.6, 112.0, 111.8, 104.8, 104.6, 104.3, 50.7, 39.3 ppm; HRMS (ESI) calcd for C₁₅H₁₂F₂N₄O₃ (M⁺) 318.0928, found 318.0933.

General procedure for click reactions and subsequent Boc deprotection: The terminal alkyne (1.0 equiv.) was dissolved in a 1:1:1 mixture of ethanol, water and methylene chloride (ca. 9 mL per 0.300 g of terminal alkyne). To this solution, the appropriate azide (1.0 equiv.) was added while stirring vigorously at room temperature. Copper (II) sulfate (15 mol%) and sodium ascorbate (45 mol%) were then added sequentially to the solution. Reaction mixtures were allowed to stir until completion via TLC analysis (12 – 24 hrs). The solvents were then removed de vacuo in which the resulting residue was dissolved in dichloromethane and purified via silica gel column chromatography (1:40 methanol: dichloromethane to 1:10 methanol: dichloromethane). To remove the Boc protecting group, the resulting product was then dissolved in a 1:4 trifluoroacetic acid: dichloromethane mixture and allowed to stir for 5 hr. Upon completion, the reaction mixture was concentrated de vacuo and then left on a high vacuum overnight. Then, methanol supplemented with HCl was added to the product forming the HCl salt of the deprotected product and then was concentrated de vacuo. The resulting residue was washed with diethyl ether and then placed on a high vacuum overnight.

N-(2-(4-(5-(2-amino-1H-imidazol-4-yl)pentyl)-1H-1,2,3-triazol-1-yl)ethyl)-2-(phenylthio)acetamide hydrochloride: tert-butyl 2-amino-4-(hept-6-ynyl)-1H-imidazole-1-carboxylate (0.112 g, 0.405 mmol) was reacted with N-(2-azidoethyl)-2-(phenylthio)acetamide (0.096 g, 0.405 mmol) following the general click procedure to give tert-butyl 2-amino-4-(5-(1-(2-(2-(phenylthio)acetamido)ethyl)-1H-1,2,3-triazol-4-yl)pentyl)-1H-imidazole-1-carboxylate ¹H NMR (300 MHz, CDCl₃) δ 7.45 (s, 1H), δ 7.19 (m, 5H), δ 7.13 (s, 1H), δ 6.43 (s, 1H), δ 6.13 (bs,

2H), δ 4.29 (s, 2H), δ 3.68 (s, 2H), δ 3.57 (s, 2H), δ 2.56 (s, 2H), δ 2.35 (s, 2H), δ 1.52 (m, 15H) ppm; ^{13}C NMR (75 MHz, CDCl_3) δ 187.8, 172.0, 169.1, 162.4, 156.9, 135.0, 129.5, 128.3, 126.8, 121.8, 85.00, 49.4, 39.9, 37.4, 29.4, 28.9, 28.4, 28.2, 27.6, 25.6 ppm; HRMS (ESI) calcd for $\text{C}_{25}\text{H}_{35}\text{N}_7\text{O}_3\text{S}$ (M^+) 513.2522, found 513.2522, which was subsequently deprotected to give

5 N-(2-(4-(5-(2-amino-1H-imidazol-4-yl)pentyl)-1H-1,2,3-triazol-1-yl)ethyl)-2-(phenylthio)acetamide hydrochloride (0.133 g, 73 % yield). ^1H NMR (300 MHz, CD_3OD) δ 8.47 (s, 1H), δ 7.30 (s, 4H), δ 7.19 (s, 1H), δ 6.58 (s, 1H), δ 4.66 (s, 2H), δ 3.76 (s, 2H), δ 3.64 (s, 2H), δ 2.84 (s, 2H), δ 2.34 (s, 2H), δ 1.73 (s, 2H), δ 1.64 (s, 2H), δ 1.26 (s, 2H) ppm; ^{13}C NMR (75 MHz, CD_3OD) δ 170.9, 159.3, 158.9, 147.2, 135.6, 128.1, 128.9, 127.6, 126.6, 108.7, 108.6, 10
52., 39.1, 37.5, 37.2, 36.6, 36.1, 30.8, 28.1, 27.9, 27.7, 24.2, 23.7, 23.6 ppm; HRMS (ESI) calcd for $\text{C}_{20}\text{H}_{27}\text{N}_7\text{OS}$ (M^+) 413.1998, found 413.1991.

N-(2-(4-(5-(2-amino-1H-imidazol-4-yl)pentyl)-1H-1,2,3-triazol-1-yl)ethyl)thiophene-2-sulfonamide hydrochloride: tert-butyl 2-amino-4-(hept-6-ynyl)-1H-imidazole-1-carboxylate (0.105 g, 0.379 mmol) was reacted with N-(2-azidoethyl)thiophene-2-sulfonamide (0.099 g,
15 0.379 mmol) following the general click procedure to give tert-butyl 2-amino-4-(5-(1-(2-(thiophene-2-sulfonamido)ethyl)-1H-1,2,3-triazol-4-yl)pentyl)-1H-imidazole-1-carboxylate ^1H NMR (300 MHz, CDCl_3) δ 7.49 (s, 3H), δ 7.39 (s, 1H), δ 6.99 (s, 1H), δ 6.41 (bm, 3H), δ 4.43 (s, 2H), δ 3.42 (s, 2H), δ 2.54 (s, 2H), δ 2.09 (s, 2H), δ 1.51 (m, 13H), δ 1.19 (s, 2H) ppm; ^{13}C NMR (75 MHz, CDCl_3) δ 150.8, 149.4, 148.1, 141.1, 138.3, 132.2, 132.1, 127.7, 122.6, 85.2, 53.8,
20 50.2, 43.2, 31.2, 29.9, 29.2, 28.8, 28.1, 25.5 ppm; HRMS (ESI) calcd for $\text{C}_{21}\text{H}_{31}\text{N}_7\text{O}_4\text{S}_2$ (M^+) 509.1879, found 509.1879, which was subsequently deprotected to give N-(2-(4-(5-(2-amino-1H-imidazol-4-yl)pentyl)-1H-1,2,3-triazol-1-yl)ethyl)thiophene-2-sulfonamide hydrochloride (0.098 g, 58 % yield). ^1H NMR (400 MHz, CD_3OD) δ 8.37 (s, 1H), δ 7.76 (s, 1H), δ 7.57 (s, 1H), δ 7.11 (s, 1H), δ 6.46 (s, 1H), δ 4.79 (s, 2H), δ 3.55 (s, 2H), δ 2.50 (s, 2H), δ 1.66-1.18 (bm, 8H)
25 ppm; ^{13}C NMR (100 MHz, CD_3OD) δ 175.6, 155.6, 147.2, 140.9, 132.7, 132.3, 127.8, 127.7, 108.8, 53.1, 42.2, 36.8, 28.1, 27.8, 27.5, 24.9, 24.3, 23.8 ppm; HRMS (ESI) calcd for $\text{C}_{16}\text{H}_{23}\text{N}_7\text{O}_2\text{S}_2$ (M^+) 409.1355, found 409.1354.

N-(2-(4-(5-(2-amino-1H-imidazol-4-yl)pentyl)-1H-1,2,3-triazol-1-yl)ethyl)-5-(dimethylamino)naphthalene-1-sulfonamide hydrochloride: tert-butyl 2-amino-4-(hept-6-
30 ynyl)-1H-imidazole-1-carboxylate (0.112 g, 0.403 mmol) was reacted with N-(2-azidoethyl)-5-(dimethylamino)naphthalene-1-sulfonamide (0.140 g, 0.403 mmol) following the general click procedure to give tert-butyl 2-amino-4-(5-(1-(2-(5-(dimethylamino)naphthalene-1-sulfonamido)ethyl)-1H-1,2,3-triazol-4-yl)pentyl)-1H-imidazole-1-carboxylate ^1H NMR (300 MHz, CDCl_3) δ 8.48 (d, J = 8.4 Hz, 1H), δ 8.24 (d, J = 8.4 Hz, 1H), δ 8.17 (d, J = 7.2 Hz, 1H), δ
35 7.44 (m, 3H), δ 7.09 (s, 1H), δ 7.07 (d, J = 7.5 Hz, 1H), δ 6.44 (s, 1H), δ 6.09 (bs, 2H), δ 4.32 (s,

2H), δ 3.36 (s, 2H), δ 2.52 (s, 2H), δ 2.22 (s, 2H), δ 1.43 (m, 13H), δ 1.22 (s, 2H) ppm; ^{13}C NMR (75 MHz, CDCl_3) δ 176.9, 152.1, 149.6, 148.0, 138.6, 135.1, 130.7, 130.1, 129.7, 129.4, 128.5, 123.3, 122.4, 119.1, 115.5, 84.9, 67.5, 50.3, 50.2, 45.6, 42.9, 37.9, 29.1, 28.9, 28.2, 28.0, 25.5 ppm; HRMS (ESI) calcd for $\text{C}_{29}\text{H}_{40}\text{N}_8\text{O}_4\text{S}$ (M^+) 596.2893, found 596.2881, which was subsequently deprotected to give N-(2-(4-(5-(2-amino-1H-imidazol-4-yl)pentyl)-1H-1,2,3-triazol-1-yl)ethyl)-5-(dimethylamino)naphthalene-1-sulfonamide hydrochloride (0.139 g, 65 % yield). ^1H NMR (400 MHz, CD_3OD) δ 8.70 (t, J = 4.8 Hz, 2H), δ 8.31 (s, 1H), δ 8.08 (s, 2H), δ 7.82 (s, 2H), δ 6.43 (s, 1H), δ 4.55 (s, 2H), δ 3.32 (s, 8H), 2.63 (s, 2H), δ 2.48 (s, 2H), δ 1.66 (s, 4H), δ 1.40 (s, 2H) ppm; ^{13}C NMR (100 MHz, CD_3OD) δ 159.9, 147.3, 140.7, 136.8, 129.9, 129.3, 127.8, 127.7, 126.5, 119.3, 108.6, 108.4, 76.7, 67.3, 51.5, 42.3, 37.4, 36.5, 28.3, 27.9, 27.6, 24.1, 24.0, 23.7 ppm; HRMS (ESI) calcd for $\text{C}_{24}\text{H}_{32}\text{N}_8\text{O}_2\text{S}$ (M^+) 496.2369, found 496.2359.

tert-butyl 2-amino-4-(5-(1-(2-(phenylsulfonamido)ethyl)-1H-1,2,3-triazol-4-yl)pentyl)-1H-imidazole-1-carboxylate hydrochloride: tert-butyl 2-amino-4-(hept-6-ynyl)-1H-imidazole-1-carboxylate (0.114 g, 0.410 mmol) was reacted with N-(2-azidoethyl)benzenesulfonamide (0.104 g, 0.410 mmol) following the general click procedure to give tert-butyl 2-amino-4-(5-(1-(2-(phenylsulfonamido)ethyl)-1H-1,2,3-triazol-4-yl)pentyl)-1H-imidazole-1-carboxylate ^1H NMR (300 MHz, CDCl_3) δ 7.81 (d, J = 6.9 Hz, 2H), δ 7.44 (m, 5H), δ 6.44 (s, 1H), δ 6.07 (s, 2H), δ 4.42 (t, J = 5.7 Hz, 2H), δ 3.36 (t, J = 5.4 Hz, 2H), δ 2.57 (t, J = 7.2 Hz, 2H), δ 2.20 (s, 2H), δ 1.47 (m, 13H), δ 1.29 (s, 2H) ppm; ^{13}C NMR (75 MHz, CDCl_3) δ 202.9, 149.5, 148.2, 140.2, 132.8, 129.4, 127.1, 122.5, 106.5, 85.0, 53.7, 50.3, 42.9, 32.7, 31.2, 29.9, 29.2, 28.9, 28.2, 26.3, 25.5 ppm; HRMS (ESI) calcd for $\text{C}_{23}\text{H}_{33}\text{N}_7\text{O}_4\text{S}$ (M^+) 503.2315, found 503.2310, which was subsequently deprotected to give tert-butyl 2-amino-4-(5-(1-(2-(phenylsulfonamido)ethyl)-1H-1,2,3-triazol-4-yl)pentyl)-1H-imidazole-1-carboxylate hydrochloride (0.142 g, 64 % yield). ^1H NMR (300 MHz, CD_3OD) δ 7.81 (d, J = 6.6 Hz, 2H), δ 7.56 (m, 4H), δ 6.47 (s, 1H), δ 4.62 (s, 2H), δ 3.41 (s, 2H), δ 2.78 (s, 2H), δ 2.48 (s, 2H), δ 1.72 (s, 2H), δ 1.63 (s, 2H), δ 1.42 (s, 2H) ppm; ^{13}C NMR (75 MHz, CD_3OD) δ 176.2, 165.6, 159.9, 147.3, 140.2, 132.8, 129.3, 127.7, 126.8, 108.5, 108.4, 515.8, 42.3, 28.1, 27.9, 27.6, 24.1 ppm; HRMS (ESI) calcd for $\text{C}_{18}\text{H}_{23}\text{N}_7\text{O}_2\text{S}$ (M^+) 403.1790, found 403.1781.

N-(2-(4-(5-(2-amino-1H-imidazol-4-yl)pentyl)-1H-1,2,3-triazol-1-yl)ethyl)-2,3,4,5,6-pentamethylbenzenesulfonamide hydrochloride: tert-butyl 2-amino-4-(hept-6-ynyl)-1H-imidazole-1-carboxylate (0.120 g, 0.434 mmol) was reacted with N-(2-azidoethyl)-2,3,4,5,6-pentamethylbenzenesulfonamide (0.141 g, 0.434 mmol) following the general click procedure to give tert-butyl 2-amino-4-(5-(1-(2-(2,3,4,5,6-pentamethylphenylsulfonamido)ethyl)-1H-1,2,3-triazol-4-yl)pentyl)-1H-imidazole-1-carboxylate ^1H NMR (300 MHz, CDCl_3) δ 7.30 (s, 1H), δ

7.01 (s, 1H), δ 6.44 (s, 1H), δ 6.17 (s, 2H), δ 4.35 (s, 2H), δ 3.34 (s, 2H), δ 2.47 (s, 2H), δ 2.48 (s, 6H), δ 2.20 (m, 11H), δ 1.43 (m, 15H) ppm; ^{13}C NMR (75 MHz, CDCl_3) δ 149.6, 148.1, 139.7, 136.2, 134.9, 134.3, 122.4, 84.9, 53.7, 50.2, 42.5, 31.1, 29.3, 28.9, 28.4, 28.2, 25.6, 19.1, 17.9, 17.2 ppm; HRMS (ESI) calcd for $\text{C}_{28}\text{H}_{43}\text{N}_7\text{O}_4\text{S}$ (M^+) 573.3097, found 573.3086, which was subsequently deprotected to give N-(2-(4-(5-(2-amino-1H-imidazol-4-yl)pentyl)-1H-1,2,3-triazol-1-yl)ethyl)-2,3,4,5,6-pentamethylbenzenesulfonamide hydrochloride (0.143 g, 76 % yield). ^1H NMR (400 MHz, CD_3OD) δ 8.11 (s, 1H), δ 6.42 (s, 1H), δ 4.48 (s, 2H), δ 3.34 (s, 2H), δ 2.65 (s, 2H), δ 2.40 (s, 8H), δ 2.12 (s, 9H), δ 1.58 (s, 4H), δ 1.19 (s, 2H) ppm; ^{13}C NMR (100 MHz, CD_3OD) δ 159.9, 147.3, 139.6, 136.1, 134.8, 134.0, 127.8, 127.6, 108.6, 108.4, 54.1, 51.5, 41.8, 36.5, 28.2, 28.1, 27.7, 24.1, 23.7, 18.2, 16.8, 16.1 ppm; HRMS (ESI) calcd for $\text{C}_{23}\text{H}_{35}\text{N}_7\text{O}_2\text{S}$ (M^+) 473.2573, found 473.2565.

(E)-N-(2-(4-(5-(2-amino-1H-imidazol-4-yl)pentyl)-1H-1,2,3-triazol-1-yl)ethyl)-2-methyl-3-phenylacrylamide hydrochloride: tert-butyl 2-amino-4-(hept-6-ynyl)-1H-imidazole-1-carboxylate (0.1103 g, 0.397 mmol) was reacted with (E)-N-(2-azidoethyl)-2-methyl-3-phenylacrylamide (0.092 g, 0.397 mmol) following the general click procedure to give (E)-tert-butyl 2-amino-4-(5-(1-(2-(2-methyl-3-phenylacrylamido)ethyl)-1H-1,2,3-triazol-4-yl)pentyl)-1H-imidazole-1-carboxylate ^1H NMR (300 MHz, CDCl_3) δ 7.28 (m, 7H), δ 6.94 (s, 2H), δ 6.06 (s, 2H), δ 4.46 (s, 2H), δ 3.79 (s, 2H), δ 2.61 (s, 2H), δ 2.27 (s, 2H), δ 1.99 (s, 3H), δ 1.49 (bs, 11H), δ 1.18 (s, 4H) ppm; ^{13}C NMR (75 MHz, CDCl_3) δ 192.4, 191.3, 170.4, 150.9, 140.2, 136.2, 134.8, 131.6, 129.6, 128.6, 128.1, 122.1, 95.8, 85.2, 74.3, 53.7, 52.6, 50.7, 49.4, 40.1, 29.9, 29.3, 28.9, 28.2, 25.6, 14.4 ppm; HRMS (ESI) calcd for $\text{C}_{27}\text{H}_{37}\text{N}_7\text{O}_3$ (M^+) 507.2958, found 507.2942, which was subsequently deprotected to give (E)-N-(2-(4-(5-(2-amino-1H-imidazol-4-yl)pentyl)-1H-1,2,3-triazol-1-yl)ethyl)-2-methyl-3-phenylacrylamide hydrochloride (0.148 g, 84 % yield). ^1H NMR (400 MHz, CD_3OD) δ 8.78 (s, 1H), δ 7.33 (s, 4H), δ 7.26 (s, 1H), δ 7.19 (s, 1H), δ 6.42 (s, 1H), δ 4.78 (s, 2H), δ 3.84 (s, 2H), δ 2.87 (s, 2H), δ 2.44 (s, 2H), δ 1.99 (s, 3H), δ 1.76 (s, 2H), δ 1.59 (s, 2H), δ 1.42 (s, 2H) ppm; ^{13}C NMR (100 MHz, CD_3OD) δ 174.7, 136.0, 134.4, 131.5, 129.3, 128.7, 128.3, 128.0, 127.6, 108.6, 94.6, 53.1, 39.2, 27.6, 27.9, 27.7, 27.6, 24.1, 23.8, 23.1, 14.3, 13.4 ppm; HRMS (ESI) calcd for $\text{C}_{22}\text{H}_{29}\text{N}_7\text{O}$ (M^+) 407.2434, found 407.2429.

N-(2-(4-(5-(2-amino-1H-imidazol-4-yl)pentyl)-1H-1,2,3-triazol-1-yl)ethyl)cinnamamide hydrochloride: tert-butyl 2-amino-4-(hept-6-ynyl)-1H-imidazole-1-carboxylate (0.106 g, 0.384 mmol) was reacted with N-(2-azidoethyl)cinnamamide (0.083 g, 0.384 mmol) following the general click procedure to give (E)-tert-butyl 2-amino-4-(5-(1-(2-cinnamamidoethyl)-1H-1,2,3-triazol-4-yl)pentyl)-1H-imidazole-1-carboxylate ^1H NMR (300 MHz, CDCl_3) δ 7.67 (s, 1H), δ 7.51 (d, $J = 11.1$ Hz, 1H), δ 7.28 (m, 6H), δ 6.45 (t, $J = 14.1$ Hz, 2H), δ 6.08 (s, 2H), δ 4.47 (s, 2H), δ 3.79 (s, 2H), δ 2.57 (s, 2H), δ 2.23 (s, 2H), δ 1.50 (bs, 13H),

δ 1.28 (s, 2H) ppm; ^{13}C NMR (75 MHz, CDCl_3) δ 166.9, 150.4, 149.5, 148.3, 141.2, 138.9, 134.9, 129.9, 128.9, 128.0, 122.1, 120.8, 106.5, 84.8, 53.7, 49.5, 39.9, 31.1, 29.3, 28.9, 28.3, 28.2, 25.6 ppm; HRMS (ESI) calcd for $\text{C}_{26}\text{H}_{35}\text{N}_7\text{O}_3$ (M⁺) 493.2801, found 493.2800, which was subsequently deprotected to give N-(2-(4-(5-(2-amino-1H-imidazol-4-yl)pentyl)-1H-1,2,3-triazol-1-yl)ethyl)cinnamamide hydrochloride (0.097 g, 59 % yield). ^1H NMR (400 MHz, CD_3OD) δ 8.10 (s, 1H), δ 7.49 (s, 3H), δ 7.32 (s, 3H), δ 6.30 d, J = 10.8 Hz, 1H), δ 6.39 (s, 1H), δ 4.66 (s, 2H), δ 3.83 (s, 2H), δ 2.73 (s, 2H), δ 2.40 (s, 2H), δ 1.57 (s, 2H), δ 1.55 (s, 2H), δ 1.36 (s, 2H) ppm; ^{13}C NMR (100 MHz, CD_3OD) δ 167.8, 160.5, 150.2, 147.3, 141.1, 134.8, 130.7, 129.9, 128.8, 127.8, 127.6, 124.9, 123.9, 120.2, 108.5, 108.3, 53.1, 39.3, 36.5, 30.6, 28.4, 28.1, 27.6, 26.5, 24.2, 24.1, 23.7 ppm; HRMS (ESI) calcd for $\text{C}_{21}\text{H}_{27}\text{N}_7\text{O}$ (M⁺) 393.2277, found 393.2273.

N-(2-(4-(5-(2-amino-1H-imidazol-4-yl)pentyl)-1H-1,2,3-triazol-1-yl)ethyl)heptadec-8-enamide hydrochloride: tert-butyl 2-amino-4-(hept-6-ynyl)-1H-imidazole-1-carboxylate (g, mmol) was reacted with N-(2-azidoethyl)octadec-9-enamide (g, mmol) following the general click procedure to give tert-butyl 2-amino-4-(5-(1-(2-heptadec-8-enamidoethyl)-1H-1,2,3-triazol-4-yl)pentyl)-1H-imidazole-1-carboxylate ^1H NMR (300 MHz, CDCl_3) δ 7.26 (s, 1H), δ 6.81 (s, 1H), δ 6.42 (s, 2H), δ 5.88 (s, 1H), δ 5.27 (s, 2H), δ 4.38 (s, 2H), δ 3.65 (s, 2H), δ 2.60 (s, 2H), δ 2.09 (s, 2H), δ 1.53-1.01 (m, 35H), 0.79 (t, J = 6.3 Hz, 3H) ppm; ^{13}C NMR (75 MHz, CDCl_3) δ 174.2, 173.8, 148.3, 130.3, 130.1, 129.9, 129.8, 121.9, 85.1, 67.1, 49.5, 39.5, 39.1, 38.0, 37.7, 36.7, 36.6, 32.8, 32.3, 32.1, 31.7, 29.9, 29.8, 29.7, 29.5, 29.3, 29.2, 28.9, 28.7, 28.1, 28.0, 27.9, 27.4, 27.3, 27.3, 26.9, 25.9, 25.6, 25.3, 22.9, 22.8, 22.6, 14.3, 14.2 ppm; HRMS (ESI) calcd for $\text{C}_{35}\text{H}_{61}\text{N}_7\text{O}_3$ (M⁺) 627.4836, found 627.4823, which was subsequently deprotected to give N-(2-(4-(5-(2-amino-1H-imidazol-4-yl)pentyl)-1H-1,2,3-triazol-1-yl)ethyl)heptadec-8-enamide hydrochloride (0.128 g, 66 % yield). ^1H NMR (400 MHz, CD_3OD) δ 8.36 (s, 1H), δ 6.96 (s, 1H), δ 6.45 (s, 1H), δ 6.09 (s, 1H), δ 4.58 (s, 2H), δ 3.58 (s, 2H), δ 2.75 (s, 2H), δ 2.10 (s, 2H), δ 1.96 (s, 4H), δ 1.69 (s, 2H), δ 1.41 (m, 4H), 1.23 (s, 24H), δ 0.84 (s, 3H) ppm; ^{13}C NMR (100 MHz, CD_3OD) δ 175.5, 158.8, 155.7, 147.3, 129.7, 129.6, 127.6, 108.5, 94.6, 76.7, 51.3, 50.4, 48.6, 48.4, 48.2, 47.9, 47.8, 47.5, 47.3, 38.9, 37.5, 36.6, 45.9, 45.8, 33.9, 32.6, 31.9, 31.5, 29.8, 29.5, 29.4, 29.3, 29.2, 29.1, 28.2, 27.8, 27.0, 25.8, 24.9, 24.2, 23.8, 22.5, 13.5, 13.3 ppm; HRMS (ESI) calcd for $\text{C}_{30}\text{H}_{53}\text{N}_7\text{O}$ (M⁺) 527.4312, found 527.4298.

N-(2-(4-(5-(2-amino-1H-imidazol-4-yl)pentyl)-1H-1,2,3-triazol-1-yl)ethyl)decanamide hydrochloride: tert-butyl 2-amino-4-(hept-6-ynyl)-1H-imidazole-1-carboxylate (0.114 g, 0.412 mmol) was reacted with N-(2-azidoethyl)decanamide (0.099 g, 0.412 mmol) following the general click procedure to give tert-butyl 2-amino-4-(5-(1-(2-decanamidoethyl)-1H-1,2,3-triazol-4-yl)pentyl)-1H-imidazole-1-carboxylate ^1H NMR (400

MHz, CDCl₃) δ 7.23 (s, 1H), δ 6.57 (s, 1H), δ 6.43 (s, 1H), δ 6.09 (s, 2H), δ 4.38 (s, 2H), δ 3.67 (s, 2H), δ 2.61 (t, J = 5.4 Hz, 2H), δ 2.08 (s, 2H), δ 2.08 (t, J = 5.7 Hz, 2H), δ 1.42 (m, 15H), δ 1.17 (m, 14H), δ 0.79 (t, J = 4.5 Hz, 3H) ppm; ¹³C NMR (100 MHz, CDCl₃) δ 174.2, 149.7, 148.4, 138.8, 121.9, 106.6, 84.9, 49.5, 39.5, 36.7, 32.1, 31.3, 30.6, 29.9, 29.7, 29.6, 29.5, 29.5, 29.3, 29.0, 28.3, 28.2, 25.9, 25.6, 25.5, 22.9, 14.3 ppm; HRMS (ESI) calcd for C₂₇H₄₇N₇O₃ (M⁺) 517.3740, found 517.3736, which was subsequently deprotected to give N-(2-(4-(5-(2-amino-1H-imidazol-4-yl)pentyl)-1H-1,2,3-triazol-1-yl)ethyl)decanamide hydrochloride (0.137 g, 73 % yield). ¹H NMR (300 MHz, CD₃OD) δ 8.55 (s, 1H), δ 6.53 (s, 1H), 4.72 (s, 2H), δ 3.75 (s, 2H), 2.89 (s, 2H), δ 2.54 (t, J = 6.8 Hz, 2H), δ 2.16 (t, J = 7.2 Hz, 2H), δ 1.81 (s, 2H), δ 1.69 (s, 2H), δ 1.49 (m, 4H), δ 1.28 (s, 12H), δ 0.89 (t, 3H) ppm; ¹³C NMR (75 MHz, CD₃OD) δ 175.7, 147.3, 128.5, 127.6, 125.1, 120.8, 108.6, 108.4, 105.6, 63.1, 59.3, 52.5, 48.6, 48.3, 48.1, 47.9, 47.7, 47.5, 57.3, 38.7, 336.6, 35.8, 36.6, 35.8, 31.8, 30.6, 29.4, 29.3, 29.3, 29.2, 28.0, 27.8, 27.7, 25.8, 24.1, 23.9, 23.3, 22.6 ppm; HRMS (ESI) calcd for C₂₂H₃₉N₇O (M⁺) 417.3216, found 417.3209.

N-(2-(4-(5-(2-amino-1H-imidazol-4-yl)pentyl)-1H-1,2,3-triazol-1-yl)ethyl)dodecanamide hydrochloride: tert-butyl 2-amino-4-(hept-6-ynyl)-1H-imidazole-1-carboxylate (0.107 g, 0.387 mmol) was reacted with N-(2-azidoethyl)dodecanamide (0.104 g, 0.387 mmol) following the general click procedure to give tert-butyl 2-amino-4-(5-(1-(2-dodecanamidoethyl)-1H-1,2,3-triazol-4-yl)pentyl)-1H-imidazole-1-carboxylate ¹H NMR (300 MHz, CDCl₃) δ 7.36 (s, 1H), δ 6.94 (s, 1H), δ 6.49 (s, 1H), δ 6.23 (s, 2H), δ 4.46 (t, J = 5.1 Hz, 2H), δ 3.74 (q, J = 5.1, 5.7 Hz, 2H), δ 2.68 (t, J = 7.5 Hz, 2H), δ 2.34 (t, J = 6.9 Hz, 2H), δ 2.16 (t, J = 7.5 Hz, 2H), δ 1.54 (m, 13H), δ 1.47 (m, 2H), δ 1.24 (m, 18H), δ 0.87 (t, J = 6.6 Hz, 3H) ppm; ¹³C NMR (75 MHz, CDCl₃) δ 174.3, 150.4, 149.5, 148.3, 138.5, 121.9, 106.5, 84.9, 49.5, 39.5, 36.6, 34.1, 32.1, 29.8, 29.8, 29.7, 29.6, 29.5, 29.3, 28.9, 28.2, 28.1, 29.0, 26.9, 25.6, 22.9, 14.3 ppm; HRMS (ESI) calcd for C₂₉H₅₁N₇O₃ (M⁺) 545.4053, found 545.4053, which was subsequently deprotected to give N-(2-(4-(5-(2-amino-1H-imidazol-4-yl)pentyl)-1H-1,2,3-triazol-1-yl)ethyl)dodecanamide hydrochloride (0.155 g, 83 % yield). ¹H NMR (400 MHz, CD₃OD) δ 8.31 (s, 1H), δ 6.49 (s, 1H), δ 4.63 (s, 2H), δ 3.97 (s, 2H), δ 2.81 (s, 2H), δ 2.51 (t, J = 7.2 Hz, 2H), δ 2.14 (t, J = 7.2 Hz, 2H), δ 1.76 (s, 2H), δ 1.67 (s, 2H), δ 1.51 (s, 2H), δ 1.44 (s, 2H), δ 1.26 (bs, 16H), δ 0.71 (s, 3H) ppm; ¹³C NMR (100 MHz, CD₃OD) δ 175.6, 159.4, 147.3, 127.6, 108.5, 51.3, 48.6, 38.8, 35.8, 31.9, 29.6, 29.5, 29.4, 29.3, 29.2, 28.2, 28.1, 27.7, 24.8, 24.2, 23.9, 22.6, 22.5, 13.4, 13.3 ppm; HRMS (ESI) calcd for C₂₄H₄₃N₇O (M⁺) 445.3429, found 445.3524.

N-(2-(4-(5-(2-amino-1H-imidazol-4-yl)pentyl)-1H-1,2,3-triazol-1-yl)ethyl)tetradecanamide hydrochloride: tert-butyl 2-amino-4-(hept-6-ynyl)-1H-imidazole-1-

carboxylate (0.111 g, 0.401 mmol) was reacted with N-(2-azidoethyl)tetradecanamide (0.130 g, 0.401 mmol) following the general click procedure to give tert-butyl 2-amino-4-(5-(1-(2-tetradecanamidoethyl)-1H-1,2,3-triazol-4-yl)pentyl)-1H-imidazole-1-carboxylate ¹H NMR (300 MHz, CDCl₃) δ 7.26 (s, 1H), δ 6.85 (s, 1H), δ 6.42 (s, 1H), δ 6.05 (s, 1H), δ 4.38 (s, 2H), δ 3.66 (s, 2H), δ 2.60 (s, 2H), δ 2.11 (s, 2H), δ 2.08 (t, J = 3.6 Hz, 2H), δ 1.41 (m, 13H), δ 1.32 (s, 2H), δ 1.17 (s, 18H), δ 0.77 (t, J = 6.6 Hz, 3H) ppm; ¹³C NMR (75 MHz, CDCl₃) δ 202.9, 174.3, 149.6, 148.3, 121.9, 100.1, 84.8, 49.5, 42.3, 39.5, 36.7, 32.1, 29.9, 29.8, 29.7, 29.6, 29.5, 29.5, 29.3, 28.9, 28.1, 25.9, 25.6, 22.9, 14.3 ppm; HRMS (ESI) calcd for C₃₁H₅₅N₇O₃ (M⁺) 573.4366, found 573.4365, which was subsequently deprotected to give N-(2-(4-(5-(2-amino-1H-imidazol-4-yl)pentyl)-1H-1,2,3-triazol-1-yl)ethyl)tetradecanamide hydrochloride (0.118 g, 87 % yield). ¹H NMR (400 MHz, CD₃OD) δ 8.51 (s, 1H), δ 6.53 (s, 1H), δ 4.71 (s, 2H), δ 3.75 (s, 2H), δ 2.89 (s, 2H), δ 2.54 (t, J = 6.8 Hz, 2H), δ 2.17 (t, J = 6.8 Hz, 2H), δ 1.80 (s, 2H), δ 1.69 (s, 2H), δ 1.54 (s, 2H), δ 1.48 (s, 2H), δ 1.28 (s, 20H), 0.88 (t, J = 6.8 Hz, 3H) ppm; ¹³C NMR (100 MHz, CD₃OD) δ 175.6, 161.8, 147.2, 127.6, 127.4, 108.5, 52.3, 38.8, 36.6, 35.8, 31.9, 30.6, 29.7, 29.6, 29.5, 29.4, 29.2, 28.0, 27.9, 27.7, 25.8, 24.1, 23.6, 23.4, 22.6, 13.4 ppm; HRMS (ESI) calcd for C₂₆H₄₇N₇O (M⁺) 473.3842, found 473.3834.

N-(2-(4-(5-(2-amino-1H-imidazol-4-yl)pentyl)-1H-1,2,3-triazol-1-yl)ethyl)palmitamide hydrochloride: tert-butyl 2-amino-4-(hept-6-ynyl)-1H-imidazole-1-carboxylate (0.087 g, 0.313 mmol) was reacted with N-(2-azidoethyl)palmitamide (0.102 g, 0.313 mmol) following the general click procedure to give tert-butyl 2-amino-4-(5-(1-(2-palmitamidoethyl)-1H-1,2,3-triazol-4-yl)pentyl)-1H-imidazole-1-carboxylate ¹H NMR (300 MHz, CDCl₃) δ 7.29 (s, 1H), δ 6.78 (t, J = 4.5 Hz, 1H), δ 6.41 (s, 1H), δ 6.19 (s, 2H), δ 4.39 (t, J = 5.4 Hz, 2H), δ 3.67 (q, J = 5.5, 4.8 Hz, 2H), δ 2.61 (t, J = 6.9 Hz, 2H), δ 2.12 (t, J = 6.5 Hz, 2H), δ 2.07 (t, J = 7.5 Hz, 2H), δ 1.42 (m, 13H), δ 1.42 (m, 2H), δ 1.32 (m, 22H), 0.81 (t, J = 5.4 Hz, 3H) ppm; ¹³C NMR (75 MHz, CDCl₃) δ 176.8, 174.3, 173.9, 149.5, 148.3, 121.9, 106.4, 85.0, 67.2, 51.1, 49.5, 39.5, 39.1, 38.1, 36.8, 36.7, 32.1, 29.9, 29.8, 29.7, 29.6, 29.5, 29.3, 28.9, 28.3, 28.2, 27.9, 25.9, 25.6, 22.9, 14.3 ppm; HRMS (ESI) calcd for C₃₃H₅₉N₇O₃ (M⁺) 601.4679, found 601.4671, which was subsequently deprotected to give N-(2-(4-(5-(2-amino-1H-imidazol-4-yl)pentyl)-1H-1,2,3-triazol-1-yl)ethyl)palmitamide hydrochloride (0.155 g, 92 % yield). ¹H NMR (300 MHz, CD₃OD) δ 8.59 (s, 1H), δ 6.53 (s, 1H), δ 4.72 (s, 2H), δ 3.75 (s, 2H), δ 2.90 (s, 2H), δ 2.54 (s, 2H), δ 2.16 (t, J = 7.2 Hz, 2H), δ 1.80 (s, 2H), δ 1.69 (s, 2H), δ 1.51 (s, 4H), δ 1.28 (bs, 24H), δ 0.71 (t, J = 6.3 Hz, 3H) ppm; ¹³C NMR (75 MHz, CD₃OD) δ 175.7, 147.2, 145.6, 128.4, 127.6, 108.6, 67.2, 52.7, 37.4, 36.5, 35.8, 31.9, 29.7, 29.5, 29.4, 29.2, 28.0, 27.9, 27.8, 27.7, 25.8, 24.1, 23.6, 23.2, 22.6 ppm; HRMS (ESI) calcd for C₂₈H₅₁N₇O (M⁺) 501.4155, found 501.4143.

N-(2-(4-(5-(2-amino-1H-imidazol-4-yl)pentyl)-1H-1,2,3-triazol-1-yl)ethyl)-4,5-dibromo-1-methyl-1H-pyrrole-2-carboxamide hydrochloride:

tert-butyl 2-amino-4-(hept-6-ynyl)-1H-imidazole-1-carboxylate (0.116 g, 0.417 mmol) was reacted with N-(2-azidoethyl)-4,5-dibromo-1-methyl-1H-pyrrole-2-carboxamide (0.145 g, 0.417 mmol) following the general click procedure to give tert-butyl 2-amino-4-(5-(1-(2-(4,5-dibromo-1-methyl-1H-pyrrole-2-carboxamido)ethyl)-1H-1,2,3-triazol-4-yl)pentyl)-1H-imidazole-1-carboxylate ¹H NMR (300 MHz, CDCl₃) δ 7.76 (s, 1H), δ 7.32 (s, 1H), δ 6.81 (s, 1H), δ 6.49 (s, 1H), δ 6.17 (s, 2H), δ 4.52 (s, 2H), δ 3.94 (s, 3H), δ 3.84 (s, 2H), δ 2.65 (t, J = 7.2 Hz, 2H), δ 2.30 (m, 2H), δ 1.58 (m, 13H), δ 1.35 (m, 2H); ¹³C NMR (75 MHz, CDCl₃) δ 161.2, 149.6, 148.2, 138.8, 127.4, 122.2, 114.9, 111.9, 106.5, 98.2, 84.5, 49.5, 39.7, 35.9, 29.3, 28.9, 28.2, 28.0, 27.7, 25.6 ppm; HRMS (ESI) calcd for C₂₃H₃₂N₈O₃ (M⁺) 626.0964, found 626.0960, which was subsequently deprotected to give N-(2-(4-(5-(2-amino-1H-imidazol-4-yl)pentyl)-1H-1,2,3-triazol-1-yl)ethyl)-4,5-dibromo-1-methyl-1H-pyrrole-2-carboxamide hydrochloride (0.151 g, 64 % yield). ¹H NMR (400 MHz, CD₃OD) δ 8.24 (s, 1H), δ 6.86 (s, 1H), δ 6.56 (s, 1H), δ 4.71 (s, 2H), δ 3.81 (s, 5H), 2.78 (s, 2H), δ 2.44 (t, J = 7.2 Hz, 2H), δ 1.69 (s, 2H), δ 1.59 (s, 2H), δ 1.37 (s, 2H) ppm; ¹³C NMR (100 MHz, CD₃OD) δ 161.4, 150.2, 159.8, 159.4, 147.3, 127.6, 127.3, 114.9, 111.6, 108.4, 97.9, 51.3, 39.0, 35.2, 28.3, 28.0, 27.6, 24.2, 23.8 ppm; HRMS (ESI) calcd for C₁₈H₂₄Br₂N₈O (M⁺) 526.0439, found 526.0425.

N-(2-(4-(5-(2-amino-1H-imidazol-4-yl)pentyl)-1H-1,2,3-triazol-1-yl)ethyl)-4-bromo-1H-pyrrole-2-carboxamide hydrochloride: tert-butyl 2-amino-4-(hept-6-ynyl)-1H-imidazole-1-carboxylate (0.104 g, 0.375 mmol) was reacted with N-(2-azidoethyl)-4-bromo-1H-pyrrole-2-carboxamide (0.131 g, 0.375 mmol) following the general click procedure to give tert-butyl 2-amino-4-(5-(1-(2-(4-bromo-1H-pyrrole-2-carboxamido)ethyl)-1H-1,2,3-triazol-4-yl)pentyl)-1H-imidazole-1-carboxylate ¹H NMR (300 MHz, CDCl₃) δ 11.44 (s, 1H), δ 7.84 (s, 1H), δ 7.28 (s, 1H), δ 6.84 (s, 1H), δ 6.74 (s, 1H), δ 6.46 (s, 1H), δ 4.89 (s, 2H), δ 3.82 (s, 2H), δ 2.57 (s, 2H), δ 2.28 (s, 2H), δ 1.55 (m, 13H), δ 1.29 (m, 2H) ppm; ¹³C NMR (75 MHz, CDCl₃) δ 186.8, 161.4, 150.7, 149.3, 148.2, 137.6, 126.3, 122.5, 122.1, 113.1, 100.4, 96.9, 85.4, 49.6, 29.1, 28.7, 28.2, 27.7, 25.5 ppm; HRMS (ESI) calcd for C₂₂H₃₁BrN₈O₃ (M⁺) 534.1703, found 534.1705, which was subsequently deprotected to give N-(2-(4-(5-(2-amino-1H-imidazol-4-yl)pentyl)-1H-1,2,3-triazol-1-yl)ethyl)-4-bromo-1H-pyrrole-2-carboxamide hydrochloride (0.129 g, 73 % yield). ¹H NMR (400 MHz, CD₃OD) δ 8.42 (s, 1H), δ 6.93 (s, 1H), δ 6.82 (s, 1H), δ 6.47 (s, 1H), δ 4.76 (s, 2H), δ 3.87 (s, 2H), δ 2.79 (s, 2H), δ 2.45 (s, 2H), δ 1.69 (s, 2H), δ 1.60 (s, 2H), δ 1.37 (s, 2H) ppm; ¹³C NMR (100 MHz, CD₃OD) δ 161.4, 159.5, 159.1, 147.3, 147.2, 127.6, 125.8, 122.3, 122.1, 112.8, 108.6, 108.5, 96.4, 51.9, 38.9, 28.0, 27.9, 27.6, 24.1, 23.6 ppm; HRMS (ESI) calcd for C₁₇H₂₃BrN₈O (M⁺) 434.1178, found 434.1171.

N-(2-(4-(5-(2-amino-1H-imidazol-4-yl)pentyl)-1H-1,2,3-triazol-1-yl)ethyl)-1H-pyrrole-2-carboxamide hydrochloride: tert-butyl 2-amino-4-(hept-6-ynyl)-1H-imidazole-1-carboxylate (0.097 g, 0.349 mmol) was reacted with N-(2-azidoethyl)-1H-pyrrole-2-carboxamide (0.063 g, .349 mmol) following the general click procedure to give tert-butyl 4-(5-(1-(2-(1H-pyrrole-2-carboxamido)ethyl)-1H-1,2,3-triazol-4-yl)pentyl)-2-amino-1H-imidazole-1-carboxylate ¹H NMR (300 MHz, CDCl₃) δ 8.18 (s, 1H), δ 7.36 (d, J = 5.7 Hz, 2H), δ 7.24 (s, 1H), δ 6.84 (t, J = 8.1 Hz, 1H), δ 6.43 (s, 1H), δ 6.04 (s, 2H), 4.49 (s, 2H), δ 3.89 (s, 2H), δ 2.53 (s, 2H), δ 2.09 (s, 2H), δ 1.44 (m, 13H), δ 1.24 (s, 2H) ppm; ¹³C NMR (75 MHz, CDCl₃) δ 165.9, 164.8, 164.6, 161.5, 137.5, 122.3, 111.0, 110.7, 107.5, 107.1, 106.8, 85.1, 49.4, 40.5, 29.3, 28.9, 28.2, 25.5 ppm; HRMS (ESI) calcd for C₂₂H₃₂N₈O₃ (M⁺) 456.2597, found 456.2590, which was subsequently deprotected to give N-(2-(4-(5-(2-amino-1H-imidazol-4-yl)pentyl)-1H-1,2,3-triazol-1-yl)ethyl)-1H-pyrrole-2-carboxamide hydrochloride (0.129 g, 94 % yield). ¹H NMR (400 MHz, CD₃OD) δ 8.65 (s, 1H), δ 7.41 (s, 2H), δ 7.17 (s, 1H), δ 6.50 (s, 1H), δ 4.74 (s, 2H), δ 3.98 (s, 2H), δ 2.88 (s, 2H), δ 2.49 (s, 2H), δ 1.78 (m, 4H), δ 1.44 (s, 2H) ppm; ¹³C NMR (100 MHz, CD₃OD) δ 166.3, 164.5, 162.0, 161.9, 137.4, 110.6, 110.3, 110.2, 108.5, 17.1, 106.8, 59.3, 52.4, 39.5, 36.5, 27.9, 27.8, 27.6, 24.1, 23.5, 23.3 ppm; HRMS (ESI) calcd for C₁₇H₂₄N₈O (M⁺) 356.2073, found 356.2080.

N-(2-(4-(5-(2-amino-1H-imidazol-4-yl)pentyl)-1H-1,2,3-triazol-1-yl)ethyl)-4-tert-butylbenzamide hydrochloride: tert-butyl 2-amino-4-(hept-6-ynyl)-1H-imidazole-1-carboxylate (0.070 g, 0.253 mmol) was reacted with N-(2-azidoethyl)-4-tert-butylbenzamide (0.062 g, 0.253 mmol) following the general click procedure to give tert-butyl 2-amino-4-(5-(1-(2-(4-tert-butylbenzamido)ethyl)-1H-1,2,3-triazol-4-yl)pentyl)-1H-imidazole-1-carboxylate ¹H NMR (300 MHz, CDCl₃) δ 7.71 (d, J = 7.5 Hz, 2H), δ 7.64 (t, J = 5.7 Hz, 1H), δ 7.36 (d, J = 7.5 Hz, 2H), δ 7.27 (s, 1H), δ 6.42 (s, 1H), δ 5.99 (s, 2H), δ 4.51 (t, J = 5.7 Hz, 2H), δ 3.88 (q, J = 5.1, 5.3 Hz, 2H), δ 2.58 (t, J = 7.5 Hz, 2H), δ 2.22 (t, J = 7.5 Hz, 2H), δ 1.52 (m, 13H), δ 1.21 (m, 11H) ppm; ¹³C NMR (75 MHz, CDCl₃) δ 168.2, 155.3, 150.4, 149.6, 148.3, 138.9, 131.2, 127.3, 125.6, 122.2, 106.5, 84.8, 49.4, 40.2, 35.1, 31.3, 31.1, 29.3, 28.9, 28.3, 28.2, 28.1, 25.6 ppm; HRMS (ESI) calcd for C₂₈H₄₁N₇O₃ (M⁺) 523.3271, found 523.3270, which was subsequently deprotected to give N-(2-(4-(5-(2-amino-1H-imidazol-4-yl)pentyl)-1H-1,2,3-triazol-1-yl)ethyl)-4-tert-butylbenzamide hydrochloride (0.105 g, 90 % yield). ¹H NMR (400 MHz, CD₃OD) δ 8.48 (s, 1H), δ 7.76 (d, J = 6.3 Hz, 2H), δ 7.48 (d, J = 6.3 Hz, 2H), δ 6.50 (s, 1H), δ 4.84 (s, 2H), δ 3.95 (s, 2H), δ 2.85 (s, 2H), δ 2.46 (t, J = 5.4 Hz, 2H), δ 1.73 (s, 2H), δ 1.57 (m, 2H), δ 1.39 (s, 2H), δ 1.31 (s, 9H) ppm; ¹³C NMR (100 MHz, CD₃OD) δ 169.2, 155.6, 147.3, 130.8, 129.9, 127.6, 127.2, 127.1, 126.7, 125.4, 125.3, 119.5, 108.6, 108.5, 52.3, 39.4, 34.6, 30.4, 27.9, 27.8, 27.6, 24.1, 23.2, 22.2 ppm; HRMS (ESI) calcd for C₂₃H₃₃N₇O (M⁺) 423.2747, found 423.2741.

N-(2-(4-(5-(2-amino-1H-imidazol-4-yl)pentyl)-1H-1,2,3-triazol-1-yl)ethyl)-2-**naphthamide hydrochloride:** tert-butyl 2-amino-4-(hept-6-ynyl)-1H-imidazole-1-carboxylate

(0.095 g, 0.344 mmol) was reacted with N-(2-azidoethyl)-2-naphthamide (0.083 g, 0.344 mmol) following the general click procedure to give tert-butyl 4-(5-(1-(2-(2-naphthamido)ethyl)-1H-

1,2,3-triazol-4-yl)pentyl)-2-amino-1H-imidazole-1-carboxylate ¹H NMR (300 MHz, CDCl₃) δ

8.24 (s, 1H), δ 8.01 (s, 1H), δ 7.72 (m, 4H), δ 7.39 (m, 2H), δ 7.26 (s, 1H), δ 6.33 (s, 1H), δ 6.04

(s, 2H), δ 4.51 (s, 2H), δ 3.88 (s, 2H), δ 2.48 (t, 2H), δ 2.12 (s, 2H), δ 1.47 (m, 13H), δ 1.17 (s,

2H) ppm; ¹³C NMR (75 MHz, CDCl₃) δ 168.5, 150.4, 149.5, 148.3, 138.6, 134.9, 132.7, 131.4,

129.2, 128.5, 128.1, 127.9, 126.8, 124.0, 122.3, 106.5, 84.9, 49.4, 40.4, 29.9, 29.2, 28.9, 28.6,

28.2, 27.9, 28.6, 27.9, 27.8, 25.6 ppm; HRMS (ESI) calcd for C₂₈H₃₅N₇O₃ (M⁺) 517.2801, found

517.2792, which was subsequently deprotected to give N-(2-(4-(5-(2-amino-1H-imidazol-4-

yl)pentyl)-1H-1,2,3-triazol-1-yl)ethyl)-2-naphthamide hydrochloride (0.140 g, 90 % yield). ¹HNMR (300 MHz, CD₃OD) δ ppm; ¹³C NMR (75 MHz, CD₃OD) δ ppm; HRMS (ESI) calcd forC₂₃H₂₇N₇O (M⁺) 417.2277, found 417.2274.**N-(2-(4-(5-(2-amino-1H-imidazol-4-yl)pentyl)-1H-1,2,3-triazol-1-yl)ethyl)-2-****iodobenzamide hydrochloride:** tert-butyl 2-amino-4-(hept-6-ynyl)-1H-imidazole-1-carboxylate

(0.083 g, 0.299 mmol) was reacted with N-(2-azidoethyl)-2-iodobenzamide (0.095 g, 0.299

mmol) following the general click procedure to give tert-butyl 2-amino-4-(5-(1-(2-(2-

iodobenzamido)ethyl)-1H-1,2,3-triazol-4-yl)pentyl)-1H-imidazole-1-carboxylate ¹H NMR (300MHz, CDCl₃) δ 7.74 (d, J = 7.5 Hz, 2H), δ 7.57 (t, J = 4.5 Hz, 1H), δ 7.36 (s, 1H), δ 7.22 (s, 2H),

δ 6.97 (m, 1H), δ 6.39 (s, 1H), δ 5.97 (s, 2H), δ 4.48 (t, J = 5.1 Hz, 2H), δ 3.85 (m, 2H), δ 2.44 (t,

J = 7.5 Hz, 2H), δ 2.18 (t, J = 6.6 Hz, 2H), δ 1.45 (m, 13H), δ 1.18 (m, 2H) ppm; ¹³C NMR (75MHz, CDCl₃) δ 170.3, 150.4, 149.5, 148.2, 141.8, 139.9, 138.7, 131.3, 128.4, 128.3, 122.3,

106.5, 92.8, 84.9, 49.2, 40.5, 29.2, 28.9, 28.3, 28.2, 28.0, 25.5 ppm; HRMS (ESI) calcd for

C₂₄H₃₂IN₇O₃ (M⁺) 593.1611, found 593.1603, which was subsequently deprotected to give N-(2-

(4-(5-(2-amino-1H-imidazol-4-yl)pentyl)-1H-1,2,3-triazol-1-yl)ethyl)-2-iodobenzamide

hydrochloride (0.125 g, 79 % yield). ¹H NMR (400 MHz, CD₃OD) δ 8.35 (s, 1H), δ 7.88 (d, J =

7.8 Hz, 1H), δ 7.45 (t, J = 6.9 Hz, 1H), δ 7.34 (d, J = 6.9 Hz, 1H), δ 7.15 (t, J = 7.5 Hz, 1H), δ

6.47 (s, 1H), δ 4.78 (s, 2H), δ 3.99 (s, 2H), δ 2.83 (s, 2H), δ 2.44 (t, J = 6.9 Hz, 2H), δ 1.76 (s,

2H), δ 1.64 (s, 2H), δ 1.29 (s, 2H) ppm; ¹³C NMR (75 MHz, CD₃OD) δ 171.6, 159.2, 147.3,

142.1, 139.8, 131.2, 128.2, 127.7, 124.4, 92.1, 52.2, 40.8, 39.4, 31.2, 28.3, 28.1, 27.7, 24.1, 20.9,

15.9, 10.7 ppm; HRMS (ESI) calcd for C₁₉H₂₄IN₇O (M⁺) 493.1087, found 493.1085.**N-(2-(4-(5-(2-amino-1H-imidazol-4-yl)pentyl)-1H-1,2,3-triazol-1-yl)ethyl)-4-****heptylbenzamide hydrochloride:** tert-butyl 2-amino-4-(hept-6-ynyl)-1H-imidazole-1-

carboxylate (0.125 g, 0.449 mmol) was reacted with N-(2-azidoethyl)-4-heptylbenzamide (0.129

g, 0.449 mmol) following the general click procedure to give tert-butyl 2-amino-4-(5-(1-(2-(4-heptylbenzamido)ethyl)-1H-1,2,3-triazol-4-yl)pentyl)-1H-imidazole-1-carboxylate ¹H NMR (400 MHz, CDCl₃) δ 7.78 (s, 1H), δ 7.65 (d, 2H), δ 7.27 (s, 1H), δ 7.09 (d, 2H), δ 6.39 (s, 1H), δ 6.22 (s, 2H), δ 4.47 (s, 2H), δ 3.81 (s, 2H), δ 3.96 (s, 4H), δ 2.20 (s, 2H), δ 2.05 (s, 2H), 1.48 (m, 12H), δ 1.17 (m, 8H), δ 0.77 (t, J = 6.4 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 168.3, 149.6, 148.2, 147.1, 138.6, 131.5, 128.6, 127.4, 122.1, 106.4, 84.8, 67.1, 49.4, 40.2, 37.9, 36.9, 31.9, 31.3, 31.1, 29.8, 28.4, 29.3, 28.9, 28.3, 28.1, 28.0, 25.5, 22.8, 14.3 ppm; HRMS (ESI) calcd for C₃₁H₄₇N₇O₃ (M⁺) 565.3704, found 656.3737, which was subsequently deprotected to give N-(2-(4-(5-(2-amino-1H-imidazol-4-yl)pentyl)-1H-1,2,3-triazol-1-yl)ethyl)-4-heptylbenzamide hydrochloride (0.194 g, 86 % yield). ¹H NMR (300 MHz, CD₃OD) δ 8.25 (s, 1H), δ 7.71 (d, J = 5.4 Hz, 2H), δ 7.24 (d, J = 5.7 Hz, 2H), δ 6.45 (s, 1H), δ 4.74 (s, 2H), 3.90 (s, 2H), δ 2.63 (s, 2H), δ 2.61 (t, 2H), δ 2.41 (s, 2H), δ 1.58 (m, 6H), δ 1.26 (m, 10 H), δ 0.85 (t, J = 4.8 Hz, 3H) ppm; ¹³C NMR (100 MHz, CD₃OD) δ 169.2, 150.2, 147.4, 131.2, 128.5, 127.6, 127.3, 108.4, 51.4, 39.6, 36.5, 35.6, 31.8, 331.3, 29.1, 29.1, 28.3, 28.0, 27.7, 24.1, 22.6, 13.4, 13.3 ppm; HRMS (ESI) calcd for C₂₆H₃₉N₇O (M⁺) 465.3216, found 465.3207.

N-(2-(4-(5-(2-amino-1H-imidazol-4-yl)pentyl)-1H-1,2,3-triazol-1-yl)ethyl)-4-hexylbenzamide hydrochloride: tert-butyl 2-amino-4-(hept-6-ynyl)-1H-imidazole-1-carboxylate (0.097 g, 0.349 mmol) was reacted with N-(2-azidoethyl)-4-hexylbenzamide (0.096 g, 0.349 mmol) following the general click procedure to give tert-butyl 2-amino-4-(5-(1-(2-(4-hexylbenzamido)ethyl)-1H-1,2,3-triazol-4-yl)pentyl)-1H-imidazole-1-carboxylate ¹H NMR (300 MHz, CDCl₃) δ 7.65 (d, J = 8.1 Hz, 2H), δ 7.55 (t, J = 6.7 Hz, 1H), δ 7.26 (s, 1H), δ 7.09 (d, J = 7.8 Hz, 2H), δ 6.39 (s, 1H), δ 6.04 (s, 2H), δ 4.48 (t, J = 4.8 Hz, 2H), δ 3.83 (q, J = 5.6, 4.8 Hz, 2H), δ 2.56 (q, J = 7.5, 8.1 Hz, 4H), δ 2.24 (t, J = 6.9 Hz, 2H), δ 1.50 (m, 13H), δ 1.21 (m, 1H), δ 0.77 (t, J = 5.7 Hz, 3H) ppm; ¹³C NMR (75 MHz, CDCl₃) δ 168.3, 148.3, 147.3, 138.7, 131.4, 128.7, 127.4, 122.2, 106.4, 84.9, 73.9, 49.5, 40.2, 36.0, 31.8, 31.3, 29.9, 29.3, 29.1, 28.9, 28.2, 28.1, 28.0, 25.6, 22.8, 14.3 ppm; HRMS (ESI) calcd for C₃₀H₄₅N₇O₃ (M⁺) 551.3584, found 551.3581, which was subsequently deprotected to give N-(2-(4-(5-(2-amino-1H-imidazol-4-yl)pentyl)-1H-1,2,3-triazol-1-yl)ethyl)-4-hexylbenzamide hydrochloride (0.164 g, 96 % yield). ¹H NMR (400 MHz, CD₃OD) δ 8.28 (s, 1H), δ 7.67 (d, J = 7.6 Hz, 2H), δ 7.21 (d, J = 7.6 Hz, 2H), δ 6.43 (s, 1H), δ 4.72 (s, 2H), δ 3.87 (s, 2H), δ 2.74 (s, 2H), δ 2.58 (t, J = 7.6 Hz, 2H), δ 2.31 (t, J = 6.8 Hz, 2H), δ 1.66 (s, 2H), δ 1.55 (s, 4H), δ 1.35 (s, 2H), δ 1.25 (s, 6H), δ 0.66 (t, J = 8.8 Hz, 3H) ppm; ¹³C NMR (100 MHz, CD₃OD) δ 169.2, 159.5, 147.4, 147.3, 131.2, 108.4, 51.2, 48.5, 48.3, 48.1, 47.9, 47.7, 47.5, 47.3, 39.5, 36.5, 35.6, 31.6, 31.7, 31.2, 28.8, 28.2, 27.9, 27.7, 24.1, 23.9, 22.6, 13.3, 9.9 ppm; HRMS (ESI) calcd for C₂₅H₃₇N₇O (M⁺) 451.3059, found 451.3058.

N-(2-(4-(5-(2-amino-1H-imidazol-4-yl)pentyl)-1H-1,2,3-triazol-1-yl)ethyl)-4-**butylbenzamide hydrochloride:** tert-butyl 2-amino-4-(hept-6-ynyl)-1H-imidazole-1-

carboxylate (0.123 g, 0.443 mmol) was reacted with N-(2-azidoethyl)-4-butylbenzamide (0.109

g, 0.443 mmol) following the general click procedure to give tert-butyl 2-amino-4-(5-(1-(2-(4-

butylbenzamido)ethyl)-1H-1,2,3-triazol-4-yl)pentyl)-1H-imidazole-1-carboxylate ¹H NMR (300MHz, CDCl₃) δ 7.67 (s, 1H), δ 7.67 (d, J = 7.5 Hz, 2H), δ 7.29 (s, 1H), δ 7.11 (d, J = 7.5 Hz,

2H), δ 6.42 (s, 1H), δ 6.25 (s, 2H), δ 4.48 (s, 2H), δ 3.83 (s, 2H), δ 2.56 (t, J = 7.2 Hz, 4H), δ

2.21 (s, 2H), 1.47 (m, 13H), δ 1.21 (m, 4H), δ 0.83 (t, J = 7.2 Hz, 3H) ppm; ¹³C NMR (75 MHz,CDCl₃) δ 168.3, 140.6, 149.5, 148.2, 147.1, 138.4, 131.5, 128.7, 127.4, 122.2, 106.4, 84.9, 49.4,

40.2, 35.7, 33.5, 29.2, 28.9, 28.3, 28.1, 27.9, 25.5, 22.4, 14.1 ppm; HRMS (ESI) calcd for

C₂₈H₄₁N₇O₃ (M⁺) 523.3271, found 523.3261, which was subsequently deprotected to give N-(2-

(4-(5-(2-amino-1H-imidazol-4-yl)pentyl)-1H-1,2,3-triazol-1-yl)ethyl)-4-butylbenzamide

hydrochloride (0.191 g, 94 % yield). ¹H NMR (300 MHz, CD₃OD) δ 8.43 (s, 1H), δ 7.69 (s, 2H),

δ 7.22 (s, 2H), δ 6.44 (s, 1H), δ 4.73 (s, 2H), δ 3.89 (s, 2H), δ 3.34 (s, 2H), δ 2.59 (s, 4H), δ 2.41

(s, 2H), δ 1.54 (s, 4H), δ 1.29 (s, 4H), δ 0.88 (t, J = 6.9 Hz, 3H) ppm; ¹³C NMR (75 MHz,CD₃OD) δ 169.2, 165.1, 160.3, 158.1, 147.4, 141.4, 131.2, 128.5, 127.7, 127.3, 126.9, 92.9, 39.6,

36.3, 33.4, 30.9, 28.0, 27.7, 24.1, 22.2, 13.2, 12.1, 10.9, 8.9, 6.2, 2.5 ppm; HRMS (ESI) calcd for

C₂₃H₃₃N₇O (M⁺) 423.2747, found 423.2738.**N-(2-(4-(5-(2-amino-1H-imidazol-4-yl)pentyl)-1H-1,2,3-triazol-1-yl)ethyl)-4-****pentylbenzamide hydrochloride:** tert-butyl 2-amino-4-(hept-6-ynyl)-1H-imidazole-1-

carboxylate (0.113 g, 0.408 mmol) was reacted with N-(2-(4-(5-(2-amino-1H-imidazol-4-

yl)pentyl)-1H-1,2,3-triazol-1-yl)ethyl)biphenyl-4-carboxamide hydrochloride (0.106 g, 0.408

mmol) following the general click procedure to give tert-butyl 2-amino-4-(5-(1-(2-(4-

pentylbenzamido)ethyl)-1H-1,2,3-triazol-4-yl)pentyl)-1H-imidazole-1-carboxylate ¹H NMR(300 MHz, CDCl₃) δ 7.75 (s, 1H), δ 7.67 (d, J = 7.5 Hz, 2H), δ 7.27 (s, 1H), δ 7.12 (d, J = 7.8

Hz, 2H), δ 4.48 (s, 2H), δ 3.84 (s, 2H), δ 2.55 (m, 4H), δ 2.22 (s, 2H), δ 1.50 (s, 13H), δ 1.23 (s,

8H), δ 0.80 (t, J = 5.7 Hz, 3H); ¹³C NMR (75 MHz, CDCl₃) δ ppm 168.3, 150.5, 149.5, 148.2,

147.2, 138.6, 131.5, 128.7, 127.4, 106.4, 84.8, 49.4, 40.2, 35.9, 31.6, 31.0, 29.3, 28.9, 28.1, 27.9,

25.6, 22.6, 14.2; HRMS (ESI) calcd for C₂₉H₄₃N₇O₃ (M⁺) 537.3427, found 537.3420, which was

subsequently deprotected to give N-(2-(4-(5-(2-amino-1H-imidazol-4-yl)pentyl)-1H-1,2,3-

triazol-1-yl)ethyl)-4-pentylbenzamide hydrochloride (0.178 g, 92 % yield). ¹H NMR (300 MHz,CD₃OD) δ 8.41 (s, 1H), δ 7.69 (d, J = 7.5 Hz, 2H), δ 7.21 (d, J = 7.2 Hz, 2H), δ 6.46 (s, 1H), δ

4.84 (s, 2H), δ 3.90 (s, 2H), δ 2.79 (s, 2H), δ 2.58 (t, J = 7.8 Hz, 2H), δ 2.39 (t, J = 7.2 Hz, 2H), δ

1.68 (s, 2H), δ 1.55 (s, 4H), δ 1.26 (s, 6H), δ 0.84 (t, J = 6.6 Hz, 3H) ppm; ¹³C NMR (75 MHz,CD₃OD) δ 169.2, 159.5, 158.9, 147.5, 147.3, 131.1, 128.5, 127.6, 127.4, 108.5, 54.2, 51.9, 39.5,

35.6, 31.4, 30.9, 30.6, 28.4, 27.9, 27.6, 26.9, 24.1, 23.5, 22.9, 22.4, 13.3 ppm; HRMS (ESI) calcd for $C_{24}H_{35}N_7O$ (M^+) 437.2903, found 437.2892.

N-(2-(4-(5-(2-amino-1H-imidazol-4-yl)pentyl)-1H-1,2,3-triazol-1-yl)ethyl)-2,4,6-trichlorobenzamide hydrochloride: tert-butyl 2-amino-4-(hept-6-ynyl)-1H-imidazole-1-

5 carboxylate (0.088 g, 0.318 mmol) was reacted with N-(2-azidoethyl)-2,4,6-trichlorobenzamide (0.093 g, 0.318 mmol) following the general click procedure to give tert-butyl 2-amino-4-(5-(1-(2-(2,4,6-trichlorobenzamido)ethyl)-1H-1,2,3-triazol-4-yl)pentyl)-1H-imidazole-1-carboxylate ^1H NMR (300 MHz, $CDCl_3$) δ 8.22 (s, 1H), δ 7.34 (s, 1H), δ 7.26 (s, 2H), δ 6.42 (s, 1H), δ 5.97 (s, 2H), δ 4.51 (s, 2H), δ 3.94 (s, 2H), δ 2.45 (t, 2H), δ 2.20 (s, 2H), δ 1.47 (m 13H), δ 1.21 (m, 10 2H) ppm; ^{13}C NMR (75 MHz, $CDCl_3$) δ 164.7, 148.0, 135.7, 134.6, 132.9, 128.7, 128.1, 127.3, 122.3, 84.9, 53.7, 49.3, 39.9, 31.8, 29.2, 28.9, 28.2, 25.5, 22.8, 14.3 ppm; HRMS (ESI) calcd for $C_{24}H_{30}Cl_3N_7O_3$ (M^+) 569.1476, found 569.1477, which was subsequently deprotected to give N-(2-(4-(5-(2-amino-1H-imidazol-4-yl)pentyl)-1H-1,2,3-triazol-1-yl)ethyl)-2,4,6-trichlorobenzamide hydrochloride (0.116 g, 72 % yield). ^1H NMR (300 MHz, CD_3OD) δ 8.36 (s, 15 1H), δ 7.45 (s, 2H), δ 6.46 (s, 1H), δ 4.77 (s, 2H), δ 3.97 (s, 2H), δ 2.80 (s, 2H), δ 2.48 (t, J = 6.9 Hz, 2H), δ 1.73 (s, 2H), δ 1.62 (s, 2H), δ 1.41 (s, 2H) ppm; ^{13}C NMR (75 MHz, CD_3OD) δ 165.5, 147.3, 136.9, 134.5, 132.7, 128.2, 128.0, 127.6, 108.6, 51.2, 39.0, 28.1, 27.7, 24.1, 23.8 ppm; HRMS (ESI) calcd for $C_{19}H_{22}Cl_3N_7O$ (M^+) 469.0951, found 469.0941.

N-(2-(4-(5-(2-amino-1H-imidazol-4-yl)pentyl)-1H-1,2,3-triazol-1-yl)ethyl)-3,5-difluorobenzamide hydrochloride: tert-butyl 2-amino-4-(hept-6-ynyl)-1H-imidazole-1-

20 carboxylate (0.123 g, 0.442 mmol) was reacted with N-(2-azidoethyl)-3,5-difluorobenzamide (0.100 g, 0.442 mmol) following the general click procedure to give tert-butyl 2-amino-4-(5-(1-(2-(3,5-difluorobenzamido)ethyl)-1H-1,2,3-triazol-4-yl)pentyl)-1H-imidazole-1-carboxylate ^1H NMR (300 MHz, $CDCl_3$) δ 8.25 (s, 1H), δ 7.39 (m, 3H), δ 6.88 (t, J = 4.8 Hz, 1H), δ 6.43 (s, 25 1H), δ 6.38 (bs, 2H), δ 4.55 (s, 2H), δ 3.93 (s, 2H), δ 2.56 (t, J = 5.1 Hz, 2H), δ 2.24 (s, 2H), δ 1.55 (m, 14H), δ 1.23 (s, 2H) ppm; ^{13}C NMR (75 MHz, $CDCl_3$) δ 165.9, 164.3, 164.2, 161.9, 150.3, 149.3, 148.2, 137.6, 122.4, 111.0, 110.8, 107.3, 107.1, 106.8, 106.6, 85.5, 49.3, 40.4, 29.9, 29.5, 29.1, 28.7, 28.2, 27.9, 27.5, 25.4 ppm; HRMS (ESI) calcd for $C_{24}H_{31}F_2N_7O_3$ (M^+) 503.2456, found 503.2458, which was subsequently deprotected to give N-(2-(4-(5-(2-amino-1H-imidazol-4-yl)pentyl)-1H-1,2,3-triazol-1-yl)ethyl)-3,5-difluorobenzamide hydrochloride 30 (0.115 g, 59 % yield). ^1H NMR (400 MHz, CD_3OD) δ 8.55 (s, 1H), δ 7.42 (d, J = 2.0 Hz, 2H), δ 7.29 (t, J = 2.40 Hz, 1H), δ 6.51 (s, 1H), δ 4.84 (t, J = 4.8 Hz, 2H), δ 3.96 (q, J = 13.2, 4.8 Hz, 2H), δ 2.89 (t, J = 7.2 Hz, 2H), δ 2.49 (t, J = 7.2 Hz, 2H), δ 1.79 (m, 2H), δ 1.75 (m, 2H), δ 1.44 (m, 2H) ppm; ^{13}C NMR (100 MHz, CD_3OD) δ 164.5, 164.4, 127.6, 126.9, 110.6, 110.6, 110.3,

108.5, 107.1, 106.8, 106.6, 52.5, 39.5, 30.4, 27.9, 27.8, 27.6, 24.0, 23.8, 23.1 ppm; HRMS (ESI) calcd for $C_{19}H_{23}F_2N_7O$ (M^+) 403.1932, found 403.1926.

N-(2-(4-(5-(2-amino-1H-imidazol-4-yl)pentyl)-1H-1,2,3-triazol-1-yl)ethyl)biphenyl-4-carboxamide hydrochloride: tert-butyl 2-amino-4-(hept-6-ynyl)-1H-imidazole-1-carboxylate

5 (0.093 g, 0.334 mmol) was reacted with N-(2-azidoethyl)biphenyl-4-carboxamide (0.089 g, 0.334 mmol) following the general click procedure to give tert-butyl 2-amino-4-(5-(1-(2-biphenyl-4-ylcarboxamidoethyl)-1H-1,2,3-triazol-4-yl)pentyl)-1H-imidazole-1-carboxylate ^1H NMR (300 MHz, $CDCl_3$) δ 7.91 (d, J = 5.1 Hz, 3H), δ 7.57 (t, J = 8.4 Hz, 4H), δ 7.44 (t, J = 6.9 Hz, 3H), δ 7.34 (1H), δ 4.60 (t, J = 5.1 Hz, 2H), δ 3.96 (q, J = 5.1, 5.4 Hz, 2H), δ 2.63 (t, J = 7.2 Hz, 2H), δ 2.27 (t, J = 7.5 Hz, 2H), δ 1.62 (m, 13H), δ 1.33 (m, 2H) ppm; ^{13}C NMR (75 MHz, $CDCl_3$) δ 168.0, 149.6, 148.3, 144.6, 140.1, 138.8, 132.8, 129.1, 128.2, 128.0, 127.4, 127.3, 122.2, 84.9, 49.4, 40.3, 29.3, 28.9, 28.3, 28.2, 28.1, 25.6 ppm; HRMS (ESI) calcd for $C_{30}H_{37}N_7O_3$ (M^+) 543.2958, found 543.2949, which was subsequently deprotected to give N-(2-(4-(5-(2-amino-1H-imidazol-4-yl)pentyl)-1H-1,2,3-triazol-1-yl)ethyl)biphenyl-4-carboxamide hydrochloride (0.093 g, 58 % yield). ^1H NMR (300 MHz, CD_3OD) δ 8.35 (s, 1H), δ 7.88 (d, J = 8.1 Hz, 2H), δ 7.65 (d, J = 8.4 Hz, 2H), δ 7.57 (d, J = 7.5 Hz, 2H), δ 7.39 (t, J = 7.5 Hz, 2H), δ 7.34 (t, J = 7.2 Hz, 1H), δ 6.39 (s, 1H), δ 4.81 (s, 2H), δ 3.96 (s, 2H), δ 2.78 (s, 2H), δ 2.37 (t, J = 7.5 Hz, 2H), δ 1.67 (s, 2H), δ 1.56 (t, 2H), δ 1.33 (s, 2H) ppm; ^{13}C NMR (75 MHz, CD_3OD) δ 203.6, 158.9, 160.1, 159.5, 159.1, 159.0, 158.5, 149.8, 147.2, 144.6, 132.4, 128.9, 127.9, 127.6, 126.9, 126.4, 108.4, 51.7, 39.5, 27.9, 27.6, 24.0, 23.5, 7.2 ppm; HRMS (ESI) calcd for $C_{25}H_{29}N_7O$ (M^+) 443.2434, found 443.2423.

N-(2-(4-(5-(2-amino-1H-imidazol-4-yl)pentyl)-1H-1,2,3-triazol-1-yl)ethyl)-4-octylbenzamide hydrochloride: tert-butyl 2-amino-4-(hept-6-ynyl)-1H-imidazole-1-carboxylate (0.078 g, 0.282 mmol) was reacted with N-(2-azidoethyl)-4-octylbenzamide (0.085

25 g, 0.282 mmol) following the general click procedure to give tert-butyl 2-amino-4-(5-(1-(2-(4-octylbenzamido)ethyl)-1H-1,2,3-triazol-4-yl)pentyl)-1H-imidazole-1-carboxylate ^1H NMR (300 MHz, $CDCl_3$) δ 7.64 (s, 2H), δ 7.47 (s, 1H), δ 7.21 (s, 1H), δ 7.09 (s, 2H), δ 6.47 (bs, 3H), δ 4.48 (s, 2H), δ 3.84 (s, 2H), δ 2.53 (s, 4H), δ 2.28 (t, J = 15.9 Hz, 2H), δ 1.51 (bs, 6H), δ 1.18 (bs, 10H), 0.77 (t, J = 6.6 Hz, 3H) ppm; ^{13}C NMR (75 MHz, $CDCl_3$) δ 180.7, 180.2, 168.3, 147.4, 142.4, 139.2, 131.4, 128.8, 127.4, 107.4, 49.6, 40.2, 36.1, 32.1, 31.4, 29.9, 29.6, 29.5, 28.8, 28.2, 28.1, 25.4, 22.9, 14.3 ppm; HRMS (ESI) calcd for $C_{32}H_{49}N_7O_3$ (M^+) 579.3896, found 579.3890, which was subsequently deprotected to give N-(2-(4-(5-(2-amino-1H-imidazol-4-yl)pentyl)-1H-1,2,3-triazol-1-yl)ethyl)-4-octylbenzamide hydrochloride (0.097 g, 67 % yield). ^1H NMR (400 MHz, CD_3OD) δ 7.65 (s, 2H), δ 7.23 (s, 3H), δ 6.45 (s, 1H), δ 3.97 (s, 4H), δ 2.60 (s, 4H), δ 2.47 (s, 2H), δ 1.57 (m, 6H), δ 1.27 (bm, 12H), δ 0.83 (t, J = 6.4 Hz, 3H) ppm; ^{13}C NMR (100 MHz,

CD₃OD) δ 183.5, 172.0, 169.3, 161.8, 153.9, 147.4, 131.2, 128.5, 127.7, 127.3, 119.1, 108.8, 104.7, 94.6, 63.4, 38.7, 35.6, 31.8, 31.3, 29.3, 29.2, 29.1, 28.1, 27.7, 27.1, 24.2, 22.5, 13.3, 12.6 ppm; HRMS (ESI) calcd for C₂₇H₄₁N₇O (M⁺) 479.3372, found 479.3378.

N-(2-(4-(5-(2-amino-1H-imidazol-4-yl)pentyl)-1H-1,2,3-triazol-1-yl)ethyl)-4-

5 **nonylbenzamide hydrochloride:** tert-butyl 2-amino-4-(hept-6-ynyl)-1H-imidazole-1-carboxylate (g, mmol) was reacted with (g, mmol) following the general click procedure to give tert-butyl 2-amino-4-(5-(1-(2-(4-nonylbenzamido)ethyl)-1H-1,2,3-triazol-4-yl)pentyl)-1H-imidazole-1-carboxylate ¹H NMR (300 MHz, CDCl₃) δ 7.70 (d, J = 7.2 Hz, 2H), δ 7.29 (s, 1H), δ 7.27 (t, J = 8.7 Hz, 1H), δ 7.16 (d, J = 7.2 Hz, 2H), δ 4.53 (s, 2H), δ 3.91 (s, 2H), δ 2.61 (m, 4H), δ 2.41 (m, 1H), δ 2.14 (s, 1H), δ 1.56 (bs, 15H), δ 1.23 (bs, 14H), δ 0.83 (t, J = 6.6 Hz, 3H) ppm; ¹³C NMR (75 MHz, CDCl₃) δ 188.1, 169.7, 168.3, 164.4, 148.0, 147.4, 131.4, 128.8, 127.4, 122.2, 51.6, 49.5, 40.2, 36.0, 32.1, 31.4, 29.7, 29.6, 29.5, 29.4, 29.3, 28.2, 25.6, 22.9, 14.3 ppm; HRMS (ESI) calcd for C₃₃H₅₁N₇O₃ (M⁺) 593.4053, found 593.4049, which was subsequently deprotected to give N-(2-(4-(5-(2-amino-1H-imidazol-4-yl)pentyl)-1H-1,2,3-triazol-1-yl)ethyl)-4-nonylbenzamide hydrochloride (0.183 g, 74 % yield). ¹H NMR (400 MHz, CD₃OD) δ 8.66 (s, 1H), δ 7.69 (s, 2H), δ 7.25 (s, 2H), δ 6.54 (s, 1H), δ 4.81 (s, 2H), δ 3.62 (s, 2H), δ 2.85 (s, 2H), δ 2.63 (s, 2H), δ 2.46 (s, 2H), δ 1.75 (s, 2H), δ 1.39 (s, 4H), δ 1.29 (m, 14H), δ 0.88 (s, 3H) ppm; ¹³C NMR (100 MHz, CD₃OD) δ 205.0, 169.2, 147.5, 147.3, 131.2, 128.5, 127.3, 108.9, 108.6, 96.0, 52.4, 51.6, 39.3, 38.9, 35.6, 31.9, 31.3, 30.6, 29.5, 29.4, 29.3, 29.2, 28.1, 27.9, 27.7, 24.1, 23.6, 22.6, 13.4 ppm; HRMS (ESI) calcd for C₂₈H₄₃N₇O (M⁺) 493.3529, found 493.3522.

N-(2-(4-(5-(2-amino-1H-imidazol-4-yl)pentyl)-1H-1,2,3-triazol-1-yl)ethyl)-2',4'-difluoro-3-hydroxybiphenyl-4-carboxamide hydrochloride: tert-butyl 2-amino-4-(hept-6-ynyl)-1H-imidazole-1-carboxylate (0.085 g, 0.306 mmol) was reacted with N-(2-azidoethyl)-2',4'-difluoro-3-hydroxybiphenyl-4-carboxamide (0.097 g, 0.306mmol) following the general click procedure to give tert-butyl 2-amino-4-(5-(1-(2-(2',4'-difluoro-3-hydroxybiphenyl-4-ylcarboxamido)ethyl)-1H-1,2,3-triazol-4-yl)pentyl)-1H-imidazole-1-carboxylate ¹H NMR (300 MHz, CDCl₃) δ 8.54 (s, 1H), δ 7.83 (s, 1H), δ 7.37 (d, J = 7.2 Hz, 1H), δ 7.28 (s, 1H), δ 7.18 (s, 1H), δ 6.89 (d, J = 7.8 Hz, 1H), δ 6.72 (s, 2H), δ 6.39 (s, 1H), δ 4.57 (s, 2H), δ 3.81 (s, 2H), δ 3.34 (s, 1H), δ 2.47 (s, 2H), δ 2.07 (s, 2H), δ 1.49 (m, 13H), δ 1.17 (s, 2H) ppm; ¹³C NMR (75 MHz, CDCl₃) δ 169.5, 163.7, 161.3, 160.5, 159.9, 150.5, 149.2, 148.2, 137.1, 136.3, 134.5, 131.4, 12.8, 126.8, 124.4, 122.3, 117.9, 116.1, 111.9, 111.6, 106.7, 104.8, 104.4, 104.1, 85.8, 50.5, 49.4, 46.6, 50.0, 37.9, 29.9, 28.9, 28.5, 28.3, 28.1, 27.4, 25.3, 25.0 ppm; HRMS (ESI) calcd for C₃₀H₃₅F₂N₄O₄ (M⁺) 595.2729, found 595.2717, which was subsequently deprotected to give (0.137 g, 84 % yield). ¹H NMR (400 MHz, CD₃OD) δ 8.60 (s, 1H), δ 7.89 (s, 1H), δ 7.52 (s, 2H), δ 6.97 (t, J = 6.3 Hz,

3H), δ 6.45 (s, 1H), δ 4.81 (s, 2H), δ 2.78 (s, 2H), δ 2.42 (s, 2H), δ 1.72 (s, 2H), δ 1.61 (d, J = 7.8 Hz, 2H), δ 1.38 (s, 2H) ppm; ^{13}C NMR (100 MHz, CD_3OD) δ 172.9, 169.8, 161.6, 159.5, 147.2, 134.4, 131.7, 128.4, 127.6, 126.0, 117.5, 115.6, 111.7, 111.5, 108.5, 104.2, 104.0, 103.7, 94.6, 51.9, 39.0, 27.9, 27.6, 26.8, 24.1, 23.8 ppm; HRMS (ESI) calcd for $\text{C}_{25}\text{H}_{27}\text{F}_2\text{N}_7\text{O}_2$ (M+)

459.2194, found 459.2190.

Procedure to Determine the Inhibitory Effect of Test Compounds on *C. albicans*, *C. neoformans* and a mixed *S. epidermidis*/*C. albicans* Biofilm Formation: Inhibition assays were performed by taking an overnight culture of yeast or yeast/bacteria strain and subculturing it at an OD_{600} of 0.05 into YPD (Yeast extract, peptone and dextrose (BD 242820)) media for the yeast alone or tryptic soy broth for the *S. epidermidis*/*C. albicans*. Stock solutions of predetermined concentrations of the test compound were then made in the necessary media. These stock solutions were aliquoted (100 μL) into the wells of the 96-well PVC microtiter plate. Sample plates were then wrapped in GLAD Press n' Seal[®] followed by an incubation under stationary conditions for 24 h at 37 °C. After incubation, the media was discarded from the wells and the plates were washed thoroughly with water. Plates were then stained with 100 μL of 0.1% solution of crystal violet (CV) and then incubated at ambient temperature for 30 min. Plates were washed with water again and the remaining stain was solubilized with 200 μL of 95% ethanol. A sample of 125 μL of solubilized CV stain from each well was transferred to the corresponding wells of a polystyrene microtiter dish. Biofilm inhibition was quantitated by measuring the OD_{540} of each well in which a negative control lane wherein no biofilm was formed served as a background and was subtracted out.


Procedure to Determine the Dispersal Effect of Test Compounds on *C. albicans* and *C. neoformans* Preformed Biofilms: Dispersion assays were performed by taking an overnight culture of bacterial strain and subculturing it at an OD_{600} of 0.01 into Yeast extract, peptone and dextrose (BD 242820) media. The resulting bacterial suspension was aliquoted (100 μL) into the wells of a 96-well PVC microtiter plate. Plates were then wrapped in GLAD Press n' Seal[®] followed by an incubation under stationary conditions at ambient temperature to establish the biofilms. After 24 h, the media was discarded from the wells and the plates were washed thoroughly with water. Stock solutions of predetermined concentrations of the test compound were then made in the necessary media. These stock solutions were aliquoted (100 μL) into the wells of the 96-well PVC microtiter plate with the established biofilms. Media alone was added to a subset of the wells to serve as a control. Sample Plates were then incubated for 24 h at 37 °C. After incubation, the media was discarded from the wells and the plates were washed thoroughly with water. Plates were then stained with 100 μL of 0.1% solution of crystal violet (CV) and then incubated at ambient temperature for 30 min. Plates were washed with water

again and the remaining stain was solubilized with 200 μ L of 95% ethanol. A sample of 125 μ L of solubilized CV stain from each well was transferred to the corresponding wells of a polystyrene microtiter dish. Biofilm dispersion was quantitated by measuring the OD₅₄₀ of each well in which a negative control lane wherein no biofilm was formed served as a background and was subtracted out.

The foregoing is illustrative of the present invention, and is not to be construed as limiting thereof. The invention is defined by the following claims, with equivalents of the claims to be included therein.

THAT WHICH IS CLAIMED IS:

1. A method of preventing, removing or inhibiting microbial biofilm formation or microbial infection in a plant or plant part thereof, comprising applying to said plant or plant part a treatment effective amount of a compound of **Formula (II)(a)**:



(II)(a)

wherein:

R^{1a} and R^{1b} are each H;

R^2 , R^3 , R^5 and are each independently H or alkyl;

each occurrence of R^x , R^y , R^u and R^v is present or absent (depending upon chain saturation), and are each independently H or alkyl;

R^6 is independently selected from the group consisting of: H, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide,


wherein R^6 is optionally substituted with one, two, three or four substituents independently selected from: halo, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide;

$n=0$ to 10; and

$m=0$ to 20;

or an agriculturally acceptable salt thereof;

or a compound of **Formula (IV)(a)**:



(IV)(a)

wherein:

R^{1a} and R^{1b} are each H;

R^2 , R^3 , R^5 and are each independently H or alkyl;

each occurrence of R^x , R^y , R^u , R^v , R^z and R^w is present or absent (depending upon chain saturation), and is each independently H or alkyl;

$n=0$ to 10;

$m=0$ to 20; and

$p=0$ to 20

or an agriculturally acceptable salt thereof.

2. The method of claim 1, wherein said plant is a fruit crop plant or a vegetable crop plant.

3. The method of claim 1, wherein said plant is a citrus tree, and said compound is applied in an amount effective to treat or control a bacterial disease selected from the group consisting of canker, bacterial spot, Black Pit (fruit), Blast, citrus variegated chlorosis, and Citrus Huanglongbing.

4. The method of claim 3, wherein said citrus tree is selected from the group consisting of orange, grapefruit, Mandarin, lemon, lime and Kumquat.

5. The method of claim 1, wherein said plant is a pome fruit, and said compound is applied in an amount effective to treat or control a bacterial disease selected from the group consisting of Fire Blight, Crown Gall, Blister spot and Hairy root.

6. The method of claim 5, wherein said pome fruit is selected from the group consisting of apple, pear, quince, Asian pear, and loquats.

7. The method of claim 1, wherein the plant is a *Musa* species such as a banana, and the compound is applied in an amount effective to treat or control *Ralstonia solanacearum*.

8. The method of claim 1, wherein the plant is a cole (Brassicaceae) such as cabbage or broccoli, and the compound is applied in an amount effective to treat or control black rot (*Xanthomonas campestris*).

9. The method of claim 1, wherein the plant is a winegrape, and the compound is applied in an amount effective to treat or control for Pierce's disease (*Xylella fastidiosa*) or crown gall (*Agrobacterium vitas*, *A. tumefaciens*).

10. The method of claim 1, wherein the plant is a stone fruit or nut (e.g., peaches, nectarines, plums, almonds, walnuts), and the compound is applied in an amount effective to treat or control bacterial spot and/or blight caused by *Xanthomonas arboricola*; blight caused by *Pseudomonas syringae*; crown gall caused by *Agrobacterium tumefaciens*; phony peach and plum; or almond leaf scorch caused by *Xylella fastidiosa*.

11. The method of claim 1, wherein the plant is a landscape and/or shade tree (e.g., oak, maple, birch, etc.) for bacterial leaf scorch disease (e.g., caused by *Xylella fastidiosa*).

12. The method of claim 1, wherein the plant is a potato, and the compound is applied in an amount effective to treat or control soft rot or black leg (*Erwinia*, *Pectobacterium*).

13. The method of claim 1, wherein said plant is a pepper plant, and said compound is applied in an amount effective to treat or control a bacterial disease selected from the group consisting of Bacterial Spot, Bacterial wilt, Bacterial canker, and *Syringae* seedling blight and leaf spot.

14. The method of claim 1, wherein said plant is a tomato plant, and said compound is applied in an amount effective to treat or control a bacterial disease selected from the group consisting of: bacterial canker, bacterial speck, bacterial spot, bacterial stem rot and fruit rot, Bacterial wilt, Pith necrosis, and *Syringae* leaf spot.

15. The method of claim 1, wherein said plant is a soybean plant, and said compound is applied in an amount effective to treat or control a bacterial disease selected from the group consisting of Bacterial blight, Bacterial pustules, Bacterial wilt, Bacterial crinkle leaf, Bacterial tan spot, and Wildfire.

16. The method of claim 1, wherein said plant is corn, and said compound is applied in an amount effective to treat or control a bacterial disease selected from the group consisting of: Bacterial leaf blight, stalk rot, bacterial stripe, chocolate spot, holcus spot all caused by

Pseudomonas species, Bacterial leaf spot caused by *Xanthomonas* species, Bacterial stalk rot, top rot and Stewart's disease caused by *Erwinia* species, seed rot-seedling blight caused by *Bacillus* species, Purple leaf sheath caused by *Hemiparasitic* species, Corn stunt caused by *Spiroplasma kunkelii*, Goss's bacterial wilt and blight caused by *Clavibacter michiganensis*.

17. The method of claim 1, wherein said plant is cotton, and said compound is applied in an amount effective to treat or control a bacterial disease selected from the group consisting of Bacterial blight caused by *Xanthomonas* species, and Crown gall caused by *Agrobacterium* species and Lint degradation caused by *Erwinia* species.

18. The method of claim 1, wherein said plant is wheat, and said compound is applied in an amount effective to treat or control a bacterial disease selected from the group consisting of Bacterial leaf blight, bacterial sheath rot and Basal glume rot caused by *Pseudomonas* species, Bacterial mosaic and Spike blight caused by *Clavibacter* species, Black chaff caused by *Xanthomonas* species, and Pink seed caused by *Erwinia* species.

19. The method of claim 1, wherein said plant is rice, and said compound is applied in an amount effective to treat or control a bacterial disease selected from the group consisting of bacterial blight and leaf streak caused by *Xanthomonas* species, Foot rot caused by *Erwinia* species, Grain rot caused by *Burkholderia* species, and Sheath brown rot caused by *Pseudomonas* species.

20. The method of claim 1, wherein said plant is pineapple, and said compound is applied in an amount effective to treat or control a bacterial disease selected from the group consisting of bacterial heart rot, fruit collapse, bacterial fruitlet brown rot, marbled fruit, pink fruit and soft rot caused by *Erwinia* species, and Acetic souring caused by Acetic acid bacteria.

21. The method of claim 1, wherein said microbial biofilm formation or microbial infection is caused by a fungi.

22. The method of claim 1, wherein said compound is applied to said plant in an amount effective to treat or control a fungal disease selected from the group consisting of rots, leaf molds, blights, wilts, damping-off, spot, root rot, stem rot, mildew, brown spot, gummosis, melanose, post-bloom fruit drop, scab, alternaria, canker, flyspeck, fruit blotch, dieback, downy mildews, ear rots, anthracnose bunts, smut, rust, eyespot and pecky rice.

23. The method of claim 1, wherein said plant is Citrus, and said compound is applied in an amount effective to treat or control a fungal disease selected from the group consisting of:

Alternaria brown spot caused by *Alternaria alternaria*,
Brown rot caused by *Phytophthora citricola*,
Greasy spot and Greasy spot rind blotch caused by *Mycosphaerella citri*,
Melanose caused by *Diaporthe citri*,
Phytophthora foot rot,
gummosis and root rot caused by *Phytophthora citrophthora*, *Phytophthora palmivora*,
and *Phytophthora syringae* and other *Phytophthora* spp.,
Post bloom fruit drop caused by *Colletotrichum acutatum*, and
Scab caused by *Elsinoe fawcettii*.

24. The method of claim 1, wherein said plant is Pome fruit, and said compound is applied in an amount effective to treat or control a fungal disease selected from the group consisting of:

Apple scab caused by *Venturia inaequalis*,
Bitter rot caused by *Colletotrichum gloeosporioides*,
Diplodia canker caused by *Dilpodia mutila*,
Phytophthora crown, collar, root and fruit rot caused by *Phytophthora* spp.,
Powdery mildew caused by *Podosphaera leucotricha*,
Pacific Coast pear rust,
Cedar apple rust,
Quince rust caused by *Gymnosporangium* spp., and
Flyspeck caused by *Schizothyrium pomi*.

25. The method of claim 1, wherein said plant is Peppers, and said compound is applied in an amount effective to treat or control a fungal disease selected from the group consisting of:

Anthrachnose caused by *Colletotrichum* spp.,
Damping-off and root rot caused by *Rhizoctonia solani*, *Phytophthora* spp., *Fusarium* spp., and *Pythium* spp.,
Phytophthora blight caused by *Phytophthora capsici*, and
Verticillium wilt caused by *Verticillium albo-atrum*.

26. The method of claim 1, wherein said plant is Tomato, and said compound is applied in an amount effective to treat or control a fungal disease selected from the group consisting of:

Alternaria stem canker caused by *Alternaria alternaria*,
Anthracnose caused by *Colletotrichum* spp.,
Fusarium crown, root rot and wilt caused by *Fusarium oxysporum*,
Gray mold caused by *Botrytis cinerea*,
Late blight caused by *Phytophthora infestans*,
Pythium damping-off and fruit rot caused by *Pythium* spp.,
Rhizoctonia damping-off and fruit rot caused by *Rhizoctonia solani*,
Septoria leaf spot caused by *Septoria lycopersici*,
Verticillium wilt caused by *Verticillium albo-atrum*, and
White mold caused by *Sclerotinia sclerotiorum*.

27. The method of claim 1, wherein said plant is Soybean, and said compound is applied in an amount effective to treat or control a fungal disease selected from the group consisting of:

Phytophthora root and stem rot caused by *Phytophthora sojae*,
Pythium root rot, damping-off and seed decay caused by *Pythium* spp.,
Brown stem rot caused by *Phialophora gregata*,
Rhizoctonia root and stem rot caused by *Rhizoctonia solani*,
Stem canker, pod and stem blight caused by *Diaporthe phaseolorum*,
Phomopsis seed decay caused by *Phomopsis longicolla*,
Charcoal rot caused by *Macrophomina phaseolina*,
Sclerotinia stem rot caused by *Sclerotinia sclerotiorum*,
Sudden death syndrome caused by *Fusarium solani*, and
Soybean Rust caused by *Phakopsora pachyrhizi*.

28. The method of claim 1, wherein said plant is Grape, and said compound is applied in an amount effective to treat or control a fungal disease selected from the group consisting of:

Alternaria rot caused by *Alternaria alternaria*,
Angular leaf spot caused by *Mycosphaerella angulata*,
Botrytis bunch rot and blight caused by *Botrytis cinerea*,
Diplodia cane dieback and bunch rot caused by *Diplodia natalensis*,
Downy mildew caused by *Plasmopara viticola*,
Phytophthora crown and root rot caused by *Phytophthora* spp.,
Powdery mildew caused by *Uncinula necator*,

Ripe rot caused by *Glomerella cingulata*,
Septoria leaf spot caused by *Septoria ampelopsidis*, and
Verticillium wilt caused by *Verticillium dahliae*.

29. The method of claim 1, wherein said plant is Potato, and said compound is applied in an amount effective to treat or control a fungal disease selected from the group consisting of:

Brown spot, Black pit and Early blight caused by *Alternaria* spp.,
Fusarium dry rot and wilt caused by *Fusarium* spp.,
Gangrene caused by *Phoma* spp.,
Late blight and Pink rot caused by *Phytophthora* spp.,
Rhizoctonia canker and black scurf caused by *Rhizoctonia solani*,
Rosellinia black rot caused by *Rosellinia* spp.,
Septoria leaf spot caused by *Septoria lycopersici*,
Stem rot caused by *Sclerotium rolfsii*,
Verticillium wilt caused by *Verticillium albo-atrum*, and
White mold caused by *Sclerotinia sclerotiorum*.

30. The method of claim 1, wherein said plant is Pineapple, and said compound is applied in an amount effective to treat or control a fungal disease selected from the group consisting of:

Anthrachnose caused by *Colletotrichum ananas*,
Butt rot and White leaf spot caused by *Chalara paradoxa*,
Leaf spot caused by *Curvularia eragrostidis*,
Phytophthora heart rot caused by *Phytophthora cinnamomi* and *Phytophthora parasitica*,
Root rot and Seedling blight caused by *Pythium* spp., and
Leaking brown ring caused by *Toffliadis dimenationa*.

31. The method of claim 1, wherein said plant is Cotton, and said compound is applied in an amount effective to treat or control a fungal disease selected from the group consisting of:

Anthrachnose caused by *Glomerella gossypii*,
Boll rot caused by *Colletotrichum gossypii*, *Fusarium* spp., *Phytophthora* spp., or *Rhizoctonia solani*,
Fusarium wilt caused by *Fusarium oxysporum*,
Leaf spot caused by *Alternaria* spp., *Cercospora gossypina*, *Rhizoctonia solani*, and *Stemphylium solani*,
Lint contamination caused by *Aspergillus flavus*,

Powdery mildew caused by *Leveillula taurica*,
 Cotton rust caused by *Puccinia schedonnardii*,
 Southwestern cotton rust caused by *Puccinia cacabata*,
 Tropical cotton rust caused by *Phakopsora gossypii*,
 Southern blight caused by *Sclerotium rolfsii*,
 Seedling disease complex caused by *Colletotrichum gossypii*, *Fusarium* spp., *Pythium* spp., *Rhizoctonia solani*, or *Thielaviopsis basicola*,
 Stem canker caused by *Phoma exigua*, and
 Verticillium wilt caused by *Verticillium dahliae*.

32. The method of claim 1, wherein said plant is Corn, and said compound is applied in an amount effective to treat or control a fungal disease selected from the group consisting of:

Anthracnose caused by *Colletotrichum graminicola*,
 Aspergillus ear and kernel rot caused by *Aspergillus flavus*,
 Banded leaf, sheath spot, root rot and stalk rot caused by *Rhizoctonia solani*,
 Brown spot, Black spot and Stalk rot caused by *Physoderma maydis*,
 Curvularia leaf spot caused by *Curvularia clavata*,
 Diplodia ear rot, stalk rot, seed rot and seedling blight caused by *Dilpodia* spp.,
 Downey mildews caused by *Sclerophthora* spp. or *Peronosclerospora* spp.,
 Ear rots caused by *Alternaria alternaria*,
 Ergot caused by *Claviceps gigantea*,
 Fusarium ear, stalk, kernel, root, seed rot, seedling blight caused by *Fusarium* spp.,
 Cercospora leaf spot caused by *Cercospora zea-maydis*,
 Helminthosporium ear rot caused by *Helminthosporium carbonum*,
 Pythium root rot and stalk rot caused by *Pythium* spp.,
 Rhizoctonia ear rot caused by *Rhizoctonia zea*,
 Common corn rust and Southern corn rust caused by *Puccinia* spp.,
 Southern blight caused by *Athelia rolfsii*,
 Common smut caused by *Ustilago zea*,
 Southern corn leaf blight and stalk rot caused by *Cochliobolus heterostrophus*, and
 storage rots caused by *Aspergillus* spp. and *Penicillium* spp.

33. The method of claim 1, wherein said plant is Rice, and said compound is applied in an amount effective to treat or control a fungal disease selected from the group consisting of:

Black kernel caused by *Curvularia lunata*,

Blast caused by *Pyricularia oryzae*,
 Brown spot caused by *Cochliobolus miyabeanus*,
 Downy mildew caused by *Sclerophthora macrospora*,
 False smut caused by *Ustilaginoidea virens*,
 Narrow brown leaf spot caused by *Cercospora janseana*,
 Pecky rice caused by *Fusarium* spp., *Microdochium oryzae*, or *Sarocladium oryzae*,
 Root rot caused by *Fusarium* spp., or *Pythium* spp.,
 Seedling blight caused by fungi (e.g., *Cochliobolus miyabeanus*, *Curvularia* spp.,
Fusarium spp., *Rhizoctonia solani*, *Sclerotium rolfsii* and *Athelia rolfsii*),
 Stackburn caused by *Alternaria padwickii*,
 Stem rot caused by *Magnaporthe salvinii*,
 Water-mold (seed-rot and seedling disease) caused by *Achlya* spp., *Fusarium* spp., or
Pythium spp.

34. The method of claim 1, wherein said plant is Wheat, and said compound is applied in an amount effective to treat or control a fungal disease selected from the group consisting of:

Alternaria leaf blight caused by *Alternaria triticina*,
 Anthracnose caused by *Colletotrichum graminicola*,
 Black head molds caused by *Cladosporium* spp., *Epicoccum* spp., *Sporobolomyces* spp.
 or *Stemphylium* spp.,
 Common bunt caused by *Tilletia* spp.,
 Crown rot, seedling blight and dryland root rot caused by *Fusarium* spp. or *Gibberella*
 spp.,
 Downey mildew caused by *Sclerophthora macrospora*,
 Dwarf bunt caused by *Tilletia controversa*,
 Ergot caused by *Claviceps purpurea*,
 Eyespot caused by *Tapesia yallundae*,
 Leaf rust caused by *Puccinia triticina*,
 Loose smut caused by *Ustilago tritici*,
 Microscopia leaf spot caused by *Phaeosphaeria microscopia*,
 Phoma spot caused by *Phoma* spp.,
 Powdery mildew caused by *Erysiphe graminis*,
 Pythium root rot,
 Snow rot caused by *Pythium* spp.,
 Rhizoctonia root rot caused by *Rhizoctonia solani*,


Scab (head blight) caused by *Fusarium* spp. or *Gibberella* spp.,
 Southern blight caused by *Sclerotium rolfii*,
 Speckled snow mold caused by *Typhula* spp.,
 Stem rust caused by *Puccinia graminis*,
 storage molds caused by *Aspergillus* spp. or *Penicillium* spp.,
 Take-all caused by *Gaeumannomyces graminis*, and
 Zoosporic root rot caused by *Lagenaria radicola*.

35. An agricultural composition comprising:

(a) an agriculturally acceptable carrier; and

(b) an antimicrobial or biofilm preventing, removing or inhibiting a compound of

Formula (II)(a):



(II)(a)

wherein:

R^{1a} and R^{1b} are each H;

R^2 , R^3 , R^5 and are each independently H or alkyl;

each occurrence of R^x , R^y , R^u and R^v is present or absent (depending upon chain saturation),
 and are each independently H or alkyl;

R^6 is independently selected from the group consisting of: H, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide,


wherein R^6 is optionally substituted with one, two, three or four substituents independently selected from: halo, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide;

$n=0$ to 10; and

$m=0$ to 20;

or an agriculturally acceptable salt thereof;

or a compound of **Formula (IV)(a)**:



(IV)(a)

wherein:

R^{1a} and R^{1b} are each H;

R^2 , R^3 , R^5 and are each independently H or alkyl;

each occurrence of R^x , R^y , R^u , R^v , R^z and R^w is present or absent (depending upon chain saturation), and is each independently H or alkyl;

$n=0$ to 10 ;

$m=0$ to 20 ; and

$p=0$ to 20

or an agriculturally acceptable salt thereof.

36. The composition of claim 35, further comprising a microbicide.

37. The composition of claim 36, wherein said microbicide comprises copper.

38. The composition of claim 36, wherein said microbicide comprises copper hydroxide.

39. The composition of claim 35, further comprising an antibiotic.


40. The composition of claim 35, further comprising a bacteriophage.

41. The composition of claim 35, further comprising a plant defense activator.

42. The composition of any of claims 35-41, wherein said carrier is an aqueous carrier or a solid particulate carrier.

43. An antimicrobial or biofilm preventing, removing or inhibiting compound of

Formula (II)(a):



(II)(a)

wherein:

R^{1a} and R^{1b} are each H;

R^2 , R^3 , R^5 and are each independently H or alkyl;

each occurrence of R^x , R^y , R^u and R^v is present or absent (depending upon chain saturation), and are each independently H or alkyl;

R^6 is independently selected from the group consisting of: H, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide,


wherein R^6 is optionally substituted with one, two, three or four substituents independently selected from: halo, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide;

$n=0$ to 10; and

$m=0$ to 20;

or an agriculturally acceptable salt thereof;

or compound of **Formula (IV)(a)**:




(IV)(a)

wherein:

R^{1a} and R^{1b} are each H;


R^2 , R^3 , R^5 and are each independently H or alkyl;
 each occurrence of R^x , R^y , R^u , R^v , R^z and R^w is present or absent (depending upon chain saturation), and is each independently H or alkyl;
 $n=0$ to 10;
 $m=0$ to 20; and
 $p=0$ to 20
 or an agriculturally acceptable salt thereof;
 for use in treating or preventing a bacterial or fungal infection in a plant or plant part.

44. The method of any of claims 1-34, wherein said compound is a compound of Formula (II)(a)(5)(D):




or an agriculturally acceptable salt thereof.

45. The composition of any of claims 35-42, wherein said compound is a compound of Formula (II)(a)(5)(D):



or an agriculturally acceptable salt thereof.

46. A method of enhancing the effects of a microbicide comprising applying a compound of **Formula (II)(a)**:



(II)(a)

wherein:

R^{1a} and R^{1b} are each H;

R^2 , R^3 , R^5 and are each independently H or alkyl;

each occurrence of R^x , R^y , R^u and R^v is present or absent (depending upon chain saturation), and are each independently H or alkyl;

R^6 is independently selected from the group consisting of: H, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide,


wherein R^6 is optionally substituted with one, two, three or four substituents independently selected from: halo, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide;

$n=0$ to 10; and

$m=0$ to 20;

or an agriculturally acceptable salt thereof;

or a compound of **Formula (IV)(a)**:



(IV)(a)

wherein:

R^{1a} and R^{1b} are each H;

R^2 , R^3 , R^5 and are each independently H or alkyl;

each occurrence of R^x , R^y , R^u , R^v , R^z and R^w is present or absent (depending upon chain saturation), and is each independently H or alkyl;

$n=0$ to 10;

$m=0$ to 20; and

$p=0$ to 20

or an agriculturally acceptable salt thereof;

in combination with said microbicide.

47. The method of claim 46, wherein said microbicide comprises copper.


48. The method of claim 46, wherein said microbicide comprises copper hydroxide.

49. The method of claim 46, wherein said microbicide is an antibiotic or a bacteriophage.

50. The method of any of claims 46-49, wherein said applying step is carried out by applying said active compound and said microbicide simultaneously.


51. The method of any of claims 46-49, wherein said applying step is carried out by applying said active compound and said microbicide sequentially.

52. The method of any of claims 46-51, wherein said active compound is a compound of Formula (II)(a)(5)(D):



or an agriculturally acceptable salt thereof.

53. A method of enhancing the effects of a plant defense activator comprising applying a compound of **Formula (II)(a)**:



(II)(a)

wherein:

R^{1a} and R^{1b} are each H;

R^2 , R^3 , R^5 and are each independently H or alkyl;

each occurrence of R^x , R^y , R^u and R^v is present or absent (depending upon chain saturation), and are each independently H or alkyl;

R^6 is independently selected from the group consisting of: H, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide,


wherein R^6 is optionally substituted with one, two, three or four substituents independently selected from: halo, hydroxy, acyl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, heteroaryl, alkoxy, amino, amide, thiol, sulfone, sulfoxide, oxo, oxy, nitro, carbonyl, carboxy, amino acid sidechain, amino acid and peptide;

$n=0$ to 10; and

$m=0$ to 20;

or an agriculturally acceptable salt thereof;

or a compound of **Formula (IV)(a)**:



(IV)(a)

wherein:

R^{1a} and R^{1b} are each H;

R^2 , R^3 , R^5 and are each independently H or alkyl;

each occurrence of R^x , R^y , R^u , R^v , R^z and R^w is present or absent (depending upon chain saturation), and is each independently H or alkyl;

$n=0$ to 10;

$m=0$ to 20; and

$p=0$ to 20


or an agriculturally acceptable salt thereof;

in combination with said plant defense activator.


54. The method of claim 53, wherein said applying step is carried out by applying said active compound and said plant defense activator simultaneously.

55. The method of claim 53, wherein said applying step is carried out by applying said active compound and said plant defense activator sequentially.

56. The method of any of claims 53-55, wherein said active compound is a compound of Formula (II)(a)(5)(D):



or an agriculturally acceptable salt thereof.







Figure 2

Effects of Bacterial Spot Disease on Peppers



No Disease



Diseased

Figure 3

Foliar Disease on Inoculated Pepper Plants




Figure 4

Pepper Fruit Yield Per Inoculated Plant

(July 1 and 9 Harvests)




Figure 5

Foliar Disease on Pepper Plants – Plot Average




Figure 6




Figure 7

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 09/66979

A. CLASSIFICATION OF SUBJECT MATTER

IPC(8) - A01N 25/08; C07D 249/04; A61K 31/4192 (2010.01)

USPC - 514/397, 424/409, 548/255, 544/139

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC(8) - A01N 25/08; C07D 249/04; A61K 31/4192 (2010.01)

USPC - 514/397, 424/409, 548/255, 544/139

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
USPC - 514/359, 385, 396, 397, 401, 402; 548/255, 300.1, 311.1, 314.7, 326.5, 331.5, 335.1, 335.5 (search term limited)

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

PubWEST DB=PGPB,USPT,EPAB; WIPO, Google Scholar

Search terms used: agricultur\$, bacteri\$, crop, diazole, fruit, fung\$, imidazole, plant, triazole, vegetable

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 2004/0235934 A1 (Fischer et al.) 25 November 2004 (25.11.2004), para [0074] compounds IIa, IIc; [0157], [0162], [0167]-[0169], [0172]-[0174], [0185], [0210], [0211], [0340]	1-44, 46-51, 53-56
Y	US 2007/0142371 A1 (Cogan et al.) 21 June 2007 (21.06.2007), pg 29, table I; para [0018]	1-44, 46-51, 53-56
Y	US 2007/0185092 A1 (Zhu et al.) 09 August 2007 (09.08.2007), pg 61, Table 2, compound 8	44, 56

☐ Further documents are listed in the continuation of Box C.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

24 January 2010 (24.01.2010)

Date of mailing of the international search report

23 FEB 2010

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US, Commissioner for Patents
P.O. Box 1450, Alexandria, Virginia 22313-1450

Facsimile No. 571-273-3201

Authorized officer:

Lee W. Young

PCT Helpdesk: 571-272-4300

PCT OSP: 571-272-7774

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 09/66979

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☒ Claims Nos.: 45, 52
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying additional fees, this Authority did not invite payment of additional fees.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- ☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- ☐ No protest accompanied the payment of additional search fees.