




(12) **DEMANDE DE BREVET CANADIEN**
CANADIAN PATENT APPLICATION
(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2017/05/31	(51) Cl.Int./Int.Cl. <i>C12N 15/52</i> (2006.01), <i>C11D 3/386</i> (2006.01), <i>C12N 9/54</i> (2006.01)
(87) Date publication PCT/PCT Publication Date: 2017/12/07	
(85) Entrée phase nationale/National Entry: 2018/11/29	(71) Demandeur/Applicant: DANISCO US INC., US
(86) N° demande PCT/PCT Application No.: US 2017/035217	(72) Inventeurs/Inventors: MARQUEZ, DAVID, US; STONER, MICHAEL, US; KELLIS, JAMES T., JR., US; GARSKE, ADAM, US; BASLER, JOSHUA ROY, US; AUGUSTYN, KATHERINE, US; ...
(87) N° publication PCT/PCT Publication No.: 2017/210295	
(30) Priorité/Priority: 2016/05/31 (US62/343,618)	(74) Agent: BERESKIN & PARR LLP/S.E.N.C.R.L.,S.R.L.

(54) Titre : VARIANTS DE PROTEASE ET LEURS UTILISATIONS
(54) Title: PROTEASE VARIANTS AND USES THEREOF



(57) **Abrégé/Abstract:**
Disclosed herein is one or more subtilisin variant, nucleic acid encoding same, and compositions and methods related to the production and use thereof, including one or more subtilisin variant that has improved stability and/or soil removal compared to one or more reference subtilisin.

(72) **Inventeurs(suite)/Inventors(continued):** CHANDRASEKARAN, PRIYANKA, US;
SCOTCHER, MILES CHRISTOPHER, US; BABE, LILIA MARIA, US; BOTT, RICHARD R., US; ESTELL, DAVID A., US;
VOGTENTANZ, GUDRUN, US; PRICELIUS, SINA, US; YAO, JIAN, US

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau(10) International Publication Number
WO 2017/210295 A1(43) International Publication Date
07 December 2017 (07.12.2017)

(51) International Patent Classification:

C12N 15/52 (2006.01) *C11D 3/386* (2006.01)
C12N 9/54 (2006.01)

(21) International Application Number:

PCT/US2017/035217

(22) International Filing Date:

31 May 2017 (31.05.2017)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/343,618 31 May 2016 (31.05.2016) US

(71) Applicant: **DANISCO US INC.** [US/US]; 925 Page Mill Road, Palo Alto, California 94304 (US).

(72) Inventors; and

(71) Applicants: **MARQUEZ, David** [US/US]; 925 Page Mill Road, Palo Alto, California 94304 (US). **STONER, Michael** [US/US]; 925 Page Mill Drive, Palo Alto, California 94304 (US). **KELLIS, Jr., James T.** [US/US]; 925 Page Mill Road, Palo Alto, California 94304 (US). **GARSKE, Adam** [US/US]; 925 Page Mill Road, Palo Alto, California 94304 (US).

(72) Inventors: **BASLER, Joshua Roy**; 925 Page Mill Road, Palo Alto, California 94304 (US). **AUGUSTYN, Katherine**; 925 Page Mill Road, Palo Alto, California 94304 (US). **CHANDRASEKARAN, Priyanka**; 925 Page Mill Road, Palo Alto, California 94304 (US). **SCOTCHER, Miles Christopher**; 925 Page Mill Road, Palo Alto, California 94304 (US). **BABE, Lilia Maria**; 925 Page Mill Road, Palo Alto, California 94304 (US). **BOTT, Richard R.**; 925 Page Mill Road, Palo Alto, California 94304 (US). **ESTELL, David A.**; 925 Page Mill Road, Palo Alto, California 94304 (US). **VOGTENTANZ, Gudrun**; 925 Page Mill Road, Palo Alto, California 94304 (US). **PRICELIUS, Sina**; 925 Page Mill Road, Palo Alto, California 94304 (US). **YAO, Jian**; 925 Page Mill Road, Palo Alto, California 94304 (US).

(74) Agent: **COHEN, Jacqueline** et al.; E. I. du Pont de Nemours and Company, Legal Patent Records Center, Chestnut Run Plaza 721/2340, 974 Centre Road, PO Box 2915 Wilmington, Delaware 19805 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KH, KN, KP, KR,

(54) Title: PROTEASE VARIANTS AND USES THEREOF

FIG. 1A

FIG. 1AA

FIG. 1AB

FIG. 1AA

FIG. 1AB

FIG. 1AA

FIG. 1AB

(57) Abstract: Disclosed herein is one or more subtilisin variant, nucleic acid encoding same, and compositions and methods related to the production and use thereof, including one or more subtilisin variant that has improved stability and/or soil removal compared to one or more reference subtilisin.

[Continued on next page]

WO 2017/210295 A1



KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- with sequence listing part of description (Rule 5.2(a))

PROTEASE VARIANTS AND USES THEREOF

CROSS-REFERENCE TO RELATED APPLICATIONS

[001] This application is related to and claims the benefit of priority from United States Provisional Patent Application Serial No. 62/343,618, filed May 31, 2016, which is hereby incorporated herein by reference in its entirety.

REFERENCE TO SEQUENCE LISTING SUBMITTED ELECTRONICALLY

[002] The content of the sequence listing electronically submitted with the application as an ASCII text file (Name: NB40785WOPCT_ST25; Size: 882KB; Created: May 26, 2017) forms part of the application and is hereby incorporated herein by reference in its entirety.

[003] Disclosed herein is one or more subtilisin variant, nucleic acid encoding same, and compositions and methods related to the production and use thereof, including one or more subtilisin variant that has improved stability and/or soil removal compared to one or more reference subtilisin.

[004] Serine proteases are enzymes (EC No. 3.4.21) possessing an active site serine that initiates hydrolysis of peptide bonds of proteins. Serine proteases comprise a diverse class of enzymes having a wide range of specificities and biological functions that are further divided based on their structure into chymotrypsin-like (trypsin-like) and subtilisin-like. The prototypical subtilisin (EC No. 3.4.21.62) was initially obtained from *Bacillus subtilis*. Subtilisins and their homologues are members of the S8 peptidase family of the MEROPS classification scheme. Members of family S8 have a catalytic triad in the order Asp, His and Ser in their amino acid sequence. Although a number of useful variant proteases have been developed for cleaning applications, there remains a need for improved protease variants.

[005] One embodiment provides one or more subtilisin variant comprising an amino acid sequence comprising two, three, or four or more variations at positions corresponding to SEQ ID NO:2, wherein said positions are selected from: (i) 1, 3, 9, 10, 15, 22, 24, 26, 27, 28, 29, 30, 31, 35, 37, 40, 43, 45, 48, 52, 68, 71, 76, 77, 78, 86, 87, 95, 96, 98, 99, 100, 101, 102, 103, 105, 108, 111, 114, 115, 117, 119, 123, 125, 126, 127, 128, 129, 130, 136, 143, 146, 147, 151, 155, 158, 160, 161, 165, 184, 187, 193, 202, 203, 204, 210, 211, 216, 217, 234, 238, 239, 242, 243, 250, 255, 258, 259, 260, 264, and 274; (ii) 1, 3, 9, 10, 15, 22, 24, 26, 28, 29, 30, 31, 35, 37, 40, 45, 48, 52, 68, 71, 76, 77, 78, 86, 87, 95, 96, 98, 99, 100, 101, 102, 105, 108, 111, 114, 115, 117, 119, 123, 125, 126, 127, 128, 129, 130, 136, 143, 146, 147, 151, 155, 158, 160, 161, 165, 184, 187, 193, 202, 203, 204, 210, 211, 216, 217, 234, 238, 239, 242, 243, 250, 255, 258, 259, 260, 264, and 274; (iii) 1, 3, 9, 10, 24, 26, 29, 30, 31, 35, 37, 40, 45, 48, 52, 68, 71, 77, 78, 86, 87, 95, 96, 98, 99, 100, 101, 102, 105, 108, 111, 115, 117, 119, 123, 127, 128, 129, 130, 136, 143, 146, 147, 151, 155, 158, 160, 161, 165, 184, 187, 193, 202, 203, 204, 210, 211, 216, 217, 234, 238, 239, 242, 243, 250, 258, 259, 260, 264, and 274; (iv) 3, 9, 10, 35, 68, 77, 78, 86, 87, 99, 101, 115, 123, 127, 128, 165, 184, 187, 202, 203, 210, 217, 242, 250, 258, and 264; or (v) 3, 68, 77, 78, 123, 127, 128, 165, 184,

202, 210, 217, and 258; with the proviso that one or more of said two, three, or four or more variations is a man-made mutation or substitution; wherein said variant has at least 85% but less than 100% amino acid sequence identity to an AprL-Clade subtilisin enzyme comprising a Motif A and/or a Motif B; wherein Motif A is selected from Motif 1, Motif 2, Motif 3, and Motif 4; wherein Motif B is selected from Motif 5, Motif 6, Motif 7, Motif 8, Motif 9, Motif 10, Motif 11, Motif 12, Motif 13, and Motif 14; and wherein the amino acid positions of the variant are numbered by correspondence with the amino acid sequence of SEQ ID NO:2.

[006] Yet another embodiment provides one or more subtilisin variant comprising an amino acid sequence comprising two, three, or four or more amino acid substitutions at two, three, or four or more positions corresponding to SEQ ID NO:2, wherein said substitutions are selected from: (i) 1Q, 3Q/V, 9E, 10M, 15I, 22Y, 24N/Q, 26Q, 27R, 28A, 29S, 30T, 31I, 35A, 37T, 40E, 43A, 45I/Q/R, 48I/T/V, 52R, 68S, 71A, 76K, 77D/E/H/N/Q/S, 78I, 86H/N/R, 87S/T, 95A/F/Q, 96S, 98H/R, 99A/E/Q/S, 100N, 101A, 102D/T, 103I, 105N, 108H/K, 111Q, 114N, 115F, 117S, 119Q, 123I, 125N, 126Q, 127E/F/Q/S/T/V, 128P/R/S, 129H/R, 130S, 136N/R, 143Q/V, 146A/S, 147L, 151G, 155E/G/N, 158D/N/Q, 160C/D/K, 161D/Q, 165A/E/Q/S, 184E/Q, 187E/P, 193D/Q, 202V, 203E/N/Q, 204I, 210E/H/I/P, 211S, 216Q, 217S, 234I/W, 238Q/R, 239K/S/T, 242G/N, 243I, 250G, 255R, 258D/E/P, 259E/P, 260W, 264H/Q/S, and 274A; (ii) 1Q, 3Q/V, 9E, 10M, 15I, 22Y, 24N/Q, 26Q, 28A, 29S, 30T, 31I, 35A, 37T, 40E, 45I/Q/R, 48I/T/V, 52R, 68S, 71A, 76K, 77D/E/H/N/S, 78I, 86H/N/R, 87S/T, 95A/F/Q, 96S, 98K/R, 99A/E/Q/S, 100N, 101A, 102D/T, 105N, 108H/K, 111Q, 114N, 115F, 117S, 119Q, 123I, 125N, 126Q, 127E/F/Q/S/T/V, 128P/R/S, 129H/R, 130S, 136N, 143Q/V, 146A/S, 147L, 151G, 155E/G/N, 158D/N/Q, 160D/K, 161D/Q, 165A/E/Q/S, 184E/Q, 187E/P, 193D/Q, 194E, 202V, 203E/N/Q, 204I, 210I/P, 211S, 216Q, 217S, 234I/W, 238Q/R, 239K/S/T, 242G/N, 243I, 250G, 255R, 258D/P, 259E/P, 260W, 264H/Q/S, and 274A; (iii) 1Q, 3V, 9E, 10M, 24Q, 26Q, 29S, 30T, 31I, 35A, 37T, 40E, 45I/Q/R, 48T/V, 52R, 68S, 71A, 77E/N, 78I, 86H/R, 87S, 95A/F, 96S, 98R, 99A/E/Q/S, 100N, 101A, 102D/T, 105N, 108H, 111Q, 115F, 117S, 119Q, 123I, 127E/Q/S/T/V, 128P/R, 129H/R, 130S, 136N, 143Q/V, 146A, 147L, 151G, 155G/N, 158D/N/Q, 160D/K, 161D/Q, 165A/E/Q, 184E/Q, 187E/P, 193Q, 202V, 203E/N/Q, 204I, 210I/P, 211S, 216Q, 217S, 234W, 238Q, 239S, 242G/N, 243I, 250G, 258D/P, 259E/P, 260W, 264H/Q/S, and 274A; (iv) 3V, 9E, 10M, 35A, 68S, 77N, 78I, 86H, 87S, 99Q/S, 101A, 115F, 123I, 127S/T, 128P/R, 165E/Q, 184Q, 187E, 202V, 203E, 210P, 217S, 242G, 250G, 258D/P, and 264H; or (v) 3V, 68S, 77N, 78I, 123I, 127S, 128P, 165Q, 184Q, 202V, 210P, 217S, and 258D/P; with the proviso that one or more of said two, three, or four or more substitutions is a man-made substitution; wherein said variant has at least 85% but less than 100% amino acid sequence identity to an AprL-Clade subtilisin enzyme comprising a Motif A and/or a Motif B; wherein Motif A is selected from Motif 1, Motif 2, Motif 3, and Motif 4; wherein Motif B is selected from Motif 5, Motif 6, Motif 7, Motif 8, Motif 9, Motif 10,

Motif 11, Motif 12, Motif 13, and Motif 14; and wherein the amino acid positions of the variant are numbered by correspondence with the amino acid sequence of SEQ ID NO:2.

[007] In one embodiment, the AprL-clade enzymes comprise Motif A, Motif B, or a combination thereof. In other embodiments, the AprL-clade enzymes comprise Motif A and Motif B. In some embodiments, one or more subtilisin variant described herein comprises Motif A, Motif B, or a combination thereof. In still other embodiments, one or more subtilisin variant described herein comprises Motif A and Motif B. In another embodiment, Motif A is selected from Motif 1, Motif 2, Motif 3, and Motif 4. In a still further embodiment, Motif B is selected from Motif 5, Motif 6, Motif 7, Motif 8, Motif 9, Motif 10, Motif 11, Motif 12, Motif 13, and Motif 14.

[008] Some embodiments are directed to a composition comprising one or more subtilisin variant described herein. Further embodiments are directed to a method of cleaning comprising contacting a surface or an item in need of cleaning with one or more subtilisin variant described herein or one or more composition described herein.

[009] Figure 1A-1D provides a CLUSTAL W sequence alignment of AprL, Blid02339, and other AprL-clade enzymes.

[0010] Figure 2 provides a phylogenetic tree of subtilisins including AprL, Blid02339, other members of the AprL-clade, and other subtilisins, including variants described herein.

[0011] Unless otherwise indicated herein, one or more subtilisin variant described herein can be made and used via conventional techniques commonly used in molecular biology, microbiology, protein purification, protein engineering, protein and DNA sequencing, recombinant DNA fields, and industrial enzyme use and development. Terms and abbreviations not defined should be accorded their ordinary meaning as used in the art. Unless defined otherwise herein, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art. Any definitions provided herein are to be interpreted in the context of the specification as a whole. As used herein, the singular “a,” “an” and “the” includes the plural unless the context clearly indicates otherwise. Unless otherwise indicated, nucleic acid sequences are written left to right in 5' to 3' orientation; and amino acid sequences are written left to right in amino to carboxy orientation. Each numerical range used herein includes every narrower numerical range that falls within such broader numerical range, as if such narrower numerical ranges were all expressly written herein.

[0012] As used herein in connection with a numerical value, the term “about” refers to a range of +/- 0.5 of the numerical value, unless the term is otherwise specifically defined in context. For instance, the phrase a “pH value of about 6” refers to pH values of from 5.5 to 6.5, unless the pH value is specifically defined otherwise.

[0013] The nomenclature of the amino acid substitutions or variations of the one or more subtilisin

variants described herein uses one or more of the following: position; position:amino acid substitutions or variation(s); or starting amino acid(s):position:amino acid substitution(s). Reference to a “position” (i.e. 5, 8, 17, 22, etc) encompasses any starting amino acid that may be present at such position, and any variation from the starting amino acid or substitution that may be present at such position. Reference to a “position: amino acid substitution(s) and/or variation(s)” (i.e. 1S/T/G, 3G, 17T, etc) encompasses any starting amino acid that may be present at such position and the one or more amino acid(s) that may vary from the starting amino acid and/or with which such starting amino acid may be substituted. An amino acid substitution excludes the starting amino acid, where the substituted amino acid and starting amino acid are the same. Reference to a starting or substituted amino acid or an amino acid variation may be further expressed as several starting, substituted, or varied amino acids separated by a foreslash (“/”). For example, D275S/K indicates position 275 is substituted with serine (S) or lysine (K) and P/S197K indicates that starting amino acid proline (P) or serine (S) at position 197 is substituted with lysine (K).

[0014] The position of an amino acid residue in a given amino acid sequence is numbered by correspondence with the amino acid sequence of SEQ ID NO:2. That is, the amino acid sequence of the AprL-clade enzyme shown in SEQ ID NO:2 serves as a reference sequence. For example, the amino acid sequence of an AprL-clade subtilisin enzyme or one or more subtilisin variant described herein is aligned with the amino acid sequence of SEQ ID NO:2 using an alignment algorithm as described herein, and each amino acid residue in the given amino acid sequence that aligns (preferably optimally aligns) with an amino acid residue in SEQ ID NO:2 is conveniently numbered by reference to the numerical position of that corresponding amino acid residue. Sequence alignment algorithms, such as, for example, described herein will identify the location where insertions or deletions occur in a subject sequence when compared to a query sequence.

[0015] The term “variant,” with respect to a polypeptide amino acid sequence, refers to a polypeptide amino acid sequence that differs from a specified wild-type, parental, or reference polypeptide amino acid sequence by including one or more man-made substitutions, insertions, or deletions of an amino acid. Similarly, the term “variant,” with respect to a polynucleotide nucleic acid sequence, refers to a polynucleotide nucleic acid sequence that differs from a specified wild-type, parental, or reference polynucleotide by including one or more man-made substitutions, insertions, or deletions of a nucleic acid acid. The identity of the wild-type, parental, or reference polypeptide amino acid sequence or polynucleotide nucleic acid sequence will be apparent from context.

[0016] The term “variation(s)” when used, for example, in the phrase “two, three, or four or more amino acid variations versus SEQ ID NO:2” or “two, three, or four or more variations at positions corresponding to SEQ ID NO:2” encompasses each amino acid that is different from the amino acid present at the corresponding position in SEQ ID NO:2. For example, the sequence of the variant of

interest is aligned with SEQ ID NO:2 and each position in the variant compared to SEQ ID NO:2 to identify the amino acids at each position that are different from the amino acid present at the corresponding positions in SEQ ID NO:2 and each amino acid that is different from the corresponding amino acid in SEQ ID NO:2 is a variation.

[0017] The terms “protease” and “proteinase” refer to an enzyme that has the ability to break down proteins and peptides. A protease has the ability to conduct “proteolysis,” by hydrolysis of peptide bonds that link amino acids together in a peptide or polypeptide chain forming the protein. This activity of a protease as a protein-digesting enzyme is referred to as “proteolytic activity.” Many well-known procedures exist for measuring proteolytic activity. For example, proteolytic activity may be ascertained by comparative assays that analyze the respective protease’s ability to hydrolyze a suitable substrate. Exemplary substrates useful in the analysis of protease or proteolytic activity, include, but are not limited to, di-methyl casein (Sigma C-9801), bovine collagen (Sigma C-9879), bovine elastin (Sigma E-1625), and bovine keratin (ICN Biomedical 902111). Colorimetric assays utilizing these substrates are well known in the art (*See e.g.*, WO99/34011 and US 6,376,450). The pNA peptidyl assay (*See e.g.*, Del Mar et al., Anal Biochem, 99:316-320, 1979) also finds use in determining the active enzyme concentration. This assay measures the rate at which p-nitroaniline is released as the enzyme hydrolyzes a soluble synthetic substrate, such as succinyl-alanine-alanine-proline-phenylalanine-p-nitroanilide (suc-AAPF-pNA). The rate of production of yellow color from the hydrolysis reaction is measured at 410 nm on a spectrophotometer and is proportional to the active enzyme concentration. In addition, absorbance measurements at 280 nanometers (nm) can be used to determine the total protein concentration in a sample of purified protein. The activity on substrate/protein concentration gives the enzyme specific activity.

[0018] The phrase “composition(s) substantially-free of boron” or “detergent(s) substantially-free of boron” refers to composition(s) or detergent(s), respectively, that contain trace amounts of boron, for example, less than about 1000 ppm (1mg/kg or liter equals 1 ppm), less than about 100 ppm, less than about 50 ppm, less than about 10 ppm, or less than about 5 ppm, or less than about 1 ppm, perhaps from other compositions or detergent constituents.

[0019] As used herein, “the genus *Bacillus*” includes all species within the genus “*Bacillus*,” as known to those of skill in the art, including but not limited to *B. subtilis*, *B. licheniformis*, *B. lentus*, *B. brevis*, *B. stearothermophilus*, *B. alkalophilus*, *B. amyloliquefaciens*, *B. clausii*, *B. halodurans*, *B. megaterium*, *B. coagulans*, *B. circulans*, *B. gibsonii*, and *B. thuringiensis*. It is recognized that the genus *Bacillus* continues to undergo taxonomical reorganization. Thus, it is intended that the genus include species that have been reclassified, including but not limited to such organisms as *B. stearothermophilus*, which is now named “*Geobacillus stearothermophilus*”, or *B. polymyxa*, which is now “*Paenibacillus*

polymyxa". The production of resistant endospores under stressful environmental conditions is considered the defining feature of the genus *Bacillus*, although this characteristic also applies to the recently named *Alicyclobacillus*, *Amphibacillus*, *Aneurinibacillus*, *Anoxybacillus*, *Brevibacillus*, *Filobacillus*, *Gracilibacillus*, *Halobacillus*, *Paenibacillus*, *Salibacillus*, *Thermobacillus*, *Ureibacillus*, and *Virgibacillus*.

[0020] As used herein, the term "mutation" refers to man-made changes to a reference amino acid or nucleic acid sequence. It is intended that the term encompass man-made substitutions, insertions and deletions.

[0021] As used herein, the term "vector" refers to a nucleic acid construct used to introduce or transfer nucleic acid(s) into a target cell or tissue. A vector is typically used to introduce foreign DNA into a cell or tissue. Vectors include plasmids, cloning vectors, bacteriophages, viruses (e.g., viral vector), cosmids, expression vectors, shuttle vectors, and the like. A vector typically includes an origin of replication, a multicloning site, and a selectable marker. The process of inserting a vector into a target cell is typically referred to as transformation. The present invention includes, in some embodiments, a vector that comprises a DNA sequence encoding a serine protease polypeptide (e.g., precursor or mature serine protease polypeptide) that is operably linked to a suitable prosequence (e.g., secretory, signal peptide sequence, etc.) capable of effecting the expression of the DNA sequence in a suitable host, and the folding and translocation of the recombinant polypeptide chain.

[0022] As used herein in the context of introducing a nucleic acid sequence into a cell, the term "introduced" refers to any method suitable for transferring the nucleic acid sequence into the cell. Such methods for introduction include but are not limited to protoplast fusion, transfection, transformation, electroporation, conjugation, and transduction. Transformation refers to the genetic alteration of a cell which results from the uptake, optional genomic incorporation, and expression of genetic material (e.g., DNA).

[0023] "Expression cassette" or "expression vector" refers to a nucleic acid construct or vector generated recombinantly or synthetically for the expression of a nucleic acid of interest (e.g., a foreign nucleic acid or transgene) in a target cell. The nucleic acid of interest typically expresses a protein of interest. An expression vector or expression cassette typically comprises a promoter nucleotide sequence that drives or promotes expression of the foreign nucleic acid. The expression vector or cassette also typically includes other specified nucleic acid elements that permit transcription of a particular nucleic acid in a target cell. A recombinant expression cassette can be incorporated into a plasmid, chromosome, mitochondrial DNA, plastid DNA, virus, or nucleic acid fragment. Some expression vectors have the ability to incorporate and express heterologous DNA fragments in a host cell or genome of the host cell. Many prokaryotic and eukaryotic expression vectors are commercially available. Selection of appropriate

expression vectors for expression of a protein from a nucleic acid sequence incorporated into the expression vector is within the knowledge of those of skill in the art.

[0024] As used herein, a nucleic acid is “operably linked” with another nucleic acid sequence when it is placed into a functional relationship with another nucleic acid sequence. For example, a promoter or enhancer is operably linked to a nucleotide coding sequence if the promoter affects the transcription of the coding sequence. A ribosome binding site may be operably linked to a coding sequence if it is positioned so as to facilitate translation of the coding sequence. Typically, “operably linked” DNA sequences are contiguous. However, enhancers do not have to be contiguous. Linking is accomplished by ligation at convenient restriction sites. If such sites do not exist, synthetic oligonucleotide adaptors or linkers may be used in accordance with conventional practice.

[0025] As used herein the term “gene” refers to a polynucleotide (e.g., a DNA segment), that encodes a polypeptide and includes regions preceding and following the coding regions. In some instances a gene includes intervening sequences (introns) between individual coding segments (exons).

[0026] As used herein, “recombinant” when used with reference to a cell typically indicates that the cell has been modified by the introduction of a foreign nucleic acid sequence or that the cell is derived from a cell so modified. For example, a recombinant cell may comprise a gene not found in identical form within the native (non-recombinant) form of the cell, or a recombinant cell may comprise a native gene (found in the native form of the cell) that has been modified and re-introduced into the cell. A recombinant cell may comprise a nucleic acid endogenous to the cell that has been modified without removing the nucleic acid from the cell; such modifications include those obtained by gene replacement, site-specific mutation, and related techniques known to those of ordinary skill in the art. Recombinant DNA technology includes techniques for the production of recombinant DNA in vitro and transfer of the recombinant DNA into cells where it may be expressed or propagated, thereby producing a recombinant polypeptide. “Recombination” and “recombining” of polynucleotides or nucleic acids refer generally to the assembly or combining of two or more nucleic acid or polynucleotide strands or fragments to generate a new polynucleotide or nucleic acid.

[0027] A nucleic acid or polynucleotide is said to “encode” a polypeptide if, in its native state or when manipulated by methods known to those of skill in the art, it can be transcribed and/or translated to produce the polypeptide or a fragment thereof. The anti-sense strand of such a nucleic acid is also said to encode the sequence.

[0028] The terms “host strain” and “host cell” refer to a suitable host for an expression vector comprising a DNA sequence of interest.

[0029] A “protein” or “polypeptide” comprises a polymeric sequence of amino acid residues. The terms “protein” and “polypeptide” are used interchangeably herein. The single and 3-letter code for

amino acids as defined in conformity with the IUPAC-IUB Joint Commission on Biochemical Nomenclature (JCBN) is used throughout this disclosure. The single letter X refers to any of the twenty amino acids. It is also understood that a polypeptide may be coded for by more than one nucleotide sequence due to the degeneracy of the genetic code.

[0030] A “prosequence” or “propeptide sequence” refers to an amino acid sequence between the signal peptide sequence and mature protease sequence that is necessary for the proper folding and secretion of the protease; they are sometimes referred to as intramolecular chaperones. Cleavage of the prosequence or propeptide sequence results in a mature active protease. Bacterial serine proteases are often expressed as pro-enzymes.

[0031] The terms “signal sequence” and “signal peptide” refer to a sequence of amino acid residues that may participate in the secretion or direct transport of the mature or precursor form of a protein. The signal sequence is typically located N-terminal to the precursor or mature protein sequence. The signal sequence may be endogenous or exogenous. A signal sequence is normally absent from the mature protein. A signal sequence is typically cleaved from the protein by a signal peptidase after the protein is transported.

[0032] The term “mature” form of a protein, polypeptide, or peptide refers to the functional form of the protein, polypeptide, or peptide without the signal peptide sequence and propeptide sequence.

[0033] The term “precursor” form of a protein or peptide refers to a mature form of the protein having a prosequence operably linked to the amino or carboxyl terminus of the protein. The precursor may also have a “signal” sequence operably linked to the amino terminus of the prosequence. The precursor may also have additional polypeptides that are involved in post-translational activity (e.g., polypeptides cleaved therefrom to leave the mature form of a protein or peptide).

[0034] The term “wild-type” in reference to an amino acid sequence or nucleic acid sequence indicates that the amino acid sequence or nucleic acid sequence is a native or naturally-occurring sequence. As used herein, the term “naturally-occurring” refers to anything (e.g., protein or polynucleotide sequences) that is found in nature. Conversely, the term “non-naturally occurring” refers to anything that is not found in nature (e.g., recombinant polynucleotide and protein sequences produced in the laboratory or modification of the wild-type sequence).

[0035] As used herein with regard to amino acid residue positions, “corresponding to” or “corresponds to” or “corresponds” refers to an amino acid residue at the enumerated position in a protein or peptide, or an amino acid residue that is analogous, homologous, or equivalent to an enumerated residue in a protein or peptide. As used herein, “corresponding region” generally refers to an analogous position in a related proteins or a reference protein.

[0036] The terms “derived from” and “obtained from” refer to not only a protein produced or

producible by a strain of the organism in question, but also a protein encoded by a DNA sequence isolated from such strain and produced in a host organism containing such DNA sequence. Additionally, the term refers to a protein which is encoded by a DNA sequence of synthetic and/or cDNA origin and which has the identifying characteristics of the protein in question. To exemplify, “proteases derived from *Bacillus*” refers to those enzymes having proteolytic activity that are naturally produced by *Bacillus*, as well as to serine proteases like those produced by *Bacillus* sources but which through the use of genetic engineering techniques are produced by other host cells transformed with a nucleic acid encoding the serine proteases.

[0037] The term “identical” in the context of two polynucleotide or polypeptide sequences refers to the nucleic acids or amino acids in the two sequences that are the same when aligned for maximum correspondence, as measured using sequence comparison or analysis algorithms described below and known in the art.

[0038] “% identity” or percent identity” or “PID” refers to protein sequence identity. Percent identity may be determined using standard techniques known in the art. The percent amino acid identity shared by sequences of interest can be determined by aligning the sequences to directly compare the sequence information, e.g., by using a program such as BLAST, MUSCLE, or CLUSTAL. The BLAST algorithm is described, for example, in Altschul et al., *J Mol Biol*, 215:403-410 (1990) and Karlin et al., *Proc Natl Acad Sci USA*, 90:5873-5787 (1993). A percent (%) amino acid sequence identity value is determined by the number of matching identical residues divided by the total number of residues of the “reference” sequence including any gaps created by the program for optimal/maximum alignment.

BLAST algorithms refer to the “reference” sequence as the “query” sequence.

[0039] As used herein, “homologous proteins” or “homologous proteases” refers to proteins that have distinct similarity in primary, secondary, and/or tertiary structure. Protein homology can refer to the similarity in linear amino acid sequence when proteins are aligned. Homology can be determined by amino acid sequence alignment, e.g., using a program such as BLAST, MUSCLE, or CLUSTAL. Homologous search of protein sequences can be done using BLASTP and PSI-BLAST from NCBI BLAST with threshold (E-value cut-off) at 0.001. (Altschul et al., “Gapped BLAST and PSI BLAST a new generation of protein database search programs”, *Nucleic Acids Res*, Set 1;25(17):3389-402(1997)). The BLAST program uses several search parameters, most of which are set to the default values. The NCBI BLAST algorithm finds the most relevant sequences in terms of biological similarity but is not recommended for query sequences of less than 20 residues (Altschul et al., *Nucleic Acids Res*, 25:3389-3402, 1997 and Schaffer et al., *Nucleic Acids Res*, 29:2994-3005, 2001). Exemplary default BLAST parameters for a nucleic acid sequence searches include: Neighboring words threshold=11; E-value cutoff=10; Scoring Matrix=NUC.3.1 (match=1, mismatch=-3); Gap Opening=5; and Gap Extension=2. Exemplary default BLAST parameters for amino acid sequence searches include: Word size = 3; E-value

cutoff=10; Scoring Matrix=BLOSUM62; Gap Opening=11; and Gap extension=1. Using this information, protein sequences can be grouped and/or a phylogenetic tree built therefrom. Amino acid sequences can be entered in a program such as the Vector NTI Advance suite and a Guide Tree can be created using the Neighbor Joining (NJ) method (Saitou and Nei, Mol Biol Evol, 4:406-425, 1987). The tree construction can be calculated using Kimura's correction for sequence distance and ignoring positions with gaps. A program such as AlignX can display the calculated distance values in parenthesis following the molecule name displayed on the phylogenetic tree.

[0040] Understanding the homology between molecules can reveal the evolutionary history of the molecules as well as information about their function; if a newly sequenced protein is homologous to an already characterized protein, there is a strong indication of the new protein's biochemical function. The most fundamental relationship between two entities is homology; two molecules are said to be homologous if they have been derived from a common ancestor. Homologous molecules, or homologs, can be divided into two classes, paralogs and orthologs. Paralogs are homologs that are present within one species. Paralogs often differ in their detailed biochemical functions. Orthologs are homologs that are present within different species and have very similar or identical functions. A protein superfamily is the largest grouping (clade) of proteins for which common ancestry can be inferred. Usually this common ancestry is based on sequence alignment and mechanistic similarity. Superfamilies typically contain several protein families which show sequence similarity within the family. The term "protein clan" is commonly used for protease superfamilies based on the MEROPS protease classification system.

[0041] The CLUSTAL W algorithm is another example of a sequence alignment algorithm (See, Thompson et al., Nucleic Acids Res, 22:4673-4680, 1994). Default parameters for the CLUSTAL W algorithm include: Gap opening penalty=10.0; Gap extension penalty=0.05; Protein weight matrix=BLOSUM series; DNA weight matrix=IUB; Delay divergent sequences %=40; Gap separation distance=8; DNA transitions weight=0.50; List hydrophilic residues=GPSNDQEKR; Use negative matrix=OFF; Toggle Residue specific penalties=ON; Toggle hydrophilic penalties=ON; and Toggle end gap separation penalty=OFF. In CLUSTAL algorithms, deletions occurring at either terminus are included. For example, a variant with a five amino acid deletion at either terminus (or within the polypeptide) of a polypeptide of 500 amino acids would have a percent sequence identity of 99% (495/500 identical residues $\times$ 100) relative to the "reference" polypeptide. Such a variant would be encompassed by a variant having "at least 99% sequence identity" to the polypeptide.

[0042] A nucleic acid or polynucleotide is "isolated" when it is at least partially or completely separated from other components, including but not limited to for example, other proteins, nucleic acids, cells, etc. Similarly, a polypeptide, protein or peptide is "isolated" when it is at least partially or completely separated from other components, including but not limited to for example, other proteins,

nucleic acids, cells, etc. On a molar basis, an isolated species is more abundant than are other species in a composition. For example, an isolated species may comprise at least about 60%, about 65%, about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, about 99%, or about 100% (on a molar basis) of all macromolecular species present. Preferably, the species of interest is purified to essential homogeneity (i.e., contaminant species cannot be detected in the composition by conventional detection methods). Purity and homogeneity can be determined using a number of techniques well known in the art, such as agarose or polyacrylamide gel electrophoresis of a nucleic acid or a protein sample, respectively, followed by visualization upon staining. If desired, a high-resolution technique, such as high performance liquid chromatography (HPLC) or a similar means can be utilized for purification of the material.

[0043] The term “purified” as applied to nucleic acids or polypeptides generally denotes a nucleic acid or polypeptide that is essentially free from other components as determined by analytical techniques well known in the art (e.g., a purified polypeptide or polynucleotide forms a discrete band in an electrophoretic gel, chromatographic eluate, and/or a media subjected to density gradient centrifugation). For example, a nucleic acid or polypeptide that gives rise to essentially one band in an electrophoretic gel is “purified.” A purified nucleic acid or polypeptide is at least about 50% pure, usually at least about 60%, about 65%, about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, about 99%, about 99.5%, about 99.6%, about 99.7%, about 99.8% or more pure (e.g., percent by weight on a molar basis). In a related sense, a composition is enriched for a molecule when there is a substantial increase in the concentration of the molecule after application of a purification or enrichment technique. The term “enriched” refers to a compound, polypeptide, cell, nucleic acid, amino acid, or other specified material or component that is present in a composition at a relative or absolute concentration that is higher than a starting composition.

[0044] As used herein, the term “functional assay” refers to an assay that provides an indication of a protein’s activity. In some embodiments, the term refers to assay systems in which a protein is analyzed for its ability to function in its usual capacity. For example, in the case of a protease, a functional assay involves determining the effectiveness of the protease to hydrolyze a proteinaceous substrate.

[0045] The term “cleaning activity” refers to a cleaning performance achieved by a serine protease polypeptide or reference subtilisin under conditions prevailing during the proteolytic, hydrolyzing, cleaning, or other process of the disclosure. In some embodiments, cleaning performance of a serine protease or reference subtilisin may be determined by using various assays for cleaning one or more enzyme sensitive stain on an item or surface (e.g., a stain resulting from food, grass, blood, ink, milk, oil, and/or egg protein). Cleaning performance of one or more subtilisin variant described herein or reference subtilisin can be determined by subjecting the stain on the item or surface to standard wash condition(s)

and assessing the degree to which the stain is removed by using various chromatographic, spectrophotometric, or other quantitative methodologies. Exemplary cleaning assays and methods are known in the art and include, but are not limited to those described in WO99/34011 and US 6,605,458, as well as those cleaning assays and methods included in the Examples provided below.

[0046] The term “cleaning effective amount” of one or more subtilisin variant described herein or reference subtilisin refers to the amount of protease that achieves a desired level of enzymatic activity in a specific cleaning composition. Such effective amounts are readily ascertained by one of ordinary skill in the art and are based on many factors, such as the particular protease used, the cleaning application, the specific composition of the cleaning composition, and whether a liquid or dry (e.g., granular, tablet, bar) composition is required, etc.

[0047] The term “cleaning adjunct material” refers to any liquid, solid, or gaseous material included in cleaning composition other than one or more subtilisin variant described herein, or recombinant polypeptide or active fragment thereof. In some embodiments, the cleaning compositions of the present disclosure include one or more cleaning adjunct materials. Each cleaning adjunct material is typically selected depending on the particular type and form of cleaning composition (e.g., liquid, granule, powder, bar, paste, spray, tablet, gel, foam, or other composition). Preferably, each cleaning adjunct material is compatible with the protease enzyme used in the composition.

[0048] Cleaning compositions and cleaning formulations include any composition that is suited for cleaning, bleaching, disinfecting, and/or sterilizing any object, item, and/or surface. Such compositions and formulations include, but are not limited to for example, liquid and/or solid compositions, including cleaning or detergent compositions (e.g., liquid, tablet, gel, bar, granule, and/or solid laundry cleaning or detergent compositions and fine fabric detergent compositions; hard surface cleaning compositions and formulations, such as for glass, wood, ceramic and metal counter tops and windows; carpet cleaners; oven cleaners; fabric fresheners; fabric softeners; and textile, laundry booster cleaning or detergent compositions, laundry additive cleaning compositions, and laundry pre-spotter cleaning compositions; dishwashing compositions, including hand or manual dishwashing compositions (e.g., “hand” or “manual” dishwashing detergents) and automatic dishwashing compositions (e.g., “automatic dishwashing detergents”). Single dosage unit forms also find use with the present invention, including but not limited to pills, tablets, gelcaps, or other single dosage units such as pre-measured powders or liquids.

[0049] Cleaning composition or cleaning formulations, as used herein, include, unless otherwise indicated, granular or powder-form all-purpose or heavy-duty washing agents, especially cleaning detergents; liquid, granular, gel, solid, tablet, paste, or unit dosage form all-purpose washing agents, especially the so-called heavy-duty liquid (HDL) detergent or heavy-duty dry (HDD) detergent types;

liquid fine-fabric detergents; hand or manual dishwashing agents, including those of the high-foaming type; hand or manual dishwashing, automatic dishwashing, or dishware or tableware washing agents, including the various tablet, powder, solid, granular, liquid, gel, and rinse-aid types for household and institutional use; liquid cleaning and disinfecting agents, including antibacterial hand-wash types, cleaning bars, mouthwashes, denture cleaners, car shampoos, carpet shampoos, bathroom cleaners; hair shampoos and/or hair-rinses for humans and other animals; shower gels and foam baths and metal cleaners; as well as cleaning auxiliaries, such as bleach additives and “stain-stick” or pre-treat types. In some embodiments, granular compositions are in “compact” form; in some embodiments, liquid compositions are in a “concentrated” form.

[0050] As used herein, “fabric cleaning compositions” include hand and machine laundry detergent compositions including laundry additive compositions and compositions suitable for use in the soaking and/or pretreatment of stained fabrics (e.g., clothes, linens, and other textile materials).

[0051] As used herein, “non-fabric cleaning compositions” include non-textile (i.e., non-fabric) surface cleaning compositions, including, but not limited to for example, hand or manual or automatic dishwashing detergent compositions, oral cleaning compositions, denture cleaning compositions, contact lens cleaning compositions, wound debridement compositions, and personal cleansing compositions.

[0052] As used herein, the term “detergent composition” or “detergent formulation” is used in reference to a composition intended for use in a wash medium for the cleaning of soiled or dirty objects, including particular fabric and/or non-fabric objects or items. In some embodiments, the detergents of the disclosure comprise one or more subtilisin variant described herein and, in addition, one or more surfactants, transferase(s), hydrolytic enzymes, oxido reductases, builders (e.g., a builder salt), bleaching agents, bleach activators, bluing agents, fluorescent dyes, caking inhibitors, masking agents, enzyme activators, antioxidants, and/or solubilizers. In some instances, a builder salt is a mixture of a silicate salt and a phosphate salt, preferably with more silicate (e.g., sodium metasilicate) than phosphate (e.g., sodium tripolyphosphate). Some embodiments are directed to cleaning compositions or detergent compositions that do not contain any phosphate (e.g., phosphate salt or phosphate builder).

[0053] As used herein, the term “bleaching” refers to the treatment of a material (e.g., fabric, laundry, pulp, etc.) or surface for a sufficient length of time and/or under appropriate pH and/or temperature conditions to effect a brightening (i.e., whitening) and/or cleaning of the material. Examples of chemicals suitable for bleaching include, but are not limited to, for example, ClO_2 , H_2O_2 , peracids, NO_2 , etc.

[0054] As used herein, “wash performance” of a protease (e.g., one or more subtilisin variant described herein, or recombinant polypeptide or active fragment thereof) refers to the contribution of one or more subtilisin variant described herein to washing that provides additional cleaning performance to

the detergent as compared to the detergent without the addition of the one or more subtilisin variant described herein to the composition. Wash performance is compared under relevant washing conditions. In some test systems, other relevant factors, such as detergent composition, sud concentration, water hardness, washing mechanics, time, pH, and/or temperature, can be controlled in such a way that condition(s) typical for household application in a certain market segment (e.g., hand or manual dishwashing, automatic dishwashing, dishware cleaning, tableware cleaning, fabric cleaning, etc.) are imitated.

[0055] The term “relevant washing conditions” is used herein to indicate the conditions, particularly washing temperature, time, washing mechanics, sud concentration, type of detergent and water hardness, actually used in households in a hand dishwashing, automatic dishwashing, or laundry detergent market segment.

[0056] As used herein, the term “disinfecting” refers to the removal of contaminants from the surfaces, as well as the inhibition or killing of microbes on the surfaces of items.

[0057] The “compact” form of the cleaning compositions herein is best reflected by density and, in terms of composition, by the amount of inorganic filler salt. Inorganic filler salts are conventional ingredients of detergent compositions in powder form. In conventional detergent compositions, the filler salts are present in substantial amounts, typically about 17 to about 35% by weight of the total composition. In contrast, in compact compositions, the filler salt is present in amounts not exceeding about 15% of the total composition. In some embodiments, the filler salt is present in amounts that do not exceed about 10%, or more preferably, about 5%, by weight of the composition. In some embodiments, the inorganic filler salts are selected from the alkali and alkaline-earth-metal salts of sulfates and chlorides. In some embodiments, the filler salt is sodium sulfate.

[0058] Disclosed herein is one or more subtilisin variant useful for cleaning applications and in methods of cleaning, as well as in a variety of industrial applications. Disclosed herein is one or more isolated, recombinant, substantially pure, or non-naturally occurring subtilisin variant. In some embodiments, one or more subtilisin variant described herein is useful in cleaning applications and can be incorporated into cleaning compositions that are useful in methods of cleaning an item or a surface in need thereof.

[0059] One embodiment provides one or more subtilisin variant comprising two, three, or four or more variations versus SEQ ID NO:2 at positions selected from: (i) 1, 3, 9, 10, 15, 17, 19, 22, 24, 25, 26, 27, 28, 30, 35, 37, 38, 43, 45, 48, 68, 71, 74, 76, 77, 78, 86, 87, 91, 95, 96, 98, 99, 100, 102, 103, 108, 111, 114, 115, 120, 123, 125, 126, 127, 128, 129, 130, 136, 143, 146, 147, 151, 155, 160, 165, 183, 184, 187, 193, 202, 203, 210, 217, 234, 238, 239, 240, 242, 247, 250, 251, 255, 258, 259, 260, and 274; (ii) 1, 3, 9, 15, 22, 24, 28, 30, 35, 37, 45, 48, 68, 71, 74, 76, 77, 78, 86, 87, 95, 96, 98, 99, 100, 108, 114, 115,

120, 123, 125, 126, 127, 128, 129, 130, 143, 146, 147, 151, 155, 165, 183, 184, 187, 193, 202, 203, 210, 217, 234, 238, 239, 240, 242, 247, 250, 255, 258, 259, 260, and 274; (iii) 1, 3, 15, 35, 68, 71, 77, 86, 87, 95, 99, 115, 123, 127, 128, 147, 155, 165, 184, 202, 210, 217, 234, 239, 242, 250, 258, and 274; or (iv) 3, 68, 77, 86, 99, 115, 123, 127, 128, 165, 184, 202, 210, 217, and 258; with the proviso that one or more of said two, three, or four or more variations is a non-naturally occurring amino acid; wherein said variant has 85% or more amino acid sequence identity to an AprL-Clade subtilisin enzyme comprising a Motif A and/or a Motif B; wherein Motif A is selected from Motif 1, Motif 2, Motif 3, and Motif 4; wherein Motif B is selected from Motif 5, Motif 6, Motif 7, Motif 8, Motif 9, Motif 10, Motif 11, Motif 12, Motif 13, and Motif 14; and wherein the amino acid positions of the variant are numbered by correspondence with the amino acid sequence of SEQ ID NO:2. Another embodiment is directed to one or more subtilisin variant comprising an amino acid sequence comprising two, three, or four or more variations at positions corresponding to SEQ ID NO:2, wherein said positions are selected from: (i) 1, 3, 9, 10, 15, 22, 24, 26, 27, 28, 29, 30, 31, 35, 37, 40, 43, 45, 48, 52, 68, 71, 76, 77, 78, 86, 87, 95, 96, 98, 99, 100, 101, 102, 103, 105, 108, 111, 114, 115, 117, 119, 123, 125, 126, 127, 128, 129, 130, 136, 143, 146, 147, 151, 155, 158, 160, 161, 165, 184, 187, 193, 202, 203, 204, 210, 211, 216, 217, 234, 238, 239, 242, 243, 250, 255, 258, 259, 260, 264, and 274; (ii) 1, 3, 9, 10, 15, 22, 24, 26, 28, 29, 30, 31, 35, 37, 40, 45, 48, 52, 68, 71, 76, 77, 78, 86, 87, 95, 96, 98, 99, 100, 101, 102, 105, 108, 111, 114, 115, 117, 119, 123, 125, 126, 127, 128, 129, 130, 136, 143, 146, 147, 151, 155, 158, 160, 161, 165, 184, 187, 193, 202, 203, 204, 210, 211, 216, 217, 234, 238, 239, 242, 243, 250, 255, 258, 259, 260, 264, and 274; (iii) 1, 3, 9, 10, 24, 26, 29, 30, 31, 35, 37, 40, 45, 48, 52, 68, 71, 77, 78, 86, 87, 95, 96, 98, 99, 100, 101, 102, 105, 108, 111, 115, 117, 119, 123, 127, 128, 129, 130, 136, 143, 146, 147, 151, 155, 158, 160, 161, 165, 184, 187, 193, 202, 203, 204, 210, 211, 216, 217, 234, 238, 239, 242, 243, 250, 258, 259, 260, 264, and 274; (iv) 3, 9, 10, 35, 68, 77, 78, 86, 87, 99, 101, 115, 123, 127, 128, 165, 184, 187, 202, 203, 210, 217, 242, 250, 258, and 264; or (v) 3, 68, 77, 78, 123, 127, 128, 165, 184, 202, 210, 217, and 258; with the proviso that one or more of said two, three, or four or more variations is a man-made mutation or substitution; wherein said variant has at least 85% but less than 100% amino acid sequence identity to an AprL-Clade subtilisin enzyme comprising a Motif A and/or a Motif B; wherein Motif A is selected from Motif 1, Motif 2, Motif 3, and Motif 4; wherein Motif B is selected from Motif 5, Motif 6, Motif 7, Motif 8, Motif 9, Motif 10, Motif 11, Motif 12, Motif 13, and Motif 14; and wherein the amino acid positions of the variant are numbered by correspondence with the amino acid sequence of SEQ ID NO:2.

[0060] Still another embodiment provides one or more subtilisin comprising an amino acid sequence comprising two, three, or four or more amino acid variations versus SEQ ID NO:2 selected from: (i) 1Q, 3Q/V, 9E/T, 10M, 15I/V, 17H, 19E, 22Y, 24N, 25D, 26R, 27R, 28A, 30T, 35A, 37T, 38G, 43A, 45I/R, 48I, 68S, 71A, 74G, 76K, 77D/H/N/Q/S, 78I, 86H/N/R, 87S/T, 91G, 95A/Q, 96S, 98H/K/R, 99L/S,

100N/R, 102R, 103I, 108H/K, 111Q, 114N, 115F, 120A, 123I, 125N, 126Q, 127F/T, 128P/S, 129T, 130S, 136A/R, 143V, 146A/S, 147L, 151G, 155A/E/G, 160C/R, 165A/E/Q/S, 183A/G, 184Q, 187P, 193D, 202V, 203E/N, 210E/H/P, 217S, 234I/W, 238R/T, 239K/S/T, 240N, 242G, 247W, 250G, 251E, 255R, 258E/H/P, 259P, 260W, and 274A/F; (ii) 1Q, 3Q/V, 9E, 15I, 22Y, 24N, 28A, 30T, 35A, 37T, 45I, 48I, 68S, 71A, 74G, 76K, 77D/H/N/S, 78I, 86H/N/R, 87S/T, 95A/Q, 96S, 98K, 99L/S, 100N, 108H/K, 114N, 115F, 120A, 123I, 125N, 126Q, 127F/T, 128P/S, 129T, 130S, 143V, 146A/S, 147L, 151G, 155A/E/G, 165A/E/Q/S, 183G, 184Q, 187P, 193D, 202V, 203E/N, 210P, 217S, 234I/W, 238R/T, 239K/S/T, 240N, 242G, 247W, 250G, 255R, 258P, 259P, 260W, and 274A; (iii) 1Q, 3V, 15I, 35A, 68S, 71A, 77N, 86H, 87S, 95A, 99S, 115F, 123I, 127T, 128P/S, 147L, 155G, 165Q/S, 184Q, 202V, 210P, 217S, 234W, 239S, 242G, 250G, 258P, and 274A; or (iv) 3V, 68S, 77N, 86H, 99S, 115F, 123I, 127T, 128P, 165Q, 184Q, 202V, 210P, 217S, and 258P; with the proviso that one or more of said two, three, or four or more variations is a non-naturally occurring amino acid; wherein said variant has 85% or more amino acid sequence identity to an AprL-Clade subtilisin enzyme comprising a Motif A and/or a Motif B; wherein Motif A is selected from Motif 1, Motif 2, Motif 3, and Motif 4; wherein Motif B is selected from Motif 5, Motif 6, Motif 7, Motif 8, Motif 9, Motif 10, Motif 11, Motif 12, Motif 13, and Motif 14; and wherein the amino acid positions of the variant are numbered by correspondence with the amino acid sequence of SEQ ID NO:2.

[0061] Yet another embodiment provides one or more subtilisin variant comprising an amino acid sequence comprising two, three, or four or more amino acid substitutions selected from: (i) A/G1, T3, P9, L/Q10, K15, H/Q17, Q19, K22, A24, N25, V26, K27, G/V28, I/V30, I35, A/S/T37, S38, N/K43, V45, A48, A68, V71, L74, N76, S/T77, I/T78, N/S86, V87, A/I91, L95, N96, S98, G99, S100, S/T102, Y103, S108, E111, T114, A/Q/T115, V120, M123, L125, G126, G127, A/P/T128, S129, G/V130, Q136, A/S143, A/I/V146, V147, A/S151, S155, N/S/Y160, G165, G/K/N/S183, N/Q184, S187, A/P/S193, K194, A/S/V202, G/S203, S/T210, S211, N217, L234, P238, A/N/T239, A242, D/N/T247, S250, R/S251, N/Y255, D/S258, S259, F260, and Q274; (ii) A/G1, T3, P9, K15, K22, A24, G/V28, I/V30, I35, A/S/T37, V45, A48, A68, V71, L74, N76, S/T77, I/T78, N/S86, V87, L95, N96, S98, G99, S100, S108, T114, A/Q/T115, V120, M123, L125, G126, G127, A/P/T128, S129, G/V130, A/S143, A/I/V146, V147, A/S151, S155, G165, K/S183, N184, S187, A/P/S193, K194, A/S/V202, G/S203, S/T210, S211, N217, L234, P238, A/N/T239, A242, D/N/T247, S250, N/Y255, D/S258, S259, F260, and Q274; (iii) A/G1, T3, K15, I35, A68, V71, S/T77, N/S86, V87, L95, G99, A/Q/T115, M123, G127, A/P/T128, V147, S155, G165, N184, A/S/V202, S/T210, N217, L234, A/N/T239, A242, S250, D/S258, and Q274; or (iv) T3, A68, S/T77, N/S86, G99, A/Q/T115, M123, G127, A/P/T128, G165, N184, A/S/V202, S/T210, N217, and D/S258; with the proviso that one or more of said two, three, or four or more substitutions is a non-naturally occurring amino acid; wherein said variant has 85% or more amino acid sequence identity to an

AprL-Clade subtilisin enzyme comprising a Motif A and/or a Motif B; wherein Motif A is selected from Motif 1, Motif 2, Motif 3, and Motif 4; wherein Motif B is selected from Motif 5, Motif 6, Motif 7, Motif 8, Motif 9, Motif 10, Motif 11, Motif 12, Motif 13, and Motif 14; and wherein the amino acid positions of the variant are numbered by correspondence with the amino acid sequence of SEQ ID NO:2. Yet still another embodiment provides one or more subtilisin variant comprising an amino acid sequence comprising two, three, or four or more amino acid substitutions selected from: (i) A/G1, T3, P9, L10, K15, H/Q17, Q19, K22, A24, N25, V26, K27, G/V28, I/V30, I35, A/S/T37, S38, N/K43, V45, A48, A68, V71, L74, N76, T77, T78, N/S86, V87, A91, L95, N96, S98, G99, S100, S/T102, Y103, S108, E111, T114, Q/T115, V120, M123, L125, G126, G127, A/P128, S129, G/V130, Q136, A143, A/I/V146, V147, A/S151, S155, N/S160, G165, K/S183, N184, S187, A/S193, K194, A/V202, G/S203, S/T210, S211, N217, L234, P238, N/T239, A242, N/T247, S250, R/S251, N/Y255, D/S258, S259, F260, and Q274; (ii) A/G1, T3, P9, K15, K22, A24, G/V28, I/V30, I35, A/S/T37, V45, A48, A68, V71, L74, N76, T77, T78, N/S86, V87, L95, N96, S98, G99, S100, S108, T114, Q/T115, V120, M123, L125, G126, G127, A/P128, S129, G/V130, A143, A/I/V146, V147, A/S151, S155, G165, K/S183, N184, S187, A/S193, K194, A/V202, G/S203, S/T210, S211, N217, L234, P238, N/T239, A242, N/T247, S250, N/Y255, D/S258, S259, F260, and Q274; (iii) A/G1, T3, K15, I35, A68, V71, T77, N/S86, V87, L95, G99, Q/T115, M123, G127, A/P128, V147, S155, G165, N184, A/V202, S/T210, N217, L234, N/T239, A242, S250, D/S258, and Q274; or (iv) T3, A68, T77, N/S86, G99, Q/T115, M123, G127, A/P128, G165, N184, A/V202, S/T210, N217, and D/S258; with the proviso that one or more of said two, three, or four or more substitutions is a non-naturally occurring amino acid; wherein said variant has 85% or more amino acid sequence identity to an AprL-Clade subtilisin enzyme comprising a Motif A and/or a Motif B; wherein Motif A is selected from Motif 1, Motif 2, Motif 3, and Motif 4; wherein Motif B is selected from Motif 5, Motif 6, Motif 7, Motif 8, Motif 9, Motif 10, Motif 11, Motif 12, Motif 13, and Motif 14; and wherein the amino acid positions of the variant are numbered by correspondence with the amino acid sequence of SEQ ID NO:2.

[0062] A further embodiment provides one or more subtilisin variant comprising an amino acid sequence comprising two, three, or four or more amino acid substitutions selected from: (i) 1Q, 3Q/V, 9E/T, 10M, 15I/V, 17H, 19E, 22Y, 24N, 25D, 26R, 27R, 28A, 30T, 35A, 37T, 38G, 43A, 45I/R, 48I, 68S, 71A, 74G, 76K, 77D/H/N/Q/S, 78I, 86H/N/R/S, 87S/T, 91G, 95A/Q, 96S, 98H/K/R, 99L/S, 100N/R, 102R, 103I, 108H/K, 111Q, 114N, 115F, 120A, 123I, 125N, 126Q, 127F/T, 128P/S, 129T, 130S, 136A/R, 143V, 146A/S, 147L, 151G, 155A/E/G, 160C/R, 165A/E/Q/S, 183A/G, 184Q, 187P, 193D, 194E, 202V, 203E/N, 210E/H/P, 211N, 217S, 234I/W, 238R/T, 239K/S/T, 242G, 247W, 250G, 251E, 255R, 258E/H/P, 259P, 260W, and 274A/F; (ii) 1Q, 3Q/V, 9E, 15I, 22Y, 24N, 28A, 30T, 35A, 37T, 45I, 48I, 68S, 71A, 74G, 76K, 77D/H/N/S, 78I, 86H/N/R/S, 87S/T, 95A/Q, 96S, 98K, 99L/S, 100N,

108H/K, 114N, 115F, 120A, 123I, 125N, 126Q, 127F/T, 128P/S, 129T, 130S, 143V, 146A/S, 147L, 151G, 155A/E/G, 165A/E/Q/S, 183G, 184Q, 187P, 193D, 194E, 202V, 203E/N, 210P, 211N, 217S, 234I/W, 238R/T, 239K/S/T, 242G, 247W, 250G, 255R, 258P, 259P, 260W, and 274A; (iii) 1Q, 3V, 15I, 35A, 68S, 71A, 77N, 86H, 87S, 95A, 99S, 115F, 123I, 127T, 128P/S, 147L, 155G, 165Q/S, 184Q, 202V, 210P, 217S, 234W, 239S, 242G, 250G, 258P, and 274A; or (iv) 3V, 68S, 77N, 86H, 99S, 115F, 123I, 127T, 128P, 165Q, 184Q, 202V, 210P, 217S, and 258P; with the proviso that one or more of said two, three, or four or more substitutions is a non-naturally occurring amino acid; wherein said variant has 85% or more amino acid sequence identity to an AprL-Clade subtilisin enzyme comprising a Motif A and/or a Motif B; wherein Motif A is selected from Motif 1, Motif 2, Motif 3, and Motif 4; wherein Motif B is selected from Motif 5, Motif 6, Motif 7, Motif 8, Motif 9, Motif 10, Motif 11, Motif 12, Motif 13, and Motif 14; and wherein the amino acid positions of the variant are numbered by correspondence with the amino acid sequence of SEQ ID NO:2.

[0063] Yet a further embodiment is directed to one or more subtilisin variant comprising an amino acid sequence comprising two, three, or four or more amino acid substitutions at two, three, or four or more positions corresponding to SEQ ID NO:2, wherein said substitutions are selected from: (i) 1Q, 3Q/V, 9E, 10M, 15I, 22Y, 24N/Q, 26Q, 27R, 28A, 29S, 30T, 31I, 35A, 37T, 40E, 43A, 45I/Q/R, 48I/T/V, 52R, 68S, 71A, 76K, 77D/E/H/N/Q/S, 78I, 86H/N/R, 87S/T, 95A/F/Q, 96S, 98H/R, 99A/E/Q/S, 100N, 101A, 102D/T, 103I, 105N, 108H/K, 111Q, 114N, 115F, 117S, 119Q, 123I, 125N, 126Q, 127E/F/Q/S/T/V, 128P/R/S, 129H/R, 130S, 136N/R, 143Q/V, 146A/S, 147L, 151G, 155E/G/N, 158D/N/Q, 160C/D/K, 161D/Q, 165A/E/Q/S, 184E/Q, 187E/P, 193D/Q, 202V, 203E/N/Q, 204I, 210E/H/I/P, 211S, 216Q, 217S, 234I/W, 238Q/R, 239K/S/T, 242G/N, 243I, 250G, 255R, 258D/E/P, 259E/P, 260W, 264H/Q/S, and 274A; (ii) 1Q, 3Q/V, 9E, 10M, 15I, 22Y, 24N/Q, 26Q, 28A, 29S, 30T, 31I, 35A, 37T, 40E, 45I/Q/R, 48I/T/V, 52R, 68S, 71A, 76K, 77D/E/H/N/S, 78I, 86H/N/R, 87S/T, 95A/F/Q, 96S, 98K/R, 99A/E/Q/S, 100N, 101A, 102D/T, 105N, 108H/K, 111Q, 114N, 115F, 117S, 119Q, 123I, 125N, 126Q, 127E/F/Q/S/T/V, 128P/R/S, 129H/R, 130S, 136N, 143Q/V, 146A/S, 147L, 151G, 155E/G/N, 158D/N/Q, 160D/K, 161D/Q, 165A/E/Q/S, 184E/Q, 187E/P, 193D/Q, 194E, 202V, 203E/N/Q, 204I, 210I/P, 211S, 216Q, 217S, 234I/W, 238Q/R, 239K/S/T, 242G/N, 243I, 250G, 255R, 258D/P, 259E/P, 260W, 264H/Q/S, and 274A; (iii) 1Q, 3V, 9E, 10M, 24Q, 26Q, 29S, 30T, 31I, 35A, 37T, 40E, 45I/Q/R, 48T/V, 52R, 68S, 71A, 77E/N, 78I, 86H/R, 87S, 95A/F, 96S, 98R, 99A/E/Q/S, 100N, 101A, 102D/T, 105N, 108H, 111Q, 115F, 117S, 119Q, 123I, 127E/Q/S/T/V, 128P/R, 129H/R, 130S, 136N, 143Q/V, 146A, 147L, 151G, 155G/N, 158D/N/Q, 160D/K, 161D/Q, 165A/E/Q, 184E/Q, 187E/P, 193Q, 202V, 203E/N/Q, 204I, 210I/P, 211S, 216Q, 217S, 234W, 238Q, 239S, 242G/N, 243I, 250G, 258D/P, 259E/P, 260W, 264H/Q/S, and 274A; (iv) 3V, 9E, 10M, 35A, 68S, 77N, 78I, 86H, 87S, 99Q/S, 101A, 115F, 123I, 127S/T, 128P/R, 165E/Q, 184Q, 187E, 202V, 203E, 210P, 217S, 242G, 250G,

258D/P, and 264H; or (v) 3V, 68S, 77N, 78I, 123I, 127S, 128P, 165Q, 184Q, 202V, 210P, 217S, and 258D/P; with the proviso that one or more of said two, three, or four or more substitutions is a man-made substitution; wherein said variant has at least 85% but less than 100% amino acid sequence identity to an AprL-Clade subtilisin enzyme comprising a Motif A and/or a Motif B; wherein Motif A is selected from Motif 1, Motif 2, Motif 3, and Motif 4; wherein Motif B is selected from Motif 5, Motif 6, Motif 7, Motif 8, Motif 9, Motif 10, Motif 11, Motif 12, Motif 13, and Motif 14; and wherein the amino acid positions of the variant are numbered by correspondence with the amino acid sequence of SEQ ID NO:2.

[0064] A still further embodiment provides one or more subtilisin variant comprising an amino acid sequence comprising two, three, or four or more amino acid substitutions selected from: (i) A/G1Q, T3Q/V, P9E/T, L/Q10M, K15I/V, Q17H, Q19E, K22Y, A24N, N25D, V26R, K27R, G/V28A, I/V30T, I35A, A/S37T, S38G, N/K43A, V45I/R, A48I, A68S, V71A, L74G, N76K, S/T77D/H/N/Q/S, T78I, N/S86H/N/R/S, V87S/T, A/I91G, L95A/Q, N96S, S98H/K/R, G99L/S, S100N/R, S/T102R, Y103I, S108H/K, E111Q, T114N, A/Q/T115F, V120A, M123I, L125N, G126Q, G127F/T, A/P/T128P/S, S129T, G/V130S, Q136A/R, A/S143V, A/I/V146A/S, V147L, A/S151G, S155A/E/G, N/S/Y160C/R, G165A/E/Q/S, G/K/N/S183A/G, N184Q, S187P, A/P/S193D, K194E, A/S202V, G/S203E/N, S/T210E/H/P, S211N, N217S, L234I/W, P238R/T, A/N/T239K/S/T, A242G, D/N/T247W, S250G, R/S251E, N/Y255R, D/S258E/H/P, S259P, F260W, and Q274A/F; (ii) A/G1Q, T3Q/V, P9E, K15I, K22Y, A24N, G/V28A, I/V30T, I35A, A/S37T, V45I, A48I, A68S, V71A, L74G, N76K, S/T77D/H/N/S, T78I, N/S86H/N/R/S, V87S/T, L95A/Q, N96S, S98K, G99L/S, S100N, S108H/K, T114N, A/Q/T115F, V120A, M123I, L125N, G126Q, G127F/T, A/P/T128P/S, S129T, G/V130S, A/S143V, A/I/V146A/S, V147L, A/S151G, S155A/E/G, G165A/E/Q/S, K/S183G, N184Q, S187P, A/P/S193D, K194E, A/S202V, G/S203E/N, S/T210P, S211N, N217S, L234I/W, P238R/T, A/N/T239K/S/T, A242G, D/N/T247W, S250G, N/Y255R, D/S258P, S259P, F260W, and Q274A; (iii) A/G1Q, T3V, K15I, I35A, A68S, V71A, S/T77N, N/S86H, V87S, L95A, G99S, A/Q/T115F, M123I, G127T, A/P/T128P/S, V147L, S155G, G165Q/S, N184Q, A/S202V, S/T210P, N217S, L234W, A/N/T239S, A242G, S250G, D/S258P, and Q274A; or (iv) T3V, A68S, S/T77N, N/S86H, G99S, A/Q/T115F, M123I, G127T, A/T128P, G165Q, N184Q, A/S202V, S/T210P, N217S, and D/S258P; with the proviso that one or more of said two, three, or four or more substitutions is a non-naturally occurring amino acid; wherein said variant has 85% or more amino acid sequence identity to an AprL-Clade subtilisin enzyme comprising a Motif A and/or a Motif B; wherein Motif A is selected from Motif 1, Motif 2, Motif 3, and Motif 4; wherein Motif B is selected from Motif 5, Motif 6, Motif 7, Motif 8, Motif 9, Motif 10, Motif 11, Motif 12, Motif 13, and Motif 14; and wherein the amino acid positions of the variant are numbered by correspondence with the amino acid sequence of SEQ ID NO:2. Yet still a further embodiment provides one or more subtilisin variant comprising an amino acid sequence comprising two, three, or four or more amino acid

substitutions selected from: (i) A/G1Q, T3Q/V, P9E/T, L10M, K15I/V, Q17H, Q19E, K22Y, A24N, N25D, V26R, K27R, G/V28A, I/V30T, I35A, A/S37T, S38G, N/K43A, V45I/R, A48I, A68S, V71A, L74G, N76K, T77D/H/N/Q/S, T78I, N/S86H/N/R/S, V87S/T, A91G, L95A/Q, N96S, S98H/K/R, G99L/S, S100N/R, S/T102R, Y103I, S108H/K, E111Q, T114N, Q/T115F, V120A, M123I, L125N, G126Q, G127F/T, A/P128P/S, S129T, G/V130S, Q136A/R, A143V, A/I/V146A/S, V147L, A/S151G, S155A/E/G, N/S160C/R, G165A/E/Q/S, K/S183A/G, N184Q, S187P, A/S193D, K194E, A202V, G/S203E/N, S/T210E/H/P, S211N, N217S, L234I/W, P238R/T, N/T239K/S/T, A242G, N/T247W, S250G, R/S251E, N/Y255R, D/S258E/H/P, S259P, F260W, and Q274A/F; (ii) A/G1Q, T3Q/V, P9E, K15I, K22Y, A24N, G/V28A, I/V30T, I35A, A/S37T, V45I, A48I, A68S, V71A, L74G, N76K, T77D/H/N/S, T78I, N/S86H/N/R/S, V87S/T, L95A/Q, N96S, S98K, G99L/S, S100N, S108H/K, T114N, Q/T115F, V120A, M123I, L125N, G126Q, G127F/T, A/P128P/S, S129T, G/V130S, A143V, A/I/V146A/S, V147L, A/S151G, S155A/E/G, G165A/E/Q/S, K/S183G, N184Q, S187P, A/S193D, K194E, A202V, G/S203E/N, S/T210P, S211N, N217S, L234I/W, P238R/T, N/T239K/S/T, A242G, N/T247W, S250G, N/Y255R, D/S258P, S259P, F260W, and Q274A; (iii) A/G1Q, T3V, K15I, I35A, A68S, V71A, T77N, N/S86H, V87S, L95A, G99S, Q/T115F, M123I, G127T, A/P128P/S, V147L, S155G, G165Q/S, N184Q, A202V, S/T210P, N217S, L234W, N/T239S, A242G, S250G, D/S258P, and Q274A; or (iv) T3V, A68S, T77N, N/S86H, G99S, Q/T115F, M123I, G127T, A128P, G165Q, N184Q, A202V, S/T210P, N217S, and D/S258P; with the proviso that one or more of said two, three, or four or more substitutions is a non-naturally occurring amino acid; wherein said variant has 85% or more amino acid sequence identity to an AprL-Clade subtilisin enzyme comprising a Motif A and/or a Motif B; wherein Motif A is selected from Motif 1, Motif 2, Motif 3, and Motif 4; wherein Motif B is selected from Motif 5, Motif 6, Motif 7, Motif 8, Motif 9, Motif 10, Motif 11, Motif 12, Motif 13, and Motif 14; and wherein the amino acid positions of the variant are numbered by correspondence with the amino acid sequence of SEQ ID NO:2.

[0065] An even further embodiment is directed to one or more subtilisin variant comprising an amino acid sequence comprising two, three, or four or more amino acid substitutions selected from: (i) A/G1Q, T3Q/V, P/S9E, L/Q10M, K15I, K22Y, A24N/Q, V26Q, K27R, G/V28A, A/G29S, I/V30T, I/L31I, I35A, A/S37T, P/S/T40E, N/K43A, S/V45I/Q/R, A48I/T/V, A/S52R, A68S, V71A, N76K, S/T77D/E/H/N/Q/S, T78I, N/S86H/N/R, V87S/T, L95A/F/Q, N96S, S98H/R, G99A/E/Q/S, S100N, G101A, S/T102D/T, Y103I, A/G105N, S108H/K, E111Q, T114N, A/Q/S/T115F, G/N117S, D119Q, M123I, L125N, G126Q, G127E/F/Q/S/T/V, A/P/S128P/R/S, S129H/R, G/V130S, Q136N/R, A/S143Q/V, A/I/V146A/S, V147L, A/S151G, S155E/G/N, S158D/N/Q, N/S/Y160C/D/K, A/S/Q/T161D/Q, G165A/E/Q/S, N/Q184E/Q, S187E/P, A/P/S193D/Q, A/S202V, A/G/S203E/N/Q, V204I, S/T210E/H/I/P, N211S, L216Q, N217S, L234I/W, P238Q/R, A/N/T239K/S/T, A242G/N, S243I, S250G, N/Y255R,

D/S258D/E/P, S259E/P, F260W, K/N/R264H/Q/S, and Q274A; (ii) A1Q, T3Q/V, P9E, L10M, K15I, K22Y, A24N/Q, V26Q, K27R, V28A, A29S, V30T, L31I, I35A, A37T, P40E, N43A, V45I/Q/R, A48I/T/V, A52R, A68S, V71A, N76K, S77D/E/H/N/Q/S, T78I, S86H/N/R, V87S/T, L95A/F/Q, N96S, S98H/R, G99A/E/Q/S, S100N, G101A, S102D/T, Y103I, G105N, S108H/K, E111Q, T114N, T115F, G117S, D119Q, M123I, L125N, G126Q, G127E/F/Q/S/T/V, A128P/R/S, S129H/R, G130S, Q136N/R, A143Q/V, V146A/S, V147L, A151G, S155E/G/N, S158D/N/Q, N160C/D/K, T161D/Q, G165A/E/Q/S, N184E/Q, S187E/P, A193D/Q, A202V, G203E/N/Q, V204I, T210E/H/I/P, N211S, L216Q, N217S, L234I/W, P238Q/R, N239K/S/T, A242G/N, S243I, S250G, Y255R, S258D/E/P, S259E/P, F260W, K264H/Q/S, and Q274A; (iii) A/G1Q, T3Q/V, P/S9E, L/Q10M, K15I, K22Y, A24N/Q, V26Q, G/V28A, A/G29S, I/V30T, I/L31I, I35A, A/S37T, P/S/T40E, S/V45I/Q/R, A48I/T/V, A/S52R, A68S, V71A, N76K, S/T77D/E/H/N/S, T78I, N/S86H/N/R, V87S/T, L95A/F/Q, N96S, S98R, G99A/E/Q/S, S100N, G101A, S/T102D/T, A/G105N, S108H/K, E111Q, T114N, A/Q/S/T115F, G/N117S, D119Q, M123I, L125N, G126Q, G127E/F/Q/S/T/V, A/P/S128P/R/S, S129H/R, G/V130S, Q136N, A/S143Q/V, A/I/V146A/S, V147L, A/S151G, S155E/G/N, S158D/N/Q, N/S/Y160D/K, A/S/Q/T161D/Q, G165A/E/Q/S, N/Q184E/Q, S187E/P, A/P/S193D/Q, A/S202V, A/G/S203E/N/Q, V204I, S/T210I/P, N211S, L216Q, N217S, L234I/W, P238Q/R, A/N/T239K/S/T, A242G/N, S243I, S250G, N/Y255R, D/S258D/P, S259E/P, F260W, K/N/R264H/Q/S, and Q274A; (iv) A1Q, T3Q/V, P9E, L10M, K15I, K22Y, A24N/Q, V26Q, V28A, A29S, V30T, L31I, I35A, A37T, P40E, V45I/Q/R, A48I/T/V, A52R, A68S, V71A, N76K, T77D/E/H/N/S, T78I, S86H/N/R, V87S/T, L95A/F/Q, N96S, S98R, G99A/E/Q/S, S100N, G101A, S102D/T, G105N, S108H/K, E111Q, T114N, T115F, G117S, D119Q, M123I, L125N, G126Q, G127E/F/Q/S/T/V, A128P/R/S, S129H/R, G130S, Q136N, A143Q/V, V146A/S, V147L, A151G, S155E/G/N, S158D/N/Q, N160D/K, T161D/Q, G165A/E/Q/S, N184E/Q, S187E/P, A193D/Q, A202V, G203E/N/Q, V204I, T210I/P, N211S, L216Q, N217S, L234I/W, P238Q/R, N239K/S/T, A242G/N, S243I, S250G, Y255R, S258D/P, S259E/P, F260W, K264H/Q/S, and Q274A; (v) A/G1Q, T3V, P/S9E, L/Q10M, A24Q, V26Q, A/G29S, I/V30T, I/L31I, I35A, A/S37T, P/S/T40E, S/V45I/Q/R, A48T/V, A/S52R, A68S, V71A, S/T77E/N, T78I, N/S86H/R, V87S, L95A/F, N96S, S98R, G99A/E/Q/S, S100N, G101A, S/T102D/T, A/G105N, S108H, E111Q, A/Q/S/T115F, G/N117S, D119Q, M123I, G127E/Q/S/T/V, A/P/S128P/R, S129H/R, G/V130S, Q136N, A/S143Q/V, A/I/V146A, V147L, A/S151G, S155G/N, S158D/N/Q, N/S/Y160D/K, A/S/Q/T161D/Q, G165A/E/Q, N184E/Q, S187E/P, A/P/S193Q, A/S202V, A/G/S203E/N/Q, V204I, S/T210I/P, N211S, L216Q, N217S, L234W, P238Q, A/N/T239S, A242G/N, S243I, S250G, D/S258D/P, S259E/P, F260W, K/N/R264H/Q/S, and Q274A; (vi) A1Q, T3V, P9E, L10M, A24Q, V26Q, A29S, V30T, L31I, I35A, A37T, P40E, V45I/Q/R, A48T/V, A52R, A68S, V71A, T77E/N, T78I, S86H/R, V87S, L95A/F, N96S, S98R, G99A/E/Q/S, S100N, G101A, S102D/T, G105N, S108H, E111Q, T115F, G117S, D119Q, M123I, G127E/Q/S/T/V, A128P/R,

S129H/R, G130S, Q136N, A143Q/V, V146A, V147L, A151G, S155G/N, S158D/N/Q, N160D/K, T161D/Q, G165A/E/Q, N184E/Q, S187E/P, A193Q, A202V, G203E/N/Q, V204I, T210I/P, N211S, L216Q, N217S, L234W, P238Q, N239S, A242G/N, S243I, S250G, S258D/P, S259E/P, F260W, K264H/Q/S, and Q274A; (vii) T3V, P/S9E, L/Q10M, I35A, A68S, S/T77N, T78I, N/S86H, V87S, G99Q/S, G101A, A/Q/S/T115F, M123I, G127S/T, A/P/S128P/R, G165E/Q, N184Q, S187E, A/S202V, A/G/S203E, S/T210P, N217S, A242G, S250G, D/S258D/P, and K/N/R264H; (viii) T3V, P9E, L10M, I35A, A68S, T77N, T78I, S86H, V87S, G99Q/S, G101A, T115F, M123I, G127S/T, A128P/R, G165E/Q, N184Q, S187E, A202V, G203E, T210P, N217S, A242G, S250G, S258D/P, and K264H; (ix) T3V, A68S, S/T77N, T78I, M123I, G127S, A/S128P, G165Q, N184Q, A/S202V, S/T210P, N217S, and D/S258D/P; or (x) T3V, A68S, T77N, T78I, M123I, G127S, A128P, G165Q, N184Q, A202V, T210P, N217S, and S258D/P; with the proviso that one or more of said two, three, or four or more substitutions is a man-made substitution; wherein said variant has at least 85% but less than 100% amino acid sequence identity to an AprL-Clade subtilisin enzyme comprising a Motif A and/or a Motif B; wherein Motif A is selected from Motif 1, Motif 2, Motif 3, and Motif 4; wherein Motif B is selected from Motif 5, Motif 6, Motif 7, Motif 8, Motif 9, Motif 10, Motif 11, Motif 12, Motif 13, and Motif 14; and wherein the amino acid positions of the variant are numbered by correspondence with the amino acid sequence of SEQ ID NO:2.

[0066] A further embodiment is directed to one or more subtilisin variant comprising an amino acid sequence comprising one or more amino acid substitution sets selected from: (i) T77N-G165Q, T77N-N217S, G165Q-N217S, A68S-T77N, N184Q-N217S, A68S-G165Q, G165Q-N184Q, T77N-A128P, T77N-N184Q, A68S-N217S, A128P-N217S, A128P-G165Q, A68S-A128P, A68S-N184Q, A128P-N184Q, T3V-N217S, A202V-N217S, T3V-T77N, T3V-N184Q, N184Q-A202V, G165Q-A202V, M123I-G165Q, T3V-G165Q, T3V-A128P, T3V-A68S, T77N-A202V, M123I-N217S, A68S-A202V, M123I-N184Q, A128P-A202V, T77N-M123I, T3V-A202V, N217S-S258P, A68S-M123I, N184Q-S258P, G165Q-S258P, M123I-A128P, T77N-S258P, T3V-M123I, A128P-S258P, A202V-S258P, A68S-S258P, T3V-S258P, M123I-A202V, M123I-S258P, A68S-G127S, G127S-A128P, T77N-G127S, G127S-A202V, G127S-N184Q, G127S-N217S, G127S-G165Q, T3V-G127S, G127S-S258P, M123I-G127S, T77N-T210P, G165Q-T210P, T210P-N217S, A128P-T210P, A68S-T210P, N184Q-T210P, T210P-S258P, A202V-T210P, M123I-T210P, T3V-T210P, G127S-T210P, G127T-N184Q, G127T-N217S, V87S-G165Q, M123I-Q136R, A68S-G127T, G127T-A128P, T77N-V87S, G127T-G165Q, T77N-G127T, T78I-A202V, A68S-S258D, G165Q-S258D, T77N-S258D, T77N-T78I, T78I-G165Q, N184Q-S258D, N217S-S258D, P9E-M123I, A128P-A242G, A202V-S258D, A68S-T78I, G165Q-A242G, T77N-A242G, T78I-A128P, T78I-G127S, P9E-G165Q, Q136R-G165Q, T78I-N184Q, V87S-A128P, V87S-N217S, A68S-V87S, G127T-A202V, I35A-T77N, T3V-T78I, T78I-N217S, A68S-A242G, A68S-G99S, G99S-A128P, G99S-G165Q, I35A-G165Q, L10M-A202V, T3V-G127T, T77N-G99S, T78I-S258P,

V45R-M123I, V87S-M123I, M123I-N239S, P9E-N184Q, S155G-G165Q, T78I-M123I, V45R-Q136R, V87S-A202V, A128P-S258D, I35A-A202V, M123I-G127T, M123I-S258D, N184Q-N239S, P9E-N217S, Q136R-S258E, S86H-S258P, T3V-A242G, T3V-S258D, G99S-N184Q, G99S-N217S, I35A-V87S, N217S-A242G, P9E-Q136R, G127T-S258P, I35A-A128P, I35A-A68S, I35A-N217S, L10M-M123I, P9E-S258P, P9E-T210P, P9E-T77N, S86H-A202V, S86H-G165Q, S86H-N184Q, T77N-S155G, T77N-V147L, T78I-T210P, V87S-S155G, A202V-A242G, A242G-S258P, A68S-G165E, G127S-A242G, G165E-N184Q, G165E-N217S, M123I-S258E, N184Q-A242G, S86H-T115F, T3V-G99S, V147L-G165Q, A128R-G165Q, A128R-S258D, A68S-A128R, A68S-V147L, G127S-S258D, G165E-A202V, G165Q-S258E, G165S-T210P, G203E-T210P, G99S-G127S, G99S-M123I, G99S-T210P, L10M-A128P, L10M-A68S, L10M-N184Q, L10M-N217S, L10M-S258P, L10M-T77N, M123I-A242G, M123I-G165E, M123I-S187P, N239S-S258P, P9E-A128P, P9E-A202V, P9E-A68S, P9E-V45R, Q136R-N184Q, Q136R-N239S, S86H-N217S, T115F-S258P, T210P-A242G, T3V-G165E, T3V-L10M, T3V-P9E, T77N-A128R, T77N-G165E, T78I-A242G, V45R-G165Q, V87T-M123I, V87T-N184Q, V87T-N239S, V87T-Q136R, A68S-G101A, A68S-V71A, G101A-A128P, G101A-G165Q, G101A-S258P, G165A-N184Q, G165E-S258P, G99S-S258P, I35A-M123I, K15I-G165S, K15I-T210P, L10M-G127S, L10M-G165Q, Q136R-S187P, S187E-A202V, S187P-S258P, S250G-S258P, T115F-A202V, T115F-G165Q, T3V-G101A, T3V-I35A, T3V-V87S, T77N-G101A, T77N-G203E, T78I-S258D, V71A-G165Q, V71A-T77N, V87S-S258P, V87S-V147L, A128P-G165E, A128P-S155G, A128P-V147L, A128R-A202V, A128R-N184Q, A128R-N217S, A1Q-S86H, A68S-G99Q, A68S-S155G, G101A-A202V, G101A-G127S, G101A-N184Q, G101A-N217S, G165Q-G203E, G165Q-N239S, G165S-N239S, G99Q-A128P, G99Q-G165Q, G99Q-N184Q, G99Q-N217S, G99S-A242G, I35A-N184Q, I35A-T78I, I35A-V147L, L10M-T210P, L10M-T78I, M123I-G165S, N184Q-G203E, N184Q-S187P, N43A-M123I, P9E-G127S, P9E-S258E, Q136R-S258P, T115F-N217S, T3V-S86H, T77H-M123I, T77H-N184Q, T77H-N239S, T77H-Q136R, T77H-V87T, T77N-G99Q, T77N-S250G, T78I-G99S, T78I-V87S, V146S-G165S, V146S-N239S, V146S-T210P, V45R-S258E, V87S-A242G, V87S-N184Q, A128S-T210P, A143V-A202V, A143V-G165Q, A143V-S258P, A1Q-N184Q, A1Q-S258P, A68S-G203E, A68S-S187E, A68S-T114N, G101A-M123I, G101A-T210P, G165A-N217S, G165Q-S187P, G99Q-A202V, G99S-A202V, I35A-A242G, I35A-S258P, L10M-I35A, L10M-V87S, M123I-A128R, M123I-V146S, N184Q-S187E, N217S-L234W, N239S-S250G, N43A-Q136R, N43A-V45R, P9E-N239S, P9E-V87T, S155G-N217S, S187E-N217S, S86H-A143V, S86H-Q274A, S86H-S250G, T114N-A128P, T114N-G165Q, T114N-N184Q, T114N-N217S, T115F-A143V, T115F-N184Q, T210P-N239S, T210P-S250G, T3V-A128R, T3V-G99Q, T77H-G165Q, T77N-S187E, T77N-T114N, V147L-N217S, V45R-N184Q, V71A-N184Q, V71A-N217S, V71A-T210P, T77N-G165Q-N217S, A68S-T77N-G165Q, T77N-N184Q-N217S, A68S-T77N-N217S, T77N-A128P-G165Q, T77N-A128P-N217S, A68S-T77N-A128P, G165Q-N184Q-

N217S, T77N-G165Q-N184Q, A68S-A128P-G165Q, A68S-G165Q-N217S, A128P-N184Q-N217S,
 A128P-G165Q-N217S, A68S-A128P-N217S, A68S-N184Q-N217S, A68S-T77N-N184Q, T77N-A128P-
 N184Q, A68S-G165Q-N184Q, A128P-G165Q-N184Q, A68S-A128P-N184Q, T3V-N184Q-N217S,
 T3V-T77N-N217S, N184Q-A202V-N217S, T3V-T77N-N184Q, T3V-A128P-N217S, T3V-T77N-
 A128P, G165Q-A202V-N217S, T3V-A128P-N184Q, T77N-A202V-N217S, T3V-G165Q-N217S, T3V-
 T77N-G165Q, T3V-A68S-N217S, T3V-A68S-T77N, A68S-A202V-N217S, T3V-G165Q-N184Q,
 A128P-A202V-N217S, G165Q-N184Q-A202V, T77N-G165Q-A202V, T77N-N184Q-A202V, T3V-
 A128P-G165Q, T3V-A68S-A128P, T3V-A68S-N184Q, A128P-N184Q-A202V, A68S-N184Q-A202V,
 A68S-T77N-A202V, T3V-A68S-G165Q, T77N-A128P-A202V, A128P-G165Q-A202V, A68S-A128P-
 A202V, A68S-G165Q-A202V, M123I-G165Q-N217S, M123I-N184Q-N217S, T77N-M123I-G165Q,
 T77N-M123I-N217S, T3V-A202V-N217S, M123I-G165Q-N184Q, T77N-M123I-N184Q, T3V-N184Q-
 A202V, A68S-M123I-N217S, T3V-T77N-A202V, T3V-A128P-A202V, T3V-G165Q-A202V, M123I-
 A128P-N217S, T3V-A68S-A202V, A68S-M123I-N184Q, A68S-T77N-M123I, M123I-A128P-N184Q,
 T3V-M123I-N217S, T77N-M123I-A128P, A68S-M123I-G165Q, M123I-A128P-G165Q, A68S-M123I-
 A128P, N184Q-N217S-S258P, T3V-M123I-N184Q, T3V-M123I-G165Q, T3V-T77N-M123I, T77N-
 N217S-S258P, G165Q-N217S-S258P, T3V-A68S-M123I, T3V-M123I-A128P, T77N-N184Q-S258P,
 A128P-N184Q-S258P, A128P-N217S-S258P, T77N-A128P-S258P, T77N-G165Q-S258P, A202V-
 N217S-S258P, G165Q-N184Q-S258P, N184Q-A202V-S258P, A68S-A128P-S258P, A68S-N184Q-
 S258P, A68S-N217S-S258P, A68S-T77N-S258P, M123I-A202V-N217S, T3V-N217S-S258P, A128P-
 G165Q-S258P, G165Q-A202V-S258P, T3V-A128P-S258P, T3V-N184Q-S258P, A68S-G165Q-S258P,
 T3V-M123I-A202V, T3V-T77N-S258P, A128P-A202V-S258P, M123I-N184Q-A202V, T77N-A202V-
 S258P, A68S-A202V-S258P, M123I-G165Q-A202V, T77N-M123I-A202V, T3V-A68S-S258P, A68S-
 M123I-A202V, M123I-A128P-A202V, T3V-A202V-S258P, T3V-G165Q-S258P, A68S-G127S-A128P,
 A68S-T77N-G127S, T77N-G127S-A128P, A68S-G127S-A202V, A68S-G127S-N184Q, A68S-G127S-
 N217S, G127S-A128P-A202V, G127S-A128P-N184Q, G127S-A128P-N217S, A68S-G127S-G165Q,
 G127S-A128P-G165Q, G127S-A202V-N217S, G127S-N184Q-A202V, G127S-N184Q-N217S, M123I-
 N217S-S258P, T77N-G127S-A202V, T77N-G127S-N184Q, T77N-G127S-N217S, M123I-N184Q-
 S258P, T77N-G127S-G165Q, T3V-A68S-G127S, T3V-G127S-A128P, G127S-G165Q-A202V, G127S-
 G165Q-N184Q, G127S-G165Q-N217S, M123I-A202V-S258P, M123I-G165Q-S258P, T3V-M123I-
 S258P, T3V-T77N-G127S, A68S-G127S-S258P, G127S-A128P-S258P, A68S-M123I-S258P, M123I-
 A128P-S258P, T3V-G127S-A202V, T3V-G127S-N184Q, T3V-G127S-N217S, T77N-M123I-S258P,
 T3V-G127S-G165Q, T77N-G127S-S258P, G127S-A202V-S258P, G127S-N184Q-S258P, G127S-
 N217S-S258P, T3V-G127S-S258P, G127S-G165Q-S258P, A68S-M123I-G127S, M123I-G127S-A128P,
 M123I-G127S-A202V, M123I-G127S-N184Q, M123I-G127S-N217S, T77N-M123I-G127S, T3V-

M123I-G127S, M123I-G127S-G165Q, M123I-G127S-S258P, T77N-G165Q-T210P, A68S-T77N-T210P, A68S-G165Q-T210P, A68S-A128P-T210P, T77N-A128P-T210P, T77N-T210P-N217S, A68S-N184Q-T210P, N184Q-T210P-N217S, T77N-N184Q-T210P, A128P-G165Q-T210P, A68S-T210P-N217S, G165Q-N184Q-T210P, G165Q-T210P-N217S, A128P-N184Q-T210P, A128P-T210P-N217S, T77N-T210P-S258P, G165Q-T210P-S258P, T3V-A128P-T210P, T3V-A68S-T210P, T3V-T77N-T210P, A128P-T210P-S258P, A202V-T210P-N217S, A68S-T210P-S258P, N184Q-A202V-T210P, T210P-N217S-S258P, T3V-G165Q-T210P, T77N-A202V-T210P, A202V-T210P-S258P, G165Q-A202V-T210P, T3V-N184Q-T210P, A128P-A202V-T210P, A68S-A202V-T210P, N184Q-T210P-S258P, T3V-A202V-T210P, T3V-T210P-N217S, T3V-T210P-S258P, M123I-T210P-N217S, T77N-M123I-T210P, A68S-M123I-T210P, M123I-A128P-T210P, M123I-N184Q-T210P, A68S-G127S-T210P, G127S-A128P-T210P, T77N-G127S-T210P, G127S-T210P-S258P, M123I-G165Q-T210P, T3V-G127S-T210P, G127S-G165Q-T210P, T3V-M123I-T210P, G127S-A202V-T210P, G127S-N184Q-T210P, G127S-T210P-N217S, G127T-N184Q-N217S, M123I-A202V-T210P, M123I-T210P-S258P, A68S-G127T-N184Q, A68S-G127T-N217S, G127T-A128P-N184Q, G127T-A128P-N217S, T77N-V87S-G165Q, A68S-G127T-A128P, G127T-G165Q-N184Q, G127T-G165Q-N217S, T77N-G127T-N184Q, T77N-G127T-N217S, A68S-G165Q-S258D, A68S-T77N-G127T, A68S-T77N-S258D, G127T-A128P-G165Q, T77N-G127T-A128P, and T77N-G165Q-S258D; or (ii) N184Q-N217S, T77N-N217S, T77N-N184Q, T77N-G165Q, G165Q-N184Q, G165Q-N217S, A128P-N217S, A68S-T77N, T77N-A128P, A68S-N217S, A128P-N184Q, A68S-N184Q, A68S-G165Q, A128P-G165Q, A68S-A128P, A202V-N217S, N184Q-A202V, G165Q-A202V, T3V-N217S, T3V-N184Q, T3V-T77N, T77N-A202V, A68S-A202V, T3V-A128P, T3V-G165Q, A128P-A202V, T3V-A68S, T3V-A202V, N184Q-S258P, N217S-S258P, M123I-N217S, M123I-N184Q, T3V-M123I, T77N-S258P, A128P-S258P, G165Q-S258P, T77N-M123I, A202V-S258P, M123I-G165Q, A68S-M123I, A68S-S258P, M123I-A128P, M123I-A202V, T3V-S258P, A68S-G127S, G127S-A128P, T77N-G127S, G127S-A202V, G127S-N184Q, G127S-N217S, G127S-G165Q, M123I-S258P, T3V-G127S, G127S-S258P, M123I-G127S, T77N-T210P, A128P-T210P, A202V-T210P, A68S-T210P, G165Q-T210P, T210P-S258P, T3V-T210P, N184Q-T210P, T210P-N217S, G127S-T210P, M123I-T210P, T78I-A202V, A68S-S258D, G165Q-S258D, T77N-S258D, T77N-T78I, T78I-G165Q, G127T-N184Q, G127T-N217S, N184Q-S258D, N217S-S258D, A202V-S258D, A68S-T78I, T78I-A128P, T78I-G127S, A68S-G127T, T78I-N184Q, G127T-A128P, G127T-A202V, T3V-T78I, T78I-N217S, G127T-G165Q, L10M-A202V, T77N-G127T, T78I-S258P, T3V-G127T, T78I-M123I, V87S-A202V, A128P-A242G, A128P-S258D, G165Q-A242G, I35A-A202V, M123I-S258D, T3V-A242G, T3V-S258D, T77N-A242G, S86H-S258P, G127T-S258P, P9E-T77N, S86H-A202V, S86H-G165Q, S86H-N184Q, T78I-T210P, A202V-A242G, A242G-S258P, A68S-A242G, A68S-G165E, G127S-A242G, G165E-N184Q, G165E-N217S, I35A-V87S, L10M-M123I, M123I-

G127T, N184Q-A242G, N217S-A242G, A128R-G165Q, A128R-S258D, A68S-A128R, A68S-G99S, G127S-S258D, G165E-A202V, G99S-A128P, G99S-G127S, G99S-G165Q, I35A-G165Q, I35A-T77N, L10M-A128P, L10M-A68S, L10M-N184Q, L10M-N217S, L10M-S258P, L10M-T77N, M123I-A242G, P9E-A128P, P9E-A202V, P9E-A68S, P9E-G165Q, P9E-N184Q, P9E-N217S, P9E-S258P, S86H-T115F, T210P-A242G, T3V-G165E, T3V-G99S, T3V-L10M, T3V-P9E, T77N-A128R, T77N-G165E, T77N-G99S, T77N-V87S, T78I-A242G, V87S-A128P, V87S-G165Q, V87S-M123I, V87S-N217S, A68S-G101A, G101A-A128P, G101A-G165Q, G101A-S258P, G165E-S258P, G99S-S258P, I35A-M123I, I35A-N217S, L10M-G127S, L10M-G165Q, M123I-G165E, P9E-T210P, S187E-A202V, S86H-N217S, T115F-A202V, T115F-G165Q, T115F-S258P, T3V-G101A, T3V-V87S, T77N-G101A, T77N-G203E, T78I-S258D, A128P-G165E, A128R-A202V, A128R-N184Q, A128R-N217S, A68S-G99Q, G101A-A202V, G101A-G127S, G101A-N184Q, G101A-N217S, G165Q-G203E, G99Q-A128P, G99Q-G165Q, G99Q-N184Q, G99Q-N217S, G99S-A242G, G99S-T210P, I35A-A128P, I35A-T78I, L10M-T210P, L10M-T78I, P9E-G127S, P9E-M123I, T3V-I35A, T77N-G99Q, T78I-G99S, T78I-V87S, V87S-N184Q, A143V-A202V, A143V-G165Q, A143V-S258P, A1Q-N184Q, A1Q-S86H, A68S-G203E, A68S-S187E, A68S-V87S, G101A-M123I, G101A-T210P, G99Q-A202V, G99S-A202V, G99S-M123I, G99S-N184Q, G99S-N217S, I35A-N184Q, L10M-I35A, L10M-V87S, M123I-A128R, N184Q-G203E, N184Q-S187E, S187E-N217S, S250G-S258P, S86H-A143V, T115F-A143V, T115F-N184Q, T115F-N217S, T3V-A128R, T3V-G99Q, T3V-S86H, T77N-S187E, T77N-N184Q-N217S, A128P-N184Q-N217S, T77N-A128P-N217S, A68S-N184Q-N217S, A68S-T77N-N217S, G165Q-N184Q-N217S, T77N-A128P-N184Q, T77N-G165Q-N217S, T77N-G165Q-N184Q, A68S-T77N-G165Q, A68S-T77N-N184Q, T77N-A128P-G165Q, A128P-G165Q-N217S, A68S-A128P-N217S, A68S-T77N-A128P, A68S-G165Q-N217S, A128P-G165Q-N184Q, A68S-A128P-N184Q, A68S-A128P-G165Q, A68S-G165Q-N184Q, N184Q-A202V-N217S, G165Q-A202V-N217S, T77N-A202V-N217S, T3V-N184Q-N217S, A68S-A202V-N217S, T3V-T77N-N217S, A128P-A202V-N217S, G165Q-N184Q-A202V, T77N-N184Q-A202V, T3V-A128P-N217S, T3V-T77N-A128P, T3V-T77N-N184Q, T77N-G165Q-A202V, A128P-N184Q-A202V, A68S-N184Q-A202V, T3V-A128P-N184Q, A68S-T77N-A202V, T77N-A128P-A202V, T3V-G165Q-N217S, T3V-T77N-G165Q, T3V-A68S-N217S, T3V-A68S-T77N, A128P-G165Q-A202V, A68S-A128P-A202V, T3V-G165Q-N184Q, A68S-G165Q-A202V, T3V-A128P-G165Q, T3V-A68S-A128P, T3V-A68S-N184Q, T3V-A68S-G165Q, T3V-A202V-N217S, T3V-N184Q-A202V, T3V-T77N-A202V, T3V-A128P-A202V, T3V-G165Q-A202V, T3V-A68S-A202V, N184Q-N217S-S258P, T3V-M123I-N217S, M123I-N184Q-N217S, T77N-M123I-N217S, A128P-N184Q-S258P, A128P-N217S-S258P, A68S-M123I-N217S, M123I-G165Q-N217S, T77N-A128P-S258P, T77N-M123I-N184Q, T77N-N184Q-S258P, M123I-A128P-N217S, T3V-M123I-N184Q, T77N-N217S-S258P, A202V-N217S-S258P, A68S-M123I-N184Q, M123I-A128P-N184Q, N184Q-A202V-S258P, T3V-M123I-G165Q, T3V-

T77N-M123I, T77N-M123I-G165Q, A68S-A128P-S258P, A68S-T77N-M123I, M123I-G165Q-N184Q,
 T3V-A68S-M123I, A68S-N184Q-S258P, G165Q-N184Q-S258P, G165Q-N217S-S258P, M123I-A202V-
 N217S, T3V-M123I-A128P, T77N-M123I-A128P, A128P-G165Q-S258P, A68S-N217S-S258P, M123I-
 A128P-G165Q, T3V-N217S-S258P, T77N-G165Q-S258P, A68S-M123I-A128P, A68S-M123I-G165Q,
 A68S-T77N-S258P, T3V-A128P-S258P, T3V-N184Q-S258P, G165Q-A202V-S258P, T3V-M123I-
 A202V, A128P-A202V-S258P, M123I-N184Q-A202V, T3V-T77N-S258P, A68S-A202V-S258P, A68S-
 G165Q-S258P, M123I-G165Q-A202V, T77N-A202V-S258P, T77N-M123I-A202V, A68S-M123I-
 A202V, M123I-A128P-A202V, T3V-A202V-S258P, T3V-A68S-S258P, T3V-G165Q-S258P, A68S-
 G127S-A128P, A68S-T77N-G127S, T77N-G127S-A128P, A68S-G127S-A202V, A68S-G127S-N184Q,
 A68S-G127S-N217S, G127S-A128P-A202V, G127S-A128P-N184Q, G127S-A128P-N217S, A68S-
 G127S-G165Q, G127S-A128P-G165Q, G127S-A202V-N217S, G127S-N184Q-A202V, G127S-N184Q-
 N217S, T77N-G127S-A202V, T77N-G127S-N184Q, T77N-G127S-N217S, T77N-G127S-G165Q, T3V-
 A68S-G127S, T3V-G127S-A128P, G127S-G165Q-A202V, G127S-G165Q-N184Q, G127S-G165Q-
 N217S, M123I-A202V-S258P, M123I-N217S-S258P, M123I-N184Q-S258P, T3V-M123I-S258P, T3V-
 T77N-G127S, A68S-G127S-S258P, G127S-A128P-S258P, M123I-A128P-S258P, T3V-G127S-A202V,
 T3V-G127S-N184Q, T3V-G127S-N217S, A68S-M123I-S258P, T3V-G127S-G165Q, T77N-G127S-
 S258P, M123I-G165Q-S258P, T77N-M123I-S258P, G127S-A202V-S258P, G127S-N184Q-S258P,
 G127S-N217S-S258P, T3V-G127S-S258P, G127S-G165Q-S258P, A68S-M123I-G127S, M123I-G127S-
 A128P, M123I-G127S-A202V, M123I-G127S-N184Q, M123I-G127S-N217S, T77N-M123I-G127S,
 T3V-M123I-G127S, M123I-G127S-G165Q, M123I-G127S-S258P, A68S-A128P-T210P, A68S-T77N-
 T210P, T77N-A128P-T210P, N184Q-A202V-T210P, T3V-A128P-T210P, T3V-A68S-T210P, T3V-
 T77N-T210P, T77N-G165Q-T210P, T77N-T210P-S258P, A128P-G165Q-T210P, A128P-T210P-S258P,
 A202V-T210P-N217S, A68S-G165Q-T210P, A68S-T210P-S258P, N184Q-T210P-N217S, T3V-G165Q-
 T210P, T77N-A202V-T210P, A128P-A202V-T210P, A128P-N184Q-T210P, A128P-T210P-N217S,
 A202V-T210P-S258P, A68S-A202V-T210P, A68S-N184Q-T210P, A68S-T210P-N217S, G165Q-
 A202V-T210P, G165Q-T210P-S258P, T3V-A202V-T210P, T3V-N184Q-T210P, T3V-T210P-S258P,
 T77N-N184Q-T210P, T77N-T210P-N217S, G165Q-N184Q-T210P, N184Q-T210P-S258P, T210P-
 N217S-S258P, T3V-T210P-N217S, G165Q-T210P-N217S, A68S-G127S-T210P, G127S-A128P-T210P,
 T77N-G127S-T210P, G127S-T210P-S258P, T3V-G127S-T210P, G127S-G165Q-T210P, A68S-M123I-
 T210P, G127S-A202V-T210P, G127S-N184Q-T210P, G127S-T210P-N217S, M123I-A128P-T210P,
 M123I-A202V-T210P, M123I-N184Q-T210P, M123I-T210P-N217S, T3V-M123I-T210P, T77N-M123I-
 T210P, M123I-G165Q-T210P, M123I-T210P-S258P, A68S-G165Q-S258D, A68S-T77N-S258D, T77N-
 G165Q-S258D, T77N-T78I-G165Q, A68S-N184Q-S258D, A68S-N217S-S258D, G127T-N184Q-
 N217S, G165Q-N184Q-S258D, G165Q-N217S-S258D, M123I-G127S-T210P, T77N-N184Q-S258D,

T77N-N217S-S258D, A68S-T77N-T78I, A68S-T78I-A128P, A68S-T78I-G127S, A68S-T78I-G165Q, T77N-T78I-A128P, T77N-T78I-G127S, T78I-A128P-G165Q, T78I-G127S-A128P, T78I-G127S-G165Q, A68S-G127T-N184Q, A68S-G127T-N217S, N184Q-N217S-S258D, T77N-T78I-N184Q, T78I-G165Q-N184Q, A202V-N217S-S258D, A68S-A202V-S258D, G127T-A128P-N184Q, G127T-A128P-N217S, G127T-A202V-N217S, G127T-N184Q-A202V, G165Q-A202V-S258D, T3V-T77N-T78I, T3V-T78I-G165Q, T77N-A202V-S258D, T77N-T78I-A202V, T77N-T78I-N217S, T78I-G165Q-A202V, T78I-G165Q-N217S, T78I-N184Q-A202V, T78I-N184Q-N217S, A68S-G127T-A128P, A68S-T78I-A202V, A68S-T78I-N184Q, A68S-T78I-N217S, G127T-G165Q-N184Q, G127T-G165Q-N217S, T3V-A68S-T78I, T3V-T78I-A128P, T3V-T78I-G127S, T77N-G127T-N184Q, T77N-G127T-N217S, T77N-T78I-S258P, T78I-A128P-A202V, T78I-A128P-N184Q, T78I-A128P-N217S, T78I-A202V-N217S, T78I-G127S-A202V, T78I-G127S-N184Q, T78I-G127S-N217S, T78I-G165Q-S258P, A68S-G127T-A202V, A68S-T77N-G127T, G127T-A128P-G165Q, N184Q-A202V-S258D, T3V-G127T-N184Q, T3V-G127T-N217S, T77N-G127T-A128P, T78I-M123I-A202V, A128P-G165Q-A242G, A128P-G165Q-S258D, A128P-N184Q-S258D, A128P-N217S-S258D, A68S-A128P-S258D, A68S-G127T-G165Q, A68S-T78I-S258P, G127T-A128P-A202V, T3V-A128P-A242G, T3V-A68S-G127T, T3V-A68S-S258D, T3V-G127T-A202V, T3V-G165Q-A242G, T3V-G165Q-S258D, T3V-T77N-A242G, T3V-T77N-S258D, T3V-T78I-N184Q, T3V-T78I-N217S, T3V-T78I-S258P, T77N-A128P-A242G, T77N-A128P-S258D, T77N-G127T-G165Q, T77N-G165Q-A242G, T78I-A128P-S258P, T78I-G127S-S258P, A68S-M123I-S258D, G127T-G165Q-A202V, M123I-G165Q-S258D, T3V-G127T-A128P, T3V-N217S-S258D, T3V-T78I-A202V, T77N-G127T-A202V, T77N-M123I-S258D, T77N-T78I-M123I, T78I-M123I-G165Q, T78I-M123I-N184Q, T78I-N184Q-S258P, A68S-T78I-M123I, A68S-T78I-T210P, G127T-N184Q-S258P, G127T-N217S-S258P, M123I-A202V-S258D, M123I-N217S-S258D, T3V-A202V-S258D, T3V-M123I-S258D, T3V-N184Q-S258D, T3V-T77N-G127T, T3V-T78I-T210P, T77N-T78I-T210P, T78I-A128P-T210P, T78I-A202V-S258P, T78I-G127S-T210P, T78I-G165Q-T210P, T78I-M123I-A128P, T78I-M123I-G127S, T78I-M123I-N217S, and T78I-N217S-S258P; with the proviso that one or more of said two, three, or four or more substitutions is a man-made substitution; wherein said variant has at least 85% but less than 100% amino acid sequence identity to an AprL-Clade subtilisin enzyme comprising a Motif A and/or a Motif B; wherein Motif A is selected from Motif 1, Motif 2, Motif 3, and Motif 4; wherein Motif B is selected from Motif 5, Motif 6, Motif 7, Motif 8, Motif 9, Motif 10, Motif 11, Motif 12, Motif 13, and Motif 14; and wherein the amino acid positions of the variant are numbered by correspondence with the amino acid sequence of SEQ ID NO:2.

[0067] Another embodiment is directed to one or more subtilisin variant comprising an amino acid sequence comprising one or more amino acid substitution sets selected from: M123I-Q136R; P9E-M123I; V45R-M123I; G127T-G165S-T210E; M123I-G165S-N239S; M123I-Q136R-G165Q; M123I-Q136R-

S258E; M123I-V146S-T210E; P9E-V45R-M123I; Q136R-G165Q-S258E; V45R-M123I-Q136R;
 G127T-V146S-G165S-N239S; L10M-K27R-S98H-M123I; M123I-Q136R-G165Q-S258E; M123I-
 V146S-G165S-T210P; M123I-V146S-T210P-S250G; P9E-M123I-Q136R-G165Q; P9E-Q136R-G165Q-
 S258E; P9E-V45R-M123I-Q136R; V45R-M123I-Q136R-G165Q; V45R-M123I-Q136R-S258E; V45R-
 Q136R-G165Q-S258E; K27R-N43A-S98H-M123I-T210H; P9E-M123I-Q136R-G165Q-S258E; P9E-
 V45R-M123I-Q136R-G165Q; P9E-V45R-M123I-Q136R-S258E; V45R-M123I-Q136R-G165Q-S258E;
 Y103I-G127T-V146S-G165S-N239S; T77H-V87T-M123I-Q136R-N184Q-N239S; V26R-V45R-S98R-
 E111Q-M123I-Q136R; T77H-V87T-M123I-Q136R-G165Q-N184Q-N239S; T77S-V87S-M123I-Q136R-
 G165Q-S187P-S258P; T77S-V87S-M123I-Q136R-N160C-S187P-S258P; T77H-V87T-M123I-Q136R-
 N184Q-S187P-N239S-S258P; T3Q-K22Y-A24N-V45I-A48I-T77N-N217S-L234I-Y255R; P9E-T77Q-
 V87T-M123I-Q136R-G165Q-N184Q-N217S-N239S-S258E; N43A-V45R-T77H-V87T-M123I-Q136R-
 G165A-N184Q-S187P-N239S-S258P; P9E-N43A-V45R-T77S-V87S-M123I-Q136R-G165Q-S187P-
 N217S-S258P; P9E-N43A-V45R-T77H-V87T-M123I-Q136R-G165Q-N184Q-S187P-N239S-S258P;
 P9E-N43A-V45R-T77Q-V87T-M123I-Q136R-G165Q-N184Q-N217S-N239S-S258E; P9E-N43A-
 V45R-T77H-V87T-M123I-Q136R-G165A-N184Q-S187P-N217S-N239S-S258E; T77N-N217S-L234W;
 T77N-N217S, T77N-G126Q-G165Q-T210P-N217S-L234W-S258P, T77N-G127F-G165Q-A202V-
 T210P-N217S-L234W-S258P; A1Q-T3V-V28A-I35A-A68S-T77N-S86H-T115F-N217S-N239S-S250G-
 S258P-Q274A; T77N-M123I-T210P-N239S-S250G-S258P; V45I-A68S-T77N-M123I-L125N-G165Q-
 N217S-S258P; A68S-V71A-T77N-L95A-G165Q-N184Q-T210P-S258P; M123I-T210P-N217S-S258P;
 G127F-G165Q-N217S-S258P; T77N-G165Q-N217S-S258P; T77S-G165A-N184Q-T210P-N217S;
 A68S-T77N-A128P-G165Q-N184Q-N217S; V71A-T77S-N184Q-S258P; T77N-M123I-G165Q-N184Q-
 N217S; V71A-L95A-A128S-T210P; V71A-M123I-A128S-T210P; V71A-M123I-A128S-G165E; T77S-
 A128P-N184Q-N217S-F260W; T77H-A128S-G165Q-T210P; T77D-S86R-S155E-G165A-N184Q;
 A68S-V71A-T77N-A128S-G165Q-T210P-N217S; V28A-I35A-A68S-T77N-A128S-G165Q-N184Q-
 G203N-T210P; A128P-T210P; T210P-S258P; A128P-T210P-S259P; A128P-S187P-T210P; A128P-
 S187P-T210P-S258P; P9E-T77S-V87T-T210P-N239K; K15I-L95Q-M123I-G165S-T210P-N217S;
 K15I-M123I-V146S-G165S-T210P-N239S; K15I-N76K-M123I-V146S-G165S-T210P-N239S; K15I-
 V146S-G165S-T210P-N239S; P009E-V146S-N184Q-G203E-T210P-P238R-N239S; M123I-G165S-
 N217S; K15I-L95Q-S108K-G165S-G203E-T210P-N217S; K15I-L95Q-S108K-G165S-G203E-T210P;
 P9E-K15I-G165S-A193D-G203E-T210P; T77N-M123I-G165Q-N217S; T77H-G165Q; T77N-G165Q-
 N217S; A68S-V71A-T77N-G165Q-N217S; T77N-V87S-M123I-S155G-G165Q-N217S; T77H-V87S-
 S155G-G165Q; T77N-V87S-S155G-G165Q-N217S; T77N-M123I-G165Q; T77N-G165Q; A68S-V71A-
 T77N-G165Q; T77N-V87S-M123I-S155G-G165Q; I35A-A68S-T77N-V87S-V147L-G165Q; A68S-
 T77N-V87S-V147L-S155E-G165Q; T77N-V87S-M123I-V147L-S155G-G165Q; A68S-T77N-M123I-

G165Q-N184Q-N217S; T77N-M123I-A128P-G165Q-N184Q-N217S; A68S-V71A-T77N-S100N-A128P-G165Q-N184Q-T210P-N217S-N239T; T77N-A128P-G165Q-N184Q-N217S; A68S-A128P-G165Q-N184Q-N217S; A68S-T77N-G165Q-N184Q-N217S; A68S-T77N-A128P-N184Q-N217S; A68S-T77N-A128P-G165Q-N217S; A68S-T77N-A128P-G165Q-N184Q; M123I-G165Q-N184Q-N217S; T77N-G165Q-N184Q-N217S; T77N-M123I-N184Q-N217S; T77N-M123I-G165Q-N184Q; T3V-A68S-T77N-A128P-G165Q-N184Q-N217S; T3V-A68S-T77N-M123I-A128P-G165Q-N184Q-N217S; T3V-T77N-M123I-G165Q-N184Q-N217S; A68S-T77N-S108H-A128P-G165Q-N184Q-N217S; A68S-T77N-G99S-A128P-G165Q-N184Q-N217S; A68S-T77N-T114N-A128P-G165Q-N184Q-N217S; A68S-T77N-M123I-A128P-G165Q-N184Q-N217S; A68S-T77N-G127T-A128P-G165Q-N184Q-N217S; A68S-T77N-A128P-G165Q-N184Q-T210P-N217S; A68S-T77N-G99S-M123I-A128P-G165Q-N184Q-N217S; A68S-T77N-G99S-G127T-A128P-G165Q-N184Q-N217S; A68S-T77N-G99S-A128P-G165Q-N184Q-T210P-N217S; A68S-T77N-T114N-M123I-A128P-G165Q-N184Q-N217S; A68S-T77N-T114N-G127T-A128P-G165Q-N184Q-N217S; A68S-T77N-T114N-A128P-G165Q-N184Q-T210P-N217S; A68S-T77N-M123I-G127T-A128P-G165Q-N184Q-N217S; A68S-T77N-M123I-A128P-G165Q-N184Q-T210P-N217S; A68S-T77N-G127T-A128P-G165Q-N184Q-T210P-N217S; A68S-T77N-G99S-M123I-A128P-G165Q-N184Q-T210P-N217S; A68S-T77N-M123I-G127T-A128P-G165Q-N184Q-T210P-N217S; T3V-A68S-T77N-G99S-M123I-A128P-G165Q-N184Q-N217S; T3V-A68S-T77N-T114N-M123I-A128P-G165Q-N184Q-N217S; T3V-A68S-T77N-M123I-G127T-A128P-G165Q-N184Q-N217S; T3V-A68S-T77N-M123I-A128P-G165Q-N184Q-T210P-N217S; S86H-T115F-A143V-G165Q-A202V-S258P; S86H-L95A-N184Q-A202V-T210P-N217S-L234W-S250G-S258P; V30T-I35A-T77N-G165Q-A202V-T210P-S250G-S258P; A1Q-T3V-I35A-T77N-T78I-S86H-T115F-G165Q-N184Q-N217S-L234W-S258P; A1Q-A37T-T77N-T78I-S86H-M123I-G165Q-N184Q-A202V-N239S-S250G-S258P-Q274A; P9E-T77N-N96S-G165Q-G203E-S250G; P9E-T77N-N96S-G165E-G203E-T210P-S250G; A1Q-T3V-S86H-G130S-S155G-G165Q-N184Q-A202V-G203N-T210P-Q274A; A1Q-T3V-S86H-G165Q-N184Q-N239S-S250G-S258P; A1Q-T77N-S86H-V147L-A151G-N184Q-N217S-S258P; S86H-T115F-S155G-G165Q-N184Q-A202V-Q274A; A68S-S86H-T115F-N184Q-A202V-G203N-N239S-S250G-S258P-Q274A; A68S-T77N-V87S-A128P-S155G-G165Q-N217S; A68S-T77N-A128P-G165Q; A68S-T77N-V87S-A128P-S155G-G165Q; I35A-A68S-T77N-V87S-A128P-V147L-G165Q-N217S; A68S-T77N-V87S-A128P-S155G-G165Q-N217S-A242G; A68S-T77N-V87S-A128P-V147L-S155E-G165Q; A68S-T77N-V87S-A128P-S155E-G165Q-N217S; A68S-T77N-S86N-V87S-A128P-S155G-G165Q-N217S-A242G; I35A-A68S-T77N-A128P-V147L-G165Q-N217S; I35A-A68S-T77N-A128P-S155G-G165Q-A242G; I35A-A68S-T77N-A128P-V147L-G165Q-A242G; A68S-T77N-V87S-A128P-S155G-G165Q-A242G; A068S-T077N-N086S-G165Q-K194E-S211N-N217S; A068S-T077N-N086S-V087S-S155G-G165Q-K194E-S211N-N217S; A068S-T077N-N086S-G165Q-K194E-S211N; A068S-

T077N-N086S-V087S-S155G-G165Q-K194E-S211N; I035A-A068S-T077N-N086S-V087S-V147L-G165Q-K194E-S211N-N217S; A068S-T077N-N086S-V087S-S155G-G165Q-K194E-S211N-N217S-A242G; A068S-T077N-N086S-V087S-V147L-S155E-G165Q-K194E-S211N; A068S-T077N-N086S-V087S-S155E-G165Q-K194E-S211N-N217S; A068S-T077N-V087S-S155G-G165Q-K194E-S211N-N217S-A242G; I035A-A068S-T077N-N086S-V147L-G165Q-K194E-S211N-N217S; I035A-A068S-T077N-N086S-S155G-G165Q-K194E-S211N-A242G; I035A-A068S-T077N-N086S-V147L-G165Q-K194E-S211N-A242G; A068S-T077N-N086S-V087S-S155G-G165Q-K194E-S211N-A242G; A68S-T77N-G127T-A128P-G165Q-N184Q-A202V-N217S; T3V-A68S-T77N-M123I-A128P-G165Q-N184Q-A202V-N217S; A68S-T77N-G127T-A128P-G165Q-N184Q-N217S-S258P; T3V-A68S-T77N-M123I-A128P-G165Q-N184Q-N217S-S258P; A68S-T77N-G127T-A128P-G165Q-N184Q-G203Q-N217S; T3V-A68S-T77N-M123I-A128P-G165Q-N184Q-G203Q-N217S; A68S-T77N-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-M123I-A128P-G165Q-N184Q-A202V-N217S-S258P; A68S-T77N-A128P-G165Q-N184Q-G203Q-N217S-S258P; T3V-A68S-T77N-M123I-A128P-G165Q-N184Q-G203Q-N217S-S258P; S86H-T115F-G127T-A143V-G165Q-N184Q-A202V-N217S-S258P; T3V-S86H-T115F-M123I-A143V-G165Q-A202V-N217S-S258P; A68S-T77N-A128P-S129H-G165Q-N184Q-A202V-N217S; T3V-A68S-T77N-M123I-A128P-S129H-G165Q-N184Q-A202V-N217S; S86H-T115F-S129H-A143V-G165Q-N184Q-A202V-N217S-S258P; T3V-S86H-T115F-M123I-S129H-A143V-G165Q-A202V-N217S-S258P; A68S-T77N-A128P-S155N-G165Q-N184Q-A202V-N217S; T3V-A68S-T77N-M123I-A128P-S155N-G165Q-N184Q-A202V-N217S; I35A-A68S-T77N-V87S-A128P-V147L-G165Q-A202V-N217S; T3V-I35A-A68S-T77N-V87S-M123I-A128P-V147L-G165Q-A202V-N217S; A68S-T77N-A128P-Q136N-S155N-G165Q-N184Q-N217S-A242N; T3V-A68S-T77N-M123I-A128P-Q136N-S155N-G165Q-N184Q-N217S-A242N; A68S-V71A-T77N-A128P-G165Q-N184Q-A202V-N217S; T3V-A68S-V71A-T77N-M123I-A128P-G165Q-N184Q-A202V-N217S; A68S-T77N-A128P-G165Q-N184Q-S187P-A202V-N217S; T3V-A68S-T77N-M123I-A128P-G165Q-N184Q-S187P-A202V-N217S; A68S-T77N-A128R-G165Q-N184Q-A202V-N217S-S258D; T3V-A68S-T77N-M123I-A128R-G165Q-N184Q-A202V-N217S-S258D; A68S-T77N-A128P-S129R-G165Q-N184Q-N217S-S258D; T3V-A68S-T77N-M123I-A128P-S129R-G165Q-N184Q-N217S-S258D; A68S-T77N-A128R-G165Q-N184Q-A202V-N217S-S258D-K264H; T3V-A68S-T77N-M123I-A128R-G165Q-N184Q-A202V-N217S-S258D-K264H; A68S-T77N-A128R-G165Q-N184Q-G203E-S258D; T3V-A68S-T77N-M123I-A128R-G165Q-N184E-A202V-N217S-S258D; T3V-A52R-A68S-T77N-M123I-A128R-G165Q-N184E-A202V-N217S-S258D; T3V-T77N-S100N-G105N-D119Q-A128P-S158N-N160D-T161Q-G165A-N184Q-N217S-S258P-K264S; T3V-T77N-S100N-G105N-A128P-S158D-T161Q-G165A-N184Q-N217S-S258P; T3V-T77N-S100N-G105N-A128P-S158N-N160D-T161Q-G165A-N184Q-N217S-S258P; A24Q-V45I-A68S-T77N-S98R-

E111Q-M123I-S158Q-G165E-N184Q-S187E-V204I-T210I-N217S-S259E; P9E-A52R-S86R-E111Q-S129R-S158Q-G165E-V204I-T210I-S258D-S259E-K264Q; T3V-A68S-T77N-G127T-A128P-G165Q-N184Q-A202V-N217S; T3V-I35A-T77N-V87S-M123I-A128P-G165Q-N184Q-A202V-N217S-A242G; T3V-A68S-T77N-G127E-A128P-G165Q-N184Q-A202V-N217S; T3V-A68S-T77N-M123I-G127V-A128P-G165Q-N184Q-N217S; T3V-A68S-T77N-M123I-A128P-A151G-G165Q-N184Q-N217S; T3V-A68S-T77N-G127T-A128P-A151G-G165Q-N184Q-N217S; T3V-T77N-M123I-A128P-A151G-G165Q-N184Q-A202V-N217S; T3V-A68S-T77N-L95F-A128P-G165Q-N184Q-A202V-N217S; T3V-A68S-T77N-L95A-M123I-A128P-G165Q-N184Q-N217S; T3V-A68S-T77N-L95F-M123I-A128P-G165Q-N184Q-N217S; T3V-A68S-T77N-G99A-A128P-G165Q-N184Q-A202V-N217S; T3V-A68S-T77N-G99E-A128P-G165Q-N184Q-A202V-N217S; T3V-A68S-T77N-G99Q-A128P-G165Q-N184Q-A202V-N217S; T3V-A68S-T77N-G99A-M123I-A128P-G165Q-N184Q-N217S; T3V-A68S-T77N-S108H-A128P-G165Q-N184Q-N217S; T3V-I35A-T77N-V87S-M123I-A128P-V146A-G165Q-N184Q-A202V-N217S-A242G; T3V-A68S-T77N-G127T-A128P-G165E-N184Q-A202V-N217S; T3V-A68S-T77N-M123I-A128P-V146A-G165Q-N184Q-N217S; T3V-A68S-T77N-M123I-A128P-T161D-G165Q-N184Q-A202V-N217S; T3V-V45R-A68S-T77N-S102D-A128P-T161D-G165Q-N184Q-A202V-N217S; T3V-A68S-T77N-G127T-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-M123I-G127T-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-M123I-G127V-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-I35A-T77N-V87S-G127T-A128P-G165Q-N184Q-A202V-N217S-A242G-S258P; T3V-T77N-M123I-G127Q-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-T77N-M123I-G127V-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-T77N-M123I-G127E-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-M123I-G127T-A128P-G165Q-N184Q-A202V-N217S; T3V-A68S-T77N-M123I-G127T-A128P-G165Q-N184Q-A202V-T210P-N217S; T3V-A68S-T77N-M123I-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P; T3V-A68S-T77N-M123I-G127T-A128P-G165Q-N184Q-A202V-N217S-S258D; L10M-I35A-T78I-V87S-M123I-A202V-S258D-F260W; L10M-I35A-T78I-V87S-M123I-A202V-F260W-K264H; L10M-I35A-T78I-V87S-A202V-S258D; L10M-I35A-T78I-V87S-S187E-A202V; L10M-T78I-V87S-S187E-A202V; L10M-I35A-T78I-S187E-A202V; T3V-L10M-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-I35A-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-T78I-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-V87S-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-M123I-G127S-A128P-G165E-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P; T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-L216Q-N217S-S258P; T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-A242G-S258P; T3V-A68S-T77N-M123I-G127S-

A128P-G165Q-N184Q-A202V-N217S-S258D; T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P-F260W; A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-L31I-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; A68S-T77N-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; A68S-T77N-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P; A68S-T77N-G127T-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-T78I-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258D; A68S-T77N-T78I-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258D; T3V-A68S-T77N-T78I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; A68S-T77N-T78I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-G127S-A128P-G165Q-N184Q-A202V-N217S-S258D; A68S-T77N-G127S-A128P-G165Q-N184Q-A202V-N217S-S258D; T3V-A68S-T77N-T78I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258D; A68S-T77N-T78I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258D; T3V-A68S-T77N-S108H-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-G99Q-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-G99Q-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P; T3V-A68S-T77N-G127S-A128P-G165E-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P; T3V-P9E-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P; T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-S187E-A202V-T210P-N217S-S258P; T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-G203E-T210P-N217S-S258P; T3V-L10M-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P; T3V-P9E-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-S187E-A202V-N217S-S258P; T3V-P9E-A68S-T77N-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P; T3V-A68S-T77N-G127S-A128P-G165Q-N184Q-S187E-A202V-T210P-N217S-S258P; T3V-A68S-T77N-G127S-A128P-G165Q-N184Q-A202V-G203E-T210P-N217S-S258P; T3V-A68S-T77N-G101A-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P; T3V-L10M-A68S-T77N-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P; T3V-P9E-A68S-T77N-M123I-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P; T3V-A68S-T77N-M123I-A128P-G165Q-N184Q-S187E-A202V-T210P-N217S-S258P; T3V-A68S-T77N-M123I-A128P-G165Q-N184Q-A202V-G203E-T210P-N217S-S258P; T3V-A68S-T77N-G101A-M123I-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P; T3V-L10M-A68S-T77N-M123I-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P; T3V-L10M-A68S-T77N-G101A-M123I-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P; T3V-P9E-L10M-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P; T3V-P9E-A68S-T77N-G99Q-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P; T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-S243I-S258P; T3V-A68S-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P;

T3V-A68S-T77N-M123I-G127S-A128P-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-M123I-G127S-A128P-G165Q-A202V-N217S-S258P; T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-S258P; T3V-A68S-T77E-M123I-G127T-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77E-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P-K264H; T3V-A68S-T77N-S102T-M123I-G127S-A128P-G165Q-N184Q-A202V-N211S-N217S-S258P; T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N211S-N217S-S258P; T3V-A68S-T77N-S102T-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P-S259P; T3V-A68S-T77N-M123I-G127S-A128P-T161Q-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-M123I-G127S-A128P-S158Q-G165Q-N184Q-A202V-N217S-S258P; T3V-V45Q-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-M123I-G127S-A128P-A143Q-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-G117S-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A48V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A48T-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A29S-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-G127S-A128P-G165Q-N184Q-A202V-N217S; T3V-A68S-T77N-G127S-A128P-G165Q-N184Q-N217S-S258P; T3V-A68S-T77N-A128P-G165Q-N184Q-A202V-N217S-S258P; A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S; A68S-T77N-M123I-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-G99Q-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-T78I-V87S-M123I-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P; T3V-A68S-T77N-T78I-M123I-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-A242G-S258P; T3V-V26Q-A68S-T77N-M123I-G127S-A128P-G165E-N184Q-A202V-T210P-N217S-S258P; T3V-P9E-L10M-A68S-T77N-M123I-G127S-A128P-G165E-N184Q-A202V-T210P-N217S-S258P; T3V-P9E-L10M-A68S-T77N-G101A-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-T78I-M123I-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P; T3V-A68S-T77N-T78I-M123I-G127S-A128P-N160K-G165Q-N184Q-A202V-T210P-N217S-A242G-S258P; T3V-A68S-T77N-V87S-M123I-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P; T3V-P40E-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77E-M123I-G127T-A128P-G165E-N184Q-A202V-N217S-P238Q-S258P; T3V-A68S-T77N-G99S-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77E-M123I-G127T-G165E-N184Q-A202V-N217S-S258P; A68S-T77N-A128P-G165Q-N184Q-N217S-S258P; T3V-A68S-T77N-G101A-S102T-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S; T3V-A68S-T77N-T78I-G99S-G127S-A128P-G165Q-T210P-A242G-S258P-

K264H; T3V-A68S-T77N-T78I-G99S-M123I-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-A242G-S258P; T3V-A68S-T77N-M123I-G127T-G165E-N184Q-A202V-N217S-P238Q-S258P; T3V-A68S-T77N-T78I-G99S-G127S-A128P-G165Q-T210P-A242G-S258P; T3V-A68S-T77N-T78I-G99S-M123I-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-A242G-S258D; I35A-M123I-G127T-T210P-S243I; T3V-A68S-T77N-G99S-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P-F260W; T3V-A68S-T77N-T78I-G99S-G101A-G127S-A128P-G165Q-T210P-A242G-S258P; A68S-T77N-G99Q-A128P-G165Q-N184Q-N217S; A68S-T77N-A128P-G165Q-N184Q-A202V-N217S; A68S-T77N-A128P-G165Q-N184Q-N217S-S258D; T3V-A68S-T77N-T78I-G99S-G101A-M123I-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-A242G-S258P; and V45R-S98R; with the proviso that one or more of said two, three, or four or more substitutions is a man-made substitution; wherein said variant has at least 85% but less than 100% amino acid sequence identity to an AprL-Clade subtilisin enzyme comprising a Motif A and/or a Motif B; wherein Motif A is selected from Motif 1, Motif 2, Motif 3, and Motif 4; wherein Motif B is selected from Motif 5, Motif 6, Motif 7, Motif 8, Motif 9, Motif 10, Motif 11, Motif 12, Motif 13, and Motif 14; and wherein the amino acid positions of the variant are numbered by correspondence with the amino acid sequence of SEQ ID NO:2.

[0068] Another embodiment provides one or more subtilisin variant comprising an amino acid sequence comprising one or more amino acid substitution sets selected from: M123I-Q136R; P9E-M123I; V45R-M123I; G127T-G165S-T210E; M123I-G165S-N239S; M123I-Q136R-G165Q; M123I-Q136R-S258E; M123I-V146S-T210E; P9E-V45R-M123I; Q136R-G165Q-S258E; V45R-M123I-Q136R; G127T-V146S-G165S-N239S; L10M-K27R-S98H-M123I; M123I-Q136R-G165Q-S258E; M123I-V146S-G165S-T210P; M123I-V146S-T210P-S250G; P9E-M123I-Q136R-G165Q; P9E-Q136R-G165Q-S258E; P9E-V45R-M123I-Q136R; V45R-M123I-Q136R-G165Q; V45R-M123I-Q136R-S258E; V45R-Q136R-G165Q-S258E; K27R-N43A-S98H-M123I-T210H; P9E-M123I-Q136R-G165Q-S258E; P9E-V45R-M123I-Q136R-G165Q; P9E-V45R-M123I-Q136R-S258E; V45R-M123I-Q136R-G165Q-S258E; Y103I-G127T-V146S-G165S-N239S; T77H-V87T-M123I-Q136R-N184Q-N239S; V26R-V45R-S98R-E111Q-M123I-Q136R; T77H-V87T-M123I-Q136R-G165Q-N184Q-N239S; T77S-V87S-M123I-Q136R-G165Q-S187P-S258P; T77S-V87S-M123I-Q136R-N160C-S187P-S258P; T77H-V87T-M123I-Q136R-N184Q-S187P-N239S-S258P; T3Q-K22Y-A24N-V45I-A48I-T77N-N217S-L234I-Y255R; P9E-T77Q-V87T-M123I-Q136R-G165Q-N184Q-N217S-N239S-S258E; N43A-V45R-T77H-V87T-M123I-Q136R-G165A-N184Q-S187P-N239S-S258P; P9E-N43A-V45R-T77S-V87S-M123I-Q136R-G165Q-S187P-N217S-S258P; P9E-N43A-V45R-T77H-V87T-M123I-Q136R-G165Q-N184Q-S187P-N239S-S258P; P9E-N43A-V45R-T77Q-V87T-M123I-Q136R-G165Q-N184Q-N217S-N239S-S258E; P9E-N43A-V45R-T77H-V87T-M123I-Q136R-G165A-N184Q-S187P-N217S-N239S-S258E; T77N-N217S-L234W; T77N-N217S, T77N-G126Q-G165Q-T210P-N217S-L234W-S258P, T77N-G127F-G165Q-A202V-

T210P-N217S-L234W-S258P; A1Q-T3V-V28A-I35A-A68S-T77N-S86H-T115F-N217S-N239S-S250G-S258P-Q274A; T77N-M123I-T210P-N239S-S250G-S258P; V45I-A68S-T77N-M123I-L125N-G165Q-N217S-S258P; A68S-V71A-T77N-L95A-G165Q-N184Q-T210P-S258P; M123I-T210P-N217S-S258P; G127F-G165Q-N217S-S258P; T77N-G165Q-N217S-S258P; T77S-G165A-N184Q-T210P-N217S; A68S-T77N-A128P-G165Q-N184Q-N217S; V71A-T77S-N184Q-S258P; T77N-M123I-G165Q-N184Q-N217S; V71A-L95A-A128S-T210P; V71A-M123I-A128S-T210P; V71A-M123I-A128S-G165E; T77S-A128P-N184Q-N217S-F260W; T77H-A128S-G165Q-T210P; T77D-S86R-S155E-G165A-N184Q; A68S-V71A-T77N-A128S-G165Q-T210P-N217S; V28A-I35A-A68S-T77N-A128S-G165Q-N184Q-G203N-T210P; A128P-T210P; T210P-S258P; A128P-T210P-S259P; A128P-S187P-T210P; A128P-S187P-T210P-S258P; P9E-T77S-V87T-T210P-N239K; K15I-L95Q-M123I-G165S-T210P-N217S; K15I-M123I-V146S-G165S-T210P-N239S; K15I-N76K-M123I-V146S-G165S-T210P-N239S; K15I-V146S-G165S-T210P-N239S; P009E-V146S-N184Q-G203E-T210P-P238R-N239S; M123I-G165S-N217S; K15I-L95Q-S108K-G165S-G203E-T210P-N217S; K15I-L95Q-S108K-G165S-G203E-T210P; P9E-K15I-G165S-A193D-G203E-T210P; T77N-M123I-G165Q-N217S; T77H-G165Q; T77N-G165Q-N217S; A68S-V71A-T77N-G165Q-N217S; T77N-V87S-M123I-S155G-G165Q-N217S; T77H-V87S-S155G-G165Q; T77N-V87S-S155G-G165Q-N217S; T77N-M123I-G165Q; T77N-G165Q; A68S-V71A-T77N-G165Q; T77N-V87S-M123I-S155G-G165Q; I35A-A68S-T77N-V87S-V147L-G165Q; A68S-T77N-V87S-V147L-S155E-G165Q; T77N-V87S-M123I-V147L-S155G-G165Q; A68S-T77N-M123I-G165Q-N184Q-N217S; T77N-M123I-A128P-G165Q-N184Q-N217S; A68S-V71A-T77N-S100N-A128P-G165Q-N184Q-T210P-N217S-N239T; T77N-A128P-G165Q-N184Q-N217S; A68S-A128P-G165Q-N184Q-N217S; A68S-T77N-G165Q-N184Q-N217S; A68S-T77N-A128P-N184Q-N217S; A68S-T77N-A128P-G165Q-N217S; A68S-T77N-A128P-G165Q-N184Q; M123I-G165Q-N184Q-N217S; T77N-G165Q-N184Q-N217S; T77N-M123I-N184Q-N217S; T77N-M123I-G165Q-N184Q; T3V-A68S-T77N-A128P-G165Q-N184Q-N217S; T3V-A68S-T77N-M123I-A128P-G165Q-N184Q-N217S; T3V-T77N-M123I-G165Q-N184Q-N217S; A68S-T77N-S108H-A128P-G165Q-N184Q-N217S; A68S-T77N-G99S-A128P-G165Q-N184Q-N217S; A68S-T77N-T114N-A128P-G165Q-N184Q-N217S; A68S-T77N-M123I-A128P-G165Q-N184Q-N217S; A68S-T77N-G127T-A128P-G165Q-N184Q-N217S; A68S-T77N-A128P-G165Q-N184Q-T210P-N217S; A68S-T77N-G99S-M123I-A128P-G165Q-N184Q-N217S; A68S-T77N-G99S-G127T-A128P-G165Q-N184Q-N217S; A68S-T77N-G99S-A128P-G165Q-N184Q-T210P-N217S; A68S-T77N-T114N-M123I-A128P-G165Q-N184Q-N217S; A68S-T77N-T114N-G127T-A128P-G165Q-N184Q-N217S; A68S-T77N-T114N-A128P-G165Q-N184Q-T210P-N217S; A68S-T77N-M123I-G127T-A128P-G165Q-N184Q-N217S; A68S-T77N-M123I-A128P-G165Q-N184Q-T210P-N217S; A68S-T77N-G127T-A128P-G165Q-N184Q-T210P-N217S; A68S-T77N-G99S-M123I-A128P-G165Q-N184Q-T210P-N217S; A68S-T77N-M123I-G127T-A128P-G165Q-N184Q-T210P-

N217S; T3V-A68S-T77N-G99S-M123I-A128P-G165Q-N184Q-N217S; T3V-A68S-T77N-T114N-M123I-A128P-G165Q-N184Q-N217S; T3V-A68S-T77N-M123I-G127T-A128P-G165Q-N184Q-N217S; T3V-A68S-T77N-M123I-A128P-G165Q-N184Q-T210P-N217S; S86H-T115F-A143V-G165Q-A202V-S258P; S86H-L95A-N184Q-A202V-T210P-N217S-L234W-S250G-S258P; V30T-I35A-T77N-G165Q-A202V-T210P-S250G-S258P; A1Q-T3V-I35A-T77N-T78I-S86H-T115F-G165Q-N184Q-N217S-L234W-S258P; A1Q-A37T-T77N-T78I-S86H-M123I-G165Q-N184Q-A202V-N239S-S250G-S258P-Q274A; P9E-T77N-N96S-G165Q-G203E-S250G; P9E-T77N-N96S-G165E-G203E-T210P-S250G; A1Q-T3V-S86H-G130S-S155G-G165Q-N184Q-A202V-G203N-T210P-Q274A; A1Q-T3V-S86H-G165Q-N184Q-N239S-S250G-S258P; A1Q-T77N-S86H-V147L-A151G-N184Q-N217S-S258P; S86H-T115F-S155G-G165Q-N184Q-A202V-Q274A; A68S-S86H-T115F-N184Q-A202V-G203N-N239S-S250G-S258P-Q274A; A68S-T77N-V87S-A128P-S155G-G165Q-N217S; A68S-T77N-A128P-G165Q; A68S-T77N-V87S-A128P-S155G-G165Q; I35A-A68S-T77N-V87S-A128P-V147L-G165Q-N217S; A68S-T77N-V87S-A128P-S155G-G165Q-N217S-A242G; A68S-T77N-V87S-A128P-V147L-S155E-G165Q; A68S-T77N-V87S-A128P-S155E-G165Q-N217S; A68S-T77N-S86N-V87S-A128P-S155G-G165Q-N217S-A242G; I35A-A68S-T77N-A128P-V147L-G165Q-N217S; I35A-A68S-T77N-A128P-S155G-G165Q-A242G; I35A-A68S-T77N-A128P-V147L-G165Q-A242G; A68S-T77N-V87S-A128P-S155G-G165Q-A242G; A068S-T077N-N086S-G165Q-K194E-S211N-N217S; A068S-T077N-N086S-V087S-S155G-G165Q-K194E-S211N-N217S; A068S-T077N-N086S-G165Q-K194E-S211N; A068S-T077N-N086S-V087S-S155G-G165Q-K194E-S211N; I035A-A068S-T077N-N086S-V087S-V147L-G165Q-K194E-S211N-N217S; A068S-T077N-N086S-V087S-S155G-G165Q-K194E-S211N-N217S-A242G; A068S-T077N-N086S-V087S-V147L-S155E-G165Q-K194E-S211N; A068S-T077N-N086S-V087S-S155E-G165Q-K194E-S211N-N217S; A068S-T077N-V087S-S155G-G165Q-K194E-S211N-N217S-A242G; I035A-A068S-T077N-N086S-V147L-G165Q-K194E-S211N-N217S; I035A-A068S-T077N-N086S-S155G-G165Q-K194E-S211N-A242G; OI35A-A068S-T077N-N086S-V147L-G165Q-K194E-S211N-A242G; and A068S-T077N-N086S-V087S-S155G-G165Q-K194E-S211N-A242G; with the proviso that one or more of said two, three, or four or more variations is a non-naturally occurring amino acid; wherein said variant has 85% or more amino acid sequence identity to an AprL-Clade subtilisin enzyme comprising a Motif A and/or a Motif B; wherein Motif A is selected from Motif 1, Motif 2, Motif 3, and Motif 4; wherein Motif B is selected from Motif 5, Motif 6, Motif 7, Motif 8, Motif 9, Motif 10, Motif 11, Motif 12, Motif 13, and Motif 14; and wherein the amino acid positions of the variant are numbered by correspondence with the amino acid sequence of SEQ ID NO:2.

[0069] Yet an even still further embodiment is directed to one or more subtilisin variant comprising an amino acid sequence comprising one or more amino acid substitution sets selected from: A68S-T77N-G127T-A128P-G165Q-N184Q-A202V-N217S; T3V-A68S-T77N-M123I-A128P-G165Q-N184Q-

A202V-N217S; A68S-T77N-G127T-A128P-G165Q-N184Q-N217S-S258P; T3V-A68S-T77N-M123I-A128P-G165Q-N184Q-N217S-S258P; A68S-T77N-G127T-A128P-G165Q-N184Q-G203Q-N217S; T3V-A68S-T77N-M123I-A128P-G165Q-N184Q-G203Q-N217S; A68S-T77N-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-M123I-A128P-G165Q-N184Q-A202V-N217S-S258P; A68S-T77N-A128P-G165Q-N184Q-G203Q-N217S-S258P; T3V-A68S-T77N-M123I-A128P-G165Q-N184Q-G203Q-N217S-S258P; S86H-T115F-G127T-A143V-G165Q-N184Q-A202V-N217S-S258P; T3V-S86H-T115F-M123I-A143V-G165Q-A202V-N217S-S258P; A68S-T77N-A128P-S129H-G165Q-N184Q-A202V-N217S; T3V-A68S-T77N-M123I-A128P-S129H-G165Q-N184Q-A202V-N217S; S86H-T115F-S129H-A143V-G165Q-N184Q-A202V-N217S-S258P; T3V-S86H-T115F-M123I-S129H-A143V-G165Q-A202V-N217S-S258P; A68S-T77N-A128P-S155N-G165Q-N184Q-A202V-N217S; T3V-A68S-T77N-M123I-A128P-S155N-G165Q-N184Q-A202V-N217S; I35A-A68S-T77N-V87S-A128P-V147L-G165Q-A202V-N217S; T3V-I35A-A68S-T77N-V87S-M123I-A128P-V147L-G165Q-A202V-N217S; A68S-T77N-A128P-Q136N-S155N-G165Q-N184Q-N217S-A242N; T3V-A68S-T77N-M123I-A128P-Q136N-S155N-G165Q-N184Q-N217S-A242N; A68S-V71A-T77N-A128P-G165Q-N184Q-A202V-N217S; T3V-A68S-V71A-T77N-M123I-A128P-G165Q-N184Q-A202V-N217S; A68S-T77N-A128P-G165Q-N184Q-S187P-A202V-N217S; T3V-A68S-T77N-M123I-A128P-G165Q-N184Q-S187P-A202V-N217S; A68S-T77N-A128R-G165Q-N184Q-A202V-N217S-S258D; T3V-A68S-T77N-M123I-A128R-G165Q-N184Q-A202V-N217S-S258D; A68S-T77N-A128P-S129R-G165Q-N184Q-N217S-S258D; T3V-A68S-T77N-M123I-A128P-S129R-G165Q-N184Q-N217S-S258D; A68S-T77N-A128R-G165Q-N184Q-A202V-N217S-S258D-K264H; T3V-A68S-T77N-M123I-A128R-G165Q-N184Q-A202V-N217S-S258D-K264H; A68S-T77N-A128R-G165Q-N184Q-G203E-S258D; T3V-A68S-T77N-M123I-A128R-G165Q-N184Q-G203E-S258D; T3V-A68S-T77N-M123I-A128R-G165Q-N184E-A202V-N217S-S258D; T3V-A52R-A68S-T77N-M123I-A128R-G165Q-N184E-A202V-N217S-S258D; T3V-T77N-S100N-G105N-D119Q-A128P-S158N-N160D-T161Q-G165A-N184Q-N217S-S258P-K264S; T3V-T77N-S100N-G105N-A128P-S158D-T161Q-G165A-N184Q-N217S-S258P; T3V-T77N-S100N-G105N-A128P-S158N-N160D-T161Q-G165A-N184Q-N217S-S258P; A24Q-V45I-A68S-T77N-S98R-E111Q-M123I-S158Q-G165E-N184Q-S187E-V204I-T210I-N217S-S259E; P9E-A52R-S86R-E111Q-S129R-S158Q-G165E-V204I-T210I-S258D-S259E-K264Q; T3V-A68S-T77N-G127T-A128P-G165Q-N184Q-A202V-N217S; T3V-I35A-T77N-V87S-M123I-A128P-G165Q-N184Q-A202V-N217S-A242G; T3V-A68S-T77N-G127E-A128P-G165Q-N184Q-A202V-N217S; T3V-A68S-T77N-M123I-G127V-A128P-G165Q-N184Q-N217S; T3V-A68S-T77N-M123I-A128P-A151G-G165Q-N184Q-N217S; T3V-A68S-T77N-G127T-A128P-A151G-G165Q-N184Q-N217S; T3V-T77N-M123I-A128P-A151G-G165Q-N184Q-A202V-N217S; T3V-A68S-T77N-L95F-A128P-G165Q-N184Q-A202V-N217S; T3V-A68S-T77N-L95A-M123I-A128P-G165Q-N184Q-N217S; T3V-A68S-T77N-L95F-M123I-A128P-G165Q-

N184Q-N217S; T3V-A68S-T77N-G99A-A128P-G165Q-N184Q-A202V-N217S; T3V-A68S-T77N-G99E-A128P-G165Q-N184Q-A202V-N217S; T3V-A68S-T77N-G99Q-A128P-G165Q-N184Q-A202V-N217S; T3V-A68S-T77N-G99A-M123I-A128P-G165Q-N184Q-N217S; T3V-A68S-T77N-S108H-A128P-G165Q-N184Q-N217S; T3V-I35A-T77N-V87S-M123I-A128P-V146A-G165Q-N184Q-A202V-N217S-A242G; T3V-A68S-T77N-G127T-A128P-G165E-N184Q-A202V-N217S; T3V-A68S-T77N-M123I-A128P-V146A-G165Q-N184Q-N217S; T3V-A68S-T77N-M123I-A128P-T161D-G165Q-N184Q-A202V-N217S; T3V-V45R-A68S-T77N-S102D-A128P-T161D-G165Q-N184Q-A202V-N217S; T3V-A68S-T77N-G127T-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-M123I-G127T-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-M123I-G127V-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-I35A-T77N-V87S-G127T-A128P-G165Q-N184Q-A202V-N217S-A242G-S258P; T3V-T77N-M123I-G127Q-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-T77N-M123I-G127V-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-T77N-M123I-G127E-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-M123I-G127T-A128P-G165Q-N184Q-A202V-N217S; T3V-A68S-T77N-M123I-G127T-A128P-G165Q-N184Q-A202V-T210P-N217S; T3V-A68S-T77N-M123I-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P; T3V-A68S-T77N-M123I-G127T-A128P-G165Q-N184Q-A202V-N217S-S258D; L10M-I35A-T78I-V87S-M123I-A202V-S258D-F260W; L10M-I35A-T78I-V87S-M123I-A202V-F260W-K264H; L10M-I35A-T78I-V87S-A202V-S258D; L10M-I35A-T78I-V87S-S187E-A202V; L10M-T78I-V87S-S187E-A202V; L10M-I35A-T78I-S187E-A202V; T3V-L10M-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-I35A-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-T78I-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-V87S-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-M123I-G127S-A128P-G165E-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P; T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-L216Q-N217S-S258P; T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-A242G-S258P; T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258D; T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P-F260W; A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-L31I-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; A68S-T77N-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; A68S-T77N-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P; A68S-T77N-G127T-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-T78I-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258D; A68S-T77N-T78I-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258D; T3V-A68S-T77N-T78I-G127S-A128P-G165Q-N184Q-

A202V-N217S-S258P; A68S-T77N-T78I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-G127S-A128P-G165Q-N184Q-A202V-N217S-S258D; A68S-T77N-G127S-A128P-G165Q-N184Q-A202V-N217S-S258D; T3V-A68S-T77N-T78I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258D; A68S-T77N-T78I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258D; T3V-A68S-T77N-S108H-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-G99Q-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-G99Q-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P; T3V-A68S-T77N-G127S-A128P-G165E-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P; T3V-P9E-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P; T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-S187E-A202V-T210P-N217S-S258P; T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-G203E-T210P-N217S-S258P; T3V-L10M-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P; T3V-P9E-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-S187E-A202V-N217S-S258P; T3V-P9E-A68S-T77N-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P; T3V-A68S-T77N-G127S-A128P-G165Q-N184Q-S187E-A202V-T210P-N217S-S258P; T3V-A68S-T77N-G127S-A128P-G165Q-N184Q-A202V-G203E-T210P-N217S-S258P; T3V-A68S-T77N-G101A-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P; T3V-L10M-A68S-T77N-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P; T3V-P9E-A68S-T77N-M123I-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P; T3V-A68S-T77N-M123I-A128P-G165Q-N184Q-S187E-A202V-T210P-N217S-S258P; T3V-A68S-T77N-M123I-A128P-G165Q-N184Q-A202V-G203E-T210P-N217S-S258P; T3V-A68S-T77N-G101A-M123I-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P; T3V-L10M-A68S-T77N-M123I-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P; T3V-L10M-A68S-T77N-G101A-M123I-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P; T3V-P9E-L10M-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P; T3V-P9E-A68S-T77N-G99Q-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P; T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-S243I-S258P; T3V-A68S-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-M123I-G127S-A128P-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-M123I-G127S-A128P-G165Q-A202V-N217S-S258P; T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-S258P; T3V-A68S-T77E-M123I-G127T-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77E-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P-K264H; T3V-A68S-T77N-S102T-M123I-G127S-A128P-G165Q-N184Q-A202V-N211S-N217S-S258P; T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N211S-N217S-S258P; T3V-A68S-T77N-S102T-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P;

S259P; T3V-A68S-T77N-M123I-G127S-A128P-T161Q-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-M123I-G127S-A128P-S158Q-G165Q-N184Q-A202V-N217S-S258P; T3V-V45Q-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A193Q-A202V-N217S-S258P; T3V-A68S-T77N-M123I-G127S-A128P-A143Q-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-G117S-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A48V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A48T-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A29S-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-G127S-A128P-G165Q-N184Q-A202V-N217S; T3V-A68S-T77N-G127S-A128P-G165Q-N184Q-N217S-S258P; T3V-A68S-T77N-A128P-G165Q-N184Q-A202V-N217S-S258P; A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S; A68S-T77N-M123I-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-G99Q-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-T78I-V87S-M123I-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P; T3V-A68S-T77N-T78I-M123I-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-A242G-S258P; T3V-V26Q-A68S-T77N-M123I-G127S-A128P-G165E-N184Q-A202V-T210P-N217S-S258P; T3V-P9E-L10M-A68S-T77N-M123I-G127S-A128P-G165E-N184Q-A202V-T210P-N217S-S258P; T3V-P9E-L10M-A68S-T77N-G101A-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-T78I-M123I-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P; T3V-A68S-T77N-T78I-M123I-G127S-A128P-N160K-G165Q-N184Q-A202V-T210P-N217S-A242G-S258P; T3V-A68S-T77N-V87S-M123I-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P; T3V-P40E-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77E-M123I-G127T-A128P-G165E-N184Q-A202V-N217S-P238Q-S258P; T3V-A68S-T77N-G99S-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77E-M123I-G127T-G165E-N184Q-A202V-N217S-S258P; A68S-T77N-A128P-G165Q-N184Q-N217S-S258P; T3V-A68S-T77N-G101A-S102T-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S; T3V-A68S-T77N-T78I-G99S-G127S-A128P-G165Q-T210P-A242G-S258P-K264H; T3V-A68S-T77N-T78I-G99S-M123I-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-A242G-S258P; T3V-A68S-T77N-M123I-G127T-G165E-N184Q-A202V-N217S-P238Q-S258P; T3V-A68S-T77N-T78I-G99S-G127S-A128P-G165Q-T210P-A242G-S258P; T3V-A68S-T77N-T78I-G99S-M123I-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-A242G-S258D; I35A-M123I-G127T-T210P-S243I; T3V-A68S-T77N-G99S-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P-F260W; T3V-A68S-T77N-T78I-G99S-G101A-G127S-A128P-G165Q-T210P-A242G-S258P; A68S-T77N-G99Q-A128P-G165Q-N184Q-N217S; A68S-T77N-A128P-G165Q-N184Q-A202V-N217S; A68S-T77N-A128P-G165Q-N184Q-N217S-S258D; T3V-A68S-T77N-T78I-G99S-G101A-M123I-

G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-A242G-S258P; and V45R-S98R; with the proviso that one or more of said two, three, or four or more substitutions is a man-made substitution; wherein said variant has at least 85% but less than 100% amino acid sequence identity to an AprL-Clade subtilisin enzyme comprising a Motif A and/or a Motif B; wherein Motif A is selected from Motif 1, Motif 2, Motif 3, and Motif 4; wherein Motif B is selected from Motif 5, Motif 6, Motif 7, Motif 8, Motif 9, Motif 10, Motif 11, Motif 12, Motif 13, and Motif 14; and wherein the amino acid positions of the variant are numbered by correspondence with the amino acid sequence of SEQ ID NO:2.

[0070] In a further embodiment, one or more subtilisin variant described herein further comprises one or more substitution selected from: 3T, 4I, 9R, 15T, 67A, 75D, 86S, 97S, 98A/D/G, 100G/M/R, 102A, 103N/Y, 105A, 117R/V, 119D/N, 122S, 127L, 128Q, 129A, 159D, 166A, 169S, 193P, 194E, 198M, 216D, 217D, 221S, 231V, 234L, 235H, 244R, 251K, and 273A, and/or one or more insertion selected from a D inserted at position 36, an E inserted at position 96, and a D inserted between positions 98 and 99, wherein the amino acid positions of the variant are numbered by correspondence with the amino acid sequence of SEQ ID NO:2.

[0071] In one embodiment one or more subtilisin variant described herein is from a parent with 70%, 75%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% amino acid sequence identity to an AprL-Clade subtilisin enzyme comprising a Motif A, Motif B, or a combination thereof. In another embodiment, one or more subtilisin variant described herein is from a parent with 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% amino acid sequence identity to an AprL-Clade subtilisin enzyme comprising a Motif A, Motif B, or a combination thereof. In yet still another embodiment, one or more subtilisin variant described herein is from a parent with 70%, 75%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% amino acid sequence identity to an AprL-Clade subtilisin enzyme comprising a Motif A and Motif B. In an even further embodiment, one or more subtilisin variant described herein is from a parent with 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% amino acid sequence identity to an AprL-Clade subtilisin enzyme comprising a Motif A and a Motif B. The AprL-clade of subtilisin enzymes is more fully described in International Patent Application No. PCT/US16/32514, filed May 13, 2016. In some embodiments, the AprL-Clade subtilisin enzyme is selected from one of the enzymes listed in Figure 1. In another embodiment, the AprL-Clade subtilisin enzyme comprises an amino acid sequence of SEQ ID NO:2, 9, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, or 205. In still another embodiment, the AprL-Clade subtilisin enzyme comprises an amino acid sequence of SEQ ID NO:2, 9, 189, 198, 202, 203, or 205. In still another embodiment, the AprL-Clade subtilisin enzyme comprises an amino acid sequence of SEQ ID NO:2 or 9.

[0072] In one embodiment, the AprL-clade enzymes comprise Motif A, Motif B, or a combination thereof. In other embodiments, the AprL-clade enzymes comprise Motif A and Motif B. In some embodiments, one or more subtilisin variant described herein comprises Motif A, Motif B, or a combination thereof. In still other embodiments, one or more subtilisin variant described herein comprises Motif A and Motif B. In another embodiment, Motif A is selected from Motif 1, Motif 2, Motif 3, and Motif 4. In a still further embodiment, Motif B is selected from Motif 5, Motif 6, Motif 7, Motif 8, Motif 9, Motif 10, Motif 11, Motif 12, Motif 13, and Motif 14. In still other embodiments, the AprL-clade enzymes comprise Motif 1 between Asp32 and His63, wherein the amino acid positions are numbered by correspondence with the amino acid sequence of SEQ ID NO:2. In even still other embodiments, the AprL-clade enzymes comprise Motif 6 or Motif 7 between Lys236 and Arg246, wherein the amino acid positions are numbered by correspondence with the amino acid sequence of SEQ ID NO:2. In yet still further embodiments, the AprL-clade enzymes comprise (i) Motif 1 between Asp32 and His63, and (ii) Motif 6 or Motif 7 between Lys236 and Arg246, wherein the amino acid positions are numbered by correspondence with the amino acid sequence of SEQ ID NO:2. In still even other embodiments, one or more subtilisin variant described herein comprises Motif 1 between Asp32 and His63, wherein the amino acid positions are numbered by correspondence with the amino acid sequence of SEQ ID NO:2. In yet still an even further embodiment, one or more subtilisin variant described herein comprises Motif 6 or Motif 7 between Lys236 and Arg246, wherein the amino acid positions are numbered by correspondence with the amino acid sequence of SEQ ID NO:2. In yet another embodiment, one or more subtilisin variant described herein comprises (i) Motif 1 between Asp32 and His63, and (ii) Motif 6 or Motif 7 between Lys236 and Arg246, wherein the amino acid positions are numbered by correspondence with the amino acid sequence of SEQ ID NO:2.

[0073] In some embodiments, the AprL-clade enzymes comprise a YNT (SEQ ID NO:166) motif ("Motif 1"). In other embodiments, the AprL-clade enzymes comprise a Y56N57T58 (SEQ ID NO:167) motif, wherein the amino acid positions are numbered by correspondence with the amino acid sequence of SEQ ID NO:2 ("Motif 2"). In further embodiments, the AprL-clade enzymes comprise an FVX₁GEX₂X₃YNT (SEQ ID NO:168) motif, wherein X₁ is A or S, X₂ is absent or any amino acid, and X₃ is A or S ("Motif 3"). In a still further embodiment, the AprL-clade enzymes comprise a F50V51X₁52G53E54X₃55Y56N57T58 (SEQ ID NO:169) motif, wherein X₁ is an A or S and X₃ is an A or S; and further wherein the amino acid positions are numbered by correspondence with the amino acid sequence of SEQ ID NO:2 ("Motif 4"). In some embodiments, the AprL-clade enzymes comprise a loop region between residues 52-60, wherein the amino acid positions are numbered by correspondence with the amino acid sequence of SEQ ID NO:2.

[0074] In other embodiments, the AprL-clade enzymes comprise an LSX_4S (SEQ ID NO: 170) motif, wherein X_4 is A or G (“Motif 5”). In still other embodiments, the AprL-clade enzymes comprise an LSAS (SEQ ID NO: 171) motif (“Motif 6”). In another embodiment, the AprL-clade enzymes comprise an LSGS (SEQ ID NO: 172) motif (“Motif 7”). In some embodiments, the AprL-clade enzymes comprise an L240S241A242S243 (SEQ ID NO: 173) motif, wherein the amino acid positions are numbered by correspondence with the amino acid sequence of SEQ ID NO: 2 (“Motif 8”). In further embodiments, the AprL-clade enzymes comprise an L240S241G242S243 (SEQ ID NO: 174) motif, wherein the amino acid positions are numbered by correspondence with the amino acid sequence of SEQ ID NO: 2 (“Motif 9”). In still further embodiments, the AprL-clade enzymes comprise a $KXXXLSX_4SQX_5R$ (SEQ ID NO: 175) motif, wherein X is any amino acid, X_4 is A or G, and X_5 is an I or V (“Motif 10”). In yet further embodiments, the AprL-clade enzymes comprise a $KXXXLSASQX_5R$ (SEQ ID NO: 176) motif, wherein X is any amino acid and X_5 is an I or V (“Motif 11”). In an even yet further embodiment, the AprL-clade enzymes comprise a $KXXXLSGSQX_5R$ (SEQ ID NO: 177) motif, wherein X is any amino acid and X_5 is an I or V (“Motif 12”). In still further embodiments, the AprL-clade enzymes comprise a K236XXXLSASQX₅R246 (SEQ ID NO: 178) motif, wherein X is any amino acid and X_5 is an I or V; and further wherein the amino acid positions are numbered by correspondence with the amino acid sequence of SEQ ID NO: 2 (“Motif 13”). In an even yet still further embodiment, the AprL-clade enzymes comprise a K236XXXLSGSQX₅R246 (SEQ ID NO: 179) motif, wherein X is any amino acid and X_5 is an I or V; and further wherein the amino acid positions are numbered by correspondence with the amino acid sequence of SEQ ID NO: 2 (“Motif 14”).

[0075] In some embodiments, one or more subtilisin variant described herein comprises a YNT (SEQ ID NO: 166) motif (“Motif 1”). In other embodiments, one or more subtilisin variant described herein comprise a Y56N57T58 (SEQ ID NO: 167) motif, wherein the amino acid positions are numbered by correspondence with the amino acid sequence of SEQ ID NO: 2 (“Motif 2”). In further embodiments, one or more subtilisin variant described herein comprises an $FVX_1GEX_2X_3YNT$ (SEQ ID NO: 168) motif, wherein X_1 is A or S, X_2 is absent or any amino acid, and X_3 is A or S (“Motif 3”). In a still further embodiment, one or more subtilisin variant described herein comprises a F50V51X₁52G53E54X355Y56N57T58 (SEQ ID NO: 169) motif, wherein X_1 is an A or S and X_3 is an A or S; and further wherein the amino acid positions are numbered by correspondence with the amino acid sequence of SEQ ID NO: 2 (“Motif 4”). In some embodiments, one or more subtilisin variant described herein comprises a loop region between residues 52-60, wherein the amino acid positions are numbered by correspondence with the amino acid sequence of SEQ ID NO: 2.

[0076] In other embodiments, one or more subtilisin variant described herein comprises an LSX_4S (SEQ ID NO: 170) motif, wherein X_4 is A or G (“Motif 5”). In still other embodiments, one or more

subtilisin variant described herein comprises an LSAS (SEQ ID NO:171) motif (“Motif 6”). In another embodiment, one or more subtilisin variant described herein comprises an LSGS (SEQ ID NO:172) motif (“Motif 7”). In some embodiments, one or more subtilisin variant described herein comprises an L240S241A242S243 (SEQ ID NO:173) motif, wherein the amino acid positions are numbered by correspondence with the amino acid sequence of SEQ ID NO:2 (“Motif 8”). In further embodiments, one or more subtilisin variant described herein comprises an L240S241G242S243 (SEQ ID NO:174) motif, wherein the amino acid positions are numbered by correspondence with the amino acid sequence of SEQ ID NO:2 (“Motif 9”). In still further embodiments, one or more subtilisin variant described herein comprises a KXXXLSX₄SQX₅R (SEQ ID NO:175) motif, wherein X is any amino acid, X₄ is A or G, and X₅ is an I or V (“Motif 10”). In yet further embodiments, one or more subtilisin variant described herein comprises a KXXXLSASQX₅R (SEQ ID NO:176) motif, wherein X is any amino acid and X₅ is an I or V (“Motif 11”). In an even yet further embodiment, one or more subtilisin variant described herein comprises a KXXXLSGSQX₅R (SEQ ID NO:177) motif, wherein X is any amino acid and X₅ is an I or V (“Motif 12”). In still further embodiments, one or more subtilisin variant described herein comprises a K236XXXLSASQX₅R246 (SEQ ID NO:178) motif, wherein X is any amino acid and X₅ is an I or V; and further wherein the amino acid positions are numbered by correspondence with the amino acid sequence of SEQ ID NO:2 (“Motif 13”). In an even yet still further embodiment, one or more subtilisin variant described herein comprises a K236XXXLSGSQX₅R246 (SEQ ID NO:179) motif, wherein X is any amino acid and X₅ is an I or V; and further wherein the amino acid positions are numbered by correspondence with the amino acid sequence of SEQ ID NO:2 (“Motif 14”).

[0077] In another embodiment, one or more subtilisin variant described herein comprise two, three, or four or more variations described herein versus SEQ ID NO:2 with the proviso (i) that said variations versus SEQ ID NO:2 are not an insertion or deletion; (ii) that the variation is not a proline when the amino acid sequence of said variant comprises a variation versus SEQ ID NO:2 at one or more of positions 37, 187, 193, and 258; (iii) that said variation is not 26R, 38S, 234R, or 250E/N when the amino acids sequence of said variant comprises a variation versus SEQ ID NO:2 at two or more of positions 26, 38, 234, and 250; and/or (iv) that said variation versus SEQ ID NO:2 is not at position 130 or 193 when the amino acid sequence of said variant comprises a variation at position 128; wherein the amino acid positions of the variant are numbered by correspondence with the amino acid sequence of SEQ ID NO:2.

[0078] In yet another embodiment, one or more subtilisin variant described herein comprise two, three, or four or more variations at positions corresponding to SEQ ID NO:2 described herein with the proviso (i) that said variations at positions corresponding to SEQ ID NO:2 are not an insertion or deletion; (ii) that the variation is not a proline when the amino acid sequence of said variant comprises a variation at one or more corresponding SEQ ID NO:2 positions selected from positions 37, 187, 193, and 258; (iii)

that said variation is not 26R, 38S, 234R, or 250E/N when the amino acid sequence of said variant comprises a variation at two or more corresponding SEQ ID NO:2 positions selected from positions 26, 38, 234, and 250; (iv) that said variation at a position corresponding to SEQ ID NO:2 is not at position 130 or 193 when the amino acid sequence of said variant comprises a variation at position 128; and/or (v) that said variant does not consist of the amino acid sequence set forth as SEQ ID NO:5 in WO2016074925 or as SEQ ID NO:110 in JP201300714.

[0079] In still another embodiment, one or more subtilisin variant described herein comprises two, three, or four or more substitutions described herein with the proviso (i) that said substitution is not proline when that amino acid sequence of said variant comprises a substitution at one or more of positions 37, 187, 193, and 258; (ii) that said substitution is not 26R, 38S, 234R, or 250E/N when the amino acid sequence of said variant comprises a substitution at two or more of positions 26, 38, 234, and 250; and/or (iii) that said substitution is not at position 130 or 193 when the amino acid sequence of said variant comprises a substitution at position 128; wherein the amino acid positions of the variant are numbered by correspondence with the amino acid sequence of SEQ ID NO:2.

[0080] In yet another embodiment, one or more subtilisin variant described herein comprises two, three, or four or more substitutions described herein with the proviso (i) that said substitution is not proline when that amino acid sequence of said variant comprises a substitution at one or more of positions 37, 187, 193, and 258; (ii) that said substitution is not 26R, 38S, 234R, or 250E/N when the amino acid sequence of said variant comprises a substitution at two or more of positions 26, 38, 234, and 250; (iii) that said substitution is not at position 130 or 193 when the amino acid sequence of said variant comprises a substitution at position 128; and/or (iv) that said variant does not consist of the amino acid sequence set forth as SEQ ID NO:5 in WO2016074925 or as SEQ ID NO:110 in JP201300714.

[0081] In one embodiment, one or more subtilisin variant described herein comprises an amino acid sequence having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% but less than 100 % amino acid sequence identity to the amino acid sequence of an AprL-Clade subtilisin enzyme identified in Figure 1. In some embodiments, one or more subtilisin variant described herein comprise an amino acid sequence having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% but less than 100% amino acid sequence identity to the amino acid sequence of SEQ ID NO:2 or 9. In other embodiments, one or more subtilisin variant described herein comprises an amino acid sequence having at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% but less than

100% amino acid sequence identity to the amino acid sequence of (i) an AprL-Clade subtilisin enzyme identified in Figure 1, or (ii) SEQ ID NO: 2 or 9. In another embodiment, one or more subtilisin variant described herein comprises an amino acid sequence having at least 89% but less than 100% amino acid sequence identity to the amino acid sequence of an AprL-Clade subtilisin enzyme identified in Figure 1. In other embodiments, one or more subtilisin variant described herein comprises an amino acid sequence having at least 89% but less than 100% amino acid sequence identity to the amino acid sequence of SEQ ID NO:2 or 9. In further embodiments, one or more subtilisin variant described herein comprises an amino acid sequence having at least 90% but less than 100% amino acid sequence identity to the amino acid sequence of an AprL-Clade subtilisin enzyme identified in Figure 1. In some embodiments, one or more subtilisin variant described herein comprises an amino acid sequence having at least 90% but less than 100% amino acid sequence identity to the amino acid sequence of SEQ ID NO:2 or 9. In still other embodiments, one or more subtilisin variant described herein comprises an amino acid sequence having at least 95%, at least 96%, at least 97%, at least 98%, at least 99% but less than 100% amino acid sequence identity to the amino acid sequence of an AprL-Clade subtilisin enzyme identified in Figure 1. In yet still other embodiments, one or more subtilisin variant described herein comprises an amino acid sequence having at least 95%, at least 96%, at least 97%, at least 98%, at least 99% but less than 100% amino acid sequence identity to the amino acid sequence of SEQ ID NO:2 or 9. In even still other embodiments, one or more subtilisin variant described herein is an isolated, recombinant, substantially pure, or non-naturally occurring subtilisin having subtilisin activity or casein hydrolysis activity (for example, dimethyl casein hydrolysis activity).

[0082] In one embodiment, one or more subtilisin variant described herein has enzymatic activity (e.g., protease activity) and thus is useful in cleaning applications, including but not limited to, methods for cleaning dishware items, tableware items, fabrics, and items having hard surfaces (e.g., the hard surface of a table, table top, wall, furniture item, floor, ceiling, etc.). Exemplary cleaning compositions comprising one or more subtilisin variant described herein are described *infra*. The enzymatic activity (e.g., protease enzyme activity) of one or more subtilisin variant described herein can be determined readily using procedures well known to those of ordinary skill in the art. The Examples presented *infra* describe methods for evaluating the enzymatic activity and cleaning performance. The performance of polypeptide enzymes of the invention in removing stains (e.g., a protein stain such as blood/milk/ink, pigment/milk/ink or egg yolk), cleaning hard surfaces, or cleaning laundry, dishware or tableware item(s) can be readily determined using procedures well known in the art and/or by using procedures set forth in the Examples.

[0083] In some embodiments, one or more subtilisin variant described herein has one or more improved property when compared to a reference subtilisin, wherein the improved property is selected

from improved protease activity, improved cleaning performance in detergent, improved thermostability in detergent, and improved aged laundry cleaning in detergent, wherein the detergent is optionally a boron-free composition. In other embodiments, one or more subtilisin variant described herein has one or more improved property when compared to a reference subtilisin, wherein the improved property is selected from improved protease activity, improved cleaning performance in detergent, improved thermostability in detergent, and improved aged laundry cleaning in detergent, wherein the detergent is optionally a boron-free composition, wherein the reference subtilisin comprises an amino acid sequence of SEQ ID NO:2, 9, or 63. In still other embodiments, one or more subtilisin variant described herein has one or more improved property when compared to a reference subtilisin, wherein the improved property is selected from improved protease activity, improved cleaning performance in detergent, improved thermostability in detergent, and improved aged laundry cleaning in detergent, wherein the detergent is optionally a boron-free composition, wherein the reference subtilisin comprises an amino acid sequence of SEQ ID NO:2 or 9. In one embodiment, one or more subtilisin variant described herein is more stable through a longer wash period as compared to a reference subtilisin. In another embodiment, one or more subtilisin variant described herein is more stable through a short, cool wash cycle or a long, hot wash-cycle as compared to a reference subtilisin. In a still yet further embodiment, the one or more improved property is (i) improved protease activity, wherein said variant has a $PI > 1$ on N-suc-AAPF-pNA or dimethyl casein substrate; (ii) improved cleaning performance in detergent, wherein said variant has a BMI, POM, egg, and/or egg yolk cleaning $PI > 1$; (iii) improved thermostability in detergent, wherein said variant has a stability $PI > 1$; and/or (iv) improved aged laundry cleaning in detergent, wherein said variant has an aged laundry cleaning $PI > 1$; wherein the detergent is optionally a boron-free composition. In yet a further embodiment, the one or more improved property is (i) improved protease activity, wherein said variant has a $PI > 1$ on N-suc-AAPF-pNA or dimethyl casein substrate; (ii) improved cleaning performance in detergent, wherein said variant has a BMI, POM, egg, and/or egg yolk cleaning $PI > 1$; (iii) improved thermostability in detergent, wherein said variant has a stability $PI > 1$ or % remaining activity $\geq 5\%$, 10% , 20% , 30% , 40% , 50% , 60% , 70% , 80% , 90% , 91% , 92% , 93% , 94% , 95% , 96% , 97% , 98% , 99% , 100% or more after 15-20 minutes at a stress temperature of 60°C , 61°C , 62°C , 63°C , 64°C , 65°C , 66°C , 67°C , 68°C , 69°C or more; and/or (iv) improved aged laundry cleaning in detergent, wherein said variant has (a) an aged laundry cleaning $PI > 1$, (b) remaining cleaning activity $>10\%$, 20% , 30% , 40% , 50% , 60% , 70% , 80% , 90% , 91% , 92% , 93% , 94% , 95% , 96% , 97% , 98% , 99% , 100% or more after 24 hours at 55°C , and/or (c) remaining cleaning activity $>10\%$, 20% , 30% , 40% , 50% , 60% , 70% , 80% , 90% , 91% , 92% , 93% , 94% , 95% , 96% , 97% , 98% , 99% , 100% or more after 3 days or 3 weeks at 37°C ; wherein the detergent is optionally a boron-free composition. In an even further embodiment, one or more subtilisin variant described herein has improved protease activity, wherein said

variant has a PI >1 on N-suc-AAPF-pNA or dimethyl casein substrate. In a still even further embodiment, one or more subtilisin variant described herein has improved cleaning performance in detergent, wherein said variant has a BMI, POM, and/or egg yolk cleaning PI >1, wherein the detergent is optionally a boron-free composition. In another embodiment, one or more subtilisin variant described herein has improved thermostability in detergent, wherein said variant has a stability PI > 1, wherein the detergent is optionally a boron-free composition. In still another embodiment, one or more subtilisin variant described herein has improved thermostability in detergent, wherein said variant has a stability PI > 1 or % remaining activity $\geq$ 5%, 10% 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 100% or more after 15-20 minutes at a stress temperature of 60°C, 61°C, 62°C, 63°C, 64°C, 65°C, 66°C, 67°C, 68°C, 69°C or more, wherein the detergent is optionally a boron-free composition. In another embodiment, one or more subtilisin variant described herein has improved aged laundry cleaning in detergent, wherein said variant has (a) an aged laundry cleaning PI > 1, (b) remaining cleaning activity >10%, 20% 30%, 40%, 50%, 60%, 70%, 80%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 100% or more after 24 hours at 55°C, and/or (c) remaining cleaning activity >10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 100% or more after 3 days or 3 weeks at 37°C, wherein the detergent is optionally a boron-free composition. In another embodiment, one or more subtilisin variant described herein has improved aged laundry cleaning in detergent, wherein said variant has an aged laundry cleaning PI > 1, wherein the detergent is optionally a boron-free composition. In another embodiment, one or more subtilisin variant described herein has improved protease activity, wherein said variant has a PI > 1 on N-suc-AAPF-pNA or dimethyl casein substrate and said PI is measured in accordance with the protease activity assay of Example 1. In a further embodiment, one or more subtilisin variant described herein has improved cleaning performance in detergent, wherein said variant has a BMI, POM, egg, and/or egg yolk cleaning PI > 1 and said PI is measured in accordance with the cleaning performance in laundry (HDL and HDD) and ADW detergents assay of Example 1. In an even further embodiment, one or more subtilisin variant described herein has improved thermostability in detergent, wherein said variant has a stability PI > 1 and said PI is measured in accordance with the general sample set-up for stability assay of Example 1. In yet an even further embodiment, one or more subtilisin variant described herein has improved thermostability in detergent, wherein said variant has a % remaining activity $\geq$ 5%, 10% 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 100% or more after 15-20 minutes at a stress temperature of 60°C, 61°C, 62°C, 63°C, 64°C, 65°C, 66°C, 67°C, 68°C, 69°C or more and said remaining activity is measured in accordance with the general sample set-up for stability assay of Example 1. In a still further embodiment, one or more subtilisin variant described herein has improved aged laundry cleaning in detergent, wherein said variant has an aged laundry cleaning PI > 1 and said PI

is measured in accordance with the aged laundry cleaning assay of Example 1. In an even still further embodiment, one or more subtilisin variant described herein has improved aged laundry cleaning in detergent, wherein said variant has remaining cleaning activity >10%, 20% 30%, 40%, 50%, 60%, 70%, 80%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 100% or more after 24 hours at 55°C, and/or remaining cleaning activity >10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 100% or more after 3 days or 3 weeks at 37°C; and said remaining cleaning activity is measured in accordance with the aged laundry cleaning assay of Example 1.

[0084] In some embodiments, the one or more subtilisin variant described herein demonstrates cleaning performance in a cleaning composition. Cleaning compositions often include ingredients harmful to the stability and performance of enzymes, making cleaning compositions a harsh environment for enzymes, e.g. serine proteases, to retain function. Thus, it is not trivial for an enzyme to be put in a cleaning composition and expect enzymatic function (e.g. serine protease activity, such as demonstrated by cleaning performance). In some embodiments, the one or more subtilisin variant described herein demonstrates cleaning performance in automatic dishwashing (ADW) detergent compositions. In some embodiments, the cleaning performance in ADW detergent compositions includes cleaning of egg yolk stains. In some embodiments, the one or more subtilisin variant described herein demonstrates cleaning performance in laundry detergent compositions. In some embodiments, the cleaning performance in laundry detergent compositions includes cleaning of blood/milk/ink, egg, egg yolk, and/or POM stains. In each of the cleaning compositions, one or more subtilisin variant described herein demonstrates cleaning performance with or without a bleach component. In an even still further embodiment, one or more ADW or laundry detergent composition described herein comprises one or more subtilisin variant described herein, wherein said variant is stable in the presence of one or more adjunct material and/or one or more additional enzyme and/or further wherein said variant is stable to autoproteolysis.

[0085] One or more subtilisin variant described herein can be subject to various changes, such as one or more amino acid insertion, deletion, and/or substitution, either conservative or non-conservative, including where such changes do not substantially alter the enzymatic activity of the variant. Similarly, a nucleic acid of the invention can also be subject to various changes, such as one or more substitution of one or more nucleotide in one or more codon such that a particular codon encodes the same or a different amino acid, resulting in either a silent variation (e.g., when the encoded amino acid is not altered by the nucleotide mutation) or non-silent variation; one or more deletion of one or more nucleic acids (or codon) in the sequence; one or more addition or insertion of one or more nucleic acids (or codon) in the sequence; and/or cleavage of, or one or more truncation, of one or more nucleic acid (or codon) in the sequence. Many such changes in the nucleic acid sequence may not substantially alter the enzymatic activity of the resulting encoded polypeptide enzyme compared to the polypeptide enzyme encoded by the

original nucleic acid sequence. A nucleic acid sequence described herein can also be modified to include one or more codon that provides for optimum expression in an expression system (e.g., bacterial expression system), while, if desired, said one or more codon still encodes the same amino acid(s).

[0086] Described herein is one or more isolated, non-naturally occurring, or recombinant polynucleotide comprising a nucleic acid sequence that encodes one or more subtilisin variant described herein, or recombinant polypeptide or active fragment thereof. One or more nucleic acid sequence described herein is useful in recombinant production (e.g., expression) of one or more subtilisin variant described herein, typically through expression of a plasmid expression vector comprising a sequence encoding the one or more subtilisin variant described herein or fragment thereof. One embodiment provides nucleic acids encoding one or more subtilisin variant described herein, wherein the variant is a mature form having proteolytic activity. In some embodiments, one or more subtilisin variant described herein is expressed recombinantly with a homologous pro-peptide sequence. In other embodiments, one or more subtilisin variant described herein is expressed recombinantly with a heterologous pro-peptide sequence (e.g., GG36 pro-peptide sequence).

[0087] One or more nucleic acid sequence described herein can be generated by using any suitable synthesis, manipulation, and/or isolation techniques, or combinations thereof. For example, one or more polynucleotide described herein may be produced using standard nucleic acid synthesis techniques, such as solid-phase synthesis techniques that are well-known to those skilled in the art. In such techniques, fragments of up to 50 or more nucleotide bases are typically synthesized, then joined (e.g., by enzymatic or chemical ligation methods) to form essentially any desired continuous nucleic acid sequence. The synthesis of the one or more polynucleotide described herein can be also facilitated by any suitable method known in the art, including but not limited to chemical synthesis using the classical phosphoramidite method (*See e.g.*, Beaucage et al. Tetrahedron Letters 22:1859-69 (1981)), or the method described in Matthes et al., EMBO J. 3:801-805 (1984) as is typically practiced in automated synthetic methods. One or more polynucleotide described herein can also be produced by using an automatic DNA synthesizer. Customized nucleic acids can be ordered from a variety of commercial sources (e.g., Midland Certified Reagent Company, Great American Gene Company, Operon Technologies Inc., and DNA 2.0). Other techniques for synthesizing nucleic acids and related principles are described by, for example, Itakura et al., Ann. Rev. Biochem. 53:323 (1984) and Itakura et al., Science 198:1056 (1984).

[0088] Recombinant DNA techniques useful in modification of nucleic acids are well known in the art, such as, for example, restriction endonuclease digestion, ligation, reverse transcription and cDNA production, and polymerase chain reaction (e.g., PCR). One or more polynucleotide described herein may also be obtained by screening cDNA libraries using one or more oligonucleotide probes that can hybridize to or PCR-amplify polynucleotides which encode one or more AprL-clade variant described

herein, or recombinant polypeptide or active fragment thereof. Procedures for screening and isolating cDNA clones and PCR amplification procedures are well known to those of skill in the art and described in standard references known to those skilled in the art. One or more polynucleotide described herein can be obtained by altering a naturally occurring polynucleotide backbone (e.g., that encodes one or more subtilisin variant described herein or reference subtilisin) by, for example, a known mutagenesis procedure (e.g., site-directed mutagenesis, site saturation mutagenesis, and in vitro recombination). A variety of methods are known in the art that are suitable for generating modified polynucleotides described herein that encode one or more subtilisin variant described herein, including, but not limited to, for example, site-saturation mutagenesis, scanning mutagenesis, insertional mutagenesis, deletion mutagenesis, random mutagenesis, site-directed mutagenesis, and directed-evolution, as well as various other recombinatorial approaches.

[0089] A further embodiment is directed to one or more vector comprising one or more subtilisin variant described herein (e.g., a polynucleotide encoding one or more subtilisin variant described herein); expression vectors or expression cassettes comprising one or more nucleic acid or polynucleotide sequence described herein; isolated, substantially pure, or recombinant DNA constructs comprising one or more nucleic acid or polynucleotide sequence described herein; isolated or recombinant cells comprising one or more polynucleotide sequence described herein; and compositions comprising one or more such vector, nucleic acid, expression vector, expression cassette, DNA construct, cell, cell culture, or any combination or mixtures thereof.

[0090] Some embodiments are directed to one or more recombinant cell comprising one or more vector (e.g., expression vector or DNA construct) described herein which comprises one or more nucleic acid or polynucleotide sequence described herein. Some such recombinant cells are transformed or transfected with such at least one vector, although other methods are available and known in the art. Such cells are typically referred to as host cells. Some such cells comprise bacterial cells, including, but not limited to *Bacillus sp.* cells, such as *B. subtilis* cells. Other embodiments are directed to recombinant cells (e.g., recombinant host cells) comprising one or more subtilisin described herein.

[0091] In some embodiments, one or more vector described herein is an expression vector or expression cassette comprising one or more polynucleotide sequence described herein operably linked to one or more additional nucleic acid segments required for efficient gene expression (e.g., a promoter operably linked to one or more polynucleotide sequence described herein). A vector may include a transcription terminator and/or a selection gene (e.g., an antibiotic resistant gene) that enables continuous cultural maintenance of plasmid-infected host cells by growth in antimicrobial-containing media.

[0092] An expression vector may be derived from plasmid or viral DNA, or in alternative embodiments, contains elements of both. Exemplary vectors include, but are not limited to pC194,

pJH101, pE194, pHP13 (See, Harwood and Cutting [eds.], Chapter 3, Molecular Biological Methods for Bacillus, John Wiley & Sons (1990); suitable replicating plasmids for *B. subtilis* include those listed on p. 92). (See also, Perego, "Integrational Vectors for Genetic Manipulations in Bacillus subtilis"; Sonenshein et al., [eds.], "Bacillus subtilis and Other Gram-Positive Bacteria: Biochemistry, Physiology and Molecular Genetics", American Society for Microbiology, Washington, D.C. (1993), pp. 615-624); and p2JM103BBI).

[0093] For expression and production of a protein of interest (e.g., one or more subtilisin variant described herein) in a cell, one or more expression vector comprising one or more copy of a polynucleotide encoding one or more subtilisin variant described herein, and in some instances comprising multiple copies, is transformed into the cell under conditions suitable for expression of the variant. In some embodiments, a polynucleotide sequence encoding one or more subtilisin variant described herein (as well as other sequences included in the vector) is integrated into the genome of the host cell, while in other embodiments, a plasmid vector comprising a polynucleotide sequence encoding one or more subtilisin variant described herein remains as autonomous extra-chromosomal element within the cell. Some embodiments provide both extrachromosomal nucleic acid elements as well as incoming nucleotide sequences that are integrated into the host cell genome. The vectors described herein are useful for production of the one or more subtilisin variant described herein. In some embodiments, a polynucleotide construct encoding one or more subtilisin variant described herein is present on an integrating vector that enables the integration and optionally the amplification of the polynucleotide encoding the variant into the host chromosome. Examples of sites for integration are well known to those skilled in the art. In some embodiments, transcription of a polynucleotide encoding one or more subtilisin variant described herein is effectuated by a promoter that is the wild-type promoter for the parent subtilisin. In some other embodiments, the promoter is heterologous to the one or more subtilisin variant described herein, but is functional in the host cell. Exemplary promoters for use in bacterial host cells include, but are not limited to the amyE, amyQ, amyL, pstS, sacB, pSPAC, pAprE, pVeg, pHpaII promoters; the promoter of the *B. stearothermophilus* maltogenic amylase gene; the *B. amyloliquefaciens* (BAN) amylase gene; the *B. subtilis* alkaline protease gene; the *B. clausii* alkaline protease gene; the *B. pumilis* xylosidase gene; the *B. thuringiensis* cryIIIA; and the *B. licheniformis* alpha-amylase gene. Additional promoters include, but are not limited to the A4 promoter, as well as phage Lambda PR or PL promoters and the *E. coli* lac, trp or tac promoters.

[0094] One or more subtilisin variant described herein can be produced in host cells of any suitable microorganism, including bacteria and fungi. In some embodiments, one or more subtilisin variant described herein can be produced in Gram-positive bacteria. In some embodiments, the host cells are *Bacillus* spp., *Streptomyces* spp., *Escherichia* spp., *Aspergillus* spp., *Trichoderma* spp., *Pseudomonas*

spp., *Corynebacterium spp.*, *Saccharomyces spp.*, or *Pichia spp.* In some embodiments, one or more subtilisin variant described herein is produced by *Bacillus sp.* host cells. Examples of *Bacillus sp.* host cells that find use in the production of the one or more subtilisin variant described herein include, but are not limited to *B. licheniformis*, *B. lentus*, *B. subtilis*, *B. amyloliquefaciens*, *B. lentus*, *B. brevis*, *B. stearothermophilus*, *B. alkalophilus*, *B. coagulans*, *B. circulans*, *B. pumilis*, *B. thuringiensis*, *B. clausii*, and *B. megaterium*, as well as other organisms within the genus *Bacillus*. In some embodiments, *B. subtilis* host cells are used to produce the variants described herein. USPNs 5,264,366 and 4,760,025 (RE 34,606) describe various *Bacillus* host strains that can be used to produce one or more subtilisin variant described herein, although other suitable strains can be used.

[0095] Several bacterial strains that can be used to produce one or more subtilisin variant described herein include non-recombinant (i.e., wild-type) *Bacillus sp.* strains, as well as variants of naturally-occurring strains and/or recombinant strains. In some embodiments, the host strain is a recombinant strain, wherein a polynucleotide encoding one or more subtilisin variant described herein has been introduced into the host. In some embodiments, the host strain is a *B. subtilis* host strain and particularly a recombinant *B. subtilis* host strain. Numerous *B. subtilis* strains are known, including, but not limited to for example, 1A6 (ATCC 39085), 168 (1A01), SB19, W23, Ts85, B637, PB1753 through PB1758, PB3360, JH642, 1A243 (ATCC 39,087), ATCC 21332, ATCC 6051, MI113, DE100 (ATCC 39,094), GX4931, PBT 110, and PEP 211 strain (See e.g., Hoch et al., Genetics 73:215–228 (1973); See also, US 4,450,235; US 4,302,544; and EP 0134048). The use of *B. subtilis* as an expression host cell is well known in the art (See e.g., Palva et al., Gene 19:81-87 (1982); Fahnestock and Fischer, J. Bacteriol., 165:796–804 (1986); and Wang et al., Gene 69:39–47 (1988)).

[0096] In some embodiments, the *Bacillus* host cell is a *Bacillus sp.* that includes a mutation or deletion in at least one of the following genes: *degU*, *degS*, *degR* and *degQ*. In some embodiments, the mutation is in a *degU* gene, and in some embodiments the mutation is *degU(Hy)32* (See e.g., Msadek et al., J. Bacteriol. 172:824-834 (1990); and Olmos et al., Mol. Gen. Genet. 253:562–567 (1997)). In some embodiments, the *Bacillus* host comprises a mutation or deletion in *scoC4* (See e.g., Caldwell et al., J. Bacteriol. 183:7329-7340 (2001)); *spoIIE* (See e.g., Arigoni et al., Mol. Microbiol. 31:1407-1415 (1999)); and/or *oppA* or other genes of the *opp* operon (See e.g., Perego et al., Mol. Microbiol. 5:173-185 (1991)). Indeed, it is contemplated that any mutation in the *opp* operon that causes the same phenotype as a mutation in the *oppA* gene will find use in some embodiments of the altered *Bacillus* strain described herein. In some embodiments, these mutations occur alone, while in other embodiments, combinations of mutations are present. In some embodiments, an altered *Bacillus* host cell strain that can be used to produce one or more subtilisin variant described herein is a *Bacillus* host strain that already includes a mutation in one or more of the above-mentioned genes. In addition, *Bacillus sp.* host cells that comprise

mutation(s) and/or deletion(s) of endogenous protease genes find use. In some embodiments, the *Bacillus* host cell comprises a deletion of the *aprE* and the *nprE* genes. In other embodiments, the *Bacillus sp.* host cell comprises a deletion of 5 protease genes, while in other embodiments the *Bacillus sp.* host cell comprises a deletion of 9 protease genes (*See e.g.*, US 2005/0202535).

[0097] Host cells are transformed with one or more nucleic acid sequence encoding one or more subtilisin variant described herein using any suitable method known in the art. Methods for introducing a nucleic acid (e.g., DNA) into *Bacillus* cells or *E. coli* cells utilizing plasmid DNA constructs or vectors and transforming such plasmid DNA constructs or vectors into such cells are well known. In some embodiments, the plasmids are subsequently isolated from *E. coli* cells and transformed into *Bacillus* cells. However, it is not essential to use intervening microorganisms such as *E. coli*, and in some embodiments, a DNA construct or vector is directly introduced into a *Bacillus* host.

[0098] Exemplary methods for introducing one or more nucleic acid sequence described herein into *Bacillus* cells are described in, for example, Ferrari et al., "Genetics," in Harwood et al. [eds.], *Bacillus*, Plenum Publishing Corp. (1989), pp. 57-72; Saunders et al., *J. Bacteriol.* 157:718-726 (1984); Hoch et al., *J. Bacteriol.* 93:1925-1937 (1967); Mann et al., *Current Microbiol.* 13:131-135 (1986); Holubova, *Folia Microbiol.* 30:97 (1985); Chang et al., *Mol. Gen. Genet.* 168:11-115 (1979); Vorobjeva et al., *FEMS Microbiol. Lett.* 7:261-263 (1980); Smith et al., *Appl. Env. Microbiol.* 51:634 (1986); Fisher et al., *Arch. Microbiol.* 139:213-217 (1981); and McDonald, *J. Gen. Microbiol.* 130:203 (1984)). Indeed, such methods as transformation, including protoplast transformation and transfection, transduction, and protoplast fusion are well known and suited for use herein. Methods known in the art to transform *Bacillus* cells include such methods as plasmid marker rescue transformation, which involves the uptake of a donor plasmid by competent cells carrying a partially homologous resident plasmid (*See*, Contente et al., *Plasmid* 2:555-571 (1979); Haima et al., *Mol. Gen. Genet.* 223:185-191 (1990); Weinrauch et al., *J. Bacteriol.* 154:1077-1087 (1983); and Weinrauch et al., *J. Bacteriol.* 169:1205-1211 (1987)). In this method, the incoming donor plasmid recombines with the homologous region of the resident "helper" plasmid in a process that mimics chromosomal transformation.

[0099] In addition to commonly used methods, in some embodiments, host cells are directly transformed with a DNA construct or vector comprising a nucleic acid encoding one or more subtilisin variant described herein (i.e., an intermediate cell is not used to amplify, or otherwise process, the DNA construct or vector prior to introduction into the host cell). Introduction of a DNA construct or vector described herein into the host cell includes those physical and chemical methods known in the art to introduce a nucleic acid sequence (e.g., DNA sequence) into a host cell without insertion into the host genome. Such methods include, but are not limited to calcium chloride precipitation, electroporation, naked DNA, and liposomes. In additional embodiments, DNA constructs or vector are co-transformed

with a plasmid, without being inserted into the plasmid. In further embodiments, a selective marker is deleted from the altered *Bacillus* strain by methods known in the art (*See*, Stahl et al., *J. Bacteriol.* 158:411-418 (1984); and Palmeros et al., *Gene* 247:255 -264 (2000)).

[00100] In some embodiments, the transformed cells are cultured in conventional nutrient media. The suitable specific culture conditions, such as temperature, pH and the like are known to those skilled in the art and are well described in the scientific literature. Some embodiments provide a culture (e.g., cell culture) comprising one or more subtilisin variant or nucleic acid sequence described herein.

[00101] In some embodiments, host cells transformed with one or more polynucleotide sequence encoding one or more subtilisin variant described herein are cultured in a suitable nutrient medium under conditions permitting the expression of the variant, after which the resulting variant is recovered from the culture. In some embodiments, the variant produced by the cells is recovered from the culture medium by conventional procedures, including, but not limited to, for example, separating the host cells from the medium by centrifugation or filtration, precipitating the proteinaceous components of the supernatant or filtrate by means of a salt (e.g., ammonium sulfate), and chromatographic purification (e.g., ion exchange, gel filtration, affinity, etc.).

[00102] In some embodiments, one or more subtilisin variant produced by a recombinant host cell is secreted into the culture medium. A nucleic acid sequence that encodes a purification facilitating domain may be used to facilitate purification of the variant. A vector or DNA construct comprising a polynucleotide sequence encoding one or more subtilisin variant described herein may further comprise a nucleic acid sequence encoding a purification facilitating domain to facilitate purification of the variant (*See e.g.*, Kroll et al., *DNA Cell Biol.* 12:441-53 (1993)). Such purification facilitating domains include, but are not limited to, for example, metal chelating peptides such as histidine-tryptophan modules that allow purification on immobilized metals (*See*, Porath, *Protein Expr. Purif.* 3:263-281 [1992]), protein A domains that allow purification on immobilized immunoglobulin, and the domain utilized in the FLAGS extension/affinity purification system. The inclusion of a cleavable linker sequence such as Factor XA or enterokinase (e.g., sequences available from Invitrogen, San Diego, CA) between the purification domain and the heterologous protein also find use to facilitate purification.

[00103] A variety of methods can be used to determine the level of production of one or more mature subtilisin variant described herein in a host cell. Such methods include, but are not limited to, for example, methods that utilize either polyclonal or monoclonal antibodies specific for the protease. Exemplary methods include, but are not limited to enzyme-linked immunosorbent assays (ELISA), radioimmunoassays (RIA), fluorescent immunoassays (FIA), and fluorescent activated cell sorting (FACS). These and other assays are well known in the art (*See e.g.*, Maddox et al., *J. Exp. Med.* 158:1211 (1983)).

[00104] Some other embodiments provide methods for making or producing one or more mature subtilisin variant described herein. A mature subtilisin variant does not include a signal peptide or a propeptide sequence. Some methods comprise making or producing one or more subtilisin variant described herein in a recombinant bacterial host cell, such as for example, a *Bacillus sp.* cell (e.g., a *B. subtilis* cell). Other embodiments provide a method of producing one or more subtilisin variant described herein, wherein the method comprises cultivating a recombinant host cell comprising a recombinant expression vector comprising a nucleic acid sequence encoding one or more subtilisin variant described herein under conditions conducive to the production of the variant. Some such methods further comprise recovering the variant from the culture.

[00105] Further embodiments provide methods of producing one or more subtilisin variant described herein, wherein the methods comprise: (a) introducing a recombinant expression vector comprising a nucleic acid encoding the variant into a population of cells (e.g., bacterial cells, such as *B. subtilis* cells); and (b) culturing the cells in a culture medium under conditions conducive to produce the variant encoded by the expression vector. Some such methods further comprise: (c) isolating the variant from the cells or from the culture medium.

[00106] Unless otherwise noted, all component or composition levels provided herein are made in reference to the active level of that component or composition, and are exclusive of impurities, for example, residual solvents or by-products, which may be present in commercially available sources. Enzyme components weights are based on total active protein. All percentages and ratios are calculated by weight unless otherwise indicated. All percentages and ratios are calculated based on the total composition unless otherwise indicated. Compositions described herein include cleaning compositions, such as detergent compositions. In the exemplified detergent compositions, the enzyme levels are expressed by pure enzyme by weight of the total composition and unless otherwise specified, the detergent ingredients are expressed by weight of the total compositions.

[00107] In one embodiment, one or more subtilisin variant described herein is useful in cleaning applications, such as, for example, but not limited to, cleaning dishware or tableware items, fabrics, medical instruments and items having hard surfaces (e.g., the hard surface of a table, table top, wall, furniture item, floor, and ceiling). In other embodiments, one or more subtilisin variant described herein is useful in disinfecting applications, such as, for example, but not limited to, disinfecting an automatic dishwashing or laundry machine.

[00108] Another embodiment is directed to a composition comprising one or more subtilisin variant described herein. In some embodiments, the composition is a cleaning composition. In other embodiments, the composition is a detergent composition. In yet other embodiments, the composition is selected from a laundry detergent composition, an automatic dishwashing (ADW) composition, a hand

(manual) dishwashing detergent composition, a hard surface cleaning composition, an eyeglass cleaning composition, a medical instrument cleaning composition, a disinfectant (e.g., malodor or microbial) composition, and a personal care cleaning composition. In still other embodiments, the composition is a laundry detergent composition, an ADW composition, or a hand (manual) dishwashing detergent composition. Even still further embodiments are directed to fabric cleaning compositions, while other embodiments are directed to non-fabric cleaning compositions. In some embodiments, the cleaning composition is boron-free. In other embodiments, the cleaning composition is phosphate-free. In still other embodiments, the composition comprises one or more subtilisin variant described herein and one or more of an excipient, adjunct material, and/or additional enzyme.

[00109] In yet still a further embodiment, the composition described herein contains phosphate, is phosphate-free, contains boron, is boron-free, or combinations thereof. In other embodiments, the composition is a boron-free composition. In some embodiments, a boron-free composition is a composition to which a borate stabilizer has not been added. In another embodiment, a boron-free composition is a composition that contains less than 5.5% boron. In a still further embodiment, a boron-free composition is a composition that contains less than 4.5% boron. In yet still another embodiment, a boron-free composition is a composition that contains less than 3.5% boron. In yet still a further embodiment, a boron-free composition is a composition that contains less than 2.5% boron. In even further embodiments, a boron-free composition is a composition that contains less than 1.5% boron. In another embodiment, a boron-free composition is a composition that contains less than 1.0% boron. In still further embodiments, a boron-free composition is a composition that contains less than 0.5% boron. In still further embodiments, a boron-free composition is a composition substantially-free of boron.

[00110] In another embodiment, one or more composition described herein is in a form selected from gel, tablet, powder, granular, solid, liquid, unit dose, and combinations thereof. In yet another embodiment, one or more composition described herein is in a form selected from a low water compact formula, low water HDL or UD, or high water formula or HDL. In some embodiments, the cleaning composition described herein is in a unit dose form. In other embodiments, the unit dose form is selected from pills, tablets, capsules, gelcaps, sachets, pouches, multi-compartment pouches, and pre-measured powders or liquids. In some embodiments, the unit dose format is designed to provide controlled release of the ingredients within a multi-compartment pouch (or other unit dose format). Suitable unit dose and controlled release formats are described, for example, in EP 2100949; WO 02/102955; US 4,765,916; US 4,972,017; and WO 04/111178. In some embodiments, the unit dose form is a tablet or powder contained in a water-soluble film or pouch.

[00111] Exemplary laundry detergent compositions include, but are not limited to, for example, liquid and powder laundry detergent compositions. Exemplary hard surface cleaning compositions include, but

not limited to, for example, compositions used to clean the hard surface of a non-dishware item, non-tableware item, table, table top, furniture item, wall, floor, and ceiling. Exemplary hard surface cleaning compositions are described, for example, in USPNs 6,610,642, 6,376,450, and 6,376,450. Exemplary personal care compositions include, but are not limited to, compositions used to clean dentures, teeth, hair, contact lenses, and skin. Exemplary components of such oral care composition include those described in, for example, US 6,376,450.

[00112] In some embodiments, one or more subtilisin variant described herein cleans at low temperatures. In other embodiments, one or more composition described herein cleans at low temperatures. In other embodiments, one or more composition described herein comprises an effective amount of one or more subtilisin variant described herein as useful or effective for cleaning a surface in need of proteinaceous stain removal

[00113] In some embodiments, adjunct materials are incorporated, for example, to assist or enhance cleaning performance; for treatment of the substrate to be cleaned; or to modify the aesthetics of the cleaning composition as is the case with perfumes, colorants, dyes or the like. One embodiment is directed to a composition comprising one or more adjunct material and one or more subtilisin variant described herein. Another embodiment is directed to a composition comprising one or more adjunct material and one or more subtilisin variant described herein, wherein the adjunct material is selected from a bleach catalyst, an additional enzyme, an enzyme stabilizer (including, for example, an enzyme stabilizing system), a chelant, an optical brightener, a soil release polymer, a dye transfer agent, a dispersants, a suds suppressor, a dye, a perfume, a colorant, a filler, a photoactivator, a fluorescer, a fabric conditioner, a hydrolyzable surfactant, a preservative, an anti-oxidant, an anti-shrinkage agent, an anti-wrinkle agent, a germicide, a fungicide, a color speckle, a silvercare agent, an anti-tarnish agent, an anti-corrosion agent, an alkalinity source, a solubilizing agent, a carrier, a processing aid, a pigment, a pH control agent, a surfactant, a builder, a chelating agent, a dye transfer inhibiting agent, a deposition aid, a dispersant, a catalytic material, a bleach activator, a bleach booster, a hydrogen peroxide, a source of hydrogen peroxide, a preformed peracid, a polymeric dispersing agent, a clay soil removal/anti-redeposition agent, a structure elasticizing agent, a fabric softener, a carrier, a hydrotrope, a processing aid, a pigment, and combinations thereof. Exemplary adjunct materials and levels of use are found in USPNs 5,576,282; 6,306,812; 6,326,348; 6,610,642; 6,605,458; 5,705,464; 5,710,115; 5,698,504; 5,695,679; 5,686,014 and 5,646,101. In embodiments in which one or more cleaning adjunct material is not compatible with one or more subtilisin variant described herein, methods are employed to keep the adjunct material and variant(s) separated (i.e., not in contact with each other) until combination of the two components is appropriate. Such separation methods include any suitable method known in the art (e.g., gelcaps, encapsulation, tablets, physical separation, etc.).

[00114] Some embodiments are directed to cleaning additive products comprising one or more subtilisin variant described herein. In some embodiments, the additive is packaged in a dosage form for addition to a cleaning process. In some embodiments, the additive is packaged in a dosage form for addition to a cleaning process where a source of peroxygen is employed and increased bleaching effectiveness is desired.

[00115] Exemplary fillers or carriers for granular compositions include, but are not limited to, for example, various salts of sulfate, carbonate and silicate; talc; and clay. Exemplary fillers or carriers for liquid compositions include, but are not limited to, for example, water or low molecular weight primary and secondary alcohols including polyols and diols (e.g., methanol, ethanol, propanol and isopropanol). In some embodiments, the compositions contain from about 5% to about 90% of such filler or carrier. Acidic fillers may be included in such compositions to reduce the pH of the resulting solution in the cleaning method or application.

[00116] In one embodiment, one or more cleaning composition described herein comprises an effective amount of one or more subtilisin variant described herein, alone or in combination with one or more additional enzyme. Typically, a cleaning composition comprises at least about 0.0001 to about 20 wt. %, from about 0.0001 to about 10 wt. %, from about 0.0001 to about 1 wt. %, from about 0.001 to about 1 wt. %, or from about 0.01 to about 0.1 wt. % of one or more protease. In another embodiment, one or more cleaning composition described herein comprises from about 0.01 to about 10 mg, about 0.01 to about 5 mg, about 0.01 to about 2 mg, about 0.01 to about 1 mg, about 0.05 to about 1 mg, about 0.5 to about 10 mg, about 0.5 to about 5 mg, about 0.5 to about 4 mg, about 0.5 to about 4 mg, about 0.5 to about 3 mg, about 0.5 to about 2 mg, about 0.5 to about 1 mg, about 0.1 to about 10 mg, about 0.1 to about 5 mg, about 0.1 to about 4 mg, about 0.1 to about 3 mg, about 0.1 to about 2 mg, about 0.1 to about 2 mg, about 0.1 to about 1 mg, or about 0.1 to about 0.5 mg of one or more protease per gram of composition.

[00117] The cleaning compositions described herein are typically formulated such that during use in aqueous cleaning operations, the wash water will have a pH of from about 4.0 to about 11.5, or even from about 5.0 to about 11.5, or even from about 5.0 to about 8.0, or even from about 7.5 to about 10.5. Liquid product formulations are typically formulated to have a pH from about 3.0 to about 9.0 or even from about 3 to about 5. Granular laundry products are typically formulated to have a pH from about 9 to about 11. In some embodiments, the cleaning compositions of the present invention can be formulated to have an alkaline pH under wash conditions, such as a pH of from about 8.0 to about 12.0, or from about 8.5 to about 11.0, or from about 9.0 to about 11.0. In some embodiments, the cleaning compositions of the present invention can be formulated to have a neutral pH under wash conditions, such as a pH of from about 5.0 to about 8.0, or from about 5.5 to about 8.0, or from about 6.0 to about 8.0, or from about 6.0 to

about 7.5. In some embodiments, the neutral pH conditions can be measured when the cleaning composition is dissolved 1:100 (wt:wt) in de-ionized water at 20°C, measured using a conventional pH meter. Techniques for controlling pH at recommended usage levels include the use of buffers, alkalis, acids, etc., and are well known to those skilled in the art.

[00118] In some embodiments, one or more subtilisin variant described herein is encapsulated to protect it during storage from the other components in the composition and/or control the availability of the variant during cleaning. In some embodiments, encapsulation enhances the performance of the variant and/or additional enzyme. In some embodiments, the encapsulating material typically encapsulates at least part of the subtilisin variant described herein. Typically, the encapsulating material is water-soluble and/or water-dispersible. In some embodiments, the encapsulating material has a glass transition temperature (T_g) of 0°C or higher. Exemplary encapsulating materials include, but are not limited to, carbohydrates, natural or synthetic gums, chitin, chitosan, cellulose and cellulose derivatives, silicates, phosphates, borates, polyvinyl alcohol, polyethylene glycol, paraffin waxes, and combinations thereof. When the encapsulating material is a carbohydrate, it is typically selected from monosaccharides, oligosaccharides, polysaccharides, and combinations thereof. In some embodiments, the encapsulating material is a starch (*See e.g.*, EP0922499, US 4,977,252, US 5,354,559, and US 5,935,826). In some embodiments, the encapsulating material is a microsphere made from plastic such as thermoplastics, acrylonitrile, methacrylonitrile, polyacrylonitrile, polymethacrylonitrile and mixtures thereof. Exemplary commercial microspheres include, but are not limited to EXPANCEL® (Akzo Nobel Chemicals International, B.V.); PM6545, PM6550, PM7220, PM7228, and EXTENDOSPHERES® (Sphere One Inc.); and LUXSIL®, Q-CEL®, and SPHERICEL® (Potters Industries LLC).

[00119] There are a variety of wash conditions including varying detergent formulations, wash water volumes, wash water temperatures, and lengths of wash time to which one or more subtilisin variant described herein may be exposed. A low detergent concentration system is directed to wash water containing less than about 800 ppm detergent components. A medium detergent concentration system is directed to wash containing between about 800 ppm and about 2000 ppm detergent components. A high detergent concentration system is directed to wash water containing greater than about 2000 ppm detergent components. In some embodiments, the “cold water washing” of the present invention utilizes “cold water detergent” suitable for washing at temperatures from about 10°C to about 40°C, from about 20°C to about 30°C, or from about 15°C to about 25°C, as well as all other combinations within the range of about 15°C to about 35°C or 10°C to 40°C.

[00120] Different geographies have different water hardness. Hardness is a measure of the amount of calcium (Ca^{2+}) and magnesium (Mg^{2+}) in the water. Water hardness is usually described in terms of the grains per gallon (gpg) mixed $\text{Ca}^{2+}/\text{Mg}^{2+}$. Most water in the United States is hard, but the degree of

hardness varies. Moderately hard (60-120 ppm) to hard (121-181 ppm) water has 60 to 181 ppm (ppm can be converted to grains per U.S. gallon by dividing ppm by 17.1) of hardness minerals.

Water	Grains per gallon	Parts per million
Soft	less than 1.0	less than 17
Slightly hard	1.0 to 3.5	17 to 60
Moderately hard	3.5 to 7.0	60 to 120
Hard	7.0 to 10.5	120 to 180
Very hard	greater than 10.5	greater than 180

[00121] Other embodiments are directed to one or more cleaning composition comprising from about 0.00001 % to about 10% by weight composition of one or more subtilisin variant described herein and from about 99.999% to about 90.0% by weight composition of one or more adjunct material. In another embodiment, the cleaning composition comprises from about 0.0001 % to about 10%, about 0.001% to about 5%, about 0.001% to about 2%, or about 0.005% to about 0.5% by weight composition of one or more subtilisin variant and from about 99.9999% to about 90.0%, about 99.999 % to about 98%, about 99.995% to about 99.5% by weight composition of one or more adjunct material.

[00122] In other embodiments, the composition described herein comprises one or more subtilisin variant described herein and one or more additional enzyme. The one or more additional enzyme is selected from acyl transferases, alpha-amylases, beta-amylases, alpha-galactosidases, arabinosidases, aryl esterases, beta-galactosidases, carrageenases, catalases, cellobiohydrolases, cellulases, chondroitinases, cutinases, DNase or nuclease, endo-beta-1, 4-glucanases, endo-beta-mannanases, esterases, exo-mannanases, galactanases, glucoamylases, hemicellulases, hyaluronidases, keratinases, laccases, lactases, ligninases, lipases, lipoxygenases, lysozymes, mannanases, metalloproteases, oxidases, pectate lyases, pectin acetyl esterases, pectinases, pentosanases, peroxidases, phenoloxidases, phosphatases, phospholipases, phytases, polygalacturonases, additional proteases, pullulanases, reductases, rhamnogalacturonases, beta-glucanases, tannases, transglutaminases, xylan acetyl-esterases, xylanases, xyloglucanases, xylosidases, and any combination or mixture thereof. Some embodiments are directed to a combination of enzymes (i.e., a “cocktail”) comprising conventional enzymes like amylase, lipase, cutinase and/or cellulase in conjunction with one or more subtilisin variant described herein and/or one or more additional protease.

[00123] In another embodiment, one or more composition described herein comprises one or more subtilisin variant described herein and one or more additional protease. In one embodiment, the additional protease is a serine protease. In another embodiment, the additional protease is an alkaline microbial protease or a trypsin-like protease. Suitable additional proteases include those of animal, vegetable or microbial origin. In some embodiments, the additional protease is a microbial protease. In other embodiments, the additional protease is a chemically or genetically modified mutant. In another

embodiment, the additional protease is an alkaline microbial protease or a trypsin-like protease. Exemplary alkaline proteases include subtilisins derived from, for example, *Bacillus* (e.g., subtilisin, *lentus*, *amyloliquefaciens*, *gibsonii*, subtilisin Carlsberg, subtilisin 309, subtilisin 147 and subtilisin 168). Exemplary metalloproteases include nprE, the recombinant form of neutral metalloprotease expressed in *B. subtilis* (See e.g., WO 07/044993), and PMN, the purified neutral metalloprotease from *B. amyloliquefaciens*. Exemplary additional proteases include but are not limited to those described in WO92/21760, WO95/23221, WO2008010925, WO09149200, WO09149144, WO09149145, WO 10056640, WO10056653, WO20100566356, WO2011072099, WO201113022, WO11140364, WO 2012151534, WO2015038792, WO2015/089441, WO2015089447, WO2015143360, WO2016001449, WO2016001450, WO2016061438, WO2016069544, WO2016069548, WO2016069552, WO 2016069557, WO2016069563, WO2016069569, WO2016087617, WO2016087619, WO2016145428, WO2016174234, WO2016183509, WO2016202835, WO2016205755, US20080090747, US 5,801,039, US 5,340,735, US 5,500,364, US 5,855,625, RE 34,606, US 5,955,340, US 5,700,676 US 6,312,936, US 6,482,628, US 8,530,219, US Provisional Appl Nos. 62/331282, 62/351649, 62/437171, 62/437174, and 62/437509, and PCT Appl Nos. PCT/CN2017/076749 and PCT/US2017/029307, as well as, metalloproteases described in WO1999014341, WO1999033960, WO1999014342, WO1999034003, WO2007044993, WO2009058303, WO 2009058661, WO2014071410, WO2014194032, WO2014194034, WO 2014194054, and WO 2014194117. Exemplary additional proteases also include, but are not limited to trypsin (e.g., of porcine or bovine origin) and the *Fusarium* protease described in WO 89/06270. Exemplary commercial proteases include, but are not limited to MAXATASE, MAXACAL, MAXAPEM, OPTICLEAN®, OPTIMASE®, PROPERASE®, PURAFECT®, PURAFECT® OXP, PURAMAX®, EXCELLASE®, PREFERENZ™ proteases (e.g. P100, P110, P280), EFFECTENZ™ proteases (e.g. P1000, P1050, P2000), EXCELLENZ™ proteases (e.g. P1000), ULTIMASE®, and PURAFAST (Danisco US); ALCALASE®, ALCALASE® ULTRA, BLAZE®, BLAZE® EVITY®, BLAZE® EVITY® 16L, CORONASE®, SAVINASE®, SAVINASE® ULTRA, SAVINASE® EVITY®, SAVINASE® EVERIS®, PRIMASE, DURAZYM, POLARZYME®, OVOZYME®, KANNASE®, LIQUANASE®, EVERIS®, NEUTRASE®, PROGRESS UNO®, RELASE® and ESPERASE® (Novozymes); BLAP™ and BLAP™ variants (Henkel); LAVERGY™ PRO 104 L (BASF), and KAP® (*B. alkalophilus* subtilisin) (Kao).

[00124] Another embodiment is directed to a composition comprising one or more subtilisin variant described herein and one or more lipase. In some embodiments, the composition comprises from about 0.00001 % to about 10%, about 0.0001 % to about 10%, about 0.001% to about 5%, about 0.001% to about 2%, or about 0.005% to about 0.5% lipase by weight composition. An exemplary lipase can be a chemically or genetically modified mutant. Exemplary lipases include, but are not limited to, e.g., those

of bacterial or fungal origin, such as, e.g., *H. lanuginosa* lipase (see, e.g., EP 258068 and EP 305216), *T. lanuginosus* lipase (see, e.g., WO 2014/059360 and WO2015/010009), *Rhizomucor miehei* lipase (see, e.g., EP 238023), *Candida* lipase, such as *C. antarctica* lipase (e.g., *C. antarctica* lipase A or B) (see, e.g., EP 214761), *Pseudomonas* lipases such as *P. alcaligenes* and *P. pseudoalcaligenes* lipase (see, e.g., EP 218272), *P. cepacia* lipase (see, e.g., EP 331376), *P. stutzeri* lipase (see, e.g., GB 1,372,034), *P. fluorescens* lipase, *Bacillus* lipase (e.g., *B. subtilis* lipase (Dartois et al., Biochem. Biophys. Acta 1131:253-260 (1993)), *B. stearoothermophilus* lipase (see, e.g., JP 64/744992), and *B. pumilus* lipase (see, e.g., WO 91/16422)). Exemplary cloned lipases include, but not limited to *Penicillium camembertii* lipase (See, Yamaguchi et al., Gene 103:61-67 (1991)), *Geotricum candidum* lipase (See, Shimada et al., J. Biochem., 106:383-388 (1989)), and various *Rhizopus* lipases, such as, *R. delemar* lipase (See, Hass et al., Gene 109:117-113 (1991)), *R. niveus* lipase (Kugimiya et al., Biosci. Biotech. Biochem. 56:716-719 (1992)) and *R. oryzae* lipase. Other lipolytic enzymes, such as cutinases, may also find use in one or more composition describe herein, including, but not limited to, e.g., cutinase derived from *Pseudomonas mendocina* (see, WO 88/09367) and/or *Fusarium solani pisi* (see, WO90/09446). Exemplary commercial lipases include, but are not limited to M1 LIPASE, LUMA FAST, and LIPOMAX (Genencor); LIPEX[®], LIPOCLEAN[®], LIPOLASE[®] and LIPOLASE[®] ULTRA (Novozymes); and LIPASE P (Amano Pharmaceutical Co. Ltd).

[00125] A still further embodiment is directed to a composition comprising one or more subtilisin variant described herein and one or more amylase. In one embodiment, the composition comprises from about 0.00001 % to about 10%, about 0.0001 % to about 10%, about 0.001% to about 5%, about 0.001% to about 2%, or about 0.005% to about 0.5% amylase by weight composition. Any amylase (e.g., alpha and/or beta) suitable for use in alkaline solutions may be useful to include in such composition. An exemplary amylase can be a chemically or genetically modified mutant. Exemplary amylases include, but are not limited to those of bacterial or fungal origin, such as, for example, amylases described in GB 1,296,839, WO91/00353, WO94/02597, WO94/183314, WO95/10603, WO95/26397, WO95/35382, WO 96/05295, WO96/23873, WO96/23874, WO96/30481, WO97/10342, WO97/41213, WO97/43424, WO 98/13481, WO98/26078, WO99/02702, WO99/09183, WO99/19467, WO99/23211, WO99/29876, WO 99/42567, WO99/43793, WO99/43794, WO99/46399, WO00/29560, WO00/60058, WO00/60059, WO 00/60060, WO01/14532, WO01/34784, WO01/64852, WO01/66712, WO01/88107, WO01/96537, WO 02/092797, WO02/10355, WO02/31124, WO2004055178, WO2004113551, WO2005001064, WO 2005003311, WO2005018336, WO2005019443, WO2005066338, WO2006002643, WO2006012899, WO2006012902, WO2006031554, WO2006063594, WO2006066594, WO2006066596, WO 2006136161, WO2008000825, WO2008088493, WO2008092919, WO2008101894, WO2008112459, WO2009061380, WO2009061381, WO2009100102, WO2009140504, WO2009149419, WO

2010059413, WO2010088447, WO2010091221, WO2010104675, WO2010115021, WO2010115028, WO2010117511, WO2011076123, WO2011076897, WO2011080352, WO2011080353, WO2011080354, WO2011082425, WO2011082429, WO2011087836, WO2011098531, WO2013063460, WO2013184577, WO2014099523, WO2014164777, and WO2015077126. Exemplary commercial amylases include, but are not limited to AMPLIFY[®], AMPLIFY PRIME[®], BAN DURAMYL[®], TERMAMYL[®], TERMAMYL[®] ULTRA, FUNGAMYL[®], STAINZYME[®], STAINZYME[®] PLUS, STAINZYME[®] ULTRA, and STAINZYME[®] EVITY[®] (Novozymes); EFFECTENZ[™] S 1000, POWERASE[®], PREFERENZ[™] S 100, PREFERENZ[™] S 110, EXCELLENZ[™] S 2000, RAPIDASE[®] and MAXAMYL[®] P (Danisco US).

[00126] Yet a still further embodiment is directed to a composition comprising one or more subtilisin variant described herein and one or more cellulase. In one embodiment, the composition comprises from about 0.00001 % to about 10%, 0.0001 % to about 10%, about 0.001% to about 5%, about 0.001% to about 2%, or about 0.005% to about 0.5% cellulase by weight of composition. Any suitable cellulase may find used in a composition described herein. An exemplary cellulase can be a chemically or genetically modified mutant. Exemplary cellulases include but are not limited, to those of bacterial or fungal origin, such as, for example, is described in WO2005054475, WO2005056787, US 7,449,318, US 7,833,773, US 4,435,307; EP 0495257; and US Provisional Appl. Nos. 62/296,678 and 62/435340. Exemplary commercial cellulases include, but are not limited to, CELLUCLEAN[®], CELLUZYME[®], CAREZYME[®], CAREZYME[®] PREMIUM, ENDOLASE[®], and RENOZYME[®] (Novozymes); REVITALENZ[®]100, REVITALENZ[®] 200/220 and REVITALENZ[®] 2000 (Danisco US); and KAC-500(B) (Kao Corporation). In some embodiments, cellulases are incorporated as portions or fragments of mature wild-type or variant cellulases, wherein a portion of the N-terminus is deleted (*see, e.g.*, US 5,874,276).

[00127] An even still further embodiment is directed to a composition comprising one or more subtilisin variant described herein and one or more mannanase. In one embodiment, the composition comprises from about 0.00001 % to about 10%, about 0.0001 % to about 10%, about 0.001% to about 5%, about 0.001% to about 2%, or about 0.005% to about 0.5% mannanase by weight composition. An exemplary mannanase can be a chemically or genetically modified mutant. Exemplary mannanases include, but are not limited to, those of bacterial or fungal origin, such as, for example, as is described in WO2016007929, WO2017/079756 and WO2017/079751; and USPNs 6566114, 6602842, and 6440991. Exemplary commercial mannanases include, but are not limited to MANNAWAY[®] (Novozymes); and EFFECTENZ[™] M 1000, PREFERENZ[®] M 100, MANNASTAR[®], and PURABRITE (Danisco US).

[00128] A yet even still further embodiment is directed to a composition comprising one or more subtilisin variant described herein and one or more peroxidase and/or oxidase enzyme. In one

embodiment, the composition comprises from about 0.00001 % to about 10%, about 0.0001 % to about 10%, about 0.001% to about 5%, about 0.001% to about 2%, or about 0.005% to about 0.5% peroxidase or oxidase by weight composition. A peroxidase may be used in combination with hydrogen peroxide or a source thereof (e.g., a percarbonate, perborate or persulfate) and an oxidase may be used in combination with oxygen. Peroxidases and oxidases are used for “solution bleaching” (i.e., to prevent transfer of a textile dye from a dyed fabric to another fabric when the fabrics are washed together in a wash liquor), alone or in combination with an enhancing agent (*see, e.g.*, WO94/12621 and WO95/01426). An exemplary peroxidase and/or oxidase can be a chemically or genetically modified mutant. Exemplary peroxidases/oxidases include, but are not limited to those of plant, bacterial, or fungal origin.

[00129] Another embodiment is directed to a composition comprising one or more subtilisin variant described herein, and one or more perhydrolase, such as, for example, is described in WO2005/056782, WO2007/106293, WO 2008/063400, WO2008/106214, and WO2008/106215.

[00130] In yet another embodiment, the one or more subtilisin variant described herein and one or more additional enzyme contained in one or more composition described herein may each independently range to about 10%, wherein the balance of the cleaning composition is one or more adjunct material.

[00131] In some embodiments, one or more composition described herein finds use as a detergent additive, wherein said additive is in a solid or liquid form. Such additive products are intended to supplement and/or boost the performance of conventional detergent compositions and can be added at any stage of the cleaning process. In some embodiments, the density of the laundry detergent composition ranges from about 400 to about 1200 g/liter, while in other embodiments it ranges from about 500 to about 950 g/liter of composition measured at 20°C.

[00132] Some embodiments are directed to a laundry detergent composition comprising one or more subtilisin variant described herein and one or more adjunct material selected from surfactants, enzyme stabilizers, builder compounds, polymeric compounds, bleaching agents, additional enzymes, suds suppressors, dispersants, lime-soap dispersants, soil suspension agents, anti-redeposition agents, corrosion inhibitors, and combinations thereof. In some embodiments, the laundry compositions also contain softening agents.

[00133] Further embodiments are directed to manual dishwashing composition comprising one or more subtilisin variant described herein and one or more adjunct material selected from surfactants, organic polymeric compounds, suds enhancing agents, group II metal ions, solvents, hydrotropes, and additional enzymes.

[00134] Other embodiments are directed to one or more composition described herein, wherein said composition is a compact granular fabric cleaning composition that finds use in laundering colored fabrics or provides softening through the wash capacity, or is a heavy duty liquid (HDL) fabric cleaning

composition. Exemplary fabric cleaning compositions and/or processes for making are described in USPNs 6,610,642 and 6,376,450. Other exemplary cleaning compositions are described, for example, in USPNs 6,605,458; 6,294,514; 5,929,022; 5,879,584; 5,691,297; 5,565,145; 5,574,005; 5,569,645; 5,565,422; 5,516,448; 5,489,392; and 5,486,303; 4,968,451; 4,597,898; 4,561,998; 4,550,862; 4,537,706; 4,515,707; and 4,515,705.

[00135] In some embodiments, the cleaning compositions comprise an acidifying particle or an amino carboxylic builder. Examples of an amino carboxylic builder include aminocarboxylic acids, salts and derivatives thereof. In some embodiment, the amino carboxylic builder is an aminopolycarboxylic builder, such as glycine-N,N-diacetic acid or derivative of general formula $\text{MOOC-CHR-N}(\text{CH}_2\text{COOM})_2$ where R is C_{1-12} alkyl and M is alkali metal. In some embodiments, the amino carboxylic builder can be methylglycine diacetic acid (MGDA), GLDA (glutamic-N,N-diacetic acid), iminodisuccinic acid (IDS), carboxymethyl inulin and salts and derivatives thereof, aspartic acid-N-monoacetic acid (ASMA), aspartic acid-N,N-diacetic acid (ASDA), aspartic acid-N-monopropionic acid (ASMP), iminodisuccinic acid (IDA), N-(2-sulfomethyl) aspartic acid (SMAS), N-(2-sulfoethyl)aspartic acid (SEAS), N-(2-sulfomethyl)glutamic acid (SMGL), N-(2-sulfoethyl) glutamic acid (SEGL), IDS (iminodiacetic acid) and salts and derivatives thereof such as N-methyliminodiacetic acid (MIDA), alpha-alanine-N,N-diacetic acid (alpha-ALDA), serine-N,N-diacetic acid (SEDA), isoserine-N,N-diacetic acid (ISDA), phenylalanine-N,N-diacetic acid (PHDA), anthranilic acid-N,N-diacetic acid (ANDA), sulfanilic acid-N,N-diacetic acid (SLDA), taurine-N,N-diacetic acid (TUDA) and sulfomethyl-N,N-diacetic acid (SMDA) and alkali metal salts and derivative thereof. In some embodiments, the acidifying particle has a weight geometric mean particle size of from about 400 μ to about 1200 μ and a bulk density of at least 550 g/L. In some embodiments, the acidifying particle comprises at least about 5% of the builder.

[00136] In some embodiments, the acidifying particle can comprise any acid, including organic acids and mineral acids. Organic acids can have one or two carboxyls and in some instances up to 15 carbons, especially up to 10 carbons, such as formic, acetic, propionic, capric, oxalic, succinic, adipic, maleic, fumaric, sebacic, malic, lactic, glycolic, tartaric and glyoxylic acids. In some embodiments, the acid is citric acid. Mineral acids include hydrochloric and sulphuric acid. In some instances, the acidifying particle is a highly active particle comprising a high level of amino carboxylic builder. Sulphuric acid has also been found to further contribute to the stability of the final particle.

[00137] Additional embodiments are directed to a cleaning composition comprising one or more subtilisin variant and one or more surfactant and/or surfactant system, wherein the surfactant is selected from nonionic surfactants, anionic surfactants, cationic surfactants, ampholytic surfactants, zwitterionic surfactants, semi-polar nonionic surfactants, and mixtures thereof. In some embodiments, the surfactant is present at a level of from about 0.1 to about 60%, while in alternative embodiments the level is from

about 1 to about 50%, while in still further embodiments the level is from about 5 to about 40%, by weight of the cleaning composition.

[00138] In some embodiments, one or more composition described herein comprises one or more detergent builders or builder systems. In one embodiment, the composition comprises from about 1%, from about 0.1% to about 80%, from about 3% to about 60%, from about 5% to about 40%, or from about 10% to about 50% builder by weight composition. Exemplary builders include, but are not limited to alkali metal; ammonium and alkanolammonium salts of polyphosphates; alkali metal silicates; alkaline earth and alkali metal carbonates; aluminosilicates; polycarboxylate compounds; ether hydroxypolycarboxylates; copolymers of maleic anhydride with ethylene or vinyl methyl ether, 1, 3, 5-trihydroxy benzene-2, 4, 6-trisulphonic acid, and carboxymethyloxysuccinic acid; ammonium and substituted ammonium salts of polyacetic acids such as ethylenediamine tetraacetic acid and nitrilotriacetic acid; polycarboxylates such as mellitic acid, succinic acid, citric acid, oxydisuccinic acid, polymaleic acid, benzene 1,3,5-tricarboxylic acid, carboxymethyloxysuccinic acid; and soluble salts thereof. In some such compositions, the builders form water-soluble hardness ion complexes (e.g., sequestering builders), such as citrates and polyphosphates, e.g., sodium tripolyphosphate, sodium tripolyphosphate hexahydrate, potassium tripolyphosphate, and mixed sodium and potassium tripolyphosphate. Exemplary builders are described in, e.g., EP 2100949. In some embodiments, the builders include phosphate builders and non-phosphate builders. In some embodiments, the builder is a phosphate builder. In some embodiments, the builder is a non-phosphate builder. In some embodiments, the builder comprises a mixture of phosphate and non-phosphate builders. Exemplary phosphate builders include, but are not limited to mono-phosphates, di-phosphates, tri-polyphosphates or oligomeric-poylphosphates, including the alkali metal salts of these compounds, including the sodium salts. In some embodiments, a builder can be sodium tripolyphosphate (STPP). Additionally, the composition can comprise carbonate and/or citrate. Other suitable non-phosphate builders include homopolymers and copolymers of polycarboxylic acids and their partially or completely neutralized salts, monomeric polycarboxylic acids and hydroxycarboxylic acids and their salts. In some embodiments, salts of the above mentioned compounds include the ammonium and/or alkali metal salts, i.e. the lithium, sodium, and potassium salts, including sodium salts. Suitable polycarboxylic acids include acyclic, alicyclic, hetero-cyclic and aromatic carboxylic acids, wherein in some embodiments, they can contain at least two carboxyl groups which are in each case separated from one another by, in some instances, no more than two carbon atoms.

[00139] In some embodiments, one or more composition described herein comprises one or more chelating agent. In one embodiment, the composition comprises from about 0.1% to about 15% or about 3% to about 10% chelating agent by weight composition. Exemplary chelating agents include, but are not

limited to, e.g., copper, iron, manganese, and mixtures thereof.

[00140] In some embodiments, one or more composition described herein comprises one or more deposition aid. Exemplary deposition aids include, but are not limited to, e.g., polyethylene glycol; polypropylene glycol; polycarboxylate; soil release polymers, such as, e.g., polytelephthalic acid; clays such as, e.g., kaolinite, montmorillonite, atapulgite, illite, bentonite, and halloysite; and mixtures thereof.

[00141] In other embodiments, one or more composition described herein comprises one or more anti-redeposition agent or non-ionic surfactant (which can prevent the re-deposition of soils) (*see, e.g.*, EP 2100949). For example, in ADW compositions, non-ionic surfactants find use for surface modification purposes, in particular for sheeting, to avoid filming and spotting and to improve shine. These non-ionic surfactants also find use in preventing the re-deposition of soils. In some embodiments, the non-ionic surfactant can be ethoxylated nonionic surfactants, epoxy-capped poly(oxyalkylated) alcohols and amine oxides surfactants.

[00142] In some embodiments, one or more composition described herein comprises one or more dye transfer inhibiting agent. Exemplary polymeric dye transfer inhibiting agents include, but are not limited to, polyvinylpyrrolidone polymers, polyamine N-oxide polymers, copolymers of N-vinylpyrrolidone and N-vinylimidazole, polyvinylloxazolidones, polyvinylimidazoles, and mixtures thereof. In one embodiment, the composition comprises from about 0.0001% to about 10%, about 0.01% to about 5%, or about 0.1% to about 3% dye transfer inhibiting agent by weight composition.

[00143] In some embodiments, one or more composition described herein comprises one or more silicate. Exemplary silicates include, but are not limited to, sodium silicates, e.g., sodium disilicate, sodium metasilicate, and crystalline phyllosilicates. In some embodiments, silicates are present at a level of from about 1% to about 20% or about 5% to about 15% by weight of the composition.

[00144] In some still additional embodiments, one or more composition described herein comprises one or more dispersant. Exemplary water-soluble organic materials include, but are not limited to, e.g., homo- or co-polymeric acids or their salts, in which the polycarboxylic acid comprises at least two carboxyl radicals separated from each other by not more than two carbon atoms.

[00145] In some further embodiments, one or more composition described herein comprises one or more enzyme stabilizer. In some embodiments, the enzyme stabilizer is water-soluble sources of calcium and/or magnesium ions. In some embodiments, the enzyme stabilizers include oligosaccharides, polysaccharides, and inorganic divalent metal salts, including alkaline earth metals, such as calcium salts. In some embodiments, the enzymes employed herein are stabilized by the presence of water-soluble sources of zinc (II), calcium (II) and/or magnesium (II) ions in the finished compositions that provide such ions to the enzymes, as well as other metal ions (e.g., barium (II), scandium (II), iron (II), manganese (II), aluminum (III), tin (II), cobalt (II), copper (II), nickel (II), and oxovanadium (IV)).

Chlorides and sulfates also find use in some embodiments. Exemplary oligosaccharides and polysaccharides (e.g., dextrans) are described, for example, in WO 07/145964. In some embodiments, reversible protease inhibitors also find use, such as boron-containing compounds (e.g., borate, 4-formyl phenyl boronic acid, and phenyl-boronic acid derivatives (such for example, those described in WO96/41859) and/or a peptide aldehyde, such as, for example, is further described in WO2009/118375 and WO2013004636.

[00146] In some embodiments, one or more composition described herein comprises one or more bleach, bleach activator, and/or bleach catalyst. In some embodiments, one or more composition described herein comprises one or more inorganic and/or organic bleaching compound. Exemplary inorganic bleaches include, but are not limited to perhydrate salts, e.g., perborate, percarbonate, perphosphate, persulfate, and persilicate salts. In some embodiments, inorganic perhydrate salts are alkali metal salts. In some embodiments, inorganic perhydrate salts are included as the crystalline solid, without additional protection, although in some other embodiments, the salt is coated. Bleach activators are typically organic peracid precursors that enhance the bleaching action in the course of cleaning at temperatures of 60°C and below. Exemplary bleach activators include compounds which, under perhydrolysis conditions, give aliphatic peroxycarboxylic acids having from about 1 to about 10 carbon atoms or about 2 to about 4 carbon atoms, and/or optionally substituted perbenzoic acid. Exemplary bleach activators are described, for example, in EP 2100949. Exemplary bleach catalysts include, but are not limited to, manganese triazacyclononane and related complexes, as well as cobalt, copper, manganese, and iron complexes. Additional exemplary bleach catalysts are described, for example, in US 4,246,612; US 5,227,084; US 4,810,410; WO 99/06521; and EP 2100949.

[00147] In some embodiments, one or more composition described herein comprises one or more catalytic metal complexes. In some embodiments, a metal-containing bleach catalyst finds use. In some embodiments, the metal bleach catalyst comprises a catalyst system comprising a transition metal cation of defined bleach catalytic activity (e.g., copper, iron, titanium, ruthenium, tungsten, molybdenum, or manganese cations), an auxiliary metal cation having little or no bleach catalytic activity (e.g., zinc or aluminum cations), and a sequester having defined stability constants for the catalytic and auxiliary metal cations, particularly ethylenediaminetetraacetic acid, ethylenediaminetetra (methylenephosphonic acid) and water-soluble salts thereof (*see, e.g.*, US 4,430,243). In some embodiments, one or more composition described herein is catalyzed by means of a manganese compound. Such compounds and levels of use are described, for example, in US 5,576,282. In additional embodiments, cobalt bleach catalysts find use and are included in one or more composition described herein. Various cobalt bleach catalysts are described, for example, in USPNs 5,597,936 and 5,595,967.

[00148] In some additional embodiments, one or more composition described herein includes a

transition metal complex of a macropolycyclic rigid ligand (MRL). As a practical matter, and not by way of limitation, in some embodiments, the compositions and cleaning processes described herein are adjusted to provide on the order of at least one part per hundred million, from about 0.005 ppm to about 25 ppm, about 0.05 ppm to about 10 ppm, or about 0.1 ppm to about 5 ppm of active MRL in the wash liquor. Exemplary MRLs include, but are not limited to special ultra-rigid ligands that are cross-bridged, such as, e.g., 5,12-diethyl-1,5,8,12-tetraazabicyclo (6.6.2) hexadecane. Exemplary metal MRLs are described, for example, in WO 2000/32601 and US 6,225,464.

[00149] In another embodiment, one or more composition described herein comprises one or more metal care agent. In some embodiments, the composition comprises from about 0.1% to about 5% metal care agent by weight composition. Exemplary metal care agents include, for example, aluminum, stainless steel, and non-ferrous metals (e.g., silver and copper). Additional exemplary metal care agents are described, for example, in EP 2100949, WO 94/26860, and WO 94/26859. In some compositions, the metal care agent is a zinc salt.

[00150] In some embodiments, the cleaning composition is a high density liquid (HDL) composition comprising one or more subtilisin variant described herein. The HDL liquid laundry detergent can comprise a deterative surfactant (10-40%) comprising anionic deterative surfactant selected from a group of linear or branched or random chain, substituted or unsubstituted alkyl sulphates, alkyl sulphonates, alkyl alkoxyated sulphate, alkyl phosphates, alkyl phosphonates, alkyl carboxylates, and/or mixtures thereof; and optionally non-ionic surfactant selected from a group of linear or branched or random chain, substituted or unsubstituted alkyl alkoxyated alcohol, for example, a C₈-C₁₈alkyl ethoxyated alcohol and/or C₆-C₁₂alkyl phenol alkoxyates, optionally wherein the weight ratio of anionic deterative surfactant (with a hydrophilic index (Hlc) of from 6.0 to 9) to non-ionic deterative surfactant is greater than 1:1. Suitable deterative surfactants also include cationic deterative surfactants (selected from alkyl pyridinium compounds, alkyl quaternary ammonium compounds, alkyl quaternary phosphonium compounds, alkyl ternary sulphonium compounds, and/or mixtures thereof); zwitterionic and/or amphoteric deterative surfactants (selected from alkanolamine sulfo-betaines); ampholytic surfactants; semi-polar non-ionic surfactants; and mixtures thereof.

[00151] In another embodiment, the cleaning composition is a liquid or gel detergent, which is not unit dosed, that may be aqueous, typically containing at least 20% and up to 95% water by weight, such as up to about 70% water by weight, up to about 65% water by weight, up to about 55% water by weight, up to about 45% water by weight, or up to about 35% water by weight. Other types of liquids, including without limitation, alkanols, amines, diols, ethers and polyols may be included in an aqueous liquid or gel. An aqueous liquid or gel detergent may contain from 0-30% organic solvent. A liquid or gel detergent may be non-aqueous.

[00152] The composition can comprise optionally, a surfactancy boosting polymer consisting of amphiphilic alkoxyated grease cleaning polymers selected from a group of alkoxyated polymers having branched hydrophilic and hydrophobic properties, such as alkoxyated polyalkylenimines in the range of 0.05wt%-10wt% and/or random graft polymers typically comprising a hydrophilic backbone comprising monomers selected from the group consisting of: unsaturated C₁-C₆carboxylic acids, ethers, alcohols, aldehydes, ketones, esters, sugar units, alkoxy units, maleic anhydride, saturated polyalcohols such as glycerol, and mixtures thereof; and hydrophobic side chain(s) selected from the group consisting of: C₄-C₂₅alkyl group, polypropylene, polybutylene, vinyl ester of a saturated C₂-C₆mono-carboxylic acid, C₁-C₆alkyl ester of acrylic or methacrylic acid, and mixtures thereof.

[00153] The composition can comprise additional polymers such as soil release polymers including, for example, anionically end-capped polyesters, for example SRP1; polymers comprising at least one monomer unit selected from saccharide, dicarboxylic acid, polyol and combinations thereof, in random or block configuration; ethylene terephthalate-based polymers and co-polymers thereof in random or block configuration, for example, Repel-o-tex SF, SF-2 and SRP6, Texcare SRA100, SRA300, SRN100, SRN170, SRN240, SRN300 and SRN325, Marloquest SL; anti-redeposition polymers (0.1 wt% to 10wt%, including, for example, carboxylate polymers, such as polymers comprising at least one monomer selected from acrylic acid, maleic acid (or maleic anhydride), fumaric acid, itaconic acid, aconitic acid, mesaconic acid, citraconic acid, methylenemalononic acid, and any mixture thereof; vinylpyrrolidone homopolymer; and/or polyethylene glycol with a molecular weight in the range of from 500 to 100,000 Da); cellulosic polymer (including, for example, alkyl cellulose; alkyl alkoxyalkyl cellulose; carboxyalkyl cellulose; alkyl carboxyalkyl cellulose, examples of which include carboxymethyl cellulose, methyl cellulose, methyl hydroxyethyl cellulose, methyl carboxymethyl cellulose; and mixtures thereof); and polymeric carboxylate (such as, for example, maleate/acrylate random copolymer or polyacrylate homopolymer).

[00154] The composition can further comprise saturated or unsaturated fatty acid, preferably saturated or unsaturated C₁₂-C₂₄fatty acid (0-10 wt%); deposition aids (including, for example, polysaccharides, cellulosic polymers, polydiallyl dimethyl ammonium halides (DADMAC), and co-polymers of DADMAC with vinyl pyrrolidone, acrylamides, imidazoles, imidazolinium halides, and mixtures thereof, in random or block configuration; cationic guar gum; cationic cellulose such as cationic hydroxyethyl cellulose; cationic starch; cationic polyacrylamides; and mixtures thereof.

[00155] The composition can further comprise dye transfer inhibiting agents examples of which include manganese phthalocyanine, peroxidases, polyvinylpyrrolidone polymers, polyamine N-oxide polymers, copolymers of N-vinylpyrrolidone and N-vinylimidazole, polyvinylloxazolidones and polyvinylimidazoles and/or mixtures thereof; chelating agents examples of which include ethylene-

diamine-tetraacetic acid (EDTA); diethylene triamine penta methylene phosphonic acid (DTPMP); hydroxy-ethane diphosphonic acid (HEDP); ethylenediamine N,N'-disuccinic acid (EDDS); methyl glycine diacetic acid (MGDA); diethylene triamine penta acetic acid (DTPA); propylene diamine tetracetic acid (PDT A); 2-hydroxypyridine-N-oxide (HPNO); or methyl glycine diacetic acid (MGDA); glutamic acid N,N-diacetic acid (N,N-dicarboxymethyl glutamic acid tetrasodium salt (GLDA); nitrilotriacetic acid (NTA); 4,5-dihydroxy-m-benzenedisulfonic acid; citric acid and any salts thereof; N-hydroxyethylethylenediaminetri-acetic acid (HEDTA), triethylenetetraaminehexaacetic acid (TTHA), N-hydroxyethyliminodiacetic acid (HEIDA), dihydroxyethylglycine (DHEG), ethylenediaminetetrapropionic acid (EDTP), and derivatives thereof.

[00156] The composition can further comprise silicone or fatty-acid based suds suppressors; an enzyme stabilizer; hueing dyes, calcium and magnesium cations, visual signaling ingredients, anti-foam (0.001 to about 4.0 wt.%), and/or structurant/thickener (0.01-5 wt.%) selected from the group consisting of diglycerides, triglycerides, ethylene glycol distearate, microcrystalline cellulose, cellulose based materials, microfiber cellulose, biopolymers, xanthan gum, gellan gum, and mixtures thereof.

[00157] In some embodiments, the cleaning composition is a high density powder (HDD) composition comprising one or more subtilisin variant described herein. The HDD powder laundry detergent can comprise a deterative surfactant including anionic deterative surfactants (selected from linear or branched or random chain, substituted or unsubstituted alkyl sulphates, alkyl sulphonates, alkyl alkoxyated sulphate, alkyl phosphates, alkyl phosphonates, alkyl carboxylates and/or mixtures thereof), non-ionic deterative surfactant (selected from 1 linear or branched or random chain, substituted or unsubstituted C₈-C₁₈ alkyl ethoxylates, and/or C₆-C₁₂ alkyl phenol alkoxyates), cationic deterative surfactants (selected from alkyl pyridinium compounds, alkyl quaternary ammonium compounds, alkyl quaternary phosphonium compounds, alkyl ternary sulphonium compounds, and mixtures thereof); zwitterionic and/or amphoteric deterative surfactants (selected from alkanolamine sulfo-betaines); ampholytic surfactants; semi-polar non-ionic surfactants and mixtures thereof; builders (phosphate free builders, e.g., zeolite builders examples of which include zeolite A, zeolite X, zeolite P and zeolite MAP in the range of 0 to less than 10 wt.%); phosphate builders, e.g., sodium tri-polyphosphate in the range of 0 to less than 10 wt.%; citric acid, citrate salts and nitrilotriacetic acid or salt thereof in the range of less than 15 wt.%; silicate salt (sodium or potassium silicate or sodium meta-silicate in the range of 0 to less than 10 wt.% or layered silicate (SKS-6)); carbonate salt (sodium carbonate and/or sodium bicarbonate in the range of 0 to less than 10 wt.%); and bleaching agents (photobleaches, e.g., sulfonated zinc phthalocyanines, sulfonated aluminum phthalocyanines, xanthenes dyes, and mixtures thereof); hydrophobic or hydrophilic bleach activators (e.g., dodecanoyl oxybenzene sulfonate, decanoyl oxybenzene sulfonate, decanoyl oxybenzoic acid or salts thereof, 3,5,5-trimethy hexanoyl oxybenzene

sulfonate, tetraacetyl ethylene diamine-TAED, and nonanoyloxybenzene sulfonate-NOBS, nitrile quats, and mixtures thereof); hydrogen peroxide; sources of hydrogen peroxide (inorganic perhydrate salts, e.g., mono or tetra hydrate sodium salt of perborate, percarbonate, persulfate, perphosphate, or persilicate); preformed hydrophilic and/or hydrophobic peracids (selected from percarboxylic acids and salts, percarbonic acids and salts, perimidic acids and salts, peroxymonosulfuric acids and salts, and mixtures thereof); and/or bleach catalyst (e.g., imine bleach boosters, such as iminium cations and polyions; iminium zwitterions; modified amines; modified amine oxides; N-sulphonyl imines; N-phosphonyl imines; N-acyl imines; thiadiazole dioxides; perfluoroimines; cyclic sugar ketones and mixtures thereof), metal-containing bleach catalyst (e.g., copper, iron, titanium, ruthenium, tungsten, molybdenum, or manganese cations along with an auxiliary metal cations such as zinc or aluminum and a sequesterant such as ethylenediaminetetraacetic acid, ethylenediaminetetra(methylenephosphonic acid) and water-soluble salts thereof).

[00158] The composition can further comprise additional detergent ingredients including perfume microcapsules, starch encapsulated perfume accord, an enzyme stabilizer, hueing agents, additional polymers including fabric integrity and cationic polymers, dye lock ingredients, fabric-softening agents, brighteners (for example C.I. Fluorescent brighteners), flocculating agents, chelating agents, alkoxylated polyamines, fabric deposition aids, and/or cyclodextrin.

[00159] In some embodiments, the cleaning composition is an ADW detergent composition comprising one or more subtilisin variant described herein. The ADW detergent composition can comprise two or more non-ionic surfactants selected from ethoxylated non-ionic surfactants, alcohol alkoxylated surfactants, epoxy-capped poly(oxyalkylated) alcohols, and amine oxide surfactants present in amounts from 0-10% by wt.; builders in the range of 5-60% by wt. comprising either phosphate (mono-phosphates, di-phosphates, tri-polyphosphates or oligomeric-polyphosphates), sodium tripolyphosphate-STPP or phosphate-free builders (amino acid based compounds, e.g., MGDA (methyl-glycine-diacetic acid) and salts and derivatives thereof, GLDA (glutamic-N,N-diacetic acid) and salts and derivatives thereof, IDS (iminodisuccinic acid) and salts and derivatives thereof, carboxy methyl inulin and salts and derivatives thereof and mixtures thereof, nitrilotriacetic acid (NTA), diethylene triamine penta acetic acid (DTPA), and B-alaninediacetic acid (B-ADA) and their salts), homopolymers and copolymers of polycarboxylic acids and their partially or completely neutralized salts, monomeric polycarboxylic acids and hydroxycarboxylic acids and their salts in the range of 0.5-50% by wt; sulfonated/carboxylated polymers (provide dimensional stability to the product) in the range of about 0.1 to about 50% by wt; drying aids in the range of about 0.1 to about 10% by wt (selected from polyesters, especially anionic polyesters optionally together with further monomers with 3- 6 functionalities which are conducive to polycondensation, specifically acid, alcohol or ester functionalities, polycarbonate-, polyurethane- and/or

polyurea-polyorganosiloxane compounds or precursor compounds thereof of the reactive cyclic carbonate and urea type); silicates in the range from about 1 to about 20% by wt (sodium or potassium silicates, e.g., sodium disilicate, sodium meta-silicate and crystalline phyllosilicates); bleach-inorganic (e.g., perhydrate salts such as perborate, percarbonate, perphosphate, persulfate and persilicate salts) and organic (e.g., organic peroxyacids including diacyl and tetraacylperoxides, especially diperoxydodecanedioic acid, diperoxytetradecanedioic acid, and diperoxyhexadecanedioic acid); bleach activator-organic peracid precursors in the range from about 0.1 to about 10% by wt.; bleach catalysts (selected from manganese triazaacyclononane and related complexes, Co, Cu, Mn and Fe bispyridylamine and related complexes, and pentamine acetate cobalt(III) and related complexes); metal care agents in the range from about 0.1-5% by wt (selected from benzotriazoles, metal salts and complexes, and silicates); enzymes in the range from about 0.01-5.0mg of active enzyme per gram of ADW detergent composition (acyl transferases, alpha-amylases, beta-amylases, alpha-galactosidases, arabinosidases, aryl esterases, beta-galactosidases, carrageenases, catalases, cellobiohydrolases, cellulases, chondroitinases, cutinases, endo-beta-1, 4-glucanases, endo-beta-mannanases, esterases, exo-mannanases, galactanases, glucoamylases, hemicellulases, hyaluronidases, keratinases, laccases, lactases, ligninases, lipases, lipoxxygenases, mannanases, oxidases, pectate lyases, pectin acetyl esterases, pectinases, pentosanases, peroxidases, phenoloxidases, phosphatases, phospholipases, phytases, polygalacturonases, proteases, pullulanases, reductases, rhamnogalacturonases, beta-glucanases, tannases, transglutaminases, xylan acetyl-esterases, xylanases, xyloglucanases, xylosidases, and mixtures thereof); and enzyme stabilizer components (selected from oligosaccharides, polysaccharides and inorganic divalent metal salts).

[00160] More embodiments are directed to compositions and methods of treating fabrics (*e.g.*, to desize a textile) using one or more subtilisin variant described herein. Fabric-treating methods are well known in the art (*see, e.g.*, US 6,077,316). For example, the feel and appearance of a fabric can be improved by a method comprising contacting the fabric with a variant described herein in a solution. The fabric can be treated with the solution under pressure.

[00161] One or more subtilisin variant described herein can be applied during or after weaving a textile, during the desizing stage, or one or more additional fabric processing steps. During the weaving of textiles, the threads are exposed to considerable mechanical strain. Prior to weaving on mechanical looms, warp yarns are often coated with sizing starch or starch derivatives to increase their tensile strength and to prevent breaking. One or more subtilisin variant described herein can be applied during or after weaving to remove the sizing starch or starch derivatives. After weaving, the variant can be used to remove the size coating before further processing the fabric to ensure a homogeneous and wash-proof result. One or more subtilisin variant described herein can be used alone or with other desizing chemical reagents and/or desizing enzymes to desize fabrics, including cotton-containing fabrics, as detergent

additives, *e.g.*, in aqueous compositions. An amylase also can be used in compositions and methods for producing a stonewashed look on indigo-dyed denim fabric and garments. For the manufacture of clothes, the fabric can be cut and sewn into clothes or garments, which are afterwards finished. In particular, for the manufacture of denim jeans, different enzymatic finishing methods have been developed. The finishing of denim garment normally is initiated with an enzymatic desizing step, during which garments are subjected to the action of proteolytic enzymes to provide softness to the fabric and make the cotton more accessible to the subsequent enzymatic finishing steps. One or more subtilisin variant described herein can be used in methods of finishing denim garments (*e.g.*, a “bio-stoning process”), enzymatic desizing and providing softness to fabrics, and/or finishing process.

[00162] In further embodiments, one or more subtilisin variant described herein can be used in industrial and institutional cleaning applications, such as, for example, in an industrial or institutional warewashing system, machine, or process; an industrial or institutional on premises laundry system, machine or process; and an industrial or institutional laundry system, machine or process.

[00163] One or more subtilisin variant described herein can be used to remove proteins from animals and their subsequent degradation or disposal, such as, *e.g.*, feathers, skin, hair, and hide. In some instances, immersion of the animal carcass in a solution comprising one or more subtilisin variant described herein can act to protect the skin from damage in comparison to the traditional immersion in scalding water or the defeathering process. In one embodiment, feathers can be sprayed with one or more subtilisin variant described herein under conditions suitable for digesting or initiating degradation of the plumage. In some embodiments, the variant can be used in combination with an oxidizing agent.

[00164] In some embodiments, the removal of the oil or fat associated with raw feathers can be assisted by one or more subtilisin variant described herein. In some embodiments, one or more subtilisin variant described herein is used in compositions for cleaning the feathers as well as to sanitize and partially dehydrate the fibers. In yet other embodiments, one or more subtilisin variant described herein finds use in recovering protein from plumage. In some other embodiments, one or more subtilisin variant described herein is applied in a wash solution in combination with 95% ethanol or other polar organic solvent with or without a surfactant at about 0.5% (v/v). In other embodiments, one or more subtilisin variant described herein may be used alone or in combination in suitable feather processing and proteolytic methods, such as those disclosed in PCT/EP2013/065362, PCT/EP2013/065363, and PCT/EP2013/065364. In some embodiments, the recovered protein can be subsequently used in animal or fish feed.

[00165] In still another embodiment, one or more animal feed composition, animal feed additive and/or pet food comprises one or more subtilisin variant described herein. Other embodiments are directed to methods for preparing such an animal feed composition, animal feed additive composition

and/or pet food comprising mixing one or more subtilisin variant described herein with one or more animal feed ingredients and/or animal feed additive ingredients and/or pet food ingredients.

[00166] The term “animal” includes all non-ruminant and ruminant animals. In a particular embodiment, the animal is a non-ruminant animal, such as a horse and a mono-gastric animal. Examples of mono-gastric animals include, but are not limited to, pigs and swine, such as piglets, growing pigs, sows; poultry such as turkeys, ducks, chicken, broiler chicks, layers; fish such as salmon, trout, tilapia, catfish and carps; and crustaceans such as shrimps and prawns. In a further embodiment, the animal is a ruminant animal including, but not limited to, cattle, young calves, goats, sheep, giraffes, bison, moose, elk, yaks, water buffalo, deer, camels, alpacas, llamas, antelope, pronghorn and nilgai.

[00167] In the present context, it is intended that the term “pet food” is understood to mean a food for a household animal such as, but not limited to, dogs, cats, gerbils, hamsters, chinchillas, fancy rats, guinea pigs; avian pets, such as canaries, parakeets, and parrots; reptile pets, such as turtles, lizards and snakes; and aquatic pets, such as tropical fish and frogs.

[00168] The terms “animal feed composition,” “feedstuff” and “fodder” are used interchangeably and can comprise one or more feed materials selected from a) cereals, such as small grains (e.g., wheat, barley, rye, oats and combinations thereof) and/or large grains such as maize or sorghum; b) by products from cereals, such as corn gluten meal, Distillers Dried Grain Solubles (DDGS) (particularly corn based Distillers Dried Grain Solubles (cDDGS), wheat bran, wheat middlings, wheat shorts, rice bran, rice hulls, oat hulls, palm kernel, and citrus pulp; c) protein obtained from sources such as soya, sunflower, peanut, lupin, peas, fava beans, cotton, canola, fish meal, dried plasma protein, meat and bone meal, potato protein, whey, copra, sesame; d) oils and fats obtained from vegetable and animal sources; and e) minerals and vitamins.

[00169] One or more subtilisin variant described herein finds further use in the enzyme aided bleaching of paper pulps such as chemical pulps, semi-chemical pulps, kraft pulps, mechanical pulps or pulps prepared by the sulfite method. In general terms, paper pulps are incubated with one or more subtilisin variant described herein under conditions suitable for bleaching the paper pulp.

[00170] In some embodiments, the pulps are chlorine free pulps bleached with oxygen, ozone, peroxide or peroxyacids. In some embodiments, one or more subtilisin variant described herein is used in enzyme aided bleaching of pulps produced by modified or continuous pulping methods that exhibit low lignin contents. In some other embodiments, one or more subtilisin variant described herein is applied alone or preferably in combination with xylanase and/or endoglucanase and/or alpha-galactosidase and/or cellobiohydrolase enzymes.

[00171] In other embodiments, one or more subtilisin variant described herein finds further use in the enzyme aided debridement of tissue. This involves the removal of dead or damaged tissue, for example, removal from wounds to aid in healing.

[00172] In even further embodiments, one or more subtilisin variant described herein finds further use in tissue culture. In particular, one or more subtilisin variant described herein can be used to suspend or resuspend cells adherent to a cell culture wall, such as during the process of harvesting cells. In another embodiment, one or more subtilisin variant described herein can be used to cleave protein bonds between cultured cells and the dish, allowing cells to become suspended in solution.

[00173] In yet another embodiment, one or more subtilisin variant described herein finds further use as a food additive, a digestive aide, and/or a food processing aid.

[00174] In still yet another embodiment, one or more subtilisin variant described herein finds further use in leather processing by removing hair from animal hides, soaking, degreasing, or bating, which is a process involving degradation of non-structural proteins during leather making.

EXAMPLE 1

Assays

[00175] The following assays were used in the examples that follow. Any deviations from the assays described herein below will be addressed in the example in which such deviation arises.

Performance Index

[00176] The performance index (PI) of an enzyme compares the performance of a variant (measured value) with a Parent enzyme selected from AprL wild-type (WT) (SEQ ID NO:2) (hereinafter “AprL Parent”), BliD02339 WT (SEQ ID NO:9) (hereinafter “BliD Parent”), or SQCBV35 variant (SEQ ID NO:63) (hereinafter SQCBV35 Parent) (theoretical value or measured value) at the same protein concentration. Theoretical concentrations for the Parent can be calculated using the parameters extracted from a Langmuir fit of a standard curve of the Parent. A PI that is greater than 1 ($PI > 1$) indicates improved performance by a variant as compared to Parent, while a PI of 1 ($PI = 1$) identifies a variant that performs the same as Parent, and a PI that is less than 1 ($PI < 1$) identifies a variant that performs worse than Parent.

Protein Determination Assay

[00177] Protein determination of AprL or BliD Variants from culture supernatant was performed with an Agilent U-HPLC or HPLC systems. For U-HPLC method, a calibration curve (0-500 ppm) of purified Parent protein was prepared and used to generate a standard curve. AprL or BliD Variants from clarified culture supernatants were diluted 10 fold in the dilution buffer (Tris 25mM, pH 7.4, 5mM $CaCl_2$) and then mixed in a 1:1 ratio with an Acetonitrile buffer (22.5mM Tris, pH 7.4, 4.5mM $CaCl_2$, 9% acetonitrile). Afterwards, the samples were filtered using a 45 μ m filterplate and loaded via an auto-

sampler onto a reverse phase column (Zorbax 300 SB-C3 column, 2.1x100 mm & 2.1x50 mm, both with 1.8 μ m beadsizes). The samples were eluted from the column with a gradient of Buffer A (0.1% Trifluoroacetic acid (TFA)) and Buffer B (0.07% Acetonitrile). The flow rate was 1 mL/min with a 4 min run and a 1 min post run column equilibration. Absorbance was measured at 220 nm, and peaks were integrated using ChemStation software (Agilent Technologies). The protein concentration of the samples was calculated based on a standard curve of the purified Parent. In some instances, the Bradford method for protein determination was used alongside with HPLC. The Bradford reagent (Bradford Quickstart 1X dye reagent, BioRad) was used and absorbance at 595/450 nm was measured on a Spectramax plate reader.

[00178] Alternately, protein determination of AprL variants from culture supernatant was performed by HPLC, using an Agilent 1100/1200 system. A calibration curve (0-1500 ppm) using purified Parent protein was prepared and used to generate a standard curve. AprL Variants from clarified culture supernatants were analysed by one of two methods. In some cases, the culture supernatants were diluted two fold in the dilution buffer (Tris 100mM, pH 8.6, 0.005% Tween 80) and loaded via an auto-sampler onto an Agilent C8 column (Agilent Poroshell 5 μ m 300SB-C8 2.1 x 75mm) pre-equilibrated at 69-70°C. The samples were eluted from the column with a gradient of Buffer A (0.1% TFA) and Buffer B (100% Acetonitrile and 0.1% TFA). The flow rate was 2 mL/min with a 2 min run. In other cases, clarified culture supernatants containing AprL Variants were pre-treated in an acid/SDS solution. 75 μ L samples were mixed with 6.4 μ L of a 4NHCl and mixed for 10 minutes on ice. 20.4 μ L 10% SDS solution was added per well and mixed, samples were filtered prior to loading into HPLC HIP-ALS Aquity 125A USEC column set at room temperature. The samples were eluted with a solution of 25 mM Na-acetate, pH 5.5, 100 mM NaCl, 0.5% SDS at a flow rate was 0.35 mL/min with an 8 min run. Absorbance was measured at 220 or 230 nm, and peaks were integrated using ChemStation (Agilent Technologies) software.

[00179] For all three protein determination methods, the protein concentration of each sample was calculated based on a standard curve of the purified Parent enzyme.

Protease Activity

[00180] The protease activity of Parent and variants thereof was tested by measuring the hydrolysis of N-suc-AAPF-pNA or dimethylcasein (DMC) substrates.

[00181] For the AAPF assay, the reagent solutions used were: 100 mM Tris pH 8.6, 10 mM CaCl_2 , 0.005% Tween®-80 (Tris/Ca buffer) and 160 mM suc-AAPF-pNA in DMSO (suc-AAPF-pNA stock solution) (Sigma: S-7388). To prepare a working solution, 1 mL suc-AAPF-pNA stock solution was added to 100 mL Tris/Ca buffer and mixed. An enzyme sample was added to a microtiter plate (MTP) containing 1 mg/mL suc-AAPF-pNA working solution and assayed for activity at 405 nm over 3-5 min

using a SpectraMax plate reader in kinetic mode at RT. The protease activity was expressed as mOD/min.

[00182] For the DMC assay, the reagent solutions used were: 2.5% DMC (Sigma) in 100 mM Sodium Carbonate pH 9.5, 0.075% TNBSA (2,4,6-trinitrobenzene sulfonic acid, Thermo Scientific) in Reagent A. Reagent A: 45.4 g $\text{Na}_2\text{B}_4\text{O}_7 \cdot 10\text{H}_2\text{O}$ (Merck) in 15 mL 4N NaOH to reach a final volume of 1000 mL in MQ water, Dilution Solution: 10 mM NaCl, 0.1 mM CaCl_2 , 0.005% Tween-80, 0.02% Na-azide. MTPs were filled with 47.5 μL DMC substrate following the addition of 2.5 μL of 20 ppm protease supernatant. 50 μL of TNBSA in Reagent A was then added with slow mixing. Activity was measured at 405 nm over 5 min using a SpectraMax plate reader in kinetic mode at RT. As stated above for the AAPF assay, activity was expressed as mOD/min.

Detergent compositions evaluated

[00183] Detergent formulas used in these studies are listed in Table 1, and include heavy duty liquid laundry (HDL), heavy duty power laundry (HDD), and automatic dishwashing (ADW) formulas. Commercial detergents were heat-inactivated to remove enzyme activity and dosed as described in Table 1. Detergent treatments for enzyme inactivation were as follows. HDL laundry detergents that contain enzymes (excludes Test HDL 1, Test HDL 2, and Persil Non-Bio HDL detergents) were inactivated by heating the neat liquid at 95°C for 16 hours in a water bath. Enzymes in HDD detergents were inactivated by preparing a 10X concentrated solution relative to what is used in the final cleaning assay and heating for 16 hours at 95°C. Commercial ADW detergent formulas were heat inactivated for 30-45 min at 65°C as follows. An ADW detergent tablet was dissolved in 700 mL 21GH water and enzyme activity was inactivated by incubating the solution for 30-45 min at 65°C. The enzyme inactivated solution was diluted 10 fold in 21GH water prior to use in the assays. Protease activity was assayed following inactivation using the AAPF substrate to ensure complete protease enzyme inactivation. Protease activity was not detectable after the heat treatment of HDD, ADW and HDL detergents. The HDL detergents: Persil Non-Bio, Test HDL 1 and Test HDL 2 are considered boron-free since they contained $\leq 5\text{mg/Kg}$ of boron, when tested for elemental boron content. GSM-B powder formula was dissolved to 3g/L for use and pH was not adjusted.

Table 1.1: List of Detergents And Conditions Used For Cleaning Performance Assays						
Assay Condition	Detergent	Type	Final Wash Conc, (g/L)	Hardness Conc. (ppm)	Buffer	*Set pH
1	Kirkland Ultra	HDD	1.09	150	2 mM NaCO_3	10.6
2	OMO Color	HDD	5.3	250	2 mM NaCO_3	10.6
3	Surf Excel	HDD	4	400	2 mM NaCO_3	10.6
4	Kirkland Ultraclean	HDL	0.17	150	5 mM sodium HEPES	8.2
5	OMO Klein & Krachtig	HDL	2.8	250	5 mM HEPES	8.2
6	Blue Moon HDL	HDL	1.33	100-250	5 mM HEPES	6.5-

						8.2
7	ALL	HDL	0.17	150	5 mM HEPES	8.2
8	Test HDL1	HDL	7.5	200-250	5 mM HEPES	8.2
9	Test HDL 2	HDL	0.84	100-150	5 mM HEPES	8.2
10	Persil Non-Bio	HDL	2.7	200-250	5 mM HEPES	8.2
11	GSM-B pH 10.5	ADW	3	374ppm	not buffered	ND
12	GSM-B pH 9	ADW	3	374ppm	1M citrate added	9
13	SUN® All in 1	ADW	1 tablet dissolved, diluted 10X	374ppm	Not buffered	ND
14	Finish® Quantum	ADW	1 tablet dissolved, diluted 10X	374ppm	Not buffered	ND
15	Finish® All-in-1	ADW	1 tablet dissolved, diluted 10X	374ppm	Not buffered	ND
16	Liby	HDL	0.89	100	5 mM HEPES	8.2
17	Custom Liquid ADW detergent	ADW	3	374ppm	20mM TRIS base	7
18	Custom Liquid ADW detergent	ADW	3	374ppm	20mM TRIS base	9

*pH of wash solution was adjusted for detergents where value is provided and not for those marked ND.

[00184] Detergent sources: ALL, Kirkland Ultra and Kirkland Ultraclean (Sun Products), OMO Color, OMO Klein & Krachtig, Surf Excel, and SUN All in One detergent tablets (Unilever), Finish All in One and Finish Quantum detergent tablets (Reckitt Benckiser), and Blue Moon (Guangzhou Blue Moon). Detergents listed above were purchased from local supermarkets in 2012. Persil Small & Mighty Non-Bio Liquid Detergent “Persil Non-Bio” (Unilever) was purchased in 2014. Liby (Guangzhou Liby Enterprise) was purchased in 2016. Test HDL 1 and Test HDL 2 detergents were custom-made in house and the composition of each is set forth in Tables 1.2 and 1.3, respectively. Table 1.4 shows the composition of the GSM-B Phosphate-free ADW detergent (purchased without enzymes, from WFK Testgewebe GmbH, Brüggen, Deutschland, www.testgewebe.de). GSM-B 9 formula was prepared by adjusting pH of purchased GSM-B Phosphate-free detergent down to pH 9. Table 1.5 shows the composition of Custom Liquid ADW Detergent Formula adjusted to pH7 and pH9 for testing.

Table 1.2: Composition Of Custom-made Test HDL 1 Detergent Formula			
Component	Ingredient	Trade name	Wt. %
Solvent	Water (total)	-	64.89
Surfactant	C12-C15 Pareth-7	Empilan KCL 7	3
	(Sodium) Dodecylbenzenesulfonate	NANSA SSA F	7.5
	K-Cocoate	NANSA PC 38F	3
	Sodium Laureth Sulfate	Empicol ESB3/M6	9
Builder	Sodium Citrate	Sodium Citrate Tribasic Dihydrate	3
Liquid properties /stability	Sorbitol	D-Sorbitol	0.8
	Propylene glycol	1,2-Propanediol	2.5

Table 1.2: Composition Of Custom-made Test HDL 1 Detergent Formula			
	Glycerin	Glycerol	0.8
	Triethanolamine	Triethanolamine	0.5
	Methylisothiazolinone	2-Methyl-4-isothiazolin-3-one	0.01
	Ethanol	-	1
Neutralizer	NaOH (4M)	-	5.27

Table 1.3: Composition Of Custom-made Test HDL 2 Detergent Formula			
Component	Ingredient	Trade name	Wt. %
Solvent	Water (total)	-	QS
	Propylene glycol	-	5
Surfactant	Sodium Alkyl Aryl Ether Sulfate	Polystep B27 (30% active)	12
	Sodium alkylbenzene sulfonate	Nacconal 90G	1.2
Builder	Trisodium citrate	-	0.5
	Na ₂ EDTA	-	0.25
Liquid properties /stability	Sodium formate	-	0.25
	Acrylic acid homopolymer	Sokalan PA 30 CL (50% active)	0.5
	Oleic acid	-	1
	Calcium chloride	-	0.1
	Brightener	Tinopal CBS-X	0.25
	Preservative	Bioban 425 (25% active)	0.04
Neutralizer	Sodium carbonate	-	0.25
	Sodium Hydroxide (50%) to pH 8.5	-	0.25

Table 1.4: Composition Of GSM-B Phosphate-Free Detergent (GSM-B, pH 10.5)	
Component	Wt %
Sodium citrate dehydrate	30.0
Maleic acid/ acrylic acid copolymer sodium Salt (SOKALAN® CP5; BASF)	12.0
Sodium perborate monohydrate	5.0
TAED	2.0
Sodium disilicate: Protol A (Cognis)	25.0
Linear fatty alcohol ethoxylate	2.0
Sodium carbonate anhydrous	add to 100

Table 1.5: Composition of Custom Liquid ADW Detergent. Formula adjusted to pH7 and pH9 for testing.	
Component	Weight %
GLDA (Dissolvine)	37
Glycerine (116)	10
Citric acid monohydrate (38)	4.0
Sodium Citrate tribasic dihydrate	3.0
Plurafac LF301 (277)	2.0
Propylene Glycol	1.0
Bayhibit S	0.25
Xanthane gom (275)	0.7
Acusol 460 N 587 (238)	0.7
Water	41.55

Cleaning Performance In Laundry (HDL And HDD) And ADW detergents

[00185] Variants were tested for cleaning performance relative to Parent on various technical soils: BMI (EMPA-116, blood/milk/ink on cotton), egg (CS-38, egg yolk with pigment aged by heat), and POM (CFT-C10, pigment/oil/milk) for laundry-based applications, and on egg yolk (PAS-38, egg yolk on polyacryl fabric, aged and colored with carbon black dye) for dish-based applications. The EMPA-116, CS-38, and CFT-C10 swatches were pre-rinsed with deionized water for 20 minutes and dried overnight at room temperature. The PAS-38 fabric was pre-rinsed with 10mM CAPS buffer (pH 11) at 60°C for 30min at 900 rpm in an iEMS shaker, rinsed with 2ml deionized water and dried overnight. For all stains, pre-punched swatches in MTP plates (Corning 3641) were prepared by Center for Testmaterials BV, Vlaardingen, Netherlands. These microswatch-containing plates were filled with detergent prior to enzyme addition.

General Sample Set-up for Stability Assays

[00186] Variants were tested for stability under various stress conditions (buffers and detergents as indicated in Table 2 below) by measuring the residual activity following incubation at elevated temperature. The elevated temperature was set to obtain approximately 30% residual activity of the stressed sample compared to the unstressed sample. Diluted enzyme sample was mixed in stressor and unstressed protease activity was measured. The diluted sample in stressor was incubated at elevated temperatures and for time periods listed in Table 2 followed by measurement of the stressed protease activity either by AAPF or DMC hydrolysis. Detergent enzymes were heat inactivated as described above, prior to use in stability assays.

Table 2: Reagents And Conditions Used For Protease Stability Assays			
Assay Condition	Detergent or Buffer	Stress temperature (°C)	Incubation time (min)
1	0.02% LAS, 2.1mM EDTA in 50mM HEPES pH8, 0.005% Tween	53.5	5
2	10% Kirkland Ultra	56	5
3	10% OMO Klein & Krachtig	51	5
4	50mM Tris pH9, 1mM EDTA, 0.005% Tween	51	5
5	10% Test HDL 1	52-55 or 60	5 or 15-20
6	10% Persil Non-Bio, Small & Mighty	62-63 or 66	15-20
7	10% Blue Moon	64 or 68	15-20
8	10% Liby	65 or 69	15-20

[00187] For the unstressed condition, enzyme was assayed immediately for activity on AAPF or DMC (see above). For the stressed condition, the PCR plate was sealed and incubated at elevated temperature for 5 min using an Eppendorf 385 MasterCycler Pro Thermocycler or PTC-200 Peltier thermal cycler, then assayed for activity. Stressed and unstressed activity was measured by either the hydrolysis of the synthetic substrate or by the DMC method, as described above. % residual activities

were calculated by taking a ratio of the stressed to unstressed activity and multiplying by 100. Stability PIs were obtained by dividing the residual activity of a variant by that of the Parent.

[00188] Aliquots of enzyme were added to detergent-filled MTPs containing microswatches to reach a final volume of 200 uL for laundry assays with a final enzyme concentration between 0.5-5 ppm. Laundry cleaning assays with HDL or HDD formulas were carried out at 25°C for 15-30 min, while ADW assays were carried out at 40°C for 30 min. Following incubation, 100-150 uL of supernatant was transferred to a fresh MTP and absorbance was read at 600 nm for EMPA-116 swatches or at 405 nm for CS-10, CS-38, and PAS-38 swatches using a SpectraMax plate reader. Absorbance results were obtained by subtracting the value for a blank control (no enzyme) from each sample value. The cleaning PI for each assay condition was obtained by dividing the absorbance values for a given variant by that of the predicted Parent at the same concentration. The Parent value was determined by fitting the standard curve of the purified Parent sample to a Langmuir fit.

Aged Laundry Cleaning Assay

[00189] (A) MTP assay: Each well of a 96-well MTP was filled with a mini stir disc followed by addition of 270 uL 100% Persil Non-Bio detergent. 18.1 uL of enzyme supernatant diluted to ~500 ppm was added to the detergent-filled MTP and mixed using a magnetic tumbling mixing apparatus (V&P Scientific, VP710 series) for 5 min. For the initial time point (T=0), 25 uL of enzyme containing detergent was transferred to a MTP filled with 125 uL of buffer (5 mM HEPES, 250 ppm 3:1 Ca:Mg water hardness, pH 8). 5 uL aliquots of enzyme diluted in detergent was then added to 185 uL buffer (5 mM HEPES, 250 ppm hardness, pH 8) previously added to a MTP containing an EMPA-116 BMI microswatch. At time point 0 (T=0), the cleaning assay was carried out in an analogous manner as described above for fresh cleaning. For the remaining time points, the detergent-enzyme plate was incubated at 37°C for 3 days or 3 weeks or 55°C for either: 1.5 h, 5 h, or 24 h; aliquots were removed at the end of each time point; and the cleaning assay carried out in an analogous manner as described above for fresh cleaning. A residual cleaning value was obtained by comparing the cleaning at the various remaining time points to the initial cleaning at T=0.

[00190] (B) Conical tube Assay: Clarified enzyme culture supernatant and 100% Test HDL 1 detergent were added to 15 mL Polystyrene centrifuge tubes at a final volume of 1.1 mL to achieve a final enzyme concentration of 88 ppm in detergent. 5 uL samples for each variant were immediately withdrawn from tube and frozen for assaying later as the initial time point (T=0). Samples were incubated at 37°C for 3 days. Following incubation, 5 uL samples were withdrawn and frozen. To prepare samples for the cleaning assay, all samples were thawed and 660 uL of buffer (5 mM HEPES, 250 ppm 3:1 Ca:Mg water hardness, pH 8) was added to the samples. 200 uL aliquots of the diluted samples were added to a MTP containing an EMPA-116 BMI microswatch. Laundry cleaning assays were carried out at 25°C for

60 min in an iEMS incubator shaking at 1400 rpm. Following incubation, 120 uL of supernatant was transferred to a fresh MTP and absorbance was read at 600 nm using the SpectraMax plate reader. Absorbance results were obtained by subtracting the value for a blank control (no enzyme) from each sample value. Residual cleaning for each variant was calculated as the ratio of cleaning remaining after 3 days to the cleaning at T=0 and reported as percent remaining cleaning

EXAMPLE 2

Combinatorial Design And Expression of AprL-clade Protease Variants

[00191] *Bacillus sp.* AprL and *B. licheniformis* Bra7 BliD02339 subtilisins, which are both members of the AprL-clade, were modified to produce AprL-Clade protease variants with improved cleaning performance and detergent stability. The AprL-clade is more fully described in International Appl. No. PCT/US16/32514, filed May 13, 2016, and published as WO2016183509.

[00192] The amino acid sequence of the mature AprL Parent enzyme is set forth as SEQ ID NO:2. A synthetic gene (SEQ ID NO:1) encoding the AprL Parent protease was synthesized by GeneArt, Life Technologies and used for generating variant sequences utilizing conventional molecular biology techniques (*see, e.g.*, Sambrook et al, "Molecular Cloning", Cold Spring Harbor Laboratory Press). Genes encoding AprL variants were also synthesized. The various AprL genes were then cloned into the pHYT expression vector (derived from pHY300PLK (Takara-Clontech)). The expression vector contained the aprE promoter (SEQ ID NO:3), aprE signal peptide (SEQ ID NO:4), AprL propeptide (SEQ ID NO:5), and BPN' terminator sequences (SEQ ID NO:6). DNA fragments encoding the various mature protease sequences of interest (wild-type and variants) were introduced using techniques known in the art. A suitable *B. subtilis* host strain was transformed with the expression plasmids containing the AprL Parent and protease variant gene sequences. The cells were grown in 96-well MTPs in cultivation medium (enriched semi-defined media based on MOPs buffer, with urea as major nitrogen source, glucose as the main carbon source, supplemented with 1% soytone for robust cell growth, containing antibiotic selection) for 2 days at 32°C, 300 rpm, with 80% humidity in shaking incubator. After centrifugation and filtration, clarified culture supernatants containing the proteases of interest were used for assays. The same process was used for construction of the BliD02339 wild-type and variant expression cassettes, subsequent *B. subtilis* host strain transformations and cultivation.

[00193] In some instances, the AprL variants were generated from an integration cassette that targets the *B. subtilis* aprE locus. Pools of DNA encoding the genes of interest were introduced into an expression cassette that was later integrated into the chromosome of a suitable *B. subtilis* strain using methods known in the art. DNA fragments containing the following elements: BPN' terminator (SEQ ID NO: 6), chloramphenicol resistance marker from *Staphylococcus aureus* (SEQ ID NO:165), *B. subtilis* aprE promoter (SEQ ID NO:3), *B. subtilis* aprE signal peptide sequence (SEQ ID NO:4), and AprL

propeptide (SEQ ID NO:5) were part of the expression cassette. The transformed cells were cultured as described above, and proteins of interest were similarly isolated

EXAMPLE 3

Performance Evaluation Of AprL Variants In Laundry Application

[00194] Variants of AprL Parent protease (SEQ ID NO:2) (hereinafter “AprL Variants”), prepared as described in Example 2 and listed in Table 3, were evaluated for cleaning performance and stability as described in Examples 3-7.

[00195] The AprL variants described in Table 3 were tested in the Laundry cleaning performance assay described in Example 1 using EMPA-116 BMI soil and the following laundry detergents: Kirkland Ultra HDD, OMO Color HDD, Surf Excel, Kirkland Ultraclean HDL, OMO Klein & Krachtig HDL, and Blue Moon HDL. The Table 3 AprL Variants with a PI >1.1 versus AprL Parent are listed in Table 4.

Table 3: AprL Variants Evaluated in Examples 3-7.		
Sample ID	Sequence Substitutions Relative To AprL Parent	SEQ ID NO:
SQM-002	M123I-Q136R	10
SQM-003	P9E-M123I	11
SQM-004	V45R-M123I	12
SQM-005	G127T-G165S-T210E	13
SQM-006	M123I-G165S-N239S	14
SQM-007	M123I-Q136R-G165Q	15
SQM-008	M123I-Q136R-S258E	16
SQM-009	M123I-V146S-T210E	17
SQM-010	P9E-V45R-M123I	18
SQM-011	Q136R-G165Q-S258E	19
SQM-012	V45R-M123I-Q136R	20
SQM-014	G127T-V146S-G165S-N239S	21
SQM-015	L10M-K27R-S98H-M123I	22
SQM-016	M123I-Q136R-G165Q-S258E	23
SQM-017	M123I-V146S-G165S-T210P	24
SQM-018	M123I-V146S-T210P-S250G	25
SQM-019	P9E-M123I-Q136R-G165Q	26
SQM-020	P9E-Q136R-G165Q-S258E	27
SQM-021	P9E-V45R-M123I-Q136R	28
SQM-022	V45R-M123I-Q136R-G165Q	29
SQM-023	V45R-M123I-Q136R-S258E	30
SQM-024	V45R-Q136R-G165Q-S258E	31
SQM-026	K27R-N43A-S98H-M123I-T210H	32
SQM-027	P9E-M123I-Q136R-G165Q-S258E	33
SQM-028	P9E-V45R-M123I-Q136R-G165Q	34
SQM-029	P9E-V45R-M123I-Q136R-S258E	35
SQM-030	V45R-M123I-Q136R-G165Q-S258E	36
SQM-031	Y103I-G127T-V146S-G165S-N239S	37
SQM-032	T77H-V87T-M123I-Q136R-N184Q-N239S	38
SQM-034	T77H-V87T-M123I-Q136R-G165Q-N184Q-N239S	40

Table 3: AprL Variants Evaluated in Examples 3-7.		
Sample ID	Sequence Substitutions Relative To AprL Parent	SEQ ID NO:
SQM-035	T77S-V87S-M123I-Q136R-G165Q-S187P-S258P	41
SQM-036	T77S-V87S-M123I-Q136R-N160C-S187P-S258P	42
SQM-037	T77H-V87T-M123I-Q136R-N184Q-S187P-N239S-S258P	43
SQM-038	T3Q-K22Y-A24N-V45I-A48I-T77N-N217S-L234I-Y255R	44
SQM-039	P9E-T77Q-V87T-M123I-Q136R-G165Q-N184Q-N217S-N239S-S258E	45
SQM-040	N43A-V45R-T77H-V87T-M123I-Q136R-G165A-N184Q-S187P-N239S-S258P	46
SQM-041	P9E-N43A-V45R-T77S-V87S-M123I-Q136R-G165Q-S187P-N217S-S258P	47
SQM-042	P9E-N43A-V45R-T77H-V87T-M123I-Q136R-G165Q-N184Q-S187P-N239S-S258P	48
SQM-043	P9E-N43A-V45R-T77Q-V87T-M123I-Q136R-G165Q-N184Q-N217S-N239S-S258E	49
SQM-044	P9E-N43A-V45R-T77H-V87T-M123I-Q136R-G165A-N184Q-S187P-N217S-N239S-S258E	50

Table 4. AprL Variants With Improved Laundry Cleaning Performance on BMI Stain in at Least One Detergent.

Sample ID	HDL			HDD		
	Blue Moon	Kirkland	OMO	Kirkland	OMO	Surf
SQM-002	0.81	1.02	0.96	1.63	1.01	1.27
SQM-003	0.96	0.97	1.03	1.25	0.62	0.62
SQM-004	0.91	1.06	1.31	1.7	0.46	1.2
SQM-007	0.54	1.22	0.93	1.65	1.51	1.29
SQM-008	0.66	1.19	0.93	1.78	1.22	1.26
SQM-010	0.91	1.15	1.01	0.94	0.5	0.66
SQM-012	0.59	0.98	0.89	2.17	1.67	1.53
SQM-016	0.48	0.58	0.46	1.56	1.32	1.3
SQM-019	0.47	0.98	0.64	1.51	0.96	1.13
SQM-021	0.85	1.32	1.15	1.88	1.26	1.22
SQM-022	0.5	0.82	0.64	1.14	0.86	1.00
SQM-023	0.76	1.34	0.98	1.65	1.65	1.24
SQM-024	0.59	0.74	0.67	1.34	1.29	1.13
SQM-028	0.46	0.99	0.76	2.21	1.78	1.49
SQM-029	0.46	0.96	0.65	1.41	1.04	1.23
SQM-030	0.55	0.87	0.7	1.26	1.01	1.03

EXAMPLE 4

Performance Evaluation Of AprL Variants In Dishwashing Application

[00196] The AprL variants described in Table 3 were tested in the ADW cleaning performance assay described in Example 1 using PAS-38 egg yolk soil and the following ADW detergents: GSM-B pH 10, GSM-B pH 9, SUN® All in One, Finish® Quantum, and Finish® All in One. AprL variants (listed on Table 3) with a PI >1.01 versus AprL Parent in at least one of these ADW detergents are listed in Table 5. The Table 3 AprL variants with PI >1.01 versus AprL Parent in all of these ADW detergents are listed in Table 6.

Table 5: AprL Variants With Improved ADW Cleaning Performance on Egg Stain in at Least One detergent.					
Sample ID	Finish[®] ALL in One	GSMB pH 10	GSMB pH 9	Finish[®] Quantum	SUN[®] All in One
SQM-002	1.17	1.12	1.2	1.15	1.26
SQM-003	0.61	0.59	1.03	0.89	0.68
SQM-004	0.97	0.89	1.38	1.21	0.94
SQM-007	1.2	1.15	1.2	1.03	1.32
SQM-008	0.75	1.02	1.14	1.15	0.97
SQM-010	0.68	0.69	1.01	1.19	0.72
SQM-012	1.1	1.03	1.3	1.12	1.18
SQM-016	1.21	1.19	1.26	1.24	1.11
SQM-019	0.98	0.85	1.07	0.88	1.1
SQM-020	0.92	0.82	0.76	1.03	0.9
SQM-021	0.96	0.92	1.25	1.07	1.07
SQM-022	1.14	1.1	1.09	1.00	1.17
SQM-023	0.84	0.87	1.04	0.9	0.89
SQM-024	1.13	1.04	1.15	1.06	1.07
SQM-028	1.38	1.31	1.45	1.18	1.44
SQM-029	0.9	0.88	1.05	1.02	1.00
SQM-030	1.07	1.12	1.12	1.03	1.17

Table 6: AprL Variants With Improved ADW Cleaning Performance on Egg Stain.					
Sample ID	Finish[®] All in One	GSMB pH10	GSMB pH 9	Finish[®] Quantum	SUN[®] All in One
SQM-002	1.17	1.12	1.2	1.15	1.26
SQM-007	1.2	1.15	1.2	1.03	1.32
SQM-012	1.1	1.03	1.3	1.12	1.18
SQM-016	1.21	1.19	1.26	1.24	1.11
SQM-022	1.14	1.1	1.09	1.00	1.17
SQM-024	1.13	1.04	1.15	1.06	1.07
SQM-028	1.38	1.31	1.45	1.18	1.44
SQM-030	1.07	1.12	1.12	1.03	1.17

EXAMPLE 5

AprL Variants With Improved Stability

[00197] The AprL variants described in Table 3 were tested in the stability assays described in Example 1 in the following detergents and buffer: LAS-EDTA buffer at 53.5°C, 10% Kirkland Ultra HDL pH 8.2 at 56°C, and OMO Klein & Krachtig HDL pH 8.2 at 51°C. The Table 3 AprL variants with (i) stability PI>1.1 versus AprL Parent in at least one of the stability assays described in Example 1, and (ii) cleaning PI>1.01 in both the Laundry and ADW cleaning performance assays described in Example 1 are listed in Table 7. The Table 3 AprL variants with PI>1.1 versus AprL Parent in at least one Laundry (HDL or HDD), ADW, or Stability assay described in Example 1 are listed in Table 8.

Table 7: AprL Variants With Improved Stability.			
Sample ID	LAS/EDTA	OMO	Kirkland
SQM-002	1.04	1.35	1.32
SQM-003	1.31	1.02	0.88

Table 7: AprL Variants With Improved Stability.			
Sample ID	LAS/EDTA	OMO	Kirkland
SQM-004	1.02	1.25	0.76
SQM-007	2.01	1.68	2.06
SQM-008	1.51	1.38	1.39
SQM-010	1.21	1.46	0.85
SQM-011	2.67	1.8	2.09
SQM-012	1.01	1.07	1.19
SQM-016	2.16	1.87	1.94
SQM-020	2.92	1.83	2.31
SQM-021	1.11	1.47	1.00
SQM-022	1.45	1.68	1.87
SQM-023	1.55	1.42	1.32
SQM-024	2.31	1.63	2.39
SQM-027	2.39	1.9	2.07
SQM-028	1.81	1.59	1.47
SQM-029	1.41	1.58	1.23
SQM-030	1.84	1.83	2.11

Table 8. AprL Variants With Improved Cleaning Performance and Stability.														
Sample ID	HDL, BMI			HDD, BMI			ADW, EGG					Stability		
	Blue Moon	Kirkland	OMO	Kirkland	OMO	Surf	Finish® All-in-1	GSMB pH 10	GSMB pH 9	Finish® Quantum	SUN	LAS/EDTA	OMO	Kirkland
SQM-002	0.81	1.02	0.96	1.63	1.01	1.27	1.17	1.12	1.2	1.15	1.26	1.04	1.35	1.32
SQM-004	0.91	1.06	1.31	1.7	0.46	1.2	0.97	0.89	1.38	1.21	0.94	1.02	1.25	0.76
SQM-007	0.54	1.22	0.93	1.65	1.51	1.29	1.2	1.15	1.2	1.03	1.32	2.01	1.68	2.06
SQM-008	0.66	1.19	0.93	1.78	1.22	1.26	0.75	1.02	1.14	1.15	0.97	1.51	1.38	1.39
SQM-010	0.91	1.15	1.01	0.94	0.5	0.66	0.68	0.69	1.01	1.19	0.72	1.21	1.46	0.85
SQM-012	0.59	0.98	0.89	2.17	1.67	1.53	1.1	1.03	1.3	1.12	1.18	1.01	1.07	1.19
SQM-016	0.48	0.58	0.46	1.56	1.32	1.3	1.21	1.19	1.26	1.24	1.11	2.16	1.87	1.94
SQM-021	0.85	1.32	1.15	1.88	1.26	1.22	0.96	0.92	1.25	1.07	1.07	1.11	1.47	1.00
SQM-022	0.5	0.82	0.64	1.14	0.86	1.00	1.14	1.1	1.09	1.00	1.17	1.45	1.68	1.87
SQM-024	0.59	0.74	0.67	1.34	1.29	1.13	1.13	1.04	1.15	1.06	1.07	2.31	1.63	2.39
SQM-028	0.46	0.99	0.76	2.21	1.78	1.49	1.38	1.31	1.45	1.18	1.44	1.81	1.59	1.47
SQM-030	0.55	0.87	0.7	1.26	1.01	1.03	1.07	1.12	1.12	1.03	1.17	1.84	1.83	2.11

EXAMPLE 6

Performance Evaluation Of Additional AprL Variants

[00198] The AprL variants described in Table 3 were tested in the protease activity assay (AAPF or DMC assay) described in Example 1, laundry cleaning performance assay described in Example 1 using EMPA-116 BMI soil and All HDL and OMO Klein & Krachtig HDL detergents, ADW cleaning performance assay described in Example 1 using PAS-38 egg yolk soil and GSM-B pH 10 and Finish® Quantum dish detergents, and the stability assays described in Example 1 in the following buffers: LAS-

EDTA buffer at 53.5°C and Tris-EDTA buffer at 51°C. The Table 3 AprL variants with a laundry cleaning PI>1.01 versus AprL parent are listed in Table 9. The Table 3 AprL variants with an ADW cleaning PI>1.01 versus AprL parent are listed in Table 10. The Table 3 AprL variants with a stability PI>1.01 versus AprL parent are listed in Table 11. The Table 3 AprL variants with laundry cleaning, ADW cleaning, and stability PI>1.01 versus AprL parent in at least one detergent are listed in Table 12.

Table 9: AprL Variants Tested in HDL Detergents and BMI soil.

Sample ID	ALL	OMO
SQM-005	1.59	1.35
SQM-006	1.88	1.22
SQM-009	1.78	1.02
SQM-014	2.59	1.22
SQM-015	2.04	1.31
SQM-017	1.83	1.44
SQM-018	2.65	1.47
SQM-026	2.47	1.10
SQM-031	1.44	1.35
SQM-032	2.40	1.21
SQM-034	1.45	1.03
SQM-035	1.25	1.04
SQM-036	1.38	1.04
SQM-037	1.69	1.17
SQM-038	1.00	1.17
SQM-041	1.19	0.85
SQM-042	1.40	0.86
SQM-043	1.31	1.11
SQM-044	1.05	0.91

Table 10: AprL Variants Tested in ADW Detergents and Egg soil.

Sample ID	GSM-B pH 10	Finish® Quantum
-----------	-------------	-----------------

Table 11: AprL Variants Tested in Stability Assays.

Sample	LAS/EDTA	Tris/EDTA
SQM-005	1.94	1.68
SQM-006	1.52	1.34
SQM-009	1.84	1.66
SQM-014	1.99	1.81
SQM-015	1.97	1.24
SQM-017	2.07	2.14
SQM-018	1.83	2.02
SQM-026	1.61	1.24
SQM-031	1.89	1.87
SQM-032	2.05	1.72
SQM-034	2.60	1.78
SQM-035	2.68	1.95
SQM-036	2.72	2.20
SQM-037	2.49	1.82
SQM-038	0.31	1.14
SQM-039	3.10	2.18
SQM-040	2.57	1.76
SQM-041	2.98	1.88
SQM-042	2.46	0.71
SQM-043	2.78	1.98
SQM-044	2.78	1.96

Table 10: AprL Variants Tested in ADW Detergents and Egg soil.

Sample ID	GSM-B pH 10	Finish® Quantum
SQM-005	2.16	1.57
SQM-006	2.59	1.83
SQM-009	1.78	1.38
SQM-014	1.95	1.46
SQM-015	1.75	1.39
SQM-017	2.20	1.56
SQM-018	2.10	1.51
SQM-026	1.71	1.41
SQM-031	2.14	1.38
SQM-032	2.49	1.42
SQM-034	2.07	1.35
SQM-035	2.04	1.24
SQM-036	1.60	1.17
SQM-037	1.87	1.30
SQM-038	1.85	1.47
SQM-039	2.09	1.41
SQM-040	1.82	1.21
SQM-041	1.90	1.29
SQM-042	2.03	1.25
SQM-043	2.24	1.47
SQM-044	1.98	1.26

Table 12: AprL Variants Tested in Cleaning Performance and Stability Assays.

Sample ID	HDL, BMI		ADW, EGG		Stability	
	ALL	OMO	GSM-B pH10	Finish® Quantum	LAS/EDTA	Tris/EDTA
SQM-005	1.59	1.35	2.16	1.57	1.94	1.68
SQM-006	1.88	1.22	2.59	1.83	1.52	1.34
SQM-009	1.78	1.02	1.78	1.38	1.84	1.66
SQM-014	2.59	1.22	1.95	1.46	1.99	1.81
SQM-015	2.04	1.31	1.75	1.39	1.97	1.24
SQM-017	1.83	1.44	2.20	1.56	2.07	2.14
SQM-018	2.65	1.47	2.10	1.51	1.83	2.02
SQM-026	2.47	1.10	1.71	1.41	1.61	1.24
SQM-031	1.44	1.35	2.14	1.38	1.89	1.87
SQM-032	2.40	1.21	2.49	1.42	2.05	1.72
SQM-034	1.45	1.03	2.07	1.35	2.60	1.78

Table 12: AprL Variants Tested in Cleaning Performance and Stability Assays.						
Sample ID	HDL, BMI		ADW, EGG		Stability	
	All	OMO	GSMB pH10	Finish [®] Quantum	LAS/EDTA	Tris/EDTA
SQM-035	1.25	1.04	2.04	1.24	2.68	1.95
SQM-036	1.38	1.04	1.60	1.17	2.72	2.20
SQM-037	1.69	1.17	1.87	1.30	2.49	1.82
SQM-038	1.00	1.17	1.85	1.47	0.31	1.14

Table 12: AprL Variants Tested in Cleaning Performance and Stability Assays.						
Sample ID	HDL, BMI		ADW, EGG		Stability	
	All	OMO	GSMB pH10	Finish [®] Quantum	LAS/EDTA	Tris/EDTA
SQM-041	1.19	0.85	1.90	1.29	2.98	1.88
SQM-042	1.40	0.86	2.03	1.25	2.46	0.71
SQM-043	1.31	1.11	2.24	1.47	2.78	1.98
SQM-044	1.05	0.91	1.98	1.26	2.78	1.96

EXAMPLE 7

Cleaning Performance Of AprL Variants In Non-boron Containing Detergents

[00199] The cleaning performance of purified AprL variants: M123I-Q136R, M123I-Q136R-G165Q, V045R-Q136R-G165Q-S258E, V045R-M123I-Q136R-G165Q-S258E, and V045R-M123I-Q136R were tested in the following detergents that did not contain detectable amounts of boron or boron-containing compounds: All 3X, Purex[®] Dirt Lift, and Persil Non Bio. These AprL variants were tested in the laundry cleaning performance assays described in Example 1. Results are shown in Table 13 as PI calculated versus AprL Parent. The proteases were tested at a dosage of 2.5 ppm. The amount of boron present in the boron-free detergents of this Example was quantified by elemental analysis, and in all instances, only trace amounts (less than 5 mg) of boron per kg of detergent were detected.

Table 13: Cleaning Performance In Boron-free Laundry Detergents, Reported As PI vs AprL Parent			
	Persil Non-Bio	ALL 3X	Purex [®] Dirt Lift
Parent	1.0	1.0	1.0
SQM-002	1.1	1.8	3.0
SQM-012	1.1	2.7	2.0
SQM-007	1.0	1.4	2.0
SQM-024	0.9	1.1	1.2
SQM-030	1.1	1.8	2.2

EXAMPLE 8

Cleaning Performance And Detergent Stability Of AprL And BliD02339 Protease Variants

[00200] AprL Variants listed in Table 14 were generated as described in Example 2. The cleaning performance and stability of the Table 14 variants were tested in one or more cleaning performance and stability assay described in Example 1.

[00201] The laundry cleaning performance of the AprL variants listed in Table 14 was tested on EMPA-116 microswatches in Test HDL 1 and Persil Non-Bio detergents (Table 1.1, assay conditions 8 and 10). Enzyme stability was tested in 10% Test HDL 1 detergent at 52.5°C and/or 54°C (Table 2, assay condition 5). Table 15 shows the results as PI values when compared to AprL Parent, wherein ND means not determined.

[00202] Table 17 lists the performance of the AprL Variants tested according to the laundry cleaning performance and stability assays described in Example 1: (i) cleaning performance on EMPA-116, CFT CS-38, and CFT-C10 microswatches in Persil Non-Bio, Test HDL 1 and Test HDL 2 detergents, and (ii) stability in 10% PNB at 63°C for 15-20 min and/or 10% Test HDL 1 at 60°C for 15-20 min. Cleaning performance results are reported as PI values versus the SQCBV35 variant (SEQ ID NO:63), and stability results are reported as % remaining activity of sample after stress versus unstressed condition. ND means the value was not determined.

Table 14: AprL Protease Variants		
Sample ID	Sequence Substitutions Relative To AprL Parent	SEQ ID NO:
SQCBV12	T77N-N217S-L234W	51
SQCBV13	T77N-N217S	52
SQCBV14	T77N-G126Q-G165Q-T210P-N217S-L234W-S258P	53
SQCBV15	T77N-G127F-G165Q-A202V-T210P-N217S-L234W-S258P	54
SQCBV18	A1Q-T3V-V28A-I35A-A68S-T77N-S86H-T115F-N217S-N239S-S250G-S258P-Q274A	55
SQCBV19	T77N-M123I-T210P-N239S-S250G-S258P	56
SQCBV20	V45I-A68S-T77N-M123I-L125N-G165Q-N217S-S258P	57
SQCBV21	A68S-V71A-T77N-L95A-G165Q-N184Q-T210P-S258P	58
SQCBV23	M123I-T210P-N217S-S258P	59
SQCBV32	G127F-G165Q-N217S-S258P	60
SQCBV33	T77N-G165Q-N217S-S258P	61
SQCBV34	T77S-G165A-N184Q-T210P-N217S	62
SQCBV35	A68S-T77N-A128P-G165Q-N184Q-N217S	63
SQCBV36	V71A-T77S-N184Q-S258P	64
SQCBV37	T77N-M123I-G165Q-N184Q-N217S	65
SQCBV38	V71A-L95A-A128S-T210P	66
SQCBV39	V71A-M123I-A128S-T210P	67
SQCBV40	V71A-M123I-A128S-G165E	68
SQCBV41	T77S-A128P-N184Q-N217S-F260W	69
SQCBV42	T77H-A128S-G165Q-T210P	70
SQCBV43	T77D-S86R-S155E-G165A-N184Q	71
SQCBV44	A068S-V071A-T077N-A128S-G165Q-T210P-N217S	72
SQCBV45	V28A-I35A-A68S-T77N-A128S-G165Q-N184Q-G203N-T210P	73
SQCBV46	A128P-T210P	74
SQCBV47	T210P-S258P	75
SQCBV49	A128P-T210P-S259P	76
SQCBV50	A128P-S187P-T210P	77
SQCBV52	A128P-S187P-T210P-S258P	78
SQCBV66	P9E-T77S-V87T-T210P-N239K	79
SQCBV70	K15I-L95Q-M123I-G165S-T210P-N217S	80
SQCBV72	K15I-M123I-V146S-G165S-T210P-N239S	81
SQCBV73	K15I-N76K-M123I-V146S-G165S-T210P-N239S	82
SQCBV74	K15I-V146S-G165S-T210P-N239S	83
SQCBV75	P9E-V146S-N184Q-G203E-T210P-P238R-N239S	84
SQCBV78	M123I-G165S-N217S	85

Table 14: AprL Protease Variants		
Sample ID	Sequence Substitutions Relative To AprL Parent	SEQ ID NO:
SQCBV85	K15I-L95Q-S108K-G165S-G203E-T210P-N217S	86
SQCBV87	K15I-L95Q-S108K-G165S-G203E-T210P	87
SQCBV88	P9E-K15I-G165S-A193D-G203E-T210P	88
SQCBV103	T77N-M123I-G165Q-N217S	89
SQCBV104	T77H-G165Q	90
SQCBV105	T77N-G165Q-N217S	91
SQCBV106	A68S-V71A-T77N-G165Q-N217S	92
SQCBV107	T77N-V87S-M123I-S155G-G165Q-N217S	93
SQCBV108	T77H-V87S-S155G-G165Q	94
SQCBV109	T77N-V87S-S155G-G165Q-N217S	95
SQCBV111	T77N-M123I-G165Q	96
SQCBV112	T77N-G165Q	97
SQCBV113	A68S-V71A-T77N-G165Q	98
SQCBV114	T77N-V87S-M123I-S155G-G165Q	99
SQCBV119	I35A-A68S-T77N-V87S-V147L-G165Q	100
SQCBV121	A68S-T77N-V87S-V147L-S155E-G165Q	101
SQCBV125	T77N-V87S-M123I-V147L-S155G-G165Q	102
SQCBV143	A68S-T77N-M123I-G165Q-N184Q-N217S	103
SQCBV161	T77N-M123I-A128P-G165Q-N184Q-N217S	104
SQCBV188	A68S-V71A-T77N-S100N-A128P-G165Q-N184Q-T210P-N217S-N239T	105
SQCBV191	T77N-A128P-G165Q-N184Q-N217S	106
SQCBV192	A68S-A128P-G165Q-N184Q-N217S	107
SQCBV193	A68S-T77N-G165Q-N184Q-N217S	108
SQCBV194	A68S-T77N-A128P-N184Q-N217S	109
SQCBV195	A68S-T77N-A128P-G165Q-N217S	110
SQCBV196	A68S-T77N-A128P-G165Q-N184Q	111
SQCBV197	M123I-G165Q-N184Q-N217S	112
SQCBV198	T77N-G165Q-N184Q-N217S	113
SQCBV199	T77N-M123I-N184Q-N217S	114
SQCBV200	T77N-M123I-G165Q-N184Q	115
SQCBV201	T3V-A68S-T77N-A128P-G165Q-N184Q-N217S	116
SQCBV202	T3V-A68S-T77N-M123I-A128P-G165Q-N184Q-N217S	117
SQCBV203	T3V-T77N-M123I-G165Q-N184Q-N217S	118
SQCBV204	A68S-T77N-S108H-A128P-G165Q-N184Q-N217S	119
SQCBV205	A68S-T77N-G99S-A128P-G165Q-N184Q-N217S	120
SQCBV206	A68S-T77N-T114N-A128P-G165Q-N184Q-N217S	121
SQCBV207	A68S-T77N-M123I-A128P-G165Q-N184Q-N217S	122
SQCBV208	A68S-T77N-G127T-A128P-G165Q-N184Q-N217S	123
SQCBV209	A68S-T77N-A128P-G165Q-N184Q-T210P-N217S	124
SQCBV211	A68S-T77N-G99S-M123I-A128P-G165Q-N184Q-N217S	125
SQCBV212	A68S-T77N-G99S-G127T-A128P-G165Q-N184Q-N217S	126
SQCBV213	A68S-T77N-G99S-A128P-G165Q-N184Q-T210P-N217S	127
SQCBV214	A68S-T77N-T114N-M123I-A128P-G165Q-N184Q-N217S	128
SQCBV215	A68S-T77N-T114N-G127T-A128P-G165Q-N184Q-N217S	129
SQCBV216	A68S-T77N-T114N-A128P-G165Q-N184Q-T210P-N217S	130
SQCBV217	A68S-T77N-M123I-G127T-A128P-G165Q-N184Q-N217S	131

Table 14: AprL Protease Variants		
Sample ID	Sequence Substitutions Relative To AprL Parent	SEQ ID NO:
SQCBV218	A68S-T77N-M123I-A128P-G165Q-N184Q-T210P-N217S	132
SQCBV219	A68S-T77N-G127T-A128P-G165Q-N184Q-T210P-N217S	133
SQCBV221	A68S-T77N-G99S-M123I-A128P-G165Q-N184Q-T210P-N217S	134
SQCBV223	A68S-T77N-M123I-G127T-A128P-G165Q-N184Q-T210P-N217S	135
SQCBV225	T3V-A68S-T77N-G99S-M123I-A128P-G165Q-N184Q-N217S	136
SQCBV226	T3V-A68S-T77N-T114N-M123I-A128P-G165Q-N184Q-N217S	137
SQCBV227	T3V-A68S-T77N-M123I-G127T-A128P-G165Q-N184Q-N217S	138
SQCBV228	T3V-A68S-T77N-M123I-A128P-G165Q-N184Q-T210P-N217S	139
SQCBV269	S86H-T115F-A143V-G165Q-A202V-S258P	140
SQCBV270	S86H-L95A-N184Q-A202V-T210P-N217S-L234W-S250G-S258P	141
SQCBV271	V30T-I35A-T77N-G165Q-A202V-T210P-S250G-S258P	142
SQCBV272	A1Q-T3V-I35A-T77N-T78I-S86H-T115F-G165Q-N184Q-N217S-L234W-S258P	143
SQCBV273	A1Q-A37T-T77N-T78I-S86H-M123I-G165Q-N184Q-A202V-N239S-S250G-S258P-Q274A	144
SQCBV274	P9E-T77N-N96S-G165Q-G203E-S250G	145
SQCBV275	P9E-T77N-N96S-G165E-G203E-T210P-S250G	146
SQCBV276	A1Q-T3V-S86H-G130S-S155G-G165Q-N184Q-A202V-G203N-T210P-Q274A	147
SQCBV277	A1Q-T3V-S86H-G165Q-N184Q-N239S-S250G-S258P	148
SQCBV278	A1Q-T77N-S86H-V147L-A151G-N184Q-N217S-S258P	149
SQCBV279	S86H-T115F-S155G-G165Q-N184Q-A202V-Q274A	150
SQCBV280	A68S-S86H-T115F-N184Q-A202V-G203N-N239S-S250G-S258P-Q274A	151

Table 15: Cleaning And Stability Of AprL Variants Compared To AprL Parent, Reported as PI				
Sample ID	Cleaning Performance		Stability	
	EMPA 116 Swatches		10% Test HDL	
	Test HDL1	Persil Non-bio	55°C	52.5°C
AprL WT	1.0	1.0	1.0	1.0
SQCBV12	0.8	ND	ND	1.1
SQCBV13	1.0	ND	ND	1.6
SQCBV14	0.6	ND	ND	1.1
SQCBV15	0.8	ND	ND	1.1
SQCBV18	0.7	ND	ND	1.2
SQCBV19	1.2	ND	ND	1.2
SQCBV20	1.1	ND	ND	1.1
SQCBV21	1.0	ND	ND	1.1
SQCBV23	1.1	ND	ND	1.4
SQCBV32	1.1	ND	ND	1.2
SQCBV33	1.0	ND	ND	1.3
SQCBV34	0.7	ND	ND	1.1
SQCBV35	0.9	ND	ND	1.4
SQCBV36	0.9	ND	ND	1.4
SQCBV37	0.9	ND	ND	1.5
SQCBV38	1.2	ND	ND	1.2
SQCBV39	1.2	ND	ND	1.4
SQCBV40	1.1	ND	ND	1.4

Table 15: Cleaning And Stability Of AprL Variants Compared To AprL Parent, Reported as PI				
Sample ID	Cleaning Performance		Stability	
	EMPA 116 Swatches		10% Test HDL	
	Test HDL1	Persil Non-bio	55°C	52.5°C
SQCBV41	1.0	ND	ND	1.4
SQCBV42	1.0	ND	ND	1.5
SQCBV43	1.0	ND	ND	1.7
SQCBV44	0.9	ND	ND	1.7
SQCBV45	0.9	ND	ND	1.7
SQCBV46	0.9	ND	ND	1.1
SQCBV47	1.0	ND	ND	1.4
SQCBV49	1.0	ND	ND	1.3
SQCBV50	0.9	ND	ND	1.6
SQCBV52	1.0	ND	ND	1.7
SQCBV66	0.8	ND	ND	1.3
SQCBV70	0.7	ND	ND	1.9
SQCBV72	1.0	ND	ND	1.5
SQCBV73	0.8	ND	ND	1.4
SQCBV74	0.8	ND	ND	1.4
SQCBV75	0.8	ND	ND	2.0
SQCBV78	1.0	ND	ND	1.9
SQCBV85	0.8	ND	ND	1.3
SQCBV87	0.7	ND	ND	1.1
SQCBV88	0.6	ND	ND	1.7
SQCBV269	0.9	0.87	7.64	ND
SQCBV270	0.9	0.87	7.65	ND
SQCBV271	0.8	0.84	7.28	ND
SQCBV272	0.9	0.86	8.06	ND
SQCBV273	0.9	0.88	7.7	ND
SQCBV274	0.8	0.85	7.37	ND
SQCBV275	0.9	0.88	7.82	ND
SQCBV276	0.7	0.7	7.85	ND
SQCBV277	0.9	0.93	6.99	ND
SQCBV278	0.9	0.83	7.5	ND
SQCBV279	0.7	0.71	7.83	ND
SQCBV280	0.7	0.73	7.46	ND

Table 16A: Aged Cleaning Performance Of AprL Variants Compared To AprL Parent, Reported As % Remaining Cleaning Activity

Sample ID	% Remaining Cleaning Activity After 24h
AprL WT	0
SQCBV13	94
SQCBV14	98
SQCBV19	89
SQCBV20	110
SQCBV21	99
SQCBV23	72
SQCBV32	72
SQCBV33	84
SQCBV35	110
SQCBV36	75
SQCBV37	110
SQCBV38	55
SQCBV39	61
SQCBV40	84

Table 16A: Aged Cleaning Performance Of AprL Variants Compared To AprL Parent, Reported As % Remaining Cleaning Activity

Sample ID	% Remaining Cleaning Activity After 24h
SQCBV41	86
SQCBV42	100
SQCBV43	99
SQCBV44	84
SQCBV45	90
SQCBV46	50
SQCBV47	72
SQCBV49	54
SQCBV50	73
SQCBV52	74
SQCBV66	96
SQCBV72	76
SQCBV73	58
SQCBV74	87
SQCBV78	51

Table 16B: Aged Cleaning Performance Of AprL Variants Compared To AprL Parent, Reported As % Remaining Cleaning Activity

Sample ID	% Remaining Cleaning Activity
AprL WT	6
SQCBV269	91
SQCBV270	87
SQCBV277	58
SQCBV276	60
SQCBV273	71
SQCBV275	61

Table 16B: Aged Cleaning Performance Of AprL Variants Compared To AprL Parent, Reported As % Remaining Cleaning Activity

Sample ID	% Remaining Cleaning Activity
SQCBV272	78
SQCBV271	84
SQCBV279	54
SQCBV280	53
SQCBV278	57
SQCBV274	63

Table 17: Cleaning Of AprL Variants Compared To SQCBV35 Variant, Reported As PI For Cleaning Assays And Stability As % Remaining Activity For Stability Assays

Sample ID	Microswatch Cleaning									Stability	
	Persil-Non Bio			Test HDL 2			Test HDL 1			10% PNB	Test HDL 1
	EMPA -116	CFT-CS-38	CFT-C10	EMPA -116	CFT-CS-38	CFT-C10	EMPA -116	CFT-CS-38	CFT-C10	63°C	60°C
AprL WT	1.27	1.33	1.43	1.81	1.51	1.23	1.33	1.45	1.49	0.1	0.1
SQCBV35	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	15.4	40.7
SQCBV103	1.12	0.90	1.17	1.64	1.38	1.16	1.15	1.26	1.17	0.2	1.7
SQCBV104	1.17	1.14	1.19	1.64	1.71	1.20	ND	ND	ND	0.2	ND
SQCBV105	1.06	0.92	1.26	1.62	1.67	1.13	ND	ND	ND	0.8	ND
SQCBV106	1.01	0.66	0.93	1.33	1.19	0.89	ND	ND	ND	2.9	ND
SQCBV107	1.16	0.92	0.99	1.56	1.56	1.02	ND	ND	ND	7.1	ND
SQCBV108	0.97	0.67	1.06	0.91	1.10	0.95	ND	ND	ND	0.5	ND

Table 17: Cleaning Of AprL Variants Compared To SQCBV35 Variant, Reported As PI For Cleaning Assays And Stability As % Remaining Activity For Stability Assays											
Sample ID	Microswatch Cleaning									Stability	
	Persil-Non Bio			Test HDL 2			Test HDL 1			10% PNB	Test HDL 1
	EMPA -116	CFT- CS-38	CFT- C10	EMPA -116	CFT- CS-38	CFT- C10	EMPA -116	CFT- CS-38	CFT- C10	63°C	60°C
SQCBV109	0.92	0.55	1.01	1.01	1.07	0.96	ND	ND	ND	0.4	ND
SQCBV111	1.25	1.21	1.09	2.50	2.13	1.32	ND	ND	ND	0.7	ND
SQCBV112	1.26	1.03	1.34	1.66	1.54	1.12	ND	ND	ND	0.2	ND
SQCBV113	1.12	1.02	1.15	1.52	1.43	1.16	ND	ND	ND	0.2	ND
SQCBV114	1.16	1.11	0.95	1.68	1.52	1.06	ND	ND	ND	0.8	ND
SQCBV119	1.20	1.32	1.40	1.48	1.66	1.41	ND	ND	ND	1.0	ND
SQCBV121	1.22	1.13	1.24	1.36	1.28	1.12	ND	ND	ND	4.8	ND
SQCBV125	1.15	0.85	0.96	1.44	1.34	0.93	ND	ND	ND	1.9	ND
SQCBV143	1.11	0.94	0.93	1.41	1.54	0.79	ND	ND	ND	3.7	ND
SQCBV161	1.02	0.94	0.76	1.46	1.02	0.82	ND	ND	ND	9.6	ND
SQCBV188	0.98	0.70	0.89	1.05	1.03	0.82	ND	ND	ND	16.0	ND
SQCBV191	0.99	1.13	1.09	1.12	0.93	1.05	1.05	1.31	1.03	10.2	34.0
SQCBV192	0.94	1.01	0.98	0.99	0.88	0.94	0.98	1.00	0.91	5.6	27.6
SQCBV193	0.96	1.06	1.02	1.03	0.98	1.03	1.04	1.08	0.97	5.6	21.0
SQCBV194	1.04	1.18	1.03	1.16	1.03	1.01	1.01	1.18	0.92	0.2	3.7
SQCBV195	0.92	0.92	0.94	1.01	0.82	0.97	0.92	0.90	0.84	10.0	31.6
SQCBV196	0.98	0.92	0.90	1.07	1.01	1.00	0.99	1.12	0.81	3.4	19.3
SQCBV197	1.14	1.42	1.18	1.58	1.33	1.11	1.21	1.60	1.13	0.1	2.0
SQCBV198	0.99	1.08	1.05	1.09	1.16	1.07	1.00	1.27	1.01	1.9	11.3
SQCBV199	1.42	1.56	1.70	2.18	2.66	1.46	1.42	1.89	1.66	0.0	0.4
SQCBV200	1.21	1.50	1.21	1.65	1.36	1.22	1.26	1.70	1.21	0.0	0.6
SQCBV201	0.90	0.95	1.07	0.99	0.91	0.97	0.96	0.92	0.84	32.2	55.9
SQCBV202	1.05	1.12	0.98	1.29	1.15	1.08	1.08	1.14	0.96	16.5	42.0
SQCBV203	1.20	1.31	1.23	1.62	1.34	1.20	1.18	1.42	1.11	1.9	10.1
SQCBV204	0.87	0.77	0.81	0.96	0.78	0.94	0.90	0.90	0.72	17.0	42.8
SQCBV205	1.17	1.12	1.00	1.44	1.11	1.07	1.29	1.44	0.94	8.3	27.8
SQCBV206	0.88	0.87	0.89	0.86	0.73	0.95	0.91	1.05	0.89	15.4	41.2
SQCBV207	1.06	1.23	1.13	1.26	1.31	1.08	1.15	1.40	1.10	5.0	24.6
SQCBV208	1.17	1.28	1.33	1.46	1.24	1.20	1.24	1.41	1.25	9.6	36.7
SQCBV209	0.99	1.04	0.97	1.11	0.88	1.02	1.02	0.91	0.99	16.8	43.2
SQCBV211	1.38	1.06	0.92	1.72	1.23	1.08	1.37	1.28	0.82	4.0	19.6
SQCBV212	1.32	0.42	0.72	1.66	1.02	0.86	1.31	0.75	0.61	3.7	22.6
SQCBV213	1.19	1.13	1.04	1.48	1.20	1.10	1.25	1.37	1.01	10.6	34.2
SQCBV214	0.99	0.97	1.00	1.16	0.93	0.99	1.06	1.18	0.95	4.2	22.5
SQCBV215	1.12	0.96	1.37	1.57	1.43	1.21	1.19	1.09	1.35	5.2	31.9
SQCBV216	1.01	0.97	1.00	1.00	0.98	1.00	1.08	1.04	1.04	14.2	41.9
SQCBV217	1.18	1.05	1.03	1.51	0.94	1.07	1.24	1.08	0.98	2.7	20.3
SQCBV218	1.06	1.15	1.05	1.40	1.18	1.14	1.11	1.12	1.02	6.3	29.5
SQCBV219	1.17	1.12	1.25	1.46	1.29	1.29	1.18	1.30	1.28	8.9	36.8
SQCBV221	1.32	0.97	0.86	1.72	0.99	1.01	1.43	1.31	0.88	6.3	29.2
SQCBV223	1.13	0.82	1.07	1.46	1.36	1.12	1.20	1.05	0.99	2.3	21.4
SQCBV225	1.30	1.11	0.94	1.70	0.97	1.08	1.41	1.29	0.99	11.5	34.7
SQCBV226	1.02	0.82	0.97	1.11	1.07	1.04	1.03	0.86	0.93	13.5	38.6

Table 17: Cleaning Of AprL Variants Compared To SQCBV35 Variant, Reported As PI For Cleaning Assays And Stability As % Remaining Activity For Stability Assays											
Sample ID	Microswatch Cleaning									Stability	
	Persil-Non Bio			Test HDL 2			Test HDL 1			10% PNB	Test HDL 1
	EMPA -116	CFT- CS-38	CFT- C10	EMPA -116	CFT- CS-38	CFT- C10	EMPA -116	CFT- CS-38	CFT- C10	63°C	60°C
SQCBV227	1.17	0.88	1.08	1.51	1.24	1.08	1.21	1.13	1.02	13.6	39.4
SQCBV228	1.03	0.77	1.10	1.38	1.21	1.04	1.07	1.10	1.01	17.1	44.0

[00203] Variants of BliD02339 Parent protease (SEQ ID NO:9) (hereinafter “BliD Variants”) set forth in Table 18 were generated as described in Example 2 and tested as described in Example 1 for cleaning performance and stability.

[00204] Table 18 lists BliD Variants that show improvement over BliD Parent in cleaning performance on 3 technical soil microswatches: EMPA-116, CFT CS-38, and CFT-C10, when tested in Persil Non-Bio, and Test HDL 2 laundry detergents, and also showed improved stability when tested in 10% Persil Non-Bio (PNB) laundry detergent (stressed at 63°C for 15-20 min). The cleaning performance and stability results are reported in Table 19. Cleaning performance results are reported as PI values versus the BliD Parent (SEQ ID NO: 9), and stability results are reported as % remaining activity of sample after stress versus unstressed condition. ND means the value was not determined.

Table 18: List of BliD02339 Protease Variants			
Sample ID	Mutations With Respect To BliD02339 Parent	Mutations With Respect To AprL Parent	SEQ ID NO:
SQBLV126	A068S-T077N-N086S-G165Q-K194E-S211N-N217S	A068S-T077N-A128P-G165Q-N217S	152
SQBLV127	A068S-T077N-N086S-V087S-S155G-G165Q-K194E-S211N-N217S	A068S-T077N-V087S-A128P-S155G-G165Q-N217S	153
SQBLV128	A068S-T077N-N086S-G165Q-K194E-S211N	A068S-T077N-A128P-G165Q	154
SQBLV129	A068S-T077N-N086S-V087S-S155G-G165Q-K194E-S211N	A068S-T077N-V087S-A128P-S155G-G165Q	155
SQBLV130	I035A-A068S-T077N-N086S-V087S-V147L-G165Q-K194E-S211N-N217S	I035A-A068S-T077N-V087S-A128P-V147L-G165Q-N217S	156
SQBLV131	A068S-T077N-N086S-V087S-S155G-G165Q-K194E-S211N-N217S-A242G	A068S-T077N-V087S-A128P-S155G-G165Q-N217S-A242G	157
SQBLV132	A068S-T077N-N086S-V087S-V147L-S155E-G165Q-K194E-S211N	A068S-T077N-V087S-A128P-V147L-S155E-G165Q	158
SQBLV133	A068S-T077N-N086S-V087S-S155E-G165Q-K194E-S211N-N217S	A068S-T077N-V087S-A128P-S155E-G165Q-N217S	159
SQBLV134	A068S-T077N-V087S-S155G-G165Q-K194E-S211N-N217S-A242G	A068S-T077N-S086N-V087S-A128P-S155G-G165Q-N217S-A242G	160
SQBLV135	I035A-A068S-T077N-N086S-V147L-G165Q-K194E-S211N-N217S	I035A-A068S-T077N-A128P-V147L-G165Q-N217S	161
SQBLV136	I035A-A068S-T077N-N086S-S155G-G165Q-K194E-S211N-A242G	I035A-A068S-T077N-A128P-S155G-G165Q-A242G	162
SQBLV137	I035A-A068S-T077N-N086S-V147L-	I035A-A068S-T077N-A128P-V147L-	163

Table 18: List of BliD02339 Protease Variants			
Sample ID	Mutations With Respect To BliD02339 Parent	Mutations With Respect To AprL Parent	SEQ ID NO:
	G165Q-K194E-S211N-A242G	G165Q-A242G	
SQBLV138	A068S-T077N-N086S-V087S-S155G-G165Q-K194E-S211N-A242G	A068S-T077N-V087S-A128P-S155G - G165Q-A242G	164

Table 19: Comparison of Cleaning of BliD Variants to BliD Parent, Reported As PI For Cleaning Assays And Stability As % Remaining Activity For Stability Assays							
Sample ID	Persil-Non Bio			Test HDL 2			Stability in 10% PNB
	EMPA-116	CFT-CS-38	CFT-C10	EMPA-116	CFT-CS-38	CFT-C10	63°C
BliD Parent	1.0	1.0	1.0	1.0	1.0	1.0	0.7
SQBLV126	1.7	2.1	2.7	1.2	1.4	3.0	10.3
SQBLV127	1.3	0.9	1.8	0.6	0.8	2.2	45.8
SQBLV128	1.6	2.3	2.4	1.3	1.3	2.8	1.7
SQBLV129	1.2	1.4	1.7	0.8	0.9	2.0	18.3
SQBLV130	1.4	1.7	1.7	1.1	0.9	1.8	37.8
SQBLV131	1.1	0.6	1.3	0.4	0.6	1.6	55.9
SQBLV132	1.5	1.6	1.8	1.2	0.9	2.1	24.2
SQBLV133	1.4	1.7	1.8	1.1	0.9	2.0	41.2
SQBLV134	1.3	2.1	2.2	0.6	0.9	2.3	48.8
SQBLV135	1.6	1.7	2.4	1.1	1.1	2.7	15.5
SQBLV136	1.2	1.1	2.0	0.7	0.9	2.1	13.8
SQBLV137	1.5	2.1	1.9	1.4	1.1	2.3	3.4
SQBLV138	1.4	2.1	2.1	0.7	0.9	2.2	29.7

EXAMPLE 9

Comparison of AprL Protease to Related Molecules

Identification of Homologous Proteases

[00205] Proteins related to the AprL protease (SEQ ID NO:2) were identified by a BLAST search (Altschul et al., Nucleic Acids Res, 25:3389-402, 1997) against the NCBI non-redundant protein database and a subset are shown in Table 20. A similar search was run against the Genome Quest Patent database with search parameters set to default values and a subset are shown in Table 21. The mature amino acid sequence (274 amino acids) for AprL protease (SEQ ID NO:2) was used as the query sequence for both searches. Percent identity (PID) for both search sets is defined as the number of identical residues divided by the number of aligned residues in the pairwise alignment. The column labeled "Sequence Length" corresponds to the length (in amino acids) of the proteins identified by accession number in the column labeled "Accession No.", while "Alignment Length" refers to the length (in amino acids) of the sequence used for alignment and PID calculation.

Table 20: List of Sequences With % Identity to AprL Protein Identified In NCBI Non-redundant Protein Database

Accession No.	PID	Organism	Sequence Length	Alignment Length
CAJ70731.1	100	<i>B. licheniformis</i>	379	274
AEQ38580	99.6	<i>B. licheniformis</i>	370	275
AAG10033/AF283295	99.6	<i>B. licheniformis</i>	310	274
AKI30031	99.6	<i>B. licheniformis</i>	379	274
WP_020450819	99.6	<i>Bacillus paralicheniformis</i>	379	274
AID16241	99.3	<i>B. licheniformis</i>	379	274
AAG00493	98.9	<i>B. licheniformis</i>	310	274
AEU17777	98.9	<i>B. licheniformis</i>	349	274
ADK11044	98.5	<i>B. licheniformis</i>	379	274
OJT55573	98.5	<i>Bacillus licheniformis</i>	378	274
EWB21773	98.2	<i>B. licheniformis</i> S 16	378	274
WP_082247249	98.2	<i>Bacillus</i>	379	274
YP_006712489/ WP_011197728.1	98	<i>B. licheniformis</i> DSM13	379	274
AEU12640	97	<i>B. licheniformis</i>	379	274
CAA62667	95	<i>B. licheniformis</i>	379	273
CAA62666	95.6	<i>B. licheniformis</i>	379	273
WP_026586291	89.1	<i>Bacillus</i> sp. NSP9.1	379	275
WP_077737098	89.1	<i>Bacillus sonorensis</i>	379	275
WP_076761713	88.4	<i>Bacillus</i> sp. NRRL B-41294	379	276
P00781	88	<i>B. licheniformis</i> (subtilisin DY)	274	274
WP_006636716	87	<i>B. sonorensis</i>	378	274
WP_082634659	85.4	<i>Bacillus glycinifermentans</i>	359	274
WP_007497196	73	<i>B. stratosphericus</i>	383	274
ADK11996	72	<i>B. pumilus</i>	383	274
ADN04910	72	<i>B. circulans</i>	275	274
AFP23380	72	<i>B. lehensis</i>	276	274
WP_017360299	72	<i>B. cereus</i>	383	274
YP_003972439/ WP_003327717.1	72	<i>B. atrophaeus</i> 1942	382	275
BAN09118	71	<i>B. subtilis</i>	381	274
WP_010192405	71	<i>Bacillus</i> sp. m3-13	379	276
WP_010333625	71	<i>B. mojavenensis</i>	381	274
AAC43580	70	<i>Bacillus</i> sp. SprC	378	274
ACJ07037	70	<i>B. subtilis</i>	381	274
WP_010329279	70	<i>B. vallismortis</i>	381	274
BAD11988	69	<i>Bacillus</i> sp. KSM-LD1	376	274
BAD21128	69	<i>Bacillus</i> sp. KSM-LD1	377	276
AAC43581	69	<i>Bacillus</i> sp. SprD	379	276
CAA74536	69	<i>B. subtilis</i> str. 168	379	275
CAA24990	69	<i>B. amyloliquefaciens</i>	376	275
AGC81872/ WP_032874368	69	<i>B. velezensis</i>	382	275
ABY25856	68	<i>Geobacillus stearothermophilus</i>	382	275
ADC49870/	62	<i>B pseudofirmus</i>	374	274

Table 20: List of Sequences With % Identity to AprL Protein Identified In NCBI Non-redundant Protein Database

Accession No.	PID	Organism	Sequence Length	Alignment Length
WP_012957236				
AAA22212	60	<i>B. alcalophilus</i>	380	274
WP_062748019	60	<i>B. clausii</i>	380	274
P29600	60	<i>B. lentus</i>	269	274
BAA25184	58	<i>Bacillus sp AprN</i>	379	273
AFK08970	58	<i>B. lehensis</i>	378	273
AAA87324/ P20724	58	<i>B. subtilis</i>	378	273
ADD64465	55	<i>Bacillus sp. JB99</i>	361	274
AGS78407	55	<i>Bacillus gibsonii</i>	375	273
BAA06157.1/ WP_003321226	54.9	Prepro-subtilisin Sendai, <i>Bacillus sp. G-825-6</i>	382	273
BAB04574.1/ WP_010897028	54.7	<i>B. halodurans C-125</i>	361	274
ABI26631	54	<i>B. clausii</i>	361	274
BAA05540/ WP_053432556	54	<i>Bacillus sp.</i>	361	274

Table 21: List Of Sequences With % Identity To AprL Protein Identified In Genome Quest Patent Database

Patent/Patent Appl Pub No-SEQ ID NO	PID	Organism	Sequence Length	Alignment Length
WO2014013080-0002	100.0	<i>B. licheniformis</i>	274	274
CN101215534-0002	99.6	<i>B. licheniformis YPIA</i>	379	274
WO2017006266-0002	99.6	<i>Bacillus licheniformis Synthetic</i>	379	274
WO9739130-0002	99.6	<i>B. licheniformis; strain PWD-1</i>	379	274
US8110391-0009	99.3	<i>B. licheniformis</i>	274	274
JP2013500714-0110	98.9	<i>B. licheniformis</i>	274	274
WO2006122655-0031	98.5	<i>B. licheniformis</i>	379	274
US7087415-0014	98.2	<i>B. licheniformis</i>	379	274
US20110171718-0120	97.8	<i>B. licheniformis</i>	274	273
US20030049619-0014	97.4	<i>B. licheniformis</i>	378	273
JP2013500714-0121	96.0	<i>B. licheniformis</i>	274	273
US7569226-0012	88.3	<i>B. subtilis DY</i>	274	274
US20050009167-0017	88.0	<i>B. subtilis DY</i>	274	274
US20110171718-0098	88.3	<i>B. licheniformis</i>	274	274
WO2008066931-6106	83.6	<i>B. licheniformis; SJ1904 (ATCC PTA-7992)</i>	333	274
JP2013500714-0106	73.0	<i>B. pumilus</i>	275	274
JP2004313043-0002	72.7	<i>Bacillus sp. KSM-9865</i>	275	275
JP2004313043-0001	72.7	<i>Bacillus sp. KSM-KP43</i>	275	275
JP2013500714-0096	70.5	<i>B. pumilus</i>	275	274
WO2011014278-0084	70.6	<i>B. subtilis</i>	275	275

Alignment of Subtilisin Sequences with AprL

[00206] The amino acid sequences of mature AprL (SEQ ID NO: 2) and BliD02339 (SEQ ID NO:9) proteins were aligned with the mature sequences of multiple subtilisins in Tables 20 and 21 using CLUSTALW software (Thompson et al., Nucleic Acids Research, 22:4673-4680, 1994) with the default parameters. This alignment is set forth in Figure 1 with a box drawn around one or more amino acid motifs selected from SEQ ID NOs: 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, and 179.

EXAMPLE 10

Phylogenetic Tree For Subtilisins Including AprL and BliD02339

[00207] A phylogenetic tree for the mature sequences of AprL protein (SEQ ID NO:2), BliD02339 protein (SEQ ID NO: 9), AprL protease variants: SQM-30 (SEQ ID NO: 36), SQCBV 20 (SEQ ID NO: 57), SQCBV 35 (SEQ ID NO:63), SQCBV37 (SEQ ID NO:65), SQCBV188 (SEQ ID NO:105), SQCBV269 (SEQ ID NO:140), SQBLV 134 (SEQ ID NO:160), SQCBV328 (SEQ ID NO:268), SQCBV419 (SEQ ID NO:302), SQCBV555 (SEQ ID NO:358), SQCBV567 (SEQ ID NO:370), and SQCBV582 (SEQ ID NO:384) was built using a subset of the sequences of subtilisins shown in Tables 20 and 21. The sequences were entered in the Vector NTI Advance suite and a Guide Tree was created using the Neighbor Joining method (NJ) (Saitou, N.; and Nei, M. (1987) Mol Biol Evol 4, 406-425). The NJ method works on a matrix of distances between all pairs of sequences to be analyzed. These distances are related to the degree of divergence between the sequences. The Guide Tree is calculated after the sequences are aligned. The tree construction was calculated using the following parameters: Kimura's correction for sequence distance and ignoring positions with gaps. The MEGA 6 program was used to display the phylogenetic tree shown in Figure 2. Based on the phylogenetic tree distribution, certain subtilisins have been grouped as shown in Figure 2 to form the AprL-clade.

EXAMPLE 11

Additional AprL Variants

[00208] AprL variants listed on Table 22 were generated as described in Example 2.

Table 22: Additional AprL Protease Variants		
Sample ID	Sequence Substitutions Relative To AprL Parent	SEQ ID NO:
SQCBV229	A68S-T77N-G127T-A128P-G165Q-N184Q-A202V-N217S	191
SQCBV230	T3V-A68S-T77N-M123I-A128P-G165Q-N184Q-A202V-N217S	193
SQCBV231	A68S-T77N-G127T-A128P-G165Q-N184Q-N217S-S258P	206
SQCBV232	T3V-A68S-T77N-M123I-A128P-G165Q-N184Q-N217S-S258P	207
SQCBV233	A68S-T77N-G127T-A128P-G165Q-N184Q-G203Q-N217S	208
SQCBV234	T3V-A68S-T77N-M123I-A128P-G165Q-N184Q-G203Q-N217S	209
SQCBV235	A68S-T77N-A128P-G165Q-N184Q-A202V-N217S-S258P	210
SQCBV236	T3V-A68S-T77N-M123I-A128P-G165Q-N184Q-A202V-N217S-S258P	211
SQCBV237	A68S-T77N-A128P-G165Q-N184Q-G203Q-N217S-S258P	212

Table 22: Additional AprL Protease Variants		
Sample ID	Sequence Substitutions Relative To AprL Parent	SEQ ID NO:
SQCBV238	T3V-A68S-T77N-M123I-A128P-G165Q-N184Q-G203Q-N217S-S258P	213
SQCBV239	S86H-T115F-G127T-A143V-G165Q-N184Q-A202V-N217S-S258P	214
SQCBV240	T3V-S86H-T115F-M123I-A143V-G165Q-A202V-N217S-S258P	215
SQCBV241	A68S-T77N-A128P-S129H-G165Q-N184Q-A202V-N217S	216
SQCBV242	T3V-A68S-T77N-M123I-A128P-S129H-G165Q-N184Q-A202V-N217S	217
SQCBV243	S86H-T115F-S129H-A143V-G165Q-N184Q-A202V-N217S-S258P	218
SQCBV244	T3V-S86H-T115F-M123I-S129H-A143V-G165Q-A202V-N217S-S258P	219
SQCBV245	A68S-T77N-A128P-S155N-G165Q-N184Q-A202V-N217S	220
SQCBV246	T3V-A68S-T77N-M123I-A128P-S155N-G165Q-N184Q-A202V-N217S	221
SQCBV249	I35A-A68S-T77N-V87S-A128P-V147L-G165Q-A202V-N217S	222
SQCBV250	T3V-I35A-A68S-T77N-V87S-M123I-A128P-V147L-G165Q-A202V-N217S	223
SQCBV251	A68S-T77N-A128P-Q136N-S155N-G165Q-N184Q-N217S-A242N	224
SQCBV252	T3V-A68S-T77N-M123I-A128P-Q136N-S155N-G165Q-N184Q-N217S-A242N	225
SQCBV253	A68S-V71A-T77N-A128P-G165Q-N184Q-A202V-N217S	226
SQCBV254	T3V-A68S-V71A-T77N-M123I-A128P-G165Q-N184Q-A202V-N217S	227
SQCBV255	A68S-T77N-A128P-G165Q-N184Q-S187P-A202V-N217S	228
SQCBV256	T3V-A68S-T77N-M123I-A128P-G165Q-N184Q-S187P-A202V-N217S	229
SQCBV257	A68S-T77N-A128R-G165Q-N184Q-A202V-N217S-S258D	230
SQCBV258	T3V-A68S-T77N-M123I-A128R-G165Q-N184Q-A202V-N217S-S258D	231
SQCBV259	A68S-T77N-A128P-S129R-G165Q-N184Q-N217S-S258D	232
SQCBV260	T3V-A68S-T77N-M123I-A128P-S129R-G165Q-N184Q-N217S-S258D	233
SQCBV261	A68S-T77N-A128R-G165Q-N184Q-A202V-N217S-S258D-K264H	234
SQCBV262	T3V-A68S-T77N-M123I-A128R-G165Q-N184Q-A202V-N217S-S258D-K264H	235
SQCBV263	A68S-T77N-A128R-G165Q-N184Q-G203E-S258D	236
SQCBV264	T3V-A68S-T77N-M123I-A128R-G165Q-N184Q-G203E-S258D	237
SQCBV266	T3V-A68S-T77N-M123I-A128R-G165Q-N184E-A202V-N217S-S258D	238
SQCBV268	T3V-A52R-A68S-T77N-M123I-A128R-G165Q-N184E-A202V-N217S-S258D	239
SQCBV300	T3V-T77N-S100N-G105N-D119Q-A128P-S158N-N160D-T161Q-G165A-N184Q-N217S-S258P-K264S	240
SQCBV301	T3V-T77N-S100N-G105N-A128P-S158D-T161Q-G165A-N184Q-N217S-S258P	241
SQCBV302	T3V-T77N-S100N-G105N-A128P-S158N-N160D-T161Q-G165A-N184Q-N217S-S258P	242
SQCBV303	A24Q-V45I-A68S-T77N-S98R-E111Q-M123I-S158Q-G165E-N184Q-S187E-V204I-T210I-N217S-S259E	243
SQCBV304	P9E-A52R-S86R-E111Q-S129R-S158Q-G165E-V204I-T210I-S258D-S259E-K264Q	244
SQCBV305	T3V-A68S-T77N-G127T-A128P-G165Q-N184Q-A202V-N217S	245
SQCBV306	T3V-I35A-T77N-V87S-M123I-A128P-G165Q-N184Q-A202V-N217S-A242G	246
SQCBV307	T3V-A68S-T77N-G127E-A128P-G165Q-N184Q-A202V-N217S	247
SQCBV308	T3V-A68S-T77N-M123I-G127V-A128P-G165Q-N184Q-N217S	248
SQCBV309	T3V-A68S-T77N-M123I-A128P-A151G-G165Q-N184Q-N217S	249
SQCBV310	T3V-A68S-T77N-G127T-A128P-A151G-G165Q-N184Q-N217S	250
SQCBV311	T3V-T77N-M123I-A128P-A151G-G165Q-N184Q-A202V-N217S	251
SQCBV312	T3V-A68S-T77N-L95F-A128P-G165Q-N184Q-A202V-N217S	252
SQCBV313	T3V-A68S-T77N-L95A-M123I-A128P-G165Q-N184Q-N217S	253
SQCBV314	T3V-A68S-T77N-L95F-M123I-A128P-G165Q-N184Q-N217S	254

Table 22: Additional AprL Protease Variants		
Sample ID	Sequence Substitutions Relative To AprL Parent	SEQ ID NO:
SQCBV315	T3V-A68S-T77N-G99A-A128P-G165Q-N184Q-A202V-N217S	255
SQCBV316	T3V-A68S-T77N-G99E-A128P-G165Q-N184Q-A202V-N217S	256
SQCBV317	T3V-A68S-T77N-G99Q-A128P-G165Q-N184Q-A202V-N217S	257
SQCBV318	T3V-A68S-T77N-G99A-M123I-A128P-G165Q-N184Q-N217S	258
SQCBV319	T3V-A68S-T77N-S108H-A128P-G165Q-N184Q-N217S	259
SQCBV320	T3V-I35A-T77N-V87S-M123I-A128P-V146A-G165Q-N184Q-A202V-N217S-A242G	260
SQCBV321	T3V-A68S-T77N-G127T-A128P-G165E-N184Q-A202V-N217S	261
SQCBV322	T3V-A68S-T77N-M123I-A128P-V146A-G165Q-N184Q-N217S	262
SQCBV323	T3V-A68S-T77N-M123I-A128P-T161D-G165Q-N184Q-A202V-N217S	263
SQCBV324	T3V-V45R-A68S-T77N-S102D-A128P-T161D-G165Q-N184Q-A202V-N217S	264
SQCBV325	T3V-A68S-T77N-G127T-A128P-G165Q-N184Q-A202V-N217S-S258P	265
SQCBV326	T3V-A68S-T77N-M123I-G127T-A128P-G165Q-N184Q-A202V-N217S-S258P	266
SQCBV327	T3V-A68S-T77N-M123I-G127V-A128P-G165Q-N184Q-A202V-N217S-S258P	267
SQCBV328	T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P	268
SQCBV329	T3V-I35A-T77N-V87S-G127T-A128P-G165Q-N184Q-A202V-N217S-A242G-S258P	269
SQCBV330	T3V-T77N-M123I-G127Q-A128P-G165Q-N184Q-A202V-N217S-S258P	270
SQCBV331	T3V-T77N-M123I-G127V-A128P-G165Q-N184Q-A202V-N217S-S258P	271
SQCBV332	T3V-T77N-M123I-G127E-A128P-G165Q-N184Q-A202V-N217S-S258P	272
SQCBV333	T3V-A68S-T77N-M123I-G127T-A128P-G165Q-N184Q-A202V-N217S	273
SQCBV334	T3V-A68S-T77N-M123I-G127T-A128P-G165Q-N184Q-A202V-T210P-N217S	274
SQCBV335	T3V-A68S-T77N-M123I-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P	275
SQCBV336	T3V-A68S-T77N-M123I-G127T-A128P-G165Q-N184Q-A202V-N217S-S258D	276
SQCBV337	L10M-I35A-T78I-V87S-M123I-A202V-S258D-F260W	277
SQCBV338	L10M-I35A-T78I-V87S-M123I-A202V-F260W-K264H	278
SQCBV339	L10M-I35A-T78I-V87S-A202V-S258D	279
SQCBV340	L10M-I35A-T78I-V87S-S187E-A202V	280
SQCBV341	L10M-T78I-V87S-S187E-A202V	281
SQCBV342	L10M-I35A-T78I-S187E-A202V	282
SQCBV400	T3V-L10M-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P	283
SQCBV401	T3V-I35A-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P	284
SQCBV402	T3V-A68S-T77N-T78I-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P	285
SQCBV403	T3V-A68S-T77N-V87S-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P	286
SQCBV404	T3V-A68S-T77N-M123I-G127S-A128P-G165E-N184Q-A202V-N217S-S258P	287
SQCBV405	T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P	288
SQCBV406	T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-L216Q-N217S-S258P	289
SQCBV407	T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-A242G-S258P	290
SQCBV408	T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258D	291

Table 22: Additional AprL Protease Variants		
Sample ID	Sequence Substitutions Relative To AprL Parent	SEQ ID NO:
SQCBV409	T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P-F260W	292
SQCBV410	A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P	293
SQCBV411	T3V-A68S-T77N-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P	294
SQCBV412	T3V-L31I-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P	295
SQCBV413	A68S-T77N-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P	296
SQCBV414	A68S-T77N-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P	297
SQCBV415	A68S-T77N-G127T-A128P-G165Q-N184Q-A202V-N217S-S258P	298
SQCBV416	T3V-A68S-T77N-T78I-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258D	299
SQCBV417	A68S-T77N-T78I-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258D	300
SQCBV418	T3V-A68S-T77N-T78I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P	301
SQCBV419	A68S-T77N-T78I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P	302
SQCBV420	T3V-A68S-T77N-G127S-A128P-G165Q-N184Q-A202V-N217S-S258D	303
SQCBV421	A68S-T77N-G127S-A128P-G165Q-N184Q-A202V-N217S-S258D	304
SQCBV422	T3V-A68S-T77N-T78I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258D	305
SQCBV423	A68S-T77N-T78I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258D	306
SQCBV424	T3V-A68S-T77N-S108H-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P	307
SQCBV500	T3V-A68S-T77N-G99Q-A128P-G165Q-N184Q-A202V-N217S-S258P	308
SQCBV501	T3V-A68S-T77N-G99Q-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P	309
SQCBV502	T3V-A68S-T77N-G127S-A128P-G165E-N184Q-A202V-N217S-S258P	310
SQCBV504	T3V-A68S-T77N-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P	311
SQCBV505	T3V-P9E-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P	312
SQCBV506	T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-S187E-A202V-T210P-N217S-S258P	313
SQCBV507	T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-G203E-T210P-N217S-S258P	314
SQCBV508	T3V-L10M-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P	315
SQCBV509	T3V-P9E-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P	316
SQCBV510	T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-S187E-A202V-N217S-S258P	317
SQCBV512	T3V-P9E-A68S-T77N-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P	318
SQCBV513	T3V-A68S-T77N-G127S-A128P-G165Q-N184Q-S187E-A202V-T210P-N217S-S258P	319
SQCBV514	T3V-A68S-T77N-G127S-A128P-G165Q-N184Q-A202V-G203E-T210P-N217S-S258P	320
SQCBV515	T3V-A68S-T77N-G101A-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P	321
SQCBV516	T3V-L10M-A68S-T77N-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P	322

Table 22: Additional AprL Protease Variants		
Sample ID	Sequence Substitutions Relative To AprL Parent	SEQ ID NO:
SQCBV517	T3V-P9E-A68S-T77N-M123I-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P	323
SQCBV518	T3V-A68S-T77N-M123I-A128P-G165Q-N184Q-S187E-A202V-T210P-N217S-S258P	324
SQCBV519	T3V-A68S-T77N-M123I-A128P-G165Q-N184Q-A202V-G203E-T210P-N217S-S258P	325
SQCBV520	T3V-A68S-T77N-G101A-M123I-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P	326
SQCBV521	T3V-L10M-A68S-T77N-M123I-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P	327
SQCBV522	T3V-L10M-A68S-T77N-G101A-M123I-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P	328
SQCBV523	T3V-P9E-L10M-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P	329
SQCBV524	T3V-P9E-A68S-T77N-G99Q-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P	330
SQCBV526	T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-S243I-S258P	331
SQCBV527	T3V-A68S-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P	332
SQCBV529	T3V-A68S-T77N-M123I-G127S-A128P-N184Q-A202V-N217S-S258P	333
SQCBV530	T3V-A68S-T77N-M123I-G127S-A128P-G165Q-A202V-N217S-S258P	334
SQCBV531	T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-S258P	335
SQCBV532	T3V-A68S-T77E-M123I-G127T-A128P-G165Q-N184Q-A202V-N217S-S258P	336
SQCBV533	T3V-A68S-T77E-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P	337
SQCBV534	T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P-K264H	338
SQCBV535	T3V-A68S-T77N-S102T-M123I-G127S-A128P-G165Q-N184Q-A202V-N211S-N217S-S258P	339
SQCBV536	T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N211S-N217S-S258P	340
SQCBV537	T3V-A68S-T77N-S102T-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P	341
SQCBV538	T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P-S259P	342
SQCBV539	T3V-A68S-T77N-M123I-G127S-A128P-T161Q-G165Q-N184Q-A202V-N217S-S258P	343
SQCBV540	T3V-A68S-T77N-M123I-G127S-A128P-S158Q-G165Q-N184Q-A202V-N217S-S258P	344
SQCBV541	T3V-V45Q-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P	345
SQCBV542	T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A193Q-A202V-N217S-S258P	346
SQCBV543	T3V-A68S-T77N-M123I-G127S-A128P-A143Q-G165Q-N184Q-A202V-N217S-S258P	347
SQCBV544	T3V-A68S-T77N-G117S-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P	348

Table 22: Additional AprL Protease Variants		
Sample ID	Sequence Substitutions Relative To AprL Parent	SEQ ID NO:
SQCBV545	T3V-A48V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P	349
SQCBV546	T3V-A48T-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P	350
SQCBV547	T3V-A29S-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P	351
SQCBV548	T3V-A68S-T77N-G127S-A128P-G165Q-N184Q-A202V-N217S	352
SQCBV549	T3V-A68S-T77N-G127S-A128P-G165Q-N184Q-N217S-S258P	353
SQCBV550	T3V-A68S-T77N-A128P-G165Q-N184Q-A202V-N217S-S258P	354
SQCBV551	A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S	355
SQCBV552	A68S-T77N-M123I-A128P-G165Q-N184Q-A202V-N217S-S258P	356
SQCBV554	T3V-A68S-T77N-G99Q-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P	357
SQCBV555	T3V-A68S-T77N-T78I-V87S-M123I-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P	358
SQCBV556	T3V-A68S-T77N-T78I-M123I-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-A242G-S258P	359
SQCBV557	T3V-V26Q-A68S-T77N-M123I-G127S-A128P-G165E-N184Q-A202V-T210P-N217S-S258P	360
SQCBV558	T3V-P9E-L10M-A68S-T77N-M123I-G127S-A128P-G165E-N184Q-A202V-T210P-N217S-S258P	361
SQCBV559	T3V-P9E-L10M-A68S-T77N-G101A-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P	362
SQCBV560	T3V-A68S-T77N-T78I-M123I-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P	363
SQCBV561	T3V-A68S-T77N-T78I-M123I-G127S-A128P-N160K-G165Q-N184Q-A202V-T210P-N217S-A242G-S258P	364
SQCBV562	T3V-A68S-T77N-V87S-M123I-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P	365
SQCBV563	T3V-P40E-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P	366
SQCBV564	T3V-A68S-T77E-M123I-G127T-A128P-G165E-N184Q-A202V-N217S-P238Q-S258P	367
SQCBV565	T3V-A68S-T77N-G99S-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P	368
SQCBV566	T3V-A68S-T77E-M123I-G127T-G165E-N184Q-A202V-N217S-S258P	369
SQCBV567	A68S-T77N-A128P-G165Q-N184Q-N217S-S258P	370
SQCBV568	T3V-A68S-T77N-G101A-S102T-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P	371
SQCBV569	T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S	372
SQCBV570	T3V-A68S-T77N-T78I-G99S-G127S-A128P-G165Q-T210P-A242G-S258P-K264H	373
SQCBV571	T3V-A68S-T77N-T78I-G99S-M123I-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-A242G-S258P	374
SQCBV572	T3V-A68S-T77N-M123I-G127T-G165E-N184Q-A202V-N217S-P238Q-S258P	375
SQCBV573	T3V-A68S-T77N-T78I-G99S-G127S-A128P-G165Q-T210P-A242G-S258P	376

Table 22: Additional AprL Protease Variants		
Sample ID	Sequence Substitutions Relative To AprL Parent	SEQ ID NO:
SQCBV574	T3V-A68S-T77N-T78I-G99S-M123I-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-A242G-S258D	377
SQCBV576	I35A-M123I-G127T-T210P-S243I	378
SQCBV577	T3V-A68S-T77N-G99S-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P-F260W	379
SQCBV578	T3V-A68S-T77N-T78I-G99S-G101A-G127S-A128P-G165Q-T210P-A242G-S258P	380
SQCBV579	A68S-T77N-G99Q-A128P-G165Q-N184Q-N217S	381
SQCBV580	A68S-T77N-A128P-G165Q-N184Q-A202V-N217S	382
SQCBV581	A68S-T77N-A128P-G165Q-N184Q-N217S-S258D	383
SQCBV582	T3V-A68S-T77N-T78I-G99S-G101A-M123I-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-A242G-S258P	384
SQCBV583	V45R-S98R	385

EXAMPLE 12

Laundry Cleaning Performance And Detergent Stability Of Additional AprL Variants

[00209] AprL Variants listed in Table 22 and SQCBV35 were tested as described in Example 1 for laundry cleaning performance and stability. BMI (EMPA-116) microswatches in Persil Non-Bio, Blue Moon, and Liby detergents (Table 2, assay conditions 6, 10, and 16) were used. Enzyme stability was tested in 10% Persil Non-Bio, Blue Moon, and Liby detergents at 62-65°C (Table 2).

[00210] Table 23 shows the results for cleaning performance in Persil Non-Bio as PI values compared to the SQCBV35 variant (SEQ ID NO:63). Table 23 also shows the results for stability in 10% PNB at 62°C for 15 min, 10% Blue Moon at 64°C for 15 min, and 10% Liby at 65°C for 15 min are reported as % remaining activity of sample after stress versus unstressed condition. Table 24 shows the results for cleaning performance in Persil Non-Bio (PNB), Blue Moon and Liby detergents as performance index (PI) values compared to the SQCBV35 variant (SEQ ID NO:63). Table 24 also shows the results for stability in 10% PNB detergent at 62°C for 15 min, 10% Blue Moon detergent at 64°C for 15 min, and 10% Liby detergent at 65°C for 15 min are reported as % remaining activity of sample after stress versus unstressed condition. Table 25 shows the results for cleaning performance in Persil Non-Bio detergent as PI values compared to the SQCBV35 variant (SEQ ID NO:63) using EMPA-116, CS-38 and CFT-10 stains. Table 25 also shows the results for stability in 10% PNB at 66°C for 20 min, 10% Blue Moon at 68°C for 20 min, and 10% Liby at 69°C for 20 min are reported as % remaining activity of sample after stress versus unstressed condition. Table 26 provides the results for laundry cleaning performance and stability in Liby, Blue Moon and PNB detergents for various AprL variants. Stability conditions we described in Example 1 Table 2. Table 27 provides the performance of various AprL Variants in an aged laundry cleaning assay (performed as described in Example 1: Aged Laundry Cleaning Assay (A)). Aged

cleaning performance is reported as % cleaning activity remaining after a 3 week incubation of samples in 100% Blue Moon detergent at 37°C.

Table 23: Laundry Cleaning Of AprL Variants Compared To SQCBV35, Reported As PI For Cleaning Assay And Stability As % Remaining Activity For Stability Assays

Sample ID	Laundry	Stability		
	PNB BMI	10% Liby 65°C	10% Blue Moon 64°C	10% PNB 62°C
SQCBV35	1.00	21	23	31
SQCBV229	1.27	66	60	63
SQCBV230	1.15	58	66	63
SQCBV231	1.53	35	40	43
SQCBV232	1.23	43	49	47
SQCBV233	1.51	19	19	29
SQCBV234	1.37	25	25	30
SQCBV235	1.13	81	92	87
SQCBV236	1.35	76	77	76
SQCBV237	1.19	60	55	47
SQCBV238	1.40	47	57	56
SQCBV239	1.45	46	45	73
SQCBV240	1.54	42	34	65
SQCBV241	1.09	65	61	73
SQCBV242	1.24	59	56	78
SQCBV243	1.16	43	50	68
SQCBV244	1.36	47	35	79
SQCBV245	1.14	67	58	88
SQCBV246	1.28	64	65	71
SQCBV249	1.08	94	78	82
SQCBV250	0.94	84	87	93
SQCBV251	1.14	39	37	42
SQCBV252	1.08	30	35	59
SQCBV253	1.05	59	53	56
SQCBV254	1.07	53	55	71
SQCBV255	1.18	67	73	92
SQCBV256	1.16	77	58	70
SQCBV257	0.88	85	85	86
SQCBV258	1.19	74	53	95
SQCBV259	1.10	53	52	71
SQCBV260	1.18	49	58	45
SQCBV261	0.96	78	100	96
SQCBV262	1.12	78	78	96
SQCBV263	1.13	59	66	60
SQCBV264	1.24	52	45	50
SQCBV266	1.06	78	89	100
SQCBV268	0.93	75	74	79

Table 24: Laundry Cleaning Of AprL Variants Compared To SQCBV35, Reported As PI For Cleaning Assays And Stability As % Remaining Activity For Stability Assays

Sample ID	HDL/BMI			Stability		
	Liby	Blue Moon	PNB	10% Liby 65°C	10% Blue Moon 64°C	10% PNB 62°C
SQCBV35	1.00	1.00	1.00	32	33	41
SQCBV300	0.52	0.74	0.81	86	89	89
SQCBV301	0.76	0.82	1.07	62	58	48
SQCBV302	0.67	0.77	0.94	71	58	74
SQCBV303	0.57	0.90	0.78	33	44	76
SQCBV304	0.69	0.63	0.59	12	29	20
SQCBV305	1.10	ND	1.21	94	77	88
SQCBV306	0.84	0.95	1.07	99	100	64
SQCBV307	0.89	ND	1.09	80	84	63
SQCBV308	0.75	0.78	0.89	33	36	56
SQCBV309	0.90	0.86	1.08	33	24	37
SQCBV310	0.80	0.82	0.98	26	20	38
SQCBV311	0.73	0.89	1.00	61	58	73
SQCBV312	1.05	0.95	0.97	52	40	66
SQCBV313	0.82	0.81	1.08	49	46	53
SQCBV314	1.03	1.06	1.05	23	11	29
SQCBV315	1.09	1.04	1.12	68	64	72
SQCBV316	1.12	0.96	1.05	80	72	81
SQCBV317	1.25	1.13	1.06	65	75	95
SQCBV318	1.51	1.16	1.06	32	22	36
SQCBV319	0.92	0.90	0.94	33	34	50
SQCBV320	0.96	1.00	1.02	100	81	99
SQCBV321	1.21	0.89	1.25	82	83	94
SQCBV322	1.06	0.99	1.10	42	32	50
SQCBV323	0.86	0.85	1.00	75	56	61
SQCBV324	0.94	0.93	0.99	84	85	85
SQCBV325	1.11	1.00	1.17	80	87	92
SQCBV326	1.04	0.99	1.12	80	93	92
SQCBV327	0.70	0.82	0.82	61	91	100
SQCBV328	1.12	1.01	1.04	100	100	100
SQCBV329	0.94	0.83	0.92	96	100	95
SQCBV330	0.87	0.91	1.07	76	89	82
SQCBV331	0.70	0.77	0.94	63	97	100
SQCBV332	0.73	0.59	0.99	88	100	100
SQCBV333	1.17	1.01	1.21	84	91	72

Table 24: Laundry Cleaning Of AprL Variants Compared To SQCBV35, Reported As PI For Cleaning Assays And Stability As % Remaining Activity For Stability Assays						
Sample ID	HDL/BMI			Stability		
	Liby	Blue Moon	PNB	10% Liby 65°C	10% Blue Moon 64°C	10% PNB 62°C
SQCBV334	1.18	1.19	1.03	84	75	83
SQCBV335	1.01	0.93	0.99	90	99	89
SQCBV336	0.81	0.94	1.15	81	100	100
SQCBV337	0.74	1.22	1.19	98	94	88
SQCBV338	1.18	ND	1.17	72	97	91
SQCBV339	0.80	ND	0.96	76	64	89
SQCBV340	0.87	ND	1.02	59	36	50
SQCBV341	0.84	0.69	1.00	37	19	34
SQCBV342	0.82	ND	1.06	56	35	41

Table 25. Laundry Cleaning Of AprL Variants Compared To SQCBV35 Variant, Reported As PI For Cleaning Assays And Stability As As % Remaining Activity For Stability Assays												
Sample ID	HDL/BMI EMPA-116			HDL/Egg CS-38			HDL/POM CFT-10			Stability		
	Liby	Blue Moon	PNB	Liby	Blue Moon	PNB	Liby	Blue Moon	PNB	10% Liby 69°C	10% Blue Moon 68°C	10% PNB 66°C
SQCBV35	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	<1%	<1%	<1%
SQCBV500	0.97	0.79	0.94	1.58	1.55	1.09	1.16	0.80	0.18	57	49	50
SQCBV501	0.79	0.80	0.89	1.41	1.41	1.01	1.11	0.72	0.12	56	51	48
SQCBV502	1.11	0.82	1.09	1.36	1.42	1.20	1.26	1.31	0.92	57	42	51
SQCBV503	0.92	0.80	0.92	1.36	1.22	0.90	1.86	1.06	0.84	69	50	59
SQCBV504	1.03	0.84	0.91	1.43	1.28	1.09	1.40	1.20	1.30	54	41	51
SQCBV505	0.86	0.64	0.78	1.23	1.33	1.09	1.40	1.02	1.33	87	69	69
SQCBV506	1.10	0.85	0.99	1.45	1.50	1.26	1.91	1.20	0.89	54	31	40
SQCBV507	0.82	0.70	0.97	1.14	1.30	1.07	1.28	0.91	0.57	71	57	58
SQCBV508	1.25	0.89	1.03	1.12	1.17	0.93	1.47	1.03	1.03	64	47	51
SQCBV509	0.92	0.78	0.95	1.15	1.20	0.94	1.47	1.15	0.95	64	51	53
SQCBV510	0.95	0.88	0.87	1.23	1.26	0.96	1.47	0.82	0.61	59	42	48
SQCBV511	1.10	0.81	1.11	1.42	1.35	1.13	1.89	1.09	1.03	61	45	49
SQCBV512	0.84	0.76	0.82	1.06	1.20	0.98	1.33	1.17	1.11	83	84	75
SQCBV513	1.09	0.91	1.07	1.31	1.27	0.99	1.75	1.10	0.95	70	54	62
SQCBV514	1.27	0.98	1.16	1.12	1.21	0.99	1.44	1.19	0.81	81	72	73
SQCBV515	0.67	0.75	0.85	0.62	0.75	0.54	1.38	0.93	0.72	53	32	42
SQCBV516	1.07	0.82	0.90	1.07	1.17	0.83	1.16	1.03	0.84	79	64	65
SQCBV517	0.76	0.72	0.81	0.98	1.07	0.87	1.56	1.02	0.80	79	73	ND

Table 25. Laundry Cleaning Of AprL Variants Compared To SQCBV35 Variant, Reported As PI For Cleaning Assays And Stability As % Remaining Activity For Stability Assays												
Sample ID	HDL/BMI EMPA-116			HDL/Egg CS-38			HDL/POM CFT-10			Stability		
	Liby	Blue Moon	PNB	Liby	Blue Moon	PNB	Liby	Blue Moon	PNB	10% Liby 69°C	10% Blue Moon 68°C	10% PNB 66°C
SQCBV518	0.68	0.64	0.91	1.22	1.20	1.03	1.52	1.07	0.78	61	38	48
SQCBV519	0.77	0.68	0.79	1.32	1.36	1.20	1.23	1.11	0.99	74	61	63
SQCBV520	0.84	0.80	0.94	1.01	1.17	0.82	1.23	1.18	1.04	36	18	26
SQCBV521	0.84	0.77	0.97	1.09	1.24	0.91	1.26	1.06	0.99	68	57	58
SQCBV522	0.63	0.72	0.87	0.81	0.85	0.46	1.60	0.93	0.71	59	39	38
SQCBV523	0.96	0.76	0.81	1.06	1.15	0.97	1.02	1.03	0.77	79	72	66
SQCBV524	0.58	0.78	0.87	1.06	1.14	0.89	1.04	0.62	0.46	86	77	67
SQCBV526	1.09	0.82	0.98	1.34	1.19	0.93	1.78	1.36	0.96	39	20	30
SQCBV527	1.13	0.92	1.04	1.38	1.24	1.21	0.81	1.04	0.76	19	10	16
SQCBV529	1.29	0.94	1.02	1.47	1.27	1.05	0.94	0.96	0.60	14	6	7
SQCBV530	1.03	0.95	1.01	1.14	1.12	1.10	1.08	0.96	0.83	32	21	27
SQCBV531	1.16	0.94	1.01	1.37	1.18	1.04	0.74	0.92	0.90	19	9	15
SQCBV532	0.90	0.78	0.94	1.01	0.94	0.73	1.31	0.71	0.80	73	62	52
SQCBV533	0.95	0.92	0.95	1.14	1.00	0.94	0.82	0.89	0.66	69	58	57
SQCBV534	1.59	1.00	1.12	1.24	1.09	1.01	1.40	1.07	1.09	64	60	55
SQCBV535	1.23	0.93	1.05	1.05	1.00	1.09	1.20	0.89	0.89	53	34	39
SQCBV536	1.07	0.96	1.05	1.02	1.04	1.01	0.63	0.92	1.05	45	36	37
SQCBV537	1.01	0.95	1.09	0.99	0.95	0.91	0.87	0.98	1.32	43	34	39
SQCBV538	0.89	0.82	1.00	1.06	0.90	0.96	0.87	0.99	1.69	59	38	43
SQCBV539	1.02	0.93	1.01	1.14	1.00	0.95	1.13	1.12	1.59	48	36	39
SQCBV540	1.06	0.86	1.10	1.13	1.07	1.04	0.70	0.89	1.03	48	35	41
SQCBV541	1.06	0.95	0.99	1.31	1.08	1.10	1.01	1.03	1.68	38	26	32
SQCBV542	1.18	0.85	0.93	1.32	1.09	1.04	1.16	0.98	1.10	44	30	38
SQCBV543	1.20	0.96	1.27	1.21	1.09	1.15	1.01	1.18	2.05	41	30	36
SQCBV544	1.12	0.87	1.05	1.09	0.95	1.03	1.31	1.13	1.24	42	28	29
SQCBV545	1.30	0.89	1.21	1.22	1.04	1.17	0.83	1.15	1.68	35	22	16
SQCBV546	1.32	0.92	1.06	0.89	0.98	1.01	1.11	0.95	1.44	37	26	30
SQCBV547	0.96	0.91	0.89	1.11	1.01	0.94	1.02	0.96	1.59	22	11	13
SQCBV548	1.25	1.07	0.88	1.33	1.20	1.09	1.28	1.03	0.96	33	19	28
SQCBV549	1.43	1.12	1.01	1.31	1.15	1.18	1.05	1.21	0.94	6	4	6
SQCBV550	0.76	0.91	0.85	0.92	0.96	0.68	0.68	0.84	0.66	71	59	62
SQCBV551	1.14	1.10	1.04	1.08	1.20	1.14	0.77	1.01	1.10	5	ND	5
SQCBV552	0.95	0.89	0.91	1.05	0.94	0.98	0.78	0.87	1.04	26	23	ND

Table 26: Laundry Cleaning Of AprL Variants Compared To SQCBV35, Reported As PI For Cleaning Assays And Stability As % Remaining Activity For Stability Assays						
SAMPLE ID	HDL/BMI EMPA-116			Stability		
	Liby	Blue Moon	PNB	10% Liby, 69°C	10% Blue Moon, 68°C	10% PNB, 66°C
SQCBV35	1.00	1.00	1.00	0.1	0.0	0.1
SQCBV400	1.07	1.05	1.05	69.8	66.9	66.1
SQCBV401	1.11	1.11	1.08	53.9	46.3	51.3
SQCBV402	1.11	1.11	1.06	85.2	86.3	84.3
SQCBV403	1.07	0.99	0.94	68.0	65.7	74.3
SQCBV404	1.07	1.05	1.11	43.3	35.5	49.8
SQCBV405	1.10	1.08	1.04	50.5	47.3	47.2
SQCBV406	1.21	1.18	1.11	21.2	18.7	25.3
SQCBV407	1.20	1.05	1.02	64.1	59.0	62.3
SQCBV408	0.88	1.02	1.05	61.2	57.1	54.1
SQCBV409	0.90	1.00	1.06	64.2	62.8	61.5
SQCBV410	1.07	1.02	1.08	41.1	34.8	41.3
SQCBV411	0.88	0.94	0.90	60.9	59.6	56.5
SQCBV412	1.15	1.00	1.01	49.9	44.6	48.3
SQCBV413	1.07	0.97	0.97	58.3	46.7	60.2
SQCBV414	0.95	0.94	0.98	63.1	49.7	43.8
SQCBV415	1.20	1.12	1.18	59.6	51.3	58.8
SQCBV416	0.88	0.98	1.05	93.3	93.8	91.7
SQCBV417	0.98	0.95	0.98	101.9	69.9	72.6
SQCBV418	0.98	0.96	1.01	71.1	75.2	81.6
SQCBV419	0.94	0.93	0.93	78.3	78.2	72.1
SQCBV420	0.87	0.82	0.90	75.6	57.8	66.2
SQCBV421	0.79	0.87	0.95	56.6	53.1	59.0
SQCBV422	0.98	0.91	0.96	86.7	72.1	83.2
SQCBV423	0.81	0.92	0.99	87.0	74.1	83.0
SQCBV424	0.86	0.86	0.89	77.8	58.5	64.7

Table 27: Aged Laundry Cleaning Performance Of AprL Variants Compared To SQCBV35 Parent, Reported As % Remaining Cleaning Activity	
Sample ID	% Remaining Cleaning Activity After 3 weeks at 37°C
SQCBV35	47
SQCBV505	69
SQCBV512	84
SQCBV514	72
SQCBV516	64

Table 27: Aged Laundry Cleaning Performance Of AprL Variants Compared To SQCBV35 Parent, Reported As % Remaining Cleaning Activity	
Sample ID	% Remaining Cleaning Activity After 3 weeks at 37°C
SQCBV517	73
SQCBV519	61
SQCBV523	72
SQCBV524	77
SQCBV532	62
SQCBV534	60
SQCBV554	67
SQCBV555	93
SQCBV556	71
SQCBV557	66
SQCBV558	65
SQCBV559	81
SQCBV560	67
SQCBV561	72
SQCBV562	62
SQCBV563	63
SQCBV564	65
SQCBV565	64
SQCBV566	83

EXAMPLE 13**ADW Cleaning Performance And Detergent Stability Of Additional AprL Variants**

[00211] The cleaning performance of AprL protease variants listed in Table 22, AprL wt and SQCBV35 was measured in dish based applications (ADW) using various detergent formulas listed on Table 1.1, and pre-punched egg yolk microswatches (PAS-38) that were either pre-rinsed or unrinsed prior to assays. To prepare pre-rinsed PAS-38 swatches, 180µl 10mM CAPS buffer pH 11 was added to MTPs containing microswatches, and plates were sealed and incubated in an iEMS incubator for 30 min at 60°C and 1100 rpm. The buffer was then removed, and the swatches were rinsed with deionized water and air dried. To measure enzyme cleaning in the ADW formulas, 175 µL of the detergent solution (3 g/L detergent prepared in 374 ppm water hardness) was added to each well of the MTPs, followed by the addition of 5 µL of enzyme. The MTP was sealed and incubated in an iEMS incubator for 30 min at 40°C and 1100 rpm. After incubation, 100 µL of wash liquid from each well was transferred to a new MTP, and the absorbance of this supernatant was measured at 405 nm on a SpectraMax plate reader. The net absorbance was obtained by subtracting the value for a blank control (no enzyme) from each sample in

the GSM-B pH 10.5 formula. When using the Custom liquid ADW Detergent Formula (at pH 7 and 9), all the supernatant was removed from the MTP wells after the enzyme incubation and transferred to another MTP to determine the absorbance at 405nm. The swatches in the MTPs were rinsed with deionized water, then 180 μ L 10mM CAPS buffer pH 11 was added per well and the plates were incubated in an iEMS incubator for another 10 min at 40°C and 1100 rpm. After this last incubation, 100 μ L of the resulting wash liquid was transferred to a new MTP, and the absorbance was measured at 405 nm. To determine total cleaning performance in the Custom liquid ADW Detergent Formula (at pH 7 and 9), the two absorbance measurements taken were added, and this total value was used to calculate the performance index (PI) for each enzyme.

[00212] In all instances, the PI was calculated for each condition and protease variant by dividing the blank subtracted total absorbance by that of the parent protease (tested at the same concentration). The value for the parent protease was determined from a standard curve of the parent protease which was included in the test and which was fitted to a Langmuir fit. Table 28 shows the ADW performance results for AprL Variants (using either rinsed or unrinsed swatches) tested in either GSM-B pH 10.5 or Custom Liquid ADW Detergent Formula (adjusted to pH 7 and pH 9). The results are reported as PI values compared to AprL wild-type protease ("AprL wt").

Table 28: ADW Cleaning Performance Of AprL Variants Tested in GSM-B and Custom ADW Formulas, compared To AprL, Reported As PI						
Sample ID	GSMB pH 10.5 Rinsed	GSMB pH 10.5 Unrinsed	Custom ADW pH 7 Rinsed	Custom ADW pH 7 Unrinsed	Custom ADW pH 9 Rinsed	Custom ADW pH 9 Unrinsed
AprL wt	1.0	1.0	1.0	1.0	1.0	1.0
SQCBV35	1.0	1.0	1.0	1.1	1.0	1.0
SQCBV236	1.2	1.4	1.1	1.4	1.0	1.2
SQCBV502	1.0	1.8	1.1	1.4	1.1	1.3
SQCBV503	1.0	1.5	1.1	1.3	1.1	1.4
SQCBV504	1.0	1.3	1.0	1.1	1.1	1.2
SQCBV527	1.0	0.9	1.0	1.3	1.0	1.0
SQCBV535	1.0	1.0	1.0	1.1	0.9	1.2
SQCBV540	0.9	0.8	1.1	1.1	1.0	1.1
SQCBV563	0.9	7.4	1.1	1.7	1.0	2.4
SQCBV564	1.1	1.1	1.0	1.1	0.9	1.1
SQCBV566	1.1	1.3	1.1	1.1	1.1	1.2
SQCBV567	1.4	1.5	1.3	1.3	1.1	1.2
SQCBV568	1.0	1.0	1.1	1.1	1.1	1.2
SQCBV569	1.2	1.2	1.2	1.4	1.0	1.3
SQCBV570	1.2	1.6	1.0	1.4	1.1	1.3

Table 28: ADW Cleaning Performance Of AprL Variants Tested in GSM-B and Custom ADW Formulas, compared To AprL, Reported As PI						
Sample ID	GSMB pH 10.5 Rinsed	GSMB pH 10.5 Unrinsed	Custom ADW pH 7 Rinsed	Custom ADW pH 7 Unrinsed	Custom ADW pH 9 Rinsed	Custom ADW pH 9 Unrinsed
SQCBV571	1.2	1.5	1.0	1.3	1.0	1.1
SQCBV572	1.1	1.3	1.0	1.2	1.0	1.2
SQCBV573	1.0	1.3	0.9	1.3	0.9	1.2
SQCBV574	1.1	1.2	1.0	1.3	1.0	1.2
SQCBV576	1.1	1.4	1.0	1.3	1.0	1.2
SQCBV577	1.1	1.1	1.0	1.3	1.0	1.2
SQCBV578	1.0	1.1	1.1	1.2	1.0	1.1
SQCBV579	0.6	1.1	1.0	1.0	0.9	1.1
SQCBV580	0.9	1.1	1.1	1.1	1.1	1.2
SQCBV581	0.9	1.0	1.0	1.1	1.1	1.1
SQCBV582	1.0	1.1	1.0	1.2	1.0	1.1
SQCBV583	1.4	1.1	0.9	0.9	1.2	1.2

CLAIMS

We claim:

1. A subtilisin variant comprising an amino acid sequence comprising two, three, or four or more variations at positions corresponding to SEQ ID NO:2, wherein said positions are selected from:

(i) 1, 3, 9, 10, 15, 22, 24, 26, 27, 28, 29, 30, 31, 35, 37, 40, 43, 45, 48, 52, 68, 71, 76, 77, 78, 86, 87, 95, 96, 98, 99, 100, 101, 102, 103, 105, 108, 111, 114, 115, 117, 119, 123, 125, 126, 127, 128, 129, 130, 136, 143, 146, 147, 151, 155, 158, 160, 161, 165, 184, 187, 193, 202, 203, 204, 210, 211, 216, 217, 234, 238, 239, 242, 243, 250, 255, 258, 259, 260, 264, and 274;

(ii) 1, 3, 9, 10, 15, 22, 24, 26, 28, 29, 30, 31, 35, 37, 40, 45, 48, 52, 68, 71, 76, 77, 78, 86, 87, 95, 96, 98, 99, 100, 101, 102, 105, 108, 111, 114, 115, 117, 119, 123, 125, 126, 127, 128, 129, 130, 136, 143, 146, 147, 151, 155, 158, 160, 161, 165, 184, 187, 193, 202, 203, 204, 210, 211, 216, 217, 234, 238, 239, 242, 243, 250, 255, 258, 259, 260, 264, and 274;

(iii) 1, 3, 9, 10, 24, 26, 29, 30, 31, 35, 37, 40, 45, 48, 52, 68, 71, 77, 78, 86, 87, 95, 96, 98, 99, 100, 101, 102, 105, 108, 111, 115, 117, 119, 123, 127, 128, 129, 130, 136, 143, 146, 147, 151, 155, 158, 160, 161, 165, 184, 187, 193, 202, 203, 204, 210, 211, 216, 217, 234, 238, 239, 242, 243, 250, 258, 259, 260, 264, and 274;

(iv) 3, 9, 10, 35, 68, 77, 78, 86, 87, 99, 101, 115, 123, 127, 128, 165, 184, 187, 202, 203, 210, 217, 242, 250, 258, and 264; or

(v) 3, 68, 77, 78, 123, 127, 128, 165, 184, 202, 210, 217, and 258;

with the proviso that one or more of said two, three, or four or more variations is a man-made mutation or substitution;

wherein said variant has at least 85% but less than 100% amino acid sequence identity to an AprL-Clade subtilisin enzyme comprising a Motif A and/or a Motif B;

wherein Motif A is selected from Motif 1, Motif 2, Motif 3, and Motif 4;

wherein Motif B is selected from Motif 5, Motif 6, Motif 7, Motif 8, Motif 9, Motif 10, Motif 11, Motif 12, Motif 13, and Motif 14; and

wherein the amino acid positions of the variant are numbered by correspondence with the amino acid sequence of SEQ ID NO:2.

2. The subtilisin variant of claim 1, wherein said variant comprises an amino acid sequence comprising two, three, or four or more amino acid substitutions at two, three, or four or more positions corresponding to SEQ ID NO:2, wherein said substitutions are selected from:

(i) 1Q, 3Q/V, 9E, 10M, 15I, 22Y, 24N/Q, 26Q, 27R, 28A, 29S, 30T, 31I, 35A, 37T, 40E, 43A, 45I/Q/R, 48I/T/V, 52R, 68S, 71A, 76K, 77D/E/H/N/Q/S, 78I, 86H/N/R, 87S/T, 95A/F/Q, 96S, 98H/R,

99A/E/Q/S, 100N, 101A, 102D/T, 103I, 105N, 108H/K, 111Q, 114N, 115F, 117S, 119Q, 123I, 125N, 126Q, 127E/F/Q/S/T/V, 128P/R/S, 129H/R, 130S, 136N/R, 143Q/V, 146A/S, 147L, 151G, 155E/G/N, 158D/N/Q, 160C/D/K, 161D/Q, 165A/E/Q/S, 184E/Q, 187E/P, 193D/Q, 202V, 203E/N/Q, 204I, 210E/H/I/P, 211S, 216Q, 217S, 234I/W, 238Q/R, 239K/S/T, 242G/N, 243I, 250G, 255R, 258D/E/P, 259E/P, 260W, 264H/Q/S, and 274A;

(ii) 1Q, 3Q/V, 9E, 10M, 15I, 22Y, 24N/Q, 26Q, 28A, 29S, 30T, 31I, 35A, 37T, 40E, 45I/Q/R, 48I/T/V, 52R, 68S, 71A, 76K, 77D/E/H/N/S, 78I, 86H/N/R, 87S/T, 95A/F/Q, 96S, 98K/R, 99A/E/Q/S, 100N, 101A, 102D/T, 105N, 108H/K, 111Q, 114N, 115F, 117S, 119Q, 123I, 125N, 126Q, 127E/F/Q/S/T/V, 128P/R/S, 129H/R, 130S, 136N, 143Q/V, 146A/S, 147L, 151G, 155E/G/N, 158D/N/Q, 160D/K, 161D/Q, 165A/E/Q/S, 184E/Q, 187E/P, 193D/Q, 194E, 202V, 203E/N/Q, 204I, 210I/P, 211S, 216Q, 217S, 234I/W, 238Q/R, 239K/S/T, 242G/N, 243I, 250G, 255R, 258D/P, 259E/P, 260W, 264H/Q/S, and 274A;

(iii) 1Q, 3V, 9E, 10M, 24Q, 26Q, 29S, 30T, 31I, 35A, 37T, 40E, 45I/Q/R, 48T/V, 52R, 68S, 71A, 77E/N, 78I, 86H/R, 87S, 95A/F, 96S, 98R, 99A/E/Q/S, 100N, 101A, 102D/T, 105N, 108H, 111Q, 115F, 117S, 119Q, 123I, 127E/Q/S/T/V, 128P/R, 129H/R, 130S, 136N, 143Q/V, 146A, 147L, 151G, 155G/N, 158D/N/Q, 160D/K, 161D/Q, 165A/E/Q, 184E/Q, 187E/P, 193Q, 202V, 203E/N/Q, 204I, 210I/P, 211S, 216Q, 217S, 234W, 238Q, 239S, 242G/N, 243I, 250G, 258D/P, 259E/P, 260W, 264H/Q/S, and 274A;

(iv) 3V, 9E, 10M, 35A, 68S, 77N, 78I, 86H, 87S, 99Q/S, 101A, 115F, 123I, 127S/T, 128P/R, 165E/Q, 184Q, 187E, 202V, 203E, 210P, 217S, 242G, 250G, 258D/P, and 264H; or

(v) 3V, 68S, 77N, 78I, 123I, 127S, 128P, 165Q, 184Q, 202V, 210P, 217S, and 258D/P.

3. The subtilisin variant of claim 1 or 2, wherein said variant comprises an amino acid sequence comprising two, three, or four or more amino acid substitutions selected from:

(i) A/G1Q, T3Q/V, P/S9E, L/Q10M, K15I, K22Y, A24N/Q, V26Q, K27R, G/V28A, A/G29S, I/V30T, I/L31I, I35A, A/S37T, P/S/T40E, N/K43A, S/V45I/Q/R, A48I/T/V, A/S52R, A68S, V71A, N76K, S/T77D/E/H/N/Q/S, T78I, N/S86H/N/R, V87S/T, L95A/F/Q, N96S, S98H/R, G99A/E/Q/S, S100N, G101A, S/T102D/T, Y103I, A/G105N, S108H/K, E111Q, T114N, A/Q/S/T115F, G/N117S, D119Q, M123I, L125N, G126Q, G127E/F/Q/S/T/V, A/P/S128P/R/S, S129H/R, G/V130S, Q136N/R, A/S143Q/V, A/I/V146A/S, V147L, A/S151G, S155E/G/N, S158D/N/Q, N/S/Y160C/D/K, A/S/Q/T161D/Q, G165A/E/Q/S, N/Q184E/Q, S187E/P, A/P/S193D/Q, A/S202V, A/G/S203E/N/Q, V204I, S/T210E/H/I/P, N211S, L216Q, N217S, L234I/W, P238Q/R, A/N/T239K/S/T, A242G/N, S243I, S250G, N/Y255R, D/S258D/E/P, S259E/P, F260W, K/N/R264H/Q/S, and Q274A;

(ii) A1Q, T3Q/V, P9E, L10M, K15I, K22Y, A24N/Q, V26Q, K27R, V28A, A29S, V30T, L31I, I35A, A37T, P40E, N43A, V45I/Q/R, A48I/T/V, A52R, A68S, V71A, N76K, S77D/E/H/N/Q/S, T78I,

S86H/N/R, V87S/T, L95A/F/Q, N96S, S98H/R, G99A/E/Q/S, S100N, G101A, S102D/T, Y103I, G105N, S108H/K, E111Q, T114N, T115F, G117S, D119Q, M123I, L125N, G126Q, G127E/F/Q/S/T/V, A128P/R/S, S129H/R, G130S, Q136N/R, A143Q/V, V146A/S, V147L, A151G, S155E/G/N, S158D/N/Q, N160C/D/K, T161D/Q, G165A/E/Q/S, N184E/Q, S187E/P, A193D/Q, A202V, G203E/N/Q, V204I, T210E/H/I/P, N211S, L216Q, N217S, L234I/W, P238Q/R, N239K/S/T, A242G/N, S243I, S250G, Y255R, S258D/E/P, S259E/P, F260W, K264H/Q/S, and Q274A;

(iii) A/G1Q, T3Q/V, P/S9E, L/Q10M, K15I, K22Y, A24N/Q, V26Q, G/V28A, A/G29S, I/V30T, I/L31I, I35A, A/S37T, P/S/T40E, S/V45I/Q/R, A48I/T/V, A/S52R, A68S, V71A, N76K, S/T77D/E/H/N/S, T78I, N/S86H/N/R, V87S/T, L95A/F/Q, N96S, S98R, G99A/E/Q/S, S100N, G101A, S/T102D/T, A/G105N, S108H/K, E111Q, T114N, A/Q/S/T115F, G/N117S, D119Q, M123I, L125N, G126Q, G127E/F/Q/S/T/V, A/P/S128P/R/S, S129H/R, G/V130S, Q136N, A/S143Q/V, A/I/V146A/S, V147L, A/S151G, S155E/G/N, S158D/N/Q, N/S/Y160D/K, A/S/Q/T161D/Q, G165A/E/Q/S, N/Q184E/Q, S187E/P, A/P/S193D/Q, A/S202V, A/G/S203E/N/Q, V204I, S/T210I/P, N211S, L216Q, N217S, L234I/W, P238Q/R, A/N/T239K/S/T, A242G/N, S243I, S250G, N/Y255R, D/S258D/P, S259E/P, F260W, K/N/R264H/Q/S, and Q274A;

(iv) A1Q, T3Q/V, P9E, L10M, K15I, K22Y, A24N/Q, V26Q, V28A, A29S, V30T, L31I, I35A, A37T, P40E, V45I/Q/R, A48I/T/V, A52R, A68S, V71A, N76K, T77D/E/H/N/S, T78I, S86H/N/R, V87S/T, L95A/F/Q, N96S, S98R, G99A/E/Q/S, S100N, G101A, S102D/T, G105N, S108H/K, E111Q, T114N, T115F, G117S, D119Q, M123I, L125N, G126Q, G127E/F/Q/S/T/V, A128P/R/S, S129H/R, G130S, Q136N, A143Q/V, V146A/S, V147L, A151G, S155E/G/N, S158D/N/Q, N160D/K, T161D/Q, G165A/E/Q/S, N184E/Q, S187E/P, A193D/Q, A202V, G203E/N/Q, V204I, T210I/P, N211S, L216Q, N217S, L234I/W, P238Q/R, N239K/S/T, A242G/N, S243I, S250G, Y255R, S258D/P, S259E/P, F260W, K264H/Q/S, and Q274A;

(v) A/G1Q, T3V, P/S9E, L/Q10M, A24Q, V26Q, A/G29S, I/V30T, I/L31I, I35A, A/S37T, P/S/T40E, S/V45I/Q/R, A48T/V, A/S52R, A68S, V71A, S/T77E/N, T78I, N/S86H/R, V87S, L95A/F, N96S, S98R, G99A/E/Q/S, S100N, G101A, S/T102D/T, A/G105N, S108H, E111Q, A/Q/S/T115F, G/N117S, D119Q, M123I, G127E/Q/S/T/V, A/P/S128P/R, S129H/R, G/V130S, Q136N, A/S143Q/V, A/I/V146A, V147L, A/S151G, S155G/N, S158D/N/Q, N/S/Y160D/K, A/S/Q/T161D/Q, G165A/E/Q, N184E/Q, S187E/P, A/P/S193Q, A/S202V, A/G/S203E/N/Q, V204I, S/T210I/P, N211S, L216Q, N217S, L234W, P238Q, A/N/T239S, A242G/N, S243I, S250G, D/S258D/P, S259E/P, F260W, K/N/R264H/Q/S, and Q274A;

(vi) A1Q, T3V, P9E, L10M, A24Q, V26Q, A29S, V30T, L31I, I35A, A37T, P40E, V45I/Q/R, A48T/V, A52R, A68S, V71A, T77E/N, T78I, S86H/R, V87S, L95A/F, N96S, S98R, G99A/E/Q/S, S100N, G101A, S102D/T, G105N, S108H, E111Q, T115F, G117S, D119Q, M123I, G127E/Q/S/T/V,

A128P/R, S129H/R, G130S, Q136N, A143Q/V, V146A, V147L, A151G, S155G/N, S158D/N/Q, N160D/K, T161D/Q, G165A/E/Q, N184E/Q, S187E/P, A193Q, A202V, G203E/N/Q, V204I, T210I/P, N211S, L216Q, N217S, L234W, P238Q, N239S, A242G/N, S243I, S250G, S258D/P, S259E/P, F260W, K264H/Q/S, and Q274A;

(vii) T3V, P/S9E, L/Q10M, I35A, A68S, S/T77N, T78I, N/S86H, V87S, G99Q/S, G101A, A/Q/S/T115F, M123I, G127S/T, A/P/S128P/R, G165E/Q, N184Q, S187E, A/S202V, A/G/S203E, S/T210P, N217S, A242G, S250G, D/S258D/P, and K/N/R264H;

(viii) T3V, P9E, L10M, I35A, A68S, T77N, T78I, S86H, V87S, G99Q/S, G101A, T115F, M123I, G127S/T, A128P/R, G165E/Q, N184Q, S187E, A202V, G203E, T210P, N217S, A242G, S250G, S258D/P, and K264H;

(ix) T3V, A68S, S/T77N, T78I, M123I, G127S, A/S128P, G165Q, N184Q, A/S202V, S/T210P, N217S, and D/S258D/P; or

(x) T3V, A68S, T77N, T78I, M123I, G127S, A128P, G165Q, N184Q, A202V, T210P, N217S, and S258D/P.

4. The subtilisin variant of claim 1, with the proviso:

(i) that said variations at positions corresponding to SEQ ID NO:2 are not an insertion or deletion;

(ii) that the variation is not a proline when the amino acid sequence of said variant comprises a variation at one or more corresponding SEQ ID NO:2 positions selected from positions 37, 187, 193, and 258;

(iii) that said variation is not 26R, 38S, 234R, or 250E/N when the amino acid sequence of said variant comprises a variation at two or more corresponding SEQ ID NO:2 positions selected from positions 26, 38, 234, and 250;

(iv) that said variation at a position corresponding to SEQ ID NO:2 is not at position 130 or 193 when the amino acid sequence of said variant comprises a variation at position 128; and/or

(v) that said variant does not consist of the amino acid sequence set forth as SEQ ID NO:5 in WO2016074925 or as SEQ ID NO:110 in JP201300714.

5. The subtilisin variant of any preceding claim, with the proviso:

(i) that said substitution is not proline when that amino acid sequence of said variant comprises a substitution at one or more of positions 37, 187, 193, and 258;

(ii) that said substitution is not 26R, 38S, 234R, or 250E/N when the amino acid sequence of said variant comprises a substitution at two or more of positions 26, 38, 234, and 250;

(iii) that said substitution is not at position 130 or 193 when the amino acid sequence of said variant comprises a substitution at position 128; and/or

(iv) that said variant does not consist of the amino acid sequence set forth as SEQ ID NO:5 in WO2016074925 or as SEQ ID NO:110 in JP201300714.

6. The subtilisin variant of any preceding claim, wherein said variant comprises an amino acid sequence comprising one or more amino acid substitution sets selected from:

(i) T77N-G165Q, T77N-N217S, G165Q-N217S, A68S-T77N, N184Q-N217S, A68S-G165Q, G165Q-N184Q, T77N-A128P, T77N-N184Q, A68S-N217S, A128P-N217S, A128P-G165Q, A68S-A128P, A68S-N184Q, A128P-N184Q, T3V-N217S, A202V-N217S, T3V-T77N, T3V-N184Q, N184Q-A202V, G165Q-A202V, M123I-G165Q, T3V-G165Q, T3V-A128P, T3V-A68S, T77N-A202V, M123I-N217S, A68S-A202V, M123I-N184Q, A128P-A202V, T77N-M123I, T3V-A202V, N217S-S258P, A68S-M123I, N184Q-S258P, G165Q-S258P, M123I-A128P, T77N-S258P, T3V-M123I, A128P-S258P, A202V-S258P, A68S-S258P, T3V-S258P, M123I-A202V, M123I-S258P, A68S-G127S, G127S-A128P, T77N-G127S, G127S-A202V, G127S-N184Q, G127S-N217S, G127S-G165Q, T3V-G127S, G127S-S258P, M123I-G127S, T77N-T210P, G165Q-T210P, T210P-N217S, A128P-T210P, A68S-T210P, N184Q-T210P, T210P-S258P, A202V-T210P, M123I-T210P, T3V-T210P, G127S-T210P, G127T-N184Q, G127T-N217S, V87S-G165Q, M123I-Q136R, A68S-G127T, G127T-A128P, T77N-V87S, G127T-G165Q, T77N-G127T, T78I-A202V, A68S-S258D, G165Q-S258D, T77N-S258D, T77N-T78I, T78I-G165Q, N184Q-S258D, N217S-S258D, P9E-M123I, A128P-A242G, A202V-S258D, A68S-T78I, G165Q-A242G, T77N-A242G, T78I-A128P, T78I-G127S, P9E-G165Q, Q136R-G165Q, T78I-N184Q, V87S-A128P, V87S-N217S, A68S-V87S, G127T-A202V, I35A-T77N, T3V-T78I, T78I-N217S, A68S-A242G, A68S-G99S, G99S-A128P, G99S-G165Q, I35A-G165Q, L10M-A202V, T3V-G127T, T77N-G99S, T78I-S258P, V45R-M123I, V87S-M123I, M123I-N239S, P9E-N184Q, S155G-G165Q, T78I-M123I, V45R-Q136R, V87S-A202V, A128P-S258D, I35A-A202V, M123I-G127T, M123I-S258D, N184Q-N239S, P9E-N217S, Q136R-S258E, S86H-S258P, T3V-A242G, T3V-S258D, G99S-N184Q, G99S-N217S, I35A-V87S, N217S-A242G, P9E-Q136R, G127T-S258P, I35A-A128P, I35A-A68S, I35A-N217S, L10M-M123I, P9E-S258P, P9E-T210P, P9E-T77N, S86H-A202V, S86H-G165Q, S86H-N184Q, T77N-S155G, T77N-V147L, T78I-T210P, V87S-S155G, A202V-A242G, A242G-S258P, A68S-G165E, G127S-A242G, G165E-N184Q, G165E-N217S, M123I-S258E, N184Q-A242G, S86H-T115F, T3V-G99S, V147L-G165Q, A128R-G165Q, A128R-S258D, A68S-A128R, A68S-V147L, G127S-S258D, G165E-A202V, G165Q-S258E, G165S-T210P, G203E-T210P, G99S-G127S, G99S-M123I, G99S-T210P, L10M-A128P, L10M-A68S, L10M-N184Q, L10M-N217S, L10M-S258P, L10M-T77N, M123I-A242G, M123I-G165E, M123I-S187P, N239S-S258P, P9E-A128P, P9E-A202V, P9E-A68S, P9E-V45R,

Q136R-N184Q, Q136R-N239S, S86H-N217S, T115F-S258P, T210P-A242G, T3V-G165E, T3V-L10M, T3V-P9E, T77N-A128R, T77N-G165E, T78I-A242G, V45R-G165Q, V87T-M123I, V87T-N184Q, V87T-N239S, V87T-Q136R, A68S-G101A, A68S-V71A, G101A-A128P, G101A-G165Q, G101A-S258P, G165A-N184Q, G165E-S258P, G99S-S258P, I35A-M123I, K15I-G165S, K15I-T210P, L10M-G127S, L10M-G165Q, Q136R-S187P, S187E-A202V, S187P-S258P, S250G-S258P, T115F-A202V, T115F-G165Q, T3V-G101A, T3V-I35A, T3V-V87S, T77N-G101A, T77N-G203E, T78I-S258D, V71A-G165Q, V71A-T77N, V87S-S258P, V87S-V147L, A128P-G165E, A128P-S155G, A128P-V147L, A128R-A202V, A128R-N184Q, A128R-N217S, A1Q-S86H, A68S-G99Q, A68S-S155G, G101A-A202V, G101A-G127S, G101A-N184Q, G101A-N217S, G165Q-G203E, G165Q-N239S, G165S-N239S, G99Q-A128P, G99Q-G165Q, G99Q-N184Q, G99Q-N217S, G99S-A242G, I35A-N184Q, I35A-T78I, I35A-V147L, L10M-T210P, L10M-T78I, M123I-G165S, N184Q-G203E, N184Q-S187P, N43A-M123I, P9E-G127S, P9E-S258E, Q136R-S258P, T115F-N217S, T3V-S86H, T77H-M123I, T77H-N184Q, T77H-N239S, T77H-Q136R, T77H-V87T, T77N-G99Q, T77N-S250G, T78I-G99S, T78I-V87S, V146S-G165S, V146S-N239S, V146S-T210P, V45R-S258E, V87S-A242G, V87S-N184Q, A128S-T210P, A143V-A202V, A143V-G165Q, A143V-S258P, A1Q-N184Q, A1Q-S258P, A68S-G203E, A68S-S187E, A68S-T114N, G101A-M123I, G101A-T210P, G165A-N217S, G165Q-S187P, G99Q-A202V, G99S-A202V, I35A-A242G, I35A-S258P, L10M-I35A, L10M-V87S, M123I-A128R, M123I-V146S, N184Q-S187E, N217S-L234W, N239S-S250G, N43A-Q136R, N43A-V45R, P9E-N239S, P9E-V87T, S155G-N217S, S187E-N217S, S86H-A143V, S86H-Q274A, S86H-S250G, T114N-A128P, T114N-G165Q, T114N-N184Q, T114N-N217S, T115F-A143V, T115F-N184Q, T210P-N239S, T210P-S250G, T3V-A128R, T3V-G99Q, T77H-G165Q, T77N-S187E, T77N-T114N, V147L-N217S, V45R-N184Q, V71A-N184Q, V71A-N217S, V71A-T210P, T77N-G165Q-N217S, A68S-T77N-G165Q, T77N-N184Q-N217S, A68S-T77N-N217S, T77N-A128P-G165Q, T77N-A128P-N217S, A68S-T77N-A128P, G165Q-N184Q-N217S, T77N-G165Q-N184Q, A68S-A128P-G165Q, A68S-G165Q-N217S, A128P-N184Q-N217S, A128P-G165Q-N217S, A68S-A128P-N217S, A68S-N184Q-N217S, A68S-T77N-N184Q, T77N-A128P-N184Q, A68S-G165Q-N184Q, A128P-G165Q-N184Q, A68S-A128P-N184Q, T3V-N184Q-N217S, T3V-T77N-N217S, N184Q-A202V-N217S, T3V-T77N-N184Q, T3V-A128P-N217S, T3V-T77N-A128P, G165Q-A202V-N217S, T3V-A128P-N184Q, T77N-A202V-N217S, T3V-G165Q-N217S, T3V-T77N-G165Q, T3V-A68S-N217S, T3V-A68S-T77N, A68S-A202V-N217S, T3V-G165Q-N184Q, A128P-A202V-N217S, G165Q-N184Q-A202V, T77N-G165Q-A202V, T77N-N184Q-A202V, T3V-A128P-G165Q, T3V-A68S-A128P, T3V-A68S-N184Q, A128P-N184Q-A202V, A68S-N184Q-A202V, A68S-T77N-A202V, T3V-A68S-G165Q, T77N-A128P-A202V, A128P-G165Q-A202V, A68S-A128P-A202V, A68S-G165Q-A202V, M123I-G165Q-N217S, M123I-N184Q-N217S, T77N-M123I-G165Q, T77N-M123I-N217S, T3V-A202V-N217S, M123I-G165Q-N184Q, T77N-M123I-

N184Q, T3V-N184Q-A202V, A68S-M123I-N217S, T3V-T77N-A202V, T3V-A128P-A202V, T3V-G165Q-A202V, M123I-A128P-N217S, T3V-A68S-A202V, A68S-M123I-N184Q, A68S-T77N-M123I, M123I-A128P-N184Q, T3V-M123I-N217S, T77N-M123I-A128P, A68S-M123I-G165Q, M123I-A128P-G165Q, A68S-M123I-A128P, N184Q-N217S-S258P, T3V-M123I-N184Q, T3V-M123I-G165Q, T3V-T77N-M123I, T77N-N217S-S258P, G165Q-N217S-S258P, T3V-A68S-M123I, T3V-M123I-A128P, T77N-N184Q-S258P, A128P-N184Q-S258P, A128P-N217S-S258P, T77N-A128P-S258P, T77N-G165Q-S258P, A202V-N217S-S258P, G165Q-N184Q-S258P, N184Q-A202V-S258P, A68S-A128P-S258P, A68S-N184Q-S258P, A68S-N217S-S258P, A68S-T77N-S258P, M123I-A202V-N217S, T3V-N217S-S258P, A128P-G165Q-S258P, G165Q-A202V-S258P, T3V-A128P-S258P, T3V-N184Q-S258P, A68S-G165Q-S258P, T3V-M123I-A202V, T3V-T77N-S258P, A128P-A202V-S258P, M123I-N184Q-A202V, T77N-A202V-S258P, A68S-A202V-S258P, M123I-G165Q-A202V, T77N-M123I-A202V, T3V-A68S-S258P, A68S-M123I-A202V, M123I-A128P-A202V, T3V-A202V-S258P, T3V-G165Q-S258P, A68S-G127S-A128P, A68S-T77N-G127S, T77N-G127S-A128P, A68S-G127S-A202V, A68S-G127S-N184Q, A68S-G127S-N217S, G127S-A128P-A202V, G127S-A128P-N184Q, G127S-A128P-N217S, A68S-G127S-G165Q, G127S-A128P-G165Q, G127S-A202V-N217S, G127S-N184Q-A202V, G127S-N184Q-N217S, M123I-N217S-S258P, T77N-G127S-A202V, T77N-G127S-N184Q, T77N-G127S-N217S, M123I-N184Q-S258P, T77N-G127S-G165Q, T3V-A68S-G127S, T3V-G127S-A128P, G127S-G165Q-A202V, G127S-G165Q-N184Q, G127S-G165Q-N217S, M123I-A202V-S258P, M123I-G165Q-S258P, T3V-M123I-S258P, T3V-T77N-G127S, A68S-G127S-S258P, G127S-A128P-S258P, A68S-M123I-S258P, M123I-A128P-S258P, T3V-G127S-A202V, T3V-G127S-N184Q, T3V-G127S-N217S, T77N-M123I-S258P, T3V-G127S-G165Q, T77N-G127S-S258P, G127S-A202V-S258P, G127S-N184Q-S258P, G127S-N217S-S258P, T3V-G127S-S258P, G127S-G165Q-S258P, A68S-M123I-G127S, M123I-G127S-A128P, M123I-G127S-A202V, M123I-G127S-N184Q, M123I-G127S-N217S, T77N-M123I-G127S, T3V-M123I-G127S, M123I-G127S-G165Q, M123I-G127S-S258P, T77N-G165Q-T210P, A68S-T77N-T210P, A68S-G165Q-T210P, A68S-A128P-T210P, T77N-A128P-T210P, T77N-T210P-N217S, A68S-N184Q-T210P, N184Q-T210P-N217S, T77N-N184Q-T210P, A128P-G165Q-T210P, A68S-T210P-N217S, G165Q-N184Q-T210P, G165Q-T210P-N217S, A128P-N184Q-T210P, A128P-T210P-N217S, T77N-T210P-S258P, G165Q-T210P-S258P, T3V-A128P-T210P, T3V-A68S-T210P, T3V-T77N-T210P, A128P-T210P-S258P, A202V-T210P-N217S, A68S-T210P-S258P, N184Q-A202V-T210P, T210P-N217S-S258P, T3V-G165Q-T210P, T77N-A202V-T210P, A202V-T210P-S258P, G165Q-A202V-T210P, T3V-N184Q-T210P, A128P-A202V-T210P, A68S-A202V-T210P, N184Q-T210P-S258P, T3V-A202V-T210P, T3V-T210P-N217S, T3V-T210P-S258P, M123I-T210P-N217S, T77N-M123I-T210P, A68S-M123I-T210P, M123I-A128P-T210P, M123I-N184Q-T210P, A68S-G127S-T210P, G127S-A128P-T210P, T77N-G127S-T210P, G127S-T210P-S258P, M123I-G165Q-T210P, T3V-

G127S-T210P, G127S-G165Q-T210P, T3V-M123I-T210P, G127S-A202V-T210P, G127S-N184Q-T210P, G127S-T210P-N217S, G127T-N184Q-N217S, M123I-A202V-T210P, M123I-T210P-S258P, A68S-G127T-N184Q, A68S-G127T-N217S, G127T-A128P-N184Q, G127T-A128P-N217S, T77N-V87S-G165Q, A68S-G127T-A128P, G127T-G165Q-N184Q, G127T-G165Q-N217S, T77N-G127T-N184Q, T77N-G127T-N217S, A68S-G165Q-S258D, A68S-T77N-G127T, A68S-T77N-S258D, G127T-A128P-G165Q, T77N-G127T-A128P, and T77N-G165Q-S258D; or

(ii) N184Q-N217S, T77N-N217S, T77N-N184Q, T77N-G165Q, G165Q-N184Q, G165Q-N217S, A128P-N217S, A68S-T77N, T77N-A128P, A68S-N217S, A128P-N184Q, A68S-N184Q, A68S-G165Q, A128P-G165Q, A68S-A128P, A202V-N217S, N184Q-A202V, G165Q-A202V, T3V-N217S, T3V-N184Q, T3V-T77N, T77N-A202V, A68S-A202V, T3V-A128P, T3V-G165Q, A128P-A202V, T3V-A68S, T3V-A202V, N184Q-S258P, N217S-S258P, M123I-N217S, M123I-N184Q, T3V-M123I, T77N-S258P, A128P-S258P, G165Q-S258P, T77N-M123I, A202V-S258P, M123I-G165Q, A68S-M123I, A68S-S258P, M123I-A128P, M123I-A202V, T3V-S258P, A68S-G127S, G127S-A128P, T77N-G127S, G127S-A202V, G127S-N184Q, G127S-N217S, G127S-G165Q, M123I-S258P, T3V-G127S, G127S-S258P, M123I-G127S, T77N-T210P, A128P-T210P, A202V-T210P, A68S-T210P, G165Q-T210P, T210P-S258P, T3V-T210P, N184Q-T210P, T210P-N217S, G127S-T210P, M123I-T210P, T78I-A202V, A68S-S258D, G165Q-S258D, T77N-S258D, T77N-T78I, T78I-G165Q, G127T-N184Q, G127T-N217S, N184Q-S258D, N217S-S258D, A202V-S258D, A68S-T78I, T78I-A128P, T78I-G127S, A68S-G127T, T78I-N184Q, G127T-A128P, G127T-A202V, T3V-T78I, T78I-N217S, G127T-G165Q, L10M-A202V, T77N-G127T, T78I-S258P, T3V-G127T, T78I-M123I, V87S-A202V, A128P-A242G, A128P-S258D, G165Q-A242G, I35A-A202V, M123I-S258D, T3V-A242G, T3V-S258D, T77N-A242G, S86H-S258P, G127T-S258P, P9E-T77N, S86H-A202V, S86H-G165Q, S86H-N184Q, T78I-T210P, A202V-A242G, A242G-S258P, A68S-A242G, A68S-G165E, G127S-A242G, G165E-N184Q, G165E-N217S, I35A-V87S, L10M-M123I, M123I-G127T, N184Q-A242G, N217S-A242G, A128R-G165Q, A128R-S258D, A68S-A128R, A68S-G99S, G127S-S258D, G165E-A202V, G99S-A128P, G99S-G127S, G99S-G165Q, I35A-G165Q, I35A-T77N, L10M-A128P, L10M-A68S, L10M-N184Q, L10M-N217S, L10M-S258P, L10M-T77N, M123I-A242G, P9E-A128P, P9E-A202V, P9E-A68S, P9E-G165Q, P9E-N184Q, P9E-N217S, P9E-S258P, S86H-T115F, T210P-A242G, T3V-G165E, T3V-G99S, T3V-L10M, T3V-P9E, T77N-A128R, T77N-G165E, T77N-G99S, T77N-V87S, T78I-A242G, V87S-A128P, V87S-G165Q, V87S-M123I, V87S-N217S, A68S-G101A, G101A-A128P, G101A-G165Q, G101A-S258P, G165E-S258P, G99S-S258P, I35A-M123I, I35A-N217S, L10M-G127S, L10M-G165Q, M123I-G165E, P9E-T210P, S187E-A202V, S86H-N217S, T115F-A202V, T115F-G165Q, T115F-S258P, T3V-G101A, T3V-V87S, T77N-G101A, T77N-G203E, T78I-S258D, A128P-G165E, A128R-A202V, A128R-N184Q, A128R-N217S, A68S-G99Q, G101A-A202V, G101A-G127S, G101A-N184Q, G101A-N217S, G165Q-

G203E, G99Q-A128P, G99Q-G165Q, G99Q-N184Q, G99Q-N217S, G99S-A242G, G99S-T210P, I35A-A128P, I35A-T78I, L10M-T210P, L10M-T78I, P9E-G127S, P9E-M123I, T3V-I35A, T77N-G99Q, T78I-G99S, T78I-V87S, V87S-N184Q, A143V-A202V, A143V-G165Q, A143V-S258P, A1Q-N184Q, A1Q-S86H, A68S-G203E, A68S-S187E, A68S-V87S, G101A-M123I, G101A-T210P, G99Q-A202V, G99S-A202V, G99S-M123I, G99S-N184Q, G99S-N217S, I35A-N184Q, L10M-I35A, L10M-V87S, M123I-A128R, N184Q-G203E, N184Q-S187E, S187E-N217S, S250G-S258P, S86H-A143V, T115F-A143V, T115F-N184Q, T115F-N217S, T3V-A128R, T3V-G99Q, T3V-S86H, T77N-S187E, T77N-N184Q-N217S, A128P-N184Q-N217S, T77N-A128P-N217S, A68S-N184Q-N217S, A68S-T77N-N217S, G165Q-N184Q-N217S, T77N-A128P-N184Q, T77N-G165Q-N217S, T77N-G165Q-N184Q, A68S-T77N-G165Q, A68S-T77N-N184Q, T77N-A128P-G165Q, A128P-G165Q-N217S, A68S-A128P-N217S, A68S-T77N-A128P, A68S-G165Q-N217S, A128P-G165Q-N184Q, A68S-A128P-N184Q, A68S-A128P-G165Q, A68S-G165Q-N184Q, N184Q-A202V-N217S, G165Q-A202V-N217S, T77N-A202V-N217S, T3V-N184Q-N217S, A68S-A202V-N217S, T3V-T77N-N217S, A128P-A202V-N217S, G165Q-N184Q-A202V, T77N-N184Q-A202V, T3V-A128P-N217S, T3V-T77N-A128P, T3V-T77N-N184Q, T77N-G165Q-A202V, A128P-N184Q-A202V, A68S-N184Q-A202V, T3V-A128P-N184Q, A68S-T77N-A202V, T77N-A128P-A202V, T3V-G165Q-N217S, T3V-T77N-G165Q, T3V-A68S-N217S, T3V-A68S-T77N, A128P-G165Q-A202V, A68S-A128P-A202V, T3V-G165Q-N184Q, A68S-G165Q-A202V, T3V-A128P-G165Q, T3V-A68S-A128P, T3V-A68S-N184Q, T3V-A68S-G165Q, T3V-A202V-N217S, T3V-N184Q-A202V, T3V-T77N-A202V, T3V-A128P-A202V, T3V-G165Q-A202V, T3V-A68S-A202V, N184Q-N217S-S258P, T3V-M123I-N217S, M123I-N184Q-N217S, T77N-M123I-N217S, A128P-N184Q-S258P, A128P-N217S-S258P, A68S-M123I-N217S, M123I-G165Q-N217S, T77N-A128P-S258P, T77N-M123I-N184Q, T77N-N184Q-S258P, M123I-A128P-N217S, T3V-M123I-N184Q, T77N-N217S-S258P, A202V-N217S-S258P, A68S-M123I-N184Q, M123I-A128P-N184Q, N184Q-A202V-S258P, T3V-M123I-G165Q, T3V-T77N-M123I, T77N-M123I-G165Q, A68S-A128P-S258P, A68S-T77N-M123I, M123I-G165Q-N184Q, T3V-A68S-M123I, A68S-N184Q-S258P, G165Q-N184Q-S258P, G165Q-N217S-S258P, M123I-A202V-N217S, T3V-M123I-A128P, T77N-M123I-A128P, A128P-G165Q-S258P, A68S-N217S-S258P, M123I-A128P-G165Q, T3V-N217S-S258P, T77N-G165Q-S258P, A68S-M123I-A128P, A68S-M123I-G165Q, A68S-T77N-S258P, T3V-A128P-S258P, T3V-N184Q-S258P, G165Q-A202V-S258P, T3V-M123I-A202V, A128P-A202V-S258P, M123I-N184Q-A202V, T3V-T77N-S258P, A68S-A202V-S258P, A68S-G165Q-S258P, M123I-G165Q-A202V, T77N-A202V-S258P, T77N-M123I-A202V, A68S-M123I-A202V, M123I-A128P-A202V, T3V-A202V-S258P, T3V-A68S-S258P, T3V-G165Q-S258P, A68S-G127S-A128P, A68S-T77N-G127S, T77N-G127S-A128P, A68S-G127S-A202V, A68S-G127S-N184Q, A68S-G127S-N217S, G127S-A128P-A202V, G127S-A128P-N184Q, G127S-A128P-N217S, A68S-G127S-G165Q, G127S-A128P-G165Q, G127S-A202V-

N217S, G127S-N184Q-A202V, G127S-N184Q-N217S, T77N-G127S-A202V, T77N-G127S-N184Q, T77N-G127S-N217S, T77N-G127S-G165Q, T3V-A68S-G127S, T3V-G127S-A128P, G127S-G165Q-A202V, G127S-G165Q-N184Q, G127S-G165Q-N217S, M123I-A202V-S258P, M123I-N217S-S258P, M123I-N184Q-S258P, T3V-M123I-S258P, T3V-T77N-G127S, A68S-G127S-S258P, G127S-A128P-S258P, M123I-A128P-S258P, T3V-G127S-A202V, T3V-G127S-N184Q, T3V-G127S-N217S, A68S-M123I-S258P, T3V-G127S-G165Q, T77N-G127S-S258P, M123I-G165Q-S258P, T77N-M123I-S258P, G127S-A202V-S258P, G127S-N184Q-S258P, G127S-N217S-S258P, T3V-G127S-S258P, G127S-G165Q-S258P, A68S-M123I-G127S, M123I-G127S-A128P, M123I-G127S-A202V, M123I-G127S-N184Q, M123I-G127S-N217S, T77N-M123I-G127S, T3V-M123I-G127S, M123I-G127S-G165Q, M123I-G127S-S258P, A68S-A128P-T210P, A68S-T77N-T210P, T77N-A128P-T210P, N184Q-A202V-T210P, T3V-A128P-T210P, T3V-A68S-T210P, T3V-T77N-T210P, T77N-G165Q-T210P, T77N-T210P-S258P, A128P-G165Q-T210P, A128P-T210P-S258P, A202V-T210P-N217S, A68S-G165Q-T210P, A68S-T210P-S258P, N184Q-T210P-N217S, T3V-G165Q-T210P, T77N-A202V-T210P, A128P-A202V-T210P, A128P-N184Q-T210P, A128P-T210P-N217S, A202V-T210P-S258P, A68S-A202V-T210P, A68S-N184Q-T210P, A68S-T210P-N217S, G165Q-A202V-T210P, G165Q-T210P-S258P, T3V-A202V-T210P, T3V-N184Q-T210P, T3V-T210P-S258P, T77N-N184Q-T210P, T77N-T210P-N217S, G165Q-N184Q-T210P, N184Q-T210P-S258P, T210P-N217S-S258P, T3V-T210P-N217S, G165Q-T210P-N217S, A68S-G127S-T210P, G127S-A128P-T210P, T77N-G127S-T210P, G127S-T210P-S258P, T3V-G127S-T210P, G127S-G165Q-T210P, A68S-M123I-T210P, G127S-A202V-T210P, G127S-N184Q-T210P, G127S-T210P-N217S, M123I-A128P-T210P, M123I-A202V-T210P, M123I-N184Q-T210P, M123I-T210P-N217S, T3V-M123I-T210P, T77N-M123I-T210P, M123I-G165Q-T210P, M123I-T210P-S258P, A68S-G165Q-S258D, A68S-T77N-S258D, T77N-G165Q-S258D, T77N-T78I-G165Q, A68S-N184Q-S258D, A68S-N217S-S258D, G127T-N184Q-N217S, G165Q-N184Q-S258D, G165Q-N217S-S258D, M123I-G127S-T210P, T77N-N184Q-S258D, T77N-N217S-S258D, A68S-T77N-T78I, A68S-T78I-A128P, A68S-T78I-G127S, A68S-T78I-G165Q, T77N-T78I-A128P, T77N-T78I-G127S, T78I-A128P-G165Q, T78I-G127S-A128P, T78I-G127S-G165Q, A68S-G127T-N184Q, A68S-G127T-N217S, N184Q-N217S-S258D, T77N-T78I-N184Q, T78I-G165Q-N184Q, A202V-N217S-S258D, A68S-A202V-S258D, G127T-A128P-N184Q, G127T-A128P-N217S, G127T-A202V-N217S, G127T-N184Q-A202V, G165Q-A202V-S258D, T3V-T77N-T78I, T3V-T78I-G165Q, T77N-A202V-S258D, T77N-T78I-A202V, T77N-T78I-N217S, T78I-G165Q-A202V, T78I-G165Q-N217S, T78I-N184Q-A202V, T78I-N184Q-N217S, A68S-G127T-A128P, A68S-T78I-A202V, A68S-T78I-N184Q, A68S-T78I-N217S, G127T-G165Q-N184Q, G127T-G165Q-N217S, T3V-A68S-T78I, T3V-T78I-A128P, T3V-T78I-G127S, T77N-G127T-N184Q, T77N-G127T-N217S, T77N-T78I-S258P, T78I-A128P-A202V, T78I-A128P-N184Q, T78I-A128P-N217S, T78I-A202V-N217S, T78I-G127S-A202V, T78I-G127S-N184Q, T78I-

G127S-N217S, T78I-G165Q-S258P, A68S-G127T-A202V, A68S-T77N-G127T, G127T-A128P-G165Q, N184Q-A202V-S258D, T3V-G127T-N184Q, T3V-G127T-N217S, T77N-G127T-A128P, T78I-M123I-A202V, A128P-G165Q-A242G, A128P-G165Q-S258D, A128P-N184Q-S258D, A128P-N217S-S258D, A68S-A128P-S258D, A68S-G127T-G165Q, A68S-T78I-S258P, G127T-A128P-A202V, T3V-A128P-A242G, T3V-A68S-G127T, T3V-A68S-S258D, T3V-G127T-A202V, T3V-G165Q-A242G, T3V-G165Q-S258D, T3V-T77N-A242G, T3V-T77N-S258D, T3V-T78I-N184Q, T3V-T78I-N217S, T3V-T78I-S258P, T77N-A128P-A242G, T77N-A128P-S258D, T77N-G127T-G165Q, T77N-G165Q-A242G, T78I-A128P-S258P, T78I-G127S-S258P, A68S-M123I-S258D, G127T-G165Q-A202V, M123I-G165Q-S258D, T3V-G127T-A128P, T3V-N217S-S258D, T3V-T78I-A202V, T77N-G127T-A202V, T77N-M123I-S258D, T77N-T78I-M123I, T78I-M123I-G165Q, T78I-M123I-N184Q, T78I-N184Q-S258P, A68S-T78I-M123I, A68S-T78I-T210P, G127T-N184Q-S258P, G127T-N217S-S258P, M123I-A202V-S258D, M123I-N217S-S258D, T3V-A202V-S258D, T3V-M123I-S258D, T3V-N184Q-S258D, T3V-T77N-G127T, T3V-T78I-T210P, T77N-T78I-T210P, T78I-A128P-T210P, T78I-A202V-S258P, T78I-G127S-T210P, T78I-G165Q-T210P, T78I-M123I-A128P, T78I-M123I-G127S, T78I-M123I-N217S, and T78I-N217S-S258P.

7. The subtilisin variant of any preceding claim, wherein said variant comprises an amino acid sequence comprising one or more substitution sets selected from: M123I-Q136R; P9E-M123I; V45R-M123I; G127T-G165S-T210E; M123I-G165S-N239S; M123I-Q136R-G165Q; M123I-Q136R-S258E; M123I-V146S-T210E; P9E-V45R-M123I; Q136R-G165Q-S258E; V45R-M123I-Q136R; G127T-V146S-G165S-N239S; L10M-K27R-S98H-M123I; M123I-Q136R-G165Q-S258E; M123I-V146S-G165S-T210P; M123I-V146S-T210P-S250G; P9E-M123I-Q136R-G165Q; P9E-Q136R-G165Q-S258E; P9E-V45R-M123I-Q136R; V45R-M123I-Q136R-G165Q; V45R-M123I-Q136R-S258E; V45R-Q136R-G165Q-S258E; K27R-N43A-S98H-M123I-T210H; P9E-M123I-Q136R-G165Q-S258E; P9E-V45R-M123I-Q136R-G165Q; P9E-V45R-M123I-Q136R-S258E; V45R-M123I-Q136R-G165Q-S258E; Y103I-G127T-V146S-G165S-N239S; T77H-V87T-M123I-Q136R-N184Q-N239S; V26R-V45R-S98R-E111Q-M123I-Q136R; T77H-V87T-M123I-Q136R-G165Q-N184Q-N239S; T77S-V87S-M123I-Q136R-G165Q-S187P-S258P; T77S-V87S-M123I-Q136R-N160C-S187P-S258P; T77H-V87T-M123I-Q136R-N184Q-S187P-N239S-S258P; T3Q-K22Y-A24N-V45I-A48I-T77N-N217S-L234I-Y255R; P9E-T77Q-V87T-M123I-Q136R-G165Q-N184Q-N217S-N239S-S258E; N43A-V45R-T77H-V87T-M123I-Q136R-G165A-N184Q-S187P-N239S-S258P; P9E-N43A-V45R-T77S-V87S-M123I-Q136R-G165Q-S187P-N217S-S258P; P9E-N43A-V45R-T77H-V87T-M123I-Q136R-G165Q-N184Q-S187P-N239S-S258P; P9E-N43A-V45R-T77Q-V87T-M123I-Q136R-G165Q-N184Q-N217S-N239S-S258E; P9E-N43A-V45R-T77H-V87T-M123I-Q136R-G165A-N184Q-S187P-N217S-N239S-S258E; T77N-N217S-L234W;

T77N-N217S, T77N-G126Q-G165Q-T210P-N217S-L234W-S258P, T77N-G127F-G165Q-A202V-
 T210P-N217S-L234W-S258P; A1Q-T3V-V28A-I35A-A68S-T77N-S86H-T115F-N217S-N239S-S250G-
 S258P-Q274A; T77N-M123I-T210P-N239S-S250G-S258P; V45I-A68S-T77N-M123I-L125N-G165Q-
 N217S-S258P; A68S-V71A-T77N-L95A-G165Q-N184Q-T210P-S258P; M123I-T210P-N217S-S258P;
 G127F-G165Q-N217S-S258P; T77N-G165Q-N217S-S258P; T77S-G165A-N184Q-T210P-N217S;
 A68S-T77N-A128P-G165Q-N184Q-N217S; V71A-T77S-N184Q-S258P; T77N-M123I-G165Q-N184Q-
 N217S; V71A-L95A-A128S-T210P; V71A-M123I-A128S-T210P; V71A-M123I-A128S-G165E; T77S-
 A128P-N184Q-N217S-F260W; T77H-A128S-G165Q-T210P; T77D-S86R-S155E-G165A-N184Q;
 A68S-V71A-T77N-A128S-G165Q-T210P-N217S; V28A-I35A-A68S-T77N-A128S-G165Q-N184Q-
 G203N-T210P; A128P-T210P; T210P-S258P; A128P-T210P-S259P; A128P-S187P-T210P; A128P-
 S187P-T210P-S258P; P9E-T77S-V87T-T210P-N239K; K15I-L95Q-M123I-G165S-T210P-N217S;
 K15I-M123I-V146S-G165S-T210P-N239S; K15I-N76K-M123I-V146S-G165S-T210P-N239S; K15I-
 V146S-G165S-T210P-N239S; P009E-V146S-N184Q-G203E-T210P-P238R-N239S; M123I-G165S-
 N217S; K15I-L95Q-S108K-G165S-G203E-T210P-N217S; K15I-L95Q-S108K-G165S-G203E-T210P;
 P9E-K15I-G165S-A193D-G203E-T210P; T77N-M123I-G165Q-N217S; T77H-G165Q; T77N-G165Q-
 N217S; A68S-V71A-T77N-G165Q-N217S; T77N-V87S-M123I-S155G-G165Q-N217S; T77H-V87S-
 S155G-G165Q; T77N-V87S-S155G-G165Q-N217S; T77N-M123I-G165Q; T77N-G165Q; A68S-V71A-
 T77N-G165Q; T77N-V87S-M123I-S155G-G165Q; I35A-A68S-T77N-V87S-V147L-G165Q; A68S-
 T77N-V87S-V147L-S155E-G165Q; T77N-V87S-M123I-V147L-S155G-G165Q; A68S-T77N-M123I-
 G165Q-N184Q-N217S; T77N-M123I-A128P-G165Q-N184Q-N217S; A68S-V71A-T77N-S100N-
 A128P-G165Q-N184Q-T210P-N217S-N239T; T77N-A128P-G165Q-N184Q-N217S; A68S-A128P-
 G165Q-N184Q-N217S; A68S-T77N-G165Q-N184Q-N217S; A68S-T77N-A128P-N184Q-N217S;
 A68S-T77N-A128P-G165Q-N217S; A68S-T77N-A128P-G165Q-N184Q; M123I-G165Q-N184Q-
 N217S; T77N-G165Q-N184Q-N217S; T77N-M123I-N184Q-N217S; T77N-M123I-G165Q-N184Q;
 T3V-A68S-T77N-A128P-G165Q-N184Q-N217S; T3V-A68S-T77N-M123I-A128P-G165Q-N184Q-
 N217S; T3V-T77N-M123I-G165Q-N184Q-N217S; A68S-T77N-S108H-A128P-G165Q-N184Q-N217S;
 A68S-T77N-G99S-A128P-G165Q-N184Q-N217S; A68S-T77N-T114N-A128P-G165Q-N184Q-N217S;
 A68S-T77N-M123I-A128P-G165Q-N184Q-N217S; A68S-T77N-G127T-A128P-G165Q-N184Q-N217S;
 A68S-T77N-A128P-G165Q-N184Q-T210P-N217S; A68S-T77N-G99S-M123I-A128P-G165Q-N184Q-
 N217S; A68S-T77N-G99S-G127T-A128P-G165Q-N184Q-N217S; A68S-T77N-G99S-A128P-G165Q-
 N184Q-T210P-N217S; A68S-T77N-T114N-M123I-A128P-G165Q-N184Q-N217S; A68S-T77N-T114N-
 G127T-A128P-G165Q-N184Q-N217S; A68S-T77N-T114N-A128P-G165Q-N184Q-T210P-N217S;
 A68S-T77N-M123I-G127T-A128P-G165Q-N184Q-N217S; A68S-T77N-M123I-A128P-G165Q-N184Q-
 T210P-N217S; A68S-T77N-G127T-A128P-G165Q-N184Q-T210P-N217S; A68S-T77N-G99S-M123I-

A128P-G165Q-N184Q-T210P-N217S; A68S-T77N-M123I-G127T-A128P-G165Q-N184Q-T210P-N217S; T3V-A68S-T77N-G99S-M123I-A128P-G165Q-N184Q-N217S; T3V-A68S-T77N-T114N-M123I-A128P-G165Q-N184Q-N217S; T3V-A68S-T77N-M123I-G127T-A128P-G165Q-N184Q-N217S; T3V-A68S-T77N-M123I-A128P-G165Q-N184Q-T210P-N217S; S86H-T115F-A143V-G165Q-A202V-S258P; S86H-L95A-N184Q-A202V-T210P-N217S-L234W-S250G-S258P; V30T-I35A-T77N-G165Q-A202V-T210P-S250G-S258P; A1Q-T3V-I35A-T77N-T78I-S86H-T115F-G165Q-N184Q-N217S-L234W-S258P; A1Q-A37T-T77N-T78I-S86H-M123I-G165Q-N184Q-A202V-N239S-S250G-S258P-Q274A; P9E-T77N-N96S-G165Q-G203E-S250G; P9E-T77N-N96S-G165E-G203E-T210P-S250G; A1Q-T3V-S86H-G130S-S155G-G165Q-N184Q-A202V-G203N-T210P-Q274A; A1Q-T3V-S86H-G165Q-N184Q-N239S-S250G-S258P; A1Q-T77N-S86H-V147L-A151G-N184Q-N217S-S258P; S86H-T115F-S155G-G165Q-N184Q-A202V-Q274A; A68S-S86H-T115F-N184Q-A202V-G203N-N239S-S250G-S258P-Q274A; A68S-T77N-V87S-A128P-S155G-G165Q-N217S; A68S-T77N-A128P-G165Q; A68S-T77N-V87S-A128P-S155G-G165Q; I35A-A68S-T77N-V87S-A128P-V147L-G165Q-N217S; A68S-T77N-V87S-A128P-S155G-G165Q-N217S-A242G; A68S-T77N-V87S-A128P-V147L-S155E-G165Q; A68S-T77N-V87S-A128P-S155E-G165Q-N217S; A68S-T77N-S86N-V87S-A128P-S155G-G165Q-N217S-A242G; I35A-A68S-T77N-A128P-V147L-G165Q-N217S; I35A-A68S-T77N-A128P-S155G-G165Q-A242G; I35A-A68S-T77N-A128P-V147L-G165Q-A242G; A68S-T77N-V87S-A128P-S155G-G165Q-A242G; A68S-T77N-N86S-G165Q-K194E-S211N-N217S; A68S-T77N-N86S-V87S-S155G-G165Q-K194E-S211N-N217S; A68S-T77N-N86S-G165Q-K194E-S211N; A68S-T77N-N86S-V87S-S155G-G165Q-K194E-S211N; I35A-A68S-T77N-N86S-V87S-V147L-G165Q-K194E-S211N-N217S; A68S-T77N-N86S-V87S-S155G-G165Q-K194E-S211N-N217S-A242G; A68S-T77N-N86S-V87S-V147L-S155E-G165Q-K194E-S211N; A68S-T77N-N86S-V087S-S155E-G165Q-K194E-S211N-N217S; A68S-T77N-V87S-S155G-G165Q-K194E-S211N-N217S-A242G; I35A-A68S-T77N-N86S-V147L-G165Q-K194E-S211N-A242G; A68S-T77N-N86S-V87S-S155G-G165Q-K194E-S211N-A242G; A68S-T77N-G127T-A128P-G165Q-N184Q-A202V-N217S; T3V-A68S-T77N-M123I-A128P-G165Q-N184Q-A202V-N217S; A68S-T77N-G127T-A128P-G165Q-N184Q-N217S-S258P; T3V-A68S-T77N-M123I-A128P-G165Q-N184Q-N217S-S258P; A68S-T77N-G127T-A128P-G165Q-N184Q-G203Q-N217S; T3V-A68S-T77N-M123I-A128P-G165Q-N184Q-G203Q-N217S; A68S-T77N-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-M123I-A128P-G165Q-N184Q-A202V-N217S-S258P; A68S-T77N-A128P-G165Q-N184Q-G203Q-N217S-S258P; T3V-A68S-T77N-M123I-A128P-G165Q-N184Q-G203Q-N217S-S258P; S86H-T115F-G127T-A143V-G165Q-N184Q-A202V-N217S-S258P; T3V-S86H-T115F-M123I-A143V-G165Q-A202V-N217S-S258P; A68S-T77N-A128P-S129H-G165Q-N184Q-A202V-N217S; T3V-A68S-T77N-M123I-

A128P-S129H-G165Q-N184Q-A202V-N217S; S86H-T115F-S129H-A143V-G165Q-N184Q-A202V-N217S-S258P; T3V-S86H-T115F-M123I-S129H-A143V-G165Q-A202V-N217S-S258P; A68S-T77N-A128P-S155N-G165Q-N184Q-A202V-N217S; T3V-A68S-T77N-M123I-A128P-S155N-G165Q-N184Q-A202V-N217S; I35A-A68S-T77N-V87S-A128P-V147L-G165Q-A202V-N217S; T3V-I35A-A68S-T77N-V87S-M123I-A128P-V147L-G165Q-A202V-N217S; A68S-T77N-A128P-Q136N-S155N-G165Q-N184Q-N217S-A242N; T3V-A68S-T77N-M123I-A128P-Q136N-S155N-G165Q-N184Q-N217S-A242N; A68S-V71A-T77N-A128P-G165Q-N184Q-A202V-N217S; T3V-A68S-V71A-T77N-M123I-A128P-G165Q-N184Q-A202V-N217S; A68S-T77N-A128P-G165Q-N184Q-S187P-A202V-N217S; T3V-A68S-T77N-M123I-A128P-G165Q-N184Q-S187P-A202V-N217S; A68S-T77N-A128R-G165Q-N184Q-A202V-N217S-S258D; T3V-A68S-T77N-M123I-A128R-G165Q-N184Q-A202V-N217S-S258D; A68S-T77N-A128P-S129R-G165Q-N184Q-N217S-S258D; T3V-A68S-T77N-M123I-A128P-S129R-G165Q-N184Q-N217S-S258D; A68S-T77N-A128R-G165Q-N184Q-A202V-N217S-S258D-K264H; T3V-A68S-T77N-M123I-A128R-G165Q-N184Q-A202V-N217S-S258D-K264H; A68S-T77N-A128R-G165Q-N184Q-G203E-S258D; T3V-A68S-T77N-M123I-A128R-G165Q-N184Q-G203E-S258D; T3V-A68S-T77N-M123I-A128R-G165Q-N184E-A202V-N217S-S258D; T3V-A52R-A68S-T77N-M123I-A128R-G165Q-N184E-A202V-N217S-S258D; T3V-T77N-S100N-G105N-D119Q-A128P-S158N-N160D-T161Q-G165A-N184Q-N217S-S258P-K264S; T3V-T77N-S100N-G105N-A128P-S158D-T161Q-G165A-N184Q-N217S-S258P; T3V-T77N-S100N-G105N-A128P-S158N-N160D-T161Q-G165A-N184Q-N217S-S258P; A24Q-V45I-A68S-T77N-S98R-E111Q-M123I-S158Q-G165E-N184Q-S187E-V204I-T210I-N217S-S259E; P9E-A52R-S86R-E111Q-S129R-S158Q-G165E-V204I-T210I-S258D-S259E-K264Q; T3V-A68S-T77N-G127T-A128P-G165Q-N184Q-A202V-N217S; T3V-I35A-T77N-V87S-M123I-A128P-G165Q-N184Q-A202V-N217S-A242G; T3V-A68S-T77N-G127E-A128P-G165Q-N184Q-A202V-N217S; T3V-A68S-T77N-M123I-G127V-A128P-G165Q-N184Q-N217S; T3V-A68S-T77N-M123I-A128P-A151G-G165Q-N184Q-N217S; T3V-A68S-T77N-G127T-A128P-A151G-G165Q-N184Q-N217S; T3V-T77N-M123I-A128P-A151G-G165Q-N184Q-A202V-N217S; T3V-A68S-T77N-L95F-A128P-G165Q-N184Q-A202V-N217S; T3V-A68S-T77N-L95A-M123I-A128P-G165Q-N184Q-N217S; T3V-A68S-T77N-L95F-M123I-A128P-G165Q-N184Q-N217S; T3V-A68S-T77N-G99A-A128P-G165Q-N184Q-A202V-N217S; T3V-A68S-T77N-G99E-A128P-G165Q-N184Q-A202V-N217S; T3V-A68S-T77N-G99Q-A128P-G165Q-N184Q-A202V-N217S; T3V-A68S-T77N-G99A-M123I-A128P-G165Q-N184Q-N217S; T3V-A68S-T77N-S108H-A128P-G165Q-N184Q-N217S; T3V-I35A-T77N-V87S-M123I-A128P-V146A-G165Q-N184Q-A202V-N217S-A242G; T3V-A68S-T77N-G127T-A128P-G165E-N184Q-A202V-N217S; T3V-A68S-T77N-M123I-A128P-V146A-G165Q-N184Q-N217S; T3V-A68S-T77N-M123I-A128P-T161D-G165Q-N184Q-A202V-N217S; T3V-V45R-A68S-T77N-S102D-A128P-T161D-G165Q-N184Q-A202V-N217S; T3V-

A68S-T77N-G127T-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-M123I-G127T-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-M123I-G127V-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-I35A-T77N-V87S-G127T-A128P-G165Q-N184Q-A202V-N217S-A242G-S258P; T3V-T77N-M123I-G127Q-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-T77N-M123I-G127V-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-T77N-M123I-G127E-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-M123I-G127T-A128P-G165Q-N184Q-A202V-N217S; T3V-A68S-T77N-M123I-G127T-A128P-G165Q-N184Q-A202V-T210P-N217S; T3V-A68S-T77N-M123I-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P; T3V-A68S-T77N-M123I-G127T-A128P-G165Q-N184Q-A202V-N217S-S258D; L10M-I35A-T78I-V87S-M123I-A202V-S258D-F260W; L10M-I35A-T78I-V87S-M123I-A202V-F260W-K264H; L10M-I35A-T78I-V87S-A202V-S258D; L10M-I35A-T78I-V87S-S187E-A202V; L10M-T78I-V87S-S187E-A202V; L10M-I35A-T78I-S187E-A202V; T3V-L10M-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-I35A-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-T78I-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-V87S-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-M123I-G127S-A128P-G165E-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P; T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-L216Q-N217S-S258P; T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-A242G-S258P; T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258D; T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P-F260W; A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-L31I-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; A68S-T77N-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; A68S-T77N-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P; A68S-T77N-G127T-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-T78I-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258D; A68S-T77N-T78I-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258D; T3V-A68S-T77N-T78I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; A68S-T77N-T78I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-G127S-A128P-G165Q-N184Q-A202V-N217S-S258D; A68S-T77N-G127S-A128P-G165Q-N184Q-A202V-N217S-S258D; T3V-A68S-T77N-T78I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258D; A68S-T77N-T78I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258D; T3V-A68S-T77N-S108H-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-G99Q-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-G99Q-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P; T3V-A68S-T77N-G127S-A128P-G165E-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-

G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P; T3V-P9E-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P; T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-S187E-A202V-T210P-N217S-S258P; T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-G203E-T210P-N217S-S258P; T3V-L10M-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P; T3V-P9E-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-S187E-A202V-N217S-S258P; T3V-P9E-A68S-T77N-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P; T3V-A68S-T77N-G127S-A128P-G165Q-N184Q-S187E-A202V-T210P-N217S-S258P; T3V-A68S-T77N-G127S-A128P-G165Q-N184Q-A202V-G203E-T210P-N217S-S258P; T3V-A68S-T77N-G101A-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P; T3V-L10M-A68S-T77N-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P; T3V-P9E-A68S-T77N-M123I-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P; T3V-A68S-T77N-M123I-A128P-G165Q-N184Q-S187E-A202V-T210P-N217S-S258P; T3V-A68S-T77N-M123I-A128P-G165Q-N184Q-A202V-G203E-T210P-N217S-S258P; T3V-A68S-T77N-G101A-M123I-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P; T3V-L10M-A68S-T77N-M123I-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P; T3V-L10M-A68S-T77N-G101A-M123I-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P; T3V-P9E-L10M-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P; T3V-P9E-A68S-T77N-G99Q-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P; T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-S243I-S258P; T3V-A68S-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-M123I-G127S-A128P-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-M123I-G127S-A128P-G165Q-A202V-N217S-S258P; T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-S258P; T3V-A68S-T77E-M123I-G127T-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77E-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P-K264H; T3V-A68S-T77N-S102T-M123I-G127S-A128P-G165Q-N184Q-A202V-N211S-N217S-S258P; T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N211S-N217S-S258P; T3V-A68S-T77N-S102T-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P-S259P; T3V-A68S-T77N-M123I-G127S-A128P-T161Q-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-M123I-G127S-A128P-S158Q-G165Q-N184Q-A202V-N217S-S258P; T3V-V45Q-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A193Q-A202V-N217S-S258P; T3V-A68S-T77N-M123I-G127S-A128P-A143Q-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-G117S-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A48V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A48T-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-

A29S-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-G127S-A128P-G165Q-N184Q-A202V-N217S; T3V-A68S-T77N-G127S-A128P-G165Q-N184Q-N217S-S258P; T3V-A68S-T77N-A128P-G165Q-N184Q-A202V-N217S-S258P; A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S; A68S-T77N-M123I-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-G99Q-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-T78I-V87S-M123I-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P; T3V-A68S-T77N-T78I-M123I-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-A242G-S258P; T3V-V26Q-A68S-T77N-M123I-G127S-A128P-G165E-N184Q-A202V-T210P-N217S-S258P; T3V-P9E-L10M-A68S-T77N-M123I-G127S-A128P-G165E-N184Q-A202V-T210P-N217S-S258P; T3V-P9E-L10M-A68S-T77N-G101A-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-T78I-M123I-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P; T3V-A68S-T77N-T78I-M123I-G127S-A128P-N160K-G165Q-N184Q-A202V-T210P-N217S-A242G-S258P; T3V-A68S-T77N-V87S-M123I-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-S258P; T3V-P40E-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77E-M123I-G127T-A128P-G165E-N184Q-A202V-N217S-P238Q-S258P; T3V-A68S-T77N-G99S-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77E-M123I-G127T-G165E-N184Q-A202V-N217S-S258P; A68S-T77N-A128P-G165Q-N184Q-N217S-S258P; T3V-A68S-T77N-G101A-S102T-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P; T3V-A68S-T77N-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S; T3V-A68S-T77N-T78I-G99S-G127S-A128P-G165Q-T210P-A242G-S258P-K264H; T3V-A68S-T77N-T78I-G99S-M123I-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-A242G-S258P; T3V-A68S-T77N-M123I-G127T-G165E-N184Q-A202V-N217S-P238Q-S258P; T3V-A68S-T77N-T78I-G99S-G127S-A128P-G165Q-T210P-A242G-S258P; T3V-A68S-T77N-T78I-G99S-M123I-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-A242G-S258D; I35A-M123I-G127T-T210P-S243I; T3V-A68S-T77N-G99S-M123I-G127S-A128P-G165Q-N184Q-A202V-N217S-S258P-F260W; T3V-A68S-T77N-T78I-G99S-G101A-G127S-A128P-G165Q-T210P-A242G-S258P; A68S-T77N-G99Q-A128P-G165Q-N184Q-N217S; A68S-T77N-A128P-G165Q-N184Q-A202V-N217S; A68S-T77N-A128P-G165Q-N184Q-N217S-S258D; T3V-A68S-T77N-T78I-G99S-G101A-M123I-G127S-A128P-G165Q-N184Q-A202V-T210P-N217S-A242G-S258P; and V45R-S98R.

8. The subtilisin variant of any preceding claim, wherein said variant further comprises:

- i. Motif A, Motif B, or a combination thereof;
- ii. Motif A and Motif B;
- iii. Motif 1 between Asp32 and His63;
- iv. Motif 6 or Motif 7 between Lys236 and Arg246; or

v. Motif 1 between Asp32 and His63 and Motif 6 or Motif 7 between Lys236 and Arg246;
wherein Motif A is selected from Motif 1, Motif 2, Motif 3, and Motif 4;
wherein Motif B is selected from Motif 5, Motif 6, Motif 7, Motif 8, Motif 9, Motif 10, Motif 11, Motif 12, Motif 13, and Motif 14; and wherein the amino acid positions are numbered by correspondence with the amino acid sequence of SEQ ID NO:2.

9. The subtilisin variant of any preceding claim, wherein said variant comprises an amino acid sequence having at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% but less than 100% amino acid sequence identity to the amino acid sequence of an AprL-Clade subtilisin enzyme identified in Figure 1.

10. The isolated subtilisin variant of any preceding claim, wherein said variant comprises an amino acid sequence having at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% but less than 100% amino acid sequence identity to the amino acid sequence of SEQ ID NO: 2 or 9.

11. The subtilisin variant of any preceding claim, wherein said variant is from a parent with 70%, 75%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% amino acid sequence identity to an AprL-Clade subtilisin enzyme comprising (i) a Motif A, Motif B, or a combination thereof; or (ii) Motif A and Motif B; wherein Motif A is selected from Motif 1, Motif 2, Motif 3, and Motif 4; and wherein Motif B is selected from Motif 5, Motif 6, Motif 7, Motif 8, Motif 9, Motif 10, Motif 11, Motif 12, Motif 13, and Motif 14.

12. The subtilisin variant of any preceding claim, wherein said variant is from a parent comprising the amino acid sequence of SEQ ID NO:2 or 9.

13. The subtilisin variant of any preceding claim, wherein said variant has protease activity and/or is isolated.

14. The subtilisin variant of any preceding claim, wherein said variant has one or more improved property when compared to a reference subtilisin; wherein the improved property is selected from improved protease activity, improved cleaning performance in detergent, improved thermostability in

detergent, and improved aged laundry cleaning in detergent; and wherein said detergent is optionally a boron-free composition.

15. The subtilisin variant of any preceding claim, wherein the reference subtilisin comprises an amino acid sequence of SEQ ID NO:2, 9, or 63.

16. The subtilisin variant of claim 13 or 14, wherein the improved property is

(i) improved protease activity, wherein said variant has a PI > 1 on N-suc-AAPF-pNA or dimethyl casein substrate;

(ii) improved cleaning performance in detergent, wherein said variant has a BMI, POM, egg, and/or egg yolk cleaning PI > 1;

(iii) improved thermostability in detergent, wherein said variant has a stability PI > 1 or % remaining activity $\geq$ 5%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 100% or more after 15-20 minutes at a stress temperature of 60°C, 61°C, 62°C, 63°C, 64°C, 65°C, 66°C, 67°C, 68°C, 69°C or more; and/or

(iv) improved aged laundry cleaning in detergent, wherein said variant has (a) an aged laundry cleaning PI > 1, (b) remaining cleaning activity > 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 100% or more after 24 hours at 55°C, and/or (c) remaining cleaning activity > 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 100% or more after 3 days or 3 weeks at 37°C.

17. The subtilisin variant of claim 16, wherein said

(i) protease activity is measured in accordance with the protease activity assay of Example 1;

(ii) cleaning performance in detergent is measured in accordance with the cleaning performance in laundry (HDL and HDD) and ADW detergents assay of Example 1;

(iii) thermostability in detergent is measured in accordance with the general sample set-up for stability assay of Example 1; and/or

(iv) aged laundry cleaning in detergent is measured in accordance with the aged laundry cleaning assay of Example 1.

18. A composition comprising one or more subtilisin variant of any preceding claim.

19. The composition of claim 18, wherein said composition is a detergent composition.

20. The composition of Claim 19, wherein the detergent composition is selected from a laundry detergent, a fabric softening detergent, a dishwashing detergent, and a hard-surface cleaning detergent.

21. The composition of any one of claims 18-20, wherein said composition further comprises one or more calcium ion and/or zinc ion; one or more enzyme stabilizer; from about 0.001% to about 1.0 weight % of said subtilisin variant; one or more bleaching agent; one or more adjunct material; and/or one or more additional enzymes or enzyme derivatives selected from the group consisting of acyl transferases, alpha-amylases, beta-amylases, alpha-galactosidases, arabinosidases, aryl esterases, beta-galactosidases, carrageenases, catalases, cellobiohydrolases, cellulases, chondroitinases, cutinases, DNase or nuclease, endo-beta-1, 4-glucanases, endo-beta-mannanases, esterases, exo-mannanases, galactanases, glucoamylases, hemicellulases, hyaluronidases, keratinases, laccases, lactases, ligninases, lipases, lipoxygenases, lysozymes, mannanases, oxidases, pectate lyases, pectin acetyl esterases, pectinases, pentosanases, peroxidases, phenoloxidases, phosphatases, phospholipases, phytases, polygalacturonases, proteases, pullulanases, reductases, rhamnogalacturonases, beta-glucanases, tannases, transglutaminases, xylan acetyl-esterases, xylanases, xyloglucanases, xylosidases, metalloproteases, additional serine proteases, and combinations thereof.
22. The composition of any one of Claims 18-21, wherein said composition contains phosphate or is phosphate-free and/or contains boron or is boron-free.
23. The composition of any one of Claims 18-22, wherein said composition is a granular, powder, solid, bar, liquid, tablet, gel, paste or unit dose composition.
24. A method of cleaning, comprising contacting a surface or an item in need of cleaning with the subtilisin variant of any one of claims 1-17 or the composition of any one of Claims 18-22; and optionally further comprising the step of rinsing said surface or item after contacting said surface or item with said variant or composition, wherein, optionally, said item is dishware or fabric.
25. A polynucleotide comprising a nucleic acid sequence that encodes the subtilisin variant of any one of claims 1-17, wherein said polynucleotide is optionally isolated.
26. An expression vector or cassette comprising the polynucleotide of claim 25.
27. The expression vector or cassette of claim 26, wherein the nucleic acid sequence is operably linked to a promoter.
28. A recombinant host cell comprising the vector or cassette of claim 26 or 27 or the polynucleotide of claim 25.

29. A composition comprising the subtilisin variant of any one of Claims 1-17, wherein said composition is a disinfectant, industrial or institutional cleaning, medical instrument cleaning, animal feed, contact lens cleaning, wound cleaning, or textile, leather or feather processing composition.

1	45	
B.lich_AprL_CAJ70731	(1)	AQTVPYGIPLIKADKVQAQGFKGANVKVAVLDTGTIQASHPDLNVV
B.licheniformis_BliD02339	(1)	AQTVPYGIPLIKADKVQAQGFKGANVKVAVLDTGTIQASHPDLNVV
B.licheniformis_AEQ38580	(1)	AQTVPYGIPLIKADKVQAQGFKGANVKVAVLDTGTIQASHPDLNVV
B.licheniformis_AKI30031	(1)	AQTVPYGIPLIKADKVQAQGFKGANVKVAVLDTGTIQASHPDLNVV
B.licheniformis_AID16241	(1)	AQTVPYGIPLIKADKVQAQGFKGANVKVAVLDTGTIQASHPDLNVV
B.licheniformis_AAG00493	(1)	AQTVPYGIPLIKADKVQAQGFKGANVKVAVLDTGTIQASHPDLNVV
B.licheniformis_AEU17777	(1)	AQTVPYGIPLIKADKVQAQGFKGANVKVAVLDTGTIQASHPDLNVV
B.licheniformis_EWH21773	(1)	AQTVPYGIPLIKADKVQAQGFKGANVKVAVLDTGTIQASHPDLNVV
B.licheniformis_AEU12640	(1)	AQTVPYGIPLIKADKLHAQGFKGANVGAVLATGPTSHPDLNVV
B.licheniformis_CAA62666	(1)	GQTVPYGIPLIKADKVQAQGFKGANVKVAVLDTGTIQASHPDLNVV
Bac.sp.NSP9.1_WP_026586291	(1)	AQTVPYGIPLIKADKVQAQGFKGANVKVGVIDTGTIQSSHSDLNVS
B.sonorensis_WP_077737098	(1)	AQTVPYGIPLIKADKVQAQGFKGANVKVGVIDTGTIQSSHSDLNVS
Bac.sp._WP_076761713	(1)	AQTVPYGISLIIKADKVQAQGFKGANVKVGVIDTGTIQSSHSDLNVV
B.glycini fermentans_WP_082634659	(1)	AQTVPYGIPQIKADKVQAQGFKGANVKVGVLDTGTIAASHSDLNVV
CN101215534-0002	(1)	AQTVPYGVPLIKADKVQAQGFKGANVKVAVLDTGTIQASHPDLNVV
WO9739130-0002	(1)	AQTVPYGIPLIKADKVQAQGFKGANVKVAVLDTGTIQASHPDLNVV
US8110391-0009	(1)	AQTVPYGIPLIKADKVQAQGFKGANVKVAVLDTGTIQASHPDLNVV
JP2013500714-0110	(1)	AQTVPYGIPLIKADKVQAQGFKGANVKVAVLDTGTIQASHPDLNVV
WO2006122655-0031	(1)	AQTVPYGIPLIKADKVQAQGFKGANVKVAVLDTGTIQASHPDLNVV
US7087415-0014	(1)	AQTVPYGIPLIKADKVQAQGFKGANVKVAVLDTGTIQASHPDLNVV
US20110171718-0120	(1)	GQTVPYGIPLIKADKVQAQGFKGANVKVAVLDTGTIQASHPDLNVV
US20030049619-0014	(1)	GQTVPYGIPLIKADKVQAQGFKGANVKVAVLDTGTIQASHPDLNVV
JP2013500714-0121	(1)	GQTVPYGIPLIKADKVQAQGFKGANVKVAVLDTGTIQASHPDLNVV
US7569226-0012	(1)	AQTVPYGIPLIKADKVQAQGFKGANVKVGII DTGTIAASHTDLKVV
US20050009167-0017	(1)	AQTVPYGIPLIKADKVQAQGFKGANVKVGII DTGTIAASHTDLKVV
US20110171718-0098	(1)	AQTVPYGIPLIKADKVQAQGFKGANVKVGII DTGTIASSHTDLKVV
WO2017006266-0002	(1)	AQTVPYGIPLIKADKVQAQGFKGANVKVAVLDTGTIQASHPDLNVV

FIG. 1A	FIG. 1AA
	FIG. 1AB

FIG. 1AA

FIG. 1A

	91	135
B.lich_AprL_CAJ70731	(90)	YAVKVLNSSGSGSYSGIVSGIEWATTNGMDVINMSLGGASGSTAM
B.licheniformis_BlId02339	(90)	YAVKVLNSSGSGSYSGIVSGIEWATTNGMDVINMSLGGPSGSTAM
B.licheniformis_AEQ38580	(90)	YAVKVLNSSGSGSYSGIVSGIEWATTNGMDVINMSLGGASGSTAM
B.licheniformis_AKI30031	(90)	YAVKVLNSSGSGSYSGIVSGIEWATTNGMDVINMSLGGASGSTAM
B.licheniformis_AID16241	(90)	YAVKVLNSSGSGSYSGIVSGIEWATTNGMDVINMSLGGASGSTAM
B.licheniformis_AAG00493	(90)	YAVKVLNSSGSGSYSGIVSGIEWATTNGMDVINMSLGGASGSTAM
B.licheniformis_AEU17777	(90)	YAVKVLNSSGSGSYSGIVSGIEWATTNGMDVINMSLGGASGSTAM
B.licheniformis_EWH21773	(90)	YAVKVLNSSGSGSYSGIVSGIEWATTNGMDVINMSLGGASGSTAM
B.licheniformis_AEU12640	(90)	YAVKVLNSSGSGSYSGIVSGIEWATTNGMDVINMSLGGASGSTAM
B.licheniformis_CAA62666	(90)	YAVKVLNSSGSGSYSGIVSGIEWATTNGMDVINMSLGGASGSTAM
Bac.sp.NSP9.1_WP_026586291	(91)	YAIKVLNSSGSGTYSAIVSGIEWATSNMGMDVINMSLGGSSGSTAL
B.sonorensis_WP_077737098	(91)	YAIKVLNSSGSGTYSAIVSGIEWATSNMGMDVINMSLGGSSGSTAL
Bac.sp._WP_076761713	(91)	YAIKVLNSSGSGTYSAIVSGIEWATANGMDVINMSLGGSSGSTAL
B.glycinifermentans_WP_082634659	(90)	YAIKVLNSSGSGTYSAIVSGIEWATANNLDVINMSLGGPSGSTAL
CN101215534-0002	(90)	YAVKVLNSSGSGSYSGIVSGIEWATTNGMDVINMSLGGASGSTAM
WO9739130-0002	(90)	YAVKVLNSSGSGSYSGIVSGIEWATTNGMDVINMSLGGASGSTAM
US8110391-0009	(90)	YAVKVLNSSGSGSYSGIVSGIEWATTNGMDVINMSLGGASGSTAM
JP2013500714-0110	(90)	YAVKVLNSSGSGTYSGIVSGIEWATTNGMDVINMSLGGPSGSTAM
WO2006122655-0031	(90)	YAVKVLNSSGSGSYSGIVSGIEWATTNGMDVINMSLGGPSGSTAM
US7087415-0014	(90)	YAVKVLNSSGSGSYSGIVSGIEWATTNGMDVINMSLGGPSGSTAM
US20110171718-0120	(90)	FAVKVLNSSGSGSYSGIVSGIEWATTNGMDVINMSLGGPSGSTAM
US20030049619-0014	(89)	FAVKVLNSSGSGSYSGIVSGIEWATTNGMDVINMSLGGPSGSTAM
JP2013500714-0121	(90)	YAVKVLNSSGSGSYSAIVSGIEWATTTGMDVINMSLGGASVSTAM
US7569226-0012	(90)	YAIKVLNSSGSGTYSAIVSGIEWATQNGLDVINMSLGGPSGSTAL
US20050009167-0017	(90)	YIAKVLNSSGSGTYSAIVSGIEWATQNGLDVINMSLGGPSGSTAL
US20110171718-0098	(90)	YAIKVLNSSGSGTYSAIVSGIEWATQNGLDVINMSLGGPSGSTAL
WO2017006266-0002	(90)	YAVKVLNSSGSGSYSGIVSGIEWATTNGMDVINMSLGGASGSTAM

FIG. 1B	FIG. 1BA
	FIG. 1BB

FIG. 1BA

136 180
B.lich_AprL_CAJ70731 (135) KQAVDNAYARGVVVAAAGNSGSSGNTNTTIGYPAKYDSVIAVGAV
B.licheniformis_BliD02339 (135) KQAVDNAYARGVVVAAAGNSGSSGNTNTTIGYPAKYDSVIAVGAV
B.licheniformis_AEQ38580 (135) KQAVDNAYARGVVVAAAGNSGSSGNTNTTIGYPAKYDSVIAVGAV
B.licheniformis_AKI30031 (135) KQAVDNAYARGVVVAAAGNSGSSGNTNTTIGYPAKYDSVIAVGAV
B.licheniformis_AID16241 (135) KQAVDNAYAKGVVVVAAAGNSGSSGNTNTTIGYPAKYDSVIAVGAV
B.licheniformis_AAG00493 (135) KQAVDNAYARGVVVVAAAGNSGSSGNTNTTIGYPAKYDSVIAVGAV
B.licheniformis_AEU17777 (135) KQAVDNAYARGVVVAAAGNSGSSGNTNTTIGYPAKYDSVIAVGAV
B.licheniformis_EWH21773 (135) KQAVDNAYAKGVVVVAAAGNSGSSGNANTTIGYPAKYDSVIAVGAV
B.licheniformis_AEU12640 (135) KQAVDNAYAKGVVVVAAAGNSGSSGNTNTTIGYPAKYDSVIAVGAV
B.licheniformis_CAA62666 (135) KQAVDNAYARGVVVAAAGNSGSSGNTNTTIGYPAKCDVIPVGGE
Bac.sp.NSP9.1_WP_026586291 (136) KQAVDNAYARGVVVAAAGNSGSSGSSNTTIGYPAKYDSVIAVGAV
B.sonorensis_WP_077737098 (136) KQAVDNAYARGVVVAAAGNSGSSGSSNTTIGYPAKYDSVIAVGAV
Bac.sp._WP_076761713 (136) KQAVDRAYS SGVVVAAAGNSGSSGSANTTIGYPAKYDSVIAVGAV
B.glycinifermentans_WP_082634659 (135) KQAVDKAYASGVVVVAAAGNSGTSGSSSTTIGYPAKYDSVIAVGAV
CN101215534-0002 (135) KQAVDNAYARGVVVAAAGNSGSSGNTNTTIGYPAKYDSVIAVGAV
WO9739130-0002 (135) KQAVDNAYARGVVVAAAGNSGSSGNTNTTIGYPAKYDSVIAVGAV
US8110391-0009 (135) KQAVDNAYARGVVVAAAGNSGNSGSTNTTIGYPAKYDSVIAVGAV
JP2013500714-0110 (135) KQAVDNAYARGVVVAAAGNSGSSGNTNTTIGYPAKYDSVIAVGAV
WO2006122655-0031 (135) KQAVDNAYARGVVVAAAGNSGSSGNTNTTIGYPAKYDSVIAVGAV
US7087415-0014 (135) KQAVDNAYARGVVVAAAGNSGSSGNTNTTIGYPAKYDSVIAVGAV
US20110171718-0120 (135) KQAVDNAYS KGVVPVAAAGNSGSSGYTNTTIGYPAKYDSVIAVGAV
US20030049619-0014 (134) KQAVDNAYS KGVVPVAAAGNSGSSGYTNTTIGYPAKYDSVIAVGAV
JP2013500714-0121 (135) KQAVDHAYARGAVVSSAGNSGSSGNTNTTIGYPAKYDSVIAVGAV
US7569226-0012 (135) KQAVDKAYASGI VVAAAGNSGSSGSQNTTIGYPAKYDSVIAVGAV
US20050009167-0017 (135) KQAVDKAYASGI VVAAAGNSGSSGSQNTTIGYPAKYDSVIAVGAV
US20110171718-0098 (135) KQAVDKAYASGI VVAAAGNSGSSGSQNTTIGYPAKYDSVIAVGAV
WO2017006266-0002 (135) KQAVDNAYAKGVVVVAAAGNSGSSGNTNTTIGYPAKYDSVIAVGAV

FIG. 1BB

FIG. 1CA	B.lich_AprL_CAJ70731	(180)	DSNSNRASFSSVGAELEVMAPGAG-VYSTYPTNTYATLNGTSMAS	181	225
	B.licheniformis_BliD02339	(180)	DSNSNRASFSSVGAKLEVMAPGAG-VYSTYPTSTYATLNGTSMAS		
FIG. 1CB	B.licheniformis_AEQ38580	(180)	DSNSNRASFSSVGAELEVMAPGAGVYSTYPTNTYATLNGTSMAS		
	B.licheniformis_AKI30031	(180)	DSNSNRASFSSVGAELEVMAPGAG-VYSTYPTNTYATLNGTSMAS		
	B.licheniformis_AID16241	(180)	DSNGNRASFSSVGAELEVMAPGAG-VYSTYPTNTYATLNGTSMAS		
	B.licheniformis_AAG00493	(180)	DSNSNRASFSSVGAELEVMAPGAG-VYSTYPTSTYATLNGTSMAS		
	B.licheniformis_AEU17777	(180)	DSNSNRASFSSVEAELEVMAPGAG-VYSTYPTNTYATLNGTSMAS		
	B.licheniformis_EWH21773	(180)	DSNSNRASFSSVGAELEVMAPGAS-VYSTYPTSTYATLNGTSMAS		
	B.licheniformis_AEU12640	(180)	DSNSNRASFSSVGAELEVMAPGAG-VYSTYPTNTYATLNGTSMAS		
	B.licheniformis_CAA62666	(180)	DSNSNRSSFSSVGAELEVMAPVSG-VYSTYPTNTYATLNGTSMAS		
	Bac.sp.NSP9.1_WP_026586291	(181)	DSNSNRASYSSVGSELEVMAPGSG-VYSTYPSNTYATLNGTSMAS		
	B.sonorensis_WP_077737098	(181)	DSNSNRASYSSVGSELEVMAPGSG-VYSTYPSNTYATLNGTSMAS		
	Bac.sp._WP_076761713	(181)	DSNSNRASFSSVGAELEVMAPGSA-VYSTYPANTYATLNGTSMAS		
	B.glycinifermentans_WP_082634659	(180)	NSSNQRASFSSVGPELDVVAPGVS-IYSTYPSNTYATLNGTSMAS		
	CN101215534-0002	(180)	DSNSNRASFSSVGAELEVMAPGAG-VYSTYPTNTYATLNGTSMAS		
	WO9739130-0002	(180)	DSNSNRASFSSVGAELEVMAPGAG-VYSTYPTNTYATLNGTSMVS		
	US8110391-0009	(180)	DSNSNRASFSSVGAELEVMAPGAG-VYSTYPTNTYATLNGTSMAS		
	JP2013500714-0110	(180)	DSNSNRASFSSVGAELEVMAPGAG-VYSTYPTSTYATLNGTSMAS		
	WO2006122655-0031	(180)	DSNSNRASFSSVGAELEVMAPGAG-VYSTYPTSTYATLNGTSMAS		
	US7087415-0014	(180)	DPNSNRASFSSVGAELEVMAPGAG-VYSTYPTSTYATLNGTSMAS		
	US20110171718-0120	(180)	DSNSNRASFSSVGAELEVMAPGAG-VYSTYPTNTYATLNGTSMAS		
	US20030049619-0014	(179)	DSNSNRASFSSVGAELEVMAPGAG-VYSTYPTNTYATLNGTSMAS		
	JP2013500714-0121	(180)	DSNSNRASFSSVGAELEVMAPGAG-VYSTYPTNTYATLNGTSMAS		
	US7569226-0012	(180)	DSNKNRASFSSVGAELEVMAPGVS-VYSTYPSNTYATLNGTSMAS		
	US20050009167-0017	(180)	DSNKNRASFSSVGAELEVMAPGVS-VYSTYPSNTYATLNGTSMAS		
	US20110171718-0098	(180)	DSNKNRASFSSVGSELEVMAPGVS-VYSTYPSNTYATLNGTSMAS		
	WO2017006266-0002	(180)	DSNSNRASFSSVGAELEVMAPGAG-VYSTYPTNTYATLNGTSMAS		

<i>FIG. 1C</i>	<i>FIG. 1CA</i>
	<i>FIG. 1CB</i>

FIG. 1CA

	226		270
B.lich_AprL_CAJ70731	(224)	PHVAGAAALILSKHPNLSASQVRNRLSSTATYLGSSFYYGKGLIN	
B.licheniformis_BliD02339	(224)	PHVAGAAALILSKHPNLSASQVRNRLSSTATYLGSSFYYGKGLIN	
B.licheniformis_AEQ38580	(225)	PHVAGAAALILSKHPNLSASQVRNRLSSTATYLGSSFYYGKGLIN	
B.licheniformis_AKI30031	(224)	PHVAGAAALILSKHPNLSASQVRNRLSSTATYLGSSFYYGKGLIN	
B.licheniformis_AID16241	(224)	PHVAGAAALILSKHPNLSASQVRNRLSSTATYLGSSFYYGKGLIN	
B.licheniformis_AAG00493	(224)	PHVAGAAALILSKHPNLSASQVRNRLSSTATYLGSSFYYGKGLIN	
B.licheniformis_AEU17777	(224)	PHVAGAAALILSKHPNLSASQVRNRLSSTATYLGSSFYYGKGLIN	
B.licheniformis_EWH21773	(224)	PHVAGAAALILSKHPNLSASQVRNRLSSTATYLGSSFYYGKGLIN	
B.licheniformis_AEU12640	(224)	PHVAGAAALILSKHPNLSASQVRNRLSSTATYLGSSFYYGKGLIN	
B.licheniformis_CAA62666	(224)	PHVAGTSAALILSKHPNLSASQVRNRLSRTATYLGSSFYYGKGLIN	
Bac.sp.NSP9.1_WP_026586291	(225)	PHVAGAAALILSKYPTLSASQVRNRLSSTATYLGSSFYYGNGLIN	
B.sonorensis_WP_077737098	(225)	PHVAGAAALILSKYPTLSASQVRNRLSSTATYLGSSFYYGNGLIN	
Bac.sp._WP_076761713	(225)	PHVAGAAALVLSKYPTLSASQVRNRLSSTATYLGSSFYYGNGLIN	
B.glycini fermentans_WP_082634659	(224)	PHVAGAAALILSKSPALASQVRDRLSSTATNLGDSFYYGKGLIN	
CN101215534-0002	(224)	PHVAGAAALILSKHPNLSASQVRNRLSSTATYLGSSFYYGKGLIN	
WO9739130-0002	(224)	PHVAGAAALILSKHPNLSASQVRNRLSSTATYLGSSFYYGKGLIN	
US8110391-0009	(224)	PHVAGAAALILSKHPNLSASQVRNRLSSTATYLGSSFYYGKGLIN	
JP2013500714-0110	(224)	PHVAGAAALILSKHPNLSASQVRNRLSSTATYLGSSFYYGKGLIN	
WO2006122655-0031	(224)	PHVAGAAALILSKHPNLSASQVRNRLSSTATYLGSSFYYGKGLIN	
US7087415-0014	(224)	PHVAGAAALILSKHPNLSASQVRNRLSSTATYLGSSFYYGKGLIN	
US20110171718-0120	(224)	PHVAGAAALILSKHPNLSASQVRNRLSSTATYLGSSFYYGKGLIN	
US20030049619-0014	(223)	PHVAGAAALILSKHPNLSASQVRNRLSSTATYLGSSFYYGKGLIN	
JP2013500714-0121	(224)	PHVAGAAALILSKHPNLSASQVRTRLSTATYLGSSFSYGRGLIN	
US7569226-0012	(224)	PHVAGAAALILSKYPTLSASQVRNRLSSTATNLGDSFYYGKGLIN	
US20050009167-0017	(224)	PHVAGAAALILSKYPTLSASQVRNRLSSTATNLGDSFYYGKGLIN	
US20110171718-0098	(224)	PHVAGAAALILSKYPTLSASQVRNRLSSTATNLGDSFYYGKGLIN	
WO2017006266-0002	(224)	PHVAGAAALILSKHPNLSASQVRNRLSSTATYLGSSFYYGKGLIN	


FIG. 1CB

7/8

271

B.lich_AprL_CAJ70731	(269)	VEAAQ	(SEQ ID NO:2)
B.licheniformis_BliD02339	(269)	VEAAQ	(SEQ ID NO:9)
B.licheniformis_AEQ38580	(270)	VEAAQ	(SEQ ID NO:180)
B.licheniformis_AKI30031	(269)	VEAAQ	(SEQ ID NO:182)
B.licheniformis_AID16241	(269)	VEAAQ	(SEQ ID NO:184)
B.licheniformis_AAG00493	(269)	VEAAQ	(SEQ ID NO:186)
B.licheniformis_AEU17777	(269)	VEAAQ	(SEQ ID NO:187)
B.licheniformis_EWH21773	(269)	VEAAQ	(SEQ ID NO:188)
B.licheniformis_AEU12640	(269)	VEAAQ	(SEQ ID NO:189)
B.licheniformis_CAA62666	(269)	VEAAQ	(SEQ ID NO:190)
Bac.sp.NSP9.1_WP_026586291	(270)	VEAAQ	(SEQ ID NO:192)
B.sonorensis_WP_077737098	(270)	VEAAQ	(SEQ ID NO:181)
Bac.sp._WP_076761713	(270)	TEAAQ	(SEQ ID NO:183)
B.glycini fermentans_WP_082634659	(269)	VEAAQ	(SEQ ID NO:185)
CN101215534-0002	(269)	VEAAQ	(SEQ ID NO:194)
WO9739130-0002	(269)	VEAAQ	(SEQ ID NO:195)
US8110391-0009	(269)	VEAAQ	(SEQ ID NO:196)
JP2013500714-0110	(269)	VEAAQ	(SEQ ID NO:197)
WO2006122655-0031	(269)	VEAAQ	(SEQ ID NO:198)
US7087415-0014	(269)	VEAAQ	(SEQ ID NO:199)
US20110171718-0120	(269)	VEAAQ	(SEQ ID NO:200)
US20030049619-0014	(268)	VEAAQ	(SEQ ID NO:201)
JP2013500714-0121	(269)	VEAAQ	(SEQ ID NO:202)
US7569226-0012	(269)	VEAAQ	(SEQ ID NO:203)
US20050009167-0017	(269)	VEAAQ	(SEQ ID NO:204)
US20110171718-0098	(269)	VEAAQ	(SEQ ID NO:205)
WO2017006266-0002	(269)	VEAAQ	(SEQ ID NO:39)

FIG. 1D

8/8**FIG. 2**

SUBSTITUTE SHEET (RULE 26)

FIG. 1A	FIG. 1AA
	FIG. 1AB

	1	45
B.lich_AprL_CAJ70731	(1)	AQTVPYGIPLIKADKVQAQGFKGANVKVAVLDTGIQASHPDLNVV
B.licheniformis_BliD02339	(1)	AQTVPYGIPLIKADKVQAQGFKGANVKVAVLDTGIQASHPDLNVV
B.licheniformis_AEQ38580	(1)	AQTVPYGIPLIKADKVQAQGFKGANVKVAVLDTGIQASHPDLNVV
B.licheniformis_AKI30031	(1)	AQTVPYGIPLIKADKVQAQGYKGANVKVAVLDTGIQASHPDLNVV
B.licheniformis_AID16241	(1)	AQTVPYGIPLIKADKVQAQGFKGANVKVAVLDTGIQASHPDLNVV
B.licheniformis_AAG00493	(1)	AQTVPYGIPLIKADKVQAQGYKGANVKVAVLDTGIQASHPDLNVV
B.licheniformis_AEU17777	(1)	AQTVPYGIPLIKADKVQAQGFKGANVKVAVLDTGIQASHPDLNVV
B.licheniformis_EWH21773	(1)	AQTVPYGIPLIKADKVQAQGYKGANVKVAVLDTGIQASHPDLNVV
B.licheniformis_AEU12640	(1)	AQTVPYGIPLIKADKLHAQGFKGANVKGAVLATGIPTSHPDNLNVV
B.licheniformis_CAA62666	(1)	GQTVPYGIPLIKADKVQAQGFKGANVKVAVLDTGIQASHPDLNVV
Bac.sp.NSP9.1_WP_026586291	(1)	AQTVPYGIPLIKADKVQAQGFKGANVKVGVIDTGIQSSHSDNLNVV
B.sonorensis_WP_077737098	(1)	AQTVPYGIPLIKADKVQAQGFKGANVKVGVIDTGIQSSHSDNLNVV
Bac.sp._WP_076761713	(1)	AQTVPYGISLIKADKVQAQGYKGANVKVGVIDTGIQSSHSDNLNVV
B.glycinifermentans_WP_082634659	(1)	AQTVPYGIPQIKADKVQAQGYKGANVKVGVLDTGIAASHSDNLNVV
CN101215534-0002	(1)	AQTVPYGVPLIKADKVQAQGFKGANVKVAVLDTGIQASHPDLNVV
WO9739130-0002	(1)	AQTVPYGIPLIKADKVQAQGFKGANVKVAVLDTGIQASHPDLNVV
US8110391-0009	(1)	AQTVPYGIPLIKADKVQAQGFKGANVKVAVLDTGIQASHPDLNVV
JP2013500714-0110	(1)	AQTVPYGIPLIKADKVQAQGFKGANVKVAVLDTGIQASHPDLNVV
WO2006122655-0031	(1)	AQTVPYGIPLIKADKVQAQGYKGANVKVAVLDTGIQASHPDLNVV
US7087415-0014	(1)	AQTVPYGIPLIKADKVQAQGYKGANVKVAVLDTGIQASHPDLNVV
US20110171718-0120	(1)	GQTVPYGIPLIKADKVQAQGFKGANVKVAVLDTGIQASHPDLNVV
US20030049619-0014	(1)	GQTVPYGIPLIKADKVQAQGFKGANVKVAVLDTGIQASHPDLNVV
JP2013500714-0121	(1)	GQTVPYGIPLIKADKVQAQGFKGANVKVAVLDTGIQASHPDLNVV
US7569226-0012	(1)	AQTVPYGIPLIKADKVQAQGYKGANVKVGIIDTGIAASHTDLKVV
US20050009167-0017	(1)	AQTVPYGIPLIKADKVQAQGYKGANVKVGIIDTGIAASHTDLKVV
US20110171718-0098	(1)	AQTVPYGIPLIKADKVQAQGYKGANVKVGIIDTGIAASHTDLKVV
WO2017006266-0002	(1)	AQTVPYGIPLIKADKVQAQGFKGANVKVAVLDTGIQASHPDLNVV

FIG. 1AA

	46	90
B.lich_AprL_CAJ70731	(46)	GGASFVAGEAYN-TDGNGHGHVAGTVAALDNTTGVLGVAPSVSL
B.licheniformis_BliD02339	(46)	GGASFVAGEAYN-TDGNGHGHVAGTVAALDNTTGVLGVAPNVSL
B.licheniformis_AEQ38580	(46)	GGASFVAGEAYN-TDGNGHGHVAGTVAALDNTTGVLGVAPSVSL
B.licheniformis_AKI30031	(46)	GGASFVAGEAYN-TDGNGHGHVAGTVAALDNTTGVLGVAPSVSL
B.licheniformis_AID16241	(46)	GGASFVAGEAYN-TDGNGHGHVAGTVAALDNTTGVLGVAPSVSL
B.licheniformis_AAG00493	(46)	GGASFVAGEAYN-TDGNGHGHVAGTVAALDNTTGVLGVAPNVSL
B.licheniformis_AEU17777	(46)	DGASFVAGEAYN-TDGNGHGHVAGTVAALDNTTGVLGVAPSVSL
B.licheniformis_EWH21773	(46)	GGASFVAGEAYN-TDGNGHGHVAGTVAALDNTTGVLGVAPSVSL
B.licheniformis_AEU12640	(46)	GGASFVAGEAYN-TDGNGHGHVAGTVAALDNTTGVLGVAPSVSL
B.licheniformis_CAA62666	(46)	GGASFVAGEAYN-TDGNGHGHVAGTVAALDNTTGVLGVAPSVSL
Bac.sp.NSP9.1_WP_026586291	(46)	GGASFVSGDSNPFIDTGNGHGHVAGTVAALDNSIGVLGVAPNVSL
B.sonorensis_WP_077737098	(46)	GGASFVSGDSNPFIDTGNGHGHVAGTVAALDNSIGVLGVAPNVSL
Bac.sp._WP_076761713	(46)	GGASFVSGDSNPYNDTGNGHGHVAGTVAALDNTTGVLGVAPNVSL
B.glycinifermentans_WP_082634659	(46)	GGASFVSGESYN-TDGNGHGHVAGTVAALDNSIGVLGVAPNVSL
CN101215534-0002	(46)	GGASFVAGEAYN-TDGNGHGHVAGTVAALDNTTGVLGVAPSVSL
WO9739130-0002	(46)	GGASFVAGEAYN-TDGNGHGHVAGTVAALDNTTGVLGVAPSVSL
US8110391-0009	(46)	GGASFVAGEAYN-TDGNGHGHVAGTVAALDNTTGVLGVAPSVSL
JP2013500714-0110	(46)	GGASFVAGEAYN-TDGNGHGHVAGTVAALDNTTGVLGVAPSVSL
WO2006122655-0031	(46)	GGASFVAGEAYN-TDGNGHGHVAGTVAALDNTTGVLGVAPNVSL
US7087415-0014	(46)	GGASFVAGEAYN-TDGNGHGHVAGTVAALDNTTGVLGVAPNVSL
US20110171718-0120	(46)	GGASFVAGEAYN-TDGNGHGHVAGTVAALDNTTGVLGVAPSVSL
US20030049619-0014	(46)	GGASFVAGEAYN-TDGNGHGHVAGTVAALDNTTG-LGVAPSVSL
JP2013500714-0121	(46)	GGASFVAGEAYN-TDGNGHGHVAGTVAALDNTTGVLGVAPSVSL
US7569226-0012	(46)	GGASFVSGESYN-TDGNGHGHVAGTVAALDNTTGVLGVAPNVSL
US20050009167-0017	(46)	GGASFVSGESYN-TDGNGHGHVAGTVAALDNTTGVLGVAPNVSL
US20110171718-0098	(46)	GGASFVSGESYN-TDGNGHGHVAGTVAALDNTTGVLGVAPNVSL
WO2017006266-0002	(46)	GGASFVAGEAYN-TDGNGHGHVAGTVAALDNTTGVLGVAPSVSL

FIG. 1AB