

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 2018222749 B2**

(54) Title
Combination therapies for treatment of BCMA-related cancers and autoimmune disorders

(51) International Patent Classification(s)
A61K 31/55 (2006.01) **A61P 35/02** (2006.01)
A61K 35/17 (2015.01) **A61P 37/00** (2006.01)
A61K 39/395 (2006.01) **A61P 37/02** (2006.01)
A61K 45/06 (2006.01) **C12N 15/62** (2006.01)
A61P 35/00 (2006.01)

(21) Application No: **2018222749** (22) Date of Filing: **2018.02.16**

(87) WIPO No: **WO18/151836**

(30) Priority Data

(31) Number	(32) Date	(33) Country
62/460,612	2017.02.17	US
62/582,270	2017.11.06	US

(43) Publication Date: **2018.08.23**

(44) Accepted Journal Date: **2024.04.18**

(71) Applicant(s)
Fred Hutchinson Cancer Center

(72) Inventor(s)
Riddell, Stanley R.;Green, Damian;Hill, Tyler

(74) Agent / Attorney
Spruson & Ferguson, GPO Box 3898, Sydney, NSW, 2001, AU

(56) Related Art
WO 2017/019496 A1
WO 2016/014565 A2
WO 2016/014789 A2
JP 2016538830 A
ANONYMOUS: "Gene Expression of Gamma Secretase (GS) Complex-Related Proteins, the Enzyme That Sheds BCMA), Among Patients with Multiple Myeloma (MM) and Effects of the GS Inhibitor LSN424354 on...", Blood, vol 128 (22), 1 Jan 2016
S. A. ALI ET AL: "T cells expressing an anti-B-cell maturation antigen chimeric antigen receptor cause remissions of multiple myeloma", BLOOD, vol. 128, no. 13, 13 July 2016, pg 1688 - 1700, DOI: 10.1182/blood-2016-04-711903
SARAH J. NAGLE ET AL: "The Promise of Chimeric Antigen Receptor Engineered T Cells in the Treatment of Hematologic Malignancies :", CANCER JOURNAL, vol. 22, no. 1, 1 January 2016 (2016-01-01), pg 27 - 33, DOI: 10.1097/PPO.0000000000000166
SARAH A. LAURENT ET AL: "[gamma]-secretase directly sheds the survival receptor BCMA from plasma cells", NATURE COMMUNICATIONS, vol. 6, 11 June 2015 (2015-06-11), pages 7333, XP055350960, DOI: 10.1038/ncomms8333

(19) World Intellectual Property
Organization
International Bureau

(43) International Publication Date
23 August 2018 (23.08.2018)



(10) International Publication Number
WO 2018/151836 A8

(51) International Patent Classification:

A61K 31/55 (2006.01) *A61P 35/02* (2006.01)
A61K 39/395 (2006.01) *A61P 37/00* (2006.01)
A61K 35/17 (2015.01) *A61K 45/06* (2006.01)
A61P 37/02 (2006.01) *C12N 15/62* (2006.01)
A61P 35/00 (2006.01)

(21) International Application Number:

PCT/US2018/000050

(22) International Filing Date:

16 February 2018 (16.02.2018)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/460,612 17 February 2017 (17.02.2017) US
62/582,270 06 November 2017 (06.11.2017) US

(71) Applicant: **FRED HUTCHINSON CANCER RESEARCH CENTER** [US/US]; 1100 Fairview Avenue North, Seattle, WA 98109 (US).

(72) Inventors: **RIDDELL, Stanley, R.**; 1763 268th Place SE, Sammamish, WA 98075 (US). **GREEN, Damian**; 2746 NE 88th Street, Seattle, WA 98115 (US). **HILL, Tyler**; 1315 N 88th Street, Seattle, WA 98103 (US).

(74) Agent: **PEPE, Jeffrey, C.** et al.; Seed Intellectual Property, Law Group LLP, 701 Fifth Avenue, Suite 5400, Seattle, WA 98104-7064 (US).

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ,

(54) Title: COMBINATION THERAPIES FOR TREATMENT OF BCMA-RELATED CANCERS AND AUTOIMMUNE DISORDERS




FIG. 7B

(57) Abstract: The present disclosure relates to methods for using BCMA-specific binding molecules (such as a BCMA-specific chimeric antigen receptor or antibody) in combination with γ -secretase inhibitors, which can be done concurrently or sequentially, to treat or prevent a B-cell related proliferative disease, such as a cancer or autoimmune disease, or the like. A BCMA-specific binding molecule in combination with γ -secretase inhibitor can be used in, for example, adoptive immunotherapy.



OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

- (84) Designated States** (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- *as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))*

Published:

- *with international search report (Art. 21(3))*
- *with sequence listing part of description (Rule 5.2(a))*

(48) Date of publication of this corrected version:

15 November 2018 (15.11.2018)

(15) Information about Correction:

see Notice of 15 November 2018 (15.11.2018)

COMBINATION THERAPIES FOR TREATMENT OF BCMA-RELATED CANCERS AND AUTOIMMUNE DISORDERS

STATEMENT OF GOVERNMENT INTEREST

This invention was made with government support under CA136551 awarded
5 by the National Institutes of Health. The government has certain rights in the invention.

STATEMENT REGARDING SEQUENCE LISTING

The Sequence Listing associated with this application is provided in text
format in lieu of a paper copy, and is hereby incorporated by reference into the
10 specification. The name of the text file containing the Sequence Listing is
360056_445WO_SEQUENCE_LISTING.txt. The text file is 14.4 KB, was created on
February 16, 2018, and is being submitted electronically via EFS-Web.

BACKGROUND

The human immune system generally protects the body from invading foreign
15 substances and pathogens. B lymphocytes, also referred to as B-cells and a component
of the immune system, produce antibodies that bind to, and in some cases mediate
destruction of, the foreign substance or pathogen. In some instances, however, the
immune system may be dysregulated and result in diseases that involve uncontrolled
proliferation of B-cells, such cancer, autoimmune disease and inflammatory disease.

20 Mature B cells and their differentiated progeny can be identified by molecules
on their cell surface, such as B cell maturation antigen (BCMA, also known as tumor
necrosis factor receptor superfamily member 17 (TNFRSF17), TNFRSF13A, and
CD269), which is expressed on plasma cells and some mature B cells. BCMA has been
shown to specifically bind to B cell activating factor (BAFF, also known as
25 TNFSF13B, TALL-1, and CD257) and a proliferation-inducing ligand (APRIL, also
known as TNFSF13, TALL-2, and CD256), which can lead to NF- κ B activation.
BCMA-targeted therapies, including adoptive transfer of BCMA-specific chimeric
antigen receptor (CAR) modified T cells, naked-BCMA specific antibodies or the
administration of BCMA-specific antibodies conjugated to a therapeutic moiety

(antibody drug conjugate, ADC, such as a radiolabel) can be used to treat some types of B cell malignancies, such as multiple myeloma, but may be limited in efficacy by the number of BCMA molecules expressed on the tumor cell surface and/or the presence of soluble BCMA in the circulation. Low BCMA surface expression on the cancer cells or soluble BCMA can limit and prevent the efficacy of therapeutic agents due to inadequate binding to the BCMA present on the surface of the tumor cells. Low levels of other target molecules on tumor cells (*e.g.*, CD19, CD20) that are targeted with antibody, antibody drug conjugates or chimeric antigen receptor T cells has been shown to limit the efficacy of these therapies, and enable cancer cells that express low levels of the target molecule to escape elimination. In the case of BCMA, the short extracellular portion of the molecule is cleaved from the cell surface and shed through the action of gamma-secretase (γ -secretase), a membrane-localized cellular enzyme involved in protein cleavage. This cleavage lowers the density of BCMA on cells such as myeloma cancer cells that express the molecule and results in elevated levels of soluble BCMA (sBCMA) in the serum of patients with certain autoimmune diseases (*e.g.*, systemic lupus erythematosis) and cancer (*e.g.*, multiple myeloma).

Currently, there remains a need in the immunotherapy field for alternative or improved compositions and methods for more efficiently treating autoimmune disease and cancer.

BRIEF SUMMARY

In a first aspect, the present invention provides a method of treating multiple myeloma in a subject, the method comprising administering to the subject a therapeutically effective amount of a T cell expressing a BCMA-specific chimeric antigen receptor (CAR) or tagged chimeric antigen receptor molecule (T-ChARM), and a therapeutically effective amount of a γ secretase inhibitor.

In a second aspect, the present invention provides a method of treating multiple myeloma in a subject, the method comprising administering to the subject a therapeutically effective amount of a T cell expressing a BCMA-specific chimeric antigen

receptor (CAR) or tagged chimeric antigen receptor molecule (T-ChARM), wherein the subject has been administered a therapeutically effective amount of a γ secretase inhibitor.

In a third aspect, the present invention provides a method of treating multiple myeloma in a subject, the method comprising administering to the subject a therapeutically effective amount of a γ secretase inhibitor, wherein the subject has been administered a therapeutically effective amount of a T cell expressing BCMA-specific chimeric antigen receptor (CAR) or tagged chimeric antigen receptor molecule (T-ChARM).

In a fourth aspect, the present invention provides use of a combination of (i) a T cell expressing a BCMA-specific chimeric antigen receptor (CAR) or tagged chimeric antigen receptor molecule (T-ChARM) and (ii) a γ secretase inhibitor in the preparation of a medicament for the treatment of multiple myeloma.

In a fifth aspect, the present invention provides use of a T cell expressing a BCMA-specific chimeric antigen receptor (CAR) or tagged chimeric antigen receptor molecule (T-ChARM) in the manufacture of a medicament for the treatment of multiple myeloma, wherein in the treatment the medicament is administered with a γ secretase inhibitor.

In a sixth aspect, the present invention provides use of a γ secretase inhibitor in the manufacture of a medicament for the treatment of multiple myeloma, wherein in the treatment the medicament is administered with a T cell expressing a BCMA-specific chimeric antigen receptor (CAR) or tagged chimeric antigen receptor molecule (T-ChARM).

In a seventh aspect, the present invention provides a kit when used for treating multiple myeloma, comprising (a) a unit dosage of a T cell expressing a BCMA-specific chimeric antigen receptor (CAR) or tagged chimeric antigen receptor molecule (T-ChARM), and (b) a unit dosage of a γ secretase inhibitor.

In certain aspects, the present disclosure is directed to treating cancer with a BCMA-specific CAR T cell and a γ secretase-inhibitor.

BRIEF DESCRIPTION OF THE DRAWINGS

Figures 1A – 1P show design and functional testing of exemplary chimeric antigen receptor (CAR) molecules of this disclosure.

(A) ILLUSTRATIONS OF EXEMPLARY CARS HAVING AN EXTRACELLULAR COMPONENT COMPRISED OF A BCMA-SPECIFIC SCFV DERIVED FROM A7D12.2 ANTIBODY ("A7") OR C11D5.3 ANTIBODY ("C11") AND A SPACER REGION (COMPRISED OF AN IGG4 HINGE REGION), A HYDROPHOBIC PORTION (COMPRISED OF A CD28 TRANSMEMBRANE DOMAIN), AND AN INTRACELLULAR COMPONENT COMPRISED OF A CD3Z EFFECTOR DOMAIN AND A 4-1BB COSTIMULATORY DOMAIN. THE SCFVS WERE CONSTRUCTED WITH THE C-TERMINAL END OF THE V_H REGION LINKED ("G₄S" VARIABLE REGION LINKER (SEQ ID NO:30)) TO THE N-TERMINAL END OF

the V_L region ("HL" orientation) or the C-terminal end of the V_L region linked to the N-terminal end of the V_H region ("LH" orientation).

(B) Flow cytometry data (carboxyfluorescein succinimidyl ester (CFSE) staining) showing proliferation of human T cells expressing the CARs shown in Figure 1A and control T cells not containing a CAR in response to BCMA-expressing tumor cells.

(C) Cytokine production (IFN- γ) by the CAR-transduced T cells shown in Figure 1B when cultured *in vitro* with the indicated BCMA⁻ (K562) or BCMA⁺ (K562 BCMA, U266, MM1.S) tumor cell lines.

(D) Specific lysis of the indicated target cell lines by the BCMA CAR-T cells.

(E) Illustration of two exemplary CARs of this disclosure, wherein the spacer region (Spacer +) can include a tag cassette, a linker module, a hinge, spacer amino acid and any combination thereof. If the CAR contains one or more tag in the spacer region, it would be referred to as a T-ChARM as described herein. The upper CAR contains an extracellular component (comprised of a BCMA-specific scFv and a spacer region that optionally contains other elements, such as a tag or linker), hydrophobic portion (comprised of a CD28 transmembrane domain), and an intracellular component (comprised of a CD3 ζ effector domain and a 4-1BB costimulatory domain). The lower CAR contains an extracellular component (comprised of a BCMA-specific scFv and a spacer region that optionally contains other elements, such as a tag or linker), hydrophobic portion (comprised of a CD28 transmembrane domain), and an intracellular component (comprised of a CD3 ζ effector domain and a CD28 costimulatory domain). Both BCMA-specific CAR/T-ChARM constructs contain a gene marker for transduction comprising a truncated human EGFR (EGFRt), which separated from the BCMA-specific CAR/T-ChARM constructs by a self-cleaving *Thoseaasigna* virus 2A (T2A) peptide sequence. Other known self-cleaving peptides can be used as well, such as porcine teschovirus-1 2A (P2A), equine rhinitis A virus (ERAV) 2A (E2A), and foot-and-mouth disease virus (FMDV) 2A (F2A).

(F) Exemplary CAR/T-ChARM constructs having different length spacer regions. sh = 12 amino acid Short spacer; 2ST = 48 amino acid spacer with two Strep tag cassettes; 3ST = 66 amino acid spacer with three Strep tag cassettes; 2ST Int = 157

amino acid intermediate length spacer with two Strep tag cassettes; and Lo = 228 amino acid spacer Long spacer. The Short and Long spacer can optionally contain a tag cassette, such as a Strep tag.

(G) Additional illustrations of exemplary CAR/T-ChARM constructs with a C11 or A7 HL scFv and having different spacer regions, optionally including STII tags.

(H) The ability of human T cells modified with BCMA-specific CARs to recognize BCMA and proliferate is measured by labeling the T cells with carboxyfluorescein (CFSE), culturing the CAR-T cells or control untransduced CFSE labeled T cells with K562 cells transduced with a polynucleotide encoding full-length BCMA (K562/BCMA), and using flow cytometry to measure the dilution of CFSE with each cell division. CFSE-labeled T cells containing different BCMA-specific CARs with different spacer lengths, but not control untransduced T cells (UT), diluted CFSE after co-culture with K562/BCMA cells. The CAR-T cells containing the 2 ST or longer spacers proliferated better than CAR-T cells expressing the short spacer.

(I, J) Cytokine release (IFN- γ , I; IL-2, J) by anti-BCMA CAR T cells with different spacer lengths in response to BCMA-expressing U266 and 8266 myeloma cells.

(K) Cell surface EGFRt and STII expression on CD8⁺ T cells transduced with anti-BCMA C11 T-ChARM or BCMA-2 CAR constructs after isolation and expansion.

(L, M) Cytokine release (IFN- γ , L; IL-2, M) by human CD4 (top panels) and CD8 (bottom panels) T cells engineered to express C11 T-ChARMs of this disclosure that include a 41BB costimulatory domain or a CD28 costimulatory domain, or by a previously disclosed anti-BCMA CAR ("BCMA-2"; see Carpenter *et al. Clin. Cancer Res.* 19:2048, 2013).

(N) Proliferation of human T cells engineered to express the disclosed C11 T-ChARMs or the BCMA-2 CAR when co-cultured with the indicated BCMA-expressing cell lines.

(O) Lysis of K562 BCMA-negative cells by CD8 T cells engineered to express C11 T-ChARMs of this disclosure (circle = 41BB costimulatory domain, square = CD28 costimulatory domain) or BCMA-2 CAR (triangle) at the indicated E:T ratios.

(P) Lysis of K562 cells transduced to express BCMA, by the engineered CD8 T cells indicated in Figure 1O, at various E:T ratios (x-axis).

Figures 2A-2Q show the production of BCMA and PD-L1 by multiple myeloma (MM) cells in culture and the effect of soluble BCMA, surface-bound BCMA, and surface-bound PD-L1 on the ability of anti-BCMA T-ChARM-T cells to recognize the tumor cells and produce IFN- γ .

(A) U266 myeloma cells were washed and plated in culture media for 1, 3, 5 and 24 hours. The media supernatant was harvested and assayed for soluble BCMA by ELISA. The data shows a time dependent increase in soluble BCMA (sBCMA) levels in the supernatant.

(B) Histograms showing BCMA expression by reference MM cells (RPMI 8226) or by exemplary patient primary MM cells having high (Pt. 1), intermediate (Pt. 2), or low/negative expression (Pt. 3) as measured by ELISA (black = anti-BCMA antibody; grey line = isotype control).

(C) Chart showing percentages of myeloma patients (n = 19) having high, intermediate, or low/negative BCMA expression in MM cells.

(D) IFN- γ production by anti-BCMA C11 T-ChARM⁺ CD8⁺ T cells in response to stimulation with patient MM cells with high (left, n = 5) or low (right, n = 4) BCMA expression (24h at 2:1 E:T (ELISA)). Significance was tested using an unpaired two-tailed T test and bars represent average IFN- γ production + SEM.

(E) Histograms showing PD-L1 expression by RPMI 8226 myeloma cells (left-most panel) and primary MM cells from 3 patients having high, low/negative, or intermediate PD-L1 expression, as measured by ELISA (diagonal shading = staining with anti-PD-L1 antibody; empty histogram = isotype control).

(F) Chart showing percentage of patients (n = 19) having high, intermediate, or low/negative PD-L1 expression in MM cells.

(G) IFN- γ production by anti-BCMA C11 T-ChARM⁺ CD8⁺ cells of this disclosure in response to patient MM cells with high (left, n = 4) or low (right, n = 5) PD-L1 expression (24h at 2:1 E:T (ELISA)). Significance was tested using an unpaired two-tailed T test and bars represent average IFN- γ production + SEM.

(H) IFN- γ production by BCMA-specific T-ChARM T cells in the presence of exogenous soluble BCMA. sBCMA was added to co-cultures of T cells expressing a BCMA-specific T-ChARM and K562 cells transduced with a polynucleotide encoding full-length BCMA (K562/BCMA⁺). There is a dose-dependent inhibition of BCMA-specific T-ChARM T cell effector function as measured by IFN- γ release into the media supernatant.

(I) Data from another dose-titration experiment showing the effect of sBCMA on IFN- γ production by BCMA-specific T-ChARM T cells recognizing K562 BCMA⁺ cells, which included adding sBCMA to culture at another, higher concentration (5000 ng/mL).

(J) Shedding of BCMA by the indicated MM cells cultured *in vitro*.

(K) sBCMA measured in bone marrow (BM) sera from patients with lower (<2% CD138⁺ cells) or higher (>2% CD138⁺ cells) disease burden.

(L) sBCMA binding to C113ST-ChARM T cells of this disclosure. T cells were incubated with the indicated levels of recombinant sBCMA (right side of diagram) and then stained with a BCMA-Fc fusion that was conjugated to APC.

(M) Flow cytometry data showing surface staining (APC-conjugated BCMA-Fc and anti-EGFRt antibody) and CD4 expression of C113ST T-ChARM and FMC63 CAR-T cells.

(N) Flow cytometry data showing IFN- γ production (y-axis) and CD4 expression (x-axis) by T cells expressing either a C113ST-ChARM ("C113ST") of this disclosure or a control anti-CD19 CAR ("FMC632"), in co-culture with target-expressing K562 cells and administered the BCMA-Fc fusion protein (left panels: 0 ng/ml BCMA-Fc; right panels: 1000 ng/ml BCMA-Fc).

(O) Dose titration showing IFN- γ release by CAR T cells in response to target cell lines as indicated in the presence or absence of exogenous recombinant BCMA (BCMA-Fc fusion). FMC63 (anti-CD19) vs. K562 CD19⁺ (control; downward-facing triangle); C113ST T-ChARM T cells vs. RPMI 8226 (upward-facing triangle), U266 (square), and K562 BCMA⁺ cells (circle). Data is representative of 2 independent experiments. Bars represent mean + SEM. P-value = < 0.05, as determined by one-way ANOVA with post-test. MFI= mean fluorescence intensity.

(P) IFN- γ production (normalized Mean Fluorescence Intensity, MFI) by the T cells shown in Figure 2O in co-culture with indicated antigen-expressing cells and in the presence or absence of BCMA-Fc (x-axis).

(Q) Cytolytic activity of CD8⁺ C113ST T-ChARM T cells against K562 BCMA⁺ target cells and of FMC63 CAR-T cells against K562 CD19⁺ target cells at varying concentrations of recombinant BCMA, analyzed by a 4h CRA at E:T ratio of 10:1. Data is representative of 2 independent experiments. Bars represent mean + SEM. P-value= $\leq$ 0.05, as determined by one way ANOVA with post-test. MFI = mean fluorescence intensity.

10 **Figures 3A-3R** show the effect of γ -secretase inhibitor (GSI) RO4929097 on levels of cell surface BCMA and other cell surface molecules on myeloma cell lines or primary myeloma cells.

(A) Cell surface BCMA was measured on four myeloma cell lines (8226, U266B1, MM1.R, H929) by flow cytometry with an anti-BCMA monoclonal antibody before, and 5 hours after (see diagrams), incubation of the myeloma cell lines with GSI RO4929097 in a range of concentrations from 0 μ M (DMSO control) to 1.0 μ M.

(B) Surface BCMA expression by MM.1R cells cultured with the indicated concentrations of RO4929097; staining with anti-BCMA antibody (black lines) compared to isotype control (grey line).

20 (C) Fold-change in surface BCMA expression by MM cell lines when cultured with the indicated concentrations of RO4929097; fold change indicated relative to untreated MM cells of the same line.

(D) Fold-change kinetics of surface BCMA expression over time by the indicated MM cells in culture with 1 μ M RO4929097.

25 (E) U266 myeloma cells were incubated for 1, 3, 5 and 24 hours in the presence of various concentrations of GSI RO429097 (0.01 μ M, 0.1 μ M and 1.0 μ M) and evaluated for surface BCMA expression by flow cytometry. BCMA expression increased in a dose-dependent manner in the presence of the GSI with the peak increase observed after 5 hours of exposure.

30 (F) Fold change of surface BCMA expression on cell lines (MM1R = triangle; U266 = square; 8226 = circle) cultured in 1 μ M GSI over time. GSI was re-

administered as a half media change every 2 days. Fold change in BCMA is defined as Treated (MFIBCMA-MFI_{iso})/Control (MFIBCMA-MFI_{iso}).. Data is representative of at least 2 independent experiments.

(G, H) Culture supernatant concentration of sBCMA in MM cell line cells
5 cultured in the presence of indicated concentrations of RO4929097.

(I) U266 myeloma cells were washed and plated in culture media in the presence of various concentrations of GSI RO4929097 (0.01 μ M, 0.1 μ M and 1.0 μ M). The media supernatant was harvested after 1, 3, 5 and 24 hours and assayed for soluble BCMA (sBCMA) by ELISA. The data shows that the amount of BCMA released from
10 the tumor cells into the supernatant over time was decreased when a GSI was present at a concentration of at least about 0.01 μ M.

(J, K) Fold change in BCMA expression (J) and supernatant sBCMA concentrations (K) at various time points after 1 μ M GSI had been removed from myeloma cell line cultures (GSI +/-) as compared to cultures with continued presence
15 of GSI (GSI +/+).

(L) Viability of indicated MM-expressing cells as measured by propidium iodide staining of cell lines cultured in 1 μ M GSI.

(M) Surface BCMA expression by primary patient MM cells cultured with the indicated concentrations of R04929097. Staining was as described in regard to Figure
20 3B.

(N) Fold change in BCMA on primary myeloma cells (n=7) cultured with vary amounts of GSI for 4 h. Primary and cell lines were cultured at 0.5×10^6 cells/mL. Fold change in BCMA is defined as Treated (MFIBCMA-MFI_{iso}) / Control (MFIBCMA-MFI_{iso}). Data is representative of 3 independent experiments with T cells derived from
25 different donors.

(O, P) Co-culture of primary myeloma cells with various concentrations of GSI for 4 hours does not affect the levels of several other cell surface molecules on tumor cells, including CS1, CD86, PD-L1, CD80 and CD38.

(Q) Staining of various surface markers on MM1R cells in the presence (black)
30 or absence (gray) of 1 μ M GSI in culture media. Isotype staining is shown as open plot.

(R) CD138+ primary myeloma cells were enriched from patient bone marrow samples, incubated for 3 hours in the presence of various concentrations of GSI RO429097 (0.01 μ M to 10 μ M) and evaluated for surface BCMA expression by flow cytometry. BCMA mean fluorescence intensity (MFI) on tumor cells is presented as
5 fold increase over that observed on tumor cells incubated without RO429097. There is an observed dose dependent upregulation of BCMA.

Figures 4A – 4C show that cytokine release following recognition of primary myeloma cells by BCMA-specific CAR-T cells is increased when the myeloma cells are pre-treated with a GSI.

10 (A) IL-2 production by BCMA CAR-T cells (BCMA-specific T-ChARM C11 3ST-CD28 and BCMA-specific T-ChARM C11 3ST-41BB) or control CD19sh CAR (short spacer)-T cells co-cultured with primary human myeloma tumor cells for 24 hours alone or with various concentrations of GSI RO429097 (0.003 μ M to 3.0 μ M).

15 (B) IFN- γ production by BMCA T-ChARM T cells co-cultured with myeloma cells at various concentrations of RO429097.

(C) Proliferation of CFSE-labeled BCMA-specific T-ChARM T cells increased in a dose-dependent manner after co-culture for 3 days with primary human myeloma tumor cells in media alone or in media containing GSI RO429097 at the indicated
20 concentrations.

Figures 5A-5R show the effect of various concentrations of GSI on CAR-T cell viability, growth, and functional activity.

(A) CD19 staining of K562 CD19⁺ and Raji cells that were cultured with or without GSI for $\pm$ 12-16 h. Isotype control shown as gray line.

25 (B) Primary human T cells were cultured in GSI RO429097 at concentrations ranging from 0.01 μ M to 100 μ M and viability was measured by trypan blue dye exclusion after 24 hours. There was no effect of GSI, at any concentration, on T cell viability.

(C) CD19 CAR-T cells were co-cultured with K562/CD19 target cells in media
30 containing various concentrations of RO429097. RO429097 inhibits CD19 CAR-T cell effector function at concentrations of $\geq$ 3 μ M when co-cultured, as determined by

measuring IL-2 (upper panel) and IFN γ (lower panel) production. Box shows the relevant therapeutic window of drug that does not inhibit CAR-T cell effector function.

(D) IL-2 production by CD19 CAR-T cells cultured with target cells in the presence of increasing concentrations of GSI RO4929097.

5 (E) IL-2 production by CD19 CAR-T cells cultured with target cells ("K562 CD19") or control cells ("K562 BCMA") in the presence of increasing concentrations of GSI RO4929097. Cells were administered the indicated amounts of the GSI and then washed (empty bars) or not (filled bars).

(F) Data from another experiment showing IL-2 production by CD19 CAR-T
10 cells after overnight co-culture with K562 BCMA⁺ or K562 CD19⁺ cells following pre-incubation with varying concentrations of GSI. After washing, GSI was either added back in (++) or left out of the co-culture (+/-) to assess the reversibility of cytokine production.

(G) Specific lysis, by CD19 CAR-T cells, of the indicated target or control cells
15 in the presence of GSI RO4929097.

(H) Proliferation of CD19 CAR-T cells cultured with CD19-expressing target cells in the presence or absence of GSI or with control cells in the absence of GSI. Cells were stained with CFSE and proliferation was measured by flow cytometry.

(I) Graphic representation of the number of cell divisions of CD19 CAR-T cells
20 in the presence of the indicated concentrations of GSI RO4929097. The width of the horizontal bars represents the proportion of CAR-T cells in culture that divided the indicated number of generations (*i.e.*, 5, 4, 3, 2, 1, or 0 generations) over the course of the experiment.

(J) Cell counts (CD8 staining) during expansion of CD19-specific CAR-T cells
25 with CD19⁺ TM LCL cells and exogenous IL-2 in the absence of GSI (circle) or the presence of GSI at 0.5 μ M (square) or 5 μ M (triangle).

(K) Cell counts of CD19-specific CAR-T cells (CD4:CD8 (1:1) expanded with CD19⁺ TM. LCL cells in the absence or presence of GSI as indicated, but without the addition of exogenous IL-2.

(L) IFN- γ concentrations in supernatants of GSI-expanded anti-CD19 CAR CD8⁺ T cells following re-stimulation with K562 cells (no antigen), K562 CD19⁺ cells, or Raji cells.

(M, N) Production of IFN- γ (M) and IL-2 (N) by the CD4:CD8 anti-CD19
5 CAR-T cell mixture following re-stimulation with the indicated cell lines in the absence of GSI or the presence of either 0.5 μ M or 5 μ M GSI.

(O) Intracellular staining showing IFN- γ production (y-axis) and CD8 expression (x-axis) by T-ChARM T cells of the present disclosure cultured with primary MM cells from 2 patients in the absence (0 μ M; left panels) or presence (1 μ M;
10 right panels) of GSI RO4929097.

(P) IFN- γ production (geometric mean fluorescence intensity (gMFI)) by T-ChARM T cells of the present disclosure cultured with primary MM cells in the presence of the indicated concentration of RO4929097 (x-axis).

(Q) IFN- γ production (normalized MFI; y-axis) by T-ChARM T cells of the
15 present disclosure cultured with primary MM cells in the presence of the indicated concentration of RO4929097 (x-axis).

(R) Proliferation of T-ChARM T cells of the present disclosure in the presence of primary MM cells that were untreated (grey shading) or treated with 1.0 μ M GSI RO4929097.

20 **Figures 6A-6C** show the effects of GSI RO4929097 on BCMA expression in a preclinical *in vivo* model of multiple myeloma.

(A) Experimental scheme for a disseminated xenograft murine myeloma model. NSG mice were irradiated (275 rad) to facilitate tumor engraftment and received human MM tumor cells (5×10^6 MM.1R) followed by GSI treatment (30 mg/kg). Subsequently,
25 mice were euthanized and blood and BM samples were taken to determine if GSI had upregulated BCMA expression on myeloma cells *in vivo*.

(B) Fold change in surface BCMA expression on myeloma cells in mice euthanized at the indicated timepoints following the second GSI administration.

(C) Levels of sBCMA in sera from mouse sacrificed at the indicated timepoints
30 after administration of RO4929097.

Figures 7A-7E show the effects of GSI RO4929097 on anti-BCMA CAR T cell therapy in the preclinical mouse MM model.

(A) Experimental scheme in which the mice received radiation followed by human MM tumor cells (5×10^6 MM.1R expressing firefly luciferase). Twenty days thereafter, mice were administered GSI (30 mg/kg) at the indicated timepoints and a single suboptimal dose of anti-BCMA T-ChARM T cells (0.33×10^6 cells, 1:1 CD4:CD8) at day 0. Bioluminescence (BLI) imaging and survival were monitored throughout.

(B) BLI images of mice taken at days 2, 17, and 16 following treatment with C113ST T-ChARM T cells (0 mg/kg 0.33×10^6 cells, 1:1 CD4:CD8, left-panels; 30 mg/kg, middle panels) or control FM63 anti-CD19 CAR T cells (0.33×10^6 cells, 1:1 CD4:CD8, 30 mg/kg, right-hand panels).

(C) Quantified luminescence data from the BLI shown in Figure 8B.

(D) Percent survival of the mice shown in Figure 8B following administration of the T-ChARM T cells.

(E) (left) Quantified luminescence data from the BLI shown in Figure 8B; (right) percent survival of the mice following administration of the T-ChARM T cells.

Figure 8 shows flow cytometric analysis of binding, by a bispecific fusion molecule with specificity for BCMA, to H929 MM cells in the presence or absence of GSI. A control bispecific fusion molecule not targeting BCMA was also tested.

DETAILED DESCRIPTION

The instant disclosure provides compositions and methods for treating autoimmune disease and cancer through the combined use of a B cell maturation antigen (BCMA)-specific binding protein, in soluble form or expressed on a cytotoxic or other cell, and a γ -secretase inhibitor (GSI). Polynucleotides encoding such BCMA-specific binding proteins can be used to generate modified host immune cells (*e.g.*, T cells) for use in, for example, adoptive immunotherapy. In certain aspects, the instant disclosure is directed to the use of such a therapy in a subject in need of treatment in combination with a GSI, which latter treatment may be administered prior to, concurrently with, or subsequent to adoptive immunotherapy (*i.e.*, immunotherapy

with a modified immune cell expressing a BCMA-specific binding protein on the cell surface). Also provided herein are immunotherapy methods comprising administering a GSI in combination with a BCMA-specific binding protein that comprises an antibody or antigen-binding portion thereof, which may in certain embodiments be conjugated or otherwise coupled to a cytotoxic drug, *e.g.*, forming an antibody-drug conjugate (ADC). The useful therapeutic applications include the treatment of a proliferative disease or disorder (such as cancer), an autoimmune disease, or aging-associated disease or disorder in the subject in which BCMA-positive cells are pathogenic.

Prior to setting forth this disclosure in more detail, it may be helpful to an understanding thereof to provide definitions of certain terms to be used herein. Additional definitions are set forth throughout this disclosure.

In the present description, any concentration range, percentage range, ratio range, or integer range is to be understood to include the value of any integer within the recited range and, when appropriate, fractions thereof (such as one tenth and one hundredth of an integer), unless otherwise indicated. Also, any number range recited herein relating to any physical feature, such as polymer subunits, size or thickness, are to be understood to include any integer within the recited range, unless otherwise indicated. As used herein, the term "about" means $\pm 20\%$ of the indicated range, value, or structure, unless otherwise indicated. It should be understood that the terms "a" and "an" as used herein refer to "one or more" of the enumerated components. The use of the alternative (*e.g.*, "or") should be understood to mean either one, both, or any combination thereof of the alternatives. As used herein, the terms "include," "have" and "comprise" are used synonymously, which terms and variants thereof are intended to be construed as non-limiting.

In addition, it should be understood that the individual compounds, or groups of compounds, derived from the various combinations of the structures and substituents described herein, are disclosed by the present application to the same extent as if each compound or group of compounds was set forth individually. Thus, selection of particular structures or particular substituents is within the scope of the present disclosure.

The term "consisting essentially of" limits the scope of a claim to the specified materials or steps, or to those that do not materially affect the basic characteristics of a claimed invention. For example, a protein domain, region, module or cassette (*e.g.*, a binding domain, hinge region, linker module, tag cassette) or a protein (which may have one or more domains, regions, modules or cassettes) "consists essentially of" a particular amino acid sequence when the amino acid sequence of a domain, region, module, cassette or protein includes extensions, deletions, mutations, or a combination thereof (*e.g.*, amino acids at the amino- or carboxy-terminus or between domains) that, in combination, contribute to at most 20% (*e.g.*, at most 15%, 10%, 8%, 6%, 5%, 4%, 3%, 2% or 1%) of the length of a domain, region, module, cassette or protein and do not substantially affect (*i.e.*, do not reduce the activity by more than 50%, such as no more than 40%, 30%, 25%, 20%, 15%, 10%, 5%, or 1%) the activity of the domain(s), region(s), module(s), cassette(s) or protein (*e.g.*, the target binding affinity of a binding protein or tag cassette).

As used herein, "proliferative disorder" refers to excessive or otherwise abnormal growth or proliferation as compared to a normal or undiseased cell. Excessive or abnormal growth includes, for example, dysregulated growth or proliferation that can occur rapidly (*e.g.*, hyperproliferation) or can occur more slowly or progressively over time (*e.g.*, multiple myeloma), within a native tissue (*e.g.*, plasmacytoma growth within bone marrow), as well as spread to/grow within a distal tissue or body site that is non-native to the diseased cell. Exemplary proliferative disorders include tumors, cancers, neoplastic tissue, carcinoma, sarcoma, malignant cells, pre malignant cells, as well as non-neoplastic or non-malignant proliferative disorders (*e.g.*, adenoma, fibroma, lipoma, leiomyoma, hemangioma, fibrosis, restenosis), or autoimmune diseases (such as rheumatoid arthritis, osteoarthritis, psoriasis, inflammatory bowel disease, or the like).

A "binding protein" (also referred to as a "binding domain," "binding region" or "binding moiety"), as used herein, refers to a molecule, such as a peptide, oligopeptide, polypeptide, or protein that possesses the ability to specifically and non-covalently associate, unite, or combine with a target molecule (*e.g.*, BCMA). A binding protein includes any naturally occurring, synthetic, semi-synthetic, or recombinantly produced

binding partner for a biological molecule, compound or other target of interest. In some embodiments, a binding protein is from an immunoglobulin or immunoglobulin-like molecule, such as an antibody or T cell receptor (TCR), which includes a functional binding domain or antigen-binding fragment thereof. Exemplary binding proteins include single chain antibody variable regions (*e.g.*, domain antibodies, sFv, scFv, Fab), BCMA ligands (*e.g.*, BAFF, APRIL and binding fragments thereof), antigen-binding regions of T cell receptors (TCRs), such as single chain TCRs (scTCRs), or synthetic polypeptides selected for the specific ability to bind to a biological molecule.

As used herein, "specifically binds" refers to an association or union of a binding domain, or a fusion protein thereof, to a target molecule with an affinity or K_a (*i.e.*, an equilibrium association constant of a particular binding interaction with units of $1/M$) equal to or greater than $10^5 M^{-1}$, while not significantly associating or uniting with any other molecules or components in a sample. Binding domains (or fusion proteins thereof) may be classified as "high affinity" binding domains (or fusion proteins thereof) or "low affinity" binding domains (or fusion proteins thereof). "High affinity" binding domains refer to those binding domains with a K_a of at least $10^7 M^{-1}$, at least $10^8 M^{-1}$, at least $10^9 M^{-1}$, at least $10^{10} M^{-1}$, at least $10^{11} M^{-1}$, at least $10^{12} M^{-1}$, or at least $10^{13} M^{-1}$. "Low affinity" binding domains refer to those binding domains with a K_a of up to $10^7 M^{-1}$, up to $10^6 M^{-1}$, up to $10^5 M^{-1}$. Alternatively, affinity may be defined as an equilibrium dissociation constant (K_d) of a particular binding interaction with units of M (*e.g.*, $10^{-5} M$ to $10^{-13} M$). In certain embodiments, a binding domain may have "enhanced affinity," which refers to a selected or engineered binding domain with stronger binding to a target antigen than a wild type (or parent) binding domain. For example, enhanced affinity may be due to a K_a (equilibrium association constant) for the target antigen that is higher than the wild type binding domain, or due to a K_d (dissociation constant) for the target antigen that is less than that of the wild type binding domain, or due to an off-rate (K_{off}) for the target antigen that is less than that of the wild type binding domain. A variety of assays are known for identifying binding domains of the present disclosure that specifically bind a particular target, as well as determining binding domain or fusion protein affinities, such as Western blot, ELISA,

and Biacore® analysis (*see also, e.g.,* Scatchard *et al., Ann. N.Y. Acad. Sci.* 51:660, 1949; and U.S. Patent Nos. 5,283,173, 5,468,614, or the equivalent).

As used herein, "heterologous" or "non-endogenous" or "exogenous" refers to any gene, protein, compound, molecule or activity that is not native to a host cell or a
5 subject, or is any gene, protein, compound, molecule or activity native to a host or host cell but has been altered or mutated such that the structure, activity or both is different as between the native and mutated molecules. In certain embodiments, heterologous, non-endogenous or exogenous molecules (*e.g.,* receptors, ligands) may not be endogenous to a host cell or subject, but instead nucleic acids encoding such molecules
10 may have been added to a host cell by conjugation, transformation, transfection, electroporation, or the like, wherein the added nucleic acid molecule may integrate into a host cell genome or can exist as extra-chromosomal genetic material (*e.g.,* as a plasmid or other self-replicating vector). The term "homologous" or "homolog" refers to a molecule or activity found in or derived from a host cell, species or strain. For
15 example, a heterologous or exogenous molecule or gene encoding the molecule may be homologous to a native host or host cell molecule or gene that encodes the molecule, respectively, but may have an altered structure, sequence, expression level or combinations thereof. A non-endogenous molecule may be from the same species, a different species or a combination thereof.

20 As used herein, the term "endogenous" or "native" refers to a gene, protein, compound, molecule or activity that is normally present in a host or host cell.

As used herein, "tag cassette" refers to a unique peptide sequence affixed to, fused to, or that is part of a protein of interest, to which a heterologous or non-endogenous cognate binding molecule (*e.g.,* receptor, ligand, antibody, or other binding
25 partner) is capable of specifically binding where the binding property can be used to detect, identify, isolate or purify, track, enrich for, or target a tagged protein or cells expressing a tagged protein, particularly when a tagged protein is part of a heterogeneous population of proteins or other material, or when cells expressing a tagged protein are part of a heterogeneous population of cells (*e.g.,* a biological sample
30 like peripheral blood). In certain embodiments, a cell expressing a tagged protein can be contacted with a heterologous or non-endogenous cognate binding molecule and

induce a biological response, such as promote cell activation, cell proliferation or cell death. In the provided fusion proteins, the ability of the tag cassette(s) to be specifically bound by the cognate binding molecule(s) is distinct from or in addition to the ability of the binding domain(s) to specifically bind to the target molecule(s). The tag cassette
5 generally is not an antigen-binding molecule, for example, is not an antibody or TCR or an antigen-binding portion thereof.

As used herein, a "hinge region" or a "hinge" refers to (a) an immunoglobulin hinge sequence (made up of, for example, upper and core regions) or a functional fragment or variant thereof, (b) a type II C-lectin interdomain (stalk) region or a
10 functional fragment or variant thereof, or (c) a cluster of differentiation (CD) molecule stalk region or a functional variant thereof. As used herein, a "wild type immunoglobulin hinge region" refers to a naturally occurring upper and middle hinge amino acid sequences interposed between and connecting the CH1 and CH2 domains (for IgG, IgA, and IgD) or interposed between and connecting the CH1 and CH3
15 domains (for IgE and IgM) found in the heavy chain of an antibody. In certain embodiments, a hinge region is human, and in particular embodiments, comprises a human IgG hinge region.

As used herein, a "spacer region" refers to one or more proteins, polypeptides, oligopeptides, peptides, domains, regions, modules, cassettes, motifs or any
20 combination thereof that join two or more proteins, polypeptides, oligopeptides, peptides, domains, regions, modules, cassettes, motifs or any combination thereof in a fusion protein. For example, a spacer region may provide a separation or spacing function to facilitate the interaction of two single chain fusion proteins, or positioning of one or more binding domains, so that the resulting polypeptide structure maintains a
25 specific binding affinity to a target molecule or maintains signaling activity (*e.g.*, effector domain activity) or both. In certain embodiments, a spacer region may comprise a "linker module" that is an amino acid sequence having from about two up to about 500 amino acids, which can provide flexibility and room for conformational movement between two regions, domains, motifs, cassettes or modules connected by a
30 linker. Exemplary linker modules include those having from one to about ten repeats of Gly_xSer_y (SEQ ID NO:31), wherein x and y are independently an integer from 0 to 10

provided that x and y are not both 0 (*e.g.*, (Gly₄Ser)₂ (SEQ ID NO:32), (Gly₃Ser)₂ (SEQ ID NO:33), Gly₂Ser, or a combination thereof such as (Gly₃Ser)₂Gly₂Ser)) (SEQ ID NO:34). In certain other embodiments, a spacer region may have a linker module that comprises one or more immunoglobulin heavy chain constant regions, such as a CH3 alone or a CH2CH3. In further embodiments, a spacer region may comprise a hinge region or a tag cassette. Each such connector component is not mutually exclusive. For example, a spacer region may comprise a hinge and one or more linker modules, or a spacer region may comprise a hinge, one or more linker modules, and one or more tag cassettes. Exemplary spacer regions can vary in length, for instance, from about five to about 500 amino acids, or from about ten to about 350 amino acids, or from about 15 to about 100 amino acids, or from about 20 to about 75 amino acids, or from about 25 to about 35 amino acids. Exemplary short spacers range from about five to about 100 amino acids (*e.g.*, 12 amino acids, 15 amino acids, 48 amino acids, 50 amino acids, 66 amino acids, 70 amino acids), intermediate spacers range from about 100 to about 200 amino acids (*e.g.*, 110 amino acids, 120 amino acids, 130 amino acids, 140 amino acids, 150 amino acids, 157 amino acids, 175 amino acids), and long spacers range from about 200 to about 500 amino acids (*e.g.*, 200 amino acids, 210 amino acids, 220 amino acids, 228 amino acids, 230 amino acids, 250 amino acids, 300 amino acids, 350 amino acids, 400 amino acids, 450 amino acids).

A "hydrophobic portion," as used herein, means any amino acid sequence having a three-dimensional structure that is thermodynamically stable in a cell membrane, and generally ranges in length from about 15 amino acids to about 30 amino acids. The structure of a hydrophobic domain may comprise an alpha helix, a beta barrel, a beta sheet, a beta helix, or any combination thereof.

As used herein, an "effector domain" is an intracellular portion of a fusion protein or receptor that can directly or indirectly promote a biological or physiological response in a cell when receiving the appropriate signal. In certain embodiments, an effector domain is part of a protein or protein complex that receives a signal when bound, or it binds directly to a target molecule, which triggers a signal from the effector domain. An effector domain may directly promote a cellular response when it contains one or more signaling domains or motifs, such as an immunoreceptor tyrosine-based

activation motif (ITAM), and such a cellular response can be aided or improved with a costimulatory domain or functional portion thereof. An exemplary protein having an effector domain is CD3 ζ . In other embodiments, an effector domain will indirectly promote a cellular response by associating with one or more other proteins that directly promote a cellular response.

A "costimulatory domain," as the term is used herein refers to a signaling moiety that provides T cells a signal which, in addition to the primary (effector) signal provided by, for instance, a CD3 ζ chain of the TCR/CD3 complex, mediates a T cell response, including activation, proliferation, differentiation, cytokine secretion, or the like. In certain embodiments, an intracellular component comprises an effector domain or functional portion thereof, a costimulatory domain or functional portion thereof, or any combination thereof.

A "variable region linker" specifically refers to a five to about 35 amino acid sequence that connects a heavy chain immunoglobulin variable region to a light chain immunoglobulin variable region or connects T cell receptor V $\alpha\beta$ and C $\alpha\beta$ chains (*e.g.*, V α -C α , V β -C β , V α -V β) or connects each V α -C α , V β -C β , V α -V β pair to a hinge or hydrophobic domain, which provides a spacer function and flexibility sufficient for interaction of the two sub-binding domains so that the resulting single chain polypeptide retains a specific binding affinity to the same target molecule as an antibody or T cell receptor. In certain embodiments, a variable region linker comprises from about ten to about 30 amino acids or from about 15 to about 25 amino acids. In particular embodiments, a variable region linker peptide comprises from one to ten repeats of Gly $_x$ Ser $_y$, wherein x and y are independently an integer from 1 to 5 (*e.g.*, Gly $_4$ Ser (SEQ ID NO:1), Gly $_3$ Ser (SEQ ID NO:2), Gly $_2$ Ser, or (Gly $_3$ Ser) $_n$ (Gly $_4$ Ser) $_1$ (SEQ ID NO:3), (Gly $_3$ Ser) $_n$ (Gly $_4$ Ser) $_n$ (SEQ ID NO:4), or (Gly $_4$ Ser) $_n$ (SEQ ID NO:5), wherein n is an integer of 1, 2, 3, 4, or 5) and wherein linked variable regions form a functional binding domain (*e.g.*, scFv, scTCR).

"Junction amino acids" or "junction amino acid residues" refer to one or more (*e.g.*, about 2-20) amino acid residues between two adjacent motifs, regions or domains of a polypeptide, such as between a binding domain and an adjacent linker region or between a hydrophobic domain and an adjacent effector domain or on one or both ends

of a linker region that links two motifs, regions or domains (*e.g.*, between a linker and an adjacent binding domain and/or between a linker and an adjacent hinge). Junction amino acids may result from the construct design of a fusion protein (*e.g.*, amino acid residues resulting from the use of a restriction enzyme site during the construction of a nucleic acid molecule encoding a fusion protein).

Terms understood by those in the art of antibody technology are each given the meaning acquired in the art, unless expressly defined differently herein. The term "antibody" refers to an intact antibody comprising at least two heavy (H) chains and two light (L) chains inter-connected by disulfide bonds, as well as an antigen-binding portion of an intact antibody that has or retains the capacity to bind a target molecule. A monoclonal antibody or antigen-binding portion thereof may be non-human, chimeric, humanized, or human, preferably humanized or human. Immunoglobulin structure and function are reviewed, for example, in Harlow *et al.*, Eds., *Antibodies: A Laboratory Manual*, Chapter 14 (Cold Spring Harbor Laboratory, Cold Spring Harbor, 1988).

For example, the terms " V_L " and " V_H " refer to the variable binding region from an antibody light and heavy chain, respectively. The variable binding regions are made up of discrete, well-defined sub-regions known as "complementarity determining regions" (CDRs) and "framework regions" (FRs). The term "CL" refers to an "immunoglobulin light chain constant region" or a "light chain constant region," *i.e.*, a constant region from an antibody light chain. The term "CH" refers to an "immunoglobulin heavy chain constant region" or a "heavy chain constant region," which is further divisible, depending on the antibody isotype into CH1, CH2, and CH3 (IgA, IgD, IgG), or CH1, CH2, CH3, and CH4 domains (IgE, IgM). A "Fab" (fragment antigen binding) is the part of an antibody that binds to antigens and includes the variable region and CH1 of the heavy chain linked to the light chain via an inter-chain disulfide bond.

As used herein, "Fc region portion" refers to the heavy chain constant region segment of the Fc fragment (the "fragment crystallizable" region or Fc region) from an antibody, which can include one or more constant domains, such as CH2, CH3, CH4, or any combination thereof. In certain embodiments, an Fc region portion includes the

CH2 and CH3 domains of an IgG, IgA, or IgD antibody or any combination thereof, or the CH3 and CH4 domains of an IgM or IgE antibody and any combination thereof. In other embodiments, a CH2CH3 or a CH3CH4 structure has sub-region domains from the same antibody isotype and are human, such as human IgG1, IgG2, IgG3, IgG4, IgA1, IgA2, IgD, IgE, or IgM (*e.g.*, CH2CH3 from human IgG1). By way of background, an Fc region is responsible for the effector functions of an immunoglobulin, such as ADCC (antibody-dependent cell-mediated cytotoxicity), CDC (complement-dependent cytotoxicity) and complement fixation, binding to Fc receptors (*e.g.*, CD16, CD32, FcRn), greater half-life *in vivo* relative to a polypeptide lacking an Fc region, protein A binding, and perhaps even placental transfer (*see* Capon *et al.*, *Nature* 337:525, 1989). In certain embodiments, an Fc region portion found in fusion proteins of the present disclosure will be capable of mediating one or more of these effector functions, or will lack one or more or all of these activities by way of, for example, one or more mutations known in the art.

In addition, antibodies have a hinge sequence that is typically situated between the Fab and Fc region (but a lower section of the hinge may include an amino-terminal portion of the Fc region). By way of background, an immunoglobulin hinge acts as a flexible spacer to allow the Fab portion to move freely in space. In contrast to the constant regions, hinges are structurally diverse, varying in both sequence and length between immunoglobulin classes and even among subclasses. For example, a human IgG1 hinge region is freely flexible, which allows the Fab fragments to rotate about their axes of symmetry and move within a sphere centered at the first of two inter-heavy chain disulfide bridges. By comparison, a human IgG2 hinge is relatively short and contains a rigid poly-proline double helix stabilized by four inter-heavy chain disulfide bridges, which restricts the flexibility. A human IgG3 hinge differs from the other subclasses by its unique extended hinge region (about four times as long as the IgG1 hinge), containing 62 amino acids (including 21 prolines and 11 cysteines), forming an inflexible poly-proline double helix and providing greater flexibility because the Fab fragments are relatively far away from the Fc fragment. A human IgG4 hinge is shorter than IgG1 but has the same length as IgG2, and its flexibility is intermediate between that of IgG1 and IgG2.

"T cell receptor" (TCR) refers to a molecule found on the surface of T cells (or T lymphocytes) that, in association with CD3, is generally responsible for recognizing antigens bound to major histocompatibility complex (MHC) molecules. The TCR has a disulfide-linked heterodimer of the highly variable α and β chains (also known as

5 TCR α and TCR β , respectively) in most T cells. In a small subset of T cells, the TCR is made up of a heterodimer of variable γ and δ chains (also known as TCR γ and TCR δ , respectively). Each chain of the TCR is a member of the immunoglobulin superfamily and possesses one N-terminal immunoglobulin variable domain, one immunoglobulin constant domain, a transmembrane region, and a short cytoplasmic tail at the C-terminal

10 end (see Janeway *et al.*, *Immunobiology: The Immune System in Health and Disease*, 3rd Ed., Current Biology Publications, p. 4:33, 1997). TCR, as used in the present disclosure, may be from various animal species, including human, mouse, rat, cat, dog, goat, horse, or other mammals. TCRs may be cell-bound (*i.e.*, have a transmembrane region or domain) or in soluble form.

15 "Major histocompatibility complex molecules" (MHC molecules) refer to glycoproteins that deliver peptide antigens to a cell surface. MHC class I molecules are heterodimers consisting of a membrane spanning α chain (with three α domains) and a non-covalently associated β 2 microglobulin. MHC class II molecules are composed of two transmembrane glycoproteins, α and β , both of which span the membrane. Each

20 chain has two domains. MHC class I molecules deliver peptides originating in the cytosol to the cell surface, where peptide:MHC complex is recognized by CD8⁺ T cells. MHC class II molecules deliver peptides originating in the vesicular system to the cell surface, where they are recognized by CD4⁺ T cells. An MHC molecule may be from various animal species, including human, mouse, rat, or other mammals.

25 A "vector" is a nucleic acid molecule that is capable of transporting another nucleic acid. Vectors may be, for example, plasmids, cosmids, viruses, or phage. An "expression vector" is a vector that is capable of directing the expression of a protein encoded by one or more genes carried by the vector when it is present in the appropriate environment.

30 "Retroviruses" are viruses having an RNA genome. "Gammaretrovirus" refers to a genus of the retroviridae family. Exemplary gammaretroviruses include mouse

stem cell virus, murine leukemia virus, feline leukemia virus, feline sarcoma virus, and avian reticuloendotheliosis viruses.

"Lentivirus" refers to a genus of retroviruses that are capable of infecting dividing and non-dividing cells. Several examples of lentiviruses include HIV (human immunodeficiency virus: including HIV type 1, and HIV type 2); equine infectious anemia virus; feline immunodeficiency virus (FIV); bovine immune deficiency virus (BIV); and simian immunodeficiency virus (SIV).

"T cells" or "cells of T cell lineage" refer to cells that show at least one phenotypic characteristic of a T cell or a precursor or progenitor thereof that distinguishes the cells from other lymphoid cells, and cells of the erythroid or myeloid lineages. Such phenotypic characteristics can include expression of one or more proteins specific for T cells (*e.g.*, CD3⁺, CD4⁺, CD8⁺), or a physiological, morphological, functional, or immunological feature specific for a T cell. For example, cells of the T cell lineage may be progenitor or precursor cells committed to the T cell lineage; CD25⁺ immature and inactivated T cells; cells that have undergone CD4 or CD8 lineage commitment; thymocyte progenitor cells that are CD4⁺CD8⁺ double positive; single positive CD4⁺ or CD8⁺; TCRαβ or TCR γδ; or mature and functional or activated T cells.

"Nucleic acid molecule", or "polynucleotide," may be in the form of RNA or DNA, which includes cDNA, genomic DNA, and synthetic DNA. A nucleic acid molecule may be double stranded or single stranded, and if single stranded, may be the coding strand or non-coding (anti-sense strand). A coding molecule may have a coding sequence identical to a coding sequence known in the art or may have a different coding sequence, which, as the result of the redundancy or degeneracy of the genetic code, or by splicing, can encode the same polypeptide.

"Treat" or "treatment" or "ameliorate" refers to medical management of a disease, disorder, or condition of a subject (*e.g.*, a human or non-human mammal, such as a primate, horse, dog, mouse, rat). For example, an appropriate dose or treatment regimen comprising BCMA-specific binding protein or a host cell expressing a BCMA-specific binding protein used in combination with a γ-secretase inhibitor (GSI) of this disclosure, and optionally an adjuvant or pre-conditioning regimen, is

administered to elicit a therapeutic or prophylactic benefit. Therapeutic or prophylactic/preventive benefit includes improved clinical outcome; lessening or alleviation of symptoms associated with a disease; decreased occurrence of symptoms; improved quality of life; longer disease-free status; diminishment of extent of disease, stabilization of disease state; delay of disease progression; remission; survival; prolonged survival; or any combination thereof.

A "therapeutically effective amount" or "effective amount" of a BCMA-specific binding protein (also referred to as a BCMA-specific or BCMA-targeted immunotherapy), a γ -secretase inhibitor, a host cell expressing a BCMA-specific binding protein, or a host cell expressing a γ -secretase inhibitor of this disclosure (e.g., BCMA-specific CAR, anti- γ -secretase antibody) refers to that amount of compound or cells sufficient to result in amelioration of one or more symptoms of the disease being treated in a statistically significant manner. When referring to an individual active ingredient or a cell expressing a single active ingredient, administered alone, a therapeutically effective dose refers to the effects of that ingredient or cell expressing that ingredient alone. When referring to a combination, a therapeutically effective dose refers to the combined amounts of active ingredients or combined adjunctive active ingredient with a cell expressing an active ingredient that results in a therapeutic effect, whether administered serially, concurrently or simultaneously. Another combination may be a cell expressing more than one active ingredient, such as two or more different BCMA-specific binding proteins or the like.

Additional definitions are provided throughout the present disclosure.

B Cell Maturation Antigen (BCMA) Binding Proteins or Molecules

In certain aspects, the present disclosure provides methods for treating a proliferative or autoimmune disease or disorder in a subject having or suspected having the disease or disorder, comprising administering to the subject a therapeutically effective amount of a BCMA-specific binding protein (or BCMA-targeted immunotherapy) and a therapeutically effective amount of a γ -secretase inhibitor. An exemplary BCMA-specific binding protein is a chimeric antigen receptor comprising an extracellular component and an intracellular component connected by a hydrophobic portion, wherein the extracellular component comprises a BCMA-specific binding

domain (e.g., BCMA-specific scFv, BCMA ligand or binding portion thereof, such as BAFF or APRIL) and optionally comprises a spacer region or hinge, and wherein the intracellular component comprises an effector domain and optionally a costimulatory domain.

5 In certain embodiments, the present disclosure provides a BCMA-targeted immunotherapy, for use with a γ -secretase inhibitor to treat a proliferative or autoimmune disease or disorder, comprising a BCMA-specific antibody or antigen-binding portion thereof, a chimeric antigen receptor (CAR), or a tagged chimeric antigen receptor molecule (T-ChARM). In certain embodiments, a BCMA-specific
10 antibody or antigen-binding portion thereof is human or humanized.

Exemplary BCMA-specific antibodies include antibodies J22.0-xi, J22.9-xi, J6M0, J6M1, J6M2, J9M0, J9M1, J9M2, 11D5-3, CA8, A7D12.2, C11 D5.3, C12A3.2, C13F12.1, 13C2, 17A5, 83A10, 13A4, 13D2, 14B11, 14E1, 29B11, 29F3, 13A7, CA7, SG1, S307118G03, S332121F02, S332126E04, S322110D07, S336105A07,
15 S335115G01, S335122F05, ET140-3, ET140-24, ET140-37, ET140-40, ET140-54, TBL-CLN1, C4.E2.1, Vicky-1, pSCHLI333, pSCHLI372, and pSCHLI373, and antigen-binding portions thereof. Various embodiments of BCMA-specific antibodies and antigen-binding portions thereof, including humanized versions, are disclosed in, for example, PCT Publication Nos. WO 2002/066516, WO 2007/062090,
20 WO 2010/104949, WO 2011/108008, WO 2012/163805, WO 2014/068079, WO 2015/166073, WO 2014/122143, WO 2014/089335, WO 2016/090327, and WO 2016/079177; Ryan *et al.*, *Mol. Cancer. Ther.* 6(11):3009, 2007; and Abbas *et al.*, *Blood* 128:1688, 2016, which BCMA-specific antibodies, antigen-binding portions thereof and humanized versions are all incorporated herein by reference in their
25 entirety. Variable domains and scFv molecules from these BCMA-specific antibodies can be used as a binding domain in any of the T-ChARM and CAR proteins mentioned herein.

Antigen binding portions or domains obtained from BCMA-specific antibodies of present disclosure and useful in the methods disclosed herein include, for example,
30 domain antibodies, sFvs, single chain Fv fragments (scFvs), Fabs, F(ab')₂, nanobodies, tandem scFvs, scFv-Fcs, scFv dimers, scFv zippers, diabodies, minibodies, triabodies,

tetrabodies, Fabs, F(ab)'2s, scFabs, miniantibodies, nanobodies, nanobody-HSAs, Bispecific T cell Engagers (BiTEs), DARs, scDiabodies, scDiabody-CH3s, or scFv-CH3 Knobs-Into-Holes (KIH) assemblies.

In certain embodiments, a BCMA-specific binding protein comprises a
5 bispecific or multispecific antibody (or an antigen-binding portion thereof) comprising a first binding region (*e.g.*, a heavy chain variable region, a light chain variable region, or both) that is specific for BCMA and at least one other binding region that is specific for a second target (*e.g.*, a BCMA epitope that is different from the epitope of the first binding region, or an epitope of a non-BCMA target, such as, for example, a tumor-
10 associated antigen that is not BCMA (for example, CD19 (*e.g.*, blinatumomab, MOR-208, SGN-19A, SAR3419, coltuximabravtansine, denituzumabmafodotin, taplitumomabpaptox, XmAb 5574, MDI-551, Merck patent anti-CD19 aka B4 aka DI-B4, XmAb 5871, MDX-1342, AFM11), CD20 (*e.g.*, rituximab, ofatumumab, ocrelizumab), CD38 (*e.g.*, daratumumab or isatuximab (SAR650984)), CD45, or a cell
15 surface protein expressed on an immune effector cell, such as a T cell (*e.g.*, CD3), an NK cell (*e.g.*, CD56), or an NK-T cell (*e.g.*, NK1.1), or another non-tumor-associated antigen or target.

In particular embodiments, the present disclosure provides a BCMA-specific binding protein alone or expressed as a T-ChARM in a cell, for use with a γ -secretase
20 inhibitor (GSI). An exemplary T-ChARM comprises an extracellular component and an intracellular component connected by a hydrophobic portion, wherein the extracellular component comprises a binding domain that specifically binds BCMA, an optional spacer region, a tag cassette, and a hinge region, and wherein the intracellular component comprises an effector domain and optionally a costimulatory domain (*e.g.*, a
25 functional domain or portion from 4-1BB, a functional domain or portion from CD28, or both). In certain embodiments, a T-ChARM binding domain comprises a BCMA-specific scFv, a BCMA-specific scTCR, or a BCMA ligand or binding portion thereof (*e.g.*, BAFF, APRIL), optionally wherein the BCMA-specific scFv is human or humanized. Various embodiments of the T-ChARMs are disclosed in PCT
30 Publication No. WO 2015/095895, which T-ChARM scaffolds are incorporated herein by reference in their entirety.

Exemplary tag cassettes include Strep tag (which refers to the original Strep® tag, Strep® tag II, or any variant thereof; *see, e.g.*, U.S. Patent No. 7,981,632, which Strep tags are incorporated herein by reference), His tag, Flag tag, Xpress tag, Avi tag, Calmodulin tag, Polyglutamate tag, HA tag, Myc tag, Nus tag, S tag, SBP tag, Softag 1, 5 Softag 3, V5 tag, CREB-binding protein (CBP), glutathione S-transferase (GST), maltose binding protein (MBP), green fluorescent protein (GFP), Thioredoxin tag, or any combination thereof. In some embodiments, a tag cassette may be a genetically engineered affinity site, such as a minimal chelation site (*e.g.*, HGGHHG, SEQ ID NO.:6). In certain embodiments, a tag cassette is a Strep tag having an amino acid 10 sequence of Trp-Ser-His-Pro-Gln-Phe-Glu-Lys (SEQ ID NO.:7) or Trp-Arg-His-Pro-Gln-Phe-Gly-Gly (SEQ ID NO.:8).

Tag cassettes may be present in single or multiple copies in fusion proteins of this disclosure. For example, a BCMA-specific binding protein of this disclosure can have one, two, three, four or five tag cassettes (*e.g.*, Strep tag). In certain embodiments, 15 an extracellular component of a BCMA-specific T-ChARM includes one tag cassette, two tag cassettes, three tag cassettes, four tag cassettes, or five tag cassettes. Each of the plurality of tag cassettes may be the same or different.

In certain embodiments, a tag cassette comprises from about five to about 500 amino acids, or from about six to about 100 amino acids, or from about seven to about 20 50 amino acids, or from about eight to about 20 amino acids. In some embodiments, a tag cassette has seven to ten amino acids. Preferably, a tag cassette is non-immunogenic or minimally immunogenic. Essentially, a tag cassette can function as a handle or beacon to allow for the identification, enrichment, isolation, promotion of proliferation, activation, tracking, or elimination of cells expressing a BCMA-specific 25 T-ChARM.

In further embodiments, the present disclosure provides a BCMA-specific binding protein, for use with a γ -secretase inhibitor, that is a chimeric antigen receptor (CAR), which comprises an extracellular component and an intracellular component connected by a hydrophobic portion, wherein the extracellular component comprises a 30 binding domain that specifically binds BCMA and a hinge region, and wherein the intracellular component comprises an effector domain and optionally a costimulatory

domain. In certain embodiments, a CAR binding domain comprises a BCMA-specific scFv, a BCMA-specific scTCR, a BCMA-specific TCR binding domain (*see, e.g.,* Walseng *et al., Scientific Reports* 7:10713 (2017), the TCR CAR constructs of which are hereby incorporated by reference in their entirety), or a BCMA ligand or binding
5 portion thereof (*e.g.,* BAFF, APRIL), optionally wherein the BCMA-specific scFv is human or humanized. In any of these embodiments, a BCMA-specific binding protein in the form of a CAR may be expressed on the surface of a cell, such as an immune system cell (*e.g.,* T cell).

A BCMA-specific T-ChARM or CAR may be cell-bound (*e.g.,* expressed on a
10 cell surface) or in soluble form. In certain embodiments, polynucleotides encoding BCMA-specific T-ChARM or CAR proteins may be codon optimized to enhance or maximize expression in a host cell, such as a T cell (Scholten *et al., Clin. Immunol.* 119:135, 2006).

In certain embodiments, a hinge present in a BCMA-specific T-ChARM or
15 CAR of this disclosure may be an immunoglobulin hinge region, such as a wild type immunoglobulin hinge region or an altered immunoglobulin hinge region thereof. In certain embodiments, a hinge is a wild type human immunoglobulin hinge region. In certain other embodiments, one or more amino acid residues may be added at the amino- or carboxy-terminus of a wild type immunoglobulin hinge region as part of a
20 fusion protein construct design. For example, one, two or three additional junction amino acid residues may be present at the hinge amino-terminus or carboxy-terminus, or a hinge may contain a terminal or internal deletion and have added back one, two or three additional junction amino acid residues.

In certain embodiments, a hinge is an altered immunoglobulin hinge in which
25 one or more cysteine residues in a wild type immunoglobulin hinge region is substituted with one or more other amino acid residues. Exemplary altered immunoglobulin hinges include an immunoglobulin human IgG1, IgG2 or IgG4 hinge region having one, two or three cysteine residues found in a wild type human IgG1, IgG2 or IgG4 hinge substituted by one, two or three different amino acid residues (*e.g.,* serine or alanine). In
30 certain embodiments, a hinge polypeptide comprises or is a sequence that is at least 80%, at least 81%, at least 82%, at least 83%, at least 84%, at least 85%, at least 86%,

at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% identical to a wild type immunoglobulin hinge region, such as a wild type human IgG1 hinge, a wild type human IgG2 hinge, or a wild type human IgG4 hinge.

5 In further embodiments, a hinge present in a BCMA-specific T-ChARM or CAR of this disclosure may be a hinge that is not based on or derived from an immunoglobulin hinge (*i.e.*, not a wild type immunoglobulin hinge or an altered immunoglobulin hinge). Examples of such hinges include peptides of about five to about 150 amino acids of the stalk region of type II C-lectins or CD molecules,
10 including peptides of about eight to about 25 amino acids or peptides of about seven to about 18 amino acids, or variants thereof.

A "stalk region" of a type II C-lectin or CD molecule refers to the portion of the extracellular domain of the type II C-lectin or CD molecule that is located between the C-type lectin-like domain (CTLCD; *e.g.*, similar to CTLCD of natural killer cell receptors)
15 and the hydrophobic portion (transmembrane domain). For example, the extracellular domain of human CD94 (GenBank Accession No. AAC50291.1) corresponds to amino acid residues 34-179, but the CTLCD corresponds to amino acid residues 61-176, so the stalk region of the human CD94 molecule comprises amino acid residues 34-60, which are located between the hydrophobic portion (transmembrane domain) and CTLCD (*see*
20 Boyington *et al.*, *Immunity* 10:75, 1999; for descriptions of other stalk regions, *see also* Beavil *et al.*, *Proc. Nat'l. Acad. Sci. USA* 89:753, 1992; and Figdor *et al.*, *Nat. Rev. Immunol.* 2:77, 2002). These type II C-lectin or CD molecules may also have junction amino acids between the stalk region and the transmembrane region or the CTLCD. In another example, the 233 amino acid human NKG2A protein (GenBank Accession No.
25 P26715.1) has a hydrophobic portion (transmembrane domain) ranging from amino acids 71-93 and an extracellular domain ranging from amino acids 94-233. The CTLCD comprises amino acids 119-231, and the stalk region comprises amino acids 99-116, which may be flanked by additional junction amino acids. Other type II C-lectin or CD molecules, as well as their extracellular ligand-binding domains, stalk regions, and
30 CTLCDs are known in the art (*see, e.g.*, GenBank Accession Nos. NP_001993.2;

AAH07037.1; NP_001773.1; AAL65234.1; CAA04925.1; for the sequences of human CD23, CD69, CD72, NKG2A and NKG2D and their descriptions, respectively).

A "derivative" of a stalk region hinge, or fragment thereof, of a type II C-lectin or CD molecule includes about an eight to about 150 amino acid sequence in which one, two, or three amino acids of the stalk region of a wild type type II C-lectin or CD molecule have a deletion, insertion, substitution, or any combination thereof. For instance, a derivative can comprise one or more amino acid substitutions and/or an amino acid deletion. In certain embodiments, a derivative of a stalk region is more resistant to proteolytic cleavage as compared to the wild-type stalk region sequence, such as those derived from about eight to about 20 amino acids of NKG2A, NKG2D, CD23, CD64, CD72, or CD94.

In certain embodiments, stalk region hinges may comprise from about seven to about 18 amino acids and can form an α -helical coiled coil structure. In certain embodiments, stalk region hinges contain 0, 1, 2, 3, or 4 cysteines. Exemplary stalk region hinges include fragments of the stalk regions, such as those portions comprising from about ten to about 150 amino acids from the stalk regions of CD69, CD72, CD94, NKG2A and NKG2D.

Alternative hinges that can be used in BCMA-specific T- ChARMs and CARs of this disclosure are from portions of cell surface receptors (interdomain regions) that connect immunoglobulin V-like or immunoglobulin C-like domains. Regions between Ig V-like domains where the cell surface receptor contains multiple Ig V-like domains in tandem and between Ig C-like domains where the cell surface receptor contains multiple tandem Ig C-like regions are also contemplated as hinges useful in BCMA-specific T-ChARMs and CARs of this disclosure. In certain embodiments, hinge sequences comprised of cell surface receptor interdomain regions may further contain a naturally occurring or added motif, such as an IgG core hinge sequence to provide one or more disulfide bonds to stabilize the BCMA-specific T-ChARM or CAR dimer formation. Examples of hinges include interdomain regions between the Ig V-like and Ig C-like regions of CD2, CD4, CD22, CD33, CD48, CD58, CD66, CD80, CD86, CD150, CD166, or CD244.

In certain embodiments, hinge sequences have from about 5 to about 150 amino acids, about 5 to about 10 amino acids, about 10 to about 20 amino acids, about 20 to about 30 amino acids, about 30 to about 40 amino acids, about 40 to about 50 amino acids, about 50 to about 60 amino acids, about 5 to about 60 amino acids, about 5 to about 40 amino acids, for instance, about 8 to about 20 amino acids or about 10 to about 15 amino acids. The hinges may be primarily flexible, but may also provide more rigid characteristics or may contain primarily α -helical structure with minimal β -sheet structure.

In certain embodiments, a hinge sequence is stable in plasma and serum, and is resistant to proteolytic cleavage. For example, the first lysine in an IgG1 upper hinge region may be mutated or deleted to minimize proteolytic cleavage, and hinges may include junction amino acids. In some embodiments, a hinge sequence may contain a naturally occurring or added motif, such as an immunoglobulin hinge core structure CPPCP (SEQ ID NO.:9) that confers the capacity to form a disulfide bond or multiple disulfide bonds to stabilize dimer formation.

A hydrophobic portion contained in a BCMA-specific binding protein of the present disclosure (*e.g.*, BCMA-specific T-ChARM or CAR) will allow a BCMA-specific binding protein of this disclosure to associate with a cellular membrane such that a portion of the binding protein will be located extracellularly (*e.g.*, tag cassette, binding domain) and a portion will be located intracellularly (*e.g.*, effector domain, costimulatory domain). A hydrophobic portion will generally be disposed within the cellular membrane phospholipid bilayer. In certain embodiments, one or more junction amino acids may be disposed between and connecting a hydrophobic portion with an effector domain, or disposed between and connecting a hydrophobic portion with a spacer region, or disposed between and connecting a hydrophobic portion with a tag cassette.

In certain embodiments, a hydrophobic domain is a transmembrane domain, such as one derived from an integral membrane protein (*e.g.*, receptor, cluster of differentiation (CD) molecule, enzyme, transporter, cell adhesion molecule, or the like). In particular embodiments, a hydrophobic portion is a transmembrane domain from CD4, CD8, CD27, or CD28.

An intracellular component contained on a BCMA-specific binding protein of the present disclosure (*e.g.*, BCMA-specific T-ChARM or CAR) will be capable of transmitting functional signals to a cell. In certain embodiments, a BCMA-specific T-ChARM or CAR will dimerize with a second single chain T-ChARM or CAR, respectively, wherein the dimerization allows the intracellular component comprising an effector domain and optionally a costimulatory domain to be in close proximity and promote signal transduction when exposed to the proper signal. In addition to forming such dimer protein complexes, the effector domains and optional costimulatory domains may further associate with other signaling factors, such as costimulatory factors, to form multiprotein complexes that produce an intracellular signal. In certain embodiments, an effector domain will indirectly promote a cellular response by associating with one or more other proteins that directly promote a cellular response. An intracellular component may include one, two, three or more receptor signaling domains (*e.g.*, effector domains), costimulatory domains, or combinations thereof. Any intracellular component comprising an effector domain or functional portion thereof, a costimulatory domain or functional portion thereof, or any combination thereof from any of a variety of signaling molecules (*e.g.*, signal transduction receptors) may be used in the BCMA-specific binding proteins of this disclosure.

An intracellular component may have an effector or costimulatory domain useful in the BCMA-specific binding proteins of this disclosure, which may be based on or from a protein of a Wnt signaling pathway (*e.g.*, LRP, Ryk, ROR2), NOTCH signaling pathway (*e.g.*, NOTCH1, NOTCH2, NOTCH3, NOTCH4), Hedgehog signaling pathway (*e.g.*, PTCH, SMO), receptor tyrosine kinases (RTKs) (*e.g.*, epidermal growth factor (EGF) receptor family, fibroblast growth factor (FGF) receptor family, hepatocyte growth factor (HGF) receptor family, Insulin receptor (IR) family, platelet-derived growth factor (PDGF) receptor family, vascular endothelial growth factor (VEGF) receptor family, tropomyosin receptor kinase (Trk) receptor family, ephrin (Eph) receptor family, AXL receptor family, leukocyte tyrosine kinase (LTK) receptor family, tyrosine kinase with immunoglobulin-like and EGF-like domains 1 (TIE) receptor family, receptor tyrosine kinase-like orphan (ROR) receptor family, discoidin domain (DDR) receptor family, rearranged during transfection (RET) receptor

family, tyrosine-protein kinase-like (PTK7) receptor family, related to receptor tyrosine kinase (RYK) receptor family, muscle specific kinase (MuSK) receptor family); G-protein-coupled receptors, GPCRs (Frizzled, Smoothed); serine/threonine kinase receptors (BMPR, TGFR); or cytokine receptors (IL-1R, IL-2R, IL-7R, IL-15R).

5 In certain embodiments, an effector domain comprises a lymphocyte receptor signaling domain or comprises an amino acid sequences having one or a plurality of immunoreceptor tyrosine-based activation motifs (ITAMs). In still further embodiments, an effector domain comprises a cytoplasmic portion that associates with a cytoplasmic signaling protein, wherein the cytoplasmic signaling protein is a
10 lymphocyte receptor or signaling domain thereof, a protein comprising a plurality of ITAMs, a costimulatory factor, or any combination thereof.

Exemplary effector and costimulatory domains include those based on or derived from 4-1BB, CD3 ϵ , CD3 δ , CD3 ζ , CD27, CD28, CD79A, CD79B, CARD11, DAP10, FcR α , FcR β , FcR γ , Fyn, HVEM, ICOS, Lck, LAG3, LAT, LRP, NOTCH1,
15 Wnt, NKG2D, OX40, ROR2, Ryk, SLAMF1, Slp76, pT α , TCR α , TCR β , TRIM, Zap70, PTCH2, or any combination thereof.

In particular embodiments, BCMA-specific binding proteins of this disclosure comprise (a) an effector domain from CD3 ζ or functional portion thereof and a costimulatory domain from CD28 or functional portion thereof, (b) an effector domain
20 from CD3 ζ or functional portion thereof and a costimulatory domain from 4-1BB or functional portion thereof, or (c) (a) an effector domain from CD3 ζ or a functional portion thereof and a costimulatory domain from CD28 and 4-1BB or functional portions thereof.

γ -Secretase Inhibitors (GSI)

25 By way of background, γ -secretase is a multi-subunit integral membrane protease complex, including presenilin (PS), nicastrin (NCT), anterior pharynx-defective 1 (APH-1), and presenilin enhancer 2 (PEN-2). PS is the catalytic subunit that is an aspartate protease capable of forming a hydrophilic catalytic pore buried within the membrane (Takasugi *et al.*, *Nature* 422:438-41, 2003), which cleaves single
30 pass transmembrane proteins within the transmembrane domain. NCT is a type I membrane glycoprotein with a large extracellular domain (ECD), which ECD captures

the amino-terminus of the substrate as a primary substrate receptor for γ -secretase (Shah *et al.*, *Cell* 122:435-47, 2005). The γ -secretase complex plays a role in a processing of a variety of substrates, including Notch, CD44, Cadherins, and ephrin B2, as well as cleaving amyloid precursor protein into amyloid beta peptide that is implicated in Alzheimer's disease. The γ -secretase complex is also known to cleave B-cell maturation antigen (BCMA) (Laurent *et al.*, *Nature Communications* 6, 2015), which is a therapeutic target in various cancers, including multiple myeloma.

Exemplary γ -secretase inhibitors (GSIs) include small molecules, peptidomimetic compounds or γ -secretase-specific binding proteins. A GSI can target any one or more of the γ -secretase complex proteins, including presenilin 1 (PS1), presenilin 2 (PS2), nicastrin (NCT), anterior pharynx-defective 1 (APH-1), and presenilin enhancer 2 (PEN-2), provided that the γ -secretase cleavage activity is reduced compared to uninhibited γ -secretase. In certain embodiments, the γ -secretase activity is reduced at least about 50%, at least about 60%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, or 100%. Assays for measuring γ -secretase activity are known in the art (*see, e.g.*, Laurent *et al.*, 2015). For example, the level of soluble BCMA can be a surrogate measure for γ -secretase activity. Representative small molecule GSIs, for use with a BCMA-targeted immunotherapy to treat a proliferative or autoimmune disease or disorder, include avagacestat, DAPT, BMS-906024, BMS-986115, LY411575, MK-0752, PF-03084014, RO4929097, semagacestat, YO-01027, and any combination thereof.

Other GSIs are γ -secretase-specific binding proteins, such as antibodies or antigen binding portions thereof that a γ -secretase complex or a γ -secretase complex protein, such as presenilin 1 (PS1), presenilin 2 (PS2), nicastrin (NCT), anterior pharynx-defective 1 (APH-1), and presenilin enhancer 2 (PEN-2). An exemplary γ -secretase-specific binding protein is a nicastrin-specific binding protein, such as antibodies scFvG9, A5226A, 2H6, 10C11, and antigen binding fragments thereof.

Binding Domains

A binding domain may be any peptide that specifically binds BCMA or specifically inhibits γ -secretase activity as described herein. Sources of binding

domains include antibody variable regions from various species (which can be in the form of antibodies, sFvs, scFvs, Fabs, scFv-based grababody, or soluble VH domain or domain antibodies), including human, rodent, avian, or ovine. Additional sources of binding domains include variable regions of antibodies from other species, such as camelid (from camels, dromedaries, or llamas; Ghahroudi *et al.*, *FEBS Lett.* 414:521, 1997; Vincke *et al.*, *J. Biol. Chem.* 284:3273, 2009; Hamers-Casterman *et al.*, *Nature* 363:446, 1993 and Nguyen *et al.*, *J. Mol. Biol.* 275:413, 1998), nurse sharks (Roux *et al.*, *Proc. Nat'l. Acad. Sci. (USA)* 95:11804, 1998), spotted ratfish (Nguyen *et al.*, *Immunogen.* 54:39, 2002), or lamprey (Herrin *et al.*, *Proc. Nat'l. Acad. Sci. (USA)* 105:2040, 2008 and Alder *et al.* *Nat. Immunol.* 9:319, 2008). These antibodies can form antigen-binding regions using only a heavy chain variable region, *i.e.*, these functional antibodies are homodimers of heavy chains only (referred to as "heavy chain antibodies") (Jespers *et al.*, *Nat. Biotechnol.* 22:1161, 2004; Cortez-Retamozo *et al.*, *Cancer Res.* 64:2853, 2004; Baral *et al.*, *Nature Med.* 12:580, 2006; and Barthelemy *et al.*, *J. Biol. Chem.* 283:3639, 2008).

An alternative source of binding domains of this disclosure includes sequences that encode random peptide libraries or sequences that encode an engineered diversity of amino acids in loop regions of alternative non-antibody scaffolds, such as scTCR (*see, e.g.*, Lake *et al.*, *Int. Immunol.* 11:745, 1999; Maynard *et al.*, *J. Immunol. Methods* 306:51, 2005; U.S. Patent No. 8,361,794), fibrinogen domains (*see, e.g.*, Weisel *et al.*, *Science* 230:1388, 1985), Kunitz domains (*see, e.g.*, US Patent No. 6,423,498), designed ankyrin repeat proteins (DARPs) (Binz *et al.*, *J. Mol. Biol.* 332:489, 2003 and Binz *et al.*, *Nat. Biotechnol.* 22:575, 2004), fibronectin binding domains (adnectins or monobodies) (Richards *et al.*, *J. Mol. Biol.* 326:1475, 2003; Parker *et al.*, *Protein Eng. Des. Selec.* 18:435, 2005 and Hackel *et al.* (2008) *J. Mol. Biol.* 381:1238-1252), cysteine-knot miniproteins (Vita *et al.* (1995) *Proc. Nat'l. Acad. Sci. (USA)* 92:6404-6408; Martin *et al.* (2002) *Nat. Biotechnol.* 21:71, 2002 and Huang *et al.* (2005) *Structure* 13:755, 2005), tetratricopeptide repeat domains (Main *et al.*, *Structure* 11:497, 2003 and Cortajarena *et al.*, *ACS Chem. Biol.* 3:161, 2008), leucine-rich repeat domains (Stumpp *et al.*, *J. Mol. Biol.* 332:471, 2003), lipocalin domains (*see, e.g.*, WO 2006/095164, Beste *et al.*, *Proc. Nat'l. Acad. Sci. (USA)* 96:1898, 1999 and Schönfeld

et al., *Proc. Nat'l. Acad. Sci. (USA)* 106:8198, 2009), V-like domains (*see, e.g.*, US Patent Application Publication No. 2007/0065431), C-type lectin domains (Zelensky and Gready, *FEBS J.* 272:6179, 2005; Beavil *et al.*, *Proc. Nat'l. Acad. Sci. (USA)* 89:753, 1992 and Sato *et al.*, *Proc. Nat'l. Acad. Sci. (USA)* 100:7779, 2003), mAb² or FcabTM (*see, e.g.*, PCT Patent Application Publication Nos. WO 2007/098934; WO 2006/072620), armadillo repeat proteins (*see, e.g.*, Madhurantakam *et al.*, *Protein Sci.* 21: 1015, 2012; PCT Patent Application Publication No. WO 2009/040338), affilin (Ebersbach *et al.*, *J. Mol. Biol.* 372: 172, 2007), affibody, avimers, knottins, fynomers, atrimers, cytotoxic T-lymphocyte associated protein-4 (Weidle *et al.*, *Cancer Gen. Proteo.* 10:155, 2013) or the like (Nord *et al.*, *Protein Eng.* 8:601, 1995; Nord *et al.*, *Nat. Biotechnol.* 15:772, 1997; Nord *et al.*, *Euro. J. Biochem.* 268:4269, 2001; Binz *et al.*, *Nat. Biotechnol.* 23:1257, 2005; Boersma and Plückthun, *Curr. Opin. Biotechnol.* 22:849, 2011).

Binding domains of this disclosure can be generated as described herein or by a variety of methods known in the art (*see, e.g.*, U.S. Patent Nos. 6,291,161 and 6,291,158). For example, binding domains of this disclosure may be identified by screening a Fab phage library for Fab fragments that specifically bind to a target of interest (*e.g.*, BCMA, γ -secretase complex component such as presenilin or nicastrin) (*see* Hoet *et al.*, *Nat. Biotechnol.* 23:344, 2005). Additionally, traditional strategies for hybridoma development using a target of interest (*e.g.*, BCMA, γ -secretase complex component such as presenilin or nicastrin) as an immunogen in convenient systems (*e.g.*, mice, HuMAb mouse®, TC mouseTM, KM-mouse®, llamas, chicken, rats, hamsters, rabbits, *etc.*) can be used to develop binding domains of this disclosure.

In some embodiments, a binding domain is a single chain Fv fragment (scFv) that comprises V_H and V_L regions specific for a target of interest (*e.g.*, BCMA, γ -secretase complex component such as presenilin or nicastrin). In certain embodiments, the V_H and V_L regions are human. Exemplary V_H and V_L regions include the segments of anti-BCMA specific antibodies J22.0-xi, J22.9-xi, J6M0, J6M1, J6M2, J9M0, J9M1, J9M2, 11D5-3, CA8, A7D12.2, C11 D5.3, C12A3.2, C13F12.1, 13C2, 17A5, 83A10, 13A4, 13D2, 14B11, 14E1, 29B11, 29F3, 13A7, CA7, SG1, S307118G03, S332121F02, S332126E04, S322110D07, S336105A07, S335115G01,

S335122F05, ET140-3, ET140-24, ET140-37, ET140-40, ET140-54, TBL-CLN1, C4.E2.1, Vicky-1, pSCHLI333, pSCHLI372, or pSCHLI373. Other exemplary V_H and V_L regions include the segments of anti-nicastrin specific antibodies or antigen binding fragments thereof from scFvG9, A5226A, 2H6, or 10C11.

5 In certain embodiments, a binding domain comprises or is a sequence that is at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, at least 99.5%, or 100% identical to an amino acid sequence of a light chain variable region (V_L) (*e.g.*, from anti-BCMA J22.0-xi, J22.9-xi, J6M0, J6M1, J6M2, J9M0, J9M1, J9M2, 11D5-3, CA8, A7D12.2, C11
10 D5.3, C12A3.2, or C13F12.1, 13C2, 17A5, 83A10, 13A4, 13D2, 14B11, 14E1, 29B11, 29F3, 13A7, CA7, S307118G03, S332121F02, S332126E04, S322110D07, S336105A07, S335115G01, S335122F05, ET140-3, ET140-24, ET140-37, ET140-40, ET140-54, TBL-CLN1, C4.E2.1, Vicky-1, pSCHLI333, pSCHLI372, or pSCHLI373; or from anti-nicastrin scFvG9, A5226A, 2H6, or 10C11) or to a heavy chain variable
15 region (V_H) (*e.g.*, from anti-BCMA J22.0-xi, J22.9-xi, J6M0, J6M1, J6M2, J9M0, J9M1, J9M2, 11D5-3, CA8, A7D12.2, C11 D5.3, C12A3.2, C13F12.1, 13C2, 17A5, 83A10, 13A4, 13D2, 14B11, 14E1, 29B11, 29F3, 13A7, CA7, SG1, S307118G03, S332121F02, S332126E04, S322110D07, S336105A07, S335115G01, S335122F05, ET140-3, ET140-24, ET140-37, ET140-40, ET140-54, TBL-CLN1, C4.E2.1, Vicky-1,
20 pSCHLI333, pSCHLI372, or pSCHLI373; or from anti-nicastrin scFvG9, A5226A, 2H6, or 10C11), or both, wherein each CDR comprises zero changes or at most one, two, or three changes, from a monoclonal antibody or fragment or derivative thereof that specifically binds to target of interest (*e.g.*, BCMA, γ -secretase complex component such as presenilin or nicastrin).

25 In certain embodiments, a binding domain V_H region of the present disclosure can be derived from or based on a V_H of a known monoclonal antibody and contains one or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10) insertions, one or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10) deletions, one or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10) amino acid substitutions (*e.g.*, conservative amino acid substitutions or non-conservative amino acid
30 substitutions), or a combination of the above-noted changes, when compared with the V_H of a known monoclonal antibody. An insertion, deletion or substitution may be

anywhere in the V_H region, including at the amino- or carboxy-terminus or both ends of this region, provided that each CDR comprises zero changes or at most one, two, or three changes and provided a binding domain containing the modified V_H region can still specifically bind its target with an affinity similar to the wild type binding domain.

5 In further embodiments, a V_L region in a binding domain of the present disclosure is derived from or based on a V_L of a known monoclonal antibody and contains one or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10) insertions, one or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10) deletions, one or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10) amino acid substitutions (*e.g.*, conservative amino acid substitutions), or a combination of the
10 above-noted changes, when compared with the V_L of the known monoclonal antibody. An insertion, deletion or substitution may be anywhere in the V_L region, including at the amino- or carboxy-terminus or both ends of this region, provided that each CDR comprises zero changes or at most one, two, or three changes and provided a binding domain containing the modified V_L region can still specifically bind its target with an
15 affinity similar to the wild type binding domain.

The V_H and V_L domains may be arranged in either orientation (*i.e.*, from amino-terminus to carboxyl terminus, V_H - V_L or V_L - V_H) and may be joined by an amino acid sequence (*e.g.*, having a length of about five to about 35 amino acids) capable of providing a spacer function such that the two sub-binding domains can interact to form
20 a functional binding domain. In certain embodiments, a variable region linker that joins the V_H and V_L domains includes those belonging to the (Gly_nSer) family, such as $(Gly_3Ser)_n(Gly_4Ser)_1$ (SEQ ID NO:3), $(Gly_3Ser)_1(Gly_4Ser)_n$ (SEQ ID NO:10), $(Gly_3Ser)_n(Gly_4Ser)_n$ (SEQ ID NO:4), or $(Gly_4Ser)_n$ (SEQ ID NO: 5), wherein n is an integer of 1 to 5. In certain embodiments, the linker is $(Gly-Gly-Gly-Gly-Ser)_3$ (SEQ
25 ID NO.:12) or $(Gly-Gly-Gly-Ser)_4$ (SEQ ID NO.:13). In certain embodiments, these (Gly_nSer) -based linkers are used to link the V_H and V_L domains in a binding domain, and these linkers may also be used to link the binding domain to a tag cassette, or to link a tag cassette to a hydrophobic portion or intracellular component.

In some embodiments, a binding domain is a single chain T cell receptor
30 (scTCR) comprising $V_{\alpha\beta}$ and $C_{\alpha\beta}$ chains (*e.g.*, V_α - C_α , V_β - C_β , V_α - V_β) or comprising V_α - C_α , V_β - C_β , V_α - V_β pair specific for a target of interest (*e.g.*, peptide-MHC complex).

In certain embodiments, a binding domain comprises or is a sequence that is at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, at least 99.5%, or 100% identical to an amino acid sequence of a TCR V_α, V_β, C_α, or C_β, wherein each CDR comprises zero changes or at most one, two, or three changes, from a TCR or fragment or derivative thereof that specifically binds to a target of interest (*e.g.*, BCMA, γ-secretase complex component such as presenilin or nicastrin).

In certain embodiments, a binding domain V_α, V_β, C_α, or C_β region of the present disclosure can be derived from or based on a V_α, V_β, C_α, or C_β of a known TCR (*e.g.*, a high-affinity TCR) and contains one or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10) insertions, one or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10) deletions, one or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10) amino acid substitutions (*e.g.*, conservative amino acid substitutions or non-conservative amino acid substitutions), or a combination of the above-noted changes, when compared with the V_α, V_β, C_α, or C_β of a known TCR. An insertion, deletion or substitution may be anywhere in a V_α, V_β, C_α, or C_β region, including at the amino- or carboxy-terminus or both ends of these regions, provided that each CDR comprises zero changes or at most one, two, or three changes and provided a binding domain containing a modified V_α, V_β, C_α, or C_β region can still specifically bind its target with an affinity similar to wild type.

BCMA, γ-secretase, or both may be found on or in association with a cell of interest ("target cell"). Exemplary target cells include a cancer cell, a cell associated with an autoimmune disease or disorder or with an inflammatory disease or disorder, and an infectious organism or cell (*e.g.*, bacteria, virus, virus-infected cell).

Cytotoxic Conjugates

Antibody-drug conjugates are used for selectively delivering a cytotoxic element to a target cell, *e.g.*, a tumor or cancer cell. Antibody-drug conjugates and related techniques and chemistries are described in, for example, Nasiri *et al.*, *J. Cell. Physiol.* (2018), Hedrich *et al.*, *Clin. Pharmacokinet.* (2017), Drake and Rabuka, *BioDrugs* 31(6):521 (2017), Meyer *et al.*, *Bioconj. Chem.* 27(12):2791 (2016), Moek *et al.*, *J. Nucl. Med.* 58:83S (2017), Nareshkumar *et al.*, *Pharm. Res.* 32:3526 (2015), Parslow *et al.*, *Biomedicines* 4:14 (2016), and Green *et al.*, *Blood* 131:611 (2018), of

which the antibody formats, cytotoxic payloads, linker and conjugation chemistries, dosing regimens, treatment methods, pharmacokinetics, and principles of ADC design are incorporated herein by reference in their entirety.

In certain embodiments of the methods provided herein, a BCMA-specific
5 antibody or antigen-binding portion thereof, chimeric antigen receptor (CAR), or tagged
chimeric antigen receptor molecule (T-ChARM) is conjugated or otherwise coupled to
a cytotoxic agent, such as a chemotherapeutic agent. A chemotherapeutic agent
includes, but is not limited to, an inhibitor of chromatin function, a topoisomerase
inhibitor, a microtubule inhibiting drug, a DNA damaging agent, an antimetabolite
10 (such as folate antagonists, pyrimidine analogs, purine analogs, and sugar-modified
analogues), a DNA synthesis inhibitor, a DNA interactive agent (such as an intercalating
agent), and a DNA repair inhibitor. Illustrative chemotherapeutic agents include,
without limitation, the following groups: anti-metabolites/anti-cancer agents, such as
pyrimidine analogs (5-fluorouracil, floxuridine, capecitabine, gemcitabine and
15 cytarabine) and purine analogs, folate antagonists and related inhibitors
(mercaptopurine, thioguanine, pentostatin and 2- chlorodeoxyadenosine (cladribine));
antiproliferative/antimitotic agents including natural products such as vinca alkaloids
(vinblastine, vincristine, and vinorelbine), microtubule disruptors such as taxane
(paclitaxel, docetaxel), vincristin, vinblastin, nocodazole, epothilones and navelbine,
20 epidipodophyllotoxins (etoposide, teniposide), DNA damaging agents (actinomycin,
amsacrine, anthracyclines, bleomycin, busulfan, camptothecin, carboplatin,
chlorambucil, cisplatin, cyclophosphamide, Cytosan, dactinomycin, daunorubicin,
doxorubicin, epirubicin, hexamethylmelamineoxaliplatin, iphosphamide, melphalan,
merchlorehtamine, mitomycin, mitoxantrone, nitrosourea, plicamycin, procarbazine,
25 taxol, taxotere, temozolamide, teniposide, triethylenethiophosphoramide and etoposide
(VP 16)); antibiotics such as dactinomycin (actinomycin D), daunorubicin, doxorubicin
(adriamycin), idarubicin, anthracyclines, mitoxantrone, bleomycins, plicamycin
(mithramycin) and mitomycin; enzymes (L-asparaginase which systemically
metabolizes L-asparagine and deprives cells which do not have the capacity to
30 synthesize their own asparagine); antiplatelet agents; antiproliferative/antimitotic
alkylating agents such as nitrogen mustards (mechlorethamine, cyclophosphamide and

analogues, melphalan, chlorambucil), ethylenimines and methylmelamines (hexamethylmelamine and thiotepa), alkyl sulfonates -busulfan, nitrosoureas (carmustine (BCNU) and analogues, streptozocin), trazenes— dacarbazine (DTIC); antiproliferative/antimitotic antimetabolites such as folic acid analogues (methotrexate);
 5 platinum coordination complexes (cisplatin, carboplatin), procarbazine, hydroxyurea, mitotane, aminoglutethimide; hormones, hormone analogues (estrogen, tamoxifen, goserelin, bicalutamide, nilutamide) and aromatase inhibitors (letrozole, anastrozole); anticoagulants (heparin, synthetic heparin salts and other inhibitors of thrombin); fibrinolytic agents (such as tissue plasminogen activator, streptokinase and urokinase),
 10 aspirin, dipyridamole, ticlopidine, clopidogrel, abciximab; antimigratory agents; antisecretory agents (breveldin); immunosuppressives (cyclosporine, tacrolimus (FK-506), sirolimus (rapamycin), azathioprine, mycophenolate mofetil); anti-angiogenic compounds (TNP470, genistein) and growth factor inhibitors (vascular endothelial growth factor (VEGF) inhibitors, fibroblast growth factor (FGF) inhibitors); angiotensin
 15 receptor blocker; nitric oxide donors; anti-sense oligonucleotides; antibodies (trastuzumab, rituximab); chimeric antigen receptors; cell cycle inhibitors and differentiation inducers (tretinoin); mTOR inhibitors, topoisomerase inhibitors (doxorubicin (adriamycin), amsacrine, camptothecin, daunorubicin, dactinomycin, eniposide, epirubicin, etoposide, idarubicin, irinotecan (CPT-11) and mitoxantrone,
 20 topotecan, irinotecan), corticosteroids (cortisone, dexamethasone, hydrocortisone, methylprednisolone, prednisone, and prednisolone); growth factor signal transduction kinase inhibitors; mitochondrial dysfunction inducers, toxins such as Cholera toxin, ricin, Pseudomonas exotoxin, Bordetella pertussis adenylate cyclase toxin, or diphtheria toxin, and caspase activators; and chromatin disruptors.

25 An exemplary ADC that specifically binds to BCMA is J6M0-mcMMAF (GSK2857916), described in Tai *et al.*, *Blood* 123(20):3128-3138 (2014), which ADC and methods of using the same are hereby incorporated by reference in their entirety. Another BCMA-specific ADC (SG1-MMAF) was described by Ryan *et al.* (*Mol. Cancer Ther.*, 6(11):3009-3018, 2007), which ADC is incorporated herein by reference.

Host Cells and Nucleic Acids

In certain aspects, the present disclosure provides nucleic acid molecules that encode any one or more of the BCMA-specific binding proteins (including multispecific and bispecific binding proteins comprising at least one BCMA binding domain) or γ -secretase inhibitors described herein. Such nucleic acid molecules can be inserted into an appropriate vector (*e.g.*, viral vector or non-viral plasmid vector) for introduction in a host cell of interest (*e.g.*, T cell).

As used herein, the term "recombinant" or "non-natural" refers to an organism, microorganism, cell, nucleic acid molecule, or vector that includes at least one genetic alteration or has been modified by introduction of an exogenous nucleic acid molecule, wherein such alterations or modifications are introduced by genetic engineering. Genetic alterations include, for example, modifications introducing expressible nucleic acid molecules encoding proteins, fusion proteins or enzymes, or other nucleic acid molecule additions, deletions, substitutions or other functional disruption of a cell's genetic material. Additional modifications include, for example, non-coding regulatory regions in which the modifications alter expression of a gene or operon. In certain embodiments, a cell, such as a T cell, obtained from a subject may be converted into a non-natural or recombinant cell (*e.g.*, a non-natural or recombinant T cell) by introducing a nucleic acid that encodes a BCMA-specific binding protein or γ -secretase inhibitor of this disclosure (*e.g.*, BCMA-specific T-ChARM or CAR; or anti- γ -secretase) as described herein and whereby the cell expresses a cell surface located BCMA specific binding protein.

A vector that encodes a core virus is referred to herein as a "viral vector." There are a large number of available viral vectors suitable for use with the compositions of the instant disclosure, including those identified for human gene therapy applications (*see* Pfeifer and Verma, *Ann. Rev. Genomics Hum. Genet.* 2:177, 2001). Suitable viral vectors include vectors based on RNA viruses, such as retrovirus-derived vectors, *e.g.*, Moloney murine leukemia virus (MLV)-derived vectors, and include more complex retrovirus-derived vectors, *e.g.*, lentivirus-derived vectors. HIV-1-derived vectors belong to this category. Other examples include lentivirus vectors derived from HIV-2, FIV, equine infectious anemia virus, SIV, and Maedi-Visna virus (ovine lentivirus).

Methods of using retroviral and lentiviral viral vectors and packaging cells for transducing mammalian host cells with viral particles containing chimeric antigen receptor transgenes are known in the art and have been previously described, for example, in U.S. Patent 8,119,772; Walchli *et al.*, *PLoS One* 6:327930, 2011; Zhao *et al.*, *J.*

5 *Immunol.* 174:4415, 2005; Engels *et al.*, *Hum. Gene Ther.* 14:1155, 2003; Frecha *et al.*, *Mol. Ther.* 18:1748, 2010; Verhoeven *et al.*, *Methods Mol. Biol.* 506:97, 2009.

Retroviral and lentiviral vector constructs and expression systems are also commercially available.

In certain embodiments, a viral vector is used to introduce a non-endogenous
10 polynucleotide encoding a BCMA-specific binding protein or γ -secretase inhibitor. A viral vector may be a retroviral vector or a lentiviral vector. A viral vector may also include polynucleotides encoding a marker for transduction. Transduction markers for viral vectors are known in the art and include selection markers, which may confer drug resistance, or detectable markers, such as fluorescent markers or cell surface proteins
15 that can be detected by methods such as flow cytometry. In particular embodiments, a viral vector further comprises a gene marker for transduction comprising green fluorescent protein, an extracellular domain of human CD2, or a truncated human EGFR (huEGFRt; *see* Wang *et al.*, *Blood* 118:1255, 2011). When a viral vector genome comprises a plurality of nucleic acid sequences to be expressed in a host cell as
20 separate transcripts, the viral vector may also comprise additional sequences between the two (or more) transcripts allowing bicistronic or multicistronic expression. Examples of such sequences used in viral vectors include internal ribosome entry sites (IRES), furin cleavage sites, viral 2A peptide (*e.g.*, P2A, T2A, E2A, F2A), or any combination thereof.

25 Other vectors also can be used for polynucleotide delivery including DNA viral vectors, including, for example adenovirus-based vectors and adeno-associated virus (AAV)-based vectors; vectors derived from herpes simplex viruses (HSVs), including amplicon vectors, replication-defective HSV and attenuated HSV (Krisky *et al.*, *Gene Ther.* 5: 1517, 1998).

30 Other vectors recently developed for gene therapy uses can also be used with the compositions and methods of this disclosure. Such vectors include those derived from

baculoviruses and α -viruses. (Jolly, D J. 1999. Emerging Viral Vectors. p. 209-40 in Friedmann T. ed. The Development of Human Gene Therapy. New York: Cold Spring Harbor Lab), or plasmid vectors (such as sleeping beauty or other transposon vectors). In some embodiments, a viral or plasmid vector further comprises a gene marker for
5 transduction (e.g. green fluorescent protein, huEGFRt).

In certain embodiments, hematopoietic progenitor cells or embryonic stem cells are modified to comprise a non-endogenous polynucleotide that encodes a BCMA-specific binding protein (e.g., BCMA-specific T-ChARM or CAR) of this disclosure. Hematopoietic progenitor cells may comprise thymocyte progenitor cells or
10 induced pluripotent stem cells, which may be derived or originate from fetal liver tissue, bone marrow, cord blood, or peripheral blood. The hematopoietic progenitor cells may be from human, mouse, rat, or other mammals. In particular embodiments, CD24^{lo} Lin⁻ CD117⁺ thymocyte progenitor cells are used.

In certain embodiments, culture conditions entail culturing hematopoietic
15 progenitor cells expressing fusion proteins of this disclosure for a sufficient time to induce proliferation or differentiation. The cells are maintained in culture generally for about 3 days to about 5 days, or about 4 to about 10 days, or about 5 to about 20 days. It will be appreciated that the cells may be maintained for an appropriate amount of time required to achieve a desired result, i.e., a desired cellular composition or level of
20 proliferation. For example, to generate a cellular composition comprising primarily immature and inactivated T cells, cells may be maintained in culture for about 5 to about 20 days. Cells may be maintained in culture for about 20 to about 30 days to generate a cellular composition comprising primarily mature T cells. Non-adherent cells may also be collected from culture at various time points, such as from about
25 several days to about 25 days. In certain embodiments, hematopoietic stem cells are co-cultured on stromal cells lines (U.S. Patent No. 7,575,925; Schmitt *et al.*, *Nat. Immunol.* 5:410, 2004; Schmitt *et al.*, *Immunity* 17:749, 2002).

One or more cytokines that promote commitment or differentiation of hematopoietic progenitor cells may be added to the culture. The cytokines may be
30 human or non-human. Representative examples of cytokines that may be used include all members of the FGF family, including FGF-4 and FGF-2; Flt-3-ligand, stem cell

factor (SCF), thrombopoietin (TPO), and IL-7. Cytokines may be used in combination with a glycosaminoglycan, such as heparin sulfate.

In some embodiments, host cells capable of expressing a BCMA-specific binding protein of this disclosure on the cell surface are T cells, including primary cells or cell lines derived from human, mouse, rat, or other mammals. If obtained from a mammal, a T cell can be obtained from numerous sources, including blood, bone marrow, lymph node, thymus, or other tissues or fluids. A T cell or subpopulations thereof (*e.g.*, naïve, central memory, effector memory) may be enriched or purified. T cell lines are well known in the art, some of which are described in Sandberg *et al.*, *Leukemia* 21:230, 2000. In certain embodiments, T cells that lack endogenous expression of TCR α and β chains are used. Such T cells may naturally lack endogenous expression of TCR α and β chains or may have been modified to block expression (*e.g.*, T cells from a transgenic mouse that does not express TCR α and β chains or cells that have been manipulated to inhibit expression of TCR α and β chains) or to knockout TCR α chain, TCR β chain, or both genes. In certain embodiments, cells capable of expressing a BCMA-specific binding protein of this disclosure on the cell surface are not T cells or cells of a T cell lineage, but cells that are progenitor cells, stem cells or cells that have been modified to express cell surface anti-CD3.

In any of the embodiments provided herein, a host cell can be a "universal donor" cell that is modified to reduce or eliminate expression of one or more endogenous genes involved in an immune response. For example, a T cell may be modified to reduce or eliminate expression of one or more polypeptides selected from PD-1, LAG-3, CTLA4, TIGIT, TIM3, an HLA complex component, or a TCR or TCR complex component. Without wishing to be bound by theory, certain endogenously expressed immune cell proteins may be recognized as foreign by an allogeneic host that receives the modified immune cells, which may result in elimination of the modified immune cells (*e.g.*, an HLA allele), or may downregulate the immune activity of the modified immune cells (*e.g.*, PD-1, LAG-3, CTLA4, TIGIT), or may interfere with the binding activity of a heterologously expressed binding protein of the present disclosure (*e.g.*, an endogenous TCR that binds to a non-tumor-associated antigen and interferes with the antigen-specific receptor of the modified immune cell specifically binding to

the tumor-associated antigen). Accordingly, decreasing or eliminating expression or activity of such endogenous genes or proteins can improve the activity, tolerance, and persistence of the modified immune cells in an allogeneic host setting, and can allow universal administration of the cells (*e.g.*, to any recipient regardless of HLA type).

5 In certain embodiments, a host cell (*e.g.*, modified immune cell) of this disclosure comprises a chromosomal gene knockout of one or more genes encoding a PD-1, LAG-3, CTLA4, TIM3, TIGIT, an HLA complex component (*e.g.*, a gene that encodes an α 1 macroglobulin, an α 2 macroglobulin, an α 3 macroglobulin, a β 1 microglobulin, or a β 2 microglobulin), a TCR component (*e.g.*, a gene that encodes a
10 TCR variable region or a TCR constant region) (*see, e.g.*, Torikai *et al.*, *Nature Sci. Rep.* 6:21757 (2016); Torikai *et al.*, *Blood* 119(24):5697 (2012); and Torikai *et al.*, *Blood* 122(8):1341 (2013) the gene editing techniques, compositions, and adoptive cell therapies of which are herein incorporated by reference in their entirety). For example, in some embodiments, a chromosomal gene knockout is produced using a
15 CRISPR/Cas9 system, and may involve transfection of the modified immune cell with a lentivirus (*e.g.*, pLentiCRISPRv2; Torikai *et al.*, *Blood* (2016)) expressing a CRISPR/Cas9 system targeting PD-1, LAG-3, CTLA4, an HLA component, or a TCR component, or any combination thereof. Primers useful for designing a lentivirus that expresses a CRISPR/Cas9 system for inhibiting an endogenously expressed immune
20 cell protein include for example, primer pairs comprising forward and reverse primers having the nucleotide sequences set forth in SEQ ID NOS:22 and 23, 24 and 25, 26 and 27, and 28 and 29.

In certain embodiments, a host T cell transfected to express a BCMA-specific binding protein (*e.g.*, T-CHARM, CAR, multispecific binding proteins comprising at
25 least one BCMA binding domain, bispecific binding proteins comprising at least one BCMA binding domain) of this disclosure is a functional T cell, such as a virus-specific T cell, a tumor antigen specific cytotoxic T cell, a naïve T cell, a memory stem T cell, a central or effector memory T cell, or a CD4⁺ CD25⁺ regulatory T cell.

One or more growth factor cytokines that promote proliferation of T cells
30 expressing a BCMA-specific binding protein of this disclosure may be added to the culture. The cytokines may be human or non-human. Exemplary growth factor

cytokines that may be used promote T cell proliferation include IL-2, IL-15, IL-21, or the like.

Uses

Diseases that may be treated with a BCMA-targeted immunotherapy or cells
5 expressing a BCMA-specific binding protein in combination with a γ -secretase inhibitor (GSI) as described in the present disclosure include cancer (*e.g.*, cancers that express BCMA), immune diseases (*e.g.*, autoimmune), or aging-related diseases (*e.g.*, senescence). Adoptive immune and gene therapy are promising treatments for various types of cancer (Morgan *et al.*, *Science* 314:126, 2006; Schmitt *et al.*, *Hum. Gene Ther.*
10 20:1240, 2009; June, *J. Clin. Invest.* 117:1466, 2007) and infectious disease (Kitchen *et al.*, *PLoS One* 4:38208, 2009; Rossi *et al.*, *Nat. Biotechnol.* 25:1444, 2007; Zhang *et al.*, *PLoS Pathog.* 6:e1001018, 2010; Luo *et al.*, *J. Mol. Med.* 89:903, 2011).

Cancers that are amenable to the compositions and methods disclosed herein are those that express, or are capable of expressing, BCMA on their cell surface.
15 Exemplary types of cancer that may be treated include myelomas (such as multiple myeloma), leukemias (such as plasma cell leukemia), lymphomas (such as lymphoplasmacytic lymphoma), plasmacytomas, Waldenström's macroglobulinemia. Other cancers that may express BCMA include adenocarcinoma of the breast and bronchogenic carcinoma of the lung. In certain embodiments, proliferative disorders
20 amenable to a combination therapy of a BCMA-specific binding protein and a GSI are certain types of B-cell cancer, including plasma cell disorders (such as, for example, multiple myeloma).

Inflammatory and autoimmune diseases include arthritis, rheumatoid arthritis, juvenile rheumatoid arthritis, osteoarthritis, polychondritis, psoriatic arthritis, psoriasis,
25 dermatitis, polymyositis/dermatomyositis, inclusion body myositis, inflammatory myositis, toxic epidermal necrolysis, systemic scleroderma and sclerosis, CREST syndrome, inflammatory bowel disease, Crohn's disease, ulcerative colitis, respiratory distress syndrome, adult respiratory distress syndrome (ARDS), meningitis, encephalitis, uveitis, colitis, glomerulonephritis, allergic conditions, eczema, asthma,
30 conditions involving infiltration of T cells and chronic inflammatory responses, atherosclerosis, autoimmune myocarditis, leukocyte adhesion deficiency, systemic

lupus erythematosus (SLE), subacute cutaneous lupus erythematosus, discoid lupus, lupus myelitis, lupus cerebritis, juvenile onset diabetes, multiple sclerosis, allergic encephalomyelitis, neuromyelitis optica, rheumatic fever, Sydenham's chorea, immune responses associated with acute and delayed hypersensitivity mediated by cytokines and
 5 T-lymphocytes, tuberculosis, sarcoidosis, granulomatosis including Wegener's granulomatosis and Churg-Strauss disease, agranulocytosis, vasculitis (including hypersensitivity vasculitis/angiitis, ANCA and rheumatoid vasculitis), aplastic anemia, Diamond Blackfan anemia, immune hemolytic anemia including autoimmune hemolytic anemia (AIHA), pernicious anemia, pure red cell aplasia (PRCA), Factor
 10 VIII deficiency, hemophilia A, autoimmune neutropenia, pancytopenia, leukopenia, diseases involving leukocyte diapedesis, central nervous system (CNS) inflammatory disorders, multiple organ injury syndrome, myasthenia gravis, antigen-antibody complex mediated diseases, anti-glomerular basement membrane disease, anti-phospholipid antibody syndrome, allergic neuritis, Behcet disease, Castleman's
 15 syndrome, Goodpasture's syndrome, Lambert-Eaton Myasthenic Syndrome, Reynaud's syndrome, Sjorgen's syndrome, Stevens-Johnson syndrome, solid organ transplant rejection, graft versus host disease (GVHD), bullous pemphigoid, pemphigus, autoimmune polyendocrinopathies, seronegative spondyloarthropathies, Reiter's disease, stiff-man syndrome, giant cell arteritis, immune complex nephritis, IgA
 20 nephropathy, IgM polyneuropathies or IgM mediated neuropathy, idiopathic thrombocytopenic purpura (ITP), thrombotic thrombocytopenic purpura (TTP), Henoch-Schonlein purpura, autoimmune thrombocytopenia, autoimmune disease of the testis and ovary including autoimmune orchitis and oophoritis, primary hypothyroidism; autoimmune endocrine diseases including autoimmune thyroiditis, chronic thyroiditis
 25 (Hashimoto's Thyroiditis), subacute thyroiditis, idiopathic hypothyroidism, Addison's disease, Grave's disease, autoimmune polyglandular syndromes (or polyglandular endocrinopathy syndromes), Type I diabetes also referred to as insulin-dependent diabetes mellitus (IDDM) and Sheehan's syndrome; autoimmune hepatitis, lymphoid interstitial pneumonitis (HIV), bronchiolitis obliterans (non-transplant), non-specific
 30 interstitial pneumonia (NSIP), Guillain-Barré Syndrome, large vessel vasculitis (including polymyalgia rheumatica and giant cell (Takayasu's) arteritis), medium vessel

vasculitis (including Kawasaki's disease and polyarteritis nodosa), polyarteritis nodosa (PAN) ankylosing spondylitis, Berger's disease (IgA nephropathy), rapidly progressive glomerulonephritis, primary biliary cirrhosis, Celiac sprue (gluten enteropathy), cryoglobulinemia, cryoglobulinemia associated with hepatitis, amyotrophic lateral sclerosis (ALS), coronary artery disease, familial Mediterranean fever, microscopic polyangiitis, Cogan's syndrome, Whiskott-Aldrich syndrome and thromboangiitis obliterans.

In particular embodiments, a method of treating a subject with the BCMA-binding protein in combination with a γ -secretase inhibitor as disclosed herein include treating multiple myeloma, plasmacytoma, plasma cell leukemia, Waldenström's macroglobulinemia, B cell lymphoma, and lymphoplasmacytic lymphoma.

A BCMA-specific binding protein, such as a CAR or T-ChARM, of this disclosure may be administered to a subject in cell-bound form (*e.g.*, gene therapy of target cell population (mature T cells (*e.g.*, CD8⁺ or CD4⁺ T cells) or other cells of T cell lineage)). In a particular embodiment, cells of T cell lineage expressing BCMA-specific binding protein (*e.g.*, BCMA specific CAR or T-ChARM) administered to a subject are syngeneic, allogeneic, or autologous cells.

A method of the present disclosure includes steps of administering a BCMA-specific binding molecule expressed on an immune cell surface (*e.g.*, T cell) and administering a γ -secretase inhibitor (GSI). In certain embodiments, the combination may be administered concurrently, together in the same pharmaceutically acceptable carrier, or in separate formulations (but concurrently). Concurrent administration means the each component is administered at the same time or within 8-12 hours of each other. Administration of a second component more than 12 hours after the first component will be considered a sequential administration. In other embodiments, a BCMA-specific immunotherapeutic and GSI can be administered sequentially (*e.g.*, one, two, three, four, five, six, seven, eight, or nine days apart; one, two, three, or four weeks apart; one, two, three, four, five, six, seven, eight, or nine weeks apart; or one, two, three, four, five, six, or more years apart; or the like), in any order and in any combination. In particular embodiments, when administered

sequentially, the GSI is administered first and the BCMA-targeted immunotherapy (soluble or cell form) is administered second. In other embodiments, a BCMA-targeted immunotherapy (soluble or cell form) is administered first and the GSI is administered second. In particular embodiments, a BCMA-targeted immunotherapy comprising a modified immune cell that specifically binds to BCMA (*e.g.*, a BCMA-specific T cell, such as a CAR-T, T-ChARM, bispecific, multispecific T cell) is administered first and a GSI is administered second (*e.g.*, within hours, days, weeks, months, or years of the BCMA immunotherapy). In further embodiments, a BCMA-targeted immunotherapy comprising a modified immune cell comprising a binding protein that specifically binds to BCMA (*e.g.*, a BCMA-specific T cell, such as a CAR-T, T-ChARM, bispecific, multispecific T cell) is (a) administered to a subject; (b) after a period of time (*e.g.*, about 5 days to about one week, about one week to about two weeks, about two weeks to about three weeks, about one week to about one month, about one month to about six months, about three months to about one year, or the longer as needed), the subject or a tissue from the subject is probed or examined for the presence or persistence of the previously administered modified immune cell comprising a binding protein that specifically binds to BCMA, and (c) administering a GSI second after the presence or persistence of the BCMA immunotherapy is detected.

In some embodiments, a GSI is administered to a subject at least once before, simultaneous with, and at least once (*e.g.*, at least two, three, or four times) after, the BCMA-targeted immunotherapy. In particular embodiments, a combination therapy of this disclosure comprises a BCMA-targeted immunotherapy and administration of from about 0.01 μM GSI to about 5 μM GSI, from about 0.03 μM GSI to about 1.5 μM GSI, from about 0.05 μM GSI to about 2.5 μM GSI, or from about 0.1 μM GSI to about 1.0 μM GSI. In certain embodiments, a combination therapy comprising a modified immune cell that specifically binds BCMA and a GSI comprises a lower amount of the immune cell that specifically binds BCMA, the GSI, or both as compared to administration of these therapies individually.

For example, by way of background and not wishing to be bound by theory, during adoptive immunotherapy engraftment of administered cells expressing the BCMA-specific binding protein may be facilitated by prior immunosuppressive

conditioning with one or more chemotherapeutic agents or treatments. In any of the embodiments described herein, the method further comprises pre-conditioning a subject concurrent with or prior to administering a BCMA-targeted immunotherapy or an immune cell expressing a BCMA-specific binding protein and a GSI. In particular
5 embodiments, a pre-conditioning regimen is an immunosuppressive conditioning comprising depleting endogenous lymphocytes (also referred to as lymphodepletion, which may be non-myeloablative or myeloablative). Exemplary lymphodepletion may be achieved with cyclophosphamide alone, cyclophosphamide in combination with fludarabine, by use of other agents or treatments that are cytotoxic to lymphocytes (*e.g.*,
10 total body irradiation), or any combination thereof.

The binding molecule, inhibitor or combination compositions may be administered orally, topically, transdermally, parenterally, by inhalation spray, vaginally, rectally, or by intracranial injection, or any combination thereof. When administered separately, a BCMA-targeted immunotherapy and a GSI may be
15 administered by the same route or by different routes. For example, in certain embodiments, the BCMA-targeted immunotherapy is administered parenterally and the GSI is administered orally, which can be concurrently or sequentially. The term “parenteral,” as used herein, includes subcutaneous injections, intravenous, intramuscular, intracisternal injection, or infusion techniques. Administration by
20 intravenous, intradermal, intramuscular, intramammary, intraperitoneal, intrathecal, retrobulbar, intrapulmonary injection and or surgical implantation at a particular site is contemplated as well. Generally, compositions are essentially free of pyrogens, as well as other impurities that could be harmful to the recipient. Injection or infusion, especially intravenous, is preferred for administering a BCMA-targeted
25 immunotherapy.

In other embodiments, BCMA-specific binding protein, GSI or both may be administered to a subject in soluble form (*e.g.*, antibody). Soluble TCRs are also known in the art (*see, e.g.*, Molloy *et al.*, *Curr. Opin. Pharmacol.* 5:438, 2005; U.S. Patent No. 6,759,243).

30 Pharmaceutical compositions including a combination therapy of a BCMA-targeted immunotherapy and a γ -secretase inhibitor of this disclosure may be

administered in a manner appropriate to the disease or condition to be treated (or prevented) as determined by persons skilled in the medical art. An appropriate dose, suitable duration, and frequency of administration of the compositions will be determined by such factors as the condition of the patient, size, type and severity of the disease, particular form of the active ingredient, and the method of administration. The present disclosure provides pharmaceutical compositions comprising cells expressing a BCMA-specific binding protein, such as a CAR or T-ChARM, and a pharmaceutically acceptable carrier, diluents, or excipient. Suitable excipients include water, saline, dextrose, glycerol, or the like and combinations thereof.

An advantage of the instant disclosure is that BCMA-specific binding protein, such as a CAR or T-ChARM, expressing cells administered to a patient can be depleted using the cognate binding partner to a tag cassette. In certain embodiments, the present disclosure provides a method for depleting a T cell expressing a BCMA-specific T-ChARM by using an antibody specific for the tag cassette, using a cognate binding partner specific for the tag cassette, or by using a second T cell expressing a CAR and having specificity for the tag cassette. In certain embodiments, a tag cassette allows for immunodepletion of a T cell expressing a BCMA-specific T-ChARM of this disclosure. Elimination of engineered T cells may be accomplished using depletion agents specific for a tag cassette. For example, if a Strep tag is used, then an anti-Strep tag antibody, anti-Strep tag scFv, or Streptactin each fused to or conjugated to a cell-toxic reagent (such as a toxin, radiometal) may be used, or an anti-Strep tag /anti-CD3 bispecific scFv, or an anti-Strep tag CAR T cell may be used.

In another aspect, the present disclosure provides a method for selectively promoting proliferation of a recombinant T cell expressing a BCMA-specific T-ChARM of this disclosure. In certain embodiments, the method comprises selective *ex vivo* proliferation of T cells expressing a BCMA-specific T-ChARM using a tag binding partner, such as an antibody. In further embodiments, the method comprises expanding functional T cells (*e.g.*, virus-specific, TAA (tumor-associated antigen) specific CTL, or specific T cell subsets, such as naïve T cells, memory stem T cells, central or effector memory T cells, CD4⁺ CD25⁺ regulatory T cells) with a tag binding

partner, such as an antibody, which may optionally be done in the presence of a costimulatory molecule binding partner (such as an anti-CD27 or antiCD28 antibody).

In still further embodiments, a BCMA-specific T-ChARM allows for selective promotion of T cell proliferation *in vivo* when expressing a BCMA-specific T-ChARM of this disclosure. In certain embodiments, a T cell expressing a CAR comprising a tag
5 cassette allows for expansion of the CAR T cells *in vivo* when contacting cells expressing a ligand (*e.g.*, including T cell suppressor cell ligands PD-L1, PD-L2). Such expanded T cells are useful in the disease treatment methods described herein. In certain embodiments, proliferation or expansion of cells expressing BCMA-specific
10 T-ChARM as disclosed herein is induced *in vivo*, which may be induced with a tag cassette binding partner (such as an anti-tag antibody) and optionally a costimulatory molecule binding partner (such as an anti-CD27 or anti-CD28 antibody).

In certain further embodiments, cells expressing BCMA-specific T-ChARM as disclosed herein are activated *in vivo*, such as at the site of a tumor. For example, a
15 composition (*e.g.*, alginate, basement membrane matrix (Matrigel®), biopolymer, or other matrix) or a carrier (*e.g.*, microbead, nanoparticle, or other solid surface) comprising a tag cassette binding partner (such as an anti-tag antibody) and a costimulatory molecule binding partner (such as an anti-CD27 or antiCD28 antibody) may be used to locally activate at the site of a tumor (*e.g.*, a solid tumor) a T cell
20 expressing a BCMA-specific T-ChARM as disclosed herein.

In certain embodiments, recombinant cells expressing a BCMA-specific T-ChARM may be detected or tracked *in vivo* by using antibodies that bind with specificity to a tag cassette (*e.g.*, anti-Tag antibodies), or by other cognate binding proteins that specifically bind the tag cassette sequence (*e.g.*, Streptactin binding to
25 Strep tag), which binding partners for the tag cassette are conjugated to a fluorescent dye, radio-tracer, iron-oxide nanoparticle or other imaging agent known in the art for detection by X-ray, CT-scan, MRI-scan, PET-scan, ultrasound, flow-cytometry, near infrared imaging systems, or other imaging modalities (*see, e.g.*, Yu *et al.*, *Theranostics* 2:3, 2012).

In further embodiments, cells expressing BCMA-specific T-ChARM of the instant disclosure may be used in diagnostic methods or imaging methods, including methods used in relation to the indications or conditions identified herein.

In some embodiments, a method comprises administering a GSI in combination
5 (e.g., concurrently, separately, or sequentially) with a BCMA-specific binding protein that may comprise, for example, a BCMA-specific antibody or antigen-binding portion thereof, such as antibody-drug conjugate, or a bispecific or a multispecific binding protein, such as those useful for pre-targeted radiation immunotherapy (*see, e.g., Green et al., Blood 131:611 (2018)*).

10

EXAMPLES

EXAMPLE 1

DESIGN AND TESTING OF BCMA-SPECIFIC CHIMERIC ANTIGEN RECEPTORS

Anti-BCMA CARs were prepared to examine their usefulness for immunotherapy of multiple myeloma and other disorders. Anti-BCMA CARs were
15 constructed with scFvs comprised of the V_H and V_L regions from C115 D5.3 ("C11") antibody and A7D12.2 ("A7") antibody, which included an IgG4 hinge region (spacer), a CD28 transmembrane domain, a 4-1BB costimulatory domain, and a CD3 ζ effector domain. The scFvs were produced in both the "HL" and "LH" orientations (*see* Figure 1A). Human T cells were transduced with expression constructs encoding the anti-
20 BCMA CARs and examined for functional characteristics. As shown in Figure 1B, T cells expressing the C11 CARs and A7 CARs proliferated when co-cultured with BCMA-expressing target cells, although the C11 CARs appeared to proliferate slightly more robustly than the A7 CAR-expressing T cells. The C11 CAR-T cells also produced more cytokines in response to target cell lines (either antigen-transfected
25 K562 cells or BCMA-expressing MM cell lines; Figure 1C) and had greater specific killing activity against target cells as compared to the A7 LH CAR-T cells (Figure 1D).

Additional A7 and C11 CARs were generated having different intracellular components, including CARs containing a 4-1BB costimulatory domain (Figure 1E,

upper illustration) and a CD28 costimulatory domain (Figure 1E, lower illustration). The length of the extracellular spacer domain was also varied to improve interaction between the CAR expressing T cell and the BCMA⁺ target cell, including short (*e.g.*, 12 amino acids, 48 amino acids, and 66 amino acids in length), intermediate (*e.g.*, 157 amino acids in length), and long (*e.g.*, 228 amino acids in length) spacers. The 48 amino acid and 157 amino acid spacer CARs each included two (2) Strep-Tag cassettes, while the 66 amino acid spacer included three (3) Strep-Tag cassettes (*see* Figure 1F). Tagged chimeric antigen receptors such as the Strep-Tag cassette-containing CARs shown in Figures 1F and 1G are referred to herein as T-ChARMs.

Human T cells were transduced to express the T-ChARMs (*see* Example 2) and analyzed for functionality. The length of the spacer had an effect on the T-ChARM constructs comprising a scFv derived from anti-BCMA antibody C11 D5.3, wherein intermediate (about 65 amino acids; *see* C11 3ST, C11 2STint and C11Lo of Figure 1H) to long (about 200 amino acids) spacers worked best for this binding domain. C113ST_4-1BB and C113ST_CD28 were selected as best-performing constructs (Figures 1H-1J) and compared to a previously disclosed anti-BCMA CAR ("BCMA-2"; Carpenter *et al. Clin. Cancer Res.* 19:2048, 2013). Expression of EGFRt on primary T cells was similar for each T-ChARM / CAR construct and surface expression of the T-ChARM was confirmed for those containing STII sequences as shown by staining with anti-STII monoclonal antibody (Figure 1K). T cells expressing either C113ST T-ChARM produced more cytokines (Figure 1L, 1M) and proliferated more robustly (Figure 1N) than T cells expressing BCMA-2 when co-cultured with target cells. It was also determined that T cells expressing either C113ST T-ChARM can recognize and lyse CD138⁺ patient MM cells (data not shown). The T cells expressing either C113ST T-ChARM or the BCMA-2 CAR did not have cytolytic activity against non-antigen-expressing K562 cells (Figure 1O), but effectively lysed K562 cells transduced to express the BCMA antigen (Figure 1P), demonstrating that the engineered T cells specifically recognize antigen.

EXAMPLE 2**PRODUCTION OF RECOMBINANT T CELLS AND
EXPRESSION OF BCMA-SPECIFIC T-CHARMS**

CD8⁺ and CD4⁺ T cells were isolated from PBMC of normal human donors
5 using CD8⁺/CD4⁺ T Cell Isolation Kit (Miltenyi Biotec), activated with anti-CD3/CD28
beads (Life Technologies) according to the manufacturer's instructions, and transduced
with a CAR-encoding lentiviral (epHIB7) supernatant generated by transient
transfection of HEK293T cells using PsPAX2 and pMD2G packaging plasmids (MOI =
3) supplemented with 0.8 µg/mL polybrene (Millipore, Bedford, MA) on day 3 after
10 activation by centrifugation at 2,100 rpm for 45 min at 32°C. T cells were expanded in
RPMI, 10% human serum, 2 mM L-glutamine and 1% penicillin-streptomycin (CTL
medium), supplemented with recombinant human (rh) IL-2 to a final concentration of
50 U/mL every 48 hours. After expansion, an aliquot of each transduced T cell line was
stained with biotin-conjugated anti-EGFR antibody and streptavidin-PE (Miltenyi,
15 Auburn, CA). The tEGFR⁺ T cells were isolated by sorting on a FACS-Aria cell sorter
(Becton Dickinson). The tEGFR⁺ T cell subset was then stimulated with irradiated
(8,000 rad) CD19⁺ B-LCL at a T cell:LCL ratio of 1:7, and expanded for 8 days in CTL
medium with addition of 50 U/mL rh IL-2 every 48 hours or using a rapid expansion
protocol for cells expressing T-ChARMS (Riddell and Greenberg, *J. Immunol. Methods*
20 *128:189, 1990*), wherein the T-ChARM comprises a scFv derived from anti-BCMA
antibody A7 or C11 D5.3.

The following conjugated antibodies were used for flow cytometric phenotyping
and analysis: CD4, CD8, CD25, CD137, CD45, Annexin V, CD62L, CD27, CD28 (BD
Biosciences), anti-Streptag II antibody (Genscript), EGFR antibody (ImClone Systems
25 Incorporated, Branchburg, NJ); streptavidin-PE (BD Biosciences, San Jose, CA).
Staining with propidium iodide (PI, BD Biosciences) was performed for live/dead cell
discrimination as directed by the manufacturer. Flow analyses were done on a FACS
Canto II, sort-purifications on a FACS AriaII (Becton Dickinson, Franklin Lakes, NJ)
and data analyzed using FlowJo software (Treestar, Ashland, OR).

30 To examine cell surface expression of BCMA-specific T-ChARM, transduced
T cells were sorted for EGFRt expression and evaluated by staining with fluorochrome

labeled anti-Streptag mAb. The mean fluorescence intensity (MFI) of EGFR staining was similar on T cells transduced with each of the BCMA-specific T-ChARM and the CD19-Short CAR. The introduction of a tag into a CAR to produce a T-ChARM did not interfere with transgene expression (data not shown). An anti-StrepTag mAb
5 specifically stained T cells transduced with the various BCMA-specific T-ChARMs, independent of the position or number of tag sequences in each T-ChARM.

EXAMPLE 3

SOLUBLE BCMA (SBCMA) INHIBITS BCMA-SPECIFIC CAR T CELL ACTIVITY

The potential effectiveness of BCMA-targeted T cell therapies against most myelomas
10 was examined. Because shedding of surface BCMA by myeloma cells may hinder T cell recognition, soluble BCMA levels were measured in the supernatants during culture of a myeloma cell line. U266 myeloma cells were washed and plated in culture media for 1, 3, 5 and 24 hours. The media supernatant was harvested and assayed for sBCMA by ELISA. The data shows a time-dependent increase in sBCMA levels in the
15 supernatant (Figure 2A). Next, the extent of BCMA expression on primary patient multiple myeloma (MM) cells and cytokine production by BCMA-specific T-ChARM T cells contacted with patient MM cells was examined. Figure 2B shows surface BCMA expression by a reference cell line (RPMI, left panel) and samples from primary patient MM cells with differing levels of BCMA expression. The pie chart of Figure
20 2C shows that BCMA expression on the surface of MM cells from nineteen (19) different patients was found to be positive for most patients, though there were some patient samples that showed intermediate to low/no surface BCMA expression (~25%). Furthermore, Figure 2D shows that the T-ChARM T cells produced more IFN- γ when cultured with patient MM cells expressing high levels of surface BCMA, as compared
25 to culture with low-BCMA patient MM cells.

Myeloma cells frequently express PD-L1, which is believed to inhibit T cell function by binding to PD-1 on T cells (Freeman *et al. J. Exp. Med.* 192(7):1027 (2000)). The possibility that PD-L1 affected BCMA-specific T-ChARM T cells was also investigated. Flow cytometry of patient samples showed that PD-L1 was also

expressed on 79% of samples, with 47% PD-L1^{hi}, 32% PD-L1^{int}, and 21% PD-L1^{low/neg} (Figures 2E and 2F). No correlation between BCMA and PD-L1 expression was observed. Figure 2G shows that T-ChARM T cells produced more IFN- γ when cultured with patient MM cells expressing low levels of PD-L1 versus high levels of PD-L1, although this trend was not statistically significant.

The effect of sBCMA was examined as well by adding sBCMA to co-cultures of T cells expressing a BCMA-specific T-ChARM and K562 cells transduced with a polynucleotide encoding full-length BCMA (K562/BCMA⁺). A dose-dependent inhibition of BCMA-specific T-ChARM T cell effector function was detected as a measure of IFN- γ release into the media supernatant upon administration of exogenous sBCMA (Figures 2H, 2I). Additionally, MM cell lines showed substantial increases in culture supernatant sBCMA within 24 hours (Figure 2J).

To examine whether high levels of sBCMA might inhibit the activity of BCMA-specific T-ChARM T cells, patient bone marrow (BM) sera were examined for sBCMA levels. Patient BM was found to have high levels of sBCMA, which roughly correlated with disease burden as determined by the percentage of CD138⁺ cells present (Figure 2K). To test whether sBCMA would bind to the T-ChARM T cells, the T cells were incubated with increasing levels of recombinant BCMA and then stained using BCMA-Fc conjugated to APC. As shown in Figure 2L, the staining decreased with higher levels of recombinant BCMA, which indicates that sBCMA could potentially have a detrimental effect BCMA-specific T-ChARM T cell function.

To confirm T-ChARM expression, C113ST T-ChARM T cells and control anti-CD19 CAR T cells ("FMC63") were incubated with Fc-BCMA and stained for EGFRt (CAR / T-ChARM transduction marker) and CD4, showing that T-ChARMs were expressed by the cells (Figure 2M). Staining of EGFRt and other T cell surface molecules was unaffected (data not shown).

To determine whether shedding and binding of sBCMA to the T-ChARMs inhibits T cell function, BCMA-Fc was added to T cell/target cell line co-cultures, resulting in much-reduced recognition (IFN- γ production and CD4 expression) by the BCMA-targeting T-ChARM T cells, but not by control anti-CD19 CAR T cells (cultured with K562 cells expressing the CD19 antigen (Figure 2N). The effect on the

BCMA T-ChARM T cells was dose-dependent (Figures 2O, 2P). Addition of recombinant BCMA not fused to the Fc also inhibited IFN- γ production in a dose-dependent fashion, but did not inhibit lysis of K562 BCMA⁺ cells by T-ChARM T cells (Figure 2Q), which may be due to high antigen density on the surface of the target cells.

5

EXAMPLE 4

EFFECT OF γ -SECRETASE INHIBITOR ON MM BCMA LEVELS

To examine the effect of γ -secretase inhibition on BCMA levels on myeloma cells, γ -secretase inhibitor (GSI) RO4929097 was used. BCMA is rapidly upregulated on various myeloma cell lines when incubated with γ -secretase inhibitor (GSI) RO4929097 (concentration used ranged from 0.001 μ M to 1.0 μ M) (Figures 3A-3D).

To examine the effect of GSI on surface BCMA expression kinetics, U266 myeloma cells were incubated for 1, 3, 5 and 24 hours in the presence of various concentrations of GSI RO429097 (0.01 μ M, 0.1 μ M and 1.0 μ M) and evaluated for cell surface BCMA expression by flow cytometry. BCMA expression increased in a dose-dependent manner in the presence of a GSI (Figure 3E). The upregulation persisted over 7 days of culture in in 1.0 μ M GSI (Figure 3F).

To test the effect of GSI on BCMA shedding in MM, three different myeloma cell lines (MM1.R, U266 and 8226) were washed and plated in culture media for 24 hours in the presence of various concentrations of GSI RO429097 (0.01 μ M, 0.1 μ M and 1.0 μ M). The media supernatant was harvested and assayed for sBCMA by ELISA. sBCMA levels decreased in the supernatant in the presence of a GSI in a dose dependent manner (Figures 3G & 3H).

To examine the effect on soluble BCMA (sBCMA) levels over time, U266 cells were washed and plated in culture media for 1, 3, 5 and 24 hours in the presence of various concentrations of GSI RO429097 (0.01 μ M, 0.1 μ M and 1.0 μ M). The media supernatant was harvested and assayed for soluble BCMA (sBCMA) by ELISA. The data shows a dose-dependent decrease in soluble BCMA levels in the supernatant when a GSI is present (Figure 3I).

Next, the effect of removing GSI from MM cell culture was examined. As shown in Figures 3J and 3K, surface BCMA levels decreased following removal of 1.0 μ M RO429097, while supernatant sBCMA increased when the GSI was removed, but did not increase when the GSI remained. These data indicate a reversible inhibitory effect of GSI on surface BCMA shedding. Finally, the viability of the tested MM cell lines was not affected by addition of 1.0 μ M RO429097 to culture (Figure 3L).

The effect of GSI on BCMA expression was then tested on primary myeloma samples from patients. CD138⁺ myeloma cells were enriched from patient bone marrow samples, incubated for 3 hours in the presence of various concentrations of GSI RO429097 (0.01 μ M to 10 μ M) and evaluated for surface BCMA expression by flow cytometry (Figures 3M and 3R). BCMA mean fluorescence intensity (MFI) on tumor cells is presented as fold increase over tumor cells incubated without RO429097. There is an observed dose-dependent upregulation of BCMA on tumor cells (Figure 3N), while there was no effect on the levels of several other cell surface molecules, including CS1, CD86, PD-L1, CD80 and CD38 (Figures 3O-3Q).

EXAMPLE 5

γ -SECRETASE INHIBITOR IMPROVES RECOGNITION OF BCMA⁺ MYELOMA CELLS BY T CELLS EXPRESSING BCMA-SPECIFIC T-CHARM

To examine the effect of a GSI on CAR T cell activity against BCMA⁺ multiple myeloma, IL-2 production by BCMA CAR-T cells (BCMA-specific T-ChARM C11 3ST-CD28 and BCMA-specific T-ChARM C11 3ST-41BB) or control CD19sh CAR (short spacer) -T cells was measured after co-culture with primary human myeloma tumor cells for 24 hours in media containing at various concentrations of GSI RO429097 (0.003 μ M to 3.0 μ M) (Figure 4A). Treatment with GSI resulted in a dose dependent increase in IL-2 production by BCMA T-ChARM T cells. Also, IFN γ production by BMCA T-ChARM T cells co-cultured with myeloma cells at various concentrations of RO429097 was measured and was also increased in the presence of GSI (Figure 4B). These data show that multiple myeloma cells stimulate BCMA-specific T-ChARM T cells better when pre-treated with a γ -secretase inhibitor than

upregulates BCMA expression on the tumor cell. Finally, proliferation of CFSE-labeled BCMA-specific T-ChARM T cells was increased in a dose dependent manner after co-culture for 3 days with primary human myeloma tumor cells in presence of GSI RO492097 (Figure 4C).

5 Without wishing to be bound by theory, a potential caveat of using GSI to augment BCMA on myeloma cells to improve CAR-T cell recognition is that high concentrations of GSI may inhibit T-cell signaling and effector functions. *See Eagar et al., Immunity 20(4):407, 2004.* To evaluate potential effects of GSI on CAR T cells in a setting where target ligand expression remains stable, CD19 CAR-T cells were co-
10 cultured with Raji or K562/CD19⁺ target cells in the presence of GSI (0.01 μ M to 100 μ M) and measured viability and effector function. Over this concentration range there was no effect on CD19 antigen expression (Figure 5A) or on CAR-T cell viability after 24 hours (Figure 5B). GSI RO4929097 was found to inhibit CD19 CAR-T cell effector function at concentrations ≥ 1 μ M when co-cultured with K562 CD19⁺ cells, as
15 determined by measuring IL-2 (Figure 5C, upper panel) and IFN γ (Figure 5C, lower panel) production (*see also* Figure 5D-5F). However, the ability of the CD19 CAR-T cells to specifically kill target cells was not affected by GSI concentration (Figure 5G). The CD19 CAR-T cells also proliferated in response to target cell stimulation in the presence of 10 μ M GSI RO4929097, though less effectively than in the absence of GSI
20 RO4929097 (Figure 5H). To further investigate the dosage effect of GSI on CAR-T cell division, a dose titration of GSI RO4929097 was added to co-cultures of CD19 CAR-T and target cells. CAR-T cell division was essentially unaffected or only slightly affected by the addition of therapeutic doses of GSI (0.01 μ M-1 μ M), whereas GSI at a higher, non-therapeutic dose (10 μ M) did detectably inhibit proliferation (Figure 5I).
25 CD19 CAR-T cells cultured in the presence of GSI did not show a decrease in expansion over 8 days in culture with (Figure 5J) or without exogenous IL-2 (Figure 5K). Also, T cells that were expanded in the presence of 5 μ M, 0.5 μ M, or absence of GSI did not show a significant difference in IFN- γ production when subsequently re-stimulated with target cells after expansion in the presence (Figure 5L) or absence
30 (Figure 5M) of exogenous IL-2. IL-2 production between the CAR-T cell groups was also not markedly different (Figure 5N). Overall, GSI at therapeutic levels does not

affect T cell function, including cytokine production, cell lysis activity, and proliferation.

Next, the effect of GSI on T cell functionality of T cells containing a BCMA-specific T-ChARM was examined. GSI treatment of primary myeloma tumor samples greatly improved IFN- γ production (Figures 5O, 5P, and 5Q) and CD8 expression (Figure 5O) by co-cultured T-ChARM T cells, with effects seen following treatment with 0.1 μ M GSI or higher, and also increased proliferation of the T cells (Figure 5R).

These data indicate the clinical utility of combining BCMA-specific CAR-T cells and a GSI in order to increase the levels of BCMA on multiple myeloma cells. This may provide an advantage or even have a synergistic effect on treating multiple myeloma by facilitating elimination of tumor cells that express low levels of BCMA. The data also show that a range of concentrations of a γ -secretase inhibitor promoted BCMA expression of tumor cells without affecting T cell activity.

15

EXAMPLE 6

IN VIVO STUDY OF GSI ACTIVITY IN A MM XENOGRAFT MODEL

To investigate whether the sBCMA-inhibiting effects of GSI RO4929097 described in Example 4 could be replicated *in vivo*, a mouse xenograft model of human MM was tested as illustrated in Figure 6A. Briefly, immunodeficient *NOD/SCID* gamma (NSG) mice (The Jackson Laboratory) received sublethal irradiation (275 rad) at day -1, followed by 5×10^6 MM.1R cells the next day (day 0). GSI RO4929097 (30 mg/kg) was administered by oral gavage at day 19 and day 20. Mice were sacrificed at different times after the second gavage administration of the GSI and blood and bone marrow samples were taken for analysis. GSI dosing was followed by a rapid increase (>3-fold by 4 hours post-second gavage) in surface BCMA expression on the tumor cells (Figure 6B) and a concurrent decrease in sBCMA (Figure 6C). The observed effects were generally reduced within 48 hours following the second gavage treatment.

EXAMPLE 7***IN VIVO* STUDY OF GSI + CAR T CELL COMBINATION THERAPY TARGETING MM**

The ability of GSI therapy to improve anti-BCMA CAR T cell therapy *in vivo* was investigated. In a similar experiment to the one described in Example 6, NSG mice were sublethally irradiated at day -21, followed by administration of human MM cells (5x10⁶ MM.1R^{fluc}) at day -20 (see Figure 7A). A first dose of GSI (30 mg/kg RO4929097) was administered by oral gavage at day -1. At day 0, the mice were injected with a suboptimal dose of C11 3ST T-ChARM T cells (0.33x10⁶ cells; CD4:CD8 1:1) and received a second dose of the GSI. Further doses of GSI were administered at day +1, day +8, and day +9. Bioluminescence imaging (BLI) was performed throughout the experiment and mice were monitored for survival. As shown in Figures 7B, 7C, and 7E (left-hand graph), the group receiving GSI + T cells had reduced luminescence and more mice with no detectable tumors as compared to the group receiving the T cells only (BLI data from T cell only group not shown). The quantified luminescence data (Figures 7C and 7E (left-hand graph)) indicated that the initial T-ChARM T cell effect began reversing at approximately day 9, when the tumors began growing out ("Mock + BCMA T cells"). The anti-tumor effect was prolonged by combination with GSI RO4929097 ("RO49 + BCMA T cells"), but the tumors in that treatment group also eventually grew out. The imaging data was consistent with survival of the mice (Figures 7D and 7E (right-hand graph)), with the mice receiving the combination therapy surviving longer than the group receiving T cells only before succumbing. In certain embodiments, GSI treatments are repeated (at least 2 to at least about 5 times up to at least about 25 times) when used in combination with anti-BCMA CAR T cells, which GSI treatments may be administered concurrently with, before, or after administration of the anti-BCMA CAR T cells.

EXAMPLE 8**GSI IMPROVES BINDING TO BCMA+ MM CELLS BY A
BISPECIFIC ANTI-BCMA FUSION PROTEIN**

A bispecific fusion protein having specificity for BCMA and for another
5 antigen, and with a YFP (yellow fluorescent protein) tag, was constructed. The
bispecific fusion protein was added to BCMA-expressing H929 MM cells in culture
(0.5×10^6 cells) either with or without GSI. Binding was assessed by flow cytometry.
As shown in Figure 8, the addition of GSI improved binding by the BCMA-binding
bispecific fusion protein, but had no effect on binding by a control bispecific fusion
10 protein that does not target BCMA. These data show that immunotherapies involving
bispecific molecules that target BCMA can be augmented or improved with GSI.

The various embodiments described above can be combined to provide further
embodiments. All of the U.S. patents, U.S. patent application publications, U.S. patent
15 applications, foreign patents, foreign patent applications and non-patent publications
referred to in this specification and/or listed in the Application Data Sheet are
incorporated herein by reference, in their entirety. Aspects of the embodiments can be
modified, if necessary to employ concepts of the various patents, applications and
publications to provide yet further embodiments.

20 These and other changes can be made to the embodiments in light of the above-
detailed description. In general, in the following claims, the terms used should not be
construed to limit the claims to the specific embodiments disclosed in the specification
and the claims, but should be construed to include all possible embodiments along with
the full scope of equivalents to which such claims are entitled. Accordingly, the claims
25 are not limited by the disclosure.

CLAIMS:

1. A method of treating multiple myeloma in a subject, the method comprising administering to the subject a therapeutically effective amount of a T cell expressing a BCMA-specific chimeric antigen receptor (CAR) or tagged chimeric antigen receptor molecule (T-ChARM), and a therapeutically effective amount of a γ -secretase inhibitor.
2. A method of treating multiple myeloma in a subject, the method comprising administering to the subject a therapeutically effective amount of a T cell expressing a BCMA-specific chimeric antigen receptor (CAR) or tagged chimeric antigen receptor molecule (T-ChARM), wherein the subject has been administered a therapeutically effective amount of a γ -secretase inhibitor.
3. A method of treating multiple myeloma in a subject, the method comprising administering to the subject a therapeutically effective amount of a γ -secretase inhibitor, wherein the subject has been administered a therapeutically effective amount of a T cell expressing BCMA-specific chimeric antigen receptor (CAR) or tagged chimeric antigen receptor molecule (T-ChARM).
4. The method of any one of claims 1-3, comprising administering a lower amount of the T cell and/or a lower amount of the γ -secretase inhibitor, respectively, as compared to a method comprising administering the T cell or the γ -secretase inhibitor individually to treat the multiple myeloma.
5. Use of a combination of (i) a T cell expressing a BCMA-specific chimeric antigen receptor (CAR) or tagged chimeric antigen receptor molecule (T-ChARM) and (ii) a γ -secretase inhibitor in the preparation of a medicament for the treatment of multiple myeloma.
6. Use of a T cell expressing a BCMA-specific chimeric antigen receptor (CAR) or tagged chimeric antigen receptor molecule (T-ChARM) in the manufacture of a medicament for

the treatment of multiple myeloma, wherein in the treatment the medicament is administered with a γ -secretase inhibitor.

7. Use of a γ -secretase inhibitor in the manufacture of a medicament for the treatment of multiple myeloma, wherein in the treatment the medicament is administered with a T cell expressing a BCMA-specific chimeric antigen receptor (CAR) or tagged chimeric antigen receptor molecule (T-ChARM).

8. The use of claim 5, wherein the medicament comprises a lower amount of the T cell and/or a lower amount of the γ -secretase inhibitor, respectively, as compared to a medicament consisting or consisting essentially of the T cell or the γ -secretase inhibitor to treat the multiple myeloma.

9. The use of claim 6, wherein the medicament comprises a lower amount of the T cell as compared to a medicament consisting or consisting essentially of the T cell and not to be administered with the γ -secretase inhibitor to treat the multiple myeloma.

10. The use of claim 7, wherein the medicament comprises a lower amount of the γ -secretase inhibitor as compared to a medicament consisting or consisting essentially of the γ -secretase inhibitor and not to be administered with the T cell to treat multiple myeloma.

11. The method of any one of claims 1-4 or the use of any one of claims 5-10, wherein the BCMA-specific CAR or T-ChARM comprises a human or humanized binding domain and/or wherein the BCMA-specific CAR or T-ChARM comprises a BCMA-specific scFv, a BCMA-specific scTCR, a BCMA-specific nanobody, a BCMA-specific domain antibody, or a BCMA ligand or binding portion thereof, wherein, optionally,

the BCMA-specific CAR or T-ChARM comprises a scFv comprising heavy chain and light chain variable regions based on BCMA antibody J22.0-xi, J22.9-xi, J6M0, J6M1, J6M2, J9M0, J9M1, J9M2, 11D5-3, CA8, A7D12.2, C11 D5.3, C12A3.2, C13F12.1, 13C2, 17A5, 83A10, 13A4, 13D2, 14B11, 14E1, 29B11, 29F3, 13A7, CA7, S307118G03, SG1, S332121F02,

S332126E04, S322110D07, S336105A07, S335115G01, S335122F05, ET140-3, ET140-24, ET140-37, ET140-40, ET140-54, TBL-CLN1, C4.E2.1, Vicky-1, pSCHLI333, pSCHLI372, or pSCHLI373.

12. The method of any one of claims 1-4 and 11 or the use of any one of claims 5-11, wherein the BCMA-specific CAR or T-ChARM comprises a hydrophobic portion disposed between an extracellular component and an intracellular component, wherein the extracellular component comprises a BCMA-specific domain, the BCMA-specific binding domain comprising (a) a BCMA-specific antigen-binding portion from BCMA antibody J22.0-xi, J22.9-xi, J6M0, 11D5-3, CA8, A7D12.2, C11 D5.3, C12A3.2, C13F12.1, 13C2, 17A5, 83A10, 13A4, 13D2, 14B11, 14E1, 29B11, 29F3, 13A7, CA7, SG1, S307118G03, S332121F02, S332126E04, S322110D07, S336105A07, S335115G01, S335122F05, ET140-3, ET140-24, ET140-37, ET140-40, ET140-54, TBL-CLN1, C4.E2.1, Vicky-1, pSCHLI333, pSCHLI372, or pSCHLI373; or (b) a BCMA ligand or binding portion thereof based on or derived from BAFF or APRIL.

13. The method of any one of claims 1-4, 11, and 12 or the use of any one of claims 5-12, wherein the BCMA-specific CAR or T-ChARM comprises a transmembrane domain disposed between an extracellular component and an intracellular component.

14. The method of claim 13 or the use of claim 13, wherein the transmembrane domain is a CD4, CD8, CD28, or CD27 transmembrane domain.

15. The method of any one of claims 1-4 and 11-14 or the use of any one of claims 5-14, wherein the BCMA-specific CAR or T-ChARM comprises:

- (i) an extracellular component comprising a spacer region comprising a hinge, wherein the spacer region has a length of from about 15 to about 100 amino acids; and/or
- (ii) an extracellular component comprising a hinge region that comprises a cluster of differentiation molecule stalk region or a functional variant thereof; and/or

(iii) an intracellular component, wherein the intracellular component comprises an effector domain or functional portion thereof, a costimulatory domain or functional portion thereof, or any combination thereof.

16. The method of claim 15 or the use of claim 15, wherein the intracellular component comprises 4-1BB (CD137), CD3 ϵ , CD3 δ , CD3 ζ , CD25, CD27, CD28, CD79A, CD79B, CARD11, DAP10, FcR α , FcR β , FcR γ , Fyn, HVEM, ICOS, Lck, LAG3, LAT, LRP, NKG2D, NOTCH1, NOTCH2, NOTCH3, NOTCH4, OX40 (CD134), ROR2, Ryk, SLAMF1, Slp76, pT α , TCR α , TCR β , TRIM, Zap70, PTCH2, or a functional portion thereof, or in any combination thereof.

17. The method of claim 15 or 16 or the use of claim 15 or 16, wherein the effector domain comprises CD3 ζ or a functional portion thereof.

18. The method of any one of claims 1-4 and 11-17 or the use of any one of claims 5-17, wherein:

the BCMA-specific CAR or T-ChARM comprises an intracellular component comprising a costimulatory domain or a functional portion thereof selected from CD27, CD28, 4-1BB (CD137), OX40 (CD134), or any combination thereof; and/or

wherein the effector domain or effector portion thereof comprises CD3 ζ or functional portion thereof, and one or more costimulatory domain or a functional portion thereof of 4-1BB (CD137), CD27, CD28, and OX40 (CD134).

19. The method of any one of claims 15-18 or the use of any one of claims 15-18, wherein:

the intracellular component comprises (a) 4-1BB or a functional portion thereof and CD3 ζ , (b) CD27 or a functional portion thereof and CD3 ζ , (c) CD28 or a functional portion thereof and CD3 ζ , (d) OX40 or a functional portion thereof and CD3 ζ , (e) CD28 or a functional portion thereof, 4-1BB or a functional portion thereof and CD3 ζ , (f) OX40 or a functional

portion thereof, 4-1BB or a functional portion thereof and CD3 ζ , or (g) CD28 or a functional portion thereof, OX40 or a functional portion thereof and CD3 ζ ; and/or

wherein the CAR or T-ChARM comprises an extracellular component comprising an immunoglobulin hinge region, a CH2 domain and a CH3 domain, a CH3 domain, or any combination thereof disposed between a BCMA-specific binding domain and a hydrophobic portion, wherein, optionally, (1) the immunoglobulin hinge region is an IgG1 hinge region and/or (2) the CH2 domain is an IgG1 CH2 domain and the CH3 domain is an IgG1 CH3 domain.

20. The method of any one of claims 1-4 and 11-19 or the use of any one of claims 5-19, wherein the T cell is a human T cell, a CD4⁺ T cell, a CD8⁺ T cell, a CD4⁻ CD8⁻ double negative T cell, a $\gamma\delta$ T cell, a naïve T cell, a central memory T cell, an effector memory T cell, bulk T cells, or any combination thereof.

21. The method of any one of claims 1-4 and 11-20 or the use of any one of claims 5-20, wherein the γ -secretase inhibitor is a small molecule or a peptidomimetic compound.

22. The method of any one of claims 1-4 and 11-21 or the use of any one of claims 5-21, wherein the γ -secretase inhibitor is avagacestat, DAPT, BMS-906024, BMS-986115, MK-0752, PF-03084014, RO4929097, or YO-01027.

23. The method of any one of claims 1-4 and 11-22 or the use of any one of claims 5-22, wherein the γ -secretase inhibitor is a nicastrin-specific binding protein, wherein, optionally, the nicastrin-specific binding protein is scFvG9, antibody A5226A, antibody 2H6, or antibody 10C11.

24. The method of any one of claims 1-4 and 11-23 or the use of any one of claims 5-23, wherein the BCMA-specific CAR or T-ChARM comprises a variable domain comprising CDR1, CDR2, and CDR3 of the amino acid sequence of any one of SEQ ID NOS.:14-16 and a variable domain comprising CDR1, CDR2, and CDR3 of the amino acid sequence of any one of SEQ ID NOS.:17-21.

25. The method of any one of claims 1-4 and 11-24, wherein the method further comprises pre-conditioning the subject with an immunosuppressive regimen prior to or concurrent with the T cell and the γ -secretase inhibitor.

26. The use of any one of claims 5-24, wherein the subject has been pre-conditioned with an immunosuppressive regimen.

27. The method of claim 25 or the use of claim 26, wherein the immunosuppressive regimen is a non-myeloablative treatment or myeloablative treatment.

28. The method of claim 27 or the use of claim 27, wherein the non-myeloablative treatment comprises cyclophosphamide or cyclophosphamide in combination with fludarabine.

29. The method of any one of claims 1-4 , 11-25, 27, and 28, wherein:
 (i) the T cell and the γ -secretase inhibitor are administered sequentially or the T cell and the γ -secretase inhibitor are administered concurrently, wherein, optionally, the T cell and the γ -secretase inhibitor are formulated together; and/or
 (ii) the T cell is administered parenterally and the γ secretase inhibitor is administered orally.

30. The use of any one of claims 5-24 and 26-28, wherein:
 (i) the T cell and/or the γ -secretase inhibitor are formulated for sequential administration or the T cell and the γ -secretase inhibitor are formulated for concurrent administration, wherein, optionally, the T cell and the γ -secretase inhibitor are formulated together; and/or
 (ii) the T cell is formulated for parenteral administration and the γ -secretase inhibitor is formulated for oral administration.

31. The method of any one of claims 1-4, 11-25, and 27-29, wherein:

the γ -secretase inhibitor is administered to the subject at least once subsequent to a first administration of the T cell, wherein, optionally, the γ -secretase inhibitor is administered at least 2, at least 3, at least 4, at least 5, at least 6, at least 7, at least 8, at least 9, at least 10, at least 15, at least 20, at least 25, at least 30, at least 35, at least 40, at least 45, or at least 50 times subsequent to the first administration of the T cell; and/or the γ -secretase inhibitor is administered at a concentration of about 30 mg/kg.

32. The use of any one of claims 5-24, 26-28, and 30, wherein the subject has previously been administered the T cell and/or wherein the medicament comprises the γ -secretase inhibitor at a concentration of about 30 mg/kg relative to the subject's weight.

33. The method of any one of claims 1-4, 11-25, and 27-29, and 31, or the use of any one of claims 5-24, 26-28, 30, and 32, wherein the BCMA-specific CAR or T-ChARM is multispecific.

34. The method of claim 33 or the use of claim 33, wherein the BCMA-specific CAR or T-ChARM is bispecific.

35. The method of any one of claims 1-4, 11-25, 27-29, 31, 33 and 34, or the use of any one of claims 5-24, 26-28, 30, and 32-34, wherein the BCMA-specific CAR or T-ChARM comprises a BCMA-specific scFv, a BCMA-specific nanobody, or a BCMA ligand or binding portion thereof.

36. The method of claim 35 or the use of claim 35, wherein the BCMA-specific CAR or T-ChARM comprises a BCMA-specific scFv.

37. The method of any one of claims 1-4, 11-25, 27-29, 31, and 33-36 or the use of any one of claims 5-24, 26-28, 30, and 32-36, wherein:

(i) the CAR comprises (1) a BCMA-specific binding domain comprising a BCMA-specific scFv, a BCMA-specific domain antibody, or a BCMA ligand or binding portion thereof,

(2) an intracellular component comprising a costimulatory domain from 4-1BB and/or CD28, and an effector domain from CD3 ζ , and/or (3) an extracellular spacer region comprising a hinge, wherein the spacer region has a length from about 15 to about 100 amino acids;

- (ii) the γ -secretase inhibitor comprises a small molecule; and
- (iii) the T cell is a human T cell, a CD4⁺ T cell, a CD8⁺ T cell, a naïve T cell, a central memory T cell, an effector memory T cell, bulk T cells, or any combination thereof.

38. The method of any one of claims 1-4, 11-25, 27-29, 31, and 33-37, or the use of any one of claims 5-24, 26-28, 30, and 32-37, wherein:



- (i) the CAR comprises (1) a BCMA-specific binding domain comprising a BCMA-specific scFv, a BCMA-specific domain antibody, or a BCMA ligand or binding portion thereof, (2) an intracellular component comprising a costimulatory domain from 4-1BB or CD28, and an effector domain from CD3 ζ , and (3) an extracellular spacer region comprising a hinge, wherein the spacer region has a length from about 15 to about 100 amino acids;
- (ii) the γ -secretase inhibitor comprises a small molecule; and
- (iii) the T cell is a human T cell, a CD4⁺ T cell, a CD8⁺ T cell, a naïve T cell, a central memory T cell, an effector memory T cell, bulk T cells, or any combination thereof.

39. The method of any one of claims 1-4, 11-25, 27-29, 31, and 33-38, or the use of any one of claims 5-24, 26-28, 30, and 32-38, wherein the T cell comprises a CD8⁺ T cell, a CD4⁺ T cell, or both.

40. A kit when used for treating multiple myeloma, comprising (a) a unit dosage of a T cell expressing a BCMA-specific chimeric antigen receptor (CAR) or tagged chimeric antigen receptor molecule (T-ChARM), and (b) a unit dosage of a γ -secretase inhibitor.

41. The kit of claim 40, further comprising a CD20-specific binding protein such as rituximab, ofatumumab, ocrelizumab; a CD19-specific binding protein; a CD45-specific binding protein; a CD38-specific binding protein; a cytokine; a chemokine; a growth factor; a chemotherapeutic agent; or a radiotherapeutic agent.

1/57

*FIG. 1A**FIG. 1B*

2/57




FIG. 1C




FIG. 1D




FIG. 1E




FIG. 1F




FIG. 1G

5/57




Anti-BCMA
scFv C11

FIG. 1H

6/57

*FIG. 1I**FIG. 1J*






FIG. 1K

8/57

*FIG. 1L*

9/57

*FIG. 1M*




FIG. 1N

11/57




FIG. 1O





FIG. 1P

12/57

*FIG. 2A*






FIG. 2B

*FIG. 2C**FIG. 2D*

15/57








FIG. 2E

*FIG. 2F**FIG. 2G*

17/57

*FIG. 2H**FIG. 2I*

18/57




FIG. 2J




FIG. 2K

19/57




FIG. 2L

20/57




FIG. 2M




FIG. 2N

22/57




FIG. 20

23/57




FIG. 2P




FIG. 2Q

24/57




FIG. 3A

25/57

MM.1R




FIG. 3B




FIG. 3C

26/57




FIG. 3D




FIG. 3E

27/57




FIG. 3F





FIG. 3G

28/57

*FIG. 3H*

29/57




FIG. 3I




FIG. 3J

30/57




FIG. 3K




FIG. 3L




FIG. 3M

32/57




FIG. 3N




FIG. 30




FIG. 3P

35/57





FIG. 3Q

36/57

*FIG. 3R*

37/57




FIG. 4A




FIG. 4B

CAR-T cell proliferation




FIG. 4C





FIG. 5A

40/57

*FIG. 5B*

41/57




FIG. 5C

42/57




FIG. 5D




FIG. 5E

43/57




FIG. 5F




FIG. 5G

44/57




FIG. 5H




FIG. 5I

45/57




FIG. 5J







FIG. 5K

46/57

*FIG. 5L*

47/57

*FIG. 5M**FIG. 5N*

48/57




FIG. 50

49/57




FIG. 5P





FIG. 5Q

50/57

*FIG. 5R*




FIG. 6A

52/57




FIG. 6B




FIG. 6C




FIG. 7A




FIG. 7B




FIG. 7C




FIG. 7D





FIG. 7E

57/57

*FIG. 8*

360056_445WO_SEQUENCE_LISTING

SEQUENCE LISTING

<110> Fred Hutchinson Cancer Research Center
Riddell, Stanley R.
Green, Damian
Hill, Tyler

<120> COMBINATION THERAPIES FOR TREATMENT OF
BCMA RELATED CANCERS AND AUTOIMMUNE DISORDERS

<130> 360056.445WO

<140> PCT

<141> 2018-02-16

<150> US 62/582,270

<151> 2017-11-06

<150> US 62/406,612

<151> 2017-02-17

<160> 34

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic sequence

<400> 1

Gly Gly Gly Gly Ser
1 5

<210> 2

<211> 4

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic sequence

360056_445W0_SEQUENCE_LISTING

<400> 2

Gly Gly Gly Ser

1

<210> 3

<211> 25

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic sequence

<220>

<221> VARIANT

<222> (5)...(20)

<223> any one or all of amino acids 5-20 can either be present or absent

<400> 3

Gly Gly Gly Ser Gly Gly Gly Ser Gly Gly Gly Ser Gly Gly Gly Ser
1 5 10 15Gly Gly Gly Ser Gly Gly Gly Gly Ser
20 25

<210> 4

<211> 45

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic sequence

<220>

<221> VARIANT

<222> (5)...(0)

<223> any one or all of amino acids 5-20 can either be present or absent.

<220>

<221> VARIANT

<222> (26)...(45)

<223> any one or all of amino acids 26-45 can either be present or absent.

<400> 4

Gly Gly Gly Ser Gly Gly Gly Ser Gly Gly Gly Ser Gly Gly Gly Ser
1 5 10 15

360056_445WO_SEQUENCE_LISTING

Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly
 20 25 30
 Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser
 35 40 45

<210> 5
 <211> 25
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Synthetic sequence

<220>
 <221> VARIANT
 <222> (6)...(20)
 <223> any one or all of amino acids 6-20 can either be
 present or absent.

<400> 5
 Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly
 1 5 10 15
 Gly Gly Gly Ser Gly Gly Gly Gly Ser
 20 25

<210> 6
 <211> 6
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Synthetic sequence

<400> 6
 His Gly Gly His His Gly
 1 5

<210> 7
 <211> 8
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Synthetic sequence

360056_445WO_SEQUENCE_LISTING

<400> 7

Trp Ser His Pro Gln Phe Glu Lys
1 5

<210> 8

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic sequence

<400> 8

Trp Arg His Pro Gln Phe Gly Gly
1 5

<210> 9

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic sequence

<400> 9

Cys Pro Pro Cys Pro
1 5

<210> 10

<211> 29

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic sequence

<220>

<221> VARIANT

<222> (10)...(29)

<223> any one or all of amino acids 10-29 can either be present or absent.

<400> 10

Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly
1 5 10 15
Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser

360056_445WO_SEQUENCE_LISTING
25

20

<210> 11
 <211> 3
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Synthetic sequence

<400> 11
 Gly Gly Gly
 1

<210> 12
 <211> 15
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Synthetic sequence

<400> 12
 Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser
 1 5 10 15

<210> 13
 <211> 16
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Synthetic sequence

<400> 13
 Gly Gly Gly Ser Gly Gly Gly Ser Gly Gly Gly Ser Gly Gly Gly Ser
 1 5 10 15

<210> 14
 <211> 111
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Synthetic sequence

360056_445WO_SEQUENCE_LISTING

<400> 14

```

Asp Ile Val Leu Thr Gln Ser Pro Ala Ser Leu Ala Val Ser Leu Gly
 1           5           10           15
Glu Arg Ala Thr Ile Asn Cys Arg Ala Ser Glu Ser Val Ser Val Ile
      20           25           30
Gly Ala His Leu Ile His Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro
      35           40           45
Lys Leu Leu Ile Tyr Leu Ala Ser Asn Leu Glu Thr Gly Val Pro Ala
      50           55           60
Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser
65           70           75           80
Ser Leu Gln Ala Glu Asp Ala Ala Ile Tyr Tyr Cys Leu Gln Ser Arg
      85           90           95
Ile Phe Pro Arg Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys
      100           105           110

```

<210> 15

<211> 111

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic sequence

<400> 15

```

Asp Ile Val Leu Thr Gln Ser Pro Ala Ser Leu Ala Val Ser Leu Gly
 1           5           10           15
Glu Arg Ala Thr Ile Asn Cys Arg Ala Ser Glu Ser Val Ser Val Ile
      20           25           30
Gly Ala His Leu Ile His Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro
      35           40           45
Lys Leu Leu Ile Tyr Leu Ala Ser Asn Leu Glu Thr Gly Val Pro Ala
      50           55           60
Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser
65           70           75           80
Ser Leu Gln Ala Glu Asp Ala Ala Ile Tyr Ser Cys Leu Gln Ser Arg
      85           90           95
Ile Phe Pro Arg Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys
      100           105           110

```

<210> 16

<211> 111

<212> PRT

<213> Artificial Sequence

360056_445W0_SEQUENCE_LISTING

<220>

<223> Synthetic sequence

<400> 16

```

Asp Ile Val Leu Thr Gln Ser Pro Ala Ser Leu Ala Met Ser Leu Gly
 1           5           10           15
Glu Arg Ala Thr Ile Ser Cys Arg Ala Ser Glu Ser Val Ser Val Ile
          20          25          30
Gly Ala His Leu Ile His Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro
          35          40          45
Lys Leu Leu Ile Tyr Leu Ala Ser Asn Leu Glu Thr Gly Val Pro Ala
          50          55          60
Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser
65          70          75          80
Arg Val Gln Ala Glu Asp Ala Ala Ile Tyr Ser Cys Leu Gln Ser Arg
          85          90          95
Ile Phe Pro Arg Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys
          100          105          110

```

<210> 17

<211> 117

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic sequence

<400> 17

```

Gln Val Gln Leu Val Gln Ser Gly Ser Glu Leu Lys Lys Pro Gly Ala
 1           5           10           15
Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Tyr
          20          25          30
Ser Ile Asn Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
          35          40          45
Gly Trp Ile Asn Thr Glu Thr Arg Glu Pro Ala Tyr Ala Tyr Asp Phe
          50          55          60
Arg Gly Arg Phe Val Phe Ser Leu Asp Thr Ser Val Ser Thr Ala Tyr
65          70          75          80
Leu Gln Ile Ser Ser Leu Lys Ala Glu Asp Thr Ala Val Tyr Tyr Cys
          85          90          95
Ala Arg Asp Tyr Ser Tyr Ala Met Asp Tyr Trp Gly Gln Gly Thr Leu
          100          105          110
Val Thr Val Ser Ser
          115

```

<210> 18

360056_445WO_SEQUENCE_LISTING

<211> 117

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic sequence

<400> 18

Gln	Val	Gln	Leu	Val	Gln	Ser	Gly	Ser	Glu	Leu	Lys	Lys	Pro	Gly	Glu
1				5					10					15	
Ser	Val	Lys	Val	Ser	Cys	Lys	Ala	Ser	Gly	Tyr	Thr	Phe	Thr	Asp	Tyr
		20						25					30		
Ser	Ile	Asn	Trp	Val	Arg	Gln	Ala	Pro	Gly	Gln	Gly	Leu	Glu	Trp	Met
		35					40					45			
Gly	Trp	Ile	Asn	Thr	Glu	Thr	Arg	Glu	Pro	Ala	Tyr	Ala	Tyr	Asp	Phe
	50					55					60				
Arg	Gly	Arg	Phe	Val	Phe	Ser	Leu	Asp	Thr	Ser	Val	Ser	Thr	Ala	Tyr
65					70					75					80
Leu	Gln	Ile	Ser	Ser	Leu	Lys	Ala	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys
			85						90					95	
Ala	Leu	Asp	Tyr	Ser	Tyr	Ala	Met	Asp	Tyr	Trp	Gly	Gln	Gly	Thr	Leu
		100						105					110		
Val	Thr	Val	Ser	Ser											
		115													

<210> 19

<211> 117

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic sequence

<400> 19

Gln	Val	Gln	Leu	Val	Gln	Ser	Gly	Ser	Glu	Leu	Lys	Lys	Pro	Gly	Glu
1				5					10					15	
Ser	Val	Lys	Ile	Ser	Cys	Lys	Ala	Ser	Gly	Tyr	Thr	Phe	Thr	Asp	Tyr
		20						25					30		
Ser	Ile	Asn	Trp	Val	Arg	Gln	Ala	Pro	Gly	Gln	Gly	Leu	Glu	Trp	Met
		35					40					45			
Gly	Trp	Ile	Asn	Thr	Glu	Thr	Arg	Glu	Pro	Ala	Tyr	Ala	Tyr	Asp	Phe
	50					55					60				
Arg	Gly	Arg	Phe	Val	Phe	Ser	Leu	Asp	Thr	Ser	Ala	Ser	Thr	Ala	Tyr
65					70					75					80
Leu	Gln	Ile	Ser	Ser	Leu	Lys	Ala	Glu	Asp	Thr	Ala	Val	Tyr	Phe	Cys
			85						90					95	

360056_445WO_SEQUENCE_LISTING

Ala Leu Asp Tyr Ser Tyr Ala Met Asp Tyr Trp Gly Gln Gly Thr Leu
 100 105 110
 Val Thr Val Ser Ser
 115

<210> 20

<211> 117

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic sequence

<400> 20

Gln Val Gln Leu Val Gln Ser Gly Ser Glu Leu Lys Lys Pro Gly Glu
 1 5 10 15
 Ser Val Lys Ile Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Tyr
 20 25 30
 Ser Ile Asn Trp Val Lys Gln Ala Pro Gly Gln Gly Leu Lys Trp Met
 35 40 45
 Gly Trp Ile Asn Thr Glu Thr Arg Glu Pro Ala Tyr Ala Tyr Asp Phe
 50 55 60
 Arg Gly Arg Phe Val Phe Ser Leu Asp Thr Ser Ala Ser Thr Ala Tyr
 65 70 75 80
 Leu Gln Ile Ser Ser Leu Lys Ala Glu Asp Thr Ala Val Tyr Phe Cys
 85 90 95
 Ala Leu Asp Tyr Ser Tyr Ala Met Asp Tyr Trp Gly Gln Gly Thr Leu
 100 105 110
 Val Thr Val Ser Ser
 115

<210> 21

<211> 117

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic sequence

<400> 21

Gln Ile Gln Leu Val Gln Ser Gly Ser Glu Leu Lys Lys Pro Gly Glu
 1 5 10 15
 Ser Val Lys Ile Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Tyr
 20 25 30
 Ser Ile Asn Trp Val Lys Gln Ala Pro Gly Gln Gly Leu Lys Trp Met
 35 40 45

360056_445WO_SEQUENCE_LISTING

Gly Trp Ile Asn Thr Glu Thr Arg Glu Pro Ala Tyr Ala Tyr Asp Phe
50 55 60
Arg Gly Arg Phe Val Phe Ser Leu Asp Thr Ser Ala Ser Thr Ala Tyr
65 70 75 80
Leu Gln Ile Ser Ser Leu Lys Ala Glu Asp Thr Ala Val Tyr Phe Cys
85 90 95
Ala Leu Asp Tyr Ser Tyr Ala Met Asp Tyr Trp Gly Gln Gly Thr Leu
100 105 110
Val Thr Val Ser Ser
115

<210> 22
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<400> 22
caccggagaa tcaaaatcgg tgaat 25

<210> 23
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<400> 23
aaacattcac cgattttgat tctcc 25

<210> 24
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<400> 24
caccgcagtt gtgtgacacg gaag 24

<210> 25

360056_445WO_SEQUENCE_LISTING

<211> 24
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Synthetic sequence

 <400> 25
 aaaccttccg tgtcacacaa ctgc 24

 <210> 26
 <211> 25
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Synthetic sequence

 <400> 26
 caccggcaaa ggtgagtgag acttt 25

 <210> 27
 <211> 25
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Synthetic sequence

 <400> 27
 aaacaaagtc tcactcacct ttgcc 25

 <210> 28
 <211> 25
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Synthetic sequence

 <400> 28
 caccggtttc tgcagccgct ttggg 25

 <210> 29

360056_445WO_SEQUENCE_LISTING

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic sequence

<400> 29

aaaccccaaa gcggctgcag aaacc

25

<210> 30

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic sequence G4S linker

<400> 30

Gly Gly Gly Gly Ser

1

5

<210> 31

<211> 20

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic sequence

<220>

<221> VARIANT

<222> (1)...(20)

<223> any one of amino acids 1-2 can be absent or can be present up 10 times, provided that amino acids 1 and 2 are not both absent

<400> 31

Gly Gly Gly Gly Gly Gly Gly Gly Gly Gly Ser Ser Ser Ser Ser Ser

1

5

10

15

Ser Ser Ser Ser

20

<210> 32

<211> 10

<212> PRT

360056_445WO_SEQUENCE_LISTING

<213> Artificial Sequence

<220>

<223> Synthetic sequence

<400> 32

Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser
1 5 10

<210> 33

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic sequence

<400> 33

Gly Gly Gly Ser Gly Gly Gly Ser
1 5

<210> 34

<211> 11

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic sequence

<400> 34

Gly Gly Gly Ser Gly Gly Gly Ser Gly Gly Ser
1 5 10