



(51) International Patent Classification:

A61K 48/00 (2006.01)

A61K 47/46 (2006.01)

C12N 15/67 (2006.01)

A61K 31/7088 (2006.01)

C07H 21/00 (2006.01)

A61P 3/00 (2006.01)

(21) International Application Number:

PCT/US2017/033402

(22) International Filing Date:

18 May 2017 (18.05.2017)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/338,457

18 May 2016 (18.05.2016)

US

(71) Applicant: MODERNATX, INC. [US/US]; 200 Technology Square, Cambridge, Massachusetts 02139 (US).

(72) Inventors: MARTINI, Paolo; c/o Modernatx, Inc., 200 Technology Square, Cambridge, Massachusetts 02139 (US). HOGE, Stephen; c/o Modernatx, Inc., 200 Technology Square, Cambridge, Massachusetts 02139 (US). BENENATO, Kerry; c/o Modernatx, Inc., 200 Technology Square, Cambridge, Massachusetts 02139 (US). PRESNYAK, Vladimir; c/o Modernatx, Inc., 200 Technology Square, Cambridge, Massachusetts 02139 (US). MCFADYEN, Iain; c/o Modernatx, Inc., 200 Technology Square, Cambridge, Massachusetts 02139 (US). KUMARASINGHE, Ellalahewage Sathyajith; c/o Modernatx, Inc., 200 Technology Square, Cambridge, Massachusetts 02139 (US). RACHLIN, Evan Lockwood; c/o Modernatx, Inc., 200 Technology Square, Cambridge, Massachusetts 02139 (US). SABNIS, Staci; c/o Modernatx, Inc., 200 Technology Square, Cambridge, Massachusetts 02139 (US).

(74) Agent: NANNENGA-COMBS, Bonnie W. et al.; Sterne, Kessler, Goldstein & Fox P.L.L.C, 1100 New York Avenue, NW, Washington, District of Columbia 20005 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG,

MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

— with sequence listing part of description (Rule 5.2(a))

(54) Title: POLYNUCLEOTIDES ENCODING ACYL-COA DEHYDROGENASE, VERY LONG-CHAIN FOR THE TREATMENT OF VERY LONG-CHAIN ACYL-COA DEHYDROGENASE DEFICIENCY

(57) Abstract: The invention relates to mRNA therapy for the treatment of VLCADD. mRNAs for use in the invention, when administered *in vivo*, encode human acyl-CoA dehydrogenase, very longchain (ACADVL), isoforms thereof, functional fragments thereof, and fusion proteins comprising ACADVL. mRNAs of the invention are preferably encapsulated in lipid nanoparticles (LNPs) to effect efficient delivery to cells and/or tissues in subjects, when administered thereto. mRNA therapies of the invention increase and/or restore deficient levels of ACADVL expression and/or activity in subjects. mRNA therapies of the invention further decrease levels of toxic metabolites associated with deficient ACADVL activity in subjects, namely acylcarnitine and acylcarnitine metabolites.



POLYNUCLEOTIDES ENCODING ACYL-COA DEHYDROGENASE, VERY LONG-CHAIN FOR THE TREATMENT OF VERY LONG-CHAIN ACYL-COA DEHYDROGENASE DEFICIENCY

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims the priority benefit of U.S. Provisional Application No. 62/338,457, filed May 18, 2016, which is hereby incorporated by reference herein in its entirety.

REFERENCE TO A SEQUENCE LISTING SUBMITTED ELECTRONICALLY VIA EFS-WEB

[0002] The content of the electronically submitted sequence listing (Name: 3529.090pc02 Sequence_Listing_ST25.txt, Size: 109,576 bytes; and Date of Creation: May 17, 2017) is herein incorporated by reference in its entirety.

BACKGROUND

[0003] Very long-chain acyl-CoA dehydrogenase deficiency (VLCADD) is an autosomal recessive metabolic disorder characterized by the abnormal buildup of very long-chain fatty acids in patients. Such buildup of fatty acids can damage internal organs, resulting in a wide-range of symptoms. Clinically, there are three different types of VLCADD, with each type exhibiting different onset and/or severity. Andresen, B. *et al.*, *Am J Hum Genet.* 64:479-494 (1999). The most severe form of the disorder is "early" VLCADD. Signs and symptoms (e.g., hypoglycemia, irritability, and lethargy) usually appear between birth and four months. Left untreated, early VLCADD results in high mortality with majority of the patients dying from cardiomyopathy. In contrast, the "childhood" and "adult" forms of VLCADD often have much milder signs and symptoms (e.g., hypoglycemia and muscle weakness) that can be exacerbated by illness or long periods of fasting. However, left untreated, childhood and adult VLCADD can also result in more dire consequences, including, but not limited to, liver failure, seizure, kidney failure, and brain damage.

[0004] VLCADD has an estimated incidence of 1 in 31,500 to 1:125,000 live births. Mendez-Figueroa, H *et al.*, *J Perinatol.* 30:558-62 (2010). Patients from all ethnic groups have been reported, and males and females are affected equally. Current treatment for VLCADD is primarily via dietary control (e.g., low-fat, high-carbohydrate diet with

frequent feedings to avoid extended periods of fasting) in order to limit the usage of metabolic pathways required for the breakdown of very long-chain fatty acids. Solis, J. *et al.*, *J Am Diet Assoc.* 102:1800-1803 (2002). However, such treatment often fails to completely or reliably control the disorder. Therefore, there is a need for improved therapy to treat VLCADD.

[0005] The principal gene associated with VLCADD is acyl-CoA dehydrogenase, very long-chain (NM_000018.3; NP_000009.1; also referred to as ACADVL, VLCAD, ACAD6, or LCACD). Moczulski, D. *et al.*, *Postepy Hig Med Dosw.* 63: 266-277 (2009). ACADVL is a metabolic enzyme (E.C. 1.3.8.9), which plays a critical role in the catabolism of long-chain fatty acids, with highest specificity for carbon lengths C14-C18. Keeler, AM *et al.*, *Mol. Ther.* 20: 1131-38 (2012). ACADVL's biological function is to catalyze the first step of the mitochondrial fatty acid beta-oxidation pathway. ACADVL localizes to the inner mitochondrial membrane, where it functions as a homodimer. Souri, M. *et al.*, *FEBS Lett.* 426:187-190 (1998). The precursor form of human ACADVL is 655 amino acids in length, while its mature form is 615 amino acids long – a 40 amino acid leader sequence is cleaved off by mitochondrial importation and processing machinery. Souri, M. *et al.*, *Am J Hum Genet.* 58:97-106 (1996). This leader sequence is referred to as ACADVL's mitochondrial transit peptide.

[0006] Mutations within the ACADVL gene can result in the complete or partial loss of ACADVL function, resulting in the abnormal buildup of very long-chain fatty acids in the plasma and the attendant signs and symptoms described above. Moczulski, D. *et al.*, *Postepy Hig Med Dosw.* 63: 266-277 (2009). Nonetheless, there is currently no available therapeutic for VLCADD that completely or reliably controls the disorder. As such, there is a need for improved therapy to treat VLCADD.

BRIEF SUMMARY

[0007] The present invention provides mRNA therapeutics for the treatment of VLCADD. The mRNA therapeutics of the invention are particularly well-suited for the treatment of VLCADD as the technology provides for the intracellular delivery of mRNA encoding ACADVL followed by *de novo* synthesis of functional ACADVL protein within target cells. The instant invention features the incorporation of modified nucleotides within therapeutic mRNAs to (1) minimize unwanted immune activation (*e.g.*, the innate immune response associated with the *in vivo* introduction of foreign

nucleic acids) and (2) optimize the translation efficiency of mRNA to protein. Exemplary aspects of the invention feature a combination of nucleotide modification to reduce the innate immune response and sequence optimization, in particular, within the open reading frame (ORF) of therapeutic mRNAs encoding ACADVL to enhance protein expression.

- [0008] In further embodiments, the mRNA therapeutic technology of the instant invention also features delivery of mRNA encoding ACADVL *via* a lipid nanoparticle (LNP) delivery system. The instant invention features novel ionizable lipid-based LNPs which have improved properties when combined with mRNA encoding ACADVL and administered *in vivo*, for example, cellular uptake, intracellular transport and/or endosomal release or endosomal escape. The LNP formulations of the invention also demonstrate reduced immunogenicity associated with the *in vivo* administration of LNPs.
- [0009] In certain aspects, the invention relates to compositions and delivery formulations comprising a polynucleotide, e.g., a ribonucleic acid (RNA), e.g., a messenger RNA (mRNA), encoding ACADVL and methods for treating VLCADD in a subject in need thereof by administering the same. In some embodiments, the invention relates to a pharmaceutical composition comprising a lipid nanoparticle encapsulated mRNA that comprises an ORF encoding an ACADVL polypeptide, wherein the composition is suitable for administration to a human subject in need of treatment for VLCADD.
- [0010] In certain aspects, the invention relates to a pharmaceutical composition comprising (a) an mRNA that comprises (i) an open reading frame (ORF) encoding an acyl-CoA dehydrogenase, very long-chain (ACADVL) polypeptide, wherein the ORF comprises at least one chemically modified nucleobase, sugar, backbone, or any combination thereof and (ii) an untranslated region (UTR) comprising a microRNA (miRNA) binding site; and (b) a delivery agent, wherein the pharmaceutical composition is suitable for administration to a human subject in need of treatment for VLCADD.
- [0011] In certain aspects, the invention relates to a pharmaceutical composition comprising an mRNA that comprises an open reading frame (ORF) encoding a human acyl-CoA dehydrogenase, very long-chain (ACADVL) polypeptide, wherein the composition when administered as a single intravenous dose to a human subject in need thereof is sufficient to (i) increase plasma ACADVL activity level to a level at or above a reference physiologic level for at least 24 hours post-administration, and/or (ii) maintain plasma ACADVL activity level at 5%, 10%, 15%, 20%, 25%, 30%, 40%, 50%, or more of a reference plasma ACADVL activity level for at least 24 hours post-administration.

[0012] In certain aspects, the invention relates to a pharmaceutical composition comprising an mRNA comprising an open reading frame (ORF) encoding a human ACADVL polypeptide, wherein the composition when administered as a single intravenous dose to a subject in need thereof is sufficient to reduce blood and/or plasma levels of an acylcarnitine by at least 5%, at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or 100% compared to the subject's baseline level or a reference acylcarnitine blood and/or plasma level, for at least 24 hours, at least 48 hours, at least 72 hours, at least 96 hours, or at least 120 hours post-administration. In some embodiments, the acylcarnitine is an acylcarnitine metabolite selected from the group consisting of C12:1 acylcarnitine, C14:1 acylcarnitine, C14:2 acylcarnitine, C14 acylcarnitine, C16 acylcarnitine, C18 acylcarnitine, C18:1 acylcarnitine, and combinations thereof.

[0013] In certain aspects, the invention relates to a pharmaceutical composition comprising an mRNA comprising an open reading frame (ORF) encoding a human ACADVL polypeptide, wherein the composition when administered as a single intravenous dose to a subject in need thereof is sufficient to reduce blood and/or plasma levels of (i) C14:1 acylcarnitine by at least 5%, at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95% or at least 100% compared to the subject's baseline level or a reference C14:1 acylcarnitine blood and/or plasma level, for at least 24 hours, at least 48 hours, at least 72 hours, at least 96 hours, or at least 120 hours post-administration; (ii) C14:2 acylcarnitine by at least 5%, at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95% or at least 100% compared to the subject's baseline level or a reference C14:2 acylcarnitine blood and/or plasma level, for at least 24 hours, at least 48 hours, at least 72 hours, at least 96 hours, or at least 120 hours post-administration; (iii) C14 acylcarnitine by at least 5%, at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95% or at least 100% compared to the subject's baseline level or a reference C14 acylcarnitine blood and/or plasma level, for at least 24 hours, at least 48 hours, at least 72 hours, at least 96 hours, or at least 120 hours post-administration; and/or (iv) C12:1 acylcarnitine by at least 5%, at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95% or at least 100% compared to the subject's baseline level or a reference C12:1 acylcarnitine

blood and/or plasma level, for at least 24 hours, at least 48 hours, at least 72 hours, at least 96 hours, or at least 120 hours post-administration.

[0014] In certain aspects, the invention relates to a pharmaceutical composition comprising an mRNA comprising an open reading frame (ORF) encoding a human ACADVL polypeptide, wherein the composition when administered as a single intravenous dose to a subject in need thereof is sufficient to reduce blood and/or plasma levels of an acylcarnitine to a concentration of less than 1.0 $\mu\text{mol/L}$ or less than 0.8 $\mu\text{mol/L}$ in a subject having very long-chain acyl-CoA dehydrogenase deficiency (VLCADD), for at least 24 hours, at least 48 hours, at least 72 hours, at least 96 hours, or at least 120 hours post-administration. In some embodiments, the acylcarnitine is an acylcarnitine metabolite selected from the group consisting of C12:1 acylcarnitine, C14:1 acylcarnitine, C14:2 acylcarnitine, C14 acylcarnitine, C16 acylcarnitine, C18 acylcarnitine, C18.1 acylcarnitine, and combinations thereof.

[0015] In some embodiments, the pharmaceutical composition further comprises a delivery agent.

[0016] In some aspects, the invention relates to a polynucleotide comprising an open reading frame (ORF) encoding an acyl-CoA dehydrogenase, very long chain (ACADVL) polypeptide, wherein the uracil or thymine content of the ORF relative to the theoretical minimum uracil or thymine content of a nucleotide sequence encoding the ACADVL polypeptide (%U_{TM} or %T_{TM}) is between 100% and about 150%. In some embodiments, the %U_{TM} or %T_{TM} of the ORF is between about 105% and about 140%, about 110% and about 140%, 115% and about 140%, about 105% and about 130%, about 110% and about 130%, about 115% and about 135%, about 105% and about 135%, about 110% and about 135%, or about 115% and about 130%. In some embodiments, the %U_{TM} or %T_{TM} of the ORF is between (i) 110%, 111%, 112%, 113%, 114%, 115%, 116%, 117%, or 118% and (ii) 128%, 129%, 130%, 131%, 132%, 133%, 134%, 135%, 136%, 137%, 138%, 139%, or 140%. In some embodiments, the uracil or thymine content of the ORF is less than the uracil or thymine content in the corresponding wild-type ORF (%U_{WT} or %T_{WT}). In some embodiments, the uracil or thymine content of the ORF is less than about 95%, less than about 90%, less than about 85%, less than 80%, less than 79%, less than 78%, less than 77%, less than 76%, less than 75%, or less than 74%. In some embodiments, the %U_{WT} or %T_{WT} of the ORF is between 69% and 75% of the %U_{WT} or %T_{WT}. In some embodiments, the uracil or thymine content of the ORF relative to the total nucleotide

content in the ORF (%U_{TL} or %T_{TL}) is less than about 50%, less than about 40%, less than about 30%, or less than about 20%.

[0017] In certain embodiments, the uracil or thymine content in the ORF relative to the total nucleotide content in the ORF (%U_{TL} or %T_{TL}) is less than about 50%, less than about 40%, less than about 30%, or less than about 20%. In some embodiments, the %U_{TL} or %T_{TL} of the ORF is less than about 16%. In some embodiments, the %U_{TL} or %T_{TL} of the ORF is between about 14% and about 16%.

[0018] In some embodiments, the guanine content of the ORF with respect to the theoretical maximum guanine content of a nucleotide sequence encoding the ACADVL polypeptide (%G_{TMX}) is at least 70%, at least 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, or about 100%. In some embodiments, the %G_{TMX} of the ORF is between about 70% and about 80%, between about 71% and about 79%, between about 71% and about 77%, or between about 72% and about 76%. In some embodiments, the cytosine content of the ORF relative to the theoretical maximum cytosine content of a nucleotide sequence encoding the ACADVL polypeptide (%C_{TMX}) is at least 58%, at least 59%, at least 60%, at least 65%, at least 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, or about 100%. In some embodiments, the %C_{TMX} of the ORF is between about 60% and about 80%, between about 62% and about 77%, between about 63% and about 78%, or between about 68% and about 75%. In some embodiments, the guanine and cytosine content (G/C) of the ORF relative to the theoretical maximum G/C content in a nucleotide sequence encoding the ACADVL polypeptide (%G/C_{TMX}) is at least 82%, at least 83%, at least 84%, at least 85%, at least 90%, at least about 95%, or about 100%. In some embodiments, the %G/C_{TMX} of the ORF is between about 80% and about 100%, between about 85% and about 99%, between about 90% and about 96%, or between about 92% and about 95%. In some embodiments, the G/C content of the ORF relative to the G/C content of the corresponding wild-type ORF (%G/C_{WT}) is at least 102%, at least 103%, at least 104%, at least 105%, at least 106%, at least 107%, at least about 110%, or at least about 112%. In some embodiments, the average G/C content in the 3rd codon position of the ORF is at least 20%, at least 21%, at least 22%, at least 23%, at least 24%, or at least 25% higher than the average G/C content in the 3rd codon position of the corresponding wild-type ORF.

[0019] In some embodiments, the invention relates to a polynucleotide comprising an ORF, (i) wherein the ORF is at least 83%, at least 84%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identical to ACADVLCO4, ACADVLCO10, or ACADVLCO17, (ii) wherein the ORF is at least 82%, at least 83%, at least 84%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identical to ACADVLCO1, ACADVLCO7, ACADVLCO15, ACADVLCO16, ACADVLCO18, ACADVLCO22, or ACADVLCO23, (iii) wherein the ORF is at least 81%, at least 82%, at least 83%, at least 84%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identical to ACADVLCO2, ACADVLCO3, ACADVLCO5, ACADVLCO6, ACADVLCO8, ACADVLCO9, ACADVLCO111, ACADVLCO12, ACADVLCO13, ACADVLCO14, ACADVLCO19, ACADVLCO20, ACADVLCO21, ACADVLCO24, or ACADVLCO25.

[0020] In some embodiments, the invention relates to a polynucleotide comprising an ORF, (i) wherein the ORF is at least 83%, at least 84%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identical to ACADVLCO4, ACADVLCO10, or ACADVLCO17, (ii) wherein the ORF is at least 82%, at least 83%, at least 84%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identical to ACADVLCO1, ACADVLCO7, ACADVLCO15, ACADVLCO16, ACADVLCO18, ACADVLCO22, or ACADVLCO23, (iii) wherein the ORF is at least 81%, at least 82%, at least 83%, at least 84%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identical to ACADVLCO2, ACADVLCO3, ACADVLCO5, ACADVLCO6, ACADVLCO8, ACADVLCO9, ACADVLCO111, ACADVLCO12, ACADVLCO13, ACADVLCO14, ACADVLCO19, ACADVLCO20, ACADVLCO21, ACADVLCO24, or ACADVLCO25.

CO14, ACADVL-CO19, ACADVL-CO20, ACADVL-CO21, ACADVL-CO24, or ACADVL-CO25.

- [0021] In some embodiments, the ORF of the polynucleotide has at least 81%, at least 82%, at least 83%, at least 84%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to a sequence selected from the group consisting of SEQ ID NOs: 7 to 31.
- [0022] In some embodiments, the ACADVL polypeptide comprises an amino acid sequence at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99%, or about 100% identical to the polypeptide sequence of wild type ACADVL, isoform 1 (SEQ ID NO: 1), and wherein the ACADVL polypeptide has acyl-CoA dehydrogenase, very long chain activity.
- [0023] In some embodiments, the ACADVL polypeptide comprises an amino acid sequence at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99%, or about 100% identical to the polypeptide sequence of wild type ACADVL, isoform 2 (SEQ ID NO: 3), and wherein the ACADVL polypeptide has acyl-CoA dehydrogenase, very long chain activity.
- [0024] In some embodiments, the ACADVL polypeptide comprises an amino acid sequence at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99%, or about 100% identical to the polypeptide sequence of wild type ACADVL, isoform 3 (SEQ ID NO: 5), and wherein the ACADVL polypeptide has acyl-CoA dehydrogenase, very long chain activity.
- [0025] In some embodiments, the ACADVL polypeptide is a variant, derivative, or mutant having an acyl-CoA dehydrogenase, very long chain activity.
- [0026] In some embodiments, the polynucleotide sequence of the invention further comprises a nucleotide sequence encoding a transit peptide.
- [0027] In some embodiments, the polynucleotide of the invention is single stranded. In some embodiments, the polynucleotide of the invention is double stranded. In some embodiments, the polynucleotide of the invention is DNA. In some embodiments, the polynucleotide of the invention is RNA. In some embodiments, the polynucleotide of the invention is mRNA.
- [0028] In some embodiments, the polynucleotide of the invention comprises at least one chemically modified nucleobase, sugar, backbone, or any combination thereof. In some

embodiments, the at least one chemically modified nucleobase of the polynucleotide of the invention is selected from the group consisting of pseudouracil (ψ), N1-methylpseudouracil (m1 ψ), 1-ethylpseudouracil, 2-thiouracil (s2U), 4'-thiouracil, 5-methylcytosine, 5-methyluracil, and any combinations thereof. In some embodiments, the at least one chemically modified nucleobase of the polynucleotide of the invention is 5-methoxyuracil. In some embodiments, at least about 25%, at least about 30%, at least about 40%, at least about 50%, at least about 60%, at least about 70%, at least about 80%, at least about 90%, at least about 95%, at least about 99%, or 100% of the uracils of the polynucleotide of the invention are 5-methoxyuracils.

- [0029] In some embodiments, the polynucleotide further comprises a miRNA binding site.
- [0030] In some embodiments, the polynucleotide comprises at least two different microRNA (miR) binding sites.
- [0031] In some embodiments, the microRNA is expressed in an immune cell of hematopoietic lineage or a cell that expresses TLR7 and/or TLR8 and secretes pro-inflammatory cytokines and/or chemokines, and wherein the polynucleotide (e.g., mRNA) comprises one or more modified nucleobases.
- [0032] In some embodiments, the mRNA comprises at least one first microRNA binding site of a microRNA abundant in an immune cell of hematopoietic lineage and at least one second microRNA binding site is of a microRNA abundant in endothelial cells.
- [0033] In some embodiments, the mRNA comprises multiple copies of a first microRNA binding site and at least one copy of a second microRNA binding site.
- [0034] In some embodiments, the mRNA comprises first and second microRNA binding sites of the same microRNA.
- [0035] In some embodiments, the microRNA binding sites are of the 3p and 5p arms of the same microRNA.
- [0036] In some embodiments, the microRNA binding site comprises one or more nucleotide sequences selected from Table 3 or Table 4.
- [0037] In some embodiments, the microRNA binding site binds to miR-126, miR-142, miR-144, miR-146, miR-150, miR-155, miR-16, miR-21, miR-223, miR-24, miR-27 or miR-26a, or any combination thereof.
- [0038] In some embodiments, the microRNA binding site binds to miR126-3p, miR-142-3p, miR-142-5p, or miR-155, or any combination thereof.

- [0039] In some embodiments, the microRNA binding site is a miR-126 binding site. In some embodiments, at least one microRNA binding site is a miR-142 binding site. In some embodiments, one microRNA binding site is a miR-126 binding site and the second microRNA binding site is for a microRNA selected from the group consisting of miR-142-3p, miR-142-5p, miR-146-3p, miR-146-5p, miR-155, miR-16, miR-21, miR-223, miR-24 and miR-27.
- [0040] In some embodiments, the mRNA comprises at least one miR-126-3p binding site and at least one miR-142-3p binding site. In some embodiments, the mRNA comprises at least one miR-142-3p binding site and at least one 142-5p binding site.
- [0041] In some embodiments, the microRNA binding sites are located in the 5' UTR, 3' UTR, or both the 5' UTR and 3' UTR of the mRNA. In some embodiments, the microRNA binding sites are located in the 3' UTR of the mRNA. In some embodiments, the microRNA binding sites are located in the 5' UTR of the mRNA. In some embodiments, the microRNA binding sites are located in both the 5' UTR and 3' UTR of the mRNA. In some embodiments, at least one microRNA binding site is located in the 3' UTR immediately adjacent to the stop codon of the coding region of the mRNA. In some embodiments, at least one microRNA binding site is located in the 3' UTR 70-80 bases downstream of the stop codon of the coding region of the mRNA. In some embodiments, at least one microRNA binding site is located in the 5' UTR immediately preceding the start codon of the coding region of the mRNA. In some embodiments, at least one microRNA binding site is located in the 5' UTR 15-20 nucleotides preceding the start codon of the coding region of the mRNA. In some embodiments, at least one microRNA binding site is located in the 5' UTR 70-80 nucleotides preceding the start codon of the coding region of the mRNA.
- [0042] In some embodiments, the mRNA comprises multiple copies of the same microRNA binding site positioned immediately adjacent to each other or with a spacer of less than 5, 5-10, 10-15, or 15-20 nucleotides.
- [0043] In some embodiments, the mRNA comprises multiple copies of the same microRNA binding site located in the 3' UTR, wherein the first microRNA binding site is positioned immediately adjacent to the stop codon and the second and third microRNA binding sites are positioned 30-40 bases downstream of the 3' most residue of the first microRNA binding site.

- [0044] In some embodiments, the microRNA binding site comprises one or more nucleotide sequences selected from SEQ ID NO: 34 and SEQ ID NO: 36. In some embodiments, the miRNA binding site binds to miR-142. In some embodiments, the miRNA binding site binds to miR-142-3p or miR-142-5p. In some embodiments, the miR-142 comprises SEQ ID NO: 32.
- [0045] In some embodiments, the microRNA binding site comprises one or more nucleotide sequences selected from SEQ ID NO: 87 and SEQ ID NO: 89. In some embodiments, the miRNA binding site binds to miR-126. In some embodiments, the miRNA binding site binds to miR-126-3p or miR-126-5p. In some embodiments, the miR-126 comprises SEQ ID NO: 85.
- [0046] In some embodiments, the mRNA comprises a 3' UTR comprising a microRNA binding site that binds to miR-142, miR-126, or a combination thereof.
- [0047] In some embodiments, the polynucleotide, e.g., mRNA, further comprises a 3' UTR. In some embodiments, the miRNA binding site is located within the 3' UTR. In some embodiments, the 3' UTR comprises a nucleic acid sequence at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99%, or 100% identical to a 3'UTR sequence selected from the group consisting of SEQ ID NOs: 39, 58 to 84, 90, 105, 108 to 115, and 120 to 130, or any combination thereof. In some embodiments, the 3' UTR comprises a nucleic acid sequence selected from the group consisting of SEQ ID NOs: 39, 58 to 84, 90, 105, 108 to 115, and 120 to 130, and any combination thereof.
- [0048] In some embodiments, the polynucleotide, e.g., mRNA, further comprises a 5' UTR. In some embodiments, the 5' UTR comprises a nucleic acid sequence at least 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99%, or about 100% identical to a 5'UTR sequence selected from the group consisting of SEQ ID NO: 38, 40 to 57, and 117 to 119, or any combination thereof. In some embodiments, the 5' UTR comprises a sequence selected from the group consisting of SEQ ID NO: 38, 40 to 57, and 117 to 119, and any combination thereof.
- [0049] In some embodiments, the polynucleotide, e.g., mRNA, further comprises a 5' terminal cap. In some embodiments, the 5' terminal cap comprises a Cap0, Cap1, ARCA, inosine, N1-methyl-guanosine, 2'-fluoro-guanosine, 7-deaza-guanosine, 8-oxo-guanosine, 2-amino-guanosine, LNA-guanosine, 2-azidoguanosine, Cap2, Cap4, -11-emphalylG cap, or an analog thereof. In some embodiments, the 5' terminal cap comprises a Cap1.

[0050] In some embodiments, the polynucleotide, e.g., mRNA, further comprises a poly-A region. In some embodiments, the poly-A region is at least about 10, at least about 20, at least about 30, at least about 40, at least about 50, at least about 60, at least about 70, at least about 80, or at least about 90 nucleotides in length. In some embodiments, the poly-A region has about 10 to about 200, about 20 to about 180, about 50 to about 160, about 70 to about 140, about 80 to about 120 nucleotides in length.

[0051] In some embodiments, the polynucleotide, e.g., mRNA, encodes an ACADVL polypeptide that is fused to one or more heterologous polypeptides. In some embodiments, the one or more heterologous polypeptides increase a pharmacokinetic property of the ACADVL polypeptide. In some embodiments, upon administration to a subject, the polynucleotide has (i) a longer plasma half-life; (ii) increased expression of an ACADVL polypeptide encoded by the ORF; (iii) greater structural stability; or (iv) any combination thereof, relative to a corresponding polynucleotide comprising SEQ ID NO: 2, 4, or 6.


[0052] In some embodiments, the polynucleotide, e.g., mRNA, comprises (i) a 5'-terminal cap; (ii) a 5'-UTR; (iii) an ORF encoding an ACADVL polypeptide; (iv) a 3'-UTR; and (v) a poly-A region. In some embodiments, the 3'-UTR comprises a miRNA binding site. In some embodiments the polynucleotide further comprises a 5'-terminal cap (e.g., Cap1) and a poly-A-tail region (e.g., about 100 nucleotides in length).

[0053] The present disclosure also provides a method of producing a polynucleotide, e.g., mRNA, of the present invention, the method comprising modifying an ORF encoding an ACADVL polypeptide by substituting at least one uracil nucleobase with an adenine, guanine, or cytosine nucleobase, or by substituting at least one adenine, guanine, or cytosine nucleobase with a uracil nucleobase, wherein all the substitutions are synonymous substitutions. In some embodiments, the method further comprises replacing at least about 90%, at least about 95%, at least about 99%, or about 100% of uracils with 5-methoxyuracils.

[0054] The present disclosure also provides a composition comprising (a) a polynucleotide, e.g., mRNA, of the invention; and (b) a delivery agent. In some embodiments, the delivery agent comprises a lipidoid, a liposome, a lipoplex, a lipid nanoparticle, a polymeric compound, a peptide, a protein, a cell, a nanoparticle mimic, a nanotube, or a conjugate. In some embodiments, the delivery agent comprises a lipid nanoparticle. In some embodiments, the lipid nanoparticle comprises a lipid selected from

the group consisting of 3-(didodecylamino)-N1,N1,4-tridodecyl-1-piperazineethanamine (KL10), N1-[2-(didodecylamino)ethyl]-N1,N4,N4-tridodecyl-1,4-piperazinediethanamine (KL22), 14,25-ditridecyl-15,18,21,24-tetraaza-octatriacontane (KL25), 1,2-dilinoleyloxy-N,N-dimethylaminopropane (DLin-DMA), 2,2-dilinoley1-4-dimethylaminomethyl-[1,3]-dioxolane (DLin-K-DMA), heptatriaconta-6,9,28,31-tetraen-19-yl 4-(dimethylamino)butanoate (DLin-MC3-DMA), 2,2-dilinoley1-4-(2-dimethylaminoethyl)-[1,3]-dioxolane (DLin-KC2-DMA), 1,2-dioleyloxy-N,N-dimethylaminopropane (DODMA), (13Z,165Z)-N,N-dimethyl-3-nonydocosa-13-16-dien-1-amine (L608), 2-({8-[(3 β)-cholest-5-en-3-yloxy]octyl}oxy)-N,N-dimethyl-3-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]propan-1-amine (Octyl-CLinDMA), (2R)-2-({8-[(3 β)-cholest-5-en-3-yloxy]octyl}oxy)-N,N-dimethyl-3-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]propan-1-amine (Octyl-CLinDMA (2R)), (2S)-2-({8-[(3 β)-cholest-5-en-3-yloxy]octyl}oxy)-N,N-dimethyl-3-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]propan-1-amine (Octyl-CLinDMA (2S)), and any combinations thereof. In some embodiments, the lipid nanoparticle comprises DLin-MC3-DMA.

[0055] In certain embodiments, the delivery agent comprises a compound having the Formula (I)



or a salt or stereoisomer thereof, wherein

R₁ is selected from the group consisting of C₅₋₃₀ alkyl, C₅₋₂₀ alkenyl, -R*YR'', -YR'', and -R''M'R';

R₂ and R₃ are independently selected from the group consisting of H, C₁₋₁₄ alkyl, C₂₋₁₄ alkenyl, -R*YR'', -YR'', and -R*OR'', or R₂ and R₃, together with the atom to which they are attached, form a heterocycle or carbocycle;

R₄ is selected from the group consisting of a C₃₋₆ carbocycle, -(CH₂)_nQ, -(CH₂)_nCHQR, -CHQR, -CQ(R)₂, and unsubstituted C₁₋₆ alkyl, where Q is selected from a carbocycle, heterocycle, -OR, -O(CH₂)_nN(R)₂, -C(O)OR, -OC(O)R, -CX₃, -CX₂H, -CXH₂, -CN, -N(R)₂, -C(O)N(R)₂, -N(R)C(O)R, -N(R)S(O)₂R, -N(R)C(O)N(R)₂, -N(R)C(S)N(R)₂, -N(R)R₈, -O(CH₂)_nOR, -N(R)C(=NR₉)N(R)₂, -N(R)C(=CHR₉)N(R)₂,

-OC(O)N(R)₂, -N(R)C(O)OR, -N(OR)C(O)R, -N(OR)S(O)₂R, -N(OR)C(O)OR, -N(OR)C(O)N(R)₂, -N(OR)C(S)N(R)₂, -N(OR)C(=NR₉)N(R)₂, -N(OR)C(=CHR₉)N(R)₂, -C(=NR₉)N(R)₂, -C(=NR₉)R, -C(O)N(R)OR, and -C(R)N(R)₂C(O)OR, and each n is independently selected from 1, 2, 3, 4, and 5;

each R₅ is independently selected from the group consisting of C₁₋₃ alkyl, C₂₋₃ alkenyl, and H;

each R₆ is independently selected from the group consisting of C₁₋₃ alkyl, C₂₋₃ alkenyl, and H;

M and M' are independently selected from -C(O)O-, -OC(O)-, -C(O)N(R')-, -N(R')C(O)-, -C(O)-, -C(S)-, -C(S)S-, -SC(S)-, -CH(OH)-, -P(O)(OR')O-, -S(O)₂-, -S-S-, an aryl group, and a heteroaryl group;

R₇ is selected from the group consisting of C₁₋₃ alkyl, C₂₋₃ alkenyl, and H;

R₈ is selected from the group consisting of C₃₋₆ carbocycle and heterocycle;

R₉ is selected from the group consisting of H, CN, NO₂, C₁₋₆ alkyl, -OR, -S(O)₂R, -S(O)₂N(R)₂, C₂₋₆ alkenyl, C₃₋₆ carbocycle and heterocycle;

each R is independently selected from the group consisting of C₁₋₃ alkyl, C₂₋₃ alkenyl, and H;

each R' is independently selected from the group consisting of C₁₋₁₈ alkyl, C₂₋₁₈ alkenyl, -R*YR'', -YR'', and H;

each R'' is independently selected from the group consisting of C₃₋₁₄ alkyl and C₃₋₁₄ alkenyl;

each R* is independently selected from the group consisting of C₁₋₁₂ alkyl and C₂₋₁₂ alkenyl;


each Y is independently a C₃₋₆ carbocycle;

each X is independently selected from the group consisting of F, Cl, Br, and I; and

m is selected from 5, 6, 7, 8, 9, 10, 11, 12, and 13; and

provided when R₄ is -(CH₂)_nQ, -(CH₂)_nCHQR, -CHQR, or -CQ(R)₂, then (i) Q is not -N(R)₂ when n is 1, 2, 3, 4 or 5, or (ii) Q is not 5, 6, or 7-membered heterocycloalkyl when n is 1 or 2.

[0056] In certain aspects, the invention relates to a composition comprising a nucleotide sequence encoding an ACADVL polypeptide and a delivery agent, wherein the delivery agent comprises a compound having the Formula (I)



(I),

or a salt or stereoisomer thereof, wherein

R_1 is selected from the group consisting of C_{5-30} alkyl, C_{5-20} alkenyl, $-R^*YR''$, $-YR''$, and $-R''M'R'$;

R_2 and R_3 are independently selected from the group consisting of H, C_{1-14} alkyl, C_{2-14} alkenyl, $-R^*YR''$, $-YR''$, and $-R^*OR''$, or R_2 and R_3 , together with the atom to which they are attached, form a heterocycle or carbocycle;

R_4 is selected from the group consisting of a C_{3-6} carbocycle, $-(CH_2)_nQ$, $-(CH_2)_nCHQR$, $-CHQR$, $-CQ(R)_2$, and unsubstituted C_{1-6} alkyl, where Q is selected from a carbocycle, heterocycle, $-OR$, $-O(CH_2)_nN(R)_2$, $-C(O)OR$, $-OC(O)R$, $-CX_3$, $-CX_2H$, $-CXH_2$, $-CN$, $-N(R)_2$, $-C(O)N(R)_2$, $-N(R)C(O)R$, $-N(R)S(O)_2R$, $-N(R)C(O)N(R)_2$, $-N(R)C(S)N(R)_2$, $-N(R)R_8$, $-O(CH_2)_nOR$, $-N(R)C(=NR_9)N(R)_2$, $-N(R)C(=CHR_9)N(R)_2$, $-OC(O)N(R)_2$, $-N(R)C(O)OR$, $-N(OR)C(O)R$, $-N(OR)S(O)_2R$, $-N(OR)C(O)OR$, $-N(OR)C(O)N(R)_2$, $-N(OR)C(S)N(R)_2$, $-N(OR)C(=NR_9)N(R)_2$, $-N(OR)C(=CHR_9)N(R)_2$, $-C(=NR_9)N(R)_2$, $-C(=NR_9)R$, $-C(O)N(R)OR$, and $-C(R)N(R)_2C(O)OR$, and each n is independently selected from 1, 2, 3, 4, and 5;

each R_5 is independently selected from the group consisting of C_{1-3} alkyl, C_{2-3} alkenyl, and H;

each R_6 is independently selected from the group consisting of C_{1-3} alkyl, C_{2-3} alkenyl, and H;

M and M' are independently selected from $-C(O)O-$, $-OC(O)-$, $-C(O)N(R')-$, $-N(R')C(O)-$, $-C(O)-$, $-C(S)-$, $-C(S)S-$, $-SC(S)-$, $-CH(OH)-$, $-P(O)(OR')O-$, $-S(O)_2-$, $-S-S-$, an aryl group, and a heteroaryl group;

R_7 is selected from the group consisting of C_{1-3} alkyl, C_{2-3} alkenyl, and H;

R_8 is selected from the group consisting of C_{3-6} carbocycle and heterocycle;

R_9 is selected from the group consisting of H, CN, NO_2 , C_{1-6} alkyl, $-OR$, $-S(O)_2R$, $-S(O)_2N(R)_2$, C_{2-6} alkenyl, C_{3-6} carbocycle and heterocycle;

each R is independently selected from the group consisting of C_{1-3} alkyl, C_{2-3} alkenyl, and H;

each R' is independently selected from the group consisting of C₁₋₁₈ alkyl, C₂₋₁₈ alkenyl, -R*YR'', -YR'', and H;

each R'' is independently selected from the group consisting of C₃₋₁₄ alkyl and C₃₋₁₄ alkenyl;

each R* is independently selected from the group consisting of C₁₋₁₂ alkyl and C₂₋₁₂ alkenyl;

each Y is independently a C₃₋₆ carbocycle;

each X is independently selected from the group consisting of F, Cl, Br, and I; and

m is selected from 5, 6, 7, 8, 9, 10, 11, 12, and 13; and

provided when R₄ is -(CH₂)_nQ, -(CH₂)_nCHQR, -CHQR, or -CQ(R)₂, then (i) Q is not -N(R)₂ when n is 1, 2, 3, 4 or 5, or (ii) Q is not 5, 6, or 7-membered heterocycloalkyl when n is 1 or 2.

[0057] In some embodiments, the delivery agent comprises a compound having the Formula (I), or a salt or stereoisomer thereof, wherein

R₁ is selected from the group consisting of C₅₋₂₀ alkyl, C₅₋₂₀ alkenyl, -R*YR'', -YR'', and -R''M'R';

R₂ and R₃ are independently selected from the group consisting of H, C₁₋₁₄ alkyl, C₂₋₁₄ alkenyl, -R*YR'', -YR'', and -R*OR'', or R₂ and R₃, together with the atom to which they are attached, form a heterocycle or carbocycle;

R₄ is selected from the group consisting of a C₃₋₆ carbocycle, -(CH₂)_nQ, -(CH₂)_nCHQR, -CHQR, -CQ(R)₂, and unsubstituted C₁₋₆ alkyl, where Q is selected from a carbocycle, heterocycle, -OR, -O(CH₂)_nN(R)₂, -C(O)OR, -OC(O)R, -CX₃, -CX₂H, -CXH₂, -CN, -N(R)₂, -C(O)N(R)₂, -N(R)C(O)R, -N(R)S(O)₂R, -N(R)C(O)N(R)₂, -N(R)C(S)N(R)₂, and -C(R)N(R)₂C(O)OR, and each n is independently selected from 1, 2, 3, 4, and 5;

each R₅ is independently selected from the group consisting of C₁₋₃ alkyl, C₂₋₃ alkenyl, and H;

each R₆ is independently selected from the group consisting of C₁₋₃ alkyl, C₂₋₃ alkenyl, and H;

M and M' are independently selected from -C(O)O-, -OC(O)-, -C(O)N(R')-, -N(R')C(O)-, -C(O)-, -C(S)-, -C(S)S-, -SC(S)-, -CH(OH)-, -P(O)(OR')O-, -S(O)₂-, an aryl group, and a heteroaryl group;

R₇ is selected from the group consisting of C₁₋₃ alkyl, C₂₋₃ alkenyl, and H;

each R is independently selected from the group consisting of C₁₋₃ alkyl, C₂₋₃ alkenyl, and H;

each R' is independently selected from the group consisting of C₁₋₁₈ alkyl, C₂₋₁₈ alkenyl, -R*YR'', -YR'', and H;

each R'' is independently selected from the group consisting of C₃₋₁₄ alkyl and C₃₋₁₄ alkenyl;


each R* is independently selected from the group consisting of C₁₋₁₂ alkyl and C₂₋₁₂ alkenyl;

each Y is independently a C₃₋₆ carbocycle;

each X is independently selected from the group consisting of F, Cl, Br, and I; and m is selected from 5, 6, 7, 8, 9, 10, 11, 12, and 13; and

provided when R₄ is -(CH₂)_nQ, -(CH₂)_nCHQR, -CHQR, or -CQ(R)₂, then (i) Q is not -N(R)₂ when n is 1, 2, 3, 4 or 5, or (ii) Q is not 5, 6, or 7-membered heterocycloalkyl when n is 1 or 2.

[0058] In certain embodiments, the compound is of Formula (IA):



(IA),

or a salt or stereoisomer thereof, wherein

l is selected from 1, 2, 3, 4, and 5;

m is selected from 5, 6, 7, 8, and 9;

M₁ is a bond or M';

R₄ is unsubstituted C₁₋₃ alkyl, or -(CH₂)_nQ, in which n is 1, 2, 3, 4, or 5 and Q is OH, -NHC(S)N(R)₂, -NHC(O)N(R)₂, -N(R)C(O)R, -N(R)S(O)₂R, -N(R)R₈, -NHC(=NR₉)N(R)₂, -NHC(=CHR₉)N(R)₂, -OC(O)N(R)₂, -N(R)C(O)OR, heteroaryl, or heterocycloalkyl;

M and M' are independently selected from -C(O)O-, -OC(O)-, -C(O)N(R')-, -P(O)(OR')O-, -S-S-, an aryl group, and a heteroaryl group; and

R₂ and R₃ are independently selected from the group consisting of H, C₁₋₁₄ alkyl, and C₂₋₁₄ alkenyl.

[0059] In certain embodiments, m is 5, 7, or 9.

[0060] In some embodiments, the compound is of Formula (IA), or a salt or stereoisomer thereof, wherein

l is selected from 1, 2, 3, 4, and 5;

m is selected from 5, 6, 7, 8, and 9;

M₁ is a bond or M';


R₄ is unsubstituted C₁₋₃ alkyl, or -(CH₂)_nQ, in which n is 1, 2, 3, 4, or 5 and Q is OH, -NHC(S)N(R)₂, or -NHC(O)N(R)₂;

M and M' are independently selected from -C(O)O-, -OC(O)-, -C(O)N(R')-, -P(O)(OR')O- an aryl group, and a heteroaryl group; and

R₂ and R₃ are independently selected from the group consisting of H, C₁₋₁₄ alkyl, and C₂₋₁₄ alkenyl.

[0061] In some embodiments, m is 5, 7, or 9.

[0062] In certain embodiments, the compound is of Formula (II):



(II),

or a salt or stereoisomer thereof, wherein

l is selected from 1, 2, 3, 4, and 5;

M₁ is a bond or M';

R₄ is unsubstituted C₁₋₃ alkyl, or -(CH₂)_nQ, in which n is 2, 3, or 4 and Q is OH, -NHC(S)N(R)₂, -NHC(O)N(R)₂, -N(R)C(O)R, -N(R)S(O)₂R, -N(R)R₈, -NHC(=NR₉)N(R)₂, -NHC(=CHR₉)N(R)₂, -OC(O)N(R)₂, -N(R)C(O)OR, heteroaryl, or heterocycloalkyl;

M and M' are independently selected from -C(O)O-, -OC(O)-, -C(O)N(R')-, -P(O)(OR')O-, -S-S-, an aryl group, and a heteroaryl group; and

R₂ and R₃ are independently selected from the group consisting of H, C₁₋₁₄ alkyl, and C₂₋₁₄ alkenyl.

[0063] In some embodiments, the compound is of Formula (II), or a salt or stereoisomer thereof, wherein

l is selected from 1, 2, 3, 4, and 5;

M₁ is a bond or M';

R_4 is unsubstituted C_{1-3} alkyl, or $-(CH_2)_nQ$, in which n is 2, 3, or 4 and Q is OH, $-NHC(S)N(R)_2$, or $-NHC(O)N(R)_2$;

M and M' are independently selected from $-C(O)O-$, $-OC(O)-$, $-C(O)N(R')$, $-P(O)(OR')O-$, an aryl group, and a heteroaryl group; and

R_2 and R_3 are independently selected from the group consisting of H, C_{1-14} alkyl, and C_{2-14} alkenyl.

[0064] In some embodiments, M_1 is M' .


[0065] In some embodiments, M and M' are independently $-C(O)O-$ or $-OC(O)-$.

[0066] In some embodiments, l is 1, 3, or 5.


[0067] In some embodiments, the compound is selected from the group consisting of Compound 1 to Compound 232, salts and stereoisomers thereof, and any combination thereof.

[0068] In some embodiments, the compound is selected from the group consisting of Compound 1 to Compound 147, salts and stereoisomers thereof, and any combination thereof.


[0069] In certain embodiments, the compound is of the Formula (IIa),




[0070] In certain embodiments, the compound is of the Formula (IIb),



[0071] In certain embodiments, the compound is of the Formula (IIc) or (IIe),






(Ile),

or a salt or stereoisomer thereof.

[0072] In certain embodiments, R_4 is as described herein. In some embodiments, R_4 is selected from $-(CH_2)_nQ$ and $-(CH_2)_nCHQR$.

[0073] In certain embodiments, the compound is of the Formula (IId),



(IId), or a salt or stereoisomer thereof,

wherein n is selected from 2, 3, and 4, and m , R' , R'' , and R_2 through R_6 are as described herein. For example, each of R_2 and R_3 may be independently selected from the group consisting of C_{5-14} alkyl and C_{5-14} alkenyl.

[0074] In some embodiments, the compound is of the Formula (IId), or a salt or stereoisomer thereof,

wherein R_2 and R_3 are independently selected from the group consisting of C_{5-14} alkyl and C_{5-14} alkenyl, n is selected from 2, 3, and 4, and R' , R'' , R_5 , R_6 and m are as defined herein.


[0075] In some embodiments, R_2 is C_8 alkyl.

[0076] In some embodiments, R_3 is C_5 alkyl, C_6 alkyl, C_7 alkyl, C_8 alkyl, or C_9 alkyl.

[0077] In some embodiments, m is 5, 7, or 9.


[0078] In some embodiments, each R_5 is H.

[0079] In some embodiments, each R_6 is H. In some embodiments, the delivery agent comprises a compound having the Formula (III)



(III),

or salts or stereoisomers thereof, wherein



t is 1 or 2;

A₁ and A₂ are each independently selected from CH or N;

Z is CH₂ or absent wherein when Z is CH₂, the dashed lines (1) and (2) each represent a single bond; and when Z is absent, the dashed lines (1) and (2) are both absent;

R₁, R₂, R₃, R₄, and R₅ are independently selected from the group consisting of C₅₋₂₀ alkyl, C₅₋₂₀ alkenyl, -R''MR', -R*YR'', -YR'', and -R*OR'';

each M is independently selected from the group consisting of -C(O)O-, -OC(O)-, -OC(O)O-, -C(O)N(R')-, -N(R')C(O)-, -C(O)-, -C(S)-, -C(S)S-, -SC(S)-, -CH(OH)-, -P(O)(OR')O-, -S(O)₂-, an aryl group, and a heteroaryl group;

X¹, X², and X³ are independently selected from the group consisting of a bond, -CH₂-, -(CH₂)₂-, -CHR-, -CHY-, -C(O)-, -C(O)O-, -OC(O)-, -C(O)-CH₂-, -CH₂-C(O)-, -C(O)O-CH₂-, -OC(O)-CH₂-, -CH₂-C(O)O-, -CH₂-OC(O)-, -CH(OH)-, -C(S)-, and -CH(SH)-;


each Y is independently a C₃₋₆ carbocycle;

each R* is independently selected from the group consisting of C₁₋₁₂ alkyl and C₂₋₁₂ alkenyl;

each R is independently selected from the group consisting of C₁₋₃ alkyl and a C₃₋₆ carbocycle;







each R' is independently selected from the group consisting of C₁₋₁₂ alkyl, C₂₋₁₂ alkenyl, and H; and

each R'' is independently selected from the group consisting of C₃₋₁₂ alkyl and C₃₋₁₂ alkenyl,




- i) at least one of X^1 , X^2 , and X^3 is not $-CH_2-$; and/or
 ii) at least one of R_1 , R_2 , R_3 , R_4 , and R_5 is $-R''MR'$.

[0080] In some embodiments, the compound is of any of Formulae (IIIa1)-(IIIa6):





[0081] The compounds of Formula (III) or any of (IIIa1)-(IIIa6) include one or more of the following features when applicable.


[0082] In some embodiments, ring A is




[0083] In some embodiments, ring A is






[0084] In some embodiments, ring A is





[0085] In some embodiments, ring A is



[0086] In some embodiments, ring A is



[0087] In some embodiments, ring A is




wherein ring, in which the N atom is connected with X^2 .

[0088] In some embodiments, Z is CH_2 .

[0089] In some embodiments, Z is absent.

[0090] In some embodiments, at least one of A_1 and A_2 is N.

- [0091] In some embodiments, each of A_1 and A_2 is N.
- [0092] In some embodiments, each of A_1 and A_2 is CH.
- [0093] In some embodiments, A_1 is N and A_2 is CH.
- [0094] In some embodiments, A_1 is CH and A_2 is N.
- [0095] In some embodiments, at least one of X^1 , X^2 , and X^3 is not $-\text{CH}_2-$. For example, in certain embodiments, X^1 is not $-\text{CH}_2-$. In some embodiments, at least one of X^1 , X^2 , and X^3 is $-\text{C}(\text{O})-$.
- [0096] In some embodiments, X^2 is $-\text{C}(\text{O})-$, $-\text{C}(\text{O})\text{O}-$, $-\text{OC}(\text{O})-$, $-\text{C}(\text{O})-\text{CH}_2-$, $-\text{CH}_2-\text{C}(\text{O})-$, $-\text{C}(\text{O})\text{O}-\text{CH}_2-$, $-\text{OC}(\text{O})-\text{CH}_2-$, $-\text{CH}_2-\text{C}(\text{O})\text{O}-$, or $-\text{CH}_2-\text{OC}(\text{O})-$. In some embodiments, X^3 is $-\text{C}(\text{O})-$, $-\text{C}(\text{O})\text{O}-$, $-\text{OC}(\text{O})-$, $-\text{C}(\text{O})-\text{CH}_2-$, $-\text{CH}_2-\text{C}(\text{O})-$, $-\text{C}(\text{O})\text{O}-\text{CH}_2-$, $-\text{OC}(\text{O})-\text{CH}_2-$, $-\text{CH}_2-\text{C}(\text{O})\text{O}-$, or $-\text{CH}_2-\text{OC}(\text{O})-$. In other embodiments, X^3 is $-\text{CH}_2-$.
- [0097] In some embodiments, X^3 is a bond or $-(\text{CH}_2)_2-$.
- [0098] In some embodiments, R_1 and R_2 are the same. In certain embodiments, R_1 , R_2 , and R_3 are the same. In some embodiments, R_4 and R_5 are the same. In certain embodiments, R_1 , R_2 , R_3 , R_4 , and R_5 are the same.
- [0099] In some embodiments, at least one of R_1 , R_2 , R_3 , R_4 , and R_5 is $-\text{R}''\text{MR}'$. In some embodiments, at most one of R_1 , R_2 , R_3 , R_4 , and R_5 is $-\text{R}''\text{MR}'$. For example, at least one of R_1 , R_2 , and R_3 may be $-\text{R}''\text{MR}'$, and/or at least one of R_4 and R_5 is $-\text{R}''\text{MR}'$. In certain embodiments, at least one M is $-\text{C}(\text{O})\text{O}-$. In some embodiments, each M is $-\text{C}(\text{O})\text{O}-$. In some embodiments, at least one M is $-\text{OC}(\text{O})-$. In some embodiments, each M is $-\text{OC}(\text{O})-$. In some embodiments, at least one M is $-\text{OC}(\text{O})\text{O}-$. In some embodiments, each M is $-\text{OC}(\text{O})\text{O}-$. In some embodiments, at least one R'' is C_3 alkyl. In certain embodiments, each R'' is C_3 alkyl. In some embodiments, at least one R'' is C_5 alkyl. In certain embodiments, each R'' is C_5 alkyl. In some embodiments, at least one R'' is C_6 alkyl. In certain embodiments, each R'' is C_6 alkyl. In some embodiments, at least one R'' is C_7 alkyl. In certain embodiments, each R'' is C_7 alkyl. In some embodiments, at least one R' is C_5 alkyl. In certain embodiments, each R' is C_5 alkyl. In other embodiments, at least one R' is C_1 alkyl. In certain embodiments, each R' is C_1 alkyl. In some embodiments, at least one R' is C_2 alkyl. In certain embodiments, each R' is C_2 alkyl.
- [0100] In some embodiments, at least one of R_1 , R_2 , R_3 , R_4 , and R_5 is C_{12} alkyl. In certain embodiments, each of R_1 , R_2 , R_3 , R_4 , and R_5 are C_{12} alkyl.
- [0101] In some embodiments, the delivery agent comprises a compound having the Formula (IV)




(IV),


or salts or stereoisomer thereof, wherein

A_1 and A_2 are each independently selected from CH or N and at least one of A_1 and A_2 is N;

Z is CH_2 or absent wherein when Z is CH_2 , the dashed lines (1) and (2) each represent a single bond; and when Z is absent, the dashed lines (1) and (2) are both absent;

R_1 , R_2 , R_3 , R_4 , and R_5 are independently selected from the group consisting of C_{6-20} alkyl and C_{6-20} alkenyl;



wherein when ring A is , then

i) R_1 , R_2 , R_3 , R_4 , and R_5 are the same, wherein R_1 is not C_{12} alkyl, C_{18} alkyl, or C_{18} alkenyl;


ii) only one of R_1 , R_2 , R_3 , R_4 , and R_5 is selected from C_{6-20} alkenyl;

iii) at least one of R_1 , R_2 , R_3 , R_4 , and R_5 have a different number of carbon atoms than at least one other of R_1 , R_2 , R_3 , R_4 , and R_5 ;

iv) R_1 , R_2 , and R_3 are selected from C_{6-20} alkenyl, and R_4 and R_5 are selected from C_{6-20} alkyl; or

v) R_1 , R_2 , and R_3 are selected from C_{6-20} alkyl, and R_4 and R_5 are selected from C_{6-20} alkenyl.

[0102] In some embodiments, the compound is of Formula (IVa):




(IVa).

- [0103] The compounds of Formula (IV) or (IVa) include one or more of the following features when applicable.
- [0104] In some embodiments, Z is CH₂.
- [0105] In some embodiments, Z is absent.
- [0106] In some embodiments, at least one of A₁ and A₂ is N.
- [0107] In some embodiments, each of A₁ and A₂ is N.
- [0108] In some embodiments, each of A₁ and A₂ is CH.
- [0109] In some embodiments, A₁ is N and A₂ is CH.
- [0110] In some embodiments, A₁ is CH and A₂ is N.
- [0111] In some embodiments, R₁, R₂, R₃, R₄, and R₅ are the same, and are not C₁₂ alkyl, C₁₈ alkyl, or C₁₈ alkenyl. In some embodiments, R₁, R₂, R₃, R₄, and R₅ are the same and are C₉ alkyl or C₁₄ alkyl.
- [0112] In some embodiments, only one of R₁, R₂, R₃, R₄, and R₅ is selected from C₆₋₂₀ alkenyl. In certain such embodiments, R₁, R₂, R₃, R₄, and R₅ have the same number of carbon atoms. In some embodiments, R₄ is selected from C₅₋₂₀ alkenyl. For example, R₄ may be C₁₂ alkenyl or C₁₈ alkenyl.
- [0113] In some embodiments, at least one of R₁, R₂, R₃, R₄, and R₅ have a different number of carbon atoms than at least one other of R₁, R₂, R₃, R₄, and R₅.
- [0114] In certain embodiments, R₁, R₂, and R₃ are selected from C₆₋₂₀ alkenyl, and R₄ and R₅ are selected from C₆₋₂₀ alkyl. In other embodiments, R₁, R₂, and R₃ are selected from C₆₋₂₀ alkyl, and R₄ and R₅ are selected from C₆₋₂₀ alkenyl. In some embodiments, R₁, R₂, and R₃ have the same number of carbon atoms, and/or R₄ and R₅ have the same number of carbon atoms. For example, R₁, R₂, and R₃, or R₄ and R₅, may have 6, 8, 9, 12, 14, or 18 carbon atoms. In some embodiments, R₁, R₂, and R₃, or R₄ and R₅, are C₁₈ alkenyl (e.g., linoleyl). In some embodiments, R₁, R₂, and R₃, or R₄ and R₅, are alkyl groups including 6, 8, 9, 12, or 14 carbon atoms.

[0115] In some embodiments, R_1 has a different number of carbon atoms than R_2 , R_3 , R_4 , and R_5 . In other embodiments, R_3 has a different number of carbon atoms than R_1 , R_2 , R_4 , and R_5 . In further embodiments, R_4 has a different number of carbon atoms than R_1 , R_2 , R_3 , and R_5 .

[0116] In other embodiments, the delivery agent comprises a compound having the Formula (V)



(V),

or salts or stereoisomers thereof, in which

A_3 is CH or N;

A_4 is CH_2 or NH; and at least one of A_3 and A_4 is N or NH;

Z is CH_2 or absent wherein when Z is CH_2 , the dashed lines (1) and (2) each represent a single bond; and when Z is absent, the dashed lines (1) and (2) are both absent;

R_1 , R_2 , and R_3 are independently selected from the group consisting of C_{5-20} alkyl, C_{5-20} alkenyl, $-R''MR'$, $-R^*YR''$, $-YR''$, and $-R^*OR''$;

each M is independently selected from $-C(O)O-$, $-OC(O)-$, $-C(O)N(R')$, $-N(R')C(O)-$, $-C(O)-$, $-C(S)-$, $-C(S)S-$, $-SC(S)-$, $-CH(OH)-$, $-P(O)(OR')O-$, $-S(O)_2-$, an aryl group, and a heteroaryl group;

X^1 and X^2 are independently selected from the group consisting of $-CH_2-$, $-(CH_2)_2-$, $-CHR-$, $-CHY-$, $-C(O)-$, $-C(O)O-$, $-OC(O)-$, $-C(O)-CH_2-$, $-CH_2-C(O)-$, $-C(O)O-CH_2-$, $-OC(O)-CH_2-$, $-CH_2-C(O)O-$, $-CH_2-OC(O)-$, $-CH(OH)-$, $-C(S)-$, and $-CH(SH)-$;

each Y is independently a C_{3-6} carbocycle;


each R^* is independently selected from the group consisting of C_{1-12} alkyl and C_{2-12} alkenyl;

each R is independently selected from the group consisting of C_{1-3} alkyl and a C_{3-6} carbocycle;

each R' is independently selected from the group consisting of C_{1-12} alkyl, C_{2-12} alkenyl, and H; and

each R'' is independently selected from the group consisting of C₃₋₁₂ alkyl and C₃₋₁₂ alkenyl.

[0117] In some embodiments, the compound is of Formula (Va):



[0118] The compounds of Formula (V) or (Va) include one or more of the following features when applicable.

[0119] In some embodiments, Z is CH₂.

[0120] In some embodiments, Z is absent.

[0121] In some embodiments, at least one of A₃ and A₄ is N or NH.

[0122] In some embodiments, A₃ is N and A₄ is NH.

[0123] In some embodiments, A₃ is N and A₄ is CH₂.


[0124] In some embodiments, A₃ is CH and A₄ is NH.

[0125] In some embodiments, at least one of X¹ and X² is not -CH₂-. For example, in certain embodiments, X¹ is not -CH₂-. In some embodiments, at least one of X¹ and X² is -C(O)-.

[0126] In some embodiments, X² is -C(O)-, -C(O)O-, -OC(O)-, -C(O)-CH₂-, -CH₂-C(O)-, -C(O)O-CH₂-, -OC(O)-CH₂-, -CH₂-C(O)O-, or -CH₂-OC(O)-.

[0127] In some embodiments, R₁, R₂, and R₃ are independently selected from the group consisting of C₅₋₂₀ alkyl and C₅₋₂₀ alkenyl. In some embodiments, R₁, R₂, and R₃ are the same. In certain embodiments, R₁, R₂, and R₃ are C₆, C₉, C₁₂, or C₁₄ alkyl. In other embodiments, R₁, R₂, and R₃ are C₁₈ alkenyl. For example, R₁, R₂, and R₃ may be linoleyl.

[0128] In other embodiments, the delivery agent comprises a compound having the Formula (VI):



or salts or stereoisomers thereof, in which

A₆ and A₇ are each independently selected from CH or N, wherein at least one of A₆ and A₇ is N;

Z is CH₂ or absent wherein when Z is CH₂, the dashed lines (1) and (2) each represent a single bond; and when Z is absent, the dashed lines (1) and (2) are both absent;

X⁴ and X⁵ are independently selected from the group consisting of -CH₂-, -(CH₂)₂-, -CHR-, -CHY-, -C(O)-, -C(O)O-, -OC(O)-, -C(O)-CH₂-, -CH₂-C(O)-, -C(O)O-CH₂-, -OC(O)-CH₂-, -CH₂-C(O)O-, -CH₂-OC(O)-, -CH(OH)-, -C(S)-, and -CH(SH)-;

R₁, R₂, R₃, R₄, and R₅ each are independently selected from the group consisting of C₅₋₂₀ alkyl, C₅₋₂₀ alkenyl, -R''MR', -R*YR'', -YR'', and -R*OR'';

each M is independently selected from the group consisting of -C(O)O-, -OC(O)-, -C(O)N(R')-, -N(R')C(O)-, -C(O)-, -C(S)-, -C(S)S-, -SC(S)-, -CH(OH)-, -P(O)(OR')O-, -S(O)₂-, an aryl group, and a heteroaryl group;

each Y is independently a C₃₋₆ carbocycle;

each R* is independently selected from the group consisting of C₁₋₁₂ alkyl and C₂₋₁₂ alkenyl;

each R is independently selected from the group consisting of C₁₋₃ alkyl and a C₃₋₆ carbocycle;

each R' is independently selected from the group consisting of C₁₋₁₂ alkyl, C₂₋₁₂ alkenyl, and H; and

each R'' is independently selected from the group consisting of C₃₋₁₂ alkyl and C₃₋₁₂ alkenyl.

[0129] In some embodiments, R₁, R₂, R₃, R₄, and R₅ each are independently selected from the group consisting of C₆₋₂₀ alkyl and C₆₋₂₀ alkenyl.

[0130] In some embodiments, R₁ and R₂ are the same. In certain embodiments, R₁, R₂, and R₃ are the same. In some embodiments, R₄ and R₅ are the same. In certain embodiments, R₁, R₂, R₃, R₄, and R₅ are the same.

[0131] In some embodiments, at least one of R₁, R₂, R₃, R₄, and R₅ is C₉₋₁₂ alkyl. In certain embodiments, each of R₁, R₂, R₃, R₄, and R₅ independently is C₉, C₁₂ or C₁₄ alkyl. In certain embodiments, each of R₁, R₂, R₃, R₄, and R₅ is C₉ alkyl.


[0132] In some embodiments, A₆ is N and A₇ is N. In some embodiments, A₆ is CH and A₇ is N.

- [0133] In some embodiments, X^4 is $-CH_2-$ and X^5 is $-C(O)-$. In some embodiments, X^4 and X^5 are $-C(O)-$.
- [0134] In some embodiments, when A_6 is N and A_7 is N, at least one of X^4 and X^5 is not $-CH_2-$, e.g., at least one of X^4 and X^5 is $-C(O)-$. In some embodiments, when A_6 is N and A_7 is N, at least one of R_1 , R_2 , R_3 , R_4 , and R_5 is $-R''MR'$.
- [0135] In some embodiments, at least one of R_1 , R_2 , R_3 , R_4 , and R_5 is not $-R''MR'$.
- [0136] In certain embodiments, the composition is a nanoparticle composition.
- [0137] In certain embodiments, the delivery agent further comprises a phospholipid.
- [0138] In certain embodiments, the phospholipid is selected from the group consisting of
 1,2-dilinoleoyl-sn-glycero-3-phosphocholine (DLPC),
 1,2-dimyristoyl-sn-glycero-phosphocholine (DMPC),
 1,2-dioleoyl-sn-glycero-3-phosphocholine (DOPC),
 1,2-dipalmitoyl-sn-glycero-3-phosphocholine (DPPC),
 1,2-distearoyl-sn-glycero-3-phosphocholine (DSPC),
 1,2-diundecanoyl-sn-glycero-phosphocholine (DUPC),
 1-palmitoyl-2-oleoyl-sn-glycero-3-phosphocholine (POPC),
 1,2-di-O-octadecenyl-sn-glycero-3-phosphocholine (18:0 Diether PC),
 1-oleoyl-2-cholesterylhemisuccinoyl-sn-glycero-3-phosphocholine (OChemPC),
 1-hexadecyl-sn-glycero-3-phosphocholine (C16 Lyso PC),
 1,2-dilinenoyl-sn-glycero-3-phosphocholine,
 1,2-diarachidonoyl-sn-glycero-3-phosphocholine,
 1,2-didocosaheptaenoyl-sn-glycero-3-phosphocholine,
 1,2-dioleoyl-sn-glycero-3-phosphoethanolamine (DOPE),
 1,2-diphytanoyl-sn-glycero-3-phosphoethanolamine (ME 16:0 PE),
 1,2-distearoyl-sn-glycero-3-phosphoethanolamine,
 1,2-dilinoleoyl-sn-glycero-3-phosphoethanolamine,
 1,2-dilinenoyl-sn-glycero-3-phosphoethanolamine,
 1,2-diarachidonoyl-sn-glycero-3-phosphoethanolamine,
 1,2-didocosaheptaenoyl-sn-glycero-3-phosphoethanolamine,
 1,2-dioleoyl-sn-glycero-3-phospho-rac-(1-glycerol) sodium salt (DOPG), sphingomyelin,
 and any mixtures thereof.
- [0139] In certain embodiments, the delivery agent further comprises a structural lipid.

[0140] In certain embodiments, the structural lipid is selected from the group consisting of cholesterol, fecosterol, sitosterol, ergosterol, campesterol, stigmasterol, brassicasterol, tomatidine, ursolic acid, alpha-tocopherol, and any mixtures thereof.

[0141] In certain embodiments, the delivery agent further comprises a PEG lipid.

[0142] In certain embodiments, the PEG lipid is selected from the group consisting of a PEG-modified phosphatidylethanolamine, a PEG-modified phosphatidic acid, a PEG-modified ceramide, a PEG-modified dialkylamine, a PEG-modified diacylglycerol, a PEG-modified dialkylglycerol, and any mixtures thereof. In some embodiments, the PEG lipid has the formula:



[0143] In certain embodiments, the delivery agent further comprises an ionizable lipid selected from the group consisting of 3-(didodecylamino)-N1,N1,4-tridodecyl-1-piperazineethanamine (KL10), N1-[2-(didodecylamino)ethyl]-N1,N4,N4-tridodecyl-1,4-piperazinediethanamine (KL22), 14,25-ditridecyl-15,18,21,24-tetraaza-octatriacontane (KL25), 1,2-dilinoleyloxy-N,N-dimethylaminopropane (DLin-DMA), 2,2-dilinoleyl-4-dimethylaminomethyl-[1,3]-dioxolane (DLin-K-DMA), heptatriaconta-6,9,28,31-tetraen-19-yl 4-(dimethylamino)butanoate (DLin-MC3-DMA), 2,2-dilinoleyl-4-(2-dimethylaminoethyl)-[1,3]-dioxolane (DLin-KC2-DMA), 1,2-dioleyloxy-N,N-dimethylaminopropane (DODMA), 2-({8-[(3β)-cholest-5-en-3-yloxy]octyl}oxy)-N,N-dimethyl-3-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]propan-1-amine (Octyl-CLinDMA), (2R)-2-({8-[(3β)-cholest-5-en-3-yloxy]octyl}oxy)-N,N-dimethyl-3-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]propan-1-amine (Octyl-CLinDMA (2R)), and (2S)-2-({8-[(3β)-cholest-5-en-3-yloxy]octyl}oxy)-N,N-dimethyl-3-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]propan-1-amine (Octyl-CLinDMA (2S)).

[0144] In certain embodiments, the delivery agent further comprises a phospholipid, a structural lipid, a PEG lipid, or any combination thereof. In some embodiments, the delivery agent comprises Compound 18, DSPC, Cholesterol, and Compound 428, e.g., with a mole ratio of about 50:10:38.5:1.5.

[0145] In certain embodiments, the composition is formulated for *in vivo* delivery.

- [0146] In certain embodiments, the composition is formulated for intramuscular, subcutaneous, or intradermal delivery.
- [0147] The present disclosure further provides a polynucleotide comprising an mRNA comprising: (i) a 5' UTR, (ii) an open reading frame (ORF) encoding a human ACADVL polypeptide, wherein the ORF comprises a sequence optimized nucleic acid sequence encoding ACADVL disclosed herein, and (iii) a 3' UTR comprising a microRNA binding site selected from miR-142, miR-126, or a combination thereof, wherein the mRNA comprises at least one chemically modified nucleobase.
- [0148] The present disclosure further provides a polynucleotide comprising an mRNA comprising: (i) a 5'-terminal cap; (ii) a 5' UTR comprising a sequence selected from the group consisting of SEQ ID NO: 38, 40 to 57, and 117 to 119, and any combination thereof; (iii) an open reading frame (ORF) encoding a human ACADVL polypeptide, wherein the ORF comprises a sequence optimized nucleic acid sequence encoding ACADVL disclosed herein (e.g., a sequence selected from the group consisting of SEQ ID NOs: 7 to 31), wherein the mRNA comprises at least one chemically modified nucleobase selected from the group consisting of pseudouracil (ψ), N1-methylpseudouracil (m1 ψ), 1-ethylpseudouracil, 2-thiouracil (s2U), 4'-thiouracil, 5-methylcytosine, 5-methyluracil, 5-methoxyuracil, and any combination thereof; and (iv) a 3' UTR comprising a nucleic acid sequence selected from the group consisting of SEQ ID NOs: 39, 58 to 84, 90, 105, 108 to 115, and 120 to 130, and any combination thereof; and (v) a poly-A-region.
- [0149] The present disclosure further provides a pharmaceutical composition comprising the polynucleotide, e.g., an mRNA, and a delivery agent. In some embodiments, the delivery agent is a lipid nanoparticle comprising Compound 18, Compound 236, a salt or a stereoisomer thereof, or any combination thereof. In some embodiments, the polynucleotide comprising a nucleotide sequence encoding a ACADVL polypeptide disclosed herein is formulated with a delivery agent comprising, e.g., a compound having the Formula (I), e.g., any of Compounds 1-232, e.g., Compound 18; a compound having the Formula (III), (IV), (V), or (VI), e.g., any of Compounds 233-342, e.g., Compound 236; or a compound having the Formula (VIII), e.g., any of Compounds 419-428, e.g., Compound 428, or any combination thereof. In some embodiments, the delivery agent comprises Compound 18, DSPC, Cholesterol, and Compound 428, e.g., with a mole ratio of about 50:10:38.5:1.5.

- [0150] In one aspect of the embodiments disclosed herein, the subject is a human subject in need of treatment or prophylaxis for VLCADD.
- [0151] In one aspect of the embodiments disclosed herein, upon administration to the subject, the mRNA has: (i) a longer plasma half-life; (ii) increased expression of an ACADVL polypeptide encoded by the ORF; (iii) a lower frequency of arrested translation resulting in an expression fragment; (iv) greater structural stability; or (v) any combination thereof, relative to a corresponding mRNA having the nucleic acid sequence of SEQ ID NO: 2 and/or administered as naked mRNA.
- [0152] In some embodiments, a pharmaceutical composition or polynucleotide, e.g., an mRNA, disclosed herein is suitable for administration as a single unit dose or a plurality of single unit doses.
- [0153] In some embodiments, a pharmaceutical composition or polynucleotide, e.g., an mRNA, disclosed herein is suitable for reducing the level of one or more biomarkers of VLCADD in the subject.
- [0154] In some embodiments, a pharmaceutical composition or polynucleotide, e.g., an mRNA, disclosed herein is for use in treating or preventing the signs and/or symptoms of VLCADD in a subject in need thereof. In some embodiments, the signs or symptoms include hypertrophic or dilated cardiomyopathy, pericardial effusion, arrhythmias, hypotonia, hepatomegaly, and intermittent hypoglycemia, or a combination thereof.
- [0155] In certain aspects, the invention relates to a host cell comprising the polynucleotide, e.g., an mRNA. In certain embodiments, the host cell is a eukaryotic cell. In certain aspects, the invention relates to a vector comprising the polynucleotide. In certain aspects, the invention relates to a method of making a polynucleotide comprising enzymatically or chemically synthesizing the polynucleotide. In certain aspects, the invention relates to a polypeptide encoded by the polynucleotide, the composition, the host cell, or the vector or produced by the method of making the polynucleotide. In certain aspects, the invention relates to a method of expressing *in vivo* an active ACADVL polypeptide in a subject in need thereof comprising administering to the subject an effective amount of the polynucleotide, the composition, the host cell, or the vector. In certain aspects, the invention relates to a method of treating or preventing VLCADD in a subject in need thereof comprising administering to the subject a therapeutically effective amount of the polynucleotide, the composition, the host cell, or

the vector, wherein the administration alleviates the signs or symptoms of VLCADD in the subject.

- [0156] The present disclosure further provides a polynucleotide comprising an mRNA comprising: (i) a 5' UTR, (ii) an open reading frame (ORF) encoding a human ACADVL polypeptide (e.g., wherein the ORF comprises a nucleic acid sequence selected from the group consisting of SEQ ID NOs: 7 to 31), and (iii) a 3' UTR comprising a microRNA binding site selected from miR-142, miR-126, or a combination thereof, wherein the mRNA comprises at least one chemically modified nucleobase.
- [0157] The present disclosure further provides a polynucleotide comprising an mRNA comprising: (i) a 5'-terminal cap; (ii) a 5' UTR comprising a sequence selected from the group consisting of SEQ ID NO: 38, 40 to 57, and 117 to 119, and any combination thereof; (iii) an open reading frame (ORF) comprising a sequence optimized nucleic acid sequence encoding ACADVL disclosed herein (e.g., a sequence selected from the group consisting of SEQ ID NOs: 7 to 31), wherein the mRNA comprises at least one chemically modified nucleobase selected from the group consisting of pseudouracil (ψ), N1-methylpseudouracil (m1 ψ), 1-ethylpseudouracil, 2-thiouracil (s2U), 4'-thiouracil, 5-methylcytosine, 5-methyluracil, 5-methoxyuracil, and any combination thereof; and (iv) a 3' UTR comprising a nucleic acid sequence selected from the group consisting of SEQ ID NOs: 39, 58 to 84, 90, 105, 108 to 115, and 120 to 130, and any combination thereof; and (v) a poly-A-region.
- [0158] The present disclosure further provides a pharmaceutical composition comprising the polynucleotide and a delivery agent. In some embodiments, the delivery agent is a lipid nanoparticle comprising a Compound selected from the group consisting of any one of Compounds 1-342, a salt or a stereoisomer thereof, or any combination thereof. In some embodiments, the delivery agent is a lipid nanoparticle comprising Compound 18, Compound 236, a salt or a stereoisomer thereof, or any combination thereof. In some embodiments, the lipid nanoparticle or the delivery agent comprises Compound 18, DSPC, Cholesterol, and Compound 428, e.g., with a mole ratio of about 50:10:38.5:1.5.
- [0159] In one aspect of the embodiments disclosed herein, the subject is a human subject in need of treatment or prophylaxis for VLCADD.
- [0160] In some embodiments, a pharmaceutical composition or polynucleotide disclosed herein is suitable for administration as a single unit dose or a plurality of single unit doses.

- [0161] In some embodiments, a pharmaceutical composition or polynucleotide disclosed herein is suitable for reducing the level of one or more biomarkers of VLCADD in the subject.
- [0162] In some embodiments, a pharmaceutical composition or polynucleotide disclosed herein is for use in treating, preventing or delaying the onset of VLCADD signs or symptoms in a subject in need thereof. In some embodiments, the signs or symptoms include hypertrophic or dilated cardiomyopathy, pericardial effusion, arrhythmias, hypotonia, hepatomegaly, and intermittent hypoglycemia, or a combination thereof.
- [0163] The present disclosure also provides a host cell comprising a polynucleotide of the invention. In some embodiments, the host cell is a eukaryotic cell. The present disclosure also provides a vector comprising a polynucleotide of the invention. Also provided is a method of making a polynucleotide of the invention comprising synthesizing the polynucleotide enzymatically or chemically. The present disclosure also provides a polypeptide encoded by a polynucleotide of the invention, a composition comprising a polynucleotide of the invention, a host cell comprising a polynucleotide of the invention, a vector comprising a polynucleotide of the invention, or produced by the method of making disclosed herein.
- [0164] Some aspects of the present invention relate to a method of expressing an ACADVL polypeptide in a human subject in need thereof comprising administering to the subject an effective amount of a pharmaceutical composition suitable for administering as a single dose or as a plurality of single unit doses to the subject. In some embodiments, the administration treats, prevents or delays the onset of one or more of the signs or symptoms of VLCADD in the subject. In some embodiments, the method comprises administering to a human subject in need of treatment for VLCADD a single intravenous dose of the pharmaceutical composition.
- [0165] Some aspects of the present invention relate to a method wherein 24 hours after the pharmaceutical composition or polynucleotide is administered to the subject the level of an acylcarnitine in the subject is reduced by at least about 100%, at least about 90%, at least about 80%, at least about 70%, at least about 60%, at least about 50%, at least about 40%, at least about 30%, at least about 20%, or at least about 10% compared to the subject's baseline acylcarnitine level. In some embodiments, the level of the acylcarnitine is reduced in the blood of the subject. In some embodiments, the acylcarnitine is an acylcarnitine metabolite selected from the group consisting of C12:1 acylcarnitine, C14:1

acylcarnitine, C14:2 acylcarnitine, C14 acylcarnitine, C16 acylcarnitine, C18 acylcarnitine, C18.1 acylcarnitine, and combinations thereof.

[0166] Some aspects of the present invention relate to a method wherein 24 hours after the pharmaceutical composition or polynucleotide is administered to the subject, the ACADVL activity in the subject is increased to at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 100%, at least 150%, at least 200%, at least 300%, at least 400%, at least 500%, or at least 600% of the ACADVL activity in a normal individual. In some embodiments, the ACADVL activity is increased in the heart, liver, brain or skeletal muscle of the subject. In some embodiments, the increased ACADVL activity persists for greater than 24, 36, 48, 60, 72, 96, 120, 144, or 168 hours.

[0167] Some aspects of the present invention relate to a method wherein 24 hours after the pharmaceutical composition or polynucleotide is administered to the subject the level of an acylcarnitine in the subject is reduced by at least about 5%, at least about 10%, at least about 20%, at least about 30%, at least about 40%, at least about 50%, at least about 60%, at least about 70%, at least about 80%, at least about 90%, or 100% compared to the subject's baseline acylcarnitine level.

[0168] In some embodiments, the pharmaceutical composition or polynucleotide is administered as a single dose of less than 1.5 mg/kg, less than 1.25 mg/kg, less than 1 mg/kg, less than 0.75 mg/kg, or less than 0.5 mg/kg.

[0169] In some embodiments, the administration to the subject is about once a week, about once every two weeks, or about once a month.

[0170] In some embodiments, the pharmaceutical composition or polynucleotide is administered intravenously.

BRIEF DESCRIPTION OF THE DRAWINGS/FIGURES

[0171] **FIG. 1** shows the protein sequence (panel A), table with domain features (panel B), graphic representation of domain structure (panel C), and nucleic acid sequence (panel D) of isoform 1 of ACADVL.

[0172] **FIG. 2** shows the protein sequence (panel A), table with domain features (panel B), graphic representation of domain structure (panel C), and nucleic acid sequence (panel D) of isoform 2 of ACADVL.

- [0173] **FIG. 3** shows the protein sequence (panel A), table with domain features (panel B), graphic representation of domain structure (panel C), and nucleic acid sequence (panel D) of isoform 3 of ACADVL.
- [0174] **FIG. 4** shows uracil (U) metrics corresponding to wild type isoform 1 of ACADVL and 25 sequence optimized ACADVL polynucleotides. The column labeled "U content (%)" corresponds to the $%U_{TL}$ parameter. The column labeled "U Content v. WT (%)" corresponds to $%U_{WT}$. The column labeled "U Content v. Theoretical Minimum (%)" corresponds to $%U_{TM}$. The column labeled "UU pairs v. WT (%)" corresponds to $%UU_{WT}$.
- [0175] **FIG. 5** shows guanine (G) metrics corresponding to wild type isoform 1 of ACADVL and 25 sequence optimized ACADVL polynucleotides. The column labeled "G Content (%)" corresponds to $%G_{TL}$. The column labeled "G Content v. WT (%)" corresponds to $%G_{WT}$. The column labeled "G Content v. Theoretical Maximum (%)" corresponds to $%G_{TMX}$.
- [0176] **FIG. 6** shows cytosine (C) metrics corresponding to wild type isoform 1 of ACADVL and 25 sequence optimized ACADVL polynucleotides. The column labeled "C Content (%)" corresponds to $%C_{TL}$. The column labeled "C Content v. WT (%)" corresponds to $%C_{WT}$. The column labeled "C Content v. Theoretical Maximum (%)" corresponds to $%C_{TMX}$.
- [0177] **FIG. 7** shows guanine plus cytosine (G/C) metrics corresponding to wild type isoform 1 of ACADVL and 25 sequence optimized ACADVL polynucleotides. The column labeled "G/C Content (%)" corresponds to $%G/C_{TL}$. The column labeled "G/C Content v. WT (%)" corresponds to $%G/C_{WT}$. The column labeled "G/C Content v. Theoretical Maximum (%)" corresponds to $%G/C_{TMX}$.
- [0178] **FIG. 8** shows a comparison between the G/C compositional bias for codon positions 1, 2, 3 corresponding to the wild type isoform 1 of ACADVL and 25 sequence optimized ACADVL polynucleotides.

DETAILED DESCRIPTION

- [0179] The present invention provides mRNA therapeutics for the treatment of VLCADD. VLCADD is an autosomal recessive metabolic disorder characterized by the abnormal buildup of very long-chain fatty acids in patients. Such buildup of fatty acids can damage internal organs, resulting in a wide-range of symptoms. The principal gene

associated with VLCADD is acyl-CoA dehydrogenase, very long-chain (ACADVL; also referred to as VLCAD, ACAD6, or LCACD). VLCADD is caused by mutations in the *ACADVL* gene. mRNA therapeutics are particularly well-suited for the treatment of VLCADD as the technology provides for the intracellular delivery of mRNA encoding ACADVL followed by *de novo* synthesis of functional ACADVL protein within target cells. After delivery of mRNA to the target cells, the desired ACADVL protein is expressed by the cells' own translational machinery, and hence, fully functional ACADVL protein replaces the defective or missing protein.

[0180] One challenge associated with delivering nucleic acid-based therapeutics (*e.g.*, mRNA therapeutics) *in vivo* stems from the innate immune response which can occur when the body's immune system encounters foreign nucleic acids. Foreign mRNAs can activate the immune system *via* recognition through toll-like receptors (TLRs), in particular TLR7/8, which is activated by single-stranded RNA (ssRNA). In nonimmune cells, the recognition of foreign mRNA can occur through the retinoic acid-inducible gene I (RIG-I). Immune recognition of foreign mRNAs can result in unwanted cytokine effects including interleukin-1 β (IL-1 β) production, tumor necrosis factor- α (TNF- α) distribution and a strong type I interferon (type I IFN) response. The instant invention features the incorporation of different modified nucleotides within therapeutic mRNAs to minimize the immune activation and optimize the translation efficiency of mRNA to protein. Particular aspects of the invention feature a combination of nucleotide modification to reduce the innate immune response and sequence optimization, in particular, within the open reading frame (ORF) of therapeutic mRNAs encoding ACADVL to enhance protein expression.

[0181] Certain embodiments of the mRNA therapeutic technology of the instant invention also features delivery of mRNA encoding ACADVL *via* a lipid nanoparticle (LNP) delivery system. Lipid nanoparticles (LNPs) are an ideal platform for the safe and effective delivery of mRNAs to target cells. LNPs have the unique ability to deliver nucleic acids by a mechanism involving cellular uptake, intracellular transport and endosomal release or endosomal escape. The instant invention features novel ionizable lipid-based LNPs combined with mRNA encoding ACADVL which have improved properties when administered *in vivo*. Without being bound in theory, it is believed that the novel ionizable lipid-based LNP formulations of the invention have improved properties, for example, cellular uptake, intracellular transport and/or endosomal release

or endosomal escape. LNPs administered by systemic route (*e.g.*, intravenous (IV) administration), for example, in a first administration, can accelerate the clearance of subsequently injected LNPs, for example, in further administrations. This phenomenon is known as accelerated blood clearance (ABC) and is a key challenge, in particular, when replacing deficient enzymes (*e.g.*, ACADVL) in a therapeutic context. This is because repeat administration of mRNA therapeutics is in most instances essential to maintain necessary levels of enzyme in target tissues in subjects (*e.g.*, subjects in need of treatment for VLCADD). Repeat dosing challenges can be addressed on multiple levels. mRNA engineering and/or efficient delivery by LNPs can result in increased levels and or enhanced duration of protein (*e.g.*, ACADVL) being expressed following a first dose of administration, which in turn, can lengthen the time between first dose and subsequent dosing. It is known that the ABC phenomenon is, at least in part, transient in nature, with the immune responses underlying ABC resolving after sufficient time following systemic administration. As such, increasing the duration of protein expression and/or activity following systemic delivery of an mRNA therapeutic of the invention in one aspect, combats the ABC phenomenon. Moreover, LNPs can be engineered to avoid immune sensing and/or recognition and can thus further avoid ABC upon subsequent or repeat dosing. Exemplary aspect of the invention feature novel LNPs which have been engineered to have reduced ABC.

1. Acyl-CoA dehydrogenase, very long-chain (ACADVL)

[0182] Acyl-CoA dehydrogenase, very long-chain (ACADVL; EC 1.3.8.9) is a metabolic enzyme that plays a critical role in the catabolism of long-chain fatty acids, with highest specificity for carbon lengths C14-C18. ACADVL's biological function is to catalyze the first step of the mitochondrial fatty acid beta-oxidation pathway. ACADVL localizes to the inner mitochondrial membrane, where it functions as a homodimer.

[0183] The most well-known health issue involving ACADVL is very long-chain acyl-CoA dehydrogenase deficiency (VLCADD), an autosomal recessive metabolic disorder characterized by the abnormal buildup of very long-chain fatty acids in a patient's plasma. Such buildup of fatty acids can damage internal organs. Mutations within the ACADVL gene can result in the complete or partial loss of VLCAD function, which left untreated, could result in dire consequences, including, *e.g.*, liver failure, seizure, kidney failure, and brain damage.

- [0184] The coding sequence (CDS) for wild type ACADVL canonical mRNA sequence, corresponding to isoform 1, is described at the NCBI Reference Sequence database (RefSeq) under accession number NM_000018.3 ("Homo sapiens acyl-CoA dehydrogenase, very long chain (ACADVL), transcript variant 1, mRNA"). The wild type ACADVL canonical protein sequence, corresponding to isoform 1, is described at the RefSeq database under accession number NP_000009.1 ("Very long-chain specific acyl-CoA dehydrogenase, mitochondrial isoform 1 precursor [Homo sapiens]"). The ACADVL isoform 1 protein is 655 amino acids long. It is noted that the specific nucleic acid sequences encoding the reference protein sequence in the RefSeq sequences are the coding sequence (CDS) as indicated in the respective RefSeq database entry.
- [0185] Isoforms 2 and 3 are produced by alternative splicing.
- [0186] The RefSeq protein and mRNA sequences for isoform 2 of ACADVL are NP_001029031.1 and NM_001033859.2, respectively. The RefSeq protein and mRNA sequences for isoform 3 of ACADVL are NP_001257376.1 and NM_001270447.1, respectively. Isoforms 2 and 3 of ACADVL are encoded by the CDS disclosed in each one of the above mentioned mRNA RefSeq entries.
- [0187] The isoform 2 polynucleotide (transcript variant 2) lacks an alternate in-frame exon in the 5' coding region, compared to variant 1. It encodes an ACADVL isoform 2 polypeptide, which has the same N and C termini but is shorter than isoform 1. The ACADVL isoform 2 protein is 633 amino acids long and lacks the amino acids corresponding to positions 47-68 in isoform 1.
- [0188] The isoform 3 polynucleotide (transcript variant 3) differs in the 5' UTR and 5' coding region, compared to variant 1. The resulting ACADVL isoform 3 polypeptide is longer and has a distinct N-terminus, compared to isoform 1. The ACADVL isoform 3 protein is 678 amino acids long and contains a different set of amino acids at positions 1-20 in isoform 1.
- [0189] In certain aspects, the invention provides a polynucleotide (e.g., a ribonucleic acid (RNA), e.g., a messenger RNA (mRNA)) comprising a nucleotide sequence (e.g., an open reading frame (ORF)) encoding an ACADVL polypeptide. In some embodiments, the ACADVL polypeptide of the invention is a wild type ACADVL isoform 1, 2, or 3 protein. In some embodiments, the ACADVL polypeptide of the invention is a variant, a peptide or a polypeptide containing a substitution, and insertion and/or an addition, a deletion and/or a covalent modification with respect to a wild-type ACADVL isoform 1,

2, or 3 sequence. In some embodiments, sequence tags or amino acids, can be added to the sequences encoded by the polynucleotides of the invention (e.g., at the N-terminal or C-terminal ends), e.g., for localization. In some embodiments, amino acid residues located at the carboxy, amino terminal, or internal regions of a polypeptide of the invention can optionally be deleted providing for fragments.

[0190] In some embodiments, the polynucleotide (e.g., a RNA, e.g., an mRNA) comprising a nucleotide sequence (e.g., an ORF) of the invention encodes a substitutional variant of an ACADVL isoform 1, 2, 3, or 4 sequence, which can comprise one, two, three or more than three substitutions. In some embodiments, the substitutional variant can comprise one or more conservative amino acids substitutions. In other embodiments, the variant is an insertional variant. In other embodiments, the variant is a deletional variant.

[0191] As recognized by those skilled in the art, ACADVL isoform 1, 2, or 3 protein fragments, functional protein domains, variants, and homologous proteins (orthologs) are also considered to be within the scope of the ACADVL polypeptides of the invention. Nonlimiting examples of polypeptides encoded by the polynucleotides of the invention are shown in FIGS. 1 to 3. For example, FIG. 1 shows the amino acid sequence of human ACADVL wild type isoform 1.

[0192] Certain compositions and methods presented in this disclosure refer to the protein or polynucleotide sequences of ACADVL isoform 1. A person skilled in the art will understand that such disclosures are equally applicable to any other isoforms of ACADVL known in the art.

2. Polynucleotides and Open Reading Frames (ORFs)

[0193] The instant invention features mRNAs for use in treating (i.e., prophylactically and/or therapeutically treating) VLCADD. The mRNAs featured for use in the invention are administered to subjects and encode human ACADVL proteins(s) *in vivo*. Accordingly, the invention relates to polynucleotides, e.g., mRNA, comprising an open reading frame of linked nucleosides encoding human ACADVL, isoforms thereof, functional fragments thereof, and fusion proteins comprising ACADVL. In some embodiments, the open reading frame is sequence-optimized. In particular embodiments, the invention provides sequence-optimized polynucleotides comprising nucleotides

encoding the polypeptide sequence of human ACADVL, or sequence having high sequence identity with those sequence optimized polynucleotides.

[0194] In certain aspects, the invention provides polynucleotides (e.g., a RNA, e.g., an mRNA) that comprise a nucleotide sequence (e.g., an ORF) encoding one or more ACADVL polypeptides. In some embodiments, the encoded ACADVL polypeptide of the invention can be selected from:

- (i) a full length ACADVL polypeptide (e.g., having the same or essentially the same length as wild-type ACADVL isoform 1, 2, or 3);
- (ii) a functional fragment of any of the ACADVL isoforms described herein (e.g., a truncated (e.g., deletion of carboxy, amino terminal, or internal regions) sequence shorter than one of wild-type isoforms 1, 2, or 3; but still retaining ACADVL enzymatic activity);
- (iii) a variant thereof (e.g., full length or truncated isoform 1, 2, or 3 proteins in which one or more amino acids have been replaced, e.g., variants that retain all or most of the ACADVL activity of the polypeptide with respect to a reference isoform (such as, e.g., T59I, D178N, or any other natural or artificial variants known in the art); or
- (iv) a fusion protein comprising (i) a full length ACADVL isoform 1, 2, or 3 protein, a functional fragment or a variant thereof, and (ii) a heterologous protein.

[0195] In certain embodiments, the encoded ACADVL polypeptide is a mammalian ACADVL polypeptide, such as a human ACADVL polypeptide, a functional fragment or a variant thereof.

[0196] In some embodiments, the polynucleotide (e.g., a RNA, e.g., an mRNA) of the invention increases ACADVL protein expression levels and/or detectable ACADVL enzymatic activity levels in cells when introduced in those cells, e.g., by at least 10%, at least 15%, at least 20%, at least 25%, at least 30%, at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 100%, compared to ACADVL protein expression levels and/or detectable ACADVL enzymatic activity levels in the cells prior to the administration of the polynucleotide of the invention. ACADVL protein expression levels and/or ACADVL enzymatic activity can be measured according to methods known in the art. In some embodiments, the polynucleotide is introduced to the cells *in vitro*. In some embodiments, the polynucleotide is introduced to the cells *in vivo*.

- [0197] In some embodiments, the polynucleotides (e.g., a RNA, e.g., an mRNA) of the invention comprise a nucleotide sequence (e.g., an ORF) that encodes a wild-type human ACADVL, e.g., wild-type isoform 1 of human ACADVL (SEQ ID NO: 1, see FIG. 1), wild-type isoform 2 of human ACADVL (SEQ ID NO: 3, see FIG. 2), or wild-type isoform 3 of human ACADVL (SEQ ID NO: 5, see FIG. 3).
- [0198] In some embodiments, the polynucleotide (e.g., a RNA, e.g., an mRNA) of the invention comprises a codon optimized nucleic acid sequence, wherein the open reading frame (ORF) of the codon optimized nucleic sequence is derived from a wild-type ACADVL sequence (e.g., wild-type isoforms 1, 2, or 3). For example, for polynucleotides of invention comprising a sequence optimized ORF encoding ACADVL isoform 2, the corresponding wild type sequence is the native ACADVL isoform 2. Similarly, for an sequence optimized mRNA encoding a functional fragment of isoform 1, the corresponding wild type sequence is the corresponding fragment from ACADVL isoform 1.
- [0199] In some embodiments, the polynucleotides (e.g., a RNA, e.g., an mRNA) of the invention comprise a nucleotide sequence encoding ACADVL isoform 1 having the full length sequence of human ACADVL isoform 1 (i.e., including the initiator methionine). In mature human ACADVL isoform 1, the initiator methionine can be removed to yield a "mature ACADVL" comprising amino acid residues of 2-655 of the translated product. The teachings of the present disclosure directed to the full sequence of human ACADVL (amino acids 1-655) are also applicable to the mature form of human ACADVL lacking the initiator methionine (amino acids 2-615). Thus, in some embodiments, the polynucleotides (e.g., a RNA, e.g., an mRNA) of the invention comprise a nucleotide sequence encoding ACADVL isoform 1 having the mature sequence of human ACADVL isoform 1 (i.e., lacking the initiator methionine). In some embodiments, the polynucleotide (e.g., a RNA, e.g., an mRNA) of the invention comprising a nucleotide sequence encoding ACADVL isoform 1 having the full length or mature sequence of human ACADVL isoform 1 is sequence optimized.
- [0200] In some embodiments, the polynucleotides (e.g., a RNA, e.g., an mRNA) of the invention comprise a nucleotide sequence (e.g., an ORF) encoding a mutant ACADVL polypeptide. In some embodiments, the polynucleotides of the invention comprise an ORF encoding an ACADVL polypeptide that comprises at least one point mutation in the ACADVL sequence and retains ACADVL enzymatic activity. In some embodiments, the

mutant ACADVL polypeptide has an ACADVL activity which is at least 10%, at least 15%, at least 20%, at least 25%, at least 30%, at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 100% of the ACADVL activity of the corresponding wild-type ACADVL (i.e., the same ACADVL isoform but without the mutation(s)). In some embodiments, the polynucleotide (e.g., a RNA, e.g., an mRNA) of the invention comprising an ORF encoding a mutant ACADVL polypeptide is sequence optimized.

[0201] In some embodiments, the polynucleotide (e.g., a RNA, e.g., an mRNA) of the invention comprises a nucleotide sequence (e.g., an ORF) that encodes an ACADVL polypeptide with mutations that do not alter ACADVL enzymatic activity. Such mutant ACADVL polypeptides can be referred to as function-neutral. In some embodiments, the polynucleotide comprises an ORF that encodes a mutant ACADVL polypeptide comprising one or more function-neutral point mutations.

[0202] In some embodiments, the mutant ACADVL polypeptide has higher ACADVL enzymatic activity than the corresponding wild-type ACADVL. In some embodiments, the mutant ACADVL polypeptide has an ACADVL activity that is at least 10%, at least 15%, at least 20%, at least 25%, at least 30%, at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 100% higher than the activity of the corresponding wild-type ACADVL (i.e., the same ACADVL isoform but without the mutation(s)).

[0203] In some embodiments, the polynucleotides (e.g., a RNA, e.g., an mRNA) of the invention comprise a nucleotide sequence (e.g., an ORF) encoding a functional ACADVL fragment, e.g., where one or more fragments correspond to a polypeptide subsequence of a wild type ACADVL polypeptide and retain ACADVL enzymatic activity. In some embodiments, the ACADVL fragment has an ACADVL activity which is at least 10%, at least 15%, at least 20%, at least 25%, at least 30%, at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 100% of the ACADVL activity of the corresponding full length ACADVL. In some embodiments, the polynucleotides (e.g., a RNA, e.g., an mRNA) of the invention comprising an ORF encoding a functional ACADVL fragment is sequence optimized.

- [0204] In some embodiments, the polynucleotide (e.g., a RNA, e.g., an mRNA) of the invention comprises a nucleotide sequence (e.g., an ORF) encoding an ACADVL fragment that has higher ACADVL enzymatic activity than the corresponding full length ACADVL. Thus, in some embodiments the ACADVL fragment has an ACADVL activity which is at least 10%, at least 15%, at least 20%, at least 25%, at least 30%, at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 100% higher than the ACADVL activity of the corresponding full length ACADVL.
- [0205] In some embodiments, the polynucleotide (e.g., a RNA, e.g., an mRNA) of the invention comprises a nucleotide sequence (e.g., an ORF) encoding an ACADVL fragment that is at least 1%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19%, 20%, 21%, 22%, 23%, 24% or 25% shorter than wild-type isoform 1, 2, 3, or 4 of ACADVL.
- [0206] In some embodiments, the polynucleotide (e.g., a RNA, e.g., an mRNA) of the invention comprises a nucleotide sequence (e.g., an ORF) encoding an ACADVL polypeptide (e.g., the wild-type sequence, functional fragment, or variant thereof), wherein the nucleotide sequence is at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identical to the sequence of SEQ ID NO:2, 4, or 6 (see, e.g., panel D in FIG. 1, 2, and 3, respectively).
- [0207] In some embodiments, the polynucleotide (e.g., a RNA, e.g., an mRNA) of the invention comprises a nucleotide sequence (e.g., an ORF) encoding an ACADVL polypeptide (e.g., the wild-type sequence, functional fragment, or variant thereof), wherein the nucleotide sequence has at least 70%, at least 71%, at least 72%, at least 73%, at least 74%, at least 75%, at least 76%, at least 77%, at least 78%, at least 79%, at least 80%, at least 81%, at least 82%, at least 83%, at least 84%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to a sequence selected from the group consisting of SEQ ID NOs: 7 to 31. See TABLE 2.
- [0208] In some embodiments, the polynucleotide (e.g., a RNA, e.g., an mRNA) of the invention comprises a nucleotide sequence (e.g., an ORF) encoding an ACADVL

polypeptide (e.g., the wild-type sequence, functional fragment, or variant thereof), wherein the nucleotide sequence has 70% to 100%, 75% to 100%, 80% to 100%, 85% to 100%, 70% to 95%, 80% to 95%, 70% to 85%, 75% to 90%, 80% to 95%, 70% to 75%, 75% to 80%, 80% to 85%, 85% to 90%, 90% to 95%, or 95% to 100%, sequence identity to a sequence selected from the group consisting of SEQ ID NOS: 7 to 31. See TABLE 2.

[0209] In some embodiments, the polynucleotide (e.g., a RNA, e.g., an mRNA) of the invention comprises a nucleotide sequence (e.g., an ORF) encoding an ACADVL polypeptide (e.g., the wild-type sequence, functional fragment, or variant thereof), wherein the nucleotide sequence is at least 75%, at least 76%, at least 77%, at least 78%, at least 79%, at least 80%, at least 81%, at least 82%, at least 83%, at least 84%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identical to the sequence of SEQ ID NO: 2, 4, or 6 (see, e.g., panel D of FIGS. 1, 2, and 3, respectively).

[0210] In some embodiments the polynucleotide (e.g., a RNA, e.g., an mRNA) of the invention comprises a nucleotide sequence (e.g., an ORF) encoding an ACADVL polypeptide (e.g., the wild-type sequence, functional fragment, or variant thereof), wherein the nucleotide sequence is between 70% and 90% identical; between 75% and 85% identical; between 76% and 84% identical; between 77% and 83% identical, between 77% and 82% identical, or between 78% and 81% identical to the sequence of SEQ ID NO: 2, 4, or 6 (see, e.g., panel D of FIGS. 1, 2, and 3, respectively).

[0211] In some embodiments, the polynucleotide (e.g., a RNA, e.g., an mRNA) of the invention comprises from about 1,500 to about 100,000 nucleotides (e.g., from 1,500 to 1,600, from 1,500 to 1,700, from 1,500 to 1,800, from 1,500 to 1,900, from 1,500 to 2,000, from 1,600 to 1,700, from 1,600 to 1,800, from 1,600 to 1,900, from 1,600 to 2,000, from 1,600 to 1,300, from 1,600 to 1,400, from 1,000 to 1,500, from 1,083 to 1,200, from 1,845 to 2,000, from 1,845 to 2,200, from 1,845 to 2,400, from 1,845 to 2,600, from 1,845 to 2,800, from 1,845 to 3,000, from 1,845 to 5,000, from 1,845 to 7,000, from 1,845 to 10,000, from 1,845 to 25,000, from 1,845 to 50,000, from 1,845 to 70,000, or from 1,845 to 100,000).

[0212] In some embodiments, the polynucleotide of the invention (e.g., a RNA, e.g., an mRNA) comprises a nucleotide sequence (e.g., an ORF) encoding an ACADVL polypeptide (e.g., the wild-type sequence, functional fragment, or variant thereof),

wherein the length of the nucleotide sequence (e.g., an ORF) is at least 600 nucleotides in length (e.g., at least or greater than about 600, 700, 80, 900, 1,000, 1,050, 1,083, 1,100, 1,200, 1,300, 1,400, 1,500, 1,600, 1,700, 1,800, 1,845, 1,900, 2,000, 2,100, 2,200, 2,300, 2,400, 2,500, 2,600, 2,700, 2,800, 2,900, 3,000, 3,100, 3,200, 3,300, 3,400, 3,500, 3,600, 3,700, 3,800, 3,900, 4,000, 4,100, 4,200, 4,300, 4,400, 4,500, 4,600, 4,700, 4,800, 4,900, 5,000, 5,100, 5,200, 5,300, 5,400, 5,500, 5,600, 5,700, 5,800, 5,900, 6,000, 7,000, 8,000, 9,000, 10,000, 20,000, 30,000, 40,000, 50,000, 60,000, 70,000, 80,000, 90,000 or up to and including 100,000 nucleotides).

[0213] In some embodiments, the polynucleotide of the invention (e.g., a RNA, e.g., an mRNA) comprises a nucleotide sequence (e.g., an ORF) encoding an ACADVL polypeptide (e.g., the wild-type sequence, functional fragment, or variant thereof) further comprises at least one nucleic acid sequence that is noncoding, e.g., a miRNA binding site. In some embodiments, the polynucleotide (e.g., a RNA, e.g., an mRNA) of the invention further comprises a 5'-UTR (e.g., selected from the sequences of SEQ ID NOs: 38, 40 to 57, and 117 to 119) and a 3'UTR (e.g., selected from the sequences of SEQ ID NOs: 39, 58 to 84, 90, 105, 108 to 115, and 120 to 130). In a further embodiment, the polynucleotide (e.g., a RNA, e.g., an mRNA) comprises a 5' terminal cap (e.g., Cap0, Cap1, ARCA, inosine, N1-methyl-guanosine, 2'-fluoro-guanosine, 7-deaza-guanosine, 8-oxo-guanosine, 2-amino-guanosine, LNA-guanosine, 2-azidoguanosine, Cap2, Cap4, 5' methylG cap, or an analog thereof) and a poly-A-tail region (e.g., about 100 nucleotides in length). In a further embodiment, the polynucleotide (e.g., a RNA, e.g., an mRNA) a comprises a 3' UTR comprising a nucleic acid sequence selected from the group consisting of SEQ ID NOs: 83 to 85 and 105, or any combination thereof.

[0214] In some embodiments, the polynucleotide of the invention (e.g., a RNA, e.g., an mRNA) comprises a nucleotide sequence (e.g., an ORF) encoding an ACADVL polypeptide is single stranded or double stranded.

[0215] In some embodiments, the polynucleotide of the invention comprising a nucleotide sequence (e.g., an ORF) encoding an ACADVL polypeptide (e.g., the wild-type sequence, functional fragment, or variant thereof) is DNA or RNA. In some embodiments, the polynucleotide of the invention is RNA. In some embodiments, the polynucleotide of the invention is, or functions as, a messenger RNA (mRNA). In some embodiments, the mRNA comprises a nucleotide sequence (e.g., an ORF) that encodes at

least one ACADVL polypeptide, and is capable of being translated to produce the encoded ACADVL polypeptide *in vitro*, *in vivo*, *in situ* or *ex vivo*.

[0216] In some embodiments, the polynucleotide of the invention (e.g., a RNA, e.g., an mRNA) comprises a sequence-optimized nucleotide sequence (e.g., an ORF) encoding an ACADVL polypeptide (e.g., the wild-type sequence, functional fragment, or variant thereof), wherein the polynucleotide comprises at least one chemically modified nucleobase, e.g., 5-methoxyuracil. In some embodiments, the polynucleotide further comprises a miRNA binding site, e.g., a miRNA binding site that binds to miR-142 and/or a miRNA binding site that binds to miR-126. In some embodiments, the polynucleotide (e.g., a RNA, e.g., an mRNA) disclosed herein is formulated with a delivery agent comprising, e.g., a compound having the Formula (I), e.g., any of Compounds 1-232, e.g., Compound 18; a compound having the Formula (III), (IV), (V), or (VI), e.g., any of Compounds 233-342, e.g., Compound 236; or a compound having the Formula (VIII), e.g., any of Compounds 419-428, e.g., Compound 428, or any combination thereof. In some embodiments, the delivery agent comprises Compound 18, DSPC, Cholesterol, and Compound 428, e.g., with a mole ratio of about 50:10:38.5:1.5.

3. Signal Sequences

[0217] The polynucleotides (e.g., a RNA, e.g., an mRNA) of the invention can also comprise nucleotide sequences that encode additional features that facilitate trafficking of the encoded polypeptides to therapeutically relevant sites. One such feature that aids in protein trafficking is the signal sequence, or targeting sequence. The peptides encoded by these signal sequences are known by a variety of names, including targeting peptides, transit peptides, and signal peptides. In some embodiments, the polynucleotide (e.g., a RNA, e.g., an mRNA) comprises a nucleotide sequence (e.g., an ORF) that encodes a signal peptide operably linked a nucleotide sequence that encodes an ACADVL polypeptide described herein.

[0218] In some embodiments, the "signal sequence" or "signal peptide" is a polynucleotide or polypeptide, respectively, which is from about 9 to 200 nucleotides (3-70 amino acids) in length that, optionally, is incorporated at the 5' (or N-terminus) of the coding region or the polypeptide, respectively. Addition of these sequences results in trafficking the encoded polypeptide to a desired site, such as the endoplasmic reticulum or the mitochondria through one or more targeting pathways. Some signal peptides are

cleaved from the protein, for example by a signal peptidase after the proteins are transported to the desired site.

[0219] In some embodiments, the polynucleotide of the invention comprises a nucleotide sequence encoding an ACADVL polypeptide, wherein the nucleotide sequence further comprises a 5' nucleic acid sequence encoding a native signal peptide. In another embodiment, the polynucleotide of the invention comprises a nucleotide sequence encoding an ACADVL polypeptide, wherein the nucleotide sequence lacks the nucleic acid sequence encoding a native signal peptide.

[0220] In some embodiments, the polynucleotide of the invention comprises a nucleotide sequence encoding an ACADVL polypeptide, wherein the nucleotide sequence further comprises a 5' nucleic acid sequence encoding a heterologous signal peptide.

4. Fusion Proteins

[0221] In some embodiments, the polynucleotide of the invention (e.g., a RNA, e.g., an mRNA) can comprise more than one nucleic acid sequence (e.g., an ORF) encoding a polypeptide of interest. In some embodiments, polynucleotides of the invention comprise a single ORF encoding an ACADVL polypeptide, a functional fragment, or a variant thereof. However, in some embodiments, the polynucleotide of the invention can comprise more than one ORF, for example, a first ORF encoding an ACADVL polypeptide (a first polypeptide of interest), a functional fragment, or a variant thereof, and a second ORF expressing a second polypeptide of interest. In some embodiments, two or more polypeptides of interest can be genetically fused, i.e., two or more polypeptides can be encoded by the same ORF. In some embodiments, the polynucleotide can comprise a nucleic acid sequence encoding a linker (e.g., a G₄S peptide linker or another linker known in the art) between two or more polypeptides of interest.

[0222] In some embodiments, a polynucleotide of the invention (e.g., a RNA, e.g., an mRNA) can comprise two, three, four, or more ORFs, each expressing a polypeptide of interest.

[0223] In some embodiments, the polynucleotide of the invention (e.g., a RNA, e.g., an mRNA) can comprise a first nucleic acid sequence (e.g., a first ORF) encoding an ACADVL polypeptide and a second nucleic acid sequence (e.g., a second ORF) encoding a second polypeptide of interest.

5. Sequence Optimization of Nucleotide Sequence Encoding an ACADVL Polypeptide

[0224] In some embodiments, the polynucleotide (e.g., a RNA, e.g., an mRNA) of the invention is sequence optimized. In some embodiments, the polynucleotide (e.g., a RNA, e.g., an mRNA) of the invention comprises a nucleotide sequence (e.g., an ORF) encoding an ACADVL polypeptide, a nucleotide sequence (e.g., an ORF) encoding another polypeptide of interest, a 5'-UTR, a 3'-UTR, a miRNA, a nucleotide sequence encoding a linker, or any combination thereof) that is sequence optimized.

[0225] A sequence-optimized nucleotide sequence, e.g., an codon-optimized mRNA sequence encoding an ACADVL polypeptide, is a sequence comprising at least one synonymous nucleobase substitution with respect to a reference sequence (e.g., a wild type nucleotide sequence encoding an ACADVL polypeptide).

[0226] A sequence-optimized nucleotide sequence can be partially or completely different in sequence from the reference sequence. For example, a reference sequence encoding polyserine uniformly encoded by TCT codons can be sequence-optimized by having 100% of its nucleobases substituted (for each codon, T in position 1 replaced by A, C in position 2 replaced by G, and T in position 3 replaced by C) to yield a sequence encoding polyserine which would be uniformly encoded by AGC codons. The percentage of sequence identity obtained from a global pairwise alignment between the reference polyserine nucleic acid sequence and the sequence-optimized polyserine nucleic acid sequence would be 0%. However, the protein products from both sequences would be 100% identical.

[0227] Some sequence optimization (also sometimes referred to codon optimization) methods are known in the art (and discussed in more detail below) and can be useful to achieve one or more desired results. These results can include, e.g., matching codon frequencies in certain tissue targets and/or host organisms to ensure proper folding; biasing G/C content to increase mRNA stability or reduce secondary structures; minimizing tandem repeat codons or base runs that can impair gene construction or expression; customizing transcriptional and translational control regions; inserting or removing protein trafficking sequences; removing/adding post translation modification sites in an encoded protein (e.g., glycosylation sites); adding, removing or shuffling protein domains; inserting or deleting restriction sites; modifying ribosome binding sites and mRNA degradation sites; adjusting translational rates to allow the various domains of the protein to fold properly; and/or reducing or eliminating problem secondary structures

within the polynucleotide. Sequence optimization tools, algorithms and services are known in the art, non-limiting examples include services from GeneArt (Life Technologies), DNA2.0 (Menlo Park CA) and/or proprietary methods.

[0228] Codon options for each amino acid are given in TABLE 1.

TABLE 1. Codon Options

Amino Acid	Single Letter Code	Codon Options
Isoleucine	I	ATT, ATC, ATA
Leucine	L	CTT, CTC, CTA, CTG, TTA, TTG
Valine	V	GTT, GTC, GTA, GTG
Phenylalanine	F	TTT, TTC
Methionine	M	ATG
Cysteine	C	TGT, TGC
Alanine	A	GCT, GCC, GCA, GCG
Glycine	G	GGT, GGC, GGA, GGG
Proline	P	CCT, CCC, CCA, CCG
Threonine	T	ACT, ACC, ACA, ACG
Serine	S	TCT, TCC, TCA, TCG, AGT, AGC
Tyrosine	Y	TAT, TAC
Tryptophan	W	TGG
Glutamine	Q	CAA, CAG
Asparagine	N	AAT, AAC
Histidine	H	CAT, CAC
Glutamic acid	E	GAA, GAG
Aspartic acid	D	GAT, GAC
Lysine	K	AAA, AAG
Arginine	R	CGT, CGC, CGA, CGG, AGA, AGG
Selenocysteine	Sec	UGA in mRNA in presence of Selenocysteine insertion element (SECIS)
Stop codons	Stop	TAA, TAG, TGA

[0229] In some embodiments, a polynucleotide (e.g., a RNA, e.g., an mRNA) of the invention comprises a sequence-optimized nucleotide sequence (e.g., an ORF) encoding an ACADVL polypeptide, a functional fragment, or a variant thereof, wherein the ACADVL polypeptide, functional fragment, or a variant thereof encoded by the sequence-optimized nucleotide sequence has improved properties (e.g., compared to an ACADVL polypeptide, functional fragment, or a variant thereof encoded by a reference nucleotide sequence that is not sequence optimized), e.g., improved properties related to expression efficacy after administration *in vivo*. Such properties include, but are not limited to, improving nucleic acid stability (e.g., mRNA stability), increasing translation efficacy in the target tissue, reducing the number of truncated proteins expressed, improving the folding or prevent misfolding of the expressed proteins, reducing toxicity

of the expressed products, reducing cell death caused by the expressed products, increasing and/or decreasing protein aggregation.

[0230] In some embodiments, the sequence-optimized nucleotide sequence is codon optimized for expression in human subjects, having structural and/or chemical features that avoid one or more of the problems in the art, for example, features which are useful for optimizing formulation and delivery of nucleic acid-based therapeutics while retaining structural and functional integrity; overcoming a threshold of expression; improving expression rates; half-life and/or protein concentrations; optimizing protein localization; and avoiding deleterious bio-responses such as the immune response and/or degradation pathways.

[0231] In some embodiments, the polynucleotides of the invention comprise a nucleotide sequence (e.g., a nucleotide sequence (e.g., an ORF) encoding an ACADVL polypeptide, a nucleotide sequence (e.g., an ORF) encoding another polypeptide of interest, a 5'-UTR, a 3'-UTR, a microRNA binding site, a nucleic acid sequence encoding a linker, or any combination thereof) that is sequence-optimized according to a method comprising:

(i) substituting at least one codon in a reference nucleotide sequence (e.g., an ORF encoding an ACADVL polypeptide) with an alternative codon to increase or decrease uridine content to generate a uridine-modified sequence;

(ii) substituting at least one codon in a reference nucleotide sequence (e.g., an ORF encoding an ACADVL polypeptide) with an alternative codon having a higher codon frequency in the synonymous codon set;

(iii) substituting at least one codon in a reference nucleotide sequence (e.g., an ORF encoding an ACADVL polypeptide) with an alternative codon to increase G/C content; or

(iv) a combination thereof.

[0232] In some embodiments, the sequence-optimized nucleotide sequence (e.g., an ORF encoding an ACADVL polypeptide) has at least one improved property with respect to the reference nucleotide sequence.

[0233] In some embodiments, the sequence optimization method is multiparametric and comprises one, two, three, four, or more methods disclosed herein and/or other optimization methods known in the art.

[0234] Features, which can be considered beneficial in some embodiments of the invention, can be encoded by or within regions of the polynucleotide and such regions

can be upstream (5') to, downstream (3') to, or within the region that encodes the ACADVL polypeptide. These regions can be incorporated into the polynucleotide before and/or after sequence-optimization of the protein encoding region or open reading frame (ORF). Examples of such features include, but are not limited to, untranslated regions (UTRs), microRNA sequences, Kozak sequences, oligo(dT) sequences, poly-A tail, and detectable tags and can include multiple cloning sites that can have XbaI recognition.

[0235] In some embodiments, the polynucleotide of the invention comprises a 5' UTR, a 3' UTR and/or a miRNA binding site. In some embodiments, the polynucleotide comprises two or more 5' UTRs and/or 3' UTRs, which can be the same or different sequences. In some embodiments, the polynucleotide comprises two or more miRNA, which can be the same or different sequences. Any portion of the 5' UTR, 3' UTR, and/or miRNA binding site, including none, can be sequence-optimized and can independently contain one or more different structural or chemical modifications, before and/or after sequence optimization.

[0236] In some embodiments, after optimization, the polynucleotide is reconstituted and transformed into a vector such as, but not limited to, plasmids, viruses, cosmids, and artificial chromosomes. For example, the optimized polynucleotide can be reconstituted and transformed into chemically competent *E. coli*, yeast, neurospora, maize, drosophila, etc. where high copy plasmid-like or chromosome structures occur by methods described herein.

6. Sequence-Optimized Nucleotide Sequences Encoding ACADVL Polypeptides

[0237] In some embodiments, the polynucleotide of the invention comprises a sequence-optimized nucleotide sequence encoding an ACADVL polypeptide disclosed herein. In some embodiments, the polynucleotide of the invention comprises an open reading frame (ORF) encoding an ACADVL polypeptide, wherein the ORF has been sequence optimized.

[0238] Exemplary sequence-optimized nucleotide sequences encoding human ACADVL isoform 1 are set forth as SEQ ID NOs: 7-31 (ACADVL-CO01, ACADVL-CO02, ACADVL-CO03, ACADVL-CO04, ACADVL-CO05, ACADVL-CO06, ACADVL-CO07, ACADVL-CO08, ACADVL-CO09, ACADVL-CO10, ACADVL-CO11, ACADVL-CO12, ACADVL-CO13, ACADVL-CO14, ACADVL-CO15, ACADVL-CO16, ACADVL-CO17, ACADVL-CO18, ACADVL-CO19, ACADVL-CO20,

ACADVL-CO21, ACADVL-CO22, ACADVL-CO23, ACADVL-CO24, and ACADVL-CO25, respectively). Exemplary sequence optimized nucleotide sequences encoding human ACADVL are shown in TABLE 2. In some embodiments, the sequence optimized ACADVL sequences set forth as SEQ ID NOs: 7-31 or shown in TABLE 2, fragments, and variants thereof are used to practice the methods disclosed herein. In some embodiments, the sequence optimized ACADVL sequences in TABLE 2, fragments and variants thereof are combined with or alternatives to the wild-type sequences disclosed in FIGS. 1-3.

- [0239] In some embodiments, a polynucleotide of the present disclosure, for example a polynucleotide comprising an mRNA nucleotide sequence encoding an ACADVL polypeptide, comprises from 5' to 3' end:
- (i) a 5' cap provided herein, for example, CAP1;
 - (ii) a 5' UTR, such as the sequences provided herein,
 - (iii) an open reading frame encoding an ACADVL polypeptide, e.g., a sequence optimized nucleic acid sequence encoding ACADVL set forth as SEQ ID NOs: 7 to 31, or shown in TABLE 2;
 - (iv) at least one stop codon;
 - (v) a 3' UTR, such as the sequences provided herein; and
 - (vi) a poly-A tail provided above.

TABLE 2: Sequence optimized sequences for human ACADVL, isoform 1

SEQ ID NO	Name	Sequence
7	ACADVL-CO01	AUGAGCGGGAACGGCAACGCCGACGCCACGCCGAGGAGAACAGCCCCAAGAUGC GGGUCAUCAGGGUGG GCACGCGGAAGUCCAGUUGGCACGCAUACAGACCGACAGCGUAGUGGCGACCCUGAAGGCCUCCUACCC CGGCCUCCAGUUCGAGAUCAUCGCCAUGAGCACCACCGGGGACAAGAUCUCGAUACCGCCUGUCCAAG AUCGGCGAGAAGUCCUUGUUUACCAAGGAGCUGGAGCAGCCUGGAGAAGAAUGAGGUGGACCUUGUGG UGCACAGCCUCAAGGACCUGCCCACCGUGCGUCCCCCGGGUUCACCAUAGGAGCCAUCUGCAAGAGGGA GAACCCGCACGACGCAGUCGUGUCCACCCCAAGUUUGUGGGUAAGACACUGGAGACCCUGCCGAGAAG AGCGUGGUGGGUACCUCAGCCUCCGCCGAGCAGCCAGCUGCAGAGGAAGUCCCGCACCUGGAAUUC GAUCCAUCAGGGGCAAUUCGAACACACAGACUGAGGAAGCUGGAUGAGCAGCAGGAAUUUCCGCGAUAAU CCUCGCGACCGCCGGGUGCAGAGGAUGGGCUGGCAUACAGGGUGGGCCAAUCCUGCACCCCGAGGAG UGCAUGUACGCCUGGGCCAGGGCGCCUCCGCGUGGAGGUGAGGGCCAAAGACCAAGACAUCUGGACC UGGUGGGGUGCCUGCACGAUCCGGAGACUCUGCUGAGGUGCAUCGCCGAGCGGGCGUUUCAGGCACCU CGAGGGCGGUGCUCGUGCCCGUGGGCGUCCAUACCGCCAUGAAGGACGGCCAGCUGUACCUCACCGGC GGCGUGUGGAGCCUGGACGGCAGCGACUCCAUCAGGAAACCAUGCAAGCGACCAUACAUGUCCCCGCCC AGCACAGGACGGGCCCCAGGAUGACCCGCGAGCUGGUCGGUAUACCGCAAGGAUAUCCCCGGGGCCC ACAGCUGGCCGCCAGAAUCUGGGCAUCUCCUGGCCAAUCUGCUGCUGUCCAAGGGCGCCAAGAACAU CUCGACGUGGCCAGGCAACUGAACGACGCACAC
8	ACADVL-CO02	AUGAGCGGCAACGGUAACGCCGCCGCCACGCCGAGGAGAACUCGCCCCAAGAUGC GGGUCAUCAGGGUGG GAACGAGGAAGUCCAGCUGGCCAGGAUCCAGACCGACAGCGUGGUGGCGACGCUAAGGCCAGCUACCC CGGCCUGCAGUUUGAGAUAAUCGCCAUGUACAACACCGGCGACAAGAUCUCGACACCGCGCUGAGCAAG AUCGGCGAGAAGUCCUUGUACCAAGGAGCUGGAGCAGCCUUGGAAAAGAACGAGGUGGAUCUGGUCG UGCACAGCCUGAAGGACCUGCCCACCGUGCGUCCCCCGGCUUCACCAUCGGGCCCAUCUGCAAGCGGGA AAAUCCCCAUGACGCCGUGGUGUCCACCCCAAGUUCGUGGGGAAGACCCUGGAGACCCUGCCGAGAAG AGCGUGGUGGGAACAGCUCACUGCGGGGGCCGCCAGCUGCAACGUAAGUCCCCCAUCUGGAGUUC

		GAUCCAUCCGAGGCAAUCUGAACACCAGGCUAGAGAAAGCUGGAUGAACACAGGAGUUCAGCGCCAUCAU CCUGGCAACCGCCGGUCUGCAGAGGAUGGGCUGGCACAACAGGGUGGGCCAAAUCCUGCACCCTGAGGAG UGCAUGUAUGCAGUGGGGACGGGCGCGCUGGGUGUCGAGGUGCGAGCGAAGGACAGGACAUCUCGACC UGGUGGGAGUGCUGCAGCAGCCCGAGACCCUGCUGCGGUGCAUCGCGGAGAGGGCCUUCUGCGCCACCU CGAGGGCGGUCUGCAGUGCCCGUGGCGGUGCAUACCGCCAUGAAGGACGGCCAGCUGUACCGACCGGC GGUGUGUGGAGCCUCGACGGCAGCGACUCCAUCAGGAAACGAUCGAGGCGACUACCGUGCCCGCAC AGCACGAGGACGGCCCGAAGACGACCCCGAGCUGGUCGGGAUACCCGCCAGGAACAUCCTCCGAGGGGCC CCAGCUGGGCGGCCAAACCUCCGCAUAUCCUGGCCAACUGCUGCUGUCCAAGGGCGCCAAGAACAU CUGGAUGUGGGCCGCCAACUGAACGACGCCAC
9	ACADVL- CO03	AUGAGCGGUAACGGCAACGCCGCGCCACAGCCGAAGAGAACUCCCCAAGAUAGGGGUGAUUAGGGUGG GUACCCGAAAGAGCCAGCUGGCCCGGAUCCAGACCGACUCCUGGGUGGCCACCCUCAAGGCAAGCUACCC CGGCCUGCAGUUCGAGAUCAUCGCCAUGAGCACCACCGGGGACAAAUUCUGCAGACCCGCCUGAGCAAG AUCGGCGAGAAUCCUUGUUCACCAAGGAGCUAGAGCACGCCUCGAAAAGAACGAGGUGGAUCUGGUCG UGCACUCCUGAAGGACCUGCCACCGUGCUGCCCCCGGCUUCACGAUCGGCGCGAUUCUGCAAGAGGGA GAACCCGCAUGACGCCGUGGUGUUAUCCCAAAUUGUGGGAAAAACCCUGGAGACUUCCTCCGAGAA AGCGUCGUGGGACCUCCUCCUGCGGGCGGGCCGCCAGCUGCAGCGCAAGUUCCTCCGACCCUGGAGUUC GCAGCAUCAGGGGAAACCUGAACACCCGGCUGAGGAAGCUGGACGAGCAGCAGGAGUUCAGCGCUAUAU CCUCGCCACCGCCGGCCUCCAGCGAAUGGGCUGGCAUAACCGGGUGGGCCAGAUCCUGCACCCTGGAAGAA UGUAUGUACGCGGUGGGCCAGGGCGCGCUGGGCGUGGAGGUGAGGGCCAAGGACCAGGACAUCUGGAUC UCGUGGGCGUGCUGCAGCAGCCCGAGACCCUGCUGAGAUGCAUCGCGAGAACGGGCCUUCUGAGGCAACCU GGAGGGCGGCGUCUCGUGCCGGUGGGCGUACAUAACCGCAAUGAAAGACGGCCAGCUCUACCGACGGGG GGGGUGUGGAGCCUCGACGGCUCGACUCCAUCAGGAGACCAUGCAGGCGCAGAUCCAUGUACCCGCC AGCACGAGGACGGCCCGAGGAUGAUCCCAACUCUGUGGGCAUACCGCCCGCAACUCCGAGGGGGCC CCAGCUGGGCGCACAGAACCUGGGAUAUACUGGCCAACCUCCUGCUCAGCAAGGGCGCCAAGAAUAUC CUCGACGUGGCCAGGCAGCUGAAUGACGCCAC
10	ACADVL- CO04	AUGAGCGGCAACGGCAACGCCGCGCCACCGCCGAGGAGAACAGCCCCAAAUGCGCGUGAUCAGGGUGG GCACCAGGAAGAGCCAGCUGGCCCGCAUCCAGACCGACAGUGUCGUGGCCACCCUCAAGGCCAGCUAUCC GGGCCUGCAAUUCGAGAUCAUCGCCAUGAGCACCACCGGGCGACAAGAUCCUGACACCGCCUCAGCAAG AUCGGCGAGAAUCCUUGUUCACCAAGGAGCUGGACGACGCCUUGGAGAAGAACGAGGUGGACCUGGUGG UGCACAGCCUCAAGGACCUGCCACCGUGCUGCCCCCGGCUUCACCAUCGGCGCCAUUCUGCGCCAGGA GAACCCCCACGACGCCGUGGUGUUAUCCCAAGUUCUGGGCAAGACCUGGAGACCCUGCCCGAGAAG AGCGUGGUCGGCACACGCUCCUCAGGAGAGCCGCCAGCUGCAGCGAAAGUUCCTCCACCUUGGAUUC GCAGCAUCCGGGGGAACCUGAACACCCGCCUGAGGAAGCUGGACGAGCAGCAAGAAUUCAGCGCCAUCAU CCUGGCCACCGCGGGCCUGCAGAGGAUGGGUGGCACAACCGCGUCGGCCAGAUCCUGCACCAGAGGAG UGCAUGUACGCGGUGGGGACAGGGGGCCUUGGGGUGGAAGUGCGCGCCAAGGACCAGGACAUCUGGACC UGGUCGGCGUGCUGCAUGACCCGAGACCCUGCUGAGGUGCAUCGCCGAGAGGGCCUUCUGCGCCACCU CGAGGGGGGUGCAGCGUGCCGUGCGCGUCCAUCAGGCCAUGAAGGACGGGCGAGCUGUACCGACCGGC GGCGUGUGGUCCUGGACGGCUCGACAGCAUCCAGGAGACCAUGCAGGCCACCAUCCAGUGCCCGCCC AACACGAGGACGGCCCGAGGACGACCCGACGUCUGCUGGCAUACCGCCAGGAUAUACCTCCGUGGGCC CCAGCUGGGCGCCAGAAUCUGCGCAUCAGCCUGGCCAAUCUGCUGCUGAGCAAGGGGGCCAAGAACAU CUGGACGUGGCCAGGCAGCUGAAUGAUGCCAC
11	ACADVL- CO05	AUGAGCGGCAACGGAAACGCCGCGCGCAGCGCCGAGGAGAACUCCCCAAAUGAGGGUCAUCAGGGUGG GCACCCGGAAGAGCCAGCUGGGCGCGGAUCCAGCAGCGUCGUGGCCACCCUCAAGGCCAGCUACCC AGGCCUGCAGUUCGAGAUCAUCGCCAUGUCGACCACCGGGGACAAGAUCCUGGAUACCGCCUCAGCAAG AUCGGCGAGAAUCCUUGUUCACCAAGGAGCUCGAGCAGGCCUUGGAGAAGAAUGAGGUGGAUCUGGUGG UGCAUUCUCCUGAAAGACCUGCCGACCGUCCUGCCCCCGGCUUCACGAUCGGAGCCAUCUGCAAACGGGA GAACCCCCACGACGCUUGGUGUUCACCCGAAGUUCUGGGUAAGACCUGGAAACACUGCCCGAGAAG UCCGUGGUGGGCACACGACGCCUGAGGGCGGGCGGCCAACUGCAGAGGAAGUUCCTCCACCUAGAUUCA GAAGCAUCCGGGGGAACCUGAUACCCGGCUGCGCAAGCUGGACGAGCAGCAGGAGUUCAGCGCCAUAU CCUGGCCACCGCUGGCCUGCAGAGGAUGGGAUGGCACAACCGUGUGGGCCAGAUCCUGCACCCTGAGGAG UGCAUGUACGCCGUGGGGACAGGGCGCGCUGGGCGUGGAGGUGCGCGCGAAGGACCAGGACAUCUGGACC UUGUCGGCGUGCUGCAGAUCCCGAGACGUCUGCUGAGGUGCAUCGCCGAGAGGGCCUUCUGAGGCAACCU GGAAGGGGGCUGUAGCGUCCCGUGGCUGUCCACCGCCAUGAAGGACGGCCAGCUGUACCGUACCGGGC GGCGUGUGGUCCUGGACGGCAGCGACAGCAUCCAGGAGACCAUGCAGGCCACCAUCCAGUCCCGCCCC AACACGAAGACGGCCCGAGGACGACCCCGAGCUGGUAGGCAUAACUGCCCGCAAUAUCCCGAGGGGGCC CCAACUGGGCGCCAGAACCUGGGCAUAAGCCUGGCGAACCUUGCUGUCCAAGGGCGCCAAGAACAU CUGGACGUCGCGCCGACGUGAACGACGCCAC
12	ACADVL- CO06	AUGAGCGGCAACGGCAACGCCGCGCCACCGCCGAGGAGAAUCCCCAAAUGAGGGUGAUCCGCGUGG GCACGAGGAAGAGCCAGCUGGCCCGGAUCCAGACCGACAGCUGGUGGCCACCCUGAAGGCCUCGUAUCC CGGGCUGCAGUUCGAGAUCAUCGCCAUGAGCACCACCGGGCGACAAGAUCCUGCAGACCCGCCUGUCCAAG AUCGGCGAGAAAAGCCUGUUCACCAAGGAACUGGAGCAGCACUGGAAAAGAACGAGGUAGACCUCGUGG UGCAUUCUCCUGAAGGACCUGCCACCGUGCUGCCCCAGGCUUCACCAUCGGCGCCAUCUGCAAACGGGA GAACCCCAUGAUGCCGUGGUGUUCACCCGAAGUUGUUGGCAAGACGUCUGGAGACCCUGCCCGAGAAG AGCGUAGUGGGCACACGUCGUGCGGAGGGGCCGCCAGCUGCAGAGGAAGUUCCTCCACCUUGGAGUUA GGAGCAUUAAGGGCAACCUGAACACCGGCUAGGAAGCUGGACGAGCAGCAGGAGUUCUCCGCUAUAU CCUGGCCACCGCCGGGCGCAGAGGAUGGGUGGCACAACAGGGUGGGGCAAAUCCUGCACCCTGAGAG UGCAUGUACGCCGUGGGCAGGGGGCCUGGGCGUGGAGGUGCGGCCAAGGAUCAGGACAUCUGGACC

		UGGUGGGGGUGCUGCAUGACCCCGAGACCCUGCUGAGGUGCAUCGCGGAAAGGGCCUUCUGCGCCACCU GGAGGGCGGCUGCAGCGUGCCCGUGGCGGUCACACCGCGAUGAAGGACGGCCAGCUGUACCGACCGGG GGGGUCUGGUCCUGGACGGCUCCGACAGCAUCCAGGAGACGAGGAGCAUGGAGAGCAUCCACGUGCCGGCC AACACGAGGACGGUCCCGAAGAUGAUCCCGAGCUGGUUGGCAUACCCGCAAGGAACAUCGCGGGGGCC CCAGCUGGGCGCACAGAACCUGGGCAUCUCGUGGCAACCUUGCUGCUGAGCAAGGGGGCCAAGAACAUC CUCGACGUGGGCAGGCAGCUAAACGACGCCCCAC
13	ACADVL- CO07	AUGUCCGGCAACGGCAACGCGCCGACGGCCGAGGAGAACAGCCCCAAGAUAGGGGUGAUCCGGGUGG GCACCAGAAAGUCCAGCUCGCGCAGGAUCCAGACUGAUAGCGUGGUCGCCACCCUCAAAGCCAGCUACCC GGGCCUGCAGUUUGAGAUCAUCGCCAUGUCCACAACCGGGCGACAAGAUCUUGGACACAGCGCUUGCGAAG AUCGGCGAAAAAAGCCUUCUACCAAGGAGCUGGAGCAGCCCCUGGAGAAGAAUGAGGUGGACCUUGGUGG UGCAUUCUCCUGAAGGACCUGCCACCGUGCUGCCCCCGGCUUACCAUCGGCGCCAUCUGCAAGAGGGA GAACCCACACGACGCGGUGGUUUCACCCCAAGUUCUGUGGGGAAAACCCUCGAAACCCUCCCCGAAAAG UCCGUGGUGGGCACCUCAGCCUGAGGCGGGCGGCCAGCUGCAGAGGAAGUUCACACCUUGCAGUUC GCAGCAUCCGGGGCAUUCUGAACACACAGGCUAGGAAGCUGGACGAGCAGCAGGAGUUCAGCGCCAUCAU CCUGGCCACCGGGGCCUCCAGAGGAUGGGCUGGCACAACAGGGUGGGCCAGAUCCUCCACCCCGAGGAA UGUAUGUACGCGUGGGGCAAGGCGCCUUGGGCGUGGAGGUCCGGGCGAAGGACAGGACAUCUCCGAGCC UGGUCGGCGUGCUGCAGCAUCCCGAGACGCUUCUAGGUGCAUCGCGGAACGCGCCUUCUCCGGCACCUC CGAGGGGGGCGCAGCGUGCCGGUGGCGGUCCACACCGCCAUGAAGGACGGCCAGCUGUACCUAGCCGGC GGCGUCUGGAGCCUGGACGGCUCGACUCCAUCAGGAGACGAUGCAGGGCCACCAUCCAUGUGCCAGCCC AGCAUGAGGACGGCCCGAAGACGACCCCCAACUGGUGGGCAUCACCGCCCGGAACAUCGCGAGGGGGCC UCAGCUCGCGGCACAGAUCUUGGCAUCAGCCUCGCCAAUUCUCCUGCUGAGCAAGGGAGCCAAGAACAUC CUCGACGUGGCCAGGCAGCUAAGCAGCCCCAC
14	ACADVL- CO08	AUGAGCGGCAACGGCAACGCGCGCCGACCGCCGAGGAGAAUAGCCCCAAGAUAGCGGGUGAUCCGCGUGG GCACCCGCAAGAGCCAGCUGGCGCGGAUCCAGACCGACUCCGUCGUGGCCACCCUGAAGGCCAGCUACCC CGGACUCCAGUUCGAGAUCAUCGCCAUGUGCACCACGGGGCGACAAGAUCUUGACACCGCCCUAGCAAA AUCGGAGAGAAUCCUGUUCACCAAGGAGCUGAACACGCCCUGGAAAAGAAGCAGGUGGACCUUGGUGG UGCACUCCUGAAGGACCUGCCACCGUGCUCUCCCCCGGAUUUACCAUCGGAGCCAUCUGCAAGCGGGA GAACCCUACGACGCGCGUGGUUCCAUCUCCCAAGUUCUGUGGGGAAGACGCUUGGAAACCCUGCCGGAGAAG AGCGUAGUGGGCACCAGCAGCUCGCGGAGGGCGCCAGCUGCAGAGGAAGUUCUCCACCUUGGAGUUUC GUAGCAUCCGCGGCAUUCUGAACACGCGGCGUCCGCAAGCUCGACGAGCAGCAGGAGUUCUCCGCGAUCAU ACUGGCCACCGCCGGCCUGCAGCGGAUGGGGUGGCACAACCGGGUGCGGCCAGAUCCUGCACCPCGAGGAG UGCAUGUACGCGUGGGCCAGGGGGCCUUGGGCGUGGAGGUAGGGGCCAAGGACCAGGACAUCUUGGACC UCGUGGGCGUCUUGCAGACACCCGAAACCCUGCUGAGGUGCAUCGCGGAGCGGGCCUUCUGAGGACCU GGAGGGCGGGUGCAGCGUGCCCGUGGCGGUCCACACCGCAAUGAAGGACGGGCAGCUGUACCUAGCCGGC GGGUGUCUGGAGCCUGGACGGCAGCAGUUCGAUACAGGAGACAUGCAAGCCACGAUCCACGUCUCCCGCCC AGCAGAGGACGGGCCGAGGACGACCCCCAACUGGUGCGGGAUACCGCCGGAACAUCGCGAGGGGGCC GCAGCUGGGCCGCCAGAACCUGGGCAUCUCCUGGCAACCUUGCUGCUGAGCAAGGGCGCCAAGAACAUCU CUGGAUGUGGGCCGCCAGCUGAACGAUGCCAC
15	ACADVL- CO09	AUGAGCGGGAAACGGCAACGCGCGCCGACCGCCGAGGAGAACAGUCCGAAGAUAGGGGUAUACAGGGUCG GGACCCGUAAGAGCCAGCUCGCGCAGGAUCCAGACCGACAGCGUGGUGGCCACCCUGAAGGCCAGCUAUCC AGGGCUGCAGUUCGAGAUCAUCGCCAUGUCCACCACCGGGGACAAAAUCCUGGACACCGCCCUAGCAAG AUAGGGCAGAAAGAGCCUGUUCACCAAGGAACUGGAGCAUGCCCUUGGAAAAGAAGCAGGUGGACCUUGGUCG UGCACAGCCUGAAGGACCUGCCACCGUGCUGCCCCCGGCUUACCAUCGCGCGCAUCUGCAAAAGGGA GAACCCCAACGACGCUUGGUUUCUACCCAAAAUUCUGUGGGGAAGACUUCGAGACCCUGCCCGAGAAG AGCGUGGUGGGUACUUCAGCCUGAGGAGGGCGCCAGCUGCAAAGGAAGUUCUCCCAUCUGGAGUUA GGAGCAUCCGGGGCAACUGAAUACCAGGCUUGCGCAAGCUGGAUGAGCAGCAGGAAUUCAGCGCCAUCAU CCUGGCCACCGCGGCGUGCAGAGGAUGGGCUGGCACAACAGGGUGCGGCCAGAUCCUGCACCPCGAGGAG UGCAUGUAGCCGUGGGCCAGGGUGCCUUGGGAGUGGAGGUAGGGCGAAGGACCAGGACAUCUUGACC UUGUCGGCGUGCUGCAGACACCCGAGACCCUGCUCAGGUGCAUCGCGGAGAGGGCCUUCUUGAGGCAUCU GGAGGGGGGUGUAGCGUGCCCGUGCAGUGCACAACCGCCAUGAAGGACGGCCACUUGUACCUAGCCGGC GGCGUGUGGAGCCUGGACGGCUCUGAUUCGAUCCAAGAGACCAUGCAAGCCACUUAACAGUGCCCGCCC AGCACGAGGACGGGCCGAGGAUGACCCACAGCUGGUUGGAAUACCGCCCGGAACAUCGCGAGGGGACC CCAGCUGGGCGCGCAGAUCUGGGCAUCUACUGGCAACCUUGCUGCUCUCCAAGGGGGCCAAGAACAUC CUGGACGUGGCUAGGCAGCUAAGCAGCCCCAC
16	ACADVL- CO10	AUGAGCGGCAACGGAAACGCGCGCCGACCGCCGAGGAAAACUCCCCAAAAUAGCGGGUUAUCAGGGUGG GCACCAGGAAGUCCAGCUCGCGCAGGAUACAAACCGACUCCUGGUAGCCACCCUGAAGGCCAGCUACCC CGGGCUCAGUUCGAGAUCAUCGCCAUGAGCACAACCGGGUGACAAGAUUUCUAGACACCGCCCUAGCAAG AUAGGGCAGAAAGAGCCUGUUCACCAAGGAGCUGGAACAUGCCCUUGGAAAAGAAGCAGGUGGACCUUGCUG UGCACAGCCUGAAGGACCUGCCACGGUGCUGCCACCGGCUUACGAUCGGCGCCAUCUGCAAAAGAGA GAAUCCCCAUGACGCGGUGGUUUUACCCCCAAGUUUGUGGGGAAGACCUUGGAGACCCUGCCCGAGAAG AGCGUGGUGGGGACGAGCAGCCUGCGCAGGGCGCCAGCUGCAGCGGAAGUUCUCCCAUCUGGAGUUUA GGAGCAUCAGGGGGAACUGAACACGAGGCUAGGAAGCUGGACGAGCAGCAGGAAUUCAGCGCCAUCAU CCUGGCGACCGCGGCCUGCAGGGAUGGGCUGGCACAACAGGGUGCGGCCAGAUCCUGCACCPCGAGGAG UGCAUGUAGCCGUGGGCCAGGGUGCCUUGGGAGUGGAGGUAGGGCGAAGGACCAGGACAUCUUGACC UUGUCGGCGUGCUGCAGACACCCGAGACCCUGCUCAGGUGCAUCGCGGAGAGGGCCUUCUUGAGGCAUCU GGAGGGGGGUGUAGCGUGCCCGUGCAGUGCACAACCGCCAUGAAGGACGGCCACUUGUACCUAGCCGGC GGCGUGUGGAGCCUGGACGGCUCUGAUUCGAUCCAAGAGACCAUGCAAGCCACUUAACAGUGCCCGCCC AGCACGAGGACGGGCCGAGGAUGACCCACAGCUGGUUGGAAUACCGCCCGGAACAUCGCGAGGGGACC CCAGCUGGGCGCGCAGAUCUGGGCAUCUACUGGCAACCUUGCUGCUCUCCAAGGGGGCCAAGAACAUC CUGGACGUGGCUAGGCAGCUAAGCAGCCCCAC

		AGCACGAAGACGGACCCGAGGAUGACCCCAAGCUGGUGGGGAUACACAGCCAGGAACAUCACAGGGGGCC CCAACUCGCCGCCAGAACCGUGGGGAUAGCCUGGCCAACCGUCUCUCCAAGGGCGCCAAGAACAU CUGGACGUGGCCAGGCAGCUGAACGAUGCCCAU
17	ACADVL- CO11	AUGUCCGGCAACGGGAACGCAGCCGCGACCGCCGAGGAGAACUCCCCAAGAUAGCGGUGAUUAGGGUCG GGACGAGGAAGAGCCAGCUGGCCAGGAUACAGACCGAUUCCGUGGUGGCCACCCUCAAGGCCAGCUACCC UGGGCUGCAAUUCGAGAUCAUAGCCAUGAGCACCACCGGCGAUAGAUACUGGACACCGCCUCAGCAAG AUCGGUGAGAGUCGUGUUAACCAAGGAACUGGAGCAGCCCGUGGAGAAGAAUGAAGUGGAUCUGGUCG UACACAGCUUGAAGGACCUGCCACCGUGUCGCCCCCGGAUUCACCAUCGGGGCCAUCUGCAAGAGGGA GAACCCCAACGACGCCGUGGUGUCCACCCCAAUUCGUCGGCAAGACCCUGGAAACCCUGCCCGAAAAG AGCGUGGUGGGCACCUCAGCCUGGCGCAGGGCCGCCAGCUGCAGAGGAAGUCCCCCACCUGGAGUJCA GGAGCAUACGGGGCAACCUGAACACCCGACUGAGGAAGCUGGACGAACAGCAAGAGUUCAGCGCCAUCAU CCUGGCAACCGCCGGCCUGCAGAGGAUGGGCUGGCACAACAGGGUGGGCCAGAUCCUGCACCACCGAGGAG UGUAUGUACGCCGUGAGGCCAGGGCGCCUCGAGUGGAGGUCAGGGCCAAGGACCAGGACAUCUCGACC UGGUGGGUGUGCUGCAGCAGCCCGAGACACUGCUGAGGUGCAUCGCGGAGAGGGCCUUCUGAGGCACCU CGAAGGCGGCGUCUCCGUGCCCGUGGGCCGUGCACAACCGCCAUAGAAAGACGGUCAGCUGUACUUGACCGGU GGCGUGUGGAGCCUCGACGGCAGCGACAGCAUCCAGGAGACAUCGAGGGCAACUCCAGCUGCCCGCAC AGCACGAAGACGGGCCUGAGGACGACCCCAAGCUGGUGGGAUACCGGCCAGGAUAUCCCCCGAGGGCC CCAGCUGGGCGGCCAGAACCUGGGCAUUCUCCUGGCCAACCUCCUGCUGAGCAAGGGUGCCAAGAAUUC CUGGACGUGGCCCGGCAGCUGAAUGACGCCCCAC
18	ACADVL- CO12	AUGAGCGGGAAACGGGAACGCCGCCACCGCCGAGGAGAAUCCCCAAGAUAGGGGUGAUUAGGGUAG GGACCAGGAAGAGCCAGCUGGCCAGGAUCCAGACUGACAGCGUGGUUGCCACCCUGAAGGCCAGCUACCC CGGUCUGCAGUUCGAGAUCAUCGCCAUUGCCACCACCGGUGACAAGAUUCUGGACACCGCCUGUCCAAG AUCGGAGAAAAGAGCCUGUUCACCAAGGAACUGGAACACGCCCUUGAGAGGAAGUAGAGUGGACCUUGUG UGCACUCCCUCAAGGACCUACCCACCGUGUCGCCCCCGGCUUUAACCAUCGGCGCCAUCUGCAAGAGGGA GAACCCCAUGACGCGGUGGUGUUCACCCCAAUUCGUGGGAAGACCCUGGAGACACUGCCCGAGAAG UCCGUGGUGGGCACCAGCAGCCUGAGGAGGGCCGCCAGCUGCAGAGGAAGUCCCCCAUCUGGAGUUA GGAGCAUCAGGGGCAACCUGAACACCCGGCUGAGGAAGCUGGAUGAACAGCAGGAGUUCAGCGCCAUCAU CCUGGCCACGGCGGGCCUGCAGCGGAUGGGGUGGCAUAACCGGGUGGGGCGAGAUCCUGCACCACCGAAGAA UGCAUGUAUGCGGUAGGGCAAGGGGCCUUGGGCGUCGAGGUGCGGGCCAAGGACCAGGACAUCUGGACC UGGUGGGGUGCUGCAUGAUCCCGAAACCCUGCCGUGCAUCGCGGAGAGGGCCUUCUCAGGCACCU GGAGGGCGGAUGCUCGGUGCCCGUCGCGUGCAUACCGCCAUGAAGGACGGCCAGCUCUACCUACCGGC GGCGUUGGUGCCUGGAUGGAAGCGACAGCAUUCAGGAACCAUGCAGGGCCACCAUCCAGUGCCAGCGC AGCACAGGAGACGGCCCCGAGGACGACCCCAAGCUGAGUGGGCAUACCGGCCAGGAACAUCCCCCGGGGCC CCAGCUGGCCGCGACAGAACCUGAGCAUACGCCUGGCCAACCUCCUGCUGAGCAAGGGCGCCAAGAACAU CUGGAUGUGGCCAGGCAGCUGAACGACGCCCCAC
19	ACADVL- CO13	AUGUCCGGUAACGGCAACCGCGCCGCCACCGCCGAGGAGAACAGCCCCAAGAUAGGGGUGAUCCGCGUGG GCACCAGGAAGAGCCAGCUGGCCAGGAUCCAAACCGACAGCGUGGUGGCCACGUGAAGGCCAGCUACCC AGGUCUGCAAUUCGAGAUCAUCGCCAUAGAGCACCACCGGGGACAAAUCUGGAUACCGCCUGUCCAAG AUCGGCGAGAAGAGCCUGUUCACCAAGGAGCUGGAACACGCCUUGGAGAAGAACGAGGUGGACCUGGUCG UGCACAGCCUGAAGGACCUGCCACCGUCCUGCCCCCGGGUUCACCAUCGGCGCCAUCUUGCAAGCGAGA GAACCCACACGACGCCGUGCUGUUCACCCCAAGUUCGUCGGCAAGACGUCGAGACCCUGCCCGAGAAG AGCGUGGUGCGGACGAGCUCUCCUGCGCGCGCCGCCAGCUCAGCGCAAAUUCUCCCAUUAAGAUUCC UGAGCAUCCGUGGCAACCUGAACACGCGUCUGAGGAAGCUGGAUGAACACAGGAGUUCAGCGCCAUCAU CCUGGCCACCGCCGGCCUGCAGAGGAUGGGCUGGCACAACCGUGUGGGCCAGAUCCUUAUCCCGAGGAG UGCAUGUACGUGUGGCCAGGGGGCUUGGGGUGGGAAGUCAGGGCCAAGGACCAGGACAUCUGGAUC UGGUCGGCGUGCUCACGACCCCGAGACUCUGCUGAGGUGUAUCGACAGAGAGGGCGUUCUCAGGCACCU CGAGGGCGGGUGCAGCGUCCUGUGGCGGUGCACACCGCCAUGAAGGACGGGCGAGCUGUACCUACCGGC GGCGUGUGGUGCCUGGAUGGCGAGCAGCAUCCAGGAGACGAUAGGAGGACCAUCCAGUGCCCGCCC AGCACGAGGACGGCCCCGAGGACGACCCCAACUGGUCGGCAUACCGCGCCGAACAUCUCCAGGGGGCC CCAACUGGCCGCCAGAAUCUGGGCAUUCUCCUGGCCAACCUCCUGAGCAAGGGGUGCCAAGAACAU CUCGACGUGGCCAGGCAGCUGAACGACGCCCCAC
20	ACADVL- CO14	AUGAGCGGCAACGGCAACCGCCGAGCCACCGCCGAGGAGAACUCCCCAAGAUAGGGGUGAUACAGGGUGG GCACCAGGAAGUCCCAACUGGCGCGGAUCCAGACCGACAGCGUGGUGGCCACCCUGAAGGGCUGCUACCC CGGCCUCCAGUUCGAGAUUAUCGCCAUAGAGCACCACCGGCGACAAGAUCCUGGAUACCGCCUGAGCAAG AUCGGGGAAAAAGCCUGUUCACCAAGGAGCUGGAGCAGCCCGUGGAGAAGAACGAGGUGGAUCUGGUGG UACAUCCUGAAGGAUUGCCACCGUGUCGCCCCCGGGUUCACCAUCGGGGCCAUCUGUAAAAAGGA GAAUCCCAACAGCGCCGUGGUGUUCACCCCAAGUUCUGGGCAAGACCCUGGAGAGCCUCCCGAGAGAG AGCGUGGUCGGAACAGCAGCCUGCGCGCGCCGUCAGCUACAACGCAAGUCCCCCACCUGGAGUUC GGAGCAUCAGGGGGAACCGAACACCCGUCUGCGGAAGCUCGACGAGCAGCAGGAGUUCAGCGCUAUCAU CCUGGCUACGGCCGGGUGCAGAGGAUGGGAUGGCACAACAGGGUCGGGCGAGAUCCUGCAUCCCGAGGAG UGCAUGUACGCCGUGGGCCAAGGGCGCCUCGAGUGGAAGUCAGGGCCAAGACCAAGACAUCUUGGACC UAGUCGGCGUGCUGCAUGACCCCGAGACCCUGCUCGCGUGCAUCGCGGAGAGGGCGUUCUGCGCCAUCU GGAGGGGGGUGCAGCGUGCCGGUGGCGUGCAUAGAGCUAUGAAAGAUAGGCAACUGUACCUACCGGC GGCGUCUGGAGCCUGGACGGCUGGACUCCAUAAGAGACGAUAGGAGGACCAUCCAGUGCCCGCGC AGCACGAGGAUGGGCCCGAGGACGAUCCCAAGCUGGUAGGCAUACCGCGCCGGAACAUCUCCCGGGGGCC GCAGCUGGCCGCCAGAACCUGGGGAUUCUCCUGGCCAACCUCCUGCUGUGCAAGGGGGCCAAGAACAU CUGGAUGUGCCAGGCAGCUGAACGACGCCCCAC

21	ACADVL- CO15	AUGUCCGGGAACGGGAACGCGGCCGCCACGGCGGAGGAGAACUCCCCAAAUGAGGGUGAUAAAGGGUGG GCACCCGGAAAAGCCAGCUGGCUCAAUCCAGACCGACAGCGUGGUGGCCACCCUGAAAGCCUCCUACCC CGGCCUCCAGUUUGAGAUAUCGCCAUGAGCACACCGGGCGACAAGAUCUCGACACCGCCUGAGCAAG AUCGGUGAAAAGUCGCGUUCACGAAGGAGCUGGAGCACGCCCUAGAGAAGAACGAGGUCGACCUGGUGG UCCACAGCCUGAAGGACCUGCCCACCGUCCUCCCCCGGAUUUACCAUUGGAGCCAUUAUGCAAACGGGA GAACCCCCAUGACGCCGUGGUCUCCACCCCAAGUUCGUCGCGCAAGACCCUGGAGACCCUCCCGGAGAAG UCCGUGGUGGGGACCAGCAGCCUGCGCAGGGGCCGCCAGCUGCGAGAGGAAAUUCCCCCAUCUGGAAUUA GGUCCAUACAGGGGGAACCUCAACACCCGGCUGCGGAAGCUCGACGAACAACAGGAAUUCAGCGCCAUAU CCUGGCCACCGCGGGGCGUGCAGAGGAUGGGGUGGCACAACAGGUGGGACAGAUCCUGCACCCTGAGGAG UGUAUGUACGCCGUGGGGCAGGGAGCCUGGGCGUCGAGGUAAGGGCGAAGGAUACAGGACAUCCUGGAUC UGGUGGGGGUGCUGCAGCACCCCGAGACCCUGCUCGUGCAUUGCCGAGAGGGCCUUCUGAGGCACCU GGAGGGCGGCUGUAGCGUGCCGUGGGCGUGCACACAGCGAUAAAGACGGCCAGCUGUAUCUGACCGGG GGCGUGUGGUCUCCUGAUGGCUAGACUCCAUCCAGGAGACGAUGCAGGCCACCAUCCAUUGGCCCGCCC AGCACGAGGACGGCCCCGAGGACGACCCCGAGCUCGUGGGGAUACCCGCCGUAACAUAACCCAGGGGGCC CCAGCUGGCCGCCAGAAUCUGGGGAUAGCCUCGCCAACCGUCGUGCUGAGCAAGGGCGCCAAGAACAUC CUGGACGUGGCCCGGCGAGCUGAACGACGCCCAU
22	ACADVL- CO16	AUGUCCGGCAACGGGAACGCCGCCGCCACUGCCGAGGAGAACAGCCCCAAGAUAGGGUGAUUAGGGUGG GCACCCGUAAGAGCCAGCUCGCCAGGAUCCAGACCGACAGCGUGGUCGCGACCCUUAAGGCCUCCUACCC CGGCCUGCAGUUCGAAAUAUCGCCAUGAGCACACGGGGGACAAAUCCUGGAUACCGCCUGAGCAAG AUAGGUGAGAAAGACCUGUUCACCAAGGAACUGGAGCACGCCCUUGGAAAAGAACGAAGUCGACCUCGUGG UGCACUCCUGAAGGACCUCGCCACCGUGCUGCCGCCCGGCUUACCAUCCGAGCCAUUCUGCAAGAGGGA GAACCCCCACGACGCCGUGGUAUUUACCCCCAAGUUCGUCGCAAGACCCUGGAGACCCUGCCCGAGAAG AGCGUGGUAGGCACAGCUCUCCUCCGACGGGCGGCCAACUCCAAAGGAAGUCCCCCACCUCGAGUUC GGAGCAUCAGGGGAAAUUCGAACACCAGGCUAGGAAGCUCGAUGAGCAGCAGGAGUUCUCCGCCAUCAU CCUGGCCACCGCCGGACUGCAAAGGAUGGGGUGGCACAACCGCGUUGGGCAAUCCUGCACCCCGAAGAG UGCAUGUACGCCGUGGGGCAGGGCGCACUCGGGUGGAGGUGAGGGCCAAGGACCAGGACAUCCUGGACC UCGUAGGCGUACUGCAGCAGCCCGAGACCCUGCUGCGCUGUAUCGUCUGAGAGGGCCUUCUGCGGCACCU GGAGGGCGGCUGCAGCGUGCCCGUCGCGUGGCACACCGCCAUGAAAGACGGCCAGCUCUAUCUGACGGGG GGGGUGUGGAGCCUGGACGGCAGCGACAGCAUCCAGGAGACAAUGCAGGCCACCAUCCACGUGCCCGCCC AGCACGAGGACGGGCGCAAGACGACCCCGAGCUCGUGGGCAUCACGGCCCGUAACAUCGCCAGGGGGCC GCAACUGGCCGCGCAGAACCUGGGCAUCUCGUGGGCAAUUCGUCUGCUGAGCAAGGGUGCCAAGAACAUC CUGGAUGUUGCCCAGCAGCUGAAUGACGCCAC
23	ACADVL- CO17	AUGAGCGGCAACGGUAACGCCGCCCGCAGCCGCGGAGGAGAACUCACCCAAGAUUCGGGUGAUACAGGGUAG GCACCAGGAAGAGCCAGCUGGCCAGGAUACAGACCGACAGCGUGGUGGCCACCGUGAAGGCGUCCUUAUC GGGCCUGCAAUUCGAAAUUAUCGCCAUGAGCACGACCGGGGACAAGAUCUGGACACCGCGCUGAGCAAA AUCGGCGAGAAGACCUCUUCACCAAGAGCUGGAGCAUGCCUGGAGAAGACGAGGACCUAGGACUUCG UGCAUUCACUGAAGGAUCUCCCCACCGUGCUGCCCGGCUUUAACCAUCGGAGCCAUUCGCAAGCGCGA AAACCCCCACGAUGCCGUGGUCUUCACCCCCAAGUUCGUAGGGAAGACCCUUGAGACCCUGCCGGAGAAG UCGGUGGUCGGGACCAGCAGCCUGCGGAGGGCCGCCAGCUCAGAGGAAGUUCGCGACCUUGAGUUA GGAGCAUCCGGGCAAUUCGAAUACAGGCUAGGAAGCUGGACGAGCAGCAAGAGUUCAGUGCCAUCAU CCUCGCCACCGCCGGCCUGCAGAGGAUGGGGUGGCACAACAGGGUGGGACAGAUCUUGCAUCCCGAGGAG UGCAUGUACGCCGUGGGCCAGGGCGCCUCCGCGUGGAAGUGCGGGCCAAGGACCAGGACAUUCUGGACC UGGUGGGCGUGCUGCACGAUCCGAGACCCUGCUGCAUGCAUCGCCGAGCGCCUUCUUAAGGCAACCU CGAGGGGGGUGCAGCGUGCCCGUGGCAGUGCACACCGCCAUGAAGGACGGCCAGCUGUACCUGACCGGG GGCGUGUGGAGCCUGGACGGCUCGGACAGCAUCCAGGAGACAAUGCAGGCCACCAUCCACGUGCCAGCUC AACAUGAGGAUGGCCCCGAGGACGAUCCGCGAGCUGGUGGGAUACCCGCCCGGAACAUCUAGGGGGCC CCAACUGGCAGCCAAAACCUUGGGCAUAAGCCUGGGCAAACCGUCUGCUCUCCAAGGGCGCCAAGAACAUC CUGGACGUCGCGAGGCAGCUGAACGACGCCAC
24	ACADVL- CO18	AUGAGCGGCAACGGCAACGCCGCCCGGACCCGAGGAGAAUAGCCCCAAAUGAGGGUGAUCCGGGUGG GCACCCGGAAGUCCAGCUGGCCCGGAUCCAGACCGACAGCGUGGUGGCCACCCUUAAGGCCAGCUACCC CGGCCUGCAAUUCGAAAUUAUCGCCAUGAGCACACCGGGCGACAAGAUCUGGAUACCGCCUGAGCAAG AUCGGCGAGAAAAGCCUCUUCACUAAGGAGCUGGAGCACGCCCUUGGAGAAAACGAAGUGGACCUGGUGG UCCAUAAGCCUUAAGGACCUGGCCACCGUGCUGCCCCCGGCUUACCAUUCGGCGCCAUCUGUAAGCGUGA GAACCCGCACGACGCCGUGGUGUUCACCCCCAAGUUUGUGGGGAAGACCCUCGAGACUUCUCCCGAGAAA AGCGUGGUGGGCAUAGCUCUCUGAGGAGGGCAGCCAGCUCAGAGGAAAUUCCCCCACCUCGAGUUA GGAGCAUCAGGGGCAACCUGAACACCGGCGUGAGGAAGCUGGACGAGCAGGAGUUCUCCGCAUUAU CCUGGCCACCGCGGGACUGCAGCAAUUGGGCUGGCACAACCGGGUGGGCAAUCCUGCACCCCGAGGAG UGCAUGUACGCCGUGGGCCAGGGGCAUAGGGGUGGAGGUGCGGGCAAAGGAUCAGGACAUCCUGGACC UGGUCGGGGUGCUGCAGCAGCCCGAGACCCUGCUCAGAUGCAUCGCCGAGCGGGCGUUCUCCGGCAUUU GGAGGGUGGCUUAGCGUGCCCGUCGCCUGCACACCGCCAUGAAGGACGGUCAGCUGUACCUACCGGG GGCGUGUGGUCUCCUGGACGGCAGCGACAGCAUCCAGGAGACCAUGCAGGCCACCAUCCAUUGGCCCGCC AGCACGAGGACGGCCCCGAGGACGACCCCGAGCUGGUGCGCAUACCCGCCAGGAACAUCGAGGGGGCC UCAGCUGGCCGCUAAAACCUAGGCAUCAGCCUGGGCAAACCUUCGUCAGCAAGGGCGCCAAGAAUAUC CUGGACGUCGCCAGGCAGCUGAACGACGCGCAU
25	ACADVL- CO19	AUGAGCGGCAACGGGAACGCCGCCGCCACCGCCGAGGAGAACAGCCCCAAGAUAGGGGUGAUCCGGGUGG GUACCAGGAAGAGCCAGCUGGCCCGAAUCCAGACCGAUAGCGUGGUGGCCACCGUGAAGGCCAGCUACCC CGGACUGCAGUUCGAGAUAUUCGCAUGAGCACAAACGGGGGACAAGAUCUGGACACCGCCUUCUCCAAG

		AUCGGCGAGAAAAGCCUCUUCACCAAAGAGCUGGAGCAUGCCCUUGGAGAAGAACGAGGUAGACCUCGUAG UCCAUAGCCUGAAGGACCUGCCACCGUGUCGCCCCCGGAUUCACCAUCCGGCCCAUCUGUAAGCGCGA GAACCCACACGACGCGUGGUUUCACCCCAAGUUCGUCGGAAGACACUGGAGACCCUGCCGAAAAG AGCGUGGUGGGCACCAGCAGCCUGAGGAGGGCCGCCAGCUGCAGCGCAAGUUCCCCCACCUAGAAUUA GGAGCAUCCGGGGCAACUUAACACCAGGCUAGGAAGCUGGAUGAGCAGCAGGAAUUCAGCGCCAUCAU CCUCGCCACGGCCGGGCGCAGAGGAUAGGCGUGGCACAACCGGGUGGGCCAGAUCCUGCAUCCGGAAGAG UGCAUGUACGCCGUGGGCCAGGGAGCCUUGGGCGUGGAGGUCCGGGCCAAGGAUCAAGAUUCCUGGAUC UGGUGGGGGUGCUGCAGCAGCCCGAGACCCUGCUGCGCUGCAUCGCCGAGAGGGCCUUCUGCGCCAUUC GGAGGGGGGUGCAGCGUGCCGUGGGCGUGCAUACCGCCAUAGAGGACGGCCAGCUUAUCUGACCGGC GGGGUUCUGGAGCCUGGACGGGAGCGACGAUCCAGGAGACCAUAGCAGCGACCAUCCAGUGCCCGCCC AGCACGAGGACGGCCCCGAAGAUAGCCCCAGCUGGUUGGGAUACCGCGAGGAACAUCGCCAGGGGGCC CCAGCUAGCCGCCAGAACCUUGGAUCUCCUGGCCAACCGUACUCAGCAAGGGUGCCAAGAAUUAUC CUGGAUGUCGCCAGGCAGCUGAACGACGCCCAU
26	ACADVL- CO20	AUGUCCGGCAACGGCAACGCCGCCGCCACCGCCGAGGAAAAUAGCCCCAAGAUCCGGGUGAUCCGGGUGG GAACACGAAAGUCCAGCUCGCCCCGAUCCAGACCGACAGCGUGGUGGCCACCCUGAAGGCCAGCUACCC UGGCGUGCAGUUCGAGAUCAUCGCAUAGGACACCGGGCGACAAGAUCCUGGAUACGGCCUGAGCCUAGCAAG AUCGGGGAGAAUCCUCUUCACCAAGGAACUGGAGCAGCCUUGGAGAAGAACGAGGUGGACCUGGUGG UCCACUCCUGAAGGACCUGCCACAGUGCUCCACCGGCUUCACCAUCGGCGCCAUCUGCAAACGCGA GAACCCCCACGACGCGGUGGUUUUACCCCCAAGUUCGUGGUAAGACCCUGGAGACCCUGCCGAAAAG AGCGUGGUGGGCACCUCAGCCUGAGGCGGGCCGCCAGCUGCAGAGGAAUUCGCCACCUGGAGUUC GGAGCAUCAGGGGCAACCUCAACACCAGGCUAGGAAGCUGGACGAGCAGCAGGAAUUUAGCGCGAUCAU CCUGGCCACCGCCGGCCUGCAGAGGAUAGGGUGGCACAACAGGGUGGGACAUAUCCUGCAUCCCGAGGAG UGCAUGUACGCGCGUGGGCCAAGGAGCUCUGGGCGUGGAAGUGAGGGCCAAAGGACCAUCCUGGACC UCGUGGGCGUACUGCAGCAGCCCGAGACCCUGCUGCGUGCAUUGCCGAGCGGGCCUUCUCCGUCACCU CGAGGGCGGCUGUUCCGUGCCCGUGGCCGUGCAUACGGCCAUGAAGGAUGGACAGCUGUAUCUGACGGGC GGCGUGUGGUCACUGGACGGGUCGGACUCCAUCCAGGAGACCAUAGCAGGCCACCAUCCAGUGCCCGCCC AGCACGAGGACGGCCAGAGACGACCCCCAGCUCUGUGGGAUAACCGCCCGCAAUAUCCUAGGGGGCC CCAGCUGGCCCGCCAGAACCUUGGGAUCAGCCUGGCCAACCUCCUGCUGUCCAAGGGCGCCAAGAACAU CUGGACGUCGCCGGCAGCUGAACGAGCCCAU
27	ACADVL- CO21	AUGAGCGGCAACGGGAACGCCGCCGCCAACCGCCGAGGAGAACUACCAAAGAUAGGGGUGAUCCGAGUGG GCACCAGGAAGUCGAGCUGGCCAGGAUCCAGACCGACAGCGUGGUGGCGACCCUCAAGGCCUCCUACCC CGGCCUGCAGUUCGAGAUAAUCGCAUUGCCACCACCGGGCGACAAGAUCCUGGAUACCGCCUCAGCAAG AUCGGCGAAAAGAGCCUGUUCACCAAGGAGCUCGAGCAGCUCUGGAGAAGAACGAGGUGGACCUGGUCG UGCACUCCUGAAAGACCUGCCACCGUGCUGCCCCCGGGCUUCACCAUAGGGGCCAUUCUGCAAGCGGGA GAACCCCCACGACGCGGUGGUUCCAUCCGAAGUUCGUGGGCAAGACCCUGGAGACCCUGCCCGAGAAG AGCGUGGUGGGCAGGACGACCUAAGAAGGGCCGACAGCUCUCCAGGGAAGUUCGCCACCUCGAGUUC GCAGCAUCAGAGGCAACCUGAACACCAGGCUCCGGAAGCUCGACGAGCAGGAGGUUAGCGCGCAUCAU CCUCGCCACCGCCGGCCUGCAGAGGAUAGGAUGGCAUAACCGGGUGGGGCAGAUCCUGCACCCTGAGGAG UGUAUGUACGCCGUGGGGCAGGGGGCCUGGGCGUGGAGGUGCGGGCCAAGGAUCAGGACAUCUCGACC UGGUGGGCGUGCUUCACGAUCCCGAGACCCUUCUGCGGUGCAUCGCCGAGCGGGCCUUCUGAGGCAUCU GGAGGGCGGGUGUAGUGGCCAGUGGCAGUUCACACCGCCAUGAAGGACGGCCAGCUGUACCUGACCGGU GGGUUUUGGAGCCUGGACGGGUCCGACAGCAUCCAGGAGACCAUAGCAGGCCACGAUCCAUUGCCCGCUC AGCAUGAGGACGGACAGAGACGACCCCGAGCUGGUGGGCAUCACGGCCGGAAACAACCCAGGGGGCC CCAGCUGGGCGGCCAGAACCUUGGCAUCAGCCUGGCCAACCGUCUGCUGAGCAAGGGCGCCAAGAACAU CUGGAUGUGGCGCGGCAGCUGAAUGAUCCAC
28	ACADVL- CO22	AUGUCCGGCAACGGCAACCGCGGCCGCCACCGCGGAGGAGAAUAGCCCCAAGAUCCGGGUGAUCCAGGGUGG GCACCAGGAAGAGCCAGCUGGCCAGGAUCCAGACCGACAGUGUGGUGGCCACCCUGAAGGCCAGCUACCC AGGGCUGCAGUUCGAGAUCAUUGCCAUUGCUACACCGGGCGACAAGAUCCUGACACCGCCUGAGCAAA AUCGGGGAGAAAGAGCCUGUUUACCAAAGAGCUGGAGCAGCCUUGGAGAAGAUAGGUGGACCUGGUCG UGCACUCCUGAAGGACCUGCCACCGUGCGCACCGGGCUUCACCAUCGGCGCCAUCUGUAAGAGGGA GAAUCCCCACGACGAGUUGUGUUCACCCCAAGUUCGUGGGCAAGACCCUGGAGACCCUGCCCGAGAAG AGCGUGGUGGGCAGUCGAGCCUGCAGCGGGCCGUCAGCUGCAGCGGAAUUCGCCACCUCGAGUUC GGUUCUAUAGGGGCAACCUCAACACCGCCUCAGGAAACUGGACGAGCAGCAGGAGUUCUGGCCAUCAU CCUCGCCACCGCCGGACUGCAAAGGAUAGGGUGGCACAACCGCGUGGGACAGAUCCUGCAUCCCGAGGAG UGCAUGUACGCCGUGGGCCAGGGCGCGUGGGAGUGGAGGUCAGGGCUAAAGACCAGGACAUAUCGACC UCGUGGGCGUGCUGCAGCAGCCGGAGACCCUGCGUGCAUAGCCGAGGGCCUUCUUAAGGACCCU GGAGGGCGGCGUGCUCGUGCCGUGGCCGUGCAGCACCGCCAUAGAGGACGGCCAAUUGUACCUGACCGGC GGGUGUGGUCCUGGACGGCUCAGACAGCAUACAGGAGACCAUGCAAGCGACCAUCCAGUGCCCGCCC AGCACGAGGACGGCCCCGAGGAUAGCCCCAGCUGGUGGGGAUCACAGCCCGCAACAUCGCCAGGGGGCC CCAGCUGGCCCGCCAGAACUUGGGCAUAGCCUGGCAACCUCCUGCUGUCCAAGGGAGCCAAAAUUAUC CUGGACGUGGCCAGGCAGCUGAACGACGCCCAU
29	ACADVL- CO23	AUGAGCGGCAACGGCAACCGCGGCCGCCACCGCGGAGGAGAACAGCCCCAAGAUAGGGGUGAUCCAGGGUGG GCACCAGGAAGUCCCAACUGGUAGGAUCCAGACCGACAGCGUGGUGGCCACCCUGAAGGAUCCUACCC CGGCCUCCAGUUCGAGAUAAUCGCAUAGGACCCACAGGCGACAAGAUCCUGGACACCGCCUGAGCAAG AUCGGCGAGAAGUCGUGUUAACCAAGGAGCUGGAGCAGCGCUGGAGAAGAAACGAGGUGGACCUGGUCG UGCACAGCCUGAAGGAUCCGACAGCCGAGCAGCCGCGGCUUCACCAUCGGCGCCAUCUGUAAGAGGGA GAAUCCCCACGACGAGUUGUGUUCACCCCAAGUUCGUGGGCAAGACCCUGGAGACCCUGCCCGAGAAG AGCGUGGUGGGCAGUCGAGCCUGCAGCGGGCCGUCAGCUGCAGCGGAAUUCGCCACCUCGAGUUC GGUUCUAUAGGGGCAACCUCAACACCGCCUCAGGAAACUGGACGAGCAGCAGGAGUUCUGGCCAUCAU CCUCGCCACCGCCGGACUGCAAAGGAUAGGGUGGCACAACCGCGUGGGACAGAUCCUGCAUCCCGAGGAG UGCAUGUACGCCGUGGGCCAGGGCGCGUGGGAGUGGAGGUCAGGGCUAAAGACCAGGACAUAUCGACC UCGUGGGCGUGCUGCAGCAGCCGGAGACCCUGCGUGCAUAGCCGAGGGCCUUCUUAAGGACCCU GGAGGGCGGCGUGCUCGUGCCGUGGCCGUGCAGCACCGCCAUAGAGGACGGCCAAUUGUACCUGACCGGC GGGUGUGGUCCUGGACGGCUCAGACAGCAUACAGGAGACCAUGCAAGCGACCAUCCAGUGCCCGCCC AGCACGAGGACGGCCCCGAGGAUAGCCCCAGCUGGUGGGGAUCACAGCCCGCAACAUCGCCAGGGGGCC CCAGCUGGCCCGCCAGAACUUGGGCAUAGCCUGGCAACCUCCUGCUGUCCAAGGGAGCCAAAAUUAUC CUGGACGUGGCCAGGCAGCUGAACGACGCCCAU

		AGCGUGGUGGGGACCAGCAGCCUGAGGCGGGGCCGCCAGCUGCAGAGGAAAUUCCCCAUCUGGAGUUA GAAGCAUCAGGGGCAACCUGAACACAGGCGUGAGGAAGCUCGACGAGCAGCAGGAAUUCAGCGCCAUCAU CCUCGCCACGGCCGGGCGUGCAGAGGAUAGGCGUGGCAUAACAGGGUGGGGAGAUCCUCCACCCCGAAGAG UGCAUGUACGCCGUGGGCCAGGGCGCCUCGGCGUGGAGGUAGGGGCCAAGGACCAGGACAUCCUGGACU UGGUGGGCGUCCUGCAUGACCCCGAGACCCUGCUGCGGUGCAUCGCCGAGCGGGCCUUCUCAGGCACCU GGAGGGCGGCGUGCAGCGUGCCCGUGGGCCGUACACACCGCCAUCAAAGAUAGGCCAGCUGUACCUGACCGGC GGCGUGUGGAGCCUGGACGGCUCGCACAGCAUCCAGGAGACCAUAGCAGGCCACCAUCCAUGUGCCCGCCC AGCACGAGGACGGGCCAGAGGACGAUCCCGAGCUUGUGGGAAUACACCGCCAGGAACAUCUAGGGGCCC ACAGCUGGCCGCCAAAUCUCGGGAUAAGCCUGGCUAACCGUGCUGAGCAAGGGCGCGAAGAACAU CUGGACGUGGGCGGACAGCUGAACGACGCCAU
30	ACADVL- CO24	AUGAGCGGCAACGGCAACGCGCGGCCACGGCCGAGGAGAACAGCCCCAAAUGAGGGUUAUCAGGGUGG GCACCCGCAAGUCACAGCUGGCCAGGAUCCAGACAGAUUCCUGGUGGGCCACCUUGAAGGCCUCCUACCC CGGCCUGCAGUUCGAGAUCAUAGCCAUUGUCAAACGACCGGGGACAAGAUCUCCAGACACCGCCUGAGCAAG AUCGGGGAGAAUCACUGUUCACAAAGGAACUGGAACACGCCUUGAGAAGAACGAGGUUGACCUGGUGG UCCACUCCUGAAGGACCGGCCAACCGUGCUGCCGCCGGGCUUACCAUCGGGGCCAUCUGCAAGCGGGA GAACCCCGACGACGCGUGCGUUCUCCACCCGAAAUUCUGUGGGGAAAACCCUCGAGACCCUGCCCGAGAAG AGCGUGGUGGGCACGUCAGCCUGCGGAGGGGCCGCCAGCUGCAGCGGAAGUUCUCCACCUUGGAGUUA GGUCCAUCCGGGGGAACCGAAUACGAGGCGUGAGGAAGCUGGACGAACAGCAGGAGUUCAGCGCCAUCAU CCUGGGCAGCCCGGCCUGCAGCGGAUGGGCUGGCACAAUCGGUGGGGCCAGAUCCUGCACCCTGGAGGAG UGCAUGUACGCCGUGGGGACAGGGCGCCUCGGCGUGCAGGUGCGGGCCAAGGACCAGGAUAUCCUGGACC UGGUAGGAGUCCUGCAGAUCCCGAAACCCUGCUCGGUGUAUAGCCGAGCGUGCCUUCUGAGGCAUCU GGAAGGCGGAUGCUCGUGCCCGUGGGCGUGCACACCGCAAUGAAAGACGGCCAGCUGUACCUGACCGGC GGCGUCUGGAGCCUGGACGGCUCAGACAGCAUCCAAGAGACCAUGCAGGCCACCAUCCACGUCUCCUGCCC AGCACGAGGACGGCCCCGAAGACGACCCGACGUGGUGGGCAUCACAGCCAGGAACAUCUCCCGGGGCCC GCAGCUGGCCGCCCAGAAUCUGGGAAUAGCCUGGCCAACCUCCUGCUGAGCAAGGGAGCCAAGAACAUC CUGGACGUGGCCCGGCAGCUGAACGACGCCAC
31	ACADVL- CO25	AUGAGCGGCAACGGCAACGCGGCCGCCACCGCCGAAGAGAACAGCCCCAAGAUAGGGGUGAUCAGGGUGG GCACGAGGAAGAGCCAGCUGGCCAGGAUCCAGACCGACAGCGUGGUGGCGACCCUGAAGGCCUUCUACCC UGGCCUGCAGUUGAGAUCAUCGCCAUGAGCACCACCGGGGAUAAGAUCUCCGACACCGCCUGUCCAAG AUUGGAGAAAAUUCGUGUACCAAGGAACUGGAGCAGCGCCUGGAGAAGAACGAGGUAGACCUUGUGG UGCAUAGCCUCAAGGAUCUGCCAACCGUGCUGCCCCCGGGUUAACCAUCGGCGCCAUAUGCAAGAGGGA GAACCCCGACGACGCGGUGUGUUCACCCCAAGUUCUGUGGGCAAAACCCUCGAGACGCGUCCCGAGAAG AGCGUCGUUGGCACCUCCAGCCUGAGGCGGGGCCGCCAGCUGCAGAGGAAGUUCUCCACCUCCAGAUUCA GAAGCAUCAGGGGCAACCUGAACACAGGCGUGAGGAAGCUGGACGAGCAGCAGGAGUUCAGCGCCAUCAU CCUGGCCACCGCGGCCUGCAGAGGAUGGGAUGGCACAAUAGGGUGGGCCAGAUCCUGCACCCTGGAGGAG UGCAUGUACGCGGUGCGACAGGGCGCGCUGGGGUGGAGGUCAGGGCGAAGGACAGGACAUCCUGGACC UGGUCGGGGUCCUGCAGCAGCCGAGACCCUGCUGAGGUGCAUCGCCGAGCGGGCCUUCUCCGGGACCU GGAGGGCGGUUGCAGCGUGCCAGUGGCCGUGGCACACCGCCAUAAAGGAUGGGCAGCUGUAUCUGACCGGC GGAGUGUGGAGCCUCGACGGUAGCGACUCCAUCAGGAACCAUAGCAGGCCACUAUCCACGUGCCCGCCC AGCACGAGGAUGGCCCGAGGACGAUCCCGAGCUGGUGCGGAUACGGCCCGGAACAUCUCCAGGGGCCC GCAGCUCGCCGACAGAACCUGGCAUCUCCUGGCCAACCUCCUGCUGCUCUCCAAGGGCGCCAAGAACAUC CUCGACGUGGCCCGGCAACUGAACGACGCCAC

[0240] The sequence-optimized nucleotide sequences disclosed herein are distinct from the corresponding wild type nucleotide acid sequences and from other known sequence-optimized nucleotide sequences, e.g., these sequence-optimized nucleic acids have unique compositional characteristics.

[0241] In some embodiments, the percentage of uracil or thymine nucleobases in a sequence-optimized nucleotide sequence (e.g., encoding an ACADVL polypeptide, a functional fragment, or a variant thereof) is modified (e.g., reduced) with respect to the percentage of uracil or thymine nucleobases in the reference wild-type nucleotide sequence. Such a sequence is referred to as a uracil-modified or thymine-modified sequence. The percentage of uracil or thymine content in a nucleotide sequence can be determined by dividing the number of uracils or thymines in a sequence by the total

number of nucleotides and multiplying by 100. In some embodiments, the sequence-optimized nucleotide sequence has a lower uracil or thymine content than the uracil or thymine content in the reference wild-type sequence. In some embodiments, the uracil or thymine content in a sequence-optimized nucleotide sequence of the invention is greater than the uracil or thymine content in the reference wild-type sequence and still maintain beneficial effects, e.g., increased expression and/or reduced Toll-Like Receptor (TLR) response when compared to the reference wild-type sequence.

[0242] The uracil or thymine content of wild-type ACADVL isoform 1 is about 21%. In some embodiments, the uracil or thymine content of a uracil- or thymine- modified sequence encoding an ACADVL polypeptide is less than 21%. In some embodiments, the uracil or thymine content of a uracil- or thymine-modified sequence encoding an ACADVL polypeptide of the invention is less than 20%, less than 19%, less than 18%, less than 17%, less than 16%, less than 15%, less than 14%, less than 13%, or less than 12%. In some embodiments, the uracil or thymine content is not less than 19%, 18%, 17%, 16%, 15%, 14%, 13%, 12%, or 11%. The uracil or thymine content of a sequence disclosed herein, i.e., its total uracil or thymine content is abbreviated herein as %U_{TL} or %T_{TL}.

[0243] In some embodiments, the uracil or thymine content (%U_{TL} or %T_{TL}) of a uracil- or thymine-modified sequence encoding an ACADVL polypeptide of the invention is between 11% and 21%, between 12% and 20%, between 13% and 19%, between 14% and 18%, between 14% and 17%, or between 14% and 16%.

[0244] In some embodiments, the uracil or thymine content (%U_{TL} or %T_{TL}) of a uracil- or thymine-modified sequence encoding an ACADVL polypeptide of the invention is between 13% and 17%, between 13% and 16%, or between 14% and 16%.

[0245] In a particular embodiment, the uracil or thymine content (%U_{TL} or %T_{TL}) of a uracil- or thymine modified sequence encoding an ACADVL polypeptide of the invention is between about 14% and about 16%.

[0246] A uracil- or thymine-modified sequence encoding an ACADVL polypeptide of the invention can also be described according to its uracil or thymine content relative to the uracil or thymine content in the corresponding wild-type nucleic acid sequence (%U_{WT} or %T_{WT}), or according to its uracil or thymine content relative to the theoretical minimum uracil or thymine content of a nucleic acid encoding the wild-type protein sequence (%U_{TM} or %T_{TM}).

- [0247] The phrases "uracil or thymine content relative to the uracil or thymine content in the wild type nucleic acid sequence," refers to a parameter determined by dividing the number of uracils or thymines in a sequence-optimized nucleic acid by the total number of uracils or thymines in the corresponding wild-type nucleic acid sequence and multiplying by 100. This parameter is abbreviated herein as %U_{WT} or %T_{WT}.
- [0248] In some embodiments, the %U_{WT} or %T_{WT} of a uracil- or thymine-modified sequence encoding an ACADVL polypeptide of the invention is above 50%, above 55%, above 60%, above 65%, above 70%, above 75%, above 80%, above 85%, above 90%, or above 95%.
- [0249] In some embodiments, the %U_{WT} or %T_{WT} of a uracil- or thymine modified sequence encoding an ACADVL polypeptide of the invention is between 60% and 85%, between 61% and 84%, between 62% and 83%, between 63% and 82%, between 64% and 81%, between 65% and 80%, between 66% and 79%, between 67% and 78%, between 68% and 77%, between 69% and 76%, or between 69% and 75%.
- [0250] In some embodiments, the %U_{WT} or %T_{WT} of a uracil- or thymine-modified sequence encoding an ACADVL polypeptide of the invention is between 67% and 77%, between 68% and 76%, or between 69% and 75%.
- [0251] In a particular embodiment, the %U_{WT} or %T_{WT} of a uracil- or thymine-modified sequence encoding an ACADVL polypeptide of the invention is between about 69% and about 75%.
- [0252] Uracil- or thymine- content relative to the uracil or thymine theoretical minimum, refers to a parameter determined by dividing the number of uracils or thymines in a sequence-optimized nucleotide sequence by the total number of uracils or thymines in a hypothetical nucleotide sequence in which all the codons in the hypothetical sequence are replaced with synonymous codons having the lowest possible uracil or thymine content and multiplying by 100. This parameter is abbreviated herein as %U_{TM} or %T_{TM}.
- [0253] For DNA it is recognized that thymine is present instead of uracil, and one would substitute T where U appears. Thus, all the disclosures related to, e.g., %U_{TM}, %U_{WT}, or %U_{TL}, with respect to RNA are equally applicable to %T_{TM}, %T_{WT}, or %T_{TL} with respect to DNA.
- [0254] In some embodiments, the %U_{TM} of a uracil-modified sequence encoding an ACADVL polypeptide of the invention is below 300%, below 295%, below 290%, below 285%, below 280%, below 275%, below 270%, below 265%, below 260%, below 255%,

below 250%, below 245%, below 240%, below 235%, below 230%, below 225%, below 220%, below 215%, below 200%, below 195%, below 190%, below 185%, below 180%, below 175%, below 170%, below 165%, below 160%, below 155%, below 150%, below 145%, below 140%, below 139%, below 138%, below 137%, below 136%, below 135%, below 134%, below 133%, below 132%, below 131%, below 130%, below 129%, below 128%, below 127%, below 126%, below 125%, below 124%, below 123%, below 122%, below 121%, below 120%, below 119%, below 118%, below 117%, below 116%, or below 115%.

[0255] In some embodiments, the %U_{TM} of a uracil-modified sequence encoding an ACADVL polypeptide of the invention is above 100%, above 101%, above 102%, above 103%, above 104%, above 105%, above 106%, above 107%, above 108%, above 109%, above 110%, above 111%, above 112%, above 113%, above 114%, above 115%, above 116%, above 117%, above 118%, above 119%, above 120%, above 121%, above 122%, above 123%, above 124%, above 125%, or above 126%, above 127%, above 128%, above 129%, or above 130%, above 135%, above 130%, above 131%, above 132%, above 133%, above 134%, or above 135%.

[0256] In some embodiments, the %U_{TM} of a uracil-modified sequence encoding an ACADVL polypeptide of the invention is between 122% and 124%, between 121% and 125%, between 120% and 126%, between 119% and 127%, between 118% and 128%, between 117% and 129%, between 116% and 130%, between 115% and 131%, between 114% and 132%, or between 113% and 133%.

[0257] In some embodiments, the %U_{TM} of a uracil-modified sequence encoding an ACADVL polypeptide of the invention is between about 118% and about 128%.

[0258] In some embodiments, a uracil-modified sequence encoding an ACADVL polypeptide of the invention has a reduced number of consecutive uracils with respect to the corresponding wild-type nucleic acid sequence. For example, two consecutive leucines can be encoded by the sequence CUUUUG, which includes a four uracil cluster. Such a subsequence can be substituted, e.g., with CUGCUC, which removes the uracil cluster.

[0259] Phenylalanine can be encoded by UUC or UUU. Thus, even if phenylalanines encoded by UUU are replaced by UUC, the synonymous codon still contains a uracil pair (UU). Accordingly, the number of phenylalanines in a sequence establishes a minimum number of uracil pairs (UU) that cannot be eliminated without altering the number of

phenylalanines in the encoded polypeptide. For example, if the polypeptide, e.g., wild type ACADVL isoform 1, has 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, or 35 phenylalanines, the absolute minimum number of uracil pairs (UU) that a uracil-modified sequence encoding the polypeptide, e.g., wild type ACADVL isoform 1, can contain is 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, or 35, respectively.

[0260] Wild type ACADVL isoform 1 contains 42 uracil pairs (UU), and 20 uracil triplets (UUU). In some embodiments, a uracil-modified sequence encoding an ACADVL polypeptide of the invention has a reduced number of uracil triplets (UUU) with respect to the wild-type nucleic acid sequence. In some embodiments, a uracil-modified sequence encoding an ACADVL polypeptide of the invention contains 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, 1 or no uracil triplets (UUU).

[0261] In some embodiments, a uracil-modified sequence encoding an ACADVL polypeptide has a reduced number of uracil pairs (UU) with respect to the number of uracil pairs (UU) in the wild-type nucleic acid sequence. In some embodiments, a uracil-modified sequence encoding an ACADVL polypeptide of the invention has a number of uracil pairs (UU) corresponding to the minimum possible number of uracil pairs (UU) in the wild-type nucleic acid sequence, e.g., 22 uracil pairs in the case of wild type ACADVL isoform 1.

[0262] In some embodiments, a uracil-modified sequence encoding an ACADVL polypeptide of the invention has at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 uracil pairs (UU) less than the number of uracil pairs (UU) in the wild-type nucleic acid sequence. In some embodiments, a uracil-modified sequence encoding an ACADVL polypeptide of the invention has between 22 and 35 uracil pairs (UU).

[0263] The phrase "uracil pairs (UU) relative to the uracil pairs (UU) in the wild type nucleic acid sequence," refers to a parameter determined by dividing the number of uracil pairs (UU) in a sequence-optimized nucleotide sequence by the total number of uracil pairs (UU) in the corresponding wild-type nucleotide sequence and multiplying by 100. This parameter is abbreviated herein as %UU_{wt}.

[0264] In some embodiments, a uracil-modified sequence encoding an ACADVL polypeptide of the invention has a %UU_{wt} less than 90%, less than 85%, less than 80%, less than 75%, less than 70%, less than 65%, less than 60%, less than 55%, less than 50%, less than 40%, less than 30%, or less than 20%.

- [0265] In some embodiments, a uracil-modified sequence encoding an ACADVL polypeptide has a %UU_{wt} between 45% and 90%. In a particular embodiment, a uracil-modified sequence encoding an ACADVL polypeptide of the invention has a %UU_{wt} between 52% and 84%.
- [0266] In some embodiments, the polynucleotide of the invention comprises a uracil-modified sequence encoding an ACADVL polypeptide disclosed herein. In some embodiments, the uracil-modified sequence encoding an ACADVL polypeptide comprises at least one chemically modified nucleobase, e.g., 5-methoxyuracil. In some embodiments, at least 95% of a nucleobase (e.g., uracil) in a uracil-modified sequence encoding an ACADVL polypeptide of the invention are modified nucleobases. In some embodiments, at least 95% of uracil in a uracil-modified sequence encoding an ACADVL polypeptide is 5-methoxyuracil. In some embodiments, the polynucleotide comprising a uracil-modified sequence further comprises a miRNA binding site, e.g., a miRNA binding site that binds to miR-142 and/or a miRNA binding site that binds to miR-126. In some embodiments, the polynucleotide comprising a uracil-modified sequence is formulated with a delivery agent comprising, e.g., a compound having the Formula (I), e.g., any of Compounds 1-232, e.g., Compound 18; a compound having the Formula (III), (IV), (V), or (VI), e.g., any of Compounds 233-342, e.g., Compound 236; or a compound having the Formula (VIII), e.g., any of Compounds 419-428, e.g., Compound 428, or any combination thereof. In some embodiments, the delivery agent comprises Compound 18, DSPC, Cholesterol, and Compound 428, e.g., with a mole ratio of about 50:10:38.5:1.5.
- [0267] In some embodiments, the "guanine content of the sequence optimized ORF encoding ACADVL with respect to the theoretical maximum guanine content of a nucleotide sequence encoding the ACADVL polypeptide," abbreviated as %G_{TMX} is at least 70%, at least 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, or about 100%. In some embodiments, the %G_{TMX} is between about 70% and about 80%, between about 71% and about 79%, between about 71% and about 77%, or between about 72% and about 76%.
- [0268] In some embodiments, the "cytosine content of the ORF relative to the theoretical maximum cytosine content of a nucleotide sequence encoding the ACADVL polypeptide," abbreviated as %C_{TMX}, is at least 58%, at least 59%, at least 60%, at least 65%, at least 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, or about 100%. In some embodiments, the %C_{TMX} is

between about 60% and about 80%, between about 62% and about 77%, between about 63% and about 78%, or between about 68% and about 75%.

[0269] In some embodiments, the "guanine and cytosine content (G/C) of the ORF relative to the theoretical maximum G/C content in a nucleotide sequence encoding the ACADVL polypeptide," abbreviated as %G/C_{TMX} is at least 82%, at least 83%, at least 84%, at least 85%, at least 90%, at least about 95%, or about 100%. The %G/C_{TMX} is between about 80% and about 100%, between about 85% and about 99%, between about 90% and about 96%, or between about 92% and about 95%.

[0270] In some embodiments, the "G/C content in the ORF relative to the G/C content in the corresponding wild-type ORF," abbreviated as %G/C_{WT} is at least 102%, at least 103%, at least 104%, at least 105%, at least 106%, at least 107%, at least about 110%, or at least about 112%.

[0271] In some embodiments, the average G/C content in the 3rd codon position in the ORF is at least 20%, at least 21%, at least 22%, at least 23%, at least 24%, or at least 25% higher than the average G/C content in the 3rd codon position in the corresponding wild-type ORF.

[0272] In some embodiments, the polynucleotide of the invention comprises an open reading frame (ORF) encoding an ACADVL polypeptide, wherein the ORF has been sequence optimized, and wherein each of %U_{TL}, %U_{WT}, %U_{TM}, %G_{TL}, %G_{WT}, %G_{TMX}, %C_{TL}, %C_{WT}, %C_{TMX}, %G/C_{TL}, %G/C_{WT}, or %G/C_{TMX}, alone or in a combination thereof is in a range between (i) a maximum corresponding to the parameter's maximum value (MAX) plus about 0.5, 1, 1.5, 2, 2.5, 3, 3.5, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5, or 10 standard deviations (STD DEV), and (ii) a minimum corresponding to the parameter's minimum value (MIN) less 0.5, 1, 1.5, 2, 2.5, 3, 3.5, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5, or 10 standard deviations (STD DEV).

7. Methods for Sequence Optimization

[0273] In some embodiments, a polynucleotide, e.g., mRNA, of the invention (e.g., a polynucleotide comprising a nucleotide sequence encoding an ACADVL polypeptide (e.g., the wild-type sequence, functional fragment, or variant thereof) is sequence optimized. A sequence optimized nucleotide sequence (nucleotide sequence is also referred to as "nucleic acid" herein) comprises at least one codon modification with respect to a reference sequence (e.g., a wild-type sequence encoding an ACADVL

polypeptide). Thus, in a sequence optimized nucleic acid, at least one codon is different from a corresponding codon in a reference sequence (e.g., a wild-type sequence).

[0274] In general, sequence optimized nucleic acids are generated by at least a step comprising substituting codons in a reference sequence with synonymous codons (i.e., codons that encode the same amino acid). Such substitutions can be effected, for example, by applying a codon substitution map (i.e., a table providing the codons that will encode each amino acid in the codon optimized sequence), or by applying a set of rules (e.g., if glycine is next to neutral amino acid, glycine would be encoded by a certain codon, but if it is next to a polar amino acid, it would be encoded by another codon). In addition to codon substitutions (i.e., "codon optimization") the sequence optimization methods disclosed herein comprise additional optimization steps which are not strictly directed to codon optimization such as the removal of deleterious motifs (destabilizing motif substitution). Compositions and formulations comprising these sequence optimized nucleic acids (e.g., a RNA, e.g., an mRNA) can be administered to a subject in need thereof to facilitate *in vivo* expression of functionally active ACADVL.

[0275] The recombinant expression of large molecules in cell cultures can be a challenging task with numerous limitations (e.g., poor protein expression levels, stalled translation resulting in truncated expression products, protein misfolding, etc.) These limitations can be reduced or avoided by administering the polynucleotides (e.g., a RNA, e.g., an mRNA), which encode a functionally active ACADVL or compositions or formulations comprising the same to a patient suffering from VLCADD, so the synthesis and delivery of the ACADVL polypeptide to treat VLCADD takes place endogenously.

[0276] Changing from an *in vitro* expression system (e.g., cell culture) to *in vivo* expression requires the redesign of the nucleic acid sequence encoding the therapeutic agent. Redesigning a naturally occurring gene sequence by choosing different codons without necessarily altering the encoded amino acid sequence can often lead to dramatic increases in protein expression levels (Gustafsson et al., 2004, Journal/Trends Biotechnol 22, 346-53). Variables such as codon adaptation index (CAI), mRNA secondary structures, cis-regulatory sequences, GC content and many other similar variables have been shown to somewhat correlate with protein expression levels (Villalobos et al., 2006, "Journal/BMC Bioinformatics 7, 285). However, due to the degeneracy of the genetic code, there are numerous different nucleic acid sequences that can all encode the same therapeutic agent. Each amino acid is encoded by up to six synonymous codons; and the

choice between these codons influences gene expression. In addition, codon usage (i.e., the frequency with which different organisms use codons for expressing a polypeptide sequence) differs among organisms (for example, recombinant production of human or humanized therapeutic antibodies frequently takes place in hamster cell cultures).

[0277] In some embodiments, a reference nucleic acid sequence can be sequence optimized by applying a codon map. The skilled artisan will appreciate that the T bases in the codon maps disclosed below are present in DNA, whereas the T bases would be replaced by U bases in corresponding RNAs. For example, a sequence optimized nucleic acid disclosed herein in DNA form, e.g., a vector or an in-vitro translation (IVT) template, would have its T bases transcribed as U based in its corresponding transcribed mRNA. In this respect, both sequence optimized DNA sequences (comprising T) and their corresponding RNA sequences (comprising U) are considered sequence optimized nucleic acid of the present invention. A skilled artisan would also understand that equivalent codon-maps can be generated by replaced one or more bases with non-natural bases. Thus, e.g., a TTC codon (DNA map) would correspond to a UUC codon (RNA map), which in turn can correspond to a ΨΨC codon (RNA map in which U has been replaced with pseudouridine).

[0278] In one embodiment, a reference sequence encoding ACADVL can be optimized by replacing all the codons encoding a certain amino acid with only one of the alternative codons provided in a codon map. For example, all the valines in the optimized sequence would be encoded by GTG or GTC or GTT.

[0279] Sequence optimized polynucleotides of the invention can be generated using one or more codon optimization methods, or a combination thereof. Sequence optimization methods which can be used to sequence optimize nucleic acid sequences are described in detail herein. This list of methods is not comprehensive or limiting.

[0280] It will be appreciated that the design principles and rules described for each one of the sequence optimization methods discussed below can be combined in many different ways, for example high G/C content sequence optimization for some regions or uridine content sequence optimization for other regions of the reference nucleic acid sequence, as well as targeted nucleotide mutations to minimize secondary structure throughout the sequence or to eliminate deleterious motifs.

[0281] The choice of potential combinations of sequence optimization methods can be, for example, dependent on the specific chemistry used to produce a synthetic

polynucleotide. Such a choice can also depend on characteristics of the protein encoded by the sequence optimized nucleic acid, e.g., a full sequence, a functional fragment, or a fusion protein comprising ACADVL, etc. In some embodiments, such a choice can depend on the specific tissue or cell targeted by the sequence optimized nucleic acid (e.g., a therapeutic synthetic mRNA).

[0282] The mechanisms of combining the sequence optimization methods or design rules derived from the application and analysis of the optimization methods can be either simple or complex. For example, the combination can be:

(i) *Sequential*: Each sequence optimization method or set of design rules applies to a different subsequence of the overall sequence, for example reducing uridine at codon positions 1 to 30 and then selecting high frequency codons for the remainder of the sequence;

(ii) *Hierarchical*: Several sequence optimization methods or sets of design rules are combined in a hierarchical, deterministic fashion. For example, use the most GC-rich codons, breaking ties (which are common) by choosing the most frequent of those codons.

(iii) *Multifactorial / Multiparametric*: Machine learning or other modeling techniques are used to design a single sequence that best satisfies multiple overlapping and possibly contradictory requirements. This approach would require the use of a computer applying a number of mathematical techniques, for example, genetic algorithms.

[0283] Ultimately, each one of these approaches can result in a specific set of rules which in many cases can be summarized in a single codon table, i.e., a sorted list of codons for each amino acid in the target protein (i.e., ACADVL), with a specific rule or set of rules indicating how to select a specific codon for each amino acid position.

a. Uridine Content Optimization

[0284] The presence of local high concentrations of uridine in a nucleic acid sequence can have detrimental effects on translation, e.g., slow or prematurely terminated translation, especially when modified uridine analogs are used in the production of synthetic mRNAs. Furthermore, high uridine content can also reduce the *in vivo* half-life of synthetic mRNAs due to TLR activation.

[0285] Accordingly, a nucleic acid sequence can be sequence optimized using a method comprising at least one uridine content optimization step. Such a step comprises, e.g., substituting at least one codon in the reference nucleic acid with an alternative codon to generate a uridine-modified sequence, wherein the uridine-modified sequence has at least one of the following properties:

- (i) increase or decrease in global uridine content;
- (ii) increase or decrease in local uridine content (i.e., changes in uridine content are limited to specific subsequences);
- (iii) changes in uridine distribution without altering the global uridine content;
- (iv) changes in uridine clustering (e.g., number of clusters, location of clusters, or distance between clusters); or
- (v) combinations thereof.

[0286] In some embodiments, the sequence optimization process comprises optimizing the global uridine content, i.e., optimizing the percentage of uridine nucleobases in the sequence optimized nucleic acid with respect to the percentage of uridine nucleobases in the reference nucleic acid sequence. For example, 30% of nucleobases can be uridines in the reference sequence and 10% of nucleobases can be uridines in the sequence optimized nucleic acid.

[0287] In other embodiments, the sequence optimization process comprises reducing the local uridine content in specific regions of a reference nucleic acid sequence, i.e., reducing the percentage of uridine nucleobases in a subsequence of the sequence optimized nucleic acid with respect to the percentage of uridine nucleobases in the corresponding subsequence of the reference nucleic acid sequence. For example, the reference nucleic acid sequence can have a 5'-end region (e.g., 30 codons) with a local uridine content of 30%, and the uridine content in that same region could be reduced to 10% in the sequence optimized nucleic acid.

[0288] In specific embodiments, codons can be replaced in the reference nucleic acid sequence to reduce or modify, for example, the number, size, location, or distribution of uridine clusters that could have deleterious effects on protein translation. Although as a general rule it is desirable to reduce the uridine content of the reference nucleic acid sequence, in certain embodiments the uridine content, and in particular the local uridine content, of some subsequences of the reference nucleic acid sequence can be increased.

- [0289] The reduction of uridine content to avoid adverse effects on translation can be done in combination with other optimization methods disclosed here to achieve other design goals. For example, uridine content optimization can be combined with ramp design, since using the rarest codons for most amino acids will, with a few exceptions, reduce the U content.
- [0290] In some embodiments, the uridine-modified sequence is designed to induce a lower Toll-Like Receptor (TLR) response when compared to the reference nucleic acid sequence. Several TLRs recognize and respond to nucleic acids. Double-stranded (ds)RNA, a frequent viral constituent, has been shown to activate TLR3. See Alexopoulou et al. (2001) *Nature*, 413:732–738 and Wang et al. (2004) *Nat. Med.*, 10:1366–1373. Single-stranded (ss)RNA activates TLR7. See Diebold et al. (2004) *Science* 303 :1529–1531. RNA oligonucleotides, for example RNA with phosphorothioate internucleotide linkages, are ligands of human TLR8. See Heil et al. (2004) *Science* 303:1526–1529. DNA containing unmethylated CpG motifs, characteristic of bacterial and viral DNA, activate TLR9. See Hemmi et al. (2000) *Nature*, 408: 740–745.
- [0291] As used herein, the term "TLR response" is defined as the recognition of single-stranded RNA by a TLR7 receptor, and in some embodiments encompasses the degradation of the RNA and/or physiological responses caused by the recognition of the single-stranded RNA by the receptor. Methods to determine and quantitate the binding of an RNA to a TLR7 are known in the art. Similarly, methods to determine whether an RNA has triggered a TLR7-mediated physiological response (e.g., cytokine secretion) are well known in the art. In some embodiments, a TLR response can be mediated by TLR3, TLR8, or TLR9 instead of TLR7.
- [0292] Suppression of TLR7-mediated response can be accomplished via nucleoside modification. RNA undergoes over hundred different nucleoside modifications in nature (see the RNA Modification Database, available at mods.rna.albany.edu). Human rRNA, for example, has ten times more pseudouridine (Ψ) and 25 times more 2'-O-methylated nucleosides than bacterial rRNA. Bacterial mRNA contains no nucleoside modifications, whereas mammalian mRNAs have modified nucleosides such as 5-methylcytidine (m5C), N6-methyladenosine (m6A), inosine and many 2'-O-methylated nucleosides in addition to N7-methylguanosine (m7G).

- [0293]** Uracil and ribose, the two defining features of RNA, are both necessary and sufficient for TLR7 stimulation, and short single-stranded RNA (ssRNA) act as TLR7 agonists in a sequence-independent manner as long as they contain several uridines in close proximity. See Diebold et al. (2006) Eur. J. Immunol. 36:3256-3267, which is herein incorporated by reference in its entirety. Accordingly, one or more of the optimization methods disclosed herein comprises reducing the uridine content (locally and/or locally) and/or reducing or modifying uridine clustering to reduce or to suppress a TLR7-mediated response.
- [0294]** In some embodiments, the TLR response (e.g., a response mediated by TLR7) caused by the uridine-modified sequence is at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, or at least about 100% lower than the TLR response caused by the reference nucleic acid sequence.
- [0295]** In some embodiments, the TLR response caused by the reference nucleic acid sequence is at least about 1-fold, at least about 1.1-fold, at least about 1.2-fold, at least about 1.3-fold, at least about 1.4-fold, at least about 1.5-fold, at least about 1.6-fold, at least about 1.7-fold, at least about 1.8-fold, at least about 1.9-fold, at least about 2-fold, at least about 3-fold, at least about 4-fold, at least about 5-fold, at least about 6-fold, at least about 7-fold, at least about 8-fold, at least about 9-fold, or at least about 10-fold higher than the TLR response caused by the uridine-modified sequence.
- [0296]** In some embodiments, the uridine content (average global uridine content) (absolute or relative) of the uridine-modified sequence is higher than the uridine content (absolute or relative) of the reference nucleic acid sequence. Accordingly, in some embodiments, the uridine-modified sequence contains at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, or at least about 100% more uridine than the reference nucleic acid sequence.
- [0297]** In other embodiments, the uridine content (average global uridine content) (absolute or relative) of the uridine-modified sequence is lower than the uridine content

(absolute or relative) of the reference nucleic acid sequence. Accordingly, in some embodiments, the uridine-modified sequence contains at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, or at least about 100% less uridine than the reference nucleic acid sequence.

[0298] In some embodiments, the uridine content (average global uridine content) (absolute or relative) of the uridine-modified sequence is less than 50%, 49%, 48%, 47%, 46%, 45%, 44%, 43%, 42%, 41%, 40%, 39%, 38%, 37%, 36%, 35%, 34%, 33%, 32%, 31%, 30%, 29%, 28%, 27%, 26%, 25%, 24%, 23%, 22%, 21%, 20%, 19%, 18%, 17%, 16%, 15%, 14%, 13%, 12%, 11%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2% or 1% of the total nucleobases in the uridine-modified sequence. In some embodiments, the uridine content of the uridine-modified sequence is between about 10% and about 20%. In some particular embodiments, the uridine content of the uridine-modified sequence is between about 12% and about 16%.

[0299] In some embodiments, the uridine content of the reference nucleic acid sequence can be measured using a sliding window. In some embodiments, the length of the sliding window is 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, or 40 nucleobases. In some embodiments, the sliding window is over 40 nucleobases in length. In some embodiments, the sliding window is 20 nucleobases in length. Based on the uridine content measured with a sliding window, it is possible to generate a histogram representing the uridine content throughout the length of the reference nucleic acid sequence and sequence optimized nucleic acids.

[0300] In some embodiments, a reference nucleic acid sequence can be modified to reduce or eliminate peaks in the histogram that are above or below a certain percentage value. In some embodiments, the reference nucleic acid sequence can be modified to eliminate peaks in the sliding-window representation which are above 65%, 60%, 55%, 50%, 45%, 40%, 35%, or 30% uridine. In another embodiment, the reference nucleic acid sequence can be modified so no peaks are over 30% uridine in the sequence optimized nucleic acid, as measured using a 20 nucleobase sliding window. In some embodiments, the reference nucleic acid sequence can be modified so no more or no less than a

predetermined number of peaks in the sequence optimized nucleic sequence, as measured using a 20 nucleobase sliding window, are above or below a certain threshold value. For example, in some embodiments, the reference nucleic acid sequence can be modified so no peaks or no more than 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 peaks in the sequence optimized nucleic acid are above 10%, 15%, 20%, 25% or 30% uridine. In another embodiment, the sequence optimized nucleic acid contains between 0 peaks and 2 peaks with uridine contents 30% of higher.

[0301] In some embodiments, a reference nucleic acid sequence can be sequence optimized to reduce the incidence of consecutive uridines. For example, two consecutive leucines could be encoded by the sequence CUUUUG, which would include a four uridine cluster. Such subsequence could be substituted with CUGCUC, which would effectively remove the uridine cluster. Accordingly, a reference nucleic sequence can be sequence optimized by reducing or eliminating uridine pairs (UU), uridine triplets (UUU) or uridine quadruplets (UUUU). Higher order combinations of U are not considered combinations of lower order combinations. Thus, for example, UUUU is strictly considered a quadruplet, not two consecutive U pairs; or UUUUUU is considered a sextuplet, not three consecutive U pairs, or two consecutive U triplets, etc.

[0302] In some embodiments, all uridine pairs (UU) and/or uridine triplets (UUU) and/or uridine quadruplets (UUUU) can be removed from the reference nucleic acid sequence. In other embodiments, uridine pairs (UU) and/or uridine triplets (UUU) and/or uridine quadruplets (UUUU) can be reduced below a certain threshold, e.g., no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 occurrences in the sequence optimized nucleic acid. In a particular embodiment, the sequence optimized nucleic acid contains less than 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 uridine pairs. In another particular embodiment, the sequence optimized nucleic acid contains no uridine pairs and/or triplets.

[0303] Phenylalanine codons, i.e., UUC or UUU, comprise a uridine pair or triples and therefore sequence optimization to reduce uridine content can at most reduce the phenylalanine U triplet to a phenylalanine U pair. In some embodiments, the occurrence of uridine pairs (UU) and/or uridine triplets (UUU) refers only to non-phenylalanine U pairs or triplets. Accordingly, in some embodiments, non-phenylalanine uridine pairs (UU) and/or uridine triplets (UUU) can be reduced below a certain threshold, e.g., no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 occurrences

in the sequence optimized nucleic acid. In a particular embodiment, the sequence optimized nucleic acid contains less than 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 non-phenylalanine uridine pairs and/or triplets. In another particular embodiment, the sequence optimized nucleic acid contains no non-phenylalanine uridine pairs and/or triplets.

[0304] In some embodiments, the reduction in uridine combinations (e.g., pairs, triplets, quadruplets) in the sequence optimized nucleic acid can be expressed as a percentage reduction with respect to the uridine combinations present in the reference nucleic acid sequence.

[0305] In some embodiments, a sequence optimized nucleic acid can contain about 1%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, or 65% of the total number of uridine pairs present in the reference nucleic acid sequence. In some embodiments, a sequence optimized nucleic acid can contain about 1%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, or 65% of the total number of uridine triplets present in the reference nucleic acid sequence. In some embodiments, a sequence optimized nucleic acid can contain about 1%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, or 65% of the total number of uridine quadruplets present in the reference nucleic acid sequence.

[0306] In some embodiments, a sequence optimized nucleic acid can contain about 1%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, or 65% of the total number of non-phenylalanine uridine pairs present in the reference nucleic acid sequence. In some embodiments, a sequence optimized nucleic acid can contain about 1%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, or 65% of the total number of non-phenylalanine uridine triplets present in the reference nucleic acid sequence.

[0307] In some embodiments, the uridine content in the sequence optimized sequence can be expressed with respect to the theoretical minimum uridine content in the sequence. The term "theoretical minimum uridine content" is defined as the uridine content of a nucleic acid sequence as a percentage of the sequence's length after all the codons in the

sequence have been replaced with synonymous codon with the lowest uridine content. In some embodiments, the uridine content of the sequence optimized nucleic acid is identical to the theoretical minimum uridine content of the reference sequence (e.g., a wild type sequence). In some aspects, the uridine content of the sequence optimized nucleic acid is about 90%, about 95%, about 100%, about 105%, about 110%, about 115%, about 120%, about 125%, about 130%, about 135%, about 140%, about 145%, about 150%, about 155%, about 160%, about 165%, about 170%, about 175%, about 180%, about 185%, about 190%, about 195% or about 200% of the theoretical minimum uridine content of the reference sequence (e.g., a wild type sequence).

[0308] In some embodiments, the uridine content of the sequence optimized nucleic acid is identical to the theoretical minimum uridine content of the reference sequence (e.g., a wild type sequence).

[0309] The reference nucleic acid sequence (e.g., a wild type sequence) can comprise uridine clusters which due to their number, size, location, distribution or combinations thereof have negative effects on translation. As used herein, the term "uridine cluster" refers to a subsequence in a reference nucleic acid sequence or sequence optimized nucleic sequence with contains a uridine content (usually described as a percentage) which is above a certain threshold. Thus, in certain embodiments, if a subsequence comprises more than about 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60% or 65% uridine content, such subsequence would be considered a uridine cluster.

[0310] The negative effects of uridine clusters can be, for example, eliciting a TLR7 response. Thus, in some implementations of the nucleic acid sequence optimization methods disclosed herein it is desirable to reduce the number of clusters, size of clusters, location of clusters (e.g., close to the 5' and/or 3' end of a nucleic acid sequence), distance between clusters, or distribution of uridine clusters (e.g., a certain pattern of cluster along a nucleic acid sequence, distribution of clusters with respect to secondary structure elements in the expressed product, or distribution of clusters with respect to the secondary structure of an mRNA).

[0311] In some embodiments, the reference nucleic acid sequence comprises at least one uridine cluster, wherein said uridine cluster is a subsequence of the reference nucleic acid sequence wherein the percentage of total uridine nucleobases in said subsequence is above a predetermined threshold. In some embodiments, the length of the subsequence is at least about 10, at least about 15, at least about 20, at least about 25, at least about 30, at

least about 35, at least about 40, at least about 45, at least about 50, at least about 55, at least about 60, at least about 65, at least about 70, at least about 75, at least about 80, at least about 85, at least about 90, at least about 95, or at least about 100 nucleobases. In some embodiments, the subsequence is longer than 100 nucleobases. In some embodiments, the threshold is 1%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19%, 20%, 21%, 22%, 23%, 24% or 25% uridine content. In some embodiments, the threshold is above 25%.

[0312] For example, an amino acid sequence comprising A, D, G, S and R could be encoded by the nucleic acid sequence GCU, GAU, GGU, AGU, CGU. Although such sequence does not contain any uridine pairs, triplets, or quadruplets, one third of the nucleobases would be uridines. Such a uridine cluster could be removed by using alternative codons, for example, by using GCC, GAC, GGC, AGC, and CGC, which would contain no uridines.

[0313] In other embodiments, the reference nucleic acid sequence comprises at least one uridine cluster, wherein said uridine cluster is a subsequence of the reference nucleic acid sequence wherein the percentage of uridine nucleobases of said subsequence as measured using a sliding window that is above a predetermined threshold. In some embodiments, the length of the sliding window is 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, or 40 nucleobases. In some embodiments, the sliding window is over 40 nucleobases in length. In some embodiments, the threshold is 1%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19%, 20%, 21%, 22%, 23%, 24% or 25% uridine content. In some embodiments, the threshold is above 25%.

[0314] In some embodiments, the reference nucleic acid sequence comprises at least two uridine clusters. In some embodiments, the uridine-modified sequence contains fewer uridine-rich clusters than the reference nucleic acid sequence. In some embodiments, the uridine-modified sequence contains more uridine-rich clusters than the reference nucleic acid sequence. In some embodiments, the uridine-modified sequence contains uridine-rich clusters which are shorter in length than corresponding uridine-rich clusters in the reference nucleic acid sequence. In other embodiments, the uridine-modified sequence contains uridine-rich clusters which are longer in length than the corresponding uridine-rich cluster in the reference nucleic acid sequence. See, Kariko et al. (2005) *Immunity* 23:165-175; Kormann et al. (2010) *Nature Biotechnology* 29:154-157; or Sahin et al.

(2014) Nature Reviews Drug Discovery | AOP, published online 19 September 2014m
doi:10.1038/nrd4278; all of which are herein incorporated by reference their entireties.

b. Guanine/Cytosine (G/C) Content

- [0315]** A reference nucleic acid sequence can be sequence optimized using methods comprising altering the Guanine/Cytosine (G/C) content (absolute or relative) of the reference nucleic acid sequence. Such optimization can comprise altering (e.g., increasing or decreasing) the global G/C content (absolute or relative) of the reference nucleic acid sequence; introducing local changes in G/C content in the reference nucleic acid sequence (e.g., increase or decrease G/C in selected regions or subsequences in the reference nucleic acid sequence); altering the frequency, size, and distribution of G/C clusters in the reference nucleic acid sequence, or combinations thereof.
- [0316]** In some embodiments, the sequence optimized nucleic acid encoding ACADVL comprises an overall increase in G/C content (absolute or relative) relative to the G/C content (absolute or relative) of the reference nucleic acid sequence. In some embodiments, the overall increase in G/C content (absolute or relative) is at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, or at least about 100% relative to the G/C content (absolute or relative) of the reference nucleic acid sequence.
- [0317]** In some embodiments, the sequence optimized nucleic acid encoding ACADVL comprises an overall decrease in G/C content (absolute or relative) relative to the G/C content of the reference nucleic acid sequence. In some embodiments, the overall decrease in G/C content (absolute or relative) is at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, or at least about 100% relative to the G/C content (absolute or relative) of the reference nucleic acid sequence.
- [0318]** In some embodiments, the sequence optimized nucleic acid encoding ACADVL comprises a local increase in Guanine/Cytosine (G/C) content (absolute or relative) in a

subsequence (i.e., a G/C modified subsequence) relative to the G/C content (absolute or relative) of the corresponding subsequence in the reference nucleic acid sequence. In some embodiments, the local increase in G/C content (absolute or relative) is by at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, or at least about 100% relative to the G/C content (absolute or relative) of the corresponding subsequence in the reference nucleic acid sequence.

[0319] In some embodiments, the sequence optimized nucleic acid encoding ACADVL comprises a local decrease in Guanine/Cytosine (G/C) content (absolute or relative) in a subsequence (i.e., a G/C modified subsequence) relative to the G/C content (absolute or relative) of the corresponding subsequence in the reference nucleic acid sequence. In some embodiments, the local decrease in G/C content (absolute or relative) is by at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, or at least about 100% relative to the G/C content (absolute or relative) of the corresponding subsequence in the reference nucleic acid sequence.

[0320] In some embodiments, the G/C content (absolute or relative) is increased or decreased in a subsequence which is at least about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, or 100 nucleobases in length.

[0321] In some embodiments, the G/C content (absolute or relative) is increased or decreased in a subsequence which is at least about 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, 200, 210, 220, 230, 240, 250, 260, 270, 280, 290, 300, 310, 320, 330, 340, 350, 360, 370, 380, 390, 400, 410, 420, 430, 440, 450, 460, 470, 480, 490, 500, 510, 520, 530, 540, 550, 560, 570, 580, 590, 600, 610, 620, 630, 640, 650, 660, 670, 680, 690, 700, 710, 720, 730, 740, 750, 760, 770, 780, 790, 800, 810, 820, 830, 840, 850, 860, 870, 880, 890, 900, 910, 920, 930, 940, 950, 960, 970, 980, 990, or 1000 nucleobases in length.

[0322] In some embodiments, the G/C content (absolute or relative) is increased or decreased in a subsequence which is at least about 1100, 1200, 1300, 1400, 1500, 1600, 1700, 1800, 1900, 2000, 2100, 2200, 2300, 2400, 2500, 2600, 2700, 2800, 2900, 3000,

3100, 3200, 3300, 3400, 3500, 3600, 3700, 3800, 3900, 4000, 4100, 4200, 4300, 4400, 4500, 4600, 4700, 4800, 4900, 5000, 5100, 5200, 5300, 5400, 5500, 5600, 5700, 5800, 5900, 6000, 6100, 6200, 6300, 6400, 6500, 6600, 6700, 6800, 6900, 7000, 7100, 7200, 7300, 7400, 7500, 7600, 7700, 7800, 7900, 8000, 8100, 8200, 8300, 8400, 8500, 8600, 8700, 8800, 8900, 9000, 9100, 9200, 9300, 9400, 9500, 9600, 9700, 9800, 9900, or 10000 nucleobases in length.

[0323] The increases or decreases in G and C content (absolute or relative) described herein can be conducted by replacing synonymous codons with low G/C content with synonymous codons having higher G/C content, or vice versa. For example, L has 6 synonymous codons: two of them have 2 G/C (CUC, CUG), 3 have a single G/C (UUG, CUU, CUA), and one has no G/C (UUA). So if the reference nucleic acid had a CUC codon in a certain position, G/C content at that position could be reduced by replacing CUC with any of the codons having a single G/C or the codon with no G/C.

[0324] See, U.S. Publ. Nos. US20140228558, US20050032730 A1; Gustafsson et al. (2012) Protein Expression and Purification 83: 37–46; all of which are incorporated herein by reference in their entireties.

c. Codon Frequency - Codon Usage Bias

[0325] Numerous codon optimization methods known in the art are based on the substitution of codons in a reference nucleic acid sequence with codons having higher frequencies. Thus, in some embodiments, a nucleic acid sequence encoding ACADVL disclosed herein can be sequence optimized using methods comprising the use of modifications in the frequency of use of one or more codons relative to other synonymous codons in the sequence optimized nucleic acid with respect to the frequency of use in the non-codon optimized sequence.

[0326] As used herein, the term "codon frequency" refers to codon usage bias, i.e., the differences in the frequency of occurrence of synonymous codons in coding DNA/RNA. It is generally acknowledged that codon preferences reflect a balance between mutational biases and natural selection for translational optimization. Optimal codons help to achieve faster translation rates and high accuracy. As a result of these factors, translational selection is expected to be stronger in highly expressed genes. In the field of bioinformatics and computational biology, many statistical methods have been proposed and used to analyze codon usage bias. See, e.g., Comeron & Aguadé (1998) J. Mol. Evol.

47: 268–74. Methods such as the 'frequency of optimal codons' (Fop) (Ikemura (1981) J. Mol. Biol. 151 (3): 389–409), the Relative Codon Adaptation (RCA) (Fox & Eril (2010) DNA Res. 17 (3): 185–96) or the 'Codon Adaptation Index' (CAI) (Sharp & Li (1987) Nucleic Acids Res. 15 (3): 1281–95) are used to predict gene expression levels, while methods such as the 'effective number of codons' (Nc) and Shannon entropy from information theory are used to measure codon usage evenness. Multivariate statistical methods, such as correspondence analysis and principal component analysis, are widely used to analyze variations in codon usage among genes (Suzuki et al. (2008) DNA Res. 15 (6): 357–65; Sandhu et al., In Silico Biol. 2008;8(2):187-92).

[0327] The nucleic acid sequence encoding an ACADVL polypeptide disclosed herein (e.g., a wild type nucleic acid sequence, a mutant nucleic acid sequence, a chimeric nucleic sequence, etc. which can be, for example, an mRNA), can be codon optimized using methods comprising substituting at least one codon in the reference nucleic acid sequence with an alternative codon having a higher or lower codon frequency in the synonymous codon set; wherein the resulting sequence optimized nucleic acid has at least one optimized property with respect to the reference nucleic acid sequence.

[0328] In some embodiments, at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 99%, or 100% of the codons in the reference nucleic acid sequence encoding ACADVL are substituted with alternative codons, each alternative codon having a codon frequency higher than the codon frequency of the substituted codon in the synonymous codon set.

[0329] In some embodiments, at least one codon in the reference nucleic acid sequence encoding ACADVL is substituted with an alternative codon having a codon frequency higher than the codon frequency of the substituted codon in the synonymous codon set, and at least one codon in the reference nucleic acid sequence is substituted with an alternative codon having a codon frequency lower than the codon frequency of the substituted codon in the synonymous codon set.

[0330] In some embodiments, at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%,

at least about 65%, at least about 70%, or at least about 75% of the codons in the reference nucleic acid sequence encoding ACADVL are substituted with alternative codons, each alternative codon having a codon frequency higher than the codon frequency of the substituted codon in the synonymous codon set.

[0331] In some embodiments, at least one alternative codon having a higher codon frequency has the highest codon frequency in the synonymous codon set. In other embodiments, all alternative codons having a higher codon frequency have the highest codon frequency in the synonymous codon set.

[0332] In some embodiments, at least one alternative codon having a lower codon frequency has the lowest codon frequency in the synonymous codon set. In some embodiments, all alternative codons having a higher codon frequency have the highest codon frequency in the synonymous codon set.

[0333] In some specific embodiments, at least one alternative codon has the second highest, the third highest, the fourth highest, the fifth highest or the sixth highest frequency in the synonymous codon set. In some specific embodiments, at least one alternative codon has the second lowest, the third lowest, the fourth lowest, the fifth lowest, or the sixth lowest frequency in the synonymous codon set.

[0334] Optimization based on codon frequency can be applied globally, as described above, or locally to the reference nucleic acid sequence encoding an ACADVL polypeptide. In some embodiments, when applied locally, regions of the reference nucleic acid sequence can be modified based on codon frequency, substituting all or a certain percentage of codons in a certain subsequence with codons that have higher or lower frequencies in their respective synonymous codon sets. Thus, in some embodiments, at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 99%, or 100% of the codons in a subsequence of the reference nucleic acid sequence are substituted with alternative codons, each alternative codon having a codon frequency higher than the codon frequency of the substituted codon in the synonymous codon set.

[0335] In some embodiments, at least one codon in a subsequence of the reference nucleic acid sequence encoding an ACADVL polypeptide is substituted with an

alternative codon having a codon frequency higher than the codon frequency of the substituted codon in the synonymous codon set, and at least one codon in a subsequence of the reference nucleic acid sequence is substituted with an alternative codon having a codon frequency lower than the codon frequency of the substituted codon in the synonymous codon set.

[0336] In some embodiments, at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, or at least about 75% of the codons in a subsequence of the reference nucleic acid sequence encoding an ACADVL polypeptide are substituted with alternative codons, each alternative codon having a codon frequency higher than the codon frequency of the substituted codon in the synonymous codon set.

[0337] In some embodiments, at least one alternative codon substituted in a subsequence of the reference nucleic acid sequence encoding an ACADVL polypeptide and having a higher codon frequency has the highest codon frequency in the synonymous codon set. In other embodiments, all alternative codons substituted in a subsequence of the reference nucleic acid sequence and having a lower codon frequency have the lowest codon frequency in the synonymous codon set.

[0338] In some embodiments, at least one alternative codon substituted in a subsequence of the reference nucleic acid sequence encoding an ACADVL polypeptide and having a lower codon frequency has the lowest codon frequency in the synonymous codon set. In some embodiments, all alternative codons substituted in a subsequence of the reference nucleic acid sequence and having a higher codon frequency have the highest codon frequency in the synonymous codon set.

[0339] In specific embodiments, a sequence optimized nucleic acid encoding an ACADVL polypeptide can comprise a subsequence having an overall codon frequency higher or lower than the overall codon frequency in the corresponding subsequence of the reference nucleic acid sequence at a specific location, for example, at the 5' end or 3' end of the sequence optimized nucleic acid, or within a predetermined distance from those region (e.g., at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95 or 100 codons from the 5' end or 3' end of the sequence optimized nucleic acid).

[0340] In some embodiments, an sequence optimized nucleic acid encoding an ACADVL polypeptide can comprise more than one subsequence having an overall codon frequency higher or lower than the overall codon frequency in the corresponding subsequence of the reference nucleic acid sequence. A skilled artisan would understand that subsequences with overall higher or lower overall codon frequencies can be organized in innumerable patterns, depending on whether the overall codon frequency is higher or lower, the length of the subsequence, the distance between subsequences, the location of the subsequences, etc. See, U.S. Pat. Nos. US5082767, US8126653, US7561973, US8401798; U.S. Publ. No. US 20080046192, US 20080076161; Int'l. Publ. No. WO2000018778; Welch et al. (2009) PLoS ONE 4(9): e7002; Gustafsson et al. (2012) Protein Expression and Purification 83: 37–46; Chung et al. (2012) BMC Systems Biology 6:134; all of which are incorporated herein by reference in their entireties.

d. Destabilizing Motif Substitution

[0341] There is a variety of motifs that can affect sequence optimization, which fall into various non-exclusive categories, for example:

- (i) *Primary sequence based motifs*: Motifs defined by a simple arrangement of nucleotides.
- (ii) *Structural motifs*: Motifs encoded by an arrangement of nucleotides that tends to form a certain secondary structure.
- (iii) *Local motifs*: Motifs encoded in one contiguous subsequence.
- (iv) *Distributed motifs*: Motifs encoded in two or more disjoint subsequences.
- (v) *Advantageous motifs*: Motifs which improve nucleotide structure or function.
- (vi) *Disadvantageous motifs*: Motifs with detrimental effects on nucleotide structure or function.

[0342] There are many motifs that fit into the category of disadvantageous motifs. Some examples include, for example, restriction enzyme motifs, which tend to be relatively short, exact sequences such as the restriction site motifs for *XbaI* (TCTAGA), *EcoRI* (GAATTC), *EcoRII* (CCWGG, wherein W means A or T, per the IUPAC ambiguity codes), or *HindIII* (AAGCTT); enzyme sites, which tend to be longer and based on consensus not exact sequence, such in the T7 RNA polymerase (GnnnnWnCRnCTCnCNNWnD, wherein n means any nucleotide, R means A or G, W means A or T, D means A or G or T but not C); structural motifs, such as GGGG repeats

(Kim et al. (1991) Nature 351(6324):331-2); or other motifs such as CUG-triplet repeats (Querido et al. (2014) J. Cell Sci. 124:1703-1714).

[0343] Accordingly, the nucleic acid sequence encoding an ACADVL polypeptide disclosed herein can be sequence optimized using methods comprising substituting at least one destabilizing motif in a reference nucleic acid sequence, and removing such disadvantageous motif or replacing it with an advantageous motif.

[0344] In some embodiments, the optimization process comprises identifying advantageous and/or disadvantageous motifs in the reference nucleic sequence, wherein such motifs are, e.g., specific subsequences that can cause a loss of stability in the reference nucleic acid sequence prior or during the optimization process. For example, substitution of specific bases during optimization can generate a subsequence (motif) recognized by a restriction enzyme. Accordingly, during the optimization process the appearance of disadvantageous motifs can be monitored by comparing the sequence optimized sequence with a library of motifs known to be disadvantageous. Then, the identification of disadvantageous motifs could be used as a post-hoc filter, i.e., to determine whether a certain modification which potentially could be introduced in the reference nucleic acid sequence should be actually implemented or not.

[0345] In some embodiments, the identification of disadvantageous motifs can be used prior to the application of the sequence optimization methods disclosed herein, i.e., the identification of motifs in the reference nucleic acid sequence encoding an ACADVL polypeptide and their replacement with alternative nucleic acid sequences can be used as a preprocessing step, for example, before uridine reduction.

[0346] In other embodiments, the identification of disadvantageous motifs and their removal is used as an additional sequence optimization technique integrated in a multiparametric nucleic acid optimization method comprising two or more of the sequence optimization methods disclosed herein. When used in this fashion, a disadvantageous motif identified during the optimization process would be removed, for example, by substituting the lowest possible number of nucleobases in order to preserve as closely as possible the original design principle(s) (e.g., low U, high frequency, etc.). See, e.g., U.S. Publ. Nos. US20140228558, US20050032730, or US20140228558, which are herein incorporated by reference in their entireties.

e. Limited Codon Set Optimization

[0347] In some particular embodiments, sequence optimization of a reference nucleic acid sequence encoding an ACADVL polypeptide can be conducted using a limited codon set, e.g., a codon set wherein less than the native number of codons is used to encode the 20 natural amino acids, a subset of the 20 natural amino acids, or an expanded set of amino acids including, for example, non-natural amino acids.

[0348] The genetic code is highly similar among all organisms and can be expressed in a simple table with 64 entries which would encode the 20 standard amino acids involved in protein translation plus start and stop codons. The genetic code is degenerate, i.e., in general, more than one codon specifies each amino acid. For example, the amino acid leucine is specified by the UUA, UUG, CUU, CUC, CUA, or CUG codons, while the amino acid serine is specified by UCA, UCG, UCC, UCU, AGU, or AGC codons (difference in the first, second, or third position). Native genetic codes comprise 62 codons encoding naturally occurring amino acids. Thus, in some embodiments of the methods disclosed herein optimized codon sets (genetic codes) comprising less than 62 codons to encode 20 amino acids can comprise 61, 60, 59, 58, 57, 56, 55, 54, 53, 52, 51, 50, 49, 48, 47, 46, 45, 44, 43, 42, 41, 40, 39, 38, 37, 36, 35, 34, 33, 32, 31, 30, 29, 28, 27, 26, 25, 24, 23, 22, 21, or 20 codons.

[0349] In some embodiments, the limited codon set comprises less than 20 codons. For example, if a protein contains less than 20 types of amino acids, such protein could be encoded by a codon set with less than 20 codons. Accordingly, in some embodiments, an optimized codon set comprises as many codons as different types of amino acids are present in the protein encoded by the reference nucleic acid sequence. In some embodiments, the optimized codon set comprises 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2 or even 1 codon.

[0350] In some embodiments, at least one amino acid selected from the group consisting of Ala, Arg, Asn, Asp, Cys, Gln, Glu, Gly, His, Ile, Leu, Lys, Phe, Pro, Ser, Thr, Tyr, and Val, i.e., amino acids which are naturally encoded by more than one codon, is encoded with less codons than the naturally occurring number of synonymous codons. For example, in some embodiments, Ala can be encoded in the sequence optimized nucleic acid by 3, 2 or 1 codons; Cys can be encoded in the sequence optimized nucleic acid by 1 codon; Asp can be encoded in the sequence optimized nucleic acid by 1 codon; Glu can be encoded in the sequence optimized nucleic acid by 1 codon; Phe can be encoded in the

sequence optimized nucleic acid by 1 codon; Gly can be encoded in the sequence optimized nucleic acid by 3 codons, 2 codons or 1 codon; His can be encoded in the sequence optimized nucleic acid by 1 codon; Ile can be encoded in the sequence optimized nucleic acid by 2 codons or 1 codon; Lys can be encoded in the sequence optimized nucleic acid by 1 codon; Leu can be encoded in the sequence optimized nucleic acid by 5 codons, 4 codons, 3 codons, 2 codons or 1 codon; Asn can be encoded in the sequence optimized nucleic acid by 1 codon; Pro can be encoded in the sequence optimized nucleic acid by 3 codons, 2 codons, or 1 codon; Gln can be encoded in the sequence optimized nucleic acid by 1 codon; Arg can be encoded in the sequence optimized nucleic acid by 5 codons, 4 codons, 3 codons, 2 codons, or 1 codon; Ser can be encoded in the sequence optimized nucleic acid by 5 codons, 4 codons, 3 codons, 2 codons, or 1 codon; Thr can be encoded in the sequence optimized nucleic acid by 3 codons, 2 codons, or 1 codon; Val can be encoded in the sequence optimized nucleic acid by 3 codons, 2 codons, or 1 codon; and, Tyr can be encoded in the sequence optimized nucleic acid by 1 codon.

[0351] In some embodiments, at least one amino acid selected from the group consisting of Ala, Arg, Asn, Asp, Cys, Gln, Glu, Gly, His, Ile, Leu, Lys, Phe, Pro, Ser, Thr, Tyr, and Val, i.e., amino acids which are naturally encoded by more than one codon, is encoded by a single codon in the limited codon set.

[0352] In some specific embodiments, the sequence optimized nucleic acid is a DNA and the limited codon set consists of 20 codons, wherein each codon encodes one of 20 amino acids. In some embodiments, the sequence optimized nucleic acid is a DNA and the limited codon set comprises at least one codon selected from the group consisting of GCT, GCC, GCA, and GCG; at least a codon selected from the group consisting of CGT, CGC, CGA, CGG, AGA, and AGG; at least a codon selected from AAT or ACC; at least a codon selected from GAT or GAC; at least a codon selected from TGT or TGC; at least a codon selected from CAA or CAG; at least a codon selected from GAA or GAG; at least a codon selected from the group consisting of GGT, GGC, GGA, and GGG; at least a codon selected from CAT or CAC; at least a codon selected from the group consisting of ATT, ATC, and ATA; at least a codon selected from the group consisting of TTA, TTG, CTT, CTC, CTA, and CTG; at least a codon selected from AAA or AAG; an ATG codon; at least a codon selected from TTT or TTC; at least a codon selected from the group consisting of CCT, CCC, CCA, and CCG; at least a codon selected from the group

consisting of TCT, TCC, TCA, TCG, AGT, and AGC; at least a codon selected from the group consisting of ACT, ACC, ACA, and ACG; a TGG codon; at least a codon selected from TAT or TAC; and, at least a codon selected from the group consisting of GTT, GTC, GTA, and GTG.

[0353] In other embodiments, the sequence optimized nucleic acid is an RNA (e.g., an mRNA) and the limited codon set consists of 20 codons, wherein each codon encodes one of 20 amino acids. In some embodiments, the sequence optimized nucleic acid is an RNA and the limited codon set comprises at least one codon selected from the group consisting of GCU, GCC, GCA, and GCG; at least a codon selected from the group consisting of CGU, CGC, CGA, CGG, AGA, and AGG; at least a codon selected from AAU or ACC; at least a codon selected from GAU or GAC; at least a codon selected from UGU or UGC; at least a codon selected from CAA or CAG; at least a codon selected from GAA or GAG; at least a codon selected from the group consisting of GGU, GGC, GGA, and GGG; at least a codon selected from CAU or CAC; at least a codon selected from the group consisting of AUU, AUC, and AUA; at least a codon selected from the group consisting of UUA, UUG, CUU, CUC, CUA, and CUG; at least a codon selected from AAA or AAG; an AUG codon; at least a codon selected from UUU or UUC; at least a codon selected from the group consisting of CCU, CCC, CCA, and CCG; at least a codon selected from the group consisting of UCU, UCC, UCA, UCG, AGU, and AGC; at least a codon selected from the group consisting of ACU, ACC, ACA, and ACG; a UGG codon; at least a codon selected from UAU or UAC; and, at least a codon selected from the group consisting of GUU, GUC, GUA, and GUG.

[0354] In some specific embodiments, the limited codon set has been optimized for *in vivo* expression of a sequence optimized nucleic acid (e.g., a synthetic mRNA) following administration to a certain tissue or cell.

[0355] In some embodiments, the optimized codon set (e.g., a 20 codon set encoding 20 amino acids) complies at least with one of the following properties:

- (i) the optimized codon set has a higher average G/C content than the original or native codon set; or,
 - (ii) the optimized codon set has a lower average U content than the original or native codon set; or,
 - (iii) the optimized codon set is composed of codons with the highest frequency;
- or,

(iv) the optimized codon set is composed of codons with the lowest frequency;
or,

(v) a combination thereof.

[0356] In some specific embodiments, at least one codon in the optimized codon set has the second highest, the third highest, the fourth highest, the fifth highest or the sixth highest frequency in the synonymous codon set. In some specific embodiments, at least one codon in the optimized codon set has the second lowest, the third lowest, the fourth lowest, the fifth lowest, or the sixth lowest frequency in the synonymous codon set.

[0357] As used herein, the term "native codon set" refers to the codon set used natively by the source organism to encode the reference nucleic acid sequence. As used herein, the term "original codon set" refers to the codon set used to encode the reference nucleic acid sequence before the beginning of sequence optimization, or to a codon set used to encode an optimized variant of the reference nucleic acid sequence at the beginning of a new optimization iteration when sequence optimization is applied iteratively or recursively.

[0358] In some embodiments, 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95% or 100% of codons in the codon set are those with the highest frequency. In other embodiments, 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95% or 100% of codons in the codon set are those with the lowest frequency.

[0359] In some embodiments, 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95% or 100% of codons in the codon set are those with the highest uridine content. In some embodiments, 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95% or 100% of codons in the codon set are those with the lowest uridine content.

[0360] In some embodiments, the average G/C content (absolute or relative) of the codon set is 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95% or 100% higher than the average G/C content (absolute or relative) of the original codon set. In some embodiments, the average G/C content (absolute or relative) of the codon set is 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95% or 100% lower than the average G/C content (absolute or relative) of the original codon set.

[0361] In some embodiments, the uracil content (absolute or relative) of the codon set is 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%,

80%, 85%, 90%, 95% or 100% higher than the average uracil content (absolute or relative) of the original codon set. In some embodiments, the uracil content (absolute or relative) of the codon set is 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95% or 100% lower than the average uracil content (absolute or relative) of the original codon set.

[0362] See also U.S. Appl. Publ. No. 2011/0082055, and Int'l. Publ. No. WO2000018778, both of which are incorporated herein by reference in their entireties.

8. Characterization of Sequence Optimized Nucleic Acids

[0363] In some embodiments of the invention, the polynucleotide (e.g., a RNA, e.g., an mRNA) comprising a sequence optimized nucleic acid disclosed herein encoding an ACADVL polypeptide can be tested to determine whether at least one nucleic acid sequence property (e.g., stability when exposed to nucleases) or expression property has been improved with respect to the non-sequence optimized nucleic acid.

[0364] As used herein, "expression property" refers to a property of a nucleic acid sequence either *in vivo* (e.g., translation efficacy of a synthetic mRNA after administration to a subject in need thereof) or *in vitro* (e.g., translation efficacy of a synthetic mRNA tested in an *in vitro* model system). Expression properties include but are not limited to the amount of protein produced by an mRNA encoding an ACADVL polypeptide after administration, and the amount of soluble or otherwise functional protein produced. In some embodiments, sequence optimized nucleic acids disclosed herein can be evaluated according to the viability of the cells expressing a protein encoded by a sequence optimized nucleic acid sequence (e.g., a RNA, e.g., an mRNA) encoding an ACADVL polypeptide disclosed herein.

[0365] In a particular embodiment, a plurality of sequence optimized nucleic acids disclosed herein (e.g., a RNA, e.g., an mRNA) containing codon substitutions with respect to the non-optimized reference nucleic acid sequence can be characterized functionally to measure a property of interest, for example an expression property in an *in vitro* model system, or *in vivo* in a target tissue or cell.

a. Optimization of Nucleic Acid Sequence Intrinsic Properties

[0366] In some embodiments of the invention, the desired property of the polynucleotide is an intrinsic property of the nucleic acid sequence. For example, the nucleotide

sequence (e.g., a RNA, e.g., an mRNA) can be sequence optimized for *in vivo* or *in vitro* stability. In some embodiments, the nucleotide sequence can be sequence optimized for expression in a particular target tissue or cell. In some embodiments, the nucleic acid sequence is sequence optimized to increase its plasma half life by preventing its degradation by endo and exonucleases.

[0367] In other embodiments, the nucleic acid sequence is sequence optimized to increase its resistance to hydrolysis in solution, for example, to lengthen the time that the sequence optimized nucleic acid or a pharmaceutical composition comprising the sequence optimized nucleic acid can be stored under aqueous conditions with minimal degradation.

[0368] In other embodiments, the sequence optimized nucleic acid can be optimized to increase its resistance to hydrolysis in dry storage conditions, for example, to lengthen the time that the sequence optimized nucleic acid can be stored after lyophilization with minimal degradation.

b. Nucleic Acids Sequence Optimized for Protein Expression

[0369] In some embodiments of the invention, the desired property of the polynucleotide is the level of expression of an ACADVL polypeptide encoded by a sequence optimized sequence disclosed herein. Protein expression levels can be measured using one or more expression systems. In some embodiments, expression can be measured in cell culture systems, e.g., CHO cells or HEK293 cells. In some embodiments, expression can be measured using *in vitro* expression systems prepared from extracts of living cells, e.g., rabbit reticulocyte lysates, or *in vitro* expression systems prepared by assembly of purified individual components. In other embodiments, the protein expression is measured in an *in vivo* system, e.g., mouse, rabbit, monkey, etc.

[0370] In some embodiments, protein expression in solution form can be desirable. Accordingly, in some embodiments, a reference sequence can be sequence optimized to yield a sequence optimized nucleic acid sequence having optimized levels of expressed proteins in soluble form. Levels of protein expression and other properties such as solubility, levels of aggregation, and the presence of truncation products (i.e., fragments due to proteolysis, hydrolysis, or defective translation) can be measured according to methods known in the art, for example, using electrophoresis (e.g., native or SDS-PAGE) or chromatographic methods (e.g., HPLC, size exclusion chromatography, etc.).

c. Optimization of Target Tissue or Target Cell Viability

[0371] In some embodiments, the expression of heterologous therapeutic proteins encoded by a nucleic acid sequence can have deleterious effects in the target tissue or cell, reducing protein yield, or reducing the quality of the expressed product (e.g., due to the presence of protein fragments or precipitation of the expressed protein in inclusion bodies), or causing toxicity.

[0372] Accordingly, in some embodiments of the invention, the sequence optimization of a nucleic acid sequence disclosed herein, e.g., a nucleic acid sequence encoding an ACADVL polypeptide, can be used to increase the viability of target cells expressing the protein encoded by the sequence optimized nucleic acid.

[0373] Heterologous protein expression can also be deleterious to cells transfected with a nucleic acid sequence for autologous or heterologous transplantation. Accordingly, in some embodiments of the present disclosure the sequence optimization of a nucleic acid sequence disclosed herein can be used to increase the viability of target cells expressing the protein encoded by the sequence optimized nucleic acid sequence. Changes in cell or tissue viability, toxicity, and other physiological reaction can be measured according to methods known in the art.

d. Reduction of Immune and/or Inflammatory Response

[0374] In some cases, the administration of a sequence optimized nucleic acid encoding ACADVL polypeptide or a functional fragment thereof can trigger an immune response, which could be caused by (i) the therapeutic agent (e.g., an mRNA encoding an ACADVL polypeptide), or (ii) the expression product of such therapeutic agent (e.g., the ACADVL polypeptide encoded by the mRNA), or (iv) a combination thereof. Accordingly, in some embodiments of the present disclosure the sequence optimization of nucleic acid sequence (e.g., an mRNA) disclosed herein can be used to decrease an immune or inflammatory response triggered by the administration of a nucleic acid encoding an ACADVL polypeptide or by the expression product of ACADVL encoded by such nucleic acid.

[0375] In some aspects, an inflammatory response can be measured by detecting increased levels of one or more inflammatory cytokines using methods known in the art, e.g., ELISA. The term "inflammatory cytokine" refers to cytokines that are elevated in an inflammatory response. Examples of inflammatory cytokines include interleukin-6 (IL-6),

CXCL1 (chemokine (C-X-C motif) ligand 1; also known as GRO α , interferon- γ (IFN γ), tumor necrosis factor α (TNF α), interferon γ -induced protein 10 (IP-10), or granulocyte-colony stimulating factor (G-CSF). The term inflammatory cytokines includes also other cytokines associated with inflammatory responses known in the art, e.g., interleukin-1 (IL-1), interleukin-8 (IL-8), interleukin-12 (IL-12), interleukin-13 (IL-13), interferon α (IFN- α), etc.

9. Modified Nucleotide Sequences Encoding ACADVL Polypeptides

[0376] In some embodiments, the polynucleotide (e.g., a RNA, e.g., an mRNA) of the invention comprises a chemically modified nucleobase, e.g., 5-methoxyuracil. In some embodiments, the mRNA is a uracil-modified sequence comprising an ORF encoding an ACADVL polypeptide, wherein the mRNA comprises a chemically modified nucleobase, e.g., 5-methoxyuracil.

[0377] In certain aspects of the invention, when the 5-methoxyuracil base is connected to a ribose sugar, as it is in polynucleotides, the resulting modified nucleoside or nucleotide is referred to as 5-methoxyuridine. In some embodiments, uracil in the polynucleotide is at least about 25%, at least about 30%, at least about 40%, at least about 50%, at least about 60%, at least about 70%, at least about 80%, at least 90%, at least 95%, at least 99%, or about 100% 5-methoxyuracil. In one embodiment, uracil in the polynucleotide is at least 95% 5-methoxyuracil. In another embodiment, uracil in the polynucleotide is 100% 5-methoxyuracil.

[0378] In embodiments where uracil in the polynucleotide is at least 95% 5-methoxyuracil, overall uracil content can be adjusted such that an mRNA provides suitable protein expression levels while inducing little to no immune response. In some embodiments, the uracil content of the ORF is between about 105% and about 140%, about 110% and about 140%, 115% and about 140%, about 105% and about 130%, about 110% and about 130%, about 115% and about 135%, about 105% and about 135%, about 110% and about 135%, or about 115% and about 130% of the theoretical minimum uracil content in the corresponding wild-type ORF (%U_{tm}). In other embodiments, the uracil content of the ORF is between about 110% and about 140% or between 118% and 128% of the %U_{tm}. In some embodiments, the uracil content of the ORF encoding an ACADVL polypeptide is about 115%, about 120%, about 125%, about 130%, about

135%, about 140%, about 145%, or about 150% of the %U_{tm}. In this context, the term "uracil" can refer to 5-methoxyuracil and/or naturally occurring uracil.

[0379] In some embodiments, the uracil content in the ORF of the mRNA encoding an ACADVL polypeptide of the invention is less than about 50%, about 40%, about 30%, or about 20% of the total nucleobase content in the ORF. In some embodiments, the uracil content in the ORF is between about 15 % and about 25% of the total nucleobase content in the ORF. In other embodiments, the uracil content in the ORF is between about 20% and about 30% of the total nucleobase content in the ORF. In one embodiment, the uracil content in the ORF of the mRNA encoding an ACADVL polypeptide is less than about 21% of the total nucleobase content in the open reading frame. In this context, the term "uracil" can refer to 5-methoxyuracil and/or naturally occurring uracil.

[0380] In further embodiments, the ORF of the mRNA encoding an ACADVL polypeptide having 5-methoxyuracil and adjusted uracil content has increased Cytosine (C), Guanine (G), or Guanine/Cytosine (G/C) content (absolute or relative). In some embodiments, the overall increase in C, G, or G/C content (absolute or relative) of the ORF is at least about 2%, at least about 3%, at least about 4%, at least about 5%, at least about 6%, at least about 7%, at least about 8%, at least about 9%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 50%, at least about 60%, at least about 70%, at least about 80%, at least about 90%, at least about 95%, or at least about 100% relative to the G/C content (absolute or relative) of the wild-type ORF. In some embodiments, the G, the C, or the G/C content in the ORF is less than about 100%, less than about 95%, less than about 90%, less than about 85%, less than about 80%, less than about 75%, or less than about 70% of the theoretical maximum G, C, or G/C content of the corresponding wild type nucleotide sequence encoding the ACADVL polypeptide (%G_{TMX}; %C_{TMX}, or %G/C_{TMX}). In other embodiments, the G, the C, or the G/C content in the ORF is between about 65% and about 98%, between about 72% and about 76%, between about 68% and about 75%, or between about 92% and about 95% of the %G_{TMX}, %C_{TMX}, or %G/C_{TMX}. In some embodiments, the increases in G and/or C content (absolute or relative) described herein can be conducted by replacing synonymous codons with low G, C, or G/C content with synonymous codons having higher G, C, or G/C content. In other embodiments, the increase in G and/or C content (absolute or relative) is

conducted by replacing a codon ending with U with a synonymous codon ending with G or C.

[0381] In further embodiments, the ORF of the mRNA encoding an ACADVL polypeptide of the invention comprises 5-methoxyuracil and has an adjusted uracil content containing less uracil pairs (UU) and/or uracil triplets (UUU) and/or uracil quadruplets (UUUU) than the corresponding wild-type nucleotide sequence encoding the ACADVL polypeptide. In some embodiments, the ORF of the mRNA encoding an ACADVL polypeptide of the invention contains no uracil pairs and/or uracil triplets and/or uracil quadruplets. In some embodiments, uracil pairs and/or uracil triplets and/or uracil quadruplets are reduced below a certain threshold, e.g., no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 occurrences in the ORF of the mRNA encoding the ACADVL polypeptide. In a particular embodiment, the ORF of the mRNA encoding the ACADVL polypeptide of the invention contains less than 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 non-phenylalanine uracil pairs and/or triplets. In another embodiment, the ORF of the mRNA encoding the ACADVL polypeptide contains no non-phenylalanine uracil pairs and/or triplets.

[0382] In further embodiments, the ORF of the mRNA encoding an ACADVL polypeptide of the invention comprises 5-methoxyuracil and has an adjusted uracil content containing less uracil-rich clusters than the corresponding wild-type nucleotide sequence encoding the ACADVL polypeptide. In some embodiments, the ORF of the mRNA encoding the ACADVL polypeptide of the invention contains uracil-rich clusters that are shorter in length than corresponding uracil-rich clusters in the corresponding wild-type nucleotide sequence encoding the ACADVL polypeptide.

[0383] In further embodiments, alternative lower frequency codons are employed. At least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 99%, or 100% of the codons in the ACADVL polypeptide-encoding ORF of the 5-methoxyuracil-comprising mRNA are substituted with alternative codons, each alternative codon having a codon frequency lower than the codon frequency of the substituted codon in the synonymous codon set. The ORF also has adjusted uracil content, as described above. In some embodiments, at least one codon

in the ORF of the mRNA encoding the ACADVL polypeptide is substituted with an alternative codon having a codon frequency lower than the codon frequency of the substituted codon in the synonymous codon set.

[0384] In some embodiments, the adjusted uracil content, ACADVL polypeptide-encoding ORF of the 5-methoxyuracil-comprising mRNA exhibits expression levels of ACADVL when administered to a mammalian cell that are higher than expression levels of ACADVL from the corresponding wild-type mRNA. In other embodiments, the expression levels of ACADVL when administered to a mammalian cell are increased relative to a corresponding mRNA containing at least 95% 5-methoxyuracil and having a uracil content of about 160%, about 170%, about 180%, about 190%, or about 200% of the theoretical minimum. In yet other embodiments, the expression levels of ACADVL when administered to a mammalian cell are increased relative to a corresponding mRNA, wherein at least about 50%, at least about 60%, at least about 70%, at least about 80%, at least about 90%, or about 100% of uracils are 1-methylpseudouracil or pseudouracil. In some embodiments, the mammalian cell is a mouse cell, a rat cell, or a rabbit cell. In other embodiments, the mammalian cell is a monkey cell or a human cell. In some embodiments, the human cell is a HeLa cell, a BJ fibroblast cell, or a peripheral blood mononuclear cell (PBMC). In some embodiments, ACADVL is expressed when the mRNA is administered to a mammalian cell *in vivo*. In some embodiments, the mRNA is administered to mice, rabbits, rats, monkeys, or humans. In one embodiment, mice are null mice. In some embodiments, the mRNA is administered to mice in an amount of about 0.01 mg/kg, about 0.05 mg/kg, about 0.1 mg/kg, or about 0.15 mg/kg. In some embodiments, the mRNA is administered intravenously or intramuscularly. In other embodiments, the ACADVL polypeptide is expressed when the mRNA is administered to a mammalian cell *in vitro*. In some embodiments, the expression is increased by at least about 2-fold, at least about 5-fold, at least about 10-fold, at least about 50-fold, at least about 500-fold, at least about 1500-fold, or at least about 3000-fold. In other embodiments, the expression is increased by at least about 10%, about 20%, about 30%, about 40%, about 50%, 60%, about 70%, about 80%, about 90%, or about 100%.

[0385] In some embodiments, adjusted uracil content, ACADVL polypeptide-encoding ORF of the 5-methoxyuracil-comprising mRNA exhibits increased stability. In some embodiments, the mRNA exhibits increased stability in a cell relative to the stability of a corresponding wild-type mRNA under the same conditions. In some embodiments, the

mRNA exhibits increased stability including resistance to nucleases, thermal stability, and/or increased stabilization of secondary structure. In some embodiments, increased stability exhibited by the mRNA is measured by determining the half-life of the mRNA (e.g., in a plasma, serum, cell, or tissue sample) and/or determining the area under the curve (AUC) of the protein expression by the mRNA over time (e.g., in vitro or in vivo). An mRNA is identified as having increased stability if the half-life and/or the AUC is greater than the half-life and/or the AUC of a corresponding wild-type mRNA under the same conditions.

[0386] In some embodiments, the mRNA of the present invention induces a detectably lower immune response (e.g., innate or acquired) relative to the immune response induced by a corresponding wild-type mRNA under the same conditions. In other embodiments, the mRNA of the present disclosure induces a detectably lower immune response (e.g., innate or acquired) relative to the immune response induced by an mRNA that encodes for an ACADVL polypeptide but does not comprise 5-methoxyuracil under the same conditions, or relative to the immune response induced by an mRNA that encodes for an ACADVL polypeptide and that comprises 5-methoxyuracil but that does not have adjusted uracil content under the same conditions. The innate immune response can be manifested by increased expression of pro-inflammatory cytokines, activation of intracellular PRRs (RIG-I, MDA5, etc.), cell death, and/or termination or reduction in protein translation. In some embodiments, a reduction in the innate immune response can be measured by expression or activity level of Type 1 interferons (e.g., IFN- α , IFN- β , IFN- κ , IFN- δ , IFN- ϵ , IFN- τ , IFN- ω , and IFN- ζ) or the expression of interferon-regulated genes such as the toll-like receptors (e.g., TLR7 and TLR8), and/or by decreased cell death following one or more administrations of the mRNA of the invention into a cell.

[0387] In some embodiments, the expression of Type-1 interferons by a mammalian cell in response to the mRNA of the present disclosure is reduced by at least 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, 99%, 99.9%, or greater than 99.9% relative to a corresponding wild-type mRNA, to an mRNA that encodes an ACADVL polypeptide but does not comprise 5-methoxyuracil, or to an mRNA that encodes an ACADVL polypeptide and that comprises 5-methoxyuracil but that does not have adjusted uracil content. In some embodiments, the interferon is IFN- β . In some embodiments, cell death frequency caused by administration of mRNA of the present disclosure to a mammalian cell is 10%, 25%, 50%, 75%, 85%, 90%, 95%, or over 95% less than the cell death

frequency observed with a corresponding wild-type mRNA, an mRNA that encodes for an ACADVL polypeptide but does not comprise 5-methoxyuracil, or an mRNA that encodes for an ACADVL polypeptide and that comprises 5-methoxyuracil but that does not have adjusted uracil content. In some embodiments, the mammalian cell is a BJ fibroblast cell. In other embodiments, the mammalian cell is a splenocyte. In some embodiments, the mammalian cell is that of a mouse or a rat. In other embodiments, the mammalian cell is that of a human. In one embodiment, the mRNA of the present disclosure does not substantially induce an innate immune response of a mammalian cell into which the mRNA is introduced.

[0388] In some embodiments, the polynucleotide is an mRNA that comprises an ORF that encodes an ACADVL polypeptide, wherein uracil in the mRNA is at least about 95% 5-methoxyuracil, wherein the uracil content of the ORF is between about 115% and about 135% of the theoretical minimum uracil content in the corresponding wild-type ORF, and wherein the uracil content in the ORF encoding the ACADVL polypeptide is less than about 21% of the total nucleobase content in the ORF. In some embodiments, the ORF that encodes the ACADVL polypeptide is further modified to increase G/C content of the ORF (absolute or relative) by at least about 40%, as compared to the corresponding wild-type ORF. In yet other embodiments, the ORF encoding the ACADVL polypeptide contains less than 20 non-phenylalanine uracil pairs and/or triplets. In some embodiments, at least one codon in the ORF of the mRNA encoding the ACADVL polypeptide is further substituted with an alternative codon having a codon frequency lower than the codon frequency of the substituted codon in the synonymous codon set. In some embodiments, the expression of the ACADVL polypeptide encoded by an mRNA comprising an ORF wherein uracil in the mRNA is at least about 95% 5-methoxyuracil, and wherein the uracil content of the ORF is between about 115% and about 130% of the theoretical minimum uracil content in the corresponding wild-type ORF, is increased by at least about 10-fold when compared to expression of the ACADVL polypeptide from the corresponding wild-type mRNA. In some embodiments, the mRNA comprises an open ORF wherein uracil in the mRNA is at least about 95% 5-methoxyuracil, and wherein the uracil content of the ORF is between about 115% and about 130% of the theoretical minimum uracil content in the corresponding wild-type ORF, and wherein the mRNA does not substantially induce an innate immune response of a mammalian cell into which the mRNA is introduced.

10. Methods for Modifying Polynucleotides

[0389] The invention includes modified polynucleotides comprising a polynucleotide described herein (e.g., a polynucleotide, e.g., an mRNA, comprising a nucleotide sequence encoding an ACADVL polypeptide). The modified polynucleotides can be chemically modified and/or structurally modified. When the polynucleotides of the present invention are chemically and/or structurally modified the polynucleotides can be referred to as "modified polynucleotides."

[0390] The present disclosure provides for modified nucleosides and nucleotides of a polynucleotide (e.g., RNA polynucleotides, such as mRNA polynucleotides) encoding an ACADVL polypeptide. A "nucleoside" refers to a compound containing a sugar molecule (e.g., a pentose or ribose) or a derivative thereof in combination with an organic base (e.g., a purine or pyrimidine) or a derivative thereof (also referred to herein as "nucleobase"). A "nucleotide" refers to a nucleoside including a phosphate group. Modified nucleotides can be synthesized by any useful method, such as, for example, chemically, enzymatically, or recombinantly, to include one or more modified or non-natural nucleosides. Polynucleotides can comprise a region or regions of linked nucleosides. Such regions can have variable backbone linkages. The linkages can be standard phosphodiester linkages, in which case the polynucleotides would comprise regions of nucleotides.

[0391] The modified polynucleotides disclosed herein can comprise various distinct modifications. In some embodiments, the modified polynucleotides contain one, two, or more (optionally different) nucleoside or nucleotide modifications. In some embodiments, a modified polynucleotide, introduced to a cell can exhibit one or more desirable properties, e.g., improved protein expression, reduced immunogenicity, or reduced degradation in the cell, as compared to an unmodified polynucleotide.

a. Structural Modifications

[0392] In some embodiments, a polynucleotide of the present invention (e.g., a polynucleotide comprising a nucleotide sequence encoding an ACADVL polypeptide) is structurally modified. As used herein, a "structural" modification is one in which two or more linked nucleosides are inserted, deleted, duplicated, inverted or randomized in a polynucleotide without significant chemical modification to the nucleotides themselves. Because chemical bonds will necessarily be broken and reformed to effect a structural

modification, structural modifications are of a chemical nature and hence are chemical modifications. However, structural modifications will result in a different sequence of nucleotides. For example, the polynucleotide "ATCG" can be chemically modified to "AT-5meC-G". The same polynucleotide can be structurally modified from "ATCG" to "ATCCCG". Here, the dinucleotide "CC" has been inserted, resulting in a structural modification to the polynucleotide.

b. Chemical Modifications

[0393] In some embodiments, the polynucleotides of the present invention are chemically modified. As used herein in reference to a polynucleotide, the terms "chemical modification" or, as appropriate, "chemically modified" refer to modification with respect to adenosine (A), guanosine (G), uridine (U), thymidine (T) or cytidine (C) ribo- or deoxyribonucleosides in one or more of their position, pattern, percent or population. Generally, herein, these terms are not intended to refer to the ribonucleotide modifications in naturally occurring 5'-terminal mRNA cap moieties.

[0394] In some embodiments, the polynucleotides of the present invention can have a uniform chemical modification of all or any of the same nucleoside type or a population of modifications produced by mere downward titration of the same starting modification in all or any of the same nucleoside type, or a measured percent of a chemical modification of all any of the same nucleoside type but with random incorporation, such as where all uridines are replaced by a uridine analog, *e.g.*, pseudouridine or 5-methoxyuridine. In another embodiment, the polynucleotides can have a uniform chemical modification of two, three, or four of the same nucleoside type throughout the entire polynucleotide (such as all uridines and all cytosines, etc. are modified in the same way).

[0395] Modified nucleotide base pairing encompasses not only the standard adenosine-thymine, adenosine-uracil, or guanosine-cytosine base pairs, but also base pairs formed between nucleotides and/or modified nucleotides comprising non-standard or modified bases, wherein the arrangement of hydrogen bond donors and hydrogen bond acceptors permits hydrogen bonding between a non-standard base and a standard base or between two complementary non-standard base structures. One example of such non-standard base pairing is the base pairing between the modified nucleotide inosine and adenine, cytosine

or uracil. Any combination of base/sugar or linker can be incorporated into polynucleotides of the present disclosure.

[0396] The skilled artisan will appreciate that, except where otherwise noted, polynucleotide sequences set forth in the instant application will recite "T"s in a representative DNA sequence but where the sequence represents RNA, the "T"s would be substituted for "U"s.

[0397] Modifications of polynucleotides (*e.g.*, RNA polynucleotides, such as mRNA polynucleotides) that are useful in the compositions, methods and synthetic processes of the present disclosure include, but are not limited to the following nucleotides, nucleosides, and nucleobases: 2-methylthio-N6-(*cis*-hydroxyisopentenyl)adenosine; 2-methylthio-N6-methyladenosine; 2-methylthio-N6-threonyl carbamoyladenosine; N6-glyciny carbamoyladenosine; N6-isopentenyladenosine; N6-methyladenosine; N6-threonyl carbamoyladenosine; 1,2'-O-dimethyladenosine; 1-methyladenosine; 2'-O-methyladenosine; 2'-O-ribosyladenosine (phosphate); 2-methyladenosine; 2-methylthio-N6 isopentenyladenosine; 2-methylthio-N6-hydroxynorvalyl carbamoyladenosine; 2'-O-methyladenosine; 2'-O-ribosyladenosine (phosphate); Isopentenyladenosine; N6-(*cis*-hydroxyisopentenyl)adenosine; N6,2'-O-dimethyladenosine; N6,2'-O-dimethyladenosine; N6,N6,2'-O-trimethyladenosine; N6,N6-dimethyladenosine; N6-acetyladenosine; N6-hydroxynorvalyl carbamoyladenosine; N6-methyl-N6-threonyl carbamoyladenosine; 2-methyladenosine; 2-methylthio-N6-isopentenyladenosine; 7-deaza-adenosine; N1-methyl-adenosine; N6, N6 (dimethyl)adenine; N6-*cis*-hydroxy-isopentenyl-adenosine; α -thio-adenosine; 2 (amino)adenine; 2 (aminopropyl)adenine; 2 (methylthio) N6 (isopentenyl)adenine; 2-(alkyl)adenine; 2-(aminoalkyl)adenine; 2-(aminopropyl)adenine; 2-(halo)adenine; 2-(halo)adenine; 2-(propyl)adenine; 2'-Amino-2'-deoxy-ATP; 2'-Azido-2'-deoxy-ATP; 2'-Deoxy-2'-a-aminoadenosine TP; 2'-Deoxy-2'-a-azidoadenosine TP; 6 (alkyl)adenine; 6 (methyl)adenine; 6-(alkyl)adenine; 6-(methyl)adenine; 7 (deaza)adenine; 8 (alkenyl)adenine; 8 (alkynyl)adenine; 8 (amino)adenine; 8 (thioalkyl)adenine; 8-(alkenyl)adenine; 8-(alkyl)adenine; 8-(alkynyl)adenine; 8-(amino)adenine; 8-(halo)adenine; 8-(hydroxyl)adenine; 8-(thioalkyl)adenine; 8-(thiol)adenine; 8-azido-adenosine; aza adenine; deaza adenine; N6 (methyl)adenine; N6-(isopentyl)adenine; 7-deaza-8-aza-adenosine; 7-methyladenine; 1-Dezaadenosine TP; 2'Fluoro-N6-Bz-deoxyadenosine TP; 2'-OMe-2-Amino-ATP; 2'O-methyl-N6-Bz-deoxyadenosine TP; 2'-a-Ethynyladenosine TP; 2-aminoadenine; 2-Aminoadenosine TP;

2-Amino-ATP; 2'-a-Trifluoromethyladenosine TP; 2-Azidoadenosine TP; 2'-b-Ethynyladenosine TP; 2-Bromoadenosine TP; 2'-b-Trifluoromethyladenosine TP; 2-Chloroadenosine TP; 2'-Deoxy-2',2'-difluoroadenosine TP; 2'-Deoxy-2'-a-mercaptopadenosine TP; 2'-Deoxy-2'-a-thiomethoxyadenosine TP; 2'-Deoxy-2'-b-aminoadenosine TP; 2'-Deoxy-2'-b-azidoadenosine TP; 2'-Deoxy-2'-b-bromoadenosine TP; 2'-Deoxy-2'-b-chloroadenosine TP; 2'-Deoxy-2'-b-fluoroadenosine TP; 2'-Deoxy-2'-b-iodoadenosine TP; 2'-Deoxy-2'-b-mercaptopadenosine TP; 2'-Deoxy-2'-b-thiomethoxyadenosine TP; 2-Fluoroadenosine TP; 2-Iodoadenosine TP; 2-Mercaptopadenosine TP; 2-methoxy-adenine; 2-methylthio-adenine; 2-Trifluoromethyladenosine TP; 3-Deaza-3-bromoadenosine TP; 3-Deaza-3-chloroadenosine TP; 3-Deaza-3-fluoroadenosine TP; 3-Deaza-3-iodoadenosine TP; 3-Deazaadenosine TP; 4'-Azidoadenosine TP; 4'-Carbocyclic adenosine TP; 4'-Ethynyladenosine TP; 5'-Homo-adenosine TP; 8-Aza-ATP; 8-bromo-adenosine TP; 8-Trifluoromethyladenosine TP; 9-Deazaadenosine TP; 2-aminopurine; 7-deaza-2,6-diaminopurine; 7-deaza-8-aza-2,6-diaminopurine; 7-deaza-8-aza-2-aminopurine; 2,6-diaminopurine; 7-deaza-8-aza-adenine, 7-deaza-2-aminopurine; 2-thiocytidine; 3-methylcytidine; 5-formylcytidine; 5-hydroxymethylcytidine; 5-methylcytidine; N4-acetylcytidine; 2'-O-methylcytidine; 2'-O-methylcytidine; 5,2'-O-dimethylcytidine; 5-formyl-2'-O-methylcytidine; Lysidine; N4,2'-O-dimethylcytidine; N4-acetyl-2'-O-methylcytidine; N4-methylcytidine; N4,N4-Dimethyl-2'-OMe-Cytidine TP; 4-methylcytidine; 5-aza-cytidine; Pseudo-iso-cytidine; pyrrolo-cytidine; α -thio-cytidine; 2-(thio)cytosine; 2'-Amino-2'-deoxy-CTP; 2'-Azido-2'-deoxy-CTP; 2'-Deoxy-2'-a-aminocytidine TP; 2'-Deoxy-2'-a-azidocytidine TP; 3 (deaza) 5 (aza)cytosine; 3 (methyl)cytosine; 3-(alkyl)cytosine; 3-(deaza) 5 (aza)cytosine; 3-(methyl)cytidine; 4,2'-O-dimethylcytidine; 5 (halo)cytosine; 5 (methyl)cytosine; 5 (propynyl)cytosine; 5 (trifluoromethyl)cytosine; 5-(alkyl)cytosine; 5-(alkynyl)cytosine; 5-(halo)cytosine; 5-(propynyl)cytosine; 5-(trifluoromethyl)cytosine; 5-bromo-cytidine; 5-iodo-cytidine; 5-propynyl cytosine; 6-(azo)cytosine; 6-aza-cytidine; aza cytosine; deaza cytosine; N4 (acetyl)cytosine; 1-methyl-1-deaza-pseudoisocytidine; 1-methyl-pseudoisocytidine; 2-methoxy-5-methyl-cytidine; 2-methoxy-cytidine; 2-thio-5-methyl-cytidine; 4-methoxy-1-methyl-pseudoisocytidine; 4-methoxy-pseudoisocytidine; 4-thio-1-methyl-1-deaza-pseudoisocytidine; 4-thio-1-methyl-pseudoisocytidine; 4-thio-pseudoisocytidine; 5-aza-zebularine; 5-methyl-zebularine; pyrrolo-pseudoisocytidine; Zebularine; (E)-5-(2-Bromo-

vinyl)cytidine TP; 2,2'-anhydro-cytidine TP hydrochloride; 2'Fluor-N4-Bz-cytidine TP; 2'Fluoro-N4-Acetyl-cytidine TP; 2'-O-Methyl-N4-Acetyl-cytidine TP; 2'O-methyl-N4-Bz-cytidine TP; 2'-a-Ethynylcytidine TP; 2'-a-Trifluoromethylcytidine TP; 2'-b-Ethynylcytidine TP; 2'-b-Trifluoromethylcytidine TP; 2'-Deoxy-2',2'-difluorocytidine TP; 2'-Deoxy-2'-a-mercaptopcytidine TP; 2'-Deoxy-2'-a-thiomethoxycytidine TP; 2'-Deoxy-2'-b-aminocytidine TP; 2'-Deoxy-2'-b-azidocytidine TP; 2'-Deoxy-2'-b-bromocytidine TP; 2'-Deoxy-2'-b-chlorocytidine TP; 2'-Deoxy-2'-b-fluorocytidine TP; 2'-Deoxy-2'-b-iodocytidine TP; 2'-Deoxy-2'-b-mercaptopcytidine TP; 2'-Deoxy-2'-b-thiomethoxycytidine TP; 2'-O-Methyl-5-(1-propynyl)cytidine TP; 3'-Ethynylcytidine TP; 4'-Azidocytidine TP; 4'-Carbocyclic cytidine TP; 4'-Ethynylcytidine TP; 5-(1-Propynyl)ara-cytidine TP; 5-(2-Chloro-phenyl)-2-thiocytidine TP; 5-(4-Amino-phenyl)-2-thiocytidine TP; 5-Aminoallyl-CTP; 5-Cyanocytidine TP; 5-Ethynylara-cytidine TP; 5-Ethynylcytidine TP; 5'-Homocytidine TP; 5-Methoxycytidine TP; 5-Trifluoromethyl-Cytidine TP; N4-Amino-cytidine TP; N4-Benzoyl-cytidine TP; Pseudoisocytidine; 7-methylguanosine; N2,2'-O-dimethylguanosine; N2-methylguanosine; Wyosine; 1,2'-O-dimethylguanosine; 1-methylguanosine; 2'-O-methylguanosine; 2'-O-ribosylguanosine (phosphate); 2'-O-methylguanosine; 2'-O-ribosylguanosine (phosphate); 7-aminomethyl-7-deazaguanosine; 7-cyano-7-deazaguanosine; Archaeosine; Methylwyosine; N2,7-dimethylguanosine; N2,N2,2'-O-trimethylguanosine; N2,N2,7-trimethylguanosine; N2,N2-dimethylguanosine; N2,7,2'-O-trimethylguanosine; 6-thio-guanosine; 7-deaza-guanosine; 8-oxo-guanosine; N1-methyl-guanosine; α -thio-guanosine; 2 (propyl)guanine; 2-(alkyl)guanine; 2'-Amino-2'-deoxy-GTP; 2'-Azido-2'-deoxy-GTP; 2'-Deoxy-2'-a-aminoguanosine TP; 2'-Deoxy-2'-a-azidoguanosine TP; 6 (methyl)guanine; 6-(alkyl)guanine; 6-(methyl)guanine; 6-methyl-guanosine; 7 (alkyl)guanine; 7 (deaza)guanine; 7 (methyl)guanine; 7-(alkyl)guanine; 7-(deaza)guanine; 7-(methyl)guanine; 8 (alkyl)guanine; 8 (alkynyl)guanine; 8 (halo)guanine; 8 (thioalkyl)guanine; 8-(alkenyl)guanine; 8-(alkyl)guanine; 8-(alkynyl)guanine; 8-(amino)guanine; 8-(halo)guanine; 8-(hydroxyl)guanine; 8-(thioalkyl)guanine; 8-(thiol)guanine; aza guanine; deaza guanine; N (methyl)guanine; N-(methyl)guanine; 1-methyl-6-thio-guanosine; 6-methoxy-guanosine; 6-thio-7-deaza-8-aza-guanosine; 6-thio-7-deaza-guanosine; 6-thio-7-methyl-guanosine; 7-deaza-8-aza-guanosine; 7-methyl-8-oxo-guanosine; N2,N2-dimethyl-6-thio-guanosine; N2-methyl-6-thio-guanosine; 1-Me-GTP; 2'Fluoro-N2-isobutyl-guanosine TP; 2'O-methyl-N2-isobutyl-guanosine TP; 2'-a-

Ethynylguanosine TP; 2'-a-Trifluoromethylguanosine TP; 2'-b-Ethynylguanosine TP; 2'-b-Trifluoromethylguanosine TP; 2'-Deoxy-2',2'-difluoroguanosine TP; 2'-Deoxy-2'-a-mercaptoguanosine TP; 2'-Deoxy-2'-a-thiomethoxyguanosine TP; 2'-Deoxy-2'-b-aminoguanosine TP; 2'-Deoxy-2'-b-azidoguanosine TP; 2'-Deoxy-2'-b-bromoguanosine TP; 2'-Deoxy-2'-b-chloroguanosine TP; 2'-Deoxy-2'-b-fluoroguanosine TP; 2'-Deoxy-2'-b-iodoguanosine TP; 2'-Deoxy-2'-b-mercaptoguanosine TP; 2'-Deoxy-2'-b-thiomethoxyguanosine TP; 4'-Azidoguanosine TP; 4'-Carbocyclic guanosine TP; 4'-Ethynylguanosine TP; 5'-Homo-guanosine TP; 8-bromo-guanosine TP; 9-Deazaguanosine TP; N2-isobutyl-guanosine TP; 1-methylinosine; Inosine; 1,2'-O-dimethylinosine; 2'-O-methylinosine; 7-methylinosine; 2'-O-methylinosine; Epoxyqueuosine; galactosyl-queuosine; Mannosylqueuosine; Queuosine; allyamino-thymidine; aza thymidine; deaza thymidine; deoxy-thymidine; 2'-O-methyluridine; 2-thiouridine; 3-methyluridine; 5-carboxymethyluridine; 5-hydroxyuridine; 5-methyluridine; 5-taurinomethyl-2-thiouridine; 5-taurinomethyluridine; Dihydrouridine; Pseudouridine; (3-(3-amino-3-carboxypropyl)uridine; 1-methyl-3-(3-amino-5-carboxypropyl)pseudouridine; 1-methylpseudouridine; 1-ethyl-pseudouridine; 2'-O-methyluridine; 2'-O-methylpseudouridine; 2'-O-methyluridine; 2-thio-2'-O-methyluridine; 3-(3-amino-3-carboxypropyl)uridine; 3,2'-O-dimethyluridine; 3-Methyl-pseudo-Uridine TP; 4-thiouridine; 5-(carboxyhydroxymethyl)uridine; 5-(carboxyhydroxymethyl)uridine methyl ester; 5,2'-O-dimethyluridine; 5,6-dihydro-uridine; 5-aminomethyl-2-thiouridine; 5-carbamoylmethyl-2'-O-methyluridine; 5-carbamoylmethyluridine; 5-carboxyhydroxymethyluridine; 5-carboxyhydroxymethyluridine methyl ester; 5-carboxymethylaminomethyl-2'-O-methyluridine; 5-carboxymethylaminomethyl-2-thiouridine; 5-carboxymethylaminomethyl-2-thiouridine; 5-carboxymethylaminomethyluridine; 5-Carbamoylmethyluridine TP; 5-methoxycarbonylmethyl-2'-O-methyluridine; 5-methoxycarbonylmethyl-2-thiouridine; 5-methoxycarbonylmethyluridine; 5-methyluridine, 5-methoxyuridine; 5-methyl-2-thiouridine; 5-methylaminomethyl-2-selenouridine; 5-methylaminomethyl-2-thiouridine; 5-methylaminomethyluridine; 5-Methyldihydrouridine; 5-Oxyacetic acid- Uridine TP; 5-Oxyacetic acid-methyl ester-Uridine TP; N1-methyl-pseudo-uracil; N1-ethyl-pseudo-uracil; uridine 5-oxyacetic acid; uridine 5-oxyacetic acid methyl ester; 3-(3-Amino-3-carboxypropyl)-Uridine TP; 5-(iso-Pentenylaminomethyl)- 2-thiouridine TP; 5-(iso-Pentenylaminomethyl)-2'-O-

methyluridine TP; 5-(iso-Pentenylaminomethyl)uridine TP; 5-propynyl uracil; α -thio-uridine; 1 (aminoalkylamino-carbonylethylenyl)-2(thio)-pseudouracil; 1 (aminoalkylaminocarbonylethylenyl)-2,4-(dithio)pseudouracil; 1 (aminoalkylaminocarbonylethylenyl)-4 (thio)pseudouracil; 1 (aminoalkylaminocarbonylethylenyl)-pseudouracil; 1 (aminocarbonylethylenyl)-2(thio)-pseudouracil; 1 (aminocarbonylethylenyl)-2,4-(dithio)pseudouracil; 1 (aminocarbonylethylenyl)-4 (thio)pseudouracil; 1 (aminocarbonylethylenyl)-pseudouracil; 1 substituted 2(thio)-pseudouracil; 1 substituted 2,4-(dithio)pseudouracil; 1 substituted 4 (thio)pseudouracil; 1 substituted pseudouracil; 1-(aminoalkylamino-carbonylethylenyl)-2-(thio)-pseudouracil; 1-Methyl-3-(3-amino-3-carboxypropyl) pseudouridine TP; 1-Methyl-3-(3-amino-3-carboxypropyl)pseudo-UTP; 1-Methyl-pseudo-UTP; 1-Ethyl-pseudo-UTP; 2 (thio)pseudouracil; 2' deoxy uridine; 2' fluorouridine; 2-(thio)uracil; 2,4-(dithio)pseudouracil; 2' methyl, 2'amino, 2'azido, 2'fluoroguanosine; 2'-Amino-2'-deoxy-UTP; 2'-Azido-2'-deoxy-UTP; 2'-Azido-deoxyuridine TP; 2'-O-methylpseudouridine; 2' deoxy uridine; 2' fluorouridine; 2'-Deoxy-2'-a-aminouridine TP; 2'-Deoxy-2'-a-azidouridine TP; 2-methylpseudouridine; 3 (3 amino-3 carboxypropyl)uracil; 4 (thio)pseudouracil; 4-(thio)pseudouracil; 4-(thio)uracil; 4-thiouracil; 5 (1,3-diazole-1-alkyl)uracil; 5 (2-aminopropyl)uracil; 5 (aminoalkyl)uracil; 5 (dimethylaminoalkyl)uracil; 5 (guanidiniumalkyl)uracil; 5 (methoxycarbonylmethyl)-2-(thio)uracil; 5 (methoxycarbonyl-methyl)uracil; 5 (methyl) 2 (thio)uracil; 5 (methyl) 2,4 (dithio)uracil; 5 (methyl) 4 (thio)uracil; 5 (methylaminomethyl)-2 (thio)uracil; 5 (methylaminomethyl)-2,4 (dithio)uracil; 5 (methylaminomethyl)-4 (thio)uracil; 5 (propynyl)uracil; 5 (trifluoromethyl)uracil; 5-(2-aminopropyl)uracil; 5-(alkyl)-2-(thio)pseudouracil; 5-(alkyl)-2,4 (dithio)pseudouracil; 5-(alkyl)-4 (thio)pseudouracil; 5-(alkyl)pseudouracil; 5-(alkyl)uracil; 5-(alkynyl)uracil; 5-(allylamino)uracil; 5-(cyanoalkyl)uracil; 5-(dialkylaminoalkyl)uracil; 5-(dimethylaminoalkyl)uracil; 5-(guanidiniumalkyl)uracil; 5-(halo)uracil; 5-(1,3-diazole-1-alkyl)uracil; 5-(methoxy)uracil; 5-(methoxycarbonylmethyl)-2-(thio)uracil; 5-(methoxycarbonyl-methyl)uracil; 5-(methyl) 2(thio)uracil; 5-(methyl) 2,4 (dithio)uracil; 5-(methyl) 4 (thio)uracil; 5-(methyl)-2-(thio)pseudouracil; 5-(methyl)-2,4 (dithio)pseudouracil; 5-(methyl)-4 (thio)pseudouracil; 5-(methyl)pseudouracil; 5-(methylaminomethyl)-2 (thio)uracil; 5-(methylaminomethyl)-2,4(dithio)uracil; 5-(methylaminomethyl)-4-(thio)uracil; 5-(propynyl)uracil; 5-(trifluoromethyl)uracil; 5-aminoallyl-uridine; 5-bromo-uridine; 5-

iodo-uridine; 5-uracil; 6 (azo)uracil; 6-(azo)uracil; 6-aza-uridine; allyamino-uracil; aza uracil; deaza uracil; N3 (methyl)uracil; P pseudo-UTP-1-2-ethanoic acid; Pseudouracil; 4-Thio-pseudo-UTP; 1-carboxymethyl-pseudouridine; 1-methyl-1-deaza-pseudouridine; 1-propynyl-uridine; 1-taurinomethyl-1-methyl-uridine; 1-taurinomethyl-4-thio-uridine; 1-taurinomethyl-pseudouridine; 2-methoxy-4-thio-pseudouridine; 2-thio-1-methyl-1-deaza-pseudouridine; 2-thio-1-methyl-pseudouridine; 2-thio-5-aza-uridine; 2-thio-dihydropseudouridine; 2-thio-dihydrouridine; 2-thio-pseudouridine; 4-methoxy-2-thio-pseudouridine; 4-methoxy-pseudouridine; 4-thio-1-methyl-pseudouridine; 4-thio-pseudouridine; 5-aza-uridine; Dihydropseudouridine; ($\pm$)1-(2-Hydroxypropyl)pseudouridine TP; (2R)-1-(2-Hydroxypropyl)pseudouridine TP; (2S)-1-(2-Hydroxypropyl)pseudouridine TP; (E)-5-(2-Bromo-vinyl)ara-uridine TP; (E)-5-(2-Bromo-vinyl)uridine TP; (Z)-5-(2-Bromo-vinyl)ara-uridine TP; (Z)-5-(2-Bromo-vinyl)uridine TP; 1-(2,2,2-Trifluoroethyl)-pseudo-UTP; 1-(2,2,3,3,3-Pentafluoropropyl)pseudouridine TP; 1-(2,2-Diethoxyethyl)pseudouridine TP; 1-(2,4,6-Trimethylbenzyl)pseudouridine TP; 1-(2,4,6-Trimethyl-benzyl)pseudo-UTP; 1-(2,4,6-Trimethyl-phenyl)pseudo-UTP; 1-(2-Amino-2-carboxyethyl)pseudo-UTP; 1-(2-Amino-ethyl)pseudo-UTP; 1-(2-Hydroxyethyl)pseudouridine TP; 1-(2-Methoxyethyl)pseudouridine TP; 1-(3,4-Bis-trifluoromethoxybenzyl)pseudouridine TP; 1-(3,4-Dimethoxybenzyl)pseudouridine TP; 1-(3-Amino-3-carboxypropyl)pseudo-UTP; 1-(3-Amino-propyl)pseudo-UTP; 1-(3-Cyclopropyl-prop-2-ynyl)pseudouridine TP; 1-(4-Amino-4-carboxybutyl)pseudo-UTP; 1-(4-Amino-benzyl)pseudo-UTP; 1-(4-Amino-butyl)pseudo-UTP; 1-(4-Amino-phenyl)pseudo-UTP; 1-(4-Azidobenzyl)pseudouridine TP; 1-(4-Bromobenzyl)pseudouridine TP; 1-(4-Chlorobenzyl)pseudouridine TP; 1-(4-Fluorobenzyl)pseudouridine TP; 1-(4-Iodobenzyl)pseudouridine TP; 1-(4-Methanesulfonylbenzyl)pseudouridine TP; 1-(4-Methoxybenzyl)pseudouridine TP; 1-(4-Methoxy-benzyl)pseudo-UTP; 1-(4-Methoxy-phenyl)pseudo-UTP; 1-(4-Methylbenzyl)pseudouridine TP; 1-(4-Methyl-benzyl)pseudo-UTP; 1-(4-Nitrobenzyl)pseudouridine TP; 1-(4-Nitro-benzyl)pseudo-UTP; 1-(4-Nitro-phenyl)pseudo-UTP; 1-(4-Thiomethoxybenzyl)pseudouridine TP; 1-(4-Trifluoromethoxybenzyl)pseudouridine TP; 1-(4-Trifluoromethylbenzyl)pseudouridine TP; 1-(5-Amino-pentyl)pseudo-UTP; 1-(6-Amino-hexyl)pseudo-UTP; 1,6-Dimethyl-pseudo-UTP; 1-[3-(2-{2-[2-(2-Aminoethoxy)-ethoxy]-ethoxy}-ethoxy)-propionyl]pseudouridine TP; 1-{3-[2-(2-Aminoethoxy)-ethoxy]-propionyl }

pseudouridine TP; 1-Acetylpsseudouridine TP; 1-Alkyl-6-(1-propynyl)-pseudo-UTP; 1-Alkyl-6-(2-propynyl)-pseudo-UTP; 1-Alkyl-6-allyl-pseudo-UTP; 1-Alkyl-6-ethynyl-pseudo-UTP; 1-Alkyl-6-homoallyl-pseudo-UTP; 1-Alkyl-6-vinyl-pseudo-UTP; 1-Allylpsseudouridine TP; 1-Aminomethyl-pseudo-UTP; 1-Benzoylpsseudouridine TP; 1-Benzyloxymethylpsseudouridine TP; 1-Benzyl-pseudo-UTP; 1-Biotinyl-PEG2-pseudouridine TP; 1-Biotinylpsseudouridine TP; 1-Butyl-pseudo-UTP; 1-Cyanomethylpsseudouridine TP; 1-Cyclobutylmethyl-pseudo-UTP; 1-Cyclobutyl-pseudo-UTP; 1-Cycloheptylmethyl-pseudo-UTP; 1-Cycloheptyl-pseudo-UTP; 1-Cyclohexylmethyl-pseudo-UTP; 1-Cyclohexyl-pseudo-UTP; 1-Cyclooctylmethyl-pseudo-UTP; 1-Cyclooctyl-pseudo-UTP; 1-Cyclopentylmethyl-pseudo-UTP; 1-Cyclopentyl-pseudo-UTP; 1-Cyclopropylmethyl-pseudo-UTP; 1-Cyclopropyl-pseudo-UTP; 1-Ethyl-pseudo-UTP; 1-Hexyl-pseudo-UTP; 1-Homoallylpsseudouridine TP; 1-Hydroxymethylpsseudouridine TP; 1-iso-propyl-pseudo-UTP; 1-Me-2-thio-pseudo-UTP; 1-Me-4-thio-pseudo-UTP; 1-Me-alpha-thio-pseudo-UTP; 1-Methanesulfonylmethylpsseudouridine TP; 1-Methoxymethylpsseudouridine TP; 1-Methyl-6-(2,2,2-Trifluoroethyl)pseudo-UTP; 1-Methyl-6-(4-morpholino)-pseudo-UTP; 1-Methyl-6-(4-thiomorpholino)-pseudo-UTP; 1-Methyl-6-(substituted phenyl)pseudo-UTP; 1-Methyl-6-amino-pseudo-UTP; 1-Methyl-6-azido-pseudo-UTP; 1-Methyl-6-bromopseudo-UTP; 1-Methyl-6-butyl-pseudo-UTP; 1-Methyl-6-chloro-pseudo-UTP; 1-Methyl-6-cyano-pseudo-UTP; 1-Methyl-6-dimethylamino-pseudo-UTP; 1-Methyl-6-ethoxy-pseudo-UTP; 1-Methyl-6-ethylcarboxylate-pseudo-UTP; 1-Methyl-6-ethyl-pseudo-UTP; 1-Methyl-6-fluoro-pseudo-UTP; 1-Methyl-6-formyl-pseudo-UTP; 1-Methyl-6-hydroxyamino-pseudo-UTP; 1-Methyl-6-hydroxy-pseudo-UTP; 1-Methyl-6-iodo-pseudo-UTP; 1-Methyl-6-iso-propyl-pseudo-UTP; 1-Methyl-6-methoxy-pseudo-UTP; 1-Methyl-6-methylamino-pseudo-UTP; 1-Methyl-6-phenyl-pseudo-UTP; 1-Methyl-6-propyl-pseudo-UTP; 1-Methyl-6-tert-butyl-pseudo-UTP; 1-Methyl-6-trifluoromethoxy-pseudo-UTP; 1-Methyl-6-trifluoromethyl-pseudo-UTP; 1-Morpholinomethylpsseudouridine TP; 1-Pentyl-pseudo-UTP; 1-Phenyl-pseudo-UTP; 1-Pivaloylpsseudouridine TP; 1-Propargylpsseudouridine TP; 1-Propyl-pseudo-UTP; 1-propynyl-pseudouridine; 1-p-tolyl-pseudo-UTP; 1-tert-Butyl-pseudo-UTP; 1-Thiomethoxymethylpsseudouridine TP; 1-Thiomorpholinomethylpsseudouridine TP; 1-Trifluoroacetylpsseudouridine TP; 1-Trifluoromethyl-pseudo-UTP; 1-Vinylpsseudouridine TP; 2,2'-anhydro-uridine TP; 2'-bromo-deoxyuridine TP; 2'-F-5-Methyl-2'-deoxy-UTP; 2'-OMe-5-Me-UTP; 2'-OMe-

pseudo-UTP; 2'-a-Ethynyluridine TP; 2'-a-Trifluoromethyluridine TP; 2'-b-Ethynyluridine TP; 2'-b-Trifluoromethyluridine TP; 2'-Deoxy-2',2'-difluorouridine TP; 2'-Deoxy-2'-a-mercaptopuridine TP; 2'-Deoxy-2'-a-thiomethoxyuridine TP; 2'-Deoxy-2'-b-aminouridine TP; 2'-Deoxy-2'-b-azidouridine TP; 2'-Deoxy-2'-b-bromouridine TP; 2'-Deoxy-2'-b-chlorouridine TP; 2'-Deoxy-2'-b-fluorouridine TP; 2'-Deoxy-2'-b-iodouridine TP; 2'-Deoxy-2'-b-mercaptopuridine TP; 2'-Deoxy-2'-b-thiomethoxyuridine TP; 2-methoxy-4-thio-uridine; 2-methoxyuridine; 2'-O-Methyl-5-(1-propynyl)uridine TP; 3-Alkyl-pseudo-UTP; 4'-Azidouridine TP; 4'-Carbocyclic uridine TP; 4'-Ethynyluridine TP; 5-(1-Propynyl)ara-uridine TP; 5-(2-Furanyl)uridine TP; 5-Cyanouridine TP; 5-Dimethylaminouridine TP; 5'-Homo-uridine TP; 5-iodo-2'-fluoro-deoxyuridine TP; 5-Phenylethynyluridine TP; 5-Trideuteromethyl-6-deuterouridine TP; 5-Trifluoromethyl-Uridine TP; 5-Vinylarauridine TP; 6-(2,2,2-Trifluoroethyl)-pseudo-UTP; 6-(4-Morpholino)-pseudo-UTP; 6-(4-Thiomorpholino)-pseudo-UTP; 6-(Substituted-Phenyl)-pseudo-UTP; 6-Amino-pseudo-UTP; 6-Azido-pseudo-UTP; 6-Bromo-pseudo-UTP; 6-Butyl-pseudo-UTP; 6-Chloro-pseudo-UTP; 6-Cyano-pseudo-UTP; 6-Dimethylamino-pseudo-UTP; 6-Ethoxy-pseudo-UTP; 6-Ethylcarboxylate-pseudo-UTP; 6-Ethyl-pseudo-UTP; 6-Fluoro-pseudo-UTP; 6-Formyl-pseudo-UTP; 6-Hydroxyamino-pseudo-UTP; 6-Hydroxy-pseudo-UTP; 6-Iodo-pseudo-UTP; 6-iso-Propyl-pseudo-UTP; 6-Methoxy-pseudo-UTP; 6-Methylamino-pseudo-UTP; 6-Methyl-pseudo-UTP; 6-Phenyl-pseudo-UTP; 6-Phenyl-pseudo-UTP; 6-Propyl-pseudo-UTP; 6-tert-Butyl-pseudo-UTP; 6-Trifluoromethoxy-pseudo-UTP; 6-Trifluoromethyl-pseudo-UTP; Alpha-thio-pseudo-UTP; Pseudouridine 1-(4-methylbenzenesulfonic acid) TP; Pseudouridine 1-(4-methylbenzoic acid) TP; Pseudouridine TP 1-[3-(2-ethoxy)]propionic acid; Pseudouridine TP 1-[3-{2-(2-[2-(2-ethoxy)-ethoxy]-ethoxy)-ethoxy}]propionic acid; Pseudouridine TP 1-[3-{2-(2-[2-{2(2-ethoxy)-ethoxy}-ethoxy]-ethoxy)-ethoxy}]propionic acid; Pseudouridine TP 1-[3-{2-(2-[2-ethoxy]-ethoxy)-ethoxy}]propionic acid; Pseudouridine TP 1-[3-{2-(2-ethoxy)-ethoxy}]propionic acid; Pseudouridine TP 1-methylphosphonic acid; Pseudouridine TP 1-methylphosphonic acid diethyl ester; Pseudo-UTP-N1-3-propionic acid; Pseudo-UTP-N1-4-butanoic acid; Pseudo-UTP-N1-5-pentanoic acid; Pseudo-UTP-N1-6-hexanoic acid; Pseudo-UTP-N1-7-heptanoic acid; Pseudo-UTP-N1-methyl-p-benzoic acid; Pseudo-UTP-N1-p-benzoic acid; Wybutosine; Hydroxywybutosine; Isowyosine; Peroxywybutosine; undermodified hydroxywybutosine; 4-demethylwyosine; 2,6-(diamino)purine; 1-(aza)-2-(thio)-3-(aza)-phenoxazin-1-yl; 1,3-(

diaza)-2-(oxo)-phenthiazin-1-yl; 1,3-(diaza)-2-(oxo)-phenoxazin-1-yl; 1,3,5-(triaz)-2,6-(diox)-naphthalene; 2 (amino)purine; 2,4,5-(trimethyl)phenyl; 2' methyl, 2'amino, 2'azido, 2'fluro-cytidine; 2' methyl, 2'amino, 2'azido, 2'fluro-adenine; 2'methyl, 2'amino, 2'azido, 2'fluro-uridine; 2'-amino-2'-deoxyribose; 2-amino-6-Chloro-purine; 2-aza-inosinyl; 2'-azido-2'-deoxyribose; 2'fluoro-2'-deoxyribose; 2'-fluoro-modified bases; 2'-O-methyl-ribose; 2-oxo-7-aminopyridopyrimidin-3-yl; 2-oxo-pyridopyrimidine-3-yl; 2-pyridinone; 3 nitropyrrole; 3-(methyl)-7-(propynyl)isocarbostyryl; 3-(methyl)isocarbostyryl; 4-(fluoro)-6-(methyl)benzimidazole; 4-(methyl)benzimidazole; 4-(methyl)indolyl; 4,6-(dimethyl)indolyl; 5 nitroindole; 5 substituted pyrimidines; 5-(methyl)isocarbostyryl; 5-nitroindole; 6-(aza)pyrimidine; 6-(azo)thymine; 6-(methyl)-7-(aza)indolyl; 6-chloro-purine; 6-phenyl-pyrrolo-pyrimidin-2-on-3-yl; 7-(aminoalkylhydroxy)-1-(aza)-2-(thio)-3-(aza)-phenthiazin-1-yl; 7-(aminoalkylhydroxy)-1-(aza)-2-(thio)-3-(aza)-phenoxazin-1-yl; 7-(aminoalkylhydroxy)-1,3-(diaza)-2-(oxo)-phenoxazin-1-yl; 7-(aminoalkylhydroxy)-1,3-(diaza)-2-(oxo)-phenthiazin-1-yl; 7-(aminoalkylhydroxy)-1,3-(diaza)-2-(oxo)-phenoxazin-1-yl; 7-(aza)indolyl; 7-(guanidiniumalkylhydroxy)-1-(aza)-2-(thio)-3-(aza)-phenoxazin-1-yl; 7-(guanidiniumalkylhydroxy)-1-(aza)-2-(thio)-3-(aza)-phenthiazin-1-yl; 7-(guanidiniumalkylhydroxy)-1-(aza)-2-(thio)-3-(aza)-phenoxazin-1-yl; 7-(guanidiniumalkylhydroxy)-1,3-(diaza)-2-(oxo)-phenoxazin-1-yl; 7-(guanidiniumalkylhydroxy)-1,3-(diaza)-2-(oxo)-phenthiazin-1-yl; 7-(guanidiniumalkylhydroxy)-1,3-(diaza)-2-(oxo)-phenoxazin-1-yl; 7-(propynyl)isocarbostyryl; 7-(propynyl)isocarbostyryl, propynyl-7-(aza)indolyl; 7-deaza-inosinyl; 7-substituted 1-(aza)-2-(thio)-3-(aza)-phenoxazin-1-yl; 7-substituted 1,3-(diaza)-2-(oxo)-phenoxazin-1-yl; 9-(methyl)-imidizopyridinyl; Aminoindolyl; Anthracenyl; bis-ortho-(aminoalkylhydroxy)-6-phenyl-pyrrolo-pyrimidin-2-on-3-yl; bis-ortho-substituted-6-phenyl-pyrrolo-pyrimidin-2-on-3-yl; Difluorotolyl; Hypoxanthine; Imidizopyridinyl; Inosinyl; Isocarbostyryl; Isoguanisine; N2-substituted purines; N6-methyl-2-amino-purine; N6-substituted purines; N-alkylated derivative; Napthalenyl; Nitrobenzimidazolyl; Nitroimidazolyl; Nitroindazolyl; Nitropyrazolyl; Nubularine; O6-substituted purines; O-alkylated derivative; ortho-(aminoalkylhydroxy)-6-phenyl-pyrrolo-pyrimidin-2-on-3-yl; ortho-substituted-6-phenyl-pyrrolo-pyrimidin-2-on-3-yl; Oxoformycin TP; para-(aminoalkylhydroxy)-6-phenyl-pyrrolo-pyrimidin-2-on-3-yl; para-substituted-6-phenyl-pyrrolo-pyrimidin-2-on-3-yl; Pentacenyl; Phenanthracenyl; Phenyl; propynyl-7-(aza)indolyl; Pyrenyl; pyridopyrimidin-3-yl; pyridopyrimidin-3-yl, 2-oxo-7-

amino-pyridopyrimidin-3-yl; pyrrolo-pyrimidin-2-on-3-yl; Pyrrolopyrimidinyl; Pyrrolopyrizinyl; Stilbenzyl; substituted 1,2,4-triazoles; Tetracenyl; Tubercidine; Xanthine; Xanthosine-5'-TP; 2-thio-zebularine; 5-aza-2-thio-zebularine; 7-deaza-2-amino-purine; pyridin-4-one ribonucleoside; 2-Amino-riboside-TP; Formycin A TP; Formycin B TP; Pyrrolosine TP; 2'-OH-ara-adenosine TP; 2'-OH-ara-cytidine TP; 2'-OH-ara-uridine TP; 2'-OH-ara-guanosine TP; 5-(2-carbomethoxyvinyl)uridine TP; and N6-(19-Amino-pentaoxanonadecyl)adenosine TP.

[0398] In some embodiments, the polynucleotide (*e.g.*, RNA polynucleotide, such as mRNA polynucleotide) includes a combination of at least two (*e.g.*, 2, 3, 4 or more) of the aforementioned modified nucleobases.

[0399] In some embodiments, the mRNA comprises at least one chemically modified nucleoside. In some embodiments, the at least one chemically modified nucleoside is selected from the group consisting of pseudouridine (ψ), 2-thiouridine (s2U), 4'-thiouridine, 5-methylcytosine, 2-thio-1-methyl-1-deaza-pseudouridine, 2-thio-1-methyl-pseudouridine, 2-thio-5-aza-uridine, 2-thio-dihydropseudouridine, 2-thio-dihydrouridine, 2-thio-pseudouridine, 4-methoxy-2-thio-pseudouridine, 4-methoxy-pseudouridine, 4-thio-1-methyl-pseudouridine, 4-thio-pseudouridine, 5-aza-uridine, dihydropseudouridine, 5-methyluridine, 5-methoxyuridine, 2'-O-methyl uridine, 1-methyl-pseudouridine (m1 ψ), 1-ethyl-pseudouridine (e1 ψ), 5-methoxy-uridine (mo5U), 5-methyl-cytidine (m5C), α -thio-guanosine, α -thio-adenosine, 5-cyano uridine, 4'-thio uridine 7-deaza-adenine, 1-methyl-adenosine (m1A), 2-methyl-adenine (m2A), N6-methyl-adenosine (m6A), and 2,6-Diaminopurine, (I), 1-methyl-inosine (m1I), wyosine (imG), methylwyosine (mimG), 7-deaza-guanosine, 7-cyano-7-deaza-guanosine (preQ0), 7-aminomethyl-7-deaza-guanosine (preQ1), 7-methyl-guanosine (m7G), 1-methyl-guanosine (m1G), 8-oxo-guanosine, 7-methyl-8-oxo-guanosine, 2,8-dimethyladenosine, 2-geranylthiouridine, 2-lysidine, 2-selenouridine, 3-(3-amino-3-carboxypropyl)-5,6-dihydrouridine, 3-(3-amino-3-carboxypropyl)pseudouridine, 3-methylpseudouridine, 5-(carboxyhydroxymethyl)-2'-O-methyluridine methyl ester, 5-aminomethyl-2-geranylthiouridine, 5-aminomethyl-2-selenouridine, 5-aminomethyluridine, 5-carbamoylhydroxymethyluridine, 5-carbamoylmethyl-2-thiouridine, 5-carboxymethyl-2-thiouridine, 5-carboxymethylaminomethyl-2-geranylthiouridine, 5-carboxymethylaminomethyl-2-selenouridine, 5-cyanomethyluridine, 5-hydroxycytidine, 5-methylaminomethyl-2-geranylthiouridine, 7-aminocarboxypropyl-demethylwyosine, 7-

aminocarboxypropylwyosine, 7-aminocarboxypropylwyosine methyl ester, 8-methyladenosine, N⁴,N⁴-dimethylcytidine, N⁶-formyladenosine, N⁶-hydroxymethyladenosine, agmatidine, cyclic N⁶-threonylcarbamoyladenosine, glutamyl-queuosine, methylated undermodified hydroxywybutosine, N⁴,N⁴,2'-O-trimethylcytidine, geranylated 5-methylaminomethyl-2-thiouridine, geranylated 5-carboxymethylaminomethyl-2-thiouridine, Qbase, preQ⁰base, preQ¹base, and two or more combinations thereof. In some embodiments, the at least one chemically modified nucleoside is selected from the group consisting of pseudouridine, 1-methyl-pseudouridine, 1-ethyl-pseudouridine, 5-methylcytosine, 5-methoxyuridine, and a combination thereof. In some embodiments, the polynucleotide (*e.g.*, RNA polynucleotide, such as mRNA polynucleotide) includes a combination of at least two (*e.g.*, 2, 3, 4 or more) of the aforementioned modified nucleobases.

(i) Base Modifications

[0400] In certain embodiments, the chemical modification is at nucleobases in the polynucleotides (*e.g.*, RNA polynucleotide, such as mRNA polynucleotide). In some embodiments, modified nucleobases in the polynucleotide (*e.g.*, RNA polynucleotide, such as mRNA polynucleotide) are selected from the group consisting of 1-methyl-pseudouridine (m1ψ), 1-ethyl-pseudouridine (e1ψ), 5-methoxy-uridine (mo5U), 5-methyl-cytidine (m5C), pseudouridine (ψ), α-thio-guanosine and α-thio-adenosine. In some embodiments, the polynucleotide includes a combination of at least two (*e.g.*, 2, 3, 4 or more) of the aforementioned modified nucleobases.

[0401] In some embodiments, the polynucleotide (*e.g.*, RNA polynucleotide, such as mRNA polynucleotide) comprises pseudouridine (ψ) and 5-methyl-cytidine (m5C). In some embodiments, the polynucleotide (*e.g.*, RNA polynucleotide, such as mRNA polynucleotide) comprises 1-methyl-pseudouridine (m1ψ). In some embodiments, the polynucleotide (*e.g.*, RNA polynucleotide, such as mRNA polynucleotide) comprises 1-ethyl-pseudouridine (e1ψ). In some embodiments, the polynucleotide (*e.g.*, RNA polynucleotide, such as mRNA polynucleotide) comprises 1-methyl-pseudouridine (m1ψ) and 5-methyl-cytidine (m5C). In some embodiments, the polynucleotide (*e.g.*, RNA polynucleotide, such as mRNA polynucleotide) comprises 1-ethyl-pseudouridine (e1ψ) and 5-methyl-cytidine (m5C). In some embodiments, the polynucleotide (*e.g.*, RNA polynucleotide, such as mRNA polynucleotide) comprises 2-thiouridine (s2U). In some

embodiments, the polynucleotide (*e.g.*, RNA polynucleotide, such as mRNA polynucleotide) comprises 2-thiouridine and 5-methyl-cytidine (m5C). In some embodiments, the polynucleotide (*e.g.*, RNA polynucleotide, such as mRNA polynucleotide) comprises methoxy-uridine (mo5U). In some embodiments, the polynucleotide (*e.g.*, RNA polynucleotide, such as mRNA polynucleotide) comprises 5-methoxy-uridine (mo5U) and 5-methyl-cytidine (m5C). In some embodiments, the polynucleotide (*e.g.*, RNA polynucleotide, such as mRNA polynucleotide) comprises 2'-O-methyl uridine. In some embodiments, the polynucleotide (*e.g.*, RNA polynucleotide, such as mRNA polynucleotide) comprises 2'-O-methyl uridine and 5-methyl-cytidine (m5C). In some embodiments, the polynucleotide (*e.g.*, RNA polynucleotide, such as mRNA polynucleotide) comprises N6-methyl-adenosine (m6A). In some embodiments, the polynucleotide (*e.g.*, RNA polynucleotide, such as mRNA polynucleotide) comprises N6-methyl-adenosine (m6A) and 5-methyl-cytidine (m5C).

- [0402] In some embodiments, the polynucleotide (*e.g.*, RNA polynucleotide, such as mRNA polynucleotide) is uniformly modified (*e.g.*, fully modified, modified throughout the entire sequence) for a particular modification. For example, a polynucleotide can be uniformly modified with 5-methyl-cytidine (m5C), meaning that all cytosine residues in the mRNA sequence are replaced with 5-methyl-cytidine (m5C). Similarly, a polynucleotide can be uniformly modified for any type of nucleoside residue present in the sequence by replacement with a modified residue such as any of those set forth above.
- [0403] In some embodiments, the chemically modified nucleosides in the open reading frame are selected from the group consisting of uridine, adenine, cytosine, guanine, and any combination thereof.
- [0404] In some embodiments, the modified nucleobase is a modified cytosine. Examples of nucleobases and nucleosides having a modified cytosine include N4-acetyl-cytidine (ac4C), 5-methyl-cytidine (m5C), 5-halo-cytidine (*e.g.*, 5-iodo-cytidine), 5-hydroxymethyl-cytidine (hm5C), 1-methyl-pseudoisocytidine, 2-thio-cytidine (s2C), 2-thio-5-methyl-cytidine.
- [0405] In some embodiments, a modified nucleobase is a modified uridine. Example nucleobases and nucleosides having a modified uridine include 5-cyano uridine or 4'-thio uridine.
- [0406] In some embodiments, a modified nucleobase is a modified adenine. Example nucleobases and nucleosides having a modified adenine include 7-deaza-adenine, 1-

methyl-adenosine (m1A), 2-methyl-adenine (m2A), N6-methyl-adenine (m6A), and 2,6-Diaminopurine.

- [0407] In some embodiments, a modified nucleobase is a modified guanine. Example nucleobases and nucleosides having a modified guanine include inosine (I), 1-methyl-inosine (m1I), wyosine (imG), methylwyosine (mimG), 7-deaza-guanosine, 7-cyano-7-deaza-guanosine (preQ0), 7-aminomethyl-7-deaza-guanosine (preQ1), 7-methyl-guanosine (m7G), 1-methyl-guanosine (m1G), 8-oxo-guanosine, 7-methyl-8-oxo-guanosine.
- [0408] In some embodiments, the nucleobase modified nucleotides in the polynucleotide (*e.g.*, RNA polynucleotide, such as mRNA polynucleotide) are 5-methoxyuridine.
- [0409] In some embodiments, the polynucleotide (*e.g.*, RNA polynucleotide, such as mRNA polynucleotide) includes a combination of at least two (*e.g.*, 2, 3, 4 or more) of modified nucleobases.
- [0410] In some embodiments, at least 95% of a type of nucleobases (*e.g.*, uracil) in a polynucleotide of the invention (*e.g.*, an mRNA polynucleotide encoding ACADVL) are modified nucleobases. In some embodiments, at least 95% of uracil in a polynucleotide of the present invention (*e.g.*, an mRNA polynucleotide encoding ACADVL) is 5-methoxyuracil.
- [0411] In some embodiments, the polynucleotide (*e.g.*, RNA polynucleotide, such as mRNA polynucleotide) comprises 5-methoxyuridine (5mo5U) and 5-methyl-cytidine (m5C).
- [0412] In some embodiments, the polynucleotide (*e.g.*, RNA polynucleotide, such as mRNA polynucleotide) is uniformly modified (*e.g.*, fully modified, modified throughout the entire sequence) for a particular modification. For example, a polynucleotide can be uniformly modified with 5-methoxyuridine, meaning that substantially all uridine residues in the mRNA sequence are replaced with 5-methoxyuridine. Similarly, a polynucleotide can be uniformly modified for any type of nucleoside residue present in the sequence by replacement with a modified residue such as any of those set forth above.
- [0413] In some embodiments, the modified nucleobase is a modified cytosine.
- [0414] In some embodiments, a modified nucleobase is a modified uracil. Example nucleobases and nucleosides having a modified uracil include 5-methoxyuracil.
- [0415] In some embodiments, a modified nucleobase is a modified adenine.
- [0416] In some embodiments, a modified nucleobase is a modified guanine.

- [0417] In some embodiments, the nucleobases, sugar, backbone, or any combination thereof in the open reading frame encoding an ACADVL polypeptide are chemically modified by at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, at least 99%, or 100%.
- [0418] In some embodiments, the uridine nucleosides in the open reading frame encoding an ACADVL polypeptide are chemically modified by at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, at least 99%, or 100%.
- [0419] In some embodiments, the adenosine nucleosides in the open reading frame encoding an ACADVL polypeptide are chemically modified by at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, at least 99%, or 100%.
- [0420] In some embodiments, the cytidine nucleosides in the open reading frame encoding an ACADVL polypeptide are chemically modified by at least at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, at least 99%, or 100%.
- [0421] In some embodiments, the guanosine nucleosides in the open reading frame encoding a polypeptide are chemically modified by at least at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, at least 99%, or 100%.
- [0422] In some embodiments, the polynucleotides can include any useful linker between the nucleosides. Such linkers, including backbone modifications, that are useful in the composition of the present disclosure include, but are not limited to the following: 3'-alkylene phosphonates, 3'-amino phosphoramidate, alkene containing backbones, aminoalkylphosphoramidates, aminoalkylphosphotriesters, boranophosphates, $-\text{CH}_2\text{-O-N}(\text{CH}_3)\text{-CH}_2\text{-}$, $-\text{CH}_2\text{-N}(\text{CH}_3)\text{-N}(\text{CH}_3)\text{-CH}_2\text{-}$, $-\text{CH}_2\text{-NH-CH}_2\text{-}$, chiral phosphonates, chiral phosphorothioates, formacetyl and thioformacetyl backbones, methylene (methylimino), methylene formacetyl and thioformacetyl backbones, methyleneimino and methylenehydrazino backbones, morpholino linkages, $-\text{N}(\text{CH}_3)\text{-CH}_2\text{-CH}_2\text{-}$, oligonucleosides with heteroatom internucleoside linkage, phosphinates, phosphoramidates, phosphorodithioates, phosphorothioate internucleoside linkages, phosphorothioates, phosphotriesters, PNA, siloxane backbones, sulfamate backbones,

sulfide sulfoxide and sulfone backbones, sulfonate and sulfonamide backbones, thionoalkylphosphonates, thionoalkylphosphotriesters, and thionophosphoramidates.

(ii) Sugar Modifications

[0423] The modified nucleosides and nucleotides (*e.g.*, building block molecules), which can be incorporated into a polynucleotide (*e.g.*, RNA or mRNA, as described herein), can be modified on the sugar of the ribonucleic acid. For example, the 2' hydroxyl group (OH) can be modified or replaced with a number of different substituents. Exemplary substitutions at the 2'-position include, but are not limited to, H, halo, optionally substituted C₁₋₆ alkyl; optionally substituted C₁₋₆ alkoxy; optionally substituted C₆₋₁₀ aryloxy; optionally substituted C₃₋₈ cycloalkyl; optionally substituted C₃₋₈ cycloalkoxy; optionally substituted C₆₋₁₀ aryloxy; optionally substituted C₆₋₁₀ aryl-C₁₋₆ alkoxy, optionally substituted C₁₋₁₂ (heterocycl)oxy; a sugar (*e.g.*, ribose, pentose, or any described herein); a polyethyleneglycol (PEG), -O(CH₂CH₂O)_nCH₂CH₂OR, where R is H or optionally substituted alkyl, and n is an integer from 0 to 20 (*e.g.*, from 0 to 4, from 0 to 8, from 0 to 10, from 0 to 16, from 1 to 4, from 1 to 8, from 1 to 10, from 1 to 16, from 1 to 20, from 2 to 4, from 2 to 8, from 2 to 10, from 2 to 16, from 2 to 20, from 4 to 8, from 4 to 10, from 4 to 16, and from 4 to 20); "locked" nucleic acids (LNA) in which the 2'-hydroxyl is connected by a C₁₋₆ alkylene or C₁₋₆ heteroalkylene bridge to the 4'-carbon of the same ribose sugar, where exemplary bridges included methylene, propylene, ether, or amino bridges; aminoalkyl, as defined herein; aminoalkoxy, as defined herein; amino as defined herein; and amino acid, as defined herein

[0424] Generally, RNA includes the sugar group ribose, which is a 5-membered ring having an oxygen. Exemplary, non-limiting modified nucleotides include replacement of the oxygen in ribose (*e.g.*, with S, Se, or alkylene, such as methylene or ethylene); addition of a double bond (*e.g.*, to replace ribose with cyclopentenyl or cyclohexenyl); ring contraction of ribose (*e.g.*, to form a 4-membered ring of cyclobutane or oxetane); ring expansion of ribose (*e.g.*, to form a 6- or 7-membered ring having an additional carbon or heteroatom, such as for anhydrohexitol, altritol, mannitol, cyclohexanyl, cyclohexenyl, and morpholino that also has a phosphoramidate backbone); multicyclic forms (*e.g.*, tricyclo; and "unlocked" forms, such as glycol nucleic acid (GNA) (*e.g.*, R-GNA or S-GNA, where ribose is replaced by glycol units attached to phosphodiester bonds), threose nucleic acid (TNA, where ribose is replaced with α -L-threofuranosyl-

(3'→2')), and peptide nucleic acid (PNA, where 2-amino-ethyl-glycine linkages replace the ribose and phosphodiester backbone). The sugar group can also contain one or more carbons that possess the opposite stereochemical configuration than that of the corresponding carbon in ribose. Thus, a polynucleotide molecule can include nucleotides containing, *e.g.*, arabinose, as the sugar. Such sugar modifications are taught International Patent Publication Nos. WO2013052523 and WO2014093924, the contents of each of which are incorporated herein by reference in their entireties.

(iii) Combinations of Modifications

[0425] The polynucleotides of the invention (*e.g.*, a polynucleotide comprising a nucleotide sequence encoding an ACADVL polypeptide or a functional fragment or variant thereof) can include a combination of modifications to the sugar, the nucleobase, and/or the internucleoside linkage. These combinations can include any one or more modifications described herein.

[0426] Combinations of modified nucleotides can be used to form the polynucleotides of the invention. Unless otherwise noted, the modified nucleotides can be completely substituted for the natural nucleotides of the polynucleotides of the invention. As a non-limiting example, the natural nucleotide uridine can be substituted with a modified nucleoside described herein. In another non-limiting example, the natural nucleotide uridine can be partially substituted or replaced (*e.g.*, about 0.1%, 1%, 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95% or 99.9%) with at least one of the modified nucleoside disclosed herein. Any combination of base/sugar or linker can be incorporated into the polynucleotides of the invention and such modifications are taught in International Patent Publications WO2013052523 and WO2014093924, and U.S. Publ. Nos. US 20130115272 and US20150307542, the contents of each of which are incorporated herein by reference in its entirety.

11. Untranslated Regions (UTRs)

[0427] Untranslated regions (UTRs) are nucleic acid sections of a polynucleotide before a start codon (5'UTR) and after a stop codon (3'UTR) that are not translated. In some embodiments, a polynucleotide (*e.g.*, a ribonucleic acid (RNA), *e.g.*, a messenger RNA (mRNA)) of the invention comprising an open reading frame (ORF) encoding an

ACADVL polypeptide further comprises UTR (e.g., a 5'UTR or functional fragment thereof, a 3'UTR or functional fragment thereof, or a combination thereof).

[0428] A UTR can be homologous or heterologous to the coding region in a polynucleotide. In some embodiments, the UTR is homologous to the ORF encoding the ACADVL polypeptide. In some embodiments, the UTR is heterologous to the ORF encoding the ACADVL polypeptide. In some embodiments, the polynucleotide comprises two or more 5'UTRs or functional fragments thereof, each of which has the same or different nucleotide sequences. In some embodiments, the polynucleotide comprises two or more 3'UTRs or functional fragments thereof, each of which has the same or different nucleotide sequences.

[0429] In some embodiments, the 5'UTR or functional fragment thereof, 3' UTR or functional fragment thereof, or any combination thereof is sequence optimized.

[0430] In some embodiments, the 5'UTR or functional fragment thereof, 3' UTR or functional fragment thereof, or any combination thereof comprises at least one chemically modified nucleobase, e.g., 1-methylpseudouridine or 5-methoxyuracil.

[0431] UTRs can have features that provide a regulatory role, e.g., increased or decreased stability, localization and/or translation efficiency. A polynucleotide comprising a UTR can be administered to a cell, tissue, or organism, and one or more regulatory features can be measured using routine methods. In some embodiments, a functional fragment of a 5'UTR or 3'UTR comprises one or more regulatory features of a full length 5' or 3' UTR, respectively.

[0432] Natural 5'UTRs bear features that play roles in translation initiation. They harbor signatures like Kozak sequences that are commonly known to be involved in the process by which the ribosome initiates translation of many genes. Kozak sequences have the consensus CCR(A/G)CCAUGG, where R is a purine (adenine or guanine) three bases upstream of the start codon (AUG), which is followed by another 'G'. 5'UTRs also have been known to form secondary structures that are involved in elongation factor binding.

[0433] By engineering the features typically found in abundantly expressed genes of specific target organs, one can enhance the stability and protein production of a polynucleotide. For example, introduction of 5'UTR of liver-expressed mRNA, such as albumin, serum amyloid A, Apolipoprotein A/B/E, transferrin, alpha fetoprotein, erythropoietin, or Factor VIII, can enhance expression of polynucleotides in hepatic cell lines or liver. Likewise, use of 5'UTR from other tissue-specific mRNA to improve

expression in that tissue is possible for muscle (e.g., MyoD, Myosin, Myoglobin, Myogenin, Herculin), for endothelial cells (e.g., Tie-1, CD36), for myeloid cells (e.g., C/EBP, AML1, G-CSF, GM-CSF, CD11b, MSR, Fr-1, i-NOS), for leukocytes (e.g., CD45, CD18), for adipose tissue (e.g., CD36, GLUT4, ACRP30, adiponectin) and for lung epithelial cells (e.g., SP-A/B/C/D).

- [0434] In some embodiments, UTRs are selected from a family of transcripts whose proteins share a common function, structure, feature or property. For example, an encoded polypeptide can belong to a family of proteins (i.e., that share at least one function, structure, feature, localization, origin, or expression pattern), which are expressed in a particular cell, tissue or at some time during development. The UTRs from any of the genes or mRNA can be swapped for any other UTR of the same or different family of proteins to create a new polynucleotide.
- [0435] In some embodiments, the 5'UTR and the 3'UTR can be heterologous. In some embodiments, the 5'UTR can be derived from a different species than the 3'UTR. In some embodiments, the 3'UTR can be derived from a different species than the 5'UTR.
- [0436] Co-owned International Patent Application No. PCT/US2014/021522 (Publ. No. WO/2014/164253, incorporated herein by reference in its entirety) provides a listing of exemplary UTRs that can be utilized in the polynucleotide of the present invention as flanking regions to an ORF.
- [0437] Exemplary UTRs of the application include, but are not limited to, one or more 5'UTR and/or 3'UTR derived from the nucleic acid sequence of: a globin, such as an α - or β -globin (e.g., a *Xenopus*, mouse, rabbit, or human globin); a strong Kozak translational initiation signal; a CYBA (e.g., human cytochrome b-245 α polypeptide); an albumin (e.g., human albumin7); a HSD17B4 (hydroxysteroid (17- β) dehydrogenase); a virus (e.g., a tobacco etch virus (TEV), a Venezuelan equine encephalitis virus (VEEV), a Dengue virus, a cytomegalovirus (CMV) (e.g., CMV immediate early 1 (IE1)), a hepatitis virus (e.g., hepatitis B virus), a sindbis virus, or a PAV barley yellow dwarf virus); a heat shock protein (e.g., hsp70); a translation initiation factor (e.g., eIF4G); a glucose transporter (e.g., hGLUT1 (human glucose transporter 1)); an actin (e.g., human α or β actin); a GAPDH; a tubulin; a histone; a citric acid cycle enzyme; a topoisomerase (e.g., a 5'UTR of a TOP gene lacking the 5' TOP motif (the oligopyrimidine tract)); a ribosomal protein Large 32 (L32); a ribosomal protein (e.g., human or mouse ribosomal protein, such as, for example, rps9); an ATP synthase (e.g., ATP5A1 or the β subunit of

mitochondrial H⁺-ATP synthase); a growth hormone e (e.g., bovine (bGH) or human (hGH)); an elongation factor (e.g., elongation factor 1 α 1 (EEF1A1)); a manganese superoxide dismutase (MnSOD); a myocyte enhancer factor 2A (MEF2A); a β -F1-ATPase, a creatine kinase, a myoglobin, a granulocyte-colony stimulating factor (G-CSF); a collagen (e.g., collagen type I, alpha 2 (Col1A2), collagen type I, alpha 1 (Col1A1), collagen type VI, alpha 2 (Col6A2), collagen type VI, alpha 1 (Col6A1)); a ribophorin (e.g., ribophorin I (RPNI)); a low density lipoprotein receptor-related protein (e.g., LRP1); a cardiotrophin-like cytokine factor (e.g., Nnt1); calreticulin (Calr); a procollagen-lysine, 2-oxoglutarate 5-dioxygenase 1 (Plod1); and a nucleobindin (e.g., Nucb1).

[0438] In some embodiments, the 5'UTR is selected from the group consisting of a β -globin 5'UTR; a 5'UTR containing a strong Kozak translational initiation signal; a cytochrome b-245 α polypeptide (CYBA) 5'UTR; a hydroxysteroid (17- β) dehydrogenase (HSD17B4) 5'UTR; a Tobacco etch virus (TEV) 5'UTR; a Venezuelan equine encephalitis virus (VEEV) 5'UTR; a 5' proximal open reading frame of rubella virus (RV) RNA encoding nonstructural proteins; a Dengue virus (DEN) 5'UTR; a heat shock protein 70 (Hsp70) 5'UTR; a eIF4G 5'UTR; a GLUT1 5'UTR; functional fragments thereof and any combination thereof.

[0439] In some embodiments, the 3'UTR is selected from the group consisting of a β -globin 3'UTR; a CYBA 3'UTR; an albumin 3'UTR; a growth hormone (GH) 3'UTR; a VEEV 3'UTR; a hepatitis B virus (HBV) 3'UTR; α -globin 3'UTR; a DEN 3'UTR; a PAV barley yellow dwarf virus (BYDV-PAV) 3'UTR; an elongation factor 1 α 1 (EEF1A1) 3'UTR; a manganese superoxide dismutase (MnSOD) 3'UTR; a β subunit of mitochondrial H(+)-ATP synthase (β -mRNA) 3'UTR; a GLUT1 3'UTR; a MEF2A 3'UTR; a β -F1-ATPase 3'UTR; functional fragments thereof and combinations thereof.

[0440] Wild-type UTRs derived from any gene or mRNA can be incorporated into the polynucleotides of the invention. In some embodiments, a UTR can be altered relative to a wild type or native UTR to produce a variant UTR, e.g., by changing the orientation or location of the UTR relative to the ORF; or by inclusion of additional nucleotides, deletion of nucleotides, swapping or transposition of nucleotides. In some embodiments, variants of 5' or 3' UTRs can be utilized, for example, mutants of wild type UTRs, or variants wherein one or more nucleotides are added to or removed from a terminus of the UTR.

- [0441] Additionally, one or more synthetic UTRs can be used in combination with one or more non-synthetic UTRs. See, e.g., Mandal and Rossi, Nat. Protoc. 2013 8(3):568-82, and sequences available at www.addgene.org/Derrick_Rossi/, the contents of each are incorporated herein by reference in their entirety. UTRs or portions thereof can be placed in the same orientation as in the transcript from which they were selected or can be altered in orientation or location. Hence, a 5' and/or 3' UTR can be inverted, shortened, lengthened, or combined with one or more other 5' UTRs or 3' UTRs.
- [0442] In some embodiments, the polynucleotide comprises multiple UTRs, e.g., a double, a triple or a quadruple 5'UTR or 3'UTR. For example, a double UTR comprises two copies of the same UTR either in series or substantially in series. For example, a double beta-globin 3'UTR can be used (see US2010/0129877, the contents of which are incorporated herein by reference in its entirety).
- [0443] In certain embodiments, the polynucleotides of the invention comprise a 5'UTR and/or a 3'UTR selected from any of the UTRs disclosed herein.
- [0444] In some embodiments, the 5'UTR comprises:

5'UTR-001 (Upstream UTR)

(GGGAAAU AAGAGAGAAAAGAAGAGUAAGAAGAAAUUAU AAGAGCCACC)
(SEQ ID NO. 40);

5'UTR-002 (Upstream UTR)

(GGGAGAU CAGAGAGAAAAGAAGAGUAAGAAGAAAUUAU AAGAGCCACC)
(SEQ ID NO. 41);

5'UTR-003 (Upstream UTR)

(GGAAUAAAAGUCUCAACACAACAUUAUACAAAACAAACGAAUCUCAAGCAA
UCAAGCAUUCUACUUCUAUUGCAGCAAUUUAAAUCAUUUCUUUUAAGCA
AAAGCAAUUUUCUGAAAAUUUUCACCAUUUACGAACGAUAGCAAC) (SEQ ID
NO. 42);

5'UTR-004 (Upstream UTR)

(GGGAGACAAGCUUGGCAUUC CGGUACUGUUGGUAAAGCCACC) (SEQ ID
NO. 43);

5'UTR-005 (Upstream UTR)

(GGGAGAUCAAGAGAGAAAAGAAGAGUAAGAAGAAAUUAAGAGCCACC)
(SEQ ID NO. 44);

5'UTR-006 (Upstream UTR)

(GGAAUAAAAGUCUCAACACAACAUUAUACAAAACAAACGAAUCUCAAGCAA
UCAAGCAUUCUACUUCUAUUGCAGCAAUUUAAAUCAUUUCUUUUAAGCA
AAAGCAAUUUUCUGAAAAUUUUCACCAUUUACGAACGAUAGCAAC) (SEQ ID
NO. 45);

5'UTR-007 (Upstream UTR)

(GGGAGACAAGCUUGGCAUUCGCUACUGUUGGUAAAGCCACC) (SEQ ID
NO. 46);

5'UTR-008 (Upstream UTR)

(GGGAAUUAACAGAGAAAAGAAGAGUAAGAAGAAAUUAAGAGCCACC)
(SEQ ID NO. 47);

5'UTR-009 (Upstream UTR)

(GGGAAAUUAGACAGAAAAGAAGAGUAAGAAGAAAUUAAGAGCCACC)
(SEQ ID NO. 48);

5'UTR-010, Upstream

(GGGAAAUAAAGAGAGUAAAGAACAGUAAGAAGAAAUUAAGAGCCACC)
(SEQ ID NO. 49);

5'UTR-011 (Upstream UTR)

(GGGAAAAAAGAGAGAAAAGAAGACUAAGAAGAAAUUAAGAGCCACC)
(SEQ ID NO. 50);

5'UTR-012 (Upstream UTR)

(GGGAAAUAAAGAGAGAGAAAAGAAGAGUAAGAAGAUUAUAAGAGCCACC)

(SEQ ID NO. 51);

5'UTR-013 (Upstream UTR)

(GGGAAAUAAAGAGACAAAACAAGAGUAAGAAGAAAUUAAGAGCCACC)

(SEQ ID NO. 52);

5'UTR-014 (Upstream UTR)

(GGGAAAUUAGAGAGUAAAGAACAGUAAGUAGAAUUAAAAGAGCCACC)

(SEQ ID NO. 53);

5'UTR-15 (Upstream UTR)

(GGGAAAUAAAGAGAGAAUAGAAGAGUAAGAAGAAAUUAAGAGCCACC)

(SEQ ID NO. 54);

5'UTR-016 (Upstream UTR)

(GGGAAAUAAAGAGAGAAAAGAAGAGUAAGAAGAAAAUUAAGAGCCACC)

(SEQ ID NO. 55);

5'UTR-017 (Upstream UTR)

(GGGAAAUAAAGAGAGAAAAGAAGAGUAAGAAGAAAUUUAAGAGCCACC)

(SEQ ID NO. 56); or

5'UTR-018 (Upstream UTR)

(UCAAGCUUUUGGACCCUCGUACAGAAGCUAAUACGACUCACUAUAGGGAA

AUAAGAGAGAAAAGAAGAGUAAGAAGAAAUUAAGAGCCACC) (SEQ ID

NO. 57).

[0445] In some embodiments, the 3'UTR comprises:

142-3p 3'UTR (UTR including miR142-3p)

(UGAUAAUAGUCCAUAAGUAGGAAACACUACAGCUGGAGCCUCGGUGGCC

AUGCUUCUUGCCCCUUGGGCCUCCCCCAGCCCCUCCUCCCCUCCUGCACC
CGUACCCCCGUGGUCUUUGAAUAAAGUCUGAGUGGGCGGC) (SEQ ID NO.
58);

142-3p 2'UTR (UTR including miR142-3p)

(UGAUAAUAGGCUGGAGCCUCGGUGGCCAUAAAGUAGGAAACACUACAC
AUGCUUCUUGCCCCUUGGGCCUCCCCCAGCCCCUCCUCCCCUCCUGCACC
CGUACCCCCGUGGUCUUUGAAUAAAGUCUGAGUGGGCGGC) (SEQ ID NO.
59);

142-3p 3'UTR (UTR including miR142-3p)

(UGAUAAUAGGCUGGAGCCUCGGUGGCCAUGCUUCUUGCCCCUCCAUAAA
GUAGGAAACACUACAUGGGCCUCCCCCAGCCCCUCCUCCCCUCCUGCAC
CCGUACCCCCGUGGUCUUUGAAUAAAGUCUGAGUGGGCGGC) (SEQ ID NO.
60);

142-3p 3'UTR (UTR including miR142-3p)

(UGAUAAUAGGCUGGAGCCUCGGUGGCCAUGCUUCUUGCCCCUUGGGCCUC
CCCCCAGUCCAUAAAGUAGGAAACACUACACCCCUCCUCCCCUCCUGCAC
CCGUACCCCCGUGGUCUUUGAAUAAAGUCUGAGUGGGCGGC) (SEQ ID NO.
61);

142-3p 3'UTR (UTR including miR142-3p)

(UGAUAAUAGGCUGGAGCCUCGGUGGCCAUGCUUCUUGCCCCUUGGGCCUC
CCCCCAGCCCCUCCUCCCCUUCUCCAUAAAGUAGGAAACACUACACUGCAC
CCGUACCCCCGUGGUCUUUGAAUAAAGUCUGAGUGGGCGGC) (SEQ ID NO.
62);

142-3p 3'UTR (UTR including miR142-3p)

(UGAUAAUAGGCUGGAGCCUCGGUGGCCAUGCUUCUUGCCCCUUGGGCCUC
CCCCCAGCCCCUCCUCCCCUCCUGCACCCGUACCCCCUCCAUAAAGUAGGA
AACACUACAGUGGUCUUUGAAUAAAGUCUGAGUGGGCGGC) (SEQ ID NO.
63);

142-3p 3'UTR (UTR including miR142-3p)

(UGAUAAUAGGCUGGAGCCUCGGUGGCCAUGCUUCUUGCCCCUUGGGCCUC
CCCCAGCCCCUCCUCCCCUCCUGCACCCGUACCCCCGUGGUCUUUGAAUA
AAGUCCAUAAAGUAGGAAACACUACACUGAGUGGGCGGC) (SEQ ID NO.
64);

3'UTR-001 (Creatine Kinase UTR) (See SEQ ID NO. 65);

3'UTR-002 (Myoglobin UTR) (See SEQ ID NO. 66);

3'UTR-003 (α -actin UTR) (See SEQ ID NO. 67);

3'UTR-004 (Albumin UTR) (See SEQ ID NO. 68);

3'UTR-005 (α -globin UTR) (See SEQ ID NO. 69);

3'UTR-006 (G-CSF UTR) (See SEQ ID NO. 70);

3'UTR-007 (Col1a2; collagen, type I, alpha 2 UTR) (See SEQ ID NO. 71);

3'UTR-008 (Col6a2; collagen, type VI, alpha 2 UTR) (See SEQ ID NO. 72);

3'UTR-009 (RPN1; ribophorin I UTR) (See SEQ ID NO. 73);

3'UTR-010 (LRP1; low density lipoprotein receptor-related protein 1 UTR) (See SEQ ID
NO. 74);

3'UTR-011 (Nnt1; cardiotrophin-like cytokine factor 1 UTR) (See SEQ ID NO. 75);

3'UTR-012 (Col6a1; collagen, type VI, alpha 1 UTR) (See SEQ ID NO. 76);

3'UTR-013 (Calr; calreticulin UTR) (See SEQ ID NO. 77);

3'UTR-014 (Col1a1; collagen, type I, alpha 1 UTR) (See SEQ ID NO. 78);

3'UTR-015 (Plod1; procollagen-lysine, 2-oxoglutarate 5-dioxygenase 1 UTR) (See SEQ ID NO. 79);

3'UTR-016 (Nucb1; nucleobindin 1 UTR) (See SEQ ID NO. 80);

3'UTR-017 (α -globin) (See SEQ ID NO. 81);

3'UTR-018 (See SEQ ID NO. 82);

3'UTR (miR142+miR126 variant 1)

(UGAUAAUAGUCCAUAAAGUAGGAAACACUACAGCUGGAGCCUCGGUGGCC
AUGCUUCUUGCCCCUUGGGCCUCCCCCAGCCCCUCCUCCCCUUCUGCACC
CGUACCCCCCGCAUUAUUACUCACGGUACGAGUGGUCUUUGAAUAAAGUCU
GAGUGGGCGGC) (SEQ ID NO. 83);

3'UTR (miR142+miR126 variant 2)

(UGAUAAUAGUCCAUAAAGUAGGAAACACUACAGCUGGAGCCUCGGUGGCC
UAGCUUCUUGCCCCUUGGGCCUCCCCCAGCCCCUCCUCCCCUUCUGCACC
CGUACCCCCCGCAUUAUUACUCACGGUACGAGUGGUCUUUGAAUAAAGUCU
GAGUGGGCGGC) (SEQ ID NO. 84); or

3'UTR (miR142 binding site)

UGAUAAUAGGCUGGAGCCUCGGUGGCCUAGCUUCUUGCCCCUUGGGCCUCC
CCCCAGCCCCUCCUCCCCUUCUGCACCCGUACCCCCUCCAUAAGUAGGAA
ACACUACAGUGGUCUUUGAAUAAAGUCUGAGUGGGCGGC (SEQ ID NO. 121).

[0446] In certain embodiments, the 5'UTR and/or 3'UTR sequence of the invention comprises a nucleotide sequence at least about 60%, at least about 70%, at least about 80%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99%, or about 100% identical to a sequence selected from

the group consisting of 5'UTR sequences comprising any of SEQ ID NOs: 38, 40 to 57, and 117 to 119 and/or 3'UTR sequences comprising any of SEQ ID NOs: 39, 58 to 84, 90, 105, 108 to 115, and 120 to 130, and any combination thereof.

[0447] The polynucleotides of the invention can comprise combinations of features. For example, the ORF can be flanked by a 5'UTR that comprises a strong Kozak translational initiation signal and/or a 3'UTR comprising an oligo(dT) sequence for templated addition of a poly-A tail. A 5'UTR can comprise a first polynucleotide fragment and a second polynucleotide fragment from the same and/or different UTRs (see, e.g., US2010/0293625, herein incorporated by reference in its entirety).

[0448] Other non-UTR sequences can be used as regions or subregions within the polynucleotides of the invention. For example, introns or portions of intron sequences can be incorporated into the polynucleotides of the invention. Incorporation of intronic sequences can increase protein production as well as polynucleotide expression levels. In some embodiments, the polynucleotide of the invention comprises an internal ribosome entry site (IRES) instead of or in addition to a UTR (see, e.g., Yakubov et al., *Biochem. Biophys. Res. Commun.* 2010 394(1):189-193, the contents of which are incorporated herein by reference in their entirety). In some embodiments, the polynucleotide comprises an IRES instead of a 5'UTR sequence. In some embodiments, the polynucleotide comprises an ORF and a viral capsid sequence. In some embodiments, the polynucleotide comprises a synthetic 5'UTR in combination with a non-synthetic 3'UTR.

[0449] In some embodiments, the UTR can also include at least one translation enhancer polynucleotide, translation enhancer element, or translational enhancer elements (collectively, "TEE," which refers to nucleic acid sequences that increase the amount of polypeptide or protein produced from a polynucleotide. As a non-limiting example, the TEE can include those described in US2009/0226470, incorporated herein by reference in its entirety, and others known in the art. As a non-limiting example, the TEE can be located between the transcription promoter and the start codon. In some embodiments, the 5'UTR comprises a TEE.

[0450] In one aspect, a TEE is a conserved element in a UTR that can promote translational activity of a nucleic acid such as, but not limited to, cap-dependent or cap-independent translation.

- [0451] In one non-limiting example, the TEE comprises the TEE sequence in the 5'-leader of the Gtx homeodomain protein. See Chappell et al., PNAS 2004 101:9590-9594, incorporated herein by reference in its entirety.
- [0452] In some embodiments, the polynucleotide of the invention comprises one or multiple copies of a TEE. The TEE in a translational enhancer polynucleotide can be organized in one or more sequence segments. A sequence segment can harbor one or more of the TEEs provided herein, with each TEE being present in one or more copies. When multiple sequence segments are present in a translational enhancer polynucleotide, they can be homogenous or heterogeneous. Thus, the multiple sequence segments in a translational enhancer polynucleotide can harbor identical or different types of the TEE provided herein, identical or different number of copies of each of the TEE, and/or identical or different organization of the TEE within each sequence segment. In one embodiment, the polynucleotide of the invention comprises a translational enhancer polynucleotide sequence. Non-limiting examples of TEE sequences are described in U.S. Publication 2014/0200261, the contents of which are incorporated herein by reference in their entirety.

12. MicroRNA (miRNA) Binding Sites

- [0453] Polynucleotides of the invention can include regulatory elements, for example, microRNA (miRNA) binding sites, transcription factor binding sites, structured mRNA sequences and/or motifs, artificial binding sites engineered to act as pseudo-receptors for endogenous nucleic acid binding molecules, and combinations thereof. In some embodiments, polynucleotides including such regulatory elements are referred to as including "sensor sequences". Non-limiting examples of sensor sequences are described in U.S. Publication 2014/0200261, the contents of which are incorporated herein by reference in their entirety.
- [0454] In some embodiments, a polynucleotide (e.g., a ribonucleic acid (RNA), e.g., a messenger RNA (mRNA)) of the invention comprises an open reading frame (ORF) encoding a polypeptide of interest and further comprises one or more miRNA binding site(s). Inclusion or incorporation of miRNA binding site(s) provides for regulation of polynucleotides of the invention, and in turn, of the polypeptides encoded therefrom, based on tissue-specific and/or cell-type specific expression of naturally-occurring miRNAs.

- [0455] The present invention also provides pharmaceutical compositions and formulations that comprise any of the polynucleotides described above. In some embodiments, the composition or formulation further comprises a delivery agent.
- [0456] In some embodiments, the composition or formulation can contain a polynucleotide comprising a sequence optimized nucleic acid sequence disclosed herein which encodes a polypeptide. In some embodiments, the composition or formulation can contain a polynucleotide (e.g., a RNA, e.g., an mRNA) comprising a polynucleotide (e.g., an ORF) having significant sequence identity to a sequence optimized nucleic acid sequence disclosed herein which encodes a polypeptide. In some embodiments, the polynucleotide further comprises a miRNA binding site, e.g., a miRNA binding site that binds.
- [0457] A miRNA, e.g., a natural-occurring miRNA, is a 19-25 nucleotide long noncoding RNA that binds to a polynucleotide and down-regulates gene expression either by reducing stability or by inhibiting translation of the polynucleotide. A miRNA sequence comprises a “seed” region, i.e., a sequence in the region of positions 2-8 of the mature miRNA. A miRNA seed can comprise positions 2-8 or 2-7 of the mature miRNA. In some embodiments, a miRNA seed can comprise 7 nucleotides (e.g., nucleotides 2-8 of the mature miRNA), wherein the seed-complementary site in the corresponding miRNA binding site is flanked by an adenosine (A) opposed to miRNA position 1. In some embodiments, a miRNA seed can comprise 6 nucleotides (e.g., nucleotides 2-7 of the mature miRNA), wherein the seed-complementary site in the corresponding miRNA binding site is flanked by an adenosine (A) opposed to miRNA position 1. See, for example, Grimson A, Farh KK, Johnston WK, Garrett-Engle P, Lim LP, Bartel DP; Mol. Cell. 2007 Jul 6;27(1):91-105. miRNA profiling of the target cells or tissues can be conducted to determine the presence or absence of miRNA in the cells or tissues. In some embodiments, a polynucleotide (e.g., a ribonucleic acid (RNA), e.g., a messenger RNA (mRNA)) of the invention comprises one or more microRNA binding sites, microRNA target sequences, microRNA complementary sequences, or microRNA seed complementary sequences. Such sequences can correspond to, e.g., have complementarity to, any known microRNA such as those taught in US Publication US2005/0261218 and US Publication US2005/0059005, the contents of each of which are incorporated herein by reference in their entirety.

[0458] microRNAs derive enzymatically from regions of RNA transcripts that fold back on themselves to form short hairpin structures often termed a pre-miRNA (precursor-miRNA). A pre-miRNA typically has a two-nucleotide overhang at its 3' end, and has 3' hydroxyl and 5' phosphate groups. This precursor-mRNA is processed in the nucleus and subsequently transported to the cytoplasm where it is further processed by DICER (a RNase III enzyme), to form a mature microRNA of approximately 22 nucleotides. The mature microRNA is then incorporated into a ribonuclear particle to form the RNA-induced silencing complex, RISC, which mediates gene silencing. Art-recognized nomenclature for mature miRNAs typically designates the arm of the pre-miRNA from which the mature miRNA derives; "5p" means the microRNA is from the 5 prime arm of the pre-miRNA hairpin and "3p" means the microRNA is from the 3 prime end of the pre-miRNA hairpin. A miR referred to by number herein can refer to either of the two mature microRNAs originating from opposite arms of the same pre-miRNA (e.g., either the 3p or 5p microRNA). All miRs referred to herein are intended to include both the 3p and 5p arms/sequences, unless particularly specified by the 3p or 5p designation.

[0459] As used herein, the term "microRNA (miRNA or miR) binding site" refers to a sequence within a polynucleotide, e.g., within a DNA or within an RNA transcript, including in the 5'UTR and/or 3'UTR, that has sufficient complementarity to all or a region of a miRNA to interact with, associate with or bind to the miRNA. In some embodiments, a polynucleotide of the invention comprising an ORF encoding a polypeptide of interest and further comprises one or more miRNA binding site(s). In exemplary embodiments, a 5'UTR and/or 3'UTR of the polynucleotide (e.g., a ribonucleic acid (RNA), e.g., a messenger RNA (mRNA)) comprises the one or more miRNA binding site(s).

[0460] A miRNA binding site having sufficient complementarity to a miRNA refers to a degree of complementarity sufficient to facilitate miRNA-mediated regulation of a polynucleotide, e.g., miRNA-mediated translational repression or degradation of the polynucleotide. In exemplary aspects of the invention, a miRNA binding site having sufficient complementarity to the miRNA refers to a degree of complementarity sufficient to facilitate miRNA-mediated degradation of the polynucleotide, e.g., miRNA-guided RNA-induced silencing complex (RISC)-mediated cleavage of mRNA. The miRNA binding site can have complementarity to, for example, a 19-25 nucleotide long miRNA sequence, to a 19-23 long nucleotide miRNA sequence, or to a 22 nucleotide long

miRNA sequence. A miRNA binding site can be complementary to only a portion of a miRNA, e.g., to a portion less than 1, 2, 3, or 4 nucleotides of the full length of a naturally-occurring miRNA sequence, or to a portion less than 1, 2, 3, or 4 nucleotides shorter than a naturally-occurring miRNA sequence. Full or complete complementarity (e.g., full complementarity or complete complementarity over all or a significant portion of the length of a naturally-occurring miRNA) is preferred when the desired regulation is mRNA degradation.

[0461] In some embodiments, a miRNA binding site includes a sequence that has complementarity (e.g., partial or complete complementarity) with an miRNA seed sequence. In some embodiments, the miRNA binding site includes a sequence that has complete complementarity with a miRNA seed sequence. In some embodiments, a miRNA binding site includes a sequence that has complementarity (e.g., partial or complete complementarity) with an miRNA sequence. In some embodiments, the miRNA binding site includes a sequence that has complete complementarity with a miRNA sequence. In some embodiments, a miRNA binding site has complete complementarity with a miRNA sequence but for 1, 2, or 3 nucleotide substitutions, terminal additions, and/or truncations.

[0462] In some embodiments, the miRNA binding site is the same length as the corresponding miRNA. In other embodiments, the miRNA binding site is one, two, three, four, five, six, seven, eight, nine, ten, eleven or twelve nucleotide(s) shorter than the corresponding miRNA at the 5' terminus, the 3' terminus, or both. In still other embodiments, the microRNA binding site is two nucleotides shorter than the corresponding microRNA at the 5' terminus, the 3' terminus, or both. The miRNA binding sites that are shorter than the corresponding miRNAs are still capable of degrading the mRNA incorporating one or more of the miRNA binding sites or preventing the mRNA from translation.

[0463] In some embodiments, the miRNA binding site binds the corresponding mature miRNA that is part of an active RISC containing Dicer. In another embodiment, binding of the miRNA binding site to the corresponding miRNA in RISC degrades the mRNA containing the miRNA binding site or prevents the mRNA from being translated. In some embodiments, the miRNA binding site has sufficient complementarity to miRNA so that a RISC complex comprising the miRNA cleaves the polynucleotide comprising the miRNA binding site. In other embodiments, the miRNA binding site has imperfect

complementarity so that a RISC complex comprising the miRNA induces instability in the polynucleotide comprising the miRNA binding site. In another embodiment, the miRNA binding site has imperfect complementarity so that a RISC complex comprising the miRNA represses transcription of the polynucleotide comprising the miRNA binding site.

[0464] In some embodiments, the miRNA binding site has one, two, three, four, five, six, seven, eight, nine, ten, eleven or twelve mismatch(es) from the corresponding miRNA.

[0465] In some embodiments, the miRNA binding site has at least about ten, at least about eleven, at least about twelve, at least about thirteen, at least about fourteen, at least about fifteen, at least about sixteen, at least about seventeen, at least about eighteen, at least about nineteen, at least about twenty, or at least about twenty-one contiguous nucleotides complementary to at least about ten, at least about eleven, at least about twelve, at least about thirteen, at least about fourteen, at least about fifteen, at least about sixteen, at least about seventeen, at least about eighteen, at least about nineteen, at least about twenty, or at least about twenty-one, respectively, contiguous nucleotides of the corresponding miRNA.

[0466] By engineering one or more miRNA binding sites into a polynucleotide of the invention, the polynucleotide can be targeted for degradation or reduced translation, provided the miRNA in question is available. This can reduce off-target effects upon delivery of the polynucleotide. For example, if a polynucleotide of the invention is not intended to be to a tissue or cell but ends up in said tissue or cell, then a miRNA abundant in the tissue or cell can inhibit the expression of the gene of interest if one or multiple binding sites of the miRNA are engineered into the 5'UTR and/or 3'UTR of the polynucleotide. Thus, in some embodiments, incorporation of one or more miRNA binding sites into an mRNA of the disclosure may reduce the hazard of off-target effects upon nucleic acid molecule delivery and/or enable tissue-specific regulation of expression of a polypeptide encoded by the mRNA. In yet other embodiments, incorporation of one or more miRNA binding sites into an mRNA of the disclosure can modulate immune responses upon nucleic acid delivery *in vivo*. In further embodiments, incorporation of one or more miRNA binding sites into an mRNA of the disclosure can modulate accelerated blood clearance (ABC) of lipid-comprising compounds and compositions described herein.

- [0467] Conversely, miRNA binding sites can be removed from polynucleotide sequences in which they naturally occur in order to increase protein expression in specific tissues. For example, a binding site for a specific miRNA can be removed from a polynucleotide to improve protein expression in tissues or cells containing the miRNA.
- [0468] Regulation of expression in multiple tissues can be accomplished through introduction or removal of one or more miRNA binding sites, e.g., one or more distinct miRNA binding sites. The decision whether to remove or insert a miRNA binding site can be made based on miRNA expression patterns and/or their profiling in tissues and/or cells in development and/or disease. Identification of miRNAs, miRNA binding sites, and their expression patterns and role in biology have been reported (e.g., Bonauer et al., *Curr Drug Targets* 2010 11:943-949; Anand and Cheresch *Curr Opin Hematol* 2011 18:171-176; Contreras and Rao *Leukemia* 2012 26:404-413 (2011 Dec 20. doi: 10.1038/leu.2011.356); Bartel *Cell* 2009 136:215-233; Landgraf et al, *Cell*, 2007 129:1401-1414; Gentner and Naldini, *Tissue Antigens*. 2012 80:393-403 and all references therein; each of which is incorporated herein by reference in its entirety).
- [0469] miRNAs and miRNA binding sites can correspond to any known sequence, including non-limiting examples described in U.S. Publication Nos. 2014/0200261, 2005/0261218, and 2005/0059005, each of which are incorporated herein by reference in their entirety.
- [0470] Examples of tissues where miRNA are known to regulate mRNA, and thereby protein expression, include, but are not limited to, liver (miR-122), muscle (miR-133, miR-206, miR-208), endothelial cells (miR-17-92, miR-126), myeloid cells (miR-142-3p, miR-142-5p, miR-16, miR-21, miR-223, miR-24, miR-27), adipose tissue (let-7, miR-30c), heart (miR-1d, miR-149), kidney (miR-192, miR-194, miR-204), and lung epithelial cells (let-7, miR-133, miR-126).
- [0471] Specifically, miRNAs are known to be differentially expressed in immune cells (also called hematopoietic cells), such as antigen presenting cells (APCs) (e.g., dendritic cells and macrophages), macrophages, monocytes, B lymphocytes, T lymphocytes, granulocytes, natural killer cells, etc. Immune cell specific miRNAs are involved in immunogenicity, autoimmunity, the immune-response to infection, inflammation, as well as unwanted immune response after gene therapy and tissue/organ transplantation. Immune cells specific miRNAs also regulate many aspects of development, proliferation, differentiation and apoptosis of hematopoietic cells (immune cells). For example, miR-

142 and miR-146 are exclusively expressed in immune cells, particularly abundant in myeloid dendritic cells. It has been demonstrated that the immune response to a polynucleotide can be shut-off by adding miR-142 binding sites to the 3'-UTR of the polynucleotide, enabling more stable gene transfer in tissues and cells. miR-142 efficiently degrades exogenous polynucleotides in antigen presenting cells and suppresses cytotoxic elimination of transduced cells (e.g., Annoni A et al., blood, 2009, 114, 5152-5161; Brown BD, et al., Nat med. 2006, 12(5), 585-591; Brown BD, et al., blood, 2007, 110(13): 4144-4152, each of which is incorporated herein by reference in its entirety).

[0472] An antigen-mediated immune response can refer to an immune response triggered by foreign antigens, which, when entering an organism, are processed by the antigen presenting cells and displayed on the surface of the antigen presenting cells. T cells can recognize the presented antigen and induce a cytotoxic elimination of cells that express the antigen.

[0473] Introducing a miR-142 binding site into the 5'UTR and/or 3'UTR of a polynucleotide of the invention can selectively repress gene expression in antigen presenting cells through miR-142 mediated degradation, limiting antigen presentation in antigen presenting cells (e.g., dendritic cells) and thereby preventing antigen-mediated immune response after the delivery of the polynucleotide. The polynucleotide is then stably expressed in target tissues or cells without triggering cytotoxic elimination.

[0474] In one embodiment, binding sites for miRNAs that are known to be expressed in immune cells, in particular, antigen presenting cells, can be engineered into a polynucleotide of the invention to suppress the expression of the polynucleotide in antigen presenting cells through miRNA mediated RNA degradation, subduing the antigen-mediated immune response. Expression of the polynucleotide is maintained in non-immune cells where the immune cell specific miRNAs are not expressed. For example, in some embodiments, to prevent an immunogenic reaction against a liver specific protein, any miR-122 binding site can be removed and a miR-142 (and/or miR-146) binding site can be engineered into the 5'UTR and/or 3'UTR of a polynucleotide of the invention.

[0475] To further drive the selective degradation and suppression in APCs and macrophage, a polynucleotide of the invention can include a further negative regulatory element in the 5'UTR and/or 3'UTR, either alone or in combination with miR-142 and/or

miR-146 binding sites. As a non-limiting example, the further negative regulatory element is a Constitutive Decay Element (CDE).

[0476] Immune cell specific miRNAs include, but are not limited to, hsa-let-7a-2-3p, hsa-let-7a-3p, hsa-7a-5p, hsa-let-7c, hsa-let-7e-3p, hsa-let-7e-5p, hsa-let-7g-3p, hsa-let-7g-5p, hsa-let-7i-3p, hsa-let-7i-5p, miR-10a-3p, miR-10a-5p, miR-1184, hsa-let-7f-1--3p, hsa-let-7f-2--5p, hsa-let-7f-5p, miR-125b-1-3p, miR-125b-2-3p, miR-125b-5p, miR-1279, miR-130a-3p, miR-130a-5p, miR-132-3p, miR-132-5p, miR-142-3p, miR-142-5p, miR-143-3p, miR-143-5p, miR-146a-3p, miR-146a-5p, miR-146b-3p, miR-146b-5p, miR-147a, miR-147b, miR-148a-5p, miR-148a-3p, miR-150-3p, miR-150-5p, miR-151b, miR-155-3p, miR-155-5p, miR-15a-3p, miR-15a-5p, miR-15b-5p, miR-15b-3p, miR-16-1-3p, miR-16-2-3p, miR-16-5p, miR-17-5p, miR-181a-3p, miR-181a-5p, miR-181a-2-3p, miR-182-3p, miR-182-5p, miR-197-3p, miR-197-5p, miR-21-5p, miR-21-3p, miR-214-3p, miR-214-5p, miR-223-3p, miR-223-5p, miR-221-3p, miR-221-5p, miR-23b-3p, miR-23b-5p, miR-24-1-5p, miR-24-2-5p, miR-24-3p, miR-26a-1-3p, miR-26a-2-3p, miR-26a-5p, miR-26b-3p, miR-26b-5p, miR-27a-3p, miR-27a-5p, miR-27b-3p, miR-27b-5p, miR-28-3p, miR-28-5p, miR-2909, miR-29a-3p, miR-29a-5p, miR-29b-1-5p, miR-29b-2-5p, miR-29c-3p, miR-29c-5p, miR-30e-3p, miR-30e-5p, miR-331-5p, miR-339-3p, miR-339-5p, miR-345-3p, miR-345-5p, miR-346, miR-34a-3p, miR-34a-5p, miR-363-3p, miR-363-5p, miR-372, miR-377-3p, miR-377-5p, miR-493-3p, miR-493-5p, miR-542, miR-548b-5p, miR-548c-5p, miR-548i, miR-548j, miR-548n, miR-574-3p, miR-598, miR-718, miR-935, miR-99a-3p, miR-99a-5p, miR-99b-3p, and miR-99b-5p. Furthermore, novel miRNAs can be identified in immune cell through micro-array hybridization and microtome analysis (e.g., Jima DD et al, Blood, 2010, 116:e118-e127; Vaz C et al., BMC Genomics, 2010, 11,288, the content of each of which is incorporated herein by reference in its entirety.)

[0477] miRNAs that are known to be expressed in the liver include, but are not limited to, miR-107, miR-122-3p, miR-122-5p, miR-1228-3p, miR-1228-5p, miR-1249, miR-129-5p, miR-1303, miR-151a-3p, miR-151a-5p, miR-152, miR-194-3p, miR-194-5p, miR-199a-3p, miR-199a-5p, miR-199b-3p, miR-199b-5p, miR-296-5p, miR-557, miR-581, miR-939-3p, and miR-939-5p. MiRNA binding sites from any liver specific miRNA can be introduced to or removed from a polynucleotide of the invention to regulate expression of the polynucleotide in the liver. Liver specific miRNA binding sites can be engineered

alone or further in combination with immune cell (e.g., APC) miRNA binding sites in a polynucleotide of the invention.

[0478] miRNAs that are known to be expressed in the lung include, but are not limited to, let-7a-2-3p, let-7a-3p, let-7a-5p, miR-126-3p, miR-126-5p, miR-127-3p, miR-127-5p, miR-130a-3p, miR-130a-5p, miR-130b-3p, miR-130b-5p, miR-133a, miR-133b, miR-134, miR-18a-3p, miR-18a-5p, miR-18b-3p, miR-18b-5p, miR-24-1-5p, miR-24-2-5p, miR-24-3p, miR-296-3p, miR-296-5p, miR-32-3p, miR-337-3p, miR-337-5p, miR-381-3p, and miR-381-5p. miRNA binding sites from any lung specific miRNA can be introduced to or removed from a polynucleotide of the invention to regulate expression of the polynucleotide in the lung. Lung specific miRNA binding sites can be engineered alone or further in combination with immune cell (e.g., APC) miRNA binding sites in a polynucleotide of the invention.

[0479] miRNAs that are known to be expressed in the heart include, but are not limited to, miR-1, miR-133a, miR-133b, miR-149-3p, miR-149-5p, miR-186-3p, miR-186-5p, miR-208a, miR-208b, miR-210, miR-296-3p, miR-320, miR-451a, miR-451b, miR-499a-3p, miR-499a-5p, miR-499b-3p, miR-499b-5p, miR-744-3p, miR-744-5p, miR-92b-3p, and miR-92b-5p. MiRNA binding sites from any heart specific microRNA can be introduced to or removed from a polynucleotide of the invention to regulate expression of the polynucleotide in the heart. Heart specific miRNA binding sites can be engineered alone or further in combination with immune cell (e.g., APC) miRNA binding sites in a polynucleotide of the invention.

[0480] miRNAs that are known to be expressed in the nervous system include, but are not limited to, miR-124-5p, miR-125a-3p, miR-125a-5p, miR-125b-1-3p, miR-125b-2-3p, miR-125b-5p, miR-1271-3p, miR-1271-5p, miR-128, miR-132-5p, miR-135a-3p, miR-135a-5p, miR-135b-3p, miR-135b-5p, miR-137, miR-139-5p, miR-139-3p, miR-149-3p, miR-149-5p, miR-153, miR-181c-3p, miR-181c-5p, miR-183-3p, miR-183-5p, miR-190a, miR-190b, miR-212-3p, miR-212-5p, miR-219-1-3p, miR-219-2-3p, miR-23a-3p, miR-23a-5p, miR-30a-5p, miR-30b-3p, miR-30b-5p, miR-30c-1-3p, miR-30c-2-3p, miR-30c-5p, miR-30d-3p, miR-30d-5p, miR-329, miR-342-3p, miR-3665, miR-3666, miR-380-3p, miR-380-5p, miR-383, miR-410, miR-425-3p, miR-425-5p, miR-454-3p, miR-454-5p, miR-483, miR-510, miR-516a-3p, miR-548b-5p, miR-548c-5p, miR-571, miR-7-1-3p, miR-7-2-3p, miR-7-5p, miR-802, miR-922, miR-9-3p, and miR-9-5p. miRNAs enriched in the nervous system further include those specifically expressed in neurons,

including, but not limited to, miR-132-3p, miR-132-5p, miR-148b-3p, miR-148b-5p, miR-151a-3p, miR-151a-5p, miR-212-3p, miR-212-5p, miR-320b, miR-320e, miR-323a-3p, miR-323a-5p, miR-324-5p, miR-325, miR-326, miR-328, miR-922 and those specifically expressed in glial cells, including, but not limited to, miR-1250, miR-219-1-3p, miR-219-2-3p, miR-219-5p, miR-23a-3p, miR-23a-5p, miR-3065-3p, miR-3065-5p, miR-30e-3p, miR-30e-5p, miR-32-5p, miR-338-5p, and miR-657. miRNA binding sites from any CNS specific miRNA can be introduced to or removed from a polynucleotide of the invention to regulate expression of the polynucleotide in the nervous system. Nervous system specific miRNA binding sites can be engineered alone or further in combination with immune cell (e.g., APC) miRNA binding sites in a polynucleotide of the invention.

[0481] miRNAs that are known to be expressed in the pancreas include, but are not limited to, miR-105-3p, miR-105-5p, miR-184, miR-195-3p, miR-195-5p, miR-196a-3p, miR-196a-5p, miR-214-3p, miR-214-5p, miR-216a-3p, miR-216a-5p, miR-30a-3p, miR-33a-3p, miR-33a-5p, miR-375, miR-7-1-3p, miR-7-2-3p, miR-493-3p, miR-493-5p, and miR-944. MiRNA binding sites from any pancreas specific miRNA can be introduced to or removed from a polynucleotide of the invention to regulate expression of the polynucleotide in the pancreas. Pancreas specific miRNA binding sites can be engineered alone or further in combination with immune cell (e.g. APC) miRNA binding sites in a polynucleotide of the invention.

[0482] miRNAs that are known to be expressed in the kidney include, but are not limited to, miR-122-3p, miR-145-5p, miR-17-5p, miR-192-3p, miR-192-5p, miR-194-3p, miR-194-5p, miR-20a-3p, miR-20a-5p, miR-204-3p, miR-204-5p, miR-210, miR-216a-3p, miR-216a-5p, miR-296-3p, miR-30a-3p, miR-30a-5p, miR-30b-3p, miR-30b-5p, miR-30c-1-3p, miR-30c-2-3p, miR30c-5p, miR-324-3p, miR-335-3p, miR-335-5p, miR-363-3p, miR-363-5p, and miR-562. miRNA binding sites from any kidney specific miRNA can be introduced to or removed from a polynucleotide of the invention to regulate expression of the polynucleotide in the kidney. Kidney specific miRNA binding sites can be engineered alone or further in combination with immune cell (e.g., APC) miRNA binding sites in a polynucleotide of the invention.

[0483] miRNAs that are known to be expressed in the muscle include, but are not limited to, let-7g-3p, let-7g-5p, miR-1, miR-1286, miR-133a, miR-133b, miR-140-3p, miR-143-3p, miR-143-5p, miR-145-3p, miR-145-5p, miR-188-3p, miR-188-5p, miR-206, miR-208a, miR-208b, miR-25-3p, and miR-25-5p. MiRNA binding sites from any muscle

specific miRNA can be introduced to or removed from a polynucleotide of the invention to regulate expression of the polynucleotide in the muscle. Muscle specific miRNA binding sites can be engineered alone or further in combination with immune cell (e.g., APC) miRNA binding sites in a polynucleotide of the invention.

[0484] miRNAs are also differentially expressed in different types of cells, such as, but not limited to, endothelial cells, epithelial cells, and adipocytes.

[0485] miRNAs that are known to be expressed in endothelial cells include, but are not limited to, let-7b-3p, let-7b-5p, miR-100-3p, miR-100-5p, miR-101-3p, miR-101-5p, miR-126-3p, miR-126-5p, miR-1236-3p, miR-1236-5p, miR-130a-3p, miR-130a-5p, miR-17-5p, miR-17-3p, miR-18a-3p, miR-18a-5p, miR-19a-3p, miR-19a-5p, miR-19b-1-5p, miR-19b-2-5p, miR-19b-3p, miR-20a-3p, miR-20a-5p, miR-217, miR-210, miR-21-3p, miR-21-5p, miR-221-3p, miR-221-5p, miR-222-3p, miR-222-5p, miR-23a-3p, miR-23a-5p, miR-296-5p, miR-361-3p, miR-361-5p, miR-421, miR-424-3p, miR-424-5p, miR-513a-5p, miR-92a-1-5p, miR-92a-2-5p, miR-92a-3p, miR-92b-3p, and miR-92b-5p. Many novel miRNAs are discovered in endothelial cells from deep-sequencing analysis (e.g., Voellenkle C et al., RNA, 2012, 18, 472-484, herein incorporated by reference in its entirety). miRNA binding sites from any endothelial cell specific miRNA can be introduced to or removed from a polynucleotide of the invention to regulate expression of the polynucleotide in the endothelial cells.

[0486] miRNAs that are known to be expressed in epithelial cells include, but are not limited to, let-7b-3p, let-7b-5p, miR-1246, miR-200a-3p, miR-200a-5p, miR-200b-3p, miR-200b-5p, miR-200c-3p, miR-200c-5p, miR-338-3p, miR-429, miR-451a, miR-451b, miR-494, miR-802 and miR-34a, miR-34b-5p, miR-34c-5p, miR-449a, miR-449b-3p, miR-449b-5p specific in respiratory ciliated epithelial cells, let-7 family, miR-133a, miR-133b, miR-126 specific in lung epithelial cells, miR-382-3p, miR-382-5p specific in renal epithelial cells, and miR-762 specific in corneal epithelial cells. miRNA binding sites from any epithelial cell specific miRNA can be introduced to or removed from a polynucleotide of the invention to regulate expression of the polynucleotide in the epithelial cells.

[0487] In addition, a large group of miRNAs are enriched in embryonic stem cells, controlling stem cell self-renewal as well as the development and/or differentiation of various cell lineages, such as neural cells, cardiac, hematopoietic cells, skin cells, osteogenic cells and muscle cells (e.g., Kuppusamy KT et al., Curr. Mol Med, 2013,

13(5), 757-764; Vidigal JA and Ventura A, *Semin Cancer Biol.* 2012, 22(5-6), 428-436; Goff LA et al., *PLoS One*, 2009, 4:e7192; Morin RD et al., *Genome Res*, 2008, 18, 610-621; Yoo JK et al., *Stem Cells Dev.* 2012, 21(11), 2049-2057, each of which is herein incorporated by reference in its entirety). MiRNAs abundant in embryonic stem cells include, but are not limited to, let-7a-2-3p, let-a-3p, let-7a-5p, let7d-3p, let-7d-5p, miR-103a-2-3p, miR-103a-5p, miR-106b-3p, miR-106b-5p, miR-1246, miR-1275, miR-138-1-3p, miR-138-2-3p, miR-138-5p, miR-154-3p, miR-154-5p, miR-200c-3p, miR-200c-5p, miR-290, miR-301a-3p, miR-301a-5p, miR-302a-3p, miR-302a-5p, miR-302b-3p, miR-302b-5p, miR-302c-3p, miR-302c-5p, miR-302d-3p, miR-302d-5p, miR-302e, miR-367-3p, miR-367-5p, miR-369-3p, miR-369-5p, miR-370, miR-371, miR-373, miR-380-5p, miR-423-3p, miR-423-5p, miR-486-5p, miR-520c-3p, miR-548e, miR-548f, miR-548g-3p, miR-548g-5p, miR-548i, miR-548k, miR-548l, miR-548m, miR-548n, miR-548o-3p, miR-548o-5p, miR-548p, miR-664a-3p, miR-664a-5p, miR-664b-3p, miR-664b-5p, miR-766-3p, miR-766-5p, miR-885-3p, miR-885-5p, miR-93-3p, miR-93-5p, miR-941, miR-96-3p, miR-96-5p, miR-99b-3p and miR-99b-5p. Many predicted novel miRNAs are discovered by deep sequencing in human embryonic stem cells (e.g., Morin RD et al., *Genome Res*, 2008, 18, 610-621; Goff LA et al., *PLoS One*, 2009, 4:e7192; Bar M et al., *Stem cells*, 2008, 26, 2496-2505, the content of each of which is incorporated herein by reference in its entirety).

[0488] In one embodiment, the binding sites of embryonic stem cell specific miRNAs can be included in or removed from the 3'UTR of a polynucleotide of the invention to modulate the development and/or differentiation of embryonic stem cells, to inhibit the senescence of stem cells in a degenerative condition (e.g. degenerative diseases), or to stimulate the senescence and apoptosis of stem cells in a disease condition.

[0489] In some embodiments, miRNAs are selected based on expression and abundance in immune cells of the hematopoietic lineage, such as B cells, T cells, macrophages, dendritic cells, and cells that are known to express TLR7/ TLR8 and/or able to secrete cytokines such as endothelial cells and platelets. In some embodiments, the miRNA set thus includes miRs that may be responsible in part for the immunogenicity of these cells, and such that a corresponding miR-site incorporation in polynucleotides of the present invention (e.g., mRNAs) could lead to destabilization of the mRNA and/or suppression of translation from these mRNAs in the specific cell type. Non-limiting representative examples include miR-142, miR-144, miR-150, miR-155 and miR-223, which are

specific for many of the hematopoietic cells; miR-142, miR150, miR-16 and miR-223, which are expressed in B cells; miR-223, miR-451, miR-26a, miR-16, which are expressed in progenitor hematopoietic cells; and miR-126, which is expressed in plasmacytoid dendritic cells, platelets and endothelial cells. For further discussion of tissue expression of miRs see e.g., Teruel-Montoya, R. et al. (2014) *PLoS One* 9:e102259; Landgraf, P. et al. (2007) *Cell* 129:1401-1414; Bissels, U. et al. (2009) *RNA* 15:2375-2384. Any one miR-site incorporation in the 3'UTR and/or 5' UTR may mediate such effects in multiple cell types of interest (e.g., miR-142 is abundant in both B cells and dendritic cells).

[0490] In some embodiments, it may be beneficial to target the same cell type with multiple miRs and to incorporate binding sites to each of the 3p and 5p arm if both are abundant (e.g., both miR-142-3p and miR142-5p are abundant in hematopoietic stem cells). Thus, in certain embodiments, polynucleotides of the invention contain two or more (e.g., two, three, four or more) miR bindings sites from: (i) the group consisting of miR-142, miR-144, miR-150, miR-155 and miR-223 (which are expressed in many hematopoietic cells); or (ii) the group consisting of miR-142, miR150, miR-16 and miR-223 (which are expressed in B cells); or the group consisting of miR-223, miR-451, miR-26a, miR-16 (which are expressed in progenitor hematopoietic cells).

[0491] In some embodiments, it may also be beneficial to combine various miRs such that multiple cell types of interest are targeted at the same time (e.g., miR-142 and miR-126 to target many cells of the hematopoietic lineage and endothelial cells). Thus, for example, in certain embodiments, polynucleotides of the invention comprise two or more (e.g., two, three, four or more) miRNA bindings sites, wherein: (i) at least one of the miRs targets cells of the hematopoietic lineage (e.g., miR-142, miR-144, miR-150, miR-155 or miR-223) and at least one of the miRs targets plasmacytoid dendritic cells, platelets or endothelial cells (e.g., miR-126); or (ii) at least one of the miRs targets B cells (e.g., miR-142, miR150, miR-16 or miR-223) and at least one of the miRs targets plasmacytoid dendritic cells, platelets or endothelial cells (e.g., miR-126); or (iii) at least one of the miRs targets progenitor hematopoietic cells (e.g., miR-223, miR-451, miR-26a or miR-16) and at least one of the miRs targets plasmacytoid dendritic cells, platelets or endothelial cells (e.g., miR-126); or (iv) at least one of the miRs targets cells of the hematopoietic lineage (e.g., miR-142, miR-144, miR-150, miR-155 or miR-223), at least one of the miRs targets B cells (e.g., miR-142, miR150, miR-16 or miR-223) and at least

one of the miRs targets plasmacytoid dendritic cells, platelets or endothelial cells (e.g., miR-126); or any other possible combination of the foregoing four classes of miR binding sites (i.e., those targeting the hematopoietic lineage, those targeting B cells, those targeting progenitor hematopoietic cells and/or those targeting plasmacytoid dendritic cells/platelets/endothelial cells).

[0492] In one embodiment, to modulate immune responses, polynucleotides of the present invention can comprise one or more miRNA binding sequences that bind to one or more miRs that are expressed in conventional immune cells or any cell that expresses TLR7 and/or TLR8 and secrete pro-inflammatory cytokines and/or chemokines (e.g., in immune cells of peripheral lymphoid organs and/or splenocytes and/or endothelial cells). It has now been discovered that incorporation into an mRNA of one or more miRs that are expressed in conventional immune cells or any cell that expresses TLR7 and/or TLR8 and secrete pro-inflammatory cytokines and/or chemokines (e.g., in immune cells of peripheral lymphoid organs and/or splenocytes and/or endothelial cells) reduces or inhibits immune cell activation (e.g., B cell activation, as measured by frequency of activated B cells) and/or cytokine production (e.g., production of IL-6, IFN- γ and/or TNF α). Furthermore, it has now been discovered that incorporation into an mRNA of one or more miRs that are expressed in conventional immune cells or any cell that expresses TLR7 and/or TLR8 and secrete pro-inflammatory cytokines and/or chemokines (e.g., in immune cells of peripheral lymphoid organs and/or splenocytes and/or endothelial cells) can reduce or inhibit an anti-drug antibody (ADA) response against a protein of interest encoded by the mRNA.

[0493] In another embodiment, to modulate accelerated blood clearance of a polynucleotide delivered in a lipid-comprising compound or composition, polynucleotides of the invention can comprise one or more miR binding sequences that bind to one or more miRNAs expressed in conventional immune cells or any cell that expresses TLR7 and/or TLR8 and secrete pro-inflammatory cytokines and/or chemokines (e.g., in immune cells of peripheral lymphoid organs and/or splenocytes and/or endothelial cells). It has now been discovered that incorporation into an mRNA of one or more miR binding sites reduces or inhibits accelerated blood clearance (ABC) of the lipid-comprising compound or composition for use in delivering the mRNA. Furthermore, it has now been discovered that incorporation of one or more miR binding sites into an mRNA reduces serum levels of anti-PEG anti-IgM (e.g., reduces or inhibits

the acute production of IgMs that recognize polyethylene glycol (PEG) by B cells) and/or reduces or inhibits proliferation and/or activation of plasmacytoid dendritic cells following administration of a lipid-comprising compound or composition comprising the mRNA.

[0494] In some embodiments, miR sequences may correspond to any known microRNA expressed in immune cells, including but not limited to those taught in US Publication US2005/0261218 and US Publication US2005/0059005, the contents of which are incorporated herein by reference in their entirety. Non-limiting examples of miRs expressed in immune cells include those expressed in spleen cells, myeloid cells, dendritic cells, plasmacytoid dendritic cells, B cells, T cells and/or macrophages. For example, miR-142-3p, miR-142-5p, miR-16, miR-21, miR-223, miR-24 and miR-27 are expressed in myeloid cells, miR-155 is expressed in dendritic cells, B cells and T cells, miR-146 is upregulated in macrophages upon TLR stimulation and miR-126 is expressed in plasmacytoid dendritic cells. In certain embodiments, the miR(s) is expressed abundantly or preferentially in immune cells. For example, miR-142 (miR-142-3p and/or miR-142-5p), miR-126 (miR-126-3p and/or miR-126-5p), miR-146 (miR-146-3p and/or miR-146-5p) and miR-155 (miR-155-3p and/or miR-155-5p) are expressed abundantly in immune cells. These microRNA sequences are known in the art and, thus, one of ordinary skill in the art can readily design binding sequences or target sequences to which these microRNAs will bind based upon Watson-Crick complementarity.

[0495] Accordingly, in various embodiments, polynucleotides of the present invention comprise at least one microRNA binding site for a miR selected from the group consisting of miR-142, miR-146, miR-155, miR-126, miR-16, miR-21, miR-223, miR-24 and miR-27. In another embodiment, the mRNA comprises at least two miR binding sites for microRNAs expressed in immune cells. In various embodiments, the polynucleotide of the invention comprises 1-4, one, two, three or four miR binding sites for microRNAs expressed in immune cells. In another embodiment, the polynucleotide of the invention comprises three miR binding sites. These miR binding sites can be for microRNAs selected from the group consisting of miR-142, miR-146, miR-155, miR-126, miR-16, miR-21, miR-223, miR-24, miR-27, and combinations thereof. In one embodiment, the polynucleotide of the invention comprises two or more (e.g., two, three, four) copies of the same miR binding site expressed in immune cells, e.g., two or more copies of a miR

binding site selected from the group of miRs consisting of miR-142, miR-146, miR-155, miR-126, miR-16, miR-21, miR-223, miR-24, miR-27.

[0496] In one embodiment, the polynucleotide of the invention comprises three copies of the same miR binding site. In certain embodiments, use of three copies of the same miR binding site can exhibit beneficial properties as compared to use of a single miR binding site. Non-limiting examples of sequences for 3' UTRs containing three miR bindings sites are shown in SEQ ID NO: 108 (three miR-142-3p binding sites) and SEQ ID NO: 110 (three miR-142-5p binding sites).

[0497] In another embodiment, the polynucleotide of the invention comprises two or more (e.g., two, three, four) copies of at least two different miR binding sites expressed in immune cells. Non-limiting examples of sequences of 3' UTRs containing two or more different miR binding sites are shown in SEQ ID NO: 83 (one miR-142-3p binding site and one miR-126-3p binding site), SEQ ID NO: 84 (one miR-142-3p binding site and one miR-126-3p binding site); SEQ ID NO: 105 (one miR 126-3p binding site), SEQ ID NO: 111 (two miR-142-5p binding sites and one miR-142-3p binding sites) and SEQ ID NO: 114 (two miR-155-5p binding sites and one miR-142-3p binding sites).

[0498] In another embodiment, the polynucleotide of the invention comprises at least two miR binding sites for microRNAs expressed in immune cells, wherein one of the miR binding sites is for miR-142-3p. In various embodiments, the polynucleotide of the invention comprises binding sites for miR-142-3p and miR-155 (miR-155-3p or miR-155-5p), miR-142-3p and miR-146 (miR-146-3 or miR-146-5p), or miR-142-3p and miR-126 (miR-126-3p or miR-126-5p).

[0499] In another embodiment, the polynucleotide of the invention comprises at least two miR binding sites for microRNAs expressed in immune cells, wherein one of the miR binding sites is for miR-126-3p. In various embodiments, the polynucleotide of the invention comprises binding sites for miR-126-3p and miR-155 (miR-155-3p or miR-155-5p), miR-126-3p and miR-146 (miR-146-3p or miR-146-5p), or miR-126-3p and miR-142 (miR-142-3p or miR-142-5p).

[0500] In another embodiment, the polynucleotide of the invention comprises at least two miR binding sites for microRNAs expressed in immune cells, wherein one of the miR binding sites is for miR-142-5p. In various embodiments, the polynucleotide of the invention comprises binding sites for miR-142-5p and miR-155 (miR-155-3p or miR-

155-5p), miR-142-5p and miR-146 (miR-146-3 or miR-146-5p), or miR-142-5p and miR-126 (miR-126-3p or miR-126-5p).

[0501] In yet another embodiment, the polynucleotide of the invention comprises at least two miR binding sites for microRNAs expressed in immune cells, wherein one of the miR binding sites is for miR-155-5p. In various embodiments, the polynucleotide of the invention comprises binding sites for miR-155-5p and miR-142 (miR-142-3p or miR-142-5p), miR-155-5p and miR-146 (miR-146-3 or miR-146-5p), or miR-155-5p and miR-126 (miR-126-3p or miR-126-5p).

[0502] miRNA can also regulate complex biological processes such as angiogenesis (e.g., miR-132) (Anand and Cheresch Curr Opin Hematol 2011 18:171-176). In the polynucleotides of the invention, miRNA binding sites that are involved in such processes can be removed or introduced, in order to tailor the expression of the polynucleotides to biologically relevant cell types or relevant biological processes. In this context, the polynucleotides of the invention are defined as auxotrophic polynucleotides.

[0503] In some embodiments, a polynucleotide of the invention comprises a miRNA binding site, wherein the miRNA binding site comprises one or more nucleotide sequences selected from TABLE 3 or TABLE 4, including one or more copies of any one or more of the miRNA binding site sequences. In some embodiments, a polynucleotide of the invention further comprises at least one, two, three, four, five, six, seven, eight, nine, ten, or more of the same or different miRNA binding sites selected from TABLE 3, including any combination thereof.

[0504] In some embodiments, the miRNA binding site binds to miR-142 or is complementary to miR-142. In some embodiments, the miR-142 comprises SEQ ID NO:32. In some embodiments, the miRNA binding site binds to miR-142-3p or miR-142-5p. In some embodiments, the miR-142-3p binding site comprises SEQ ID NO:34. In some embodiments, the miR-142-5p binding site comprises SEQ ID NO:36. In some embodiments, the miRNA binding site comprises a nucleotide sequence at least 80%, at least 85%, at least 90%, at least 95%, or 100% identical to SEQ ID NO:34 or SEQ ID NO:36.

[0505] In some embodiments, the miRNA binding site binds to miR-126 or is complementary to miR-126. In some embodiments, the miR-126 comprises SEQ ID NO: 85. In some embodiments, the miRNA binding site binds to miR-126-3p or miR-126-5p. In some embodiments, the miR-126-3p binding site comprises SEQ ID NO: 87. In some

embodiments, the miR-126-5p binding site comprises SEQ ID NO: 89. In some embodiments, the miRNA binding site comprises a nucleotide sequence at least 80%, at least 85%, at least 90%, at least 95%, or 100% identical to SEQ ID NO: 87 or SEQ ID NO: 89.

[0506] In one embodiment, the 3' UTR comprises two miRNA binding sites, wherein a first miRNA binding site binds to miR-142 and a second miRNA binding site binds to miR-126. In a specific embodiment, the 3' UTR binding to miR-142 and miR-126 comprises, consists, or consists essentially of the sequence of SEQ ID NO: 83 or 84.

TABLE 3. miR-142, miR-126, and miR-142 and miR-126 binding sites

SEQ ID NO.	Description	Sequence
32	miR-142	GACAGUGCAGUCACCCAUAAAGUAGAAAGCACUA CUAACAGCACUGGAGGGUGUAGUGUUUCCUACUU UAUGGAUGAGUGUACUGUG
33	miR-142-3p	UGUAGUGUUUCCUACUUUAUGGA
34	miR-142-3p binding site	UCCAUAAAGUAGGAAACACUACA
35	miR-142-5p	CAUAAAGUAGAAAGCACUACU
36	miR-142-5p binding site	AGUAGUGC UUUCUACUUUAUG
85	miR-126	CGCUGGCGACGGGACAUUAUUACUUUUGGUACGC GCUGUGACACUCAAACUCGUACCGUGAGUAAUA AUGCGCCGUCCACGGCA
86	miR-126-3p	UCGUACCGUGAGUAAUAAUGCG
87	miR-126-3p binding site	CGCAUUAUUACUCACGGUACGA
88	miR-126-5p	CAUUAUUACUUUUGGUACGCG
89	miR-126-5p binding site	CGCGUACCAAAAGUAAUAAUG

[0507] In some embodiments, a miRNA binding site is inserted in the polynucleotide of the invention in any position of the polynucleotide (e.g., the 5'UTR and/or 3'UTR). In some embodiments, the 5'UTR comprises a miRNA binding site. In some embodiments, the 3'UTR comprises a miRNA binding site. In some embodiments, the 5'UTR and the

3'UTR comprise a miRNA binding site. The insertion site in the polynucleotide can be anywhere in the polynucleotide as long as the insertion of the miRNA binding site in the polynucleotide does not interfere with the translation of a functional polypeptide in the absence of the corresponding miRNA; and in the presence of the miRNA, the insertion of the miRNA binding site in the polynucleotide and the binding of the miRNA binding site to the corresponding miRNA are capable of degrading the polynucleotide or preventing the translation of the polynucleotide.

[0508] In some embodiments, a miRNA binding site is inserted in at least about 30 nucleotides downstream from the stop codon of an ORF in a polynucleotide of the invention comprising the ORF. In some embodiments, a miRNA binding site is inserted in at least about 10 nucleotides, at least about 15 nucleotides, at least about 20 nucleotides, at least about 25 nucleotides, at least about 30 nucleotides, at least about 35 nucleotides, at least about 40 nucleotides, at least about 45 nucleotides, at least about 50 nucleotides, at least about 55 nucleotides, at least about 60 nucleotides, at least about 65 nucleotides, at least about 70 nucleotides, at least about 75 nucleotides, at least about 80 nucleotides, at least about 85 nucleotides, at least about 90 nucleotides, at least about 95 nucleotides, or at least about 100 nucleotides downstream from the stop codon of an ORF in a polynucleotide of the invention. In some embodiments, a miRNA binding site is inserted in about 10 nucleotides to about 100 nucleotides, about 20 nucleotides to about 90 nucleotides, about 30 nucleotides to about 80 nucleotides, about 40 nucleotides to about 70 nucleotides, about 50 nucleotides to about 60 nucleotides, about 45 nucleotides to about 65 nucleotides downstream from the stop codon of an ORF in a polynucleotide of the invention.

[0509] In some embodiments, a miRNA binding site is inserted within the 3' UTR immediately following the stop codon of the coding region within the polynucleotide of the invention, e.g., mRNA. In some embodiments, if there are multiple copies of a stop codon in the construct, a miRNA binding site is inserted immediately following the final stop codon. In some embodiments, a miRNA binding site is inserted further downstream of the stop codon, in which case there are 3' UTR bases between the stop codon and the miR binding site(s). In some embodiments, three non-limiting examples of possible insertion sites for a miR in a 3' UTR are shown in SEQ ID NOs: 58, 59, and 115, which show a 3' UTR sequence with a miR-142-3p site inserted in one of three different possible insertion sites, respectively, within the 3' UTR.

[0510] In some embodiments, one or more miRNA binding sites can be positioned within the 5' UTR at one or more possible insertion sites. For example, three non-limiting examples of possible insertion sites for a miR in a 5' UTR are shown in SEQ ID NOs: 117, 118, and 119, which show a 5' UTR sequence with a miR-142-3p site inserted into one of three different possible insertion sites, respectively, within the 5' UTR.

[0511] In one embodiment, a codon optimized open reading frame encoding a polypeptide of interest comprises a stop codon and the at least one microRNA binding site is located within the 3' UTR 1-100 nucleotides after the stop codon. In one embodiment, the codon optimized open reading frame encoding the polypeptide of interest comprises a stop codon and the at least one microRNA binding site for a miR expressed in immune cells is located within the 3' UTR 30-50 nucleotides after the stop codon. In another embodiment, the codon optimized open reading frame encoding the polypeptide of interest comprises a stop codon and the at least one microRNA binding site for a miR expressed in immune cells is located within the 3' UTR at least 50 nucleotides after the stop codon. In other embodiments, the codon optimized open reading frame encoding the polypeptide of interest comprises a stop codon and the at least one microRNA binding site for a miR expressed in immune cells is located within the 3' UTR immediately after the stop codon, or within the 3' UTR 15-20 nucleotides after the stop codon or within the 3' UTR 70-80 nucleotides after the stop codon. In other embodiments, the 3'UTR comprises more than one miRNA binding site (e.g., 2-4 miRNA binding sites), wherein there can be a spacer region (e.g., of 10-100, 20-70 or 30-50 nucleotides in length) between each miRNA binding site. In another embodiment, the 3' UTR comprises a spacer region between the end of the miRNA binding site(s) and the poly A tail nucleotides. For example, a spacer region of 10-100, 20-70 or 30-50 nucleotides in length can be situated between the end of the miRNA binding site(s) and the beginning of the poly A tail.

[0512] In one embodiment, a codon optimized open reading frame encoding a polypeptide of interest comprises a start codon and the at least one microRNA binding site is located within the 5' UTR 1-100 nucleotides before (upstream of) the start codon. In one embodiment, the codon optimized open reading frame encoding the polypeptide of interest comprises a start codon and the at least one microRNA binding site for a miR expressed in immune cells is located within the 5' UTR 10-50 nucleotides before (upstream of) the start codon. In another embodiment, the codon optimized open reading

frame encoding the polypeptide of interest comprises a start codon and the at least one microRNA binding site for a miR expressed in immune cells is located within the 5' UTR at least 25 nucleotides before (upstream of) the start codon. In other embodiments, the codon optimized open reading frame encoding the polypeptide of interest comprises a start codon and the at least one microRNA binding site for a miR expressed in immune cells is located within the 5' UTR immediately before the start codon, or within the 5' UTR 15-20 nucleotides before the start codon or within the 5' UTR 70-80 nucleotides before the start codon. In other embodiments, the 5'UTR comprises more than one miRNA binding site (e.g., 2-4 miRNA binding sites), wherein there can be a spacer region (e.g., of 10-100, 20-70 or 30-50 nucleotides in length) between each miRNA binding site.

[0513] In one embodiment, the 3' UTR comprises more than one stop codon, wherein at least one miRNA binding site is positioned downstream of the stop codons. For example, a 3' UTR can comprise 1, 2 or 3 stop codons. Non-limiting examples of triple stop codons that can be used include: UGAUAAUAG, UGAUAGUAA, UAAUGAUAG, UGAUAAUAA, UGAUAGUAG, UAAUGAUGA, UAAUAGUAG, UGAUGAUGA, UAAUAAUAA and UAGUAGUAG. Within a 3' UTR, for example, 1, 2, 3 or 4 miRNA binding sites, e.g., miR-142-3p binding sites, can be positioned immediately adjacent to the stop codon(s) or at any number of nucleotides downstream of the final stop codon. When the 3' UTR comprises multiple miRNA binding sites, these binding sites can be positioned directly next to each other in the construct (i.e., one after the other) or, alternatively, spacer nucleotides can be positioned between each binding site.

[0514] In one embodiment, the 3' UTR comprises three stop codons with a single miR-142-3p binding site located downstream of the 3rd stop codon. Non-limiting examples of sequences of 3' UTR having three stop codons and a single miR-142-3p binding site located at different positions downstream of the final stop codon are shown in **Table 4** below.

TABLE 4. 3'UTRs, miR Sequences, and miR Binding Sites

SEQ ID NO:	Sequence
33	UGUAGUGUUUCCUACUUUAUGGA (miR 142-3p sequence)
34	UCCAUAAGUAGGAAACACUACA (miR 142-3p binding site)
35	CAUAAAGUAGAAAGCACUACU (miR 142-5p sequence)

87	CGCAUUAUUACUCACGGUACGA (miR 126-3p binding site)
91	CCUCUGAAAUUCAGUUCUUCAG (miR 146-3p sequence)
92	UGAGAACUGAAUCCAUGGGUU (miR 146-5p sequence)
93	CUCCUACAUAUUAGCAUUAACA (miR 155-3p sequence)
94	UUA AUGCUAAUCGUGAUAGGGGU (miR 155-5p sequence)
86	UCGUACCGUGAGUAAUAAUGCG (miR 126-3p sequence)
88	CAUUAUUACUUUUGGUACGCG (miR 126-5p sequence)
95	CCAGUAUUAACUGUGCUGCUGA (miR 16-3p sequence)
96	UAGCAGCACGUAAAUUUGGCG (miR 16-5p sequence)
97	CAACACCAGUCGAUGGGCUGU (miR 21-3p sequence)
98	UAGCUUAUCAGACUGAUGUUGA (miR 21-5p sequence)
99	UGUCAGUUUGUCAAUACCCCA (miR 223-3p sequence)
100	CGUGUAUUUGACAAGCUGAGUU (miR 223-5p sequence)
101	UGGCUCAGUUCAGCAGGAACAG (miR 24-3p sequence)
102	UGCCUACUGAGCUGAUUUCAGU (miR 24-5p sequence)
103	UUCACAGUGGCUAAGUCCGC (miR 27-3p sequence)
104	AGGGCUUAGCUGCUUGUGAGCA (miR 27-5p sequence)
106	UUA AUGCUAAUUGUGAUAGGGGU (miR 155-5p sequence)
107	ACCCCUAUCACAAUAGCAUUA (miR 155-5p binding site)
58	<u>UGAUAAUAG</u> UCCAUAAGUAGGAAACACUACAGCUGGAGCCUCGGUGGCCA UGCUUCUUGCCCCUUGGGCCUCCCCCAGCCCCUCCUCCCCUCCUGCACC CGUACCCCCGUGGUCUUUGAAUAAAGUCUGAGUGGGCGGC (3'UTR with miR 142-3p binding site, P1 insertion)
59	<u>UGAUAAUAG</u> GCUGGAGCCUCGGUGGCCUCCAUAAGUAGGAAACACUACACA UGCUUCUUGCCCCUUGGGCCUCCCCCAGCCCCUCCUCCCCUCCUGCACC CGUACCCCCGUGGUCUUUGAAUAAAGUCUGAGUGGGCGGC (3'UTR with miR 142-3p binding site, P2 insertion)
60	<u>UGAUAAUAG</u> GCUGGAGCCUCGGUGGCCAUGCUUCUUGCCCCUCCAUAAG UAGGAAACACUACAUGGGCCUCCCCCAGCCCCUCCUCCCCUCCUGCACC CGUACCCCCGUGGUCUUUGAAUAAAGUCUGAGUGGGCGGC

	(3' UTR including miR142-3p binding site)
61	UGAUAAUAGGCUGGAGCCUCGGUGGCCAUGCUUCUUGCCCCUUGGGCCUCC CCCCAGUCCAUAAAGUAGGAAACACUACACCCCUCCUCCCCUUCUGCACC CGUACCCCGUGGUCUUUGAAUAAAGUCUGAGUGGGCGGC (3'UTR including miR142-3p binding site)
62	UGAUAAUAGGCUGGAGCCUCGGUGGCCAUGCUUCUUGCCCCUUGGGCCUCC CCCCAGCCCCUCCUCCCCUUCUCCAUAAAGUAGGAAACACUACACUGCACC CGUACCCCGUGGUCUUUGAAUAAAGUCUGAGUGGGCGGC (3'UTR including miR142-3p binding site)
63	UGAUAAUAGGCUGGAGCCUCGGUGGCCAUGCUUCUUGCCCCUUGGGCCUCC CCCCAGCCCCUCCUCCCCUUCUGCACC CGUACCCCUCCAUAAAGUAGGA AACACUACAGUGGUCUUUGAAUAAAGUCUGAGUGGGCGGC (3' UTR with miR 142-3p binding site)
64	UGAUAAUAGGCUGGAGCCUCGGUGGCCAUGCUUCUUGCCCCUUGGGCCUCC CCCCAGCCCCUCCUCCCCUUCUGCACC CGUACCCCGUGGUCUUUGAAUA AAGUCCAUAAAGUAGGAAACACUACACUGAGUGGGCGGC (3'UTR including miR142-3p binding site)
82	UGAUAAUAGGCUGGAGCCUCGGUGGCCAUGCUUCUUGCCCCUUGGGCCUCC CCCCAGCCCCUCCUCCCCUUCUGCACC CGUACCCCGUGGUCUUUGAAUA AAGUCUGAGUGGGCGGC (3' UTR, no miR binding sites)
83	UGAUAAUAGUCCAUAAAGUAGGAAACACUACAGCUGGAGCCUCGGUGGCCA UGCUUCUUGCCCCUUGGGCCUCCCCCAGCCCCUCCUCCCCUUCUGCACC CGUACCCCGCAUUAUACUCACGGUACGAGUGGUCUUUGAAUAAAGUCU GAGUGGGCGGC (3' UTR with miR 142-3p and miR 126-3p binding sites)
84	UGAUAAUAGUCCAUAAAGUAGGAAACACUACAGCUGGAGCCUCGGUGGCCU AGCUUCUUGCCCCUUGGGCCUCCCCCAGCCCCUCCUCCCCUUCUGCACC CGUACCCCGCAUUAUACUCACGGUACGAGUGGUCUUUGAAUAAAGUCU GAGUGGGCGGC (3' UTR with miR 142-3p and miR 126-3p binding sites variant 2)
90	GCUGGAGCCUCGGUGGCCAUGCUUCUUGCCCCUUGGGCCUCCCCCAGCCC CUCCUCCCCUUCUGCACC CGUACCCCUCCAUAAAGUAGGAAACACUACA GUGGUCUUUGAAUAAAGUCUGAGUGGGCGGC (3'UTR with miR 142-3p binding site)
105	UGAUAAUAGGCUGGAGCCUCGGUGGCCAUGCUUCUUGCCCCUUGGGCCUCC CCCCAGCCCCUCCUCCCCUUCUGCACC CGUACCCCGCAUUAUACUCA CGGUACGAGUGGUCUUUGAAUAAAGUCUGAGUGGGCGGC (3'UTR with miR 126-3p binding site)
108	UGAUAAUAGUCCAUAAAGUAGGAAACACUACAGCUGGAGCCUCGGUGGCCA UGCUUCUUGCCCCUUGGGCCUCCAUAAAGUAGGAAACACUACAUCCCCCA GCCCCUCCUCCCCUUCUGCACC CGUACCCCUCCAUAAAGUAGGAAACAC UACAGUGGUCUUUGAAUAAAGUCUGAGUGGGCGGC (3' UTR with 3 miR 142-3p binding sites)
109	UGAUAAUAGGCUGGAGCCUCGGUGGCCAUGCUUCUUGCCCCUUGGGCCUCC CCCCAGCCCCUCCUCCCCUUCUGCACC CGUACCCCGAGUAGUGCUUUCUA CUUUAUGGUGGUCUUUGAAUAAAGUCUGAGUGGGCGGC (3'UTR with miR 142-5p binding site)
110	UGAUAAUAGAGUAGUGCUUUCUACUUAUGGCUGGAGCCUCGGUGGCCAUG

	CUUCUUGCCCCUUGGGCC <u>AGUAGUGCUUUCUACUUUAUG</u> UCCCCCAGCCC CUCCUCCCCUUCUGCACCCGUACCCCC <u>AGUAGUGCUUUCUACUUUAUG</u> GU GGUCUUUGAAUAAAGUCUGAGUGGGCGGC (3'UTR with 3 miR 142-5p binding sites)
111	<u>UGAUAAUAGAGUAGUGCUUUCUACUUUAUG</u> GCUGGAGCCUCGGUGGCCAUG CUUCUUGCCCCUUGGGCCUCCAUAAGUAGGAAACACUACAUCCCCCCAGC CCCUCUCCCCUUCUGCACCCGUACCCCC <u>AGUAGUGCUUUCUACUUUAUG</u> GUGGUCUUUGAAUAAAGUCUGAGUGGGCGGC (3'UTR with 2 miR 142-5p binding sites and 1 miR 142-3p binding site)
112	<u>UGAUAAUAG</u> GCUGGAGCCUCGGUGGCCAUGCUUCUUGCCCCUUGGGCCUCC CCCCAGCCCCUCCUCCCCUUCUGCACCCGUACCCCC <u>ACCCCUAUCACAAU</u> <u>UAGCAUUAAG</u> UGGUCUUUGAAUAAAGUCUGAGUGGGCGGC (3'UTR with miR 155-5p binding site)
113	<u>UGAUAAUAG</u> ACCCCUAUCACAAUUAAGCAUUAAGCUGGAGCCUCGGUGGCCA UGCUCUUGCCCCUUGGGCCACCCCUAUCACAAUUAAGCAUUAUCCCCCA GCCCCUCCUCCCCUUCUGCACCCGUACCCCCACCCCUAUCACAAUUAAGCA UUAAGUGGUCUUUGAAUAAAGUCUGAGUGGGCGGC (3' UTR with 3 miR 155-5p binding sites)
114	<u>UGAUAAUAG</u> ACCCCUAUCACAAUUAAGCAUUAAGCUGGAGCCUCGGUGGCCA UGCUCUUGCCCCUUGGGCCUCCAUAAGUAGGAAACACUACAUCCCCCA GCCCCUCCUCCCCUUCUGCACCCGUACCCCCACCCCUAUCACAAUUAAGCA UUAAGUGGUCUUUGAAUAAAGUCUGAGUGGGCGGC (3'UTR with 2 miR 155-5p binding sites and 1 miR 142-3p binding site)
115	<u>UGAUAAUAG</u> GCUGGAGCCUCGGUGGCCAUGCUUCUUGCCCCUUGGGCCUCC AUAAAGUAGGAAACACUACAUCCCCCCAGCCCCUCCUCCCCUUCUGCACC CGUACCCCCGUGGUCUUUGAAUAAAGUCUGAGUGGGCGGC (3'UTR with miR 142-3p binding site, P3 insertion)
116	AGUAGUGCUUUCUACUUUAUG (miR-142-5p binding site)
117	GGGAAUAAGAGUCCAUAAGUAGGAAACACUACAAGAAAAGAAGAGUAAG AAGAAUAUAAGAGCCACC (5' UTR with miR142-3p binding site at position p1)
118	GGGAAUAAGAGAGAAAAGAAGAGUAAUCCAUAAGUAGGAAACACUACAG AAGAAUAUAAGAGCCACC (5' UTR with miR142-3p binding site at position p2)
119	GGGAAUAAGAGAGAAAAGAAGAGUAGAAGAAUAUAUCCAUAAGUAG GAAACACUACAGAGCCACC (5' UTR with miR142-3p binding site at position p3)
120	<u>UGAUAAUAGAGUAGUGCUUUCUACUUUAUG</u> GCUGGAGCCUCGGUGGCCAUG CUUCUUGCCCCUUGGGCC <u>AGUAGUGCUUUCUACUUUAUG</u> UCCCCCAGCCC CUCUCCCCUUCUGCACCCGUACCCCC <u>AGUAGUGCUUUCUACUUUAUG</u> GUG GUCUUUGAAUAAAGUCUGAGUGGGCGGC (3'UTR with 3 miR 142-5p binding sites)
121	<u>UGAUAAUAG</u> GCUGGAGCCUCGGUGGCCUAGCUUCUUGCCCCUUGGGCCUCC CCCCAGCCCCUCCUCCCCUUCUGCACCCGUACCCCCUCCAUAAGUAGGA <u>AACACUACAGUGGUCUUUGAAUAAAGUCUGAGUGGGCGGC</u> (3' UTR with miR 142-3p binding site variant 2)
122	<u>UGAUAAUAG</u> GCUGGAGCCUCGGUGGCCUAGCUUCUUGCCCCUUGGGCCUCC CCCCAGCCCCUCCUCCCCUUCUGCACCCGUACCCCCGUGGUCUUUGAAUA AAGUCUGAGUGGGCGGC

	(3' UTR, no miR binding sites variant 2)
123	UGAUAAUAG GCUGGAGCCUCGGUGGCCUAGCUUCUUGCCCCUUGGGCCUCC CCCCAGCCCCUCCUCCCCUUCUGCACC CGUACCCCCCGCAUUAUACUCA <u>CGGUACGAGUGGUCUUUGAAUAAAGUCUGAGUGGGCGGC</u> (3' UTR with miR 126-3p binding site variant 3)
124	UGAUAAUAG UCCAUAAGUAGGAAACACUACAGCUGGAGCCUCGGUGGCCU AGCUUCUUGCCCCUUGGGCCUCCAUAAGUAGGAAACACUACAUCCCCCA GCCCCUCCUCCCCUUCUGCACC CGUACCCCCUCCAUAAGUAGGAAACAC <u>UACAGUGGUCUUUGAAUAAAGUCUGAGUGGGCGGC</u> (3' UTR with 3 miR 142-3p binding sites variant 2)
125	UGAUAAUAG UCCAUAAGUAGGAAACACUACAGCUGGAGCCUCGGUGGCCU AGCUUCUUGCCCCUUGGGCCUCCCCCAGCCCCUCCUCCCCUUCUGCACC CGUACCCCCGUGGUCUUUGAAUAAAGUCUGAGUGGGCGGC (3'UTR with miR 142-3p binding site, P1 insertion variant 2)
126	UGAUAAUAG GCUGGAGCCUCGGUGGCCUCCAUAAGUAGGAAACACUACACU AGCUUCUUGCCCCUUGGGCCUCCCCCAGCCCCUCCUCCCCUUCUGCACC CGUACCCCCGUGGUCUUUGAAUAAAGUCUGAGUGGGCGGC (3'UTR with miR 142-3p binding site, P2 insertion variant 2)
127	UGAUAAUAG GCUGGAGCCUCGGUGGCCUAGCUUCUUGCCCCUUGGGCCUCC AUAAAGUAGGAAACACUACAUCCCCCAGCCCCUCCUCCCCUUCUGCACC CGUACCCCCGUGGUCUUUGAAUAAAGUCUGAGUGGGCGGC (3'UTR with miR 142-3p binding site, P3 insertion variant 2)
128	UGAUAAUAG GCUGGAGCCUCGGUGGCCUAGCUUCUUGCCCCUUGGGCCUCC CCCCAGCCCCUCCUCCCCUUCUGCACC CGUACCCCCACCCCUAUCACAAU <u>UAGCAUUAAGUGGUCUUUGAAUAAAGUCUGAGUGGGCGGC</u> (3'UTR with miR 155-5p binding site variant 2)
129	UGAUAAUAG ACCCCUAUCACAAUUAAGCAUUAAGCUGGAGCCUCGGUGGCCU AGCUUCUUGCCCCUUGGGCCACCCCUAUCACAAUUAAGCAUUAUCCCCCA GCCCCUCCUCCCCUUCUGCACC CGUACCCCCACCCCUAUCACAAUUAAGCA <u>UUAAGUGGUCUUUGAAUAAAGUCUGAGUGGGCGGC</u> (3' UTR with 3 miR 155-5p binding sites variant 2)
130	UGAUAAUAG ACCCCUAUCACAAUUAAGCAUUAAGCUGGAGCCUCGGUGGCCU AGCUUCUUGCCCCUUGGGCCUCCAUAAGUAGGAAACACUACAUCCCCCA GCCCCUCCUCCCCUUCUGCACC CGUACCCCCACCCCUAUCACAAUUAAGCA <u>UUAAGUGGUCUUUGAAUAAAGUCUGAGUGGGCGGC</u> (3'UTR with 2 miR 155-5p binding sites and 1 miR 142-3p binding site variant 2)

Stop codon = bold

miR 142-3p binding site = underline

miR 126-3p binding site = bold underline

miR 155-5p binding site = shaded

miR 142-5p binding site = shaded and bold underline

[0515] In one embodiment, the polynucleotide of the invention comprises a 5' UTR, a codon optimized open reading frame encoding a polypeptide of interest, a 3' UTR comprising the at least one miRNA binding site for a miR expressed in immune cells, and a 3' tailing region of linked nucleosides. In various embodiments, the 3' UTR comprises

1-4, at least two, one, two, three or four miRNA binding sites for miRs expressed in immune cells, preferably abundantly or preferentially expressed in immune cells.

[0516] In one embodiment, the at least one miRNA expressed in immune cells is a miR-142-3p microRNA binding site. In one embodiment, the miR-142-3p microRNA binding site comprises the sequence shown in SEQ ID NO: 34. In one embodiment, the 3' UTR of the mRNA comprising the miR-142-3p microRNA binding site comprises the sequence shown in SEQ ID NO: 90.

[0517] In one embodiment, the at least one miRNA expressed in immune cells is a miR-126 microRNA binding site. In one embodiment, the miR-126 binding site is a miR-126-3p binding site. In one embodiment, the miR-126-3p microRNA binding site comprises the sequence shown in SEQ ID NO: 87. In one embodiment, the 3' UTR of the mRNA of the invention comprising the miR-126-3p microRNA binding site comprises the sequence shown in SEQ ID NO: 105.

[0518] Non-limiting exemplary sequences for miRs to which a microRNA binding site(s) of the disclosure can bind include the following: miR-142-3p (SEQ ID NO: 33), miR-142-5p (SEQ ID NO: 35), miR-146-3p (SEQ ID NO: 91), miR-146-5p (SEQ ID NO: 92), miR-155-3p (SEQ ID NO: 93), miR-155-5p (SEQ ID NO: 94), miR-126-3p (SEQ ID NO: 86), miR-126-5p (SEQ ID NO: 88), miR-16-3p (SEQ ID NO: 95), miR-16-5p (SEQ ID NO: 96), miR-21-3p (SEQ ID NO: 97), miR-21-5p (SEQ ID NO: 98), miR-223-3p (SEQ ID NO: 99), miR-223-5p (SEQ ID NO: 100), miR-24-3p (SEQ ID NO: 101), miR-24-5p (SEQ ID NO: 102), miR-27-3p (SEQ ID NO: 103) and miR-27-5p (SEQ ID NO: 104). Other suitable miR sequences expressed in immune cells (e.g., abundantly or preferentially expressed in immune cells) are known and available in the art, for example at the University of Manchester's microRNA database, miRBase. Sites that bind any of the aforementioned miRs can be designed based on Watson-Crick complementarity to the miR, typically 100% complementarity to the miR, and inserted into an mRNA construct of the disclosure as described herein.

[0519] In another embodiment, a polynucleotide of the present invention (e.g., and mRNA, e.g., the 3' UTR thereof) can comprise at least one miRNA binding site to thereby reduce or inhibit accelerated blood clearance, for example by reducing or inhibiting production of IgMs, e.g., against PEG, by B cells and/or reducing or inhibiting proliferation and/or activation of pDCs, and can comprise at least one miRNA binding site for modulating tissue expression of an encoded protein of interest.

- [0520] The distance between the miRNA binding sites can vary considerably; a number of different constructs have been tested with differing placement of the two miRNA binding sites and all have been functional. In certain embodiments, a nucleotide spacer is positioned between the two miRNA binding sites of a sufficient length to allow binding of RISC to each one. In one embodiment, the two miRNA binding sites are positioned about 40 bases apart from each other and the overall length of the 3' UTR is approximately 100-110 bases.
- [0521] miRNA gene regulation can be influenced by the sequence surrounding the miRNA such as, but not limited to, the species of the surrounding sequence, the type of sequence (e.g., heterologous, homologous, exogenous, endogenous, or artificial), regulatory elements in the surrounding sequence and/or structural elements in the surrounding sequence. The miRNA can be influenced by the 5'UTR and/or 3'UTR. As a non-limiting example, a non-human 3'UTR can increase the regulatory effect of the miRNA sequence on the expression of a polypeptide of interest compared to a human 3'UTR of the same sequence type.
- [0522] In one embodiment, other regulatory elements and/or structural elements of the 5'UTR can influence miRNA mediated gene regulation. One example of a regulatory element and/or structural element is a structured IRES (Internal Ribosome Entry Site) in the 5'UTR, which is necessary for the binding of translational elongation factors to initiate protein translation. EIF4A2 binding to this secondarily structured element in the 5'-UTR is necessary for miRNA mediated gene expression (Meijer HA et al., Science, 2013, 340, 82-85, herein incorporated by reference in its entirety). The polynucleotides of the invention can further include this structured 5'UTR in order to enhance microRNA mediated gene regulation.
- [0523] At least one miRNA binding site can be engineered into the 3'UTR of a polynucleotide of the invention. In this context, at least two, at least three, at least four, at least five, at least six, at least seven, at least eight, at least nine, at least ten, or more miRNA binding sites can be engineered into a 3'UTR of a polynucleotide of the invention. For example, 1 to 10, 1 to 9, 1 to 8, 1 to 7, 1 to 6, 1 to 5, 1 to 4, 1 to 3, 2, or 1 miRNA binding sites can be engineered into the 3'UTR of a polynucleotide of the invention. In one embodiment, miRNA binding sites incorporated into a polynucleotide of the invention can be the same or can be different miRNA sites. A combination of different miRNA binding sites incorporated into a polynucleotide of the invention can

include combinations in which more than one copy of any of the different miRNA sites are incorporated. In another embodiment, miRNA binding sites incorporated into a polynucleotide of the invention can target the same or different tissues in the body. As a non-limiting example, through the introduction of tissue-, cell-type-, or disease-specific miRNA binding sites in the 3'-UTR of a polynucleotide of the invention, the degree of expression in specific cell types (e.g., myeloid cells, endothelial cells, etc.) can be reduced.

[0524] In one embodiment, a miRNA binding site can be engineered near the 5' terminus of the 3'UTR, about halfway between the 5' terminus and 3' terminus of the 3'UTR and/or near the 3' terminus of the 3'UTR in a polynucleotide of the invention. As a non-limiting example, a miRNA binding site can be engineered near the 5' terminus of the 3'UTR and about halfway between the 5' terminus and 3' terminus of the 3'UTR. As another non-limiting example, a miRNA binding site can be engineered near the 3' terminus of the 3'UTR and about halfway between the 5' terminus and 3' terminus of the 3'UTR. As yet another non-limiting example, a miRNA binding site can be engineered near the 5' terminus of the 3'UTR and near the 3' terminus of the 3'UTR.

[0525] In another embodiment, a 3'UTR can comprise 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 miRNA binding sites. The miRNA binding sites can be complementary to a miRNA, miRNA seed sequence, and/or miRNA sequences flanking the seed sequence.

[0526] In some embodiments, the expression of a polynucleotide of the invention can be controlled by incorporating at least one sensor sequence in the polynucleotide and formulating the polynucleotide for administration. As a non-limiting example, a polynucleotide of the invention can be targeted to a tissue or cell by incorporating a miRNA binding site and formulating the polynucleotide in a lipid nanoparticle comprising an ionizable lipid, including any of the lipids described herein.

[0527] A polynucleotide of the invention can be engineered for more targeted expression in specific tissues, cell types, or biological conditions based on the expression patterns of miRNAs in the different tissues, cell types, or biological conditions. Through introduction of tissue-specific miRNA binding sites, a polynucleotide of the invention can be designed for optimal protein expression in a tissue or cell, or in the context of a biological condition.

[0528] In some embodiments, a polynucleotide of the invention can be designed to incorporate miRNA binding sites that either have 100% identity to known miRNA seed

sequences or have less than 100% identity to miRNA seed sequences. In some embodiments, a polynucleotide of the invention can be designed to incorporate miRNA binding sites that have at least: 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identity to known miRNA seed sequences. The miRNA seed sequence can be partially mutated to decrease miRNA binding affinity and as such result in reduced down modulation of the polynucleotide. In essence, the degree of match or mis-match between the miRNA binding site and the miRNA seed can act as a rheostat to more finely tune the ability of the miRNA to modulate protein expression. In addition, mutation in the non-seed region of a miRNA binding site can also impact the ability of a miRNA to modulate protein expression.

- [0529] In one embodiment, a miRNA sequence can be incorporated into the loop of a stem loop.
- [0530] In another embodiment, a miRNA seed sequence can be incorporated in the loop of a stem loop and a miRNA binding site can be incorporated into the 5' or 3' stem of the stem loop.
- [0531] In one embodiment, a translation enhancer element (TEE) can be incorporated on the 5' end of the stem of a stem loop and a miRNA seed can be incorporated into the stem of the stem loop. In another embodiment, a TEE can be incorporated on the 5' end of the stem of a stem loop, a miRNA seed can be incorporated into the stem of the stem loop and a miRNA binding site can be incorporated into the 3' end of the stem or the sequence after the stem loop. The miRNA seed and the miRNA binding site can be for the same and/or different miRNA sequences.
- [0532] In one embodiment, the incorporation of a miRNA sequence and/or a TEE sequence changes the shape of the stem loop region which can increase and/or decrease translation. (see e.g., Kedde et al., "A Pumilio-induced RNA structure switch in p27-3'UTR controls miR-221 and miR-22 accessibility." *Nature Cell Biology*. 2010, incorporated herein by reference in its entirety).
- [0533] In one embodiment, the 5'-UTR of a polynucleotide of the invention can comprise at least one miRNA sequence. The miRNA sequence can be, but is not limited to, a 19 or 22 nucleotide sequence and/or a miRNA sequence without the seed.
- [0534] In one embodiment the miRNA sequence in the 5'UTR can be used to stabilize a polynucleotide of the invention described herein.

[0535] In another embodiment, a miRNA sequence in the 5'UTR of a polynucleotide of the invention can be used to decrease the accessibility of the site of translation initiation such as, but not limited to a start codon. See, e.g., Matsuda et al., PLoS One. 2010 11(5):e15057; incorporated herein by reference in its entirety, which used antisense locked nucleic acid (LNA) oligonucleotides and exon-junction complexes (EJC) around a start codon (-4 to +37 where the A of the AUG codons is +1) in order to decrease the accessibility to the first start codon (AUG). Matsuda showed that altering the sequence around the start codon with an LNA or EJC affected the efficiency, length and structural stability of a polynucleotide. A polynucleotide of the invention can comprise a miRNA sequence, instead of the LNA or EJC sequence described by Matsuda et al, near the site of translation initiation in order to decrease the accessibility to the site of translation initiation. The site of translation initiation can be prior to, after or within the miRNA sequence. As a non-limiting example, the site of translation initiation can be located within a miRNA sequence such as a seed sequence or binding site.

[0536] In some embodiments, a polynucleotide of the invention can include at least one miRNA in order to dampen the antigen presentation by antigen presenting cells. The miRNA can be the complete miRNA sequence, the miRNA seed sequence, the miRNA sequence without the seed, or a combination thereof. As a non-limiting example, a miRNA incorporated into a polynucleotide of the invention can be specific to the hematopoietic system. As another non-limiting example, a miRNA incorporated into a polynucleotide of the invention to dampen antigen presentation is miR-142-3p.

[0537] In some embodiments, a polynucleotide of the invention can include at least one miRNA in order to dampen expression of the encoded polypeptide in a tissue or cell of interest. As a non-limiting example, a polynucleotide of the invention can include at least one miR-142-3p binding site, miR-142-3p seed sequence, miR-142-3p binding site without the seed, miR-142-5p binding site, miR-142-5p seed sequence, miR-142-5p binding site without the seed, miR-146 binding site, miR-146 seed sequence and/or miR-146 binding site without the seed sequence.

[0538] In some embodiments, a polynucleotide of the invention can comprise at least one miRNA binding site in the 3'UTR in order to selectively degrade mRNA therapeutics in the immune cells to subdue unwanted immunogenic reactions caused by therapeutic delivery. As a non-limiting example, the miRNA binding site can make a polynucleotide

of the invention more unstable in antigen presenting cells. Non-limiting examples of these miRNAs include mir-142-5p, mir-142-3p, mir-146a-5p, and mir-146-3p.

- [0539] In one embodiment, a polynucleotide of the invention comprises at least one miRNA sequence in a region of the polynucleotide that can interact with a RNA binding protein.
- [0540] In some embodiments, the polynucleotide of the invention (e.g., a RNA, e.g., an mRNA) comprising (i) a sequence-optimized nucleotide sequence (e.g., an ORF) encoding an ACADVL polypeptide (e.g., the wild-type sequence, functional fragment, or variant thereof) and (ii) a miRNA binding site (e.g., a miRNA binding site that binds to miR-142).
- [0541] In some embodiments, the polynucleotide of the invention comprises a uracil-modified sequence encoding an ACADVL polypeptide disclosed herein and a miRNA binding site disclosed herein, e.g., a miRNA binding site that binds to miR-142 and/or a miRNA binding site that binds to miR-126. In some embodiments, the polynucleotide of the invention comprises a uracil-modified sequence encoding a polypeptide disclosed herein and a miRNA binding site disclosed herein, e.g., a miRNA binding site that binds to miR-142miR-126, miR-142, miR-144, miR-146, miR-150, miR-155, miR-16, miR-21, miR-223, miR-24, miR-27 or miR-26a. In some embodiments, the miRNA binding site binds to miR126-3p, miR-142-3p, miR-142-5p, or miR-155. In some embodiments, the polynucleotide of the invention comprises a uracil-modified sequence encoding a polypeptide disclosed herein and at least two different microRNA binding sites, wherein the microRNA is expressed in an immune cell of hematopoietic lineage or a cell that expresses TLR7 and/or TLR8 and secretes pro-inflammatory cytokines and/or chemokines, and wherein the polynucleotide comprises one or more modified nucleobases. In some embodiments, the uracil-modified sequence encoding an ACADVL polypeptide comprises at least one chemically modified nucleobase, e.g., 5-methoxyuracil. In some embodiments, at least 95% of a type of nucleobase (e.g., uracil) in a uracil-modified sequence encoding an ACADVL polypeptide of the invention are modified nucleobases. In some embodiments, at least 95% of uracil in a uracil-modified sequence encoding an ACADVL polypeptide is 5-methoxyuridine. In some embodiments, the polynucleotide comprising a nucleotide sequence encoding an ACADVL polypeptide disclosed herein and a miRNA binding site is formulated with a delivery agent comprising, e.g., a compound having the Formula (I), e.g., any of Compounds 1-232, e.g.,

Compound 18; a compound having the Formula (III), (IV), (V), or (VI), e.g., any of Compounds 233-342, e.g., Compound 236; or a compound having the Formula (VIII), e.g., any of Compounds 419-428, e.g., Compound 428, or any combination thereof. In some embodiments, the delivery agent comprises Compound 18, DSPC, Cholesterol, and Compound 428, e.g., with a mole ratio of about 50:10:38.5:1.5.

13. 3' UTR

[0542] In certain embodiments, a polynucleotide of the present invention (e.g., a polynucleotide comprising a nucleotide sequence encoding an ACADVL polypeptide of the invention) further comprises a 3' UTR.

[0543] 3'-UTR is the section of mRNA that immediately follows the translation termination codon and often contains regulatory regions that post-transcriptionally influence gene expression. Regulatory regions within the 3'-UTR can influence polyadenylation, translation efficiency, localization, and stability of the mRNA. In one embodiment, the 3'-UTR useful for the invention comprises a binding site for regulatory proteins or microRNAs.

[0544] In certain embodiments, the 3' UTR useful for the polynucleotides of the invention comprises a 3'UTR selected from the group consisting of SEQ ID NO: 39, 58 to 84, 90, 105, 108 to 115, and 120 to 130, or any combination thereof. In some embodiments, the 3' UTR comprises a nucleic acid sequence selected from the group consisting of SEQ ID NOs: 83, 84, and 105, or any combination thereof.

[0545] In certain embodiments, the 3' UTR sequence useful for the invention comprises a nucleotide sequence at least about 60%, at least about 70%, at least about 80%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99%, or about 100% identical to a sequence selected from the group consisting of SEQ ID NO: 39, 58 to 84, 90, 105, 108 to 115, and 120 to 130, or any combination thereof.

14. Regions having a 5' Cap

[0546] The invention also includes a polynucleotide that comprises both a 5' Cap and a polynucleotide of the present invention (e.g., a polynucleotide comprising a nucleotide sequence encoding an ACADVL polypeptide).

- [0547] The 5' cap structure of a natural mRNA is involved in nuclear export, increasing mRNA stability and binds the mRNA Cap Binding Protein (CBP), which is responsible for mRNA stability in the cell and translation competency through the association of CBP with poly(A) binding protein to form the mature cyclic mRNA species. The cap further assists the removal of 5' proximal introns during mRNA splicing.
- [0548] Endogenous mRNA molecules can be 5'-end capped generating a 5'-ppp-5'-triphosphate linkage between a terminal guanosine cap residue and the 5'-terminal transcribed sense nucleotide of the mRNA molecule. This 5'-guanylate cap can then be methylated to generate an N7-methyl-guanylate residue. The ribose sugars of the terminal and/or anteterminal transcribed nucleotides of the 5' end of the mRNA can optionally also be 2'-O-methylated. 5'-decapping through hydrolysis and cleavage of the guanylate cap structure can target a nucleic acid molecule, such as an mRNA molecule, for degradation.
- [0549] In some embodiments, the polynucleotides of the present invention (e.g., a polynucleotide comprising a nucleotide sequence encoding an ACADVL polypeptide) incorporate a cap moiety.
- [0550] In some embodiments, polynucleotides of the present invention (e.g., a polynucleotide comprising a nucleotide sequence encoding an ACADVL polypeptide) comprise a non-hydrolyzable cap structure preventing decapping and thus increasing mRNA half-life. Because cap structure hydrolysis requires cleavage of 5'-ppp-5' phosphodiester linkages, modified nucleotides can be used during the capping reaction. For example, a Vaccinia Capping Enzyme from New England Biolabs (Ipswich, MA) can be used with α -thio-guanosine nucleotides according to the manufacturer's instructions to create a phosphorothioate linkage in the 5'-ppp-5' cap. Additional modified guanosine nucleotides can be used such as α -methyl-phosphonate and seleno-phosphate nucleotides.
- [0551] Additional modifications include, but are not limited to, 2'-O-methylation of the ribose sugars of 5'-terminal and/or 5'-anteterminal nucleotides of the polynucleotide (as mentioned above) on the 2'-hydroxyl group of the sugar ring. Multiple distinct 5'-cap structures can be used to generate the 5'-cap of a nucleic acid molecule, such as a polynucleotide that functions as an mRNA molecule. Cap analogs, which herein are also referred to as synthetic cap analogs, chemical caps, chemical cap analogs, or structural or functional cap analogs, differ from natural (*i.e.*, endogenous, wild-type or physiological) 5'-caps in their chemical structure, while retaining cap function. Cap analogs can be

chemically (*i.e.*, non-enzymatically) or enzymatically synthesized and/or linked to the polynucleotides of the invention.

- [0552] For example, the Anti-Reverse Cap Analog (ARCA) cap contains two guanines linked by a 5'-5'-triphosphate group, wherein one guanine contains an N7 methyl group as well as a 3'-O-methyl group (*i.e.*, N7,3'-O-dimethyl-guanosine-5'-triphosphate-5'-guanosine (m⁷G-3'mppp-G; which can equivalently be designated 3' O-Me-m⁷G(5')ppp(5')G). The 3'-O atom of the other, unmodified, guanine becomes linked to the 5'-terminal nucleotide of the capped polynucleotide. The N7- and 3'-O-methylated guanine provides the terminal moiety of the capped polynucleotide.
- [0553] Another exemplary cap is mCAP, which is similar to ARCA but has a 2'-O-methyl group on guanosine (*i.e.*, N7,2'-O-dimethyl-guanosine-5'-triphosphate-5'-guanosine, m⁷Gm-ppp-G).
- [0554] In some embodiments, the cap is a dinucleotide cap analog. As a non-limiting example, the dinucleotide cap analog can be modified at different phosphate positions with a boranophosphate group or a phosphoroselenoate group such as the dinucleotide cap analogs described in U.S. Patent No. US 8,519,110, the contents of which are herein incorporated by reference in its entirety.
- [0555] In another embodiment, the cap is a cap analog is a N7-(4-chlorophenoxyethyl) substituted dinucleotide form of a cap analog known in the art and/or described herein. Non-limiting examples of a N7-(4-chlorophenoxyethyl) substituted dinucleotide form of a cap analog include a N7-(4-chlorophenoxyethyl)-G(5')ppp(5')G and a N7-(4-chlorophenoxyethyl)-m^{3'-O}G(5')ppp(5')G cap analog (*See, e.g.*, the various cap analogs and the methods of synthesizing cap analogs described in Kore et al. Bioorganic & Medicinal Chemistry 2013 21:4570-4574; the contents of which are herein incorporated by reference in its entirety). In another embodiment, a cap analog of the present invention is a 4-chloro/bromophenoxyethyl analog.
- [0556] While cap analogs allow for the concomitant capping of a polynucleotide or a region thereof, in an *in vitro* transcription reaction, up to 20% of transcripts can remain uncapped. This, as well as the structural differences of a cap analog from an endogenous 5'-cap structures of nucleic acids produced by the endogenous, cellular transcription machinery, can lead to reduced translational competency and reduced cellular stability.
- [0557] Polynucleotides of the invention (e.g., a polynucleotide comprising a nucleotide sequence encoding an ACADVL polypeptide) can also be capped post-manufacture

(whether IVT or chemical synthesis), using enzymes, in order to generate more authentic 5'-cap structures. As used herein, the phrase "more authentic" refers to a feature that closely mirrors or mimics, either structurally or functionally, an endogenous or wild type feature. That is, a "more authentic" feature is better representative of an endogenous, wild-type, natural or physiological cellular function and/or structure as compared to synthetic features or analogs, etc., of the prior art, or which outperforms the corresponding endogenous, wild-type, natural or physiological feature in one or more respects. Non-limiting examples of more authentic 5'cap structures of the present invention are those that, among other things, have enhanced binding of cap binding proteins, increased half-life, reduced susceptibility to 5' endonucleases and/or reduced 5'decapping, as compared to synthetic 5'cap structures known in the art (or to a wild-type, natural or physiological 5'cap structure). For example, recombinant Vaccinia Virus Capping Enzyme and recombinant 2'-O-methyltransferase enzyme can create a canonical 5'-5'-triphosphate linkage between the 5'-terminal nucleotide of a polynucleotide and a guanine cap nucleotide wherein the cap guanine contains an N7 methylation and the 5'-terminal nucleotide of the mRNA contains a 2'-O-methyl. Such a structure is termed the Cap1 structure. This cap results in a higher translational-competency and cellular stability and a reduced activation of cellular pro-inflammatory cytokines, as compared, *e.g.*, to other 5'cap analog structures known in the art. Cap structures include, but are not limited to, 7mG(5')ppp(5')N,pN2p (cap 0), 7mG(5')ppp(5')NlmpNp (cap 1), and 7mG(5')-ppp(5')NlmpN2mp (cap 2).

[0558] As a non-limiting example, capping chimeric polynucleotides post-manufacture can be more efficient as nearly 100% of the chimeric polynucleotides can be capped. This is in contrast to ~80% when a cap analog is linked to a chimeric polynucleotide in the course of an *in vitro* transcription reaction.

[0559] According to the present invention, 5' terminal caps can include endogenous caps or cap analogs. According to the present invention, a 5' terminal cap can comprise a guanine analog. Useful guanine analogs include, but are not limited to, inosine, N1-methyl-guanosine, 2'fluoro-guanosine, 7-deaza-guanosine, 8-oxo-guanosine, 2-amino-guanosine, LNA-guanosine, and 2-azido-guanosine.

15. Poly-A Tails

- [0560] In some embodiments, the polynucleotides of the present disclosure (e.g., a polynucleotide comprising a nucleotide sequence encoding an ACADVL polypeptide) further comprise a poly-A tail. In further embodiments, terminal groups on the poly-A tail can be incorporated for stabilization. In other embodiments, a poly-A tail comprises des-3' hydroxyl tails.
- [0561] During RNA processing, a long chain of adenine nucleotides (poly-A tail) can be added to a polynucleotide such as an mRNA molecule in order to increase stability. Immediately after transcription, the 3' end of the transcript can be cleaved to free a 3' hydroxyl. Then poly-A polymerase adds a chain of adenine nucleotides to the RNA. The process, called polyadenylation, adds a poly-A tail that can be between, for example, approximately 80 to approximately 250 residues long, including approximately 80, 90, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, 200, 210, 220, 230, 240 or 250 residues long.
- [0562] PolyA tails can also be added after the construct is exported from the nucleus.
- [0563] According to the present invention, terminal groups on the poly A tail can be incorporated for stabilization. Polynucleotides of the present invention can include des-3' hydroxyl tails. They can also include structural moieties or 2'-O-methyl modifications as taught by Junjie Li, et al. (Current Biology, Vol. 15, 1501–1507, August 23, 2005, the contents of which are incorporated herein by reference in its entirety).
- [0564] The polynucleotides of the present invention can be designed to encode transcripts with alternative polyA tail structures including histone mRNA. According to Norbury, "Terminal uridylation has also been detected on human replication-dependent histone mRNAs. The turnover of these mRNAs is thought to be important for the prevention of potentially toxic histone accumulation following the completion or inhibition of chromosomal DNA replication. These mRNAs are distinguished by their lack of a 3' poly(A) tail, the function of which is instead assumed by a stable stem-loop structure and its cognate stem-loop binding protein (SLBP); the latter carries out the same functions as those of PABP on polyadenylated mRNAs" (Norbury, "Cytoplasmic RNA: a case of the tail wagging the dog," Nature Reviews Molecular Cell Biology; AOP, published online 29 August 2013; doi:10.1038/nrm3645) the contents of which are incorporated herein by reference in its entirety.

[0565] Unique poly-A tail lengths provide certain advantages to the polynucleotides of the present invention. Generally, the length of a poly-A tail, when present, is greater than 30 nucleotides in length. In another embodiment, the poly-A tail is greater than 35 nucleotides in length (*e.g.*, at least or greater than about 35, 40, 45, 50, 55, 60, 70, 80, 90, 100, 120, 140, 160, 180, 200, 250, 300, 350, 400, 450, 500, 600, 700, 800, 900, 1,000, 1,100, 1,200, 1,300, 1,400, 1,500, 1,600, 1,700, 1,800, 1,900, 2,000, 2,500, and 3,000 nucleotides).

[0566] In some embodiments, the polynucleotide or region thereof includes from about 30 to about 3,000 nucleotides (*e.g.*, from 30 to 50, from 30 to 100, from 30 to 250, from 30 to 500, from 30 to 750, from 30 to 1,000, from 30 to 1,500, from 30 to 2,000, from 30 to 2,500, from 50 to 100, from 50 to 250, from 50 to 500, from 50 to 750, from 50 to 1,000, from 50 to 1,500, from 50 to 2,000, from 50 to 2,500, from 50 to 3,000, from 100 to 500, from 100 to 750, from 100 to 1,000, from 100 to 1,500, from 100 to 2,000, from 100 to 2,500, from 100 to 3,000, from 500 to 750, from 500 to 1,000, from 500 to 1,500, from 500 to 2,000, from 500 to 2,500, from 500 to 3,000, from 1,000 to 1,500, from 1,000 to 2,000, from 1,000 to 2,500, from 1,000 to 3,000, from 1,500 to 2,000, from 1,500 to 2,500, from 1,500 to 3,000, from 2,000 to 3,000, from 2,000 to 2,500, and from 2,500 to 3,000).

[0567] In some embodiments, the poly-A tail is designed relative to the length of the overall polynucleotide or the length of a particular region of the polynucleotide. This design can be based on the length of a coding region, the length of a particular feature or region or based on the length of the ultimate product expressed from the polynucleotides.

[0568] In this context, the poly-A tail can be 10, 20, 30, 40, 50, 60, 70, 80, 90, or 100% greater in length than the polynucleotide or feature thereof. The poly-A tail can also be designed as a fraction of the polynucleotides to which it belongs. In this context, the poly-A tail can be 10, 20, 30, 40, 50, 60, 70, 80, or 90% or more of the total length of the construct, a construct region or the total length of the construct minus the poly-A tail. Further, engineered binding sites and conjugation of polynucleotides for Poly-A binding protein can enhance expression.

[0569] Additionally, multiple distinct polynucleotides can be linked together via the PABP (Poly-A binding protein) through the 3'-end using modified nucleotides at the 3'-terminus of the poly-A tail. Transfection experiments can be conducted in relevant cell

lines at and protein production can be assayed by ELISA at 12hr, 24hr, 48hr, 72hr and day 7 post-transfection.

[0570] In some embodiments, the polynucleotides of the present invention are designed to include a polyA-G Quartet region. The G-quartet is a cyclic hydrogen bonded array of four guanine nucleotides that can be formed by G-rich sequences in both DNA and RNA. In this embodiment, the G-quartet is incorporated at the end of the poly-A tail. The resultant polynucleotide is assayed for stability, protein production and other parameters including half-life at various time points. It has been discovered that the polyA-G quartet results in protein production from an mRNA equivalent to at least 75% of that seen using a poly-A tail of 120 nucleotides alone.

16. Start codon region

[0571] The invention also includes a polynucleotide that comprises both a start codon region and the polynucleotide described herein (e.g., a polynucleotide comprising a nucleotide sequence encoding an ACADVL polypeptide). In some embodiments, the polynucleotides of the present invention can have regions that are analogous to or function like a start codon region.

[0572] In some embodiments, the translation of a polynucleotide can initiate on a codon that is not the start codon AUG. Translation of the polynucleotide can initiate on an alternative start codon such as, but not limited to, ACG, AGG, AAG, CTG/CUG, GTG/GUG, ATA/AUA, ATT/AUU, TTG/UUG (see Touriol et al. *Biology of the Cell* 95 (2003) 169-178 and Matsuda and Mauro *PLoS ONE*, 2010 5:11; the contents of each of which are herein incorporated by reference in its entirety).

[0573] As a non-limiting example, the translation of a polynucleotide begins on the alternative start codon ACG. As another non-limiting example, polynucleotide translation begins on the alternative start codon CTG or CUG. As yet another non-limiting example, the translation of a polynucleotide begins on the alternative start codon GTG or GUG.

[0574] Nucleotides flanking a codon that initiates translation such as, but not limited to, a start codon or an alternative start codon, are known to affect the translation efficiency, the length and/or the structure of the polynucleotide. (*See, e.g.,* Matsuda and Mauro *PLoS ONE*, 2010 5:11; the contents of which are herein incorporated by reference in its entirety). Masking any of the nucleotides flanking a codon that initiates translation can be

used to alter the position of translation initiation, translation efficiency, length and/or structure of a polynucleotide.

[0575] In some embodiments, a masking agent can be used near the start codon or alternative start codon in order to mask or hide the codon to reduce the probability of translation initiation at the masked start codon or alternative start codon. Non-limiting examples of masking agents include antisense locked nucleic acids (LNA) polynucleotides and exon-junction complexes (EJCs) (*See, e.g.,* Matsuda and Mauro describing masking agents LNA polynucleotides and EJCs (PLoS ONE, 2010 5:11); the contents of which are herein incorporated by reference in its entirety).

[0576] In another embodiment, a masking agent can be used to mask a start codon of a polynucleotide in order to increase the likelihood that translation will initiate on an alternative start codon. In some embodiments, a masking agent can be used to mask a first start codon or alternative start codon in order to increase the chance that translation will initiate on a start codon or alternative start codon downstream to the masked start codon or alternative start codon.

[0577] In some embodiments, a start codon or alternative start codon can be located within a perfect complement for a miRNA binding site. The perfect complement of a miRNA binding site can help control the translation, length and/or structure of the polynucleotide similar to a masking agent. As a non-limiting example, the start codon or alternative start codon can be located in the middle of a perfect complement for a miRNA binding site. The start codon or alternative start codon can be located after the first nucleotide, second nucleotide, third nucleotide, fourth nucleotide, fifth nucleotide, sixth nucleotide, seventh nucleotide, eighth nucleotide, ninth nucleotide, tenth nucleotide, eleventh nucleotide, twelfth nucleotide, thirteenth nucleotide, fourteenth nucleotide, fifteenth nucleotide, sixteenth nucleotide, seventeenth nucleotide, eighteenth nucleotide, nineteenth nucleotide, twentieth nucleotide or twenty-first nucleotide.

[0578] In another embodiment, the start codon of a polynucleotide can be removed from the polynucleotide sequence in order to have the translation of the polynucleotide begin on a codon that is not the start codon. Translation of the polynucleotide can begin on the codon following the removed start codon or on a downstream start codon or an alternative start codon. In a non-limiting example, the start codon ATG or AUG is removed as the first 3 nucleotides of the polynucleotide sequence in order to have translation initiate on a downstream start codon or alternative start codon. The polynucleotide sequence where the

start codon was removed can further comprise at least one masking agent for the downstream start codon and/or alternative start codons in order to control or attempt to control the initiation of translation, the length of the polynucleotide and/or the structure of the polynucleotide.

17. Stop Codon Region

[0579] The invention also includes a polynucleotide that comprises both a stop codon region and the polynucleotide described herein (e.g., a polynucleotide comprising a nucleotide sequence encoding an ACADV L polypeptide). In some embodiments, the polynucleotides of the present invention can include at least two stop codons before the 3' untranslated region (UTR). The stop codon can be selected from TGA, TAA and TAG in the case of DNA, or from UGA, UAA and UAG in the case of RNA. In some embodiments, the polynucleotides of the present invention include the stop codon TGA in the case of DNA, or the stop codon UGA in the case of RNA, and one additional stop codon. In a further embodiment the additional stop codon can be TAA or UAA. In another embodiment, the polynucleotides of the present invention include three consecutive stop codons, four stop codons, or more.

18. Insertions and Substitutions

[0580] The invention also includes a polynucleotide of the present disclosure that further comprises insertions and/or substitutions.

[0581] In some embodiments, the 5'UTR of the polynucleotide can be replaced by the insertion of at least one region and/or string of nucleosides of the same base. The region and/or string of nucleotides can include, but is not limited to, at least 3, at least 4, at least 5, at least 6, at least 7 or at least 8 nucleotides and the nucleotides can be natural and/or unnatural. As a non-limiting example, the group of nucleotides can include 5-8 adenine, cytosine, thymine, a string of any of the other nucleotides disclosed herein and/or combinations thereof.

[0582] In some embodiments, the 5'UTR of the polynucleotide can be replaced by the insertion of at least two regions and/or strings of nucleotides of two different bases such as, but not limited to, adenine, cytosine, thymine, any of the other nucleotides disclosed herein and/or combinations thereof. For example, the 5'UTR can be replaced by inserting 5-8 adenine bases followed by the insertion of 5-8 cytosine bases. In another example, the

5'UTR can be replaced by inserting 5-8 cytosine bases followed by the insertion of 5-8 adenine bases.

[0583] In some embodiments, the polynucleotide can include at least one substitution and/or insertion downstream of the transcription start site that can be recognized by an RNA polymerase. As a non-limiting example, at least one substitution and/or insertion can occur downstream of the transcription start site by substituting at least one nucleic acid in the region just downstream of the transcription start site (such as, but not limited to, +1 to +6). Changes to region of nucleotides just downstream of the transcription start site can affect initiation rates, increase apparent nucleotide triphosphate (NTP) reaction constant values, and increase the dissociation of short transcripts from the transcription complex curing initial transcription (Briebe et al, *Biochemistry* (2002) 41: 5144-5149; herein incorporated by reference in its entirety). The modification, substitution and/or insertion of at least one nucleoside can cause a silent mutation of the sequence or can cause a mutation in the amino acid sequence.

[0584] In some embodiments, the polynucleotide can include the substitution of at least 1, at least 2, at least 3, at least 4, at least 5, at least 6, at least 7, at least 8, at least 9, at least 10, at least 11, at least 12 or at least 13 guanine bases downstream of the transcription start site.

[0585] In some embodiments, the polynucleotide can include the substitution of at least 1, at least 2, at least 3, at least 4, at least 5 or at least 6 guanine bases in the region just downstream of the transcription start site. As a non-limiting example, if the nucleotides in the region are GGGAGA, the guanine bases can be substituted by at least 1, at least 2, at least 3 or at least 4 adenine nucleotides. In another non-limiting example, if the nucleotides in the region are GGGAGA the guanine bases can be substituted by at least 1, at least 2, at least 3 or at least 4 cytosine bases. In another non-limiting example, if the nucleotides in the region are GGGAGA the guanine bases can be substituted by at least 1, at least 2, at least 3 or at least 4 thymine, and/or any of the nucleotides described herein.

[0586] In some embodiments, the polynucleotide can include at least one substitution and/or insertion upstream of the start codon. For the purpose of clarity, one of skill in the art would appreciate that the start codon is the first codon of the protein coding region whereas the transcription start site is the site where transcription begins. The polynucleotide can include, but is not limited to, at least 1, at least 2, at least 3, at least 4, at least 5, at least 6, at least 7 or at least 8 substitutions and/or insertions of nucleotide

bases. The nucleotide bases can be inserted or substituted at 1, at least 1, at least 2, at least 3, at least 4 or at least 5 locations upstream of the start codon. The nucleotides inserted and/or substituted can be the same base (*e.g.*, all A or all C or all T or all G), two different bases (*e.g.*, A and C, A and T, or C and T), three different bases (*e.g.*, A, C and T or A, C and T) or at least four different bases.

[0587] As a non-limiting example, the guanine base upstream of the coding region in the polynucleotide can be substituted with adenine, cytosine, thymine, or any of the nucleotides described herein. In another non-limiting example the substitution of guanine bases in the polynucleotide can be designed so as to leave one guanine base in the region downstream of the transcription start site and before the start codon (see Esvelt *et al.* Nature (2011) 472(7344):499-503; the contents of which is herein incorporated by reference in its entirety). As a non-limiting example, at least 5 nucleotides can be inserted at 1 location downstream of the transcription start site but upstream of the start codon and the at least 5 nucleotides can be the same base type.

19. Polynucleotide Comprising an mRNA Encoding an ACADVL Polypeptide

[0588] In certain embodiments, a polynucleotide of the present disclosure, for example a polynucleotide comprising an mRNA nucleotide sequence encoding an ACADVL polypeptide, comprises from 5' to 3' end:

- (i) a 5' cap provided above;
- (ii) a 5' UTR, such as the sequences provided above;
- (iii) an open reading frame encoding an ACADVL polypeptide, *e.g.*, a sequence optimized nucleic acid sequence encoding ACADVL disclosed herein;
- (iv) at least one stop codon;
- (v) a 3' UTR, such as the sequences provided above; and
- (vi) a poly-A tail provided above.

[0589] In some embodiments, the polynucleotide further comprises a miRNA binding site, *e.g.*, a miRNA binding site that binds to miRNA-142. In some embodiments, the 5'UTR comprises the miRNA binding site.

[0590] In some embodiments, a polynucleotide of the present disclosure comprises a nucleotide sequence encoding a polypeptide sequence at least 70%, at least 80%, at least 81%, at least 82%, at least 83%, at least 84%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least

94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identical to the protein sequence of a wild type ACADVL (e.g., isoform 1, 2, 3, or 4).

[0591] In some embodiments, a polynucleotide of the present disclosure, for example a polynucleotide comprising an mRNA nucleotide sequence encoding an ACADVL polypeptide, comprises (1) a 5' cap provided above, for example, CAP1, (2) a sequence optimized nucleic acid sequence encoding ACADVL disclosed herein (e.g., a nucleotide sequence selected from the group consisting of SEQ ID NO: 7 to 31), and (3) a poly-A tail provided above, for example, a poly A tail of approximately 100 residues.

20. Methods of Making Polynucleotides

[0592] The present disclosure also provides methods for making a polynucleotide of the invention (e.g., a polynucleotide comprising a nucleotide sequence encoding an ACADVL polypeptide) or a complement thereof.

[0593] In some aspects, a polynucleotide (e.g., a RNA, e.g., an mRNA) disclosed herein, and encoding an ACADVL polypeptide, can be constructed using *in vitro* transcription. In other aspects, a polynucleotide (e.g., a RNA, e.g., an mRNA) disclosed herein, and encoding an ACADVL polypeptide, can be constructed by chemical synthesis using an oligonucleotide synthesizer.

[0594] In other aspects, a polynucleotide (e.g., a RNA, e.g., an mRNA) disclosed herein, and encoding an ACADVL polypeptide is made by using a host cell. In certain aspects, a polynucleotide (e.g., a RNA, e.g., an mRNA) disclosed herein, and encoding an ACADVL polypeptide is made by one or more combination of the IVT, chemical synthesis, host cell expression, or any other methods known in the art.

[0595] Naturally occurring nucleosides, non-naturally occurring nucleosides, or combinations thereof, can totally or partially naturally replace occurring nucleosides present in the candidate nucleotide sequence and can be incorporated into a sequence-optimized nucleotide sequence (e.g., a RNA, e.g., an mRNA) encoding an ACADVL polypeptide. The resultant polynucleotides, e.g., mRNAs, can then be examined for their ability to produce protein and/or produce a therapeutic outcome.

a. In Vitro Transcription / Enzymatic Synthesis

[0596] The polynucleotides of the present invention disclosed herein (e.g., a polynucleotide comprising a nucleotide sequence encoding an ACADVL polypeptide)

can be transcribed using an *in vitro* transcription (IVT) system. The system typically comprises a transcription buffer, nucleotide triphosphates (NTPs), an RNase inhibitor and a polymerase. The NTPs can be selected from, but are not limited to, those described herein including natural and unnatural (modified) NTPs. The polymerase can be selected from, but is not limited to, T7 RNA polymerase, T3 RNA polymerase and mutant polymerases such as, but not limited to, polymerases able to incorporate polynucleotides disclosed herein. See U.S. Publ. No. US20130259923, which is herein incorporated by reference in its entirety.

[0597] Any number of RNA polymerases or variants can be used in the synthesis of the polynucleotides of the present invention. RNA polymerases can be modified by inserting or deleting amino acids of the RNA polymerase sequence. As a non-limiting example, the RNA polymerase can be modified to exhibit an increased ability to incorporate a 2'-modified nucleotide triphosphate compared to an unmodified RNA polymerase (see International Publication WO2008078180 and U.S. Patent 8,101,385; herein incorporated by reference in their entireties).

[0598] Variants can be obtained by evolving an RNA polymerase, optimizing the RNA polymerase amino acid and/or nucleic acid sequence and/or by using other methods known in the art. As a non-limiting example, T7 RNA polymerase variants can be evolved using the continuous directed evolution system set out by Esvelt et al. (Nature 472:499-503 (2011); herein incorporated by reference in its entirety) where clones of T7 RNA polymerase can encode at least one mutation such as, but not limited to, lysine at position 93 substituted for threonine (K93T), I4M, A7T, E63V, V64D, A65E, D66Y, T76N, C125R, S128R, A136T, N165S, G175R, H176L, Y178H, F182L, L196F, G198V, D208Y, E222K, S228A, Q239R, T243N, G259D, M267I, G280C, H300R, D351A, A354S, E356D, L360P, A383V, Y385C, D388Y, S397R, M401T, N410S, K450R, P451T, G452V, E484A, H523L, H524N, G542V, E565K, K577E, K577M, N601S, S684Y, L699I, K713E, N748D, Q754R, E775K, A827V, D851N or L864F. As another non-limiting example, T7 RNA polymerase variants can encode at least mutation as described in U.S. Pub. Nos. 20100120024 and 20070117112; herein incorporated by reference in their entireties. Variants of RNA polymerase can also include, but are not limited to, substitutional variants, conservative amino acid substitution, insertional variants, and/or deletional variants.

- [0599] In one aspect, the polynucleotide can be designed to be recognized by the wild type or variant RNA polymerases. In doing so, the polynucleotide can be modified to contain sites or regions of sequence changes from the wild type or parent chimeric polynucleotide.
- [0600] Polynucleotide or nucleic acid synthesis reactions can be carried out by enzymatic methods utilizing polymerases. Polymerases catalyze the creation of phosphodiester bonds between nucleotides in a polynucleotide or nucleic acid chain. Currently known DNA polymerases can be divided into different families based on amino acid sequence comparison and crystal structure analysis. DNA polymerase I (pol I) or A polymerase family, including the Klenow fragments of *E. coli*, *Bacillus* DNA polymerase I, *Thermus aquaticus* (Taq) DNA polymerases, and the T7 RNA and DNA polymerases, is among the best studied of these families. Another large family is DNA polymerase α (pol α) or B polymerase family, including all eukaryotic replicating DNA polymerases and polymerases from phages T4 and RB69. Although they employ similar catalytic mechanism, these families of polymerases differ in substrate specificity, substrate analog-incorporating efficiency, degree and rate for primer extension, mode of DNA synthesis, exonuclease activity, and sensitivity against inhibitors.
- [0601] DNA polymerases are also selected based on the optimum reaction conditions they require, such as reaction temperature, pH, and template and primer concentrations. Sometimes a combination of more than one DNA polymerases is employed to achieve the desired DNA fragment size and synthesis efficiency. For example, Cheng et al. increase pH, add glycerol and dimethyl sulfoxide, decrease denaturation times, increase extension times, and utilize a secondary thermostable DNA polymerase that possesses a 3' to 5' exonuclease activity to effectively amplify long targets from cloned inserts and human genomic DNA. (Cheng et al., PNAS 91:5695-5699 (1994), the contents of which are incorporated herein by reference in their entirety). RNA polymerases from bacteriophage T3, T7, and SP6 have been widely used to prepare RNAs for biochemical and biophysical studies. RNA polymerases, capping enzymes, and poly-A polymerases are disclosed in the co-pending International Publication No. WO2014028429, the contents of which are incorporated herein by reference in their entirety.
- [0602] In one aspect, the RNA polymerase which can be used in the synthesis of the polynucleotides of the present invention is a Syn5 RNA polymerase. (see Zhu et al. Nucleic Acids Research 2013, doi:10.1093/nar/gkt1193, which is herein incorporated by

reference in its entirety). The Syn5 RNA polymerase was recently characterized from marine cyanophage Syn5 by Zhu et al. where they also identified the promoter sequence (see Zhu et al. Nucleic Acids Research 2013, the contents of which is herein incorporated by reference in its entirety). Zhu et al. found that Syn5 RNA polymerase catalyzed RNA synthesis over a wider range of temperatures and salinity as compared to T7 RNA polymerase. Additionally, the requirement for the initiating nucleotide at the promoter was found to be less stringent for Syn5 RNA polymerase as compared to the T7 RNA polymerase making Syn5 RNA polymerase promising for RNA synthesis.

- [0603]** In one aspect, a Syn5 RNA polymerase can be used in the synthesis of the polynucleotides described herein. As a non-limiting example, a Syn5 RNA polymerase can be used in the synthesis of the polynucleotide requiring a precise 3'-terminus.
- [0604]** In one aspect, a Syn5 promoter can be used in the synthesis of the polynucleotides. As a non-limiting example, the Syn5 promoter can be 5'-ATTGGGCACCCGTAAGGG-3' (SEQ ID NO: 37) as described by Zhu et al. (Nucleic Acids Research 2013).
- [0605]** In one aspect, a Syn5 RNA polymerase can be used in the synthesis of polynucleotides comprising at least one chemical modification described herein and/or known in the art (see e.g., the incorporation of pseudo-UTP and 5Me-CTP described in Zhu et al. Nucleic Acids Research 2013).
- [0606]** In one aspect, the polynucleotides described herein can be synthesized using a Syn5 RNA polymerase which has been purified using modified and improved purification procedure described by Zhu et al. (Nucleic Acids Research 2013).
- [0607]** Various tools in genetic engineering are based on the enzymatic amplification of a target gene which acts as a template. For the study of sequences of individual genes or specific regions of interest and other research needs, it is necessary to generate multiple copies of a target gene from a small sample of polynucleotides or nucleic acids. Such methods can be applied in the manufacture of the polynucleotides of the invention.
- [0608]** For example, polymerase chain reaction (PCR), strand displacement amplification (SDA), nucleic acid sequence-based amplification (NASBA), also called transcription mediated amplification (TMA), and/or rolling-circle amplification (RCA) can be utilized in the manufacture of one or more regions of the polynucleotides of the present invention.
- [0609]** Assembling polynucleotides or nucleic acids by a ligase is also widely used.

b. Chemical synthesis

[0610] Standard methods can be applied to synthesize an isolated polynucleotide sequence encoding an isolated polypeptide of interest, such as a polynucleotide of the invention (e.g., a polynucleotide comprising a nucleotide sequence encoding an ACADVL polypeptide). For example, a single DNA or RNA oligomer containing a codon-optimized nucleotide sequence coding for the particular isolated polypeptide can be synthesized. In other aspects, several small oligonucleotides coding for portions of the desired polypeptide can be synthesized and then ligated. In some aspects, the individual oligonucleotides typically contain 5' or 3' overhangs for complementary assembly.

[0611] A polynucleotide disclosed herein (e.g., a RNA, e.g., an mRNA) can be chemically synthesized using chemical synthesis methods and potential nucleobase substitutions known in the art. See, for example, International Publication Nos. WO2014093924, WO2013052523; WO2013039857, WO2012135805, WO2013151671; U.S. Publ. No. US20130115272; or U.S. Pat. Nos. US8999380 or US8710200, all of which are herein incorporated by reference in their entireties.

c. Purification of Polynucleotides Encoding ACADVL

[0612] Purification of the polynucleotides described herein (e.g., a polynucleotide comprising a nucleotide sequence encoding an ACADVL polypeptide) can include, but is not limited to, polynucleotide clean-up, quality assurance and quality control. Clean-up can be performed by methods known in the arts such as, but not limited to, AGENCOURT® beads (Beckman Coulter Genomics, Danvers, MA), poly-T beads, LNATM oligo-T capture probes (EXIQON® Inc., Vedbaek, Denmark) or HPLC based purification methods such as, but not limited to, strong anion exchange HPLC, weak anion exchange HPLC, reverse phase HPLC (RP-HPLC), and hydrophobic interaction HPLC (HIC-HPLC).

[0613] The term "purified" when used in relation to a polynucleotide such as a "purified polynucleotide" refers to one that is separated from at least one contaminant. As used herein, a "contaminant" is any substance that makes another unfit, impure or inferior. Thus, a purified polynucleotide (e.g., DNA and RNA) is present in a form or setting different from that in which it is found in nature, or a form or setting different from that which existed prior to subjecting it to a treatment or purification method.

- [0614] In some embodiments, purification of a polynucleotide of the invention (e.g., a polynucleotide comprising a nucleotide sequence encoding an ACADVL polypeptide) removes impurities that can reduce or remove an unwanted immune response, *e.g.*, reducing cytokine activity.
- [0615] In some embodiments, the polynucleotide of the invention (e.g., a polynucleotide comprising a nucleotide sequence encoding an ACADVL polypeptide) is purified prior to administration using column chromatography (*e.g.*, strong anion exchange HPLC, weak anion exchange HPLC, reverse phase HPLC (RP-HPLC), and hydrophobic interaction HPLC (HIC-HPLC), or (LCMS)).
- [0616] In some embodiments, the polynucleotide of the invention (e.g., a polynucleotide comprising a nucleotide sequence an ACADVL polypeptide) purified using column chromatography (*e.g.*, strong anion exchange HPLC, weak anion exchange HPLC, reverse phase HPLC (RP-HPLC, hydrophobic interaction HPLC (HIC-HPLC), or (LCMS)) presents increased expression of the encoded ACADVL protein compared to the expression level obtained with the same polynucleotide of the present disclosure purified by a different purification method.
- [0617] In some embodiments, a column chromatography (*e.g.*, strong anion exchange HPLC, weak anion exchange HPLC, reverse phase HPLC (RP-HPLC), hydrophobic interaction HPLC (HIC-HPLC), or (LCMS)) purified polynucleotide comprises a nucleotide sequence encoding an ACADVL polypeptide comprising one or more of the point mutations known in the art.
- [0618] In some embodiments, the use of RP-HPLC purified polynucleotide increases ACADVL protein expression levels in cells when introduced into those cells, *e.g.*, by 10-100%, i.e., at least about 10%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 90%, at least about 95%, or at least about 100% with respect to the expression levels of ACADVL protein in the cells before the RP-HPLC purified polynucleotide was introduced in the cells, or after a non-RP-HPLC purified polynucleotide was introduced in the cells.
- [0619] In some embodiments, the use of RP-HPLC purified polynucleotide increases functional ACADVL protein expression levels in cells when introduced into those cells, *e.g.*, by 10-100%, i.e., at least about 10%, at least about 20%, at least about 25%, at least

about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 90%, at least about 95%, or at least about 100% with respect to the functional expression levels of ACADVL protein in the cells before the RP-HPLC purified polynucleotide was introduced in the cells, or after a non-RP-HPLC purified polynucleotide was introduced in the cells.

[0620] In some embodiments, the use of RP-HPLC purified polynucleotide increases detectable ACADVL activity in cells when introduced into those cells, *e.g.*, by 10-100%, *i.e.*, at least about 10%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 90%, at least about 95%, or at least about 100% with respect to the activity levels of functional ACADVL in the cells before the RP-HPLC purified polynucleotide was introduced in the cells, or after a non-RP-HPLC purified polynucleotide was introduced in the cells.

[0621] In some embodiments, the purified polynucleotide is at least about 80% pure, at least about 85% pure, at least about 90% pure, at least about 95% pure, at least about 96% pure, at least about 97% pure, at least about 98% pure, at least about 99% pure, or about 100% pure.

[0622] A quality assurance and/or quality control check can be conducted using methods such as, but not limited to, gel electrophoresis, UV absorbance, or analytical HPLC. In another embodiment, the polynucleotide can be sequenced by methods including, but not limited to reverse-transcriptase-PCR.

d. Quantification of Expressed Polynucleotides Encoding ACADVL

[0623] In some embodiments, the polynucleotides of the present invention (*e.g.*, a polynucleotide comprising a nucleotide sequence encoding an ACADVL polypeptide), their expression products, as well as degradation products and metabolites can be quantified according to methods known in the art.

[0624] In some embodiments, the polynucleotides of the present invention can be quantified in exosomes or when derived from one or more bodily fluid. As used herein "bodily fluids" include peripheral blood, serum, plasma, ascites, urine, cerebrospinal fluid (CSF), sputum, saliva, bone marrow, synovial fluid, aqueous humor, amniotic fluid,

cerumen, breast milk, bronchoalveolar lavage fluid, semen, prostatic fluid, cowper's fluid or pre-ejaculatory fluid, sweat, fecal matter, hair, tears, cyst fluid, pleural and peritoneal fluid, pericardial fluid, lymph, chyme, chyle, bile, interstitial fluid, menses, pus, sebum, vomit, vaginal secretions, mucosal secretion, stool water, pancreatic juice, lavage fluids from sinus cavities, bronchopulmonary aspirates, blastocyl cavity fluid, and umbilical cord blood. Alternatively, exosomes can be retrieved from an organ selected from the group consisting of lung, heart, pancreas, stomach, intestine, bladder, kidney, ovary, testis, skin, colon, breast, prostate, brain, esophagus, liver, and placenta.

[0625] In the exosome quantification method, a sample of not more than 2mL is obtained from the subject and the exosomes isolated by size exclusion chromatography, density gradient centrifugation, differential centrifugation, nanomembrane ultrafiltration, immunoabsorbent capture, affinity purification, microfluidic separation, or combinations thereof. In the analysis, the level or concentration of a polynucleotide can be an expression level, presence, absence, truncation or alteration of the administered construct. It is advantageous to correlate the level with one or more clinical phenotypes or with an assay for a human disease biomarker.

[0626] The assay can be performed using construct specific probes, cytometry, qRT-PCR, real-time PCR, PCR, flow cytometry, electrophoresis, mass spectrometry, or combinations thereof while the exosomes can be isolated using immunohistochemical methods such as enzyme linked immunosorbent assay (ELISA) methods. Exosomes can also be isolated by size exclusion chromatography, density gradient centrifugation, differential centrifugation, nanomembrane ultrafiltration, immunoabsorbent capture, affinity purification, microfluidic separation, or combinations thereof.

[0627] These methods afford the investigator the ability to monitor, in real time, the level of polynucleotides remaining or delivered. This is possible because the polynucleotides of the present invention differ from the endogenous forms due to the structural or chemical modifications.

[0628] In some embodiments, the polynucleotide can be quantified using methods such as, but not limited to, ultraviolet visible spectroscopy (UV/Vis). A non-limiting example of a UV/Vis spectrometer is a NANODROP® spectrometer (ThermoFisher, Waltham, MA). The quantified polynucleotide can be analyzed in order to determine if the polynucleotide can be of proper size, check that no degradation of the polynucleotide has occurred. Degradation of the polynucleotide can be checked by methods such as, but not

limited to, agarose gel electrophoresis, HPLC based purification methods such as, but not limited to, strong anion exchange HPLC, weak anion exchange HPLC, reverse phase HPLC (RP-HPLC), and hydrophobic interaction HPLC (HIC-HPLC), liquid chromatography-mass spectrometry (LCMS), capillary electrophoresis (CE) and capillary gel electrophoresis (CGE).

21. Pharmaceutical Compositions and Formulations

[0629] The present invention provides pharmaceutical compositions and formulations that comprise any of the polynucleotides described above. In some embodiments, the composition or formulation further comprises a delivery agent.

[0630] In some embodiments, the composition or formulation can contain a polynucleotide comprising a sequence optimized nucleic acid sequence disclosed herein which encodes an ACADVL polypeptide. In some embodiments, the composition or formulation can contain a polynucleotide (e.g., a RNA, e.g., an mRNA) comprising a polynucleotide (e.g., an ORF) having significant sequence identity to a sequence optimized nucleic acid sequence disclosed herein which encodes an ACADVL polypeptide. In some embodiments, the polynucleotide further comprises a miRNA binding site, e.g., a miRNA binding site that binds miR-126, miR-142, miR-144, miR-146, miR-150, miR-155, miR-16, miR-21, miR-223, miR-24, miR-27 and miR-26a.

[0631] Pharmaceutical compositions or formulation can optionally comprise one or more additional active substances, e.g., therapeutically and/or prophylactically active substances. Pharmaceutical compositions or formulation of the present invention can be sterile and/or pyrogen-free. General considerations in the formulation and/or manufacture of pharmaceutical agents can be found, for example, in *Remington: The Science and Practice of Pharmacy* 21st ed., Lippincott Williams & Wilkins, 2005 (incorporated herein by reference in its entirety). In some embodiments, compositions are administered to humans, human patients or subjects. For the purposes of the present disclosure, the phrase "active ingredient" generally refers to polynucleotides to be delivered as described herein.

[0632] Formulations and pharmaceutical compositions described herein can be prepared by any method known or hereafter developed in the art of pharmacology. In general, such preparatory methods include the step of associating the active ingredient with an excipient and/or one or more other accessory ingredients, and then, if necessary and/or desirable, dividing, shaping and/or packaging the product into a desired single- or multi-dose unit.

- [0633] A pharmaceutical composition or formulation in accordance with the present disclosure can be prepared, packaged, and/or sold in bulk, as a single unit dose, and/or as a plurality of single unit doses. As used herein, a "unit dose" refers to a discrete amount of the pharmaceutical composition comprising a predetermined amount of the active ingredient. The amount of the active ingredient is generally equal to the dosage of the active ingredient that would be administered to a subject and/or a convenient fraction of such a dosage such as, for example, one-half or one-third of such a dosage.
- [0634] Relative amounts of the active ingredient, the pharmaceutically acceptable excipient, and/or any additional ingredients in a pharmaceutical composition in accordance with the present disclosure can vary, depending upon the identity, size, and/or condition of the subject being treated and further depending upon the route by which the composition is to be administered.
- [0635] In some embodiments, the compositions and formulations described herein can contain at least one polynucleotide of the invention. As a non-limiting example, the composition or formulation can contain 1, 2, 3, 4 or 5 polynucleotides of the invention. In some embodiments, the compositions or formulations described herein can comprise more than one type of polynucleotide. In some embodiments, the composition or formulation can comprise a polynucleotide in linear and circular form. In another embodiment, the composition or formulation can comprise a circular polynucleotide and an in vitro transcribed (IVT) polynucleotide. In yet another embodiment, the composition or formulation can comprise an IVT polynucleotide, a chimeric polynucleotide and a circular polynucleotide.
- [0636] Although the descriptions of pharmaceutical compositions and formulations provided herein are principally directed to pharmaceutical compositions and formulations that are suitable for administration to humans, it will be understood by the skilled artisan that such compositions are generally suitable for administration to any other animal, *e.g.*, to non-human animals, *e.g.* non-human mammals.
- [0637] The present invention provides pharmaceutical formulations that comprise a polynucleotide described herein (*e.g.*, a polynucleotide comprising a nucleotide sequence encoding an ACADVL polypeptide). The polynucleotides described herein can be formulated using one or more excipients to: (1) increase stability; (2) increase cell transfection; (3) permit the sustained or delayed release (*e.g.*, from a depot formulation of the polynucleotide); (4) alter the biodistribution (*e.g.*, target the polynucleotide to specific

tissues or cell types); (5) increase the translation of encoded protein *in vivo*; and/or (6) alter the release profile of encoded protein *in vivo*. In some embodiments, the pharmaceutical formulation further comprises a delivery agent comprising, e.g., a compound having the Formula (I), e.g., any of Compounds 1-232, e.g., Compound 18; a compound having the Formula (III), (IV), (V), or (VI), e.g., any of Compounds 233-342, e.g., Compound 236; or a compound having the Formula (VIII), e.g., any of Compounds 419-428, e.g., Compound 428, or any combination thereof. In some embodiments, the delivery agent comprises Compound 18, DSPC, Cholesterol, and Compound 428, e.g., with a mole ratio of about 50:10:38.5:1.5.

[0638] A pharmaceutically acceptable excipient, as used herein, includes, but are not limited to, any and all solvents, dispersion media, or other liquid vehicles, dispersion or suspension aids, diluents, granulating and/or dispersing agents, surface active agents, isotonic agents, thickening or emulsifying agents, preservatives, binders, lubricants or oil, coloring, sweetening or flavoring agents, stabilizers, antioxidants, antimicrobial or antifungal agents, osmolality adjusting agents, pH adjusting agents, buffers, chelants, cyoprotectants, and/or bulking agents, as suited to the particular dosage form desired. Various excipients for formulating pharmaceutical compositions and techniques for preparing the composition are known in the art (see Remington: The Science and Practice of Pharmacy, 21st Edition, A. R. Gennaro (Lippincott, Williams & Wilkins, Baltimore, MD, 2006; incorporated herein by reference in its entirety).

[0639] Exemplary diluents include, but are not limited to, calcium or sodium carbonate, calcium phosphate, calcium hydrogen phosphate, sodium phosphate, lactose, sucrose, cellulose, microcrystalline cellulose, kaolin, mannitol, sorbitol, etc., and/or combinations thereof.

[0640] Exemplary granulating and/or dispersing agents include, but are not limited to, starches, pregelatinized starches, or microcrystalline starch, alginic acid, guar gum, agar, poly(vinyl-pyrrolidone), (providone), cross-linked poly(vinyl-pyrrolidone) (crospovidone), cellulose, methylcellulose, carboxymethyl cellulose, cross-linked sodium carboxymethyl cellulose (croscarmellose), magnesium aluminum silicate (VEEGUM®), sodium lauryl sulfate, etc., and/or combinations thereof.

[0641] Exemplary surface active agents and/or emulsifiers include, but are not limited to, natural emulsifiers (e.g., acacia, agar, alginic acid, sodium alginate, tragacanth, chondrux, cholesterol, xanthan, pectin, gelatin, egg yolk, casein, wool fat, cholesterol, wax, and

lecithin), sorbitan fatty acid esters (e.g., polyoxyethylene sorbitan monooleate [TWEEN®80], sorbitan monopalmitate [SPAN®40], glyceryl monooleate, polyoxyethylene esters, polyethylene glycol fatty acid esters (e.g., CREMOPHOR®), polyoxyethylene ethers (e.g., polyoxyethylene lauryl ether [BRIJ®30]), PLUORINC®F 68, POLOXAMER®188, etc. and/or combinations thereof.

[0642] Exemplary binding agents include, but are not limited to, starch, gelatin, sugars (e.g., sucrose, glucose, dextrose, dextrin, molasses, lactose, lactitol, mannitol), amino acids (e.g., glycine), natural and synthetic gums (e.g., acacia, sodium alginate), ethylcellulose, hydroxyethylcellulose, hydroxypropyl methylcellulose, etc., and combinations thereof.

[0643] Oxidation is a potential degradation pathway for mRNA, especially for liquid mRNA formulations. In order to prevent oxidation, antioxidants can be added to the formulations. Exemplary antioxidants include, but are not limited to, alpha tocopherol, ascorbic acid, acorbyl palmitate, benzyl alcohol, butylated hydroxyanisole, m-cresol, methionine, butylated hydroxytoluene, monothioglycerol, sodium or potassium metabisulfite, propionic acid, propyl gallate, sodium ascorbate, etc., and combinations thereof.

[0644] Exemplary chelating agents include, but are not limited to, ethylenediaminetetraacetic acid (EDTA), citric acid monohydrate, disodium edetate, fumaric acid, malic acid, phosphoric acid, sodium edetate, tartaric acid, trisodium edetate, etc., and combinations thereof.

[0645] Exemplary antimicrobial or antifungal agents include, but are not limited to, benzalkonium chloride, benzethonium chloride, methyl paraben, ethyl paraben, propyl paraben, butyl paraben, benzoic acid, hydroxybenzoic acid, potassium or sodium benzoate, potassium or sodium sorbate, sodium propionate, sorbic acid, etc., and combinations thereof.

[0646] Exemplary preservatives include, but are not limited to, vitamin A, vitamin C, vitamin E, beta-carotene, citric acid, ascorbic acid, butylated hydroxyanisol, ethylenediamine, sodium lauryl sulfate (SLS), sodium lauryl ether sulfate (SLES), etc., and combinations thereof.

[0647] In some embodiments, the pH of polynucleotide solutions are maintained between pH 5 and pH 8 to improve stability. Exemplary buffers to control pH can include, but are

not limited to sodium phosphate, sodium citrate, sodium succinate, histidine (or histidine-HCl), sodium malate, sodium carbonate, etc., and/or combinations thereof.

[0648] Exemplary lubricating agents include, but are not limited to, magnesium stearate, calcium stearate, stearic acid, silica, talc, malt, hydrogenated vegetable oils, polyethylene glycol, sodium benzoate, sodium or magnesium lauryl sulfate, etc., and combinations thereof.

[0649] The pharmaceutical composition or formulation described here can contain a cryoprotectant to stabilize a polynucleotide described herein during freezing. Exemplary cryoprotectants include, but are not limited to mannitol, sucrose, trehalose, lactose, glycerol, dextrose, etc., and combinations thereof.

[0650] The pharmaceutical composition or formulation described here can contain a bulking agent in lyophilized polynucleotide formulations to yield a "pharmaceutically elegant" cake, stabilize the lyophilized polynucleotides during long term (e.g., 36 month) storage. Exemplary bulking agents of the present invention can include, but are not limited to sucrose, trehalose, mannitol, glycine, lactose, raffinose, and combinations thereof.

[0651] In some embodiments, the pharmaceutical composition or formulation further comprises a delivery agent. The delivery agent of the present disclosure can include, without limitation, liposomes, lipid nanoparticles, lipidoids, polymers, lipoplexes, microvesicles, exosomes, peptides, proteins, cells transfected with polynucleotides, hyaluronidase, nanoparticle mimics, nanotubes, conjugates, and combinations thereof.

22. Delivery Agents

a. Lipid Compound


[0652] The present disclosure provides pharmaceutical compositions with advantageous properties. The lipid compositions described herein may be advantageously used in lipid nanoparticle compositions for the delivery of therapeutic and/or prophylactic agents, e.g., mRNAs, to mammalian cells or organs. For example, the lipids described herein have little or no immunogenicity. For example, the lipid compounds disclosed herein have a lower immunogenicity as compared to a reference lipid (e.g., MC3, KC2, or DLinDMA). For example, a formulation comprising a lipid disclosed herein and a therapeutic or prophylactic agent, e.g., mRNA, has an increased therapeutic index as compared to a

corresponding formulation which comprises a reference lipid (*e.g.*, MC3, KC2, or DLinDMA) and the same therapeutic or prophylactic agent.

[0653] In certain embodiments, the present application provides pharmaceutical compositions comprising:

- (a) a polynucleotide comprising a nucleotide sequence encoding an ACADVL polypeptide; and
- (b) a delivery agent.

[0654] In some embodiments, the delivery agent comprises a lipid compound having the Formula (I)



(I),

wherein

R₁ is selected from the group consisting of C₅₋₃₀ alkyl, C₅₋₂₀ alkenyl, -R*YR'', -YR'', and -R''M'R';

R₂ and R₃ are independently selected from the group consisting of H, C₁₋₁₄ alkyl, C₂₋₁₄ alkenyl, -R*YR'', -YR'', and -R*OR'', or R₂ and R₃, together with the atom to which they are attached, form a heterocycle or carbocycle;

R₄ is selected from the group consisting of a C₃₋₆ carbocycle, -(CH₂)_nQ, -(CH₂)_nCHQR, -CHQR, -CQ(R)₂, and unsubstituted C₁₋₆ alkyl, where Q is selected from a carbocycle, heterocycle, -OR, -O(CH₂)_nN(R)₂, -C(O)OR, -OC(O)R, -CX₃, -CX₂H, -CXH₂, -CN, -N(R)₂, -C(O)N(R)₂, -N(R)C(O)R, -N(R)S(O)₂R, -N(R)C(O)N(R)₂, -N(R)C(S)N(R)₂, -N(R)R₈, -O(CH₂)_nOR, -N(R)C(=NR₉)N(R)₂, -N(R)C(=CHR₉)N(R)₂, -OC(O)N(R)₂, -N(R)C(O)OR, -N(OR)C(O)R, -N(OR)S(O)₂R, -N(OR)C(O)OR, -N(OR)C(O)N(R)₂, -N(OR)C(S)N(R)₂, -N(OR)C(=NR₉)N(R)₂, -N(OR)C(=CHR₉)N(R)₂, -C(=NR₉)N(R)₂, -C(=NR₉)R, -C(O)N(R)OR, and -C(R)N(R)₂C(O)OR, and each n is independently selected from 1, 2, 3, 4, and 5;

each R₅ is independently selected from the group consisting of C₁₋₃ alkyl, C₂₋₃ alkenyl, and H;

each R₆ is independently selected from the group consisting of C₁₋₃ alkyl, C₂₋₃ alkenyl, and H;

M and M' are independently selected from -C(O)O-, -OC(O)-, -C(O)N(R')-, -N(R')C(O)-, -C(O)-, -C(S)-, -C(S)S-, -SC(S)-, -CH(OH)-, -P(O)(OR')O-, -S(O)₂-, -S-S-, an aryl group, and a heteroaryl group;

R₇ is selected from the group consisting of C₁₋₃ alkyl, C₂₋₃ alkenyl, and H;

R₈ is selected from the group consisting of C₃₋₆ carbocycle and heterocycle;

R₉ is selected from the group consisting of H, CN, NO₂, C₁₋₆ alkyl, -OR, -S(O)₂R, -S(O)₂N(R)₂, C₂₋₆ alkenyl, C₃₋₆ carbocycle and heterocycle;

each R is independently selected from the group consisting of C₁₋₃ alkyl, C₂₋₃ alkenyl, and H;

each R' is independently selected from the group consisting of C₁₋₁₈ alkyl, C₂₋₁₈ alkenyl, -R*YR'', -YR'', and H;

each R'' is independently selected from the group consisting of C₃₋₁₄ alkyl and C₃₋₁₄ alkenyl;

each R* is independently selected from the group consisting of C₁₋₁₂ alkyl and C₂₋₁₂ alkenyl;

each Y is independently a C₃₋₆ carbocycle;

each X is independently selected from the group consisting of F, Cl, Br, and I; and m is selected from 5, 6, 7, 8, 9, 10, 11, 12, and 13, or salts or stereoisomers thereof.

[0655] In some embodiments, a subset of compounds of Formula (I) includes those in which

R₁ is selected from the group consisting of C₅₋₂₀ alkyl, C₅₋₂₀ alkenyl, -R*YR'', -YR'', and -R''M'R';

R₂ and R₃ are independently selected from the group consisting of H, C₁₋₁₄ alkyl, C₂₋₁₄ alkenyl, -R*YR'', -YR'', and -R*OR'', or R₂ and R₃, together with the atom to which they are attached, form a heterocycle or carbocycle;

R₄ is selected from the group consisting of a C₃₋₆ carbocycle, -(CH₂)_nQ, -(CH₂)_nCHQR, -CHQR, -CQ(R)₂, and unsubstituted C₁₋₆ alkyl, where Q is selected from a carbocycle, heterocycle, -OR, -O(CH₂)_nN(R)₂, -C(O)OR, -OC(O)R, -CX₃, -CX₂H, -CXH₂, -CN, -N(R)₂, -C(O)N(R)₂, -N(R)C(O)R, -N(R)S(O)₂R, -N(R)C(O)N(R)₂, -N(R)C(S)N(R)₂, and -C(R)N(R)₂C(O)OR, and each n is independently selected from 1, 2, 3, 4, and 5;

each R₅ is independently selected from the group consisting of C₁₋₃ alkyl, C₂₋₃ alkenyl, and H;

each R₆ is independently selected from the group consisting of C₁₋₃ alkyl, C₂₋₃ alkenyl, and H;

M and M' are independently selected from -C(O)O-, -OC(O)-, -C(O)N(R')-, -N(R')C(O)-, -C(O)-, -C(S)-, -C(S)S-, -SC(S)-, -CH(OH)-, -P(O)(OR')O-, -S(O)₂-, an aryl group, and a heteroaryl group;

R₇ is selected from the group consisting of C₁₋₃ alkyl, C₂₋₃ alkenyl, and H;

each R is independently selected from the group consisting of C₁₋₃ alkyl, C₂₋₃ alkenyl, and H;

each R' is independently selected from the group consisting of C₁₋₁₈ alkyl, C₂₋₁₈ alkenyl, -R*YR'', -YR'', and H;

each R'' is independently selected from the group consisting of C₃₋₁₄ alkyl and C₃₋₁₄ alkenyl;

each R* is independently selected from the group consisting of C₁₋₁₂ alkyl and C₂₋₁₂ alkenyl;

each Y is independently a C₃₋₆ carbocycle;

each X is independently selected from the group consisting of F, Cl, Br, and I; and m is selected from 5, 6, 7, 8, 9, 10, 11, 12, and 13,

or salts or stereoisomers thereof, wherein alkyl and alkenyl groups may be linear or branched.

[0656] In some embodiments, a subset of compounds of Formula (I) includes those in which when R₄ is -(CH₂)_nQ, -(CH₂)_nCHQR, -CHQR, or -CQ(R)₂, then (i) Q is not -N(R)₂ when n is 1, 2, 3, 4 or 5, or (ii) Q is not 5, 6, or 7-membered heterocycloalkyl when n is 1 or 2.

[0657] In other embodiments, another subset of compounds of Formula (I) includes those in which

R₁ is selected from the group consisting of C₅₋₃₀ alkyl, C₅₋₂₀ alkenyl, -R*YR'', -YR'', and -R''M'R';

R₂ and R₃ are independently selected from the group consisting of H, C₁₋₁₄ alkyl, C₂₋₁₄ alkenyl, -R*YR'', -YR'', and -R*OR'', or R₂ and R₃, together with the atom to which they are attached, form a heterocycle or carbocycle;

R_4 is selected from the group consisting of a C_{3-6} carbocycle, $-(CH_2)_nQ$, $-(CH_2)_nCHQR$, $-CHQR$, $-CQ(R)_2$, and unsubstituted C_{1-6} alkyl, where Q is selected from a C_{3-6} carbocycle, a 5- to 14-membered heteroaryl having one or more heteroatoms selected from N, O, and S, $-OR$, $-O(CH_2)_nN(R)_2$, $-C(O)OR$, $-OC(O)R$, $-CX_3$, $-CX_2H$, $-CXH_2$, $-CN$, $-C(O)N(R)_2$, $-N(R)C(O)R$, $-N(R)S(O)_2R$, $-N(R)C(O)N(R)_2$, $-N(R)C(S)N(R)_2$, $-CRN(R)_2C(O)OR$, $-N(R)R_8$, $-O(CH_2)_nOR$, $-N(R)C(=NR_9)N(R)_2$, $-N(R)C(=CHR_9)N(R)_2$, $-OC(O)N(R)_2$, $-N(R)C(O)OR$, $-N(OR)C(O)R$, $-N(OR)S(O)_2R$, $-N(OR)C(O)OR$, $-N(OR)C(O)N(R)_2$, $-N(OR)C(S)N(R)_2$, $-N(OR)C(=NR_9)N(R)_2$, $-N(OR)C(=CHR_9)N(R)_2$, $-C(=NR_9)N(R)_2$, $-C(=NR_9)R$, $-C(O)N(R)OR$, and a 5- to 14-membered heterocycloalkyl having one or more heteroatoms selected from N, O, and S which is substituted with one or more substituents selected from oxo ($=O$), OH, amino, and C_{1-3} alkyl, and each n is independently selected from 1, 2, 3, 4, and 5;

each R_5 is independently selected from the group consisting of C_{1-3} alkyl, C_{2-3} alkenyl, and H;

each R_6 is independently selected from the group consisting of C_{1-3} alkyl, C_{2-3} alkenyl, and H;

M and M' are independently selected from $-C(O)O-$, $-OC(O)-$, $-C(O)N(R')-$, $-N(R')C(O)-$, $-C(O)-$, $-C(S)-$, $-C(S)S-$, $-SC(S)-$, $-CH(OH)-$, $-P(O)(OR')O-$, $-S(O)_2-$, $-S-S-$, an aryl group, and a heteroaryl group;

R_7 is selected from the group consisting of C_{1-3} alkyl, C_{2-3} alkenyl, and H;

R_8 is selected from the group consisting of C_{3-6} carbocycle and heterocycle;

R_9 is selected from the group consisting of H, CN, NO_2 , C_{1-6} alkyl, $-OR$, $-S(O)_2R$, $-S(O)_2N(R)_2$, C_{2-6} alkenyl, C_{3-6} carbocycle and heterocycle;

each R is independently selected from the group consisting of C_{1-3} alkyl, C_{2-3} alkenyl, and H;

each R' is independently selected from the group consisting of C_{1-18} alkyl, C_{2-18} alkenyl, $-R^*YR''$, $-YR''$, and H;

each R'' is independently selected from the group consisting of C_{3-14} alkyl and C_{3-14} alkenyl;

each R^* is independently selected from the group consisting of C_{1-12} alkyl and C_{2-12} alkenyl;

each Y is independently a C_{3-6} carbocycle;

each X is independently selected from the group consisting of F, Cl, Br, and I; and

m is selected from 5, 6, 7, 8, 9, 10, 11, 12, and 13,
or salts or stereoisomers thereof.

[0658] In other embodiments, another subset of compounds of Formula (I) includes those in which

R_1 is selected from the group consisting of C_{5-20} alkyl, C_{5-20} alkenyl, $-R^*YR''$, $-YR''$, and $-R''M'R'$;

R_2 and R_3 are independently selected from the group consisting of H, C_{1-14} alkyl, C_{2-14} alkenyl, $-R^*YR''$, $-YR''$, and $-R^*OR''$, or R_2 and R_3 , together with the atom to which they are attached, form a heterocycle or carbocycle;

R_4 is selected from the group consisting of a C_{3-6} carbocycle, $-(CH_2)_nQ$, $-(CH_2)_nCHQR$, $-CHQR$, $-CQ(R)_2$, and unsubstituted C_{1-6} alkyl, where Q is selected from a C_{3-6} carbocycle, a 5- to 14-membered heteroaryl having one or more heteroatoms selected from N, O, and S, $-OR$, $-O(CH_2)_nN(R)_2$, $-C(O)OR$, $-OC(O)R$, $-CX_3$, $-CX_2H$, $-CXH_2$, $-CN$, $-C(O)N(R)_2$, $-N(R)C(O)R$, $-N(R)S(O)_2R$, $-N(R)C(O)N(R)_2$, $-N(R)C(S)N(R)_2$, $-CRN(R)_2C(O)OR$, and a 5- to 14-membered heterocycloalkyl having one or more heteroatoms selected from N, O, and S which is substituted with one or more substituents selected from oxo ($=O$), OH, amino, and C_{1-3} alkyl, and each n is independently selected from 1, 2, 3, 4, and 5;

each R_5 is independently selected from the group consisting of C_{1-3} alkyl, C_{2-3} alkenyl, and H;

each R_6 is independently selected from the group consisting of C_{1-3} alkyl, C_{2-3} alkenyl, and H;

M and M' are independently selected from $-C(O)O-$, $-OC(O)-$, $-C(O)N(R')$, $-N(R')C(O)-$, $-C(O)-$, $-C(S)-$, $-C(S)S-$, $-SC(S)-$, $-CH(OH)-$, $-P(O)(OR')O-$, $-S(O)_2-$, an aryl group, and a heteroaryl group;

R_7 is selected from the group consisting of C_{1-3} alkyl, C_{2-3} alkenyl, and H;

each R is independently selected from the group consisting of C_{1-3} alkyl, C_{2-3} alkenyl, and H;

each R' is independently selected from the group consisting of C_{1-18} alkyl, C_{2-18} alkenyl, $-R^*YR''$, $-YR''$, and H;

each R'' is independently selected from the group consisting of C_{3-14} alkyl and C_{3-14} alkenyl;

each R* is independently selected from the group consisting of C₁₋₁₂ alkyl and C₂₋₁₂ alkenyl;

each Y is independently a C₃₋₆ carbocycle;

each X is independently selected from the group consisting of F, Cl, Br, and I; and

m is selected from 5, 6, 7, 8, 9, 10, 11, 12, and 13,

or salts or stereoisomers thereof.

[0659] In yet other embodiments, another subset of compounds of Formula (I) includes those in which

R₁ is selected from the group consisting of C₅₋₃₀ alkyl, C₅₋₂₀ alkenyl, -R*YR'', -YR'', and -R''M'R';

R₂ and R₃ are independently selected from the group consisting of H, C₁₋₁₄ alkyl, C₂₋₁₄ alkenyl, -R*YR'', -YR'', and -R*OR'', or R₂ and R₃, together with the atom to which they are attached, form a heterocycle or carbocycle;

R₄ is selected from the group consisting of a C₃₋₆ carbocycle, -(CH₂)_nQ, -(CH₂)_nCHQR, -CHQR, -CQ(R)₂, and unsubstituted C₁₋₆ alkyl, where Q is selected from a C₃₋₆ carbocycle, a 5- to 14-membered heterocycle having one or more heteroatoms selected from N, O, and S, -OR, -O(CH₂)_nN(R)₂, -C(O)OR, -OC(O)R, -CX₃, -CX₂H, -CXH₂, -CN, -C(O)N(R)₂, -N(R)C(O)R, -N(R)S(O)₂R, -N(R)C(O)N(R)₂, -N(R)C(S)N(R)₂, -CRN(R)₂C(O)OR, -N(R)R₈, -O(CH₂)_nOR, -N(R)C(=NR₉)N(R)₂, -N(R)C(=CHR₉)N(R)₂, -OC(O)N(R)₂, -N(R)C(O)OR, -N(OR)C(O)R, -N(OR)S(O)₂R, -N(OR)C(O)OR, -N(OR)C(O)N(R)₂, -N(OR)C(S)N(R)₂, -N(OR)C(=NR₉)N(R)₂, -N(OR)C(=CHR₉)N(R)₂, -C(=NR₉)R, -C(O)N(R)OR, and -C(=NR₉)N(R)₂, and each n is independently selected from 1, 2, 3, 4, and 5; and when Q is a 5- to 14-membered heterocycle and (i) R₄ is -(CH₂)_nQ in which n is 1 or 2, or (ii) R₄ is -(CH₂)_nCHQR in which n is 1, or (iii) R₄ is -CHQR, and -CQ(R)₂, then Q is either a 5- to 14-membered heteroaryl or 8- to 14-membered heterocycloalkyl;

each R₅ is independently selected from the group consisting of C₁₋₃ alkyl, C₂₋₃ alkenyl, and H;

each R₆ is independently selected from the group consisting of C₁₋₃ alkyl, C₂₋₃ alkenyl, and H;

M and M' are independently selected from -C(O)O-, -OC(O)-, -C(O)N(R')-, -N(R')C(O)-, -C(O)-, -C(S)-, -C(S)S-, -SC(S)-, -CH(OH)-, -P(O)(OR')O-, -S(O)₂-, -S-S-, an aryl group, and a heteroaryl group;

R₇ is selected from the group consisting of C₁₋₃ alkyl, C₂₋₃ alkenyl, and H;

R₈ is selected from the group consisting of C₃₋₆ carbocycle and heterocycle;

R₉ is selected from the group consisting of H, CN, NO₂, C₁₋₆ alkyl, -OR, -S(O)₂R, -S(O)₂N(R)₂, C₂₋₆ alkenyl, C₃₋₆ carbocycle and heterocycle;

each R is independently selected from the group consisting of C₁₋₃ alkyl, C₂₋₃ alkenyl, and H;

each R' is independently selected from the group consisting of C₁₋₁₈ alkyl, C₂₋₁₈ alkenyl, -R*YR'', -YR'', and H;

each R'' is independently selected from the group consisting of C₃₋₁₄ alkyl and C₃₋₁₄ alkenyl;

each R* is independently selected from the group consisting of C₁₋₁₂ alkyl and C₂₋₁₂ alkenyl;

each Y is independently a C₃₋₆ carbocycle;

each X is independently selected from the group consisting of F, Cl, Br, and I; and

m is selected from 5, 6, 7, 8, 9, 10, 11, 12, and 13,

or salts or stereoisomers thereof.

[0660] In yet other embodiments, another subset of compounds of Formula (I) includes those in which

R₁ is selected from the group consisting of C₅₋₂₀ alkyl, C₅₋₂₀ alkenyl, -R*YR'', -YR'', and -R''M'R';

R₂ and R₃ are independently selected from the group consisting of H, C₁₋₁₄ alkyl, C₂₋₁₄ alkenyl, -R*YR'', -YR'', and -R*OR'', or R₂ and R₃, together with the atom to which they are attached, form a heterocycle or carbocycle;

R₄ is selected from the group consisting of a C₃₋₆ carbocycle, -(CH₂)_nQ, -(CH₂)_nCHQR, -CHQR, -CQ(R)₂, and unsubstituted C₁₋₆ alkyl, where Q is selected from a C₃₋₆ carbocycle, a 5- to 14-membered heterocycle having one or more heteroatoms selected from N, O, and S, -OR, -O(CH₂)_nN(R)₂, -C(O)OR, -OC(O)R, -CX₃, -CX₂H, -CXH₂, -CN, -C(O)N(R)₂, -N(R)C(O)R, -N(R)S(O)₂R, -N(R)C(O)N(R)₂, -N(R)C(S)N(R)₂, -CRN(R)₂C(O)OR, and each n is independently selected from 1, 2, 3, 4, and 5; and when Q is a 5- to 14-membered heterocycle and (i) R₄ is -(CH₂)_nQ in which n is 1 or 2, or (ii) R₄ is -(CH₂)_nCHQR in which n is 1, or (iii) R₄ is -CHQR, and -CQ(R)₂, then Q is either a 5- to 14-membered heteroaryl or 8- to 14-membered heterocycloalkyl;

each R₅ is independently selected from the group consisting of C₁₋₃ alkyl, C₂₋₃ alkenyl, and H;

each R₆ is independently selected from the group consisting of C₁₋₃ alkyl, C₂₋₃ alkenyl, and H;

M and M' are independently selected from -C(O)O-, -OC(O)-, -C(O)N(R')-, -N(R')C(O)-, -C(O)-, -C(S)-, -C(S)S-, -SC(S)-, -CH(OH)-, -P(O)(OR')O-, -S(O)₂-, an aryl group, and a heteroaryl group;

R₇ is selected from the group consisting of C₁₋₃ alkyl, C₂₋₃ alkenyl, and H;

each R is independently selected from the group consisting of C₁₋₃ alkyl, C₂₋₃ alkenyl, and H;

each R' is independently selected from the group consisting of C₁₋₁₈ alkyl, C₂₋₁₈ alkenyl, -R*YR'', -YR'', and H;

each R'' is independently selected from the group consisting of C₃₋₁₄ alkyl and C₃₋₁₄ alkenyl;

each R* is independently selected from the group consisting of C₁₋₁₂ alkyl and C₂₋₁₂ alkenyl;

each Y is independently a C₃₋₆ carbocycle;

each X is independently selected from the group consisting of F, Cl, Br, and I; and m is selected from 5, 6, 7, 8, 9, 10, 11, 12, and 13,

or salts or stereoisomers thereof.

[0661] In still another embodiments, another subset of compounds of Formula (I) includes those in which

R₁ is selected from the group consisting of C₅₋₃₀ alkyl, C₅₋₂₀ alkenyl, -R*YR'', -YR'', and -R''M'R';

R₂ and R₃ are independently selected from the group consisting of H, C₁₋₁₄ alkyl, C₂₋₁₄ alkenyl, -R*YR'', -YR'', and -R*OR'', or R₂ and R₃, together with the atom to which they are attached, form a heterocycle or carbocycle;

R₄ is selected from the group consisting of a C₃₋₆ carbocycle, -(CH₂)_nQ, -(CH₂)_nCHQR, -CHQR, -CQ(R)₂, and unsubstituted C₁₋₆ alkyl, where Q is selected from a C₃₋₆ carbocycle, a 5- to 14-membered heteroaryl having one or more heteroatoms selected from N, O, and S, -OR, -O(CH₂)_nN(R)₂, -C(O)OR, -OC(O)R, -CX₃, -CX₂H, -CXH₂, -CN, -C(O)N(R)₂, -N(R)C(O)R, -N(R)S(O)₂R, -N(R)C(O)N(R)₂, -N(R)C(S)N(R)₂, -CRN(R)₂C(O)OR, -N(R)R₈, -O(CH₂)_nOR, -N(R)C(=NR₉)N(R)₂, -N(R)C(=CHR₉)N(R)₂,

-OC(O)N(R)₂, -N(R)C(O)OR, -N(OR)C(O)R, -N(OR)S(O)₂R, -N(OR)C(O)OR, -N(OR)C(O)N(R)₂, -N(OR)C(S)N(R)₂, -N(OR)C(=NR₉)N(R)₂, -N(OR)C(=CHR₉)N(R)₂, -C(=NR₉)R, -C(O)N(R)OR, and -C(=NR₉)N(R)₂, and each n is independently selected from 1, 2, 3, 4, and 5;

each R₅ is independently selected from the group consisting of C₁₋₃ alkyl, C₂₋₃ alkenyl, and H;

each R₆ is independently selected from the group consisting of C₁₋₃ alkyl, C₂₋₃ alkenyl, and H;

M and M' are independently selected from -C(O)O-, -OC(O)-, -C(O)N(R')-, -N(R')C(O)-, -C(O)-, -C(S)-, -C(S)S-, -SC(S)-, -CH(OH)-, -P(O)(OR')O-, -S(O)₂-, -S-S-, an aryl group, and a heteroaryl group;

R₇ is selected from the group consisting of C₁₋₃ alkyl, C₂₋₃ alkenyl, and H;

R₈ is selected from the group consisting of C₃₋₆ carbocycle and heterocycle;

R₉ is selected from the group consisting of H, CN, NO₂, C₁₋₆ alkyl, -OR, -S(O)₂R, -S(O)₂N(R)₂, C₂₋₆ alkenyl, C₃₋₆ carbocycle and heterocycle;

each R is independently selected from the group consisting of C₁₋₃ alkyl, C₂₋₃ alkenyl, and H;

each R' is independently selected from the group consisting of C₁₋₁₈ alkyl, C₂₋₁₈ alkenyl, -R*YR'', -YR'', and H;

each R'' is independently selected from the group consisting of C₃₋₁₄ alkyl and C₃₋₁₄ alkenyl;

each R* is independently selected from the group consisting of C₁₋₁₂ alkyl and C₂₋₁₂ alkenyl;

each Y is independently a C₃₋₆ carbocycle;

each X is independently selected from the group consisting of F, Cl, Br, and I; and

m is selected from 5, 6, 7, 8, 9, 10, 11, 12, and 13,

or salts or stereoisomers thereof.

[0662] In still another embodiments, another subset of compounds of Formula (I) includes those in which

R₁ is selected from the group consisting of C₅₋₂₀ alkyl, C₅₋₂₀ alkenyl, -R*YR'', -YR'', and -R''M'R';

R_2 and R_3 are independently selected from the group consisting of H, C_{1-14} alkyl, C_{2-14} alkenyl, $-R^*YR''$, $-YR''$, and $-R^*OR''$, or R_2 and R_3 , together with the atom to which they are attached, form a heterocycle or carbocycle;

R_4 is selected from the group consisting of a C_{3-6} carbocycle, $-(CH_2)_nQ$, $-(CH_2)_nCHQR$, $-CHQR$, $-CQ(R)_2$, and unsubstituted C_{1-6} alkyl, where Q is selected from a C_{3-6} carbocycle, a 5- to 14-membered heteroaryl having one or more heteroatoms selected from N, O, and S, $-OR$, $-O(CH_2)_nN(R)_2$, $-C(O)OR$, $-OC(O)R$, $-CX_3$, $-CX_2H$, $-CXH_2$, $-CN$, $-C(O)N(R)_2$, $-N(R)C(O)R$, $-N(R)S(O)_2R$, $-N(R)C(O)N(R)_2$, $-N(R)C(S)N(R)_2$, $-CRN(R)_2C(O)OR$, and each n is independently selected from 1, 2, 3, 4, and 5;

each R_5 is independently selected from the group consisting of C_{1-3} alkyl, C_{2-3} alkenyl, and H;

each R_6 is independently selected from the group consisting of C_{1-3} alkyl, C_{2-3} alkenyl, and H;

M and M' are independently selected from $-C(O)O-$, $-OC(O)-$, $-C(O)N(R')-$, $-N(R')C(O)-$, $-C(O)-$, $-C(S)-$, $-C(S)S-$, $-SC(S)-$, $-CH(OH)-$, $-P(O)(OR')O-$, $-S(O)_2-$, an aryl group, and a heteroaryl group;

R_7 is selected from the group consisting of C_{1-3} alkyl, C_{2-3} alkenyl, and H;

each R is independently selected from the group consisting of C_{1-3} alkyl, C_{2-3} alkenyl, and H;

each R' is independently selected from the group consisting of C_{1-18} alkyl, C_{2-18} alkenyl, $-R^*YR''$, $-YR''$, and H;

each R'' is independently selected from the group consisting of C_{3-14} alkyl and C_{3-14} alkenyl;

each R^* is independently selected from the group consisting of C_{1-12} alkyl and C_{2-12} alkenyl;

each Y is independently a C_{3-6} carbocycle;

each X is independently selected from the group consisting of F, Cl, Br, and I; and

m is selected from 5, 6, 7, 8, 9, 10, 11, 12, and 13,

or salts or stereoisomers thereof.

[0663] In yet other embodiments, another subset of compounds of Formula (I) includes those in which

R_1 is selected from the group consisting of C_{5-30} alkyl, C_{5-20} alkenyl, $-R^*YR''$, $-YR''$, and $-R''M'R'$;

R_2 and R_3 are independently selected from the group consisting of H, C_{2-14} alkyl, C_{2-14} alkenyl, $-R^*YR''$, $-YR''$, and $-R^*OR''$, or R_2 and R_3 , together with the atom to which they are attached, form a heterocycle or carbocycle;

R_4 is $-(CH_2)_nQ$ or $-(CH_2)_nCHQR$, where Q is $-N(R)_2$, and n is selected from 3, 4, and 5;

each R_5 is independently selected from the group consisting of C_{1-3} alkyl, C_{2-3} alkenyl, and H;

each R_6 is independently selected from the group consisting of C_{1-3} alkyl, C_{2-3} alkenyl, and H;

M and M' are independently selected from $-C(O)O-$, $-OC(O)-$, $-C(O)N(R')$, $-N(R')C(O)-$, $-C(O)-$, $-C(S)-$, $-C(S)S-$, $-SC(S)-$, $-CH(OH)-$, $-P(O)(OR')O-$, $-S(O)_2-$, $-S-S-$, an aryl group, and a heteroaryl group;

R_7 is selected from the group consisting of C_{1-3} alkyl, C_{2-3} alkenyl, and H;

each R is independently selected from the group consisting of C_{1-3} alkyl, C_{2-3} alkenyl, and H;

each R' is independently selected from the group consisting of C_{1-18} alkyl, C_{2-18} alkenyl, $-R^*YR''$, $-YR''$, and H;

each R'' is independently selected from the group consisting of C_{3-14} alkyl and C_{3-14} alkenyl;

each R^* is independently selected from the group consisting of C_{1-12} alkyl and C_{1-12} alkenyl;

each Y is independently a C_{3-6} carbocycle;

each X is independently selected from the group consisting of F, Cl, Br, and I; and

m is selected from 5, 6, 7, 8, 9, 10, 11, 12, and 13,

or salts or stereoisomers thereof.

[0664] In yet other embodiments, another subset of compounds of Formula (I) includes those in which

R_1 is selected from the group consisting of C_{5-20} alkyl, C_{5-20} alkenyl, $-R^*YR''$, $-YR''$, and $-R''M'R'$;

R_2 and R_3 are independently selected from the group consisting of H, C_{2-14} alkyl, C_{2-14} alkenyl, $-R^*YR''$, $-YR''$, and $-R^*OR''$, or R_2 and R_3 , together with the atom to which they are attached, form a heterocycle or carbocycle;

R_4 is $-(CH_2)_nQ$ or $-(CH_2)_nCHQR$, where Q is $-N(R)_2$, and n is selected from 3, 4, and 5;

each R_5 is independently selected from the group consisting of C_{1-3} alkyl, C_{2-3} alkenyl, and H;

each R_6 is independently selected from the group consisting of C_{1-3} alkyl, C_{2-3} alkenyl, and H;

M and M' are independently selected from $-C(O)O-$, $-OC(O)-$, $-C(O)N(R')$, $-N(R')C(O)-$, $-C(O)-$, $-C(S)-$, $-C(S)S-$, $-SC(S)-$, $-CH(OH)-$, $-P(O)(OR')O-$, $-S(O)_2-$, an aryl group, and a heteroaryl group;

R_7 is selected from the group consisting of C_{1-3} alkyl, C_{2-3} alkenyl, and H;

each R is independently selected from the group consisting of C_{1-3} alkyl, C_{2-3} alkenyl, and H;

each R' is independently selected from the group consisting of C_{1-18} alkyl, C_{2-18} alkenyl, $-R^*YR''$, $-YR''$, and H;

each R'' is independently selected from the group consisting of C_{3-14} alkyl and C_{3-14} alkenyl;

each R^* is independently selected from the group consisting of C_{1-12} alkyl and C_{1-12} alkenyl;

each Y is independently a C_{3-6} carbocycle;

each X is independently selected from the group consisting of F, Cl, Br, and I; and

m is selected from 5, 6, 7, 8, 9, 10, 11, 12, and 13,

or salts or stereoisomers thereof.

[0665] In still other embodiments, another subset of compounds of Formula (I) includes those in which

R_1 is selected from the group consisting of C_{5-30} alkyl, C_{5-20} alkenyl, $-R^*YR''$, $-YR''$, and $-R''M'R'$;

R_2 and R_3 are independently selected from the group consisting of C_{1-14} alkyl, C_{2-14} alkenyl, $-R^*YR''$, $-YR''$, and $-R^*OR''$, or R_2 and R_3 , together with the atom to which they are attached, form a heterocycle or carbocycle;

R_4 is selected from the group consisting of $-(CH_2)_nQ$, $-(CH_2)_nCHQR$, $-CHQR$, and $-CQ(R)_2$, where Q is $-N(R)_2$, and n is selected from 1, 2, 3, 4, and 5;

each R_5 is independently selected from the group consisting of C_{1-3} alkyl, C_{2-3} alkenyl, and H;

each R₆ is independently selected from the group consisting of C₁₋₃ alkyl, C₂₋₃ alkenyl, and H;

M and M' are independently selected from -C(O)O-, -OC(O)-, -C(O)N(R')-, -N(R')C(O)-, -C(O)-, -C(S)-, -C(S)S-, -SC(S)-, -CH(OH)-, -P(O)(OR')O-, -S(O)₂-, -S-S-, an aryl group, and a heteroaryl group;

R₇ is selected from the group consisting of C₁₋₃ alkyl, C₂₋₃ alkenyl, and H;

each R is independently selected from the group consisting of C₁₋₃ alkyl, C₂₋₃ alkenyl, and H;

each R' is independently selected from the group consisting of C₁₋₁₈ alkyl, C₂₋₁₈ alkenyl, -R*YR'', -YR'', and H;

each R'' is independently selected from the group consisting of C₃₋₁₄ alkyl and C₃₋₁₄ alkenyl;

each R* is independently selected from the group consisting of C₁₋₁₂ alkyl and C₁₋₁₂ alkenyl;

each Y is independently a C₃₋₆ carbocycle;

each X is independently selected from the group consisting of F, Cl, Br, and I; and m is selected from 5, 6, 7, 8, 9, 10, 11, 12, and 13,

or salts or stereoisomers thereof.

[0666] In still other embodiments, another subset of compounds of Formula (I) includes those in which

R₁ is selected from the group consisting of C₅₋₂₀ alkyl, C₅₋₂₀ alkenyl, -R*YR'', -YR'', and -R''M'R';

R₂ and R₃ are independently selected from the group consisting of C₁₋₁₄ alkyl, C₂₋₁₄ alkenyl, -R*YR'', -YR'', and -R*OR'', or R₂ and R₃, together with the atom to which they are attached, form a heterocycle or carbocycle;

R₄ is selected from the group consisting of -(CH₂)_nQ, -(CH₂)_nCHQR, -CHQR, and -CQ(R)₂, where Q is -N(R)₂, and n is selected from 1, 2, 3, 4, and 5;

each R₅ is independently selected from the group consisting of C₁₋₃ alkyl, C₂₋₃ alkenyl, and H;

each R₆ is independently selected from the group consisting of C₁₋₃ alkyl, C₂₋₃ alkenyl, and H;

M and M' are independently selected from -C(O)O-, -OC(O)-, -C(O)N(R')-, -N(R')C(O)-, -C(O)-, -C(S)-, -C(S)S-, -SC(S)-, -CH(OH)-, -P(O)(OR')O-, -S(O)₂-, an aryl group, and a heteroaryl group;

R₇ is selected from the group consisting of C₁₋₃ alkyl, C₂₋₃ alkenyl, and H;

each R is independently selected from the group consisting of C₁₋₃ alkyl, C₂₋₃ alkenyl, and H;

each R' is independently selected from the group consisting of C₁₋₁₈ alkyl, C₂₋₁₈ alkenyl, -R*YR'', -YR'', and H;

each R'' is independently selected from the group consisting of C₃₋₁₄ alkyl and C₃₋₁₄ alkenyl;

each R* is independently selected from the group consisting of C₁₋₁₂ alkyl and C₁₋₁₂ alkenyl;


each Y is independently a C₃₋₆ carbocycle;

each X is independently selected from the group consisting of F, Cl, Br, and I; and

m is selected from 5, 6, 7, 8, 9, 10, 11, 12, and 13,

or salts or stereoisomers thereof.

[0667] In certain embodiments, a subset of compounds of Formula (I) includes those of Formula (IA):



or a salt or stereoisomer thereof, wherein l is selected from 1, 2, 3, 4, and 5; m is selected from 5, 6, 7, 8, and 9; M₁ is a bond or M'; R₄ is unsubstituted C₁₋₃ alkyl, or -(CH₂)_nQ, in which Q is OH, -NHC(S)N(R)₂, -NHC(O)N(R)₂, N(R)C(O)R, -N(R)S(O)₂R, -N(R)R₈, -NHC(=NR₉)N(R)₂, -NHC(=CHR₉)N(R)₂, -OC(O)N(R)₂, -N(R)C(O)OR, heteroaryl, or heterocycloalkyl; M and M' are independently selected from -C(O)O-, -OC(O)-, -C(O)N(R')-, -P(O)(OR')O-, -S-S-, an aryl group, and a heteroaryl group; and

R₂ and R₃ are independently selected from the group consisting of H, C₁₋₁₄ alkyl, and C₂₋₁₄ alkenyl.

[0668] In some embodiments, a subset of compounds of Formula (I) includes those of Formula (IA), or a salt or stereoisomer thereof, wherein

l is selected from 1, 2, 3, 4, and 5; m is selected from 5, 6, 7, 8, and 9;


M_1 is a bond or M' ;

R_4 is unsubstituted C_{1-3} alkyl, or $-(CH_2)_nQ$, in which Q is OH, $-NHC(S)N(R)_2$, or $-NHC(O)N(R)_2$;

M and M' are independently selected from $-C(O)O-$, $-OC(O)-$, $-C(O)N(R')$ -, $-P(O)(OR')O-$, an aryl group, and a heteroaryl group; and

R_2 and R_3 are independently selected from the group consisting of H, C_{1-14} alkyl, and C_{2-14} alkenyl.

[0669] In certain embodiments, a subset of compounds of Formula (I) includes those of Formula (II):



or a salt or stereoisomer thereof, wherein l is selected from 1, 2, 3, 4, and 5; M_1 is a bond or M' ; R_4 is unsubstituted C_{1-3} alkyl, or $-(CH_2)_nQ$, in which n is 2, 3, or 4, and Q is OH, $-NHC(S)N(R)_2$, $-NHC(O)N(R)_2$, $-N(R)C(O)R$, $-N(R)S(O)_2R$, $-N(R)R_8$, $-NHC(=NR_9)N(R)_2$, $-NHC(=CHR_9)N(R)_2$, $-OC(O)N(R)_2$, $-N(R)C(O)OR$, heteroaryl, or heterocycloalkyl; M and M' are independently selected from $-C(O)O-$, $-OC(O)-$, $-C(O)N(R')$ -, $-P(O)(OR')O-$, $-S-S-$, an aryl group, and a heteroaryl group; and

R_2 and R_3 are independently selected from the group consisting of H, C_{1-14} alkyl, and C_{2-14} alkenyl.

[0670] In some embodiments, a subset of compounds of Formula (I) includes those of Formula (II), or a salt or stereoisomer thereof, wherein

l is selected from 1, 2, 3, 4, and 5;


M_1 is a bond or M' ;

R_4 is unsubstituted C_{1-3} alkyl, or $-(CH_2)_nQ$, in which n is 2, 3, or 4, and Q is OH, $-NHC(S)N(R)_2$, or $-NHC(O)N(R)_2$;

M and M' are independently selected from $-C(O)O-$, $-OC(O)-$, $-C(O)N(R')$ -, $-P(O)(OR')O-$, an aryl group, and a heteroaryl group; and

R_2 and R_3 are independently selected from the group consisting of H, C_{1-14} alkyl, and C_{2-14} alkenyl.


[0671] In some embodiments, the compound of Formula (I) is of the Formula (IIa),



(IIa),

or a salt thereof, wherein R_4 is as described above.


[0672] In some embodiments, the compound of Formula (I) is of the Formula (IIb),



(IIb),

or a salt thereof, wherein R_4 is as described above.


[0673] In some embodiments, the compound of Formula (I) is of the Formula (IIc),



(IIc),

or a salt thereof, wherein R_4 is as described above.

[0674] In some embodiments, the compound of Formula (I) is of the Formula (IIE):



(IIE),


or a salt thereof, wherein R_4 is as described above.

[0675] In some embodiments, the compound of Formula (IIa), (IIb), (IIc), or (IIE) comprises an R_4 which is selected from $-(CH_2)_nQ$ and $-(CH_2)_nCHQR$, wherein Q , R and n are as defined above.

[0676] In some embodiments, Q is selected from the group consisting of $-OR$, $-OH$, $-O(CH_2)_nN(R)_2$, $-OC(O)R$, $-CX_3$, $-CN$, $-N(R)C(O)R$, $-N(H)C(O)R$, $-N(R)S(O)_2R$, $-N(H)S(O)_2R$, $-N(R)C(O)N(R)_2$, $-N(H)C(O)N(R)_2$, $-N(H)C(O)N(H)(R)$, $-N(R)C(S)N(R)_2$, $-N(H)C(S)N(R)_2$, $-N(H)C(S)N(H)(R)$, and a heterocycle, wherein R is as defined above.

In some aspects, n is 1 or 2. In some embodiments, Q is OH, -NHC(S)N(R)₂, or -NHC(O)N(R)₂.

[0677] In some embodiments, the compound of Formula (I) is of the Formula (II_d),


$$(II_d),$$

or a salt thereof, wherein R₂ and R₃ are independently selected from the group consisting of C₅₋₁₄ alkyl and C₅₋₁₄ alkenyl, n is selected from 2, 3, and 4, and R', R'', R₅, R₆ and m are as defined above.

[0678] In some aspects of the compound of Formula (IIId), R₂ is C₈ alkyl. In some aspects of the compound of Formula (IIId), R₃ is C₅-C₉ alkyl. In some aspects of the compound of Formula (IIId), m is 5, 7, or 9. In some aspects of the compound of Formula (IIId), each R₅ is H. In some aspects of the compound of Formula (IIId), each R₆ is H.

[0679] In another aspect, the present application provides a lipid composition (e.g., a lipid nanoparticle (LNP)) comprising: (1) a compound having the Formula (I); (2) optionally a helper lipid (e.g. a phospholipid); (3) optionally a structural lipid (e.g. a sterol); and (4) optionally a lipid conjugate (e.g. a PEG-lipid). In exemplary embodiments, the lipid composition (e.g., LNP) further comprises a polynucleotide encoding an ACADVL polypeptide, e.g., a polynucleotide encapsulated therein.

[0680] As used herein, the term “alkyl” or “alkyl group” means a linear or branched, saturated hydrocarbon including one or more carbon atoms (e.g., one, two, three, four, five, six, seven, eight, nine, ten, eleven, twelve, thirteen, fourteen, fifteen, sixteen, seventeen, eighteen, nineteen, twenty, or more carbon atoms).

[0681] The notation “C₁₋₁₄ alkyl” means a linear or branched, saturated hydrocarbon including 1-14 carbon atoms. An alkyl group can be optionally substituted.

[0682] As used herein, the term “alkenyl” or “alkenyl group” means a linear or branched hydrocarbon including two or more carbon atoms (e.g., two, three, four, five, six, seven, eight, nine, ten, eleven, twelve, thirteen, fourteen, fifteen, sixteen, seventeen, eighteen, nineteen, twenty, or more carbon atoms) and at least one double bond.

- [0683]** The notation “C₂₋₁₄ alkenyl” means a linear or branched hydrocarbon including 2-14 carbon atoms and at least one double bond. An alkenyl group can include one, two, three, four, or more double bonds. For example, C₁₈ alkenyl can include one or more double bonds. A C₁₈ alkenyl group including two double bonds can be a linoleyl group. An alkenyl group can be optionally substituted.
- [0684]** As used herein, the term “carbocycle” or “carbocyclic group” means a mono- or multi-cyclic system including one or more rings of carbon atoms. Rings can be three, four, five, six, seven, eight, nine, ten, eleven, twelve, thirteen, fourteen, or fifteen membered rings.
- [0685]** The notation “C₃₋₆ carbocycle” means a carbocycle including a single ring having 3-6 carbon atoms. Carbocycles can include one or more double bonds and can be aromatic (e.g., aryl groups). Examples of carbocycles include cyclopropyl, cyclopentyl, cyclohexyl, phenyl, naphthyl, and 1,2-dihydronaphthyl groups. Carbocycles can be optionally substituted.
- [0686]** As used herein, the term “heterocycle” or “heterocyclic group” means a mono- or multi-cyclic system including one or more rings, where at least one ring includes at least one heteroatom. Heteroatoms can be, for example, nitrogen, oxygen, or sulfur atoms. Rings can be three, four, five, six, seven, eight, nine, ten, eleven, or twelve membered rings. Heterocycles can include one or more double bonds and can be aromatic (e.g., heteroaryl groups). Examples of heterocycles include imidazolyl, imidazolidinyl, oxazolyl, oxazolidinyl, thiazolyl, thiazolidinyl, pyrazolidinyl, pyrazolyl, isoxazolidinyl, isoxazolyl, isothiazolidinyl, isothiazolyl, morpholinyl, pyrrolyl, pyrrolidinyl, furyl, tetrahydrofuryl, thiophenyl, pyridinyl, piperidinyl, quinolyl, and isoquinolyl groups. Heterocycles can be optionally substituted.
- [0687]** As used herein, a “biodegradable group” is a group that can facilitate faster metabolism of a lipid in a subject. A biodegradable group can be, but is not limited to, -C(O)O-, -OC(O)-, -C(O)N(R’)-, -N(R’)C(O)-, -C(O)-, -C(S)-, -C(S)S-, -SC(S)-, -CH(OH)-, -P(O)(OR’)O-, -S(O)₂-, an aryl group, and a heteroaryl group.
- [0688]** As used herein, an “aryl group” is a carbocyclic group including one or more aromatic rings. Examples of aryl groups include phenyl and naphthyl groups.
- [0689]** As used herein, a “heteroaryl group” is a heterocyclic group including one or more aromatic rings. Examples of heteroaryl groups include pyrrolyl, furyl, thiophenyl, imidazolyl, oxazolyl, and thiazolyl. Both aryl and heteroaryl groups can be optionally

substituted. For example, M and M' can be selected from the non-limiting group consisting of optionally substituted phenyl, oxazole, and thiazole. In the formulas herein, M and M' can be independently selected from the list of biodegradable groups above.

[0690] Alkyl, alkenyl, and cyclyl (e.g., carbocyclyl and heterocyclyl) groups can be optionally substituted unless otherwise specified. Optional substituents can be selected from the group consisting of, but are not limited to, a halogen atom (e.g., a chloride, bromide, fluoride, or iodide group), a carboxylic acid (e.g., -C(O)OH), an alcohol (e.g., a hydroxyl, -OH), an ester (e.g., -C(O)OR or -OC(O)R), an aldehyde (e.g., -C(O)H), a carbonyl (e.g., -C(O)R, alternatively represented by C=O), an acyl halide (e.g., -C(O)X, in which X is a halide selected from bromide, fluoride, chloride, and iodide), a carbonate (e.g., -OC(O)OR), an alkoxy (e.g., -OR), an acetal (e.g., -C(OR)₂R^{""}, in which each OR are alkoxy groups that can be the same or different and R^{""} is an alkyl or alkenyl group), a phosphate (e.g., P(O)₄³⁻), a thiol (e.g., -SH), a sulfoxide (e.g., -S(O)R), a sulfinic acid (e.g., -S(O)OH), a sulfonic acid (e.g., -S(O)₂OH), a thial (e.g., -C(S)H), a sulfate (e.g., S(O)₄²⁻), a sulfonyl (e.g., -S(O)₂-), an amide (e.g., -C(O)NR₂, or -N(R)C(O)R), an azido (e.g., -N₃), a nitro (e.g., -NO₂), a cyano (e.g., -CN), an isocyano (e.g., -NC), an acyloxy (e.g., -OC(O)R), an amino (e.g., -NR₂, -NRH, or -NH₂), a carbamoyl (e.g., -OC(O)NR₂, -OC(O)NRH, or -OC(O)NH₂), a sulfonamide (e.g., -S(O)₂NR₂, -S(O)₂NRH, -S(O)₂NH₂, -N(R)S(O)₂R, -N(H)S(O)₂R, -N(R)S(O)₂H, or -N(H)S(O)₂H), an alkyl group, an alkenyl group, and a cyclyl (e.g., carbocyclyl or heterocyclyl) group.

[0691] In any of the preceding, R is an alkyl or alkenyl group, as defined herein. In some embodiments, the substituent groups themselves can be further substituted with, for example, one, two, three, four, five, or six substituents as defined herein. For example, a C₁₋₆ alkyl group can be further substituted with one, two, three, four, five, or six substituents as described herein.

[0692] The compounds of any one of Formulae (I), (IA), (II), (IIa), (IIb), (IIc), (IId), and (IIe) include one or more of the following features when applicable.

[0693] In some embodiments, R₄ is selected from the group consisting of a C₃₋₆ carbocycle, -(CH₂)_nQ, -(CH₂)_nCHQR, -CHQR, and -CQ(R)₂, where Q is selected from a C₃₋₆ carbocycle, 5- to 14- membered aromatic or non-aromatic heterocycle having one or more heteroatoms selected from N, O, S, and P, -OR, -O(CH₂)_nN(R)₂, -C(O)OR, -OC(O)R, -CX₃, -CX₂H, -CXH₂, -CN, -N(R)₂, -C(O)N(R)₂, -N(R)C(O)R, -N(R)S(O)₂R,

$-\text{N}(\text{R})\text{C}(\text{O})\text{N}(\text{R})_2$, $-\text{N}(\text{R})\text{C}(\text{S})\text{N}(\text{R})_2$, and $-\text{C}(\text{R})\text{N}(\text{R})_2\text{C}(\text{O})\text{OR}$, and each n is independently selected from 1, 2, 3, 4, and 5.

[0694] In another embodiment, R_4 is selected from the group consisting of a C_{3-6} carbocycle, $-(\text{CH}_2)_n\text{Q}$, $-(\text{CH}_2)_n\text{CHQR}$, $-\text{CHQR}$, and $-\text{CQ}(\text{R})_2$, where Q is selected from a C_{3-6} carbocycle, a 5- to 14-membered heteroaryl having one or more heteroatoms selected from N, O, and S, $-\text{OR}$, $-\text{O}(\text{CH}_2)_n\text{N}(\text{R})_2$, $-\text{C}(\text{O})\text{OR}$, $-\text{OC}(\text{O})\text{R}$, $-\text{CX}_3$, $-\text{CX}_2\text{H}$, $-\text{CXH}_2$, $-\text{CN}$, $-\text{C}(\text{O})\text{N}(\text{R})_2$, $-\text{N}(\text{R})\text{C}(\text{O})\text{R}$, $-\text{N}(\text{R})\text{S}(\text{O})_2\text{R}$, $-\text{N}(\text{R})\text{C}(\text{O})\text{N}(\text{R})_2$, $-\text{N}(\text{R})\text{C}(\text{S})\text{N}(\text{R})_2$, $-\text{C}(\text{R})\text{N}(\text{R})_2\text{C}(\text{O})\text{OR}$, and a 5- to 14-membered heterocycloalkyl having one or more heteroatoms selected from N, O, and S which is substituted with one or more substituents selected from oxo ($=\text{O}$), OH, amino, and C_{1-3} alkyl, and each n is independently selected from 1, 2, 3, 4, and 5.


[0695] In another embodiment, R_4 is selected from the group consisting of a C_{3-6} carbocycle, $-(\text{CH}_2)_n\text{Q}$, $-(\text{CH}_2)_n\text{CHQR}$, $-\text{CHQR}$, and $-\text{CQ}(\text{R})_2$, where Q is selected from a C_{3-6} carbocycle, a 5- to 14-membered heterocycle having one or more heteroatoms selected from N, O, and S, $-\text{OR}$, $-\text{O}(\text{CH}_2)_n\text{N}(\text{R})_2$, $-\text{C}(\text{O})\text{OR}$, $-\text{OC}(\text{O})\text{R}$, $-\text{CX}_3$, $-\text{CX}_2\text{H}$, $-\text{CXH}_2$, $-\text{CN}$, $-\text{C}(\text{O})\text{N}(\text{R})_2$, $-\text{N}(\text{R})\text{C}(\text{O})\text{R}$, $-\text{N}(\text{R})\text{S}(\text{O})_2\text{R}$, $-\text{N}(\text{R})\text{C}(\text{O})\text{N}(\text{R})_2$, $-\text{N}(\text{R})\text{C}(\text{S})\text{N}(\text{R})_2$, $-\text{C}(\text{R})\text{N}(\text{R})_2\text{C}(\text{O})\text{OR}$, and each n is independently selected from 1, 2, 3, 4, and 5; and when Q is a 5- to 14-membered heterocycle and (i) R_4 is $-(\text{CH}_2)_n\text{Q}$ in which n is 1 or 2, or (ii) R_4 is $-(\text{CH}_2)_n\text{CHQR}$ in which n is 1, or (iii) R_4 is $-\text{CHQR}$, and $-\text{CQ}(\text{R})_2$, then Q is either a 5- to 14-membered heteroaryl or 8- to 14-membered heterocycloalkyl.

[0696] In another embodiment, R_4 is selected from the group consisting of a C_{3-6} carbocycle, $-(\text{CH}_2)_n\text{Q}$, $-(\text{CH}_2)_n\text{CHQR}$, $-\text{CHQR}$, and $-\text{CQ}(\text{R})_2$, where Q is selected from a C_{3-6} carbocycle, a 5- to 14-membered heteroaryl having one or more heteroatoms selected from N, O, and S, $-\text{OR}$, $-\text{O}(\text{CH}_2)_n\text{N}(\text{R})_2$, $-\text{C}(\text{O})\text{OR}$, $-\text{OC}(\text{O})\text{R}$, $-\text{CX}_3$, $-\text{CX}_2\text{H}$, $-\text{CXH}_2$, $-\text{CN}$, $-\text{C}(\text{O})\text{N}(\text{R})_2$, $-\text{N}(\text{R})\text{C}(\text{O})\text{R}$, $-\text{N}(\text{R})\text{S}(\text{O})_2\text{R}$, $-\text{N}(\text{R})\text{C}(\text{O})\text{N}(\text{R})_2$, $-\text{N}(\text{R})\text{C}(\text{S})\text{N}(\text{R})_2$, $-\text{C}(\text{R})\text{N}(\text{R})_2\text{C}(\text{O})\text{OR}$, and each n is independently selected from 1, 2, 3, 4, and 5.

[0697] In another embodiment, R_4 is unsubstituted C_{1-4} alkyl, e.g., unsubstituted methyl.

[0698] In certain embodiments, the disclosure provides a compound having the Formula (I), wherein R_4 is $-(\text{CH}_2)_n\text{Q}$ or $-(\text{CH}_2)_n\text{CHQR}$, where Q is $-\text{N}(\text{R})_2$, and n is selected from 3, 4, and 5.

[0699] In certain embodiments, the disclosure provides a compound having the Formula (I), wherein R_4 is selected from the group consisting of $-(\text{CH}_2)_n\text{Q}$, $-(\text{CH}_2)_n\text{CHQR}$, $-\text{CHQR}$, and $-\text{CQ}(\text{R})_2$, where Q is $-\text{N}(\text{R})_2$, and n is selected from 1, 2, 3, 4, and 5.


- [0700] In certain embodiments, the disclosure provides a compound having the Formula (I), wherein R_2 and R_3 are independently selected from the group consisting of C_{2-14} alkyl, C_{2-14} alkenyl, $-R^*YR''$, $-YR''$, and $-R^*OR''$, or R_2 and R_3 , together with the atom to which they are attached, form a heterocycle or carbocycle, and R_4 is $-(CH_2)_nQ$ or $-(CH_2)_nCHQR$, where Q is $-N(R)_2$, and n is selected from 3, 4, and 5.
- [0701] In certain embodiments, R_2 and R_3 are independently selected from the group consisting of C_{2-14} alkyl, C_{2-14} alkenyl, $-R^*YR''$, $-YR''$, and $-R^*OR''$, or R_2 and R_3 , together with the atom to which they are attached, form a heterocycle or carbocycle.
- [0702] In some embodiments, R_1 is selected from the group consisting of C_{5-20} alkyl and C_{5-20} alkenyl.
- [0703] In other embodiments, R_1 is selected from the group consisting of $-R^*YR''$, $-YR''$, and $-R''M'R'$.
- [0704] In certain embodiments, R_1 is selected from $-R^*YR''$ and $-YR''$. In some embodiments, Y is a cyclopropyl group. In some embodiments, R^* is C_8 alkyl or C_8 alkenyl. In certain embodiments, R'' is C_{3-12} alkyl. For example, R'' can be C_3 alkyl. For example, R'' can be C_{4-8} alkyl (e.g., C_4 , C_5 , C_6 , C_7 , or C_8 alkyl).
- [0705] In some embodiments, R_1 is C_{5-20} alkyl. In some embodiments, R_1 is C_6 alkyl. In some embodiments, R_1 is C_8 alkyl. In other embodiments, R_1 is C_9 alkyl. In certain embodiments, R_1 is C_{14} alkyl. In other embodiments, R_1 is C_{18} alkyl.
- [0706] In some embodiments, R_1 is C_{5-20} alkenyl. In certain embodiments, R_1 is C_{18} alkenyl. In some embodiments, R_1 is linoleyl.
- [0707] In certain embodiments, R_1 is branched (e.g., decan-2-yl, undecan-3-yl, dodecan-4-yl, tridecan-5-yl, tetradecan-6-yl, 2-methylundecan-3-yl, 2-methyldecan-2-yl, 3-methylundecan-3-yl, 4-methyldodecan-4-yl, or heptadeca-9-yl). In certain embodiments, R_1 is .
- [0708] In certain embodiments, R_1 is unsubstituted C_{5-20} alkyl or C_{5-20} alkenyl. In certain embodiments, R' is substituted C_{5-20} alkyl or C_{5-20} alkenyl (e.g., substituted with a C_{3-6} carbocycle such as 1-cyclopropylnonyl).
- [0709] In other embodiments, R_1 is $-R''M'R'$.
- [0710] In some embodiments, R' is selected from $-R^*YR''$ and $-YR''$. In some embodiments, Y is C_{3-8} cycloalkyl. In some embodiments, Y is C_{6-10} aryl. In some

embodiments, Y is a cyclopropyl group. In some embodiments, Y is a cyclohexyl group. In certain embodiments, R* is C₁ alkyl.

[0711] In some embodiments, R'' is selected from the group consisting of C₃₋₁₂ alkyl and C₃₋₁₂ alkenyl. In some embodiments, R'' adjacent to Y is C₁ alkyl. In some embodiments, R'' adjacent to Y is C₄₋₉ alkyl (*e.g.*, C₄, C₅, C₆, C₇ or C₈ or C₉ alkyl).

[0712] In some embodiments, R' is selected from C₄ alkyl and C₄ alkenyl. In certain embodiments, R' is selected from C₅ alkyl and C₅ alkenyl. In some embodiments, R' is selected from C₆ alkyl and C₆ alkenyl. In some embodiments, R' is selected from C₇ alkyl and C₇ alkenyl. In some embodiments, R' is selected from C₉ alkyl and C₉ alkenyl.

[0713] In other embodiments, R' is selected from C₁₁ alkyl and C₁₁ alkenyl. In other embodiments, R' is selected from C₁₂ alkyl, C₁₂ alkenyl, C₁₃ alkyl, C₁₃ alkenyl, C₁₄ alkyl, C₁₄ alkenyl, C₁₅ alkyl, C₁₅ alkenyl, C₁₆ alkyl, C₁₆ alkenyl, C₁₇ alkyl, C₁₇ alkenyl, C₁₈ alkyl, and C₁₈ alkenyl. In certain embodiments, R' is branched (*e.g.*, decan-2-yl, undecan-3-yl, dodecan-4-yl, tridecan-5-yl, tetradecan-6-yl, 2-methylundecan-3-yl, 2-methyldecan-2-yl, 3-methylundecan-3-yl, 4-methyldodecan-4-yl or heptadeca-9-yl). In

certain embodiments, R' is .

[0714] In certain embodiments, R' is unsubstituted C₁₋₁₈ alkyl. In certain embodiments, R' is substituted C₁₋₁₈ alkyl (*e.g.*, C₁₋₁₅ alkyl substituted with a C₃₋₆ carbocycle such as 1-cyclopropylnonyl).

[0715] In some embodiments, R'' is selected from the group consisting of C₃₋₁₄ alkyl and C₃₋₁₄ alkenyl. In some embodiments, R'' is C₃ alkyl, C₄ alkyl, C₅ alkyl, C₆ alkyl, C₇ alkyl, or C₈ alkyl. In some embodiments, R'' is C₉ alkyl, C₁₀ alkyl, C₁₁ alkyl, C₁₂ alkyl, C₁₃ alkyl, or C₁₄ alkyl.

[0716] In some embodiments, M' is -C(O)O-. In some embodiments, M' is -OC(O)-.

[0717] In other embodiments, M' is an aryl group or heteroaryl group. For example, M' can be selected from the group consisting of phenyl, oxazole, and thiazole.

[0718] In some embodiments, M is -C(O)O-. In some embodiments, M is -OC(O)-. In some embodiments, M is -C(O)N(R')-. In some embodiments, M is -P(O)(OR')O-.

[0719] In other embodiments, M is an aryl group or heteroaryl group. For example, M can be selected from the group consisting of phenyl, oxazole, and thiazole.

[0720] In some embodiments, M is the same as M'. In other embodiments, M is different from M'.


- [0721] In some embodiments, each R_5 is H. In certain such embodiments, each R_6 is also H.
- [0722] In some embodiments, R_7 is H. In other embodiments, R_7 is C_{1-3} alkyl (*e.g.*, methyl, ethyl, propyl, or *i*-propyl).
- [0723] In some embodiments, R_2 and R_3 are independently C_{5-14} alkyl or C_{5-14} alkenyl.
- [0724] In some embodiments, R_2 and R_3 are the same. In some embodiments, R_2 and R_3 are C_8 alkyl. In certain embodiments, R_2 and R_3 are C_2 alkyl. In other embodiments, R_2 and R_3 are C_3 alkyl. In some embodiments, R_2 and R_3 are C_4 alkyl. In certain embodiments, R_2 and R_3 are C_5 alkyl. In other embodiments, R_2 and R_3 are C_6 alkyl. In some embodiments, R_2 and R_3 are C_7 alkyl.
- [0725] In other embodiments, R_2 and R_3 are different. In certain embodiments, R_2 is C_8 alkyl. In some embodiments, R_3 is C_{1-7} (*e.g.*, C_1 , C_2 , C_3 , C_4 , C_5 , C_6 , or C_7 alkyl) or C_9 alkyl.
- [0726] In some embodiments, R_7 and R_3 are H.
- [0727] In certain embodiments, R_2 is H.
- [0728] In some embodiments, m is 5, 7, or 9.
- [0729] In some embodiments, R_4 is selected from $-(CH_2)_nQ$ and $-(CH_2)_nCHQR$.
- [0730] In some embodiments, Q is selected from the group consisting of $-OR$, $-OH$, $-O(CH_2)_nN(R)_2$, $-OC(O)R$, $-CX_3$, $-CN$, $-N(R)C(O)R$, $-N(H)C(O)R$, $-N(R)S(O)_2R$, $-N(H)S(O)_2R$, $-N(R)C(O)N(R)_2$, $-N(H)C(O)N(R)_2$, $-N(H)C(O)N(H)(R)$, $-N(R)C(S)N(R)_2$, $-N(H)C(S)N(R)_2$, $-N(H)C(S)N(H)(R)$, $-C(R)N(R)_2C(O)OR$, a carbocycle, and a heterocycle.
- [0731] In certain embodiments, Q is $-OH$.
- [0732] In certain embodiments, Q is a substituted or unsubstituted 5- to 10- membered heteroaryl, *e.g.*, Q is an imidazole, a pyrimidine, a purine, 2-amino-1,9-dihydro-6*H*-purin-6-one-9-yl (or guanine-9-yl), adenine-9-yl, cytosine-1-yl, or uracil-1-yl. In certain embodiments, Q is a substituted 5- to 14-membered heterocycloalkyl, *e.g.*, substituted with one or more substituents selected from oxo ($=O$), OH, amino, and C_{1-3} alkyl. For example, Q is 4-methylpiperazinyl, 4-(4-methoxybenzyl)piperazinyl, or isoindolin-2-yl-1,3-dione.
- [0733] In certain embodiments, Q is an unsubstituted or substituted C_{6-10} aryl (such as phenyl) or C_{3-6} cycloalkyl.

- [0734] In some embodiments, n is 1. In other embodiments, n is 2. In further embodiments, n is 3. In certain other embodiments, n is 4. For example, R_4 can be $-(CH_2)_2OH$. For example, R_4 can be $-(CH_2)_3OH$. For example, R_4 can be $-(CH_2)_4OH$. For example, R_4 can be benzyl. For example, R_4 can be 4-methoxybenzyl.
- [0735] In some embodiments, R_4 is a C_{3-6} carbocycle. In some embodiments, R_4 is a C_{3-6} cycloalkyl. For example, R_4 can be cyclohexyl optionally substituted with *e.g.*, OH, halo, C_{1-6} alkyl, etc. For example, R_4 can be 2-hydroxycyclohexyl.
- [0736] In some embodiments, R is H.
- [0737] In some embodiments, R is unsubstituted C_{1-3} alkyl or unsubstituted C_{2-3} alkenyl. For example, R_4 can be $-CH_2CH(OH)CH_3$ or $-CH_2CH(OH)CH_2CH_3$.
- [0738] In some embodiments, R is substituted C_{1-3} alkyl, *e.g.*, CH_2OH . For example, R_4 can be $-CH_2CH(OH)CH_2OH$.
- [0739] In some embodiments, R_2 and R_3 , together with the atom to which they are attached, form a heterocycle or carbocycle. In some embodiments, R_2 and R_3 , together with the atom to which they are attached, form a 5- to 14- membered aromatic or non-aromatic heterocycle having one or more heteroatoms selected from N, O, S, and P. In some embodiments, R_2 and R_3 , together with the atom to which they are attached, form an optionally substituted C_{3-20} carbocycle (*e.g.*, C_{3-18} carbocycle, C_{3-15} carbocycle, C_{3-12} carbocycle, or C_{3-10} carbocycle), either aromatic or non-aromatic. In some embodiments, R_2 and R_3 , together with the atom to which they are attached, form a C_{3-6} carbocycle. In other embodiments, R_2 and R_3 , together with the atom to which they are attached, form a C_6 carbocycle, such as a cyclohexyl or phenyl group. In certain embodiments, the heterocycle or C_{3-6} carbocycle is substituted with one or more alkyl groups (*e.g.*, at the same ring atom or at adjacent or non-adjacent ring atoms). For example, R_2 and R_3 , together with the atom to which they are attached, can form a cyclohexyl or phenyl group bearing one or more C_5 alkyl substitutions. In certain embodiments, the heterocycle or C_{3-6} carbocycle formed by R_2 and R_3 , is substituted with a carbocycle groups. For example, R_2 and R_3 , together with the atom to which they are attached, can form a cyclohexyl or phenyl group that is substituted with cyclohexyl. In some embodiments, R_2 and R_3 , together with the atom to which they are attached, form a C_{7-15} carbocycle, such as a cycloheptyl, cyclopentadecanyl, or naphthyl group.
- [0740] In some embodiments, R_4 is selected from $-(CH_2)_nQ$ and $-(CH_2)_nCHQR$. In some embodiments, Q is selected from the group consisting of $-OR$, $-OH$, $-O(CH_2)_nN(R)_2$,


-OC(O)R, -CX₃, -CN, -N(R)C(O)R, -N(H)C(O)R, -N(R)S(O)₂R, -N(H)S(O)₂R, -N(R)C(O)N(R)₂, -N(H)C(O)N(R)₂, -N(H)C(O)N(H)(R), -N(R)C(S)N(R)₂, -N(H)C(S)N(R)₂, -N(H)C(S)N(H)(R), and a heterocycle. In other embodiments, Q is selected from the group consisting of an imidazole, a pyrimidine, and a purine.

[0741] In some embodiments, R₂ and R₃, together with the atom to which they are attached, form a heterocycle or carbocycle. In some embodiments, R₂ and R₃, together with the atom to which they are attached, form a C₃₋₆ carbocycle, such as a phenyl group. In certain embodiments, the heterocycle or C₃₋₆ carbocycle is substituted with one or more alkyl groups (*e.g.*, at the same ring atom or at adjacent or non-adjacent ring atoms). For example, R₂ and R₃, together with the atom to which they are attached, can form a phenyl group bearing one or more C₅ alkyl substitutions.


[0742] In some embodiments, the pharmaceutical compositions of the present disclosure, the compound of Formula (I) is selected from the group consisting of:




(Compound 1),




(Compound 2),




(Compound 3),




(Compound 4),




(Compound 5),




(Compound 6),




(Compound 7),




(Compound 8),




(Compound 9),




(Compound 10),




(Compound 11),




(Compound 12),




(Compound 13),




(Compound 14),




(Compound 15),




(Compound 16),




(Compound 17),




(Compound 18),




(Compound 19),




(Compound 20),




(Compound 21),




(Compound 22),




(Compound 23),




(Compound 24),




(Compound 25),




(Compound 26),




(Compound 27),




(Compound 28),




(Compound 29),




(Compound 30),




(Compound 31),




(Compound 32),




(Compound 33),




(Compound 34),




(Compound 35),




(Compound 36),




(Compound 37),




(Compound 38),




(Compound 39),




(Compound 40),




(Compound 41),




(Compound 42),




(Compound 43),




(Compound 44),




(Compound 45),




(Compound 46),




(Compound 47),




(Compound 48),




(Compound 49),




(Compound 50),




(Compound 51),




(Compound 52),




(Compound 53),




(Compound 54),




(Compound 55),




(Compound 56),




(Compound 57),




(Compound 58),




(Compound 59),




(Compound 60),




(Compound 61),




(Compound 62),




(Compound 63),




(Compound 64),




(Compound 65),




(Compound 66),




(Compound 67),




(Compound 68),




(Compound 69),




(Compound 70),




(Compound 71),




(Compound 72),




(Compound 73),




(Compound 74),




(Compound 75),




(Compound 76),




(Compound 77),




(Compound 78),




(Compound 79),




(Compound 80),




(Compound 81),




(Compound 82),




(Compound 83),




(Compound 84),




(Compound 85),




(Compound 86),




(Compound 87),




(Compound 88),




(Compound 89),




(Compound 90),




(Compound 91),




(Compound 92),




(Compound 93),




(Compound 94),




(Compound 95),




(Compound 96),




(Compound 97),




(Compound 98),




(Compound 99),




(Compound 100),




(Compound 101),




(Compound 102),




(Compound 103),




(Compound 104),




(Compound 105),




(Compound 106),




(Compound 107),




(Compound 108),




(Compound 109),




(Compound 110),




(Compound 111),




(Compound 112),




(Compound 113),




(Compound 114),




(Compound 115),




(Compound 116),




(Compound 117),




(Compound 118),




(Compound 119),




(Compound 120),




(Compound 121),




(Compound 122),




(Compound 123),




(Compound 124),




(Compound 125),




(Compound 126),




(Compound 127),




(Compound 128),




(Compound 129),




(Compound 130),




(Compound 131),




(Compound 132),




(Compound 133),




(Compound 134),




(Compound 135),




(Compound 136),




(Compound 137),




(Compound 138),




(Compound 139),




(Compound 140),




(Compound 141),




(Compound 142),




(Compound 143),




(Compound 144),




(Compound 145),




(Compound 146),




(Compound 147),




(Compound 148),




(Compound 149),




(Compound 150),




(Compound 151),




(Compound 152),




(Compound 153),




(Compound 154),




(Compound 155),




(Compound 156),




(Compound 157),




(Compound 158),




(Compound 159),




(Compound 160),




(Compound 161),




(Compound 162),




(Compound 163),




(Compound 164),




(Compound 165),




(Compound 166),




(Compound 167),




(Compound 168),




(Compound 169),




(Compound 170),




(Compound 171),




(Compound 172),




(Compound 173),




(Compound 174),




(Compound 175),




(Compound 176),




(Compound 177),




(Compound 178),




(Compound 179),




(Compound 180),




(Compound 181),




(Compound 182),




(Compound 183),




(Compound 184),




(Compound 185),




(Compound 186),




(Compound 187),




(Compound 188),




(Compound 189),




(Compound 190),




(Compound 191),




(Compound 192),




(Compound 193),




(Compound 194),




(Compound 195),




(Compound 196),




(Compound 197),




(Compound 198),




(Compound 199),




(Compound 200),




(Compound 201),




(Compound 202),




(Compound 203),




(Compound 204),




(Compound 205),




(Compound 206),




(Compound 207),




(Compound 208),




(Compound 209),




(Compound 210),




(Compound 211),




(Compound 212),




(Compound 213),




(Compound 214),




(Compound 215),




(Compound 216),




(Compound 217),




(Compound 218),




(Compound 219),




(Compound 220),




(Compound 221),




(Compound 222),




(Compound 223),




(Compound 224),




(Compound 225),




(Compound 226),




(Compound 227),




(Compound 228),




(Compound 229),



(Compound 230),



(Compound 231),




(Compound 232),

and salts or stereoisomers thereof.

[0743] In other embodiments, the compound of Formula (I) is selected from the group consisting of Compound 1-Compound 147, or salt or stereoisomers thereof.


[0744] In some embodiments ionizable lipids including a central piperazine moiety are provided.

[0745] In some embodiments, the delivery agent comprises a lipid compound having the Formula (III)



(III),

or salts or stereoisomers thereof, wherein



ring A is

t is 1 or 2;

A₁ and A₂ are each independently selected from CH or N;

Z is CH₂ or absent wherein when Z is CH₂, the dashed lines (1) and (2) each represent a single bond; and when Z is absent, the dashed lines (1) and (2) are both absent;

R_1 , R_2 , R_3 , R_4 , and R_5 are independently selected from the group consisting of C_{5-20} alkyl, C_{5-20} alkenyl, $-R''MR'$, $-R^*YR''$, $-YR''$, and $-R^*OR''$;

each M is independently selected from the group consisting of $-C(O)O-$, $-OC(O)-$, $-OC(O)O-$, $-C(O)N(R')-$, $-N(R')C(O)-$, $-C(O)-$, $-C(S)-$, $-C(S)S-$, $-SC(S)-$, $-CH(OH)-$, $-P(O)(OR')O-$, $-S(O)_2-$, an aryl group, and a heteroaryl group;

X^1 , X^2 , and X^3 are independently selected from the group consisting of a bond, $-CH_2-$, $-(CH_2)_2-$, $-CHR-$, $-CHY-$, $-C(O)-$, $-C(O)O-$, $-OC(O)-$, $-C(O)-CH_2-$, $-CH_2-C(O)-$, $-C(O)O-CH_2-$, $-OC(O)-CH_2-$, $-CH_2-C(O)O-$, $-CH_2-OC(O)-$, $-CH(OH)-$, $-C(S)-$, and $-CH(SH)-$;


each Y is independently a C_{3-6} carbocycle;


each R^* is independently selected from the group consisting of C_{1-12} alkyl and C_{2-12} alkenyl;

each R is independently selected from the group consisting of C_{1-3} alkyl and a C_{3-6} carbocycle;

each R' is independently selected from the group consisting of C_{1-12} alkyl, C_{2-12} alkenyl, and H; and

each R'' is independently selected from the group consisting of C_{3-12} alkyl and C_{3-12} alkenyl,




wherein when ring A is , then


- i) at least one of X^1 , X^2 , and X^3 is not $-CH_2-$; and/or
- ii) at least one of R_1 , R_2 , R_3 , R_4 , and R_5 is $-R''MR'$.

[0746]


In some embodiments, the compound is of any of Formulae (IIIa1)-(IIIa6):




(IIIa1),




(IIIa2),




(IIIa3),



(IIIa4),




(IIIa5), or





(IIIa6).

[0747] The compounds of Formula (III) or any of (IIIa1)-(IIIa6) include one or more of the following features when applicable.

[0748] In some embodiments, ring A is




[0749] In some embodiments, ring A is




or



[0750] In some embodiments, ring A is




[0751] In some embodiments, ring A is



[0752] In some embodiments, ring A is



[0753] In some embodiments, ring A is



wherein ring, in which the N atom is connected with X^2 .

[0754] In some embodiments, Z is CH_2 .

[0755] In some embodiments, Z is absent.

[0756] In some embodiments, at least one of A_1 and A_2 is N.

[0757] In some embodiments, each of A_1 and A_2 is N.

[0758] In some embodiments, each of A_1 and A_2 is CH.


[0759] In some embodiments, A_1 is N and A_2 is CH.

[0760] In some embodiments, A_1 is CH and A_2 is N.


[0761] In some embodiments, at least one of X^1 , X^2 , and X^3 is not $-\text{CH}_2-$. For example, in certain embodiments, X^1 is not $-\text{CH}_2-$. In some embodiments, at least one of X^1 , X^2 , and X^3 is $-\text{C}(\text{O})-$.

[0762] In some embodiments, X^2 is $-\text{C}(\text{O})-$, $-\text{C}(\text{O})\text{O}-$, $-\text{OC}(\text{O})-$, $-\text{C}(\text{O})\text{CH}_2-$, $-\text{CH}_2\text{C}(\text{O})-$, $-\text{C}(\text{O})\text{OCH}_2-$, $-\text{OC}(\text{O})\text{CH}_2-$, $-\text{CH}_2\text{C}(\text{O})\text{O}-$, or $-\text{CH}_2\text{OC}(\text{O})-$.


- [0763] In some embodiments, X^3 is $-C(O)-$, $-C(O)O-$, $-OC(O)-$, $-C(O)-CH_2-$, $-CH_2-C(O)-$, $-C(O)O-CH_2-$, $-OC(O)-CH_2-$, $-CH_2-C(O)O-$, or $-CH_2-OC(O)-$. In other embodiments, X^3 is $-CH_2-$.
- [0764] In some embodiments, X^3 is a bond or $-(CH_2)_2-$.
- [0765] In some embodiments, R_1 and R_2 are the same. In certain embodiments, R_1 , R_2 , and R_3 are the same. In some embodiments, R_4 and R_5 are the same. In certain embodiments, R_1 , R_2 , R_3 , R_4 , and R_5 are the same.
- [0766] In some embodiments, at least one of R_1 , R_2 , R_3 , R_4 , and R_5 is $-R''MR'$. In some embodiments, at most one of R_1 , R_2 , R_3 , R_4 , and R_5 is $-R''MR'$. For example, at least one of R_1 , R_2 , and R_3 may be $-R''MR'$, and/or at least one of R_4 and R_5 is $-R''MR'$. In certain embodiments, at least one M is $-C(O)O-$. In some embodiments, each M is $-C(O)O-$. In some embodiments, at least one M is $-OC(O)-$. In some embodiments, each M is $-OC(O)-$. In some embodiments, at least one M is $-OC(O)O-$. In some embodiments, each M is $-OC(O)O-$. In some embodiments, at least one R'' is C_3 alkyl. In certain embodiments, each R'' is C_3 alkyl. In some embodiments, at least one R'' is C_5 alkyl. In certain embodiments, each R'' is C_5 alkyl. In some embodiments, at least one R'' is C_6 alkyl. In certain embodiments, each R'' is C_6 alkyl. In some embodiments, at least one R'' is C_7 alkyl. In certain embodiments, each R'' is C_7 alkyl. In some embodiments, at least one R' is C_5 alkyl. In certain embodiments, each R' is C_5 alkyl. In other embodiments, at least one R' is C_1 alkyl. In certain embodiments, each R' is C_1 alkyl. In some embodiments, at least one R' is C_2 alkyl. In certain embodiments, each R' is C_2 alkyl.
- [0767] In some embodiments, at least one of R_1 , R_2 , R_3 , R_4 , and R_5 is C_{12} alkyl. In certain embodiments, each of R_1 , R_2 , R_3 , R_4 , and R_5 are C_{12} alkyl.
- [0768] In certain embodiments, the compound is selected from the group consisting of:




(Compound 233),




(Compound 234),




(Compound 235),




(Compound 236),




(Compound 237),




(Compound 238),




(Compound 239),




(Compound 240),




(Compound 241),




(Compound 242),




(Compound 243),




(Compound 244),




(Compound 245),




(Compound 246),




(Compound 247),




(Compound 248),




(Compound 274),




(Compound 275),




(Compound 276),




(Compound 277),




(Compound 278),




(Compound 279),




(Compound 280),




(Compound 281),




(Compound 282),




(Compound 283),




(Compound 284),




(Compound 285),




(Compound 286),




(Compound 287),




(Compound 288),




(Compound 289),




(Compound 290),




(Compound 291),




(Compound 292),




(Compound 293),




(Compound 294),




(Compound 295),




(Compound 296),




(Compound 297),




(Compound 298),




(Compound 300),




(Compound 301),




(Compound 302),




(Compound 303),




(Compound 304),




(Compound 305),




(Compound 306),




(Compound 307),




(Compound 308),




(Compound 310),




(Compound 311)




(Compound 312),




(Compound 313),




(Compound 314),




(Compound 315),




(Compound 316),




(Compound 317),




(Compound 318),




(Compound 319),




(Compound 320),




(Compound 321),




(Compound 322),




(Compound 323),




(Compound 324),




(Compound 325),




(Compound 326),




(Compound 327),




(Compound 328),




(Compound 329),




(Compound 330),




(Compound 331),




(Compound 332),




(Compound 333),




(Compound 334),




(Compound 335),




(Compound 336),




(Compound 337),




(Compound 338),



(Compound 339),




(Compound 340), and



(Compound 341).

[0769] In some embodiments, the delivery agent comprises Compound 236.

[0770] In some embodiments, the delivery agent comprises a compound having the Formula (IV)




(IV),


or salts or stereoisomer thereof, wherein

A_1 and A_2 are each independently selected from CH or N and at least one of A_1 and A_2 is N;

Z is CH₂ or absent wherein when Z is CH₂, the dashed lines (1) and (2) each represent a single bond; and when Z is absent, the dashed lines (1) and (2) are both absent;


R₁, R₂, R₃, R₄, and R₅ are independently selected from the group consisting of C₆₋₂₀ alkyl and C₆₋₂₀ alkenyl;



wherein when ring A is , then

- i) R₁, R₂, R₃, R₄, and R₅ are the same, wherein R₁ is not C₁₂ alkyl, C₁₈ alkyl, or C₁₈ alkenyl;
- ii) only one of R₁, R₂, R₃, R₄, and R₅ is selected from C₆₋₂₀ alkenyl;
- iii) at least one of R₁, R₂, R₃, R₄, and R₅ have a different number of carbon atoms than at least one other of R₁, R₂, R₃, R₄, and R₅;
- iv) R₁, R₂, and R₃ are selected from C₆₋₂₀ alkenyl, and R₄ and R₅ are selected from C₆₋₂₀ alkyl; or
- v) R₁, R₂, and R₃ are selected from C₆₋₂₀ alkyl, and R₄ and R₅ are selected from C₆₋₂₀ alkenyl.

[0771] In some embodiments, the compound is of Formula (IVa):



(IVa).

[0772] The compounds of Formula (IV) or (IVa) include one or more of the following features when applicable.

[0773] In some embodiments, Z is CH₂.


[0774] In some embodiments, Z is absent.

[0775] In some embodiments, at least one of A₁ and A₂ is N.


[0776] In some embodiments, each of A₁ and A₂ is N.

[0777] In some embodiments, each of A₁ and A₂ is CH.


- [0778] In some embodiments, A₁ is N and A₂ is CH.
- [0779] In some embodiments, A₁ is CH and A₂ is N.
- [0780] In some embodiments, R₁, R₂, R₃, R₄, and R₅ are the same, and are not C₁₂ alkyl, C₁₈ alkyl, or C₁₈ alkenyl. In some embodiments, R₁, R₂, R₃, R₄, and R₅ are the same and are C₉ alkyl or C₁₄ alkyl.
- [0781] In some embodiments, only one of R₁, R₂, R₃, R₄, and R₅ is selected from C₆₋₂₀ alkenyl. In certain such embodiments, R₁, R₂, R₃, R₄, and R₅ have the same number of carbon atoms. In some embodiments, R₄ is selected from C₅₋₂₀ alkenyl. For example, R₄ may be C₁₂ alkenyl or C₁₈ alkenyl.
- [0782] In some embodiments, at least one of R₁, R₂, R₃, R₄, and R₅ have a different number of carbon atoms than at least one other of R₁, R₂, R₃, R₄, and R₅.
- [0783] In certain embodiments, R₁, R₂, and R₃ are selected from C₆₋₂₀ alkenyl, and R₄ and R₅ are selected from C₆₋₂₀ alkyl. In other embodiments, R₁, R₂, and R₃ are selected from C₆₋₂₀ alkyl, and R₄ and R₅ are selected from C₆₋₂₀ alkenyl. In some embodiments, R₁, R₂, and R₃ have the same number of carbon atoms, and/or R₄ and R₅ have the same number of carbon atoms. For example, R₁, R₂, and R₃, or R₄ and R₅, may have 6, 8, 9, 12, 14, or 18 carbon atoms. In some embodiments, R₁, R₂, and R₃, or R₄ and R₅, are C₁₈ alkenyl (e.g., linoleyl). In some embodiments, R₁, R₂, and R₃, or R₄ and R₅, are alkyl groups including 6, 8, 9, 12, or 14 carbon atoms.
- [0784] In some embodiments, R₁ has a different number of carbon atoms than R₂, R₃, R₄, and R₅. In other embodiments, R₃ has a different number of carbon atoms than R₁, R₂, R₄, and R₅. In further embodiments, R₄ has a different number of carbon atoms than R₁, R₂, R₃, and R₅.
- [0785] In some embodiments, the compound is selected from the group consisting of:




(Compound 249),




(Compound 250),




(Compound 251),




(Compound 252),




(Compound 253),




(Compound 254),




(Compound 255),




(Compound 256),




(Compound 257),




(Compound 258),




(Compound 259),




(Compound 260),




(Compound 261),




(Compound 262),




(Compound 263),



(Compound 264),




(Compound 265), and



(Compound 266).

[0786] In other embodiments, the delivery agent comprises a compound having the Formula (V)



(V),

or salts or stereoisomers thereof, in which

A₃ is CH or N;

A₄ is CH₂ or NH; and at least one of A₃ and A₄ is N or NH;

Z is CH₂ or absent wherein when Z is CH₂, the dashed lines (1) and (2) each represent a single bond; and when Z is absent, the dashed lines (1) and (2) are both absent;

R₁, R₂, and R₃ are independently selected from the group consisting of C₅₋₂₀ alkyl, C₅₋₂₀ alkenyl, -R''MR', -R*YR'', -YR'', and -R*OR'';

each M is independently selected from -C(O)O-, -OC(O)-, -C(O)N(R')-, -N(R')C(O)-, -C(O)-, -C(S)-, -C(S)S-, -SC(S)-, -CH(OH)-, -P(O)(OR')O-, -S(O)₂-, an aryl group, and a heteroaryl group;

X¹ and X² are independently selected from the group consisting of -CH₂-, -(CH₂)₂-, -CHR-, -CHY-, -C(O)-, -C(O)O-, -OC(O)-, -C(O)-CH₂-, -CH₂-C(O)-, -C(O)O-CH₂-, -OC(O)-CH₂-, -CH₂-C(O)O-, -CH₂-OC(O)-, -CH(OH)-, -C(S)-, and -CH(SH)-;

each Y is independently a C₃₋₆ carbocycle;


each R* is independently selected from the group consisting of C₁₋₁₂ alkyl and C₂₋₁₂ alkenyl;

each R is independently selected from the group consisting of C₁₋₃ alkyl and a C₃₋₆ carbocycle;

each R' is independently selected from the group consisting of C₁₋₁₂ alkyl, C₂₋₁₂ alkenyl, and H; and


each R'' is independently selected from the group consisting of C₃₋₁₂ alkyl and C₃₋₁₂ alkenyl.

[0787] In some embodiments, the compound is of Formula (Va):




(Va).


- [0788] The compounds of Formula (V) or (Va) include one or more of the following features when applicable.
- [0789] In some embodiments, Z is CH₂.
- [0790] In some embodiments, Z is absent.
- [0791] In some embodiments, at least one of A₃ and A₄ is N or NH.
- [0792] In some embodiments, A₃ is N and A₄ is NH.
- [0793] In some embodiments, A₃ is N and A₄ is CH₂.
- [0794] In some embodiments, A₃ is CH and A₄ is NH.
- [0795] In some embodiments, at least one of X¹ and X² is not -CH₂-. For example, in certain embodiments, X¹ is not -CH₂-. In some embodiments, at least one of X¹ and X² is -C(O)-.
- [0796] In some embodiments, X² is -C(O)-, -C(O)O-, -OC(O)-, -C(O)-CH₂-, -CH₂-C(O)-, -C(O)O-CH₂-, -OC(O)-CH₂-, -CH₂-C(O)O-, or -CH₂-OC(O)-.
- [0797] In some embodiments, R₁, R₂, and R₃ are independently selected from the group consisting of C₅₋₂₀ alkyl and C₅₋₂₀ alkenyl. In some embodiments, R₁, R₂, and R₃ are the same. In certain embodiments, R₁, R₂, and R₃ are C₆, C₉, C₁₂, or C₁₄ alkyl. In other embodiments, R₁, R₂, and R₃ are C₁₈ alkenyl. For example, R₁, R₂, and R₃ may be linoleyl.
- [0798] In some embodiments, the compound is selected from the group consisting of:




(Compound 267),




(Compound 268)




(Compound 269),




(Compound 270),




(Compound 271),



(Compound 272),




(Compound 273), and



(Compound 309).

[0799] In other embodiments, the delivery agent comprises a compound having the Formula (VI):



(VI),

or salts or stereoisomers thereof, in which

A_6 and A_7 are each independently selected from CH or N, wherein at least one of A_6 and A_7 is N;

Z is CH_2 or absent wherein when Z is CH_2 , the dashed lines (1) and (2) each represent a single bond; and when Z is absent, the dashed lines (1) and (2) are both absent;

X^4 and X^5 are independently selected from the group consisting of $-\text{CH}_2-$, $-(\text{CH}_2)_2-$, $-\text{CHR}-$, $-\text{CHY}-$, $-\text{C(O)}-$, $-\text{C(O)O}-$, $-\text{OC(O)}-$, $-\text{C(O)-CH}_2-$, $-\text{CH}_2-\text{C(O)}-$, $-\text{C(O)O-CH}_2-$, $-\text{OC(O)-CH}_2-$, $-\text{CH}_2-\text{C(O)O}-$, $-\text{CH}_2-\text{OC(O)}-$, $-\text{CH(OH)}-$, $-\text{C(S)}-$, and $-\text{CH(SH)}-$;

R_1 , R_2 , R_3 , R_4 , and R_5 each are independently selected from the group consisting of C_{5-20} alkyl, C_{5-20} alkenyl, $-\text{R}''\text{MR}'$, $-\text{R}^*\text{YR}''$, $-\text{YR}''$, and $-\text{R}^*\text{OR}''$;

each M is independently selected from the group consisting of $-\text{C(O)O}-$, $-\text{OC(O)}-$, $-\text{C(O)N(R}')$, $-\text{N(R}')C(O)-$, $-\text{C(O)}-$, $-\text{C(S)}-$, $-\text{C(S)S}-$, $-\text{SC(S)}-$, $-\text{CH(OH)}-$, $-\text{P(O)(OR}')O}-$, $-\text{S(O)}_2-$, an aryl group, and a heteroaryl group;

each Y is independently a C_{3-6} carbocycle;

each R* is independently selected from the group consisting of C₁₋₁₂ alkyl and C₂₋₁₂ alkenyl;

each R is independently selected from the group consisting of C₁₋₃ alkyl and a C₃₋₆ carbocycle;

each R' is independently selected from the group consisting of C₁₋₁₂ alkyl, C₂₋₁₂ alkenyl, and H; and

each R'' is independently selected from the group consisting of C₃₋₁₂ alkyl and C₃₋₁₂ alkenyl.

[0800] In some embodiments, R₁, R₂, R₃, R₄, and R₅ each are independently selected from the group consisting of C₆₋₂₀ alkyl and C₆₋₂₀ alkenyl.

[0801] In some embodiments, R₁ and R₂ are the same. In certain embodiments, R₁, R₂, and R₃ are the same. In some embodiments, R₄ and R₅ are the same. In certain embodiments, R₁, R₂, R₃, R₄, and R₅ are the same.

[0802] In some embodiments, at least one of R₁, R₂, R₃, R₄, and R₅ is C₉₋₁₂ alkyl. In certain embodiments, each of R₁, R₂, R₃, R₄, and R₅ independently is C₉, C₁₂ or C₁₄ alkyl. In certain embodiments, each of R₁, R₂, R₃, R₄, and R₅ is C₉ alkyl.


[0803] In some embodiments, A₆ is N and A₇ is N. In some embodiments, A₆ is CH and A₇ is N.

[0804] In some embodiments, X⁴ is -CH₂- and X⁵ is -C(O)-. In some embodiments, X⁴ and X⁵ are -C(O)-.

[0805] In some embodiments, when A₆ is N and A₇ is N, at least one of X⁴ and X⁵ is not -CH₂-, e.g., at least one of X⁴ and X⁵ is -C(O)-. In some embodiments, when A₆ is N and A₇ is N, at least one of R₁, R₂, R₃, R₄, and R₅ is -R''MR'.


[0806] In some embodiments, at least one of R₁, R₂, R₃, R₄, and R₅ is not -R''MR'.

[0807] In some embodiments, the compound is



(Compound 299).

[0808] In other embodiments, the delivery agent comprises a compound having the Formula:



(Compound 342).

- [0809] Amine moieties of the lipid compounds disclosed herein can be protonated under certain conditions. For example, the central amine moiety of a lipid according to Formula (I) is typically protonated (i.e., positively charged) at a pH below the pKa of the amino moiety and is substantially not charged at a pH above the pKa. Such lipids can be referred to ionizable amino lipids.
- [0810] In one specific embodiment, ionizable amino lipid is Compound 18. In another embodiment, the ionizable amino lipid is Compound 236.
- [0811] In some embodiments, the amount the ionizable amino lipid, e.g., compound of Formula (I), ranges from about 1 mol % to 99 mol % in the lipid composition.
- [0812] In one embodiment, the amount of the ionizable amino lipid, e.g., the compound of Formula (I), is at least about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, or 99 mol % in the lipid composition.
- [0813] In one embodiment, the amount of the ionizable amino lipid, e.g., the compound of Formula (I) ranges from about 30 mol % to about 70 mol %, from about 35 mol % to about 65 mol %, from about 40 mol % to about 60 mol %, and from about 45 mol % to about 55 mol % in the lipid composition.
- [0814] In one specific embodiment, the amount of the ionizable amino lipid, e.g., the compound of Formula (I), is about 50 mol % in the lipid composition.
- [0815] In addition to the ionizable amino lipid disclosed herein, e.g., compound of Formula (I), the lipid composition of the pharmaceutical compositions disclosed herein can comprise additional components such as phospholipids, structural lipids, PEG-lipids, and any combination thereof.

b. Additional Components in the Lipid Composition

(i) **Phospholipids**

[0816] The lipid composition of the pharmaceutical composition disclosed herein can comprise one or more phospholipids, for example, one or more saturated or (poly)unsaturated phospholipids or a combination thereof. In general, phospholipids comprise a phospholipid moiety and one or more fatty acid moieties.

[0817] A phospholipid moiety can be selected, for example, from the non-limiting group consisting of phosphatidyl choline, phosphatidyl ethanolamine, phosphatidyl glycerol, phosphatidyl serine, phosphatidic acid, 2-lysophosphatidyl choline, and a sphingomyelin.

[0818] A fatty acid moiety can be selected, for example, from the non-limiting group consisting of lauric acid, myristic acid, myristoleic acid, palmitic acid, palmitoleic acid, stearic acid, oleic acid, linoleic acid, alpha-linolenic acid, erucic acid, phytanoic acid, arachidic acid, arachidonic acid, eicosapentaenoic acid, behenic acid, docosapentaenoic acid, and docosahexaenoic acid.


[0819] Particular phospholipids can facilitate fusion to a membrane. For example, a cationic phospholipid can interact with one or more negatively charged phospholipids of a membrane (e.g., a cellular or intracellular membrane). Fusion of a phospholipid to a membrane can allow one or more elements (e.g., a therapeutic agent) of a lipid-containing composition (e.g., LNPs) to pass through the membrane permitting, e.g., delivery of the one or more elements to a target tissue.


[0820] Non-natural phospholipid species including natural species with modifications and substitutions including branching, oxidation, cyclization, and alkynes are also contemplated. For example, a phospholipid can be functionalized with or cross-linked to one or more alkynes (e.g., an alkenyl group in which one or more double bonds is replaced with a triple bond). Under appropriate reaction conditions, an alkyne group can undergo a copper-catalyzed cycloaddition upon exposure to an azide. Such reactions can be useful in functionalizing a lipid bilayer of a nanoparticle composition to facilitate membrane permeation or cellular recognition or in conjugating a nanoparticle composition to a useful component such as a targeting or imaging moiety (e.g., a dye).

[0821] Phospholipids include, but are not limited to, glycerophospholipids such as phosphatidylcholines, phosphatidylethanolamines, phosphatidylserines,

phosphatidylinositols, phosphatidyl glycerols, and phosphatidic acids. Phospholipids also include phosphosphingolipid, such as sphingomyelin.

[0822] Examples of phospholipids include, but are not limited to, the following:





[0823] In certain embodiments, a phospholipid useful or potentially useful in the present invention is an analog or variant of DSPC. In certain embodiments, a phospholipid useful or potentially useful in the present invention is a compound of Formula (IX):




or a salt thereof, wherein:

each R¹ is independently optionally substituted alkyl; or optionally two R¹ are joined together with the intervening atoms to form optionally substituted monocyclic carbocyclyl or optionally substituted monocyclic heterocyclyl; or optionally three R¹ are joined together with the intervening atoms to form optionally substituted bicyclic carbocyclyl or optionally substituted bicyclic heterocyclyl;

n is 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10;

m is 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10;



each instance of L^2 is independently a bond or optionally substituted C_{1-6} alkylene, wherein one methylene unit of the optionally substituted C_{1-6} alkylene is optionally replaced with $-O-$, $-N(R^N)-$, $-S-$, $-C(O)-$, $-C(O)N(R^N)-$, $-NR^NC(O)-$, $-C(O)O-$, $-OC(O)-$, $-OC(O)O-$, $-OC(O)N(R^N)-$, $-NR^NC(O)O-$, or $-NR^NC(O)N(R^N)-$;


each instance of R^2 is independently optionally substituted C_{1-30} alkyl, optionally substituted C_{1-30} alkenyl, or optionally substituted C_{1-30} alkynyl; optionally wherein one or more methylene units of R^2 are independently replaced with optionally substituted carbocyclylene, optionally substituted heterocyclylene, optionally substituted arylene, optionally substituted heteroarylene, $-N(R^N)-$, $-O-$, $-S-$, $-C(O)-$, $-C(O)N(R^N)-$, $-NR^NC(O)-$, $-NR^NC(O)N(R^N)-$, $-C(O)O-$, $-OC(O)-$, $-OC(O)O-$, $-OC(O)N(R^N)-$, $-NR^NC(O)O-$, $-C(O)S-$, $-SC(O)-$, $-C(=NR^N)-$, $-C(=NR^N)N(R^N)-$, $-NR^NC(=NR^N)-$, $-NR^NC(=NR^N)N(R^N)-$, $-C(S)-$, $-C(S)N(R^N)-$, $-NR^NC(S)-$, $-NR^NC(S)N(R^N)-$, $-S(O)-$, $-OS(O)-$, $-S(O)O-$, $-OS(O)O-$, $-OS(O)_2-$, $-S(O)_2O-$, $-OS(O)_2O-$, $-N(R^N)S(O)-$, $-S(O)N(R^N)-$, $-N(R^N)S(O)N(R^N)-$, $-OS(O)N(R^N)-$, $-N(R^N)S(O)O-$, $-S(O)_2-$, $-N(R^N)S(O)_2-$, $-S(O)_2N(R^N)-$, $-N(R^N)S(O)_2N(R^N)-$, $-OS(O)_2N(R^N)-$, or $-N(R^N)S(O)_2O-$;

each instance of R^N is independently hydrogen, optionally substituted alkyl, or a nitrogen protecting group;

Ring B is optionally substituted carbocyclyl, optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl; and

p is 1 or 2;

provided that the compound is not of the formula:




wherein each instance of R^2 is independently unsubstituted alkyl, unsubstituted alkenyl, or unsubstituted alkynyl.

Phospholipid Head Modifications

[0824] In certain embodiments, a phospholipid useful or potentially useful in the present invention comprises a modified phospholipid head (*e.g.*, a modified choline group). In

certain embodiments, a phospholipid with a modified head is DSPC, or analog thereof, with a modified quaternary amine. For example, in embodiments of Formula (IX), at least one of R^1 is not methyl. In certain embodiments, at least one of R^1 is not hydrogen or methyl. In certain embodiments, the compound of Formula (IX) is of one of the following formulae:




or a salt thereof, wherein:

each t is independently 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10;

each u is independently 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10; and


each v is independently 1, 2, or 3.

[0825] In certain embodiments, the compound of Formula (IX) is of one of the following formulae:




or a salt thereof.


[0826] In certain embodiments, a compound of Formula (IX) is one of the following:




(Compound 400)




(Compound 401)




(Compound 402)




(Compound 403)




(Compound 404)




(Compound 405)




(Compound 406)



(Compound 407)




(Compound 408)



(Compound 409),

or a salt thereof.


[0827] In certain embodiments, a compound of Formula (IX) is of Formula (IX-a):




(IX-a),

or a salt thereof.

[0828] In certain embodiments, phospholipids useful or potentially useful in the present invention comprise a modified core. In certain embodiments, a phospholipid with a modified core described herein is DSPC, or analog thereof, with a modified core structure. For example, in certain embodiments of Formula (IX-a), group A is not of the following formula:




[0829] In certain embodiments, the compound of Formula (IX-a) is of one of the following formulae:




or a salt thereof.

[0830] In certain embodiments, a compound of Formula (IX) is one of the following:




or salts thereof.

[0831] In certain embodiments, a phospholipid useful or potentially useful in the present invention comprises a cyclic moiety in place of the glyceride moiety. In certain embodiments, a phospholipid useful in the present invention is DSPC, or analog thereof, with a cyclic moiety in place of the glyceride moiety. In certain embodiments, the compound of Formula (IX) is of Formula (IX-b):



or a salt thereof.


[0832] In certain embodiments, the compound of Formula (IX-b) is of Formula (IX-b-1):



or a salt thereof, wherein:


w is 0, 1, 2, or 3.

[0833] In certain embodiments, the compound of Formula (IX-b) is of Formula (IX-b-2):




or a salt thereof.

[0834] In certain embodiments, the compound of Formula (IX-b) is of Formula (IX-b-3):




or a salt thereof.

[0835] In certain embodiments, the compound of Formula (IX-b) is of Formula (IX-b-4):



or a salt thereof.

[0836] In certain embodiments, the compound of Formula (IX-b) is one of the following:




or salts thereof.

Phospholipid Tail Modifications

[0837] In certain embodiments, a phospholipid useful or potentially useful in the present invention comprises a modified tail. In certain embodiments, a phospholipid useful or potentially useful in the present invention is DSPC, or analog thereof, with a modified tail. As described herein, a “modified tail” may be a tail with shorter or longer aliphatic

chains, aliphatic chains with branching introduced, aliphatic chains with substituents introduced, aliphatic chains wherein one or more methylenes are replaced by cyclic or heteroatom groups, or any combination thereof. For example, in certain embodiments, the compound of (IX) is of Formula (IX-a), or a salt thereof, wherein at least one instance of R^2 is each instance of R^2 is optionally substituted C_{1-30} alkyl, wherein one or more methylene units of R^2 are independently replaced with optionally substituted carbocyclylene, optionally substituted heterocyclylene, optionally substituted arylene, optionally substituted heteroarylene, $-N(R^N)-$, $-O-$, $-S-$, $-C(O)-$, $-C(O)N(R^N)-$, $-NR^NC(O)-$, $-NR^NC(O)N(R^N)-$, $-C(O)O-$, $-OC(O)-$, $-OC(O)O-$, $-OC(O)N(R^N)-$, $-NR^NC(O)O-$, $-C(O)S-$, $-SC(O)-$, $-C(=NR^N)-$, $-C(=NR^N)N(R^N)-$, $-NR^NC(=NR^N)-$, $-NR^NC(=NR^N)N(R^N)-$, $-C(S)-$, $-C(S)N(R^N)-$, $-NR^NC(S)-$, $-NR^NC(S)N(R^N)-$, $-S(O)-$, $-OS(O)-$, $-S(O)O-$, $-OS(O)O-$, $-OS(O)_2-$, $-S(O)_2O-$, $-OS(O)_2O-$, $-N(R^N)S(O)-$, $-S(O)N(R^N)-$, $-N(R^N)S(O)N(R^N)-$, $-OS(O)N(R^N)-$, $-N(R^N)S(O)O-$, $-S(O)_2-$, $-N(R^N)S(O)_2-$, $-S(O)_2N(R^N)-$, $-N(R^N)S(O)_2N(R^N)-$, $-OS(O)_2N(R^N)-$, or $-N(R^N)S(O)_2O-$.

[0838] In certain embodiments, the compound of Formula (IX) is of Formula (IX-c):




(IX-c),

or a salt thereof, wherein:

each x is independently an integer between 0-30, inclusive; and

each instance is G is independently selected from the group consisting of optionally substituted carbocyclylene, optionally substituted heterocyclylene, optionally substituted arylene, optionally substituted heteroarylene, $-N(R^N)-$, $-O-$, $-S-$, $-C(O)-$, $-C(O)N(R^N)-$, $-NR^NC(O)-$, $-NR^NC(O)N(R^N)-$, $-C(O)O-$, $-OC(O)-$, $-OC(O)O-$, $-OC(O)N(R^N)-$, $-NR^NC(O)O-$, $-C(O)S-$, $-SC(O)-$, $-C(=NR^N)-$, $-C(=NR^N)N(R^N)-$, $-NR^NC(=NR^N)-$, $-NR^NC(=NR^N)N(R^N)-$, $-C(S)-$, $-C(S)N(R^N)-$, $-NR^NC(S)-$, $-NR^NC(S)N(R^N)-$, $-S(O)-$, $-OS(O)-$, $-S(O)O-$, $-OS(O)O-$, $-OS(O)_2-$, $-S(O)_2O-$, $-OS(O)_2O-$, $-N(R^N)S(O)-$, $-S(O)N(R^N)-$, $-N(R^N)S(O)N(R^N)-$, $-OS(O)N(R^N)-$, $-N(R^N)S(O)O-$, $-S(O)_2-$, $-N(R^N)S(O)_2-$, $-S(O)_2N(R^N)-$, $-N(R^N)S(O)_2N(R^N)-$, $-OS(O)_2N(R^N)-$, or $-N(R^N)S(O)_2O-$. Each possibility represents a separate embodiment of the present invention.

[0839] In certain embodiments, the compound of Formula (IX-c) is of Formula (IX-c-1):




(IX-c-1),

or salt thereof, wherein:

each instance of v is independently 1, 2, or 3.


[0840] In certain embodiments, the compound of Formula (IX-c) is of Formula (IX-c-2):



(IX-c-2),


or a salt thereof.

[0841] In certain embodiments, the compound of Formula (IX-c) is of the following formula:




or a salt thereof.

[0842] In certain embodiments, the compound of Formula (IX-c) is the following:



or a salt thereof.


[0843] In certain embodiments, the compound of Formula (IX-c) is of Formula (IX-c-3):



(IX-c-3),


or a salt thereof.

[0844] In certain embodiments, the compound of Formula (IX-c) is of the following formulae:




or a salt thereof.

[0845] In certain embodiments, the compound of Formula (IX-c) is the following:




or a salt thereof.


[0846] In certain embodiments, a phospholipid useful or potentially useful in the present invention comprises a modified phosphocholine moiety, wherein the alkyl chain linking the quaternary amine to the phosphoryl group is not ethylene (*e.g.*, *n* is not 2). Therefore, in certain embodiments, a phospholipid useful or potentially useful in the present invention is a compound of Formula (IX), wherein *n* is 1, 3, 4, 5, 6, 7, 8, 9, or 10. For example, in certain embodiments, a compound of Formula (IX) is of one of the following formulae:




or a salt thereof.

[0847] In certain embodiments, a compound of Formula (IX) is one of the following:






(Compound 412)



(Compound 413)





(Compound 414),

or salts thereof.

(ii) Alternative lipids

[0848] In certain embodiments, an alternative lipid is used in place of a phospholipid of the invention. Non-limiting examples of such alternative lipids include the following:






(iii) Structural Lipids

[0849] The lipid composition of a pharmaceutical composition disclosed herein can comprise one or more structural lipids. As used herein, the term “structural lipid” refers to sterols and also to lipids containing sterol moieties.

[0850] Incorporation of structural lipids in the lipid nanoparticle may help mitigate aggregation of other lipids in the particle. Structural lipids can be selected from the group including but not limited to, cholesterol, fecosterol, sitosterol, ergosterol, campesterol, stigmasterol, brassicasterol, tomatidine, tomatine, ursolic acid, alpha-tocopherol, hopanoids, phytosterols, steroids, and mixtures thereof. In some embodiments, the structural lipid is a sterol. As defined herein, “sterols” are a subgroup of steroids consisting of steroid alcohols. In certain embodiments, the structural lipid is a steroid. In certain embodiments, the structural lipid is cholesterol. In certain embodiments, the structural lipid is an analog of cholesterol. In certain embodiments, the structural lipid is alpha-tocopherol. Examples of structural lipids include, but are not limited to, the following:



[0851] In one embodiment, the amount of the structural lipid (e.g., an sterol such as cholesterol) in the lipid composition of a pharmaceutical composition disclosed herein ranges from about 20 mol % to about 60 mol %, from about 25 mol % to about 55 mol %, from about 30 mol % to about 50 mol %, or from about 35 mol % to about 45 mol %.

[0852] In one embodiment, the amount of the structural lipid (e.g., an sterol such as cholesterol) in the lipid composition disclosed herein ranges from about 25 mol % to about 30 mol %, from about 30 mol % to about 35 mol %, or from about 35 mol % to about 40 mol %.

[0853] In one embodiment, the amount of the structural lipid (e.g., a sterol such as cholesterol) in the lipid composition disclosed herein is about 24 mol %, about 29 mol %, about 34 mol %, or about 39 mol %.

[0854] In some embodiments, the amount of the structural lipid (e.g., an sterol such as cholesterol) in the lipid composition disclosed herein is at least about 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, or 60 mol %.

(iv) Polyethylene Glycol (PEG)-Lipids

[0855] The lipid composition of a pharmaceutical composition disclosed herein can comprise one or more a polyethylene glycol (PEG) lipid.

[0856] As used herein, the term “PEG-lipid” refers to polyethylene glycol (PEG)-modified lipids. Non-limiting examples of PEG-lipids include PEG-modified phosphatidylethanolamine and phosphatidic acid, PEG-ceramide conjugates (e.g., PEG-CerC14 or PEG-CerC20), PEG-modified dialkylamines and PEG-modified 1,2-diacyloxypropan-3-amines. Such lipids are also referred to as PEGylated lipids. For example, a PEG lipid can be PEG-c-DOMG, PEG-DMG, PEG-DLPE, PEG-DMPE, PEG-DPPC, or a PEG-DSPE lipid.

[0857] In some embodiments, the PEG-lipid includes, but not limited to 1,2-dimyristoyl-sn-glycerol methoxypolyethylene glycol (PEG-DMG), 1,2-distearoyl-sn-glycero-3-phosphoethanolamine-N-[amino(polyethylene glycol)] (PEG-DSPE), PEG-disteryl glycerol (PEG-DSG), PEG-dipalmetoleyl, PEG-dioleyl, PEG-distearyl, PEG-diacylglycamide (PEG-DAG), PEG-dipalmitoyl phosphatidylethanolamine (PEG-DPPE), or PEG-1,2-dimyristyloxypropyl-3-amine (PEG-c-DMA).

[0858] In one embodiment, the PEG-lipid is selected from the group consisting of a PEG-modified phosphatidylethanolamine, a PEG-modified phosphatidic acid, a PEG-modified ceramide, a PEG-modified dialkylamine, a PEG-modified diacylglycerol, a PEG-modified dialkylglycerol, and mixtures thereof.

[0859] In some embodiments, the lipid moiety of the PEG-lipids includes those having lengths of from about C₁₄ to about C₂₂, preferably from about C₁₄ to about C₁₆. In some embodiments, a PEG moiety, for example an mPEG-NH₂, has a size of about 1000, 2000, 5000, 10,000, 15,000 or 20,000 Daltons. In one embodiment, the PEG-lipid is PEG_{2k}-DMG.


[0860] In one embodiment, the lipid nanoparticles described herein can comprise a PEG lipid which is a non-diffusible PEG. Non-limiting examples of non-diffusible PEGs include PEG-DSG and PEG-DSPE.

[0861] PEG-lipids are known in the art, such as those described in U.S. Patent No. 8158601 and International Publ. No. WO 2015/130584 A2, which are incorporated herein by reference in their entirety.

[0862] In general, some of the other lipid components (e.g., PEG lipids) of various formulae, described herein may be synthesized as described International Patent Application No. PCT/US2016/000129, filed December 10, 2016, entitled “Compositions and Methods for Delivery of Therapeutic Agents,” which is incorporated by reference in its entirety.

[0863] The lipid component of a lipid nanoparticle composition may include one or more molecules comprising polyethylene glycol, such as PEG or PEG-modified lipids. Such species may be alternately referred to as PEGylated lipids. A PEG lipid is a lipid modified with polyethylene glycol. A PEG lipid may be selected from the non-limiting group including PEG-modified phosphatidylethanolamines, PEG-modified phosphatidic acids, PEG-modified ceramides, PEG-modified dialkylamines, PEG-modified diacylglycerols, PEG-modified dialkylglycerols, and mixtures thereof. For example, a PEG lipid may be PEG-c-DOMG, PEG-DMG, PEG-DLPE, PEG-DMPE, PEG-DPPC, or a PEG-DSPE lipid.

[0864] In some embodiments the PEG-modified lipids are a modified form of PEG DMG. PEG-DMG has the following structure:



[0865] In one embodiment, PEG lipids useful in the present invention can be PEGylated lipids described in International Publication No. WO2012099755, the contents of which is herein incorporated by reference in its entirety. Any of these exemplary PEG lipids described herein may be modified to comprise a hydroxyl group on the PEG chain. In certain embodiments, the PEG lipid is a PEG-OH lipid. As generally defined herein, a “PEG-OH lipid” (also referred to herein as “hydroxy-PEGylated lipid”) is a PEGylated lipid having one or more hydroxyl (–OH) groups on the lipid. In certain embodiments, the PEG-OH lipid includes one or more hydroxyl groups on the PEG chain. In certain embodiments, a PEG-OH or hydroxy-PEGylated lipid comprises an –OH group at the terminus of the PEG chain. Each possibility represents a separate embodiment of the present invention.

[0866] In certain embodiments, a PEG lipid useful in the present invention is a compound of Formula (VII). Provided herein are compounds of Formula (VII):



or salts thereof, wherein:

R^3 is ---OR^O ;


R^O is hydrogen, optionally substituted alkyl, or an oxygen protecting group;

r is an integer between 1 and 100, inclusive;

L^1 is optionally substituted C_{1-10} alkylene, wherein at least one methylene of the optionally substituted C_{1-10} alkylene is independently replaced with optionally substituted carbocyclylene, optionally substituted heterocyclylene, optionally substituted arylene, optionally substituted heteroarylene, ---O--- , $\text{---N(R}^N\text{)---}$, ---S--- , ---C(O)--- , $\text{---C(O)N(R}^N\text{)---}$, $\text{---NR}^N\text{C(O)---}$, ---C(O)O--- , ---OC(O)--- , ---OC(O)O--- , $\text{---OC(O)N(R}^N\text{)---}$, $\text{---NR}^N\text{C(O)O---}$, or $\text{---NR}^N\text{C(O)N(R}^N\text{)---}$;

D is a moiety obtained by click chemistry or a moiety cleavable under physiological conditions;

m is 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10;



each instance of L^2 is independently a bond or optionally substituted C_{1-6} alkylene, wherein one methylene unit of the optionally substituted C_{1-6} alkylene is optionally replaced with ---O--- , $\text{---N(R}^N\text{)---}$, ---S--- , ---C(O)--- , $\text{---C(O)N(R}^N\text{)---}$, $\text{---NR}^N\text{C(O)---}$, ---C(O)O--- , ---OC(O)--- , ---OC(O)O--- , $\text{---OC(O)N(R}^N\text{)---}$, $\text{---NR}^N\text{C(O)O---}$, or $\text{---NR}^N\text{C(O)N(R}^N\text{)---}$;

each instance of R^2 is independently optionally substituted C_{1-30} alkyl, optionally substituted C_{1-30} alkenyl, or optionally substituted C_{1-30} alkynyl; optionally wherein one or more methylene units of R^2 are independently replaced with optionally substituted carbocyclylene, optionally substituted heterocyclylene, optionally substituted arylene, optionally substituted heteroarylene, $\text{---N(R}^N\text{)---}$, ---O--- , ---S--- , ---C(O)--- , $\text{---C(O)N(R}^N\text{)---}$, $\text{---NR}^N\text{C(O)---}$, $\text{---NR}^N\text{C(O)N(R}^N\text{)---}$, ---C(O)O--- , ---OC(O)--- , ---OC(O)O--- , $\text{---OC(O)N(R}^N\text{)---}$, $\text{---NR}^N\text{C(O)O---}$, ---C(O)S--- , ---SC(O)--- , $\text{---C(=NR}^N\text{)---}$, $\text{---C(=NR}^N\text{)N(R}^N\text{)---}$, $\text{---NR}^N\text{C(=NR}^N\text{)---}$, $\text{---NR}^N\text{C(=NR}^N\text{)N(R}^N\text{)---}$, ---C(S)--- , $\text{---C(S)N(R}^N\text{)---}$, $\text{---NR}^N\text{C(S)---}$, $\text{---NR}^N\text{C(S)N(R}^N\text{)---}$, ---S(O)--- , ---OS(O)--- , ---S(O)O--- , ---OS(O)O--- , $\text{---OS(O)}_2\text{---}$, $\text{---S(O)}_2\text{O---}$, $\text{---OS(O)}_2\text{O---}$, $\text{---N(R}^N\text{)S(O)---}$, $\text{---S(O)N(R}^N\text{)---}$, $\text{---N(R}^N\text{)S(O)N(R}^N\text{)---}$, $\text{---OS(O)N(R}^N\text{)---}$, $\text{---N(R}^N\text{)S(O)O---}$, $\text{---S(O)}_2\text{---}$,

$-N(R^N)S(O)_2-$, $-S(O)_2N(R^N)-$, $-N(R^N)S(O)_2N(R^N)-$, $-OS(O)_2N(R^N)-$, or $-N(R^N)S(O)_2O-$;

each instance of R^N is independently hydrogen, optionally substituted alkyl, or a nitrogen protecting group;

Ring B is optionally substituted carbocyclyl, optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl; and


p is 1 or 2.

[0867] In certain embodiments, the compound of Formula (VII) is a PEG-OH lipid (*i.e.*, R^3 is $-OR^O$, and R^O is hydrogen). In certain embodiments, the compound of Formula (VII) is of Formula (VII-OH):




or a salt thereof.

[0868] In certain embodiments, D is a moiety obtained by click chemistry (*e.g.*, triazole). In certain embodiments, the compound of Formula (VII) is of Formula (VII-a-1) or (VII-a-2):



or a salt thereof.


[0869] In certain embodiments, the compound of Formula (VII) is of one of the following formulae:



or a salt thereof, wherein


s is 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10.

[0870] In certain embodiments, the compound of Formula (VII) is of one of the following formulae:




or a salt thereof.

[0871] In certain embodiments, a compound of Formula (VII) is of one of the following formulae:




or a salt thereof.


[0872] In certain embodiments, a compound of Formula (VII) is of one of the following formulae:




(Compound 415),



(Compound 416),




(Compound 417),




(Compound 418),

or a salt thereof.

[0873] In certain embodiments, D is a moiety cleavable under physiological conditions (*e.g.*, ester, amide, carbonate, carbamate, urea). In certain embodiments, a compound of Formula (VII) is of Formula (VII-b-1) or (VII-b-2):




(VII-b-1)




(VII-b-2),

or a salt thereof.

[0874] In certain embodiments, a compound of Formula (VII) is of Formula (VII-b-1-OH) or (VII-b-2-OH):




(VII-b-1-OH)



(VII-b-2-OH),


or a salt thereof.

[0875] In certain embodiments, the compound of Formula (VII) is of one of the following formulae:




or a salt thereof.

[0876] In certain embodiments, a compound of Formula (VII) is of one of the following formulae:




or a salt thereof.

[0877] In certain embodiments, a compound of Formula (VII) is of one of the following formulae:



or a salt thereof.

[0878] In certain embodiments, a compound of Formula (VII) is of one of the following formulae:



or salts thereof.

[0879] In certain embodiments, a PEG lipid useful in the present invention is a PEGylated fatty acid. In certain embodiments, a PEG lipid useful in the present invention is a compound of Formula (VIII). Provided herein are compounds of Formula (VIII):



or a salts thereof, wherein:

R^3 is $-\text{OR}^{\text{O}}$;


R^{O} is hydrogen, optionally substituted alkyl or an oxygen protecting group;

r is an integer between 1 and 100, inclusive;

R^5 is optionally substituted C_{10-40} alkyl, optionally substituted C_{10-40} alkenyl, or optionally substituted C_{10-40} alkynyl; and optionally one or more methylene groups of R^5 are replaced with optionally substituted carbocyclylene, optionally substituted heterocyclylene, optionally substituted arylene, optionally substituted heteroarylene, $-\text{N}(\text{R}^{\text{N}})-$, $-\text{O}-$, $-\text{S}-$, $-\text{C}(\text{O})-$, $-\text{C}(\text{O})\text{N}(\text{R}^{\text{N}})-$, $-\text{NR}^{\text{N}}\text{C}(\text{O})-$, $-\text{NR}^{\text{N}}\text{C}(\text{O})\text{N}(\text{R}^{\text{N}})-$, $-\text{C}(\text{O})\text{O}-$, $-\text{OC}(\text{O})-$, $-\text{OC}(\text{O})\text{O}-$, $-\text{OC}(\text{O})\text{N}(\text{R}^{\text{N}})-$, $-\text{NR}^{\text{N}}\text{C}(\text{O})\text{O}-$, $-\text{C}(\text{O})\text{S}-$, $-\text{SC}(\text{O})-$, $-\text{C}(=\text{NR}^{\text{N}})-$, $-\text{C}(=\text{NR}^{\text{N}})\text{N}(\text{R}^{\text{N}})-$, $-\text{NR}^{\text{N}}\text{C}(=\text{NR}^{\text{N}})-$, $-\text{NR}^{\text{N}}\text{C}(=\text{NR}^{\text{N}})\text{N}(\text{R}^{\text{N}})-$, $-\text{C}(\text{S})-$, $-\text{C}(\text{S})\text{N}(\text{R}^{\text{N}})-$, $-\text{NR}^{\text{N}}\text{C}(\text{S})-$, $-\text{NR}^{\text{N}}\text{C}(\text{S})\text{N}(\text{R}^{\text{N}})-$, $-\text{S}(\text{O})-$, $-\text{OS}(\text{O})-$, $-\text{S}(\text{O})\text{O}-$, $-\text{OS}(\text{O})\text{O}-$, $-\text{OS}(\text{O})_2-$, $-\text{S}(\text{O})_2\text{O}-$, $-\text{OS}(\text{O})_2\text{O}-$, $-\text{N}(\text{R}^{\text{N}})\text{S}(\text{O})-$, $-\text{S}(\text{O})\text{N}(\text{R}^{\text{N}})-$, $-\text{N}(\text{R}^{\text{N}})\text{S}(\text{O})\text{N}(\text{R}^{\text{N}})-$, $-\text{OS}(\text{O})\text{N}(\text{R}^{\text{N}})-$, $-\text{N}(\text{R}^{\text{N}})\text{S}(\text{O})\text{O}-$, $-\text{S}(\text{O})_2-$, $-\text{N}(\text{R}^{\text{N}})\text{S}(\text{O})_2-$, $-\text{S}(\text{O})_2\text{N}(\text{R}^{\text{N}})-$, $-\text{N}(\text{R}^{\text{N}})\text{S}(\text{O})_2\text{N}(\text{R}^{\text{N}})-$, $-\text{OS}(\text{O})_2\text{N}(\text{R}^{\text{N}})-$, or $-\text{N}(\text{R}^{\text{N}})\text{S}(\text{O})_2\text{O}-$; and

each instance of R^{N} is independently hydrogen, optionally substituted alkyl, or a nitrogen protecting group.


[0880] In certain embodiments, the compound of Formula (VIII) is of Formula (VIII-OH):




(VIII-OH),

or a salt thereof. In some embodiments, r is 45.


[0881] In certain embodiments, a compound of Formula (VIII) is of one of the following formulae:




(Compound 419),




Compound 420),




(Compound 421),




(Compound 422),




(Compound 423),



(Compound 424),




(Compound 425),



(Compound 426),

or a salt thereof. In some embodiments, r is 45.


[0882] In yet other embodiments the compound of Formula (VIII) is:



(Compound 427),

or a salt thereof.

[0883] In one embodiment, the compound of Formula (VIII) is



(Compound 428).


- [0884]** In one embodiment, the amount of PEG-lipid in the lipid composition of a pharmaceutical composition disclosed herein ranges from about 0.1 mol % to about 5 mol %, from about 0.5 mol % to about 5 mol %, from about 1 mol % to about 5 mol %, from about 1.5 mol % to about 5 mol %, from about 2 mol % to about 5 mol %, from about 0.1 mol % to about 4 mol %, from about 0.5 mol % to about 4 mol %, from about 1 mol % to about 4 mol %, from about 1.5 mol % to about 4 mol %, from about 2 mol % to about 4 mol %, from about 0.1 mol % to about 3 mol %, from about 0.5 mol % to about 3 mol %, from about 1 mol % to about 3 mol %, from about 1.5 mol % to about 3 mol %, from about 2 mol % to about 3 mol %, from about 0.1 mol % to about 2 mol %, from about 0.5 mol % to about 2 mol %, from about 1 mol % to about 2 mol %, from about 1.5 mol % to about 2 mol %, from about 0.1 mol % to about 1.5 mol %, from about 0.5 mol % to about 1.5 mol %, or from about 1 mol % to about 1.5 mol %.
- [0885]** In one embodiment, the amount of PEG-lipid in the lipid composition disclosed herein is about 2 mol %.
- [0886]** In one embodiment, the amount of PEG-lipid in the lipid composition disclosed herein is at least about 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1, 1.1, 1.2, 1.3, 1.4, 1.5, 1.6, 1.7, 1.8, 1.9, 2, 2.1, 2.2, 2.3, 2.4, 2.5, 2.6, 2.7, 2.8, 2.9, 3, 3.1, 3.2, 3.3, 3.4, 3.5, 3.6, 3.7, 3.8, 3.9, 4, 4.1, 4.2, 4.3, 4.4, 4.5, 4.6, 4.7, 4.8, 4.9, or 5 mol %.
- [0887]** In some aspects, the lipid composition of the pharmaceutical compositions disclosed herein does not comprise a PEG-lipid.

(v) Other Ionizable Amino Lipids

- [0888]** The lipid composition of the pharmaceutical composition disclosed herein can comprise one or more ionizable amino lipids in addition to a lipid according to Formula (I), (III), (IV), (V), or (VI).
- [0889]** Ionizable lipids can be selected from the non-limiting group consisting of 3-(didodecylamino)-N1,N1,4-tridodecyl-1-piperazineethanamine (KL10), N1-[2-(didodecylamino)ethyl]-N1,N4,N4-tridodecyl-1,4-piperazinediethanamine (KL22), 14,25-ditridecyl-15,18,21,24-tetraaza-octatriacontane (KL25), 1,2-dilinoleyloxy-N,N-dimethylaminopropane (DLin-DMA), 2,2-dilinoley1-4-dimethylaminomethyl-[1,3]-dioxolane (DLin-K-DMA), heptatriaconta-6,9,28,31-tetraen-19-yl 4-(dimethylamino)butanoate (DLin-MC3-DMA), 2,2-dilinoley1-4-(2-dimethylaminoethyl)-[1,3]-dioxolane (DLin-KC2-DMA),


1,2-dioleoyloxy-N,N-dimethylaminopropane (DODMA), (13Z,16Z)-N,N-dimethyl-3-nonydocosa-13-16-dien-1-amine (L608),
 2-({8-[(3 β)-cholest-5-en-3-yloxy]octyl}oxy)-N,N-dimethyl-3-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]propan-1-amine (Octyl-CLinDMA),
 (2R)-2-({8-[(3 β)-cholest-5-en-3-yloxy]octyl}oxy)-N,N-dimethyl-3-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]propan-1-amine (Octyl-CLinDMA (2R)), and
 (2S)-2-({8-[(3 β)-cholest-5-en-3-yloxy]octyl}oxy)-N,N-dimethyl-3-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]propan-1-amine (Octyl-CLinDMA (2S)). In addition to these, an ionizable amino lipid can also be a lipid including a cyclic amine group.


[0890] Ionizable lipids can also be the compounds disclosed in International Publication No. WO 2017/075531 A1, hereby incorporated by reference in its entirety. For example, the ionizable amino lipids include, but not limited to:



and any combination thereof.


[0891] Ionizable lipids can also be the compounds disclosed in International Publication No. WO 2015/199952 A1, hereby incorporated by reference in its entirety. For example, the ionizable amino lipids include, but not limited to:





and any combination thereof.

[0892] Ionizable lipids can further include, but are not limited to:



and any combination thereof.

(vi) Other Lipid Composition Components

[0893] The lipid composition of a pharmaceutical composition disclosed herein can include one or more components in addition to those described above. For example, the lipid composition can include one or more permeability enhancer molecules, carbohydrates, polymers, surface altering agents (e.g., surfactants), or other components. For example, a permeability enhancer molecule can be a molecule described by U.S. Patent Application Publication No. 2005/0222064. Carbohydrates can include simple sugars (e.g., glucose) and polysaccharides (e.g., glycogen and derivatives and analogs thereof).

[0894] A polymer can be included in and/or used to encapsulate or partially encapsulate a pharmaceutical composition disclosed herein (e.g., a pharmaceutical composition in lipid

nanoparticle form). A polymer can be biodegradable and/or biocompatible. A polymer can be selected from, but is not limited to, polyamines, polyethers, polyamides, polyesters, polycarbamates, polyureas, polycarbonates, polystyrenes, polyimides, polysulfones, polyurethanes, polyacetylenes, polyethylenes, polyethyleneimines, polyisocyanates, polyacrylates, polymethacrylates, polyacrylonitriles, and polyarylates.

[0895] The ratio between the lipid composition and the polynucleotide range can be from about 10:1 to about 60:1 (wt/wt).

[0896] In some embodiments, the ratio between the lipid composition and the polynucleotide can be about 10:1, 11:1, 12:1, 13:1, 14:1, 15:1, 16:1, 17:1, 18:1, 19:1, 20:1, 21:1, 22:1, 23:1, 24:1, 25:1, 26:1, 27:1, 28:1, 29:1, 30:1, 31:1, 32:1, 33:1, 34:1, 35:1, 36:1, 37:1, 38:1, 39:1, 40:1, 41:1, 42:1, 43:1, 44:1, 45:1, 46:1, 47:1, 48:1, 49:1, 50:1, 51:1, 52:1, 53:1, 54:1, 55:1, 56:1, 57:1, 58:1, 59:1 or 60:1 (wt/wt). In some embodiments, the wt/wt ratio of the lipid composition to the polynucleotide encoding a therapeutic agent is about 20:1 or about 15:1.

[0897] In some embodiments, the pharmaceutical composition disclosed herein can contain more than one polypeptides. For example, a pharmaceutical composition disclosed herein can contain two or more polynucleotides (e.g., RNA, e.g., mRNA).

[0898] In one embodiment, the lipid nanoparticles described herein can comprise polynucleotides (e.g., mRNA) in a lipid:polynucleotide weight ratio of 5:1, 10:1, 15:1, 20:1, 25:1, 30:1, 35:1, 40:1, 45:1, 50:1, 55:1, 60:1 or 70:1, or a range or any of these ratios such as, but not limited to, 5:1 to about 10:1, from about 5:1 to about 15:1, from about 5:1 to about 20:1, from about 5:1 to about 25:1, from about 5:1 to about 30:1, from about 5:1 to about 35:1, from about 5:1 to about 40:1, from about 5:1 to about 45:1, from about 5:1 to about 50:1, from about 5:1 to about 55:1, from about 5:1 to about 60:1, from about 5:1 to about 70:1, from about 10:1 to about 15:1, from about 10:1 to about 20:1, from about 10:1 to about 25:1, from about 10:1 to about 30:1, from about 10:1 to about 35:1, from about 10:1 to about 40:1, from about 10:1 to about 45:1, from about 10:1 to about 50:1, from about 10:1 to about 55:1, from about 10:1 to about 60:1, from about 10:1 to about 70:1, from about 15:1 to about 20:1, from about 15:1 to about 25:1, from about 15:1 to about 30:1, from about 15:1 to about 35:1, from about 15:1 to about 40:1, from about 15:1 to about 45:1, from about 15:1 to about 50:1, from about 15:1 to about 55:1, from about 15:1 to about 60:1 or from about 15:1 to about 70:1.

[0899] In one embodiment, the lipid nanoparticles described herein can comprise the polynucleotide in a concentration from approximately 0.1 mg/ml to 2 mg/ml such as, but not limited to, 0.1 mg/ml, 0.2 mg/ml, 0.3 mg/ml, 0.4 mg/ml, 0.5 mg/ml, 0.6 mg/ml, 0.7 mg/ml, 0.8 mg/ml, 0.9 mg/ml, 1.0 mg/ml, 1.1 mg/ml, 1.2 mg/ml, 1.3 mg/ml, 1.4 mg/ml, 1.5 mg/ml, 1.6 mg/ml, 1.7 mg/ml, 1.8 mg/ml, 1.9 mg/ml, 2.0 mg/ml or greater than 2.0 mg/ml.

(vii) Nanoparticle Compositions

[0900] In some embodiments, the pharmaceutical compositions disclosed herein are formulated as lipid nanoparticles (LNP). Accordingly, the present disclosure also provides nanoparticle compositions comprising (i) a lipid composition comprising a delivery agent such as a compound of Formula (I) or (III) as described herein, and (ii) a polynucleotide encoding an ACADVL polypeptide. In such nanoparticle composition, the lipid composition disclosed herein can encapsulate the polynucleotide encoding an ACADVL polypeptide.

[0901] Nanoparticle compositions are typically sized on the order of micrometers or smaller and can include a lipid bilayer. Nanoparticle compositions encompass lipid nanoparticles (LNPs), liposomes (e.g., lipid vesicles), and lipoplexes. For example, a nanoparticle composition can be a liposome having a lipid bilayer with a diameter of 500 nm or less.

[0902] Nanoparticle compositions include, for example, lipid nanoparticles (LNPs), liposomes, and lipoplexes. In some embodiments, nanoparticle compositions are vesicles including one or more lipid bilayers. In certain embodiments, a nanoparticle composition includes two or more concentric bilayers separated by aqueous compartments. Lipid bilayers can be functionalized and/or crosslinked to one another. Lipid bilayers can include one or more ligands, proteins, or channels.

[0903] In some embodiments, the nanoparticle compositions of the present disclosure comprise at least one compound according to Formula (I), (III), (IV), (V), or (VI). For example, the nanoparticle composition can include one or more of Compounds 1-147, or one or more of Compounds 1-342. Nanoparticle compositions can also include a variety of other components. For example, the nanoparticle composition may include one or more other lipids in addition to a lipid according to Formula (I), (III), (IV), (V), or (VI), such as (i) at least one phospholipid, (ii) at least one structural lipid, (iii) at least one PEG-lipid,

or (iv) any combination thereof. Inclusion of structural lipid can be optional, for example when lipids according to Formula III are used in the lipid nanoparticle compositions of the invention.

- [0904] Nanoparticle compositions of the present disclosure comprise at least one compound according to Formula (I). For example, the nanoparticle composition can include one or more of Compounds 1-147. Nanoparticle compositions can also include a variety of other components. For example, the nanoparticle composition can include one or more other lipids in addition to a lipid according to Formula (I) or (II), for example (i) at least one phospholipid, (ii) at least one structural lipid, (iii) at least one PEG-lipid, or (iv) any combination thereof.
- [0905] In some embodiments, the nanoparticle composition comprises a compound of Formula (I), (e.g., Compounds 18, 25, 26 or 48). In some embodiments, the nanoparticle composition comprises a compound of Formula (I) (e.g., Compounds 18, 25, 26 or 48) and a phospholipid (e.g., DOPE or DSPC).
- [0906] In some embodiments, the nanoparticle composition comprises a compound of Formula (III) (e.g., Compound 236). In some embodiments, the nanoparticle composition comprises a compound of Formula (III) (e.g., Compound 236) and a phospholipid (e.g., DOPE or DSPC).
- [0907] In some embodiments, the nanoparticle composition comprises a lipid composition consisting or consisting essentially of compound of Formula (I) (e.g., Compounds 18, 25, 26 or 48). In some embodiments, the nanoparticle composition comprises a lipid composition consisting or consisting essentially of a compound of Formula (I) (e.g., Compounds 18, 25, 26 or 48) and a phospholipid (e.g., DSPC).
- [0908] In one embodiment, a lipid nanoparticle comprises an ionizable lipid, a structural lipid, a phospholipid, and mRNA. In some embodiments, the LNP comprises an ionizable lipid, a PEG-modified lipid, a sterol and a structural lipid. In some embodiments, the LNP has a molar ratio of about 20-60% ionizable lipid: about 5-25% structural lipid: about 25-55% sterol; and about 0.5-15% PEG-modified lipid. In some embodiments, the LNP comprises a molar ratio of about 50% ionizable lipid, about 1.5% PEG-modified lipid, about 38.5% cholesterol and about 10% structural lipid. In some embodiments, the LNP comprises a molar ratio of about 55% ionizable lipid, about 2.5% PEG lipid, about 32.5% cholesterol and about 10% structural lipid. In some embodiments, the ionizable lipid is an ionizable amino lipid and the structural lipid is a neutral lipid, and the sterol is

a cholesterol. In some embodiments, the LNP has a molar ratio of 50:38.5:10:1.5 of ionizable lipid: cholesterol: DSPC: PEG lipid. In some embodiments, the ionizable lipid is Compound 18 or Compound 236, and the PEG lipid is Compound 428.

[0909] In some embodiments, the LNP has a molar ratio of 50:38.5:10:1.5 of Compound 18 : Cholesterol : Phospholipid : Compound 428. In some embodiments, the LNP has a molar ratio of 50:38.5:10:1.5 of Compound 18 : Cholesterol : DSPC : Compound 428.

[0910] In some embodiments, the LNP has a molar ratio of 50:38.5:10:1.5 of Compound 236 : Cholesterol : Phospholipid : Compound 428. In some embodiments, the LNP has a molar ratio of 50:38.5:10:1.5 of Compound 236 : Cholesterol : DSPC : Compound 428.

[0911] In some embodiments, the LNP has a polydispersity value of less than 0.4. In some embodiments, the LNP has a net neutral charge at a neutral pH. In some embodiments, the LNP has a mean diameter of 50-150 nm. In some embodiments, the LNP has a mean diameter of 80-100 nm.

[0912] As generally defined herein, the term “lipid” refers to a small molecule that has hydrophobic or amphiphilic properties. Lipids may be naturally occurring or synthetic. Examples of classes of lipids include, but are not limited to, fats, waxes, sterol-containing metabolites, vitamins, fatty acids, glycerolipids, glycerophospholipids, sphingolipids, saccharolipids, and polyketides, and prenol lipids. In some instances, the amphiphilic properties of some lipids leads them to form liposomes, vesicles, or membranes in aqueous media.

[0913] In some embodiments, a lipid nanoparticle (LNP) may comprise an ionizable lipid. As used herein, the term “ionizable lipid” has its ordinary meaning in the art and may refer to a lipid comprising one or more charged moieties. In some embodiments, an ionizable lipid may be positively charged or negatively charged. An ionizable lipid may be positively charged, in which case it can be referred to as “cationic lipid.” In certain embodiments, an ionizable lipid molecule may comprise an amine group, and can be referred to as an ionizable amino lipid. As used herein, a “charged moiety” is a chemical moiety that carries a formal electronic charge, e.g., monovalent (+1, or -1), divalent (+2, or -2), trivalent (+3, or -3), etc. The charged moiety may be anionic (i.e., negatively charged) or cationic (i.e., positively charged). Examples of positively-charged moieties include amine groups (e.g., primary, secondary, and/or tertiary amines), ammonium groups, pyridinium group, guanidine groups, and imidazolium groups. In a particular embodiment, the charged moieties comprise amine groups. Examples of negatively-

charged groups or precursors thereof, include carboxylate groups, sulfonate groups, sulfate groups, phosphonate groups, phosphate groups, hydroxyl groups, and the like. The charge of the charged moiety may vary, in some cases, with the environmental conditions, for example, changes in pH may alter the charge of the moiety, and/or cause the moiety to become charged or uncharged. In general, the charge density of the molecule may be selected as desired.

[0914] It should be understood that the terms “charged” or “charged moiety” does not refer to a “partial negative charge” or “partial positive charge” on a molecule. The terms “partial negative charge” and “partial positive charge” are given its ordinary meaning in the art. A “partial negative charge” may result when a functional group comprises a bond that becomes polarized such that electron density is pulled toward one atom of the bond, creating a partial negative charge on the atom. Those of ordinary skill in the art will, in general, recognize bonds that can become polarized in this way.

[0915] In some embodiments, the ionizable lipid is an ionizable amino lipid, sometimes referred to in the art as an “ionizable cationic lipid”. In one embodiment, the ionizable amino lipid may have a positively charged hydrophilic head and a hydrophobic tail that are connected via a linker structure.

[0916] In addition to these, an ionizable lipid may also be a lipid including a cyclic amine group.

[0917] In one embodiment, the ionizable lipid may be selected from, but not limited to, an ionizable lipid described in International Publication Nos. WO2013086354 and WO2013116126; the contents of each of which are herein incorporated by reference in their entirety.

[0918] In yet another embodiment, the ionizable lipid may be selected from, but not limited to, Formula CLI-CLXXXXII of US Patent No. 7,404,969; each of which is herein incorporated by reference in their entirety.

[0919] In one embodiment, the lipid may be a cleavable lipid such as those described in International Publication No. WO2012170889, herein incorporated by reference in its entirety. In one embodiment, the lipid may be synthesized by methods known in the art and/or as described in International Publication Nos. WO2013086354; the contents of each of which are herein incorporated by reference in their entirety.

[0920] Nanoparticle compositions can be characterized by a variety of methods. For example, microscopy (e.g., transmission electron microscopy or scanning electron

microscopy) can be used to examine the morphology and size distribution of a nanoparticle composition. Dynamic light scattering or potentiometry (e.g., potentiometric titrations) can be used to measure zeta potentials. Dynamic light scattering can also be utilized to determine particle sizes. Instruments such as the Zetasizer Nano ZS (Malvern Instruments Ltd, Malvern, Worcestershire, UK) can also be used to measure multiple characteristics of a nanoparticle composition, such as particle size, polydispersity index, and zeta potential.

[0921] The size of the nanoparticles can help counter biological reactions such as, but not limited to, inflammation, or can increase the biological effect of the polynucleotide.

[0922] As used herein, “size” or “mean size” in the context of nanoparticle compositions refers to the mean diameter of a nanoparticle composition.

[0923] In one embodiment, the polynucleotide encoding an ACADVL polypeptide are formulated in lipid nanoparticles having a diameter from about 10 to about 100 nm such as, but not limited to, about 10 to about 20 nm, about 10 to about 30 nm, about 10 to about 40 nm, about 10 to about 50 nm, about 10 to about 60 nm, about 10 to about 70 nm, about 10 to about 80 nm, about 10 to about 90 nm, about 20 to about 30 nm, about 20 to about 40 nm, about 20 to about 50 nm, about 20 to about 60 nm, about 20 to about 70 nm, about 20 to about 80 nm, about 20 to about 90 nm, about 20 to about 100 nm, about 30 to about 40 nm, about 30 to about 50 nm, about 30 to about 60 nm, about 30 to about 70 nm, about 30 to about 80 nm, about 30 to about 90 nm, about 30 to about 100 nm, about 40 to about 50 nm, about 40 to about 60 nm, about 40 to about 70 nm, about 40 to about 80 nm, about 40 to about 90 nm, about 40 to about 100 nm, about 50 to about 60 nm, about 50 to about 70 nm, about 50 to about 80 nm, about 50 to about 90 nm, about 50 to about 100 nm, about 60 to about 70 nm, about 60 to about 80 nm, about 60 to about 90 nm, about 60 to about 100 nm, about 70 to about 80 nm, about 70 to about 90 nm, about 70 to about 100 nm, about 80 to about 90 nm, about 80 to about 100 nm and/or about 90 to about 100 nm.

[0924] In one embodiment, the nanoparticles have a diameter from about 10 to 500 nm. In one embodiment, the nanoparticle has a diameter greater than 100 nm, greater than 150 nm, greater than 200 nm, greater than 250 nm, greater than 300 nm, greater than 350 nm, greater than 400 nm, greater than 450 nm, greater than 500 nm, greater than 550 nm, greater than 600 nm, greater than 650 nm, greater than 700 nm, greater than 750 nm,

greater than 800 nm, greater than 850 nm, greater than 900 nm, greater than 950 nm or greater than 1000 nm.

[0925] In some embodiments, the largest dimension of a nanoparticle composition is 1 μm or shorter (e.g., 1 μm , 900 nm, 800 nm, 700 nm, 600 nm, 500 nm, 400 nm, 300 nm, 200 nm, 175 nm, 150 nm, 125 nm, 100 nm, 75 nm, 50 nm, or shorter).

[0926] A nanoparticle composition can be relatively homogenous. A polydispersity index can be used to indicate the homogeneity of a nanoparticle composition, e.g., the particle size distribution of the nanoparticle composition. A small (e.g., less than 0.3) polydispersity index generally indicates a narrow particle size distribution. A nanoparticle composition can have a polydispersity index from about 0 to about 0.25, such as 0.01, 0.02, 0.03, 0.04, 0.05, 0.06, 0.07, 0.08, 0.09, 0.10, 0.11, 0.12, 0.13, 0.14, 0.15, 0.16, 0.17, 0.18, 0.19, 0.20, 0.21, 0.22, 0.23, 0.24, or 0.25. In some embodiments, the polydispersity index of a nanoparticle composition disclosed herein can be from about 0.10 to about 0.20.

[0927] The zeta potential of a nanoparticle composition can be used to indicate the electrokinetic potential of the composition. For example, the zeta potential can describe the surface charge of a nanoparticle composition. Nanoparticle compositions with relatively low charges, positive or negative, are generally desirable, as more highly charged species can interact undesirably with cells, tissues, and other elements in the body. In some embodiments, the zeta potential of a nanoparticle composition disclosed herein can be from about -10 mV to about +20 mV, from about -10 mV to about +15 mV, from about -10 mV to about +10 mV, from about -10 mV to about +5 mV, from about -10 mV to about 0 mV, from about -10 mV to about -5 mV, from about -5 mV to about +20 mV, from about -5 mV to about +15 mV, from about -5 mV to about +10 mV, from about -5 mV to about +5 mV, from about -5 mV to about 0 mV, from about 0 mV to about +20 mV, from about 0 mV to about +15 mV, from about 0 mV to about +10 mV, from about 0 mV to about +5 mV, from about +5 mV to about +20 mV, from about +5 mV to about +15 mV, or from about +5 mV to about +10 mV.

[0928] In some embodiments, the zeta potential of the lipid nanoparticles can be from about 0 mV to about 100 mV, from about 0 mV to about 90 mV, from about 0 mV to about 80 mV, from about 0 mV to about 70 mV, from about 0 mV to about 60 mV, from about 0 mV to about 50 mV, from about 0 mV to about 40 mV, from about 0 mV to about 30 mV, from about 0 mV to about 20 mV, from about 0 mV to about 10 mV, from about

10 mV to about 100 mV, from about 10 mV to about 90 mV, from about 10 mV to about 80 mV, from about 10 mV to about 70 mV, from about 10 mV to about 60 mV, from about 10 mV to about 50 mV, from about 10 mV to about 40 mV, from about 10 mV to about 30 mV, from about 10 mV to about 20 mV, from about 20 mV to about 100 mV, from about 20 mV to about 90 mV, from about 20 mV to about 80 mV, from about 20 mV to about 70 mV, from about 20 mV to about 60 mV, from about 20 mV to about 50 mV, from about 20 mV to about 40 mV, from about 20 mV to about 30 mV, from about 30 mV to about 100 mV, from about 30 mV to about 90 mV, from about 30 mV to about 80 mV, from about 30 mV to about 70 mV, from about 30 mV to about 60 mV, from about 30 mV to about 50 mV, from about 30 mV to about 40 mV, from about 40 mV to about 100 mV, from about 40 mV to about 90 mV, from about 40 mV to about 80 mV, from about 40 mV to about 70 mV, from about 40 mV to about 60 mV, and from about 40 mV to about 50 mV. In some embodiments, the zeta potential of the lipid nanoparticles can be from about 10 mV to about 50 mV, from about 15 mV to about 45 mV, from about 20 mV to about 40 mV, and from about 25 mV to about 35 mV. In some embodiments, the zeta potential of the lipid nanoparticles can be about 10 mV, about 20 mV, about 30 mV, about 40 mV, about 50 mV, about 60 mV, about 70 mV, about 80 mV, about 90 mV, and about 100 mV.

[0929] The term “encapsulation efficiency” of a polynucleotide describes the amount of the polynucleotide that is encapsulated by or otherwise associated with a nanoparticle composition after preparation, relative to the initial amount provided. As used herein, “encapsulation” can refer to complete, substantial, or partial enclosure, confinement, surrounding, or encasement.

[0930] Encapsulation efficiency is desirably high (e.g., close to 100%). The encapsulation efficiency can be measured, for example, by comparing the amount of the polynucleotide in a solution containing the nanoparticle composition before and after breaking up the nanoparticle composition with one or more organic solvents or detergents.

[0931] Fluorescence can be used to measure the amount of free polynucleotide in a solution. For the nanoparticle compositions described herein, the encapsulation efficiency of a polynucleotide can be at least 50%, for example 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100%. In some embodiments, the encapsulation efficiency can be at least 80%. In certain embodiments, the encapsulation efficiency can be at least 90%.

- [0932] The amount of a polynucleotide present in a pharmaceutical composition disclosed herein can depend on multiple factors such as the size of the polynucleotide, desired target and/or application, or other properties of the nanoparticle composition as well as on the properties of the polynucleotide.
- [0933] For example, the amount of an mRNA useful in a nanoparticle composition can depend on the size (expressed as length, or molecular mass), sequence, and other characteristics of the mRNA. The relative amounts of a polynucleotide in a nanoparticle composition can also vary.
- [0934] The relative amounts of the lipid composition and the polynucleotide present in a lipid nanoparticle composition of the present disclosure can be optimized according to considerations of efficacy and tolerability. For compositions including an mRNA as a polynucleotide, the N:P ratio can serve as a useful metric.
- [0935] As the N:P ratio of a nanoparticle composition controls both expression and tolerability, nanoparticle compositions with low N:P ratios and strong expression are desirable. N:P ratios vary according to the ratio of lipids to RNA in a nanoparticle composition.
- [0936] In general, a lower N:P ratio is preferred. The one or more RNA, lipids, and amounts thereof can be selected to provide an N:P ratio from about 2:1 to about 30:1, such as 2:1, 3:1, 4:1, 5:1, 6:1, 7:1, 8:1, 9:1, 10:1, 12:1, 14:1, 16:1, 18:1, 20:1, 22:1, 24:1, 26:1, 28:1, or 30:1. In certain embodiments, the N:P ratio can be from about 2:1 to about 8:1. In other embodiments, the N:P ratio is from about 5:1 to about 8:1. In certain embodiments, the N:P ratio is between 5:1 and 6:1. In one specific aspect, the N:P ratio is about 5.67:1.
- [0937] In addition to providing nanoparticle compositions, the present disclosure also provides methods of producing lipid nanoparticles comprising encapsulating a polynucleotide. Such method comprises using any of the pharmaceutical compositions disclosed herein and producing lipid nanoparticles in accordance with methods of production of lipid nanoparticles known in the art. *See, e.g.,* Wang et al. (2015) "Delivery of oligonucleotides with lipid nanoparticles" *Adv. Drug Deliv. Rev.* 87:68-80; Silva et al. (2015) "Delivery Systems for Biopharmaceuticals. Part I: Nanoparticles and Microparticles" *Curr. Pharm. Technol.* 16: 940-954; Naseri et al. (2015) "Solid Lipid Nanoparticles and Nanostructured Lipid Carriers: Structure, Preparation and Application"

Adv. Pharm. Bull. 5:305-13; Silva et al. (2015) "Lipid nanoparticles for the delivery of biopharmaceuticals" Curr. Pharm. Biotechnol. 16:291-302, and references cited therein.

23. Other Delivery Agents

a. Liposomes, Lipoplexes, and Lipid Nanoparticles

[0938] In some embodiments, the compositions or formulations of the present disclosure comprise a delivery agent, e.g., a liposome, a lipoplex, a lipid nanoparticle, or any combination thereof. The polynucleotides described herein (e.g., a polynucleotide comprising a nucleotide sequence encoding an ACADVL polypeptide) can be formulated using one or more liposomes, lipoplexes, or lipid nanoparticles. Liposomes, lipoplexes, or lipid nanoparticles can be used to improve the efficacy of the polynucleotides directed protein production as these formulations can increase cell transfection by the polynucleotide; and/or increase the translation of encoded protein. The liposomes, lipoplexes, or lipid nanoparticles can also be used to increase the stability of the polynucleotides.

[0939] Liposomes are artificially-prepared vesicles that can primarily be composed of a lipid bilayer and can be used as a delivery vehicle for the administration of pharmaceutical formulations. Liposomes can be of different sizes. A multilamellar vesicle (MLV) can be hundreds of nanometers in diameter, and can contain a series of concentric bilayers separated by narrow aqueous compartments. A small unilamellar vesicle (SUV) can be smaller than 50 nm in diameter, and a large unilamellar vesicle (LUV) can be between 50 and 500 nm in diameter. Liposome design can include, but is not limited to, opsonins or ligands to improve the attachment of liposomes to unhealthy tissue or to activate events such as, but not limited to, endocytosis. Liposomes can contain a low or a high pH value in order to improve the delivery of the pharmaceutical formulations.

[0940] The formation of liposomes can depend on the pharmaceutical formulation entrapped and the liposomal ingredients, the nature of the medium in which the lipid vesicles are dispersed, the effective concentration of the entrapped substance and its potential toxicity, any additional processes involved during the application and/or delivery of the vesicles, the optimal size, polydispersity and the shelf-life of the vesicles for the intended application, and the batch-to-batch reproducibility and scale up production of safe and efficient liposomal products, etc.

[0941] As a non-limiting example, liposomes such as synthetic membrane vesicles can be prepared by the methods, apparatus and devices described in U.S. Pub. Nos. US20130177638, US20130177637, US20130177636, US20130177635, US20130177634, US20130177633, US20130183375, US20130183373, and US20130183372. In some embodiments, the polynucleotides described herein can be encapsulated by the liposome and/or it can be contained in an aqueous core that can then be encapsulated by the liposome as described in, e.g., Intl. Pub. Nos. WO2012031046, WO2012031043, WO2012030901, WO2012006378, and WO2013086526; and U.S. Pub. Nos. US20130189351, US20130195969 and US20130202684. Each of the references in herein incorporated by reference in its entirety.

[0942] In some embodiments, the polynucleotides described herein can be formulated in a cationic oil-in-water emulsion where the emulsion particle comprises an oil core and a cationic lipid that can interact with the polynucleotide anchoring the molecule to the emulsion particle. In some embodiments, the polynucleotides described herein can be formulated in a water-in-oil emulsion comprising a continuous hydrophobic phase in which the hydrophilic phase is dispersed. Exemplary emulsions can be made by the methods described in Intl. Pub. Nos. WO2012006380 and WO201087791, each of which is herein incorporated by reference in its entirety.

[0943] In some embodiments, the polynucleotides described herein can be formulated in a lipid-polycation complex. The formation of the lipid-polycation complex can be accomplished by methods as described in, e.g., U.S. Pub. No. US20120178702. As a non-limiting example, the polycation can include a cationic peptide or a polypeptide such as, but not limited to, polylysine, polyornithine and/or polyarginine and the cationic peptides described in Intl. Pub. No. WO2012013326 or U.S. Pub. No. US20130142818. Each of the references is herein incorporated by reference in its entirety.

[0944] In some embodiments, the polynucleotides described herein can be formulated in a lipid nanoparticle (LNP) such as those described in Intl. Pub. Nos. WO2013123523, WO2012170930, WO2011127255 and WO2008103276; and U.S. Pub. No. US20130171646, each of which is herein incorporated by reference in its entirety.

[0945] Lipid nanoparticle formulations typically comprise one or more lipids. In some embodiments, the lipid is an ionizable lipid (e.g., an ionizable amino lipid), sometimes referred to in the art as an “ionizable cationic lipid.” In some embodiments, lipid nanoparticle formulations further comprise other components, including a phospholipid, a

structural lipid, and a molecule capable of reducing particle aggregation, for example a PEG or PEG-modified lipid.

[0946] Exemplary ionizable lipids include, but not limited to, any one of Compounds 1-342 disclosed herein, DLin-MC3-DMA (MC3), DLin-DMA, DLenDMA, DLin-D-DMA, DLin-K-DMA, DLin-M-C2-DMA, DLin-K-DMA, DLin-KC2-DMA, DLin-KC3-DMA, DLin-KC4-DMA, DLin-C2K-DMA, DLin-MP-DMA, DODMA, 98N12-5, C12-200, DLin-C-DAP, DLin-DAC, DLinDAP, DLinAP, DLin-EG-DMA, DLin-2-DMAP, KL10, KL22, KL25, Octyl-CLinDMA, Octyl-CLinDMA (2R), Octyl-CLinDMA (2S), and any combination thereof. Other exemplary ionizable lipids include, (13Z,16Z)-N,N-dimethyl-3-nonyldocosa-13,16-dien-1-amine (L608), (20Z,23Z)-N,N-dimethylnonacos-20,23-dien-10-amine, (17Z,20Z)-N,N-dimethylhexacos-17,20-dien-9-amine, (16Z,19Z)-N,N-dimethylpentacos-16,19-dien-8-amine, (13Z,16Z)-N,N-dimethyldocosa-13,16-dien-5-amine, (12Z,15Z)-N,N-dimethylhenicos-12,15-dien-4-amine, (14Z,17Z)-N,N-dimethyltricos-14,17-dien-6-amine, (15Z,18Z)-N,N-dimethyltetracos-15,18-dien-7-amine, (18Z,21Z)-N,N-dimethylheptacos-18,21-dien-10-amine, (15Z,18Z)-N,N-dimethyltetracos-15,18-dien-5-amine, (14Z,17Z)-N,N-dimethyltricos-14,17-dien-4-amine, (19Z,22Z)-N,N-dimethyloctacos-19,22-dien-9-amine, (18Z,21Z)-N,N-dimethylheptacos-18,21-dien-8-amine, (17Z,20Z)-N,N-dimethylhexacos-17,20-dien-7-amine, (16Z,19Z)-N,N-dimethylpentacos-16,19-dien-6-amine, (22Z,25Z)-N,N-dimethylhentriacont-22,25-dien-10-amine, (21Z,24Z)-N,N-dimethyltriacont-21,24-dien-9-amine, (18Z)-N,N-dimethylheptacos-18-en-10-amine, (17Z)-N,N-dimethylhexacos-17-en-9-amine, (19Z,22Z)-N,N-dimethyloctacos-19,22-dien-7-amine, N,N-dimethylheptacos-10-amine, (20Z,23Z)-N-ethyl-N-methylnonacos-20,23-dien-10-amine, 1-[(11Z,14Z)-1-nonylicos-11,14-dien-1-yl]pyrrolidine, (20Z)-N,N-dimethylheptacos-20-en-10-amine, (15Z)-N,N-dimethylheptacos-15-en-10-amine, (14Z)-N,N-dimethylnonacos-14-en-10-amine, (17Z)-N,N-dimethylnonacos-17-en-10-amine, (24Z)-N,N-dimethyltritriacont-24-en-10-amine, (20Z)-N,N-dimethylnonacos-20-en-10-amine, (22Z)-N,N-dimethylhentriacont-22-en-10-amine, (16Z)-N,N-dimethylpentacos-16-en-8-amine, (12Z,15Z)-N,N-dimethyl-2-nonylhenicos-12,15-dien-1-amine, N,N-dimethyl-1-[(1S,2R)-2-octylcyclopropyl]heptadecan-8-amine, 1-[(1S,2R)-2-hexylcyclopropyl]-N,N-dimethylnonadecan-10-amine, N,N-dimethyl-1-[(1S,2R)-2-octylcyclopropyl]nonadecan-10-amine, N,N-dimethyl-21-[(1S,2R)-2-octylcyclopropyl]henicosan-10-amine, N,N-dimethyl-1-[(1S,2S)-2-[(1R,2R)-2-

pentylcyclopropyl)methyl}cyclopropyl]nonadecan-10-amine, N,N-dimethyl-1-[(1S,2R)-2-octylcyclopropyl]hexadecan-8-amine, N,N-dimethyl-[(1R,2S)-2-undecylcyclopropyl]tetradecan-5-amine, N,N-dimethyl-3-{7-[(1S,2R)-2-octylcyclopropyl]heptyl}dodecan-1-amine, 1-[(1R,2S)-2-heptylcyclopropyl]-N,N-dimethyloctadecan-9-amine, 1-[(1S,2R)-2-decylcyclopropyl]-N,N-dimethylpentadecan-6-amine, N,N-dimethyl-1-[(1S,2R)-2-octylcyclopropyl]pentadecan-8-amine, R-N,N-dimethyl-1-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]-3-(octyloxy)propan-2-amine, S-N,N-dimethyl-1-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]-3-(octyloxy)propan-2-amine, 1-{2-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]-1-[(octyloxy)methyl]ethyl}pyrrolidine, (2S)-N,N-dimethyl-1-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]-3-[(5Z)-oct-5-en-1-yloxy]propan-2-amine, 1-{2-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]-1-[(octyloxy)methyl]ethyl}azetidine, (2S)-1-(hexyloxy)-N,N-dimethyl-3-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]propan-2-amine, (2S)-1-(heptyloxy)-N,N-dimethyl-3-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]propan-2-amine, N,N-dimethyl-1-(nonyloxy)-3-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]propan-2-amine, N,N-dimethyl-1-[(9Z)-octadec-9-en-1-yloxy]-3-(octyloxy)propan-2-amine; (2S)-N,N-dimethyl-1-[(6Z,9Z,12Z)-octadeca-6,9,12-trien-1-yloxy]-3-(octyloxy)propan-2-amine, (2S)-1-[(11Z,14Z)-icosa-11,14-dien-1-yloxy]-N,N-dimethyl-3-(pentyloxy)propan-2-amine, (2S)-1-(hexyloxy)-3-[(11Z,14Z)-icosa-11,14-dien-1-yloxy]-N,N-dimethylpropan-2-amine, 1-[(11Z,14Z)-icosa-11,14-dien-1-yloxy]-N,N-dimethyl-3-(octyloxy)propan-2-amine, 1-[(13Z,16Z)-docosa-13,16-dien-1-yloxy]-N,N-dimethyl-3-(octyloxy)propan-2-amine, (2S)-1-[(13Z,16Z)-docosa-13,16-dien-1-yloxy]-3-(hexyloxy)-N,N-dimethylpropan-2-amine, (2S)-1-[(13Z)-docos-13-en-1-yloxy]-3-(hexyloxy)-N,N-dimethylpropan-2-amine, 1-[(13Z)-docos-13-en-1-yloxy]-N,N-dimethyl-3-(octyloxy)propan-2-amine, 1-[(9Z)-hexadec-9-en-1-yloxy]-N,N-dimethyl-3-(octyloxy)propan-2-amine, (2R)-N,N-dimethyl-H(1-metoyloctyl)oxy]-3-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]propan-2-amine, (2R)-1-[(3,7-dimethyloctyl)oxy]-N,N-dimethyl-3-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]propan-2-amine, N,N-dimethyl-1-(octyloxy)-3-({8-[(1S,2S)-2-{[(1R,2R)-2-pentylcyclopropyl]methyl}cyclopropyl]octyl}oxy)propan-2-amine, N,N-dimethyl-1-{[8-(2-octylcyclopropyl)octyl]oxy}-3-(octyloxy)propan-2-amine, and (11E,20Z,23Z)-N,N-dimethylnonacos-11,20,2-trien-10-amine, and any combination thereof.

[0947] Phospholipids include, but are not limited to, glycerophospholipids such as phosphatidylcholines, phosphatidylethanolamines, phosphatidylserines,

phosphatidylinositols, phosphatidyl glycerols, and phosphatidic acids. Phospholipids also include phosphosphingolipid, such as sphingomyelin. In some embodiments, the phospholipids are DLPC, DMPC, DOPC, DPPC, DSPC, DUPC, 18:0 Diether PC, DLnPC, DAPC, DHAPC, DOPE, 4ME 16:0 PE, DSPE, DLPE, DLnPE, DAPE, DHAPE, DOPG, and any combination thereof. In some embodiments, the phospholipids are MPPC, MSPC, PMPC, PSPC, SMPC, SPPC, DHAPE, DOPG, and any combination thereof. In some embodiments, the amount of phospholipids (e.g., DSPC) in the lipid composition ranges from about 1 mol% to about 20 mol%.

[0948] The structural lipids include sterols and lipids containing sterol moieties. In some embodiments, the structural lipids include cholesterol, fecosterol, sitosterol, ergosterol, campesterol, stigmasterol, brassicasterol, tomatidine, tomatine, ursolic acid, alpha-tocopherol, and mixtures thereof. In some embodiments, the structural lipid is cholesterol. In some embodiments, the amount of the structural lipids (e.g., cholesterol) in the lipid composition ranges from about 20 mol% to about 60 mol%.

[0949] The PEG-modified lipids include PEG-modified phosphatidylethanolamine and phosphatidic acid, PEG-ceramide conjugates (e.g., PEG-CerC14 or PEG-CerC20), PEG-modified dialkylamines and PEG-modified 1,2-diacyloxypropan-3-amines. Such lipids are also referred to as PEGylated lipids. For example, a PEG lipid can be PEG-c-DOMG, PEG-DMG, PEG-DLPE, PEG DMPE, PEG-DPPC, or a PEG-DSPE lipid. In some embodiments, the PEG-lipid are 1,2-dimyristoyl-sn-glycerol methoxypolyethylene glycol (PEG-DMG), 1,2-distearoyl-sn-glycero-3-phosphoethanolamine-N-[amino(polyethylene glycol)] (PEG-DSPE), PEG-disteryl glycerol (PEG-DSG), PEG-dipalmitoleyl, PEG-dioleyl, PEG-distearyl, PEG-diacylglycamide (PEG-DAG), PEG-dipalmitoyl phosphatidylethanolamine (PEG-DPPE), or PEG-1,2-dimyristyloxypropyl-3-amine (PEG-c-DMA). In some embodiments, the PEG moiety has a size of about 1000, 2000, 5000, 10,000, 15,000 or 20,000 Daltons. In some embodiments, the amount of PEG-lipid in the lipid composition ranges from about 0.1 mol% to about 5 mol%.

[0950] In some embodiments, the LNP formulations described herein can additionally comprise a permeability enhancer molecule. Non-limiting permeability enhancer molecules are described in U.S. Pub. No. US20050222064, herein incorporated by reference in its entirety.

[0951] The LNP formulations can further contain a phosphate conjugate. The phosphate conjugate can increase in vivo circulation times and/or increase the targeted delivery of

the nanoparticle. Phosphate conjugates can be made by the methods described in, e.g., Intl. Pub. No. WO2013033438 or U.S. Pub. No. US20130196948. The LNP formulation can also contain a polymer conjugate (e.g., a water soluble conjugate) as described in, e.g., U.S. Pub. Nos. US20130059360, US20130196948, and US20130072709. Each of the references is herein incorporated by reference in its entirety.

[0952] The LNP formulations can comprise a conjugate to enhance the delivery of nanoparticles of the present invention in a subject. Further, the conjugate can inhibit phagocytic clearance of the nanoparticles in a subject. In some embodiments, the conjugate can be a "self" peptide designed from the human membrane protein CD47 (e.g., the "self" particles described by Rodriguez et al, Science 2013 339, 971-975, herein incorporated by reference in its entirety). As shown by Rodriguez et al. the self peptides delayed macrophage-mediated clearance of nanoparticles which enhanced delivery of the nanoparticles.

[0953] The LNP formulations can comprise a carbohydrate carrier. As a non-limiting example, the carbohydrate carrier can include, but is not limited to, an anhydride-modified phytoglycogen or glycogen-type material, phytoglycogen octenyl succinate, phytoglycogen beta-dextrin, anhydride-modified phytoglycogen beta-dextrin (e.g., Intl. Pub. No. WO2012109121, herein incorporated by reference in its entirety).

[0954] The LNP formulations can be coated with a surfactant or polymer to improve the delivery of the particle. In some embodiments, the LNP can be coated with a hydrophilic coating such as, but not limited to, PEG coatings and/or coatings that have a neutral surface charge as described in U.S. Pub. No. US20130183244, herein incorporated by reference in its entirety.

[0955] The LNP formulations can be engineered to alter the surface properties of particles so that the lipid nanoparticles can penetrate the mucosal barrier as described in U.S. Pat. No. 8,241,670 or Intl. Pub. No. WO2013110028, each of which is herein incorporated by reference in its entirety.

[0956] The LNP engineered to penetrate mucus can comprise a polymeric material (i.e., a polymeric core) and/or a polymer-vitamin conjugate and/or a tri-block co-polymer. The polymeric material can include, but is not limited to, polyamines, polyethers, polyamides, polyesters, polycarbamates, polyureas, polycarbonates, poly(styrenes), polyimides, polysulfones, polyurethanes, polyacetylenes, polyethylenes, polyethyleneimines, polyisocyanates, polyacrylates, polymethacrylates, polyacrylonitriles, and polyarylates.

- [0957] LNP engineered to penetrate mucus can also include surface altering agents such as, but not limited to, polynucleotides, anionic proteins (e.g., bovine serum albumin), surfactants (e.g., cationic surfactants such as for example dimethyldioctadecylammonium bromide), sugars or sugar derivatives (e.g., cyclodextrin), nucleic acids, polymers (e.g., heparin, polyethylene glycol and poloxamer), mucolytic agents (e.g., N-acetylcysteine, mugwort, bromelain, papain, clerodendrum, acetylcysteine, bromhexine, carbocisteine, eprazinone, mesna, ambroxol, sobrerol, domiodol, letosteine, stepronin, tiopronin, gelsolin, thymosin β 4 dornase alfa, neltenexine, erdosteine) and various DNases including rhDNase.
- [0958] In some embodiments, the mucus penetrating LNP can be a hypotonic formulation comprising a mucosal penetration enhancing coating. The formulation can be hypotonic for the epithelium to which it is being delivered. Non-limiting examples of hypotonic formulations can be found in, e.g., Intl. Pub. No. WO2013110028, herein incorporated by reference in its entirety.
- [0959] In some embodiments, the polynucleotide described herein is formulated as a lipoplex, such as, without limitation, the ATUPLEXTM system, the DACC system, the DBTC system and other siRNA-lipoplex technology from Silence Therapeutics (London, United Kingdom), STEMFECTTM from STEMGENT® (Cambridge, MA), and polyethylenimine (PEI) or protamine-based targeted and non-targeted delivery of nucleic acids (Aleku et al. Cancer Res. 2008 68:9788-9798; Strumberg et al. Int J Clin Pharmacol Ther 2012 50:76-78; Santel et al., Gene Ther 2006 13:1222-1234; Santel et al., Gene Ther 2006 13:1360-1370; Gutbier et al., Pulm Pharmacol. Ther. 2010 23:334-344; Kaufmann et al. Microvasc Res 2010 80:286-293; Weide et al. J Immunother. 2009 32:498-507; Weide et al. J Immunother. 2008 31:180-188; Pascolo Expert Opin. Biol. Ther. 4:1285-1294; Fotin-Mleczek et al., 2011 J. Immunother. 34:1-15; Song et al., Nature Biotechnol. 2005, 23:709-717; Peer et al., Proc Natl Acad Sci U S A. 2007 6;104:4095-4100; deFougerolles Hum Gene Ther. 2008 19:125-132; all of which are incorporated herein by reference in its entirety).
- [0960] In some embodiments, the polynucleotides described herein are formulated as a solid lipid nanoparticle (SLN), which can be spherical with an average diameter between 10 to 1000 nm. SLN possess a solid lipid core matrix that can solubilize lipophilic molecules and can be stabilized with surfactants and/or emulsifiers. Exemplary SLN can

be those as described in Intl. Pub. No. WO2013105101, herein incorporated by reference in its entirety.

[0961] In some embodiments, the polynucleotides described herein can be formulated for controlled release and/or targeted delivery. As used herein, "controlled release" refers to a pharmaceutical composition or compound release profile that conforms to a particular pattern of release to effect a therapeutic outcome. In one embodiment, the polynucleotides can be encapsulated into a delivery agent described herein and/or known in the art for controlled release and/or targeted delivery. As used herein, the term "encapsulate" means to enclose, surround or encase. As it relates to the formulation of the compounds of the invention, encapsulation can be substantial, complete or partial. The term "substantially encapsulated" means that at least greater than 50, 60, 70, 80, 85, 90, 95, 96, 97, 98, 99, 99.9, 99.9 or greater than 99.999% of the pharmaceutical composition or compound of the invention can be enclosed, surrounded or encased within the delivery agent. "Partially encapsulation" means that less than 10, 10, 20, 30, 40 50 or less of the pharmaceutical composition or compound of the invention can be enclosed, surrounded or encased within the delivery agent.

[0962] Advantageously, encapsulation can be determined by measuring the escape or the activity of the pharmaceutical composition or compound of the invention using fluorescence and/or electron micrograph. For example, at least 1, 5, 10, 20, 30, 40, 50, 60, 70, 80, 85, 90, 95, 96, 97, 98, 99, 99.9, 99.99 or greater than 99.99% of the pharmaceutical composition or compound of the invention are encapsulated in the delivery agent.

[0963] In some embodiments, the polynucleotide controlled release formulation can include at least one controlled release coating (e.g., OPADRY®, EUDRAGIT RL®, EUDRAGIT RS® and cellulose derivatives such as ethylcellulose aqueous dispersions (AQUACOAT® and SURELEASE®)). In some embodiments, the polynucleotide controlled release formulation can comprise a polymer system as described in U.S. Pub. No. US20130130348, or a PEG and/or PEG related polymer derivative as described in U.S. Pat. No. 8,404,222, each of which is incorporated by reference in its entirety.

[0964] In some embodiments, the polynucleotides described herein can be encapsulated in a therapeutic nanoparticle, referred to herein as "therapeutic nanoparticle polynucleotides." Therapeutic nanoparticles can be formulated by methods described in, e.g., Intl. Pub. Nos. WO2010005740, WO2010030763, WO2010005721,

WO2010005723, and WO2012054923; and U.S. Pub. Nos. US20110262491, US20100104645, US20100087337, US20100068285, US20110274759, US20100068286, US20120288541, US20120140790, US20130123351 and US20130230567; and U.S. Pat. Nos. 8,206,747, 8,293,276, 8,318,208 and 8,318,211, each of which is herein incorporated by reference in its entirety.

[0965] In some embodiments, the therapeutic nanoparticle polynucleotide can be formulated for sustained release. As used herein, "sustained release" refers to a pharmaceutical composition or compound that conforms to a release rate over a specific period of time. The period of time can include, but is not limited to, hours, days, weeks, months and years. As a non-limiting example, the sustained release nanoparticle of the polynucleotides described herein can be formulated as disclosed in Intl. Pub. No. WO2010075072 and U.S. Pub. Nos. US20100216804, US20110217377, US20120201859 and US20130150295, each of which is herein incorporated by reference in their entirety.

[0966] In some embodiments, the therapeutic nanoparticle polynucleotide can be formulated to be target specific, such as those described in Intl. Pub. Nos. WO2008121949, WO2010005726, WO2010005725, WO2011084521 and WO2011084518; and U.S. Pub. Nos. US20100069426, US20120004293 and US20100104655, each of which is herein incorporated by reference in its entirety.

[0967] The LNPs can be prepared using microfluidic mixers or micromixers. Exemplary microfluidic mixers can include, but are not limited to, a slit interdigital micromixer including, but not limited to those manufactured by Microinnova (Allerheiligen bei Wildon, Austria) and/or a staggered herringbone micromixer (SHM) (see Zhigaltsevet al., "Bottom-up design and synthesis of limit size lipid nanoparticle systems with aqueous and triglyceride cores using millisecond microfluidic mixing," *Langmuir* 28:3633-40 (2012); Belliveau et al., "Microfluidic synthesis of highly potent limit-size lipid nanoparticles for in vivo delivery of siRNA," *Molecular Therapy-Nucleic Acids* 1:e37 (2012); Chen et al., "Rapid discovery of potent siRNA-containing lipid nanoparticles enabled by controlled microfluidic formulation," *J. Am. Chem. Soc.* 134(16):6948-51 (2012); each of which is herein incorporated by reference in its entirety). Exemplary micromixers include Slit Interdigital Microstructured Mixer (SIMM-V2) or a Standard Slit Interdigital Micro Mixer (SSIMM) or Caterpillar (CPMM) or Impinging-jet (IJMM,) from the Institut für Mikrotechnik Mainz GmbH, Mainz Germany. In some embodiments,

methods of making LNP using SHM further comprise mixing at least two input streams wherein mixing occurs by microstructure-induced chaotic advection (MICA). According to this method, fluid streams flow through channels present in a herringbone pattern causing rotational flow and folding the fluids around each other. This method can also comprise a surface for fluid mixing wherein the surface changes orientations during fluid cycling. Methods of generating LNPs using SHM include those disclosed in U.S. Pub. Nos. US20040262223 and US20120276209, each of which is incorporated herein by reference in their entirety.

[0968] In some embodiments, the polynucleotides described herein can be formulated in lipid nanoparticles using microfluidic technology (see Whitesides, George M., "The Origins and the Future of Microfluidics," *Nature* 442: 368-373 (2006); and Abraham et al., "Chaotic Mixer for Microchannels," *Science* 295: 647-651 (2002); each of which is herein incorporated by reference in its entirety). In some embodiments, the polynucleotides can be formulated in lipid nanoparticles using a micromixer chip such as, but not limited to, those from Harvard Apparatus (Holliston, MA) or Dolomite Microfluidics (Royston, UK). A micromixer chip can be used for rapid mixing of two or more fluid streams with a split and recombine mechanism.

[0969] In some embodiments, the polynucleotides described herein can be formulated in lipid nanoparticles having a diameter from about 1 nm to about 100 nm such as, but not limited to, about 1 nm to about 20 nm, from about 1 nm to about 30 nm, from about 1 nm to about 40 nm, from about 1 nm to about 50 nm, from about 1 nm to about 60 nm, from about 1 nm to about 70 nm, from about 1 nm to about 80 nm, from about 1 nm to about 90 nm, from about 5 nm to about 100 nm, from about 5 nm to about 10 nm, about 5 nm to about 20 nm, from about 5 nm to about 30 nm, from about 5 nm to about 40 nm, from about 5 nm to about 50 nm, from about 5 nm to about 60 nm, from about 5 nm to about 70 nm, from about 5 nm to about 80 nm, from about 5 nm to about 90 nm, about 10 to about 20 nm, about 10 to about 30 nm, about 10 to about 40 nm, about 10 to about 50 nm, about 10 to about 60 nm, about 10 to about 70 nm, about 10 to about 80 nm, about 10 to about 90 nm, about 20 to about 30 nm, about 20 to about 40 nm, about 20 to about 50 nm, about 20 to about 60 nm, about 20 to about 70 nm, about 20 to about 80 nm, about 20 to about 90 nm, about 20 to about 100 nm, about 30 to about 40 nm, about 30 to about 50 nm, about 30 to about 60 nm, about 30 to about 70 nm, about 30 to about 80 nm, about 30 to about 90 nm, about 30 to about 100 nm, about 40 to about 50 nm, about 40 to about

60 nm, about 40 to about 70 nm, about 40 to about 80 nm, about 40 to about 90 nm, about 40 to about 100 nm, about 50 to about 60 nm, about 50 to about 70 nm about 50 to about 80 nm, about 50 to about 90 nm, about 50 to about 100 nm, about 60 to about 70 nm, about 60 to about 80 nm, about 60 to about 90 nm, about 60 to about 100 nm, about 70 to about 80 nm, about 70 to about 90 nm, about 70 to about 100 nm, about 80 to about 90 nm, about 80 to about 100 nm and/or about 90 to about 100 nm.

[0970] In some embodiments, the lipid nanoparticles can have a diameter from about 10 to 500 nm. In one embodiment, the lipid nanoparticle can have a diameter greater than 100 nm, greater than 150 nm, greater than 200 nm, greater than 250 nm, greater than 300 nm, greater than 350 nm, greater than 400 nm, greater than 450 nm, greater than 500 nm, greater than 550 nm, greater than 600 nm, greater than 650 nm, greater than 700 nm, greater than 750 nm, greater than 800 nm, greater than 850 nm, greater than 900 nm, greater than 950 nm or greater than 1000 nm.

[0971] In some embodiments, the polynucleotides can be delivered using smaller LNPs. Such particles can comprise a diameter from below 0.1 μm up to 100 nm such as, but not limited to, less than 0.1 μm , less than 1.0 μm , less than 5 μm , less than 10 μm , less than 15 μm , less than 20 μm , less than 25 μm , less than 30 μm , less than 35 μm , less than 40 μm , less than 50 μm , less than 55 μm , less than 60 μm , less than 65 μm , less than 70 μm , less than 75 μm , less than 80 μm , less than 85 μm , less than 90 μm , less than 95 μm , less than 100 μm , less than 125 μm , less than 150 μm , less than 175 μm , less than 200 μm , less than 225 μm , less than 250 μm , less than 275 μm , less than 300 μm , less than 325 μm , less than 350 μm , less than 375 μm , less than 400 μm , less than 425 μm , less than 450 μm , less than 475 μm , less than 500 μm , less than 525 μm , less than 550 μm , less than 575 μm , less than 600 μm , less than 625 μm , less than 650 μm , less than 675 μm , less than 700 μm , less than 725 μm , less than 750 μm , less than 775 μm , less than 800 μm , less than 825 μm , less than 850 μm , less than 875 μm , less than 900 μm , less than 925 μm , less than 950 μm , or less than 975 μm .

[0972] The nanoparticles and microparticles described herein can be geometrically engineered to modulate macrophage and/or the immune response. The geometrically engineered particles can have varied shapes, sizes and/or surface charges to incorporate the polynucleotides described herein for targeted delivery such as, but not limited to, pulmonary delivery (see, e.g., Intl. Pub. No. WO2013082111, herein incorporated by reference in its entirety). Other physical features the geometrically engineering particles

can include, but are not limited to, fenestrations, angled arms, asymmetry and surface roughness, charge that can alter the interactions with cells and tissues.

[0973] In some embodiment, the nanoparticles described herein are stealth nanoparticles or target-specific stealth nanoparticles such as, but not limited to, those described in U.S. Pub. No. US20130172406, herein incorporated by reference in its entirety. The stealth or target-specific stealth nanoparticles can comprise a polymeric matrix, which can comprise two or more polymers such as, but not limited to, polyethylenes, polycarbonates, polyanhydrides, polyhydroxyacids, polypropylfumerates, polycaprolactones, polyamides, polyacetals, polyethers, polyesters, poly(orthoesters), polycyanoacrylates, polyvinyl alcohols, polyurethanes, polyphosphazenes, polyacrylates, polymethacrylates, polycyanoacrylates, polyureas, polystyrenes, polyamines, polyesters, polyanhydrides, polyethers, polyurethanes, polymethacrylates, polyacrylates, polycyanoacrylates, or combinations thereof.

b. Lipidoids

[0974] In some embodiments, the compositions or formulations of the present disclosure comprise a delivery agent, e.g., a lipidoid. The polynucleotides described herein (e.g., a polynucleotide comprising a nucleotide sequence encoding an ACADVL polypeptide) can be formulated with lipidoids. Complexes, micelles, liposomes or particles can be prepared containing these lipidoids and therefore to achieve an effective delivery of the polynucleotide, as judged by the production of an encoded protein, following the injection of a lipidoid formulation via localized and/or systemic routes of administration. Lipidoid complexes of polynucleotides can be administered by various means including, but not limited to, intravenous, intramuscular, or subcutaneous routes.

[0975] The synthesis of lipidoids is described in literature (see Mahon et al., *Bioconjug. Chem.* 2010 21:1448-1454; Schroeder et al., *J Intern Med.* 2010 267:9-21; Akinc et al., *Nat Biotechnol.* 2008 26:561-569; Love et al., *Proc Natl Acad Sci U S A.* 2010 107:1864-1869; Siegwart et al., *Proc Natl Acad Sci U S A.* 2011 108:12996-3001; all of which are incorporated herein in their entireties).

[0976] Formulations with the different lipidoids, including, but not limited to penta[3-(1-laurylaminopropionyl)]-triethylenetetramine hydrochloride (TETA-5LAP; also known as 98N12-5, see Murugaiah et al., *Analytical Biochemistry*, 401:61 (2010)), C12-200 (including derivatives and variants), and MD1, can be tested for in vivo activity. The

lipidoid "98N12-5" is disclosed by Akinc et al., *Mol Ther.* 2009 17:872-879. The lipidoid "C12-200" is disclosed by Love et al., *Proc Natl Acad Sci U S A.* 2010 107:1864-1869 and Liu and Huang, *Molecular Therapy.* 2010 669-670. Each of the references is herein incorporated by reference in its entirety.

[0977] In one embodiment, the polynucleotides described herein can be formulated in an aminoalcohol lipidoid. Aminoalcohol lipidoids can be prepared by the methods described in U.S. Patent No. 8,450,298 (herein incorporated by reference in its entirety).

[0978] The lipidoid formulations can include particles comprising either 3 or 4 or more components in addition to polynucleotides. Lipidoids and polynucleotide formulations comprising lipidoids are described in Intl. Pub. No. WO 2015051214 (herein incorporated by reference in its entirety).

c. Hyaluronidase

[0979] In some embodiments, the polynucleotides described herein (e.g., a polynucleotide comprising a nucleotide sequence encoding an ACADVL polypeptide) and hyaluronidase for injection (e.g., intramuscular or subcutaneous injection). Hyaluronidase catalyzes the hydrolysis of hyaluronan, which is a constituent of the interstitial barrier. Hyaluronidase lowers the viscosity of hyaluronan, thereby increases tissue permeability (Frost, *Expert Opin. Drug Deliv.* (2007) 4:427-440). Alternatively, the hyaluronidase can be used to increase the number of cells exposed to the polynucleotides administered intramuscularly or subcutaneously.

d. Nanoparticle Mimics

[0980] In some embodiments, the polynucleotides described herein (e.g., a polynucleotide comprising a nucleotide sequence encoding an ACADVL polypeptide) is encapsulated within and/or absorbed to a nanoparticle mimic. A nanoparticle mimic can mimic the delivery function organisms or particles such as, but not limited to, pathogens, viruses, bacteria, fungus, parasites, prions and cells. As a non-limiting example, the polynucleotides described herein can be encapsulated in a non-viron particle that can mimic the delivery function of a virus (see e.g., Intl. Pub. No. WO2012006376 and U.S. Pub. Nos. US20130171241 and US20130195968, each of which is herein incorporated by reference in its entirety).

e. Nanotubes

[0981] In some embodiments, the compositions or formulations of the present disclosure comprise the polynucleotides described herein (e.g., a polynucleotide comprising a nucleotide sequence encoding an ACADV L polypeptide) attached or otherwise bound to (e.g., through steric, ionic, covalent and/or other forces) at least one nanotube, such as, but not limited to, rosette nanotubes, rosette nanotubes having twin bases with a linker, carbon nanotubes and/or single-walled carbon nanotubes. Nanotubes and nanotube formulations comprising a polynucleotide are described in, e.g., Intl. Pub. No. WO2014152211, herein incorporated by reference in its entirety.

f. Self-Assembled Nanoparticles, or Self-Assembled Macromolecules

[0982] In some embodiments, the compositions or formulations of the present disclosure comprise the polynucleotides described herein (e.g., a polynucleotide comprising a nucleotide sequence encoding an ACADV L polypeptide) in self-assembled nanoparticles, or amphiphilic macromolecules (AMs) for delivery. AMs comprise biocompatible amphiphilic polymers that have an alkylated sugar backbone covalently linked to poly(ethylene glycol). In aqueous solution, the AMs self-assemble to form micelles. Nucleic acid self-assembled nanoparticles are described in Intl. Appl. No. PCT/US2014/027077, and AMs and methods of forming AMs are described in U.S. Pub. No. US20130217753, each of which is herein incorporated by reference in its entirety.

g. Inorganic Nanoparticles, Semi-Conductive and Metallic Nanoparticles

[0983] In some embodiments, the compositions or formulations of the present disclosure comprise the polynucleotides described herein (e.g., a polynucleotide comprising a nucleotide sequence encoding an ACADV L polypeptide) in inorganic nanoparticles, or water-dispersible nanoparticles comprising a semiconductive or metallic material. The inorganic nanoparticles can include, but are not limited to, clay substances that are water swellable. The water-dispersible nanoparticles can be hydrophobic or hydrophilic nanoparticles. As a non-limiting example, the inorganic, semi-conductive and metallic nanoparticles are described in, e.g., U.S. Pat. Nos. 5,585,108 and 8,257,745; and U.S. Pub. Nos. US20120228565, US 20120265001 and US 20120283503, each of which is herein incorporated by reference in their entirety.

h. Surgical Sealants: Gels and Hydrogels

[0984] In some embodiments, the compositions or formulations of the present disclosure comprise the polynucleotides described herein (e.g., a polynucleotide comprising a nucleotide sequence encoding an ACADVL polypeptide) in a surgical sealant. Surgical sealants such as gels and hydrogels are described in Intl. Appl. No. PCT/US2014/027077, herein incorporated by reference in its entirety.

i. Suspension formulations

[0985] In some embodiments, the compositions or formulations of the present disclosure comprise the polynucleotides described herein (e.g., a polynucleotide comprising a nucleotide sequence encoding an ACADVL polypeptide) in suspensions. In some embodiments, suspensions comprise a polynucleotide, water immiscible oil depots, surfactants and/or co-surfactants and/or co-solvents. Suspensions can be formed by first preparing an aqueous solution of a polynucleotide and an oil-based phase comprising one or more surfactants, and then mixing the two phases (aqueous and oil-based).

[0986] Exemplary oils for suspension formulations can include, but are not limited to, sesame oil and Miglyol (comprising esters of saturated coconut and palm kernel oil-derived caprylic and capric fatty acids and glycerin or propylene glycol), corn oil, soybean oil, peanut oil, beeswax and/or palm seed oil. Exemplary surfactants can include, but are not limited to Cremophor, polysorbate 20, polysorbate 80, polyethylene glycol, transcitol, Capmul®, labrasol, isopropyl myristate, and/or Span 80. In some embodiments, suspensions can comprise co-solvents including, but not limited to ethanol, glycerol and/or propylene glycol.

[0987] In some embodiments, suspensions can provide modulation of the release of the polynucleotides into the surrounding environment by diffusion from a water immiscible depot followed by resolubilization into a surrounding environment (e.g., an aqueous environment).

[0988] In some embodiments, the polynucleotides can be formulated such that upon injection, an emulsion forms spontaneously (e.g., when delivered to an aqueous phase), which can provide a high surface area to volume ratio for release of polynucleotides from an oil phase to an aqueous phase. In some embodiments, the polynucleotide is formulated in a nanoemulsion, which can comprise a liquid hydrophobic core surrounded by or coated with a lipid or surfactant layer. Exemplary nanoemulsions and their preparations

are described in, e.g., U.S. Pat. No. 8,496,945, herein incorporated by reference in its entirety.

j. Cations and Anions

[0989] In some embodiments, the compositions or formulations of the present disclosure comprise the polynucleotides described herein (e.g., a polynucleotide comprising a nucleotide sequence encoding an ACADVL polypeptide) and a cation or anion, such as Zn^{2+} , Ca^{2+} , Cu^{2+} , Mg^{2+} and combinations thereof. Exemplary formulations can include polymers and a polynucleotide complexed with a metal cation as described in, e.g., U.S. Pat. Nos. 6,265,389 and 6,555,525, each of which is herein incorporated by reference in its entirety. In some embodiments, cationic nanoparticles can contain a combination of divalent and monovalent cations. The delivery of polynucleotides in cationic nanoparticles or in one or more depot comprising cationic nanoparticles can improve polynucleotide bioavailability by acting as a long-acting depot and/or reducing the rate of degradation by nucleases.

k. Molded Nanoparticles and Microparticles

[0990] In some embodiments, the compositions or formulations of the present disclosure comprise the polynucleotides described herein (e.g., a polynucleotide comprising a nucleotide sequence encoding an ACADVL polypeptide) in molded nanoparticles in various sizes, shapes and chemistry. For example, the nanoparticles and/or microparticles can be made using the PRINT® technology by LIQUIDA TECHNOLOGIES® (Morrisville, NC) (e.g., International Pub. No. WO2007024323, herein incorporated by reference in its entirety).

[0991] In some embodiments, the polynucleotides described herein (e.g., a polynucleotide comprising a nucleotide sequence encoding an ACADVL polypeptide) is formulated in microparticles. The microparticles can contain a core of the polynucleotide and a cortex of a biocompatible and/or biodegradable polymer, including but not limited to, poly(α -hydroxy acid), a polyhydroxy butyric acid, a polycaprolactone, a polyorthoester and a polyanhydride. The microparticle can have adsorbent surfaces to adsorb polynucleotides. The microparticles can have a diameter of from at least 1 micron to at least 100 microns (e.g., at least 1 micron, at least 10 micron, at least 20 micron, at least 30 micron, at least 50 micron, at least 75 micron, at least 95 micron, and at least 100 micron). In some embodiment, the compositions or formulations of the present disclosure

are microemulsions comprising microparticles and polynucleotides. Exemplary microparticles, microemulsions and their preparations are described in, e.g., U.S. Pat. Nos. 8,460,709, 8,309,139 and 8,206,749; U.S. Pub. Nos. US20130129830, US2013195923 and US20130195898; and Intl. Pub. No. WO2013075068, each of which is herein incorporated by reference in its entirety.

l. NanoJackets and NanoLiposomes

[0992] In some embodiments, the compositions or formulations of the present disclosure comprise the polynucleotides described herein (e.g., a polynucleotide comprising a nucleotide sequence encoding an ACADVL polypeptide) in NanoJackets and NanoLiposomes by Keystone Nano (State College, PA). NanoJackets are made of materials that are naturally found in the body including calcium, phosphate and can also include a small amount of silicates. Nanojackets can have a size ranging from 5 to 50 nm.

[0993] NanoLiposomes are made of lipids such as, but not limited to, lipids that naturally occur in the body. NanoLiposomes can have a size ranging from 60-80 nm. In some embodiments, the polynucleotides disclosed herein are formulated in a NanoLiposome such as, but not limited to, Ceramide NanoLiposomes.

m. Cells or Minicells

[0994] In some embodiments, the compositions or formulations of the present disclosure comprise the polynucleotides described herein (e.g., a polynucleotide comprising a nucleotide sequence encoding an ACADVL polypeptide) that is transfected ex vivo into cells, which are subsequently transplanted into a subject. Cell-based formulations of the polynucleotide disclosed herein can be used to ensure cell transfection (e.g., in the cellular carrier), alter the biodistribution of the polynucleotide (e.g., by targeting the cell carrier to specific tissues or cell types), and/or increase the translation of encoded protein.

[0995] Exemplary cells include, but are not limited to, red blood cells, virosomes, and electroporated cells (see e.g., Godfrin et al., Expert Opin Biol Ther. 2012 12:127-133; Fang et al., Expert Opin Biol Ther. 2012 12:385-389; Hu et al., Proc Natl Acad Sci U S A. 2011 108:10980-10985; Lund et al., Pharm Res. 2010 27:400-420; Huckriede et al., J Liposome Res. 2007;17:39-47; Cusi, Hum Vaccin. 2006 2:1-7; de Jonge et al., Gene Ther. 2006 13:400-411; all of which are herein incorporated by reference in its entirety).

[0996] A variety of methods are known in the art and are suitable for introduction of nucleic acid into a cell, including viral and non-viral mediated techniques. Examples of

typical non-viral mediated techniques include, but are not limited to, electroporation, calcium phosphate mediated transfer, nucleofection, sonoporation, heat shock, magnetofection, liposome mediated transfer, microinjection, microprojectile mediated transfer (nanoparticles), cationic polymer mediated transfer (DEAE-dextran, polyethylenimine, polyethylene glycol (PEG) and the like) or cell fusion.

- [0997] In some embodiments, the polynucleotides described herein can be delivered in synthetic virus-like particles (VLPs) synthesized by the methods as described in Intl. Pub Nos. WO2011085231 and WO2013116656; and U.S. Pub. No. 20110171248, each of which is herein incorporated by reference in its entirety.
- [0998] The technique of sonoporation, or cellular sonication, is the use of sound (e.g., ultrasonic frequencies) for modifying the permeability of the cell plasma membrane. Sonoporation methods are known to deliver nucleic acids in vivo (Yoon and Park, *Expert Opin Drug Deliv.* 2010 7:321-330; Postema and Gilja, *Curr Pharm Biotechnol.* 2007 8:355-361; Newman and Bettinger, *Gene Ther.* 2007 14:465-475; U.S. Pub. Nos. US20100196983 and US20100009424; all herein incorporated by reference in their entirety).
- [0999] In some embodiments, the polynucleotides described herein can be delivered by electroporation. Electroporation techniques are known to deliver nucleic acids in vivo and clinically (Andre et al., *Curr Gene Ther.* 2010 10:267-280; Chiarella et al., *Curr Gene Ther.* 2010 10:281-286; Hojman, *Curr Gene Ther.* 2010 10:128-138; all herein incorporated by reference in their entirety). Electroporation devices are sold by many companies worldwide including, but not limited to BTX® Instruments (Holliston, MA) (e.g., the AgilePulse In Vivo System) and Inovio (Blue Bell, PA) (e.g., Inovio SP-5P intramuscular delivery device or the CELLECTRA® 3000 intradermal delivery device).
- [1000] In some embodiments, the cells are selected from the group consisting of mammalian cells, bacterial cells, plant, microbial, algal and fungal cells. In some embodiments, the cells are mammalian cells, such as, but not limited to, human, mouse, rat, goat, horse, rabbit, hamster or cow cells. In a further embodiment, the cells can be from an established cell line, including, but not limited to, HeLa, NS0, SP2/0, KEK 293T, Vero, Caco, Caco-2, MDCK, COS-1, COS-7, K562, Jurkat, CHO-K1, DG44, CHOK1SV, CHO-S, Huvec, CV-1, Huh-7, NIH3T3, HEK293, 293, A549, HepG2, IMR-90, MCF-7, U-20S, Per.C6, SF9, SF21 or Chinese Hamster Ovary (CHO) cells.

- [1001] In certain embodiments, the cells are fungal cells, such as, but not limited to, Chrysosporium cells, Aspergillus cells, Trichoderma cells, Dictyostelium cells, Candida cells, Saccharomyces cells, Schizosaccharomyces cells, and Penicillium cells.
- [1002] In certain embodiments, the cells are bacterial cells such as, but not limited to, E. coli, B. subtilis, or BL21 cells. Primary and secondary cells to be transfected by the methods of the invention can be obtained from a variety of tissues and include, but are not limited to, all cell types that can be maintained in culture. The primary and secondary cells include, but are not limited to, fibroblasts, keratinocytes, epithelial cells (e.g., mammary epithelial cells, intestinal epithelial cells), endothelial cells, glial cells, neural cells, formed elements of the blood (e.g., lymphocytes, bone marrow cells), muscle cells and precursors of these somatic cell types. Primary cells can also be obtained from a donor of the same species or from another species (e.g., mouse, rat, rabbit, cat, dog, pig, cow, bird, sheep, goat, horse).
- [1003] In some embodiments, the compositions or formulations of the present disclosure comprise the polynucleotides described herein in bacterial minicells. As a non-limiting example, bacterial minicells can be those described in Intl. Pub. No. WO2013088250 or U.S. Pub. No. US20130177499, each of which is herein incorporated by reference in its entirety.
- n. Semi-solid Compositions
- [1004] In some embodiments, the compositions or formulations of the present disclosure comprise the polynucleotides described herein (e.g., a polynucleotide comprising a nucleotide sequence encoding an ACADV L polypeptide) in a hydrophobic matrix to form a semi-solid or paste-like composition. As a non-limiting example, the semi-solid or paste-like composition can be made by the methods described in Intl. Pub. No. WO201307604, herein incorporated by reference in its entirety.
- o. Exosomes
- [1005] In some embodiments, the compositions or formulations of the present disclosure comprise the polynucleotides described herein (e.g., a polynucleotide comprising a nucleotide sequence encoding an ACADV L polypeptide) in exosomes, which can be loaded with at least one polynucleotide and delivered to cells, tissues and/or organisms. As a non-limiting example, the polynucleotides can be loaded in the exosomes as

described in Intl. Pub. No. WO2013084000, herein incorporated by reference in its entirety.

p. Silk-Based Delivery

[1006] In some embodiments, the compositions or formulations of the present disclosure comprise the polynucleotides described herein (e.g., a polynucleotide comprising a nucleotide sequence encoding an ACADVL polypeptide) that is formulated for silk-based delivery. The silk-based delivery system can be formed by contacting a silk fibroin solution with a polynucleotide described herein. As a non-limiting example, a sustained release silk-based delivery system and methods of making such system are described in U.S. Pub. No. US20130177611, herein incorporated by reference in its entirety.

q. Amino Acid Lipids

[1007] In some embodiments, the compositions or formulations of the present disclosure comprise the polynucleotides described herein (e.g., a polynucleotide comprising a nucleotide sequence encoding an ACADVL polypeptide) that is formulation with an amino acid lipid. Amino acid lipids are lipophilic compounds comprising an amino acid residue and one or more lipophilic tails. Non-limiting examples of amino acid lipids and methods of making amino acid lipids are described in U.S. Pat. No. 8,501,824. The amino acid lipid formulations can deliver a polynucleotide in releasable form that comprises an amino acid lipid that binds and releases the polynucleotides. As a non-limiting example, the release of the polynucleotides described herein can be provided by an acid-labile linker as described in, e.g., U.S. Pat. Nos. 7,098,032, 6,897,196, 6,426,086, 7,138,382, 5,563,250, and 5,505,931, each of which is herein incorporated by reference in its entirety.

r. Microvesicles

[1008] In some embodiments, the compositions or formulations of the present disclosure comprise the polynucleotides described herein (e.g., a polynucleotide comprising a nucleotide sequence encoding an ACADVL polypeptide) in a microvesicle formulation. Exemplary microvesicles include those described in U.S. Pub. No. US20130209544 (herein incorporated by reference in its entirety). In some embodiments, the microvesicle is an ARRDC1-mediated microvesicles (ARMMs) as described in Intl. Pub. No. WO2013119602 (herein incorporated by reference in its entirety).

s. Interpolyelectrolyte Complexes

[1009] In some embodiments, the compositions or formulations of the present disclosure comprise the polynucleotides described herein (e.g., a polynucleotide comprising a nucleotide sequence encoding an ACADVL polypeptide) in an interpolyelectrolyte complex. Interpolyelectrolyte complexes are formed when charge-dynamic polymers are complexed with one or more anionic molecules. Non-limiting examples of charge-dynamic polymers and interpolyelectrolyte complexes and methods of making interpolyelectrolyte complexes are described in U.S. Pat. No. 8,524,368, herein incorporated by reference in its entirety.

t. Crystalline Polymeric Systems

[1010] In some embodiments, the compositions or formulations of the present disclosure comprise the polynucleotides described herein (e.g., a polynucleotide comprising a nucleotide sequence encoding an ACADVL polypeptide) in crystalline polymeric systems. Crystalline polymeric systems are polymers with crystalline moieties and/or terminal units comprising crystalline moieties. Exemplary polymers are described in U.S. Pat. No. 8,524,259 (herein incorporated by reference in its entirety).

u. Polymers, Biodegradable Nanoparticles, and Core-Shell Nanoparticles

[1011] In some embodiments, the compositions or formulations of the present disclosure comprise the polynucleotides described herein (e.g., a polynucleotide comprising a nucleotide sequence encoding an ACADVL polypeptide) and a natural and/or synthetic polymer. The polymers include, but not limited to, polyethenes, polyethylene glycol (PEG), poly(L-lysine)(PLL), PEG grafted to PLL, cationic lipopolymer, biodegradable cationic lipopolymer, polyethyleneimine (PEI), cross-linked branched poly(alkylene imines), a polyamine derivative, a modified poloxamer, elastic biodegradable polymer, biodegradable copolymer, biodegradable polyester copolymer, biodegradable polyester copolymer, multiblock copolymers, poly[α -(4-aminobutyl)-L-glycolic acid] (PAGA), biodegradable cross-linked cationic multi-block copolymers, polycarbonates, polyanhydrides, polyhydroxyacids, polypropylfumerates, polycaprolactones, polyamides, polyacetals, polyethers, polyesters, poly(orthoesters), polycyanoacrylates, polyvinyl alcohols, polyurethanes, polyphosphazenes, polyureas, polystyrenes, polyamines, polylysine, poly(ethylene imine), poly(serine ester), poly(L-lactide-co-L-lysine), poly(4-

hydroxy-L-proline ester), amine-containing polymers, dextran polymers, dextran polymer derivatives or combinations thereof.

- [1012] Exemplary polymers include, DYNAMIC POLYCONJUGATE® (Arrowhead Research Corp., Pasadena, CA) formulations from MIRUS® Bio (Madison, WI) and Roche Madison (Madison, WI), PHASERXTM polymer formulations such as, without limitation, SMARTT POLYMER TECHNOLOGY™ (PHASERX®, Seattle, WA), DMRI/DOPE, poloxamer, VAXFECTIN® adjuvant from Vical (San Diego, CA), chitosan, cyclodextrin from Calando Pharmaceuticals (Pasadena, CA), dendrimers and poly(lactic-co-glycolic acid) (PLGA) polymers. RONDEL™ (RNAi/Oligonucleotide Nanoparticle Delivery) polymers (Arrowhead Research Corporation, Pasadena, CA) and pH responsive co-block polymers such as PHASERX® (Seattle, WA).
- [1013] The polymer formulations allow a sustained or delayed release of the polynucleotide (e.g., following intramuscular or subcutaneous injection). The altered release profile for the polynucleotide can result in, for example, translation of an encoded protein over an extended period of time. The polymer formulation can also be used to increase the stability of the polynucleotide. Sustained release formulations can include, but are not limited to, PLGA microspheres, ethylene vinyl acetate (EVAc), poloxamer, GELSITE® (Nanotherapeutics, Inc. Alachua, FL), HYLENEX® (Halozyme Therapeutics, San Diego CA), surgical sealants such as fibrinogen polymers (Ethicon Inc. Cornelia, GA), TISSELL® (Baxter International, Inc. Deerfield, IL), PEG-based sealants, and COSEAL® (Baxter International, Inc. Deerfield, IL).
- [1014] As a non-limiting example modified mRNA can be formulated in PLGA microspheres by preparing the PLGA microspheres with tunable release rates (e.g., days and weeks) and encapsulating the modified mRNA in the PLGA microspheres while maintaining the integrity of the modified mRNA during the encapsulation process. EVAc are non-biodegradable, biocompatible polymers that are used extensively in pre-clinical sustained release implant applications (e.g., extended release products Ocusert a pilocarpine ophthalmic insert for glaucoma or progestasert a sustained release progesterone intrauterine device; transdermal delivery systems Testoderm, Duragesic and Selegiline; catheters). Poloxamer F-407 NF is a hydrophilic, non-ionic surfactant triblock copolymer of polyoxyethylene-polyoxypropylene-polyoxyethylene having a low viscosity at temperatures less than 5°C and forms a solid gel at temperatures greater than 15°C.

- [1015] As a non-limiting example, the polynucleotides described herein can be formulated with the polymeric compound of PEG grafted with PLL as described in U.S. Pat. No. 6,177,274. As another non-limiting example, the polynucleotides described herein can be formulated with a block copolymer such as a PLGA-PEG block copolymer (see e.g., U.S. Pub. No. US20120004293 and U.S. Pat. Nos. 8,236,330 and 8,246,968), or a PLGA-PEG-PLGA block copolymer (see e.g., U.S. Pat. No. 6,004,573). Each of the references is herein incorporated by reference in its entirety.
- [1016] In some embodiments, the polynucleotides described herein can be formulated with at least one amine-containing polymer such as, but not limited to polylysine, polyethylene imine, poly(amidoamine) dendrimers, poly(amine-co-esters) or combinations thereof. Exemplary polyamine polymers and their use as delivery agents are described in, e.g., U.S. Pat. Nos. 8,460,696, 8,236,280, each of which is herein incorporated by reference in its entirety.
- [1017] In some embodiments, the polynucleotides described herein can be formulated in a biodegradable cationic lipopolymer, a biodegradable polymer, or a biodegradable copolymer, a biodegradable polyester copolymer, a biodegradable polyester polymer, a linear biodegradable copolymer, PAGA, a biodegradable cross-linked cationic multi-block copolymer or combinations thereof as described in, e.g., U.S. Pat. Nos. 6,696,038, 6,517,869, 6,267,987, 6,217,912, 6,652,886, 8,057,821, and 8,444,992; U.S. Pub. Nos. US20030073619, US20040142474, US20100004315, US2012009145 and US20130195920; and Intl Pub. Nos. WO2006063249 and WO2013086322, each of which is herein incorporated by reference in its entirety.
- [1018] In some embodiments, the polynucleotides described herein can be formulated in or with at least one cyclodextrin polymer as described in U.S. Pub. No. US20130184453. In some embodiments, the polynucleotides described herein can be formulated in or with at least one crosslinked cation-binding polymers as described in Intl. Pub. Nos. WO2013106072, WO2013106073 and WO2013106086. In some embodiments, the polynucleotides described herein can be formulated in or with at least PEGylated albumin polymer as described in U.S. Pub. No. US20130231287. Each of the references is herein incorporated by reference in its entirety.
- [1019] In some embodiments, the polynucleotides disclosed herein can be formulated as a nanoparticle using a combination of polymers, lipids, and/or other biodegradable agents, such as, but not limited to, calcium phosphate. Components can be combined in a core-

shell, hybrid, and/or layer-by-layer architecture, to allow for fine-tuning of the nanoparticle for delivery (Wang et al., *Nat Mater.* 2006 5:791-796; Fuller et al., *Biomaterials.* 2008 29:1526-1532; DeKoker et al., *Adv Drug Deliv Rev.* 2011 63:748-761; Endres et al., *Biomaterials.* 2011 32:7721-7731; Su et al., *Mol Pharm.* 2011 Jun 6;8(3):774-87; herein incorporated by reference in their entireties). As a non-limiting example, the nanoparticle can comprise a plurality of polymers such as, but not limited to hydrophilic-hydrophobic polymers (e.g., PEG-PLGA), hydrophobic polymers (e.g., PEG) and/or hydrophilic polymers (Intl. Pub. No. WO20120225129, herein incorporated by reference in its entirety).

[1020] The use of core-shell nanoparticles has additionally focused on a high-throughput approach to synthesize cationic cross-linked nanogel cores and various shells (Siegwart et al., *Proc Natl Acad Sci U S A.* 2011 108:12996-13001; herein incorporated by reference in its entirety). The complexation, delivery, and internalization of the polymeric nanoparticles can be precisely controlled by altering the chemical composition in both the core and shell components of the nanoparticle. For example, the core-shell nanoparticles can efficiently deliver siRNA to mouse hepatocytes after they covalently attach cholesterol to the nanoparticle.

[1021] In some embodiments, a hollow lipid core comprising a middle PLGA layer and an outer neutral lipid layer containing PEG can be used to delivery of the polynucleotides as described herein. In some embodiments, the lipid nanoparticles can comprise a core of the polynucleotides disclosed herein and a polymer shell, which is used to protect the polynucleotides in the core. The polymer shell can be any of the polymers described herein and are known in the art. The polymer shell can be used to protect the polynucleotides in the core.

[1022] Core-shell nanoparticles for use with the polynucleotides described herein are described in U.S. Pat. No. 8,313,777 or Intl. Pub. No. WO2013124867, each of which is herein incorporated by reference in their entirety.

v. Peptides and Proteins

[1023] In some embodiments, the compositions or formulations of the present disclosure comprise the polynucleotides described herein (e.g., a polynucleotide comprising a nucleotide sequence encoding an ACADV L polypeptide) that is formulated with peptides and/or proteins to increase transfection of cells by the polynucleotide, and/or to alter the

biodistribution of the polynucleotide (e.g., by targeting specific tissues or cell types), and/or increase the translation of encoded protein (e.g., Intl. Pub. Nos. WO2012110636 and WO2013123298. In some embodiments, the peptides can be those described in U.S. Pub. Nos. US20130129726, US20130137644 and US20130164219. Each of the references is herein incorporated by reference in its entirety.

w. Conjugates

[1024] In some embodiments, the compositions or formulations of the present disclosure comprise the polynucleotides described herein (e.g., a polynucleotide comprising a nucleotide sequence encoding an ACADVL polypeptide) that is covalently linked to a carrier or targeting group, or including two encoding regions that together produce a fusion protein (e.g., bearing a targeting group and therapeutic protein or peptide) as a conjugate. The conjugate can be a peptide that selectively directs the nanoparticle to neurons in a tissue or organism, or assists in crossing the blood-brain barrier.

[1025] The conjugates include a naturally occurring substance, such as a protein (e.g., human serum albumin (HSA), low-density lipoprotein (LDL), high-density lipoprotein (HDL), or globulin); an carbohydrate (e.g., a dextran, pullulan, chitin, chitosan, inulin, cyclodextrin or hyaluronic acid); or a lipid. The ligand can also be a recombinant or synthetic molecule, such as a synthetic polymer, e.g., a synthetic polyamino acid, an oligonucleotide (e.g., an aptamer). Examples of polyamino acids include polyamino acid is a polylysine (PLL), poly L-aspartic acid, poly L-glutamic acid, styrene-maleic acid anhydride copolymer, poly(L-lactide-co-glycolide) copolymer, divinyl ether-maleic anhydride copolymer, N-(2-hydroxypropyl)methacrylamide copolymer (HMPA), polyethylene glycol (PEG), polyvinyl alcohol (PVA), polyurethane, poly(2-ethylacrylic acid), N-isopropylacrylamide polymers, or polyphosphazene. Example of polyamines include: polyethylenimine, polylysine (PLL), spermine, spermidine, polyamine, pseudopeptide-polyamine, peptidomimetic polyamine, dendrimer polyamine, arginine, amidine, protamine, cationic lipid, cationic porphyrin, quaternary salt of a polyamine, or an alpha helical peptide.

[1026] In some embodiments, the conjugate can function as a carrier for the polynucleotide disclosed herein. The conjugate can comprise a cationic polymer such as, but not limited to, polyamine, polylysine, polyalkylenimine, and polyethylenimine that can be grafted to with poly(ethylene glycol). Exemplary conjugates and their preparations

are described in U.S. Pat. No. 6,586,524 and U.S. Pub. No. US20130211249, each of which herein is incorporated by reference in its entirety.

- [1027] The conjugates can also include targeting groups, e.g., a cell or tissue targeting agent, e.g., a lectin, glycoprotein, lipid or protein, e.g., an antibody, that binds to a specified cell type such as a kidney cell. A targeting group can be a thyrotropin, melanotropin, lectin, glycoprotein, surfactant protein A, Mucin carbohydrate, multivalent lactose, multivalent galactose, N-acetyl-galactosamine, N-acetyl-glucosamine multivalent mannose, multivalent fucose, glycosylated polyaminoacids, multivalent galactose, transferrin, bisphosphonate, polyglutamate, polyaspartate, a lipid, cholesterol, a steroid, bile acid, folate, vitamin B12, biotin, an RGD peptide, an RGD peptide mimetic or an aptamer.
- [1028] Targeting groups can be proteins, e.g., glycoproteins, or peptides, e.g., molecules having a specific affinity for a co-ligand, or antibodies e.g., an antibody, that binds to a specified cell type such as an endothelial cell or bone cell. Targeting groups can also include hormones and hormone receptors. They can also include non-peptidic species, such as lipids, lectins, carbohydrates, vitamins, cofactors, multivalent lactose, multivalent galactose, N-acetyl-galactosamine, N-acetyl-glucosamine multivalent mannose, multivalent fructose, or aptamers. The ligand can be, for example, a lipopolysaccharide, or an activator of p38 MAP kinase.
- [1029] The targeting group can be any ligand that is capable of targeting a specific receptor. Examples include, without limitation, folate, GalNAc, galactose, mannose, mannose-6P, aptamers, integrin receptor ligands, chemokine receptor ligands, transferrin, biotin, serotonin receptor ligands, PSMA, endothelin, GCPII, somatostatin, LDL, and HDL ligands. In particular embodiments, the targeting group is an aptamer. The aptamer can be unmodified or have any combination of modifications disclosed herein. As a non-limiting example, the targeting group can be a glutathione receptor (GR)-binding conjugate for targeted delivery across the blood-central nervous system barrier as described in, e.g., U.S. Pub. No. US2013021661012 (herein incorporated by reference in its entirety).
- [1030] In some embodiments, the conjugate can be a synergistic biomolecule-polymer conjugate, which comprises a long-acting continuous-release system to provide a greater therapeutic efficacy. The synergistic biomolecule-polymer conjugate can be those described in U.S. Pub. No. US20130195799. In some embodiments, the conjugate can be

an aptamer conjugate as described in Intl. Pat. Pub. No. WO2012040524. In some embodiments, the conjugate can be an amine containing polymer conjugate as described in U.S. Pat. No. 8,507,653. Each of the references is herein incorporated by reference in its entirety. In some embodiments, the polynucleotides can be conjugated to SMARTT POLYMER TECHNOLOGY® (PHASERX®, Inc. Seattle, WA).

[1031] In some embodiments, the polynucleotides described herein are covalently conjugated to a cell penetrating polypeptide, which can also include a signal sequence or a targeting sequence. The conjugates can be designed to have increased stability, and/or increased cell transfection; and/or altered the biodistribution (e.g., targeted to specific tissues or cell types).

[1032] In some embodiments, the polynucleotides described herein can be conjugated to an agent to enhance delivery. In some embodiments, the agent can be a monomer or polymer such as a targeting monomer or a polymer having targeting blocks as described in Intl. Pub. No. WO2011062965. In some embodiments, the agent can be a transport agent covalently coupled to a polynucleotide as described in, e.g., U.S. Pat. Nos. 6,835,393 and 7,374,778. In some embodiments, the agent can be a membrane barrier transport enhancing agent such as those described in U.S. Pat. Nos. 7,737,108 and 8,003,129. Each of the references is herein incorporated by reference in its entirety.

x. Micro-Organs

[1033] In some embodiments, the compositions or formulations of the present disclosure comprise the polynucleotides described herein (e.g., a polynucleotide comprising a nucleotide sequence encoding an ACADVL polypeptide) in a micro-organ that can then express an encoded polypeptide of interest in a long-lasting therapeutic formulation. Exemplary micro-organs and formulations are described in Intl. Pub. No. WO2014152211 (herein incorporated by reference in its entirety).

y. Pseudovirions

[1034] In some embodiments, the compositions or formulations of the present disclosure comprise the polynucleotides described herein (e.g., a polynucleotide comprising a nucleotide sequence encoding an ACADVL polypeptide) in pseudovirions (e.g., pseudovirions developed by Aura Biosciences, Cambridge, MA).

[1035] In some embodiments, the pseudovirion used for delivering the polynucleotides can be derived from viruses such as, but not limited to, herpes and papillomaviruses as

described in, e.g., U.S. Pub. Nos. US20130012450, US20130012566, US21030012426 and US20120207840; and Intl. Pub. No. WO2013009717, each of which is herein incorporated by reference in its entirety.

[1036] The pseudovirion can be a virus-like particle (VLP) prepared by the methods described in U.S. Pub. Nos. US20120015899 and US20130177587, and Intl. Pub. Nos. WO2010047839, WO2013116656, WO2013106525 and WO2013122262. In one aspect, the VLP can be bacteriophages MS, Q β , R17, fr, GA, Sp, MI, I, MXI, NL95, AP205, f2, PP7, and the plant viruses Turnip crinkle virus (TCV), Tomato bushy stunt virus (TBSV), Southern bean mosaic virus (SBMV) and members of the genus Bromovirus including Broad bean mottle virus, Brome mosaic virus, Cassia yellow blotch virus, Cowpea chlorotic mottle virus (CCMV), Melandrium yellow fleck virus, and Spring beauty latent virus. In another aspect, the VLP can be derived from the influenza virus as described in U.S. Pub. No. US20130177587 and U.S. Pat. No. 8,506,967. In one aspect, the VLP can comprise a B7-1 and/or B7-2 molecule anchored to a lipid membrane or the exterior of the particle such as described in Intl. Pub. No. WO2013116656. In one aspect, the VLP can be derived from norovirus, rotavirus recombinant VP6 protein or double layered VP2/VP6 such as the VLP as described in Intl. Pub. No. WO2012049366. Each of the references is herein incorporated by reference in its entirety.

[1037] In some embodiments, the pseudovirion can be a human papilloma virus-like particle as described in Intl. Pub. No. WO2010120266 and U.S. Pub. No. US20120171290. In some embodiments, the virus-like particle (VLP) can be a self-assembled particle. In one aspect, the pseudovirions can be virion derived nanoparticles as described in U.S. Pub. Nos. US20130116408 and US20130115247; and Intl. Pub. No. WO2013119877. Each of the references is herein incorporated by reference in their entirety.

[1038] Non-limiting examples of formulations and methods for formulating the polynucleotides described herein are also provided in Intl. Pub. No WO2013090648 (incorporated herein by reference in their entirety).

24. Accelerated Blood Clearance

[1039] The invention provides compounds, compositions and methods of use thereof for reducing the effect of ABC on a repeatedly administered active agent such as a biologically active agent. As will be readily apparent, reducing or eliminating altogether

the effect of ABC on an administered active agent effectively increases its half-life and thus its efficacy.

[1040] In some embodiments the term reducing ABC refers to any reduction in ABC in comparison to a positive reference control ABC inducing LNP such as an MC3 LNP. ABC inducing LNPs cause a reduction in circulating levels of an active agent upon a second or subsequent administration within a given time frame. Thus a reduction in ABC refers to less clearance of circulating agent upon a second or subsequent dose of agent, relative to a standard LNP. The reduction may be, for instance, at least 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98%, or 100%. In some embodiments the reduction is 10-100%, 10-50%, 20-100%, 20-50%, 30-100%, 30-50%, 40%-100%, 40-80%, 50-90%, or 50-100%. Alternatively the reduction in ABC may be characterized as at least a detectable level of circulating agent following a second or subsequent administration or at least a 2 fold, 3 fold, 4 fold, 5 fold increase in circulating agent relative to circulating agent following administration of a standard LNP. In some embodiments the reduction is a 2-100 fold, 2-50 fold, 3-100 fold, 3-50 fold, 3-20 fold, 4-100 fold, 4-50 fold, 4-40 fold, 4-30 fold, 4-25 fold, 4-20 fold, 4-15 fold, 4-10 fold, 4-5 fold, 5 -100 fold, 5-50 fold, 5-40 fold, 5-30 fold, 5-25 fold, 5-20 fold, 5-15 fold, 5-10 fold, 6-100 fold, 6-50 fold, 6-40 fold, 6-30 fold, 6-25 fold, 6-20 fold, 6-15 fold, 6-10 fold, 8-100 fold, 8-50 fold, 8-40 fold, 8-30 fold, 8-25 fold, 8-20 fold, 8-15 fold, 8-10 fold, 10-100 fold, 10-50 fold, 10-40 fold, 10-30 fold, 10-25 fold, 10-20 fold, 10-15 fold, 20-100 fold, 20-50 fold, 20-40 fold, 20-30 fold, or 20-25 fold.

[1041] The disclosure provides lipid-comprising compounds and compositions that are less susceptible to clearance and thus have a longer half-life in vivo. This is particularly the case where the compositions are intended for repeated including chronic administration, and even more particularly where such repeated administration occurs within days or weeks.

[1042] Significantly, these compositions are less susceptible or altogether circumvent the observed phenomenon of accelerated blood clearance (ABC). ABC is a phenomenon in which certain exogenously administered agents are rapidly cleared from the blood upon second and subsequent administrations. This phenomenon has been observed, in part, for a variety of lipid-containing compositions including but not limited to lipidated agents, liposomes or other lipid-based delivery vehicles, and lipid-encapsulated agents. Heretofore, the basis of ABC has been poorly understood and in some cases attributed to

a humoral immune response and accordingly strategies for limiting its impact in vivo particularly in a clinical setting have remained elusive.

- [1043] This disclosure provides compounds and compositions that are less susceptible, if at all susceptible, to ABC. In some important aspects, such compounds and compositions are lipid-comprising compounds or compositions. The lipid-containing compounds or compositions of this disclosure, surprisingly, do not experience ABC upon second and subsequent administration in vivo. This resistance to ABC renders these compounds and compositions particularly suitable for repeated use in vivo, including for repeated use within short periods of time, including days or 1-2 weeks. This enhanced stability and/or half-life is due, in part, to the inability of these compositions to activate B1a and/or B1b cells and/or conventional B cells, pDCs and/or platelets.
- [1044] This disclosure therefore provides an elucidation of the mechanism underlying accelerated blood clearance (ABC). It has been found, in accordance with this disclosure and the inventions provided herein, that the ABC phenomenon at least as it relates to lipids and lipid nanoparticles is mediated, at least in part an innate immune response involving B1a and/or B1b cells, pDC and/or platelets. B1a cells are normally responsible for secreting natural antibody, in the form of circulating IgM. This IgM is poly-reactive, meaning that it is able to bind to a variety of antigens, albeit with a relatively low affinity for each.
- [1045] It has been found in accordance with the invention that some lipidated agents or lipid-comprising formulations such as lipid nanoparticles administered in vivo trigger and are subject to ABC. It has now been found in accordance with the invention that upon administration of a first dose of the LNP, one or more cells involved in generating an innate immune response (referred to herein as sensors) bind such agent, are activated, and then initiate a cascade of immune factors (referred to herein as effectors) that promote ABC and toxicity. For instance, B1a and B1b cells may bind to LNP, become activated (alone or in the presence of other sensors such as pDC and/or effectors such as IL6) and secrete natural IgM that binds to the LNP. Pre-existing natural IgM in the subject may also recognize and bind to the LNP, thereby triggering complement fixation. After administration of the first dose, the production of natural IgM begins within 1-2 hours of administration of the LNP. Typically by about 2-3 weeks the natural IgM is cleared from the system due to the natural half-life of IgM. Natural IgG is produced beginning around 96 hours after administration of the LNP. The agent, when administered in a naïve

setting, can exert its biological effects relatively unencumbered by the natural IgM produced post-activation of the B1a cells or B1b cells or natural IgG. The natural IgM and natural IgG are non-specific and thus are distinct from anti-PEG IgM and anti-PEG IgG.

[1046] Although Applicant is not bound by mechanism, it is proposed that LNPs trigger ABC and/or toxicity through the following mechanisms. It is believed that when an LNP is administered to a subject the LNP is rapidly transported through the blood to the spleen. The LNPs may encounter immune cells in the blood and/or the spleen. A rapid innate immune response is triggered in response to the presence of the LNP within the blood and/or spleen. Applicant has shown herein that within hours of administration of an LNP several immune sensors have reacted to the presence of the LNP. These sensors include but are not limited to immune cells involved in generating an immune response, such as B cells, pDC, and platelets. The sensors may be present in the spleen, such as in the marginal zone of the spleen and/or in the blood. The LNP may physically interact with one or more sensors, which may interact with other sensors. In such a case the LNP is directly or indirectly interacting with the sensors. The sensors may interact directly with one another in response to recognition of the LNP. For instance many sensors are located in the spleen and can easily interact with one another. Alternatively one or more of the sensors may interact with LNP in the blood and become activated. The activated sensor may then interact directly with other sensors or indirectly (e.g., through the stimulation or production of a messenger such as a cytokine e.g., IL6).

[1047] In some embodiments the LNP may interact directly with and activate each of the following sensors: pDC, B1a cells, B1b cells, and platelets. These cells may then interact directly or indirectly with one another to initiate the production of effectors which ultimately lead to the ABC and/or toxicity associated with repeated doses of LNP. For instance, Applicant has shown that LNP administration leads to pDC activation, platelet aggregation and activation and B cell activation. In response to LNP platelets also aggregate and are activated and aggregate with B cells. pDC cells are activated. LNP has been found to interact with the surface of platelets and B cells relatively quickly. Blocking the activation of any one or combination of these sensors in response to LNP is useful for dampening the immune response that would ordinarily occur. This dampening of the immune response results in the avoidance of ABC and/or toxicity.

- [1048] The sensors once activated produce effectors. An effector, as used herein, is an immune molecule produced by an immune cell, such as a B cell. Effectors include but are not limited to immunoglobulin such as natural IgM and natural IgG and cytokines such as IL6. B1a and B1b cells stimulate the production of natural IgMs within 2-6 hours following administration of an LNP. Natural IgG can be detected within 96 hours. IL6 levels are increased within several hours. The natural IgM and IgG circulate in the body for several days to several weeks. During this time the circulating effectors can interact with newly administered LNPs, triggering those LNPs for clearance by the body. For instance, an effector may recognize and bind to an LNP. The Fc region of the effector may be recognized by and trigger uptake of the decorated LNP by macrophage. The macrophage are then transported to the spleen. The production of effectors by immune sensors is a transient response that correlates with the timing observed for ABC.
- [1049] If the administered dose is the second or subsequent administered dose, and if such second or subsequent dose is administered before the previously induced natural IgM and/or IgG is cleared from the system (e.g., before the 2-3 window time period), then such second or subsequent dose is targeted by the circulating natural IgM and/or natural IgG or Fc which trigger alternative complement pathway activation and is itself rapidly cleared. When LNP are administered after the effectors have cleared from the body or are reduced in number, ABC is not observed.
- [1050] Thus, it is useful according to aspects of the invention to inhibit the interaction between LNP and one or more sensors, to inhibit the activation of one or more sensors by LNP (direct or indirect), to inhibit the production of one or more effectors, and/or to inhibit the activity of one or more effectors. In some embodiments the LNP is designed to limit or block interaction of the LNP with a sensor. For instance the LNP may have an altered PC and/or PEG to prevent interactions with sensors. Alternatively or additionally an agent that inhibits immune responses induced by LNPs may be used to achieve any one or more of these effects.
- [1051] It has also been determined that conventional B cells are also implicated in ABC. Specifically, upon first administration of an agent, conventional B cells, referred to herein as CD19(+), bind to and react against the agent. Unlike B1a and B1b cells though, conventional B cells are able to mount first an IgM response (beginning around 96 hours after administration of the LNPs) followed by an IgG response (beginning around 14 days after administration of the LNPs) concomitant with a memory response. Thus

conventional B cells react against the administered agent and contribute to IgM (and eventually IgG) that mediates ABC. The IgM and IgG are typically anti-PEG IgM and anti-PEG IgG.

[1052] It is contemplated that in some instances, the majority of the ABC response is mediated through B1a cells and B1a-mediated immune responses. It is further contemplated that in some instances, the ABC response is mediated by both IgM and IgG, with both conventional B cells and B1a cells mediating such effects. In yet still other instances, the ABC response is mediated by natural IgM molecules, some of which are capable of binding to natural IgM, which may be produced by activated B1a cells. The natural IgMs may bind to one or more components of the LNPs, e.g., binding to a phospholipid component of the LNPs (such as binding to the PC moiety of the phospholipid) and/or binding to a PEG-lipid component of the LNPs (such as binding to PEG-DMG, in particular, binding to the PEG moiety of PEG-DMG). Since B1a expresses CD36, to which phosphatidylcholine is a ligand, it is contemplated that the CD36 receptor may mediate the activation of B1a cells and thus production of natural IgM. In yet still other instances, the ABC response is mediated primarily by conventional B cells.

[1053] It has been found in accordance with the invention that the ABC phenomenon can be reduced or abrogated, at least in part, through the use of compounds and compositions (such as agents, delivery vehicles, and formulations) that do not activate B1a cells. Compounds and compositions that do not activate B1a cells may be referred to herein as B1a inert compounds and compositions. It has been further found in accordance with the invention that the ABC phenomenon can be reduced or abrogated, at least in part, through the use of compounds and compositions that do not activate conventional B cells. Compounds and compositions that do not activate conventional B cells may in some embodiments be referred to herein as CD19-inert compounds and compositions. Thus, in some embodiments provided herein, the compounds and compositions do not activate B1a cells and they do not activate conventional B cells. Compounds and compositions that do not activate B1a cells and conventional B cells may in some embodiments be referred to herein as B1a/CD19-inert compounds and compositions.

[1054] These underlying mechanisms were not heretofore understood, and the role of B1a and B1b cells and their interplay with conventional B cells in this phenomenon was also not appreciated.

- [1055] Accordingly, this disclosure provides compounds and compositions that do not promote ABC. These may be further characterized as not capable of activating B1a and/or B1b cells, platelets and/or pDC, and optionally conventional B cells also. These compounds (e.g., agents, including biologically active agents such as prophylactic agents, therapeutic agents and diagnostic agents, delivery vehicles, including liposomes, lipid nanoparticles, and other lipid-based encapsulating structures, etc.) and compositions (e.g., formulations, etc.) are particularly desirable for applications requiring repeated administration, and in particular repeated administrations that occur within with short periods of time (e.g., within 1-2 weeks). This is the case, for example, if the agent is a nucleic acid based therapeutic that is provided to a subject at regular, closely-spaced intervals. The findings provided herein may be applied to these and other agents that are similarly administered and/or that are subject to ABC.
- [1056] Of particular interest are lipid-comprising compounds, lipid-comprising particles, and lipid-comprising compositions as these are known to be susceptible to ABC. Such lipid-comprising compounds particles, and compositions have been used extensively as biologically active agents or as delivery vehicles for such agents. Thus, the ability to improve their efficacy of such agents, whether by reducing the effect of ABC on the agent itself or on its delivery vehicle, is beneficial for a wide variety of active agents.
- [1057] Also provided herein are compositions that do not stimulate or boost an acute phase response (ARP) associated with repeat dose administration of one or more biologically active agents.
- [1058] The composition, in some instances, may not bind to IgM, including but not limited to natural IgM.
- [1059] The composition, in some instances, may not bind to an acute phase protein such as but not limited to C-reactive protein.
- [1060] The composition, in some instances, may not trigger a CD5(+) mediated immune response. As used herein, a CD5(+) mediated immune response is an immune response that is mediated by B1a and/or B1b cells. Such a response may include an ABC response, an acute phase response, induction of natural IgM and/or IgG, and the like.
- [1061] The composition, in some instances, may not trigger a CD19(+) mediated immune response. As used herein, a CD19(+) mediated immune response is an immune response that is mediated by conventional CD19(+), CD5(-) B cells. Such a response may include induction of IgM, induction of IgG, induction of memory B cells, an ABC response, an

anti-drug antibody (ADA) response including an anti-protein response where the protein may be encapsulated within an LNP, and the like.

[1062] B1a cells are a subset of B cells involved in innate immunity. These cells are the source of circulating IgM, referred to as natural antibody or natural serum antibody. Natural IgM antibodies are characterized as having weak affinity for a number of antigens, and therefore they are referred to as “poly-specific” or “poly-reactive”, indicating their ability to bind to more than one antigen. B1a cells are not able to produce IgG. Additionally, they do not develop into memory cells and thus do not contribute to an adaptive immune response. However, they are able to secrete IgM upon activation. The secreted IgM is typically cleared within about 2-3 weeks, at which point the immune system is rendered relatively naïve to the previously administered antigen. If the same antigen is presented after this time period (e.g., at about 3 weeks after the initial exposure), the antigen is not rapidly cleared. However, significantly, if the antigen is presented within that time period (e.g., within 2 weeks, including within 1 week, or within days), then the antigen is rapidly cleared. This delay between consecutive doses has rendered certain lipid-containing therapeutic or diagnostic agents unsuitable for use.

[1063] In humans, B1a cells are CD19(+), CD20(+), CD27(+), CD43(+), CD70(-) and CD5(+). In mice, B1a cells are CD19(+), CD5(+), and CD45 B cell isoform B220(+). It is the expression of CD5 which typically distinguishes B1a cells from other convention B cells. B1a cells may express high levels of CD5, and on this basis may be distinguished from other B-1 cells such as B-1b cells which express low or undetectable levels of CD5. CD5 is a pan-T cell surface glycoprotein. B1a cells also express CD36, also known as fatty acid translocase. CD36 is a member of the class B scavenger receptor family. CD36 can bind many ligands, including oxidized low density lipoproteins, native lipoproteins, oxidized phospholipids, and long-chain fatty acids.

[1064] B1b cells are another subset of B cells involved in innate immunity. These cells are another source of circulating natural IgM. Several antigens, including PS, are capable of inducing T cell independent immunity through B1b activation. CD27 is typically upregulated on B1b cells in response to antigen activation. Similar to B1a cells, the B1b cells are typically located in specific body locations such as the spleen and peritoneal cavity and are in very low abundance in the blood. The B1b secreted natural IgM is typically cleared within about 2-3 weeks, at which point the immune system is rendered relatively naïve to the previously administered antigen. If the same antigen is presented

after this time period (e.g., at about 3 weeks after the initial exposure), the antigen is not rapidly cleared. However, significantly, if the antigen is presented within that time period (e.g., within 2 weeks, including within 1 week, or within days), then the antigen is rapidly cleared. This delay between consecutive doses has rendered certain lipid-containing therapeutic or diagnostic agents unsuitable for use.

[1065] In some embodiments it is desirable to block B1a and/or B1b cell activation. One strategy for blocking B1a and/or B1b cell activation involves determining which components of a lipid nanoparticle promote B cell activation and neutralizing those components. It has been discovered herein that at least PEG and phosphatidylcholine (PC) contribute to B1a and B1b cell interaction with other cells and/or activation. PEG may play a role in promoting aggregation between B1 cells and platelets, which may lead to activation. PC (a helper lipid in LNPs) is also involved in activating the B1 cells, likely through interaction with the CD36 receptor on the B cell surface. Numerous particles have PEG-lipid alternatives, PEG-less, and/or PC replacement lipids (e.g. oleic acid or analogs thereof) have been designed and tested. Applicant has established that replacement of one or more of these components within an LNP that otherwise would promote ABC upon repeat administration, is useful in preventing ABC by reducing the production of natural IgM and/or B cell activation. Thus, the invention encompasses LNPs that have reduced ABC as a result of a design which eliminates the inclusion of B cell triggers.

[1066] Another strategy for blocking B1a and/or B1b cell activation involves using an agent that inhibits immune responses induced by LNPs. These types of agents are discussed in more detail below. In some embodiments these agents block the interaction between B1a/B1b cells and the LNP or platelets or pDC. For instance the agent may be an antibody or other binding agent that physically blocks the interaction. An example of this is an antibody that binds to CD36 or CD6. The agent may also be a compound that prevents or disables the B1a/B1b cell from signaling once activated or prior to activation. For instance, it is possible to block one or more components in the B1a/B1b signaling cascade the results from B cell interaction with LNP or other immune cells. In other embodiments the agent may act one or more effectors produced by the B1a/B1b cells following activation. These effectors include for instance, natural IgM and cytokines.

[1067] It has been demonstrated according to aspects of the invention that when activation of pDC cells is blocked, B cell activation in response to LNP is decreased.

Thus, in order to avoid ABC and/or toxicity, it may be desirable to prevent pDC activation. Similar to the strategies discussed above, pDC cell activation may be blocked by agents that interfere with the interaction between pDC and LNP and/or B cells/platelets. Alternatively agents that act on the pDC to block its ability to get activated or on its effectors can be used together with the LNP to avoid ABC.

[1068] Platelets may also play an important role in ABC and toxicity. Very quickly after a first dose of LNP is administered to a subject platelets associate with the LNP, aggregate and are activated. In some embodiments it is desirable to block platelet aggregation and/or activation. One strategy for blocking platelet aggregation and/or activation involves determining which components of a lipid nanoparticle promote platelet aggregation and/or activation and neutralizing those components. It has been discovered herein that at least PEG contribute to platelet aggregation, activation and/or interaction with other cells. Numerous particles have PEG-lipid alternatives and PEG-less have been designed and tested. Applicant has established that replacement of one or more of these components within an LNP that otherwise would promote ABC upon repeat administration, is useful in preventing ABC by reducing the production of natural IgM and/or platelet aggregation. Thus, the invention encompasses LNPs that have reduced ABC as a result of a design which eliminates the inclusion of platelet triggers. Alternatively agents that act on the platelets to block its activity once it is activated or on its effectors can be used together with the LNP to avoid ABC.

(i) Measuring ABC Activity and related activities

[1069] Various compounds and compositions provided herein, including LNPs, do not promote ABC activity upon administration in vivo. These LNPs may be characterized and/or identified through any of a number of assays, such as but not limited to those described below, as well as any of the assays disclosed in the Examples section, include the methods subsection of the Examples.

[1070] In some embodiments the methods involve administering an LNP without producing an immune response that promotes ABC. An immune response that promotes ABC involves activation of one or more sensors, such as B1 cells, pDC, or platelets, and one or more effectors, such as natural IgM, natural IgG or cytokines such as IL6. Thus administration of an LNP without producing an immune response that promotes ABC, at a minimum involves administration of an LNP without significant activation of one or

more sensors and significant production of one or more effectors. Significant used in this context refers to an amount that would lead to the physiological consequence of accelerated blood clearance of all or part of a second dose with respect to the level of blood clearance expected for a second dose of an ABC triggering LNP. For instance, the immune response should be dampened such that the ABC observed after the second dose is lower than would have been expected for an ABC triggering LNP.

(ii) B1a or B1b activation assay

[1071] Certain compositions provided in this disclosure do not activate B cells, such as B1a or B1b cells (CD19⁺ CD5⁺) and/or conventional B cells (CD19⁺ CD5⁻). Activation of B1a cells, B1b cells, or conventional B cells may be determined in a number of ways, some of which are provided below. B cell population may be provided as fractionated B cell populations or unfractionated populations of splenocytes or peripheral blood mononuclear cells (PBMC). If the latter, the cell population may be incubated with the LNP of choice for a period of time, and then harvested for further analysis. Alternatively, the supernatant may be harvested and analyzed.

(iii) Upregulation of activation marker cell surface expression

[1072] Activation of B1a cells, B1b cells, or conventional B cells may be demonstrated as increased expression of B cell activation markers including late activation markers such as CD86. In an exemplary non-limiting assay, unfractionated B cells are provided as a splenocyte population or as a PBMC population, incubated with an LNP of choice for a particular period of time, and then stained for a standard B cell marker such as CD19 and for an activation marker such as CD86, and analyzed using for example flow cytometry. A suitable negative control involves incubating the same population with medium, and then performing the same staining and visualization steps. An increase in CD86 expression in the test population compared to the negative control indicates B cell activation.

(iv) Pro-inflammatory cytokine release

[1073] B cell activation may also be assessed by cytokine release assay. For example, activation may be assessed through the production and/or secretion of cytokines such as IL-6 and/or TNF-alpha upon exposure with LNPs of interest.

[1074] Such assays may be performed using routine cytokine secretion assays well known in the art. An increase in cytokine secretion is indicative of B cell activation.

(v) LNP binding/association to and/or uptake by B cells

[1075] LNP association or binding to B cells may also be used to assess an LNP of interest and to further characterize such LNP. Association/binding and/or uptake/internalization may be assessed using a detectably labeled, such as fluorescently labeled, LNP and tracking the location of such LNP in or on B cells following various periods of incubation.

[1076] The invention further contemplates that the compositions provided herein may be capable of evading recognition or detection and optionally binding by downstream mediators of ABC such as circulating IgM and/or acute phase response mediators such as acute phase proteins (e.g., C-reactive protein (CRP)).

(vi) Methods of use for reducing ABC

[1077] Also provided herein are methods for delivering LNPs, which may encapsulate an agent such as a therapeutic agent, to a subject without promoting ABC.

[1078] In some embodiments, the method comprises administering any of the LNPs described herein, which do not promote ABC, for example, do not induce production of natural IgM binding to the LNPs, do not activate B1a and/or B1b cells. As used herein, an LNP that “does not promote ABC” refers to an LNP that induces no immune responses that would lead to substantial ABC or a substantially low level of immune responses that is not sufficient to lead to substantial ABC. An LNP that does not induce the production of natural IgMs binding to the LNP refers to LNPs that induce either no natural IgM binding to the LNPs or a substantially low level of the natural IgM molecules, which is insufficient to lead to substantial ABC. An LNP that does not activate B1a and/or B1b cells refer to LNPs that induce no response of B1a and/or B1b cells to produce natural IgM binding to the LNPs or a substantially low level of B1a and/or B1b responses, which is insufficient to lead to substantial ABC.

[1079] In some embodiments the terms do not activate and do not induce production are a relative reduction to a reference value or condition. In some embodiments the reference value or condition is the amount of activation or induction of production of a molecule such as IgM by a standard LNP such as an MC3 LNP. In some embodiments the relative

reduction is a reduction of at least 30%, for example at least 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100%. In other embodiments the terms do not activate cells such as B cells and do not induce production of a protein such as IgM may refer to an undetectable amount of the active cells or the specific protein.

(vii) Platelet effects and toxicity

[1080] The invention is further premised in part on the elucidation of the mechanism underlying dose-limiting toxicity associated with LNP administration. Such toxicity may involve coagulopathy, disseminated intravascular coagulation (DIC, also referred to as consumptive coagulopathy), whether acute or chronic, and/or vascular thrombosis. In some instances, the dose-limiting toxicity associated with LNPs is acute phase response (APR) or complement activation-related pseudoallergy (CARPA).

[1081] As used herein, coagulopathy refers to increased coagulation (blood clotting) in vivo. The findings reported in this disclosure are consistent with such increased coagulation and significantly provide insight on the underlying mechanism. Coagulation is a process that involves a number of different factors and cell types, and heretofore the relationship between and interaction of LNPs and platelets has not been understood in this regard. This disclosure provides evidence of such interaction and also provides compounds and compositions that are modified to have reduced platelet effect, including reduced platelet association, reduced platelet aggregation, and/or reduced platelet aggregation. The ability to modulate, including preferably down-modulate, such platelet effects can reduce the incidence and/or severity of coagulopathy post-LNP administration. This in turn will reduce toxicity relating to such LNP, thereby allowing higher doses of LNPs and importantly their cargo to be administered to patients in need thereof.

[1082] CARPA is a class of acute immune toxicity manifested in hypersensitivity reactions (HSRs), which may be triggered by nanomedicines and biologicals. Unlike allergic reactions, CARPA typically does not involve IgE but arises as a consequence of activation of the complement system, which is part of the innate immune system that enhances the body's abilities to clear pathogens. One or more of the following pathways, the classical complement pathway (CP), the alternative pathway (AP), and the lectin

pathway (LP), may be involved in CARPA. Szebeni, *Molecular Immunology*, 61:163-173 (2014).

[1083] The classical pathway is triggered by activation of the C1-complex, which contains C1q, C1r, C1s, or C1qr2s2. Activation of the C1-complex occurs when C1q binds to IgM or IgG complexed with antigens, or when C1q binds directly to the surface of the pathogen. Such binding leads to conformational changes in the C1q molecule, which leads to the activation of C1r, which in turn, cleave C1s. The C1r2s2 component now splits C4 and then C2, producing C4a, C4b, C2a, and C2b. C4b and C2b bind to form the classical pathway C3-convertase (C4b2b complex), which promotes cleavage of C3 into C3a and C3b. C3b then binds the C3 convertase to form the C5 convertase (C4b2b3b complex). The alternative pathway is continuously activated as a result of spontaneous C3 hydrolysis. Factor P (properdin) is a positive regulator of the alternative pathway. Oligomerization of properdin stabilizes the C3 convertase, which can then cleave much more C3. The C3 molecules can bind to surfaces and recruit more B, D, and P activity, leading to amplification of the complement activation.

[1084] Acute phase response (APR) is a complex systemic innate immune responses for preventing infection and clearing potential pathogens. Numerous proteins are involved in APR and C-reactive protein is a well-characterized one.

[1085] It has been found, in accordance with the invention, that certain LNP are able to associate physically with platelets almost immediately after administration in vivo, while other LNP do not associate with platelets at all or only at background levels. Significantly, those LNPs that associate with platelets also apparently stabilize the platelet aggregates that are formed thereafter. Physical contact of the platelets with certain LNPs correlates with the ability of such platelets to remain aggregated or to form aggregates continuously for an extended period of time after administration. Such aggregates comprise activated platelets and also innate immune cells such as macrophages and B cells.

25. Methods of Use

[1086] The polynucleotides, pharmaceutical compositions and formulations described herein are used in the preparation, manufacture and therapeutic use to treat and/or prevent ACADVL-related diseases, disorders or conditions. In some embodiments, the

polynucleotides, compositions and formulations of the invention are used to treat and/or prevent VLCADD.

[1087] In some embodiments, the polynucleotides, pharmaceutical compositions and formulations of the invention are used in methods for reducing the levels of acylcarnitine and/or an acylcarnitine metabolite in a subject in need thereof. For instance, one aspect of the invention provides a method of alleviating the signs and symptoms of VLCADD in a subject comprising the administration of a composition or formulation comprising a polynucleotide encoding ACADVL to that subject (e.g., an mRNA encoding an ACADVL polypeptide).

[1088] In some embodiments, the polynucleotides, pharmaceutical compositions and formulations of the invention are used to reduce the level of a metabolite associated with VLCADD (e.g., the substrate or product, i.e., acylcarnitine and/or an acylcarnitine metabolite), the method comprising administering to the subject an effective amount of a polynucleotide encoding an ACADVL polypeptide.

[1089] In some embodiments, the administration of an effective amount of a polynucleotide, pharmaceutical composition or formulation of the invention reduces the levels of a biomarker of VLCADD, e.g., acylcarnitine or an acylcarnitine metabolite. In some embodiments, the administration of the polynucleotide, pharmaceutical composition or formulation of the invention results in reduction in the level of one or more biomarkers of VLCADD, e.g., acylcarnitine or an acylcarnitine metabolite, within a short period of time (e.g., within about 6 hours, within about 8 hours, within about 12 hours, within about 16 hours, within about 20 hours, or within about 24 hours) after administration of the polynucleotide, pharmaceutical composition or formulation of the invention.

[1090] Replacement therapy is a potential treatment for VLCADD. Thus, in certain aspects of the invention, the polynucleotides, e.g., mRNA, disclosed herein comprise one or more sequences encoding an ACADVL polypeptide that is suitable for use in gene replacement therapy for VLCADD. In some embodiments, the present disclosure treats a lack of ACADVL or ACADVL activity, or decreased or abnormal ACADVL activity in a subject by providing a polynucleotide, e.g., mRNA, that encodes an ACADVL polypeptide to the subject. In some embodiments, the polynucleotide is sequence-optimized. In some embodiments, the polynucleotide (e.g., an mRNA) comprises a nucleic acid sequence (e.g., an ORF) encoding an ACADVL polypeptide, wherein the nucleic acid is sequence-optimized, e.g., by modifying its G/C, uridine, or thymidine

content, and/or the polynucleotide comprises at least one chemically modified nucleoside. In some embodiments, the polynucleotide comprises a miRNA binding site, e.g., a miRNA binding site that binds miRNA-142 and/or a miRNA binding site that binds miRNA-126.

[1091] In some embodiments, the administration of a composition or formulation comprising polynucleotide, pharmaceutical composition or formulation of the invention to a subject results in a decrease in acylcarnitine and/or an acylcarnitine metabolite in cells to a level at least 10%, at least 15%, at least 20%, at least 25%, at least 30%, at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or to 100% lower than the level observed prior to the administration of the composition or formulation.

[1092] In some embodiments, the administration of the polynucleotide, pharmaceutical composition or formulation of the invention results in expression of ACADVL protein in cells of the subject. In some embodiments, administering the polynucleotide, pharmaceutical composition or formulation of the invention results in an increase of ACADVL enzymatic activity in the subject. For example, in some embodiments, the polynucleotides of the present invention are used in methods of administering a composition or formulation comprising an mRNA encoding an ACADVL polypeptide to a subject, wherein the method results in an increase of ACADVL enzymatic activity in at least some cells of a subject.

[1093] In some embodiments, the administration of a composition or formulation comprising an mRNA encoding an ACADVL polypeptide to a subject results in an increase of ACADVL enzymatic activity in cells subject to a level at least 10%, at least 15%, at least 20%, at least 25%, at least 30%, at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or to 100% or more of the activity level expected in a normal subject, e.g., a human not in need of treatment for VLCADD.

[1094] In some embodiments, the administration of the polynucleotide, pharmaceutical composition or formulation of the invention results in expression of ACADVL protein in at least some of the cells of a subject that persists for a period of time sufficient to allow significant acylcarnitine metabolism to occur.

- [1095] In some embodiments, the expression of the encoded polypeptide is increased. In some embodiments, the polynucleotide increases ACADVL expression levels in cells when introduced into those cells, *e.g.*, by at least 10%, at least 15%, at least 20%, at least 25%, at least 30%, at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or to 100% with respect to the ACADVL expression level in the cells before the polypeptide is introduced in the cells.
- [1096] In some embodiments, the method or use comprises administering a polynucleotide, *e.g.*, mRNA, comprising a nucleotide sequence having sequence similarity to a polynucleotide selected from the group of SEQ ID NOs: 7 to 31 (See Table 2), wherein the polynucleotide encodes an ACADVL polypeptide.
- [1097] Other aspects of the present disclosure relate to transplantation of cells containing polynucleotides to a mammalian subject. Administration of cells to mammalian subjects is known to those of ordinary skill in the art, and includes, but is not limited to, local implantation (*e.g.*, topical or subcutaneous administration), organ delivery or systemic injection (*e.g.*, intravenous injection or inhalation), and the formulation of cells in pharmaceutically acceptable carriers.
- [1098] The present disclosure also provides methods to increase ACADVL activity in a subject in need thereof, *e.g.*, a subject with VLCADD, comprising administering to the subject a therapeutically effective amount of a composition or formulation comprising mRNA encoding an ACADVL polypeptide disclosed herein, *e.g.*, a human ACADVL polypeptide, a mutant thereof, or a fusion protein comprising a human ACADVL.
- [1099] In some aspects, the ACADVL activity measured after administration to a subject in need thereof, *e.g.*, a subject with VLCADD, is at least the normal ACADVL activity level observed in healthy human subjects. In some aspects, the ACADVL activity measured after administration is at higher than the ACADVL activity level observed in VLCADD patients, *e.g.*, untreated VLCADD patients. In some aspects, the increase in ACADVL activity in a subject in need thereof, *e.g.*, a subject with VLCADD, after administering to the subject a therapeutically effective amount of a composition or formulation comprising mRNA encoding an ACADVL polypeptide disclosed herein is at least about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, or greater than 100 percent of the normal ACADVL activity level observed in healthy human subjects. In some aspects, the increase in ACADVL activity above the ACADVL

activity level observed in VLCADD patients after administering to the subject a composition or formulation comprising an mRNA encoding an ACADVL polypeptide disclosed herein (e.g., after a single dose administration) is maintained for at least 1 day, 2 days, 3 days, 4 days, 5 days, 6 days, 7 days, 8 days, 9 days, 10 days, 12 days, 14 days, 21 days, or 28 days.

[1100] Acylcarnitine or acylcarnitine metabolite levels can be measured in the plasma or tissues (e.g., heart, liver, brain or skeletal muscle tissue) using methods known in the art. The present disclosure also provides a method to decrease Acylcarnitine or acylcarnitine metabolite levels in a subject in need thereof, e.g., untreated VLCADD patients, comprising administering to the subject a therapeutically effective amount of a composition or formulation comprising mRNA encoding an ACADVL polypeptide disclosed herein.

[1101] The present disclosure also provides a method to treat, prevent, or ameliorate the symptoms of VLCADD (e.g., hypertrophic or dilated cardiomyopathy, pericardial effusion, arrhythmias, hypotonia, hepatomegaly, or intermittent hypoglycemia) in an VLCADD patient comprising administering to the subject a therapeutically effective amount of a composition or formulation comprising mRNA encoding an ACADVL polypeptide disclosed herein. In some aspects, the administration of a therapeutically effective amount of a composition or formulation comprising mRNA encoding an ACADVL polypeptide disclosed herein to subject in need of treatment for VLCADD results in reducing the symptoms of VLCADD.

[1102] In some aspects, the dose of mRNA encoding an ACADVL polypeptide disclosed herein is at least about 0.1 nmol/kg, at least about 0.2 nmol/kg, at least about 0.3 nmol/kg, at least about 0.4 nmol/kg, at least about 0.5 nmol/kg, at least about 0.6 nmol/kg, at least about 0.7 nmol/kg, at least about 0.8 nmol/kg, at least about 0.9 nmol/kg, at least about 1 nmol/kg, at least about 1.1 nmol/kg, at least about 1.2 nmol/kg, at least about 1.3 nmol/kg, at least about 1.4 nmol/kg, at least about 1.5 nmol/kg, at least about 1.6 nmol/kg, at least about 1.7 nmol/kg, at least about 1.8 nmol/kg, at least about 1.9 nmol/kg, at least about 2 nmol/kg, at least about 2.5 nmol/kg, at least about 3 nmol/kg, at least about 3.5 nmol/kg, at least about 4 nmol/kg, at least about 4.5 nmol/kg, or at least about 5 nmol/kg. In some aspects, the dose of mRNA encoding an ACADVL polypeptide disclosed herein is at least about 0.05 mg/kg, at least about 0.1 mg/kg, at least about 0.15 mg/kg, at least about 0.2 mg/kg, at least about 0.25 mg/kg, at least about 0.3 mg/kg, at

least about 0.35 mg/kg, at least about 0.4 mg/kg, at least about 0.45 mg/kg, at least about 0.5 mg/kg, at least about 0.55 mg/kg, at least about 0.6 mg/kg, at least about 0.7 mg/kg, at least about 0.75 mg/kg, at least about 0.8 mg/kg, at least about 0.85 mg/kg, at least about 0.9 mg/kg, at least about 0.95 mg/kg, or at least about 1 mg/kg.

[1103] In some embodiments, the polynucleotides (e.g., mRNA), pharmaceutical compositions and formulations used in the methods of the invention comprise a uracil-modified sequence encoding an ACADVL polypeptide disclosed herein and a miRNA binding site disclosed herein, e.g., a miRNA binding site that binds to miR-142 and/or a miRNA binding site that binds to miR-126. In some embodiments, the uracil-modified sequence encoding an ACADVL polypeptide comprises at least one chemically modified nucleobase, e.g., 5-methoxyuracil. In some embodiments, at least 95% of a type of nucleobase (e.g., uracil) in a uracil-modified sequence encoding an ACADVL polypeptide of the invention are modified nucleobases. In some embodiments, at least 95% of uracil in a uracil-modified sequence encoding an ACADVL polypeptide is 5-methoxyuridine. In some embodiments, the polynucleotide comprising a nucleotide sequence encoding an ACADVL polypeptide disclosed herein and a miRNA binding site is formulated with a delivery agent comprising, e.g., a compound having the Formula (I), e.g., any of Compounds 1-232, e.g., Compound 18; a compound having the Formula (III), (IV), (V), or (VI), e.g., any of Compounds 233-342, e.g., Compound 236; or a compound having the Formula (VIII), e.g., any of Compounds 419-428, e.g., Compound 428, or any combination thereof. In some embodiments, the delivery agent comprises Compound 18, DSPC, Cholesterol, and Compound 428, e.g., with a mole ratio of about 50:10:38.5:1.5.

[1104] The skilled artisan will appreciate that the therapeutic effectiveness of a drug or a treatment of the instant invention can be characterized or determined by measuring the level of expression of an encoded protein (e.g., enzyme) in a sample or in samples taken from a subject (e.g., from a preclinical test subject (rodent, primate, etc.) or from a clinical subject (human)). Likewise, the therapeutic effectiveness of a drug or a treatment of the instant invention can be characterized or determined by measuring the level of activity of an encoded protein (e.g., enzyme) in a sample or in samples taken from a subject (e.g., from a preclinical test subject (rodent, primate, etc.) or from a clinical subject (human)). Furthermore, the therapeutic effectiveness of a drug or a treatment of the instant invention can be characterized or determined by measuring the level of an appropriate biomarker in sample(s) taken from a subject. Levels of protein and/or

biomarkers can be determined post-administration with a single dose of an mRNA therapeutic of the invention or can be determined and/or monitored at several time points following administration with a single dose or can be determined and/or monitored throughout a course of treatment, e.g., a multi-dose treatment.

(i) ACADVL Protein Expression Levels

[1105] Certain aspects of the invention feature measurement, determination and/or monitoring of the expression level or levels of α -galactosidase A (ACADVL) protein in a subject, for example, in an animal (e.g., rodents, primates, and the like) or in a human subject. Animals include normal, healthy or wild type animals, as well as animal models for use in understanding VLCADD and treatments thereof. Exemplary animal models include rodent models, for example, ACADVL deficient mice also referred to as ACADVL $-/-$ knockout mice.

[1106] ACADVL protein expression levels can be measured or determined by any art-recognized method for determining protein levels in biological samples, e.g., blood samples or needle tissue biopsy. The term "level" or "level of a protein" as used herein, preferably means the weight, mass or concentration of the protein within a sample or a subject. It will be understood by the skilled artisan that in certain embodiments the sample may be subjected, e.g., to any of the following: purification, precipitation, separation, e.g. centrifugation and/or HPLC, and subsequently subjected to determining the level of the protein, e.g., using mass and/or spectrometric analysis. In exemplary embodiments, enzyme-linked immunosorbent assay (ELISA) can be used to determine protein expression levels. In other exemplary embodiments, protein purification, separation and LC-MS can be used as a means for determining the level of a protein according to the invention. In some embodiments, an mRNA therapy of the invention (e.g., a single intravenous dose) results in increased ACADVL protein expression levels in the tissue (e.g., heart, liver, brain or skeletal muscle) of the subject (e.g., 2-fold, 3-fold, 4-fold, 5-fold, 6-fold, 7-fold, 8-fold, 9-fold, 10-fold, 20-fold, 30-fold, 40-fold, 50-fold increase and/or increased to at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 75%, 80%, at least 85%, at least 90%, at least 95%, or at least 100% or a reference level or normal level) for at least 6 hours, at least 12 hours, at least 24 hours, at least 36 hours, at least 48 hours, at least 60 hours, at least 72 hours, at least 84 hours, at least 96 hours, at least 108 hours, at least 122 hours, at

least 144 hours, or at least 168 hours after administration of a single dose of the mRNA therapy.

(ii) ACADVL Protein Activity

[1107] In VLCADD patients, ACADVL enzymatic activity is reduced, and VLCADD patients typically have little to no ACADVL protein activity. Further aspects of the invention feature measurement, determination and/or monitoring of the activity level(s) (*i.e.*, enzymatic activity level(s)) of ACADVL protein in a subject, for example, in an animal (e.g., rodent, primate, and the like) or in a human subject. Activity levels can be measured or determined by any art-recognized method for determining enzymatic activity levels in biological samples. The term "activity level" or "enzymatic activity level" as used herein, preferably means the activity of the enzyme per volume, mass or weight of sample or total protein within a sample. In exemplary embodiments, the "activity level" or "enzymatic activity level" is described in terms of units per milliliter of fluid (e.g., bodily fluid, e.g., serum, plasma, urine and the like) or is described in terms of units per weight of tissue or per weight of protein (*e.g.*, total protein) within a sample. Units ("U") of enzyme activity can be described in terms of weight or mass of substrate hydrolyzed per unit time. Exemplary embodiments of the invention feature ACADVL activity described in terms of U/ml plasma or U/mg protein (tissue), where units ("U") are described in terms of nmol substrate hydrolyzed per hour (or nmol/hr).

[1108] In exemplary embodiments, an mRNA therapy of the invention features a pharmaceutical composition comprising a dose of mRNA effective to result in at least 5 U/mg, at least 10 U/mg, at least 20 U/mg, at least 30 U/mg, at least 40 U/mg, at least 50 U/mg, at least 60 U/mg, at least 70 U/mg, at least 80 U/mg, at least 90 U/mg, at least 100 U/mg, or at least 150 U/mg of ACADVL activity in plasma or tissue (e.g., heart, liver, brain or skeletal muscle) between 6 and 12 hours, or between 12 and 24 hours, between 24 and 48 hours, between 48 and 72 hours, more than 3 days, more than 4 days, more than 5 days, more than 6 days, more than 7 days, more than 1 week, more than 2 weeks, more than 3 weeks, more than 4 weeks, more than 5 weeks, or more than 6 weeks post administration.

[1109] In exemplary embodiments, the invention features a pharmaceutical composition comprising a dose of mRNA effective to result in an increase of plasma ACADVL activity level to a level at or above a reference ACADVL activity level or a normal

physiologic level for at least 12 hours, at least 18 hours, at least 24 hours, at least 36 hours, at least 48 hours, or at least 72 hours. In some embodiments, the invention features a pharmaceutical composition comprising a dose of mRNA sufficient to maintain at least 50% of a reference plasma ACADVL activity level or a normal physiologic plasma ACADVL activity 24 hours, 48 hours, 72 hours, 96 hours, 120 hours, 144 hours, or 168 hours post-administration. In some embodiments, the invention features a pharmaceutical composition comprising a dose of mRNA effective to result in an increase of plasma ACADVL activity for at least 24 hours, for at least 48 hours, for at least 72 hours, for at least 96 hours, for at least 120 hours, for at least 144 hours, or for at least 168 hours post-administration, wherein the increased plasma ACADVL activity levels are at least 5%, at least 10%, at least 15%, at least 20%, at least 25%, at least 30%, at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or more than 100% of a corresponding plasma ACADVL activity level or normal physiological ACADVL activity level.

[1110] In exemplary embodiments, the invention features a pharmaceutical composition comprising a dose of mRNA effective to result in an increase of liver, heart, kidney, or spleen ACADVL activity level to a level at or above a reference tissue ACADVL activity level or a normal physiologic level for at least 12 hours, at least 18 hours, at least 24 hours, at least 36 hours, at least 48 hours, or at least 72 hours. In some embodiments the invention features a pharmaceutical composition comprising a dose of mRNA sufficient to maintain at least 5%, at least 10%, at least 15%, at least 20%, at least 25%, at least 30%, at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 100% of a reference heart, liver, brain or skeletal muscle ACADVL activity or normal physiologic tissue ACADVL activity 24 hours, 48 hours, 72 hours, 96 hours, 120 hours, 144 hours, or 168 hours post-administration. In some embodiments, the invention features a pharmaceutical composition comprising a dose of mRNA effective to result in an increase of tissue (e.g., heart, liver, brain or skeletal muscle) ACADVL activity level for at least 24 hours, for at least 48 hours, for at least 72 hours, for at least 96 hours, for at least 120 hours, for at least 144 hours, or for at least 168 hours post-administration, wherein the increased tissue ACADVL activity level is at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least

80%, at least 90%, or more than 100% of a corresponding reference tissue ACADVL activity level or normal physiological tissue ACADVL activity level.

[1111] In exemplary embodiments, an mRNA therapy of the invention features a pharmaceutical composition comprising a single intravenous dose of mRNA that results in the above-described levels of activity. In another embodiment, an mRNA therapy of the invention features a pharmaceutical composition which can be administered in multiple single unit intravenous doses of mRNA that maintain the above-described levels of activity.

(iii) ACADVL Biomarkers

[1112] In some embodiments, the administration of an effective amount of a polynucleotide, pharmaceutical composition or formulation of the invention reduces the levels of a biomarker of VLCADD, e.g., acylcarnitine or an acylcarnitine metabolite. In some embodiments, the administration of the polynucleotide, pharmaceutical composition or formulation of the invention results in reduction in the level of one or more biomarkers of VLCADD, e.g., acylcarnitine or an acylcarnitine metabolite, within a short period of time after administration of the polynucleotide, pharmaceutical composition or formulation of the invention.

[1113] In some embodiments, the level of one or more biomarkers of VLCADD, e.g., acylcarnitine, is measured in blood. In some embodiments, the level of one or more biomarkers is measured in a component of blood, for example in plasma or serum. Methods of obtaining blood and components of blood and for measuring the level of biomarkers in blood or a component of blood are known in the art. In some embodiments, the level of one or more biomarkers of VLCADD, e.g., acylcarnitine, is measured in a dried blood spot. Methods of determining the level of biomarkers such as acylcarnitine are known in the art, for example, the methods in Rinaldo et al., *Genet Med* 10(2):151–156 (2008), which is hereby incorporated by reference herein, may be used in determining the level of acylcarnitine. In embodiments where levels of acylcarnitine are determined in dried blood spots, the method of determination can be mass spectrometry methods known in the art, for example, the mass spectrometry methods in Rashed et al., *Pediatric Research* 38:324-331 (1995), which is hereby incorporated by reference herein, may be used.

- [1114] In some embodiments, the level of one or more biomarkers of VLCADD, e.g., acylcarnitine, is measured in urine. Methods of obtaining urine and for measuring the level of biomarkers in urine are known in the art. In some embodiments, the level of one or more biomarkers of VLCADD, e.g., acylcarnitine, is measured in bile. Methods of obtaining bile and for measuring the level of biomarkers in bile are known in the art. In some embodiments, the level of one or more biomarkers of VLCADD, e.g., acylcarnitine, is measured in liver tissue. Methods of obtaining liver tissue, e.g. biopsy, and for measuring the level of biomarkers in liver tissue are known in the art.
- [1115] In some embodiments, the blood, plasma or serum level of acylcarnitine is reduced to less than about 0.6 $\mu\text{mol/L}$, about 0.7 $\mu\text{mol/L}$, about 0.8 $\mu\text{mol/L}$, about 0.9 $\mu\text{mol/L}$, about 1.0 $\mu\text{mol/L}$, about 1.1 $\mu\text{mol/L}$ or about 1.2 $\mu\text{mol/L}$ in a subject having VLCADD, for at least 24 hours, at least 48 hours, at least 72 hours, at least 96 hours, or at least 120 hours post-administration of a pharmaceutical composition or polynucleotide as described herein. Reference levels of acylcarnitine in the blood, plasma or serum or subjects having VLCADD and in subjects that do not have VLCADD can be found in the art, for example, in Leslie et al., Very Long-Chain Acyl-Coenzyme A Dehydrogenase Deficiency, GeneReviews 2009 (available at <https://www.ncbi.nlm.nih.gov/books/NBK6816/>) and in McHugh et al., Genet Med 13(3)230-254 (2011), both of which are hereby incorporated by reference herein.
- [1116] In embodiments where acylcarnitine is the biomarker measured, the acylcarnitine measured is an acylcarnitine metabolite. In some embodiments, the acylcarnitine metabolite measured is C12:1, C14:1, C14:2, C14, C16, C18, C18:1 acylcarnitine, or combinations thereof (where the fatty acid chains of the acylcarnitine are named according to the usual fatty acid naming conventions, e.g., C:D where C is the number of carbons in the fatty acid group and D is the number of double bonds, e.g., C12:1 has a fatty acid chain of 12 carbons with one double bond). In some embodiments, the acylcarnitine metabolite measured is C12:1, C14:1, C14:2, C14 acylcarnitine, or combinations thereof.
- [1117] Further aspects of the invention feature determining the level (or levels) of a biomarker, e.g., acylcarnitine and/or acylcarnitine metabolites, determined in a sample as compared to a level (e.g., a reference level) of the same or another biomarker in another sample, e.g., from the same patient, from another patient, from a control and/or from the same or different time points, and/or a physiologic level, and/or an elevated level, and/or

a supraphysiologic level, and/or a level of a control. The skilled artisan will be familiar with physiologic levels of biomarkers, for example, levels in normal or wild type animals, normal or healthy subjects, and the like, in particular, the level or levels characteristic of subjects who are healthy and/or normal functioning. As used herein, the phrase “elevated level” means amounts greater than normally found in a normal or wild type preclinical animal or in a normal or healthy subject, e.g. a human subject. As used herein, the term “supraphysiologic” means amounts greater than normally found in a normal or wild type preclinical animal or in a normal or healthy subject, e.g. a human subject, optionally producing a significantly enhanced physiologic response. As used herein, the term “comparing” or “compared to” preferably means the mathematical comparison of the two or more values, e.g., of the levels of the biomarker(s). It will thus be readily apparent to the skilled artisan whether one of the values is higher, lower or identical to another value or group of values if at least two of such values are compared with each other. Comparing or comparison to can be in the context, for example, of comparing to a control value in said subject prior to administration (e.g., in a person in need of treatment for VLCADD) or in a normal or healthy subject. Comparing or comparison to can also be in the context, for example, of comparing to a control value, e.g., as compared to a reference level in said subject prior to administration (e.g., in a person in need of treatment for VLCADD) or in a normal or healthy subject.

[1118] As used herein, a “control” is preferably a sample from a subject wherein the VLCADD status of said subject is known. In one embodiment, a control is a sample of a healthy patient. In another embodiment, the control is a sample from at least one subject having a known VLCADD status, for example, a severe, mild, or healthy VLCADD status, e.g. a control patient. In another embodiment, the control is a sample from a subject not being treated for VLCADD. In a still further embodiment, the control is a sample from a single subject or a pool of samples from different subjects and/or samples taken from the subject(s) at different time points.

[1119] The term “level” or “level of a biomarker” as used herein, preferably means the mass, weight or concentration of a biomarker of the invention within a sample or a subject. Biomarkers of the invention include, for example, acylcarnitine and/or acylcarnitine metabolites. It will be understood by the skilled artisan that in certain embodiments the sample may be subjected to, e.g., one or more of the following: substance purification, precipitation, separation, e.g. centrifugation and/or HPLC and

subsequently subjected to determining the level of the biomarker, e.g. using an ELISA assay or mass spectrometric analysis. In exemplary embodiments, LC-MS can be used as a means for determining the level of a biomarker according to the invention.

[1120] In some embodiments, the blood, plasma or serum level of acylcarnitine is reduced at least 1.5-fold, at least 2-fold, at least 3-fold, at least 5-fold, at least 10-fold, at least 12 fold, at least 15 fold, or at least 20-fold compared to a reference ammonia blood, plasma or serum level in a subject having VLCADD, for at least 24 hours, at least 48 hours, at least 72 hours, at least 96 hours, or at least 120 hours post-administration of a pharmaceutical composition or polynucleotide as described herein. Certain embodiments an mRNA of the invention can be used express an ACADVL polypeptide at a level sufficient to reduce the plasma or tissue levels of acylcarnitine and/or acylcarnitine metabolites in a VLCADD patient to less than about 95, 90, 80, 70, 60, 50, 45, 40, 35, 30, 25, 20, 15, 10, or 5 percent of the a reference level in said subject for at least 24 hours, at least 48 hours, at least 72 hours, at least 96 hours, at least 120 hours, at least 144 hours, at least one week, at least two weeks, at least three weeks, at least four weeks, at least 5 weeks, at least 6 weeks, at least 7 weeks, at least eight weeks, at least 9 weeks, or at least 10 weeks post-administration.

[1121] The term "determining the level" of a biomarker as used herein can mean methods which include quantifying an amount of at least one substance in a sample from a subject, for example, in a bodily fluid from the subject (e.g., serum, plasma, urine, blood, lymph, fecal, etc.) or in a tissue of the subject (e.g., heart, liver, brain or skeletal muscle, etc.).

[1122] The term "reference level" as used herein can refer to levels (e.g., of a biomarker) in a subject prior to administration of an mRNA therapy of the invention (e.g., in a person in need of treatment for VLCADD) or in a normal or healthy subject.

[1123] As used herein, the term "normal subject" or "healthy subject" refers to a subject not suffering from symptoms associated with VLCADD. Moreover, a subject will be considered to be normal (or healthy) if it has no mutation of the functional portions or domains of the *ACADVL* gene and/or no mutation of the *ACADVL* gene resulting in a reduction of or deficiency of the enzyme ACADVL or the activity thereof, resulting in symptoms associated with VLCADD. Said mutations will be detected if a sample from the subject is subjected to a genetic testing for such ACADVL mutations. In exemplary embodiments of the present invention, a sample from a healthy subject is used as a

control sample, or the known or standardized value for the level of biomarker from samples of healthy or normal subjects is used as a control.

[1124] In some embodiments, comparing the level of the biomarker in a sample from a subject in need of treatment for VLCADD, a subject in need of prevention of VLCADD, or a subject being treated for VLCADD to a control level of the biomarker comprises comparing the level of the biomarker in the sample from the subject (in need of treatment or being treated for VLCADD) to a baseline or reference level, wherein if a level of the biomarker in the sample from the subject (in need of treatment or being treated for VLCADD) is elevated, increased or higher compared to the baseline or reference level, this is indicative that the subject is in need of treatment for VLCADD and/or is in need of treatment; and/or wherein if a level of the biomarker in the sample from the subject (in need of treatment or being treated for VLCADD) is decreased or lower compared to the baseline level this is indicative that the subject is not suffering from, is successfully being treated for VLCADD, or is not in need of treatment for VLCADD. The stronger the reduction (e.g., at least 2-fold, at least 3-fold, at least 4-fold, at least 5-fold, at least 6-fold, at least 7-fold, at least 8-fold, at least 10-fold, at least 20-fold, at least 30 fold, at least 40-fold, at least 50-fold reduction and/or at least 5%, at least 10%, at least 20%, at least 30% at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, or at least 100% reduction) of the level of a biomarker within a certain time period, e.g., within 6 hours, within 12 hours, 24 hours, 36 hours, 48 hours, 60 hours, or 72 hours, and/or for a certain duration of time, e.g., 48 hours, 72 hours, 96 hours, 120 hours, 144 hours, 1 week, 2 weeks, 3 weeks, 4 weeks, 1 month, 2 months, 3 months, 4 months, 5 months, 6 months, 7 months, 8 months, 9 months, 10 months, 11 months, 12 months, 18 months, 24 months, etc. the more successful is a therapy, such as for example an mRNA therapy of the invention (e.g., a single dose or a multiple regimen).

[1125] A reduction of at least about 5%, at least about 10%, at least about 20%, at least about 30%, at least about 40%, at least about 50%, at least about 60%, at least about 70%, at least about 80%, at least about 90%, at least 100% or more of the level of biomarker, in particular, in bodily fluid (e.g., plasma) or in tissue(s) in a subject (e.g., heart, liver, brain or skeletal muscle), for example acylcarnitine and/or acylcarnitine metabolites, within 1, 2, 3, 4, 5, 6 or more days following administration is indicative of a dose suitable for successful treatment VLCADD, wherein reduction as used herein, preferably means that the level of biomarker determined at the end of a specified time period (e.g., post-

administration, for example, of a single intravenous dose) is compared to the level of the same biomarker determined at the beginning of said time period (e.g., pre-administration of said dose). Exemplary time periods include 12, 24, 48, 72, 96, 120, 144, or 168 hours post administration, in particular 24, 48, 72 or 96 hours post administration.

[1126] A sustained reduction in substrate levels (e.g., biomarkers such as acylcarnitine and/or acylcarnitine metabolites) is particularly indicative of mRNA therapeutic dosing and/or administration regimens successful for treatment of VLCADD. Such sustained reduction can be referred to herein as “duration” of effect. In exemplary embodiments, a reduction of at least about 10%, at least about 20%, at least about 30%, at least about 40%, at least about 50%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, or at least about 95% or more of the level of biomarker, in particular, in a bodily fluid (e.g., plasma) or in tissue(s) in a subject (e.g., heart, liver, brain or skeletal muscle) within 1, 2, 3, 4, 5, 6, 7, 8 or more days following administration is indicative of a successful therapeutic approach. In exemplary embodiments, sustained reduction in substrate (e.g., biomarker) levels in one or more samples (e.g., fluids and/or tissues) is preferred. For example, mRNA therapies resulting in sustained reduction in acylcarnitine and/or acylcarnitine metabolites (as defined herein), optionally in combination with sustained reduction of said biomarker in at least one tissue, preferably two, three, four, five or more tissues, is indicative of successful treatment.

[1127] In some embodiments, a single dose of an mRNA therapy of the invention is about 0.2 to about 0.8 mpk, about 0.3 to about 0.7 mpk, about 0.4 to about 0.8 mpk, or about 0.5 mpk. In another embodiment, a single dose of an mRNA therapy of the invention is less than 1.5 mpk, less than 1.25 mpk, less than 1 mpk, less than 0.75 mpk, or less than 0.5 mpk.

26. Compositions and Formulations for Use

[1128] Certain aspects of the invention are directed to compositions or formulations comprising any of the polynucleotides disclosed above.

[1129] In some embodiments, the composition or formulation comprises:

(i) a polynucleotide (e.g., a RNA, e.g., an mRNA) comprising a sequence-optimized nucleotide sequence (e.g., an ORF) encoding an ACADVL polypeptide (e.g., the wild-type sequence, functional fragment, or variant thereof), wherein the

polynucleotide comprises at least one chemically modified nucleobase, e.g., 5-methoxyuracil (e.g., wherein at least about 25%, at least about 30%, at least about 40%, at least about 50%, at least about 60%, at least about 70%, at least about 80%, at least about 90%, at least about 95%, at least about 99%, or 100% of the uracils are 5-methoxyuracils), and wherein the polynucleotide further comprises a miRNA binding site, e.g., a miRNA binding site that binds to miR-142 (e.g., a miR-142-3p or miR-142-5p binding site) and/or a miRNA binding site that binds to miR-126 (e.g., a miR-126-3p or miR-126-5p binding site); and

(ii) a delivery agent comprising, e.g., a compound having the Formula (I), e.g., any of Compounds 1-232, e.g., Compound 18; a compound having the Formula (III), (IV), (V), or (VI), e.g., any of Compounds 233-342, e.g., Compound 236; or a compound having the Formula (VIII), e.g., any of Compounds 419-428, e.g., Compound 428, or any combination thereof.

[1130] In some embodiments, the uracil or thymine content of the ORF relative to the theoretical minimum uracil or thymine content of a nucleotide sequence encoding the ACADVL polypeptide (%U_{TM} or %T_{TM}), is between about 100% and about 150%.

[1131] In some embodiments, the polynucleotides, compositions or formulations above are used to treat and/or prevent an ACADVL-related diseases, disorders or conditions, e.g., VLCADD.

27. Forms of Administration

[1132] The polynucleotides, pharmaceutical compositions and formulations of the invention described above can be administered by any route that results in a therapeutically effective outcome. These include, but are not limited to enteral (into the intestine), gastroenteral, epidural (into the dura matter), oral (by way of the mouth), transdermal, peridural, intracerebral (into the cerebrum), intracerebroventricular (into the cerebral ventricles), epicutaneous (application onto the skin), intradermal, (into the skin itself), subcutaneous (under the skin), nasal administration (through the nose), intravenous (into a vein), intravenous bolus, intravenous drip, intraarterial (into an artery), intramuscular (into a muscle), intracardiac (into the heart), intraosseous infusion (into the bone marrow), intrathecal (into the spinal canal), intraperitoneal, (infusion or injection into the peritoneum), intravesical infusion, intravitreal, (through the eye), intracavernous injection (into a pathologic cavity) intracavitary (into the base of the penis), intravaginal

administration, intrauterine, extra-amniotic administration, transdermal (diffusion through the intact skin for systemic distribution), transmucosal (diffusion through a mucous membrane), transvaginal, insufflation (snorting), sublingual, sublabial, enema, eye drops (onto the conjunctiva), in ear drops, auricular (in or by way of the ear), buccal (directed toward the cheek), conjunctival, cutaneous, dental (to a tooth or teeth), electro-osmosis, endocervical, endosinusal, endotracheal, extracorporeal, hemodialysis, infiltration, interstitial, intra-abdominal, intra-amniotic, intra-articular, intrabiliary, intrabronchial, intrabursal, intracartilaginous (within a cartilage), intracaudal (within the cauda equine), intracisternal (within the cisterna magna cerebellomedularis), intracorneal (within the cornea), dental intracornal, intracoronary (within the coronary arteries), intracorporus cavernosum (within the dilatable spaces of the corporus cavernosa of the penis), intradiscal (within a disc), intraductal (within a duct of a gland), intraduodenal (within the duodenum), intradural (within or beneath the dura), intraepidermal (to the epidermis), intraesophageal (to the esophagus), intragastric (within the stomach), intragingival (within the gingivae), intraileal (within the distal portion of the small intestine), intralesional (within or introduced directly to a localized lesion), intraluminal (within a lumen of a tube), intralymphatic (within the lymph), intramedullary (within the marrow cavity of a bone), intrameningeal (within the meninges), intraocular (within the eye), intraovarian (within the ovary), intrapericardial (within the pericardium), intrapleural (within the pleura), intraprostatic (within the prostate gland), intrapulmonary (within the lungs or its bronchi), intrasinal (within the nasal or periorbital sinuses), intraspinal (within the vertebral column), intrasynovial (within the synovial cavity of a joint), intratendinous (within a tendon), intratesticular (within the testicle), intrathecal (within the cerebrospinal fluid at any level of the cerebrospinal axis), intrathoracic (within the thorax), intratubular (within the tubules of an organ), intratympanic (within the aurus media), intravascular (within a vessel or vessels), intraventricular (within a ventricle), iontophoresis (by means of electric current where ions of soluble salts migrate into the tissues of the body), irrigation (to bathe or flush open wounds or body cavities), laryngeal (directly upon the larynx), nasogastric (through the nose and into the stomach), occlusive dressing technique (topical route administration that is then covered by a dressing that occludes the area), ophthalmic (to the external eye), oropharyngeal (directly to the mouth and pharynx), parenteral, percutaneous, periarticular, peridural, perineural, periodontal, rectal, respiratory (within the respiratory tract by inhaling orally or nasally for local or systemic

effect), retrobulbar (behind the pons or behind the eyeball), intramyocardial (entering the myocardium), soft tissue, subarachnoid, subconjunctival, submucosal, topical, transplacental (through or across the placenta), transtracheal (through the wall of the trachea), transtympanic (across or through the tympanic cavity), ureteral (to the ureter), urethral (to the urethra), vaginal, caudal block, diagnostic, nerve block, biliary perfusion, cardiac perfusion, photopheresis or spinal. In specific embodiments, compositions can be administered in a way that allows them cross the blood-brain barrier, vascular barrier, or other epithelial barrier. In some embodiments, a formulation for a route of administration can include at least one inactive ingredient.

[1133] The polynucleotides of the present invention (e.g., a polynucleotide comprising a nucleotide sequence encoding an ACADVL polypeptide or a functional fragment or variant thereof) can be delivered to a cell naked. As used herein in, "naked" refers to delivering polynucleotides free from agents that promote transfection. The naked polynucleotides can be delivered to the cell using routes of administration known in the art and described herein.

[1134] The polynucleotides of the present invention (e.g., a polynucleotide comprising a nucleotide sequence encoding an ACADVL polypeptide or a functional fragment or variant thereof) can be formulated, using the methods described herein. The formulations can contain polynucleotides that can be modified and/or unmodified. The formulations can further include, but are not limited to, cell penetration agents, a pharmaceutically acceptable carrier, a delivery agent, a bioerodible or biocompatible polymer, a solvent, and a sustained-release delivery depot. The formulated polynucleotides can be delivered to the cell using routes of administration known in the art and described herein.

[1135] A pharmaceutical composition for parenteral administration can comprise at least one inactive ingredient. Any or none of the inactive ingredients used can have been approved by the US Food and Drug Administration (FDA). A non-exhaustive list of inactive ingredients for use in pharmaceutical compositions for parenteral administration includes hydrochloric acid, mannitol, nitrogen, sodium acetate, sodium chloride and sodium hydroxide.

[1136] Injectable preparations, for example, sterile injectable aqueous or oleaginous suspensions can be formulated according to the known art using suitable dispersing agents, wetting agents, and/or suspending agents. Sterile injectable preparations can be sterile injectable solutions, suspensions, and/or emulsions in nontoxic parenterally

acceptable diluents and/or solvents, for example, as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that can be employed are water, Ringer's solution, U.S.P., and isotonic sodium chloride solution. Sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil can be employed including synthetic mono- or diglycerides. Fatty acids such as oleic acid can be used in the preparation of injectables. The sterile formulation can also comprise adjuvants such as local anesthetics, preservatives and buffering agents.

[1137] Injectable formulations can be sterilized, for example, by filtration through a bacterial-retaining filter, and/or by incorporating sterilizing agents in the form of sterile solid compositions that can be dissolved or dispersed in sterile water or other sterile injectable medium prior to use.

[1138] Injectable formulations can be for direct injection into a region of a tissue, organ and/or subject. As a non-limiting example, a tissue, organ and/or subject can be directly injected a formulation by intramyocardial injection into the ischemic region. (*See, e.g., Zangi et al. Nature Biotechnology 2013; the contents of which are herein incorporated by reference in its entirety*).

[1139] In order to prolong the effect of an active ingredient, it is often desirable to slow the absorption of the active ingredient from subcutaneous or intramuscular injection. This can be accomplished by the use of a liquid suspension of crystalline or amorphous material with poor water solubility. The rate of absorption of the drug then depends upon its rate of dissolution which, in turn, can depend upon crystal size and crystalline form. Alternatively, delayed absorption of a parenterally administered drug form is accomplished by dissolving or suspending the drug in an oil vehicle. Injectable depot forms are made by forming microencapsule matrices of the drug in biodegradable polymers such as polylactide-polyglycolide. Depending upon the ratio of drug to polymer and the nature of the particular polymer employed, the rate of drug release can be controlled. Examples of other biodegradable polymers include poly(orthoesters) and poly(anhydrides). Depot injectable formulations are prepared by entrapping the drug in liposomes or microemulsions that are compatible with body tissues.

28. Kits and Devices**a. Kits**

[1140] The invention provides a variety of kits for conveniently and/or effectively using the claimed nucleotides of the present invention. Typically kits will comprise sufficient amounts and/or numbers of components to allow a user to perform multiple treatments of a subject(s) and/or to perform multiple experiments.

[1141] In one aspect, the present invention provides kits comprising the molecules (polynucleotides) of the invention.

[1142] Said kits can be for protein production, comprising a first polynucleotides comprising a translatable region. The kit can further comprise packaging and instructions and/or a delivery agent to form a formulation composition. The delivery agent can comprise a saline, a buffered solution, a lipidoid or any delivery agent disclosed herein.

[1143] In some embodiments, the buffer solution can include sodium chloride, calcium chloride, phosphate and/or EDTA. In another embodiment, the buffer solution can include, but is not limited to, saline, saline with 2mM calcium, 5% sucrose, 5% sucrose with 2mM calcium, 5% Mannitol, 5% Mannitol with 2mM calcium, Ringer's lactate, sodium chloride, sodium chloride with 2mM calcium and mannose (*See, e.g.*, U.S. Pub. No. 20120258046; herein incorporated by reference in its entirety). In a further embodiment, the buffer solutions can be precipitated or it can be lyophilized. The amount of each component can be varied to enable consistent, reproducible higher concentration saline or simple buffer formulations. The components can also be varied in order to increase the stability of modified RNA in the buffer solution over a period of time and/or under a variety of conditions. In one aspect, the present invention provides kits for protein production, comprising: a polynucleotide comprising a translatable region, provided in an amount effective to produce a desired amount of a protein encoded by the translatable region when introduced into a target cell; a second polynucleotide comprising an inhibitory nucleic acid, provided in an amount effective to substantially inhibit the innate immune response of the cell; and packaging and instructions.

[1144] In one aspect, the present invention provides kits for protein production, comprising a polynucleotide comprising a translatable region, wherein the polynucleotide exhibits reduced degradation by a cellular nuclease, and packaging and instructions.

[1145] In one aspect, the present invention provides kits for protein production, comprising a polynucleotide comprising a translatable region, wherein the polynucleotide exhibits reduced degradation by a cellular nuclease, and a mammalian cell suitable for translation of the translatable region of the first nucleic acid.

b. Devices

[1146] The present invention provides for devices that can incorporate polynucleotides that encode polypeptides of interest. These devices contain in a stable formulation the reagents to synthesize a polynucleotide in a formulation available to be immediately delivered to a subject in need thereof, such as a human patient

[1147] Devices for administration can be employed to deliver the polynucleotides of the present invention according to single, multi- or split-dosing regimens taught herein. Such devices are taught in, for example, International Application Publ. No. WO2013151666, the contents of which are incorporated herein by reference in their entirety.

[1148] Method and devices known in the art for multi-administration to cells, organs and tissues are contemplated for use in conjunction with the methods and compositions disclosed herein as embodiments of the present invention. These include, for example, those methods and devices having multiple needles, hybrid devices employing for example lumens or catheters as well as devices utilizing heat, electric current or radiation driven mechanisms.

[1149] According to the present invention, these multi-administration devices can be utilized to deliver the single, multi- or split doses contemplated herein. Such devices are taught for example in, International Application Publ. No. WO2013151666, the contents of which are incorporated herein by reference in their entirety.

[1150] In some embodiments, the polynucleotide is administered subcutaneously or intramuscularly via at least 3 needles to three different, optionally adjacent, sites simultaneously, or within a 60 minutes period (*e.g.*, administration to 4, 5, 6, 7, 8, 9, or 10 sites simultaneously or within a 60 minute period).

c. Methods and Devices utilizing catheters and/or lumens

[1151] Methods and devices using catheters and lumens can be employed to administer the polynucleotides of the present invention on a single, multi- or split dosing schedule. Such methods and devices are described in International Application Publication No.

WO2013151666, the contents of which are incorporated herein by reference in their entirety.

d. Methods and Devices utilizing electrical current

[1152] Methods and devices utilizing electric current can be employed to deliver the polynucleotides of the present invention according to the single, multi- or split dosing regimens taught herein. Such methods and devices are described in International Application Publication No. WO2013151666, the contents of which are incorporated herein by reference in their entirety.

29. Definitions

[1153] In order that the present disclosure can be more readily understood, certain terms are first defined. As used in this application, except as otherwise expressly provided herein, each of the following terms shall have the meaning set forth below. Additional definitions are set forth throughout the application.

[1154] The invention includes embodiments in which exactly one member of the group is present in, employed in, or otherwise relevant to a given product or process. The invention includes embodiments in which more than one, or all of the group members are present in, employed in, or otherwise relevant to a given product or process.

[1155] In this specification and the appended claims, the singular forms "a", "an" and "the" include plural referents unless the context clearly dictates otherwise. The terms "a" (or "an"), as well as the terms "one or more," and "at least one" can be used interchangeably herein. In certain aspects, the term "a" or "an" means "single." In other aspects, the term "a" or "an" includes "two or more" or "multiple."

[1156] Furthermore, "and/or" where used herein is to be taken as specific disclosure of each of the two specified features or components with or without the other. Thus, the term "and/or" as used in a phrase such as "A and/or B" herein is intended to include "A and B," "A or B," "A" (alone), and "B" (alone). Likewise, the term "and/or" as used in a phrase such as "A, B, and/or C" is intended to encompass each of the following aspects: A, B, and C; A, B, or C; A or C; A or B; B or C; A and C; A and B; B and C; A (alone); B (alone); and C (alone).

[1157] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this

disclosure is related. For example, the Concise Dictionary of Biomedicine and Molecular Biology, Juo, Pei-Show, 2nd ed., 2002, CRC Press; The Dictionary of Cell and Molecular Biology, 3rd ed., 1999, Academic Press; and the Oxford Dictionary Of Biochemistry And Molecular Biology, Revised, 2000, Oxford University Press, provide one of skill with a general dictionary of many of the terms used in this disclosure.

[1158] Wherever aspects are described herein with the language "comprising," otherwise analogous aspects described in terms of "consisting of" and/or "consisting essentially of" are also provided.

[1159] Units, prefixes, and symbols are denoted in their Système International de Unites (SI) accepted form. Numeric ranges are inclusive of the numbers defining the range. Where a range of values is recited, it is to be understood that each intervening integer value, and each fraction thereof, between the recited upper and lower limits of that range is also specifically disclosed, along with each subrange between such values. The upper and lower limits of any range can independently be included in or excluded from the range, and each range where either, neither or both limits are included is also encompassed within the invention. Where a value is explicitly recited, it is to be understood that values which are about the same quantity or amount as the recited value are also within the scope of the invention. Where a combination is disclosed, each subcombination of the elements of that combination is also specifically disclosed and is within the scope of the invention. Conversely, where different elements or groups of elements are individually disclosed, combinations thereof are also disclosed. Where any element of an invention is disclosed as having a plurality of alternatives, examples of that invention in which each alternative is excluded singly or in any combination with the other alternatives are also hereby disclosed; more than one element of an invention can have such exclusions, and all combinations of elements having such exclusions are hereby disclosed.

[1160] Nucleotides are referred to by their commonly accepted single-letter codes. Unless otherwise indicated, nucleic acids are written left to right in 5' to 3' orientation. Nucleobases are referred to herein by their commonly known one-letter symbols recommended by the IUPAC-IUB Biochemical Nomenclature Commission. Accordingly, A represents adenine, C represents cytosine, G represents guanine, T represents thymine, U represents uracil.

- [1161] Amino acids are referred to herein by either their commonly known three letter symbols or by the one-letter symbols recommended by the IUPAC-IUB Biochemical Nomenclature Commission. Unless otherwise indicated, amino acid sequences are written left to right in amino to carboxy orientation.
- [1162] *About:* The term "about" as used in connection with a numerical value throughout the specification and the claims denotes an interval of accuracy, familiar and acceptable to a person skilled in the art; such interval of accuracy is $\pm 10\%$.
- [1163] Where ranges are given, endpoints are included. Furthermore, unless otherwise indicated or otherwise evident from the context and understanding of one of ordinary skill in the art, values that are expressed as ranges can assume any specific value or subrange within the stated ranges in different embodiments of the invention, to the tenth of the unit of the lower limit of the range, unless the context clearly dictates otherwise.
- [1164] *ACADVL Associated Disease:* As used herein the terms "ACADVL-associated disease" or "ACADVL-associated disorder" refer to diseases or disorders, respectively, which result from aberrant ACADVL activity (e.g., decreased activity or increased activity). As a non-limiting example, very long-chain acyl-CoA dehydrogenase deficiency is an ACADVL-associated disease. Numerous clinical variants of VLCADD are known in the art. See, e.g., www.omim.org/entry/609575.
- [1165] The terms "ACADVL enzymatic activity," "ACADVL activity," and "acylcarnitine metabolic activity" are used interchangeably in the present disclosure and refer to ACADVL's ability to catalyze fatty acid oxidation. Accordingly, a fragment or variant retaining or having ACADVL enzymatic activity or ACADVL activity refers to a fragment or variant that has measurable enzymatic catalytic activity in fatty acid oxidation.
- [1166] *Administered in combination:* As used herein, the term "administered in combination" or "combined administration" means that two or more agents are administered to a subject at the same time or within an interval such that there can be an overlap of an effect of each agent on the patient. In some embodiments, they are administered within about 60, 30, 15, 10, 5, or 1 minute of one another. In some embodiments, the administrations of the agents are spaced sufficiently closely together such that a combinatorial (e.g., a synergistic) effect is achieved.
- [1167] *Amino acid substitution:* The term "amino acid substitution" refers to replacing an amino acid residue present in a parent or reference sequence (e.g., a wild type ACADVL

sequence) with another amino acid residue. An amino acid can be substituted in a parent or reference sequence (e.g., a wild type ACADVL polypeptide sequence), for example, via chemical peptide synthesis or through recombinant methods known in the art.

Accordingly, a reference to a "substitution at position X" refers to the substitution of an amino acid present at position X with an alternative amino acid residue. In some aspects, substitution patterns can be described according to the schema AnY, wherein A is the single letter code corresponding to the amino acid naturally or originally present at position n, and Y is the substituting amino acid residue. In other aspects, substitution patterns can be described according to the schema An(YZ), wherein A is the single letter code corresponding to the amino acid residue substituting the amino acid naturally or originally present at position X, and Y and Z are alternative substituting amino acid residues.

[1168] In the context of the present disclosure, substitutions (even when they referred to as amino acid substitution) are conducted at the nucleic acid level, *i.e.*, substituting an amino acid residue with an alternative amino acid residue is conducted by substituting the codon encoding the first amino acid with a codon encoding the second amino acid.

[1169] *Animal:* As used herein, the term "animal" refers to any member of the animal kingdom. In some embodiments, "animal" refers to humans at any stage of development. In some embodiments, "animal" refers to non-human animals at any stage of development. In certain embodiments, the non-human animal is a mammal (*e.g.*, a rodent, a mouse, a rat, a rabbit, a monkey, a dog, a cat, a sheep, cattle, a primate, or a pig). In some embodiments, animals include, but are not limited to, mammals, birds, reptiles, amphibians, fish, and worms. In some embodiments, the animal is a transgenic animal, genetically-engineered animal, or a clone.

[1170] *Approximately:* As used herein, the term "approximately," as applied to one or more values of interest, refers to a value that is similar to a stated reference value. In certain embodiments, the term "approximately" refers to a range of values that fall within 25%, 20%, 19%, 18%, 17%, 16%, 15%, 14%, 13%, 12%, 11%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1%, or less in either direction (greater than or less than) of the stated reference value unless otherwise stated or otherwise evident from the context (except where such number would exceed 100% of a possible value).

[1171] *Associated with:* As used herein with respect to a disease, the term "associated with" means that the symptom, measurement, characteristic, or status in question is linked

to the diagnosis, development, presence, or progression of that disease. As association can, but need not, be causatively linked to the disease. For example, signs and symptoms, sequelae, or any effects causing a decrease in the quality of life of a patient of VLCADD are considered associated with VLCADD and in some embodiments of the present invention can be treated, ameliorated, or prevented by administering the polynucleotides of the present invention to a subject in need thereof.

[1172] When used with respect to two or more moieties, the terms "associated with," "conjugated," "linked," "attached," and "tethered," when used with respect to two or more moieties, means that the moieties are physically associated or connected with one another, either directly or via one or more additional moieties that serves as a linking agent, to form a structure that is sufficiently stable so that the moieties remain physically associated under the conditions in which the structure is used, *e.g.*, physiological conditions. An "association" need not be strictly through direct covalent chemical bonding. It can also suggest ionic or hydrogen bonding or a hybridization based connectivity sufficiently stable such that the "associated" entities remain physically associated.

[1173] *Bifunctional*: As used herein, the term "bifunctional" refers to any substance, molecule or moiety that is capable of or maintains at least two functions. The functions can affect the same outcome or a different outcome. The structure that produces the function can be the same or different. For example, bifunctional modified RNAs of the present invention can encode an ACADVL peptide (a first function) while those nucleosides that comprise the encoding RNA are, in and of themselves, capable of extending the half-life of the RNA (second function). In this example, delivery of the bifunctional modified RNA to a subject suffering from a protein deficiency would produce not only a peptide or protein molecule that can ameliorate or treat a disease or conditions, but would also maintain a population modified RNA present in the subject for a prolonged period of time. In other aspects, a bifunctional modified mRNA can be a chimeric or quimeric molecule comprising, for example, an RNA encoding an ACADVL peptide (a first function) and a second protein either fused to first protein or co-expressed with the first protein.

[1174] *Biocompatible*: As used herein, the term "biocompatible" means compatible with living cells, tissues, organs or systems posing little to no risk of injury, toxicity or rejection by the immune system.

- [1175] *Biodegradable*: As used herein, the term "biodegradable" means capable of being broken down into innocuous products by the action of living things.
- [1176] *Biologically active*: As used herein, the phrase "biologically active" refers to a characteristic of any substance that has activity in a biological system and/or organism. For instance, a substance that, when administered to an organism, has a biological effect on that organism, is considered to be biologically active. In particular embodiments, a polynucleotide of the present invention can be considered biologically active if even a portion of the polynucleotide is biologically active or mimics an activity considered biologically relevant.
- [1177] *Chimera*: As used herein, "chimera" is an entity having two or more incongruous or heterogeneous parts or regions. For example, a chimeric molecule can comprise a first part comprising an ACADVL polypeptide, and a second part (e.g., genetically fused to the first part) comprising a second therapeutic protein (e.g., a protein with a distinct enzymatic activity, an antigen binding moiety, or a moiety capable of extending the plasma half life of ACADVL, for example, an Fc region of an antibody).
- [1178] *Sequence Optimization*: The term "sequence optimization" refers to a process or series of processes by which nucleobases in a reference nucleic acid sequence are replaced with alternative nucleobases, resulting in a nucleic acid sequence with improved properties, e.g., improved protein expression or decreased immunogenicity.
- [1179] In general, the goal in sequence optimization is to produce a synonymous nucleotide sequence than encodes the same polypeptide sequence encoded by the reference nucleotide sequence. Thus, there are no amino acid substitutions (as a result of codon optimization) in the polypeptide encoded by the codon optimized nucleotide sequence with respect to the polypeptide encoded by the reference nucleotide sequence.
- [1180] *Codon substitution*: The terms "codon substitution" or "codon replacement" in the context of sequence optimization refer to replacing a codon present in a reference nucleic acid sequence with another codon. A codon can be substituted in a reference nucleic acid sequence, for example, via chemical peptide synthesis or through recombinant methods known in the art. Accordingly, references to a "substitution" or "replacement" at a certain location in a nucleic acid sequence (e.g., an mRNA) or within a certain region or subsequence of a nucleic acid sequence (e.g., an mRNA) refer to the substitution of a codon at such location or region with an alternative codon.

- [1181] As used herein, the terms "coding region" and "region encoding" and grammatical variants thereof, refer to an Open Reading Frame (ORF) in a polynucleotide that upon expression yields a polypeptide or protein.
- [1182] *Compound*: As used herein, the term "compound," is meant to include all stereoisomers and isotopes of the structure depicted. As used herein, the term "stereoisomer" means any geometric isomer (e.g., cis- and trans- isomer), enantiomer, or diastereomer of a compound. The present disclosure encompasses any and all stereoisomers of the compounds described herein, including stereomerically pure forms (e.g., geometrically pure, enantiomerically pure, or diastereomerically pure) and enantiomeric and stereoisomeric mixtures, e.g., racemates. Enantiomeric and stereomeric mixtures of compounds and means of resolving them into their component enantiomers or stereoisomers are well-known. "Isotopes" refers to atoms having the same atomic number but different mass numbers resulting from a different number of neutrons in the nuclei. For example, isotopes of hydrogen include tritium and deuterium. Further, a compound, salt, or complex of the present disclosure can be prepared in combination with solvent or water molecules to form solvates and hydrates by routine methods.
- [1183] *Contacting*: As used herein, the term "contacting" means establishing a physical connection between two or more entities. For example, contacting a mammalian cell with a nanoparticle composition means that the mammalian cell and a nanoparticle are made to share a physical connection. Methods of contacting cells with external entities both in vivo and ex vivo are well known in the biological arts. For example, contacting a nanoparticle composition and a mammalian cell disposed within a mammal can be performed by varied routes of administration (e.g., intravenous, intramuscular, intradermal, and subcutaneous) and can involve varied amounts of nanoparticle compositions. Moreover, more than one mammalian cell can be contacted by a nanoparticle composition.
- [1184] *Conservative amino acid substitution*: A "conservative amino acid substitution" is one in which the amino acid residue in a protein sequence is replaced with an amino acid residue having a similar side chain. Families of amino acid residues having similar side chains have been defined in the art, including basic side chains (e.g., lysine, arginine, or histidine), acidic side chains (e.g., aspartic acid or glutamic acid), uncharged polar side chains (e.g., glycine, asparagine, glutamine, serine, threonine, tyrosine, or cysteine), nonpolar side chains (e.g., alanine, valine, leucine, isoleucine, proline, phenylalanine,

methionine, or tryptophan), beta-branched side chains (*e.g.*, threonine, valine, isoleucine) and aromatic side chains (*e.g.*, tyrosine, phenylalanine, tryptophan, or histidine). Thus, if an amino acid in a polypeptide is replaced with another amino acid from the same side chain family, the amino acid substitution is considered to be conservative. In another aspect, a string of amino acids can be conservatively replaced with a structurally similar string that differs in order and/or composition of side chain family members.

- [1185] *Non-conservative amino acid substitution*: Non-conservative amino acid substitutions include those in which (i) a residue having an electropositive side chain (*e.g.*, Arg, His or Lys) is substituted for, or by, an electronegative residue (*e.g.*, Glu or Asp), (ii) a hydrophilic residue (*e.g.*, Ser or Thr) is substituted for, or by, a hydrophobic residue (*e.g.*, Ala, Leu, Ile, Phe or Val), (iii) a cysteine or proline is substituted for, or by, any other residue, or (iv) a residue having a bulky hydrophobic or aromatic side chain (*e.g.*, Val, His, Ile or Trp) is substituted for, or by, one having a smaller side chain (*e.g.*, Ala or Ser) or no side chain (*e.g.*, Gly).
- [1186] Other amino acid substitutions can be readily identified by workers of ordinary skill. For example, for the amino acid alanine, a substitution can be taken from any one of D-alanine, glycine, beta-alanine, L-cysteine and D-cysteine. For lysine, a replacement can be any one of D-lysine, arginine, D-arginine, homo-arginine, methionine, D-methionine, ornithine, or D- ornithine. Generally, substitutions in functionally important regions that can be expected to induce changes in the properties of isolated polypeptides are those in which (i) a polar residue, *e.g.*, serine or threonine, is substituted for (or by) a hydrophobic residue, *e.g.*, leucine, isoleucine, phenylalanine, or alanine; (ii) a cysteine residue is substituted for (or by) any other residue; (iii) a residue having an electropositive side chain, *e.g.*, lysine, arginine or histidine, is substituted for (or by) a residue having an electronegative side chain, *e.g.*, glutamic acid or aspartic acid; or (iv) a residue having a bulky side chain, *e.g.*, phenylalanine, is substituted for (or by) one not having such a side chain, *e.g.*, glycine. The likelihood that one of the foregoing non-conservative substitutions can alter functional properties of the protein is also correlated to the position of the substitution with respect to functionally important regions of the protein: some non-conservative substitutions can accordingly have little or no effect on biological properties.
- [1187] *Conserved*: As used herein, the term "conserved" refers to nucleotides or amino acid residues of a polynucleotide sequence or polypeptide sequence, respectively, that are

those that occur unaltered in the same position of two or more sequences being compared. Nucleotides or amino acids that are relatively conserved are those that are conserved amongst more related sequences than nucleotides or amino acids appearing elsewhere in the sequences.

[1188] In some embodiments, two or more sequences are said to be "completely conserved" if they are 100% identical to one another. In some embodiments, two or more sequences are said to be "highly conserved" if they are at least 70% identical, at least 80% identical, at least 90% identical, or at least 95% identical to one another. In some embodiments, two or more sequences are said to be "highly conserved" if they are about 70% identical, about 80% identical, about 90% identical, about 95%, about 98%, or about 99% identical to one another. In some embodiments, two or more sequences are said to be "conserved" if they are at least 30% identical, at least 40% identical, at least 50% identical, at least 60% identical, at least 70% identical, at least 80% identical, at least 90% identical, or at least 95% identical to one another. In some embodiments, two or more sequences are said to be "conserved" if they are about 30% identical, about 40% identical, about 50% identical, about 60% identical, about 70% identical, about 80% identical, about 90% identical, about 95% identical, about 98% identical, or about 99% identical to one another. Conservation of sequence can apply to the entire length of a polynucleotide or polypeptide or can apply to a portion, region or feature thereof.

[1189] *Controlled Release:* As used herein, the term "controlled release" refers to a pharmaceutical composition or compound release profile that conforms to a particular pattern of release to effect a therapeutic outcome.

[1190] *Cyclic or Cyclized:* As used herein, the term "cyclic" refers to the presence of a continuous loop. Cyclic molecules need not be circular, only joined to form an unbroken chain of subunits. Cyclic molecules such as the engineered RNA or mRNA of the present invention can be single units or multimers or comprise one or more components of a complex or higher order structure.

[1191] *Cytotoxic:* As used herein, "cytotoxic" refers to killing or causing injurious, toxic, or deadly effect on a cell (*e.g.*, a mammalian cell (*e.g.*, a human cell)), bacterium, virus, fungus, protozoan, parasite, prion, or a combination thereof.

[1192] *Delivering:* As used herein, the term "delivering" means providing an entity to a destination. For example, delivering a polynucleotide to a subject can involve administering a nanoparticle composition including the polynucleotide to the subject

(e.g., by an intravenous, intramuscular, intradermal, or subcutaneous route).

Administration of a nanoparticle composition to a mammal or mammalian cell can involve contacting one or more cells with the nanoparticle composition.

- [1193] *Delivery Agent*: As used herein, "delivery agent" refers to any substance that facilitates, at least in part, the *in vivo*, *in vitro*, or *ex vivo* delivery of a polynucleotide to targeted cells.
- [1194] *Destabilized*: As used herein, the term "destable," "destabilize," or "destabilizing region" means a region or molecule that is less stable than a starting, wild-type or native form of the same region or molecule.
- [1195] *Diastereomer*: As used herein, the term "diastereomer," means stereoisomers that are not mirror images of one another and are non-superimposable on one another.
- [1196] *Digest*: As used herein, the term "digest" means to break apart into smaller pieces or components. When referring to polypeptides or proteins, digestion results in the production of peptides.
- [1197] *Distal*: As used herein, the term "distal" means situated away from the center or away from a point or region of interest.
- [1198] *Domain*: As used herein, when referring to polypeptides, the term "domain" refers to a motif of a polypeptide having one or more identifiable structural or functional characteristics or properties (e.g., binding capacity, serving as a site for protein-protein interactions).
- [1199] *Dosing regimen*: As used herein, a "dosing regimen" or a "dosing regimen" is a schedule of administration or physician determined regimen of treatment, prophylaxis, or palliative care.
- [1200] *Effective Amount*: As used herein, the term "effective amount" of an agent is that amount sufficient to effect beneficial or desired results, for example, clinical results, and, as such, an "effective amount" depends upon the context in which it is being applied. For example, in the context of administering an agent that treats a protein deficiency (e.g., an ACADVL deficiency), an effective amount of an agent is, for example, an amount of mRNA expressing sufficient ACADVL to ameliorate, reduce, eliminate, or prevent the symptoms associated with the ACADVL deficiency, as compared to the severity of the symptom observed without administration of the agent. The term "effective amount" can be used interchangeably with "effective dose," "therapeutically effective amount," or "therapeutically effective dose."

- [1201] *Enantiomer*: As used herein, the term "enantiomer" means each individual optically active form of a compound of the invention, having an optical purity or enantiomeric excess (as determined by methods standard in the art) of at least 80% (*i.e.*, at least 90% of one enantiomer and at most 10% of the other enantiomer), at least 90%, or at least 98%.
- [1202] *Encapsulate*: As used herein, the term "encapsulate" means to enclose, surround or encase.
- [1203] *Encapsulation Efficiency*: As used herein, "encapsulation efficiency" refers to the amount of a polynucleotide that becomes part of a nanoparticle composition, relative to the initial total amount of polynucleotide used in the preparation of a nanoparticle composition. For example, if 97 mg of polynucleotide are encapsulated in a nanoparticle composition out of a total 100 mg of polynucleotide initially provided to the composition, the encapsulation efficiency can be given as 97%. As used herein, "encapsulation" can refer to complete, substantial, or partial enclosure, confinement, surrounding, or encasement.
- [1204] *Encoded protein cleavage signal*: As used herein, "encoded protein cleavage signal" refers to the nucleotide sequence that encodes a protein cleavage signal.
- [1205] *Engineered*: As used herein, embodiments of the invention are "engineered" when they are designed to have a feature or property, whether structural or chemical, that varies from a starting point, wild type or native molecule.
- [1206] *Enhanced Delivery*: As used herein, the term "enhanced delivery" means delivery of more (e.g., at least 1.5 fold more, at least 2-fold more, at least 3-fold more, at least 4-fold more, at least 5-fold more, at least 6-fold more, at least 7-fold more, at least 8-fold more, at least 9-fold more, at least 10-fold more) of a polynucleotide by a nanoparticle to a target tissue of interest (e.g., mammalian liver) compared to the level of delivery of a polynucleotide by a control nanoparticle to a target tissue of interest (e.g., MC3, KC2, or DLinDMA). The level of delivery of a nanoparticle to a particular tissue can be measured by comparing the amount of protein produced in a tissue to the weight of said tissue, comparing the amount of polynucleotide in a tissue to the weight of said tissue, comparing the amount of protein produced in a tissue to the amount of total protein in said tissue, or comparing the amount of polynucleotide in a tissue to the amount of total polynucleotide in said tissue. It will be understood that the enhanced delivery of a

nanoparticle to a target tissue need not be determined in a subject being treated, it can be determined in a surrogate such as an animal model (e.g., a rat model).

- [1207] *Exosome*: As used herein, "exosome" is a vesicle secreted by mammalian cells or a complex involved in RNA degradation.
- [1208] *Expression*: As used herein, "expression" of a nucleic acid sequence refers to one or more of the following events: (1) production of an mRNA template from a DNA sequence (e.g., by transcription); (2) processing of an mRNA transcript (e.g., by splicing, editing, 5' cap formation, and/or 3' end processing); (3) translation of an mRNA into a polypeptide or protein; and (4) post-translational modification of a polypeptide or protein.
- [1209] *Ex Vivo*: As used herein, the term "ex vivo" refers to events that occur outside of an organism (e.g., animal, plant, or microbe or cell or tissue thereof). Ex vivo events can take place in an environment minimally altered from a natural (e.g., in vivo) environment.
- [1210] *Feature*: As used herein, a "feature" refers to a characteristic, a property, or a distinctive element. When referring to polypeptides, "features" are defined as distinct amino acid sequence-based components of a molecule. Features of the polypeptides encoded by the polynucleotides of the present invention include surface manifestations, local conformational shape, folds, loops, half-loops, domains, half-domains, sites, termini or any combination thereof.
- [1211] *Formulation*: As used herein, a "formulation" includes at least a polynucleotide and one or more of a carrier, an excipient, and a delivery agent.
- [1212] *Fragment*: A "fragment," as used herein, refers to a portion. For example, fragments of proteins can comprise polypeptides obtained by digesting full-length protein isolated from cultured cells. In some embodiments, a fragment is a subsequences of a full length protein (e.g., ACADVL) wherein N-terminal, and/or C-terminal, and/or internal subsequences have been deleted. In some preferred aspects of the present invention, the fragments of a protein of the present invention are functional fragments.
- [1213] *Functional*: As used herein, a "functional" biological molecule is a biological molecule in a form in which it exhibits a property and/or activity by which it is characterized. Thus, a functional fragment of a polynucleotide of the present invention is a polynucleotide capable of expressing a functional ACADVL fragment. As used herein, a functional fragment of ACADVL refers to a fragment of wild type ACADVL (i.e., a fragment of any of its naturally occurring isoforms), or a mutant or variant thereof, wherein the fragment retains a least about 10%, at least about 15%, at least about 20%, at

least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, or at least about 95% of the biological activity of the corresponding full length protein.

[1214] *Helper Lipid*: As used herein, the term “helper lipid” refers to a compound or molecule that includes a lipidic moiety (for insertion into a lipid layer, e.g., lipid bilayer) and a polar moiety (for interaction with physiologic solution at the surface of the lipid layer). Typically the helper lipid is a phospholipid. A function of the helper lipid is to “complement” the amino lipid and increase the fusogenicity of the bilayer and/or to help facilitate endosomal escape, e.g., of nucleic acid delivered to cells. Helper lipids are also believed to be a key structural component to the surface of the LNP.

[1215] *Homology*: As used herein, the term "homology" refers to the overall relatedness between polymeric molecules, e.g. between nucleic acid molecules (e.g. DNA molecules and/or RNA molecules) and/or between polypeptide molecules. Generally, the term "homology" implies an evolutionary relationship between two molecules. Thus, two molecules that are homologous will have a common evolutionary ancestor. In the context of the present invention, the term homology encompasses both to identity and similarity.

[1216] In some embodiments, polymeric molecules are considered to be "homologous" to one another if at least 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 99% of the monomers in the molecule are identical (exactly the same monomer) or are similar (conservative substitutions). The term "homologous" necessarily refers to a comparison between at least two sequences (polynucleotide or polypeptide sequences).

[1217] *Identity*: As used herein, the term "identity" refers to the overall monomer conservation between polymeric molecules, e.g., between polynucleotide molecules (e.g. DNA molecules and/or RNA molecules) and/or between polypeptide molecules. Calculation of the percent identity of two polynucleotide sequences, for example, can be performed by aligning the two sequences for optimal comparison purposes (e.g., gaps can be introduced in one or both of a first and a second nucleic acid sequences for optimal alignment and non-identical sequences can be disregarded for comparison purposes). In certain embodiments, the length of a sequence aligned for comparison purposes is at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at

least 95%, or 100% of the length of the reference sequence. The nucleotides at corresponding nucleotide positions are then compared. When a position in the first sequence is occupied by the same nucleotide as the corresponding position in the second sequence, then the molecules are identical at that position. The percent identity between the two sequences is a function of the number of identical positions shared by the sequences, taking into account the number of gaps, and the length of each gap, which needs to be introduced for optimal alignment of the two sequences. The comparison of sequences and determination of percent identity between two sequences can be accomplished using a mathematical algorithm. When comparing DNA and RNA, thymine (T) and uracil (U) can be considered equivalent.

[1218] Suitable software programs are available from various sources, and for alignment of both protein and nucleotide sequences. One suitable program to determine percent sequence identity is bl2seq, part of the BLAST suite of program available from the U.S. government's National Center for Biotechnology Information BLAST web site (blast.ncbi.nlm.nih.gov). Bl2seq performs a comparison between two sequences using either the BLASTN or BLASTP algorithm. BLASTN is used to compare nucleic acid sequences, while BLASTP is used to compare amino acid sequences. Other suitable programs are, e.g., Needle, Stretcher, Water, or Matcher, part of the EMBOSS suite of bioinformatics programs and also available from the European Bioinformatics Institute (EBI) at www.ebi.ac.uk/Tools/psa.

[1219] Sequence alignments can be conducted using methods known in the art such as MAFFT, Clustal (ClustalW, Clustal X or Clustal Omega), MUSCLE, etc.

[1220] Different regions within a single polynucleotide or polypeptide target sequence that aligns with a polynucleotide or polypeptide reference sequence can each have their own percent sequence identity. It is noted that the percent sequence identity value is rounded to the nearest tenth. For example, 80.11, 80.12, 80.13, and 80.14 are rounded down to 80.1, while 80.15, 80.16, 80.17, 80.18, and 80.19 are rounded up to 80.2. It also is noted that the length value will always be an integer.

[1221] In certain aspects, the percentage identity "%ID" of a first amino acid sequence (or nucleic acid sequence) to a second amino acid sequence (or nucleic acid sequence) is calculated as $\%ID = 100 \times (Y/Z)$, where Y is the number of amino acid residues (or nucleobases) scored as identical matches in the alignment of the first and second sequences (as aligned by visual inspection or a particular sequence alignment program)

and Z is the total number of residues in the second sequence. If the length of a first sequence is longer than the second sequence, the percent identity of the first sequence to the second sequence will be higher than the percent identity of the second sequence to the first sequence.

[1222] One skilled in the art will appreciate that the generation of a sequence alignment for the calculation of a percent sequence identity is not limited to binary sequence-sequence comparisons exclusively driven by primary sequence data. It will also be appreciated that sequence alignments can be generated by integrating sequence data with data from heterogeneous sources such as structural data (e.g., crystallographic protein structures), functional data (e.g., location of mutations), or phylogenetic data. A suitable program that integrates heterogeneous data to generate a multiple sequence alignment is T-Coffee, available at www.tcoffee.org, and alternatively available, e.g., from the EBI. It will also be appreciated that the final alignment used to calculate percent sequence identity can be curated either automatically or manually.

[1223] *Immune response*: The term “immune response” refers to the action of, for example, lymphocytes, antigen presenting cells, phagocytic cells, granulocytes, and soluble macromolecules produced by the above cells or the liver (including antibodies, cytokines, and complement) that results in selective damage to, destruction of, or elimination from the human body of invading pathogens, cells or tissues infected with pathogens, cancerous cells, or, in cases of autoimmunity or pathological inflammation, normal human cells or tissues. In some cases, the administration of a nanoparticle comprising a lipid component and an encapsulated therapeutic agent can trigger an immune response, which can be caused by (i) the encapsulated therapeutic agent (e.g., an mRNA), (ii) the expression product of such encapsulated therapeutic agent (e.g., a polypeptide encoded by the mRNA), (iii) the lipid component of the nanoparticle, or (iv) a combination thereof.

[1224] *Inflammatory response*: “Inflammatory response” refers to immune responses involving specific and non-specific defense systems. A specific defense system reaction is a specific immune system reaction to an antigen. Examples of specific defense system reactions include antibody responses. A non-specific defense system reaction is an inflammatory response mediated by leukocytes generally incapable of immunological memory, e.g., macrophages, eosinophils and neutrophils. In some aspects, an immune

response includes the secretion of inflammatory cytokines, resulting in elevated inflammatory cytokine levels.

- [1225] *Inflammatory cytokines*: The term “inflammatory cytokine” refers to cytokines that are elevated in an inflammatory response. Examples of inflammatory cytokines include interleukin-6 (IL-6), CXCL1 (chemokine (C-X-C motif) ligand 1; also known as GRO α , interferon- γ (IFN γ), tumor necrosis factor α (TNF α), interferon γ -induced protein 10 (IP-10), or granulocyte-colony stimulating factor (G-CSF). The term inflammatory cytokines includes also other cytokines associated with inflammatory responses known in the art, e.g., interleukin-1 (IL-1), interleukin-8 (IL-8), interleukin-12 (IL-12), interleukin-13 (IL-13), interferon α (IFN- α), etc.
- [1226] *In Vitro*: As used herein, the term “in vitro” refers to events that occur in an artificial environment, e.g., in a test tube or reaction vessel, in cell culture, in a Petri dish, etc., rather than within an organism (e.g., animal, plant, or microbe).
- [1227] *In Vivo*: As used herein, the term “in vivo” refers to events that occur within an organism (e.g., animal, plant, or microbe or cell or tissue thereof).
- [1228] *Insertional and deletional variants*: “Insertional variants” when referring to polypeptides are those with one or more amino acids inserted immediately adjacent to an amino acid at a particular position in a native or starting sequence. “Immediately adjacent” to an amino acid means connected to either the alpha-carboxy or alpha-amino functional group of the amino acid. “Deletional variants” when referring to polypeptides are those with one or more amino acids in the native or starting amino acid sequence removed. Ordinarily, deletional variants will have one or more amino acids deleted in a particular region of the molecule.
- [1229] *Intact*: As used herein, in the context of a polypeptide, the term “intact” means retaining an amino acid corresponding to the wild type protein, e.g., not mutating or substituting the wild type amino acid. Conversely, in the context of a nucleic acid, the term “intact” means retaining a nucleobase corresponding to the wild type nucleic acid, e.g., not mutating or substituting the wild type nucleobase.
- [1230] *Ionizable amino lipid*: The term “ionizable amino lipid” includes those lipids having one, two, three, or more fatty acid or fatty alkyl chains and a pH-titratable amino head group (e.g., an alkylamino or dialkylamino head group). An ionizable amino lipid is typically protonated (i.e., positively charged) at a pH below the pKa of the amino head group and is substantially not charged at a pH above the pKa. Such ionizable amino lipids

include, but are not limited to DLin-MC3-DMA (MC3) and (13Z,165Z)-N,N-dimethyl-3-nonydocosa-13-16-dien-1-amine (L608).

[1231] *Isolated*: As used herein, the term “isolated” refers to a substance or entity that has been separated from at least some of the components with which it was associated (whether in nature or in an experimental setting). Isolated substances (*e.g.*, polynucleotides or polypeptides) can have varying levels of purity in reference to the substances from which they have been isolated. Isolated substances and/or entities can be separated from at least about 10%, about 20%, about 30%, about 40%, about 50%, about 60%, about 70%, about 80%, about 90%, or more of the other components with which they were initially associated. In some embodiments, isolated substances are more than about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, about 99%, or more than about 99% pure. As used herein, a substance is “pure” if it is substantially free of other components.

[1232] *Substantially isolated*: By “substantially isolated” is meant that the compound is substantially separated from the environment in which it was formed or detected. Partial separation can include, for example, a composition enriched in the compound of the present disclosure. Substantial separation can include compositions containing at least about 50%, at least about 60%, at least about 70%, at least about 80%, at least about 90%, at least about 95%, at least about 97%, or at least about 99% by weight of the compound of the present disclosure, or salt thereof.

[1233] A polynucleotide, vector, polypeptide, cell, or any composition disclosed herein which is “isolated” is a polynucleotide, vector, polypeptide, cell, or composition which is in a form not found in nature. Isolated polynucleotides, vectors, polypeptides, or compositions include those which have been purified to a degree that they are no longer in a form in which they are found in nature. In some aspects, a polynucleotide, vector, polypeptide, or composition which is isolated is substantially pure.

[1234] *Isomer*: As used herein, the term “isomer” means any tautomer, stereoisomer, enantiomer, or diastereomer of any compound of the invention. It is recognized that the compounds of the invention can have one or more chiral centers and/or double bonds and, therefore, exist as stereoisomers, such as double-bond isomers (*i.e.*, geometric E/Z isomers) or diastereomers (*e.g.*, enantiomers (*i.e.*, (+) or (-)) or cis/trans isomers). According to the invention, the chemical structures depicted herein, and therefore the compounds of the invention, encompass all of the corresponding stereoisomers, that is,

both the stereomerically pure form (*e.g.*, geometrically pure, enantiomerically pure, or diastereomerically pure) and enantiomeric and stereoisomeric mixtures, *e.g.*, racemates. Enantiomeric and stereoisomeric mixtures of compounds of the invention can typically be resolved into their component enantiomers or stereoisomers by well-known methods, such as chiral-phase gas chromatography, chiral-phase high performance liquid chromatography, crystallizing the compound as a chiral salt complex, or crystallizing the compound in a chiral solvent. Enantiomers and stereoisomers can also be obtained from stereomerically or enantiomerically pure intermediates, reagents, and catalysts by well-known asymmetric synthetic methods.

[1235] *Linker*: As used herein, a "linker" refers to a group of atoms, *e.g.*, 10-1,000 atoms, and can be comprised of the atoms or groups such as, but not limited to, carbon, amino, alkylamino, oxygen, sulfur, sulfoxide, sulfonyl, carbonyl, and imine. The linker can be attached to a modified nucleoside or nucleotide on the nucleobase or sugar moiety at a first end, and to a payload, *e.g.*, a detectable or therapeutic agent, at a second end. The linker can be of sufficient length as to not interfere with incorporation into a nucleic acid sequence. The linker can be used for any useful purpose, such as to form polynucleotide multimers (*e.g.*, through linkage of two or more chimeric polynucleotides molecules or IVT polynucleotides) or polynucleotides conjugates, as well as to administer a payload, as described herein. Examples of chemical groups that can be incorporated into the linker include, but are not limited to, alkyl, alkenyl, alkynyl, amido, amino, ether, thioether, ester, alkylene, heteroalkylene, aryl, or heterocyclyl, each of which can be optionally substituted, as described herein. Examples of linkers include, but are not limited to, unsaturated alkanes, polyethylene glycols (*e.g.*, ethylene or propylene glycol monomeric units, *e.g.*, diethylene glycol, dipropylene glycol, triethylene glycol, tripropylene glycol, tetraethylene glycol, or tetraethylene glycol), and dextran polymers and derivatives thereof. Other examples include, but are not limited to, cleavable moieties within the linker, such as, for example, a disulfide bond (-S-S-) or an azo bond (-N=N-), which can be cleaved using a reducing agent or photolysis. Non-limiting examples of a selectively cleavable bond include an amido bond can be cleaved for example by the use of tris(2-carboxyethyl)phosphine (TCEP), or other reducing agents, and/or photolysis, as well as an ester bond can be cleaved for example by acidic or basic hydrolysis.

[1236] *Methods of Administration*: As used herein, "methods of administration" can include intravenous, intramuscular, intradermal, subcutaneous, or other methods of

delivering a composition to a subject. A method of administration can be selected to target delivery (*e.g.*, to specifically deliver) to a specific region or system of a body.

- [1237] *Modified*: As used herein "modified" refers to a changed state or structure of a molecule of the invention. Molecules can be modified in many ways including chemically, structurally, and functionally. In some embodiments, the mRNA molecules of the present invention are modified by the introduction of non-natural nucleosides and/or nucleotides, *e.g.*, as it relates to the natural ribonucleotides A, U, G, and C. Noncanonical nucleotides such as the cap structures are not considered "modified" although they differ from the chemical structure of the A, C, G, U ribonucleotides.
- [1238] *Mucus*: As used herein, "mucus" refers to the natural substance that is viscous and comprises mucin glycoproteins.
- [1239] *Nanoparticle Composition*: As used herein, a "nanoparticle composition" is a composition comprising one or more lipids. Nanoparticle compositions are typically sized on the order of micrometers or smaller and can include a lipid bilayer. Nanoparticle compositions encompass lipid nanoparticles (LNPs), liposomes (*e.g.*, lipid vesicles), and lipoplexes. For example, a nanoparticle composition can be a liposome having a lipid bilayer with a diameter of 500 nm or less.
- [1240] *Naturally occurring*: As used herein, "naturally occurring" means existing in nature without artificial aid.
- [1241] *Non-human vertebrate*: As used herein, a "non-human vertebrate" includes all vertebrates except *Homo sapiens*, including wild and domesticated species. Examples of non-human vertebrates include, but are not limited to, mammals, such as alpaca, banteng, bison, camel, cat, cattle, deer, dog, donkey, gayal, goat, guinea pig, horse, llama, mule, pig, rabbit, reindeer, sheep water buffalo, and yak.
- [1242] *Nucleic acid sequence*: The terms "nucleic acid sequence," "nucleotide sequence," or "polynucleotide sequence" are used interchangeably and refer to a contiguous nucleic acid sequence. The sequence can be either single stranded or double stranded DNA or RNA, *e.g.*, an mRNA.
- [1243] The term "nucleic acid," in its broadest sense, includes any compound and/or substance that comprises a polymer of nucleotides. These polymers are often referred to as polynucleotides. Exemplary nucleic acids or polynucleotides of the invention include, but are not limited to, ribonucleic acids (RNAs), deoxyribonucleic acids (DNAs), threose nucleic acids (TNAs), glycol nucleic acids (GNAs), peptide nucleic acids (PNAs), locked

nucleic acids (LNAs, including LNA having a β -D-ribo configuration, α -LNA having an α -L-ribo configuration (a diastereomer of LNA), 2'-amino-LNA having a 2'-amino functionalization, and 2'-amino- α -LNA having a 2'-amino functionalization), ethylene nucleic acids (ENA), cyclohexenyl nucleic acids (CeNA) or hybrids or combinations thereof.

- [1244] The phrase "nucleotide sequence encoding" refers to the nucleic acid (*e.g.*, an mRNA or DNA molecule) coding sequence which encodes a polypeptide. The coding sequence can further include initiation and termination signals operably linked to regulatory elements including a promoter and polyadenylation signal capable of directing expression in the cells of an individual or mammal to which the nucleic acid is administered. The coding sequence can further include sequences that encode signal peptides.
- [1245] *Off-target*: As used herein, "off target" refers to any unintended effect on any one or more target, gene, or cellular transcript.
- [1246] *Open reading frame*: As used herein, "open reading frame" or "ORF" refers to a sequence which does not contain a stop codon in a given reading frame.
- [1247] *Operably linked*: As used herein, the phrase "operably linked" refers to a functional connection between two or more molecules, constructs, transcripts, entities, moieties or the like.
- [1248] *Optionally substituted*: Herein a phrase of the form "optionally substituted X" (*e.g.*, optionally substituted alkyl) is intended to be equivalent to "X, wherein X is optionally substituted" (*e.g.*, "alkyl, wherein said alkyl is optionally substituted"). It is not intended to mean that the feature "X" (*e.g.*, alkyl) *per se* is optional.
- [1249] *Part*: As used herein, a "part" or "region" of a polynucleotide is defined as any portion of the polynucleotide that is less than the entire length of the polynucleotide.
- [1250] *Patient*: As used herein, "patient" refers to a subject who can seek or be in need of treatment, requires treatment, is receiving treatment, will receive treatment, or a subject who is under care by a trained professional for a particular disease or condition. In some embodiments, the treatment is needed, required, or received to prevent or decrease the risk of developing acute disease, *i.e.*, it is a prophylactic treatment.
- [1251] *Pharmaceutically acceptable*: The phrase "pharmaceutically acceptable" is employed herein to refer to those compounds, materials, compositions, and/or dosage forms that are, within the scope of sound medical judgment, suitable for use in contact

with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other problem or complication, commensurate with a reasonable benefit/risk ratio.

[1252] *Pharmaceutically acceptable excipients:* The phrase "pharmaceutically acceptable excipient," as used herein, refers any ingredient other than the compounds described herein (for example, a vehicle capable of suspending or dissolving the active compound) and having the properties of being substantially nontoxic and non-inflammatory in a patient. Excipients can include, for example: antiadherents, antioxidants, binders, coatings, compression aids, disintegrants, dyes (colors), emollients, emulsifiers, fillers (diluent), film formers or coatings, flavors, fragrances, glidants (flow enhancers), lubricants, preservatives, printing inks, sorbents, suspending or dispersing agents, sweeteners, and waters of hydration. Exemplary excipients include, but are not limited to: butylated hydroxytoluene (BHT), calcium carbonate, calcium phosphate (dibasic), calcium stearate, croscarmellose, crosslinked polyvinyl pyrrolidone, citric acid, crospovidone, cysteine, ethylcellulose, gelatin, hydroxypropyl cellulose, hydroxypropyl methylcellulose, lactose, magnesium stearate, maltitol, mannitol, methionine, methylcellulose, methyl paraben, microcrystalline cellulose, polyethylene glycol, polyvinyl pyrrolidone, povidone, pregelatinized starch, propyl paraben, retinyl palmitate, shellac, silicon dioxide, sodium carboxymethyl cellulose, sodium citrate, sodium starch glycolate, sorbitol, starch (corn), stearic acid, sucrose, talc, titanium dioxide, vitamin A, vitamin E, vitamin C, and xylitol.

[1253] *Pharmaceutically acceptable salts:* The present disclosure also includes pharmaceutically acceptable salts of the compounds described herein. As used herein, "pharmaceutically acceptable salts" refers to derivatives of the disclosed compounds wherein the parent compound is modified by converting an existing acid or base moiety to its salt form (*e.g.*, by reacting the free base group with a suitable organic acid). Examples of pharmaceutically acceptable salts include, but are not limited to, mineral or organic acid salts of basic residues such as amines; alkali or organic salts of acidic residues such as carboxylic acids; and the like. Representative acid addition salts include acetate, acetic acid, adipate, alginate, ascorbate, aspartate, benzenesulfonate, benzoic sulfonic acid, benzoate, bisulfate, borate, butyrate, camphorate, camphorsulfonate, citrate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, fumarate, glucoheptonate, glycerophosphate, hemisulfate, heptonate, hexanoate, hydrobromide,

hydrochloride, hydroiodide, 2-hydroxy-ethanesulfonate, lactobionate, lactate, laurate, lauryl sulfate, malate, maleate, malonate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, nitrate, oleate, oxalate, palmitate, pamoate, pectinate, persulfate, 3-phenylpropionate, phosphate, picrate, pivalate, propionate, stearate, succinate, sulfate, tartrate, thiocyanate, toluenesulfonate, undecanoate, valerate salts, and the like.

Representative alkali or alkaline earth metal salts include sodium, lithium, potassium, calcium, magnesium, and the like, as well as nontoxic ammonium, quaternary ammonium, and amine cations, including, but not limited to ammonium, tetramethylammonium, tetraethylammonium, methylamine, dimethylamine, trimethylamine, triethylamine, ethylamine, and the like. The pharmaceutically acceptable salts of the present disclosure include the conventional non-toxic salts of the parent compound formed, for example, from non-toxic inorganic or organic acids. The pharmaceutically acceptable salts of the present disclosure can be synthesized from the parent compound that contains a basic or acidic moiety by conventional chemical methods. Generally, such salts can be prepared by reacting the free acid or base forms of these compounds with a stoichiometric amount of the appropriate base or acid in water or in an organic solvent, or in a mixture of the two; generally, nonaqueous media like ether, ethyl acetate, ethanol, isopropanol, or acetonitrile are used. Lists of suitable salts are found in *Remington's Pharmaceutical Sciences*, 17th ed., Mack Publishing Company, Easton, Pa., 1985, p. 1418, *Pharmaceutical Salts: Properties, Selection, and Use*, P.H. Stahl and C.G. Wermuth (eds.), Wiley-VCH, 2008, and Berge et al., *Journal of Pharmaceutical Science*, 66, 1-19 (1977), each of which is incorporated herein by reference in its entirety.

[1254] *Pharmaceutically acceptable solvate*: The term "pharmaceutically acceptable solvate," as used herein, means a compound of the invention wherein molecules of a suitable solvent are incorporated in the crystal lattice. A suitable solvent is physiologically tolerable at the dosage administered. For example, solvates can be prepared by crystallization, recrystallization, or precipitation from a solution that includes organic solvents, water, or a mixture thereof. Examples of suitable solvents are ethanol, water (for example, mono-, di-, and tri-hydrates), *N*-methylpyrrolidinone (NMP), dimethyl sulfoxide (DMSO), *N,N'*-dimethylformamide (DMF), *N,N'*-dimethylacetamide (DMAC), 1,3-dimethyl-2-imidazolidinone (DMEU), 1,3-dimethyl-3,4,5,6-tetrahydro-2-(1H)-pyrimidinone (DMPU), acetonitrile (ACN), propylene glycol, ethyl acetate, benzyl

alcohol, 2-pyrrolidone, benzyl benzoate, and the like. When water is the solvent, the solvate is referred to as a "hydrate."

- [1255] *Pharmacokinetic*: As used herein, "pharmacokinetic" refers to any one or more properties of a molecule or compound as it relates to the determination of the fate of substances administered to a living organism. Pharmacokinetics is divided into several areas including the extent and rate of absorption, distribution, metabolism and excretion. This is commonly referred to as ADME where: (A) Absorption is the process of a substance entering the blood circulation; (D) Distribution is the dispersion or dissemination of substances throughout the fluids and tissues of the body; (M) Metabolism (or Biotransformation) is the irreversible transformation of parent compounds into daughter metabolites; and (E) Excretion (or Elimination) refers to the elimination of the substances from the body. In rare cases, some drugs irreversibly accumulate in body tissue.
- [1256] *Physicochemical*: As used herein, "physicochemical" means of or relating to a physical and/or chemical property.
- [1257] *Polynucleotide*: The term "polynucleotide" as used herein refers to polymers of nucleotides of any length, including ribonucleotides, deoxyribonucleotides, analogs thereof, or mixtures thereof. This term refers to the primary structure of the molecule. Thus, the term includes triple-, double- and single-stranded deoxyribonucleic acid ("DNA"), as well as triple-, double- and single-stranded ribonucleic acid ("RNA"). It also includes modified, for example by alkylation, and/or by capping, and unmodified forms of the polynucleotide. More particularly, the term "polynucleotide" includes polydeoxyribonucleotides (containing 2-deoxy-D-ribose), polyribonucleotides (containing D-ribose), including tRNA, rRNA, hRNA, siRNA and mRNA, whether spliced or unspliced, any other type of polynucleotide which is an N- or C-glycoside of a purine or pyrimidine base, and other polymers containing non-nucleotidic backbones, for example, polyamide (*e.g.*, peptide nucleic acids "PNAs") and polymorpholino polymers, and other synthetic sequence-specific nucleic acid polymers providing that the polymers contain nucleobases in a configuration which allows for base pairing and base stacking, such as is found in DNA and RNA. In particular aspects, the polynucleotide comprises an mRNA. In other aspect, the mRNA is a synthetic mRNA. In some aspects, the synthetic mRNA comprises at least one unnatural nucleobase. In some aspects, all nucleobases of a certain class have been replaced with unnatural nucleobases (*e.g.*, all uridines in a

polynucleotide disclosed herein can be replaced with an unnatural nucleobase, *e.g.*, 5-methoxyuridine). In some aspects, the polynucleotide (*e.g.*, a synthetic RNA or a synthetic DNA) comprises only natural nucleobases, *i.e.*, A (adenosine), G (guanosine), C (cytidine), and T (thymidine) in the case of a synthetic DNA, or A, C, G, and U (uridine) in the case of a synthetic RNA.

[1258] The skilled artisan will appreciate that the T bases in the codon maps disclosed herein are present in DNA, whereas the T bases would be replaced by U bases in corresponding RNAs. For example, a codon-nucleotide sequence disclosed herein in DNA form, *e.g.*, a vector or an in-vitro translation (IVT) template, would have its T bases transcribed as U based in its corresponding transcribed mRNA. In this respect, both codon-optimized DNA sequences (comprising T) and their corresponding mRNA sequences (comprising U) are considered codon-optimized nucleotide sequence of the present invention. A skilled artisan would also understand that equivalent codon-maps can be generated by replaced one or more bases with non-natural bases. Thus, *e.g.*, a TTC codon (DNA map) would correspond to a UUC codon (RNA map), which in turn would correspond to a ΨΨC codon (RNA map in which U has been replaced with pseudouridine).

[1259] Standard A-T and G-C base pairs form under conditions which allow the formation of hydrogen bonds between the N3-H and C4-oxy of thymidine and the N1 and C6-NH₂, respectively, of adenosine and between the C2-oxy, N3 and C4-NH₂, of cytidine and the C2-NH₂, N'—H and C6-oxy, respectively, of guanosine. Thus, for example, guanosine (2-amino-6-oxy-9-β-D-ribofuranosyl-purine) can be modified to form isoguanosine (2-oxy-6-amino-9-β-D-ribofuranosyl-purine). Such modification results in a nucleoside base which will no longer effectively form a standard base pair with cytosine. However, modification of cytosine (1-β-D-ribofuranosyl-2-oxy-4-amino-pyrimidine) to form isocytosine (1-β-D-ribofuranosyl-2-amino-4-oxy-pyrimidine-) results in a modified nucleotide which will not effectively base pair with guanosine but will form a base pair with isoguanosine (U.S. Pat. No. 5,681,702 to Collins et al.). Isocytosine is available from Sigma Chemical Co. (St. Louis, Mo.); isocytidine can be prepared by the method described by Switzer et al. (1993) Biochemistry 32:10489-10496 and references cited therein; 2'-deoxy-5-methyl-isocytidine can be prepared by the method of Tor et al., 1993, J. Am. Chem. Soc. 115:4461-4467 and references cited therein; and isoguanine nucleotides can be prepared using the method described by Switzer et al., 1993, *supra*,

and Mantsch et al., 1993, Biochem. 14:5593-5601, or by the method described in U.S. Pat. No. 5,780,610 to Collins et al. Other non-natural base pairs can be synthesized by the method described in Piccirilli et al., 1990, Nature 343:33-37, for the synthesis of 2,6-diaminopyrimidine and its complement (1-methylpyrazolo-[4,3]pyrimidine-5,7-(4H,6H)-dione. Other such modified nucleotide units which form unique base pairs are known, such as those described in Leach et al. (1992) J. Am. Chem. Soc. 114:3675-3683 and Switzer et al., supra.

- [1260] *Polypeptide*: The terms "polypeptide," "peptide," and "protein" are used interchangeably herein to refer to polymers of amino acids of any length. The polymer can comprise modified amino acids. The terms also encompass an amino acid polymer that has been modified naturally or by intervention; for example, disulfide bond formation, glycosylation, lipidation, acetylation, phosphorylation, or any other manipulation or modification, such as conjugation with a labeling component. Also included within the definition are, for example, polypeptides containing one or more analogs of an amino acid (including, for example, unnatural amino acids such as homocysteine, ornithine, p-acetylphenylalanine, D-amino acids, and creatine), as well as other modifications known in the art.
- [1261] The term, as used herein, refers to proteins, polypeptides, and peptides of any size, structure, or function. Polypeptides include encoded polynucleotide products, naturally occurring polypeptides, synthetic polypeptides, homologs, orthologs, paralogs, fragments and other equivalents, variants, and analogs of the foregoing. A polypeptide can be a monomer or can be a multi-molecular complex such as a dimer, trimer or tetramer. They can also comprise single chain or multichain polypeptides. Most commonly disulfide linkages are found in multichain polypeptides. The term polypeptide can also apply to amino acid polymers in which one or more amino acid residues are an artificial chemical analogue of a corresponding naturally occurring amino acid. In some embodiments, a "peptide" can be less than or equal to 50 amino acids long, e.g., about 5, 10, 15, 20, 25, 30, 35, 40, 45, or 50 amino acids long.
- [1262] *Polypeptide variant*: As used herein, the term "polypeptide variant" refers to molecules that differ in their amino acid sequence from a native or reference sequence. The amino acid sequence variants can possess substitutions, deletions, and/or insertions at certain positions within the amino acid sequence, as compared to a native or reference sequence. Ordinarily, variants will possess at least about 50% identity, at least about 60%

identity, at least about 70% identity, at least about 80% identity, at least about 90% identity, at least about 95% identity, at least about 99% identity to a native or reference sequence. In some embodiments, they will be at least about 80%, or at least about 90% identical to a native or reference sequence.

[1263] *Polypeptide per unit drug (PUD)*: As used herein, a PUD or product per unit drug, is defined as a subdivided portion of total daily dose, usually 1 mg, pg, kg, etc., of a product (such as a polypeptide) as measured in body fluid or tissue, usually defined in concentration such as pmol/mL, mmol/mL, etc. divided by the measure in the body fluid.

[1264] *Preventing*: As used herein, the term "preventing" refers to partially or completely delaying onset of an infection, disease, disorder and/or condition; partially or completely delaying onset of one or more signs and symptoms, features, or clinical manifestations of a particular infection, disease, disorder, and/or condition; partially or completely delaying onset of one or more signs and symptoms, features, or manifestations of a particular infection, disease, disorder, and/or condition; partially or completely delaying progression from an infection, a particular disease, disorder and/or condition; and/or decreasing the risk of developing pathology associated with the infection, the disease, disorder, and/or condition.

[1265] *Proliferate*: As used herein, the term "proliferate" means to grow, expand or increase or cause to grow, expand or increase rapidly. "Proliferative" means having the ability to proliferate. "Anti-proliferative" means having properties counter to or inapposite to proliferative properties.

[1266] *Prophylactic*: As used herein, "prophylactic" refers to a therapeutic or course of action used to prevent the spread of disease.

[1267] *Prophylaxis*: As used herein, a "prophylaxis" refers to a measure taken to maintain health and prevent the spread of disease. An "immune prophylaxis" refers to a measure to produce active or passive immunity to prevent the spread of disease.

[1268] *Protein cleavage site*: As used herein, "protein cleavage site" refers to a site where controlled cleavage of the amino acid chain can be accomplished by chemical, enzymatic or photochemical means.

[1269] *Protein cleavage signal*: As used herein "protein cleavage signal" refers to at least one amino acid that flags or marks a polypeptide for cleavage.

- [1270] *Protein of interest*: As used herein, the terms "proteins of interest" or "desired proteins" include those provided herein and fragments, mutants, variants, and alterations thereof.
- [1271] *Proximal*: As used herein, the term "proximal" means situated nearer to the center or to a point or region of interest.
- [1272] *Pseudouridine*: As used herein, pseudouridine (ψ) refers to the C-glycoside isomer of the nucleoside uridine. A "pseudouridine analog" is any modification, variant, isoform or derivative of pseudouridine. For example, pseudouridine analogs include but are not limited to 1-carboxymethyl-pseudouridine, 1-propynyl-pseudouridine, 1-taurinomethyl-pseudouridine, 1-taurinomethyl-4-thio-pseudouridine, 1-methylpseudouridine ($m^1\psi$), 1-methyl-4-thio-pseudouridine ($m^1s^4\psi$), 4-thio-1-methyl-pseudouridine, 3-methyl-pseudouridine ($m^3\psi$), 2-thio-1-methyl-pseudouridine, 1-methyl-1-deaza-pseudouridine, 2-thio-1-methyl-1-deaza-pseudouridine, dihydropseudouridine, 2-thio-dihydropseudouridine, 2-methoxyuridine, 2-methoxy-4-thio-uridine, 4-methoxy-pseudouridine, 4-methoxy-2-thio-pseudouridine, N1-methyl-pseudouridine, 1-methyl-3-(3-amino-3-carboxypropyl)pseudouridine ($acp^3\psi$), and 2'-O-methyl-pseudouridine (ψm).
- [1273] *Purified*: As used herein, "purify," "purified," "purification" means to make substantially pure or clear from unwanted components, material defilement, admixture or imperfection.
- [1274] *Reference Nucleic Acid Sequence*: The term "reference nucleic acid sequence" or "reference nucleic acid" or "reference nucleotide sequence" or "reference sequence" refers to a starting nucleic acid sequence (e.g., a RNA, e.g., an mRNA sequence) that can be sequence optimized. In some embodiments, the reference nucleic acid sequence is a wild type nucleic acid sequence, a fragment or a variant thereof. In some embodiments, the reference nucleic acid sequence is a previously sequence optimized nucleic acid sequence.
- [1275] *Salts*: In some aspects, the pharmaceutical composition for delivery disclosed herein comprises salts of some of their lipid constituents. The term "salt" includes any anionic and cationic complex. Non-limiting examples of anions include inorganic and organic anions, e.g., fluoride, chloride, bromide, iodide, oxalate (e.g., hemioxalate), phosphate, phosphonate, hydrogen phosphate, dihydrogen phosphate, oxide, carbonate, bicarbonate, nitrate, nitrite, nitride, bisulfite, sulfide, sulfite, bisulfate, sulfate, thiosulfate, hydrogen sulfate, borate, formate, acetate, benzoate, citrate, tartrate, lactate, acrylate,

polyacrylate, fumarate, maleate, itaconate, glycolate, gluconate, malate, mandelate, tiglate, ascorbate, salicylate, polymethacrylate, perchlorate, chlorate, chlorite, hypochlorite, bromate, hypobromite, iodate, an alkylsulfonate, an arylsulfonate, arsenate, arsenite, chromate, dichromate, cyanide, cyanate, thiocyanate, hydroxide, peroxide, permanganate, and mixtures thereof.

- [1276] *Sample:* As used herein, the term "sample" or "biological sample" refers to a subset of its tissues, cells or component parts (*e.g.*, body fluids, including but not limited to blood, mucus, lymphatic fluid, synovial fluid, cerebrospinal fluid, saliva, amniotic fluid, amniotic cord blood, urine, vaginal fluid and semen). A sample further can include a homogenate, lysate or extract prepared from a whole organism or a subset of its tissues, cells or component parts, or a fraction or portion thereof, including but not limited to, for example, plasma, serum, spinal fluid, lymph fluid, the external sections of the skin, respiratory, intestinal, and genitourinary tracts, tears, saliva, milk, blood cells, tumors, or organs. A sample further refers to a medium, such as a nutrient broth or gel, which can contain cellular components, such as proteins or nucleic acid molecule.
- [1277] *Signal Sequence:* As used herein, the phrases "signal sequence," "signal peptide," and "transit peptide" are used interchangeably and refer to a sequence that can direct the transport or localization of a protein to a certain organelle, cell compartment, or extracellular export. The term encompasses both the signal sequence polypeptide and the nucleic acid sequence encoding the signal sequence. Thus, references to a signal sequence in the context of a nucleic acid refer in fact to the nucleic acid sequence encoding the signal sequence polypeptide.
- [1278] *Signal transduction pathway:* A "signal transduction pathway" refers to the biochemical relationship between a variety of signal transduction molecules that play a role in the transmission of a signal from one portion of a cell to another portion of a cell. As used herein, the phrase "cell surface receptor" includes, for example, molecules and complexes of molecules capable of receiving a signal and the transmission of such a signal across the plasma membrane of a cell.
- [1279] *Similarity:* As used herein, the term "similarity" refers to the overall relatedness between polymeric molecules, *e.g.* between polynucleotide molecules (*e.g.* DNA molecules and/or RNA molecules) and/or between polypeptide molecules. Calculation of percent similarity of polymeric molecules to one another can be performed in the same

manner as a calculation of percent identity, except that calculation of percent similarity takes into account conservative substitutions as is understood in the art.

- [1280] *Single unit dose*: As used herein, a "single unit dose" is a dose of any therapeutic administered in one dose/at one time/single route/single point of contact, *i.e.*, single administration event.
- [1281] *Split dose*: As used herein, a "split dose" is the division of single unit dose or total daily dose into two or more doses.
- [1282] *Specific delivery*: As used herein, the term "specific delivery," "specifically deliver," or "specifically delivering" means delivery of more (*e.g.*, at least 1.5 fold more, at least 2-fold more, at least 3-fold more, at least 4-fold more, at least 5-fold more, at least 6-fold more, at least 7-fold more, at least 8-fold more, at least 9-fold more, at least 10-fold more) of a polynucleotide by a nanoparticle to a target tissue of interest (*e.g.*, mammalian liver) compared to an off-target tissue. The level of delivery of a nanoparticle to a particular tissue can be measured by comparing the amount of protein produced in a tissue to the weight of said tissue, comparing the amount of polynucleotide in a tissue to the weight of said tissue, comparing the amount of protein produced in a tissue to the amount of total protein in said tissue, or comparing the amount of polynucleotide in a tissue to the amount of total polynucleotide in said tissue. For example, for renovascular targeting, a polynucleotide is specifically provided to a mammalian kidney as compared to the liver and spleen if 1.5, 2-fold, 3-fold, 5-fold, 10-fold, 15 fold, or 20 fold more polynucleotide per 1 g of tissue is delivered to a kidney compared to that delivered to the liver or spleen following systemic administration of the polynucleotide. It will be understood that the ability of a nanoparticle to specifically deliver to a target tissue need not be determined in a subject being treated, it can be determined in a surrogate such as an animal model (*e.g.*, a rat model).
- [1283] *Stable*: As used herein "stable" refers to a compound that is sufficiently robust to survive isolation to a useful degree of purity from a reaction mixture, and in some cases capable of formulation into an efficacious therapeutic agent.
- [1284] *Stabilized*: As used herein, the term "stabilize," "stabilized," "stabilized region" means to make or become stable.
- [1285] *Stereoisomer*: As used herein, the term "stereoisomer" refers to all possible different isomeric as well as conformational forms that a compound can possess (*e.g.*, a compound of any formula described herein), in particular all possible stereochemically

and conformationally isomeric forms, all diastereomers, enantiomers and/or conformers of the basic molecular structure. Some compounds of the present invention can exist in different tautomeric forms, all of the latter being included within the scope of the present invention.

- [1286] *Subject*: By "subject" or "individual" or "animal" or "patient" or "mammal," is meant any subject, particularly a mammalian subject, for whom diagnosis, prognosis, or therapy is desired. Mammalian subjects include, but are not limited to, humans, domestic animals, farm animals, zoo animals, sport animals, pet animals such as dogs, cats, guinea pigs, rabbits, rats, mice, horses, cattle, cows; primates such as apes, monkeys, orangutans, and chimpanzees; canids such as dogs and wolves; felids such as cats, lions, and tigers; equids such as horses, donkeys, and zebras; bears, food animals such as cows, pigs, and sheep; ungulates such as deer and giraffes; rodents such as mice, rats, hamsters and guinea pigs; and so on. In certain embodiments, the mammal is a human subject. In other embodiments, a subject is a human patient. In a particular embodiment, a subject is a human patient in need of treatment.
- [1287] *Substantially*: As used herein, the term "substantially" refers to the qualitative condition of exhibiting total or near-total extent or degree of a characteristic or property of interest. One of ordinary skill in the biological arts will understand that biological and chemical characteristics rarely, if ever, go to completion and/or proceed to completeness or achieve or avoid an absolute result. The term "substantially" is therefore used herein to capture the potential lack of completeness inherent in many biological and chemical characteristics.
- [1288] *Substantially equal*: As used herein as it relates to time differences between doses, the term means plus/minus 2%.
- [1289] *Substantially simultaneous*: As used herein and as it relates to plurality of doses, the term means within 2 seconds.
- [1290] *Suffering from*: An individual who is "suffering from" a disease, disorder, and/or condition has been diagnosed with or displays one or more signs and symptoms of the disease, disorder, and/or condition.
- [1291] *Susceptible to*: An individual who is "susceptible to" a disease, disorder, and/or condition has not been diagnosed with and/or can not exhibit signs and symptoms of the disease, disorder, and/or condition but harbors a propensity to develop a disease or its signs and symptoms. In some embodiments, an individual who is susceptible to a disease,

disorder, and/or condition (for example, VLCADD) can be characterized by one or more of the following: (1) a genetic mutation associated with development of the disease, disorder, and/or condition; (2) a genetic polymorphism associated with development of the disease, disorder, and/or condition; (3) increased and/or decreased expression and/or activity of a protein and/or nucleic acid associated with the disease, disorder, and/or condition; (4) habits and/or lifestyles associated with development of the disease, disorder, and/or condition; (5) a family history of the disease, disorder, and/or condition; and (6) exposure to and/or infection with a microbe associated with development of the disease, disorder, and/or condition. In some embodiments, an individual who is susceptible to a disease, disorder, and/or condition will develop the disease, disorder, and/or condition. In some embodiments, an individual who is susceptible to a disease, disorder, and/or condition will not develop the disease, disorder, and/or condition.

[1292] *Sustained release*: As used herein, the term "sustained release" refers to a pharmaceutical composition or compound release profile that conforms to a release rate over a specific period of time.

[1293] *Synthetic*: The term "synthetic" means produced, prepared, and/or manufactured by the hand of man. Synthesis of polynucleotides or other molecules of the present invention can be chemical or enzymatic.

[1294] *Targeted Cells*: As used herein, "targeted cells" refers to any one or more cells of interest. The cells can be found *in vitro*, *in vivo*, *in situ* or in the tissue or organ of an organism. The organism can be an animal, for example a mammal, a human, a subject or a patient.

[1295] *Target tissue*: As used herein "target tissue" refers to any one or more tissue types of interest in which the delivery of a polynucleotide would result in a desired biological and/or pharmacological effect. Examples of target tissues of interest include specific tissues, organs, and systems or groups thereof. In particular applications, a target tissue can be a liver, a kidney, a lung, a spleen, or vascular endothelium in vessels (e.g., intra-coronary or intra-femoral). An "off-target tissue" refers to any one or more tissue types in which the expression of the encoded protein does not result in a desired biological and/or pharmacological effect.

[1296] The presence of a therapeutic agent in an off-target issue can be the result of: (i) leakage of a polynucleotide from the administration site to peripheral tissue or distant off-target tissue via diffusion or through the bloodstream (e.g., a polynucleotide intended to

express a polypeptide in a certain tissue would reach the off-target tissue and the polypeptide would be expressed in the off-target tissue); or (ii) leakage of a polypeptide after administration of a polynucleotide encoding such polypeptide to peripheral tissue or distant off-target tissue via diffusion or through the bloodstream (e.g., a polynucleotide would express a polypeptide in the target tissue, and the polypeptide would diffuse to peripheral tissue).

[1297] *Targeting sequence*: As used herein, the phrase "targeting sequence" refers to a sequence that can direct the transport or localization of a protein or polypeptide.

[1298] *Terminus*: As used herein the terms "termini" or "terminus," when referring to polypeptides, refers to an extremity of a peptide or polypeptide. Such extremity is not limited only to the first or final site of the peptide or polypeptide but can include additional amino acids in the terminal regions. The polypeptide based molecules of the invention can be characterized as having both an N-terminus (terminated by an amino acid with a free amino group (NH₂)) and a C-terminus (terminated by an amino acid with a free carboxyl group (COOH)). Proteins of the invention are in some cases made up of multiple polypeptide chains brought together by disulfide bonds or by non-covalent forces (multimers, oligomers). These sorts of proteins will have multiple N- and C-termini. Alternatively, the termini of the polypeptides can be modified such that they begin or end, as the case can be, with a non-polypeptide based moiety such as an organic conjugate.

[1299] *Therapeutic Agent*: The term "therapeutic agent" refers to an agent that, when administered to a subject, has a therapeutic, diagnostic, and/or prophylactic effect and/or elicits a desired biological and/or pharmacological effect. For example, in some embodiments, an mRNA encoding an ACADVL polypeptide can be a therapeutic agent.

[1300] *Therapeutically effective amount*: As used herein, the term "therapeutically effective amount" means an amount of an agent to be delivered (e.g., nucleic acid, drug, therapeutic agent, diagnostic agent, prophylactic agent, etc.) that is sufficient, when administered to a subject suffering from or susceptible to an infection, disease, disorder, and/or condition, to treat, improve signs and symptoms of, diagnose, prevent, and/or delay the onset of the infection, disease, disorder, and/or condition.

[1301] *Therapeutically effective outcome*: As used herein, the term "therapeutically effective outcome" means an outcome that is sufficient in a subject suffering from or susceptible to an infection, disease, disorder, and/or condition, to treat, improve signs and

symptoms of, diagnose, prevent, and/or delay the onset of the infection, disease, disorder, and/or condition.

- [1302] *Total daily dose:* As used herein, a "total daily dose" is an amount given or prescribed in 24 hr. period. The total daily dose can be administered as a single unit dose or a split dose.
- [1303] *Transcription factor:* As used herein, the term "transcription factor" refers to a DNA-binding protein that regulates transcription of DNA into RNA, for example, by activation or repression of transcription. Some transcription factors effect regulation of transcription alone, while others act in concert with other proteins. Some transcription factor can both activate and repress transcription under certain conditions. In general, transcription factors bind a specific target sequence or sequences highly similar to a specific consensus sequence in a regulatory region of a target gene. Transcription factors can regulate transcription of a target gene alone or in a complex with other molecules.
- [1304] *Transcription:* As used herein, the term "transcription" refers to methods to produce mRNA (e.g., an mRNA sequence or template) from DNA (e.g., a DNA template or sequence).
- [1305] *Transfection:* As used herein, "transfection" refers to the introduction of a polynucleotide (e.g., exogenous nucleic acids) into a cell wherein a polypeptide encoded by the polynucleotide is expressed (e.g., mRNA) or the polypeptide modulates a cellular function (e.g., siRNA, miRNA). As used herein, "expression" of a nucleic acid sequence refers to translation of a polynucleotide (e.g., an mRNA) into a polypeptide or protein and/or post-translational modification of a polypeptide or protein. Methods of transfection include, but are not limited to, chemical methods, physical treatments and cationic lipids or mixtures.
- [1306] *Treating, treatment, therapy:* As used herein, the term "treating" or "treatment" or "therapy" refers to partially or completely alleviating, ameliorating, improving, relieving, delaying onset of, inhibiting progression of, reducing severity of, and/or reducing incidence of one or more signs and symptoms or features of a disease, e.g., VLCADD. For example, "treating" VLCADD can refer to diminishing signs and symptoms associate with the disease, prolong the lifespan (increase the survival rate) of patients, reducing the severity of the disease, preventing or delaying the onset of the disease, etc. Treatment can be administered to a subject who does not exhibit signs of a disease, disorder, and/or condition and/or to a subject who exhibits only early signs of a disease, disorder, and/or

condition for the purpose of decreasing the risk of developing pathology associated with the disease, disorder, and/or condition.

[1307] *Unmodified*: As used herein, "unmodified" refers to any substance, compound or molecule prior to being changed in some way. Unmodified can, but does not always, refer to the wild type or native form of a biomolecule. Molecules can undergo a series of modifications whereby each modified molecule can serve as the "unmodified" starting molecule for a subsequent modification.

[1308] *Uracil*: Uracil is one of the four nucleobases in the nucleic acid of RNA, and it is represented by the letter U. Uracil can be attached to a ribose ring, or more specifically, a ribofuranose via a β -N₁-glycosidic bond to yield the nucleoside uridine. The nucleoside uridine is also commonly abbreviated according to the one letter code of its nucleobase, i.e., U. Thus, in the context of the present disclosure, when a monomer in a polynucleotide sequence is U, such U is designated interchangeably as a "uracil" or a "uridine."

[1309] *Uridine Content*: The terms "uridine content" or "uracil content" are interchangeable and refer to the amount of uracil or uridine present in a certain nucleic acid sequence. Uridine content or uracil content can be expressed as an absolute value (total number of uridine or uracil in the sequence) or relative (uridine or uracil percentage respect to the total number of nucleobases in the nucleic acid sequence).

[1310] *Uridine-Modified Sequence*: The terms "uridine-modified sequence" refers to a sequence optimized nucleic acid (e.g., a synthetic mRNA sequence) with a different overall or local uridine content (higher or lower uridine content) or with different uridine patterns (e.g., gradient distribution or clustering) with respect to the uridine content and/or uridine patterns of a candidate nucleic acid sequence. In the content of the present disclosure, the terms "uridine-modified sequence" and "uracil-modified sequence" are considered equivalent and interchangeable.

[1311] A "high uridine codon" is defined as a codon comprising two or three uridines, a "low uridine codon" is defined as a codon comprising one uridine, and a "no uridine codon" is a codon without any uridines. In some embodiments, a uridine-modified sequence comprises substitutions of high uridine codons with low uridine codons, substitutions of high uridine codons with no uridine codons, substitutions of low uridine codons with high uridine codons, substitutions of low uridine codons with no uridine codons, substitution of no uridine codons with low uridine codons, substitutions of no

uridine codons with high uridine codons, and combinations thereof. In some embodiments, a high uridine codon can be replaced with another high uridine codon. In some embodiments, a low uridine codon can be replaced with another low uridine codon. In some embodiments, a no uridine codon can be replaced with another no uridine codon. A uridine-modified sequence can be uridine enriched or uridine rarefied.

[1312] *Uridine Enriched:* As used herein, the terms "uridine enriched" and grammatical variants refer to the increase in uridine content (expressed in absolute value or as a percentage value) in a sequence optimized nucleic acid (e.g., a synthetic mRNA sequence) with respect to the uridine content of the corresponding candidate nucleic acid sequence. Uridine enrichment can be implemented by substituting codons in the candidate nucleic acid sequence with synonymous codons containing less uridine nucleobases. Uridine enrichment can be global (i.e., relative to the entire length of a candidate nucleic acid sequence) or local (i.e., relative to a subsequence or region of a candidate nucleic acid sequence).

[1313] *Uridine Rarefied:* As used herein, the terms "uridine rarefied" and grammatical variants refer to a decrease in uridine content (expressed in absolute value or as a percentage value) in a sequence optimized nucleic acid (e.g., a synthetic mRNA sequence) with respect to the uridine content of the corresponding candidate nucleic acid sequence. Uridine rarefaction can be implemented by substituting codons in the candidate nucleic acid sequence with synonymous codons containing less uridine nucleobases. Uridine rarefaction can be global (i.e., relative to the entire length of a candidate nucleic acid sequence) or local (i.e., relative to a subsequence or region of a candidate nucleic acid sequence).

[1314] *Variant:* The term variant as used in present disclosure refers to both natural variants (e.g., polymorphisms, isoforms, etc.) and artificial variants in which at least one amino acid residue in a native or starting sequence (e.g., a wild type sequence) has been removed and a different amino acid inserted in its place at the same position. These variants can be described as "substitutional variants." The substitutions can be single, where only one amino acid in the molecule has been substituted, or they can be multiple, where two or more amino acids have been substituted in the same molecule. If amino acids are inserted or deleted, the resulting variant would be an "insertional variant" or a "deletional variant" respectively.

30. Embodiments

- [1315] Throughout this section, the term embodiment is abbreviated as 'E' followed by an ordinal. For example, E1 is equivalent to Embodiment 1.
- [1316] E1. A polynucleotide comprising an open reading frame (ORF) encoding an acyl-CoA dehydrogenase, very long chain (ACADVL) polypeptide, wherein the uracil or thymine content of the ORF relative to the theoretical minimum uracil or thymine content of a nucleotide sequence encoding the ACADVL polypeptide (%U_{TM} or %T_{TM}) is between 100% and about 150%.
- [1317] E2. The polynucleotide of E1, wherein the %U_{TM} or %T_{TM} is between about 105% and about 140%, about 110% and about 140%, 115% and about 140%, about 105% and about 130%, about 110% and about 130%, about 115% and about 135%, about 105% and about 135%, about 110% and about 135%, or about 115% and about 130%.
- [1318] E3. The polynucleotide of E2, wherein the %U_{TM} or %T_{TM} is between (i) 110%, 111%, 112%, 113%, 114%, 115%, 116%, 117%, or 118% and (ii) 128%, 129%, 130%, 131%, 132%, 133%, 134%, 135%, 136%, 137%, 138%, 139%, or 140%.
- [1319] E4. The polynucleotide of any one of E1 to E3, wherein the uracil or thymine content in the ORF is less than the uracil or thymine content in the corresponding wild-type ORF (%U_{WT} or %T_{WT}).
- [1320] E5. The polynucleotide of E4, wherein the uracil or thymine content in the ORF is less than about 95%, less than about 90%, less than about 85%, less than 80%, less than 79%, less than 78%, less than 77%, less than 76%, less than 75%, or less than 74%.
- [1321] E6. The polynucleotide of E4, wherein the %U_{WT} or %T_{WT} is between 69% and 75% of the %U_{WT} or %T_{WT}.
- [1322] E7. The polynucleotide of any one of E1 to E6, wherein the uracil or thymine content in the ORF relative to the total nucleotide content in the ORF (%U_{TL} or %T_{TL}) is less than about 50%, less than about 40%, less than about 30%, or less than about 20%.
- [1323] E8. The polynucleotide of E7, wherein the %U_{TL} or %T_{TL} is less than about 16%.
- [1324] E9. The polynucleotide of any one of E1 to E8, wherein the %U_{TL} or %T_{TL} is between about 14% and about 16%.
- [1325] E10. The polynucleotide of any one of E1 to E9, wherein the guanine content of the ORF with respect to the theoretical maximum guanine content of a nucleotide sequence encoding the ACADVL polypeptide (%G_{TMX}) is at least 70%, at least 75%, at

least about 80%, at least about 85%, at least about 90%, at least about 95%, or about 100%.

- [1326] E11. The polynucleotide of E10, wherein the %G_{TMX} is between about 70% and about 80%, between about 71% and about 79%, between about 71% and about 77%, or between about 72% and about 76%.
- [1327] E12. The polynucleotide of any one of E1 to E11, wherein the cytosine content of the ORF relative to the theoretical maximum cytosine content of a nucleotide sequence encoding the ACADVL polypeptide (%C_{TMX}) is at least 58%, at least 59%, at least 60%, at least 65%, at least 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, or about 100%.
- [1328] E13. The polynucleotide of E12, wherein the %C_{TMX} is between about 60% and about 80%, between about 62% and about 77%, between about 63% and about 78%, or between about 68% and about 75%.
- [1329] E14. The polynucleotide of any one of E1 to E13, wherein the guanine and cytosine content (G/C) of the ORF relative to the theoretical maximum G/C content in a nucleotide sequence encoding the ACADVL polypeptide (%G/C_{TMX}) is at least 82%, at least 83%, at least 84%, at least 85%, at least 90%, at least about 95%, or about 100%.
- [1330] E15. The polynucleotide of any one of E1 to E13, wherein the %G/C_{TMX} is between about 80% and about 100%, between about 85% and about 99%, between about 90% and about 96%, or between about 92% and about 95%.
- [1331] E16. The polynucleotide of any one of E1 to E15, wherein the G/C content in the ORF relative to the G/C content in the corresponding wild-type ORF (%G/C_{WT}) is at least 102%, at least 103%, at least 104%, at least 105%, at least 106%, at least 107%, at least about 110%, or at least about 112%.
- [1332] E17. The polynucleotide of any one of E1 to E15, wherein the average G/C content in the 3rd codon position in the ORF is at least 20%, at least 21%, at least 22%, at least 23%, at least 24%, or at least 25% higher than the average G/C content in the 3rd codon position in the corresponding wild-type ORF.
- [1333] E18. The polynucleotide of any one of E1 to E17, wherein the ORF further comprises at least one low-frequency codon.
- [1334] E19. The polynucleotide of any one of E1 to E18,
(i) wherein the ORF is at least 83%, at least 84%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least

93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identical to ACADV-L-CO4, ACADV-L-CO10, or ACADV-L-CO17,

(ii) wherein the ORF is at least 82%, at least 83%, at least 84%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identical to ACADV-L-CO1, ACADV-L-CO7, ACADV-L-CO15, ACADV-L-CO16, ACADV-L-CO18, ACADV-L-CO22, or ACADV-L-CO23,

(iii) wherein the ORF is at least 81%, at least 82%, at least 83%, at least 84%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identical to ACADV-L-CO2, ACADV-L-CO3, ACADV-L-CO5, ACADV-L-CO6, ACADV-L-CO8, ACADV-L-CO9, ACADV-L-CO111, ACADV-L-CO12, ACADV-L-CO13, ACADV-L-CO14, ACADV-L-CO19, ACADV-L-CO20, ACADV-L-CO21, ACADV-L-CO24, or ACADV-L-CO25.

[1335] E20. A polynucleotide comprising an ORF,

(i) wherein the ORF is at least 83%, at least 84%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identical to ACADV-L-CO4, ACADV-L-CO10, or ACADV-L-CO17,

(ii) wherein the ORF is at least 82%, at least 83%, at least 84%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identical to ACADV-L-CO1, ACADV-L-CO7, ACADV-L-CO15, ACADV-L-CO16, ACADV-L-CO18, ACADV-L-CO22, or ACADV-L-CO23,

(iii) wherein the ORF is at least 81%, at least 82%, at least 83%, at least 84%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identical to ACADV-L-CO2, ACADV-L-CO3, ACADV-L-CO5, ACADV-L-CO6, ACADV-L-CO8, ACADV-L-CO9, ACADV-L-CO111, ACADV-L-CO12, ACADV-L-CO13, ACADV-L-CO14, ACADV-L-CO19, ACADV-L-CO20, ACADV-L-CO21, ACADV-L-CO24, or ACADV-L-CO25.

[1336] E21A. The polynucleotide of any one of E1 to E20, wherein the ORF has at least 81%, at least 82%, at least 83%, at least 84%, at least 85%, at least 86%, at least 87%, at

least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to a sequence selected from the group consisting of SEQ ID NOs: 7-31.

- [1337] E21B. The polynucleotide of any one of E1 to E20, wherein the ACADVL polypeptide comprises an amino acid sequence at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99%, or about 100% identical to the polypeptide sequence of wild type ACADVL, isoform 1 (SEQ ID NO: 1), and wherein the ACADVL polypeptide has acyl-CoA dehydrogenase, very long chain activity.
- [1338] E22. The polynucleotide of any one of E1 to E20, wherein the ACADVL polypeptide comprises an amino acid sequence at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99%, or about 100% identical to the polypeptide sequence of wild type ACADVL, isoform 2 (SEQ ID NO: 3), and wherein the ACADVL polypeptide has acyl-CoA dehydrogenase, very long chain activity.
- [1339] E23. The polynucleotide of any one of E1 to E20, wherein the ACADVL polypeptide comprises an amino acid sequence at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99%, or about 100% identical to the polypeptide sequence of wild type ACADVL, isoform 3 (SEQ ID NO: 5), and wherein the ACADVL polypeptide has acyl-CoA dehydrogenase, very long chain activity.
- [1340] E24. The polynucleotide of E23, wherein the ACADVL polypeptide is a variant, derivative, or mutant having an acyl-CoA dehydrogenase, very long chain activity.
- [1341] E25. The polynucleotide of any one of E1 to E24, wherein the polynucleotide sequence further comprises a nucleotide sequence encoding a transit peptide.
- [1342] E26. The polynucleotide of any one of E1 to E25, wherein the polynucleotide is single stranded.
- [1343] E27. The polynucleotide of any one of E1 to E25, wherein the polynucleotide is double stranded.
- [1344] E28. The polynucleotide of any one of E1 to E27, wherein the polynucleotide is DNA.
- [1345] E29. The polynucleotide of any one of E1 to E27, wherein the polynucleotide is RNA.
- [1346] E30. The polynucleotide of E29, wherein the polynucleotide is mRNA.


- [1347] E31. The polynucleotide of any one of E1 to E30, wherein the polynucleotide comprises at least one chemically modified nucleobase, sugar, backbone, or any combination thereof.
- [1348] E32. The polynucleotide of E31, wherein the at least one chemically modified nucleobase is selected from the group consisting of pseudouracil (ψ), N1-methylpseudouracil (m1 ψ), 2-thiouracil (s2U), 4'-thiouracil, 5-methylcytosine, 5-methyluracil, and any combinations thereof.
- [1349] E33. The polynucleotide of E32, wherein the at least one chemically modified nucleobase is 5-methoxyuracil.
- [1350] E34. The polynucleotide of E33, wherein at least about 25%, at least about 30%, at least about 40%, at least about 50%, at least about 60%, at least about 70%, at least about 80%, at least about 90%, at least about 95%, at least about 99%, or 100% of the uracils are 5-methoxyuracils.
- [1351] E35. The polynucleotide of any one of E1 to E34, wherein the polynucleotide further comprises a miRNA binding site.
- [1352] E36. The polynucleotide of E35, wherein the miRNA binding site comprises one or more nucleotide sequences selected from SEQ ID NO: 34 and SEQ ID NO: 36.
- [1353] E37. The polynucleotide of E36, wherein the miRNA binding site binds to miR-142.
- [1354] E38. The polynucleotide of E36 or E37, wherein the miRNA binding site binds to miR-142-3p or miR-142-5p.
- [1355] E39. The polynucleotide of E37 or E38, wherein the miR-142 comprises SEQ ID NO: 32.
- [1356] E40. The polynucleotide of any one of E1 to E39, wherein the polynucleotide further comprises a 5' UTR.
- [1357] E41. The polynucleotide of E40, wherein the 5' UTR comprises a nucleic acid sequence at least 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99%, or about 100% identical to a 5' UTR sequence selected from the group consisting of SEQ ID NO: 38, 40 to 57, and 117 to 119, or any combination thereof.
- [1358] E42. The polynucleotide of any one of E1 to E41, wherein the polynucleotide further comprises a 3' UTR.

- [1359] E43. The polynucleotide of E42, wherein the 3' UTR comprises a nucleic acid sequence at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99%, or about 100% identical to a 3' UTR sequence selected from the group consisting of SEQ ID NO: 39, 58 to 84, 90, 105, 108 to 115, and 120 to 130, or any combination thereof.
- [1360] E44. The polynucleotide of E42 or E43, wherein the miRNA binding site is located within the 3' UTR.
- [1361] E45. The polynucleotide of any one of E1 to E44, wherein the polynucleotide further comprises a 5' terminal cap.
- [1362] E46. The polynucleotide of E45, wherein the 5' terminal cap comprises a Cap0, Cap1, ARCA, inosine, N1-methyl-guanosine, 2'-fluoro-guanosine, 7-deaza-guanosine, 8-oxo-guanosine, 2-amino-guanosine, LNA-guanosine, 2-azidoguanosine, Cap2, Cap4, 5' methylG cap, or an analog thereof.
- [1363] E47. The polynucleotide of any one of E1 to E46, wherein the polynucleotide further comprises a poly-A region.
- [1364] E48. The polynucleotide of E47, wherein the poly-A region is at least about 10, at least about 20, at least about 30, at least about 40, at least about 50, at least about 60, at least about 70, at least about 80, or at least about 90 nucleotides in length.
- [1365] E49. The polynucleotide of E48, wherein the poly-A region has about 10 to about 200, about 20 to about 180, about 50 to about 160, about 70 to about 140, or about 80 to about 120 nucleotides in length.
- [1366] E50. The polynucleotide of any one of E1 to E49, wherein the polynucleotide encodes an ACADVL polypeptide that is fused to one or more heterologous polypeptides.
- [1367] E51. The polynucleotide of E50, wherein the one or more heterologous polypeptides increase a pharmacokinetic property of the polypeptide.
- [1368] E52. The polynucleotide of any one of E1 to E51, wherein upon administration to a subject, the polynucleotide has:
- (i) a longer plasma half-life;
 - (ii) increased expression of an ACADVL polypeptide encoded by the ORF;
 - (iii) a lower frequency of arrested translation resulting in an expression fragment;
 - (iv) greater structural stability; or
 - (v) any combination thereof,
- relative to a corresponding polynucleotide comprising SEQ ID NOs: 2, 4, or 6.

- [1369] E53. The polynucleotide of any one of E1 to E52, wherein the polynucleotide comprises:
- (i) a 5'-terminal cap;
 - (ii) a 5'-UTR;
 - (iii) an ORF encoding an ACADVL polypeptide;
 - (iv) a 3'-UTR; and
 - (v) a poly-A region.
- [1370] E54. The polynucleotide of E54, wherein the 3'-UTR comprises a miRNA binding site.
- [1371] E55. A method of producing the polynucleotide of any one of E1 to E54, the method comprising modifying an ORF encoding an ACADVL polypeptide by substituting at least one uracil nucleobase with an adenine, guanine, or cytosine nucleobase, or by substituting at least one adenine, guanine, or cytosine nucleobase with a uracil nucleobase, wherein all the substitutions are synonymous substitutions.
- [1372] E56. The method of E55, wherein the method further comprises replacing at least about 90%, at least about 95%, at least about 99%, or about 100% of uracils with 5-methoxyuracils.
- [1373] E57. A composition comprising the polynucleotide of any one of E1 to E54; and a delivery agent.
- [1374] E58. The composition of E57, wherein the delivery agent comprises a lipidoid, a liposome, a lipoplex, a lipid nanoparticle, a polymeric compound, a peptide, a protein, a cell, a nanoparticle mimic, a nanotube, or a conjugate.
- [1375] E59. The composition of E57, wherein the delivery agent comprises a lipid nanoparticle.
- [1376] E60. The composition of E59, wherein the lipid nanoparticle comprises a lipid selected from the group consisting of
- 3-(didodecylamino)-N1,N1,4-tridodecyl-1-piperazineethanamine (KL10),
 - N1-[2-(didodecylamino)ethyl]-N1,N4,N4-tridodecyl-1,4-piperazinediethanamine (KL22),
 - 14,25-ditridecyl-15,18,21,24-tetraaza-octatriacontane (KL25),
 - 1,2-dilinoleyloxy-N,N-dimethylaminopropane (DLin-DMA),
 - 2,2-dilinoleyloxy-4-dimethylaminomethyl-[1,3]-dioxolane (DLin-K-DMA),
 - heptatriaconta-6,9,28,31-tetraen-19-yl 4-(dimethylamino)butanoate (DLin-MC3-DMA),
 - 2,2-dilinoleyloxy-4-(2-dimethylaminoethyl)-[1,3]-dioxolane (DLin-KC2-DMA),

1,2-dioleoyloxy-N,N-dimethylaminopropane (DODMA), (13Z,165Z)-N,N-dimethyl-3-nonydocosa-13-16-dien-1-amine (L608),
 2-({8-[(3 β)-cholest-5-en-3-yloxy]octyl}oxy)-N,N-dimethyl-3-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]propan-1-amine (Octyl-CLinDMA),
 (2R)-2-({8-[(3 β)-cholest-5-en-3-yloxy]octyl}oxy)-N,N-dimethyl-3-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]propan-1-amine (Octyl-CLinDMA (2R)),
 (2S)-2-({8-[(3 β)-cholest-5-en-3-yloxy]octyl}oxy)-N,N-dimethyl-3-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]propan-1-amine (Octyl-CLinDMA (2S)), and any combinations thereof.

[1377] E61. The composition of any one of E57 to E60, wherein the delivery agent comprises a compound having the Formula (I)



(I),

or a salt or stereoisomer thereof, wherein

R₁ is selected from the group consisting of C₅₋₂₀ alkyl, C₅₋₂₀ alkenyl, -R*YR'', -YR'', and -R''M'R';

R₂ and R₃ are independently selected from the group consisting of H, C₁₋₁₄ alkyl, C₂₋₁₄ alkenyl, -R*YR'', -YR'', and -R*OR'', or R₂ and R₃, together with the atom to which they are attached, form a heterocycle or carbocycle;

R₄ is selected from the group consisting of a C₃₋₆ carbocycle, -(CH₂)_nQ, -(CH₂)_nCHQR, -CHQR, -CQ(R)₂, and unsubstituted C₁₋₆ alkyl, where Q is selected from a carbocycle, heterocycle, -OR, -O(CH₂)_nN(R)₂, -C(O)OR, -OC(O)R, -CX₃, -CX₂H, -CXH₂, -CN, -N(R)₂, -C(O)N(R)₂, -N(R)C(O)R, -N(R)S(O)₂R, -N(R)C(O)N(R)₂, -N(R)C(S)N(R)₂, and -C(R)N(R)₂C(O)OR, and each n is independently selected from 1, 2, 3, 4, and 5;


each R₅ is independently selected from the group consisting of C₁₋₃ alkyl, C₂₋₃ alkenyl, and H;

each R₆ is independently selected from the group consisting of C₁₋₃ alkyl, C₂₋₃ alkenyl, and H;

M and M' are independently selected from -C(O)O-, -OC(O)-, -C(O)N(R')-, -N(R')C(O)-, -C(O)-, -C(S)-, -C(S)S-, -SC(S)-, -CH(OH)-, -P(O)(OR')O-, -S(O)₂-, an aryl group, and a heteroaryl group;

R_7 is selected from the group consisting of C_{1-3} alkyl, C_{2-3} alkenyl, and H;
 each R is independently selected from the group consisting of C_{1-3} alkyl, C_{2-3} alkenyl, and H;
 each R' is independently selected from the group consisting of C_{1-18} alkyl, C_{2-18} alkenyl, $-R^*YR''$, $-YR''$, and H;
 each R'' is independently selected from the group consisting of C_{3-14} alkyl and C_{3-14} alkenyl;
 each R^* is independently selected from the group consisting of C_{1-12} alkyl and C_{2-12} alkenyl;
 each Y is independently a C_{3-6} carbocycle;
 each X is independently selected from the group consisting of F, Cl, Br, and I; and
 m is selected from 5, 6, 7, 8, 9, 10, 11, 12, and 13; and
 provided when R_4 is $-(CH_2)_nQ$, $-(CH_2)_nCHQR$, $-CHQR$, or $-CQ(R)_2$, then (i) Q is not $-N(R)_2$ when n is 1, 2, 3, 4 or 5, or (ii) Q is not 5, 6, or 7-membered heterocycloalkyl when n is 1 or 2.

[1378] E62. A composition comprising a nucleotide sequence encoding an ACADVL polypeptide and a delivery agent, wherein the delivery agent comprises a compound having the Formula (I)



or a salt or stereoisomer thereof, wherein

R_1 is selected from the group consisting of C_{5-20} alkyl, C_{5-20} alkenyl, $-R^*YR''$, $-YR''$, and $-R''M'R'$;

R_2 and R_3 are independently selected from the group consisting of H, C_{1-14} alkyl, C_{2-14} alkenyl, $-R^*YR''$, $-YR''$, and $-R^*OR''$, or R_2 and R_3 , together with the atom to which they are attached, form a heterocycle or carbocycle;

R_4 is selected from the group consisting of a C_{3-6} carbocycle, $-(CH_2)_nQ$, $-(CH_2)_nCHQR$, $-CHQR$, $-CQ(R)_2$, and unsubstituted C_{1-6} alkyl, where Q is selected from a carbocycle, heterocycle, $-OR$, $-O(CH_2)_nN(R)_2$, $-C(O)OR$, $-OC(O)R$, $-CX_3$, $-CX_2H$, $-CXH_2$, $-CN$, $-N(R)_2$, $-C(O)N(R)_2$, $-N(R)C(O)R$, $-N(R)S(O)_2R$, $-N(R)C(O)N(R)_2$,

-N(R)C(S)N(R)₂, and -C(R)N(R)₂C(O)OR, and each n is independently selected from 1, 2, 3, 4, and 5;

each R₅ is independently selected from the group consisting of C₁₋₃ alkyl, C₂₋₃ alkenyl, and H;

each R₆ is independently selected from the group consisting of C₁₋₃ alkyl, C₂₋₃ alkenyl, and H;

M and M' are independently selected from -C(O)O-, -OC(O)-, -C(O)N(R')-, -N(R')C(O)-, -C(O)-, -C(S)-, -C(S)S-, -SC(S)-, -CH(OH)-, -P(O)(OR')O-, -S(O)₂-, an aryl group, and a heteroaryl group;

R₇ is selected from the group consisting of C₁₋₃ alkyl, C₂₋₃ alkenyl, and H;

each R is independently selected from the group consisting of C₁₋₃ alkyl, C₂₋₃ alkenyl, and H;

each R' is independently selected from the group consisting of C₁₋₁₈ alkyl, C₂₋₁₈ alkenyl,

-R*YR'', -YR'', and H:

each R'' is independently selected from the group consisting of C₃₋₁₄ alkyl and C₃₋₁₄ alkenyl;


each R* is independently selected from the group consisting of C₁₋₁₂ alkyl and C₂₋₁₂ alkenyl;

each Y is independently a C₃₋₆ carbocycle;

each X is independently selected from the group consisting of F, Cl, Br, and I; and m is selected from 5, 6, 7, 8, 9, 10, 11, 12, and 13; and

provided when R₄ is -(CH₂)_nQ, -(CH₂)_nCHQR, -CHQR, or -CQ(R)₂, then (i) Q is not -N(R)₂ when n is 1, 2, 3, 4 or 5, or (ii) Q is not 5, 6, or 7-membered heterocycloalkyl when n is 1 or 2.

[1379] E63. The composition of E61 or E62, wherein the compound is of Formula (IA):

 $(IA)_i$

or a salt or stereoisomer thereof, wherein

1 is selected from 1, 2, 3, 4, and 5:

m is selected from 5, 6, 7, 8, and 9;

M_1 is a bond or M' ;


R_4 is unsubstituted C_{1-3} alkyl, or $-(CH_2)_nQ$, in which n is 1, 2, 3, 4, or 5 and Q is OH, $-NHC(S)N(R)_2$, or $-NHC(O)N(R)_2$;

M and M' are independently selected from $-C(O)O-$, $-OC(O)-$, $-C(O)N(R')$, $-P(O)(OR')O-$, an aryl group, and a heteroaryl group; and

R_2 and R_3 are independently selected from the group consisting of H, C_{1-14} alkyl, and C_{2-14} alkenyl.

[1380] E64. The composition of any one of E61 to E63, wherein m is 5, 7, or 9.

[1381] E65. The composition of any one of E61 to E64, wherein the compound is of Formula (II):



or a salt or stereoisomer thereof, wherein

l is selected from 1, 2, 3, 4, and 5;

M_1 is a bond or M' ;

R_4 is unsubstituted C_{1-3} alkyl, or $-(CH_2)_nQ$, in which n is 2, 3, or 4 and Q is OH, $-NHC(S)N(R)_2$, or $-NHC(O)N(R)_2$;

M and M' are independently selected from $-C(O)O-$, $-OC(O)-$, $-C(O)N(R')$, $-P(O)(OR')O-$, an aryl group, and a heteroaryl group; and

R_2 and R_3 are independently selected from the group consisting of H, C_{1-14} alkyl, and C_{2-14} alkenyl.


[1382] E66. The composition of any one of E63 to E65, wherein M_1 is M' .

[1383] E67. The composition of E66, wherein M and M' are independently $-C(O)O-$ or $-OC(O)-$.

[1384] E68. The composition of any one of E63 to E67, wherein l is 1, 3, or 5.


[1385] E69. The composition of E61 or E62, wherein the compound is selected from the group consisting of Compound 1 to Compound 147, salts and isomers thereof, and any combination thereof.

[1386] E70. The composition of E61 or E62, wherein the compound is of the Formula (IIa),




(IIa), or a salt or stereoisomer thereof.

[1387] E71. The composition of E61 or E62, wherein the compound is of the Formula (IIb),




(IIb), or a salt or stereoisomer thereof.

[1388] E72. The composition of E61 or E62, wherein the compound is of the Formula (IIc) or (Ile),



(IIc),




(Ile),

or a salt or stereoisomer thereof.

[1389] E73. The composition of any one of E70 to E72, wherein R_4 is selected from $-(CH_2)_nQ$ and $-(CH_2)_nCHQR$.

[1390] E74. The composition of E61 or E62, wherein the compound is of the Formula (IIId),



(IIId), or a salt or stereoisomer thereof,

wherein R_2 and R_3 are independently selected from the group consisting of C_{5-14} alkyl and C_{5-14} alkenyl, n is selected from 2, 3, and 4, and R' , R'' , R_5 , R_6 and m are as defined in E61 or 62.

- [1391] E75. The composition of E74, wherein R_2 is C_8 alkyl.
- [1392] E76. The composition of E75, wherein R_3 is C_5 alkyl, C_6 alkyl, C_7 alkyl, C_8 alkyl, or C_9 alkyl.
- [1393] E77. The composition of any one of E74 to E76, wherein m is 5, 7, or 9.
- [1394] E78. The composition of any one of E74 to E77, wherein each R_5 is H.
- [1395] E79. The composition of E78, wherein each R_6 is H.
- [1396] E80. The composition of any one of E61 to E79, which is a nanoparticle composition.
- [1397] E81. The composition of E80, wherein the delivery agent further comprises a phospholipid.
- [1398] E82. The composition of E81, wherein the phospholipid is selected from the group consisting of 1,2-dilinoleoyl-sn-glycero-3-phosphocholine (DLPC), 1,2-dimyristoyl-sn-glycero-phosphocholine (DMPC), 1,2-dioleoyl-sn-glycero-3-phosphocholine (DOPC), 1,2-dipalmitoyl-sn-glycero-3-phosphocholine (DPPC), 1,2-distearoyl-sn-glycero-3-phosphocholine (DSPC), 1,2-diundecanoyl-sn-glycero-phosphocholine (DUPC), 1-palmitoyl-2-oleoyl-sn-glycero-3-phosphocholine (POPC), 1,2-di-O-octadecenyl-sn-glycero-3-phosphocholine (18:0 Diether PC), 1-oleoyl-2-cholesterylhemisuccinoyl-sn-glycero-3-phosphocholine (OChemPC), 1-hexadecyl-sn-glycero-3-phosphocholine (C16 Lyso PC), 1,2-dilinenoyl-sn-glycero-3-phosphocholine, 1,2-diarachidonoyl-sn-glycero-3-phosphocholine, 1,2-didocosahexaenoyl-sn-glycero-3-phosphocholine, 1,2-dioleoyl-sn-glycero-3-phosphoethanolamine (DOPE), 1,2-diphytanoyl-sn-glycero-3-phosphoethanolamine (ME 16:0 PE), 1,2-distearoyl-sn-glycero-3-phosphoethanolamine, 1,2-dilinoleoyl-sn-glycero-3-phosphoethanolamine, 1,2-dilinenoyl-sn-glycero-3-phosphoethanolamine, 1,2-diarachidonoyl-sn-glycero-3-phosphoethanolamine,

1,2-didocosahexaenoyl-sn-glycero-3-phosphoethanolamine, 1,2-dioleoyl-sn-glycero-3-phospho-rac-(1-glycerol) sodium salt (DOPG), sphingomyelin, and any mixtures thereof.

- [1399] E83. The composition of any one of E61 to E82, wherein the delivery agent further comprises a structural lipid.
- [1400] E84. The composition of E83, wherein the structural lipid is selected from the group consisting of cholesterol, fecosterol, sitosterol, ergosterol, campesterol, stigmasterol, brassicasterol, tomatidine, ursolic acid, alpha-tocopherol, and any mixtures thereof.
- [1401] E85. The composition of any one of E61 to E84, wherein the delivery agent further comprises a PEG lipid.
- [1402] E86. The composition of E85, wherein the PEG lipid is selected from the group consisting of a PEG-modified phosphatidylethanolamine, a PEG-modified phosphatidic acid, a PEG-modified ceramide, a PEG-modified dialkylamine, a PEG-modified diacylglycerol, a PEG-modified dialkylglycerol, and any mixtures thereof.
- [1403] E87. The composition of any one of E61 to E86, wherein the delivery agent further comprises an ionizable lipid selected from the group consisting of 3-(didodecylamino)-N1,N1,4-tridodecyl-1-piperazineethanamine (KL10), N1-[2-(didodecylamino)ethyl]-N1,N4,N4-tridodecyl-1,4-piperazinediethanamine (KL22), 14,25-ditridecyl-15,18,21,24-tetraaza-octatriacontane (KL25), 1,2-dilinoleyloxy-N,N-dimethylaminopropane (DLin-DMA), 2,2-dilinoleyloxy-4-dimethylaminomethyl-[1,3]-dioxolane (DLin-K-DMA), heptatriaconta-6,9,28,31-tetraen-19-yl 4-(dimethylamino)butanoate (DLin-MC3-DMA), 2,2-dilinoleyloxy-4-(2-dimethylaminoethyl)-[1,3]-dioxolane (DLin-KC2-DMA), 1,2-dioleyloxy-N,N-dimethylaminopropane (DODMA), 2-({8-[(3 β)-cholest-5-en-3-yloxy]octyl}oxy)-N,N-dimethyl-3-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]propan-1-amine (Octyl-CLinDMA), (2R)-2-({8-[(3 β)-cholest-5-en-3-yloxy]octyl}oxy)-N,N-dimethyl-3-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]propan-1-amine (Octyl-CLinDMA (2R)), and (2S)-2-({8-[(3 β)-cholest-5-en-3-yloxy]octyl}oxy)-N,N-dimethyl-3-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]propan-1-amine (Octyl-CLinDMA (2S)).

- [1404] E88. The composition of any one of E61 to E87, wherein the delivery agent further comprises a phospholipid, a structural lipid, a PEG lipid, or any combination thereof.
- [1405] E89. The composition of any one of E61 to E88, wherein the composition is formulated for *in vivo* delivery.
- [1406] E90. The composition according any one of E61 to E89, which is formulated for intramuscular, subcutaneous, or intradermal delivery.
- [1407] E91. A host cell comprising the polynucleotide of any one of E1 to E54.
- [1408] E92. The host cell of E91, wherein the host cell is a eukaryotic cell.
- [1409] E93. A vector comprising the polynucleotide of any one of E1 to E54.
- [1410] E94. A method of making a polynucleotide comprising enzymatically or chemically synthesizing the polynucleotide of any one of E1 to E54.
- [1411] E95. A polypeptide encoded by the polynucleotide of any one of E1 to E54, the composition of any one of E57 to E90, the host cell of E91 or E92, or the vector of E93 or produced by the method of E94.
- [1412] E96. A method of expressing *in vivo* an active ACADVL polypeptide in a subject in need thereof comprising administering to the subject an effective amount of the polynucleotide of any one of E1 to E54, the composition of any one of E57 to E90, the host cell of E91 or E92, or the vector of E93.
- [1413] E97. A method of treating very long-chain acyl-CoA dehydrogenase (VLCAD) deficiency in a subject in need thereof comprising administering to the subject a therapeutically effective amount of the polynucleotide of any one of E1 to E54, the composition of any one of E57 to E90, the host cell of E91 or E92, or the vector of E93, wherein the administration alleviates the signs or symptoms of VLCAD deficiency in the subject.
- [1414] E98. A method to prevent or delay the onset of VLCAD deficiency signs or symptoms in a subject in need thereof comprising administering to the subject a prophylactically effective amount of the polynucleotide of any one of E1 to E54, the composition of any one of E57 to E90, the host cell of E91 or E92, or the vector of E93 before VLCAD deficiency signs or symptoms manifest, wherein the administration prevents or delays the onset of VLCAD deficiency signs or symptoms in the subject.
- [1415] E99. A method to ameliorate the signs or symptoms of VLCAD deficiency in a subject in need thereof comprising administering to the subject a therapeutically effective

amount of the polynucleotide of any one of E1 to E54, the composition of any one of E57 to E90, the host cell of E91 or E92, or the vector of E93 before VLCAD deficiency signs or symptoms manifest, wherein the administration ameliorates VLCAD deficiency signs or symptoms in the subject.

31. Equivalents and Scope

[1416] Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments in accordance with the invention described herein. The scope of the present invention is not intended to be limited to the above Description, but rather is as set forth in the appended claims.

[1417] In the claims, articles such as "a," "an," and "the" can mean one or more than one unless indicated to the contrary or otherwise evident from the context. Claims or descriptions that include "or" between one or more members of a group are considered satisfied if one, more than one, or all of the group members are present in, employed in, or otherwise relevant to a given product or process unless indicated to the contrary or otherwise evident from the context. The invention includes embodiments in which exactly one member of the group is present in, employed in, or otherwise relevant to a given product or process. The invention includes embodiments in which more than one, or all of the group members are present in, employed in, or otherwise relevant to a given product or process.

[1418] It is also noted that the term "comprising" is intended to be open and permits but does not require the inclusion of additional elements or steps. When the term "comprising" is used herein, the term "consisting of" is thus also encompassed and disclosed.

[1419] Where ranges are given, endpoints are included. Furthermore, it is to be understood that unless otherwise indicated or otherwise evident from the context and understanding of one of ordinary skill in the art, values that are expressed as ranges can assume any specific value or subrange within the stated ranges in different embodiments of the invention, to the tenth of the unit of the lower limit of the range, unless the context clearly dictates otherwise.

[1420] In addition, it is to be understood that any particular embodiment of the present invention that falls within the prior art can be explicitly excluded from any one or more of the claims. Since such embodiments are deemed to be known to one of ordinary skill in

the art, they can be excluded even if the exclusion is not set forth explicitly herein. Any particular embodiment of the compositions of the invention (*e.g.*, any nucleic acid or protein encoded thereby; any method of production; any method of use; etc.) can be excluded from any one or more claims, for any reason, whether or not related to the existence of prior art.

- [1421] All cited sources, for example, references, publications, databases, database entries, and art cited herein, are incorporated into this application by reference, even if not expressly stated in the citation. In case of conflicting statements of a cited source and the instant application, the statement in the instant application shall control.
- [1422] Section and table headings are not intended to be limiting.

Examples

EXAMPLE 1

Chimeric Polynucleotide Synthesis

A. *Triphosphate route*

- [1423] Two regions or parts of a chimeric polynucleotide can be joined or ligated using triphosphate chemistry. According to this method, a first region or part of 100 nucleotides or less can be chemically synthesized with a 5' monophosphate and terminal 3'desOH or blocked OH. If the region is longer than 80 nucleotides, it can be synthesized as two strands for ligation.
- [1424] If the first region or part is synthesized as a non-positionally modified region or part using *in vitro* transcription (IVT), conversion the 5'monophosphate with subsequent capping of the 3' terminus can follow. Monophosphate protecting groups can be selected from any of those known in the art.
- [1425] The second region or part of the chimeric polynucleotide can be synthesized using either chemical synthesis or IVT methods. IVT methods can include an RNA polymerase that can utilize a primer with a modified cap. Alternatively, a cap of up to 80 nucleotides can be chemically synthesized and coupled to the IVT region or part.
- [1426] It is noted that for ligation methods, ligation with DNA T4 ligase, followed by treatment with DNase should readily avoid concatenation.

[1427] The entire chimeric polynucleotide need not be manufactured with a phosphate-sugar backbone. If one of the regions or parts encodes a polypeptide, then such region or part can comprise a phosphate-sugar backbone.

[1428] Ligation can then be performed using any known click chemistry, orthoclick chemistry, solulink, or other bioconjugate chemistries known to those in the art.

B. Synthetic route

[1429] The chimeric polynucleotide can be made using a series of starting segments. Such segments include:

(a) Capped and protected 5' segment comprising a normal 3'OH (SEG. 1)

(b) 5' triphosphate segment which can include the coding region of a polypeptide and comprising a normal 3'OH (SEG. 2)

(c) 5' monophosphate segment for the 3' end of the chimeric polynucleotide (*e.g.*, the tail) comprising cordycepin or no 3'OH (SEG. 3)

[1430] After synthesis (chemical or IVT), segment 3 (SEG. 3) can be treated with cordycepin and then with pyrophosphatase to create the 5'monophosphate.

[1431] Segment 2 (SEG. 2) can then be ligated to SEG. 3 using RNA ligase. The ligated polynucleotide can then be purified and treated with pyrophosphatase to cleave the diphosphate. The treated SEG.2-SEG. 3 construct is then purified and SEG. 1 is ligated to the 5' terminus. A further purification step of the chimeric polynucleotide can be performed.

[1432] Where the chimeric polynucleotide encodes a polypeptide, the ligated or joined segments can be represented as: 5'UTR (SEG. 1), open reading frame or ORF (SEG. 2) and 3'UTR+PolyA (SEG. 3).

[1433] The yields of each step can be as much as 90-95%.

EXAMPLE 2

PCR for cDNA Production

[1434] PCR procedures for the preparation of cDNA can be performed using 2x KAPA HIFI™ HotStart ReadyMix by Kapa Biosystems (Woburn, MA). This system includes 2x KAPA ReadyMix 12.5 µl; Forward Primer (10 µM) 0.75 µl; Reverse Primer (10 µM) 0.75 µl; Template cDNA -100 ng; and dH₂O diluted to 25.0 µl. The PCR reaction

conditions can be: at 95° C for 5 min. and 25 cycles of 98° C for 20 sec, then 58° C for 15 sec, then 72° C for 45 sec, then 72° C for 5 min. then 4° C to termination.

[1435] The reverse primer of the instant invention can incorporate a poly-T₁₂₀ for a poly-A₁₂₀ in the mRNA. Other reverse primers with longer or shorter poly(T) tracts can be used to adjust the length of the poly(A) tail in the polynucleotide mRNA.

[1436] The reaction can be cleaned up using Invitrogen's PURELINK™ PCR Micro Kit (Carlsbad, CA) per manufacturer's instructions (up to 5 µg). Larger reactions will require a cleanup using a product with a larger capacity. Following the cleanup, the cDNA can be quantified using the NANODROP™ and analyzed by agarose gel electrophoresis to confirm the cDNA is the expected size. The cDNA can then be submitted for sequencing analysis before proceeding to the *in vitro* transcription reaction.

EXAMPLE 3

In vitro Transcription (IVT)

[1437] The *in vitro* transcription reactions can generate polynucleotides containing uniformly modified polynucleotides. Such uniformly modified polynucleotides can comprise a region or part of the polynucleotides of the invention. The input nucleotide triphosphate (NTP) mix can be made using natural and un-natural NTPs.

[1438] A typical *in vitro* transcription reaction can include the following:

- 1 Template cDNA—1.0 µg
- 2 10x transcription buffer (400 mM Tris-HCl pH 8.0, 190 mM MgCl₂, 50 mM DTT, 10 mM Spermidine)—2.0 µl
- 3 Custom NTPs (25mM each) —7.2 µl
- 4 RNase Inhibitor—20 U
- 5 T7 RNA polymerase —3000 U
- 6 dH₂O—Up to 20.0 µl. and
- 7 Incubation at 37° C for 3 hr-5 hrs.

[1439] The crude IVT mix can be stored at 4° C overnight for cleanup the next day. 1 U of RNase-free DNase can then be used to digest the original template. After 15 minutes of incubation at 37° C, the mRNA can be purified using Ambion's MEGACLEAR™ Kit (Austin, TX) following the manufacturer's instructions. This kit can purify up to 500 µg of RNA. Following the cleanup, the RNA can be quantified using the NanoDrop and

analyzed by agarose gel electrophoresis to confirm the RNA is the proper size and that no degradation of the RNA has occurred.

EXAMPLE 4

Enzymatic Capping

[1440] Capping of a polynucleotide can be performed with a mixture includes: IVT RNA 60 µg-180µg and dH₂O up to 72 µl. The mixture can be incubated at 65° C for 5 minutes to denature RNA, and then can be transferred immediately to ice.

[1441] The protocol can then involve the mixing of 10x Capping Buffer (0.5 M Tris-HCl (pH 8.0), 60 mM KCl, 12.5 mM MgCl₂) (10.0 µl); 20 mM GTP (5.0 µl); 20 mM S-Adenosyl Methionine (2.5 µl); RNase Inhibitor (100 U); 2'-O-Methyltransferase (400U); Vaccinia capping enzyme (Guanylyl transferase) (40 U); dH₂O (Up to 28 µl); and incubation at 37° C for 30 minutes for 60 µg RNA or up to 2 hours for 180 µg of RNA.

[1442] The polynucleotide can then be purified using Ambion's MEGACLEAR™ Kit (Austin, TX) following the manufacturer's instructions. Following the cleanup, the RNA can be quantified using the NANODROP™ (ThermoFisher, Waltham, MA) and analyzed by agarose gel electrophoresis to confirm the RNA is the proper size and that no degradation of the RNA has occurred. The RNA product can also be sequenced by running a reverse-transcription-PCR to generate the cDNA for sequencing.

EXAMPLE 5

PolyA Tailing Reaction

[1443] Without a poly-T in the cDNA, a poly-A tailing reaction must be performed before cleaning the final product. This can be done by mixing Capped IVT RNA (100 µl); RNase Inhibitor (20 U); 10x Tailing Buffer (0.5 M Tris-HCl (pH 8.0), 2.5 M NaCl, 100 mM MgCl₂)(12.0 µl); 20 mM ATP (6.0 µl); Poly-A Polymerase (20 U); dH₂O up to 123.5 µl and incubating at 37° C for 30 min. If the poly-A tail is already in the transcript, then the tailing reaction can be skipped and proceed directly to cleanup with Ambion's MEGACLEAR™ kit (Austin, TX) (up to 500 µg). Poly-A Polymerase is, in some cases, a recombinant enzyme expressed in yeast.

[1444] It should be understood that the processivity or integrity of the polyA tailing reaction does not always result in an exact size polyA tail. Hence polyA tails of approximately between 40-200 nucleotides, *e.g.*, about 40, 50, 60, 70, 80, 90, 91, 92, 93,

94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 150-165, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164 or 165 are within the scope of the invention.

EXAMPLE 6

Natural 5' Caps and 5' Cap Analogues

[1445] 5'-capping of polynucleotides can be completed concomitantly during the *in vitro*-transcription reaction using the following chemical RNA cap analogs to generate the 5'-guanosine cap structure according to manufacturer protocols: 3'-O-Me-m7G(5')ppp(5') G [the ARCA cap]; G(5')ppp(5')A; G(5')ppp(5')G; m7G(5')ppp(5')A; m7G(5')ppp(5')G (New England BioLabs, Ipswich, MA). 5'-capping of modified RNA can be completed post-transcriptionally using a Vaccinia Virus Capping Enzyme to generate the "Cap 0" structure: m7G(5')ppp(5')G (New England BioLabs, Ipswich, MA). Cap 1 structure can be generated using both Vaccinia Virus Capping Enzyme and a 2'-O methyl-transferase to generate: m7G(5')ppp(5')G-2'-O-methyl. Cap 2 structure can be generated from the Cap 1 structure followed by the 2'-O-methylation of the 5'-antepenultimate nucleotide using a 2'-O methyl-transferase. Cap 3 structure can be generated from the Cap 2 structure followed by the 2'-O-methylation of the 5'-preantepenultimate nucleotide using a 2'-O methyl-transferase. Enzymes can be derived from a recombinant source.

[1446] When transfected into mammalian cells, the modified mRNAs can have a stability of between 12-18 hours or more than 18 hours, *e.g.*, 24, 36, 48, 60, 72 or greater than 72 hours.

EXAMPLE 7

Capping Assays

A. Protein Expression Assay

[1447] Polynucleotides encoding a polypeptide, containing any of the caps taught herein, can be transfected into cells at equal concentrations. After 6, 12, 24 and 36 hours post-transfection, the amount of protein secreted into the culture medium can be assayed by ELISA. Synthetic polynucleotides that secrete higher levels of protein into the medium would correspond to a synthetic polynucleotide with a higher translationally-competent Cap structure.

B. Purity Analysis Synthesis

[1448] Polynucleotides encoding a polypeptide, containing any of the caps taught herein, can be compared for purity using denaturing Agarose-Urea gel electrophoresis or HPLC analysis. Polynucleotides with a single, consolidated band by electrophoresis correspond to the higher purity product compared to polynucleotides with multiple bands or streaking bands. Synthetic polynucleotides with a single HPLC peak would also correspond to a higher purity product. The capping reaction with a higher efficiency would provide a more pure polynucleotide population.

C. Cytokine Analysis

[1449] Polynucleotides encoding a polypeptide, containing any of the caps taught herein, can be transfected into cells at multiple concentrations. After 6, 12, 24 and 36 hours post-transfection the amount of pro-inflammatory cytokines such as TNF-alpha and IFN-beta secreted into the culture medium can be assayed by ELISA. Polynucleotides resulting in the secretion of higher levels of pro-inflammatory cytokines into the medium would correspond to polynucleotides containing an immune-activating cap structure.

D. Capping Reaction Efficiency

[1450] Polynucleotides encoding a polypeptide, containing any of the caps taught herein, can be analyzed for capping reaction efficiency by LC-MS after nuclease treatment. Nuclease treatment of capped polynucleotides would yield a mixture of free nucleotides and the capped 5'-5-triphosphate cap structure detectable by LC-MS. The amount of capped product on the LC-MS spectra can be expressed as a percent of total polynucleotide from the reaction and would correspond to capping reaction efficiency. The cap structure with higher capping reaction efficiency would have a higher amount of capped product by LC-MS.

EXAMPLE 8**Agarose Gel Electrophoresis of Modified RNA or RT PCR Products**

[1451] Individual polynucleotides (200-400 ng in a 20 µl volume) or reverse transcribed PCR products (200-400 ng) can be loaded into a well on a non-denaturing 1.2% Agarose E-Gel (Invitrogen, Carlsbad, CA) and run for 12-15 minutes according to the manufacturer protocol.

EXAMPLE 9**Nanodrop Modified RNA Quantification and UV Spectral Data**

- [1452] Modified polynucleotides in TE buffer (1 μ l) can be used for Nanodrop UV absorbance readings to quantitate the yield of each polynucleotide from an chemical synthesis or *in vitro* transcription reaction.

EXAMPLE 10**Formulation of Modified mRNA Using Lipidoids**

- [1453] Polynucleotides can be formulated for *in vitro* experiments by mixing the polynucleotides with the lipidoid at a set ratio prior to addition to cells. *In vivo* formulation can require the addition of extra ingredients to facilitate circulation throughout the body. To test the ability of these lipidoids to form particles suitable for *in vivo* work, a standard formulation process used for siRNA-lipidoid formulations can be used as a starting point. After formation of the particle, polynucleotide can be added and allowed to integrate with the complex. The encapsulation efficiency can be determined using a standard dye exclusion assays.

EXAMPLE 11**Method of Screening for Protein Expression****A. *Electrospray Ionization***

- [1454] A biological sample that can contain proteins encoded by a polynucleotide administered to the subject can be prepared and analyzed according to the manufacturer protocol for electrospray ionization (ESI) using 1, 2, 3 or 4 mass analyzers. A biologic sample can also be analyzed using a tandem ESI mass spectrometry system.
- [1455] Patterns of protein fragments, or whole proteins, can be compared to known controls for a given protein and identity can be determined by comparison.

B. *Matrix-Assisted Laser Desorption/Ionization*

- [1456] A biological sample that can contain proteins encoded by one or more polynucleotides administered to the subject can be prepared and analyzed according to the manufacturer protocol for matrix-assisted laser desorption/ionization (MALDI).

- [1457] Patterns of protein fragments, or whole proteins, can be compared to known controls for a given protein and identity can be determined by comparison.

C. *Liquid Chromatography-Mass spectrometry-Mass spectrometry*

- [1458] A biological sample, which can contain proteins encoded by one or more polynucleotides, can be treated with a trypsin enzyme to digest the proteins contained within. The resulting peptides can be analyzed by liquid chromatography-mass spectrometry-mass spectrometry (LC/MS/MS). The peptides can be fragmented in the mass spectrometer to yield diagnostic patterns that can be matched to protein sequence databases via computer algorithms. The digested sample can be diluted to achieve 1 ng or less starting material for a given protein. Biological samples containing a simple buffer background (*e.g.*, water or volatile salts) are amenable to direct in-solution digest; more complex backgrounds (*e.g.*, detergent, non-volatile salts, glycerol) require an additional clean-up step to facilitate the sample analysis.
- [1459] Patterns of protein fragments, or whole proteins, can be compared to known controls for a given protein and identity can be determined by comparison.

EXAMPLE 12

Synthesis of mRNA Encoding ACADVL

- [1460] Sequence optimized polynucleotides encoding ACADVL polypeptides, *i.e.*, SEQ ID NOs: 1, 3, or 5 are synthesized and characterized as described in Examples 1 to 11. mRNA's encoding both human ACADVL are prepared for Examples 13-19 described below, and are synthesized and characterized as described in Examples 1 to 11.
- [1461] An mRNA encoding human ACADVL is constructed, *e.g.*, by using the ORF sequence provided in SEQ ID NO: 2, 4, or 6. The mRNA sequence includes both 5' and 3' UTR regions (see, *e.g.*, SEQ ID NOs: 38 39, and 83). In a construct, the 5'UTR and 3'UTR sequences are:

5' UTR

UCAAGCUUUUGGACCCUCGUACAGAAGCUAAUACGACUCACUAUAGGGAAA
UAAGAGAGAAAAGAAGAGUAAGAAGAAAUAUAAGAGCCACC (SEQ ID NO: 38)

3' UTR

UGAUAUAGGCUGGAGCCUCGGUGGCCAUGCUUCUUGCCCCUUGGGCCUCC
CCCCAGCCCCUCCUCCCCUCCUGCACCCGUACCCCGUGGUCUUUGAAUA

AAGUCUGAGUGGGCGGC (SEQ ID NO: 39)

or

3' UTR

UGAUAAUAGUCCAUAAAGUAGGAAACACUACAGCUGGAGCCUCGGUGGCCAUGCUUCUUGCCCCUU
GGGCCUCCCCCAGCCCCUCCUCCCUUCCUGCACCCGUACCCCCCGCAUUAUUACUCACGGUACG
AGUGGUCUUUGAAUAAAGUCUGAGUGGGCGGC (SEQ ID NO: 83)

- [1462]** The ACADVL mRNA sequence was prepared as modified mRNA. Specifically, during in vitro translation, modified mRNA can be generated using 5-methoxy-UTP to ensure that the mRNAs contain 100% 5-methoxy-uridine instead of uridine. Further, ACADVL-mRNA can be synthesized with a primer that introduces a polyA-tail, and a Cap 1 structure is generated on both mRNAs using Vaccinia Virus Capping Enzyme and a 2'-O methyl-transferase to generate: m7G(5')ppp(5')G-2'-O-methyl.

EXAMPLE 13

Detecting Endogenous ACADVL Expression *In Vitro*

- [1463]** ACADVL expression is characterized in a variety of cell lines derived from both mice and human sources. Cell are cultured in standard conditions and cell extracts are obtained by placing the cells in lysis buffer. For comparison purposes, appropriate controls are also prepared. To analyze ACADVL expression, lysate samples are prepared from the tested cells and mixed with lithium dodecyl sulfate sample loading buffer and subjected to standard Western blot analysis. For detection of ACADVL, the antibody used is a commercial anti-ACADVL antibody. For detection of a load control, the antibody used is anti-citrate synthase (rabbit polyclonal; PA5-22126; Thermo-Fisher Scientific®). To examine the localization of endogenous ACADVL, immunofluorescence analysis is performed on cells. ACADVL expression is detected using a commercial anti-ACADVL. The location of specific organelles can be detected with existing commercial products. For example, mitochondria can be detected using Mitotracker, and the nucleus can be stained with DAPI. Image analysis is performed on a Zeiss ELYRA imaging system.

- [1464] Endogenous ACADVL expression can be used as a base line to determine changes in ACADVL expression resulting from transfection with mRNAs comprising nucleic acids encoding ACADVL.

EXAMPLE 14

In Vitro Expression of ACADVL in HeLa Cells

- [1465] To measure in vitro expression of human ACADVL in HeLa cells, those cells are seeded on 12-well plates (BD Biosciences, San Jose, USA) one day prior to transfection. mRNA formulations comprising human ACADVL or a GFP control are transfected using 800 ng mRNA and 2 μ L Lipofectamin 2000 in 60 μ L OPTI-MEM per well and incubated.
- [1466] After 24 hours, the cells in each well are lysed using a consistent amount of lysis buffer. Appropriate controls are used. Protein concentrations of each are determined using a BCA assay according to manufacturer's instructions. To analyze ACADVL expression, equal loads of each lysate (24 μ g) are prepared in a loading buffer and subjected to standard Western blot analysis. For detection of ACADVL, a commercial anti-ACADVL antibody is used according to the manufacturer's instructions.

EXAMPLE 15

In Vitro ACADVL Activity in HeLa Cells

- [1467] An in vitro ACADVL activity assay is performed to determine whether ACADVL exogenously-expressed after introduction of mRNA comprising an ACADVL sequence is active.

A. Expression Assay

- [1468] HeLa cells are transfected with mRNA formulations comprising human ACADVL or a GFP control. Cells are transfected with Lipofectamin 2000 and lysed as described in Example 14 above. Appropriate controls are also prepared.

B. Activity Assay

- [1469] To assess whether exogenous ACADVL can function, an in vitro activity assay is performed using transfected HeLa cell lysates as the source of enzymatic activity. To begin, lysate is mixed ACADVL substrate. The reaction is stopped by adding 100 g/L

TCA and vortexing. The reaction tubes are then centrifuged at 13,000 g for 1 min, and the supernatant is analyzed for the presence of labeled enzymatic products resulting from the activity of ACADVL using HPLC-based separation and quantification. Specifically, 20 μ L of each activity reaction supernatant are analyzed using a HPLC system equipped with a Quaternary-Pump, a Multi-sampler, a Thermostated Column-Compartment, a Poroshell EC-C18 120 HPLC-column and a Radiometric Detector controlled by OpenLAB Chromatography Data System, all used according to the manufacturers' recommendations.

EXAMPLE 16

Measuring *In Vitro* Expression of ACADVL in Cells

[1470] Cells from normal subjects and VLCADD patients are examined for their capacity to express exogenous ACADVL. Cells are transfected with mRNA formulations comprising human ACADVL, mouse ACADVL, or a GFP control via electroporation using a standard protocol. Each construct is tested separately. After incubation, cells are lysed and protein concentration in each lysate is measured using a suitable assay, e.g., by BCA assay. To analyze ACADVL expression, equal loads of each lysate are prepared in a loading buffer and subjected to standard Western blot analysis. For detection of ACADVL, an anti-ACADVL is used. For detection of a load control, the antibody used is anti-citrate synthase (rabbit polyclonal; MA5-17625; Pierce®).

EXAMPLE 17

Measuring *In Vitro* ACADVL Activity in Lysates

A. Expression

[1471] Cells from normal human subjects and VLCADD patients are cultured. Cells are transfected with mRNA formulations comprising human ACADVL, mouse ACADVL, or a GFP control via electroporation using a standard protocol.

B. Activity Assay

[1472] To assess whether exogenous ACADVL function, an *in vitro* activity assay is performed using transfected cell lysates as the source of enzymatic activity. Lysate containing expressed ACADVL protein is incubated with labeled ACADVL substrate,

and the activity of ACADVL is quantified by measuring the levels of labeled products resulting from the enzymatic activity of ACADVL.

EXAMPLE 18

In Vivo ACADVL expression in Animal Models

[1473] To assess the ability of ACADVL-containing mRNA's to facilitate ACADVL expression *in vivo*, mRNA encoding human ACADVL is introduced into C57B/L6 mice. C57B/L6 mice are injected intravenously with either control mRNA (NT-FIX) or human ACADVL mRNA. The mRNA is formulated in lipid nanoparticles for delivery into the mice. Mice are sacrificed after 24 or 48 hrs. and ACADVL protein levels in liver lysates are determined by capillary electrophoresis (CE). Citrate synthase expression is examined for use as a load control. For control NT-FIX injections, 4 mice are tested for each time point. For human ACADVL mRNA injections, 6 mice are tested for each time point. Treatment with mRNA encoding ACADVL is expected to reliably induce expression of ACADVL.

EXAMPLE 19

Human ACADVL Mutant and Chimeric Constructs

[1474] A polynucleotide of the present invention can comprise at least a first region of linked nucleosides encoding human ACADVL, which can be constructed, expressed, and characterized according to the examples above. Similarly, the polynucleotide sequence can contain one or more mutations that results in the expression of an ACADVL with increased or decreased activity. Furthermore, the polynucleotide sequence encoding ACADVL can be part of a construct encoding a chimeric fusion protein.

EXAMPLE 20

Production of nanoparticle compositions

A. Production of nanoparticle compositions

[1475] Nanoparticles can be made with mixing processes such as microfluidics and T-junction mixing of two fluid streams, one of which contains the polynucleotide and the other has the lipid components.

- [1476] Lipid compositions are prepared by combining a lipid according to Formula (I), such as Compound 18 or a lipid according to Formula (III) such as Compound 236, a phospholipid (such as DOPE or DSPC, obtainable from Avanti Polar Lipids, Alabaster, AL), a PEG lipid (such as 1,2-dimyristoyl-*sn*-glycerol methoxypolyethylene glycol, also known as PEG-DMG, obtainable from Avanti Polar Lipids, Alabaster, AL or Compound 428), and a structural lipid (such as cholesterol, obtainable from Sigma-Aldrich, Taufkirchen, Germany, or a corticosteroid (such as prednisolone, dexamethasone, prednisone, and hydrocortisone), or a combination thereof) at concentrations of about 50 mM in ethanol. Solutions should be refrigerated for storage at, for example, -20° C. Lipids are combined to yield desired molar ratios and diluted with water and ethanol to a final lipid concentration of between about 5.5 mM and about 25 mM.
- [1477] Nanoparticle compositions including a polynucleotide and a lipid composition are prepared by combining the lipid solution with a solution including the a polynucleotide at lipid composition to polynucleotide wt:wt ratios between about 5:1 and about 50:1. The lipid solution is rapidly injected using a NanoAssemblr microfluidic based system at flow rates between about 10 ml/min and about 18 ml/min into the polynucleotide solution to produce a suspension with a water to ethanol ratio between about 1:1 and about 4:1.
- [1478] For nanoparticle compositions including an RNA, solutions of the RNA at concentrations of 0.1 mg/ml in deionized water are diluted in 50 mM sodium citrate buffer at a pH between 3 and 4 to form a stock solution.
- [1479] Nanoparticle compositions can be processed by dialysis to remove ethanol and achieve buffer exchange. Formulations are dialyzed twice against phosphate buffered saline (PBS), pH 7.4, at volumes 200 times that of the primary product using Slide-A-Lyzer cassettes (Thermo Fisher Scientific Inc., Rockford, IL) with a molecular weight cutoff of 10 kD. The first dialysis is carried out at room temperature for 3 hours. The formulations are then dialyzed overnight at 4° C. The resulting nanoparticle suspension is filtered through 0.2 µm sterile filters (Sarstedt, Nümbrecht, Germany) into glass vials and sealed with crimp closures. Nanoparticle composition solutions of 0.01 mg/ml to 0.10 mg/ml are generally obtained.
- [1480] The method described above induces nano-precipitation and particle formation. Alternative processes including, but not limited to, T-junction and direct injection, can be used to achieve the same nano-precipitation.

B. Characterization of nanoparticle compositions

- [1481] A Zetasizer Nano ZS (Malvern Instruments Ltd, Malvern, Worcestershire, UK) can be used to determine the particle size, the polydispersity index (PDI) and the zeta potential of the nanoparticle compositions in 1×PBS in determining particle size and 15 mM PBS in determining zeta potential.
- [1482] Ultraviolet-visible spectroscopy can be used to determine the concentration of a polynucleotide (*e.g.*, RNA) in nanoparticle compositions. 100 µL of the diluted formulation in 1×PBS is added to 900 µL of a 4:1 (v/v) mixture of methanol and chloroform. After mixing, the absorbance spectrum of the solution is recorded, for example, between 230 nm and 330 nm on a DU 800 spectrophotometer (Beckman Coulter, Beckman Coulter, Inc., Brea, CA). The concentration of polynucleotide in the nanoparticle composition can be calculated based on the extinction coefficient of the polynucleotide used in the composition and on the difference between the absorbance at a wavelength of, for example, 260 nm and the baseline value at a wavelength of, for example, 330 nm.
- [1483] For nanoparticle compositions including an RNA, a QUANT-IT™ RIBOGREEN® RNA assay (Invitrogen Corporation Carlsbad, CA) can be used to evaluate the encapsulation of an RNA by the nanoparticle composition. The samples are diluted to a concentration of approximately 5 µg/mL in a TE buffer solution (10 mM Tris-HCl, 1 mM EDTA, pH 7.5). 50 µL of the diluted samples are transferred to a polystyrene 96 well plate and either 50 µL of TE buffer or 50 µL of a 2% Triton X-100 solution is added to the wells. The plate is incubated at a temperature of 37° C for 15 minutes. The RIBOGREEN® reagent is diluted 1:100 in TE buffer, and 100 µL of this solution is added to each well. The fluorescence intensity can be measured using a fluorescence plate reader (Wallac Victor 1420 Multilabel Counter; Perkin Elmer, Waltham, MA) at an excitation wavelength of, for example, about 480 nm and an emission wavelength of, for example, about 520 nm. The fluorescence values of the reagent blank are subtracted from that of each of the samples and the percentage of free RNA is determined by dividing the fluorescence intensity of the intact sample (without addition of Triton X-100) by the fluorescence value of the disrupted sample (caused by the addition of Triton X-100).
- [1484] Exemplary formulations of the nanoparticle compositions are presented in the TABLE 5 below. The term "Compound" refers to an ionizable lipid such as MC3,

Compound 18, or Compound 236. "Phospholipid" can be DSPC or DOPE. "PEG-lipid" can be PEG-DMG or Compound 428.

TABLE 5. Exemplary Formulations of Nanoparticles

Composition (mol %)	Components
40:20:38.5:1.5	Compound:Phospholipid:Chol:PEG-lipid
45:15:38.5:1.5	Compound:Phospholipid:Chol:PEG-lipid
50:10:38.5:1.5	Compound:Phospholipid:Chol:PEG-lipid
55:5:38.5:1.5	Compound:Phospholipid:Chol:PEG-lipid
60:5:33.5:1.5	Compound:Phospholipid:Chol:PEG-lipid
45:20:33.5:1.5	Compound:Phospholipid:Chol:PEG-lipid
50:20:28.5:1.5	Compound:Phospholipid:Chol:PEG-lipid
55:20:23.5:1.5	Compound:Phospholipid:Chol:PEG-lipid
60:20:18.5:1.5	Compound:Phospholipid:Chol:PEG-lipid
40:15:43.5:1.5	Compound:Phospholipid:Chol:PEG-lipid
50:15:33.5:1.5	Compound:Phospholipid:Chol:PEG-lipid
55:15:28.5:1.5	Compound:Phospholipid:Chol:PEG-lipid
60:15:23.5:1.5	Compound:Phospholipid:Chol:PEG-lipid
40:10:48.5:1.5	Compound:Phospholipid:Chol:PEG-lipid
45:10:43.5:1.5	Compound:Phospholipid:Chol:PEG-lipid
55:10:33.5:1.5	Compound:Phospholipid:Chol:PEG-lipid
60:10:28.5:1.5	Compound:Phospholipid:Chol:PEG-lipid
40:5:53.5:1.5	Compound:Phospholipid:Chol:PEG-lipid
45:5:48.5:1.5	Compound:Phospholipid:Chol:PEG-lipid
50:5:43.5:1.5	Compound:Phospholipid:Chol:PEG-lipid
40:20:40:0	Compound:Phospholipid:Chol:PEG-lipid
45:20:35:0	Compound:Phospholipid:Chol:PEG-lipid
50:20:30:0	Compound:Phospholipid:Chol:PEG-lipid
55:20:25:0	Compound:Phospholipid:Chol:PEG-lipid
60:20:20:0	Compound:Phospholipid:Chol:PEG-lipid
40:15:45:0	Compound:Phospholipid:Chol:PEG-lipid
45:15:40:0	Compound:Phospholipid:Chol:PEG-lipid
50:15:35:0	Compound:Phospholipid:Chol:PEG-lipid
55:15:30:0	Compound:Phospholipid:Chol:PEG-lipid
60:15:25:0	Compound:Phospholipid:Chol:PEG-lipid
40:10:50:0	Compound:Phospholipid:Chol:PEG-lipid
45:10:45:0	Compound:Phospholipid:Chol:PEG-lipid
50:10:40:0	Compound:Phospholipid:Chol:PEG-lipid
55:10:35:0	Compound:Phospholipid:Chol:PEG-lipid
60:10:30:0	Compound:Phospholipid:Chol:PEG-lipid

[1485] It is to be appreciated that the Detailed Description section, and not the Summary and Abstract sections, is intended to be used to interpret the claims. The Summary and

Abstract sections can set forth one or more but not all exemplary embodiments of the present invention as contemplated by the inventor(s), and thus, are not intended to limit the present invention and the appended claims in any way.

[1486] The present invention has been described above with the aid of functional building blocks illustrating the implementation of specified functions and relationships thereof. The boundaries of these functional building blocks have been arbitrarily defined herein for the convenience of the description. Alternate boundaries can be defined so long as the specified functions and relationships thereof are appropriately performed.

[1487] The foregoing description of the specific embodiments will so fully reveal the general nature of the invention that others can, by applying knowledge within the skill of the art, readily modify and/or adapt for various applications such specific embodiments, without undue experimentation, without departing from the general concept of the present invention. Therefore, such adaptations and modifications are intended to be within the meaning and range of equivalents of the disclosed embodiments, based on the teaching and guidance presented herein. It is to be understood that the phraseology or terminology herein is for the purpose of description and not of limitation, such that the terminology or phraseology of the present specification is to be interpreted by the skilled artisan in light of the teachings and guidance.

[1488] The breadth and scope of the present invention should not be limited by any of the above-described exemplary embodiments, but should be defined only in accordance with the following claims and their equivalents.

[1489] All publications, patent applications, patents, and other references mentioned herein are incorporated by reference in their entirety. In case of conflict, the present specification, including definitions, will control.

WHAT IS CLAIMED IS:

1. A pharmaceutical composition comprising a lipid nanoparticle encapsulated mRNA that comprises an open reading frame (ORF) encoding an acyl-CoA dehydrogenase, very long-chain (ACADVL) polypeptide, wherein the composition is suitable for administration to a human subject in need of treatment for very long-chain acyl-CoA dehydrogenase deficiency (VLCADD).

2. A pharmaceutical composition comprising:

(a) an mRNA that comprises (i) an open reading frame (ORF) encoding an acyl-CoA dehydrogenase, very long-chain (ACADVL) polypeptide, wherein the ORF comprises at least one chemically modified nucleobase, sugar, backbone, or any combination thereof and (ii) an untranslated region (UTR) comprising a microRNA (miRNA) binding site; and

(b) a delivery agent,

wherein the pharmaceutical composition is suitable for administration to a human subject in need of treatment for VLCADD.

3. A pharmaceutical composition comprising an mRNA that comprises an open reading frame (ORF) encoding a human acyl-CoA dehydrogenase, very long-chain (ACADVL) polypeptide, wherein the composition when administered as a single intravenous dose to a human subject in need thereof is sufficient to (i) increase plasma ACADVL activity level to a level at or above a reference physiologic plasma level for at least 24 hours post-administration, and/or (ii) maintain plasma ACADVL activity level at 5%, 10%, 15%, 20%, 25%, 30%, 40%, 50%, or more of a reference plasma ACADVL activity level for at least 24 hours post-administration.

4. A pharmaceutical composition comprising an mRNA comprising an open reading frame (ORF) encoding a human ACADVL polypeptide, wherein the composition when administered as a single intravenous dose to a subject in need thereof is sufficient to reduce blood and/or plasma levels of:

(i) an acylcarnitine by at least 5%, at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or 100% compared to the subject's baseline blood and/or plasma acylcarnitine level or a reference

acylcarnitine blood and/or plasma level, for at least 24 hours, at least 48 hours, at least 72 hours, at least 96 hours, or at least 120 hours post-administration.

5. The pharmaceutical composition of claim 4, wherein the acylcarnitine is an acylcarnitine metabolite selected from the group consisting of C12:1 acylcarnitine, C14:1 acylcarnitine, C14:2 acylcarnitine, C14 acylcarnitine, C16 acylcarnitine, C18 acylcarnitine, C18:1 acylcarnitine, and combinations thereof.

6. A pharmaceutical composition comprising an mRNA comprising an open reading frame (ORF) encoding a human ACADVL polypeptide, wherein the composition when administered as a single intravenous dose to a subject in need thereof is sufficient to reduce blood and/or plasma levels of:

(i) C14:1 acylcarnitine by at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95% or at least 100% compared to the subject's baseline or a reference C14:1 acylcarnitine blood and/or plasma level, for at least 24 hours, at least 48 hours, at least 72 hours, at least 96 hours, or at least 120 hours post-administration;

(ii) C14:2 acylcarnitine by at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95% or at least 100% compared to the subject's baseline or a reference C14:2 acylcarnitine blood and/or plasma level, for at least 24 hours, at least 48 hours, at least 72 hours, at least 96 hours, or at least 120 hours post-administration;

(iii) C14 acylcarnitine by at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95% or at least 100% compared to the subject's baseline or a reference C14 acylcarnitine blood and/or plasma level, for at least 24 hours, at least 48 hours, at least 72 hours, at least 96 hours, or at least 120 hours post-administration; and/or

(iv) C12:1 acylcarnitine by at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95% or at least 100% compared to a subject's baseline or a reference C12:1 acylcarnitine blood and/or plasma level, for at least 24 hours, at least 48 hours, at least 72 hours, at least 96 hours, or at least 120 hours post-administration.

7. A pharmaceutical composition comprising an mRNA comprising an open reading frame (ORF) encoding a human ACADVL polypeptide, wherein the composition when administered as a single intravenous dose to a subject in need thereof is sufficient to reduce blood and/or plasma levels of an acylcarnitine to a concentration of less than 1.0 $\mu\text{mol/L}$ or less than 0.8 $\mu\text{mol/L}$ in a subject having very long-chain acyl-CoA dehydrogenase deficiency (VLCADD), for at least 24 hours, at least 48 hours, at least 72 hours, at least 96 hours, or at least 120 hours post-administration.

8. The pharmaceutical composition of claim 7, wherein the acylcarnitine is an acylcarnitine metabolite selected from the group consisting of C12:1 acylcarnitine, C14:1 acylcarnitine, C14:2 acylcarnitine, C14 acylcarnitine, C16 acylcarnitine, C18 acylcarnitine, C18:1 acylcarnitine, and combinations thereof.

9. The pharmaceutical composition of any one of claims 1 or 3-8 further comprising a delivery agent.

10. A polynucleotide comprising an open reading frame (ORF) encoding an acyl-CoA dehydrogenase, very long-chain (ACADVL) polypeptide, wherein the uracil or thymine content of the ORF relative to the theoretical minimum uracil or thymine content of a nucleotide sequence encoding the ACADVL polypeptide (%U_{TM} or %T_{TM}), is between about 100% and about 150%.

11. The polynucleotide of claim 10, wherein the %U_{TM} or %T_{TM} is between about 105% and about 140%, about 110% and about 140%, 115% and about 140%, about 105% and about 130%, about 110% and about 130%, about 115% and about 135%, about 105% and about 135%, about 110% and about 135%, or about 115% and about 130%.

12. The polynucleotide of claim 10 or 11, wherein the uracil or thymine content of the ORF relative to the uracil or thymine content of the corresponding wild-type ORF (%U_{WT} or %T_{WT}) is less than 100%.

13. The polynucleotide of claim 12, wherein the %U_{WT} or %T_{WT} is less than about 95%, less than about 90%, less than about 85%, less than 80%, less than 79%, less than 78%, less than 77%, less than 76%, less than 75%, or less than 74%.

14. The polynucleotide of any one of claims 10-13, wherein the uracil or thymine content in the ORF relative to the total nucleotide content in the ORF (%U_{TL} or %T_{TL}) is less than about 50%, less than about 40%, less than about 30%, or less than about 20%.

15. The polynucleotide of claim 14, wherein the %U_{TL} or %T_{TL} is less than about 16%.

16. The polynucleotide of any one of claims 10-15, wherein the guanine content of the ORF with respect to the theoretical maximum guanine content of a nucleotide sequence encoding the ACADVL polypeptide (%G_{TMX}) is at least 70%, at least 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, or about 100%.

17. The polynucleotide of claim 16, wherein the %G_{TMX} is between about 70% and about 80%, between about 71% and about 79%, between about 71% and about 77%, or between about 72% and about 76%.

18. The polynucleotide of any one of claims 10-17, wherein the cytosine content of the ORF relative to the theoretical maximum cytosine content of a nucleotide sequence encoding the ACADVL polypeptide (%C_{TMX}) is at least 58%, at least 59%, at least 60%, at least 65%, at least 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, or about 100%.

19. The polynucleotide of claim 18, wherein the %C_{TMX} is between about 60% and about 80%, between about 62% and about 77%, between about 63% and about 78%, or between about 68% and about 75%.

20. The polynucleotide of any one of claims 10-19, wherein the guanine and cytosine content (G/C) of the ORF relative to the theoretical maximum G/C content in a nucleotide

sequence encoding the ACADVL polypeptide (%G/C_{TMX}) is at least about 82%, at least 83%, at least 84%, at least 85%, at least 90%, at least about 95%, or about 100%.

21. The polynucleotide of claim 20, wherein the %G/C_{TMX} is between about 80% and about 100%, between about 85% and about 99%, between about 90% and about 96%, or between about 92% and about 95%.

22. The polynucleotide of any one of claims 10-21, wherein the G/C content in the ORF relative to the G/C content in the corresponding wild-type ORF (%G/C_{WT}) is at least 102%, at least 103%, at least 104%, at least 105%, at least 106%, at least 107%, at least about 110%, or at least about 112%.

23. The polynucleotide of any one of claims 10-22, wherein the ORF has at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to a nucleic acid sequence selected from the group consisting of SEQ ID NOs: 7 to 31.

24. The polynucleotide of any one of claims 10-23 which is an mRNA.

25. A pharmaceutical composition comprising the mRNA of claim 24 and a delivery agent.


26. The pharmaceutical composition of any one of claims 2, 9, and 25, wherein the delivery agent comprises a lipidoid, a liposome, a lipoplex, a lipid nanoparticle, a polymeric compound, a peptide, a protein, a cell, a nanoparticle mimic, a nanotube, or a conjugate.

27. The pharmaceutical composition of any one of claims 2, 9, 25, and 26, wherein the lipid nanoparticle or the delivery agent comprises a lipid selected from the group consisting of 3-(didodecylamino)-N1,N1,4-tridodecyl-1-piperazineethanamine (KL10), N1-[2-(didodecylamino)ethyl]-N1,N4,N4-tridodecyl-1,4-piperazinediethanamine (KL22), 14,25-ditridecyl-15,18,21,24-tetraaza-octatriacontane (KL25), 1,2-dilinoleyloxy-N,N-dimethylaminopropane (DLin-DMA), 2,2-dilinoleyloxy-4-dimethylaminomethyl-[1,3]-dioxolane (DLin-K-DMA),

heptatriaconta-6,9,28,31-tetraen-19-yl 4-(dimethylamino)butanoate (DLin-MC3-DMA), 2,2-dilinoleyl-4-(2-dimethylaminoethyl)-[1,3]-dioxolane (DLin-KC2-DMA), 1,2-dioleyloxy-N,N-dimethylaminopropane (DODMA), (13Z,165Z)-N,N-dimethyl-3-nonydocosa-13-16-dien-1-amine (L608), 2-({8-[(3 β)-cholest-5-en-3-yloxy]octyl}oxy)-N,N-dimethyl-3-[(9Z,12Z)-octadeca-9,12-dien-1-yl oxy]propan-1-amine (Octyl-CLinDMA), (2R)-2-({8-[(3 β)-cholest-5-en-3-yloxy]octyl}oxy)-N,N-dimethyl-3-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]propan-1-amine (Octyl-CLinDMA (2R)), (2S)-2-({8-[(3 β)-cholest-5-en-3-yloxy]octyl}oxy)-N,N-dimethyl-3-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]propan-1-amine (Octyl-CLinDMA (2S)), and any combination thereof.

28. The pharmaceutical composition of claim 27, wherein the lipid nanoparticle comprises DLin-MC3-DMA.

29. The pharmaceutical composition of any one of claims 2, 9, 25, and 26, wherein the lipid nanoparticle or the delivery agent comprises a compound having the Formula (I)



(I),

or a salt or stereoisomer thereof, wherein

R₁ is selected from the group consisting of C₅₋₃₀ alkyl, C₅₋₂₀ alkenyl, -R*YR'', -YR'', and -R''M'R';

R₂ and R₃ are independently selected from the group consisting of H, C₁₋₁₄ alkyl, C₂₋₁₄ alkenyl, -R*YR'', -YR'', and -R*OR'', or R₂ and R₃, together with the atom to which they are attached, form a heterocycle or carbocycle;

R₄ is selected from the group consisting of a C₃₋₆ carbocycle, -(CH₂)_nQ, -(CH₂)_nCHQR, -CHQR, -CQ(R)₂, and unsubstituted C₁₋₆ alkyl, where Q is selected from a carbocycle, heterocycle, -OR, -O(CH₂)_nN(R)₂, -C(O)OR, -OC(O)R, -CX₃, -CX₂H, -CXH₂, -CN, -N(R)₂, -C(O)N(R)₂, -N(R)C(O)R, -N(R)S(O)₂R, -N(R)C(O)N(R)₂, -N(R)C(S)N(R)₂, -N(R)R₈, -O(CH₂)_nOR, -N(R)C(=NR₉)N(R)₂, -N(R)C(=CHR₉)N(R)₂, -OC(O)N(R)₂, -N(R)C(O)OR, -N(OR)C(O)R, -N(OR)S(O)₂R, -N(OR)C(O)OR, -N(OR)C(O)N(R)₂, -N(OR)C(S)N(R)₂,

$-\text{N}(\text{OR})\text{C}(=\text{NR}_9)\text{N}(\text{R})_2$, $-\text{N}(\text{OR})\text{C}(=\text{CHR}_9)\text{N}(\text{R})_2$, $-\text{C}(=\text{NR}_9)\text{N}(\text{R})_2$, $-\text{C}(=\text{NR}_9)\text{R}$, $-\text{C}(\text{O})\text{N}(\text{R})\text{OR}$, and $-\text{C}(\text{R})\text{N}(\text{R})_2\text{C}(\text{O})\text{OR}$, and each n is independently selected from 1, 2, 3, 4, and 5;

each R_5 is independently selected from the group consisting of C_{1-3} alkyl, C_{2-3} alkenyl, and H;

each R_6 is independently selected from the group consisting of C_{1-3} alkyl, C_{2-3} alkenyl, and H;

M and M' are independently selected from $-\text{C}(\text{O})\text{O}-$, $-\text{OC}(\text{O})-$, $-\text{C}(\text{O})\text{N}(\text{R}')$, $-\text{N}(\text{R}')\text{C}(\text{O})-$, $-\text{C}(\text{O})-$, $-\text{C}(\text{S})-$, $-\text{C}(\text{S})\text{S}-$, $-\text{SC}(\text{S})-$, $-\text{CH}(\text{OH})-$, $-\text{P}(\text{O})(\text{OR}')\text{O}-$, $-\text{S}(\text{O})_2-$, $-\text{S}-\text{S}-$, an aryl group, and a heteroaryl group;

R_7 is selected from the group consisting of C_{1-3} alkyl, C_{2-3} alkenyl, and H;

R_8 is selected from the group consisting of C_{3-6} carbocycle and heterocycle;

R_9 is selected from the group consisting of H, CN, NO_2 , C_{1-6} alkyl, $-\text{OR}$, $-\text{S}(\text{O})_2\text{R}$, $-\text{S}(\text{O})_2\text{N}(\text{R})_2$, C_{2-6} alkenyl, C_{3-6} carbocycle and heterocycle;

each R is independently selected from the group consisting of C_{1-3} alkyl, C_{2-3} alkenyl, and H;

each R' is independently selected from the group consisting of C_{1-18} alkyl, C_{2-18} alkenyl, $-\text{R}^*\text{YR}''$, $-\text{YR}''$, and H;

each R'' is independently selected from the group consisting of C_{3-14} alkyl and C_{3-14} alkenyl;


each R^* is independently selected from the group consisting of C_{1-12} alkyl and C_{2-12} alkenyl;

each Y is independently a C_{3-6} carbocycle;

each X is independently selected from the group consisting of F, Cl, Br, and I; and m is selected from 5, 6, 7, 8, 9, 10, 11, 12, and 13; and

provided that when R_4 is $-(\text{CH}_2)_n\text{Q}$, $-(\text{CH}_2)_n\text{CHQR}$, $-\text{CHQR}$, or $-\text{CQ}(\text{R})_2$, then (i) Q is not $-\text{N}(\text{R})_2$ when n is 1, 2, 3, 4 or 5, or (ii) Q is not 5, 6, or 7-membered heterocycloalkyl when n is 1 or 2.

30. The pharmaceutical composition of claim 29, wherein the lipid nanoparticle or the delivery agent comprises the compound is of Formula (IA):



(IA),

or a salt or stereoisomer thereof, wherein

l is selected from 1, 2, 3, 4, and 5;

m is selected from 5, 6, 7, 8, and 9;

M₁ is a bond or M';


R₄ is unsubstituted C₁₋₃ alkyl, or -(CH₂)_nQ, in which Q is OH, -NHC(S)N(R)₂, -NHC(O)N(R)₂, -N(R)C(O)R, -N(R)S(O)₂R, -N(R)R₈, -NHC(=NR₉)N(R)₂, -NHC(=CHR₉)N(R)₂, -OC(O)N(R)₂, -N(R)C(O)OR, heteroaryl or heterocycloalkyl;

M and M' are independently selected from -C(O)O-, -OC(O)-, -C(O)N(R')-, -P(O)(OR')O-, -S-S-, an aryl group, and a heteroaryl group; and

R₂ and R₃ are independently selected from the group consisting of H, C₁₋₁₄ alkyl, and C₂₋₁₄ alkenyl.

31. The pharmaceutical composition of any one of claims 29 to 30, wherein m is 5, 7, or 9.

32. The pharmaceutical composition of any one of claims 29 to 30, wherein the compound is of Formula (II)



(II)

or a salt or stereoisomer thereof, wherein

l is selected from 1, 2, 3, 4, and 5;


M₁ is a bond or M';

R₄ is unsubstituted C₁₋₃ alkyl, or -(CH₂)_nQ, in which n is 2, 3, or 4, and Q is OH, -NHC(S)N(R)₂, -NHC(O)N(R)₂, -N(R)C(O)R, -N(R)S(O)₂R, -N(R)R₈, -NHC(=NR₉)N(R)₂, -NHC(=CHR₉)N(R)₂, -OC(O)N(R)₂, -N(R)C(O)OR, heteroaryl, or heterocycloalkyl;

M and M' are independently selected from -C(O)O-, -OC(O)-, -C(O)N(R')-, -P(O)(OR')O-, -S-S-, an aryl group, and a heteroaryl group; and


R_2 and R_3 are independently selected from the group consisting of H, C_{1-14} alkyl, and C_{2-14} alkenyl.

33. The pharmaceutical composition of any one of claims 30 to 32, wherein M_1 is M' .
34. The pharmaceutical composition of claim 32, wherein M and M' are independently $-C(O)O-$ or $-OC(O)-$.
35. The pharmaceutical composition of any one of claims 30 to 34, wherein l is 1, 3, or 5.
36. The pharmaceutical composition of claim 29, wherein the compound is selected from the group consisting of Compound 1 to Compound 232, salts and stereoisomers thereof, and any combination thereof.
37. The pharmaceutical composition of claim 36, wherein the compound is selected from the group consisting of Compound 1 to Compound 147, salts and stereoisomers thereof, and any combination thereof.
38. The pharmaceutical composition of claim 37, wherein the compound is Compound 18, a salt or a stereoisomer thereof, or any combination thereof.
39. The pharmaceutical composition of any one of claims 2, 9, 25, and 26, wherein the lipid nanoparticle or the delivery agent comprises a compound having the Formula (III)



(III),

or salts or stereoisomers thereof, wherein



t is 1 or 2;

A₁ and A₂ are each independently selected from CH or N;

Z is CH₂ or absent wherein when Z is CH₂, the dashed lines (1) and (2) each represent a single bond; and when Z is absent, the dashed lines (1) and (2) are both absent;

R₁, R₂, R₃, R₄, and R₅ are independently selected from the group consisting of C₅₋₂₀ alkyl, C₅₋₂₀ alkenyl, -R''MR', -R*YR'', -YR'', and -R*OR'';

each M is independently selected from the group consisting of -C(O)O-, -OC(O)-, -OC(O)O-, -C(O)N(R')-, -N(R')C(O)-, -C(O)-, -C(S)-, -C(S)S-, -SC(S)-, -CH(OH)-, -P(O)(OR')O-, -S(O)₂-, an aryl group, and a heteroaryl group;

X¹, X², and X³ are independently selected from the group consisting of a bond, -CH₂-, -CH₂)₂-, -CHR-, -CHY-, -C(O)-, -C(O)O-, -OC(O)-, -C(O)-CH₂-, -CH₂-C(O)-, -C(O)O-CH₂-, -OC(O)-CH₂-, -CH₂-C(O)O-, -CH₂-OC(O)-, -CH(OH)-, -C(S)-, and -CH(SH)-;


each Y is independently a C₃₋₆ carbocycle;

each R* is independently selected from the group consisting of C₁₋₁₂ alkyl and C₂₋₁₂ alkenyl;

each R is independently selected from the group consisting of C₁₋₃ alkyl and a C₃₋₆ carbocycle;

each R' is independently selected from the group consisting of C₁₋₁₂ alkyl, C₂₋₁₂ alkenyl, and H; and







each R'' is independently selected from the group consisting of C₃₋₁₂ alkyl and C₃₋₁₂ alkenyl,




wherein when ring A is , then

- i) at least one of X¹, X², and X³ is not -CH₂-; and/or
- ii) at least one of R₁, R₂, R₃, R₄, and R₅ is -R''MR'.

40. The pharmaceutical composition of claim 39, wherein the compound has the formula:



41. The pharmaceutical composition of any one of claims 2, 9, 25, and 26, wherein the lipid nanoparticle or the delivery agent comprises a compound having the Formula (IV)




(IV),


or salts or stereoisomer thereof, wherein

A₁ and A₂ are each independently selected from CH or N and at least one of A₁ and A₂ is N;

Z is CH₂ or absent wherein when Z is CH₂, the dashed lines (1) and (2) each represent a single bond; and when Z is absent, the dashed lines (1) and (2) are both absent;

R₁, R₂, R₃, R₄, and R₅ are independently selected from the group consisting of C₆₋₂₀ alkyl and C₆₋₂₀ alkenyl;



wherein when ring A is , then

i) R₁, R₂, R₃, R₄, and R₅ are the same, wherein R₁ is not C₁₂ alkyl, C₁₈ alkyl, or C₁₈ alkenyl;


ii) only one of R₁, R₂, R₃, R₄, and R₅ is selected from C₆₋₂₀ alkenyl;

iii) at least one of R₁, R₂, R₃, R₄, and R₅ have a different number of carbon atoms than at least one other of R₁, R₂, R₃, R₄, and R₅;

iv) R₁, R₂, and R₃ are selected from C₆₋₂₀ alkenyl, and R₄ and R₅ are selected from C₆₋₂₀ alkyl; or


v) R₁, R₂, and R₃ are selected from C₆₋₂₀ alkyl, and R₄ and R₅ are selected from C₆₋₂₀ alkenyl.

42. The pharmaceutical composition of claim 41, wherein the compound is of Formula (IVa):



(IVa).

43. The pharmaceutical composition of any one of claims 2, 9, 25, and 26, wherein the lipid nanoparticle or the delivery agent comprises a compound having the Formula (V)



(V),

or salts or stereoisomers thereof, in which

A₃ is CH or N;

A₄ is CH₂ or NH; and at least one of A₃ and A₄ is N or NH;

Z is CH₂ or absent wherein when Z is CH₂, the dashed lines (1) and (2) each represent a single bond; and when Z is absent, the dashed lines (1) and (2) are both absent;

R₁, R₂, and R₃ are independently selected from the group consisting of C₅₋₂₀ alkyl, C₅₋₂₀ alkenyl, -R''MR', -R*YR'', -YR'', and -R*OR'';

each M is independently selected from -C(O)O-, -OC(O)-, -C(O)N(R')-, -N(R')C(O)-, -C(O)-, -C(S)-, -C(S)S-, -SC(S)-, -CH(OH)-, -P(O)(OR')O-, -S(O)₂-, an aryl group, and a heteroaryl group;

X¹ and X² are independently selected from the group consisting of -CH₂-, -(CH₂)₂-, -CHR-, -CHY-, -C(O)-, -C(O)O-, -OC(O)-, -C(O)-CH₂-, -CH₂-C(O)-, -C(O)O-CH₂-, -OC(O)-CH₂-, -CH₂-C(O)O-, -CH₂-OC(O)-, -CH(OH)-, -C(S) - and -CH(SH) -

each Y is independently a C₃₋₆ carbocycle;


each R* is independently selected from the group consisting of C₁₋₁₂ alkyl and C₂₋₁₂ alkenyl;

each R is independently selected from the group consisting of C₁₋₃ alkyl and a C₃₋₆ carbocycle;


each R' is independently selected from the group consisting of C₁₋₁₂ alkyl, C₂₋₁₂ alkenyl, and H; and

each R'' is independently selected from the group consisting of C₃₋₁₂ alkyl and C₃₋₁₂ alkenyl.

44. The pharmaceutical composition of claim 43, wherein the compound is of Formula (Va):



45. The pharmaceutical composition of any one of claims 2, 9, 25, and 26, wherein the lipid nanoparticle or the delivery agent comprises a compound having the Formula (VI):



or salts or stereoisomers thereof, in which

A₆ and A₇ are each independently selected from CH or N, wherein at least one of A₆ and A₇ is N;

Z is CH₂ or absent wherein when Z is CH₂, the dashed lines (1) and (2) each represent a single bond; and when Z is absent, the dashed lines (1) and (2) are both absent;

X⁴ and X⁵ are independently selected from the group consisting of -CH₂-, -(CH₂)₂-, -CHR-, -CHY-, -C(O)-, -C(O)O-, -OC(O)-, -C(O)-CH₂-, -CH₂-C(O)-, -C(O)O-CH₂-, -OC(O)-CH₂-, -CH₂-C(O)O-, -CH₂-OC(O)-, -CH(OH)-, -C(S)-, and -CH(SH)-;

R₁, R₂, R₃, R₄, and R₅ each are independently selected from the group consisting of C₅₋₂₀ alkyl, C₅₋₂₀ alkenyl, -R''MR', -R''YR'', -YR'', and -R''OR'';

each M is independently selected from the group consisting of -C(O)O-, -OC(O)-, -C(O)N(R')-, -N(R')C(O)-, -C(O)-, -C(S)-, -C(S)S-, -SC(S)-, -CH(OH)-, -P(O)(OR')O-, -S(O)₂-, an aryl group, and a heteroaryl group;

each Y is independently a C₃₋₆ carbocycle;

each R* is independently selected from the group consisting of C₁₋₁₂ alkyl and C₂₋₁₂ alkenyl;


each R is independently selected from the group consisting of C₁₋₃ alkyl and a C₃₋₆ carbocycle;

each R' is independently selected from the group consisting of C₁₋₁₂ alkyl, C₂₋₁₂ alkenyl, and H; and each R'' is independently selected from the group consisting of C₃₋₁₂ alkyl and C₃₋₁₂ alkenyl.

46. The pharmaceutical composition of any one of claims 2, 9, 25, and 26, wherein the lipid nanoparticle or the delivery agent comprises a compound selected from the group consisting of Compound 233 to Compound 342, salts and stereoisomers thereof, and any combination thereof.

47. The pharmaceutical composition of claim 46, wherein the compound is Compound 236, a salt or a stereoisomer thereof, or any combination thereof.

48. A pharmaceutical composition comprising an mRNA comprising an open reading frame (ORF) encoding an acyl-CoA dehydrogenase, very long-chain (ACADVL) polypeptide and a delivery agent comprising a compound having the Formula (I)



(I),

or a salt or stereoisomer thereof, wherein

R₁ is selected from the group consisting of C₅₋₃₀ alkyl, C₅₋₂₀ alkenyl, -R*YR'', -YR'', and -R''M'R';

R₂ and R₃ are independently selected from the group consisting of H, C₁₋₁₄ alkyl, C₂₋₁₄ alkenyl, -R*YR'', -YR'', and -R*OR'', or R₂ and R₃, together with the atom to which they are attached, form a heterocycle or carbocycle;

R₄ is selected from the group consisting of a C₃₋₆ carbocycle, -(CH₂)_nQ, -(CH₂)_nCHQR, -CHQR, -CQ(R)₂, and unsubstituted C₁₋₆ alkyl, where Q is selected from a carbocycle, heterocycle, -OR, -O(CH₂)_nN(R)₂, -C(O)OR, -OC(O)R, -CX₃, -CX₂H, -CXH₂, -CN, -N(R)₂, -C(O)N(R)₂, -N(R)C(O)R, -N(R)S(O)₂R, -N(R)C(O)N(R)₂, -N(R)C(S)N(R)₂, -N(R)R₈,

-O(CH₂)_nOR, -N(R)C(=NR₉)N(R)₂, -N(R)C(=CHR₉)N(R)₂, -OC(O)N(R)₂, -N(R)C(O)OR, -N(OR)C(O)R, -N(OR)S(O)₂R, -N(OR)C(O)OR, -N(OR)C(O)N(R)₂, -N(OR)C(S)N(R)₂, -N(OR)C(=NR₉)N(R)₂, -N(OR)C(=CHR₉)N(R)₂, -C(=NR₉)N(R)₂, -C(=NR₉)R, -C(O)N(R)OR, and -C(R)N(R)₂C(O)OR, and each n is independently selected from 1, 2, 3, 4, and 5;

each R₅ is independently selected from the group consisting of C₁₋₃ alkyl, C₂₋₃ alkenyl, and H;

each R₆ is independently selected from the group consisting of C₁₋₃ alkyl, C₂₋₃ alkenyl, and H;

M and M' are independently selected from -C(O)O-, -OC(O)-, -C(O)N(R')-, -N(R')C(O)-, -C(O)-, -C(S)-, -C(S)S-, -SC(S)-, -CH(OH)-, -P(O)(OR')O-, -S(O)₂-, -S-S-, an aryl group, and a heteroaryl group;

R₇ is selected from the group consisting of C₁₋₃ alkyl, C₂₋₃ alkenyl, and H;

R₈ is selected from the group consisting of C₃₋₆ carbocycle and heterocycle;

R₉ is selected from the group consisting of H, CN, NO₂, C₁₋₆ alkyl, -OR, -S(O)₂R, -S(O)₂N(R)₂, C₂₋₆ alkenyl, C₃₋₆ carbocycle and heterocycle;

each R is independently selected from the group consisting of C₁₋₃ alkyl, C₂₋₃ alkenyl, and H;

each R' is independently selected from the group consisting of C₁₋₁₈ alkyl, C₂₋₁₈ alkenyl, -R*YR'', -YR'', and H;

each R'' is independently selected from the group consisting of C₃₋₁₄ alkyl and C₃₋₁₄ alkenyl;


each R* is independently selected from the group consisting of C₁₋₁₂ alkyl and C₂₋₁₂ alkenyl;

each Y is independently a C₃₋₆ carbocycle;

each X is independently selected from the group consisting of F, Cl, Br, and I; and m is selected from 5, 6, 7, 8, 9, 10, 11, 12, and 13; and

provided that when R₄ is -(CH₂)_nQ, -(CH₂)_nCHQR, -CHQR, or -CQ(R)₂, then (i) Q is not -N(R)₂ when n is 1, 2, 3, 4 or 5, or (ii) Q is not 5, 6, or 7-membered heterocycloalkyl when n is 1 or 2.

49. The pharmaceutical composition of claim 48, wherein the delivery agent comprises the compound is of Formula (IA):



(IA),

or a salt or stereoisomer thereof, wherein

l is selected from 1, 2, 3, 4, and 5;

m is selected from 5, 6, 7, 8, and 9;

M₁ is a bond or M';


R₄ is unsubstituted C₁₋₃ alkyl, or -(CH₂)_nQ, in which Q is OH, -NHC(S)N(R)₂, -NHC(O)N(R)₂, -N(R)C(O)R, -N(R)S(O)₂R, -N(R)R₈, -NHC(=NR₉)N(R)₂, -NHC(=CHR₉)N(R)₂, -OC(O)N(R)₂, -N(R)C(O)OR, heteroaryl or heterocycloalkyl;

M and M' are independently selected from -C(O)O-, -OC(O)-, -C(O)N(R')-, -P(O)(OR')O-, -S-S-, an aryl group, and a heteroaryl group; and

R₂ and R₃ are independently selected from the group consisting of H, C₁₋₁₄ alkyl, and C₂₋₁₄ alkenyl.

50. The pharmaceutical composition of any one of claims 48 to 49, wherein m is 5, 7, or 9.

51. The pharmaceutical composition of any one of claims 48 to 50, wherein the compound is of Formula (II)



(II)

or a salt or stereoisomer thereof, wherein

l is selected from 1, 2, 3, 4, and 5;

M₁ is a bond or M';

R₄ is unsubstituted C₁₋₃ alkyl, or -(CH₂)_nQ, in which n is 2, 3, or 4, and Q is OH, -NHC(S)N(R)₂, or -NHC(O)N(R)₂, -N(R)C(O)R, -N(R)S(O)₂R, -N(R)R₈, -NHC(=NR₉)N(R)₂, -NHC(=CHR₉)N(R)₂, -OC(O)N(R)₂, -N(R)C(O)OR, heteroaryl or heterocycloalkyl;

M and M' are independently selected from -C(O)O-, -OC(O)-, -C(O)N(R')-, -P(O)(OR')O-, -S-S-, an aryl group, and a heteroaryl group; and

R_2 and R_3 are independently selected from the group consisting of H, C_{1-14} alkyl, and C_{2-14} alkenyl.

52. The pharmaceutical composition of any one of claims 49 to 51, wherein M_1 is M' .

53. The pharmaceutical composition of claim 52, wherein M and M' are independently $-C(O)O-$ or $-OC(O)-$.


54. The pharmaceutical composition of any one of claims 49 to 53, wherein l is 1, 3, or 5.

55. The pharmaceutical composition of claim 48, wherein the compound is selected from the group consisting of Compound 1 to Compound 232, salts and stereoisomers thereof, and any combination thereof.

56. The pharmaceutical composition of claim 55, wherein the compound is selected from the group consisting of Compound 1 to Compound 147, salts and stereoisomers thereof, and any combination thereof.


57. The pharmaceutical composition of claim 56, wherein the compound is Compound 18, a salt or a stereoisomer thereof, or any combination thereof.

58. A pharmaceutical composition comprising an mRNA comprising an open reading frame (ORF) encoding an acyl-CoA dehydrogenase, very long-chain (ACADVL) polypeptide and a delivery agent, wherein the delivery agent comprises a compound having the Formula (III)



(III),

or salts or stereoisomers thereof, wherein



t is 1 or 2;

A₁ and A₂ are each independently selected from CH or N;

Z is CH₂ or absent wherein when Z is CH₂, the dashed lines (1) and (2) each represent a single bond; and when Z is absent, the dashed lines (1) and (2) are both absent;

R₁, R₂, R₃, R₄, and R₅ are independently selected from the group consisting of C₅₋₂₀ alkyl, C₅₋₂₀ alkenyl, -R''MR', -R*YR'', -YR'', and -R*OR'';

each M is independently selected from the group consisting of -C(O)O-, -OC(O)-, -OC(O)O-, -C(O)N(R')-, -N(R')C(O)-, -C(O)-, -C(S)-, -C(S)S-, -SC(S)-, -CH(OH)-, -P(O)(OR')O-, -S(O)₂-, an aryl group, and a heteroaryl group;

X¹, X², and X³ are independently selected from the group consisting of a bond, -CH₂-, -CH₂)₂-, -CHR-, -CHY-, -C(O)-, -C(O)O-, -OC(O)-, -C(O)-CH₂-, -CH₂-C(O)-, -C(O)O-CH₂-, -OC(O)-CH₂-, -CH₂-C(O)O-, -CH₂-OC(O)-, -CH(OH)-, -C(S)-, and -CH(SH)-;


each Y is independently a C₃₋₆ carbocycle;

each R* is independently selected from the group consisting of C₁₋₁₂ alkyl and C₂₋₁₂ alkenyl;

each R is independently selected from the group consisting of C₁₋₃ alkyl and a C₃₋₆ carbocycle;

each R' is independently selected from the group consisting of C₁₋₁₂ alkyl, C₂₋₁₂ alkenyl, and H; and







each R'' is independently selected from the group consisting of C₃₋₁₂ alkyl and C₃₋₁₂ alkenyl,



wherein when ring A is , then


- i) at least one of X¹, X², and X³ is not -CH₂-; and/or
- ii) at least one of R₁, R₂, R₃, R₄, and R₅ is -R''MR'.

59. The pharmaceutical composition of claim 58, wherein the compound has the formula:



60. The pharmaceutical composition comprising an mRNA comprising an open reading frame (ORF) encoding an acyl-CoA dehydrogenase, very long-chain (ACADVL)

polypeptide and a delivery agent, wherein the delivery agent comprises a compound having the Formula (IV)




(IV),

or salts or stereoisomer thereof, wherein

A₁ and A₂ are each independently selected from CH or N and at least one of A₁ and A₂ is N;

Z is CH₂ or absent wherein when Z is CH₂, the dashed lines (1) and (2) each represent a single bond; and when Z is absent, the dashed lines (1) and (2) are both absent;

R₁, R₂, R₃, R₄, and R₅ are independently selected from the group consisting of C₆₋₂₀ alkyl and C₆₋₂₀ alkenyl;



wherein when ring A is , then

i) R₁, R₂, R₃, R₄, and R₅ are the same, wherein R₁ is not C₁₂ alkyl, C₁₈ alkyl, or C₁₈ alkenyl;


ii) only one of R₁, R₂, R₃, R₄, and R₅ is selected from C₆₋₂₀ alkenyl;

iii) at least one of R₁, R₂, R₃, R₄, and R₅ have a different number of carbon atoms than at least one other of R₁, R₂, R₃, R₄, and R₅;

iv) R₁, R₂, and R₃ are selected from C₆₋₂₀ alkenyl, and R₄ and R₅ are selected from C₆₋₂₀ alkyl; or


v) R₁, R₂, and R₃ are selected from C₆₋₂₀ alkyl, and R₄ and R₅ are selected from C₆₋₂₀ alkenyl.

61. The pharmaceutical composition of claim 60, wherein the compound is of Formula (IVa):



(IVa).

62. A pharmaceutical composition comprising an mRNA comprising an open reading frame (ORF) encoding an acyl-CoA dehydrogenase, very long-chain (ACADVL) polypeptide and a delivery agent, wherein the delivery agent comprises a compound having the Formula (V)



(V),

or salts or stereoisomers thereof, in which

A₃ is CH or N;

A₄ is CH₂ or NH; and at least one of A₃ and A₄ is N or NH;

Z is CH₂ or absent wherein when Z is CH₂, the dashed lines (1) and (2) each represent a single bond; and when Z is absent, the dashed lines (1) and (2) are both absent;

R₁, R₂, and R₃ are independently selected from the group consisting of C₅₋₂₀ alkyl, C₅₋₂₀ alkenyl, -R''MR', -R*YR'', -YR'', and -R*OR'';

each M is independently selected from -C(O)O-, -OC(O)-, -C(O)N(R')-, -N(R')C(O)-, -C(O)-, -C(S)-, -C(S)S-, -SC(S)-, -CH(OH)-, -P(O)(OR')O-, -S(O)₂-, an aryl group, and a heteroaryl group;

X¹ and X² are independently selected from the group consisting of -CH₂-, -(CH₂)₂-, -CHR-, -CHY-, -C(O)-, -C(O)O-, -OC(O)-, -C(O)-CH₂-, -CH₂-C(O)-, -C(O)O-CH₂-, -OC(O)-CH₂-, -CH₂-C(O)O-, -CH₂-OC(O)-, -CH(OH)-, -C(S) - and -CH(SH) -

each Y is independently a C₃₋₆ carbocycle;


each R* is independently selected from the group consisting of C₁₋₁₂ alkyl and C₂₋₁₂ alkenyl;

each R is independently selected from the group consisting of C₁₋₃ alkyl and a C₃₋₆ carbocycle;


each R' is independently selected from the group consisting of C₁₋₁₂ alkyl, C₂₋₁₂ alkenyl, and H; and

each R'' is independently selected from the group consisting of C₃₋₁₂ alkyl and C₃₋₁₂ alkenyl.

63. The pharmaceutical composition of claim 62, wherein the compound is of Formula (Va):



64. A pharmaceutical composition comprising an mRNA comprising an open reading frame (ORF) encoding an acyl-CoA dehydrogenase, very long-chain (ACADVL) polypeptide and a delivery agent, wherein the delivery agent comprises a compound having the Formula (VI):



or salts or stereoisomers thereof, in which

A₆ and A₇ are each independently selected from CH or N, wherein at least one of A₆ and A₇ is N;

Z is CH₂ or absent wherein when Z is CH₂, the dashed lines (1) and (2) each represent a single bond; and when Z is absent, the dashed lines (1) and (2) are both absent;

X⁴ and X⁵ are independently selected from the group consisting of -CH₂-, -(CH₂)₂-, -CHR-, -CHY-, -C(O)-, -C(O)O-, -OC(O)-, -C(O)-CH₂-, -CH₂-C(O)-, -C(O)O-CH₂-, -OC(O)-CH₂-, -CH₂-C(O)O-, -CH₂-OC(O)-, -CH(OH)-, -C(S)-, and -CH(SH)-;

R₁, R₂, R₃, R₄, and R₅ each are independently selected from the group consisting of C₅₋₂₀ alkyl, C₅₋₂₀ alkenyl, -R''MR', -R''YR'', -YR'', and -R''OR'';

each M is independently selected from the group consisting of -C(O)O-, -OC(O)-, -C(O)N(R')-, -N(R')C(O)-, -C(O)-, -C(S)-, -C(S)S-, -SC(S)-, -CH(OH)-, -P(O)(OR')O-, -S(O)₂-, an aryl group, and a heteroaryl group;

each Y is independently a C₃₋₆ carbocycle;

each R* is independently selected from the group consisting of C₁₋₁₂ alkyl and C₂₋₁₂ alkenyl;

each R is independently selected from the group consisting of C₁₋₃ alkyl and a C₃₋₆ carbocycle;

each R' is independently selected from the group consisting of C₁₋₁₂ alkyl, C₂₋₁₂ alkenyl, and H; and each R'' is independently selected from the group consisting of C₃₋₁₂ alkyl and C₃₋₁₂ alkenyl.

65. A pharmaceutical composition comprising an mRNA comprising an open reading frame (ORF) encoding an acyl-CoA dehydrogenase, very long-chain (ACADVL) polypeptide and a delivery agent, wherein the delivery agent comprises a compound selected from the group consisting of Compound 233 to Compound 342, salts and stereoisomers thereof, and any combination thereof.

66. The pharmaceutical composition of claim 65, wherein the compound is Compound 236, a salt or a stereoisomer thereof, or any combination thereof.

67. The pharmaceutical composition of any one of claims 25-66, wherein the lipid nanoparticle or the delivery agent further comprises a PEG lipid.

68. The pharmaceutical composition of claim 67, wherein the PEG lipid has the Formula (VII):



or a salt thereof, wherein:

R³ is -OR⁰;


R⁰ is hydrogen, optionally substituted alkyl, or an oxygen protecting group;

r is an integer between 1 and 100, inclusive;

L^1 is optionally substituted C_{1-10} alkylene, wherein at least one methylene of the optionally substituted C_{1-10} alkylene is independently replaced with optionally substituted carbocyclylene, optionally substituted heterocyclylene, optionally substituted arylene, optionally substituted heteroarylene, $-O-$, $-N(R^N)-$, $-S-$, $-C(O)-$, $-C(O)N(R^N)-$, $-NR^NC(O)-$, $-C(O)O-$, $-OC(O)-$, $-OC(O)O-$, $-OC(O)N(R^N)-$, $-NR^NC(O)O-$, or $-NR^NC(O)N(R^N)-$;

D is a moiety obtained by click chemistry or a moiety cleavable under physiological conditions;

m is 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10;




each instance of L^2 is independently a bond or optionally substituted C_{1-6} alkylene, wherein one methylene unit of the optionally substituted C_{1-6} alkylene is optionally replaced with $-O-$, $-N(R^N)-$, $-S-$, $-C(O)-$, $-C(O)N(R^N)-$, $-NR^NC(O)-$, $-C(O)O-$, $-OC(O)-$, $-OC(O)O-$, $-OC(O)N(R^N)-$, $-NR^NC(O)O-$, or $-NR^NC(O)N(R^N)-$;

each instance of R^2 is independently optionally substituted C_{1-30} alkyl, optionally substituted C_{1-30} alkenyl, or optionally substituted C_{1-30} alkynyl; optionally wherein one or more methylene units of R^2 are independently replaced with optionally substituted carbocyclylene, optionally substituted heterocyclylene, optionally substituted arylene, optionally substituted heteroarylene, $-N(R^N)-$, $-O-$, $-S-$, $-C(O)-$, $-C(O)N(R^N)-$, $-NR^NC(O)-$, $-NR^NC(O)N(R^N)-$, $-C(O)O-$, $-OC(O)-$, $-OC(O)O-$, $-OC(O)N(R^N)-$, $-NR^NC(O)O-$, $-C(O)S-$, $-SC(O)-$, $-C(=NR^N)-$, $-C(=NR^N)N(R^N)-$, $-NR^NC(=NR^N)-$, $-NR^NC(=NR^N)N(R^N)-$, $-C(S)-$, $-C(S)N(R^N)-$, $-NR^NC(S)-$, $-NR^NC(S)N(R^N)-$, $-S(O)-$, $-OS(O)-$, $-S(O)O-$, $-OS(O)O-$, $-OS(O)_2-$, $-S(O)_2O-$, $-OS(O)_2O-$, $-N(R^N)S(O)-$, $-S(O)N(R^N)-$, $-N(R^N)S(O)N(R^N)-$, $-OS(O)N(R^N)-$, $-N(R^N)S(O)O-$, $-S(O)_2-$, $-N(R^N)S(O)_2-$, $-S(O)_2N(R^N)-$, $-N(R^N)S(O)_2N(R^N)-$, $-OS(O)_2N(R^N)-$, or $-N(R^N)S(O)_2O-$;

each instance of R^N is independently hydrogen, optionally substituted alkyl, or a nitrogen protecting group;

Ring B is optionally substituted carbocyclyl, optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl; and
p is 1 or 2.

69. The pharmaceutical composition of claim 67, wherein the PEG lipid has the Formula (VIII):



(VIII),

or a salts thereof, wherein:

R^3 is $-\text{OR}^{\text{O}}$;


R^{O} is hydrogen, optionally substituted alkyl or an oxygen protecting group;

r is an integer between 1 and 100, inclusive;

R^5 is optionally substituted C_{10-40} alkyl, optionally substituted C_{10-40} alkenyl, or optionally substituted C_{10-40} alkynyl; and optionally one or more methylene groups of R^5 are replaced with optionally substituted carbocyclylene, optionally substituted heterocyclylene, optionally substituted arylene, optionally substituted heteroarylene, $-\text{N}(\text{R}^{\text{N}})-$, $-\text{O}-$, $-\text{S}-$, $-\text{C}(\text{O})-$, $-\text{C}(\text{O})\text{N}(\text{R}^{\text{N}})-$, $-\text{NR}^{\text{N}}\text{C}(\text{O})-$, $-\text{NR}^{\text{N}}\text{C}(\text{O})\text{N}(\text{R}^{\text{N}})-$, $-\text{C}(\text{O})\text{O}-$, $-\text{OC}(\text{O})-$, $-\text{OC}(\text{O})\text{O}-$, $-\text{OC}(\text{O})\text{N}(\text{R}^{\text{N}})-$, $-\text{NR}^{\text{N}}\text{C}(\text{O})\text{O}-$, $-\text{C}(\text{O})\text{S}-$, $-\text{SC}(\text{O})-$, $-\text{C}(=\text{NR}^{\text{N}})-$, $-\text{C}(=\text{NR}^{\text{N}})\text{N}(\text{R}^{\text{N}})-$, $-\text{NR}^{\text{N}}\text{C}(=\text{NR}^{\text{N}})-$, $-\text{NR}^{\text{N}}\text{C}(=\text{NR}^{\text{N}})\text{N}(\text{R}^{\text{N}})-$, $-\text{C}(\text{S})-$, $-\text{C}(\text{S})\text{N}(\text{R}^{\text{N}})-$, $-\text{NR}^{\text{N}}\text{C}(\text{S})-$, $-\text{NR}^{\text{N}}\text{C}(\text{S})\text{N}(\text{R}^{\text{N}})-$, $-\text{S}(\text{O})-$, $-\text{OS}(\text{O})-$, $-\text{S}(\text{O})\text{O}-$, $-\text{OS}(\text{O})\text{O}-$, $-\text{OS}(\text{O})_2-$, $-\text{S}(\text{O})_2\text{O}-$, $-\text{OS}(\text{O})_2\text{O}-$, $-\text{N}(\text{R}^{\text{N}})\text{S}(\text{O})-$, $-\text{S}(\text{O})\text{N}(\text{R}^{\text{N}})-$, $-\text{N}(\text{R}^{\text{N}})\text{S}(\text{O})\text{N}(\text{R}^{\text{N}})-$, $-\text{OS}(\text{O})\text{N}(\text{R}^{\text{N}})-$, $-\text{N}(\text{R}^{\text{N}})\text{S}(\text{O})\text{O}-$, $-\text{S}(\text{O})_2-$, $-\text{N}(\text{R}^{\text{N}})\text{S}(\text{O})_2-$, $-\text{S}(\text{O})_2\text{N}(\text{R}^{\text{N}})-$, $-\text{N}(\text{R}^{\text{N}})\text{S}(\text{O})_2\text{N}(\text{R}^{\text{N}})-$, $-\text{OS}(\text{O})_2\text{N}(\text{R}^{\text{N}})-$, or $-\text{N}(\text{R}^{\text{N}})\text{S}(\text{O})_2\text{O}-$; and

each instance of R^{N} is independently hydrogen, optionally substituted alkyl, or a nitrogen protecting group.

70. The pharmaceutical composition of claim 69, wherein the PEG lipid has the formula:



71. The pharmaceutical composition of claim 70, wherein the PEG lipid is Compound 428.

72. The pharmaceutical composition of any one of claims 25-71, wherein the lipid nanoparticle or the delivery agent further comprises a phospholipid.

73. The pharmaceutical composition of claim 72, wherein the phospholipid has the Formula (IX):




or a salt thereof, wherein:

each R^1 is independently optionally substituted alkyl; or optionally two R^1 are joined together with the intervening atoms to form optionally substituted monocyclic carbocyclyl or optionally substituted monocyclic heterocyclyl; or optionally three R^1 are joined together with the intervening atoms to form optionally substituted bicyclic carbocyclyl or optionally substituted bicyclic heterocyclyl;

n is 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10;

m is 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10;



each instance of L^2 is independently a bond or optionally substituted C_{1-6} alkylene, wherein one methylene unit of the optionally substituted C_{1-6} alkylene is optionally replaced with $-\text{O}-$, $-\text{N}(\text{R}^{\text{N}})-$, $-\text{S}-$, $-\text{C}(\text{O})-$, $-\text{C}(\text{O})\text{N}(\text{R}^{\text{N}})-$, $-\text{NR}^{\text{N}}\text{C}(\text{O})-$, $-\text{C}(\text{O})\text{O}-$, $-\text{OC}(\text{O})-$, $-\text{OC}(\text{O})\text{O}-$, $-\text{OC}(\text{O})\text{N}(\text{R}^{\text{N}})-$, $-\text{NR}^{\text{N}}\text{C}(\text{O})\text{O}-$, or $-\text{NR}^{\text{N}}\text{C}(\text{O})\text{N}(\text{R}^{\text{N}})-$;


each instance of R^2 is independently optionally substituted C_{1-30} alkyl, optionally substituted C_{1-30} alkenyl, or optionally substituted C_{1-30} alkynyl; optionally wherein one or more methylene units of R^2 are independently replaced with optionally substituted carbocyclylene, optionally substituted heterocyclylene, optionally substituted arylene, optionally substituted heteroarylene, $-\text{N}(\text{R}^{\text{N}})-$, $-\text{O}-$, $-\text{S}-$, $-\text{C}(\text{O})-$, $-\text{C}(\text{O})\text{N}(\text{R}^{\text{N}})-$, $-\text{NR}^{\text{N}}\text{C}(\text{O})-$, $-\text{NR}^{\text{N}}\text{C}(\text{O})\text{N}(\text{R}^{\text{N}})-$, $-\text{C}(\text{O})\text{O}-$, $-\text{OC}(\text{O})-$, $-\text{OC}(\text{O})\text{O}-$, $-\text{OC}(\text{O})\text{N}(\text{R}^{\text{N}})-$, $-\text{NR}^{\text{N}}\text{C}(\text{O})\text{O}-$, $-\text{C}(\text{O})\text{S}-$, $-\text{SC}(\text{O})-$, $-\text{C}(=\text{NR}^{\text{N}})-$, $-\text{C}(=\text{NR}^{\text{N}})\text{N}(\text{R}^{\text{N}})-$, $-\text{NR}^{\text{N}}\text{C}(=\text{NR}^{\text{N}})-$, $-\text{NR}^{\text{N}}\text{C}(=\text{NR}^{\text{N}})\text{N}(\text{R}^{\text{N}})-$, $-\text{C}(\text{S})-$, $-\text{C}(\text{S})\text{N}(\text{R}^{\text{N}})-$, $-\text{NR}^{\text{N}}\text{C}(\text{S})-$, $-\text{NR}^{\text{N}}\text{C}(\text{S})\text{N}(\text{R}^{\text{N}})-$, $-\text{S}(\text{O})-$, $-\text{OS}(\text{O})-$, $-\text{S}(\text{O})\text{O}-$, $-\text{OS}(\text{O})\text{O}-$, $-\text{OS}(\text{O})_2-$, $-\text{S}(\text{O})_2\text{O}-$, $-\text{OS}(\text{O})_2\text{O}-$, $-\text{N}(\text{R}^{\text{N}})\text{S}(\text{O})-$, $-\text{S}(\text{O})\text{N}(\text{R}^{\text{N}})-$, $-\text{N}(\text{R}^{\text{N}})\text{S}(\text{O})\text{N}(\text{R}^{\text{N}})-$, $-\text{OS}(\text{O})\text{N}(\text{R}^{\text{N}})-$, $-\text{N}(\text{R}^{\text{N}})\text{S}(\text{O})\text{O}-$, $-\text{S}(\text{O})_2-$, $-\text{N}(\text{R}^{\text{N}})\text{S}(\text{O})_2-$, $-\text{S}(\text{O})_2\text{N}(\text{R}^{\text{N}})-$, $-\text{N}(\text{R}^{\text{N}})\text{S}(\text{O})_2\text{N}(\text{R}^{\text{N}})-$, $-\text{OS}(\text{O})_2\text{N}(\text{R}^{\text{N}})-$, or $-\text{N}(\text{R}^{\text{N}})\text{S}(\text{O})_2\text{O}-$;

each instance of R^{N} is independently hydrogen, optionally substituted alkyl, or a nitrogen protecting group;

Ring B is optionally substituted carbocyclyl, optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl; and


p is 1 or 2;

provided that the compound is not of the formula:



wherein each instance of R^2 is independently unsubstituted alkyl, unsubstituted alkenyl, or unsubstituted alkynyl.

74. The pharmaceutical composition of claim 73, wherein the phospholipid has the formulae:



or a salt thereof, wherein:

each t is independently 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10;

each u is independently 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10; and

each v is independently 1, 2, or 3.

75. The pharmaceutical composition of any one of claims 2, 9, 25, and 26, wherein the lipid nanoparticle or the delivery agent comprises Compound 18, DSPC, Cholesterol, and Compound 428 with a mole ratio of about 50:10:38.5:1.5.

76. The pharmaceutical composition of any one of claims 1-9 and 25-75 or the polynucleotide of claim 24, wherein the mRNA comprises a microRNA (miR) binding site.

77. The pharmaceutical composition or polynucleotide of claim 76, wherein the mRNA comprises at least two different microRNA (miR) binding sites, wherein the microRNA is expressed in an immune cell of hematopoietic lineage or a cell that expresses TLR7 and/or TLR8 and secretes pro-inflammatory cytokines and/or chemokines, and wherein the mRNA comprises one or more modified nucleobases.

78. The pharmaceutical composition or polynucleotide of claim 77, wherein the mRNA comprises at least one first microRNA binding site of a microRNA abundant in an immune cell of hematopoietic lineage and at least one second microRNA binding site is of a microRNA abundant in endothelial cells.

79. The pharmaceutical composition or polynucleotide of any one of claims 76-78, wherein the mRNA comprises multiple copies of a first microRNA binding site and at least one copy of a second microRNA binding site.

80. The pharmaceutical composition or polynucleotide of any one of claims 76-79, wherein the mRNA comprises first and second microRNA binding sites of the same microRNA.

81. The pharmaceutical composition or polynucleotide of claim 80, wherein the microRNA binding sites are of the 3p and 5p arms of the same microRNA.

82. The pharmaceutical composition or polynucleotide of any one of claims 76-81, wherein the microRNA binding site is for a microRNA selected from the group consisting of miR-126, miR-142, miR-144, miR-146, miR-150, miR-155, miR-16, miR-21, miR-223, miR-24, miR-27, miR-26a, or any combination thereof.

83. The pharmaceutical composition or polynucleotide of claim 82, wherein the microRNA binding site is for a microRNA selected from the group consisting of miR126-3p, miR-142-3p, miR-142-5p, miR-155, or any combination thereof.

84. The pharmaceutical composition or polynucleotide of claim 76, wherein at least one microRNA binding site is a miR-126 binding site.

85. The pharmaceutical composition or polynucleotide of claim 76, wherein at least one microRNA binding site is a miR-142 binding site.

86. The pharmaceutical composition or polynucleotide of claim 76, wherein one microRNA binding site is a miR-126 binding site and a second microRNA binding site is for a microRNA selected from the group consisting of miR-142-3p, miR-142-5p, miR-146-3p, miR-146-5p, miR-155, miR-16, miR-21, miR-223, miR-24 and miR-27.

87. The pharmaceutical composition or polynucleotide of claim 76, comprising at least one miR-126-3p binding site and at least one miR-142-3p binding site.

88. The pharmaceutical composition or polynucleotide of claim 76, comprising at least one miR-142-3p binding site and at least one 142-5p binding site.

89. The pharmaceutical composition or polynucleotide of any one of claims 76-88, wherein the microRNA binding sites are located in the 5' UTR, 3' UTR, or both the 5' UTR and 3' UTR of the mRNA.

90. The pharmaceutical composition or polynucleotide of claim 89, wherein the microRNA binding sites are located in the 3' UTR of the mRNA.

91. The pharmaceutical composition or polynucleotide of claim 89, wherein the microRNA binding sites are located in the 5' UTR of the mRNA.

92. The pharmaceutical composition or polynucleotide of claim 89, wherein the microRNA binding sites are located in both the 5' UTR and 3' UTR of the mRNA.

93. The pharmaceutical composition or polynucleotide of claim 89, wherein at least one microRNA binding site is located in the 3' UTR immediately adjacent to the stop codon of the coding region of the mRNA.

94. The pharmaceutical composition or polynucleotide of claim 89, wherein at least one microRNA binding site is located in the 3' UTR 70-80 bases downstream of the stop codon of the coding region of the mRNA.

95. The pharmaceutical composition or polynucleotide of claim 89, wherein at least one microRNA binding site is located in the 5' UTR immediately preceding the start codon of the coding region of the mRNA.

96. The pharmaceutical composition or polynucleotide of claim 89, wherein at least one microRNA binding site is located in the 5' UTR 15-20 nucleotides preceding the start codon of the coding region of the mRNA.

97. The pharmaceutical composition or polynucleotide of claim 89, wherein at least one microRNA binding site is located in the 5' UTR 70-80 nucleotides preceding the start codon of the coding region of the mRNA.

98. The pharmaceutical composition or polynucleotide of claim 89, wherein the mRNA comprises multiple copies of the same microRNA binding site positioned immediately adjacent to each other or with a spacer of less than 5, 5-10, 10-15, or 15-20 nucleotides.

99. The pharmaceutical composition or polynucleotide of claim 89, wherein the mRNA comprises multiple copies of the same microRNA binding site located in the 3' UTR, wherein the first microRNA binding site is positioned immediately adjacent to the stop codon and the second and third microRNA binding sites are positioned 30-40 bases downstream of the 3' most residue of the first microRNA binding site.

100. The pharmaceutical composition of any one of claims 1-9 and 25-99 or the polynucleotide of claim 24 and 76-99, wherein the mRNA comprises a 3' UTR comprising a microRNA binding site selected from miR-142, miR-126, or a combination thereof.

101. The pharmaceutical composition of any one of claims 1-9 and 25-100 or the polynucleotide of claim 24 and 76-100, wherein the mRNA comprises a 3' UTR comprising a nucleic acid sequence at least about 90%, at least about 95%, at least about 96%, at least about

97%, at least about 98%, at least about 99%, or 100% identical to a 3' UTR sequence selected from the group consisting of SEQ ID NOs: 39, 58 to 84, 90, 105, 108 to 115, and 120 to 130, and any combination thereof.

102. The pharmaceutical composition or the polynucleotide of claim 101, wherein the 3' UTR comprises a nucleic acid sequence selected from the group consisting of SEQ ID NOs: 39, 58 to 84, 90, 105, 108 to 115, and 120 to 130, and any combination thereof.

103. The pharmaceutical composition of any one of claims 1-9 and 25-102 or the polynucleotide of any one of claims 24 and 76-102, wherein the mRNA comprises a 5' UTR.

104. The pharmaceutical composition or the polynucleotide of claim 103, wherein the 5' UTR comprises a nucleic acid sequence at least 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99%, or 100% identical to a 5' UTR sequence selected from the group consisting of SEQ ID NO: 38, 40 to 57, and 117 to 119, and any combination thereof.

105. The pharmaceutical composition or the polynucleotide of claim 104, wherein the 5' UTR comprising a sequence selected from the group consisting of SEQ ID NO: 38, 40 to 57, and 117 to 119, and any combination thereof.

106. The pharmaceutical composition of any one of claims 1-9 and 25-105 or the polynucleotide of any one of claims 24 and 76-105, wherein the mRNA further comprises a 5' terminal cap.

107. The pharmaceutical composition or the polynucleotide of claim 106, wherein the 5' terminal cap comprises a Cap0, Cap1, ARCA, inosine, N1-methyl-guanosine, 2'-fluoro-guanosine, 7-deaza-guanosine, 8-oxo-guanosine, 2-amino-guanosine, LNA-guanosine, 2-azidoguanosine, Cap2, Cap4, 5' methylG cap, or an analog thereof.

108. The pharmaceutical composition or the polynucleotide of claim 107, wherein the 5' terminal cap comprises a Cap1.

109. The pharmaceutical composition of any one of claims 1-9 and 25-108 or the polynucleotide of any one of claims 24 and 76-108, wherein the mRNA further comprises a poly-A region.

110. The pharmaceutical composition or the polynucleotide of claim 109, wherein the poly-A region is at least about 10, at least about 20, at least about 30, at least about 40, at least about 50, at least about 60, at least about 70, at least about 80, or at least about 90 nucleotides in length.

111. The pharmaceutical composition or the polynucleotide of claim 110, wherein the poly-A region has about 10 to about 200, about 20 to about 180, about 50 to about 160, about 70 to about 140, or about 80 to about 120 nucleotides in length.

112. The pharmaceutical composition of any one of claims 1-9 and 25-111 or the polynucleotide of any one of claims 24 and 76-111, wherein the mRNA comprises at least one chemically modified nucleobase, sugar, backbone, or any combination thereof.

113. The pharmaceutical composition or the polynucleotide of claim 112, wherein the at least one chemically modified nucleobase is selected from the group consisting of pseudouracil (ψ), N1-methylpseudouracil (m1 ψ), 1-ethylpseudouracil, 2-thiouracil (s2U), 4'-thiouracil, 5-methylcytosine, 5-methyluracil, 5-methoxyuracil, and any combination thereof.

114. The pharmaceutical composition or the polynucleotide of claim 112 or 113, wherein at least about 25%, at least about 30%, at least about 40%, at least about 50%, at least about 60%, at least about 70%, at least about 80%, at least about 90%, at least about 95%, at least about 99%, or 100% of the uracils or thymines are chemically modified.

115. The pharmaceutical composition or the polynucleotide of any one of claims 112-114, wherein at least about 25%, at least about 30%, at least about 40%, at least about 50%, at least about 60%, at least about 70%, at least about 80%, at least about 90%, at least about 95%, at least about 99%, or 100% of the guanines are chemically modified.

116. The pharmaceutical composition or the polynucleotide of any one of claims 112-115, wherein at least about 25%, at least about 30%, at least about 40%, at least about 50%, at least about 60%, at least about 70%, at least about 80%, at least about 90%, at least about 95%, at least about 99%, or 100% of the cytosines are chemically modified.

117. The pharmaceutical composition or the polynucleotide of any one of claims 112-116, wherein at least about 25%, at least about 30%, at least about 40%, at least about 50%, at least about 60%, at least about 70%, at least about 80%, at least about 90%, at least about 95%, at least about 99%, or 100% of the adenines are chemically modified.

118. The pharmaceutical composition of any one of claims 1-9 and 25-117 or the polynucleotide of any one of claims 24 and 76-117, wherein the mRNA is purified.

119. The pharmaceutical composition or the polynucleotide of any one of the preceding claims, wherein the ACADVL polypeptide is a wild type, variant, or mutant having an ACADVL activity.

120. The pharmaceutical composition or the polynucleotide of claim 119, wherein the ACADVL polypeptide comprises a wild type polypeptide sequence consisting of SEQ ID NO: 1.

121. The pharmaceutical composition or the polynucleotide of any one of the preceding claims, wherein the ACADVL polypeptide is an ACADVL fusion protein.

122. The pharmaceutical composition or the polynucleotide of claim 121, wherein the ACADVL fusion protein comprises heterologous protein moiety.

123. A polynucleotide comprising an mRNA comprising:

- (i) a 5' UTR,
- (ii) an open reading frame (ORF) encoding a human acyl-CoA dehydrogenase, very long-chain (ACADVL) polypeptide, and
- (iii) a 3' UTR comprising a microRNA binding site selected from miR-142, miR-126, or a combination thereof,

wherein the mRNA comprises at least one chemically modified nucleobase.

124. A polynucleotide comprising an mRNA comprising:
- (i) a 5'-terminal cap;
 - (ii) a 5' UTR comprising a sequence selected from the group consisting of SEQ ID NOs: 38, 40 to 57, and 117 to 119, and any combination thereof;
 - (iii) an open reading frame (ORF) encoding a human acyl-CoA dehydrogenase, very long-chain (ACADVL) polypeptide,
 - (iv) a 3' UTR comprising a nucleic acid sequence selected from the group consisting of SEQ ID NOs: 39, 58 to 84, 90, 105, 108 to 115, and 120 to 130, and any combination thereof; and
 - (v) a poly-A-region;

wherein the mRNA comprises at least one chemically modified nucleobase selected from the group consisting of pseudouracil (ψ), N1-methylpseudouracil (m1 ψ), 1-ethylpseudouracil, 2-thiouracil (s2U), 4'-thiouracil, 5-methylcytosine, 5-methyluracil, 5-methoxyuracil, and any combination thereof.

125. A pharmaceutical composition comprising the polynucleotide of any one of claims 118-124, and a delivery agent.

126. The pharmaceutical composition of claim 125, wherein the delivery agent is a lipid nanoparticle comprising a Compound selected from the group consisting of any one of Compounds 1-342 or Compound 419-428, a salt or a stereoisomer thereof, or any combination thereof.

127. The pharmaceutical composition of claim 125, wherein the delivery agent is a lipid nanoparticle comprising Compound 18 or Compound 236, Compound 428, a salt or a stereoisomer thereof, or any combination thereof.

128. The pharmaceutical composition of claim 125, wherein the delivery agent comprises Compound 18, DSPC, Cholesterol, and Compound 428 with a mole ratio of about 50:10:38.5:1.5.

129. The pharmaceutical composition of any one of claims 1-9, 25-122, and 125-128, wherein the subject is a human subject in need of treatment or prophylaxis for VLCADD.

130. The pharmaceutical composition of any one of claim 1-9, 25-122, and 125-129, wherein upon administration to the subject, the mRNA has:

- (i) a longer plasma half-life;
- (ii) increased expression of an ACADVL polypeptide encoded by the ORF;
- (iii) greater structural stability; or
- (iv) any combination thereof,

relative to a corresponding mRNA having the nucleic acid sequence of SEQ ID NO: 2 and/or administered as naked mRNA.

131. The pharmaceutical composition or polynucleotide of any one of the preceding claims, which is suitable for administration as a single unit dose or a plurality of single unit doses.

132. The pharmaceutical composition or polynucleotide of any one of the preceding claims, which is suitable for reducing the level of one or more biomarkers of VLCADD in the subject.

133. The pharmaceutical composition or polynucleotide of any one of the preceding claims for use in treating, preventing or delaying the onset of VLCADD signs or symptoms in the subject.

134. The pharmaceutical composition or polynucleotide of claim 133, wherein the signs or symptoms include hypertrophic or dilated cardiomyopathy, pericardial effusion, arrhythmias, hypotonia, hepatomegaly, and intermittent hypoglycemia, or a combination thereof.

135. A method of expressing an acyl-CoA dehydrogenase, very long-chain (ACADVL) polypeptide in a human subject in need thereof comprising administering to the subject an effective amount of the pharmaceutical composition of any one of claims 1-9, 25-122, and 125-134 or the polynucleotide of any one of claims 10-24, 76-124, and 131-134, wherein the

pharmaceutical composition or polynucleotide is suitable for administering as a single dose or as a plurality of single unit doses to the subject.

136. A method of treating, preventing or delaying the onset of VLCADD signs or symptoms in a human subject in need thereof comprising administering to the subject an effective amount of the pharmaceutical composition of any one of claims 1-9, 25-122, and 125-134 or the polynucleotide of any one of claims 10-24, 76-124, and 131-134, wherein the administration treats, prevents or delays the onset of one or more of the signs or symptoms of VLCADD in the subject.

137. A method for the treatment of VLCADD, comprising administering to a human subject in need of treatment for VLCADD a single intravenous dose of the pharmaceutical composition of any one of claims 1-9, 25-122, and 125-134 or the polynucleotide of any one of claims 10-24, 76-124, and 131-134.

138. The method of any one of the claims 135-137, wherein 24 hours after the pharmaceutical composition or polynucleotide is administered to the subject the level of an acylcarnitine in the subject is reduced by at least about 100%, at least about 90%, at least about 80%, at least about 70%, at least about 60%, at least about 50%, at least about 40%, at least about 30%, at least about 20%, or at least about 10% compared to the subject's baseline acylcarnitine level.

139. The method of claim 138, wherein the level of the acylcarnitine is reduced in the blood of the subject.

140. The method of claim 138 or 139, wherein the acylcarnitine is an acylcarnitine metabolite selected from the group consisting of C12:1 acylcarnitine, C14:1 acylcarnitine, C14:2 acylcarnitine, C14 acylcarnitine, C16 acylcarnitine, C18 acylcarnitine, C18.1 acylcarnitine, and combinations thereof.

141. The method of any one of claims 135-140, wherein 24 hours after the pharmaceutical composition or polynucleotide is administered to the subject, the ACADVL activity in the subject is increased to at least 10%, at least 20%, at least 30%, at least 40%, at

least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 100%, at least 150%, at least 200%, at least 300%, at least 400%, at least 500%, or at least 600% of the ACADVL activity in a normal individual.

142. The method of claim 141, wherein the ACADVL activity is increased in the heart, liver, brain or skeletal muscle of the subject.

143. The method of claim 141 or 142, wherein the increased ACADVL activity persists for greater than 24, 36, 48, 60, 72, 96, 120, 144, or 168 hours.

144. The method of any one of the claims 135-143, wherein 24 hours after the pharmaceutical composition or polynucleotide is administered to the subject the level of an acylcarnitine in the subject is reduced by at least 10%, at least 20%, at least about 20%, at least about 30%, at least about 40%, at least about 50%, at least about 60%, at least about 70%, at least about 80%, at least about 90%, or 100% compared to the subject's baseline acylcarnitine level.

145. The method of any one of claims 135-144, wherein the pharmaceutical composition or polynucleotide is administered as a single dose of less than 1.5 mg/kg, less than 1.25 mg/kg, less than 1 mg/kg, less than 0.75 mg/kg, or less than 0.5 mg/kg.

146. The method of any one of claims 135-145, wherein the administration to the subject is about once a week, about once every two weeks, or about once a month.

147. The method of any one of claims 135-146, wherein the pharmaceutical composition or polynucleotide is administered intravenously.

A **SEQ ID NO: 1**
(ACADVL; very long-chain specific acyl-CoA dehydrogenase, wt, isoform 1)

MQAARMAASLGRQLLRLGGGSSRLTALLGQPRPGPARRPYAGGAAQLALDKSDSHPSDA
LTRKKPAKAESKSFVGMFKGQLTTDQVFPYPSVLNEEQTOFLKELVEPVSRRFFEEVND
PAKNDALEMVEETTWQGLKELGAFGLQVPSELGGVGLCNTQYARLVEIVGMHDLGVGIT
LGAHQSIGFKGILLFGTKAQKEKYLKPLASGETVAAFCLTEPSSGSDAASIRTSAPVSP
CGKYITLNGSKLWISNGGLADIFTVFAKTPVTDPATGAVKEKITAFFVVERGFGGITHGP
PEKKMGIKASNTAEVFFDGVVRPSENVLGEVSGGFKVAMHILNNGRFGMAAALAGTMRG
IIAKAVDHATNRTQFGEKIHNFGLIQEKLARMVMLQYVTESMAYMVSANMDQGATDFQI
EAAISKIFGSEAAWKVTDECIQIMGGMGFMKEPGVERVLRDLRIFRIFEGTNDILRLFV
ALQGCMDKGKELSGLSALKNPFGNAGLLLGEAGKQLRRRAGLGSGLSLSGLVHPELSR
SGELAVRALEQFATVVEAKLIKHKKGIVNEQFLLQRLADGAIDLAMVVLSRASRSLS
EGHPTAQHEKMLCDTWCIEAAARIREGMAALQSDPWQQELYRNFKSI SKALVERGGVVT
SNPLGF

See very long-chain specific acyl-CoA dehydrogenase, Uniprot Acc. No. P49748. This isoform has been chosen as the 'canonical' sequence. All positional information in this entry refers to it.

B

Feature	Position	Length	Description
Binding site	223-223	1	Substrate; via carbonyl oxygen
Binding site	366-366	1	FAD
Active site	462-462	1	Proton acceptor
Binding site	463-463	1	Substrate; via amide nitrogen



FIG. 1

D

SEQ ID NO: 2

ATGCAGGCGGCACGGATGGCAGCGAGCTTGGGGCGGCAGCTGCTGAGGCTCGGGGGAGGAAGCAGTC
GGCTGACTGCGCTCTTAGGGCAACCCCGGCCCGGCCCTGCCCGGCGGCCCTATGCCGGGGGTGCCGC
TCAGCTGGCTCTGGACAAGTCAGATTCCCACCCCTCTGACGCTCTGACCAGGAAAAACCGGCCAAG
GCGGAATCTAAGTCCTTTGCTGTGGGAATGTTCAAAGGCCAGTTAACCACAGATCAGGTGTTCCCAT
ACCCGTCCGTGCTCAACGAAGAGCAGACACAGTTCCTTAAAGAGCTGGTGGAGCCTGTGTCGCGTTT
CTTCGAAGAAGTGAACGATCCCGCCAAGAATGACGCTTTAGAGATGGTTGAGGAGACCACTTGGCAG
GGCCTCAAGGAAGTGGGGGCTTTGGTCTGCAAGTGCCAGTGAGCTGGGTGGTGTAGGCCTTTGCA
ACACCCAGTACGCCCGTTTGGTGGAGATCGTGGGCATGCATGACCTTGGCGTGGGCATTACCTGGG
GGCCCATCAGAGCATCGGTTTTAAAGGCATCCTGCTCTTTGGCACAAAGGCCCAGAAAGAAAAATAC
CTCCCAAGCTGGCATCTGGGGAGACTGTGGCCGCTTTCTGTCTAACCGAGCCCTCAAGCGGGTCAG
ATGCAGCCTCCATCCGAACCTCTGCTGTGCCCAGCCCCCTGTGGAAAATACTATAACCTCAATGGAAG
CAAGCTTTGGATCAGTAATGGGGGCTTAGCAGACATCTTCACGGTCTTTGCCAAGACACCAGTTACA
GATCCAGCCACAGGAGCCGTGAAAGAGAAGATCACAGCTTTTGTGGTGGAAAGGGGCTTCGGGGGCA
TTACCCATGGGGCCCCCTGAGAAGAAGATGGGCATCAAGGCTTCAAACACAGCAGAGGTGTTCTTTGA
TGGAGTCCGGGTGCCAAGTGAGAACGTGCTGGGTGAAGTTGGGAGTGGCTTCAAGGTTGCCATGCAC
ATCCTCAACAATGGAAGGTTTTGGCATGGCTGCGGCCCTGGCAGGTACCATGAGAGGCATCATTGCTA
AGGCGGTAGATCATGCCACTAATCGTACCCAGTTTGGGGAGAAAATTCACAACTTTGGGCTGATCCA
GGAGAAGCTGGCACGGATGGTTATGCTGCAGTATGTAAGTGAAGTCCATGGCTTACATGGTGAGTGCT
AACATGGACCAGGGAGCCACGGACTTCCAGATAGAGGCCGCCATCAGCAAAATCTTTGGCTCGGAGG
CAGCCTGGAAGGTGACAGATGAATGCATCCAAATCATGGGGGATGGGCTTCATGAAGGAACCTGG
AGTAGAGCGTGTGCTCCGAGATCTTCGCATCTTCCGGATCTTTGAGGGAACAAATGACATTCTTCGG
CTGTTTGTGGCTCTGCAGGGCTGTATGGACAAAGGAAAGGAGCTCTCTGGGCTTGGCAGTGCTCTAA
AGAATCCCTTCGGGAATGCTGGCCTCCTGCTAGGAGAGGCGAGGCAACAGCTTAGGCGGCGGGCAGG
GCTGGGCAGCGGCCTGAGTTTGGAGCGGACTTGTCCACCCGGAGTTGAGTCGGTCTGGCGAGCTGGCA
GTACGGGCTCTGGAGCAGTTTGCCACTGTGGTGGAGGCCAAGCTGATAAAACACAAGAAGGGGATTG
TCAATGAACAGTTTCTGCTGCAGCGGCTGGCAGACGGGGCCATCGACCTCTATGCCATGGTGGTGGT
TTTGTGCGAGGGCCTCAAGATCCCTGAGTGAGGGCCACCCACGGCCCAGCATGAGAAAATGCTCTGT
GACACCTGGTGTATCGAAGCTGCAGCTCGGATCCGAGAGGGCATGGCCGCCCTGCAGTCTGACCCCT
GGCAGCAAGAGCTCTATCGCAACTTCAAATCTATCTCAAGGCCTTGGTGGAGCGGGGTGGTGTGGT
CACCAGTAACCCACTTGGCTTC

FIG. 1 (cont)

A SEQ ID NO: 3
(ACADVL; very long-chain specific anyl-CoA dehydrogenase, wt, isoform 2)

MQAARMAASLGRQLRLRLGGGSSRLTALLGQPRPGPARRPYAGGAAQLALDKSDSHPSDA
LTRKKPAKAESKSFVGMFKGQLTTDQVFPYPSVLNEEQTOFLKELVEPVSRRFFEEVND
PAKNDALEMVEETTWQGLKELGAFGLQVPSELGGVGLCNTQYARLVEIVGMHDLGVGIT
LGAHQSIGFKGILLFGTKAQKEKYLPKLASGETVAAFCLTEPSSGSDAASIRTSAVPSP
CGKYYTLNGSKLWISNGGLADIFTVFAKTPVTDPATGAVKEKITAFAVVERGFGGITHGP
PEKKMGIKASNTAEVFFDGVVRVPSENVLGEVSGGFKVAMHILNNGRFGMAAALAGTMRG
IIAKAVDHATNRTQFGEKIHNFGLIQEKLARMVMLQYVTESMAYMVSANMDQGATDFQI
EAAISKIFGSEAAWKVTDECIQIMGGMGFMKEPGVERVLRDLRIFRIFEGTNDILRLFV
ALQGCMDDKGKELSGLSALKNPFNGAGLLLGEGAGQLRRRAGLGSGLSLSGLVHPELSR
SGELAVRALEQFATVVEAKLIKHHKGIVNEQFLLQRLADGAIDLYAMVVLSRASRSLS
EGHPTAQHEKMLCDTWCIEAAARIREGMAALQSDPWQQELYRNFKSISKALVERGGVVT
SNPLGF

See very long-chain specific anyl-CoA dehydrogenase, Uniprot Acc. No.
P49748. The sequence of this isoform differs from the canonical sequence
as follows:
47-68: Missing

B

Feature	Position	Length	Description
Binding site	223-223	1	Substrate; via carbonyl oxygen
Binding site	366-366	1	FAD
Active site	462-462	1	Proton acceptor
Binding site	463-463	1	Substrate; via amide nitrogen

C




FIG. 2

D

SEQ ID NO: 4

ATGCAGGCGGCACGGATGGCAGCGAGCTTGGGGCGGCAGCTGCTGAGGCTCGGGGGAGGAAGCAGTC
GGCTGACTGCGCTCTTAGGGCAACCCCGGCCCGGCCCTGCCCGGCGGCCCTATGCCGGGGGTGCCGC
TCAGCTGGCTCTGGACAAGTCAGATTCCCACCCCTCTGACGCTCTGACCAGGAAAAAACGGCCAAG
GCGGAATCTAAGTCCTTTGCTGTGGGAATGTTCAAAGGCCAGTTAACCACAGATCAGGTGTTCCCAT
ACCCGTCCGTGCTCAACGAAGAGCAGACACAGTTCTTTAAAGAGCTGGTGGAGCCTGTGTCGCGTTT
CTTCGAAGAAGTGAACGATCCCGCCAAGAATGACGCTTTAGAGATGGTTGAGGAGACCCTTGGCAG
GGCCTCAAGGAAGTGGGGGCCCTTTGGTCTGCAAGTGCCAGTGAGCTGGGTGGTGTAGGCCTTTGCA
ACACCCAGTACGCGCGTTTGGTGGAGATCGTGGGCATGCATGACCTTGGCGTGGGCATTACCCTGGG
GGCCCATCAGAGCATCGGTTTTAAAGGCATCCTGCTCTTTGGCACAAAGGCCCAGAAAGAAAAATAC
CTCCCCAAGCTGGCATCTGGGGAGACTGTGGCCGCTTTCTGTCTAACCGAGCCCTCAAGCGGGTCAG
ATGCAGCCTCCATCCGAACCTCTGCTGTGCCAGCCCCCTGTGGAAAAATACTATACCCTCAATGGAAG
CAAGCTTTGGATCAGTAATGGGGGCCTAGCAGACATCTTCACGGTCTTTGCCAAGACACCAGTTACA
GATCCAGCCACAGGAGCCGTGAAAGAGAAGATCACAGCTTTTGTGGTGGAAAGGGGCTTCGGGGGCA
TTACCCATGGGGCCCCCTGAGAAGAAGATGGGCATCAAGGCTTCAAACACAGCAGAGGTGTTCTTTGA
TGGAGTCCGGGTGCCAAGTGAGAACGTGCTGGGTGAAGTTGGGAGTGGCTTCAAGGTTGCCATGCAC
ATCCTCAACAATGGAAGGTTTGGCATGGCTGCGGCCCTGGCAGGTACCATGAGAGGCATCATTGCTA
AGGCGGTAGATCATGCCACTAATCGTACCCAGTTTGGGGAGAAAAATTCACAACCTTTGGGCTGATCCA
GGAGAAGCTGGCACGGATGGTTATGCTGCAGTATGTAAGTGAAGTCCATGGCTTACATGGTGAGTGCT
AACATGGACCAGGGAGCCACGGACTTCCAGATAGAGGGCCGCCATCAGCAAAATCTTTGGCTCGGAGG
CAGCCTGGAAGGTGACAGATGAATGCATCCAAATCATGGGGGGTATGGGCTTCATGAAGGAACCTGG
AGTAGAGCGTGCTCCGAGATCTTCGCATCTTCCGGATCTTTGAGGGAACAAATGACATTCTTCGG
CTGTTTGTGGCTCTGCAGGGCTGTATGGACAAAGGAAAGGAGCTCTCTGGGCTTGGCAGTGCTCTAA
AGAATCCCTTCGGGAATGCTGGCCTCCTGCTAGGAGAGGCAGGCAAACAGCTTAGGCGGCGGGCAGG
GCTGGGCAGCGGCCTGAGTTTGAAGCGGACTTGTCCACCCGGAGTTGAGTCGGTCTGGCGAGCTGGCA
GTACGGGCTCTGGAGCAGTTTGGCACTGTGGTGGAGGCCAAGCTGATAAAACACAAGAAGGGGATTG
TCAATGAACAGTTTCTGCTGCAGCGGCTGGCAGACGGGGCCATCGACCTCTATGCCATGGTGGTGGT
TTTGTGAGGGGCTCAAGATCCCTGAGTGAGGGCCACCCACGGCCAGCATGAGAAAATGCTCTGT
GACACCTGGTGTATCGAAGCTGCAGCTCGGATCCGAGAGGGCATGGCCGCCCTGCAGTCTGACCCCT
GGCAGCAAGAGCTCTATCGCAACTTCAAATCTATCTCCAAGGCCTTGGTGGAGCGGGGTGGTGTGGT
CACCAGTAACCCACTTGGCTTC

FIG. 2 (cont)

A SEQ ID NO: 5 (ACADVL; very long-chain specific acyl-CoA dehydrogenase, wt, isoform 3)

MLGGLAAAAGTRIMGKEIEAEAQRPLRQTWRPGQPPAMTAKTMSSRLTALLGQPRPGPA
RRPYAGGAAQLALDKSDSHPSDALTRKKPAKAESKSFVGMFKGQLTTDQVFPYPSVLN
EEQTQFLKELVEPVSRFFEEVNDPAKNDALEMVEETTWQGLKELGAFGLQVPSELGGVG
LCNTQYARLVEIVGMHDLGVGITLGAHQSIGFKGILLFGTKAQKEKYLPKLASGETVAA
FCLTEPSSGSDAASIRTSAVPSPCGKYITLNGSKLWISNGGLADIFTVFAKTPVTDPAT
GAVKEKITAFVVERGFGGITHGPPEKKMGIKASNTAEVFFDGVVRVPSENVLGEVSGGFK
VAMHILNNGRFGMAAALAGTMRGIIAKAVDHATNRTQFGEKIHNFGLIQEKLARMVMLQ
YVTESMAYMVSANMDQGATDFQIEAAISKIFGSEAAWKVTDECIQIMGGMGFMKEPGVE
RVLRLDLRIFRIFEGTNDILRLFVALQGCMDKGKELSGLSALKNPFGNAGLLLGAEAGKQ
LRRRAGLGSLSLSGLVHPELSRSGELAVRALEQFATVVEAKLIKHKKGIVNEQFLLQR
LADGAIDLAMVVVLSRASRSLSEGHPTAQHEKMLCDTWCIEAAARIREGMAALQSDPW
QQELYRNFKSISKALVERGGVVTSNPLGF

See very long-chain specific acyl-CoA dehydrogenase, Uniprot Acc. No. P49748. The sequence of this isoform differs from the canonical sequence as follows:

1-20: MQAARMAASLGRQLRLGGG →
MLGGLAAAAGTRIMGKEIEAEAQRPLRQTWRPGQPPAMTAKTM

B

Feature	Position	Length	Description
Binding site	223-223	1	Substrate; via carbonyl oxygen
Binding site	366-366	1	FAD
Active site	462-462	1	Proton acceptor
Binding site	463-463	1	Substrate; via amide nitrogen



FIG. 3

D

SEQ ID NO: 6

ATGTTGGGGGGCCTGGCCGCGGCGGGGAAACCCGGATAATGGGAAAGGAGATAGAAGCAGAAGCAC
AGAGGCCCCCTGAGGCAAACATGGAGACCTGGCCAGCCACCAGCGATGACAGCAAAGACGATGAGCTC
GCGGCTCACGGCGCTCCTGGGGCAGCCCCGGCCCGGCCCTGCCCGGCGGCCCTATGCCGGGGGTGCC
GCTCAACTGGCTCTGGACAAGTCAGATTCCCACCCCTCTGACGCTCTGACCAGGAAAAAGCCGGCCA
AGGCGGAATCTAAGTCCTTTGCTGTGGGAATGTTCAAAGGCCAGCTCACCACAGATCAGGTGTTCCC
ATACCCGTCCGTGCTCAACGAAGAGCAGACACAGTTTCTTAAAGAGCTGGTGGAGCCTGTGTCCCGT
TTCTTCGAGGAAGTGAACGATCCCGCCAAGAATGACGCTCTGGAGATGGTGGAGGAGACCACTTGGC
AGGGCCTCAAGGAGCTGGGGGCCTTTGGTCTGCAAGTGCCCACTGAGCTGGGTGGTGTGGGCCTTTG
CAACACCCAGTACGCCCCGTTTGGTGGAGATCGTGGGCATGCATGACCTTGGCGTGGGCATCACCCTG
GGGGCCCATCAGAGCATCGGTTTCAAAGGCATCCTGCTTTTGGCACAAAGGCCCAGAAAGAAAAAT
ACCTCCCCAAGCTGGCATCTGGGGAGACTGTGGCCGCTTCTGTCTAACCAGGCCCTCAAGCGGGTC
AGATGCAGCCTCCATCCGAACCTCTGCTGTGCCAGCCCCCTGTGGAAAAATACTATACCCCTCAATGGA
AGCAAGCTTTGGATCAGTAATGGGGGCCTAGCAGACATTTTACGGTCTTTGCCAAGACACCAAGTTA
CAGATCCAGCCACAGGAGCCGTGAAGGAGAAGATCACAGCTTTTGTGGTGGAGAGGGGCTTCGGGGG
CATTACCCATGGGCCCCCTGAGAAGAAGATGGGCATCAAGGCTTCAAACACAGCAGAGGTGTTCTTT
GATGGAGTACGGGTGCCATCGGAGAACGTGCTGGGTGAGGTTGGGAGTGGCTTCAAGGTTGCCATGC
ACATCCTCAACAATGGAAGGTTTGGCATGGCTGCGGCCCTGGCAGGTACCATGAGAGGCATCATTGC
TAAGGCGGTAGATCATGCCACTAATCGTACCCAGTTTGGGGAAAAAATTCACAACTTTGGGCTGATC
CAGGAGAAGCTGGCACGGATGGTTATGCTGCAGTATGTAAGTGAAGTCCATGGCTTACATGGTGAAGT
CTAACATGGACCAGGGAGCCACGGACTTCCAGATAGAGGCCGCCATCAGCAAAATCTTTGGCTCGGA
GGCAGCCTGGAAGGTGACAGATGAATGCATCCAAATCATGGGGGGTATGGGCTTCATGAAGGAACCT
GGAGTAGAGCGTGTGCTCCGAGATCTTCGCATCTTCCGGATCTTTGAGGGGACAAATGACATTCTTC
GGCTGTTTGTGGCTCTGCAGGGCTGTATGGACAAAGGAAAGGAGCTCTCTGGGCTTGGCAGTGCTCT
AAAGAATCCCTTCGGGAATGCTGGCCTCCTGCTAGGAGAGGCGAGGCAACAGCTGAGGCGGCGGGCA
GGGCTGGGCAGCGGCCTGAGTCTCAGCGGACTTGTCCACCCGAGTTGAGTCGGAGTGGCGAGCTGG
CAGTACGGGCTCTGGAGCAGTTTGCCACTGTTGTGGAGGCCAAGCTGATAAAACACAAGAAGGGGAT
TGTCATGAACAGTTTCTGCTGCAGCGGCTGGCAGACGGGGCCATCGACCTCTATGCTATGGTGGTG
GTTCTCTCGAGGGCCTCAAGATCCCTGAGTGAGGGGCCACCCACGGCCCAGCATGAGAAAAATGCTCT
GTGACACCTGGTGTATCGAGGCTGCAGCTCGGATCCGAGAGGGCATGGCCGCCCTGCAGTCTGACCC
CTGGCAGCAAGAGCTCTACCGCAACTTCAAAAGCATCTCCAAGGCCTTGGTGGAGCGGGGTGGTGTG
GTCACCAGCAACCCACTTGGCTTC

FIG. 3 (cont)

Protein	Length	Theoretical Minimum U (%)	Theoretical Minimum U (abs)	U Content (%)	U Content v WT (%)	U Content v Theoretical Minimum (%)	UU pairs	UU pairs v WT (%)	UUU	UUUU	UUUUU
ACADVL-PROTEIN	1965	439	20.87%	100.00%	100.00%	100.00%	23	100.00%	20	3	0
ACADVL-CO01	1965	300	15.27%	73.17%	73.17%	125.52%	23	64.59%	9	1	0
ACADVL-CO02	1965	303	15.42%	73.90%	73.90%	126.76%	27	64.59%	5	0	2
ACADVL-CO03	1965	303	15.42%	73.90%	73.90%	126.78%	35	83.33%	2	2	1
ACADVL-CO04	1965	299	15.22%	72.93%	72.93%	125.10%	22	52.58%	7	1	1
ACADVL-CO05	1965	291	14.81%	70.98%	70.98%	121.76%	27	64.29%	5	0	0
ACADVL-CO06	1965	297	15.11%	72.44%	72.44%	124.27%	30	71.43%	5	0	0
ACADVL-CO07	1965	292	14.86%	71.02%	71.02%	122.16%	31	73.61%	5	0	0
ACADVL-CO08	1965	297	15.11%	72.44%	72.44%	124.27%	26	61.90%	6	1	2
ACADVL-CO09	1965	299	15.22%	72.93%	72.93%	125.10%	31	73.61%	6	0	0
ACADVL-CO10	1965	294	14.96%	71.71%	71.71%	123.01%	27	64.29%	7	1	0
ACADVL-CO11	1965	295	15.01%	71.95%	71.95%	123.43%	29	69.05%	6	0	1
ACADVL-CO12	1965	303	15.42%	73.90%	73.90%	126.76%	26	61.90%	6	1	1
ACADVL-CO13	1965	284	14.56%	71.71%	71.71%	123.01%	25	59.52%	5	2	1
ACADVL-CO14	1965	306	15.52%	74.59%	74.59%	127.62%	29	69.05%	6	0	0
ACADVL-CO15	1965	291	14.81%	70.98%	70.98%	121.76%	22	52.39%	9	2	0
ACADVL-CO16	1965	298	14.66%	70.24%	70.24%	120.50%	29	69.05%	4	0	0
ACADVL-CO17	1965	296	15.17%	72.68%	72.68%	124.69%	28	66.67%	5	0	2
ACADVL-CO18	1965	287	15.11%	73.44%	73.44%	124.37%	30	71.43%	4	1	0
ACADVL-CO19	1965	284	14.45%	69.27%	69.27%	118.83%	24	57.14%	7	0	1
ACADVL-CO20	1965	284	14.45%	69.27%	69.27%	118.83%	34	80.95%	1	0	0
ACADVL-CO21	1965	293	14.91%	71.46%	71.46%	122.59%	25	59.52%	7	1	1
ACADVL-CO22	1965	294	14.96%	71.71%	71.71%	123.01%	27	64.59%	7	0	1
ACADVL-CO23	1965	287	14.61%	70.00%	70.00%	120.08%	31	73.81%	4	0	1
ACADVL-CO24	1965	289	14.71%	70.49%	70.49%	120.92%	22	53.38%	7	1	1
ACADVL-CO25	1965	295	15.01%	71.95%	71.95%	123.43%	34	60.95%	5	1	0
	MAX	305	15.52%	74.59%	74.59%	127.62%	35	83.33%	9	2	2
	MIN	284	14.45%	69.27%	69.27%	118.83%	22	52.58%	1	0	0
	AVERAGE	294.68	15.01%	71.92%	71.92%	123.58%	27.76	66.10%	5.64	0.6	0.64
	MEDIAN	295	15.01%	71.95%	71.95%	123.43%	27	64.29%	6	0	1
	STD DEV	5.80	0.20%	1.42%	1.42%	2.43%	3.74	8.92%			

FIG. 4

Protein	Length	Theoretical Maximum G (%)	Theoretical Maximum G (abs)		
ACADVL Protein	655	45.55%	855		
Nucleic Acid	Length	G Content(abs)	G Content(%)	G Content v WT (%)	G Content v Theoretical Maximum (%)
ACADVL-WT	1965	618	31.45%	100.00%	69.05%
ACADVL-CO01	1965	650	33.08%	105.18%	72.63%
ACADVL-CO02	1965	648	32.98%	104.85%	72.40%
ACADVL-CO03	1965	649	33.03%	105.02%	72.51%
ACADVL-CO04	1965	653	33.23%	105.66%	72.96%
ACADVL-CO05	1965	667	33.94%	107.93%	74.53%
ACADVL-CO06	1965	651	33.13%	105.34%	72.74%
ACADVL-CO07	1965	649	33.03%	105.02%	72.51%
ACADVL-CO08	1965	653	33.23%	105.66%	72.96%
ACADVL-CO09	1965	655	33.33%	105.99%	73.18%
ACADVL-CO10	1965	663	33.74%	107.28%	74.09%
ACADVL-CO11	1965	654	33.28%	105.83%	73.07%
ACADVL-CO12	1965	667	33.94%	107.93%	74.53%
ACADVL-CO13	1965	657	33.44%	106.31%	73.41%
ACADVL-CO14	1965	661	33.64%	106.96%	73.85%
ACADVL-CO15	1965	669	34.05%	108.25%	74.75%
ACADVL-CO16	1965	676	34.40%	109.39%	75.53%
ACADVL-CO17	1965	667	33.94%	107.93%	74.53%
ACADVL-CO18	1965	663	33.74%	107.28%	74.08%
ACADVL-CO19	1965	660	33.59%	106.80%	73.74%
ACADVL-CO20	1965	650	33.08%	105.18%	72.63%
ACADVL-CO21	1965	672	34.20%	108.74%	75.08%
ACADVL-CO22	1965	662	33.69%	107.12%	73.97%
ACADVL-CO23	1965	660	33.59%	106.80%	73.74%
ACADVL-CO24	1965	668	33.99%	108.09%	74.64%
ACADVL-CO25	1965	660	33.59%	106.80%	73.74%
	MAX	676	34.40%	109.39%	75.53%
	MIN	648	32.98%	104.85%	72.40%
	MEAN	659.36	33.56%	106.69%	73.67%
	MEDIAN	660	33.59%	106.80%	73.74%
	STD DEV	8.01	0.41%	1.30%	0.89%

FIG. 5

Protein	Length	Theoretical Maximum C	Theoretical Maximum C (abs)		
ACADVL Protein	655	43.46%	854		
Nucleic Acid	Length	C Content(abs)	C Content(%)	C Content v WT (%)	C Content v Theoretical Maximum (%)
ACADVL-WT	1965	435	25.19%	100.00%	57.56%
ACADVL-CO01	1965	609	30.99%	123.03%	71.31%
ACADVL-CO02	1965	609	30.99%	123.03%	71.31%
ACADVL-CO03	1965	606	30.84%	122.42%	70.96%
ACADVL-CO04	1965	610	31.04%	123.23%	71.43%
ACADVL-CO05	1965	610	31.04%	123.23%	71.43%
ACADVL-CO06	1965	621	31.60%	125.45%	72.72%
ACADVL-CO07	1965	631	32.11%	127.47%	73.89%
ACADVL-CO08	1965	621	31.60%	125.45%	72.72%
ACADVL-CO09	1965	613	31.20%	123.84%	71.78%
ACADVL-CO10	1965	613	31.20%	123.84%	71.78%
ACADVL-CO11	1965	617	31.40%	124.65%	72.25%
ACADVL-CO12	1965	587	29.87%	118.59%	68.74%
ACADVL-CO13	1965	608	30.94%	122.83%	71.19%
ACADVL-CO14	1965	602	30.64%	121.62%	70.49%
ACADVL-CO15	1965	603	30.69%	121.82%	70.61%
ACADVL-CO16	1965	592	30.13%	119.60%	69.32%
ACADVL-CO17	1965	607	30.89%	122.63%	71.08%
ACADVL-CO18	1965	593	30.18%	119.80%	69.44%
ACADVL-CO19	1965	614	31.25%	124.04%	71.90%
ACADVL-CO20	1965	621	31.60%	125.45%	72.72%
ACADVL-CO21	1965	600	30.53%	121.21%	70.26%
ACADVL-CO22	1965	602	30.64%	121.62%	70.49%
ACADVL-CO23	1965	615	31.30%	124.24%	72.01%
ACADVL-CO24	1965	608	30.94%	122.83%	71.19%
ACADVL-CO25	1965	604	30.74%	122.02%	70.73%
	MAX	631	32.11%	127.47%	73.89%
	MIN	587	29.87%	118.59%	68.74%
	AVERAGE	608.64	30.97%	122.96%	71.27%
	MEDIAN	609	30.99%	123.03%	71.31%
	STD DEV	9.91	0.50%	2.00%	1.16%

FIG. 6

Protein	Length	Theoretical Maximum GC	Theoretical Maximum GC (abs)		
ACADVL Protein	655	69.26%	1361		
Nucleic Acid	Length	GC Content(abs)	GC Content(%)	GC Content v WT (%)	GC Content v Theoretical Maximum (%)
ACADVL-WT	1965	1113	56.64%	100.00%	81.78%
ACADVL-CO01	1965	1259	64.07%	113.12%	92.51%
ACADVL-CO02	1965	1257	63.97%	112.94%	92.36%
ACADVL-CO03	1965	1255	63.87%	112.76%	92.21%
ACADVL-CO04	1965	1263	64.27%	113.48%	92.80%
ACADVL-CO05	1965	1277	64.99%	114.73%	93.83%
ACADVL-CO06	1965	1272	64.73%	114.29%	93.46%
ACADVL-CO07	1965	1280	65.14%	115.00%	94.05%
ACADVL-CO08	1965	1274	64.83%	114.47%	93.61%
ACADVL-CO09	1965	1268	64.53%	113.93%	93.17%
ACADVL-CO10	1965	1276	64.94%	114.65%	93.75%
ACADVL-CO11	1965	1271	64.68%	114.20%	93.39%
ACADVL-CO12	1965	1254	63.82%	112.67%	92.14%
ACADVL-CO13	1965	1265	64.38%	113.66%	92.95%
ACADVL-CO14	1965	1263	64.27%	113.48%	92.80%
ACADVL-CO15	1965	1272	64.73%	114.29%	93.46%
ACADVL-CO16	1965	1268	64.53%	113.93%	93.17%
ACADVL-CO17	1965	1274	64.83%	114.47%	93.61%
ACADVL-CO18	1965	1256	63.92%	112.85%	92.29%
ACADVL-CO19	1965	1274	64.83%	114.47%	93.61%
ACADVL-CO20	1965	1271	64.68%	114.20%	93.39%
ACADVL-CO21	1965	1272	64.73%	114.29%	93.46%
ACADVL-CO22	1965	1264	64.33%	113.57%	92.87%
ACADVL-CO23	1965	1275	64.89%	114.56%	93.68%
ACADVL-CO24	1965	1276	64.94%	114.65%	93.75%
ACADVL-CO25	1965	1264	64.33%	113.57%	92.87%
	MAX	1280	65.14%	115.00%	94.05%
	MIN	1254	63.82%	112.67%	92.14%
	AVERAGE	1269	64.53%	113.93%	93.17%
	MEDIAN	1271	64.68%	114.20%	93.39%
	STD DEV	7.58	0.39%	0.68%	0.56%

FIG. 7

Sequence	GC	GC 1st	GC 2nd	GC 3rd
ACADVL-WT	56.64	61.22	43.97	64.73
ACADVL-CO12	63.82	60.76	43.97	86.72
ACADVL-CO03	63.87	61.07	43.97	86.56
ACADVL-CO18	63.92	59.54	43.97	88.24
ACADVL-CO02	63.97	60.61	43.97	87.33
ACADVL-CO01	64.07	60.46	43.97	87.79
ACADVL-CO14	64.27	61.07	43.97	87.79
ACADVL-CO04	64.27	60.92	43.97	87.94
ACADVL-CO22	64.33	60.61	43.97	88.4
ACADVL-CO25	64.33	60.46	43.97	88.55
ACADVL-CO13	64.38	61.22	43.97	87.94
ACADVL-CO16	64.53	60.31	43.97	89.31
ACADVL-CO09	64.53	60.46	43.97	89.16
ACADVL-CO20	64.68	60.76	43.97	89.31
ACADVL-CO11	64.68	60.92	43.97	89.16
ACADVL-CO21	64.73	60.15	43.97	90.08
ACADVL-CO06	64.73	60.92	43.97	89.31
ACADVL-CO15	64.73	61.37	43.97	88.85
ACADVL-CO17	64.83	61.22	43.97	89.31
ACADVL-CO08	64.83	61.53	43.97	89.01
ACADVL-CO19	64.83	61.22	43.97	89.31
ACADVL-CO23	64.89	61.37	43.97	89.31
ACADVL-CO10	64.94	60.46	43.97	90.38
ACADVL-CO24	64.94	60.46	43.97	90.38
ACADVL-CO05	64.99	61.07	43.97	89.92
ACADVL-CO07	65.14	61.53	43.97	89.92
Overall	64.23	60.83	43.97	87.87

FIG. 8

INTERNATIONAL SEARCH REPORT

International application No

PCT/US2017/033402

A. CLASSIFICATION OF SUBJECT MATTER

INV. A61K48/00 C12N15/67 C07H21/00 A61K47/46 A61K31/7088
A61P3/00

ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

A61K C12N C07H

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, FSTA, BIOSIS, EMBASE, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2013/151665 A2 (MODERNA THERAPEUTICS [US]) 10 October 2013 (2013-10-10) abstract claims 1, 2, 4, 6, 38-44 Pages 222-223, Table 6, entries 660-664 pages 10-12, paragraph 31-36 pages 135-146, paragraph 146-181 sequence GS_NUC_ALERT:W02013151665.250468 ----- -/-	1-147



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

23 August 2017

Date of mailing of the international search report

29/08/2017

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Gohlke, Pascale

INTERNATIONAL SEARCH REPORT

International application No

PCT/US2017/033402

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	ANDRESEN B S ET AL: "CLONING AND CHARACTERIZATION OF HUMAN VERY-LONG-CHAIN ACYL-COA DEHYDROGENASE CDNA, CHROMOSOMAL ASSIGNMENT OF THE GENE AND IDENTIFICATION IN FOUR PATIENTS OF NINE DIFFERENT MUTATIONS WITHIN THE VLCAD GENE", HUMAN MOLECULAR GENETICS, OXFORD UNIVERSITY PRESS, GB, vol. 5, no. 4, 1 January 1990 (1990-01-01), pages 461-472, XP002949600, ISSN: 0964-6906, DOI: 10.1093/HMG/5.4.461 the whole document	3-8, 10-24, 101-111, 118-124, 129-132
Y	----- DREW WEISSMAN ET AL: "mRNA: Fulfilling the Promise of Gene Therapy", MOLECULAR THERAPY, vol. 23, no. 9, 1 September 2015 (2015-09-01), pages 1416-1417, XP055391450, US ISSN: 1525-0016, DOI: 10.1038/mt.2015.138 the whole document	1-147
Y	----- UGUR SAHIN ET AL: "mRNA-based therapeutics - developing a new class of drugs", NATURE REVIEWS DRUG DISCOVERY, vol. 13, no. 10, 19 September 2014 (2014-09-19), pages 759-780, XP055159205, ISSN: 1474-1776, DOI: 10.1038/nrd4278 abstract Scheme "Timeline"; page 760 page 761, right-hand column - page 762 page 764	1-147
Y	----- ANDREAS THESS ET AL: "Sequence-engineered mRNA Without Chemical Nucleoside Modifications Enables an Effective Protein Therapy in Large Animals", MOLECULAR THERAPY, vol. 23, no. 9, 8 June 2015 (2015-06-08), pages 1456-1464, XP055316910, US ISSN: 1525-0016, DOI: 10.1038/mt.2015.103 abstract	1-147
Y	----- WO 2014/144196 A1 (SHIRE HUMAN GENETIC THERAPIES [US]) 18 September 2014 (2014-09-18) abstract; claims ----- -/--	1-147

INTERNATIONAL SEARCH REPORT

International application No

PCT/US2017/033402

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 2015/167017 A1 (ROY ATANU [US] ET AL) 18 June 2015 (2015-06-18) abstract; claims	1-147
X,P	----- WO 2017/062513 A1 (MODERNATX INC [US]) 13 April 2017 (2017-04-13) abstract; claims -----	1-147

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2017/033402

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2013151665 A2	10-10-2013	AU 2013243834 A1	30-10-2014
		AU 2013243946 A1	30-10-2014
		AU 2013243947 A1	30-10-2014
		AU 2013243948 A1	30-10-2014
		AU 2013243950 A1	30-10-2014
		AU 2013243951 A1	30-10-2014
		AU 2013243952 A1	30-10-2014
		AU 2013243953 A1	30-10-2014
		AU 2013243954 A1	30-10-2014
		AU 2013243955 A1	30-10-2014
		CA 2868393 A1	10-10-2013
		CA 2868398 A1	10-10-2013
		CA 2868418 A1	10-10-2013
		CA 2868422 A1	10-10-2013
		CA 2868429 A1	10-10-2013
		CA 2868434 A1	10-10-2013
		CA 2868438 A1	10-10-2013
		CA 2868440 A1	10-10-2013
		CA 2868996 A1	10-10-2013
		CA 2869005 A1	10-10-2013
		CN 104870022 A	26-08-2015
		EP 2833892 A2	11-02-2015
		EP 2833894 A1	11-02-2015
		EP 2833920 A2	11-02-2015
		EP 2833921 A2	11-02-2015
		EP 2833922 A2	11-02-2015
		EP 2833923 A1	11-02-2015
		EP 2834259 A1	11-02-2015
		EP 2834260 A1	11-02-2015
		EP 2834358 A2	11-02-2015
		EP 2847329 A1	18-03-2015
		EP 2849799 A2	25-03-2015
		HK 1206601 A1	15-01-2016
		HK 1206635 A1	15-01-2016
		HK 1206636 A1	15-01-2016
		HK 1206638 A1	15-01-2016
		HK 1206639 A1	15-01-2016
		HK 1206748 A1	15-01-2016
		HK 1206779 A1	15-01-2016
		HK 1206780 A1	12-02-2016
		HK 1207306 A1	29-01-2016
		JP 2015513912 A	18-05-2015
		JP 2015513913 A	18-05-2015
		JP 2015513914 A	18-05-2015
		JP 2015513916 A	18-05-2015
		JP 2015516143 A	08-06-2015
		JP 2015517995 A	25-06-2015
		JP 2015518704 A	06-07-2015
		JP 2015518816 A	06-07-2015
		JP 2015519040 A	09-07-2015
		JP 2015519881 A	16-07-2015
		JP 2017121239 A	13-07-2017
		JP 2017121240 A	13-07-2017
		JP 2017121241 A	13-07-2017
		JP 2017121243 A	13-07-2017
		JP 2017123847 A	20-07-2017
		JP 2017123853 A	20-07-2017
		US 2013259923 A1	03-10-2013

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2017/033402

Patent document cited in search report	Publication date	Patent family member(s)	Publication date	
		US 2013259924 A1	03-10-2013	
		US 2014010861 A1	09-01-2014	
		US 2014105964 A1	17-04-2014	
		US 2014105965 A1	17-04-2014	
		US 2014105966 A1	17-04-2014	
		US 2014107189 A1	17-04-2014	
		US 2014113959 A1	24-04-2014	
		US 2014141067 A1	22-05-2014	
		US 2014148502 A1	29-05-2014	
		US 2014155472 A1	05-06-2014	
		US 2014155473 A1	05-06-2014	
		US 2014155474 A1	05-06-2014	
		US 2014155475 A1	05-06-2014	
		US 2014161873 A1	12-06-2014	
		US 2014171485 A1	19-06-2014	
		US 2014179771 A1	26-06-2014	
		US 2014186432 A1	03-07-2014	
		US 2014193482 A1	10-07-2014	
		US 2014194494 A1	10-07-2014	
		US 2014200262 A1	17-07-2014	
		US 2014200263 A1	17-07-2014	
		US 2014200264 A1	17-07-2014	
		US 2014206755 A1	24-07-2014	
		US 2014221465 A1	07-08-2014	
		US 2014249208 A1	04-09-2014	
		US 2014255467 A1	11-09-2014	
		US 2014255468 A1	11-09-2014	
		US 2015044277 A1	12-02-2015	
		US 2016075733 A1	17-03-2016	
		WO 2013151663 A1	10-10-2013	
		WO 2013151664 A1	10-10-2013	
		WO 2013151665 A2	10-10-2013	
		WO 2013151667 A1	10-10-2013	
		WO 2013151668 A2	10-10-2013	
		WO 2013151669 A1	10-10-2013	
		WO 2013151670 A2	10-10-2013	
		WO 2013151671 A1	10-10-2013	
		WO 2013151672 A2	10-10-2013	
		WO 2013151736 A2	10-10-2013	

WO 2014144196	A1	18-09-2014	EP 2972360 A1	20-01-2016
			US 2016151409 A1	02-06-2016
			WO 2014144196 A1	18-09-2014

US 2015167017	A1	18-06-2015	CA 2933568 A1	18-06-2015
			CN 106170553 A	30-11-2016
			EP 2918275 A1	16-09-2015
			EP 3053585 A1	10-08-2016
			EP 3080289 A2	19-10-2016
			JP 2017500034 A	05-01-2017
			US 2015167017 A1	18-06-2015
			US 2016304552 A1	20-10-2016
			WO 2015089511 A2	18-06-2015

WO 2017062513	A1	13-04-2017	NONE	
