



(51) International Patent Classification:
C12Q 1/00 (2006.01)

(21) International Application Number:
PCT/US2014/029495

(22) International Filing Date:
14 March 2014 (14.03.2014)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
61/799,276 15 March 2013 (15.03.2013) US

(71) Applicant (for all designated States except US): **THE TRUSTEES OF COLUMBIA UNIVERSITY IN THE CITY OF NEW YORK** [US/US]; West 116th Street And Broadway, New York, NY 10027 (US).

(72) Inventors; and

(71) Applicants (for US only): **JU, Jingyue** [US/US]; 267 Marietta Street, Englewood Cliffs, NJ 07632 (US). **KUMAR, Shiv** [US/US]; 21 Muirhead Court, Belle Mead, NJ 08502 (US). **TAO, Chuanjuan** [CN/US]; 605 West 112 Street, Apt 6I, New York, NY 10025 (US). **KALACHIKOV, Sergey** [US/US]; 620 West 171 Street, Apt 4D, New York, NY 10032 (US). **RUSO, James, J.** [US/US]; 60 Haven Avenue, Apt. 25E, New York, NY 10032 (US).

(74) Agent: **WHITE, John, P.**; Cooper & Dunham LLP, 30 Rockefeller Plaza, New York, NY 10112 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))

(54) Title: METHOD FOR DETECTING MULTIPLE PREDETERMINED COMPOUNDS IN A SAMPLE

(57) Abstract: This invention provides methods for detecting the presence of a plurality of predetermined compounds in a sample using a plurality of tag moieties and at least one nanopore. This invention also provides methods for determining the quantity of each of a plurality of predetermined compounds in a sample using a plurality of tag moieties and at least one nanopore. This invention further provides methods for detecting interaction of at least two predetermined compounds using at least one tag moiety and at least one nanopore.



METHOD FOR DETECTING MULTIPLE PREDETERMINED COMPOUNDS IN A SAMPLE

5 Throughout this application, certain patents and publications are referenced, the latter by authors and publication year. Full citations for these publications may be found immediately preceding the claims. The disclosures of these patents and publications in their entireties are hereby incorporated by reference into this
10 application in order to describe more fully the state of the art to which this invention relates.

FIELD OF INVENTION

This invention provides methods for detecting the presence of a
15 plurality of predetermined compounds in a sample using a plurality of tag moieties and at least one nanopore. This invention also provides methods for determining the quantity of each of a plurality of predetermined compounds in a sample using a plurality of tag moieties and at least one nanopore. This invention further provides
20 methods for detecting interaction of at least two predetermined compounds using a tag moiety and at least one nanopore.

BACKGROUND OF THE INVENTION

Classic immunological approaches to detection of proteins using
25 antibodies, receptors, or other binding partners include, among others, enzyme-linked immunosorbent (ELISA) assay (generally in the form of an antibody sandwich method), radioimmunoassays, and immunoblotting methods [Burnette 1981, Engvall et al. 1971, and Yalow et al. 1960], with equivalent biochemical approaches being
30 used for protein-receptor and protein-ligand reactions. Majority of these methods rely on examining one protein at a time. Moreover, the protein target typically must be present in large amounts and at relatively high concentrations to assure a reliably detectable signal.

35 Beginning in 1975 but largely over the last 15 years, methods that attempt to simultaneously examine many different proteins have

- 2 -

appeared. These include 2-D gel electrophoresis, tandem mass spectrometry (MS-2 or MS-3) systems with intermediate protein cleavage, isotope-coded affinity tag (ICAT)-MS, MudPIT (LC-2/MS-2), and combinations of these approaches [Guerrera et al. 2005, Gygi et al. 1999, Klose 1975, and O'Farrell 1975]. For instance, in the ICAT approach, proteins from different tissues are labeled with tags containing either hydrogen or deuterium, and the differential patterns are observed by mass spectrometry. While many of these methods allow multiple samples to be compared concurrently, due to the cost of associated technologies such as mass spectroscopy, many of these approaches have not found general utility.

A relatively recent addition to the repertoire, protein arrays, in which fluorescently labeled proteins are allowed to bind to numerous spots, each containing covalently attached antibodies for a specific protein (antigen), are an appealing solution, as they can be mass produced and data analysis standardized [Angenendt 2005, Bussow et al. 1998, Cahill 2001, de Wildt et al. 2000]. While limited by the number of available specific antibodies that can function on a solid phase, a more important shortcoming of this method is the relative binding ability of the antibodies. Unlike DNA probes on gene expression microarrays, where probes can be selected to be fairly uniform in their binding affinity for mRNA targets, different antibodies may bind their fluorescently labeled antigens with very different affinities. Because of this variable affinity, quantification from spot to spot (antigen to antigen) becomes difficult, especially when combined with the likelihood that the fluorescent signal can lie outside of the linear range of detection. For example, low copy number proteins in the sample will not be seen, unless their binding is stronger than the average antigen-antibody interaction elsewhere on the chip, in which case they will be over-represented. Moreover, the effective concentration range will have the same floors and ceilings as other fluorescent methods on microarrays, and small changes in protein levels will be difficult to distinguish using protein arrays. Another common issue with fluorescent labeling is the existence of overlapping emission

- 3 -

spectra, which limits the number of differentially labeled samples that can be applied to the arrays.

In summary, gene regulation analysis at the level of protein
5 synthesis, like proteomics in general, lags behind nucleic acid
analysis in its throughput, sensitivity and automation. This is due
to the relatively poor stability of proteins, their high
heterogeneity, and the requirement for a much wider dynamic range of
detection with increased demand for sensitivity approaching the
10 single molecule detection level, a need not easily met by
fluorescent or colorimetric measurements. While protein arrays
based on antibody interactions with fluorescently labeled antigens
or secondary antibodies have gained some degree of popularity over
the last decade [Cahill 2001], some of the drawbacks of this
15 approach, including fluorescence saturation and overlap in
fluorescent emission, make accurate quantification difficult. While
there are currently several examples of application of nanopore-
based analytics, no existing technology allows quantification of
protein-protein interactions with a plurality of tags.

20

- 4 -

SUMMARY OF THE INVENTION

This invention provides a method for detecting the presence of a plurality of predetermined compounds in a sample which comprises:

- 5 (a) contacting the sample with a plurality of detectable tag moieties, wherein each detectable tag moiety comprises at least one detectable component, under conditions permitting the tag moieties to attach to each of the predetermined compounds present in the sample to form detectably tagged
10 predetermined compounds, wherein each tag moiety uniquely attaches to one of the predetermined compounds present in the sample and is distinguishable from any tag moiety attached to any other predetermined compound present in the sample;
- 15 (b) treating the sample from step (a) to form an electrolyte solution containing the detectably tagged predetermined compounds;
- 20 (c) contacting the electrolyte solution from step (b) with at least one nanopore, wherein said at least one nanopore comprises a means for binding each of the detectably tagged predetermined compounds present in the electrolyte solution, under conditions such that the detectably tagged predetermined compounds bind to said at least one nanopore;
- 25 (d) treating the detectably tagged predetermined compounds bound to said at least one nanopore so as to separate the tag moieties from the predetermined compounds; and
- 30 (e) detecting said at least one detectable component of each of the tag moieties so released as the tag moieties pass through said at least one nanopore so as to thereby detecting the presence of the plurality of predetermined compounds in the sample.

This invention provides a method for detecting the presence of a plurality of predetermined compounds in a sample which comprises:

- 35 (a) contacting the sample with a plurality of detectable tag moieties, wherein each detectable tag moiety comprises at least one detectable component, under conditions permitting

- 5 -

- the tag moieties to attach to each of the predetermined compounds present in the sample to form detectably tagged predetermined compounds, wherein each tag moiety uniquely attaches to one of the predetermined compounds present in the sample and is distinguishable from any tag moiety attached to any other predetermined compound present in the sample;
- 5 (b) treating the sample from step (a) to form an electrolyte solution containing the detectably tagged predetermined compounds;
- 10 (c) contacting the electrolyte solution from step (b) with at least one nanopore, wherein said at least one nanopore comprises a means for binding each of the detectably tagged predetermined compounds present in the electrolyte solution, under conditions such that the detectably tagged predetermined
- 15 compounds bind to said at least one nanopore, and wherein each of the tag moieties of said bound detectably tagged predetermined compounds localizes within said at least one nanopore;
- 20 (d) detecting said at least one detectable component of each of the tag moieties with said at least one nanopore so as to thereby detecting the presence of the plurality of predetermined compounds in the sample.

This invention also provides a method for determining the quantity of each of a plurality of predetermined compounds in a sample which comprises:

- 25 (a) contacting the sample with a plurality of detectable tag moieties, wherein each detectable tag moiety comprises at least one detectable component, under conditions permitting
- 30 the tag moieties to attach to each of the predetermined compounds present in the sample to form detectably tagged predetermined compounds, wherein each tag moiety uniquely attaches to one of the predetermined compounds present in the sample and is distinguishable from any tag moiety attached to
- 35 any other predetermined compound present in the sample;

- 6 -

- (b) treating the sample from step (a) to form an electrolyte solution containing the detectably tagged predetermined compounds;
 - (c) contacting the electrolyte solution from step (b) with at least one nanopore, wherein said at least one nanopore comprises a means for binding each of the detectably tagged predetermined compounds present in the electrolyte solution, under conditions such that the detectably tagged predetermined compounds bind to said at least one nanopore;
 - (d) treating the detectably tagged predetermined compounds bound to said at least one nanopore so as to separate the tag moieties from the predetermined compounds;
 - (e) determining the amount of said at least one detectable component of each of the tag moieties so released as the tag moieties pass through said at least one nanopore; and
 - (f) determining, from the amounts of said detectable components determined in step (e), the quantity of each of the plurality of predetermined compounds present in the sample.
- 20 This invention also provides a method for determining the quantity of each of a plurality of predetermined compounds in a sample which comprises:
- (a) contacting the sample with a plurality of detectable tag moieties, wherein each detectable tag moiety comprises at least one detectable component, under conditions permitting the tag moieties to attach to each of the predetermined compounds present in the sample to form detectably tagged predetermined compounds, wherein each tag moiety uniquely attaches to one of the predetermined compounds present in the sample and is distinguishable from any tag moiety attached to any other predetermined compound present in the sample;
 - (b) treating the sample from step (a) to form an electrolyte solution containing the detectably tagged predetermined compounds;
 - (c) contacting the electrolyte solution from step (b) with at least one nanopore, wherein said at least one nanopore

- 7 -

- comprises a means for binding each of the detectably tagged predetermined compounds present in the electrolyte solution, under conditions such that the detectably tagged predetermined compounds bind to said at least one nanopore, and wherein each of the tag moieties of said bound detectably tagged predetermined compounds localizes within said at least one nanopore;
- (d) determining the amount of said at least one detectable component of each of the tag moieties with said at least one nanopore; and
- (e) determining, from the amounts of said detectable components determined in step (d), the quantity of each of the plurality of predetermined compounds present in the sample.
- 15 This invention further provides a method for detecting interaction of at least two predetermined compounds which comprises:
- (a) contacting at least a first predetermined compound present in a sample with a detectable tag moiety, wherein the detectable tag moiety comprises at least one detectable component, under conditions permitting the tag moiety to attach to the predetermined compound present in the sample to form a detectably tagged predetermined compound;
- (b) treating the sample from step (a) to form an electrolyte solution containing the detectably tagged predetermined compounds;
- (c) contacting the electrolyte solution from step (b) with at least one nanopore, wherein at least a second predetermined compound is immobilized on said at least one nanopore, under conditions permitting the binding of the detectably tagged predetermined compound with the second predetermined compound;
- (d) treating the detectably tagged predetermined compound bound to said at least one nanopore so as to separate the tag moiety from the predetermined compound; and
- (e) detecting said at least one detectable component of the tag moiety so released as the tag moiety passes through said at

- 8 -

least one nanopore so as to thereby detecting the interaction of the predetermined compounds.

- This invention even further provides a method for detecting interaction of at least two predetermined compounds which comprises:
- 5 (a) contacting at least a first predetermined compound present in a sample with a detectable tag moiety, wherein the detectable tag moiety comprises at least one detectable component, under conditions permitting the tag moiety to attach to the predetermined compound present in the sample to form a
10 detectably tagged predetermined compound;
 - (b) treating the sample from step (a) to form an electrolyte solution containing the detectably tagged predetermined compound;
 - 15 (c) contacting the electrolyte solution from step (b) with at least one nanopore, wherein at least a second predetermined compound is immobilized on said at least one nanopore, under conditions permitting the binding of the detectably tagged predetermined compound with the second predetermined compound,
20 and wherein the tag moiety of said detectably tagged predetermined compound bound to the second predetermined compound localizes within said at least one nanopore;
 - (d) detecting said at least one detectable component of the tag moiety with said at least one nanopore so as to thereby
25 detecting the interaction of the predetermined compounds.

This invention even further provides a method for detecting the presence of a plurality of predetermined compounds in a plurality of samples, comprising:

- 30 (a) contacting each sample with a nanopore-detectable tag moiety, under conditions permitting the tag moiety to attach to the plurality of predetermined compounds present in the sample to form detectably tagged predetermined compounds for that sample, which are distinguishable from any nanopore-detectable
35 tag moiety attached to similar predetermined compounds from any other sample;

- 9 -

- (b) mixing the resulting tagged compounds from the plurality of samples from step (a) and treating them in such a way as to form an electrolyte solution containing the set of detectably and distinguishably tagged predetermined compounds;
- 5 (c) contacting the electrolyte solution from step (b) with at least one nanopore, wherein said at least one nanopore comprises a means for specific analyte moiety binding to each of the detectably tagged predetermined compounds in the electrolyte solution, under conditions such that the
- 10 detectably tagged predetermined compounds bind to said at least one nanopore;
- (d) treating the detectably tagged predetermined compounds bound to said at least one nanopore so as to separate the tag moieties from the predetermined compounds; and
- 15 (e) detecting said at least one detectable component of each of the tag moieties so released as the tag moieties pass through said at least one nanopore so as to thereby detect the presence of the plurality of predetermined compounds in the plurality of samples.

20

- 10 -

BRIEF DESCRIPTION OF THE FIGURES

Figure 1. Multiplex quantitative protein sample analysis by detection of protein-protein interactions using nanopores and cleavable tags. **(A)** The analyte consists of a protein conjugated with a reporter tag via a cleavable linker (e.g., an antigen, antibody, protein A, or other molecule). **(B)** Detection of tagged proteins using α -hemolysin nanopores. (1) A mixture of two protein samples labeled with distinct tags is added to the nanopore-containing chip; (2) The analyte protein is captured on the *cis* side in the vicinity of the α -hemolysin nanopore via protein-protein interactions with a bound primary antibody, receptor, etc. (3) After protein binding and washing, the tags are cleaved off the proteins according to the nature of the cleavable linker; the cleaved tags are electrophoretically driven through the pore to produce unique current blockade signals which are analyzed to quantitatively and qualitatively analyze protein presence.

Figure 2. Correlation between conductance signatures and mass distribution of a polydisperse mixture of PEG oligomers (red) during their transit through a single nanopore as compared with a 29-ethylene glycol unit PEG polymer (blue). The conductance based mass spectrum clearly resolves the repeat units of PEG within the 30-50 monomer unit range [Robertson et al. 2007].

Figure 3. Synthesis of chemically cleavable linkers for protein labeling. Dotted lines indicate the cleavage site on treatment with TCEP.

Figure 4. Synthesis of a photochemically cleavable linker for protein labeling using a nitrobenzyl based linker as the starting material. Dotted lines indicate the cleavage site on treatment with TCEP.

Figure 5. Monodisperse coumarin-PEG derivatives of different size.

- 11 -

Figure 6. Discrimination of a mixture of coumarin-PEG tags of different sizes in a α -hemolysin nanopore at single molecule level. Left: time series data indicates that when PEG tags enter a single α -hemolysin channel, they cause current blockades that are characteristic in their duration and amplitude. Right: frequency histogram of the mean current blockade caused by individual molecules and expressed as ratio of current blocked vs open channel, which shows baseline resolution with a 10 kHz measurement bandwidth. The colored bars at the top represent the 6 σ distribution of the data, which suggests that a single tag could be discriminated with accuracy better than 1 in 300,000 events [Kumar et al. 2012].

Figure 7. Detection of interactions (an antibody and antigen interaction is used as an example) of tagged compounds using nanopores. **(A)** An analyte consists of a target protein conjugated with a reporter tag (e.g., an antigen, antibody, protein A, or other molecule). The tagged analyte binds to an antibody immobilized on top of the nanopore, the tag which has been designed to have a positive charge will enter the nanopore, causing a unique electrical current blockade, which can be used to identify the target molecule. **(B)** An analyte consists of a target protein conjugated with a reporter tag (e.g., an antigen, antibody, protein A, or other molecule) through a cleavable linker. The tagged analyte binds to an antibody immobilized on top of the nanopore, upon cleavage of the linker, the tag which has been designed to have a positive charge will pass through the nanopore, causing a unique electrical current blockade, which can be used to identify the target molecules.

- 12 -

DETAILED DESCRIPTION OF THE INVENTION

This invention provides a method for detecting the presence of a plurality of predetermined compounds in a sample which comprises:

- 5 (a) contacting the sample with a plurality of detectable tag moieties, wherein each detectable tag moiety comprises at least one detectable component, under conditions permitting the tag moieties to attach to each of the predetermined compounds present in the sample to form detectably tagged
10 predetermined compounds, wherein each tag moiety uniquely attaches to one of the predetermined compounds present in the sample and is distinguishable from any tag moiety attached to any other predetermined compound present in the sample;
- 15 (b) treating the sample from step (a) to form an electrolyte solution containing the detectably tagged predetermined compounds;
- 20 (c) contacting the electrolyte solution from step (b) with at least one nanopore, wherein said at least one nanopore comprises a means for binding each of the detectably tagged predetermined compounds present in the electrolyte solution, under conditions such that the detectably tagged predetermined compounds bind to said at least one nanopore;
- 25 (d) treating the detectably tagged predetermined compounds bound to said at least one nanopore so as to separate the tag moieties from the predetermined compounds; and
- 30 (e) detecting said at least one detectable component of each of the tag moieties so released as the tag moieties pass through said at least one nanopore so as to thereby detecting the presence of the plurality of predetermined compounds in the sample.

This invention also provides a method for detecting the presence of a plurality of predetermined compounds in a sample which comprises:

- 35 (a) contacting the sample with a plurality of detectable tag moieties, wherein each detectable tag moiety comprises at least one detectable component, under conditions permitting

- 13 -

- the tag moieties to attach to each of the predetermined compounds present in the sample to form detectably tagged predetermined compounds, wherein each tag moiety uniquely attaches to one of the predetermined compounds present in the sample and is distinguishable from any tag moiety attached to any other predetermined compound present in the sample;
- 5 (b) treating the sample from step (a) to form an electrolyte solution containing the detectably tagged predetermined compounds;
- 10 (c) contacting the electrolyte solution from step (b) with at least one nanopore, wherein said at least one nanopore comprises a means for binding each of the detectably tagged predetermined compounds present in the electrolyte solution, under conditions such that the detectably tagged predetermined
- 15 compounds bind to said at least one nanopore, and wherein each of the tag moieties of said bound detectably tagged predetermined compounds localizes within said at least one nanopore;
- 20 (d) detecting said at least one detectable component of each of the tag moieties with said at least one nanopore so as to thereby detecting the presence of the plurality of predetermined compounds in the sample.

This invention further provides a method for determining the quantity of each of a plurality of predetermined compounds in a sample which comprises:

- 25 (a) contacting the sample with a plurality of detectable tag moieties, wherein each detectable tag moiety comprises at least one detectable component, under conditions permitting
- 30 the tag moieties to attach to each of the predetermined compounds present in the sample to form detectably tagged predetermined compounds, wherein each tag moiety uniquely attaches to one of the predetermined compounds present in the sample and is distinguishable from any tag moiety attached to
- 35 any other predetermined compound present in the sample;

- 14 -

- (b) treating the sample from step (a) to form an electrolyte solution containing the detectably tagged predetermined compounds;
- 5 (c) contacting the electrolyte solution from step (b) with at least one nanopore, wherein said at least one nanopore comprises a means for binding each of the detectably tagged predetermined compounds present in the electrolyte solution, under conditions such that the detectably tagged predetermined compounds bind to said at least one nanopore;
- 10 (d) treating the detectably tagged predetermined compounds bound to said at least one nanopore so as to separate the tag moieties from the predetermined compounds;
- (e) determining the amount of said at least one detectable component of each of the tag moieties so released as the tag
- 15 moieties pass through said at least one nanopore; and
- (f) determining, from the amounts of said detectable components determined in step (e), the quantity of each of the plurality of predetermined compounds present in the sample.
- 20 This invention even further provides a method for determining the quantity of each of a plurality of predetermined compounds in a sample which comprises:
- (a) contacting the sample with a plurality of detectable tag moieties, wherein each detectable tag moiety comprises at
- 25 least one detectable component, under conditions permitting the tag moieties to attach to each of the predetermined compounds present in the sample to form detectably tagged predetermined compounds, wherein each tag moiety uniquely attaches to one of the predetermined compounds present in the sample and is distinguishable from any tag moiety attached to
- 30 any other predetermined compound present in the sample;
- (b) treating the sample from step (a) to form an electrolyte solution containing the detectably tagged predetermined compounds;
- 35 (c) contacting the electrolyte solution from step (b) with at least one nanopore, wherein said at least one nanopore

- 15 -

- comprises a means for binding each of the detectably tagged predetermined compounds present in the electrolyte solution, under conditions such that the detectably tagged predetermined compounds bind to said at least one nanopore, and wherein each of the tag moieties of said bound detectably tagged predetermined compounds localizes within said at least one nanopore;
- 5 (d) determining the amount of said at least one detectable component of each of the tag moieties with said at least one nanopore; and
- 10 (e) determining, from the amounts of said detectable components determined in step (d), the quantity of each of the plurality of predetermined compounds present in the sample.

15 In an embodiment of the invention, the methods further comprising a washing step to remove any detectably tagged predetermined compound which is not bound to said at least one nanopore prior to step (d).

In an embodiment of the invention, the predetermined compound is a protein. In a preferred embodiment, the predetermined compound is an antibody. In another embodiment of the invention, the predetermined compound is non-proteinaceous.

20

In an embodiment of the invention, the tag moiety comprises more than one detectable component. It is contemplated that each such detectable component is independently detectable.

25

In a further embodiment of the invention, the detectable component is selected from the group consisting of ethylene glycol, an amino acid, a carbohydrate, a peptide, a dye, a fluorescent compound, a chemiluminescent compound, a mononucleotide, a dinucleotide, a trinucleotide, a tetranucleotide, a pentanucleotide, a hexanucleotide, a polynucleotide, a nucleotide monophosphate, a nucleotide diphosphate, a nucleotide polyphosphate, an aliphatic acid, an aromatic acid, an unsubstituted alcohol or thiol, an alcohol or a thiol substituted with one or more halogens, a cyano

30

35

- 16 -

group, a nitro group, an alkyl group, an alkenyl group, an alkynyl group, and an azido group.

5 In certain embodiments, the detectable component of said tag moieties comprises a multiplicity of ethylene glycol units. In a further embodiment, the multiplicity of ethylene glycol units comprises 16, 20, 24, or 36 ethylene glycol units.

10 In an embodiment of the invention disclosed herein, the tag moiety attaches to the predetermined compound via a cleavable linker. In further embodiments of the invention, the cleavable linker is a photocleavable linker or a chemically cleavable linker.

15 In one embodiment, the photocleavable linker is a 2-nitrobenzyl linker. In another embodiment, the chemically cleavable linker is an azido linker. In an embodiment, UV light is used to cleave the photocleavable linker. Methods for production of cleavably capped and/or cleavably linked molecules are disclosed in U.S. Patent No. 6,664,079, which is hereby incorporated by reference.

20 In an embodiment of the claimed method, at least one of the predetermined compounds present in the sample is a protein and at least one of said tag moieties attaches to the carboxy or amino terminus of said protein.

25 In another embodiment, at least one of the predetermined compounds present in the sample is a protein and at least one of said tag moieties attaches to a lysine, an arginine, or a cysteine residue of said protein.

30 In a further embodiment of invention, the nanopore is a biological nanopore, a modified biological nanopore, or a synthetic nanopore. In certain embodiments, the nanopore is proteinaceous, in particular an alpha hemolysin (α -hemolysin).

35

- 17 -

In yet another embodiment, the nanopore is a solid-state nanopore. In a specific embodiment, the nanopore comprises graphene. It is contemplated that in certain embodiment the nanopore is in a membrane.

5 In an embodiment of the invention, the nanopore is part of an array of nanopores. In certain embodiments, each nanopore in said array comprises identical means for binding the detectably tagged predetermined compounds. In certain other embodiments, each nanopore in said array comprises different means for binding the
10 detectably tagged predetermined compounds.

In an embodiment, the means for binding the detectably tagged predetermined compounds is a protein, in particular an antibody. In another embodiment, the means for binding the detectably tagged
15 predetermined compounds is non-proteinaceous.

In an embodiment of the methods disclosed herein, a tag moiety is distinguishable from any other tag moiety based on blockade signature of said tag moiety detectable with said at least one
20 nanopore. In certain embodiments, the blockade signature is result of a change in current amplitude or conductance of said at least one nanopore.

In an embodiment, said at least one nanopore further comprising a
25 mean for ejecting said tag moiety from the nanopore.

This invention also provides a method for detecting interaction of at least two predetermined compounds which comprises:

(a) contacting at least a first predetermined compound present in
30 a sample with a detectable tag moiety, wherein the detectable tag moiety comprises at least one detectable component, under conditions permitting the tag moiety to attach to the predetermined compound present in the sample to form a detectably tagged predetermined compound;

- 18 -

- (b) treating the sample from step (a) to form an electrolyte solution containing the detectably tagged predetermined compounds;
- 5 (c) contacting the electrolyte solution from step (b) with at least one nanopore, wherein at least a second predetermined compound is immobilized on said at least one nanopore, under conditions permitting the binding of the detectably tagged predetermined compound with the second predetermined compound;
- 10 (d) treating the detectably tagged predetermined compound bound to said at least one nanopore so as to separate the tag moiety from the predetermined compound; and
- (e) detecting said at least one detectable component of the tag moiety so released as the tag moiety passes through said at least one nanopore so as to thereby detecting the interaction
15 of the predetermined compounds.

This invention further provides a method for detecting interaction of at least two predetermined compounds which comprises:

- 20 (a) contacting at least a first predetermined compound present in a sample with a detectable tag moiety, wherein the detectable tag moiety comprises at least one detectable component, under conditions permitting the tag moiety to attach to the predetermined compound present in the sample to form a detectably tagged predetermined compound;
- 25 (b) treating the sample from step (a) to form an electrolyte solution containing the detectably tagged predetermined compound;
- 30 (c) contacting the electrolyte solution from step (b) with at least one nanopore, wherein at least a second predetermined compound is immobilized on said at least one nanopore, under conditions permitting the binding of the detectably tagged predetermined compound with the second predetermined compound, and wherein the tag moiety of said detectably tagged predetermined compound bound to the second predetermined
35 compound localizes within said at least one nanopore;

- 19 -

(d) detecting said at least one detectable component of the tag moiety with said at least one nanopore so as to thereby detecting the interaction of the predetermined compounds.

5 In an embodiment, the method further comprises a washing step to remove any detectably tagged predetermined compound which is not bound to the second predetermined compound prior to step (d).

In an embodiment of the invention, at least one of the predetermined
10 compounds is a protein. In a specific embodiment, the protein is an antibody. In another embodiment of the invention, at least one of the predetermined compounds is non-proteinaceous.

In an embodiment, the tag moiety comprises more than one detectable
15 component. In a specific embodiment, each detectable component is independently detectable.

In an embodiment of the invention, each of said at least one detectable component of said tag moiety is selected from the group
20 consisting of ethylene glycol, an amino acid, a carbohydrate, a peptide, a dye, a fluorescent compound, a chemilluminiscent compound, a mononucleotide, a dinucleotide, a trinucleotide, a tetranucleotide, a pentanucleotide, a hexanucleotide, a polynucleotide, a nucleotide monophosphate, a nucleotide
25 diphosphate, a nucleotide polyphosphate, an aliphatic acid, an aromatic acid, an unsubstituted alcohol or thiol, an alcohol or a thiol substituted with one or more halogens, a cyano group, a nitro group, an alkyl group, an alkenyl group, an alkynyl group, and an azido group.

30 In an embodiment of the invention, said at least one detectable component of said tag moiety comprises a multiplicity of ethylene glycol units. In certain specific embodiments, said multiplicity of ethylene glycol units comprises 16, 20, 24, or 36 ethylene glycol
35 units.

- 20 -

In an embodiment of the invention, said tag moiety attaches to the predetermined compound via a cleavable linker. The cleavable linker could be a photocleavable linker or a chemically cleavable linker.

5 In an embodiment, the predetermined compound present in the sample is a protein and said tag moiety attaches to the carboxy or amino terminus of said protein. In another embodiment, the predetermined compound present in the sample is a protein and said tag moiety attaches to a lysine, an arginine, or a cysteine residue of said
10 protein.

In certain embodiment, said at least one nanopore is a biological nanopore, a modified biological nanopore, or a synthetic nanopore. In a specific embodiment, said at least one nanopore is
15 proteinaceous. In a preferred embodiment, said at least one nanopore is an alpha hemolysin (α -hemolysin).

In another embodiment, said at least one nanopore is a solid-state nanopore. In a further embodiment, the solid-state nanopore
20 comprises graphene.

In an embodiment, said at least one nanopore is in a membrane.

In another embodiment, said at least one nanopore is part of an array of nanopores. In some embodiments, each nanopore in said
25 array comprises an identical immobilized second predetermined compound. In other embodiments, each nanopore in said array comprises a different immobilized second predetermined compound.

In an embodiment, the second predetermined compound is a protein.
30 In a specific embodiment, the protein is an antibody.

In another embodiment, the second predetermined compound is non-proteinaceous.

- 21 -

In an embodiment, said tag moiety produces a blockade signature detectable with said at least one nanopore. In certain embodiment, the blockade signature is result of a change in current amplitude of said at least one nanopore. In certain other embodiment, the
5 blockade signature is result of a change in conductance of said at least one nanopore.

In an embodiment, the sample comprises a plurality of predetermined compounds. In such an embodiment, step (a) further comprises
10 contacting the plurality of predetermined compounds present in the sample with a plurality of detectable tag moieties, under conditions permitting the tag moieties to attach to each of the predetermined compounds present in the sample to form detectably tagged
15 predetermined compounds, wherein each tag moiety uniquely attaches to one of the predetermined compounds present in the sample and is distinguishable from any tag moiety attached to any other predetermined compound present in the sample.

In an embodiment, said at least one nanopore comprises a plurality
20 of predetermined compounds capable of binding to the plurality of detectably tagged predetermined compounds present in the sample.

In an embodiment, said at least one nanopore further comprising a means for ejecting said tag moiety from the nanopore. In a specific
25 embodiment, the means for ejecting said tag moiety from the nanopore consisting of a means to adjust electric field of said nanopore.

This invention also provides a detectable tag moiety for use in the methods disclosed herein which comprises a protein reactive group,
30 at least one detectable component, and a linker connecting the protein reactive group and the at least one detectable component.

In an embodiment, the linker is a cleavable linker. The cleavable linker could be a photocleavable linker or a chemically cleavable
35 linker.

- 22 -

This invention further provides a method for detecting the presence of a plurality of predetermined compounds in a plurality of samples, comprising:

- 5 (a) contacting each sample with a nanopore-detectable tag moiety, under conditions permitting the tag moiety to attach to the plurality of predetermined compounds present in the sample to form detectably tagged predetermined compounds for that sample, which are distinguishable from any nanopore-detectable tag moiety attached to similar predetermined compounds from
10 any other sample;
- (b) mixing the resulting tagged compounds from the plurality of samples from step (a) and treating them in such a way as to form an electrolyte solution containing the set of detectably and distinguishably tagged predetermined compounds;
- 15 (c) contacting the electrolyte solution from step (b) with at least one nanopore, wherein said at least one nanopore comprises a means for specific analyte moiety binding to each of the detectably tagged predetermined compounds in the electrolyte solution, under conditions such that the
20 detectably tagged predetermined compounds bind to said at least one nanopore;
- (d) treating the detectably tagged predetermined compounds bound to said at least one nanopore so as to separate the tag moieties from the predetermined compounds; and
- 25 (e) detecting said at least one detectable component of each of the tag moieties so released as the tag moieties pass through said at least one nanopore so as to thereby detect the presence of the plurality of predetermined compounds in the plurality of samples.

30

Each method and process described herein can be performed using compound with cleavable or noncleavable tags.

For the foregoing embodiments, each embodiment disclosed herein is
35 contemplated as being applicable to each of the other disclosed embodiments.

- 23 -

As used herein, and unless stated otherwise, each of the following terms shall have the definition set forth below.

"Antibody" shall include, without limitation, (a) an immunoglobulin molecule comprising two heavy chains and two light chains and which recognizes an antigen; (b) a polyclonal or monoclonal immunoglobulin molecule; and (c) a monovalent or divalent fragment thereof. Immunoglobulin molecules may derive from any of the commonly known classes, including but not limited to IgA, secretory IgA, IgG, IgE and IgM. IgG subclasses are well known to those in the art and include, but are not limited to, human IgG1, IgG2, IgG3 and IgG4. Antibodies can be both naturally occurring and non-naturally occurring. Furthermore, antibodies include chimeric antibodies, wholly synthetic antibodies, single chain antibodies, and fragments thereof. Antibodies may be human or nonhuman. Antibody fragments include, without limitation, Fab fragments, Fv fragments and other antigen-binding fragments.

"Nanopore" includes, for example, a structure comprising (a) a first and a second compartment separated by a physical barrier, which barrier has at least one pore with a diameter, for example, of from about 1 to 10 nm, and (b) a means for applying an electric field across the barrier so that a charged molecule such as DNA, nucleotide, nucleotide analogue, or tag, can pass from the first compartment through the pore to the second compartment. The nanopore ideally further comprises a means for measuring the electronic signature of a molecule passing through its barrier. The nanopore barrier may be synthetic or naturally occurring in part. Barriers can include, for example, lipid bilayers having therein α -hemolysin, oligomeric protein channels such as porins, and synthetic peptides and the like. Barriers can also include inorganic plates having one or more holes of a suitable size. Herein "nanopore", "nanopore barrier" and the "pore" in the nanopore barrier are sometimes used equivalently. It is understood that the electric field of a nanopore may be adjustable. It is also understood that a charged molecule such as DNA, nucleotide, nucleotide analogue, or tag, does

- 24 -

not need to pass from the first compartment through the pore to the second compartment in order to produce an electronic signature. Such electronic signature may be produced by localization of the molecule within the pore.

5

Nanopore devices are known in the art and nanopores and methods employing them are disclosed in U.S. Patent Nos. 7,005,264 B2; 7,846,738; 6,617,113; 6,746,594; 6,673,615; 6,627,067; 6,464,842; 6,362,002; 6,267,872; 6,015,714; 5,795,782; and U.S. Publication
10 Nos. 2004/0121525, 2003/0104428, and 2003/0104428, each of which are hereby incorporated by reference in their entirety.

"Blockade signature" of a molecule passing through a pore via application of an electronic field shall include, for example, the
15 duration of the nucleotide's passage through the pore together with the observed amplitude of current during that passage. Blockade signature for a molecule is envisioned and can be, for example, a plot of current (e.g. pA) versus time for the molecule to pass through the pore via application of an electric field.
20 Alternatively, blockade signature is also determinable for a molecule which does not pass through a pore. Blockade signature of such a molecule is also envisioned and can be, for example, a plot of current (e.g. pA) versus time for the molecule to enter into or pass adjacent to the pore. Herein "blockade signature", "blockade
25 signal", and "electronic signature" are sometime used equivalently.

A specific event diagram is constructed which is the plot of translocation time versus blockade current. This specific event diagram (also referred to as an blockade signature) is used to
30 distinguish molecules by single-channel recording techniques based on characteristic parameters such as translocation current, translocation duration, and their corresponding dispersions in the diagram.

35 As used herein, a "tag" or a "tag moiety" is any chemical group or molecule that is capable of producing a unique blockade signature

- 25 -

detectable with a nanopore. In some cases, a tag comprises one or more of ethylene glycol, an amino acid, a carbohydrate, a peptide, a dye, a fluorescent compound, a chemilluminiscent compound, a mononucleotide, a dinucleotide, a trinucleotide, a tetranucleotide, 5 a pentanucleotide, a hexanucleotide, a polynucleotide, a nucleotide monophosphate, a nucleotide diphosphate, a nucleotide polyphosphate, an aliphatic acid, an aromatic acid, an unsubstituted alcohol or thiol, an alcohol or a thiol substituted with one or more halogens, a cyano group, a nitro group, an alkyl group, an alkenyl group, an alkynyl group, an azido group, or a combination thereof. 10

As used herein, unless otherwise specified, a tag moiety which is different or distinguishable from the tag moiety of a referenced molecule means that the tag moiety has a different chemical 15 structure from the chemical structure of the other/referenced tag moiety. A tag moiety is different or distinguishable from the tag moiety of a referenced molecule could also mean that the tag moiety has a different blockade signature from the blockade signature of the other/referenced tag moiety. 20

As used herein, a tag which "localizes" within a pore is a tag located inside or adjacent to the pore. A tag which localizes within a pore does not necessarily pass through or translocate the pore. 25

As used herein, "proteinaceous" compound means any biopolymer formed from amino acids, such as peptides, proteins, antibodies, antigens, or a fragment or portion thereof. Such compound may be naturally occurring or non-naturally occurring. 30

As used herein, "alkyl" includes both branched and straight-chain saturated aliphatic hydrocarbon groups having the specified number of carbon atoms and may be unsubstituted or substituted. Thus, C1-Cn as in "C1-Cn alkyl" includes groups having 1, 2, ..., n-1 or n 35 carbons in a linear or branched arrangement. For example, a "C1-C5 alkyl" includes groups having 1, 2, 3, 4, or 5 carbons in a linear

- 26 -

or branched arrangement, and specifically includes methyl, ethyl, n-propyl, isopropyl, n-butyl, t-butyl, and pentyl

As used herein, "alkenyl" refers to a non-aromatic hydrocarbon group, straight or branched, containing at least 1 carbon to carbon double bond, and up to the maximum possible number of non-aromatic carbon-carbon double bonds may be present, and may be unsubstituted or substituted. For example, "C2-C5 alkenyl" means an alkenyl group having 2, 3, 4, or 5, carbon atoms, and up to 1, 2, 3, or 4, carbon-carbon double bonds respectively. Alkenyl groups include ethenyl, propenyl, and butenyl.

The term "alkynyl" refers to a hydrocarbon group straight or branched, containing at least 1 carbon to carbon triple bond, and up to the maximum possible number of non-aromatic carbon-carbon triple bonds may be present, and may be unsubstituted or substituted. Thus, "C2-C5 alkynyl" means an alkynyl group having 2 or 3 carbon atoms and 1 carbon-carbon triple bond, or having 4 or 5 carbon atoms and up to 2 carbon-carbon triple bonds. Alkynyl groups include ethynyl, propynyl and butynyl.

The term "substituted" refers to a functional group as described above such as an alkyl, or a hydrocarbyl, in which at least one bond to a hydrogen atom contained therein is replaced by a bond to non-hydrogen or non-carbon atom, provided that normal valencies are maintained and that the substitution(s) result(s) in a stable compound. Substituted groups also include groups in which one or more bonds to a carbon(s) or hydrogen(s) atom are replaced by one or more bonds, including double or triple bonds, to a heteroatom. Non-limiting examples of substituents include the functional groups described above, and for example, N, e.g. so as to form -CN.

It is understood that substituents and substitution patterns on the compounds of the instant invention can be selected by one of ordinary skill in the art to provide compounds that are chemically stable and that can be readily synthesized by techniques known in

- 27 -

the art, as well as those methods set forth below, from readily available starting materials. If a substituent is itself substituted with more than one group, it is understood that these multiple groups may be on the same carbon or on different carbons, so long as a stable structure results.

In choosing the compounds of the present invention, one of ordinary skill in the art will recognize that the various substituents, i.e. R_1 , R_2 , etc. are to be chosen in conformity with well-known principles of chemical structure connectivity.

In the compound structures depicted herein, hydrogen atoms, except on ribose and deoxyribose sugars, are generally not shown. However, it is understood that sufficient hydrogen atoms exist on the represented carbon atoms to satisfy the octet rule.

Where a range of values is provided, unless the context clearly dictates otherwise, it is understood that each intervening integer of the value, and each tenth of each intervening integer of the value, unless the context clearly dictates otherwise, between the upper and lower limit of that range, and any other stated or intervening value in that stated range, is encompassed within the invention. The upper and lower limits of these smaller ranges may independently be included in the smaller ranges, and are also encompassed within the invention, subject to any specifically excluded limit in the stated range. Where the stated range includes one or both of the limits, ranges excluding (i) either or (ii) both of those included limits are also included in the invention.

All combinations of the various elements described herein are within the scope of the invention. All sub-combinations of the various elements described herein are also within the scope of the invention.

This invention will be better understood by reference to the Experimental Details which follow, but those skilled in the art will

- 28 -

readily appreciate that the specific experiments detailed are only illustrative of the invention as described more fully in the claims which follow thereafter.

5 **Experimental Details And Discussions**

Building on the success of recent work on developing nanopore-based DNA sequencing, a multiplex digital protein detection and quantification technique using proteins tagged with cleavable PEG
10 molecules of different lengths and nanopore detection is developed. In this approach, a specific antibody is covalently attached in the vicinity of a nanopore. Protein analytes from different samples are labeled with cleavable tags that each produces a distinct current blockade signal in the α -hemolysin nanopore. After capture of the
15 tagged protein samples by the antibodies, the tags are cleaved off the proteins, and identified at the single molecule level as they traverse the nanopore.

20 *I. Design and Synthesize of Cleavable Protein-Labeling Tags*

A set of modified poly(ethylene glycol) (PEG) tags has been previously developed for single-molecule sequencing and demonstrated to generate unique current blockade signatures when traversing a nanopore. Synthetic derivatives of these tags are developed, which
25 are capable of efficiently conjugating with protein analytes or antibodies via photocleavable or chemically cleavable linkers to allow their subsequent identification in the nanopore.

Design of the cleavable tag structure is shown in **Fig. 1A** and
30 consists of three distinct components: a protein reactive group, a cleavable linker and a nanopore tag. A This design is based on studies of the transit of PEG molecules of different molecular size through the α -hemolysin nanopore [Robertson et al. 2007] and experience in designing cleavable linkers for DNA sequencing by
35 synthesis with reversible terminators bearing fluorescent tags [Guo et al. 2008 and Seo et al. 2005]. Robertson et al. demonstrated

- 29 -

that the transit of polydisperse PEG molecules through a *Staphylococcus aureus* α -hemolysin nanopore causes distinct current blockade states that generate characteristic conductance signatures at single-molecule level (**Fig. 2**), which correlate with the
5 molecular weight of individual PEG molecules. In addition to conductance reduction, the residence time within the nanopore increases linearly with PEG molecular weight [Robertson et al. 2007]. Both of these characteristics can be employed to design recognizable tags for protein labeling as demonstrated below.

10

A scheme of the method for detection of tagged proteins using α -hemolysin nanopores is shown in **Fig. 1B**. In this scheme, a mixture of two protein samples labeled with distinct tags is added to the nanopore-containing chip; the analyte protein is captured on the *cis*
15 side in the vicinity of the α -hemolysin nanopore via protein-protein interactions with a bound primary antibody or receptor; after protein binding and washing, the tags are cleaved off the proteins according to the nature of the cleavable linker; the cleaved tags are electrophoretically driven through the pore to produce unique
20 current blockade signals which are analyzed to quantitatively and qualitatively determine protein presence. A given tag traversing the nanopore causes a current blockade signature of characteristic amplitude and duration, resulting in tag recognition and quantification. As each blockade event occurs within a millisecond,
25 tens of thousands of tags can be analyzed in a short time, their counts providing a digital measure of the analytes' relative abundance in the samples at the single molecule level.

There are a number of chemical transformations which can be used for
30 protein conjugation with the nanopore tags. Beside the C- or N-terminus of proteins, other modification sites include lysines and arginines, which can be modified by treatment with NHS esters or isothiocyanates, and cysteine thiols modified with maleimides or α -halo-carbonyl compounds.

35

- 30 -

In this instance, the amino group on lysines or the thiol group of cysteines is modified with a cleavable linker-attached PEG molecule. Since most proteins have several lysines, but relatively few unreacted cysteines, conditions can be optimized for a range of label densities. The linker can be either a chemically or photochemically cleavable linker that would react with the lysines or cysteines on proteins and release PEG molecules after cleavage with tris(2-carboxyethyl)phosphine (TCEP) or with light at 350 nm, respectively; released PEG molecules are identified by the current blockade signal on the nanopore.

a) Chemically Cleavable Linkers

An example of synthesis of a chemically cleavable linker is shown in **Fig. 3**. The azido-linker (1) [Guo et al. 2008] is reacted with an appropriate size PEG-NHS ester (2) to provide an azido-cleavable linker-PEG molecule (3). The acid group in (3) is activated with DSC/DMF to provide the PEG-cleavable linker-NHS ester (4) which is reacted with the lysine amino group in the protein to provide a labeled protein similar to (7).

Alternatively, compound (4) can be derivatized with functional groups (maleimides or iodoacetamides) which react efficiently with the thiol group on cysteine. Thus compound (4), on reaction with amino-maleimide or iodoacetamide-ethylamine, yields compounds (5) and (6), respectively. These compounds, upon reaction with the cysteine thiol group, provide labeled proteins similar in structure to (8) and (9), respectively. Upon treatment with TCEP, the azido linked PEG-protein molecules cleave to release PEG molecules of structure (10), which pass through the nanopore and produce the resulting current blockade signal.

b) Photochemically Cleavable Linkers

Photochemically cleavable linkers are synthesized using similar methodology as the synthesis of chemically cleavable linkers, except that a nitrobenzyl based linker (11) [Seo et al. 2005] is used as the starting material, as shown in **Fig. 4**. Remainder of the

- 31 -

synthesis scheme is similar to that of chemically cleavable linkers, and provides photochemically cleavable linkers **(13)-(15)**. These linkers, after labeling lysines or cysteines on proteins and photochemical cleavage, provide PEG-linkers similar to structure **(16)**.

II. Test Tags For Coupling, Cleavage, And Nanopore Discrimination

Attaching the tags to either streptavidin or to the Fc portion of a rabbit IgG antibody allows use of biotin or anti-rabbit antibodies as capture agents in later analyses. Gel shift assays are used to confirm attachment and cleavage of the tags. Different molecular tags synthesized are evaluated for ability to quantitatively attach to protein samples and allow subsequent cleavage at high efficiency in order to capture signatures in the nanopore. Based on the results of these experiments, the best tag-linker configurations are selected for labeling and detection, or redesigned to improve their performance, if necessary.

In particular, PEG tags of various lengths coupled with protein-reactive groups via different cleavable linkers are tested for performance efficiency in coupling and cleavage, as well as for discrimination of their current blockade signals in a nanopore. The goal of these experiments is to select the optimal tags with coupling and cleavage chemistries effective for the analyte proteins and chemically compatible with the nanopore environment and electronic detection.

To test protein-protein interactions, tag cleavage, and nanopore detection, commercially available streptavidin (Life Technologies) and polyclonal rabbit IgG (Thermo Fisher Scientific) are used as the protein analytes. Exclusion of any particular antigen from these studies simplifies the analysis. By using highly standardized reactions with primary and secondary antibody interactions instead, the conditions found here can be immediately adapted for full antigen-antibody sandwich methods.

- 32 -

PEG tags are attached to the amino group on lysine of the analyte proteins using standard NHS chemistry, and the optimal conjugation conditions are determined using protein gel shift assays. The protein labeling extent is found using PEG tags derivatized with coumarin to allow spectrophotometric detection at 350 nm where protein absorption of light is low. Several coumarin-PEG derivative compounds have been successfully synthesized for nanopore blockade evaluation (**Fig. 5**).

10

Cleavage reaction conditions are determined in a similar way. Tag-labeled proteins (streptavidin or rabbit IgG) are treated with TCEP or, in the case of photocleavable tags, irradiated with near-UV light ($\lambda \sim 365$ nm). Cleavage is evaluated for completion by protein gel shift and spectroscopic measurements.

15

To evaluate the discrimination of the cleaved tags with the nanopore the tagged proteins are captured on a solid phase biotin-coated 96-well plates (Thermo Fisher Scientific) for PEG-tagged streptavidin; and on a solid phase goat-anti-rabbit polyclonal antibody-coated 96-well plates (Thermo Fisher Scientific) for PEG-tagged rabbit IgG. Follow by inducing release of the tags and counting of the released tags in bulk at single molecule level using a solitary α -hemolysin nanopore patch clamp electrode assembly by dipping the patch clamp directly into the wells of the plate.

25

Discrimination for four coumarin-PEG derivatives of the sizes of 16, 20, 24, and 36 monomer units is shown in **Fig. 6**, an 8-second current trace produced by a mixture of these PEG oligomers transiting the solitary nanopore. A time series data shown on left side indicates that when PEG tags enter a single α -hemolysin channel, they cause current blockades that are characteristic in their duration and amplitude. On right side, a frequency histogram of the mean current blockade caused by individual molecules and expressed as ratio of current blocked vs open channel shows baseline resolution with a 10 kHz measurement bandwidth. The colored bars at the top represent

30

35

- 33 -

the 6 σ distribution of the data, which suggests that a single tag could be discriminated with accuracy better than 1 in 300,000 events.

5 As the analysis of the blockage signals illustrates, data show excellent discrimination of the selected PEG tags over a 6 σ confidence interval and also demonstrate the speed and accuracy of the quantitative analysis of the released tags. In particular, there is a substantial gap between PEG sizes of 24 and 36 ethylene
10 glycol units. Based on this result, one could select different length PEGs (available commercially from Quanta Biodesign Ltd. or other suppliers) and monitor their nanopore blockade signals for additional tags within this molecular size range that retain a similar level of discrimination.

15 Noteworthy as well is the remarkably low expected noise originating from the transit of non-tag molecules through the nanopore due to the high selectivity of the nanopore aperture. Majority of the polymers, including proteins, are incapable of entering the pore,
20 and the smaller molecules produce barely detectable blockage of the pore. Thus, rational chemical design of cleavable linkers and selection of appropriate PEG polymers generate a series of optimal tags suitable for discrimination of multiplex samples using nanopore.

25

III. Multiplex Quantification Of Protein Samples Labeled With Different Tags Using Nanopores

30 A library of PEG tags that yield distinct nanopore signals is used to test the multiplex quantification scheme by capturing protein samples labeled with different tags using ligands or antibodies. Sensitivity and dynamic range of the nanopore detection method is
35 also determined and compared with fluorescent and colorimetric detection.

- 34 -

Sample multiplexing can also be accomplished by binding such a library of tags that yield distinct nanopore signals to the same protein derived from different samples, different individuals, or
5 different samples or individuals before, during and following some change, treatment or perturbation. Again, as for protein multiplexing, sensitivity and dynamic range can be determined and compared with fluorescent and colorimetric methods.

10 The experiments involve selecting a series of well discriminated tags, attaching the tags to streptavidin (SA) or rabbit antibodies, mixing the tagged proteins in different ratios, and preparing serial dilutions of the mixtures. The nanopore tags are also mixed with SA-alkaline phosphatase (Thermo Fisher Scientific), SA-AlexaFluor
15 488 (Life Technologies) or, in the case of the antibody reaction, with similarly labeled rabbit IgG antibodies (available from Thermo Fischer Scientific).

Biotin or anti-rabbit IgG antibodies are attached to the wells of
20 96-well plates to perform the binding and collection of the cleaved tags for detection in the nanopore. After performing the analyte capture using appropriately coated 96-well plates and induce the tag cleavage, tag quantification is performed using the patch clamp technique with a single nanopore and quantitation is performed in
25 the same wells using fluorescent or colorimetric detection.

Comparing the tag counts on the *trans* side of the nanopore with the predefined proportions on the *cis* side allows evaluation of the concentration estimation using the nanopore, estimate the dynamic
30 and quantitative (free of Poisson noise) range of the method in the presence of different tags, and directly compare them with the sensitivity and the performance of other non-nanopore based methods, which also serve as internal controls for analyte capture.

35 Mixtures of proteins with different tags are detected in the expected ratios. Differences in tag-labeling efficiencies

- 35 -

associated with variations in amino acid compositions among different antigens do not present an obstacle for the quantification because differences in the number of tags coupled with each antigen type can be inferred from their amino acid sequence, verified
5 experimentally, and corrected in downstream analysis.

IV. Detection of Molecular Interactions

Figure 7 illustrates detection of molecular interactions, using an
10 antibody and antigen interaction as an example, with nanopores and tags. As shown in **Figure 7(A)**, an analyte consists of a target protein conjugated with a reporter tag (e.g., an antigen, antibody, protein A, or other molecule). When the tagged analyte binds to the antibody immobilized on the top of the nanopore, the tag which has
15 been designed to have a positive charge will enter the nanopore, causing a unique electrical current blockade, which can be used to identify the target molecules. As shown in **Figure 7(B)**, an analyte also could consist of a target protein conjugated with a reporter tag (e.g., an antigen, antibody, protein A, or other molecule)
20 through a cleavable linker. When the tagged analyte binds to the antibody immobilized on the top of the nanopore, upon cleavage of the linker, the tag which has been designed to have a positive charge will pass through the nanopore, causing a unique electrical current blockade, which can be used to identify the target
25 molecules.

Discussion

The explosive development of massively parallel nucleic acid
30 sequencing and quantification methodology and associated cost reductions during the last decade have revolutionized mutation and polymorphism detection, transcriptional regulation of gene expression, and numerous other genome scale studies. Indeed, the high-throughput nature of these technologies has created entire new
35 disciplines in functional genomics and systems biology. However, analysis of gene regulation at the level of protein synthesis, like

- 36 -

proteomics in general, lags behind nucleic acid analysis in its throughput, sensitivity and automation. Among the major complications in developing high throughput proteomics are relatively poor stability of proteins as analytes, high
5 heterogeneity of protein analytes, and the requirement for a much wider dynamic range of detection with increased demand for sensitivity approaching the single molecule detection level. This need is not easily met by existing technologies which use fluorescent or colorimetric measurements.

10

Building on the success of recent work on developing nanopore-based DNA sequencing, a technique for multiplex digital protein detection and quantification using PEG-tagged proteins and nanopore array detection is developed. In an example of this approach, a specific
15 antibody is covalently attached in the vicinity of a nanopore. Analyte proteins from different samples are labeled with cleavable tags that produce a distinct current blockade signal in the α -hemolysin nanopore. After capture of the tagged protein samples by the antibodies, the tags are cleaved off the proteins and identified
20 at the single molecule level electronically as they traverse the nanopore. The advantage of this approach is generation of a digital output for the protein quantification with single molecule detection capability; by setting arbitrary upper detection limits, the technique permits characterization of either a wide range of protein
25 concentrations or discrimination of very small changes in protein levels. Thus, the technique has scalability and miniaturization comparable to the most advanced current single molecule genomic methods. The nanopore detection technique based on cleavable molecular tags addresses shortcomings of existing protein arrays
30 approach, which is based on antibody interaction with fluorescently labeled antigens or secondary antibodies. While there are currently several applications of nanopore-based analytics, application of nanopore for quantification of protein-protein interactions with multiplex tags is novel.

35

- 37 -

These experiments elucidate major parameters for this protein detection method, estimate the method's sensitivity, and define operational conditions utilizing a library of cleavable molecular tags and a single α -hemolysin nanopore. The detection system use a streptavidin-biotin system and a sandwich antibody detection scheme in which the primary antibody is attached to a surface and the cleavable poly(ethylene glycol) (PEG) tag is attached to the antigen (or to a secondary antibody for more efficient multi-nanopore arrays). Result obtained with this method is compared with result of ELISAs with fluorescent and colorimetric detection.

Developed based on existing method using nanopores and cleavable tags for DNA sequencing, this protein detection method consists of (1) labeling protein analytes, antigens or secondary antibodies with photocleavable or chemically cleavable tags; (2) reacting the resulting molecules with nanopore arrays in which antibodies specific to each protein analyte are associated with specific individual nanopores; and (3) after binding and washing, cleaving the tags and identifying them during their traversal of the adjacent nanopore. Different tags producing distinct current blockade signatures define the sample source, while registering the number of blockade events quantifies the number of tags of each type transiting the nanopore. All other assay components either do not enter the nanopores and/or are washed away between assay steps.

The three steps outlined above: cleavable tag design, capture of tagged protein samples, and detection using electronic identification of cleaved tags at single molecule resolution with a nanopore, bring the analysis to the nanoscale while at the same time permitting scale-up of the overall throughput for simplified whole-proteome quantitation.

This protein detection method has significant advantages over current fluorescent antibody-based proteomics arrays and may form a basis for creating protein detection sensors with the advantages of traditional semiconductor technologies using nanopore-based sensor

- 38 -

arrays using inexpensive and highly scalable standardized semiconductor technologies being developed.

5 Additionally, regardless of the type of tag, an important aspect of this method involves labeling different compounds with tags having distinguishable current blockade signatures generated as result of the tags traverse the nanopore. This differs substantially from existing protein detection system in which an aptamer binds to a protein to form a complex in solution on the *cis* side of a nanopore,
10 where the complex are attracted to the nanopore by the aptamer, and generates a two-step blockade current, first due to the aptamer entering the pore and the second due to the protein which blocks but does not enter the pore and is subsequently cleaved to allow the next aptamer to enter. The existing process requires measuring, and
15 account for, current blockade signatures for unbound aptamer and aptamer-free protein in any quantitative measurement schemes.

This approach generates a digital output for the protein quantification with near-single molecule per cell detection
20 capability. By setting arbitrary upper detection limits, the technique permits characterization of a wide range of protein concentrations or discrimination of very small changes in protein abundance. The technique has scalability and miniaturization comparable to the most advanced current single molecule genomic
25 methods (e.g., direct RNA-Seq). Eventually, the approach can be adopted to use nanopore arrays capable of characterizing hundreds to thousands of proteins at once from multiple tissues at the single-molecule level rapidly and at low cost, and follow changes in these proteins during different pathological states.

30 These experiments establish the method and exploring its range of performance. Once established, the method can be applied as a detection method of choice with existing protein microarrays and for other proteomics methods where demands for single molecule detection
35 and a simplified electronic readout are high. Potential obstacles inherent to proteomics beyond detection, such as differences in

- 39 -

labeling extent and variable affinities, can be overcome using more reliable and uniform binding counterparts such as aptamers or affibodies, build experimentally based correction methods for differences in labeling, design secondary detection methods, and
5 more elaborate sandwich schemes to enhance sensitivity. This method has advantages over fluorescent and colorimetric detection in that it uses standard semiconductor arrays with digital rather than analog detection output permitting counting of events at the single molecule level. It has virtually unrestricted dynamic range, low
10 cost, and high scalability from detection of a handful of specific proteins to whole proteome analysis. This method is also capable of complementing, at the level of proteomics, the advances in high throughput DNA and RNA techniques in such diverse areas of analysis as translational regulation of gene expression, immune response
15 biomarker identification, protein-protein interaction profiling, translational regulation by miRNAs, and non-PCR methods for detection of pathogens.

- 40 -

REFERENCES

- Angenendt P., "Progress in protein and antibody microarray technology", Drug Discov. Today 2005; 10(7):503-511.
- 5 Burnette WN. "'Western blotting': electrophoretic transfer of proteins from sodium dodecyl sulfate--polyacrylamide gels to unmodified nitrocellulose and radiographic detection with antibody and radioiodinated protein A", Anal. Biochem. 1981; 112(2):195-203.
- 10 Bussow K, Cahill D, Nietfeld W, Bancroft D, Scherzinger E, Lehrach H, Walter G. "A method for global protein expression and antibody screening on high-density filters of an arrayed cDNA library", Nucleic Acids Res. 1998; 26(21):5007-5008.
- 15 Cahill DJ. "Protein and antibody arrays and their medical applications", J. Immunol. Methods, 2001; 250(1-2):81-91.
- de Wildt RM, Mundy CR, Gorick BD, Tomlinson IM. "Antibody arrays for high-throughput screening of antibody-antigen interactions", Nat. Biotechnol. 2000; 18(9):989-994.
- 20 Engvall E, Perlmann P. "Enzyme-linked immunosorbent assay (ELISA), Quantitative assay of immunoglobulin", G. Immunochemistry 1971; 8(9):871-874.
- 25 Gygi SP, Rist B, Gerber SA, Turecek F, Gelb MH, Aebersold R. "Quantitative analysis of complex protein mixtures using isotope-coded affinity tags", Nat. Biotechnol. 1999; 17(10):994-999.
- 30 Guerrera IC, Kleiner O. "Application of mass spectrometry in proteomics", Biosci. Rep. 2005; 25(1-2):71-93.
- 35 Guo J, Xu N, Li Z, Zhang S, Wu J, Kim DH, Sano Marma M, Meng Q, Cao H, Li X, Shi S, Yu L, Kalachikov S, Russo JJ, Turro NJ, Ju J. "Four-color DNA sequencing with 3'-O-modified nucleotide reversible

- 41 -

terminators and chemically cleavable fluorescent dideoxynucleotides", Proc. Natl. Acad. Sci. USA 2008; 105(27):9145-9150

- 5 Klose J. "Protein mapping by combined isoelectric focusing and electrophoresis of mouse tissues: a novel approach to testing for induced point mutations in mammals", Humangenetik 1975; 26(3):231-243.
- 10 Kumar S, Tao C, Chien M, Hellner B, Balijepalli A, Robertson JW, Li Z, Russo JJ, Reiner JE, Kasianowicz JJ, Ju J. "PEG-labeled nucleotides and nanopore detection for single molecule DNA sequencing by synthesis", Sci. Rep. 2012; 2:684.
- 15 O'Farrell PH. "High resolution two-dimensional electrophoresis of proteins", J. Biol. Chem. 1975; 250(10):4007-4021.
- Robertson JW, Rodrigues CG, Stanford VM, Robinson KA, Krasilnikov OV, Kasianowicz JJ. "Single-molecule mass spectrometry in solution
- 20 using a solitary nanopore", Proc. Natl. Acad. Sci. USA 2007; 104(20):8207-8211.
- Seo TS, Bai X, Kim DH, Meng Q, Shi S, Ruparel H, Li Z, Turro NJ, Ju J. "Four-color DNA sequencing by synthesis on a chip using
- 25 photocleavable fluorescent nucleotides", Proc. Natl. Acad. Sci. USA 2005; 102(17):5926-5931.
- Yalow RS, Berson SA. "Immunoassay of endogenous plasma insulin in man", J. Clin. Invest. 1960; 39:1157-1175.

30

- 42 -

WHAT IS CLAIMED IS:

1. A method for detecting the presence of a plurality of predetermined compounds in a sample which comprises:
 - (a) contacting the sample with a plurality of detectable tag moieties, wherein each detectable tag moiety comprises at least one detectable component, under conditions permitting the tag moieties to attach to each of the predetermined compounds present in the sample to form detectably tagged predetermined compounds, wherein each tag moiety uniquely attaches to one of the predetermined compounds present in the sample and is distinguishable from any tag moiety attached to any other predetermined compound present in the sample;
 - (b) treating the sample from step (a) to form an electrolyte solution containing the detectably tagged predetermined compounds;
 - (c) contacting the electrolyte solution from step (b) with at least one nanopore, wherein said at least one nanopore comprises a means for binding each of the detectably tagged predetermined compounds present in the electrolyte solution, under conditions such that the detectably tagged predetermined compounds bind to said at least one nanopore;
 - (d) treating the detectably tagged predetermined compounds bound to said at least one nanopore so as to separate the tag moieties from the predetermined compounds; and
 - (e) detecting said at least one detectable component of each of the tag moieties so released as the tag moieties pass through said at least one nanopore so as to thereby detecting the presence of the plurality of predetermined compounds in the sample.
2. A method for detecting the presence of a plurality of predetermined compounds in a sample which comprises:

- 43 -

- (a) contacting the sample with a plurality of detectable tag moieties, wherein each detectable tag moiety comprises at least one detectable component, under conditions permitting the tag moieties to attach to each of the predetermined compounds present in the sample to form detectably tagged predetermined compounds, wherein each tag moiety uniquely attaches to one of the predetermined compounds present in the sample and is distinguishable from any tag moiety attached to any other predetermined compound present in the sample;
 - (b) treating the sample from step (a) to form an electrolyte solution containing the detectably tagged predetermined compounds;
 - (c) contacting the electrolyte solution from step (b) with at least one nanopore, wherein said at least one nanopore comprises a means for binding each of the detectably tagged predetermined compounds present in the electrolyte solution, under conditions such that the detectably tagged predetermined compounds bind to said at least one nanopore, and wherein each of the tag moieties of said bound detectably tagged predetermined compounds localizes within said at least one nanopore;
 - (d) detecting said at least one detectable component of each of the tag moieties with said at least one nanopore so as to thereby detecting the presence of the plurality of predetermined compounds in the sample.
3. A method for determining the quantity of each of a plurality of predetermined compounds in a sample which comprises:
- (a) contacting the sample with a plurality of detectable tag moieties, wherein each detectable tag moiety comprises at least one detectable component, under conditions permitting the tag moieties to attach to each of the predetermined compounds present in the sample to form detectably tagged predetermined compounds, wherein each tag moiety uniquely attaches to one of the predetermined

- 44 -

- compounds present in the sample and is distinguishable from any tag moiety attached to any other predetermined compound present in the sample;
- (b) treating the sample from step (a) to form an electrolyte solution containing the detectably tagged predetermined compounds;
 - (c) contacting the electrolyte solution from step (b) with at least one nanopore, wherein said at least one nanopore comprises a means for binding each of the detectably tagged predetermined compounds present in the electrolyte solution, under conditions such that the detectably tagged predetermined compounds bind to said at least one nanopore;
 - (d) treating the detectably tagged predetermined compounds bound to said at least one nanopore so as to separate the tag moieties from the predetermined compounds;
 - (e) determining the amount of said at least one detectable component of each of the tag moieties so released as the tag moieties pass through said at least one nanopore; and
 - (f) determining, from the amounts of said detectable components determined in step (e), the quantity of each of the plurality of predetermined compounds present in the sample.
4. A method for determining the quantity of each of a plurality of predetermined compounds in a sample which comprises:
- (a) contacting the sample with a plurality of detectable tag moieties, wherein each detectable tag moiety comprises at least one detectable component, under conditions permitting the tag moieties to attach to each of the predetermined compounds present in the sample to form detectably tagged predetermined compounds, wherein each tag moiety uniquely attaches to one of the predetermined compounds present in the sample and is distinguishable from any tag moiety attached to any other predetermined compound present in the sample;

- 45 -

- (b) treating the sample from step (a) to form an electrolyte solution containing the detectably tagged predetermined compounds;
 - (c) contacting the electrolyte solution from step (b) with at least one nanopore, wherein said at least one nanopore comprises a means for binding each of the detectably tagged predetermined compounds present in the electrolyte solution, under conditions such that the detectably tagged predetermined compounds bind to said at least one nanopore, and wherein each of the tag moieties of said bound detectably tagged predetermined compounds localizes within said at least one nanopore;
 - (d) determining the amount of said at least one detectable component of each of the tag moieties with said at least one nanopore; and
 - (e) determining, from the amounts of said detectable components determined in step (d), the quantity of each of the plurality of predetermined compounds present in the sample.
5. The method of any one of claims 1-4, further comprising a washing step to remove any detectably tagged predetermined compound which is not bound to said at least one nanopore prior to step (d).
6. The method of any one of claims 1-5, wherein the predetermined compound is a protein.
7. The method of claim 6, wherein the protein is an antibody.
8. The method of any one of claims 1-5, wherein the predetermined compound is non-proteinaceous.
9. The method of any one of claims 1-8, wherein at least one of said tag moieties comprises more than one detectable component.

- 46 -

10. The method of any one of claims 1-9, wherein each detectable component is independently detectable.
11. The method of any one of claims 1-10, wherein each of said at least one detectable component of said tag moieties is selected from the group consisting of ethylene glycol, an amino acid, a carbohydrate, a peptide, a dye, a fluorescent compound, a chemilluminiscent compound, a mononucleotide, a dinucleotide, a trinucleotide, a tetranucleotide, a pentanucleotide, a hexanucleotide, a polynucleotide, a nucleotide monophosphate, a nucleotide diphosphate, a nucleotide polyphosphate, an aliphatic acid, an aromatic acid, an unsubstituted alcohol or thiol, an alcohol or a thiol substituted with one or more halogens, a cyano group, a nitro group, an alkyl group, an alkenyl group, an alkynyl group, and an azido group.
12. The method of any one of claims 1-11, wherein said at least one detectable component of said tag moieties comprises a multiplicity of ethylene glycol units.
13. The method of any one of claims 1-12, wherein said multiplicity of ethylene glycol units comprises 16, 20, 24, or 36 ethylene glycol units.
14. The method of any one of claims 1-13, wherein at least one of said tag moieties attaches to the predetermined compound via a cleavable linker.
15. The method of claim 14, wherein the cleavable linker is a photocleavable linker.
16. The method of claim 14, wherein the cleavable linker is a chemically cleavable linker.

- 47 -

17. The method of any one of claims 1-16, wherein at least one of the predetermined compounds present in the sample is a protein and at least one of said tag moieties attaches to the carboxy or amino terminus of said protein.
18. The method of any one of claims 1-17, wherein at least one of the predetermined compounds present in the sample is a protein and at least one of said tag moieties attaches to a lysine, an arginine, or a cysteine residue of said protein.
19. The method of any one of claims 1-18, wherein said at least one nanopore is a biological nanopore, a modified biological nanopore, or a synthetic nanopore.
20. The method of claim 19, wherein said at least one nanopore is proteinaceous.
21. The method of any one of claims 1-20, wherein said at least one nanopore is an alpha hemolysin (α -hemolysin).
22. The method of any one of claims 1-18, wherein said at least one nanopore is a solid-state nanopore.
23. The method of claim 22, wherein the solid-state nanopore comprises graphene.
24. The method of any one of claims 1-23, wherein said at least one nanopore is in a membrane.
25. The method of any one of claims 1-24, wherein said at least one nanopore is part of an array of nanopores.
26. The method of claim 25, wherein each nanopore in said array comprises identical means for binding the detectably tagged predetermined compounds.

- 48 -

27. The method of claim 25, wherein each nanopore in said array comprises different means for binding the detectably tagged predetermined compounds.
28. The method of any one of claims 1-27, wherein the means for binding the detectably tagged predetermined compounds is a protein.
29. The method of claim 28, wherein the protein is an antibody.
30. The method of any one of claim 1-27, wherein the means for binding the detectably tagged predetermined compounds is non-proteinaceous.
31. The method of any one of claims 1-30, wherein each tag moiety is distinguishable from any other tag moiety based on blockade signature of said tag moiety detectable with said at least one nanopore.
32. The method of claim 31, wherein the blockade signature is result of a change in current amplitude of said at least one nanopore.
33. The method of claim 31, wherein the blockade signature is result of a change in conductance of said at least one nanopore.
34. The method of any one of claims 1-33, wherein said at least one nanopore further comprising a means for ejecting said tag moiety from the nanopore.
35. A method for detecting interaction of at least two predetermined compounds which comprises:
 - (a) contacting at least a first predetermined compound present in a sample with a detectable tag moiety, wherein the detectable tag moiety comprises at least one

- 49 -

detectable component, under conditions permitting the tag moiety to attach to the predetermined compound present in the sample to form a detectably tagged predetermined compound;

- (b) treating the sample from step (a) to form an electrolyte solution containing the detectably tagged predetermined compounds;
- (c) contacting the electrolyte solution from step (b) with at least one nanopore, wherein at least a second predetermined compound is immobilized on said at least one nanopore, under conditions permitting the binding of the detectably tagged predetermined compound with the second predetermined compound;
- (d) treating the detectably tagged predetermined compound bound to said at least one nanopore so as to separate the tag moiety from the predetermined compound; and
- (e) detecting said at least one detectable component of the tag moiety so released as the tag moiety passes through said at least one nanopore so as to thereby detecting the interaction of the predetermined compounds.

36. A method for detecting interaction of at least two predetermined compounds which comprises:

- (a) contacting at least a first predetermined compound present in a sample with a detectable tag moiety, wherein the detectable tag moiety comprises at least one detectable component, under conditions permitting the tag moiety to attach to the predetermined compound present in the sample to form a detectably tagged predetermined compound;
- (b) treating the sample from step (a) to form an electrolyte solution containing the detectably tagged predetermined compound;
- (c) contacting the electrolyte solution from step (b) with at least one nanopore, wherein at least a second predetermined compound is immobilized on said at least

- 50 -

one nanopore, under conditions permitting the binding of the detectably tagged predetermined compound with the second predetermined compound, and wherein the tag moiety of said detectably tagged predetermined compound bound to the second predetermined compound localizes within said at least one nanopore;

- (d) detecting said at least one detectable component of the tag moiety with said at least one nanopore so as to thereby detecting the interaction of the predetermined compounds.
37. The method of claim 35 or 36, further comprising a washing step to remove any detectably tagged predetermined compound which is not bound to the second predetermined compound prior to step (d).
38. The method of any one of claims 35-37, wherein at least one of the predetermined compounds is a protein.
39. The method of claim 38, wherein the protein is an antibody.
40. The method of any one of claims 35-37, wherein at least one of the predetermined compounds is non-proteinaceous.
41. The method of any one of claims 35-40, wherein said tag moiety comprises more than one detectable component.
42. The method of any one of claims 35-41, wherein each detectable component is independently detectable.
43. The method of any one of claims 35-42, wherein each of said at least one detectable component of said tag moiety is selected from the group consisting of ethylene glycol, an amino acid, a carbohydrate, a peptide, a dye, a fluorescent compound, a chemiluminescent compound, a mononucleotide, a dinucleotide, a trinucleotide, a tetranucleotide, a pentanucleotide, a

- 51 -

hexanucleotide, a polynucleotide, a nucleotide monophosphate, a nucleotide diphosphate, a nucleotide polyphosphate, an aliphatic acid, an aromatic acid, an unsubstituted alcohol or thiol, an alcohol or a thiol substituted with one or more halogens, a cyano group, a nitro group, an alkyl group, an alkenyl group, an alkynyl group, and an azido group.

44. The method of any one of claims 35-43, wherein said at least one detectable component of said tag moiety comprises a multiplicity of ethylene glycol units.
45. The method of any one of claims 35-44, wherein said multiplicity of ethylene glycol units comprises 16, 20, 24, or 36 ethylene glycol units.
46. The method of any one of claims 35-45, wherein said tag moiety attaches to the predetermined compound via a cleavable linker.
47. The method of claim 46, wherein the cleavable linker is a photocleavable linker.
48. The method of claim 46, wherein the cleavable linker is a chemically cleavable linker.
49. The method of any one of claims 35-48, wherein the predetermined compound present in the sample is a protein and said tag moiety attaches to the carboxy or amino terminus of said protein.
50. The method of any one of claims 35-49, wherein the predetermined compound present in the sample is a protein and said tag moiety attaches to a lysine, an arginine, or a cysteine residue of said protein.

- 52 -

51. The method of any one of claims 35-50, wherein said at least one nanopore is a biological nanopore, a modified biological nanopore, or a synthetic nanopore.
52. The method of claim 51, wherein said at least one nanopore is proteinaceous.
53. The method of any one of claims 35-52, wherein said at least one nanopore is an alpha hemolysin (α -hemolysin).
54. The method of any one of claims 35-50, wherein said at least one nanopore is a solid-state nanopore.
55. The method of claim 54, wherein the solid-state nanopore comprises graphene.
56. The method of any one of claims 35-55, wherein said at least one nanopore is in a membrane.
57. The method of any one of claims 35-56, wherein said at least one nanopore is part of an array of nanopores.
58. The method of claim 57, wherein each nanopore in said array comprises an identical immobilized second predetermined compound.
59. The method of claim 57, wherein each nanopore in said array comprises a different immobilized second predetermined compound.
60. The method of any one of claims 35-59, wherein the second predetermined compound is a protein.
61. The method of claim 60, wherein the protein is an antibody.

- 53 -

62. The method of any one of claim 35-59, wherein the second predetermined compound is non-proteinaceous.
63. The method of any one of claims 35-62, wherein said tag moiety produces a blockade signature detectable with said at least one nanopore.
64. The method of claim 63, wherein the blockade signature is result of a change in current amplitude of said at least one nanopore.
65. The method of claim 63, wherein the blockade signature is result of a change in conductance of said at least one nanopore.
66. The method of any one of claims 35-65, wherein said at least one nanopore further comprising a means for ejecting said tag moiety from the nanopore.
67. The method of any one of claims 35-66, wherein the sample comprises a plurality of predetermined compounds.
68. The method of claim 67, wherein in step (a) contacting the plurality of predetermined compounds present in the sample with a plurality of detectable tag moieties, under conditions permitting the tag moieties to attach to each of the predetermined compounds present in the sample to form detectably tagged predetermined compounds, wherein each tag moiety uniquely attaches to one of the predetermined compounds present in the sample and is distinguishable from any tag moiety attached to any other predetermined compound present in the sample.
69. The method of any one of claims 35-68, wherein said at least one nanopore comprises a plurality of predetermined compounds

- 54 -

capable of binding to the plurality of detectably tagged predetermined compounds present in the sample.

70. The method of any one of claims 1-68, wherein the means for ejecting said tag moiety from the nanopore consisting of a means to adjust electric field of said nanopore.
71. A detectable tag moiety for use in the method of any one of claims 1-70 which comprises a protein reactive group, at least one detectable component, and a linker connecting the protein reactive group and the at least one detectable component.
72. The tag moiety of claim 71, wherein the linker is a cleavable linker.
73. The tag moiety of claim 72, wherein the cleavable linker is a photocleavable linker.
74. The tag moiety of claim 72, wherein the cleavable linker is a chemically cleavable linker.
75. The tag moiety of any one of claims 71-74, wherein the at one detectable component is selected from the group consisting of ethylene glycol, an amino acid, a carbohydrate, a peptide, a dye, a fluorescent compound, a chemilluminiscent compound, a mononucleotide, a dinucleotide, a trinucleotide, a tetranucleotide, a pentanucleotide, a hexanucleotide, a polynucleotide, a nucleotide monophosphate, a nucleotide diphosphate, a nucleotide polyphosphate, an aliphatic acid, an aromatic acid, an unsubstituted alcohol or thiol, an alcohol or a thiol substituted with one or more halogens, a cyano group, a nitro group, an alkyl group, an alkenyl group, an alkynyl group, and an azido group.

- 55 -

76. A method for detecting the presence of a plurality of predetermined compounds in a plurality of samples, comprising:

- (a) contacting each sample with a nanopore-detectable tag moiety, under conditions permitting the tag moiety to attach to the plurality of predetermined compounds present in the sample to form detectably tagged predetermined compounds for that sample, which are distinguishable from any nanopore-detectable tag moiety attached to similar predetermined compounds from any other sample;
- (b) mixing the resulting tagged compounds from the plurality of samples from step (a) and treating them in such a way as to form an electrolyte solution containing the set of detectably and distinguishably tagged predetermined compounds;
- (c) contacting the electrolyte solution from step (b) with at least one nanopore, wherein said at least one nanopore comprises a means for specific analyte moiety binding to each of the detectably tagged predetermined compounds in the electrolyte solution, under conditions such that the detectably tagged predetermined compounds bind to said at least one nanopore;
- (d) treating the detectably tagged predetermined compounds bound to said at least one nanopore so as to separate the tag moieties from the predetermined compounds; and
- (e) detecting said at least one detectable component of each of the tag moieties so released as the tag moieties pass through said at least one nanopore so as to thereby detect the presence of the plurality of predetermined compounds in the plurality of samples.

FIGURE 1




FIGURE 2




FIGURE 3




FIGURE 4




FIGURE 5







FIGURE 6





FIGURE 7



INTERNATIONAL SEARCH REPORT

International application no.

PCT/US 14/29495

A. CLASSIFICATION OF SUBJECT MATTER

IPC(8) - C12Q 1/00 (2014.01)

USPC - 435/4; 435/6.11; 426/501

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC(8) - C12Q 1/00 (2014.01)

USPC - 435/4; 435/6.11; 426/501

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

IPC(8) - C12Q 1/00 (2014.01)

USPC - 435/4; 435/6.11; 426/501

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Minesoft Patbase, Google Scholar, keywords: nanopore and tags and dissociation and interaction, pore and PA = (THE TRUSTEES OF COLUMBIA UNIVERSITY IN THE CITY OF NEW YORK); (nanopore* or nanochannel*) and (dissociat* or detach*) w20 (tag or label); TAC = sequence w2 pool* and tag*

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 2013/016486 A1 (Bashir et al.) 31 January 2013 (31.01.2013) para [0051], [0052], [0090], [0091]; Figure 5, 6	1-5, 35-37, 76
Y	Wei et al. "Stochastic sensing of proteins with receptor-modified solid-state nanopores." Nature nanotechnology, 11 March 2012, Vol.7, no. 4, pp 257-263 (page 2 top portion; Figure 1a-h)	1-5, 35-37, 76
Y	US 2010/0227414 A1 (Ervin) 09 September 2010 (09.09.2010) para [0072], [0075]	5, 35-37
Y	US 2012 /0214162 A1 (Oliver) 23 August 2012 (23.08.2012) para [0024], [0030]	76

☐ Further documents are listed in the continuation of Box C.


* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

01 July 2014 (01.07.2014)

Date of mailing of the international search report

30 JUL 2014

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US, Commissioner for Patents
P.O. Box 1450, Alexandria, Virginia 22313-1450

Facsimile No. 571-273-3201

Authorized officer:

Lee W. Young

PCT Helpdesk: 571-272-4300

PCT OSP: 571-272-7774

INTERNATIONAL SEARCH REPORT

International application no.

PCT/US 14/29495

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☒ Claims Nos.: 6-34 and 38-75
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying additional fees, this Authority did not invite payment of additional fees.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- ☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- ☐ No protest accompanied the payment of additional search fees.