

(54) Title
Active substance combinations with insecticidal and acaricidal properties

(51)⁷ International Patent Classification(s)
A01N 043/12 A01N 047/02
A01N 037/28 A01N 047/34
A01N 043/16 A01N 047/38
A01N 043/24 A01N 063/00
A01N 043/28 A01N 063/04
A01N 043/68 A01N 065/00
A01N 043/90 A01N 043/08

(21) Application No: **2001258303** (22) Date of Filing: **2001.04.02**

(87) WIPO No: **WO01/78511**

(30) Priority Data

(31) Number (32) Date (33) Country
10018370 2000.04.14 DE

(43) Publication Date: **2001.10.30**

(43) Publication Journal Date: **2002.01.17**

(44) Accepted Journal Date: **2005.03.10**

(71) Applicant(s)
Bayer Aktiengesellschaft

(72) Inventor(s)
Fischer, Reiner; Bretschneider, Thomas; Erdelen, Christoph

(74) Agent / Attorney
Davies Collison Cave, Level 10 10 Barrack Street, Sydney, NSW, 2000

(56) Related Art
AU 12765/01
AU 52680/99

(12) NACH DEM VERTRAG ÜBER DIE INTERNATIONALE ZUSAMMENARBEIT AUF DEM GEBIET DES
PATENTWESENS (PCT) VERÖFFENTLICHTE INTERNATIONALE ANMELDUNG

(19) Weltorganisation für geistiges Eigentum
Internationales Büro



(43) Internationales Veröffentlichungsdatum
25. Oktober 2001 (25.10.2001)

PCT

(10) Internationale Veröffentlichungsnummer
WO 01/78511 A1

(51) Internationale Patentklassifikation⁷: A01N 43/12,
43/08 // (A01N 43/12, 65:00, 63:04, 63:00, 47:38, 47:34,
47:02, 43:90, 43:68, 43:28, 43:24, 43:16, 37:28)

(21) Internationales Aktenzeichen: PCT/EP01/03713

(22) Internationales Anmeldedatum:
2. April 2001 (02.04.2001)

(25) Einreichungssprache: Deutsch

(26) Veröffentlichungssprache: Deutsch

(30) Angaben zur Priorität:
100 18 370.0 14. April 2000 (14.04.2000) DE

(71) Anmelder (für alle Bestimmungsstaaten mit Ausnahme
von US): BAYER AKTIENGESELLSCHAFT [DE/DE];
51368 Leverkusen (DE).

(72) Erfinder; und

(75) Erfinder/Anmelder (nur für US): FISCHER, Reiner
[DE/DE]; Nelly-Sachs-Str. 23, 40789 Monheim (DE).
ERDELEN, Christoph [DE/DE]; Unterbüscherhof 15,
42799 Leichlingen (DE). BRETSCHNEIDER, Thomas
[DE/DE]; Talstr. 29 b, 53797 Lohmar (DE).

(74) Gemeinsamer Vertreter: BAYER AKTIENGE-
SELLSCHAFT; 51368 Leverkusen (DE).

(81) Bestimmungsstaaten (national): AE, AG, AL, AM, AT,
AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR,
CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK,
SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA,
ZW.

(84) Bestimmungsstaaten (regional): ARIPO-Patent (GH,
GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW),
curasisches Patent (AM, AZ, BY, KG, KZ, MD, RU, TJ,
TM), europäisches Patent (AT, BE, CH, CY, DE, DK,
ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR),
OAPI-Patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML,
MR, NE, SN, TD, TG).

Veröffentlicht:

- mit internationalem Recherchenbericht
- vor Ablauf der für Änderungen der Ansprüche geltenden
Frist; Veröffentlichung wird wiederholt, falls Änderungen
eintreffen

Zur Erklärung der Zweibuchstaben-Codes, und der anderen
Abkürzungen wird auf die Erklärungen ("Guidance Notes on
Codes and Abbreviations") am Anfang jeder regulären Ausgabe
der PCT-Gazette verwiesen.



WO 01/78511 A1

(54) Title: ACTIVE SUBSTANCE COMBINATIONS WITH INSECTICIDAL AND ACARICIDAL PROPERTIES

(54) Bezeichnung: WIRKSTOFFKOMBINATIONEN MIT INSEKTIZIDEN UND AKARIZIDEN EIGENSCHAFTEN

(57) Abstract: The novel active substance combinations made from given cyclic ketoenols and the active substances (1 to 22) cited in the description have very good insecticidal and acaricidal properties.

(57) Zusammenfassung: Die neuen Wirkstoffkombinationen aus bestimmten cyclischen Ketoenolen und den in der Beschreibung aufgeführten Wirkstoffe (1) bis (22) besitzen sehr gute insektizide und akarizide Eigenschaften.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization

International Bureau



(43) International publication date

25 October 2001 (25.10.2001)

PCT

(10) International publication number

WO 01/78511 A1

- (51) International patent classification⁷: A01N 43/12, 43/08 // (A01N 43/12, 65:00, 63:04, 63:00, 47:38, 47:34, 47:02, 43:90, 43:68, 43:28, 43:24, 43:16, 37:28)
- (21) International application number: PCT/EP01/03713
- (22) International filing date: 2 April 2001 (02.04.2001)
- (25) Language of filing: German
- (26) Language of publication: German
- (30) Data relating to the priority: 100 18 370.0 14 April 2000 (14.04.2000) DE
- (71) Applicant (for all designated States except US): BAYER AKTIENGESELLSCHAFT [DE/DE]; 51368 Leverkusen (DE).
- (72) Inventors; and
- (75) Inventors/Applicants (US only): FISCHER, Reiner [DE/DE]; Nelly-Sachs-Str. 23, 40789 Monheim (DE). ERDELEN, Christoph [DE/DE]; Unterbüscherhof 15, 42799 Leichlingen (DE). BRETSCHNEIDER, Thomas [DE/DE]; Talstr. 29 b, 53797 Lohmar (DE).
- (74) Joint Representative: BAYER AKTIENGESELLSCHAFT; 51368 Leverkusen (DE).
- (81) Designated states (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.
- (84) Designated states (regional): ARIPO Patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian Patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European Patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI Patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

- With the International Search Report.
- Before expiry of the period provided for amending the claims, will be republished if such amendments are received.

For an explanation of the two-letter codes and the other abbreviations, reference is made to the explanations ("Guidance Notes on Codes and Abbreviations") at the beginning of each regular edition of the PCT Gazette.

As printed

WO 01/78511 A1

(54) Title: ACTIVE SUBSTANCE COMBINATIONS WITH INSECTICIDAL AND ACARICIDAL PROPERTIES

(54) Bezeichnung: WIRKSTOFFKOMBINATIONEN MIT INSEKTIZIDEN UND AKARIZIDEN EIGENSCHAFTEN

(57) Abstract: The novel active substance combinations made from given cyclic ketoenols and the active substances (1 to 22) cited in the description have very good insecticidal and acaricidal properties.

(57) Zusammenfassung: Die neuen Wirkstoffkombinationen aus bestimmten cyclischen Ketoenolen und den in der Beschreibung aufgeführten Wirkstoffe (1) bis (22) besitzen sehr gute insektizide und akarizide Eigenschaften.

- 1 -


Active substance combinations with insecticidal and acaricidal properties

The present invention relates to novel active compound combinations comprising known cyclic ketoenols and further known insecticidally active compounds, which combinations are highly suitable for controlling animal pests such as insects and undesirable acarids.

It is already known that certain cyclic ketoenols have insecticidal and acaricidal properties (EP-A-528 156). WO 95/01971, EP-A-647 637, WO 96/16061, WO 96/20196, WO 96/25395, WO 96/35664, WO 97/02243, WO 97/01535, WO 97/36868, WO 97/43275, WO 98/05638, WO 98/06721, WO 99/16748, WO 99/43649, WO 99/48869 and WO 99/55673 describe further ketoenols having insecticidal and acaricidal properties. The activity of these substances is good; however, at low application rates it is sometimes unsatisfactory.

Furthermore, it is already known that numerous heterocycles, organotin compounds, benzoyl ureas and pyrethroids have insecticidal and acaricidal properties (cf. WO 93-22 297, WO 93-10 083, DE-A 2 641 343, EP-A-347 488, EP-A-210 487, US-A 3 264 177 and EP-A-234 045). However, the activity of these substances is not always satisfactory.

It has now been found that compounds of the formula (I)



in which

X represents C₁-C₆-alkyl, halogen, C₁-C₆-alkoxy or C₁-C₃-halogenoalkyl,

- Y represents hydrogen, C₁-C₆-alkyl, halogen, C₁-C₆-alkoxy or C₁-C₃-halogenoalkyl,
- 5 Z represents C₁-C₆-alkyl, halogen or C₁-C₆-alkoxy,
- n represents a number from 0 to 3,
- 10 A represents hydrogen or in each case optionally halogen-substituted straight-chain or branched C₁-C₁₂-alkyl, C₃-C₈-alkenyl, C₃-C₈-alkinyl, C₁-C₁₀-alkoxy-C₂-C₈-alkyl, C₁-C₈-polyalkoxy-C₂-C₈-alkyl, C₁-C₁₀-alkylthio-C₂-C₈-alkyl or cycloalkyl having 3-8 ring atoms, which may be interrupted by oxygen and/or sulphur or represents in each case optionally halogen-, C₁-C₆-alkyl-, C₁-C₆-halogenoalkyl-, C₁-C₆-alkoxy, C₁-C₆-halogenoalkoxy- or nitro-substituted phenyl or
- 15 phenyl-C₁-C₆-alkyl.
- B represents hydrogen, C₁-C₆-alkyl or C₁-C₆-alkoxy- C₂-C₄-alkyl


or in which

20

A and B together with the carbon atom to which they are attached form a saturated or unsaturated 3- to 8-membered ring which is optionally interrupted by oxygen and/or sulfur and optionally substituted by halogen, C₁-C₆-alkyl, C₁-C₆-alkoxy, C₁-C₄-halogenoalkyl, C₁-C₄-halogenoalkoxy, C₁-C₄-alkylthio or optionally substituted phenyl or is

25 optionally benzo-fused,

G represents hydrogen (a) or represents the groups



in which

5 R^1 represents in each case optionally halogen-substituted C_1 - C_{20} -alkyl, C_2 - C_{20} -alkenyl, C_1 - C_8 -alkoxy- C_2 - C_8 -alkyl, C_1 - C_8 -alkylthio- C_2 - C_8 -alkyl, C_1 - C_8 -polyalkoxy- C_2 - C_8 -alkyl or cycloalkyl having 3-8 ring atoms which may be interrupted by oxygen and/or sulfur atoms,

10 represents optionally halogen-, nitro-, C_1 - C_6 -alkyl-, C_1 - C_6 -alkoxy-, C_1 - C_6 -halogenoalkyl- or C_1 - C_6 -halogenoalkoxy-substituted phenyl,

15 represents optionally halogen-, C_1 - C_6 -alkyl-, C_1 - C_6 -alkoxy-, C_1 - C_6 -halogenoalkyl- or C_1 - C_6 -halogenoalkoxy-substituted phenyl- C_1 - C_6 -alkyl,

20 represents in each case optionally halogen- and/or C_1 - C_6 -alkyl-substituted pyridyl, pyrimidyl, thiazolyl or pyrazolyl,

25 represents optionally halogen- and/or C_1 - C_6 -alkyl-substituted phenoxy- C_1 - C_6 -alkyl,

R^2 represents in each case optionally halogen-substituted C_1 - C_{20} -alkyl, C_2 - C_{20} -alkenyl, C_1 - C_8 -alkoxy- C_2 - C_8 -alkyl or C_1 - C_8 -polyalkoxy- C_2 - C_8 -alkyl,

30 represents in each case optionally halogen-, nitro-, C_1 - C_6 -alkyl-, C_1 - C_6 -alkoxy- or C_1 - C_6 -halogenoalkyl-substituted phenyl or benzyl,

5 R³ represents optionally halogen-substituted C₁-C₈-alkyl, represents in each case optionally C₁-C₄-alkyl-, halogen-, C₁-C₄-halogenoalkyl-, C₁-C₄-alkoxy-, C₁-C₄-halogenoalkoxy-, nitro- or cyano-substituted phenyl or benzyl,


10 R⁴ and R⁵ independently of one another represent in each case optionally halogen-substituted C₁-C₈-alkyl, C₁-C₈-alkoxy, C₁-C₈-alkylamino, di-(C₁-C₈)-alkylamino, C₁-C₈-alkylthio, C₂-C₅-alkenylthio, C₂-C₅-alkinylthio or C₃-C₇-cycloalkylthio, represent in each case optionally halogen-, nitro-, cyano-, C₁-C₄-alkoxy-, C₁-C₄-halogenoalkoxy-, C₁-C₄-alkylthio-, C₁-C₄-halogenoalkylthio-, C₁-C₄-alkyl- or C₁-C₄-halogenoalkyl-substituted phenyl, phenoxy or phenylthio,

15 R⁶ and R⁷ independently of one another represent in each case optionally halogen-substituted C₁-C₁₀-alkyl, C₁-C₁₀-alkoxy, C₃-C₈-alkenyl or C₁-C₈-alkoxy-C₁-C₈-alkyl, represent optionally halogen-, C₁-C₆-halogenoalkyl-, C₁-C₆-alkyl- or C₁-C₆-alkoxy-substituted phenyl, represent optionally halogen-, C₁-C₆-alkyl-, C₁-C₆-halogenoalkyl- or
20 C₁-C₆-alkoxy-substituted benzyl or together represent a 5- or 6-membered ring which may optionally be interrupted by oxygen or sulfur and which may optionally be substituted by C₁-C₆-alkyl,

and

A) benzoylureas, preferably

1. chlorfluazuron




5 known from DE-A-2 818 830

and/or

2. diflubenzuron

10




known from DE-A 2 123 236

and/or

15


3. lufenuron



known from EP-A-179 022

and/or


4. teflubenzuron



5 known from EP-A-052 833

and/or

5. triflumuron




10

known from DE-A-2 601 780

and/or

15 6. novaluron




known from US 4,980,376

20

and/or

7. flufenoxuron




5 known from EP-A 161 019

and/or

8. hexaflumuron

10



known from EP-A 71 279

and/or

15

B) macrolides, preferably

9. emamectin


20

known from EP-A-089 202

and/or

C) diacylhydrazines, preferably


10. methoxyfenozide



known from EP-A-639 559

and/or


10 11. tebufenozide



known from EP-A-339 854

15 and/or

12. halofenozide




known from EP-A 228 564

5

and/or

- D) halogenocycloalkanes, preferably

- 10 13. endosulfan



known from DE-A-1 015 797

15 and/or

14. *Trichogramma* spp.

known from The Pesticide Manual, 11th Edition, 1997, p. 1236

20 and/or


15. *Verticillium lecanii*

known from The Pesticide Manual, 11th Edition, 1997, p. 1266

and/or

16. fipronil

5




known from EP-A-295 117

and/or

10

17. cyromazine



known from DE-A-2 736 876

15

and/or


18. azadirachtin

known from The Pesticide Manual, 11th Edition, 1997, p. 59

20

and/or

19. diofenolan




known from DE-A 2 655 910

5

and/or

20. indoxacarb




10

known from WO 92/11249

and/or


15 21. chromafenozide (ANS-118)



known from EP-A-496 342

and/or

5 22. bistrifluoron (DBI-3204)



known from WO 98/00 394

10 have very good insecticidal and acaricidal properties.

Surprisingly, the insecticidal and acaricidal activity of the active compound combination according to the invention is considerably higher than the sum of the activities of the individual active compounds. An unforeseeable true synergistic effect is present, not just an addition of activities.

15

In addition to at least one active compound of the formula (I), the active compound combinations according to the invention comprise at least one active compound from the group of compounds 1 to 22.

20

Preference is given to active compound combinations comprising compounds of the formula (I)

in which

25

X represents C₁-C₄-alkyl, halogen, C₁-C₄-alkoxy or C₁-C₂-halogenoalkyl,

Y represents hydrogen, C₁-C₄-alkyl, halogen, C₁-C₄-alkoxy or C₁-C₂-halogenoalkyl,

5 Z represents C₁-C₄-alkyl, halogen or C₁-C₄-alkoxy,

n represents 0 or 1,

10 A and B together with the carbon atom to which they are attached form a saturated 5- or 6-membered ring which is optionally substituted by C₁-C₄-alkyl or C₁-C₄-alkoxy,

G represents hydrogen (a) or represents the groups


15 in which

20 R¹ represents in each case optionally halogen-substituted C₁-C₁₆-alkyl, C₂-C₁₆-alkenyl, C₁-C₆-alkoxy-C₂-C₆-alkyl or cycloalkyl having 3-7 ring atoms which may be interrupted by 1 or 2 oxygen and/or sulfur atoms, represents optionally halogen-, nitro-, C₁-C₄-alkyl-, C₁-C₄-alkoxy-, C₁-C₃-halogenoalkyl-, C₁-C₃-halogenoalkoxy-substituted phenyl;

25 R² represents in each case optionally halogen-substituted C₁-C₁₆-alkyl, C₂-C₁₆-alkenyl or C₁-C₆-alkoxy-C₂-C₆-alkyl,

30 represents in each case optionally halogen-, nitro-, C₁-C₄-alkyl-, C₁-C₄-alkoxy- or C₁-C₄-halogenoalkyl-substituted phenyl or benzyl, and at least one active compound from the group of compounds 1 to 22.

Particular preference is given to active compound combinations comprising the dihydrofuranone derivative of the formula (I-b-1)



5 and at least one active compound from the group of compounds 1 to 22.

In addition, the active compound combinations may also comprise further fungicidally, acaricidally or insecticidally active components which may be admixed.

10 If the active compounds are present in the active compound combinations according to the invention in certain weight ratios, the synergistic effect is particularly pronounced. However, the weight ratios of the active compounds in the active compound combinations may be varied within a relatively wide range. In general, the combinations according to the invention comprise active compounds of the formula

15 (I) and the co-components in the preferred and particularly preferred mixing ratios indicated in the table below.

* the mixing ratios are based on weight ratios. The ratio is to be understood as meaning active compound of the formula (I): co-component

20

Co-component	Preferred mixing ratio	Particularly preferred mixing ratio
Chlorfluazuron	10:1 to 1:10	5:1 to 1:5
Diflubenzuron	10:1 to 1:10	5:1 to 1:5
Lufenuron	20:1 to 1:5	10:1 to 1:2

Co-component	Preferred mixing ratio	Particularly preferred mixing ratio
Teflubenzuron	20:1 to 1:5	10:1 to 1:2
Triflumuron	10:1 to 1:10	5:1 to 1:5
Novaluron	10:1 to 1:10	5:1 to 1:5
Flufenoxuron	10:1 to 1:10	5:1 to 1:5
Hexaflumuron	20:1 to 1:5	5:1 to 1:2
Emamectin	50:1 to 1:5	10:1 to 1:1
Methoxyfenozide	10:1 to 1:10	5:1 to 1:5
Tebufenozide	10:1 to 1:10	5:1 to 1:5
Halofenozide	2:1 to 1:100	1:1 to 1:30
Endosulfan	10:1 to 1:10	5:1 to 1:5
Fipronil	10:1 to 1:10	5:1 to 1:5
Cyromazine	10:1 to 1:10	5:1 to 1:5
Azadirachtin	50:1 to 1:5	10:1 to 1:1
Diofenolan	100:1 to 1:2	20:1 to 1:1
Indoxacarb	50:1 to 1:5	20:1 to 1:2
Trichogramma spp.	1 000 g a.i./ha:20 000 wasps/ha to 10 g a.i./ha : 500 000 wasps/ha	300 g a.i./ha:50 000 wasps/ha to 50 g a.i./ha : 300 000 wasps/ha
Verticillium lecanii	0.05 % a.i.:0.05 % F ^(*) to 0.001 % a.i.:0.5 % F ^(*)	0.03 % a.i.:0.1 % F ^(*) to 0.005 % a.i.:0.2 % F ^(*)
chromafenozide	10:1 to 1:10	5:1 to 1:5
bistrifluoron	10:1 to 1:10	5:1 to 1:5

F^(*) Formulation comprising 10⁹ to 10¹⁰ spores/g

The active compound combinations according to the invention are suitable for controlling animal pests, preferably arthropods and nematodes, in particular insects and arachnids found in agriculture, in forests, in the protection of stored products and materials and in the hygiene sector. They are active against normally sensitive and resistant species, and against all or individual developmental stages. The abovementioned pests include:

From the order of the Isopoda, for example, *Oniscus asellus*, *Armadillidium vulgare*,
10 *Porcellio scaber*.

From the order of the Diplopoda, for example, *Blaniulus guttulatus*.

From the order of the Chilopoda, for example, *Geophilus carpophagus*, *Scutigera*
spp.

From the order of the Symphyla, for example, *Scutigereilla immaculata*.

15 From the order of the Thysanura, for example, *Lepisma saccharina*.

From the order of the Collembola, for example, *Onychiurus armatus*.

From the order of the Orthoptera, for example, *Acheta domesticus*, *Gryllotalpa* spp.,
Locusta migratoria migratorioides, *Melanoplus* spp., *Schistocerca gregaria*.

20 From the order of the Blattaria, for example, *Blatta orientalis*, *Periplaneta americana*,
Leucophaea maderae, *Blattella germanica*.

From the order of the Dermaptera, for example, *Forficula auricularia*.

From the order of the Isoptera, for example, *Reticulitermes* spp.

25 From the order of the Phthiraptera, for example, *Pediculus humanus corporis*,
Haematopinus spp., *Linognathus* spp., *Trichodectes* spp., *Damalinea* spp.

From the order of the Thysanoptera, for example, *Hercinothrips femoralis*, *Thrips*
tabaci, *Thrips palmi*, *Frankliniella occidentalis*.

From the order of the Heteroptera, for example, *Eurygaster* spp., *Dysdercus*
intermedius, *Piesma quadrata*, *Cimex lectularius*, *Rhodnius prolixus*, *Triatoma* spp.

30 From the order of the Homoptera, for example, *Aleurodes brassicae*, *Bemisia tabaci*,
Trialeurodes vaporariorum, *Aphis gossypii*, *Brevicoryne brassicae*, *Cryptomyzus*

ribis, *Aphis fabae*, *Aphis pomi*, *Eriosoma lanigerum*, *Hyalopterus arundinis*,
Phylloxera vastatrix, *Pemphigus* spp., *Macrosiphum avenae*, *Myzus* spp., *Phorodon*
humuli, *Rhopalosiphum padi*, *Empoasca* spp., *Euscelis bilobatus*, *Nephotettix*
cincticeps, *Lecanium corni*, *Saissetia oleae*, *Laodelphax striatellus*, *Nilaparvata*
5 *lugens*, *Aonidiella aurantii*, *Aspidiotus hederae*, *Pseudococcus* spp., *Psylla* spp.

From the order of the Lepidoptera, for example, *Pectinophora gossypiella*, *Bupalus*
piniarius, *Cheimatobia brumata*, *Lithocolletis blancardella*, *Hyponomeuta padella*,
Plutella xylostella, *Malacosoma neustria*, *Euproctis chrysorrhoea*, *Lymantria* spp.,
Bucculatrix thurberiella, *Phyllocnistis citrella*, *Agrotis* spp., *Euxoa* spp., *Feltia* spp.,
10 *Earias insulana*, *Heliothis* spp., *Mamestra brassicae*, *Panolis flammea*, *Spodoptera*
spp., *Trichoplusia ni*, *Carpocapsa pomonella*, *Pieris* spp., *Chilo* spp., *Pyrausta*
nubilalis, *Ephestia kuehniella*, *Galleria mellonella*, *Tineola bisselliella*, *Tinea*
pellionella, *Hofmannophila pseudospretella*, *Cacoecia podana*, *Capua reticulana*,
Choristoneura fumiferana, *Clysia ambiguella*, *Homona magnanima*, *Tortrix viridana*,
15 *Cnaphalocerus* spp., *Oulema oryzae*.

From the order of the Coleoptera, for example, *Anobium punctatum*, *Rhizophthera*
dominica, *Bruchidius obtectus*, *Acanthoscelides obtectus*, *Hylotrupes bajulus*,
Agelastica alni, *Leptinotarsa decemlineata*, *Phaedon cochleariae*, *Diabrotica* spp.,
Psylliodes chrysocephala, *Epilachna varivestis*, *Atomaria* spp., *Oryzaephilus*
20 *surinamensis*, *Anthonomus* spp., *Sitophilus* spp., *Otiorrhynchus sulcatus*,
Cosmopolites sordidus, *Ceuthorrhynchus assimilis*, *Hypera postica*, *Dermestes* spp.,
Trogoderma spp., *Anthrenus* spp., *Attagenus* spp., *Lyctus* spp., *Meligethes aeneus*,
Ptinus spp., *Niptus hololeucus*, *Gibbium psylloides*, *Tribolium* spp., *Tenebrio*
molitor, *Agriotes* spp., *Conoderus* spp., *Melolontha melolontha*, *Amphimallon solsti-*
25 *tialis*, *Costelytra zealandica*, *Lissorhoptrus oryzophilus*.

From the order of the Hymenoptera, for example, *Diprion* spp., *Hoplocampa* spp.,
Lasius spp., *Monomorium pharaonis*, *Vespa* spp.

From the order of the Diptera, for example, *Aedes* spp., *Anopheles* spp., *Culex* spp.,
Drosophila melanogaster, *Musca* spp., *Fannia* spp., *Calliphora erythrocephala*,
30 *Lucilia* spp., *Chrysomyia* spp., *Cuterebra* spp., *Gastrophilus* spp., *Hyppobosca* spp.,
Stomoxys spp., *Oestrus* spp., *Hypoderma* spp., *Tabanus* spp., *Tannia* spp., *Bibio*

hortulanus, *Oscinella frit*, *Phorbia* spp., *Pegomyia hyoscyami*, *Ceratitis capitata*, *Dacus oleae*, *Tipula paludosa*, *Hylemyia* spp., *Liriomyza* spp.

From the order of the Siphonaptera, for example, *Xenopsylla cheopis*, *Ceratophyllus* spp.

- 5 From the class of the Arachnida, for example, *Scorpio maurus*, *Latrodectus mactans*, *Acarus siro*, *Argas* spp., *Ornithodoros* spp., *Dermanyssus gallinae*, *Eriophyes ribis*, *Phyllocoptura oleivora*, *Boophilus* spp., *Rhipicephalus* spp., *Amblyomma* spp., *Hyalomma* spp., *Ixodes* spp., *Psoroptes* spp., *Chorioptes* spp., *Sarcoptes* spp., *Tarsonemus* spp., *Bryobia praetiosa*, *Panonychus* spp., *Tetranychus* spp.,
10 *Hemitarsonemus* spp., *Brevipalpus* spp.

The plant-parasitic nematodes include, for example, *Pratylenchus* spp., *Radopholus similis*, *Ditylenchus dipsaci*, *Tylenchulus semipenetrans*, *Heterodera* spp., *Globodera* spp., *Meloidogyne* spp., *Aphelenchoides* spp., *Longidorus* spp., *Xiphinema* spp.,
15 *Trichodorus* spp., *Bursaphelenchus* spp.

The active compound combinations can be converted into the customary formulations such as solutions, emulsions, wettable powders, suspensions, powders, dusts, pastes, soluble powders, granules, suspension-emulsion concentrates, natural and synthetic materials impregnated with active compound, and microencapsulations in polymeric materials.
20

These formulations are produced in a known manner, for example by mixing the active compounds with extenders, that is, liquid solvents and/or solid carriers, optionally with the use of surfactants, that is, emulsifiers and/or dispersants, and/or foam formers.
25

If the extender used is water, it is also possible, for example, to use organic solvents as cosolvents. The following are essentially suitable as liquid solvents: aromatics such as xylene, toluene or alkyl-naphthalenes, chlorinated aromatics and chlorinated aliphatic hydrocarbons such as chlorobenzenes, chloroethylenes or methylene
30

chloride, aliphatic hydrocarbons such as cyclohexane or paraffins, for example mineral oil fractions, mineral and vegetable oils, alcohols such as butanol or glycol and their ethers and esters, ketones such as acetone, methyl ethyl ketone, methyl isobutyl ketone or cyclohexanone, strongly polar solvents such as dimethylformamide and dimethyl sulphoxide, or else water.

Suitable solid carriers are:

for example ammonium salts and ground natural minerals such as kaolins, clays, talc, chalk, quartz, attapulgite, montmorillonite or diatomaceous earth, and ground synthetic materials such as highly disperse silica, alumina and silicates; suitable solid carriers for granules are: for example crushed and fractionated natural rocks such as calcite, marble, pumice, sepiolite and dolomite, or else synthetic granules of inorganic and organic meals, and granules of organic material such as sawdust, coconut shells, maize cobs and tobacco stalks; suitable emulsifiers and/or foam formers are: for example nonionic and anionic emulsifiers such as polyoxyethylene fatty acid esters, polyoxyethylene fatty alcohol ethers, for example alkylaryl polyglycol ethers, alkylsulphonates, alkyl sulphates, arylsulphonates, or else protein hydrolysates; suitable dispersants are: for example lignin-sulphite waste liquors and methylcellulose.

Tackifiers such as carboxymethylcellulose and natural and synthetic polymers in the form of powders, granules or latices, such as gum arabic, polyvinyl alcohol and polyvinyl acetate, or else natural phospholipids such as cephalins and lecithins and synthetic phospholipids can be used in the formulations. Other additives can be mineral and vegetable oils.

It is possible to use colorants such as inorganic pigments, for example iron oxide, titanium oxide and Prussian Blue, and organic colorants such as alizarin colorants, azo colorants and metal phthalocyanine colorants, and trace nutrients such as salts of iron, manganese, boron, copper, cobalt, molybdenum and zinc.

The formulations generally comprise between 0.1 and 95% by weight of active compound, preferably between 0.5 and 90%.

5 The active compound combinations according to the invention can be present in their commercially available formulations and in the use forms, prepared from these formulations, as a mixture with other active compounds, such as insecticides, attractants, sterilants, bactericides, acaricides, nematocides, fungicides, growth-regulating substances or herbicides. The insecticides include, for example, phosphates, carbamates, carboxylates, chlorinated hydrocarbons, phenylureas and
10 substances produced by microorganisms.

Mixtures with other known active compounds such as herbicides or with fertilizers and growth regulators are also possible.

15 When used as insecticides, the active compound combinations according to the invention can furthermore be present in their commercially available formulations and in the use forms, prepared from these formulations, as a mixture with synergists. Synergists are compounds which increase the action of the active compounds, without it being necessary for the synergist added to be active itself.
20

The active compound content of the use forms prepared from the commercially available formulations can vary within wide limits. The active compound concentration of the use forms can be from 0.0000001 to 95% by weight of active compound, preferably between 0.0001 and 1% by weight.
25

The compounds are employed in a customary manner appropriate for the use forms.

30 When used against hygiene pests and stored-product pests, the active compound combinations are distinguished by an excellent residual action on wood and clay as well as good stability to alkali on limed substrates.

The active compound combinations according to the invention are not only active against plant pests, hygiene pests and stored-product pests, but also, in the veterinary medicine sector, against animal parasites (ectoparasites) such as hard ticks, soft ticks, mange mites, harvest mites, flies (stinging and licking), parasitizing fly larvae, lice, hair lice, bird lice and fleas. These parasites include:

5

From the order of the Anoplurida, for example, *Haematopinus* spp., *Linognathus* spp., *Pediculus* spp., *Phtirus* spp., *Solenopotes* spp.

10

From the order of the Mallophagida and the suborders Amblycerina and Ischnocerina, for example, *Trimenopon* spp., *Menopon* spp., *Trinoton* spp., *Bovicola* spp., *Werneckiella* spp., *Lepikentron* spp., *Damalina* spp., *Trichodectes* spp., *Felicola* spp.

15

From the order Diptera and the suborders Nematocerina and Brachycerina, for example, *Aedes* spp., *Anopheles* spp., *Culex* spp., *Simulium* spp., *Eusimulium* spp., *Phlebotomus* spp., *Lutzomyia* spp., *Culicoides* spp., *Chrysops* spp., *Hybomitra* spp., *Atylotus* spp., *Tabanus* spp., *Haematopota* spp., *Philipomyia* spp., *Braula* spp., *Musca* spp., *Hydrotaea* spp., *Stomoxys* spp., *Haematobia* spp., *Morellia* spp., *Fannia* spp., *Glossina* spp., *Calliphora* spp., *Lucilia* spp., *Chrysomyia* spp., *Wohlfahrtia* spp., *Sarcophaga* spp., *Oestrus* spp., *Hypoderma* spp., *Gasterophilus* spp., *Hippobosca* spp., *Lipoptena* spp., *Melophagus* spp.

20

From the order of the Siphonaptera, for example, *Pulex* spp., *Ctenocephalides* spp., *Xenopsylla* spp., *Ceratophyllus* spp.

From the order of the Heteroptera, for example, *Cimex* spp., *Triatoma* spp., *Rhodnius* spp., *Panstrongylus* spp.

25

From the order of the Blattaria, for example, *Blatta orientalis*, *Periplaneta americana*, *Blattella germanica*, *Supella* spp.

30

From the subclass of the Acaria (Acarida) and the orders of the Meta- and Mesostigmata, for example, *Argas* spp., *Ornithodoros* spp., *Otobius* spp., *Ixodes* spp., *Amblyomma* spp., *Boophilus* spp., *Dermacentor* spp., *Haemophysalis* spp., *Hyalomma* spp., *Rhipicephalus* spp., *Dermanyssus* spp., *Raillietia* spp., *Pneumonyssus* spp., *Sternostoma* spp., *Varroa* spp.

From the order of the Actinedida (Prostigmata) and Acaridida (Astigmata), for example, Acarapis spp., Cheyletiella spp., Ornithocheyletia spp., Myobia spp., Psorergates spp., Demodex spp., Trombicula spp., Listrophorus spp., Acarus spp., Tyrophagus spp., Caloglyphus spp., Hypodectes spp., Pterolichus spp., Psoroptes spp., Chorioptes spp., Otodectes spp., Sarcoptes spp., Notoedres spp., Knemidocoptes spp., Cytodites spp., Laminosioptes spp.

The active compound combinations according to the invention are also suitable for controlling arthropods which attack agricultural livestock such as, for example, cattle, sheep, goats, horses, pigs, donkeys, camels, buffaloes, rabbits, chickens, turkeys, ducks, geese, honey-bees, other domestic animals such as, for example, dogs, cats, caged birds, aquarium fish and so-called experimental animals such as, for example, hamsters, guinea pigs, rats and mice. By controlling these arthropods, cases of death and reductions in productivity (for meat, milk, wool, hides, eggs, honey and the like) should be diminished, so that more economical and simpler animal husbandry is possible by the use of the active compound combinations according to the invention.

The active compound combinations according to the invention are used in the veterinary sector in a known manner by enteral administration in the form of, for example, tablets, capsules, potions, drenches, granules, pastes, boluses, the feed-through method, suppositories, by parenteral administration such as, for example, by injections (intramuscularly, subcutaneously, intravenously, intraperitoneally and the like), implants, by nasal administration, by dermal administration in the form of, for example, immersing or dipping, spraying, pouring-on, spotting-on, washing, dusting, and with the aid of active-compound-comprising moulded articles such as collars, ear tags, tail tags, limb bands, halters, marking devices and the like.

When used for cattle, poultry, domestic animals and the like, the active compound combinations can be applied as formulations (for example powders, emulsions, flowables) comprising the active compounds in an amount of 1 to 80% by weight,

either directly or after 100- to 10,000-fold dilution, or they may be used as a chemical dip.

5 Moreover, it has been found that the active compound combinations according to the invention show a potent insecticidal action against insects which destroy industrial materials.

The following insects may be mentioned by way of example and with preference, but not by way of limitation:

10 Beetles such as

Hylotrupes bajulus, Chlorophorus pilosis, Anobium punctatum, Xestobium rufovillosum, Ptilinus pecticornis, Dendrobium pertinex, Ernobius mollis, Priobium carpini, Lyctus brunneus, Lyctus africanus, Lyctus planicollis, Lyctus linearis, Lyctus pubescens, Trogoxylon aequale, Minthes rugicollis, Xyleborus spec., Tryptodendron spec., Apate monachus, Bostrychus capucins, Heterobostrychus brunneus, Sinoxylon spec., Dinoderus minutus.

Dermapterans such as

Sirex juvenus, Urocerus gigas, Urocerus gigas taignus, Urocerus augur.

Termites such as

20 Kaloterme flavicollis, Cryptoterme brevis, Heteroterme indicola, Reticuliterme flavipes, Reticuliterme santonensis, Reticuliterme lucifugus, Mastoterme darwiniensis, Zootermopsis nevadensis, Coptoterme formosanus.

Bristle-tails such as *Lepisma saccharina*.

25 Industrial materials in the present context are understood as meaning non-living materials such as, preferably, polymers, adhesives, glues, paper and board, leather, wood, timber products and paints.

30 The material which is to be protected from insect attack is very especially preferably wood and timber products.

Wood and timber products which can be protected by the composition according to the invention, or mixtures comprising it, are to be understood as meaning, for example:

5 Construction timber, wooden beams, railway sleepers, bridge components, jetties, vehicles made of wood, boxes, pallets, containers, telephone poles, wood lagging, windows and doors made of wood, plywood, chipboard, joinery, or timber products which quite generally are used in house construction or building joinery.

10 The active compound combinations can be used as such, in the form of concentrates or generally customary formulations such as powders, granules, solutions, suspensions, emulsions or pastes.

15 The abovementioned formulations can be prepared in a manner known per se, for example by mixing the active compounds with at least one solvent or diluent, emulsifier, dispersant and/or binder or fixative, water repellent, if desired desiccants and UV stabilizers, and if desired colorants and pigments and other processing auxiliaries.

20 The insecticidal compositions or concentrates used for protecting wood and timber products comprise the active compound according to the invention in a concentration of 0.0001 to 95% by weight, in particular 0.001 to 60% by weight.

25 The amount of composition or concentrate employed depends on the species and the abundance of the insects and on the medium. The optimal quantity to be employed can be determined in each case by test series upon application. In general, however, it will suffice to employ 0.0001 to 20% by weight, preferably 0.001 to 10% by weight, of the active compound, based on the material to be protected.

30 A suitable solvent and/or diluent is an organochemical solvent or solvent mixture and/or an oily or oil-type organochemical solvent or solvent mixture of low volatility

and/or a polar organochemical solvent or solvent mixture and/or water and, if appropriate, an emulsifier and/or wetter.

5 Organochemical solvents which are preferably employed are oily or oil-type solvents with an evaporation number of above 35 and a flash point of above 30°C, preferably above 45°C. Such oily and oil-type solvents which are insoluble in water and of low volatility and which are used are suitable mineral oils or their aromatic fractions or mineral-oil-containing solvent mixtures, preferably white spirit, petroleum and/or alkylbenzene.

10

Mineral oils with a boiling range of 170 to 220°C, white spirit with a boiling range of 170 to 220°C, spindle oil with a boiling range of 250 to 350°C, petroleum and aromatics with a boiling range of 160 to 280°C, oil of turpentine, and the like are advantageously used.

15

In a preferred embodiment, liquid aliphatic hydrocarbons with a boiling range of 180 to 210°C or high-boiling mixtures of aromatic and aliphatic hydrocarbons with a boiling range of 180 to 220°C and/or spindle oil and/or monochloronaphthalene, preferably α -monochloronaphthalene are used.

20

The organic oily or oil-type solvents of low volatility and with an evaporation number of above 35 and a flash point of above 30°C, preferably above 45°C, can be replaced in part by organochemical solvents of high or medium volatility, with the proviso that the solvent mixture also has an evaporation number of above 35 and a flash point of above 30°C, preferably above 45°C, and that the mixture is soluble or emulsifiable in this solvent mixture.

25

In a preferred embodiment, some of the organochemical solvent or solvent mixture or an aliphatic polar organochemical solvent or solvent mixture is replaced. Aliphatic organochemical solvents which contain hydroxyl and/or ester and/or ether groups are preferably used, such as, for example, glycol ethers, esters or the like.

30

Organochemical binders used for the purposes of the present invention are the synthetic resins and/or binding drying oils which are known per se and which can be diluted in water and/or dissolved or dispersed or emulsified in the organochemical solvents employed, in particular binders composed of, or comprising, an acrylate resin, a vinyl resin, for example polyvinyl acetate, polyester resin, polycondensation or polyaddition resin, polyurethane resin, alkyd resin or modified alkyd resin, phenol resin, hydrocarbon resin such as indene/coumarone resin, silicone resin, drying vegetable and/or drying oils and/or physically drying binders based on a natural and/or synthetic resin.

The synthetic resin employed as binder can be employed in the form of an emulsion, dispersion or solution. Bitumen or bituminous substances may also be used as binders, in amounts of up to 10% by weight. In addition, colorants, pigments, water repellants, odour-masking agents, and inhibitors or anticorrosive agents and the like, all of which are known per se, can be employed.

In accordance with the invention, the composition or the concentrate preferably comprises, as organochemical binders, at least one alkyd resin or modified alkyd resin and/or a drying vegetable oil. Alkyd resins which are preferably used in accordance with the invention are those with an oil content of over 45% by weight, preferably 50 to 68% by weight.

Some or all of the abovementioned binder can be replaced by a fixative (mixture) or plasticizer (mixture). These additives are intended to prevent volatilization of the active compounds, and also crystallization or precipitation. They preferably replace 0.01 to 30% of the binder (based on 100% of binder employed).

The plasticizers are from the chemical classes of the phthalic esters, such as dibutyl phthalate, dioctyl phthalate or benzyl butyl phthalate, phosphoric esters such as tributyl phosphate, adipic esters such as di-(2-ethylhexyl) adipate, stearates such as

butyl stearate or amyl stearate, oleates such as butyl oleate, glycerol ethers or higher-molecular-weight glycol ethers, glycerol esters and p-toluenesulphonic esters.

5 Fixatives are based chemically on polyvinyl alkyl ethers such as, for example, polyvinyl methyl ether, or ketones such as benzophenone and ethylenebenzophenone.

Other suitable solvents or diluents are, in particular, water, if appropriate as a mixture with one or more of the abovementioned organochemical solvents or diluents, emulsifiers and dispersants.

10

Particularly effective timber protection is achieved by industrial-scale impregnating processes, for example the vacuum, double-vacuum or pressure processes.

15

The active compound combinations according to the invention can at the same time be employed for protecting objects which come into contact with saltwater or brackish water, such as hulls, screens, nets, buildings, moorings and signalling systems, against fouling.

20

Fouling by sessile Oligochaeta, such as Serpulidae, and by shells and species from the Ledamorpha group (goose barnacles), such as various Lepas and Scalpellum species, or by species from the Balanomorpha group (acorn barnacles), such as Balanus or Pollicipes species, increases the frictional drag of ships and, as a consequence, leads to a marked increase in operation costs owing to higher energy consumption and additionally frequent residence in the dry dock.

25

Apart from fouling by algae, for example Ectocarpus sp. and Ceramium sp., fouling by sessile Entomostraka groups, which come under the generic term Cirripedia (cirriped crustaceans), is of particular importance.

30

Surprisingly, it has now been found that the active compound combinations according to the invention have an outstanding antifouling action.

Using the active compound combinations according to the invention, allows the use of heavy metals such as, for example, in bis(trialkyltin) sulphides, tri-*n*-butyltin laurate, tri-*n*-butyltin chloride, copper(I) oxide, triethyltin chloride, tri-*n*-butyl(2-phenyl-4-chlorophenoxy)tin, tributyltin oxide, molybdenum disulphide, antimony oxide, polymeric butyl titanate, phenyl-(bispyridine)-bismuth chloride, tri-*n*-butyltin fluoride, manganese ethylenebisthiocarbamate, zinc dimethyldithiocarbamate, zinc ethylenebisthiocarbamate, zinc salts and copper salts of 2-pyridinethiol 1-oxide, bisdimethyldithiocarbamoylzinc ethylenebisthiocarbamate, zinc oxide, copper(I) ethylene-bisdithiocarbamate, copper thiocyanate, copper naphthenate and tributyltin halides to be dispensed with, or the concentration of these compounds to be substantially reduced.

If appropriate, the ready-to-use antifouling paints can additionally comprise other active compounds, preferably algicides, fungicides, herbicides, molluscicides, or other antifouling active compounds.

Preferably suitable components in combinations with the antifouling compositions according to the invention are:

algicides such as

2-*tert*-butylamino-4-cyclopropylamino-6-methylthio-1,3,5-triazine, dichlorophen, diuron, endothal, fentin acetate, isoproturon, methabenzthiazuron, oxyfluorfen, quinoclamine and terbutryn;

fungicides such as

benzo[*b*]thiophenecarboxylic acid cyclohexylamide S,S-dioxide, dichlofluanid, fluorfolpet, 3-iodo-2-propinyl butylcarbamate, tolylfluanid and azoles such as azaconazole, cyproconazole, epoxyconazole, hexaconazole, metconazole, propiconazole and tebuconazole;

molluscicides such as

fentin acetate, metaldehyde, methiocarb, niclosamid, thiodicarb and trimethacarb;

or conventional antifouling active compounds such as

5 4,5-dichloro-2-octyl-4-isothiazolin-3-one, diiodomethylparatryl sulphone, 2-(N,N-dimethylthiocarbamoylthio)-5-nitrothiazyl, potassium, copper, sodium and zinc salts of 2-pyridinethiol 1-oxide, pyridine-triphenylborane, tetrabutylstannoxane, 2,3,5,6-tetrachloro-4-(methylsulphonyl)-pyridine, 2,4,5,6-tetrachloroisophthalonitrile, tetramethylthiuram disulphide and 2,4,6-trichlorophenylmaleimide.

10 The antifouling compositions used comprise the active compound combinations according to the invention in a concentration of 0.001 to 50% by weight, in particular 0.01 to 20% by weight.

15 Moreover, the antifouling compositions according to the invention comprise the customary components such as, for example, those described in Ungerer, *Chem. Ind.* 1985, 37, 730-732 and Williams, *Antifouling Marine Coatings*, Noyes, Park Ridge, 1973.

20 Besides the algicidal, fungicidal, molluscicidal active compounds and insecticidal active compounds according to the invention, antifouling paints comprise, in particular, binders.

25 Examples of recognized binders are polyvinyl chloride in a solvent system, chlorinated rubber in a solvent system, acrylic resins in a solvent system, in particular in an aqueous system, vinyl chloride/vinyl acetate copolymer systems in the form of aqueous dispersions or in the form of organic solvent systems, butadiene/styrene/acrylonitrile rubbers, drying oils such as linseed oil, resin esters or modified hardened resins in combination with tar or bitumens, asphalt and epoxy compounds, small amounts of chlorine rubber, chlorinated polypropylene and vinyl
30 resins.

If appropriate, paints also comprise inorganic pigments, organic pigments or colorants which are preferably insoluble in saltwater. Paints may furthermore comprise materials such as colophony to allow controlled release of the active compounds. Furthermore, the paints may comprise plasticizers, modifiers which affect the rheological properties and other conventional constituents. The compounds according to the invention or the abovementioned mixtures may also be incorporated into self-polishing antifouling systems.

The active compound combinations are also suitable for controlling animal pests, in particular insects, arachnids and mites, which are found in enclosed spaces such as, for example, dwellings, factory halls, offices, vehicle cabins and the like. They can be employed in domestic insecticide products for controlling these pests. They are active against sensitive and resistant species and against all developmental stages. These pests include:

From the order of the Scorpionidea, for example, *Buthus occitanus*.

From the order of the Acarina, for example, *Argas persicus*, *Argas reflexus*, *Bryobia* spp., *Dermanyssus gallinae*, *Glyciphagus domesticus*, *Ornithodoros moubat*, *Rhipicephalus sanguineus*, *Trombicula alfreddugesi*, *Neutrombicula autumnalis*, *Dermatophagoides pteronissimus*, *Dermatophagoides forinae*.

From the order of the Araneae, for example, *Aviculariidae*, *Araneidae*.

From the order of the Opiliones, for example, *Pseudoscorpiones chelifer*, *Pseudoscorpiones cheiridium*, *Opiliones phalangium*.

From the order of the Isopoda, for example, *Oniscus asellus*, *Porcellio scaber*.

From the order of the Diplopoda, for example, *Blaniulus guttulatus*, *Polydesmus* spp.

From the order of the Chilopoda, for example, *Geophilus* spp.

From the order of the Zygentoma, for example, *Ctenolepisma* spp., *Lepisma saccharina*, *Lepismodes inquilinus*.

From the order of the Blattaria, for example, *Blatta orientalis*, *Blattella germanica*, *Blattella asahinai*, *Leucophaea maderae*, *Panchlora* spp., *Parcoblatta* spp., *Periplaneta australasiae*, *Periplaneta americana*, *Periplaneta brunnea*, *Periplaneta fuliginosa*, *Supella longipalpa*.

From the order of the Saltatoria, for example, *Acheta domesticus*.

From the order of the Dermaptera, for example, *Forficula auricularia*.

From the order of the Isoptera, for example, *Kalotermes* spp., *Reticulitermes* spp.

From the order of the Psocoptera, for example, *Lepinatus* spp., *Liposcelis* spp.

- 5 From the order of the Coleoptera, for example, *Anthrenus* spp., *Attagenus* spp., *Dermestes* spp., *Latheticus oryzae*, *Necrobia* spp., *Ptinus* spp., *Rhizopertha dominica*, *Sitophilus granarius*, *Sitophilus oryzae*, *Sitophilus zeamais*, *Stegobium paniceum*.

- From the order of the Diptera, for example, *Aedes aegypti*, *Aedes albopictus*, *Aedes taeniorhynchus*, *Anopheles* spp., *Calliphora erythrocephala*, *Chrysozona pluvialis*,
10 *Culex quinquefasciatus*, *Culex pipiens*, *Culex tarsalis*, *Drosophila* spp., *Fannia canicularis*, *Musca domestica*, *Phlebotomus* spp., *Sarcophaga carnaria*, *Simulium* spp., *Stomoxys calcitrans*, *Tipula paludosa*.

From the order of the Lepidoptera, for example, *Achroia grisella*, *Galleria mellonella*, *Plodia interpunctella*, *Tinea cloacella*, *Tinea pellionella*, *Tineola bisselliella*.

- 15 From the order of the Siphonaptera, for example, *Ctenocephalides canis*, *Ctenocephalides felis*, *Pulex irritans*, *Tunga penetrans*, *Xenopsylla cheopis*.

From the order of the Hymenoptera, for example, *Camponotus herculeanus*, *Lasius fuliginosus*, *Lasius niger*, *Lasius umbratus*, *Monomorium pharaonis*, *Paravespula* spp., *Tetramorium caespitum*.

- 20 From the order of the Anoplura, for example, *Pediculus humanus capitis*, *Pediculus humanus corporis*, *Phthirus pubis*.

From the order of the Heteroptera, for example, *Cimex hemipterus*, *Cimex lectularius*, *Rhodinus prolixus*, *Triatoma infestans*.

- 25 They are used as aerosols, pressure-free spray products, for example pump and atomizer sprays, automatic fogging systems, foggers, foams, gels, evaporator products with evaporator tablets made of cellulose or polymer, liquid evaporators, gel and membrane evaporators, propeller-driven evaporators, energy-free, or passive, evaporation systems, moth papers, moth bags and moth gels, as granules or dusts, in baits for spreading or in bait stations.

5 All plants and plant parts can be treated in accordance with the invention. Plants are to be understood as meaning in the present context all plants and plant populations such as desired and undesired wild plants or crop plants (inclusive of naturally occurring crop plants). Crop plants can be plants which can be obtained by conventional plant breeding and optimization methods or by biotechnological and genetical engineering or by combinations of these methods, inclusive of the transgenic plants and inclusive of the plant varieties protectable or not protectable by plant breeders' rights, such as shoot, leaf, flower and root, examples which may be mentioned being leaves, needles, stalks, stems, flowers, fruit bodies, fruits, seeds, 10 roots, tubers and rhizomes. The plant parts also include harvested material, and vegetative and generative propagation material, for example cuttings, tubers, rhizomes, offsets and seeds.

15 Treatment according to the invention of the plants and plant parts with the active compounds is carried out directly or by allowing the compounds to act on their surroundings, environment or storage space by the customary treatment methods, for example by immersion, spraying, evaporation, fogging, scattering, painting on and, in the case of propagation material, in particular in the case of seed, also by applying one or more coats.

20 As already mentioned above, it is possible to treat all plants and their parts according to the invention. In a preferred embodiment, wild plant species and plant varieties, or those obtained by conventional biological breeding methods, such as crossing or protoplast fusion, and parts thereof, are treated. In a further preferred embodiment, transgenic plants and plant varieties obtained by genetical engineering, if appropriate 25 in combination with conventional methods (Genetic Modified Organisms), and parts thereof are treated. The term "parts" or "parts of plants" or "plant parts" has been explained above.

30 Particularly preferably, plants of the plant varieties which are in each case commercially available or in use are treated according to the invention.

Depending on the plant species or plant varieties, their location and growth conditions (soils, climate, vegetation period, diet), the treatment according to the invention may also result in superadditive ("synergistic") effects. Thus, for example, reduced application rates and/or a widening of the activity spectrum and/or an increase in the activity of the substances and compositions which can be used according to the invention, better plant growth, increased tolerance to high or low temperatures, increased tolerance to drought or to water or soil salt content, increased flowering performance, easier harvesting, accelerated maturation, higher harvest yields, better quality and/or a higher nutritional value of the harvested products, better storage stability and/or processability of the harvested products are possible which exceed the effects which were actually to be expected.

The transgenic plants or plant varieties (i.e. those obtained by genetical engineering) which are preferred and to be treated according to the invention include all plants which, in the genetic modification, received genetic material which imparts particularly advantageous useful properties ("traits") to these plants. Examples of such properties are better plant growth, increased tolerance to high or low temperatures, increased tolerance to drought or to water or soil salt content, increased flowering performance, easier harvesting, accelerated maturation, higher harvest yields, better quality and/or a higher nutritional value of the harvested products, better storage stability and/or processability of the harvested products. Further and particularly emphasized examples of such properties are a better defence of the plants against animal and microbial pests, such as against insects, mites, phytopathogenic fungi, bacteria and/or viruses, and also increased tolerance of the plants to certain herbicidally active compounds. Examples of transgenic plants which may be mentioned are the important crop plants, such as cereals (wheat, rice), maize, soya beans, potatoes, cotton, oilseed rape and also fruit plants (with the fruits apples, pears, citrus fruits and grapevines), and particular emphasis is given to maize, soya beans, potatoes, cotton and oilseed rape. Traits that are particularly emphasized are increased defence of the plants against insects by toxins formed in the plants, in particular those formed by the genetic material from *Bacillus thuringiensis* (for

example by the genes CryIA(a), CryIA(b), CryIA(c), CryIIA, CryIIIA, CryIIIB2, Cry9c, Cry2Ab, Cry3Bb and CryIF and also combinations thereof) (hereinbelow referred to as "Bt plants"). Traits that are furthermore particularly emphasized are the increased tolerance of the plants to certain herbicidally active compounds, for example imidazolinones, sulphonylureas, glyphosate or phosphinotricin (for example the "PAT" gene). The genes which impart the desired traits in question can also be present in combination with one another in the transgenic plants. Examples of "Bt plants" which may be mentioned are maize varieties, cotton varieties, soya bean varieties and potato varieties which are sold under the trade names YIELD GARD® (for example maize, cotton, soya beans), KnockOut® (for example maize), StarLink® (for example maize), Bollgard® (cotton), Nucotn® (cotton) and NewLeaf® (potato). Examples of herbicide-tolerant plants which may be mentioned are maize varieties, cotton varieties and soya bean varieties which are sold under the trade names Roundup Ready® (tolerance to glyphosate, for example maize, cotton, soya beans), Liberty Link® (tolerance to phosphinotricin, for example oilseed rape), IMI® (tolerance to imidazolinones) and STS® (tolerance to sulphonylureas, for example maize). Herbicide-resistant plants (plants bred in a conventional manner for herbicide tolerance) which may be mentioned include the varieties sold under the name Clearfield® (for example maize). Of course, these statements also apply to plant varieties having these traits or genetic traits still to be developed, which plants will be developed and/or marketed in the future.

The plants listed can be treated according to the invention in a particularly advantageous manner with the active compound mixtures according to the invention. The preferred ranges stated above for the mixtures also apply to the treatment of these plants. Particular emphasis is given to the treatment of plants with the mixtures specifically mentioned in the present text.

The good insecticidal and acaricidal action of the active compound combinations according to the invention can be seen from the examples which follow. While the

individual active compounds show weaknesses in their action, the combinations show an action which exceeds a simple sum of actions.

5 A synergistic effect in insecticides and acaricides is always present when the action of the active compound combinations exceeds the total of the actions of the active compounds when applied individually.

The expected action for a given combination of two active compounds can be calculated as follows, using the formula of S.R. Colby, Weeds 15 (1967), 20-22:

10

If

X is the kill rate, expressed as % of the untreated control, when employing active compound A at an application rate of m g/ha or in a concentration of m ppm,

15

Y is the kill rate, expressed as % of the untreated control, when employing active compound B at an application rate of n g/ha or in a concentration of n ppm and

20

E is the kill rate, expressed as % of the untreated control, when employing active compounds A and B at application rates of m and n g/ha or in a concentration of m and n ppm,

25 then

$$E = X + Y - \frac{X \cdot Y}{100}$$

If the actual insecticidal kill rate exceeds the calculated value, the action of the combination is superadditive, i.e. a synergistic effect is present. In this case, the

30

actually observed kill rate must exceed the value calculated using the above formula for the expected kill rate (E).

5 After the desired period of time, the kill in % is determined. 100% means that all animals have been killed; 0% means that none of the animals have been killed.

Throughout this specification and the claims which follow, unless the context requires otherwise, the word "comprise", or variations such as "comprises" or "comprising", will be understood to imply the inclusion of a stated integer or group of integers or steps but not the exclusion of any other integer or group of integers or steps.

Persons skilled in the art will appreciate that numerous variations and modifications will become apparent. All such variations and modifications which become apparent to persons skilled in the art, should be considered to fall within the spirit and scope that the invention broadly appearing before described.

The reference to any prior art in this specification is not, and should not be taken as, an acknowledgement or any form of suggestion that the prior art forms part of the common general knowledge in Australia.

In order that the invention may be readily understood and put into practical effect, particular preferred embodiments will now be described by way of the following, non-limiting Examples.

Example A

Plutella Test (normally sensitive)

- 5 Solvent: 7 parts by weight of dimethylformamide
Emulsifier: 2 parts by weight of alkylaryl polyglycol ether

10 To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amounts of solvent and emulsifier, and the concentrate is diluted with emulsifier-containing water to the desired concentration.

15 Cabbage leaves (*Brassica oleracea*) are treated by being dipped into the preparation of active compound of the desired concentration, and are populated with normally sensitive caterpillars of the diamondback moth (*Plutella xylostella*) while the leaves are still moist.

After the desired period of time, the kill in % is determined. 100% means that all caterpillars have been killed; 0% means that none of the caterpillars have been killed. The kill rates determined are calculated using Colby's formula.

20 In this test, the following active compound combination according to the present application showed a synergistically enhanced activity compared to the active compounds applied individually:

Table A
plant-damaging insects
Plutella Test (normally sensitive)

Active compounds	Active compound concentration in ppm	Kill rate in % after 3 ^d	
Ex. I-b-1			
known	100	40	
Indoxacarb			
known	0.16	15	
Ex. I-b-1 + indoxacarb			
(625:1)	100 + 0.16	<u>found*</u>	<u>calc.**</u>
according to the invention		70	49

5

found* = Activity found

calc.** = Activity calculated using Colby's formula

Example B

Plutella Test (resistant)

- 5 Solvent: 7 parts by weight of dimethylformamide
Emulsifier: 2 parts by weight of alkylaryl polyglycol ether

10 To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amounts of solvent and emulsifier, and the concentrate is diluted with emulsifier-containing water to the desired concentration.

15 Cabbage leaves (*Brassica oleracea*) are treated by being dipped into the preparation of active compound of the desired concentration and are populated with resistant caterpillars of the diamondback moth (*Plutella xylostella*) while the leaves are still moist.

20 After the desired period of time, the kill in % is determined. 100% means that all caterpillars have been killed; 0% means that none of the caterpillars have been killed. The kill rates determined are calculated using Colby's formula.

In this test, the following active compound combination according to the present application showed a synergistically enhanced activity compared to the active compounds applied individually:

Table B-1
 plant-damaging insects
Plutella Test (resistant)

Active compounds	Active compound concentration in ppm	Kill rate in % after 7 ^d	
Ex. I-b-1			
Known	100	35	
Fipronil			
Known	0.032	40	
Ex. I-b-1 + fipronil			
(3125:1)	100 + 0.032	<u>found*</u>	<u>calc.**</u>
According to the invention		90	61

5

Table B-2
 plant-damaging insects
Plutella Test (resistant)

Active compounds	Active compound concentration in ppm	Kill rate in % after 7 ^d	
Ex. I-b-1			
Known	500	40	
Indoxacarb			
Known	0.16	90	
Ex. I-b-1 + indoxacarb			
(3125:1)	500 + 0.16	<u>found*</u>	<u>calc.**</u>
According to the invention		100	94

10

found* = Activity found

calc.** = Activity calculated using Colby's formula

Example C

Spodoptera frugiperda Test

5

Solvent: 7 parts by weight of dimethylformamide
Emulsifier: 2 parts by weight of alkylaryl polyglycol ether

10 To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amounts of solvent and emulsifier, and the concentrate is diluted with emulsifier-containing water to the desired concentration.

15 Cabbage leaves (*Brassica oleracea*) are treated by being dipped into the preparation of active compound of the desired concentration and are populated with caterpillars of the army worm (*Spodoptera frugiperda*) while the leaves are still moist.

After the desired period of time, the kill in % is determined. 100% means that all caterpillars have been killed; 0% means that none of the caterpillars have been killed. The kill rates determined are calculated using Colby's formula.

20

In this test, the following active compound combination according to the present application showed a synergistically enhanced activity compared to the active compounds applied individually:

Table C-1
 plant-damaging insects
Spodoptera frugiperda Test

Active compounds	Active compound concentration in ppm	Kill rate in % after 7 ^d	
Ex. I-b-1			
Known	100	0	
Emamectin-benzoate			
Known	0.16	55	
Ex. I-b-1 + emamectin-benzoate (625:1)	100 + 0.16	<u>found*</u>	<u>calc.**</u>
According to the invention		80	55

5

Table C-2
 plant-damaging insects
Spodoptera frugiperda Test

Active compounds	Active compound concentration in ppm	Kill rate in % after 7 ^d	
Ex. I-b-1			
Known	100	0	
Fipronil			
Known	0.8	20	
Ex. I-b-1 + fipronil (125:1)	100 + 0.8	<u>found*</u>	<u>calc.**</u>
According to the invention		55	20

10

found* = Activity found

calc.** = Activity calculated using Colby's formula

Example D

Critical concentration test/soil insects – treatment of transgenic plants

- 5 Test insect: **Diabrotica balteata – larvae in soil**
- Solvent: 7 parts by weight of acetone
- Emulsifier: 1 part by weight of alkylaryl polyglycol ether
- 10 To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amount of solvent, the stated amount of emulsifier is added and the concentrate is diluted with water to the desired concentration.
- 15 The preparation of active compound is poured onto the soil. Here, the concentration of the active compound in the preparation is virtually irrelevant, only the amount by weight of active compound per volume unit of soil, which is stated in ppm (mg/l), matters. The soil is filled into 0.25 l pots and these are allowed to stand at 20°C.
- 20 Immediately after preparation, 5 pre-germinated maize cobs of the variety YIELD GUARD (trade mark of Monsanto Comp., USA) are placed into each pot. After 2 days, the test insects in question are placed into the treated soil. After a further 7 days, the efficacy of the active compound is determined by counting the maize plants that have emerged (1 plant = 20% efficacy).

Example E

Heliothis virescens test - treatment of transgenic plants

- 5 Solvent: 7 parts by weight of acetone
Emulsifier: 1 part by weight of alkylaryl polyglycol ether


10 To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amount of solvent and the stated amount of emulsifier, and the concentrate is diluted with water to the desired concentration.

15 Soya bean shoots (glycine max) of the variety Roundup Ready (trade mark of Monsanto Comp. USA) are treated by being dipped into the preparation of active compound of the desired concentration and are populated with the tobacco budworm *Heliothis virescens* while the leaves are still moist.

After the desired period of time, the kill of the insects is determined.

The claims defining the invention are as follows:

1. Composition, comprising mixtures of compounds of the formula



5

and at least one of the following compounds

chlorfluazuron

diflubenzuron

10 lufenuron

teflubenzuron

triflumuron

novaluron

flufenoxuron

15 hexaflumuron

emamectin

methoxyfenozide

tebufenozide

halofenozide

20 endosulfan

Trichlogramma spp.

Verticillium lecanii

fipronil

cyromazine

2001258303 17 Feb 2005

- 46 -

azadirachtin
diofenolan
indoxacarb
chromafenozide
5 bistrifluoron

2. Use of compositions as defined in Claim 1 for controlling animal pests.
3. Method for controlling animal pests, wherein a composition as defined
10 in Claim 1 is allowed to act on animal pests and/or their habitat.
4. Process for preparing insecticidal and acaricidal compositions, wherein
a composition as defined in Claim 1 are mixed with extenders and/or
15 surfactants.
5. Composition as defined in Claim 1, use thereof, or a process of
preparing said composition, substantially as hereinbefore described,
with reference to the accompanying Examples.

20 DATED this 16th day of February, 2005

BAYER AKTIENGESELLSCHAFT

by its Patent Attorneys

DAVIES COLLISON CAVE