




(86) Date de dépôt PCT/PCT Filing Date: 1998/12/16  
 (87) Date publication PCT/PCT Publication Date: 1999/07/01  
 (45) Date de délivrance/Issue Date: 2010/02/23  
 (85) Entrée phase nationale/National Entry: 2000/06/15  
 (86) N° demande PCT/PCT Application No.: IT 1998/000364  
 (87) N° publication PCT/PCT Publication No.: 1999/032516  
 (30) Priorité/Priority: 1997/12/19 (IT RM97A000796)

(51) Cl.Int./Int.Cl. *C12N 15/12* (2006.01),  
*A61K 38/17* (2006.01), *A61K 48/00* (2006.01),  
*C07K 14/47* (2006.01), *C12N 15/10* (2006.01),  
*C12N 15/86* (2006.01), *C12P 21/02* (2006.01),  
*A61K 38/00* (2006.01)  
 (72) Inventeurs/Inventors:  
 BOTTAZZI, BARBARA, IT;  
 INTRONA, MARTINO, IT;  
 MANTOVANI, ALBERTO, IT;  
 VECCHI, ANNUNCIATA, IT  
 (73) Propriétaire/Owner:  
 SIGMA-TAU INDUSTRIE FARMACEUTICHE RIUNITE  
 S.P.A., IT  
 (74) Agent: FETHERSTONHAUGH & CO.

(54) Titre : COMPOSITIONS PHARMACEUTIQUES CONTENANT UNE PENTRAXINE LONGUE PTX3  
 (54) Title: PHARMACEUTICAL COMPOSITIONS CONTAINING THE LONG PENTRAXIN PTX3



(57) Abrégé/Abstract:

Pharmaceutical compositions are described containing a long pentraxin PTX3, particularly human PTX3, for the therapy of infectious and inflammatory or tumour diseases; expression vectors containing cDNA coding for PTX3; recombinant host cells transfected with such vectors; a method for producing substantial amounts of PTX3 involving the culturing of such cells, and the use of said vectors in the gene therapy of tumours.

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification <sup>6</sup> : C07K 14/47, C12N 15/12, A61K 38/17, C12N 15/10, C12P 21/02, A61K 48/00</p>	A3	<p>(11) International Publication Number: <b>WO 99/32516</b></p> <p>(43) International Publication Date: 1 July 1999 (01.07.99)</p>
<p>(21) International Application Number: PCT/IT98/00364</p> <p>(22) International Filing Date: 16 December 1998 (16.12.98)</p> <p>(30) Priority Data: RM97A000796 19 December 1997 (19.12.97) IT</p> <p>(71) Applicant (for all designated States except US): SIGMA-TAU INDUSTRIE FARMACEUTICHE RIUNITE S.P.A. [IT/IT]; Viale Shakespeare, 47, I-00144 Rome (IT).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): BOTTAZZI, Barbara [IT/IT]; Via Cascina Bianca, 28/2, I-20142 Milan (IT). INTRONA, Martino [IT/IT]; Via Balzarotti, 12/I, I-20017 Rho (IT). MANTOVANI, Alberto [IT/IT]; Largo Brasilia, 4, I-20100 Milan (IT). VECCHI, Annunciata [IT/IT]; Via Spontini, 11, I-20131 Milan (IT).</p> <p>(74) Common Representative: SIGMA-TAU INDUSTRIE FAR- MACEUTICHE RIUNITE S.P.A.; Viale Skakespeare, 47, I-00144 Rome (IT).</p>		<p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p><b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p> <p>(88) Date of publication of the international search report: 28 October 1999 (28.10.99)</p>
<p>(54) Title: PHARMACEUTICAL COMPOSITIONS CONTAINING THE LONG PENTRAXIN PTX3</p>		
<p>(57) Abstract</p> <p>Pharmaceutical compositions are described containing a long pentraxin PTX3, particularly human PTX3, for the therapy of infectious and inflammatory or tumour diseases; expression vectors containing cDNA coding for PTX3; recombinant host cells transfected with such vectors; a method for producing substantial amounts of PTX3 involving the culturing of such cells, and the use of said vectors in the gene therapy of tumours.</p>		

**Pharmaceutical compositions containing the long pentraxin PTX3**

5           The present invention relates to pharmaceutical compositions containing the long pentraxin PTX3 (PTX3) or one of its functional derivatives. In particular, the invention relates to the aforesaid compositions for the therapy of infectious and inflammatory diseases or tumours.

10           The invention also relates to expression vectors containing the complete cDNA sequence coding for PTX3 or one of its functional derivatives, recombinant host cells transfected with such expression vectors and a method for producing PTX3 or one of its functional derivatives. Further, the invention relates to gene therapy methods for  
15 the treatment of tumours, based on the use of the aforesaid expression vectors.

          To date, we have yet to fully understand the biological function of PTX3, a protein which is expressed in various types of cells, most notably in mononuclear phagocytes and endothelial cells, after  
20 exposure to the inflammatory cytokines Interleukin 1beta (IL-1beta) and Tumour Necrosis Factor alpha (TNF-alpha).

          To date, there has also been no description of any therapeutic use of PTX3 or of its functional derivatives.

PTX3 consists of two structural domains, an N-terminal unrelated to any known molecule, and a C-terminal similar to the short pentraxins such as C-reactive protein (CRP). A substantial degree of similarity has been found between human PTX3 (hPTX3) and  
5 animal PTX3s.

The PTX3 gene is located on chromosome 3 of the mouse in a region similar to the human 3q region (q24-28), in agreement with the documented location of hPTX3 in the 3q 25 region. Furthermore, mouse PTX3 (mPTX3) (Introna M., Vidal Alles V., Castellano M., Picardi  
10 G., De Gioia L., Bottazzi B., Peri G., Breviario F., Salmona M., De Gregorio L., Dragani T.A., Srinivasan N., Blundell T.L., Hamilton T.A. and Mantovani A.: Cloning of mouse PTX3, a new member of the pentraxin gene family expressed at extrahepatic sites. *Blood* 87 (1996) 1862-1872) is very similar to hPTX3 in terms of organisation, location  
15 and sequence (Breviario F., d'Aniello E.M., Golay J., Peri G., Bottazzi B., Bairoch A., Saccone S., Marzella R., Predazzi V., Rocchi M., Della Valle G., Dejana E., Mantovani A., Introna M.: Interleukin-1-inducible genes in endothelial cells. Cloning of a new gene related to C-reactive protein and serum amyloid P component. *J. Biol. Chem.* 267:22190,  
20 1992).

In particular, the degree of identity between the sequences is 82% between the human and mouse gene and reaches 90% if conservative substitutions are considered.

The high degree of similarity between the hPTX3 and mPTX3 sequences is a sign of the high degree of conservation of pentraxin during evolution (Pepys M.B., Baltz M.L.: Acute phase proteins with special reference to C-reactive protein and related proteins (pentraxins) and serum amyloid A protein. Adv. Immunol. 34: 141, 1983).

CRP is a marker for immuno-inflammatory and infectious disease. After a trauma, a lesion or infection of a tissue triggers off, in the affected subject, a complex series of reactions aimed at preventing extension of the damage, at destroying the infecting organism and at activating the repair process in order to restore normal function. This process constitutes the so-called acute-phase response, and the main marker currently used for the acute-phase response is CRP. In normal human serum, in fact, it is present in concentrations of less than 10 µg/ml, but can increase more than 1,000-fold in response to a trauma or inflammation (Koj A.: "Acute phase reactants" in "Structure and Function of Plasma Proteins". Allison A., ed. Plenum Press, New York, 1974, pp. 73-131).

Previous therapeutic uses of CRP are already known. For instance, US Patent 4,857,314 dated 15.08.1989 discloses the use of CRP in combination with TNF for the treatment of tumours.

PCT publication WO 94/18999 filed 24.02.1994 discloses mutants of CRP which are useful for the preparation of diagnostic kits for determining immunocomplexes in

biological fluids and for the treatment of viral and microbial diseases, tumours and endotoxic shock.

International patent application PCT/US94/09729 dated 26.08.1994 also discloses mutants of CRP which are useful for the preparation of diagnostic kits and for the treatment of viral and microbial diseases and tumours.

The ability of PTX3 to recognise and bind specifically to ligands which are also recognised by short pentraxins has been evaluated *in vitro* using purified recombinant PTX3. Short pentraxins such as CRP and SAP (serum amyloid P component) are characterised by their ability to recognise and bind in a calcium-dependent manner to a broad spectrum of ligands, including phosphocholine, phosphoethanolamine, many sugars, the best characterised of which is an agarose derivative rich in pyruvate [methyl 4-6-O-(1-carboxyethylidene)-beta-D-galacto-pyranoside] or MO $\beta$ DG, complement fragments and proteins of the extracellular matrix, particularly fibronectin and type IV collagen. Unlike the short pentraxins, PTX3 is unable to bind either calcium (assessed by Inductive Coupled Plasma/Atomic Emission Spectroscopy) or phosphocholine, phosphoethanolamine or MO $\beta$ DG. Moreover, PTX3 is unable to bind extracellular matrix proteins such as fibronectin or type IV collagen. On the other hand, PTX3 is capable of binding the C1q complement fragment which is also recognised by the short pentraxins (Table 1). It

should be stressed, however, that, whereas CRP and SAP have to be cross-linked to bind C1q, PTX3 is capable of recognising and binding this complement fragment in the naturally occurring form.

Surprisingly, it has now been found that the long pentraxin PTX3 or its  
5 functional derivatives are useful therapeutic agents, particularly for the therapy of infectious and inflammatory diseases or tumours.

In one aspect, there is provided orally, parenterally, transdermally or subcutaneously administrable pharmaceutical composition comprising a polypeptide comprising the amino acid sequence of human long pentraxin PTX3  
10 as set forth in SEQ ID NO: 1, and a pharmacologically acceptable excipient, wherein the polypeptide binds to C1q and recruits leukocytes.

In another aspect, there is provided use of a tumour cell from a patient suffering from a cancer, said cell transfected with an expression vector comprising a sequence coding for a polypeptide comprising the amino acid  
15 sequence of human long pentraxin PTX3 as set forth in SEQ ID NO: 1, for gene therapy of a tumour in said patient, wherein the polypeptide binds to C1q and recruits leukocytes.

In another aspect, there is provided use of a tumour cell from a patient suffering from a cancer, said cell transfected with an expression vector  
20 comprising a sequence coding for a polypeptide comprising the amino acid sequence of human long pentraxin PTX3 as set forth in SEQ ID NO: 1, for preparing a medicament for gene therapy of a tumour in said patient, wherein the

- 5a -

polypeptide binds to C1q and recruits leukocytes.

In another aspect, there is provided use of a viral expression vector comprising a sequence coding for a polypeptide comprising the amino acid sequence of human long pentraxin PTX3 as set forth in SEQ ID NO: 1, for gene  
5 therapy of a tumour condition, wherein the polypeptide binds to C1q and recruits leukocytes.

In another aspect, there is provided use of a viral expression vector comprising a sequence coding for a polypeptide comprising the amino acid sequence of human long pentraxin PTX3 as set forth in SEQ ID NO: 1, for  
10 preparing a medicament for gene therapy of a tumour condition, wherein the polypeptide binds to C1q and recruits leukocytes.

In another aspect, there is provided use of human long pentraxin PTX3 having the sequence as set forth in SEQ ID NO: 1 for the treatment of an infectious, or tumour disease, wherein the long pentraxin PTX3 binds to C1q and  
15 recruits leukocytes.

In another aspect, there is provided use of human long pentraxin PTX3 having the sequence as set forth in SEQ ID NO: 1 for the preparation of a medicament for the treatment of an infectious, or tumour disease, wherein the long pentraxin PTX3 binds to C1q and recruits leukocytes.

20 A convenient method of producing substantial amounts of long pentraxin PTX3 or one of its functional derivatives consists in preparing expression vectors (e.g. plasmids) containing the complete cDNA sequence coding for

- 5b -

PTX3 or one of its functional derivatives and in using these to transfer eukaryotic cells in culture (e.g. Chinese hamster ovary cells, CHO). After cloning the recombinant host cells thus transfected, the cell clone capable of producing the highest levels of PTX3 is selected.

- 5           According to the present invention, the above-mentioned expression vectors containing the cDNA sequence coding for long pentraxin PTX3 are also utilised in gene therapy methods for the treatment of tumour conditions.

A first gene therapy method consists in:

- a) taking samples of cells from a patient suffering from a tumour;
- b) transfecting these cells with an expression vector containing the complete cDNA sequence coding for long pentraxin PTX3 or one of its functional derivatives; and
- c) inoculating the tumour patient with these transfected cells.

A second gene therapy method for the treatment of tumours consists in:

- a) preparing an expression vector of viral origin (such as an adenovirus or retrovirus) containing the complete cDNA sequence coding for long pentraxin PTX3 or one of its functional derivatives; and
- b) injecting the tumour affected patient with the expression vector thus obtained.

Though the mechanism of action of PTX3 or its functional derivatives has yet to be definitively clarified, their anticancer activity in any event is not attributable to a direct cytolytic or cytostatic effect on the tumour cells, but rather to mechanisms mediated by the host and related to the leukocyte recruitment ability exerted by these compounds, as will be described below.

- 7 -

There now follows a description of the experimental procedures and results are reported demonstrating the unexpected activity of the compounds according to the invention described herein.

**Production of recombinant PTX3:** a fragment containing the complete  
5 cDNA sequence of human PTX3 (sequence 2 and Fig. 6) was subcloned in the Bam H1 site of the expression vector pSG5 (Fig. 1) (Stratagene, La Jolla, CA, USA) and transfected in CHO cells using the precipitated calcium procedure. A clone selected in G418, capable of producing large amounts of PTX3, was used  
10 as a source from which the protein was purified by chromatography by means of ion exchange and gel filtration.

**Binding of PTX3 to C1q:** the binding of PTX3 to C1q was assessed in an ELISA system. A 96-well plate was covered with 250-500 ng of C1q per well (one night at 4°C) and then washed with PBS with Ca<sup>++</sup> and Mg<sup>++</sup> containing 0.05% Tween<sup>™</sup> 20 (PBS). The wells were then blocked with 5% milk in PBS (2  
15 h at room temperature) and subsequently incubated with variable concentrations of PTX3 (30 min at 37°C). After a further series of washings, the plate was incubated with a rat monoclonal antibody to PTX3 (1 h at room temperature) and then with the second antibody, a peroxidase-conjugated rat anti-IgG antibody (1 h at room temperature). After washing, chromogen was added and  
20 absorbance was read at 405 nm using an automatic plate

reader. In a number of experiments, the wells were covered with PTX3 and C1q binding was evaluated using an anti-C1q antibody.

Biotinylated protein was used to determine the C1q binding affinity. PTX3 was biotinylated according to standard procedures using  
5 an activated biotin modified by the addition of a "spacer arm". (SPA - Società Prodotti Antibiotici).

Figures 2(A) and 2(B) give the C1q binding and binding affinity results. These results show the very substantial degree of C1q binding and binding affinity of PTX3.

10 **Leukocyte recruitment:** the leukocyte recruitment induced by PTX3 was studied *in vivo* in the subcutaneous pocket system. The subcutaneous pocket was induced in the experimental animal by means of two subcutaneous injections of 5 mL of air into the animal's back with an intervening interval of three days. On day 6, 1 µg of PTX3  
15 in 0.5% carboxymethylcellulose was administered into the pocket. After 4 h, the animals were anaesthetised and the pocket was washed with 1 mL of saline solution. The washing liquid was recovered and was submitted to a total count and a differential count of the cells present.

20 The results obtained are reported in Figure 3 and show the substantial leukocyte recruitment capacity of PTX3 in normal animals, whereas Figure 4 shows the results obtained in genetically modified

animals, without C1q, in which the leukocyte recruitment is significantly lower.

**Anticancer activity:** a line of murine mastocytoma P815 was co-transfected by electroporation with the expression vector pSG5 containing the cDNA of human PTX3 or its antisense and the vector pSV2 which endows the transfected cells with neomycin resistance. After selection with neomycin 0.5 mg/mL, the cells were cloned by limit dilution.

To assess the production of PTX3,  $2.5 \times 10^5$  cells were cultivated in 200  $\mu$ L of RPMI + 3% FCS for 24 h and the supernatant was tested by ELISA. The clones obtained produced protein levels ranging from 1 to 35 ng/mL, while the clones containing the antisense produced no measurable levels of PTX3. The clones considered showed the same growth rate *in vivo*.

Male DBA/2N CrlBR mice aged 8-10 weeks were subcutaneously injected with  $1 \times 10^5$  cells of P815 PTX3-producing clones or with clones containing the antisense gene. The mice were monitored 3 times daily for occurrence of tumours and once daily for survival.

The results obtained are reported in Table 2 and show the efficacy of PTX3, in this experimental model of gene therapy, in bringing about healing of the animals and complete rejection of the tumour after the take of the inoculated tumour cells.

These results are statistically significant with  $p < 0.01$  (Fisher test) both as compared to controls and to the group treated with the antisense.

In the light of these results it is clear that the anticancer activity reported above correlates closely with the leukocyte recruitment which occurs in the mouse as a result of recognition of the C1q by PTX3. In genetically modified mice, no such leukocyte recruitment occurs. The leukocyte recruitment capacity, on the basis of the anticancer activity of the compounds according to the invention, indicates that these compounds may also have a useful application in the treatment of diseases caused by pathogens such as bacteria, fungi, protozoa or viruses.

**TABLE 1** PENTRAXIN BINDING ABILITY TO VARIOUS LIGANDS

	CRP	SAP	PTX3
Ca <sup>2+</sup>	+	+	-
Phosphocholine	+	-	-
Phosphoethanolamine	+	+	-
MO $\beta$ DG	-	+	-
C1q	+	+	+
Type IV collagen	ND	+	-
Fibronectin	ND	+	-

ND: test not performed

**TABLE 2** *IN VIVO* ANTICANCER ACTIVITY OF PTX3

	Clone <sup>1</sup>	Reject <sup>2</sup>
5	Parent P815 (control)	4/25
	P815-AS1 (antisense)	3/8
10	P815-PTX3-1 (sense)	14/14*

1 :  $1 \times 10^5$  cells of the clone indicated were injected subcutaneously.

15 2 : Number of animals that definitely reject the tumour out of total number of animals in which it took.

\* :  $p < 0.01$  as compared both to mice treated with parent cells and to mice treated with cells of the antisense clones (Fisher test).

WO 99/32516

PCT/IT98/00364

12

**Brief description of drawings**

**Figure 2:** PTX3 binding to C1q. Panel A shows the binding of the supernatant of the culture containing PTX3 (sense) and of the purified protein to C1q and C1s immobilised on plate. The binding is assessed as optical density (O.D.) at 405 nm. Panel B shows the saturation curve obtained with the biotinylated protein. The kinetic parameters were calculated using the non-linear fitting statistical method.

**Figure 3:** PTX3-induced leukocyte recruitment: 1  $\mu$ g of FTX3 is injected into a subcutaneous pocket induced in the back of CD 1 mice by inoculation of 5 ml of air.

**Figure 4:** PTX3-induced leukocyte recruitment in normal animals and in genetically modified animals without, C1q. PTX3 is injected into a subcutaneous induced on the back of the animals.

**Sequence SEQ. ID. NO 1:** Amino acid sequence of human FITX3.

The underlined amino acids constitute the peptide signal.

Mature hPTX3 consists of 364 amino acids.

**Sequence SEQ. ID. NO 2:** Nucleotide sequence of fragment of cDNA of human PTX3. Upper case letters denote nucleotides coding for the protein, while lower case letters denote regions at 3' and 5' not translated but present in the construct.

AMENDED SHEET

WO 99/32516

PCT/IT98/00364

1

## SEQUENCE LISTING

<110> sigma tau industrie farmaceutiche riunite s.p.a.

<120> Pharmaceutical compositions containing the long pentraxin PTX3 for the therapy of infectious and inflammatory diseases or tumours, expression vectors containing cDNA coding for PTX3, and use of such v

<130> long human pentraxin

<140>

<141>

<150> RM97A000796

<151> 1997-12-19

<160> 2

<170> PatentIn Ver. 2.0

<210> 1

<211> 381

<212> PRT

<213> HUMAN LONG PENTRAXIN PTX3

<400> 1

Met His Leu Leu Ala Ile Leu Phe Cys Ala Leu Trp Ser Ala Val Leu  
 1 5 10 15

Ala Glu Asn Ser Asp Asp Tyr Asp Leu Met Tyr Val Asn Leu Asp Asn  
 20 25 30

Glu Ile Asp Asn Gly Leu His Pro Thr Glu Asp Pro Thr Pro Cys Asp  
 35 40 45

Cys Gly Gln Glu His Ser Glu Trp Asp Lys Leu Phe Ile Met Leu Glu  
 50 55 60

Asn Ser Gln Met Arg Glu Arg Met Leu Leu Gln Ala Thr Asp Asp Val  
 65 70 75 80

Leu Arg Gly Glu Leu Gln Arg Leu Arg Glu Glu Leu Gly Arg Leu Ala  
 85 90 95

Glu Ser Leu Ala Arg Pro Cys Ala Pro Gly Ala Pro Ala Glu Ala Arg  
 100 105 110

WO 99/32516

PCT/IT98/00364

2

Leu Thr Ser Ala Leu Asp Glu Leu Leu Gln Ala Thr Arg Asp Ala Gly  
 115 120 125

Arg Arg Leu Ala Arg Met Glu Gly Ala Glu Ala Gln Arg Pro Glu Glu  
 130 135 140

Ala Gly Arg Ala Leu Ala Ala Val Leu Glu Glu Leu Arg Gln Thr Arg  
 145 150 155 160

Ala Asp Leu His Ala Val Gln Gly Trp Ala Ala Arg Ser Trp Leu Pro  
 165 170 175

Ala Gly Cys Glu Thr Ala Ile Leu Phe Pro Met Arg Ser Lys Lys Ile  
 180 185 190

Phe Gly Ser Val His Pro Val Arg Pro Met Arg Leu Glu Ser Phe Ser  
 195 200 205

Ala Cys Ile Trp Val Lys Ala Thr Asp Val Leu Asn Lys Thr Ile Leu  
 210 215 220

Phe Ser Tyr Gly Thr Lys Arg Asn Pro Tyr Glu Ile Gln Leu Tyr Leu  
 225 230 235 240

Ser Tyr Gln Ser Ile Val Phe Val Val Gly Gly Glu Glu Asn Lys Leu  
 245 250 255

Val Ala Glu Ala Met Val Ser Leu Gly Arg Trp Thr His Leu Cys Gly  
 260 265 270

Thr Trp Asn Ser Glu Glu Gly Leu Thr Ser Leu Trp Val Asn Gly Glu  
 275 280 285

Leu Ala Ala Thr Thr Val Glu Met Ala Thr Gly His Ile Val Pro Glu  
 290 295 300

Gly Gly Ile Leu Gln Ile Gly Gln Glu Lys Asn Gly Cys Cys Val Gly  
 305 310 315 320

Gly Gly Phe Asp Glu Thr Leu Ala Phe Ser Gly Arg Leu Thr Gly Phe  
 325 330 335

Asn Ile Trp Asp Ser Val Leu Ser Asn Glu Glu Ile Arg Glu Thr Gly  
 340 345 350

Gly Ala Glu Ser Cys His Ile Arg Gly Asn Ile Val Gly Trp Gly Val  
 355 360 365

WO 99/32516

PCT/IT98/00364

3

Thr Glu Ile Gln Pro His Gly Gly Ala Gln Tyr Val Ser  
 370 375 380

&lt;210&gt; 2

&lt;211&gt; 1404

&lt;212&gt; DNA

&lt;213&gt; HUMAN LONG PENTRAXIN PTX3

&lt;400&gt; 2

```

ctcaaactca gctcacttga gagtctcctc ccgccagctg tggaaagaac tttgcgtctc 60
tccagcaatg catctccttg cgattctggt ttgtgctctc tggctctgcag tgttggccga 120
gaactcggat gattatgatc tcatgtatgt gaatttggac aacgaaatag acaatggact 180
ccatcccact gaggaccca cgccgtgcga ctgcggtcag gagcactcgg aatgggacaa 240
gctcttcata atgctggaga actcgcagat gagagagcgc atgctgctgc aagccacgga 300
cgacgtcctg cggggcgagc tgcagaggct gcgggaggag ctgggcccggc tcgcggaaag 360
cctggcgagg ccgtgcgcgc cgggggctcc cgcagaggcc aggctgacca gtgctctgga 420
cgagctgctg caggcgacc gcgacgcggg ccgcaggctg gcgcgtatgg agggcgcgga 480
ggcgcagcgc ccagaggagg cggggcgcgc cctggcccgc gtgctagagg agctgcggca 540
gacgcgagcc gacctgcacg cgggtgcagg ctgggctgcc cggagctggc tgccggcagg 600
ttgtgaaaca gctattttat tccaatgcg ttccaagaag atttttggaa gcgtgcatcc 660
agtgagacca atgaggcttg agtcttttag tgctctcatt tgggtcaaag ccacagatgt 720
attaaacaaa accatcctgt ttcctatgg cacaagagg aatccatag aatccagct 780
gtatctcagc taccaatcca tagtgtttgt ggtgggtgga gaggagaaca aactggttgc 840
tgaagccatg gtttccctgg gaaggtggac ccacctgtgc ggcacctgga attcagagga 900
agggctcaca tccttggtgg taaatggtga actggcggct accactggtg agatggccac 960
aggtcacatt gttcctgagg gaggaatcct gcagattggc caagaaaaga atggctgctg 1020
tgtgggtggt ggctttgatg aacattagc cttctctggg agactcacag gcttcaatat 1080
ctgggatagt gttcttagca atgaagagat aagagagacc ggaggagcag agtcttgtca 1140
catccggggg aatattggtg ggtggggagt cacagagatc cagccacatg gaggagctca 1200
gtatgtttca taaatggtgt gaaactccac ttgaagcaa agaaagaaac tcacacttaa 1260
aacacatgcc agttgggaag gtctgaaaac tcagtgcata ataggaacac ttgagactaa 1320
tgaaagagag agttgagacc aatctttatt tgtactggcc aaatactgaa taaacagttg 1380
aaggaaagac attggaaaaa gctt

```

1404

**CLAIMS**

1. Orally, parenterally, transdermally or subcutaneously administrable pharmaceutical composition comprising a polypeptide comprising the amino acid sequence of human long pentraxin PTX3 as set forth in SEQ ID  
5 NO: 1, and a pharmacologically acceptable excipient, wherein the polypeptide binds to C1q and recruits leukocytes.
2. The composition according to claim 1, for the treatment of an infectious  
10 disease or tumour.
3. The composition according to claim 2, for the treatment of a disease caused  
by a bacterium, a fungus, a protozoa or a virus.
4. Use of a tumour cell from a patient suffering from a cancer, said cell  
15 transfected with an expression vector comprising a sequence coding for a polypeptide comprising the amino acid sequence of human long pentraxin PTX3 as set forth in SEQ ID NO: 1, for gene therapy of a tumour in said patient, wherein the polypeptide binds to C1q and recruits leukocytes.
- 20 5. Use of a tumour cell from a patient suffering from a cancer, said cell transfected with an expression vector comprising a sequence coding for a polypeptide comprising the amino acid sequence of human long pentraxin

- 14 -

PTX3 as set forth in SEQ ID NO: 1, for preparing a medicament for gene therapy of a tumour in said patient, wherein the polypeptide binds to C1q and recruits leukocytes.

5 6. Use of a viral expression vector comprising a sequence coding for a polypeptide comprising the amino acid sequence of human long pentraxin PTX3 as set forth in SEQ ID NO: 1, for gene therapy of a tumour condition, wherein the polypeptide binds to C1q and recruits leukocytes.

10 7. Use of a viral expression vector comprising a sequence coding for a polypeptide comprising the amino acid sequence of human long pentraxin PTX3 as set forth in SEQ ID NO: 1, for preparing a medicament for gene therapy of a tumour condition, wherein the polypeptide binds to C1q and recruits leukocytes.

15

8. Use according to claim 6 or claim 7, in which the viral expression vector is an adenovirus or retrovirus.


9. Use of human long pentraxin PTX3 having the sequence as set forth in  
20 SEQ ID NO: 1 for the treatment of an infectious, or tumour disease, wherein the long pentraxin PTX3 binds to C1q and recruits leukocytes.

- 15 -



10. Use of human long pentraxin PTX3 having the sequence as set forth in  
SEQ ID NO: 1 for the preparation of a medicament for the treatment of an  
infectious, or tumour disease, wherein the long pentraxin PTX3 binds to  
5 C1q and recruits leukocytes.
  
11. Use according to claim 9 or 10, wherein the infectious or tumour disease is  
caused by a bacterium, a fungus, a protozoa or a virus.

10


**FIGURE 1**




**FIGURE 2**



**FIGURE 3**



**FIGURE 4**



04-02-2000

PCT/IT98/00364

SEQ

WO 99/32516

PCT/IT98/00364

5/6

SEQ. ID. NO:  
SequenceVI

HLAALFECAL	WSAVLAENSD	DYDLMYVNL	NEIDNGLHPT	M . 1	41
EDPTPCDCGQ	EHSEWDKLF	MLENSQMRER	MLLQATDDVL		81
RGELQRLRE	LGRLAESLAR	PCAPGAPAEA	RLTSALDELL		121
QATSDAGRRL	ARMEGAEAQR	PEEAGRALAA	VLEELRQTRA		161
DLHAVQGWAA	RSWLFAGCET	AILFPMRSKK	IFGSVHPVRF		201
MRLESFSACT	WVKATDVLNK	TILFSYGTKR	NPVEIQLYLS		241
YQSIVFVGG	EENKLVAEAM	VSLGRWTHLC	GTWNSEEGLT		281
SLWVNGELAA	TTVEMATGHI	VPEGGILOIG	QEKNGCCVGG		321
GFDETLAFSG	RLTGFNIWDS	VLSNEEIRET	GGAESCHIRG		361
NIVGWGVTEI	QPHGGAQYVS				381

01-02-2000

PCT/TT98/00364

SEQ. ID. NO.

WO 99/32516

PCT/TT98/00364

6/6  
SEQ. ID. NO:  
SequenceV2

	CTCA AACCCAGCTC ACCGAGAGAGT CCCCCCCCCG CAGCCGCGGA AAGAACCTCG	54
5	CGCCCCCCCC GCAATGCATC TCCTTCCGAT TCTGTTTTGT GCTCTCTCGT CTGCAGTGTT	114
	GCCCCGAGAAC TCGGATGATT ATGATCTCAT GTATGTGAAT TTGGACAAAG AAATAGACAA	174
	TGGACTCCAT CCCACTGAGG ACCCCACGCC GTGCGACTGC GGTCCAGGAGC ACTCCGAATG	234
	CSACAAGETC TTCATCATGC TGGAGAACTC GCAGATGAGA GAGCCCATGC TGCTGCAAGC	294
	CACCGACGAC GTCTCTCGGG GCGAGCTGCA GAGGCTGCGG GAGGAGCTGG GCGGCTCCG	354
10	CGAAAGCCTG GCGAGCCCTT GCGGCCCCGG GCTCCCGCA GAGGCCAGGC TGACCAGTCC	414
	TCTGGACGAG CTGCTGCAGG CSACCCCGGA CCGGGGCGGC AGGCTGGCCG GTATGGAGGG	474
	CGCGGAGGCG CAGCGCCGAG AGGAGCGCGG GCGCGCCCTG GCGCGCGTGC TAGAGGAGCT	534
	GCGGCAGACG CGAGCCGACC TGCACGCGGT GCAGGGCTGG GCTGCCCGGA GCTGGCTGCC	594
	GCCAGCTTGT GAAACAGCTA TTTTATTCGC AATGCGTTCC AAGAAGATTT TTGGAAGCGT	654
15	GCATCCAGTG AGACCAATGA GGCTTGAGTC TTTTAGTGCC TGCATTTGGG TCAAASCCAC	714
	AGATGTATTA AACAAAACCA TCCTGTTTTT CTATGGCACA AAGAGGAATC CATATGAAAT	774
	CCAGCTGTAT CTCAGCTACC AATCCATAGT GTTTGTGGTG GGTGGAGAGG AGAACAAACT	834
	GGTTGCTGAA GCCATGGTTT CCCTGGGAAG GTGGACCCAC CTGTCCGGCA CCTGGGAATC	894
	AGAGGAAGGG CTCACATCCT TGTGGGTAAA TGGTGAAGTC GCGGCTACCA CTGTTGAGAT	954
20	GCCCCACAGG CACATTCTTC CTGAGGGAGG AATCCTGCAG AITGGCCAAG AAAAGAATGG	1014
	CTGCTGTGTG GGTGGTGGCT TTGATGAAAC ATTAGCCTTC TCTGGGAGAC TCACAGCCTT	1074
	CAATATCTGG GATAGTGTTC TTAGCAATGA AGAGATAAGA GAGACCGGAG GAGCAGAGTC	1134
	TTGTACATC CGGGGGAAAT TTGTTGGGTG GCGAGTCACA GAGATCCAGC CACATGGAGG	1194
	AGCTCACTAT GTTTGATAAA TGTCTGAAA CCCCACCTGA AGCCAAAGAAA GAACCTCAC	1254
25	ACCCAAAACA CATGCCAGTC GGGAAAGTCC GAAAACCTCAG TGCATCACTG GAACCTCTGA	1314
	GACCAACTGA AGAGAGAGTC GAGACCAATC CCCCCTCTGA CTGGCCAAAT ACTGAAATAA	1374
	CAGCTGAAAG AAGACACTCG GAAAAGCTCC	1404

**peptide  
signal**

**N-terminal  
domain**

**179**

**pentraxin  
domain**

**Hind 3**

**Bam H1**



**Bam H1**

**hPTX3**

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification <sup>6</sup> : C07K 14/47, C12N 15/12, A61K 38/17, C12N 15/10, C12P 21/02, A61K 48/00</p>	A3	<p>(11) International Publication Number: <b>WO 99/32516</b></p> <p>(43) International Publication Date: 1 July 1999 (01.07.99)</p>
<p>(21) International Application Number: PCT/IT98/00364</p> <p>(22) International Filing Date: 16 December 1998 (16.12.98)</p> <p>(30) Priority Data: RM97A000796 19 December 1997 (19.12.97) IT</p> <p>(71) Applicant (for all designated States except US): SIGMA-TAU INDUSTRIE FARMACEUTICHE RIUNITE S.P.A. [IT/IT]; Viale Shakespeare, 47, I-00144 Rome (IT).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): BOTTAZZI, Barbara [IT/IT]; Via Cascina Bianca, 28/2, I-20142 Milan (IT). INTRONA, Martino [IT/IT]; Via Balzarotti, 12/I, I-20017 Rho (IT). MANTOVANI, Alberto [IT/IT]; Largo Brasilia, 4, I-20100 Milan (IT). VECCHI, Annunciata [IT/IT]; Via Spontini, 11, I-20131 Milan (IT).</p> <p>(74) Common Representative: SIGMA-TAU INDUSTRIE FARMACEUTICHE RIUNITE S.P.A.; Viale Skakespeare, 47, I-00144 Rome (IT).</p>		<p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p><b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p> <p>(88) Date of publication of the international search report: 28 October 1999 (28.10.99)</p>
<p>(54) Title: PHARMACEUTICAL COMPOSITIONS CONTAINING THE LONG PENTRAXIN PTX3</p>		
<p>(57) Abstract</p> <p>Pharmaceutical compositions are described containing a long pentraxin PTX3, particularly human PTX3, for the therapy of infectious and inflammatory or tumour diseases; expression vectors containing cDNA coding for PTX3; recombinant host cells transfected with such vectors; a method for producing substantial amounts of PTX3 involving the culturing of such cells, and the use of said vectors in the gene therapy of tumours.</p>		