




(12) **DEMANDE DE BREVET CANADIEN
CANADIAN PATENT APPLICATION**
(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2019/11/07
(87) Date publication PCT/PCT Publication Date: 2020/05/14
(85) Entrée phase nationale/National Entry: 2021/05/07
(86) N° demande PCT/PCT Application No.: US 2019/060285
(87) N° publication PCT/PCT Publication No.: 2020/097344
(30) Priorités/Priorities: 2018/11/08 (US62/757,679);
2019/09/12 (US62/899,584)

(51) Cl.Int./Int.Cl. *C07K 14/315* (2006.01),
A61K 48/00 (2006.01), *C12N 15/63* (2006.01),
C12N 15/90 (2006.01), *C12N 9/22* (2006.01)
(71) Demandeur/Applicant:
ARIZONA BOARD OF REGENTS ON BEHALF OF
ARIZONA STATE UNIVERSITY, US
(72) Inventeurs/Inventors:
KIANI, SAMIRA, US;
EBRAHIMKHANI, MO REZA, US;
MOGHADAM, FARZANEH, US
(74) Agent: TORYS LLP

(54) Titre : IMMUNOMODULATION SYNTHETIQUE AVEC UN SUPER-REPRESSEUR DE CRISPR IN VIVO
(54) Title: SYNTHETIC IMMUNOMODULATION WITH A CRISPR SUPER-REPRESSOR IN VIVO



(57) Abrégé/Abstract:

Provided herein are CRISPR-based synthetic repression systems as well as methods and compositions using the synthetic repression systems to treat septicemia, an adverse immune response in a subject and Waldenström macroglobulinemia.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau

(43) International Publication Date
14 May 2020 (14.05.2020)



(10) International Publication Number
WO 2020/097344 A1

(51) International Patent Classification:

C07K 14/315 (2006.01) C12N 15/90 (2006.01)
C12N 9/22 (2006.01) A61K 48/00 (2006.01)
C12N 15/63 (2006.01)

(21) International Application Number:

PCT/US2019/060285

(22) International Filing Date:

07 November 2019 (07.11.2019)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/757,679 08 November 2018 (08.11.2018) US
62/899,584 12 September 2019 (12.09.2019) US

(71) Applicant: **ARIZONA BOARD OF REGENTS ON BEHALF ARIZONA STATE UNIVERSITY** [US/US]; 1475 N. Scottsdale Road, Suite 200, Scottsdale, AZ 85257 (US).

(72) Inventors: **KIANI, Samira**; 10240 N. 77th Street, Scottsdale, AZ 85258 (US). **EBRAHIMKHANI, Mohammed, R.**; 10240 N. 77th Street, Scottsdale, AZ 85258 (US). **MOGHADAM, Farzaneh**; 834 S. Casitas Dr, Unit B, Tempe, AZ 85281 (US).


(74) Agent: **HELMICH, Kate, E.**; Quarles & Braddy LLP, 33 East Main Street, Suite 900, Madison, WI 53703 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

(54) Title: SYNTHETIC IMMUNOMODULATION WITH A CRISPR SUPER-REPRESSOR IN VIVO

(57) Abstract: Provided herein are CRISPR-based synthetic repression systems as well as methods and compositions using the synthetic repression systems to treat septicemia, an adverse immune response in a subject and Waldenstrom macroglobulinemia.



WO 2020/097344 A1

WO 2020/097344 A1 

Published:

- *with international search report (Art. 21(3))*
- *with sequence listing part of description (Rule 5.2(a))*

SYNTHETIC IMMUNOMODULATION WITH A CRISPR SUPER- REPRESSOR IN VIVO

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims priority to U.S. Provisional Application No. 62/757,679, filed November 8, 2018, and U.S. Provisional Application No. 62/899,584, filed September 12, 2019, each of which is incorporated herein in its entirety.

STATEMENT REGARDING FEDERALLY SPONSORED RESEARCH

[0002] This invention was made with government support under R01 EB024562 awarded by the National Institutes of Health and HR0011-16-23657 awarded by DARPA. The government has certain rights in the invention.

REFERENCE TO A SEQUENCE LISTING SUBMITTED VIA EFS-WEB

[0003] The content of the ASCII text file of the sequence listing named "112624_01135_ST25.txt" which is 48.4 kb in size was created on November 7, 2019 and electronically submitted via EFS-Web herewith the application is incorporated herein by reference in its entirety.

BACKGROUND

[0004] Compared to the common gene delivery methods, adeno-associated viral (AAV) vectors possess a broad potential for therapeutic gene delivery due to their remarkable capacity to target a variety of tissue types at high efficiency. AAV vectors are currently the prevalent type of DNA viruses used in clinic. Long-term and stable expression of the transgene achieved with AAV is necessary for the treatment of chronic diseases. However, it has been shown that AAV vectors or the delivered transgenes evoke an adaptive immune response, which leads to the production of antibodies and cytotoxic T cell response against cells expressing viral capsids or the transgenes. Subsequently, this adaptive host immune response compromises the effective expression of the transgene and have posed a major challenge in AAV-mediated gene therapy *in vivo*.

[0005] In parallel, the Clustered, Regulatory Interspaced, Short Palindromic Repeats (CRISPR)-Cas9 system is providing unprecedented opportunities for gene therapies through facilitating gene editing and gene modulation. Although extremely promising, Cas9 protein from *Streptococcus Pyogenes* (Sp-Cas9), the most common form of CRISPR studied so far, faces a number of challenges for clinical translation including potential immune response and genomic specificity. Recent studies on human samples hint to pre-existing humoral and cellular immune response against Sp-Cas9 in a subset of patients tested, raising the notion that Cas9 protein can also elicit cellular and humoral immune response in new patients. Moreover, it has been demonstrated that temporal limitation of Cas9 expression inside the cells is beneficial towards reducing the off-target activity of CRISPR in the genome. All together, these pieces of evidence suggest that strategies to modulate adverse consequences of CRISPR and its delivery vehicle, including AAV virus, on demand will be of great importance for the safety of clinical translation.

SUMMARY OF THE INVENTION

[0006] In a first aspect, provided herein is a synthetic repression system for repressing myeloid differentiation primary response 88 (MyD88) expression *in vivo* comprising a single amplicon comprising: (a) a guide RNA (gRNA) comprising (i) a guide sequence complementary to a portion of MyD88 and (ii) an aptamer target site specific for an RNA binding protein; (b) a nucleotide sequence encoding the aptamer RNA binding protein fused to one or more repression domain; and (c) a nucleotide sequence encoding a multifunctional Cas nuclease; wherein the single amplicon is packaged in a vector for DNA-based viral delivery. In some embodiments the gRNA guide sequence is 14 nucleotides in length. In some embodiments, the gRNA guide sequence targets within 100 base pairs (bp) upstream of TATA box region of MyD88. In some embodiments, gRNA guide sequence is selected from the group consisting of SEQ ID NOs:1-4. In some embodiments, the aptamer target site is an MS2 aptamer target site and the RNA binding protein is bacteriophage MS2 coat protein. In some embodiments, the gRNA comprises the MS2 aptamer target site and the guide sequence is 14 nucleotides in length.

[0007] In some embodiments, the repression domain is selected from the group consisting of a Kruppel associated box (KRAB) domain, methyl-CpG (mCpG) binding domain 2 (meCP2), Switch independent 3 transcription regulator family member A (SIN3A), histone deacetylase HDT1 (HDT1), n-terminal truncation of methyl-CpG-binding domain containing protein 2

(MBD2B), nuclear inhibitor of protein phosphatase-1 (NIPP1), and heterochromatin protein 1 (HP1A). In some embodiments, the nucleotide of (b) encodes MS2 fused to at least two repression domains. In some embodiments, the nucleotide sequence encoding MS2 fused to a repression domain comprises a sequence selected from the group consisting of SEQ ID NOs:55, 59, 61, and 63.

[0008] In some embodiments, the system comprises a nucleotide sequence encoding a multifunctional Cas nuclease fused to the aptamer RNA binding protein and a repression domain. In some embodiments, the Cas nuclease is an *S. aureus* Cas9 nuclease or an *S. pyrogenes* Cas9 nuclease. In some embodiments, the vector is an AAV2/1 delivery vector.

[0009] In a second aspect, provided herein is a pharmaceutical composition comprising the synthetic repression system described herein and a pharmaceutically acceptable delivery vehicle.

[0010] In a third aspect, provided herein is a method of treating a subject with septicemia comprising administering to the subject with septicemia a therapeutically effective amount of a composition comprising the synthetic repression system described herein. In some embodiments, levels of immune related markers Icam-1, Tnfa, Ncf, Il6, Ifn- α , Ifn- β , Ifn- γ , and Stat4 are reduced in the subject following administration. In some embodiments, the method additionally comprises the step of monitoring the level of at least one immune related marker selected from the group consisting of Icam-1, Tnfa, Ncf, Il6, Ifn- α , Ifn- β , Ifn- γ , and Stat4 and administering to the subject additional doses of the synthetic repression system composition until the level of the at least one immune related marker is reduced. In some embodiments, serum lactate is reduced in the subject following administration.

[0011] In a fourth aspect, provided herein is a method of preventing an adverse immune response in a subject comprising administering to the subject a therapeutically effective amount of a composition comprising the synthetic repression system described herein prior to or concurrently with administration of an AAV vector based gene therapy to the subject, wherein the level of an immune related marker selected from the group consisting of Icam-1, Tnfa, Ncf, Il6, Ifn- α , Ifn- β , Ifn- γ , and Stat4 is reduced in the subject relative to subject who received the AAV vector based gene therapy but did not receive the therapeutic synthetic repression system composition. In some embodiments, the subject previously received at least one AAV vector based gene therapy. In some embodiments, the method additionally comprises the step of monitoring the level of at least one immune related marker selected from the group consisting of

Icam-1, Tnfa, Ncf, Il6, Ifn- α , Ifn- β , Ifn- γ , and Stat4 and administering to the subject additional doses of the synthetic repression system composition until the level of the at least one immune related marker is reduced.

[0012] In a fifth aspect, provided herein is a method of treating Waldenström macroglobulinemia in a subject in need thereof comprising administering to a subject with Waldenström macroglobulinemia a therapeutically effective amount of a composition comprising the synthetic repression system described herein. In some embodiments, the method additionally comprises administering to the subject a composition comprising a gRNA configured to cleave MyD88 at L265 locus and a homology directed repair template encoding a portion of wild-type MyD88 at the L265 locus, whereby levels of endogenous L265P mutant MyD88 are reduced in the subject.

BRIEF DESCRIPTION OF DRAWINGS

[0013] FIGS. 1A-1H show CRISPR-based targeted *Myd88* repression *in vitro* and *in vivo* demonstrates efficacy in modulating host response against AAV-based gene therapies. (FIG. 1A) Top: Schematic representation of the gRNA binding sites targeted to the promoter of *Myd88* locus; Bottom: Mouse Neuroblastoma (N2A) cell lines were transfected with MyD88 gRNA pairs or mock gRNA control together with Cas9 nuclease and either MS2-KRAB or MS2-HP1aKRAB cassettes. 72 hours post-transfection cells were treated with Lipopolysaccharides (*E. coli* LPS, serotype 0111:B4) at a concentration of 10 μ g/ml to induce *Myd88* expression. (FIG. 1B) qRT-PCR analysis of *Myd88* expression mRNA level post LPS treatment. Fold changes were quantified relative to the expression level of cells receiving non-targeting Mockg RNA (N = 4 independent transfections). (FIG. 1C) Schematic of experiments involving retro-orbital injection of AAV-MyD88 or Mock repressors to Cas9 nuclease transgenic mice. (FIG. 1D) qRT-PCR analysis of *Myd88* expression level in lung, blood, and bone marrow of Cas9 transgenic mice 3 weeks post retro-orbital injections of 1E+12 GC of AAV/Myd88 or AAV/Mock vectors carrying MS2- HP1a -KRAB or MS2-KRAB (N = 4-5 for injected groups and N = 2 for Not Injected group). Fold changes are relative to universal control (FIGS. 1E and 1F) Fold-change in the expression level of *Icam-1* (FIG. 1E), and *Tnfa* (FIG. 1F) mRNAs relative to the universal control. (N = 4-5 for injected groups and N = 2 for Not Injected group). Universal control is the level of the desired transcript in the blood sample collected from a not injected mouse, which did

not receive any AAV injection. **P <0.01 and *P <0.05 indicate statistical significance measured by Students' t-test. (FIG. 1G) Volcano plot showing significance versus expression of differentially expressed genes between bone marrow samples collected from mice treated with Myd88-Ms2-HP1aKRAB versus Myd88-MS2-KRAB. Points above the dotted line represent genes significantly (adj. p-value <0.05) up and down regulated. Highly downregulated genes in the presence of MS2-HP1aKRAB are a family of immunoglobulin heavy and light chains. (FIG. 1H) Analysis of anti-AAV1 IgG2A antibody measured by ELISA, reveals AAV1 specific antibody is markedly repressed upon Myd88 pre-treatment. **P <0.01 and *P <0.05 indicate statistical significance measured by Students' t-test.

[0014] FIGS. 2A-2H show CRISPR-based modulation of host inflammatory response confers protection following LPS-mediated septicemia. (FIG. 2A) Schematic of the experiments to assess the protective effect of CRISPR mediated MyD88 repression in septicemia. 1E+12 GC of AAV vectors were injected to Cas9 expressing mice via retro-orbital injection and approx. 3 weeks later they were treated i.p. with LPS (5mg/kg). At 6 hours post LPS injection, the mice were sacrificed. (FIGS. 2B-2D) Endogenous repression of inflammatory genes. qRT-PCR analysis of *in vivo* Myd88 (FIG. 2B), *Tnfa* (FIG. 2C), and *Icam-1* (FIG. 2D) expression relative to the universal control following LPS injection (N = 5-6 for injected groups and N = 2 for Not Injected group). (FIG. 2E) Measurement of a panel of inflammatory cytokines in plasma and lung using multiplex-ELISA assay; values are displayed in the heat maps as log base 10 of the measured concentration. (FIG. 2F) Circulating L-lactate in plasma samples collected from mice 6 hours post LPS injection (N=4). (FIG. 2G) qRT-PCR analysis of *in vivo* Myd88 expression in liver samples 6 hours post LPS injection. Fold change expression levels were quantified relative to the universal control (N = 4-6 for Injected groups and N = 2 for Not Injected group). (FIG. 2H) Plasma concentration of Cholesterol, HDL, LDL and ALT (N = 3-4). "Universal control "is the level of the transcript of interest in the blood sample collected from a Not Injected mouse, which did not receive any AAV injection. Note for 2B-D and 2C, data of "Not Injected" mouse are also reported in Fig. 1D-F. **P <0.01 and *P <0.05 indicate statistical significance measured by Students' t-test.

[0015] FIGS. 3A-3B show aptamer-mediated CRISPR repression *in vitro*. (FIG. 3A) Schematic of aptamer-mediated recruitment of repressor domains to CRISPR complex. (FIG. 3B) mRNA expression of targeted genes following aptamer-mediated recruitment of repressor

domains to CRISPR complex in HEK293FT cells. Fold changes were quantified relative to dCas9 only control group. N=3 independent transfections.

[0016] FIG. 4 shows *in vivo* analysis of AAV2/1 tropism towards different tissues. AAV2/1-GFP was delivered to C57BL/6 mice via retro-orbital injection. GFP expression was assessed in different tissues by qRT-PCR. Average fold change expression levels are indicated above each group and are quantified relative to not injected mice (N=3 for not injected group and N=4-6 for AAV-GFP group).

[0017] FIGS. 5A-5C show RNA-seq analyses of bone marrow samples collected from mice treated with AAV- Myd88- MS2-HP1aKrab versus AAV- Myd88-MS2-Krab. (FIG. 5A) Scatter plot comparing expression of genes (Fragments Per Kilobase of transcript per Million mapped reads FPKM) in two replicates of bone marrows from Myd88-MS2-HP1aKRAB versus Myd88-MS2-KRAB. *Myd88*, *Il1b*, *Icam1*, *Tnfa* and *Il6* are highlighted in red and the most down regulated genes in Myd88-MS2-HP1a-KRAB groups as compared to MyD88-MS2-KRAB are highlighted in Cyan. (FIG. 5B) GO enrichment Histogram comparing bone marrow samples collected from mice treated with AAV- Myd88- MS2-HP1aKrab versus AAV- Myd88-MS2-Krab. The top 20 significantly enriched terms in the GO enrichment analysis are displayed. * demonstrates padj value <0.05 and significance. Note that pathways such as defense response to bacteria, which are associated with Myd88 signaling are mostly down regulated when HP1aKRAB was used. (FIG. 5C) Reactome Enrichment bar graph displaying the top 20 enriched genes in the Reactome database comparing in the BM samples of Myd88- MS2-HP1aKRAB versus Myd88-MS2-KRAB. * demonstrate significances padj value of <0.05 and significance.

[0018] FIG. 6 shows analysis of a set of immune-related transcripts following LPS injury. qRT-PCR analysis of *Ncf*, *Il6*, *Ifny*, and *Il1β* mRNA expression in Lung, Blood, and Bone marrow quantified relative to the universal control following LPS injection (N = 5-6 for injected groups and N = 2 for Not Injected group). Universal control is the level of the desired transcript in a blood sample collected from a not injected Cas9 transgenic mouse. *P <0.05; significant

[0019] FIG. 7 shows an assessment of the level of a panel of immune related genes in lung and bone marrow following LPS injection. qRT-PCR analysis of *in vivo* *CD68* (Macrophage marker), *Infa*, *Infb*, *CD4*, *Cxcl1*, and *Stat4* relative to the universal control following LPS injection in lung and bone marrow. (N = 5-6 for injected groups and N = 2 for Not Injected

group). "Universal control" is the level of the transcript of interest in a blood sample collected from a not injected Cas9 transgenic mouse. *P <0.05; significant

[0020] FIGS. 8A-8C show an assessment of the repression efficiency of a set of gRNAs targeting a different region of *Myd88* locus. (FIG. 8A) Schematic representation of the gRNA binding sites targeted to the promoter of *Myd88* locus. (FIG. 8B) N2A cells were transfected with indicated gRNAs and other components of CRISPR machinery including MS2-HP1aKRAB cassette. Levels of *Myd88* repression were analyzed via qRT-PCR 3 days after transfections (N = 2). Fold changes are relative to the group that did not receive any gRNA (No Guide). (FIG. 8C) qRT-PCR analysis of *in vivo Myd88* expression 3 weeks post retro-orbital injection of AAV. Gene expression fold-change was quantified relative to the universal control (N = 4-6 for Injected groups and N = 2 for Not Injected group). Universal control is the level of the desired transcript in a blood sample collected from a not injected Cas9 transgenic mouse. Note for (FIG. 8C) AAV-Mock-MS2-HP1aKRAB and AAV-Myd88set1-MS2-HP1aKRAB data are also reported in Fig. 1D. *P <0.05; significant

[0021] FIG. 9 shows analysis of serum markers of tissue damage following LPS injury. Serum Blood Urea Nitrogen (BUN), a marker of injury to kidney, was increased in serum after LPS injury, and *Myd88* repression prevented such increase. LPS treatment and *Myd88* repression did not have any significant effect on serum Lipase level, which generally indicates injury to the pancreas. *P <0.05; significant

[0022] FIGS. 10A-10C show endogenous repression of *Cxcr4*. (FIG. 10A) Schematic representation of the gRNA binding sites targeted to the promoter of *Cxcr4* locus. (FIG. 10B) Raw 264.7 cells were transduced with AAV-CXCR4 along with AAV-Cas9. Expression levels of *Cxcr4* mRNA was analyzed using qRT-PCR 5 days post transduction. Fold changes were quantified relative to the expression level of cells receiving the same dosage of AAV-Mock. (N = 2). (FIG. 10C) qRT-PCR analysis of *in vivo Cxcr4* expression in lung samples 3 weeks post AAV injections to Cas9 transgenic mice. Fold change expression levels were quantified relative to mice receiving the same dosage of AAV-Mock. (N=4) *P <0.05; significant.

[0023] FIG. 11 shows the differences between MS2-HP1aKRAB and Cas9-HP1aKRAB. Data shows similar level of repression using Cas9 plus MS2 fused Hp1a-Krab repressor and Cas9 fused to HP1a-Krab.

[0024] FIG. 12 shows repression of Myd88 and downstream cytokines followed by CRISPR delivery to wild type mice.

[0025] FIG. 13 shows repression of Myd88 and downstream cytokines followed by CRISPR delivery to wild type mice injected with LPS.

[0026] FIG. 14 shows a comparison of the repression efficiency of Cas9 with 14nt versus 20nt gRNA guide sequences targeting different loci. Data shows similar level of repression using Cas9+MS2-Hp1a-Krab repressor in combination with 14 nt gRNA or 20ntgRNA.

[0027] FIG. 15 shows qRT-PCR analysis of *in vivo Myd88* expression 24 hours post CRISPR delivery. Mice received 2.5mg/kg LPS. Two hours post LPS injection mice received the CRISPR synthetic repression system using pepjet. qRT-PCR analysis of *Myd88* expression shows decreased level of *Myd88* in lung, liver, and bone marrow.

[0028] FIG. 16 shows IDEXX liver panel data. Mice received 2.5mg/kg LPS. Two hours post LPS injection mice received the CRISPR synthetic repression system using pepjet.

INCORPORATION BY REFERENCE

[0029] All publications, patents, and patent applications mentioned in this specification are herein incorporated by reference to the same extent as if each individual publication, patent, and patent application was specifically and individually indicated to be incorporated by reference.

DETAILED DESCRIPTION OF THE INVENTION

[0030] The present disclosure describes compositions and methods for repressing expression of myeloid differentiation primary response 88 (MyD88) *in vivo*. The use of the compositions and methods described herein is beneficial for treating or preventing the immune response associated with administration of adeno-associated viral vectors and CRISPR gene editing constructs. Further, the compositions and methods described herein are useful in treating or preventing the aberrant immune response associated with septicemia.

[0031] The CRISPR synthetic repressor systems provided herein are configured to inhibit transcription of MyD88. MyD88 is a key node in innate and adaptive immune responses, acting as an essential adaptor molecule for a number of signaling pathways, including Toll-like receptor (TLR) and Interleukin (IL)-1 family signal transduction. Developed using multifunctional Cas nucleases and gRNA aptamers, the synthetic repressor systems modulate endogenous MyD88

levels *in vivo* and contribute to immunomodulation of the innate and adaptive immune response of the target cell. For example, by introducing into the target cell a multifunctional Cas nuclease along with (i) a gRNA-MS2 aptamer binding target complex complementary to MyD88 and (ii) a repression domain linked to MS2, the synthetic repressor system reduces transcription of MyD88 and downregulates signaling pathways associated with MyD88 function and implicated in the cellular immune response against foreign substances, such as bacteria and viruses.

[0032] Accordingly, in some aspects provided herein is a synthetic repression system for modulating *in vivo* expression of MyD88. In certain embodiments, the system comprises (a) a guide RNA (gRNA) including (i) a guide sequence of 15 or less nucleotides (nt) in length that is complementary to at least a portion of the MyD88 gene and (ii) an aptamer target site specific for an RNA binding protein, (b) a nucleotide sequence encoding the aptamer RNA binding protein fused to a repression domain, and (c) a nucleotide sequence encoding a multifunctional Cas nuclease. In some embodiments, the gRNA nucleotide, the repression domain nucleotide, and the Cas nuclease nucleotide comprise a single amplicon. In certain embodiments, the single amplicon is packaged in a vector for DNA-based viral delivery.

[0033] In preferred embodiments, the CRISPR-based synthetic repression system is a single amplicon that can be packaged for *in vivo* delivery into a mammalian cell using a delivery vector such as an exosome, virus, viral particle, virus-like particle, or nanoparticle. In certain embodiments, provided herein is a synthetic regulatory system comprising, in a single amplicon, a multifunctional Cas nuclease, which is in some cases fused to a repression domain, and a gRNA comprising a guide sequence of 14 nucleotides in length, wherein the synthetic regulatory system modulates cleavage and/or transcription in a mammalian cell.

[0034] As used herein, a "guide RNA" (gRNA) refers to a nucleotide sequence including a guide sequence that is complementary to at least a portion of a target gene and a scaffold sequence providing secondary structure and a primary sequence (e.g., tracrRNA) to recruit the Cas nuclease to the target gene. In some embodiments, the gRNA includes secondary structural elements, such as, but not limited to, a hairpin loop, a tetraloop, a stemloop, and combinations thereof. Variations of the gRNA sequence and structure are known in the art. One of skill in the art will recognize the variability tolerance of the gRNA sequence applicable to Cas9 mediated repression that is suitable for the disclosed synthetic repression system. A gRNA target gene also comprises a Protospacer Adjacent Motif (PAM) located immediately downstream from the target site of the target gene.

Examples of PAM sequence are known (see, e.g., Shah et al., *RNA Biology* 10 (5): 891-899, 2013). In some embodiments, the PAM sequence is dependent upon the species of Cas nuclease used in the architecture.

[0035] In some embodiments, the guide sequence of the gRNA that is complementary to at least a portion of the target gene is a sequence of about 20 nucleotides (nt). In some embodiments, the guide sequence of the gRNA that is complementary to at least a portion of the target gene is a truncated guide sequence of 15 or less nucleotides (nt). In some embodiments, the gRNA includes a 15-nt guide sequence, a 14-nt guide sequence, or a 13-nt guide sequence. Without being bound by any particular mechanism, theory, or mode of action, the synthetic repression system described herein exploits the ablation of Cas nuclease activity upon binding of the nuclease to a gRNA with 14-nt guide sequence rather than a 20-nt guide sequence. By recruiting the Cas complex to the target site of the target gene but inhibiting the nuclease activity, the Cas complex will not cleave the target gene but will inhibit and repress expression of the target gene. Additional recruitment of repression domains to the stalled Cas complex will augment the repression of the target gene.

[0036] In certain embodiments, the gRNA is configured to target one or more endogenous genes. By way of example, endogenous genes targeted for repression can include, without limitation, growth factors, cytokines, genes involved in homology directed repair, genes involved in non-homologous end joining (NHEJ), metabolic enzymes, cell cycle progression enzymes, and genes involved in the pathways of wound healing and tissue repair. In a preferred embodiment, the endogenous gene is MyD88. In some embodiments, the gRNA is configured to target within 200 base pairs (bp) upstream or downstream of the transcription start site of MyD88. In some embodiments, the gRNA is configured to target within 100 bp upstream of the TATA box region in the MyD88 gene. In some embodiments, the guide sequence of the gRNA is selected from the group consisting of SEQ ID NOs: 1-4.

[0037] To improve gene repression effectiveness and scalability of the system, guide RNAs (gRNAs) are in some cases engineered to comprise a hairpin aptamer target site specific for an RNA binding protein. In some cases, the aptamer target site is appended to the tetraloop and stem loop of a gRNA. RNA recognition motifs are known in the art, such as, but not limited to, the MS2 binding motif, the COM binding motif, or the PP7 binding motif to which certain proteins, MS2 coat protein, COM, or PP7, respectively, bind. In some cases, the aptamer target site is capable of

binding to the dimerized MS2 bacteriophage coat proteins. PP7 is the RNA-binding coat protein of the bacteriophage Pseudomonas. Like MS2, it binds a specific RNA sequence and secondary structure. The PP7 RNA-recognition motif is distinct from that of MS2. Consequently, PP7 and MS2 can be multiplexed to mediate distinct effects at different genomic loci simultaneously.

[0038] In some embodiments, the gRNA is expressed under the control of a RNA Pol II promoter or an RNA Pol III promoter. Examples of pol III promoters include, but are not limited to, U6 and H1 promoters. Examples of pol II promoters include, but are not limited to, the retroviral Rous sarcoma virus (RSV) LTR promoter (optionally with the RSV enhancer), the cytomegalovirus (CMV) promoter (optionally with the CMV enhancer), the SV40 promoter, the dihydrofolate reductase promoter, the β -actin promoter, the phosphoglycerol kinase (PGK) promoter, and the EF1 α promoter. In some cases, CRISPR-responsive promoters are used. As used herein, the term "CRISPR-responsive promoter" encompasses eukaryotic promoters as well as synthetic gene regulatory devices and circuits for regulated gene expression. In some embodiments, a CRISPR-responsive promoter comprises a RNA Pol II promoter or an RNA Pol III promoter. The CRISPR-responsive promoter can be a CRISPR-repressible promoter (CRP) or a CRISPR-activatable promoter (CAP).

[0039] Standard Pol III promoters, such as U6 and H1, which have been used to express gRNAs and short hairpin RNAs (shRNA), do not allow spatial or temporal control of downstream genes. Spatial limitation of CRISPR gRNAs have several important benefits, particularly with regards to human therapeutics: 1) limiting gRNA expression to tissue/cell type of interest, which eliminates the concern that delivery tools distribute the CRISPR coding cassette systematically; 2) avoiding germline mutations by eliminating concern over CRISPR activity in germ cells; 3) differential gene editing/modulation in neighboring or cross-talking cells: gRNA regulation by distinct cell type-specific promoters enables scientists to modulate the expression of different set of genes in distinct cell types. Accordingly, in certain embodiments, the CRISPR-based genetic circuit is configured to drive expression of a guide RNA (gRNA) using a cell type-specific promoter. In some cases, a CRISPR-based genetic circuit is configured to comprise an endogenous or cell type-specific promoter in place of a synthetic promoter. For example, a synthetic RNA Pol II or Pol III promoter can be swapped with a cell type- or context-specific promoter and interfaced with intracellular signaling, enabling multistep sensing and modulation of cellular behavior. In some cases, a transcriptional repression cascade comprises

two, three, or four interconnected CRISPR transcriptional repression circuits (NAND logic gates).

[0040] A promoter, generally, is a region of nucleic acid that initiates transcription of a nucleic acid encoding a product. A promoter may be located upstream (e.g., 0 bp to -100 bp, -30 bp, -75 bp, or -90 bp) from the transcriptional start site of a nucleic acid encoding a product, or a transcription start site may be located within a promoter. A promoter may have a length of 100-1000 nucleotide base pairs, or 50-2000 nucleotide base pairs. In some embodiments, promoters have a length of at least 2 kilobases (e.g., 2-5 kb, 2-4 kb, or 2-3 kb).

[0041] In certain embodiments, gRNA expression from RNA pol II promoters can be modulated using Csy4 endoribonuclease-mediated cleavage. In some cases, multiple gRNAs are placed in tandem from a single coding region processed by Csy4. In some cases, high ON/OFF ratios are achieved by using a modified U6-driven 14-nt gRNA cassette, where 20-nt gRNA target sites are inserted within both the U6 promoter site and body of the gRNA. This structure forms a second generation kill switch which enables full destruction of the 14-nt gRNA cassette upon expression of 20-nt gRNAs *in vitro*, *in vivo*, and *ex vivo*. In some cases, such kill switches are expressed *in vivo* in, for example, in tissues that are frequent targets in gene therapy and are tolerogenic immune environments. Additional embodiments of Cas-mediated kill switches suitable for use in the synthetic repressor systems described herein are described in PCT Publication No. WO 2019/005856, which is incorporated herein by reference in its entirety.

[0042] In some cases, the RNA binding protein (e.g., MS2, COM, or PCP) specific for the aptamer target site incorporated in the gRNA is fused to a repression domain such as, for example, a Kruppel associated box (KRAB) domain. Other suitable repression domains are known in the art including, but not limited to, methyl-CpG (mCpG) binding domain 2 (meCP2), Switch independent 3 transcription regulator family member A (SIN3A), histone deacetylase HDT1 (HDT1), n-terminal truncation of methyl-CpG-binding domain containing protein 2 (MBD2B), nuclear inhibitor of protein phosphatase-1 (NIPPI1), and heterochromatin protein 1 (HP1A). In some embodiments, the RNA binding protein is fused to KRAB and at least one additional repression domain. In some embodiments, the RNA binding protein MS2 is fused to KRAB and at least one additional repression domain. In some embodiments, MS2 is fused to KRAB and HP1A. In some embodiments, MS2 is fused to KRAB and MECP2. In some embodiments, MS2 is fused to MECP2, KRAB, and MBD2B. In some embodiments, MS2 is

fused to MBD2B and HP1A. In some embodiments, the RNA binding protein - repression domain fusion is encoded from an engineered nucleic acid.

[0043] In some embodiments, the RNA binding protein and at least one repression domain are fused to a multifunctional Cas nuclease. In some embodiments, multifunctional Cas nuclease is encoded from an engineered nucleic acid. For example, in certain embodiments, transcriptional modifiers are fused to a multifunctional Cas nuclease to enable site-specific transcriptional modifications. Various strategies can be used to engineer such fusion molecules. In some cases, transcriptional modulators are directly fused to a multifunctional Cas nuclease. In other cases, the modulator is fused to another RNA binding protein such as MS2 bacteriophage coat protein in order to recruit the modulator to the Cas/gRNA/DNA complex.

[0044] In some cases, the multifunctional Cas nuclease is fused to a repression domain. In other cases, repression is achieved without the use of any repression domain but, rather, through Cas nuclease-mediated steric hindrance. The repression domain can comprise an RNA binding protein (e.g., MS2) fused to a repression domain such as, for example, a Kruppel associated box (KRAB) domain. Other suitable repression domains are described herein.

[0045] The components of synthetic repression systems described herein are preferably provided in a single amplicon. In some cases, however, the components may be in the form of two or more polynucleotide sequences. In such cases, a synthetic repression system can comprise introducing into a single cell three cassettes comprising components of a CRISPR-based synthetic repression system, where the first cassette comprises a nucleotide sequence encoding a polypeptide of a multifunctional Cas9 nuclease, a second cassette comprises the gRNA construct, and a third cassette encodes the RNA binding protein -repression domain fusion. In some embodiments, the synthetic repression system comprises introducing into a single cell two cassettes comprising components of a CRISPR-based synthetic repression system, where the first cassette comprises a nucleotide encoding a polypeptide of a multifunctional Cas9 nuclease fused to a repression domain and the second cassette comprises the gRNA construct. In some embodiments, one or more of the cassettes may be under the control of an inducible promoter. In some cases, it will be advantageous to fuse a multifunctional Cas9 nuclease to a reporter polypeptide or other polypeptide of interest (e.g., a therapeutic protein). The reporter polypeptide may be a fluorescent polypeptide such as near infrared fluorescent protein (iRFP) (to monitor Cas9 protein dynamics).

[0046] In some cases, an activator of the inducible promoter is provided by a cassette comprising the safety construct, such as those described in PCT Publication No. WO 2019/005856, which is incorporated herein by reference in its entirety. Activators can mediate or promote recruitment of polymerase machinery to the CRISPR-Cas complex. The activator can be a zinc-finger protein fused to an activation domain such as a VP16 transcription activation domain or VP64 transcription activation domain. In certain embodiments, orthogonally acting protein-binding RNA aptamers such as MS2 are used for aptamer-mediated recruitment of an activator to the CRISPR-Cas complex. For example, an MS2-VPR fusion protein can be used to aid CRISPR-CAS/14-nt gRNA-mediated gene activation by means of aptamer-mediated recruitment of an activator to the CRISPR-Cas complex. In the presence of an inducer, a safety 20-nt gRNA is expressed, resulting in destruction of the 14-nt gRNA cassette. The synthetic regulatory circuit can be an engineered polynucleotide.

[0047] As used herein, the terms "engineered nucleic acid" and "engineered polynucleotide" are used interchangeably and refer to a nucleic acid that has been designed and made using known *in vitro* techniques in the art. In some embodiments, an engineered polynucleotide, also referred to as a circuit herein, is a nucleic acid that is not isolated from the genome of an organism. In some embodiments, the engineered polynucleotide is introduced to a cell, plurality of cells, an organ or an organism to perform diverse functions (e.g., differentiation of cells, as sensors within cells, program a cell to act as a sensor, and delivery of selective cell-based therapies).

[0048] In some embodiments, the engineered polynucleotide comprises one or more promoters, such as an inducible promoter, constitutive promoter, or a tissue-specific or cell type-specific promoter. Inducible promoters allow regulation of gene expression and can be regulated by exogenously supplied compounds, environmental factors such as temperature, or the presence of a specific physiological state, e.g., acute phase, a particular differentiation state of the cell, or in replicating cells only. In some cases, the cell type specific promoter is a germ cell-specific promoter such as, for example, phosphoglycerate kinase 2 (Pgk2) promoter. Inducible promoters and inducible systems are known and available from a variety of commercial sources, including, without limitation, Invitrogen, Clontech, and Ariad. Many other systems have been described and can be readily selected by one of skill in the art. Examples of inducible promoters regulated by exogenously supplied promoters include the zinc-inducible sheep metallothioneine (MT) promoter, the dexamethasone (Dex) -inducible mouse mammary tumor virus (MMTV) promoter, the T7

polymerase promoter system [WO 98/10088]; the ecdysone insect promoter [No et al, Proc. Natl. Acad. Sci. USA, 93:3346- 3351 (1996)], the tetracycline-repressible system [Gossen et al, Proc. Natl. Acad. Sci. USA, 89:5547-5551 (1992)], the tetracycline-inducible system [Gossen et al, Science, 268: 1766- 1769 (1995), see also Harvey et al, Curr. Opin. Chem. Biol., 2:512-518 (1998)], the RU486- inducible system [Wang et al, Nat. Biotech., 15:239-243 (1997) and Wang et al, Gene Ther., 4:432-441 (1997)] and the rapamycin-inducible system [Magari et al, J. Clin. Invest., 100:2865-2872 (1997)]. Still other types of inducible promoters which may be useful in this context are those which are regulated by a specific physiological state, e.g., temperature, acute phase, a particular differentiation state of the cell, or in replicating cells only. Non-limiting examples of control elements include promoters, activators, repressor elements, insulators, silencers, response elements, introns, enhancers, transcriptional start sites, termination signals, linkers and poly(A) tails. Any combination of such control elements is contemplated herein (e.g., a promoter and an enhancer).

[0049] In some embodiments, the promoter is a lung specific promoter. Suitable lung specific promoters are known in the art and include, but are not limited to an albumin promoter and a human alpha-1 anti-trypsin promoter. In some embodiments, the promoter is a blood specific promoter. In some embodiments, the promoter is a bone marrow specific promoter. In some embodiments, the promoter is a macrophage specific promoter, such as CD11B. In some embodiments the promoter is a B cell specific promoter, such as CD19. In some embodiments, the promoter is a dendritic cell specific promoter, such as CD11C. In some embodiments, the promoter is a T cell specific promoter, such as CD4 and CD8.

[0050] CRISPR systems belong to different classes, with different repeat patterns, sets of genes, and species ranges. A CRISPR enzyme is typically a type I or III CRISPR enzyme. The CRISPR system is derived advantageously from a type II CRISPR system. The type II CRISPR enzyme may be any Cas enzyme. The terms “Cas” and “CRISPR-associated Cas” are used interchangeably herein. The Cas enzyme can be any naturally-occurring nuclease as well as any chimeras, mutants, homologs, or orthologs. In some embodiments, one or more elements of a CRISPR system is derived from a particular organism comprising an endogenous CRISPR system, such as *Streptococcus pyogenes* (SP) CRISPR systems or *Staphylococcus aureus* (SA) CRISPR systems. The CRISPR system is a type II CRISPR system and the Cas enzyme is Cas9 or a catalytically inactive Cas9 (dCas9). Other non-limiting examples of Cas proteins include

Cas1, Cas1B, Cas2, Cas3, Cas4, Cas5, Cas6, Cas7, Cas8, Cas9 (also known as Csn1 and Csx12), Cas10, Csy1, Csy2, Csy3, Cse1, Cse2, Csc1, Csc2, Csa5, Csn2, Csm2, Csm3, Csm4, Csm5, Csm6, Cmr1, Cmr3, Cmr4, Cmr5, Cmr6, Csb1, Csb2, Csb3, Csx17, Csx14, Csx10, Csx16, CsaX, Csx3, Csx1, Csx15, Csf1, Csf2, Csf3, Csf4, homologues thereof, or modified versions thereof. A comprehensive review of the Cas protein family is presented in Haft et al. (2005) *Computational Biology, PLoS Comput. Biol.* 1:e60. At least 41 CRISPR-associated (Cas) gene families have been described to date.

[0051] It will be understood that the CRISPR-Cas system as described herein is non-naturally occurring in a cell, i.e. engineered or exogenous to the cell. The CRISPR-Cas system as referred to herein has been introduced in a cell. Methods for introducing the CRISPR-Cas system in a cell are known in the art, and are further described herein elsewhere. The cell comprising the CRISPR-Cas system, or having the CRISPR-Cas system introduced, according to the invention comprises or is capable of expressing the individual components of the CRISPR-Cas system to establish a functional CRISPR complex, capable of modifying (such as cleaving) a target DNA sequence. Accordingly, as referred to herein, the cell comprising the CRISPR-Cas system can be a cell comprising the individual components of the CRISPR-Cas system to establish a functional CRISPR complex, capable of modifying (such as cleaving) a target DNA sequence. Alternatively, as referred to herein, and preferably, the cell comprising the CRISPR-Cas system can be a cell comprising one or more nucleic acid molecule encoding the individual components of the CRISPR-Cas system, which can be expressed in the cell to establish a functional CRISPR complex, capable of modifying (such as cleaving) a target DNA sequence.

[0052] In some embodiments, a synthetic CRISPR-based repression system as described herein may be introduced into a biological system (e.g., a virus, prokaryotic or eukaryotic cell, zygote, embryo, plant, or animal, e.g., non-human animal). A prokaryotic cell may be a bacterial cell. A eukaryotic cell may be, e.g., a fungal (e.g., yeast), invertebrate (e.g., insect, worm), plant, vertebrate (e.g., mammalian, avian) cell. A mammalian cell may be, e.g., a mouse, rat, non-human primate, or human cell. A cell may be of any type, tissue layer, tissue, or organ of origin. In some embodiments a cell may be, e.g., an immune system cell such as a lymphocyte or macrophage, a fibroblast, a muscle cell, a fat cell, an epithelial cell, or an endothelial cell. A cell may be a member of a cell line, which may be an immortalized mammalian cell line capable of proliferating indefinitely in culture.

[0053] To achieve endogenous repression of multiple genes, repression domains are included in multiple gRNAs. In some cases, the gRNAs further comprise one or more aptamers such as MS2, COM, or PP7 to which certain proteins (MS2 coat protein, COM, or PCP, respectively) bind. The repression domain is fused to the aptamer binding protein and, thus, will be recruited to the Cas9/gRNA complex to achieve repression of transcription of endogenous genes. PP7 is the RNA-binding coat protein of the bacteriophage *Pseudomonas*. Like MS2, it binds a specific RNA sequence and secondary structure. The PP7 RNA-recognition motif is distinct from that of MS2. Consequently, PP7 and MS2 can be multiplexed to mediate distinct effects at different genomic loci simultaneously.

[0054] In some cases, SMASh or another degradation/destabilization domain is fused to Cas9 nuclease or other protein fusion described herein. In general, destabilizing domains are small protein domains that are unstable and degraded in the absence of ligand, but whose stability is rescued by binding to a high-affinity cell-permeable ligand. Genetic fusion of a destabilizing domain to a protein of interest results in degradation of the entire fusion. Addition of a ligand for the destabilizing domain protects the fusion from degradation and, in this manner, adds ligand-dependent stability to a protein of interest.

[0055] In some cases, a viral or plasmid vector system is employed for delivery of a synthetic CRISPR-based repression system described herein. Preferably, the vector is a viral vector, such as a lenti- or baculo- or preferably adeno-viral/adeno-associated viral vectors, but other means of delivery are known (such as yeast systems, microvesicles, gene guns/means of attaching vectors to gold nanoparticles) and are provided. In some embodiments, one or more of the viral or plasmid vectors may be delivered via liposomes, nanoparticles, exosomes, microvesicles, or a gene-gun. In certain preferred embodiments, CRISPR-based repression system or components thereof (e.g., gRNAs) are packaged for delivery to a cell in one or more viral delivery vectors. Suitable viral delivery vectors include, without limitation, adeno-viral/adeno-associated viral (AAV) vectors, lentiviral vectors, and Herpes Simplex Virus 1 (HSV-1) vectors. For example, a synthetic CRISPR-based repression system can be introduced into one or more Herpes simplex amplicon vectors.

[0056] In some embodiments, the synthetic CRISPR-based repression system is introduced into one or more AAV vectors. In some embodiments, the AAV vector is a AAV2/1 vector. As used herein, "AAV2/1 vector" refers to a recombinant adeno-associated viral vector including

AAV2 inverted terminal repeats and AAV1 *Rep* and *Cap* genes. Other suitable AAV vectors are known in the art, such as, but not limited to AAV1.

[0057] Viral vectors and viral particles are commonly used viral delivery platforms for gene therapy. Therefore, for faster clinical translation, CRISPR-based genetic circuits described herein can be constructed using embedded kill switches and incorporated into Herpes simplex amplicon vectors. Preferably, a single carrier vector is used to achieve transfection of all synthetic genetic components to target cells. Without being bound to any particular theory or mode of action, it is believed that the ability to control CRISPR functionality, by adding only cleaving 20-nt gRNAs, provide an ideal safety kill switch mechanism due to the small DNA footprint of gRNA and limited payload capacity of viral particles. In some cases, a Herpes simplex viral (HSV) delivery system is used for delivery of engineered polynucleotides. In other cases, non-viral particles can be used for delivery. For example, controlled spatial transfection and/or destruction of CRISPR genetic circuits is achieved by assembling and packaging of all CRISPR-based genetic circuit elements in HSV-1 amplicon vectors having large payload capacity (150 kb) and capable of infecting multiple target cells.

[0058] In another aspect, provided herein is a co-virus strategy, where the method comprises introducing into a single cell two AAV vectors. The first vector carries (i) a nucleotide sequence encoding a fusion polypeptide of a Cas9 nuclease, and (ii) a gRNA configured for gene modulation as described herein. In some cases, the Cas9 nuclease is fused to a reporter polypeptide. The reporter polypeptide may be a fluorescent polypeptide such as near infrared fluorescent protein (iRFP) (to monitor Cas9 protein dynamics). The second vector is the “safety vector” and carries a 20-nt controllable gRNA as described in PCT Publication No. WO 2019/005856, which is incorporated herein by reference in its entirety. To ensure that expression of the Cas9-iRFP fusion polypeptide carried by the first virus occurs only in cells into which both AAV vectors are introduced, expression of the fusion polypeptide is under the control of an inducible promoter. In some cases, an activator of the inducible promoter is carried by the safety virus. Activators can mediate or promote recruitment of the Pol II machinery to the CRISPR-Cas complex. The activator can be a zinc-finger protein fused to an activation domain such as a VP16 transcription activation domain or VP64 transcription activation domain. In certain embodiments, orthogonally acting protein-binding RNA aptamers such as MS2 are used for aptamer-mediated recruitment of an activator to the CRISPR-Cas complex. For example, the

second virus (safety virus) can carry an MS2-VPR fusion protein that aids in CRISPR-CAS and 14-nt gRNA-mediated gene activation by means of aptamer-mediated recruitment of an activator to the CRISPR-Cas complex. In the presence of an inducer, the safety 20-nt gRNA is expressed, resulting in destruction of the 14-nt gRNA cassette.

[0059] Applications of the CRISPR-based genetic circuits described herein include, without limitation, *in vivo* CRISPR-based precision gene therapies for treating chronic and acute conditions in a variety of cell types; a platform of CRISPR cell classifiers can be applied to various cells, tissues, or organs for various therapeutic and/or preventative applications; and *in vivo* interrogation of endogenous genes using CRISPR activators and repressors, including CRISPR-mediated endogenous gene activation. By way of example, therapeutic applications include, without limitation, treating septicemia in subject, preventing or treating an immune response in a subject, and treating Waldenström macroglobulinemia.

[0060] In certain aspects, provided herein are methods for treating septicemia, methods for treating or preventing an immune response in a subject, and methods for treating Waldenström macroglobulinemia. The methods of treatment or prevention described herein include administration to a subject in need thereof a pharmaceutical composition comprising a therapeutically effective amount of the CRISPR-based synthetic repression system described herein or the polynucleotides and vectors encoding the CRISPR-based synthetic repression system described herein. As used herein, the term “pharmaceutical composition” refers to a chemical or biological composition suitable for administration to a mammal. Such compositions typically include the active agent and a pharmaceutically acceptable carrier. As used herein the term “pharmaceutically acceptable carrier” includes saline, solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like, compatible with pharmaceutical administration. Supplementary active compounds can also be incorporated into the compositions. Examples of compositions appropriate for such therapeutic applications include preparations for parenteral, subcutaneous, transdermal, intradermal, intramuscular, intracoronarial, intramyocardial, intraperitoneal, intravenous or intraarterial (*e.g.*, injectable), or intratracheal administration, such as sterile suspensions, emulsions, and aerosols. In some cases, pharmaceutical compositions appropriate for therapeutic applications may be in admixture with one or more pharmaceutically acceptable excipients, diluents, or carriers such as sterile water, physiological saline, glucose or the like. For example, compositions of the

CRISPR-based synthetic repression system described herein can be administered to a subject as a pharmaceutical composition comprising a carrier solution.

[0061] Formulations may be designed or intended for oral, rectal, nasal, systemic, topical or transmucosal (including buccal, sublingual, ocular, vaginal and rectal) and parenteral (including subcutaneous, intramuscular, intravenous, intraarterial, intradermal, intraperitoneal, intrathecal, intraocular and epidural) administration. In general, aqueous and non-aqueous liquid or cream formulations are delivered by a parenteral, oral or topical route. In other embodiments, the compositions may be present as an aqueous or a non-aqueous liquid formulation or a solid formulation suitable for administration by any route, *e.g.*, oral, topical, buccal, sublingual, parenteral, aerosol, a depot such as a subcutaneous depot or an intraperitoneal or intramuscular depot. In some cases, pharmaceutical compositions are lyophilized. In other cases, pharmaceutical compositions as provided herein contain auxiliary substances such as wetting or emulsifying agents, pH buffering agents, gelling or viscosity enhancing additives, preservatives, flavoring agents, colors, and the like, depending upon the route of administration and the preparation desired. The pharmaceutical compositions may be formulated according to conventional pharmaceutical practice (see, *e.g.*, Remington: *The Science and Practice of Pharmacy*, 20th edition, 2000, ed. A. R. Gennaro, Lippincott Williams & Wilkins, Philadelphia, and *Encyclopedia of Pharmaceutical Technology*, eds. J. Swarbrick and J. C. Boylan, 1988-1999, Marcel Dekker, New York). For intravenous administration, suitable carriers include physiological saline, bacteriostatic water, Cremophor EL[®] (BASF, Parsippany, N.J., USA) or phosphate buffered saline (PBS). In all cases, a composition for parenteral administration must be sterile and should be formulated for ease of injectability. The composition should be stable under the conditions of manufacture and storage, and must be shielded from contamination by microorganisms such as bacteria and fungi.

[0062] In some embodiments, solutions or suspensions used for parenteral, intradermal, or subcutaneous application can include the following components: a sterile diluent such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerin, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl parabens; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid (EDTA); buffers such as acetates, citrates or phosphates, and agents for the adjustment of tonicity, such as sodium chloride or dextrose. The pH can be adjusted with acids or bases, such

as hydrochloric acid or sodium hydroxide. The preparation can be enclosed in ampoules, disposable syringes or multiple-dose vials made of glass or plastic. For convenience of the patient or treating physician, the dosing formulation can be provided in a kit containing all necessary equipment (*e.g.*, vials of drug, vials of diluent, syringes and needles) for a course of treatment (*e.g.*, 7 days of treatment).

[0063] Sterile injectable solutions can be prepared by incorporating the active compound in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle, which contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, typical methods of preparation include vacuum drying and freeze drying, which can yield a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

[0064] The preferred route may vary with, for example, the subject's pathological condition or age or the subject's response to therapy or that is appropriate to the circumstances. The formulations can also be administered by two or more routes, where the delivery methods are essentially simultaneous or they may be essentially sequential with little or no temporal overlap in the times at which the composition is administered to the subject.

[0065] Suitable regimes for initial administration and further doses or for sequential administrations also are variable, may include an initial administration followed by subsequent administrations, but nonetheless, may be ascertained by the skilled artisan from this disclosure, the documents cited herein, and the knowledge in the art.

[0066] In some embodiments, compositions of the CRISPR-based synthetic repression system described herein are administered to a subject in need thereof using an infusion, topical application, surgical transplantation, or implantation. In an exemplary embodiment, administration is systemic. In such cases, compositions of the CRISPR-based synthetic repression system described herein can be provided to a subject in need thereof in a pharmaceutical composition adapted for intravenous administration to subjects. Typically, compositions for intravenous administration are solutions in sterile isotonic aqueous buffer. The use of such buffers and diluents is well known in the art. Where necessary, the composition may

also include a local anesthetic to ameliorate any pain at the site of the injection. Generally, the ingredients are supplied either separately or mixed together in unit dosage form, for example, as a cryopreserved concentrate in a hermetically sealed container such as an ampoule indicating the quantity of active agent. Where the composition is to be administered by infusion, it can be dispensed with an infusion bottle containing sterile pharmaceutical grade water or saline. Where the composition is administered by injection, an ampoule of sterile water for injection or saline can be provided so that the ingredients may be mixed prior to administration. In some cases, compositions of the CRISPR-based synthetic repression system described herein are lyophilized prior to administration.

[0067] In one embodiment, compositions of the CRISPR-based synthetic repression system described herein are administered intravenously. For example, an agent of the present technology may be administered *via* rapid intravenous bolus injection. In some embodiments, the agent of the present technology is administered as a constant-rate intravenous infusion.

[0068] The agent of the present technology may also be administered orally, topically, intranasally, intramuscularly, subcutaneously, or transdermally. In one embodiment, transdermal administration is by iontophoresis, in which the charged composition is delivered across the skin by an electric current.

[0069] Other routes of administration include intracranio-ventricular, intracerebroventricularly or intrathecally. Intracerebroventricularly refers to administration into the ventricular system of the brain. Intrathecally refers to administration into the space under the arachnoid membrane of the spinal cord. Thus, in some embodiments, intracerebroventricular or intrathecal administration is used for those diseases and conditions which affect the organs or tissues of the central nervous system.

[0070] For systemic, intracerebroventricular, intrathecal, topical, intranasal, subcutaneous, or transdermal administration, formulations of the agents of the present technology may utilize conventional diluents, carriers, or excipients *etc.*, such as those known in the art to deliver the agents of the present technology. For example, the formulations may comprise one or more of the following: a stabilizer, a surfactant, such as a nonionic surfactant, and optionally a salt and/or a buffering agent. The agents of the present technology may be delivered in the form of an aqueous solution, or in a lyophilized form.

[0071] Therapeutically effective amounts of compositions of the CRISPR-based synthetic repression system described herein are administered to a subject in need thereof. An effective dose or amount is an amount sufficient to effect a beneficial or desired clinical result. With regard to methods of the present invention, the effective dose or amount, which can be administered in one or more administrations, is the amount of compositions of the CRISPR-based synthetic repression system described herein sufficient to elicit a therapeutic effect in a subject to whom the agent is administered. The dosage ranges described herein are exemplary and are not intended to be limiting. Dosage, toxicity, and therapeutic efficacy of the agents of the present technology can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, *e.g.*, for determining the LD50 (the dose lethal to 50% of the population) and the ED50 (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio LD50/ED50.

[0072] The data obtained from the cell culture assays and animal studies can be used in formulating a range of dosage for use in humans. The dosage of such compounds lies preferably within a range of circulating concentrations that include the ED50 with little or no toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration utilized. For any agent of the present technology used in the methods described herein, the therapeutically effective dose can be estimated initially from cell culture assays. A dose can be formulated in animal models to achieve a circulating plasma concentration range that includes the IC₅₀ (*i.e.*, the concentration of the test compound which achieves a half-maximal inhibition of symptoms) as determined in cell culture. Such information can be used to more accurately determine useful doses in humans. Levels in plasma may be measured, for example, by high performance liquid chromatography.

[0073] Typically, an effective amount of the compositions of the CRISPR-based synthetic repression system described herein, sufficient for achieving a therapeutic or prophylactic effect, ranges from about 1×10^9 genome copies (GC)/kilogram (kg) body weight to about 1×10^{14} GC/kg body weight. In some embodiments, the dosage ranges will be from about 1×10^9 GC/kg body weight per day to about 1×10^{14} GC/kg body weight per day. For example dosages between about 1×10^9 GC/kg body weight to about 1×10^{14} GC/kg body weight can be administered every day, every two days or every three days or within the range of 1×10^9 GC/kg body weight to about

1×10^{14} GC/kg body weight every week, every two weeks or every three weeks. In one embodiment, a single dosage of the agent of the present technology ranges from 1×10^9 GC/kg body weight to about 1×10^{14} GC/kg body weight. An exemplary treatment regimen entails administration once per day or once a week. In therapeutic applications, a relatively high dosage at relatively short intervals is sometimes required until progression of the disease is reduced or terminated, or until the subject shows partial or complete amelioration of symptoms of disease. Thereafter, the patient can be administered a prophylactic regimen. The schedule of doses is optimized to maintain the therapeutic concentration at the target tissue, such as by single daily or weekly administration, but also including continuous administration (*e.g.*, parenteral infusion or transdermal application).

[0074] The skilled artisan will appreciate that certain factors may influence the dosage and timing required to effectively treat a subject, including but not limited to, the severity of the disease or disorder, previous treatments, the general health and/or age of the subject, and the presence of other diseases. Moreover, treatment of a subject with a therapeutically effective amount of the therapeutic compositions described herein can include a single treatment or a series of treatments.

[0075] In some aspects, provided herein are methods for treating septicemia in a subject. Methods of treating septicemia include administering to a subject with or at risk for septicemia a therapeutically effective amount of the compositions of the CRISPR-based synthetic repression system described herein and specific for repression of MyD88. Due to its role in innate and adaptive immunity, repression of MyD88 will result in reduction of the aberrant immune response associated with septicemia. A subject treated with the compositions described herein may have a reduction in the levels of immune related markers Icam-1, Tnf α , Ncf, and Il6, Ifn- α , Ifn- β , Ifn- γ , and Stat4 as compared to the level of the immune related marker prior to treatment. In some embodiments, the levels of immune related markers Icam-1, Tnf α , Ncf, and Il6, Ifn- α , Ifn- β , Ifn- γ , and Stat4 are compared to an internal or external control or reference standard. A subject treated with the compositions described herein may have a reduction in the levels of lung and blood serum cytokines as compared to the level of the cytokines prior to treatment. A subject treated with the compositions described herein may have a reduction in serum lactate levels as compared to the level measured prior to treatment.

[0076] In some aspects, provided herein are methods for treating or preventing an adverse immune response in a subject receiving an adeno-associated viral vector gene therapy or a CRISPR-based gene therapy. Methods of preventing an adverse immune response in a subject include administering to a subject a therapeutically effective amount of the compositions of the CRISPR-based synthetic repression system described herein and specific for repression of MyD88 prior to administration of an AAV vector therapy to the subject. The compositions described herein may be administered prior to or at the same time as the AAV vector therapy. In some embodiments, the subject has received or will receive multiple courses of an AAV vector therapy treatment and the compositions described herein are administered prior to, immediately after or concurrently with the AAV vector treatment. A subject treated with the compositions described herein may have a reduction in the levels of immune related markers Icam-1, Tnf α , Ncf, Il6, Ifn- α , Ifn- β , Ifn- γ , and Stat4 as compared to the level of the immune related markers observed in a subject who received the AAV vector therapy but not the compositions of the CRISPR-based synthetic repression system described herein and specific for repression of MyD88. A subject treated with the compositions described herein may have a reduction in the levels of lung and blood serum cytokines as compared to the level of the immune related markers observed in a subject who received the AAV vector therapy but not the compositions of the CRISPR-based synthetic repression system described herein and specific for repression of MyD88.

[0077] In some embodiments, the methods of treatment include monitoring the level of at least one immune related marker selected from the group consisting of Icam-1, Tnf α , Ncf, Il6, Ifn- α , Ifn- β , Ifn- γ , and Stat4 and re-administering the compositions of the CRISPR-based synthetic repression system described herein and specific for repression of MyD88 to the subject until the level of the at least one marker decreases or returns to the level prior to the subject receiving the AAV vector therapy or prior to the subject having septicemia.

[0078] In some aspects, provided herein are methods for the treatment of Waldanstrom macroglobulinemia. Waldanstrom macroglobulinemia is a type of non-Hodgkin lymphoma characterized by unchecked lymphocyte proliferation and macroglobulin production. Myd88 mutation L265P supports malignant growth through NF- κ B signaling and its inhibition is associated with decreased survival of cancer cells. Methods for treating Waldanstrom macroglobulinemia in a subject include administering to a subject in need thereof the compositions of the CRISPR-based synthetic repression system described herein and specific for

repression of MyD88. In some embodiments, the CRISPR-based synthetic repression system described herein represses endogenous, mutant MyD88 expression and is designed to use homology directed repair (HDR) to correct the L265P mutation. In some embodiments, the composition additionally includes gRNA specific for Cas nuclease cleavage near the L265 locus of MyD88 and includes an HDR template encoding the correct wild-type sequence at the L265 locus.

[0079] So that the compositions, methods, and systems provided herein may more readily be understood, certain terms are defined:

[0080] As used in this specification and the appended claims, the singular forms “a,” “an,” and “the” include plural references unless the context clearly dictates otherwise. Any reference to “or” herein is intended to encompass “and/or” unless otherwise stated.

[0081] The terms "comprising", "comprises" and "comprised of" as used herein are synonymous with "including", "includes" or "containing", "contains", and are inclusive or open-ended and do not exclude additional, non-recited members, elements, or method steps. The phraseology and terminology used herein is for the purpose of description and should not be regarded as limiting. The use of "including," "comprising," "having," "containing," "involving," and variations thereof, is meant to encompass the items listed thereafter and additional items. Use of ordinal terms such as "first," "second," "third," etc., in the claims to modify a claim element does not by itself connote any priority, precedence, or order of one claim element over another or the temporal order in which acts of a method are performed. Ordinal terms are used merely as labels to distinguish one claim element having a certain name from another element having a same name (but for use of the ordinal term), to distinguish the claim elements.

[0082] As used herein, the terms “synthetic” and “engineered” are used interchangeably and refer to the aspect of having been manipulated by the hand of man.

[0083] As used herein, “modifying” (“modify”) one or more target nucleic acid sequences refers to changing all or a portion of a (one or more) target nucleic acid sequence and includes the cleavage, introduction (insertion), replacement, and/or deletion (removal) of all or a portion of a target nucleic acid sequence. All or a portion of a target nucleic acid sequence can be completely or partially modified using the methods provided herein. For example, modifying a target nucleic acid sequence includes replacing all or a portion of a target nucleic acid sequence with one or more nucleotides (e.g., an exogenous nucleic acid sequence) or removing or deleting all or a portion (e.g., one or more nucleotides) of a target nucleic acid sequence. Modifying the one or more target

nucleic acid sequences also includes introducing or inserting one or more nucleotides (e.g., an exogenous sequence) into (within) one or more target nucleic acid sequences.

[0084] Unless otherwise defined, all terms used in this disclosure, including technical and scientific terms, have the meaning as commonly understood by one of ordinary skill in the art to which this invention belongs.

[0085] Although this description refers to certain aspects or embodiments, such aspects or embodiments are illustrative and non-exhaustive in nature. Having reviewed the present disclosure, persons of ordinary skill in the art will readily recognize and appreciate that numerous other possible variations or alternative configurations or aspects are possible and were contemplated within the scope of the present disclosure.

Example 1

[0086] The embodiments described here demonstrate use of a CRISPR-based synthetic repression system to repress endogenous MyD88 expression *in vitro* and *in vivo*.

[0087] The embodiment described here employs the notion of CRISPR Multifunctionality, reported previously (see Dahlman, J. E. *et al.* "Orthogonal gene knockout and activation with a catalytically active Cas9 nuclease," *Nature biotechnology* 33, 1159 (2015), and Kiani, S. *et al.* "Cas9 gRNA engineering for genome editing, activation and repression," *Nature methods* 12, 1051 (2015), each of which is incorporated herein by reference). The ability to switch between nuclease-dependent and -independent functions of a single Cas9 protein, offers exciting possibilities to combine gene editing with epigenetic manipulations. Inventors previously reported that truncation of guide RNA (gRNA) from 5' end, enables the application of a nuclease competent Cas9 protein for transcriptional modulation of genes. Liao *et al.* recently demonstrated the utility of this system *in vivo* for transcriptional activation of genes. (Liao, H.-K. *et al.* "In vivo target gene activation via CRISPR/Cas9-mediated trans-epigenetic modulation," *Cell* 171, 1495-1507. e1415 (2017)). However, the wider utility of this concept for transcriptional repression or simultaneous DNA cleavage/transcriptional modulation and in clinically relevant conditions remain to be examined. This disclosure describes technologies that exploit this notion in the context of safety and controllability of CRISPR-based *in vivo* therapies.

[0088] Myeloid differentiation primary response 88 (MyD88) is a key node in innate and adaptive immune responses, acting as an essential adaptor molecule for a number of signaling

pathways, including Toll-like receptor (TLR) and Interleukin (IL)-1 family signal transduction. Given the central role of MyD88 signaling in innate and adaptive immunity, the inventors set out to determine whether the inventors can achieve synthetic immunomodulation through regulation of endogenous *Myd88* levels *in vivo*.

[0089] The inventors previously reported enhanced CRISPR-based transcriptional repressors *in vitro* developed by direct fusion of a set of modulators to catalytically dead Cas9 protein (MeCP2, MBD2 or HP1a). The inventors first devised an experiment to determine the repression domains from our previously reported candidates that can lead to efficient transcriptional repression when fused to the MS2 coat protein (referred here to as MS2) and recruited to the CRISPR complex by gRNA aptamer binding (FIG. 3A). Quantitative real time polymerase chain reaction (qRT-PCR) analysis of a set of target genes in Human Embryonic Kidney 293 (HEK293FT) cells established that the MS2-HP1aKRAB [heterochromatin protein 1(HP1a)-Krüppel associated box (KRAB)] enabled efficient repression across the genes the inventors tested (FIG. 3B).

[0090] To translate these finding *in vivo*, the inventors set out to utilize nuclease competent *Streptococcus Pyogenes* (Sp) -Cas9 protein and truncated 14nt gRNAs for transcriptional repression of *Myd88* *in vivo*. The inventors first examined MS2-HP1aKRAB-mediated repression of endogenous mouse *Myd88* level *in vitro* and compared the efficiency with commonly used KRAB-based transcriptional repression. The inventors rationally designed two 14nt gRNAs that target within 100bp upstream of the TATA box region in the endogenous mouse *Myd88* locus or used a previously reported non-targeting mock gRNA as a control (Fig. 1A). qRT-PCR for *Myd88* demonstrates the *in vitro* functionality of the gRNAs and superiority of MS2-HP1aKRAB in repression of endogenous *Myd88* (Fig. 1B).

[0091] To test these repressors *in vivo*, the inventors pursued delivery through packaging gRNAs and MS2-repression cassettes within Adeno-Associated Virus (AAV) vectors. AAV serotypes have been used to deliver CRISPR *in vivo*, the most common serotype being AAV9, which has high affinity to numerous parenchymal cell populations. Here, the inventors employed AAV2/1, which is a recombinant AAV vector consisting of AAV2 inverted terminal repeats, and *AAV1 Rep* and *Cap* genes. AAV1 has been shown to be effective in transduction of components of the immune system and non-parenchymal cells such as dendritic cells, and endothelial cells. Moreover, AAV1 capsid can induce MyD88 signaling as part of the pathways

of immunity against AAVs in the host. Our assessment of AAV2/1 tissue affinity revealed the highest expression in blood, lung and bone marrow (FIG. 4). Subsequently, the inventors performed systemic delivery of AAV/Myd88 gRNA or control AAV/Mock gRNA with MS2-HP1aKRAB or MS2-KRAB cassettes to Cas9 nuclease transgenic mice (Fig. 1C). To test whether CRISPR mediated transcriptional repression of *Myd88* can create physiologically relevant effect on the immune response against AAV, the inventors specifically chose Cas9-expressing mice as they enable us to eliminate potential confounding effects associated with delivery of Cas9. Three weeks after injections, blood, lung, and bone marrow were harvested and *Myd88* expression was assessed by qRT-PCR (Fig. 1D). Compared to uninjected controls, AAV vector delivery led to an increase in *Myd88* across different tissues the inventors tested. Treatment with CRISPR to repress endogenous *Myd88* with HP1aKRAB led to a significant reduction in the level of *Myd88* in blood (~84%), lung (~75%) and bone marrow (~63%) as compared to the mock gRNA treated group, in agreement with high affinity of AAV2/1 for these tissues. Administration of the KRAB domain alone led to a less pronounced repression of *Myd88* in lung (~52%), blood (~59%) and bone marrow (~34%), with slightly higher variation among the animals tested (Fig. 1D and Table 1).

[0092] Table 1: Repression levels of *Myd88*, *Icam-1*, and *Tnfa* were assessed by qRT-PCR in lung, blood, and bone marrow 3 weeks post retro-orbital injection of AAV. Gene expression fold-change was quantified relative to the universal control. The repression levels are reported as percentage of fold change of AAV-Myd88 group divided by the fold change of AAV-Mock group for each gene. NC=No Change

	No LPS	Lung	Blood	Bone Marrow
		Mean of repression (%) \pm STD	Mean of repression (%) \pm STD	Mean of repression (%) \pm STD
Myd88-HP1a-KRAB	Myd88	75% \pm 0.389	84% \pm 0.124	63% \pm 0.227
	Icam-1	74% \pm 0.1	88% \pm 0.05	61% \pm 0.24
	Tnfa	63% \pm 0.19	58% \pm 0.09	57% \pm 0.18
Myd88-KRAB	Myd88	52% \pm 0.7	59% \pm 0.185	34% \pm 0.276
	Icam-1	NC \pm 0.35	70% \pm 0.29	41% \pm 0.21
	Tnfa	37% \pm 0.46	45% \pm 0.57	56% \pm 0.19

[0093] To assess the potency of repression in altering the downstream gene regulatory network, the inventors evaluated the levels of tumor necrosis factor- α (TNF- α) and intercellular adhesion molecule-1 (ICAM-1), two signaling elements directly modulated by the MyD88 signaling pathway. *Myd88* targeting with MS2-HP1a-KRAB led to a significant reduction in

Icam-1 and *TNF α* expression across multiple tissues, whereas targeting with MS2-KRAB did not lead to a similar consistent effect (FIGS. 1E-1F and Table 1).

[0094] To perform a systemic assessment of the repression efficiency of MS2- HP1aKRAB system as compared to MS2-KRAB, the inventors performed next generation RNA sequencing on the bone marrow of mice treated with these constructs. MS2-HP1aKRAB treated mice expressed lower MyD88 levels compared to MS2-KRAB treated ones (FIG. 5A), which is accompanied with changes in downstream signaling pathways such as *Il1 β* . Of note, GO Enrichment analysis revealed that Myd88-MS2-HP1aKRAB treated mice had significant downregulation of signaling pathways implicated in the immune and defense response against foreign organisms and bacteria, which are pathways associated with Myd88 function (FIG. 5B). Similarly, Reactome database revealed the TLR pathway as one of the highly significant pathways that are downregulated in the presence of MS2-HP1aKRAB (FIG. 5C). This evidence suggests that modulation of *Myd88* and its downstream immune pathways is most effective with the MS2- HP1a KRAB repressor *in vivo*.

[0095] Interestingly, volcano plotting of differentially expressed genes revealed the constant region of heavy chain of immunoglobulin G1 and G2(*Ighg1* and *Ighg2b*) and other immunoglobulin related heavy and light chain genes as most down-regulated with HP1a KRAB relative to KRAB (Fig. 1G).

[0096] Prior studies demonstrate that viral DNA stimulates TLR (i.e., TLR9), which in turn activates MyD88 and initiates downstream signaling events leading to adaptive immunity, and antibody production against AAVs. In light of prior evidence and the observed repression of the immunoglobulin pathway, the inventors asked whether there was a decrease in AAV-specific humoral response in Myd88 repressed groups. The inventors measured immunoglobulin G (IgG) response against AAV1 capsid. Compared to the un-injected controls, the inventors detected an 11-fold increase in plasma IgG2a levels against AAV1 in AAV/Mock treated animals. However, those who received AAV/Myd88 had significantly lower (60%) plasma IgG2a against AAV-1 (Fig. 1H). This finding presents an exciting opportunity to modulate humoral immunity against AAV, a significant challenge in current AAV-based gene therapies, with a tool inherently suited to perform concurrent gene editing (nuclease competent CRISPR).

[0097] Having identified a more potent CRISPR repressor for transcriptional modulation of *MyD88*, the inventors next asked whether this repression is sufficient to modulate downstream

host response and confer protection when there is an augmented systemic transcriptional response, such as in septicemia. Septicemia is a pressing medical issue due to the emergence of antibiotic-resistance and rising longevity of patients suffering from chronic diseases. Moreover, high mortality rate due to septicemia still remains a medical challenge following trauma in the battlefield, highlighting the need for novel prevention strategies.

[0098] The inventors pre-treated Cas9 mice with AAV/Myd88-Ms2-HP1aKRAB or AAV/Mock and three weeks later subjected them to systemic lipopolysaccharides (LPS) (from *Escherichia coli* 0127:B8) treatment. Six hours following LPS treatment, the inventors harvested lung, blood and bone marrow and assessed the transcript levels of *Myd88* and major inflammatory cytokines (Fig. 2A). The inventors observed significant repression of *Myd88* in lung (61%), blood (80%) and bone marrow (76%) compared to AAV/Mock-treated mice (Fig. 2B). In response to LPS, *Myd88* repression prevented upregulation of a wide range of inflammatory and immune-related markers that are directly or indirectly downstream of Myd88 signaling such as *Icam-1*, *Tnfa*, *Ncf*, and *Il6*, *Ifn- α* , *Ifn- β* , *Ifn- γ* , and *Stat4* (FIGS. 2C-2D, FIG. 6, and FIG. 7). Analysis of serum and lung cytokine levels using a quantitative ELISA-based chemiluminescent assay revealed lower level of cytokines in Myd88-repressed mice (Fig. 2E). Additionally, serum lactate level, a systemic marker associated with septicemia and tissue damage, was significantly lower when mice were pretreated with the AAV/Myd88 repression cassette before LPS exposure, indicating a reduced systemic injury (Fig. 2F).

[0099] Next, the inventors explored *Myd88* repression levels through targeting a different region of the *Myd88* promoter. The inventors constructed gRNAs targeting within 100bp upstream of the transcription start site (Set 2) and detected superior in vitro efficacy compared to the previous set of gRNAs (Set 1) (FIGS. 8A-8B). Strikingly, the new *Myd88* gRNAs led to superior repression in the liver and significant but less pronounced repression of *Myd88* in blood, lung and bone marrow when compared to previous set of gRNAs (FIG. 8C). This was also observed following LPS injury (Fig. 2G), demonstrating a potential organ and cell type-specific effect of gRNA target site within the endogenous promoters. Next, the inventors asked if this CRISPR-based *Myd88* repression is sufficient to induce a patho-physiologically relevant phenotype after LPS-induced damage. Analysis of serum markers of systemic inflammatory response and tissue damage of the liver showed that pretreatment with CRISPR-Set 2 attenuates the detrimental effects of LPS injection (Fig. 2H). In particular, high density lipoprotein (HDL)

has been shown to increase following LPS treatment to eliminate systemic LPS to protect tissues from damage and has been associated with MyD88 signaling. In accordance with this, the inventors found that LPS induced an elevation in HDL, low density lipoproteins (LDL), and cholesterol in mock-treated groups but not under MyD88 repression. In addition, MyD88 repression suppressed the elevation in Alanine Transaminase (ALT), a marker of hepatocyte damage, and Blood Urea Nitrogen (BUN), a marker of acute kidney injury, which further suggests that Myd88 repression could ameliorate liver and systemic tissue damage following septicemia (FIG. 2H and FIG. 9). Finally, the inventors showed the extensibility of MS2-HP1aKRAB super-repression for synthetic control of other immunologically relevant loci such as *CXCR4* *in vitro* and *in vivo* (FIGS. 10A-10C).

[00100] In summary, the inventors provide compositions, systems, and methods for synthetic control of immune response *in vivo* using a newly developed CRISPR-based transcriptional super-repressor against endogenous *Myd88*. The inventors show that this system is effective in modulating layers of downstream signaling and can create a visible protective phenotype *in vivo*. This notion is especially attractive in case of delivery using a less common AAV serotype (AAV2/1) known to target smaller cellular populations *in vivo* (e.g. non-parenchymal cells).

[00101] The inventors demonstrate that targeting the *Myd88* locus with AAV/CRISPR generated less IgG2a against AAV1 and modulated general immunoglobulin expression patterns, consistent with prior reports on failure of generation of antigen specific *IgG2a* response in *Myd88*^{-/-} animals. Additionally, AAV-1 was shown to be able to target dendritic cells, and *Myd88* signaling in these cells was important to mount adaptive immune response against AAVs. The ability to control *Myd88* levels using a CRISPR-based synthetic repressor is of significance in light of the common challenges involved with AAV-based clinical gene therapies, as this pathway has been demonstrated as a key node to induce humoral immunity against many AAV serotypes and not just AAV2/1 *in vivo*. Employment of a nuclease competent Cas9 and a truncated gRNA in this study opens up an opportunity for simultaneous application of CRISPR for targeted gene editing while modulating the immune response, which makes CRISPR-mediated gene repression superior to previously used systems such as shRNAs.

[00102] This strategy was also effective in modulating systemic inflammatory response against (LPS)-induced endotoxemia. CRISPR-mediated endogenous repression of *Myd88* prevented upregulation of a wide range of inflammatory markers and conferred a protective phenotype.

This notion is promising for the application of CRISPR-based transcriptional regulation as a readily programmable tool for modulating inflammatory conditions and protecting against septicemia. Finally, activating somatic mutation in Myd88 (L265P) is implicated in Waldenström macroglobulinemia, a type of non-Hodgkin lymphoma characterized by unchecked lymphocyte proliferation and macroglobulin production. Myd88 mutation supports malignant growth through NF- κ B signaling and its inhibition is associated with decreased survival of cancer cells. The ability to systemically repress Myd88 (and CXCR4) as shown in this study, can therefore present an exciting therapeutic opportunity for this group of patients.

Methods

[00103] *MS2 Fusion repressors* - To construct the MS2 fused transcriptional repressors, the specific domains of interest were amplified from vectors previously published in our group and subsequently cloned into pcDNA3-MS2-VP64 backbone (Addgene plasmid ID: 79371)¹⁰. The pcDNA3-MS2-VP64 vector was digested with NotI and AgeI to remove the VP64 domain and then the amplified repressors were cloned into this backbone via Gibson Assembly method.

[00104] *U6-gRNA-MS2 plasmids* - To generate these plasmids, 14bp or 20bp spacers of gRNAs were inserted into sgRNA-MS2 cloning backbone (Addgene plasmid ID: 61424) at BbsI site via golden gate reaction. All the gRNA sequences are listed below.

[00105] *AAV vectors* - Following cloning of the gRNAs into a U6-sgRNA-MS2 backbone, the U6-gRNA encoding region was amplified from this vector and inserted within gateway entry vectors using golden gate reaction. Using the same method, the repressor domain and a shefla promoter were cloned into gateway entry vectors. The designed safety switch sgRNA sequence was synthesized as gblocks (Integrated DNA Technologies) and inserted into a gateway entry vector (Addgene plasmid ID #62084) digested with HindIII and SphI. Further sub-cloning of all these components into AAV backbone via LR reaction (Invitrogen) made the AAV vectors. All the primers and gRNA sequences are listed below.

[00106] *AAV packaging and purification* - Constructed AAV vectors were digested by SmaI digest to test the integrity of ITR regions before virus production. Verified AAV vectors were used to generate AAV2/1-Myd88, AAV2/1-CXCR4, AAV2/DJ-TTN, AAV2/DJ-Cas9, and AAV2/1-Cas9 by PackGene® Biotech, LLC. The virus titers were quantified via Real-time SYBR Green PCR at 1.5E+13 GC/ml against standard curves using linearized parental AAV

vectors.

[00107] AVV2/1-Myd88 (SEQ ID NO:64)

CCTGCAGGCAGCTGCGCGCTCGCTCGCTCACTGAGGCCGCCCGGGCGTCGGGGCGACCTTTGGTCGCCC
GGCCTCAGTGAGCGAGCGAGCGCGCAGAGAGGGAGTGGCCAACTCCATCACTAGGGGTTCTGCGGC
CGCACGCGTTGATATCAACTTTGTATAGAAAAGTTTTCCCATGATTCCTTCATATTTGCATATACGATA
CAAGGCTGTTAGAGAGATAAATTGGAATTAATTTGACTGTAAACACAAAGATATTAGTACAAAATACGT
GACGTAGAAAAGTAATAATTTCTTGGGTAGTTTGCAGTTTTAAAATTATGTTTTAAAATGGACTATCATA
TGCTTACCGTAACTTGAAAGTATTTTCGATTTCTTGGCTTTATATATCTTGTGGAAAGGACGAAACACCT
GCGGGTCTGCACTGTTTTAGAGCTAGGCCAACATGAGGATCACCCATGTCTGCAGGGCCTAGCAAGT
TAAAATAAGGCTAGTCCGTTATCAACTTGGCCAACATGAGGATCACCCATGTCTGCAGGGCCAAGTGG
CACCGAGTCGGTGCTTTTTTTGAATTCTGATGCGGTATTTTCTCCTTACGCATCTGTGCGGTATTTTACA
CCGCATACGTCAAAGCAACCATAGTACGCGCCCTGTATTTCCCATGATTCCTTCATATTTGCATATACG
ATACAAGGCTGTTAGAGAGATAATTGGAATTAATTTGACTGTAAACACAAAGATATTAGTACAAAATA
CGTGACGTAGAAAAGTAATAATTTCTTGGGTAGTTTGCAGTTTTAAAATTATGTTTTAAAATGGACTATC
ATATGCTTACCGTAACTTGAAAGTATTTTCGATTTCTTGGCTTTATATATCTTGTGGAAAGGACGAAACA
CCGAAGCTTCGGGAGGGTTTTAGAGCTAGGCCAACATGAGGATCACCCATGTCTGCAGGGCCTAGCAA
GTTAAAATAAGGCTAGTCCGTTATCAACTTGGCCAACATGAGGATCACCCATGTCTGCAGGGCCAAGT
GGCACCGAGTCGGTGCTTTTTTTGAATTCTGATGCGGTATTTTCTCCTTACGCATCTGTGCGGTATTTCA
CACCGCATACGTCAAAGCAACCATAGTACGCGCCCTGTAGCGGACCCAAGTTTGTACAAAAAAGCAG
GCTAGGCTCCGGTGCCCGTCAGTGGGCAGAGCGCACATCGCCCACAGTCCCCGAGAAGTTGGGGGGA
GGGGTCCGCAATTGAACCGGTGCTAGAGAAGGTGGCGCGGGGTAACCTGGGAAAGTGATGTGCGTGT
ACTGGCTCCGCTTTTTTCCCGAGGGTGGGGGAGAACCGTATATAAGTGCAGTAGTCGCCGTGAACGTT
CTTTTTCGCAACGGGTTTGGCCGACAGCTTAAGCTTGGTACCGAGCTCGGATCCACTAGTAA
CGGCCCGCAGTGTGCTGGAATTCTGCAGATATCCATCACACTGGCGGCCGCTCGAGCATGCATCTAGA
GGACAACCTTTGTATACAAAAGTTGAGGCTGGATCCCGGTACCCTCGACACCTGCTGACAGGTCCACCA
TGGGACCTAAGAAAAAGAGGAAGGTGGCGGCCGCTGACTACAAGGATGACGACGATAAATCTAGAAT
GGCTTCTAACTTTACTCAGTTTCGTTCTCGTCGACAATGGCGGAACCTGGCGACGTGACTGTGCCCCAAG
CAACTTCGCTAACGGGATCGTGAATGGATCAGCTCTAAGCTCGGTTTACAGGCTTACAAAAGTAACCT
GTAGGTTTCGTCAGAGCTCTGCGCAGATAACGCAATAACCCATCAAAGTTCGAGGTTGCCTAAAGGCCCC
TGGCGTTTCGTAATAATATGGAACCTAACCTTCCAATTTTCGCCACGAATTCGACTGCGAGCTTATT
GTTAAGGCAATGCAAAGTCTCCTAAAAGATGGAAACCCGATTCCCTCAGCAATCGCAGCAAACCTCCGG
CATCTACGAGGCCAGCCCCAAGAAGAAGAGGAAGGTGAGTGGTGGAGGAAGTGGCGGGTCAAGGGTC
GAGCCCCAAGAAAAAACGGAAGTGGAGGCATCAATGAAGGAGGGAGAAAAACAACAAACCCAGGGA
GAAAAGTGAAGGGAATAAGAGAAAAAGCTCCTTCTCTAACAGTGCAGACGATATCAAGTCCAAGAAA
AAGCGGGAGCAGTCTAATGACATTGCTAGGGGCTTCGAGAGAGGACTGGAGCCAGAAAAAATCATTG
GGGCAACCGACAGCTGCGGCGATCTGATGTTTCTCATGAAATGGAAGGACACAGATGAGGCCGACCT
GGTGTCTGCCAAAGAAGCTAACGTGAAGTGTCCCCAGATCGTCATTGCTTTTTACGAGGAAAGGCTCA
CCTGGCACGCATATCCTGAGGATGCCGAAAACAAGGAGAAGGAATCAGCTAAGAGCTCGGGAGGTGG
TTCGGGTGGCTCTGGATCAATGGACGCGAAATCACTTACGGCATGGTCGAGAACACTGGTTACGTTCA
AGGACGTGTTTGTGGACTTTACACGTGAGGAGTGGAAATTGCTGGATACTGCGCAACAAATTGTGTAT
CGAAATGTCATGCTTGAGAATTACAAGAACCTCGTCAGTCTCGGATACCAGTTGACGAAACCGGATGT
GATCCTTAGGCTCGAAAAGGGGGAAGAACCTTGGCTGGTATAGAAGAATTGATCCCGCTTCAGTGCA
GGTGAATTCTACCACCAGCTTAAGCTTGGTACCGAGCTCGGATCCACTAGTAAACGGCCCGCAGTGTG
CTGGAATTCTGCAGATATCCATCACACTGGCGGCCGCTCGAGCATGCATCTAGAGGTACCCAGCTTTCT
TGTAACAAAGTGGTAAGCTTGCCTCGAGCAGCGTGTCTCGAGAGATCTACGGGTGGCATCCCTGTGACC
CCTCCCAGTGCCTCTCCTGGCCCTGGAAGTTGCCACTCCAGTGCCCAACAGCCTTGTCTAATAAAAAT
TAAGTTGCATCATTTTGTCTGACTAGGTGTCTTCTATAATATTATGGGGTGGAGGGGGTGGTATGGA
GCAAGGGGCAAGTTGGGAAGACAACCTGTAGGGCCTGCGGGGTCTATTGGGAACCAAGCTGGAGTGC
AGTGGACAATCTTGGCTCACTGCAATCTCCGCCTCTGGGTTCAAGCGATTCTCCTGCCTCAGCCTCC
CGAGTTGTTGGGATTCCAGGCATGCATGACCAGGCTCAGCTAATTTTTGTTTTTTTGGTAGAGACGGGG
TTTACCATATTGGCCAGGCTGGTCTCAACTCCTAATCTCAGGTGATCTACCCACCTTGGCCTCCCAA
ATTGCTGGGATTACAGGCGTGAACCACTGCTCCCTTCCCTGTCTTCTGATTTTGTAGGTAACCAACGTG
CGGACCGAGCGGCCGAGGAACCCCTAGTGATGGAGTTGGCCACTCCCTCTCTGCGCGTCTGCTCGCT
CACTGAGGCCGGGCGACCAAAGGTGCGCCGACGCCCGGGCTTTGCCCGGGCGGCTCAGTGAGCGAG

CGAGCGCGCAGCTGCCTGCAGGGGCGCCTGATGCGGTATTTTCTCCTTACGCATCTGTGCGGTATTTCA
 CACCGCATAACGTCAAAGCAACCATAGTACGCGCCCTGTAGCGGCGCATTAAAGCGCGGGCGGGTGTGGTG
 GTTACGCGCAGCGTGACCGCTACACTTGCCAGCGCCCTAGCGCCCCGCTCCTTTTCGCTTCTTCCCTTCT
 TTCTCGCCACGTTTCGCCGGCTTTCCCCGTCAAGCTCTAAATCGGGGGCTCCCTTTAGGGTTCCGATTTA
 GTGCTTTACGGCACCTCGACCCCAAAAACTTGATTTGGGTGATGGTTCACGTAGTGGGCCATCGCCCT
 GATAGACGGTTTTTTCGCCCTTTGACGTTGGAGTCCACGTTCTTTAATAGTGGACTCTTGTTCAAACTG
 GAACAACACTCAACCCTATCTCGGGCTATTCTTTTGATTTATAAGGGATTTTGCCGATTTTCGGCCTATT
 GGTTAAAAAATGAGCTGATTTAACAAAAATTTAACCGGAATTTAACAAAAATTTAACGTTTACAATT
 TTATGGTGCACCTCTCAGTACAATCTGCTCTGATGCCGCATAGTTAAGCCAGCCCCGACACCCGCCAAC
 ACCCGCTGACGCGCCCTGACGGGCTTGTCTGCTCCCGGCATCCGCTTACAGACAAGCTGTGACCGTCTC
 CGGGAGCTGCATGTGTCAGAGGTTTTACCGTCAACCGAAACGCGCGAGACGAAAGGGCCTCGTGA
 TACGCCTATTTTTATAGGTTAATGTCATGATAATAATGGTTTCTTAGACGTCAGGTGGCACTTTTCGGG
 GAAATGTGCGCGGAACCCCTATTTGTTTATTTTTCTAAATACATTCAAATATGTATCCGCTCATGAGAC
 AATAACCCTGATAAATGCTTCAATAATTTGAAAAAGGAAGAGTATGAGTATTC AACATTTCCGTGTC
 GCCCTTATCCCTTTTTTTCGGCATTTTGCCTTCTGTTTTTGCTCACCCAGAAACGCTGGTGAAAGTAA
 AAGATGCTGAAGATCAGTTGGGTGCACGAGTGGGTACATCGAACTGGATCTCAACAGCGGTAAGTAA
 CTTGAGAGTTTTTCGCCGAAGAACGTTTCCAATGATGAGCACTTTTAAAGTTCTGCTATGTGGCGCG
 GTATTATCCCGTATTGACGCCGGGCAAGGCAACTCGGTCGCCGCATACACTATTCTCAGAATGACTT
 GGTTGAGTACTACCAGTCACAGAAAAGCATCTTACGGATGGCATGACAGTAAGAGAATTATGCAGTG
 CTGCCATAACCATGAGTGATAACACTGCGGCCAACTTACTTCTGACAACGATCGGAGGACCGAAGGAG
 CTAACCGCTTTTTTGCACAACATGGGGGATCATGTAACCTCGCCTTGATCGTTGGGAACCGGAGCTGAA
 TGAAGCCATAACAAACGACGAGCGTGACACCACGATGCCTGTAGCAATGGCAACAACGTTGCGCAAA
 CTATTAACCTGGCGAACTACTTACTCTAGCTTCCCGCAACAATTAATAGACTGGATGGAGGCGGATAA
 AGTTGCAGGACCCTTCTGCGCTCGGCCCTTCCGGCTGGCTGGTTTATTGCTGATAAATCTGGAGCCGG
 TGAGCGTGGGTCTCGCGGTATCATTGCAGCACTGGGGCCAGATGGTAAGCCCTCCCGTATCGTAGTTA
 TCTACACGACGGGGAGTCAGGCAACTATGGATGAACGAAATAGACAGATCGCTGAGATAGGTGCCTC
 ACTGATTAAGCATTGGTAACTGTCAGACCAAGTTTACTCATATATACTTTAGATTGATTTAAAACCTCA
 TTTTTAATTTAAAAGGATCTAGGTGAAGATCCTTTTTGATAATCTCATGACCAAAATCCCTTAAACGTGA
 GTTTTCGTTCCACTGAGCGTCAGACCCCGTAGAAAAGATCAAAGGATCTTCTTGAGATCCTTTTTTTCT
 GCGCGTAATCTGCTGCTTGCAAACAAAAAACCACCGCTACCAGCGGTGGTTGTTTTCGGGATCAAG
 AGCTACCAACTCTTTTTCCGAAGGTAACCTGGCTTACGACAGAGCGCAGATACCAAATACTGTCCTTCTAG
 TGTAGCCGTAGTTAGGCCACCACTTCAAGAACTCTGTAGCACCCGCTACATACCTCGCTCTGCTAATCC
 TGTTACCAGTGGCTGCTGCCAGTGGCGATAAGTCGTGTCTTACCGGGTTGGACTCAAGACGATAGTTA
 CCGGATAAGGCGCAGCGGTCCGGCTGAACGGGGGGTTCGTGCACACAGCCCAGCTTGGAGCGAACGA
 CCTACACCGAACTGAGATACCTACAGCGTGAGCTATGAGAAAGCGCCACGCTTCCCGAAGGGAGAAA
 GGCGGACAGGTATCCGGTAAGCGGCAGGGTCGGAACAGGAGAGCGCACGAGGGAGCTTCCAGGGGG
 AAACGCCTGGTATCTTTATAGTCTGTCCGGTTTTCCACCTCTGACTTGAGCGTCGATTTTTGTGATG
 CTCGTCAGGGGGCGGAGCCTATGGA AAAACGCCAGCAACGCGGCCTTTTACGGTCTGTCCTTTT
 GCTGGCCTTTTGCTCACATGT

[00108] AAV2/1-CXCR4 (SEQ ID NO:65)

CCTGCAGGCAGCTGCGCGCTCGCTCGCTCACTGAGGCCCGCCGGGCGTCGGGCGACCTTTGGTTCGCC
 GGCCTCAGTGAGCGAGCGAGCGCGCAGAGAGGGAGTGGCCAACTCCATCACTAGGGGTTCTGCGGC
 CGCACGCGTTGATCAACTTTGTATAGAAAAGTTTTCCCATGATTCCTTCATATTTGCATATACGATA
 CAAGGCTGTTAGAGAGATAAATTGGAATTAATTTGACTGTAAACACAAAGATATTAGTACAAAATACGT
 GACGTAGAAAAGTAATAATTTCTTGGGTAGTTTGCAGTTTTAAAATTATGTTTTAAAATGGACTATCATA
 TGCTTACCGTAACTTGAAGTATTTTCGATTTCTTGGCTTTATATATCTTGTGGAAGGACGAAACACCA
 TAAAAGTCCGGTTGTTTTAGAGCTAGGCCAACATGAGGATCACCCATGTCTGCAGGGCCTAGCAAGTT
 AAAATAAGGCTAGTCCGTTATCAACTTGGCCAACATGAGGATCACCCATGTCTGCAGGGCCAAGTGGC
 ACCGAGTCCGGTGTTTTTTGAATTCTGATGCGGTATTTTCTCCTTACGCATCTGTGCGGTATTTACAC
 CGCATAACGTCAAAGCAACCATAGTACGCGCCCTGATTTCCCATGATTCCTTCATATTTGCATATACGA
 TACAAGGCTGTTAGAGAGATAAATTGGAATTAATTTGACTGTAAACACAAAGATATTAGTACAAAATAC
 GTGACGTAGAAAAGTAATAATTTCTTGGGTAGTTTGCAGTTTTAAAATTATGTTTTAAAATGGACTATCA
 TATGCTTACCGTAACTTGAAGTATTTTCGATTTCTTGGCTTTATATATCTTGTGGAAGGACGAAACAC
 CTCCGGTAACCAAGTTTTAGAGCTAGGCCAACATGAGGATCACCCATGTCTGCAGGGCCTAGCAAG
 TAAAATAAGGCTAGTCCGTTATCAACTTGGCCAACATGAGGATCACCCATGTCTGCAGGGCCAAGTG

GCACCGAGTCGGTGCTTTTTTTGAATTCTGATGCGGTATTTTCTCCTTACGCATCTGTGCGGTATTTAC
 ACCGCATACGTCAAAGCAACCATAGTACGCGCCCTGTAGCGGACCCAAGTTTGTACAAAAAGCAGG
 CTAGGCTCCGGTGCCCGTCAGTGGGCAGAGCGCACATCGCCACAGTCCCCGAGAAGTTGGGGGGAG
 GGGTCGGCAATTGAACCGGTGCCTAGAGAAGGTGGCGCGGGGTAAACTGGGAAAGTGATGTCGTGTA
 CTGGCTCCGCCTTTTTCCCGAGGGTGGGGGAGAACCGTATATAAGTGCAGTAGTCGCCGTGAACGTTT
 TTTTCGCAACGGGTTTGCCGCCAGAACACAGCTTAAGCTTGGTACCGAGCTCGGATCCACTAGTAAC
 GGCCGCCAGTGTGCTGGAATTCTGCAGATATCCATCACACTGGCGGCCGCTCGAGCATGCATCTAGAG
 GACAACTTTGTATACAAAAGTTGAGGCTGGATCCCGGTACCCTCGACACCTGCTGACAGGTCCACCAT
 GGGACCTAAGAAAAAGAGGAAGGTGGCGGCCGCTGACTACAAGGATGACGACGATAAATCTAGAATG
 GCTTCTAACTTTACTCAGTTTCGTTCTCGTCGACAATGGCGGAACTGGCGACGTGACTGTCGCCCCAAGC
 AACTTCGCTAACGGGATCGCTGAATGGATCAGCTCTAACTCGCGTTTACAGGCTTACAAAGTAACCTG
 TAGCGTTCGTAGAGCTCTGCGCAGAATCGCAAATACACCATCAAAGTCGAGGTGCCTAAAGGCGCCT
 GGCGTTCGTACTTAAATATGGAATAACCAATTTTCGCCACGAATCCGACTGCGAGCTTATTG
 TTAAGGCAATGCAAAGTCTCCTAAAAGATGGAACCCGATTCCCTCAGCAATCGCAGCAAACCTCCGGC
 ATCTACGAGGCCAGCCCCAAGAAGAAGGAAGGAGGAGTGTGAGTGGAGGAAAGTGGCGGTCAGGCTCG
 AGCCCCAAGAAAAACGGAAAGTGGAGGCAATGAAGGAGGGAGAAAAACAACAAACCCAGGGAG
 AAAAGTGAAGGGAATAAGAGAAAAAGCTCCTTCTCTAACAGTGCAGACGATATCAAGTCCAAGAAAA
 AGCGGGAGCAGTCTAATGACATTGCTAGGGGCTTCGAGAGAGGACTGGAGCCAGAAAAAATCATTGG
 GGCAACCGACAGCTGCGGCGATCTGATGTTTCTCATGAAATGGAAGGACACAGATGAGGCCGACCTG
 GTGCTCGCCAAAGAAGCTAACGTGAAGTGTCCCCAGATCGTCATTGCTTTTTACGAGGAAAGGCTCAC
 CTGGCACGCATATCCTGAGGATGCCGAAAACAAGGAGAAGGAATCAGCTAAGAGCTCGGGAGGTGGT
 TCGGGTGGCTCTGGATCAATGGACGCGAAATCACTTACGGCATGGTCGAGAACAACCTGGTTACGTTCAA
 GGACGTGTTTGTGGACTTTACACGTGAGGAGTGGAAATTGCTGGATACTGCGCAACAAATTGTGTATC
 GAAATGTCATGCTTGAGAATTACAAGAACCTCGTCAGTCTCGGATACCAGTTGACGAAACCGGATGTG
 ATCCTTAGGCTCGAAAAGGGGGAAGAACCTTGGCTGGTATAGAAGAATTCGATCCCGCTTCAGTGCAG
 GTGAATTCTACCACCCAGCTTAAGCTTGGTACCGAGCTCGGATCCACTAGTAACGGCCGCCAGTGTGC
 TGGAATTCTGCAGATATCCATCACACTGGCGGCCGCTCGAGCATGCATCTAGAGGTACCCAGCTTTCTT
 GTACAAAGTGGTAAGCTTGCCTCGAGCAGCGCTGCTCGAGAGATCTACGGGTGGCATCCCTGTGACCC
 CTCCCCAGTGCCTCTCCTGGCCCTGGAAGTTGCCACTCCAGTGGCCACCAGCCTTGTCTAATAAAATT
 AAGTTGCATCATTTTGTCTGACTAGGTGTCTTCTATAAATATTATGGGGTGGAGGGGGGTGGTATGGAG
 CAAGGGGCAAGTTGGGAAGACAACCTGTAGGGCTGCGGGGTCTATTGGGAACCAAGCTGGAGTGCA
 GTGGCACAATCTTGGCTCACTGCAATCTCCGCCTCCTGGGTTCAAGCGATTCTCCTGCCTCAGCCTCCC
 GAGTTGTTGGGATTCCAGGCATGCATGACCAGGCTCAGCTAATTTTTGTTTTTTGGTAGAGACGGGGT
 TTCACCATATTGGCCAGGCTGGTCTCCAATCCTAATCTCAGGTGATCTACCCACCTTGGCCTCCAAA
 TTGCTGGGATTACAGGCGTGAACCACTGCTCCCTTCCCTGTCTTCTGATTTTGTAGGTAACCACGTGC
 GGACCGAGCGGCCGACGAAACCCCTAGTGATGGAGTTGGCCACTCCCTCTCTGCGCGCTCGCTCGCTC
 ACTGAGGCCGGGCGACCAAAGGTCGCCCCGACGCCGGGCTTTGCCGGGCGGCCTCAGTGAGCGAGC
 GAGCGCGCAGCTGCCTGCAGGGGCGCCTGATGCGGTATTTTCTCCTTACGCATCTGTGCGGTATTTAC
 ACCGCATCGTCAAAGCAACCATAGTACGCGCCCTGTAGCGGCGCATTAAAGCGCGGCGGTGTGGTG
 GTTACGCGCAGCTGACCGCTACACTTCCAGCGCCCTAGCGCCGCTCCTTTCCGCTTTCTCCCTTCTC
 TTCTCGCCACGTTCCCGGCTTTCCCGTCAAGCTCTAAATCGGGGGCTCCCTTTAGGGTTCCGATTTA
 GTGCTTTACGGCACCTCGACCCCAAAAACTTGATTTGGGTGATGTTTACGTAAGTGGGCCATCGCCCT
 GATAGACGGTTTTTCGCCCTTTGACGTTGGAGTCCACGTTCTTTAATAGTGGACTCTTGTTCCAAACCTG
 GAACAACACTCAACCCTATCTCGGGCTATTCTTTTATTATAAAGGATTTTGCCGATTTCCGGCCTATT
 GGTAAAAAATGAGCTGATTTAACAAAAATTTAACCGCAATTTTAAACAAAATATTAACGTTTACAATT
 TTATGGTGCACCTCTCAGTACAATCTGCTCTGATGCCGCATAGTTAAGCCAGCCCCGACACCCGCCAAC
 ACCCGCTGACGCGCCCTGACGGCTTGTCTGCTCCCGGCATCCGCTTACAGACAAGCTGTGACCGTCTC
 CGGGAGCTGCATGTGTCAGAGGTTTTACCGTTCATACCGAAACGCGCGAGACGAAAGGGCCTCGTGA
 TACGCCTATTTTTATAGGTTAATGTCATGATAATAATGGTTTTCTTAGACGTGAGGTGGCACTTTTCGGG
 GAAATGTGCGCGGAACCCCTATTTGTTATTTTTCTAAATACATTCAAATATGTATCCGCTCATGAGAC
 AATAACCCTGATAAATGCTTCAATAATATTGAAAAAGGAAGAGTATGAGTATTCAACATTTCCGTGTC
 GCCCTTATTCCCTTTTTTGGCGCATTTTGCCTTCTGTTTTTGCTCACCCAGAAACGCTGGTGAAAGTAA
 AAGATGCTGAAGATCAGTTGGGTGCACGAGTGGGTACATCGAACTGGATCTCAACAGCGGTAAGATC
 CTTGAGAGTTTTCGCCCCGAAGAACGTTTTCCAATGATGAGCACTTTTAAAGTTCTGCTATGTGGCGCG
 GTATTATCCCGTATTGACGCCGGGCAAGAGCAACTCGGTCGCCGCATACACTATTCTCAGAATGACTT
 GGTTGAGTACTACCAGTCACAGAAAAGCATCTTACGGATGGCATGACAGTAAGAGAATTATGCAGTG

CTGCCATAACCATGAGTGATAAACTGCGGCCAACTTACTTCTGACAACGATCGGAGGACCGAAGGAG
 CTAACCGCTTTTTTGCACAACATGGGGGATCATGTAACCTCGCCTTGATCGTTGGGAACCGGAGCTGAA
 TGAAGCCATACCAAACGACGAGCGTGACACCACGATGCCTGTAGCAATGGCAACAACGTTGCGCAAA
 CTATTAACCTGGCGAACTACTTACTCTAGCTTCCCGGCAACAATTAATAGACTGGATGGAGGCGGATAA
 AGTTGCAGGACCACTTCTGCGCTCGGCCCTTCCGGCTGGCTGGTTTATTGCTGATAAATCTGGAGCCGG
 TGAGCGTGGGTCTCGCGGTATCATTGCAGCACTGGGGCCAGATGGTAAGCCCTCCCGTATCGTAGTTA
 TCTACACGACGGGGAGTCAGGCAACTATGGATGAACGAAATAGACAGATCGCTGAGATAGGTGCCTC
 ACTGATTAAGCATTGGTAACTGTCAGACCAAGTTTACTCATATATACTTTAGATTGATTTAAAACCTCA
 TTTTAAATTTAAAAGGATCTAGGTGAAGATCCTTTTTGATAATCTCATGACCAAAATCCCTTAAACGTGA
 GTTTTCGTTCCACTGAGCGTCAGACCCCGTAGAAAAGATCAAAGGATCTTCTTGAGATCCTTTTTTTCT
 GCGCGTAATCTGCTGCTTGCAAACAAAAAACCACCGCTACCAGCGGTGGTTTGTGGCCGATCAAG
 AGCTACCAACTCTTTTTCCGAAGGTAACCTGGCTTACGACAGCGCAGATACCAAATACTGTCCTTCTAG
 TGTAGCCGTAGTTAGGCCACCACTTCAAGAACTCTGTAGCACCGCCTACATACCTCGCTCTGCTAATCC
 TGTTACCAGTGGCTGCTGCCAGTGGCGATAAGTCGTGTCTTACCGGGTTGGACTCAAGACGATAGTTA
 CCGGATAAGGCGCAGCGGTTCGGGCTGAACGGGGGTTTCGTGCACACAGCCAGCTTGGAGCGAACGA
 CCTACACCAAGTATCCGTAAGCGGACGGTGTAGCTATGAGAAAGCGCCACGCTTCCCGAAGGGAGAA
 GGCGACAGACTTCCGTAAGCGGACGGTTCGGAACAGGAGAGCGCACGAGGGGACTTCCAGGGGG
 AAACGCCTGGTATCTTTATAGTCCTGTTCGGGTTTCGCCACCTCTGACTTGAGCGTTCGATTTTGTGATG
 CTCGTCAGGGGGCGGAGCCTATGGAAAACGCCAGCAACGCGGCCTTTTTACGGTTCCTGGCCTTTT
 GCTGGCCTTTTGCTCACATGT

[00109] AAV2/DJ-Cas9 and AAV/2/1 Cas9 (SEQ ID NO:66)

CCTGCAGGCAGCTGCGCGCTCGCTCGCTCACTGAGGCCGCCCGGGCAAAGCCCGGGCGTTCGGGCGACC
 TTTGGTGCCCCGCCTCAGTGAGCGAGCGAGCGCGCAGAGAGGGAGTGGCCAACTCCATCACTAGGG
 GTTCTCGGGCCTTAGAAATGTAGTCTTATGCAATACTCTTGTAGTCTTGCAACATGGTAACGATGAG
 TTAGCAACATGCCTTACAAGGAGAGAAAAAGCACCGTGCATGCCGATTGGTGGAAAGTAAGGTGGTAC
 GATCGTGCCTTATTAGGAAGGCAACAGACGGGTCTGACATGGATTGGACGAACCACTGAATTGCCGCA
 TTGCAGAGATATTGTATTTAAGTGCCTAGCTCGATACATAAAACCGGTGCCACCATGTACCCATACGA
 TGTTCCAGATTACGCTTCGCCGAAGAAAAAGCGCAAGGTTCGAAGCGTCCGACAAGAAGTACAGCATC
 GGCCTGGACATCGGCACCAACTCTGTGGGCTGGGCCGTGATCACCGACGAGTACAAGGTGCCAGCA
 AGAAATTCAAGGTGCTGGGCAACACCGACCGGCACAGCATCAAGAAGAACCTGATCGGAGCCCTGCT
 GTTCGACAGCGGCGAAACAGCCGAGGCCACCCGGCTGAAGAGAACCAGCAGAAGAAGATACACCAG
 ACGGAAGAACCAGTCTGCTATCTGCAAGAGATCTTCAGCAACGAGATGGCCAAGGTGGACGACAGC
 TTCTTCCACAGACTGGAAGAGTCTTCTGTTGGAAGAGGATAAGAAGCACGAGCGGCACCCCTCTT
 CGGCAACATCGTGGACGAGGTGGCCTACCACGAGAAGTACCCACCATCTACCACCTGAGAAAGAAA
 CTGGTGGACAGCACCGACAAGGCCGACCTGCGGTAGTCTATCTGGCCCTGGCCACATGATCAAGTT
 CCGGGGCCACTTCTGATCGAGGGCGACCTGAACCCCGACAACAGCGACGTGGACAAGCTGTTTCATCC
 AGCTGGTGCAGACCTACAACCAGCTGTTTCGAGGAAAACCCCATCAACGCCAGCGGCGTGGACGCCAA
 GGCCATCCTGTCTGCCAGACTGAGCAAGAGCAGACGGCTGGAAAATCTGATCGCCAGCTGCCCGGCG
 AGAAGAAGAATGGCTGTTTCGGCAACCTGATTGCCCTGAGCCTGGGCCTGACCCCAACTTCAAGAGC
 AACTTCGACCTGGCCGAGGATGCCAACTGCAGCTGAGCAAGGACACCTACGACGACGACCTGGACA
 ACCTGCTGGCCAGATCGGCGACCAGTACGCCGACCTGTTTCTGGCCGCAAGAACCTGTCCGACGCC
 ATCCTGCTGAGCGACATCCTGAGAGTGAACACCGAGATACCAAGGCCCCCTGAGCGCCTCTATGAT
 CAAGAGATACGACGAGCACCACCAGGACCTGACCCTGCTGAAAGCTCTCGTGCGGCAGCAGCTGCCT
 GAGAAGTACAAAGAGATTTTCTTCGACCAGAGCAAGAACGGCTACGCCGGCTACATTGACGGCGGAG
 CCAGCCAGGAAGAGTTCTACAAGTTCATCAAGCCATCCTGGAAAAGATGGACGGCACCAGGAACT
 GCTCGTGAAGCTGAACAGAGAGGACCTGCTGCGGAAGCAGCGGACCTTCGACAACGGCAGCATCCCC
 CACCAGATCCACCTGGGAGAGCTGCACGCCATTCTGCGGCGGCAGGAAGATTTTTACCCATTCTGAA
 GGACAACCGGAAAAGATCGAGAAGATCCTGACCTTCCGCATCCCCTACTACGTGGGCCCTCTGGCCA
 GGGGAAAACAGCAGATTCGCTGGATGACCAGAAAAGAGCGAGGAAACCATCACCCCTGGAACCTCGA
 GGAAGTGGTGGACAAGGGCGCTTCCGCCAGAGCTTCATCGAGCGGATGACCAACTTCGATAAGAAC
 CTGCCAACGAGAAGGTGCTGCCAAGCACAGCCTGCTGTACGAGTACTTCACCGTGTATAACGAGCT
 GACCAAAGTGAAATACGTGACCGAGGGAATGAGAAAAGCCCGCCTTCTGAGCGGCGAGCAGAAAAG
 GCCATCGTGGACCTGCTGTTCAAGACCAACCGGAAAGTGACCGTGAAGCAGCTGAAAGAGGACTACT
 TCAAGAAAATCGAGTCTTCGACTCCGTGGAAAATCTCCGGCGTGGAAAGATCGGTTCAACGCCTCCCTG
 GGCACATACCACGATCTGCTGAAAATTATCAAGGACAAGGACTTCTGGACAATGAGGAAAACGAGG

ACATTCTGGAAGATATCGTGCTGACCCTGACACTGTTTGAGGACAGAGAGATGATCGAGGAACGGCTG
AAAACCTATGCCACCTGTTCGACGACAAAAGTGATGAAGCAGCTGAAGCGGCGGAGATACACCGGCT
GGGGCAGGCTGAGCCGGAAGCTGATCAACGGCATCCGGGACAAGCAGTCCGGCAAGACAATCCTGGA
TTTCCTGAAGTCCGACGGCTTCGCCAACAGAACTTCATGCAGCTGATCCACGACGACAGCCTGACCT
TTAAAGAGGACATCCAGAAAGCCCAGGTGTCCGGCCAGGGCGATAGCCTGCACGAGCACATTGCCAA
TCTGGCCGGCAGCCCCGCCATTAAGAAGGGCATCCTGCAGACAGTGAAGGTGGTGGACGAGCTCGTG
AAAGTGATGGGCCGGCACAAGCCCCGAGAACATCGTGATCGAAATGGCCAGAGAGAACCAGACCACC
AGAAGGGACAGAAGAACAGCCGCGAGAGAATGAAGCGGATCGAAGAGGGCATCAAAGAGCTGGGCA
GCCAGATCCTGAAAGAACACCCCGTGGAAAACACCCAGCTGCAGAACGAGAAGCTGTACCTGTACTA
CCTGCAGAAATGGGCGGGATATGTACGTGGACCAGGAAGTGGACATCAACCGGCTGTCCGACTACGAT
GTGGACCATATCGTGCCTCAGAGCTTTCTGAAGGACGACTCCATCGACAACAAGGTGCTGACCAGAAG
CGACAAGAACCGGGGCAAGAGCGACAACGTGCCCTCCGAAGAGGTCTGTAAGAAGATGAAGAACTA
CTGGCGGCAGCTGCTGAACGCCAAGCTGATTACCCAGAGAAAAGTTCGACAATCTGACCAAGGCCGAG
AGAGGCGGCTGAGCGAACTGGATAAAGGCCGGCTTCATCAAGAGACAGCTGGTGGAAACCCGGCAGA
TCACAAAGCAGTGGCACAGATCCTGGACTCCCGGATGAACACTAAGTACGACGAGAATGACAAGCT
GATCCGGGAAGTGAAGTGATCACCTGAAGTCCAAGCTGGTGTCCGATTTCCGGAAGGATTTCCAGT
TTTACAAAGTGCAGAGATCAACAACACTACCACACGCCACGACGCCTACCTGAACCGCTGTGGGA
ACCGCCTGATCAAAAAGTACCCTAAGCTGGAAAGCGAGTTCGTGTACGGCGACTACAAGGTGTACG
ACGTGCGGAAGATGATCGCCAAGAGCGAGCAGGAAATCGGCAAGGCTACCGCCAAGTACTTCTTCTA
CAGCAACATCATGAACTTTTTCAAGACCGAGATTACCCTGGCCAACGGCGAGATCCGGAAGCGGCCTC
TGATCGAGACAAACGGCGAAACCGGGGAGATCGTGTGGGATAAAGGGCCGGGATTTTGCACCCGTGCG
GAAAGTGCTGAGCATGCCCAAGTGAATATCGTGAAAAAGACCGAGGTGCAGACAGGCGGCTTCAGC
AAAGAGTCTATCCTGCCAAGAGGAACAGCGATAAGCTGATCGCCAGAAAGAAGGACTGGGACCCTA
AGAAGTACGGCGGCTTCGACAGCCCCACCGTGGCCTATTCTGTGCTGGTGGTGGCCAAAGTGGAAAAG
GGCAAGTCCAAGAACTGAAGAGTGTGAAAGAGCTGCTGGGGATCACCATCATGGAAAGAAGCAGCT
TCGAGAAGAATCCCATCGACTTTCTGGAAGCCAAGGGCTACAAAGAAGTGA AAAAGGACCTGATCAT
CAAGCTGCCTAAGTACTCCCTGTTTCGAGCTGGAAAACGGCCGGAAGAGAATGCTGGCCTCTGCCGGC
AACTGCAGAAGGGAAACGAACTGGCCCTGCCCTCCAAATATGTGAACTTCTGTACCTGGCCAGCCAC
TATGAGAAGCTGAAGGGCTCCCCGAGGATAATGAGCAGAAACAGCTGTTTGTGGAACAGCACAAAGC
ACTACCTGGACGAGATCATCGAGCAGATCAGCGAGTTCTCCAAGAGAGTGATCCTGGCCGACGCTAAT
CTGGACAAAGTGCTGTCCGCCTACAACAAGCACCCGGGATAAGCCCATCAGAGAGCAGGCCGAGAATA
TCATCCACCTGTTTACCCTGACCAATCTGGGAGCCCTGCCGCTTCAAGTACTTTGACACCACCATCG
ACCGGAAGAGGTACACCAGCACCAAAAGAGGTGCTGGACGCCACCCTGATCCACCAGAGCATCACCGG
CCTGTACGAGACACGGATCGACCTGTCTCAGCTGGGAGGCGACAGCCCCAAGAAGAAGAGAAAGGTG
GAGGCCAGCTAAGAATTCAGCGGCGCGCTTCCCAAAACCCACCAAACTAAGTAACTGCTACTTCTCT
CAGCAACACCAAGATCAATGAAAGAGGCAAGGTGGGTCTTCGAGAAGACCTGCTTAATAAAAAGATCT
TTATTTTCATTAGATCTGTGTGTTGGTTTTTTGTGTGGCGCCGCAGGAACCCCTAGTGATGGAGTTGG
CCACTCCCTCTCTGCGCGCTCGCTCGCTCACTGAGGCCGGCGACCAAAGGTGCGCCGACGCCCGGGC
TTTGCCCGGGCGGCCTCAGTGAGCGAGCGAGCGCAGCTGCCTGCAGGGGCGCCTGATGCGGTATTT
TCTCCTTACGATCTGTGCGGTATTTACACCCGATACGTCGTAAGCAACCATAGTACCGCCCTGTAGC
GGCGCATTAAAGCGCGGGGTGTGGTGTACGCGCAGCGTGACCGGTACACTTGGCAGCGCCCTAGC
GCCCGCTCCTTTGCTTTCTCCCTTCTTCTCGCCACGTTGCGCGGCTTTCCCGTCAAGCTCTAAAT
CGGGGGCTCCCTTAGGGTTCCGATTTAGTGCTTTACGGCACCTCGACCCCCAAAAAATTGATTTGGGT
GATGGTTCACGTAGTGGGCCATCGCCCTGATAGACGGTTTTTTGCGCCCTTTGACGTTGGAGTCCACGTT
TTAATAGTGGACTCTTGTTCCAAACTGGAACAACACTCAACCCTATCTCGGGCTATTCTTTTGATTTAT
AAGGGATTTTGCAGATTTCCGGCCTATTGGTTAAAAAATGAGCTGATTTAACAAAAATTTAACGCGAAT
TTAACAAAAATATTAACGTTTACAATTTTATGGTGCCTCTCAGTACAATCTGCTCTGATGCCGCATAG
TTAAGCCAGCCCCGACACCCGCCAACACCCGCTGACGCGCCCTGACGGGCTTGTCTGCTCCCGGCATC
CGCTTACAGACAAGCTGTGACCGTCTCCGGGAGCTGCATGTGTGACAGAGTTTTACCGTCTATCCGA
AACGCGGAGACGAAAGGGCCTCGTGATACGCTATTTTTATAGGTTAATGTCATGATAATAATGGTT
TCTTAGACGTGAGTGGCACTTTTCGGGGAATGTGCGCGGAACCCCTATTTGTTTATTTTTCTAAATA
CATTCAAATATGTATCCGCTCATGAGACAATAACCCTGATAAATGCTTCAATAATATTGAAAAAGGAA
GAGTATGAGTATTAACATTTCCGTGTGCCCTTATTCCCTTTTTGCGGCATTTTGCTTCCTGTTTTG
CTACCCAGAAACGCTGGTGAAGTAAAAGATGCTGAAGATCAGTTGGGTGCACGAGTGGTTACAT
CGAACTGGATCTCAACAGCGGTAAGATCCTTGAGAGTTTTCGCCCCGAAGAAGTTTTCCAATGATGA
GCACTTTTAAAGTTCTGCTATGTGGCGCGGTATTATCCCGTATTGACGCCGGGCAAGAGCAACTCGGTC

GCCGCATACACTATTCTCAGAATGACTTGGTTGAGTACTCACCAGTCACAGAAAAGCATCTTACGGAT
 GGCATGACAGTAAGAGAATTATGCAGTGCTGCCATAACCATGAGTGATAAACTGCGGCCAACTTACT
 TCTGACAACGATCGGAGGACCGAAGGAGCTAACCGCTTTTTTGCACAACATGGGGGATCATGTAATC
 GCCTTGATCGTTGGGAACCGGAGCTGAATGAAGCCATAACAAACGACGAGCGTGACACCACGATGCC
 TGTAGCAATGGCAACAACGTTGCGCAAATACTAATACTGGCGAACTACTTACTCTAGCTTCCCGGCAAC
 AATTAATAGACTGGATGGAGGCGGATAAAGTTGCAGGACCACTTCTGCGCTCGGCCCTTCCGGCTGGC
 TGGTTTATTGCTGATAAATCTGGAGCCGGTGAGCGTGGAAGCCGCGGTATCATTGCAGCACTGGGGCC
 AGATGGTAAGCCCTCCCGTATCGTAGTTATCTACACGACGGGGAGTCAGGCAACTATGGATGAACGAA
 ATAGACAGATCGCTGAGATAGGTGCCTCACTGATTAAGCATTGGTAACTGTCAGACCAAGTTTACTCA
 TATACTTTAGATTGATTTAAAACCTTCATTTTTAATTTAAAAGGATCTAGGTGAAGATCCTTTTTGATA
 ATCTCATGACAAAATCCCTTAAACGTGAGTTTTTCGTTCCACTGAGCGTCAGACCCCGTAGAAAAGATC
 AAAGGATCTTCTTGAGATCCTTTTTTCTGCGCGTAATCTGCTGCTTGCAAACAAAAAACCACCGCTA
 CCAGCGGTGGTTTGTGGCCGATCAAGAGCTACCAACTCTTTTTCCGAAGGTAAGTGGCTTACAGCAGA
 GCGCAGATAACCAATACTGTCCTTCTAGTGTAGCCGTAGTTAGGCCACCACTTCAAGAACTCTGTAGC
 ACCGCCTACATACTCGCTCTGCTAATCCTGTTACCAGTGGCTGCTGCCAGTGGCGATAAGTTCGTGTCT
 TACCGGTTGGACTCAAGACGATAGTTACCGGATAAGGCGCAGCGGTGCGGGTGAACGGGGGGTTTCG
 TGCACACAGCCCAGCTTGGAGCGAACGACCTACACCGAACTGAGATACCTACAGCTGAGCTATGAG
 AAAGCGCCACGCTTCCCGAAGGGAGAAAGGCGGACAGGTATCCGGTAAGCGGCAGGGTTCGGAACAG
 GAGAGCGCACGAGGGAGCTTCCAGGGGGAAACGCCTGGTATCTTTATAGTCCTGTCCGGTTTCGCCAC
 CTCTGACTTGAGCGTCGATTTTTGTGATGCTCGTCAGGGGGGCGGAGCCTATGGAAAACGCCAGCAA
 CGCGGCCTTTTTACGGTTCCTGGCCTTTTGCTGGCCTTTTGCTCACATGT

[00110] *Cell culture for endogenous target repression* - HEK293FT, Neuro-2a, and Raw 264.7 cell lines (purchased from ATCC) were maintained in Dulbecco's modified Eagle's medium (DMEM - Life Technologies) with 10% fetal bovine serum (FBS - Life Technologies), 2mM glutamine, 1.0 mM sodium pyruvate (Life Technologies) and 1% penicillin- streptomycin (Life Technologies) in incubators at 37 °C and 5% CO₂.

[00111] *Transfection of in vitro cultured cells* - HEK293FT cells were seeded approximately 50,000 cells per well in 24-well plates and transfected the next day. HEK293FT cells were co-transfected with plasmid DNAs encoding gRNA (10 ng), dCas9 (200 ng), MS2-fused repressor (100 ng), puromycin resistant gene (50 ng), and Enhance Blue Fluorescent Protein (EBFP) as a transfection control (25 ng). Polyethylenimine (PEI) (Polysciences) was used to transfect HEK293FT cells. Transfection complexes were prepared according to manufacturer's instructions. Cells were treated with 0.5ug/ml puromycin (Gibco-life tech) at 24 hours post-transfection. Cells were collected 72 hours post-transfection and total RNA was collected from cells using RNAeasy Plus mini kit (Qiagen).

[00112] Neuro-2a cells were seeded approximately 50,000 cells per well in 24-well plates and transfected the next day. Cells were co-transfected with plasmids encoding gRNA (100 ng), Cas9 nuclease (70 ng), and EBFP as a transfection control (25 ng), and a Puromycin resistance gene (50ng). Plasmids were delivered to Neuro-2a cells with Lipofectamine LTX. Cells were treated with 0.5ug/ml puromycin (Gibco-life tech) at 24 hours post-transfection. 5 days later, cells were

treated with LPS at the concentration of 10ug/ml and after 5 hours total RNA was collected from cells using RNAeasy Plus mini kit (Qiagen).

[00113] For CXCR4 repression, Raw 264.7 cells were seeded approximately 50,000 cells per well in 24-well plates and transduced the next day. Cell culture media was removed and AAV was added to the culture at 1E+9 GC/cell in 250 microliter media. Media was refreshed 12 hours post-transduction. 5 days later, cells were treated with LPS at the concentration of 0.1ug/ml for 5 hours and then total RNA was collected using RNAeasy Plus mini kit (Qiagen).

[00114] *Quantitative RT-PCR Analysis* - Cells or tissues were lysed and RNA was extracted using RNeasy Plus mini kit (Qiagen) or trizol (Life Technologies) followed by cDNA synthesis using the High-Capacity RNA-to-cDNA Kit (Thermo Fisher) or iScript cDNA synthesis kit (Bio-Rad). qRT-PCR was performed using SYBR Green PCR Master Mix (Thermo Fisher). All analyses were normalized to 18s rRNA (ΔCt) and fold-changes were calculated against untransfected controls or Cas9 only groups for *in vitro* experiments and AAV-GFP injected group for *in vivo* experiments ($2^{-\Delta\Delta Ct}$). Primer sequences for qPCR are listed below.

[00115] *DNA isolation and real-time PCR For AAV genomic copies in tissues* - DNA isolation from tissues was performed using DNeasy blood and tissue kit (Qiagen) according to the manufacturer's protocol. qPCR reactions consist of 100ng template DNA from each sample and primers and probes for viral inverted terminal repeats (ITRs) using Probe Based qPCR Master Mix (IDT) according to the manufacturer's protocol. Data was normalized to mouse Acvr2b control gene.

[00116] *ELISA-based chemiluminescent assay* - After harvesting mice, plasma samples were collected and stored in one-time aliquots at -80 °C. Lung samples were lysed using 1x cell lysis buffer (Cell Signaling) (ratio of 100 mg of tissue to 1 ml of buffer) followed by homogenization and sonication of the lysed tissue. The assay was performed using the Q-Plex™ Mouse Cytokine – Screen (16-Plex) kit (Quansys Biosciences) following the manufacturer's protocol. Briefly, samples or calibrators were added into wells of a 96 well plate arrayed with analyte specific antibodies that capture GM-CSF, IL-1 α , IL-1 β , IL-4, IL-5, IL-6, IL-12p70, IL-17, MCP-1, MIP-1 α , RANTES, and TNF α . Plates were washed and biotinylated analyte specific antibodies were added. After washing, streptavidin-horseradish peroxidase (SHRP) was added. Following an additional wash, the amount of SHRP remaining on each location of the array was measured with the addition of a chemiluminescent substrate.

[00117] *Antibody ELISA* - Fifty microliter AAV particles diluted in 1x coating buffer (13 mM sodium carbonate, 35 mM sodium bicarbonate buffer, pH 9.2) containing 2×10^9 per viral particles were added to each well in a Microlon® high protein binding 96-well plate (Greiner). Wells were washed three times with 1× Tris Buffered Saline + Tween-20 (TBST, Bethyl) and blocked with 1x Tris Buffered Saline + 1% BSA (Bethyl) for 1 hr. at RT. Wells were washed three times with TBST. The standard curve was generated using purified mouse host IgG2a anti-AAV1 (Fitzgerald) antibody in twofold dilutions in TBST + 1% BSA + 1:500 negative control mouse serum, beginning from a concentration of 10,000 ng/ml AAV1 antibody. The standards were added to the plate followed by the serum samples at a dilution of 1:500 and incubated for 1 hr. at RT. Wells were washed four times with TBST and then goat anti-mouse HRP antibody was added at a concentration of 1:500 and incubated for 1 hr. at RT. Wells were washed four times with wash buffer and TMB substrate was added to the wells. Reactions were terminated by adding 0.18 M H₂SO₄ after development of the standard curve (15min). Finally, absorbance was measured at 450 wavelengths using a plate reader (BioTek). Absorbance results were exported and analyzed in Excel.

[00118] *Lactate assay* - Blood samples were collected using EDTA coated tubes. Samples were centrifuged at 1,000 x g for 10 minutes. Plasma was collected and stored at -80°C. Lactate assay was performed following the manufacturer's protocol (L-Lactate Assay Kit, Cayman Chemicals). Briefly, samples were deproteinated by adding 0.5 M MPA. After pelleting the protein, supernatant was added to Potassium Carbonate and centrifuged at 10,000 x g for five minutes at 4°C. Samples were diluted four-fold and added to the designated wells. Next, assay buffer cofactor mixture, Fluorometric Substrate, and Enzyme Mixture were added to each well. Plate was incubated for 20 minutes at RT and the fluorescence was measured using an excitation wavelength of 530-540 nm and an emission wavelength of 585-595 nm. Absorbance results were exported and analyzed in Excel according to the manufacturer's protocol.

[00119] *Examination of liver injury after LPS injection* - Plasma samples were sent to IDEXX Laboratories to measure a panel of tissue injury markers including ALT, Cholesterol, LDL, HDL, BUN, and Lipase.

[00120] *Animals* - All the experiments with animals were approved by the Institutional Animal Care and Use Committee (IACUC) at Arizona State University and have been performed according to institutional guidelines. All the experiments were performed on at least 3 mice of 6-

8 weeks old per group. Both male and female were included in each experiment. The sample size in each group is indicated in each figure legend. All mice were handled equally. Male C57BL/6 mice (JAX Stock number: 000664) were used for studying AAV2/1-GFP, tropism towards different organs and AAV-CRISPR activation experiments. Both male and female Rosa26-Cas9 knockin mice (JAX Stock number 026179) were used for AAV-CRISPR repression experiments.

[00121] *Retro-Orbital injections* - AAV particles were delivered to mice through Retro-orbital injection of the venous sinus. Animals were anesthetized with 3% isoflurane and virus particles were injected to the left eye with 100 microliters of AAV solution (1E11 to 1E12 genome copy per mouse for experiments with Cas9 expressing mice and a total of 1.01E+12 GC AAV-TTN and AAV-Cas9 (100:1 ratio) for C57BL/6 mice).

[00122] *Tissue harvest* - Mice were euthanized via CO₂ inhalation 3 weeks following injection. Tissue samples taken from liver, spleen, lung, testis, bone marrow and blood were collected in RLT (Life Technologies) or snap frozen for RNA analysis.

[00123] *In vivo LPS Induction* - For LPS induction, mice were treated by intraperitoneal (i.p.) injection of lipopolysaccharides (from Escherichia coli 0127:B8 (Sigma-Aldrich, St. Louis, MO, USA) dissolved in phosphate-buffered saline (PBS) at a concentration of 50 mg/ml. Mice were euthanized six hours post LPS injection via CO₂ inhalation.

[00124] *RNA Sequencing and Data Analysis* - RNA was extracted from mice bone marrow samples using RNeasy Plus mini kit (Qiagen) followed by globin mRNA depletion using GLOBINclear™ Kit, mouse/rat kit (Thermofisher). None-directional library preparation were performed at Novogene Corporation Inc. followed by RNA sequencing using Illumina Nova Platform with paired-end 150 run (2×150 bases). Coverage was minimum 25 million reads per sample. FASTQ files were then aligned to mouse genome sequence using STAR software and uniquely mapped read counts were visualized with Integrative Genomics Viewer (IGV). Gene expression level was calculated by the number of mapped reads. According to all gene expression level (RPKM or FPKM) of each sample, correlation coefficient of sample between groups was calculated. Readcount obtained from Gene Expression Analysis were used for differential expression analysis and differential expression analysis of different groups was performed using the DESeq2 R package. Hierarchical clustering analysis was carried out of log₁₀ (FPKM+1) of union differential expression genes, within all comparison groups. ClusterProfiler software was used for enrichment analysis, including GO Enrichment, DO

Enrichment, KEGG Enrichment and Reactome Enrichment.

[00125] *Statistical analysis* - Statistical analyses are included in the figure legends. All of the data are presented as the mean \pm SEM or mean \pm SD. N = number of individual transfections for *in vitro* experiments and N = number of animals for *in vivo* experiments. Statistical analyses were performed using Prism 7 Software (GraphPad). The inventors performed two-tailed Student's t-test. A value of $p < 0.05$ was considered significant, represented as * ($p < 0.05$), ** ($p < 0.01$), *** ($p < 0.001$) or n.s. (not significant).

[00126] Table 2: 14 nt guide sequences within gRNAs described herein - 5'-3'

Mouse	Myd88-1	GCCTAGTCCATCCA	(SEQ ID NO:1)
Mouse	Myd88-2	CCACCGATCAAGGT	(SEQ ID NO:2)
Mouse	Myd88-3	GCGAGCGTACTGGA	(SEQ ID NO:3)
Mouse	Myd88-4	GTGGACGGCACCGG	(SEQ ID NO:4)
Mouse	Mock	AGCTTAGGGATAAC	(SEQ ID NO:5)
Mouse	CXCR4-1	ATAAAAGTCCGGTT	(SEQ ID NO:6)
Mouse	CXCR4-2	TCCGGTAACCACCA	(SEQ ID NO:7)

[00127] Table 3: 20 nt guide sequences within gRNAs described herein gRNAs - 5'-3'

Human	SEL1L-1	GGCAGGAAGAGCAGCGGCGAGG	(SEQ ID NO:8)
Human	SEL1L-5	GGGGGGCGGATACTGACCCG	(SEQ ID NO:9)
Human	SEL1L-7	GGATACTGACCCGAGGACGCCG	(SEQ ID NO:10)
Human	SYVN1-10	GGGCGCTGGGTTCTTGGTGAGT	(SEQ ID NO:11)
Human	SYVN1-4	GGTTGCGGGCGTCGCAGGCA	(SEQ ID NO:12)
Human	SYVN1-3	GGCACCGGCGTCTGAGGTCTC	(SEQ ID NO:13)
Human	NEAT1-1	GGCGACAGGGAGGGATGCGCGCC	(SEQ ID NO:14)
Human	NEAT1-2	GGCGCGCCTGGGTGTAGTTGT	(SEQ ID NO:15)
Human	NEAT1-3	GGAAGTGGCTAGCTCAGGGCTTC	(SEQ ID NO:16)
Human	XIST-1	GGCAGCGCTTTAAGAACTGAA	(SEQ ID NO:17)
Human	XIST-2	GGACTGAAGATCTCTCTGCACTT	(SEQ ID NO:18)
Human	XIST-3	GGCCATATTTCTTACTCTCTCG	(SEQ ID NO:19)
Human	CXCR4	GCAGGTAGCAAAGTGACGCCGA	(SEQ ID NO:20)

[00128] Table 4: Mouse qPCR primers - 5'-3'. IDT™ PRIMETIME™ predesigned qPCR primer and probe kits are commercially available for the targets noted below and therefore the sequences of these primers are not separately provided herein.

GFP-FW1:	CAACCACTACCTGAGCACCC	(SEQ ID NO:21)
GFP-RV1:	GTCCATGCCGAGAGTGATCC	(SEQ ID NO:22)

Myd88 primer FW	GTGAGGATATACTGAAGGAGCTG	(SEQ ID NO:23)
Myd88 primer RV	CTGTAAAGGCTTCTCGGACTC	(SEQ ID NO:24)
Stat4-FW	CCTGACATTCCCAAAGACAAAGC	(SEQ ID NO:25)
Stat4-RV	TCTCTCAGCACAGCATATGCAC	(SEQ ID NO:26)
Cxcl1-FW	GACCATGGCTGGGATTCACC	(SEQ ID NO:27)
Cxcl1-RV	CCAAGGGAGCTTCAGGGTCA	(SEQ ID NO:28)
IFN α -fw	GGACT T TGGAT TCCCGCAGGAGA AG	(SEQ ID NO:29)
IFN α -RV	GCTGCATCAGACAGCCT TGCAGGTC	(SEQ ID NO:30)
IFN- γ -PRIMETIME	IDT:MM.PT.58.41769240	
IFN- β -PRIMETIME	IDT:MM.PT.58.30132453.G	
Icam-1-FW	CAATTTCTCATGCCGCACAG	(SEQ ID NO:31)
Icam-1-RV	AGCTGGAAGATCGAAAGTCCG	(SEQ ID NO:32)
TNF α -FW	AGGCTGCCCCGACTACGT	(SEQ ID NO:33)
TNF α -RV	GACTTTCTCCTGGTATGAGATAGCAAA	(SEQ ID NO:34)
NCF-FW	GCTGCGTGAACACTATCCTGG	(SEQ IDNO:35)
NCF-RV	AGGTCGTA CTCTCCATTCTGTA	(SEQ ID NO:36)
18S-FW	GGCCGTTCTTAGTTGGTGGAGCG	(SEQ ID NO:37)
18S-RV	CTGAACGCCACTTGTCCTC	(SEQ ID NO:38)
CD68-FW	TGCGGCTCCCTGTGTGT	(SEQ ID NO:39)
CD68-RV	TCTTCCTCTGTTCCCTGGGCTAT	(SEQ ID NO:40)
CXCR4-PRIMETIME	IDT:MM.PT.58.41597935	
CD8-FW	GCTCAGTCATCAGCAACTCG	(SEQ ID NO:41)
CD8-RV	TCACAGGCGAAGTCCATC	(SEQ ID NO:42)
CD4-FW	GAGAGTCAGCGGAGTTCTC	(SEQ ID NO:43)
CD4-RV	CTCACAGGTCAAAGTATTGTTG	(SEQ ID NO:44)
IL6-PRIMETIME	IDT: Mm.PT.58.100	
IL1B-FW	TCGCTCAGGGTCACAAGAAA	(SEQ ID NO:45)
IL1B-RV	CATCAGAGGCAAGGAGGAAAAC	(SEQ ID NO:46)

[00129] Table 5: Human qPCR primers - 5'-3'

SEL1L-FW	GTGGCTGTTGGAGTCGGTAT	(SEQ ID NO:47)
SEL1L-RV	ATTCCTCCCCACCCTCTCT	(SEQ ID NO:48)
NEAT1-FW	AGGTCAGGCAGAGGAAGTCA	(SEQ ID NO:49)
NEAT1-RV	CTGCCTCCCGATAACAACAT	(SEQ ID NO:50)
XIST-1-FW	AGGTCAGGCAGAGGAAGTCA	(SEQ ID NO:51)
XIST-1-RV	CTGCCTCCCGATAACAACAT	(SEQ ID NO:52)
SYVN1-FW	ACCAGCATCCCTAGCTCAGA	(SEQ ID NO:53)
SYVN1-RV	TCCTCAGGCATCTCCTCTGT	(SEQ ID NO:54)

[00130] MS2-*krab*-Mecp2 (SEQ ID NO:55), MS2 (SEQ ID NO:56), *krab* (SEQ ID NO:57),

and Mecp2 (SEQ ID NO:58):

CCACCATGGGACCTAAGAAAAAGAGGAAGGTGGCGGCCGCTGACTACAAGGATG
 ACGACGATAAATCTAGAATGGCTTCTAACTTTACTCAGTTCGTTCTCGTCGACA
 ATGGCGGAACTGGCGACGTGACTGTCGCCCCAAGCAACTTCGCTAACGGGATC
 GCTGAATGGATCAGCTCTAACTCGCGTTCACAGGCTTACAAAGTAACCTGTAGC
 GTTCGTCAGAGCTCTGCGCAGAATCGCAAATACACCATCAAAGTCGAGGTGCC
 TAAAGGCGCCTGGCGTTCGTACTTAAATATGGAATAACCATTTCCAATTTTCGC
 CACGAATTCCGACTGCGAGCTTATTGTTAAGGCAATGCAAGGTCTCCTAAAAGA
 TGAAACCCGATTCCCTCAGCAATCGCAGCAAACCTCCGGCATCTACGAGGCCA
 GCCCAAGAAGAAGAGGAAGGTGAGTGGTGGAGGAAGTGGCGGGTCAGGGTCGTC
 CATGGACGCGAAATCACTTACGGCATGGTTCGAGAACAACCTGGTTACGTTCAAGGACGTGTT
 TGTGGACTTTACACGTGAGGAGTGGAAATTGCTGGATACTGCGCAACAAATTGTGTATCGA
 AATGTCATGCTTGAGAATTACAAGAACCTCGTCAGTCTCGGATAACCAGTTGACGAAACCGG
 ATGTGATCCTTAGGCTCGAAAAGGGGGAAGAACCTTGGCTGGTATCGGGAGGTGGTTCG
 GGTGGCTCTGGATCAAGCCCAAAGAAGAAACGGAAGGTGGAAGCCTCAGTGCAGGT
GAAAAGGGTGCTGGAAAAATCCCCCGCAAACCTCCTCGTGAAGATGCCCTTCCAGG
CTTCCCCTGGCGGAAAAGGTGAAGGGGGTGGCGCAACCACATCTGCCAGGTCATG
GTCATCAAGCGACCTGGAAGGAAAAGAAAGGCCGAGGCTGACCCTCAGGCCATTCC
AAAGAAACGGGGACGCAAGCCAGGGTCCGTGGTTCGCAGCTGCAGCAGCTGAGGCT
AAGAAAAGGCAGTGAAGGAAAGCTCCATCCGCAGTGTGCAGGAGACTGTCCTGCC
CATCAAGAAGAGGAAGACTAGGGAGACCGTGTCCATCGAGGTCAAAGAAGTGGTCA
AGCCCCTGCTCGTGTCCACCCTGGGCGAAAAATCTGGAAAGGGGCTCAAACATGC
AAGTCACCTGGACGGAAAAGCAAGGAGTCTAGTCCAAAGGGGCGCTCAAGCTCCGC
TTCTAGTCCCCCTAAAAAGGAACACCATCACCATCACCATCACGCCGAGTCTCCTAA
GGCTCCTATGCCACTGCTCCCACCACCTCCACCACCTGAGCCACAGTCAAGCGAAGA
CCCCATCAGCCCACCCGAGCCTCAGGATCTGTCCTCTAGTATTTGCAAAGAGGAAAA
GATGCCCAGAGCAGGCAGCCTGGAGAGTGATGGCTGTCCAAAAGAACCCGCCAAGA
CCCAGCCTATGGTGGCAGCCGCTGCAACTACCACCACAACCACAACCTACCACAGTG
GCCGAAAAATAACAAGCATCGCGGCGAGGGCGAACGAAAGGACATTGTGTCAAGCTC
CATGCCCAGACCTAACCGGGAGGAACCAGTCGATAGTAGGACACCCGTGACTGAGA
GAGTCTCATGA

[00131] MS2-HP1A-*krab* (SEQ ID NO:59), MS2 (SEQ ID NO:56), HP1A (SEQ ID NO:60), *krab* (SEQ ID NO:57)

CCACCATGGGACCTAAGAAAAAGAGGAAGGTGGCGGCCGCTGACTACAAGGATG
 ACGACGATAAATCTAGAATGGCTTCTAACTTTACTCAGTTCGTTCTCGTCGACA
 ATGGCGGAACTGGCGACGTGACTGTCGCCCAAGCAACTTCGCTAACGGGATC
 GCTGAATGGATCAGCTCTAACTCGCGTTCACAGGCTTACAAAGTAACCTGTAGC
 GTTCGTCAGAGCTCTGCGCAGAATCGCAAATACACCATCAAAGTCGAGGTGCC
 TAAAGGCGCCTGGCGTTCGTACTTAAATATGGAATAACCATTCCAATTTTCGC
 CACGAATTCCGACTGCGAGCTTATTGTTAAGGCAATGCAAGGTCTCCTAAAAGA
 TGGAACCCGATTCCCTCAGCAATCGCAGCAAACCTCCGGCATCTACGAGGCCA
 GCCCAAGAAGAAGAGGAAGGTGAGTGGTGGAGGAAGTGGCGGGTCAGGGTTCGTC
 CAGCCCCAAGAAAAACGGAAAGTGGAGGCATCAATGAAGGAGGGAGAAAACAAC
AAACCCAGGGAGAAAAGTGAAGGGAATAAGAGAAAAAGCTCCTTCTCTAACAGTGC
AGACGATATCAAGTCCAAGAAAAAGCGGGAGCAGTCTAATGACATTGCTAGGGGCT
TCGAGAGAGGACTGGAGCCAGAAAAATCATTGGGGCAACCGACAGCTGCGGCGAT
CTGATGTTTCTCATGAAATGGAAGGACACAGATGAGGCCGACCTGGTGCTCGCCAA
AGAAGCTAACGTGAAGTGTCCCAGATCGTCATTGCTTTTTACGAGGAAAGGCTCAC
CTGGCACGCATATCCTGAGGATGCCGAAAACAAGGAGAAGGAATCAGCTAAGAGCC
 TCGGGAGGTGGTTCGGGTGGCTCTGGATCAATGGACGCGAAATCACTTACGGCATGGT
 CGAGAACACTGGTTACGTTCAAGGACGTGTTTGTGGACTTTACACGTGAGGAGTGGAAATT
 GCTGGATACTGCGCAACAAATTGTGTATCGAAATGTCATGCTTGAGAATTACAAGAACCTC
 GTCAGTCTCGGATACCAGTTGACGAAACCGGATGTGATCCTTAGGCTCGAAAAGGGGGAA
 GAACCTTGGCTGGTATGA

[00132] MS2-Mecp2-*krab*-MBD2B (SEQ ID NO:61), MS2 (SEQ ID NO:56), Mecp2 (SEQ ID NO:58), *krab* (SEQ ID NO:57), MBD2B (SEQ DI NO:62)

CCACCATGGGACCTAAGAAAAAGAGGAAGGTGGCGGCCGCTGACTACAAGGATG
 ACGACGATAAATCTAGAATGGCTTCTAACTTTACTCAGTTCGTTCTCGTCGACA
 ATGGCGGAACTGGCGACGTGACTGTCGCCCAAGCAACTTCGCTAACGGGATC
 GCTGAATGGATCAGCTCTAACTCGCGTTCACAGGCTTACAAAGTAACCTGTAGC

GTTCGTCAGAGCTCTGCGCAGAATCGCAAATACACCATCAAAGTCGAGGTGCC
TAAAGGCGCCTGGCGTTCGTACTTAAATATGGAAC TAACCATTCCAATTTTCGC
CACGAATTCCGACTGCGAGCTTATTGTTAAGGCAATGCAAGGTCTCCTAAAAGA
TGGAAACCCGATTCCCTCAGCAATCGCAGCAA ACTCCGGCATCTACGAGGCCA
GCCCCAAGAAGAAGAGGAAGGTGAGTGGTGGAGGAAGTGGCGGGTCAGGGTCGTC
CAGCCCAAAGAAGAAACGGAAGGTGGAAGCCTCAGTGCAGGTGAAAAGGGTGCTG
GAAAAATCCCCGGCAA ACTCCTCGTGAAGATGCCCTTCCAGGCTTCCCCTGGCGGA
AAAGGTGAAGGGGGTGGCGCAACCACATCTGCCCAGGTCATGGTCATCAAGCGACC
TGGAAGGAAAAGAAAGGCCGAGGCTGACCCTCAGGCCATTCCAAAGAAACGGGGA
CGCAAGCCAGGGTCCGTGGTCGCAGCTGCAGCAGCTGAGGCTAAGAAAAAGGCAGT
GAAGGAAAGCTCCATCCGCAGTGTGCAGGAGACTGTCCTGCCCATCAAGAAGAGGA
AGACTAGGGAGACCGTGTCCATCGAGGTC AAAGAAGTGGTCAAGCCCCTGCTCGTG
TCCACCCTGGGCGAAAAATCTGGAAAGGGGCTCAAACATGCAAGTCACCTGGACG
GAAAAGCAAGGAGTCTAGTCCAAAGGGGCGCTCAAGCTCCGCTTCTAGTCCCCCTA
AAAAGGAACACCATCACCATCACCATCACGCCGAGTCTCCTAAGGCTCCTATGCCAC
TGCTCCCACCACCTCCACCACCTGAGCCACAGTCAAGCGAAGACCCCATCAGCCCAC
CCGAGCCTCAGGATCTGTCCTCTAGTATTTGCAAAGAGGAAAAGATGCCCAGAGCA
GGCAGCCTGGAGAGTGATGGCTGTCCAAAAGAACCCGCCAAGACCCAGCCTATGGT
GGCAGCCGCTGCAACTACCACCACAACCACA ACTACCACAGTGGCCGAAAAATACA
AGCATCGCGGCGAGGGCGAACGAAAGGACATTGTGTCAAGCTCCATGCCCAGACCT
AACCGGGAGGAACCAGTCGATAGTAGGACACCCGTGACTGAGAGAGTCTCATCGGG
AGGTGGTTCGGGTGGCTCTGGATCAATGGACGCGAAATCACTTACGGCATGGTCGAGA
ACACTGGTTACGTTCAAGGACGTGTTTGTGGACTTTACACGTGAGGAGTGGAATTGCTG
GATACTGCGCAACAAATTGTGTATCGAAATGTCATGCTTGAGAATTACAAGAACCTCGTCA
GTCTCGGATACCAGTTGACGAAACCGGATGTGATCCTTAGGCTCGAAAAGGGGGAAGAAC
CTTGGCTGGTATCGGGAGGTGGTTCGGGTGGCTCTGGATCATCCCCAAAAAGAAAC
GCAAGGTGAGGCATCTGCTCGCTATCTCGGTAACACTGTGCATCTGTCATCATT
CGATTTCCGCACTGGCAAGATGATGCCTTCTAAACTGCAGAAGAACAACAGAA
GGCTGAGAAATGACCCACTCAACCAGAATAAGGGAAAACCCGATCTGAACACC
AACTCCCTATCCGGCAGACAGCTAGTATTTTCAAGCAGCCTGTGACTAAAGTC
ACCAACCACCCATCCAATAAGGTGAAATCTGACCCACAGAGGATGAATGAGCA

GCCCAGACAGCTGTTTTGGGAAAAGCGCCTGCAGGGTCTCTCTGCAAGTGATG
TGACAGAGCAGATCATTAAAGACTATGGAAGTGCCTAAAAGGACTCCAGGGAGTG
GGACCTGGGTCTAACGACGAGACTCTGCTCTCAGCTGTCGCAAGCGCACTGCA
TACCAGCTCCGCACCCATCACAGGACAGGTGAGTGCCGCTGTCGAGAAGAACC
CAGCCGTGTGGCTGAATACTTCACAGCCCCTCTGCAAGGCTTTCATCGTCACCG
ACGAGGATATTCGGAAGCAGGAGGAACGCGTGCAGCAGGTCCGAAAAATCCTG
GAAGACGCTCTCATGGCAGATATTCTGAGCAGAGCAGCCGACACCGAGGAAAT
GGATATTGAGATGGACTCCGGGGATGAAGCCTGAA

[00133] MS2-*MBD2B*-*HP1A* (SEQ ID NO:63), MS2 (SEQ ID NO:56), *MBD2B* (SEQ ID NO:62), *HP1A* (SEQ ID NO:60)

CCACCATGGGACCTAAGAAAAAGAGGAAGGTGGCGGCCGCTGACTACAAGGATG
ACGACGATAAATCTAGAATGGCTTCTAACTTTACTCAGTTCGTTCTCGTCGACA
ATGGCGGAACTGGCGACGTGACTGTCGCCCAAGCAACTTCGCTAACGGGATC
GCTGAATGGATCAGCTCTAACTCGCGTTCACAGGCTTACAAAGTAACCTGTAGC
GTTTCGTCAGAGCTCTGCGCAGAATCGCAAATACACCATCAAAGTCGAGGTGCC
TAAAGGCGCCTGGCGTTCGTACTTAAATATGGAACTAACCATTCCAATTTTCGC
CACGAATTCCGACTGCGAGCTTATTGTTAAGGCAATGCAAGGTCTCCTAAAAGA
TGAAACCCGATTCCCTCAGCAATCGCAGCAAACCTCCGGCATCTACGAGGCCA
GCCCCAAGAAGAAGAGGAAGGTGAGTGGTGGAGGAAGTGGCGGGTCAGGGTTCGTC
CTCCCCAAAAAGAAACGCAAGGTGGAGGCATCTGCTCGCTATCTCGGTAACACTGTC
GATCTGTCATCATTTCGATTTCCGCACTGGCAAGATGATGCCTTCTAAACTGCAGAAGAACA
AACAGAGGCTGAGAAATGACCCACTCAACCAGAATAAGGGAAAACCCGATCTGAACACCA
CACTCCCTATCCGGCAGACAGCTAGTATTTTCAAGCAGCCTGTGACTAAAGTCACCAACCA
CCCATCCAATAAGGTGAAATCTGACCCACAGAGGATGAATGAGCAGCCAGACAGCTGTT
TTGGGAAAAGCGCCTGCAGGGTCTCTCTGCAAGTGATGTGACAGAGCAGATCATTAAAGAC
TATGAAACTGCCAAAAGGACTCCAGGGAGTGGGACCTGGGTCTAACGACGAGACTCTGCT
CTCAGCTGTCGCAAGCGCACTGCATACCAGCTCCGCACCCATCACAGGACAGGTGAGTG
CCGCTGTCGAGAAGAACCAGCCGTGTGGCTGAATACTTCACAGCCCCTCTGCAAGGCTT
TCATCGTCACCGACGAGGATATTCGGAAGCAGGAGGAACGCGTGCAGCAGGTCCGAAAA
ATCCTGGAAGACGCTCTCATGGCAGATATTCTGAGCAGAGCAGCCGACACCGAGGAAATG

GATATTGAGATGGACTCCGGGGATGAAGCCTCGGGAGGTGGTTCGGGTGGCTCTGG
ATCAAGCCCCAAGAAAAACGGAAAGTGGAGGCATCAATGAAGGAGGGAGAAAAC
AACAAACCCAGGGAGAAAAGTGAAGGGAATAAGAGAAAAAGCTCCTTCTCTAACA
GTGCAGACGATATCAAGTCCAAGAAAAAGCGGGAGCAGTCTAATGACATTGCTAGG
GGCTTCGAGAGAGGACTGGAGCCAGAAAAATCATTGGGGCAACCGACAGCTGCGG
CGATCTGATGTTTCTCATGAAATGGAAGGACACAGATGAGGCCGACCTGGTGCTCGC
CAAAGAAGCTAACGTGAAGTGTCCCCAGATCGTCATTGCTTTTTACGAGGAAAGGCT
CACCTGGCACGCATATCCTGAGGATGCCGAAAACAAGGAGAAGGAATCAGCTAAGA
GCTGA

CLAIMS

We claim:

1. A synthetic repression system for repressing myeloid differentiation primary response 88 (MyD88) expression *in vivo* comprising a single amplicon comprising:
 - (a) a guide RNA (gRNA) comprising (i) a guide sequence complementary to a portion of MyD88 and (ii) an aptamer target site specific for an RNA binding protein;
 - (b) a nucleotide sequence encoding the aptamer RNA binding protein fused to one or more repression domain; and
 - (c) a nucleotide sequence encoding a multifunctional Cas nuclease;wherein the single amplicon is packaged in a vector for DNA-based viral delivery.
2. The synthetic repression system of claim 1, wherein the gRNA guide sequence is 15 or fewer nucleotides in length.
3. The synthetic repression system of claim 1, wherein the gRNA guide sequence targets within 100 base pairs (bp) upstream of TATA box region of MyD88.
4. The synthetic repression system of claim 1, wherein gRNA guide sequence is selected from the group consisting of SEQ ID NOs: 1-4.
5. The synthetic repression system of claim 1, wherein the aptamer target site is an MS2 aptamer target site and the RNA binding protein is bacteriophage MS2 coat protein.
6. The synthetic repression system of claim 5, wherein the gRNA comprises the MS2 aptamer target site and the guide sequence is 15 or fewer nucleotides in length.
7. The synthetic repression system of claim 5, wherein the repression domain is selected from the group consisting of a Kruppel associated box (KRAB) domain, methyl-CpG (mCpG) binding domain 2 (meCP2), Switch independent 3 transcription regulator family member A (SIN3A), histone deacetylase HDT1 (HDT1), n-terminal truncation of methyl-CpG-binding

domain containing protein 2 (MBD2B), nuclear inhibitor of protein phosphatase-1 (NIPP1), and heterochromatin protein 1 (HP1A).

8. The synthetic repression system of claim 7, wherein the nucleotide of (b) encodes MS2 fused to at least two repression domains.

9. The synthetic repression system of claim 8, wherein the nucleotide sequence encoding MS2 fused to a repression domain comprises a sequence selected from the group consisting of SEQ ID NOs:55, 59, 61, and 63.

10. The synthetic repression system of claim 1, wherein the system comprises a nucleotide sequence encoding a multifunctional Cas nuclease fused to the aptamer RNA binding protein and a repression domain.

11. The synthetic repression system of claim 1, wherein the Cas nuclease is an *S. aureus* Cas9 nuclease or an *S. pyrogenes* Cas9 nuclease.

12. The synthetic repression system of claim 1, wherein the vector is an AAV2/1 delivery vector.

13. A pharmaceutical composition comprising the synthetic repression system of claim 1 and a pharmaceutically acceptable delivery vehicle.

14. A method of treating a subject with septicemia comprising administering to the subject with septicemia a therapeutically effective amount of the composition of claim 13.

15. The method of claim 14, wherein levels of immune related markers Icam-1, Tnf α , Ncf, Il6, Ifn- α , Ifn- β , Ifn- γ , and Stat4 are reduced in the subject following administration.

16. The method of claim 14, additionally comprising the step of monitoring the level of at least one immune related marker selected from the group consisting of Icam-1, Tnf α , Ncf, Il6, Ifn-

α , Ifn- β , Ifn- γ , and Stat4 and administering to the subject additional doses of the synthetic repression system composition until the level of the at least one immune related marker is reduced.

17. The method of claim 14, wherein serum lactate is reduced in the subject following administration.

18. A method of preventing an adverse immune response in a subject comprising administering to the subject a therapeutically effective amount of the composition of claim 13 prior to or concurrently with administration of an AAV vector based gene therapy to the subject, wherein the level of an immune related marker selected from the group consisting of Icam-1, Tnf α , Ncf, Il6, Ifn- α , Ifn- β , Ifn- γ , and Stat4 is reduced in the subject relative to subject who received the AAV vector based gene therapy but did not receive the therapeutic synthetic repression system composition.

19. The method of claim 18, wherein the subject previously received at least one AAV vector based gene therapy.


20. The method of claim 18, additionally comprising the step of monitoring the level of at least one immune related marker selected from the group consisting of Icam-1, Tnf α , Ncf, Il6, Ifn- α , Ifn- β , Ifn- γ , and Stat4 and administering to the subject additional doses of the synthetic repression system composition until the level of the at least one immune related marker is reduced.

21. A method of treating Waldenström macroglobulinemia in a subject in need thereof comprising administering to a subject with Waldenström macroglobulinemia a therapeutically effective amount of the composition of claim 13.


22. The method of treatment of claim 21, additionally comprising administering to the subject a composition comprising a gRNA configured to cleave MyD88 at L265 locus and a

homology directed repair template encoding a portion of wild-type MyD88 at the L265 locus, whereby levels of endogenous L265P mutant MyD88 are reduced in the subject.


FIGS. 1A-1H






FIGS. 1A-1H CONTINUED




FIGS. 1A-1H CONTINUED



FIGS. 2 A-2H




FIGS. 2A-2H CONTINUED




6/23

FIGS. 2A-2H CONTINUED



FIGS. 3A-3B

A



B






FIG. 4




FIGS. 5A-5C CONTINUED

B



FIGS. 5A-5C CONTINUED

C








FIG. 7



FIGS. 8A-8C







FIG. 9



FIGS. 10A-10C




FIG. 11




FIG. 12




FIG. 13




FIG. 14




FIG. 15










FIG. 15 CONTINUED




22/23


FIG. 16



FIGS. 1A-1H



FIGS. 1A-1H CONTINUED



FIGS. 1A-1H CONTINUED

