

(51) International Patent Classification:
C07D 403/14 (2006.01) **C07D 405/14** (2006.01)(21) International Application Number:
PCT/US2010/060077(22) International Filing Date:
13 December 2010 (13.12.2010)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
61/286,942 16 December 2009 (16.12.2009) US
12/957,512 1 December 2010 (01.12.2010) US(71) Applicant (for all designated States except US): **BRISTOL-MYERS SQUIBB COMPANY** [US/US]; P.O. Box 4000, Route 206 and ProvinceLine Road, Princeton, New Jersey 08543-4000 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **BELEMA, Makenen** [US/US]; c/o Bristol-Myers Squibb Company, 5 Research Parkway, Wallingford, CT 06492 (US). **ROMINE, Jeffrey, Lee** [US/US]; c/o Bristol-Myers Squibb Company, 5 Research Parkway, Wallingford, CT 06492 (US). **NGUYEN, Van, N.** [US/US]; c/o Bristol-Myers Squibb Company, 5 Research Parkway, Wallingford, Connecticut 06492 (US). **WANG, Gan** [US/US]; c/o Bristol-Myers Squibb Company, 5 Research Parkway, Wallingford, Connecticut 06492 (US). **LOPEZ, Omar, D.** [US/US]; c/o Bristol-Myers Squibb Company, 5 Research Parkway, Wallingford, Connecticut 06492 (US). **ST. LAURENT, Denis, R.** [US/US]; c/o Bristol-Myers Squibb Company, 5 Research Parkway, Wallingford, Connecticut 06492 (US). **CHEN, Qi** [CN/US]; c/o Bristol-Myers Squibb Company, 5 Research Parkway, Wallingford, Connecticut 06492 (US). **BENDER, John, A.** [US/US]; c/o Bristol-Myers Squibb Company, 5 Research Parkway, Wallingford, Connecticut 06492 (US). **YANG, Zhong** [CN/US]; c/o Bristol-Myers Squibb Company, 5 Research Parkway, Wallingford, Connecticut 06492 (US). **HEWAWASAM, Piyasena** [US/US]; c/o Bristol-Myers Squibb Company, 5Research Parkway, Wallingford, Connecticut 06492 (US). **XU, Ningning** [CN/US]; c/o Bristol-Myers Squibb Company, 5 Research Parkway, Wallingford, Connecticut 06492 (US). **MEANWELL, Nicholas, A.** [US/US]; c/o Bristol-Myers Squibb Company, 5 Research Parkway, Wallingford, Connecticut 06492 (US). **EASTER, John, A.** [US/US]; c/o Bristol-Myers Squibb Company, 5 Research Parkway, Wallingford, Connecticut 06492 (US). **SU, Bao-Ning** [CN/US]; c/o Bristol-Myers Squibb Company, 1 Squibb Drive, New Brunswick, New Jersey 08903 (US). **SMITH, Michael, J.** [US/US]; c/o Bristol-Myers Squibb Company, 1 Squibb Drive, New Brunswick, New Jersey 08903 (US).(74) Agents: **MINGO, Pamela, A.** et al.; Bristol-Myers Squibb Company, P.O. Box 4000, Princeton, New Jersey 08543-4000 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.


(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))

[Continued on next page]

(54) Title: HEPATITIS C VIRUS INHIBITORS



(57) Abstract: The present disclosure relates to compounds, compositions and methods for the treatment of Hepatitis C virus (HCV) infection. Also disclosed are pharmaceutical compositions containing such compounds and methods for using these compounds in the treatment of HCV infection.



— *as to the applicant's entitlement to claim the priority of
the earlier application (Rule 4.17(iii))*

Published:

— *with international search report (Art. 21(3))*

HEPATITIS C VIRUS INHIBITORS

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims the benefit of U.S. Provisional Application Serial
5 Number 61/286,942 filed December 16, 2009.

[0002] The present disclosure is generally directed to antiviral compounds, and more specifically directed to compounds which can inhibit the function of the NS5A protein encoded by Hepatitis C virus (HCV), compositions comprising such compounds, and
10 methods for inhibiting the function of the NS5A protein.

[0003] HCV is a major human pathogen, infecting an estimated 170 million persons worldwide - roughly five times the number infected by human immunodeficiency virus type 1. A substantial fraction of these HCV infected individuals develop serious progressive liver disease, including cirrhosis and hepatocellular carcinoma.

15 **[0004]** The current standard of care for HCV, which employs a combination of pegylated-interferon and ribavirin, has a non-optimal success rate in achieving sustained viral response and causes numerous side effects. Thus, there is a clear and long-felt need to develop effective therapies to address this undermet medical need.


[0005] HCV is a positive-stranded RNA virus. Based on a comparison of the
20 deduced amino acid sequence and the extensive similarity in the 5' untranslated region, HCV has been classified as a separate genus in the Flaviviridae family. All members of the Flaviviridae family have enveloped virions that contain a positive stranded RNA genome encoding all known virus-specific proteins via translation of a single, uninterrupted, open reading frame.

25 **[0006]** Considerable heterogeneity is found within the nucleotide and encoded amino acid sequence throughout the HCV genome due to the high error rate of the encoded RNA dependent RNA polymerase which lacks a proof-reading capability. At least six major genotypes have been characterized, and more than 50 subtypes have been described with distribution worldwide. The clinical significance of the genetic heterogeneity of
30 HCV has demonstrated a propensity for mutations to arise during monotherapy treatment, thus additional treatment options for use are desired. The possible modulator effect of genotypes on pathogenesis and therapy remains elusive.

[0007] The single strand HCV RNA genome is approximately 9500 nucleotides in length and has a single open reading frame (ORF) encoding a single large polypeptide of about 3000 amino acids. In infected cells, this polypeptide is cleaved at multiple sites by cellular and viral proteases to produce the structural and non-structural (NS) proteins. In the case of HCV, the generation of mature non-structural proteins (NS2, NS3, NS4A, NS4B, NS5A, and NS5B) is effected by two viral proteases. The first one is believed to be a metalloprotease and cleaves at the NS2-NS3 junction; the second one is a serine protease contained within the N-terminal region of NS3 (also referred to herein as NS3 protease) and mediates all the subsequent cleavages downstream of NS3, both in cis, at the NS3-NS4A cleavage site, and in trans, for the remaining NS4A-NS4B, NS4B-NS5A, NS5A-NS5B sites. The NS4A protein appears to serve multiple functions by both acting as a cofactor for the NS3 protease and assisting in the membrane localization of NS3 and other viral replicase components. The formation of a NS3-NS4A complex is necessary for proper protease activity resulting in increased proteolytic efficiency of the cleavage events. The NS3 protein also exhibits nucleoside triphosphatase and RNA helicase activities. NS5B (also referred to herein as HCV polymerase) is a RNA-dependent RNA polymerase that is involved in the replication of HCV with other HCV proteins, including NS5A, in a replicase complex.

[0008] Compounds useful for treating HCV-infected patients are desired which selectively inhibit HCV viral replication. In particular, compounds which are effective to inhibit the function of the NS5A protein are desired. The HCV NS5A protein is described, for example, in the following references: Tan, S.L. et al., *Virology*, 284:1-12 (2001); Park, K.-J. et al., *J. Biol. Chem.*, 307:11-30718 (2003); Tellinghuisen, T.L. et al., *Nature*, 435:374 (2005); Love, R.A. et al., *J. Virol.*, 83:4395 (2009); Appel, N. et al., *J. Biol. Chem.*, 281:9833 (2006); Huang, L., *J. Biol. Chem.*, 280:36417 (2005); Rice, C. et al., WO 2006/093867.

[0009] In one aspect the present disclosure provides a compound of Formula (I)



(I),

or a pharmaceutically acceptable salt thereof, wherein:

n is 0, 1, or 2;

X is selected from hydrogen, alkenyl, cyano, cycloalkyl, (cycloalkyl)alkyl, halo, and heterocyclyl;

5 R¹ is selected from hydrogen and halo;

R² is selected from hydrogen, alkenyl, cyano, cycloalkyl, (cycloalkyl)alkyl, halo, and heterocyclyl; or

R¹ and R², together with the carbon atoms to which they are attached, form a six-membered aromatic ring optionally substituted with one halo group;

10 provided that at least one of X and R² is selected from alkenyl, cyano, cycloalkyl, (cycloalkyl)alkyl, halo, and heterocyclyl;

each R³ is alkyl, wherein the alkyl can optionally form a fused three- or four-membered ring with an adjacent carbon atom or a spirocyclic three- or four-membered ring with the carbon atom to which it is attached; wherein the fused and spirocyclic rings

15 are optionally substituted with one or two alkyl groups;

each R⁴ is independently selected from hydrogen and -C(O)R⁵; and

each R⁵ is independently selected from alkoxy, alkyl, arylalkoxy, arylalkyl, cycloalkyl, heterocyclyl, heterocyclylalkyl, (NR^cR^d)alkenyl, and (NR^cR^d)alkyl.

20 **[0010]** In a first embodiment of the first aspect the present disclosure provides a compound of Formula (I), or a pharmaceutically acceptable salt thereof, wherein each R⁵ is independently selected from alkoxy, heterocyclyl, and (NR^cR^d)alkyl.

[0011] In a second embodiment of the first aspect the present disclosure provides a compound of Formula (I), or a pharmaceutically acceptable salt thereof, wherein X is
25 halo. In a third embodiment of the first aspect R² is halo.

[0012] In a fourth embodiment of the first aspect the present disclosure provides a compound of Formula (I), or a pharmaceutically acceptable salt thereof, wherein R¹ and R², together with the carbon atoms to which they are attached, form a six-membered aromatic ring optionally substituted with one halo group.

30 **[0013]** In a fifth embodiment of the first aspect the present disclosure provides a compound of Formula (I), or a pharmaceutically acceptable salt thereof, wherein X is hydrogen.

[0014] In a second aspect the present disclosure provides a composition comprising a compound of Formula (I), or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier. In a first embodiment of the second aspect the composition further comprises at least one additional compound having anti-HCV activity. In a second embodiment of the second aspect at least one of the additional compounds is an interferon or a ribavirin. In a third embodiment of the second aspect the interferon is selected from interferon alpha 2B, pegylated interferon alpha, consensus interferon, interferon alpha 2A, and lymphoblastoid interferon tau.

[0015] In a fourth embodiment of the second aspect the present disclosure provides a composition comprising a compound of Formula (I), or a pharmaceutically acceptable salt thereof, a pharmaceutically acceptable carrier, and at least one additional compound having anti-HCV activity, wherein at least one of the additional compounds is selected from interleukin 2, interleukin 6, interleukin 12, a compound that enhances the development of a type 1 helper T cell response, interfering RNA, anti-sense RNA, Imiqimod, ribavirin, an inosine 5'-monophosphate dehydrogenase inhibitor, amantadine, and rimantadine.

[0016] In a fifth embodiment of the second aspect the present disclosure provides a composition comprising a compound of Formula (I), or a pharmaceutically acceptable salt thereof, a pharmaceutically acceptable carrier, and at least one additional compound having anti-HCV activity, wherein at least one of the additional compounds is effective to inhibit the function of a target selected from HCV metalloprotease, HCV serine protease, HCV polymerase, HCV helicase, HCV NS4B protein, HCV entry, HCV assembly, HCV egress, HCV NS5A protein, and IMPDH for the treatment of an HCV infection.

[0017] In a third aspect the present disclosure provides a method of treating an HCV infection in a patient, comprising administering to the patient a therapeutically effective amount of a compound of Formula (I), or a pharmaceutically acceptable salt thereof. In a first embodiment of the third aspect the method further comprises administering at least one additional compound having anti-HCV activity prior to, after or simultaneously with the compound of Formula (I), or a pharmaceutically acceptable salt thereof. In a second embodiment of the third aspect at least one of the additional compounds is an interferon or a ribavirin. In a third embodiment of the third aspect the interferon is selected from

interferon alpha 2B, pegylated interferon alpha, consensus interferon, interferon alpha 2A, and lymphoblastoid interferon tau.

[0018] In a fourth embodiment of the third aspect the present disclosure provides a method of treating an HCV infection in a patient, comprising administering to the patient a therapeutically effective amount of a compound of Formula (I), or a pharmaceutically acceptable salt thereof, and at least one additional compound having anti-HCV activity prior to, after or simultaneously with the compound of Formula (I), or a pharmaceutically acceptable salt thereof, wherein at least one of the additional compounds is selected from interleukin 2, interleukin 6, interleukin 12, a compound that enhances the development of a type 1 helper T cell response, interfering RNA, anti-sense RNA, Imiquimod, ribavirin, an inosine 5'-monophosphate dehydrogenase inhibitor, amantadine, and rimantadine.

[0019] In a fifth embodiment of the third aspect the present disclosure provides a method of treating an HCV infection in a patient, comprising administering to the patient a therapeutically effective amount of a compound of Formula (I), or a pharmaceutically acceptable salt thereof, and at least one additional compound having anti-HCV activity prior to, after or simultaneously with the compound of Formula (I), or a pharmaceutically acceptable salt thereof, wherein at least one of the additional compounds is effective to inhibit the function of a target selected from HCV metalloprotease, HCV serine protease, HCV polymerase, HCV helicase, HCV NS4B protein, HCV entry, HCV assembly, HCV egress, HCV NS5A protein, and IMPDH for the treatment of an HCV infection.

[0020] Other aspects of the present disclosure may include suitable combinations of embodiments disclosed herein.

[0021] Yet other aspects and embodiments may be found in the description provided herein.

[0022] The description of the present disclosure herein should be construed in congruity with the laws and principals of chemical bonding. In some instances it may be necessary to remove a hydrogen atom in order to accommodate a substituent at any given location.

[0023] It should be understood that the compounds encompassed by the present disclosure are those that are suitably stable for use as pharmaceutical agent.

[0024] It is intended that the definition of any substituent or variable at a particular location in a molecule be independent of its definitions elsewhere in that molecule. For

example, when R¹ and R² both contain an R⁴ group, the two R⁴ groups may be the same or different.

[0025] All patents, patent applications, and literature references cited in the specification are herein incorporated by reference in their entirety. In the case of
5 inconsistencies, the present disclosure, including definitions, will prevail.

[0026] As used in the present specification, the following terms have the meanings indicated:

[0027] As used herein, the singular forms “a”, “an”, and “the” include plural reference unless the context clearly dictates otherwise.

10 [0028] Unless stated otherwise, all aryl, cycloalkyl, and heterocyclyl groups of the present disclosure may be substituted as described in each of their respective definitions. For example, the aryl part of an arylalkyl group may be substituted as described in the definition of the term “aryl”.

[0029] The term “alkenyl,” as used herein, refers to a straight or branched chain
15 group of two to six carbon atoms containing at least one carbon-carbon double bond.

[0030] The term “alkenyloxy,” as used herein, refers to an alkenyl group attached to the parent molecular moiety through an oxygen atom.

[0031] The term “alkenyloxy carbonyl,” as used herein, refers to an alkenyloxy group attached to the parent molecular moiety through a carbonyl group.


20 [0032] The term “alkoxy,” as used herein, refers to an alkyl group attached to the parent molecular moiety through an oxygen atom.

[0033] The term “alkoxyalkyl,” as used herein, refers to an alkyl group substituted with one, two, or three alkoxy groups.


[0034] The term “alkoxyalkyl carbonyl,” as used herein, refers to an alkoxyalkyl
25 group attached to the parent molecular moiety through a carbonyl group.

[0035] The term “alkoxycarbonyl,” as used herein, refers to an alkoxy group attached to the parent molecular moiety through a carbonyl group.

[0036] The term “alkyl,” as used herein, refers to a group derived from a straight or branched chain saturated hydrocarbon containing from one to six carbon atoms. In the
30 compounds of the present disclosure, when R³ is alkyl, the alkyl can optionally form a fused three- or four-membered ring with an adjacent carbon atom to provide the structure shown below:



wherein m is selected from 1 and 2, wherein z is 0, 1, or 2, and wherein R^s is alkyl; or wherein the alkyl can optionally form a spirocyclic three- or four-membered ring with the carbon atom to which it is attached to provide the structure shown below:



5

wherein m is selected from 1 and 2; wherein z is 0, 1, or 2, and wherein R^s is alkyl.

[0037] The term “alkylcarbonyl,” as used herein, refers to an alkyl group attached to the parent molecular moiety through a carbonyl group.

10 **[0038]** The term “alkylcarbonyloxy,” as used herein, refers to an alkylcarbonyl group attached to the parent molecular moiety through an oxygen atom.

[0039] The term “alkylsulfanyl,” as used herein, refers to an alkyl group attached to the parent molecular moiety through a sulfur atom.

15 **[0040]** The term “alkylsulfonyl,” as used herein, refers to an alkyl group attached to the parent molecular moiety through a sulfonyl group.

20 **[0041]** The term “aryl,” as used herein, refers to a phenyl group, or a bicyclic fused ring system wherein one or both of the rings is a phenyl group. Bicyclic fused ring systems consist of a phenyl group fused to a four- to six-membered aromatic or non-aromatic carbocyclic ring. The aryl groups of the present disclosure can be attached to the parent molecular moiety through any substitutable carbon atom in the group.

Representative examples of aryl groups include, but are not limited to, indanyl, indenyl, naphthyl, phenyl, and tetrahydronaphthyl. The aryl groups of the present disclosure are optionally substituted with one, two, three, four, or five substituents independently selected from alkenyl, alkoxy, alkoxyalkyl, alkoxycarbonyl, alkyl, alkylcarbonyl, a second aryl group, arylalkoxy, arylalkyl, arylcarbonyl, cyano, halo, haloalkoxy,

25

haloalkyl, heterocyclyl, heterocyclylalkyl, heterocyclylcarbonyl, hydroxy, hydroxyalkyl, nitro, $-NR^xR^y$, $(NR^xR^y)alkyl$, oxo, and $-P(O)OR_2$, wherein each R is independently selected from hydrogen and alkyl; and wherein the alkyl part of the arylalkyl and the heterocyclylalkyl are unsubstituted and wherein the second aryl group, the aryl part of the arylalkyl, the aryl part of the arylcarbonyl, the heterocyclyl, and the heterocyclyl part of the heterocyclylalkyl and the heterocyclylcarbonyl are further optionally substituted with one, two, or three substituents independently selected from alkoxy, alkyl, cyano, halo, haloalkoxy, haloalkyl, and nitro.

[0042] The term “arylalkoxy,” as used herein, refers to an aryl group attached to the parent molecular moiety through an alkoxy group.

[0043] The term “arylalkoxycarbonyl,” as used herein, refers to an arylalkoxy group attached to the parent molecular moiety through a carbonyl group.

[0044] The term “arylalkyl,” as used herein, refers to an alkyl group substituted with one, two, or three aryl groups. The alkyl part of the arylalkyl is further optionally substituted with one or two additional groups independently selected from alkoxy, alkylcarbonyloxy, halo, haloalkoxy, haloalkyl, heterocyclyl, hydroxy, and $-NR^cR^d$, wherein the heterocyclyl is further optionally substituted with one or two substituents independently selected from alkoxy, alkyl, unsubstituted aryl, unsubstituted arylalkoxy, unsubstituted arylalkoxycarbonyl, halo, haloalkoxy, haloalkyl, hydroxy, $-NR^xR^y$, and oxo.

[0045] The term “arylalkylcarbonyl,” as used herein, refers to an arylalkyl group attached to the parent molecular moiety through a carbonyl group.

[0046] The term “arylcarbonyl,” as used herein, refers to an aryl group attached to the parent molecular moiety through a carbonyl group.

[0047] The term “aryloxy,” as used herein, refers to an aryl group attached to the parent molecular moiety through an oxygen atom.

[0048] The term “aryloxycarbonyl,” as used herein, refers to an aryloxy group attached to the parent molecular moiety through a carbonyl group.

[0049] The term “arylsulfonyl,” as used herein, refers to an aryl group attached to the parent molecular moiety through a sulfonyl group.

[0050] The term “carbonyl,” as used herein, refers to $-C(O)-$.

[0051] The term “carboxy,” as used herein, refers to $-CO_2H$.

[0052] The term “cyano,” as used herein, refers to $-CN$.

- [0053] The term “cycloalkyl,” as used herein, refers to a saturated monocyclic, hydrocarbon ring system having three to seven carbon atoms and zero heteroatoms. Representative examples of cycloalkyl groups include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, and cyclohexyl. The cycloalkyl groups of the present disclosure are optionally substituted with one, two, three, four, or five substituents independently selected from alkoxy, alkyl, aryl, cyano, halo, haloalkoxy, haloalkyl, heterocyclyl, hydroxy, hydroxyalkyl, nitro, and -NR^xR^y, wherein the aryl and the heterocyclyl are further optionally substituted with one, two, or three substituents independently selected from alkoxy, alkyl, cyano, halo, haloalkoxy, haloalkyl, hydroxy, and nitro.
- 10 [0054] The term “(cycloalkyl)alkyl,” as used herein, refers to an alkyl group substituted with one, two, or three cycloalkyl groups.
- [0055] The term “cycloalkyloxy,” as used herein, refers to a cycloalkyl group attached to the parent molecular moiety through an oxygen atom.
- [0056] The term “cycloalkyloxycarbonyl,” as used herein, refers to a cycloalkyloxy group attached to the parent molecular moiety through a carbonyl group.
- 15 [0057] The term “cycloalkylsulfonyl,” as used herein, refers to a cycloalkyl group attached to the parent molecular moiety through a sulfonyl group.
- [0058] The term “formyl,” as used herein, refers to -CHO.
- [0059] The term “halo,” as used herein, refers to Cl, Br, F, or I.
- 20 [0060] The term “haloalkoxy,” as used herein, refers to a haloalkyl group attached to the parent molecular moiety through an oxygen atom.
- [0061] The term “haloalkoxycarbonyl,” as used herein, refers to a haloalkoxy group attached to the parent molecular moiety through a carbonyl group.
- [0062] The term “haloalkyl,” as used herein, refers to an alkyl group substituted with one, two, three, or four halogen atoms.
- 25 [0063] The term “heterocyclyl,” as used herein, refers to a four-, five-, six-, or seven-membered ring containing one, two, three, or four heteroatoms independently selected from nitrogen, oxygen, and sulfur. The four-membered ring has zero double bonds, the five-membered ring has zero to two double bonds, and the six- and seven-membered rings have zero to three double bonds. The term “heterocyclyl” also includes bicyclic groups in which the heterocyclyl ring is fused to another monocyclic heterocyclyl group, or a four- to six-membered aromatic or non-aromatic carbocyclic ring; as well as bridged
- 30

bicyclic groups such as 7-azabicyclo[2.2.1]hept-7-yl, 2-azabicyclo[2.2.2]oct-2-yl, 2,5-diazabicyclo[2.2.1]heptan-2-yl, and 2-azabicyclo[2.2.2]oct-3-yl. The heterocyclyl groups of the present disclosure can be attached to the parent molecular moiety through any carbon atom or nitrogen atom in the group. Examples of heterocyclyl groups include, but
5 are not limited to, benzothienyl, furyl, imidazolyl, indolinyl, indolyl, isoquinolinyl, isothiazolyl, isoxazolyl, morpholinyl, oxazolyl, oxetanyl, piperazinyl, piperidinyl, pyrazolyl, pyridinyl, pyrrolidinyl, pyrrolopyridinyl, pyrrolyl, quinolinyl, tetrahydrofuranyl, tetrahydropyranyl, thiazolyl, thienyl, and thiomorpholinyl. The heterocyclyl groups of the present disclosure are optionally substituted with one, two,
10 three, four, or five substituents independently selected from alkenyl, alkoxy, alkoxyalkyl, alkoxycarbonyl, alkyl, alkylcarbonyl, aryl, arylalkoxycarbonyl, arylalkyl, arylcarbonyl, cyano, halo, haloalkoxy, haloalkyl, a second heterocyclyl group, heterocyclylalkyl, heterocyclylcarbonyl, hydroxy, hydroxyalkyl, nitro, $-NR^xR^y$, $(NR^xR^y)alkyl$, and oxo, wherein the alkyl part of the arylalkyl and the heterocyclylalkyl are unsubstituted and
15 wherein the aryl, the aryl part of the arylalkyl, the aryl part of the arylcarbonyl, the second heterocyclyl group, and the heterocyclyl part of the heterocyclylalkyl and the heterocyclylcarbonyl are further optionally substituted with one, two, or three substituents independently selected from alkoxy, alkyl, cyano, halo, haloalkoxy, haloalkyl, and nitro.

[0064] The term “heterocyclylalkoxy,” as used herein, refers to a heterocyclyl group
20 attached to the parent molecular moiety through an alkoxy group.

[0065] The term “heterocyclylalkoxycarbonyl,” as used herein, refers to a heterocyclylalkoxy group attached to the parent molecular moiety through a carbonyl group.

[0066] The term “heterocyclylalkyl,” as used herein, refers to an alkyl group
25 substituted with one, two, or three heterocyclyl groups. The alkyl part of the heterocyclylalkyl is further optionally substituted with one or two additional groups independently selected from alkoxy, alkylcarbonyloxy, aryl, halo, haloalkoxy, haloalkyl, hydroxy, and $-NR^cR^d$, wherein the aryl is further optionally substituted with one or two substituents independently selected from alkoxy, alkyl, unsubstituted aryl, unsubstituted
30 arylalkoxy, unsubstituted arylalkoxycarbonyl, halo, haloalkoxy, haloalkyl, hydroxy, and $-NR^xR^y$.

[0067] The term “heterocyclalkylcarbonyl,” as used herein, refers to a heterocyclalkyl group attached to the parent molecular moiety through a carbonyl group.

[0068] The term “heterocyclcarbonyl,” as used herein, refers to a heterocycl group attached to the parent molecular moiety through a carbonyl group.

[0069] The term “heterocycloxy,” as used herein, refers to a heterocycl group attached to the parent molecular moiety through an oxygen atom.

[0070] The term “heterocycloxycarbonyl,” as used herein, refers to a heterocycloxy group attached to the parent molecular moiety through a carbonyl group.

[0071] The term “hydroxy,” as used herein, refers to -OH.

[0072] The term “hydroxyalkyl,” as used herein, refers to an alkyl group substituted with one, two, or three hydroxy groups.


[0073] The term “hydroxyalkylcarbonyl,” as used herein, refers to a hydroxyalkyl group attached to the parent molecular moiety through a carbonyl group.

[0074] The term “nitro,” as used herein, refers to -NO₂.

[0075] The term “-NR^cR^d,” as used herein, refers to two groups, R^c and R^d, which are attached to the parent molecular moiety through a nitrogen atom. R^c and R^d are independently selected from hydrogen, alkenyloxycarbonyl, alkoxyalkylcarbonyl, alkoxy carbonyl, alkyl, alkylcarbonyl, alkylsulfonyl, aryl, arylalkoxycarbonyl, arylalkyl, arylalkylcarbonyl, arylcarbonyl, aryloxycarbonyl, arylsulfonyl, cycloalkyl, cycloalkyloxy, cycloalkyloxycarbonyl, cycloalkylsulfonyl, formyl, haloalkoxycarbonyl, heterocycl, heterocyclalkoxycarbonyl, heterocyclalkyl, heterocyclalkylcarbonyl, heterocyclcarbonyl, heterocycloxycarbonyl, hydroxyalkylcarbonyl, (NR^eR^f)alkyl, (NR^eR^f)alkylcarbonyl, (NR^eR^f)carbonyl, (NR^eR^f)sulfonyl, -C(NCN)OR', and -C(NCN)NR^xR^y, wherein R' is selected from alkyl and unsubstituted phenyl, and wherein the alkyl part of the arylalkyl, the arylalkylcarbonyl, the heterocyclalkyl, and the heterocyclalkylcarbonyl are further optionally substituted with one -NR^eR^f group; and wherein the aryl, the aryl part of the arylalkoxycarbonyl, the arylalkyl, the arylalkylcarbonyl, the arylcarbonyl, the aryloxycarbonyl, and the arylsulfonyl, the heterocycl, and the heterocycl part of the heterocyclalkoxycarbonyl, the heterocyclalkyl, the heterocyclalkylcarbonyl, the heterocyclcarbonyl, and the heterocycloxycarbonyl are further optionally substituted with one, two, or three

substituents independently selected from alkoxy, alkyl, cyano, halo, haloalkoxy, haloalkyl, and nitro.

[0076] The term “(NR^cR^d)alkenyl,” as used herein, refers to



- 5 wherein R^c and R^d are as defined herein and each R^q is independently hydrogen or C₁₋₃ alkyl.

- [0077]** The term “(NR^cR^d)alkyl,” as used herein, refers to an alkyl group substituted with one, two, or three -NR^cR^d groups. The alkyl part of the (NR^cR^d)alkyl is further optionally substituted with one or two additional groups selected from alkoxy,
- 10 alkoxyalkylcarbonyl, alkoxycarbonyl, alkylsulfanyl, arylalkoxycarbonyl, carboxy, cycloalkyl, heterocyclyl, heterocyclylcarbonyl, hydroxy, and (NR^eR^f)carbonyl; wherein the heterocyclyl is further optionally substituted with one, two, three, four, or five substituents independently selected from alkoxy, alkyl, cyano, halo, haloalkoxy, haloalkyl, and nitro.
- 15 **[0078]** The term “-NR^eR^f,” as used herein, refers to two groups, R^e and R^f, which are attached to the parent molecular moiety through a nitrogen atom. R^e and R^f are independently selected from hydrogen, alkyl, unsubstituted aryl, unsubstituted arylalkyl, unsubstituted cycloalkyl, unsubstituted (cycloalkyl)alkyl, unsubstituted heterocyclyl, unsubstituted heterocyclylalkyl, (NR^xR^y)alkyl, and (NR^xR^y)carbonyl.
- 20 **[0079]** The term “(NR^eR^f)alkyl,” as used herein, refers to an alkyl group substituted with one, two, or three -NR^eR^f groups.
- [0080]** The term “(NR^eR^f)alkylcarbonyl,” as used herein, refers to an (NR^eR^f)alkyl group attached to the parent molecular moiety through a carbonyl group.
- [0081]** The term “(NR^eR^f)carbonyl,” as used herein, refers to an -NR^eR^f group
- 25 attached to the parent molecular moiety through a carbonyl group.
- [0082]** The term “(NR^eR^f)sulfonyl,” as used herein, refers to an -NR^eR^f group attached to the parent molecular moiety through a sulfonyl group.
- [0083]** The term “-NR^xR^y,” as used herein, refers to two groups, R^x and R^y, which are attached to the parent molecular moiety through a nitrogen atom. R^x and R^y are

independently selected from hydrogen, alkoxycarbonyl, alkyl, alkylcarbonyl, unsubstituted aryl, unsubstituted arylalkoxycarbonyl, unsubstituted arylalkyl, unsubstituted cycloalkyl, unsubstituted heterocyclyl, and $(\text{NR}^{\text{x}'}\text{R}^{\text{y}'})\text{carbonyl}$, wherein $\text{R}^{\text{x}'}$ and $\text{R}^{\text{y}'}$ are independently selected from hydrogen and alkyl.

5 **[0084]** The term “ $(\text{NR}^{\text{x}'}\text{R}^{\text{y}'})\text{alkyl}$,” as used herein, refers to an alkyl group substituted with one, two, or three $-\text{NR}^{\text{x}'}\text{R}^{\text{y}'}$ groups.

[0085] The term “ $(\text{NR}^{\text{x}'}\text{R}^{\text{y}'})\text{carbonyl}$,” as used herein, refers to an $-\text{NR}^{\text{x}'}\text{R}^{\text{y}'}$ group attached to the parent molecular moiety through a carbonyl group.

[0086] The term “ $-\text{NR}^{\text{x}'}\text{R}^{\text{y}'}$,” as used herein, refers to two groups, R^{x} and R^{y} , which are
10 attached to the parent molecular moiety through a nitrogen atom. R^{x} and R^{y} are independently selected from hydrogen and alkyl.

[0087] The term “ $(\text{NR}^{\text{x}'}\text{R}^{\text{y}'})\text{carbonyl}$,” as used herein, refers to an $-\text{NR}^{\text{x}'}\text{R}^{\text{y}'}$ group attached to the parent molecular moiety through a carbonyl group.

[0088] The term “oxo,” as used herein, refers to $=\text{O}$.

15 **[0089]** The term “sulfonyl,” as used herein, refers to $-\text{SO}_2-$.

[0090] Asymmetric centers exist in the compounds of the present disclosure. These centers are designated by the symbols “R” or “S”, depending on the configuration of substituents around the chiral carbon atom. It should be understood that the disclosure encompasses all stereochemical isomeric forms, or mixtures thereof, which possess the
20 ability to inhibit NS5A. Individual stereoisomers of compounds can be prepared synthetically from commercially available starting materials which contain chiral centers or by preparation of mixtures of enantiomeric products followed by separation such as conversion to a mixture of diastereomers followed by separation or recrystallization, chromatographic techniques, or direct separation of enantiomers on chiral
25 chromatographic columns. Starting compounds of particular stereochemistry are either commercially available or can be made and resolved by techniques known in the art.

[0091] Certain compounds of the present disclosure may also exist in different stable conformational forms which may be separable. Torsional asymmetry due to restricted rotation about an asymmetric single bond, for example because of steric hindrance or ring
30 strain, may permit separation of different conformers. The present disclosure includes each conformational isomer of these compounds and mixtures thereof.

[0092] The compounds of the present disclosure also exist as tautomers; therefore the present disclosure also encompasses all tautomeric forms.

[0093] The term “compounds of the present disclosure”, and equivalent expressions, are meant to embrace compounds of Formula (I), and pharmaceutically acceptable enantiomers, diastereomers, and salts thereof. Similarly, references to intermediates are meant to embrace their salts where the context so permits.

[0094] The present disclosure is intended to include all isotopes of atoms occurring in the present compounds. Isotopes include those atoms having the same atomic number but different mass numbers. By way of general example and without limitation, isotopes of hydrogen include deuterium and tritium. Isotopes of carbon include ^{13}C and ^{14}C . Isotopically-labeled compounds of the invention can generally be prepared by conventional techniques known to those skilled in the art or by processes analogous to those described herein, using an appropriate isotopically-labeled reagent in place of the non-labeled reagent otherwise employed. Such compounds may have a variety of potential uses, for example as standards and reagents in determining biological activity. In the case of stable isotopes, such compounds may have the potential to favorably modify biological, pharmacological, or pharmacokinetic properties.

[0095] The compounds of the present disclosure can exist as pharmaceutically acceptable salts. The term “pharmaceutically acceptable salt,” as used herein, represents salts or zwitterionic forms of the compounds of the present disclosure which are water or oil-soluble or dispersible, which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of patients without excessive toxicity, irritation, allergic response, or other problem or complication commensurate with a reasonable benefit/risk ratio, and are effective for their intended use. The salts can be prepared during the final isolation and purification of the compounds or separately by reacting a suitable nitrogen atom with a suitable acid. Representative acid addition salts include acetate, adipate, alginate, citrate, aspartate, benzoate, benzenesulfonate, bisulfate, butyrate, camphorate, camphorsulfonate; digluconate, glycerophosphate, hemisulfate, heptanoate, hexanoate, formate, fumarate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxyethanesulfonate, lactate, maleate, mesitylenesulfonate, methanesulfonate, naphthylenesulfonate, nicotinate, 2-naphthalenesulfonate, oxalate, palmoate, pectinate, persulfate, 3-phenylpropionate, picrate, pivalate, propionate, succinate, tartrate,

trichloroacetate, trifluoroacetate, phosphate, glutamate, bicarbonate, para-toluenesulfonate, and undecanoate. Examples of acids which can be employed to form pharmaceutically acceptable addition salts include inorganic acids such as hydrochloric, hydrobromic, sulfuric, and phosphoric, and organic acids such as oxalic, maleic, succinic, and citric.

[0096] Basic addition salts can be prepared during the final isolation and purification of the compounds by reacting a carboxy group with a suitable base such as the hydroxide, carbonate, or bicarbonate of a metal cation or with ammonia or an organic primary, secondary, or tertiary amine. The cations of pharmaceutically acceptable salts include lithium, sodium, potassium, calcium, magnesium, and aluminum, as well as nontoxic amine cations such as ammonium, tetramethylammonium, tetraethylammonium, methylamine, dimethylamine, trimethylamine, triethylamine, diethylamine, ethylamine, tributylamine, pyridine, N,N-dimethylaniline, N-methylpiperidine, N-methylmorpholine, dicyclohexylamine, procaine, dibenzylamine, N,N-dibenzylphenethylamine, and N,N'-dibenzylethylenediamine. Other representative organic amines useful for the formation of base addition salts include ethylenediamine, ethanolamine, diethanolamine, piperidine, and piperazine.

[0097] When it is possible that, for use in therapy, therapeutically effective amounts of a compound of Formula (I), as well as pharmaceutically acceptable salts thereof, may be administered as the raw chemical, it is possible to present the active ingredient as a pharmaceutical composition. Accordingly, the disclosure further provides pharmaceutical compositions, which include therapeutically effective amounts of compounds of Formula (I) or pharmaceutically acceptable salts thereof, and one or more pharmaceutically acceptable carriers, diluents, or excipients. The term "therapeutically effective amount," as used herein, refers to the total amount of each active component that is sufficient to show a meaningful patient benefit, *e.g.*, a sustained reduction in viral load. When applied to an individual active ingredient, administered alone, the term refers to that ingredient alone. When applied to a combination, the term refers to combined amounts of the active ingredients that result in the therapeutic effect, whether administered in combination, serially, or simultaneously. The compounds of Formula (I) and pharmaceutically acceptable salts thereof, are as described above. The carrier(s), diluent(s), or excipient(s) must be acceptable in the sense of being compatible with the

other ingredients of the formulation and not deleterious to the recipient thereof. In accordance with another aspect of the present disclosure there is also provided a process for the preparation of a pharmaceutical formulation including admixing a compound of Formula (I), or a pharmaceutically acceptable salt thereof, with one or more

5 pharmaceutically acceptable carriers, diluents, or excipients. The term “pharmaceutically acceptable,” as used herein, refers to those compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of patients without excessive toxicity, irritation, allergic response, or other problem or complication commensurate with a reasonable benefit/risk ratio, and
10 are effective for their intended use.

[0098] Pharmaceutical formulations may be presented in unit dose forms containing a predetermined amount of active ingredient per unit dose. Dosage levels of between about 0.01 and about 250 milligram per kilogram (“mg/kg”) body weight per day, preferably between about 0.05 and about 100 mg/kg body weight per day of the compounds of the
15 present disclosure are typical in a monotherapy for the prevention and treatment of HCV mediated disease. Typically, the pharmaceutical compositions of this disclosure will be administered from about 1 to about 5 times per day or alternatively, as a continuous infusion. Such administration can be used as a chronic or acute therapy. The amount of active ingredient that may be combined with the carrier materials to produce a single
20 dosage form will vary depending on the condition being treated, the severity of the condition, the time of administration, the route of administration, the rate of excretion of the compound employed, the duration of treatment, and the age, gender, weight, and condition of the patient. Preferred unit dosage formulations are those containing a daily dose or sub-dose, as herein above recited, or an appropriate fraction thereof, of an active
25 ingredient. Generally, treatment is initiated with small dosages substantially less than the optimum dose of the compound. Thereafter, the dosage is increased by small increments until the optimum effect under the circumstances is reached. In general, the compound is most desirably administered at a concentration level that will generally afford anti-virally effective results without causing any harmful or deleterious side effects.

30 [0099] When the compositions of this disclosure comprise a combination of a compound of the present disclosure and one or more additional therapeutic or prophylactic agent, both the compound and the additional agent are usually present at

dosage levels of between about 10 to 150%, and more preferably between about 10 and 80% of the dosage normally administered in a monotherapy regimen.

[00100] Pharmaceutical formulations may be adapted for administration by any appropriate route, for example by the oral (including buccal or sublingual), rectal, nasal, 5 topical (including buccal, sublingual, or transdermal), vaginal, or parenteral (including subcutaneous, intracutaneous, intramuscular, intra-articular, intrasynovial, intrasternal, intrathecal, intralesional, intravenous, or intradermal injections or infusions) route. Such formulations may be prepared by any method known in the art of pharmacy, for example by bringing into association the active ingredient with the carrier(s) or excipient(s). Oral 10 administration or administration by injection are preferred.

[00101] Pharmaceutical formulations adapted for oral administration may be presented as discrete units such as capsules or tablets; powders or granules; solutions or suspensions in aqueous or non-aqueous liquids; edible foams or whips; or oil-in-water liquid emulsions or water-in-oil emulsions.

15 [00102] For instance, for oral administration in the form of a tablet or capsule, the active drug component can be combined with an oral, non-toxic pharmaceutically acceptable inert carrier such as ethanol, glycerol, water, and the like. Powders are prepared by comminuting the compound to a suitable fine size and mixing with a similarly comminuted pharmaceutical carrier such as an edible carbohydrate, as, for 20 example, starch or mannitol. Flavoring, preservative, dispersing, and coloring agent can also be present.

[00103] Capsules are made by preparing a powder mixture, as described above, and filling formed gelatin sheaths. Glidants and lubricants such as colloidal silica, talc, magnesium stearate, calcium stearate, or solid polyethylene glycol can be added to the 25 powder mixture before the filling operation. A disintegrating or solubilizing agent such as agar-agar, calcium carbonate, or sodium carbonate can also be added to improve the availability of the medicament when the capsule is ingested.

[00104] Moreover, when desired or necessary, suitable binders, lubricants, disintegrating agents, and coloring agents can also be incorporated into the mixture. 30 Suitable binders include starch, gelatin, natural sugars such as glucose or beta-lactose, corn sweeteners, natural and synthetic gums such as acacia, tragacanth or sodium alginate, carboxymethylcellulose, polyethylene glycol, and the like. Lubricants used in

these dosage forms include sodium oleate, sodium chloride, and the like. Disintegrators include, without limitation, starch, methyl cellulose, agar, bentonite, xanthan gum, and the like. Tablets are formulated, for example, by preparing a powder mixture, granulating or slugging, adding a lubricant and disintegrant, and pressing into tablets. A powder

5 mixture is prepared by mixing the compound, suitably comminuted, with a diluent or base as described above, and optionally, with a binder such as carboxymethylcellulose, an algininate, gelatin, or polyvinyl pyrrolidone, a solution retardant such as paraffin, a resorption accelerator such as a quaternary salt and/or an absorption agent such as bentonite, kaolin, or dicalcium phosphate. The powder mixture can be granulated by

10 wetting with a binder such as syrup, starch paste, acacia mucilage, or solutions of cellulosic or polymeric materials and forcing through a screen. As an alternative to granulating, the powder mixture can be run through the tablet machine and the result is imperfectly formed slugs broken into granules. The granules can be lubricated to prevent sticking to the tablet forming dies by means of the addition of stearic acid, a stearate salt,

15 talc, or mineral oil. The lubricated mixture is then compressed into tablets. The compounds of the present disclosure can also be combined with a free flowing inert carrier and compressed into tablets directly without going through the granulating or slugging steps. A clear or opaque protective coating consisting of a sealing coat of shellac, a coating of sugar or polymeric material, and a polish coating of wax can be

20 provided. Dyestuffs can be added to these coatings to distinguish different unit dosages.

[00105] Oral fluids such as solution, syrups, and elixirs can be prepared in dosage unit form so that a given quantity contains a predetermined amount of the compound. Syrups can be prepared by dissolving the compound in a suitably flavored aqueous solution, while elixirs are prepared through the use of a non-toxic vehicle. Solubilizers and

25 emulsifiers such as ethoxylated isostearyl alcohols and polyoxyethylene sorbitol ethers, preservatives, flavor additive such as peppermint oil or natural sweeteners, or saccharin or other artificial sweeteners, and the like can also be added.

[00106] Where appropriate, dosage unit formulations for oral administration can be microencapsulated. The formulation can also be prepared to prolong or sustain the

30 release as for example by coating or embedding particulate material in polymers, wax, or the like.

[00107] The compounds of Formula (I), and pharmaceutically acceptable salts thereof, can also be administered in the form of liposome delivery systems, such as small unilamellar vesicles, large unilamellar vesicles, and multilamellar vesicles. Liposomes can be formed from a variety of phospholipids, such as cholesterol, stearylamine, or
5 phosphatidylcholines.

[00108] The compounds of Formula (I) and pharmaceutically acceptable salts thereof may also be delivered by the use of monoclonal antibodies as individual carriers to which the compound molecules are coupled. The compounds may also be coupled with soluble polymers as targetable drug carriers. Such polymers can include polyvinylpyrrolidone,
10 pyran copolymer, polyhydroxypropylmethacrylamidephenol, polyhydroxyethylaspartamidephenol, or polyethyleneoxidepolylysine substituted with palitoyl residues. Furthermore, the compounds may be coupled to a class of biodegradable polymers useful in achieving controlled release of a drug, for example, polylactic acid, polyepsilon caprolactone, polyhydroxy butyric acid, polyorthoesters,
15 polyacetals, polydihydropyrans, polycyanoacrylates, and cross-linked or amphipathic block copolymers of hydrogels.

[00109] Pharmaceutical formulations adapted for transdermal administration may be presented as discrete patches intended to remain in intimate contact with the epidermis of the recipient for a prolonged period of time. For example, the active ingredient may be
20 delivered from the patch by iontophoresis as generally described in *Pharmaceutical Research*, 3(6):318 (1986).

[00110] Pharmaceutical formulations adapted for topical administration may be formulated as ointments, creams, suspensions, lotions, powders, solutions, pastes, gels, sprays, aerosols, or oils.

25 [00111] For treatments of the eye or other external tissues, for example mouth and skin, the formulations are preferably applied as a topical ointment or cream. When formulated in an ointment, the active ingredient may be employed with either a paraffinic or a water-miscible ointment base. Alternatively, the active ingredient may be formulated in a cream with an oil-in-water cream base or a water-in oil base.

30 [00112] Pharmaceutical formulations adapted for topical administrations to the eye include eye drops wherein the active ingredient is dissolved or suspended in a suitable carrier, especially an aqueous solvent.

[00113] Pharmaceutical formulations adapted for topical administration in the mouth include lozenges, pastilles, and mouth washes.

[00114] Pharmaceutical formulations adapted for rectal administration may be presented as suppositories or as enemas.

- 5 [00115] Pharmaceutical formulations adapted for nasal administration wherein the carrier is a solid include a coarse powder having a particle size for example in the range 20 to 500 microns which is administered in the manner in which snuff is taken, *i.e.*, by rapid inhalation through the nasal passage from a container of the powder held close up to the nose. Suitable formulations wherein the carrier is a liquid, for administration as a
10 nasal spray or nasal drops, include aqueous or oil solutions of the active ingredient.

[00116] Pharmaceutical formulations adapted for administration by inhalation include fine particle dusts or mists, which may be generated by means of various types of metered, dose pressurized aerosols, nebulizers, or insufflators.

- [00117] Pharmaceutical formulations adapted for vaginal administration may be
15 presented as pessaries, tampons, creams, gels, pastes, foams, or spray formulations.

- [00118] Pharmaceutical formulations adapted for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats, and solutes which render the formulation isotonic with the blood of the intended recipient; and aqueous and non-aqueous sterile suspensions which may
20 include suspending agents and thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example sealed ampoules and vials, and may be stored in a freeze-dried (lyophilized) condition requiring only the addition of the sterile liquid carrier, for example water for injections, immediately prior to use.

- Extemporaneous injection solutions and suspensions may be prepared from sterile
25 powders, granules, and tablets.

[00119] It should be understood that in addition to the ingredients particularly mentioned above, the formulations may include other agents conventional in the art having regard to the type of formulation in question, for example those suitable for oral administration may include flavoring agents.

- 30 [00120] The term "patient" includes both human and other mammals.

[00121] The term "treating" refers to: (i) preventing a disease, disorder or condition from occurring in a patient that may be predisposed to the disease, disorder, and/or

condition but has not yet been diagnosed as having it; (ii) inhibiting the disease, disorder, or condition, *i.e.*, arresting its development; and (iii) relieving the disease, disorder, or condition, *i.e.*, causing regression of the disease, disorder, and/or condition.

[00122] The compounds of the present disclosure can also be administered with a
 5 cyclosporin, for example, cyclosporin A. Cyclosporin A has been shown to be active against HCV in clinical trials (*Hepatology*, 38:1282 (2003); *Biochem. Biophys. Res. Commun.*, 313:42 (2004); *J. Gastroenterol.*, 38:567 (2003)).

[00123] Table 1 below lists some illustrative examples of compounds that can be
 10 administered with the compounds of this disclosure. The compounds of the disclosure can be administered with other anti-HCV activity compounds in combination therapy, either jointly or separately, or by combining the compounds into a composition.

Table 1

Brand Name	Physiological Class	Type of Inhibitor or Target	Source Company
NIM811		Cyclophilin	Novartis
Debio-025		inhibitors	Debiopharm
Zadaxin		Immunomodulator	SciClone
Suvus		Methylene blue	Bioenvision
Actilon (CPG10101)		TLR9 agonist	Coley
Batabulin (T67)	Anticancer	β -Tubulin inhibitor	Tularik Inc., South San Francisco, CA
ISIS 14803	Antiviral	Antisense	ISIS Pharmaceuticals Inc, Carlsbad, CA / Elan Pharmaceuticals Inc., New York, NY
Summetrel	Antiviral	Antiviral	Endo Pharmaceuticals Holdings Inc., Chadds Ford, PA
GS-9132 (ACH-806)	Antiviral	HCV inhibitor	Achillion / Gilead

Brand Name	Physiological Class	Type of Inhibitor or Target	Source Company
Pyrazolopyrimidine compounds and salts From WO 2005/047288 26 May 2005	Antiviral	HCV inhibitors	Arrow Therapeutics Ltd.
Levovirin	Antiviral	IMPDH inhibitor	Ribapharm Inc., Costa Mesa, CA
Merimepodib (VX-497)	Antiviral	IMPDH inhibitor	Vertex Pharmaceuticals Inc., Cambridge, MA
XTL-6865 (XTL-002)	Antiviral	Monoclonal antibody	XTL Biopharmaceuticals Ltd., Rehovot, Israel
Telaprevir (VX-950, LY-570310)	Antiviral	NS3 serine protease inhibitor	Vertex Pharmaceuticals Inc., Cambridge, MA / Eli Lilly and Co. Inc., Indianapolis, IN
HCV-796	Antiviral	NS5B replicase inhibitor	Wyeth / Viropharma
NM-283	Antiviral	NS5B replicase inhibitor	Idenix / Novartis
GL-59728	Antiviral	NS5B replicase inhibitor	Gene Labs / Novartis
GL-60667	Antiviral	NS5B replicase inhibitor	Gene Labs / Novartis
2'C MeA	Antiviral	NS5B replicase inhibitor	Gilead
PSI 6130	Antiviral	NS5B replicase inhibitor	Roche

Brand Name	Physiological Class	Type of Inhibitor or Target	Source Company
R1626	Antiviral	NS5B replicase inhibitor	Roche
2'C Methyl adenosine	Antiviral	NS5B replicase inhibitor	Merck
JTK-003	Antiviral	RdRp inhibitor	Japan Tobacco Inc., Tokyo, Japan
Levovirin	Antiviral	Ribavirin	ICN Pharmaceuticals, Costa Mesa, CA
Ribavirin	Antiviral	Ribavirin	Schering-Plough Corporation, Kenilworth, NJ
Viramidine	Antiviral	Ribavirin prodrug	Ribapharm Inc., Costa Mesa, CA
Heptazyme	Antiviral	Ribozyme	Ribozyme Pharmaceuticals Inc., Boulder, CO
BILN-2061	Antiviral	Serine protease inhibitor	Boehringer Ingelheim Pharma KG, Ingelheim, Germany
SCH 503034	Antiviral	Serine protease inhibitor	Schering-Plough
Zadazim	Immune modulator	Immune modulator	SciClone Pharmaceuticals Inc., San Mateo, CA
Ceplene	Immunomodulator	Immune modulator	Maxim Pharmaceuticals Inc., San Diego, CA

Brand Name	Physiological Class	Type of Inhibitor or Target	Source Company
CELLCEPT®	Immunosuppressant	HCV IgG immuno-suppressant	F. Hoffmann-La Roche LTD, Basel, Switzerland
Civacir	Immunosuppressant	HCV IgG immuno-suppressant	Nabi Biopharmaceuticals Inc., Boca Raton, FL
Albuferon - α	Interferon	Albumin IFN- α 2b	Human Genome Sciences Inc., Rockville, MD
Infergen A	Interferon	IFN alfacon-1	InterMune Pharmaceuticals Inc., Brisbane, CA
Omega IFN	Interferon	IFN- ω	Intarcia Therapeutics
IFN- β and EMZ701	Interferon	IFN- β and EMZ701	Transition Therapeutics Inc., Ontario, Canada
REBIF®	Interferon	IFN- β 1a	Serono, Geneva, Switzerland
Roferon A	Interferon	IFN- α 2a	F. Hoffmann-La Roche Ltd., Basel, Switzerland
Intron A	Interferon	IFN- α 2b	Schering-Plough Corporation, Kenilworth, NJ

Brand Name	Physiological Class	Type of Inhibitor or Target	Source Company
Intron A and Zadaxin	Interferon	IFN- α 2b/ α 1-thymosin	RegeneRx Biopharma. Inc., Bethesda, MD / SciClone Pharmaceuticals Inc, San Mateo, CA
Rebetron	Interferon	IFN- α 2b / ribavirin	Schering-Plough Corporation, Kenilworth, NJ
Actimmune	Interferon	INF- γ	InterMune Inc., Brisbane, CA
Interferon- β	Interferon	Interferon- β -1a	Serono
Multiferon	Interferon	Long lasting IFN	Viragen / Valentis
Wellferon	Interferon	Lymphoblastoid IFN- α n1	GlaxoSmithKline plc, Uxbridge, UK
Omniferon	Interferon	natural IFN- α	Viragen Inc., Plantation, FL
Pegasys	Interferon	PEGylated IFN- α 2a	F. Hoffmann-La Roche LTD, Basel, Switzerland
Pegasys and Ceplene	Interferon	PEGylated IFN- α 2a / immune modulator	Maxim Pharmaceuticals Inc., San Diego, CA
Pegasys and Ribavirin	Interferon	PEGylated IFN- α 2a / ribavirin	F. Hoffmann-La Roche LTD, Basel, Switzerland
PEG-Intron	Interferon	PEGylated IFN- α 2b	Schering-Plough Corporation, Kenilworth, NJ

Brand Name	Physiological Class	Type of Inhibitor or Target	Source Company
PEG-Intron / Ribavirin	Interferon	PEGylated IFN- α 2b / ribavirin	Schering-Plough Corporation, Kenilworth, NJ
IP-501	Liver protection	Antifibrotic	Indevus Pharmaceuticals Inc., Lexington, MA
IDN-6556	Liver protection	Caspase inhibitor	Idun Pharmaceuticals Inc., San Diego, CA
ITMN-191 (R-7227)	Antiviral	Serine protease inhibitor	InterMune Pharmaceuticals Inc., Brisbane, CA
GL-59728	Antiviral	NS5B replicase inhibitor	Genelabs
ANA-971	Antiviral	TLR-7 agonist	Anadys
Boceprevir	Antiviral	Serine protease inhibitor	Schering-Plough
TMS-435	Antiviral	Serine protease inhibitor	Tibotec BVBA, Mechelen, Belgium
BI-201335	Antiviral	Serine protease inhibitor	Boehringer Ingelheim Pharma KG, Ingelheim, Germany
MK-7009	Antiviral	Serine protease inhibitor	Merck
PF-00868554	Antiviral	Replicase inhibitor	Pfizer
ANA598	Antiviral	Non-Nucleoside NS5B polymerase inhibitor	Anadys Pharmaceuticals, Inc., San Diego, CA, USA

Brand Name	Physiological Class	Type of Inhibitor or Target	Source Company
IDX375	Antiviral	Non-Nucleoside replicase inhibitor	Idenix Pharmaceuticals, Cambridge, MA, USA
BILB 1941	Antiviral	NS5B polymerase inhibitor	Boehringer Ingelheim Canada Ltd R&D, Laval, QC, Canada
PSI-7851	Antiviral	Nucleoside polymerase inhibitor	Pharmasset, Princeton, NJ, USA
VCH-759	Antiviral	NS5B polymerase inhibitor	Vir °Chem Pharma
VCH-916	Antiviral	NS5B polymerase inhibitor	Vir °Chem Pharma
GS-9190	Antiviral	NS5B polymerase inhibitor	Gilead
Peg-interferon lamda	Antiviral	Interferon	ZymoGenetics / Bristol-Myers Squibb

[00124] The compounds of the present disclosure may also be used as laboratory reagents. Compounds may be instrumental in providing research tools for designing of viral replication assays, validation of animal assay systems and structural biology studies to further enhance knowledge of the HCV disease mechanisms. Further, the compounds of the present disclosure are useful in establishing or determining the binding site of other antiviral compounds, for example, by competitive inhibition.

[00125] The compounds of this disclosure may also be used to treat or prevent viral contamination of materials and therefore reduce the risk of viral infection of laboratory or medical personnel or patients who come in contact with such materials, *e.g.*, blood, tissue,

surgical instruments and garments, laboratory instruments and garments, and blood collection or transfusion apparatuses and materials.

[00126] This disclosure is intended to encompass compounds having Formula (I) when prepared by synthetic processes or by metabolic processes including those occurring in the human or animal body (*in vivo*) or processes occurring *in vitro*.

[00127] The abbreviations used in the present application, including particularly in the illustrative examples which follow, are well-known to those skilled in the art. Some of the abbreviations used are as follows: ca. for about; min or mins for minutes; h or hr or hrs for hours; rt or RT for room temperature or retention time (context will dictate); R_t for

retention time; TFA for trifluoroacetic acid; DMSO for dimethylsulfoxide; Me for methyl; THF for tetrahydrofuran; *t*-Bu or *t*-Bu for *tert*-butyl; EDCI for 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride; DMAP for 4-dimethylaminopyridine; DBU for 1,8-diazabicycloundec-7-ene; Ph for phenyl; DEA for diethylamine; Et for ethyl; DMF for N,N-dimethylformamide; OAc for acetate; HMDS for hexamethyldisilane; pTsOH for para-toluenesulfonic acid; iPr₂EtN, DIEA, or DIPEA for diisopropylethylamine; EtOAc or EtOAc or EA for ethyl acetate; Et₃SiH for triethylsilane; MeOH for methanol; TMSCHN₂ for trimethylsilyldiazomethane; H-D-Ser-OBzl.HCl for D-serine benzyl ester hydrochloride; EtOH for ethanol; Me₂S for dimethylsulfide; TEA or Et₃N for triethylamine; LiHMDS for lithium

hexamethyldisilazide; DIBAL for diisobutylaluminum hydride; TBDMS-Cl for *tert*-butyldimethylsilyl chloride; *i*-PrOH for isopropanol; Boc, boc, or BOC for *tert*-butoxycarbonyl; Cbz-Cl for benzyl chloroformate; Bn for benzyl; DEAD for diethylazodicarboxylate; mCPBA for meta-chloroperoxybenzoic acid; DCM for dichloromethane; TMSCN for trimethylsilyl cyanide; ACN or ME CN for acetonitrile; dpppe for 1,5-bis(diphenylphosphino)pentane; TMEDA for tetramethylethylenediamine; DMA for N,N-dimethylacetamide; MeOD for CD₃OD; Hex for hexanes; NaOEt for sodium ethoxide; MTBE for methyl *tert* butyl ether; NCS for N-chlorosuccinimide; Et₂O for diethyl ether; DME for 1,2-dimethoxyethane; and EEDQ for N-ethoxycarbonyl-2-ethoxy-1,2-dihydroquinoline.

[00128] The present disclosure will now be described in connection with certain embodiments which are not intended to limit its scope. On the contrary, the present disclosure covers all alternatives, modifications, and equivalents as can be included

within the scope of the claims. Thus, the following examples, which include specific embodiments, will illustrate one practice of the present disclosure, it being understood that the examples are for the purposes of illustration of certain embodiments and are presented to provide what is believed to be the most useful and readily understood description of its procedures and conceptual aspects.

[00129] Starting materials can be obtained from commercial sources or prepared by well-established literature methods known to those of ordinary skill in the art.

SYNTHESIS OF COMMON CAPS

Compound Analysis Conditions

[00130] Purity assessment and low resolution mass analysis were conducted on a Shimadzu LC system coupled with Waters MICROMASS® ZQ MS system. It should be noted that retention times may vary slightly between machines. Additional LC conditions applicable to the current section, unless noted otherwise.

Cond.-MS-W1

Column = XTERRA® 3.0 X 50 mm S7

Start %B = 0

Final %B = 100

Gradient time = 2 min

Stop time = 3 min

Flow Rate = 5 mL/min

Wavelength = 220 nm

Solvent A = 0.1% TFA in 10% methanol/90% H₂O

Solvent B = 0.1% TFA in 90% methanol/10% H₂O

Cond.-MS-W2

Column = XTERRA® 3.0 X 50 mm S7

Start %B = 0

Final %B = 100

Gradient time = 3 min

Stop time = 4 min
Flow Rate = 4 mL/min
Wavelength = 220 nm
Solvent A = 0.1% TFA in 10% methanol/90% H₂O
5 Solvent B = 0.1% TFA in 90% methanol/10% H₂O

Cond.-MS-W5

Column = XTERRA® 3.0 X 50 mm S7
Start %B = 0
10 Final %B = 30
Gradient time = 2 min
Stop time = 3 min
Flow Rate = 5 mL/min
Wavelength = 220 nm
15 Solvent A = 0.1% TFA in 10% methanol/90% H₂O
Solvent B = 0.1% TFA in 90% methanol/10% H₂O

Cond.-D1

Column = XTERRA® C18 3.0 X 50 mm S7
20 Start %B = 0
Final %B = 100
Gradient time = 3 min
Stop time = 4 min
Flow Rate = 4 mL/min
25 Wavelength = 220 nm
Solvent A = 0.1% TFA in 10% methanol/90% H₂O
Solvent B = 0.1% TFA in 90% methanol/10% H₂O

Cond.-D2

30 Column = PHENOMENEX® Luna 4.6 X 50 mm S10
Start %B = 0
Final %B = 100

Gradient time = 3 min

Stop time = 4 min

Flow Rate = 4 mL/min

Wavelength = 220 nm

5 Solvent A = 0.1% TFA in 10% methanol/90% H₂O

Solvent B = 0.1% TFA in 90% methanol/10% H₂O

Cond.-M3

Column = XTERRA® C18 3.0 X 50 mm S7

10 Start %B = 0

Final %B = 40

Gradient time = 2 min

Stop time = 3 min

Flow Rate = 5 mL/min

15 Wavelength = 220 nm

Solvent A = 0.1% TFA in 10% methanol/90% H₂O

Solvent B = 0.1% TFA in 90% methanol/10% H₂O

Condition I

20 Column = PHENOMENEX® Luna 3.0 X 50 mm S10

Start %B = 0

Final %B = 100

Gradient time = 2 min

Stop time = 3 min

25 Flow Rate = 4 mL/min

Wavelength = 220 nm

Solvent A = 0.1% TFA in 10% methanol/90% H₂O

Solvent B = 0.1% TFA in 90% methanol/10% H₂O

30 Condition II

Column = PHENOMENEX® Luna 4.6 X 50 mm S10

Start %B = 0

Final %B = 100

Gradient time = 2 min

Stop time = 3 min

Flow Rate = 5 mL/min

5 Wavelength = 220 nm

Solvent A = 0.1% TFA in 10% methanol/90% H₂O

Solvent B = 0.1% TFA in 90% methanol/10% H₂O

Condition III

10 Column = XTERRA® C18 3.0 x 50mm S7

Start %B = 0

Final %B = 100

Gradient time = 3 min

Stop time = 4 min

15 Flow Rate = 4 mL/min


Wavelength = 220 nm

Solvent A = 0.1% TFA in 10% methanol/90% H₂O

Solvent B = 0.1% TFA in 90% methanol/10% H₂O

20

Cap-1




(R)-2-(Dimethylamino)-2-phenylacetic acid

[00131] A suspension of 10% Pd/C (2.0g) in methanol (10 mL) was added to a mixture of (R)-2-phenylglycine (10g, 66.2 mmol), formaldehyde (33 mL of 37% wt. in water),

25 1N HCl (30 mL) and methanol (30 mL), and exposed to H₂ (60 psi) for 3 hours. The reaction mixture was filtered through diatomaceous earth (CELITE®), and the filtrate was concentrated *in vacuo*. The resulting crude material was recrystallized from isopropanol to provide the HCl salt of Cap-1 as a white needle (4.0 g). Optical rotation: -117.1° [c=9.95 mg/mL in H₂O; λ=589 nm]. ¹H NMR (DMSO-d₆, δ=2.5 ppm, 500 MHz):
 30 δ 7.43-7.34 (m, 5H), 4.14 (s, 1H), 2.43 (s, 6H); LC (Condition I): RT=0.25; LC-MS:

Anal. Calcd. for $[M+H]^+$ $C_{10}H_{14}NO_2$ 180.10; found 180.17; HRMS: Anal. Calcd. for $[M+H]^+$ $C_{10}H_{14}NO_2$ 180.1025; found 180.1017.

Cap-2




5

(R)-2-(Diethylamino)-2-phenylacetic acid

[00132] $NaBH_3CN$ (6.22g, 94 mmol) was added in portions over a few minutes to a cooled (ice/water) mixture of (R)-2-Phenylglycine (6.02 g, 39.8 mmol) and methanol (100 mL), and stirred for 5 minutes. Acetaldehyde (10 mL) was added dropwise over 10 minutes and stirring was continued at the same cooled temperature for 45 minutes and at ambient temperature for ~6.5 hours. The reaction mixture was cooled back with ice-water bath, treated with water (3 mL) and then quenched with a dropwise addition of concentrated HCl over ~45 minutes until the pH of the mixture was ~1.5 - 2.0. The cooling bath was removed and the stirring was continued while adding concentrated HCl in order to maintain the pH of the mixture around 1.5-2.0. The reaction mixture was stirred overnight, filtered to remove the white suspension, and the filtrate was concentrated *in vacuo*. The crude material was recrystallized from ethanol to afford the HCl salt of Cap-2 as a shining white solid in two crops (crop-1: 4.16 g; crop-2: 2.19 g). 1H NMR (DMSO- d_6 , δ =2.5 ppm, 400 MHz): 10.44 (1.00, br s, 1H), 7.66 (m, 2H), 7.51 (m, 3H), 5.30 (s, 1H), 3.15 (br m, 2H), 2.98 (br m, 2H), 1.20 (app br s, 6H). Crop-1: $[\alpha]^{25}_{-102.21^\circ}$ ($c=0.357$, H_2O); crop-2: $[\alpha]^{25}_{-99.7^\circ}$ ($c=0.357$, H_2O). LC (Condition I): RT=0.43 min; LC-MS: Anal. Calcd. for $[M+H]^+$ $C_{12}H_{18}NO_2$: 208.13; found 208.26.

20

Cap-3




25

[00133] Acetaldehyde (5.0 mL, 89.1 mmol) and a suspension of 10% Pd/C (720 mg) in methanol/ H_2O (4mL/1 mL) was sequentially added to a cooled (~15 °C) mixture of (R)-

- 2-phenylglycine (3.096g, 20.48 mmol), 1N HCl (30 mL) and methanol (40 mL). The cooling bath was removed and the reaction mixture was stirred under a balloon of H₂ for 17 hours. An additional acetaldehyde (10 mL, 178.2 mmol) was added and stirring continued under H₂ atmosphere for 24 hours [Note: the supply of H₂ was replenished as needed throughout the reaction]. The reaction mixture was filtered through diatomaceous earth (CELITE®), and the filtrate was concentrated *in vacuo*. The resulting crude material was recrystallized from isopropanol to provide the HCl salt of (R)-2-(ethylamino)-2-phenylacetic acid as a shining white solid (2.846g). ¹H NMR (DMSO-d₆, δ=2.5 ppm, 400 MHz): δ 14.15 (br s, 1H), 9.55 (br s, 2H), 7.55-7.48 (m, 5H), 2.88 (br m, 1H), 2.73 (br m, 1H), 1.20 (app t, *J*=7.2, 3H). LC (Condition I): RT=0.39 min; >95 % homogeneity index; LC-MS: Anal. Calcd. for [M+H]⁺ C₁₀H₁₄NO₂: 180.10; found 180.18.
- [00134]** A suspension of 10% Pd/C (536 mg) in methanol/H₂O (3 mL/1 mL) was added to a mixture of (R)-2-(ethylamino)-2-phenylacetic acid/HCl (1.492g, 6.918 mmol), formaldehyde (20 mL of 37% wt. in water), 1N HCl (20 mL) and methanol (23 mL).
- The reaction mixture was stirred under a balloon of H₂ for ~72 hours, where the H₂ supply was replenished as needed. The reaction mixture was filtered through diatomaceous earth (CELITE®) and the filtrate was concentrated *in vacuo*. The resulting crude material was recrystallized from isopropanol (50 mL) to provide the HCl salt of Cap-3 as a white solid (985 mg). ¹H NMR (DMSO-d₆, δ=2.5 ppm, 400 MHz): δ 10.48 (br s, 1H), 7.59-7.51 (m, 5H), 5.26 (s, 1H), 3.08 (app br s, 2H), 2.65 (br s, 3H), 1.24 (br m, 3H). LC (Condition I): RT=0.39 min; >95 % homogeneity index; LC-MS: Anal. Calcd. for [M+H]⁺ C₁₁H₁₆NO₂: 194.12; found 194.18; HRMS: Anal. Calcd. for [M+H]⁺ C₁₁H₁₆NO₂: 194.1180; found 194.1181.

25

Cap-4




(R)-2-(Methoxycarbonylamino)-2-phenylacetic acid

[00135] ClCO₂Me (3.2 mL, 41.4 mmol) was added dropwise to a cooled (ice/water) THF (410 mL) semi-solution of (R)-*tert*-butyl 2-amino-2-phenylacetate/HCl (9.877 g,

- 40.52 mmol) and diisopropylethylamine (14.2 mL, 81.52 mmol) over 6 min, and stirred at similar temperature for 5.5 hours. The volatile component was removed *in vacuo*, and the residue was partitioned between water (100 mL) and ethyl acetate (200 mL). The organic layer was washed with 1N HCl (25 mL) and saturated NaHCO₃ solution (30 mL), dried
- 5 (MgSO₄), filtered, and concentrated *in vacuo*. The resultant colorless oil was triturated from hexanes, filtered and washed with hexanes (100 mL) to provide (R)-*tert*-butyl 2-(methoxycarbonylamino)-2-phenylacetate as a white solid (7.7 g). ¹H NMR (DMSO-d₆, δ=2.5 ppm, 400 MHz): 7.98 (d, *J*=8.0, 1H), 7.37-7.29 (m, 5H), 5.09 (d, *J*=8, 1H), 3.56 (s, 3H), 1.33 (s, 9H). LC (Condition I): RT=1.53 min; ~90 % homogeneity index; LC-MS:
- 10 Anal. Calcd. for [M+Na]⁺ C₁₄H₁₉NNaO₄: 288.12; found 288.15.
- [00136]** TFA (16 mL) was added dropwise to a cooled (ice/water) CH₂Cl₂ (160 mL) solution of the above product over 7 minutes, and the cooling bath was removed and the reaction mixture was stirred for 20 hours. Since the deprotection was still not complete, an additional TFA (1.0 mL) was added and stirring continued for an additional 2 hours.
- 15 The volatile component was removed *in vacuo*, and the resulting oil residue was treated with diethyl ether (15 mL) and hexanes (12 mL) to provide a precipitate. The precipitate was filtered and washed with diethyl ether/hexanes (~1:3 ratio; 30 mL) and dried *in vacuo* to provide Cap-4 as a fluffy white solid (5.57 g). Optical rotation: -176.9° [c=3.7 mg/mL in H₂O; λ=589 nm]. ¹H NMR (DMSO-d₆, δ=2.5 ppm, 400 MHz): δ 12.84 (br s,
- 20 1H), 7.96 (d, *J*=8.3, 1H), 7.41-7.29 (m, 5H), 5.14 (d, *J*=8.3, 1H), 3.55 (s, 3H). LC (Condition I): RT=1.01 min; >95 % homogeneity index; LC-MS: Anal. Calcd. for [M+H]⁺ C₁₀H₁₂NO₄ 210.08; found 210.17; HRMS: Anal. Calcd. for [M+H]⁺ C₁₀H₁₂NO₄ 210.0766; found 210.0756.

25

Cap-5




[00137] A mixture of (R)-2-phenylglycine (1.0 g, 6.62 mmol), 1,4-dibromobutane (1.57 g, 7.27 mmol) and Na₂CO₃ (2.10 g, 19.8 mmol) in ethanol (40 mL) was heated at 100 °C for 21 hours. The reaction mixture was cooled to ambient temperature and

filtered, and the filtrate was concentrated *in vacuo*. The residue was dissolved in ethanol and acidified with 1N HCl to pH 3-4, and the volatile component was removed *in vacuo*. The resulting crude material was purified by a reverse phase HPLC (water/methanol/TFA) to provide the TFA salt of Cap-5 as a semi-viscous white foam (1.0 g). ¹H NMR (DMSO-d₆, δ=2.5, 500 MHz) δ 10.68 (br s, 1H), 7.51 (m, 5H), 5.23 (s, 1H), 3.34 (app br s, 2H), 3.05 (app br s, 2H), 1.95 (app br s, 4H); RT=0.30 minutes (Condition I); >98% homogeneity index; LC-MS: Anal. Calcd. for [M+H]⁺ C₁₂H₁₆NO₂: 206.12; found 206.25.

10


Cap-6



[00138] The TFA salt of Cap-6 was synthesized from (R)-2-phenylglycine and 1-bromo-2-(2-bromoethoxy)ethane by using the method of preparation of Cap-5. ¹H NMR (DMSO-d₆, δ=2.5, 500 MHz) δ 12.20 (br s, 1H), 7.50 (m, 5H), 4.92 (s, 1H), 3.78 (app br s, 4H), 3.08 (app br s, 2H), 2.81 (app br s, 2H); RT=0.32 minutes (Condition I); >98%; LC-MS: Anal. Calcd. for [M+H]⁺ C₁₂H₁₆NO₃: 222.11; found 222.20; HRMS: Anal. Calcd. for [M+H]⁺ C₁₂H₁₆NO₃: 222.1130; found 222.1121.

15

Cap-7



Cap-7a: enantiomer-1
Cap-7b: enantiomer-2

20

[00139] A CH₂Cl₂ (200 mL) solution of p-toluenesulfonyl chloride (8.65 g, 45.4 mmol) was added dropwise to a cooled (-5 °C) CH₂Cl₂ (200 mL) solution of (S)-benzyl 2-hydroxy-2-phenylacetate (10.0 g, 41.3 mmol), triethylamine (5.75 mL, 41.3 mmol) and 4-dimethylaminopyridine (0.504 g, 4.13 mmol), while maintaining the temperature between -5 °C and 0 °C. The reaction was stirred at 0 °C for 9 hours, and then stored in a

25


freezer (-25 °C) for 14 hours. It was allowed to thaw to ambient temperature and washed with water (200 mL), 1N HCl (100 mL) and brine (100 mL), dried (MgSO₄), filtered, and concentrated *in vacuo* to provide benzyl 2-phenyl-2-(tosyloxy)acetate as a viscous oil which solidified upon standing (16.5 g). The chiral integrity of the product was not checked and that product was used for the next step without further purification. ¹H NMR (DMSO-d₆, δ=2.5, 500 MHz) δ 7.78 (d, *J*= 8.6, 2H), 7.43-7.29 (m, 10H), 7.20 (m, 2H), 6.12 (s, 1H), 5.16 (d, *J*=12.5, 1H), 5.10 (d, *J*=12.5, 1H), 2.39 (s, 3H). RT=3.00 (Condition III); >90% homogeneity index; LC-MS: Anal. Calcd. for [M+H]⁺ C₂₂H₂₀NaO₅S: 419.09; found 419.04.

- 10 **[00140]** A THF (75 mL) solution of benzyl 2-phenyl-2-(tosyloxy)acetate (6.0 g, 15.1 mmol), 1-methylpiperazine (3.36 mL, 30.3 mmol) and N,N-diisopropylethylamine (13.2 mL, 75.8 mmol) was heated at 65 °C for 7 hours. The reaction was allowed to cool to ambient temperature and the volatile component was removed *in vacuo*. The residue was partitioned between ethylacetate and water, and the organic layer was washed with water and brine, dried (MgSO₄), filtered, and concentrated *in vacuo*. The resulting crude material was purified by flash chromatography (silica gel, ethyl acetate) to provide benzyl 2-(4-methylpiperazin-1-yl)-2-phenylacetate as an orangish-brown viscous oil (4.56 g). Chiral HPLC analysis (CHIRALCEL® OD-H) indicated that the sample is a mixture of stereoisomers in a 38.2 to 58.7 ratio. The separation of the stereoisomers were effected as follow: the product was dissolved in 120 mL of ethanol/heptane (1:1) and injected (5 mL/injection) on chiral HPLC column (Chiracel OJ, 5 cm ID x 50 cm L, 20 μm) eluting with 85:15 Heptane/ethanol at 75 mL/min, and monitored at 220 nm. Stereoisomer-1 (1.474 g) and stereoisomer-2 (2.2149 g) were retrieved as viscous oil. ¹H NMR (CDCl₃, δ=7.26, 500 MHz) 7.44-7.40 (m, 2H), 7.33-7.24 (m, 6H), 7.21-7.16 (m, 2H), 5.13 (d, *J*=12.5, 1H), 5.08 (d, *J*=12.5, 1H), 4.02 (s, 1H), 2.65-2.38 (app br s, 8H), 2.25 (s, 3H). RT=2.10 (Condition III); >98% homogeneity index; LC-MS: Anal. Calcd. for [M+H]⁺ C₂₀H₂₅N₂O₂: 325.19; found 325.20.

- 30 **[00141]** A methanol (10 mL) solution of either stereoisomer of benzyl 2-(4-methylpiperazin-1-yl)-2-phenylacetate (1.0 g, 3.1 mmol) was added to a suspension of 10% Pd/C (120 mg) in methanol (5.0 mL). The reaction mixture was exposed to a balloon of hydrogen, under a careful monitoring, for <50 minutes. Immediately after the completion of the reaction, the catalyst was filtered through diatomaceous earth


(CELITE®) and the filtrate was concentrated *in vacuo* to provide Cap-7, contaminated with phenylacetic acid as a tan foam (867.6 mg; mass is above the theoretical yield). The product was used for the next step without further purification. ¹H NMR (DMSO-d₆, δ=2.5, 500 MHz) δ 7.44-7.37 (m, 2H), 7.37-7.24 (m, 3H), 3.92 (s, 1H), 2.63-2.48 (app. br s, 2H), 2.48-2.32 (m, 6H), 2.19 (s, 3H); RT=0.31 (Condition II); >90% homogeneity index; LC-MS: Anal. Calcd. for [M+H]⁺ C₁₃H₁₉N₂O₂: 235.14; found 235.15; HRMS: Anal. Calcd. for [M+H]⁺ C₁₃H₁₉N₂O₂: 235.1447; found 235.1440.

[00142] The synthesis of Cap-8 and Cap-9 was conducted according to the synthesis of Cap-7 by using appropriate amines for the S_N2 displacement step (*i.e.*, 4-hydroxypiperidine for Cap-8 and (S)-3-fluoropyrrolidine for Cap-9) and modified conditions for the separation of the respective stereoisomeric intermediates, as described below.




[00143] The stereoisomeric separation of the intermediate benzyl 2-(4-hydroxypiperidin-1-yl)-2-phenyl acetate was effected by employing the following conditions: the compound (500 mg) was dissolved in ethanol/heptane (5 mL/45 mL). The resulting solution was injected (5 mL/injection) on a chiral HPLC column (Chiracel OJ, 2 cm ID x 25 cm L, 10 μm) eluting with 80:20 heptane/ethanol at 10 mL/min, monitored at 220 nm, to provide 186.3 mg of stereoisomer-1 and 209.1 mg of stereoisomer-2 as light-yellow viscous oils. These benzyl ester was hydrogenolysed according to the preparation of Cap-7 to provide Cap-8: ¹H NMR (DMSO-d₆, δ=2.5, 500 MHz) 7.40 (d, J=7, 2H), 7.28-7.20 (m, 3H), 3.78 (s 1H), 3.46 (m, 1H), 2.93 (m, 1H), 2.62 (m, 1H), 2.20 (m, 2H), 1.70 (m, 2H), 1.42 (m, 2H). RT=0.28 (Condition II); >98% homogeneity index; LC-MS: Anal. Calcd. for [M+H]⁺ C₁₃H₁₈NO₃: 236.13; found 236.07; HRMS: Calcd. for [M+H]⁺ C₁₃H₁₈NO₃: 236.1287; found 236.1283.

Cap-9



- [00144] The diastereomeric separation of the intermediate benzyl 2-((S)-3-fluoropyrrolidin-1-yl)-2-phenylacetate was effected by employing the following
- 5 conditions: the ester (220 mg) was separated on a chiral HPLC column (Chiracel OJ-H, 0.46 cm ID x 25 cm L, 5 μ m) eluting with 95% CO₂ / 5% methanol with 0.1% TFA, at 10 bar pressure, 70 mL/min flow rate, and a temperature of 35 °C. The HPLC elute for the respective stereoisomers was concentrated, and the residue was dissolved in CH₂Cl₂ (20 mL) and washed with an aqueous medium (10 mL water + 1 mL saturated NaHCO₃
- 10 solution). The organic phase was dried (MgSO₄), filtered, and concentrated *in vacuo* to provide 92.5 mg of fraction-1 and 59.6 mg of fraction-2. These benzyl esters were hydrogenolysed according to the preparation of Cap-7 to prepare Cap-9a and Cap-9b.
- Cap-9a (diastereomer-1; the sample is a TFA salt as a result of purification on a reverse phase HPLC using H₂O/methanol/TFA solvent): ¹H NMR (DMSO-d₆, δ =2.5, 400 MHz) 7.55-7.48 (m, 5H), 5.38 (d of m, *J*=53.7, 1H), 5.09 (br s, 1H), 3.84-2.82 (br m, 4H), 2.31-2.09 (m, 2H). RT=0.42 (Condition I); >95% homogeneity index; LC-MS: Anal. Calcd. for [M+H]⁺ C₁₂H₁₅FNO₂: 224.11; found 224.14; Cap-9b (diastereomer-2): ¹H NMR (DMSO-d₆, δ =2.5, 400 MHz) 7.43-7.21 (m, 5H), 5.19 (d of m, *J*=55.9, 1H), 3.97 (s, 1H), 2.95-2.43 (m, 4H), 2.19-1.78 (m, 2H). RT=0.44 (Condition I); LC-MS: Anal. Calcd. for
- 20 [M+H]⁺ C₁₂H₁₅FNO₂: 224.11; found 224.14.

Cap-10




- [00145] To a solution of D-proline (2.0 g, 17 mmol) and formaldehyde (2.0 mL of
- 25 37% wt. in H₂O) in methanol (15 mL) was added a suspension of 10% Pd/C (500 mg) in methanol (5 mL). The mixture was stirred under a balloon of hydrogen for 23 hours. The reaction mixture was filtered through diatomaceous earth (CELITE®) and concentrated

in vacuo to provide Cap-10 as an off-white solid (2.15 g). ^1H NMR (DMSO- d_6 , δ =2.5, 500 MHz) 3.42 (m, 1H), 3.37 (dd, J =9.4, 6.1, 1H), 2.85-2.78 (m, 1H), 2.66 (s, 3H), 2.21-2.13 (m, 1H), 1.93-1.84 (m, 2H), 1.75-1.66 (m, 1H). RT=0.28 (Condition II); >98% homogeneity index; LC-MS: Anal. Calcd. for $[\text{M}+\text{H}]^+$ $\text{C}_6\text{H}_{12}\text{NO}_2$: 130.09; found 129.96.

5


Cap-11



[00146] A mixture of (2S,4R)-4-fluoropyrrolidine-2-carboxylic acid (0.50 g, 3.8 mmol), formaldehyde (0.5 mL of 37% wt. in H_2O), 12 N HCl (0.25 mL) and 10% Pd/C (50 mg) in methanol (20 mL) was stirred under a balloon of hydrogen for 19 hours. The reaction mixture was filtered through diatomaceous earth (CELITE®) and the filtrate was concentrated *in vacuo*. The residue was recrystallized from isopropanol to provide the HCl salt of Cap-11 as a white solid (337.7 mg). ^1H NMR (DMSO- d_6 , δ =2.5, 500 MHz) 5.39 (d m, J =53.7, 1H), 4.30 (m, 1H), 3.90 (ddd, J =31.5, 13.5, 4.5, 1H), 3.33 (dd, J =25.6, 13.4, 1H), 2.85 (s, 3H), 2.60-2.51 (m, 1H), 2.39-2.26 (m, 1H). RT=0.28 (Condition II); >98% homogeneity index; LC-MS: Anal. Calcd. for $[\text{M}+\text{H}]^+$ $\text{C}_6\text{H}_{11}\text{FNO}_2$: 148.08; found 148.06.

15

Cap-12 (same as Cap-52)



20

(S)-2-(Methoxycarbonylamino)propanoic acid


[00147] L-Alanine (2.0 g, 22.5 mmol) was dissolved in 10% aqueous sodium carbonate solution (50 mL), and a THF (50 mL) solution of methyl chloroformate (4.0 mL) was added to it. The reaction mixture was stirred under ambient conditions for 4.5 hours and concentrated *in vacuo*. The resulting white solid was dissolved in water and acidified with 1N HCl to a pH ~ 2-3. The resulting solutions was extracted with ethyl acetate (3 x 100 mL), and the combined organic phase was dried (Na_2SO_4), filtered, and concentrated *in vacuo* to provide a colorless oil (2.58 g). 500 mg of this material was

25

purified by a reverse phase HPLC (H₂O/methanol/TFA) to provide 150 mg of Cap-12 as a colorless oil. ¹H NMR (DMSO-d₆, δ=2.5, 500 MHz) 7.44 (d, *J*=7.3, 0.8H), 7.10 (br s, 0.2H), 3.97 (m, 1H), 3.53 (s, 3H), 1.25 (d, *J*=7.3, 3H).

5


Cap-13



[00148] A mixture of L-alanine (2.5 g, 28 mmol), formaldehyde (8.4 g, 37 wt. %), 1N HCl (30 mL) and 10% Pd/C (500 mg) in methanol (30 mL) was stirred under a hydrogen atmosphere (50 psi) for 5 hours. The reaction mixture was filtered through diatomaceous earth (CELITE®) and the filtrate was concentrated *in vacuo* to provide the HCl salt of Cap-13 as an oil which solidified upon standing under vacuum (4.4 g; the mass is above theoretical yield). The product was used without further purification. ¹H NMR (DMSO-d₆, δ=2.5, 500 MHz) δ 12.1 (br s, 1H), 4.06 (q, *J*=7.4, 1H), 2.76 (s, 6H), 1.46 (d, *J*=7.3, 3H).

15

Cap-14



(R)-2-Phenyl-2-(piperidin-1-yl)acetic acid

[00149] Step 1: A mixture of (R)-(-)-D-phenylglycine *tert*-butyl ester (3.00 g, 12.3 mmol), NaBH₃CN (0.773 g, 12.3 mmol), KOH (0.690 g, 12.3 mmol) and acetic acid (0.352 mL, 6.15 mmol) were stirred in methanol at 0 °C. To this mixture was added glutaric dialdehyde (2.23 mL, 12.3 mmol) dropwise over 5 minutes. The reaction mixture was stirred as it was allowed to warm to ambient temperature and stirring was continued at the same temperature for 16 hours. The solvent was subsequently removed and the residue was partitioned with 10% aqueous NaOH and ethyl acetate. The organic phase was separated, dried (MgSO₄), filtered and concentrated to dryness to provide a clear oil. This material was purified by reverse-phase preparative HPLC (Primesphere C-18, 30 x

100mm; CH₃CN-H₂O-0.1% TFA) to give the intermediate ester (2.70 g, 56%) as a clear oil. ¹H NMR (400 MHz, CDCl₃) δ 7.53-7.44 (m, 3H), 7.40-7.37 (m, 2H), 3.87 (d, *J*=10.9 Hz, 1H), 3.59 (d, *J*=10.9 Hz, 1H), 2.99 (t, *J*=11.2 Hz, 1H), 2.59 (t, *J*=11.4 Hz, 1H), 2.07-2.02 (m, 2H), 1.82 (d, *J*=1.82 Hz, 3H), 1.40 (s, 9H). LC-MS: Anal. Calcd. for

5 C₁₇H₂₅NO₂: 275; found: 276 (M+H)⁺.


[00150] Step 2: To a stirred solution of the intermediate ester (1.12g, 2.88mmol) in dichloromethane (10 mL) was added TFA (3 mL). The reaction mixture was stirred at ambient temperature for 4 hours and then it was concentrated to dryness to give a light yellow oil. The oil was purified using reverse-phase preparative HPLC (Primesphere C-

10 18, 30 x 100mm; CH₃CN-H₂O-0.1% TFA). The appropriate fractions were combined and concentrated to dryness *in vacuo*. The residue was then dissolved in a minimum amount of methanol and applied to applied to MCX LP extraction cartridges (2 x 6 g). The cartridges were rinsed with methanol (40 mL) and then the desired compound was eluted using 2M ammonia in methanol (50 mL). Product-containing fractions were

15 combined and concentrated and the residue was taken up in water. Lyophilization of this solution provided the title compound (0.492 g, 78%) as a light yellow solid. ¹H NMR (DMSO-d₆) δ 7.50 (s, 5H), 5.13 (s, 1H), 3.09 (br s, 2H), 2.92-2.89 (m, 2H), 1.74 (m, 4H), 1.48 (br s, 2H). LC-MS: Anal. Calcd. for C₁₃H₁₇NO₂: 219; found: 220 (M+H)⁺.

20

Cap-15




[00151] Step 1: (S)-1-Phenylethyl 2-bromo-2-phenylacetate. To a mixture of α-bromophenylacetic acid (10.75 g, 0.050 mol), (S)-(-)-1-phenylethanol (7.94 g, 0.065 mol) and DMAP (0.61 g, 5.0 mmol) in dry dichloromethane (100 mL) was added solid EDCI

25 (12.46 g, 0.065 mol) all at once. The resulting solution was stirred at room temperature under Ar for 18 hours and then it was diluted with ethyl acetate, washed (H₂O x 2, brine), dried (Na₂SO₄), filtered, and concentrated to give a pale yellow oil. Flash chromatography (SiO₂/ hexane-ethyl acetate, 4:1) of this oil provided the title compound

(11.64 g, 73%) as a white solid. ^1H NMR (400 MHz, CDCl_3) δ 7.53-7.17 (m, 10H), 5.95 (q, $J=6.6$ Hz, 0.5H), 5.94 (q, $J=6.6$ Hz, 0.5H), 5.41 (s, 0.5H), 5.39 (s, 0.5H), 1.58 (d, $J=6.6$ Hz, 1.5H), 1.51 (d, $J=6.6$ Hz, 1.5H).

[00152] Step 2: (S)-1-Phenylethyl (R)-2-(4-hydroxy-4-methylpiperidin-1-yl)-2-phenylacetate. To a solution of (S)-1-phenylethyl 2-bromo-2-phenylacetate (0.464 g, 1.45 mmol) in THF (8 mL) was added triethylamine (0.61 mL, 4.35 mmol), followed by tetrabutylammonium iodide (0.215 g, 0.58 mmol). The reaction mixture was stirred at room temperature for 5 minutes and then a solution of 4-methyl-4-hydroxypiperidine (0.251 g, 2.18 mmol) in THF (2 mL) was added. The mixture was stirred for 1 hour at room temperature and then it was heated at 55-60 °C (oil bath temperature) for 4 hours. The cooled reaction mixture was then diluted with ethyl acetate (30 mL), washed (H_2O x2, brine), dried (MgSO_4), filtered and concentrated. The residue was purified by silica gel chromatography (0-60% ethyl acetate-hexane) to provide first the (S,R)-isomer of the title compound (0.306 g, 60%) as a white solid and then the corresponding (S,S)-isomer (0.120 g, 23%), also as a white solid. (S,R)-isomer: ^1H NMR (CD_3OD) δ 7.51-7.45 (m, 2H), 7.41-7.25 (m, 8H), 5.85 (q, $J=6.6$ Hz, 1H), 4.05 (s, 1H), 2.56-2.45 (m, 2H), 2.41-2.29 (m, 2H), 1.71-1.49 (m, 4H), 1.38 (d, $J=6.6$ Hz, 3H), 1.18 (s, 3H). LC-MS: Anal. Calcd. for $\text{C}_{22}\text{H}_{27}\text{NO}_3$: 353; found: 354 ($\text{M}+\text{H}$) $^+$. (S,S)-isomer: ^1H NMR (CD_3OD) δ 7.41-7.30 (m, 5H), 7.20-7.14 (m, 3H), 7.06-7.00 (m, 2H), 5.85 (q, $J=6.6$ Hz, 1H), 4.06 (s, 1H), 2.70-2.60 (m, 1H), 2.51 (dt, $J=6.6, 3.3$ Hz, 1H), 2.44-2.31 (m, 2H), 1.75-1.65 (m, 1H), 1.65-1.54 (m, 3H), 1.50 (d, $J=6.8$ Hz, 3H), 1.20 (s, 3H). LC-MS: Anal. Calcd. for $\text{C}_{22}\text{H}_{27}\text{NO}_3$: 353; found: 354 ($\text{M}+\text{H}$) $^+$.

[00153] Step 3: (R)-2-(4-Hydroxy-4-methylpiperidin-1-yl)-2-phenylacetic acid. To a solution of (S)-1-phenylethyl (R)-2-(4-hydroxy-4-methylpiperidin-1-yl)-2-phenylacetate (0.185 g, 0.52 mmol) in dichloromethane (3 mL) was added trifluoroacetic acid (1 mL) and the mixture was stirred at room temperature for 2 hours. The volatiles were subsequently removed *in vacuo* and the residue was purified by reverse-phase preparative HPLC (Primesphere C-18, 20 x 100mm; CH_3CN - H_2O -0.1% TFA) to give the title compound (as TFA salt) as a pale bluish solid (0.128 g, 98%). LC-MS: Anal. Calcd. for $\text{C}_{14}\text{H}_{19}\text{NO}_3$: 249; found: 250 ($\text{M}+\text{H}$) $^+$.




[00154] Step 1: (S)-1-Phenylethyl 2-(2-fluorophenyl)acetate. A mixture of 2-fluorophenylacetic acid (5.45 g, 35.4 mmol), (S)-1-phenylethanol (5.62 g, 46.0 mmol), EDCI (8.82 g, 46.0 mmol) and DMAP (0.561 g, 4.60 mmol) in CH₂Cl₂ (100 mL) was stirred at room temperature for 12 hours. The solvent was then concentrated and the residue partitioned with H₂O-ethyl acetate. The phases were separated and the aqueous layer back-extracted with ethyl acetate (2x). The combined organic phases were washed (H₂O, brine), dried (Na₂SO₄), filtered, and concentrated *in vacuo*. The residue was purified by silica gel chromatography (BIOTAGE®/ 0-20% ethyl acetate-hexane) to provide the title compound as a colorless oil (8.38 g, 92%). ¹H NMR (400 MHz, CD₃OD) δ 7.32-7.23 (m, 7H), 7.10-7.04 (m, 2), 5.85 (q, *J*=6.5 Hz, 1H), 3.71 (s, 2H), 1.48 (d, *J*=6.5 Hz, 3H).

[00155] Step 2: (R)-((S)-1-Phenylethyl 2-(2-fluorophenyl)-2-(piperidin-1-yl)acetate. To a solution of (S)-1-phenylethyl 2-(2-fluorophenyl)acetate (5.00 g, 19.4 mmol) in THF (1200 mL) at 0 °C was added DBU (6.19 g, 40.7 mmol) and the solution was allowed to warm to room temperature while stirring for 30 minutes. The solution was then cooled to -78 °C and a solution of CBr₄ (13.5 g, 40.7 mmol) in THF (100 mL) was added and the mixture was allowed to warm to -10 °C and stirred at this temperature for 2 hours. The reaction mixture was quenched with saturated aq. NH₄Cl and the layers were separated. The aqueous layer was back-extracted with ethyl acetate (2x) and the combined organic phases were washed (H₂O, brine), dried (Na₂SO₄), filtered, and concentrated *in vacuo*. To the residue was added piperidine (5.73 mL, 58.1 mmol) and the solution was stirred at room temperature for 24 hours. The volatiles were then concentrated *in vacuo* and the residue was purified by silica gel chromatography (BIOTAGE®/ 0-30% diethyl ether-hexane) to provide a pure mixture of diastereomers (2:1 ratio by ¹H NMR) as a yellow oil (2.07 g, 31%), along with unreacted starting material (2.53 g, 51%). Further chromatography of the diastereomeric mixture (BIOTAGE®/ 0-10% diethyl ether-toluene) provided the title compound as a colorless oil (0.737 g, 11%). ¹H NMR (400

MHz, CD₃OD) δ 7.52 (ddd, $J=9.4, 7.6, 1.8$ Hz, 1H), 7.33 - 7.40 (m, 1), 7.23 - 7.23 (m, 4H), 7.02 - 7.23 (m, 4H), 5.86 (q, $J=6.6$ Hz, 1H), 4.45 (s, 1H), 2.39 - 2.45 (m, 4H), 1.52 - 1.58 (m, 4H), 1.40 - 1.42 (m, 1H), 1.38 (d, $J=6.6$ Hz, 3H). LC-MS: Anal. Calcd. for C₂₁H₂₄FNO₂: 341; found: 342 (M+H)⁺.

- 5 **[00156]** Step 3: (R)-2-(2-Fluorophenyl)-2-(piperidin-1-yl)acetic acid. A mixture of (R)-((S)-1-phenylethyl 2-(2-fluorophenyl)-2-(piperidin-1-yl)acetate (0.737 g, 2.16 mmol) and 20% Pd(OH)₂/C (0.070 g) in ethanol (30 mL) was hydrogenated at room temperature and atmospheric pressure (H₂ balloon) for 2 hours. The solution was then purged with Ar, filtered through diatomaceous earth (CELITE®), and concentrated *in vacuo*. This provided the title compound as a colorless solid (0.503 g, 98%). ¹H NMR (400 MHz, CD₃OD) δ 7.65 (ddd, $J=9.1, 7.6, 1.5$ Hz, 1H), 7.47-7.53 (m, 1H), 7.21-7.30 (m, 2H), 3.07-3.13 (m, 4H), 1.84 (br s, 4H), 1.62 (br s, 2H). LC-MS: Anal. Calcd. for C₁₃H₁₆FNO₂: 237; found: 238 (M+H)⁺.


15






- [00157]** Step 1: (S)-1-Phenylethyl (R)-2-(4-hydroxy-4-phenylpiperidin-1-yl)-2-phenylacetate. To a solution of (S)-1-phenylethyl 2-bromo-2-phenylacetate (1.50 g, 4.70 mmol) in THF (25 mL) was added triethylamine (1.31 mL, 9.42 mmol), followed by tetrabutylammonium iodide (0.347 g, 0.94 mmol). The reaction mixture was stirred at room temperature for 5 minutes and then a solution of 4-phenyl-4-hydroxypiperidine (1.00 g, 5.64 mmol) in THF (5 mL) was added. The mixture was stirred for 16 hours and then it was diluted with ethyl acetate (100 mL), washed (H₂O x2, brine), dried (MgSO₄), filtered and concentrated. The residue was purified on a silica gel column (0-60% ethyl acetate-hexane) to provide an approximately 2:1 mixture of diastereomers, as judged by ¹H NMR. Separation of these isomers was performed using supercritical fluid chromatography (CHIRALCEL® OJ-H, 30 x 250mm; 20% ethanol in CO₂ at 35 °C), to give first the (R)-isomer of the title compound (0.534 g, 27%) as a yellow oil and then the

- corresponding (S)-isomer (0.271 g, 14%), also as a yellow oil. (S,R)-isomer: ^1H NMR (400 MHz, CD_3OD) δ 7.55-7.47 (m, 4H), 7.44-7.25 (m, 10H), 7.25-7.17 (m, 1H), 5.88 (q, $J=6.6$ Hz, 1H), 4.12 (s, 1H), 2.82-2.72 (m, 1H), 2.64 (dt, $J=11.1$, 2.5 Hz, 1H), 2.58-2.52 (m, 1H), 2.40 (dt, $J=11.1$, 2.5 Hz, 1H), 2.20 (dt, $J=12.1$, 4.6 Hz, 1H), 2.10 (dt, $J=12.1$, 4.6 Hz, 1H), 1.72-1.57 (m, 2H), 1.53 (d, $J=6.5$ Hz, 3H). LC-MS: Anal. Calcd. for $\text{C}_{27}\text{H}_{29}\text{NO}_3$: 415; found: 416 ($\text{M}+\text{H}$) $^+$; (S,S)-isomer: ^1H NMR (400 MHz, CD_3OD) δ 7.55-7.48 (m, 2H), 7.45-7.39 (m, 2H), 7.38-7.30 (m, 5H), 7.25-7.13 (m, 4H), 7.08-7.00 (m, 2H), 5.88 (q, $J=6.6$ Hz, 1H), 4.12 (s, 1H), 2.95-2.85 (m, 1H), 2.68 (dt, $J=11.1$, 2.5 Hz, 1H), 2.57-2.52 (m, 1H), 2.42 (dt, $J=11.1$, 2.5 Hz, 1H), 2.25 (dt, $J=12.1$, 4.6 Hz, 1H), 2.12 (dt, $J=12.1$, 4.6 Hz, 1H), 1.73 (dd, $J=13.6$, 3.0 Hz, 1H), 1.64 (dd, $J=13.6$, 3.0 Hz, 1H), 1.40 (d, $J=6.6$ Hz, 3H). LC-MS: Anal. Calcd. for $\text{C}_{27}\text{H}_{29}\text{NO}_3$: 415; found: 416 ($\text{M}+\text{H}$) $^+$.

[00158] The following esters were prepared in similar fashion:

Intermediate-17a		<p>Diastereomer 1: ^1H NMR (500 MHz, $\text{DMSO}-d_6$) δ ppm 1.36 (d, $J=6.41$ Hz, 3H) 2.23-2.51 (m, 4H) 3.35 (s, 4H) 4.25 (s, 1H) 5.05 (s, 2H) 5.82 (d, $J=6.71$ Hz, 1H) 7.15-7.52 (m, 15H). LC-MS: Anal. Calcd. for: $\text{C}_{28}\text{H}_{30}\text{N}_2\text{O}_4$ 458.22; found: 459.44 ($\text{M}+\text{H}$)$^+$.</p> <p>Diastereomer 2: ^1H NMR (500 MHz, $\text{DMSO}-d_6$) δ ppm 1.45 (d, $J=6.71$ Hz, 3H) 2.27-2.44 (m, 4H) 3.39 (s, 4H) 4.23 (s, 1H) 5.06 (s, 2H) 5.83 (d, $J=6.71$ Hz, 1H) 7.12 (dd, $J=6.41$, 3.05 Hz, 2H) 7.19-7.27 (m, 3H) 7.27-7.44 (m, 10H). LC-MS: Anal. Calcd. for: $\text{C}_{28}\text{H}_{30}\text{N}_2\text{O}_4$ 458.22; found: 459.44 ($\text{M}+\text{H}$)$^+$.</p>
------------------	--	---

Intermediate -17b		Diastereomer 1: RT=11.76 minutes (Condition II); LC-MS: Anal. Calcd. for: $C_{20}H_{22}N_2O_3$ 338.16; found: 339.39 (M+H) ⁺ . Diastereomer 2: RT=10.05 minutes (Condition II). LC-MS: Anal. Calcd. for: $C_{20}H_{22}N_2O_3$ 338.16; found: 339.39 (M+H) ⁺ .
Intermediate -17c		Diastereomer 1: T _R =4.55 minutes (Condition I); LC-MS: Anal. Calcd. for: $C_{21}H_{26}N_2O_2$ 338.20; found: 339.45 (M+H) ⁺ . Diastereomer 2: T _R =6.00 minutes (Condition I). LC-MS: Anal. Calcd. for: $C_{21}H_{26}N_2O_2$ 338.20; found: 339.45 (M+H) ⁺ .
Intermediate -17d		Diastereomer 1: RT=7.19 minutes (Condition I); LC-MS: Anal. Calcd. for: $C_{27}H_{29}NO_2$ 399.22; found: 400.48 (M+H) ⁺ . Diastereomer 2: RT=9.76 minutes (Condition I); LC-MS: Anal. Calcd. for: $C_{27}H_{29}NO_2$ 399.22; found: 400.48 (M+H) ⁺ .

Chiral SFC Conditions for determining retention time:

Condition I

- 5 Column: CHIRALPAK® AD-H Column, 4.62x50 mm, 5μm
Solvents: 90% CO₂-10% methanol with 0.1%DEA
Temp: 35 °C
Pressure: 150 bar
Flow rate: 2.0 mL/min.
- 10 UV monitored at 220 nm
Injection: 1.0 mg/3mL methanol

Condition II

Column: CHIRALCEL® OD-H Column, 4.62x50 mm, 5µm

Solvents: 90% CO₂-10% methanol with 0.1%DEA

Temp: 35 °C

Pressure: 150 bar

5 Flow rate: 2.0 mL/min.

UV monitored at 220 nm



Injection: 1.0 mg/mL methanol



[00159] Cap-17, Step 2: (R)-2-(4-Hydroxy-4-phenylpiperidin-1-yl)-2-phenylacetic acid. To a solution of (S)-1-phenylethyl (R)-2-(4-hydroxy-4-phenylpiperidin-1-yl)-2-phenylacetate (0.350 g, 0.84 mmol) in dichloromethane (5 mL) was added trifluoroacetic acid (1 mL) and the mixture was stirred at room temperature for 2 hours. The volatiles were subsequently removed *in vacuo* and the residue was purified by reverse-phase preparative HPLC (Primesphere C-18, 20 x 100mm; CH₃CN-H₂O-0.1% TFA) to give the title compound (as TFA salt) as a white solid (0.230 g, 88%). LC-MS: Anal. Calcd. for C₁₉H₂₁NO₃: 311.15; found: 312 (M+H)⁺.

10
15

[00160] The following carboxylic acids were prepared in optically pure form in a similar fashion:

20

Cap-17a		RT=2.21 (Condition II); ¹ H NMR (500 MHz, DMSO-d ₆) δ ppm 2.20-2.35 (m, 2H) 2.34-2.47 (m, 2H) 3.37 (s, 4H) 3.71 (s, 1H) 5.06 (s, 2H) 7.06-7.53 (m, 10H). LC-MS: Anal. Calcd. for: C ₂₀ H ₂₂ N ₂ O ₄ 354.16; found: 355.38 (M+H) ⁺ .
Cap-17b		RT=0.27 (Condition III); LC-MS: Anal. Calcd. for: C ₁₂ H ₁₄ N ₂ O ₃ 234.10; found: 235.22 (M+H) ⁺ .

Cap-17c		RT=0.48 (Condition II); LC-MS: Anal. Calcd. for: $C_{13}H_{18}N_2O_2$ 234.14; found: 235.31 (M+H) ⁺ .
Cap-17d		RT=2.21 (Condition I); LC-MS: Anal. Calcd. for: $C_{19}H_{21}NO_2$ 295.16; found: 296.33 (M+H) ⁺ .

LC-MS Conditions for determining retention time:

Condition I

5 Column: PHENOMENEX® Luna 4.6 X 50 mm S10

Start % B=0

Final % B=100

Gradient Time=4 min

Flow Rate=4 mL/min

10 Wavelength=220

Solvent A=10% methanol - 90% H₂O - 0.1% TFA

Solvent B=90% methanol - 10% H₂O - 0.1% TFA

Condition II

15 Column: Waters SunFire 4.6 X 50 mm S5

Start % B=0

Final % B=100

Gradient Time=2 min

Flow Rate=4 mL/min

20 Wavelength=220

Solvent A=10% methanol - 90% H₂O - 0.1% TFA

Solvent B=90% methanol - 10% H₂O - 0.1% TFA

Condition III

Column: PHENOMENEX® 10 μ 3.0 X 50 mm

5 Start % B=0

Final % B=100


Gradient Time=2 min

Flow Rate=4 mL/min

Wavelength=220




10 Solvent A=10% methanol - 90% H₂O - 0.1% TFA











Solvent B=90% methanol - 10% H₂O - 0.1% TFA









- 15 **[00161]** Step 1: (R,S)-Ethyl 2-(4-pyridyl)-2-bromoacetate. To a solution of ethyl 4-pyridylacetate (1.00 g, 6.05 mmol) in dry THF (150 mL) at 0 °C under argon was added DBU (0.99 mL, 6.66 mmol). The reaction mixture was allowed to warm to room temperature over 30 minutes and then it was cooled to -78 °C. To this mixture was added CBr₄ (2.21 g, 6.66 mmol) and stirring was continued at -78 °C for 2 hours. The reaction
- 20 mixture was then quenched with sat. aq. NH₄Cl and the phases were separated. The organic phase was washed (brine), dried (Na₂SO₄), filtered, and concentrated *in vacuo*. The resulting yellow oil was immediately purified by flash chromatography (SiO₂/hexane-ethyl acetate, 1:1) to provide the title compound (1.40 g, 95%) as a somewhat unstable yellow oil. ¹H NMR (400 MHz, CDCl₃) δ 8.62 (dd, *J*=4.6, 1.8 Hz, 2H), 7.45
- 25 (dd, *J*=4.6, 1.8 Hz, 2H), 5.24 (s, 1H), 4.21-4.29 (m, 2H), 1.28 (t, *J*=7.1 Hz, 3H). LC-MS: Anal. Calcd. for C₉H₁₀BrNO₂: 242, 244; found: 243, 245 (M+H)⁺.

- [00162]** Step 2: (R,S)-Ethyl 2-(4-pyridyl)-2-(N,N-dimethylamino)acetate. To a solution of (R,S)-ethyl 2-(4-pyridyl)-2-bromoacetate (1.40 g, 8.48 mmol) in DMF (10 mL) at room temperature was added dimethylamine (2M in THF, 8.5 mL, 17.0 mmol). After completion of the reaction (as judged by thin layer chromatography) the volatiles were removed *in vacuo* and the residue was purified by flash chromatography (BIOTAGE®, 40+M SiO₂ column; 50%-100% ethyl acetate-hexane) to provide the title compound (0.539 g, 31%) as a light yellow oil. ¹H NMR (400 MHz, CDCl₃) δ 8.58 (d, *J*=6.0 Hz, 2H), 7.36 (d, *J*=6.0 Hz, 2H), 4.17 (m, 2H), 3.92 (s, 1H), 2.27 (s, 6H), 1.22 (t, *J*=7.0 Hz). LC-MS: Anal. Calcd. for C₁₁H₁₆N₂O₂: 208; found: 209 (M+H)⁺.
- [00163]** Step 3: (R,S)-2-(4-Pyridyl)-2-(N,N-dimethylamino)acetic acid. To a solution of (R,S)-ethyl 2-(4-pyridyl)-2-(N,N-dimethylamino)acetate (0.200 g, 0.960 mmol) in a mixture of THF-methanol-H₂O (1:1:1, 6 mL) was added powdered LiOH (0.120 g, 4.99 mmol) at room temperature. The solution was stirred for 3 hours and then it was acidified to pH 6 using 1N HCl. The aqueous phase was washed with ethyl acetate and then it was lyophilized to give the dihydrochloride of the title compound as a yellow solid (containing LiCl). The product was used as such in subsequent steps. ¹H NMR (400 MHz, DMSO-d₆) δ 8.49 (d, *J*=5.7 Hz, 2H), 7.34 (d, *J*=5.7 Hz, 2H), 3.56 (s, 1H), 2.21 (s, 6H).
- [00164]** The following examples were prepared in similar fashion using the method described above:

Cap-19		LC-MS: Anal. Calcd. for C ₉ H ₁₂ N ₂ O ₂ : 180; found: 181 (M+H) ⁺ .
Cap-20		LC-MS: no ionization. ¹ H NMR (400 MHz, CD ₃ OD) δ 8.55 (d, <i>J</i> =4.3 Hz, 1H), 7.84 (app t, <i>J</i> =5.3 Hz, 1H), 7.61 (d, <i>J</i> =7.8 Hz, 1H), 7.37 (app t, <i>J</i> =5.3 Hz, 1H), 4.35 (s, 1H), 2.60 (s, 6H).
Cap-21		LC-MS: Anal. Calcd. for C ₉ H ₁₁ ClN ₂ O ₂ : 214, 216; found: 215, 217 (M+H) ⁺ .

Cap-22		LC-MS: Anal. Calcd. for C ₁₀ H ₁₂ N ₂ O ₄ : 224; found: 225 (M+H) ⁺ .
Cap-23		LC-MS: Anal. Calcd. for C ₁₄ H ₁₅ NO ₂ : 229; found: 230 (M+H) ⁺ .
Cap-24		LC-MS: Anal. Calcd. for C ₁₁ H ₁₂ F ₃ NO ₂ : 247; found: 248 (M+H) ⁺ .
Cap-25		LC-MS: Anal. Calcd. for C ₁₁ H ₁₂ F ₃ NO ₂ : 247; found: 248 (M+H) ⁺ .
Cap-26		LC-MS: Anal. Calcd. for C ₁₀ H ₁₂ FNO ₂ : 197; found: 198 (M+H) ⁺ .
Cap-27		LC-MS: Anal. Calcd. for C ₁₀ H ₁₂ FNO ₂ : 247; found: 248 (M+H) ⁺ .
Cap-28		LC-MS: Anal. Calcd. for C ₁₀ H ₁₂ ClNO ₂ : 213; found: 214 (M+H) ⁺ .
Cap-29		LC-MS: Anal. Calcd. for C ₁₀ H ₁₂ ClNO ₂ : 213; found: 214 (M+H) ⁺ .
Cap-30		LC-MS: Anal. Calcd. for C ₁₀ H ₁₂ ClNO ₂ : 213; found: 214 (M+H) ⁺ .
Cap-31		LC-MS: Anal. Calcd. for C ₈ H ₁₂ N ₂ O ₂ S: 200; found: 201 (M+H) ⁺ .


Cap-32		LC-MS: Anal. Calcd. for C ₈ H ₁₁ NO ₂ S: 185; found: 186 (M+H) ⁺ .
Cap-33		LC-MS: Anal. Calcd. for C ₈ H ₁₁ NO ₂ S: 185; found: 186 (M+H) ⁺ .
Cap-34		LC-MS: Anal. Calcd. for C ₁₁ H ₁₂ N ₂ O ₃ : 220; found: 221 (M+H) ⁺ .
Cap-35		LC-MS: Anal. Calcd. for C ₁₂ H ₁₃ NO ₂ S: 235; found: 236 (M+H) ⁺ .
Cap-36		LC-MS: Anal. Calcd. for C ₁₂ H ₁₄ N ₂ O ₂ S: 250; found: 251 (M+H) ⁺ .



- [00165]** Step 1: (R,S)-Ethyl 2-(quinolin-3-yl)-2-(N,N-dimethylamino)-acetate. A mixture of ethyl N,N-dimethylaminoacetate (0.462 g, 3.54 mmol), K₃PO₄ (1.90 g, 8.95 mmol), Pd(t-Bu₃P)₂ (0.090 g, 0.176 mmol), 3-bromoquinoline and toluene (10 mL) was degassed with a stream of Ar bubbles for 15 minutes. The reaction mixture was then heated at 100 °C for 12 hours, after which it was cooled to room temperature and poured into H₂O. The mixture was extracted with ethyl acetate (2x) and the combined organic phases were washed (H₂O, brine), dried (Na₂SO₄), filtered, and concentrated *in vacuo*. The residue was purified first by reverse-phase preparative HPLC (Primesphere C-18, 30 x 100mm; CH₃CN-H₂O-5 mM NH₄OAc) and then by flash chromatography (SiO₂/hexane-ethyl acetate, 1:1) to provide the title compound (0.128 g, 17%) as an orange oil.

^1H NMR (400 MHz, CDCl_3) δ 8.90 (d, $J=2.0$ Hz, 1H), 8.32 (d, $J=2.0$ Hz, 1H), 8.03-8.01 (m, 2H), 7.77 (ddd, $J=8.3, 6.8, 1.5$ Hz, 1H), 7.62 (ddd, $J=8.3, 6.8, 1.5$ Hz, 1H), 4.35 (s, 1H), 4.13 (m, 2H), 2.22 (s, 6H), 1.15 (t, $J=7.0$ Hz, 3H). LC-MS: Anal. Calcd. for $\text{C}_{15}\text{H}_{18}\text{N}_2\text{O}_2$: 258; found: 259 ($\text{M}+\text{H}$) $^+$.

- 5 **[00166]** Step 2: (R,S) 2-(Quinolin-3-yl)-2-(N,N-dimethylamino)acetic acid. A mixture of (R,S)-ethyl 2-(quinolin-3-yl)-2-(N,N-dimethylamino)acetate (0.122 g, 0.472 mmol) and 6M HCl (3 mL) was heated at 100 °C for 12 hours. The solvent was removed *in vacuo* to provide the dihydrochloride of the title compound (0.169 g, >100%) as a light yellow foam. The unpurified material was used in subsequent steps without further
10 purification. LC-MS: Anal. Calcd. for $\text{C}_{13}\text{H}_{14}\text{N}_2\text{O}_2$: 230; found: 231 ($\text{M}+\text{H}$) $^+$.




- [00167]** Step 1: (R)-((S)-1-Phenylethyl) 2-(dimethylamino)-2-(2-fluorophenyl)acetate and (S)-((S)-1-Phenylethyl) 2-(dimethylamino)-2-(2-fluorophenyl)acetate. To a mixture
15 of (RS)-2-(dimethylamino)-2-(2-fluorophenyl)acetic acid (2.60 g, 13.19 mmol), DMAP (0.209 g, 1.71 mmol) and (S)-1-phenylethanol (2.09 g, 17.15 mmol) in CH_2Cl_2 (40 mL) was added EDCI (3.29 g, 17.15 mmol) and the mixture was allowed to stir at room temperature for 12 hours. The solvent was then removed *in vacuo* and the residue
20 partitioned with ethyl acetate- H_2O . The layers were separated, the aqueous layer was back-extracted with ethyl acetate (2x) and the combined organic phases were washed (H_2O , brine), dried (Na_2SO_4), filtered, and concentrated *in vacuo*. The residue was purified by silica gel chromatography (BIOTAGE®/ 0-50% diethyl ether-hexane). The resulting pure diastereomeric mixture was then separated by reverse-phase preparative
25 HPLC (Primesphere C-18, 30 x 100mm; $\text{CH}_3\text{CN}-\text{H}_2\text{O}-0.1\%$ TFA) to give first (S)-1-

phenethyl (R)-2-(dimethylamino)-2-(2-fluorophenyl)acetate (0.501 g, 13%) and then (S)-1-phenethyl (S)-2-(dimethylamino)-2-(2-fluorophenyl)-acetate (0.727 g, 18%), both as their TFA salts. (S,R)-isomer: ^1H NMR (400 MHz, CD_3OD) δ 7.65-7.70 (m, 1H), 7.55-7.60 (ddd, $J=9.4, 8.1, 1.5$ Hz, 1H), 7.36-7.41 (m, 2H), 7.28-7.34 (m, 5H), 6.04 (q, $J=6.5$ Hz, 1H), 5.60 (s, 1H), 2.84 (s, 6H), 1.43 (d, $J=6.5$ Hz, 3H). LC-MS: Anal. Calcd. for $\text{C}_{18}\text{H}_{20}\text{FNO}_2$: 301; found: 302 ($\text{M}+\text{H}$) $^+$; (S,S)-isomer: ^1H NMR (400 MHz, CD_3OD) δ 7.58-7.63 (m, 1H), 7.18-7.31 (m, 6H), 7.00 (dd, $J=8.5, 1.5$ Hz, 2H), 6.02 (q, $J=6.5$ Hz, 1H), 5.60 (s, 1H), 2.88 (s, 6H), 1.54 (d, $J=6.5$ Hz, 3H). LC-MS: Anal. Calcd. for $\text{C}_{18}\text{H}_{20}\text{FNO}_2$: 301; found: 302 ($\text{M}+\text{H}$) $^+$.

10 **[00168]** Step 2: (R)-2-(Dimethylamino)-2-(2-fluorophenyl)acetic acid. A mixture of (R)-((S)-1-phenylethyl) 2-(dimethylamino)-2-(2-fluorophenyl)acetate TFA salt (1.25 g, 3.01 mmol) and 20% $\text{Pd}(\text{OH})_2/\text{C}$ (0.125 g) in ethanol (30 mL) was hydrogenated at room temperature and atmospheric pressure (H_2 balloon) for 4 hours. The solution was then purged with Ar, filtered through diatomaceous earth (CELITE®), and concentrated *in vacuo*. This gave the title compound as a colorless solid (0.503 g, 98%). ^1H NMR (400 MHz, CD_3OD) δ 7.53-7.63 (m, 2H), 7.33-7.38 (m, 2H), 5.36 (s, 1H), 2.86 (s, 6H). LC-MS: Anal. Calcd. for $\text{C}_{10}\text{H}_{12}\text{FNO}_2$: 197; found: 198 ($\text{M}+\text{H}$) $^+$.

[00169] The S-isomer could be obtained from (S)-((S)-1-phenylethyl) 2-(dimethylamino)-2-(2-fluorophenyl)acetate TFA salt in similar fashion.

20




[00170] A mixture of (R)-2-(2-chlorophenyl)glycine (0.300 g, 1.62 mmol), formaldehyde (35% aqueous solution, 0.80 mL, 3.23 mmol) and 20% $\text{Pd}(\text{OH})_2/\text{C}$ (0.050 g) was hydrogenated at room temperature and atmospheric pressure (H_2 balloon) for 4 hours. The solution was then purged with Ar, filtered through diatomaceous earth (CELITE®) and concentrated *in vacuo*. The residue was purified by reverse-phase preparative HPLC (Primesphere C-18, 30 x 100mm; $\text{CH}_3\text{CN}-\text{H}_2\text{O}-0.1\%$ TFA) to give the TFA salt of the title compound (R)-2-(dimethylamino)-2-(2-chlorophenyl)acetic acid as a colorless oil

(0.290 g, 55%). ^1H NMR (400 MHz, CD_3OD) δ 7.59-7.65 (m, 2H), 7.45-7.53 (m, 2H), 5.40 (s, 1H), 2.87 (s, 6H). LC-MS: Anal. Calcd. for $\text{C}_{10}\text{H}_{12}\text{ClNO}_2$: 213; found: 214 ($\text{M}+\text{H}$) $^+$.

5


Cap-40



[00171] To an ice-cold solution of (R)-2-chlorophenylglycine (1.00 g, 5.38 mmol) and NaOH (0.862 g, 21.6 mmol) in H_2O (5.5 mL) was added methyl chloroformate (1.00 mL, 13.5 mmol) dropwise. The mixture was allowed to stir at 0 °C for 1 hour and then it was acidified by the addition of conc. HCl (2.5 mL). The mixture was extracted with ethyl acetate (2x) and the combined organic phase was washed (H_2O , brine), dried (Na_2SO_4), filtered, and concentrated *in vacuo* to give the title compound (R)-2-(methoxycarbonylamino)-2-(2-chlorophenyl)acetic acid as a yellow-orange foam (1.31 g, 96%). ^1H NMR (400 MHz, CD_3OD) δ 7.39 - 7.43 (m, 2H), 7.29 - 7.31 (m, 2H), 5.69 (s, 1H), 3.65 (s, 3H). LC-MS: Anal. Calcd. for $\text{C}_{10}\text{H}_{10}\text{ClNO}_4$: 243; found: 244 ($\text{M}+\text{H}$) $^+$.

15

Cap-41








[00172] To a suspension of 2-(2-(chloromethyl)phenyl)acetic acid (2.00 g, 10.8 mmol) in THF (20 mL) was added morpholine (1.89 g, 21.7 mmol) and the solution was stirred at room temperature for 3 hours. The reaction mixture was then diluted with ethyl acetate and extracted with H_2O (2x). The aqueous phase was lyophilized and the residue was purified by silica gel chromatography (BIOTAGE®/ 0-10% methanol- CH_2Cl_2) to give the title compound 2-(2-(morpholinomethyl)phenyl)acetic acid as a colorless solid (2.22 g,

20

87%). ^1H NMR (400 MHz, CD_3OD) δ 7.37-7.44 (m, 3H), 7.29-7.33 (m, 1H), 4.24 (s, 2H), 3.83 (br s, 4H), 3.68 (s, 2H), 3.14 (br s, 4H). LC-MS: Anal. Calcd. for $\text{C}_{13}\text{H}_{17}\text{NO}_3$: 235; found: 236 ($\text{M}+\text{H}$) $^+$.

- 5 [00173] The following examples were similarly prepared using the method described for Cap-41:

Cap-42		LC-MS: Anal. Calcd. for $\text{C}_{14}\text{H}_{19}\text{NO}_2$: 233; found: 234 ($\text{M}+\text{H}$) $^+$.
Cap-43		LC-MS: Anal. Calcd. for $\text{C}_{13}\text{H}_{17}\text{NO}_2$: 219; found: 220 ($\text{M}+\text{H}$) $^+$.
Cap-44		LC-MS: Anal. Calcd. for $\text{C}_{11}\text{H}_{15}\text{NO}_2$: 193; found: 194 ($\text{M}+\text{H}$) $^+$.
Cap-45		LC-MS: Anal. Calcd. for $\text{C}_{14}\text{H}_{20}\text{N}_2\text{O}_2$: 248; found: 249 ($\text{M}+\text{H}$) $^+$.




[00174] HMDS (1.85 mL, 8.77 mmol) was added to a suspension of (R)-2-amino-2-phenylacetic acid p-toluenesulfonate (2.83 g, 8.77 mmol) in CH₂Cl₂ (10 mL) and the mixture was stirred at room temperature for 30 minutes. Methyl isocyanate (0.5 g, 8.77 mmol) was added in one portion stirring continued for 30 minutes. The reaction was

5 quenched by addition of H₂O (5 mL) and the resulting precipitate was filtered, washed with H₂O and n-hexanes, and dried under vacuum. (R)-2-(3-methylureido)-2-phenylacetic acid (1.5 g; 82 %). was recovered as a white solid and it was used without further purification. ¹H NMR (500 MHz, DMSO-d₆) δ ppm 2.54 (d, *J*=4.88 Hz, 3H) 5.17 (d, *J*=7.93 Hz, 1H) 5.95 (q, *J*=4.48 Hz, 1H) 6.66 (d, *J*=7.93 Hz, 1H) 7.26-7.38 (m, 5H)

10 12.67 (s, 1H). LC-MS: Anal. Calcd. for C₁₀H₁₂N₂O₃ 208.08 found 209.121 (M+H)⁺; HPLC PHENOMENEX® C-18 3.0 × 46 mm, 0 to 100% B over 2 minutes, 1 minute hold time, A=90% water, 10% methanol, 0.1% TFA, B=10% water, 90% methanol, 0.1% TFA, RT=1.38 min, 90% homogeneity index.

15




[00175] The desired product was prepared according to the method described for Cap-45a. ¹H NMR (500 MHz, DMSO-d₆) δ ppm 0.96 (t, *J*=7.17 Hz, 3H) 2.94-3.05 (m, 2H) 5.17 (d, *J*=7.93 Hz, 1H) 6.05 (t, *J*=5.19 Hz, 1H) 6.60 (d, *J*=7.63 Hz, 1H) 7.26-7.38 (m, 5H) 12.68 (s, 1H). LC-MS: Anal. Calcd. for C₁₁H₁₄N₂O₃ 222.10 found 223.15 (M+H)⁺.

20 HPLC XTERRA® C-18 3.0 × 506 mm, 0 to 100% B over 2 minutes, 1 minute hold time, A=90% water, 10% methanol, 0.2% H₃PO₄, B=10% water, 90% methanol, 0.2% H₃PO₄, RT=0.87 min, 90% homogeneity index.


25

Cap-47



[00176] Step 1: (R)-*tert*-Butyl 2-(3,3-dimethylureido)-2-phenylacetate. To a stirred solution of (R)-*tert*-butyl-2-amino-2-phenylacetate (1.0 g, 4.10 mmol) and Hunig's base (1.79 mL, 10.25 mmol) in DMF (40 mL) was added dimethylcarbamoyl chloride (0.38 mL, 4.18 mmol) dropwise over 10 minutes. After stirring at room temperature for 3 hours, the reaction was concentrated under reduced pressure and the resulting residue was dissolved in ethyl acetate. The organic layer was washed with H₂O, 1N aq. HCl and brine, dried (MgSO₄), filtered and concentrated under reduced pressure. (R)-*tert*-butyl 2-(3,3-dimethylureido)-2-phenylacetate was obtained as a white solid (0.86 g; 75%) and used without further purification. ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 1.33 (s, 9H) 2.82 (s, 6H) 5.17 (d, *J*=7.63 Hz, 1H) 6.55 (d, *J*=7.32 Hz, 1H) 7.24-7.41 (m, 5H). LC-MS: Anal. Calcd. for C₁₅H₂₂N₂O₃ 278.16 found 279.23 (M+H)⁺; HPLC PHENOMENEX® Luna C-18 4.6 × 50 mm, 0 to 100% B over 4 minutes, 1 minute hold time, A=90% water, 10% methanol, 0.1% TFA, B=10% water, 90% methanol, 0.1% TFA, RT=2.26 min, 97% homogeneity index.

[00177] Step 2: (R)-2-(3,3-Dimethylureido)-2-phenylacetic acid. To a stirred solution of ((R)-*tert*-butyl 2-(3,3-dimethylureido)-2-phenylacetate (0.86 g, 3.10 mmol) in CH₂Cl₂ (250 mL) was added TFA (15 mL) dropwise and the resulting solution was stirred at rt for 3 hours. The desired compound was then precipitated out of solution with a mixture of EtOAc:Hexanes (5:20), filtered off and dried under reduced pressure. (R)-2-(3,3-dimethylureido)-2-phenylacetic acid was isolated as a white solid (0.59g, 86%) and used without further purification. ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 2.82 (s, 6H) 5.22 (d, *J*=7.32 Hz, 1H) 6.58 (d, *J*=7.32 Hz, 1H) 7.28 (t, *J*=7.17 Hz, 1H) 7.33 (t, *J*=7.32 Hz, 2H) 7.38-7.43 (m, 2H) 12.65 (s, 1H). LC-MS: Anal. Calcd. for C₁₁H₁₄N₂O₃: 222.24; found: 223.21 (M+H)⁺. HPLC XTERRA® C-18 3.0 × 50 mm, 0 to 100% B over 2 minutes, 1 minute hold time, A=90% water, 10% methanol, 0.2% H₃PO₄, B=10% water, 90% methanol, 0.2% H₃PO₄, RT=0.75 min, 93% homogeneity index.




[00178] Step 1: (R)-*tert*-Butyl 2-(3-cyclopentylureido)-2-phenylacetate. To a stirred solution of (R)-2-amino-2-phenylacetic acid hydrochloride (1.0 g, 4.10 mmol) and Hunig's base (1.0 mL, 6.15 mmol) in DMF (15 mL) was added cyclopentyl isocyanate (0.46 mL, 4.10 mmol) dropwise and over 10 minutes. After stirring at room temperature for 3 hours, the reaction was concentrated under reduced pressure and the resulting residue was taken up in ethyl acetate. The organic layer was washed with H₂O and brine, dried (MgSO₄), filtered, and concentrated under reduced pressure. (R)-*tert*-butyl 2-(3-cyclopentylureido)-2-phenylacetate was obtained as an opaque oil (1.32 g; 100 %) and used without further purification. ¹H NMR (500 MHz, CD₃Cl-D) δ ppm 1.50-1.57 (m, 2H) 1.58-1.66 (m, 2H) 1.87-1.97 (m, 2H) 3.89-3.98 (m, 1H) 5.37 (s, 1H) 7.26-7.38 (m, 5H). LC-MS: Anal. Calcd. for C₁₈H₂₆N₂O₃ 318.19 found 319.21 (M+H)⁺; HPLC XTERRA® C-18 3.0 × 50 mm, 0 to 100% B over 4 minutes, 1 minute hold time, A=90% water, 10% methanol, 0.1% TFA, B=10% water, 90% methanol, 0.1% TFA, RT=2.82 min, 96% homogeneity index.

[00179] Step 2: (R)-2-(3-Cyclopentylureido)-2-phenylacetic acid. To a stirred solution of (R)-*tert*-butyl 2-(3-cyclopentylureido)-2-phenylacetate (1.31 g, 4.10 mmol) in CH₂Cl₂ (25 mL) was added TFA (4 mL) and triethylsilane (1.64 mL; 10.3 mmol) dropwise, and the resulting solution was stirred at room temperature for 6 hours. The volatile components were removed under reduced pressure and the crude product was recrystallized in ethyl acetate/pentanes to yield (R)-2-(3-cyclopentylureido)-2-phenylacetic acid as a white solid (0.69 g, 64%). ¹H NMR (500 MHz, DMSO-d₆) δ ppm 1.17-1.35 (m, 2H) 1.42-1.52 (m, 2H) 1.53-1.64 (m, 2H) 1.67-1.80 (m, 2H) 3.75-3.89 (m, 1H) 5.17 (d, *J*=7.93 Hz, 1H) 6.12 (d, *J*=7.32 Hz, 1H) 6.48 (d, *J*=7.93 Hz, 1H) 7.24-7.40


(m, 5H) 12.73 (s, 1H). LC-MS: Anal. Calcd. for $C_{14}H_{18}N_2O_3$: 262.31; found: 263.15 (M+H)⁺. HPLC XTERRA® C-18 3.0 × 50 mm, 0 to 100% B over 2 minutes, 1 minute hold time, A=90% water, 10% methanol, 0.2% H_3PO_4 , B=10% water, 90% methanol, 0.2% H_3PO_4 , RT=1.24 min, 100% homogeneity index.

5



[00180] To a stirred solution of 2-(benzylamino)acetic acid (2.0 g, 12.1 mmol) in formic acid (91 mL) was added formaldehyde (6.94 mL, 93.2 mmol). After five hours at 70 °C, the reaction mixture was concentrated under reduced pressure to 20 mL and a white solid precipitated. Following filtration, the mother liquors were collected and further concentrated under reduced pressure providing the crude product. Purification by reverse-phase preparative HPLC (XTERRA® 30 X 100 mm, detection at 220 nm, flow rate 35 mL/min, 0 to 35% B over 8 min; A= 90% water, 10 % methanol, 0.1% TFA, B=10% water, 90 % methanol, 0.1% TFA) provided the title compound 2-(benzyl(methyl)-amino)acetic acid as its TFA salt (723 mg, 33%) as a colorless wax. ¹H NMR (300 MHz, DMSO-d₆) δ ppm 2.75 (s, 3H) 4.04 (s, 2H) 4.34 (s, 2H) 7.29-7.68 (m, 5H). LC-MS: Anal. Calcd. for: $C_{10}H_{13}NO_2$ 179.09; found: 180.20 (M+H)⁺.


20



[00181] To a stirred solution of 3-methyl-2-(methylamino)butanoic acid (0.50 g, 3.81 mmol) in water (30 mL) was added K_2CO_3 (2.63 g, 19.1 mmol) and benzyl chloride (1.32 g, 11.4 mmol). The reaction mixture was stirred at ambient temperature for 18 hours. The reaction mixture was extracted with ethyl acetate (30 mL x 2) and the aqueous layer was concentrated under reduced pressure providing the crude product which was purified

by reverse-phase preparative HPLC (XTERRA® 30 x 100mm, detection at 220 nm, flow rate 40 mL/min, 20 to 80% B over 6 min; A= 90% water, 10 % methanol, 0.1% TFA, B=10% water, 90 % methanol, 0.1% TFA) to provide 2-(benzyl(methyl)amino)-3-methylbutanoic acid, TFA salt (126 mg, 19%) as a colorless wax. ¹H NMR (500 MHz, DMSO-d₆) δ ppm 0.98 (d, 3H) 1.07 (d, 3H) 2.33-2.48 (m, 1H) 2.54-2.78 (m, 3H) 3.69 (s, 1H) 4.24 (s, 2H) 7.29-7.65 (m, 5H). LC-MS: Anal. Calcd. for: C₁₃H₁₉NO₂ 221.14; found: 222.28 (M+H)⁺.


Cap-51



(S)-2-(Methoxycarbonylamino)-3-methylbutanoic acid

[00182] Na₂CO₃ (1.83g, 17.2 mmol) was added to NaOH (33 mL of 1M/H₂O, 33 mmol) solution of L-valine (3.9 g, 33.29 mmol) and the resulting solution was cooled with ice-water bath. Methyl chloroformate (2.8 mL, 36.1 mmol) was added dropwise over 15 min, the cooling bath was removed and the reaction mixture was stirred at ambient temperature for 3.25 hr. The reaction mixture was washed with ether (50 mL, 3x), and the aqueous phase was cooled with ice-water bath and acidified with concentrated HCl to a pH region of 1-2, and extracted with CH₂Cl₂ (50 mL, 3x). The organic phase was dried (MgSO₄) and evaporated *in vacuo* to afford Cap-51 as a white solid (6 g). ¹H NMR for the dominant rotamer (DMSO-d₆, δ=2.5 ppm, 500 MHz): 12.54 (s, 1H), 7.33 (d, *J*=8.6, 1H), 3.84 (dd, *J*=8.4, 6.0, 1H), 3.54 (s, 3H), 2.03 (m, 1H), 0.87 (m, 6H). HRMS: Anal. Calcd. for [M+H]⁺ C₇H₁₄NO₄: 176.0923; found 176.0922.

Cap-51 (alternate route)




(S)-2-(Methoxycarbonylamino)-3-methylbutanoic acid

- [00183] DIEA (137.5 mL, 0.766 mol) was added to a suspension of (*S*)-*tert*-butyl 2-amino-3-methylbutanoate hydrochloride (75.0 g, 0.357 mol) in THF (900 mL), and the mixture was cooled to 0 °C (ice/water bath). Methyl chloroformate (29.0 mL, 0.375 mol) was added dropwise over 45 min, the cooling bath was removed and the heterogeneous mixture was stirred at ambient temperature for 3 h. The solvent was removed under diminished pressure and the residue partitioned between EtOAc and water (1 L each). The organic layer was washed with H₂O (1 L) and brine (1 L), dried (MgSO₄), filtered and concentrated under diminished pressure. The crude material was passed through a plug of silica gel (1 kg), eluting with hexanes (4 L) and 15:85 EtOAc/hexanes (4 L) to afford (*S*)-*tert*-butyl 2-(methoxycarbonylamino)-3-methylbutanoate as a clear oil (82.0 g, 99% yield). ¹H NMR (500 MHz, DMSO-d₆, δ = 2.5 ppm) 7.34 (d, *J* = 8.6, 1 H), 3.77 (dd, *J* = 8.6, 6.1, 1 H), 3.53 (s, 3 H), 1.94 - 2.05 (m, 1 H), 1.39 (s, 9 H), 0.83 - 0.92 (m, 6 H). ¹³C-NMR (126 MHz, DMSO-d₆, δ = 39.2 ppm) 170.92, 156.84, 80.38, 60.00, 51.34, 29.76, 27.62, 18.92, 17.95. LC-MS: [M+Na]⁺ 254.17.
- [00184] Trifluoroacetic acid (343 mL, 4.62 mol) and Et₃SiH (142 mL, 0.887 mol) were added sequentially to a solution of (*S*)-*tert*-butyl 2-(methoxycarbonylamino)-3-methylbutanoate (82.0 g, 0.355 mol) in CH₂Cl₂ (675 mL), and the mixture was stirred at ambient temperature for 4 h. The volatile component was removed under diminished pressure and the resultant oil triturated with petroleum ether (600 mL) to afford a white solid, which was filtered and washed with hexanes (500 mL) and petroleum ether (500 mL). Recrystallization from EtOAc/petroleum ether afforded Cap-51 as white flaky crystals (54.8 g, 88 % yield). MP = 108.5-109.5 °C. ¹H NMR (500 MHz, DMSO-d₆, δ = 2.5 ppm) 12.52 (s, 1 H), 7.31 (d, *J* = 8.6, 1 H), 3.83 (dd, *J* = 8.6, 6.1, 1 H), 3.53 (s, 3 H), 1.94 - 2.07 (m, 1 H), 0.86 (dd, *J* = 8.9, 7.0, 6 H). ¹³C NMR (126 MHz, DMSO-d₆, δ = 39.2 ppm) 173.30, 156.94, 59.48, 51.37, 29.52, 19.15, 17.98. LC-MS: [M+H]⁺ = 176.11. Anal. Calcd. for C₇H₁₃NO₄: C, 47.99; H, 7.48; N, 7.99. Found: C, 48.17; H, 7.55; N, 7.99. Optical Rotation: [α]_D = -4.16 (12.02 mg/mL; MeOH). Optical purity: >99.5 % ee. Note: the optical purity assessment was made on the methyl ester derivative of Cap-51, which was prepared under a standard TMSCHN₂ (benzene/MeOH) esterification protocol. HPLC analytical conditions: column, CHIRALPAK® AD-H (4.6 x 250mm, 5μm); solvent, 95% heptane / 5% IPA (isocratic); flow rate, 1 mL/min; temperature, 35 °C; UV monitored at 205 nm.

[Note: Cap-51 could also be purchased from Flamm.]

Cap-52 (same as Cap-12)



5 (S)-2-(Methoxycarbonylamino)propanoic acid

[00185] Cap-52 was synthesized from L-alanine according to the procedure described for the synthesis of Cap-51. For characterization purposes, a portion of the crude material was purified by a reverse phase HPLC (H₂O/methanol/TFA) to afford Cap-52 as a colorless viscous oil. ¹H NMR (DMSO-d₆, δ=2.5 ppm, 500 MHz): 12.49 (br s, 1H), 7.43 (d, *J*=7.3, 0.88H), 7.09 (app br s, 0.12H), 3.97 (m, 1H), 3.53 (s, 3H), 1.25 (d, *J*=7.3, 3H).

10





Cap-53 to Cap-64




[00186] Cap-53 to Cap-64 were prepared from appropriate starting materials according to the procedure described for the synthesis of Cap-51, with noted modifications if any.

15


Cap	Structure	Data
Cap-53a: (R) Cap-53b: (S) ((S)-2-(methoxy-carbonyl-amino)butanoic acid)		¹ H NMR (DMSO-d ₆ , δ = 2.5 ppm, 500 MHz): δ 12.51 (br s, 1H), 7.4 (d, <i>J</i> = 7.9, 0.9H), 7.06 (app s, 0.1H), 3.86-3.82 (m, 1H), 3.53 (s, 3H), 1.75-1.67 (m, 1H), 1.62-1.54 (m, 1H), 0.88 (d, <i>J</i> = 7.3, 3H). RT = 0.77 minutes (Cond. 2); LC-MS: Anal. Calcd. for [M+Na] ⁺ C ₆ H ₁₁ NNaO ₄ : 184.06; found 184.07. HRMS Calcd. for [M+Na] ⁺ C ₆ H ₁₁ NNaO ₄ : 184.0586; found 184.0592.

Cap	Structure	Data
Cap-54a: (R) Cap-54b: (S) ((S)-2-cyclopropyl-2-(methoxy-carbonyl-amino)acetic acid)		^1H NMR (DMSO- d_6 , δ = 2.5 ppm, 500 MHz): δ 12.48 (s, 1H), 7.58 (d, J = 7.6, 0.9H), 7.25 (app s, 0.1H), 3.52 (s, 3H), 3.36-3.33 (m, 1H), 1.10-1.01 (m, 1H), 0.54-0.49 (m, 1H), 0.46-0.40 (m, 1H), 0.39-0.35 (m, 1H), 0.31-0.21 (m, 1H). HRMS Calcd. for $[\text{M}+\text{H}]^+$ $\text{C}_7\text{H}_{12}\text{NO}_4$: 174.0766; found 174.0771
Cap-55		^1H NMR (DMSO- d_6 , δ = 2.5 ppm, 500 MHz): δ 12.62 (s, 1H), 7.42 (d, J = 8.2, 0.9H), 7.07 (app s, 0.1H), 5.80-5.72 (m, 1H), 5.10 (d, J = 17.1, 1H), 5.04 (d, J = 10.4, 1H), 4.01-3.96 (m, 1H), 3.53 (s, 3H), 2.47-2.42 (m, 1H), 2.35-2.29 (m, 1H).
Cap-56 (S)-3-methoxy-2-(methoxy-carbonyl-amino)propanoic acid		^1H NMR (DMSO- d_6 , δ = 2.5 ppm, 500 MHz): δ 12.75 (s, 1H), 7.38 (d, J = 8.3, 0.9H), 6.96 (app s, 0.1H), 4.20-4.16 (m, 1H), 3.60-3.55 (m, 2H), 3.54 (s, 3H), 3.24 (s, 3H).
Cap-57		^1H NMR (DMSO- d_6 , δ = 2.5 ppm, 500 MHz): δ 12.50 (s, 1H), 8.02 (d, J = 7.7, 0.08H), 7.40 (d, J = 7.9, 0.76H), 7.19 (d, J = 8.2, 0.07H), 7.07 (d, J = 6.7, 0.09H), 4.21-4.12 (m, 0.08H), 4.06-3.97 (m, 0.07H), 3.96-3.80 (m, 0.85H), 3.53 (s, 3H), 1.69-1.51 (m, 2H), 1.39-1.26 (m, 2H), 0.85 (t, J = 7.4, 3H). LC (Cond. 2): RT = 1.39 LC-MS: Anal. Calcd. for $[\text{M}+\text{H}]^+$ $\text{C}_7\text{H}_{14}\text{NO}_4$: 176.09; found 176.06.

Cap	Structure	Data
Cap-58		^1H NMR (DMSO- d_6 , δ = 2.5 ppm, 500 MHz): δ 12.63 (br s, 1H), 7.35 (s, 1H), 7.31 (d, J = 8.2, 1H), 6.92 (s, 1H), 4.33-4.29 (m, 1H), 3.54 (s, 3H), 2.54 (dd, J = 15.5, 5.4, 1H), 2.43 (dd, J = 15.6, 8.0, 1H). RT = 0.16 min (Cond. 2); LC-MS: Anal. Calcd. for $[\text{M}+\text{H}]^+$ $\text{C}_6\text{H}_{11}\text{N}_2\text{O}_5$: 191.07; found 191.14.
Cap-59a: (R) Cap-59b: (S)		^1H NMR (DMSO- d_6 , δ = 2.5 ppm, 400 MHz): δ 12.49 (br s, 1H), 7.40 (d, J = 7.3, 0.89H), 7.04 (br s, 0.11H), 4.00-3.95 (m, 3H), 1.24 (d, J = 7.3, 3H), 1.15 (t, J = 7.2, 3H). HRMS: Anal. Calcd. for $[\text{M}+\text{H}]^+$ $\text{C}_6\text{H}_{12}\text{NO}_4$: 162.0766; found 162.0771.
Cap-60		The crude material was purified with a reverse phase HPLC ($\text{H}_2\text{O}/\text{MeOH}/\text{TFA}$) to afford a colorless viscous oil that crystallized to a white solid upon exposure to high vacuum. ^1H NMR (DMSO- d_6 , δ = 2.5 ppm, 400 MHz): δ 12.38 (br s, 1H), 7.74 (s, 0.82H), 7.48 (s, 0.18H), 3.54/3.51 (two s, 3H), 1.30 (m, 2H), 0.98 (m, 2H). HRMS: Anal. Calcd. for $[\text{M}+\text{H}]^+$ $\text{C}_6\text{H}_{10}\text{NO}_4$: 160.0610; found 160.0604.
Cap-61		^1H NMR (DMSO- d_6 , δ = 2.5 ppm, 400 MHz): δ 12.27 (br s, 1H), 7.40 (br s, 1H), 3.50 (s, 3H), 1.32 (s, 6H). HRMS: Anal. Calcd. for $[\text{M}+\text{H}]^+$ $\text{C}_6\text{H}_{12}\text{NO}_4$: 162.0766; found 162.0765.

Cap	Structure	Data
Cap-62		^1H NMR (DMSO- d_6 , δ = 2.5 ppm, 400 MHz): δ 12.74 (br s, 1H), 4.21 (d, J = 10.3, 0.6H), 4.05 (d, J = 10.0, 0.4H), 3.62/3.60 (two singlets, 3H), 3.0 (s, 3H), 2.14-2.05 (m, 1H), 0.95 (d, J = 6.3, 3H), 0.81 (d, J = 6.6, 3H). LC-MS: Anal. Calcd. for $[\text{M-H}]^-$ $\text{C}_8\text{H}_{14}\text{NO}_4$: 188.09; found 188.05.
Cap-63		[Note: the reaction was allowed to run for longer than what was noted for the general procedure.] ^1H NMR (DMSO- d_6 , δ = 2.5 ppm, 400 MHz): 12.21 (br s, 1H), 7.42 (br s, 1H), 3.50 (s, 3H), 2.02-1.85 (m, 4H), 1.66-1.58 (m, 4H). LC-MS: Anal. Calcd. for $[\text{M+H}]^+$ $\text{C}_8\text{H}_{14}\text{NO}_4$: 188.09; found 188.19.
Cap-64		[Note: the reaction was allowed to run for longer than what was noted for the general procedure.] ^1H NMR (DMSO- d_6 , δ = 2.5 ppm, 400 MHz): 12.35 (br s, 1H), 7.77 (s, 0.82H), 7.56/7.52 (overlapping br s, 0.18H), 3.50 (s, 3H), 2.47-2.40 (m, 2H), 2.14-2.07 (m, 2H), 1.93-1.82 (m, 2H).

Cap-65




- [00187] Methyl chloroformate (0.65 mL, 8.39 mmol) was added dropwise over 5 min to a cooled (ice-water) mixture of Na_2CO_3 (0.449 g, 4.23 mmol), NaOH (8.2 mL of 1M/ H_2O , 8.2 mmol) and (*S*)-2-amino-3-hydroxy-3-methylbutanoic acid (1.04 g, 7.81

mmol). The reaction mixture was stirred for 45 min, and then the cooling bath was removed and stirring was continued for an additional 3.75 hr. The reaction mixture was washed with CH₂Cl₂, and the aqueous phase was cooled with ice-water bath and acidified with concentrated HCl to a pH region of 1-2. The volatile component was removed *in vacuo* and the residue was taken up in a 2:1 mixture of MeOH/CH₂Cl₂ (15 mL) and filtered, and the filtrate was rotovaped to afford Cap-65 as a white semi-viscous foam (1.236 g). ¹H NMR (DMSO-d₆, δ = 2.5 ppm, 400 MHz): δ 6.94 (d, *J* = 8.5, 0.9 H), 6.53 (br s, 0.1H), 3.89 (d, *J* = 8.8, 1H), 2.94 (s, 3H), 1.15 (s, 3H), 1.13 (s, 3H).


10 [00188] Cap-66 and Cap-67 were prepared from appropriate commercially available starting materials by employing the procedure described for the synthesis of Cap-65.

Cap-66




15 [00189] ¹H NMR (DMSO-d₆, δ = 2.5 ppm, 400 MHz): δ 12.58 (br s, 1H), 7.07 (d, *J* = 8.3, 0.13H), 6.81 (d, *J* = 8.8, 0.67H), 4.10-4.02 (m, 1.15H), 3.91 (dd, *J* = 9.1, 3.5, 0.85H), 3.56 (s, 3H), 1.09 (d, *J* = 6.2, 3H). [Note: only the dominant signals of NH were noted].

Cap-67



20 [00190] ¹H NMR (DMSO-d₆, δ = 2.5 ppm, 400 MHz): 12.51 (br s, 1H), 7.25 (d, *J* = 8.4, 0.75H), 7.12 (br d, *J* = 0.4, 0.05H), 6.86 (br s, 0.08H), 3.95-3.85 (m, 2H), 3.54 (s, 3H), 1.08 (d, *J* = 6.3, 3H). [Note: only the dominant signals of NH were noted].


25 Cap-68



[00191] Methyl chloroformate (0.38 ml, 4.9 mmol) was added drop-wise to a mixture of 1N NaOH (aq) (9.0 ml, 9.0 mmol), 1M NaHCO₃ (aq) (9.0 ml, 9.0 mol), L-aspartic acid β-benzyl ester (1.0 g, 4.5 mmol) and dioxane (9 ml). The reaction mixture was stirred at ambient conditions for 3 hr, and then washed with ethyl acetate (50 ml, 3x). The aqueous layer was acidified with 12N HCl to a pH ~ 1-2, and extracted with ethyl acetate (3 x 50 ml). The combined organic layers were washed with brine, dried (Na₂SO₄), filtered, and concentrated *in vacuo* to afford Cap-68 as a light yellow oil (1.37g; mass is above theoretical yield, and the product was used without further purification). ¹H NMR (DMSO-d₆, δ = 2.5 ppm, 500 MHz): δ 12.88 (br s, 1H), 7.55 (d, *J* = 8.5, 1H), 7.40-7.32 (m, 5H), 5.13 (d, *J* = 12.8, 1H), 5.10 (d, *J* = 12.9, 1H), 4.42-4.38 (m, 1H), 3.55 (s, 3H), 2.87 (dd, *J* = 16.2, 5.5, 1H), 2.71 (dd, *J* = 16.2, 8.3, 1H). LC (Cond. 2): RT = 1.90 min; LC-MS: Anal. Calcd. for [M+H]⁺ C₁₃H₁₆NO₆: 282.10; found 282.12.

15

Cap-69a and Cap-69b








Cap-69a: (*R*)-enantiomer
Cap-69b: (*S*)-enantiomer


[00192] NaCNBH₃ (2.416 g, 36.5 mmol) was added in batches to a chilled (~15 °C) water (17 mL)/MeOH (10 mL) solution of alanine (1.338 g, 15.0 mmol). A few minutes later acetaldehyde (4.0 mL, 71.3 mmol) was added drop-wise over 4 min, the cooling bath was removed, and the reaction mixture was stirred at ambient condition for 6 hr. An additional acetaldehyde (4.0 mL) was added and the reaction was stirred for 2 hr. Concentrated HCl was added slowly to the reaction mixture until the pH reached ~ 1.5, and the resulting mixture was heated for 1 hr at 40 °C. Most of the volatile component was removed *in vacuo* and the residue was purified with a DOWEX® 50WX8-100 ion-exchange resin (column was washed with water, and the compound was eluted with dilute NH₄OH, prepared by mixing 18 ml of NH₄OH and 282 ml of water) to afford Cap-69 (2.0 g) as an off-white soft hygroscopic solid. ¹H NMR (DMSO-d₆, δ = 2.5 ppm, 400 MHz): δ

3.44 (q, $J = 7.1$, 1H), 2.99-2.90 (m, 2H), 2.89-2.80 (m, 2H), 1.23 (d, $J = 7.1$, 3H), 1.13 (t, $J = 7.3$, 6H).


Cap-70 to Cap-74x

- 5 [00193] Cap-70 to Cap-74x were prepared according to the procedure described for the synthesis of Cap-69 by employing appropriate starting materials.

Cap-70a: (R) Cap-70b: (S)		^1H NMR (DMSO- d_6 , $\delta = 2.5$ ppm, 400 MHz): δ 3.42 (q, $J = 7.1$, 1H), 2.68-2.60 (m, 4H), 1.53-1.44 (m, 4H), 1.19 (d, $J = 7.3$, 3H), 0.85 (t, $J = 7.5$, 6H). LC-MS: Anal. Calcd. for $[\text{M}+\text{H}]^+$ $\text{C}_9\text{H}_{20}\text{NO}_2$: 174.15; found 174.13.
Cap-71a: (R) Cap-71b: (S)		^1H NMR (DMSO- d_6 , $\delta = 2.5$ ppm, 500 MHz): δ 3.18-3.14 (m, 1H), 2.84-2.77 (m, 2H), 2.76-2.68 (m, 2H), 1.69-1.54 (m, 2H), 1.05 (t, $J = 7.2$, 6H), 0.91 (t, $J = 7.3$, 3H). LC-MS: Anal. Calcd. for $[\text{M}+\text{H}]^+$ $\text{C}_8\text{H}_{18}\text{NO}_2$: 160.13; found 160.06.
Cap-72		^1H NMR (DMSO- d_6 , $\delta = 2.5$ ppm, 400 MHz): δ 2.77-2.66 (m, 3H), 2.39-2.31 (m, 2H), 1.94-1.85 (m, 1H), 0.98 (t, $J = 7.1$, 6H), 0.91 (d, $J = 6.5$, 3H), 0.85 (d, $J = 6.5$, 3H). LC-MS: Anal. Calcd. for $[\text{M}+\text{H}]^+$ $\text{C}_9\text{H}_{20}\text{NO}_2$: 174.15; found 174.15.
Cap-73		^1H NMR (DMSO- d_6 , $\delta = 2.5$ ppm, 500 MHz): δ 9.5 (br s, 1H), 3.77 (dd, $J = 10.8, 4.1$, 1H), 3.69-3.61 (m, 2H), 3.26 (s, 3H), 2.99-2.88 (m, 4H), 1.13 (t, $J = 7.2$, 6H).
Cap-74		^1H NMR (DMSO- d_6 , $\delta = 2.5$ ppm, 500 MHz): δ 7.54 (s, 1H), 6.89 (s, 1H), 3.81 (t, $J = 6.6$, 1H), 2.82-2.71 (m, 4H), 2.63 (dd, $J = 15.6, 7.0$, 1H), 2.36 (dd, $J = 15.4, 6.3$, 1H), 1.09 (t, $J = 7.2$, 6H). RT = 0.125 minutes (Cond. 2); LC-MS: Anal. Calcd. for $[\text{M}+\text{H}]^+$ $\text{C}_8\text{H}_{17}\text{N}_2\text{O}_3$: 189.12; found 189.13.


Cap-74x		LC-MS: Anal. Calcd. for $[M+H]^+$ C ₁₀ H ₂₂ NO ₂ : 188.17; found 188.21
---------	---	---

Cap-75



5

Cap-75, Step a



[00194] NaBH₃CN (1.6 g, 25.5 mmol) was added to a cooled (ice/water bath) water (25 ml)/methanol (15 ml) solution of H-D-Ser-OBzl HCl (2.0 g, 8.6 mmol).

Acetaldehyde (1.5 ml, 12.5 mmol) was added drop-wise over 5 min, the cooling bath was

10 removed, and the reaction mixture was stirred at ambient condition for 2 hr. The reaction was carefully quenched with 12N HCl and concentrated *in vacuo*. The residue was

dissolved in water and purified with a reverse phase HPLC (MeOH/H₂O/TFA) to afford the TFA salt of (R)-benzyl 2-(diethylamino)-3-hydroxypropanoate as a colorless viscous oil (1.9g). ¹H NMR (DMSO-d₆, δ = 2.5 ppm, 500 MHz): δ 9.73 (br s, 1H), 7.52-7.36 (m,

15 5H), 5.32 (d, *J* = 12.2, 1H), 5.27 (d, *J* = 12.5, 1H), 4.54-4.32 (m, 1H), 4.05-3.97 (m, 2H), 3.43-3.21 (m, 4H), 1.23 (t, *J* = 7.2, 6H). LC-MS (Cond. 2): RT = 1.38 min; LC-MS:


Anal. Calcd. for $[M+H]^+$ C₁₄H₂₂NO₃: 252.16; found 252.19.

Cap-75

20 [00195] NaH (0.0727 g, 1.82 mmol, 60%) was added to a cooled (ice-water) THF (3.0 mL) solution of the TFA salt (R)-benzyl 2-(diethylamino)-3-hydroxypropanoate (0.3019 g, 0.8264 mmol) prepared above, and the mixture was stirred for 15 min. Methyl iodide (56 μL, 0.90 mmol) was added and stirring was continued for 18 hr while allowing the

bath to thaw to ambient condition. The reaction was quenched with water and loaded onto a MeOH pre-conditioned MCX (6 g) cartridge, and washed with methanol followed by compound elution with 2N NH₃/Methanol. Removal of the volatile component *in vacuo* afforded Cap-75, contaminated with (R)-2-(diethylamino)-3-hydroxypropanoic acid, as a yellow semi-solid (100 mg). The product was used as is without further purification.

Cap-76




- 10 [00196] NaCNBH₃ (1.60 g, 24.2 mmol) was added in batches to a chilled (~15 °C) water/MeOH (12 mL each) solution of (*S*)-4-amino-2-(*tert*-butoxycarbonylamino) butanoic acid (2.17 g, 9.94 mmol). A few minutes later acetaldehyde (2.7 mL, 48.1 mmol) was added drop-wise over 2 min, the cooling bath was removed, and the reaction mixture was stirred at ambient condition for 3.5 hr. An additional acetaldehyde (2.7 mL,
- 15 48.1 mmol) was added and the reaction was stirred for 20.5 hr. Most of the MeOH component was removed *in vacuo*, and the remaining mixture was treated with concentrated HCl until its pH reached ~ 1.0 and then heated for 2 hr at 40 °C. The volatile component was removed *in vacuo*, and the residue was treated with 4 M HCl/dioxane (20 mL) and stirred at ambient condition for 7.5 hr. The volatile component
- 20 was removed *in vacuo* and the residue was purified with DOWEX® 50WX8-100 ion-exchange resin (column was washed with water and the compound was eluted with dilute NH₄OH, prepared from 18 ml of NH₄OH and 282 ml of water) to afford intermediate (*S*)-2-amino-4-(diethylamino)butanoic acid as an off-white solid (1.73 g).
- [00197] Methyl chloroformate (0.36 mL, 4.65 mmol) was added drop-wise over 11
- 25 min to a cooled (ice-water) mixture of Na₂CO₃ (0.243 g, 2.29 mmol), NaOH (4.6 mL of 1M/H₂O, 4.6 mmol) and the above product (802.4 mg). The reaction mixture was stirred for 55 min, and then the cooling bath was removed and stirring was continued for an additional 5.25 hr. The reaction mixture was diluted with equal volume of water and

washed with CH₂Cl₂ (30 mL, 2x), and the aqueous phase was cooled with ice-water bath and acidified with concentrated HCl to a pH region of 2. The volatile component was then removed *in vacuo* and the crude material was free-based with MCX resin (6.0g; column was washed with water, and sample was eluted with 2.0 M NH₃/MeOH) to afford
 5 impure Cap-76 as an off-white solid (704 mg). ¹H NMR (MeOH-d₄, δ = 3.29 ppm, 400 MHz): δ 3.99 (dd, *J* = 7.5, 4.7, 1H), 3.62 (s, 3H), 3.25-3.06 (m, 6H), 2.18-2.09 (m, 1H), 2.04-1.96 (m, 1H), 1.28 (t, *J* = 7.3, 6H). LC-MS: Anal. Calcd. for [M+H]⁺ C₁₀H₂₁N₂O₄: 233.15; found 233.24.

10

Cap-77a and Cap-77b




Cap-77a: enantiomer-1
 Cap-77b: enantiomer-2

[00198] The synthesis of Cap-77 was conducted according to the procedure described for Cap-7 by using 7-azabicyclo[2.2.1]heptane for the S_N2 displacement step, and by effecting the stereoisomeric separation of the intermediate benzyl 2-(7-

15 azabicyclo[2.2.1]heptan-7-yl)-2-phenylacetate using the following condition: the intermediate (303.7 mg) was dissolved in ethanol, and the resulting solution was injected on a chiral HPLC column (Chiracel AD-H column, 30 x 250 mm, 5 μm) eluting with 90% CO₂-10% EtOH at 70 mL/min, and a temperature of 35 °C to provide 124.5 mg of stereoisomer-1 and 133.8 mg of stereoisomer-2. These benzyl esters were
 20 hydrogenolysed according to the preparation of Cap-7 to provide Cap-77: ¹H NMR (DMSO-d₆, δ = 2.5 ppm, 400 MHz): δ 7.55 (m, 2H), 7.38-7.30 (m, 3H), 4.16 (s, 1H), 3.54 (app br s, 2H), 2.08-1.88 (m, 4 H), 1.57-1.46 (m, 4H). LC (Cond. 1): RT = 0.67 min; LC-MS: Anal. Calcd. for [M+H]⁺ C₁₄H₁₈NO₂: 232.13; found 232.18. HRMS: Anal. Calcd. for [M+H]⁺ C₁₄H₁₈NO₂: 232.1338; found 232.1340.

25


Cap-78



[00199] NaCNBH₃ (0.5828 g, 9.27 mmol) was added to a mixture of the HCl salt of (R)-2-(ethylamino)-2-phenylacetic acid (an intermediate in the synthesis of Cap-3; 0.9923 mg, 4.60 mmol) and (1-ethoxycyclopropoxy)trimethylsilane (1.640 g, 9.40 mmol) in MeOH (10 mL), and the semi-heterogeneous mixture was heated at 50 °C with an oil bath for 20 hr. More (1-ethoxycyclopropoxy)trimethylsilane (150 mg, 0.86 mmol) and NaCNBH₃ (52 mg, 0.827 mmol) were added and the reaction mixture was heated for an additional 3.5 hr. It was then allowed to cool to ambient temperature and acidified to a ~ pH region of 2 with concentrated HCl, and the mixture was filtered and the filtrate was rotovaped. The resulting crude material was taken up in *i*-PrOH (6 mL) and heated to effect dissolution, and the non-dissolved part was filtered off and the filtrate concentrated *in vacuo*. About 1/3 of the resultant crude material was purified with a reverse phase HPLC (H₂O/MeOH/TFA) to afford the TFA salt of Cap-78 as a colorless viscous oil (353 mg). ¹H NMR (DMSO-*d*₆, δ = 2.5 ppm, 400 MHz; after D₂O exchange): δ 7.56-7.49 (m, 5H), 5.35 (s, 1H), 3.35 (m, 1H), 3.06 (app br s, 1H), 2.66 (m, 1H), 1.26 (t, *J* = 7.3, 3H), 0.92 (m, 1H), 0.83-0.44 (m, 3H). LC (Cond. 1): RT = 0.64 min; LC-MS: Anal. Calcd. for [M+H]⁺ C₁₃H₁₈NO₂: 220.13; found 220.21. HRMS: Anal. Calcd. for [M+H]⁺ C₁₃H₁₈NO₂: 220.1338; found 220.1343.

20

Cap-79




[00200] Ozone was bubbled through a cooled (-78 °C) CH₂Cl₂ (5.0 mL) solution Cap-55 (369 mg, 2.13 mmol) for about 50 min until the reaction mixture attained a tint of blue color. Me₂S (10 pipette drops) was added, and the reaction mixture was stirred for 35 min. The -78 °C bath was replaced with a -10 °C bath and stirring continued for an

additional 30 min, and then the volatile component was removed *in vacuo* to afford a colorless viscous oil.

[00201] NaBH₃CN (149 mg, 2.25 mmol) was added to a MeOH (5.0 mL) solution of the above crude material and morpholine (500 μ L, 5.72 mmol) and the mixture was stirred at ambient condition for 4 hr. It was cooled to ice-water temperature and treated with concentrated HCl to bring its pH to ~2.0, and then stirred for 2.5 hr. The volatile component was removed *in vacuo*, and the residue was purified with a combination of MCX resin (MeOH wash; 2.0 N NH₃/MeOH elution) and a reverse phase HPLC (H₂O/MeOH/TFA) to afford Cap-79 containing unknown amount of morpholine.

[00202] In order to consume the morpholine contaminant, the above material was dissolved in CH₂Cl₂ (1.5 mL) and treated with Et₃N (0.27 mL, 1.94 mmol) followed by acetic anhydride (0.10 mL, 1.06 mmol) and stirred at ambient condition for 18 hr. THF (1.0 mL) and H₂O (0.5 mL) were added and stirring continued for 1.5 hr. The volatile component was removed *in vacuo*, and the resultant residue was passed through MCX resin (MeOH wash; 2.0 N NH₃/MeOH elution) to afford impure Cap-79 as a brown viscous oil, which was used for the next step without further purification.

Cap-80a and Cap-80b



[00203] SOCl₂ (6.60 mL, 90.5 mmol) was added drop-wise over 15 min to a cooled (ice-water) mixture of (S)-3-amino-4-(benzyloxy)-4-oxobutanoic acid (10.04g, 44.98 mmol) and MeOH (300 mL), the cooling bath was removed and the reaction mixture was stirred at ambient condition for 29 hr. Most of the volatile component was removed *in vacuo* and the residue was carefully partitioned between EtOAc (150 mL) and saturated NaHCO₃ solution. The aqueous phase was extracted with EtOAc (150 mL, 2x), and the combined organic phase was dried (MgSO₄), filtered, and concentrated *in vacuo* to afford (S)-1-benzyl 4-methyl 2-aminosuccinate as a colorless oil (9.706g). ¹H NMR (DMSO-d₆, δ = 2.5 ppm, 400 MHz): δ 7.40-7.32 (m, 5H), 5.11 (s, 2H), 3.72 (app t, *J* = 6.6, 1H), 3.55

(s, 3H), 2.68 (dd, $J = 15.9, 6.3$, 1H), 2.58 (dd, $J = 15.9, 6.8$, 1H), 1.96 (s, 2H). LC (Cond. 1): RT = 0.90 min; LC-MS: Anal. Calcd. for $[M+H]^+ C_{12}H_{16}NO_4$: 238.11; found 238.22.

[00204] $Pb(NO_3)_2$ (6.06 g, 18.3 mmol) was added over 1 min to a CH_2Cl_2 (80 mL)

solution of (S)-1-benzyl 4-methyl 2-aminosuccinate (4.50 g, 19.0 mmol), 9-bromo-9-

5 phenyl-9H-fluorene (6.44 g, 20.0 mmol) and Et_3N (3.0 mL, 21.5 mmol), and the heterogeneous mixture was stirred at ambient condition for 48 hr. The mixture was filtered and the filtrate was treated with $MgSO_4$ and filtered again, and the final filtrate was concentrated. The resulting crude material was submitted to a BIOTAGE®

purification (350 g silica gel, CH_2Cl_2 elution) to afford (S)-1-benzyl 4-methyl 2-(9-

10 phenyl-9H-fluoren-9-ylamino)succinate as highly viscous colorless oil (7.93 g). 1H NMR (DMSO- d_6 , $\delta = 2.5$ ppm, 400 MHz): δ 7.82 (m, 2H), 7.39-7.13 (m, 16H), 4.71 (d, $J = 12.4$, 1H), 4.51 (d, $J = 12.6$, 1H), 3.78 (d, $J = 9.1$, NH), 3.50 (s, 3H), 2.99 (m, 1H), 2.50-2.41 (m, 2H, partially overlapped with solvent). LC (Cond. 1): RT = 2.16 min; LC-MS: Anal. Calcd. for $[M+H]^+ C_{31}H_{28}NO_4$: 478.20; found 478.19.

15 **[00205]** LiHMDS (9.2 mL of 1.0 M/THF, 9.2 mmol) was added drop-wise over 10 min to a cooled ($-78^\circ C$) THF (50 mL) solution of (S)-1-benzyl 4-methyl 2-(9-phenyl-9H-fluoren-9-ylamino)succinate (3.907 g, 8.18 mmol) and stirred for ~ 1 hr. MeI (0.57 mL, 9.2 mmol) was added drop-wise over 8 min to the mixture, and stirring was continued for 16.5 hr while allowing the cooling bath to thaw to room temperature. After quenching

20 with saturated NH_4Cl solution (5 mL), most of the organic component was removed *in vacuo* and the residue was partitioned between CH_2Cl_2 (100 mL) and water (40 mL). The organic layer was dried ($MgSO_4$), filtered, and concentrated *in vacuo*, and the resulting crude material was purified with a BIOTAGE® (350 g silica gel; 25% EtOAc/hexanes) to afford 3.65 g of a 2S/3S and 2S/3R diastereomeric mixtures of 1-benzyl 4-methyl 3-

25 methyl-2-(9-phenyl-9H-fluoren-9-ylamino)succinate in $\sim 1.0:0.65$ ratio (1H NMR). The stereochemistry of the dominant isomer was not determined at this juncture, and the mixture was submitted to the next step without separation. Partial 1H NMR data (DMSO- d_6 , $\delta = 2.5$ ppm, 400 MHz): major diastereomer, δ 4.39 (d, $J = 12.3$, 1H of CH_2), 3.33 (s, 3H, overlapped with H_2O signal), 3.50 (d, $J = 10.9$, NH), 1.13 (d, $J = 7.1$, 3H); minor diastereomer, δ 4.27 (d, $J = 12.3$, 1H of CH_2), 3.76 (d, $J = 10.9$, NH), 3.64 (s, 3H), 0.77 (d, $J = 7.0$, 3H). LC (Cond. 1): RT = 2.19 min; LC-MS: Anal. Calcd. for $[M+H]^+ C_{32}H_{30}NO_4$: 492.22; found 492.15.

- [00206] Diisobutylaluminum hydride (20.57 ml of 1.0 M in hexanes, 20.57 mmol) was added drop-wise over 10 min to a cooled (-78 °C) THF (120 mL) solution of (2S)-1-benzyl 4-methyl 3-methyl-2-(9-phenyl-9H-fluoren-9-ylamino)succinate (3.37 g, 6.86 mmol) prepared above, and stirred at -78 °C for 20 hr. The reaction mixture was removed from the cooling bath and rapidly poured into ~1M H₃PO₄/H₂O (250 mL) with stirring, and the mixture was extracted with ether (100 mL, 2x). The combined organic phase was washed with brine, dried (MgSO₄), filtered and concentrated *in vacuo*. A silica gel mesh of the crude material was prepared and submitted to chromatography (25% EtOAc/hexanes; gravity elution) to afford 1.1g of (2S,3S)-benzyl 4-hydroxy-3-methyl-2-(9-phenyl-9H-fluoren-9-ylamino)butanoate, contaminated with benzyl alcohol, as a colorless viscous oil and (2S,3R)-benzyl 4-hydroxy-3-methyl-2-(9-phenyl-9H-fluoren-9-ylamino)butanoate containing the (2S,3R) stereoisomer as an impurity. The later sample was resubmitted to the same column chromatography purification conditions to afford 750 mg of purified material as a white foam. [Note: the (2S, 3S) isomer elutes before the (2S,3R) isomer under the above condition]. (2S, 3S) isomer: ¹H NMR (DMSO-d₆, δ = 2.5 ppm, 400 MHz): 7.81 (m, 2H), 7.39-7.08 (m, 16H), 4.67 (d, *J* = 12.3, 1H), 4.43 (d, *J* = 12.4, 1H), 4.21 (app t, *J* = 5.2, OH), 3.22 (d, *J* = 10.1, NH), 3.17 (m, 1H), 3.08 (m, 1H), ~2.5 (m, 1H, overlapped with the solvent signal), 1.58 (m, 1H), 0.88 (d, *J* = 6.8, 3H). LC (Cond. 1): RT = 2.00 min; LC-MS: Anal. Calcd. for [M+H]⁺ C₃₁H₃₀NO₃: 464.45; found 464.22. (2S, 3R) isomer: ¹H NMR (DMSO-d₆, δ = 2.5 ppm, 400 MHz): 7.81 (d, *J* = 7.5, 2H), 7.39-7.10 (m, 16H), 4.63 (d, *J* = 12.1, 1H), 4.50 (app t, *J* = 4.9, 1H), 4.32 (d, *J* = 12.1, 1H), 3.59-3.53 (m, 2H), 3.23 (m, 1H), 2.44 (dd, *J* = 9.0, 8.3, 1H), 1.70 (m, 1H), 0.57 (d, *J* = 6.8, 3H). LC (Cond. 1): RT = 1.92 min; LC-MS: Anal. Calcd. for [M+H]⁺ C₃₁H₃₀NO₃: 464.45; found 464.52.
- [00207] The relative stereochemical assignments of the DIBAL-reduction products were made based on NOE studies conducted on lactone derivatives prepared from each isomer by employing the following protocol: LiHMDS (50 μL of 1.0 M/THF, 0.05 mmol) was added to a cooled (ice-water) THF (2.0 mL) solution of (2S,3S)-benzyl 4-hydroxy-3-methyl-2-(9-phenyl-9H-fluoren-9-ylamino)butanoate (62.7 mg, 0.135 mmol), and the reaction mixture was stirred at similar temperature for ~2 hr. The volatile component was removed *in vacuo* and the residue was partitioned between CH₂Cl₂ (30 mL), water (20 mL) and saturated aqueous NH₄Cl solution (1 mL). The organic layer was dried

(MgSO₄), filtered, and concentrated *in vacuo*, and the resulting crude material was submitted to a BIOTAGE® purification (40 g silica gel; 10-15% EtOAc/hexanes) to afford (3S,4S)-4-methyl-3-(9-phenyl-9H-fluoren-9-ylamino)dihydrofuran-2(3H)-one as a colorless film of solid (28.1 mg). (2S,3R)-benzyl 4-hydroxy-3-methyl-2-(9-phenyl-9H-fluoren-9-ylamino)butanoate was elaborated similarly to (3S,4R)-4-methyl-3-(9-phenyl-9H-fluoren-9-ylamino)dihydrofuran-2(3H)-one. (3S,4S)-lactone isomer: ¹H NMR (DMSO-d₆, δ = 2.5 ppm, 400 MHz), 7.83 (d, *J* = 7.5, 2H), 7.46-7.17 (m, 11H), 4.14 (app t, *J* = 8.3, 1H), 3.60 (d, *J* = 5.8, NH), 3.45 (app t, *J* = 9.2, 1H), ~2.47 (m, 1H, partially overlapped with solvent signal), 2.16 (m, 1H), 0.27 (d, *J* = 6.6, 3H). LC (Cond. 1): RT = 1.98 min; LC-MS: Anal. Calcd. for [M+Na]⁺ C₂₄H₂₁NNaO₂: 378.15; found 378.42. (3S,4R)-lactone isomer: ¹H NMR (DMSO-d₆, δ = 2.5 ppm, 400 MHz), 7.89 (d, *J* = 7.6, 1H), 7.85 (d, *J* = 7.3, 1H), 7.46-7.20 (m, 11H), 3.95 (dd, *J* = 9.1, 4.8, 1H), 3.76 (d, *J* = 8.8, 1H), 2.96 (d, *J* = 3.0, NH), 2.92 (dd, *J* = 6.8, 3, NCH), 1.55 (m, 1H), 0.97 (d, *J* = 7.0, 3H). LC (Cond. 1): RT = 2.03 min; LC-MS: Anal. Calcd. for [M+Na]⁺ C₂₄H₂₁NNaO₂: 378.15; found 378.49.

[00208] TBDMS-Cl (48 mg, 0.312 mmol) followed by imidazole (28.8 mg, 0.423 mmol) were added to a CH₂Cl₂ (3 ml) solution of (2S,3S)-benzyl 4-hydroxy-3-methyl-2-(9-phenyl-9H-fluoren-9-ylamino)butanoate (119.5 mg, 0.258 mmol), and the mixture was stirred at ambient condition for 14.25 hr. The reaction mixture was then diluted with CH₂Cl₂ (30 mL) and washed with water (15 mL), and the organic layer was dried (MgSO₄), filtered, and concentrated *in vacuo*. The resultant crude material was purified with a BIOTAGE® (40 g silica gel; 5% EtOAc/hexanes) to afford (2S,3S)-benzyl 4-(*tert*-butyldimethylsilyloxy)-3-methyl-2-(9-phenyl-9H-fluoren-9-ylamino)butanoate, contaminated with TBDMS based impurities, as a colorless viscous oil (124.4 mg). (2S,3R)-benzyl 4-hydroxy-3-methyl-2-(9-phenyl-9H-fluoren-9-ylamino)butanoate was elaborated similarly to (2S,3R)-benzyl 4-(*tert*-butyldimethylsilyloxy)-3-methyl-2-(9-phenyl-9H-fluoren-9-ylamino)butanoate. (2S,3S)-silyl ether isomer: ¹H NMR (DMSO-d₆, δ = 2.5 ppm, 400 MHz), 7.82 (d, *J* = 4.1, 1H), 7.80 (d, *J* = 4.0, 1H), 7.38-7.07 (m, 16 H), 4.70 (d, *J* = 12.4, 1H), 4.42 (d, *J* = 12.3, 1H), 3.28-3.19 (m, 3H), 2.56 (dd, *J* = 10.1, 5.5, 1H), 1.61 (m, 1H), 0.90 (d, *J* = 6.8, 3H), 0.70 (s, 9H), -0.13 (s, 3H), -0.16 (s, 3H). LC (Cond. 1, where the run time was extended to 4 min): RT = 3.26 min; LC-MS: Anal. Calcd. for [M+H]⁺ C₃₇H₄₄NO₃Si: 578.31; found 578.40. (2S,3R)-silyl ether isomer: ¹H


NMR (DMSO- d_6 , δ = 2.5 ppm, 400 MHz), 7.82 (d, J = 3.0, 1H), 7.80 (d, J = 3.1, 1H), 7.39-7.10 (m, 16H), 4.66 (d, J = 12.4, 1H), 4.39 (d, J = 12.4, 1H), 3.61 (dd, J = 9.9, 5.6, 1H), 3.45 (d, J = 9.5, 1H), 3.41 (dd, J = 10, 6.2, 1H), 2.55 (dd, J = 9.5, 7.3, 1H), 1.74 (m, 1H), 0.77 (s, 9H), 0.61 (d, J = 7.1, 3H), -0.06 (s, 3H), -0.08 (s, 3H).

- 5 **[00209]** A balloon of hydrogen was attached to a mixture of (2S,3S)-benzyl 4-(*tert*-butyldimethylsilyloxy)-3-methyl-2-(9-phenyl-9H-fluoren-9-ylamino)butanoate (836 mg, 1.447 mmol) and 10% Pd/C (213 mg) in EtOAc (16 mL) and the mixture was stirred at room temperature for ~ 21 hr, where the balloon was recharged with H₂ as necessary. The reaction mixture was diluted with CH₂Cl₂ and filtered through a pad of diatomaceous earth (CELITE®-545), and the pad was washed with EtOAc (200 mL), EtOAc/MeOH (1:1 mixture, 200 mL) and MeOH (750 mL). The combined organic phase was concentrated, and a silica gel mesh was prepared from the resulting crude material and submitted to a flash chromatography (8:2:1 mixture of EtOAc/i-PrOH/H₂O) to afford (2S,3S)-2-amino-4-(*tert*-butyldimethylsilyloxy)-3-methylbutanoic acid as a white fluffy solid (325 mg). (2S,3R)-benzyl 4-(*tert*-butyldimethylsilyloxy)-3-methyl-2-(9-phenyl-9H-fluoren-9-ylamino)butanoate was similarly elaborated to (2S,3R)-2-amino-4-(*tert*-butyldimethylsilyloxy)-3-methylbutanoic acid. (2S,3S)-amino acid isomer: ¹H NMR (methanol- d_4 , δ = 3.29 ppm, 400 MHz), 3.76 (dd, J = 10.5, 5.2, 1H), 3.73 (d, J = 3.0, 1H), 3.67 (dd, J = 10.5, 7.0, 1H), 2.37 (m, 1H), 0.97 (d, J = 7.0, 3H), 0.92 (s, 9H), 0.10 (s, 6H). LC-MS: Anal. Calcd. for [M+H]⁺ C₁₁H₂₆NO₃Si: 248.17; found 248.44. (2S,3R)-amino acid isomer: ¹H NMR (methanol- d_4 , δ = 3.29 ppm, 400 MHz), 3.76-3.75 (m, 2H), 3.60 (d, J = 4.1, 1H), 2.16 (m, 1H), 1.06 (d, J = 7.3, 3H), 0.91 (s, 9H), 0.09 (s, 6H). Anal. Calcd. for [M+H]⁺ C₁₁H₂₆NO₃Si: 248.17; found 248.44.

- 25 **[00210]** Water (1 mL) and NaOH (0.18 mL of 1.0 M/H₂O, 0.18 mmol) were added to a mixture of (2S,3S)-2-amino-4-(*tert*-butyldimethylsilyloxy)-3-methylbutanoic acid (41.9 mg, 0.169 mmol) and Na₂CO₃ (11.9 mg, 0.112 mmol), and sonicated for about 1 min to effect dissolution of reactants. The mixture was then cooled with an ice-water bath, methyl chloroformate (0.02 mL, 0.259 mmol) was added over 30 s, and vigorous stirring was continued at similar temperature for 40 min and then at ambient temperature for 2.7 hr. The reaction mixture was diluted with water (5 mL), cooled with ice-water bath and treated drop-wise with 1.0 N HCl aqueous solution (~0.23 mL). The mixture was further diluted with water (10 mL) and extracted with CH₂Cl₂ (15 mL, 2x). The combined

organic phase was dried (MgSO_4), filtered, and concentrated *in vacuo* to afford Cap-80a as an off-white solid. (2S,3R)-2-amino-4-(*tert*-butyldimethylsilyloxy)-3-methylbutanoic acid was similarly elaborated to Cap-80b. Cap-80a: ^1H NMR ($\text{DMSO}-d_6$, $\delta = 2.5$ ppm, 400 MHz), 12.57 (br s, 1H), 7.64 (d, $J = 8.3$, 0.3H), 7.19 (d, $J = 8.8$, 0.7H), 4.44 (dd, $J = 8.1$, 4.6, 0.3H), 4.23 (dd, $J = 8.7$, 4.4, 0.7H), 3.56/3.53 (two singlets, 3H), 3.48-3.40 (m, 2H), 2.22-2.10 (m, 1H), 0.85 (s, 9H), ~ 0.84 (d, 0.9H, overlapped with *t*-Bu signal), 0.79 (d, $J = 7$, 2.1H), 0.02/0.01/0.00 (three overlapping singlets, 6H). LC-MS: Anal. Calcd. for $[\text{M}+\text{Na}]^+ \text{C}_{13}\text{H}_{27}\text{NNaO}_5\text{Si}$: 328.16; found 328.46. Cap-80b: ^1H NMR (CDCl_3 , $\delta = 7.24$ ppm, 400 MHz), 6.00 (br d, $J = 6.8$, 1H), 4.36 (dd, $J = 7.1$, 3.1, 1H), 3.87 (dd, $J = 10.5$, 3.0, 1H), 3.67 (s, 3H), 3.58 (dd, $J = 10.6$, 4.8, 1H), 2.35 (m, 1H), 1.03 (d, $J = 7.1$, 3H), 0.90 (s, 9H), 0.08 (s, 6H). LC-MS: Anal. Calcd. for $[\text{M}+\text{Na}]^+ \text{C}_{13}\text{H}_{27}\text{NNaO}_5\text{Si}$: 328.16; found 328.53. The crude products were utilized without further purification.


Cap-81




[00211] Prepared according to the protocol described by Falb et al., *Synthetic Communications*, 23:2839 (1993).

Cap-82 to Cap-85


[00212] Cap-82 to Cap-85 were synthesized from appropriate starting materials according to the procedure described for Cap-51 or Cap-13. The samples exhibited similar spectral profiles as that of their stereoisomers (*i.e.*, Cap-4, Cap-13, Cap-51 and Cap-52, respectively).




Cap-82



Cap-83




Cap-84



Cap-85


Cap-86



(2S,3R)-3-Methoxy-2-(methoxycarbonylamino)butanoic acid


[00213] To a mixture of O-methyl-L-threonine (3.0 g, 22.55 mmol), NaOH (0.902 g, 22.55 mmol) in H₂O (15 mL) was added ClCO₂Me (1.74 mL, 22.55 mmol) dropwise at 0 °C. The mixture was allowed to stir for 12 h and acidified to pH 1 using 1N HCl. The aqueous phase was extracted with EtOAc and (2x250 mL) and 10% MeOH in CH₂Cl₂ (250 mL) and the combined organic phases were concentrated under *in vacuo* to afford a colorless oil (4.18 g, 97%) which was of sufficient purity for use in subsequent steps. ¹H NMR (400 MHz, CDCl₃) δ 4.19 (s, 1H), 3.92-3.97 (m, 1H), 3.66 (s, 3H), 1.17 (d, *J* = 7.7 Hz, 3H). LC-MS: Anal. Calcd. for C₇H₁₃NO₅: 191; found: 190 (M-H)⁻.

Cap-87



[00214] To a mixture of L-homoserine (2.0 g, 9.79 mmol), Na₂CO₃ (2.08 g, 19.59 mmol) in H₂O (15 mL) was added ClCO₂Me (0.76 mL, 9.79 mmol) dropwise at 0 °C. The mixture was allowed to stir for 48 h and acidified to pH 1 using 1N HCl. The aqueous phase was extracted with EtOAc and (2X250 mL) and the combined organic phases were concentrated *in vacuo* to afford a colorless solid (0.719 g, 28%) which was of sufficient purity for use in subsequent steps. ¹H NMR (400 MHz, CDCl₃) δ 4.23 (dd, *J* = 4.5, 9.1 Hz, 1H), 3.66 (s, 3H), 3.43-3.49 (m, 2H), 2.08 - 2.14 (m, 1H), 1.82 - 1.89 (m, 1H). LC-MS: Anal. Calcd. for C₇H₁₃NO₅: 191; found: 192 (M+H)⁺.


Cap-88



[00215] A mixture of L-valine (1.0 g, 8.54 mmol), 3-bromopyridine (1.8 mL, 18.7 mmol), K₂CO₃ (2.45 g, 17.7 mmol) and CuI (169 mg, 0.887 mmol) in DMSO (10 mL)


was heated at 100 °C for 12h. The reaction mixture was cooled to rt, poured into H₂O (ca. 150 mL) and washed with EtOAc (x2). The organic layers were extracted with a small amount of H₂O and the combined aq phases were acidified to ca. pH 2 with 6N HCl. The volume was reduced to about one-third and 20g of cation exchange resin (Strata) was added. The slurry was allowed to stand for 20 min and loaded onto a pad of cation exchange resin (Strata) (ca. 25g). The pad was washed with H₂O (200 mL), MeOH (200 mL), and then NH₃ (3M in MeOH, 2X200 mL). The appropriate fractions was concentrated *in vacuo* and the residue (ca. 1.1 g) was dissolved in H₂O, frozen and lyophyllized. The title compound was obtained as a foam (1.02 g, 62%). ¹H NMR (400 MHz, DMSO-d₆) δ 8.00 (s, br, 1H), 7.68 - 7.71 (m, 1H), 7.01 (s, br, 1H), 6.88 (d, *J* = 7.5 Hz, 1H), 5.75 (s, br, 1H), 3.54 (s, 1H), 2.04 - 2.06 (m, 1H), 0.95 (d, *J* = 6.0 Hz, 3H), 0.91 (d, *J* = 6.6 Hz, 3H). LC-MS: Anal. Calcd. for C₁₀H₁₄N₂O₂: 194; found: 195 (M+H)⁺.

Cap-89



[00216] A mixture of L-valine (1.0 g, 8.54 mmol), 5-bromopyrimidine (4.03 g, 17.0 mmol), K₂CO₃ (2.40 g, 17.4 mmol) and CuI (179 mg, 0.94 mmol) in DMSO (10 mL) was heated at 100 °C for 12h. The reaction mixture was cooled to RT, poured into H₂O (ca. 150 mL) and washed with EtOAc (x2). The organic layers were extracted with a small amount of H₂O and the combined aq phases were acidified to ca. pH 2 with 6N HCl. The volume was reduced to about one-third and 20g of cation exchange resin (Strata) was added. The slurry was allowed to stand for 20 min and loaded onto a pad of cation exchange resin (Strata) (ca. 25g). The pad was washed with H₂O (200 mL), MeOH (200 mL), and then NH₃ (3M in MeOH, 2x200 mL). The appropriate fractions was concentrated *in vacuo* and the residue (ca. 1.1 g) was dissolved in H₂O, frozen and lyophyllized. The title compound was obtained as a foam (1.02 g, 62%). ¹H NMR (400 MHz, CD₃OD) showed the mixture to contain valine and the purity could not be estimated. The material was used as is in subsequent reactions. LC-MS: Anal. Calcd. for C₉H₁₃N₃O₂: 195; found: 196 (M+H)⁺.

Cap-90














- [00217] Cap-90 was prepared according to the method described for the preparation of
 5 Cap-1. The crude material was used as is in subsequent steps. LC-MS: Anal. Calcd. for $C_{11}H_{15}NO_2$: 193; found: 192 (M-H)⁻.






Cap-91 to Cap-116







- [00218] The following Caps were prepared according to the method used for
 10 preparation of Cap-51 unless noted otherwise:

Cap	Structure	LC-MS
Cap-91		LC-MS: Anal. Calcd. for $C_{11}H_{13}NO_4$: 223; found: 222 (M-H) ⁻ .
Cap-92		LC-MS: Anal. Calcd. for $C_{11}H_{13}NO_4$: 223; found: 222 (M-H) ⁻ .
Cap-93		LC-MS: Anal. Calcd. for $C_{10}H_{12}N_2O_4$: 224; found: 225 (M+H) ⁺ .
Cap-94		LC-MS: Anal. Calcd. for $C_8H_{11}N_3O_4$: 213; found: 214 (M+H) ⁺ .
Cap-95		LC-MS: Anal. Calcd. for $C_{13}H_{17}NO_4$: 251; found: 250 (M-H) ⁻ .

Cap	Structure	LC-MS
Cap-96		LC-MS: Anal. Calcd. for C ₁₂ H ₁₅ NO ₄ : 237; found: 236 (M-H) ⁻ .
Cap-97		LC-MS: Anal. Calcd. for C ₉ H ₁₅ NO ₄ : 201; found: 200 (M-H) ⁻ .
Cap-98		LC-MS: Anal. Calcd. for C ₉ H ₁₅ NO ₄ : 201; found: 202 (M+H) ⁺ .
Cap-99		¹ H NMR (400 MHz, CD ₃ OD) δ 3.88 - 3.94 (m, 1H), 3.60, 3.61 (s, 3H), 2.80 (m, 1H), 2.20 (m 1H), 1.82 - 1.94 (m, 3H), 1.45 - 1.71 (m, 2H).
Cap-99a		¹ H NMR (400 MHz, CD ₃ OD) δ 3.88 - 3.94 (m, 1H), 3.60, 3.61 (s, 3H), 2.80 (m, 1H), 2.20 (m 1H), 1.82 - 1.94 (m, 3H), 1.45 - 1.71 (m, 2H).
Cap-100		LC-MS: Anal. Calcd. for C ₁₂ H ₁₄ NO ₄ F: 255; found: 256 (M+H) ⁺ .






Cap	Structure	LC-MS
Cap-101		LC-MS: Anal. Calcd. for C ₁₁ H ₁₃ NO ₄ : 223; found: 222 (M-H) ⁻ .
Cap-102		LC-MS: Anal. Calcd. for C ₁₁ H ₁₃ NO ₄ : 223; found: 222 (M-H) ⁻ .
Cap-103		LC-MS: Anal. Calcd. for C ₁₀ H ₁₂ N ₂ O ₄ : 224; found: 225 (M+H) ⁺ .
Cap-104		¹ H NMR (400 MHz, CD ₃ OD) δ 3.60 (s, 3H), 3.50 - 3.53 (m, 1H), 2.66 - 2.69 and 2.44 - 2.49 (m, 1H), 1.91 - 2.01 (m, 2H), 1.62 - 1.74 (m, 4H), 1.51 - 1.62 (m, 2H).
Cap-105		¹ H NMR (400 MHz, CD ₃ OD) δ 3.60 (s, 3H), 3.33 - 3.35 (m, 1H, partially obscured by solvent), 2.37 - 2.41 and 2.16 - 2.23 (m, 1H), 1.94 - 2.01 (m, 4H), 1.43 - 1.53 (m, 2H), 1.17 - 1.29 (m, 2H).



Cap	Structure	LC-MS
Cap-106	 <p>Prepared from cis-4-aminocyclohexane carboxylic acid and acetaldehyde by employing a similar procedure described for the synthesis of Cap-2. The crude HCl salt was passed through MCX (MeOH/H₂O/CH₂Cl₂ wash; 2 N NH₃/MeOH elution) to afford an oil, which was dissolved in CH₃CN/H₂O and lyophilized to afford a tan solid.</p>	¹ H NMR (400 MHz, CD ₃ OD) δ 3.16 (q, <i>J</i> = 7.3 Hz, 4H), 2.38 - 2.41 (m, 1H), 2.28 - 2.31 (m, 2H), 1.79 - 1.89 (m, 2H), 1.74 (app, ddd <i>J</i> = 3.5, 12.5, 15.9 Hz, 2H), 1.46 (app dt <i>J</i> = 4.0, 12.9 Hz, 2H), 1.26 (t, <i>J</i> = 7.3 Hz, 6H)
Cap-107		LC-MS: Anal. Calcd. for C ₈ H ₁₀ N ₂ O ₄ S: 230; found: 231 (M+H) ⁺ .
Cap-108		LC-MS: Anal. Calcd. for C ₁₅ H ₁₇ N ₃ O ₄ : 303; found: 304 (M+H) ⁺ .
Cap-109		LC-MS: Anal. Calcd. for C ₁₀ H ₁₂ N ₂ O ₄ : 224; found: 225 (M+H) ⁺ .
Cap-110		LC-MS: Anal. Calcd. for C ₁₀ H ₁₂ N ₂ O ₄ : 224; found: 225 (M+H) ⁺ .

Cap	Structure	LC-MS
Cap-111		LC-MS: Anal. Calcd. for C ₁₂ H ₁₆ NO ₈ P: 333; found: 334 (M+H) ⁺ .
Cap-112		LC-MS: Anal. Calcd. for C ₁₃ H ₁₄ N ₂ O ₄ : 262; found: 263 (M+H) ⁺ .
Cap-113		LC-MS: Anal. Calcd. for C ₁₈ H ₁₉ NO ₅ : 329; found: 330 (M+H) ⁺ .
Cap-114		¹ H NMR (400 MHz, CDCl ₃) δ 4.82 - 4.84 (m, 1H), 4.00 - 4.05 (m, 2H), 3.77 (s, 3H), 2.56 (s, br, 2H)
Cap-115		¹ H NMR (400 MHz, CDCl ₃) δ 5.13 (s, br, 1H), 4.13 (s, br, 1H), 3.69 (s, 3H), 2.61 (d, <i>J</i> = 5.0 Hz, 2H), 1.28 (d, <i>J</i> = 9.1 Hz, 3H).
Cap-116		¹ H NMR (400 MHz, CDCl ₃) δ 5.10 (d, <i>J</i> = 8.6 Hz, 1H), 3.74 - 3.83 (m, 1H), 3.69 (s, 3H), 2.54 - 2.61 (m, 2H), 1.88 (sept, <i>J</i> = 7.0 Hz, 1H), 0.95 (d, <i>J</i> = 7.0 Hz, 6H).


Cap-117 to Cap-123

[00219] For the preparation of Cap-117 to Cap-123 the Boc amino acids were obtained from commercially sources and were deprotected by treatment with 25% TFA in CH_2Cl_2 . After complete reaction as judged by LC-MS the solvents were removed *in vacuo* and the corresponding TFA salt of the amino acid was carbamoylated with methyl chloroformate according to the procedure described for Cap-51.

Cap	Structure	LC-MS
Cap-117		LC-MS: Anal. Calcd. for $\text{C}_{12}\text{H}_{15}\text{NO}_4$: 237; found: 238 (M+H) ⁺ .
Cap-118		LC-MS: Anal. Calcd. for $\text{C}_{10}\text{H}_{13}\text{NO}_4\text{S}$: 243; found: 244 (M+H) ⁺ .
Cap-119		LC-MS: Anal. Calcd. for $\text{C}_{10}\text{H}_{13}\text{NO}_4\text{S}$: 243; found: 244 (M+H) ⁺ .
Cap-120		LC-MS: Anal. Calcd. for $\text{C}_{10}\text{H}_{13}\text{NO}_4\text{S}$: 243; found: 244 (M+H) ⁺ .
Cap-121		¹ H NMR (400 MHz, CDCl_3) δ 4.06 - 4.16 (m, 1H), 3.63 (s, 3H), 3.43 (s, 1H), 2.82 and 2.66 (s, br, 1H), 1.86 - 2.10 (m, 3H), 1.64 - 1.76 (m, 2H), 1.44 - 1.53 (m, 1H).


Cap	Structure	LC-MS
Cap-122		¹ H NMR profile is similar to that of its stereoisomer, Cap-121.
Cap-123		LC-MS: Anal. Calcd. for C ₂₇ H ₂₆ N ₂ O ₆ : 474; found: 475 (M+H) ⁺ .

Cap-124



[00220] The hydrochloride salt of L-threonine *tert*-butyl ester was carbamoylated according to the procedure for Cap-51. The crude reaction mixture was acidified with 1N HCl to pH~1 and the mixture was extracted with EtOAc (2X50 mL). The combined organic phases were concentrated *in vacuo* to give a colorless oil which solidified on standing. The aqueous layer was concentrated *in vacuo* and the resulting mixture of product and inorganic salts was triturated with EtOAc-CH₂Cl₂-MeOH (1:1:0.1) and then the organic phase concentrated *in vacuo* to give a colorless oil which was shown by LC-MS to be the desired product. Both crops were combined to give 0.52 g of a solid. ¹H NMR (400 MHz, CD₃OD) δ 4.60 (m, 1H), 4.04 (d, *J* = 5.0 Hz, 1H), 1.49 (d, *J* = 6.3 Hz, 3H). LC-MS: Anal. Calcd. for C₅H₇NO₄: 145; found: 146 (M+H)⁺.


Cap-125



[00221] To a suspension of Pd(OH)₂, (20%, 100 mg), aqueous formaldehyde (37% wt, 4 ml), acetic acid, (0.5 mL) in methanol (15 mL) was added (S)-4-amino-2-(*tert*-butoxycarbonylamino)butanoic acid (1 g, 4.48 mmol). The reaction was purged several

times with hydrogen and was stirred overnight with an hydrogen balloon room temperature. The reaction mixture was filtered through a pad of diatomaceous earth (CELITE®), and the volatile component was removed *in vacuo*. The resulting crude material was used as is for the next step. LC-MS: Anal. Calcd. for C₁₁H₂₂N₂O₄: 246;

5 found: 247 (M+H)⁺.



[00222] This procedure is a modification of that used to prepare Cap-51. To a


10 suspension of 3-methyl-L-histidine (0.80 g, 4.70 mmol) in THF (10mL) and H₂O (10 mL) at 0 °C was added NaHCO₃ (0.88 g, 10.5 mmol). The resulting mixture was treated with ClCO₂Me (0.40 mL, 5.20 mmol) and the mixture allowed to stir at 0 °C. After stirring for ca. 2h LC-MS showed no starting material remaining. The reaction was acidified to pH 2 with 6 N HCl.

15 [00223] The solvents were removed *in vacuo* and the residue was suspended in 20 mL of 20% MeOH in CH₂Cl₂. The mixture was filtered and concentrated to give a light yellow foam (1.21 g.). LC-MS and ¹H NMR showed the material to be a 9:1 mixture of the methyl ester and the desired product. This material was taken up in THF (10mL) and H₂O (10mL), cooled to 0 °C and LiOH (249.1 mg, 10.4 mmol) was added. After stirring

20 ca. 1h LC-MS showed no ester remaining. Therefore the mixture was acidified with 6N HCl and the solvents removed *in vacuo*. LC-MS and ¹H NMR confirm the absence of the ester. The title compound was obtained as its HCl salt contaminated with inorganic salts (1.91 g, >100%). The compound was used as is in subsequent steps without further purification. ¹H NMR (400 MHz, CD₃OD) δ 8.84, (s, 1H), 7.35 (s, 1H), 4.52 (dd, *J* =

25 5.0, 9.1 Hz, 1H), 3.89 (s, 3H), 3.62 (s, 3H), 3.35 (dd, *J* = 4.5, 15.6 Hz, 1H, partially obscured by solvent), 3.12 (dd, *J* = 9.0, 15.6 Hz, 1H). LC-MS: Anal. Calcd. for C₉H₁₃N₃O₄: 227.09; found: 228.09 (M+H)⁺.


Cap-127




[00224] Cap-127 was prepared according to the method for Cap-126 above starting from (S)-2-amino-3-(1-methyl-1H-imidazol-4-yl)propanoic acid (1.11 g, 6.56 mmol), NaHCO₃ (1.21 g, 14.4 mmol) and ClCO₂Me (0.56 mL, 7.28 mmol). The title compound was obtained as its HCl salt (1.79 g, >100%) contaminated with inorganic salts. LC-MS and ¹H NMR showed the presence of ca. 5% of the methyl ester. The crude mixture was used as is without further purification. ¹H NMR (400 MHz, CD₃OD) δ 8.90 (s, 1H), 7.35 (s, 1H), 4.48 (dd, *J* = 5.0, 8.6 Hz, 1H), 3.89 (s, 3H), 3.62 (s, 3H), 3.35 (m, 1H), 3.08 (m, 1H); LC-MS: Anal. Calcd. for C₉H₁₃N₃O₄: 227.09; found: 228 (M+H)⁺.

10

Preparation of Cap-128



Step 1. Preparation of (S)-benzyl 2-(*tert*-butoxycarbonylamino)pent-4-ynoate (cj-27b).




15

[00225] To a solution of cj-27a (1.01 g, 4.74 mmol), DMAP (58 mg, 0.475 mmol) and iPr₂NEt (1.7 mL, 9.8 mmol) in CH₂Cl₂ (100 mL) at 0 °C was added Cbz-Cl (0.68 mL,


4.83 mmol). The solution was allowed to stir for 4 h at 0 °C, washed (1N KHSO₄, brine), dried (Na₂SO₄), filtered, and concentrated *in vacuo*. The residue was purified by flash column chromatography (TLC 6:1 hex:EtOAc) to give the title compound (1.30 g, 91%) as a colorless oil. ¹H NMR (400 MHz, CDCl₃) δ 7.35 (s, 5H), 5.35 (d, br, *J* = 8.1 Hz, 1H), 5.23 (d, *J* = 12.2 Hz, 1H), 5.17 (d, *J* = 12.2 Hz, 1H), 4.48 - 4.53 (m, 1H), 2.68 - 2.81 (m, 2H), 2.00 (t, *J* = 2.5 Hz, 1H), 1.44 (s, 9H). LC-MS: Anal. Calcd. for C₁₇H₂₁NO₄: 303; found: 304 (M+H)⁺.

Step 2. Preparation of (S)-benzyl 3-(1-benzyl-1H-1,2,3-triazol-4-yl)-2-(*tert*-butoxycarbonylamino)propanoate (cj-28).




[00226] To a mixture of (S)-benzyl 2-(*tert*-butoxycarbonylamino)pent-4-ynoate (0.50 g, 1.65 mmol), sodium ascorbate (0.036 g, 0.18 mmol), CuSO₄·5H₂O (0.022 g, 0.09 mmol) and NaN₃ (0.13 g, 2.1 mmol) in DMF-H₂O (5 mL, 4:1) at rt was added BnBr (0.24 mL, 2.02 mmol) and the mixture was warmed to 65 °C. After 5h LC-MS indicated low conversion. A further portion of NaN₃ (100 mg) was added and heating was continued for 12h. The reaction was poured into EtOAc and H₂O and shaken. The layers were separated and the aqueous layer extracted 3x with EtOAc and the combined organic phases washed (H₂O x3, brine), dried (Na₂SO₄), filtered, and concentrated. The residue was purified by flash (BIOTAGE®, 40+M 0-5% MeOH in CH₂Cl₂; TLC 3% MeOH in CH₂Cl₂) to afford a light yellow oil which solidified on standing (748.3 mg, 104%). The NMR was consistent with the desired product but suggests the presence of DMF. The material was used as is without further purification. ¹H NMR (400 MHz, DMSO-d₆) δ 7.84 (s, 1H), 7.27 - 7.32 (m, 10H), 5.54 (s, 2H), 5.07 (s, 2H), 4.25 (m, 1H), 3.16 (dd, *J* = 1.0, 5.3 Hz, 1H), 3.06 (dd, *J* = 5.3, 14.7 Hz), 2.96 (dd, *J* = 9.1, 14.7 Hz, 1H), 1.31 (s, 9H). LC-MS: Anal. Calcd. for C₂₄H₂₈N₄O₄: 436; found: 437 (M+H)⁺.

Step 3. Preparation of (S)-benzyl 3-(1-benzyl-1H-1,2,3-triazol-4-yl)-2-(methoxycarbonylamino)propanoate (cj-29).




[00227] A solution of (S)-benzyl 3-(1-benzyl-1H-1,2,3-triazol-4-yl)-2-(*tert*-
 5 butoxycarbonylamino)propanoate (0.52 g, 1.15 mmol) in CH₂Cl₂ was added TFA (4 mL). The mixture was allowed to stir at room temperature for 2h. The mixture was concentrated *in vacuo* to give a colorless oil which solidified on standing. This material was dissolved in THF-H₂O and cooled to 0 °C. Solid NaHCO₃ (0.25 g, 3.00 mmol) was added followed by ClCO₂Me (0.25 mL, 3.25 mmol). After stirring for 1.5h the mixture
 10 was acidified to pH~2 with 6N HCl and then poured into H₂O-EtOAc. The layers were separated and the aq phase extracted 2x with EtOAc. The combined org layers were washed (H₂O, brine), dried (Na₂SO₄), filtered, and concentrated *in vacuo* to give a colorless oil (505.8 mg, 111%, NMR suggested the presence of an unidentified impurity) which solidified while standing on the pump. The material was used as is without further
 15 purification. ¹H NMR (400 MHz, DMSO-d₆) δ 7.87 (s, 1H), 7.70 (d, *J* = 8.1 Hz, 1H), 7.27 - 7.32 (m, 10H), 5.54 (s, 2H), 5.10 (d, *J* = 12.7 Hz, 1H), 5.06 (d, *J* = 12.7 Hz, 1H), 4.32 - 4.37 (m, 1H), 3.49 (s, 3H), 3.09 (dd, *J* = 5.6, 14.7 Hz, 1H), 2.98 (dd, *J* = 9.6, 14.7 Hz, 1H). LC-MS: Anal. Calcd. for C₂₁H₂₂N₄O₄: 394; found: 395 (M+H)⁺.

20 Step 4. Preparation of (S)-2-(methoxycarbonylamino)-3-(1H-1,2,3-triazol-4-yl)propanoic acid (Cap-128).




[00228] (S)-Benzyl 3-(1-benzyl-1H-1,2,3-triazol-4-yl)-2-(methoxycarbonylamino)propanoate (502 mg, 1.11 mmol) was hydrogenated in the
 25 presence of Pd-C (82 mg) in MeOH (5 mL) at atmospheric pressure for 12h. The mixture

was filtered through diatomaceous earth (CELITE®) and concentrated *in vacuo*. (S)-2-(methoxycarbonylamino)-3-(1H-1,2,3-triazol-4-yl)propanoic acid was obtained as a colorless gum (266 mg, 111%) which was contaminated with ca. 10% of the methyl ester. The material was used as is without further purification. ¹H NMR (400 MHz, DMSO-d₆) δ 12.78 (s, br, 1H), 7.59 (s, 1H), 7.50 (d, *J* = 8.0 Hz, 1H), 4.19 - 4.24 (m, 1H), 3.49 (s, 3H), 3.12 (dd, *J* = 4.8 Hz, 14.9 Hz, 1H), 2.96 (dd, *J* = 9.9, 15.0 Hz, 1H). LC-MS: Anal. Calcd. for C₇H₁₀N₄O₄: 214; found: 215 (M+H)⁺.




Step 1. Preparation of (S)-2-(benzyloxycarbonylamino)-3-(1H-pyrazol-1-yl)propanoic acid (cj-31).



15 **[00229]** A suspension of (S)-benzyl 2-oxooxetan-3-ylcarbamate (0.67 g, 3.03 mmol), and pyrazole (0.22 g, 3.29 mmol) in CH₃CN (12 mL) was heated at 50 °C for 24h. The mixture was cooled to rt overnight and the solid filtered to afford (S)-2-(benzyloxycarbonylamino)-3-(1H-pyrazol-1-yl)propanoic acid (330.1 mg). The filtrate was concentrated *in vacuo* and then triturated with a small amount of CH₃CN (ca. 4 mL) to afford a second crop (43.5 mg). Total yield 370.4 mg (44%). m.p. 165.5 - 168 °C. lit m.p. 168.5 - 169.5 [Vederas et al., *J. Am. Chem. Soc.*, 107:7105 (1985)]. ¹H NMR (400 MHz, CD₃OD) δ 7.51 (d, *J* = 2.0, 1H), 7.48 (s, *J* = 1.5 Hz, 1H), 7.24 - 7.34 (m, 5H), 6.23 m, 1H), 5.05 (d, 12.7 Hz, 1H), 5.03 (d, *J* = 12.7 Hz, 1H), 4.59 - 4.66 (m, 2H), 4.42 - 4.49 (m, 1H). LC-MS: Anal. Calcd. for C₁₄H₁₅N₃O₄: 289; found: 290 (M+H)⁺.


25

Step 2. Preparation of (S)-2-(methoxycarbonylamino)-3-(1H-pyrazol-1-yl)propanoic acid (Cap-129).




[00230] (S)-2-(Benzyloxycarbonylamino)-3-(1H-pyrazol-1-yl)propanoic acid (0.20 g, 0.70 mmol) was hydrogenated in the presence of Pd-C (45 mg) in MeOH (5 mL) at atmospheric pressure for 2h. The product appeared to be insoluble in MeOH, therefore the reaction mixture was diluted with 5mL H₂O and a few drops of 6N HCl. The homogeneous solution was filtered through diatomaceous earth (CELITE®), and the MeOH removed *in vacuo*. The remaining solution was frozen and lyophilized to give a yellow foam (188.9 mg). This material was suspended in THF-H₂O (1:1, 10mL) and then cooled to 0 °C. To the cold mixture was added NaHCO₃ (146.0 mg, 1.74 mmol) carefully (evolution of CO₂). After gas evolution had ceased (ca. 15 min) ClCO₂Me (0.06 mL, 0.78 mmol) was added dropwise. The mixture was allowed to stir for 2h and was acidified to pH~2 with 6N HCl and poured into EtOAc. The layers were separated and the aqueous phase extracted with EtOAc (x5). The combined organic layers were washed (brine), dried (Na₂SO₄), filtered, and concentrated to give the title compound as a colorless solid (117.8 mg, 79%). ¹H NMR (400 MHz, DMSO-d₆) δ 13.04 (s, 1H), 7.63 (d, *J* = 2.6 Hz, 1H), 7.48 (d, *J* = 8.1 Hz, 1H), 7.44 (d, *J* = 1.5 Hz, 1H), 6.19 (app t, *J* = 2.0 Hz, 1H), 4.47 (dd, *J* = 3.0, 12.9 Hz, 1H), 4.29 - 4.41 (m, 2H), 3.48 (s, 3H). LC-MS: Anal. Calcd. for C₈H₁₁N₃O₄: 213; found: 214 (M+H)⁺.

Cap-130




[00231] Cap-130 was prepared by acylation of commercially available (R)-phenylglycine analogous to the procedure given in: Calmes, M. et al., *Tetrahedron*, 43(10):2285 (1987).




[00232] Step a: Dimethylcarbamoyl chloride (0.92 mL, 10 mmol) was added slowly to a solution of (S)-benzyl 2-amino-3-methylbutanoate hydrochloride (2.44 g; 10 mmol) and Hunig's base (3.67 mL, 21 mmol) in THF (50 mL). The resulting white suspension was stirred at room temperature overnight (16 hours) and concentrated under reduced pressure. The residue was partitioned between ethyl acetate and water. The organic layer was washed with brine, dried (MgSO₄), filtered, and concentrated under reduced pressure. The resulting yellow oil was purified by flash chromatography, eluting with ethyl acetate:hexanes (1:1). Collected fractions were concentrated under vacuum providing 2.35 g (85%) of clear oil. ¹H NMR (300 MHz, DMSO-d₆) δ ppm 0.84 (d, *J*=6.95 Hz, 3H), 0.89 (d, *J*=6.59 Hz, 3H), 1.98-2.15 (m, 1H), 2.80 (s, 6H), 5.01-5.09 (m, *J*=12.44 Hz, 1H), 5.13 (d, *J*=12.44 Hz, 1H), 6.22 (d, *J*=8.05 Hz, 1H), 7.26-7.42 (m, 5H). LC (Cond. 1): RT = 1.76 min; MS: Anal. Calcd. for [M+H]⁺ C₁₆H₂₂N₂O₃: 279.17; found 279.03.

[00233] Step b: To an MeOH (50 mL) solution of the intermediate prepared above (2.35 g; 8.45 mmol) was added Pd/C (10%; 200 mg) and the resulting black suspension was flushed with N₂ (3x) and placed under 1 atm of H₂. The mixture was stirred at room temperature overnight and filtered through a microfiber filter to remove the catalyst. The resulting clear solution was then concentrated under reduced pressure to obtain 1.43 g (89%) of Cap-131 as a white foam, which was used without further purification. ¹H NMR (500 MHz, DMSO-d₆) δ ppm 0.87 (d, *J*=4.27 Hz, 3H), 0.88 (d, *J*=3.97 Hz, 3H), 1.93-2.11 (m, 1H), 2.80 (s, 6H), 3.90 (dd, *J*=8.39, 6.87 Hz, 1H), 5.93 (d, *J*=8.54 Hz, 1H), 12.36 (s, 1H). LC (Cond. 1): RT = 0.33 min; MS: Anal. Calcd. for [M+H]⁺ C₈H₁₇N₂O₃: 189.12; found 189.04.


Cap-132




- [00234]** Cap-132 was prepared from (S)-benzyl 2-aminopropanoate hydrochloride according to the method described for Cap-131. ^1H NMR (500 MHz, DMSO- d_6) δ ppm 1.27 (d, $J=7.32$ Hz, 3H), 2.80 (s, 6H), 4.06 (qt, 1H), 6.36 (d, $J=7.32$ Hz, 1H), 12.27 (s, 1H). LC (Cond. 1): RT = 0.15 min; MS: Anal. Calcd. for $[\text{M}+\text{H}]^+$ $\text{C}_6\text{H}_{13}\text{N}_2\text{O}_3$: 161.09; found 161.00.




- [00235]** Cap-133 was prepared from (S)-*tert*-butyl 2-amino-3-methylbutanoate hydrochloride and 2-fluoroethyl chloroformate according to the method described for Cap-47. ^1H NMR (500 MHz, DMSO- d_6) δ ppm 0.87 (t, $J = 6.71$ Hz, 6H), 1.97-2.10 (m, 1H), 3.83 (dd, $J=8.39, 5.95$ Hz, 1H), 4.14-4.18 (m, 1H), 4.20-4.25 (m, 1H), 4.50-4.54 (m, 1H), 4.59-4.65 (m, 1H), 7.51 (d, $J = 8.54$ Hz, 1H), 12.54 (s, 1H).



- [00236]** Cap-134 was prepared from (S)-diethyl alanine and methyl chloroformate according to the method described for Cap-51. ^1H NMR (500 MHz, DMSO- d_6) δ ppm 0.72-0.89 (m, 6H), 1.15-1.38 (m, 4H), 1.54-1.66 (m, 1H), 3.46-3.63 (m, 3H), 4.09 (dd, $J = 8.85, 5.19$ Hz, 1H), 7.24 (d, $J = 8.85$ Hz, 1H), 12.55 (s, 1H). LC (Cond. 2): RT = 0.66 min; LC-MS: Anal. Calcd. for $[\text{M}+\text{H}]^+$ $\text{C}_9\text{H}_{18}\text{NO}_4$: 204.12; found 204.02.




- [00237] A solution of D-2-amino-(4-fluorophenyl)acetic acid (338 mg, 2.00 mmol), 1N HCl in diethylether (2.0 mL, 2.0 mmol) and formalin (37%, 1 mL) in methanol (5 mL) was subjected to balloon hydrogenation over 10% palladium on carbon (60 mg) for 16 h at 25 °C. The mixture was then filtered through CELITE® to afford the HCl salt of Cap-135 as a white foam (316 mg, 80%). ¹H NMR (300 MHz, MeOH-d₄) δ 7.59 (dd, *J* = 8.80, 5.10 Hz, 2H), 7.29 (t, *J* = 8.6 Hz, 2H), 5.17 (s, 1H), 3.05 (v br s, 3H), 2.63 (v br s, 3H); *R*_t = 0.19 min (Cond.-MS-W5); 95% homogeneity index; LRMS: Anal. Calcd. for [M+H]⁺ C₁₀H₁₃FNO₂: 198.09; found: 198.10.



- [00238] To a cooled (-50 °C) suspension of 1-benzyl-1*H*-imidazole (1.58 g, 10.0 mmol) in anhydrous diethyl ether (50 mL) under nitrogen was added *n*-butyl lithium (2.5 M in hexanes, 4.0 mL, 10.0 mmol) dropwise. After being stirred for 20 min at -50 °C, dry carbon dioxide (passed through Drierite) was bubbled into the reaction mixture for 10 min before it was allowed to warm up to 25 °C. The heavy precipitate which formed on addition of carbon dioxide to the reaction mixture was filtered to yield a hygroscopic, white solid which was taken up in water (7 mL), acidified to pH = 3, cooled, and induced to crystallize with scratching. Filtration of this precipitate gave a white solid which was suspended in methanol, treated with 1N HCl/diethyl ether (4 mL) and concentrated *in vacuo*. Lyophilization of the residue from water (5 mL) afforded the HCl salt of Cap-136 as a white solid (817 mg, 40%). ¹H NMR (300 MHz, DMSO-d₆) δ 7.94 (d, *J* = 1.5 Hz, 1H), 7.71 (d, *J* = 1.5 Hz, 1H), 7.50-7.31 (m, 5H), 5.77 (s, 2H); *R*_t = 0.51 min (Cond.-MS-


W5); 95% homogeneity index; LRMS: Anal. Calc. for $[M+H]^+$ $C_{11}H_{12}N_2O_2$: 203.08; found: 203.11.

Cap-137




5

Cap-137, Step a



[00239] A suspension of 1-chloro-3-cyanoisoquinoline (188 mg, 1.00 mmol; prepared according to the procedure in WO 2003/099274) (188 mg, 1.00 mmol), cesium fluoride (303.8 mg, 2.00 mmol), bis(tri-*tert*-butylphosphine)palladium dichloride (10 mg, 0.02 mmol) and 2-(tributylstannyl)furan (378 μ L, 1.20 mmol) in anhydrous dioxane (10 mL) under nitrogen was heated at 80 °C for 16 h before it was cooled to 25 °C and treated with saturated, aqueous potassium fluoride solution with vigorous stirring for 1 h. The mixture was partitioned between ethyl acetate and water and the organic phase was separated, washed with brine, dried over Na_2SO_4 , filtered and concentrated. Purification of the residue on silica gel (elution with 0% to 30% ethyl acetate/hexanes) afforded Cap-137, Step a as a white solid which was used as is (230 mg, 105%). R_t = 1.95 min (Cond.-MS-W2); 90% homogeneity index; LRMS: Anal. Calc. for $[M+H]^+$ $C_{14}H_8N_2O$: 221.07; found: 221.12.


Cap-137




[00240] To a suspension of Cap-137, Step a (110 mg, 0.50 mmol) and sodium periodate (438 mg, 2.05 mmol) in carbon tetrachloride (1 mL), acetonitrile (1 mL) and water (1.5 mL) was added ruthenium trichloride hydrate (2 mg, 0.011 mmol). The mixture was stirred at 25 °C for 2 h and then partitioned between dichloromethane and water. The aqueous layer was separated, extracted twice more with dichloromethane and the combined dichloromethane extracts were dried over Na₂SO₄, filtered and concentrated. Trituration of the residue with hexanes afforded Cap-137 (55 mg, 55%) as a grayish-colored solid. *R*_t = 1.10 min (Cond.-MS-W2); 90% homogeneity index; LC-MS: Anal. Calc. for [M+H]⁺ C₁₁H₈N₂O₂: 200.08; found: 200.08.

Cap-138 to Cap-158


Synthetic Strategy. Method A.



Cap-138



Cap-138, Step a




[00241] To a stirred suspension of 5-hydroxyisoquinoline (prepared according to the procedure in WO 2003/099274) (2.0 g, 13.8 mmol) and triphenylphosphine (4.3 g, 16.5 mmol) in dry tetrahydrofuran (20 mL) was added dry methanol (0.8 mL) and diethyl azodicarboxylate (3.0 mL, 16.5 mmol) portionwise. The mixture was stirred at room temperature for 20 h before it was diluted with ethyl acetate and washed with brine, dried over Na₂SO₄, filtered and concentrated. The residue was preabsorbed onto silica gel and

purified (elution with 40% ethyl acetate/hexanes) to afford Cap-138, Step a as a light yellow solid (1.00 g, 45%). ^1H NMR (CDCl_3 , 500 MHz) δ 9.19 (s, 1H), 8.51 (d, $J = 6.0$ Hz, 1H), 7.99 (d, $J = 6.0$ Hz, 1H), 7.52-7.50 (m, 2H), 7.00-6.99 (m, 1H), 4.01 (s, 3H); $R_t = 0.66$ min (Cond. D2); 95% homogeneity index; LC-MS: Anal. Calc. for $[\text{M}+\text{H}]^+$

5 $\text{C}_{10}\text{H}_{10}\text{NO}$: 160.08; found 160.10.


Cap-138, Step b



[00242] To a stirred solution of Cap-138, Step a (2.34 g, 14.7 mmol) in anhydrous
 10 dichloromethane (50 mL) at room temperature was added *meta*-chloroperbenzoic acid (77%, 3.42 g, 19.8 mmol) in one portion. After being stirred for 20 h, powdered potassium carbonate (2.0 g) was added and the mixture was stirred for 1 h at room temperature before it was filtered and concentrated to afford Cap-138, Step b as a pale, yellow solid which was sufficiently pure to carry forward (2.15 g, 83.3%). ^1H NMR
 15 (CDCl_3 , 400 MHz) δ 8.73 (d, $J = 1.5$ Hz, 1H), 8.11 (dd, $J = 7.3, 1.7$ Hz, 1H), 8.04 (d, $J = 7.1$ Hz, 1H), 7.52 (t, $J = 8.1$ Hz, 1H), 7.28 (d, $J = 8.3$ Hz, 1H), 6.91 (d, $J = 7.8$ Hz, 1H), 4.00 (s, 3H); $R_t = 0.92$ min, (Cond.-D1); 90% homogeneity index; LC-MS: Anal. Calc. for $[\text{M}+\text{H}]^+$ $\text{C}_{10}\text{H}_{10}\text{NO}_2$: 176.07; found: 176.0.

20


Cap-138, Step c



[00243] To a stirred solution of Cap-138, Step b (0.70 g, 4.00 mmol) and triethylamine (1.1 mL, 8.00 mmol) in dry acetonitrile (20 mL) at room temperature under nitrogen was added trimethylsilylcyanide (1.60 mL, 12.00 mmol). The mixture was heated at 75 °C for
 25 20 h before it was cooled to room temperature, diluted with ethyl acetate and washed with saturated sodium bicarbonate solution and brine prior to drying over Na_2SO_4 and solvent

concentration. The residue was flash chromatographed on silica gel (elution with 5% ethyl acetate/hexanes) to 25% ethyl acetate/hexanes to afford Cap-138, Step c (498.7 mg) as a white, crystalline solid along with 223 mg of additional Cap-138, Step c recovered from the filtrate. ¹H NMR (CDCl₃, 500 MHz) δ 8.63 (d, *J* = 5.5 Hz, 1H), 8.26 (d, *J* = 5.5 Hz, 1H), 7.88 (d, *J* = 8.5 Hz, 1H), 7.69 (t, *J* = 8.0 Hz, 1H), 7.08 (d, *J* = 7.5 Hz, 1H), 4.04 (s, 3H); *R*_t = 1.75 min, (Cond.-D1); 90% homogeneity index; LC-MS: Anal. Calc. for [M+H]⁺ C₁₁H₉N₂O: 185.07; found: 185.10.

Cap-138




10


[00244] Cap-138, Step c (0.45 g, 2.44 mmol) was treated with 5*N* sodium hydroxide solution (10 mL) and the resulting suspension was heated at 85 °C for 4 h, cooled to 25 °C, diluted with dichloromethane and acidified with 1*N* hydrochloric acid. The organic phase was separated, washed with brine, dried over Na₂SO₄, concentrated to ¼ volume and filtered to afford Cap-138 as a yellow solid (0.44g, 88.9%). ¹H NMR (DMSO-*d*₆, 400 MHz) δ 13.6 (br s, 1H), 8.56 (d, *J* = 6.0 Hz, 1H), 8.16 (d, *J* = 6.0 Hz, 1H), 8.06 (d, *J* = 8.8 Hz, 1H), 7.71-7.67 (m, 1H), 7.30 (d, *J* = 8.0 Hz, 1H), 4.02 (s, 3H); *R*_t = 0.70 min (Cond.-D1); 95% homogeneity index; LC-MS: Anal. Calc. for [M+H]⁺ C₁₁H₁₀NO₃: 204.07; found: 204.05.

20


Synthetic Strategy. Method B (derived from *Tetrahedron Letters*, 42:6707 (2001)).



Cap-139




Cap-139, Step a



- 5 [00245] To a thick-walled, screw-top vial containing an argon-degassed suspension of 1-chloro-6-methoxyisoquinoline (1.2 g, 6.2 mmol; prepared according to the procedure in WO 2003/099274), potassium cyanide (0.40 g, 6.2 mmol), 1,5-bis(diphenylphosphino)pentane (0.27 g, 0.62 mmol) and palladium (II) acetate (70 mg, 0.31 mmol) in anhydrous toluene (6 mL) was added *N,N,N',N'*-
- 10 tetramethylethylenediamine (0.29 mL, 2.48 mmol). The vial was sealed, heated at 150 °C for 22 h and then allowed to cool to 25 °C. The reaction mixture was diluted with ethyl acetate, washed with water and brine, dried over Na₂SO₄, filtered and concentrated. The residue was purified on silica gel eluting with 5% ethyl acetate/hexanes to 25% ethyl acetate/hexanes to afford Cap-139, Step a as a white solid (669.7 mg). ¹H NMR (CDCl₃,
- 15 500 MHz) δ 8.54 (d, *J* = 6.0 Hz, 1H), 8.22 (d, *J* = 9.0 Hz, 1H), 7.76 (d, *J* = 5.5 Hz, 1H), 7.41-7.39 (m, 1H), 7.13 (d, *J* = 2.0 Hz, 1H), 3.98 (s, 3H); *R*_t = 1.66 min (Cond.-D1); 90% homogeneity index; LC-MS: Anal. Calc. for [M+H]⁺ C₁₁H₉N₂O: 185.07; found: 185.20.

Cap-139




20

- [00246] Cap-139 was prepared from the basic hydrolysis of Cap-139, Step a with 5*N* NaOH according to the procedure described for Cap-138. ¹H NMR (400 MHz, DMSO-*d*₆) δ 13.63 (v br s, 1H), 8.60 (d, *J* = 9.3 Hz, 1H), 8.45 (d, *J* = 5.6 Hz, 1H), 7.95 (d, *J* = 5.9 Hz, 1H), 7.49 (d, *J* = 2.2 Hz, 1H), 7.44 (dd, *J* = 9.3, 2.5 Hz, 1H), 3.95 (s, 3H); *R*_t = 0.64


min (Cond.-D1); 90% homogeneity index; LC-MS: Anal. Calc. for $[M+H]^+$ $C_{11}H_{10}NO_3$: 204.07; found: 204.05.

Cap-140




5

Cap-140, Step a



- [00247]** To a vigorously-stirred mixture of 1,3-dichloro-5-ethoxyisoquinoline (482 mg, 2.00 mmol; prepared according to the procedure in WO 2005/051410), palladium (II) acetate (9 mg, 0.04 mmol), sodium carbonate (223 mg, 2.10 mmol) and 1,5-bis(diphenylphosphino)pentane (35 mg, 0.08 mmol) in dry dimethylacetamide (2 mL) at 25 °C under nitrogen was added N,N,N',N'-tetramethylethylenediamine (60 mL, 0.40 mmol). After 10 min, the mixture was heated to 150 °C, and then a stock solution of acetone cyanohydrin (prepared from 457 μ L of acetone cyanohydrin in 4.34 mL DMA) was added in 1 mL portions over 18 h using a syringe pump. The mixture was then partitioned between ethyl acetate and water and the organic layer was separated, washed with brine, dried over Na_2SO_4 , filtered and concentrated. The residue was purified on silica gel eluting with 10% ethyl acetate/hexanes to 40% ethyl acetate/hexanes to afford Cap-140, Step a as a yellow solid (160 mg, 34%). R_t = 2.46 min (Cond.-MS-W2); 90% homogeneity index; LC-MS: Anal. Calc. for $[M+H]^+$ $C_{12}H_9ClN_2O$: 233.05; found: 233.08.


Cap-140



[00248] Cap-140 was prepared by the acid hydrolysis of Cap-140, Step a with 12*N* HCl as described in the procedure for the preparation of Cap-141, described below. $R_t = 2.24$ min (Cond.-MS-W2); 90% homogeneity index; LC-MS: Anal. Calc. for $[M+H]^+$


5 $C_{12}H_{11}ClNO_3$: 252.04; found: 252.02.

Cap-141



10


Cap-141, Step a



[00249] Cap-141, Step a was prepared from 1-bromo-3-fluoroisoquinoline (prepared from 3-amino-1-bromoisoquinoline using the procedure outlined in *J. Med. Chem.*, 13:613 (1970)) as described in the procedure for the preparation of Cap-140, Step a (*vide supra*). 1H NMR (500 MHz, $CDCl_3$) δ 8.35 (d, $J = 8.5$ Hz, 1H), 7.93 (d, $J = 8.5$ Hz, 1H), 7.83 (t, $J = 7.63$ Hz, 1H), 7.77-7.73 (m, 1H), 7.55 (s, 1H); $R_t = 1.60$ min (Cond.-D1); 90% homogeneity index; LC-MS: Anal. Calc. for $[M+H]^+$ $C_{10}H_6FN_2$: 173.05; found: 172.99.

15

Cap-141




20

[00250] Cap-141, Step a (83 mg, 0.48 mmol) was treated with 12*N* HCl (3 mL) and the resulting slurry was heated at 80 °C for 16 h before it was cooled to room temperature


and diluted with water (3 mL). The mixture was stirred for 10 min and then filtered to afford Cap-141 as an off-white solid (44.1 mg, 47.8%). The filtrate was diluted with dichloromethane and washed with brine, dried over Na₂SO₄, and concentrated to afford additional Cap-141 which was sufficiently pure to be carried forward directly (29.30 mg, 31.8%). ¹H NMR (DMSO-d₆, 500 MHz) δ 14.0 (br s, 1H), 8.59-8.57 (m, 1H), 8.10 (d, *J* = 8.5 Hz, 1H), 7.88-7.85 (m, 2H), 7.74-7.71 (m, 1H); *R*_t = 1.33 min (Cond.-D1); 90% homogeneity index; LC-MS: Anal. Calc. for [M+H]⁺ C₁₀H₇FNO₂: 192.05; found: 191.97.

Cap-142



10


Cap-142, Step a



[00251] Cap-142, Step a was prepared from 4-bromoisquinoline N-oxide as described in the two-step procedure for the preparation of Cap-138, steps b and c. *R*_t = 1.45 min (Cond.-MS-W1); 90% homogeneity index; LC-MS: Anal. Calc. for [M+H]⁺ C₁₀H₆BrN₂: 232.97; found: 233.00.

15

Cap-142, Step b




20

[00252] To an argon-degassed suspension of Cap-142, Step a (116 mg, 0.50 mmol), potassium phosphate tribasic (170 mg, 0.80 mmol), palladium (II) acetate (3.4 mg, 0.015 mmol) and 2-(dicyclohexylphosphino)biphenyl (11 mg, 0.03 mmol) in anhydrous toluene (1 mL) was added morpholine (61 μ L, 0.70 mmol). The mixture was heated at 100 °C for 16 h, cooled to 25 °C and filtered through diatomaceous earth (CELITE®). Purification of the residue on silica gel, eluting with 10% to 70% ethyl acetate/hexanes afforded Cap-142, Step b (38 mg, 32%) as a yellow solid, which was carried forward directly. R_t = 1.26 min (Cond.-MS-W1); 90% homogeneity index; LC-MS: Anal. Calc. for $[M+H]^+$ $C_{14}H_{14}N_3O$: 240.11; found: 240.13.

10


Cap-142



[00253] Cap-142 was prepared from Cap-142, Step b with 5*N* sodium hydroxide as described in the procedure for Cap-138. R_t = 0.72 min (Cond.-MS-W1); 90% homogeneity index; LC-MS: Anal. Calc. for $[M+H]^+$ $C_{14}H_{15}N_2O_3$: 259.11; found: 259.08.


15

Cap-143




20

Cap-143, Step a



[00254] To a stirred solution of 3-amino-1-bromoisoquinoline (444 mg, 2.00 mmol) in anhydrous dimethylformamide (10 mL) was added sodium hydride (60%, unwashed, 96 mg, 2.4 mmol) in one portion. The mixture was stirred at 25 °C for 5 min before 2-bromoethyl ether (90%, 250 µL, 2.00 mmol) was added. The mixture was stirred further at 25 °C for 5 h and at 75 °C for 72 h before it was cooled to 25 °C, quenched with saturated ammonium chloride solution and diluted with ethyl acetate. The organic layer was separated, washed with water and brine, dried over Na₂SO₄, filtered and concentrated. Purification of the residue on silica gel eluting with 0% to 70% ethyl acetate/hexanes afforded Cap-143, Step a as a yellow solid (180 mg, 31%). *R*_t = 1.75 min (Cond.-MS-W1); 90% homogeneity index; LC-MS: Anal. Calc. for [M+H]⁺ C₁₃H₁₄BrN₂O: 293.03; found: 293.04.


Cap-143




[00255] To a cold (-60 °C) solution of Cap-143, Step a (154 mg, 0.527 mmol) in anhydrous tetrahydrofuran (5 mL) was added a solution of *n*-butyllithium in hexanes (2.5 M, 0.25 mL, 0.633 mmol). After 10 min, dry carbon dioxide was bubbled into the reaction mixture for 10 min before it was quenched with 1N HCl and allowed to warm to 25 °C. The mixture was then extracted with dichloromethane (3 x 30 mL) and the combined organic extracts were concentrated *in vacuo*. Purification of the residue by a reverse phase HPLC (MeOH/water/TFA) afforded Cap-144 (16 mg, 12%). *R*_t = 1.10 min (Cond.-MS-W1); 90% homogeneity index; LC-MS: Anal. Calc. for [M+H]⁺ C₁₄H₁₅N₂O₃: 259.11; found: 259.08.

25

Cap-144




Cap-144, Step a



- 5 [00256] 1,3-Dichloroisoquinoline (2.75 g, 13.89 mmol) was added in small portions to a cold (0 °C) solution of fuming nitric acid (10 mL) and concentrated sulfuric acid (10 mL). The mixture was stirred at 0 °C for 0.5 h before it was gradually warmed to 25 °C where it stirred for 16 h. The mixture was then poured into a beaker containing chopped ice and water and the resulting suspension was stirred for 1 h at 0 °C before it was filtered
- 10 to afford Cap-144, Step a (2.73 g, 81%) as a yellow solid which was used directly. $R_t = 2.01$ min. (Cond.-D1); 95% homogeneity index; LC-MS: Anal. Calc. for $[M+H]^+$ $C_9H_5Cl_2N_2O_2$: 242.97; found: 242.92.


Cap-144, Step b



- 15 [00257] Cap-144, Step a (0.30 g, 1.23 mmol) was taken up in methanol (60 mL) and treated with platinum oxide (30 mg), and the suspension was subjected to Parr hydrogenation at 7 psi H_2 for 1.5 h. Then formalin (5 mL) and additional platinum oxide (30 mg) were added, and the suspension was resubjected to Parr hydrogenation at 45 psi
- 20 H_2 for 13 h. It was then suction-filtered through diatomaceous earth (CELITE®) and concentrated down to $\frac{1}{4}$ volume. Suction-filtration of the ensuing precipitate afforded the

title compound as a yellow solid which was flash chromatographed on silica gel eluting with 5% ethyl acetate in hexanes to 25% ethyl acetate in hexanes to afford Cap-144, Step b (231 mg, 78%) as a pale yellow solid. $R_t = 2.36$ min (Cond.-D1); 95% homogeneity index; ^1H NMR (400 MHz, CDCl_3) δ 8.02 (s, 1H), 7.95 (d, $J = 8.6$ Hz, 1H), 7.57-7.53 (m, 1H), 7.30 (d, $J = 7.3$ Hz, 1H), 2.88 (s, 6H); LC-MS: Anal. Calc. for $[\text{M}+\text{H}]^+$ $\text{C}_{11}\text{H}_{11}\text{Cl}_2\text{N}_2$: 241.03; found: 241.02. HRMS: Anal. Calc. for $[\text{M}+\text{H}]^+$ $\text{C}_{11}\text{H}_{11}\text{Cl}_2\text{N}_2$: 241.0299; found: 241.0296.

Cap-144, Step c




10

[00258] Cap-144, Step c was prepared from Cap-144, Step b according to the procedure described for the preparation of Cap-139, Step a. $R_t = 2.19$ min (Cond.-D1); 95% homogeneity index; LC-MS: Anal. Calc. for $[\text{M}+\text{H}]^+$ $\text{C}_{12}\text{H}_{11}\text{ClN}_3$: 232.06; found: 232.03. HRMS: Anal. Calc. for $[\text{M}+\text{H}]^+$ $\text{C}_{12}\text{H}_{11}\text{ClN}_3$: 232.0642; found: 232.0631.

15

Cap-144






[00259] Cap-144 was prepared according to the procedure described for Cap-141. $R_t = 2.36$ min (Cond.-D1); 90%; LC-MS: Anal. Calc. for $[\text{M}+\text{H}]^+$ $\text{C}_{12}\text{H}_{12}\text{ClN}_2\text{O}_2$: 238.01; found: 238.09.




20




Cap-145 to Cap-162




[00260] Cap-145 to Cap-162 were prepared from the appropriate 1-chloroisoquinolines according to the procedure described for the preparation of Cap-138 (Method A) or Cap-139 (Method B) unless noted otherwise as outlined below.




25




Cap-#	Cap	Method	Hydrolysis	R _t (LC-Cond.); % homogeneity index; MS data
Cap-145	 <p>Prepared from commercially available 1,3- dichloroisoquinoline</p>	B	12 <i>N</i> HCl	1.14 min (Cond.- MS-W1); 90%; LC-MS: Anal. Calc. for [M+H] ⁺ C ₁₀ H ₇ ClNO ₂ : 208.02; found: 208.00.
Cap-146	 <p>Prepared from commercially available 3-hydroxyisoquinoline</p>	A	5 <i>N</i> NaOH	1.40 min (Cond.- D1); 95%; LC- MS: Anal. Calc. for [M+H] ⁺ C ₁₁ H ₁₀ NO ₃ : 204.07; found: 204.06.
Cap-147	 <p>Prepared from commercially available 1-chloro-4- hydroxyisoquinoline</p>	B	5 <i>N</i> NaOH	0.87 min (Cond.- D1); 95%; LC- MS: Anal. Calc. for [M+H] ⁺ C ₁₁ H ₁₀ NO ₃ : 204.07; found: 204.05.

Cap-#	Cap	Method	Hydrolysis	R _t (LC-Cond.); % homogeneity index; MS data
Cap-148	 <p>Prepared from commercially available 7-hydroxyisoquinoline</p>	A	5 <i>N</i> NaOH	0.70 min (Cond.-D1); 95%; LC-MS: Anal. Calc. for [M+H] ⁺ C ₁₁ H ₁₀ NO ₃ : 204.07; found: 204.05.
Cap-149	 <p>Prepared from commercially available 5-hydroxyisoquinoline</p>	A	5 <i>N</i> NaOH	0.70 min (Cond.-D1); 95%; LC-MS: Anal. Calc. for [M+H] ⁺ C ₁₁ H ₁₀ NO ₃ : 204.07; found: 204.05.
Cap-150	 <p>Prepared from 8-methoxy-1-chloroisoquinoline, which can be synthesized following the procedure in WO 2003/099274</p>	A	12 <i>N</i> HCl	0.26 min (Cond.-D1); 95%; LC-MS: Anal. Calc. for [M+H] ⁺ C ₁₁ H ₁₀ NO ₃ : 204.07; found: 204.04.


Cap-#	Cap	Method	Hydrolysis	R _t (LC-Cond.); % homogeneity index; MS data
Cap-151 3-chloro-5-methoxyisoquinoline-1-carboxylic acid	 <p>Prepared from 5-methoxy-1,3-dichloroisoquinoline, which can be synthesized following the procedure in WO 2005/051410</p>	B	12N HCl	1.78 min (Cond.-D1); 90%; LC-MS: Anal. Calc. for [M+H] ⁺ C ₁₁ H ₉ ClNO ₃ : 238.03; found: 238.09.
Cap-152	 <p>Prepared from commercially available 6-methoxy-1,3-dichloroisoquinoline</p>	B	12N HCl	1.65 min (Cond.-D1); 95%; LC-MS: Anal. Calc. for [M+H] ⁺ C ₁₁ H ₉ ClNO ₃ : 238.00; found: 238.09.
Cap-153	 <p>Prepared from 4-bromoisoquinoline, which can be synthesized following the procedure in WO 2003/062241</p>	A	6N HCl	1.18 min (Cond.-MS-W1); 95%; LC-MS: Anal. Calc. for [M+H] ⁺ C ₁₀ H ₇ BrNO ₂ : 251.97; found: 251.95.

Cap-#	Cap	Method	Hydrolysis	R _t (LC-Cond.); % homogeneity index; MS data
Cap-154	 <p>Prepared from 7-fluoro-1-chloroisoquinoline, which can be synthesized following the procedure in WO 2003/099274</p>	B	5N NaOH	0.28 min (Cond.-MS-W1); 90%; LC-MS: Anal. Calc. for [M+H] ⁺ C ₁₀ H ₇ FNO ₂ : 192.05; found: 192.03.
Cap-155	 <p>Prepared from 1,7-dichloroisoquinoline, which can be synthesized following the procedure in WO 2003/099274</p>	B	5N NaOH	0.59 min (Cond.-MS-W1); 90%; LC-MS: Anal. Calc. for [M+H] ⁺ C ₁₀ H ₇ ClNO ₂ : 208.02; found: 208.00.
Cap-156	 <p>Prepared from 1,6-dichloroisoquinoline, which can be synthesized following the procedure in WO 2003/099274</p>	B	5N NaOH	0.60 min (Cond.-MS-W1); 90%; LC-MS: Anal. Calc. for [M+H] ⁺ C ₁₀ H ₇ ClNO ₂ : 208.02; found: 208.03.

Cap-#	Cap	Method	Hydrolysis	R _t (LC-Cond.); % homogeneity index; MS data
Cap-157	 <p>Prepared from 1,4-dichloroisoquinoline, which can be synthesized following the procedure in WO 2003/062241</p>	B	12N HCl	1.49 min (Cond.-D1); 95%; LC-MS: Anal. Calc. for [M+H] ⁺ C ₁₀ H ₇ ClNO ₂ : 208.02; found: 208.00.
Cap-158	 <p>Prepared from 1,5-dichloroisoquinoline, which can be synthesized following the procedure in WO 2003/099274</p>	B	5N NaOH	0.69 min (Cond.-MS-W1); 90%; LC-MS: Anal. Calc. for [M+H] ⁺ C ₁₀ H ₇ ClNO ₂ : 208.02; found: 208.01.
Cap-159	 <p>Prepared from 5-fluoro-1-chloroisoquinoline, which can be synthesized following the procedure in WO 2003/099274</p>	B	5N NaOH	0.41 min (Cond.-MS-W1); 90%; LC-MS: Anal. Calc. for [M+H] ⁺ C ₁₀ H ₇ FNO ₂ : 192.05; found: 192.03.


Cap-#	Cap	Method	Hydrolysis	R _t (LC-Cond.); % homogeneity index; MS data
Cap-160	 <p>Prepared from 6-fluoro-1-chloroisoquinoline, which can be synthesized following the procedure in WO 2003/099274</p>	B	5N NaOH	0.30 min (Cond.-MS-W1); 90%; LC-MS: Anal. Calc. for [M+H] ⁺ C ₁₀ H ₇ FNO ₂ : 192.05; found: 192.03.
Cap-161	 <p>Prepared from 4-bromoquinoline-2-carboxylic acid and dimethylamine (DMSO, 100 °C)</p>	--	--	0.70 min (Cond. D1); 95%; LC-MS: Anal. Calc. for [M+H] ⁺ C ₁₂ H ₁₃ N ₂ O ₂ : 217.10; found: 217.06.
Cap-162	 <p>Prepared from m-anisidine following the procedure described in <i>J. Hetero. Chem.</i>, 17 (1993) and <i>Heterocycles</i>, 60:953 (2003).</p>	--	--	0.65 min (Cond.-M3); 95%; LC-MS: Anal. Calc. for [M+H] ⁺ C ₁₁ H ₁₀ NO ₃ : 204.07; found: 203.94.

Cap-163




[00261] To a solution of 2-ketobutyric acid (1.0 g, 9.8 mmol) in diethylether (25 ml) was added phenylmagnesium bromide (22 ml, 1M in THF) dropwise. The reaction was stirred at ~25 °C under nitrogen for 17.5h. The reaction was acidified with 1N HCl and the product was extracted with ethyl acetate (3 x 100 ml). The combined organic layer was washed with water followed by brine and dried over MgSO₄. After concentration *in vacuo*, a white solid was obtained. The solid was recrystallized from hexanes/ethyl acetate to afford Cap-163 as white needles (883.5 mg). ¹H NMR (DMSO-d₆, δ = 2.5 ppm, 500 MHz): 12.71 (br s, 1 H), 7.54-7.52 (m, 2H), 7.34-7.31 (m, 2H), 7.26-7.23 (m, 1H), 5.52-5.39 (br s, 1H), 2.11 (m, 1H), 1.88 (m, 1H), 0.79 (app t, J = 7.4 Hz, 3H).

Cap-164



[00262] A mixture of 2-amino-2-phenylbutyric acid (1.5 g, 8.4 mmol), formaldehyde (14 mL, 37% in water), 1N HCl (10 mL) and 10% Pd/C (0.5 mg) in MeOH (40 mL) was exposed to H₂ at 50 psi in a Parr bottle for 42 h. The reaction was filtered over CELITE® and concentrated *in vacuo*, the residue was taken up in MeOH (36 mL) and the product was purified with a reverse phase HPLC (MeOH/H₂O/TFA) to afford the TFA salt of Cap-164 as a white solid (1.7 g). ¹H NMR (DMSO-d₆, δ = 2.5 ppm, 500 MHz) 7.54-7.47 (m, 5H), 2.63 (m, 1H), 2.55 (s, 6H), 2.31 (m, 1H), 0.95 (app t, J = 7.3 Hz, 3H).


Cap-165



[00263] To a mixture of 2-amino-2-indanecarboxylic acid (258.6 mg, 1.46 mmol) and formic acid (0.6 ml, 15.9 mmol) in 1,2-dichloroethane (7 ml) was added formaldehyde (0.6 ml, 37% in water). The mixture was stirred at ~25 °C for 15 min then heated at 70

°C for 8h. The volatile component was removed *in vacuo*, and the residue was dissolved in DMF (14 mL) and purified by a reverse phase HPLC (MeOH/H₂O/TFA) to afford the TFA salt of Cap-165 as a viscous oil (120.2 mg). ¹H NMR (DMSO-d₆, δ = 2.5 ppm, 500 MHz): 7.29-7.21 (m, 4 H), 3.61 (d, J = 17.4 Hz, 2H), 3.50 (d, J = 17.4 Hz, 2H), 2.75 (s, 6H). LC-MS: Anal. Calcd. for [M+H]⁺ C₁₂H₁₆NO₂: 206.12; found: 206.07.


Cap-166a and Cap-166b



Cap-166a: Diastereomer-1
Cap-166b: Diastereomer-2

[00264] Cap-166a and Cap-166b were prepared from (1S, 4S)-(+)-2-methyl-2,5-diazabicyclo[2.2.1]heptane (2HBr) according to the method described for the synthesis of Cap-7a and Cap-7b, with the exception that the benzyl ester intermediate was separated using a semi-prep Chiralcel OJ column, 20 x 250 mm, 10 μm eluting with 85:15 heptane/ethanol mixture at 10 mL/min elution rate for 25 min. Cap-166b: ¹H NMR (DMSO-d₆, δ = 2.5 ppm, 500 MHz): 7.45 (d, J = 7.3 Hz, 2H), 7.27-7.19 (m, 3H), 4.09 (s, 1H), 3.34 (app br s, 1H), 3.16 (app br s, 1H), 2.83 (d, J=10.1 Hz, 1H), 2.71 (m, 2H), 2.46 (m, 1H), 2.27 (s, 3H), 1.77 (d, J = 9.8 Hz, 1H), 1.63 (d, J = 9.8 Hz, 1H). LC-MS: Anal. Calcd. for [M+H]⁺ C₁₄H₁₉N₂O₂: 247.14; found: 247.11.

Cap-167




[00265] A solution of racemic Boc-1,3-dihydro-2H-isoindole carboxylic acid (1.0g, 3.8 mmol) in 20% TFA/CH₂Cl₂ was stirred at ~25 °C for 4h. All the volatile component was removed *in vacuo*. A mixture of the resultant crude material, formaldehyde (15 mL, 37% in water), 1N HCl (10 mL) and 10% Pd/C (10 mg) in MeOH was exposed to H₂ (40 PSI) in a Parr bottle for 23 h. The reaction mixture was filtered over CELITE® and concentrated *in vacuo* to afford Cap-167 as a yellow foam (873.5 mg). ¹H NMR (DMSO-

d_6 , $\delta = 2.5$ ppm, 500 MHz) 7.59-7.38 (m, 4H), 5.59 (s, 1H), 4.84 (d, $J = 14$ Hz, 1H), 4.50 (d, $J = 14.1$ Hz, 1H), 3.07 (s, 3H). LC-MS: Anal. Calcd. for $[M+H]^+$ $C_{10}H_{12}NO_2$: 178.09; found: 178.65.

5


Cap-168



[00266] Racemic Cap-168 was prepared from racemic Boc-aminoindane-1-carboxylic acid according to the procedure described for the preparation of Cap-167. The crude material was employed as such.


10

Cap-169



[00267] A mixture of 2-amino-2-phenylpropanoic acid hydrochloride (5.0 g, 2.5 mmol), formaldehyde (15 ml, 37% in water), 1N HCl (15 ml), and 10% Pd/C (1.32 g) in MeOH (60 mL) was placed in a Parr bottle and shaken under hydrogen (55 PSI) for 4 days. The reaction mixture was filtered over CELITE® and concentrated *in vacuo*. The residue was taken up in MeOH and purified by reverse phase prep-HPLC (MeOH/water/TFA) to afford the TFA salt of Cap-169 as a viscous semi-solid (2.1 g). 1H NMR ($CDCl_3$, $\delta = 7.26$ ppm, 500 MHz): 7.58-7.52 (m, 2 H), 7.39-7.33 (m, 3H), 2.86 (br s, 3H), 2.47 (br s, 3H), 1.93 (s, 3H). LC-MS: Anal. Calcd. for $[M+H]^+$ $C_{11}H_{16}NO_2$: 194.12; found: 194.12.


Cap-170



(S)-2-(Methoxycarbonylamino)-2-(tetrahydro-2H-pyran-4-yl)acetic acid

[00268] To (S)-2-amino-2-(tetrahydro-2H-pyran-4-yl)acetic acid (505mg; 3.18mmol; obtained from Astatech) in water (15ml) was added sodium carbonate (673mg; 6.35mmol), and the resultant mixture was cooled to 0 °C and then methyl chloroformate (0.26ml; 3.33mmol) was added dropwise over 5 minutes. The reaction was allowed to stir for 18 hours while allowing the bath to thaw to ambient temperature. The reaction mixture was then partitioned between 1N HCl and ethyl acetate. The organic layer was removed and the aqueous layer was further extracted with 2 additional portions of ethyl acetate. The combined organic layers were washed with brine, dried over magnesium sulfate, filtered and concentrated *in vacuo* to afford Cap-170 a colorless residue. ¹H NMR (500 MHz, DMSO-d₆) δ ppm 12.65 (1 H, br s), 7.44 (1 H, d, *J*=8.24 Hz), 3.77 - 3.95 (3 H, m), 3.54 (3 H, s), 3.11 - 3.26 (2 H, m), 1.82 - 1.95 (1 H, m), 1.41 - 1.55 (2 H, m), 1.21 - 1.39 (2 H, m); LC-MS: Anal. Calcd. for [M+H]⁺ C₉H₁₆NO₅: 218.1; found 218.1.

Cap-171




[00269] A solution of methyl 2-(benzyloxycarbonylamino)-2-(oxetan-3-ylidene)acetate (200 mg, 0.721 mmol; *Il Farmaco*, 56:609-613 (2001)) in ethyl acetate (7 ml) and CH₂Cl₂ (4.00 ml) was degassed by bubbling nitrogen for 10min. Dimethyl dicarbonate (0.116 ml, 1.082 mmol) and Pd/C (20 mg, 0.019 mmol) were then added, the reaction mixture was fitted with a hydrogen balloon and allowed to stir at ambient temperature overnight at which time TLC (95:5 CH₂Cl₂ / MeOH: visualized with stain made from 1g Ce(NH₄)₂SO₄, 6g ammonium molybdate, 6ml sulfuric acid, and 100ml water) indicated complete conversion. The reaction was filtered through CELITE® and concentrated. The residue was purified via BIOTAGE® (load with dichloromethane on 25 samplet; elute on 25S column with dichloromethane for 3CV then 0 to 5% MeOH / dichloromethane over 250ml then hold at 5% MeOH / dichloromethane for 250ml; 9ml fractions). Collected fractions containing desired material and concentrated to 120mg

(81%) of methyl 2-(methoxycarbonylamino)-2-(oxetan-3-yl)acetate as a colorless oil. ¹H NMR (500 MHz, chloroform-d) δ ppm 3.29 - 3.40 (m, *J*=6.71 Hz, 1 H) 3.70 (s, 3 H) 3.74 (s, 3 H) 4.55 (t, *J*=6.41 Hz, 1 H) 4.58 - 4.68 (m, 2 H) 4.67 - 4.78 (m, 2 H) 5.31 (br s, 1 H). LC-MS: Anal. Calcd. for [M+H]⁺ C₈H₁₄NO₅: 204.2; found 204.0.


- 5 **[00270]** To methyl 2-(methoxycarbonylamino)-2-(oxetan-3-yl)acetate (50 mg, 0.246 mmol) in THF (2 mL) and water (0.5 mL) was added lithium hydroxide monohydrate (10.33 mg, 0.246 mmol). The resultant solution was allowed to stir overnight at ambient temperature. TLC (1:1 EA / Hex; Hanessian stain [1g Ce(NH₄)₂SO₄, 6g ammonium molybdate, 6ml sulfuric acid, and 100ml water]) indicated ~10% starting material
- 10 remaining. Added an additional 3mg LiOH and allowed to stir overnight at which time TLC showed no starting material remaining. Concentrated *in vacuo* and placed on high vac overnight providing 55 mg lithium 2-(methoxycarbonylamino)-2-(oxetan-3-yl)acetate as a colorless solid. ¹H NMR (500 MHz, MeOD) δ ppm 3.39 - 3.47 (m, 1 H) 3.67 (s, 3 H) 4.28 (d, *J*=7.93 Hz, 1 H) 4.64 (t, *J*=6.26 Hz, 1 H) 4.68 (t, *J*=7.02 Hz, 1 H) 4.73 (d,
- 15 *J*=7.63 Hz, 2 H).

Cap-172



20

Cap-172, Step a




- [00271]** The following diazotization step was adapted from Barton, A. et al., *J.C.S. Perkin Trans I*, 159-164 (1982): A solution of NaNO₂ (166 mg, 2.4 mmol) in water (0.6 mL) was added slowly to a stirred, cold (0 °C) solution of methyl 2-amino-5-ethyl-1,3-thiazole-4-carboxylate (186 mg, 1.0 mmol), CuSO₄•5H₂O (330 mg, 1.32 mmol), NaCl (260 mg, 4.45 mmol) and H₂SO₄ (5.5 mL) in water (7.5 mL). The mixture was stirred at
- 25 0 °C for 45 min and allowed to warm up to room temperature where it stirred further for 1 h before CuCl (118 mg) was added. This mixture was stirred further at room temperature

for 16 h before it was diluted with brine and extracted with ether twice. The organic layers were combined, dried over MgSO_4 and concentrated to give methyl 2-chloro-5-ethylthiazole-4-carboxylate (*i.e.*, Cap-172, Step a) (175 mg, 85%) as an orange oil (80% pure) which was used directly in the next reaction. $R_t = 1.99$ min (Cond.-MD1); LC-MS: Anal. Calcd. for $[\text{M}+\text{H}]^+$ $\text{C}_7\text{H}_9\text{ClNO}_2\text{S}$: 206.01; found: 206.05.


Cap-172

[00272] To a solution of methyl 2-chloro-5-ethylthiazole-4-carboxylate (175 mg) in THF/ H_2O /MeOH (20 mL/ 3 mL/ 12 mL) was added LiOH (305 mg, 12.76 mmol). The mixture was stirred at room temperature overnight before it was concentrated down and neutralized with 1N HCl in ether (25 mL). The residue was extracted twice with ethyl acetate and the organic layers were combined, dried over MgSO_4 and evaporated to yield Cap-172 (60 mg, 74%) as a red solid which was used without further purification. ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ ppm 13.03-13.42 (1 H, m), 3.16 (2 H, q, $J = 7.4$ Hz), 1.23 (3 H, t, $J = 7.5$ Hz). $R_t = 1.78$ min (Cond.-MD1); LC-MS: Anal. Calcd. for $[\text{M}+\text{H}]^+$ $\text{C}_6\text{H}_7\text{ClNO}_2\text{S}$: 191.99; found: 191.99.

Cap-173



Cap-173, Step a




[00273] The following diazotization step was adapted from Barton, A. et al., *J.C.S. Perkin Trans I*, 159-164 (1982): A solution of NaNO_2 (150 mg, 2.17 mmol) in water (1.0 mL) was added dropwise to a stirred, cold (0°C) solution of methyl 2-amino-5-ethyl-1,3-thiazole-4-carboxylate (186 mg, 1.0 mmol) in 50% H_3PO_2 (3.2 mL). The mixture was stirred at 0°C for 1 h and allowed to warm up to room temperature where it stirred further for 2h. After recooling to 0°C , the mixture was treated slowly with a solution of NaOH

(85 mg) in water (10 mL). The mixture was then diluted with saturated NaHCO₃ solution and extracted twice with ether. The organic layers were combined, dried over MgSO₄ and concentrated to give methyl 5-ethylthiazole-4-carboxylate (*i.e.*, Cap-173, Step a) (134 mg, 78%) as an orange oil (85% pure) which was used directly in the next reaction. R_t = 1.58 min (Cond.-MD1); LC-MS: Anal. Calcd. for [M+H]⁺ C₇H₁₀NO₂S: 172.05; found: 172.05.

Cap-173


[00274] To a solution of methyl 5-ethylthiazole-4-carboxylate (134 mg) in THF/H₂O/MeOH (18 mL/ 2.7 mL/ 11 mL) was added LiOH (281 mg, 11.74 mmol). The mixture was stirred at room temperature overnight before it was concentrated down and neutralized with 1N HCl in ether (25 mL). The residue was extracted twice with ethyl acetate and the organic layers were combined, dried over MgSO₄ and evaporated to yield Cap-173 (90 mg, 73%) as an orange solid which was used without further purification. ¹H NMR (300 MHz, DMSO-d₆) δ ppm 12.74-13.04 (1 H, m), 3.20 (2 H, q, *J* = 7.3 Hz), 1.25 (3 H, t, *J* = 7.5 Hz). R_t = 1.27 min (Cond.-MD1); LC-MS: Anal. Calcd. for [M+H]⁺ C₆H₈NO₂S: 158.03; found: 158.04.

Cap-174



20

Cap-174, Step a



[00275] Triflic anhydride (5.0 g, 18.0 mmol) was added dropwise to a cold (0 °C) solution of methyl 3-hydroxypicolinate (2.5 g, 16.3 mmol) and TEA (2.5 mL, 18.0 mmol) in CH₂Cl₂ (80 mL). The mixture was stirred at 0 °C for 1h before it was allowed to warm up to room temperature where it stirred for an additional 1 h. The mixture was then


quenched with saturated NaHCO₃ solution (40 mL) and the organic layer was separated, washed with brine, dried over MgSO₄ and concentrated to give methyl 3-(trifluoromethylsulfonyloxy)picolinate (*i.e.*, Cap-174, Step a) (3.38 g, 73%) as a dark brown oil (>95% pure) which was used directly without further purification. ¹H NMR (300 MHz, CDCl₃) δ ppm 8.72-8.79 (1 H, m), 7.71 (1 H, d, *J* = 1.5 Hz), 7.58-7.65 (1 H, m), 4.04 (3 H, s). *R*_t = 1.93 min (Cond.-MD1); LC-MS: Anal. Calcd. for [M+H]⁺ C₈H₇F₃NO₅S: 286.00; found: 286.08.

Cap-174


10 [00276] To a solution of methyl 3-(trifluoromethylsulfonyloxy)picolinate (570 mg, 2.0 mmol) in DMF (20 mL) was added LiCl (254 mg, 6.0 mmol), tributyl(vinyl)stannane (761 mg, 2.4 mmol) and bis(triphenylphosphine)palladium dichloride (42 mg, 0.06 mmol). The mixture was heated at 100 °C overnight before a saturated solution of KF (20 mL) was added to the reaction mixture at room temperature. This mixture was stirred for 15 4 h before it was filtered through CELITE® and the pad of CELITE® was washed with ethyl acetate. The aqueous phase of the filtrate was then separated and concentrated down *in vacuo*. The residue was treated with 4*N* HCl in dioxanes (5 mL) and the resulting mixture was extracted with methanol, filtered and evaporated to afford Cap-174 (260 mg) as a green solid which was slightly contaminated with inorganic salts but was 20 used without further purification. ¹H NMR (300 MHz, DMSO-*d*₆) δ ppm 8.21 (1 H, d, *J* = 3.7 Hz), 7.81-7.90 (1 H, m), 7.09 (1 H, dd, *J* = 7.7, 4.8 Hz), 6.98 (1 H, dd, *J* = 17.9, 11.3 Hz), 5.74 (1 H, dd, *J* = 17.9, 1.5 Hz), 5.20 (1 H, d, *J* = 11.0 Hz). *R*_t = 0.39 min (Cond.-MD1); LC-MS: Anal. Calcd. for [M+H]⁺ C₈H₈NO₂: 150.06; found: 150.07.

25

Cap-175




Cap-175, Step a



- [00277] To a solution of methyl 3-(trifluoromethylsulfonyloxy)picolinate (*i.e.*, Cap-174, Step a) (570 mg, 2.0 mmol), an intermediate in the preparation of Cap-174, in DMF (20 mL) was added LiCl (254 mg, 6.0 mmol), tributyl(vinyl)stannane (761 mg, 2.4 mmol) and bis(triphenylphosphine)palladium dichloride (42 mg, 0.06 mmol). The mixture was heated at 100 °C for 4 h before the solvent was removed *in vacuo*. The residue was taken up in acetonitrile (50 mL) and hexanes (50 mL) and the resulting mixture was washed twice with hexanes. The acetonitrile layer was then separated, filtered through CELITE®, and evaporated. Purification of the residue by flash chromatography on a Horizon instrument (gradient elution with 25% ethyl acetate in hexanes to 65% ethyl acetate in hexanes) afforded methyl 3-vinylpicolinate (*i.e.*, Cap-175, Step a) (130 mg, 40%) as a yellow oil. ¹H NMR (300 MHz, CDCl₃) δ ppm 8.60 (1 H, dd, *J* = 4.6, 1.7 Hz), 7.94 (1 H, d, *J* = 7.7 Hz), 7.33-7.51 (2 H, m), 5.72 (1 H, d, *J* = 17.2 Hz), 5.47 (1 H, d, *J* = 11.0 Hz), 3.99 (3 H, s). *R*_t = 1.29 min (Cond.-MD1); LC-MS: Anal. Calcd. for [M+H]⁺ C₉H₁₀NO₂: 164.07; found: 164.06.


Cap-175, Step b




- [00278] Palladium on carbon (10%, 25 mg) was added to a solution of methyl 3-vinylpicolinate (120 mg, 0.74 mmol) in ethanol (10 mL). The suspension was stirred at room temperature under an atmosphere of hydrogen for 1 h before it was filtered through CELITE® and the pad of CELITE® was washed with methanol. The filtrate was concentrated down to dryness to yield methyl 3-ethylpicolinate (*i.e.*, Cap-175, Step b) which was taken directly into the next reaction. *R*_t = 1.15 min (Cond.-MD1); LC-MS: Anal. Calcd. for [M+H]⁺ C₉H₁₂NO₂: 166.09; found: 166.09.

Cap-175

[00279] To a solution of methyl 3-ethylpicolinate in THF/H₂O/MeOH (5 mL/ 0.75 mL/ 3 mL) was added LiOH (35 mg, 1.47 mmol). The mixture was stirred at room temperature for 2 d before additional LiOH (80 mg) was added. After an additional 24 h at room temperature, the mixture was filtered and the solvent was removed *in vacuo*. The residue was then treated with 4*N* HCl in dioxanes (5 mL) and the resulting suspension was concentrated down to dryness to yield Cap-175 as a yellow solid which was used without further purification. ¹H NMR (300 MHz, DMSO-d₆) δ ppm 8.47 (1 H, dd, *J* = 4.8, 1.5 Hz), 7.82-7.89 (1 H, m), 7.53 (1 H, dd, *J* = 7.7, 4.8 Hz), 2.82 (2 H, q, *J* = 7.3 Hz), 1.17 (3 H, t, *J* = 7.5 Hz). *R*_t = 0.36 min (Cond.-MD1); LC-MS: Anal. Calcd. for [M+H]⁺ C₈H₁₀NO₂: 152.07; found: 152.10.



(S)-2-(4,4-Difluorocyclohexyl)-2-(methoxycarbonylamino)acetic acid




[00280] A solution of 1,4-dioxaspiro[4.5]decan-8-one (15 g, 96 mmol) in EtOAc (150 mL) was added to a solution of methyl 2-(benzyloxycarbonylamino)-2-(dimethoxyphosphoryl)acetate (21.21 g, 64.0 mmol) in 1,1,3,3-tetramethylguanidine (10.45 mL, 83 mmol) and EtOAc (150 mL). The resulting solution was stirred at ambient temperature for 72 h and then it was diluted with EtOAc (25 mL). The organic layer was washed with 1*N* HCl (75 mL), H₂O (100 mL) and brine (100 mL), dried (MgSO₄), filtered and concentrated. The residue was purified via BIOTAGE® (5% to 25% EtOAc/Hexanes; 300g column). The combined fractions containing the product were then concentrated under vacuum and the residue was re-crystallized from hexanes/EtOAc

to give white crystals that corresponded to methyl 2-(benzyloxycarbonylamino)-2-(1,4-dioxaspiro[4.5]decan-8-ylidene)acetate (6.2 g) ^1H NMR (400 MHz, CDCl_3 -d) δ ppm 7.30 - 7.44 (5 H, m), 6.02 (1 H, br. s.), 5.15 (2 H, s), 3.97 (4 H, s), 3.76 (3 H, br. s.), 2.84 - 2.92 (2 H, m), 2.47 (2 H, t, $J=6.40$ Hz), 1.74 - 1.83 (4 H, m). LC (Cond. OL1): $R_t = 2.89$ min.


5 LC-MS: Anal. Calcd. for $[\text{M}+\text{Na}]^+ \text{C}_{19}\text{H}_{23}\text{NNaO}_6$: 745.21; found: 745.47.

Cap-176, Step b



[00281] Ester Cap-176, Step b was prepared from alkene Cap-176, Step a according to the method of Burk, M.J. et al. (*J. Am. Chem. Soc.*, 117:9375-9376 (1995)) and references therein): A 500 mL high-pressure bottle was charged with alkene Cap-176, Step a (3.5 g, 9.68 mmol) in degassed MeOH (200 mL) under a blanket of N_2 . The solution was then charged with (-)-1,2-Bis((2S,5S)-2,5-dimethylphospholano)ethane(cyclooctadiene) rhodium (I) tetrafluoroborate (0.108 g, 0.194 mmol) and the resulting mixture was flushed with N_2 (3x) and charged with H_2 (3x). The solution was shaken vigorously under 70 psi of H_2 at ambient temperature for 72 h. The solvent was removed under reduced pressure and the remaining residue was taken up in EtOAc. The brownish solution was then filtered through a plug of Silica Gel and eluted with EtOAc. The solvent was concentrated under vacuum to afford a clear oil corresponding to ester Cap-176, Step b (3.4 g). ^1H NMR (500 MHz, CDCl_3 -d) δ ppm 7.28 - 7.43 (5 H, m), 5.32 (1 H, d, $J=9.16$ Hz), 5.06 - 5.16 (2 H, m), 4.37 (1 H, dd, $J=9.00, 5.04$ Hz), 3.92 (4 H, t, $J=3.05$ Hz), 3.75 (3 H, s), 1.64 - 1.92 (4 H, m), 1.37 - 1.60 (5 H, m). LC (Cond. OL1): $R_t = 1.95$ min. LC-MS: Anal. Calcd. for $[\text{M}+\text{H}]^+ \text{C}_{19}\text{H}_{26}\text{NO}_6$: 364.18; found: 364.27.


Cap-176, Step c



[00282] Ester Cap-176, Step b (4.78 g, 13.15 mmol) was dissolved in THF (15 mL) followed by sequential addition of water (10 mL), glacial acetic acid (26.4 mL, 460 mmol) and dichloroacetic acid (5.44 mL, 65.8 mmol). The resulting mixture was stirred for 72 h at ambient temperature, and the reaction was quenched by slow addition of solid Na₂CO₃ with vigorous stirring until the release of gas was no longer visible. Crude product was extracted into 10% ethyl acetate-dichloromethane and the organic layers were combined, dried (MgSO₄) filtered and concentrated. The resulting residue was purified via BIOTAGE® (0 to 30% EtOAc/Hex; 25 g column) to afford ketone Cap-176, Step c (3.86g) as a clear oil. ¹H NMR (400 MHz, CDCl₃-d) δ ppm 7.28 - 7.41 (5 H, m), 5.55 (1 H, d, *J*=8.28 Hz), 5.09 (2 H, s), 4.46 (1 H, dd, *J*=8.16, 5.14 Hz), 3.74 (3 H, s), 2.18 - 2.46 (5 H, m), 1.96 - 2.06 (1 H, m), 1.90 (1 H, ddd, *J*=12.99, 5.96, 2.89 Hz), 1.44 - 1.68 (2 H, m, *J*=12.36, 12.36, 12.36, 12.36, 4.77 Hz). LC (Cond. OL1): *R*_t = 1.66 min. LC-MS: Anal. Calcd. for [M+Na]⁺ C₁₇H₂₁NNaO₅: 342.13; found: 342.10.

15


Cap-176, Step d



[00283] DEOXO-FLUOR® (3.13 mL, 16.97 mmol) was added to a solution of ketone Cap-176, Step c (2.71 g, 8.49 mmol) in CH₂Cl₂ (50 mL) followed by addition of a catalytic amount of EtOH (0.149 mL, 2.55 mmol). The resulting yellowish solution was stirred at rt overnight. The reaction was quenched by addition of sat. aq. NaHCO₃ (25 mL) and the mixture was extracted with EtOAc (3X75 mL)). The combined organic layers were dried (MgSO₄), filtered and dried to give a yellowish oil. The residue was purified via BIOTAGE® chromatography (2% to 15% EtOAc/Hex; 90g column) and a white solid corresponding to the difluoro amino acid difluoride Cap-176, Step d (1.5 g) was recovered. ¹H NMR (400 MHz, CDCl₃-d) δ ppm 7.29 - 7.46 (5 H, m), 5.34 (1 H, d, *J*=8.28 Hz), 5.12 (2 H, s), 4.41 (1 H, dd, *J*=8.66, 4.89 Hz), 3.77 (3 H, s), 2.06 - 2.20 (2 H, m), 1.83 - 1.98 (1 H, m), 1.60 - 1.81 (4 H, m), 1.38 - 1.55 (2 H, m). ¹⁹F NMR (376 MHz, CDCl₃-d) δ ppm -92.15 (1 F, d, *J*=237.55 Hz), -102.44 (1 F, d, *J*=235.82 Hz). LC (Cond.

OL1): $R_t = 1.66$ min. LC-MS: Anal. Calcd. for $[2M+Na]^+ C_{34}H_{42}F_4N_2NaO_8$: 705.28; found: 705.18.

Cap-176, Step e




5

[00284] Difluoride Cap-176, Step d (4 g, 11.72 mmol) was dissolved in MeOH (120 mL) and charged with Pd/C (1.247 g, 1.172 mmol). The suspension was flushed with N_2 (3x) and the reaction mixture was placed under 1 atm of H_2 (balloon). The mixture was stirred at ambient temperature for 48 h. The suspension was then filtered through a plug of CELITE® and concentrated under vacuum to give an oil that corresponded to amino acid Cap-176, Step e (2.04 g) and that was used without further purification. 1H NMR (400 MHz, DMSO- d_6) δ ppm 3.62 (3 H, s), 3.20 (1 H, d, $J=5.77$ Hz), 1.91 - 2.09 (2 H, m), 1.50 - 1.88 (7 H, m), 1.20 - 1.45 (2 H, m). ^{19}F NMR (376 MHz, DMSO- d_6) δ ppm -89.39 (1 F, d, $J=232.35$ Hz), -100.07 (1 F, d, $J=232.35$ Hz). ^{13}C NMR (101 MHz, DMSO- d_6) δ ppm 175.51 (1 C, s), 124.10 (1 C, t, $J=241.21$, 238.90 Hz), 57.74 (1 C, s), 51.39 (1 C, s), 39.23 (1 C, br. s.), 32.02 - 33.83 (2 C, m), 25.36 (1 C, d, $J=10.02$ Hz), 23.74 (1 C, d, $J=9.25$ Hz). LC (Cond. OL2): $R_t = 0.95$ min. LC-MS: Anal. Calcd. for $[2M+H]^+ C_{18}H_{31}F_4N_2O_2$: 415.22; found: 415.40.

20


Cap-176, Step f



[00285] Methyl chloroformate (1.495 mL, 19.30 mmol) was added to a solution of amino acid Cap-176, Step e (2 g, 9.65 mmol) and DIEA (6.74 mL, 38.6 mmol) in CH_2Cl_2 (100 mL). The resulting solution was stirred at rt for 3 h and volatiles were removed under reduced pressure. The residue was purified via BIOTAGE® (0% to 20% EtOAc/Hex; 90g column). A clear oil that solidified upon standing under vacuum and corresponding to carbamate Cap-176, Step f (2.22 g) was recovered. 1H NMR (500 MHz,

CDCl₃-d) δ ppm 5.27 (1 H, d, $J=8.55$ Hz), 4.39 (1 H, dd, $J=8.85$, 4.88 Hz), 3.77 (3 H, s), 3.70 (3 H, s), 2.07 - 2.20 (2 H, m), 1.84 - 1.96 (1 H, m), 1.64 - 1.82 (4 H, m), 1.39 - 1.51 (2 H, m). ¹⁹F NMR (471 MHz, CDCl₃-d) δ ppm -92.55 (1 F, d, $J=237.13$ Hz), -102.93 (1 F, d, $J=237.12$ Hz). ¹³C NMR (126 MHz, CDCl₃-d) δ ppm 171.97 (1 C, s), 156.69 (1 C, s), 119.77 - 125.59 (1 C, m), 57.24 (1 C, br. s.), 52.48 (1 C, br. s.), 52.43 (1 C, s), 39.15 (1 C, s), 32.50 - 33.48 (2 C, m), 25.30 (1 C, d, $J=9.60$ Hz), 24.03 (1 C, d, $J=9.60$ Hz). LC (Cond. OL1): R_t = 1.49 min. LC-MS: Anal. Calcd. for [M+Na]⁺ C₁₁H₁₇F₂NNaO₄: 288.10; found: 288.03.

10




(S)-2-(4,4-Difluorocyclohexyl)-2-(methoxycarbonylamino)acetic acid



[00286] A solution of LiOH (0.379 g, 15.83 mmol) in water (25 mL) was added to a solution of carbamate Cap-176, Step f (2.1 g, 7.92 mmol) in THF (75 mL) and the resulting mixture was stirred at ambient temperature for 4 h. THF was removed under vacuum and the remaining aqueous phase was acidified with 1N HCl solution (2 mL) and then extracted with EtOAc (2 X 50 mL). The combined organic layers were dried (MgSO₄), filtered and concentrated to give a white foam corresponding to Cap-176 (1.92 g). ¹H NMR (400 MHz, DMSO-d₆) δ ppm 12.73 (1 H, s), 7.50 (1 H, d, $J=8.78$ Hz), 3.97 (1 H, dd, $J=8.53$, 6.02 Hz), 3.54 (3 H, s), 1.92 - 2.08 (2 H, m), 1.57 - 1.90 (5 H, m), 1.34 - 1.48 (1 H, m), 1.27 (1 H, qd, $J=12.72$, 3.26 Hz). ¹⁹F NMR (376 MHz, DMSO-d₆) δ ppm -89.62 (1 F, d, $J=232.35$ Hz), -99.93 (1 F, d, $J=232.35$ Hz). LC (Cond. OL2): R_t = 0.76 min. LC-MS: Anal. Calcd. for [M-H]⁺ C₁₀H₁₄F₂NO₄: 250.09; found: 250.10.

25

Cap-177a-d




Cap-177a-d, Step a



[00287] 1,1,3,3-Tetramethylguanidine (0.985 mL, 7.85 mmol) was added to a stirred solution of methyl 2-(benzyloxycarbonylamino)-2-(dimethoxyphosphoryl)acetate (2.0 g, 6.0 mmol) in EtOAc (40 mL) and the mixture was stirred at rt under N₂ for 10 min. Then dihydro-2H-pyran-3(4H)-one [23462-75-1] (0.604 g, 6.04 mmol) was added and the mixture was stirred at rt for 16 h. The reaction mixture was then cooled in freezer for 10 min and neutralized with aq. citric acid (1.5 g in 20 mL water). The two phases were partitioned and the organic layer was washed with 0.25 N aq.HCl and brine, and then dried (MgSO₄) and concentrated to a colorless oil. The crude material was purified by flash silica chromatography (loading solvent: DCM, eluted with EtOAc/Hexanes, gradient from 20% to 30% EtOAc) to yield two isomeric products: The first eluted product was (Z)-methyl 2-(benzyloxycarbonylamino)-2-(2H-pyran-3(4H,5H,6H)-ylidene)acetate (490 mg) (white solid), and the second was (E)-methyl 2-(benzyloxycarbonylamino)-2-(2H-pyran-3(4H,5H,6H)-ylidene)acetate (433 mg) (white solid). LC-MS retention time 1.398 min (for Z-isomer) and 1.378min (for E-isomer); m/z 304.08 (for Z-isomer) and 304.16 (for E-isomer) (MH⁺). LC data was recorded on a Shimadzu LC-10AS liquid chromatograph equipped with a PHENOMENEX® Luna 10u C18 3.0x50mm column using a SPD-10AV UV-Vis detector at a detector wave length of 220 nm. The elution conditions employed a flow rate of 4 mL/min, a gradient of 100% Solvent A / 0% Solvent B to 0% Solvent A / 100% Solvent B, a gradient time of 3 min, a hold time of 1 min, and an analysis time of 4 min where Solvent A was 5% MeOH / 95% H₂O / 10 mM ammonium acetate and Solvent B was 5% H₂O / 95% MeOH / 10 mM ammonium acetate. MS data was determined using a MICROMASS® Platform for LC in electrospray mode. ¹H NMR (400 MHz, chloroform-d) (for Z-isomer) δ ppm 7.30 - 7.44 (m, 5 H), 6.18 (br. s., 1 H), 5.10 - 5.17 (m, 2 H), 4.22 (s, 2 H), 3.78 (br. s., 3 H), 2.93 - 3.02 (m, 2 H), 1.80 (dt, J=11.7, 5.8 Hz, 2 H), 1.62 (s, 2 H). ¹H NMR (400 MHz, chloroform-d) (for E-isomer) δ ppm 7.31 - 7.44 (m, 5 H), 6.12 (br. s., 1 H), 5.13 - 5.17 (m, 2 H), 4.64 (br. s., 2 H), 3.70 - 3.82 (m, 5 H), 2.49 (t, J=6.5 Hz, 2 H), 1.80 (br. s., 2 H).

(Note: the absolute regiochemistry was determined by ^1H NMR shifts and coupling constants).

Cap-177a-d, Step b



5

[00288] (-)-1,2-Bis((2S,5S)-2,5-dimethylphospholano)ethane(cyclooctadiene)-rhodium(I)tetrafluoroborate (28.2 mg, 0.051 mmol) was added to a stirred solution of (Z)-methyl 2-(benzyloxycarbonylamino)-2-(2H-pyran-3(4H,5H,6H)-ylidene)acetate (310 mg, 1.015 mmol) in MeOH (10 mL) and the mixture was vacuum flushed with N_2 , followed by H_2 , and then the reaction was stirred under H_2 (60 psi) at rt for 2d. The reaction mixture was concentrated and the residue was purified by flash silica chromatography (loading solvent: DCM, eluted with 20% EtOAc in hexanes) to yield (S)-methyl 2-(benzyloxycarbonylamino)-2-((S)-tetrahydro-2H-pyran-3-yl)acetate (204 mg) as clear colorless oil. LC-MS retention time 1.437 min; m/z 307.89 (MH⁺). LC data was recorded on a Shimadzu LC-10AS liquid chromatograph equipped with a PHENOMENEX® Luna 10u C18 3.0x50mm column using a SPD-10AV UV-Vis detector at a detector wave length of 220 nm. The elution conditions employed a flow rate of 4 mL/min, a gradient of 100% Solvent A / 0% Solvent B to 0% Solvent A / 100% Solvent B, a gradient time of 3 min, a hold time of 1 min, and an analysis time of 4 min where Solvent A was 5% MeOH / 95% H_2O / 10 mM ammonium acetate and Solvent B was 5% H_2O / 95% MeOH / 10 mM ammonium acetate. MS data was determined using a MICROMASS® Platform for LC in electrospray mode. ^1H NMR (400 MHz, chloroform-d) δ ppm 7.30 - 7.46 (m, 5 H), 5.32 (d, $J=8.8$ Hz, 1 H), 5.12 (s, 2 H), 4.36 (dd, $J=8.9$, 5.6 Hz, 1 H), 3.84 - 3.98 (m, 2 H), 3.77 (s, 3 H), 3.28 - 3.37 (m, 1 H), 3.23 (dd, $J=11.3$, 10.5 Hz, 1 H), 2.04 - 2.16 (m, 1 H), 1.61 - 1.75 (m, 3 H), 1.31 - 1.43 (m, 1 H).


25

[00289] The other stereoisomer ((E)-methyl 2-(benzyloxycarbonylamino)-2-(2H-pyran-3(4H,5H,6H)-ylidene)acetate) (360 mg, 1.18 mmol) was reduced in a similar manner to yield (S)-methyl 2-(benzyloxycarbonylamino)-2-((R)-tetrahydro-2H-pyran-3-yl)acetate (214 mg) as clear colorless oil. LC-MS retention time 1.437 min; m/z 308.03

(MH⁺). LC data was recorded on a Shimadzu LC-10AS liquid chromatograph equipped with a PHENOMENEX® Luna 10u C18 3.0x50mm column using a SPD-10AV UV-Vis detector at a detector wave length of 220 nm. The elution conditions employed a flow rate of 4 mL/min, a gradient of 100% Solvent A / 0% Solvent B to 0% Solvent A / 100% Solvent B, a gradient time of 3 min, a hold time of 1 min, and an analysis time of 4 min where Solvent A was 5% MeOH / 95% H₂O / 10 mM ammonium acetate and Solvent B was 5% H₂O / 95% MeOH / 10 mM ammonium acetate. MS data was determined using a MICROMASS® Platform for LC in electrospray mode. ¹H NMR (400 MHz, chloroform-d) δ ppm 7.30 - 7.44 (m, 5 H), 5.31 (d, *J*=9.0 Hz, 1 H), 5.12 (s, 2 H), 4.31 (dd, *J*=8.7, 6.9 Hz, 1 H), 3.80 - 3.90 (m, 2 H), 3.77 (s, 3 H), 3.37 (td, *J*=10.8, 3.5 Hz, 1 H), 3.28 (dd, *J*=11.3, 9.8 Hz, 1 H), 1.97 - 2.10 (m, 1 H), 1.81 (d, *J*=11.5 Hz, 1 H), 1.61 - 1.72 (m, 2 H), 1.33 - 1.46 (m, 1 H).

[00290] The individual enantiomers of Cap-177a, Step b (Cap-177c, Step b) and Cap-177b, Step b (Cap-177d, Step b) were prepared in the same manner and in similar yields utilizing (-)-1,2-Bis((2R,5R)-2,5-dimethylphospholano)ethane (cyclooctadiene)-rhodium(I)tetrafluoroborate as the hydrogenation catalyst for the olefin reductions of the individual stereoisomer starting materials.

Cap-177a and Cap-177b, Step c




[00291] 10% Pd/C (69.3 mg, 0.065 mmol) was added to a solution of (S)-methyl 2-(benzyloxycarbonylamino)-2-((S)-tetrahydro-2H-pyran-3-yl)acetate (200 mg, 0.651 mmol) and dimethyl dicarbonate [4525-33-1] (0.104 mL, 0.976 mmol) in MeOH (10 mL). The reaction mixture was vacuum flushed with N₂, followed by H₂, and then the reaction was stirred under H₂ (55 psi) at rt for 5 h. The reaction mixture was filtered through CELITE®/silica pad and the filtrate was concentrated to a colorless oil. The crude oil was purified by flash silica chromatography (loading solvent: DCM, eluted with 30% EtOAc in hexanes) to yield product (S)-methyl 2-(methoxycarbonylamino)-2-((S)-tetrahydro-2H-pyran-3-yl)acetate (132 mg) as colorless oil. LC-MS retention time 0.92

min; m/z 231.97 (MH⁺). LC data was recorded on a Shimadzu LC-10AS liquid chromatograph equipped with a PHENOMENEX® Luna 10u C18 3.0x50mm column using a SPD-10AV UV-Vis detector at a detector wave length of 220 nM. The elution conditions employed a flow rate of 4 mL/min, a gradient of 100% Solvent A / 0% Solvent B to 0% Solvent A / 100% Solvent B, a gradient time of 3 min, a hold time of 1 min, and an analysis time of 4 min where Solvent A was 5% MeOH / 95% H₂O / 10 mM ammonium acetate and Solvent B was 5% H₂O / 95% MeOH / 10 mM ammonium acetate. MS data was determined using a MICROMASS® Platform for LC in electrospray mode. ¹H NMR (400 MHz, chloroform-d) δ ppm 5.24 (d, J=8.5 Hz, 1 H), 4.34 (dd, J=8.9, 5.6 Hz, 1 H), 3.84 - 3.97 (m, 2 H), 3.77 (s, 3 H), 3.70 (s, 3 H), 3.29 - 3.38 (m, 1 H), 3.23 (dd, J=11.2, 10.4 Hz, 1 H), 2.03 - 2.14 (m, 1 H), 1.56 - 1.75 (m, 3 H), 1.32 - 1.43 (m, 1 H).

[00292] Another diastereomer ((S)-methyl 2-(benzyloxycarbonylamino)-2-((R)-tetrahydro-2H-pyran-3-yl)acetate) was transformed in a similar manner to yield (S)-methyl 2-(methoxycarbonylamino)-2-((R)-tetrahydro-2H-pyran-3-yl)acetate as clear colorless oil. LC-MS retention time 0.99 min; m/z 231.90 (MH⁺). LC data was recorded on a Shimadzu LC-10AS liquid chromatograph equipped with a PHENOMENEX® Luna 10u C18 3.0x50mm column using a SPD-10AV UV-Vis detector at a detector wave length of 220 nM. The elution conditions employed a flow rate of 4 mL/min, a gradient of 100% Solvent A / 0% Solvent B to 0% Solvent A / 100% Solvent B, a gradient time of 3 min, a hold time of 1 min, and an analysis time of 4 min where Solvent A was 5% MeOH / 95% H₂O / 10 mM ammonium acetate and Solvent B was 5% H₂O / 95% MeOH / 10 mM ammonium acetate. MS data was determined using a MICROMASS® Platform for LC in electrospray mode. ¹H NMR (400 MHz, chloroform-d) δ ppm 5.25 (d, J=8.0 Hz, 1 H), 4.29 (dd, J=8.4, 7.2 Hz, 1 H), 3.82 - 3.90 (m, 2 H), 3.77 (s, 3 H), 3.70 (s, 3 H), 3.37 (td, J=10.8, 3.3 Hz, 1 H), 3.28 (t, J=10.5 Hz, 1 H), 1.96 - 2.08 (m, 1 H), 1.81 (dd, J=12.9, 1.6 Hz, 1 H), 1.56 - 1.72 (m, 2 H), 1.33 - 1.46 (m, 1 H).

[00293] The individual enantiomers of Cap-177a, Step c (Cap-177c, Step c) and Cap-177b, Step c (Cap-177d, Step c) were prepared in a similar manner and is similar yields using the appropriate starting materials from Cap-177a-d, Step b.

Cap-177a and Cap-177b, Step d




- [00294]** To a solution of (S)-methyl 2-(methoxycarbonylamino)-2-((S)-tetrahydro-2H-pyran-3-yl)acetate (126 mg, 0.545 mmol) in THF (4 mL) stirring at rt was added a solution of 1M LiOH (1.090 mL, 1.090 mmol) in water. The reaction was stirred at rt for 3h, neutralized with 1M HCl (1.1 mL) and extracted with EtOAc (3x 10 mL). The organics were dried, filtered and concentrated to yield (S)-2-(methoxycarbonylamino)-2-((S)-tetrahydro-2H-pyran-3-yl)acetic acid (Cap-177a) (125 mg) as a clear colorless oil. LC-MS retention time 0.44 min; m/z 218.00 (MH⁺). LC data was recorded on a Shimadzu LC-10AS liquid chromatograph equipped with a PHENOMENEX® Luna 10u C18 3.0x50mm column using a SPD-10AV UV-Vis detector at a detector wave length of 220 nM. The elution conditions employed a flow rate of 4 mL/min, a gradient of 100% Solvent A / 0% Solvent B to 0% Solvent A / 100% Solvent B, a gradient time of 3 min, a hold time of 1 min, and an analysis time of 4 min where Solvent A was 5% MeOH / 95% H₂O / 10 mM ammonium acetate and Solvent B was 5% H₂O / 95% MeOH / 10 mM ammonium acetate. MS data was determined using a MICROMASS® Platform for LC in electrospray mode. ¹H NMR (400 MHz, chloroform-d) δ ppm 5.28 (d, *J*=8.8 Hz, 1 H), 4.38 (dd, *J*=8.7, 5.6 Hz, 1 H), 3.96 - 4.04 (m, 1 H), 3.91 (d, *J*=11.0 Hz, 1 H), 3.71 (s, 3 H), 3.33 - 3.41 (m, 1 H), 3.24 - 3.32 (m, 1 H), 2.10 - 2.24 (m, 1 H), 1.74 - 1.83 (m, 1 H), 1.63 - 1.71 (m, 2 H), 1.35 - 1.49 (m, 1 H).
- [00295]** Another diastereomer ((S)-methyl 2-(methoxycarbonylamino)-2-((R)-tetrahydro-2H-pyran-3-yl)acetate) was transformed in a similar manner to yield (S)-2-(methoxycarbonylamino)-2-((R)-tetrahydro-2H-pyran-3-yl)acetic acid (Cap-177b) as clear colorless oil. LC-MS retention time 0.41 min; m/z 217.93 (MH⁺). LC data was recorded on a Shimadzu LC-10AS liquid chromatograph equipped with a PHENOMENEX® Luna 10u C18 3.0x50mm column using a SPD-10AV UV-Vis detector at a detector wave length of 220 nM. The elution conditions employed a flow rate of 4 mL/min, a gradient of 100% Solvent A / 0% Solvent B to 0% Solvent A / 100% Solvent B, a gradient time of 3 min, a hold time of 1 min, and an analysis time of 4 min where Solvent A was 5% MeOH / 95% H₂O / 10 mM ammonium acetate and Solvent B

was 5% H₂O / 95% MeOH / 10 mM ammonium acetate. MS data was determined using a MICROMASS® Platform for LC in electrospray mode. ¹H NMR (400 MHz, chloroform-d) δ ppm 6.18 (br. s., 1 H), 5.39 (d, *J*=8.5 Hz, 1 H), 4.27 - 4.37 (m, 1 H), 3.82 - 3.96 (m, 2 H), 3.72 (s, 3 H), 3.42 (td, *J*=10.8, 3.3 Hz, 1 H), 3.35 (t, *J*=10.4 Hz, 1 H),
 5 2.01 - 2.18 (m, 1 H), 1.90 (d, *J*=11.8 Hz, 1 H), 1.59 - 1.76 (m, 2 H), 1.40 - 1.54 (m, 1 H).


[00296] The individual enantiomers of Cap-177a (Cap-177c) and Cap-177b (Cap-177d) were prepared in a similar manner and is similar yields using the appropriate starting materials from Cap-177a-d, Step c.

10

Cap-178




Cap-178, Step a




- 15 **[00297]** To a solution of (2S,3S,4S)-2-methyl-3,4-dihydro-2H-pyran-3,4-diyl diacetate (5 g, 23.34 mmol) in 20 mL of MeOH in a hydrogenation tank was added Pd/C (150 mg, 0.141 mmol). The resulting mixture was hydrogenated at 40 psi on Parr Shaker for 1 hour. The mixture was then filtered and the filtrate was concentrated to afford Cap-178, Step a (5.0 g) as a clear oil, which solidified while standing. ¹H NMR (500 MHz, CDCl₃)
 20 δ ppm 4.85 - 4.94 (1 H, m), 4.69 (1 H, t, *J*=9.46 Hz), 3.88 - 3.94 (1 H, m), 3.44 (1 H, td, *J*=12.21, 1.83 Hz), 3.36 (1 H, dq, *J*=9.42, 6.12 Hz), 2.03 - 2.08 (1 H, m), 2.02 (3 H, s), 2.00 (3 H, s), 1.70 - 1.80 (1 H, m), 1.16 (3 H, d, *J*=6.10 Hz).

Cap-178, Step b



[00298] To a solution of Cap-178, Step a (5.0 g, 23 mmol) in 50 mL of MeOH was added several drops of sodium methoxide. After stirring at room temperature for 30 min, sodium methoxide (0.1 mL, 23.12 mmol) was added and the solution was stirred at room temperature overnight. The solvent was then removed under vacuum. The residue was diluted with benzene and concentrated to afford the corresponding diol as a yellow solid. The solid was dissolved in 50 mL of pyridine and to this solution at -35 °C was added benzoyl chloride (2.95 mL, 25.4 mmol) dropwise. The resulting mixture was stirred at -35 °C for 1 hour then at room temperature overnight. The mixture was diluted with Et₂O and washed with water. The aqueous layer was extracted with EtOAc (2X). The combined organic layers were dried with MgSO₄ and concentrated. The crude product was purified by flash chromatography (silica gel, 5%-15% EtOAc/Hex) to afford Cap-178, Step b (4.5 g) as clear oil which slowly crystallized upon prolonged standing. LC-MS: Anal. Calcd. for [M+Na]⁺ C₁₃H₁₆NaO₄ 259.09; found 259.0; ¹H NMR (500 MHz, CDCl₃) δ ppm 8.02 - 8.07 (2 H, m), 7.55 - 7.61 (1 H, m), 7.45 (2 H, t, *J*=7.78 Hz), 5.01 (1 H, ddd, *J*=11.44, 8.70, 5.49 Hz), 3.98 (1 H, ddd, *J*=11.90, 4.88, 1.53 Hz), 3.54 (1 H, td, *J*=12.36, 2.14 Hz), 3.41 (1 H, t, *J*=9.00 Hz), 3.31 - 3.38 (1 H, m), 2.13 - 2.19 (1 H, m), 1.83 - 1.94 (1 H, m), 1.36 (3 H, d, *J*=5.80 Hz).

20 Cap-178, Step c




[00299] To a mixture of NaH (1.143 g, 28.6 mmol) (60% in mineral oil) in 6 mL of CS₂ was added Cap-178, Step b (4.5 g, 19 mmol) in 40 mL of CS₂ dropwise over 15 min. The resulting mixture was stirred at room temperature for 30 min. The mixture turned light orange with some solid. MeI (14.29 mL, 229 mmol) was then added dropwise over 20 min. The mixture was then stirred at room temperature overnight. The reaction was carefully quenched with saturated NH₄Cl solution. The mixture was extracted with

EtOAc (3X). The combined organic layers were dried with MgSO_4 and concentrated. The crude product was purified by flash chromatography (silica gel, 6% EtOAc/Hex) to afford Cap-178, Step c (3.13 g) as clear oil. LC-MS: Anal. Calcd. for $[\text{M}+\text{Na}]^+$

5 $\text{C}_{15}\text{H}_{18}\text{NaO}_4\text{S}_2$ 349.05; found 349.11; ^1H NMR (500 MHz, CDCl_3) δ ppm 7.94 - 8.00 (2 H, m), 7.50 - 7.58 (1 H, m), 7.41 (2 H, t, $J=7.78$ Hz), 5.96 (1 H, t, $J=9.46$ Hz), 5.28 (1 H, ddd, $J=11.37, 9.38, 5.49$ Hz), 4.02 (1 H, ddd, $J=11.98, 4.96, 1.68$ Hz), 3.54 - 3.68 (2 H, m), 2.48 (3 H, s), 2.31 (1 H, dd), 1.88 - 1.99 (1 H, m), 1.28 (3 H, d).

Cap-178, Step d



10


[00300] To a mixture of Cap-178, Step c (3.13 g, 9.59 mmol) and AIBN (120 mg, 0.731 mmol) in 40 mL of benzene at 80 °C was added tri-n-butyltin hydride (10.24 mL, 38.4 mmol). The resulting mixture was stirred at reflux temperature for 20 min then cooled to room temperature. The mixture was diluted with diethyl ether and 100 mL of

15 KF (10 g) aqueous solution was added and the mixture was stirred vigorously for 30 min. The two layers were then separated and the aqueous phase was extracted with EtOAc (2X). The organic layer was dried with MgSO_4 and concentrated. The crude product was purified by flash chromatography (silica gel, deactivated with 3% Et_3N in Hexanes and flushed with 3% Et_3N in Hexanes to remove tributyltin derivative and then eluted with

20 15% EtOAc/Hex) to afford Cap-178, Step d (1.9 g) as clear oil. ^1H NMR (500 MHz, CDCl_3) δ ppm 7.98 - 8.07 (2 H, m), 7.52 - 7.58 (1 H, m), 7.43 (2 H, t, $J=7.63$ Hz), 5.08 - 5.17 (1 H, m), 4.06 (1 H, ddd, $J=11.90, 4.88, 1.53$ Hz), 3.50 - 3.59 (2 H, m), 2.08 - 2.14 (1 H, m), 1.99 - 2.06 (1 H, m), 1.69 - 1.80 (1 H, m), 1.41 - 1.49 (1 H, m), 1.24 (3 H, d, $J=6.10$ Hz).

25


Cap-178, Step e



[00301] To a mixture of Cap-178, Step d (1.9 g, 8.63 mmol) in 10 mL of MeOH was added sodium methoxide (2 mL, 4.00 mmol) (2 M in methanol). The resulting mixture was stirred at room temperature for 5 hours. The solvent was removed under vacuum. The mixture was neutralized with saturated NH_4Cl solution and extracted with EtOAc (3X). The organic layers were dried with MgSO_4 and concentrated to afford Cap-178, Step e (0.8 g) as clear oil. The product was used in the next step without further purification. ^1H NMR (400 MHz, CDCl_3) δ ppm 4.01 (1 H, ddd, $J=11.80, 5.02, 1.76$ Hz), 3.73 - 3.83 (1 H, m), 3.36 - 3.46 (2 H, m), 1.92 - 2.00 (1 H, m), 1.88 (1 H, m), 1.43 - 1.56 (1 H, m), 1.23 (3 H, d), 1.15 - 1.29 (1 H, m).

10


Cap-178, Step f



[00302] Tosyl-Cl (2.63 g, 13.77 mmol) was added to a solution of Cap-178, Step e (0.8 g, 6.89 mmol) and pyridine (2.23 mL, 27.5 mmol) in 100 mL of CH_2Cl_2 . The resulting mixture was stirred at room temperature for 3 days. 10 mL of water was then added into the reaction mixture and the mixture was stirred at room temperature for an hour. The two layers were separated and the organic phase was washed with water and 1 N HCl aq. solution. The organic phase was dried with MgSO_4 and concentrated to afford Cap-178, Step f (1.75 g) as a light yellow solid. The product was used in the next step without further purification. Anal. Calcd. for $[\text{M}+\text{H}]^+$ $\text{C}_{13}\text{H}_{19}\text{O}_4\text{S}$ 271.10; found 270.90; ^1H NMR (500 MHz, CDCl_3) δ ppm 7.79 (2 H, d, $J=8.24$ Hz), 7.34 (2 H, d, $J=7.93$ Hz), 4.53 - 4.62 (1 H, m), 3.94 (1 H, ddd, $J=12.13, 4.96, 1.83$ Hz), 3.29 - 3.41 (2 H, m), 2.45 (3 H, s), 1.90 - 1.97 (1 H, m), 1.79 - 1.85 (1 H, m), 1.64 - 1.75 (1 H, m), 1.38 - 1.48 (1 H, m), 1.17 (3 H, d, $J=6.10$ Hz).


25

Cap-178, Step g



- [00303]** To a microwave tube was placed ethyl 2-(diphenylmethyleneamino)acetate (1.6 g, 5.92 mmol) and Cap-178, Step f (1.6 g, 5.92 mmol). 10 mL of toluene was added. The tube was sealed and LiHMDS (7.1 mL, 7.10 mmol) (1 N in toluene) was added
- 5 dropwise under N₂. The resulting dark brown solution was heated at 100 °C under microwave radiation for 6 hours. To the mixture was then added water and the mixture was extracted with EtOAc (3X). The combined organic layers were washed with brine, dried with MgSO₄ and concentrated to afford a diastereomeric mixture of Cap-3, Step g (3.1 g) as an orange oil. The crude mixture was submitted to the next step without
- 10 separation. LC-MS: Anal. Calcd. for [M+H]⁺ C₂₃H₂₈NO₃ 366.21; found 366.3.


Cap-178, Step h



- [00304]** To a solution of the diastereomeric mixture of ethyl Cap-178, Step g in 20 mL
- 15 of THF was added HCl (30 ml, 60.0 mmol) (2 N aqueous). The resulting mixture was stirred at room temperature for 1 hour. The mixture was extracted with EtOAc and the aqueous layer was concentrated to afford an HCl salt of Cap-178, Step h (1.9 g) as an orange oil. The salt was used in the next step without further purification. LC-MS: Anal. Calcd. for [M+H]⁺ C₁₀H₂₀NO₃ 202.14; found 202.1.

20


Cap-178, Step i



[00305] A solution of 1.9 g Cap-178, Step h (HCl salt), DiPEA (4.19 mL, 24.0 mmol) and methyl chloroformate (1.24 mL, 16.0 mmol) in 20 mL of CH₂Cl₂ was stirred at room temperature for 1 hour. The mixture was diluted with CH₂Cl₂ and washed with water.


- 5 The organic layer was dried with Na₂SO₄ and concentrated. The crude product was purified by flash chromatography (silica gel, 0-20% EtOAc/Hex) to afford Cap-178, Step i (1.1 g) as a yellow oil. Anal. Calcd. for [M+Na]⁺ C₁₂H₂₁NNaO₅ 282.13; found 282.14; ¹H NMR (400 MHz, CDCl₃) δ ppm 5.16 (1 H, br. s.), 4.43 - 4.58 (1 H, m), 4.17 - 4.28 (2 H, m), 3.89 - 4.03 (1 H, m), 3.72 - 3.78 (2 H, m), 3.67 - 3.72 (3 H, m), 2.07 - 2.19 (1 H, m), 1.35 - 1.77 (4 H, m), 1.30 (3 H, td, *J*=7.09, 2.89 Hz), 1.19 (3 H, d, *J*=6.53 Hz).
- 10

Cap-178, Step j



- [00306]** To a mixture of Cap-178, Step i (1.1 g, 4.2 mmol) in 5 mL of THF and 2 mL of water was added LiOH (6.36 mL, 12.7 mmol) (2 N aq.). The resulting mixture was stirred at room temperature overnight. The mixture was then neutralized with 1 N HCl aq. and extracted with EtOAc (3X). The combined organic layers were dried with MgSO₄ and concentrated to afford Cap-178, Step j (0.8 g) as a clear oil. LC-MS: Anal. Calcd. for [M+H]⁺ C₁₀H₁₈NO₅ 232.12; found 232.1; ¹H NMR (400 MHz, CDCl₃) δ ppm 5.20 (1 H, d, *J*=8.28 Hz), 4.54 (1 H, t, *J*=8.16 Hz), 3.95 - 4.10 (1 H, m), 3.66 - 3.85 (5 H, m), 2.15 - 2.29 (1 H, m), 1.41 - 1.85 (4 H, m), 1.23 (3 H, dd, *J*=6.53, 1.76 Hz).
- 15
- 20

Cap-178, Step k




- [00307] To a solution of Cap-178, Step j (240 mg, 1.04 mmol), (S)-1-phenylethanol (0.141 mL, 1.142 mmol) and EDC (219 mg, 1.14 mmol) in 10 mL of CH₂Cl₂ was added DMAP (13.95 mg, 0.114 mmol). The resulting solution was stirred at room temperature overnight and the solvent was removed under vacuum. The residue was taken up into EtOAc, washed with water, dried with MgSO₄ and concentrated. The crude product was purified by chromatography (silica gel, 0-15% EtOAc/Hexanes) to afford Cap-178, Step k as a mixture of two diastereomers. The mixture was separated by chiral HPLC (CHIRALPAK® AS column, 21 x 250 mm, 10 μm) eluting with 90% 0.1% diethylamine/Heptane-10% EtOH at 15 mL/min to afford Cap-178, Step k stereoisomer 1 (eluted first) and Cap-178, Step k stereoisomer 2 (eluted second) as white solids. The stereochemistry of the isomers was not assigned.

- [00308] Cap-178, Step k stereoisomer 1 (130 mg): LC-MS: Anal. Calcd. for [M+Na]⁺ C₁₈H₂₅NNaO₅ 358.16; found 358.16; ¹H NMR (500 MHz, CDCl₃) δ ppm 7.28 - 7.38 (5 H, m), 5.94 (1 H, q, *J*=6.71 Hz), 5.12 (1 H, d, *J*=9.16 Hz), 4.55 (1 H, t, *J*=9.00 Hz), 3.72 - 3.81 (1 H, m), 3.67 (3 H, s), 3.60 - 3.70 (2 H, m), 1.98 - 2.08 (1 H, m), 1.59 (3 H, d, *J*=6.71 Hz), 1.38 - 1.47 (2 H, m), 1.30 (2 H, t, *J*=5.34 Hz), 0.93 (3 H, d, *J*=6.41 Hz).

Cap-178, Stereoisomer 1


- [00309] To a solution of Cap-178, Step k stereoisomer 1 ((S)-2-(methoxycarbonylamino)-2-((2S,4R)-2-methyltetrahydro-2H-pyran-4-yl)acetic acid) (150 mg, 0.447 mmol) in 10 mL of EtOH was added Pd/C (20 mg, 0.188 mmol) and the mixture was hydrogenated on Parr shaker at 40 psi overnight. The mixture was then filtered and the filtrate was concentrated to afford Cap-178, stereoisomer 1 (100 mg) as a sticky white solid. LC-MS: Anal. Calcd. for [M+H]⁺ C₁₀H₁₈NO₅ 232.12; found 232.1; ¹H NMR (500 MHz, CDCl₃) δ ppm 5.14 - 5.27 (1 H, m), 4.51 (1 H, t, *J*=8.39 Hz), 3.90 - 4.07 (1 H, m), 3.60 - 3.83 (5 H, m), 2.06 - 2.27 (1 H, m), 1.45 - 1.77 (4 H, m), 1.21 (3 H, d, *J*=6.41 Hz).

Cap-179 (Enantiomer-1 and Enantiomer-2)




5

Cap-179, Step a




- [00310] 2,6-Dimethyl-4H-pyran-4-one (10 g, 81 mmol) was dissolved in ethanol (125 mL) and Pd/C (1 g, 0.94 mmol) was added. The mixture was hydrogenated in a Parr shaker under H₂ (0.325 g, 161 mmol) (70 psi) at room temperature for 12 hrs. The catalyst was filtered through a pad of CELITE® and washed with ethanol. The filtrate was concentrated in vacuum and the residue was purified via BIOTAGE® (2% to 25 % EtOAc/Hex; 160g column). Two fractions of clear oils were isolated. The first eluting one corresponded to (2R,6S)-2,6-dimethyldihydro-2H-pyran-4(3H)-one (1.8 g) while the second one corresponded to Cap-179, Step a (1.8 g).
- 15 [00311] (2R,6S)-2,6-Dimethyldihydro-2H-pyran-4(3H)-one data: ¹H NMR (500 MHz, CDCl₃) δ ppm 3.69 (2 H, ddd, *J*=11.29, 5.95, 2.29 Hz), 2.24 - 2.36 (2 H, m), 2.08 - 2.23 (2 H, m), 1.18 - 1.34 (6 H, m); ¹³C NMR (126 MHz, CDCl₃) δ ppm 206.96 (1 C, br. s.), 72.69 (2 C, s), 48.70 (2 C, s), 21.72 (2 C, s).
- [00312] Cap-179, Step a data: ¹H NMR (500 MHz, CDCl₃) δ ppm 3.69 - 3.78 (1 H, m), 3.36 - 3.47 (2 H, m), 2.10 (1 H, br. s.), 1.88 (2 H, dd, *J*=12.05, 4.73 Hz), 1.19 (6 H, d, *J*=6.10 Hz), 1.10 (2 H, q, *J*=10.70 Hz); ¹³C NMR (126 MHz, CDCl₃) δ ppm 71.44 (2 C, s), 67.92 (1 C, s), 42.59 (2 C, s), 21.71 (2 C, s).
- 20

Cap-179, Step b




- [00313] DEAD (2.311 mL, 14.59 mmol) was added drop wise to a solution of Cap-179, Step a (0.38 g, 2.92 mmol), 4-nitrobenzoic acid (2.195 g, 13.14 mmol) and Ph₃P (3.83 g, 14.59 mmol) in benzene (25 mL). Heat evolution was detected and the resulting
- 5 amber solution was stirred at ambient temperature for 6 h. Solvent was removed under reduced pressure and the residue was purified via BIOTAGE® (0 to 15% EtOAc/Hex; 80g column). A white solid corresponding to Cap-179, Step b (0.77 g) was isolated. LC-MS: Anal. Calcd. for [M]⁺ C₁₄H₁₇NO₅: 279.11; found 279.12. ¹H NMR (500 MHz, CDCl₃) δ ppm 8.27 - 8.32 (2 H, m), 8.20 - 8.24 (2 H, m), 5.45 (1 H, quin, *J*=2.82 Hz),
- 10 3.92 (2 H, dqd, *J*=11.90, 6.10, 6.10, 6.10, 1.53 Hz), 1.91 (2 H, dd, *J*=14.80, 2.29 Hz), 1.57 (3 H, dt, *J*=14.65, 3.05 Hz), 1.22 (6 H, d, *J*=6.10 Hz).

Cap-179, Step c



- 15 [00314] A solution LiOH (0.330 g, 13.8 mmol) in water (8 mL) was added to a solution of Cap-179, Step b (0.77 g, 2.76 mmol) in THF (30 mL) and the resulting mixture was stirred at ambient temperature for 16 h. THF was removed under reduced pressure and the aqueous layer was diluted with more water (20 mL) and extracted with EtOAc (3 X 15 mL). The combined organic layers were dried (MgSO₄), filtered and
- 20 concentrated under vacuum. An oily residue with a white solid was recovered. The mixture was triturated with hexanes and the solid was filtered off to yield a clear oil corresponding to Cap-179, Step c (0.34 g). ¹H NMR (500 MHz, CDCl₃) δ ppm 4.21 (1 H, quin, *J*=2.82 Hz), 3.87 - 3.95 (2 H, m), 1.72 (1 H, br. s.), 1.63 (2 H, dd, *J*=14.34, 2.14 Hz), 1.39 - 1.47 (2 H, m), 1.17 (6 H, d, *J*=6.41 Hz).


Cap-179, Step d



- [00315] p-Tosyl chloride (3.98 g, 20.89 mmol) was added to a solution of Cap-179, Step c (1.36 g, 10.5 mmol) and Pyridine (3.38 mL, 41.8 mmol) in CH₂Cl₂ (150 mL) at room temperature and stirred for 24h and then concentrated to a yellow oil. The remaining residue was added to pyridine (20 mL) and water (30 mL) and the resulting mixture was stirred at ambient temperature for 1 1/2 h. The mixture was extracted with Et₂O (75 mL) and the separated organic layer was the washed thoroughly with 1 N aq. HCl (4 X 50 mL). The organic layer was then dried (MgSO₄), filtered and concentrated.
- 10 A white solid corresponding to Cap-179, Step d (2.2 g) was isolated. LC-MS: Anal. Calcd. for [2M+H]⁺ C₂₈H₄₁O₈S₂ : 569.22; found 569.3. ¹H NMR (400 MHz, CDCl₃) δ ppm 7.80 (2 H, d, *J*=8.28 Hz), 7.35 (2 H, d, *J*=8.03 Hz), 4.89 (1 H, quin, *J*=2.82 Hz), 3.77 - 3.88 (2 H, m), 2.46 (3 H, s), 1.77 (2 H, dd, *J*=14.93, 2.89 Hz), 1.36 (2 H, ddd, *J*=14.31, 11.54, 2.76 Hz), 1.13 (6 H, d, *J*=6.27 Hz).

15

Cap-179, Step e




- [00316] LiHMDS (4.30 mL, 4.30 mmol) was added to a solution of Cap-179, Step d (1.02 g, 3.59 mmol) and benzyl 2-(diphenylmethyleneamino)acetate (1.181 g, 3.59 mmol) in toluene (25 mL) at room temperature in a sealed microwave vial and the resulting mixture was then stirred for 5 h at 100 °C under microwave radiation. The reaction was quenched with water (10 mL), extracted with EtOAc, washed with water, dried over MgSO₄, filtrated, and concentrated in vacuum. The residue was purified via BIOTAGE® (0% to 6% EtOAc/Hex; 80g column) and a yellow oil corresponding to Cap-179, Step e
- 20 (1.2 g) was isolated. Anal. Calcd. for [2M+Na]⁺ C₅₈H₆₂N₂NaO₆ : 905.45; found 905.42. ¹H NMR (400 MHz, CDCl₃) δ ppm 7.64 - 7.70 (4 H, m), 7.29 - 7.44 (29 H, m), 7.06 (4 H, dd, *J*=7.65, 1.63 Hz), 5.18 (2 H, d, *J*=2.01 Hz), 3.89 (2 H, d, *J*=6.53 Hz), 3.79 - 3.87 (1 H,

25

m), 3.46 (5 H, dq, $J=11.26, 5.87, 5.87, 5.87, 5.87, 1.88$ Hz), 2.47 (2 H, s), 2.35 - 2.46 (2 H, m), 1.78 (1 H, dd, $J=14.81, 3.01$ Hz), 1.62 - 1.65 (1 H, m), 1.61 (2 H, s), 1.36 - 1.43 (3 H, m), 1.19 (7 H, d, $J=6.27$ Hz), 1.14 (11 H, dd, $J=6.15, 2.89$ Hz), 0.86 - 0.96 (3 H, m).

5

Cap-179, Step f (Enantiomer-1 and Enantiomer-2)



- [00317]** Cap-179, Step e (2.08 g, 4.71 mmol) was dissolved in THF (100 mL) and treated with 2 N HCl (9.42 mL, 18.84 mmol). The resulting clear solution was stirred at ambient temperature for 4 h and then THF was removed under reduced pressure. The remaining aqueous layer was extracted with hexanes (3 X 20 mL) and after diluting with H₂O (20 mL), the aqueous phase was basified with 1 N NaOH to pH = 10 and extracted with EtOAc (3 x 10 mL). The combined organic layers were dried (MgSO₄), filtered and concentrated under vacuum. The resulting residue was taken up in CH₂Cl₂ (100 mL) and charged with DIEA (2.468 mL, 14.13 mmol) and methyl chloroformate (0.401 mL, 5.18 mmol). The resulting solution was stirred at ambient temperature for 2 h. The reaction mixture was quenched with water (10 mL) and the organic layer was removed under reduced pressure. The aqueous layer was then extracted with EtOAc (3 x 10 mL) and the combined organic layers were dried (MgSO₄), filtered and concentrated. The residue was purified via BIOTAGE® (10% EtOAc/Hex; 25 g column). A clear colorless oil corresponding to Cap-179, Step f (1.05 g) was recovered. LC-MS: Anal. Calcd. for [M+H]⁺ C₁₈H₂₆NO₅ : 336.18; found 336.3. ¹H NMR (500 MHz, CDCl₃) δ ppm 7.32 - 7.40 (5 H, m), 5.26 (1 H, d, $J=8.24$ Hz), 5.13 - 5.24 (2 H, m), 4.36 (1 H, dd, $J=8.85, 4.88$ Hz), 3.68 (3 H, s), 3.32 - 3.46 (2 H, m), 2.02 - 2.14 (1 H, m), 1.52 (1 H, d, $J=12.82$ Hz), 1.32 (1 H, d, $J=12.51$ Hz), 1.11 - 1.18 (6 H, m), 0.89 - 1.07 (2 H, m).
- [00318]** A chiral SFC method was developed to separate the racemic mixture by using 12% methanol as the modifier on a CHIRALPAK® AD-H column (30 x 250mm, 5μm) (Temp = 35 °C, Pressure = 150 bar, Wavelength = 210 nm, Flow rate = 70 mL/min for 8 min, Solvent A = CO₂, Solvent B = MeOH). The two separated isomers, Cap-179 Step f


(Enantiomer-1) (first eluting) and Cap-179 Step f (Enantiomer-2) (second eluting) exhibited the same analytical data as the corresponding mixture (see above).

Cap-179 (Enantiomer-1 and Enantiomer-2)


- 5 **[00319]** Cap-179 Step f (Enantiomer-1) (0.35 g, 1.044 mmol) was dissolved in MeOH (50 mL) in a Parr bottle and charged with Pd/C (0.111 g, 1.044 mmol). The suspension was then placed in a Parr shaker and the mixture was flushed with N₂ (3X), placed under 40 psi of H₂ (2.104 mg, 1.044 mmol) and shaken at room temperature for 2 h. The catalyst was filtered off through a pad of CELITE® and the solvent was removed under
- 10 reduced pressure, to yield an amber solid corresponding to Cap-179 Enantiomer-1 (0.25 g). ¹H NMR (500 MHz, DMSO-d₆) δ ppm 12.74 (4 H, br. s.), 7.35 (4 H, d, *J*=6.10 Hz), 3.85 (4 H, br. s.), 3.53 (3 H, s), 3.35 (2 H, ddd, *J*=15.95, 9.99, 6.10 Hz), 1.97 (1 H, br. s.), 1.48 (2 H, t, *J*=13.28 Hz), 1.06 (6 H, d, *J*=6.10 Hz), 0.82 - 1.00 (2 H, m).
- [00320]** Cap-179 Enantiomer-2 was prepared similarly: ¹H NMR (500 MHz, DMSO-
- 15 d₆) δ ppm 12.50 (1 H, br. s.), 7.31 (1 H, br. s.), 3.84 (1 H, t, *J*=7.32 Hz), 3.53 (3 H, s), 3.29 - 3.41 (2 H, m), 1.99 (1 H, s), 1.48 (2 H, t, *J*=14.34 Hz), 1.06 (6 H, d, *J*=6.10 Hz), 0.95 (1 H, q, *J*=12.21 Hz), 0.87 (1 H, q, *J*=11.80 Hz). [Note: the minor variation in the ¹H NMR profile of the enantiomers is likely a result of a difference in sample concentration.]

20

Cap-180 (racemic mixture)



Cap-180, Step a




25

- [00321]** p-Tosyl-Cl (4.39 g, 23.0 mmol) was added to a solution of Cap-179, Step a (1.50 g, 11.5 mmol) and pyridine (3.73 mL, 46.1 mmol) in CH₂Cl₂ (50 mL) at room

temperature and stirred for 2 days. The reaction was diluted with CH_2Cl_2 , washed with water, then 1 N HCl. The organic layer was dried (MgSO_4) and concentrated to a yellow oil which was purified via BIOTAGE® (5% to 20% EtOAc/Hex; 40g column). A clear oil that solidified under vacuum and corresponding to Cap-180, Step a (2.89 g) was isolated. LC-MS: Anal. Calcd. for $[2\text{M}+\text{Na}]^+ \text{C}_{28}\text{H}_{40}\text{NaO}_8\text{S}_2$: 591.21; found 591.3. ^1H NMR (500 MHz, CDCl_3) δ ppm 7.80 (2 H, d, $J=8.24$ Hz), 7.35 (2 H, d, $J=7.93$ Hz), 4.59 (1 H, tt, $J=11.37, 4.96$ Hz), 3.36 - 3.46 (2 H, m), 2.46 (3 H, s), 1.91 (2 H, dd, $J=12.05, 5.04$ Hz), 1.37 (2 H, dt, $J=12.67, 11.52$ Hz), 1.19 (6 H, d, $J=6.10$ Hz).

10


Cap-180, Step b



[00322] LiHMDS 1 N (7.09 mL, 7.09 mmol) was added to a solution of Cap-180, Step a (1.68 g, 5.91 mmol) and ethyl 2-(diphenylmethyleneamino)acetate (1.579 g, 5.91 mmol) in toluene (30 mL) at room temperature and the resulting mixture was then stirred for 16 h at 85 °C. The reaction was quenched with water (50 mL), extracted with EtOAc, washed with water, dried over MgSO_4 , filtrated, and concentrated *in vacuo*. The residue was purified via BIOTAGE® (0% to 15% EtOAc/Hex; 40g column). A clear yellowish oil corresponding to Cap-180, Step b (racemic mixture; 0.64 g) was isolated. LC-MS: Anal. Calcd. for $[\text{M}+\text{H}]^+ \text{C}_{24}\text{H}_{30}\text{NO}_3$: 380.22; found 380.03. ^1H NMR (400 MHz, CDCl_3) δ ppm 7.64 - 7.70 (2 H, m), 7.45 - 7.51 (3 H, m), 7.38 - 7.44 (1 H, m), 7.31 - 7.37 (2 H, m), 7.13 - 7.19 (2 H, m), 4.39 (1 H, d, $J=10.54$ Hz), 4.16 - 4.26 (2 H, m), 3.29 - 3.39 (1 H, m), 2.93 - 3.03 (1 H, m), 2.70 (1 H, m, $J=9.41, 4.14$ Hz), 1.42 - 1.49 (2 H, m), 1.31 - 1.37 (1 H, m), 1.29 (4 H, t, $J=7.15$ Hz), 1.04 (6 H, dd, $J=7.78, 6.27$ Hz).

25

Cap-180, Step c




[00323] Cap-180, Step b (0.36 g, 0.949 mmol) was dissolved in THF (10 mL) and treated with 2 N HCl (1.897 mL, 3.79 mmol). The resulting clear solution was stirred at ambient temperature for 20 h and THF was removed under reduced pressure. The remaining aqueous layer was extracted with hexanes (3 X 20 mL) and after diluting with H₂O (20 mL), the aqueous phase was basified with 1 N NaOH to pH = 10 and extracted with EtOAc (3 x 10 mL). The combined organic layers were dried (MgSO₄), filtered and concentrated under vacuum. The resulting residue was taken up in CH₂Cl₂ (10.00 mL) and charged with DIEA (0.497 mL, 2.85 mmol) and methyl chloroformate (0.081 mL, 1.044 mmol). The resulting solution was stirred at ambient temperature for 2 h and the reaction mixture was quenched with water (10 mL) and the organic layer was removed under reduced pressure. Aqueous layer was extracted with EtOAc (3 x 10 mL) and the combined organic layers were dried (MgSO₄), filtered and concentrated. An amber oil corresponding to Cap-180, Step c (0.21 g) was recovered and it was used without further purification. LC-MS: Anal. Calcd. for [M+H]⁺ C₁₃H₂₄NO₅ : 273.17; found 274.06. ¹H NMR (400 MHz, CDCl₃) δ ppm 5.20 (1 H, d, *J*=8.03 Hz), 4.59 (1 H, t, *J*=10.16 Hz), 4.11 - 4.27 (3 H, m), 3.69 - 3.82 (2 H, m), 3.64 (3 H, s), 1.95 - 2.07 (1 H, m), 1.63 (1 H, d, *J*=13.80 Hz), 1.41 (2 H, dd, *J*=8.03, 4.02 Hz), 1.31 - 1.37 (1 H, m), 1.26 (3 H, t, *J*=7.15 Hz), 1.16 (1 H, d, *J*=6.27 Hz), 1.12 (6 H, dd, *J*=6.15, 3.89 Hz).


20 Cap-180 (racemic mixture)

[00324] Cap-180, Step c (0.32 g, 1.2 mmol) was dissolved in THF (10 mL) and charged with LiOH (0.056 g, 2.342 mmol) in water (3.33 mL) at 0 °C. The resulting solution was stirred at rt for 2 h. THF was removed under reduced pressure and the remaining residue was diluted with water (15 mL) and washed with Et₂O (2 x 10 mL). The aqueous layer was then acidified with 1N HCl to pH ~ 2 and extracted with EtOAc (3 x 15 mL). The combined organic layers were dried (MgSO₄), filtered and concentrated under vacuum to yield Cap-180 (racemic mixture) (0.2 g) as a white foam. LC-MS: Anal. Calcd. for [M+H]⁺ C₁₁H₂₀NO₅ : 246.13; found 246.00. ¹H NMR (400 MHz, CDCl₃) δ ppm 5.14 (1 H, d, *J*=9.03 Hz), 4.65 (1 H, t, *J*=9.91 Hz), 3.63 - 3.89 (5 H, m), 1.99 - 2.13 (1 H, m), 1.56 - 1.73 (2 H, m), 1.48 - 1.55 (1 H, m), 1.35 - 1.48 (1 H, m), 1.27 (1 H, br. s.), 1.17 (6 H, d, *J*=6.02 Hz).

Cap-185 (Enantiomer-1 and Enantiomer-2)



Cap-185, Step a



5

[00325] To a mixture of furan (1.075 mL, 14.69 mmol) and zinc (1.585 g, 24.24 mmol) in 1 mL of THF was added 1,1,3,3-tetrabromopropan-2-one (8.23 g, 22.03 mmol) and triethyl borate (5.25 mL, 30.8 mmol) in 4 mL of THF dropwise during 1 hour in dark. The resulting mixture was stirred at room temperature in dark for 17 hours. The resulting dark


10 brown mixture was cooled to -15 °C, and 6 mL of water was added. The mixture was warmed to 0 °C and stirred at this temperature for 30 min. The mixture was then filtered and washed with ether. The filtrate was diluted with water and extracted with ether (3x). The combined organic layers were dried with MgSO₄ and concentrated to afford dark brown oil. The dark brown oil was dissolved in 6 mL of MeOH and the solution was

15 added dropwise to a mixture of zinc (4.99 g, 76 mmol), copper (I) chloride (0.756 g, 7.64 mmol) and ammonium chloride (5.4 g, 101 mmol) in 20 mL of MeOH. The reaction temperature was maintained below 15 °C during addition. The mixture was then stirred at room temperature for 20 hours, filtered, and the filtrate was diluted with water and

20 extracted with CH₂Cl₂ (3X). The combined organic layers were dried with MgSO₄ and concentrated. The crude product was purified by flash chromatography (silica gel, 0-14% EtOAc/Hex) to afford Cap-185, Step a as a white solid (1.0 g) as a white solid, which turned yellow soon. ¹H NMR (500 MHz, CDCl₃) δ ppm 6.24 (2 H, s), 5.01 (2 H, d, *J*=4.88 Hz), 2.73 (2 H, dd, *J*=16.94, 5.04 Hz), 2.31 (2 H, d, *J*=16.79 Hz).


25

Cap-185, Step b




- [00326] To a solution of Cap-185, Step a (240 mg, 1.933 mmol) in 2 mL of THF at -78 °C was added L-selectride (3.87 mL, 3.87 mmol) (1 M in THF) dropwise over 100 min. The resulting mixture was stirred at -78 °C for 1 hour and then at room temperature overnight. The mixture was then cooled to 0 °C, 4 mL of 20% NaOH aqueous solution was added, followed by 2 mL of H₂O₂ (30% water solution) dropwise. The resulting mixture was stirred for 1 hour and then neutralized with 6N HCl (~5 mL). The aqueous layer was saturated with NaCl and extracted with CH₂Cl₂ (3X). The combined organic layers were dried with MgSO₄ and concentrated. The crude product was purified by flash chromatography (silica gel, 0-40% EtOAc/Hex) to afford Cap-185, Step b (180 mg) as clear oil. ¹H NMR (400 MHz, CDCl₃) δ ppm 6.49 (2 H, s), 4.76 (2 H, d, *J*=4.27 Hz), 3.99 (1 H, t, *J*=5.77 Hz), 2.29 (2 H, ddd, *J*=15.18, 5.65, 4.02 Hz), 1.70 - 1.78 (2 H, m).

Cap-185, Step c



- [00327] p-Tosyl-Cl (544 mg, 2.85 mmol) was added to a solution of Cap-185, Step b (180 mg, 1.427 mmol) and pyridine (0.462 mL, 5.71 mmol) in 5 mL of CH₂Cl₂ (5 mL) and the mixture was stirred at room temperature for 2 days. The reaction was diluted with CH₂Cl₂ and washed with 1 N aq. HCl. The aqueous layer was extracted with CH₂Cl₂ (2X). The combined organic layers were dried with MgSO₄ and concentrated. The crude product was purified by flash chromatography (silica gel, 0-15% EtOAc/Hex) to afford Cap-185, Step c (210 mg) as a white solid. ¹H NMR (500 MHz, CDCl₃) δ ppm 7.73 (2 H, d, *J*=8.24 Hz), 7.32 (2 H, d, *J*=8.24 Hz), 6.25 (2 H, s), 4.76 (1 H, t, *J*=5.65 Hz), 4.64 (2 H, d, *J*=3.66 Hz), 2.44 (3 H, s), 2.18 (2 H, td, *J*=10.07, 5.49 Hz), 1.71 (2 H, d, *J*=15.56 Hz).


Cap-185, Step d



- [00328] A microwave tube was charged with benzyl 2-(diphenylmethyleneamino)acetate (1.5 g, 4.57 mmol) and Cap-185, Step c (1.28 g, 4.57

mmol) in 5 mL of toluene. The tube was sealed and LiHMDS (5.5 mL, 5.5 mmol) (1 N in toluene) was added dropwise under N₂. The resulting dark brown solution was heated at 100 °C in microwave for 5 hours. To the mixture was then added water and EtOAc. The layers were separated and the water phase was extracted with EtOAc (2X). The combined organic layers were concentrated to afford Cap-185, Step d as a racemic mixture of. The crude mixture was submitted to the next step without purification or separation. LC-MS: Anal. Calcd. for [M+H]⁺ C₂₉H₂₈NO₃ 438.21; found 438.4.


Cap-185, Step e



10

[00329] To a solution of the racemic mixture of Cap-185, Step d in 30 mL of THF was added HCl (20 mL) (2 N aq.). The resulting mixture was stirred at room temperature for 2 hours. After the reaction was done as judged by TLC, the two layers were separated. The aqueous layer was washed with EtOAc, neutralized with sat. NaHCO₃ aq. solution and then extracted with EtOAc (3X). The combined organic layers were dried with MgSO₄ and concentrated to afford Cap-185, Step e. LC-MS: Anal. Calcd. for [M+H]⁺ C₁₆H₂₀NO₃ 274.14; found 274.12.

Cap-185, Step f



20

[00330] A solution of the crude Cap-185, Step e, DiPEA (1.24 mL, 7.1 mmol) and methyl chloroformate (0.55 mL, 7.1 mmol) in 5 mL of CH₂Cl₂ was stirred at room temperature for 1 hour. The mixture was then diluted with CH₂Cl₂ and washed with water. The organic layer was dried with Na₂SO₄ and concentrated. The crude product was purified by flash chromatography (silica gel, 0-40% EtOAc/Hex) to afford 700 mg of the racemic mixture. The mixture was then separated by chiral HPLC (CHIRALPAK® AD-H column, 30 x 250 mm, 5 um) eluting with 88% CO₂-12% EtOH at 70 mL/min to afford

25

240 mg of Enantiomer-1 and 310 mg of Enantiomer-2 of Cap-1, Step f as white solids.

Enantiomer-1: LC-MS: Anal. Calcd. for $[M+H]^+$ $C_{18}H_{22}NO_5$ 332.15; found 332.3. 1H

NMR (500 MHz, $CDCl_3$) δ ppm 7.30 - 7.40 (5 H, m), 6.03 - 6.16 (2 H, m), 5.09 - 5.26 (3 H, m), 4.65 - 4.74 (2 H, m), 4.33 (1 H, dd, $J=9.16, 4.88$ Hz), 3.67 (3 H, s), 2.27 - 2.38 (1


5 H, m), 1.61 - 1.69 (1 H, m), 1.45 - 1.56 (1 H, m), 1.34 (1 H, dd, $J=13.43, 5.19$ Hz), 1.07 (1 H, dd, $J=13.12, 5.19$ Hz). Enantiomer-2: LC-MS: Anal. Calcd. for $[M+H]^+$ $C_{18}H_{22}NO_5$ 332.15; found 332.06.

Cap-185 (Enantiomer-1 and Enantiomer-2)

10 **[00331]** To a hydrogenation bottle containing a solution Cap-185, Step f (Enantiomer-2) (300 mg, 0.905 mmol) in 10 mL of MeOH was added Pd/C (15 mg, 0.141 mmol) under a cover of nitrogen. The mixture was hydrogenated on a Parr shaker at 40 psi for 3 hours. The mixture was then filtered and the filtrate was concentrated to afford Cap-185 (Enantiomer-2) (200 mg) as a white solid. LC-MS: Anal. Calcd. for $[M+H]^+$ $C_{11}H_{18}NO_5$ 244.12; found 244.2. 1H NMR (500 MHz, $CDCl_3$) δ ppm 5.33 (1 H, br. s.), 4.46 (2 H, d), 4.28 (1 H, br. s.), 3.68 (3 H, s), 2.35 (1 H, br. s.), 1.91 - 2.03 (2 H, m), 1.56 - 1.80 (4 H, m), 1.36 - 1.55 (2 H, m). [Note: Cap-185 (Enantiomer-1) can be obtained in a similar fashion.]

20


Cap-186



[00332] To a solution of the ester Cap-185, Step f (Enantiomer-2) (150 mg, 0.453 mmol) in 4 mL of MeOH was added NaOH (4 mL of 1 N in water, 4.00 mmol). The resulting mixture was stirred at room temperature for 3 hours. The methanol was then removed under vacuum, and the residue was neutralized with 1 N HCl solution and extracted with EtOAc (3X). The combined organic layers were dried with $MgSO_4$ and concentrated to afford Cap-186 that was contaminated with some benzyl alcohol (sticky white solid; 115 mg). LC-MS: Anal. Calcd. for $[M+H]^+$ $C_{11}H_{16}NO_5$ 242.10; found 242.1. 1H NMR (500 MHz, $CDCl_3$) δ ppm 6.10 - 6.19 (2 H, m), 5.36 (1 H, d, $J=8.85$ Hz), 4.75 -


4.84 (2 H, m), 4.28 (1 H, dd, $J=8.55, 4.58$ Hz), 3.68 (3 H, s), 2.33 - 2.45 (1 H, m), 1.60 - 1.72 (2 H, m), 1.30 - 1.48 (2 H, m).

Cap-187



5


Cap-187, Step a



- [00333] To a solution of Cap-178, Step e (2.2 g, 18.94 mmol), PPh_3 (24.84 g, 95 mmol) and 4-nitrobenzoic acid (14.24 g, 85 mmol) in 30 mL of benzene was added DEAD (42.9 mL, 95 mmol) dropwise. The resulting light orange solution was stirred at room temperature overnight. The solvent was then removed under vacuum and the residue was purified by flash chromatography (silica gel, 0-15% EtOAc/Hex) to afford Cap-187, Step a (2.3 g) as a white solid. ^1H NMR (500 MHz, CDCl_3) δ ppm 8.27 - 8.34 (2 H, m), 8.20 - 8.26 (2 H, m), 5.45 (1 H, t, $J=2.90$ Hz), 3.83 - 3.96 (3 H, m), 1.90 - 2.03 (2 H, m), 1.80 - 1.88 (1 H, m), 1.61 - 1.70 (1 H, m), 1.21 (3 H, d, $J=6.10$ Hz).

15

Cap-187, Step b



- [00334] To a solution of Cap-187, Step a (2.3 g, 8.67 mmol) in 10 mL of MeOH was added sodium methoxide (2.372 mL, 8.67 mmol) (25% in Methanol). The resulting mixture was stirred at room temperature for 3 hours. Water was added, and the mixture was extracted with EtOAc (5X). The combined organic layers were dried with MgSO_4 and concentrated. The crude product was purified by flash chromatography (silica gel, 0-

20

15% EtOAc/Hex, then 15-50% EtOAc/Hex) to afford Cap-187, Step b (0.85 g) as clear oil. ¹H NMR (500 MHz, CDCl₃) δ ppm 4.19 - 4.23 (1 H, m), 3.82 - 3.91 (2 H, m), 3.73 - 3.79 (1 H, m), 1.79 - 1.88 (1 H, m), 1.62 - 1.68 (1 H, m), 1.46 - 1.58 (2 H, m), 1.14 (3 H, d, *J*=6.10 Hz).


5

Cap-187

[00335] The individual enantiomers of Cap-187 were synthesized from Cap-187, Step b according to the procedure described for Cap-178. LC-MS: Anal. Calcd. for [M+H]⁺ C₁₀H₁₈NO₅ 232.12; found 232.1. ¹H NMR (400 MHz, CDCl₃) δ ppm 5.26 (1 H, d, *J*=7.78 Hz), 4.32 - 4.43 (1 H, m), 4.07 (1 H, dd, *J*=11.54, 3.51 Hz), 3.72 (3 H, s), 3.39 - 3.50 (2 H, m), 2.08 - 2.23 (1 H, m), 1.54 - 1.68 (1 H, m), 1.38 - 1.52 (1 H, m), 1.11 - 1.32 (5 H, m).


10

Cap-188 (four stereoisomers)



15

Cap-188, Step a




[00336] To a solution of 2,2-dimethyldihydro-2H-pyran-4(3H)-one (2 g, 15.60 mmol) in 50 mL of MeOH was slowly added sodium borohydride (0.649 g, 17.16 mmol). The resulting mixture was stirred at room temperature for 3 hours. To the mixture was then added 1 N HCl aqueous solution until it crosses into acidic pH range and then extracted with EtOAc (3X). The combined organic layers were dried with MgSO₄ and concentrated to afford Cap-188, Step a (1.9 g) as clear oil. The product was used in the next step without purification. ¹H NMR (400 MHz, CDCl₃) δ ppm 3.91 - 4.02 (1 H, m), 3.79 - 3.86 (1 H, m), 3.63 (1 H, td, *J*=12.05, 2.51 Hz), 1.82 - 1.93 (2 H, m), 1.40 - 1.53 (1 H, m), 1.29 - 1.38 (1 H, m), 1.27 (3 H, s), 1.20 (3 H, s).

20

25

Cap-188.1 and Cap-188.2, Step b




[00337] p-Tosyl-Cl (5.56 g, 29.2 mmol) was added to a solution of Cap-188, Step a
 5 (1.9 g, 14.59 mmol) and pyridine (4.72 mL, 58.4 mmol) in 100 mL of CH₂Cl₂. The
 resulting mixture was stirred at room temperature for 3 days. To the reaction was added
 10 mL of water, and the mixture was stirred at room temperature for an additional hour.
 The two layers were separated and the organic phase was washed with water and 1 N HCl
 aqueous solution. The organic phase was dried with MgSO₄ and concentrated to afford
 10 the mixture of two enantiomers as a light yellow solid. The mixture was then separated by
 chiral HPLC (CHIRALPAK® AD column, 21 x 250 mm, 10 um) eluting with 92%
 0.1% diethylamine/Heptane-8% EtOH at 15 mL/min to afford Cap-188.1, Step b (1.0 g)
 and Cap-188.2, Step b (1.0 g). The absolute stereochemistry of the two enantiomers was
 not assigned. Cap-188.1, Step b: LC-MS: Anal. Calcd. for [2M+Na]⁺ C₂₈H₄₀NaO₈S₂
 15 591.21; found 591.3. ¹H NMR (500 MHz, CDCl₃) δ ppm 7.79 (2 H, d, *J*=8.24 Hz), 7.34
 (2 H, d, *J*=8.24 Hz), 4.72 - 4.81 (1 H, m), 3.78 (1 H, dt, *J*=12.44, 4.16 Hz), 3.53 - 3.61 (1
 H, m), 2.45 (3 H, s), 1.75 - 1.86 (2 H, m), 1.61 - 1.71 (1 H, m), 1.52 - 1.60 (1 H, m), 1.22
 (3 H, s), 1.14 (3 H, s). Cap-188.2, Step b: LC-MS: Anal. Calcd. for [2M+Na]⁺
 C₂₈H₄₀NaO₈S₂ 591.21; found 591.3;

20

Cap-188


[00338] The four stereoisomers of Cap-188 could be synthesized from Cap-188.1, Step
 b and Cap-188.2, Step b, according to the procedure described for the preparation of Cap-
 178. Cap-188 (Stereoisomer-1): LC-MS: Anal. Calcd. for [M+Na]⁺ C₁₁H₁₉NNaO₅ 268.12;
 25 found 268.23. ¹H NMR (500 MHz, CDCl₃) δ ppm 5.32 (1 H, d, *J*=8.55 Hz), 4.26 - 4.35 (1
 H, m), 3.57 - 3.82 (5 H, m), 2.11 - 2.34 (1 H, m), 1.25 - 1.58 (4 H, m), 1.21 (6 H, d,
J=6.10 Hz). Cap-188 (Stereoisomer-2): LC-MS: Anal. Calcd. for [M+H]⁺ C₁₁H₂₀NO₅
 246.13; found 246.1. ¹H NMR (500 MHz, CDCl₃) δ ppm 5.25 (1 H, d, *J*=8.55 Hz), 4.33
 (1 H, dd, *J*=8.39, 5.04 Hz), 3.80 (1 H, dd, *J*=11.90, 3.97 Hz), 3.62 - 3.76 (4 H, m), 2.20 -
 30 2.32 (1 H, m), 1.52 - 1.63 (1 H, m), 1.27 - 1.49 (3 H, m), 1.22 (6 H, d, *J*=14.04 Hz).

Cap-189




5

Cap-189, Step a



- [00339]** To a solution of phenylmagnesium bromide (113 mL, 340 mmol) (3 M in ether) in 100 mL of ether was added dropwise exo-2,3-epoxynorbornane (25 g, 227 mmol) in 50 mL of ether. After the initial exotherm, the mixture was heated to reflux overnight. The reaction was then cooled to room temperature and quenched carefully with water (~ 10 mL). The mixture was diluted with ether and washed with a 3 N HCl aqueous solution (~ 160 mL). The aqueous layer was extracted with ether (2X) and the combined organic layers were dried with MgSO₄ and concentrated. The crude product was purified by flash chromatography (silica gel, 0-18% EtOAc/Hex) to afford Cap-189, Step a (11 g).
- ¹H NMR (400 MHz, CDCl₃) δ ppm 6.03 - 6.11 (2 H, m), 3.76 (1 H, d, *J*=11.29 Hz), 2.72 - 2.81 (2 H, m), 1.98 (1 H, d, *J*=11.29 Hz), 1.67 - 1.76 (2 H, m), 0.90 - 0.97 (2 H, m).

Cap-189, Step b




- [00340]** To a solution of oxalyl chloride (59.9 mL, 120 mmol) in 200 mL of CH₂Cl₂ at -78 °C was added DMSO (17.01 mL, 240 mmol) in 100 mL of CH₂Cl₂. The mixture was stirred for 10 min, and Cap-189, Step a (11 g, 100 mmol) in 150 mL of CH₂Cl₂ was added followed by Et₃N (72.4 mL, 519 mmol) in 30 mL of CH₂Cl₂. The mixture was stirred at -78 °C for 30 min and then warmed to room temperature. Water (150 mL) was added and the mixture was stirred at room temperature for 30 mins. The two layers were then separated, and the aqueous layer was extracted with CH₂Cl₂ (2X). The organic layers

were combined, dried with MgSO_4 and concentrated. The crude product was purified by flash chromatography (silica gel, 0-5% EtOAc/Hex) to afford Cap-189, Step b (5.3 g) as a light yellow oil. ^1H NMR (500 MHz, CDCl_3) δ ppm 6.50 - 6.55 (2 H, m), 2.78 - 2.84 (2 H, m), 1.92 - 1.99 (2 H, m), 1.17 - 1.23 (2 H, m).

5


Cap-189, Step c



- [00341] A mixture of Cap-189, Step b (5.3 g, 49.0 mmol), *p*-toluenesulfonic acid monohydrate (1.492 g, 7.84 mmol) and ethylene glycol (4.10 mL, 73.5 mmol) in 100 mL of benzene was refluxed for 4 hours and then stirred at room temperature overnight. The reaction was partitioned between Et_2O and aqueous sat. NaHCO_3 solution and the two layers were separated. The organic layer was washed with brine, dried with MgSO_4 and concentrated. The crude product was purified by flash chromatography (silica gel, 0-6% EtOAc/Hex) to afford Cap-189, Step c (5.2 g) as a clear oil. ^1H NMR (400 MHz, CDCl_3) δ ppm 6.20 (2 H, t, $J=2.13$ Hz), 3.90 - 3.97 (2 H, m), 3.81 - 3.89 (2 H, m), 2.54 (2 H, m), 1.89 - 1.99 (2 H, m), 0.95 - 1.03 (2 H, m).

15


Cap-189, Step d



- [00342] A solution of Cap-189, Step c (5.2 g, 34.2 mmol) in 60 mL of MeOH and 50 mL of CH_2Cl_2 was cooled to -78°C and treated with ozone gas until a light blue color was apparent. The reaction was then bubbled with N_2 to remove the excess ozone gas (blue color disappeared) and sodium borohydride (1.939 g, 51.3 mmol) was added into the reaction. The reaction was then warmed to 0°C . Acetone was added into the mixture to quench the excess sodium borohydride. The mixture was concentrated and the residue was purified by flash chromatography (silica gel, 100% EtOAc) to afford Cap-189, Step d (5.0 g) as a clear oil. ^1H NMR (400 MHz, CDCl_3) δ ppm 3.99 - 4.09 (4 H, m), 3.68 (4 H, m), 2.17 - 2.29 (2 H, m), 1.92 - 2.10 (2 H, m), 1.77 - 1.88 (2 H, m), 1.57 - 1.70 (2 H, m).


25

Cap-189, Step e



- [00343] To a solution of Cap-189, Step d (1 g, 5.31 mmol) in 20 mL of CH₂Cl₂ was added silver oxide (3.8 g), p-Ts-Cl (1.215 g, 6.38 mmol) and KI (0.176 g, 1.063 mmol). The resulting solution was stirred at room temperature for 3 days. The mixture was then filtered and the filtrate was concentrated. The crude product was purified by flash chromatography (silica gel, 60% EtOAc/Hex) to afford Cap-189, Step e (0.79 g) as clear oil. LC-MS: Anal. Calcd. for [M+Na]⁺ C₁₆H₂₂NaO₆S 365.10; found 365.22. ¹H NMR (400 MHz, CDCl₃) δ ppm 7.80 (2 H, d, *J*=8.28 Hz), 7.36 (2 H, d, *J*=8.03 Hz), 4.11 - 4.17 (1 H, m), 3.85 - 4.06 (5 H, m), 3.64 - 3.71 (1 H, m), 3.55 - 3.63 (1 H, m), 2.47 (3 H, s), 2.32 - 2.43 (1 H, m), 2.15 - 2.27 (1 H, m), 1.70 - 1.89 (2 H, m), 1.52 - 1.66 (1 H, m), 1.35 - 1.47 (1 H, m).

15 Cap-189, Step f




- [00344] To a solution of Cap-189, Step e (2.2 g, 6.43 mmol) in 40 mL of MeOH was added potassium carbonate (1.776 g, 12.85 mmol). The resulting mixture was stirred at room temperature overnight. The mixture was then diluted with water and EtOAc. The two layers were separated. The aqueous layer was extracted with EtOAc (2X). The combined organic layers were washed with brine, dried with MgSO₄ and concentrated. The crude product was purified by flash chromatography (silica gel, 0-15% EtOAc/Hex) to afford Cap-189, Step f (0.89 g, 5.23 mmol, 81%) as clear oil. ¹H NMR (400 MHz, CDCl₃) δ ppm 3.89 - 4.02 (6 H, m), 3.58 (2 H, dd, *J*=10.79, 2.51 Hz), 1.69 - 1.89 (6 H, m).

Cap-189, Step g



[00345] To the solution of Cap-189, Step f (890 mg, 5.23 mmol) in 15 mL of THF was added HCl (15 mL, 45.0 mmol) (3 M aqueous). The resulting mixture was stirred at room temperature overnight. The mixture was then diluted with ether and the two layers were separated. The aqueous phase was extracted with ether (2X) and the combined organic layers were dried with MgSO₄ and concentrated to afford Cap-189, Step g (0.95 g, containing some residual solvents). The product was used in the next step without purification. ¹H NMR (500 MHz, CDCl₃) δ ppm 3.95 - 4.00 (2 H, m), 3.85 (2 H, d, *J*=10.68 Hz), 2.21 - 2.28 (2 H, m), 1.99 - 2.04 (2 H, m), 1.90 - 1.96 (2 H, m).

Cap-189, Step h (Enantiomer-1 and Enantiomer-2)




[00346] To a solution of (+/-)-benzyloxycarbonyl-α-phosphonoglycine trimethyl ester (1733 mg, 5.23 mmol) in 6 mL of THF at -20 °C was added 1,1,3,3-tetramethylguanidine (0.723 mL, 5.75 mmol). The resultant light yellow mixture was stirred at -20 °C for 1 hour, and Cap-189, Step g (660 mg, 5.23 mmol) in 3 mL of THF was added and mixture was then stirred at room temperature for 3 days. The reaction mixture was then diluted with EtOAc, washed with a 0.1 N HCl aq. solution. The aqueous layer was extracted with EtOAc (2X) and the combined organic layers were dried with MgSO₄ and concentrated.

The crude product was purified by flash chromatography (silica gel, 0-4% EtOAc/CH₂Cl₂) to afford 960 mg of the racemic mixture. The mixture was separated by chiral HPLC (CHIRALPAK® AD column, 21 x 250 mm, 10 μm) eluting with 90% 0.1% diethylamine/Heptane-10% EtOH at 15 mL/min to afford Cap-189, Step h (Enantiomer-1; 300 mg) and Cap-189, Step h (Enantiomer-2; 310 mg) as white solids. Cap-189, Step h (Enantiomer-1): LC-MS: Anal. Calcd. for [M+H]⁺ C₁₈H₂₂NO₅ 332.15; found 332.2. ¹H NMR (500 MHz, CDCl₃) δ ppm 7.29 - 7.41 (5 H, m), 6.00 (1 H, br. s.), 5.13 (2 H, s), 3.63 - 3.87 (8 H, m), 2.84 (1 H, br. s.), 1.84 - 2.02 (2 H, m), 1.63 - 1.84 (2 H, m). Cap-189,

Step h (Enantiomer-2): LC-MS: Anal. Calcd. for $[M+H]^+$ $C_{18}H_{22}NO_5$ 332.15; found 332.2.

Cap-189, Step i




5

[00347] N_2 was bubbled through a solution of Cap-189, Step h (Enantiomer-2; 290 mg, 0.875 mmol) in 10 mL of MeOH in a 500 mL hydrogenation bottle for 30 mins. To the solution was added (S,S)-Me-BPE-Rh (9.74 mg, 0.018 mmol), and the mixture was then hydrogenated at 60 psi for 6 days. The mixture was concentrated, and chiral analytical

10 HPLC (CHIRALPAK® OJ column) indicated that there were a small amount of remaining starting material and one major product. The residue was then separated by chiral HPLC (CHIRALPAK® OJ column, 21 x 250 mm, 10 μ m) eluting with 70% 0.1% diethylamine/ Heptane-30% EtOH at 15 mL/min to afford Cap-189, Step i, (150 mg) as clear oil. LC-MS: Anal. Calcd. for $[M+H]^+$ $C_{18}H_{24}NO_5$ 334.17; found 334.39. 1H NMR (500 MHz, $CDCl_3$) δ ppm 7.28 - 7.41 (5 H, m), 5.12 - 5.18 (1 H, m), 5.09 (2 H, s), 4.05 (1 H, t, $J=10.07$ Hz), 3.75 (3 H, s), 3.60 - 3.72 (2 H, m), 3.41 - 3.50 (2 H, m), 2.10 (1 H, br. s.), 1.72 - 1.99 (6 H, m).

15

Cap-189, Step j



20

[00348] To a solution of Cap-189, Step i (150 mg, 0.450 mmol) in 10 mL of MeOH in a hydrogenation bottle were added dimethyl dicarbonate (0.072 mL, 0.675 mmol) and 10% Pd/C (23.94 mg, 0.022 mmol) under a cover of nitrogen cover. The mixture was then hydrogenated on Parr-shaker at 45 psi overnight. The mixture was filtered and the filtrate


25 was concentrated to afford Cap-189, Step j (110 mg) as a clear oil. LC-MS: Anal. Calcd. for $[M+H]^+$ $C_{12}H_{20}NO_5$ 258.13; found 258.19. 1H NMR (500 MHz, $CDCl_3$) δ ppm 5.08 (1

H, d, $J=9.16$ Hz), 4.03 (1 H, t, $J=10.07$ Hz), 3.75 (3 H, s), 3.60 - 3.72 (5 H, m), 3.46 (2 H, t, $J=10.38$ Hz), 2.11 (1 H, br. s.), 1.72 - 1.99 (6 H, m).

Cap-189


- 5 [00349] To a mixture of Cap-189, Step j (110 mg, 0.428 mmol) in 2 mL of THF and 1 mL of water was added LiOH (0.641 mL, 1.283 mmol) (2 N aq.). The resulting mixture was stirred at room temperature overnight. The mixture was neutralized with a 1 N HCl aq. solution and extracted with EtOAc (3X). The combined organic layers were dried with MgSO_4 and concentrated to afford Cap-189 (100 mg) as a white solid. LC-MS: Anal.
- 10 Calcd. for $[\text{M}+\text{Na}]^+ \text{C}_{11}\text{H}_{17}\text{NNaO}_5$ 266.10; found 266.21. ^1H NMR (500 MHz, CDCl_3) δ ppm 5.10 (1 H, d, $J=9.16$ Hz), 4.02 (1 H, t, $J=10.07$ Hz), 3.62 - 3.78 (5 H, m), 3.49 (2 H, d, $J=10.68$ Hz), 2.07 - 2.22 (2 H, m), 1.72 - 1.98 (6 H, m).

Cap-190 (diastereomeric mixture)




15

Cap-190, Step a




- [00350] To a mixture of cyclopent-3-enol (2.93 g, 34.8 mmol) and imidazole (5.22 g, 77 mmol) in 30 mL of DMF at 0 °C was added *t*-butyldimethylchlorosilane (6.30 g, 41.8 mmol). The resulting colorless mixture was stirred at room temperature overnight. Hexanes and water were then added to the mixture and the two layers were separated. The aqueous layer was extracted with EtOAc (2X) and the combined organic layers were washed with brine, dried with MgSO_4 and concentrated. The crude product was purified
- 20 by flash chromatography (silica gel, 2% EtOAc/Hex) to afford Cap-190, Step a (6.3 g) as a clear oil. ^1H NMR (500 MHz, CDCl_3) δ ppm 5.65 (2 H, s), 4.49 - 4.56 (1 H, m), 2.56 (2 H, dd, $J=15.26, 7.02$ Hz), 2.27 (2 H, dd, $J=15.26, 3.36$ Hz), 0.88 (9 H, s), 0.06 (6 H, s).
- 25

Cap-190, Step b




[00351] To a solution of Cap-190, Step a (2.3 g, 11.59 mmol) in 40 mL of CH₂Cl₂ at 0 °C was added m-CPBA (5.60 g, 16.23 mmol) in 5 portions. The reaction mixture was stirred at room temperature overnight. Hexanes and water were then added to the mixture and the two layers were separated. The organic layer was washed with 50 mL aq. 10% NaHSO₃ and brine, dried with MgSO₄ and concentrated. The crude product was purified by flash chromatography (silica gel, 3%-6% EtOAc/Hex) to afford Cap-190, Step b (1.42 g) and its trans diastereomer (0.53 g) as clear oils. Cap-190, Step b (*cis*): ¹H NMR (400 MHz, CDCl₃) δ ppm 4.39 - 4.47 (1 H, m), 3.47 (2 H, s), 2.01 - 2.10 (2 H, m), 1.93 - 2.00 (2 H, m), 0.88 (9 H, s), 0.04 (6 H, s). Cap-190, Step b (*trans*): ¹H NMR (400 MHz, CDCl₃) δ ppm 4.04 - 4.14 (1 H, m), 3.47 (2 H, s), 2.41 (2 H, dd, *J*=14.05, 7.28 Hz), 1.61 (2 H, dd, *J*=14.18, 6.90 Hz), 0.87 (9 H, s), 0.03 (6 H, s).

Cap-190, Step c




[00352] To a solution of (S)-1,2'-methylenedipyrrolidine (0.831 g, 5.39 mmol) in 15 mL of benzene at 0 °C was added dropwise *n*-butyllithium (4.90 mL, 4.90 mmol) (1 M in hexane). The solution turned bright yellow. The mixture was stirred at 0 °C for 30 min. Cap-190, Step b (*cis*-isomer; 0.7 g, 3.27 mmol) in 10 mL of benzene was then added and the resulting mixture was stirred at 0 °C for 3 hours. EtOAc and sat. NH₄Cl aq. solution were added into the mixture, and the two layers were separated. The organic layer was washed with water and brine, dried with MgSO₄ and concentrated. The crude product was purified by flash chromatography (silica gel, 15% EtOAc/Hex) to afford Cap-190, Step c (400 mg) as a light yellow oil. ¹H NMR (500 MHz, CDCl₃) δ ppm 5.84 - 5.98 (2 H, m), 4.53 - 4.69 (2 H, m), 2.63 - 2.73 (1 H, m), 1.51 (1 H, dt, *J*=13.73, 4.43 Hz), 0.89 (9 H, s), 0.08 (6 H, s).

Cap-190, Step d




[00353] To a solution of Cap-190, Step c (400 mg, 1.866 mmol), MeI (1.866 mL, 3.73 mmol) (2 M in *t*-butyl methyl ether) in 5 mL of THF at 0 °C was added NaH (112 mg, 2.80 mmol) (60% in mineral oil). The resulting mixture was allowed to warm up to room temperature and stirred at room temperature overnight. The reaction was then quenched with water and extracted with EtOAc (3X). The combined organic layers were washed with brine, dried with MgSO₄ and concentrated. The crude product was purified by flash chromatography (silica gel, 5% EtOAc/Hex) to afford Cap-190, Step d (370 mg) as light yellow oil. ¹H NMR (500 MHz, CDCl₃) δ ppm 5.92 - 5.96 (1 H, m), 5.87 - 5.91 (1 H, m), 4.64 - 4.69 (1 H, m), 4.23 - 4.28 (1 H, m), 3.32 (3 H, s), 2.62 - 2.69 (1 H, m), 1.54 (1 H, dt, *J*=13.12, 5.49 Hz), 0.89 (9 H, s), 0.07 (5 H, d, *J*=1.83 Hz).

Cap-190, Step e



[00354] To a solution of Cap-190, Step d (400 mg, 1.751 mmol) in 10 mL of EtOAc in a hydrogenation bottle was added platinum(IV) oxide (50 mg, 0.220 mmol). The resulting mixture was hydrogenated at 50 psi on Parr shaker for 2 hours. The mixture was then filtered through CELITE®, and the filtrate was concentrated to afford Cap-190, Step e (400 mg) as a clear oil. LC-MS: Anal. Calcd. for [M+H]⁺ C₁₂H₂₇O₂Si 231.18; found 231.3. ¹H NMR (500 MHz, CDCl₃) δ ppm 4.10 - 4.17 (1 H, m), 3.65 - 3.74 (1 H, m), 3.27 (3 H, s), 1.43 - 1.80 (6 H, m), 0.90 (9 H, s), 0.09 (6 H, s).

Cap-190, Step f



[00355] To a solution of Cap-190, Step e (400 mg, 1.736 mmol) in 5 mL of THF was added TBAF (3.65 mL, 3.65 mmol) (1 N in THF). The color of the mixture turned brown after several min., and it was stirred at room temperature overnight. The volatile


5 component was removed under vacuum, and the residue was purified by flash chromatography (silica gel, 0-25% EtOAc/Hex) to afford Cap-190, Step f (105 mg) as light yellow oil. ¹H NMR (500 MHz, CDCl₃) δ ppm 4.25 (1 H, br. s.), 3.84 - 3.92 (1 H, m), 3.29 (3 H, s), 1.67 - 2.02 (6 H, m).

10 Cap-190


[00356] Cap-190 was then synthesized from Cap-190, Step f according to the procedure described for Cap-182. LC-MS: Anal. Calcd. for [M+Na]⁺ C₁₀H₁₇NNaO₅ 254.10; found 254.3. ¹H NMR (500 MHz, CDCl₃) δ ppm 5.25 (1 H, d, *J*=8.55 Hz), 4.27 - 4.41 (1 H, m), 3.81 - 3.90 (1 H, m), 3.69 (3 H, s), 3.26 (3 H, s), 2.46 - 2.58 (1 H, m), 1.76 - 1.99 (3 H, m), 1.64 - 1.73 (1 H, m), 1.40 - 1.58 (1 H, m), 1.22 - 1.38 (1 H, m).

15

Cap-191 (Enantiomer-1)



20 Cap-191, Step a




[00357] To a solution of diisopropylamine (3 mL, 21.05 mmol) in THF (3 mL) at -78 °C under nitrogen was added n-butyl lithium (2.5 M in hexanes; 8.5 mL, 21.25 mmol). The reaction was stirred at -78 °C for 10 min then brought up to 0 °C for 25 min. The

25 reaction was cooled down again to -78 °C, methyl tetrahydro-2H-pyran-4-carboxylate (3 g, 20.81 mmol) in THF (3 mL) was added. The reaction was stirred at -78 °C for 15 min

then brought up to 0 °C for 30 min. The reaction was cooled down to -78 °C, methyl iodide (1.301 ml, 20.81 mmol) was added. After the addition, the cold bath was removed and the reaction was allowed to slowly warm up to ~25 °C and stirred for 22h. Ethyl acetate and aqueous HCl (0.1N) were added, and the organic layer was separated and washed with brine and dried (MgSO₄), filtered, and concentrated *in vacuo*. The residue was loaded on a Thomson's silica gel cartridge eluting with 10% ethyl acetate/hexanes to afford a light yellow oil (2.83 g). ¹H NMR (400 MHz, DMSO-d₆) δ ppm 3.73-3.66 (m, 2H), 3.66 (s, 3H), 3.40-3.30 (m, 2H), 1.95-1.93 (dm, 1H), 1.92-1.90 (dm, 1H), 1.43 (ddd, J = 13.74, 9.72, 3.89, 2H), 1.18 (s, 3H).

10


Cap-191, Step b



[00358] To a solution of Cap-191, Step a (3 g, 18.96 mmol) in toluene (190 ml) at -78 °C under nitrogen was added diisobutylaluminum hydride (1.5M in toluene; 26.5 ml, 39.8 mmol) dropwise. The reaction was continued to stir at -78 °C for 1.5h., and the bath was removed and was stirred for 18h. The reaction was quenched with MeOH (20 mL). HCl (1M, 150 mL) was added and the mixture was extracted with EtOAc (4 x 40 mL). The combined organic phases were washed with brine, dried (MgSO₄), filtered, and concentrated *in vacuo*. The residue was purified with flash chromatography (silica gel; 40% ethyl acetate/hexanes) to afford a colorless oil (1.36 g). ¹H NMR (400 MHz, CDCl₃) δ ppm 3.77 (dt, J = 11.73, 4.55, 2H), 3.69-3.60 (m, 2H), 3.42 (s, 2H), 1.71-1.40 (bs, 1H) 1.59 (ddd, J = 13.74, 9.72, 4.39, 2H), 1.35-1.31 (m, 1H), 1.31-1.27 (m, 1H), 1.06 (s, 3H).

25


Cap-191, Step c



[00359] To a solution of DMSO (5.9 ml, 83 mmol) in CH₂Cl₂ (85 ml) at -78 °C under nitrogen was added oxalyl chloride (3.8 ml, 43.4 mmol) and stirred for 40 min. A solution of Cap-191, Step b (4.25 g, 32.6 mmol) in CH₂Cl₂ (42.5 ml) was then added.

The reaction was continued to be stirred at -78 °C under nitrogen for 2h. The reaction was quenched with cold 20% K₂HPO₄ (aq) (10 mL) and water. The mixture was stirred at ~25 °C for 15 min, diluted with diethyl ether (50 mL) and the layers were separated. The aqueous layer was extracted with diethyl ether (2 x 50 mL). The combined organic layers were washed with brine, dried (MgSO₄), filtered, and concentrated *in vacuo*. The residue was taken up in CH₂Cl₂ (4 mL) and purified with flash chromatography (silica gel, eluting with CH₂Cl₂) to afford a colorless oil (2.1 g). ¹H NMR (400 MHz, CDCl₃) δ ppm 9.49 (s, 1H), 3.80 (dt, J = 11.98, 4.67, 2H), 3.53 (ddd, J = 12.05, 9.41, 2.89, 2H), 1.98 (ddd, J = 4.71, 3.20, 1.38, 1H), 1.94 (ddd, J = 4.71, 3.20, 1.38, 1H), 1.53 (ddd, J = 13.87, 9.60, 4.14, 2H), 1.12 (s, 3H).


Cap-191, Step d



[00360] To a solution of Cap-191c (2.5 g, 19.51 mmol) in CHCl₃ (20 mL) under nitrogen at ~25 °C was added (*R*)-2-amino-2-phenylethanol (2.94 g, 21.46 mmol) and stirred for 5h. The reaction was cooled to 0 °C, trimethylsilyl cyanide (3.8 mL, 30.4 mmol) was added dropwise. The cold bath was removed and the reaction was allowed to stir at ~25 °C under nitrogen for 15.5 h. The reaction was treated with 3N HCl (20 mL) and water (20 mL), and the product was extracted with CHCl₃ (3 x 50 mL). The combined organic layers were dried (NaSO₄), filtered, and concentrated *in vacuo*. The residue was purified with flash chromatography (silica gel; 40% ethyl acetate/hexanes) to afford two diastereomers: Cap-191, Step d1 (diastereomer 1) as a colorless oil which solidified into a white solid upon standing (3 g). ¹H NMR (400 MHz, DMSO-d₆) δ ppm 7.42-7.26 (m, 5H), 5.21 (t, J = 5.77, 1H), 3.87 (dd, J = 8.53, 4.52, 1H), 3.61-3.53 (m, 1H), 3.53-3.37 (m, 5H), 3.10 (d, J = 13.05, 1H), 2.65 (d, J = 13.05, 1H), 1.64-1.55 (m, 1H), 1.55-1.46 (m, 1H), 1.46-1.39 (m, 1H), 1.31-1.23 (m, 1H), 1.11 (s, 3H). LC-MS: Anal. Calcd. for [M+H]⁺ C₁₆H₂₃N₂O₂: 275.18; found 275.20. Cap-191, Step d2 (diastereomer 2) as a light yellow oil (0.5 g). ¹H NMR (400 MHz, DMSO-d₆) δ ppm 7.44-7.21 (m, 5H), 4.82 (t, J = 5.40, 1H), 3.82-3.73 (m, 1H), 3.73-3.61 (m, 3H), 3.61-3.37 (m, 5H), 2.71 (dd,

$J = 9.29, 4.77, 1\text{H}$), $1.72\text{--}1.55$ (m, 2H), $1.48\text{--}1.37$ (m, 1H), $1.35\text{--}1.25$ (m, 1H), 1.10 (s, 3H). LC-MS: Anal. Calcd. for $[\text{M}+\text{H}]^+ \text{C}_{16}\text{H}_{23}\text{N}_2\text{O}_2$: 275.18; found 275.20.

Cap-191, Step e



5


[00361] To a solution of Cap-191, Step d2 (diastereomer 2) (0.4472 g, 1.630 mmol) in CH_2Cl_2 (11 ml) and MeOH (5.50 ml) at 0°C under nitrogen was added lead tetraacetate (1.445 g, 3.26 mmol). The reaction was stirred for 1.5h, the cold bath was removed and stirring was continued for 20h. The reaction was treated with a phosphate buffer ($\text{pH} = 7$; 6 mL) and stirred for 45 min. The reaction was filtered over CELITE®, washed with CH_2Cl_2 and the layers were separated. The aqueous layer was extracted with CH_2Cl_2 (3 X 25 mL), and the combined organic layers was washed with brine, dried (MgSO_4), filtered and concentrated *in vacuo*. The residue was purified with flash chromatography (silica gel; 15% ethyl acetate/hexanes) to afford the imine intermediate as a colorless oil (181.2 mg). ^1H NMR (400 MHz, DMSO-d_6) δ ppm 8.55 (d, $J = 1.00$, 1H), 7.89–7.81 (m, 2H), 7.61–7.46 (m, 3H), 4.80 (d, $J = 1.00$, 1H), 3.74 (tt, $J = 11.80, 4.02$, 2H), 3.62–3.46 (m, 2H), 1.79–1.62 (m, 2H), 1.46–1.30 (m, 2H), 1.15 (s, 3H).

[00362] The imine intermediate was taken up in 6N HCl (10 mL) and heated at 90°C for 10 days. The reaction was removed from the heat, allowed to cool to room temperature and extracted with ethyl acetate (3 x 25 mL). The aqueous layer was concentrated *in vacuo* to afford an off-white solid. The solid was taken up in MeOH and loaded on a pre-conditioned MCX (6 g) cartridge, washed with MeOH followed by elution with 2N NH_3/MeOH solution and concentrated *in vacuo* to afford an off-white solid (79.8 mg). ^1H NMR (400 MHz, DMSO-d_6) δ ppm 14.33–13.51 (bs, 1H), 8.30 (bs, 3H), 3.82–3.75 (m, 1H), 3.70 (dt, $J = 11.80, 4.02$, 2H), 3.58–3.43 (m, 2H), 1.76–1.60 (m, 2H), 1.47–1.36 (m, 1H), 1.36–1.27 (m, 1H), 1.08 (s, 3H). LC-MS: Anal. Calcd. for $[\text{M}+\text{H}]^+ \text{C}_8\text{H}_{16}\text{NO}_3$: 174.11; found 174.19.

Cap-191 (Enantiomer-1)


[00363] To a solution of Cap-191, Step e (0.0669 g, 0.386 mmol) and sodium carbonate (0.020 g, 0.193 mmol) in sodium hydroxide (1M aq.; 0.4 ml, 0.40 mmol) at 0 °C was added methyl chloroformate (0.035 ml, 0.453 mmol) dropwise. The reaction was removed from the cold bath and allowed to stir at ~25 °C for 3h. The reaction was washed with diethyl ether (3 x 20 mL). The aqueous layer was acidified with 12 N HCl (pH ~ 1-2), and extracted with ethyl acetate (2 x 20 mL). The combined organic layers were dried (MgSO₄), filtered, and concentrated *in vacuo* to afford Cap-191 as a colorless film (66.8 mg). ¹H NMR (400 MHz, DMSO-d₆) δ ppm 13.10-12.37 (bs, 1H), 7.37 (d, J = 9.04, 1H), 4.02 (d, J = 9.29, 1H), 3.72-3.57 (m, 2H), 3.56 (s, 3H), 3.54-3.44 (m, 2H), 1.65 (ddd, J = 13.61, 9.72, 4.27, 1H), 1.53 (ddd, J = 13.68, 9.66, 4.27, 1H), 1.41-1.31 (m, 1H), 1.31-1.22 (m, 1H), 1.00 (s, 3H). LC-MS: Anal. Calcd. for [M+Na]⁺ C₁₀H₁₇NO₅Na: 254.10; found 254.11.

Cap-192 (Enantiomer-2)




[00364] Cap-192 (Enantiomer-2) was prepared from Cap-191, Step d1 according to the procedure described for the preparation of its enantiomer Cap-191.

Cap-193




Cap-193, Step a



[00365] To a solution of methyl 2-(benzyloxycarbonylamino)-2-(dimethoxyphosphoryl)acetate (1.45 g, 4.2 mmol) in DCM was added DBU (0.70 ml, 4.7 mmol). The reaction mixture was stirred for 10 min, followed by addition of a solution of 1,3-dimethoxypropan-2-one (0.5 g, 4.2 mmol) in DCM. The reaction mixture was stirred at room temperature for 18 hrs. The reaction mixture was charged to an 80 g silica gel cartridge which was eluted with an 18 min gradient of 0-70% EtOAc in hexane to afford Cap-193, Step a (0.8 g) as a thick oil. ¹H NMR (400 MHz, MeOD) ppm 7.23 - 7.43 (5 H, m), 4.99 - 5.18 (2 H, m), 4.16 (2 H, s), 4.06 (2 H, s), 3.66 - 3.78 (3 H, s), 3.26 (3 H, s), 3.23 (3 H, s). LC-MS: Anal. Calcd. For [M+Na]⁺ C₁₆H₂₁NNaO₆: 346.14; found: 346.12.

10


Cap-193, Step b



[00366] A reaction mixture of ester Cap-193, Step a (0.5 g) and (+)-1,2-bis((2S,5S)-2,5-diethylphospholano)benzene(cyclooctadiene)rhodium (I) tetrafluoroborate (0.1 g) in MeOH was stirred under 55 psi of H₂ for 18 hrs. The reaction mixture was concentrated to dryness. The residue was charged to a 25 g silica gel cartridge and eluted with an 18 min gradient of 0-80% EtOAc in hexane to afford Cap-193, Step b (0.49 g) as a clear oil. LC-MS: Anal. Calcd. For [M+Na]⁺ C₁₆H₂₃NNaO₆: 348.15; found: 348.19.

20

Cap-193, Step c



[00367] A reaction mixture of Cap-193, Step b (0.16 g), dimethyl dicarbonate (0.13 g) and 10% Pd/C (0.026 g) in EtOAc was stirred under H₂ at room temperature for 2 hrs. The reaction mixture was filtered and concentrated to yield the methyl carbamate Cap-193, Step c. LC-MS: Anal. Calcd. For [M+Na]⁺ C₁₀H₁₉NNaO₆: 272.12; found: 272.07.


25

Cap-193

[00368] To a solution of ester Cap-193, Step c in THF (1 mL) and MeOH (0.25 mL) was added 1 N NaOH (1 mL). The reaction mixture was stirred at room temperature for 2 hrs. The reaction mixture was concentrated and diluted with EtOAc and 1 N HCl. The aqueous phase was extracted with EtOAc, and the combined organic phase was washed with sat. NaCl, dried over anhydrous Na₂SO₄, filtered and concentrated to yield Cap-193 (0.082 g). ¹H NMR (400 MHz, CDCl₃) 5.99 (1 H, d, *J*=8.56 Hz), 4.57 (1 H, dd, *J*=8.56, 3.27 Hz), 3.67 (3 H, s), 3.49 (2 H, d, *J*=4.28 Hz), 3.45 - 3.44 (2 H, m), 3.26 - 3.35 (6 H, m). LC-MS: Anal. Calcd. For [M+Na]⁺ C₉H₁₇NNaO₆: 258.11; found: 258.13.

10

Cap-194



[00369] Piperidine (1.0 mL, 10 mmol) was added to a solution of (S)-2-(((9H-fluoren-9-yl)methoxy)carbonylamino)-4-methoxybutanoic acid (0.355 g, 1 mmol) in DMF (3 mL), and the mixture was stirred at rt for 3 h. The volatiles were removed and the residue was partitioned between sat. NaHCO₃ (aq.) (5 mL) and EtOAc (5 mL). The aqueous layer was further washed with EtOAc and Et₂O. To the aqueous solution was added Na₂CO₃ (212 mg, 2.0 mmol) followed by methyl chloroformate (0.16 mL, 2.0 mmol) and the reaction mixture was stirred at rt for 16 h. The reaction mixture was acidified with 1 N HCl (aq.) until pH < 7 and then extracted with EtOAc (2 x 10 mL). The combined organic layers were dried (Na₂SO₄), filtered and concentrated. The residue was purified by flash silica chromatography (EtOAc/hexanes, gradient from 20% to 70%) to yield (S)-4-methoxy-2-(methoxycarbonylamino)butanoic acid (Cap-194) (91.5 mg) as viscous colorless oil. LC-MS retention time = 0.61 min; *m/z* 214 [M+Na]⁺. (Column:

25 PHENOMENEX® Luna 3.0 x 50mm S10. Solvent A = 90% Water:10% Methanol: 0.1% TFA. Solvent B = 10% Water :90% Methanol: 0.1% TFA. Flow Rate = 4 mL/min. Start % B = 0. Final % B = 100. Gradient Time = 3 min. Wavelength = 220). ¹H NMR (400 MHz, chloroform-*d*) δ ppm 7.41 (br. s., 1 H), 5.74 - 6.02 (m, 1 H), 4.32 - 4.56 (m, 1 H), 3.70 (s, 3 H), 3.54 (t, *J*=5.0 Hz, 2 H), 3.34 (s, 3 H), 1.99 - 2.23 (m, 2 H).

Example Section

[00370] Low resolution mass analysis was conducted on a Shimadzu LC system coupled with Waters MICROMASS® ZQ MS system (Cond. 1) or Waters Acquity HPLC with Waters PDA UV-Vis detection and Waters ZQ MS (Cond. 2). Retention time (R_t) were derived by employing the following conditions, and it should be noted that retention times may vary slightly between instruments:

Condition 1a

- 10 Column = PHENOMENEX® Luna 4.6X 30 mm S10
Start %B = 0
Final %B = 100
Gradient time = 3 min
Stop time = 4 min
15 Flow Rate = 4 mL/min
Wavelength = 220 nm
Solvent A = 0.1% TFA in 10% methanol/90% H₂O
Solvent B = 0.1% TFA in 90% methanol/10% H₂O

20 Condition 1b

- Column = Waters Acquity BEH C18; 1.7 μ m; 150 X 2.1 mm ID; (at 35C)
Hold 10%B = 0-1min
10-50%B = 0-25 min
50-98%B = 25-33 min
25 Hold 98%B = 32-35 min
98-10%B = 35.0-35.5 min
Hold 10%B = 35.5-40 min
Flow rate = 0.35 ml/min
Wavelength = 254 nm
30 Solvent A = 0.05% TFA in water
Solvent B = 0.05% TFA in CH₃CN

Condition 1c

- Column = Waters Acquity BEH C18; 1.7 μ m; 150 X 2.1 mm ID; (at 35 °C)
- Hold 10%B = 0-1min
- 10-60%B = 1-4 min
- 5 60-98%B = 4-21 min
- Hold 98%B = 21-21.5 min
- 98-10%B = 21.5-22 min
- Hold 10%B = 22-25 min
- Flow rate = 0.35 ml/min
- 10 Wavelength = 315 nm
- Solvent A = 0.05% TFA in water
- Solvent B = 0.05% TFA in CH₃CN

Condition 2a

- 15 Column = Waters SunFire C18, 4.6X150 mm, 3.5 μ m
- Start %B = 10
- Final %B = 50
- Gradient time = 20 min
- Stop time = varies 25 to 40 min
- 20 Flow Rate = 1 mL/min
- Wavelength = 220 & 254 nm
- Solvent A = 0.1% TFA in 5% CH₃CN/95%H₂O
- Solvent B = 0.1% TFA in 95% CH₃CN/5% H₂O

25 Condition 2b

- Column = Waters Xbridge phenyl, 4.6X150 mm, 3 μ m
- Start %B = 10
- Final %B = 50
- Gradient time = 20 min
- 30 Stop time = varies 25 to 40 min
- Flow Rate = 1 mL/min
- Wavelength = 220 & 254 nm

Solvent A = 0.1% TFA in 5% CH₃CN/95%H₂O

Solvent B = 0.1% TFA in 95% CH₃CN/5% H₂O

Condition 3

5 Column = PHENOMENEX® Luna C18 (2), 3 u, 150x4.6mm

Start %B = 0

Final %B = 100

Gradient time: = 10 min

Flow rate = 1 mL/min

10 Wavelength = 220 and 256 nm

Solvent A = H₂O/CH₃CN (95:5) + 0.05% TFA

Solvent B = H₂O/CH₃CN (5:95) + 0.05% TFA

Condition OL1

15 Column = PHENOMENEX® Luna 3.0 X 50 mm S10

Start %B = 0

Final %B = 100

Gradient time = 4 min

Stop time = 5 min

20 Flow Rate = 4 mL/min

Wavelength = 220 nm

Solvent A = 0.1% TFA in 10% methanol/90%H₂O

Solvent B = 0.1% TFA in 90% methanol/10% H₂O

25 Condition OL2

Column = PHENOMENEX® Luna 50X 2 mm 3 u

Start %B = 0

Final %B = 100

Gradient time = 4 min

30 Stop time = 5 min

Flow Rate = 0.8 mL/min

Oven Temp = 40 °C

Wavelength = 220 nm

Solvent A = 0.1% TFA in 10% Acetonitrile/90% H₂O

Solvent B = 0.1% TFA in 90% Acetonitrile/10% H₂O

5 Condition OL3

Column = Waters Acquity BEH C18; 1.7 µm; 150 X 2.1 mm ID; (at 35 °C)

Hold 10%B = 0-1 min

10-60%B = 1-4 min

60-98%B = 4-21 min

10 Hold 98%B = 21-21.5 min

98-10%B = 21.5-22 min

Hold 10%B = 22-25 min

Flow rate = 0.35 ml/min

Wavelength = 315 nm

15 Solvent A = 0.05% TFA in water

Solvent B = 0.05% TFA in CH₃CN

Condition OL4a

Column = Waters SunFire C18, 4.6X150 mm, 3.5 µm

20 Start %B = 10

Final %B = 100

Gradient time = 15 min

Stop time = 18 min

Flow Rate = 1 mL/min

25 Wavelength = 220 & 254 nm

Solvent A = 0.1% TFA in 5% CH₃CN/95% H₂O

Solvent B = 0.1% TFA in 95% CH₃CN/5% H₂O

Condition OL4b

30 Column = Waters Xbridge phenyl, 4.6X150 mm, 3 µm

Start %B = 10

Final %B = 50

Gradient time = 15 min

Stop time = 18 min

Flow Rate = 1 mL/min

Wavelength = 220 & 254 nm

- 5 Solvent A = 0.1% TFA in 5% CH₃CN/95%H₂O
Solvent B = 0.1% TFA in 95% CH₃CN/5% H₂O

Condition OL4c

Column = Waters Acquity BEH C18; 1.7 µm; 50 X 2.1 mm ID; (at 35 °C)

- 10 Hold 2%B = 0-1 min
2-98%B = 1-1.5 min
98%B = 1.5-2.2 min
Flow rate = 0.8 ml/min
Wavelength = 2 nm
15 Solvent A = 0.05% TFA in water
Solvent B = 0.05% TFA in CH₃CN

Condition OL5a

Column = PHENOMENEX® Luna 3.0 X 50 mm S10

- 20 Start %B = 0
Final %B = 100
Gradient time = 3 min
Stop time = 4 min
Flow Rate = 4 mL/min
25 Wavelength = 220 nm
Solvent A = 0.1% TFA in 10% methanol/90%H₂O
Solvent B = 0.1% TFA in 90% methanol/10% H₂O

Condition OL5b

- 30 Column = PHENOMENEX® Luna 3.0 X 50 mm S10
Start %B = 0
Final %B = 100

Gradient time = 3 min

Stop time = 4 min

Flow Rate = 4 mL/min

Wavelength = 220 nm

- 5 Solvent A = 10 mM NH₄OAc in 5% methanol/95% H₂O
Solvent B = 10 mM NH₄OAc in 95% methanol/5% H₂O

Condition-D4

Column = PHENOMENEX® Luna, 3.0 X 50 mm S10

- 10 Start %B = 0

Final %B = 100

Gradient time = 3 min

Stop time = 4 min

Flow Rate = 4 mL/min

- 15 Wavelength = 220 nm

Solvent A = 0.1% TFA in 10% methanol/90% H₂O

Solvent B = 0.1% TFA in 90% methanol/10% H₂O

Condition J4

- 20 Column = PHENOMENEX® Luna 4.6X 50 mm S10

Start %B = 0

Final %B = 100

Gradient time = 4 min

Stop time = 5 min

- 25 Flow Rate = 4 mL/min

Wavelength = 220 nm

Solvent A = 0.1% TFA in 10% methanol/90% H₂O

Solvent B = 0.1% TFA in 90% methanol/10% H₂O

- 30 Condition PY1

Column = PHENOMENEX®, 2.0 X 50 mm, 3 µm

Start %B = 0

Final %B = 100

Gradient time = 4 min

Stop time = 5 min

Flow Rate = 0.8 mL/min

5 Wavelength = 220 nm


Solvent A = 0.1 % TFA in 10% methanol/90% water

Solvent B = 0.1 % TFA in 90% methanol/10% water


Oven temp. = 40 °C

10

Examples 1 and 2




Example 1, Step a



15 [00371] Amide 1a ($\text{CH}_3\text{SO}_3\text{H}$) was prepared according to the procedure described for the synthesis of its enantiomer in patent WO 2004/052850.


Example 1, Step b



20 [00372] A 1 L round bottom flask equipped with a nitrogen inlet, overhead agitator, thermocouple and heating mantle was charged with 50 g (225 mmol) amide 1a ($\text{CH}_3\text{SO}_3\text{H}$) and 250 mL isopropanol. The resulting slurry was then charged with 252 mL of 23 wt% NaOEt in EtOH (2.68 M, 675 mmol, 3.0 equiv) and stirred at 50 °C for ca. 1h. The mixture was charged with 12.2 mL (675 mmol, 3 equiv) of water and heated to

- 60 °C. The resulting slurry was allowed to stir at 60 °C for ca. 18h. The slurry was cooled to rt and charged with 250 mL water and 98.2 g (450 mmol, 2.0 equiv) di-*t*-butyldicarbonate. Ethanol and isopropanol were removed *via* vacuum distillation and the aqueous mixture cooled to 0 °C. The mixture was neutralized with 76 mL (456 mmol)
- 5 6M aqueous HCl while maintaining an internal temperature < 5 °C. The product was extracted with 500 mL MTBE and the rich organic layer was washed with 100 mL water. The clear solution was concentrated down to 150 mL *via* vacuum distillation and the resulting slurry was charged with 600 mL heptane while maintaining an internal
- 10 temperature > 45 °C. The slurry was cooled to rt over ca. 30 min and allowed to stir at rt for ca. 2h. The product was filtered, washed with 250 mL 4:1 heptane:MTBE and dried under vacuum at 70 °C to give 40.5 g (178 mmol, 79% yield, 99.8 AP at 205 nm) of acid 1b: ¹H NMR (400 MHz, DMSO-*d*₆) δ 12.48 (s, 1H), 4.02-3.80 (m, 1H), 3.45-3.15 (m, 1H), 2.40-2.19 (m, 1H), 2.19-2.0 (m, 1H), 1.70-1.50 (m, 1H), 1.50-1.20 (m, 9H), 0.83-0.60 (m, 1H), 0.33-0.55 (m, 1H); ¹³C NMR (100 MHz, DMSO-*d*₆) δ 173.7, 173.2, 155.0,
- 15 154.3, 79.4, 60.5, 60.2, 37.6, 32.6, 31.8, 28.4, 28.2, 15.6, 15.2, 14.4; HRMS calcd for C₁₁H₁₈NO₄ (M + H; ESI⁺): 228.1236. Found: 228.1234.

Alternative synthesis of acid 1b




- 20 [00373] In a 500-mL reactor ester 1b.1 (commercially available, 17.5 g, 1.00 equiv) was dissolved in THF (87.5 mL). The resulting solution was cooled to -75 °C and 1.5M

- DIBAL-H in toluene (61.3 mL, 1.5 equiv) was charged while maintaining the temperature below -70 °C. The resulting solution was stirred at -70 °C for 1 hour. Trifluoroacetic acid (2.3 mL, 0.5 equiv) was charged over 10 minutes maintaining the internal temperature below -70 °C. Triethylamine (51.3 mL, 6 equiv) was then charged over 15 minutes maintaining the internal temperature below -70 °C. Trifluoroacetic anhydride (11.2 mL, 1.3 equiv) was charged over 10 minutes maintaining the internal temperature below -70 °C. The reaction was then allowed to warm to room temperature over 90 minutes and quenched *via* inverse addition to a solution of 20 wt % aqueous citric acid monohydrate (96.6 g, 1.5 equiv) while maintaining a temperature below 15 °C. The resulting mixture was stirred at room temperature for 2 hours then the lower aqueous layer was discarded. The product rich organic layer was washed twice with 70 mL saturated aqueous sodium bicarbonate. Solid sodium bicarbonate (1.7 g, 0.1 g/g Example 146) was charged and the solution was solvent exchanged into pure toluene under vacuum to provide 1b.2 as a solution in 2 L/kg toluene.
- 15 **[00374]** A solution of 1b.2 (16.5g theoretical from Example 151) in 33 mL toluene was polish filtered into a 250 mL reactor. Trifluorotoluene (50 mL) and chloriodomethane (43.2 g, 4.0 equiv) were then charged and the resulting solution cooled to -20 °C. 1.1M Diethylzinc in toluene (111 mL, 2.0 equiv) was charged while maintaining the internal temperature < -8 °C. The resulting solution was stirred at -15 to -20 °C for 14 hours. The reaction mixture was warmed to 0 °C then quenched via inverse addition to a solution of 20 wt % aqueous citric acid (135.7 g, 2.3 equiv). The reactor was rinsed with toluene (82 mL) and the rinse added to the quench solution. The resulting biphasic mixture was stirred for 20 minutes then the lower aqueous layer was split and discarded. The rich organic was washed twice with 60 mL 13 wt % aqueous NaCl followed by 60 mL saturated NaHCO₃. The resulting solution was solvent exchanged into pure IPA under vacuum to provide 1b.3 as a solution in 10 L/kg IPA.
- 25 **[00375]** A 250mL reactor was charged with a solution of 1b.3 (147 mL, 14.7g theoretical from ester 1b.1) in IPA. The solution was warmed to 35 °C and solid sodium hydroxide (6.2 g, 3.0 equiv) was added. The resulting mixture was stirred at 35 °C overnight. Water (44 mL) was added and the organic solvents removed under vacuum. MTBE (145 ml) was added and the pH adjusted to 3.0 with 6N aqueous HCl. The aqueous layer was split and discarded. The product rich organic was washed with 60 mL
- 30

water then azeotropically dried under vacuum *via* constant volume addition of MTBE. The solution was concentrated to 55 mL and stirred at 50 °C for 30 minutes. The solution was cooled to room temperature over 1 hour during which time a slurry formed. Heptane (90 mL) was charged over 90 min and the resulting slurry aged for 1h. The solids were collected on a medium glass frit and washed with 22.5 mL 3:1 heptane:MTBE followed by 22.5 mL heptane. The tan solid was dried in a 50 °C vacuum oven to provide 5.48 g (46%) acid 1b with 94.9 LCAP purity. The crude acid 1b was dissolved in 55 mL MTBE at 50 °C. The resulting solution was concentrated to 20 mL and cooled to room temperature over 1 hour. Heptane (33mL) was then added over 90 minutes. The resulting solids were collected on a medium glass frit, washed with heptane (15mL), and dried in a 50 °C vacuum oven to provide 4.45 g of acid 1b as a tan powder (98.8 AP, 98.8% chiral purity, 37% from ester 1b.1).


Example 1, Step c



[00376] A 250 mL round bottom flask equipped with a nitrogen inlet, overhead agitator and thermocouple was charged with 20.0 g (88.0 mmol, 2.11 equiv) acid 1b, 16.5 g (41.7 mmol, 1 equiv) 1,1'-(biphenyl-4,4'-diyl)bis(2-bromoethanone), 110 mL acetonitrile and 55 mL tetrahydrofuran. Diisopropylethylamine (15.1 mL, 86.6 mmol, 2.08 equiv) was then charged while maintaining an internal temperature < 25 °C. The mixture was allowed to stir at 20-25 °C for ca. 5h and charged with 83 mL ethyl acetate and 90 mL 13 wt% aqueous NaCl. The resulting biphasic mixture was separated and the rich organic layer was washed with an additional 90 mL 13 wt% aqueous NaCl. The rich organic layer was diluted with 20 mL tetrahydrofuran and washed with an aqueous mixture of NaHCO₃ and NaCl (45 mL 1M aqueous NaHCO₃ and 45 mL 26% aqueous NaCl). The rich organic layer was solvent exchanged into toluene *via* vacuum distillation to a target volume of ca. 160 mL. The resulting toluene solution of ketoester 1c was used as is in the next step.


[00377] [Note: for the preparation of 1,1'-(biphenyl-4,4'-diyl)bis(2-bromoethanone), see process patent application WO 2009/020825].

Example 1, Step d



[00378] The above toluene solution of ketoester 1c was charged with 64.2 g (832.9
5 mmol, 20.0 equiv) NH_4OAc and allowed to stir at 90-100 °C for ca. 18h. Cooled to 60 °C and charged 255 mL 2:1 AcOH:water. The resulting biphasic mixture was separated and the toluene layer was washed with 58 mL 1:1 AcOH:water. The rich aqueous layers were combined and residual toluene was removed *via* vacuum distillation. The aqueous solution was diluted with 60 mL methanol and heated to 50-60 °C. Charged 106 mL
10 (1060 mmol, 25.4 equiv) 10 N NaOH while maintaining an internal temperature < 60 °C. The resulting slurry was then cooled to rt. The slurry was filtered and washed with 100 mL water followed by 400 mL methanol to give 26.1 g of crude imidazole 1d. The wet, crude imidazole 1d was then charged into a 500 mL round bottom flask equipped with a nitrogen inlet, overhead agitator and thermocouple. Charged 165 mL N-methyl-2-
15 pyrrolidinone and heated to 50 °C. The resulting clear solution was charged with 30 mL Methanol and allowed to stir at 50 °C for ca. 18h. The resulting slurry was charged with an additional 130 mL methanol while maintaining an internal temperature > 45 °C. The slurry was allowed to stir at 50 °C for ca. 30 min and cooled to rt. The slurry was filtered and the solids were washed with 90 mL 1:1 methanol:N-methyl-2-pyrrolidinone
20 followed by 200 mL methanol. The solids were dried under vacuum at 70 °C to give 22.7 g (31.5 mmol, 76% yield, 95 AP at 254nm) imidazole 1d: ^1H NMR (400 MHz, DMSO-d_6) δ 11.95 (s, 2H), 7.89-7.76 (d, 4H), 7.74-7.60 (d, 4H), 7.50 (s, 2H) 4.62 (s, 2H), 3.55-3.30 (m, 2H), 2.45-2.20 (m, 4H), 1.70-1.59 (m, 2H), 1.59-0.90 (s, 18H), 0.83-0.69 (m, 2H), 0.65-0.49 (m, 2H); HRMS calcd for $\text{C}_{38}\text{H}_{45}\text{N}_6\text{O}_4$ ($\text{M} + \text{H}$; ESI^+): 649.3502. Found:
25 649.3524.


Example 1, Step e



[00379] A 250 mL round bottom flask equipped with a nitrogen inlet and overhead agitator was charged with 7.0 g (10.8 mmol) imidazole 1d, 105 mL methanol and 3.7 mL (20.8 mmol, 1.93 equiv) 5.6 M HCl in isopropanol. The resulting solution was treated with charcoal and filtered. The charcoal was washed with 140 mL methanol and combined with the filtrate. The rich organic stream was concentrated down to ca. 70 mL and charged into a round bottom flask equipped with a nitrogen inlet, overhead agitator and thermocouple. The solution was then charged with 14.75 mL (88.5 mmol, 8.2 equiv) 6M HCl and allowed to stir at 50 °C. After ca. 12h at 50 °C, the mixture was charged with 50 mL isopropanol and the resulting slurry was allowed to stir at 50 °C for 1h. The slurry was cooled to rt and aged for ca. 15h. The product was filtered and washed with 35 mL 4:1 isopropanol:methanol followed by 70 mL isopropanol. The solids were dried at 55 °C under vacuum to give 5.3 g (8.9 mmol, 83%, 99.8 AP at 254 nm) pyrrolidine 1e/4HCl: ¹H NMR (400 MHz, DMSO-d₆) δ 14.00-9.38 (bs, 8H), 8.31 (s, 2H), 8.06-7.96 (d, 4H), 7.94-7.84 (d, 4H), 5.05-4.89 (dd, 2H), 3.55-3.42 (m, 2H), 2.87-2.69 (dt, 2H), 2.64-2.53 (dd, 2H), 2.05-1.89 (m, 2H), 1.17-0.98 (m, 2H), 0.96-0.82 (dd, 2H); ¹³C NMR (100 MHz, DMSO-d₆) δ 140.0, 139.3, 133.6, 127.2, 126.6, 125.8, 116.6, 49.8, 34.3, 29.9, 15.2, 5.3; HRMS calcd for C₂₈H₂₉N₆ (M + H; ESI⁺): 449.2454. Found: 449.2470.

20

Example 1, Step f



[00380] A 250 mL jacketed reactor equipped with a nitrogen inlet, overhead agitator and thermocouple was charged with 4.24 g (24.2 mmol, 2.4 equiv) (S)-2-(methoxycarbonylamino)-3-methylbutanoic acid, 3.86 g (25.21 mmol, 2.5 equiv) 1-

hydroxybenzotriazole monohydrate, 4.55 g (23.73 mmol, 2.35 equiv) 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride and 60 mL acetonitrile. The mixture was stirred for ca. 1h and charged with 6 g (10.1 mmol) pyrrolidine 1e/4HCl. The resulting slurry was cooled to 10 °C and charged with 7.92 mL (45.41 mmol, 4.5 equiv) diisopropylethylamine. The mixture was allowed to warm to rt and stirred for ca. 19h. The resulting organic solution was washed with 36 mL 13 wt% aqueous NaCl. The rich organic was charged with 12 mL acetonitrile and washed with 36 mL of an aqueous solution containing 13 wt% aqueous NaCl and 1 M NaOH. The rich organic was then charged with 12 mL methanol and heated to 50 °C. Water (60 mL) was added over a period of 2h and the resulting slurry was cooled to rt and aged for ca. 2h. The solids were filtered, washed with 36 mL 1:1 acetonitrile:water and dried under vacuum. A 250 mL jacketed reactor equipped with a nitrogen inlet, overhead agitator and thermocouple was charged with the above solids and dissolved with 240 mL SDA3A grade ethanol. The solution was concentrated down to ca. 50 mL *via* vacuum distillation and charged with 4.24 mL (23.19 mmol, 2.30 equiv) 5.47 M HCl in isopropanol and an additional 30 mL SDA3A ethanol. The mixture was concentrated to ca. 50 mL, diluted with 40 mL SDA3A ethanol and treated with charcoal. The charcoal was filtered and washed with 90 mL SDA3A ethanol. The rich filtrate and wash were combined and concentrated down to 40 mL. Charged 16 mL ethyl acetate and heated to 40 °C. Charged 60 mg amide 1f/2HCl seed crystals and stirred at 40 °C for 1h. Charged an additional 68 mL ethyl acetate over 1.5h while maintaining an internal temperature of 40 °C. Stirred the resulting slurry at 40 °C for ca. 18h and cooled to rt. The slurry was filtered and washed with 24 mL 3:1 ethyl acetate:SDA3A ethanol and 30 mL ethyl acetate. The solids were dried under vacuum at 50 °C to give 6.83 g (8.17 mmol, 81%, 99.5 AP at 300 nm) amide 1f/2HCl: ¹H NMR (600 MHz, DMSO-d₆) δ 15.51 (s, 2H), 14.95 (s, 2H) 8.19 (s, 2H), 8.05 (d, 4H), 7.91 (d, 4H), 7.25 (d, 2H), 5.18 (t, 2H), 4.44 (t, 2H), 3.77 (s, 2H), 3.55 (s, 6H), 2.50 (m, 2H), 2.39 (m, 2H), 2.24 (m, 2H), 1.91 (m, 2H), 0.95 (m, 2H), 0.92 (d, 6H), 0.81 (d, 6H), 0.75 (s, 2H); ¹³C NMR (125 MHz, DMSO-d₆) δ 172.7, 156.9, 148.5, 139.1, 131.7, 127.1, 126.5, 125.9, 115.1, 57.8, 54.5, 51.5, 37.3, 32.9, 29.2, 19.7, 17.5, 17.5, 15.7; Calculated Elemental Analysis (corrected for 0.81% water): C = 59.86%, H = 6.30%, N = 13.29%, Cl = 8.41%; Found: C = 59.99%, H = 6.27%, N = 13.12%, Cl = 8.42%.

[00381] Preparation of Seed for amide 1f/2HCl: Amide 1f was prepared according to the basic procedure outlined above using 504 mg (0.8 mmol) pyrrolidine 1e.4HCl. After reaction completion, the rich acetonitrile solution was washed with 3 mL 13% aqueous NaCl, 2x 3 mL of an aqueous solution containing 13% NaCl and 1M NaOH, and 3 mL 13% aqueous NaCl. The rich organic was concentrated down to a residue and diluted with 10 mL acetonitrile. The hazy mixture was filtered and the clear filtrate was concentrated down to a residue. The residue was diluted with 10 mL SDA3A ethanol and charged with 2.1 mL (1.9 mmol, 2.4 equiv) 0.88 M HCl in ethanol. The mixture was concentrated down to a residue and diluted with 1.8 mL isopropanol. The resulting solution was heated to 50 °C and allowed to stir for ca. 18h. The resulting slurry was cooled to rt, filtered and washed with 2:1 acetone:ethanol to give 476 mg (0.57 mmol, 78%) amide 1f/2HCl.


[00382] [Note: (S)-2-(methoxycarbonylamino)-3-methylbutanoic acid was purchased from Flamma.]

Examples 1 and 2

[00383] A sample of amide 1f/2HCl (106.9 mg) was free-based (2 g MCX column; MeOH wash; 2 N NH₃/MeOH elution) and dried *in vacuo*. NCS (0.0195 g, 0.146 mmol) was added to a DMF (2.5 mL) solution of the resultant material and heated with an oil bath at 50 °C for 16.5 hr. Most of the volatile component was removed *in vacuo* and the residue was dissolved in MeOH and submitted to a reverse phase HPLC purification (MeOH/water/TFA; column: PHENOMENEX® Luna, 30X100 mm S10 Axia) to retrieve the TFA salts of Example 1 (white foam; 56.1 mg) and Example 2 (white foam; 22.3 mg). Example 1: ¹H NMR (DMSO-d₆, δ = 2.50 ppm, 400 MHz): 12.61 (br s, 1H), 8.13 (s, 1H), 7.93 (d, *J* = 8.5, 2H), 7.90-7.86 (m, 4H), 7.82 (d, *J* = 8.5, 2H), 7.27 (d, *J* = 8.3, 1H), 7.17 (d, *J* = 8.5, 1H), 5.02-4.94 (m, 2H), 4.44-4.38 (m, 1.80H), 4.31 (app br s, 0.2H), 3.75 (m, 1H), 3.62 (m, 1H), 3.55 (s, 3H), 3.54 (s, 3H), 2.56-2.50 ('m' partially overlapped with solvent signal, 1H), 2.41-2.23 (m, 3H), 2.17-2.09 (m, 1H), 2.07-2.00 (m, 1H), 1.97-1.91 (m, 1H), 1.88-1.81 (m, 1H), 1.01-0.81 (m, 15H), 0.71 (m, 1H). LC (Cond. 2a and 2b): >95% homogeneity index. LC-MS (Cond. 1a): *R*_t = 1.80 min. LC-MS: Anal. Calcd. for [M+H]⁺ C₄₂H₅₀ClN₈O₆: 797.35; found 797.33. Example 2: ¹H NMR (DMSO-d₆, δ = 2.50 ppm, 400 MHz): 7.86 (d, *J* = 8.6, 4H), 7.80 (d, *J* = 8.5, 4H), 7.17 (d, *J* = 8.5, 2H), 4.96


(dd, $J = 7.5, 6.0, 2\text{H}$), 4.40 (app t, $J = 7.8, 2\text{H}$), 3.61 (m, 2H), 3.54 (s, 6H), 2.33-2.23 (m, 4H), 2.09-1.98 (m, 2H), 1.90-1.82 (m, 2H), 1.01-0.82 (overlapped 'm' and two 'd' at 0.96 ppm and 0.87 ppm with $J = 6.8$ and $J = 6.5$, respectively; 14H); 0.71 (m, 1.7H), 0.62 (m, 0.3H). LC (Cond. 2a and 2b): >95% homogeneity index. LC-MS (Cond. 1a): $R_t = 2.79$ min. LC-MS: Anal. Calcd. for $[\text{M}+\text{H}]^+ \text{C}_{42}\text{H}_{49}\text{Cl}_2\text{N}_8\text{O}_6$: 831.32; found 831.26.

Example 3



10

Example 3, Step a




[00384] For the synthesis of amide 3a/2HCl, see the process patent application WO 2009/020825.

15

Example 3

[00385] To a solution of amide 3a (32 mg, 0.043 mmol) in acetic acid (2 mL) was added 2 eqv. of 1M bromine in acetic acid and the reaction was stirred for 30 min. An additional 1 eqv. bromine was added (solids formed), and the reaction was neutralized with 5% NaHCO_3 and extracted with EtOAc (2x10 mL). The combined extracts were washed with sat'd $\text{Na}_2\text{S}_2\text{O}_3$ soln and brine and concentrated. The residue was dried under high vacuum to afford Example 3 (41 mg). LC-MS (Cond.-J4): $R_t = 3.54$ min. LC-MS Anal. Calcd. for $[\text{M}+\text{H}]^+ \text{C}_{40}\text{H}_{49}\text{Br}_2\text{N}_8\text{O}_6$: 897.22; found 897.28.

Examples 4 and 5




[00386] Diisopropylethyl amine (100 ul, 74.2 mg, 0.574 mmol) was added to a mixture of amide 3a (197mg, 0.243 mmol) and N-chlorosuccinamide (39 mg, 0.292 mmol) in acetonitrile (2 mL) and stirred for overnight. An additional portion of N-

chlorosuccinamide (66mg, 0.494 mmol) was then added to the reaction mixture and stirred until the consumption of the starting material was confirmed by HPLC analysis. The crude reaction mixture was then purified by reverse phase HPLC (Solvent A = H₂O/CH₃CN (95:5) + 0.05% TFA; Solvent B = H₂O/CH₃CN (5:95) + 0.05% TFA; Column: Luna C18, 5u, 100 × 21.5 mm; Flow rate: 20 mL/min) to afford the TFA salts of

Example 4 (yellowish amorphous powder, 60 mg) and Example 3 (yellowish amorphous powder, 115 mg). Example 5: LC-MS (Cond. 3): R_t = 6.13 min. ¹H NMR (DMSO-d₆, TMS, 600 MHz): 8.16 (s, 1H), 7.95-7.91 (m, 7H), 7.85 (d, *J* = 8.8, 2H), 7.35 (d, *J* = 8.4, 1H), 7.32 (d, *J* = 8.4, 1H), 5.19 (t, *J* = 7.3, 1H), 5.05 (m, 1H), 4.15 (t, *J* = 7.7, 1H). 4.08 (t, *J* = 8.1, 1H), 3.88-3.81 (m, 4H), 3.56 (s, 6H), 2.43 (m, 1H), 2.21-2.11 (m, 4H), 2.08-2.05 (m, 2H), 1.97 (m, 3H), 0.91 (d, *J* = 6.5, 3H), 0.87 (d, *J* = 7.0, 3H), 0.85 (d, *J* = 7.0, 3H), 0.80 (d, *J* = 6.6, 3H). HRMS: Calcd. for [M+H]⁺ C₄₀H₅₀ClN₈O₆: 773.3542; found 773.3530. Example 4: LC-MS (Cond. 3): R_t = 8.61 min. ¹H NMR (DMSO-d₆, TMS, 600 MHz): 7.89 (d, *J* = 8.3, 4H), 7.83 (d, *J* = 8.3, 4H), 7.33 (d, *J* = 8.4, 2H), 5.05 (m, 2H), 4.09 (t, *J* = 8.3, 2H), 3.84-3.81 (m, 4H), 3.56 (s, 6H), 2.24-2.15 (m, 4H), 2.00-1.95 (m, 6H), 0.91 (d, *J* = 6.7, 6H), 0.87 (d, *J* = 6.6, 6H). HRMS: Calcd. for [M+H]⁺ C₄₀H₄₉Cl₂N₈O₆: 807.3152; found 807.3138.


Examples 6 and 7



Example-6: R = H

Example-7: R = Cl

Example 6, Step a




- 5 **[00387]** To a mixture of pyrrolidine 1e/4HCl (0.104 g, 0.174 mmol), HATU (0.133 g, 0.349 mmol) and Cap-193 (0.082 g, 0.349 mmol) in DMF was added DIEA (0.183 mL). The reaction mixture was stirred at room temperature for 3 hrs. The reaction mixture was purified by preparatory HPLC (PHENOMENEX® Axia 5 u column, 35 min gradient from 0-90%B. A = H₂O/CH₃CN/10 mM NH₄OAc, 95:5. B = CH₃CN/H₂O/10 mM
- 10 NH₄OAc 95:5) to afford Example 6, Step a (0.064 g). LC-MS: Anal. Calcd. For [M+H]⁺ C₄₆H₅₉N₈O₁₀: 883.43; found: 883.26. ¹H NMR (400 MHz, MeOD) ppm 7.90 (2 H, s), 7.78 - 7.86 (8 H, m), 5.13 (2 H, dd, *J*=9.2, 6.9 Hz), 4.96 (2 H, d, *J*=5.0 Hz), 3.73 - 3.79 (2 H, m), 3.63 (6 H, s), 3.39 - 3.49 (6 H, m), 3.35 (1 H, d, *J*=4.8 Hz), 3.32 - 3.34 (1 H, m), 3.30 (6 H, s), 3.15 (6 H, s), 2.66 (2 H, dd, *J*=13.5, 9.4 Hz), 2.37 - 2.51 (4 H, m), 2.02 -
- 15 2.09 (2 H, m), 1.03 - 1.14 (2 H, m), 0.81 - 0.93 (2 H, m).

Examples 6 and 7

- [00388]** To a solution of Example 6, Step a (0.057 g) in DMF was added NCS (10.34 mg, 0.077 mmol). The reaction mixture was heated at 60 °C for 3 hrs. The reaction
- 20 mixture was purified by preparatory HPLC (PHENOMENEX® Axia 5 u 30 x 100 mm column, 20 min gradient from 0-100%B. A = H₂O/CH₃OH/TFA 90:10:0.1. B = CH₃OH/H₂O/TFA 90:10:0.1) to afford Example 6 (0.022 g) and Example 7 (0.017 g). Example 6: LC (Cond.-J5): R_t = 3.46 min. LC-MS: Anal. Calcd. For [M+H]⁺


$C_{46}H_{58}ClN_8O_{10}$: 917.39; found: 917.12. 1H NMR (400 MHz, MeOD) ppm 7.87 - 7.91 (1 H, m), 7.77 - 7.86 (8 H, m), 5.14 (1 H, dd, $J=9.2, 6.9$ Hz), 5.05 (1 H, dd, $J=8.1, 6.3$ Hz), 4.96 (2 H, t, $J=5.7$ Hz), 3.73 - 3.79 (1 H, m), 3.67 - 3.72 (1 H, m), 3.57 - 3.66 (6 H, m), 3.32 - 3.51 (8 H, m), 3.29 (6 H, d, $J=1.8$ Hz), 3.19 (3 H, s), 3.12 - 3.16 (3 H, m), 2.60 - 2.72 (1 H, m), 2.32 - 2.54 (5 H, m), 2.02 (2 H, d, $J=6.3$ Hz), 1.00 - 1.19 (2 H, m), 0.88 (1 H, br. s.), 0.82 (1 H, br. s.). Example 7: LC (Cond. PY1): $R_t = 4.083$ min. LC-MS: Anal. Calcd. For $[M+H]^+$ $C_{46}H_{56}Cl_2N_8O_{10}$: 951.35; found: 951.09. 1H NMR (400 MHz, MeOD) ppm 7.74 - 7.84 (8 H, m), 5.05 (2 H, t, $J=7.2$ Hz), 4.95 (2 H, d, $J=5.5$ Hz), 3.66 - 3.72 (2 H, m), 3.63 (6 H, s), 3.37 - 3.53 (8 H, m), 3.30 (6 H, s), 3.19 (6 H, s), 2.32 - 2.51 (6 H, m), 1.92 - 2.06 (2 H, m), 1.08 (2 H, dt, $J=8.6, 5.8$ Hz), 0.81 (2 H, d, $J=1.8$ Hz).

Example V1



15

Example V1, Step a




[00389] A solution of carbamate 1d (0.80 g, 1.23 mmol) and NCS (0.214 g, 1.603 mmol) in DMF (12 mL) was heated at 50 °C for 17h. After it was allowed to cool to ambient temperature, volatile components were removed *in vacuo*. The residue was dissolved in MeOH (36 mL) and the two products were separated by prep-HPLC (Column: XTERRA®, 30 X 100 mm, S5; Start %B = 50, Final %B = 100; Gradient time = 10 min; Stop time = 12 min; Flow rate = 30 ml/min; Wavelength = 220; Solvent A = 10% MeOH – 90% H₂O – 0.1% TFA; Solvent B = 90% MeOH – 10% H₂O – 0.1% TFA). Each of the two fractions were neutralized with an excess solution of 2N NH₃/MeOH and

concentrated *in vacuo* to remove most of the methanol, and the residue was partitioned between 20% MeOH/CHCl₃ and water. The organic layer was dried (MgSO₄), filtered, and concentrated *in vacuo* to afford chloride V1a-1 (light yellow foam; 288.4 mg) and dichloride V1a-2 (light yellow solid; 400.4 mg). Chloride V1a-1: ¹H NMR (DMSO-d₆, δ = 2.5 ppm, 400 MHz): 12.62-12.60 (m, 1H), 12.23 (s, 0.2H), 11.92 (s, 0.8H), 7.85-7.32 (m, 9H), 4.68-4.42 (m, 2H), 3.47-3.38 (m, 2H), 2.40-2.35 (m, 2H), 2.26-2.20 (m, 2H), 1.71-1.58 (m, 2H), 1.55-0.96 (app br s, 18H), 0.82-0.69 (m, 2H), 0.64-0.51 (m, 2H). LC (V-Cond. 1): R_t = 2.44 min. LC-MS: Anal. Calcd. for [M+H]⁺ C₃₈H₄₄ClN₆O₄: 683.31; found: 683.31. Dichloride V1a-2: ¹H NMR (DMSO-d₆, δ = 2.5 ppm, 400 MHz): 12.62 (s, 2H), 7.85 (d, *J* = 8.8, 4H), 7.82 (d, *J* = 8.9, 4H), 4.61-4.42 (m, 2H), 3.53-3.38 (m, 2H), 2.41-2.35 (m, 2H), 2.27-2.20 (m, 2H), 1.70-1.59 (m, 2H), 1.53-0.94 (app br s, 18H), 0.80-0.70 (m, 2H), 0.64-0.53 (m, 2H). LC (Cond. 1a): R_t = 2.89 min. LC-MS: Anal. Calcd. For [M+H]⁺ C₃₈H₄₃Cl₂N₆O₄: 717.27; found: 717.31.

15

Example V1, Step b




[00390] To a suspension of chloride V1a-1 (0.2741 g, 0.401 mmol) in MeOH (1 mL) was added 4M HCl in dioxane (4 mL). The mixture was stirred at room temperature for 7h, and then the volatile component was removed *in vacuo* to afford the HCl salt of V1b as a tan solid (231.7 mg). The product was used without further purification. ¹H NMR (DMSO-d₆, δ = 2.5 ppm, 400 MHz): 13.51 (br s, 1H), 10.53-10.30 (m, 2H), 9.87-9.72 (m, 1H), 8.02-7.85 (m, 9H), 4.72-4.68 (m, 1H), 4.65-4.53 (m, 1H), 3.42-3.30 (m, 2H), 2.68-2.37 ('m' partially overlapped with solvent signal, 4H), 1.97-1.87 (m, 2H), 1.16-1.07 (m, 2H), 0.89-0.80 (m, 2H). LC (Cond. 1a): R_t = 1.66 min. LC-MS: Anal. Calcd. for [M+H]⁺ C₂₈H₂₈ClN₆: 483.21; found: 483.20. [Note: the exact HCl content of the product was not determined].


Example V1


[00391] To a solution of chloride V1b/4HCl (0.045 g, 0.072 mmol), (S)-2-(methoxycarbonylamino)butanoic acid (0.025 g, 0.157 mmol), and N,N-diisopropylethylamine (0.075 mL, 0.429 mmol) in DMF (1.5 mL) was added HATU (0.057 g, 0.150 mmol). The reaction mixture was stirred at ~25 °C for 1h. It was diluted with MeOH (2.5 mL) and the product was purified by prep-HPLC (Column: XTERRA®, 30 X 100 mm, S5; Start %B = 30, Final %B = 90; Gradient time = 10 min; Stop time = 12 min; Flow rate = 30 ml/min; Wavelength = 220; Solvent A = 10% MeOH – 90% H₂O – 0.1% TFA; Solvent B = 90% MeOH – 10% H₂O – 0.1% TFA) to afford the TFA salts of Example V1 as a light yellow solid (39.4 mg). ¹H NMR (DMSO-d₆, δ = 2.5 ppm, 400 MHz): 14.80-14.36 (br m, 1.4H), 12.58 (app br s, 0.6H), 8.14 (s, 1H), 7.94 (d, *J* = 8.5, 2H), 7.89 (d, *J* = 8.3, 2H), 7.87 (d, *J* = 8.4, 2H), 7.82 (d, *J* = 8.6, 2H), 7.38 (d, *J* = 7.8, 0.9H), 7.28 (d, *J* = 7.8, 0.9H), 7.07 (app br s, 0.1H), 6.96 (app br s, 0.1H), 5.02 (dd, *J* = 8.8, 6.8, 1H), 4.96 (dd, *J* = 7.0, 6.0, 1H), 4.50-4.39 (m, 2H), 3.74-3.67 (m, 1H), 3.59-3.48 (m, 7H), 2.60-2.47 ('m' partially overlapped with solvent signal, 1H), 2.42-2.22 (m, 3H), 2.01-1.65 (m, 4H), 1.65-1.52 (m, 2H), 1.06-0.84 (m, 8H), 0.83-0.77 (m, 1H), 0.74-0.66 (m, 1H). LC (Cond. 2a and 2b): >95% homogeneity index. LC (Cond. 1a): R_t = 1.66 min. LC-MS: Anal. Calcd. for [M+H]⁺ C₄₀H₄₆ClN₈O₆: 769.32; found: 769.35.

Example V2


[00392] Example V2 (TFA salt) was prepared from pyrrolidine V1b/4HCl and (S)-2-(methoxycarbonylamino)-2-(tetrahydro-2H-pyran-4-yl)acetic acid according to the procedure described for the synthesis of Example V1.



Example		R _t (LC-Cond.); LC (Cond. 2a and 2b): % homogeneity index; MS data
---------	---	--


V2		2.06 min (Cond. 1a); >95%; LC-MS: Anal. Calcd. for $[M+H]^+$ $C_{46}H_{54}ClN_8O_8$: 881.38; found: 881.36.
----	---	--

Example V3



5

Example V3, Step a



[00393] Intermediate V3a was prepared as HCl salt from carbamate V1a-2 according to the procedure described for the preparation of pyrrolidine V1b. 1H NMR (DMSO- d_6 , δ = 2.5 ppm, 400 MHz): 10.62-10.48 (m, 2H), 9.92-9.75 (m, 2H), 7.92 (d, J = 8.8, 4H), 7.89 (d, J = 9.1, 4H), 4.68-4.57 (m, 2H), 3.40-3.31 (m, 2H), 2.54-2.50 ('m' partially overlapped with solvent signal, 2H), 2.46-2.39 (m, 2H), 1.91-1.85 (m, 2H), 1.16-1.07 (m, 2H), 0.85-0.80 (m, 2H). LC (Cond. 1a): R_t = 2.01 min. LC-MS: Anal. Calcd. for $[M+H]^+$ $C_{28}H_{27}Cl_2N_6$: 517.17; found: 517.06. [Note: the exact HCl salt content of the product was not determined].

15

Example V3

[00394] Example V3 (TFA salt) was prepared from the HCl salt of pyrrolidine V3a according to the procedure described for the synthesis of Example V1. 1H NMR (DMSO- d_6 , δ = 2.5 ppm, 400 MHz): 12.59 (very br s, ~2H), 7.86 (d, J = 8.5, 4H), 7.80 (d, J = 8.3, 4H), 7.46 (app br s, 0.13H), 7.37 (d, J = 7.6, 1.73H), 6.97 (app br s, 0.14H), 4.97-4.94 (m, 2H), 4.52-4.35 (m, 2H), 3.63-3.45 (m, 8H), 2.36-2.23 (m, 4H), 1.96-1.83 (m, 2H),


20

1.82-1.67 (m, 2H), 1.64-1.49 (m, 2H), 1.13-0.82 (m, 8H), 0.78-0.61 (m, 2H). LC (Cond. 2a and 2b): >95% homogeneity index. LC (Cond. 1a): $R_t = 2.66$ min. LC-MS: Anal. Calcd. for $[M+H]^+$ $C_{40}H_{45}Cl_2N_8O_6$: 803.28; found: 803.28.

5

Examples V4 to V6

[00395] Examples V4 to V6 (TFA salt) were prepared from the HCl salt of pyrrolidine V3a and the appropriate acids according to the procedure described for the synthesis of Example V1.




10

Example		R_t (LC-Cond.); LC (Cond. 2a and 2b): % homogeneity index; MS data
V4		2.64 min (Cond. 1a); >95%; LC-MS: Anal. Calcd. for $[M+H]^+$ $C_{46}H_{53}Cl_2N_8O_8$: 915.34; found: 915.70. 1H NMR (DMSO- d_6 , $\delta = 2.5$ ppm, 400 MHz): 12.63 (very br s, ~2H), 7.87 (d, $J = 8.6$, 4H), 7.78 (d, $J = 8.5$, 4H), 7.42 (app br s, 0.1H), 7.26 (d, $J = 8.6$, 1.75H), 6.97 (app br s, 0.14H), 4.95 (dd, $J = 8.4$, 5.1, 2H), 4.48 (app t, $J = 8.1$, 2H), 3.85 (m, 4H), 3.73-3.64 (m, 2H), 3.54 (s, 6H), 3.32-3.18 (m, 4H), 2.37-2.19 (m, 4H), 2.05-1.91 (m, 2H), 1.91-1.78 (m, 2H), 1.71-1.58 (m, 2H), 1.56-1.39 (m, 4H), 1.39-1.25 (m, 2H), 1.05-0.93 (m, 2H), 0.77-0.66 (m, 2H)


Example		R _t (LC-Cond.); LC (Cond. 2a and 2b): % homogeneity index; MS data
V5		2.59 min (Cond. 1a); >95%; LC-MS: Anal. Calcd. for [M+H] ⁺ C ₃₈ H ₄₁ Cl ₂ N ₈ O ₆ : 775.25; found: 775.29. ¹ H NMR (DMSO-d ₆ , δ = 2.5 ppm, 400 MHz): 12.56 (very br s, ~2H), 7.86 (d, <i>J</i> = 8.5, 4H), 7.80 (d, <i>J</i> = 8.5, 4H), 7.37 (d, <i>J</i> = 7.6, 1.70H), 7.02 (app br s, 0.3H), 4.95 (app t, <i>J</i> = 6.8, 2H), 4.63-4.54 (m, 2H), 3.62-3.34 (overlapped 'm' and 's'; s is at 3.52; 8H), 2.31-2.28 (m, 4H), 1.91-1.84 (m, 2H), 1.28 (s, 3H), 1.27 (d, <i>J</i> = 7, 6H), 1.02-0.97 (m, 2H), 0.76-0.61 (m, 2H).
V5.1		2.65 min (Cond. 1a); >95%; LC-MS: Anal. Calcd. for [M+H] ⁺ C ₃₈ H ₄₁ Cl ₂ N ₈ O ₆ : 775.25; found: 775.35.
V5.2		2.72 min (Cond. 1a); >95%; LC-MS: Anal. Calcd. for [M+H] ⁺ C ₄₀ H ₄₅ Cl ₂ N ₈ O ₆ : 803.28; found: 803.38.
V5.3		2.50 min (Cond. 1a); >95%; LC-MS: Anal. Calcd. for [M+H] ⁺ C ₃₈ H ₄₁ Cl ₂ N ₈ O ₄ : 743.26; found: 743.36.
V6		2.75 min (Cond. 1a); >95%; LC-MS: Anal. Calcd. for [M+H] ⁺ C ₄₂ H ₄₅ Cl ₂ N ₈ O ₆ : 827.28; found: 827.33.

Examples V7 and V8



V7 (R₁ = H, R₂ = Cl)
V8 (R₁ = Cl, R₂ = H)

Example V7, Step a




[00396] To a solution of pyrrolidine 1e/4HCl (0.350 g, 0.589 mmol), (S)-2-(methoxycarbonylamino)-3-methylbutanoic acid (0.103 g, 0.589 mmol), (S)-2-(methoxycarbonylamino)-2-(tetrahydro-2H-pyran-4-yl)acetic acid (0.128 g, 0.589 mmol) and N,N-diisopropylethylamine (0.720 mL, 4.12 mmol) in DMF (3 mL) was added HATU (0.493 g, 1.295 mmol). The reaction mixture was stirred at ~25 °C for 4 h, and then was diluted with MeOH and purified by prep-HPLC (Column: XTERRA®, 30 X 100 mm, S5; Start %B = 30, Final %B = 75; Gradient time = 15 min; Stop time = 15 min; Flow rate = 30 ml/min; Wavelength = 220; Solvent A = 10% MeOH – 90% H₂O – 0.1% TFA; Solvent B = 90% MeOH – 10% H₂O – 0.1% TFA) to isolate product V7a out of the three possible products. Product V7a was dissolved in MeOH and free-based (6g MCX cartridge; MeOH wash; 2N NH₃/MeOH elution) and concentrated *in vacuo* to afford a tan solid (103.2 mg). LC (Cond. 1a): R_t = 1.63 min. LC-MS: Anal. Calcd. for [M+H]⁺ C₄₄H₅₃N₈O₇: 805.40; found: 805.49.

Examples V7 and V8

[00397] A solution of product V7a (0.103 g, 0.128 mmol) and NCS (0.022 g, 0.167 mmol) in DMF (1.5 mL) was heated at 50 °C for 24h. After it was allowed to cool to ambient temperature, the reaction mixture was diluted with MeOH and purified by prep-HPLC (Column: XTERRA®, 30 X 100 mm, S5; Start %B = 40, Final %B = 100; Gradient time = 15 min; Stop time = 17 min; Flow rate = 30 ml/min; Wavelength = 220; Solvent A = 10% MeOH – 90% H₂O – 0.1% TFA; Solvent B = 90% MeOH – 10% H₂O – 0.1% TFA) to isolate a mixture of regioisomers V7 and V8. The mixture was dissolved in MeOH and submitted to a different prep-HPLC purification condition (Column: Waters SunFire, 30 X 100 mm, S5; Start %B = 10, Final %B = 50; Gradient time = 20 min; Stop time = 20 min; Flow rate = 30 ml/min; Wavelength = 220; Solvent A = 10% Acetonitrile – 90% H₂O – 0.1% TFA; Solvent B = 90% Acetonitrile – 10% H₂O – 0.1%


- TFA) to separate the two regioisomers as TFA salts. Example V7 (off-white solid, 17.3 mg): ^1H NMR (DMSO- d_6 , δ = 2.5 ppm, 400 MHz): 14.52 (app br s, too broad to integrate), 12.64 (app br s, too broad to integrate), 8.14 (s, 1H), 7.95-7.79 (m, 8H), 7.26 (two overlapped 'd', J = \sim 8.6, 1.70H), 7.10 (app br s, 0.08H), 6.98 (app br s, 0.22H),
- 5 5.01-4.97 (m, 1H), 4.96-4.92 (m, 1H), 4.50-4.46 (m, 1H), 4.43-4.40 (m, 1H), 3.90-3.81 (m, 2H), 3.79-3.72 (m, 1H), 3.71-3.64 (m, 1H), 3.54 (app s, 6H), 3.31-3.18 (m, 2H), 2.65-2.50 ('m' partially overlapped with solvent, 1H), 2.45-2.19 (m, 3H), 2.18-2.08 (m, 1H), 2.04-1.89 (m, 2H), 1.89-1.79 (m, 1H), 1.69-1.59 (m, 1H), 1.55-1.41 (m, 2H), 1.38-1.23 (m, 1H), 1.04-0.77 (m, 9H), 0.74-0.67 (m, 1H). LC (Cond. 2a and 2b): >95%
- 10 homogeneity index. LC (Cond. 1a): R_t = 2.15 min. LC-MS: Anal. Calcd. for $[\text{M}+\text{H}]^+$ $\text{C}_{44}\text{H}_{51}\text{Cl}_1\text{N}_8\text{O}_7$: 839.36; found: 839.35. Example V8 (off-white solid, 16.8 mg): ^1H NMR (DMSO- d_6 , δ = 2.5 ppm, 400 MHz): 14.53 (br s, too broad to integrate), 12.61 (br s, too broad to integrate), 8.13 (s, 1H), 7.94 (d, J = 8.3, 2H), 7.90 (d, J = 8.6, 2H), 7.85 (d, J = 8.5, 2H), 7.82 (d, J = 8.6, 2H), 7.36 (d, J = 8.8, \sim 1H), 7.17 (d, J = 8.5, \sim 1H),
- 15 5.00-4.94 (m, 2H), 4.49 (app t, J = 7.7, 1H), 4.40 (app t, J = 7.9, 1H), 3.90-3.78 (m, 3H), 3.65-3.59 (m, 1H), 3.55 (s, 3H), 3.54 (s, 3H), 3.35-3.19 (m, 2H), 2.64-2.50 ('m' partially overlapped with solvent signal, 1H), 2.41-2.34 (m, 1H), 2.33-2.22 (m, 2H), 2.12-1.99 (m, 2H), 1.99-1.90 (m, 1H), 1.89-1.81 (m, 1H), 1.52-1.27 (m, 4H), 1.07-0.84 (m, 8H), 0.84-0.77 (m, 1H), 0.75-0.65 (m, 1H). LC (Cond. 2a and 2b): >95% homogeneity index. LC
- 20 (Cond. 1a): R_t = 2.15 min. LC-MS: Anal. Calcd. for $[\text{M}+\text{H}]^+$ $\text{C}_{44}\text{H}_{51}\text{Cl}_1\text{N}_8\text{O}_7$: 839.36; found: 839.38.

Example V9



25


Example V9, Step a



[00398] Boc₂O (0.734 g, 3.36 mmol) followed DMAP (0.021 g, 0.168 mmol) were added to a mixture of pyrrolidine 1e/4HCl (2.00 g, 3.36 mmol) and Et₃N (2.3 ml, 16.82 mmol) in DMF (60 ml), and stirred at ambient condition for 4.5 h. The volatile component was removed *in vacuo*, and the residue was partitioned between 1.0 N NaOH (20 mL) and 20% MeOH/CHCl₃ (50 mL). The aqueous phase was washed with 20% MeOH/CHCl₃ (50 mL, 2x). The combined organic phase was dried (MgSO₄) and concentrated *in vacuo*. A silica gel mesh was prepared from the resultant crude material and submitted to a BIOTAGE® purification (160 g silica gel) where the column was first eluted with EtOAc until all of the higher R_f spot (*i.e.*, bis-Boc derivative 1d; 0.28 g) came out, and then the column was eluted with 5-10% MeOH/CH₂Cl₂ over 2.5 L to elute residual bis-Boc x (followed by mono-Boc V9a (0.81 g; ~containing 1.1 mol equiv of Et₃N). LC-MS: Anal. Calcd. for [M+H]⁺ C₃₃H₃₇N₆O₂: 549.30; found 549.45.

15


Example V9, Step b



[00399] To a solution of carbamate V9a (1 g, 1.823 mmol), (S)-2-(methoxycarbonylamino)-2-(tetrahydro-2H-pyran-4-yl)acetic acid (0.400 g, 1.841 mmol), and N,N-diisopropylethylamine (0.65 mL, 3.72 mmol) in DMF (8 mL) was added HATU (0.770 g, 2.026 mmol), and the reaction mixture was stirred at ~25 °C for 3 h. The volatile component was removed *in vacuo*, and the residue was taken up in 20% MeOH/CHCl₃ (250 mL), washed with water (3 x 40 mL), dried over MgSO₄, filtered, and concentrated *in vacuo*. The residue was taken up in CHCl₃ (4 mL) and loaded onto a Thomson's silica gel cartridge and eluted with 10% MeOH/EtOAc to afford product V9b as a tan foam (1.15 g). LC (Cond. 1a): R_t = 1.70 min; LC-MS: Anal. Calcd. for [M+H]⁺ C₄₂H₅₀N₇O₆: 748.38; found: 748.40.


25

Example V9, Step c



[00400] A solution of V9b (0.9387 g, 1.255 mmol) and NCS (0.335 g, 2.51 mmol) in
 5 DMF (13 mL) was heated at 50 °C for 24.5 h. The volatile component was removed *in vacuo* and the residue was taken up in CH₂Cl₂ (200 mL), washed with water (3 x 50 mL), followed by brine (50 mL), dried over MgSO₄, filtered, and concentrated *in vacuo*. The residue was taken up in CH₂Cl₂ (6 mL) and submitted to a silica gel purification (100% ethyl acetate) to afford compound V9c as a yellow solid (720.7 mg). LC (Cond. 1a): R_t =
 10 3.19 min. LC-MS: Anal. Calcd. for [M+H]⁺ C₄₂H₄₈Cl₂N₇O₆: 816.30; found: 816.35.

Example V9, Step d



[00401] Pyrroldine V9d was prepared as HCl salt from carbamate V9c according to the
 15 procedure described for the synthesis of V1b. ¹H NMR (DMSO-d₆, δ = 2.5 ppm, 400 MHz): 10.49-10.38 (m, 1H), 9.84-9.72 (m, 1H), 7.90-7.81 (m, 8H), 7.27 (d, J = 8.6, 0.89H), 6.99 (app br s, 0.11H), 4.98-4.95 (m, 1H), 4.65-4.55 (m, 1H), 4.50-4.46 (m, 1H), 3.90-3.80 (m, 2H), 3.76-3.64 (m, 1H), 3.54 (s, 3H), 3.41-3.32 (m, 1H), 3.32-3.19 (m, 2H), 2.54-2.40 ('m' partially overlapped with solvent signal, 2H), 2.36-2.24 (m, 2H), 2.06-1.95
 20 (m, 1H), 1.94-1.78 (m, 2H), 1.69-1.57 (m, 1H), 1.55-1.40 (m, 2H), 1.39-1.27 (m, 1H), 1.17-1.07 (m, 1H), 1.04-0.92 (m, 1H), 0.88-0.78 (m, 1H), 0.76-0.66 (m, 1H). LC (Cond. 1a): R_t = 2.66 min; >95% homogeneity index. LC-MS: Anal. Calcd. for [M+H]⁺ C₃₇H₄₀Cl₂N₇O₄: 716.25; found: 716.28.

25

Example V9


[00402] Example V9 (TFA salt) was prepared from the HCl salt of intermediate V9d and (S)-2-(methoxycarbonylamino)butanoic acid according to the procedure described for the synthesis of Example V1. ¹H NMR (DMSO-d₆, δ = 2.5 ppm, 400 MHz): 12.59 (br s, ~2H), 7.87-7.77 (m, 8H), 7.29/7.26 (two partially overlapped 'd', J = 8.6/7.8, 1.82H), 6.97 (app br s, 0.18H), 4.97-4.93 (m, 2H), 4.50-4.42 (m, 2H), 3.90-3.81 (m, 2H), 3.72-3.65 (m, 1H), 3.60-3.46 (overlapping m and two s, s are at 3.54 and 3.53, 7H), 3.32-3.18 (m, 2H), 2.38-2.18 (m, 4H), 2.05-1.23 (collection of 'm', 9H), 1.09-0.85 (m, 5H), 0.76-0.59 (m, 2H). LC (Cond. 2a and 2b): >95% homogeneity index. LC (Cond. 1a): R_t = 2.66 min. LC-MS: Anal. Calcd. for [M+H]⁺ C₄₃H₄₉Cl₂N₈O₇: 859.31; found: 859.43.




10

Examples V10 to V16

[00403] Examples V10 to V16 (TFA salt) were prepared from the HCl salt of pyrrolidine V9d and appropriate acids according to the procedure described for the preparation of Example V1. Examples V13 to V16 required an additional reverse phase HPLC purification which employed the following condition: Column: Waters SunFire, 30 X 100 mm, S5; Start %B = 10, Final %B = 50; Gradient time = 20 min; Stop time = 20 min; Flow rate = 30 ml/min; Wavelength = 220; Solvent A = 10% Acetonitrile – 90% H₂O – 0.1% TFA; Solvent B = 90% Acetonitrile – 10% H₂O – 0.1% TFA.


20




Example		R _t (LC-Cond.); LC (Cond. 2a and 2b): % homogeneity index; MS data
V10		2.59 min (Cond. 1a); >95%; LC-MS: Anal. Calcd. for [M+H] ⁺ C ₄₁ H ₄₅ Cl ₂ N ₈ O ₇ : 831.28; found: 831.31.
V11		2.54 min (Cond. 1a); >95%; LC-MS: Anal. Calcd. for [M+H] ⁺ C ₄₂ H ₄₇ Cl ₂ N ₈ O ₇ : 845.29; found: 845.17.

Example		R _t (LC-Cond.); LC (Cond. 2a and 2b): % homogeneity index; MS data
V12		2.69 min (Cond. 1a); >95%; LC-MS: Anal. Calcd. for [M+H] ⁺ C ₄₄ H ₅₁ Cl ₂ N ₈ O ₇ : 873.33; found: 873.27.
V13		2.73 min (Cond. 1a); >95%; LC-MS: Anal. Calcd. for [M+H] ⁺ C ₄₄ H ₄₉ Cl ₂ N ₈ O ₇ : 871.31; found: 871.41.
V14		2.83 min (Cond. 1a); >95%; LC-MS: Anal. Calcd. for [M+H] ⁺ C ₄₅ H ₅₁ Cl ₂ N ₈ O ₇ : 885.33; found: 885.45.
V15		2.70 min (Cond. 1a); >95%; LC-MS: Anal. Calcd. for [M+H] ⁺ C ₄₄ H ₅₁ Cl ₂ N ₈ O ₈ : 889.32; found: 889.44.
V16		2.52 min (Cond. 1a); >95%; LC-MS: Anal. Calcd. for [M+H] ⁺ C ₄₂ H ₄₆ Cl ₂ N ₇ O ₆ : 814.29; found: 814.18.

Examples GW1-1 to GW1-3






[00404] For the preparation of carbamate GW1a, see U.S. Patent Application 2009/0068140.

5


Example GW1, Step b



[00405] To a solution of carbamate GQ1a (0.830 g, 1.271 mmol) in DMF (9 mL) was added NCS (0.221 g, 1.653 mmol). The reaction mixture was heated at 50 °C for 16 hr. More NCS (0.1 g, 0.75 mmol) was added and heating was continued for another 4 hr. It was allowed to cool to ambient temperature and partitioned between DCM and water (20 mL each). The aqueous layer was extracted with DCM (20 mL), and the combined organic phase was dried (Na₂SO₄) and concentrated *in vacuo*. The resulting crude material was dissolved in MeOH and submitted to a reverse phase HPLC purification (MeOH/water/TFA; column: PHENOMENEX® Luna, 30X100 mm S10 Axia) to retrieve the TFA salts of GW1b as light yellow foam (0.25g). ¹H NMR (MeOD, δ = 3.30 ppm, 400 MHz): 7.82 (app s, 8H), 4.95-4.85 ('m' partially overlapped with HOD signal, 2H), 4.04 (m, 2H), 2.36-2.14 (m, 6H), 1.77 (m, 2H), 1.50-1.20 (overlap of 'br s' and 'd', 24H). LC (Cond. Ia): RT = 3.0 min. LC-MS: Anal. Calcd. for [M+H]⁺ C₃₈H₄₇Cl₂N₆O₄: 721.30; found 721.32.

20

Example GW1, Step c




[00406] 4N HCl in dioxane (10 mL) was added to carbamate GW1b (0.29 g, 0.306 mmol) and the reaction mixture was stirred at room temperature for 4 hrs. Solvent was removed and dried under vacuum overnight to give the HCl salt of product GW1c as brown solid (0.21g). ¹H NMR (MeOD, δ = 3.30 ppm, 400 MHz): 7.85 (d, J = 8.6, 4H), 7.80 (d, J = 8.8, 4H), 4.95-4.85 (m, 2H), 3.89-3.80 (m, 2H), 2.58-2.31 (m, 6H), 2.01-1.91 (m, 2H), 1.50 (d, 6.5 Hz, 6H). LC (Cond. 1a): RT = 3.0 min. LC-MS: Anal. Calcd. for [M+H]⁺ C₂₈H₃₁Cl₂N₆: 521.20; found 521.20.


Examples GW1-1 to GW1-3

[00407] To a solution of the HCl salt of GW1c (150 mg, 0.225 mmol), (S)-2-(methoxycarbonylamino)-2-(tetrahydro-2H-pyran-4-yl)acetic acid (57.6 mg, 0.265 mmol) and (S)-2-(methoxycarbonylamino)-3-methylbutanoic acid (48.0 mg, 0.274 mmol) in DMF (10 mL) were added DIEA (0.236 mL, 1.349 mmol) and HATU (176 mg, 0.463 mmol), and the mixture was stirred at room temperature for 1 hr. Most of the volatile component was removed *in vacuo* and the residue was dissolved in MeOH and submitted to a reverse phase HPLC purification (MeOH/water/TFA; column: PHENOMENEX® Luna, 30X100 mm S10 Axia) to isolate the three products as TFA salts. Example GW1-1 (light yellow foam, 25 mg) was contaminated with unidentified impurity. ¹H NMR (MeOD, δ = 3.30 ppm, 400 MHz): 8.05-7.93 (m, 1.54H), 7.87-7.56 (m, 6.46H), 5.42 (br d, J = 6.6, 0.68H), 4.97 (m, 1.32H), 4.76-4.69 (m, 1.2H), 4.31-4.14 (m, 2.8H), 3.98-3.84 (m, 4H), 3.68/3.67/3.63 (three 's', 6H), 3.40-3.21 ('m' partially overlapped with solvent signal, 4H), 2.72 (m, 0.7H), 2.45-1.14 (overlapped many 'm' and two 'd', 'd' at 1.51 and 1.18 with J = 6.5 and 6.3, respectively; 23.3H). LC (Cond. 2a and 2b): >91.8% homogeneity index. LC (Cond. 1c): RT = 6.22 min. LC-MS: Anal. Calcd. for [M+H]⁺ C₄₆H₅₇Cl₂N₈O₈: 919.37; found 919.9. Example GW1-2 (light yellow foam, 39 mg); LC (Cond. 2a and 2b): >95% homogeneity index. LC (Cond. 1a): RT = 2.97 min. LC-MS: Anal. Calcd. for [M+H]⁺ C₄₄H₅₅Cl₂N₈O₇: 877.36; found 877.35. Example GW1-3 (light yellow foam, 22 mg), LC (Cond. 2a and 2b): >95% homogeneity index. LC (Cond. 1a): RT = 3.07 min. LC-MS: Anal. Calcd. for [M+H]⁺ C₄₂H₅₃Cl₂N₈O₆: 835.35; found 835.34.

Example GW2




Example GW2, Step a



- 5 **[00408]** To a solution of carbamate GW1a (0.830 g, 1.271 mmol) in DMF (9 mL) was added NCS (0.221 g, 1.653 mmol). The reaction mixture was heated at 50 °C for 16 hr. Additional NCS (0.1g, 0.75 mmol) was added and heating was continued at 50 °C for an additional 4 hr. The mixture was partitioned between DCM and water (20 mL each). The aqueous phase was extracted with DCM (20 mL), and the combined phase was dried
- 10 with Na₂SO₄ and concentrated *in vacuo*. The resulting crude product was dissolved in MeOH and submitted to a reverse phase HPLC purification (MeOH/water/TFA; column: PHENOMENEX® Luna, 30X100 mm S10 Axia) to retrieve carbamate GW2a as light yellow foam (0.03g). ¹H NMR (MeOD, δ = 3.30 ppm, 400 MHz): 7.76-7.64 (m, 8H), 7.34 (s, 1H), 5.10-4.65 (overlapping with HOD signal, m, 2H), 3.99 (m, 2H), 2.31-2.01
- 15 (m, 6H), 1.78-1.76 (m, 2H), 1.55-1.15 (app br m, 24H). LC (Cond. 1a): RT = 2.55 min. LC-MS: Anal. Calcd. for [M+H]⁺ C₃₈H₄₈ClN₆O₄: 687.34; found 687.39.

Example GW2, Step b



- 20 **[00409]** 4N HCl in dioxane (10 mL) was added to carbamate GW2a (0.19 g, 0.208 mmol) and it was stirred at room temperature for 4 hr. Solvent was removed and it was dried under vacuum overnight to afford the HCl salt of pyrrolidine GW2b as brown solid


(0.16g). ^1H NMR (MeOD, δ = 3.30 ppm, 400 MHz): 8.11 (s, 1H), 7.95 (d, J = 8.5, 2H), 7.87 (appears 'd', 4H), 7.82 (d, J = 8.8, 2H), 5.22 (m, 1H), 4.92 – 4.83 (over lapping with HOD signal, m, 1H), 3.99 (m, 1H), 3.84 (m, 1H), 2.78-2.62 (m, 2H), 2.57-2.31 (m, 4H), 2.12-1.91 (m, 2H), 1.55 (d, J = 6.8, 3H), 1.50 (d, J = 6.8, 3H). LC (Cond. 1a): RT = 1.74 min. LC-MS: Anal. Calcd. for $[\text{M}+\text{H}]^+ \text{C}_{28}\text{H}_{32}\text{ClN}_6$: 487.24; found 487.21.



Example GW2

[00410] To a solution of the HCl salt of pyrrolidine GW2b (80 mg, 0.126 mmol) and (S)-2-(methoxycarbonylamino)-3-methylbutanoic acid (48.7 mg, 0.278 mmol) in DMF (3 mL), DIEA (0.132 mL, 0.758 mmol) and HATU (99 mg, 0.260 mmol) were added and the mixture was stirred at room temperature for 1 hr. Most of the volatile component was removed *in vacuo* and the residue was dissolved in MeOH and submitted to a reverse phase HPLC purification (MeOH/water/TFA; column: PHENOMENEX® Luna, 30X100 mm S10 Axia) to retrieve the TFA salts of GW2 (light yellow foam; 50 mg). ^1H NMR (MeOD, δ = 3.30 ppm, 400 MHz): 8.00-7.75 (m, 9H), 5.71 (app br m, 0.2H), 5.40 (d, J = 6.3, 0.45 H), 5.15-5.11 (m, 0.87H), 5.00-4.96 (m, 0.92H), 4.81-4.68 (m, 1.56H), 4.26-4.03 (m, 2H), 3.71/3.70/3.64/3.63 (four 's', 6H), 2.77-1.87 (m, 9.24H), 1.65-1.48 (overlapped 'm' and 'd', 4.66 H), 1.28 (d, J = 0.62H), 1.17 (d, J = 6.3H, 1.23H), 1.06-0.82 (m, 12.15H). LC (Cond. 2a and 2b): >95% homogeneity index. LC-MS (Cond. 1a): RT = 2.46 min. LC-MS Anal. Calcd. for $[\text{M}+\text{H}]^+ \text{C}_{42}\text{H}_{54}\text{ClN}_8\text{O}_5$: 801.39; found 801.41.


Example GW2-1

[00411] Example GW2-1 (TFA salt) was prepared from pyrrolidine GW2b and (S)-2-(methoxycarbonylamino)-2-(tetrahydro-2H-pyran-4-yl)acetic acid according to the procedure described for the preparation of Example GW2.




Example		RT (LC-Cond.); % homogeneity index; MS data
GW2-1		2.23 min (Cond. 1a); >98%; LC-MS: Anal. Calcd. for $[M+H]^+$ C ₄₆ H ₅₈ ClN ₈ O ₈ : 885.41; found 885.37

Example GW3




5


Example GW3, Step a



GW3a-1



GW3a-2




GW3a-3

[00412] The above three esters were prepared from (*S*)-1-*tert*-butyl 2-methyl 5-oxopyrrolidine-1,2-dicarboxylate according to the procedure described in *Tetrahedron Letters*, 3203-3205 (2003).

10


Example GW3, Step b



[00413] Borane-methyl sulfide complex (5.44 mL, 10.88 mmol) was added to a solution of ester GW3a-2 (1.4 g, 5.44 mmol) in THF (25 mL), and the reaction mixture


was heated at 40 °C for 7 hr. The volatile component was removed *in vacuo* and the residue was partitioned between EtOAc and water (50 mL each). The aqueous layer was extracted with EtOAc (30 mL), and the combined organic phase was dried with Na₂SO₄, and concentrated *in vacuo*. The resultant colorless oil was purified with a flash chromatography (0-50% EtOAc/Hexane) to afford ester GW3b as a colorless oil (0.77 g).
¹H NMR (CDCl₃, δ = 7.24 ppm, 400 MHz): 4.29-4.18 (m, 1H), 3.78-3.66 (m, 4H), 2.99 (app t, *J* = 10.1, 1H), 2.43-2.97 (m, 1H), 2.43-2.37 (m, 1H), 2.30-2.18 (m, 1H), 1.60-1.52 (m, 1H), 1.47/1.42 (two 's', 9H), 1.08-1.05 (m, 3H).

Example GW3, Step c



To a solution of ester GW3b (1.69 g, 6.95 mmol) in ethanol (10 mL) was added solution of LiOH (0.250 g, 10.42 mmol) in water (5.00 mL), and the reaction mixture was stirred at room temperature for 5 hr. The organic solvent was evaporated *in vacuo* and the residue was diluted with water (10 mL) and washed with ether (10 mL). It was chilled in ice-water bath, and acidified to a pH range of ~2 with 1N HCl. It was then extracted with EtOAc (20 mL, 3x). The organic layer was dried with Na₂SO₄ and concentrated *in vacuo* to afford acid GW3c as a colorless oil, which became a white solid upon extended exposure to high vacuum (1.38g).
¹H NMR (CDCl₃, δ = 7.24 ppm, 400 MHz): 4.39-4.22 (m, 1H), 3.80-3.69 (m, 0.91H), 3.59-3.35 (m, 0.18H), 3.03-2.89 (m, 0.91H), 2.51-2.22 (m, 2H), 1.98-1.91 (m, 0.71H), 1.68-1.60 (0.29H), 1.50/1.44 (two 's', 9H), 1.09 (app m, 3H).

Example GW3, Step d




To a suspension of GW3c (1.83 g, 7.98 mmol) and 1,1'-(biphenyl-4,4'-diyl)bis(2-bromoethanone) (1.581 g, 3.99 mmol) in CH₃CN (30 mL) was added DIEA

(1.436 mL, 8.22 mmol), and the mixture was stirred at room temperature for 4 hr. Solvent was evaporated *in vacuo* and the residue was partitioned between EtOAc and water (50 mL each). The organic layer was washed with sat. NaHCO₃ (20 mL), dried with Na₂SO₄, and concentrated *in vacuo* to afford diester GW3d as light yellow solid


5 (2.67g), which was used as is for the next step.

Example GW3, Step e



[00416] To a solution of diketoester GW3d (2.67 g, 3.85 mmol) in xylenes (30 mL)
 10 was added ammonium acetate (2.97 g, 38.5 mmol), and the mixture was heated at 140 °C for 6 hr in a sealed tube. The volatile component was removed *in vacuo* and the residue was partitioned between DCM (50 mL) and water (50 mL). The organic layer was washed with sat. NaHCO₃ (20 mL), dried with Na₂SO₄, and concentrated *in vacuo*. The resulting crude material was purified with a flash chromatography (50-100%
 15 EtOAc/Hexane, 100% EtOAc-10% MeOH/EtOAc)) to afford imidazole GW3e as orange solid (1.3g). ¹H NMR (DMSO-d₆, δ = 2.5 ppm, 400 MHz): 12.21-11.87 (three br s, 2H), 7.82-7.51 (m, 9.2H), 7.30 (m, 0.6H), 6.88-6.81 (m, 0.2H), 4.9-4.67 (m, 1.81H), 3.98-3.93 (m, 0.19H), 3.77-3.54 (m, 2H), 3.07-2.76 (m, 2H), 2.43-2.04 (m, 4H), 1.80-1.56 (m, 2H), 1.41-1.33 (m, 8H), 1.10-1.04 (m, 16H). LC-MS (Cond. 1a): RT = 2.09 min. LC-MS
 20 Anal. Calcd. for [M+H]⁺ C₃₈H₄₉N₆O₄: 653.38; found 653.51.


Example GW3, Step f



[00417] 4N HCl in dioxane (12.10 mL, 48.4 mmol) was added to carbamate GW3e (1.3 g, 1.99 mmol) and the mixture was stirred at room temperature for 5 hr. The volatile component was removed *in vacuo*, and the product was dried under vacuum overnight to afford HCl salt of pyrrolidine GW3f as a yellow solid (1.14g). ¹H NMR (DMSO-d₆, δ = 2.5 ppm, 400 MHz): 10.32 (app br s, 2H), 9.79 (app br s, 2H), 8.13 (s, 2H), 8.02 (d, J=8.3, 4H), 7.99 (d, J=8.3, 4H), 5.01 (br m, 2H), 3.47 (br m, 2H), 3.05 (br m, 2H), 2.62 (m, 2H), 2.45 (m, 2H), 2.21 (m, 2H), 1.13 (d, J = 6.3, 6H). LC-MS (Cond. 1a): RT = 2.09 min. LC-MS Anal. Calcd. for [M+H]⁺ C₂₈H₃₃N₆: 453.28; found 453.17.

10

Example GW3, Step g



[00418] To a suspension of the HCl salt of pyrrolidine GW3f (1.14 g, 1.905 mmol), (S)-2-(methoxycarbonylamino)-2-(tetrahydro-2H-pyran-4-yl)acetic acid (0.488 g, 2.248 mmol) and (S)-2-(methoxycarbonylamino)-3-methylbutanoic acid (0.407 g, 2.324 mmol) in DMF (15 mL) was added DIEA (1.996 mL, 11.43 mmol). Total dissolution was effected with the help of sonication, and then HATU (1.478 g, 3.89 mmol) was added, and the reaction mixture was stirred at room temperature for 2 hr. Most of the volatile component was removed *in vacuo* and the residue was dissolved in MeOH and submitted to a reverse phase HPLC purification (MeOH/water/TFA; column: PHENOMENEX® Luna, 30X100 mm S10 Axia) to afford TFA salts of three products: GW3g-1 (0.28g), GW3g-2 (0.64g) and GW3g-3 (0.36g) as light yellow foam. Product GW3g-3: ¹H NMR (DMSO, δ = 2.5 ppm, 400 MHz): 14.74 (app br s, not integratable), 8.16 (s, 2H), 7.97 (d, J=8.6, 4H), 7.88 (d, J=8.3, 4H), 7.29 (d, J=8.0, < 2H), 5.23 (br m, 0.1H), 5.07 (dd, J = 10.6, 7, 1.9H), 4.18 (m, 2H), 4.09 (m, 2H), 3.53 (s, 6H), 3.39 (m, 2H), 2.56-2.35 ('m'

partially overlapped with solvent signal, 2H), 1.95 (m, 2H), 1.82 (m, 2H), 1.13 (d, $J=6.3$, 6H), 0.79 (d, $J=6.5$, 6H), 0.76 (d, $J=6.5$, 6H) [Note: the last three integrations include minor rotamers with signals in the 1.1-0.85 ppm region that were not peak-picked]. LC-MS (Cond. 1a): RT = 1.79 min. LC-MS Anal. Calcd. for $[M+H]^+$ C₄₂H₅₅N₈O₆: 767.42; found 767.30. Product GW3g-1, LC-MS (Cond. 1a): RT = 1.62 min. LC-MS Anal. Calcd. for $[M+H]^+$ C₄₆H₅₉N₈O₈: 851.45; found 851.33. Product GW3g-2, LC-MS (Cond. 1a): RT = 1.70 min. LC-MS Anal. Calcd. for $[M+H]^+$ C₄₄H₅₇N₈O₇: 809.44; found 809.30.


10

Example GW3

[00419] To a solution of GW3g-3/2TFA (160mg, 0.174 mmol) in DMF (2 mL) was added NCS (30.6 mg, 0.229 mmol). The mixture was heated to 50 °C for 5.5 hr. The residue was dissolved in MeOH and submitted to a reverse phase HPLC purification (MeOH/water/TFA; column: PHENOMENEX® Luna, 30X100 mm S10 Axia) to retrieve the TFA salts of GW3 as light yellow foam (40 mg). [Note: for analogous cases the chlorination was conducted on a starting material that was in a free base form.] ¹H NMR (MeOD, δ = 3.30 ppm, 400 MHz): 7.88 (s, 1H), 7.84-7.76 (m, 8H), 5.19 (dd, J = 11.1, 7.1, 1H), 4.98 (dd, J = 10.4, 7.3, 1H), 4.32 (m, 1H), 4.24-4.17 (m, 3H), 3.63 (s, 3H), 3.62 (s, 3H), 3.40-3.33 (m, 2H), 2.65 (m, 1H), 2.50 (m, 2H), 2.37 (m, 1H), 1.99 (m, 2H), 1.82 (m, 2H), 1.22 (d, $J=6.3$, 2.85H), 1.18 (d, $J=6.3$, 2.85H), 1.09 (m, 0.3H), 0.92-0.85 (m, 12H). LC (Cond. 2a and 2b): >95% homogeneity index. LC-MS (Cond. 1a): RT = 2.32 min. LC-MS Anal. Calcd. for $[M+H]^+$ C₄₂H₅₄ClN₈O₆: 801.39; found 801.25.

20

Example GW4




25

[00420] Example GW4 (TFA salt) was prepared from GW3g-1 according to the procedure described for the preparation of Example GW3. LC (Cond. 2a and 2b): >95%

homogeneity index. LC-MS (Cond. 1a): RT = 2.17 min (Cond. 1a). LC-MS: Anal. Calcd. for $[M+H]^+$ C₄₆H₅₈ClN₈O₈: 885.41; found 885.26.

Examples GW5 to GW7




5

[00421] To a solution of GW3g-2 (565mg, 0.588 mmol) in DMF (2 mL) was added NCS (0.021 g, 0.161 mmol), and the reaction mixture was heated at 50 °C for 17 hr. Most of the volatile component was removed *in vacuo* and the residue was dissolved in MeOH and submitted to a reverse phase HPLC purification (MeOH/water/TFA; column: PHENOMENEX® Luna, 30X100 mm S10 Axia) to afford two fractions (one is GW5 and the second one is mixture of GW6 and GW7). The fractions were dried by *in vacuo* and retrieved as light yellow solid. Both fractions were dissolved separately in methanol and submitted to a reverse phase HPLC purification (ACN/TFA/Water, Water-SunFire 30X100 mm S5) to retrieve the TFA salts of Example GW5, GW6 and GW7. Example GW5: ¹H NMR (MeOD, δ = 3.30 ppm, 400 MHz): 7.74 (m, 8H), 5.02 (m, 2H), 4.29-4.18 (m, 4H), 3.90 (m, 2H), 3.63 (s, 6H), 3.41-3.33 (m, 4H), 2.52 (m, 2H), 2.39 (br m, 2H), 2.01-1.79 (m, 4H), 1.58-1.27 (m, 4H), 1.18 (d, *J*=6.5, 5.63H), 1.09 (m, 0.37H), 0.93 (d, *J*=6.8, 3H), 0.88 (d, *J*=6.5, 3H). LC (Cond. 2a and 2b): >95% homogeneity index. LC-MS (Cond. 1a): RT = 2.81 min. LC-MS Anal. Calcd. for [M+H]⁺ C₄₄H₅₅Cl₂N₈O₇: 877.36; found 877.30. Example GW6: ¹H NMR (MeOD, δ = 3.30 ppm, 400 MHz): 7.86 (s, 1H), 7.82-7.73 (m, 8H), 5.19 (dd, *J* = 11, 7, 1H), 4.98 (dd, *J* = 10.4, 7.3, 1H), 4.33-4.19 (m, 4H), 3.90 (m, 2H), 3.67 (m, 0.35H), 3.63/3.62 (two overlapped 's', 5.65H), 3.41-3.31 (m, 4H), 2.65 (m, 1H), 2.49 (m, 2H), 2.38 (m, 1H), 2.00-1.79 (m, 4H), 1.57-1.27 (m, 4H), 1.22/1.18 (two overlapping 'd', *J* = 6.5/6.3, respectively, 5.69H), 1.09 (m, 0.31H), 0.90/0.86 (two overlapping 'd', *J* = 6.8/6.8, 6H). LC (Cond. 2a and 2b): >95% homogeneity index. LC-MS (Cond. 1a): RT = 2.25 min. LC-MS Anal. Calcd. for [M+H]⁺ C₄₄H₅₆ClN₈O₇: 843.40; found 843.24. Example GW7: ¹H NMR (MeOD, δ =

3.30 ppm, 400 MHz): 7.87-7.75 (m, 9H), 5.19 (dd, $J=11$, 7.0, 1H), 4.99 (dd, $J=10.3$, 7.3, 1H), 4.38-4.17 (m, 4H), 3.90 (m, 2H), 3.64/3.62 (two 's', 6H), 3.41-3.26 ('m' partially overlapped with solvent signal, 4H), 2.64 (m, 1H), 2.50 (m, 2H), 2.37 (m, 1H), 2.03-1.79 (m, 4H), 1.60-1.08 (m, 10H), 0.92 (d, $J=6.8$, 3H), 0.86 (d, $J=6.8$, 3H). LC
 5 (Cond. 2a and 2b): >95% homogeneity index. LC-MS (Cond. 1a): RT = 2.24 min. LC-MS Anal. Calcd. for $[M+H]^+$ C₄₄H₅₆ClN₈O₇: 843.40; found 843.24.

Example GW8




10

Example GW8, Step a



[00422] To a solution of GW3e (0.360 g, 0.551 mmol) in DMF (9 mL) was added NCS (0.096 g, 0.717 mmol), and the reaction mixture was heated at 50 °C for 15 hr. The
 15 residue was diluted with MeOH and submitted to a reverse phase HPLC purification (MeOH/water/TFA; column: PHENOMENEX® Luna, 30X100 mm S10 Axia) to retrieve the TFA salt of carbamate GW8a as light yellow foam (20 mg). TFA salts of GW3e (60 mg) and the mono-chloro analog (70 mg) were also isolated. Carbamate GW8a: ¹H NMR (MeOD, δ = 3.30 ppm, 400 MHz): 7.83 (app s, 8H), 4.95-4.79 ('m' partially overlapped with HOD signal, 2H), 3.85-3.75 (m, 2H), 3.14 (m, 2H), 2.58-2.48 (m, 2H), 2.44-2.30 (m, 2H), 1.80-1.71 (m, 2H), 1.43 (s, 6H), 1.23 (s, 12H), 1.14/1.12 (two overlapping 's', 6H).
 20 LC-MS (Cond. 1a): RT = 2.98 min. LC-MS Anal. Calcd. for $[M+H]^+$ C₃₈H₄₇ClN₆O₄: 721.30; found 721.39.

Example GW8, Step b




[00423] 4N HCl in dioxane (3 mL) was added to the TFA salt of GW8a (55 mg, 0.063 mmol), and the mixture was stirred at room temperature for 5 hr. Solvent was removed *in vacuo* and the residue was dried under vacuum overnight to afford the HCl salt of pyrrolidine GW8b as a brown solid (50mg). LC-MS (Cond. 1a): RT = 2.20 min. LC-MS Anal. Calcd. for $[M+H]^+$ C₂₈H₃₁Cl₂N₆: 521.20; found 521.20.


Example GW8

[00424] To a suspension of the HCl salt of GW8b (50 mg, 0.075 mmol) in DMF (3 mL) was added (S)-2-(methoxycarbonylamino)-3-methylbutanoic acid (28.9 mg, 0.165 mmol), DIEA (0.079 mL, 0.450 mmol), and HATU (58.7 mg, 0.154 mmol), and the mixture was stirred at room temperature for 2 hr. Most of the volatile component was removed *in vacuo* and the residue was dissolved in MeOH and submitted to a reverse phase HPLC purification (MeOH/water/TFA; column: PHENOMENEX® Luna, 30X100 mm S10 Axia) to retrieve the TFA salts of GW8 (light yellow foam; 40 mg). ¹H NMR (MeOD, δ = 3.30 ppm, 400 MHz): 7.80-7.75 (m, 8H), 5.01 (dd, *J* = 10.4, 7.4, 2H), 4.26-4.17 (m, 4H), 3.63 (s, 6H), 3.38-3.33 (m, 2H), 2.52-2.49 (m, 2H), 2.45-2.33 (m, 2H), 2.03-1.95 (m, 2H), 1.87-1.79 (m, 2H), 1.19 (d, *J*=6.3, 6H), 0.92 (d, *J*=6.8, 6H), 0.87 (d, *J*=6.8, 6H). LC (Cond. 2a and 2b): >95% homogeneity index. LC-MS (Cond. 1a): RT = 2.89 min. LC-MS Anal. Calcd. for $[M+H]^+$ C₄₂H₅₃Cl₂N₈O₆: 835.35; found 835.41.

Examples GW9 and GW10





Example GW9, Step a



- 5 [00425] To a solution of GW3a-2 (1.31 g, 5.09 mmol) in THF (25 mL) was added methylmagnesium bromide (2.037 mL, 6.11 mmol) dropwise at -40 °C (dry ice/acetone bath). It was stirred at the same temperature for 2 hr, and then the bath was removed and stirring was continued for an additional 1 hr. Acetic acid (1 mL), water (10 mL) and ether (50 mL) were added, the mixture was shaken and the organic layer was separated.
- 10 The organic layer was washed with water, dried with Na₂SO₄, and concentrated *in vacuo*. The resultant crude material was purified with a flash chromatography (5-95% EtOAc/Hexane) to afford ester GW9a as colorless oil (0.6g). ¹H NMR (CDCl₃, δ = 7.24 ppm, 400 MHz): 4.97 (br d, *J* = 8.3, 1H), 4.29 (m, 1H), 3.74/3.73 (overlapping 's', 3H), 2.66 (m, 1H), 2.27-2.22 (m, 0.21H), 2.17 (s, 2.58H), 2.10-2.02 (m, 1.22H), 1.80-1.1.74
- 15 (m, 1H), 1.43/1.40 (overlapping 's', 9H), 1.16 (d, *J*=7.1, 2.6H), 1.05(d, *J*=6.8, 0.4H).

Example GW9, Step b




- [00426] To a solution of ketone GW9a (0.6 g, 2.195 mmol) in DCM (15 mL) was
- 20 added TFA (0.846 mL, 10.98 mmol), and the mixture was stirred at room temperature for 7 hr. The volatile component was removed *in vacuo*, and dried under vacuum overnight to afford a light yellow oil (0.63g).

[00427] A solution of the above crude product (0.5g, 1.8 mmol) in methanol (20 mL) was added to a 500 mL Parr shaker bottle containing Pd/C (0.025 g, 0.234 mmol). After evacuation and refilling with nitrogen was conducted (3x), the mixture was shaken under 60 psi for 24 hr. The reaction mixture was filtered through a filter paper and the volatile component was removed *in vacuo* to afford a light yellow oil (0.41g).

[00428] To a solution of the above crude product (0.48g, 1.77 mmol) in CH₂Cl₂ (7 mL) was added DMAP (10.81 mg, 0.088 mmol), triethylamine (0.740 mL, 5.31 mmol). Di-*tert*-butyl dicarbonate (0.386g, 1.77 mmol) was added in portions over 15 min and the mixture was stirred at room temperature for 18 hr. After the volatile component was removed *in vacuo*, the crude material was purified with a flash chromatography (0-22% EtOAc/Hexane) to afford two major products. The first elute was GW9b-1 (0.54g). ¹H NMR (CDCl₃, δ = 7.24 ppm, 400 MHz) 4.41/4.30 (br m, 1H), 3.74 (s, 3H), 3.48/3.33 (br m, 1H), 2.37-2.22 (m, 0.2H), 2.10-2.04 (m, 0.9H), 1.98-1.19 (m, 0.9H), 1.79 (m, 1H), 1.46-1.32 (m, 12H), 1.05/1.01/0.97 (three overlapping 'd', *J* = 6.8/6.6/6.3, respectively, 3H). The second elute was GW9b-2 contaminated with unidentified impurity (0.48 g). Clean fractions of GW9b-2 was used to acquire the following spectral data: ¹H NMR (CDCl₃, δ = 7.24 ppm, 400 MHz) 4.23-4.12 (two overlapping 'dd', 1H), 4.00-3.85 (two overlapping 'm', 1H), 3.72/3.71 (overlapping 's', 3H), 2.34-2.20 (m, 2H), 1.69-1.55 (1H), 1.45/1.39 (overlapping 's', 9H), 1.13-1.01 (m, 3.13H), 0.97 (d, *J* = 6.3, 2.87H).


Example GW9, Step c



[00429] To a solution of GW9b-1 (3.73 g, 14.50 mmol) in ethanol (40 mL) was added solution of LiOH (0.417 g, 17.39 mmol) in water (20.00 mL), and the mixture was stirred at room temperature for 6 hr. More LiOH (0.1g, 4.3 mmol) was added, and stirring was continued for an additional 2 hr. Most of the organic component was evaporated, and the remaining portion was washed with ether (20 mL). The aqueous layer was chilled with ice-water bath, acidified with 1N HCl to a pH of 2-3, extracted with EtOAc (50 mL, 4x), dried with Na₂SO₄, and concentrated to afford a colorless oil, which became a white solid upon exposure to high vacuum (3.43g). The solid was dissolved in a minimum amount of

EtOAc with the help of heating gun that brought it to a refluxing condition. After cooling down to room temperature, 5 drops of hexane were added and allowed to stand at ambient temperature overnight to afford acid GW9c as white needles, which was filtered and washed with hexane, dried under vacuum (2.02g). ¹H NMR (CDCl₃, δ = 7.24 ppm, 400 MHz) 4.40 (app dd, *J*=8.3, 2.8, 1H), 3.30 (br m, 1H), 2.50 (br m, 1H), 2.01-1.93 (m, 1H), 1.68-1.59 (m, 1H), 1.52 (s, 9H), 1.29 (d, *J*=6.0, 3H), 1.07 (d, *J*=6.8, 3H).

Example GW9, Step d



[00430] Compound GW9d (TFA salt; light yellow foam) was prepared from acid GW9c according to the procedure described for the preparation of precursors GW3g from acid GW3c with the exception that (S)-2-(methoxycarbonylamino)-3-methylbutanoic acid was employed for the final step. ¹H NMR (MeOD, δ = 3.30 ppm, 400 MHz): 7.97-7.96 (m, 2H), 7.88-7.82 (m, 8H), 5.70 (br d, *J* = 6.7, 0.6H), 5.29 (dd, *J* = 10.3, 1.2H), 5.19-5.14 (m, 0.2H), ~4.90 (overlapped with HOD signal, 1H), 4.38 (m, 1H), 4.12-4.06 (m, 2H), 3.71/3.67/3.64 (overlapping 's', 6H), 2.73-2.68 (m, 0.6H), 2.53-2.45 (m, 1.4H), 2.35-2.18 (m, 2.5H), 2.07-1.82 (m, 3.5H), 1.53 (d, *J* = 6.8, 3.6H), 1.40 (d, *J* = 7.1, 0.5H), 1.30 (d, *J* = 7.1, 1.9H), 1.17-0.83 (collection of overlapping 'd', 18H). LC (Cond. 2a and 2b): >95% homogeneity index. LC-MS (Cond. 1a): RT = 2.07 min. LC-MS Anal. Calcd. for [M+H]⁺ C₄₄H₅₉N₈O₆: 795.46; found 795.44.


Examples GW9 and GW10

[00431] To a solution of GW9d (TFA salt; 310 mg, 0.328 mmol) in DMF (6 mL) was added NCS (62.5 mg, 0.468 mmol), and the mixture was heated at 50 °C for 9 hr. Most of the volatile component was removed *in vacuo* and the residue was dissolved in MeOH and submitted to a reverse phase HPLC purification (MeOH/water/TFA; column: PHENOMENEX® Luna, 30X100 mm S10 Axia) to separate the mono- and di-chloro

products. The mono-chlorinated product was further purified with a different reverse phase HPLC condition (ACN/TFA/Water, Water-SunFire 30X100 mm S5). Example GW9 (light yellow foam; 36 mg) and Example GW10 (light yellow foam; 60 mg) were retrieved as TFA salts. Example GW9: ^1H NMR (DMSO, δ = 2.50 ppm, 400 MHz):

- 5 12.71-12.64 (br m, 1H), 8.10 (br s, 1H), 7.95-7.83 (m, 8H), 7.64-7.58 (m, 1.4H), 7.51 (d, J =8.6, 0.6H), 5.56 (m, 0.22H), 5.33 (m, 0.33H), 5.16 (m, 0.78H), 5.04 (m, 0.67H), 4.27-4.11 (m, 2H), 3.93-3.86 ('m' overlapped with H_2O signal, 2H), 3.56-3.53 (m, 6H), 2.62-1.64 (collection of 'm', 8H), 1.50-0.75 (collection of 'd', 24H). LC (Cond. 2a and 2b): >95% homogeneity index. LC-MS (Cond. 1a): RT = 2.66 min. LC-MS Anal. Calcd. for
- 10 $[\text{M}+\text{H}]^+ \text{C}_{44}\text{H}_{58}\text{ClN}_8\text{O}_6$: 829.42; found 829.44. Example GW10: ^1H NMR (DMSO, δ = 2.50 ppm, 400 MHz): 12.7 (br m, 2H), 8.00-7.80 (m, 8H), 7.63 (m, 0.76H), 7.51 (d, J =8.6, 1.09H), 7.08 (m, 0.09H), 6.57 (m, 0.05H), 5.33 (d, J =6.3, 0.74 H), 5.05 (app t, 1.26H), 4.25-4.12 (m, 2H), 3.90-3.85 ('m' overlapped with H_2O signal, 2H), 3.58-3.49 (m, 6H), 2.59-2.55 (m, 0.5H), 2.33-2.13 (m, 2.7H), 1.98-1.88 (m, 4.05H), 1.71-1.63 (m,
- 15 0.75H), 1.43 (d, J =6.6, 3.7H), 1.18 (d, J =6.3, 2.05H), 1.05-0.76 (collection of 'd', 18.25H). LC (Cond. 2a and 2b): >95% homogeneity index. LC-MS (Cond. 1a): RT = 3.29 min. LC-MS Anal. Calcd. for $[\text{M}+\text{H}]^+ \text{C}_{44}\text{H}_{57}\text{Cl}_2\text{N}_8\text{O}_6$: 863.38; found 863.38.

Examples GW11 and GW12




20

[00432] Examples GW11 and GW12 were prepared according to the procedure described for the preparation of Examples GW9 and GW10 with the exception that (S)-2-(methoxycarbonylamino)butanoic acid was used in place of (S)-2-(methoxycarbonylamino)-3-methylbutanoic acid. Example GW11(TFA salt): LC-MS


25 (Cond. 1a): RT = 2.57 min. LC-MS Anal. Calcd. for $[\text{M}+\text{H}]^+ \text{C}_{42}\text{H}_{54}\text{ClN}_8\text{O}_6$: 801.39; found 801.38. Example GW12 (TFA salt): LC-MS (Cond. 1a): RT = 3.24 min. LC-MS Anal. Calcd. for $[\text{M}+\text{H}]^+ \text{C}_{42}\text{H}_{53}\text{Cl}_2\text{N}_8\text{O}_6$: 835.35; found 835.31.

Examples OL1 and OL2



5

Example OL1, Step a




- [00433]** Pyrrolidine 1c/4HCl (455 mg, 0.766 mmol), (S)-2-(4,4-difluorocyclohexyl)-2-(methoxycarbonylamino)acetic acid (385 mg, 1.532 mmol), HATU (612 mg, 1.609 mmol) and DIEA (0.803 mL, 4.60 mmol) were combined in DMF (30 mL) and the resulting yellow solution was stirred at ambient temperature for 3 h. Solvent was removed under reduced pressure and the residue was re-dissolved in methanol and purified by preparatory HPLC. (Solvent A: 10% Acetonitrile / 90% water / 10 mM NH₄OAc; Solvent B: 90% Acetonitrile / 10% water / 10 mM NH₄OAc; Column: SunFire Prep MS C18 30 x 150mm S10; Wavelength: 220nm; Flow rate: 40 ml/min; Gradient: 10% B to 75% B over 30 min with a 30 min hold time). An off-white solid corresponding to product OL1a (0.37 g, 0.396 mmol) was recovered. ¹H NMR (500 MHz, DMSO-d₆) δ ppm 11.67 - 12.26 (2 H, m), 7.80 (4 H, d, *J*=7.93 Hz), 7.67 - 7.76 (1 H, m), 7.65 (4 H, d, *J*=8.24 Hz), 7.54 (2 H, s), 7.23 - 7.35 (2 H, m), 5.07 (2 H, dd, *J*=8.24, 3.66 Hz), 4.54 (2 H, t, *J*=7.93 Hz), 3.62 (2 H, d, *J*=4.27 Hz), 3.55 (6 H, s), 2.36 (2 H, t, *J*=10.99 Hz), 2.20 - 2.30 (2 H, m), 2.01 (4 H, br. s.), 1.57 - 1.85 (9 H, m), 1.19 - 1.53 (6 H, m), 0.92 - 1.08 (2 H, m), 0.71 (2 H, br. s.). LC (Cond. OL3): *R*_t = 0.79 min. LC-MS: Anal. Calcd. for [M+H]⁺ C₄₈H₅₅F₄N₈O₆: 915.42; found: 915.8. HPLC purity assessment (Cond. OL4a): *R*_t = 23.12 min, homogeneity index = 98.4 %.


Examples OL1 and OL2

[00434] NCS (6.83 mg, 0.051 mmol) was added to a solution of amide OL1a (39 mg, 0.043 mmol) in DMF (2 mL) and the resulting mixture was heated to 50 °C for 5 h. The reaction mixture was then purified by preparatory HPLC. Solvent A: 10%MeOH / 90% water / 0.1% TFA; Solvent B: 90% MeOH / 10% water / 0.1% TFA; Column: SunFire Prep MS C18 30 x 100mm 5u; Wavelength: 220nm; Flow rate: 40 ml/min; Gradient: 0% B to 75% B over 30 min with a 2 min hold time. After concentration of the fractions the TFA salts of Example OL1 (15 mg) and Example OL2 (16 mg) were isolated. Example OL1: ¹H NMR (500 MHz, DMSO-d₆) δ ppm 8.11 (1 H, s), 7.74 - 8.00 (8 H, m), 7.40 (1 H, d, *J*=8.55 Hz), 7.28 (1 H, d, *J*=8.24 Hz), 4.91 - 5.05 (2 H, m), 4.48 - 4.59 (2 H, m), 3.74 - 3.87 (2 H, m), 3.62 - 3.71 (1 H, m), 3.55 (3 H, s), 3.54 (3 H, s), 2.19 - 2.43 (6 H, m), 1.66 - 2.07 (19 H, m), 1.56 (1 H, d, *J*=10.38 Hz), 1.26 - 1.48 (5 H, m), 0.89 - 1.02 (2 H, m), 0.81 (1 H, s), 0.70 (1 H, br. s.). LC (Cond. OL3): *R*_t = 0.87 min. LC-MS: Anal. Calcd. For [M+H]⁺ C₄₈H₅₄ClF₄N₈O₆: 949.38; found: 949.6. HPLC purity assessment (Cond. OL4a): *R*_t = 9.16 min, homogeneity index = 94.7 %. Example OL2: ¹H NMR (500 MHz, DMSO-d₆) δ ppm 7.86 (4 H, d, *J*=8.55 Hz), 7.79 (4 H, d, *J*=8.24 Hz), 7.29 (2 H, d, *J*=8.55 Hz), 4.95 (2 H, dd, *J*=8.55, 4.88 Hz), 4.52 (2 H, t, *J*=7.63 Hz), 3.67 (2 H, t, *J*=4.58 Hz), 3.54 (6 H, s), 2.21 - 2.40 (4 H, m), 2.01 (4 H, br. s.), 1.67 - 1.95 (13 H, m), 1.27 - 1.46 (4 H, m), 0.94 - 1.02 (2 H, m), 0.70 (2 H, d). LC (Cond. OL3): *R*_t = 1.01 min. LC-MS: Anal. Calcd. For [M+H]⁺ C₄₈H₅₃Cl₂F₄N₈O₆: 983.34; found: 983.7. HPLC purity assessment (Cond. OL4a): *R*_t = 12.25 min, homogeneity index = 94.7 %.

Examples OL3 to OL5




Example OL3: *R*₁ = *R*₂ = Cl
 Example OL4: *R*₁/*R*₂ = Cl/H
 Example OL5: *R*₁/*R*₂ = Cl/H } Regiochemistry was not determined



- [00435]** Pyrrolidine V9a (207 mg, 0.377 mmol), (S)-2-(methoxycarbonylamino)-2-(tetrahydro-2H-pyran-4-yl)acetic acid (82 mg, 0.377 mmol), HATU (158 mg, 0.415 mmol) and DIEA (0.132 mL, 0.755 mmol) were combined in DMF (5 mL) and the
- 5 resulting yellow solution was stirred at ambient temperature for 3 hr. The mixture was purified by preparatory HPLC. Solvent A: 10% MeOH / 90% water / 0.1% TFA; Solvent B: 90% MeOH / 10% water / 0.1% TFA; Column: SunFire Prep MS C18 30 x 100mm 5u; Wavelength: 220nm; Flow rate: 40 mL/min; Gradient: 20% B to 80% B over 30 min. with a 2 min hold time. A white solid corresponding to the TFA salt of carbamate OL3a
- 10 (0.19 g) was recovered [Note: carbamate OL3a and V9b are the same besides their form status]. ¹H NMR (500 MHz, DMSO-d₆) δ ppm 8.18 (1 H, br. s.), 8.12 (1 H, s), 7.80 - 8.00 (8 H, m), 7.34 (1 H, d, *J*=8.55 Hz), 5.01 (1 H, t, *J*=8.09 Hz), 4.79 - 4.90 (1 H, m), 4.49 (1 H, t, *J*=7.48 Hz), 3.74 - 3.87 (3 H, m), 3.55 (4 H, s), 3.18 - 3.32 (2 H, m), 2.52 - 2.59 (2 H, m), 2.37 (2 H, ddd, *J*=19.99, 13.43, 6.26 Hz), 2.07 (1 H, br. s.), 1.92 (1 H, dt, *J*=13.20, 6.68 Hz), 1.73 (1 H, ddd, *J*=13.20, 6.49, 6.26 Hz), 1.06 - 1.52 (17 H, m), 0.80
- 15 (2 H, br. s.), 0.74 (1 H, br. s.). LC (Cond. OL1): *R*_t = 1.98 min. LC-MS: Anal. Calcd. For [M+H]⁺ C₄₂H₅₀N₇O₆: 748.38; found: 748.52.

Example OL3, Step b




20

- [00436]** Carbamate OL3a (0.19 g, 0.195 mmol) was dissolved in CH₂Cl₂ (15 mL) and charged with 4N HCl in dioxanes (3 mL, 12.00 mmol). The resulting suspension was stirred at ambient temperature for 2 h and volatiles were removed under reduced pressure.

A yellowish solid corresponding to the HCl salt (3X) of pyrrolidine OL3b (0.12 g) was recovered and used without further purification. ^1H NMR (500 MHz, DMSO- d_6) δ ppm 8.14 (1 H, s), 7.83 - 8.01 (10 H, m), 7.34 (1 H, d, $J=8.55$ Hz), 5.06 (1 H, t, $J=7.93$ Hz), 4.73 (1 H, t, $J=8.24$ Hz), 4.50 (1 H, t, $J=7.48$ Hz), 3.78 - 3.87 (3 H, m), 3.64 - 3.74 (2 H, m), 3.56 (3 H, s), 3.44 - 3.51 (1 H, m), 3.39 - 3.44 (1 H, m), 3.20 - 3.35 (2 H, m), 2.55 - 2.64 (2 H, m), 2.34 - 2.43 (1 H, m), 2.06 - 2.16 (1 H, m), 1.87 - 1.97 (2 H, m), 1.27 - 1.51 (4 H, m), 1.12 (1 H, d, $J=4.58$ Hz), 0.90 - 0.98 (1 H, m), 0.82 - 0.89 (1 H, m), 0.78 (1 H, br. s.). LC (Cond. OL2): $R_t = 2.51$ min. LC-MS: Anal. Calcd. For $[\text{M}+\text{H}]^+ \text{C}_{37}\text{H}_{42}\text{N}_7\text{O}_4$: 648.33; found: 648.4.

10

Example OL3, Step c



[00437] (S)-2-(4,4-Difluorocyclohexyl)-2-(methoxycarbonylamino)acetic acid (87 mg, 0.345 mmol), the HCl salt (3X) of pyrrolidine OL3b (261 mg, 0.345 mmol), HATU (144 mg, 0.379 mmol) and DIEA (0.301 mL, 1.724 mmol) were combined in DMF (10 mL) and the resulting brownish solution was stirred at ambient temperature for 2 h. Sample was purified directly by preparatory HPLC. Solvent A: 10%MeOH / 90% water / 0.1% TFA; Solvent B: 90% MeOH / 10% water / 0.1% TFA; Column: SunFire Prep MS C18 30 x 100mm 5u; Wavelength: 220nm; Flow rate: 40 mL/min; Gradient: 30% B to 70% B over 30 min with a 2 min hold time. A white solid corresponding to the TFA salt (2x) of amide OL3c (132 mg, 0.117 mmol, 33.8 % yield) was recovered. ^1H NMR (500 MHz, DMSO- d_6) δ ppm 11.68 - 12.18 (2 H, m), 7.78 (3 H, t, $J=8.39$ Hz), 7.73 (1 H, d, $J=9.46$ Hz), 7.66 (4 H, d, $J=8.55$ Hz), 7.54 (2 H, dd, $J=4.12, 1.68$ Hz), 7.28 (2 H, dd, $J=19.68, 8.39$ Hz), 5.08 (2 H, ddd, $J=8.39, 4.12, 3.97$ Hz), 4.46 - 4.58 (2 H, m), 3.82 - 3.94 (2 H, m), 3.61 - 3.67 (2 H, m), 3.54 (6 H, d, $J=1.53$ Hz), 3.24 (2 H, t, $J=11.29$ Hz), 2.33 - 2.41 (2 H, m), 2.21 - 2.31 (2 H, m), 1.91 - 2.08 (4 H, m), 1.75 - 1.89 (6 H, m), 1.58 (1 H, d, $J=13.43$ Hz), 1.40 - 1.52 (2 H, m), 1.26 - 1.40 (3 H, m), 0.97 - 1.07 (2 H, m), 0.67 - 0.79

(2 H, m). LC (Cond. OL1): $R_t = 1.94$ min. LC-MS: Anal. Calcd. For $[M+H]^+$
 $C_{47}H_{55}N_8O_7$: 881.42; found: 881.58.

Examples OL3 to OL5

- 5 **[00438]** NCS (22.66 mg, 0.170 mmol) was added to a solution of the TFA salt of Example OL3c (115 mg, 0.131 mmol) in DMF (3 mL) and the resulting mixture was heated to 50 °C for 5 h. The mixture was purified by preparatory HPLC. Solvent A: 05% MeCN / 95% water / 10 mM NH_4Ac ; Solvent B: 95% MeCN / 5% water / 10 mM NH_4Ac ; Column: SunFire Prep MS C18 30 X 100mm S10; Wavelength: 220nm; Flow rate: 35 ml/min; Gradient: 10% B to 100% B over 30 min with a 2 min hold time. Two fractions were isolated, where the first elute corresponded to a mixture of mono-chlorinated analogs (Examples OL4 and OL5) and the second one corresponded to the bis-chlorinated analog (Example OL3). After concentration of the corresponding fractions, the mixture was re-dissolved in methanol and the monochlorinated
- 10 regioisomers were separated by preparatory HPLC. Solvent A: 10% Acetonitrile/ 90% water / 0.1% TFA; Solvent B: 90% Acetonitrile / 10% water / 0.1% TFA; Column: PHENOMENEX® Luna 21 x 100mm S10; Wavelength: 220nm; Flow rate: 25 ml/min; Gradient: 10% B to 50% B over 60 min. with a 2 min hold time. Two fractions corresponding to Example OL4 and Example OL5 were isolated as TFA and that their
- 20 relative regiochemistry was not determined.
- [00439]** Example OL3 (36 mg) was recovered as a white solid. 1H NMR (500 MHz, DMSO- d_6) δ ppm 12.55 (2 H, d, $J=1.22$ Hz), 7.87 (4 H, d, $J=8.24$ Hz), 7.78 (4 H, d, $J=7.02$ Hz), 7.27 (2 H, dd, $J=18.62, 8.55$ Hz), 5.76 (1 H, s), 4.91 - 5.01 (2 H, m), 4.45 - 4.58 (2 H, m), 3.85 (2 H, d, $J=9.77$ Hz), 3.64 - 3.71 (2 H, m), 3.54 (6 H, s), 3.20 - 3.31 (2
- 25 H, m), 2.21 - 2.35 (4 H, m), 2.02 (3 H, d, $J=3.97$ Hz), 1.59 - 1.93 (9 H, m), 1.21 - 1.55 (7 H, m), 0.93 - 1.02 (2 H, m), 0.71 (2 H, m). LC (Cond. OL3): $R_t = 0.95$ min. LC-MS: Anal. Calcd. For $[M+H]^+$ $C_{47}H_{53}Cl_2F_2N_8O_7$: 949.34; found: 949.8. HPLC purity assessment (Cond. OL4b): $R_t = 11.10$ min, homogeneity index = 100 %.
- [00440]** Example OL4 (10 mg) was recovered as a white solid. 1H NMR (500 MHz, DMSO- d_6) δ ppm 14.62 (1 H, br. s.), 12.59 (1 H, br. s.), 8.12 (1 H, br. s.), 7.87 (8 H, dd, $J=19.38, 8.09$ Hz), 7.80 (2 H, d, $J=7.63$ Hz), 7.40 (1 H, d, $J=8.85$ Hz), 7.26 (1 H, d, $J=8.24$ Hz), 5.00 (1 H, t, $J=7.93$ Hz), 4.95 (1 H, dd, $J=8.39, 5.04$ Hz), 4.54 (1 H, t,
- 30

$J=7.48$ Hz), 4.49 (1 H, t, $J=7.93$ Hz), 3.86 (3 H, dd, $J=11.44$, 2.59 Hz), 3.79 (1 H, br. s.), 3.65 - 3.72 (2 H, m), 3.55 (6 H, d, $J=5.19$ Hz), 3.19 - 3.28 (2 H, m), 2.20 - 2.43 (4 H, m), 1.89 - 2.07 (5 H, m), 1.30 - 1.89 (10 H, m), 0.92 - 1.03 (2 H, m), 0.80 (1 H, br. s.), 0.72 (1 H, br. s.) LC (Cond. OL3): $R_t = 0.85$ min. LC-MS: Anal. Calcd. For $[M+H]^+$


- 5 $C_{47}H_{54}ClF_2N_8O_7$: 915.38; found: 915.9. HPLC purity assessment (Cond. OL4a): $R_t = 9.03$ min, homogeneity index = 100 %.

[00441] Example OL5 (12 mg, 7.96 % yield) was recovered as a white solid. 1H NMR (500 MHz, DMSO- d_6) δ ppm 14.62 (1 H, br. s.), 12.59 (1 H, br. s.), 8.09 (1 H, br. s.), 7.87 - 7.98 (4 H, m), 7.84 (2 H, d, $J=8.24$ Hz), 7.80 (2 H, d, $J=8.24$ Hz), 7.35 (1 H, d, $J=8.24$ Hz), 7.29 (1 H, d, $J=8.55$ Hz), 4.91 - 5.03 (2 H, m), 4.51 (2 H, ddd, $J=15.64$, 7.71, 7.48 Hz), 3.78 - 3.90 (4 H, m), 3.64 - 3.71 (1 H, m), 3.54 (6 H, s), 3.16 - 3.31 (2 H, m), 2.19 - 2.44 (3 H, m), 2.02 (3 H, br. s.), 1.64 - 1.95 (7 H, m), 1.27 - 1.54 (6 H, m), 0.96 (2 H, dd, $J=12.21$, 7.32 Hz), 0.81 (1 H, br. s.), 0.70 (1 H, br. s.). LC (Cond. OL3): $R_t = 0.86$ min. LC-MS: Anal. Calcd. For $[M+H]^+$ $C_{47}H_{54}ClF_2N_8O_7$: 915.38; found: 915.9. HPLC purity

10 assessment (Cond. OL4a): $R_t = 8.11$ min, homogeneity index = 100 %.


15

Example OL6




- [00442] Example OL6 (TFA salt) can be prepared starting from pyrrolidine V1b/4HCl
- 20 and Cap-179 (Enantiomer-1) according to the procedure described for the synthesis of Example V1. 1H NMR (400 MHz, DMSO- d_6) δ ppm 8.11 (1 H, br. s.), 7.74 - 8.01 (8 H, m), 7.22 - 7.35 (1 H, m), 7.14 (1 H, d, $J=7.53$ Hz), 4.87 - 5.15 (2 H, m), 4.47 (2 H, d, $J=7.03$ Hz), 3.80 (1 H, br. s.), 3.65 (1 H, br. s.), 3.55 (6 H, br. s.), 3.36 (4 H, br. s.), 2.20 - 2.37 (3 H, m), 2.13 (1 H, br. s.), 1.90 - 2.05 (2 H, m), 1.85 (1 H, br. s.), 1.63 - 1.80 (1 H, m), 1.38 - 1.61 (2 H, m), 0.62 - 1.14 (25 H, m). LC (Cond. 2a and 2b): >95%
- 25 homogeneity index. LC (Cond. OL4c): $R_t = 0.85$ min. LC-MS: Anal. Calcd. for $[M+H]^+$ $C_{50}H_{62}ClN_8O_8$: 937.44; found: 937.5.

Example OL7



5


Example OL7, Step a



[00443] To a solution of (2S,2'S)-*tert*-butyl 2,2'-(5,5'-(biphenyl-4,4'-diyl)bis(1H-imidazole-5,2-diyl))dipyrrolidine-1-carboxylate (500 mg, 0.800 mmol) in AcOH (30 mL) was added bromine (0.041 mL, 0.800 mmol) in AcOH (0.8 mL) dropwise over 10 min.

- 10 The resulting reaction mixture was stirred at room temperature overnight, neutralized with sat. NaHCO₃, then extracted with CH₂Cl₂. The organic layers was dried with MgSO₄ and concentrated *in vacuo*. The resultant crude material was purified by flash chromatography (silica gel; 1:1 EtOAc/Hex then 2:1 EtOAc/Hex) to afford bromide OL7a.1 (200 mg) and dibromide OL7a.2 (140 mg) as white solids (note: the dibromide
- 15 eluted first from column).


Example OL7, Step b.1



- [00444] A mixture of cyclopropylboronic acid (13.49 mg, 0.157 mmol), potassium phosphate tribasic (90 mg, 0.423 mmol) in water (0.03 mL) was stirred at room temperature for 15 min. Bromide OL7a.1 (85 mg, 0.121 mmol), Pd(II)acetate (2.71 mg, 0.012 mmol), tricyclohexylphosphine (6.8 mg, 0.024 mmol) and toluene (1 mL) were

then added, and the reaction mixture was stirred at 110 °C overnight. The volatile component was removed *in vacuo* and the residue was purified by preparative HPLC using the following condition. Column: Waters SunFire OBD 19 X 100 mm S5; Solvent A = 10%ACN-90%H₂O-0.1%TFA; Solvent B = 90%ACN-10%H₂O-0.1%TFA; Start %B = 10; Final %B = 75; Flow rate = 20 ml/min; Gradient time = 20 min. The TFA salt of the coupled product OL7b.1 was retrieved as a light yellow solid (79 mg).


Example OL7, Step b.2



10 [00445] A mixture of dibromide OL7a.2 (140 mg, 0.179 mmol), cyclopropylboronic acid (40.0 mg, 0.465 mmol), Pd(II)acetate (8.03 mg, 0.036 mmol), potassium phosphate tribasic (266 mg, 1.252 mmol), tricyclohexylphosphine (20.07 mg, 0.072 mmol) in toluene (1.5 mL) and Water (0.05 mL) was heated with a microwave at 110 °C for 1.5 hours. LCMS showed that there is no reaction. The mixture was filtered, and the filtrate was concentrated *in vacuo* and the resultant crude product was purified by prep. HPLC: Column = Waters SunFire OBD 19 X 100 mm S5; Solvent A = 10%ACN-90%H₂O-0.1%TFA; Solvent B = 90%ACN-10%H₂O-0.1%TFA; Start %B = 10. Final %B = 70; Flow rate = 20 ml/min; Gradient time = 20 min. The TFA salt of the coupled product OL7b.2 was obtained as a light yellow solid.

20

Example OL7, Step c



[00446] To a solution of carbamate OL7b prepared above (79 mg) in CH₂Cl₂ (2 mL) was added HCl (2 mL, 8.00 mmol) (4 N in dioxane), and the resulting mixture was stirred at room temperature for 30 mins. The volatile component was removed *in vacuo* to afford

25


the HCl salt of OL7c (69 mg), which was used in the next step without further purification.


Example OL7













- 5 [00447] To a mixture of (S)-2-(methoxycarbonylamino)-3-methylbutanoic acid (20.09 mg, 0.115 mmol), pyrrolidine OL7/4 HCl (35 mg, 0.057 mmol) in DMF (1 mL) was added DiPEA (0.06 mL, 0.34 mmol) and HATU (43.6 mg, 0.115 mmol). The reaction mixture was stirred at room temperature for 1 hour, and then ammonia (2 mL, 2 M in Methanol) was added, and stirring was continued for an additional 1 hour. The volatile component was removed *in vacuo*, and the residue was purified by preparative HPLC: Column = Waters SunFire C18 19 X 100 mm 5u; Solvent A = 10%MeOH-90%H₂O-0.1%TFA; Solvent B = 90%MeOH-10%H₂O-0.1%TFA; Start %B = 10; Final %B = 75; Flow rate = 25 ml/min; Gradient time = 18 min. The TFA salt of Example OL7 was obtained as an off-white solid (24 mg). Rt = 1.68 min (Cond. OL5a). (M+H)⁺ 779.47. ¹H NMR (500 MHz, DMSO-d₆) δ ppm 8.12 (1 H, br. s.), 7.97 (4 H, dd, *J*=12.97, 8.39 Hz), 7.86 - 7.92 (2 H, m), 7.80 (2 H, d, *J*=8.24 Hz), 7.32 (2 H, t, *J*=8.70 Hz), 5.14 (1 H, t, *J*=7.02 Hz), 5.05 (1 H, t, *J*=7.48 Hz), 4.12 (2 H, t, *J*=6.71 Hz), 3.76 - 3.92 (4 H, m), 3.54 (6 H, s), 2.39 (2 H, d, *J*=6.10 Hz), 1.95 - 2.21 (9 H, m), 1.08 (2 H, d, *J*=8.24 Hz), 0.84 (7 H, t, *J*=7.17 Hz), 0.79 (6 H, d, *J*=6.41 Hz), 0.67 - 0.74 (1 H, m)

Examples OL8 to OL14


[00448] Examples OL8 to OL14 were prepared as TFA salts by employing appropriate precursors and the procedure described for the preparation of Example OL7.




Example		R ₂	Rt(Condition). LC-MS (M+H) ⁺ observed
---------	---	----------------	--

Example		R ₂	Rt(Condition). LC-MS (M+H) ⁺ observed
OL8		H	Rt = 1.49 min (Cond. OL5a). (M+H) ⁺ 863.48
OL9			Rt = 1.77 min (Cond. OL5a). (M+H) ⁺ 819.55
OL10			Rt = 1.54 min (Cond. OL5a). (M+H) ⁺ 883.51
OL11		H	Rt = 1.74 min (Cond. OL5a). (M+H) ⁺ 803.41
OL12		H	Rt = 2.40 min (Cond. OL5b). (M+H) ⁺ 887.60
OL13			Rt = 1.87 min (Cond. OL5a). (M+H) ⁺ 843.39
OL14			Rt = 2.55 min (Cond. OL5b). [(M+H)/2] ⁺ = 464.36

Example OL15





[00449] A mixture of bromide OL7a.1 (200 mg, 0.284 mmol), CsF (0.021 mL, 0.568 mmol), Pd(Ph₃P)₄ (19.71 mg, 0.017 mmol) and 2-allyl-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (0.107 mL, 0.568 mmol) in THF (3 mL) was heated at 140 °C in microwave for 1 hour. The solvent was removed *in vacuo*, and the residue was purified by preparative HPLC. Column, PHENOMENEX® Luna 10u (30 X 100 mm); Solvent A = 5%CH₃CN-95%H₂O-10mmNH₄Oac; Solvent B = 95%CH₃CN-5%H₂O-10mmNH₄Oac; Start %B = 30; Final %B = 100; Flow rate = 25 ml/min; Gradient time = 48 min. Product OL15a was obtained as a white solid (90 mg).


10

Example OL15

[00450] Carbamate OL15a was elaborated to the TFA salt of Example OL15 according to the procedure described for the synthesis of Example OL7. Rt = 1.73 min (Cond. OL5a). (M+H)⁺ 779.53.

15


Example OL16



[00451] Example OL16 (TFA salt) was prepared from appropriate precursors according to the procedure described for the synthesis of Example OL15. Rt = 1.73 min (Cond. OL5a). (M+H)⁺ 863.11.

20

Examples OL17 and OL18



[00452] To a solution of OL15a (70 mg, 0.105 mmol), diiodomethane (0.102 mL, 1.263 mmol) in toluene (0.5 mL) at 0 °C was quickly added diethylzinc (1.263 mL, 1.263 mmol) (1 M in heptane). The resulting mixture was stirred at 0 °C for 1.5 hour then at


5 room temperature for 4 hours. To the mixture was added a saturated aqueous NaHCO₃ solution and the mixture was extracted with CH₂Cl₂. The combined organic layers were dried under MgSO₄ and concentrated. The residue was purified by preparatory HPLC: Column, PHENOMENEX® Luna 10u (30 X 100 mm); Solvent A = 5%CH₃CN-95%H₂O-10mm NH₄Oac; Solvent B = 95%CH₃CN-5%H₂O-10mmNH₄OAc; Start %B =

10 30; Final %B = 100; Flow rate = 25 ml/min. Gradient time = 25 min. The resultant material (white solid, 40 mg) was dissolved in CH₂Cl₂ (1 mL) and treated with HCl (1 mL, 4.00 mmol) (4 N in dioxane). The mixture was stirred at room temperature for 1 hour. The volatile component was removed *in vacuo* to afford a product (35 mg), which was coupled with (S)-2-(methoxycarbonylamino)-3-methylbutanoic acid according to the


15 procedure described for the synthesis of Example OL7. The obtained crude residue was purified by preparative HPLC: Column = Waters SunFire C18 19 X 100 mm 5u; Solvent A = 10%MeOH-90%H₂O-0.1%TFA; Solvent B = 90%MeOH-10%H₂O-0.1%TFA; Start %B = 20; Final %B = 85; Flow rate = 25 ml/min; Gradient time = 27 min. Two fractions were isolated, where the first one to elute corresponded to the TFA salt of Example OL18

20 (Rt = 1.74 min (Cond. OL5a). (M+H)⁺ 793.54) and the second one to elute corresponded to the TFA salt of Example OL17 (Rt = 1.77 min (Cond. OL5a). (M+H)⁺ 793.53).

Example OL19




Example OL19, Step a



[00453] A mixture of 1-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole (59.1 mg, 0.284 mmol), potassium phosphate tribasic (121 mg, 0.568 mmol) in water (0.2 mL) was stirred at room temperature for 15 min. Bromide OL7a.1 (100 mg, 0.142 mmol), Pd(Ph₃P)₄ (9.85 mg, 0.008 mmol) and DMF (2 mL) were then added, and the reaction mixture was stirred for 40 min. at 130 °C under microwave radiation. The volatile component was removed *in vacuo* and the residue was purified by preparative HPLC using the following condition. Column: PHENOMENEX® Luna 10u (30 X 100 mm); Solvent A = 10%ACN-90%H₂O-10 mm NH₄OAc; Solvent B = 90%ACN-10%H₂O-10mmNH₄OAc; Start %B = 10; Final %B = 80; Flow rate = 25 ml/min; Gradient time = 15 min. The coupled product OL19a was retrieved as a light yellow solid (120 mg).

15

Example OL19, Step b



[00454] To a solution of carbamate OL19a prepared above (120 mg) in CH₂Cl₂ (1 mL) was added HCl (1 mL, 4.00 mmol) (4 N in dioxane), and the resulting mixture was stirred at room temperature for 1h. The volatile component was removed *in vacuo* to afford the HCl salt of OL19b (87 mg), which was used in the next step without further purification.

20

Example OL19

[00455] To a mixture of (S)-2-(methoxycarbonylamino)-3-methylbutanoic acid (15.30 mg, 0.087 mmol), pyrrolidine OL19b/4 HCl (30 mg, 0.044 mmol) in DMF (1 mL) was


25

added DiPEA (0.053 mL, 0.306 mmol) and HATU (33.2 mg, 0.087 mmol). The reaction mixture was stirred at room temperature for 1 hour, the volatile component was removed *in vacuo*, and the residue was purified by preparative HPLC: Column = Waters Atlantis OBD 30 X 100 mm 5u; Solvent A = 10%MeOH-90%H₂O-0.1%TFA; Solvent B =

- 5 90%MeOH-10%H₂O-0.1%TFA; Start %B = 20; Final %B = 75; Flow rate = 25 ml/min; Gradient time = 25 min. The TFA salt of Example OL19 was obtained as a white solid (33 mg). Rt = 1.63 min (Cond. OL5a). (M+H)⁺ 819.40. ¹H NMR (500 MHz, DMSO-d₆) δ ppm 8.13 (1 H, s), 7.88 - 7.96 (2 H, m), 7.78 - 7.88 (4 H, m), 7.57 (1 H, br. s.), 7.43 - 7.51 (2 H, m), 7.33 (2 H, dd, *J*=8.39, 3.51 Hz), 6.42 (1 H, br. s.), 5.10 - 5.18 (2 H, m), 4.11 (2
10 H, td, *J*=7.93, 2.75 Hz), 3.78 - 3.91 (4 H, m), 3.59 - 3.66 (3 H, m), 3.54 (6 H, d, *J*=1.53 Hz), 2.35 - 2.45 (1 H, m), 2.30 (1 H, d, *J*=7.32 Hz), 2.13 - 2.22 (2 H, m), 1.93 - 2.13 (6 H, m), 0.89 (5 H, d, *J*=6.71 Hz), 0.83 (6 H, t, *J*=6.87 Hz), 0.78 (3 H, d, *J*=6.71 Hz).








Examples OL20 to OL25








- 15 [00456] Examples OL20 to OL25 were prepared as TFA salts by employing appropriate precursors and the procedure described for the preparation of Example OL19.




Example OL20-OL23

Example OL24-OL25

Example		R ₂	Rt(Condition). LC-MS (M+H) ⁺ observed
OL20			Rt = 2.19 min (Cond. OL5b). (M+H) ⁺ 903.65
OL21			Rt = 1.56 min (Cond. OL5a). (M+H) ⁺ 819.39
OL22			Rt = 1.43 min (Cond. OL5a). (M+H) ⁺ 816.66


Example		R ₂	Rt(Condition). LC-MS (M+H) ⁺ observed
OL23			Rt = 1.27 min (Cond. OL5a). (M+H) ⁺ 903.61
OL24			Rt = 1.83 min (Cond. OL5a). (M+H) ⁺ 830.50
OL25			Rt = 1.62 min (Cond. OL5a). [(M+H)/2] ⁺ 457.6

Example DSTL-1



5

Example DSTL-1, Step a




[00457] Diphenylphosphoryl azide (17.09 mL, 79 mmol) was added to a solution of 6-bromo-2-naphthoic acid (16.5 g, 65.7 mmol), triethylamine (18.32 mL, 131 mmol), and *tert*-butylalcohol (7.54 mL, 79 mmol) in toluene (225 mL) and stirred for 4 h at 100 °C.

- 10 The volatiles were removed by rotary evaporation and the residue taken up in EtOAc (500 mL) and washed with water and brine. A precipitate formed upon concentration which was isolated by filtration and washed with 1:1 Et₂O/Hex to give Example DSTL-1, Step a (10.5 g). A second crop of less pure product was isolated upon concentration of

the mother liquor (9.8 g); combined yield (93%). LC-MS (Cond.-J4): RT = 3.44 min.

LC-MS Anal. Calcd. for $[M+Na]^+$ C₁₅H₁₆BrNO₂: 345.02; found 345.03.

Example DSTL-1, Step b




5

[00458] Example DSTL-1, Step a (5 g, 15.52 mmol) was diluted in acetic acid (50 mL) and fuming nitric acid (2.3 mL) was added dropwise over 20 min. The reaction was stirred for 2 h and the product, isolated by filtration, was partitioned between DCM and sat'd NaHCO₃ soln. The organic layer was concentrated and Example DSTL-1, Step b

10 was obtained 5.7 g (quant). LC-MS (Cond.-J4): RT = 3.52 min. LC-MS Anal. Calcd. for $[M+Na]^+$ C₁₅H₁₅BrN₂O₄: 390.02; found 390.99.

Example DSTL-1, Step c




HCl salt

15 **[00459]** Tin(II)chloride dehydrate (3 g, 16.34 mmol) was added to a solution of Example DSTL-1, Step b (2 g, 5.47 mmol) in MeOH (100 mL) and the solution was stirred for 18 h at 70 °C. The solvent was removed by rotary evaporation and Example DSTL-1, Step c (assume theoretical 1.25 g) was dried under high vacuum. LC-MS (Cond.-J4): RT = 1.49 min. LC-MS Anal. Calcd. for $[M+H]^+$ C₁₀H₉BrN₂: 237.00; found


20 236.96.

Example DSTL-1, Step d




[00460] HATU (2.085 g, 5.48 mmol) was added to a solution of Example DSTL-1, Step c (1.25 g, 5.48 mmol), Example 1, Step b (1.246 g, 5.48 mmol), and Hunig's base (7.09 g, 54.8 mmol) in DMF (70 mL) and the reaction mixture was stirred for 6 h before being partitioned between EtOAc (500 mL) and sat'd NaHCO₃ (150 mL). The tin salts that precipitated were removed by filtration through CELITE®, and the organic phase was concentrated to yield a residue which was taken up in AcOH (100 mL) and heated at 60 °C for 18 h. The solvent was removed by rotary evaporation under high vacuum, and residue taken up in CH₂Cl₂ and washed with sat'd NaHCO₃ soln. After concentration, the crude product was charged (DCM) to a Thompson silica gel cartridge (110 g) and subject to gradient elution; 15 - 100% B over 1 L to give Example DSTL-1, Step d (420 mg, 44 %). LC-MS (Cond.-J4): RT = 2.51 min. LC-MS Anal. Calcd. for [M+H]⁺ C₂₁H₂₂BrN₃O₂: 430.10.; found 430.06.

Example DSTL-1, Step d.1



[00461] Example DSTL-1, Step d.1 was prepared from Example 1, Step b according to the procedure described for the synthesis of its desmethano analog in patent application WO 2008/021927.


Example DSTL-1, Step e



[00462] To a solution of Example DSTL-1, Step d.1 (560 mg, 1.39 mmol) in DMF (15 mL) was added NCS (203 mg, 1.52 mmol). The reaction mixture was heated at 50 °C for 16 h. The solution was purged with nitrogen to evaporate DMF, and the crude residue was charged (DCM) to a Thompson SiO₂ column (80 g) and gradient elution performed (BIOTAGE®). Segment 1: 10 - 100% B over 1.5 L. Segment 2: Hold 100% B 300 mL. (A/B Hexanes/EtOAc). There was isolated Example DSTL-1, Step e (589.1 mg, 92 %

yield). LC-MS (Cond.-D4): RT = 2.61 min. LC-MS Anal. Calcd. for $[M+Na]^+$
 $C_{19}H_{22}ClN_3O_2$: 462.04; found 462.07.

Example DSTL-1, Step f




5

[00463] Tetrakis(triphenylphosphine) palladium (76 mg, 0.066 mmol) was added in one portion to a stirred suspension of Example DSTL-1, Step e (580 mg, 1.32 mmol), bis(pinacolato)diboron (671 mg, 2.64 mmol) and potassium acetate (324 mg, 3.30 mmol) in a solution of dioxane (12 mL) in a screw-capped pressure vessel. The reaction mixture was evacuated and flushed with argon (3x) and placed into a pre-heated oil bath (80 °C) and stirred for 16 h. Upon cooling, the reaction mixture was diluted with EtOAc, washed with sat'd $NaHCO_3$ soln, brine and dried over Na_2SO_4 . The crude residue was charged (DCM) to a Thompson SiO_2 column (80 g) and gradient elution performed (BIOTAGE®). Segment 1: 5 - 100% B over 1.5 L. Segment 2: Hold 100% B 300 mL. (A/B CH_2Cl_2 /EtOAc). There was isolated Example DTSL-1, Step f (714.7 mg, 95%) as yellow foam. LC-MS (Cond.-D4): RT = 2.69 min. LC-MS Anal. Calcd. for $[M+H]^+$
 $C_{25}H_{34}BClN_3O_4$: 486.24; found 486.26.

10

15

Example DSTL-1, Step g




20

[00464] Tetrakis(triphenylphosphine) palladium (27.0 mg, 0.023 mmol) was added in one portion to a stirred suspension of Example DSTL-1, Step d (200 mg, 0.467 mmol), Example DSTL-1, Step f (250 mg, 0.514 mmol) and $NaHCO_3$ (196 mg, 2.335 mmol) in a degassed solution of DME (4 mL) and water (1 mL) under argon in a screw-capped pressure vessel. The solution was evacuated and charged with argon (3x) and placed into a pre-heated oil bath (80 °C) and stirred for 14 h. The mixture was diluted with EtOAc

25

- (10 mL), THF (2 mL) and MeOH (1 mL), washed with brine and dried over Na₂SO₄. After concentration to remove solvent, the residue was charged to a Thompson SiO₂ column (80 g) and eluted (BIOTAGE®) by gradient. Segment 1: 20 - 100% B over 1.5 L, Segment 2: hold 100% B for 300 mL; A/B Hexanes/EtOAc. There was isolated
- 5 DSTL-1, Step g (195.0 mg, 52.6%) as a tan solid. LC-MS (Cond.-D4): RT = 2.40 min. LC-MS Anal. Calcd. for [M+H]⁺ C₄₀H₄₄ClN₆O₄: 707.31; found 707.43.

Example DSTL-1, Step h



HCl salt

- 10 **[00465]** Cold (0 °C) 4*N* HCl in dioxane (4 mL) was added to a stirred solution of Example DSTL-1, Step g (65 mg, 0.092 mmol) in MeOH (1 mL). The mixture was stirred at rt for 3 h, concentrated, and placed under high vacuum to provide DSTL-1, Step h as an orange-tan solid and as a tetra HCl salt. LC-MS (Cond.-D4): RT = 1.63 min. LC-MS Anal. Calcd. for [M+H]⁺ C₃₀H₂₈ClN₆: 507.21; found 507.26.


15

Example DSTL-1

- [00466]** Example DSTL-1 was prepared from Example DSTL-1, Step h according to the procedure described for the preparation of Example GW2. Purification was accomplished by preparative HPLC (0 - 50% B) over a 25 min gradient (at 40ml/min)
- 20 using a Waters SunFire column (30 x 100 mm, S5) where Solvent B = 90% CH₃CN - 10% H₂O - 0.1% TFA and A = 5% CH₃CN - 95% H₂O - 0.1% TFA. %. There was isolated Example DSTL-1 (70.4 mg, 67.7%) as a white solid. ¹H NMR (MeOD, 500 MHz, δ): 8.47 (d, *J* = 8.6 Hz, 1H), 8.44 (s, 1H), 8.15 (s, 1H), 8.13 (s, 1H), 7.91 (d, *J* = 8.6 Hz, 2H), 7.85 (d, *J* = 8.6 Hz, 2H), 7.80 (d, *J* = 8.6 Hz, 1H), 5.34-5.31 (m, 1H), 5.07 (t, *J* = 7.32 Hz, 1H), 4.61 - 4.58 (m, 2H), 3.39-3.31 (m, 1H), 3.75-3.72 (m, 1H), 3.69 (s, 3H), 3.68 (s, 3H), 2.83-2.81 (m, 1H), 2.64-2.59 (m, 1H), 2.49-2.48 (m, 2H), 2.23-2.14 (m, 3H), 2.05-2.03 (m, 1H), 1.18-1.12 (m, 2H), 1.05-0.92 (m, 13H), 0.83-0.81 (m, 1H). LC-MS
- 25

(Cond.-D4): RT = 2.24 min. LC-MS Anal. Calcd. for $[M+H]^+$ $C_{44}H_{50}ClN_8O_6$: 821.36; found 821.54.

Example DSTL-2




5

[00467] Example DSTL-2 was prepared from Example DSTL-1, Step h according to the procedure described for the preparation of Example GW2 except using (S)-2-(methoxycarbonylamino)-2-(tetrahydro-2H-pyran-4-yl)acetic acid instead of Cap-51.


Purification was accomplished by preparative HPLC (0 - 50% B) over a 25 min gradient (at 40ml/min) using a Waters SunFire column (30 x 100 mm, S5) where Solvent B = 90% CH_3CN - 10% H_2O - 0.1% TFA and A = 5% CH_3CN - 95% H_2O - 0.1% TFA. There was isolated Example DSTL-2 (18.0 mg) as a white solid. LC-MS (Cond.-D4): RT = 2.04 min. LC-MS Anal. Calcd. for $[M+H]^+$ $C_{48}H_{54}ClN_8O_6$: 905.38; found 905.60.

15

Examples DSTL-3 and DSTL-4




Examples DSTL-3, Step a and DSTL-4, Step a



- [00468] A solution of bromine (0.079 mL, 1.54 mmol) in AcOH (1 mL) was added portionwise to a stirred solution of Example 1, Step d (1.0 g, 1.541 mmol) in AcOH (40 mL), and the mixture was stirred at rt for 3 h. The solvent was removed *in vacuo* and the residue was taken up in DCM and washed with sat'd NaHCO₃ soln. The aqueous layer was extracted twice more with DCM and the combined organic extracts were washed with brine and dried (Na₂SO₄). The residue was charged (DCM) to a Thompson silica gel cartridge (160 g) and eluted with 30 - 100% B over 2 L. Segment 2: hold 100% B for 800 mL. A/B Hexanes/EtOAc. There was isolated 3 components; Starting material (303.6 mg); Example DSTL-3, Step a (X = H; 382.1 mg); yellow solid. LC-MS (Cond.-D4): RT = 2.21 min. LC-MS Anal. Calcd. for [M+H]⁺ C₃₈H₄₄BrN₆O₄: 729.26; found . 729.28; Example DTSL-4, Step a (X = Br; 287.2 mg); light, yellow solid. LC-MS (Cond.-D4): RT = 2.62 min. LC-MS Anal. Calcd. for [M+H]⁺ C₃₈H₄₃Br₂N₆O₄: 807.17; found: 807.33.

- 15 Examples DSTL-3, Step b and DSTL-4, Step b



- [00469] The products of Example DSTL-3, Step b and DSTL-4, Step b were brought forward separately. Removal of the protecting groups was performed as described in the procedure from Example DSTL-1, Step h. X = H; Example DTSL-3, Step b: Yellow solid, 33.1 mg (97% pure); LC-MS (Cond.-D4): RT = 1.49 min. LC-MS Anal. Calcd. for [M+H]⁺ C₂₈H₂₈BrN₆: 529.16; found 529.24. X = Br; Example DTSL-4, Step b: Yellow solid, 33 mg (96% pure); LC-MS (Cond.-D4): RT = 1.82 min. LC-MS Anal. Calcd. for [M+H]⁺ C₂₈H₂₇Br₂N₆: 607.07; found 607.16.


Example DSTL-3

[00470] Example DSTL-3 was prepared from Example DSTL-3, Step b according to the procedure described for the preparation of Example GW2. Purification was accomplished by preparative HPLC; 0% - 50%B over a 30 min gradient (at 40ml/min) using a Waters SunFire column (30 x 100 mm, S5) B = 90% CH₃CN - 10% H₂O - 0.1% TFA and A = 5% CH₃CN - 95% H₂O - 0.1%TFA. There was isolated Example DSTL-3 (29.0 mg) as a white solid. LC-MS (Cond.-D4): RT = 2.07 min. LC-MS Anal. Calcd. for [M+H]⁺ C₄₂H₅₀BrN₈O₆: 843.30; found 843.39.


Example DSTL-4

[00471] Example DSTL-4 was prepared from Example DSTL-4, Step b according to the procedure described for the preparation of Example GW2. Purification was accomplished by preparative HPLC; 0% - 50%B over a 30 min gradient at 40ml/min using a Waters SunFire column (30 x 100 mm, S5) B = 90% CH₃CN - 10% H₂O - 0.1% TFA and A = 5% CH₃CN - 95% H₂O - 0.1%TFA. There was isolated Example DSTL-4 (30.2 mg) as a white solid. ¹H NMR (MeOD, 500 MHz, δ): 7.82 (s, 8H), 5.08-5.05 (m, 2H), 4.58 (app d, *J* = 6.7 Hz, 2H), 3.76-3.74 (m, 2H), 3.67 (s, 6H), 2.57-2.52 (m, 2H), 2.50-2.44 (m, 2 H), 2.22-2.17 (m, 2H), 2.06-2.01 (m, 2H), 1.12-1.08 (m, 2H), 1.04 (d, *J* = 6.7 Hz, 6H), 0.95 (d, *J* = 6.7 Hz, 6H), 0.84-0.82 (m, 2H). LC-MS (Cond.-D4): RT = 2.56 min. LC-MS Anal. Calcd. for [M+H]⁺ C₄₂H₄₉BrN₈O₆: 921.31; found 921.30.

Example DSTL-5



Example DSTL-5, Step a




[00472] Accufluor (45-50% on alumina) (345 mg, 1.2 mmol) was added in one portion to a stirred suspension of Example 1, Step d, (1R,1'R,3S,3'S,5R,5'R)-*tert*-butyl 3,3'-(5,5'-(biphenyl-4,4'-diyl)bis(1H-imidazole-5,2-diyl))bis(2-azabicyclo[3.1.0]hexane-2-carboxylate) (500 mg, 0.77 mmol) in dry DMF (5 mL) at rt. The mixture was placed into a preheated oil bath at 60 °C and stirred 4 h before the solvent was removed by rotary evaporation under high vacuum. The crude product was charged to a Thompson 90 g silica gel cartridge and eluted 10 – 100 % B over 2 L, and hold 100% B for 1 L (Solvent B = EtOAc; Solvent A = hexanes). A second gradient was applied eluting 0 – 100 % B over 2 L (Solvent B = methanol; Solvent A = EtOAc). Three fractions were collected: Example DSTL-5, Step a (X=F), (1R,1'R,3S,3'S,5R,5'R)-*tert*-butyl 3,3'-(5,5'-(biphenyl-4,4'-diyl)bis(4-fluoro-1H-imidazole-5,2-diyl))bis(2-azabicyclo[3.1.0]hexane-2-carboxylate) (45.0 mg, 7 %) as a yellow solid; Example DSTL-5, Step a (X=H) ((1R,3S,5R)-*tert*-butyl 3-(5-(4'-(2-((1R,3S,5R)-2-(*tert*-butoxycarbonyl)-2-azabicyclo[3.1.0]hexan-3-yl)-1H-imidazol-5-yl)biphenyl-4-yl)-4-fluoro-1H-imidazol-2-yl)-2-azabicyclo[3.1.0]hexane-2-carboxylate (193 mg, 32 %) as a yellow solid, and recovered starting material (153 mg, 24 %).

[00473] Example DSTL-5, Step a X = F: A sample (~10 mg) of the first eluting compound was further purified by preparative HPLC (0 - 100% B) over a 30 min gradient (at 40ml/min) using a PHENOMENEX® Luna column (30 x 100 mm, 10u) where Solvent B = 90% CH₃CN - 10% H₂O - 0.1% TFA and A = 5% CH₃CN - 95% H₂O - 0.1% TFA. There was isolated (4.6 mg) as a light yellow solid. LC-MS (Cond.-D4): RT = 4.43 min. LC-MS Anal. Calcd. for [M+H]⁺ C₃₈H₄₃F₂N₆O₄: 685.33; found 685.38.

[00474] Example DSTL-5, Step a X = H: A sample (~15 mg) of the second eluting compound was further purified by preparative HPLC (0 - 100% B) over a 30 min gradient (at 40ml/min) using a PHENOMENEX® Luna column (30 x 100 mm, 10u) where Solvent B = 90% CH₃CN - 10% H₂O - 0.1% TFA and A = 5% CH₃CN - 95% H₂O - 0.1%

TFA. There was isolated (11.2 mg) as an off-white solid. LC-MS (Cond.-D4): RT = 4.43 min. LC-MS Anal. Calcd. for $[M+H]^+$ $C_{38}H_{44}FN_6O_4$: 667.34; found 667.40.

Example DSTL-5, Step b



5

[00475] The products of Example DSTL-5, Step a, (X = F) and (X = H), were brought forward separately. Removal of the protecting groups was performed as described in the procedure from Example DSTL-1, Step h.


[00476] Example DTSL-5, Step b (X = F): Yellow solid, 23.4 mg; LC-MS (Cond.-D4): RT = 2.89 min. LC-MS Anal. Calcd. for $[M+H]^+$ $C_{28}H_{27}F_2N_6$: 485.22; found 485.25.

[00477] Example DTSL-5, Step b (X = H): 18 mg (95% pure); LC-MS (Cond.-D4): RT = 2.55 min. LC-MS Anal. Calcd. for $[M+H]^+$ $C_{28}H_{28}FN_6$: 467.25; found 467.27.

Example DSTL-5


[00478] Example DSTL-5 was prepared from Example DSTL-5, Step b (X = H) according to the procedure described for the preparation of Example GW2. Purification was accomplished by preparative HPLC; 0% - 50%B over a 30 min gradient at 40ml/min using a Waters SunFire column (30 x 100 mm, S5) B = 90% CH_3CN - 10% H_2O - 0.1% TFA and A = 5% CH_3CN - 95% H_2O - 0.1%TFA. There was isolated Example DSTL-5 (37.3 mg, 39%) as a pale yellow solid. 1H NMR (MeOD, 500 MHz, δ): 7.88 (s, 1 H), 7.84 (d, $J=8.5$ Hz, 2H), 7.80 (d, $J=8.5$ Hz, 2H), 7.77 (d, $J=8.5$ Hz, 2H), 7.68 (d, $J=8.5$, 2H), 5.13 (t, $J=7.0$ Hz, 1H), 5.04 (t, $J=6.1$ Hz, 1H), 4.56 (d, $J=6.7$ Hz, 2H) 3.82 (br. s, 1H), 3.68-3.67 (m, 7H), 2.72-2.68 (m, 1H), 2.51-2.43 (m, 3H), 2.21-2.14 (m, 2H), 2.09 (br. s, 1H), 2.03 (br. s, 1H), 1.15-1.09 (m, 2H), 1.02 (d, $J=6.7$ Hz, 6H), 0.94 (t, $J=7.0$ Hz, 6H), 0.90 (br. s, 1H), 0.81 (br. s, 1H). LC-MS (Cond.-D4): RT = 3.45 min. LC-MS Anal. Calcd. for $[M+H]^+$ $C_{42}H_{50}FN_8O_6$: 781.39; found 781.54.

Example DSTL-6



[00479] Example DSTL-6 was prepared from Example DSTL-5, Step b (X = H) according to the procedure described for the preparation of Example GW2 except using (S)-2-(methoxycarbonylamino)-2-(tetrahydro-2H-pyran-4-yl)acetic acid instead of Cap-51. Purification was accomplished by preparative HPLC (0 - 50% B) over a 25 min gradient (at 40ml/min) using a Waters SunFire column (30 x 100 mm, S5) where Solvent B = 90% CH₃CN - 10% H₂O - 0.1% TFA and A = 5% CH₃CN - 95% H₂O - 0.1% TFA. There was isolated Example DSTL-6 (37.1 mg, 36%) as a white solid. LC-MS (Cond.-D4): RT = 3.17 min. LC-MS Anal. Calcd. for [M+H]⁺ C₄₈H₅₄FN₈O₆: 865.41; found 865.55.


Example DSTL-7



[00480] Example DSTL-7 was prepared from Example DSTL-5, Step b (X = F) according to the procedure described for the preparation of Example GW2 except using (S)-2-(methoxycarbonylamino)-2-(tetrahydro-2H-pyran-4-yl)acetic acid instead of Cap-51. Purification was accomplished by preparative HPLC (0 - 50% B) over a 25 min gradient (at 40ml/min) using a Waters SunFire column (30 x 100 mm, S5) where Solvent B = 90% CH₃CN - 10% H₂O - 0.1% TFA and A = 5% CH₃CN - 95% H₂O - 0.1% TFA. There was isolated Example DSTL-7 (14 mg, 25%) as a white solid. ¹H NMR (MeOD,


500 MHz, δ) 7.72 (d, $J=8.5$ Hz, 4H), 7.64 (d, $J=8.5$ Hz, 4H), 5.03 (t, $J=6.7$ Hz, 2H), 4.63 (d, $J=7.6$ Hz, 2H), 3.97-3.93 (m, 4H), 3.72 (br. s, 2H), 3.67 (s, 6H), 3.42-3.37 (m, 4H), 2.44-2.42 (m, 4H), 2.02 (br. s, 4H), 1.63-1.57 (m, 6H), 1.46-1.43 (m, 2H), 1.14-1.19 (m, 2H), 0.80 (br. s, 2H); LC-MS (Cond.-D4): RT = 3.17 min. LC-MS Anal. Calcd. for [M+H]⁺ C₄₆H₅₃F₂N₈O₆: 883.40; found 883.57.

Example DSTL-8



10

Example DSTL-8, Step a




[00481] Accufluor (45-50% on alumina) (274 mg, 0.92 mmol) was added in one portion to a stirred suspension of Example GW1, Step a, (2S,2'S,5S,5'S)-*tert*-butyl 5,5'-(5,5'-(biphenyl-4,4'-diyl)bis(1H-imidazole-5,2-diyl))bis(2-methylpyrrolidine-1-carboxylate) (400 mg, 0.6 mmol) in dry DMF (4 mL) at rt. The mixture was placed into a preheated oil bath at 60 °C and stirred 4 h before an additional 135 mg of Accufluor was added and stirring was continued at 60 °C for 16 h. The solvent was removed by rotary evaporation under high vacuum, and the crude product was charged to a Thompson 90 g silica gel cartridge and eluted 10 – 100 % B over 2 L, and hold 100% B for 1 L (Solvent B = EtOAc; Solvent A = hexanes). A second gradient was applied eluting 0 – 100 % B over 2 L (Solvent B = methanol; Solvent A = EtOAc). Three fractions were collected: Example DSTL-8, Step a (X = F), (2S,2'S,5S,5'S)-*tert*-butyl 5,5'-(5,5'-

(biphenyl-4,4'-diyl)bis(4-fluoro-1H-imidazole-5,2-diyl))bis(2-methylpyrrolidine-1-carboxylate) (33 mg, 5.1 %) as a yellow-orange solid LC-MS (Cond.-D4): RT = 4.64 min. LC-MS Anal. Calcd. for $[M+H]^+$ $C_{38}H_{47}F_2N_6O_4$: 689.36; found 689.50. Example DSTL-8, Step a (X = H) (2S,5S)-*tert*-butyl 2-(5-(4'-(2-((2S,5S)-1-(*tert*-butoxycarbonyl)-5-methylpyrrolidin-2-yl)-1H-imidazol-5-yl)biphenyl-4-yl)-4-fluoro-1H-imidazol-2-yl)-5-methylpyrrolidine-1-carboxylate (108.6 mg, 17 %) as a yellow-orange solid. LC-MS (Cond.-D4): RT = 3.83 min. LC-MS Anal. Calcd. for $[M+H]^+$ $C_{38}H_{48}FN_6O_4$: 671.37; found 671.48 and recovered starting material (321.2 mg, 80%) as a reddish-orange solid.

10

Example DSTL-8, Step b



[00482] The products of Example DSTL-8, Step a, (X = F) and (X = H), were brought forward separately. Removal of the protecting groups was performed as described in the procedure from Example DSTL-1, Step h.

15 **[00483]** Example DTSL-8, Step b (X = F): Yellow solid, 22.7 mg; LC-MS (Cond.-D4): RT = 3.03 min. LC-MS Anal. Calcd. for $[M+H]^+$ $C_{28}H_{30}F_2N_6$: 489.26; found 489.35.

[00484] Example DTSL-8, Step b (X = H): 35 mg (95% pure); LC-MS (Cond.-D4): RT = 2.63 min. LC-MS Anal. Calcd. for $[M+H]^+$ $C_{28}H_{31}FN_6$: 471.27; found 471.33.


20

Example DSTL-8

[00485] Example DSTL-5 was prepared from Example DSTL-8, Step b (X = H) according to the procedure described for the preparation of Example GW2. Purification was accomplished by preparative HPLC; 0% - 50%B over a 30 min gradient at 40ml/min using a Waters SunFire column (30 x 100 mm, S5) B = 90% CH_3CN - 10% H_2O - 0.1% TFA and A = 5% CH_3CN - 95% H_2O - 0.1%TFA. There was isolated Example DSTL-8 (31.9 mg, 41%) as a light peach-colored solid. LC-MS (Cond.-D4): RT = 3.66 min. LC-MS Anal. Calcd. for $[M+H]^+$ $C_{42}H_{54}FN_8O_6$: 785.42; found 785.48.


25

Example DSTL-9



- [00486] Example DSTL-9 was prepared from Example DSTL-8, Step b (X = H) according to the procedure described for the preparation of Example GW2 except using (S)-2-(methoxycarbonylamino)-2-(tetrahydro-2H-pyran-4-yl)acetic acid instead of Cap-51. Purification was accomplished by preparative HPLC (0 - 50% B) over a 25 min gradient (at 40ml/min) using a Waters SunFire column (30 x 100 mm, S5) where Solvent B = 90% CH₃CN - 10% H₂O - 0.1% TFA and A = 5% CH₃CN - 95% H₂O - 0.1% TFA. There was isolated Example DSTL-9 (27.2 mg, 32%) as a pale yellow solid. LC-MS (Cond.-D4): RT = 3.37 min. LC-MS Anal. Calcd. for [M+H]⁺ C₄₆H₅₈FN₈O₈: 869.44; found 869.54.

Example DSTL-10




- [00487] Example DSTL-10 was prepared from Example DSTL-8, Step b (X = F) according to the procedure described for the preparation of Example GW2 except using (S)-2-(methoxycarbonylamino)-2-(tetrahydro-2H-pyran-4-yl)acetic acid instead of Cap-51. Purification was accomplished by preparative HPLC (0 - 50% B) over a 25 min gradient (at 40ml/min) using a Waters SunFire column (30 x 100 mm, S5) where Solvent B = 90% CH₃CN - 10% H₂O - 0.1% TFA and A = 5% CH₃CN - 95% H₂O - 0.1% TFA.


There was isolated Example DSTL-10 (13.8 mg, 25%) as a pale yellow solid. LC-MS (Cond.-D4): RT = 4.28 min. LC-MS Anal. Calcd. for $[M+H]^+$ C₄₆H₅₇F₂N₈O₈: 887.43; found 887.50.

5

Example DSTL-11




Example DSTL-11, Step a



- 10 [00488] The preparation of DSTL-11, Step a, was conducted in analogous fashion to compound J5 in patent application 20080213 US 10889A USCIP. LC-MS (Cond.-D4): RT = 2.22 min. LC-MS Anal. Calcd. for $[M+H]^+$ C₂₂H₂₇BrN₃O₄: 476.12; found 478.17.

Example DSTL-11, Step b




15

- [00489] 5N sodium hydroxide (1.469 mL, 7.4 mmol) was added to a solution of DSTL-11, Step a (1R,3S,5R)-*tert*-butyl 3-(4-(4-bromophenyl)-5-(ethoxycarbonyl)-1H-imidazol-2-yl)-2-azabicyclo[3.1.0]hexane-2-carboxylate (1.75 g, 3.7 mmol) in ethanol (40 mL). The mixture was stirred at 50 °C for 16 h and additional 5N NaOH (1.5 mL) was added. The mixture was stirred at 60 °C for 6h and additional 5N NaOH (0.75 mL)
- 20

and stirring was continued at 60 °C for 16h. The ethanol was removed by rotary evaporation and the residue was taken up in ethyl acetate and neutralized to pH = 5 (1N HCl approx 20mL until close to pH = 7, then with pH =5 phosphate buffer). The organic layer was separated, washed with brine, dried and concentrated. There was isolated


- 5 Example DSTL-11, Step b (1.6 g, 92 %) as a white solid. LC-MS (Cond.-D4): RT = 1.78 min. LC-MS Anal. Calcd. for $[M+H]^+$ C₂₀H₂₂BrN₃O₄: 448.09; found 447.97.

Example DSTL-11, Step c



- 10 **[00490]** CDI (0.58 g, 3.57 mmol) was added in one portion to a stirred soln of DSTL-11, Step b, 4-(4-bromophenyl)-2-((1R,3S,5R)-2-(*tert*-butoxycarbonyl)-2-azabicyclo[3.1.0]hexan-3-yl)-1H-imidazole-5-carboxylic acid (1.6 g, 3.57 mmol) in dry THF (30 mL). The mixture was stirred at 50 °C for 4 h before being cooled to rt and NH₄OH (conc) (4.96 mL, 35.7 mmol) was added. The mixture was stirred for 16 h
- 15 before, diluted with EtOAc, washed with sat'd NaHCO₃ soln, brine, and dried. The crude product was charged to a Thompson 160 g silica gel cartridge and eluted 40 - 100% B over 3 L, followed by a hold at 100% B for 1L . (Solvent B = EtOAc; Solvent A = hexanes). There was isolated Example DSTL-11, Step c, (1.76 g 99%). LC-MS (Cond.-D4): RT = 1.75 min. LC-MS Anal. Calcd. for $[M+H]^+$ C₂₀H₂₄BrN₄O₃ 449.10; found:
- 20 449.06.

Example DSTL-11, Step d




- [00491]** 2,4,6-Trichloro-1,3,5-triazine (cyanuric chloride) (168 mg, 0.91 mmol) was
- 25 added in one portion to a stirred solution of Example DSTL-11, Step c, (1R,3S,5R)-*tert*-butyl 3-(4-(4-bromophenyl)-5-carbamoyl-1H-imidazol-2-yl)-2-azabicyclo[3.1.0]hexane-

2-carboxylate (903 mg, 1.82 mmol) in dry DMF (4 mL). The mixture was stirred at for 2 h and then warmed to 50 °C for 16 h. Additional cyanuric chloride (20mg) was added and the mixture was stirred further at 50 °C for 24 h, diluted with EtOAc and water, and the organic layer was washed with sat'd NaHCO₃ soln, brine, and dried (Na₂SO₄). The crude product was charged to a Thompson 90 g silica gel cartridge and eluted 25-100% B over 1.5 L, followed by a hold at 100% B 500 mL. (Solvent B = EtOAc; Solvent A = hexanes). There was isolated Example DSTL-11, Step d, (346 g 60%) as a yellowish-tan foam. LC-MS (Cond.-D4): RT = 3.93 min. LC-MS Anal. Calcd. for [M+H]⁺ C₂₀H₂₂BrN₄O₂ 431.09; found: 431.01.

10

Example DSTL-11, Step e




[00492] Tetrakis(triphenyl)palladium (45.8 mg, 0.04 mmol) was added to a argon purged suspension of Example DSTL-11, Step d, (1R,3S,5R)-*tert*-butyl 3-(5-(4-bromophenyl)-4-cyano-1H-imidazol-2-yl)-2-azabicyclo[3.1.0]hexane-2-carboxylate (340 mg, 0.79 mmol). (1R,3S,5R)-*tert*-butyl 3-(4-(4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)-1H-imidazol-2-yl)-2-azabicyclo[3.1.0]hexane-2-carboxylate (393 mg, 0.871 mmol), and NaHCO₃ (333 mg, 3.96 mmol) in DME (6 mL) and Water (1.5 mL) in a thick-walled, screw-topped tube. The reaction mixture was flushed with argon, sealed, and immersed in a pre-heated oil bath (80 °C) and stirred 14 h. The mixture was diluted with EtOAc, washed with sat'd NaHCO₃ soln, brine, dried over (Na₂SO₄). The crude product was charged to a Thompson 90 g silica gel cartridge and eluted with 35 – 100 % B over 1.5 L, followed by hold at 100 % B for 0.5 L. (Solvent B = EtOAc; Solvent A = hexanes). There was isolated Example DSTL-11, Step e, (200 mg, 36%) as a light yellow foam.

[00493] A small sample (~20 mg) was subject to further purification by prep. HPLC (0%B to 100%B over a 25 min gradient at 40ml/min) using a PHENOMENEX® Gemini column (30 x 100 mm, 10u) where Solvent B = 95% CH₃CN - 5% H₂O- 10mM NH₄OAc

and A = 5% CH₃CN - 95% H₂O - 10mM NH₄OAc. (12.9 mg) LC-MS (Cond.-D4): RT = 3.51 min. LC-MS Anal. Calcd. for [M+H]⁺ C₃₉H₄₄N₇O₄ 674.35; found: 674.29.

Example DSTL-11, Step f




[00494] The product of Example DSTL-11, Step f was obtained upon removal of the protecting groups as described in the procedure from Example DSTL-1, Step h. LC-MS (Cond.-D4): RT = 2.36 min. LC-MS Anal. Calcd. for [M+H]⁺ C₂₉H₂₈N₇: 474.24; found 474.13.

Example DSTL-11

[00495] Example DSTL-11 was prepared from Example DSTL-11, Step f according to the procedure described for the preparation of Example GW2. Purification was accomplished by preparative HPLC; 0% - 50%B over a 30 min gradient at 40ml/min using a Waters SunFire column (30 x 100 mm, S5) B = 90% CH₃CN - 10% H₂O - 0.1% TFA and A = 5% CH₃CN - 95% H₂O - 0.1%TFA. There was isolated Example DSTL-11 (21.5 mg, 31%) as an off white solid. ¹H NMR (MeOD, 500 MHz, δ): 7.93-7.82 (m, 9 H), 5.15-5.08 (m, 2H), 4.58 – 4.56 (m, 2H), 3.82 (br. s, 1H), 3.72 (br. s, 1H) 3.67 (s, 6H), 2.73-2.67 (m, 1H), 2.50-2.46 (m, 3H), 2.21-2.14 (m, 2H), 2.09 (br. s, 1H), 2.03 (br. s, 1H), 1.14-1.10 (m, 2H), 1.05 – 1.02 (m, 6H), 0.97-0.93 (m, 6H), 0.90 (br. s, 1H), 0.83 (br. s, 1H). LC-MS (Cond.-D4): RT = 3.25 min. LC-MS Anal. Calcd. for [M+H]⁺ C₄₃H₅₀N₉O₆: 780.39; found 788.31.

Example DSTL-12




[00496] Example DSTL-12 was prepared from Example DSTL-11, Step f according to the procedure described for the preparation of Example GW2 except using (S)-2-(methoxycarbonylamino)-2-(tetrahydro-2H-pyran-4-yl)acetic acid instead of Cap-51.


- 5 Purification was accomplished by preparative HPLC (0 - 50% B) over a 25 min gradient (at 40ml/min) using a Waters SunFire column (30 x 100 mm, S5) where Solvent B = 90% CH₃CN - 10% H₂O - 0.1% TFA and A = 5% CH₃CN - 95% H₂O - 0.1% TFA. There was isolated Example DSTL-12 (26.7 mg, 35%) as an off white solid. LC-MS (Cond.-D4): RT = 2.98 min. LC-MS Anal. Calcd. for [M+H]⁺ C₄₇H₅₄N₉O₈: 872.41; found 872.35.

10

Example JLR-1




Example JLR-1, Step a



15


[00497] For the synthesis of Example JLR-1, Step a, see the process patent application WO 2009/020825.

Example JLR-1, Step b



[00498] A solution of a 1 M bromine in AcOH (0.4 mL) was added to a solution of
 5 Example JLR-1, Step a (250 mg, 0.40 mmol) in acetic acid (15 mL) and stirred for 3 h. The reaction mixture was poured onto DCM (100 mL) and treated with sat'd NaHCO₃ soln and solid NaHCO₃, until pH = neutral. The organic phase was concentrated and dried under vacuum. The reaction gave a 1:1:1 mixture of starting material : mono
 bromide: dibromide which were separated upon application (DCM) to a Thompson SiO₂
 10 column (25 g). Elution (BIOTAGE®) by gradient 15 - 100% B over 500 mL gave Example JLR-1, Step b (180 mg). LC-MS (Cond.-J4): RT = 2.75 min. LC-MS Anal. Calcd. for [M+H]⁺ C₃₆H₄₄BrN₆O₄: 705.26; found 705.22.

Example JLR-1, Step c



15 [00499] A solution of 4N HCl in dioxane (4 mL) was added to Example JLR-1, Step b (38 mg, 0.054 mmol) in MeOH (4 mL) and stirred for 4 h. The solvent was removed under vacuum and the tetra HCl salt was dried under high vacuum. LC-MS (Cond.-J4): RT = 1.75 min. LC-MS Anal. Calcd. for [M+H]⁺ C₂₆H₂₈BrN₆: 503.16; found 503.26.


20

Example JLR-1

[00500] Example JLR-1 was prepared from Example JLR-1, Step c according to the procedure described for the preparation of Example GW2. Purification by preparative HPLC PHENOMENEX® Luna column (30 x 100 mm S10) running 25 min gradient
 25 from 15-100% B (at 40ml/min) where Solvent B = 90% MeOH - 10% H₂O - 0.1% TFA


and A = 5% MeOH - 95% H₂O - 0.1% TFA. LC-MS (Cond.-J4): RT = 2.61 min. LC-MS Anal. Calcd. for [M+H]⁺ C₄₀H₅₀BrN₈O₆: 819.31; found 819.23.

Example JLR-2




5

Example JLR-2, Step a



- [00501] Iodine (162 mg, 0.640 mmol) was added to a solution of Example JLR-1, Step a (400 mg, 0.640 mmol) and NaHCO₃ (161 mg, 1.921 mmol) in dioxane (15 mL) and water (15 mL) and the reaction stirred for 6 h, quenched with 10% Na₂S₂O₃ solution (40 mL) and diluted with EtOAc/THF (2:1). The organic phase was concentrated and the crude product was applied (DCM) to a Thompson silica gel column (40 g) and eluted with 35-100% B over 1 L (A/B DCM/EtOAc) to give a mixture of starting material, diiodide (255 mg), and Example JLR-2, Step a (46 mg, 9%). LC-MS (Cond.-J4): RT = 2.45 min. LC-MS Anal. Calcd. for [M+H]⁺ C₃₆H₄₄IN₆O₄: 751.24; found 751.18.

Example JLR-2, Step b



HCl salt

[00502] A solution of 4N HCl in dioxane (5 mL) was added to JLR-2, Step a (46 mg, 0.061 mmol) in MeOH (5 mL) and stirred for 4 h. The solvent was removed under vacuum and the tetra HCl salt was dried under high vacuum. LC-MS (Cond.-J4): RT = 1.70 min. LC-MS Anal. Calcd. for $[M+H]^+$ C₂₆H₂₈IN₆: 551.13; found 551.09.


5

Example JLR-2

[00503] Example JLR-2 was prepared from Example JLR-2, Step b according to the procedure described for the preparation of Example GW2. Purification by semi-prep HPLC; Dynamax 6A semi-prep C8 column; 5% - 95% B over 30 min (at 20ml/min) where Solvent B = 90% CH₃CN - 10% H₂O - 1% NH₄OAc and A = 5% CH₃CN - 95% H₂O - 1% NH₄OAc. LC-MS (Cond.-J4): RT = 2.35 min. LC-MS Anal. Calcd. for $[M+H]^+$ C₄₀H₅₀IN₈O₆: 865.29; found 865.31.


10

Example JLR-3



15

Example JLR-3, Step a




[00504] NCS (175 mg, 1.310 mmol) was added to a solution of Example DSTL-1, Step d (510 mg, 1.191 mmol) in DMF (10 mL) and stirred for 18 h at 50 °C. The solvent was removed by rotary evaporation under high vacuum, and the residue charged (DCM) to a Thomson silica gel cartridge (80 g). Gradient elution was performed from 25 - 100% B over 750 mL (A/B Hex/EtOAc) to give Example JLR-3, Step a (250 mg, 46%). ¹H NMR

20

(MeOD, 500 MHz, δ): 8.58 (s, 1H), 8.39 (d, J = 8.6 Hz, 1H), 7.97 (s, 1H), 7.93 (d, J = 7.6 Hz, 1H), 4.98 (s, 1H), 3.70 (s, 1H), 2.76-2.71 (m, 1H), 2.50-2.45 (m, 1H), 1.86 (s, 1H), 1.48/1.12 (s, 9H), 0.96-0.92 (m, 1H), 0.79-0.77 (m, 1H). LC-MS (Cond.-J4): RT = 3.13 min. LC-MS Anal. Calcd. for $[M+H]^+$ $C_{21}H_{22}BrClN_3O_2$: 464.06; found 464.05.

5


Example JLR-3, Step a.1



[00505] Example JLR-3, Step a.1 was prepared from Example DSTL-1, Step d.1 according to a procedure described in patent application WO 2008/021927 for the preparation of its desmethano analog.

10


Example JLR-3, Step b



[00506] Example JLR-3, Step b was prepared from Example JLR-3, Step a and Example JLR-3, Step a.1 according to the procedure described for the preparation Example DSTL-1, Step g to give Example JLR-3, Step b. LC-MS (Cond.-J4): RT = 2.72 min. LC-MS Anal. Calcd. for $[M+H]^+$ $C_{40}H_{44}ClN_6O_4$: 707.31; found 707.44.

15

Example JLR-3, Step c




20

[00507] Example JLR-3, Step c was prepared from Example JLR-3, Step b according to the procedure described for the preparation DSTL-3, Step a to give Example JLR-3,

Step c. LC-MS (Cond.-J4): RT = 3.30 min. LC-MS Anal. Calcd. for $[M+H]^+$
 $C_{40}H_{43}BrClN_6O_4$: 787.21; found 787.34.

Example JLR-3, Step d




HCl salt

[00508] Example JLR-3, Step d was prepared from Example JLR-3, Step c according to the procedure described in the procedure from Example JLR-1, Step b. LC-MS (Cond.-J4): RT = 2.37 min. LC-MS Anal. Calcd. for $[M+H]^+$ $C_{30}H_{27}BrClN_6$: 587.12; found 587.14.


Example JLR-3

[00509] Example JLR-3 was prepared from Example JLR-3, Step d according to the procedure described for the preparation of Example GW2. Purification by preparative HPLC PHENOMENEX® Luna column (30 x 100 mm S10) running 18 min gradient from 15-100% B (at 40ml/min) where Solvent B = 90% MeOH - 10% H₂O - 0.1% TFA and A = 5% MeOH - 95% H₂O - 0.1% TFA. ¹H NMR (MeOD, 500 MHz, δ): 8.69-8.68 (m, 1H), 8.52-8.50 (m, 1H), 8.18-8.16 (m, 1H), 7.98-7.96 (m, 1H), 7.91-7.90 (m, 4H), 5.34-5.34 (m, 1H), 5.09-5.06 (m, 1 H), 4.62-4.58 (m, 2H), 3.91 (br s, 1H), 3.76 (br s, 1H) 3.69 (s, 6H), 2.77-2.73 (m, 1H), 2.64-2.59 (m, 1H), 2.51-2.48 (m, 2H), 2.23-2.16 (m, 3H), 2.06 (br s, 1H), 1.17-1.12 (m, 2H), 1.06-0.92 (m, 13H), 0.83 (m, 1H). LC-MS (Cond.-J4): RT = 3.14 min. LC-MS Anal. Calcd. for $[M+H]^+$ $C_{44}H_{44}BrClN_8O_6$: 901.27; found 901.45.

Example JLR-4




Example JLR-4, Step a



- 5 **[00510]** Hunig's base (1.2 g, 9 mmol) was added to a solution of 2-bromo-1-(4-bromophenyl)ethanone (2.5 g, 8.99 mmol) and (2S,5S)-1-(*tert*-butoxycarbonyl)-5-methylpyrrolidine-2-carboxylic acid (1.9 g, 8.29 mmol) in acetonitrile (75 mL) and stirred for 6 h at 24 °C. The reaction mixture was concentrated and the residue taken up in EtOAc and washed with water. Concentration gave a white solid which was taken up
- 10 in xylene (90 mL). Ammonium acetate (4.23 g, 70.4 mmol) was added, and the solution was stirred in a screw-capped pressure vessel at 135 °C for 3.5 h. The reaction mixture was diluted with EtOAc (400 mL) and washed with sat'd NaHCO₃ soln and concentrated. The crude product was charged (DCM) to a 80g Thompson silica gel cartridge and gradient elution was performed from 15% to 100% B over 1 L (A/B Hex/EtOAc) gave
- 15 Example JLR-4, Step a (yield not determined). LC-MS (Cond.-J4): RT = 2.21 min. LC-MS Anal. Calcd. for [M+H]⁺ C₁₉H₂₅BrN₃O₂; 406.12; found 406.10. [Note: for the synthesis of the starting carboxylic acid, see U.S. Patent Application 2009/0068140).

Example JLR-4, Step b




20

- [00511]** NCS (161 mg, 1.21 mmol) was added to a solution of Example JLR-4, Step a (700 mg, 1.7 mmol) in DMF (10 mL) and stirred for 18 h at 50 °C. Additional NCS (50

mg) and the reaction continued for 6 h before removing the solvent by nitrogen purge. The crude product was charged (DCM) to a 40g Thompson silica gel cartridge, and gradient elution was performed from 15% to 100% B over 750 mL (A/B Hex/EtOAc) to give Example JLR-4, Step b (580 mg, 70 %). ¹H NMR (MeOD, 500 MHz, δ): 7.62 (app. s, 4H), 4.77 (br. s, 1H), 4.03 (br. s, 1H), 2.24 (br s, 2H), 2.16-2.11 (m, 1H) 1.79 (br. s, 1H), 1.47-1.25 (m, 12H). LC-MS (Cond.-J4): RT = 3.33 min. LC-MS Anal. Calcd. for [M+H]⁺ C₁₉H₂₄BrClN₃O₂: 440.08; found 440.0.


Example JLR-4, Step c



[00512] EEDQ (1.67 g, 6.75 mmol) was added to a solution of Example DSTL-1, Step c (1.6 g, 6.75 mmol) and (2S,5S)-1-(*tert*-butoxycarbonyl)-5-methylpyrrolidine-2-carboxylic acid (1.55 g, 6.75 mmol) in DCM (100 mL) and stirred for 6h. (Note: The dianiline was not completely soluble). The reaction mixture was diluted with DCM (1 vol) and washed with half sat'd NaHCO₃ soln. Concentration gave a solid (2.5 g). LC-MS (Cond.-J4): RT = 3.07 min. LC-MS Anal. Calcd. for [M+H]⁺ C₂₁H₂₇BrN₃O₃: 448.13; found 448.11.


[00513] The crude solid (2.5 g, 5.58 mmol) was taken up in AcOH (200 mL) and stirred for 18 h at 60 °C. Concentration under high vacuum removed the solvent. The residue was taken up in DCM, washed with sat'd NaHCO₃ soln, and concentrated. The residue was charged (DCM) to a 80 g Thompson silica gel cartridge and gradient elution was performed from 15% to 100% B over 750 mL. (A/B Hex/EtOAc) to give Example JLR-4, Step c (2.6 g). ¹H NMR (MeOD, 500 MHz, δ): 8.36-8.35 (m, 2H), 8.0 (d, *J* = 9 Hz, 1H), 7.91 (dd, *J* = 9, 2 Hz, 1H), 7.87 (d, *J* = 9 Hz, 1H), 5.31-5.28 (m, 1H), 4.17 (br. s, 1H), 2.59-2.56 (m, 1H), 2.39-2.31 (m, 2H) 1.86-1.83 (m, 1H), 1.52-1.19 (m, 12H). LC-MS (Cond.-J4): RT = 2.57 min. LC-MS Anal. Calcd. for [M+H]⁺ C₂₁H₂₅BrN₃O₂: 430.12; found 430.09.

Example JLR-4, Step d




[00514] Example JLR-4, Step d was prepared from Example JLR-4, Step c according to the procedure described for the preparation of Example DSTL-1 Step f. LC-MS (Cond.-J4): RT = 2.86 min. LC-MS Anal. Calcd. for $[M+H]^+$ $C_{27}H_{37}BN_3O_4$: 478.29; found 478.25.

Example JLR-4, Step e



[00515] Example JLR-4, Step e was prepared from Example JLR-4, Step d and Example JLR-4, Step b according to the procedure described for the preparation DSTL-1, Step g to give Example JLR-4, Step e. 1H NMR (MeOD, 500 MHz, δ): 8.53 (d, J = 8.6 Hz, 1H), 8.49 (s, 1H), 8.20 (dd, J = 8.6, 2 Hz, 2H), 7.97 (d, J = 8.6 Hz, 2H), 7.92 (d, J = 8.6 Hz, 2H), 7.88 (d, J = 8.9 Hz, 1H). 5.32 (t, J = 6.4 Hz, 1H), 4.85 (br. s, 1H), 4.20 (br. s, 1H), 4.06 (br. s, 1H). 2.59-2.58 (m, 1H), 2.44-2.28 (m, 4H), 2.19-2.15 (m, 1H). 1.89-1.84 (m, 1H), 1.83-1.79 (m, 1H), 1.54-1.20 (m, 24H). LC-MS (Cond.-J4): RT = 3.18 min. LC-MS Anal. Calcd. for $[M+H]^+$ $C_{40}H_{48}ClN_6O_4$: 711.34; found 711.31.

Example JLR-4, Step f



HCl salt


[00516] Example JLR-4, Step f was prepared from Example JLR-4, Step e according to the procedure described for the preparation Example JLR-1, Step b to give Example

JLR-4, Step f. LC-MS (Cond.-J4): RT = 2.19 min. LC-MS Anal. Calcd. for $[M+H]^+$ $C_{30}H_{32}ClN_6$: 511.24; found 711.23.

Example JLR-4


- 5 **[00517]** Example JLR-4 was prepared from Example JLR-4, Step f according to the procedure described for the preparation of Example GW2. Purification by preparative HPLC Luna Axia column (30 x 100 mm C18) running 25 min gradient from 15-100% B (at 40ml/min) where Solvent B = 90% MeOH - 10% H₂O - 0.1% TFA and A = 5% MeOH - 95% H₂O - 0.1% TFA. LC-MS (Cond.-J4): RT = 3.12 min. LC-MS Anal.
- 10 Calcd. for $[M+H]^+$ $C_{44}H_{54}ClN_8O_6$: 825.39; found 825.31.

Example JLR-5




- [00518]** Example JLR-5 was prepared from Example JLR-4, Step f according to the procedure described for the preparation of Example GW2 except using and (S)-2-(methoxycarbonylamino)-2-(tetrahydro-2H-pyran-4-yl)acetic acid. Purification by preparative HPLC Luna Axia column (30 x 100 mm C18) running 25 min gradient from 15-100% B (at 40ml/min) where Solvent B = 90% MeOH - 10% H₂O - 0.1% TFA and A = 5% MeOH - 95% H₂O - 0.1% TFA. LC-MS (Cond.-J4): RT = 2.85 min. LC-MS
- 20 Anal. Calcd. for $[M+H]^+$ $C_{48}H_{58}ClN_8O_6$: 909.41; found 909.39.

Example ZY1



Example ZY1, Step a



- [00519] HATU (151 mg, 0.40 mmol) was added to a solution of an HCl salt of 4,4'-bis(2-((1R,3S,5R)-2-azabicyclo[3.1.0]hexan-3-yl)-1H-imidazol-4-yl)biphenyl (Example 1e) (107 mg, 0.18 mmol) and (S)-2-(methoxycarbonylamino)-2-((S)-tetrahydro-2H-pyran-3-yl)acetic acid (cap-177a) (86 mg, 0.40 mmol) in DMF (1.5 mL) and DIPEA (0.25 mL, 1.4 mmol) and the mixture was stirred at rt for 2 h. The reaction mixture was diluted with MeOH and purified by prep HPLC (H₂O-MeOH with 0.1% TFA buffer) to yield a TFA salt of dimethyl (S,1S,1'S)-2,2'-((1R,1'R,3S,3'S,5R,5'R)-3,3'-(4,4'-(biphenyl-4,4'-diyl))bis(1H-imidazole-4,2-diyl))bis(2-azabicyclo[3.1.0]hexane-3,2-diyl))bis(2-oxo-1-((S)-tetrahydro-2H-pyran-3-yl)ethane-2,1-diyl)dicarbamate (148.5 mg) as light yellow solid. LC-MS retention time = 1.61 min; m/z = 847 [M+H]⁺. (Column: PHENOMENEX® Luna 3.0 x 50mm S10. Solvent A = 90% Water:10% Methanol: 0.1% TFA. Solvent B = 10% Water :90% Methanol: 0.1% TFA. Flow Rate = 4 mL/min. Start % B = 0. Final % B = 100. Gradient Time = 3 min. Wavelength = 220). ¹H NMR (400 MHz, MeOD) δ ppm 7.80 - 7.91 (m, 10 H), 5.13 (dd, J=9.3, 7.0 Hz, 2 H), 4.79 (d, J=8.3 Hz, 2 H), 3.84 - 3.97 (m, 2 H), 3.70 - 3.80 (m, 4 H), 3.68 (s, 6 H), 3.52 - 3.63 (m, 2 H), 3.34 - 3.42 (m, 2 H), 2.61 - 2.75 (m, 2 H), 2.43 - 2.54 (m, 2 H), 1.98 - 2.15 (m, 4 H), 1.67 - 1.84 (m, 4 H), 1.48 - 1.65 (m, 4 H), 0.98 - 1.16 (m, 2 H), 0.77 - 0.93 (m, 2 H).


Example ZY1

- [00520] NCS (8.2 mg, 0.061 mmol) was added to a solution of a TFA salt of dimethyl (S,1S,1'S)-2,2'-((1R,1'R,3S,3'S,5R,5'R)-3,3'-(4,4'-(biphenyl-4,4'-diyl))bis(1H-imidazole-4,2-diyl))bis(2-azabicyclo[3.1.0]hexane-3,2-diyl))bis(2-oxo-1-((S)-tetrahydro-2H-pyran-3-yl)ethane-2,1-diyl)dicarbamate (Example ZY1, Step a) (33 mg, 0.031 mmol) in DMF (1 mL) and the mixture was stirred at 50 °C for 16 h. The reaction mixture was diluted with MeOH, filtered and purified by prep HPLC (H₂O-MeOH with 0.1% TFA buffer) to yield a TFA salt of dimethyl (S,1S,1'S)-2,2'-((1R,1'R,3S,3'S,5R,5'R)-3,3'-(4,4'-(biphenyl-

4,4'-diyl)bis(5-chloro-1H-imidazole-4,2-diyl))bis(2-azabicyclo[3.1.0]hexane-3,2-diyl))bis(2-oxo-1-((S)-tetrahydro-2H-pyran-3-yl)ethane-2,1-diyl)dicarbamate, (Example ZY1) (25 mg) as white solid. LC-MS retention time = 1.92 min; m/z = 458 $[M/2+H]^+$. (Column: PHENOMENEX® Luna 3.0 x 50mm S10. Solvent A = 95% Water/ 5%


- 5 Methanol/10 mM Ammonium Acetate. Solvent B = 5% Water/ 95% Methanol/10 mM Ammonium Acetate. Flow Rate = 4 mL/min. Start % B = 0. Final % B = 100. Gradient Time = 2 min. Wavelength = 220). 1H NMR (400 MHz, MeOD) δ ppm 7.77 - 7.85 (m, 8 H), 5.03 (dd, $J=8.8, 6.0$ Hz, 2 H), 4.75 - 4.86 (m, 2 H), 3.71 - 3.89 (m, 6 H), 3.67 (s, 6 H), 3.36 - 3.62 (m, 4 H), 2.38 - 2.57 (m, 4 H), 1.94 - 2.12 (m, 4 H), 1.69 - 1.87 (m, 4 H), 1.46
- 10 - 1.65 (m, 4 H), 1.03 - 1.17 (m, 2 H), 0.77 - 0.85 (m, 2 H).

Example ZY2



15

Example ZY2, Step a




- [00521] HATU (140 mg, 0.37 mmol) was added to a solution of an HCl salt of 4,4'-bis(2-((1R,3S,5R)-2-azabicyclo[3.1.0]hexan-3-yl)-1H-imidazol-4-yl)biphenyl, (Example 1e) (100 mg, 0.17 mmol) and (S)-2-(methoxycarbonylamino)-2-((R)-tetrahydro-2H-pyran-3-yl)acetic acid (cap117b) (80 mg, 0.37 mmol) in DMF (1.5 mL) and DIPEA (0.23 mL, 1.3 mmol) and the mixture was stirred at rt for 3 h. Then the reaction mixture was
- 20 diluted with MeOH and purified by prep HPLC (H₂O-MeOH with 0.1% TFA buffer) to yield a TFA salt of dimethyl (R,1S,1'S)-2,2'-((1R,1'R,3S,3'S,5R,5'R)-3,3'-(4,4'-(biphenyl-4,4'-diyl)bis(1H-imidazole-4,2-diyl))bis(2-azabicyclo[3.1.0]hexane-3,2-diyl))bis(2-oxo-
- 25 1-((R)-tetrahydro-2H-pyran-3-yl)ethane-2,1-diyl)dicarbamate (Example ZY2, Step a) (76.2 mg) as white solid. LC-MS retention time = 1.53 min; m/z = 847 $[M+H]^+$. (Column: PHENOMENEX® Luna 3.0 x 50mm S10. Solvent A = 90% Water:10%

Methanol: 0.1% TFA. Solvent B = 10% Water:90% Methanol: 0.1% TFA. Flow Rate = 4 mL/min. Start % B = 0. Final % B = 100. Gradient Time = 3 min. Wavelength = 220). ¹H NMR (400 MHz, MeOD) δ ppm 7.75 - 7.96 (m, 10 H), 5.13 (dd, *J*=9.3, 7.0 Hz, 2 H), 4.60 (d, *J*=8.3 Hz, 2 H), 3.77 - 3.95 (m, 6 H), 3.67 (s, 6 H), 3.24 - 3.47 (m, 4 H), 2.62 - 2.76 (m, 2 H), 2.40 - 2.57 (m, 2 H), 1.97 - 2.19 (m, 4 H), 1.54 - 1.82 (m, 6 H), 1.36 - 1.54 (m, 2 H), 1.00 - 1.17 (m, 2 H), 0.79 - 0.93 (m, 2 H).


Example ZY2

[00522] NCS (5.2 mg, 0.039 mmol) was added to a solution of a TFA salt of dimethyl (R,1S,1'S)-2,2'-((1R,1'R,3S,3'S,5R,5'R)-3,3'-(4,4'-(biphenyl-4,4'-diyl)bis(1H-imidazole-4,2-diyl))bis(2-azabicyclo[3.1.0]hexane-3,2-diyl))bis(2-oxo-1-((R)-tetrahydro-2H-pyran-3-yl)ethane-2,1-diyl)dicarbamate (Example ZY2, Step a) (21 mg, 0.020 mmol) in DMF (1 mL) and the mixture was stirred at 50 °C for 16 h. Then the reaction mixture was diluted with MeOH, filtered and purified by prep HPLC (H₂O-MeOH with 0.1% TFA buffer) to yield a TFA salt of dimethyl (R,1S,1'S)-2,2'-((1R,1'R,3S,3'S,5R,5'R)-3,3'-(4,4'-(biphenyl-4,4'-diyl)bis(5-chloro-1H-imidazole-4,2-diyl))bis(2-azabicyclo[3.1.0]hexane-3,2-diyl))bis(2-oxo-1-((R)-tetrahydro-2H-pyran-3-yl)ethane-2,1-diyl)dicarbamate (Example ZY2) (10.8 mg) as white solid. LC-MS retention time = 1.92 min; *m/z* = 458 [M/2 + H]⁺. (Column: PHENOMENEX® Luna 3.0 x 50mm S10. Solvent A = 95% Water/ 5% Methanol/10 mM Ammonium Acetate. Solvent B = 5% Water/ 95% Methanol/10 mM Ammonium Acetate. Flow Rate = 4 mL/min. Start % B = 0. Final % B = 100. Gradient Time = 2 min. Wavelength = 220). ¹H NMR (400 MHz, MeOD) δ ppm 7.76 - 7.87 (m, 8 H), 5.04 (dd, *J*=8.8, 6.0 Hz, 2 H), 4.61 (d, *J*=8.3 Hz, 2 H), 3.75 - 3.95 (m, 6 H), 3.66 (s, 6 H), 3.35 - 3.48 (m, 4 H), 2.39 - 2.58 (m, 4 H), 1.97 - 2.16 (m, 4 H), 1.73 - 1.87 (m, 2 H), 1.40 - 1.73 (m, 6 H), 1.01 - 1.16 (m, 2 H), 0.81 (d, *J*=1.8 Hz, 2 H).

Example ZY3



Example ZY3, Step a




- [00523] HATU (82 mg, 0.22 mmol) was added to a solution of an HCl salt of 4,4'-bis(2-((1R,3S,5R)-2-azabicyclo[3.1.0]hexan-3-yl)-1H-imidazol-4-yl)biphenyl (Example 1e) (58.3 mg, 0.098 mmol) and (R)-2-(methoxycarbonylamino)-2-((R)-tetrahydro-2H-pyran-3-yl)acetic acid (cap 117c) (49 mg, 0.23 mmol) in DMF (1 mL) and DIPEA (0.14 mL, 0.79 mmol) and the mixture was stirred at rt for 3 h. Then the reaction mixture was diluted with MeOH, filtered and purified by prep HPLC (H₂O-MeOH with 0.1% TFA buffer) to yield a TFA salt of dimethyl (R,1R,1'R)-2,2'-((1R,1'R,3S,3'S,5R,5'R)-3,3'-(4,4'-(biphenyl-4,4'-diyl)bis(1H-imidazole-4,2-diyl))bis(2-azabicyclo[3.1.0]hexane-3,2-diyl))bis(2-oxo-1-((R)-tetrahydro-2H-pyran-3-yl)ethane-2,1-diyl)dicarbamate (Example ZY3, Step a) (79.8 mg) as white solid. LC-MS retention time = 1.63 min; m/z = 847 [M+H]⁺. (Column: PHENOMENEX® Luna 3.0 x 50mm S10. Solvent A = 90% Water:10% Methanol: 0.1% TFA. Solvent B = 10% Water:90% Methanol: 0.1% TFA. Flow Rate = 4 mL/min. Start % B = 0. Final % B = 100. Gradient Time = 3 min. Wavelength = 220).

Example ZY3

- [00524] NCS (6.7 mg, 0.050 mmol) was added to a solution of a TFA salt of dimethyl (R,1R,1'R)-2,2'-((1R,1'R,3S,3'S,5R,5'R)-3,3'-(4,4'-(biphenyl-4,4'-diyl)bis(1H-imidazole-4,2-diyl))bis(2-azabicyclo[3.1.0]hexane-3,2-diyl))bis(2-oxo-1-((R)-tetrahydro-2H-pyran-3-yl)ethane-2,1-diyl)dicarbamate (Example ZY3, Step a) (27 mg, 0.025 mmol) in DMF (1 mL) and the mixture was stirred at 50 °C for 16 h. Then the reaction mixture was diluted with MeOH, filtered and purified by prep HPLC (H₂O-MeOH with 0.1% TFA buffer) to yield a TFA salt of dimethyl (R,1R,1'R)-2,2'-((1R,1'R,3S,3'S,5R,5'R)-3,3'-(4,4'-(biphenyl-4,4'-diyl)bis(5-chloro-1H-imidazole-4,2-diyl))bis(2-azabicyclo[3.1.0]hexane-3,2-diyl))bis(2-oxo-1-((R)-tetrahydro-2H-pyran-3-yl)ethane-2,1-diyl)dicarbamate (Example ZY3) (15.8 mg) as white solid. LC-MS retention time = 1.93 min; m/z = 458 [M/2 + H]⁺. (Column: PHENOMENEX® Luna 3.0 x 50mm S10. Solvent A = 95% Water/ 5% Methanol/10 mM Ammonium Acetate. Solvent B = 5% Water/ 95%


Methanol/10 mM Ammonium Acetate. Flow Rate = 4 mL/min. Start % B = 0. Final % B = 100. Gradient Time = 2 min. Wavelength = 220).

Example ZY4



5

Example ZY4, Step a




- [00525] HATU (75 mg, 0.20 mmol) was added to a solution of an HCl salt of 4,4'-bis(2-((1R,3S,5R)-2-azabicyclo[3.1.0]hexan-3-yl)-1H-imidazol-4-yl)biphenyl (Example 1e) (53.5 mg, 0.090 mmol) and (R)-2-(methoxycarbonylamino)-2-((S)-tetrahydro-2H-pyran-3-yl)acetic acid (cap117d) (45 mg, 0.21 mmol) in DMF (1 mL) and DIPEA (0.13 mL, 0.72 mmol) and the mixture was stirred at rt for 3 h. Then the reaction mixture was diluted with MeOH, filtered and purified by prep HPLC (H₂O-MeOH with 0.1% TFA buffer) to yield a TFA salt of dimethyl (S,1R,1'R)-2,2'-((1R,1'R,3S,3'S,5R,5'R)-3,3'-(4,4'-(biphenyl-4,4'-diyl)bis(1H-imidazole-4,2-diyl))bis(2-azabicyclo[3.1.0]hexane-3,2-diyl))bis(2-oxo-1-((S)-tetrahydro-2H-pyran-3-yl)ethane-2,1-diyl)dicarbamate (Example ZY4, Step a) (78.5 mg) as white solid. LC-MS retention time = 1.57 min; m/z = 847 [M+H]⁺. (Column: PHENOMENEX® Luna 3.0 x 50mm S10. Solvent A = 90% Water:10% Methanol: 0.1% TFA. Solvent B = 10% Water :90% Methanol: 0.1% TFA. Flow Rate = 4 mL/min. Start % B = 0. Final % B = 100. Gradient Time = 3 min. Wavelength = 220).

Example ZY4

- [00526] NCS (9.2 mg, 0.069 mmol) was added to a solution of a TFA salt of dimethyl (S,1R,1'R)-2,2'-((1R,1'R,3S,3'S,5R,5'R)-3,3'-(4,4'-(biphenyl-4,4'-diyl)bis(1H-imidazole-4,2-diyl))bis(2-azabicyclo[3.1.0]hexane-3,2-diyl))bis(2-oxo-1-((S)-tetrahydro-2H-pyran-


- 3-yl)ethane-2,1-diyl)dicarbamate (Example ZY4, Step a) (37 mg, 0.034 mmol) in DMF (1 mL) and the mixture was stirred at 50 °C for 16 h. Then the reaction mixture was diluted with MeOH, filtered and purified by prep HPLC (H₂O-MeOH with 0.1% TFA buffer) to yield a TFA salt of dimethyl (S,1R,1'R)-2,2'-((1R,1'R,3S,3'S,5R,5'R)-3,3'-(4,4'-(biphenyl-4,4'-diyl)bis(5-chloro-1H-imidazole-4,2-diyl))bis(2-azabicyclo[3.1.0]hexane-3,2-diyl))bis(2-oxo-1-((S)-tetrahydro-2H-pyran-3-yl)ethane-2,1-diyl)dicarbamate (Example ZY4) (25 mg) as white solid. LC-MS retention time = 1.93 min; m/z = 458 [M/2 + H]⁺. (Column: PHENOMENEX® Luna 3.0 x 50mm S10. Solvent A = 95% Water/ 5% Methanol/10 mM Ammonium Acetate. Solvent B = 5% Water/ 95% Methanol/10 mM Ammonium Acetate. Flow Rate = 4 mL/min. Start % B = 0. Final % B = 100. Gradient Time = 2 min. Wavelength = 220).

Example ZY5



- 15 [00527] NCS (5.3 mg, 0.039 mmol) was added to a solution of a TFA salt of dimethyl (S,1S,1'S)-2,2'-((1R,1'R,3S,3'S,5R,5'R)-3,3'-(4,4'-(biphenyl-4,4'-diyl)bis(1H-imidazole-4,2-diyl))bis(2-azabicyclo[3.1.0]hexane-3,2-diyl))bis(2-oxo-1-((S)-tetrahydro-2H-pyran-3-yl)ethane-2,1-diyl)dicarbamate (Example ZY1, Step a) (42.4 mg, 0.039 mmol) in DMF (1 mL) and the mixture was stirred at 50 °C for 16 h. Then the reaction mixture was
- 20 diluted with MeOH, filtered and purified by prep HPLC (H₂O-MeOH with 0.1% TFA buffer) to yield Example ZY5 as a TFA salt (11.7 mg, white solid). LC-MS retention time = 2.04 min; m/z = 881 [M+H]⁺. (Column: PHENOMENEX® Luna 3.0 x 50mm S10. Solvent A = 90% Water:10% Methanol: 0.1% TFA. Solvent B = 10% Water:90% Methanol: 0.1% TFA. Flow Rate = 4 mL/min. Start % B = 0. Final % B = 100. Gradient
- 25 Time = 3 min. Wavelength = 220). ¹H NMR (400 MHz, MeOD) δ ppm 7.76 - 7.92 (m, 9 H), 5.13 (dd, J=9.2, 6.9 Hz, 1 H), 5.04 (dd, J=8.0, 6.5 Hz, 1 H), 4.74 - 4.80 (m, 2 H), 3.71 - 3.94 (m, 6 H), 3.68 (s, 3 H), 3.67 (s, 3 H), 3.36 - 3.63 (m, 4 H), 2.63 - 2.75 (m, 1 H), 2.40 - 2.55 (m, 3 H), 1.95 - 2.15 (m, 4 H), 1.67 - 1.86 (m, 4 H), 1.45 - 1.65 (m, 4 H), 1.03 - 1.18 (m, 2 H), 0.84 - 0.91 (m, 1 H), 0.75 - 0.83 (m, 1 H).


Example ZY6




[00528] NCS (1.1 mg, 8.4 μ mol) was added to a solution of a TFA salt of dimethyl
 5 (R,1S,1'S)-2,2'-((1R,1'R,3S,3'S,5R,5'R)-3,3'-(4,4'-(biphenyl-4,4'-diyl)bis(1H-imidazole-
 4,2-diyl))bis(2-azabicyclo[3.1.0]hexane-3,2-diyl))bis(2-oxo-1-((R)-tetrahydro-2H-pyran-
 3-yl)ethane-2,1-diyl)dicarbamate (Example ZY2, Step a) (8.1 mg, 8.4 μ mol) in DMF (1
 mL) and the mixture was stirred at 50 °C for 16 h. Then the reaction was diluted with
 MeOH, filtered and purified by prep HPLC (H₂O-MeOH with 0.1% TFA buffer) to yield
 10 a TFA salt of Example ZY6 (2.2 mg) as white solid. LC-MS retention time = 2.01 min;
 m/z = 881 [M+H]⁺. (Column: PHENOMENEX® Luna 3.0 x 50mm S10. Solvent A =
 90% Water:10% Methanol: 0.1% TFA. Solvent B = 10% Water:90% Methanol: 0.1%
 TFA. Flow Rate = 4 mL/min. Start % B = 0. Final % B = 100. Gradient Time = 3 min.
 Wavelength = 220). ¹H NMR (400 MHz, MeOD) δ ppm 7.77 - 7.90 (m, 9 H), 5.12 (dd,
 15 J=9.2, 6.9 Hz, 1 H), 5.05 (dd, J=7.9, 6.1 Hz, 1 H), 4.58 - 4.96 (m, 2 H), 3.71 - 3.94 (m, 6
 H), 3.68 (s, 3 H), 3.66 (br. s., 3 H), 3.34 - 3.46 (m, 4 H), 2.64 - 2.75 (m, 1 H), 2.40 - 2.54
 (m, 3 H), 1.96 - 2.16 (m, 4 H), 1.40 - 1.85 (m, 8 H), 1.04 - 1.18 (m, 2 H), 0.84 - 0.92 (m, 1
 H), 0.79 (m, 1 H).

20

Example ZY7



Example ZY7, Step a



[00529] HATU (95 mg, 0.25 mmol) was added to a solution of an HCl salt of 4,4'-bis(2-((1R,3S,5R)-2-azabicyclo[3.1.0]hexan-3-yl)-1H-imidazol-4-yl)biphenyl (Example 1e) (67.6 mg, 0.114 mmol) and (S)-4-methoxy-2-(methoxycarbonylamino)butanoic acid (Example ZY7, Step a) (50 mg, 0.26 mmol) in DMF (1 mL) and DIPEA (0.16 mL, 0.91 mmol) and the mixture was stirred at rt for 3 h. The reaction mixture was diluted with MeOH, filtered and purified by prep HPLC (H₂O-MeOH with 0.1% TFA buffer) to yield a TFA salt of dimethyl (2S,2'S)-1,1'-((1R,1'R,3S,3'S,5R,5'R)-3,3'-(4,4'-(biphenyl-4,4'-diyl)bis(1H-imidazole-4,2-diyl))bis(2-azabicyclo[3.1.0]hexane-3,2-diyl))bis(4-methoxy-1-oxobutane-2,1-diyl)dicarbamate (Example ZY7, Step a) (96.4 mg) as white solid. LC-MS retention time = 1.53 min; m/z 795 [M+H]⁺. (Column: PHENOMENEX® Luna 3.0 x 50mm S10. Solvent A = 90% Water:10% Methanol: 0.1% TFA. Solvent B = 10% Water:90% Methanol: 0.1% TFA. Flow Rate = 4 mL/min. Start % B = 0. Final % B = 100. Gradient Time = 3 min. Wavelength = 220). ¹H NMR (400 MHz, MeOD) δ ppm 7.78 - 7.97 (m, 10 H), 5.16 (dd, J=9.2, 6.7 Hz, 2 H), 4.76 - 4.83 (m, 2 H), 3.75 - 3.84 (m, 2 H), 3.66 (s, 6 H), 3.45 - 3.56 (m, 4 H), 3.32 (s, 6 H), 2.68 (dd, J=13.4, 9.4 Hz, 2 H), 2.42 - 2.53 (m, 2 H), 2.01 - 2.23 (m, 4 H), 1.80 - 1.95 (m, 2 H), 1.05 - 1.18 (m, 2 H), 0.78 - 0.98 (m, 2 H).

20


Example ZY7

[00530] NCS (9.7 mg, 0.073 mmol) was added to a solution of a TFA salt of dimethyl (2S,2'S)-1,1'-((1R,1'R,3S,3'S,5R,5'R)-3,3'-(4,4'-(biphenyl-4,4'-diyl)bis(1H-imidazole-4,2-diyl))bis(2-azabicyclo[3.1.0]hexane-3,2-diyl))bis(4-methoxy-1-oxobutane-2,1-diyl)dicarbamate (Example ZY7, Step a) (29 mg, 0.036 mmol) in DMF (1 mL) and the mixture was stirred at 50 °C for 16 h. The reaction mixture was diluted with MeOH, filtered and purified by prep HPLC (H₂O-MeOH with 0.1% TFA buffer) to yield a TFA salt of dimethyl (2S,2'S)-1,1'-((1R,1'R,3S,3'S,5R,5'R)-3,3'-(4,4'-(biphenyl-4,4'-diyl)bis(5-chloro-1H-imidazole-4,2-diyl))bis(2-azabicyclo[3.1.0]hexane-3,2-diyl))bis(4-methoxy-1-oxobutane-2,1-diyl)dicarbamate (Example ZY7) (7.9 mg) as white solid. LC-MS

retention time = 2.51 min; m/z = 863 $[M+H]^+$. (Column: PHENOMENEX® Luna 3.0 x 50mm S10. Solvent A = 90% Water:10% Methanol: 0.1% TFA. Solvent B = 10% Water:90% Methanol: 0.1% TFA. Flow Rate = 4 mL/min. Start % B = 0. Final % B = 100. Gradient Time = 3 min. Wavelength = 220). 1H NMR (400 MHz, MeOD) δ ppm

7.82 (s, 8 H), 5.07 (dd, J =8.5, 5.8 Hz, 2 H), 4.75 - 4.83 (m, 2 H), 3.70 - 3.76 (m, 2 H), 3.66 (s, 6 H), 3.44 - 3.54 (m, 4 H), 3.32 (s, 6 H), 2.39 - 2.57 (m, 4 H), 1.98 - 2.21 (m, 4 H), 1.82 - 1.94 (m, 2 H), 1.05 - 1.19 (m, 2 H), 0.78 - 0.92 (m, 2 H).

Example ZY8



[00531] NCS (5.6 mg, 0.042 mmol) was added to a solution of a TFA salt of dimethyl (2S,2'S)-1,1'-((1R,1'R,3S,3'S,5R,5'R)-3,3'-(4,4'-(biphenyl-4,4'-diyl)bis(1H-imidazole-4,2-diyl))bis(2-azabicyclo[3.1.0]hexane-3,2-diyl))bis(4-methoxy-1-oxobutane-2,1-diyl)dicarbamate (Example ZY7, Step b) (33.3 mg, 0.042 mmol) in DMF (1 mL) and the mixture was stirred at 50 °C for 16 h. The reaction mixture was diluted with MeOH, filtered and purified by prep HPLC (H₂O-MeOH with 0.1% TFA buffer) to yield a TFA salt of Example ZY8 (9.4 mg) as a white solid. LC-MS retention time = 1.99 min; m/z = 829 $[M+H]^+$. (Column: PHENOMENEX® Luna 3.0 x 50mm S10. Solvent A = 90% Water:10% Methanol: 0.1% TFA. Solvent B = 10% Water:90% Methanol: 0.1% TFA. Flow Rate = 4 mL/min. Start % B = 0. Final % B = 100. Gradient Time = 3 min. Wavelength = 220). 1H NMR (400 MHz, MeOD) δ ppm 7.77 - 7.91 (m, 9 H), 5.15 (dd, J =9.2, 6.7 Hz, 1 H), 5.07 (t, J =7.0 Hz, 1 H), 4.77 - 4.83 (m, 2 H), 3.76 - 3.83 (m, 1 H), 3.68 - 3.75 (m, 1 H), 3.66 (s, 6 H), 3.44 - 3.56 (m, 4 H), 3.33 (s, 6 H), 2.63 - 2.75 (m, 1 H), 2.41 - 2.56 (m, 3 H), 1.97 - 2.23 (m, 4 H), 1.80 - 1.95 (m, 2 H), 1.06 - 1.19 (m, 2 H), 0.87 - 0.96 (m, 1 H), 0.84 (m, 1 H).

BIOLOGICAL ACTIVITY

[00532] An HCV Replicon assay was utilized in the present disclosure, and was prepared, conducted and validated as described in commonly owned

PCT/US2006/022197 and in O'Boyle et al., *Antimicrob. Agents Chemother.*, 49(4):1346-1353 (Apr. 2005). Assay methods incorporating luciferase reporters have also been used as described (Apath.com).

[00533] HCV-neo replicon cells and replicon cells containing resistance substitutions
 5 in the NS5A region were used to test the currently described family of compounds. The compounds were determined to have differing degrees of reduced inhibitory activity on cells containing mutations vs. the corresponding inhibitory potency against wild-type cells. Thus, the compounds of the present disclosure can be effective in inhibiting the function of the HCV NS5A protein and are understood to be as effective in combinations
 10 as previously described in application PCT/US2006/022197 and commonly owned WO 04/014852. It should be understood that the compounds of the present disclosure can inhibit multiple genotypes of HCV. Table 2 shows the EC₅₀ (Effective 50% inhibitory concentration) values of representative compounds of the present disclosure against the HCV 1b genotype. In one embodiment, compounds of the present disclosure are
 15 inhibitory versus 1a, 1b, 2a, 2b, 3a, 4a, and 5a genotypes. EC₅₀ values against HCV 1b are as follows. EC₅₀ values against HCV 1b are as follows: A = 0.2 pM -<1 pM; B = 1 pM - <10 pM; and C = 10 pM -< 100 pM; and D = 100 pM-0.15 μM.

[00534] The compounds of the present disclosure may inhibit HCV by mechanisms in addition to or other than NS5A inhibition. In one embodiment the compounds of the
 20 present disclosure inhibit HCV replicon and in another embodiment the compounds of the present disclosure inhibit NS5A.

Table 2

Example	EC ₅₀ (μM)	EC ₅₀ -Range
M1		B
M2		B
M4		B
M5		B
M6	4.08E-06	B
M7	2.13E-06	B
V1		B

Example	EC ₅₀ (μM)	EC ₅₀ -Range
V2	8.88E-05	C
V3		B
V4	3.22E-05	C
V5	6.58E-07	A
V5.1	3.90E-06	B
V5.2	3.06E-05	C
V5.3	1.33E-04	D
V6		A
V7		B
V8		B
V9	2.70E-06	B
V10		B
V11		B
V12		B
V13		B
V14		B
V15		C
V16	9.66E-07	A
GW1-1		B
GW1-2		B
GW1-3		B
GW2		B
GW2-1	1.27E-05	C
GW3		B
GW4		C
GW5		B
GW6		B
GW7		B
GW8		B
GW9		A

Example	EC ₅₀ (μM)	EC ₅₀ -Range
GW10		B
GW11		A
GW12		A
OL1	3.06E-06	B
OL2		B
OL3		B
OL4		B
OL5		B
OL6		B
OL7		B
OL8	2.09E-04	D
OL9	6.08E-06	B
OL10		D
OL11		B
OL12		C
OL13		B
OL14		C
OL15		B
OL16		C
OL17		B
OL18		B
OL19		C
OL20		D
OL21		C
OL22	4.08E-04	D
OL23		D
OL24		B
OL25		D
DSTL-1		A
DSTL-2		C

Example	EC ₅₀ (μM)	EC ₅₀ -Range
DSTL-3	1.25E-06	B
DSTL-4		A
DSTL-5	5.59E-07	A
DSTL-6		B
DSTL-7		B
DSTL-8		A
DSTL-9		B
DSTL-10		B
DSTL-11		A
DSTL-12		C
JLR-1		B
JLR-2		B
JLR-3		A
JLR-4	7.69E-07	A
JLR-5		B
ZY1		B
ZY2		B
ZY3		B
ZY4		B
ZY5		B
ZY6	1.08E-05	C
ZY7		B
ZY8		B


[00535] It will be evident to one skilled in the art that the present disclosure is not limited to the foregoing illustrative examples, and that it can be embodied in other specific forms without departing from the essential attributes thereof. It is therefore
5 desired that the examples be considered in all respects as illustrative and not restrictive, reference being made to the appended claims, rather than to the foregoing examples, and

all changes which come within the meaning and range of equivalency of the claims are therefore intended to be embraced therein.

CLAIMS

WHAT IS CLAIMED IS:

- 5 1. A compound of Formula (I)



(I),

or a pharmaceutically acceptable salt thereof, wherein:

- n is 0, 1, or 2;
- 10 X is selected from hydrogen, alkenyl, cyano, cycloalkyl, (cycloalkyl)alkyl, halo, and heterocyclyl;
- R¹ is selected from hydrogen and halo;
- R² is selected from hydrogen, alkenyl, cyano, cycloalkyl, (cycloalkyl)alkyl, halo, and heterocyclyl; or
- 15 R¹ and R², together with the carbon atoms to which they are attached, form a six-membered aromatic ring optionally substituted with one halo group;
- provided that at least one of X and R² is selected from alkenyl, cyano, cycloalkyl, (cycloalkyl)alkyl, halo, and heterocyclyl;
- each R³ is alkyl, wherein the alkyl can optionally form a fused three- or four-
- 20 membered ring with an adjacent carbon atom or a spirocyclic three- or four-membered ring with the carbon atom to which it is attached; wherein the fused and spirocyclic rings are optionally substituted with one or two alkyl groups;
- each R⁴ is independently selected from hydrogen and -C(O)R⁵; and
- each R⁵ is independently selected from alkoxy, alkyl, arylalkoxy, arylalkyl,
- 25 cycloalkyl, heterocyclyl, heterocyclylalkyl, (NR^cR^d)alkenyl, and (NR^cR^d)alkyl.

2. A compound of claim 1, or a pharmaceutically acceptable salt thereof, wherein each R⁵ is independently selected from alkoxy, heterocyclyl, and (NR^cR^d)alkyl.

3. A compound of claim 1, or a pharmaceutically acceptable salt thereof,
wherein X is halo.


4. A compound of claim 3, or a pharmaceutically acceptable salt thereof,
5 wherein R² is halo.

5. A compound of claim 1, or a pharmaceutically acceptable salt thereof,
wherein R¹ and R², together with the carbon atoms to which they are attached, form a six-
membered aromatic ring optionally substituted with one halo group.



10






6. A compound of claim 1, or a pharmaceutically acceptable salt thereof,
wherein X is hydrogen.

7. A compound selected from:










15










5
















5













5


























































































5






5







;




;



;



; and




;

5

or a pharmaceutically acceptable salt thereof.

8. A compound selected from




Example OL8-OL10

Example OL11-OL14














Example		R ₂
OL8		H
OL9		
OL10		
OL11		H
OL12		H
OL13		
OL14		

;



Example OL20-OL23

Example OL24-OL25

Example		R ₂
OL20		
OL21		
OL22		
OL23		
OL24		 ; and
OL25		

;

or a pharmaceutically acceptable salt thereof.

9. A composition comprising a compound of claim 1, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

5

10. The composition of claim 9 further comprising at least one additional compound having anti-HCV activity.

11. The composition of claim 10 wherein at least one of the additional compounds is an interferon or a ribavirin.

10

12. The composition of claim 11 wherein the interferon is selected from interferon alpha 2B, pegylated interferon alpha, consensus interferon, interferon alpha 2A, and lymphoblastoid interferon tau.

5 13. The composition of claim 10 wherein at least one of the additional compounds is selected from interleukin 2, interleukin 6, interleukin 12, a compound that enhances the development of a type 1 helper T cell response, interfering RNA, anti-sense RNA, Imiquimod, ribavirin, an inosine 5'-monophosphate dehydrogenase inhibitor, amantadine, and rimantadine.

10

14. The composition of claim 10 wherein at least one of the additional compounds is effective to inhibit the function of a target selected from HCV metalloprotease, HCV serine protease, HCV polymerase, HCV helicase, HCV NS4B protein, HCV entry, HCV assembly, HCV egress, HCV NS5A protein, and IMPDH for
15 the treatment of an HCV infection.

15. A method of treating an HCV infection in a patient, comprising administering to the patient a therapeutically effective amount of a compound of claim 1, or a pharmaceutically acceptable salt thereof.

20

16. The method of claim 15 further comprising administering at least one additional compound having anti-HCV activity prior to, after or simultaneously with the compound of claim 1, or a pharmaceutically acceptable salt thereof.

25 17. The method of claim 16 wherein at least one of the additional compounds is an interferon or a ribavirin.

18. The method of claim 17 wherein the interferon is selected from interferon alpha 2B, pegylated interferon alpha, consensus interferon, interferon alpha 2A, and
30 lymphoblastoid interferon tau.

19. The method of claim 16 wherein at least one of the additional compounds is selected from interleukin 2, interleukin 6, interleukin 12, a compound that enhances the development of a type 1 helper T cell response, interfering RNA, anti-sense RNA, Imiqimod, ribavirin, an inosine 5'-monophosphate dehydrogenase inhibitor, amantadine,
5 and rimantadine.

20. The method of claim 16 wherein at least one of the additional compounds is effective to inhibit the function of a target selected from HCV metalloprotease, HCV serine protease, HCV polymerase, HCV helicase, HCV NS4B protein, HCV entry, HCV
10 assembly, HCV egress, HCV NS5A protein, and IMPDH for the treatment of an HCV infection.

INTERNATIONAL SEARCH REPORT

International application No

PCT/US2010/060077

A. CLASSIFICATION OF SUBJECT MATTER

INV. C07D403/14 C07D405/14
ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X,P	WO 2010/062821 A1 (GLAXOSMITHKLINE LLC [US]; SCHMITZ FRANK ULRICH [US]; RAI ROOPA [US]; R) 3 June 2010 (2010-06-03) claim 1	1-20
X,P	----- WO 2010/111673 A1 (PRESIDIO PHARMACEUTICALS INC [US]; ZHONG MIN [US]; LI LEPING [US]) 30 September 2010 (2010-09-30) claim 1	1-20
A	----- WO 2009/102325 A1 (SQUIBB BRISTOL MYERS CO [US]; BACHAND CAROL [CA]; BELEMA MAKONEN [US];) 20 August 2009 (2009-08-20) claims 1, 3, 15 page 2, lines 21-29 page 3, lines 1-20 pages 512-537; examples J.1-J51 ----- -/-	1-20

☒ Further documents are listed in the continuation of Box C.

☒ See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

28 January 2011

Date of mailing of the international search report

07/02/2011

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Guazzelli, Giuditta

INTERNATIONAL SEARCH REPORT

International application No

PCT/US2010/060077

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 2009/202478 A1 (BACHAND CAROL [CA] ET AL) 13 August 2009 (2009-08-13) claims 1, 9-11, 17 -----	1-20

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2010/060077

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2010062821 A1	03-06-2010	NONE	
WO 2010111673 A1	30-09-2010	NONE	
WO 2009102325 A1	20-08-2009	AU 2008350334 A1 CA 2715839 A1 EP 2242752 A1	20-08-2009 20-08-2009 27-10-2010
US 2009202478 A1	13-08-2009	AR 070366 A1 AU 2009214997 A1 CA 2715559 A1 EP 2245027 A1 KR 20100124290 A PE 13812009 A1 WO 2009102633 A1	31-03-2010 20-08-2009 20-08-2009 03-11-2010 26-11-2010 18-09-2009 20-08-2009