



(51) International Patent Classification:

C07K 16/30 (2006.01) **A61P 35/00** (2006.01)
A61K 39/395 (2006.01)

(21) International Application Number:

PCT/US2009/049139

(22) International Filing Date:

30 June 2009 (30.06.2009)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/077,041 30 June 2008 (30.06.2008) US
61/097,034 15 September 2008 (15.09.2008) US

(71) Applicant (for all designated States except US): **MORPHOTEK, INC.** [US/US]; 210 Welsh Pool Road, Exton, PA 19341 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **NICOLAIDES, Nicholas, C.** [US/US]; 4 Cider Mill Court, Garnett Valley, PA 19061 (US). **SASS, Philip, M.** [US/US]; 1903 Black Hawk Circle, Audubon, PA 19403 (US). **GRASSO, Luigi** [IT/US]; 403 Barbara Lane, Bryn Mawr, PA 19010 (US). **EBEL, Wolfgang** [DE/US]; 302 Killybegs Lane, Plymouth Meeting, PA 19462 (US). **ROUTHIER,**

Eric [US/US]; 547 Concord Road, Glen Mills, PA 19342 (US). **YAO, Jun** [CN/US]; 4403 Franklin Court, Chester Springs, PA 19425 (US). **ZHOU, Yuhong** [US/US]; 285 Shelbourne Lane, Phoenixville, PA 19460 (US). **ZUO, Xun** [US/US]; 27 Sandy Lane, Malvern, PA 19355 (US).

(74) Agents: **CALDWELL, John, W.** et al.; Woodcock Washburn LLP, Cira Centre, 12th Floor, 2929 Arch Street, Philadelphia, PA 19104-2891 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,

[Continued on next page]

(54) Title: ANTI-GD2 ANTIBODIES AND METHODS AND USES RELATED THERETO




Fig. 18

(57) Abstract: Described herein are antibodies that specifically bind ganglioside GD2. Also described are nucleotides encoding such antibodies, cells expressing such antibodies, methods of use for such antibodies, and methods for using the antibodies to treat diseases associated with ganglioside GD2. In addition, tissue culture media supplements are described as are methods of use for the supplements. Described herein are albumin-ganglioside conjugates and corresponding methods for producing such conjugates. Methods of purifying or isolating antibodies are also described.



MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR),
OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML,
MR, NE, SN, TD, TG).

— *before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments (Rule 48.2(h))*

Published:

— *with international search report (Art. 21(3))*

— *with sequence listing part of description (Rule 5.2(a))*

ANTI-GD2 ANTIBODIES AND METHODS AND USES RELATED THERETO

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims the benefit of U.S. Provisional Application No. 61/077,041, filed June 30, 2008, and of U.S. Provisional Application No. 61/097,034, filed September 15, 2008, both of which are incorporated by reference herein.

TECHNICAL FIELD

[0002] The invention generally relates to the field of immunotherapeutic agents. More specifically, the invention relates to monoclonal antibodies that bind gangliosides and methods for using such antibodies to treat subjects in need of such treatment. Also provided are methods for using the described antibodies for diagnostic and therapeutic purposes. In addition, tissue culture media compositions and methods that relate to culturing cells expressing the described antibodies, as well as methods for purifying or isolating the described antibodies are also provided.

BACKGROUND

[0003] Cancer is the second leading cause of death in the United States. The National Cancer Institute (NCI) of the U.S. National Institutes of Health (NIH) estimates that over 500,000 individuals will die from cancer and over 1.4 million individuals will be diagnosed with cancer in 2008. There is a global research effort to develop methods of treatment for cancer and to enhance the general understanding of how to prevent cancer, with countries like the United Kingdom, the United States, and others making cancer-related research a top priority. For example, the NIH has earmarked over 5 billion dollars in funding for cancer-related research for the 2008 fiscal year, almost twice the amount of funding provided for other serious diseases, such as HIV and heart disease.

[0004] There are over 100 different types of cancer. Despite the wide variety of cancers, most cancer-related deaths are caused by a few common cancers, such as lung cancer, colorectal cancer, and breast cancer. The NCI estimates that these three forms of cancer will cause over 250,000 deaths in 2008. While not as deadly as those cancers mentioned previously, skin cancer represents about half of all newly diagnosed cancers in the U.S., making it the most common type of cancer. Of the various types of skin cancer, melanoma is the rarest and most

deadly. The American Cancer Society estimates that while melanoma accounts for only 4% of all diagnosed skin cancers in the U.S., it causes 79% of all skin cancer-related deaths.

[0005] Successful cancer treatment is often attributed to early diagnosis and treatment. A powerful method of diagnosing cancer is detecting tumor markers known to be associated with cancer. Tumor markers are substances, often proteins, which are produced by tumor cells, or other cells in the body in response to cancer. Tumor markers can be found in the blood, urine, on the surface of tumor cells, or on (or in) other, non-cancerous cells and tissues. Gangliosides have been identified as one type of marker associated with many tumors (Hakomori, 99 Proc. Natl. Acad. Sci. U.S.A. 10718 (2002)).

[0006] Gangliosides are glycosphingolipids with at least one sugar-linked sialic acid. These compounds are components of cell plasma membranes and are thought to play a central role in a variety of biological functions, such as cell growth, cell differentiation, cell signaling, and serving as receptors for microbial toxins (Jacques, *et al.* 4 Org. Biomol. Chem. 142 (2006)). Over 40 different gangliosides have been identified; however, a certain subset of these, GM3, GM2, GD3, and GD2, are commonly over-expressed by tumor cells (*Id*). Furthermore, gangliosides are reported to be highly immunogenic in humans. The combination of high immunogenicity and over-expression by tumor cells makes gangliosides a potential target for cancer therapeutics.

[0007] Ganglioside-specific monoclonal antibodies have been developed and, in some cases, examined for efficacy in humans (*See generally*, Azuma *et al.*, 13 Clin. Cancer Res. 2745 (2007); Irie, *et al.*, 53 Cancer Immunol. Immunother. 110 (2004); Irie and Morton, 83 Proc. Natl. Acad. Sci. U.S.A. 8694 (1986)). One of the earliest reports concerning a human, GD2-specific antibody was by Cahan, *et al.* (79 Proc. Natl. Acad. Sci. U. S. A. 7629 (1982)). Immunotherapy studies have shown that higher survival rates correlate with ganglioside-specific IgM, rather than IgG, for patients vaccinated with a melanoma vaccine (Jones *et al.*, 66 J. Natl. Cancer Inst. 249 (1981)). Consistent with this finding, human studies involving IgM antibodies to gangliosides GD2 and GD3 have provided encouraging results for the use of such antibodies as a possible treatment for melanoma (Irie, *et al.*, 53 Cancer Immunol. Immunother. 110; Irie and Morton, 83 Proc. Natl. Acad. Sci. U.S.A. 8694 (1986)).

[0008] IgM is one of five human antibody isotypes; IgG, IgA, IgE, and IgD are the others. IgM is typically the first antibody produced in a humoral immune response because the position of IgM heavy chain constant genes allows it to be produced without isotype switching (Charles A. Janeway, *et al.* Immunobiology 9-12 (5th ed. 2001)). Because IgM is often produced before genetic maturation occurs, this class of antibody isotype typically has lower affinity for a

given antigen than other isotypes (*Id.*). IgM compensates for low affinity by forming polymers, which increase the avidity of the antibody molecule (*Id.*). Polymeric IgM usually forms as a pentamer associated with J-chain (a ~15 kD molecule that promotes antibody polymerization); however, it can also polymerize as a pentamer or hexamer in the absence of J-chain (*Id.* at 4-19). In some instances, the polymeric state of IgM allows it to mediate highly effective activation of the complement pathway in the presence of pathogens. For example, J-chain-containing pentameric IgM is not usually effective in activating complement unless it undergoes a structural change when bound to an antigen (*Id.* at 9-17). In contrast, J-chain-free or hexameric IgM has been shown to activate complement up to 100-fold better than pentameric IgM (Weirisma *et al.*, 160 J. Immunol. 5979 (1998)).

[0009] Cancer is a global health problem. Though progress has been made in treating various forms of this disease, improved therapeutics are needed. Immunotherapeutics are thought to have great potential for cancer therapy. An IgM antibody specific for an antigen expressed by cancer cells may prove to be an effective cancer therapeutic.

SUMMARY

[0010] Described herein are isolated antibodies and antigen-binding fragments that specifically bind to ganglioside GD2. In some embodiments, the antibodies or antigen-binding fragments are IgMs. In some embodiments, the antibodies or antigen-binding fragments are pentameric or hexameric. While the antibodies or antigen-binding fragments may be human, humanized, or chimeric, the antibodies or antigen-binding fragments are preferably human. The antibodies or antigen-binding fragments may include a heavy chain having an amino acid sequence substantially the same as or identical to SEQ ID NO: 40 and a light chain having an amino acid sequence substantially the same as or identical to SEQ ID NO: 42. In some embodiments, antibodies or antigen-binding fragments may include a heavy chain CDR1 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 10. In some embodiments, antibodies or antigen-binding fragments may include a heavy chain CDR2 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 11. In some embodiments, antibodies or antigen-binding fragments may include a heavy chain CDR3 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 12. In some embodiments, antibodies or antigen-binding fragments may include a light chain CDR1 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 26. In some embodiments, antibodies or antigen-binding fragments may include a light chain CDR2 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 27. In some embodiments, antibodies or antigen-binding fragments

may include a light chain CDR3 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 28. The antibodies or antigen-binding fragments may include a heavy chain having a CDR1 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 10; a CDR2 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 11; and a CDR3 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 12. The antibodies or antigen-binding fragments may include a light chain having a CDR1 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 26; a CDR2 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 27; and a CDR3 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 28. The antibodies or antigen-binding fragments may include a heavy chain having a CDR1 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 10; a CDR2 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 11; and a CDR3 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 12, and also have a light chain having a CDR1 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 26; a CDR2 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 27; and a CDR3 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 28.

[0011] The antibodies or antigen-binding fragments may include a heavy chain FWR1 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 13. In some embodiments, the antibodies or antigen-binding fragments include a heavy chain FWR2 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 14. In some embodiments, the antibodies or antigen-binding fragments include a heavy chain FWR3 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 15. In some embodiments, the antibodies or antigen-binding fragments include a light chain FWR1 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 29. In some embodiments, the antibodies or antigen-binding fragments include a light chain FWR2 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 30. In some embodiments, the antibodies or antigen-binding fragments include a light chain FWR3 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 31. In some embodiments, the antibodies or antigen-binding fragments include a heavy chain having a FWR1 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 13; a FWR2 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 14; and a FWR3 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 15. In some embodiments, the antibodies or antigen-binding fragments include a light chain having a FWR1 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 29; a FWR2 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 30; and a FWR3

amino acid sequence substantially the same as, or identical to, SEQ ID NO: 31. In some embodiments the antibodies or antigen-binding fragments may include a heavy chain and a light chain, wherein the heavy chain includes a FWR1 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 13; a FWR2 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 14; and a FWR3 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 15; and the light chain includes a FWR1 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 29; a FWR2 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 30; and a FWR3 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 31.

[0012] In some embodiments, the antibodies or antigen-binding fragments include a heavy chain having a CDR1 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 10; a CDR2 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 11; and a CDR3 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 12; a FWR1 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 13; a FWR2 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 14; and a FWR3 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 15. In some embodiments, the antibodies or antigen-binding fragments include a light chain having a CDR1 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 26; a CDR2 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 27; and a CDR3 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 28; a FWR1 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 29; a FWR2 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 30; and a FWR3 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 31. In some embodiments, the antibodies or antigen-binding fragments have a heavy and a light chain, wherein the heavy chain has a CDR1 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 10; a CDR2 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 11; and a CDR3 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 12; a FWR1 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 13; a FWR2 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 14; and a FWR3 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 15; and the light chain has a CDR1 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 26; a CDR2 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 27; and a CDR3 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 28; a FWR1 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 29; a FWR2 amino acid

sequence substantially the same as, or identical to, SEQ ID NO: 30; and a FWR3 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 31. Antigen-binding arrangements of CDRs and FWRs may also be engineered using antibody-like proteins as CDR scaffolding. Such engineered antigen-binding proteins are within the scope of the disclosure.

[0013] The described antibodies or antigen-binding fragments may include a heavy chain that has an amino acid sequence substantially the same as, or identical to, SEQ ID NO: 40. In some embodiments, a polynucleotide substantially the same as, or identical to, SEQ ID NO: 39 may encode this heavy chain amino acid sequence. The described antibodies or antigen-binding fragments may include a light chain that has an amino acid sequence substantially the same as, or identical to, SEQ ID NO: 42. In some embodiments, a polynucleotide substantially the same as, or identical to, SEQ ID NO: 41 may encode this light chain amino acid sequence. The described antibodies or antigen-binding fragments may include a heavy and a light chain, wherein the heavy chain has an amino acid sequence substantially the same as, or identical to, SEQ ID NO: 40, and the light chain has an amino acid sequence substantially the same as, or identical to, SEQ ID NO: 42.

[0014] In some embodiments, the antibodies are produced by antibody-producing cells deposited with the Amer. Type Cult. Coll. (10801 University Blvd., Manassas, Virginia 20110-2209) on July 16, 2008 and have been assigned Accession No. PTA-9376. In some embodiments, the antibodies, or antigen-binding fragments thereof, have the binding affinity for GD2 of the antibodies produced by the deposited antibody-producing cells. In some embodiments, the disclosed antibodies, or antigen-binding fragments thereof, comprise the heavy and light chain CDRs of the antibodies produced by the deposited antibody-producing cells. In some embodiments, the antibodies, or antigen-binding fragments thereof, comprise the heavy and light chain variable regions of the antibodies produced by the deposited antibody-producing cells. In some embodiments, the antibodies, or antigen-binding fragments thereof, exhibit substantially the same, or greater, level of complement-dependent cytotoxic activity as antibodies produced by the deposited antibody-producing cells. In addition, antibodies, or antigen-binding fragments thereof, that compete with the antibodies produced by the deposited cells for binding to GD2 are contemplated to be within the scope of the antibodies described herein. For instance, an antibody may compete with another antibody, or an antigen-binding fragment, if it prevents or hinders the other antibody from binding to an antigenic site to which the antibody would otherwise bind in the absence of the competing antibody.

[0015] Also disclosed are polynucleotides that encode antibodies or antigen-binding fragments that specifically bind to GD2. In some embodiments, the polynucleotides encode an

antibody or antigen-binding fragment thereof having a heavy chain CDR1 sequence substantially the same as, or identical to, SEQ ID NO: 10, for example SEQ ID NO: 2. In some embodiments, the polynucleotides encode an antibody or antigen-binding fragment thereof having a heavy chain CDR2 substantially the same as, or identical to, SEQ ID NO: 11, for example SEQ ID NO: 3. In some embodiments, the polynucleotides encode an antibody or antigen-binding fragment thereof having a heavy chain CDR3 substantially the same as, or identical to, SEQ ID NO: 12, for example SEQ ID NO: 4. In some embodiments, the polynucleotides encode an antibody or antigen-binding fragment thereof having a light chain CDR1 substantially the same as, or identical to, SEQ ID NO: 26, for example SEQ ID NO: 18. In some embodiments, the polynucleotides encode an antibody or antigen-binding fragment thereof having a light chain CDR2 substantially the same as, or identical to, SEQ ID NO: 27, for example SEQ ID NO: 19. In some embodiments, the polynucleotides encode an antibody or antigen-binding fragment thereof having a light chain CDR3 substantially the same as, or identical to, SEQ ID NO: 28, for example SEQ ID NO: 20. The polynucleotides may encode an antibody or antigen-binding fragment thereof having a heavy chain with a CDR1 substantially the same as, or identical to, SEQ ID NO: 10, for example SEQ ID NO: 2; a CDR2 substantially the same as, or identical to, SEQ ID NO: 11, for example SEQ ID NO: 3; and a CDR3 substantially the same as, or identical to, SEQ ID NO: 12, for example SEQ ID NO: 4. The polynucleotides may encode an antibody or antigen-binding fragment thereof having a light chain CDR1 substantially the same as, or identical to, SEQ ID NO: 26, for example SEQ ID NO: 18; a CDR2 substantially the same as, or identical to, SEQ ID NO: 27, for example SEQ ID NO: 19; and a CDR3 substantially the same as, or identical to, SEQ ID NO: 28, for example SEQ ID NO: 20. The polynucleotides may encode an antibody or antigen-binding fragment thereof having a heavy chain CDR1 substantially the same as, or identical to, SEQ ID NO: 10, for example SEQ ID NO: 2; a CDR2 encoded by a nucleotide sequence substantially the same as, or identical to, SEQ ID NO: 11, for example SEQ ID NO: 3; and a CDR3 encoded by a nucleotide sequence substantially the same as, or identical to, SEQ ID NO: 12, for example SEQ ID NO: 4; and a light chain CDR1 substantially the same as, or identical to, SEQ ID NO: 26, for example SEQ ID NO: 18; a CDR2 substantially the same as, or identical to, SEQ ID NO: 27, for example SEQ ID NO: 19; and a CDR3 substantially the same as, or identical to, SEQ ID NO: 28, for example SEQ ID NO: 20. Antigen-binding arrangements of CDRs may also be engineered using antibody-like proteins as CDR scaffolding. Such engineered antigen-binding proteins are within the scope of the disclosure.

[0016] In some embodiments, the polynucleotides encode an antibody or antigen-binding fragment thereof having a heavy chain FWR1 substantially the same as, or identical to, SEQ ID NO: 13, for example SEQ ID NO: 5. In some embodiments, the polynucleotides encode an antibody or antigen-binding fragment thereof having a heavy chain FWR2 substantially the same as, or identical to, SEQ ID NO: 14, for example SEQ ID NO: 6. In some embodiments, the polynucleotides encode an antibody or antigen-binding fragment thereof having a heavy chain FWR3 substantially the same as, or identical to, SEQ ID NO: 15, for example SEQ ID NO: 7. In some embodiments, the polynucleotides encode an antibody or antigen-binding fragment thereof having a light chain FWR1 substantially the same as, or identical to, SEQ ID NO: 29, for example SEQ ID NO: 21. In some embodiments, the polynucleotides encode an antibody or antigen-binding fragment thereof having a light chain FWR2 substantially the same as, or identical to, SEQ ID NO: 30, for example SEQ ID NO: 22. In some embodiments, the polynucleotides encode an antibody or antigen-binding fragment thereof having a light chain FWR3 substantially the same as, or identical to, SEQ ID NO: 31, for example SEQ ID NO: 23. In some embodiments, the polynucleotides encode an antibody or antigen-binding fragment thereof having a heavy chain FWR1 substantially the same as, or identical to, SEQ ID NO: 13, for example SEQ ID NO: 5; a FWR2 substantially the same as, or identical to, SEQ ID NO: 14, for example SEQ ID NO: 6; and a FWR3 substantially the same as, or identical to, SEQ ID NO: 15, for example SEQ ID NO: 7. In some embodiments, the polynucleotides encode an antibody or antigen-binding fragment thereof having a light FWR1 substantially the same as, or identical to, SEQ ID NO: 29, for example SEQ ID NO: 21; a FWR2 substantially the same as, or identical to, SEQ ID NO: 30, for example SEQ ID NO: 22; and a FWR3 substantially the same as, or identical to, SEQ ID NO: 31, for example SEQ ID NO: 23. In some embodiments, the polynucleotides encode an antibody or antigen-binding fragment thereof having a heavy chain and a light chain, wherein a heavy chain FWR1 is substantially the same as, or identical to, SEQ ID NO: 13, for example SEQ ID NO: 5; a heavy chain FWR2 is substantially the same as, or identical to, SEQ ID NO: 14, for example SEQ ID NO: 6; and a heavy chain FWR3 is substantially the same as, or identical to, SEQ ID NO: 15, for example SEQ ID NO: 7; and a light chain FWR1 is substantially the same as, or identical to, SEQ ID NO: 29, for example SEQ ID NO: 21; a light chain FWR2 is substantially the same as, or identical to, SEQ ID NO: 30, for example SEQ ID NO: 22; and a light chain FWR3 substantially the same as, or identical to, SEQ ID NO: 31, for example SEQ ID NO: 23.

[0017] In some embodiments, the polynucleotides encode an antibody or antigen-binding fragment thereof having a heavy chain CDR1 substantially the same as, or identical to,

SEQ ID NO: 10, for example SEQ ID NO: 2; a heavy chain CDR2 substantially the same as, or identical to, SEQ ID NO: 11, for example SEQ ID NO: 3; and a heavy chain CDR3 substantially the same as, or identical to, SEQ ID NO: 12, for example SEQ ID NO: 4; a heavy chain FWR1 substantially the same as, or identical to, SEQ ID NO: 13, for example SEQ ID NO: 5; a heavy chain FWR2 substantially the same as, or identical to, SEQ ID NO: 14, for example SEQ ID NO: 6; and a heavy chain FWR3 substantially the same as, or identical to, SEQ ID NO: 15, for example SEQ ID NO: 7. In some embodiments, the polynucleotides encode an antibody or antigen-binding fragment thereof having a light chain CDR1 substantially the same as, or identical to, SEQ ID NO: 26, for example SEQ ID NO: 18; a light chain CDR2 substantially the same as, or identical to, SEQ ID NO: 27, for example SEQ ID NO: 19; and a light chain CDR3 substantially the same as, or identical to, SEQ ID NO: 28, for example SEQ ID NO: 20; a light chain FWR1 substantially the same as, or identical to, SEQ ID NO: 29, for example SEQ ID NO: 21; a light chain FWR2 substantially the same as, or identical to, SEQ ID NO: 30, for example SEQ ID NO: 22; and a light chain FWR3 substantially the same as, or identical to, SEQ ID NO: 31, for example SEQ ID NO: 23.

[0018] In some embodiments, the polynucleotides encode an antibody or antigen-binding fragment thereof having a heavy and a light chain, wherein the polynucleotides encode a heavy chain CDR1 substantially the same as, or identical to, SEQ ID NO: 10, for example SEQ ID NO: 2; a heavy chain CDR2 substantially the same as, or identical to, SEQ ID NO: 11, for example SEQ ID NO: 3; a heavy chain CDR3 substantially the same as, or identical to, SEQ ID NO: 12, for example SEQ ID NO: 4; a heavy chain FWR1 substantially the same as, or identical to, SEQ ID NO: 13, for example SEQ ID NO: 5; a heavy chain FWR2 substantially the same as, or identical to, SEQ ID NO: 14, for example SEQ ID NO: 6; and a heavy chain FWR3 substantially the same as, or identical to, SEQ ID NO: 15, for example SEQ ID NO: 7; and a light chain CDR1 substantially the same as, or identical to, SEQ ID NO: 26, for example SEQ ID NO: 18; a light chain CDR2 substantially the same as, or identical to, SEQ ID NO: 27, for example SEQ ID NO: 19; a light chain CDR3 substantially the same as, or identical to, SEQ ID NO: 28, for example SEQ ID NO: 20; a light chain FWR1 substantially the same as, or identical to, SEQ ID NO: 29, for example SEQ ID NO: 21; a light chain FWR2 substantially the same as, or identical to, SEQ ID NO: 30, for example SEQ ID NO: 22; and a light chain FWR3 substantially the same as, or identical to, SEQ ID NO: 31, for example SEQ ID NO: 23. Polynucleotides encoding engineered antigen-binding proteins also are within the scope of the disclosure.

[0019] Vectors comprising the antibody- and antigen-binding fragment-encoding polynucleotides are provided, as are cells expressing the antibodies or antigen-binding fragments that specifically bind to GD2.

[0020] Described herein are methods for treating or preventing GD2-associated disease in a subject in need of such treatment. In some embodiments, the GD2-associated disease is cancer. In some embodiments, the GD2-associated disease is melanoma. The methods comprise administering to the subject an antibody or antigen-binding fragment thereof that specifically binds to GD2 in an amount effective to treat or prevent the GD2-associated disease. In some embodiments the methods comprise administering a pharmaceutical composition including an antibody or antigen-binding fragment thereof and a pharmaceutically acceptable carrier. In some aspects, the antibody or antigen-binding fragment thereof is an IgM antibody or antigen-binding fragment. In some embodiments, the described methods of treating or preventing GD2-associated disease in a subject may be carried out with an antibody, or antigen-binding fragment, that competes for binding with the GD2-specific antibodies described herein.

[0021] Described herein are methods for detecting GD2-expressing cells *in vivo* or *in vitro*. The methods may include administering to a subject an antibody or antigen-binding fragment thereof described herein that specifically binds to GD2 to allow for the detection or localization in a subject. In some embodiments, the described antibodies or antigen-binding fragments may be detectably labeled. Such detectable labels may include fluorescent labels, radiolabels, biotin, and the like. Alternatively, in some embodiments the GD2-specific antibodies or antigen-binding fragments are not labeled and, instead are detected by a secondary antibody which is detectably labeled. In some embodiments, the described methods of detection may be carried out with an antibody, or antigen-binding fragment, that competes for binding with the GD2-specific antibodies and antigen-binding fragments described herein.

[0022] Described herein are methods of making and purifying or isolating IgM antibodies or antigen-binding fragments thereof. The methods include culturing a host cell under conditions suitable to produce the antibodies or an antigen-binding fragments, recovering the antibodies or antigen-binding fragments from the cell culture, and purifying or isolating the recovered antibodies or antigen-binding fragments. Also featured are methods of purifying or isolating IgM antibodies or antigen-binding fragments, including optionally washing the antibodies or an antigen-binding fragments with a detergent, applying a solution comprising the antibodies or antigen-binding fragments to an affinity chromatography column, applying eluate from the affinity chromatography column to a cation exchange chromatography column, and applying eluate from the cation exchange chromatography column to a hydroxyapatite

chromatography column and recovering the eluate from the hydroxyapatite chromatography column.

[0023] Additional methods include enhancing the viability of cultured eukaryotic cells by supplementing cell culture media with compositions comprising amino acids, sugars, and vitamins. In some embodiments the compositions may comprise glucose, glutamate, aspartate, serine, histidine, threonine, arginine, tyrosine, cysteine, valine, methionine, tryptophan, phenylalanine, isoleucine, leucine, lysine, proline, nicotinic acid amide, pyridoxine HCl, folic acid, vitamin B-12, riboflavin, and thiamine HCl. Described herein are also methods for enhancing protein production by cells modified to produce proteins. In one aspect, these methods include supplementing the growth media of cells with valeric acid.

[0024] Described herein are protein conjugates of albumin, for example bovine serum albumin (BSA), and a ganglioside and methods for producing such protein conjugates. In one embodiment, the conjugated ganglioside may be GD2. In one embodiment, the conjugated ganglioside may be GM2. In one embodiment, the conjugated ganglioside may be GM3. In one embodiment, albumin is conjugated to a reducing end of a carbohydrate moiety of a ganglioside by reductive amination. In one embodiment, the reductive amination is catalyzed by sodium cyanoborohydride.

BRIEF DESCRIPTION OF THE DRAWINGS

[0025] Figure 1 shows variable antigen specificity of antibodies present in the supernatant of an Epstein-Barr Virus-transformed human lymphoblast pool (HLP) as determined by ELISA. Figure 1A demonstrates that concentrated cell culture supernatants from the HLP, at cell passage 2 (8.1X concentrated) or 4 (14.2X concentrated), contain IgM antibodies specific for GD2. Figure 1B demonstrates that cell culture supernatants from the HLP contain IgM antibodies that bind gangliosides GD1a, GD2, GM2, and GM3.

[0026] Figure 2 shows binding levels of IgM antibodies produced by the HLP cell cultures on GD2-expressing 1205LU melanoma cells, as analyzed by fluorescence activated cell sorting (FACS). Figure 2A serves as negative control, showing binding in the absence of GD2-specific primary antibody. Figure 2B and 2C show binding of HLP-derived IgM obtained from cell culture supernatants concentrated to either 8.1X or 14.2X, respectively. Figure 2D shows binding of a murine GD2-specific IgM antibody produced by murine hybridoma HB-8568TM (ATCC #HB 8568TM) and serves as a positive control.

[0027] Figure 3 shows an antigen-specific ELISA using spent medium from hybridoma clones generated by the HLP-derived hybridomas. Figure 3A shows binding characteristics of

IgM antibodies contained in media of hybridomas 3B2, 5D7, and 10B4 for gangliosides GD2, GD1a, GM2, and GM3. Figure 3B shows binding characteristics of IgM antibodies contained in media of the 1470 hybridoma subclone for gangliosides GD2, GD1a, GM2, and GM3.

[0028] Figure 4 shows binding levels of IgM antibodies produced by hybridoma 3B2 and 1470 cell cultures on GD2-expressing 1205LU melanoma cells, as analyzed by FACS. Figure 4A serves as negative control, showing lack of binding in the absence of GD2-specific primary antibody. Figure 4B, which serves as a positive control, shows binding of a murine GD2-specific IgM antibody produced by HB-8568™ cells. Figs. 4C and D show binding of IgM produced by 3B2 and 1470 hybridoma cells, respectively.

[0029] Figure 5 shows cell death due to complement-dependent cytotoxicity (CDC) mediated by IgM obtained from hybridoma 3B2 tissue culture supernatant or non-GD2-specific human IgM (nhIgM).

[0030] Figure 6 provides a schematic representation of the cloning of AB527 antibody heavy and light chain segments.

[0031] Figure 7 provides a schematic representation of the AB527 expression vector.

[0032] Figure 8 illustrates the binding activity of AB527 for the gangliosides GD2, GM2, and GD3, as measured by surface plasmon resonance.

[0033] Figure 9 provides a graphical representation of AB527 binding equilibrium as measured by surface plasmon resonance for differing concentrations of antibody bound to GD2.

[0034] Figure 10 shows the staining patterns of various gangliosides separated by thin layer chromatography. Orcinol staining shows non-specific ganglioside staining, which provides a basis for comparison with AB527 staining.

[0035] Figure 11 shows flow cytometry (columns I and II) and comparative light microscopy and immunofluorescence microscopy images (columns III and IV) of AB527 binding to human melanoma cell lines that express GD2 at the cell surface (cell lines: M14, M0023, M101, M18, M10-Vac, PM0496) or lack GD2 expression (GD2-negative lines: M21, M238, IMCD0023, MG1055).

[0036] Figure 12 provides a graphical representation of a saturation binding curve and corresponding Scatchard plot that characterizes the binding of AB527 to EL4 cells.

[0037] Figure 13 shows AB527 staining of GD2 positive (M14) and negative (MG1055, RPMI7951 and JS0592) melanoma cell lines. Immunohistochemical analysis was performed using biotinylated AB527 (Figure 13A) or biotinylated, non-GD2-specific, control IgM (Figure 13B).

[0038] Figure 14 shows that M14 tumor sections are not immunoreactive with biotinylated hIgM (14a), but do show immunoreactivity with biotinylated AB527 (14b).

[0039] Figure 15 shows a comparison of complement-dependent cytotoxicity (CDC) of GD2 positive or GD2 negative cells mediated by either IgM obtained from hybridoma AB527-HYB-3B2-3C9 or AB527 IgM.

[0040] Figure 16 is a graphical representation of AB527-mediated CDC in the presence or absence of complement, for human tumor cells that either express GD2 (M14) or do not express GD2 (1205LU).

[0041] Figure 17 illustrates the relative degree of CDC activity for non-specific human IgM, AB527, non-specific hIgG, and isotype-switched AB527 having an IgG1 constant region.

[0042] Figure 18 shows the relative degree of CDC activity for AB527 antibodies with or without J-chain (JC).

[0043] Figure 19 shows a reducing SDS-PAGE gel of detergent-treated conditioned culture supernatant (CCS) applied to a protein A affinity chromatography column (Load) and the material collected from pooled fractions after elution by 3M MgCl₂ (Elute). Arrows point to the IgM heavy chain (~70kD) and light chain (~25kD).

[0044] Figure 20 shows a reducing SDS-PAGE gel of AB527 captured by differing amounts of MacroCap™ SP cation exchange resin. Arrows point to the IgM heavy chain (~70kD) and light chain (~25kD).

[0045] Figure 21 shows relative amounts of various forms of AB527 IgM in the flowthrough (Figure 21A) or eluate (figure 21B) collected after application of differing volumes of AB527 intermediate to MacroCap™ SP cation exchange resin.

[0046] Figure 22 depicts chromatograms from size exclusion-HPLC, verifying the size distribution of material present in the flowthrough (Figure 22A) and elution (Figure 22B) from the MacroCap™ SP column.

[0047] Figure 23 provides a graphical representation of AB527 binding to the GD2-BSA conjugate.

[0048] Figure 24 provides a graphical representation comparing AB527 binding to GD2-BSA, GM2-BSA, and GM3-BSA conjugates.

[0049] Figure 25 shows a comparison of viable cell density over time for AB527-expressing cells cultured in media with (culture media supplement (CMS)) and without (CD CHO control) the addition of an embodiment of the CMS.

[0050] Figure 26 shows the relative percentage of viable cells over time for AB527-expressing cells cultured in media with (CMS) and without (CD CHO control) the addition of an embodiment of the CMS.

[0051] Figure 27 shows the integral viable cell density (IVC) over time for AB527-expressing cells cultured in media with (CMS) and without (CD CHO control) the addition of an embodiment of the CMS.

[0052] Figure 28 shows the total antibody production, after 14 days in culture, for AB527-expressing cells cultured in media with (CMS) and without (CD CHO control) the addition of an embodiment of the CMS.

DETAILED DESCRIPTION OF ILLUSTRATIVE EMBODIMENTS

[0053] Various terms relating to aspects of the description are used throughout the specification and claims. Such terms are to be given their ordinary meaning in the art unless otherwise indicated. Other specifically defined terms are to be construed in a manner consistent with the definitions provided herein.

[0054] As used in this specification and the appended claims, the singular forms “a,” “an,” and “the” include plural referents unless the content clearly dictates otherwise. Thus, for example, reference to “a cell” includes a combination of two or more cells, and the like.

[0055] The term “about” as used herein when referring to a measurable value such as an amount, a temporal duration, and the like, is meant to encompass variations of up to $\pm 20\%$ from the specified value, as such variations are appropriate to perform the disclosed methods. Unless otherwise indicated, all numbers expressing quantities of ingredients, properties such as molecular weight, reaction conditions, and so forth used in the specification and claims are to be understood as being modified in all instances by the term “about.” Accordingly, unless indicated to the contrary, the numerical parameters set forth in the following specification and attached claims are approximations that may vary depending upon the desired properties sought to be obtained by the present invention. At the very least, and not as an attempt to limit the application of the doctrine of equivalents to the scope of the claims, each numerical parameter should at least be construed in light of the number of reported significant digits and by applying ordinary rounding techniques.

[0056] Notwithstanding that the numerical ranges and parameters setting forth the broad scope of the invention are approximations, the numerical values set forth in the specific examples are reported as precisely as possible. Any numerical value, however, inherently contain

certain errors necessarily resulting from the standard deviation found in their respective testing measurements.

[0057] As used herein, the term “cytotoxic” or “cytostatic” agent refers to an agent that inhibits the biological processes of a cell, or reduces the viability or proliferative potential of a cell. Cytotoxic or cytostatic agents may function in a variety of ways, for example, but not by way of limitation, by inducing DNA damage, inducing cell cycle arrest, inhibiting DNA synthesis, inhibiting transcription, inhibiting translation or protein synthesis, inhibiting cell division, or inducing apoptosis. As used herein, the term “chemotherapeutic agent” refers to cytotoxic, cytostatic, and antineoplastic agents that preferentially kill, inhibit the growth of, or inhibit the metastasis of neoplastic cells or disrupt the cell cycle of rapidly proliferating cells. Chemotherapeutic agents include, but are not limited to, synthetic compounds, natural and recombinant bacterial toxins, natural and recombinant fungal toxins, natural and recombinant plant toxins, fissionable nuclides, and radionuclides. Specific examples of chemotherapeutic agents include, but are not limited to, pokeweed antiviral protein, abrin, ricin and each of their A chains, momordin, saporin, bryodin 1, bouganin, gelonin, Diphtheria toxin, Pseudomonas exotoxin, Shiga toxin, calicheamicin, maytansinoid, lead-212, bismuth-212, astatine-211, iodine-131, scandium-47, rhenium-186, rhenium-188, yttrium-90, iodine-123, iodine-124, iodine-125, bromine-77, indium-111, boron-10, actinide, altretamine, actinomycin D, plicamycin, puromycin, gramicidin D, doxorubicin, colchicine, cytochalasin B, cyclophosphamide, emetine, maytansine, amsacrine, cisplatin, etoposide, etoposide orthoquinone, teniposide, daunorubicin, gemcitabine, doxorubicin, mitoxantraone, bisanthrene, Bleomycin, methotrexate, vindesine, adriamycin, vincristine, vinblastine, BCNU, taxol, tarceva, avastin, mitomycin, 5-fluorouracil, cyclophosphamide and certain cytokines such as TNF-alpha and TNF-beta.

[0058] “Isolated” means altered “by the hand of man” from the natural state. If a molecule or composition occurs in nature, it has been “isolated” if it has been changed or removed from its original environment, or both. For example, a polynucleotide or a polypeptide naturally present in a living plant or animal is not “isolated,” but the same polynucleotide or polypeptide separated from the coexisting materials of its natural state is “isolated” as the term is employed herein.

[0059] “Polynucleotide,” synonymously referred to as “nucleic acid molecule” or “nucleic acids,” refers to any polyribonucleotide or polydeoxyribonucleotide, which may be unmodified RNA or DNA or modified RNA or DNA. “Polynucleotides” include, without limitation single- and double-stranded DNA, DNA that is a mixture of single- and double-stranded regions, single- and double-stranded RNA, and RNA that is mixture of single- and

double-stranded regions, hybrid molecules comprising DNA and RNA that may be single-stranded or, more typically, double-stranded or a mixture of single- and double-stranded regions. In addition, “polynucleotide” refers to triple-stranded regions comprising RNA or DNA or both RNA and DNA. The term polynucleotide also includes DNAs or RNAs containing one or more modified bases and DNAs or RNAs with backbones modified for stability or for other reasons. “Modified” bases include, for example, tritylated bases and unusual bases such as inosine. A variety of modifications may be made to DNA and RNA; thus, “polynucleotide” embraces chemically, enzymatically or metabolically modified forms of polynucleotides as typically found in nature, as well as the chemical forms of DNA and RNA characteristic of viruses and cells. “Polynucleotide” also embraces relatively short nucleic acid chains, often referred to as oligonucleotides.

[0060] “Substantially the same” with respect to nucleic acid or amino acid sequences, means at least 65% identity between two or more sequences. Preferably, the term refers to at least 70% identity between two or more sequences, more preferably at least 75% identity, more preferably at least 80% identity, more preferably at least 85% identity, more preferably at least 90% identity, more preferably at least 91% identity, more preferably at least 92% identity, more preferably at least 93% identity, more preferably at least 94% identity, more preferably at least 95% identity, more preferably at least 96% identity, more preferably at least 97% identity, more preferably at least 98% identity, and more preferably at least 99% or greater identity. Such identity may be determined using mBLAST algorithm (Altschul *et al.* (1990) Proc. Natl. Acad. Sci. USA 87:2264-8; Karlin and Altschul (1993) Proc. Natl. Acad. Sci. USA 90:5873-7).

[0061] A “vector” is a replicon, such as plasmid, phage, cosmid, or virus in which another nucleic acid segment may be operably inserted so as to bring about the replication or expression of the segment.

[0062] The term “operably linked” or “operably inserted” means that the regulatory sequences necessary for expression of the coding sequence are placed in a nucleic acid molecule in the appropriate positions relative to the coding sequence so as to enable expression of the coding sequence. By way of example, a promoter is operably linked with a coding sequence when the promoter is capable of controlling the transcription or expression of that coding sequence. Coding sequences may be operably linked to promoters or regulatory sequences in a sense or antisense orientation. The term “operably linked” is sometimes applied to the arrangement of other transcription control elements (*e.g.*, enhancers) in an expression vector.

[0063] A cell has been “transformed” when exogenous or heterologous nucleic acids such as DNA have been introduced inside the cell. The transforming DNA may or may not be

integrated (covalently linked) into the genome of the cell. In prokaryotes, yeast, and mammalian cells for example, the transforming DNA may be maintained on an episomal element such as a plasmid. With respect to eukaryotic cells, a stably transformed cell, or “stable cell” is demonstrated by the ability of the eukaryotic cell to establish cell lines or clones comprised of a population of daughter cells containing the transforming DNA. A “clone” is a population of cells derived from a single cell or common ancestor by mitosis. A “cell line” is a clone of a primary cell that is capable of stable growth *in vitro* for many generations. In some examples provided herein, cells are transformed by transfecting the cells with DNA.

[0064] “Polypeptide” refers to any peptide or protein comprising two or more amino acids joined to each other by peptide bonds or modified peptide bonds, i.e., peptide isosteres. “Polypeptide” refers to both short chains, commonly referred to as peptides, oligopeptides or oligomers, and to longer chains, generally referred to as proteins. Polypeptides may contain amino acids other than the 20 gene-encoded amino acids. “Polypeptides” include amino acid sequences modified either by natural processes, such as post-translational processing, or by chemical modification techniques which are well known in the art. Such modifications are well described in basic texts, monographs, and research literature. Modifications may occur anywhere in a polypeptide, including the peptide backbone, the amino acid side-chains and the amino or carboxyl termini. The same type of modification may be present in the same or varying degrees at several sites in a given polypeptide. Also, a given polypeptide may contain many types of modifications. Polypeptides may be branched as a result of ubiquitination, and they may be cyclic, with or without branching. Cyclic, branched and branched cyclic polypeptides may result from natural posttranslational processes or may be made by synthetic methods. Modifications include acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent cross-links, formation of cystine, formation of pyroglutamate, formylation, gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, proteolytic processing, phosphorylation, prenylation, racemization, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins such as arginylation, and ubiquitination (See, for instance, *Proteins - Structure and Molecular Properties*, 2nd Ed., T. E. Creighton, W. H. Freeman and Company, New York, 1993 and Wold, F., *Posttranslational Protein Modifications: Perspectives and Prospects*, pgs. 1-12 in *Posttranslational Covalent Modification of Proteins*, B. C. Johnson, Ed., Academic Press,

New York, 1983; Seifter *et al.*, Analysis for Protein Modifications and Nonprotein Cofactors, Meth Enzymol (1990) 182:626-646 and Rattan *et al.*, Protein Synthesis: Posttranslational Modifications and Aging, Ann NY Acad Sci (1992) 663:48-62).

[0065] “Biomolecules” include proteins, polypeptides, nucleic acids, lipids, monosaccharides, polysaccharides, and all fragments, analogs, homologs, conjugates, and derivatives thereof.

[0066] The terms “express” and “produce” are used synonymously herein, and refer to the biosynthesis of a gene product. These terms encompass the transcription of a gene into RNA. These terms also encompass translation of RNA into one or more polypeptides, and further encompass all naturally occurring post-transcriptional and post-translational modifications. The expression or production of an antibody or antigen-binding fragment thereof may be within the cytoplasm of the cell, or into the extracellular milieu such as the growth medium of a cell culture.

[0067] The terms “treating” or “treatment” refer to any success or indicia of success in the attenuation or amelioration of an injury, pathology or condition, including any objective or subjective parameter such as abatement, remission, diminishing of symptoms or making the injury, pathology, or condition more tolerable to the patient, slowing in the rate of degeneration or decline, making the final point of degeneration less debilitating, improving a subject’s physical or mental well-being, or prolonging the length of survival. The treatment may be assessed by objective or subjective parameters; including the results of a physical examination, neurological examination, or psychiatric evaluations.

[0068] “Effective amount” and “therapeutically effective amount” are used interchangeably herein, and refer to an amount of an antibody, antigen-binding fragment, or antibody composition, as described herein, effective to achieve a particular biological or therapeutic result such as, but not limited to, biological or therapeutic results disclosed, described, or exemplified herein. A therapeutically effective amount of the antibody or antigen-binding fragment thereof may vary according to factors such as the disease state, age, sex, and weight of the individual, and the ability of the antibody or antigen-binding fragment thereof to elicit a desired response in the individual. Such results may include, but are not limited to, the treatment of cancer, as determined by any means suitable in the art.

[0069] “Pharmaceutically acceptable” refers to those properties and substances which are acceptable to the patient from a pharmacological/toxicological point of view and to the manufacturing pharmaceutical chemist from a physical/chemical point of view regarding composition, formulation, stability, patient acceptance, and bioavailability.

[0070] “Pharmaceutically acceptable carrier” refers to a medium that does not interfere with the effectiveness of the biological activity of the active ingredient(s) and is not toxic to the host to which it is administered.

[0071] “Antibody” refers to all isotypes of immunoglobulins (IgG, IgA, IgE, IgM, IgD, and IgY) including various monomeric and polymeric forms of each isotype, unless otherwise specified.

[0072] Antigen-binding fragments are any proteinaceous structure that may exhibit binding affinity for a particular antigen. Some antigen-binding fragments are composed of portions of intact antibodies that retain antigen-binding specificity of the parent antibody molecule. For example, antigen-binding fragments may comprise at least one variable region (either a heavy chain or light chain variable region) or one or more CDRs of an antibody known to bind a particular antigen. Examples of suitable antigen-binding fragments include, without limitation diabodies and single-chain molecules as well as Fab, F(ab')₂, Fc, Fabc, and Fv molecules, single chain (Sc) antibodies, individual antibody light chains, individual antibody heavy chains, chimeric fusions between antibody chains or CDRs and other proteins, protein scaffolds, or molecules, heavy chain monomers or dimers, light chain monomers or dimers, dimers consisting of one heavy and one light chain, and the like. All antibody isotypes may be used to produce antigen-binding fragments. Additionally, antigen-binding fragments may include non-antibody proteinaceous frameworks that may successfully incorporate polypeptide segments in an orientation that confers affinity for a given antigen of interest, such as protein scaffolds. Antigen-binding fragments may be recombinantly produced or produced by enzymatic or chemical cleavage of intact antibodies. The phrase “an antibody or antigen-binding fragment thereof” may be used to denote that a given antigen-binding fragment incorporates one or more amino acid segments of the antibody referred to in the phrase.

[0073] “Antibody compositions” refer to antibodies or binding fragments thereof that are coupled with at least one pharmaceutically acceptable carrier, chemotherapeutic agent, or diagnostic moiety, such as ¹¹¹In-1,4,7,10-tetraazacyclododecane-N,N',N'',N'''-tetraacetic acid (¹¹¹In-DOTA).

[0074] “Specific binding” refers to the ability of an antibody, or antigen-binding fragment, to bind to a particular biomolecule with an affinity that is greater than that with which it may bind other biomolecules.

[0075] The embodiments described herein are not limited to particular methods, reagents, compounds, compositions or biological systems, which can, of course, vary.

Furthermore, the terminology used herein is for the purpose of describing particular antibodies or antigen-binding fragments only, and is not intended to be limiting.

GD2-Specific Antibodies

[0076] Described herein are isolated antibodies or antigen-binding fragments that specifically bind the ganglioside GD2. In one embodiment, the antibody or antigen-binding fragment thereof is a monoclonal antibody or antigen-binding fragment; however, other embodiments include GD2-specific polyclonal antibodies and derivatives or fragments of antibodies that retain specificity for GD2. The general structure of an antibody molecule comprises an antigen binding domain, which includes heavy and light chains, and the Fc domain, which serves a variety of functions, including complement fixation. In some embodiments, the antibodies or antigen-binding fragments mediate complement-dependent cytotoxicity.

[0077] There are five classes of immunoglobulins wherein the primary structure of the heavy chain, in the Fc region, determines the immunoglobulin class. Specifically, the alpha, delta, epsilon, gamma, and mu chains correspond to IgA, IgD, IgE, IgG and IgM isotypes, respectively. The described antibodies or antigen-binding fragments include all isotypes and synthetic multimers of the four-chain immunoglobulin structure. The described antibodies or antigen-binding fragments also include the IgY isotype generally found in hen or turkey serum and hen or turkey egg yolk. Antibodies or antigen-binding fragments non-covalently, specifically, and reversibly bind an antigen.

[0078] The antibodies or antigen-binding fragments of the disclosed subject matter may be derived from any species. For example, the antibodies or antigen-binding fragments may be mouse, rat, goat, horse, swine, bovine, chicken, rabbit, donkey, human, and the like. For use in the treatment of humans, non-human derived antibodies or antigen-binding fragments may be genetically or structurally altered to be less antigenic upon administration to a human patient.

[0079] In some embodiments, the antibodies or antigen-binding fragments are chimeric. As used herein, the term “chimeric” antibody, or antigen-binding fragment, means an antibody, or antigen-binding fragment thereof, having at least some portion of at least one variable domain derived from the antibody amino acid sequence of a non-human mammal, a rodent, or a reptile, while the remaining portions of the antibody, or antigen-binding fragment thereof, are derived from a human. For example, a chimeric antibody may comprise a mouse antigen binding domain with a human Fc or other such structural domain.

[0080] In some embodiments, the antibodies are humanized antibodies. Humanized antibodies may be chimeric immunoglobulins, immunoglobulin chains or fragments thereof

(such as Fv, Fab, Fab', F(ab')₂ or other antigen-binding subsequences of antibodies) that contain minimal sequence derived from non-human immunoglobulin. For the most part, humanized antibodies are human immunoglobulins (recipient antibody) in which residues from a complementary-determining region (CDR) of the recipient are replaced by residues from a CDR of a non-human species (donor antibody) such as mouse, rat or rabbit having the desired specificity, affinity, and capacity. In some instances, Fv framework region (FWR) residues of the human immunoglobulin are replaced by corresponding non-human residues. Furthermore, humanized antibodies may comprise residues which are found neither in the recipient antibody nor in the imported CDR or framework sequences. These modifications are made to further refine and optimize antibody performance. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the CDR regions correspond to those of a non-human immunoglobulin and all or substantially all of the FWR regions are those of a human immunoglobulin sequence. The humanized antibody optimally also will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin. For further details, see Jones *et al.*, 321 Nature 522-5 (1986); Reichmann *et al.*, 332 Nature 323-9 (1988); and Presta, 2 Curr. Op. Struct. Biol. 593-6 (1992).

[0081] In some cases, the described antibodies are human antibodies or antigen binding fragments thereof. As used herein, the term "human antibody" means that the antibody is either solely from human origin or an antibody in which the variable and constant domain sequences are human sequences. The term encompasses antibodies with sequences derived from (i.e., that utilize) human genes, but which have been changed, *e.g.*, to decrease possible immunogenicity, increase affinity, eliminate cysteines that may cause undesirable folding, etc. The term encompasses such antibodies produced recombinantly in non-human cells, which may impart glycosylation not typical of human cells.

[0082] The antibodies or antigen-binding fragments described herein may be labeled or otherwise conjugated to various chemical or biomolecule moieties, for example, for therapeutic or diagnostic applications. The moieties may be cytotoxic, for example, bacterial toxins, viral toxins, radioisotopes, and the like. The moieties may be detectable labels, for example, fluorescent labels, radiolabels, biotin, and the like, for example, ¹¹¹In-DOTA, ¹¹¹In-diethylenetriaminepentaacetic acid (DTPA), or radionuclides, such as, but not limited to, lead-212, bismuth-212, astatine-211, iodine-131, scandium-47, rhenium-186, rhenium-188, yttrium-90, iodine-123, iodine-124, iodine-125, bromine-77, indium-111, and fissionable nuclides such as boron-10 or an actinide.

[0083] In some embodiments, the antibodies or antigen-binding fragments are conjugated to one or more chemotherapeutic agents such as, but not limited to radionuclides, toxins, and cytotoxic and cytostatic agents. In other embodiments the antibodies or antigen-binding fragments are used in combination with one or more chemotherapeutic agents. The antibodies or antigen-binding fragments described herein may be used alone or with (*e.g.*, coadministered or conjugated to) a biomolecule or chemotherapeutic agent such as a cytotoxic or cytostatic agent. In some embodiments, the chemotherapeutic agent is a radionuclide, including, but not limited to lead-212, bismuth-212, astatine-211, iodine-131, scandium-47, rhenium-186, rhenium-188, yttrium-90, iodine-123, iodine-124, iodine-125, bromine-77, indium-111, and fissionable nuclides such as boron-10 or an actinide. In other embodiments, the chemotherapeutic agent is a toxin or cytotoxic drug, pokeweed antiviral protein, abrin, ricin and each of their A chains, momordin, saporin, bryodin 1, bouganin, gelonin, Diphtheria toxin, Pseudomonas exotoxin, Shiga toxin, calicheamicin, maytansinoid, altretamine, actinomycin D, plicamycin, puromycin, gramicidin D, doxorubicin, colchicine, cytochalasin B, cyclophosphamide, emetine, maytansine, amsacrine, cisplatin, etoposide, etoposide orthoquinone, teniposide, daunorubicin, gemcitabine, doxorubicin, mitoxantraone, bisanthrene, Bleomycin, methotrexate, pemetrexed, cisplatin, vindesine, adriamycin, vincristine, vinblastine, BCNU, a taxane (*e.g.*, Taxol™), tarceva, avastin, mitomycin, 5-fluorouracil, cyclophosphamide, certain cytokines such as TNF-alpha and TNF-beta, and the like. Methods of conjugation of antibodies or antigen-binding fragments to such agents are known in the literature.

[0084] Antibody specificity is primarily determined by the six CDR regions, especially H chain CDR3 (Kala *et al.*, 132 J. Biochem. 535-41 (2002); Morea *et al.*, 275 J. Mol. Biol. 269-94 (1998); and, Chothia *et al.*, 196 J. Mol. Biol. 901-17 (1987)). Antibody framework regions, however, can play a role in antigen-antibody interactions (Panka *et al.*, 85 Proc. Natl. Acad. Sci. USA 3080-4 (1988)), particularly in influencing the conformation of CDR loops (Foote *et al.*, 224 J. Mol. Biol. 487-99 (1992)). Thus, the described antibodies or antigen-binding fragments may comprise any combination of H or L chain CDR or FWR regions that confer specificity to GD2. Domain shuffling experiments, which are routinely carried out in the art (Jirholt *et al.*, 215 Gene 471-6 (1998); Söderlind *et al.*, 18 Nature Biotechnology 852-6 (2000)), may be employed to generate antibodies that specifically bind GD2 according to the specifications described and exemplified herein. Antibodies or antigen-binding fragments generated by such domain shuffling experiments are within the scope of the antibodies or antigen-binding fragments described herein. Furthermore, CDRs may also be arranged to bind a given antigen by

engineering antibody-like proteins to serve as CDR scaffolding (Nicaise et al., 13 Protein Sci. 1882 (2004)). Such antigen-binding proteins are within the scope of the antibodies described herein.

[0085] The antibodies or antigen-binding fragments described herein can occur in a variety of forms, but will include one or more of the antibody segments shown in Table 1.

Table 1. Antibody segments of the described antibodies and antigen-binding fragments thereof (“Lc” denotes light chain and “Hc” denotes heavy chain).

AB527 Antibody Segment	Amino Acid SEQ ID NO.	DNA SEQ ID NO.
Lc CDR1	26	18
Lc CDR2	27	19
Lc CDR3	28	20
Lc FWR1	29	21
Lc FWR2	30	22
Lc FWR3	31	23
Lc variable domain	32	24
Light chain	42	41
Hc CDR1	10	2
Hc CDR2	11	3
Hc CDR3	12	4
Hc FWR1	13	5
Hc FWR2	14	6
Hc FWR3	15	7
Hc variable domain	16	8
Heavy chain	40	39

[0086] In some embodiments, the antibodies or antigen-binding fragments may include a heavy chain CDR1 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 10. In some embodiments, the antibodies or antigen-binding fragments may include a heavy chain CDR2 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 11. In some embodiments, antibodies or antigen-binding fragments may include a heavy chain CDR3 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 12. In some embodiments, antibodies or antigen-binding fragments may include a light chain CDR1 amino

acid sequence substantially the same as, or identical to, SEQ ID NO: 26. In some embodiments, antibodies or antigen-binding fragments may include a light chain CDR2 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 27. In some embodiments, antibodies or antigen-binding fragments may include a light chain CDR3 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 28. In some embodiments, antibodies or antigen-binding fragments may include a heavy chain CDR1 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 10; a CDR2 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 11; and a CDR3 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 12. In some embodiments, antibodies or antigen-binding fragments may include a light chain CDR1 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 26; a CDR2 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 27; and a CDR3 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 28. In some embodiments, antibodies or antigen-binding fragments may include a heavy chain and a light chain, wherein the heavy chain has a CDR1 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 10; a CDR2 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 11; and a CDR3 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 12; and the light chain has a CDR1 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 26; a CDR2 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 27; and a CDR3 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 28. Antigen-binding arrangements of CDRs may also be engineered using antibody-like proteins as CDR scaffolding. Such engineered antigen-binding proteins are within the scope of the disclosure.

[0087] In some embodiments, antibodies or antigen-binding fragments may include a heavy chain FWR1 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 13. In some embodiments, antibodies or antigen-binding fragments may include a heavy chain FWR2 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 14. In some embodiments, antibodies or antigen-binding fragments may include a heavy chain FWR3 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 15. In some embodiments, antibodies or antigen-binding fragments may include a light chain FWR1 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 29. In some embodiments, antibodies or antigen-binding fragments may include a light chain FWR2 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 30. In some embodiments, antibodies or antigen-binding fragments may include a light chain FWR3 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 31. In some embodiments, antibodies or antigen-binding

fragments may include a heavy chain having a FWR1 amino acid sequence that is substantially the same as, or identical to, SEQ ID NO: 13; a FWR2 amino acid sequence that is substantially the same as, or identical to, SEQ ID NO: 14; and a FWR3 amino acid sequence that is substantially the same as, or identical to, SEQ ID NO: 15. In some embodiments, antibodies or antigen-binding fragments may include a light chain having a FWR1 amino acid sequence that is substantially the same as, or identical to, SEQ ID NO: 29; a FWR2 amino acid sequence that is substantially the same as, or identical to, SEQ ID NO: 30; and a FWR3 amino acid sequence that is substantially the same as, or identical to, SEQ ID NO: 31. In some embodiments, antibodies or antigen-binding fragments may include a heavy chain and a light chain, wherein the heavy chain includes a FWR1 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 13; a FWR2 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 14; and a FWR3 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 15; and the light chain includes a FWR1 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 29; a FWR2 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 30; and a FWR3 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 31.

[0088] In some embodiments, antibodies or antigen-binding fragments may include a heavy chain having a CDR1 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 10; a CDR2 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 11; and a CDR3 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 12; a FWR1 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 13; a FWR2 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 14; and a FWR3 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 15. In some embodiments, the antibodies or antigen-binding fragments include a light chain having a CDR1 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 26; a CDR2 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 27; and a CDR3 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 28; a FWR1 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 29; a FWR2 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 30; and a FWR3 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 31. In some embodiments, the antibodies or antigen-binding fragments include a heavy and a light chain, wherein the heavy chain includes a CDR1 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 10; a CDR2 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 11; and a CDR3 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 12; a

FWR1 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 13; a FWR2 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 14; and a FWR3 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 15; and the light chain includes a CDR1 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 26; a CDR2 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 27; and a CDR3 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 28; a FWR1 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 29; a FWR2 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 30; and a FWR3 amino acid sequence substantially the same as, or identical to, SEQ ID NO: 31. Antigen-binding arrangements of CDRs and FWRs may also be engineered using antibody-like proteins as CDR scaffolding. Such engineered antigen-binding proteins are within the scope of the disclosure.

[0089] In some embodiments, the antibodies or antigen-binding fragments described herein have a heavy chain that includes the amino acid sequence substantially the same as, or identical to, SEQ ID NO: 40. In some embodiments, the antibodies or antigen-binding fragments described herein have a light chain that includes the amino acid sequence substantially the same as, or identical to, SEQ ID NO: 42. The described antibodies or antigen-binding fragments may have a heavy and a light chain, wherein the heavy chain includes the amino acid sequence substantially the same as, or identical to, SEQ ID NO: 40 and the light chain includes the amino acid sequence substantially the same as, or identical to, SEQ ID NO: 42.

[0090] Also described are polynucleotides that encode antibodies or antigen-binding fragments that specifically bind to GD2. In some embodiments, the polynucleotides encode an antibody or antigen-binding fragment thereof having a heavy chain CDR1 sequence substantially the same as, or identical to, SEQ ID NO: 10, for example SEQ ID NO: 2. In some embodiments, the polynucleotides encode an antibody or antigen-binding fragment thereof having a heavy chain CDR2 substantially the same as, or identical to, SEQ ID NO: 11, for example SEQ ID NO: 3. In some embodiments, the polynucleotides encode an antibody or antigen-binding fragment thereof having a heavy chain CDR3 substantially the same as, or identical to, SEQ ID NO: 12, for example SEQ ID NO: 4. In some embodiments, the polynucleotides encode an antibody or antigen-binding fragment thereof having a light chain CDR1 substantially the same as, or identical to, SEQ ID NO: 26, for example SEQ ID NO: 18. In some embodiments, the polynucleotides encode an antibody or antigen-binding fragment thereof having a light chain CDR2 substantially the same as, or identical to, SEQ ID NO: 27, for example SEQ ID NO: 19. In some embodiments, the polynucleotides encode an antibody or antigen-binding fragment thereof having a light chain CDR3 substantially the same as, or identical to, SEQ ID NO: 28, for

example SEQ ID NO: 20. The polynucleotides may encode an antibody or antigen-binding fragment thereof having a heavy chain with a CDR1 substantially the same as, or identical to, SEQ ID NO: 10, for example SEQ ID NO: 2; a CDR2 substantially the same as, or identical to, SEQ ID NO: 11, for example SEQ ID NO: 3; and a CDR3 substantially the same as, or identical to, SEQ ID NO: 12, for example SEQ ID NO: 4. The polynucleotides may encode an antibody or antigen-binding fragment thereof having a light chain CDR1 substantially the same as, or identical to, SEQ ID NO: 26, for example SEQ ID NO: 18; a CDR2 substantially the same as, or identical to, SEQ ID NO: 27, for example SEQ ID NO: 19; and a CDR3 substantially the same as, or identical to, SEQ ID NO: 28, for example SEQ ID NO: 20. The polynucleotides may encode an antibody or antigen-binding fragment thereof having a heavy chain CDR1 substantially the same as, or identical to, SEQ ID NO: 10, for example SEQ ID NO: 2; a CDR2 encoded by a nucleotide sequence substantially the same as, or identical to, SEQ ID NO: 11, for example SEQ ID NO: 3; and a CDR3 encoded by a nucleotide sequence substantially the same as, or identical to, SEQ ID NO: 12, for example SEQ ID NO: 4; and a light chain CDR1 substantially the same as, or identical to, SEQ ID NO: 26, for example SEQ ID NO: 18; a CDR2 substantially the same as, or identical to, SEQ ID NO: 27, for example SEQ ID NO: 19; and a CDR3 substantially the same as, or identical to, SEQ ID NO: 28, for example SEQ ID NO: 20.

[0091] In some embodiments, the polynucleotides encode an antibody or antigen-binding fragment thereof having a heavy chain FWR1 substantially the same as, or identical to, SEQ ID NO: 13, for example SEQ ID NO: 5. In some embodiments, the polynucleotides encode an antibody or antigen-binding fragment thereof having a heavy chain FWR2 substantially the same as, or identical to, SEQ ID NO: 14, for example SEQ ID NO: 6. In some embodiments, the polynucleotides encode an antibody or antigen-binding fragment thereof having a heavy chain FWR3 substantially the same as, or identical to, SEQ ID NO: 15, for example SEQ ID NO: 7. In some embodiments, the polynucleotides encode an antibody or antigen-binding fragment thereof having a light chain FWR1 substantially the same as, or identical to, SEQ ID NO: 29, for example SEQ ID NO: 21. In some embodiments, the polynucleotides encode an antibody or antigen-binding fragment thereof having a light chain FWR2 substantially the same as, or identical to, SEQ ID NO: 30, for example SEQ ID NO: 22. In some embodiments, the polynucleotides encode an antibody or antigen-binding fragment thereof having a light chain FWR3 substantially the same as, or identical to, SEQ ID NO: 31, for example SEQ ID NO: 23. In some embodiments, the polynucleotides encode an antibody or antigen-binding fragment thereof having a heavy chain FWR1 substantially the same as, or identical to, SEQ ID NO: 13, for example SEQ ID NO: 5; a FWR2 substantially the same as, or identical to, SEQ ID NO: 14,

for example SEQ ID NO: 6; and a FWR3 substantially the same as, or identical to, SEQ ID NO: 15, for example SEQ ID NO: 7. In some embodiments, the polynucleotides encode an antibody or antigen-binding fragment thereof having a light FWR1 substantially the same as, or identical to, SEQ ID NO: 29, for example SEQ ID NO: 21; a FWR2 substantially the same as, or identical to, SEQ ID NO: 30, for example SEQ ID NO: 22; and a FWR3 substantially the same as, or identical to, SEQ ID NO: 31, for example SEQ ID NO: 23. In some embodiments, the polynucleotides encode an antibody or antigen-binding fragment thereof having a heavy chain and a light chain, wherein a heavy chain FWR1 is substantially the same as, or identical to, SEQ ID NO: 13, for example SEQ ID NO: 5; a heavy chain FWR2 is substantially the same as, or identical to, SEQ ID NO: 14, for example SEQ ID NO: 6; and a heavy chain FWR3 is substantially the same as, or identical to, SEQ ID NO: 15, for example SEQ ID NO: 7; and a light chain FWR1 is substantially the same as, or identical to, SEQ ID NO: 29, for example SEQ ID NO: 21; a light chain FWR2 is substantially the same as, or identical to, SEQ ID NO: 30, for example SEQ ID NO: 22; and a light chain FWR3 substantially the same as, or identical to, SEQ ID NO: 31, for example SEQ ID NO: 23.

[0092] In some embodiments, the polynucleotides encode an antibody or antigen-binding fragment thereof having a heavy chain CDR1 substantially the same as, or identical to, SEQ ID NO: 10, for example SEQ ID NO: 2; a heavy chain CDR2 substantially the same as, or identical to, SEQ ID NO: 11, for example SEQ ID NO: 3; and a heavy chain CDR3 substantially the same as, or identical to, SEQ ID NO: 12, for example SEQ ID NO: 4; a heavy chain FWR1 substantially the same as, or identical to, SEQ ID NO: 13, for example SEQ ID NO: 5; a heavy chain FWR2 substantially the same as, or identical to, SEQ ID NO: 14, for example SEQ ID NO: 6; and a heavy chain FWR3 substantially the same as, or identical to, SEQ ID NO: 15, for example SEQ ID NO: 7. In some embodiments, the polynucleotides encode an antibody or antigen-binding fragment thereof having a light chain CDR1 substantially the same as, or identical to, SEQ ID NO: 26, for example SEQ ID NO: 18; a light chain CDR2 substantially the same as, or identical to, SEQ ID NO: 27, for example SEQ ID NO: 19; and a light chain CDR3 substantially the same as, or identical to, SEQ ID NO: 28, for example SEQ ID NO: 20; a light chain FWR1 substantially the same as, or identical to, SEQ ID NO: 29, for example SEQ ID NO: 21; a light chain FWR2 substantially the same as, or identical to, SEQ ID NO: 30, for example SEQ ID NO: 22; and a light chain FWR3 substantially the same as, or identical to, SEQ ID NO: 31, for example SEQ ID NO: 23.

[0093] In some embodiments, the polynucleotides encode an antibody or antigen-binding fragment thereof having a heavy and a light chain, wherein the polynucleotides encode a

heavy chain CDR1 substantially the same as, or identical to, SEQ ID NO: 10, for example SEQ ID NO: 2; a heavy chain CDR2 substantially the same as, or identical to, SEQ ID NO: 11, for example SEQ ID NO: 3; a heavy chain CDR3 substantially the same as, or identical to, SEQ ID NO: 12, for example SEQ ID NO: 4; a heavy chain FWR1 substantially the same as, or identical to, SEQ ID NO: 13, for example SEQ ID NO: 5; a heavy chain FWR2 substantially the same as, or identical to, SEQ ID NO: 14, for example SEQ ID NO: 6; and a heavy chain FWR3 substantially the same as, or identical to, SEQ ID NO: 15, for example SEQ ID NO: 7; and a light chain CDR1 substantially the same as, or identical to, SEQ ID NO: 26, for example SEQ ID NO: 18; a light chain CDR2 substantially the same as, or identical to, SEQ ID NO: 27, for example SEQ ID NO: 19; a light chain CDR3 substantially the same as, or identical to, SEQ ID NO: 28, for example SEQ ID NO: 20; a light chain FWR1 substantially the same as, or identical to, SEQ ID NO: 29, for example SEQ ID NO: 21; a light chain FWR2 substantially the same as, or identical to, SEQ ID NO: 30, for example SEQ ID NO: 22; and a light chain FWR3 substantially the same as, or identical to, SEQ ID NO: 31, for example SEQ ID NO: 23.

[0094] Polynucleotides encoding engineered antigen-binding proteins also are within the scope of the disclosure.

[0095] In some embodiments, the polynucleotides described (and the peptides they encode) include a leader sequence. Any leader sequence known in the art may be employed. The leader sequence may include, but is not limited to, a restriction site or a translation start site. In some embodiments, the leader sequence has the nucleic acid sequence ATGGGATGGAGCTGTATCATCCTCTTCTTGGTAGCAACAGCTACAGGTGTACACAGC (SEQ ID NO: 43). In some embodiments, the leader sequence encodes the amino acid sequence MGWSCILFLVATATGVHS (SEQ ID NO: 44).

[0096] Because of the natural sequence variation likely to exist among heavy and light chains and the genes encoding them, one would expect to find some level of variation within the amino acid sequences or the genes encoding the antibodies or antigen-binding fragments described herein, with little or no impact on their unique binding properties (*e.g.*, specificity and affinity). Such an expectation is due in part to the degeneracy of the genetic code, as well as to the evolutionary success of conservative amino acid sequence variations, which do not appreciably alter the nature of the encoded protein. Accordingly, some embodiments include antibodies or antigen-binding fragments having 90%, 95%, 96%, 97%, 98%, or 99% homology to the antibodies or antigen-binding fragments herein. Other embodiments include GD-2 specific antibodies or antigen-binding fragments, that have framework, scaffold, or other non-binding regions that do not share significant homology with the antibodies and antigen-binding

fragments described herein, but do incorporate one or more CDRs or other sequences needed to confer binding that are 90%, 95%, 96%, 97%, 98%, or 99% homologous to such sequences described herein.

[0097] The antibodies or antigen-binding fragments described herein include variants having single or multiple amino acid substitutions, deletions, or additions that retain the biological properties (*e.g.*, binding affinity or immune effector activity) of the described antibodies or antigen-binding fragments. The skilled person may produce variants having single or multiple amino acid substitutions, deletions, or additions. These variants may include: (a) variants in which one or more amino acid residues are substituted with conservative or nonconservative amino acids, (b) variants in which one or more amino acids are added to or deleted from the polypeptide, (c) variants in which one or more amino acids include a substituent group, and (d) variants in which the polypeptide is fused with another peptide or polypeptide such as a fusion partner, a protein tag or other chemical moiety, that may confer useful properties to the polypeptide, such as, for example, an epitope for an antibody, a polyhistidine sequence, a biotin moiety and the like. Antibodies or antigen-binding fragments described herein may include variants in which amino acid residues from one species are substituted for the corresponding residue in another species, either at the conserved or nonconserved positions. In other embodiments, amino acid residues at nonconserved positions are substituted with conservative or nonconservative residues. The techniques for obtaining these variants, including genetic (suppressions, deletions, mutations, etc.), chemical, and enzymatic techniques, are known to the person having ordinary skill in the art.

[0098] The antibodies or antigen-binding fragments described herein may embody several antibody isotypes, such as IgM, IgD, IgG, IgA and IgE. Antibody or antigen-binding fragment thereof specificity is largely determined by the amino acid sequence, and arrangement, of the CDRs. Therefore, the CDRs of one isotype may be transferred to another isotype without altering antigen specificity. Alternatively, techniques have been established to cause hybridomas to switch from producing one antibody isotype to another (isotype switching) without altering antigen specificity. Accordingly, such antibody isotypes are within the scope of the described antibodies or antigen-binding fragments.

[0099] Altering the isotype of an antibody may be useful for combining different effector functions, conferred by the antibody constant region, with antigen specificity. For example, IgG is produced in monomeric form, is effective at activating complement, and is commonly found in the plasma and intracellular compartments of the body. IgA is produced as a monomer and a J-chain-associated dimer. IgA is most commonly found in the luminal spaces of

the body and breast milk, because the dimeric form may be transported across epithelial cell barriers. Like IgA, IgM can also form polymers, however, it tends to form pentamers, when associated with J-chain, and pentamers and hexamers not associated with J-chain. These highly polymeric antibodies serve to increase overall antigen-binding capability by increasing avidity without reducing affinity. Both polymeric forms of IgM can activate complement efficiently, which can result in complement-dependent cytotoxicity (CDC).

[0100] The antibodies or antigen-binding fragments described herein, in some embodiments, are polymeric IgM antibodies or antigen-binding fragments. In some embodiments, the antibodies or antigen-binding fragments are pentameric IgM antibodies or antigen-binding fragments that are associated with J-chain. In other embodiments, the antibodies or antigen-binding fragments are hexameric IgM antibodies or antigen-binding fragments that are not associated with J-chain. In some embodiments, the antibodies or antigen-binding fragments are pentameric IgM antibodies or antigen-binding fragments that are not associated with J-chain. In some embodiments, the antibodies or antigen-binding fragments are hexameric IgM antibodies or antigen-binding fragments that are associated with J-chain.

[0101] The antibodies or antigen-binding fragments described herein have binding affinities (in M) for GD2 that include a dissociation constant (K_D) of less than 1×10^{-2} . In some embodiments, the K_D is less than 1×10^{-3} . In other embodiments, the K_D is less than 1×10^{-4} . In some embodiments, the K_D is less than 1×10^{-5} . In still other embodiments, the K_D is less than 1×10^{-6} , 2×10^{-6} , 3×10^{-6} , 4×10^{-6} , 5×10^{-6} , 6×10^{-6} , 7×10^{-6} , 8×10^{-6} , or 9×10^{-6} . In other embodiments, the K_D is less than 1×10^{-7} , 2×10^{-7} , or 3×10^{-7} , 2×10^{-7} , 3×10^{-7} , 4×10^{-7} , 5×10^{-7} , 6×10^{-7} , 7×10^{-7} , 8×10^{-7} , or 9×10^{-7} . In other embodiments, the K_D is less than 1×10^{-8} , 2×10^{-8} , 3×10^{-8} , 4×10^{-8} , 5×10^{-8} , 6×10^{-8} , 7×10^{-8} , 8×10^{-8} , or 9×10^{-8} . In other embodiments, the K_D is less than 1×10^{-9} , 2×10^{-9} , 3×10^{-9} , 4×10^{-9} , 5×10^{-9} , 6×10^{-9} , 7×10^{-9} , 8×10^{-9} , or 9×10^{-9} . In other embodiments, the K_D is less than 1×10^{-10} , 2×10^{-10} , 3×10^{-10} , 2×10^{-10} , 3×10^{-10} , 4×10^{-10} , 5×10^{-10} , 6×10^{-10} , 7×10^{-10} , 8×10^{-10} , or 9×10^{-10} . In still other embodiments, the K_D is less than 1×10^{-11} , 2×10^{-11} , 3×10^{-11} , 4×10^{-11} , 5×10^{-11} , 6×10^{-11} , 7×10^{-11} , 8×10^{-11} , or 9×10^{-11} . In some embodiments, the K_D is less than 1×10^{-12} . In other embodiments, the K_D is less than 1×10^{-13} . In other embodiments, the K_D is less than 1×10^{-14} . In still other embodiments, the K_D is less than 1×10^{-15} . In preferred embodiments, the K_D is 4.5×10^{-9} or less.

[0102] The antibodies or antigen-binding fragments described herein, in some embodiments, have specific binding affinities for GD2 in contrast with each of GD1a, GM2 or GM3. In some embodiments, the K_D for GD2 differs from the K_D for each of GD1a, GM2 or GM3 by at least 3-fold, preferably 10-fold.

[0103] The antibodies or antigen-binding fragments described herein may be modified, *e.g.*, by the covalent attachment of any type of molecule to the antibody or antigen-binding fragment thereof such that covalent attachment does not prevent the antibody or antigen-binding fragment thereof from binding to its epitope. Examples of suitable modifications include, but are not limited to glycosylation, acetylation, pegylation, phosphorylation, amidation, and the like. In some embodiments the antibodies or antigen-binding fragments may themselves be derivatized by known protecting/blocking groups, proteolytic cleavage, linkage to a cellular ligand or other proteins, and the like. The antibodies or antigen-binding fragments may have post-translational moieties that improve upon antibody or antigen-binding fragment thereof activity or stability. These moieties include sulfur, methyl, carbohydrate, phosphorus as well as other chemical groups commonly found on immunoglobulin molecules. Furthermore, the antibodies or antigen-binding fragments may contain one or more non-classical amino acids.

[0104] Antibodies or antigen-binding fragments described herein may be labeled with or conjugated to toxic or non-toxic moieties. Toxic moieties include, for example, bacterial toxins, viral toxins, plant toxins, fungal toxins, radioisotopes, and the like. Antibodies or antigen-binding fragments may be labeled for use in biological assays (*e.g.*, radioisotope labels, fluorescent labels) to aid in detection of the antibody or antigen-binding fragment. Antibodies or antigen-binding fragments may also be labeled or conjugated for diagnostic or therapeutic purposes, *e.g.*, with radioactive isotopes that deliver radiation directly to a desired site for applications such as radioimmunotherapy (Garmestani *et al.*, 28 Nucl. Med. Biol. 409 (2001)), imaging techniques and radioimmunoguided surgery or labels that allow for *in vivo* imaging or detection of specific antibody/antigen complexes. Antibodies or antigen-binding fragments may also be conjugated with toxins to provide an immunotoxin (*see*, Kreitman, R.J., 31 Adv. Drug Del. Rev. 53 (1998)).

[0105] Described herein are compositions comprising at least one described antibody or antigen-binding fragment thereof and a pharmaceutically acceptable carrier. Such compositions are useful, for example, for administration to patients to treat cancer, such as those described and exemplified herein. The compositions may be formulated as any of various preparations that are known and suitable in the art, including those described and exemplified herein. In some embodiments, the compositions are aqueous formulations. Aqueous solutions may be prepared by admixing the antibodies or antigen-binding fragments in water or suitable physiologic buffer, and optionally adding suitable colorants, flavors, preservatives, stabilizing and thickening agents and the like as desired. Aqueous suspensions may also be made by dispersing the antibodies or antigen-binding fragments in water or physiologic buffer with viscous material, such as natural

or synthetic gums, resins, methylcellulose, sodium carboxymethylcellulose, and other well-known suspending agents.

[0106] Also included are liquid formulations and solid form preparations which are intended to be converted, shortly before use, to liquid preparations. Such liquids include solutions, suspensions, syrups, slurries, and emulsions. Liquid preparations may be prepared by conventional means with pharmaceutically acceptable additives such as suspending agents (*e.g.*, sorbitol syrup, cellulose derivatives or hydrogenated edible fats or oils); emulsifying agents (*e.g.*, lecithin or acacia); non-aqueous vehicles (*e.g.*, almond oil, oily esters, or fractionated vegetable oils); and preservatives (*e.g.*, methyl or propyl-p-hydroxybenzoates or sorbic acid). These preparations may contain, in addition to the active agent, colorants, flavors, stabilizers, buffers, artificial and natural sweeteners, dispersants, thickeners, solubilizing agents, and the like. The compositions may be in powder or lyophilized form for constitution with a suitable vehicle such as sterile water, physiological buffer, saline solution, or alcohol, before use.

[0107] The compositions may be formulated for injection into a subject. For injection, the compositions described may be formulated in aqueous solutions such as water or alcohol, or in physiologically compatible buffers such as Hanks's solution, Ringer's solution, or physiological saline buffer. The solution may contain one or more formulatory agents such as suspending, stabilizing or dispersing agents. Injection formulations may also be prepared as solid form preparations which are intended to be converted, shortly before use, to liquid form preparations suitable for injection, for example, by constitution with a suitable vehicle, such as sterile water, saline solution, or alcohol, before use.

[0108] The compositions may be formulated in sustained release vehicles or depot preparations. Such long-acting formulations may be administered by implantation (for example subcutaneously or intramuscularly) or by intramuscular injection. Thus, for example, the compositions may be formulated with suitable polymeric or hydrophobic materials (for example, as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt. Liposomes and emulsions are well-known examples of delivery vehicles suitable for use as carriers for hydrophobic drugs.

[0109] Disclosed herein are methods for detecting GD-2-expressing cells using the described antibodies, or antigen-binding fragments, either *in vivo* or *in vitro*. Some embodiments make use of disclosed antibodies, or antigen-binding fragments, that are conjugated to detectable labels such as fluorescent labels, radiolabels, biotin, enzymes and the like, for example ¹¹¹In-DOTA, ¹¹¹In-DTPA, or radionuclides, including, but not limited to, lead-212, bismuth-212, astatine-211, iodine-131, scandium-47, rhenium-186, rhenium-188, yttrium-90, iodine-123,

iodine-124, iodine-125, bromine-77, indium-111, and fissionable nuclides such as boron-10 or an actinide. For example, a detectably labeled antibody, or antigen-binding fragment, may be administered to a subject to detect and localize cells in the subject that express GD2. Such methods could also be used to detect cells or tissues where GD2 is highly expressed relative to other cells or tissues in the subject. Alternatively, one embodiment of the disclosed method for detecting GD-2-expressing cells may include using a detectably labeled antibody or antigen-binding fragment, described herein, to detect or quantify the expression of GD2 by cells obtained from a subject, such as cells obtained from a blood sample or tissue biopsy.

[0110] Alternatively, the described methods for detecting GD-2-expressing cells using the described antibodies, or antigen-binding fragments, may be performed using GD2-specific antibodies that are not labeled. For example, in one embodiment, GD2-expressing cells may be detected in a subject by first administering to the subject a GD2-specific antibody, or antigen-binding fragment, described herein followed by administration of a detectably labeled secondary antibody capable of binding the GD2-specific antibody, or antigen-binding fragment, administered initially. A similar methodology could be used to detect GD2-expressing cells *in vitro*.

[0111] Also described herein are methods for treating or preventing diseases in subjects in need of such treatment or prevention. In some aspects, the methods can include identifying a subject in need of treatment or prevention of GD2-associated diseases, such as cancer, *e.g.*, melanoma. Other embodiments include an antibody or antigen-binding fragment thereof that is labeled or conjugated for diagnostic purposes, *e.g.*, with radioactive isotopes that deliver radiation directly to a desired site, such as imaging techniques or detection of specific antibody/antigen complexes. In one embodiment, the methods comprise administering to the subject a GD2-specific antibody or antigen-binding fragment, such as a recombinant, human, GD2-specific IgM antibody in an amount effective to treat or prevent disease. In one aspect the methods include administering to the subject a composition, such as those described and exemplified herein, the composition comprising a pharmaceutically acceptable carrier and at least one antibody or antigen-binding fragment thereof that specifically binds to GD2, in an amount effective to treat or prevent disease. In one embodiment, the methods comprise administering to the subject at least one antibody or antigen-binding fragment, such as the antibodies or antigen-binding fragments described and exemplified herein, that specifically binds to GD2, in an amount effective to treat or prevent disease. In one embodiment, the methods comprise administering to the subject at least one GD2-specific antibody or antigen-binding fragment thereof labeled or conjugated to toxic or non-toxic moieties exemplified herein.

[0112] The antibodies or antigen-binding fragments described herein may be administered orally in any acceptable dosage form such as capsules, tablets, aqueous suspensions, solutions or the like. The antibodies or antigen-binding fragments may also be administered parenterally including but not limited to: subcutaneous, intravenous, intramuscular, intra-articular, intra-synovial, intrasternal, intranasal, topically, intrathecal, intrahepatic, intralesional, and intracranial injection or infusion techniques. Alternatively, the antibodies or antigen-binding fragments will be intravenously or intraperitoneally, for example, by injection.

[0113] The subject may be any animal, and preferably is a mammal such as a mouse, rat, hamster, guinea pig, rabbit, cat, dog, monkey, donkey, cow, horse, pig, and the like. Most preferably, the mammal is a human. In some embodiments, subjects may be administered at least one anti-GD2 antibody or antigen-binding fragment thereof in a daily dose range of 0.01 μ g to 500 mg of antibody or antigen-binding fragment thereof per kg of the weight of the subject. The dose administered to the subject may also be measured in terms of total amount of the at least one anti-GD2 antibody or antigen-binding fragment thereof administered per day. In some embodiments, a subject is administered 5 to 5000 milligrams of at least one anti-GD2 antibody or antigen-binding fragment thereof per day. In some embodiments, a subject is administered up to 10 milligrams of at least one anti-GD2 antibody or antigen-binding fragment thereof per day. In some embodiments, a subject is administered up to 100 milligrams of at least one anti-GD2 antibody or antigen-binding fragment thereof per day. In some embodiments, a subject is administered up to 250 milligrams of at least one anti-GD2 antibody or antigen-binding fragment thereof per day. In some embodiments, a subject is administered up to 500 milligrams of at least one anti-GD2 antibody or antigen-binding fragment thereof per day. In some embodiments, a subject is administered up to 750 milligrams of at least one anti-GD2 antibody or antigen-binding fragment thereof per day. In some embodiments, a subject is administered up to 1000 milligrams of at least one anti-GD2 antibody or antigen-binding fragment thereof per day. In some embodiments, a subject is administered up to 1500 milligrams of at least one anti-GD2 antibody or antigen-binding fragment thereof per day. In some embodiments, a subject is administered up to 2000 milligrams of at least one anti-GD2 antibody or antigen-binding fragment thereof per day. In some embodiments, a subject is administered up to 2500 milligrams of at least one anti-GD2 antibody or antigen-binding fragment thereof per day. In some embodiments, a subject is administered up to 3000 milligrams of at least one anti-GD2 antibody or antigen-binding fragment thereof per day. In some embodiments, a subject is administered up to 3500 milligrams of at least one anti-GD2 antibody or antigen-binding fragment thereof per day. In some embodiments, a subject is administered up to 4000 milligrams of at least one anti-

GD2 antibody or antigen-binding fragment thereof per day. In some embodiments, a subject is administered up to 4500 milligrams of at least one anti-GD2 antibody or antigen-binding fragment thereof per day. In some embodiments, a subject is administered up to 5000 milligrams of at least one anti-GD2 antibody or antigen-binding fragment thereof per day. In some embodiments, antibody or antigen-binding fragment thereof is administered to a subject weekly or bi-weekly.

[0114] Treatment may be initiated with smaller dosages that are less than the optimum dose of the at least one anti-GD2 antibody or antigen-binding fragment, followed by an increase in dosage over the course of the treatment until the optimum effect under the circumstances is reached. If indicated, the total daily dosage may be divided and administered in portions throughout the day.

[0115] For effective treatment of GD2-associated diseases, one skilled in the art may recommend a dosage schedule and dosage amount adequate for the subject being treated. Dosing may occur one to four or more times daily for as long as needed. The dosing may occur less frequently if the compositions are formulated in sustained delivery vehicles. The dosage schedule may also vary depending on the active drug concentration, which may depend on the needs of the subject.

[0116] Also provided are vectors comprising the polynucleotides described herein. The vectors can be expression vectors. Recombinant expression vectors containing a sequence encoding a polypeptide of interest are thus provided. The expression vector may contain one or more additional sequences such as but not limited to regulatory sequences (*e.g.*, promoter, enhancer), a selection marker, and a polyadenylation signal. Vectors for transforming a wide variety of host cells are well known and include, but are not limited to, plasmids, phagemids, cosmids, baculoviruses, bacmids, bacterial artificial chromosomes (BACs), yeast artificial chromosomes (YACs), as well as other bacterial, yeast and viral vectors.

[0117] Recombinant expression vectors within the scope of the description include synthetic, genomic, or cDNA-derived nucleic acid fragments that encode at least one recombinant protein which may be operably linked to suitable regulatory elements. Such regulatory elements may include a transcriptional promoter, sequences encoding suitable mRNA ribosomal binding sites, and sequences that control the termination of transcription and translation. Expression vectors, especially mammalian expression vectors, may also include one or more nontranscribed elements such as an origin of replication, a suitable promoter and enhancer linked to the gene to be expressed, other 5' or 3' flanking nontranscribed sequences, 5' or 3' nontranslated sequences (such as necessary ribosome binding sites), a polyadenylation site,

splice donor and acceptor sites, or transcriptional termination sequences. An origin of replication that confers the ability to replicate in a host may also be incorporated.

[0118] The transcriptional and translational control sequences in expression vectors to be used in transforming vertebrate cells may be provided by viral sources. Exemplary vectors may be constructed as described by Okayama and Berg, 3 Mol. Cell. Biol. 280 (1983).

[0119] In some embodiments, the antibody or antigen-binding fragment-coding sequence is placed under control of a powerful constitutive promoter, such as the promoters for the following genes: hypoxanthine phosphoribosyl transferase (HPRT), adenosine deaminase, pyruvate kinase, beta-actin, human myosin, human hemoglobin, human muscle creatine, and others. In addition, many viral promoters function constitutively in eukaryotic cells and are suitable for use with the described embodiments. Such viral promoters include without limitation, Cytomegalovirus (CMV) immediate early promoter, the early and late promoters of SV40, the Mouse Mammary Tumor Virus (MMTV) promoter, the long terminal repeats (LTRs) of Moloney leukemia virus, Human Immunodeficiency Virus (HIV), Epstein Barr Virus (EBV), Rous Sarcoma Virus (RSV), and other retroviruses, and the thymidine kinase promoter of Herpes Simplex Virus. In one embodiment, the antibody or antigen-binding fragment thereof coding sequence is placed under control of an inducible promoter such as the metallothionein promoter, tetracycline-inducible promoter, doxycycline-inducible promoter, promoters that contain one or more interferon-stimulated response elements (ISRE) such as protein kinase R 2',5'-oligoadenylate synthetases, Mx genes, ADAR1, and the like.

[0120] Vectors described herein may contain one or more Internal Ribosome Entry Site(s) (IRES). Inclusion of an IRES sequence into fusion vectors may be beneficial for enhancing expression of some proteins. In some embodiments the vector system will include one or more polyadenylation sites (*e.g.*, SV40), which may be upstream or downstream of any of the aforementioned nucleic acid sequences. Vector components may be contiguously linked, or arranged in a manner that provides optimal spacing for expressing the gene products (*i.e.*, by the introduction of "spacer" nucleotides between the ORFs), or positioned in another way. Regulatory elements, such as the IRES motif, may also be arranged to provide optimal spacing for expression.

[0121] The vectors may comprise selection markers, which are well known in the art. Selection markers include positive and negative selection markers, for example, antibiotic resistance genes (*e.g.*, neomycin resistance gene, a hygromycin resistance gene, a kanamycin resistance gene, a tetracycline resistance gene, a penicillin resistance gene), glutamate synthase genes, HSV-TK, HSV-TK derivatives for ganciclovir selection, or bacterial purine nucleoside

phosphorylase gene for 6-methylpurine selection (Gadi *et al.*, 7 Gene Ther. 1738-1743 (2000)). A nucleic acid sequence encoding a selection marker or the cloning site may be upstream or downstream of a nucleic acid sequence encoding a polypeptide of interest or cloning site.

[0122] The vectors described herein may be used to transform various cells with the genes encoding the described antibodies or antigen-binding fragments. For example, the vectors may be used to generate antibody or antigen-binding fragment-producing cells. Thus, another aspect features host cells transformed with vectors comprising a nucleic acid sequence encoding an antibody or antigen-binding fragment thereof that specifically binds GD2, such as the antibodies or antigen-binding fragments described and exemplified herein.

[0123] Numerous techniques are known in the art for the introduction of foreign genes into cells and may be used to construct the recombinant cells for purposes of carrying out the described methods, in accordance with the various embodiments described and exemplified herein. The technique used should provide for the stable transfer of the heterologous gene sequence to the host cell, such that the heterologous gene sequence is heritable and expressible by the cell progeny, and so that the necessary development and physiological functions of the recipient cells are not disrupted. Techniques which may be used include but are not limited to chromosome transfer (*e.g.*, cell fusion, chromosome mediated gene transfer, micro cell mediated gene transfer), physical methods (*e.g.*, transfection, spheroplast fusion, microinjection, electroporation, liposome carrier), viral vector transfer (*e.g.*, recombinant DNA viruses, recombinant RNA viruses) and the like (described in Cline, 29 Pharmac. Ther. 69-92 (1985)). Calcium phosphate precipitation and polyethylene glycol (PEG)-induced fusion of bacterial protoplasts with mammalian cells may also be used to transform cells.

[0124] Cells suitable for use in the expression of the antibodies or antigen-binding fragments described herein are preferably eukaryotic cells, more preferably cells of plant, rodent, or human origin, for example but not limited to NSO, CHO, perC.6, Tk-ts13, BHK, HEK293 cells, COS-7, T98G, CV-1/EBNA, L cells, C127, 3T3, HeLa, NS1, Sp2/0 myeloma cells, and BHK cell lines, among others. In addition, expression of antibodies may be accomplished using hybridoma cells. Methods for producing hybridomas are well established in the art.

[0125] Cells transformed with expression vectors described herein may be selected or screened for recombinant expression of the antibodies or antigen-binding fragments described herein. Recombinant-positive cells are expanded and screened for subclones exhibiting a desired phenotype, such as high level expression, enhanced growth properties, or the ability to yield proteins with desired biochemical characteristics, for example, due to protein modification or altered post-translational modifications. These phenotypes may be due to inherent properties of

a given subclone or to mutation. Mutations may be effected through the use of chemicals, UV-wavelength light, radiation, viruses, insertional mutagens, inhibition of DNA mismatch repair, or a combination of such methods.

[0126] Once a cell expressing the desired protein is identified, it can be expanded and selected. Transformed cells may be selected in a number of ways. For example, cells may be selected for expression of the polypeptide of interest. Cells transformed with a vector that contains a selectable marker, such as production of fluorescent protein, may be positively selected for expression of the marker. In other embodiments, the cells containing a vector with a drug resistance gene may be positively selected for the ability to grow under selective conditions.

Kits

[0127] A kit is provided for inhibiting or reducing growth of cancer cells in a patient. Also provided are kits for identifying the presence of dysplastic cells *in vitro* or *in vivo*.

[0128] The kits described herein may comprise an antibody, antigen-binding fragments thereof, or an antibody composition described herein and instructions for using the kit in a method for inhibiting or reducing growth of tumor cells in the patient or in a method for identifying the presence of dysplastic cells, for example, in a biological sample. The kit may comprise at least one chemotherapeutic or cytotoxic reagent. The kit may comprise an antifolate compound. The kit may comprise at least one diagnostic reagent. An example of a diagnostic reagent is a detectable label, for example but not limited to a radioactive, fluorescent, or chromophoric agent (*e.g.*, ^{111}In -DOTA). The detectable label may comprise an enzyme. The kit may comprise instructions and a means for administering the antibody or antibody composition, for example, by injection.

Uses and Methods of Making BSA-Ganglioside Conjugates

[0129] Described herein are albumin-ganglioside conjugates and methods for producing such conjugates. To determine whether an antibody, or antigen-binding fragment, of interest is effective in mediating binding to a given antigen, one must characterize antibody binding. There are numerous ways to characterize antibody binding, such as dot-blot, western blot, immunoprecipitation assay, ELISA, FACS analysis/Flow cytometry, and immunofluorescence detection of bound antibody. While a variety of assays are available to test antibody binding, the particular assay used must be selected with an understanding of the antibody's purpose in mind. For example, if an antibody or antigen-binding fragment thereof is to be used in a clinical setting, it should be characterized by an assay that will present the antigen of interest in a form most

likely to be found *in vivo*. In this instance, and in the case of a cell-surface protein, flow cytometry may be a useful assay because it allows the interaction of the antibody or antigen-binding fragment thereof to be assessed in the context of the native antigen.

[0130] One way to identify cells containing a vector that encodes an antigen-specific binding protein is using an Enzyme-Linked ImmunoSorbent Assay (ELISA) that makes use of the antigen of interest. There are two main approaches used to perform an ELISA: an indirect ELISA and a sandwich ELISA. To carry out an indirect ELISA, the antigen of interest is adsorbed or fixed to the surface of a microtiter plate and then the antibody or antigen-binding fragment thereof of interest is added to the microtiter plate to detect the antigen. Conversely, a sandwich ELISA makes use of an antibody known to be specific for the antigen of interest, in addition to the antibody being characterized. The known antibody is used to coat the microtiter plate, such that it will bind, or capture, the antigen of interest when it is added to the microtiter plate. Once the antigen of interest has been captured, the antibody or antigen-binding fragment thereof of interest is added to the microtiter plate to assess its binding characteristics for the antigen. In either case, bound antibody or antigen-binding fragment thereof is typically detected by an enzyme-conjugated secondary antibody that is specific for the antibody or antigen-binding fragment thereof of interest.

[0131] An ELISA useful for characterizing ganglioside-specific antibodies or antigen-binding fragments would be useful in determining the potential applications of such antibodies or antigen-binding fragments. The use of gangliosides as ELISA antigens is complicated, however, by the fact that free gangliosides are poorly immunogenic (Jacques *et al.*, 4 Org. Biomol. Chem. 142-154 (2006)).

Ganglioside Conjugates

[0132] Disclosed herein are ganglioside-albumin conjugates that not only retain the antigenic properties of the conjugated ganglioside, but increase the stability of the ganglioside and allow better adherence to an ELISA microtiter plate. A ganglioside is conjugated to albumin through reductive amination of the ganglioside by primary amines in albumin in the presence of sodium cyanoborohydride. Conjugation proceeds through Schiff base formation to the reducing end(s) of the carbohydrate moiety on the ganglioside, followed by reduction with sodium cyanoborohydride. Accordingly, some embodiments described herein include a ganglioside conjugated to a carrier protein via a reducing end(s) of the carbohydrate moiety on the ganglioside. Some embodiments comprise a ganglioside conjugated to a carrier protein, wherein the carrier is BSA. In some embodiments, the ganglioside is GD2. In some embodiments, the

ganglioside is GM3. In some embodiments, the ganglioside is GM2. In another embodiment, GD2 is conjugated to BSA. In some embodiments, the reductive amination of a ganglioside is catalyzed by sodium cyanoborohydride.

[0133] In some embodiments the ganglioside-albumin conjugate retains ganglioside antigenicity such that a ganglioside-specific monoclonal antibody or antigen-binding fragment thereof can bind the conjugate. In some embodiments the GD2-BSA conjugate retains GD2 antigenicity such that a GD2-specific monoclonal antibody or antigen-binding fragment thereof can bind the conjugate. In some embodiments the GM2-BSA conjugate retains GM2 antigenicity such that a GM2-specific monoclonal antibody or antigen-binding fragment thereof can bind the conjugate. In some embodiments the GM3-BSA conjugate retains GM3 antigenicity such that a GM3-specific monoclonal antibody or antigen-binding fragment thereof can bind the conjugate.

[0134] The albumin-ganglioside conjugates described herein may be used in an ELISA to characterize the specificity of an antibody or antigen-binding fragment thereof for a particular ganglioside. In one aspect, the albumin-ganglioside conjugates described provide an enhanced ability to adsorb the conjugated gangliosides to ELISA plates relative to unconjugated gangliosides. Once adsorbed, the conjugates may be used as ganglioside antigens in a direct ELISA to characterize binding aspects of ganglioside specific antibodies or antigen-binding fragments. In one embodiment, the albumin-ganglioside conjugate may be a BSA-GD2 conjugate. In one embodiment, the BSA-ganglioside conjugate may be a BSA-GM2 conjugate. In one embodiment, the BSA-ganglioside conjugate may be a BSA-GM3 conjugate. Other such gangliosides may be conjugated to BSA using the methods described and then used for the purposes described herein. Such conjugates and uses thereof are within the scope of this disclosure.

[0135] Embodiments of the protein conjugates include the following.

[0136] Embodiment A1 provides a protein conjugate comprising albumin and a ganglioside.

[0137] Embodiment A2 provides the conjugate of embodiment A1 wherein the albumin is bovine serum albumin.

[0138] Embodiment A3 provides the conjugate of embodiment A1, wherein the ganglioside is conjugated to albumin via a reducing end of the carbohydrate moiety on the ganglioside.

[0139] Embodiment A4 provides the conjugate of embodiment A3 wherein the albumin is bovine serum albumin.

[0140] Embodiment A5 provides the conjugate of embodiment A3 wherein the ganglioside is GD2.

[0141] Embodiment A6 provides the conjugate of embodiment A1 wherein the conjugate is immunoreactive with a GD2-specific antibody.

[0142] Embodiment A7 provides the conjugate of embodiment A1, wherein the ganglioside is GM2.

[0143] Embodiment A8 provides the conjugate of embodiment A7, wherein the conjugate is immunoreactive with a GM2-specific antibody.

[0144] Embodiment A9 provides the conjugate of embodiment A1, wherein the ganglioside is GM3.

[0145] Embodiment A10 provides the conjugate of embodiment A9, wherein the conjugate is immunoreactive with a GM3-specific antibody.

[0146] Embodiment A11 provides a method of conjugating a ganglioside to albumin, the method comprising reductively aminating a reducing end of a carbohydrate moiety on the ganglioside.

[0147] Embodiment A12 provides the method of embodiment A11, wherein the albumin is bovine serum albumin.

[0148] Embodiment A13 provides the method of embodiment A11, wherein the reductive amination is catalyzed by sodium cyanoborohydride.

[0149] Embodiment A14 provides the method of embodiment A13, wherein the ganglioside is GD2.

[0150] Embodiment A15 provides the method of embodiment A13, wherein the ganglioside is GM2.

[0151] Embodiment A16 provides the method of embodiment A13, wherein the ganglioside is GM3.

Cell Culture Compositions and Methods

[0152] Described herein are compositions and methods for culturing eukaryotic cells. The cells may be any type of eukaryotic cell, but, in one embodiment, are preferably of mammalian origin. In some embodiments, the cells are Chinese Hamster ovary (CHO) cells. In some embodiments, the eukaryotic cells are genetically modified to produce IgM antibodies or antigen-binding fragments. The eukaryotic cells may be clinical isolates, transformed with foreign DNA or RNA, immortalized, virus-infected, or modified by other generally known biological or chemical means. The eukaryotic cells may be grown in monolayers or in

suspended culture, which may occur in a cell culture incubator, bioreactor, shake-flask, or other similar tissue culture device. In addition, the eukaryotic cells can be cultured at a temperature above 32 °C and below 50 °C, in between 2% to 8% CO₂, with or without shaking. In one embodiment, the cells can be cultured at a temperature of 37 °C, in 5% CO₂, with shaking at 120 rpm. The eukaryotic cells can be cultured in tissue culture media, such as GIBCO-CD-CHO complete medium (1 L GIBCO-CD-CHO medium + 25µM MSX) or similar complete medium, in combination with the compositions and methods disclosed herein. The cultured eukaryotic cells may be engineered to produce a foreign protein, which the compositions and methods described herein can, but need not necessarily, enhance.

[0153] Disclosed herein are compositions suitable for use with eukaryotic cell culture media. The composition includes, but is not limited to, various essential and nonessential amino acids, for example, alanine, arginine, asparagine, aspartic acid, cysteine, glutamine, glutamic acid, glycine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine, and valine; sugars, for example, glucose, fructose, mannose, and galactose; and vitamins, such as folic acid, vitamin B-12, vitamin D, and riboflavin that may influence growth characteristics of cultured eukaryotic cells. In embodiment B1, the composition includes glucose, glutamate, aspartate, serine, histidine, threonine, arginine, tyrosine, cysteine, valine, methionine, tryptophan, phenylalanine, isoleucine, leucine, lysine, proline, nicotinic acid amide, pyridoxine HCl, folic acid, vitamin B-12, riboflavin, and thiamine HCl.

[0154] In embodiment B2, the composition comprises the cell culture of embodiment B1, wherein the concentration of glucose is 50 g/L, the concentration of glutamate is from 2.5 to 3.75 g/L, the concentration of aspartate is from 1.5 to 2.0 g/L, the concentration of serine is from 0.3 to 0.5 g/L, the concentration of histidine is from 1.1 to 1.5 g/L, the concentration of threonine is from 2.0 to 3.0 g/L, the concentration of arginine is from 1.0 to 1.5 g/L, the concentration of tyrosine is from 1.8 to 2.2 g/L, the concentration of cysteine is from 0.9 to 1.1 g/L, the concentration of valine is from 1.0 to 3.0, the concentration of methionine is from 0.8 to 1.2 g/L, the concentration of Tryptophan is from 0.5 to 0.8 g/L, the concentration of phenylalanine 1.3 to 1.7 g/L, the concentration of isoleucine is from 0.8 to 2.4 g/L, the concentration of leucine is from 1.5 to 4.5 g/L, the concentration of lysine is from 3.5 to 5.0 g/L, the concentration of proline is from 0.5 to 0.7 g/L, the concentration of nicotinic acid amide is from 30 to 40 mg/L, the concentration of pyridoxine HCl is from 200 to 250 mg/L, the concentration of folic acid 100 to 130 mg/L, the concentration of vitamin B-12 20 to 40 mg/L,

the concentration of riboflavin is from 20 to 40 mg/L, and the concentration of thiamine HCl is from 100 to 150 mg/L.

[0155] Also disclosed herein are methods for culturing eukaryotic cells in which a composition described in the preceding paragraph may be used to enhance the viability of eukaryotic cells. For example, the compositions may be added to cultured CHO cells and to cultured IgM-producing eukaryotic (e.g., CHO) cells. The composition may be used to enhance the viability of eukaryotic tissue culture cells by adding the composition to growth media, as a media supplement, either before or after growth media is applied to eukaryotic cells.

Alternatively, growth media components may be added to the composition to produce a growth media that contains the composition.

[0156] Embodiment B3 provides a method of culturing eukaryotic cells comprising adding to the cells a primary tissue culture media and then supplementing the primary tissue culture media with a cell culture composition comprising, glucose, glutamate, aspartate, serine, histidine, threonine, arginine, tyrosine, cysteine, valine, methionine, tryptophan, phenylalanine, isoleucine, leucine, lysine, proline, nicotinic acid amide, pyridoxine HCl, folic acid, vitamin B-12, riboflavin, and thiamine HCl.

[0157] Embodiment B4 provides the method of embodiment B3 wherein the volume of the cell culture composition that is added to the primary tissue culture media is between 1.0% and 20% of the volume of the primary tissue culture media.

[0158] Embodiment B5 provides the method of embodiment B3, wherein the cell culture composition is not added to the primary tissue culture media before the third day following addition of the primary tissue culture media.

[0159] Embodiment B6 provides the method of embodiment B3, wherein the eukaryotic cells have been transformed to produce one or more proteins.

[0160] Embodiment B7 provides the method of embodiment B6, wherein the one or more proteins is an antibody, or an antigen-binding fragment thereof.

[0161] Embodiment B8 provides the method of embodiment B7 wherein the antibody is an IgM.

[0162] Embodiment B9 provides the method of embodiment B3, wherein the cell culture composition is added to the primary tissue culture media on consecutive days.

[0163] In another embodiment, the composition may be used to increase the amount of protein produced by a eukaryotic cell that is genetically modified to produce a recombinant protein. For example, the composition may be added to cultured cells engineered to express a recombinant human antibody or antigen-binding fragment thereof (e.g., cells modified to express

IgM and more specifically CHO cells modified to express IgM). The composition may be used to increase recombinant protein expression by adding the composition to growth media, as a media supplement, either before or after growth media is applied to eukaryotic cells.

Alternatively, growth media components may be added to the composition to produce a growth media that contains the composition.

[0164] The composition may be used as growth media or as a supplement for growth media. As a supplement the composition may be added to cell culture media either before or after the culture media is added to tissue culture containers. The composition described herein may be used to supplement eukaryotic cell culture media to a total concentration of 1.5% to 2.5%. These same concentrations may be used to produce cell culture media already containing the composition.

[0165] Also featured are methods for increasing the amount of protein produced by a eukaryotic cell that is genetically modified to produce a recombinant protein by adding valeric acid to cell culture media. For example, the valeric acid may be added to media for cultured CHO, hybridoma, and NS0 cells engineered to express a recombinant human antibody or antigen-binding fragment. In one embodiment, valeric acid may be added to cell culture media to a final concentration from 0.1 mM to 10 mM. In one embodiment, valeric acid may be added at specific points in the cell growth period, for example, valeric acid may be added to the culture media on specific days following revival of cell stocks or after cell cultures are split.

[0166] Embodiment B10 provides a method of enhancing protein production by cells, comprising supplementing the growth media of the cells with valeric acid.

[0167] Embodiment B11 provides the method of embodiment B10, wherein the cells are antibody-producing cells.

[0168] Embodiment B12 provides the method of embodiment B10, wherein the growth media is supplemented to have a concentration of valeric acid from 0.1 mM to 10 mM.

Antibody Isolation or Purification

[0169] The antibodies or antigen-binding fragments described herein can be separated from a substantial portion of the cellular growth media from which they are recovered to allow for a more purified or isolated form of the antibody or antigen-binding fragment. Described herein are methods to purify or isolate antibodies or antigen-binding fragments, such as human IgM antibodies or antigen-binding fragments, from a solution, *e.g.*, conditioned culture supernatant (CCS). Increasing the purity of a solution containing the antibodies or antigen-binding fragments may be accomplished in a number of ways, including, but not limited to,

dialysis, size exclusion chromatography, centrifugation, ion-exchange chromatography, gradient centrifugation, filtration with a size-exclusion filter, affinity chromatography, immunoaffinity chromatography, and high performance liquid chromatography. The described antibodies or antigen-binding fragments may also be modified, genetically or chemically, to include an affinity tag, such as polyhistidine, that may be used to increase antibody or antigen-binding fragment thereof purity through affinity purification techniques.

[0170] Accordingly, some embodiments include an antibody or antigen-binding fragment thereof that is separated from a solution by affinity chromatography. Some embodiments comprise an antibody or antigen-binding fragment thereof that is substantially separated from a solution by affinity chromatography, wherein affinity for the antibody or antigen-binding fragment thereof is mediated by protein A. Some embodiments comprise an antibody or antigen-binding fragment thereof that is substantially separated from a solution by ion-exchange chromatography. Some embodiments comprise an antibody or antigen-binding fragment thereof that is substantially separated from a solution by cation-exchange chromatography, wherein the cation-exchange chromatography column comprises an acrylamide-dextran copolymer resin. Some embodiments comprise an antibody or antigen-binding fragment thereof that is substantially separated from a solution by hydroxyapatite chromatography. Some embodiments comprise an antibody or antigen-binding fragment thereof that is substantially separated from a solution by hydroxyapatite chromatography, wherein the hydroxyapatite chromatography column incorporates a ceramic calcium phosphate resin.

[0171] The process by which the described antibodies or antigen-binding fragments can be purified or isolated may vary as the requirements for the resulting antibody or antigen-binding fragment thereof may differ depending on the application for which the antibodies or antigen-binding fragments will be used. For example, antibodies or antigen-binding fragments for use in a pharmaceutical composition would require stringent purification, while antibodies or antigen-binding fragments for use in an *in vitro* diagnostic assay can be purified to a lesser degree. Accordingly some embodiments include contacting an antibody or antigen-binding fragment-containing sample with a protein A matrix. Some embodiments include contacting an antibody or antigen-binding fragment-containing sample with an ion-exchange matrix. In one embodiment, the ion-exchange matrix can be a cation-exchange chromatography matrix. In a more preferable embodiment the cation-exchange chromatography matrix can include acrylamide-dextran copolymer resin. Some embodiments include contacting an antibody or antigen-binding fragment-containing sample with a hydroxyapatite matrix. In one embodiment, the hydroxyapatite matrix can include a calcium phosphate resin.

[0172] Serial affinity chromatography is a process that yields an isolated or purified protein. The process may employ two or more rounds of differing affinity chromatography techniques to produce a purified or isolated protein of interest (Friedrichs and Grose, 49 J. Virol. 992 (1984)). Accordingly, in some embodiments an antibody or antigen-binding fragment-containing sample may be contacted with a protein A matrix under conditions that promote binding of antibodies or antigen-binding fragments to the matrix; the matrix may be washed to remove unbound proteins; the material bound by the protein A matrix may be eluted; the eluted material may be contacted by a cation-exchange chromatography matrix under conditions that promote binding of antibodies or antigen-binding fragments to the matrix; the matrix may be washed to remove unbound proteins; the material bound by the cation-exchange chromatography matrix may be eluted; the eluted material may be contacted by a hydroxyapatite resin under conditions that promote binding of antibodies or antigen-binding fragments to the matrix; the matrix may be washed to remove unbound proteins; and the bound material may be eluted. In addition, optional steps may be added to various points in this process to allow for more stringent purification of the antibody or antigen-binding fragment thereof. For example, the antibody or antigen-binding fragment-containing solution may be supplemented with detergent, such as Triton®-X 100 or Tween® 80, to inactivate microbes, such as bacteria, viruses, and parasites.

[0173] Affinity-based chromatography of proteins is a multi-step process that generally involves equilibrating a chromatography column, contacting a sample solution with the matrix of a column, washing away the unbound material in the column, and eluting the desired material. This general process may be repeated one or more times, under varying conditions, to increase the purity of a sample, as needed. In some embodiments a protein A matrix may be used to contact the sample to be purified. In one embodiment, the protein A matrix may be porous, having pores ranging from 1000 to 5000 angstroms in diameter. In another embodiment the protein A matrix has pores with an average diameter of 3000 angstroms.

[0174] In some embodiments the affinity matrix is washed and equilibrated before use. For example, the affinity matrix may be washed with purified water to remove any contaminants. In some embodiments the affinity matrix may be washed with 3 to 10 column volumes of purified water. In another embodiment the affinity matrix may be washed with 5 column volumes of purified water. In some embodiments the affinity matrix may be washed with at least one acidic buffer. For example, the affinity matrix may be washed with 1 to 5 column volumes of 20 mM HCl at pH 1.5. In another embodiment the affinity matrix is washed with 3 column volumes of 20 mM HCl at pH 1.5.

[0175] In addition, the affinity matrix may be washed with at least one additional buffer, such as 6 M guanidine-HCl. Before contacting the affinity matrix with a sample the matrix may be equilibrated with a mildly basic or neutral pH buffer. In one embodiment, the affinity matrix may be equilibrated with 2 to 8 column volumes of a buffer at pH 7.5. In another embodiment the affinity matrix may be equilibrated with 5 column volumes of a 10 mM sodium phosphate buffer containing 200 mM NaCl and 0.01% Tween®-80, at pH 7.5. Sodium phosphate buffers described herein may be prepared from a mixture of monobasic and dibasic sodium phosphate. Mixed in the correct ratios, these solutions may yield phosphate buffers within the range of pH 4 to pH 10.

[0176] Samples to be purified or isolated via contacting the affinity matrix should be substantially free of large particulate matter. Such particulate matter may be removed in a variety of ways, such as by centrifugation or filtration. In another embodiment the conditioned culture supernatant (CCS) described herein is made substantially free of large particulate matter before contact with an affinity matrix. In yet another embodiment the CCS is filtered to remove particulate matter. A filter with an appropriate pore size may be used, such as a filter having pores with an average diameter of 1 μm or a filter having pores with an average diameter of 0.75 μm or a filter having pores with an average diameter of 0.5 μm or a filter having pores with an average diameter of 0.22 μm , or a filter having pores with an average diameter of 0.1 μm .

[0177] Before contacting the affinity matrix with a solution containing a sample to be purified or isolated, the solution may be treated with detergent to inactivate at least one microbial contaminant, such as bacteria, viruses, and parasites. In one embodiment, the CCS is supplemented with detergent to inactivate at least one microbial contaminant. In one embodiment, the detergent is Triton® X-100. The concentration of detergent added to the solution containing a sample to be purified or isolated may vary to meet the needs of a particular purification scheme. In one embodiment, the solution containing a sample to be purified or isolated may have 10% detergent. In another embodiment, the solution containing a sample to be purified or isolated may have 7% detergent. In one embodiment, the solution containing a sample to be purified or isolated may have 5% detergent. In one embodiment, the solution containing a sample to be purified or isolated may have 3% detergent. In one embodiment, the solution containing a sample to be purified or isolated may have 2% detergent. In one embodiment, the solution containing a sample to be purified or isolated may have 1% detergent. In one embodiment, the solution containing a sample to be purified or isolated may have 0.5% detergent. In one embodiment, the solution containing a sample to be purified or isolated may have 0.1% detergent.

[0178] Once applied to the affinity matrix, the sample may be processed at a desired flow rate, for example, to facilitate processing or binding of the sample. While any flow rate may be used, common flow rates are between 1 and 200 cm/h. In some embodiments the flow rate of the sample may be 1 cm/h. In other embodiments the flow rate of the sample may be 10 cm/h. In one embodiment, the flow rate of the sample may be 25 cm/h. In one embodiment, the flow rate of the sample may be 50 cm/h. In another embodiment the flow rate of the sample may be 76 cm/h. In another embodiment the flow rate of the sample may be 100 cm/h. In some embodiments the flow rate of the sample may be 125 cm/h. In another embodiment the flow rate of the sample may be 150 cm/h. In one embodiment, the flow rate of the sample may be 175 cm/h. In one embodiment, the flow rate of the sample may be 200 cm/h.

[0179] After the affinity matrix has been contacted by the sample material, unbound sample material may be removed by washing with a buffer. In one embodiment, the affinity matrix may be washed with a buffer with an acidic pH. In one embodiment, the affinity matrix may be washed with a buffer with a neutral pH. In one embodiment, the affinity matrix may be washed with a buffer with a basic pH. In another embodiment the affinity matrix may be washed with 5 to 15 column volumes of a 10 mM sodium phosphate buffer containing 200 mM NaCl and 0.01% Tween®-80, at pH 7.5.

[0180] After unbound proteins have been removed from the affinity matrix by washing, proteins bound to the affinity matrix may be eluted by washing the affinity matrix with an elution buffer. The elution buffer should be selected carefully to assure that it will disrupt the interaction between the protein of interest, *e.g.*, a human IgM antibody, and the matrix, but will not denature or otherwise deteriorate the condition of the protein of interest. In one embodiment, the affinity matrix may be washed with a buffer with an acidic pH to elute the protein of interest. In one embodiment, the affinity matrix may be washed with a buffer with a neutral pH to elute the protein of interest. In one embodiment, the affinity matrix may be washed with a buffer with a basic pH to elute the protein of interest. Some embodiments include a buffer having sodium phosphate and magnesium chloride for eluting material bound to a chromatography column, where up to 10 column volumes of this buffer are used to elute bound material from the column. In some embodiments, a buffer having between 5 and 10 mM sodium phosphate and 1 and 5 M magnesium chloride may be used for eluting material bound to an affinity matrix. Another embodiment, includes 3 column volumes of a buffer comprising 5 mM sodium phosphate and 3 M magnesium chloride to elute material bound to an affinity matrix. In some embodiments IgM may be eluted from a protein A matrix using a 5 mM sodium phosphate buffer supplemented with 3 M MgCl₂ at pH 6.8.

[0181] Disclosed herein are methods for purifying or isolating proteins, *e.g.*, IgM antibodies or antigen-binding fragments, using ion-exchange chromatography. Ion-exchange chromatography of proteins is a multi-step process that generally involves equilibrating a chromatography column, contacting a sample solution with the matrix of a column, washing away the unbound material in the column, and eluting the desired material. This general process may be repeated one or more times, under varying conditions, to increase the purity of a sample, as needed. In some embodiments the cation exchange chromatography matrix may be an acrylamide-dextran copolymer resin. In one embodiment, the cation exchange chromatography matrix may be MacroCap™ SP resin.

[0182] In some embodiments the ion-exchange matrix is washed and equilibrated before use. For example, the ion-exchange matrix may be washed with purified water to remove any contaminants. In one embodiment, the ion-exchange matrix may be washed with 1 to 10 column volumes of purified water. In one embodiment, the ion-exchange matrix may be washed with 2 column volumes of purified water. In some embodiments the ion-exchange matrix may be washed with at least one buffer. For example, the ion-exchange matrix may be washed with 1 to 5 column volumes of a buffer having 0.5 M NaOH. In another embodiment the ion-exchange matrix is washed with 3 column volumes of a buffer having 0.5 M NaOH. In some embodiments the ion-exchange matrix is washed with 3 column volumes of a buffer having 2 M NaCl. Before contacting the ion-exchange matrix with a sample the matrix may be equilibrated with an acidic, basic, or neutral pH buffer. In one embodiment, the ion-exchange matrix may be equilibrated with 2 to 8 column volumes of a buffer at pH 6.8. In one embodiment, the ion-exchange matrix may be equilibrated with 5 column volumes of a 10 mM sodium phosphate buffer, containing 75 mM NaCl and 0.01% Tween®-80, at pH 6.8.

[0183] Samples to be purified or isolated via contacting the ion-exchange matrix should be substantially free of large particulate matter. Such particulate matter may be removed in a variety of ways, such as by centrifugation or filtration. In some embodiments the conditioned culture supernatant (CCS) described herein is made substantially free of large particulate matter before having contact with an ion-exchange matrix. In another embodiment the CCS is filtered to remove particulate matter. A filter with any pore size may be used, such as a filter having pores with an average diameter of 1 µm or a filter having pores with an average diameter of 0.75 µm or a filter having pores with an average diameter of 0.5 µm or a filter having pores with an average diameter of 0.22 µm, or a filter having pores with an average diameter of 0.1 µm.

[0184] Before contacting the ion-exchange matrix with a solution containing a sample to be purified, the solution may be treated with detergent to inactivate at least one microbial

contaminant, such as bacteria, viruses, and parasites, or to help maintain the target protein in a soluble state. In some embodiments the sample may be supplemented with detergent to inactivate at least one microbial contaminant or to help maintain the target protein in a soluble state. In one embodiment, the detergent is Triton® X-100. In one embodiment, the detergent is Tween®-80. The concentration of detergent added to the solution containing a sample to be purified or isolated may vary to meet the needs of a particular purification scheme. In one embodiment, the solution containing a sample to be purified or isolated may have 10% detergent. In one embodiment, the solution containing a sample to be purified or isolated may have 7% detergent. In one embodiment, the solution containing a sample to be purified or isolated may have 5% detergent. In one embodiment, the solution containing a sample to be purified or isolated may have 3% detergent. In one embodiment, the solution containing a sample to be purified or isolated may have 2% detergent. In one embodiment, the solution containing a sample to be purified or isolated may have 1% detergent. In one embodiment, the solution containing a sample to be purified or isolated may have 0.5% detergent. In one embodiment, the solution containing a sample to be purified or isolated may have 0.1% detergent.

[0185] Once applied to the ion-exchange matrix, the sample may be processed at a desired flow rate, for example, to facilitate processing or binding of the sample. While any flow rate may be used, common flow rates are between 1 and 200 cm/h. In one embodiment, the flow rate of the sample may be 1 cm/h. In one embodiment, the flow rate of the sample may be 10 cm/h. In one embodiment, the flow rate of the sample may be 25 cm/h. In one embodiment, the flow rate of the sample may be 50 cm/h. In one embodiment, the flow rate of the sample may be 76 cm/h. In one embodiment, the flow rate of the sample may be 100 cm/h. In one embodiment, the flow rate of the sample may be 125 cm/h. In one embodiment, the flow rate of the sample may be 150 cm/h. In one embodiment, the flow rate of the sample may be 175 cm/h. In another embodiment the flow rate of the sample may be 200 cm/h.

[0186] After the ion-exchange matrix has been contacted by the sample material, unbound sample material may be removed by washing with a buffer. In one embodiment, the ion-exchange matrix may be washed with a buffer with an acidic pH. In one embodiment, the ion-exchange matrix may be washed with a buffer with a neutral pH. In one embodiment, the ion-exchange matrix may be washed with a buffer with a basic pH. In one embodiment, the ion-exchange matrix may be washed with 5 to 15 column volumes of a buffer containing 10 mM sodium phosphate, 75 mM NaCl, and 0.01% Tween®-80, at pH 6.8.

[0187] After unbound proteins have been removed from the ion-exchange matrix by washing, proteins bound to the ion-exchange matrix may be eluted by washing the ion-exchange

matrix with an elution buffer. The elution buffer should be selected carefully to assure that it will disrupt the interaction between the protein of interest, *e.g.*, a human IgM antibody, and the matrix, but will not denature or otherwise deteriorate the condition of the protein of interest. In one embodiment, the ion-exchange matrix may be washed with a buffer with an acidic pH to elute the protein of interest. In one embodiment, the ion-exchange matrix may be washed with a buffer with a neutral pH to elute the protein of interest. In one embodiment, the ion-exchange matrix may be washed with a buffer with a basic pH to elute the protein of interest. Some embodiments include a buffer having sodium phosphate and sodium chloride for eluting material bound to a chromatography column, where up to 10 column volumes of this buffer are used to elute bound material from the column. In some embodiments, a buffer having between 5 and 50 mM sodium phosphate and 150 and 500M sodium chloride may be used for eluting material bound to an ion-exchange matrix. Another embodiment includes 4 column volumes of a buffer comprising 10 mM sodium phosphate buffer containing 200 mM NaCl to elute material bound to an ion-exchange matrix. In some embodiments IgM may be eluted from a MacroCap™ SP matrix using a 10 mM sodium phosphate buffer containing 200 mM NaCl and 0.01% Tween®-80, at pH 6.8.

[0188] Disclosed herein are methods for purifying or isolating proteins, *e.g.*, IgM antibodies or antigen-binding fragments, using hydroxyapatite chromatography. Hydroxyapatite chromatography of proteins is a multi-step process that generally involves equilibrating a chromatography column, contacting a sample solution with the matrix of a column, washing away the unbound material in the column, and eluting the desired material. This general process may be repeated one or more times, under varying conditions, to increase the purity of a sample, as needed. In some embodiments the hydroxyapatite chromatography matrix may be a calcium phosphate matrix. In one embodiment, the hydroxyapatite chromatography matrix may be CHT® II Ceramic Hydroxyapatite, 80 µm bead size.

[0189] In some embodiments the hydroxyapatite matrix is hydrated before use. In one embodiment, the hydroxyapatite matrix is hydrated with a solution having 200 mM potassium phosphate, at pH 9.0, using 0.54g of dry matrix for each mL of desired column-bed volume. Before contacting the hydroxyapatite matrix with a sample the matrix may be equilibrated with an acidic, basic, or neutral pH buffer. In one embodiment, the hydroxyapatite matrix may be equilibrated with 2 to 8 column volumes of a buffer at pH 6.8. In another embodiment the hydroxyapatite matrix may be equilibrated with 5 column volumes of a buffer containing 10 mM sodium phosphate, 100 mM NaCl, and 0.01% Tween®-80, at pH 6.8.

[0190] Samples to be purified or isolated via contacting the hydroxyapatite matrix should be substantially free of large particulate matter. Such particulate matter may be removed in a variety of ways, such as by centrifugation or filtration. In one embodiment, the conditioned culture supernatant (CCS) is made substantially free of large particulate matter before having contact with a hydroxyapatite matrix. In one embodiment, the CCS is filtered to remove particulate matter. A filter with any pore size may be used, such as a filter having pores with an average diameter of 1 μm or a filter having pores with an average diameter of 0.75 μm or a filter having pores with an average diameter of 0.5 μm or a filter having pores with an average diameter of 0.22 μm , or a filter having pores with an average diameter of 0.1 μm .

[0191] Before contacting the hydroxyapatite matrix with a solution containing a sample to be purified, the solution may be treated with detergent to inactivate at least one microbial contaminant, such as bacteria, viruses, and parasites, or to help maintain the target protein in a soluble state. In some embodiments the sample may be supplemented with detergent to inactivate at least one microbial contaminant or to help maintain the target protein in a soluble state. In one embodiment, the detergent is Triton® X-100. In one embodiment, the detergent is Tween®-80. The concentration of detergent added to the solution containing a sample to be purified or isolated may vary to meet the needs of a particular purification scheme. In one embodiment, the solution containing a sample to be purified or isolated may have 10% detergent. In one embodiment, the solution containing a sample to be purified or isolated may have 7% detergent. In one embodiment, the solution containing a sample to be purified or isolated may have 5% detergent. In one embodiment, the solution containing a sample to be purified or isolated may have 3% detergent. In one embodiment, the solution containing a sample to be purified or isolated may have 2% detergent. In one embodiment, the solution containing a sample to be purified or isolated may have 1% detergent. In one embodiment, the solution containing a sample to be purified or isolated may have 0.5% detergent. In one embodiment, the solution containing a sample to be purified or isolated may have 0.1% detergent.

[0192] Once applied to the hydroxyapatite matrix, the sample may be processed at a desired flow rate, for example, to facilitate processing or binding of the sample. While any flow rate may be used, common flow rates are between 1 and 200 cm/h. In one embodiment, the flow rate of the sample may be 1 cm/h. In one embodiment, the flow rate of the sample may be 10 cm/h. In one embodiment, the flow rate of the sample may be 25 cm/h. In one embodiment, the flow rate of the sample may be 50 cm/h. In one embodiment, the flow rate of the sample may be 76 cm/h. In one embodiment, the flow rate of the sample may be 100 cm/h. In one embodiment, the flow rate of the sample may be 125 cm/h. In one embodiment, the flow rate of the sample

may be 150 cm/h. In one embodiment, the flow rate of the sample may be 175 cm/h. In one embodiment, the flow rate of the sample may be 200 cm/h.

[0193] After the hydroxyapatite matrix has been contacted by the sample material, unbound sample material may be removed by washing with a buffer. In one embodiment, the hydroxyapatite matrix may be washed with a buffer with an acidic pH. In one embodiment, the hydroxyapatite matrix may be washed with a buffer with a neutral pH. In one embodiment, the hydroxyapatite matrix may be washed with a buffer with a basic pH. In one embodiment, the hydroxyapatite matrix may be washed with 5 to 15 column volumes of a buffer containing 10 mM sodium phosphate, 100 mM NaCl, and 0.01% Tween®-80, at pH 6.8.

[0194] After unbound proteins have been removed from the hydroxyapatite matrix by washing, proteins bound to the hydroxyapatite matrix may be eluted by washing the hydroxyapatite matrix with an elution buffer. The elution buffer should be selected carefully to assure that it will disrupt the interaction between the protein of interest, *e.g.*, a human IgM antibody, and the matrix, but will not denature or otherwise deteriorate the condition of the protein of interest. In one embodiment, the hydroxyapatite matrix may be washed with a buffer with an acidic pH to elute the protein of interest. In one embodiment, the hydroxyapatite matrix may be washed with a buffer with a neutral pH to elute the protein of interest. In one embodiment, the hydroxyapatite matrix may be washed with a buffer with a basic pH to elute the protein of interest. Some embodiments include a buffer having sodium phosphate and sodium chloride for eluting material bound to a hydroxyapatite column, where up to 10 column volumes of this buffer are used to elute bound material from the column. In some embodiments, a buffer having between 150 and 500 mM sodium phosphate and 50 and 200 M sodium chloride may be used for eluting material bound to a hydroxyapatite matrix. In one embodiment, 4 column volumes of a buffer comprising 175 mM sodium phosphate buffer containing 100 mM NaCl may be used to elute material bound to an hydroxyapatite matrix. In another embodiment IgM may be eluted from a CHT® II Ceramic Hydroxyapatite matrix using a buffer containing 175 mM sodium phosphate, 100 mM NaCl, and 0.01% Tween®-80, at pH 6.8.

[0195] Buffers of varying composition, with respect to salts used, salt concentration, detergents used, detergent concentration, pH, and the like, may be used to substantially perform the methods described herein. Accordingly, such variations are contemplated to be within the scope of the disclosure provided, which is not intended to be exclusive or limiting.

[0196] Embodiment C1 provides a method of purifying or isolating a protein, the method comprising applying a solution comprising the protein to an affinity chromatography column, applying eluate from the affinity chromatography column to a cation exchange

chromatography column, and applying eluate from the cation exchange chromatography column to a hydroxyapatite chromatography column, and obtaining the eluate from the hydroxyapatite chromatography column.

[0197] Embodiment C2 provides the method of embodiment C1, wherein the affinity chromatography column comprises at least one substrate coated with protein A.

[0198] Embodiment C3 provides the method of embodiment C2, wherein the at least one substrate coated with protein A is porous.

[0199] Embodiment C4 provides the method of embodiment C3, wherein the porous substrate comprises pores of 3000 angstroms.

[0200] Embodiment C5 provides the method of embodiment C1, wherein the at least one substance coated with protein A binds a protein.

[0201] Embodiment C6 provides the method of embodiment C5, wherein the protein bound by at least one substance coated with protein A is eluted with a solution comprising magnesium chloride.

[0202] Embodiment C7 provides the method of embodiment C6, wherein the solution comprising magnesium chloride is a solution of 1M to 5M magnesium chloride.

[0203] Embodiment C8 provides the method of embodiment C6, wherein the solution comprising magnesium chloride is a solution of 3M magnesium chloride.

[0204] Embodiment C9 provides the method of embodiment C5, wherein the protein bound by the at least one substance coated with protein A is an antibody.

[0205] Embodiment C10 provides the method of embodiment C9, wherein the antibody is IgM.

[0206] Embodiment C11 provides the method of embodiment C1, wherein the cation exchange chromatography column comprises an acrylamide-dextran copolymer resin.

[0207] Embodiment C12 provides the method of embodiment C11, wherein the acrylamide-dextran copolymer resin binds a protein.

[0208] Embodiment C13 provides the method of embodiment C12, wherein the protein bound by the acrylamide-dextran copolymer resin is eluted with a solution comprising 10 mM sodium phosphate, 200 mM sodium chloride, and 0.01% polysorbate 80.

[0209] Embodiment C14 provides the method of embodiment C13, wherein the protein bound by the acrylamide-dextran copolymer resin is an antibody.

[0210] Embodiment C15 provides the method of embodiment C14, wherein the antibody is IgM.

[0211] Embodiment C16 provides the method of embodiment C1, wherein the hydroxyapatite chromatography column comprises a calcium phosphate resin.

[0212] Embodiment C17 provides the method of embodiment C16, wherein the calcium phosphate resin binds a protein.

[0213] Embodiment C18 provides the method of embodiment C17, wherein the protein bound by the calcium phosphate resin is eluted with a solution comprising 175 mM sodium phosphate, 100 mM sodium chloride, and 0.01% polysorbate 80.

[0214] Embodiment C19 provides the method of embodiment C18, wherein the protein bound by the calcium phosphate resin is an antibody.

[0215] Embodiment C20 provides the method of embodiment C19, wherein the antibody is IgM.

[0216] Embodiment C21 provides a method of purifying or isolating a protein, comprising

- a. applying a solution containing the protein to an affinity chromatography column comprising at least one substrate coated with protein A, and eluting any proteins bound by the at least one substance coated with protein A with a solution comprising 3M magnesium chloride;
- b. applying the eluate from the affinity chromatography column to a cation exchange chromatography column comprising an acrylamide-dextran copolymer resin, and eluting any proteins bound by the acrylamide-dextran copolymer resin with a solution comprising 10 mM sodium phosphate, 200 mM sodium chloride, and 0.01% polysorbate 80;
- c. applying the eluate from the cation exchange chromatography column to a hydroxyapatite comprising a calcium phosphate resin, and eluting any proteins bound by the calcium phosphate resin with a solution comprising 175 mM sodium phosphate, 100 mM sodium chloride, and 0.01% polysorbate 80.

[0217] The following examples are provided to describe the embodiments described herein with greater detail. They are intended to illustrate, not to limit, the embodiments.

EXAMPLE 1 GD-2 Specificity of CV-3-Derived IgM

[0218] A pool of human lymphoblast cells transformed with Epstein-Barr Virus (HLP) (Cahan, *et al.*) was tested for the ability to produce GD2-specific IgM antibodies. The cells were

cultured in RMPI complete media (10% heat-inactivated FBS, 1% L-Glu, 1% antibiotics (Sigma)). Spent growth media was collected at passages two through four and assayed for IgM concentration and reactivity with GD2. The HLP produced 6-8 $\mu\text{g/mL}$ IgM, (quantified by ELISA), after a culture of 2.5×10^5 cells/mL was grown for 3 days.

[0219] To determine whether the IgM produced by the HLP was specific for GD2, spent media from each culture was analyzed by ELISA. Briefly, gangliosides were coated to ELISA plates by evaporating 200 μl of ethanol containing 25 ng of GD2, GD1a, GM2 or GM3. Plates were then blocked and binding of IgM-containing cell culture media or human IgM were assessed according to regular ELISA procedures. Spent media from the HLP (concentrated either 8.1-fold or 14.2-fold) contained GD2-reactive IgM (Figure 1A); however, the antibodies also recognized gangliosides GD1a, GM2, and GM3 (Figure 1B). These data indicate that IgM produced by the HLP is polyreactive.

[0220] FACS analysis was conducted to determine whether IgM produced by the HLP could bind GD2 expressed on the surface of melanoma cells. Briefly, 1205LU cells, a human melanoma cell line that expresses GD2 on its surface, were incubated with spent media from HLP cultures, washed, and then labeled with secondary antibody FITC-goat-anti-human Ig (Jackson Laboratories). As shown in Figure 2, only culture media from very early passages of the HLP (passage 2) produced IgM that could recognize cell surface GD2 (Figure 2B). However, this activity was quickly lost after subculturing the HLP for a few weeks (Figure 2C). These results suggest that the cells producing specific GD2-reactive IgM are slow growers in the HLP and are quickly overpopulated by non-producers or nonspecific antibody producers, or that the production of this IgM is not stable. The murine anti-GD2 IgM antibody MAb-126 (ATCC# HB-8568TM) was used as a positive control and demonstrated robust GD2-specific binding (Figure 2D).

EXAMPLE 2

Hybridomas Derived from the HLP Produce GD-2 Specific IgM that Mediates CDC Activity on GD2-Positive Cells

[0221] In order to generate EBV-null hybridoma lines that produce anti-GD2 specific IgM, cells from early passages of the HLP were fused with fusion partners A6 (ATCC CRL-8192) or K6H6/B5 (ATCC CRL-1823) to form hybridomas. Due to the apparently low abundance of GD2-specific subclones in the HLP, 80,000 clones were screened by GD2-specific ELISA. Approximately 5–10% of antibody preparations from these hybridoma lines showed positive GD2 reactivity. However, over 90% of the GD2 positive clones showed reactivity to

multiple gangliosides when screened for specificity using a panel of gangliosides (GD1, GM2 and GM3) (Figure 3A – clones 5D7 and 10B4). Moreover, the majority of polyreactive antibodies cross-reacted to a panel of non-related control proteins (data not shown). These results suggest that the lymphoblast cells producing GD2-specific IgM are rare in the HLP.

[0222] Subcloning and screening efforts led to the isolation of GD2-specific IgM-producing hybridoma cell lines. Two GD2 specific hybridoma lines, 3B2 and 1470, were isolated. Cultures of both cell lines were seeded at $0.3\text{--}0.4 \times 10^6/\text{mL}$ of cells of seeding density and split every 3 to 4 days, which yielded $3\text{--}8 \mu\text{g/mL}$ of IgM in spent medium. Ganglioside-specific ELISA revealed positive reactivity with GD2 only and no reactivity to three control gangliosides (GD1a, GM2, and GM3) (Figure 3A and B), suggesting that these cells produce GD2-specific antibodies. In addition, FACS analysis showed positive staining of 1205LU human melanoma cells with 3B2 and 1470 antibodies, suggesting that these antibodies can recognize GD2 on the cell surface (Figure 4). Molecular analyses of clones 3B2 and 1470 were performed to determine clone origin. Light chain specific ELISA showed that both clones secreted IgM containing a kappa light chain (data not shown). Sequence analysis of cDNA from these cells revealed identical heavy and light chain sequences, suggesting that the two clones were derived from the same HLP lymphoblast clone (data not shown).

[0223] Partially purified antibody from the 3B2 culture was used to assess the ability of this antibody to mediate complement-dependent cytotoxicity (CDC) on GD2 expressing melanoma cell lines, as this activity is largely dependent on the conformation that the antibody assumes when bound to its antigen (Janeway et al., Immunobiology, 9-12 (5th ed. 2001)). Melanoma target cell lines (GD2-positive cell lines LF0023 and M14, and GD2-negative cell lines PM0496 and JS0592) were cultured in RPMI1640 supplemented with 10% FBS, 2 mM L-Glutamine, non-essential amino acids, and 6 μM HEPES and harvested with trypsin before use. Target cells were incubated with 3B2 IgM-containing supernatant in the presence of 20% human serum. After incubation for 1 hour at 37°C, live cells were identified with Cell Titer Glo® reagent, (Promega Corp., Madison, WI). Percent killing was determined as the ratio of signal from treated versus untreated cells. The data in Figure 5 suggest that IgM produced by the 3B2 clone can elicit a potent CDC reaction on certain GD2 positive melanoma cell lines.

[0224] Molecular analyses were performed to determine whether clones 3B2 and 1470 were infected with EBV. Six PCR primer pairs ((EBNA2-1141f (SEQ ID NO: 45) and EBNA2-1440r (SEQ ID NO: 46); EBV2001f (SEQ ID NO: 47) and EBV2622r (SEQ ID NO: 48); EBV1901F (SEQ ID NO: 49) and EBV2822R (SEQ ID NO: 50); EBV169461f (SEQ ID NO: 51) and EBV170100r (SEQ ID NO: 52); EBV169480f (SEQ ID NO: 53) and EBV170080r (SEQ

ID NO: 54); EBV8491F (SEQ ID NO: 55) and EBV9020r (SEQ ID NO: 56)) were designed to verify the existence and intactness of the EBV genome. One of the primer pairs amplified the open reading frames of EBNA-2; two overlapping sets each amplified the 5' and 3' end of the EBV genome; and one pair amplified the origin of replication. Initially, both 3B2 and 1470 clones were EBV positive as demonstrated by genomic PCR (data not shown). However, further subcloning by limiting dilution plating at less than one cell per well generated subclones of the 3B2 and 1470 lines that did not contain the EBV genome, as determined by EBV-specific PCR analysis (data not shown).

[0225] EBV-negative 3B2 subclones (AB527-HYB-3B2-EBVnull) have been placed with the Amer. Type Cult. Coll. (10801 University Blvd., Manassas, Virginia 20110-2209) on July 16, 2008 and have been assigned Accession No. PTA-9376. AB527-HYB-3B2-EBVnull subclones were subjected to a second round of subcloning to identify clones that produced a single form of immunoglobulin J-chain, as the original hybridoma may produce human or mouse J-chain. To identify clones possessing a single form of J-chain, or J-chain-null clones, RT-PCR was performed using primers specific for human J-chain (hu-285F (SEQ ID NO: 57) and hu-418R (SEQ ID NO: 58)), or mouse J-chain (m-230F (SEQ ID NO: 59) and m-370R (SEQ ID NO: 60)). The results indicated that the 3B2 subclone, AB527-HYB-3B2-3C9, is EBV null, human J-chain null, murine J-chain positive and secretes approximately 30 mg/L of monoclonal, GD2-specific IgM in static culture.

EXAMPLE 3

Production of a Transfectoma Cell Line Expressing GD2-Specific IgM

[0226] Total RNA isolated from AB527-HYB-3B2-3C9, the EBV null, human J-chain null, murine J-chain positive hybridoma, was reverse transcribed into cDNA and used as template for the PCR amplification of the IgM heavy chain, using primers corresponding to SEQ ID NOs: 33 and 35, and light chain, using primers corresponding to SEQ ID NOs: 36 and 38. The heavy chain nucleotide sequence amplified from AB527-HYB-3B2-3C9 encodes a polypeptide having the amino acid sequence of SEQ ID NO: 40, and the light chain nucleotide sequence amplified from AB527-HYB-3B2-3C9 encodes a polypeptide having the amino acid sequence of SEQ ID NO: 42. A second round of PCR amplification was performed to add a human leader sequence, along with 5' and 3' restriction endonuclease recognition sites were introduced in order to facilitate cloning of the amplicons into pEE6.4 (heavy chain) and pEE14.4 (light chain), both vectors are supplied as part of Lonza Biologics Glutamine Synthetase

Expression System (GS System). The second round PCR amplification of the IgM heavy chain was conducted using primers corresponding to SEQ ID NOs: 34 and 35, while the light chain amplification used primers corresponding to SEQ ID NOs: 37 and 38.

[0227] PCR amplified light chain cDNA was digested with HindIII and EcoRI and ligated into similarly cut pEE14.4, while heavy chain cDNA was digested with HindIII and MfeI and ligated into pEE6.4 digested with HindIII and EcoRI (MfeI and EcoRI leave compatible “sticky” ends). After verification of the nucleotide sequence of the heavy and light chains, the heavy chain expression cassette was introduced into pEE14.4-AB527-LC as a NotI/PvuII fragment to construct the final expression vector pEE14.4-AB527-LC-HC (p0311) (Figure 6). pEE14.4 is a 10 Kb vector which contains a glutamine synthase mini gene, allowing for selection of stable transfectants in the presence of the GS inhibitor L-methionine sulfoximine (MSX). Transcription of the cDNA inserts in this vector is controlled by the human CMV immediate early promoter, upstream of the cDNA insert is the hCMV-MIE 5’ untranslated region including introns 1, downstream of the cDNA insert is the SV40 polyadenylation signal allowing for efficient polyadenylation of the transcript (Figure 7).

[0228] CHO cells were transfected with linearized, double-gene vector p0311 by electroporation to allow for expression of recombinant GD2-specific IgM. Following transfection the cells were plated in 96-well plates using non-selective medium containing dialyzed fetal bovine serum (dFBS) and glutamine synthase supplement. The following day selective medium containing dFBS, glutamine synthase supplement and MSX (10 μ M final concentration) was added to each well, enabling selection of cells containing the expression vector. Only colonies perceived to have come from a single transfected cell were pursued.

[0229] Transfectants were screened for the presence of human IgM by ELISA. Colonies were expanded in serum-containing selection medium and surviving colonies were examined by quantitative secondary assay to identify the highest-producing colonies. The highest-producing transfectants were adapted to chemically defined IS-CHO-CD™ medium (Irvine Scientific, Santa Ana, CA) and seeded at 2,500 cells/well in flat bottom 96 well plates (80 plates) in media containing 10 μ M MSX. Drug-resistant colonies were analyzed for IgM production by ELISA. Colonies positive for IgM production were scaled up and analyzed for antibody productivity in subsequent secondary (6 well) and tertiary (20 mL shake flask) assays. This resulted in the identification of an IgM-producing CHO-K1SV transfectoma clone, 127C8. The 127C8 clone does not express J-chain, therefore, the recombinant AB527 IgM produced by this cell line is J-chain-deficient. The amino acid sequences of the heavy and light chains of AB527 are represented by SEQ ID NOs: 40 and 42, respectively.

EXAMPLE 4

AB527 Binds Specifically to GD2 *In Vitro*

[0230] Several experiments were conducted to determine whether recombinant AB527 could bind to purified GD2 *in vitro*. Initially, surface plasmon resonance experiments were performed to not only determine whether recombinant AB527 would bind to GD2, but also to determine whether it exhibited affinity for GM2 or GD3 (Figure 8). The experiments were performed as follows: 40 μ L of 0.3 mg/mL solutions of GD2 (BioDesign), GD3 (HyTest), and GM2 (USBiological) in 10 mM HEPES, 150 mM NaCl, 3 mM EDTA (HBS-E) containing 30% ethanol were injected over flow cells 2, 3, and 4, respectively, of a CM5 chip at a flow rate of 5 μ L/minute using a BIAcore 3000 instrument. Five repeated injections of 20 μ L 10 mM NaOH were done following immobilization of each ganglioside until a stable baseline was achieved. Next, 80 μ L of 200 nM recombinant AB527 in HBS-E buffer was injected over all 4 flow cells (flow cell 1 was used as a reference flow cell) at a flow rate of 20 μ L/minute. Dissociation of recombinant AB527 was followed for 6 minutes. The chip surface was regenerated with 50 μ L of 10 mM NaOH.

[0231] To better characterize the binding kinetics of AB527, additional surface plasmon resonance experiments were conducted using serial dilutions of AB527 and GD2 immobilized on a BIAcore CM5 chip. Briefly, two-fold serial dilutions of AB527, starting at 800 nM, were injected over a GD2-containing CM5 chip, described above, at a flow rate of 20 μ L/minute in HBS-E buffer. The chip surface was regenerated between each cycle with 50 μ L of 10 mM NaOH. Equilibrium binding at each concentration (R_{eq}) was determined for each concentration tested in BIAevaluation software and a plot of R_{eq} vs. concentration was then done. The resulting data were fitted to a non-linear steady-state binding model using GraphPad Prism® software. When the R_{eq} is plotted as a function of AB527 concentration, the steady state K_D , or the concentration of AB527 at which one-half maximal binding is achieved, can be determined (Figure 9). For AB527 binding to GD2, the steady state K_D value is 4.5×10^{-9} M.

[0232] The specificity of recombinant AB527 was also tested by immunostaining various gangliosides separated by thin layer chromatography (TLC). The following gangliosides were used in the immunoTLC analysis: GM mix (Matreya, 0.5 mg/mL in ethanol contains GM1, GM2, and GM3), GD mix (Matreya, 0.5 mg/mL in ethanol, contains GD1a, GD1b, and GD3), GM4 (Matreya, 1 mg/mL in ethanol), GD2 (BioDesign, 1 mg/mL in ethanol), and GM3 (USBiological, 1 mg/mL in ethanol). A total of 2 μ L of each ganglioside was spotted on to a

5x10 cm silica TLC plate (EMD Chemicals, Inc.) and developed in a solution having a ratio of 58:37:8 of CHCl_3 , MeOH, and 0.1M CaCl_2 . TLC plates were stained for total ganglioside by spraying with 0.1% orcinol in 1.36N H_2SO_4 and heating at 100 °C. For immunoTLC analysis, developed plates were first treated with 0.1% polyisobutylmethacrylate in H_2O , and then dried. The plates were then blocked for 1 hour with 1% BSA in 1x phosphate-buffered saline (PBS). The blocked plate was then incubated in 2 $\mu\text{g/mL}$ AB527 in 1x PBS containing 1% BSA overnight at 4 °C. The plates were subsequently washed with 1x PBS, then incubated with goat anti-human IgG/IgM-horseradish peroxidase conjugate diluted 1:4000 in PBS containing 1% BSA. Plates were washed, and bound antibody was detected by incubating the plates in 0.3 mg/mL 4-chloro 1-naphthol and 0.03% H_2O_2 . The comparative staining results are shown in Figure 10.

Example 5

AB527 Labels GD2-Expressing Cells

[0233] The ability of AB527 to specifically label GD2-expressing melanoma cell lines was assessed. GD2-positive lines: M14, M0023, M101, M18, M10-Vac, and PM0496, and GD2-negative lines: M21, M238, IMCD0023, MG1055 were cultured separately in RPMI1640 supplemented with 10% FBS, 2 mM L-Glutamine, non-essential amino acids, and 6 μM HEPES and harvested with trypsin. Trypsinized cells were incubated with either 10 $\mu\text{g/mL}$ of control IgM (human IgM, Pierce cat# 31146) (column I) or 10 $\mu\text{g/mL}$ of AB527 (column II) for 1 hour on ice. After 3 washes with 1x PBS, cells were incubated with 10 $\mu\text{g/mL}$ FITC-labeled goat anti human Ig (H+L) antibody (Southern Biotech, Birmingham, AL) for 1 hour on ice, followed by either flow cytometry (columns I and II, respectively) or light microscopy and immunofluorescence microscopy analysis (columns III and IV, respectively). As shown in Figure 11, AB527 labeled cells express GD2 on their surface (column II). Conversely, none of the cell lines examined showed staining with non-GD2-specific human IgM (column I). Immunofluorescent studies showed a direct correlation with flow cytometry data (column IV), light microscopy images are also shown for comparative purposes (column III).

[0234] Cell-based ELISAs were performed to assess AB527 binding to the GD2-expressing cell lines M14, LLC-MK2, EL4 cells and to CHO-K1 cells (negative control), which provided the data to perform corresponding Scatchard analyses. Figure 12 provides a representative set of Scatchard data for EL4 cells. AB527 binding to M14 and LLC-MK2 cells was determined to be nearly identical (0.13 ± 0.02 nM and 0.15 ± 0.04 nM, respectively). Binding to EL4 cells was somewhat lower (0.93 ± 0.16 nM). This difference may be due to the markedly

lower expression of GD2 observed on EL4 cells relative to M14 and LLC-MK2 (data not shown). For these experiments, M14, LLC-MK2, EL4, and CHO-K1 cells were obtained from the American Tissue Type Collection (ATCC, Manassas, VA). M14, EL4, and CHO-K1 cells were cultured in RPMI supplemented with 10% fetal bovine serum. LLC-MK2 cells were cultured in Medium 199 supplemented with 1% horse serum. Cells were suspended in 1X PBS and added to a black, U-bottomed microtiter plate (Grenier cat. # 665209) at 2.5×10^6 cells/well, then pelleted at 1500 rpm for 5 minutes at room temperature. Supernatant was discarded, and cells were resuspended in a solution containing AB527 starting at 20 $\mu\text{g/mL}$ serially diluted 1:2 in PBS containing 2% BSA. Cells were incubated with antibody for 1 hr at room temperature. Cells were washed three times with 200 μL PBS containing 2% BSA and 0.05% Tween®-20, centrifuging at 1500 rpm for 5 minutes to collect cells between wash steps. Cells were then incubated in a PBS solution containing 1 $\mu\text{g/mL}$ horseradish peroxidase conjugated goat anti-human IgG/IgM (Jackson ImmunoResearch) and 2% BSA for 1 hr at room temperature. Cells were washed as above. Bound AB527 was detected using QuantaBlu™ fluorescent substrate (Pierce), with excitation/emission wavelengths of 325 nm/420 nm, in a SpectraMax® M5 multimodal plate reader (Molecular Devices). Raw data was plotted in SoftMax Pro® (ver. 5.2) and fitted to a 4-parameter model. Data analysis was performed in GraphPad Prism® software (ver. 4.03). Non-specific background was first subtracted (binding to CHO-K1 cells), and data were then analyzed by Scatchard analysis.

[0235] Immunohistochemical (IHC) experiments were conducted to determine whether AB527 exhibited cross-reactivity to melanoma cells that do not express GD2. Purified AB527, as well as a non-specific human IgM (Pierce cat# 31146), were biotinylated using NHS ester coupling chemistry, at a molar ratio of 50:1 biotinylation reagent (EZ-Link™ sulfo-NHS-LC-biotin, Pierce cat# 21335) in phosphate-buffered saline, according to the provided instructions, and used to stain various GD2-positive (M14) or GD2-negative (MG1055, RPMI7951 and JS0592) cell lines. Briefly, each melanoma cell line was seeded on chamber slides and cultured overnight in RPMI1640 supplemented with 10% FBS, 2 mM L-Glutamine, non-essential amino acids, and 6 μM HEPES. Cells were washed once with 1x PBS and formalin-fixed. Fixed cells were incubated overnight with 1 $\mu\text{g/mL}$ biotinylated AB527 or biotinylated nonspecific human IgM (negative control) and labeled cells were detected using BioGenex Super Sensitive™ Link-Label IHC Detection System. As shown in Figure 13A, AB527 specifically binds to GD2 positive cells and has minimal cross reactivity with GD2 negative cells, while the non-specific isotope control antibody does not stain either cell type (Figure 13B).

[0236] Binding characteristics of AB527 were also assessed using tumor and normal tissue samples. AB527 and a nonspecific normal human IgM (hIgM) were biotinylated at Covance Research Products and used for IHC studies at Charles River Laboratories. Human melanoma cancer cells, M14, were implanted in nude rats to establish xenograft tumors, which were then used as GD2 positive control tissue for IHC method development. Preliminary studies using a direct avidin–biotin–peroxidase complex (ABC) procedure yielded positive staining of M14 tumor sections with biotinylated AB527 (Figure 14(b)) but not with biotinylated hIgM (Figure 14(a)). As expected, lymphocytes in cryosections of normal human spleen did not stain with either antibody. After titration of the antibodies, the staining concentrations for biotinylated AB527 and biotinylated hIgM selected for the GLP human tissue cross-reactivity study were set at 3 and 15 $\mu\text{g/mL}$. Using this method, a study was conducted to determine AB527 binding in 35 normal human tissues (3 samples/donors per tissue). As expected, high affinity membranous staining (positive cell membrane staining at the lower concentration of AB527 - 3 $\mu\text{g/mL}$) could be detected in M14 tumor sections. As shown in table 2, high affinity membranous staining was only found in the epithelial cells of esophagus (2 of 3 samples), fallopian tube (1 of 3 samples), and ureter (1 of 3 samples), as well as in decidual cells of placenta (3 of 3 samples), and reticular cells of thymus (2 of 3 samples). All the other tissues showed no high affinity membranous staining (table 2).

Table 2: AB527 Immunohistochemistry in Normal Human Tissues

Tissue	Positive/Total (Cell Type)	Tissue	Positive/Total (Cell Type)	Tissue	Positive/Total (Cell Type)
Adrenal	Negative	Liver	Negative	Skin	Negative
Blood Cells	Negative	Lung	Negative	Spinal Cord	Negative
Blood Vessels	Negative	Lymph Node	Negative	Spleen	Negative
Bone Marrow	Negative	Ovary	Negative	Striated Muscle	Negative
Brain (Cortex)	Negative	Fallopian Tube	1/3 (Epithelium)	Testis	Negative
Brain (Cerebellum)	Negative	Pancreas	Negative	Thymus	2/3 (Reticular Cells)
Breast	Negative	Parathyroid	Negative	Thyroid	Negative
Eye	Negative	Peripheral Nerve	Negative	Tonsil	Negative
Esophagus	2/3 Epithelium	Pituitary	Negative	Ureter	1/3 (Epithelium)
GI Tract	Negative	Placenta	3/3 (Decidual Cells)	Urinary Bladder	Negative
Heart	Negative	Prostate	Negative	Uterus (Body)	Negative
Kidney	Negative	Salivary Gland	Negative	Uterus (Cervix)	Negative

EXAMPLE 6

AB527 Facilitates Complement Dependent Cytotoxicity of Cells Expressing GD2

[0237] To determine whether AB527 mediates CDC, the melanoma cell lines described above for Figure 11 were incubated with partially purified antibody from the AB527-HYB-3B2-3C9 clone or AB527. Target cells were cultured in RPMI1640 supplemented with 10% FBS, 2 mM L-Glutamine, non-essential amino acids, and 6 μ M HEPES and harvested with trypsin before use. Target cells were incubated with 10 μ g/mL of either AB527 or AB527-HYB-3B2-3C9 clone IgM in the presence of 20% human serum. After incubation for 1 hour at 37 °C, live cells were identified with Cell Titer Glo® reagent (Promega Corp., Madison, WI). Percent killing was determined as the ratio of signal from treated versus untreated cells. The data in Figure 15 show that both antibodies can elicit a potent CDC reaction on certain melanoma cell lines, with AB527 consistently causing a greater degree of CDC. The results of the experiments using AB527 described in this Example are summarized in table 3.

Table 3: Comparison of FACS, Immunofluorescence Microscopy and CDC analysis of human melanoma cell lines treated with AB527.

Cell Line	AB527 FACS Staining	AB527 IHC	AB527-mediated CDC Killing
M14	+	+	+
LF0023	+	+	+
M101	+	+	+
M18	+	+	+
M10VAC	+	+	+
M21	+/-	-	-
M238	-	-	-
PM0496	+	+	-
IMCD0023	+/-	-	-
MG1055	-	-	-

[0238] Additional work was done to show that AB527 facilitates CDC only of cells that express GD2 in the presence of complement. As shown in Figure 16, AB527-mediated CDC activity only occurred to appreciable levels in the presence of complement and GD2-expressing cells, such as M14 cells, as opposed to 1205LU cells, which are human tumor cells that do not express GD2.

[0239] To assess the *in vitro* ability of IgG1 isotype-switched AB527 to induce CDC, the HindIII/BamHI variable region DNA fragment of AB527 was cloned in-frame into a HindIII/BamHI digested pEE6.4 vector containing the IgG1 constant region. Co-expression with the AB527 light chain caused the production of an IgG1 version of AB527. The ability of this antibody to mediate CDC of M14 cells was compared to the IgM version of AB527 at 1-100 $\mu\text{g/mL}$, as shown in Figure 17.

[0240] Studies were also conducted to assess the role that J-chain might play in AB527-mediated CDC. To produce AB527 IgM molecules having J-chain, plasmid p0362, which encodes J-chain, was transfected into CHOK1-SV cells and a stable, J-chain-expressing clonal line was generated. The AB527 expression plasmid was then introduced into this line to generate clones expressing AB527 and J-chain. Three clones named 5.F2-JC, 5.F8-JC, and 6.C3-JC were isolated. The antibodies were purified in parallel to compare their relative CDC activity against a non-J-chain containing version of AB527. Dose response curves are shown in

Figure 18. The calculated 90% of effective dose (ED90) for AB527, 5.F2-JC, 5.F8-JC, and 6.C3-JC was 1.21, 4.35, 2.52, and 6.81 $\mu\text{g/mL}$, respectively. IgM containing J-chain (5.F2-JC, 5.F8-C, and 6.C3-JC) appeared to be less effective at mediating CDC compared to AB527 without J-chain.

EXAMPLE 7

Capture of IgM from Conditioned Culture Supernatant by Protein A Affinity Chromatography

[0241] A column was packed with CPG3000A-protein A resin (Millipore). The volume of resin used to capture IgM was estimated, with the understanding that 1 mL of resin binds approximately 8 mg of IgM at a flow rate of 76 cm/h (table 4). Once the column was loaded with the desired amount of resin, it was connected to an FPLC apparatus and washed with 5 column volumes of purified water; 3 column volumes of 20 mM HCl, pH 1.5 (buffer A2); and 3 column volumes of 6 M guanidine·HCl (buffer A3). Before loading sample, the column was equilibrated with 5 column volumes of a 10 mM sodium phosphate buffer containing 200 mM NaCl and 0.01% Tween®-80, pH 7.5 (buffer A1).

[0242] Conditioned culture supernatant (CCS) was clarified by filtration through a .22 μm membrane filter. Detergent was added to the filtered CCS resulting in a solution containing 1% Triton® X-100 and 0.1% TNBP and incubated at 4°C for more than 2 hours. Detergent-treated CCS was applied to an equilibrated CPG3000A protein A column and processed at a flow rate of 76 cm/h. After washing away unbound proteins with 10 column volumes of buffer A1, IgM was eluted from the column using a 5 mM sodium phosphate buffer supplemented with 3 M MgCl_2 , pH 6.8 (buffer B). Remaining bound proteins were removed from the column by applying 3 column volumes of buffer A2 and 3 column volumes of buffer A3. To assess purity, a reducing SDS-PAGE gel comparing detergent-treated CCS applied to the protein A column (Load) and AB527 collected from pooled fractions after elution (Elute) was performed (Figure 19 - arrows point to the IgM μ chain (~70kD) and light chain (~25kD)). Protein A affinity chromatography allowed for recovery of $\geq 90\%$ of input IgM, which is substantially free of contaminants.

Table 4 - Dynamic binding capacity of IgM by CPG3000A protein A resin. Varying volumes of CCS containing AB527 were run through a 10X25 mm CPG3000A column at different linear flow rates. The concentration of AB527 loaded into the column and present in the flowthrough were measured by determining their relative concentrations as compared to that of a standard

curve for known IgM standards determined on a protein A column (POROS A, Applied Biosystems), fitted on a HPLC system. The ratio of AB527 in each flowthrough to the amount of loaded antibody was computed for each injection to determine the dynamic binding capacity of the column at 5% and 10% IgM breakthrough.

Linear Flow Rate (cm·hr ⁻¹)	Dynamic Binding Capacity (DBC) ¹ (mg AB527/L CPG3000A)		AB527 in Flowthrough (FT) (mg)	
	5% Breakthrough	10% Breakthrough	5% Breakthrough	10% Breakthrough
23	5780	6420	11559	12840
38	6869	7630	13737	15260
76	4395	5437	8789	10873
152	3126	3910	6252	7819

¹DBC was determined on a 1.0 cm x 2.5 cm bed (2ml bed volume)

EXAMPLE 8

Separation of IgM from Conditioned Culture Supernatant Contaminants by Cation Exchange Chromatography

[0243] To remove contaminants, IgM suspended in buffer B was concentrated by diafiltration using a Prep/Scale-TFF 1 cartridge (Millipore) and a 10-fold volume of a buffer containing 10 mM sodium phosphate, 75 mM NaCl. Diafiltration was conducted at a flow rate of approximately 200 mL/minute, with inlet pressure kept at less than 20 PSI and permeate flow at approximately 60 mL/minute. A solution of 10% Triton® X-100 was added to the diafiltered sample to yield a 1% final concentration of Triton® X-100, and the sample was incubated at room temperature for 1 hour and filtered through a 0.2 µm membrane.

[0244] A cation exchange column was packed with MacroCap™ SP resin (GE Healthcare). The volume of resin used to capture IgM was estimated, with the understanding that 1 mL of resin binds approximately 9 mg of IgM at a flow rate of 76 cm/h (Figure 20). Once the column was loaded with the desired amount of resin, it was connected to an FPLC apparatus and washed with 2 column volumes of purified water; 3 column volumes of 0.5 M NaOH (buffer A6), and 3 column volumes of 2 M NaCl. Before loading sample, the column was equilibrated with 5 column volumes of a buffer containing 10 mM sodium phosphate, 75 mM NaCl, 0.01% Tween®-80, pH 6.8 (buffer A4).

[0245] The diafiltered sample was applied to the column and processed at a flow rate of 76 cm/h. After washing away unbound proteins with 5 column volumes of buffer A4, IgM was eluted from the column using 4 column volumes of a 10 mM sodium phosphate buffer containing 200 mM NaCl and 0.01% Tween®-80, pH 6.8 (buffer C). Remaining bound proteins were removed from the column by applying 3 column volumes of buffer A6 and 3 column

volumes of 2 M NaCl As shown in Figs. 13A and B, the flowthrough material largely contained a mixture of IgM trimers and dimers (Figure 21A); however, the eluted material was almost exclusively pentameric IgM (Figure 21B). Cation exchange chromatography provides $\geq 95\%$ yield of pentameric IgM that is substantially free of contaminants and incompletely assembled IgM (Figure 22).

EXAMPLE 9

Separation of IgM from Conditioned Culture Supernatant Contaminants by Hydroxyapatite Chromatography

[0246] The hydroxyapatite column was prepared using CHT® II Ceramic Hydroxyapatite, 80 μm bead size (Bio-Rad Laboratories, #157-8100). The volume of resin used to capture IgM was estimated, with the understanding that 1 mL of resin binds approximately 20 mg of IgM at a flow rate of 76 cm/h, and at this flow rate there is less than 5% loss of IgM in the column flow-through. The column was packed by hydrating the matrix with 200 mM potassium phosphate pH 9.0, using 0.54g of dry matrix for each mL of desired column-bed volume. The hydrated matrix was added to the column, allowing it to settle before the resin was equilibrated with 5 column volumes of a buffer containing 10 mM sodium phosphate, 100 mM NaCl, 0.01% Tween®-80, pH 6.8 (buffer A8)

[0247] To reduce contaminants, IgM suspended in 10 mM sodium phosphate, 200 mM NaCl, 0.01% Tween®-80, pH 6.8 was diluted with an equivalent volume of 0.01% Tween®-80 and added to the equilibrated column. The IgM suspension was processed through the column at a flow rate of 76 cm/h. The column was washed with 10 column volumes of buffer A8, to remove unbound substances. Bound IgM was eluted with 4 column volumes of a 175 mM sodium phosphate buffer containing 100 mM NaCl and 0.01% Tween®-80, pH 6.8 (data not shown). Following elution, remaining bound proteins were removed from the column with a 500 mM sodium phosphate buffer containing 100 mM NaCl, pH 6.8. The resulting IgM solution was substantially free of contaminants. The solution eluted from the hydroxyapatite matrix, when preceded by protein A affinity chromatography and cation exchange chromatography, described herein, was over 99% pure as compared to the clarified cell culture supernatant.

EXAMPLE 10

Production of a Ganglioside-BSA Conjugates

[0248] A GD2-BSA conjugate was produced to provide an effective GD2 antigen to examine the specificity of GD2 specific antibodies by immunoassay. In addition, GM2-BSA and

GM3-BSA conjugates were produced to serve as controls for GD2-BSA binding specificity. The covalent conjugates were prepared by site-specific reductive amination of the gangliosides by amines present on BSA, utilizing sodium cyanoborohydride as a mild reducing agent. Coupling is thought to occur mainly to terminal ϵ -amino groups on the side chains of lysine residues present in BSA.

[0249] To prepare the conjugates, 400 μ l of a 2mg/ml solution of BSA dissolved in 0.05 M carbonate-bicarbonate buffer, pH 9.6, was combined with 500 μ l of a 1 mg/ml solution of GD2 (Bioscience International, #A86168H), GM2 (Axxora, #ALX-302-005-M001), or GM3 (Axxora, #ALX-302-003-M002) dissolved in absolute ethanol; 2070 μ l of 0.05 M carbonate-bicarbonate buffer, pH 9.6; and 30 μ l of a 5 M sodium cyanoborohydride solution of 10 mM sodium hydroxide. The reaction mixture was incubated at 25°C for two hours before adding 1200 μ l of 1M ethanolamine, pH 8.0 and incubating at 25°C for an additional 15 minutes. The reaction solution was placed in dialysis tubing (Pierce Biotechnology, # 68100) and dialyzed twice in 1L of a 50 mM sodium carbonate-bicarbonate solution, pH 9.6, at 2-8°C for 6-18 hours.

EXAMPLE 11

AB527 Binds to BSA-Conjugated GD2

[0250] Antigenicity of the GD2-BSA conjugate was assessed by ELISA. ELISA coating buffer was prepared by dissolving one carbonate/bicarbonate capsule (Sigma) in 100mL ultrapure water. GD2-BSA conjugate was diluted to a concentration of 2 μ g/ml, in ELISA coating buffer, and added to wells of a 96-well ELISA plate (Greiner Bio-One cat # 655081). The plates were sealed and incubated at 2-8°C for 16-24 hours. Following incubation, the coating buffer was removed from the wells and 0.3 mL/well of ELISA blocking solution (PBS supplemented with 2.5% BSA (w/v) and 0.05% Tween®-20) was added. Plates were incubated, while shaking on a microplate shaker, for 2 hours at 21-25 °C before adding AB527 antibody. Two-fold serial dilutions of AB527 (initial working concentration of 2 μ g/ml) were added to the blocked ELISA plates and incubated with shaking on a microplate shaker at 21-25°C for 1 hour. Plates were washed three times on the Dynex Ultrawash™ plate washer, using 0.3mL/well ELISA wash buffer. Bound AB527 was detected with HRP-conjugated goat anti-human IgG + IgM (H+L), diluted 1:10,000 in ELISA blocking buffer. The HRP-conjugate was removed and wells were washed three times with 0.3mL/well ELISA wash buffer. SureBlue™ TMB substrate (50 μ l/well) was added to washed wells and plates were incubated at 21-25°C for 15 minutes, before AB527 binding was assessed by light absorption (450 nm) (Figure 23). Nonspecific absorption at 450 nm was determined to be 0.065 in the absence of AB527.

[0251] In contrast to the GD2-BSA conjugate, AB527 did not recognize other ganglioside-BSA conjugates. The gangliosides GM2 and GM3 were conjugated to BSA, by the method described for GD2, and both were assessed, by ELISA, for the ability to elicit AB527 binding. As shown in Figure 24, AB527 recognized only the GD2-BSA conjugate. It should be noted that a GM2-specific antibody was able to detect the GM2-BSA conjugate; however, an appropriate GM3-specific antibody was not available to assess the antigenic nature of the GM3-BSA conjugate (data not shown).

EXAMPLE 12

Effect of CMS on AB527-Expressing Cells Grown in GIBCO-CD CHO Complete Medium

[0252] To determine the effect of culture media supplement (CMS) on cell growth and antibody production, cells were revived and cultured in a shake flask in 20 mL GIBCO-CD-CHO complete medium. All cultures were incubated in 125 mL shake flask at 37 °C, 5% CO₂, 120 rpm. The seeding density was 3.5×10^5 cells/mL at > 95% viability. Cells were cultured for 14 days in either CD-CHO media or in CD-CHO media supplemented on days 3-7 with 2% (vol/vol) CMS. Following addition of CMS on day 7 a total of 10% of the initial culture volume had been supplemented with CMS. Viable cell density (Figure 25), percent viability (Figure 26), and integral viable cell density (IVC) (Figure 27) were determined using a CEDEX automated cell counter on days 0, 3, 5, 7, 10, 13, and 14. Antibody production was determined for each group of cultured cells on day 14 by protein A HPLC (Figure 28).

[0253] As shown in Figure 25, cultured cells provided with 2% CMS from day 3 to day 7 achieved a maximum viable cell density of 12.8×10^6 cells/mL on day 7. The control cell culture, without CMS, reached a maximum cell density of 10.7×10^6 cells/mL on day 7. The use of CMS replenishes limiting medium components such as glucose, amino acids and vitamins and helps the culture reach a higher cell density. In both cell culture conditions, cells maintained a good viability (> 90%) until day 7 (Figure 26); however, cultured cells provided CMS from day 3 to day 7 demonstrated a better viability than the control culture after day 7. It is possible that some key nutrients were depleted in the control culture after day 7 while the CMS added back those key nutrients and helped the cell maintain a better viability.

[0254] On day 14, total antibody production was assessed for each culture. Cells cultured with CMS reached a maximum antibody titer of 7240 mg/L, much higher than that observed for the control culture (3036 mg/L) (Figure 28). The higher antibody titer could be a result of a higher specific antibody production rate after the addition of the nutrients.

Investigation of the cell growth and antibody production profiles for each growth condition demonstrated that the culture provided CMS achieved a 140% increase in antibody production (table 5). Even though the doubling time was the same for both cultures, the CMS did help cultures achieve a 50% improvement in IVC and another 50% increase in specific antibody production rate. The 50% improvement in IVC was due to a higher maximum cell density achieved in the culture with CMS. Since the final antibody titer is equal to the IVC multiple specific production rate, it suggested that both IVC and specific antibody production rate contributed to the 140% increase in antibody production in the cell culture with CMS. Table 5 summarizes the effects the CMS on AB527-expressing cells.

Table 5 - Summary of the Effects of CMS on Cell Growth and Antibody Production of AB527-Expressing Cells.

Culture Conditions	Maximum Viable Cell Density (10^5 /mL)	Final Antibody Titer (mg/L)	Doubling Time (DT)	Integral of Viable Cell Density (IVC) (cells·day/mL)	Specific Antibody Production Rate (Qp) (mg/ 10^9 cells/day)
CD-CHO Control	107.00	3036	24.17	6.85E+7	44.08
CMS	128.40	7239.9	24.17	1.04E+8	66.95

EXAMPLE 13

Effect of Valeric Acid on AB527-Expressing Cells Grown in GIBCO-CD CHO Complete Medium

[0255] To determine the effect of valeric acid on cell growth, AB527-expressing cells were cultured in GIBCO-CD-CHO complete medium supplemented with valeric acid to a concentration of 0 mM, 1 mM, 2 mM, 4 mM, or 8 mM. Cells were cultured for 14 days and viable cell density, percent viability, and integral viable cell density were determined using a CEDEX automated cell counter on days 0, 3, 5, 7, 10, 13, and 14. Antibody production was determined for each group of cultured cells on day 14 by protein A HPLC.

[0256] Cells cultured without valeric acid achieved the maximum cell density of 10.8×10^6 cells/mL at day 7; however, cells cultured with valeric acid grew much slower as the concentration of valeric acid increased. For example, cells cultured in 8 mM valeric acid, reached a maximum cell density of 5.4×10^6 cells/mL at day 7, which is the lowest maximum cell density of all cultures. Cell viability studies demonstrated that cells cultured with valeric acid had lower viability before day 7, with the lowest viability at day 7 (67.4%) observed for cells cultured in 8 mM valeric acid. These results indicate that higher concentration of the

valeric acid inhibited cell proliferation. After day 7, the viability of the control culture dropped much quicker; however, this was likely due to consumption of key nutrients since this cell culture had a much higher cell density than those cultured in valeric acid (table 6).

[0257] Unexpectedly, however, valeric acid had a positive effect on antibody production. All cells cultured in the presence of valeric acid, with exception of the highest concentration of 8 mM, had higher antibody titers than the control culture at day 14. Cell cultured in 1 mM valeric acid produced the highest antibody titer (4137 mg/L) at day 14, which is 36% higher than that of the control culture (3036 mg/L) (table 6).

[0258] Even though the valeric acid inhibited the cell growth, it did not demonstrate the significant negative effect on the integral of viable cell density, except at the highest concentration of 8 mM. The IVC values at day 14 were quite similar for the other four culture conditions. It is likely that the inhibition of valeric acid on the cell proliferation led to a slower consumption of the nutrients, thus avoiding late-stage nutrient-depleted media, which was encountered by the control culture.

[0259] In summary, the cell culture can achieve a better antibody production in presence of the valeric acid. The specific antibody production rate ranging from 52 to 58 mg/10⁹ cells/day was observed in cultures with valeric acid (table 6), which are much higher than the value of 44 mg/10⁹ cells/day observed for the control culture. Thus, the highest antibody production achieved by cells cultured in 1 mM valeric acid was mainly due to the increases in the specific antibody production (table 6).

Table 6 - Summary of the Effects of Valeric Acid on Cell Growth and Antibody Production of AB527-Expressing Cells.

Culture Conditions	Maximum Viable Cell Density (10 ⁵ /mL)	Final Antibody Titer (mg/L)	Doubling Time (DT)	Integral of Viable Cell Density (IVC) (cells·day/mL)	Specific Antibody Production Rate (Qp) (mg/10 ⁹ cells/day)
CD-CHO Control	107.00	3036	24.17	6.85E+7	44.08
Valeric Acid (1 mM)	94.60	4137.1	24.17	6.74 E+7	58.44
Valeric Acid (2 mM)	81.00	3706.7	24.17	6.55 E+7	54.45
Valeric Acid (4 mM)	77.40	3600.1	24.17	6.72 E+7	52.24
Valeric Acid (8 mM)	54.40	2760.4	24.17	4.76 E+7	56.16

What is Claimed:

1. An isolated antibody, or an antigen-binding fragment thereof, that specifically binds to human GD2, wherein the heavy chain CDR3 comprises the amino acid sequence of SEQ ID NO: 12.
2. The antibody or antigen-binding fragment thereof of claim 1, wherein the heavy chain CDR1 comprises the amino acid sequence of SEQ ID NO: 10, the heavy chain CDR2 comprises the amino acid sequence of SEQ ID NO: 11, the light chain CDR1 comprises the amino acid sequence of SEQ ID NO: 26, the light chain CDR2 comprises the amino acid sequence of SEQ ID NO: 27, and the light chain CDR3 comprises the amino acid sequence of SEQ ID NO: 28.
3. The antibody or antigen-binding fragment thereof of claim 1, wherein the antibody is a human antibody.
4. The antibody or antigen-binding fragment thereof of claim 1, wherein the heavy chain variable domain comprises the amino acid sequence of SEQ ID NO: 16.
5. The antibody or antigen-binding fragment thereof of claim 1 wherein the light chain variable domain comprises the amino acid sequence of SEQ ID NO: 32.
6. The antibody or antigen-binding fragment thereof of claim 1, wherein the heavy chain comprises the amino acid sequence of SEQ ID NO: 40.
7. The antibody or antigen-binding fragment thereof of claim 1, wherein the light chain comprises the amino acid sequence of SEQ ID NO: 42.
8. The antibody or antigen-binding fragment thereof of claim 1, where the antibody or antigen-binding fragment thereof is of the IgM isotype.
9. The antibody or antigen-binding fragment thereof of claim 8, wherein the antibody, or antigen-binding fragment thereof is polymeric.
10. The antibody or antigen-binding fragment thereof of claim 9, wherein the polymeric state of the antibody or antigen-binding fragment thereof is pentameric.

11. The antibody or antigen-binding fragment thereof of claim 10, wherein the pentameric antibody or antigen-binding fragment thereof does not incorporate a J-chain.
12. An antibody or antigen-binding fragment thereof of claim 1 which mediates complement dependent cytotoxicity.
13. A polynucleotide encoding the antibody, or antigen binding fragment, of claim 1.
14. The polynucleotide of claim 13, comprising SEQ ID NO: 4.
15. The polynucleotide of claim 14, comprising the nucleic acid sequences of SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 18, SEQ ID NO: 19, and SEQ ID NO: 20
16. The polynucleotide of claim 14, comprising the nucleic acid sequence of SEQ ID NO: 8.
17. The polynucleotide of claim 14, comprising the nucleic acid sequence of SEQ ID NO: 24.
18. The polynucleotide of claim 14, comprising the nucleic acid sequence of SEQ ID NO: 39.
19. The polynucleotide of claim 14, comprising the nucleic acid sequence of SEQ ID NO: 41.
20. A composition comprising the antibody, or antigen-binding fragment, of claim 1 and a pharmaceutically acceptable carrier.
21. A vector comprising the polynucleotide sequence of claim 14.
22. A cell comprising the vector of claim 21.
23. The cell of claim 22, wherein the cell is a bacterial cell.
24. The cell of claim 22, wherein the cell is a eukaryotic cell.
25. The eukaryotic cell of claim 24, wherein the eukaryotic cell is a Chinese Hamster Ovary (CHO) cell.
26. A cell that expresses the antibody or antigen-binding fragment thereof of claim 1.

27. A cell that expresses the antibody or antigen-binding fragment thereof of claim 11.
28. The cell of claim 26, wherein the cell is a hybridoma.
29. The cell of claim 27, wherein the cell is a transfectoma.
30. A method for treating or preventing a GD2-associated disease in a subject, comprising administering to the subject an antibody or an antigen-binding fragment thereof that specifically binds to GD2 in an amount effective to treat or prevent the disease.
31. The method of claim 30, wherein the GD2-associated disease is cancer.
32. The method of claim 31, wherein the GD2-associated disease is melanoma.
33. The method of claim 30, wherein the subject is a mammal.
34. The method of claim 30, wherein the mammal is a human.
35. A method of making an antibody or antigen-binding of fragment, comprising culturing a host cell of claim 26 under conditions suitable to produce the antibody or antigen-binding fragment, and recovering the antibody or antigen-binding fragment thereof from the cell culture.
36. A method of making an antibody or antigen-binding of fragment, comprising culturing a host cell of claim 27 under conditions suitable to produce the antibody or antigen-binding fragment, and recovering the antibody or antigen-binding fragment thereof from the cell culture.
37. An antibody, or antigen-binding fragment thereof, produced by an AB527-HYB-3B2-3C9 cell, wherein such antibody, or antigen-binding fragment thereof, does not incorporate J-chain.
38. A method of detecting a GD2-expressing cell in a subject, comprising administering to the subject the antibody, or antigen-binding fragment, of claim 1 and detecting cells expressing GD2.

39. The method of claim 38, wherein, the antibody, or antigen-binding fragment thereof, is detectably labeled.

40. The method of claim 39, wherein the detectable label is iodine-131.

41. The antibody of claim 2, or antigen-binding fragment thereof, wherein the antibody, or antigen-binding fragment, exhibits a steady state dissociation constant (K_D) for GD2 of 4.5×10^{-9} molar.

42. An antibody, or antigen-binding fragment thereof, comprising the heavy chain and light chain CDRs of the antibody produced by a cell having ATCC accession No. PTA-9376.

43. An antibody, or antigen-binding fragment thereof, comprising the heavy chain and light chain variable domains of the antibody produced by a cell having ATCC accession No. PTA-9376.

1/28




Fig. 1A




Fig. 1B

2/28




Fig. 2

3/28




Fig. 3A




Fig. 3B

4/28




Fig. 4

5/28




Fig. 5

6/28




Fig. 6

7/28




Fig. 7

8/28




Fig. 8

9/28




Fig. 9

10/28




Fig. 10

11/28




Fig. 11

12/28




Fig. 12

13/28




Fig. 13

14/28




Fig. 14

15/28




Fig. 15

16/28




Fig. 16

17/28




Fig. 17

18/28




Fig. 18




Fig. 19

20/28




Fig. 20

21/28




Fig. 21

22/28




Fig. 22

23/28




Fig. 23

24/28




Fig. 24

25/28




Fig. 25

26/28




Fig. 26

27/28




Fig. 27

28/28




Fig. 28

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US2009/049139

Box No. I Nucleotide and/or amino acid sequence(s) (Continuation of item 1.b of the first sheet)

1. With regard to any nucleotide and/or amino acid sequence disclosed in the international application and necessary to the claimed invention, the international search was carried out on the basis of:

a. type of material

☒

a sequence listing

☐

table(s) related to the sequence listing

b. format of material

☒

on paper

☒

in electronic form

c. time of filing/furnishing

☒

contained in the international application as filed

☒

filed together with the international application in electronic form

☐

furnished subsequently to this Authority for the purpose of search

2. ☐ In addition, in the case that more than one version or copy of a sequence listing and/or table relating thereto has been filed or furnished, the required statements that the information in the subsequent or additional copies is identical to that in the application as filed or does not go beyond the application as filed, as appropriate, were furnished.

3. Additional comments:

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2009/049139

A. CLASSIFICATION OF SUBJECT MATTER

INV. C07K16/30 A61K39/395 A61P35/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, BIOSIS, EMBASE, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	IRIE R ET AL: "REGRESSION OF CUTANEOUS METASTATIC MELANOMA BY INTRALESIONAL INJECTION WITH HUMAN MONOCLONAL ANTIBODY TO GANGLIOSIDE GD2" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA, NATIONAL ACADEMY OF SCIENCE, WASHINGTON, DC, US, vol. 83, no. 22, 1 November 1986 (1986-11-01), pages 8694-8698, XP002017115 ISSN: 0027-8424	30-34
Y	abstract ----- -/--	1-29, 35-43

☒ Further documents are listed in the continuation of Box C.

☒ See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

16 October 2009

Date of mailing of the international search report

30/10/2009

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Loubradou, Gabriel

INTERNATIONAL SEARCH REPORT

International application No

PCT/US2009/049139

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2004/055056 A (MERCK PATENT GMBH [DE]; GILLIES STEPHEN D [US]; LO KIN-MING [US]) 1 July 2004 (2004-07-01)	30-34
Y	abstract; claims; examples; sequences 1,2	1-29, 35-43
Y	----- KATANO M ET AL: "Human monoclonal antibody to a neuroectodermal tumor antigen (OFA-I-2)." ANNALS OF THE NEW YORK ACADEMY OF SCIENCES 1983, vol. 417, 1983, pages 427-434, XP002550651 ISSN: 0077-8923 abstract	1-43
Y	----- MUJOO K ET AL: "DISIALOGANGLIOSIDE GD2 ON HUMAN NEUROBLASTOMA CELLS: TARGET ANTIGEN FOR MONOCLONAL ANTIBODY-MEDIATED CYTOLYSIS AND SUPPRESSION OF TUMOR GROWTH" CANCER RESEARCH, AMERICAN ASSOCIATION FOR CANCER RESEARCH, BALTIMORE, MD., US, vol. 47, no. 4, 15 February 1987 (1987-02-15), pages 1098-1104, XP009038918 ISSN: 0008-5472 abstract	1-43
Y	----- WO 97/34634 A (SLOAN KETTERING INST CANCER [US]; CHEUNG NAI KONG V [US]; LARSON STEVE) 25 September 1997 (1997-09-25) page 2, line 1 - line 6; table 1 page 11 - page 13	1-43
Y	----- WO 2008/043777 A (UNIV NANTES [FR]; BIRKLE STEPHANE [FR]; MUSSINI JEAN-MARIE [FR]; AUBRY) 17 April 2008 (2008-04-17) claims; examples	1-43
Y	----- CERATO E ET AL: "VARIABLE REGION GENE SEGMENTS OF NINE MONOCLONAL ANTIBODIES SPECIFIC TO DISIALOGANGLIOSIDES (GD2, GD3) AND THEIR O-ACETYLATED DERIVATIVES" HYBRIDOMA, LIEBERT, NEW YORK, NY, US, vol. 16, no. 4, 1 August 1997 (1997-08-01), pages 307-316, XP008078446 ISSN: 0272-457X abstract; table 3 ----- -/--	1-43

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2009/049139

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	ALVAREZ-RUEDA NIDIA ET AL: "Binding activities and antitumor properties of a new mouse/human chimeric antibody specific for GD2 ganglioside antigen." CLINICAL CANCER RESEARCH : AN OFFICIAL JOURNAL OF THE AMERICAN ASSOCIATION FOR CANCER RESEARCH 15 SEP 2007, vol. 13, no. 18 Pt 2, 15 September 2007 (2007-09-15), pages 5613s-5620s, XP002550652 ISSN: 1078-0432 abstract	1-43
Y	HOOGENBOOM H R: "Selecting and screening recombinant antibody libraries" NATURE BIOTECHNOLOGY, NATURE PUBLISHING GROUP, NEW YORK, NY, US, vol. 23, no. 9, 1 September 2005 (2005-09-01), pages 1105-1116, XP002348401 ISSN: 1087-0156 page 1111, left-hand column, paragraph 2 - page 1112, left-hand column, paragraph 1; table 2	41,43
Y	EP 1 652 925 A (CHUGAI PHARMACEUTICAL CO LTD [JP]; IRIE REIKO [US] CHUGAI PHARMACEUTIC) 3 May 2006 (2006-05-03) the whole document	8-12,29
A	WO 94/19457 A (UNIV CALIFORNIA [US]) 1 September 1994 (1994-09-01) the whole document	
A	IMAI MASAKI ET AL: "Complement-mediated mechanisms in anti-GD2 monoclonal antibody therapy of murine metastatic cancer." CANCER RESEARCH, vol. 65, no. 22, November 2005 (2005-11), pages 10562-10568, XP002550653 ISSN: 0008-5472 abstract	

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2009/049139

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 2004055056	A	01-07-2004	AU 2003298187 A1 BR 0317376 A CA 2510180 A1 CN 1726227 A EP 1572748 A1 JP 2006521085 T KR 20050085775 A MX PA05006384 A ZA 200505681 A	09-07-2004 16-11-2005 01-07-2004 25-01-2006 14-09-2005 21-09-2006 29-08-2005 29-08-2005 26-04-2006
WO 9734634	A	25-09-1997	NONE	
WO 2008043777	A	17-04-2008	AU 2007306384 A1 EP 2076542 A1 FR 2906808 A1	17-04-2008 08-07-2009 11-04-2008
EP 1652925	A	03-05-2006	AU 2004256354 A1 CA 2532966 A1 WO 2005005636 A1	20-01-2005 20-01-2005 20-01-2005
WO 9419457	A	01-09-1994	AT 255633 T AU 687791 B2 AU 6238094 A CA 2156857 A1 DE 69433378 D1 DE 69433378 T2 DK 687295 T3 EP 0687295 A1 ES 2211877 T3 JP 8507209 T PT 687295 E	15-12-2003 05-03-1998 14-09-1994 01-09-1994 15-01-2004 14-10-2004 05-04-2004 20-12-1995 16-07-2004 06-08-1996 30-04-2004