

(19)



(11)

EP 1 305 285 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the grant of the patent:
16.05.2007 Bulletin 2007/20

(51) Int Cl.:
C07D 209/08 ^(2006.01) **A61K 31/404** ^(2006.01)
A61K 31/405 ^(2006.01) **C07D 209/12** ^(2006.01)
A61P 3/10 ^(2006.01)

(21) Application number: **01954836.1**

(86) International application number:
PCT/US2001/022979

(22) Date of filing: **20.07.2001**

(87) International publication number:
WO 2002/008188 (31.01.2002 Gazette 2002/05)

(54) N-SUBSTITUTED INDOLES USEFUL IN THE TREATMENT OF DIABETES

N-SUBSTITUIERTE INDOLE MIT ANWENDUNG IN DER BEHANDLUNG VON DIABETES

INDOLES N-SUBSTITUES UTILES POUR LE TRAITEMENT DU DIABETE

(84) Designated Contracting States:
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE TR**
Designated Extension States:
AL LT LV MK RO SI

- **JONES, Anthony Brian**
Rahway, NJ 07065-0907 (US)
- **WOOD, Harold Blair**
Rahway, NJ 07065-0907 (US)

(30) Priority: **25.07.2000 US 220778 P**

(43) Date of publication of application:
02.05.2003 Bulletin 2003/18

(73) Proprietor: **Merck & Co., Inc.**
Rahway, New Jersey 07065-0907 (US)

(74) Representative: **Horgan, James Michael Frederic
et al**
Merck & Co., Inc.
European Patent Department
Merck Sharp & Dohme Limited,
Hertford Road,
Hoddesdon
Hertfordshire, EN11 9BU (GB)

(72) Inventors:

- **ACTON, John, J. III**
Rahway, NJ 07065-0907 (US)
- **BLACK, Regina Marie**
Rahway, NJ 07065-0907 (US)

(56) References cited:
WO-A-94/14434 **WO-A-99/43654**
WO-A2-00/35886 **US-A- 5 728 709**

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 1 305 285 B1

Description

FIELD OF THE INVENTION

[0001] The instant invention is concerned with N-substituted indoles having aryloxyalkanoic acid substituents, and pharmaceutically acceptable salts and prodrugs thereof, which are useful as therapeutic compounds, particularly in the treatment of Type 2 diabetes mellitus, often referred to as non-insulin dependent diabetes (NIDDM), of conditions that are often associated with this disease, and of lipid disorders.

BACKGROUND OF THE INVENTION

[0002] Diabetes refers to a disease process derived from multiple causative factors and characterized by elevated levels of plasma glucose or hyperglycemia in the fasting state or after administration of glucose during an oral glucose tolerance test. Persistent or uncontrolled hyperglycemia is associated with increased and premature morbidity and mortality. Often abnormal glucose homeostasis is associated both directly and indirectly with alterations of the lipid, lipoprotein and apolipoprotein metabolism and other metabolic and hemodynamic disease. Therefore patients with Type 2 diabetes mellitus are at especially increased risk of macrovascular and microvascular complications, including coronary heart disease, stroke, peripheral vascular disease, hypertension, nephropathy, neuropathy, and retinopathy. Therefore, therapeutical control of glucose homeostasis, lipid metabolism and hypertension are critically important in the clinical management and treatment of diabetes mellitus.

[0003] There are two generally recognized forms of diabetes. In type 1 diabetes, or insulin-dependent diabetes mellitus (IDDM), patients produce little or no insulin, the hormone which regulates glucose utilization. In type 2 diabetes, or noninsulin dependent diabetes mellitus (NIDDM), patients often have plasma insulin levels that are the same or even elevated compared to nondiabetic subjects; however, these patients have developed a resistance to the insulin stimulating effect on glucose and lipid metabolism in the main insulin-sensitive tissues, which are muscle, liver and adipose tissues, and the plasma insulin levels, while elevated, are insufficient to overcome the pronounced insulin resistance.

[0004] Insulin resistance is not primarily due to a diminished number of insulin receptors but to a post-insulin receptor binding defect that is not yet understood. This resistance to insulin responsiveness results in insufficient insulin activation of glucose uptake, oxidation and storage in muscle and inadequate insulin repression of lipolysis in adipose tissue and of glucose production and secretion in the liver.

[0005] The available treatments for type 2 diabetes, which have not changed substantially in many years, have recognized limitations. While physical exercise and reductions in dietary intake of calories will dramatically improve the diabetic condition, compliance with this treatment is very poor because of well-entrenched sedentary lifestyles and excess food consumption, especially of foods containing high amounts of saturated fat. Increasing the plasma level of insulin by administration of sulfonylureas (e.g. tolbutamide and glipizide), which stimulate the pancreatic β -cells to secrete more insulin, and/or by injection of insulin after the response to sulfonylureas fails, will result in high enough insulin concentrations to stimulate the very insulin-resistant tissues. However, dangerously low levels of plasma glucose can result from these last two treatments, and increasing insulin resistance due to the even higher plasma insulin levels can occur. The biguanides increase insulin sensitivity resulting in some correction of hyperglycemia. However, the two biguanides, phenformin and metformin, can induce lactic acidosis and nausea/diarrhea, respectively.

[0006] The glitazones (i.e. 5-benzylthiazolidine-2,4-diones) are a more recently described class of compounds with potential for a novel mode of action in preventing or ameliorating many symptoms of type 2 diabetes. These agents substantially increase insulin sensitivity in muscle, liver and adipose tissue in several animal models of type 2 diabetes resulting in partial or complete correction of the elevated plasma levels of glucose without occurrence of hypoglycemia.

[0007] Disorders of lipid metabolism or dyslipidemias include various conditions characterized by abnormal concentrations of one or more lipids (i.e. cholesterol and triglycerides), and/or apolipoproteins (i.e., apolipoproteins A, B, C and E), and/or lipoproteins (i.e., the macromolecular complexes formed by the lipid and the apolipoprotein that allow lipids to circulate in blood, such as LDL, VLDL and IDL). Cholesterol is mostly carried in Low Density Lipoproteins (LDL), and this component is commonly known as the "bad" cholesterol because it has been shown that elevations in LDL-cholesterol correlate closely to the risk of coronary heart disease. A smaller component of cholesterol is carried in the High Density Lipoproteins and is commonly known as the "good" cholesterol. In fact, it is known that the primary function of HDL is to accept cholesterol deposited in the arterial wall and to transport it back to the liver for disposal through the intestine. Although it is desirable to lower elevated levels of LDL cholesterol, it is also desirable to increase levels of HDL cholesterol. Generally, it has been found that increased levels of HDL are associated with lower risk for coronary heart disease (CHD). See, for example, Gordon, et al., Am. J. Med., 62, 707-714 (1977); Stampfer, et al., N. England J. Med., 325, 373-381 (1991); and Kannel, et al., Ann. Internal Med., 90, 85-91 (1979). An example of an HDL raising agent is nicotinic acid, a drug with limited utility because doses that achieve HDL raising are associated with undesirable effects, such as flushing.

[0008] Dyslipidemias were originally classified by Fredrickson according to the combination of alterations mentioned above. The Fredrickson classification includes 6 phenotypes (i.e., I, IIa, IIb, III, IV and V) with the most common being the isolated hypercholesterolemia (or type IIa) which is usually accompanied by elevated concentrations of total and LDL cholesterol. The initial treatment for hypercholesterolemia is often to modify the diet to one low in fat and cholesterol, coupled with appropriate physical exercise, followed by drug therapy when LDL-lowering goals are not met by diet and exercise alone

[0009] A second common form of dyslipidemia is the mixed or combined hyperlipidemia or type IIb and III of the Fredrickson classification. This dyslipidemia is often prevalent in patients with type 2 diabetes, obesity and the metabolic syndrome. In this dyslipidemia there are modest elevations of LDL-cholesterol, accompanied by more pronounced elevations of small dense LDL-cholesterol particles, VLDL and/or IDL (i.e., triglyceride rich lipoproteins), and total triglycerides. In addition, concentrations of HDL are often low.

[0010] Peroxisome proliferators are a structurally diverse group of compounds that when administered to rodents elicit dramatic increases in the size and number of hepatic and renal peroxisomes, as well as concomitant increases in the capacity of peroxisomes to metabolize fatty acids via increased expression of the enzymes of the beta-oxidation cycle. Compounds of this group include but are not limited to the fibrate class of lipid modulating drugs, herbicides and phthalate plasticizers. Peroxisome proliferation is also triggered by dietary or physiological factors such as a high-fat diet and cold acclimatization.

[0011] Three sub-types of peroxisome proliferator activated receptor (PPAR) have been discovered and described; they are peroxisome proliferator activated receptor alpha (PPAR α), peroxisome proliferator activated receptor gamma (PPAR γ) and peroxisome proliferator activated receptor delta (PPAR δ). Identification of PPAR α , a member of the nuclear hormone receptor superfamily activated by peroxisome proliferators, has facilitated analysis of the mechanism by which peroxisome proliferators exert their pleiotropic effects. PPAR α is activated by a number of medium and long-chain fatty acids, and it is involved in stimulating β -oxidation of fatty acids. PPAR α is also associated with the activity of fibrates and fatty acids in rodents and humans. Fibric acid derivatives such as clofibrate, fenofibrate, benzaifibrate, ciprofibrate, beclofibrate and etofibrate, as well as gemfibrozil, each of which are PPAR α ligands and/or activators, produce a substantial reduction in plasma triglycerides as well as some increase in HDL. The effects on LDL cholesterol are inconsistent and might depend upon the compound and/or the dyslipidemic phenotype. For these reasons, this class of compounds has been primarily used to treat hypertriglyceridemia (i.e., Fredrickson Type IV and V) and/or mixed hyperlipidemia.

[0012] The PPAR γ receptor subtypes are involved in activating the program of adipocyte differentiation and are not involved in stimulating peroxisome proliferation in the liver. There are two known protein isoforms of PPAR γ : PPAR γ 1 and PPAR γ 2 which differ only in that PPAR γ 2 contains an additional 28 amino acids present at the amino terminus. The DNA sequences for the human isotypes are described in Elbrecht, et al., BBRC 224:431-437 (1996). In mice, PPAR γ 2 is expressed specifically in fat cells. Tontonoz *et al.*, *Cell* 79: 1147-1156 (1994) provide evidence to show that one physiological role of PPAR γ 2 is to induce adipocyte differentiation. As with other members of the nuclear hormone receptor superfamily, PPAR γ 2 regulates the expression of genes through interaction with other proteins and binding to hormone response elements, for example in the 5' flanking regions of responsive genes. An example of a PPAR γ 2 responsive gene is the tissue-specific adipocyte P2 gene. Although peroxisome proliferators, including the fibrates and fatty acids, activate the transcriptional activity of PPAR's, only prostaglandin J₂ derivatives have been identified as potential natural ligands of the PPAR γ subtype, which also binds thiazolidinedione antidiabetic agents with high affinity.

[0013] The human nuclear receptor gene PPAR δ (hPPAR δ) has been cloned from a human osteosarcoma cell cDNA library and is fully described in A. Schmidt et al., *Molecular Endocrinology*, 6:1634-1641 (1992). It should be noted that PPAR δ is also referred to in the literature as PPAR β and as NUC1, and each of these names refers to the same receptor; in Schmidt *et al.* the receptor is referred to as NUC1.

[0014] In WO96/01430, a human PPAR subtype, hNUC1B, is disclosed. The amino acid sequence of hNUC1B differs from human PPAR δ (referred to therein as hNUC1) by one amino acid, i.e., alanine at position 292. Based on in vivo experiments described therein, the authors suggest that hNUC1B protein represses hPPAR α and thyroid hormone receptor protein activity.

[0015] It has been disclosed in WO97/28149 that agonists of PPAR δ are useful in raising HDL plasma levels. WO97/27857, 97/28115, 97/28137 and 97/27847 disclose compounds that are useful as antidiabetic, antiobesity, anti-atherosclerosis and antihyperlipidemic agents, and which may exert their effect through activation of PPARs.

[0016] It is generally believed that glitazones exert their effects by binding to the peroxisome proliferator activated receptor (PPAR) family of receptors, controlling certain transcription elements having to do with the biological entities listed above. See Hulin et al., *Current Pharm. Design* (1996) 2, 85-102. In particular, PPAR γ has been implicated as the major molecular target for the glitazone class of insulin sensitizers.

[0017] A number of glitazones that are PPAR agonists have been approved for use in the treatment of diabetes. These are troglitazone, rosiglitazone and pioglitazone, all of which are primarily or exclusively PPAR γ agonists. Although glitazones are beneficial in the treatment of NIDDM, there have been some serious adverse events associated with the

use of the compounds. The most serious of these has been liver toxicity, which has resulted in a number of deaths. The most serious problems have occurred using troglitazone, which was recently withdrawn from the market because of toxicity concerns.

[0018] In addition to potential hepatotoxicity, there are several shortcomings associated with the glitazones: (1) Monotherapy for NIDDM produces modest efficacy - reductions in average plasma glucose of $\approx 20\%$ or a decline from $\approx 9.0\%$ to $\approx 8.0\%$ in HemoglobinA1C. (2) There is room for improvement in lipid effects; troglitazone causes a slight increase in LDL cholesterol, and triglyceride lowering is modest relative to the effect of fibrates; results reported to date with rosiglitazone suggest no effect on triglycerides and a possible net increase in the LDL:HDL ratio. Currently available data on pioglitazone appear to indicate that it lowers triglycerides modestly and may also have a neutral or positive effect on LDL vs. HDL (i.e. slight HDL raising with no effect on LDL). (3) All three glitazones have been associated with significant weight gain as well as other AE's (mild edema and mild anemia). These shortcomings provide an opportunity to develop better insulin sensitizers for Type 2 diabetes which function via similar mechanism(s) of action.


[0019] Because of the problems that have occurred with the glitazones, researchers in a number of laboratories have been investigating classes of PPAR agonists that are not glitazones and do not contain 1,3-thiazolidinedione moieties, but that modulate the three known PPAR subtypes, in concert or in isolation, to varying degrees (as measured by intrinsic potency, maximal extent of functional response or spectrum of changes in gene expression). Such classes of compounds are expected to be useful in the treatment of diabetes and associated conditions, dyslipidemias and associated conditions and several other indications and may be free of some of the side effects that have been found in many of the glitazones.

SUMMARY OF THE INVENTION

[0020] The class of compounds described herein is a new class of PPAR agonists that do not contain a 1,3-thiazolidinedione moiety and therefore are not glitazones. The class of compounds includes compounds that are primarily PPAR γ agonists and PPAR γ partial agonists. Some compounds may also have PPAR α activity in addition to the PPAR γ activity, so that the compounds are mixed PPAR α/γ agonists. These compounds are useful in the treatment, control and/or prevention of diabetes, hyperglycemia, and insulin resistance. The compounds of the invention exhibit reduced side effects relating to body and heart weight gain in preclinical animal studies compared with other PPAR γ compounds including rosiglitazone.

[0021] The compounds may also be useful in the treatment of mixed or diabetic dyslipidemia and other lipid disorders (including isolated hypercholesterolemia as manifested by elevations in LDL-C and/or non-HDL-C and/or hyperapobliproteinemia, hypertriglyceridemia and/or increase in triglyceride-rich-lipoproteins, or low HDL cholesterol concentrations), atherosclerosis, obesity, vascular restenosis, inflammatory conditions, neoplastic conditions, psoriasis, polycystic ovary syndrome and other PPAR mediated diseases, disorders and conditions.

[0022] The present invention is directed to compounds of formula I:



I

wherein:

R¹ is methyl, optionally substituted with 1-3 F;

R², R³ and R⁴ are each independently selected from the group consisting of H, halogen, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, aryl, OC₁-C₆ alkyl, OC₂-C₆ alkenyl, OC₂-C₆ alkynyl, O-aryl, OH, SC₁-C₆ alkyl, SC₂-C₆ alkenyl, SC₂-C₆ alkynyl, SO₂C₁-C₆ alkyl, SO₂C₂-C₆ alkenyl, SO₂C₂-C₆ alkynyl, OCON(R₅)₂, OCO

(C₁-C₆-alkyl) and CN, wherein all instances of alkyl, alkenyl and alkynyl are optionally linear or branched and all instances of alkyl, alkenyl, alkynyl, cycloalkyl and aryl are optionally substituted with 1-5 substituents independently selected from the group consisting of halogen, aryl, O-aryl and OMe;

R⁵ and R⁶ are, at each occurrence, independently selected from the group consisting of H, F, OH and C₁-C₅ alkyl, and R⁵ and R⁶ groups that are on the same carbon atom optionally may be joined to form a C₃-C₆ cycloalkyl group;

R⁷ and R⁸ are each independently selected from the group consisting of H, F, and C₁₋₅ alkyl, or R⁷ and R⁸ optionally may be joined to form a C₃-C₆ cycloalkyl group;

R⁹ is selected from the group consisting of H and C₁-C₅ alkyl, said alkyl being optionally linear or branched;

Ar¹ is phenyl, 1-naphthyl, 2-naphthyl, pyridyl or quinolyl wherein Ar¹ is substituted with 1-3 groups independently selected from R⁴;

X is selected from the group consisting of C=O, S(O)₂, CH₂, CH(CH₃), C(CH₃)₂, CF₂, and cyclopropylidene;

Y is O or S; and

n is 0-5;

and pharmaceutically acceptable salts and prodrugs thereof.

[0023] The present compounds are effective in lowering glucose, lipids, and insulin in diabetic animals and lipids in non-diabetic animals. The compounds are expected to be efficacious in the treatment, control and/or prevention of non-insulin dependent diabetes mellitus (NIDDM) in humans and in the treatment, control, and/or prevention of conditions associated with NIDDM, including hyperlipidemia, dyslipidemia, obesity, hypercholesterolemia, hypertriglyceridemia, atherosclerosis, vascular restenosis, inflammatory conditions, neoplastic conditions, and other PPAR mediated diseases, disorders and conditions.

DETAILED DESCRIPTION OF THE INVENTION

[0024] The invention has numerous embodiments. It provides compounds of formula I, including pharmaceutically acceptable salts of these compounds, prodrugs of these compounds, and pharmaceutical compositions comprising any of the compounds described and a pharmaceutically acceptable carrier.

[0025] In one embodiment, in compounds having the formula I, R¹ is CH₃.

[0026] In another embodiment of compounds having the formula I, R¹ is CH₃;

[0027] R², R³, and R⁴ are each independently selected from the group consisting of H, halogen, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, aryl, OC₁-C₆ alkyl, OC₂-C₆ alkenyl, OC₂-C₆ alkynyl, O-aryl, OH, SC₁-C₆ alkyl, SC₂-C₆ alkenyl, SC₂-C₆ alkynyl, OCON(R₅)₂, OCO(C₁-C₆-alkyl) and CN, wherein all instances of alkyl, alkenyl and alkynyl are optionally linear or branched and all instances of alkyl, alkenyl, alkynyl, cycloalkyl and aryl are optionally substituted with 1-5 substituents independently selected from the group consisting of halogen, aryl, O-aryl and OMe; and

[0028] X is selected from the group consisting of C=O, CH₂, CH(CH₃), C(CH₃)₂, CF₂, and cyclopropylidene.

[0029] In another embodiment, in compounds having the formula I, R², R³, and R⁴ are each independently selected from the group consisting of H, OCH₃, OCF₃, F, Cl and CH₃, where CH₃ is optionally substituted with 1-3 groups independently selected from F, Cl, and OCH₃. In more specific embodiments, R², R³, and R⁴ are each independently selected from the group consisting of H, OCH₃, OCF₃, and Cl.

[0030] In another group of compounds having the formula I, R⁵ and R⁶ are H.

[0031] In another group of compounds having the formula I, R⁷ and R⁸ are each independently CH₃ or H.

[0032] In preferred groups of compounds having the formula I, R⁹ is H.

[0033] In other compounds having formula I, X is C=O.

[0034] In other compounds having formula I, Y is O.

[0035] In another group of compounds having formula I, n is 0, 1, or 2. In a more specific subset of this group of compounds, n is 1.

[0036] Another group of compounds having formula I includes compounds in which Ar¹ is phenyl, 1-naphthyl or 2-naphthyl. A subset of this group of compounds includes compounds in which Ar¹ is phenyl or 2-naphthyl. In either case, Ar¹ is substituted with 1-3 groups independently selected from R⁴.

[0037] In preferred groups of compounds, aryl substituents are phenyl groups.

[0038] A preferred set of compounds having formula I has the following substituents:

R¹ is CH₃;

R² is selected from the group consisting of H, OCH₃, and OCF₃;

R³, R⁵, R⁶, and R⁹ are H;

R⁴ is selected from the group consisting of H, Cl, and OCH₃;

R⁷ and R⁸ are each independently selected from the group consisting of H and CH₃;

X is- C=O;

Y is O;

and n is 1.

[0039] Specific examples of compounds of this invention are provided as Examples 1-31, named below:

Example 1: (2S)-2-(3-([1-(4-Methoxybenzoyl)-2-methyl-5-(trifluoromethoxy)-1*H*-indol-3-yl]methyl}phenoxy)propanoic acid

Example 2: 2-(2-([1-(4-Chlorobenzoyl)-2-methyl-5-(methoxy)-1*H*-indol-3-yl]methyl}phenoxy)acetic acid

Example 3: 2-(3-([1-(4-Chlorobenzoyl)-2-Methyl-5-(methoxy)-1*H*-indol-3-yl]methyl}phenoxy)acetic acid

Example 4: 2-(4-([1-(4-Chlorobenzoyl)-2-methyl-5-(methoxy)-1*H*-indol-3-yl]methyl}phenoxy)acetic acid

Example 5: 2-(2-([1-(4-Chlorobenzoyl)-2-methyl-5-(methoxy)-1*H*-indol-3-yl]methyl}phenoxy)propanoic acid

Example 6: 2-(3-([1-(4-Chlorobenzoyl)-2-methyl-5-(methoxy)-1*H*-indol-3-yl]methyl}phenoxy)propanoic acid

Example 7: 2-(4-([1-(4-Chlorobenzoyl)-2-methyl-5-(methoxy)-1*H*-indol-3-yl]methyl}phenoxy)propanoic acid

Example 8: 2-(2-([1-(4-Chlorobenzoyl)-2-methyl-5-(methoxy)-1*H*-indol-3-yl]methyl}phenoxy)-2-methylpropanoic acid

Example 9: 2-(3-([1-(4-Chlorobenzoyl)-2-methyl-5-(methoxy)-1*H*-indol-3-yl]methyl}phenoxy)-2-methylpropanoic acid

Example 10: 2-(4-([1-(4-Chlorobenzoyl)-2-methyl-5-(methoxy)-1*H*-indol-3-yl]methyl}phenoxy)-2-methylpropanoic acid

Example 11: 2-(2-([1-(4-Methoxybenzoyl)-2-methyl-5-(trifluoromethoxy)-1*H*-indol-3-yl]methyl}phenoxy)acetic acid

Example 12: 2-(3-([1-(4-Methoxybenzoyl)-2-Methyl-5-(trifluoromethoxy)-1*H*-indol-3-yl]methyl}phenoxy)acetic acid

Example 13: 2-(2-([1-(4-methoxybenzoyl)-2-Methyl-5-(trifluoromethoxy)-1*H*-indol-3-yl]methyl}phenoxy)propanoic acid

Example 14: 2-(2-([1-(2-naphthoyl)-2-Methyl-5-(trifluoromethoxy)-1*H*-indol-3-yl]methyl}phenoxy)propanoic acid

Example 15: (2R)-2-(2-([1-(4-methoxybenzoyl)-2-Methyl-5-(trifluoromethoxy)-1*H*-indol-3-yl]methyl}phenoxy)propanoic acid

Example 16: (2S)-2-(2-([1-(4-Methoxybenzoyl)-2-methyl-5-(trifluoromethoxy)-1*H*-indol-3-yl]methyl}phenoxy)propanoic acid






Example 17: 2-(3-([1-(4-Methoxybenzoyl)-2-methyl-5-(trifluoromethoxy)-1*H*-indol-3-yl]methyl}phenoxy)propanoic acid

acid






Example 18: 2-(3-[[1-(2-Naphthoyl)-2-methyl-5-(trifluoromethoxy)-1*H*-indol-3-yl]methyl]phenoxy)propanoic acidExample 19: 2-(3-[[1-(4-Chlorobenzoyl)-2-methyl-5-(trifluoromethoxy)-1*H*-indol-3-yl]methyl]phenoxy)propanoic acidExample 20: 2-(3-[[1-(2,4-Dichlorobenzoyl)-2-methyl-5-(trifluoromethoxy)-1*H*-indol-3-yl]methyl]phenoxy)propanoic acidExample 21: (2*R*)-2-(3-[[1-(4-Methoxybenzoyl)-2-methyl-5-(trifluoromethoxy)-1*H*-indol-3-yl]methyl]phenoxy)propanoic acidExample 22: (2*R*)-2-(3-[[1-(2-Naphthoyl)-2-methyl-5-(trifluoromethoxy)-1*H*-indol-3-yl]methyl]phenoxy)propanoic acidExample 23: (2*S*)-2-(3-[[1-(2-Naphthoyl)-2-methyl-5-(trifluoromethoxy)-1*H*-indol-3-yl]methyl]phenoxy)propanoic acidExample 24: 2-(2-[[1-(4-Methoxybenzoyl)-2-methyl-5-(trifluoromethoxy)-1*H*-indol-3-yl]methyl]phenoxy)-2-methylpropanoic acidExample 25: 2-(3-[[1-(4-Methoxybenzoyl)-2-methyl-2-(trifluoromethoxy)-1*H*-indol-3-yl]methyl]phenoxy)-2-methylpropanoic acidExample 26: 2-(2-[[1-(2-Naphthoyl)-2-methyl-5-(trifluoromethoxy)-1*H*-indol-3-yl]methyl]phenoxy)-2-methylpropanoic acidExample 27: 2-(3-[[1-(2-Naphthoyl)-2-methyl-5-(trifluoromethoxy)-1*H*-indol-3-yl]methyl]phenoxy)-2-methylpropanoic acidExample 28: (2*R*)-2-(3-{2-[1-(4-Methoxybenzoyl)-2-methyl-5-(trifluoromethoxy)-1*H*-indol-3-yl]ethyl}phenoxy)propionic acidExample 29: (2*S*)-2-{3-[1-(4-methoxybenzoyl)-2-methyl-5-(trifluoromethoxy)-1*H*-indol-3-yl]phenoxy}propionic acidExample 30: (2*S*)-2-(3-{1-[1-(4-Methoxybenzoyl)-2-methyl-5-(trifluoromethoxy)-1*H*-indol-3-yl]cyclopropyl}phenoxy)propanoic acidExample 31: 2-{3-[1-(4-Methoxybenzoyl)-2-methyl-5-(trifluoromethoxy)-1*H*-indol-3-yl]phenoxy}-2-methylpropanoic acid.**[0040]** The structures of these specific compounds are shown in the following Table of Examples:**TABLE OF EXAMPLES**

Example 1	
-----------	--





(continued)

Example 2	
Example 3	
Example 4	
Example 5	
Example 6	






(continued)

Example 7	
Example 8	
Example 9	
Example 10	
Example 11	






(continued)

Example 12	
Example 13	
Example 14	
Example 15	






(continued)

Example 16	
Example 17	
Example 18	
Example 19	
Example 20	


(continued)

Example 21	
Example 22	
Example 23	
Example 24	
Example 25	

(continued)

Example 26	
Example 27	
Example 28	
Example 29	
Example 30	

(continued)



[0041] The compounds as defined above are useful in the following methods of treating, controlling, and preventing diseases, as well as other diseases not listed below:

(1) a method for treating, controlling or preventing diabetes mellitus, and particularly non-insulin dependent diabetes mellitus, in a mammalian patient in need of such treatment which comprises administering to the patient a therapeutically effective amount of a compound of Formula I;

(2) a method for treating, controlling, or preventing hyperglycemia in a mammalian patient in need of such treatment which comprises administering to the patient a therapeutically effective amount of a compound of Formula I;

(3) a method for treating, controlling, or preventing lipid disorders, hyperlipidemia, or low HDL in a mammalian patient in need of such treatment which comprises administering to the patient a therapeutically effective amount of a compound of Formula I;

(4) a method for treating, controlling, or preventing obesity in a mammalian patient in need of such treatment which comprises administering to the patient a therapeutically effective amount of a compound of Formula I;

(5) a method for treating, controlling, or preventing hypercholesterolemia in a mammalian patient in need of such treatment which comprises administering to the patient a therapeutically effective amount of a compound of Formula I;

(6) a method for treating, controlling, or preventing hypertriglyceridemia in a mammalian patient in need of such treatment which comprises administering to the patient a therapeutically effective amount of a compound of Formula I;

(7) a method for treating, controlling, or preventing dyslipidemia, including low HDL cholesterol, in a mammalian patient in need of such treatment which comprises administering to the patient a therapeutically effective amount of a compound of Formula I;

(8) a method for treating, controlling, or preventing atherosclerosis in a mammalian patient in need of such treatment which comprises administering to the patient a therapeutically effective amount of a compound of Formula I. It is understood that the sequelae of atherosclerosis (angina, claudication, heart attack, stroke, etc.) are thereby treated.

Definitions

[0042] "Ac" is acetyl, which is $\text{CH}_3\text{C}(\text{O})-$.

[0043] "Alkyl", as well as other groups having the prefix "alk", such as alkoxy or alkanoyl, means carbon chains which may be linear or branched or combinations thereof, unless the carbon chain is defined otherwise. Examples of alkyl groups include methyl, ethyl, propyl, isopropyl, butyl, sec- and tert-butyl, pentyl, hexyl, heptyl, octyl, nonyl, and the like.

[0044] "Alkenyl" means carbon chains which contain at least one carbon-carbon double bond, and which may be linear or branched or combinations thereof. Examples of alkenyl include vinyl, allyl, isopropenyl, pentenyl, hexenyl, heptenyl, 1-propenyl, 2-butenyl, 2-methyl-2-butenyl, and the like.

[0045] "Alkynyl" means carbon chains which contain at least one carbon-carbon triple bond, and which may be linear or branched or combinations thereof. Examples of alkynyl include ethynyl, propargyl, 3-methyl-1-pentynyl, 2-heptynyl and the like.

[0046] "Cycloalkyl" means mono- or bicyclic saturated carbocyclic rings, each having from 3 to 10 carbon atoms, unless otherwise stated. The term also includes a monocyclic ring fused to an aryl group in which the point of attachment is on the nonaromatic portion. Examples of cycloalkyl include cyclopropyl, cyclopentyl, cyclohexyl, cycloheptyl, and the like.

[0047] "Aryl" (and "arylene") when used to describe a substituent or group in a structure means a monocyclic, bicyclic or tricyclic compound in which all the rings are aromatic and which contains only carbon ring atoms. The term "aryl" can also refer to an aryl group fused to a monocyclic cycloalkyl or monocyclic heterocycle in which the point(s) of attachment is on the aromatic portion. "Heterocyclyl," "heterocycle," and "heterocyclic" means a fully or partially saturated monocyclic, bicyclic or tricyclic ring system containing at least one heteroatom selected from N, S and O, each of said rings having

from 3 to 10 atoms. Examples of aryl substituents include phenyl and naphthyl. Aryl rings fused to cycloalkyls are found in indanyl, indenyl, and tetrahydronaphthyl. Examples of aryl fused to heterocyclic groups are found in 2,3-dihydrobenzofuranyl, benzopyranyl, 1,4-benzodioxanyl, and the like. Examples of heterocycles include tetrahydrofuran, piperazine, and morpholine. Preferred aryl groups are phenyl rings.

[0048] "Heteroaryl" (and heteroarylene) means a mono-, bi- or tricyclic aromatic ring containing at least one ring heteroatom selected from N, O and S (including SO and SO₂), with each ring containing 5 to 6 atoms. Examples of heteroaryl include pyrrolyl, isoxazolyl, isothiazolyl, pyrazolyl, pyridyl, oxazolyl, oxadiazolyl, thiadiazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, furanyl, triazinyl, thienyl, pyrimidyl, pyridazinyl, pyrazinyl, benzisoxazolyl, benzoxazolyl, benzothiazolyl, benzimidazolyl, benzofuranyl, benzothiophenyl (including S-oxide and dioxide), furo(2,3-b)pyridyl, quinolyl, indolyl, isoquinolyl, dibenzofuran and the like.

[0049] "Halogen" includes fluorine, chlorine, bromine and iodine.

[0050] "Me" represents methyl.

[0051] The term "composition," as in pharmaceutical composition, is intended to encompass a product comprising the active ingredient(s), and the inert ingredient(s) that make up the carrier, as well as any product which results, directly or indirectly, from combination, complexation or aggregation of any two or more of the ingredients, or from dissociation of one or more of the ingredients, or from other types of reactions or interactions of one or more of the ingredients. Accordingly, the pharmaceutical compositions of the present invention encompass any composition made by admixing a compound of the present invention and a pharmaceutically acceptable carrier.

Optical Isomers - Diastereomers - Geometric Isomers - Tautomers

[0052] Compounds of Formula I may contain one or more asymmetric centers and can thus occur as racemates and racemic mixtures, single enantiomers, diastereomeric mixtures and individual diastereomers. The present invention is meant to comprehend all such isomeric forms of the compounds of Formula I.

[0053] Some of the compounds described herein may contain olefinic double bonds, and unless specified otherwise, are meant to include both E and Z geometric isomers.

[0054] Some of the compounds described herein may exist with different points of attachment of hydrogen, referred to as tautomers. Such an example may be a ketone and its enol form, known as keto-enol tautomers. The individual tautomers as well as mixtures thereof are encompassed with compounds of Formula I.

[0055] Compounds of the Formula I having two asymmetric centers may be separated into diastereoisomeric pairs of enantiomers by, for example, fractional crystallization from a suitable solvent, for example methanol or ethyl acetate or a mixture thereof. The pair of enantiomers thus obtained, and enantiomeric pairs in general, may be separated into individual stereoisomers by conventional means, for example by the use of an optically active acid or base as a resolving agent or chiral separation columns.

[0056] Alternatively, any enantiomer of a compound of the general Formula I or Ia may be obtained by stereospecific synthesis using optically pure starting materials or reagents of known configuration.

Salts

[0057] The term "pharmaceutically acceptable salts" refers to salts prepared from pharmaceutically acceptable non-toxic bases or acids including inorganic or organic bases and inorganic or organic acids. Salts derived from inorganic bases include aluminum, ammonium, calcium, copper, ferric, ferrous, lithium, magnesium, manganic salts, manganous, potassium, sodium, zinc, and the like. Particularly preferred are the ammonium, calcium, magnesium, potassium, and sodium salts. Salts in the solid form may exist in more than one crystal structure, and may also be in the form of hydrates. Salts derived from pharmaceutically acceptable organic non-toxic bases include salts of primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines, and basic ion exchange resins, such as arginine, betaine, caffeine, choline, N,N'-dibenzylethylenediamine, diethylamine, 2-diethylaminoethanol, 2-dimethylaminoethanol, ethanolamine, ethylenediamine, N-ethyl-morpholine, N-ethylpiperidine, glucamine, glucosamine, histidine, hydrabamine, isopropylamine, lysine, methylglucamine, morpholine, piperazine, piperidine, polyamine resins, procaine, purines, theobromine, triethylamine, trimethylamine, tripropylamine, tromethamine, and the like.

[0058] When the compound of the present invention is basic, salts may be prepared from pharmaceutically acceptable non-toxic acids, including inorganic and organic acids. Such acids include acetic, benzenesulfonic, benzoic, camphorsulfonic, citric, ethanesulfonic, fumaric, gluconic, glutamic, hydrobromic, hydrochloric, isethionic, lactic, maleic, malic, mandelic, methanesulfonic, mucic, nitric, pamoic, pantothenic, phosphoric, succinic, sulfuric, tartaric, p-toluenesulfonic acid, and the like. Particularly preferred are citric, hydrobromic, hydrochloric, maleic, phosphoric, sulfuric, and tartaric acids.

[0059] It will be understood that, as used herein, references to the compounds of Formula I are meant to also include the pharmaceutically acceptable salts.

Metabolites - Prodrugs

[0060] Therapeutically active metabolites of other compounds, where the metabolites themselves fall within the scope of the claims herein, are also claimed. Prodrugs, which are compounds that are converted to the claimed compounds as they are being administered to a patient or after they have been administered to a patient, are also claimed as part of this invention. A non-limiting example of a prodrug of the carboxylic acids of this invention would be an ester of the carboxylic acid group, for example a C₁ to C₆ ester, which may be linear or branched, which metabolizes to a compound claimed herein. An ester which has functionality that makes it more easily hydrolyzed after administration to a patient may also be a prodrug.

[0061] Prodrugs of the class of compounds of this invention may be described as compounds having the Formula I, wherein R⁹ is now defined as a group that is easily removed under physiological conditions during or after administration to a mammalian patient to yield a compound having Formula I, where R⁹ is H, or the carboxylate anion thereof (in solution), or a pharmaceutically acceptable salt thereof, where the substituents and groups and values of n are as defined above for compounds having Formula I.

[0062] Examples of prodrugs of Formula I include compounds in which OR⁹ of the CO₂R⁹ group is selected from the group consisting of -OR¹⁰, -OCH₂OR¹⁰, -OCH(CH₃)OR¹⁰, -OCH₂OC(O)R¹⁰, -OCH(CH₃)OC(O)R¹⁰, -OCH₂OC(O)OR¹⁰, -OCH(CH₃)OC(O)OR¹⁰, -NR¹¹R¹¹, and -ONR¹¹R¹¹, where each R¹⁰ is independently selected from C₁₋₆ alkyl optionally substituted with one or two groups selected from -CO₂H, -CONH₂, -NH₂, -OH, -OAc, NHAc, and phenyl; and wherein each R¹¹ is independently selected from H and R¹⁰. Compounds having Formula I, where R⁹ has the chemical structure described above, are described as prodrugs. However, regardless of whether they are active as prodrugs, yielding -compounds or salts of Formula I, or whether they have a different means of exhibiting pharmaceutical activity, such compounds are included in this invention. Such compounds are claimed herein, regardless of the mechanism leading to their activity.

Utilities

[0063] Compounds of the present invention are potent ligands with agonist or partial agonist activity on the various peroxisome proliferator activator receptor subtypes, particularly PPAR γ . The compounds may also be ligands or agonists of the PPAR α subtype as well, resulting in mixed PPAR α/γ agonism or in agonism of mainly the PPAR α subtype. These compounds are useful in treating, controlling or preventing diseases, disorders or conditions, wherein the treatment is mediated by the activation of an individual PPAR subtype (γ or α) or a combination of PPAR subtypes (e.g. α/γ), and particularly the PPAR γ subtype.

[0064] Compounds of the present invention may be useful in treating, controlling or preventing many PPAR mediated diseases and conditions, including, but are not limited to, (1) diabetes mellitus, and especially non-insulin dependent diabetes mellitus (NIDDM), (2) hyperglycemia, (3) low glucose tolerance, (4) insulin resistance, (5) obesity, (6) lipid disorders, (7) dyslipidemia, (8) hyperlipidemia, (9) hypertriglyceridemia, (10) hypercholesterolemia, (11) low HDL levels, (12) high LDL levels, (13) atherosclerosis and its sequelae, (14) vascular restenosis, (15) irritable bowel syndrome, (16) inflammatory bowel disease, including Crohn's disease and ulcerative colitis, (17) other inflammatory conditions, (18) pancreatitis, (19) abdominal obesity, (20) neurodegenerative disease, (21) retinopathy, (22) neoplastic conditions, (23) adipose cell tumors, (24) adipose cell carcinomas, such as liposarcoma, (25) prostate cancer and other cancers, including gastric, breast, bladder and colon cancers, (26) angiogenesis, (27) Alzheimer's disease, (28) psoriasis, (29) high blood pressure, (30) Syndrome X, (31) ovarian hyperandrogenism (polycystic ovarian syndrome), and other disorders where insulin resistance is a component.

[0065] Another utility of the compounds of the invention is a method for the treatment, control, or prevention of hypercholesterolemia, atherosclerosis, low HDL levels, high LDL levels, hyperlipidemia, hypertriglyceridemia, and/or dyslipidemia, which comprises administering to a mammal in need of such treatment a therapeutically effective amount of a PPAR agonist or partial agonist having formula 1. The PPAR agonist may be used alone or advantageously may be administered with a cholesterol biosynthesis inhibitor, particularly an HMG-CoA reductase inhibitor such as lovastatin, simvastatin, pravastatin, fluvastatin, atorvastatin, rivastatin, itavastatin, or ZD-4522. The PPAR agonist may also be used advantageously in combination with other lipid lowering drugs such as cholesterol absorption inhibitors (for example stanol esters, sterol glycosides such as tiqueside, and azetidinones such as ezetimibe), ACAT inhibitors (such as avasimibe), and with niacin, bile acid sequestrants, microsomal triglyceride transport inhibitors, and bile acid reuptake inhibitors. These combination treatments may also be effective for the treatment, control or prevention of one or more related conditions selected from the group consisting of hypercholesterolemia, atherosclerosis, hyperlipidemia, hypertriglyceridemia, dyslipidemia, high LDL, and low HDL.

[0066] Another utility of the compounds of the invention is a method of treating inflammatory conditions, including inflammatory bowel disease, Crohn's disease, and ulcerative colitis by administering an effective amount of a PPAR agonist, which may be a PPAR α agonist, a PPAR γ agonist, or a PPAR α/γ dual agonist. Additional inflammatory diseases

that may be treated with the instant invention include gout, rheumatoid arthritis, osteoarthritis, multiple sclerosis, asthma, ARDS, psoriasis, vasculitis, ischemia/reperfusion injury, frostbite, and related diseases.

Administration and Dose Ranges

[0067] Any suitable route of administration may be employed for providing a mammal, especially a human, with an effective dose of a compound of the present invention. For example, oral, rectal, topical, parenteral, ocular, pulmonary, nasal, and the like may be employed. Dosage forms include tablets, troches, dispersions, suspensions, solutions, capsules, creams, ointments, aerosols, and the like. Preferably compounds of Formula I are administered orally.

[0068] The effective dosage of active ingredient employed may vary depending on the particular compound employed, the mode of administration, the condition being treated and the severity of the condition being treated. Such dosage may be ascertained readily by a person skilled in the art.

[0069] When treating or preventing diabetes mellitus and/or hyperglycemia or hypertriglyceridemia or other diseases for which compounds of Formula I are indicated, generally satisfactory results are obtained when the compounds of the present invention are administered at a daily dosage of from about 0.1 milligram to about 100 milligram per kilogram of animal body weight, preferably given as a single daily dose or in divided doses two to six times a day, or in sustained release form. For most large mammals, the total daily dosage is from about 1.0 milligrams to about 1000 milligrams; preferably from about 1 milligrams to about 50 milligrams. In the case of a 70 kg adult human, the total daily dose will generally be from about 7 milligrams to about 350 milligrams. This dosage regimen may be adjusted to provide the optimal therapeutic response.

Pharmaceutical Compositions

[0070] Another aspect of the present invention provides pharmaceutical compositions which comprise a compound of Formula I and a pharmaceutically acceptable carrier. The pharmaceutical compositions of the present invention comprise a compound of Formula I or a pharmaceutically acceptable salt or prodrug thereof as an active ingredient, as well as a pharmaceutically acceptable carrier and optionally other therapeutic ingredients. The term "pharmaceutically acceptable salts" refers to salts prepared from pharmaceutically acceptable non-toxic bases or acids including inorganic bases or acids and organic bases or acids.

[0071] The compositions include compositions suitable for oral, rectal, topical, parenteral (including subcutaneous, intramuscular, and intravenous), ocular (ophthalmic), pulmonary (nasal or buccal inhalation), or nasal administration, although the most suitable route in any given case will depend on the nature and severity of the conditions being treated and on the nature of the active ingredient. They may be conveniently presented in unit dosage form and prepared by any of the methods well-known in the art of pharmacy.

[0072] In practical use, the compounds of Formula I can be combined as the active ingredient in intimate admixture with a pharmaceutical carrier according to conventional pharmaceutical compounding techniques. The carrier may take a wide variety of forms depending on the form of preparation desired for administration, e.g., oral or parenteral (including intravenous). In preparing the compositions for oral dosage form, any of the usual pharmaceutical media may be employed, such as, for example, water, glycols, oils, alcohols, flavoring agents, preservatives, coloring agents and the like in the case of oral liquid preparations, such as, for example, suspensions, elixirs and solutions; or carriers such as starches, sugars, microcrystalline cellulose, diluents, granulating agents, lubricants, binders, disintegrating agents and the like in the case of oral solid preparations such as, for example, powders, hard and soft capsules and tablets, with the solid oral preparations being preferred over the liquid preparations.

[0073] Because of their ease of administration, tablets and capsules represent the most advantageous oral dosage unit form in which case solid pharmaceutical carriers are obviously employed. If desired, tablets may be coated by standard aqueous or nonaqueous techniques. Such compositions and preparations should contain at least 0.1 percent of active compound. The percentage of active compound in these compositions may, of course, be varied and may conveniently be between about 2 percent to about 60 percent of the weight of the unit. The amount of active compound in such therapeutically useful compositions is such that an effective dosage will be obtained. The active compounds can also be administered intranasally as, for example, liquid drops or spray.

[0074] The tablets, pills, capsules, and the like may also contain a binder such as gum tragacanth, acacia, corn starch or gelatin; excipients such as dicalcium phosphate; a disintegrating agent such as corn starch, potato starch, alginic acid; a lubricant such as magnesium stearate; and a sweetening agent such as sucrose, lactose or saccharin. When a dosage unit form is a capsule, it may contain, in addition to materials of the above type, a liquid carrier such as a fatty oil.

[0075] Various other materials may be present as coatings or to modify the physical form of the dosage unit. For instance, tablets may be coated with shellac, sugar or both. A syrup or elixir may contain, in addition to the active ingredient, sucrose as a sweetening agent, methyl and propylparabens as preservatives, a dye and a flavoring such as cherry or orange flavor.

[0076] Compounds of formula I may also be administered parenterally. Solutions or suspensions of these active compounds can be prepared in water suitably mixed with a surfactant such as hydroxy-propylcellulose. Dispersions can also be prepared in glycerol, liquid polyethylene glycols and mixtures thereof in oils. Under ordinary conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms.

[0077] The pharmaceutical forms suitable for injectable use include sterile aqueous solutions or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions. In all cases, the form must be sterile and must be fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (e.g. glycerol, propylene glycol and liquid polyethylene glycol), suitable mixtures thereof, and vegetable oils.

Combination Therapy

[0078] Compounds of Formula I may be used in combination with other drugs that may also be useful in the treatment, prevention, suppression or amelioration of the diseases or conditions for which compounds of Formula I are useful. Such other drugs may be administered, by a route and in an amount commonly used therefor, contemporaneously or sequentially with a compound of Formula I. When a compound of Formula I is used contemporaneously with one or more other drugs, a pharmaceutical composition in unit dosage form containing such other drugs and the compound of Formula I is preferred. However, the combination therapy also includes therapies in which the compound of Formula I and one or more other drugs are administered on different overlapping schedules. It is also contemplated that when used in combination with one or more other active ingredients, the compound of the present invention and the other active ingredients may be used in lower doses than when each is used singly. Accordingly, the pharmaceutical compositions of the present invention include those that contain one or more other active ingredients, in addition to a compound of Formula I.

[0079] Examples of other active ingredients that may be administered in combination with a compound of Formula I, and either administered separately or in the same pharmaceutical composition, include, but are not limited to:

- (a) (i) other PPAR agonists such as the glitazones (e.g. troglitazone, pioglitazone, englitazone, MCC-555, rosiglitazone, and the like), and compounds disclosed in WO97/27857, 97/28115, 97/28137 and 97/27847; (ii) biguanides such as metformin and phenformin; (iii) protein tyrosine phosphatase-1B (PTP-1B) inhibitors, and (iv) dipeptidyl peptidase IV (DP-N) inhibitors;
- (b) insulin or insulin mimetics;
- (c) sulfonylureas such as tolbutamide and glipizide, or related materials;
- (d) α -glucosidase inhibitors (such as acarbose);
- (e) cholesterol lowering agents such as (i) HMG-CoA reductase inhibitors (lovastatin, simvastatin, pravastatin, fluvastatin, atorvastatin, rivastatin, itavastatin, ZD-4522 and other statins), (ii) sequestrants (cholestyramine, colestipol, and dialkylaminoalkyl derivatives of a cross-linked dextran), (iii) nicotinic alcohol, nicotinic acid or a salt thereof, (iv) PPAR α agonists such as fenofibric acid derivatives (gemfibrozil, clofibrate, fenofibrate and benzafibrate), (v) PPAR α/γ dual agonists, such as KRP-297, (vi) inhibitors of cholesterol absorption, such as for example beta-sitosterol, (vii) acyl CoA:cholesterol acyltransferase inhibitors, such as for example avasimibe, and (viii) anti-oxidants, such as probucol;
- (f) PPAR δ agonists such as those disclosed in WO97/28149;
- (g) antiobesity compounds such as fenfluramine, dexfenfluramine, phentiramine, sulbitramine, orlistat, neuropeptide Y5 inhibitors, and β_3 adrenergic receptor agonists;
- (h) an ileal bile acid transporter inhibitor; and
- (i) agents intended for use in inflammatory conditions such as aspirin, non-steroidal anti-inflammatory drugs, glucocorticoids, azulfidine, and cyclooxygenase 2 selective inhibitors.

[0080] The above combinations include combinations of a compound of the present invention not only with one other active compound, but also with two or more other active compounds. Non-limiting examples include combinations of compounds having Formula I with two or more active compounds selected from biguanides, sulfonylureas, HMG-CoA reductase inhibitors, other PPAR agonists, PTP-1B inhibitors, DP-IV inhibitors, and anti-obesity compounds.

BIOLOGICAL ASSAYS

A) PPAR Binding Assays

[0081] For preparation of recombinant human PPAR γ , PPAR δ , and PPAR α : Human PPAR γ_2 , human PPAR δ and

human PPAR α were expressed as gst-fusion proteins in *E. coli*. The full length human cDNA for PPAR γ_2 was subcloned into the pGEX-2T expression vector (Pharmacia). The full length human cDNAs for PPAR δ and PPAR α were subcloned into the pGEX-KT expression vector (Pharmacia). *E. coli* containing the respective plasmids were propagated, induced, and harvested by centrifugation. The resuspended pellet was broken in a French press and debris was removed by centrifugation at 12,000 X g. Recombinant human PPAR receptors were purified by affinity chromatography on glutathione sepharose. After application to the column, and one wash, receptor was eluted with glutathione. Glycerol (10%) was added to stabilize the receptor and aliquots were stored at -80°C. For binding to PPAR γ , an aliquot of receptor was incubated in TEGM (10 mM Tris, pH 7.2, 1 mM EDTA, 10% glycerol, 7 μ L/100 mL β -mercaptoethanol, 10 mM Na molybdate, 1 mM dithiothreitol, 5 μ g/mL aprotinin, 2 μ g/mL leupeptin, 2 μ g/mL benzamidine and 0.5 mM PMSF) containing 0.1% non-fat dry milk and 10 nM [3 H $_2$] AD5075, (21 Ci/mmol), \pm test compound as described in Berger et al (Novel peroxisome proliferator-activated receptor (PPAR γ) and PPAR δ ligands produce distinct biological effects. J. Biol. Chem. (1999), 274: 6718-6725. Assays were incubated for ~16 hr at 4°C in a final volume of 150 μ L. Unbound ligand was removed by incubation with 100 μ L dextran/gelatin-coated charcoal, on ice, for ~10 min. After centrifugation at 3000 rpm for 10 min at 4°C, 50 μ L of the supernatant fraction was counted in a Topcount.

[0082] For binding to PPAR δ , an aliquot of receptor was incubated in TEGM (10 mM Tris, pH 7.2, 1 mM EDTA, 10% glycerol, 7 μ L/100 mL β -mercaptoethanol, 10 mM Na molybdate, 1 mM dithiothreitol, 5 μ g/mL aprotinin, 2 μ g/mL leupeptin, 2 μ g/mL benzamide and 0.5 mM PMSF) containing 0.1% non-fat dry milk and 2.5 nM [3 H $_2$]L-783483, (17 Ci/mmol), \pm test compound as described in Berger et al (Novel peroxisome proliferator-activated receptor (PPAR γ) and PPAR δ ligands produce distinct biological effects. 1999 J Biol Chem 274: 6718-6725). (L-783483 is 3-chloro-4-(3-(7-propyl-3-trifluoromethyl-6-benz-[4,5]-isoxazoloxy)propylthio)phenylacetic acid, Ex. 20 in WO 97/28137). Assays were incubated for ~16 hr at 4°C in a final volume of 150 μ L. Unbound ligand was removed by incubation with 100 μ L dextran/gelatin-coated charcoal, on ice, for ~10 min. After centrifugation at 3000 rpm for 10 min at 4°C, 50 μ L of the supernatant fraction was counted in a Topcount.

[0083] For binding to PPAR α , an aliquot of receptor was incubated in TEGM (10 mM Tris, pH 7.2, 1 mM EDTA, 10% glycerol, 7 μ L/100 mL β -mercaptoethanol, 10 mM Na molybdate, 1 mM dithiothreitol, 5 μ g/mL aprotinin, 2 μ g/mL leupeptin, 2 μ g/mL benzamide and 0.5 mM PMSF) containing 0.1% non-fat dry milk and 5.0 nM [3 H $_2$]L-797773, (34 Ci/mmol), \pm test compound. (L-797773 is (3-(4-(3-phenyl-7-propyl-6-benz-[4,5]-isoxazoloxy)butyloxy))phenylacetic acid, Ex.62 in WO 97/28137). Assays were incubated for ~16 hr at 4°C in a final volume of 150 μ L. Unbound ligand was removed by incubation with 100 μ L dextran/gelatin-coated charcoal, on ice, for ~10 min. After centrifugation at 3000 rpm for 10 min at 4°C, 50 μ L of the supernatant fraction was counted in a Topcount.

B) Gal-4 hPPAR Transactivation Assays

[0084] The chimeric receptor expression constructs, pcDNA3-hPPAR γ /GAL4, pcDNA3-hPPAR δ /GAL4, pcDNA3-hPPAR α /GAL4 were prepared by inserting the yeast GAL4 transcription factor DBD adjacent to the ligand binding domains (LBDs) of hPPAR γ , hPPAR δ , hPPAR α , respectively. The reporter construct, pUAS(5X)-tk-luc was generated by inserting 5 copies of the GALA response element upstream of the herpes virus minimal thymidine kinase promoter and the luciferase reporter gene. pCMV-lacZ contains the galactosidase Z gene under the regulation of the cytomegalovirus promoter. COS-1 cells were seeded at 12 X 10³ cells/well in 96 well cell culture plates in high glucose Dulbecco's modified Eagle medium (DMEM) containing 10% charcoal stripped fetal calf serum (Gemini Bio-Products, Calabasas, CA), nonessential amino acids, 100 units/ml Penicillin G and 100 mg/ml Streptomycin sulfate at 37 °C in a humidified atmosphere of 10% CO₂. After 24 h, transfections were performed with Lipofectamine (GIBCO BRL, Gaithersburg, MD) according to the instructions of the manufacturer. Briefly, transfection mixes for each well contained 0.48 μ L of Lipofectamine, 0.00075 μ g of pcDNA3-PPAR/GAL4 expression vector, 0.045 μ g of pUAS(5X)-tk-luc reporter vector and 0.0002 μ g of pCMV-lacZ as an internal control for transactivation efficiency. Cells were incubated in the transfection mixture for 5 h at 37° C in an atmosphere of 10% CO₂. The cells were then incubated for ~48 h in fresh high glucose DMEM containing 5% charcoal stripped fetal calf serum, nonessential amino acids, 100 units/ml Penicillin G and 100 mg/ml Streptomycin sulfate \pm increasing concentrations of test compound. Since the compounds were solubilized in DMSO, control cells were incubated with equivalent concentrations of DMSO; final DMSO concentrations were \leq 0.1%, a concentration which was shown not to effect transactivation activity. Cell lysates were produced using Reporter Lysis Buffer (Promega, Madison, WI) according to the manufacturer's instructions. Luciferase activity in cell extracts was determined using Luciferase Assay Buffer (Promega, Madison, WI) in an ML3000 luminometer (Dynatech Laboratories, Chantilly, VA). β -galactosidase activity was determined using β -D-galactopyranoside (Calbiochem, San Diego, CA). Partial agonism was determined by comparison of maximal transactivation activity with standard PPAR agonists such as rosiglitazone and pioglitazone. If the maximal stimulation of transactivation was less than 50% of the effect observed with standard compounds, then the compound was designated as a partial agonist.


C) *In Vivo* Studies

[0085] Male db/db mice (10-11 week old C57B1/KFJ, Jackson Labs, Bar Harbor, ME) were housed 5/cage and allowed *ad lib.* access to ground Purina rodent chow and water. The animals, and their food, were weighed every 2 days and were dosed daily by gavage with vehicle (0.5% carboxymethylcellulose) \pm test compound at the indicated dose. Drug suspensions were prepared daily. Plasma glucose, and triglyceride concentrations were determined from blood obtained by tail bleeds at 3-5 day intervals during the study period. Glucose, and triglyceride, determinations were performed on a Boehringer Mannheim Hitachi 911 automatic analyzer (Boehringer Mannheim, Indianapolis, IN) using heparinized plasma diluted 1:6 (v/v) with normal saline. Lean animals were age-matched heterozygous mice maintained in the same manner.

EXAMPLES


[0086] The following Examples are provided to illustrate the invention, including methods of making the compounds of the invention, and are not to be construed as limiting the invention in any manner. The scope of the invention is defined by the appended claims.

Scheme1. Synthesis of Example 1



Example 1


[0087]



(2S)-2-(3-([1-(4-Methoxybenzoyl)-2-methyl-5-(trifluoromethoxy)-1*H*-indol-3-yl]methyl)phenoxy)propanoic acid

Step 1. (3E)-4-(3-Hydroxyphenyl)-3-buten-2-one (2):

[0088]




3-Hydroxybenzaldehyde (4.0 g, 32.8 mmole) was dissolved in THF (165 mL) and 1-triphenylphosphoranylidene-2-propanone (20.9 g, 65.6 mmole) was added. The solution was heated to reflux until TLC monitoring determined reaction was complete. Silica gel chromatography with 20% ethyl acetate in hexanes as eluent was used to isolate the title compound in 65% yield.

¹H NMR (400 MHz, CDCl₃): δ 7.50 (d, 1H), 7.30 (t, 1H), 7.14 (d, 1H), 7.09 (s, 1H), 6.93 (dd, 1H), 6.72 (d, 1H), 5.70 (s, 1H), 2.42 (s, 3H).

Step 2. 4-(3-Hydroxyphenyl)-2-butanone (3):

[0089]




Compound 2 from Step 1 (2.0 g, 12.3 mmole) was dissolved in ethyl acetate (120 mL). The reaction vessel was evacuated and charged with nitrogen gas. Then 10% palladium on activated charcoal was added (200 mg). The reaction vessel was then evacuated and charged with hydrogen gas and the reaction monitored by TLC. After 1 hour the reaction was filtered over celite and the filtrate evaporated to give the title compound in nearly quantitative yield.

¹H NMR (400 MHz, CDCl₃): δ 7.17 (t, 1H), 6.78 (d, 1H), 6.70 (s, 1H), 6.69 (d, 1H), 5.00 (br s, 1H), 2.88 (t, 2H), 2.78 (t, 2H), 2.18 (s, 3H).

Step 3. 3-([2-Methyl-5-(trifluoromethoxy)-1*H*-indol-3-yl]methyl)phenol (4):

[0090]




p-Trifluoromethoxyphenyl hydrazine hydrochloride (2.58 gr, 11.3 mmole) and Compound 3 (1.86 gl, 11.3 mmole) were stirred in acetic acid at 110°C for 45 minutes, at which time reaction was complete by HPLC. Acetic acid was removed by rotary evaporation and the resulting residue was purified by normal phase chromatography to give an orange oil (2.83 gr, 78%).

¹H NMR (400MHz, CDCl₃): δ 7.91 (br s, 1H), 7.1-7.25 (m, 3H), 6.99 (m, 1H), 6.83 (d, 1H), 6.62 (m, 2H), 5.05 (br s, 1H), 3.99 (s, 2H), 2.38 (s, 3H).

Step 4. Allyl (2S)-2-(3-([2-methyl-5-(trifluoromethoxy)-1H-indol-3-yl]methyl)phenoxy)propanoate (5)

[0091]




The phenolic indole (4) (50 mg, 0.16 mmole) was dissolved in dichloromethane (2 mL). To the phenol solution was added (S)-allyl lactate (24 mg, 0.19 mmole), triphenylphosphine (50 mg, 0.19 mmole), and diethylazodicarboxylate (DEAD) (0.030 mL, 0.19 mmole) and the reaction was monitored by TLC. Once complete the reaction was purified by silica gel chromatography to give the title compound (44.1 mg, 64%).

¹H NMR (400MHz, CDCl₃): δ 7.93 (s, 1H), 7.24 (d, 1H), 7.21 (s, 1H), 7.17 (t, 1H), 6.98 (d, 1H), 6.86 (d, 1H), 6.73 (s, 1H), 6.68 (d, 1H), 5.84 (m, 1H), 5.22 (m, 2H), 4.72 (q, 1H), 4.58 (m, 2H), 4.01 (s, 2H), 2.39 (s, 3H), 1.59 (d, 3H)

Step 5. Allyl (2S)-2-(3-([1-(4-Methoxybenzoyl)-2-methyl-5-(trifluoromethoxy)-1H-indol-3-yl]methyl)phenoxy)propanoate (6)

[0092]




Compound 5 (467 mg, 1.1 mmole) was dissolved in tetrahydrofuran (11 mL) and cooled to -78°C. Sodium bis(trimethylsilyl) amide 1.3 mL of a 1.0N solution in THF) was added and the reaction mixture was stirred for 10 minutes. p-Anisoyl chloride (221 mg, 1.3 mmole) was then added. The reaction was warmed to 0°C then quenched with saturated ammonium chloride and diluted with ether (100 ml). The ether layer was washed with water (2X), brine (1X) and dried over sodium sulfate followed by filtration and evaporation of the filtrate giving the title compound after silica gel chromatography (490 mg, 79%).

¹H NMR (400MHz, CDCl₃): δ 7.77 (d, 2H), 7.21 (t, 1H), 7.18 (s, 1H), 7.24 (d, 1H), 7.01 (d, 2H), 6.91 (d, 1H), 6.87 (d,

1H), 6.76 (s, 1H), 6.71 (d, 1H), 5.84 (m, 1H), 5.22 (m, 2H), 4.72 (q, 1H), 4.58 (m, 2H), 4.02 (s, 2H), 3.93 (s, 3H), 2.40 (s, 3H), 1.61 (d, 3H).

Step 6. (2S)-2-(3-[[1-(4-Methoxybenzoyl)-2-methyl-5-(trifluoromethoxy)-1H-indol-3-yl]methyl]phenoxy)propanoic acid (7)

[0093]



Compound 6 (490 mg, 0.86 mmole) was dissolved in DMF (9 mL). 5,5-Dimethyl-1,3-cyclohexanedione (181 mg, 1.29 mmole), N,N-diisopropylethylamine (0.225 mL, 1.29mmole) and (tetrakis(phenylphosphine)palladium (50 mg, 0.043 mmole) were then added and the solution stirred for 2 hours. Then aqueous ammonium chloride was added and the solution was extracted repeatedly with dichloromethane. The combined organics were dried over sodium sulfate, filtered and the filtrate was evaporated. The crude isolate was then purified by silica gel chromatography to give the title compound (395 mg, 87%).


¹H NMR (400MHz, CDCl₃): 7.76 (d, 2H), 7.23 (t, 1H), 7.18 (s, 1H), 7.02 (d, 1H), 7.01 (d, 2H), 6.92 (d, 1H), 6.91 (d, 1H), 6.76 (s, 1H), 6.75 (d, 1H), 4.76 (q, 1H), 4.05 (s, 2H), 3.94 (s, 3H), 2.41 (s, 3H), 1.63 (d, 3H).

Examples 2-31

[0094] The following compounds were prepared in a similar fashion to that shown in the above scheme and in Example 1 from commercially available starting materials.

Example 2

[0095]




2-(2-[[1-(4-Chlorobenzoyl)-2-methyl-5-(methoxy)-1H-indol-3-yl]methyl]phenoxy)acetic acid

¹H NMR (400 MHz, CDCl₃): δ 7.69(d,2H), 7.49(d,2H), 7.21 (dt,1H), 7.05 (dd, 1H), 6.93 (m,2H), 6.83 (m,2H), 6.66 (dd, 1H), 4.78 (s, 2H), 4.10 (s, 2H), 3.74 (s, 3H), 2.38 (s, 3H).

Example 3

[0096]




2-(3-([1-(4-Chlorobenzoyl)-2-Methyl-5-(methoxy)-1*H*-indol-3-yl]methyl)phenoxy)acetic acid

^1H NMR (400 MHz, CDCl_3): δ 7.69(d, 2H), 7.49(d, 2H), 7.24 (t, 1H), 6.94 (d, 1H), 6.88 (d, 1H), 6.80 (d, 2H), 6.76 (dd, 1H), 6.66 (dd, 1H), 4.64 (s, 2H), 4.03 (s, 2H), 3.77 (s, 3H), 2.39 (s, 3H).

Example 4

[0097]




2-(4-([1-(4-Chlorobenzoyl)-2-methyl-5-(methoxy)-1*H*-indol-3-yl]methyl)phenoxy)acetic acid

^1H NMR (400 MHz, CDCl_3): δ 7.68(d, 2H), 7.49(d, 2H), 7.16 (m, 2H), 6.85 (m, 3H), 6.79 (d, 1H), 6.66 (dd, 1H), 4.64 (s, 2H), 3.99 (s, 2H), 3.76 (s, 3H), 2.39 (s, 3H).

Example 5

[0098]




2-(2-([1-(4-Chlorobenzoyl)-2-methyl-5-(methoxy)-1*H*-indol-3-yl]methyl)phenoxy)propanoic acid

^1H NMR (400 MHz, CDCl_3): δ 7.69(d, 2H), 7.48(d, 2H), 7.18(m, 1H), 7.05 (d, 1H), 6.91 (m, 2H), 6.82 (m, 2H), 6.65 (dd, 1H), 4.92 (q, 1H), 4.09 (q, 2H), 3.74 (s, 3H), 2.37 (s, 3H), 1.71 (d, 3H).

Example 6

[0099]




2-(3-([1-(4-Chlorobenzoyl)-2-methyl-5-(methoxy)-1*H*-indol-3-yl]methyl)phenoxy)propanoic acid

^1H NMR (400 MHz, CDCl_3): δ 7.69(d, 2H), 7.49(d, 2H), 7.21(t, 1H), 6.90 (d, 1H), 6.88 (d, 1H), 6.81 (d, 1H), 6.78 (br t, 1H), 6.73 (dd, 1H), 6.65 (dd, 1H), 4.75 (q, 1H), 4.01 (s, 2H), 3.77 (s, 3H), 2.38 (s, 3H) 1.64 (d, 3H).

Example 7

[0100]




2-(4-([1-(4-Chlorobenzoyl)-2-methyl-5-(methoxy)-1*H*-indol-3-yl]methyl)phenoxy)propanoic acid

^1H NMR (300 MHz, CDCl_3): δ 7.66 (d, 2H), 7.48 (d, 2H), 7.17 (m, 2H), 6.80 (m, 4H), 6.62 (d, 1H), 4.74 (q, 1H), 3.96 (s, 2H), 3.75 (s, 3H), 2.38 (s, 3H), 1.61 (d, 3H); ES-MS ($M+1$) 478, 480.

Example 8

[0101]




2-(2-([1-(4-Chlorobenzoyl)-2-methyl-5-(methoxy)-1*H*-indol-3-yl]methyl)phenoxy)-2-methylpropanoic acid

^1H NMR (400 MHz, CDCl_3): δ 7.69(d, 2H), 7.49(d, 2H), 7.13(m, 1H), 7.01 (d, 1H), 6.89 (m, 3H), 6.77 (d, 1H), 6.66 (dd, 1H), 4.04 (s, 2H), 3.73(s, 3H), 2.36 (s, 3H) 1.69(s, 6H).

Example 9


[0102]



2-(3-([1-(4-Chlorobenzoyl)-2-methyl-5-(methoxy)-1*H*-indol-3-yl]methyl)phenoxy)-2-methylpropanoic acid
¹H NMR (400 MHz, CDCl₃): δ 7.69(d, 2H), 7.49(d, 2H), 7.19(m, 2H), 6.96 (d, 1H), 6.88 (d, 1H), 6.79 (m, 2H), 6.66 (dd, 1H), 4.00 (s, 2H), 3.76 (s, 3H), 2.38 (s, 3H) 1.55 (s, 6H).

Example 10


[0103]



2-(4-([1-(4-Chlorobenzoyl)-2-methyl-5-(methoxy)-1*H*-indol-3-yl]methyl)phenoxy)-2-methylpropanoic acid
¹H NMR (400 MHz, CDCl₃): δ 7.69(d, 2H), 7.49(d, 2H), 7.24(d, 2H), 6.88 (m, 3H), 6.79 (d, 1H), 6.66 (d, 1H), 4.00 (s, 2H), 3.76 (s, 3H), 2.39 (s, 3H) 1.57 (s, 6H).

Example 11


[0104]



2-(2-([1-(4-Methoxybenzoyl)-2-methyl-5-(trifluoromethoxy)-1*H*-indol-3-yl]methyl)phenoxy)acetic acid
¹H NMR (400 MHz, CDCl₃): δ 7.74 (d, 2H), 7.28 (s, 1H), 7.21 (dt, 1H), 7.05 (m, 2H), 7.01 (d, 2H), 6.92 (t, 1H), 6.89 (dd, 1H), 6.82 (d, 1H), 4.78 (s, 2H), 4.13 (s, 2H), 3.93 (s, 3H), 2.41(s, 3H).

Example 12

[0105]




2-(3-([1-(4-methoxybenzoyl)-2-methyl-5-(trifluoromethoxy)-1H-indol-3-yl]methyl)phenoxy)acetic acid

^1H NMR (400 MHz, CDCl_3): δ 7.78 (d, 2H), 7.22 (t, 1H), 7.18 (s, 1H), 7.01 (m, 3H), 6.92 (m, 2H), 6.80 (s, 1H), 6.78 (d, 1H), 4.63 (s, 2H), 4.15 (s, 2H), 3.93 (s, 3H), 2.41 (s, 3H).

Example 13

[0106]




2-(2-([1-(4-methoxybenzoyl)-2-methyl-5-(trifluoromethoxy)-1H-indol-3-yl]methyl)phenoxy)propanoic acid

^1H NMR (400 MHz, CDCl_3): δ 7.74 (d, 2H), 7.25 (s, 1H), 7.19 (dt, 1H), 7.05 (d, 1H), 7.03 (d, 1H), 7.00 (d, 2H), 6.90 (m, 2H), 6.82 (d, 1H), 4.93 (q, 1H), 4.12 (q, 2H), 3.93 (s, 3H), 2.41 (s, 3H), 1.72 (d, 3H).

Example 14

[0107]




2-(2-([1-(2-naphthoyl)-2-methyl-5-(trifluoromethoxy)-1H-indol-3-yl]methyl)phenoxy)propanoic acid

^1H NMR (500 MHz, $\text{DMSO}-d_6$): δ 12.96 (s, 1H), 8.35 (s, 1H), 8.07 (m, 3H), 7.76 (dd, 1H), 7.70 (t, 1H), 7.63 (t, 1H), 7.46 (s, 1H), 7.13 (m, 2H), 7.09 (d, 1H), 6.99 (s, 1H), 6.85 (t, 2H), 4.91 (q, 1H), 4.06 (s, 2H), 2.31 (s, 3H), 1.51 (d, 3H).

Example 15

[0108]




(2R)-2-(2-([1-(4-methoxybenzoyl)-2-methyl-5-(trifluoromethoxy)-1H-indol-3-yl]methyl)phenoxy)propanoic acid

^1H NMR (500 MHz, CDCl_3): δ 7.74 (d, 2H), 7.25 (s, 1H), 7.19 (dt, 1H), 7.05 (d, 1H), 7.03 (d, 1H), 7.00 (d, 2H), 6.90 (m, 2H), 6.82 (d, 1H), 4.93 (q, 1H), 4.12 (q, 2H), 3.93 (s, 3H), 2.41 (s, 3H), 1.72 (d, 3H).

Example 16

[0109]




(2S)-2-(2-([1-(4-methoxybenzoyl)-2-methyl-5-(trifluoromethoxy)-1H-indol-3-yl]methyl)phenoxy)propanoic acid

^1H NMR (500 MHz, CDCl_3): δ 7.74 (d, 2H), 7.25 (s, 1H), 7.19 (dt, 1H), 7.05 (d, 1H), 7.03 (d, 1H), 7.00 (d, 2H), 6.90 (m, 2H), 6.82 (d, 1H), 4.93 (q, 1H), 4.12 (q, 2H), 3.93 (s, 3H), 2.41 (s, 3H), 1.72 (d, 3H).

Example 17

[0110]




EP 1 305 285 B1

2-(3-([1-(4-Methoxybenzoyl)-2-methyl-5-(trifluoromethoxy)-1*H*-indol-3-yl]methyl}phenoxy)propanoic acid

¹HNMR (400MHz, CDCl₃): 7.76 (d, 2H), 7.23 (t, 1H), 7.18 (s, 1H), 7.02 (d, 1H), 7.01 (d, 2H), 6.92 (d, 1H), 6.91 (d, 1H), 6.76 (s, 1H), 6.75 (d, 1H), 4.76 (q, 1H), 4.05 (s, 2H), 3.94 (s, 3H), 2.41 (s, 3H), 1.63 (d, 3H).

Example 18

[0111]




2-(3-([1-(2-Naphthoyl)-2-methyl-5-(trifluoromethoxy)-1*H*-indol-3-yl]methyl}phenoxy)propanoic acid

¹HNMR (400MHz, CDCl₃): δ 8.30 (s, 1H), 7.97 (m, 3H), 7.80 (d, 1H), 7.68 (t, 1H), 7.61 (t, 1H), 7.24 (t, 1H), 7.20 (s, 1H), 7.03 (d, 1H), 6.92 (d, 1H), 6.86 (d, 1H), 6.80 (s, 1H), 6.75 (d, 1H), 4.78 (q, 1H), 4.06 (s, 2H), 2.41 (s, 3H), 1.65 (d, 3H).

Example 19

[0112]




2-(3-([1-(4-Chlorobenzoyl)-2-methyl-5-(trifluoromethoxy)-1*H*-indol-3-yl]methyl}phenoxy)propanoic acid

¹H NMR (400 MHz, CDCl₃): δ 6.69 (d, 2H), 7.50 (d, 2H), 7.24 (s, 1H), 7.15 (t, 1H), 7.03 (m, 2H), 6.90 (m, 2H), 6.79 (d, 1H), 4.91 (q, 1H), 4.07 (dd, 2H), 2.38 (s, 3H), 1.72 (d, 3H); ES-MS (M+1) 532, 534.

Example 20

[0113]




EP 1 305 285 B1

2-(3-([1-(2,4-Dichlorobenzoyl)-2-methyl-5-(trifluoromethoxy)-1*H*-indol-3-yl]methyl)phenoxy)propanoic acid

¹H NMR (400 MHz, CDCl₃): δ 7.49 (m, 4H), 7.24 (s, 1H), 7.18 (t, 1H), 6.99 (m, 2H), 6.87 (t, 1H), 6.78 (d, 1H), 4.93 (q, 1H), 4.08 (dd, 2H), 2.21 (s, 3H), 1.71 (d, 3H); ES-MS (M+1) 566, 568, 570.

Example 21

[0114]




(2R)-2-(3-([1-(4-Methoxybenzoyl)-2-methyl-5-(trifluoromethoxy)-1*H*-indol-3-yl]methyl)phenoxy)propanoic acid

¹H NMR (400MHz, CDCl₃): 7.76 (d, 2H), 7.23 (t, 1H), 7.18 (s, 1H), 7.02 (d, 1H), 7.01 (d, 2H), 6.92 (d, 1H), 6.91 (d, 1H), 6.76 (s, 1H), 6.75 (d, 1H), 4.76 (q, 1H), 4.05 (s, 2H), 3.94 (s, 3H), 2.41 (s, 3H), 1.63 (d, 3H).

Example 22

[0115]




(2R)-2-(3-([1-(2-Naphthoyl)-2-methyl-5-(trifluoromethoxy)-1*H*-indol-3-yl]methyl)phenoxy)propanoic acid

¹H NMR (400MHz, CDCl₃): d 8.30 (s, 1H), 7.97 (m, 3H); 7.80 (d, 1H), 7.68 (t, 1H), 7.61 (t, 1H), 7.24 (t, 1H), 7.20 (s, 1H), 7.03 (d, 1H), 6.92 (d, 1H), 6.86 (d, 1H), 6.80 (s, 1H), 6.75 (d, 1H), 4.78 (q, 1H), 4.06 (s, 2H), 2.41 (s, 3H), 1.65 (d, 3H).

Example 23

[0116]




EP 1 305 285 B1

(2S)-2-(3-([1-(2-Naphthoyl)-2-methyl-5-(trifluoromethoxy)-1*H*-indol-3-yl]methyl)phenoxy)propanoic acid

¹H NMR (400MHz, CDCl₃): δ 8.30 (s, 1H), 7.97 (m, 3H), 7.80 (d, 1H), 7.68 (t, 1H), 7.61 (t, 1H), 7.24 (t, 1H), 7.20 (s, 1H), 7.03 (d, 1H), 6.92 (d, 1H), 6.86 (d, 1H), 6.80 (s, 1H), 6.75 (d, 1H), 4.78 (q, 1H), 4.06 (s, 2H), 2.41 (s, 3H), 1.65 (d, 3H).

Example 24

[0117]




2-(2-([1-(4-Methoxybenzoyl)-2-methyl-5-(trifluoromethoxy)-1*H*-indol-3-yl]methyl)phenoxy)-2-methylpropanoic acid

¹H NMR (400 MHz, CDCl₃): δ 7.74 (d, 2H), 7.15 (m, 2H), 7.01 (m, 4H), 6.85 (m, 3H), 4.08 (s, 2H), 3.93 (s, 3H), 2.39 (s, 3H), 1.68 (s, 6H); ES-MS (M+1) 542.

Example 25

[0118]




2-(3-([1-(4-Methoxybenzoyl)-2-methyl-5-(trifluoromethoxy)-1*H*-indol-3-yl]methyl)phenoxy)-2-methylpropanoic acid

¹H NMR (400 MHz, CDCl₃): δ 7.76 (d, 2H), 7.19 (m, 2H), 7.00 (m, 4H), 6.82 (d, 2H), 6.78 (m, 2H), 4.22 (s, 2H), 3.91 (s, 3H), 2.21 (s, 3H), 1.57 (s, 6H); ES-MS (M+1) 542.

Example 26


[0119]



2-(2-([1-(2-Naphthoyl)-2-methyl-5-(trifluoromethoxy)-1*H*-indol-3-yl]methyl)phenoxy)-2-methylpropanoic acid
¹H NMR (400 MHz, CDCl₃): δ 8.29 (s, 1H), 9.98 (m, 3H), 7.79 (d, 1H), 6.15 (m, 2H), 7.20 (s, 1H), 7.17 (t, 1H), 7.05 (m, 2H), 6.91 (t, 1H), 6.84 (m, 2H), 4.09 (s, 2H), 2.39 (s, 3H), 1.70 (s, 6H); ES-MS (M+1) 562.

Example 27


[0120]



2-(3-([1-(2-Naphthoyl)-2-methyl-5-(trifluoromethoxy)-1*H*-indol-3-yl]methyl)phenoxy)-2-methylpropanoic acid
¹H NMR (400 MHz, CDCl₃): δ 8.31 (s, 1H), 7.95 (m, 3H), 7.80 (d, 1H), 7.65 (m, 2H), 7.20 (m, 2H), 7.03 (d, 1H), 6.97 (d, 1H), 6.83 (d, 1H), 6.79 (m, 2H), 4.07 (s, 2H), 2.21 (s, 3H), 1.58 (s, 6H); ES-MS (M+1) 363.

Example 28

[0121]




(2*R*)-2-(3-{2-[1-(4-Methoxybenzoyl)-2-methyl-5-(trifluoromethoxy)-1*H*-indol-3-yl]ethyl}phenoxy)propionic acid
¹H NMR (400 MHz, CDCl₃): δ 7.75-7.60 (br m, 2H), 7.33-7.29 (m, 2H), 7.21 (t, 1H), 7.02 (d, 1H), 6.97-6.95 (m, 2H), 6.88 (d, 1H), 6.80 (d, 1H), 6.01 (s, 1H), 4.51 (br m, 1H), 3.92 (s, 3H), 3.11-3.02 (m, 2H), 2.81-2.75 (m, 2H), 1.65 (s,

3H), 1.56 (d, 3H); ES-MS (M+1) 542.

Example 29

[0122]




(2S)-2-{3-[1-(4-methoxybenzoyl)-2-methyl-5-(trifluoromethoxy)-1H-indol-3-yl]phenoxy}propionic acid

¹H NMR (500 MHz, CDCl₃): δ 7.81 (d, 2H), 7.46 (t, 1H), 7.39 (br s, 1H), 7.16 (d, 1H), 7.11 (d, 1H), 7.04 (m, 3H), 6.97 (m, 2H), 4.91 (q, 1H), 3.95 (s, 3H), 2.44 (s, 3H), 1.74 (d, 3H).

Example 30

[0123]




(2S)-2-{3-[1-(4-Methoxybenzoyl)-2-methyl-5-(trifluoromethoxy)-1H-indol-3-yl]cyclopropyl}phenoxy}propanoic acid

¹H NMR (500 MHz, CDCl₃): δ 7.77 (d, 2H), 7.46, (s, 1H), 7.18 (t, 1H), 7.02 (m, 3H), 6.92 (d, 1H), 6.75 (d, 2H), 6.69-6.66 (m, 2H), 4.70 (q, 1H), 3.94 (s, 3H), 2.47 (s, 3H), 1.63 (d, 3H), 1.50 (m, 2H), 1.33 (m, 2H); ES-MS (M+1) 554.

Example 31

[0124]




2-{3-[1-(4-Methoxybenzoyl)-2-methyl-5-(trifluoromethoxy)-1H-indol-3-yl]phenoxy}-2-methylpropanoic acid

¹H NMR (500 MHz, CDCl₃): δ 7.80 (d, 2H), 7.43 (t, 1H), 7.38 (br s, 1H), 7.20 (d, 1H), 7.10 (d, 1H), 7.02(m, 5H), 3.94 (s, 3H), 2.43 (s, 3H), 1.69 (s, 6H).

Claims

1. A compound of formula I:



wherein:

R¹ is methyl, optionally substituted with 1-3 F;

R², R³ and R⁴ are each independently selected from the group consisting of H, halogen, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, aryl, OC₁-C₆ alkyl, OC₂-C₆ alkenyl, OC₂-C₆ alkynyl, O-aryl, OH, SC₁-C₆ alkyl, SC₂-C₆ alkenyl, SC₂-C₆ alkynyl, SO₂C₁-C₆ alkyl, SO₂C₂-C₆ alkenyl, SO₂C₂-C₆ alkynyl, OCON(R₅)₂, OCO (C₁-C₆-alkyl) and CN, wherein all instances of alkyl, alkenyl and alkynyl are optionally linear or branched and all instances of alkyl, alkenyl, alkynyl, cycloalkyl and aryl are optionally substituted with 1-5 substituents independently selected from the group consisting of halogen, aryl, O-aryl and OMe;

R⁵ and R⁶ are, at each occurrence, independently selected from the group consisting of H, F, OH and C₁-C₅ alkyl, and R⁵ and R⁶ groups that are on the same carbon atom optionally may be joined to form a C₃-C₆ cycloalkyl group;

R⁷ and R⁸ are each independently selected from the group consisting of H, F, and C₁-₅ alkyl, or R⁷ and R⁸ optionally may be joined to form a C₃-C₆ cycloalkyl group;

R⁹ is selected from the group consisting of H and C₁-C₅ alkyl, said alkyl being optionally linear or branched;

Ar¹ is phenyl, 1-naphthyl, 2-naphthyl, pyridyl or quinolyl wherein Ar¹ is substituted with 1-3 groups independently selected from R⁴;

X is selected from the group consisting of C=O, S(O)₂, CH₂, CH(CH₃), C(CH₃)₂, CF₂, and cyclopropylidene;

Y is O or S; and

n is 0-5;

and pharmaceutically acceptable salts and prodrugs thereof.

2. The compound of Claim 1 wherein:

R¹ is CH₃;

R², R³ and R⁴ are each independently selected from the group consisting of H, halogen, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, aryl, OC₁-C₆ alkyl, OC₂-C₆ alkenyl, OC₂-C₆ alkynyl, O-aryl, OH, SC₁-C₆ alkyl, SC₂-C₆ alkenyl, SC₂-C₆ alkynyl, OCON(R₅)₂, OCO(C₁-C₆-alkyl) and CN, wherein all instances of alkyl, alkenyl and alkynyl are optionally linear or branched and all instances of alkyl, alkenyl, alkynyl, cycloalkyl and aryl are optionally substituted with 1-5 substituents independently selected from the group consisting of halogen, aryl, O-aryl and OMe; and

X is selected from the group consisting of C=O, CH₂, CH(CH₃), C(CH₃)₂, CF₂, and cyclopropylidene.

3. The compound of Claim 1 wherein R², R³, and R⁴ are each independently selected from the group consisting of H, OCH₃, OCF₃, F, Cl and CH₃, wherein CH₃ is optionally substituted with 1-3 groups independently selected from F, Cl, and OCH₃.

4. The compound of Claim 1 wherein R^2 , R^3 , and R^4 are each independently selected from the group consisting of H, OCH_3 , OCF_3 , and Cl.

5. The compound of Claim 1 wherein R^5 and R^6 are H.

6. The compound of Claim 1 wherein R^7 and R^8 are each independently CH_3 or H.

7. The compound of Claim 1 wherein R^9 is H.

8. The compound of Claim 1 wherein X is C=O.

9. The compound of Claim 1 wherein Y is O.

10. The compound of Claim 1 wherein n is 0, 1 or 2.

11. The compound of Claim 1 wherein n is 1.

12. The compound of Claim 1 wherein Ar^1 is phenyl, 1-naphthyl or 2-naphthyl.

13. The compound of Claim 1 wherein Ar^1 is phenyl or 2-naphthyl, Ar^1 being substituted with 1-3 groups independently selected from R^4 .

14. The compound of Claim 1 wherein aryl is phenyl.

15. The compound of Claim 1 wherein:

R^1 is CH_3 ;

R^2 is selected from the group consisting of H, OCH_3 , and OCF_3 ;

R^3 , R^5 , R^6 , and R^9 are H;

R^4 is selected from the group consisting of H, Cl, and OCH_3 ;

R^7 and R^8 are each independently selected from the group consisting of H and CH_3 ;

X is C=O;

Y is O;


and n is 1.

16. The compound of Claim 1 wherein R^9 is defined as a group that is easily removed under physiological conditions during or after administration to a mammalian patient to yield a carboxylic acid in which R^9 is H, or the carboxylate anion thereof, or a pharmaceutically acceptable salt thereof, and R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , Ar^1 , X, Y, and n are as defined in Claim 1.


17. The compound of Claim 16, wherein OR^9 of the group $-CO_2R^9$ is selected from the group consisting of $-OR^{10}$, $-OCH_2OR^{10}$, $-OCH(CH_3)OR^{10}$, $-OCH_2OC(O)R^{10}$, $-OCH(CH_3)OC(O)R^{10}$, $-OCH_2OC(O)OR^{10}$, $-OCH(CH_3)OC(O)OR^{10}$, $-NR^{11}R^{11}$, and $-ONR^{11}R^{11}$,

wherein each R^{10} is independently selected from C_1 - C_6 alkyl optionally substituted with one or two groups selected from $-CO_2H$, $-CONH_2$, NH_2 , $-OH$, $-OAc$, $NHAc$ and phenyl; and wherein each R^{11} is independently selected from H and R^{10} .

18. A compound which is selected from the group consisting of:




5




10

15




20

25




30

35








40





45



50

55





5

10

15

20

25

30






35






40

45

50

55





5

10

15

20

25

30





35



40

45

50

55





19. A compound which is selected from the group consisting of:

- (2S)-2-(3-([1-(4-Methoxybenzoyl)-2-methyl-5-(trifluoromethoxy)-1H-indol-3-yl]methyl)phenoxy)propanoic acid;
 2-(2-([1-(4-Chlorobenzoyl)-2-methyl-5-(methoxy)-1H-indol-3-yl]methyl)phenoxy)acetic acid;
 2-(3-([1-(4-Chlorobenzoyl)-2-Methyl-5-(methoxy)-1H-indol-3-yl]methyl)phenoxy)acetic acid;
 2-(4-([1-(4-Chlorobenzoyl)-2-methyl-5-(methoxy)-1H-indol-3-yl]methyl)phenoxy)acetic acid;
 2-(2-([1-(4-Chlorobenzoyl)-2-methyl-5-(methoxy)-1H-indol-3-yl]methyl)phenoxy)propanoic acid;
 2-(3-([1-(4-Chlorobenzoyl)-2-methyl-5-(methoxy)-1H-indol-3-yl]methyl)phenoxy)propanoic acid;
 2-(4-([1-(4-Chlorobenzoyl)-2-methyl-5-(methoxy)-1H-indol-3-yl]methyl)phenoxy)propanoic acid;
 2-(2-([1-(4-Chlorobenzoyl)-2-methyl-5-(methoxy)-1H-indol-3-yl]methyl)phenoxy)-2-methylpropanoic acid;
 2-(3-([1-(4-Chlorobenzoyl)-2-methyl-5-(methoxy)-1H-indol-3-yl]methyl)phenoxy)-2-methylpropanoic acid;
 2-(4-([1-(4-Chlorobenzoyl)-2-methyl-5-(methoxy)-1H-indol-3-yl]methyl)phenoxy)-2-methylpropanoic acid;
 2-(2-([1-(4-Methoxybenzoyl)-2-methyl-5-(trifluoromethoxy)-1H-indol-3-yl]methyl)phenoxy)acetic acid;
 2-(3-([1-(4-Methoxybenzoyl)-2-Methyl-5-(trifluoromethoxy)-1H-indol-3-yl]methyl)phenoxy)acetic acid;
 2-(2-([1-(4-methoxybenzoyl)-2-Methyl-5-(trifluoromethoxy)-1H-indol-3-yl]methyl)phenoxy)propanoic acid;
 2-(2-([1-(2-naphthoyl)-2-Methyl-5-(trifluoromethoxy)-1H-indol-3-yl]methyl)phenoxy)propanoic acid;
 (2R)-2-(2-([1-(4-methoxybenzoyl)-2-Methyl-5-(trifluoromethoxy)-1H-indol-3-yl]methyl)phenoxy)propanoic acid;
 (2S)-2-(2-([1-(4-Methoxybenzoyl)-2-methyl-5-(trifluoromethoxy)-1H-indol-3-yl]methyl)phenoxy)propanoic acid;
 2-(3-([1-(4-Methoxybenzoyl)-2-methyl-5-(trifluoromethoxy)-1H-indol-3-yl]methyl)phenoxy)propanoic acid;
 2-(3-([1-(2-Naphthoyl)-2-methyl-5-(trifluoromethoxy)-1H-indol-3-yl]methyl)phenoxy)propanoic acid;
 2-(3-([1-(4-Chlorobenzoyl)-2-methyl-5-(trifluoromethoxy)-1H-indol-3-yl]methyl)phenoxy)propanoic acid;
 2-(3-([1-(2,4-Dichlorobenzoyl)-2-methyl-5-(trifluoromethoxy)-1H-indol-3-yl]methyl)phenoxy)propanoic acid;
 (2R)-2-(3-([1-(4-Methoxybenzoyl)-2-methyl-5-(trifluoromethoxy)-1H-indol-3-yl]methyl)phenoxy)propanoic acid;
 (2R)-2-(3-([1-(2-Naphthoyl)-2-methyl-5-(trifluoromethoxy)-1H-indol-3-yl]methyl)phenoxy)propanoic acid;
 (2S)-2-(3-([1-(2-Naphthoyl)-2-methyl-5-(trifluoromethoxy)-1H-indol-3-yl]methyl)phenoxy)propanoic acid;
 2-(2-([1-(4-Methoxybenzoyl)-2-methyl-5-(trifluoromethoxy)-1H-indol-3-yl]methyl)phenoxy)-2-methylpropanoic acid;
 2-(3-([1-(4-Methoxybenzoyl)-2-methyl-5-(trifluoromethoxy)-1H-indol-3-yl]methyl)phenoxy)-2-methylpropanoic acid;
 2-(2-([1-(2-Naphthoyl)-2-methyl-5-(trifluoromethoxy)-1H-indol-3-yl]methyl)phenoxy)-2-methylpropanoic acid;
 2-(3-([1-(2-Naphthoyl)-2-methyl-5-(trifluoromethoxy)-1H-indol-3-yl]methyl)phenoxy)-2-methylpropanoic acid;
 (2R)-2-(3-([1-(4-Methoxybenzoyl)-2-methyl-5-(trifluoromethoxy)-1H-indol-3-yl]methyl)phenoxy)propionic acid;
 (2S)-2-(3-([1-(4-methoxybenzoyl)-2-methyl-5-(trifluoromethoxy)-1H-indol-3-yl]phenoxy)propionic acid;
 (2S)-2-(3-([1-(4-Methoxybenzoyl)-2-methyl-5-(trifluoromethoxy)-1H-indol-3-yl]cyclopropyl)phenoxy)propanoic acid; and

2-{3-[1-(4-Methoxybenzoyl)-2-methyl-5-(trifluoromethoxy)-1*H*-indol-3-yl]phenoxy}-2-methylpropanoic acid.

20. A pharmaceutical composition comprising a compound of any one of Claims 1 to 19 or a pharmaceutically acceptable salt thereof and a pharmaceutically acceptable carrier.

21. A compound of any one of Claims 1 to 19 or a pharmaceutically acceptable salt thereof for use in a method of treatment of the human body by therapy.

22. Use of a compound of any one of claims 1 to 19 or a pharmaceutically acceptable salt thereof for the manufacture of a medicament for treating, controlling or preventing one or more diseases, disorders, or conditions selected from (1) non-insulin dependent diabetes mellitus (NIDDM), (2) hyperglycemia, (3) low glucose tolerance, (4) insulin resistance, (5) obesity, (6) lipid disorders, (7) dyslipidemia, (8) hyperlipidemia, (9) hypertriglyceridemia, (10) hypercholesterolemia, (11) low HDL levels, (12) high LDL levels, (13) atherosclerosis and its sequelae, (14) vascular restenosis, (15) irritable bowel syndrome, (16) inflammatory bowel disease, including Crohn's disease and ulcerative colitis, (17) other inflammatory conditions, (18) pancreatitis, (19) abdominal obesity, (20) neurodegenerative disease, (21) retinopathy, (22) neoplastic conditions, (23) adipose cell tumors, (24) adipose cell carcinomas, such as liposarcoma, (25) prostate cancer and other cancers, including gastric, breast, bladder and colon cancers, (26) angiogenesis, (27) Alzheimer's disease, (28) psoriasis, (29) high blood pressure, (30) Syndrome X, (31) ovarian hyperandrogenism (polycystic ovarian syndrome), and other disorders where insulin resistance is a component.

23. A pharmaceutical composition comprising (1) a compound of any one of Claims 1 to 19 or a pharmaceutically acceptable salt thereof, (2) one or more compounds selected from the group consisting of:

(a) insulin sensitizers including (i) PPAR γ agonists such as the glitazones (e.g. troglitazone, pioglitazone, englitazone, MCC-555, rosiglitazone, and the like), and compounds disclosed in WO97/27857, 97/28115, 97/28137 and 97/27847; (ii) biguanides such as metformin and phenformin; (iii) protein tyrosine phosphatase-1B (PTP-1B) inhibitors, and (iv) dipeptidyl peptidase IV (DP-IV) inhibitors;

(b) insulin or insulin mimetics;

(c) sulfonylureas such as tolbutamide and glipizide, or related materials;

(d) α -glucosidase inhibitors (such as acarbose);

(e) cholesterol lowering agents such as (i) HMG-CoA reductase inhibitors (lovastatin, simvastatin, pravastatin, fluvastatin, atorvastatin, rivastatin, itavastatin, ZD-4522 and other statins), (ii) sequestrants (cholestyramine, colestipol, and dialkylaminoalkyl derivatives of a cross-linked dextran), (iii) nicotinic alcohol, nicotinic acid or a salt thereof, (iv) PPAR α agonists such as fenofibric acid derivatives (gemfibrozil, clofibrate, fenofibrate and benzafibrate), (v) PPAR α/γ dual agonists, such as KRP-297, (vi) inhibitors of cholesterol absorption, such as for example beta-sitosterol, (vii) acyl CoA:cholesterol acyltransferase inhibitors, such as for example avasimibe, and (viii) anti-oxidants, such as probucol;

(f) PPAR δ agonists such as those disclosed in WO97/28149;

(g) antiobesity compounds such as fenfluramine, dexfenfluramine, phentiramine, sulbitramine, orlistat, neuropeptide Y5 inhibitors, and β_3 adrenergic receptor agonists;


(h) an ileal bile acid transporter inhibitor; and

(i) agents intended for use in inflammatory conditions such as aspirin, non-steroidal anti-inflammatory drugs, glucocorticoids, azulfidine, and cyclooxygenase 2 selective inhibitors; and (3) a pharmaceutically acceptable carrier.

24. A pharmaceutical composition according to Claim 23 where component (2) is an HMG-CoA reductase inhibitor.

Patentansprüche

1. Eine Verbindung der Formel I:



wobei:

R¹ Methyl ist, gegebenenfalls substituiert mit 1-3 F,

R², R³ und R⁴ jeweils unabhängig ausgewählt sind aus der Gruppe, bestehend aus H, Halogen, C₁-C₆-Alkyl, C₂-C₆-Alkenyl, C₂-C₆-Alkinyl, C₃-C₈-Cycloalkyl, Aryl, OC₁-C₆-Alkyl, OC₂-C₆-Alkenyl, OC₂-C₆-Alkinyl, O-Aryl, OH, SC₁-C₆-Alkyl, SC₂-C₆-Alkenyl, SC₂-C₆-Alkinyl, SO₂C₁-C₆-Alkyl, SO₂C₂-C₆-Alkenyl, SO₂C₂-C₆-Alkinyl, OCON(R₅)₂, OCO(C₁-C₆-Alkyl) und CN, wobei alle Fälle von Alkyl, Alkenyl und Alkinyl gegebenenfalls linear oder verzweigt sind und alle Fälle von Alkyl, Alkenyl, Alkinyl, Cycloalkyl und Aryl gegebenenfalls substituiert sind mit 1-5 Substituenten, unabhängig ausgewählt aus der Gruppe, bestehend aus Halogen, Aryl, O-Aryl und OMe,

R⁵ und R⁶ bei jedem Auftreten unabhängig ausgewählt sind aus der Gruppe, bestehend aus H, F, OH und C₁-C₅-Alkyl, und R⁵- und R⁶-Gruppen, die sich am selben Kohlenstoffatom befinden, gegebenenfalls verbunden sein können, um eine C₃-C₆-Cycloalkylgruppe zu bilden,

R⁷ und R⁸ jeweils unabhängig ausgewählt sind aus der Gruppe, bestehend aus H, F und C₁₋₅-Alkyl, oder R⁷ und R⁸ gegebenenfalls verbunden sein können, um eine C₃-C₆-Cycloalkylgruppe zu bilden,

R⁹ ausgewählt ist aus der Gruppe, bestehend aus H und C₁-C₅-Alkyl, wobei das Alkyl gegebenenfalls linear oder verzweigt ist,

Ar¹ Phenyl, 1-Naphthyl, 2-Naphthyl, Pyridyl oder Chinolyl ist, wobei Ar¹ substituiert ist mit 1-3 Gruppen, unabhängig ausgewählt aus R⁴,

X ausgewählt ist aus der Gruppe, bestehend aus C=O, S(O)₂, CH₂, CH(CH₃), C(CH₃)₂, CF₂ und Cyclopropylen,

Y O oder S ist und

n 0-5 ist,

und pharmazeutisch annehmbare Salze und Prodrugs davon.

2. Die Verbindung nach Anspruch 1, wobei:

R¹ CH₃ ist,


R², R³ und R⁴ jeweils unabhängig ausgewählt sind aus der Gruppe, bestehend aus H, Halogen, C₁-C₆-Alkyl, C₂-C₆-Alkenyl, C₂-C₆-Alkinyl, C₃-C₈-Cycloalkyl, Aryl, OC₁-C₆-Alkyl, OC₂-C₆-Alkenyl, OC₂-C₆-Alkinyl, O-Aryl, OH, SC₁-C₆-Alkyl, SC₂-C₆-Alkenyl, SC₂-C₆-Alkinyl, OCON(R₅)₂, OCO(C₁-C₆-Alkyl) und CN, wobei alle Fälle von Alkyl, Alkenyl und Alkinyl gegebenenfalls linear oder verzweigt sind und alle Fälle von Alkyl, Alkenyl, Alkinyl, Cycloalkyl und Aryl gegebenenfalls substituiert sind mit 1-5 Substituenten, unabhängig ausgewählt aus der Gruppe, bestehend aus Halogen, Aryl, O-Aryl und OMe, und

X ausgewählt ist aus der Gruppe, bestehend aus C=O, CH₂, CH(CH₃), C(CH₃)₂, CF₂ und Cyclopropylen.


3. Die Verbindung nach Anspruch 1, wobei R², R³ und R⁴ jeweils unabhängig ausgewählt sind aus der Gruppe, bestehend aus H, OCH₃, OCF₃, F, Cl und CH₃, wobei CH₃ gegebenenfalls substituiert ist mit 1-3 Gruppen, unabhängig ausgewählt aus F, Cl und OCH₃.

4. Die Verbindung nach Anspruch 1, wobei R², R³ und R⁴ jeweils unabhängig ausgewählt sind aus der Gruppe, bestehend aus H, OCH₃, OCF₃ und Cl.

5. Die Verbindung nach Anspruch 1, wobei R^5 und R^6 H sind.
6. Die Verbindung nach Anspruch 1, wobei R^7 und R^8 jeweils unabhängig CH_3 oder H sind.
7. Die Verbindung nach Anspruch 1, wobei R^9 H ist.
8. Die Verbindung nach Anspruch 1, wobei X C=O ist.
9. Die Verbindung nach Anspruch 1, wobei Y O ist.
10. Die Verbindung nach Anspruch 1, wobei n 0, 1 oder 2 ist.
11. Die Verbindung nach Anspruch 1, wobei n 1 ist.
12. Die Verbindung nach Anspruch 1, wobei Ar^1 Phenyl, 1-Naphthyl oder 2-Naphthyl ist.
13. Die Verbindung nach Anspruch 1, wobei Ar^1 Phenyl oder 2-Naphthyl ist und Ar^1 substituiert ist mit 1-3 Gruppen, unabhängig ausgewählt aus R^4 .
14. Die Verbindung nach Anspruch 1, wobei Aryl Phenyl ist.
15. Die Verbindung nach Anspruch 1, wobei:
 R^1 CH_3 ist,
 R^2 ausgewählt ist aus der Gruppe, bestehend aus H, OCH_3 und OCF_3 ,
 R^3 , R^5 , R^6 und R^9 H sind,
 R^4 ausgewählt ist aus der Gruppe, bestehend aus H, Cl und OCH_3 ,
 R^7 und R^8 jeweils unabhängig ausgewählt sind aus der Gruppe, bestehend aus H und CH_3 ,
 X C=O ist,
Y O ist
und n 1 ist.
16. Die Verbindung nach Anspruch 1, wobei R^9 als eine Gruppe definiert ist, die leicht unter physiologischen Bedingungen während oder nach der Verabreichung an einen Säuger-Patienten entfernt wird, um eine Carbonsäure zu ergeben, bei der R^9 H ist, oder das Carboxylatanion davon oder ein pharmazeutisch annehmbares Salz davon, und R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , Ar^1 , X, Y und n wie in Anspruch 1 definiert sind.
17. Die Verbindung nach Anspruch 16, wobei OR^9 der Gruppe $-CO_2R^9$ ausgewählt ist aus der Gruppe, bestehend aus $-OR^{10}$, $-OCH_2OR^{10}$, $-OCH(CH_3)OR^{10}$, $-OCH_2OC(O)R^{10}$, $-OCH(CH_3)OC(O)R^{10}$, $-OCH_2OC(O)OR^{10}$, $-OCH(CH_3)OC(O)OR^{10}$, $-NR^{11}R^{11}$ und $-ONR^{11}R^{11}$, wobei jedes R^{10} unabhängig ausgewählt ist aus C_1 - C_6 -Alkyl, gegebenenfalls substituiert mit einer oder zwei Gruppen, ausgewählt aus $-CO_2H$, $-CONH_2$, NH_2 , $-OH$, $-OAc$, $NHAc$ und Phenyl, und wobei jedes R^{11} unabhängig ausgewählt ist aus H und R^{10} .
18. Eine Verbindung, die ausgewählt ist aus der Gruppe, bestehend aus:




5




10

15




20

25




30

35








40






45




50

55






5




10




15

20




25

30




35

40












45

50



55





19. Eine Verbindung, die ausgewählt ist aus der Gruppe, bestehend aus:

(2S)-2-(3-([1-(4-Methoxybenzoyl)-2-methyl-5-(trifluormethoxy)-1H-indol-3-yl]methyl)phenoxy)propansäure,
 2-(2-([1-(4-Chlorbenzoyl)-2-methyl-5-(methoxy)-1H-indol-3-yl]methyl)phenoxy)essigsäure,
 2-(3-([1-(4-Chlorbenzoyl)-2-methyl-5-(methoxy)-1H-indol-3-yl]methyl)phenoxy)essigsäure,
 2-(4-([1-(4-Chlorbenzoyl)-2-methyl-5-(methoxy)-1H-indol-3-yl]methyl)phenoxy)essigsäure,
 2-(2-([1-(4-Chlorbenzoyl)-2-methyl-5-(methoxy)-1H-indol-3-yl]methyl)phenoxy)propansäure,
 2-(3-([1-(4-Chlorbenzoyl)-2-methyl-5-(methoxy)-1H-indol-3-yl]methyl)phenoxy)propansäure,
 2-(4-([1-(4-Chlorbenzoyl)-2-methyl-5-(methoxy)-1H-indol-3-yl]methyl)phenoxy)propansäure,
 2-(2-([1-(4-Chlorbenzoyl)-2-methyl-5-(methoxy)-1H-indol-3-yl]methyl)phenoxy)-2-methylpropansäure,
 2-(3-([1-(4-Chlorbenzoyl)-2-methyl-5-(methoxy)-1H-indol-3-yl]methyl)phenoxy)-2-methylpropansäure,
 2-(4-([1-(4-Chlorbenzoyl)-2-methyl-5-(methoxy)-1H-indol-3-yl]methyl)phenoxy)-2-methylpropansäure,
 2-(2-([1-(4-Methoxybenzoyl)-2-methyl-5-(trifluormethoxy)-1H-indol-3-yl]methyl)phenoxy)essigsäure,
 2-(3-([1-(4-Methoxybenzoyl)-2-methyl-5-(trifluormethoxy)-1H-indol-3-yl]methyl)phenoxy)essigsäure,
 2-(2-([1-(4-Methoxybenzoyl)-2-methyl-5-(trifluormethoxy)-1H-indol-3-yl]methyl)phenoxy)propansäure,

2-(2-[[1-(2-Naphthoyl)-2-methyl-5-(trifluormethoxy)-1*H*-indol-3-yl]methyl]phenoxy)propansäure,
 (2*R*)-2-(2-[[1-(4-Methoxybenzoyl)-2-methyl-5-(trifluormethoxy)-1*H*-indol-3-yl]methyl]phenoxy)propansäure,
 (2*S*)-2-(2-[[1-(4-Methoxybenzoyl)-2-methyl-5-(trifluormethoxy)-1*H*-indol-3-yl]methyl]phenoxy)propansäure,
 2-(3-[[1-(4-Methoxybenzoyl)-2-methyl-5-(trifluormethoxy)-1*H*-indol-3-yl]methyl]phenoxy)propansäure,
 2-(3-[[1-(2-Naphthoyl)-2-methyl-5-(trifluormethoxy)-1*H*-indol-3-yl]methyl]phenoxy)propansäure,
 2-(3-[[1-(4-Chlorbenzoyl)-2-methyl-5-(trifluormethoxy)-1*H*-indol-3-yl]methyl]phenoxy)propansäure,
 2-(3-[[1-(2,4-Dichlorbenzoyl)-2-methyl-5-(trifluormethoxy)-1*H*-indol-3-yl]methyl]phenoxy)propansäure,
 (2*R*)-2-(3-[[1-(4-Methoxybenzoyl)-2-methyl-5-(trifluormethoxy)-1*H*-indol-3-yl]methyl]phenoxy)propansäure,
 (2*R*)-2-(3-[[1-(2-Naphthoyl)-2-methyl-5-(trifluormethoxy)-1*H*-indol-3-yl]methyl]phenoxy)propansäure,
 (2*S*)-2-(3-[[1-(2-Naphthoyl)-2-methyl-5-(trifluormethoxy)-1*H*-indol-3-yl]methyl]phenoxy)propansäure,
 2-(2-[[1-(4-Methoxybenzoyl)-2-methyl-5-(trifluormethoxy)-1*H*-indol-3-yl]methyl]phenoxy)-2-methylpropansäure,
 2-(3-[[1-(4-Methoxybenzoyl)-2-methyl-5-(trifluormethoxy)-1*H*-indol-3-yl]methyl]phenoxy)-2-methylpropansäure,
 2-(2-[[1-(2-Naphthoyl)-2-methyl-5-(trifluormethoxy)-1*H*-indol-3-yl]methyl]phenoxy)-2-methylpropansäure,
 2-(3-[[1-(2-Naphthoyl)-2-methyl-5-(trifluormethoxy)-1*H*-indol-3-yl]methyl]phenoxy)-2-methylpropansäure,
 (2*R*)-2-(3-{2-[1-(4-Methoxybenzoyl)-2-methyl-5-(trifluormethoxy)-1*H*-indol-3-yl]ethyl}phenoxy)propionsäure,
 (2*S*)-2-(3-{2-[1-(4-Methoxybenzoyl)-2-methyl-5-(trifluormethoxy)-1*H*-indol-3-yl]phenoxy})propionsäure,
 (2*S*)-2-(3-{1-(4-Methoxybenzoyl)-2-methyl-5-(trifluormethoxy)-1*H*-indol-3-yl}cyclopropyl)phenoxy)propansäure und
 2-(3-[1-(4-Methoxybenzoyl)-2-methyl-5-(trifluormethoxy)-1*H*-indol-3-yl]phenoxy)-2-methylpropansäure.

20. Eine pharmazeutische Zusammensetzung, die eine Verbindung nach irgendeinem der Ansprüche 1 bis 19 oder ein pharmazeutisch annehmbares Salz davon und einen pharmazeutisch annehmbaren Träger enthält.

21. Eine Verbindung nach irgendeinem der Ansprüche 1 bis 19 oder ein pharmazeutisch annehmbares Salz davon zur Verwendung bei einem Verfahren zur Behandlung des menschlichen Körpers durch Therapie.

22. Verwendung einer Verbindung nach irgendeinem der Ansprüche 1 bis 19 oder eines pharmazeutisch annehmbaren Salzes davon zur Herstellung eines Medikaments zur Behandlung, Kontrolle oder Prävention von ein oder mehreren Erkrankungen, Störungen oder Zuständen, ausgewählt aus (1) nichtinsulinabhängigem Diabetes mellitus (NIDDM), (2) Hyperglykämie, (3) niedriger Glukosetoleranz, (4) Insulinresistenz, (5) Fettleibigkeit, (6) Lipidstörungen, (7) Dyslipidämie, (8) Hyperlipidämie, (9) Hypertriglyceridämie, (10) Hypercholesterinämie, (11) niedrigen HDL-Spiegeln, (12) hohen LDL-Spiegeln, (13) Atherosklerose und deren Folgekrankheiten, (14) vaskulärer Restenose, (15) Colon irritabile, (16) entzündlicher Darmerkrankung, einschließlich Morbus Crohn und Colitis ulcerosa, (17) anderen Entzündungszuständen, (18) Pankreatitis, (19) abdominaler Fettleibigkeit, (20) neurodegenerativer Erkrankung, (21) Retinopathie, (22) neoplastischen Zuständen, (23) adipösen Zelltumoren, (24) adipösen Zellkarzinomen, wie z.B. Liposarkom, (25) Prostatakrebs und anderen Karzinomen, einschließlich Magen-, Brust-, Blasen- und Kolonkarzinome, (26) Angiogenese, (27) Alzheimer-Krankheit, (28) Psoriasis, (29) hohem Blutdruck, (30) Syndrom X, (31) Eierstock-Hyperandrogenismus (polyzystischem Ovarialsyndrom) und anderen Störungen, bei denen die Insulinresistenz eine Komponente ist.

23. Eine pharmazeutische Zusammensetzung, die (1) eine Verbindung nach irgendeinem der Ansprüche 1 bis 19 oder ein pharmazeutisch annehmbares Salz davon, (2) eine oder mehrere Verbindungen, ausgewählt aus der Gruppe, bestehend aus:


- (a) Insulinsensibilisatoren, einschließlich (i) PPAR γ -Agonisten, wie z.B. die Glitazone (z.B. Troglitazon, Pioglitazon, Englitazon, MCC-555, Rosiglitazon und dergleichen) und Verbindungen, die in der WO97/27857, 97/28115, 97/28137 und 97/27847 offenbart sind, (ii) Biguanide, wie z.B. Metformin und Phenformin, (iii) Protein-Tyrosin-Phosphatase-1B-(PTP-1B)-Inhibitoren und (iv) Dipeptidylpeptidase-IV-(DP-IV)-Inhibitoren,
- (b) Insulin oder Insulin-Mimetika,
- (c) Sulfonylharnstoffen, wie z.B. Tolbutamid und Glipizid, oder verwandten Materialien,
- (d) α -Glucosidaseinhibitoren (wie z.B. Acarbose),
- (e) Cholesterinsenkungsmittel, wie z.B. (i) HMG-CoA-Reduktaseinhibitoren (Lovastatin, Simvastatin, Pravastatin, Fluvastatin, Atorvastatin, Rivastatin, Itavastatin, ZD-4522 und andere Statine), (ii) Sequestriermittel (Cholestyramin, Colestipol und Dialkylaminoalkylderivate eines vernetzten Dextrans), (iii) Nicotinyalkohol, Nicotinsäure oder ein Salz davon, (iv) PPAR α -Agonisten, wie z.B. Fenofibrinsäurederivate (Gemfibrozil, Clofibrat, Fenofibrat und Benzafibrat), (v) duale PPAR α/γ -Agonisten, wie z.B. KRP-297, (vi) Cholesterinabsorptionsinhi-

bitoren, wie zum Beispiel beta-Sitosterol, (vii) Acyl-CoA:Cholesterinacyltransferaseinhibitoren, wie zum Beispiel Avasimibe, und (viii) Antioxidantien, wie z.B. Probucol,
 (f) PPAR δ -Agonisten, wie z.B. diejenigen, die in der WO97/28149 offenbart sind,
 (g) Verbindungen gegen Fettleibigkeit, wie z.B. Fenfluramin, Dexfenfluramin, Phentiramin, Sulbitramin, Orlistat,
 (h) einem Ileum-Gallensäure-Transporter-Inhibitor und
 (i) Mitteln, die zur Verwendung bei Entzündungszuständen gedacht sind, wie z.B. Aspirin, nichtsteroidale Antiphlogistika, Glukokortikoide, Azulfidin und cyclooxygenase-2-selektive Inhibitoren, und
 (3) einen pharmazeutisch annehmbaren Träger enthält.

24. Eine pharmazeutische Zusammensetzung gemäß Anspruch 23, wobei die Komponente (2) ein HMG-CoA-Reduktase-Inhibitor ist.

Revendications

1. Composé de formule I :



dans laquelle :

R¹ est un groupe méthyle facultativement substitué par 1 à 3 F ;

R², R³ et R⁴ sont chacun indépendamment choisis dans le groupe consistant en H, un atome d'halogène, un groupe alkyle en C₁ à C₆, alcényle en C₂ à C₆, alcynyle en C₂ à C₆, cycloalkyle en C₃ à C₈, aryle, O-alkyle en C₁ à C₆, O-alcényle en C₂ à C₆, O-alcynyle en C₂ à C₆, O-aryle, OH, S-alkyle en C₁ à C₆, S-alcényle en C₂ à C₆, S-alcynyle en C₂ à C₆, SO₂-alkyle en C₁ à C₆, SO₂-alcényle en C₂ à C₆, SO₂-alcynyle en C₂ à C₆, OCON (R₅)₂, OCO(alkyle en C₁ à C₆) et CN, où tous les cas d'alkyle, d'alcényle et d'alcynyle sont facultativement linéaires ou ramifiés, et tous les cas d'alkyle, d'alcényle, d'alcynyle, de cycloalkyle et d'aryle sont facultativement substitués par 1 à 5 substituants indépendamment choisis dans le groupe consistant en un atome d'halogène, un groupe aryle, O-aryle et OMe ;

R⁵ et R⁶ sont, à chaque occurrence, indépendamment choisis dans le groupe consistant en H, F, OH et un groupe alkyle en C₁ à C₅, et les groupes R⁵ et R⁶ qui sont sur le même atome de carbone peuvent facultativement être joints pour former un groupe cycloalkyle en C₃ à C₆ ;

R⁷ et R⁸ sont chacun indépendamment choisis dans le groupe consistant en H, F et un groupe alkyle en C₁ à C₅, ou R⁷ et R⁸ peuvent facultativement être joints pour former un groupe cycloalkyle en C₃ à C₆ ;

R⁹ est choisi dans le groupe consistant en H et un groupe alkyle en C₁ à C₅, ledit groupe alkyle étant facultativement linéaire ou ramifié ;

Ar¹ est un groupe phényle, 1-naphtyle, 2-naphtyle, pyridyle ou quinoléyle où Ar¹ est substitué par 1 à 3 groupes indépendamment choisis parmi R⁴ ;

X est choisi dans le groupe consistant en C=O, S(O)₂, CH₂, CH(CH₃), C(CH₃)₂, CF₂ et cyclopropylidène ;

Y est O ou S ; et

n vaut 0 à 5 ;

et les sels et promédicaments pharmaceutiquement acceptables de ceux-ci.

2. Composé selon la revendication 1, dans lequel :

R^1 est CH_3 ;

R^2 , R^3 et R^4 sont chacun indépendamment choisis dans le groupe consistant en H, un atome d'halogène, un groupe alkyle en C_1 à C_6 , alcényle en C_2 à C_6 , alcynyle en C_2 à C_6 , cycloalkyle en C_3 à C_8 , aryle, O-alkyle en C_1 à C_6 , O-alcényle en C_2 à C_6 , O-alcynyle en C_2 à C_6 , O-aryle, OH, S-alkyle en C_1 à C_6 , S-alcényle en C_2 à C_6 , S-alcynyle en C_2 à C_6 , $OCN(R^5)_2$, OCO (alkyle en C_1 à C_6) et CN, où tous les cas d'alkyle, d'alcényle et d'alcynyle sont facultativement linéaires ou ramifiés, et tous les cas d'alkyle, d'alcényle, d'alcynyle, de cycloalkyle et d'aryle sont facultativement substitués par 1 à 5 substituants indépendamment choisis dans le groupe consistant en un atome d'halogène, un groupe aryle, O-aryle et OMe ; et

X est choisi dans le groupe consistant en $C=O$, CH_2 , $CH(CH_3)$, $C(CH_3)_2$, CF_2 et cyclopropylidène ;

3. Composé selon la revendication 1, dans lequel R^2 , R^3 et R^4 sont chacun indépendamment choisis dans le groupe consistant en H, OCH_3 , OCF_3 , F, Cl et CH_3 , où CH_3 est facultativement substitué par 1 à 3 groupes indépendamment choisis parmi F, Cl et OCH_3 .

4. Composé selon la revendication 1, dans lequel R^2 , R^3 et R^4 sont chacun indépendamment choisis dans le groupe consistant en H, OCH_3 , OCF_3 et Cl.

5. Composé selon la revendication 1, dans lequel R^5 et R^6 sont H.

6. Composé selon la revendication 1, dans lequel R^7 et R^8 sont chacun indépendamment CH_3 ou H.

7. Composé selon la revendication 1, dans lequel R^9 est H.

8. Composé selon la revendication 1, dans lequel X est $C=O$.

9. Composé selon la revendication 1, dans lequel Y est O.

10. Composé selon la revendication 1, dans lequel n vaut 0, 1 ou 2.

11. Composé selon la revendication 1, dans lequel n vaut 1.

12. Composé selon la revendication 1, dans lequel Ar^1 est un groupe phényle, 1-naphtyle ou 2-naphtyle.

13. Composé selon la revendication 1, dans lequel Ar^1 est un groupe phényle ou 2-naphtyle, Ar^1 étant substitué par 1 à 3 groupes indépendamment choisis parmi R^4 .

14. Composé selon la revendication 1, où aryle est un groupe phényle.

15. Composé selon la revendication 1, dans lequel :

R^1 est CH_3 ;

R^2 est choisi dans le groupe consistant en H, OCH_3 et OCF_3 ;

R^3 , R^5 , R^6 et R^9 sont H ;

R^4 est choisi dans le groupe consistant en H, Cl et OCH_3 .

R^7 et R^8 sont chacun indépendamment choisis dans le groupe consistant en H et CH_3 ;

X est $C=O$;





Y est O ;

et n vaut 1.


16. Composé selon la revendication 1, dans lequel R^9 est défini comme un groupe qui est facilement éliminé dans des conditions physiologiques pendant et après administration à un patient mammifère pour donner un acide carboxylique dans lequel R^9 est H, ou l'anion carboxylate de celui-ci, ou un sel pharmaceutiquement acceptable de celui-ci, et R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , Ar^1 , X, Y et n sont comme défini dans la revendication 1.

17. Composé selon la revendication 16, dans lequel OR^9 du groupe $-CO_2R^9$ est choisi dans le groupe consistant en $-OR^{10}$, $-OCH_2OR^{10}$, $-OCH(CH_3)OR^{10}$, $-OCH_2OC(O)R^{10}$, $-OCH(CH_3)OC(O)R^{10}$, $-OCH_2OC(O)OR^{10}$, $-OCH(CH_3)OC(O)OR^{10}$, $-NR^{11}R^{11}$ et $-ONR^{11}R^{11}$,
où chaque R^{10} est indépendamment choisi parmi un groupe alkyle en C_1 à C_6 facultativement substitué par un ou deux groupes choisis parmi $-CO_2H$, $-CONH_2$, NH_2 , $-OH$, $-OAc$, $NHAc$ et phényle ; et
où chaque R^{11} est indépendamment choisi parmi H et R^{10} .

18. Composé qui est choisi dans le groupe consistant en :




5




10

15




20

25




30

35








40






45













50



55











19. Composé qui est choisi dans le groupe consistant en :

l'acide (2S)-2-(3-([1-(4-méthoxybenzoyl)-2-méthyl-5-(trifluorométhoxy)-1H-indol-3-yl]méthyl)phénoxy)propanoïque;

l'acide 2-(2-([1-(4-chlorobenzoyl)-2-méthyl-5-(méthoxy)-1H-indol-3-yl]méthyl)phénoxy)acétique;

l'acide 2-(3-([1-(4-chlorobenzoyl)-2-méthyl-5-(méthoxy)-1H-indol-3-yl]méthyl)phénoxy)acétique

l'acide 2-(4-([1-(4-chlorobenzoyl)-2-méthyl-5-(méthoxy)-1H-indol-3-yl]méthyl)phénoxy)acétique;

l'acide 2-(2-([1-(4-chlorobenzoyl)-2-méthyl-5-(méthoxy)-1H-indol-3-yl]méthyl)phénoxy)propanoïque ;

l'acide 2(3-([1-(4-chlorobenzoyl)-2-méthyl-5-(méthoxy)-1H-indol-3-yl]méthyl)phénoxy)propanoïque ;

l'acide 2-(4-([1-(4-chlorobenzoyl)-2-méthyl-5-(méthoxy)-1H-indol-3-yl]méthyl)phénoxy)propanoïque ;

l'acide 2-(2-([1-(4-chlorobenzoyl)-2-méthyl-5-(méthoxy)-1H-indol-3-yl]méthyl)phénoxy)-2-méthylpropanoïque ;

l'acide 2-(3-([1-(4-chlorobenzoyl)-2-méthyl-5-(méthoxy)-1H-indol-3-yl]méthyl)phénoxy)-2-méthylpropanoïque ;

l'acide 2-(4-([1-(4-chlorobenzoyl)-2-méthyl-5-(méthoxy)-1H-indol-3-yl]méthyl)phénoxy)-2-méthylpropanoïque ;

l'acide 2-(2-([1-(4-méthoxybenzoyl)-2-méthyl-5-(trifluorométhoxy)-1H-indol-3-yl]méthyl)phénoxy)acétique ;

l'acide 2-(3-([1-(4-méthoxybenzoyl)-2-méthyl-5-(trifluorométhoxy)-1H-indol-3-yl]méthyl)phénoxy)acétique ;

l'acide 2-(2-([1-(4-méthoxybenzoyl)-2-méthyl-5-(trifluorométhoxy)-1H-indol-3-yl]méthyl)phénoxy) propanoïque ;

l'acide 2-(2-([1-(2-naphtoyl)-2-méthyl-5-(trifluorométhoxy)-1H-indol-3-yl]méthyl)phénoxy) propanoïque ;

l'acide (2R)-2-(2-([1-(4-méthoxybenzoyl)-2-méthyl-5-(trifluorométhoxy)-1H-indol-3-yl]méthyl)phénoxy) propanoïque ;

l'acide (2S)-2-(2-([1-(4-méthoxybenzoyl)-2-méthyl-5-(trifluorométhoxy)-1H-indol-3-yl]méthyl)phénoxy) propanoïque ;

l'acide 2-(3-([1-(4-méthoxybenzoyl)-2-méthyl-5-(trifluorométhoxy)-1H-indol-3-yl]méthyl)phénoxy) propanoïque ;

l'acide 2-(3-([1-(2-naphtoyl)-2-méthyl-5-(trifluorométhoxy)-1H-indol-3-yl]méthyl)phénoxy) propanoïque ;

l'acide 2-(3-([1-(4-chlorobenzoyl)-2-méthyl-5-(trifluorométhoxy)-1H-indol-3-yl]méthyl)phénoxy) propanoïque ;

l'acide 2-(3-([1-(2,4-dichlorobenzoyl)-2-méthyl-5-(trifluorométhoxy)-1H-indol-3-yl]méthyl)phénoxy) propanoïque ;

l'acide (2R)-2-(3-([1-(4-méthoxybenzoyl)-2-méthyl-5-(trifluorométhoxy)-1H-indol-3-yl]méthyl)phénoxy) propanoïque ;

l'acide (2R)-2-(3-([1-(2-naphtoyl)-2-méthyl-5-(trifluorométhoxy)-1H-indol-3-yl]méthyl)phénoxy) propanoïque ;

l'acide (2S)-2-(3-([1-(2-naphtoyl)-2-méthyl-5-(trifluorométhoxy)-1H-indol-3-yl]méthyl)phénoxy) propanoïque ;

l'acide 2-(2-([1-(4-méthoxybenzoyl)-2-méthyl-5-(trifluorométhoxy)-1H-indol-3-yl]méthyl)phénoxy)-2-méthylpropanoïque ;

l'acide 2-(3-([1-(4-méthoxybenzoyl)-2-méthyl-5-(trifluorométhoxy)-1H-indol-3-yl]méthyl)phénoxy)-2-méthylpropanoïque ;

l'acide 2-(2-([1-(2-naphtoyl)-2-méthyl-5-(trifluorométhoxy)-1H-indol-3-yl]méthyl)phénoxy)-2-méthylpropanoïque ;

l'acide 2-(3-([1-(2-naphtoyl)-2-méthyl-5-(trifluorométhoxy)-1H-indol-3-yl]méthyl)phénoxy)-2-méthylpropanoïque ;

l'acide (2R)-2-(3-{2-[1-(4-méthoxybenzoyl)-2-méthyl-5-(trifluorométhoxy)-1*H*-indol-3-yl]éthyl}phénoxy) propionique ;

l'acide (2S)-2-(3-[1-(4-méthoxybenzoyl)-2-méthyl-5-(trifluorométhoxy)-1*H*-indol-3-yl]phénoxy)propionique

l'acide (2S)-2-(3-(1-{1-(4-méthoxybenzoyl)-2-méthyl-5-(trifluorométhoxy)-1*H*-indol-3-yl}cyclopropyl)phénoxy)propanoïque; et

l'acide 2-{3-[[1-(4-méthoxybenzoyl)-2-méthyl-5-(trifluorométhoxy)-1*H*-indol-3-yl]phénoxy]-2-méthylpropanoïque ;

20. Composition pharmaceutique comprenant un composé selon l'une quelconque des revendications 1 à 19 ou un sel pharmaceutiquement acceptable de celui-ci et un support pharmaceutiquement acceptable.

21. Composé selon l'une quelconque des revendications 1 à 19 ou sel pharmaceutiquement acceptable de celui-ci pour une utilisation dans un procédé de traitement du corps humain par thérapie.

22. Utilisation d'un composé selon l'une quelconque des revendications 1 à 19 ou un sel pharmaceutiquement acceptable de celui-ci pour la fabrication d'un médicament destiné à traiter, réguler ou prévenir un(e) ou plusieurs maladies, troubles ou affections choisi(e)s parmi (1) le diabète sucré non dépendant de l'insuline (NIDDM), (2) l'hyperglycémie, (3) une faible tolérance au glucose, (4) la résistance à l'insuline, (5) l'obésité, (6) les troubles lipidiques, (7) la dyslipidémie, (8) l'hyperlipidémie, (9) l'hypertriglycéridémie, (10) l'hypercholestérolémie, (11) de faibles taux de HDL, (12) des taux de LDL élevés, (13) l'athérosclérose et ses séquelles, (14) la resténose vasculaire, (15) le syndrome du côlon irritable, (16) l'affection abdominale inflammatoire, comprenant la maladie de Crohn et la recto-colite hémorragique, (17) d'autres affections inflammatoires, (18) la pancréatite, (19) l'obésité abdominale, (20) la maladie neurodégénérative, (21) la rétinopathie, (22) les affections néoplasiques, (23) les tumeurs des adipocytes, (24) le carcinome des adipocytes, tel que le liposarcome, (25) le cancer de la prostate et d'autres cancers, comprenant les cancers de l'estomac, du sein, de la vessie et du côlon, (26) l'angiogenèse, (27) la maladie d'Alzheimer, (28) le psoriasis, (29) une tension artérielle élevée, (30) le syndrome X, (31) l'hyperandrogénisme ovarien (syndrome ovarien polykystique) et d'autres troubles où la résistance à l'insuline est une composante.

23. Composition pharmaceutique comprenant (1) un composé selon l'une quelconque des revendications 1 à 19 ou un sel pharmaceutiquement acceptable de celui-ci, (2) un ou plusieurs composés choisis dans le groupe consistant en :

(a) les sensibilisateurs à l'insuline comprenant (i) les agonistes PPAR γ tels que les glitazones (par exemple la troglitazone, la pioglitazone, l'englitazone, MCC-555, la rosiglitazone et similaires) et les composés révélés dans les documents WO 97/27 857, 97/28 115, 97/28 137 et 97/27 847; (ii) les biguanides tels que la metformine et la phenformine; (iii) les inhibiteurs de protéine tyrosine phosphatase-IB (PTP-1B), et (iv) les inhibiteurs de dipeptidyl peptidase IV (DP-IV);

(b) l'insuline ou les substances mimétiques de l'insuline ;

(c) les sulfonurées telles que le tolbutamide et le glipizide, ou les matières apparentées ;

(d) les inhibiteurs d' α -glucosidase (tels que l'acarbose) ;

(e) les agents hypocholestérolémiants tels que (i) les inhibiteurs de HMG-CoA réductase (la lovastatine, la simvastatine, la pravastatine, la fluvastatine, l'atorvastatine, la rivastatine, l'itavastatine, ZD-4522 et d'autres statines), (ii) les séquestrants (la cholestyramine, le colestipol et les dérivés dialkylaminoalkyle d'un dextrane réticulé), (iii) l'alcool nicotinique, l'acide nicotinique ou un sel de celui-ci, (iv) les agonistes de PPAR α tels que les dérivés d'acide fénofibrique (le gemfibrozil, le clofibrate, le fénofibrate et le benzafibrate), (v) les agonistes doubles PPAR α/γ tels que KRP-297, (vi) les inhibiteurs de l'absorption de cholestérol, tels que par exemple le bêta-sitostérol, (vii) les inhibiteurs d'acyl CoA-cholestérol acyltransférase, tels que par exemple l'avasimibc, et (viii) les antioxydants, tels que le probucol;

(f) les agonistes de PPAR δ tels que ceux révélés dans le document WO 97/28 149;

(g) les composés anti-obésité tels que la fenfluramine, la dexfenfluramine, la phentramine, la sulbitramine, l'orlistat, les inhibiteurs de neuropeptide Y5, et les agonistes de récepteur adrénergique $\beta 3$;

(h) un inhibiteur de transporteur d'acide biliaire iléal ; et

(i) les agents destinés à une utilisation dans les affections inflammatoires tels que l'aspirine, des médicaments anti-inflammatoires non stéroïdiens, des glucocorticoïdes, l'azulfidine et les inhibiteurs sélectifs de la cyclooxygénase 2 ; et

(3) un support pharmaceutiquement acceptable.

- 24.** Composition pharmaceutique selon la revendication 23, dans laquelle le composant (2) est un inhibiteur de la HMG-CoA réductase.

5

10

15

20

25

30

35

40

45

50

55