



Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2207323 C 2006/10/10

(11)(21) **2 207 323**

(12) **BREVET CANADIEN
CANADIAN PATENT**

(13) **C**

(86) Date de dépôt PCT/PCT Filing Date: 1995/12/12
(87) Date publication PCT/PCT Publication Date: 1996/06/20
(45) Date de délivrance/Issue Date: 2006/10/10
(85) Entrée phase nationale/National Entry: 1997/06/06
(86) N° demande PCT/PCT Application No.: US 1995/015613
(87) N° publication PCT/PCT Publication No.: 1996/018395
(30) Priorité/Priority: 1994/12/15 (US357,121)

(51) Cl.Int./Int.Cl. **A61K 31/428** (2006.01),
A61P 25/28 (2006.01)
(72) Inventeurs/Inventors:
HALL, EDWARD DALLAS, US;
VON VOIGTLANDER, PHILIP F., US;
ROHDE, FRANK A., DE
(73) Propriétaires/Owners:
BOEHRINGER INGELHEIM KG, DE;
PHARMACIA & UPJOHN COMPANY, US
(74) Agent: MACRAE & CO.

(54) Titre : UTILISATION DE PRAMIPEXOLE EN TANT QU'AGENT NEUROPROTECTEUR
(54) Title: USE OF PRAMIPEXOLE AS A NEUROPROTECTIVE AGENT

(57) Abrégé/Abstract:

The present invention provides the use of pramipexole as a neuroprotective agent.



PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ :**A61K 31/425****A1**

(11) International Publication Number:

WO 96/18395

(43) International Publication Date:

20 June 1996 (20.06.96)

(21) International Application Number: PCT/US95/15613

(22) International Filing Date: 12 December 1995 (12.12.95)

(30) Priority Data:

08/357,121

15 December 1994 (15.12.94) US

(60) Parent Application or Grant

(63) Related by Continuation

US

Filed on

08/357,121 (CIP)

15 December 1994 (15.12.94)

(71) Applicants (for all designated States except US): THE UPJOHN COMPANY [US/US]; 301 Henrietta Street, Kalamazoo, MI 49001 (US). BOEHRINGER INGELHEIM KG [DE/DE]; Postfach 200, D-55216 Ingelheim am Rhein (DE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): HALL, Edward, Dallas [US/US]; 1432 Woodland Avenue, Portage, MI 49002 (US). VON VOIGTLANDER, Philip, F. [US/US]; 1 South Lake Doster, Plainwell, MI 49080 (US). ROHDE, Frank, A. [DE/DE]; Am Geistert 4, D-55413 Weiler (DE).

(74) Agent: WELCH, Lawrence, T.; The Upjohn Company, Corporate Intellectual Property Law, 301 Henrietta Street, Kalamazoo, MI 49001 (US).

(81) Designated States: AL, AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TT, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, LS, MW, SD, SZ, UG).

Published*With international search report.**Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.*

(54) Title: USE OF PRAMIPEXOLE AS A NEUROPROTECTIVE AGENT

(57) Abstract

The present invention provides the use of pramipexole as a neuroprotective agent.

USE OF PRAMIPEXOLE AS A NEUROPROTECTIVE AGENT

FIELD OF THE INVENTION

The present invention relates to the use of pramipexole or 2-amino-6-n-propylamino-4,5,6,7-tetrahydrobenzo-thiazole, or the (+)- or (-)-enantiomers thereof,
5 and the pharmacologically acceptable salts thereof, as a neuroprotective agent.

BACKGROUND OF THE INVENTION

A number of central nervous system diseases and conditions result in neuronal damage. These conditions which can lead to nerve damage include:

Primary neurodegenerative disease; Huntington's Chorea; Stroke and other
10 hypoxic or ischemic processes; neurotrauma; metabolically induced neurological damage; sequelae from cerebral seizures; hemorrhagic stroke; secondary neurodegenerative disease (metabolic or toxic); Parkinson's disease; Alzheimer's disease, Senile Dementia of Alzheimer's Type (SDAT); age associated cognitive dysfunctions; or vascular dementia, multi-infarct dementia, Lewy body dementia, or
15 neurodegenerative dementia.

Pramipexole is a dopamine-D₃/D₂ agonist the synthesis of which is described in European Patent 186 087 and its counterpart, U.S. Patent 4,886,812. It is known primarily for the treatment of schizophrenia and Parkinson's disease. It is known from German patent application DE 38 43 227 that pramipexole lowers the plasma
20 level of prolactin. Also, this European patent application discloses the use of pramipexole in the treatment of drug dependency. Further, it is known from German patent application DE 39 33 738 that pramipexole can be used to decrease abnormal high levels of thyroid stimulating hormone (TSH). U.S. patent 5,112,842 discloses the transdermal administration of the compounds and transdermal systems
25 containing these active compounds. WO 94/12387 describes the use of pramipexole as an antidepressant agent.

Up to now there is no commercially available drug for the therapeutic treatment of stroke with proven evidence of efficacy.

Surprisingly and unexpectedly, it has been found that pramipexole and its
30 (+)-enantiomer also has a neuroprotective effect.

INFORMATION DISCLOSURE

Piribedil, a vasodilator which binds to a multitude of receptors including dopamine receptors, is reported to have an effect on functional and biochemical parameters in a gerbil model of global cerebral ischemia. See, e.g., Society for
35 Neuroscience Abstracts, 19:673 (1993); *id.*, at 1645.

Lisuride binds to several different receptors including dopamine D₂ and 5-HT_{1a} receptors. It is reported that Lisuride, when administered before the event, reduced brain edema and prolonged survival time in a rat model of cerebral infarction. Miya Zawa, et al. Nippon-Yakurigaku-Zasshi 98(6):449-561, (1991).

5

SUMMARY OF THE INVENTION

The present invention particularly provides a method for preventing neuronal damage in a patient suffering from or susceptible to such neuronal damage comprising the administration of an effective amount of 2-amino-6-n-propylamino-4,5,6,7-tetrahydrobenzothiazole, its (-)-enantiomer or (+)-enantiomer thereof, and
10 pharmacologically acceptable salts thereof especially an effective amount of pramipexole which is the (-)-enantiomer of 2-amino-6-n-propylamino-4,5,6,7-tetrahydrobenzothiazole-dihydrochloride or an effective amount of the (+)-enantiomer of 2-amino-6-n-propylamino-4,5,6,7-tetrahydrobenzothiazole dihydrochloride.

15

Conditions which can cause nerve damage are well-known to an ordinarily skilled neurologist or similar physician and include:

Primary neurogenerative disease;

Huntington's Chorea;

Stroke and other hypoxic or ischemic processes;

20

Neurotrauma;

Metabolically induced neurological damage;

Sequelae from cerebral seizures;

Hemorrhagic stroke;

Secondary neurodegenerative disease (metabolic or toxic);

25

Parkinson's disease;

Alzheimer's disease, other memory disorders; or

Vascular dementia, multi-infarct dementia, Lewy body dementia, or neurogenerative dementia.

The preferred indication for pramipexole, in the context of the present
30 invention, is Parkinson's disease which is characterized by progressive degeneration of nigrostriatal dopamine neurons. In this sense, the term Parkinson's disease also comprises the term Parkinson's syndrome. In addition to pramipexole's palliative action (i.e. replacement of the lost dopamine neurotransmitter function), the compound may slow the degeneration of surviving dopamine neurons and thereby
35 slows the progression of the disease.

The prophylactic use of the compound of this invention includes use as monotherapy in early or pre-symptomatic stages of Parkinson's disease and prevention of neurodegenerative disorders based on ischaemia.

The synthesis, formulation and administration of pramipexole is described in
 5 U.S. patents 4,843,086; 4,886,812; and 5,112,842.

2-Amino-6-n-propyl-amino-4,5,6,7-tetrahydrobenzothiazole, particularly the (-)-enantiomer thereof, and the pharmacologically acceptable acid addition salts thereof can be given for preventing of neuronal damage. The form of conventional
 10 galenic preparations consist essentially of an inert pharmaceutical carrier and an effective dose of the active substance; e.g., plain or coated tablets, capsules, lozenges, powders, solutions, suspensions, emulsions, syrups, suppositories, etc.

L-dopa is shown to be toxic to cerebellar granule cells in culture. Pramipexole and the (+) enantiomer blocked L-dopa toxicity. The EC_{50} for both was
 15 between 0.3 and 1 μ M and 10 μ M provided viability measurements that were equal to control cells not exposed to L-dopa. The mechanism of protection does not appear to involve receptor activation given that the (+) enantiomer is less active in monoamine receptor binding assays. The possibility exists that 2-amino-6-n-propylamino-4,5,6,7-tetrahydrobenzothiazole, its (+) and (-) enantiomers and the
 20 pharmacologically acceptable acid addition salts thereof, especially pramipexole and the (+) enantiomer are acting as antioxidants toward reactive oxygen species known to be generated from L-dopa incubation.

The effective dose range is 0.01 to 2.0 mg/kg. More preferred is a dose of 1-2 mg/kg *per os* (PO). The preferred total dose level for neuroprotection is 0.5-20
 25 mg/kg/day PO. The preferred human dose is 0.1-6.0 mg/day total dose, divided in 2 or 3 administrations. In addition to being administered by oral or intravenous route, pramipexole may be administered transdermally.

DESCRIPTION OF THE PREFERRED EMBODIMENTS

The present invention is seen more fully by the examples given below:

30 Example 1:

We examined histologically retrograde degeneration of the dopaminergic nigrostriatal (NS) tract over 28 days in gerbils subjected to a 10-minute period of forebrain ischemia via bilateral carotid occlusion. There was in control animals a 39% loss of NS cell bodies in the zona compacta (as judged by tyroxine hydroxylase
 35 immunocytochemistry) over that time period. Daily post-ischemic oral dosing (1

mg/kg PO BID beginning at 1 hr after insult) with pramipexole attenuated 28-day post-ischemic loss of NS DA neurons by 36% ($p < 0.01$ vs vehicle-treated).

Example 2:

We have successfully replicated the finding of Example 1; i.e., that
 5 pramipexole can protect nigrostriatal dopamine neurons in a gerbil model of 10 minutes of bilateral carotid occlusion-induced forebrain ischemia plus 28 days of post-ischemic survival. The administration of pramipexole was via an initial dose of 1 mg/kg PO 60 minutes after ischemia and again at the end of day 1 followed by twice daily dosing for the next 26 days. A twice daily dose of 0.3 mg/kg produced a
 10 threshold effect (14% increase in nigrostriatal dopamine neurons compared to vehicle). A twice daily dose of 1 mg/kg produced a significant 38% improvement in 28 day post-ischemic dopamine neuronal survival ($p < 0.0001$ vs vehicle-treated). This action appears to be specific for dopamine neurons since the post-ischemic loss of the non-dopaminergic neurons in the CA1 area of the hippocampus was not
 15 significantly affected.

Example 3:

Primary Cultures of Cerebellar Granular Cells

Primary cultures of cerebellar granular cells were prepared from 8-day-old Sprague Dawley rats (Charles River, Portage, MI) as described previously (1).
 20 Neurons were grown on 6-well, 35 mm culture dishes (Nunc, Denmark) for 8-9 days, 2 ml/well, at a density of 1×10^6 cells/ml. Glial cell proliferation was prevented by adding cytosine-arabino-furanoside-monophosphate (Sigma, St. Louis, MO), 19 hrs after plating at a final concentration of 10 μ M. Cultures generated by this method have been characterized and shown to contain more than 90% granule cells (2).

Cell Toxicity Models

Experiments were started after cells were 8 days in vitro (8 DIV). Cells were washed with two ml of serum free growth medium. Concentrations of 100X stock solutions of PPX and the (+) enantiomer in serum free growth medium were made. These were delivered to cells by adding 20 μ l to cells in growth medium per well.
 30 The final concentration of drugs ranged from 1 nM to 10 μ M. After 5 min 20 μ l of a 10X stock solution of L-dopa in serum free growth medium was added to each well so that the final concentration was 100 μ M. Cells were incubated under the above conditions for 24 hrs. For viability measurements cells were then washed twice and pulsed with 1 μ Ci/ml of α -(methyl)- 14 C-aminoisobutyric acid (New England
 35 Nuclear) in Locke's buffer for 1 hr. Cells were solubilized, following a wash, with

4822.P CP

0.5% Triton[®] X-100. The solubilized cells were then counted for radioactivity on a scintillation counter. Data was expressed as the mean of triplicates $\pm$ S.D. for each point.

Results from the table show that pramipexole was neuroprotective toward the toxicity (64.8 %) associated with L-dopa. At 10 μ M pramipexole, L-dopa treated cells were not different from controls. Pramipexole is shown to decrease slightly but significantly cAMP levels in cerebellar granular cells (3). This suggests the possible involvement of dopamine receptors (D_2 family) in the mechanism of neuroprotection. To test this hypothesis, the (+) enantiomer was tested in a parallel experiment with pramipexole. The (+) enantiomer has been shown to be inactive in a battery of binding assays that involve adrenergic and serotonergic receptors, and less active in dopaminergic receptors. The results show that the (+) enantiomer is equally as potent and effective compared to pramipexole as a neuroprotective agent in this assay.

The neuroprotective effects of pramipexole in L-dopa mediated toxicity in cerebellar granule cell does not appear to involve the activation of dopamine receptors.

The (+) enantiomer of pramipexole shows utility as a neuroprotectant despite the fact that it shows little ability to bind to monoamine receptors.

The possibility exists that pramipexole and it's (+) enantiomer can act as an antioxidant toward L-dopa toxicity which is known to involve the generation of reactive oxygen species.

TABLE 1
Toxicity (64.8%) of 100 μ M L-dopa:
Effects of pramipexole and (+) enantiomer.

	Dose (nM)	Pramipexole (% of control*)	(+) enantiomer (% of control*)
30	0	35.2 $\pm$ 8.5	35.2 $\pm$ 8.5
	1	30.0 $\pm$ 6.5	51.3 $\pm$ 5.8
	10	39.2 $\pm$ 9.4	57.8 $\pm$ 4.9
	100	64.8 $\pm$ 4.7	58.3 $\pm$ 10.3
	300	76.2 $\pm$ 7.3	81.0 $\pm$ 18.3
35	1000	65.5 $\pm$ 14.8	86.9 $\pm$ 7.0
	3000	84.9 $\pm$ 0.6	95.2 $\pm$ 2.5
	10000	90.0 $\pm$ 11.5	99.8 $\pm$ 8.9

* Control is defined as the radioactivity associated with cells that were exposed to vehicle buffer rather than 100 μ M L-dopa. Control value was equal to an average of 129547 CPMs for triplicate counts.

[®]trade-mark

CLAIMS

1. Use of a compound selected from 2-amino-6-n-propylamino-4,5,6,7-tetrahydrobenzothiazole or a pharmacologically-acceptable acid addition salt thereof for the preparation of a medicament for preventing neuronal damage or the progression of neuronal damage.
5
2. Use according to claim 1 wherein the compound is the (+) enantiomer.
3. Use according to claim 1 wherein the compound is the (-) enantiomer.
4. Use according to claim 1 , 2 or 3 wherein the compound is the dihydrochloride.
10
5. Use according to claim 1, 2, 3 or 4 wherein the neuronal damage results from Parkinson's disease, primary neurodegenerative disease, Huntingdon's Chorea, stroke and other hypoxic or ischemic processes, neurotrauma, metabolically induced neurological damage, sequelae from cerebral seizures, haemorrhagic stroke, secondary neurodegenerative disease (metabolic or toxic),
15 Alzheimer's disease, vascular dementia, multi-infarct dementia, Lewy body dementia, or neurogenerative dementia.
6. Use according to claim 5 wherein the neuronal damage results from Parkinson's disease.
7. Use according to any one of claims 1 to 6 wherein the dosage is 0.1 to 6 mg/day.
20
8. Use according to any one of claims 1 to 6 wherein the dosage is 0.5 to 20 mg/kg/day.

9. Use according to any one of claims 1 to 6 wherein the dosage is 1 to 2 mg/kg/day.
10. Use according to claim 8 or claim 9 wherein the medicament is adapted for *per os* administration.
- 5 11. Use of an effective amount of the compound 2-amino-6-n-propylamino-4,5,6,7- tetrahydrobenzothiazole or a pharmacologically-acceptable acid addition salt thereof to prevent neuronal damage or the progression of neuronal damage in a patient suffering from or susceptible to such neuronal damage.
- 10 12. The use of Claim 11 where the condition is selected from Parkinson's disease, primary neurodegenerative disease, Huntingdon's Chorea, stroke and other hypoxic or ischemic processes, neurotrauma, metabolically induced neurological damage, sequelae from cerebral seizures, haemorrhagic stroke, secondary neurodegenerative disease (metabolic or toxic), Alzheimer's disease, vascular dementia, multi-infarct dementia, Lewy body dementia, or
- 15 neurogenerative dementia.
13. The use of claim 12, where the condition is Parkinson's disease.
14. The use of claim 13, where the total dose of the compound is about 0.1 to 6 mg/day.
- 15 20 15. The use of claim 11, 12, 13 or 14 wherein the compound is the form of the (+) enantiomer.
16. The use of claim 11, 12, 13 or 14 wherein the compound is the form of the (-) enantiomer.

17. The use of claim 11, 12, 13, 14, 15 or 16 wherein the compound is the dihydrochloride.

18. The use of claim 11, 12, 13, 14, 15, 16 or 17 wherein said use is oral use.

5