



US012203100B2

(12) **United States Patent**  
**Whitehead et al.**

(10) **Patent No.:** **US 12,203,100 B2**

(45) **Date of Patent:** **\*Jan. 21, 2025**

(54) **DEVELOPMENT OF DENGUE VIRUS  
VACCINE COMPONENTS**

(71) Applicant: **The Government of the USA as  
represented by the Secretary, Dept. of  
Health and Human Services,**  
Bethesda, MD (US)

(72) Inventors: **Stephen S. Whitehead,** Bethesda, MD  
(US); **Joseph E. Blaney,** Gettysburg,  
PA (US); **Brian R. Murphy,** Bethesda,  
MD (US); **Ching-Juh Lai,** Bethesda,  
MD (US)

(73) Assignee: **The Government of the USA as  
represented by the Secretary, Dept. of  
Health and Human Services,**  
Bethesda, MD (US)

(\* ) Notice: Subject to any disclaimer, the term of this  
patent is extended or adjusted under 35  
U.S.C. 154(b) by 0 days.

This patent is subject to a terminal dis-  
claimer.

(21) Appl. No.: **18/355,265**

(22) Filed: **Jul. 19, 2023**

(65) **Prior Publication Data**

US 2024/0084269 A1 Mar. 14, 2024

**Related U.S. Application Data**

(60) Continuation of application No. 17/724,037, filed on  
Apr. 19, 2022, now Pat. No. 11,746,335, which is a  
continuation of application No. 16/912,359, filed on  
Jun. 25, 2020, now Pat. No. 11,332,722, which is a  
continuation of application No. 16/173,217, filed on  
Oct. 29, 2018, now Pat. No. 10,724,007, which is a  
continuation of application No. 14/742,533, filed on  
Jun. 17, 2015, now Pat. No. 10,160,957, which is a  
continuation of application No. 13/692,557, filed on  
Dec. 3, 2012, now Pat. No. 9,090,873, which is a  
division of application No. 12/376,756, filed as  
application No. PCT/US2007/076004 on Aug. 15,  
2007, now Pat. No. 8,337,860.

(60) Provisional application No. 60/837,723, filed on Aug.  
15, 2006.

(51) **Int. Cl.**

**C12N 7/04** (2006.01)  
**A61K 39/00** (2006.01)  
**C07K 14/005** (2006.01)  
**C07K 14/18** (2006.01)  
**C12N 7/00** (2006.01)

(52) **U.S. Cl.**

CPC ..... **C12N 7/045** (2013.01); **C07K 14/005**  
(2013.01); **C07K 14/1825** (2013.01); **C12N**  
**7/00** (2013.01); **A61K 2039/5254** (2013.01);  
**C12N 2770/24122** (2013.01); **C12N**  
**2770/24162** (2013.01); **Y02A 50/30** (2018.01)

(58) **Field of Classification Search**

CPC ..... **C12N 7/00**; **C12N 2770/24134**; **C12N**  
**2770/24161**; **C12N 2770/24162**; **C12N**  
**2770/24122**; **C12N 2770/24171**; **C12N**  
**2770/24121**; **C12N 7/045**; **C12N 15/86**;  
**C12N 2770/24143**; **C12N 2770/24151**;  
**A61K 2039/5254**; **A61K 39/12**; **A61K**  
**2039/53**; **C07K 14/005**; **C07K 14/1825**

See application file for complete search history.

(56) **References Cited**

**U.S. PATENT DOCUMENTS**

7,517,531 B2 4/2009 Whitehead et al.  
7,851,194 B2 12/2010 Markoff et al.  
8,075,903 B2 12/2011 Whitehead et al.  
8,337,860 B2 12/2012 Whitehead et al.  
9,090,873 B2 7/2015 Whitehead et al.  
RE46,042 E 6/2016 Whitehead et al.  
9,783,787 B2 10/2017 Whitehead et al.  
RE46,631 E 12/2017 Whitehead et al.

(Continued)

**FOREIGN PATENT DOCUMENTS**

WO WO 2008/022196 2/2008  
WO WO 03/092592 11/2013

**OTHER PUBLICATIONS**

Whitehead SS, Falgout B, Hanley KA, Blaney JE Jr, Markoff L,  
Murphy Br. A live, attenuated dengue virus type 1 vaccine candidate  
with a 30-nucleotide deletion in the 3' untranslated region is highly  
attenuated and immunogenic in monkeys. *J Virol.* Jan. 2003;77(2):1653-  
7. (Year: 2003).\*

Alvarez et al., "Role of RNA structures present at the 3' UTR of  
dengue virus on translation, RNA synthesis, and viral replication,"  
*Virology*, vol. 339, pp. 200-212, 2005.

Blaney et al., "Dengue virus type 3 vaccine candidates generated by  
introduction of deletions in the 3' untranslated region (3'-UTR) or  
by exchange of the DENV-3 3'-UTR with that of DENV-4,"  
*Vaccine*, vol. 26, pp. 817-828, 2008.

(Continued)

*Primary Examiner* — Rachel B Gill

(74) *Attorney, Agent, or Firm* — Klarquist Sparkman,  
LLP

(57) **ABSTRACT**

The invention is related to a dengue virus or chimeric  
dengue virus that contains a mutation in the 3' untranslated  
region (3'-UTR) comprising a  $\Delta 30$  mutation that removes  
the TL-2 homologous structure in each of the dengue virus  
serotypes 1, 2, 3, and 4, and nucleotides additional to the  
 $\Delta 30$  mutation deleted from the 3'-UTR that removes  
sequence in the 5' direction as far as the 5' boundary of the  
TL-3 homologous structure in each of the dengue serotypes  
1, 2, 3, and 4, or a replacement of the 3'-UTR of a dengue  
virus of a first serotype with the 3'-UTR of a dengue virus  
of a second serotype, optionally containing the  $\Delta 30$  mutation  
and nucleotides additional to the  $\Delta 30$  mutation deleted from  
the 3'-UTR; and immunogenic compositions, methods of  
inducing an immune response, and methods of producing a  
dengue virus or chimeric dengue virus.

**3 Claims, 22 Drawing Sheets**

**Specification includes a Sequence Listing.**

(56)

**References Cited**

## U.S. PATENT DOCUMENTS

RE46,641 E 12/2017 Whitehead et al.  
2018/0010099 A1 1/2018 Whitehead et al.

## OTHER PUBLICATIONS

Blaney Jr., et al., "Development of a Live Attenuated Dengue Virus Vaccine Using Reverse Genetics," *Viral Immunology*, vol. 19, No. 1, pp. 10-32, 2006.

Blaney Jr., et al., "Genetically Modified, Live Attenuated Dengue Virus Type 3 Vaccine Candidates," *Am. J. Trop. Med. Hyg.*, vol. 71, pp. 811-821, 2004.

Durbin et al., "rDEN2/4Δ30(ME), A Live Attenuated Chimeric Dengue Serotype 2 Vaccine Is Safe and Highly Immunogenic in Healthy Dengue-Naive Adults," *Human Vaccines*, vol. 2, No. 6, pp. 255-260, 2006.

International Search Report dated Jan. 13, 2009 for PCT/US2007/076004.

Men et al., "Dengue Type 4 Virus Mutants Containing Deletions in the 3' Noncoding Region of the RNA Genome: Analysis of Growth Restriction in Cell Culture and Altered Viremia Pattern and

Immunogenicity in Rhesus Monkeys," *Journal of Virology*, vol. 70, No. 5, pp. 3930-3937, 1996.

Proutski et al., "Secondary structure of the 3' untranslated region of flaviviruses: similarities and differences," *Nucleic Acids Research*, vol. 25, No. 6, pp. 1194-1202, 1997.

Whitehead et al., "Substitution of the structural genes of dengue virus type 4 with those of type 2 results in chimeric vaccine candidates which are attenuated for mosquitoes, mice, and rhesus monkeys," *Vaccine*, vol. 21, pp. 4307-4316, 2003.

Worobey et al., "Widespread intra-serotype recombination in natural populations of dengue virus," *Proc. Natl. Acad. Sci.*, vol. 96, pp. 7352-7357, 1999.


Yu et al., "The topology of bulges in the long stem of the flavivirus 3' stem-loop is a major determinant of RNA replication competence." *Journal of Virology*, vol. 79, pp. 2309-2324, 2005.

Zeng et al., "Identification of specific nucleotide sequences within the conserved 3'-SL in the dengue type 2 virus genome required for replication," *Journal of Virology*, vol. 72, pp. 7510-7522, 1988.


Zhou et al., "Comparative analysis reveals no consistent association between the secondary structure of the 3'-untranslated region of dengue viruses and disease syndrome," *Journal of General Virology*, vol. 87, pp. 2595-2603, 2006.

\* cited by examiner

**A. Delete additional nucleotides from the 3' UTR:**



**B. Replace the 3' UTR of DEN3 with the UTR from DEN4 $\Delta 30$ :**



**Figure 1**






Figure 1b




Figure 1c




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6




Figure 7




Figure 8





Figure 9



DEN1Δ30/31  
ΔG=-71.3  
(mfold v3.2: 1 0 10, P 114 0 10)

Figure 10




Figure 11




Figure 12




Figure 13




Figure 14




Figure 15




Figure 16




Figure 17




Figure 18




Figure 19

## DEVELOPMENT OF DENGUE VIRUS VACCINE COMPONENTS

### CROSS REFERENCE TO RELATED APPLICATIONS

This application is a continuation of U.S. patent application Ser. No. 17/724,037, filed Apr. 19, 2022, which is a continuation of U.S. patent application Ser. No. 16/912,359, filed Jun. 25, 2020, which is a continuation of U.S. patent application Ser. No. 16/173,217, filed Oct. 29, 2018, which is a continuation of U.S. patent application Ser. No. 14/742,533, filed Jun. 17, 2015, which is a continuation of U.S. patent application Ser. No. 13/692,557, filed Dec. 3, 2012, which is a divisional application of U.S. patent application Ser. No. 12/376,756, filed Dec. 17, 2009, which is a National Phase of International Application No. PCT/US2007/076004, filed on Aug. 15, 2007, which claims the benefit of U.S. Provisional Application No. 60/837,723, filed Aug. 15, 2006. The disclosures of each of these applications are hereby expressly incorporated by reference in their entirety.

### FIELD OF THE INVENTION

The invention relates to mutations in the 3' untranslated region of the genome of dengue virus serotypes 1, 2, 3, and 4 that are useful in attenuating the growth characteristics of dengue virus vaccines.

### DESCRIPTION OF THE RELATED ART

There are four serotypes of dengue virus (dengue virus type 1 [DEN1], DEN2, DEN3, and DEN4) that annually cause an estimated 50 to 100 million cases of dengue fever and 500,000 cases of the more severe form of dengue virus infection known as dengue hemorrhagic fever/dengue shock syndrome (Gubler, D. J. and M. Meltzer 1999 *Adv Virus Res* 53:35-70). Dengue virus is widely distributed throughout the tropical and semitropical regions of the world, and the number of dengue virus infections continues to increase due to the expanding range of its *Aedes aegypti* mosquito vector. A vaccine is not available for the control of dengue disease despite its importance as a reemerging disease. The goal of immunization is to protect against dengue virus disease by the induction of a long-lived neutralizing antibody response against each of the four serotypes. Simultaneous protection against all four serotypes is required, since an increase in disease severity can occur in persons with preexisting antibodies to a heterotypic dengue virus. Such immunization can be achieved economically with a live, attenuated virus vaccine.

Dengue viruses are positive-sense RNA viruses belonging to the Flavivirus genus. The approximately 11,000-base genome contains a single open reading frame encoding a polyprotein which is processed by proteases of both viral and cellular origin into three structural proteins (C, prM, and E) and at least seven nonstructural (NS) proteins. Both ends of the dengue virus genome contain an untranslated region (UTR), and the overall genome organization is 5MJTR-C-prM-E-NSI-NS2A-NS2B-NS3-NS4A-NS4B-NS5-UTR-3'. The 3' UTR is nearly 400 bases in length and is predicted to contain several stem-loop structures conserved among dengue virus serotypes (Brinton, M. A. et al. 1986 *Virology* 153:113-121, Hahn, C. S. et al. 1987 *J Mol Biol* 198:33-41, Proutski, V. et al. 1997 *Nucleic Acids Res* 25:1194-1202, Rauscher, S. et al. 1997 *RNA* 3:779-791, Shurtleff, A. et al. 2001 *Virology* 281:75-87). One such stem-loop structure,

identified as TL-2 in the proposed secondary structure of the 3' UTR (Proutski, V. et al. 1997 *Nucleic Acids Res* 25:1194-1202), was previously removed by deletion of 30 nucleotides from the DEN4 genome (3' nucleotides 172 to 143) (Men, R. et al. 1996 *J Viral* 70:3930-3937) and has subsequently been designated as the  $\Delta 30$  mutation (Durbin, A. P. et al. 2001 *Am J Trop Med Hyg* 65:405-413). The resulting virus, rDEN4 $\Delta 30$ , was shown to be attenuated in rhesus monkeys compared to parental viruses containing an intact TL-2 sequence and is attenuated in humans (Durbin, A. P. et al. 2001 *Am J Trop Med Hyg* 65:405-413).

### SUMMARY OF THE INVENTION

The invention is related to a dengue virus or chimeric dengue virus comprising a mutation in the 3' untranslated region (3'-UTR) selected from the group consisting of:

- a  $\Delta 30$  mutation that removes the TL-2 homologous structure in each of the dengue virus serotypes 1, 2, 3, and 4, and nucleotides additional to the  $\Delta 30$  mutation deleted from the 3'-UTR that removes sequence in the 5' direction as far as the 5' boundary of the TL-3 homologous structure in each of the dengue virus serotypes 1, 2, 3, and 4; and
- (b) a replacement of the 3'-UTR of a dengue virus of a first serotype with the 3'-UTR of a dengue virus of a second serotype, optionally containing the 430 mutation and nucleotides additional to the 430 mutation deleted from the 3'-UTR;

and immunogenic compositions, methods of inducing an immune response, and methods of producing a dengue virus or chimeric dengue virus.

### BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1. Two approaches to attenuate dengue viruses. A) (a-c) Deletion of additional nucleotides from the 3'-UTR (DEN3 wt Sleman/78, SEQ ID NO: 1). B) Replacement of the 3'-UTR of a dengue virus of a first serotype with the 3'-UTR of a dengue virus of a second serotype.

FIG. 1a is a magnified version of FIG. 1 in the portion designated as A(a).

FIG. 1b is a magnified version of FIG. 1 in the portion designated as A(b).

FIG. 1c is a magnified version of FIG. 1 in the portion designated as A(c).

FIG. 2. Predicted secondary structure of the TL-1, TL-2 and TL-3 region of the 3'-UTR of each DEN serotype. The GenBank accession number of the sequence used for construction of each secondary structure model is indicated. Only the last 278, 281, 276 and 281 nucleotides of DEN1, DEN2, DEN3, and DEN4, respectively, which comprise TL-1, TL-2 and TL-3, are used to avoid circularization of the structure and subsequent misfolding of known and experimentally-verified structural elements. The mfold program constraints specific for each structure model are indicated. Nucleotides that border the principle deletions are circled and numbered, with nucleotide numbering beginning at the 3' genome end (reverse-direction numbering system). The nucleotide sequence shown in FIG. 2.—SEQ ID NO: 2.

FIG. 3. Predicted secondary structure of the TL-1, TL-2 and TL-3 region of the 3'-UTR of each DEN serotype. The GenBank accession number of the sequence used for construction of each secondary structure model is indicated. Only the last 278, 281, 276 and 281 nucleotides of DEN1, DEN2, DEN3, and DEN4, respectively, which comprise TL-1, TL-2 and TL-3, are used to avoid circularization of the

3

structure and subsequent misfolding of known and experimentally-verified structural elements. The refold program constraints specific for each structure model are indicated. Nucleotides that border the principle deletions are circled and numbered, with nucleotide numbering beginning at the 3' genome end (reverse-direction numbering system). The nucleotide sequence shown in FIG. 3:—SEQ ID NO: 3.

FIG. 4. Predicted secondary structure of the TL-1, TL-2 and TL-3 region of the 3'-UTR of each DEN serotype. The GenBank accession number of the sequence used for construction of each secondary structure model is indicated. Only the last 278, 281, 276 and 281 nucleotides of DEN1, DEN2, DEN3, and DEN4, respectively, which comprise TL-1, TL-2 and TL-3, are used to avoid circularization of the structure and subsequent misfolding of known and experimentally-verified structural elements. The mfold program constraints specific for each structure model are indicated. Nucleotides that border the principle deletions are circled and numbered, with nucleotide numbering beginning at the 3' genome end (reverse-direction numbering system). The nucleotide sequence shown in FIG. 4:—SEQ ID NO: 4.

FIG. 5. Predicted secondary structure of the TL-1, TL-2 and TL-3 region of the 3'-LITR of each DEN serotype. The GenBank accession number of the sequence used for construction of each secondary structure model is indicated. Only the last 278, 281, 276 and 281 nucleotides of DEN1, DEN2, DEN3, and DEN4, respectively, which comprise n-1, TL-2 and TL-3, are used to avoid circularization of the structure and subsequent misfolding of known and experimentally-verified structural elements. The ifold program constraints specific for each structure model are indicated. Nucleotides that border the principle deletions are circled and numbered, with nucleotide numbering beginning at the 3' genome end (reverse-direction numbering system). The nucleotide sequence shown in FIG. 5:—SEQ ID NO: 5.

FIG. 6.  $\Delta 30$  deletion mutation depicted for each of the dengue virus serotypes DEN1, DEN2, DEN3 and DEN4. The  $\Delta 30$  mutation deletes nt 174 to 145 of DEN1, nt 173 to 144 of DEN2, nt 173 to 143 of DEN3, and nt 172 to 143 of DEN4, with reverse-direction numbering system. The deleted region is indicated by the  $\Delta$  symbol. The nucleotide sequence shown in FIG. 6:—SEQ ID NO: 6.

FIG. 7.  $\Delta 30$  deletion mutation depicted for each of the dengue virus serotypes DEN1, DEN2, DEN3 and DEN4. The  $\Delta 30$  mutation deletes nt 174 to 145 of DEN1, nt 173 to 144 of DEN2, nt 173 to 143 of DEN3, and nt 172 to 143 of DEN4, with reverse-direction numbering system. The deleted region is indicated by the  $\Delta$  symbol. The nucleotide sequence shown in FIG. 7:—SEQ ID NO: 7.

FIG. 8.  $\Delta 30$  deletion mutation depicted for each of the dengue virus serotypes DEN1, DEN2, DEN3 and DEN4. The  $\Delta 30$  mutation deletes nt 174 to 145 of DEN1, nt 173 to 144 of DEN2, nt 173 to 143 of DEN3, and nt 172 to 143 of DEN4, with reverse-direction numbering system. The deleted region is indicated by the  $\Delta$  symbol. The nucleotide sequence shown in FIG. 8:—SEQ ID NO: 8.

FIG. 9.  $\Delta 30$  deletion mutation depicted for each of the dengue virus serotypes DEN1, DEN2, DEN3 and DEN4. The  $\Delta 30$  mutation deletes nt 174 to 145 of DEN1, nt 173 to 144 of DEN2, nt 173 to 143 of DEN3, and nt 172 to 143 of DEN4, with reverse-direction numbering system. The deleted region is indicated by the  $\Delta$  symbol. The nucleotide sequence shown in FIG. 9:—SEQ ID NO: 9.

FIG. 10.  $\Delta 30/31$  deletion mutation depicted for each of the dengue virus serotypes DEN1, DEN2, DEN3 and DEN4. In addition to the deletion of the nucleotides comprising the  $\Delta 30$  mutation, the  $\Delta 31$  mutation deletes nt 258 to 228 of

4

DEN1, DEN2, DEN3, and DEN4, with reverse-direction numbering system. The deleted region is indicated by the  $\Delta$  symbol. The nucleotide sequence shown in FIG. 10:—SEQ ID NO: 10.

FIG. 11.  $\Delta 30/31$  deletion mutation depicted for each of the dengue virus serotypes DEN1, DEN2, DEN3 and DEN4. In addition to the deletion of the nucleotides comprising the  $\Delta 30$  mutation, the  $\Delta 31$  mutation deletes nt 258 to 228 of DEN1, DEN2, DEN3, and DEN4, with reverse-direction numbering system. The deleted region is indicated by the  $\Delta$  symbol. The nucleotide sequence shown in FIG. 11:—SEQ ID NO: 11.

FIG. 12.  $\Delta 30/31$  deletion mutation depicted for each of the dengue virus serotypes DEN1, DEN2, DEN3 and DEN4. In addition to the deletion of the nucleotides comprising the  $\Delta 30$  mutation, the  $\Delta 31$  mutation deletes nt 258 to 228 of DEN1, DEN2, DEN3, and DEN4, with reverse-direction numbering system. The deleted region is indicated by the  $\Delta$  symbol. The nucleotide sequence shown in FIG. 12:—SEQ ID NO: 12.

FIG. 13.  $\Delta 30/31$  deletion mutation depicted for each of the dengue virus serotypes DEN1, DEN2, DEN3 and DEN4. In addition to the deletion of the nucleotides comprising the  $\Delta 30$  mutation, the  $\Delta 31$  mutation deletes nt 258 to 228 of DEN1, DEN2, DEN3, and DEN4, with reverse-direction numbering system. The deleted region is indicated by the  $\Delta$  symbol. The nucleotide sequence shown in FIG. 13:—SEQ ID NO: 13.

FIG. 14.  $\Delta 86$  deletion mutation depicted for each of the dengue virus serotypes DEN1, DEN2, DEN3 and DEN4. The  $\Delta 86$  mutation deletes nt 228 to 145 of DEN1, nt 228 to 144 of DEN2, nt 228 to 143 of DEN3, and nt 228 to 143 of DEN4, with reverse-direction numbering system. The deleted region is indicated by the  $\Delta$  symbol. The nucleotide sequence shown in FIG. 14:—SEQ ID NO: 14.

FIG. 15.  $\Delta 86$  deletion mutation depicted for each of the dengue virus serotypes DEN1, DEN2, DEN3 and DEN4. The  $\Delta 86$  mutation deletes nt 228 to 145 of DEN1, nt 228 to 144 of DEN2, nt 228 to 143 of DEN3, and nt 228 to 143 of DEN4, with reverse-direction numbering system. The deleted region is indicated by the  $\Delta$  symbol. The nucleotide sequence shown in FIG. 15:—SEQ ID NO: 15.

FIG. 16.  $\Delta 86$  deletion mutation depicted for each of the dengue virus serotypes DEN1, DEN2, DEN3 and DEN4. The  $\Delta 86$  mutation deletes nt 228 to 145 of DEN1, nt 228 to 144 of DEN2, nt 228 to 143 of DEN3, and nt 228 to 143 of DEN4, with reverse-direction numbering system. The deleted region is indicated by the  $\Delta$  symbol. The nucleotide sequence shown in FIG. 16:—SEQ ID NO: 16.

FIG. 17.  $\Delta 86$  deletion mutation depicted for each of the dengue virus serotypes DEN1, DEN2, DEN3 and DEN4. The  $\Delta 86$  mutation deletes nt 228 to 145 of DEN1, nt 228 to 144 of DEN2, nt 228 to 143 of DEN3, and nt 228 to 143 of DEN4, with reverse-direction numbering system. The deleted region is indicated by the  $\Delta$  symbol. The nucleotide sequence shown in FIG. 17:—SEQ ID NO: 17.

FIG. 18. Chimerization of rDEN3 with the rDEN4 or rDEN4  $\Delta 30$  ( ) 3'-UTR. A) recombinant 3'-UTR chimeric dengue viruses were constructed by replacing the 3'-urR of rDEN3 with regions derived from either rDEN4 or rDEN4  $\Delta 30$ . The relative location of the  $\Delta 30$  mutation in the 3'-UTR is indicated by an arrow. The junctions between the ORF and UTR for rDEN3 and rDEN4 are indicated as junctions 1 and 2, respectively. Intertypic junction 3 is also indicated for the resulting chimeric viruses. B) nucleotide and amino acid sequence of the junction regions are shown. For junction 3, nucleotide substitutions used to introduce a unique hpaI

restriction enzyme recognition site are shown in lower case. Junction 1—SEQ ID NOS: 18 (nucleotide) and 19 (amino acid); junction 2—SEQ ID NOS: 20 (nucleotide) and 21 (amino acid); junction 3—SEQ NOS: 22 (nucleotide) and 23 (amino acid).

FIG. 19. Replication in vero cells and C6/36 cells. Four mutant viruses were compared to wild type rDEN3 for replication in vero cells and C6/36 cells. 75 cm<sup>2</sup> flasks of confluent cells were infected at a multiplicity of infection of 0.01. Aliquots of 0.5 ml were removed from flasks daily for seven days. After addition of spg to a concentration of 1×, samples were frozen on dry ice and stored at -80° C. Virus titer was determined by plaque assay on vero cells for all samples. The limit of detection is 1.0 log<sub>10</sub>pfu/ml.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENT

Definitions

Unless defined otherwise, technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. See, e.g., Singleton P and Sainsbury D., Dictionary of Microbiology and Molecular Biology 3rd ed., J. Wiley & Sons, Chichester, New York, 2001, and Fields Virology 4th ed., Knipe D.M. and Howley P.M. eds, Lippincott Williams & Wilkins, Philadelphia 2001.

The term “about” means within 1.2, or 3 nucleotides. Mutant Dengue Viruses and Chimeric Dengue Viruses

A goal of the invention is to develop a set of type-specific, live attenuated dengue vaccine components that can be formulated into a safe, effective, and economical tetravalent dengue vaccine. The Δ30 mutation attenuates DEN4 in rhesus monkeys (Men. R. et al. 1996 *J Viral* 70:3930-3937)). The Δ30 mutation removes a homologous structure (TL-2) in each of the dengue virus serotypes 1, 2, 3, and 4 (FIGS. 2-5). However, the Δ30 mutation was found to not attenuate DEN 3 in rhesus monkeys.

An embodiment of the invention provides dengue viruses and chimeric dengue viruses having one or more mutations that result in attenuation, methods of making such dengue viruses, and methods for using these dengue viruses to prevent or treat dengue virus infection. The mutation (or mutations) in the dengue virus of the invention is present in the 3' untranslated region (3'-UTR) formed by the most downstream approximately 384 nucleotides of the viral RNA, which have been shown to play a role in determining attenuation. The viruses and methods of the invention are described further, as follows.

One example of a dengue virus that can be used in the invention is the serotype 3, Sleman/78 strain. The applicability of the invention to all members of the dengue virus taxonomic group is inferred by the observation that the properties of other dengue virus strains are similar to that of any one dengue virus strain. Dengue viruses have been grouped into four serotypes (DEN1, DEN2, DENS and DEN4). Numerous strains have been identified for each of the four serotypes. The complete genomic sequences of various dengue virus strains are provided as Genbank accession numbers in Table A.

TABLE A

Examples of Dengue Virus Strains		
Serotype	Strain	Accession No.
1	02-20	AB178040
1	16007	AF180817
1	16007 PDK-13	AF180818

TABLE A-continued

Examples of Dengue Virus Strains			
Serotype	Strain	Accession No.	
5	1	259par00	AF514883
	1	280par00	AF514878
	1	293arg00	AY206457
	1	295arg00	AF514885
	1	297arg00	AF514889
	10	301arg00	AF514876
	1	98901518	AB189120
	1	98901530	AB189121
	1	A88	AB074761
	1	Abidjan	AF298807
	1	ARG0028	AY277665
	15	ARG0048	AY277666
	1	ARG9920	AY277664
	1	BR-90	AF226685
	1	BR-01-MR	AF513110
	1	BR-97-111	AF311956
	1	BR-97-233	AF311958
	1	BR-97-409	AF311957
	20	Cambodia	AF309641
	1	FGA-89	AF226687
	1	FGA-NA d1d	AF226686
	1	Fj231-04	DQ193572
	1	GD05-99	AY376738
	1	GD23-95	AY373427
	25	GZ-80	AF350498
	1	D1-hu-Yap-NIID27-2004	AB204803
	1	D1-H-IMTSSA-98-606	AF298808
	1	Mochizuki	AB074760
	1	D1.Myanmar.059-01	AY708047
	1	D1.Myanmar.194-01	AY713474
	30	D1.Myanmar.206-01	AY713475
	1	D1.Myanmar.23819-96	AY722802
	1	D1.Myanmar.305-01	AY713476
	1	D1.Myanmar.31459-98	AY726555
	1	D1.Myanmar.31987-98	AY726554
	1	D1.Myanmar.32514-98	AY722803
	35	D1.Myanmar.37726-01	AY726549
	1	D1.Myanmar.38862-01	AY726550
	1	D1.Myanmar.40553-71	AY713473
	1	D1.Myanmar.40568-76	AY722801
	1	D1.Myanmar.44168-01	AY726551
	1	D1.Myanmar.44988-02	AY726552
	40	D1.Myanmar.49440-02	AY726553
	1	rWestern Pacific-delta30	AY145123
	1	Western Pacific rDEN1mutF	AY145122
	1	S275-90	A75711
	1	D1-hu-Seychelles-NIID41-2003	AB195673
	1	Singapore 8114-93	AY762084
	45	Singapore S275-90	M87512
	1	ThD1_0008_81	AY732483
	1	ThD1_0049_01	AY732482
	1	ThD1_0081_82	AY732481
	1	ThD1_0097_94	AY732480
	1	ThD1_0102_01	AY732479
	50	ThD1_0323_91	AY732478
	1	ThD1_0336_91	AY732477
	1	ThD1_0442_80	AY732476
	1	ThD1_0488_94	AY732475
	1	ThD1_0673_80	AY732474
	1	Recombinant Western Pacific	AY145121
	55	Nauru Island Western Pacific 45AZ5	NC_001477
	1	Nauru Island, Western Pacific Bethesda	U88535
	1	Nauru Island Western Pacific 45AZ5-PDK27	U88537
	2	131	AF100469
	2	16681-PDK53	M84728
	2	16681 Blok	M84727
	60	16681 Kinney	U87411
	2	43	AF204178
	2	44	AF204177
	2	98900663	AB189122
	2	98900665	AB189123
	2	98900666	AB189124
	65	BA05i	AY858035
	2	Bangkok 1974	AJ487271

TABLE A-continued

Examples of Dengue Virus Strains		
Serotype	Strain	Accession No.
2	BR64022	AF489932
2	C0166	AF100463
2	C0167	AF100464
2	C0371	AF100461
2	C0390	AF100462
2	China 04	AF119661
2	Cuba115-97	AY702036
2	Cuba13-97	AY702034
2	Cuba165-97	AY702038
2	Cuba205-97	AY702039
2	Cuba58-97	AY702035
2	Cuba89-97	AY702037
2	DR23-01	AB122020
2	DR31-01	AB122021
2	DR59-01	AB122022
2	FJ-10	AF276619
2	FJ11-99	AF359579
2	I348600	AY702040
2	IQT1797	AF100467
2	IQT2913	AF100468
2	Jamaica-N.1409	M20558
2	K0008	AF100459
2	K0010	AF100460
2	Mar4	AF100466
2	DEN2-H-IMTSSA-MART-98-703	AF208496
2	New Guinea C	AF038403
2	New Guinea C-PUO-218 hybrid	AF038402
2	New Guinea-C	M29095
2	PDK-53	U87412
2	S1 vaccine	NC_001474
2	TB16i	AY858036
2	ThD2_0017_98	DQ181799
2	ThD2_0026_88	DQ181802
2	ThD2_0038_74	DQ181806
2	ThD2_0055_99	DQ181798
2	ThD2_0078_01	DQ181797
2	ThD2_0168_79	DQ181805
2	ThD2_0263_95	DQ181800
2	ThD2_0284.90	DQ181801
2	ThD2_0433_85	DQ181803
2	ThD2_0498_84	DQ181804
2	ThNH-28-93	AF022435
2	ThNH29-93	AF169678
2	ThNH36-93	AF169679
2	ThNH45-93	AF169680
2	ThNH-52-93	AF022436
2	ThNH54-93	AF169682
2	ThNH55-93	AF169681
2	ThNH62-93	AF169683
2	ThNH63-93	AF169684
2	ThNH69-93	AF169685
2	ThNH73-93	AF169686
2	ThNH76-93	AF169687
2	ThNH81-93	AF169688
2	ThNH-p36-93	AF022441
2	ThNH-7-93	AF022434
2	ThNH-p11-93	AF022437
2	ThNH-p12-93	AF022438
2	ThNH-p14-93	AF022439
2	ThNH-p16-93	AF022440
2	Tonga-74	AY744147
2	TSV01	AY037116
2	Taiwan-1008DHF	AY776328
2	Ven2	AF100465
3	D3-H-IMTSSA-MART-1999-1243	AY099337
3	D3-H-IMTSSA-SRI-2000-1266	AY099336
3	80-2	AF317645
3	98901403	AB189125
3	98901437	AB189126
3	98901517	AB189127
3	98902890	AB189128
3	BA51	AY858037
3	BDH02-1	AY496871
3	BDH02-3	AY496873
3	BDH02-4	AY496874
3	BDH02-7	AY496877

TABLE A-continued

Examples of Dengue Virus Strains		
Serotype	Strain	Accession No.
3	BR74886-02	AY679147
3	C0331-94	AY876494
3	C0360-94	AY923865
3	den3_88	AY858038
3	den3_98	AY858039
10	FW01	AY858040
3	FW06	AY858041
3	H87	NC_001475
3	D3-Hu-TL018NIID-2005	AB214879
3	D3-Hu-TL029NIID-2005	AB214880
3	D3-Hu-TL109NIID-2005	AB214881
3	D3-Hu-TL129NIID-2005	AB214882
15	InJ_16_82	DQ401690
3	KJ30i	AY858042
3	KJ46	AY858043
3	KJ71	AY858044
3	mutant BDH02_01	DQ401689
3	mutant BDH02_03	DQ401691
20	mutant BDH02_04	DQ401692
3	mutant BDH02_07	DQ401693
3	mutant InJ_I6_82	DQ401694
3	mutant PhMH_J1_97	DQ401695
3	PF89-27643	AY744677
3	PF89-320219	AY744678
25	PF90-3050	AY744679
3	PF90-3056	AY744680
3	PF90-6056	AY744681
3	PF92-2956	AY744682
3	PF92-2986	AY744683
3	PH86	AY858045
30	PhMH-J1-97	AY496879
3	PI64	AY858046
3	Singapore	AY662691
3	Singapore 8120-95	AY766104
3	Sleman-78	AY648961
3	TB16	AY858047
35	TB55i	AY858048
3	ThD3 0007 87	AY676353
3	ThD3 0010 87	AY676353
3	ThD3 0055 93	AY676351
3	ThD3 0104 93	AY676350
3	ThD3 1283 98	AY676349
3	ThD3 1687 98	AY676348
40	PF92-4190	AY744684
3	PF94-136116	AY744685
3	Taiwan-739079A	AY776329
4	2A	AF375822
4	Recombinant clone rDEN4	AF326825
4	2Adel30	AF326826
45	814669	AF326573
4	B5	AF289029
4	rDEN4del30	AF326827
4	H241	AY947539
4	rDEN4	NC_002640
4	Singapore 8976-95	AY762085
50	SW38i	AY858050
4	ThD4_0017_97	AY618989
4	ThD4_0087_77	AY618991
4	ThD4_0348_91	AY618990
1	ThD4_0476_97	AY618988
4	ThD4_0485_01	AY618992
4	ThD4_0734_00	AY618993
55	Taiwan-2K0713	AY776330
4	Unknown	M14931

Mutations can be made in the 3'-UTR of a wild type infectious clone, e.g., dengue virus serotype 3, strain Sle-  
 marif78 or an infectious clone of another wild type, virulent  
 dengue virus, and the mutants can then be tested in an animal  
 model system (e.g., in mouse and/or monkey model sys-  
 tems) to identify sites that cause attenuation. Attenuation is  
 judged by, for example, detection of decreased viremia. One  
 or more additional mutations found to attenuate the wild-  
 type virus are optionally introduced into a wild type dengue

virus, and these mutants are tested in an animal model system (e.g., in a mouse and/or a monkey model system) to determine whether the resulting mutants are attenuated. Mutants that are found to be attenuated can then be used as new vaccine strains that have increased safety, due to attenuation.

In addition to the viruses listed above, dengue viruses including chimeric dengue viruses that include one or more attenuating mutations are included in the invention. These chimeras can consist of a dengue virus of a first serotype (i.e., a background dengue virus) in which a structural protein (or proteins) has been replaced with a corresponding structural protein (or proteins) of a dengue virus of a second serotype. For example, the chimeras can consist of a background dengue virus in which the prM and E proteins of the dengue virus of the first serotype have been replaced with the prM and E proteins of the dengue virus of the second serotype. The chimeric viruses can be made from any combination of dengue viruses of different serotypes. The dengues against which immunity is sought is the source of the inserted structural protein(s).

As is noted above, mutations that are included in the viruses of the present invention are attenuating. These mutations are present in the dengue virus 3'-UTR structure to attenuate the virus. Mutations can be made in the 3'-UTR using standard methods, such as site-directed mutagenesis. One example of the type of mutation present in the viruses of the invention is substitutions, but other types of mutations, such as deletions and insertions, can be used as well. In addition, as is noted above, the mutations can be present singly or in the context of one or more additional mutations.

Referring to FIG. 1, two approaches were taken to attenuate dengue virus. In one aspect, nucleotides additional to the  $\Delta 30$  mutation were deleted from the 3'-UTR. In another aspect, the 3'-UTR of a dengue virus of a first serotype was replaced with the 3'-UTR from a dengue virus of a second serotype (optionally containing the  $\Delta 30$  mutation and nucleotides additional to the  $\Delta 30$  mutation deleted from the 3'-UTR).

Deletion of Nucleotides Additional to the  $\Delta 30$  Mutation from the 3'-UTR

Referring to FIGS. 2-5, using the first approach, the 3'-UTR of dengue viruses contain various conserved sequence motifs. The sequence of the DEN4 3'-UTR is illustrated in FIG. 5. The genome of DEN4 strain 814669 contains 10,646 nucleotides, of which the last 384 nt at the 3' terminus are untranslated (non-coding). The locations of various sequence components in this region are designated with the reverse-direction numbering system. These sequences include the 3' distal secondary structure (nt 1 to 93), predicted to form stem-loop 1 (SL-1), which contains terminal loop 1 (TL-1). Nucleotides 117-183 form stem-loop 2 (SL-2) which contains TL-2. Nucleotides 201-277 form a pair of stem-loops (SL-3) which in part contains TL-3. Although the primary sequence of stem-loop 1 differs slightly among the dengue serotypes, the secondary structure is strictly conserved (compare FIGS. 2-5). Although the nucleotide spacing between SL-2 and neighboring SL-1 and SL-3 differ among the dengue virus serotypes, the overall structure of SL-2 is well-conserved. In addition, the exposed 9 nucleotides that comprise TL-2 are identical within all 4 dengue serotypes. It is TL-2 and its supporting stem structure that are removed by the  $\Delta 30$  mutation (about nt 143-172). Removal of these 30 nucleotides results in formation of a new predicted structural element (SL-2 $\Delta 30$ ) which has

a primary sequence and secondary structure which is identical for each of the dengue virus serotypes (compare FIGS. 6-9).

FIGS. 10-13 illustrate the approach where nucleotides additional to the  $\Delta 30$  mutation are deleted from the 3'-UTR. The  $\Delta 30$  mutation removes the TL-2 homologous structure in each of the dengue virus serotypes 1, 2, 3, and 4. The approach where nucleotides additional to the  $\Delta 30$  mutation are deleted from the 3'-ITR removes the TL-2 homologous structure and sequence up to and optionally including the TL-3 homologous structure so that the deletion extends as far as the 5' boundary of the TL-3 homologous structure in each of the dengue virus serotypes 1, 2, 3, and 4. In the approach illustrated in FIGS. 10-14, an additional deletion of about 31 nucleotides from TL-3 results in formation of a new predicted structural element (SL-3 $\Delta 31$ ).

Referring to FIGS. 14-17, the  $\Delta 86$  mutation removes the TL-2 homologous structure and removes sequence up to the TL-3 homologous structure in each of the dengue virus serotypes DEN1, DEN2, DEN3 and DEN4. This deletion results in the formation of a new predicted structural element (SL-2 $\Delta 86$ ).

In some embodiments that involve deletion of nucleotides additional to the  $\Delta 30$  mutation, nucleic acid deletions are made to the 3'-UTR structure of the dengue virus genome to attenuate the virus while maintaining its immunogenicity. The deletions include the  $\Delta 30$  deletion (nt 173-143 of the serotype 3 Slemant78 strain in an exemplary manner or corresponding thereto in other strains of DEN1, DEN2, DEN3, or DEN4; numbering is from the 3' end of the viral genome) in addition to deletion of additional 3'-UTR sequence that is contiguous or non-contiguous to the  $\Delta 30$  deletion. The  $\Delta 30$  deletion corresponds to the TL-2 structure of the 3'-UTR. One type of embodiment, termed rDEN1 $\Delta 30/31$ , rDEN2 $\Delta 30/31$ , rDEN3 $\Delta 30/31$ , or rDEN4 $\Delta 30/31$  includes the original  $\Delta 30$  deletion and a non-contiguous 31 nt deletion that removes both the original TL-2 and TL-3 structures. Another type of embodiment, termed rDEN I  $\Delta 61$ , rDEN2 $\Delta 61$ , rDEN3 $\Delta 61$ , or rDEN4 $\Delta 61$  includes the  $\Delta 30$  deletion and deletion of 31 contiguous nucleotides extending 3' from the  $\Delta 30$  deletion. Another type of embodiment, termed rDEN1M&, rDEN2 $\Delta 86$ , rDEN3 $\Delta 86$ , or rDEN4 $\Delta 86$ , includes the  $\Delta 30$  deletion and deletion of 56 contiguous nucleotides extending 5' from the  $\Delta 30$  deletion. For DEN3, a complete list of mutant viruses constructed to contain 3'-UTR deletion mutations is presented below in Table 2.

Replacement of the 3'-UTR of a Dengue Virus of a First Serotype with the 3'-UTR from a Dengue Virus of a Second Serotype

Using the second approach, the 3'-UTR of rDEN3 may be replaced with the 3'-UTR of rDEN4, optionally containing the  $\Delta 30$  mutation and nucleotides additional to the  $\Delta 30$  mutation deleted from the 3'-UTR. Other examples include replacement of the 3'-UTR of rDEN3 with the 3'-UTR of dengue virus serotypes 1 and 2, optionally containing the  $\Delta 30$  mutation and nucleotides additional to the  $\Delta 30$  mutation deleted from the 3'-UTR. Other examples include: replacement of the 3'-UTR of rDEN1 with the 3'-UTR of dengue virus serotypes 2, 3, and 4, optionally containing the  $\Delta 30$  mutation and nucleotides additional to the  $\Delta 30$  mutation deleted from the 3'-UTR; replacement of the 3'-UTR of rDEN2 with the 3'-UTR of dengue virus serotypes 1, 3, and 4, optionally containing the  $\Delta 30$  mutation and nucleotides additional to the  $\Delta 30$  mutation deleted from the 3'-UTR; and, replacement of the 3'-UTR of rDEN4 with the 3'-UTR of

dengue virus serotypes 1, 2, and 3, optionally containing the  $\Delta 30$  mutation and nucleotides additional to the  $\Delta 30$  mutation deleted from the 3'-UTR.

Embodiments that involve replacement of the 3'-UTR of a dengue virus of a first serotype with the 3'-UTR of dengue virus of a second serotype include:

- a) rDEN1-3'D2, rDEN1-3'172x, rDEN1-3'D3, rDEN1-3'D3x, rDEN-3'D4, rDEN1-3'D4x; rDEN1/2-3'D1, rDEN1/2-3'1)x, rDEN1/2-3'D3, rDEN1/2-3'D3x, rDEN1/2-3'D4, rDEN1/2-3'D4x; rDEN1/3-3'D3, rDEN1/3-3'D1x, rDEN1/3-3'D2, rDEN1/3-3'D2x, rDEN1/3-3'D4, rDEN1/3-3'D-4x; rDEN1/4-3'D1, rDEN1/4-3'D1x, rDEN1/4-3'D2, rDEN1/4-3'D2x, rDEN1/4-3'D3, rDEN1/4-3'D3x;
- b) rDEN2-3'D1, rDEN2-3'D1x, rDEN2-3'D3, rDEN2-3'D3x, rDEN2-3'D4, rDEN2-3'D4x; rDEN2/1-3'D2, rDEN2/1-3'12x, rDEN 2/1-3'D3, rDEN2/1-3'D3x, rDEN2/1-3'D4, rDEN2/1-3'D4x; rDEN2/3-3'D1, rDEN2/3-3'D1x, rDEN2/3-3'D2, rDEN2/3-3'D2x, rDEN2/3-3'D4, rDEN2/3-3'D4x; rDEN2/4-3'D1, rDEN2/4-3'D1x, rDEN2/4-3'D2, rDEN2/4-3'D2x, rDEN2/4-3'D3, rDEN2/4-3'D3x;
- c) rDEN3-3'D1, rDEN3-3'D1x, rDEN3-3'D2, rDEN3-3'D2x, rDEN3-3'D4, rDEN3-3'134x; rDEN3/1-3'D2, rDEN3/1-3'D2x, rDEN3/1-3'D3, rDEN3/1-3'D3x, rDEN3/1-3'D4, rDEN3/1-3'D4x; rDEN3/2-3'D1, rDEN3/2-3'D1x, rDEN3/2-3'D3, rDEN3/2-3'D3x, rDEN3/2-3'D4, rDEN3/2-3'D4x; rDEN3/4-3'D1, rDEN3/4-3'D1x, rDEN3/4-3'D2, rDEN3/4-3'D2x, rDEN3/4-3'12x, rDEN3/4-3'13x, rDEN3/4-3'D3x; and
- d) rDEN4-3'1, rDEN4-3'131x, rDEN4-3'D2, rDEN4-3'D2x, rDEN4-3'D3, rDEN4-3'13x; rDEN4/1-3'D2, rDEN4/1-3'D2x, rDEN4/1-3'D3, rDEN4/1-3'D3x, rDEN4/1-3'D4, rDEN4/1-3'14x; rDEN4/2-3'D1, rDEN4/2-3'D1x, rDEN4/2-3'D3, rDEN4/2-3'D3x, rDEN4/2-3'D4, rDEN4/2-3'D4x; rDEN4/3-3'D1, rDEN4/3-3'D1x, rDEN4/3-3'D2, rDEN4/3-3'D2x, rDEN4/3-3'D4, rDEN4/3-3'D4x;
- where x is a mutation listed in Table 2.

Method of Making and Using Dengue or Chimeric Dengue Viruses

The viruses (including chimeric viruses) of the present invention can be made using standard methods in the art. For example, an RNA molecule corresponding to the genome of a virus can be introduced into host cells, e.g., Vero cells, from which (or the supernatants of which) progeny virus can then be purified. In this method, a nucleic acid molecule (e.g., an RNA molecule) corresponding to the genome of a virus is introduced into the host cells, virus is harvested from the medium in which the cells have been cultured, and the virus is formulated for the purposes of vaccination.

The viruses of the invention can be administered as primary prophylactic agents in adults or children at risk of infection, or can be used as secondary agents for treating infected patients. For example, in the case of DEN virus and chimeric DEN viruses, the vaccines can be used in adults or children at risk of DEN virus infection, or can be used as secondary agents for treating DEN virus-infected patients. Examples of patients who can be treated using the DEN virus--related vaccines and methods of the invention include (i) children in areas in which DEN virus is endemic, (ii) foreign travelers, (iii) military personnel, and (iv) patients in areas of a DEN virus epidemic. Moreover, inhabitants of regions into which the disease has been observed to be expanding (e.g., beyond Sri Lanka,

East Africa and Latin America), or regions in which it may be observed to expand in the future can be treated according to the invention.

Formulation of the invention can be carried out using methods that are standard in the art. Numerous pharmaceutically acceptable solutions for use in vaccine preparation are well known and can readily be adapted for use in the present invention by those of skill in this art (see, e.g., Remington's Pharmaceutical Sciences (18th edition), ed. A. Gennaro. 1990, Mack Publishing Co., Easton, PA). The viruses can be diluted in a physiologically acceptable solution, such as sterile saline, sterile buffered saline, or L-15 medium. In another example, the viruses can be administered and formulated, for example, as a fluid harvested from cell cultures infected with dengue virus or chimeric dengue virus.

The vaccines of the invention can be administered using methods that are well known in the art, and appropriate amounts of the vaccines administered can readily be determined by those of skill in the art. For example, the viruses of the invention can be formulated as sterile aqueous solutions containing between  $10^2$  and  $10^7$  infectious units (e.g., plaque-forming units or tissue culture infectious doses) in a dose volume of 0.1 to 1.0 ml, to be administered by, for example, intramuscular, subcutaneous, or intradermal routes. Further, the vaccines of the invention can be administered in a single dose or, optionally, administration can involve the use of a priming dose followed by a booster dose that is administered, e.g., 2-6 months later, as determined to be appropriate by those of skill in the art.

Optionally, adjuvants that are known to those skilled in the art can be used in the administration of the viruses of the invention. Adjuvants that can be used to enhance the immunogenicity of the viruses include, for example, liposomal formulations, synthetic adjuvants, such as (e.g., QS21), inuramyl dipeptide, monophosphoryl lipid A, or polyphosphazine. Although these adjuvants are typically used to enhance immune responses to inactivated vaccines, they can also be used with live vaccines.

#### Nucleic Acid Sequences

Nucleic acid sequences of DEN viruses are useful for designing nucleic acid probes and primers for the detection of deletion or chimeric 3'-UTRs in a sample or specimen with high sensitivity and specificity. Probes or primers corresponding to deletion or chimeric 3'-UTRs can be used to detect the presence of deletion or chimeric 3'-UTRs in general in the sample, to quantify the amount of deletion or chimeric 3'-UTRs in the sample, or to monitor the progress of therapies used to treat DEN virus infection. The nucleic acid and corresponding amino acid sequences are useful as laboratory tools to study the organisms and diseases and to develop therapies and treatments for the diseases.

Nucleic acid probes and primers selectively hybridize with nucleic acid molecules encoding deletion or chimeric 3'-UTRs or complementary sequences thereof. By "selective" or "selectively" is meant a sequence which does not hybridize with other nucleic acids to prevent adequate detection of the deletion or chimeric 3'-UTRs. Therefore, in the design of hybridizing nucleic acids, selectivity will depend upon the other components present in the sample. The hybridizing nucleic acid should have at least 70% complementarity with the segment of the nucleic acid to which it hybridizes. As used herein to describe nucleic acids, the term "selectively hybridizes" excludes the occasional randomly hybridizing nucleic acids, and thus has the same meaning as "specifically hybridizing." The selectively hybridizing nucleic acid probes and primers of this invention

can have at least 70%, 80%, 85%, 90%, 95%, 97%, 98% and 99% complementarity with the segment of the sequence to which it hybridizes, preferably 85% or more.

The present invention also contemplates sequences, probes and primers that selectively hybridize to the encoding nucleic acid or the complementary, or opposite, strand of the nucleic acid. Specific hybridization with nucleic acid can occur with minor modifications or substitutions in the nucleic acid, so long as functional species-species hybridization capability is maintained. By "probe" or "primer" is meant nucleic acid sequences that can be used as probes or primers for selective hybridization with complementary nucleic acid sequences for their detection or amplification, which probes or primers can vary in length from about 5 to 100 nucleotides, or preferably from about 10 to 50 nucleotides, or most preferably about 18-24 nucleotides. Isolated nucleic acids are provided herein that selectively hybridize with the species-specific nucleic acids under stringent conditions and should have at least five nucleotides complementary to the sequence of interest as described in *Molecular Cloning: A Laboratory Manual*, 2<sup>nd</sup> ed., Sambrook, Fritsch and Mardatis, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY, 1989.

If used as primers, the composition preferably includes at least two nucleic acid molecules which hybridize to different regions of the target molecule so as to amplify a desired region. Depending on the length of the probe or primer, the target region can range between 70% complementary bases and full complementarity and still hybridize under stringent conditions. For example, for the purpose of detecting the presence of deletion or chimeric 3'-UTRs, the degree of complementarity between the hybridizing nucleic acid (probe or primer) and the sequence to which it hybridizes is at least enough to distinguish hybridization with a nucleic acid from other organisms.

The nucleic acid sequences of the invention include a diagnostic probe that serves to report the detection of a cDNA amplicon amplified from the viral genomic RNA template by using a reverse-transcription/polymerase chain reaction (RT-PCR), as well as forward and reverse amplimers that are designed to amplify the cDNA amplicon. In certain instances, one of the amplimers is designed to contain a vaccine virus-specific mutation at the 3'-terminal end of the amplimer, which effectively makes the test even more specific for the vaccine strain because extension of the primer at the target site, and consequently amplification, will occur only if the viral RNA template contains that specific mutation.

Automated PCR-based nucleic acid sequence detection systems have been recently developed. TaqMan assay (Applied Biosystems) is widely used. A more recently developed strategy for diagnostic genetic testing makes use of molecular beacons (Tyagi and Kramer 1996 *Nature Biotechnology* 18:303-308). Molecular beacon assays employ quencher and reporter dyes that differ from those used in the TaqMan assay. These and other detection systems may be used by one skilled in the art.

Dengue Virus Type 3 (DEN3) Vaccine Components Generated by Introduction of Deletions in the 3' Untranslated Region (UTR) or Exchange of the DEN3 3'-UTR with that of DEN4

There are four dengue virus serotypes (DEN1, DEN2, DEN3, and DEN4) which circulate in tropical and subtropical regions of the world inhabited by more than 2.5 billion people (Gubler D J 1998 *Clin Microbiol Rev* 11:480-496). DEN viruses are endemic in at least 100 countries and cause more human disease than any other arbovirus. Annually,

there are an estimated 50-100 million dengue infections and hundreds of thousands of cases of dengue hemorrhagic fever/ shock syndrome (DHF/DSS), with children bearing much of the disease burden (Gubler D J and Meltzer M 1999 *Adv Virus Res* 53:35-70). DHF/DSS remains a leading cause of hospitalization and death of children in at least eight southeast Asian countries (World Health Organization 1997 *Dengue Haemorrhagic Fever: Diagnosis, Treatment, Prevention and Control*, 2<sup>nd</sup> edition, WHO, Geneva). The dramatic increase in both the incidence and severity of disease caused by the four DEN serotypes over the past two decades is due in large part to the geographic expansion of the mosquito vectors, *Aedes aegypti* and *Aedes albopictus*, and the increased prevalence of the four DEN serotypes (Gubler D J 1998 *Clin Microbiol Rev* 11:480-496). The dengue viruses are maintained in a life cycle of transmission from mosquito to human to mosquito with no other apparent viral reservoir participating in its life cycle in urban settings (Rice C M, 1996 in *Flaviviridae: The viruses and their replication*, Fields R N, Knipe D M, Howley P M, Chanock R M, Melnick J L., Monath T P, Roizman B, Straus S E, eds. Fields Virology. Philadelphia: Lippincott-Raven Publishers, pp. 931-959).

The DEN viruses, members of the Flaviviridae family, have spherical virions of approximately 40 to 60 nm, which contain a single-stranded positive-sense RNA genome. A single polypeptide is co-translationally processed by viral and cellular proteases generating three structural proteins (capsid C, membrane M, and envelope E) and at least seven non-structural (NS) proteins. The genome organization of the DEN viruses is 5' UTR-C-prM-E-NS1-NS2A-NS2B-NS3-NS4A-NS4B-NS5-UTR-3' (UTR-untranslated region, prM-membrane precursor) (Rice C M, 1996 in *Flaviviridae: The viruses and their replication*, Fields R N, Knipe D M, Howley P M, Chanock R M, Melnick J L., Monath T P, Roizman B, Straus S E, eds. Fields Virology. Philadelphia: Lippincott-Raven Publishers, pp. 931-959).

In response to the increasing incidence and severity of DEN infection, development of vaccines is being pursued to prevent DEN virus disease. An economical vaccine that prevents disease caused by the DEN viruses has become a global public health priority. The cost-effectiveness, safety, and long-term efficacy associated with the live attenuated vaccine against yellow fever (YE) virus, another mosquito-borne flavivirus, serves as a model for the feasibility of developing a live attenuated DEN virus vaccine (Monath T P, 1999 in *Yellow fever*, Plotkin S A, Orenstein W A, eds. Vaccines, Philadelphia: W. B. Saunders Co., 815-879). Additionally, an effective live attenuated Japanese encephalitis (JE) virus vaccine is used in Asia, and inactivated virus vaccines are available for JE and tick-borne encephalitis virus. The need for a vaccine against the DEN viruses is mounting, and, despite much effort, the goal of developing a safe and efficacious DEN virus vaccine has yet to be attained. An effective DEN virus vaccine must confer protection from each serotype, because all four serotypes commonly circulate in endemic regions and secondary infection with a heterologous serotype is associated with increased disease severity.

We have employed two strategies for generating live attenuated vaccine components against each serotype that can then be combined into tetravalent formulations (Blaney J E et al. 2006 *Viral Immunol.* 19:10-32). First, reverse genetics has been used to introduce an attenuating 30 nucleotide deletion ( $\Delta$ 30) mutation into the 3'-UTR of (DNA clones of each DEN serotype (Durbin, A P et al. 2001 *Am J Trop Med Hyg* 65:405-413; Whitehead S S et al. 2003 *J Virol*

77:1653-1657; Blaney J E et al. 2004 *Am J Trop Med Hyg* 71:811-821; Blaney J E et al 2004 *BMC Inf Dis* 4:39). Initially, the rDEN4Δ30 vaccine component was found to be attenuated in rhesus monkeys (Table 1) and phase I clinical trials in humans have demonstrated that virus infection results in low viremia, is strongly immunogenic, and exhibits minimal reactogenicity with no observation of serious adverse events (Durbin, A. P. et al, 2001 *Am J Trop Med Hyg* 65:405-413; Durbin et al. 2005 *J Inf Dis* 191:710-718). Recently, the rDEN1Δ30 vaccine component, which was also attenuated in rhesus monkeys (Table 1), has been found to share a similar phenotype in clinical trials as that observed for rDEN4Δ30; low viremia, strong immunogenicity; and minimal reactogenicity in 20 volunteers (Whitehead S S et al. 2003 *J Virol* 77:1653-1657; Blaney J E et al. 2006 *Viral Immunol.* 19:10-32). Unfortunately, the rDEN2Δ30 and rDEN3Δ30 vaccine components did not appear to be satisfactorily attenuated in rhesus monkeys during pre-clinical testing and there is no plan to test these in humans (Table 1) (Blaney J E et al. 2004 *Am J Trop Med Hyg* 71:811-821; Blaney J E et al. 2004 *BMC Inf Dis* 4:39). Consequently, an alternative strategy for vaccine development has been generation of antigenic chimeric viruses by replacement of structural proteins of the attenuated rDEN4Δ30 vaccine component with those from DEN2 or DEN3 yielding the rDEN2/4Δ30 and rDEN3/4Δ30 vaccine components, respectively (Whitehead S S et al. 2003 *Vaccine* 21:4307-4316; Blaney J E et al. 2004 *Am Trop Med Hyg* 71:811-821). The rDEN2/4Δ30 vaccine virus has been tested in humans and appears safe and strongly immunogenic, while clinical evaluation of the rDEN3/4Δ30 virus is currently planned.

TABLE 1

Effects of the Δ30 mutation on the four DEN serotypes in rhesus monkeys					
Virus	% of viremic monkeys	Viremia <sup>a</sup>			Reference
		Mean no. of viremic days per monkey	Mean peak virus titer (log <sub>10</sub> PFU/ml) ±SE	Geometric mean neutralizing antibody titer <sup>b</sup>	
rDEN1	100	2.8	2.1 ± 0.1	1,230	Whitehead et al.
rDEN1Δ30	50	0.5	0.8 ± 0.1	780	<i>J. Virol.</i> , 2003, 77: 1653
rDEN2	100	4.0	1.9 ± 0.1	173	Blaney et al. <i>BMC Inf Dis.</i> , 2004, 4: 39
rDEN2Δ30	100	2.8	1.7 ± 0.2	91	
rDEN3	100	2.3	1.4 ± 0.2	363	Blaney et al. <i>Am. J. Trop. Med. Hyg.</i> , 2004 71: 811
rDEN3Δ30	100	2.0	1.5 ± 0.2	265	
rDEN4	100	3.0	2.2 ± 0.2	322	Hanley et al.
rDEN4Δ30	100	2.0	1.4 ± 0.2	154	<i>Vaccine</i> , 2004 22: 3440

<sup>a</sup>Groups of rhesus monkeys were inoculated subcutaneously with 5.0 log<sub>10</sub> PFU of the indicated virus in a 1 ml dose. Serum was collected daily for 10 days. Virus titer in serum was determined by plaque assay in Vero cells.

<sup>b</sup>Plaque reduction (60%) neutralizing antibody titers were determined on day 28 serum using indicated wild type virus. Reciprocal dilution of geometric mean is indicated.

Here, we describe novel vaccine components for the DEN3 serotype generated by genetic modification of the 3'-UTR of the DEN3 cDNA clone (Blaney J E et al. 2004 *Am J Trop Med Hyg* 71:811-821). Development of these DEN3 vaccine components, which possess the full complement of wild type DEN3 proteins, is important for two reasons. First, the present vaccine component for DEN3, rDEN3/4Δ30, may be found to be under- or over-attenuated in

clinical trials. Second, an optimal vaccine for conferring protection from disease caused by DEN3 may require induction of T cell responses against the entire set of DEN3 proteins, rather than just the M and E which are the only DEN3 sequences present in the rDEN3/4Δ30 chimeric virus. To generate additional DEN3 vaccine components, novel deletions which encompass or border the Δ30 deletion in the 3'-UTR were introduced into the rDEN3 cDNA clone. Alternatively, the 3'-UTR of the rDEN3 cDNA clone was replaced with that of rDEN4 or rDEN4Δ30. Viable viruses were analyzed for attenuation phenotypes in tissue culture, SCID mice transplanted with HuH-7 cells, and rhesus monkeys. Three mutant viruses (rDEN3Δ30/31, rDEN3Δ86, and rDEN3-3'D4Δ30) have preclinical phenotypes which suggest they may be safe and immunogenic in humans.

#### Generation of rDEN3 Deletion Mutants

We sought to generate expanded deletion mutations which include the original Δ30 (nt 173-143) mutation. Table 2 lists seven deletion mutations which encompass the original Δ30 mutation including Δ50, Δ61, Δ80, Δ86, Δ116A, Δ116B and Δ146. In addition, the Δ30/31 mutation includes the original Δ30 mutation and a non-contiguous 31 nt deletion. The Δ31 mutation was also generated alone to discern the contribution of either Δ30 or Δ31 in the combined Δ30/31 deletion mutation. The location of bordering nucleotides of deletions in the predicted secondary structure of the DEN3 3'-UTR are indicated in FIG. 4. In addition, the predicted secondary structure of the DEN3 3'-UTR for rDEN3Δ30, rDEN3Δ30/31, and rDEN3Δ86 are indicated in FIGS. 8, 12, and 16, respectively.

TABLE 2

Deletion mutations created in the 3'-UTR of DEN3 Sleman/78		
Mutation	Deleted nucleotides <sup>a</sup>	Deletion junction
Δ30	173-143	-CCAAAGACU-
Δ31	258-228	-CUGCAGACU-

TABLE 2-continued

Deletion mutations created in the 3'-UTR of DEN3 Sleman/78		
Mutation	Deleted nucleotides <sup>a</sup>	Deletion junction
Δ50	192-143	-CACAAAGACU-
Δ61	173-113	-CCGAAUAAA-
Δ80	192-113	-CACAAUAAA-
Δ86	228-143	-UAGCAGACU-
Δ116 (A)	228-113	-UAGCAUAAA-
Δ116 (B)	258-143	-CUGCAGACU-
Δ146	258-113	-CUGCADAAA-
Δ30/31	173-143	-CCAAAGACU-
	258-228	-CUGCAGACU-

<sup>a</sup>Numbering is from the 3'-end of viral genome

PCR mutagenesis was used to introduce the nine new deletion mutations into the DEN3 Sleman/78 eDNA plasmid, p3, which was previously used to generate the rDEN3Δ30 vaccine component (Blaney J E et al. 2004 *Am J Trop Med Hyg* 71:811-821). The p3-frag.4 cDNA subclone was used as the template for PCR reactions with indicated pairs of mutagenic oligonucleotides listed in Table 3, except for the Δ30/31 deletion mutation which used p3-frag.4Δ30 cDNA subclone as a template. PCR products were ligated and used to transform competent bacterial cells, Plasmid DNA was isolated from bacterial clones and the presence of the appropriate deletion mutation was confirmed by

TABLE 3-continued

Mutagenic primer sequences for construction of 3'-UTR deletions	
Primer name	Sequence (5'→3')
143F	GACTAGAGGTTAGAGGAGAC (SEQ ID NO: 25)
228F	GACTAGCGGTTAGAGGAGACCCC (SEQ ID NO: 26)
173R	TCGGGCCCCGCTGCTGCGTTG (SEQ ID NO: 27)
173R (for Δ30)	TTGGGCCCCGCTGCTGCGTTG (SEQ ID NO: 28)
192R	TGTGTCATGGGAGGGGTCCTC (SEQ ID NO: 29)
228R	GCTACACCGTGCGTACAGCTTCC (SEQ ID NO: 30)
258R	GCAGCCTCCCAGGTTTTACGTCC (SEQ ID NO: 31)

TABLE 4

Mutations in addition to the engineered deletions that were identified in the rDEN3Δ30/31 and rDEN3Δ86 viruses when compared to the DEN3 p3 plasmid cDNA (Genbank # AY656169)					
Virus	Gene	Nucleotide position	Nucleotide substitution	Amino acid position	Amino acid change
rDEN3Δ30/31	NS4B	7164 <sup>a</sup>	U→C	115	Val→Ala
	NS4B	7398	C→U	193	Ala→Val
rDEN3Δ86	M	512	A→G	26	Lys→Glu
	NS3	6076	C→U	521	silent
	NS4B	7164 <sup>a</sup>	U→C	115	Val→Ala
	NS5	8623	U→C	353	silent
	NS5	10267 <sup>b</sup>	A→U	END	end→Tyr
	3'-UTR	10455	G→C	-	-

<sup>a</sup>The 7164 mutation is a Vero cell adaptation mutation which was engineered into the cDNA construct.

<sup>b</sup>There is a mixed population at this nt position (A→A/U) that changes the stop codon (UAA) at the end of NS5 to UAU which encodes Tyr. This would serve to extend NS5 by 2 amino acids (Tyr-Thr-End) because a stop codon remains at nts 10271-10273.

sequence analysis. To generate intact DEN3 eDNA plasmids containing the deletion mutations, the KpnI-PstI fragment (963 nt) from the mutated p3-frag.4 eDNA subclones were introduced into the p3-7164 cDNA plasmid. The p3-7164 plasmid encodes the 7164 Vero cell adaptation mutation which had previously been shown to enhance growth and transfection efficiency in Vero cells (Blaney J E et al. 2004 *Am J Trop Med Hyg* 71:811-821). Full length p3 plasmids containing the deletion mutations were confirmed to contain the correct 3'-UTR sequence. Mutations in addition to the engineered deletions were identified in the rDEN3Δ30/31 and rDEN3Δ86 viruses when compared to the DEN3 p3 plasmid cDNA (Genbank #AY656169) (Table 4).

TABLE 3

Mutagenic primer sequences for construction of 3'-UTR deletions	
Primer name	Sequence (5'→3')
113F	TAAAAACAGCATATTGACGCTGGGAG (SEQ ID NO: 24)

For recovery of viruses, 5'-capped RNA transcripts were synthesized in vitro from cDNA plasmids and transfected into either Vero cells or C6/36 cells. Prior to transcription and generation of infectious virus, the linker sequences were removed from eDNA plasmids by digestion with SpeI. Plasmids were then recircularized by ligation, linearized with Acc651 (isoschizomer of KpnI which cleaves leaving only a single 3' nucleotide), and transcribed in vitro using SP6 polymerase. Purified transcripts were then transfected into Vero or C6/36 cells.

Recombinant viruses encoding each of the nine mutations, Δ30/31, Δ31, Δ50, MI, Δ80, Δ86, Δ116A, Δ146, were successfully recovered in C6/36 cells, while only rDENΔ31 was recovered in Vero cells. The rDEN3 deletion mutant viruses were then passaged once in Vero cells followed by biological cloning by two terminal dilutions in Vero cells. Cloned viruses were then passaged two to seven times in Vero cells in an attempt to reach a stock titer of at least 6.0 log<sub>10</sub> PFU/ml which is considered sufficient to allow for cost-effective manufacture. Three recombinant viruses (rDEN3Δ50, rDEN3Δ116A, and rDEN3Δ146) were found

to be excessively restricted for replication in Vero cells, despite being viable. Therefore, these three viruses were not studied further. The genetic sequence of the 3'-UTR was determined for the six remaining deletion mutant viruses that reached peak virus titers of at least  $6.0 \log_{10}$  PFU/ml. The correct 3'-UTR sequence with the appropriate deletion was found for rDEN3Δ61, rDEN3Δ80, rDEN3Δ86 and rDEN3Δ30/31. However, two mutant viruses were found to contain additional deletions or mutations and were deemed to potentially have unstable genotypes. First, rDEN3Δ31 had the correct 3'-UTR deletion of nt 258-228 but also contained a 25 nt deletion of nt 222-198. Second, rDEN3Δ116B had the correct 3'-UTR deletion ant 258-143 but also contained a 8 nt deletion of nt 430-423 and a single A->G substitution at nt 265. The potential of genetic instability with these viruses precludes their use as vaccine components so they were not further studied. Therefore, of the nine original deletions constructed, four mutant viruses were found to replicate efficiently in Vero cells and were studied further; rDEN3Δ61, rDEN3Δ80, rDEN3Δ86 and rDEN3Δ30/31.

Generation of rDEN3 Chimeric Viruses with the 3'-UTR Derived from rDEN4 or rDEN4Δ30

Another strategy was employed to generate novel rDEN3 vaccine components; replacement of the 3'-UTR of the rDEN3 cDNA clone with that of rDEN4 or rDEN4Δ30 (FIG. 18A). The 3'-UTR chimeric virus, rDEN3-3'D4Δ30, was designed to be a vaccine component for inclusion in tetravalent formulations which share the Δ30 deletion mutation among all four serotypes. The rDEN3-3'D4 virus was designed to discern the contribution of the 3'-UTR chimerization and the Δ30 mutation to any observed phenotypes.

The p3-3'D4Δ30 plasmid was generated as follows. First, PCR mutagenesis was used to introduce a HpaI restriction

make the full length cDNA clone, p3-3'D4Δ30. The sequence of the 3'-UTR and N55 junction were confirmed to be correct.

To generate p3-3'D4, the 30 deleted nucleotides of the Δ30 deletion mutation were introduced into the p3-frag.4-3'D4Δ30 subclone. Briefly, the kflul---KpnI fragment of p3-frag.4-3'D4Δ30, which encompasses the Δ30 region, was replaced with the corresponding fragment of p4 to make the plasmid, p3-frag.4-3'D4. To generate a full length p3 genome, the PstI-KpnI fragment of p3 was replaced with the corresponding fragment of p3-frag.4-3'D4. The 3'-UTR sequence of the p3-3'D4 plasmid was determined to be correct and contained the missing 30 nt of the Δ30 mutation.

For recovery of viruses, 5'-capped RNA transcripts were synthesized in vitro from cDNA plasmids and transfected into either Vero cells or C6/36 cells. Prior to transcription and generation of infectious virus, the linker sequences were removed from cDNA plasmids by digestion with SpeI. Plasmids were then recircularized by ligation, linearized with Acc651 (isoschizomer of KpnI which cleaves leaving only a single 3' nucleotide), and transcribed in vitro using SP6 polymerase. Purified transcripts were then transfected into Vero or C6/36 cells.

rDEN3-3'D4 was recovered in C6/36 cells and Vero cells, while rDEN3-3'D4Δ30 could only be recovered in Vero cells. Mutant viruses were then passaged once in Vero cells followed by biological cloning by two terminal dilutions in Vero cells. rDEN3-3'D4 and rDEN3-3'D4Δ30 were then passaged four or six times in Vero cells, respectively. The genetic sequence of the NS5 — 3'-UTR junction and 3'-15fR was found to be correct for rDEN3-3'D4 and rDEN3-3'D4Δ30. Therefore, both viruses were studied further.

Mutations were also identified in the rDEN3-3'D4Δ30 virus compared to the DEN3 p3 plasmid cDNA clone (5'-UTR and genes) and DEN4 p4 cDNA clone (3'-UTR) (Table 5).

TABLE 5

Mutations in the rDEN3-3D4Δ30 virus compared to the DEN3 p3 plasmid cDNA clone (5'-UTR and genes) and DEN4 p4 cDNA clone (3'-UTR)					
Virus	Gene	Nucleotide position	Nucleotide substitution	Amino acid position	Amino acid change
rDEN3-3'D4Δ30	C	250	U → C	52	silent
	NS3	5899	U → C	462	silent
	NS4B <sup>a</sup>	7164	U → C	115	Val → Ala
	3'-UTR	10534	A → G	—	—

<sup>a</sup>The 7164 mutation is a Vero cell adaptation mutation which was engineered into the cDNA construct.

site into the p3-frag.4 cDNA subclone (FIG. 18B). PCR products were ligated and used to transform competent bacterial Plasmid DNA was isolated from bacterial clones and the presence of the appropriate deletion mutation was confirmed by sequence analysis. To introduce the rDEN4Δ30 3'-UTR into the p3-frag.4(HpaI) cDNA subclone, a 364 nt fragment encompassing the p4Δ30 3'-UTR was amplified by PCR using a forward primer (5'-AACAAACAACAACACCAAAGGCTATTG-3', SEQ ID NO: 32) and reverse primer (5'-CC-TACCGGTACCAGAACCTGTG-3', SEQ ID NO: 33). To generate the p3-frag.4-3'D4Δ30 cDNA subclone, the HpaI-KpnI fragment was removed from p3-frag.4(HpaI) and replaced with the p4Δ30 3'-UTR PCR fragment which had been cleaved by KpnI. The PdyI-KpnI fragment of p3-frag.4-3'D4Δ30 was introduced into the p3 plasmid to

50 Replication of DEN3 Mutant Viruses in SCID-HuH-7 Mice

The four deletion mutant viruses (rDEN3Δ30/31, rDEN3Δ61, rDEN3Δ80, and rDEN3Δ86) which were found to replicate to high titer in Vero cells and were confirmed to have the correct 3'-UTR sequence and the rDEN3-3'D4 and rDEN3-3'D4Δ30 viruses were first evaluated in SCID-HuH-7 mice. The rDEN3-3'D4 and rDEN3-3'D4Δ30 were compared to determine the effect on replication of the 3'-UTR chimerization and any further attenuation conferred by the Δ30 mutation. SCID-HuH-7 mice contain solid tumors of the HuH-7 human hepatoma cell line, and analysis of virus replication in this mouse model serves as a surrogate for DEN virus replication in the human liver. Numerous DEN virus mutant viruses have been identified by evaluation in SCID-HuH-7 mice (Blaney J E et al. 2002 *Virology* 300:125-139; Hanley et al. 2004 *Vaccine* 22:3440-3448; Blaney J E et al. 2006 *Viral Immunol.* 19:10-32). This mouse

model provided the original evidence that the rDEN3Δ30 virus was not attenuated compared to parent virus rDEN3, while the antigenic chimeric virus, rDEN3/4Δ30, was approximately 100-fold restricted in replication in the SCID-Ilufl-7 mice when compared to wild type parent viruses (Blaney 1E et al. 2004 *Am J Trop Med Hyg* 71:811-821).

For analysis of virus replication in SCID-HuH-7 mice, four to six week-old SCID mice (Tac:ler:Ha(ICR)-Prkdc<sup>scid</sup>) (Taconic Farms) were injected intraperitoneally with 0.1 mL phosphate-buffered saline containing 10<sup>7</sup> HuH-7 cells which had been propagated in tissue culture. Tumors were detected in the peritoneum five to six weeks after transplantation, and tumor-bearing mice were infected by direct inoculation into the tumor with 10<sup>4</sup> FEU of virus in 50 μl Opti-MEM I. Serum was collected from infected mice on day 7 post-infection and frozen at -80° C. The virus titer was determined by plaque assay in Vero cells.

As indicated in Table 6, wild type DEN3 Sleman/78 replicated to a mean peak virus titer of nearly 10<sup>6.9</sup> PFU/ml. Although a decreased level of replication was observed for each of the six mutant viruses, the differences in replication were not statistically significant. However, rDEN3Δ86 and rDEN3-3'D4Δ30 were more than 10-fold restricted in replication compared to wild type DEN3 virus, while the replication of rDEN3Δ30/3 was slightly less than 10-fold restricted. On the basis of this arbitrary cut-off, these three viruses were selected for further evaluation. It is important to note that the rDEN4LΔ30 virus which has a well-characterized, attenuation and non-reactogenic phenotype in humans was found to be only 6-fold restricted in replication in SCID-HuH-7 mice compared to wild type rDEN4 virus (Hanley et al. 2004 *Vaccine* 22:3440-3448).

TABLE 6

Replication of mutant DEN3 viruses in HuH-7-SCID mice.		
Virus <sup>1</sup>	Mean peak No. of virus titer mice (log <sub>10</sub> pfu/ml ± SE)	Fold-reduction compared to DEN3 (Sleman/78)
DEN3 (Sleman/78)	8 6.9 ± 0.1	—
rDEN3Δ30/31	8 6.0 ± 0.3	8
rDEN3Δ61	9 6.3 ± 0.2	4
rDEN3Δ80	9 6.3 ± 0.3	4
rDEN3Δ86	10 5.6 ± 0.4	20
rDEN3-3'D4	11 6.5 ± 0.4	3
rDEN3-3'D4Δ30	9 5.7 ± 0.2	16

<sup>1</sup>Groups of HuH-7-SCID mice were inoculated into the tumor with 4.0 log<sub>10</sub> PFU of the indicated virus. Serum was collected on day 7 and virus titer was determined in Vero cells.

Because the rDEN3-3'D4Δ30 virus and the rDEN3Δ30/31 and rDEN3Δ86 viruses encode the full set of DEN3 structural and non-structural proteins, they would be expected to induce the full complement of humoral and cellular immunity. This more complete immune induction would be advantageous compared to that induced by the chimeric rDEN3/4,Δ30, which encodes only the structural proteins of DEN3.

#### Replication of DEN3 Mutant Viruses in Tissue Culture

The level of virus replication in Vero cells and mosquito C6/36 cells was assessed for the rDEN3Δ30/31 and rDEN3Δ86 deletion mutant viruses and the rDEN3-3'D4 viruses with and without Δ30. Replication in Vero cells was analyzed because these cells are the substrate for manufacture, while growth in C6/36 cells was assessed because attenuation phenotypes in these mosquito cells may be associated with restricted replication in *Aedes* mosquitoes

which serve as the vector for DEN virus transmission (Hanley et al. 2003 *Virology* 312:222-232).

Growth kinetics were evaluated as follows. Confluent monolayers of Vero cells and C6/36 cells in 75 cm<sup>2</sup> flasks were infected at a multiplicity of infection of 0.01. Aliquots of 0.5 ml tissue culture supernatant were removed daily for seven days, combined with SPO stabilizer, and frozen at -80° C. Virus titer of all samples was determined by plaque assay in Vero cells. The limit of detection for the plaque assay is 1.0 log<sub>10</sub>PFU/ml.

The replication kinetics of each virus in both cell lines is shown in FIG. 19. In Vero cells, iDEN3Δ30/31, rDEN3Δ86, and rDEN3-3'D4Δ30 replicated to a peak level that approximated that of wild type rDEN3 and with similar kinetics to that of wild type rDEN3. These three vaccine components reached peak virus titers of 6.5 to 6.7 log<sub>10</sub> PFU/ml which demonstrates the feasibility of manufacture for each of these viruses. In Vero cells, the rDEN3-3'D4 virus replicated to a peak titer of 7.8 log<sub>10</sub>PFLU/ml which is nearly 100-fold higher than that observed for wild type rDEN3 indicating that inclusion of the DEN4 3'-UTR may augment replication in Vero cells. This could also be attributed to more efficient Vero cell adaptation of the rDEN3-3'D4 virus. rDEN4 replicates to a peak titer of approximately 8.0 log<sub>10</sub>PFU/ml which indicates that the chimeric virus achieves a peak titer that does not exceed that of either of its parent viruses (Blaney J E et al. 2006 *Viral Immunol.* 19:10-32).

Analysis of virus replication in C6/36 cells demonstrated that rDEN3Δ86 and rDEN3-3'D4Δ30 reached peak titers approximately 10-fold lower than the peak virus titer of wild type rDEN3 virus, 6.9 log<sub>10</sub>PFU/ml (FIG. 19). The rDEN3-3'D4 virus replicated to a peak titer similar to that observed for wild type rDEN3. The most striking result was the lack of replication observed in C6/36 cells for the rDEN3Δ30/31 virus. After day 1, virus was not detected in culture medium from C6/36 cells infected with rDEN3Δ30/31 virus despite the efficient replication observed in Vero cells. These results were confirmed in a second independent growth curve experiment and indicate a host range attenuation phenotype in tissue culture which is envisioned as being accompanied by an attenuation phenotype in mosquitoes as well.

#### Replication and Immunogenicity of DEN3 Mutant Viruses in Rhesus Monkeys

Based on the slight attenuation in SCID-HUH-7 mice and efficient growth in Vero cells, rDEN3Δ30/31, rDEN3Δ86, and rDEN3-3'D4Δ30 were evaluated in rhesus monkeys. The mutant viruses were compared with wild type DEN3 for level and duration of viremia, neutralizing antibody induction, and the ability to confer protection from wild type DEN3 virus challenge. The rDEN3-3'D4 virus was also evaluated to discern the contribution of the 3'-UTR chimerization upon attenuation with and without the Δ30 mutation. An attenuation phenotype in rhesus monkeys has generally been a strong predictor of safety for vaccine components in clinical trials including rDEN4Δ30, rDEN1Δ30, and rDEN2/4Δ30 (Blaney J E et al. 2006 *Viral Immunol.* 19:10-32).

Groups of four rhesus monkeys were inoculated subcutaneously with 10<sup>5</sup> PR of the indicated viruses (Table 7). Two monkeys were mock infected with virus diluent. For detection of viremia, serum was collected on days 0.8 and on day 10 and frozen at -80° C. Virus titer in serum samples was determined by plaque assay in Vero cells. Serum was collected on day 28 for detection of neutralizing antibodies directed against DEN3. Levels of neutralizing antibodies were determined using a plaque reduction neutralization

assay in Vero cells against wild type DEN3 virus. On day 35 post-infection, all monkeys were challenged by subcutaneous infection with 10<sup>5</sup> DEN3 wild type virus. Serum was collected on days 0-8 and on day 10 and frozen at -80° C. Virus titer in serum samples was determined by plaque assay in Vero cells.

TABLE 7

Replication and immunogenicity of rDEN3 mutant viruses in rhesus monkeys.								
Virus <sup>1</sup>	No. of monkeys	Mean		Geometric mean		Post-challenge <sup>4</sup>		
		% of monkeys with viremia	no. of days per monkey	Mean peak virus titer <sup>2</sup> (log <sub>10</sub> pfu/ml ± SE)	serum neutralizing antibody titer (reciprocal dilution) <sup>3</sup>	% of monkeys with viremia	Mean peak virus titer <sup>2</sup> (log <sub>10</sub> pfu/ml ± SE)	
DEN3 (Sleman/78)	4	100	3.5	1.8 ± 0.1	<5	253	0	<1.0
rDEN3Δ30/31	4	0	0	<1.0	<5	304	0	<1.0
rDEN3Δ86	4	0	0	<1.0	<5	224	0	<1.0
rDEN3-3'D4	4	75	1.5	1.3 ± 0.2	<5	229	0	<1.0
rDEN3-3'D4Δ30	4	0	0	<1.0	<5	77	0	<1.0
mock infected	2	0	0	<1.0	<5	<5	100	1.8 ± 0.2

<sup>1</sup>Groups of rhesus monkeys were inoculated subcutaneously on day 0 with 5.0 log<sub>10</sub> PFU of the indicated virus in a 1 ml dose. Serum was collected daily on days 0-8 and 10 and once on day 28.  
<sup>2</sup>Virus titer in serum was determined by plaque assay in Vero cells.  
<sup>3</sup>Plaque reduction (60%) neutralizing antibody titers were determined using DEN3 (Sleman/78).  
<sup>4</sup>Monkeys were challenged after 35 days with DEN3 (Sleman/78) administered subcutaneously in a 1 ml dose containing 5.0 log<sub>10</sub> PFU. Serum was collected daily on days 0-8 and 10.

Wild type DEN3 Sleman/78 virus replicated in rhesus monkeys to a mean peak virus titer of 1.8 log<sub>10</sub> PFU/ml serum with all monkeys developing viremia (Table 7). These results parallel previous studies of DEN3 in rhesus monkeys (Blaney J E et al. 2004 *Am J Trop Med Hyg* 71:811-821). No viremia was detected in any monkey infected with any of the three vaccine components, rDEN3Δ30/31, rDEN3Δ86, or rDEN3-3'D4Δ30 demonstrating a clear attenuation phenotype for each of these viruses in rhesus monkeys. Interestingly, the rDEN3-3'D4 virus was detected in 75% of monkeys with a mean peak virus titer of 1.3 log<sub>10</sub>PFU/ml serum suggesting that the presence of the Δ30 mutation is critical for attenuation of the 3'-UTR chimeric virus. Despite the lack of detectable viremia, mean neutralizing antibody levels in monkeys infected with rDEN3Δ30/31 and rDEN3Δ86 reached levels similar to that of wild type DEN3 virus, 1:253 (Table 7). In contrast, the rDEN3-3'D4Δ30 virus induced mean neutralizing antibody levels approximately three-fold lower than DEN3. However, 100% of monkeys immunized with each vaccine component seroconverted as defined by a four-fold or greater rise in serum neutralizing antibody levels after infection. Thus all monkeys were deemed to be infected by each of the vaccine components despite the lack of detectable viremia. Determination of virus titer in serum after challenge with DEN3 virus indicated that immunization with each of the vaccine components induced complete protection from detectable viremia as would be expected given the observed neutralizing antibody levels.

Replication in Mosquitoes

Replication of rDEN3 and rDEN3Δ30/31 was studied in *Toxorhynchites' ambainensis* mosquitoes. Intrathoracic inoculation of serial ten-fold dilutions of test virus was performed as described previously (Troyer J. M. e al, 2001 *Am. J. Trap. Med. Hyg.* 65:414-9). After a 14 day incubation, heads were separated and homogenized in diluent. Virus titer in head homogenates was determined by plaque assay in Vero cells.

Based on the attenuation of rDEN3Δ30/31 in rhesus monkeys and its restricted replication in C6/36 mosquito cells, rDEN3Δ30/31 was compared to wild type rDEN3 fbr infectivity and level of replication in highly sensitive *Toxorhynchites amboinensis* mosquitoes (Table 8). Ten-fold serial dilutions of virus were inoculated intrathoracically, and the

ability to infect head tissues was evaluated by performing a plaque assay on mosquito head homogenates after a 14 day incubation. The infectivity of rDEN3 and rDEN3Δ30/31 was very similar as the 50% mosquito infectious dose was approximately 10<sup>1-3</sup> PFU for both viruses (Table 8). However, the level of replication of rDEN3Δ30/31 in the heads of infected mosquitoes was about 5- to 50-fold reduced. This reduction was significant at the 10<sup>2-3</sup> and 10<sup>1-3</sup> PFU doses tested. This finding indicates that although rDEN3Δ30/31 has infectivity for *Toxorhynchites* by intrathoracic infection similar to that of wild type rDEN3, there is a statistically significant restriction in the level of replication in mosquitoes afforded by the Δ30/31 mutation.

TABLE 8

Replication of rDEN3 and rDEN3Δ30/31 in <i>Toxorhynchites amboinensis</i>					
Virus	Dose <sup>a</sup> (log <sub>10</sub> PFU)	No. tested		Mean virus titer <sup>c</sup> (log <sub>10</sub> PFU/head)	Reduction compared to same dose of wt virus (log <sub>10</sub> )
		infected <sup>b</sup>	not infected		
rDEN3 wt	2.3	20	90	4.2 ± 0.1 <sup>d</sup>	
	1.3	19	53	4.2 ± 0.1 <sup>e</sup>	
rDEN3Δ30/31	0.3	17	18	4.3 ± 0.3	
	2.3	12	83	2.7 ± 0.3 <sup>d</sup>	1.5
	1.3	16	44	3.1 ± 0.3 <sup>e</sup>	1.1
	0.3	8	13	3.6 ± 0.0	0.7

<sup>a</sup>Virus titer administered intrathoracically in a 0.2 μl inoculum.  
<sup>b</sup>Percentage of mosquitoes with detectable virus at day 14 post-inoculation was determined by plaque assay on mosquito head homogenates in Vero cells.  
<sup>c</sup>Calculated using only values of virus-positive heads.  
<sup>d</sup>For 10<sup>2.3</sup> PFU dose of rDEN3 and rDEN3Δ30/31, mean virus titers were significantly different as determined by a Tukey-Kramer post-hoc test (P < 0.001).  
<sup>e</sup>For 10<sup>1.3</sup> PFU dose of rDEN3 and rDEN3Δ30/31, mean virus titers were significantly different as determined by a Tukey-Kramer post-hoc test (P < 0.005).

While the present invention has been described in some detail for purposes of clarity and understanding, one skilled in the art will appreciate that various changes in form and detail can be made without departing from the true scope of the invention. All figures, tables, and appendices, as well as

patents, applications, and publications, referred to above, are hereby incorporated by reference.

## SEQUENCE LISTING

Sequence total quantity: 33

SEQ ID NO: 1 moltype = RNA length = 276  
 FEATURE Location/Qualifiers  
 source 1..276  
 mol\_type = unassigned RNA  
 organism = Dengue virus

SEQUENCE: 1  
 taaaacctgg gaggctgcaa actgtggaag ctgtacgcac ggtgtagcag actagcgggt 60  
 agaggagacc cctcccatga cacaaacgcag cagcggggcc cgagctctga ggaagctgt 120  
 acctccttgc aaaggactag aggttagagg agaccccccg caaataaaaa cagcatattg 180  
 acgctgggag agaccagaga tcctgctgtc tcctcagcat cattccaggc acagaacgcc 240  
 agaaaatgga atggtgctgt tgaatcaaca gtttct 276

SEQ ID NO: 2 moltype = RNA length = 278  
 FEATURE Location/Qualifiers  
 source 1..278  
 mol\_type = unassigned RNA  
 organism = Dengue virus

SEQUENCE: 2  
 taaaaccctg gaggctgcaa aaccatgga gctgtacgca tgggtagca gactagtgtg 60  
 tagaggagac ccctcccaag acacaacgca gcagcggggc ccaacaccag ggaagctgt 120  
 accctggtgg taaggactag aggttagagg agaccccccg cacacaaca aacagcatat 180  
 tgacgctggg agaccagaga gatcctgctg tcctcagcag atcattccag gcacagaacg 240  
 ccagaaaatg gaatggtgct gttgaaatca caggttct 278

SEQ ID NO: 3 moltype = RNA length = 281  
 FEATURE Location/Qualifiers  
 source 1..281  
 mol\_type = unassigned RNA  
 organism = Dengue virus

SEQUENCE: 3  
 taaaaaatcc gagggccac aaaccatgga agctgtacgc atggcgtagt ggactagcgg 60  
 ttagaggaga cccctccctt acagatcgca gcaacaatgg gggcccaagg tgagatgaag 120  
 ctgtagtctc actggaagga ctgagaggtta gaggagacc ccccaaaaca aaaaacagca 180  
 tattgacgct gggaaagacc agagatcctg ctgtctctc agcatcattc caggcacagg 240  
 acgccagaaa atggaatggt gctgttgaat caacaggttc t 281

SEQ ID NO: 4 moltype = RNA length = 276  
 FEATURE Location/Qualifiers  
 source 1..276  
 mol\_type = unassigned RNA  
 organism = Dengue virus

SEQUENCE: 4  
 taaaacctgg gaggctgcaa actgtggaag ctgtacgcac ggtgtagcag actagcgggt 60  
 agaggagacc cctcccatga cacaaacgcag cagcggggcc cgagctctga ggaagctgt 120  
 acctccttgc aaaggactag aggttagagg agaccccccg caaataaaaa cagcatattg 180  
 acgctgggag agaccagaga tcctgctgtc tcctcagcat cattccaggc acagaacgcc 240  
 agaaaatgga atggtgctgt tgaatcaaca gtttct 276

SEQ ID NO: 5 moltype = RNA length = 281  
 FEATURE Location/Qualifiers  
 source 1..281  
 mol\_type = unassigned RNA  
 organism = Dengue virus

SEQUENCE: 5  
 atccccaggg agggcctgcg ccacggaagc tgtacgcgtg gcatattgga ctagcgggta 60  
 gaggagacc cctcccatcag tgacaaaacg cagcaaaagg gggcccgaag ccaggaggaa 120  
 gctgtactcc tgggtggaag actagaggtt agaggagacc ccccaaacac aaaaacagca 180  
 tattgacgct gggaaagacc agagatcctg ctgtctctgc aacatcaatc caggcacaga 240  
 gcgccgaag atggaatggt gttgttgatc caacaggttc t 281

SEQ ID NO: 6 moltype = RNA length = 248  
 FEATURE Location/Qualifiers  
 misc\_feature 1..248  
 note = Dengue virus construct  
 source 1..248  
 mol\_type = other RNA  
 organism = synthetic construct

SEQUENCE: 6  
 taaaaccctg gaggctgcaa aaccatgga gctgtacgca tgggtagca gactagtgtg 60  
 tagaggagac ccctcccaag acacaacgca gcagcggggc ccaagactag aggttagagg 120  
 agaccccccg cacacaaca aacagcatat tgacgctggg agagaccaga gatcctgctg 180  
 tcctcagcag atcattccag gcacagaacg ccagaaaatg gaatggtgct gttgaaatca 240  
 caggttct 248

-continued

---

SEQ ID NO: 7                   moltype = RNA   length = 251  
FEATURE                        Location/Qualifiers  
misc\_feature                   1..251  
                                note = Dengue virus construct  
source                         1..251  
                                mol\_type = other RNA  
                                organism = synthetic construct

SEQUENCE: 7  
taaaaaatcc gggaggccac aaacctgga agctgtacgc atggcgtagt ggactagcgg 60  
ttagaggaga cccctccctt acagatcgca gcaacaatgg gggcccaaga ctagaggtta 120  
gaggagacc ccccaaaaca aaaaacagca tattgacgct gggaaagacc agagatcctg 180  
ctgtctcttc agcatcattc caggcacagg acgccagaaa atggaatggt gctggtgaat 240  
caacaggttc t 251

SEQ ID NO: 8                   moltype = RNA   length = 245  
FEATURE                        Location/Qualifiers  
misc\_feature                   1..245  
                                note = Dengue virus construct  
source                         1..245  
                                mol\_type = other RNA  
                                organism = synthetic construct

SEQUENCE: 8  
taaaaactgg gaggctgcaa actgtggaag ctgtacgcac ggtgtagcag actagcggtt 60  
agaggagacc cctcccatga cacaaacgag cagcggggcc caagactaga ggtagaggga 120  
gacccccgc aaataaaaaa agcatattga cgctgggaga gaccagagat cctgctgtct 180  
cctcagcatc attccaggca cagaacgcca gaaaatggaa tgggtgctgt gaatcaacag 240  
gttct 245

SEQ ID NO: 9                   moltype = RNA   length = 250  
FEATURE                        Location/Qualifiers  
misc\_feature                   1..250  
                                note = Dengue virus construct  
source                         1..250  
                                mol\_type = other RNA  
                                organism = synthetic construct

SEQUENCE: 9  
atccccaggg aggccatgcg ccacggaagc tgtacgcgtg gcatattgga ctagcggtta 60  
gaggagacc cctcccatca tgacaaaaag cagcaaaaagg gggcccaaga ctagaggtta 120  
gaggagacc ccccaacaca aaaacagcat attgacgctg ggaagacca gagatcctgc 180  
tgtctctgca acatcaatcc aggcacagag cgccgcaaga tggattggtg ttgttgatcc 240  
aacaggttct 250

SEQ ID NO: 10                  moltype = RNA   length = 217  
FEATURE                        Location/Qualifiers  
misc\_feature                   1..217  
                                note = Dengue virus construct  
source                         1..217  
                                mol\_type = other RNA  
                                organism = synthetic construct

SEQUENCE: 10  
taaaaaccgg gaggctgca actagtgggt agaggagacc cctccaaga cacaaacgag 60  
cagcggggcc caagactaga ggtagaggga gacccccgc acaacaaca acagcatatt 120  
gacgctggga gagaccagag atcctgctgt ctctacagca tcattccagg cacagaacgc 180  
cagaaaatgg aatggtgctg ttgaatcaac aggttct 217

SEQ ID NO: 11                  moltype = RNA   length = 220  
FEATURE                        Location/Qualifiers  
misc\_feature                   1..220  
                                note = Dengue virus construct  
source                         1..220  
                                mol\_type = other RNA  
                                organism = synthetic construct

SEQUENCE: 11  
taaaaaatcc gggaggccac aaatagcggg tagaggagac ccctccctta cagatcgag 60  
caacaatggg ggccaagac tagaggttag aggagacccc ccaaaaaca aaaacagcat 120  
attgacgctg ggaagacca gagatcctgc tgtctctca gcatcattcc aggcacagga 180  
cgccagaaaa tggaatggtg ctgttgaatc aacaggttct 220

SEQ ID NO: 12                  moltype = RNA   length = 214  
FEATURE                        Location/Qualifiers  
misc\_feature                   1..214  
                                note = Dengue virus construct  
source                         1..214  
                                mol\_type = other RNA  
                                organism = synthetic construct

SEQUENCE: 12  
taaaaactgg gaggctgcca ctacggggtg gaggagacc ctcccatgac acaacgcagc 60  
agcggggccc aagactagag gtttagaggag accccccgca aataaaaaa gcatattgac 120

-continued

---

gctgggagag accagagatc ctgctgtctc ctcagcatca ttccaggcac agaacgccag 180  
 aaaatggaat ggtgctgttg aatcaacagg ttct 214

SEQ ID NO: 13           moltype = RNA length = 219  
 FEATURE                Location/Qualifiers  
 misc\_feature           1..219  
                           note = Dengue virus construct  
 source                  1..219  
                           mol\_type = other RNA  
                           organism = synthetic construct

SEQUENCE: 13  
 atccccaggg aggccatgcg ccacggtttag aggagacccc tcccatcact gacaaaacgc 60  
 agcaaaaagg ggcccaagac tagaggtttag aggagacccc cccaacacaa aaacagcata 120  
 ttgacgctgg gaaagaccag agatcctgct gtctctgcaa catcaatcca ggcacagagc 180  
 gccgcaagat ggattgtgtg tgttgatcca acaggttct 219

SEQ ID NO: 14           moltype = RNA length = 194  
 FEATURE                Location/Qualifiers  
 misc\_feature           1..194  
                           note = Dengue virus construct  
 source                  1..194  
                           mol\_type = other RNA  
                           organism = synthetic construct

SEQUENCE: 14  
 taaaaaccgg ggaggctgca aaccatgga gctgtacgca tgggtagca gactagaggt 60  
 tagaggagac ccccgcaaca acaacaaaca gcatattgac gctgggagag accagagatc 120  
 ctgctgtctc tacagcatca ttccaggcac agaacgccag aaaatggaat ggtgctgttg 180  
 aatcaacagg ttct 194

SEQ ID NO: 15           moltype = RNA length = 196  
 FEATURE                Location/Qualifiers  
 misc\_feature           1..196  
                           note = Dengue virus construct  
 source                  1..196  
                           mol\_type = other RNA  
                           organism = synthetic construct

SEQUENCE: 15  
 taaaaaatcc gggaggccac aaaccatgga agctgtacgc atggcgtagt ggagactaga 60  
 ggtagagaga gaccccccca aaacaaaaa cagcatattg acgctgggaa agaccagaga 120  
 tcctgctgtc tcctcagcat cattccaggc acaggacgcc agaaaatgga atggtgctgt 180  
 tgaatcaaca ggttct 196

SEQ ID NO: 16           moltype = RNA length = 190  
 FEATURE                Location/Qualifiers  
 misc\_feature           1..190  
                           note = Dengue virus construct  
 source                  1..190  
                           mol\_type = other RNA  
                           organism = synthetic construct

SEQUENCE: 16  
 taaaacctgg gaggctgcaa actgtggaag ctgtacgcac ggtgtagcga ctagaggta 60  
 gagagagacc cccgcaata aaaacagcat attgacgctg ggagagacca gagatcctgc 120  
 tgtctctca gcatcattcc aggcacagaa cgccagaaaa tggaatggtg ctgttgaatc 180  
 aacaggttct 190

SEQ ID NO: 17           moltype = RNA length = 195  
 FEATURE                Location/Qualifiers  
 misc\_feature           1..195  
                           note = Dengue virus construct  
 source                  1..195  
                           mol\_type = other RNA  
                           organism = synthetic construct

SEQUENCE: 17  
 atccccaggg aggccatgcg ccacggaagc tgtacgctg gcatattgga ctagactaga 60  
 ggtagagaga gaccccccca acacaaaaac agcatattga cgctgggaaa gaccagagat 120  
 cctgctgctc ctgcaacatc aatccaggca cagagcgccg caagatggat tgggtgtgtt 180  
 gatccaacag gttct 195

SEQ ID NO: 18           moltype = DNA length = 39  
 FEATURE                Location/Qualifiers  
 misc\_feature           1..39  
                           note = Junction region  
 source                  1..39  
                           mol\_type = other DNA  
                           organism = synthetic construct

SEQUENCE: 18  
 gagggagcca tttgtaaac gtaggagtg aaaaagagg 39

-continued

---

SEQ ID NO: 19	moltype = AA length = 5	
FEATURE	Location/Qualifiers	
REGION	1..5	
source	note = Junction region	
	1..5	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 19		5
EGAIW		
SEQ ID NO: 20	moltype = DNA length = 39	
FEATURE	Location/Qualifiers	
misc_feature	1..39	
source	note = Junction region	
	1..39	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 20		39
agtgaaggag ttctgtaatt accaacaaca aacacacaaa		
SEQ ID NO: 21	moltype = AA length = 5	
FEATURE	Location/Qualifiers	
REGION	1..5	
source	note = Junction region	
	1..5	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 21		5
SEGLV		
SEQ ID NO: 22	moltype = DNA length = 39	
FEATURE	Location/Qualifiers	
misc_feature	1..39	
source	note = Junction region	
	1..39	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 22		39
gaggagacca ttggtagtt aacaacaaca aacacacaaa		
SEQ ID NO: 23	moltype = AA length = 5	
FEATURE	Location/Qualifiers	
REGION	1..5	
source	note = Junction region	
	1..5	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 23		5
EGAIW		
SEQ ID NO: 24	moltype = DNA length = 26	
FEATURE	Location/Qualifiers	
misc_feature	1..26	
source	note = synthetic primer	
	1..26	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 24		26
taaaaacagc atattgacgc tgggag		
SEQ ID NO: 25	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
misc_feature	1..20	
source	note = synthetic primer	
	1..20	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 25		20
gactagaggt tagaggagac		
SEQ ID NO: 26	moltype = DNA length = 23	
FEATURE	Location/Qualifiers	
misc_feature	1..23	
source	note = Synthetic primer	
	1..23	
	mol_type = other DNA	
	organism = synthetic construct	

-continued

---

SEQUENCE: 26			
gactagcggg tagaggagac ccc			23
SEQ ID NO: 27	moltype = DNA	length = 21	
FEATURE	Location/Qualifiers		
misc_feature	1..21		
	note = synthetic primer		
source	1..21		
	mol_type = other DNA		
	organism = synthetic construct		
SEQUENCE: 27			
tcgggccccg ctgctgcggt g			21
SEQ ID NO: 28	moltype = DNA	length = 21	
FEATURE	Location/Qualifiers		
misc_feature	1..21		
	note = synthetic primer		
source	1..21		
	mol_type = other DNA		
	organism = synthetic construct		
SEQUENCE: 28			
ttgggccccg ctgctgcggt g			21
SEQ ID NO: 29	moltype = DNA	length = 20	
FEATURE	Location/Qualifiers		
misc_feature	1..20		
	note = synthetic primer		
source	1..20		
	mol_type = other DNA		
	organism = synthetic construct		
SEQUENCE: 29			
tgtgtcatgg gaggggtctc			20
SEQ ID NO: 30	moltype = DNA	length = 23	
FEATURE	Location/Qualifiers		
misc_feature	1..23		
	note = synthetic primer		
source	1..23		
	mol_type = other DNA		
	organism = synthetic construct		
SEQUENCE: 30			
gctacacggt gcgtacagct tcc			23
SEQ ID NO: 31	moltype = DNA	length = 23	
FEATURE	Location/Qualifiers		
misc_feature	1..23		
	note = synthetic primer		
source	1..23		
	mol_type = other DNA		
	organism = synthetic construct		
SEQUENCE: 31			
gcagcctccc aggttttacg tcc			23
SEQ ID NO: 32	moltype = DNA	length = 27	
FEATURE	Location/Qualifiers		
misc_feature	1..27		
	note = synthetic primer		
source	1..27		
	mol_type = other DNA		
	organism = synthetic construct		
SEQUENCE: 32			
aacaacaaca aacaccaaag gctattg			27
SEQ ID NO: 33	moltype = DNA	length = 23	
FEATURE	Location/Qualifiers		
misc_feature	1..23		
	note = synthetic primer		
source	1..23		
	mol_type = other DNA		
	organism = synthetic construct		
SEQUENCE: 33			
ectaccgta ccagaacctg ttg			23

---

We claim:

1. An immunogenic composition that is tetravalent for dengue serotypes 1, 2, 3, and 4, wherein the composition comprises:

a nucleic acid encoding a dengue 1 virus comprising a  $\Delta 30/31$  mutation, wherein the  $\Delta 30$  mutation deletes nucleotides from about 174 to about 145 of the dengue 1 3' untranslated region (UTR) and the  $\Delta 31$  mutation deletes nucleotides from about 258 to about 228 of the dengue 1 3' UTR, designated with reverse-order numbering system; and

a nucleic acid encoding a dengue 3 virus comprising a  $\Delta 30/31$  mutation, wherein the  $\Delta 30$  mutation deletes nucleotides from about 173 to about 143 of the dengue 3 3' UTR and the  $\Delta 31$  mutation deletes nucleotides from about 258 to about 228 of the dengue 3 3' UTR, designated with reverse-order numbering system.

2. A method of inducing an immune response to a dengue virus in a patient, comprising administering the immunogenic composition of claim 1 to a patient, thereby inducing the immune response.

3. The method of claim 2, wherein the patient is an adult or child at risk of dengue infection or a person infected with dengue virus.

\* \* \* \* \*